



(51) International Patent Classification:

A61K 31/436 (2006.01) **A61K 31/519** (2006.01)
A61K 31/337 (2006.01) **A61K 31/555** (2006.01)
A61K 31/404 (2006.01) **A61K 31/704** (2006.01)
A61K 31/407 (2006.01) **A61K 31/7068** (2006.01)
A61K 31/4745 (2006.01) **A61K 33/24** (2006.01)
A61K 31/513 (2006.01) **A61P 35/00** (2006.01)
A61K 31/517 (2006.01)

(21) International Application Number:

PCT/US2018/033482

(22) International Filing Date:

18 May 2018 (18.05.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/507,962 18 May 2017 (18.05.2017) US
 15/807,011 08 November 2017 (08.11.2017) US

(71) Applicant: **RENOVORX, INC.** [US/US]; 4546 El Camino Real, Suite 203, Los Altos, California 94022 (US).

(72) Inventors; and

(71) Applicants: **AGAH, Ramtin** [US/US]; 1011 Seymour Lane, Menlo Park, California 94025 (US). **NAJMABADI,**

Kamran [US/US]; 3114 David Avenue, Palo Alto, California 94303 (US). **BAGAI, Shaun R.** [US/US]; 2047 Cecelia Way, Mountain View, California 94040 (US).

(74) Agent: **MAI, Kassity L.** et al.; Cooley LLP, 1299 Pennsylvania Avenue, NW, Suite 700, Washington, District of Columbia 20004 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: METHODS FOR TREATING CANCEROUS TUMORS




FIG. 5

(57) Abstract: Apparatuses and methods described herein relate to treating cancerous tumors using radiation therapy and chemotherapy. In some embodiments, a method of treatment includes administering radiation therapy targeting a tumor, isolating a segment of a vessel proximate to the tumor, and administering a dose of a chemotherapeutic agent to the segment of the vessel. The method can further include waiting a period of time after administering the radiation therapy before administering the dose of the chemotherapeutic agent. In some embodiments, a catheter device including first and second occluding elements can be used to isolate the segment of the vessel.

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*

METHODS FOR TREATING CANCEROUS TUMORS

Cross Reference to Related Applications

[0001] This application claims priority to and the benefit of U.S. Provisional Patent Application No. 62/507,962, titled “Methods for Treating Cancerous Tumors,” filed May 18, 2017, and U.S. Nonprovisional Patent Application No. 15/807,011, titled “Methods for Treating Cancerous Tumors,” filed November 8, 2017, the disclosure of each of which is incorporated herein by reference in its entirety.

Background

[0002] Cancer begins when a cell begins dividing uncontrollably. Eventually, these cells form a visible mass or tumor. Solid tumors are masses of abnormal tissue that originate in organs or soft tissues that typically do not include fluid areas. Some examples of solid tumors include: pancreatic cancer, lung cancer, brain cancer, liver cancer, uterine cancer, and colon cancer.

[0003] Traditionally, tumors have been treated with surgical resection, radiation, and/or chemotherapy. Surgical resection involves the removal of tumor tissue. Radiation uses beams of intense energy to kill cancer cells and to shrink tumors. And chemotherapy involves the use of therapeutic agents or drugs to treat cancer. But surgical resection may not completely remove a tumor. Radiation and chemotherapy can have undesirable systemic side effects, including extreme fatigue, hair loss, infection, nausea and vomiting, and others that limit their usefulness. More recently, direct activation of the patient’s immune system to attack cancerous cells has shown promise in treating certain solid tumors, but not all. Thus, the need for an improvement in both the safety and the efficacy of current therapy still exists.

[0004] Use of localized intra-arterial therapies, including trans-arterial chemo-delivery (TAC) or trans-arterial chemo-embolization (TACE), has been shown to be clinically beneficial for a certain subset of solid tumors. TAC or TACE can involve imaging an organ having a tumor using angiography, isolating a branch of the artery that feeds the tumor or portion of the organ containing the tumor, and then locally injecting chemotherapy in a bolus fashion via the isolated artery. Localized intra-arterial therapies allow higher drug concentration to reach the tumor, overcoming the problem of poor blood flow to tumor mass in comparison to healthy tissue. Furthermore, localized intra-arterial therapies can also take advantage of the first pass effect of chemotherapeutics by generating higher level drug concentrations at the tumor cell

membrane and therefore enhancing cellular drug uptake as compared to non-localized infusion. Lastly, local delivery can reduce systemic side effects of chemotherapy.

[0005] One of the limitations of TAC and TACE is the need for selective cannulation and isolation of the tumor feeder vessel or arterial branch that can target the smallest portion of the organ containing the tumor. But it may be difficult to target and limit drug delivery to a small portion of the organ containing the tumor while achieving desired efficacy levels with the cancer treatment. On the one hand, limiting drug delivery to a small portion of the organ can reduce the potential impact of the administered drug on surrounding healthy tissue. But on the other hand, when the isolated region becomes too small, drug uptake levels by the tumor may decrease and reduce the efficacy of the cancer treatment. Given these limitations, a method to deliver a sufficient dose of a chemotherapeutic drug in addition to and independent of the need to cannulate and isolate to a specific feeding/supplying branch of a tumor feeder vessel is highly desirable.

Pancreatic Cancer

[0006] In 2016, pancreatic cancer ranked as the fourth leading cause of cancer death in the United States, and the tenth most commonly diagnosed tumor type in men and women. Estimates of incidence and deaths caused by pancreatic cancer are approximately 53,070 and 41,780, respectively (American Cancer Society: Cancer Facts and Figures, American Cancer Society, 2016). Projections based on the changing demographics of the United States population and changes in incidence and death rates reveal that, unless earlier diagnosis is made possible or better treatment options become available, pancreatic cancer is anticipated to move from the fourth to the second leading cause of cancer death in the United States by 2020.

[0007] Systemic chemotherapy as treatment for pancreatic cancer may be modestly effective due to low drug penetration in the pancreas because a drug infused systemically only moderately penetrates the pancreas, which may generally increase toxicity within a patient's body but not have an effect on the cancer. In many instances, tumors located in the pancreas are located in tissue surrounding an artery but not in a region of an artery that can be targeted and isolated. Accordingly, it may be difficult for a biologic agent or drug to reach and treat the tumors. Among solid tumors, drug delivery to pancreatic tumors is especially difficult due to the hypo-vascular and poorly perfused nature of the pancreas. The unique environment of the pancreas lends itself to reduced drug levels within the organ tissue, which reduces the effectiveness of systemic chemotherapy that relies on a functional vasculature for delivery to

tumor cells. Also, the effect of chemotherapy is concentration dependent, and systemic infusion oftentimes results in low concentrations. Aside from dosing limitations in treating pancreatic cancer, many systemic side effects of chemotherapeutic agents can result from the treatment.

[0008] In an attempt to increase the effectiveness of chemotherapeutic agents on pancreatic tumors while decreasing systemic toxicity, various researchers have delivered drugs directly to the pancreas using traditional endovascular catheters. These initial attempts have been limited due to the redundant nature of blood supply to the pancreas and its adjacent organs. Non-selective engagement of the pancreatic vessels can also lead to the wash through of chemotherapy to other adjacent organs. Most of the arterial branches to the pancreas are small; thus, selective engagement of these small branches via conventional catheters is difficult. Thus, there is a need to address these and other deficiencies.

Lung Cancer

[0009] Lung cancer is another deadly cancer that is difficult to treat. Lung cancer is responsible for 23% of total cancer deaths. Long-term exposure to tobacco smoke causes 80 to 90% of lung cancers. Nonsmokers account for 10 to 15% of lung cancer cases, and these cases are often attributed to a combination of genetic factors or other environmental exposures (Vogl, T.J., *et al.*, *Seminars in Interventional Radiology*, 2013, 30(2):176-184).

[0010] Common treatments for lung cancer depend on the cancer's specific pathology, staging, and the patient's performance status (*e.g.*, ability to breath). Traditional treatment options are surgery, chemotherapy, immunotherapy, radiation therapy, and palliative care. Intravascular techniques for localized delivery of chemotherapeutic agents have also been used to treat lung cancer, and include cancer therapy such as arterial chemoembolization, bronchial artery infusion (BAI), isolated lung perfusion (ILP), and lung suffusion. Chemotherapeutics approved for the treatment of non-small cell lung cancer in the United States include methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, afatinib dimaleate, everolimus, alectinib, pemetrexed disodium, atezolizumab, bevacizumab, carboplatin, ceritinib, crizotinib, ramucirumab, docetaxel, erlotinib hydrochloride, gefitinib, afatinib dimaleate, gemcitabine hydrochloride, pembrolizumab, mechlorethamine hydrochloride, methotrexate, vinorelbine tartrate, necitumumab, nivolumab, paclitaxel, ramucirumab, and osimertinib, and the combinations carboplatin-taxol and gemcitabine-cisplatin (<https://www.cancer.gov/aboutcancer>). Drugs approved for the treatment of small cell lung

cancer include methotrexate, everolimus, doxorubicin hydrochloride, etoposide phosphate, topotecan hydrochloride, mechlorethamine hydrochloride, and topotecan (<https://www.cancer.gov/aboutcancer>). Lung cancer such as small cell lung cancer can sometimes be treated with a combination of radiation therapy and one or more chemotherapeutics. But other types of lung cancer such as non-small cell lung cancer may not be sensitive to current chemotherapeutics. In many instances, current treatment methods are not effective at providing meaningful treatment or palliative care. Thus, it is desirable to have a more effective method for treating lung cancer tumors.

Brain Cancer

[0011] Malignant gliomas comprise up to 80% of primary malignant brain tumors in the adults. Among these, glioblastomas are the most deadly and account for 82% of all malignant gliomas (Suryadevra, C.M., *et al.*, *Surg. Neurol. Int.*, 2015, 6(1):S68-S77). The current standard of care includes surgical resection, followed by adjuvant external beam radiation and chemotherapy with drugs such as temozolomide. Conventional therapy is nonspecific and often results in a tragic destruction of healthy brain tissue. These treatments can be incapacitating and produce a median overall survival of just twelve to fifteen months. In addition, the invasive properties of glioblastomas make complete resection difficult and the glioblastomas may recur following initial treatment. Malignant gliomas are also highly vascularized tumors, and their unique capacities for regulating angiogenesis contribute to their resistance against known therapies.

[0012] Malignant gliomas, including glioblastoma multiforme, have been treated with inter-arterial chemotherapy. Typically, a catheter is inserted in the femoral artery and ends in the carotid artery, while a separate microcatheter is also inserted into the femoral artery and used to explore the specific vessels feeding the tumor for administration of the chemotherapy (Burkhardt, J-K., *et al.*, *Interventional Radiology*, 2011, 17:286-295). But such methods are not always effective and can be improved.

Liver Cancer

[0013] Liver cancer is another difficult-to-treat cancer characterized by solid tumors. In 2016, an estimated 39,230 adults (28,410 men and 10,820 women) in the United States will be diagnosed with primary liver cancer. Liver cancer also commonly metastasizes to other parts of the body. It is estimated that 27,170 deaths (18,280 men and 8,890 women) from this disease will occur this year. Liver cancer is the tenth most common cancer and the fifth most common cause of cancer death among men. It is also the eighth most common cause of cancer death

among women (American Cancer Society: Cancer Facts and Figures, American Cancer Society, 2016). When compared with the United States, liver cancer is much more common in developing countries within Africa and East Asia. In some countries, it is the most common cancer type. The one-year survival rate for people with liver cancer is 44%. The five-year survival rate is 17%. For the 43% of people who are diagnosed at an early stage, the five-year survival rate is 31%, while it is only 11% if the cancer has spread to surrounding tissues or organs and/or the regional lymph nodes. If the cancer has spread to a distant part of the body, the 5-year survival rate is only 3% (<http://www.cancer.net/cancer-types/liver-cancer/statistics>).

[0014] Currently, patients with hepatocellular carcinoma and cirrhosis are frequently treated with non-specific trans-arterial therapy using techniques that deliver treatments directly into the liver (Lewandowski, R. J., *et al.*, *Radiology*, 2011, 259(3):641-657). Physicians use the femoral artery to gain access to the hepatic artery, one of two blood vessels that feed the liver. Trans-arterial therapy such as TACE involves delivery of chemotherapy directly to the liver, followed by a process to embolize the chemotherapy. In this therapy, a thick, oily substance (for example, Lipiodol) is mixed with chemotherapy (for example, floxuridine, sorafenib tosylate or a mixture of platinol, mitomycin, and adriamycin) and injected under radiological guidance directly into the artery supplying the tumor via a catheter. The Lipiodol, or other particles, helps to contain the chemotherapy within the tumor and blocks further blood flow, thus cutting off the tumor's food and oxygen supply. TACE with doxorubicin-filled beads delivers the beads directly to the liver, which releases chemotherapy slowly over time and also blocks the blood flow to the tumor. In a similar therapy, radioactive yttrium beads are delivered via a catheter into the hepatic artery. The beads deliver radiation to the tumor, which kills the tumor cells, although other unintended areas of the liver may also receive radiation, creating undesirable destruction of healthy tissue. Thus, there is a need to improve current treatment methods.

Uterine Cancer

[0015] In 2016, an estimated 60,050 women in the United States were diagnosed with uterine endometrial cancer, with an estimated 10,470 deaths occurring (<http://www.cancer.net/cancer-types/uterine-cancer/statistics>). Uterine cancer is the fourth most common cancer for women in the United States. The incidence of endometrial cancer is rising, mainly due to a rise in obesity, which is an important risk factor for this disease. It is the sixth most common cause of cancer death among women in the United States with the 5-year survival rate being 82%.

[0016] Concurrent chemoradiotherapy (CCRT) is the main treatment for locally advanced cervical cancer. Neoadjuvant chemotherapy (NAC) was widely employed until CCRT became the standard, and conflicting results have been reported. Neoadjuvant intra-arterial chemotherapy (IANAC) is another method for delivering NAC as an alternative to systemic chemotherapy. IANAC has been reported to achieve beneficial results that cannot be obtained by systemic chemotherapy or CCRT. Kawaguchi *et al.* have reported that IANAC with cisplatin followed by radical hysterectomy or radiotherapy afforded similar results to concurrent chemoradiotherapy for stage IIIB cervical cancer (Kawaguchi *et al.*, *World Journal of Oncology*, 2013, 4(6):221-229). Drugs approved for use in the United States for the treatment of cervical cancer include bevacizumab, bleomycin, and topotecan hydrochloride, and the combination gemcitabine-cisplatin. Uterine cancer of endometrial origin may be treated with, for example, megestrol acetate. But many systemic side effects of chemotherapeutic agents can result from current treatment methods. It is desirable to have a specific means of targeting uterine tumors.

Colon Cancer

[0017] In the United States, colorectal cancer is the fourth most common cancer diagnosed each year for all adults combined. Separately, it is the third most common cancer in men and third most common cancer in women. In 2016, an estimated 134,490 adults in the United States were diagnosed with colorectal cancer, with 95,270 new cases of colon cancer and 39,220 new cases of rectal cancer. It is estimated that 49,190 deaths (26,020 men and 23,170 women) were attributed to colon or rectal cancer in 2016. Colorectal cancer is the second leading cause of cancer death in the United States, although when it is detected early, it can often be cured. The death rate from this type of cancer has been declining since the mid-1980s, probably because of an improvement in early diagnosis. The 5-year survival rate colorectal cancer is 65%, while the 10-year survival rate is 58% (<http://www.cancer.net/node/18707>).

[0018] When possible, surgical removal of colorectal tumors is the treatment of choice as it can eliminate the cancer completely. However, metastasis to other organs, particularly the liver and the lung, is common and complicates the treatment of colon and rectal cancer dramatically. It is therefore desirable to have a method of treating metastasized colon and rectal cancers that are present in other organs of the body. Drugs approved for use in treating colon cancer in the United States include bevacizumab, irinotecan hydrochloride, capecitabine, cetuximab, ramucirumab, oxaliplatin, 5-FU, fluorouracil, leucovorin calcium, trifluridine, tipiracil

hydrochloride, oxaliplatin, panitumumab, ramucirumab, regorafenib, ziv-aflibercept and the combinations capox, folfiri-bevacizumab, folfiri-cetuximab, FU-LV, xeliri and xelox.

Summary of the Invention

[0019] Apparatuses and methods are described herein that relate to, for example, the treatment of cancerous tumors. In some embodiments, the method comprises: a) first administering a course of radiation therapy targeting an area including a solid tumor; b) second waiting a period of time for the radiation to take effect on the microvasculature in the area; and c) third administering a therapeutically effective dose of a chemotherapeutic agent to an isolated arterial section near the solid tumor.

[0020] In some embodiments, the method comprises: a) first administering a targeted dose of radiation to an area including a solid tumor; b) second waiting a period of time; c) third isolating an area containing a cancerous tumor by, for example, isolating an arterial segment proximate to the tumor; and d) fourth administering a localized therapeutically effective dose of a chemotherapeutic agent.

[0021] In some embodiments, the method comprises: a) administering a course of radiation therapy to an area including a solid tumor; b) isolating the proximal and the distal part of the vasculature closest to the tumor to produce an isolated arterial segment; c) decreasing the intraluminal pressure of the isolated arterial segment to the level of the interstitium; and d) administering a therapeutically effective dose of a chemotherapeutic drug. In one embodiment, the method comprises an additional step of waiting a period of time following the step of administering the course of radiation therapy.

[0022] In some embodiments, the method includes delivering radiation therapy to a target area including a tumor; and inserting a catheter device into an artery where the catheter device includes a first occlusion member, a second occlusion member, and a body defining a lumen in fluid communication with an infusion port. The infusion port is disposed between the first occlusion member and the second occlusion member. The first occlusion member and the second occlusion member are moved to an area of the artery disposed proximate to the target area. The first occlusion member and the second occlusion member are deployed to isolate the area of the artery disposed proximate to the target area. A dose of chemotherapeutic agent is then delivered to the isolated area of the artery via the lumen and the infusion port. The

chemotherapeutic agent permeates to the target area including the tumor from the isolated area of the artery.

[0023] In some embodiments, the method includes administering a dose of radiation to a target area including a tumor; inserting a catheter device into a vessel, the catheter device including a first occlusion element and a second occlusion element; isolating a segment of the vessel proximate to the target area using the first occlusion element and the second occlusion element; and delivering a dose of an agent to the segment via the catheter device.

[0024] In some embodiments, the method includes administering a dose of radiation to a target area including a tumor; isolating a segment of the vessel proximate to the target area; adjusting an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area; and delivering a dose of an agent to the segment via the catheter device.

[0025] In some embodiments, use of a catheter device inserted into a vessel for delivering a dose of an agent to a segment of the vessel proximate to a target area including a tumor that has been administered a dose of radiation is disclosed. The catheter device can include a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

[0026] In some embodiments, use of a system including: (i) a radiation source for administration of a dose of radiation to a target area including a tumor; and (ii) a catheter device inserted into a vessel for delivering a dose of an agent to a segment of the vessel proximate to the target area that has been administered the dose of radiation, is disclosed. The catheter device can include a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

[0027] In some embodiments, use of a catheter device for delivering a dose of an agent to a segment of a vessel proximate a target area including a tumor that has been administered a dose of radiation is disclosed. The catheter device can isolate the segment of the vessel proximate to the target area and adjust an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area.

[0028] In some embodiments, a catheter device inserted into a vessel for use in delivering a dose of an agent to a segment of the vessel proximate to a target area including a tumor that

has been administered a dose of radiation is disclosed. The catheter device can include a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

[0029] In some embodiments, a system comprising: (i) a radiation source for use in administration of a dose of radiation to a target area including a tumor; and (ii) a catheter device inserted into a vessel for use in delivering a dose of an agent to a segment of the vessel proximate to the target area that has been administered the dose of radiation, is disclosed. The catheter device can include a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

[0030] In some embodiments, a catheter device for use in delivering a dose of an agent to a segment of a vessel proximate a target area including a tumor that has been administered a dose of radiation is disclosed. The catheter device can isolate the segment of the vessel proximate to the target area and adjust an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area.

[0031] In some embodiments, a chemotherapeutic agent comprising one or more compounds selected from any of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, fluorouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate, for use in treatment of a tumor, is disclosed. The chemotherapeutic agent can be administered subsequent to a course of radiation therapy applied to tissue surrounding the tumor after a period of time sufficient for the radiation therapy to reduce or devascularize the tissue.

[0032] In some embodiments, a chemotherapeutic agent comprising one or more compounds selected from any of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, fluorouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate, for use in treatment of a tumor, is disclosed. The chemotherapeutic agent can be

administered to a vessel proximate to tissue surrounding the tumor after administering a course of radiation therapy to the tissue.

[0033] In some embodiments, a method includes administering a course of radiation to tissue surrounding a tumor; inserting a catheter device into a vessel, the catheter device including at least one occlusion element; isolating a segment of the vessel proximate to the tissue using the at least one occlusion element; and delivering a chemotherapeutic agent to the segment via the catheter device.

[0034] In some embodiments, a method includes administering, at a first time, a dose of radiation to a target tissue including a tumor; and delivering, at a second time, a dose of a chemotherapeutic agent to the target tissue, the second time being after the first time by a period of time, the dose of radiation being effective to reduce the microvasculature within the target tissue during the period of time.

[0035] Other objects of the invention may be apparent to one skilled in the art upon reading the following specification and claims.

Brief Description of the Drawings

[0036] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[0037] FIG. 1 is an illustration of a catheter device disposed within a vessel, according to an embodiment.

[0038] FIG. 2 is a flowchart illustrating a method for treating a cancerous tumor, according to embodiments described herein.

[0039] FIG. 3 is a graph showing a change in pressure (mmHg) in a vessel over time while undergoing treatment, according to an embodiment.

[0040] FIG. 4A is schematic illustration of a catheter device shown in a dilated configuration disposed within a vessel, according to an embodiment.

[0041] FIG. 4B is a schematic illustration of dispersal of an infused agent into tissue surrounding a vessel, according to an embodiment.

[0042] FIG. 5 is a flowchart illustrating a method for treating a cancerous tumor, according to embodiments described herein.

[0043] FIG. 6A is an illustration of dispersal of an infused agent into tissue surrounding a vessel without application of radiation therapy, and FIG. 6B is an illustration of dispersal of an infused agent into tissue surrounding a vessel with application of radiation therapy, according to embodiments described herein.

[0044] FIG. 7 is an image of a pancreatic tumor after undergoing treatment according to methods described herein.

[0045] FIG. 8 is an image showing penetration of infused agents into tissue surrounding a vessel via the microvasculature, according to an embodiment.

[0046] FIG. 9 is a graph comparing survival rates of patients treated according to different methods described herein.

[0047] FIG. 10 is a bar chart comparing survival rates of patients treated according to different methods described herein.

[0048] FIG. 11 is an image of a catheter device disposed in a vessel in a patient's groin area, according to an embodiment.

[0049] FIG. 12 is an image showing penetration of an infused agent into tissue surrounding a vessel after undergoing treatment, according to an embodiment.

[0050] FIG. 13 is a flowchart illustrating a method for treating a tumor, according to embodiments described herein.

[0051] FIG. 14 is a schematic illustration of radiation being administered to target tissue including a tumor, according to embodiments described herein.

[0052] FIG. 15 is a schematic illustration of a dose of chemotherapeutic agent being delivered to target tissue including a tumor, according to embodiments described herein.

Detailed Description of the Invention

[0053] This application is not limited to particular methodologies or the specific compositions described. It is also understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present application will be limited only by the appended claims and their equivalents.

[0054] Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present application, the preferred methods and materials are now described.

[0055] As used herein and in the appended claims, the singular forms “a,” “and,” and “the,” include plural referents unless the context clearly dictates otherwise. Thus, for example, the term “chemotherapeutic” is intended to mean a single chemotherapeutic or a combination of chemotherapeutics; “a course of radiation therapy” is intended to mean one or more courses of radiation therapies, or combinations thereof; the term “agent” is intended to mean a single agent or a combination of agents, and so on and so forth.

[0056] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0057] The words “proximal” and “distal” refer to direction closer to and away from, respectively, an operator (*e.g.*, surgeon, physician, nurse, technician, etc.) who would insert the medical device into the patient, with the tip-end (*i.e.*, distal end) of the device inserted inside a patient’s body first. Thus, for example, the implant end first inserted inside the patient’s body would be the distal end of the implant, while the implant end to last enter the patient’s body would be the proximal end of the implant.

[0058] “Treat”, “treating” and “treatment” of cancerous tumors refer to reducing the frequency of symptoms of cancer (including eliminating them entirely), avoiding the occurrence of cancer, and/or reducing the severity of symptoms of cancer.

[0059] “Therapeutically effective amount” and “therapeutically effective dose” means the amount or dosage of a compound that, when administered to a patient for treating cancerous tumors, is sufficient to effect such treatment. The “therapeutically effective amount” or “therapeutically effective dose” will vary depending on, for example, the compound, the size of the tumor, and the age, weight, etc., of the patient to be treated.

Overview of the Invention

[0060] The present application provides a method for treating or ameliorating solid cancerous tumors, wherein a course of targeted radiation therapy is first administered to an area including one or more tumors. A period of time is allowed to elapse in order for the radiation to take

effect in down-sizing the tumor(s). The radiation may also reduce the microvasculature in the tissue in the area including the tumor(s). This period is followed by the administration of a therapeutically effective amount of a chemotherapeutic agent to an isolated arterial section near the solid tumor. Isolation of the arterial section may be accomplished by isolating the proximal and the distal part of the vasculature closest to the tumor whereby the intraluminal pressure is then decreased to the level of the interstitium. The therapeutically effective dose of the chemotherapeutic agent may then be administered via infusion. Combination of radiation therapy followed by properly administered chemotherapy is complementary and has a synergistic clinical effect when combined.

[0061] Intra-arterial delivery of chemotherapy, including TAC and TACE, has been shown to be effective and safe in treatment of certain solid tumors. A prerequisite for effective TAC or TACE is the selective engagement of nearby arterial vessels and, more commonly, the vessels feeding the tumor itself. The precise engagement of the feeding or branch vessel remains a major limitation for expanding the use of TACE and TAC in solid tumors, including but not limited to, pancreatic adenocarcinoma. The isolation of the artery supplying the tumor or the relevant tissue can be a technical challenge for a number of reasons, for example: a) there are organs with no dedicated single blood vessel supplying those specific organs; b) side and terminal branches of an artery can cause collateral flow to tissues and organs beyond the area of interest; and c) the tumor feeder vessels may be too small for detection by angiography; and d) the feeding branch/artery cannot be cannulated.

[0062] To address these problems, methods disclosed herein may involve administering radiation therapy to an area including a tumor. The radiation may reduce the microvasculature in the tissue in the area including the tumor. After the radiation therapy, the proximal and the distal part of the vasculature (*e.g.*, an artery) closest to the tumor is isolated using a double balloon catheter. Both the side and the terminal branches are excluded, which prevents drug washout. The reduced microvasculature in the tissue in the area also reduces drug washout. Upon inflation of both balloons in the isolated arterial segment, the intra-luminal pressure is reduced to the level of interstitium (typically, 10-20 mmHg). A therapeutic agent such as, for example, a chemotherapeutic drug, can be infused into the isolated arterial segment. The infusion of the chemotherapeutic drug in the isolated region, without any major runoff, leads to an increase in the intra-luminal pressure of at least about 30 mmHg in the isolated vessel segment. The pressure gradient forces the infused agent to traverse the arterial wall and enter the surrounding tissue, especially the vasa vasorum surrounding the vessel wall, with

subsequent influx of the therapeutic agent into the tissue. This technique is referred to herein as “trans-arterial micro-perfusion” or TAMP.

[0063] According to certain embodiments described herein, TAMP is not dependent on angiographic identification and cannulization of the tumor arterial supply or feeding vessels and thus overcomes deficiencies of current techniques. In TAMP, the drug traverses the arterial wall (*e.g.*, endothelium and media) before entering into the adventitia and interstitium. The interstitial concentration achieved is dependent on both the influx of the drug into the tissue across the artery wall and the efflux of the drug out of the interstitium via capillaries in the tissue area and the venous system. Hence, one can increase localized tissue concentration by both increasing the influx and reducing the efflux using the approach described above. The infusion parameters that determine the influx of the drug via TAMP include, but are not limited to, the intraluminal pressure achieved between the balloons, the intraluminal drug concentration, and the duration of infusion. By varying these parameters, one can change the drug influx and interstitial concentration.

[0064] In some embodiments, catheter devices such as those described in U.S. Patent Application No. 14/293,603, filed June 2, 2014, titled “Devices, methods and kits for delivery of therapeutic materials to a target artery,” now issued as U.S. Patent No. 9,457,171, and U.S. Patent Application No. 14/958,428, filed December 3, 2015, titled “Occlusion catheter system and methods of use,” the disclosures of which are incorporated herein by reference, can be used and/or adapted for use with TAMP techniques described herein. FIG. 1 depicts an example catheter device 100. The catheter device 100 includes a first occlusion element 102 and a second occlusion element 104. The occlusion elements 102, 104 can be any suitable devices or mechanisms that are configured to selectively limit, block, obstruct, or otherwise occlude a bodily lumen (*e.g.*, artery) in which the occlusion elements 102, 104 are disposed. For example, in some embodiments, the occlusion elements 102, 104 can be inflatable balloons or the like that can be transitioned between a collapsed (*e.g.*, deflated) configuration and an expanded (*e.g.*, inflated) configuration. The first occlusion element 102 can be coupled to a distal end portion of a first catheter, and the second occlusion element 104 can be coupled to the distal end portion of a second catheter. Alternatively, in some embodiments, the first occlusion element 102 and the second occlusion element 104 can be coupled to a single catheter at different points along the catheter. The catheter device 100 can be used to isolate a segment 120 of a bodily lumen (*e.g.*, artery) within the space defined between the first occlusion element 102 and the second occlusion element 104. After the segment 120 is isolated, a procedure can

be performed within the isolated segment 120 such as, for example, delivering a therapeutic agent to the isolated segment 120 and surrounding tissue 110.

[0065] While the catheter device 100 is depicted as having two occlusion elements 102, 104, other catheter devices that can be used with methods disclosed herein can include a single occlusion element or more than two occlusion elements (e.g., three occlusion elements), as needed to isolate a portion of a bodily lumen. For example, a catheter device with a single occlusion element can be used to isolate a portion of a vessel that is adjacent to or near a closed or terminating end of the vessel. Alternatively, a catheter device with three or more occlusion elements can be used to isolate a segment of a vessel that splits into one or more branches.

[0066] FIG. 2 illustrates a method 200 for performing a TAMP procedure. The method includes introducing a catheter (e.g., the catheter device 100) into a mammalian body into a bodily lumen (e.g., artery), at 202. The catheter can be advanced to a target area, at 204, and used to isolate the target area, at 206. In some embodiments, the catheter can include two occlusion members (e.g., occlusion elements 102, 104) that can be deployed (e.g., inflated) to isolate a segment of the bodily lumen to exclude the segment from its side and terminal branches. For example, a first occlusion member (e.g., a distal occlusion element) can be inflated, at 207, and a second occlusion member can be inflated (e.g., a proximal occlusion element), at 208. After the occlusion elements are deployed, an agent can be injected through an injection port of the catheter device to the isolated segment disposed between the two occlusion members, at 210. In some embodiments, the catheter can include a single occlusion member, and the single occlusion member can be used to isolate a segment of a bodily lumen such as a portion of a vessel near a terminating end or a dead-end. In some embodiments, a contrast dye can be injected into the isolated segment and the surrounding area can be visualized to determine whether the segment has been correctly isolated. For example, the injection of contrast through the infusion port can ensure that no extra vessels or bodily lumens are included in the isolated area. If desired, the catheter can be moved and the procedure repeated until the clinician can confirm that the catheter is correctly positioned. After the positioning of the catheter is confirmed, a therapeutic cell/biologic/agent can be introduced to the isolated segment through the infusion port.

[0067] FIG. 3 graphically illustrates how pressure (mmHg) in a bodily lumen (e.g., artery) changes over time as a TAMP procedure is performed (e.g., method 200). As shown in FIG. 3, the pressure in the bodily lumen drops when a first balloon or occlusion element is inflated and

continues to drop until a second balloon or occlusion element is inflated. The pressure then increases when an agent (*e.g.*, a contrast dye, a therapeutic agent) is infused into the segment isolated by the first balloon and the second balloon.

[0068] FIGS. 4A and 4B schematically depict an example of a catheter device 300 disposed within a bodily lumen 310 (*e.g.*, artery) and the dispersal of an infused agent 360 through the bodily lumen 310 into surrounding tissue. According to methods described herein (*e.g.*, method 200), the infused agent 360 can be injected into an isolated segment 320 and allowed to infuse into the surrounding tissue via, for example, a concentration gradient. As shown in FIG. 4B, the infused agent 360 can infuse through a wall 312 of the bodily lumen 310 into the surrounding tissue. As shown, the concentration of the agent 360 decreases as the distance (shown in millimeters (mm)) from the isolated segment 320 of the bodily lumen 310 increases.

[0069] In combination with the techniques described above, if one can decrease the tissue efflux of the chemotherapeutic drug, the drug concentration near an isolated segment of a bodily lumen may be advantageously increased. When a tumor is located in this region, the increased concentration can increase the effect of the chemotherapeutic drug on the tumor. One technique that can decrease tissue efflux is to radiate the tissue prior to treatment. Radiation can decrease tissue microvasculature in tissue containing cancerous tumors. Thus, combining prior radiation to decrease tissue microvasculature with TAMP can have a synergistic effect. Combining the steps of radiation of the cancerous tissue prior to the treatment, waiting two or more weeks for the microvasculature to decrease, followed by use of the TAMP technique to deliver chemotherapy in the isolated segment of the bodily lumen (*e.g.*, artery) closest to the tumor, produces a synergistic effect that the use of the TAMP technique alone does not.

[0070] Methods described herein can be used to treat solid cancerous tumors arising from any organ of the body where the tumor has its own or a proximate blood supply provided by a bodily lumen (*e.g.*, artery) that can be isolated. Examples of cancers that can be treated using methods described herein can be, but are not limited to, pancreatic cancer, lung cancer, liver cancer, uterine cancer, colon cancer, or brain cancer.

[0071] For example, apparatuses and methods described herein can be used to isolate a targeted region in a patient's pancreas. Studies have shown that a course of radiation prior to TAMP treatment has significant clinical benefit in patients with locally advanced pancreatic cancer. Combining these two modalities led to a significant increase in median survival, a reduction of tumor markers, and downsizing of the tumor. A similar combination therapy administered by

methods described herein may have clinical benefit in solid tumors in other organs and tissue areas where TAMP may be considered as a treatment option. Such tumors include, but are not limited to, pancreatic tumors, lung tumors, brain tumors, liver tumors, uterine tumors, and colon tumors.

[0072] In some embodiments, a method of treating a cancerous tumor can involve: first administering a course of radiation therapy targeting tissue including a solid cancerous tumor; second waiting a period of time for the destructive effect of the radiation on the vasculature to take effect; and third administering a therapeutically effective dose of a chemotherapeutic agent to an isolated section of a bodily lumen near the solid tumor. The targeted solid tumor can be, for example, a pancreatic tumor, a lung tumor, a brain tumor, a liver tumor, a uterine tumor, or a colon tumor. The administration of radiation on the targeted tissue area can include, for example, delivering approximately 20 to 50 Gy of radiation over approximately one to five weeks in approximately one to 25 sessions. The period of time between administration of the radiation therapy and administration of the chemotherapeutic agent can be selected to maximize the devascularization of the tissue surrounding the tumor. Depending on various factors including the specific course of radiation and the specific tissue area or organ, this period of time can be, for example, approximately one to six months, as short as two weeks, or as long as six months. Examples of suitable chemotherapeutic agents include doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, fluorouracil, folfinirix, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozitrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate or a combination of these drugs. In some embodiments, the section of the bodily lumen near the cancerous tumor can be isolated by the use of a catheter device to deliver the chemotherapeutic agent. In some embodiments, the catheter device can be used to increase the intraluminal pressure in the isolated section of the bodily lumen to achieve increased tissue penetration.

[0073] In some embodiments, a method of treating a cancerous tumor can involve: first administering a targeted dose of radiation to tissue including a solid tumor; second waiting a period of time; third isolating an area containing a cancerous tumor; and fourth administering a localized therapeutically effective dose of a chemotherapeutic agent. Similar to other methods described herein, the targeted solid tumor may be, for example, a pancreatic tumor, a lung tumor, a brain tumor, a liver tumor, a uterine tumor, or a colon tumor. The administration of radiation on the targeted tissue area can include, for example, delivering approximately 20 to

50 Gy of radiation over approximately one to five weeks in approximately one to 25 sessions. The period of time between administration of the radiation therapy and administration of the chemotherapeutic agent can be selected to maximize the devascularization of the tissue surrounding the tumor. Depending on various factors including the specific course of radiation and the specific tissue area or organ, this period of time can be, for example, at least a month. The isolated area can be, for example, an artery that is in proximity to the tumor. In some embodiments, a catheter device can be used to isolate the area. The catheter device can be used to increase the intraluminal pressure in the isolated artery. The isolated area can be, for example, the area of tissue involving the tumor. Examples of suitable chemotherapeutic agents include doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, fluorouracil, folfirinix, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate or a combination of these drugs.

[0074] In some embodiments, a method of treating a cancerous tumor can involve: administering a course of radiation therapy to tissue including a solid tumor; isolating the proximal and the distal part of the vasculature closest to the tumor to produce an isolated arterial segment; decreasing the intraluminal pressure of the isolated arterial segment to the level of the interstitium; and administering a therapeutically effective dose of a chemotherapeutic drug. The course of the radiation therapy can decrease tissue efflux of the chemotherapeutic drug. In some embodiments, the vasculature can be isolated using a double balloon catheter positioned to exclude both the side and terminal branches of the artery. The chemotherapeutic drug can pass across the artery wall and into the surrounding tissue via a pressure gradient generated by the increase in the intraluminal pressure above the interstitial pressure. In some embodiments, the method can additionally include waiting a period of time following the step of administering the course of radiation therapy. The period of time between administration of the radiation therapy and administration of the chemotherapeutic agent can be selected to maximize the devascularization of the tissue surrounding the tumor. For example, depending on various factors including the specific course of administering the radiation therapy and the specific tissue region, this period of time can be at least two weeks. The targeted solid tumor can be, for example, a pancreatic tumor, a lung tumor, a brain tumor, a liver tumor, a uterine tumor, or a colon tumor. The chemotherapeutic drug can be, for example, a single chemotherapeutic or a combination of chemotherapeutic drugs.

[0075] FIG. 5 is a flowchart illustrating a method 500 of treating a tumor involving the use of radiation. In particular, the method involves administering a course of radiation therapy to a target area, at 502. For example, an amount of radiation (*e.g.*, 20-50 Gy) can be administered to a patient in multiple sessions (*e.g.* 1-25 sessions) over a period of time (*e.g.*, a few days to six months). The target area can be a tissue area including a tumor. The method then optionally includes waiting a period of time for the radiation therapy to devascularize the tissue in the target area, at 504.

[0076] The method 500 further includes introducing a catheter (*e.g.*, the catheter device 100) into a mammalian body into a bodily lumen (*e.g.*, artery), at 506. The catheter can be advanced to a target area, at 508, and used to isolate the target area, at 510. In some embodiments, the catheter can include two occlusion members (*e.g.*, occlusion elements 102, 104) that can be deployed (*e.g.*, inflated) to isolate a segment of the bodily lumen to exclude the segment from its side and terminal branches. After the occlusion elements are deployed, an agent can be injected through an injection port of the catheter device to the isolated segment disposed between the two occlusion members, at 512. In some embodiments, a contrast dye can be injected into the isolated segment and the surrounding area can be visualized to determine whether the segment has been correctly isolated. For example, the injection of contrast through the infusion port can ensure that no extra vessels or bodily lumens are included in the isolated area. If desired, the catheter can be moved and the procedure repeated until the clinician can confirm that the catheter is correctly positioned. After the positioning of the catheter is confirmed, a therapeutic cell/biologic/agent can be introduced to the isolated segment through the infusion port.

[0077] In some embodiments, the step of administering the radiation therapy (502) can occur during and/or after the steps of introducing the catheter into the mammalian body (506), advancing the catheter to the target area (508), isolating the target area (510), and/or injecting a therapeutic agent into the target area (512). In some embodiments, one or more steps of the method 500 can be repeated before, during, and/or after other steps of the method 500.

[0078] FIGS. 6A and 6B schematically illustrate the effects of radiation on the vasa vasorum microvasculature in tissue surrounding an isolated segment 620 of a bodily lumen 610. By reducing the microvasculature, the radiation therapy reduces drug washout and increases drug tissue concentration when a drug is delivered to the area using methods described herein, such as, for example, TAMP. FIG. 6A depicts an area of tissue surrounding an isolated segment 620

of a bodily lumen 610 prior to radiation therapy. FIG. 6B depicts the area of tissue after radiation therapy. After the radiation, the number of microvasculature connections 632 (*e.g.*, micro-vessels extending from the isolated section 620 to the venous system 630) is reduced, thereby allowing a greater concentration of an infused drug 660 to remain in the tissue area.

[0079] As depicted in FIGS. 6A and 6B, a catheter device 600 can be used to deliver the infused drug 660 to the target area. The catheter device 600 can be similar to other catheter devices described herein (*e.g.*, catheter device 100 and catheter device 300). For example, the catheter device 600 has a first occlusion element 602 and a second occlusion element 604, which are coupled to distal end portions of a first catheter 601 and a second catheter 603, respectively. The catheter device 600 also includes a port 605 for delivering the infused drug 660 to the isolated segment 620 between the first occlusion element 602 and the second occlusion element 604. Once the infused drug 660 is delivered to the isolated segment 620, it can pass through a wall 612 of the bodily lumen 610 into surrounding tissue.

Radiation Therapy

[0080] In methods described herein, radiation therapy can include, for example, external-beam radiation therapy delivered by X-rays, gamma rays, proton beams, or other appropriate sources. Radiation therapy damages cells by destroying the genetic material that controls how cells grow and divide. While both healthy and cancerous cells are damaged by radiation therapy, the goal of radiation therapy is to destroy as few normal, healthy cells as possible. The radiation therapy described herein can be targeted as narrowly as possible to the solid tumor(s) being treated or the tissue closely surrounding the solid tumor(s).

[0081] Typically, a radiation treatment plan is individualized for a patient, based upon detailed imaging scans showing the location of a patient's tumor(s) and the normal areas around it. The amount of radiation that normal tissue in different parts of the body can safely receive is known to one skilled in the art. Computed tomography (CT) scans are most frequently employed, but magnetic resonance imaging (MRI), positron emission tomography (PET), and ultrasound scans may also be used. A radiation oncologist determines the exact area that will be treated, the total radiation dose that will be delivered to the tumor, how much dose will be allowed for the normal tissues around the tumor, and the safest angles (paths) for radiation delivery. Radiation doses for cancer treatment are measured in Gy, which is a measure of the amount of radiation energy absorbed by one kilogram of human tissue. Different doses of radiation are needed to kill different types of cancer cells. Patients can receive external-beam

radiation therapy in daily treatment sessions over the course of several weeks. The number of treatment sessions depends on many factors, including the total radiation dose that will be given. For example, one dose, which constitutes a fraction of the total planned dose of radiation, can be given each day. In a different instance, two treatments a day can be given.

[0082] As will be appreciated by one skilled in the art, the course of radiation therapy appropriate for use in the method of the present invention will depend on the specific cancerous tumor being treated. The specific dose of radiation, the duration of the radiation, and the number of treatments for any particular individual will depend upon a variety of factors including the type of cancer, the size of the tumor(s), and the patient's age and medical history including, for example, the amount of radiation previously received. Concurrent chemotherapy may also impact the dose of radiation given.

[0083] When treating a pancreatic cancer, for example, the course of radiation therapy can be approximately 20 to 50 Gy of radiation delivered in approximately one to 25 treatments over approximately one to five weeks. Alternatively, two to five sessions of radiation can be given over a period of approximately a week. For certain types of cancer, the amount of radiation therapy delivered may be as low as one Gy. In preferred embodiments, the course of radiation therapy can be approximately 40 to 50 Gy of radiation delivered in approximately 22 to 25 treatments over approximately four to five weeks. As may be appreciated by one skilled in the art, the amount of radiation therapy useful in methods described herein is that necessary to devascularize the solid tumor of interest thus allowing the TAMP technique to be used advantageously.

[0084] In methods described herein, after administering the radiation therapy, a physician may wait for a period of time before administering chemotherapy (e.g., a chemotherapeutic agent) such that the tumorous tissue can die (e.g., necrosis) or become devascularized. For example, a chemotherapeutic agent for use in treatment of a tumor can be administered subsequent to a course of radiation therapy applied to tissue surrounding the tumor after a period of time sufficient for the radiation therapy to reduce or devascularize the tissue. Alternatively or additionally, a chemotherapeutic agent for use in treatment of a tumor can be administered to a vessel proximate to tissue surrounding the tumor after administering a course of radiation therapy to the tissue.

[0085] In some embodiments, this period of time can be selected to maximize devascularization of the solid tumor and/or tissue containing the solid tumor. In some

embodiments, this period of time can be selected to maximize the effect of the chemotherapy based on a sufficient amount of devascularization. For example, in some embodiments, the reduction in the microvasculature of the tissue can enhance the efficacy of the later-delivered chemotherapeutic agent by reducing drug washout. In certain instances, the time period that elapses before administering chemotherapy can be at least a month. In other instances, the period of time is approximately two weeks to six months.

Chemotherapeutics

[0086] Specific chemotherapeutics can be selected based on the particular solid cancerous tumor that is to be treated. For example, the following chemotherapeutic agents and others may be used in the treatment of pancreatic cancer: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, fluorouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate. In some embodiments, a combination of agents may be employed. For example, when treating pancreatic cancer, a combination of gemcitabine hydrochloride (Gemzar®) and paclitaxel albumin-stabilized nanoparticle formulation (Abraxane®) may be used.

[0087] The above-described chemotherapeutic agents are available from a variety of corporate sources licensed to provide such agents for human use. Generic formulations of non-proprietary chemotherapeutics are typically available from a variety of manufacturers. A list of these licensed suppliers is available from the U.S. Food and Drug Administration's "Approved Drug Products with Therapeutic Evaluations," commonly known as the "Orange Book" (<http://www.accessdata.fda.gov/scripts/cder/ob/>). Proprietary chemotherapeutics are typically available from one manufacturer, also identifiable in the Orange Book. For example, the corporate source for Gemzar® is Eli Lilly and Company (Indianapolis, IN) and Celgene Corporation (Summit, NJ) supplies Abraxane®.

[0088] Methods described herein can use an amount of chemotherapeutic agent that is known to be therapeutically effective at treating a tumor. For example, the amount of chemotherapeutic agent that is used can be based on the Prescribing Information for a particular chemotherapeutic drug. A physician can adjust the amount of the chemotherapeutic agent to an amount that is appropriate for use with the TAMP techniques described herein.

[0089] In methods described herein, a therapeutic agent (*e.g.*, chemotherapy drug) can be delivered via rapid infusion (*e.g.*, injected directly into an artery over a period of minutes, intravenous infusion (*e.g.*, through a drip or pump over a period of approximately 20 minutes to a few hours), or continuous infusion (*e.g.*, through a continuation infusion pump over a period of weeks to months). The infusion of the drug into an isolated space increases the intraluminal or interior pressure of the vessel to above the interstitial pressure of the surrounding tissue and the pressure gradient forces the drug across a vessel wall and into the surrounding tissue.

Catheter Device

[0090] In some embodiments, methods described herein can use a catheter device such as, for example, a double occlusion balloon catheter to isolate a segment of a bodily lumen (*e.g.*, artery) and allow infusion of a therapeutic agent (*e.g.*, chemotherapy drug) into the isolated segment between the balloon after they are inflated. For example, methods disclosed herein may use catheter devices such as those described in U.S. Patent Application No. 14/293,603, filed June 2, 2014, titled “Devices, methods and kits for delivery of therapeutic materials to a target artery,” now issued as U.S. Patent No. 9,457,171, and U.S. Patent Application No. 14/958,428, filed December 3, 2015, titled “Occlusion catheter system and methods of use,” which are incorporated herein by reference. Briefly, a catheter device suitable for isolating a section of a bodily lumen near a solid tumor includes, but is not limited to, features and functions such as, for example: (1) selective isolation of the targeted portion of the portion of the artery for targeted delivery of the therapeutic agent to the solid tumor; (2) an infusion port allowing first, injection of contrast into the isolated segment to allow direct visualization of the origin of the branches of the artery supplying the cancerous tissue, and second, introduction of chemotherapeutic drugs; and (3) a self-contained assembly unit with easy retrieval after completion of the procedure. In one embodiment, the catheter device includes expandable occlusion elements in the form of inflatable balloons that can be used to isolate a proximal and distal end of a bodily lumen of interest.

[0091] Methods described herein can include, for example, introducing a catheter device into a splenic artery of the pancreas. The catheter device can have, for example, two lumens—one for inflation/deployment of the balloons/occluding elements and a second for introduction of the infusate (*e.g.*, therapeutic agent) to the space between the two balloons. The catheter can be advanced to a target portion of the splenic artery. A region of the target portion of the

splenic artery is selectively isolated and the infusate is injected into the isolated region. In some embodiments, the method can include advancing at least a portion of the catheter device to an ostium of a celiac artery, its hepatic branch (and its branches), or if necessary, the superior mesenteric artery, depending on a patient's anatomy. In some embodiments, a contrast dye is injected into the isolated region to confirm exclusion of side branches before injecting the infusate.

[0092] In some embodiments, the catheter device can have one or more features to achieve a desired effect on a specific anatomy of tumors. For example, there may be: (1) a separate inflation lumen for the proximal and the distal occluders/balloons to allow different size occluders/balloons proximally and distally; (2) slidable catheters to allow the distance between the occluders/balloons to be adjusted; and (3) a sensor at the tip to monitor pressure in the isolated segment of the bodily lumen.

[0093] In some embodiments, a catheter device inserted into a vessel can be used for delivering a dose of an agent to a segment of the vessel proximate to a target area including a tumor that has been administered a dose of radiation. The catheter device can include a first occlusion element and a second occlusion element for isolating the segment of the vessel proximate to the target area. In some embodiments, the catheter device can include a single occlusion element, where the single occlusion element can isolate a segment of a vessel adjacent to or near a closed or terminating end of the vessel.

[0094] In some embodiments, a system comprising: (i) a radiation source for administration of a dose of radiation to a target area including a tumor; and (ii) a catheter device inserted into a vessel, can be used for delivering a dose of an agent to a segment of the vessel proximate to the target area that has been administered the dose of radiation. The catheter device can include a first occlusion element and a second occlusion element for isolating the segment of the vessel proximate to the target area.

[0095] In some embodiments, a catheter device can be used for delivering a dose of an agent to a segment of a vessel proximate a target area including a tumor that has been administered a dose of radiation. The catheter device can isolate the segment of the vessel proximate to the target area and adjust an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area.

[0096] FIGS. 4A and 4B schematically depict an example of a catheter device 300 disposed within a bodily lumen 310 (e.g., artery) and the dispersal of an infused substance 360 through

the bodily lumen 310 into surrounding tissue. The catheter device 300 can be similar to other catheter devices (*e.g.*, catheter device 100) described herein. For example, catheter device 300 includes a first occlusion element 302 and a second occlusion element 304 for occluding a portion 320 of bodily lumen 310. The first occlusion element 302 is coupled to a distal end portion of a first catheter 301, and the second occlusion element 304 is coupled to a distal end portion of a second catheter 303. The occlusion elements 302, 304 are filter elements that can be moved between a collapsed configuration for insertion of the catheter device 300 into a body of a patient (*e.g.*, into an artery) and an expanded or dilated configuration, as shown in FIGS. 4A and 4B, for occluding a portion of a bodily lumen. The occlusion elements 302, 304 when in the collapsed configuration have a smaller outer perimeter (or diameter) than when in the expanded configuration.

[0097] As depicted in FIGS. 4A and 4B, the catheter device 300 can be used to isolate a segment 320 of a bodily lumen 310 within the space defined between the first occlusion element 302 and the second occlusion element 304. The catheter device 300 can include a lumen in fluid communication with port or opening 305 for delivering an agent 360 (*e.g.*, a dye or a chemotherapy drug) to the space between the first occlusion element 302 and the second occlusion element 304. The first catheter 301 can define the lumen and the opening 305. The opening 305 can be disposed on the distal end portion of the first catheter 301 distal to the first occlusion element 302. The second catheter 303 can be movably disposed within a lumen defined by the first catheter 301 such that the second catheter 303 can be moved relative to the first catheter 301 to move the second occlusion element 304 relative to the first occlusion element 302. According to some embodiments of the disclosure, the second occlusion element 304 can be moved toward the first occlusion element 302 to increase pressure within the isolated segment 320. The increased pressure can be used, for example, to drive delivery of the agent 360 through the wall 312 of the bodily lumen 310 and into the surrounding tissue.

[0098] In some embodiments, the catheter device 310 can have a sensor such as a pressure transducer 306 that may assist with achieving an optimal pressure within an occluded arterial segment for optimizing trans-arterial diffusion of an infused substance during a method of cancer treatment (*e.g.*, a TAMP procedure). The pressure transducer 306 may be disposed along the catheter device 300 in the isolated arterial segment 320 (*e.g.*, disposed between the first occlusion element 302 and the second occlusion element 304 (as depicted in FIGS. 4A and 4B)). The pressure transducer 306 can be disposed on one of the catheters 301, 303, or disposed on one of the occlusion elements 302, 304. The pressure transducer 306 can be designed to

measure an intraluminal pressure of the isolated segment 320. The pressure measurements may be used to adjust the intraluminal pressure of the isolated segment 320 to a predetermined or optimal pressure level. A physician may use the pressure measurements to determine a rate of infusing a drug or other therapeutic material into the isolated segment 320 in order to decrease or increase the intraluminal pressure of the isolated segment 320. For example, a physician can increase the rate of infusion of a drug to increase the intraluminal pressure of the isolated segment 320 above the pressure of tissue surrounding the isolated segment 320 (*e.g.*, above the pressure of the interstitium) to create a pressure gradient between the intraluminal space and the surrounding tissue to increase permeation of the infused drug through the arterial wall and into the tissue. Additionally or alternatively, a physician can increase or decrease the intraluminal pressure of the isolated segment 320 by adjusting the position of the two occlusion elements 302, 304 relative to one another (*e.g.*, moving the two occlusion elements 302, 304 closer or further apart from one another).

Determination of Therapeutic Effectiveness

[0099] The efficacy of the methods of the present invention in the treatment of solid cancerous tumors can be evaluated in human clinical trials conducted under appropriate standards and ethical guidelines as set forth by the U.S. Food and Drug Administration (FDA). Such studies are conducted according to U.S. and International Standards of Good Clinical Practice. Typically, such trials are comparison trials, in that the method of the present invention is utilized in one cohort of patients, while one or more other cohorts receive alternative methods of treating the tumors. The alternative methods can include, for example, treatment with systemic chemotherapy alone.

[0100] A clinical trial for the treatment of cancerous tumors may have a primary objective of evaluating survival in patients who undergo radiation therapy followed by intra-arterial delivery of a chemotherapeutic agent to an isolated arterial section near the solid tumor after a suitable interval of time elapses. The second objective of such a trial is to assess tumor response by known imaging techniques at the primary site of application of the chemotherapeutic agent. In particular, the size of the tumor before and after treatment can be determined and evaluated across different treatment methods. In addition, the conversion rate from unresectable or borderline resectable to potentially resectable or resectable tumors can be determined. The results may be analyzed using standard statistical techniques known to those skilled in the art.

[0101] The following examples, including clinical studies, are offered by way of illustration and not by way of limitation.

Examples: Experiments with Pig Tissue

[0102] FIG. 7 is an image 700 of a pancreatic tumor 710 of a pig after treatment with the TAMP method. The method involved occluding the celiac artery of the pig with a double balloon catheter, rapidly infusing dye at six milliliters per minute for ten minutes into the isolated segment of the celiac artery, and harvesting the tissue next to the celiac artery. As depicted, the infused dye has permeated into the harvested tissue.

[0103] FIG. 8 is an image 800 of tissue surrounding the celiac artery of a pig after treatment with the TAMP method. The method involved occluding the celiac artery of the pig with a double balloon catheter (*i.e.*, a balloon catheter with occlusion elements or balloons 802) and rapidly infusing dye at six milliliters per minute for ten minutes into the isolated segment of the celiac artery. The image shows the tissue surrounding the celiac artery in situ within ten sections after initiating the rapid infusion of the dye. As depicted in FIG. 8, the dye 804 has penetrated into the surrounding tissue via the vasa vasorum microvasculature.

[0104] FIGS. 11 and 12 relate to an experiment conducted on a pig to evaluate the effect of radiation on tissue penetration of drugs/molecules using the TAMP technique. The experiment involved administering radiation treatment to a left groin area in a pig and comparing the penetration of a dye introduced using the TAMP technique in the left groin area (referred to herein as the test 1200) versus the penetration of the dye in the right groin area (referred to herein as the control 1210). A Yucatan pig was anesthetized for a CT scan of the femoral artery to plan the radiation treatment. Both femoral arteries were accessed for the placement of sterile gold fiducial markers to mark the areas of interest for comparison following the radiation treatment and infusion of dye. The left groin area was treated with a single radiation session using the CyberKnife® system at 15 Gy outside the artery adjacent to the gold marker. FIG. 11 depicts an image 1100 of the left groin area with the gold marker 1106. One month following radiation treatment, with the animal under general anesthesia, a percutaneous exposure of the left and right femoral artery was completed. From the left carotid artery, a double balloon catheter 1110 was advanced to the area of the left femoral artery and a proximal balloon 1102 and a distal balloon 1104 of the catheter 1110 were positioned and inflated adjacent to the gold fiducial marker 1106, as shown in FIG. 11. The catheter 1110 isolated the relevant segment of the left femoral artery adjacent to the gold fiducial marker 1106 to ensure exclusion of any

large side branches and to achieve optimal intravascular pressure in the isolated segment. A syringe pump was then used to inject a dye at six milliliters per minute for approximately 30 seconds through an infusion port between the two balloon catheters. The same procedure was repeated for the right femoral artery. At the conclusion of the procedures in the left and right femoral arteries, the area of dye penetration around the blood vessels were measured. As shown in FIG. 12, there was approximately a three-fold increase in penetration on the irradiated left side (*i.e.*, test 1200) as compared to the control right side (*i.e.*, control 1210).

Examples: Clinical Studies of Treating Pancreatic Cancer

[0105] A post-market registry study was conducted to assess patient survival and clinical outcomes using the RenovoCath™ RC120 catheter (RenovoRx, Los Altos, CA) in a clinical, prospective observational setting when used to deliver a chemotherapeutic to the pancreas as described in Table 1.

Table 1

Study Title:	Inter-Arterial Treatment of Pancreatic Cancer Using the RenovoCath™ RC120 Catheter
Development Phase:	Post market
Study Type:	Global Multicenter, Prospective, Observational Registry
Product Description:	The RenovoCath™ RC120 Catheter is an endovascular multi-lumen, two handled catheter designed to isolate variable segments of arteries supplying the target organ using two slideable, compliant balloons.
Study Population:	Patients with pancreatic cancer, with and without prior radiation therapy
Chemotherapeutic agent:	Gemcitabine injection (Gemzar®)
Primary Objectives	1. Evaluate survival in patients diagnosed with pancreatic cancer who undergo intra-arterial delivery of chemotherapeutic agents to the pancreas 2. Assess tumor response in the primary site of application as assessed by imaging
Primary Endpoints	1. Survival 2. Tumor response 3. Performance of RenovoCath in defined population (pancreatic cancer) in a clinical setting
Secondary Objective/ Endpoints	1. Assess conversion rate from unresectable or borderline resectable to potentially resectable or resectable pancreatic cancer 2. Further define and analyze potential selection criteria for patients who present with locally advanced pancreatic cancer that may benefit from the intra-arterial procedure
Study Sites:	Multicenter

[0106] After patient screening and enrollment, eligible patients underwent selective catheterization introduced via the femoral artery into the celiac axis into the splenic, hepatic and/or superior mesenteric artery(ies) using the TAMP technique as described above for the localized delivery of gemcitabine followed by embolization agent lipiodol. An interventional radiologist used the RenovoCath™ RC 120 catheter to optimize drug delivery to the tumor(s). At the conclusion of the case, the femoral artery arteriotomy was sealed and patient monitored as per standard institutional protocol.

[0107] All patients enrolled during the two-year registry enrollment period were followed periodically for survival outcome. Patients were contacted by telephone at the following

intervals after the final intra-arterial treatment: 6 months $\pm$ 30 days, 1 year $\pm$ 30 days and 2 years $\pm$ 30 days. Patients were assessed for serious adverse events with specific attention to events related to local delivery of chemotherapeutic agents to the pancreas and device performance.

[0108] FIG. 9 shows the superiority of the TAMP technique as compared to standard systemic chemotherapy. Graph 900 shows the increase in survival benefits in patients treated with TAMP compared to those treated with a systemic intravenous infusion of the same drug (*i.e.*, gemcitabine). Line 902 represents the survival percentage of patients completing eight treatments of gemcitabine using the TAMP technique. Line 904 represents the survival percentage of patients completing more than two treatments of gemcitabine using the TAMP technique. And line 906 represents the survival percentage of patients given systemic infusion of gemcitabine, the results of which are taken from Chauffert *et al.*, *Ann. Oncol.*, 2008, 19:1592-9. As depicted in FIG. 9, the survival rates of the patients treated with the TAMP technique (*i.e.*, lines 902, 904) were greater than the survival rates of the patients given systemic infusion of the same drug (*i.e.*, line 906).

[0109] FIG. 10 is a graph 1000 illustrating the effect of the TAMP technique on patient survival after radiation therapy compared with the effect of the TAMP technique with no radiation therapy. Fifteen patients with locally advanced pancreatic cancer were treated with gemcitabine in a dose-escalated protocol administered in four cycles using the TAMP technique. Each cycle consisted of two treatments two weeks apart. The efficacy data for the fifteen patients who received more than one cycle of TAMP treatment are shown in FIG. 10. Of these fifteen patients, five had no prior treatment of any kind, five had prior systemic chemotherapy, and five received radiation in addition to systemic chemotherapy prior to entering the study. On average, patients received radiation one to six months prior to enrolling in the study and receiving TAMP therapy. For the three groups of patients, the most pronounced survival benefit was seen in patients who had received radiation prior to the initiation of TAMP therapy. Specifically, patients with prior radiation had a significant improvement in survival compared to the ones that had either no prior treatment or only prior systemic chemotherapy with no radiation therapy. Patients were treated with TAMP therapy, regardless of their prior history. The dotted portion of the bars (1020, 1030) indicates the average time from diagnosis to the first TAMP therapy, while the clear portion of the bars (1010, 1022, 1032) indicates the average time from the first TAMP therapy to death.

[0110] This study demonstrated that a course of radiation prior to chemotherapy treatment administered via the TAMP technique has significant clinical benefit in patients with locally advanced pancreatic cancer. Combining these two modalities led to significant increase in median survival, reduction of tumor markers, and downsizing of the tumor. It is expected that a similar combination therapy would have clinical benefit in any solid tumors where diffusion-dependent infusion of a chemotherapeutic may be considered as a treatment option.

[0111] Although some of the methods have been described above as including the delivery of a therapeutic agent using a catheter having two occlusion elements (such as the catheter 100), in some embodiments, a method can include delivering a dose of a chemotherapeutic agent using any suitable mechanism. For example, in some embodiments, a method of treating a tumor can include first administering a dose of radiation and then, after a time period, delivering a dose of a chemotherapeutic agent. As described herein, by delivering the dose of chemotherapeutic agent after the time period has elapsed, the effectiveness of the chemotherapeutic agent can be enhanced by reducing washout from the targeted tissue. The chemotherapeutic agent can be delivered using any suitable mechanism, such as, for example, a catheter with a vessel proximate the target tissue, an injector, an implanted reservoir (e.g., a microsphere or the like), or the like. For example, FIG. 13 is a flow chart of a method 10 of treating a cancerous tumor according to an embodiment. The method is described along with the schematic illustrations of the tumor T in the target tissue (identified as TISSUE) in FIGS. 14 and 15.

[0112] The method 10 includes administering, at a first time, a dose of radiation (identified as RAD) to the target tissue including a tumor T, at 11. The target tissue can be any tissue of the types shown and described herein. For example, in some embodiments, the target tissue can be the pancreas, the liver, the brain, lung tissue, uterine tissue, or any other tissue that includes an undesirable tumor T. The radiation RAD can be delivered by any suitable system 20 (e.g., a system including a radiation source) and according to any suitable protocol as described herein. For example, in some embodiments, the radiation system 20 can be an internal radiation system that generates the radiation RAD within the body. In other embodiments, the radiation system 20 can be an external radiation system that generates the radiation RAD outside of the body. The radiation is delivered in a dose that is effective to reduce the microvasculature within the target tissue, similar to that described above with reference to FIGS. 6A and 6B. For example, in some embodiments, the method includes delivering approximately 20 to 50 Gy of radiation over approximately one to five weeks in approximately one to 25 sessions.

[0113] The method 10 further includes administering, at a second time, a dose of a chemotherapeutic agent (identified as CHEM) to the target tissue, at 12. The chemotherapeutic agent CHEM can be delivered by any suitable system 30 and according to any suitable protocol as described herein. For example, in some embodiments, the delivery system 30 can be a catheter of the types shown and described herein. In other embodiments, the delivery system 30 can be a catheter having two occlusion elements that are in a fixed position relative to each other. In such embodiments, the distance between the two occlusion elements determines the area (or length) of the vessel that is isolated for delivery of the agent. In yet other embodiments, the delivery system 30 can be a catheter having a single occlusion element. For example, in such embodiments, the catheter can be conveyed into a terminal vessel adjacent the target tissue and can be sufficient to isolate a segment of the terminal vessel. In yet other embodiments, the delivery system 30 need not be a catheter, but rather, can be an injector, a reservoir, or other delivery system.

[0114] The second time is after the first time by a period of time. As described herein, the dose of radiation is effective to reduce the microvasculature within the target tissue during the period of time. As described herein, the reduction in the microvasculature of the tissue enhances the efficacy of the later-delivered chemotherapeutic agent by reducing drug washout. The period of time can be, for example, between two weeks and six months.

[1001] While various embodiments have been described above, it should be understood that they have been presented by way of example only, and not limitation. Where schematics and/or embodiments described above indicate certain components arranged in certain orientations or positions, the arrangement of components may be modified. While the embodiments have been particularly shown and described, it will be understood that various changes in form and details may be made. Although various embodiments have been described as having particular features and/or combinations of components, other embodiments are possible having a combination of any features and/or components from any of embodiments as discussed above. For example, the size and specific shape of the various components can be different from the embodiments shown, while still providing the functions as described herein. Furthermore, each feature disclosed herein may be replaced by alternative features serving the same, equivalent or similar purpose, unless expressly stated otherwise. Thus, unless expressly stated otherwise, each feature disclosed is one example only of a generic series of equivalent or similar features.

[1002] Where methods and/or events described above indicate certain events and/or procedures occurring in certain order, the ordering of certain events and/or procedures may be modified. Additionally, certain events and/or procedures may be performed concurrently in a parallel process when possible, as well as performed sequentially as described above.

What is claimed is:

1. A chemotherapeutic agent comprising one or more compounds selected from any of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate, for use in treatment of a tumor, wherein the chemotherapeutic agent is administered subsequent to a course of radiation therapy applied to tissue surrounding the tumor after a period of time sufficient for the radiation therapy to reduce or devascularize the tissue.
2. A chemotherapeutic agent comprising one or more compounds selected from any of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate, for use in treatment of a tumor, wherein the chemotherapeutic agent is administered to a vessel proximate to tissue surrounding the tumor after administering a course of radiation therapy to the tissue.
3. The chemotherapeutic agent of any one of claims 1 and 2, wherein the chemotherapeutic agent includes at least two compounds selected from any of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, or sunitinab malate.
4. The chemotherapeutic agent of any one of claims 1 and 2, wherein the administration is via a catheter device.
5. The chemotherapeutic agent of claim 4, wherein the catheter device defines a lumen and an infusion portion, the lumen in communication with the infusion port and configured to administer the chemotherapeutic agent.

6. The chemotherapeutic agent of claim 5, wherein the catheter device includes at least one occlusion element configured to isolate a vessel segment, and the lumen is configured to administer the chemotherapeutic agent to the vessel segment isolated by the at least one occlusion element.
7. The chemotherapeutic agent of any one of claims 1 and 2, wherein the tumor is a pancreatic tumor.
8. The chemotherapeutic agent of claim 2, wherein the chemotherapeutic agent is administered to the vessel after a period of time following the administration of the course of radiation therapy to the tissue, such that the course of radiation can devascularize the tissue.
9. The chemotherapeutic agent of any one of claims 1 and 8, wherein the period of time is between two weeks and six months.
10. The chemotherapeutic agent of any one of claims 1 and 2, wherein the course of radiation therapy includes an amount of radiation between 20 and 50 gray (Gy) being administered over one or more sessions and a period of one to five weeks.
11. The chemotherapeutic agent of claim 10, wherein the amount of radiation is selected based on one or more characteristics of the tumor, the one or more characteristics comprising at least one of: a location of the tumor, and a size of the tumor.
12. The chemotherapeutic agent of any one of claims 1 and 2, wherein the chemotherapeutic agent is administered subsequent to administering a first agent.
13. The chemotherapeutic agent of claim 12, wherein the first agent is a dye.
14. The chemotherapeutic agent of claim 12, wherein the first agent is a first chemotherapeutic agent and the chemotherapeutic agent is a second chemotherapeutic agent different from the first chemotherapeutic agent.

15. The chemotherapeutic agent of claim 2, wherein the chemotherapeutic agent is administered subsequent to adjusting an intraluminal pressure of a segment of the vessel to a level of pressure of an interstitial space between the vessel and the tissue.
16. A method, comprising:
administering, at a first time, a dose of radiation to a target tissue including a tumor;
and
delivering, at a second time, a dose of a chemotherapeutic agent to the target tissue, the second time being after the first time by a period of time, the dose of radiation being effective to reduce the microvasculature within the target tissue during the period of time such that the washout of the chemotherapeutic agent is reduced.
17. The method of claim 16, wherein the delivering the dose of the chemotherapeutic agent includes:
isolating a segment of a vessel proximate to the target tissue;
adjusting an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target tissue; and
delivering the dose of the chemotherapeutic agent to the segment via a catheter device.
18. The method of claim 17, wherein the catheter device includes an occlusion element configured to isolate the segment of the vessel.
19. The method of claim 17, wherein:
the catheter device includes a first occlusion element and a second occlusion element;
and
the isolating the segment includes expanding the first occlusion element and the second occlusion element.
20. The method of claim 16, wherein the delivering the dose of the chemotherapeutic agent includes diffusing the chemotherapeutic agent through any one of a reservoir or a wall of a vessel associated with the target tissue.

21. The method of claim 16, wherein the chemotherapeutic agent includes one or more compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.
22. The method of claim 16, wherein the period of time is between two weeks and six months.
23. A method, comprising:
administering a dose of radiation to a target area including a tumor;
inserting a catheter device into a vessel, the catheter device including a first occlusion element and a second occlusion element;
isolating a segment of the vessel proximate to the target area using the first occlusion element and the second occlusion element;
delivering a dose of an agent to the segment via the catheter device.
24. The method of claim 23, wherein the agent is a chemotherapeutic agent.
25. The method of claim 24, wherein the chemotherapeutic agent includes one or more compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.
26. The method of claim 24, wherein the chemotherapeutic agent includes at least two compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

27. The method of claim 23, wherein the catheter device defines a lumen and an infusion port, the lumen in communication with the infusion port and configured to deliver the dose of the agent to the segment.
28. The method of claim 27, wherein the infusion port is disposed on the catheter device between the first occlusion element and the second occlusion element such that the infusion port can deliver the dose of the agent to the segment isolated between the first occlusion element and the second occlusion element.
29. The method of claim 23, wherein the tumor is a pancreatic tumor.
30. The method of claim 23, wherein the insertion of the catheter device into the vessel occurs after the administering of the dose of radiation.
31. The method of claim 23, wherein the delivery of the dose of the agent occurs after a predefined period of time following the administering of the dose of radiation.
32. The method of claim 31, wherein the predefined period of time allows the dose of radiation to devascularize the target area.
33. The method of claim 31, wherein the predefined period of time is between two weeks and six months.
34. The method of claim 23, further comprising administering one or more additional doses of radiation to the target area,
wherein the dose of radiation and the additional doses of radiation are administered during a period of one to five weeks, and
wherein the dose of radiation and the additional doses of radiation include an amount of radiation totaling between 20 and 50 gray (Gy).

35. The method of claim 34, wherein the amount of radiation is selected based on one or more characteristics of the tumor, the one or more characteristics including at least one of: a location of the tumor, and a size of the tumor.
36. The method of claim 23, wherein the dose of the agent is a first dose of a first agent, and further comprising delivering a second dose of a second agent to the isolated area of the artery.
37. The method of claim 36, wherein the first agent is a dye and the second agent is a chemotherapeutic agent.
38. The method of claim 36, wherein the first agent is a first chemotherapeutic agent and the second agent is a second chemotherapeutic agent different from the first chemotherapeutic agent.
39. A method, comprising:
administering a dose of radiation to a target area including a tumor;
isolating a segment of a vessel proximate to the target area;
adjusting an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area; and
delivering a dose of an agent to the segment via the catheter device.
40. The method of claim 39, wherein the segment of the vessel is isolated using the catheter device including a first occluding element and a second occluding element.
41. The method of claim 40, wherein the catheter device further includes a pressure sensor configured to measure the intraluminal pressure of the segment, and
wherein the adjusting of the intraluminal pressure is based on measurements made by the pressure sensor.
42. The method of claim 39, wherein the adjusting of the intraluminal pressure includes adjusting the intraluminal pressure of the segment to a level greater than a pressure of the target

area such that a pressure gradient is created between the segment and the target area, the pressure gradient configured to increase a permeation of the agent into the target area.

43. Use of a catheter device inserted into a vessel for delivering a dose of an agent to a segment of the vessel proximate to a target area including a tumor that has been administered a dose of radiation, the catheter device comprising a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

44. Use of a system comprising: (i) a radiation source for administration of a dose of radiation to a target area including a tumor; and (ii) a catheter device inserted into a vessel for delivering a dose of an agent to a segment of the vessel proximate to the target area that has been administered the dose of radiation, the catheter device comprising a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

45. The use of claim 43 or 44, wherein the agent is a chemotherapeutic agent.

46. The use of claim 45, wherein the chemotherapeutic agent consists of one or more compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

47. The use of claim 45, wherein the chemotherapeutic agent consists of at least two compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

48. The use of claim 43 or 44, wherein the catheter device defines a lumen and an infusion port, the lumen in communication with the infusion port and configured to deliver the dose of the agent to the segment.
49. The use of claim 48, wherein the infusion port is disposed on the catheter device between the first occlusion element and the second occlusion element such that the infusion port can deliver the dose of the agent to the segment isolated between the first occlusion element and the second occlusion element.
50. The use of claim 43 or 44, wherein the tumor is a pancreatic tumor.
51. The use of claim 43 or 44, wherein the insertion of the catheter device into the vessel occurs after the administration of the dose of radiation.
52. The use of claim 43 or 44, wherein the catheter device is for delivering the dose of the agent after a predefined period of time following the administration of the dose of radiation.
53. The use of claim 52, wherein the predefined period of time allows the dose of radiation to devascularize the target area.
54. The use of claim 52, wherein the predefined period of time is between two weeks and six months.
55. The use of claim 43 or 44, wherein the target area has been administered one or more additional doses of radiation,
wherein the dose of radiation and the additional doses of radiation have been administered during a period of one to five weeks, and
wherein the dose of radiation and the additional doses of radiation include an amount of radiation totaling between 20 and 50 gray (Gy).
56. The use of claim 55, wherein the amount of radiation is selected based on one or more characteristics of the tumor, the one or more characteristics comprising at least one of: a location of the tumor, and a size of the tumor.

57. The use of claim 43 or 44, wherein the dose of the agent is a first dose of a first agent, and further comprises delivery of a second dose of a second agent to the isolated area of the artery.

58. The use of claim 57, wherein the first agent is a dye and the second agent is a chemotherapeutic agent.

59. The use of claim 57, wherein the first agent is a first chemotherapeutic agent and the second agent is a second chemotherapeutic agent different from the first chemotherapeutic agent.

60. Use of a catheter device for delivering a dose of an agent to a segment of a vessel proximate a target area including a tumor that has been administered a dose of radiation, the catheter device configured to isolate the segment of the vessel proximate to the target area, and to adjust an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area.

61. The use of claim 60, wherein the catheter device comprises a first occluding element and a second occluding element, wherein the first occluding element and the second occluding element are for isolating the segment of the vessel.

62. The use of claim 61, wherein the catheter device further comprises a pressure sensor configured to measure the intraluminal pressure of the segment, and wherein the adjustment of the intraluminal pressure is based on measurements made by the pressure sensor.

63. The use of claim 60, wherein the adjustment of the intraluminal pressure comprises adjusting the intraluminal pressure of segment to a level greater than a pressure of the target area such that a pressure gradient is created between the segment and the target area, the pressure gradient configured to increase a permeation of the agent into the target area.

64. A catheter device inserted into a vessel for use in delivering a dose of an agent to a segment of the vessel proximate to a target area including a tumor that has been administered

a dose of radiation, the catheter device comprising a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

65. The catheter device of claim 64, wherein the agent is a chemotherapeutic agent.

66. The catheter device of claim 65, wherein the chemotherapeutic agent consists of one or more compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

67. The catheter device of claim 65, wherein the chemotherapeutic agent consists of at least two compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

68. The catheter device of claim 64, wherein the catheter device defines a lumen and an infusion port, the lumen in communication with the infusion port and configured to deliver the dose of the agent to the segment.

69. The catheter device of claim 68, wherein the infusion port is disposed on the catheter device between the first occlusion element and the second occlusion element such that the infusion port can deliver the dose of the agent to the segment isolated between the first occlusion element and the second occlusion element.

70. The catheter device of claim 64, wherein the tumor is a pancreatic tumor.

71. The catheter device of claim 64, wherein the insertion of the catheter device into the vessel occurs after the administration of the dose of radiation.

72. The catheter device of claim 64, wherein the catheter device is for delivering the dose of the agent after a predefined period of time following the administration of the dose of radiation.

73. The catheter device of claim 72, wherein the predefined period of time allows the dose of radiation to devascularize the target area.

74. The catheter device of claim 72, wherein the predefined period of time is between two weeks and six months.

75. The catheter device of claim 74, wherein the target area has been administered one or more additional doses of radiation,

wherein the dose of radiation and the additional doses of radiation have been administered during a period of one to five weeks, and

wherein the dose of radiation and the additional doses of radiation include an amount of radiation totaling between 20 and 50 gray (Gy).

76. The catheter device of claim 75, wherein the amount of radiation is selected based on one or more characteristics of the tumor, the one or more characteristics comprising at least one of: a location of the tumor, and a size of the tumor.

77. The catheter device of claim 74, wherein the dose of the agent is a first dose of a first agent, and further comprises delivery of a second dose of a second agent to the isolated area of the artery.

78. The catheter device of claim 77, wherein the first agent is a dye and the second agent is a chemotherapeutic agent.

79. The catheter device of claim 77, wherein the first agent is a first chemotherapeutic agent and the second agent is a second chemotherapeutic agent different from the first chemotherapeutic agent.

80. A system comprising: (i) a radiation source for use in administration of a dose of radiation to a target area including a tumor; and (ii) a catheter device inserted into a vessel for use in delivering a dose of an agent to a segment of the vessel proximate to the target area that has been administered the dose of radiation, the catheter device comprising a first occlusion element and a second occlusion element, wherein the first occlusion element and the second occlusion element are for isolating the segment of the vessel proximate to the target area.

81. The system of claim 80, wherein the agent is a chemotherapeutic agent.

82. The system of claim 81, wherein the chemotherapeutic agent consists of one or more compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

83. The system of claim 81, wherein the chemotherapeutic agent consists of at least two compounds selected from a group consisting of: doxorubicin, erlotinib hydrochloride, everolimus, 5-FU, flurouracil, folfirinox, gemcitabine hydrochloride, gemcitabine-cisplatin, gemcitabine-oxaliplatin, irinotecan hydrochloride liposome, leucovorin, mitomycin C, mitozytrex, mutamycin, oxaliplatin, paclitaxel, paclitaxel albumin-stabilized nanoparticle formulation, and sunitinab malate.

84. The system of claim 80, wherein the catheter device defines a lumen and an infusion port, the lumen in communication with the infusion port and configured to deliver the dose of the agent to the segment.

85. The system of claim 84, wherein the infusion port is disposed on the catheter device between the first occlusion element and the second occlusion element such that the infusion port can deliver the dose of the agent to the segment isolated between the first occlusion element and the second occlusion element.

86. The system of claim 80, wherein the tumor is a pancreatic tumor.

87. The system of claim 80, wherein the insertion of the catheter device into the vessel occurs after the administration of the dose of radiation.

88. The system of claim 80, wherein the catheter device is for delivering the dose of the agent after a predefined period of time following the administration of the dose of radiation.

89. The system of claim 88, wherein the predefined period of time allows the dose of radiation to devascularize the target area.

90. The system of claim 88, wherein the predefined period of time is between two weeks and six months.

91. The system of claim 80, wherein the target area has been administered one or more additional doses of radiation,

wherein the dose of radiation and the additional doses of radiation have been administered during a period of one to five weeks, and

wherein the dose of radiation and the additional doses of radiation include an amount of radiation totaling between 20 and 50 gray (Gy).

92. The system of claim 91, wherein the amount of radiation is selected based on one or more characteristics of the tumor, the one or more characteristics comprising at least one of: a location of the tumor, and a size of the tumor.

93. The system of claim 80, wherein the dose of the agent is a first dose of a first agent, and further comprises delivery of a second dose of a second agent to the isolated area of the artery.

94. The system of claim 93, wherein the first agent is a dye and the second agent is a chemotherapeutic agent.

95. The system of claim 93, wherein the first agent is a first chemotherapeutic agent and the second agent is a second chemotherapeutic agent different from the first chemotherapeutic agent.

96. A catheter device for use in delivering a dose of an agent to a segment of a vessel proximate a target area including a tumor that has been administered a dose of radiation, the catheter device configured to isolate the segment of the vessel proximate to the target area, and to adjust an intraluminal pressure of the segment to a level of pressure of an interstitial space between the vessel and the target area.

97. The use of claim 96, wherein the catheter device comprises a first occluding element and a second occluding element, wherein the first occluding element and the second occluding element are for isolating the segment of the vessel.

98. The use of claim 97, wherein the catheter device further comprises a pressure sensor configured to measure the intraluminal pressure of the segment, and wherein the adjustment of the intraluminal pressure is based on measurements made by the pressure sensor.

99. The use of claim 96, wherein the adjustment of the intraluminal pressure comprises adjusting the intraluminal pressure of segment to a level greater than a pressure of the target area such that a pressure gradient is created between the segment and the target area, the pressure gradient configured to increase a permeation of the agent into the target area.




FIG. 1

2/16




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B

6/16


500

FIG. 5

7/16




FIG. 6A




FIG. 6B

8/16

700




FIG. 7

9/16




FIG. 8




FIG. 9




FIG. 10

12/16


1100

FIG. 11




FIG. 12

10

14/16




FIG. 13




FIG. 14

16/16




FIG. 15

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2018/033482

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K31/436 A61K31/337 A61K31/404 A61K31/407 A61K31/4745 A61K31/513 A61K31/517 A61K31/519 A61K31/555 A61K31/704 A61K31/7068 A61K33/24 A61P35/00								
According to International Patent Classification (IPC) or to both national classification and IPC								
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K A61P Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, BIOSIS, EMBASE, CHEM ABS Data								
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>X</td> <td> Anonymous: "Researchers Report Survival Benefits with Use of RenovoCath™ in Patients with Locally Advanced Pancreatic Tumors", RenovoRx , 19 April 2017 (2017-04-19), XP002783340, Retrieved from the Internet: URL: http://renovorx.com/researchers-report-survival-benefits-use-renovocath-patients-locally-advanced-pancreatic-tumors/ [retrieved on 2018-07-24] the whole document ----- -/-- </td> <td> 1-8, 15-21, 23-32, 39-53, 60-73, 80-89, 96-99 </td> </tr> </tbody> </table>			Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	Anonymous: "Researchers Report Survival Benefits with Use of RenovoCath™ in Patients with Locally Advanced Pancreatic Tumors", RenovoRx , 19 April 2017 (2017-04-19), XP002783340, Retrieved from the Internet: URL: http://renovorx.com/researchers-report-survival-benefits-use-renovocath-patients-locally-advanced-pancreatic-tumors/ [retrieved on 2018-07-24] the whole document ----- -/--	1-8, 15-21, 23-32, 39-53, 60-73, 80-89, 96-99
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.						
X	Anonymous: "Researchers Report Survival Benefits with Use of RenovoCath™ in Patients with Locally Advanced Pancreatic Tumors", RenovoRx , 19 April 2017 (2017-04-19), XP002783340, Retrieved from the Internet: URL: http://renovorx.com/researchers-report-survival-benefits-use-renovocath-patients-locally-advanced-pancreatic-tumors/ [retrieved on 2018-07-24] the whole document ----- -/--	1-8, 15-21, 23-32, 39-53, 60-73, 80-89, 96-99						
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.								
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family								
Date of the actual completion of the international search		Date of mailing of the international search report						
8 August 2018		24/08/2018						
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Allnutt, Sarah						

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/033482

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MAHADEVAN A ET AL: "Stereotactic Body Radiotherapy and Gemcitabine for Locally Advanced Pancreatic Cancer", INTERNATIONAL JOURNAL OF RADIATION: ONCOLOGY BIOLOGY PHYSICS, vol. 78, no. 3, 1 November 2010 (2010-11-01), pages 735-742, XP027325922, PERGAMON PRESS, USA ISSN: 0360-3016 [retrieved on 2010-02-18] page 736, column 1, paragraph 4 - column 2, paragraphs 1,3 -----	1-3, 7-11,14, 16, 20-22, 33-35, 38, 54-56, 59, 90-92,95
X	WO 2014/197362 A1 (AGAH RAMTIN [US]; NAJMABADI KAMRAN [US]; ZANCHI MARTA [US]) 11 December 2014 (2014-12-11) the whole document -----	12,13, 36,37, 57,58, 64,65, 68-79, 93,94, 96-99
X	US 2014/276135 A1 (AGAH R; NAJMABADI K; ZANCHI M; ZANCHI M G) 18 September 2014 (2014-09-18) the whole document -----	12,13, 36,37, 57,58, 64,65, 68-79, 93,94, 96-99
A	RenovoRx: "RenovoCath Animation", 17 December 2014 (2014-12-17), XP054978562, Retrieved from the Internet: URL:https://www.youtube.com/watch?v=LFZ7tvCu2a4&feature=youtu.be [retrieved on 2018-08-07] the whole document -----	1-99
A	Anonymous: "RenovoCath for Targeted Fluid Delivery Into Peripheral Vasculature Cleared in Europe", Medgadget 23 October 2015 (2015-10-23), XP002783342, Retrieved from the Internet: URL:https://www.medgadget.com/2015/10/renovocath-targeted-fluid-delivery-peripheral-vasculature-cleared-europe.html [retrieved on 2018-07-25] the whole document -----	1-99

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2018/033482

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014197362 A1	11-12-2014	US 2017157370 A1 WO 2014197362 A1	08-06-2017 11-12-2014
US 2014276135 A1	18-09-2014	US 2014276135 A1 US 2016015948 A1	18-09-2014 21-01-2016