

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12) Rettet oversættelse af
europæisk patentskrift

(10) **DK/EP 2885408 T5**

-
- (51) Int.Cl.: **C 12 N 15/00 (2006.01)** **A 61 P 31/00 (2006.01)** **C 07 K 14/01 (2006.01)**
C 12 N 15/09 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2024-10-14**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-01-25**
- (86) Europæisk ansøgning nr.: **13829140.6**
- (86) Europæisk indleveringsdag: **2013-08-16**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-06-24**
- (86) International ansøgning nr.: **US2013055454**
- (87) Internationalt publikationsnr.: **WO2014028895**
- (30) Prioritet: **2012-08-16 US 201261684076 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Synthetic Genomics, Inc., 11149 N. Torrey Pines Rd, Suite 100, La Jolla, CA 92037, USA**
- (72) Opfinder: **GIBSON, Daniel, G., 5241 Shelley Place, Carlsbad, CA 92008, USA**
GILL, John, E., 1432 Dolphin Court, San Marcos, CA 92078, USA
VENTER, J., Craig, 7510 Hillside Drive, La Jolla, CA 92037-3940, USA
- (74) Fuldmægtig i Danmark: **Patrade A/S, Ceresbyen 75, 6., 8000 Århus C, Danmark**
- (54) Benævnelse: **DIGITAL TIL BIOLOGISK KONVERTER**
- (56) Fremdragne publikationer:
WO-A1-99/14318
WO-A2-2008/028024
US-A1- 2004 223 885
US-A1- 2005 267 971
US-B2- 7 118 883
US-B2- 7 923 533
SIYING MA ET AL: "DNA synthesis, assembly and applications in synthetic biology", CURRENT OPINION IN CHEMICAL BIOLOGY, vol. 16, no. 3-4, 1 August 2012 (2012-08-01), pages 260-267, XP055212288, ISSN: 1367-5931, DOI: 10.1016/j.cbpa.2012.05.001
DANIEL G GIBSON ET AL: "Enzymatic assembly of DNA molecules up to several hundred kilobases", NATURE METHODS, vol. 6, no. 5, 12 April 2009 (2009-04-12), pages 343-345, XP055224105, GB ISSN: 1548-7091, DOI: 10.1038/nmeth.1318
DE ROCQUIGNY, H ET AL: 'First Large Scale Chemical Synthesis Of The 72 Amino Acid HIV-1 Nucleocapsid Protein NCp7 In An Active Form.' BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS. vol. 180, no. 2, 31 October 1991, pages 1010 - 1018, XP029118880
D. G. Gibson ET AL: "Creation of a Bacterial Cell Controlled by a Chemically Synthesized Genome", Science,

Fortsættes ...

vol. 329, no. 5987, 2 July 2010 (2010-07-02), pages 52-56, XP055082599, ISSN: 0036-8075, DOI: 10.1126/science.1190719

NOTKA FRANK ET AL: "Industrial scale gene synthesis", METHODS IN ENZYMOLOGY, ACADEM. PRESS, USA, vol. 498, 1 January 2011 (2011-01-01), pages 247-275, XP009192694, ISSN: 1557-7988

Tom Ellis ET AL: "DNA assembly for synthetic biology: from parts to pathways and beyond", Integrative Biology, vol. 3, no. 2, 1 January 2011 (2011-01-01), page 109, XP055047340, ISSN: 1757-9694, DOI: 10.1039/c0ib00070a

Nathan J. Hillson ET AL: "j5 DNA Assembly Design Automation Software", ACS synthetic biology, vol. 1, no. 1, 20 January 2012 (2012-01-20), pages 14-21, XP055325880, USA ISSN: 2161-5063, DOI: 10.1021/sb2000116

Daniel G Gibson ET AL: "Chemical synthesis of the mouse mitochondrial genome", Nature Methods, vol. 7, no. 11, 1 November 2010 (2010-11-01), pages 901-903, XP055173787, ISSN: 1548-7091, DOI: 10.1038/nmeth.1515

FIKRI E. ALEMDAROGLU ET AL: "Generation of Multiblock Copolymers by PCR: Synthesis, Visualization and Nanomechanical Properties", NANO LETTERS, vol. 9, no. 10, 14 October 2009 (2009-10-14), pages 3658-3662, XP055600046, US ISSN: 1530-6984, DOI: 10.1021/nl901899t

SIYING MA ET AL: "DNA synthesis, assembly and applications in synthetic biology", CURRENT OPINION IN CHEMICAL BIOLOGY, vol. 16, no. 3-4, 1 August 2012 (2012-08-01), pages 260 - 267, XP055212288, ISSN: 1367-5931, DOI: 10.1016/j.cbpa.2012.05.001

DANIEL G GIBSON ET AL: "Enzymatic assembly of DNA molecules up to several hundred kilobases", NATURE METHODS, vol. 6, no. 5, 12 April 2009 (2009-04-12), GB, pages 343 - 345, XP055224105, ISSN: 1548-7091, DOI: 10.1038/nmeth.1318

D. G. GIBSON ET AL: "Creation of a Bacterial Cell Controlled by a Chemically Synthesized Genome", SCIENCE, vol. 329, no. 5987, 2 July 2010 (2010-07-02), pages 52 - 56, XP055082599, ISSN: 0036-8075, DOI: 10.1126/science.1190719

NOTKA FRANK ET AL: "Industrial scale gene synthesis", METHODS IN ENZYMOLOGY, ACADEM. PRESS, USA, vol. 498, 1 January 2011 (2011-01-01), pages 247 - 275, XP009192694, ISSN: 1557-7988

TOM ELLIS ET AL: "DNA assembly for synthetic biology: from parts to pathways and beyond", INTEGRATIVE BIOLOGY, vol. 3, no. 2, 1 January 2011 (2011-01-01), pages 109, XP055047340, ISSN: 1757-9694, DOI: 10.1039/c0ib00070a

NATHAN J. HILLSON ET AL: "j5 DNA Assembly Design Automation Software", ACS SYNTHETIC BIOLOGY, vol. 1, no. 1, 20 January 2012 (2012-01-20), USA, pages 14 - 21, XP055325880, ISSN: 2161-5063, DOI: 10.1021/sb2000116

DANIEL G GIBSON ET AL: "Chemical synthesis of the mouse mitochondrial genome", NATURE METHODS, vol. 7, no. 11, 1 November 2010 (2010-11-01), pages 901 - 903, XP055173787, ISSN: 1548-7091, DOI: 10.1038/nmeth.1515

FIKRI E. ALEMDAROGLU ET AL: "Generation of Multiblock Copolymers by PCR: Synthesis, Visualization and Nanomechanical Properties", NANO LETTERS, vol. 9, no. 10, 14 October 2009 (2009-10-14), US, pages 3658 - 3662, XP055600046, ISSN: 1530-6984, DOI: 10.1021/nl901899t

DE ROCQUIGNY, H ET AL: "First Large Scale Chemical Synthesis Of The 72 Amino Acid HIV-1 Nucleocapsid Protein NCp7 In An Active Form.", BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS., vol. 180, no. 2, 31 October 1991 (1991-10-31), pages 1010 - 1018, XP029118880

DESCRIPTION

FIELD OF THE INVENTION

[0001] The invention pertains to the automated conversion of biological information, such as a biological sequence of interest, into a final biological entity. The invention also pertains to providing for, e.g., a rapid response to biological threats.

BACKGROUND OF THE INVENTION

[0002] The following description of the background of the invention is provided to aid in understanding the invention, but is not admitted to be nor to describe, prior art to the invention.

[0003] Genetic information is stored in the form of a sequence of nucleotides that form a DNA molecule, which thus encodes the information necessary for the biological synthesis of proteins and peptides necessary for all cellular processes. Digital technology allows the transmission of digital information across enormous distances within seconds. This information is useful for any number of processes that convert the digital information into a useful function. It would be very desirable to have a system that allows the transmission of biological information in digital form across great distances, and then the conversion of that digital information into any of a wide variety of biological entities. Such biological entities would then be useful for the performance of a wide variety of biological functions such as, for example, the response to a biological threat to a community. A desirable system would also allow for the use of the biological information for the synthesis of DNA molecules, RNA molecules, proteins, virus and phage particles, vaccines, and synthetic cells.

[0004] With respect to responding to a biological threat, one important aspect can be the provision of a sufficient quantity of vaccine to inoculate a sufficient number of the members of the population against the threat. A critical item of information necessary to manufacture a vaccine is the biological sequence information associated with the biological threat. When the biological threat is a virus, that sequence information can be obtained by deriving the sequence information from a viral sample. Once the sequence is determined, a vaccine can be manufactured. This can take a variety of formats, one of which is to manufacture the protein coat or portion thereof of the viral threat, or another antigenic component of the biological threat, which will provide the antigen to stimulate a response in inoculated individuals against the biological threat.

[0005] Thus, for example, a nucleic acid sequence coding for the antigen can be synthesized. This can involve up to several days of work to coordinate the synthesis of oligonucleotides, which can then be assembled to form the final nucleic acid sequence. This can also involve the participation of several laboratories. After being obtained the final nucleic acid sequence can

then be translated, whether *in vitro* or *in vivo*, to synthesize the antigenic protein.

[0006] In the response to a biological threat, time can be of the utmost importance, so that members of medical response teams who will implement a response plan or work directly with infected persons can be inoculated and thus obtain immunity from the threat and be available to continue to carry out their duties unimpeded by threats of illness. Delays in vaccine preparation also lead to insufficient quantities of vaccine being ready at critical times and are also an important limiting factor in responding to a biological threat.

[0007] Furthermore, the specifics of a viral or other biological threat often differ from one locale to another. Thus, a vaccine that might be maximally effective in one locale may not be as effective in another locale due to rapid virus mutation.

[0008] There is therefore a need for systems that can transmit biological information in digital form, and allow for the conversion of that digital information into a variety of biological final products. Having a system that is automated would also contribute greatly to achievement of these goals. Such a system will meet various biological challenges, such as rapidly and effectively responding to viral and other biological threats for which time may be critical. It would also allow for a response that is tailored from one locale to another to meet the specific threats present in various locales.

SUMMARY OF THE INVENTION

[0009] The invention is as set out in the claims.

[0010] The present invention provides systems for receiving biological sequence information from a remote location and the automated synthesis of a biological entity based on the biological sequence information. Also described are methods for receiving biological sequence information from a remote location and the automated synthesis of a biological entity based on the biological sequence information. The systems and methods enable a rapid response to viral and other biological threats that are specific to a particular locale. Kits are also provided that include materials and reagents for performing the methods on systems of the invention. The system is able to produce a biological entity such as a DNA molecule, an RNA molecule, protein, or peptide from the biological sequence information in an automated fashion, and is further able to produce viral particles *in vivo*, also in automated fashion. The system is thus useful in the production of vaccines, such as an influenza vaccine. In a first aspect the invention provides an automated system for the synthesis of a dsDNA molecule according to provided sequence information, comprising:

an assembly unit for assembling the dsDNA molecule according to the provided sequence information,

wherein the assembly unit comprises or is connected to vessels containing nucleotide

derivative or nucleoside phosphoramidites and containing components for transporting reagents within the system and for executing steps in an automated method for assembling the dsDNA molecule through multiple levels of biological assembly, wherein the multiple levels of biological assembly comprise 1) the synthesis of a plurality of oligonucleotides by linking nucleotide derivatives or nucleoside phosphoramidites, and 2) the assembly of one or more dsDNA molecule(s) by joining the plurality of oligonucleotides; wherein no human intervention occurs after the method is initiated until the method is completed; and

wherein the assembly unit further comprises software programming instructions which direct manipulations of the parts of the assembly unit to synthesize the plurality of oligonucleotides from nucleotide derivative or nucleoside phosphoramidites according to the sequence information and assemble the oligonucleotides into the dsDNA molecule in the automated method;

the automated method comprising performing on the plurality of oligonucleotides a step of PCR amplification in a first reaction zone of a reaction container, a step of error correction on the PCR-amplified oligonucleotides in a second reaction zone of a reaction container, and a second PCR step in a third reaction zone of a reaction container;

wherein the error correction is performed using a combination of a member of the CEL nuclease family of mismatch-specific nucleases and exonuclease III;

wherein the assembly unit employs robotics to transport samples between reaction zones such that the automated method can be conducted in one reaction container;

wherein the oligonucleotides are fragments of the dsDNA molecule or variants thereof;

and further wherein the assembly unit performs assembly in an automated method that comprises the synthesis of the dsDNA molecule that is greater than 100 bp by joining said oligonucleotides.

[0011] Also described herein is a system for receiving biological sequence information and activating the synthesis of a functional biological entity. The system has a receiving unit for receiving a signal encoding biological sequence information transmitted from a transmitting unit, the transmitting unit present at a remote location from the receiving unit. The system also has an assembly unit connected to the receiving unit, for assembling the biological entity according to the biological sequence information. The assembly unit has or is connected to reagent vessels containing biological building block molecules and/or already synthesized oligonucleotides, and also has components for transporting reagents within the system and for executing steps in an automated method for synthesizing the functional biological entity. The receiving unit of the system receives the biological sequence information and provides it to the assembly unit.

[0012] In one embodiment transmitting unit and the receiving unit are computers that are part

of a computer network. The biological building block molecules can be dNTPs or nucleoside phosphoramidites. The system can also have programming instructions for assembling the dNTPs or nucleoside phosphoramidites into a set of oligonucleotides according to the biological sequence information. It can also have programming instructions for assembling the set of oligonucleotides into a DNA molecule. In one embodiment the functional biological entity is a DNA molecule. The oligonucleotides can also be provided to the system of the invention (or to the methods described herein) already synthesized and the system or method is for assembling the oligonucleotides into a DNA molecule.

[0013] The system in one embodiment has one or more vessels containing reagents for the transcription and/or translation of the functional DNA molecule into a biological product. The biological sequence information can encode a nucleic acid encoding for at least a portion of a virus particle or bacteriophage. The assembly unit can also convert the biological entity into a biological product, and in some embodiments the biological product can be a virus particle or a portion of a virus particle or a phage or cells. The virus particle or portion of a virus particle can be a protein antigen. In some embodiments the assembly unit also has, or is connected, one or more vessels containing a host cell. The systems of the invention can also have one or more vessels and reagents for the transcription of the DNA molecule into an RNA molecule, and/or for the translation of an RNA molecule into a functional peptide or protein, which translation can be done *in vitro* or *in vivo*. The system can also have programming instructions for performing steps of an automated method in distinct reaction zones of a reaction container.

[0014] Also described herein are methods of synthesizing a functional biological entity. The method involves receiving biological sequence information from a transmitting unit in a system for synthesizing the functional biological entity. The system of the methods is a system of the invention described herein. The method further involves synthesizing the functional biological entity in an automated method.

[0015] In the methods the assembly unit can be prepared for assembly of the functional biological entity prior to receiving the biological sequence information. The preparation can involve charging reaction containers of the system with biological building blocks or building block polymers, such as dNTPs or nucleoside phosphoramidites or with already synthesized oligonucleotides or peptides. In some methods described herein the biological sequence information encodes for a nucleic acid molecule encoding for at least a portion of a virus particle. In methods where a functional protein or peptide is produced, it can be a viral protein or peptide. The functional biological entity can also be a DNA molecule larger than 5 kb.

[0016] In one embodiment of the methods the building block molecules are linked in an *in vitro* reaction to form building block polymers. The building block molecules can be nucleotides or nucleoside phosphoramidites, and the building block polymers can be oligonucleotides or peptides or ribonucleotides or derivatives of any.

[0017] In some embodiments the methods involve transporting oligonucleotides from a first zone of a reaction container to a second zone of the reaction container. The methods can also

involve performing one step of the method in a first reaction zone of the reaction container and a second step of the method in a second reaction zone of the reaction container. In one embodiment the first step is the assembly of a DNA molecule and the second step is the transcription of the DNA molecule into an RNA. The methods can also have a third step performed in a third reaction zone of the reaction container, the third step comprising the *in vivo* production of viral particles. In other embodiments synthesizing the biological entity comprises a DNA assembly step. The steps of the method can also involve multiple levels of biological assembly. Thus the methods can have a DNA assembly step and a RNA transcription step. The DNA assembly step can be performed in a first reaction zone of the reaction container and the RNA transcription step can be performed in a second reaction zone of the reaction container. The method can also have a step involving the *in vivo* production of viral particles in a third reaction zone of the reaction container.

[0018] The summary of the invention described above is not limiting and other features and advantages of the invention will be apparent from the following detailed description of the invention, and from the claims.

DETAILED DESCRIPTION OF THE DRAWINGS

[0019]

Figure 1 provides a graphical illustration of a use of the present invention, broadly depicting the transmission of biological information and its conversion into a useful biological entity.

Figure 2 illustrates a 0.8% pre-cast agarose gel showing the assembly of nucleic acid constructs HA (H) and NA (N) from various influenza virus strains, each assembled from 96 pooled oligonucleotides in a system of the invention using the methods described herein. Both Constructs HA and NA are of approximately 3 kb. Fig. 2a - Lane 1: A/Brisbane/10/2010(H1N1)_HA; Lane 2: A/Brisbane/10/2010(H1N1)_NA; Lane 3: X179A_TD(H1N1)_HA; Lane 4: X179A(H1N1)_NA; Lane 5: A/Victoria/361/2011_CDC/E3(H3N2)_HA ; Lane 6: A/Victoria/361/2011(H3N2)_NA; Lane 7: A/Brisbane/256/2011_P2/E3(H3N2)_HA; Lane 8: A/Brisbane/256/2011_P2/E3(H3N2)_NA; Standards lane. Fig 2b - Standards lane; Lane 1: B/Texas/06/2011_BX-45_HA; Lane 2: B/Texas/06/2011_BX-49_NA; Lane 3: B/New_Hampshire/1/2012_HA; Lane 4: B/New_Hampshire/1/2012_NA; Lane 5: B/Brisbane/60/08_HA; Lane 6: B/Brisbane/60/08_NA; Lane 7: B/Nevada/03/2011_v2_HA; Lane 8: B/Nevada/03/2011_v2_NA.

DETAILED DESCRIPTION OF THE INVENTION

[0020] The present invention provides a system for the transmission of biological information in

digital form across great distances, and the subsequent and rapid conversion of that information into a final biological entity. The biological sequence information can be that information necessary for the synthesis of a DNA molecule, an RNA molecule, a protein or peptide, virus and/or phage particles, vaccines, synthetic cells, or any biological entity. The biological entities can be synthesized by a system of the invention located at the particular locale where the biological entity is to be assembled or synthesized.

[0021] The invention provides numerous advantages with respect to rapidly responding to biological threats. In one embodiment the biological threat is a viral threat. The invention allows for biological sequence information to be rapidly transmitted to a location under a biological threat, for example a remote location, and immediately be used in the process for preparing the vaccine or other materials necessary for an appropriate response. The threat can be an influenza or other viral threat.

[0022] In one embodiment the biological sequence information is the sequence information necessary for the formulation of a vaccine to the virus, i.e., the sequence of one or more varying regions of the virus particle. The invention therefore allows for a system of the invention to be set up and prepared with necessary reagents ahead of time so that the system can receive biological sequence information and immediately and in an automated manner begin preparing a vaccine or other appropriate biological response to deal with the threat immediately upon receiving the sequence information. The community under threat can therefore have multiple or large-scale systems prepared to receive the sequence information in order to maximize the supply of vaccine that will be available within a short timeframe for inoculating a population against the biological threat. The invention is versatile and allows a vaccine to be customized for dealing with the specific viral threat in a particular locale. Thus, where different strains of virus are affecting different locales, each locale can tailor a vaccine response to the particular strain affecting that locale in order to respond most accurately and effectively to the threat. By reducing or eliminating the number of hours that pass prior to beginning preparation of the vaccine and by preparing the most effective vaccine for the subject locale, the effectiveness of the response to the biological threat is greatly improved.

[0023] The present invention also has applicability to responding to biological threats related to bioterrorism where the rapidity and accuracy of the response may impart a critical and lifesaving effectiveness to the response. Thus, locales facing such threats can prepare by setting up a system disclosed herein in order to be able to mount the fastest and most accurate response, and thus have the greatest chance of defeating the threat with minimal damage.

[0024] The systems of the invention can be designed so that the methods can be performed on any reaction containers containing a plurality of reaction areas where samples can be manipulated. The reaction areas can be either defined (i.e. by physical boundaries on the container) or undefined (i.e., defined only by the space occupied by a sample). In one embodiment the reaction container is a standard 96-well plate, which contains a plurality of small reaction areas accommodating samples to be manipulated, i.e., each well of the plate.

This is convenient because thermocyclers used in the systems of the invention (and in the methods described herein) are typically designed to hold a 96 well plate. But it will be realized that any type of reaction container that holds a plurality of reaction areas containing samples to be manipulated will also function in the invention. For example a reaction plate having other than 96 wells will be a suitable reaction container in the methods, or a petri dish having a plurality of reaction areas where sample can be manipulated is also suitable even if the reaction areas are not defined by physical boundaries. Even a biochip can be a reaction container since it has a plurality of reaction areas on the chip.

[0025] By defining and utilizing reaction zones on the reaction containers it was discovered that a system can be designed so that a multi-level process from synthesizing oligonucleotides to the production of a protein or vaccine and additional methods can be performed in an automated method. This is accomplished by performing particular steps in reaction areas located in a specific reaction zone on the reaction container.

[0026] By employing robotics in the invention samples can be transported from particular reaction zones of a reaction container to other reaction zones, and an automated method can be conducted in one reaction container. Samples are not harmed or destroyed by conditions (e.g., temperature) for carrying out subsequent steps in a process in the same reaction container because the appropriate conditions for subsequent steps can be established on the reaction container prior to the system moving the sample into the reaction zone where the next step will occur. Thus, the reaction samples can be moved across the reaction container and into a new zone as each subsequent step is performed and the conditions created on the reaction container for that subsequent step, and the entire process can be completed in an automated method in the same reaction container. Thus, the systems of the invention perform automated methods.

[0027] An automated method is one where no human intervention is necessary after the method is initiated - the method goes to completion from that point without a human having to perform any action, and the method produces a functional biological entity. For example, in one embodiment an automated method is a method where no human intervention occurs after sequence instructions are received by the system of the invention, and the system executes the method to produce a functional nucleic acid molecule. Human intervention is any action taken by a person to move the method to completion. Human intervention includes, but is not limited to, the manual movement of samples or of the reaction container from one location to another or an action involving the preparation or movement of samples or reagents, the provision or re-supply of reagents or reactants, the turning on or off of any component of the system, or any other action necessary to complete execution of the method. In one example of an automated method is initiated with the receipt of biological sequence information from a transmitting unit. The method can proceed as an automated method to result in production of a DNA molecule, and in one embodiment has multiple levels of biological assembly and proceeds to the transcription of the DNA to produce mRNA, and can also proceed further to the translation of the mRNA into a protein biological product.

[0028] The system of the invention can have multiple levels of biological assembly so that a synthesized DNA molecule, for example, can be transcribed into an RNA molecule, and the RNA molecule further translated into a protein or peptide of interest, or combined with a cell culture to result in production of a drug product or synthesis of a whole vaccine. Translated proteins can also be further processed by the system to produce, for example, a viral particle or vaccine, or portions thereof. Levels of assembly include 1) synthesis of a dsDNA or ssDNA molecule from linking nucleotides or nucleoside phosphoramidites; 2) assembly of a dsDNA molecule by joining oligonucleotides (a DNA assembly step); 3) synthesis/transcription of an RNA molecule according to a DNA molecule (a transcription step); 4) synthesis of a protein or peptide from joining amino acids or synthesis of a protein from joining peptides; 5) synthesis/translation of a protein or peptide according to the sequence of an RNA molecule (a translation step); 6) assembly of two or more proteins (or protein sub-units) into a protein having quaternary structure (e.g., a virus particle or portion thereof). The systems of the invention (and of the methods described herein) can perform any of these levels of assembly. A system or method that performs at least two of these levels of synthesis is a system or method having multiple levels of biological assembly, but systems and methods can also be devised having at least 3 levels or at least 4 levels or at least 5 levels of biological assembly. The systems of the invention (and the methods described herein) synthesize a dsDNA molecule from nucleotides (including derivatives) or nucleoside phosphoramidites (including derivatives). The systems of the invention (and the methods described herein) synthesize a dsDNA molecule from oligonucleotides. In another embodiment the systems of the invention (and the methods described herein) synthesize a protein or peptide from amino acids. The oligonucleotides or peptides described herein can also be modified, derivatized, or labeled, as described herein.

[0029] With reference to Figure 1 there is depicted an embodiment of the invention. A transmitting unit 1 transmits biological sequence information 3 from a location to the receiving units 2 of various systems of the invention located in locations remote from the transmitting unit 1. The receiving units 2 receive the signal encoding biological sequence information 3 and provide the biological sequence information to an assembly unit. The assembly unit assembles the biological entity 4, which in this embodiment is a DNA molecule. In this embodiment the assembly unit further converts the biological entity into a biological product 5, which can be a virus coat or portion thereof 5a, or a virus 5b, or a protein antigen 5c. The biological product is useful as a vaccine 6.

Receiving Unit

[0030] The receiving unit receives a signal encoding biological sequence information. In one embodiment the receiving unit is a computer that receives digitally encoded biological sequence information from the transmitting unit. The receiving unit can therefore be connected to the internet and/or can be otherwise connected to a computer network, which allows information to be exchanged between computers connected to the network. The internet is also a computer network as the term is used herein. In one embodiment the transmitting unit is

located in a location remote from the receiving unit, for example in another city or another country or nation. The receiving unit can even be located in a place off of the Earth or outside of the Earth's atmosphere such as, for example, in an orbiting space station or even on the Moon or on a planet other than Earth. In some embodiments the transmitting unit is located in a remote location that is at least 10 miles or at least 25 miles or at least 50 miles or at least 100 miles or at least 250 miles or at least 1000 miles from the receiving unit. In some embodiments the receiving unit receives the biological sequence information and provides it to the assembly unit in a form that the assembly unit can convert into programming instructions for the assembly of the biological sequence. The receiving unit in one embodiment is therefore connected to the transmitting unit through a computer network. The receiving unit is also connected to the assembly unit. In various embodiments the receiving unit can also be a computer or circuit board or part of a computer integrated into the assembly unit, or a sub-unit of the assembly unit. The connections between the receiving unit and transmitting unit can be indirect, i.e., through one or more additional computers, routers, or other electronic devices interposed between them. The connection can also be through a wireless or satellite connection. The connection can also be direct, as in a direct connection between the transmitting unit and receiving unit through the computer network. However, the receiving unit will receive the signal transmitted by the transmitting unit. In one embodiment the receiving unit receives the biological sequence in the form of electromagnetic waves and provides the sequence to the assembly unit for assembly of the biological entity or biological product.

Transmitting Unit

[0031] The transmitting unit transmits biological sequence information to the receiving unit of the system. A digital signal can represent information in the form of a series of binary digits. Persons of ordinary skill in the art understand that virtually any kind of information can be readily transmitted as a digital signal, including biological sequence information. These types of information are readily transmitted through computer networks, which can span the globe. In one embodiment of the invention the transmitting unit is a computer that is connected to a computer network that allows information to be transmitted from one computer to another, or exchanged between computers connected to the network (e.g., the internet). Thus, in one embodiment the transmitting unit and the receiving unit are computers that are part of a computer network, and the transmitting unit has hardware and/or software for acquiring or converting the sequence information into a digital format and transmitting the sequence information to a receiving unit, for example through a computer network. In other embodiments the transmitting unit is a telephone or a keyboard and the operator can manually key in the biological sequence information. The transmitting unit can also be an electronic device that sends the message to the receiving unit in any format (e.g., HTML). The format of data encoding and transmission can be any convenient format that can be converted by the receiving unit into instructions for synthesis of the biological sequence. The transmitting unit can obtain the information from a person that enters the information or can obtain it in an automated fashion from an instrument that determines the sequence. The source of the biological sequence information can be a sample of the virus or other biological entity.

[0032] The biological sequence information can be a series of nucleotides, phosphoramidites (e.g., nucleoside phosphoramidites), amino acids, or any sequence providing the information necessary to synthesize the primary structure of a biological entity. The biological sequence information can also be provided as a code that can be deciphered to arrive at the information necessary to synthesize the primary structure of the biological entity.

[0033] It is often the case that the sequence of a biological entity (e.g., a virus) will be deciphered at a scientific facility with special expertise in this technical area. DNA sequencing technology is widely known in the art and various methods of isolating DNA and performing sequencing are available. Thus, the sequence can be deciphered using standard, known technologies such as, for example, sequencing by synthesis, the dideoxy or chain termination (Sanger) method, the chemical degradation method, thermal cycle sequencing, pyrosequencing, or sequencing by hybridization. These are only some examples of DNA sequencing methods as persons of ordinary skill in the art are aware of numerous additional methods. The precise method of sequencing is not important, only that the sequence of the biological entity is acquired by the transmitting unit for transmission to a receiving unit from the transmitting unit. The sequence can be deciphered and can also be a predicted sequence, which predicted sequence can be based on trends or information derived from year to year changes in a virus. The year to year changes in the virus can be, for example, changes over a period of at least 2 years or at least 3 years or at least 5 years or at least 10 years.

[0034] In one embodiment the transmitting unit can also comprise or be associated with a DNA sequencer, or have the capacity to sequence DNA. Therefore a transmitting unit can have the capacity to determine the DNA sequence of an organism or biological sample and transmit the biological sequence information to a receiving unit at a remote location. Therefore the system of the invention (and the methods described herein) can have a sample return capability from remote locations, including locations outside of the Earth's atmosphere, such as the Moon or other planets.

Assembly Unit

[0035] The assembly unit synthesizes the biological sequence according to the information received by the receiving unit from the transmitting unit. The assembly unit deployed in the system of the invention can be connected to the receiving unit. As with the other system components the connection can be either a direct connection between the units, or can be an indirect connection through one or more additional computers, routers, or other electronic devices interposed between the units. The assembly unit can be in communication with the receiving unit so that, when the receiving unit receives the signal containing the biological sequence information from the transmitting unit, the information is provided to the assembly unit, which can immediately begin to synthesize the sequence in an automated method. The assembly unit can have various functions including, but not limited to, one or more of the synthesis of a nucleic acid molecule, the amplification of a nucleic acid molecule, and the

transcription and/or translation of a nucleic acid molecule into a peptide or protein.

[0036] In some embodiments the assembly unit has the ability to perform a second (or third or fourth) level of biological assembly, e.g., the assembly of biological entities produced into larger biological entities. Thus, in one embodiment the assembly unit can synthesize oligonucleotides in an automated method, and then subsequently assemble the oligonucleotides into a larger dsDNA molecule. In one embodiment the oligonucleotides synthesized can be fragments of a DNA molecule of interest, or portions or variants of a DNA molecule of interest. The oligonucleotides can be assembled in combinatorial fashion, either sequentially or simultaneously as desired. A typical oligonucleotide can be from 40-100 nucleotides or from 30-110 nucleotides or from 50-90 nucleotides or about 60 nucleotides. But in various other embodiments the DNA molecules produced by the assembly unit can be greater than 100 bp, or greater than 200 bp, or greater than 300 bp, or greater than 400 bp, or greater than 500 bp, or greater than 600 bp, or greater than 700 bp, or greater than 800 bp, or greater than 900 bp, or greater than 1000 bp or from 100-1000 bp or from 200-1000 bp or from 300-1000 bp or from 400-1000 bp or from 500-1000 bp or from 600-1000 bp or from 700-1000 bp. The lengths recited can be the lengths with or without single-stranded overhangs (i.e., "sticky ends").

[0037] In various embodiments the assembly unit comprises one or more sub-units of a nucleic acid synthesizer, a protein and peptide synthesizer, a PCR thermocycler, and one or more sub-units for performing *in vitro* transcription and/or translation on a sample. Another sub-unit can be provided for incubating a biological entity in a cell culture and/or for maintaining cells in a cell culture.

[0038] Any one or any combination of the sub-units or all of the sub-units of the assembly unit can be automated. The systems of the invention can include an automated nucleic acid and/or amino acid synthesizer. The assembly unit of the invention can contain a plurality of vessels or containers having the chemicals, reagents, biological building blocks, or building block polymers necessary to perform nucleic acid or protein and peptide synthesis, and valves necessary to permit or restrict the flow of reagents, as well as vessels for conducting certain chemical reactions. These can also be present outside the assembly unit with passages providing the assembly unit access to them. The biological building block molecules can be, for example, amino acids, dNTPs, and/or nucleoside phosphoramidites necessary for synthesis of the desired biological sequence. The biological building blocks can also include monosaccharides, disaccharides, or polysaccharides. When the biological entity or biological product comprises more than one type of molecule, for example a glycoprotein, the biological building blocks will comprise amino acids and mono-, di-, or poly-saccharides as desired. Oligonucleotides or other building block polymers that are already assembled can also be provided in the vessels or containers.

[0039] The biological building blocks can also be modified, derivatized, or labeled. For example the biological building blocks can be nucleoside phosphoramidites or nucleoside triphosphates (NTPs), or derivatives of either. In various embodiments the modified,

derivatized, or labeled building blocks can be aminoallyl, biotin, or 2' fluoro- modified or labeled NTPs or phosphoramidites. Additional modified, derivatized, or labeled building blocks include 2-aminopurine, 2,6-diaminopurine, 5-bromo deoxyuridine, inverted dT (prevents unwanted 5' ligations), dideoxycytidine, 5-methyl dC, deoxyinosine, 5-nitroindole, hydroxymethyl dC, Iso dC and iso dG, or locked nucleic acids to improve stability where desired. The assembly unit can contain the vessels that contain the chemicals, reagents, or biological building blocks, or the vessels can be outside the assembly unit and be connected to the assembly unit, for example via tubes or other pathways that carry the materials. The instruments typically contain software programming instructions necessary to control the instrument and synthesize a nucleic acid or protein/peptide in an automated fashion. Thus, the software programming instructions can be instructions directing the manipulations of the parts of the assembly unit to synthesize a set of oligonucleotides from biological building blocks (e.g., dNTPs or nucleoside phosphoramidites) according to the biological sequence information, and/or can include instructions for assembling the oligonucleotides into larger DNA molecules, and can further include instructions for transcribing the DNA molecules into RNA and for translating the RNA into protein molecules, and can further include instructions for assembling protein molecules into larger protein structures. In one embodiment the instructions are for assembling oligonucleotides into DNA or RNA molecules. In one embodiment the instructions are for assembling protein molecules into a virus or portion of a virus. And in another embodiment the system includes programming instructions for performing steps of an automated method in distinct zones of a reaction container.

[0040] The assembly unit can be a single, unitary unit, or can have various sub-units that perform the different functions. Thus, in one embodiment a sub-unit of the assembly unit performs oligonucleotide synthesis, for example with an oligonucleotide synthesizer. A robotic arm can be included to perform transfer of the reaction container (e.g. a 96 well plate) from the synthesizer to a liquid handler with thermocycling capabilities. Synthesized oligonucleotides can also be provided to another reaction container, or another zone of the same reaction container or to another sub-unit of the assembly unit for amplification, for example to a PCR thermocycler, and/or assembly. In one embodiment the system is provided with already synthesized oligonucleotides and assembles the oligonucleotides into one or more DNA molecules. And one or more other sub-units can perform transcription and/or translation of the synthesized nucleic acid into protein product, such as an automated *in vitro* translation system, or these functions too can be performed in another zone of the same reaction container. In each case the sub-units are prepared with the necessary reagents, chemical, and biological building blocks to perform the function in an automated manner. Thus, the assembly unit can begin to perform one or all of the functions immediately upon being provided with the biological sequence information from the receiving unit.

[0041] The assembly unit synthesizes the biological sequence according to the information received by the receiving unit from the transmitting unit or according to sequence information otherwise provided to the assembly unit. The assembly unit can thus in an automated fashion direct the synthesis of the desired biological entity. All of the units in the system, including the assembly unit, can be prepared with necessary software and pre-charged with necessary

reagents and chemicals in the required vessels so that the biological sequence information can be received by the receiving unit and the assembly unit can instantly begin synthesizing the desired biological entity. Thus, the assembly unit can be prepared for assembly or synthesis of the biological entity prior to receiving the biological sequence information.

[0042] Following synthesis of oligonucleotides or peptides the assembly unit can also have the capability to assemble oligonucleotides into larger nucleic acid molecules and, respectively, peptides into polypeptides and proteins. The systems and methods can also assemble DNA molecules from oligonucleotides. Thus, the nucleic acid molecule or protein or peptide can be assembled through a series of reactions. In one embodiment for nucleic acid molecules the GIBSON ASSEMBLY[®] (Synthetic Genomics, San Diego, CA) reaction is used to assemble the nucleic acids, but in other embodiments any suitable method of assembling oligonucleotides (or peptides) into larger nucleic acids (or polypeptides) can be employed. The oligonucleotides can also be assembled into double-stranded nucleic acid molecules. Persons of ordinary skill are also aware of methods of assembling peptides into polypeptides and proteins, which can also be applied in the invention.

[0043] The assembly unit, or sub-units thereof, comprise one or more vessels that contain reagents, chemicals, biological building blocks, or building block polymers useful for performing the function of the assembly unit or sub-unit thereof. The vessels can be containers made of glass or another suitable material and can be pre-filled by the user prior to the receiving unit receiving the biological sequence information from the transmitting unit. Thus, the assembly unit can be set up and waiting to receive the biological sequence information from the transmitting unit. In some embodiments the functions of the assembly unit or sub-sets thereof will be managed and orchestrated by appropriate software that will direct the combination of materials from the various vessels. Each sub-unit of the assembly unit can operate from its own software upon receipt of the required information or the assembly unit can have one program that directs the various sub-units that may comprise the assembly unit.

[0044] *In vitro* and/or *in vivo* transcription and translation can be performed using a variety of methods known to persons of ordinary skill. In different protocols extracts of rabbit reticulocytes, wheat germ, *E. coli*, as well as human cell lysates are used in the synthesis of proteins. Human *in vitro* translation systems can be useful when the biological entity is a vaccine for use against human pathogens. The extracts can be prepared and contain the essential components of cellular translational machinery such as, for example, ribosomes, tRNAs, aminoacyl-tRNA synthetases, initiation, elongation, and termination factors, energy sources such as ATPs, necessary co-factors and other proteins. In *in vitro* transcription and translation reactions can be performed at great speed and at the microliter or nanoliter level. A variety of commercially available kits can also be used for convenient setup and have been used with success but the essential components can also be assembled generically. Whether kits or generic materials are used, in some embodiments a cell-free solution containing essential components of the cellular translational machinery is utilized.

[0045] In one *in vitro* translation method extracts of an immortalized human cell line are used

to provide the ribosomes, initiation and elongation factors, tRNAs and the other basic components required for protein synthesis. Some systems utilize proprietary accessory proteins, ATP, and an energy regenerating system to sustain the synthesis of target proteins from DNA templates. The extracts can be prepared and pre-charged into a vessel of the assembly unit and be available to perform translation of synthesized nucleic acid when needed. The assembly unit can maintain these extracts at required temperatures to extend their useful life. The vessel or vessels for performing transcription and/or translation can also be linked to the vessels where DNA will be synthesized and amplified using PCR, and then transcribed into RNA. Upon transcription of RNA a portion of the reaction can be transferred into the vessel for performing translation, or the transcription and translation reactions can be performed in a single vessel. In some embodiments the biological entity requires protein modifications, such as glycosylated residues, a refolding step, or enzymatic modifications that must occur prior to the biological entity having activity or full activity. Reagents and programming protocols can be included within the assembly unit for this purpose as well.

[0046] All of these reactions can be performed in the reaction container itself within particular reaction areas or zones. The systems of the invention can also include sub-units, reaction areas, or reaction zones for the purification of biological products.

[0047] In one aspect the invention provides a system for the synthesis of a functional biological entity according to provided biological sequence information. The system has an assembly unit for assembling the biological entity according to the provided biological sequence information. The assembly unit contains or is connected to vessels containing biological building block molecules or building block polymers and has components for transporting reagents within the system and for executing steps in an automated method of the invention for synthesizing the functional biological entity. These systems can also have any of the components or features described herein. The biological entity can be any as described herein (e.g., a DNA molecule). The system can have a receiving unit for receiving a signal or data having the biological sequence information. The biological sequence information can be provided by manually keying in the sequence via an electronic interface, or by transmitting the sequence to a receiving unit from a transmitting unit, as described herein.

Biological Entity

[0048] The biological entity is a polynucleotide or DNA molecule, or a polyribonucleotide or an RNA molecule, or a peptide or polypeptide or protein. In some embodiments the biological entity is a nucleic acid or a DNA molecule. When the biological entity is a DNA or RNA molecule it can be convertible into a biological product. The DNA or RNA can be single-stranded DNA or RNA, or double-stranded DNA or RNA. When the biological entity is a DNA molecule it can be a functional DNA molecule, meaning that it can be transcribed into an RNA that can be translated into a functional protein or peptide, or is directly useful as a DNA molecule (e.g., as a DNA vaccine). A functional RNA molecule can be translated into a functional protein or peptide. A functional protein or peptide is one that has a clinical use in the treatment of a disease or

disorder, or is directly useful in some biological context (e.g., as a structural protein or is useful as an enzyme). In various embodiments the functional protein or peptide can be the domain of a protein, a binding subunit, an enzyme or enzyme subunit that has enzymatic activity, a protein or peptide that has antigenic activity that is useful in the generation of a vaccine, a viral protein or a subunit thereof, a viral coat protein (e.g., an HA or NA protein), or a plurality of proteins or peptides that form a viral particle when combined *in vitro* or when combined *in vitro* in host cells. The clinical use can thus be the generation of an antigenic response or the binding of a protein or peptide to a specific binding molecule or receptor but in one embodiment the antigenic response is one that furthers the treatment of a disease or disorder (e.g. the generation of a vaccine to influenza) or finds use in an assay to identify the presence of an epitope. The functional protein or peptide can also provide a desirable property such as, for example, a desirable taste, texture, scent, or another property. In one embodiment a functional protein or peptide has a function other than the generation of an immune or antigenic response.

[0049] The biological entity can be of any size. Oligonucleotides of up to about 200 bp can be assembled but greater accuracy can often be achieved by synthesizing smaller oligonucleotides. In various embodiments the biological entity is a single-stranded or double-stranded DNA molecule of greater than 100 base pairs in length, or greater than 200 bp, or greater than 300 bp, or greater than 400 bp, or greater than 500 bp, or greater than 1 kb, or greater than 2.5 kb or greater than 3 kb or greater than 4 kb or greater than 5 kb or greater than 6 kb or greater than 7 kb or greater than 8 kb or greater than 9 kb or 1-10 kb or 1-9 kb or 1-8 kb or 2-10 kb or 2-9 kb or 2-8 kb. Using additional techniques the person of ordinary skill with resort to the present disclosure will realize that DNA sequences synthesized as oligonucleotides can also be joined to assemble a DNA molecule using the method described herein of greater than 1 kb, or greater than 2 kb, or greater than 3 kb, or greater than 5 kb, or greater than 6 kb, or greater than 7 kb. In other embodiments the system of the invention (and the methods described herein) can be used to assemble a DNA molecule of greater than 100 kb, or greater than 200 kb, or less than 200 kb, or greater than 300 kb, or less than 300 kb, or greater than 500 kb, or greater than 800 kb or greater than 1 mega-base or less than 1 mega-base.

[0050] Biological sequence information is that sequence information necessary for the synthesis of a biological entity, for example a nucleic acid, peptide, or protein. In some embodiments the biological sequence information is the sequence (or order) of nucleotides, amino acids, or other building block that comprises the primary structure of the nucleic acid or peptide or protein that is the biological entity. In different embodiments this information may be provided to the receiving unit in a binary form or in another encoded form. Biological Product

[0051] A biological product is a biological molecule that is synthesized or assembled using the sequence information provided by the biological entity. Thus, in different embodiments the biological entity is a DNA molecule or an RNA molecule and the biological product can be any biological product made therefrom such as, for example, a viral genome or a portion thereof, a viral particle or viral coat or a portion of either, a bacteriophage or portion thereof, an antigenic

portion of a viral particle or viral coat, a bacterial genome or portion thereof, a gene, a nucleic acid sequence, a single-stranded DNA molecule (ssDNA), a double-stranded DNA molecule (dsDNA), an RNA molecule, an anti-sense RNA moiety, an siRNA moiety, an RNAi moiety, a double-stranded RNA moiety, a protein molecule or protein moiety, a protein antigen or portion thereof, an enzyme, a structural protein, a regulatory protein, a nutritional protein, a binding protein, a transport protein, a peptide molecule, a gene or genome of a fungus or portion thereof, or a synthetic cell. The biological product can also be a protein, peptide, or polypeptide that has undergone modification such as, for example, glycosylation of certain amino acid residues to produce a glycoprotein, or another molecule formed of two or more biological building blocks. In other embodiments the biological product is a genome of a synthetic bacteria or a portion thereof. In a particular embodiment the biological product is an influenza virus and the sequence information is information that is used to prepare a vaccine against the biological threat. When the biological product is a virus or virus particle, it can be an attenuated virus or a killed virus or a harmless virus. In some embodiments the biological entity itself can also be the biological product. An example of when the biological entity is the biological product is a DNA vaccine, where the DNA molecule itself is useful as a vaccine. DNA vaccines are comprised of pieces of DNA that code for pathogen proteins. After injection into the body the host cells synthesize the pathogen proteins, stimulating an immune response. In one embodiment the biological entity is a plurality of single-stranded DNA (ssDNA) or oligonucleotides that can be assembled into a double-stranded DNA (dsDNA). A "portion" of a molecule, virus, or other biological entity can be at least 10% or at least 20% or at least 30% or at least 40% or at least 50% or at least 60% or at least 70% or at least 80% or at least 90% of the native molecule or biological entity.

[0052] In some embodiments the action of a host cell can be utilized to arrive at the biological product. In systems and methods where host cells are used they can be located within a subunit of the assembly unit that can receive a biological entity from another subunit for further processing, or can simply be maintained in a zone of the reaction container. For example the host cells can be maintained in a vessel within the subunit. Host cells are useful in the systems of the invention (and the methods described herein) for a variety of purposes. While *in vitro* translation can be used in the systems of the invention (and the methods described herein), transcription, translation, and assembly of molecular sub-units can also be performed by host cells. Host cells can also be used to receive a biological entity and use it to synthesize a biological product. In one embodiment host cells are transfected with multiple DNAs, which are biological entities synthesized by a subunit of the assembly unit. The host cells can use the DNAs to make proteins within the host cells and/or to assemble the proteins into a biological product, for example a viral particle or portion thereof, or a protein or enzyme having multiple subunits. Host cells may also be used to replicate phage or viral particles following infection or transfection.

Methods

[0053] Described herein are methods of synthesizing a functional biological entity. In the

invention a digital DNA sequence or coded sequence can be entered into a software program that automatically designs a synthesis paradigm for the received DNA sequence (e.g., ARCHETYPE®, Synthetic Genomics, San Diego, CA). The software can break the sequence into designed overlapping oligonucleotides of, for example, about 50-80 bases or about 40-90 bases or about 30-100 bases or about 30-60 bases or about 30-80 bases or 20-100 bases or 20-80 bases or 10-100 bases or 10-90 bases or 10-80 bases, and any of these sizes can be used as the oligonucleotides described herein. The oligonucleotide sequences are then transmitted to the sub-units of the system for assembly. In a particular embodiment the software can design the oligonucleotides to contain an overlap of about 30 bp ($\pm 5\%$ or $\pm 10\%$) between adjacent oligonucleotides (ssDNA) of about 60 bp. The oligonucleotides can also be designed to have universal primer-binding domains for PCR amplification and restriction sites to release the primer-binding domains following the PCR amplification. The software used in the invention can also have the ability to modify received or desired sequence into codon-optimized sequences tailored to a particular host organism to be used in the method. The software can be present in any unit or sub-unit of the system.

[0054] The methods can be conducted on one or more samples that are situated in one or more reaction areas of a reaction container. In some embodiments the reaction container is a reaction plate. An example of a reaction plate is a 96 well plate, but it will be realized that the methods can be conducted on a reaction container having any number of reaction areas. When the reaction container is a reaction plate it can have any convenient number of reaction wells (areas), such as the 96 reaction wells on a standard 96 well plate. In some embodiments the methods involve one or more steps of transporting a sample from one zone of a vessel to a second zone of the vessel.

[0055] In some embodiments the method can be completed on a single reaction container. These embodiments can involve dividing the reaction container into reaction zones, and transporting one or more samples from one zone of the reaction container to a second zone of the reaction container to perform distinct steps of a method, but can also involve leaving all or a portion of the sample in place and moving the reaction container so that one or more zones of the reaction container are exposed to a different environment. For example, in a step involving PCR the zone can be located at a point in the system where it is exposed to a thermocycling schedule. At a DNA assembly step the reaction container can be located at another point in the system where it is exposed to the conditions appropriate for DNA assembly. The change of location can be accomplished by a physical movement of the reaction container or, alternatively, by a movement of one or more units of the system. The reaction container is thus moved relative to the system of the invention.

[0056] A zone of a reaction container is a distinct area of a reaction container containing one or more reaction areas where sample is collected. Normally each zone of a reaction container will contain multiple reaction areas, such as the wells in a reaction plate. But a reaction area does not require a physical barrier or boundary - it requires only that a distinct reaction can be carried out in the reaction area relative to other reaction areas. A reaction container can have any number of zones but each zone has one or more reaction areas for collecting and holding

a sample. Each zone of a reaction container is treated in the same manner or subjected to the same treatment or process in a given time period. For example the samples present in a zone may all be cycled at the same temperature of a PCR cycle being executed on that zone of the reaction container, or all reaction samples in a reaction zone may be picked up in automated fashion and moved to another zone on the same reaction container.

[0057] Thus in one embodiment of the methods a PCR step is performed in a first reaction zone of the reaction container and an error correction step is performed in a second zone of the reaction container, and a 2nd PCR step is performed in a third reaction zone of a reaction container. The methods can also have a step of DNA assembly performed in another reaction zone of the reaction container, and a transcription step performed in another reaction zone of the reaction container, and a translation step performed in another reaction zone of the reaction container, and a transfection step performed in another reaction zone of the reaction container. Thus, each step of any of the methods described herein can be performed in a distinct reaction zone of a reaction container.

Kits

[0058] Also described herein are kits for use in the systems of the invention (and in the methods described herein). The kits can contain reagents and components necessary to assemble a biological product, such as any combination of biological building blocks, building block polymers, buffers, or other reagents for carrying out a method of the invention on a system of the invention. Either alone or with any combination of the above reagents and components, the kits can also contain a reaction container for performing the methods described herein. Any combination of the reagents or reaction components or reaction containers can be provided in a container having the members of the kit. Reagents and reagent components can also be provided in a reagent vessel that fits onto an interface of a system of the invention.

[0059] The kits can also include one or more reaction containers that have been designed to fit onto an interface of a system of the invention and the reaction containers can be prepared or pre-charged with reagents and/or biological building blocks and/or building block polymers necessary for the synthesis of a biological entity using a system of the invention. The kits can also contain instructions for attaching the reaction containers or reagent vessels to a system of the invention and/or instructions for using the reaction containers or reagent vessels in a method described herein. The instructions can also be provided on a website and the kit can include a link to the website, either instead of or in addition to the instructions provided with the kit.

[0060] Any of the kits described herein can also include instructions for conducting an assay described herein and/or a link to a website providing the same information or information for preparing and/or using a system of the invention to synthesize a biological entity and/or for

preparing or pre-charging vessels with reagents, components, and/or biological building blocks necessary to synthesize a biological entity. For example, the instructions can provide guidance on the type and quantity of reagents to be used in the systems and methods. The kits can be provided or packaged in a single container or in multiple containers. The vessels can be made of glass, plastic or any suitable material, and may also be sterile. One or more vessels can be provided in a sterile container within the container comprising the kit.

Additional Aspects of the Systems

[0061] The systems of the invention produce a biological entity by carrying out an automated method. The biological entity can be any desired sequence. In one embodiment the systems of the invention can be pre-programmed to include desired sequences or constructs as part of the biological entity. For example the systems of the invention can be pre-programmed to enable the operator to append a particular sequence to be synthesized to a plasmid or other vector desirable for the further use of the synthesized sequence. In other embodiments the systems can synthesize a requested sequence and append regulatory sequences or promoter sequences or binding elements for trans-acting factors, or signal sequences to the requested sequence as part of the biological entity. Thus, when the operator knows that a sequence will be synthesized for later use in a particular host organism, the system can synthesize the requested sequences with the regulatory sequences or other desirable sequences particular to the host organism, thus further simplifying and speeding the preparation of the biological entity or biological product. In addition to being synthesized by the systems of the invention (and by the methods described herein) the regulatory sequences, promoter sequences, signal sequences, and binding elements for trans-acting factors can also be pre-charged into reagent vessels and/or reaction containers described herein as pre-made sequences for use as desired. These pre-made sequences can also be included in any kit described herein.

[0062] In one embodiment the systems of the invention are programmed to perform biosecurity screening in order to identify prohibited sequences. The biosecurity screening is done by comparing the sequence the system has received (in all 6 reading frames) and is requested to synthesize against a pre-programmed database of prohibited sequences. The prohibited sequences can be derived from a list of pathogens, biological weapons, or other biosecurity threats that are prohibited by a local government from being synthesized. The prohibited sequences can be one or more of DNA or RNA or proteins or peptides. The system can be programmed so that if a sequence is received that is identified on the list of prohibited sequences the system will be rendered inoperative or will otherwise not synthesize a prohibited sequence. Prohibited sequences can also be obtained from the International Gene Synthesis Consortium. The system can also screen against protein sequences derived from a prohibited sequences list.

[0063] Some embodiments of the systems of the invention perform modifications to DNA after synthesis, for example in a sub-unit of the system. In other embodiments the systems can also perform post-transcriptional and/or post-translational modifications. In one embodiment the

system produces a DNA molecule, which can then be modified to produce a modified DNA molecule. In one embodiment the modification is the methylation of DNA at particular nucleotides or nucleotide analogs. In another embodiment a sub-unit of a system of the invention can contain methyltransferases and/or have the capacity to perform phosphorylation or dephosphorylation on biological entities or biological products described herein. 2' fluoro and 2' O-methyl NTPs can also be produced enzymatically or chemically by the system to improve *in vivo* stability of DNA or RNA. In another embodiment the system produces a requested RNA molecule and adds a 7-methylguanosine residue to the 5' end as a 5' cap or a 5'-5' phosphate linkage as a cap, which the system can then methylate to form mature mCAP. Guanosine-5'-triphosphate-5'guanosine can also be used as an RNA cap. RNA bases can also be post-transcriptionally modified by the systems using a 2' O-methyl group to increase melting temperature and increase stability. In another embodiment the system produces a requested RNA molecule having a poly-A tail, with any number of A residues. Other embodiments include the inclusion of cleavage signals or sequences, or GUrich sequences in the biological entity, which also can be tailored to a particular host cell to be used in a subsequent procedure. In still more embodiments the systems of the invention produce a requested peptide, poly-peptide or protein sequence with

Example 1 - Synthesis of a Functional HA and NA DNA Molecules and Protein Moieties

[0064] This example illustrates the automated assembly of DNA constructs of HA and NA genes from an oligonucleotide pool and their transfection into a host cell that produces viral particles. Influenza viruses are made of a viral envelope containing glycoproteins wrapped around a central core. The central core contains the viral RNA genome and other viral proteins that package and protect the RNA. The influenza genome typically contains eight pieces of RNA with each containing one or two genes encoding viral proteins. In the case of influenza A, the genome contains 11 genes on eight pieces of RNA, encoding for 11 proteins, including hemagglutinin (HA) and neuraminidase (NA). Other proteins include nucleoprotein (NP), M1, M2, NS1, NS2, PA, PB 1, PB1-F2 and PB2.

[0065] Hemagglutinin (HA) and neuraminidase (NA) are glycoproteins present on the outside of the viral particles. These glycoproteins have key functions in the life cycle of the virus, including assisting in binding to host cells and reproduction of viral particles. The assembled virus containing these proteins is therefore useful in the production of a vaccine.

Oligonucleotide Synthesis and Assembly

[0066] A pool of 96 oligonucleotides representing the sequence of DNA constructs of the HA and NA genes were provided to an assembly unit of a system of the invention. The HA and NA constructs were approximately 3 kb in length and were assembled from 96 oligonucleotides in the method. The first and last oligonucleotides contained primer binding domains for PCR

amplification and NotI restriction sites to release the primer binding domains following amplification and expose overlapping regions for DNA assembly, if necessary to assemble larger fragments.

[0067] The assembly unit utilized a BIOMEK[®] NXP, Span-8 laboratory automation workstation (Beckman Instruments Inc., Fullerton, CA) with integrated thermal-cycling capabilities.

[0068] The assembly unit was programmed to perform several different steps in the process namely, 1) PCR1 amplification to amplify oligonucleotides; 2) an error correction step on the PCR1 amplified oligonucleotides; 3) a PCR2 step to amplify the corrected oligonucleotides; 4) a PCR product purification step to provide pure amplified oligonucleotides; 5) an assembly step to assemble the oligonucleotide products into a gene. Each process can be performed in a distinct reaction zone of the reaction container (which is a 96 well plate), and the reaction zone can be one or more columns on the 96 well plate. Assembly reaction is at 50 °C for 30-60 minutes and the reaction is temperature shifted and held at 10 °C thereafter.

1st PCR and error correction

[0069]

1. 1. For each assembled product PCR reactions were performed in automated fashion:

25 µl 2X PHUSION[®] Hot-Start Master Mix (Thermo Fisher Scientific Oy, Oy, FI)

2 µl 1% PEG 8000

0.25 µl Terminal Primer 1 (100 uM)

0.25 µl Terminal Primer 2 (100 uM)

20 µl MBG water

2.5 µl of the oligo pool above was transferred at 50 nM as template to a reaction zone of the reaction container containing PCR master mix (or combine subsequently).

2. 2. Thermal-cycle occurred using the following parameters:

98 °C for 1 min

30X (98 °C 30 sec, 65 C 6 minutes and extending that by 15 sec/cycle

72 °C for 5 min

10 °C forever

Error Correction

[0070] Denaturation/annealing by thermocycling using the following parameters:

98 °C for 2 min

2 °C/sec to 85 °C

85 °C for 2 min

0.1 °C/sec to 25 C

25 °C 2 min

10 °C forever

2.7 µl was removed and added to 8.3 ul of error correction mix comprising 5.3 ul water/2 ul SURVEYOR™ nuclease (Transgenomic, Inc., Omaha, NE) /1 ul 1:4000 diluted Exo III.

[0071] Thermo-cycled at 42 °C for 1 hour, 10 °C forever

2nd PCR

[0072] See above for Master Mix recipe.

[0073] 2.5 µl of error-corrected sample was used in a 50 ul total reaction volume.

[0074] See above for the thermal cycler conditions.

PCR Purification

[0075] PCR product was purified using the AMPURE® XP technology (Agencourt, Bioscience Corp. Beverly, MA)

GIBSON ASSEMBLY® (Synthetic Genomics, San Diego, CA) to combine subassemblies into HA and NA genes Within Plasmid Vectors.

[0076] Nucleic acid constructs of approximately 3 kb were produced. The electrophoretic gels

are shown in Fig. 2. These genes already include promoter regions (pol I and pol II) for expression following transfection into mammalian cells.

[0077] Genes were cloned into pUC19 using GIBSON ASSEMBLY[®] mix with the vector.

[0078] The following were combined:

5 µl of mix

5 µl of DNA amplicon

[0079] The mix was incubated within the reaction container on the thermo-cycler 50 °C for 1 hour, then 10 °C forever.

Transform Plasmid Vector DNA to Host Cells

[0080] The next step is to combine the genes with the constant influenza genes to provide the complete viral genome. The genes are then transfected into MDCK cells to produce a rescued virus. The cells then produce the virus product, which is ready for harvest.

Example 2 - Generation of Immunological Response

[0081] In a further experiment sequences were synthesized in a manner as described in Example 1 for the influenza A/Shanghai/2/2013 HA gene. The synthetic HA sequences were cloned into a DNA template containing RNA-dependent RNA polymerase genes, genetic control elements, and a T7 polymerase promoter according to previously described self-amplifying RNA vaccine technology (Geall et al., Nonviral delivery of self-amplifying RNA vaccines, PNAS, Vol. 109, No. 36, pp. 14604-09 (2012)). After verification of the construct *in vitro* transcription of the RNA, capping of the RNA, and transfection of BHK cells with the RNA was carried out. The integrity of the transcribed RNA was demonstrated by agarose gel electrophoresis (not shown). The cells expressed the H7 HA, as demonstrated by gel electrophoresis and Western blot with an H7-specific antibody. The RNA was encapsulated in a lipid nanoparticle (LNP) delivery system and the vaccine was used to immunize mice (Geall *et al.*, and Hekele *et al.*, Emerging Microbes and Infections (2013) 2, e52). Two weeks after the first injection, six of seven H7/LNP -immunized mice seroconverted with H7-specific hemagglutinin inhibition titers ranging from 1:20 to 1:80 (data not shown).

Example 3 - Synthesis of a Functional Phage Particle and Oligonucleotide design

[0082] The present invention can be applied to the production of a phage, which can be useful in phage therapy. In one embodiment the phage is a bacteriophage. Phage therapy is useful for the treatment of pathogenic bacterial infections, in humans, animals, and plants. Phage therapy can be particularly useful in applications where the bacteria does not respond to convention methods of control.

[0083] The 5,386 bp PhiX DNA biological sequence is entered into a transmitting unit of a system of the invention using ARCHETYPE[®] software (Synthetic Genomics, Inc., San Diego, CA) (Ref. Seq: NC_001422, sequence below) at a laboratory. The software divides the sequence into four overlapping fragments of about 1.4 kb each (sub-assemblies of the PhiX genome sequence). Each fragment overlaps the next by 40 bp to form a circular molecule when assembled. Each of the four fragments is further divided into 43-45 overlapping oligonucleotides. These oligonucleotides are about 64 bases in length with 32 bp overlaps. The first and last oligonucleotides of each of the four overlapping fragments contain primer binding domains for PCR amplification and NotI restriction sites to release the primer binding domains, following amplification, and expose overlapping regions for DNA assembly. Following entry the biological sequence information is transmitted to a receiving unit located in a laboratory in a remote city.

Oligonucleotide Synthesis

[0084] The biological sequence information is received by a receiving unit of a system of the invention, which in this embodiment is a computer connected to the same computer network as the transmitting unit. The receiving unit is connected to an assembly unit, which in this embodiment also has a BIOAUTOMATION[™] 192E oligonucleotide synthesizer (BioAutomation Corp., Plano, TX) as one sub-unit. All sub-units of the assembly unit are set up prior to receiving the biological sequence information. Setup includes, but is not limited to, charging all vessels with requisite chemicals, reagents, and biological building blocks, as well as preparing all software programming prior to receiving the biological sequence information so that activation of the synthesis of the biological entity can begin immediately upon receiving the sequence information. Each step of the method is performed in a distinct reaction zone of the reaction container.

[0085] After receipt of the biological sequence information, software within the assembly unit directs the synthesis of the oligonucleotides, which are synthesized using the biological building blocks previously provided to the system (in this case nucleoside phosphoramidite building blocks). This synthesis is performed on a sub-unit such as an oligonucleotide synthesizer.

[0086] After synthesis the oligonucleotides are cleaved from the solid support by the addition of concentrated ammonium hydroxide and incubated at 55 °C, each step done in automated fashion without further human intervention after setup. Oligonucleotides making up each fragment are separately pooled and purified by passing through a column containing

POLYPAK™ packing (Glen Research, Sterling, VA). Other packings can also be used that are stable in the pH range 1-13. Thus, the ammonium hydroxide solution, diluted with water, is loaded directly onto the packing. After elution of failure sequences, the trityl protecting group is removed and washed from the support-bound oligonucleotide. The fully deprotected product can then be eluted and isolated by lyophilization. Deprotection procedures suitable for automation are used, for example on-column deprotection using Ethylenediamine (EDA)/Toluene solution. A paramagnetic bead-based purification system can also be used, an example of which is AGENCOURT® COSMCPREP® (Agencourt Bioscience Corp., Beverly, MA). Following the above procedure the four pools of oligonucleotides making up each of the four fragments required to produce overlapping dsDNA fragments of PhiX are produced.

GIBSON ASSEMBLY® (Synthetic Genomics, San Diego, CA) to produce four overlapping PhiX subassemblies

[0087] The following steps are also performed in automated fashion by the assembly unit, which can have as many sub-units as necessary and convenient. Each sub-unit of the assembly unit is pre-charged with reagents as necessary to perform all steps. After oligonucleotide assembly a robotic arm is used to transfer the reaction container from the oligo synthesizer to a liquid handler with thermo-cycling capabilities. In a fully automated fashion, 10 µl of each of the four oligonucleotide pools above are added to 10 µl of PCR assembly reagents - one can utilize a dilution of about 10 µM. A convenient form of PCR reagents is 2X HF PHUSION® Mix (Thermo Fisher Scientific Oy, Oy, FI). The protocol is set forth below.

[0088] Assembly reaction is at 50 °C for 30-60 minutes and the reaction is temperature shifted and held at 10 °C thereafter.

1st PCR and error correction

[0089]

1. 1. For each assembled product PCR reactions are performed in automated fashion:

25.00 µl 2X HF PHUSION® Mix (Thermo Fisher Scientific Oy, Oy, FI)

0.25 µl Terminal Primer 1 (100 µM)

0.25 µl Terminal Primer 2 (100 µM)

22 µl Water

2.5 µl Template (GIBSON ASSEMBLY® (Synthetic Genomics, Inc., San Diego, CA) of the reaction product from above)

2. 2. Thermal-cycle occurs using the following parameters:

98 °C for 1 min

Cycle 25X {98 °C for 10 sec, 60 °C for 30 sec., 72 °C for 1.5 min)

72 °C for 5 min

98 °C for 2 min

Increase 2 °C /sec to 85 °C

85 °C for 2 min

Decrease 0.1 °C /sec to 25 °C

25 °C for 2 min

Hold at 10°C thereafter

3. 3. 6.25 µl of S/E error correction mix is added (protocol below) to each sample

4. 4. Error correction reaction is at 42 °C for 1 hour and then the reaction is temperature shifted and held at 10 °C thereafter.

2nd PCR

[0090]

1. 1. The following four PCR reactions are also performed in automated fashion by the thermo-cycling sub-unit of the assembly unit:

25.00 µl 2X HOTSTART-IT® Taq Master Mix (Affymetrix, Inc., Santa Clara, CA) or alternative (PHUSION® (Thermo Fisher Scientific Oy, Oy, FI) per reaction

0.25 µl Primer 1 (100 µM)

0.25 µl Primer 2 (100 µM)

22.00 µl MBG Water

2.50 µl Error corrected template from above.

GIBSON ASSEMBLY™ {Synthetic Genomics, San Diego, CA) to combine four subassemblies into phiX genome

[0091] The following procedure is performed in automated fashion:

1. 1. 2.5 µl of each of the four amplicons is removed from the 2nd PCR above and pooled into another coordinate of a 96-well plate.
2. 2. 2 µl of NotI restriction enzyme is added and incubated at 37 °C for 1 hour, then 65 °C for 20 minutes to inactivate the NotI enzyme.
3. 3. 12 µl 2X GIBSON ASSEMBLY™ (Synthetic Genomics, Inc., San Diego, CA) master mix is added and incubated at 50 °C for 1 hour.

Production of active PhiX virus

[0092] In this step active virus particles are produced. These steps are also performed in automated fashion. Following DNA synthesis and assembly as described above, the assembled ΦX174 genome is combined with chemically-competent *E. coli* strain C or another ΦX174-sensitive *E. coli* strain (e.g., DH10). Following the heat-shock at 42°C, cells are recovered in SOC medium.

1. 1. 5 µl of assembly reaction above is combined with 50 µl of chemically competent DH10 *E. coli* cells in a vessel.
2. 2. Thermal-cycling is performed according to the following conditions:
 - 4 °C for 2 min
 - 42 °C for 30 sec
 - 4 °C for 2 min
3. 3. 150 µl SOC medium is added and cells are incubated at 37 °C for several hours

[0093] Active phage particles are purified from *E. coli* and tested for capacity to form plaques when spread on LB agar plates containing *E. coli* cells (Smith et al, PNAS 2003).

[0094] As an alternative approach to transforming *E. coli* cells, it is also possible to test for the capacity to form active phage particles in a cell-free system. In this embodiment, synthetic genomes are incubated for 2 hours in one of the cell-free expression kits described above.

REAGENT COMPONENTS

[0095] Components for GIBSON ASSEMBLY® (Synthetic Genomics, Inc., San Diego, CA).

1. 1. 5X isothermal (ISO) reaction buffer (25% PEG-8000, 500 mM Tris-HCl pH 8.0, 50 mM MgCl₂, 50 mM DTT, 1 mM each of the 4 dNTPs, and 5 mM NAD). This is prepared as described below.
2. 2. T5 exonuclease
3. 3. PHUSION® High Fidelity DNA polymerase (Thermo Fisher Scientific Oy, Oy, FI)
4. 4. Taq DNA ligase

Procedure

[0096]

1. 1. 5X ISO buffer is prepared. Six ml of this buffer can be prepared by combining the following:

3 ml of 1 M Tris-HCl pH 8.0

300 µl of 1 M MgCl₂

600 µl of 10 mM dNTPs

300 µl of 1 M DTT (1.54 g dissolved in dH₂O up to 10 ml)

1.5 g PEG-8000

300 µl of 100 mM NAD (0.66 g dissolved in dH₂O up to 10 ml; resuspend by heating at 50 °C followed by continuous vortexing)

Water is added to 6 ml. Aliquot 1 ml and store at -20 °C.

2. 2. 800 µl of the assembly master mixture is prepared, sufficient for 80 reactions. This can be prepared by combining the following:

320 µl 5X ISO buffer

6.4 µl of 1 U/µl T5 exo (diluted 1:10 from enzyme stock in 1x T5 exo buffer) 20 µl of 2 U/µl PHUSION® High Fidelity DNA polymerase (Thermo Fisher

Scientific Oy, Oy, FI)

80 µl of 40 U/µl Taq ligase

374 µl dH₂O

The mixture is mixed well and stored at -20 °C, or on ice if to be used immediately.

3. 3. The assembly mixture can be stored at -20 °C for at least one year. The enzymes remain active following at least 10 freeze-thaw cycles.

[0097] The mixture is ideal for the assembly of DNA molecules with 20-150 bp overlaps.

ERROR CORRECTION MIX COMPONENTS-SURVEYOR NUCLEASE+ EXONUCLEASE III (S/E)

[0098] 250 µl of SURVEYOR™ Nuclease (Transgenomic Inc., Omaha, NE) + 0.03125 µl Exonuclease III

PhiX genome sequence:

[0099] SEQ ID NO: 5

GAGTTTTATCGCTTCCATGACGCAGAAGTTAACTTTCGGATATTTCTGATGAG
TCGAAAAATTATCTTGATAAAGCAGGAATTACTACTGCTTGTTACGAATTAAAT
CGAAGTGGACTGCTGGCGGAAAATGAGAAAATTCGACCTATCCTTGCGCAGCTC
GAGAAGCTCTTACTTTGCGACCTTTCGCCATCAACTAACGATTCTGTCAAAAAC
GACGCGTTGGATGAGGAGAAGTGGCTTAATATGCTTGGCACGTTTCGTCAAGGAC
TGGTTTAGATATGAGTCACATTTTGTTCATGGTAGAGATTCTCTTGTTGACATTTT
AAAAGAGCGTGGATTACTATCTGAGTCCGATGCTGTTCAACCACTAATAGGTAAG
AAATCATGAGTCAAGTTACTGAACAATCCGTACGTTTCCAGACCGCTTTGGCCTC
TATTAAGCTCATTGAGGCTTCTGCCGTTTTGGATTAAACCGAAGATGATTCGATT
TTCTGACGAGTAACAAAGTTTGGATTGCTACTGACCGCTCTCGTGCTCGTCGCTG
CGTTGAGGCTTGCGTTTATGGTACGCTGGACTTTGTGGGATACCCTCGCTTTCCTG
CTCCTGTTGAGTTTATTGCTGCCGTCATTGCTTATTATGTTTCATCCCGTCAACATT

CAAACGGCCTGTCTCATCATGGAAGGCGCTGAATTTACGGAAAACATTATTAATG
GCGTCGAGCGTCCGGTTAAAGCCGCTGAATTGTTTCGCGTTTACCTTGCGTGTACG
CGCAGGAAACACTGACGTTCTTACTGACGCAGAAGAAAACGTGCGTCAAAAATT
ACGTGCGGAAGGAGTGATGTAATGTCTAAAGGTAAAAACGTTCTGGCGCTCGC
CCTGGTCGTCCGCAGCCGTTGCGAGGTACTAAAGGCAAGCGTAAAGGCGCTCGT
CTTTGGTATGTAGGTGGTCAACAATTTTAATTGCAGGGGCTTCGGCCCCCTTACTTG
AGGATAAATTATGTCTAATATTCAAACCTGGCGCCGAGCGTATGCCGCATGACCTT
TCCCATCTTGCTTCCCTTGCTGGTCAGATTGGTCGTCTTATTACCATTTCAACTAC
TCCGGTTATCGCTGGCGACTCCTTCGAGATGGACGCCGTTGGCGCTCTCCGTCTTT

CTCCATTGCGTCGTGGCCTTGCTATTGACTCTACTGTAGACATTTTACTTTTTAT
GTCCCTCATCGTCACGTTTATGGTGAACAGTGGATTAAAGTTCATGAAGGATGGTG
TTAATGCCACTCCTCTCCCCGACTGTAAACACTACTGGTTATATTGACCATGCCGCT
TTTCTTGGCACGATTAACCCTGATACCAATAAAATCCCTAAGCATTTGTTTCAGG
GTTATTTGAATATCTATAACAACATTTTTAAAGCGCCGTGGATGCCTGACCGTAC
CGAGGCTAACCCTAATGAGCTTAATCAAGATGATGCTCGTTATGGTTTCCGTTGC
TGCCATCTCAAAAACATTTGGACTGCTCCGCTTCCTCCTGAGACTGAGCTTTCTCG
CCAAATGACGACTTCTACCACATCTATTGACATTATGGGTCTGCAAGCTGCTTAT
GCTAATTTGCATACTGACCAAGAACGTGATTACTTCATGCAGCGTTACCATGATG
TTATTTCTTCATTTGGAGGTAAAACCTCTTATGACGCTGACAACCGTCCTTTACTT
GTCATGCGCTCTAATCTCTGGGCATCTGGCTATGATGTTGATGGAACGACCAAA
CGTCGTTAGGCCAGTTTTCTGGTCGTGTTCAACAGACCTATAAACATTCTGTGCC
GCGTTTCTTTGTTCCCTGAGCATGGCACTATGTTTACTCTTGCGCTTGTTTCGTTTTCC
GCCTACTGCGACTAAAGAGATTTCAGTACCTTAACGCTAAAGGTGCTTTGACTTAT
ACCGATATTGCTGGCGACCCTGTTTTGTATGGCAACTTGCCGCCGCGTGAAATTT
CTATGAAGGATGTTTTCCGTTCTGGTGATTCGTCTAAGAAGTTTAAGATTGCTGA
GGGTCAGTGGTATCGTTATGCGCCTTCGTATGTTTCTCCTGCTTATCACCTTCTTG
AAGGCTTCCCATTCAATCAGGAACCGCCTTCTGGTGATTTGCAAGAACGCGTACT
TATTCGCCACCATGATTATGACCAGTGTTCAGTCCGTTTCAGTTGTTGCAGTGGA
ATAGTCAGGTAAATTTAATGTGACCGTTTATCGCAATCTGCCGACCACTCGCGA
TTCAATCATGACTTCGTGATAAAAGATTGAGTGTGAGGTTATAACGCCGAAGCGG
TAAAAATTTTAATTTTTGCCGCTGAGGGGTTGACCAAGCGAAGCGCGGTAGGTTT
TCTGCTTAGGAGTTTAATCATGTTTCAGACTTTTATTTCTCGCCATAATTCAAAC
TTTTTTCTGATAAGCTGGTTCTCACTTCTGTTACTCCAGCTTCTTCGGCACCTGTTT

TACAGACACCTAAAGCTACATCGTCAACGTTATATTTTGATAGTTTGACGGTTAA
TGCTGGTAAATGGTGGTTTTCTTCATTGCATTCAGATGGATACATCTGTCAACGCCG
CTAATCAGGTTGTTTCTGTTGGTGCTGATATTGCTTTTGATGCCGACCCTAAATTT
TTTGCTGTTTGGTTTCGCTTTGAGTCTTCTTCGGTCCGACTACCCTCCCAGCTGC
CTATGATGTTTATCCTTTGAATGGTCGCCATGATGGTGGTTATTATACCGTCAAGG
ACTGTGTGACTATTGACGTCCTTCCCCGTACGCCGGGCAATAACGTTTATGTTGG
TTTCATGGTTTGGTCTAACTTTACCGCTACTAAATGCCGCGGATTGGTTTCGCTGA
ATCAGGTTATTAAAGAGATTATTTGTCTCCAGCCACTTAAGTGAGGTGATTTATG
TTTGGTGCTATTGCTGGCGGTATTGCTTCTGCTCTTGCTGGTGGCGCCATGTCTAA
ATTGTTTGGAGGCGGTCAAAAAGCCGCCTCCGGTGGCATTCAAGGTGATGTGCTT
GCTACCGATAACAATACTGTAGGCATGGGTGATGCTGGTATTAAATCTGCCATTC
AAGGCTCTAATGTTCCCTAACCCCTGATGAGGCCGCCCTAGTTTTGTTTCTGGTGCT
ATGGCTAAAGCTGGTAAAGGACTTCTTGAAGGTACGTTGCAGGCTGGCACTTCTG
CCGTTTCTGATAAGTTGCTTGATTGTTGGTGGACTTGGTGGCAAGTCTGCCGCTGAT
AAAGGAAAGGATACTCGTGATTATCTTGCTGCTGCATTTCTGAGCTTAATGCTT
GGGAGCGTGCTGGTGCTGATGCTTCCTCTGCTGGTATGGTTGACGCCGGATTGTA

GAATCAAAAAGAGCTTACTAAAATGCAACTGGACAATCAGAAAGAGATTGCCGA
GATGCAAAATGAGACTCAAAAAGAGATTGCTGGCATTTCAGTCGGCGACTTCACG
CCAGAATACGAAAGACCAGGTATATGCACAAAATGAGATGCTTGCTTATCAACA
GAAGGAGTCTACTGCTCGCGTTGCGTCTATTATGGAAAACACCAATCTTTCCAAG
CAACAGCAGGTTTCCGAGATTATGCGCCAAATGCTTACTCAAGCTCAAACGGCTG
GTCAGTATTTTACCAATGACCAAATCAAAGAAATGACTCGCAAGGTTAGTGCTGA
GGTTGACTTAGTTCATCAGCAAACGCAGAATCAGCGGTATGGCTCTTCTCATATT
GGCGCTACTGCAAAGGATATTTCTAATGTCGTCCTGATGCTGCTTCTGGTGTGG
TTGATATTTTTTCATGGTATTGATAAAGCTGTTGCCGATACTTGGAACAATTTCTGG
AAAGACGGTAAAGCTGATGGTATTGGCTCTAATTTGTCTAGGAAATAACCGTCAG
GATTGACACCCTCCCAATTGTATGTTTTTCATGCCTCCAAATCTTGAGGCTTTTTT
ATGGTTCGTTCTTATTACCCTTCTGAATGTCACGCTGATTATTTTGACTTTGAGCG
TATCGAGGCTCTTAAACCTGCTATTGAGGCTTGTTGGCATTCTACTCTTCTCAAT
CCCCAATGCTTGGCTTCCATAAGCAGATGGATAACCGCATCAAGCTCTTGGAAGA
GATTCTGTCTTTTCGTATGCAGGGCGTTGAGTTCGATAATGGTGATATGTATGTTG
ACGGCCATAAGGCTGCTTCTGACGTTCTGTGATGAGTTTGTATCTGTTACTGAGAA
GTTAATGGATGAATTGGCACAATGCTACAATGTGCTCCCCCACTTGATATTAAT

AACACTATAGACCACCGCCCCGAAGGGGACGAAAAATGGTTTTTAGAGAACGAG
AAGACGGTTACGCAGTTTTGCCGCAAGCTGGCTGCTGAACGCCCTCTTAAGGATA
TTCGCGATGAGTATAATTACCCCAAAAAGAAAGGTATTAAGGATGAGTGTTCAA
GATTGCTGGAGGCCTCCACTATGAAATCGCGTAGAGGCTTTGCTATTCAGCGTTT
GATGAATGCAATGCGACAGGCTCATGCTGATGGTTGGTTTATCGTTTTTGACACT
CTCACGTTGGCTGACGACCGATTAGAGGCGTTTTATGATAATCCCAATGCTTTCG
GTGACTATTTTCGTGATATTGGTTCGTATGGTTCTTGCTGCCGAGGGTCGCAAGGC
TAATGATTCACACGCCGACTGCTATCAGTATTTTTGTGTGCCTGAGTATGGTACA
GCTAATGGCCGTCTTCATTTCCATGCGGTGCACTTTATGCGGACACTTCCTACAG
GTAGCGTTGACCCTAATTTTGGTTCGTGCGGTACGCAATCGCCGCCAGTTAAATAG
CTTGCAAAATACGTGGCCTTATGGTTACAGTATGCCCATCGCAGTTCGCTACACG
CAGGACGCTTTTTTCACGTTCTGGTTGGTTGTGGCCTGTTGATGCTAAAGGTGAGC
CGCTTAAAGCTACCAGTTATATGGCTGTTGGTTTCTATGTGGCTAAATACGTAA
CAAAAAGTCAGATATGGACCTTGCTGCTAAAGGTCTAGGAGCTAAAGAATGGAA
CAACTCACTAAAAACCAAGCTGTCGCTACTTCCCAAGAAGCTGTTCAGAATCAGA
ATGAGCCGCAACTTCGGGATGAAAATGCTCACAATGACAAATCTGTCCACGGAG
TGCTTAATCCAACTTACCAAGCTGGGTTACGACGCGACGCCGTTCAACCAGATAT
TGAAGCAGAACGCAAAAAGAGAGATGAGATTGAGGCTGGGAAAAGTTACTGTA
GCCGACGTTTTTGGCGGGCGCAACCTGTGACGACAAATCTGCTCAAATTTATGCGCG
CTTCGATAAAAATGATTGGCGTATCCAACCTGCA

[0100] SEQ ID NO: 1 - Fragment # 1:

GAGTTTTATCGCTTCCATGACGCAGAAGTTAACACTTTCGGATATTTCTGATGAG
TCGAAAAATTATCTTGATAAAGCAGGAATTACTACTGCTTGTTTACGAATTAAAT
CGAAGTGGACTGCTGGCGGAAAATGAGAAAATTCGACCTATCCTTGCGCAGCTC
GAGAAGCTCTTACTTTGCGACCTTTCGCCATCAACTAACGATTCTGTCAAAAAC
GACGCGTTGGATGAGGAGAAGTGGCTTAATATGCTTGGCACGTTTCGTCAAGGAC
TGGTTTAGATATGAGTCACATTTTGTTCATGGTAGAGATTCTCTTGTTGACATTTT
AAAAGAGCGTGGATTACTATCTGAGTCCGATGCTGTTCAACCACTAATAGGTAAG
AAATCATGAGTCAAGTTACTGAACAATCCGTACGTTTCCAGACCGCTTTGGCCTC
TATTAAGCTCATTACAGGCTTCTGCCGTTTTGGATTAAACCGAAGATGATTTTCGATT
TTCTGACGAGTAACAAAGTTTGGATTGCTACTGACCGCTCTCGTGCTCGTCGCTG
CGTTGAGGCTTGCCTTATGGTACGCTGGACTTTGTGGGATACCCTCGCTTTCCTG
CTCCTGTTGAGTTTATTGCTGCCGTCATTGCTTATTATGTTTCATCCCGTCAACATT

CAAACGGCCTGTCTCATCATGGAAGGCGCTGAATTTACGGAACATTATTAATG
GCGTCGAGCGTCCGTTAAAGCCGCTGAATTGTTTCGCGTTTACCTTGCGTGTACG
CGCAGGAAACACTGACGTTCTTACTGACGCAGAAGAAAACGTGCGTCAAAAATT
ACGTGCGGAAGGAGTGATGTAATGTCTAAAGGTAAAAACGTTCTGGCGCTCGC
CCTGGTCGTCCGCAGCCGTTGCGAGGTAATAAGGCAAGCGTAAAGGCGCTCGT
CTTTGGTATGTAGGTGGTCAACAATTTTAATTGCAGGGGCTTCGGCCCCCTTACTTG
AGGATAAATTATGTCTAATATTCAAACCTGGCGCCGAGCGTATGCCGCATGACCTT
TCCCATCTTGCTTTCCTTGCTGGTCAGATTGGTCGTCTTATTACCATTTCAACTAC
TCCGGTTATCGCTGGCGACTCCTTCGAGATGGACGCCGTTGGCGCTCTCCGTCTTT
CTCCATTGCGTCGTGGCCTTGCTATTGACTCTACTGTAGACATTTTTACTTTTTAT
GTCCCTCATCGTCACGTTTATGGTGAACAGTGGATTAAGTTCATGAAGGATGGTG
TTAATGCCACTCCTCTCCCGACTGTAAACACTACTGGTTATATTGACCATGCCGCT
TTTCTTGGCACGATTAACCCTGATACCAATAAAATCCCTAAGCATTGTTTCAGG
GTTATTTGAATATCTATAACA

[0101] SEQ ID NO: 2 - Fragment # 2:

CCTAAGCATTTGTTTCAGGGTTATTTGAATATCTATAACAACCTATTTTAAAGCGCC
GTGGATGCCTGACCGTACCGAGGCTAACCTAATGAGCTTAATCAAGATGATGCT
CGTTATGGTTTCCGTTGCTGCCATCTCAAAAACATTTGGACTGCTCCGCTTCCTCC
TGAGACTGAGCTTCTCGCCAAATGACGACTTCTACCACATCTATTGACATTATG
GGTCTGCAAGCTGCTTATGCTAATTTGCATACTGACCAAGAACGTGATTACTTCA
TGCAGCGTTACCATGATGTTATTTCTTCATTTGGAGGTAAAACCTCTTATGACGCT
GACAACCGTCCTTTACTTGTCATGCGCTCTAATCTCTGGGCATCTGGCTATGATGT
TGATGGAACCTGACCAAACGTCGTTAGGCCAGTTTTCTGGTCGTGTTCAACAGACC
TATAAACATTCTGTGCCGCGTTTCTTTGTTTCTGAGCATGGCACTATGTTTACTCT
TGCGCTTGTTTCGTTTTCCGCCTACTGCGACTAAAGAGATTCAGTACCTTAACGCTA

AAGGTGCTTTGACTTATACCGATATTGCTGGCGACCCTGTTTTGTATGGCAACTTG
CCGCCGCGTGAAATTTCTATGAAGGATGTTTTCCGTTCTGGTGATTTCGTCTAAGA
AGTTTAAGATTGCTGAGGGTCAGTGGTATCGTTATGCGCCTTCGTATGTTTCTCCT
GCTTATCACCTTCTTGAAGGCTTCCCATTCATTCAGGAACCGCCTTCTGGTGATT
GCAAGAACGCGTACTTATTCGCCACCATGATTATGACCAGTGTTTCCAGTCCGTT
CAGTTGTTGCAGTGGAATAGTCAGGTTAAATTTAATGTGACCGTTTATCGCAATC
TGCCGACCACTCGCGATTCAATCATGACTTCGTGATAAAAGATTGAGTGTGAGGT
TATAACGCCGAAGCGGTAAAAATTTAATTTTTGCCGCTGAGGGGTTGACCAAGC
GAAGCGCGGTAGGTTTTCTGCTTAGGAGTTTAATCATGTTTCAGACTTTTATTTCT
CGCCATAATTCAAACCTTTTTTCTGATAAGCTGGTTCCTCACTTCTGTTACTCCAGC
TTCTTCGGCACCTGTTTTACAGACACCTAAAGCTACATCGTCAACGTTATATTTTG
ATAGTTTGACGGTTAATGCTGGTAATGGTGGTTTTCTTCATTGCATTGAGATGGAT
ACATCTGTCAACGCCGCTAATCAGGTTGTTTCTGTTGGTGCTGATATTGCTTTTGA
TGCCGACCCTAAATTTTTTGCCTGTTTGGTTCGCTTTGAGTCTTCTTCGGTTCGGA
CTACCCCTCCCGACTGCCTATGATGTTTATCCTTTGAATGGTCGCCATGATGGTGGT
TATTATACC

[0102] SEQ ID NO: 3 - Fragment # 3:

TTATCCTTTGAATGGTCGCCATGATGGTGGTTATTATACCGTCAAGGACTGTGTG
ACTATTGACGTCCTTCCCCGTACGCCGGGCAATAACGTTTATGTTGGTTTTCATGGT
TTGGTCTAACTTTACCGCTACTAAATGCCGCGGATTGGTTTCGCTGAATCAGGTT
ATTAAAGAGATTATTTGTCTCCAGCCACTTAAGTGAGGTGATTTATGTTTGGTGCT
ATTGCTGGCGGTATTGCTTCTGCTCTTGCTGGTGGCGCCATGTCTAAATTGTTTGG
AGGCGGTCAAAAAGCCGCCTCCGGTGGCATTCAAGGTGATGTGCTTGCTACCGAT
AACAACTACTGTAGGCATGGGTGATGCTGGTATTAAATCTGCCATTCAAGGCTCTA
ATGTTCCCTAACCCCTGATGAGGCCGCCCTAGTTTTGTTTCTGGTGCTATGGCTAAA
GCTGGTAAAGGACTTCTTGAAGGTACGTTGCAGGCTGGCACTTCTGCCGTTTCTG
ATAAGTTGCTTGATTTGGTTGGACTTGGTGGCAAGTCTGCCGCTGATAAAGGAAA
GGATACTCGTGATTATCTTGCTGCTGCATTCCTGAGCTTAATGCTTGGGAGCGTG
CTGGTGCTGATGCTTCCTCTGCTGGTATGGTTGACGCCGGATTTGAGAATCAAAA
AGAGCTTACTAAAATGCAACTGGACAATCAGAAAGAGATTGCCGAGATGCAAAA
TGAGACTCAAAAAGAGATTGCTGGCATTGAGTCGGCGACTTCACGCCAGAATAC
GAAAGACCAGGTATATGCACAAAATGAGATGCTTGCTTATCAACAGAAGGAGTC
TACTGCTCGCGTTGCGTCTATTATGGAAAACACCAATCTTCCAAGCAACAGCAG
GTTTCCGAGATTATGCGCCAAATGCTTACTCAAGCTCAAACGGCTGGTCAGTATT
TTACCAATGACCAAATCAAAGAAATGACTCGCAAGGTTAGTGCTGAGGTTGACTT
AGTTCATCAGCAAACGCAGAATCAGCGGTATGGCTCTTCTCATATTGGCGCTACT
GCAAAGGATATTTCTAATGTCGTCACTGATGCTGCTTCTGGTGTGGTTGATATTTT

TCATGGTATTGATAAAGCTGTTGCCGATACTTGGAACAATTTCTGGAAAGACGGT
 AAAGCTGATGGTATTGGCTCTAATTTGTCTAGGAAATAACCGTCAGGATTGACAC
 CCTCCCAATTGTATGTTTTTCATGCCTCCAAATCTTGGAGGCTTTTTTATGGTTCGT
 TCTTATTACCCTTCTGAATGTCACGCTGATTATTTTGACTTTGAGCGTATCGAGGC
 TCTTAAACCTGCTATTGAGGCTTGTGGCATTCTACTCTTTCTCAATCCCCAATGC
 TTGGCTTCCATAAGCAGAT

[0103] SEQ ID NO: 4 - Fragment #4:

CTCTTTCTCAATCCCCAATGCTTGGCTTCCATAAGCAGATGGATAACCGCATCAA
 GCTCTTGGAAGAGATTCTGTCTTTTCGTATGCAGGGCGTTGAGTTCGATAATGGT
 GATATGTATGTTGACGGCCATAAGGCTGCTTCTGACGTTTCGTGATGAGTTTGTAT
 CTGTTACTGAGAAGTTAATGGATGAATTGGCACAATGCTACAATGTGCTCCCCCA
 ACTTGATATTAATAAACAATAGACCACCGCCCCGAAGGGGACGAAAAATGGTT
 TTAGAGAACGAGAAGACGGTTACGCAGTTTTGCCGCAAGCTGGCTGCTGAACG
 CCCTCTTAAGGATATTCGCGATGAGTATAATTACCCCAAAAAGAAAGGTATTAAG
 GATGAGTGTTCAAGATTGCTGGAGGCCTCCACTATGAAATCGCGTAGAGGCTTTG
 CTATTCAGCGTTTGATGAATGCAATGCGACAGGCTCATGCTGATGGTTGGTTTAT
 CGTTTTTGACACTCTCACGTTGGCTGACGACCGATTAGAGGCGTTTTATGATAAT
 CCCAATGCTTTGCGTGACTATTTTCGTGATATTGGTCGTATGGTTCTTGCTGCCGA
 GGGTCGCAAGGCTAATGATTCACACGCCGACTGCTATCAGTATTTTTGTGTGCCT
 GAGTATGGTACAGCTAATGGCCGTCTTCATTTCCATGCGGTGCACTTTATGCGGA
 CACTTCCTACAGGTAGCGTTGACCCTAATTTTGGTCGTCGGGTACGCAATCGCCG
 CCAGTTAAATAGCTTGCAAATAACGTGGCCTTATGGTTACAGTATGCCCATCGCA
 GTTCGCTACACGCAGGACGCTTTTTACGTTCTGGTTGGTTGTGGCCTGTTGATGC
 TAAAGGTGAGCCGCTTAAAGCTACCAGTTATATGGCTGTTGGTTTCTATGTGGCT
 AAATACGTTAACAAAAAGTCAGATATGGACCTTGCTGCTAAAGGTCTAGGAGCT
 AAAGAATGGAACAACCTCACTAAAAACCAAGCTGTCGCTACTTCCCAAGAAGCTG
 TTCAGAATCAGAATGAGCCGCAACTTCGGGATGAAAATGCTCACAATGACAAAT
 CTGTCCACGGAGTGCTTAATCCAACCTACCAAGCTGGGTTACGACGCGACGCCGT
 TCAACCAGATATTGAAGCAGAACGCAAAAAGAGAGATGAGATTGAGGCTGGGA
 AAAGTTACTGTAGCCGACGTTTTGGCGGCGCAACCTGTGACGACAAATCTGCTCA
 AATTTATGCGCGCTTCGATAAAAAATGATTGGCGTATCCAACCTGCAGAGTTTTAT
 CGCTTCCATGACGCAGAAGTTAACACTTTCG

Example 4 - Synthesis of a Functional DNA Molecule and Protein Moiety

[0104] The nucleic acid sequence encoding a functional green fluorescent protein (GFP) and

appropriate regulatory sequences for expression is entered into ARCHETYPE[®] (Synthetic Genomics, Inc., San Diego, CA) or another software that divides the sequence into about 60 bp oligonucleotides with about 30 bp overlaps. The first and last oligonucleotides contain primer binding domains for PCR amplification and NotI restriction sites to release the primer binding domains following amplification and expose overlapping regions for DNA assembly, if necessary to assemble larger fragments.

[0105] The system includes a BIOMEK[®] NXP, Span-8 laboratory automation workstation (Beckman Instruments Inc., Fullerton, CA) with integrated thermal-cycling capabilities as one sub-unit. Additional sub-units include an automated *in vitro* translation system containing vessels with reagents for carrying out a cell-free translation of nucleic acid into protein. Following entry into the transmitting unit the biological sequence information is transmitted to a receiving unit located in a laboratory in a remote city.

Oligonucleotide Synthesis

[0106] Following a synthesis plan similar to those described above the biological sequence information is received by a receiving unit of a system of the invention, as described in the Examples. In this embodiment the receiving unit is a computer connected to the same computer network as the transmitting unit. The receiving unit is connected to an assembly unit, which in this embodiment has a BIOAUTOMATION[™] 192E oligonucleotide synthesizer (BioAutomation Corp., Plano, TX) as one sub-unit. The additional sub-units of the assembly unit, including the sub-unit for *in vitro* translation, are all set up prior to receiving the biological sequence information by charging all vessels with requisite chemicals, reagents, and biological building blocks, as well as preparing all software programming prior to receiving the biological sequence information so that activation of the synthesis of the biological entity can begin immediately upon receiving the sequence information.

[0107] After receipt of the biological sequence information software within the assembly unit directs the synthesis of the oligonucleotides, which are synthesized using dNTPs or phosphoramidites and other reagents previously provided to the system. This step is done in the same manner as described above in the Examples and the set of oligonucleotides are produced, which are then assembled and amplified as described in the Examples to produce the nucleic acid sequence encoding the GFP gene with appropriate regulatory sequences necessary for expression.

[0108] *In vitro* and automated transcription and translation can be performed on the assembled nucleic acid sequence in a distinct reaction zone of the reaction container prepared for that purpose. Various kits are commercially available for performing *in vitro* translation and/or cell-free protein expression and can be conveniently employed in the present invention, e.g., PIERCE[®] (Pierce Biotechnology Inc., Rockford, IL) *In Vitro* Protein Expression Systems

available from THERMO SCIENTIFIC® (Thermo Fisher Scientific, Inc., Waltham, MA) or the T7 RiboMAX™ Express kit from PROMEGA® (Promega Corp, Madison, WI), which can generate 27 kb *in vitro* transcripts in minutes. Different kits are available depending on the type and size of protein to be translated. The reaction zone of the reaction container where the reaction is to be performed is pre-charged with all necessary reagents. The GFP construct was also designed to contain a T7 or a strong *E. coli* promoter upstream of the coding sequence, a ribosome binding site with a spacing sequence of 5-7 bp upstream of the AUG translational start point and an untranslated region downstream of the stop coding.

[0109] The synthetic PCR product is added to the protein expression components in the designated reaction zone. Following the expression reaction, a black light is held to the reaction mixture and the presence of translated GFP is confirmed by the emission of green light.

Example 5 - Synthesis of an Antigenic Biological Product

[0110] HER-2 (human epidermal growth factor receptor 2) is a protein for which amplification or over-expression play a role in the progression of certain breast cancers. The protein is encoded by the ERBB2 gene. A number of antibody therapies are available that target HER-2. A vaccine can be made containing the HER-2 antigen, which can be administered to a patient who will then form an antigenic response to the vaccine and produce antibodies to HER-2. The vaccine can be useful in cancer treatments, and in any therapy where the production of HER-2 antibodies finds useful application.

[0111] The sequence of HER-2 protein is entered into a software program that divides the sequence into an appropriate number of overlapping fragments representing subassemblies of the gene. Suitable regulatory sequences are included in the sequence. Each fragment is of appropriate length for gene synthesis and is designed to overlap the next fragment by an appropriate number of base pairs such as, for example, about 40 bp. The biological sequence information is transmitted by a transmitting unit to a receiving unit of a system of the invention. As explained herein the gene can be assembled as a series of overlapping oligonucleotides, and a sub-unit of the assembly unit assembles the oligonucleotides into the final dsDNA molecule, or the dsDNA molecule can be assembled in a different zone of the reaction container. Suitable regulatory sequences for expression are included. The assembled gene is transferred to another sub-unit of the assembly unit or to another zone of the reaction container and cell-free, *in vitro* transcription is performed, and then to another sub-unit or another zone for translation of the DNA sequence, as explained herein. A purification step is performed if desired. The harvested HER-2 may be formulated and is utilized as an antigenic vaccine.

Example 6 - Synthesis of a DNA Vaccine

[0112] The invention also finds use in the production of a DNA vaccine. The DNA sequence encoding for one or more HIV proteins is entered into a software program that separates the DNA sequence into overlapping oligonucleotide fragments of appropriate length, including DNA sequences of an appropriate plasmid (e.g., pGA2/JS2) with appropriate regulatory sequences for expression.

[0113] The sequence is transmitted by a transmitting unit of a system of the invention to a receiving unit of a system of the invention. As described herein, the receiving unit provides the sequence to an assembly unit of a system of the invention and the assembly unit begins synthesis of overlapping oligonucleotides, which will be assembled by a subunit of the assembly unit (or by the same sub-unit in a different zone of the reaction container) into the full plasmid vaccine. DNA may then be formulated and used as a DNA vaccine. When injected into the patient the DNA vaccine will provide instructions to cells to make one or more HIV proteins, which will then provoke an immune response. The technique is applicable to any DNA virus that can be incorporated into a plasmid or other vector that can be injected into the patient to be treated to provide immunity.

[0114] West Nile virus is another example of a pathogen that is susceptible to prevention through the use of a DNA vaccine using the prM and E genes.

Example 7 - Synthesis of a Vaccine From Protein Sub-Units

[0115] Sub-unit vaccines offer an opportunity to present an antigen to the immune system without the need to introduce whole viral particles. One example is the OPTAFLU® (Novartis Vaccines and Diagnostics, GmbH, Marburg, Germany) vaccine for providing immunity to the influenza virus. The present invention can be applied in the rapid production of sub-unit vaccines.

[0116] The A/H3N2, A/H1N1 and B strains of the current flu virus are obtained from the World Health Organization strain recommendations and are used in the preparation of a trivalent vaccine using the viral surface antigens. The nucleic acid sequences are separated into fragments as described above using an appropriate software program, with appropriate regulatory sequences included. After synthesis of the nucleic acid, as described above, the nucleic acid is transferred to a sub-unit of the assembly unit (or to another zone of the reaction container) for *in vitro* translation. The vaccine can be produced in a sub-unit where a cell culture is being maintained, which sub-unit has been previously prepared with an established culture of MDCK cells (a canine kidney cell line) in appropriate media.

Example 8 - Synthesis of a Virus-Like Particle Vaccine

[0117] Non-infectious virus-like particles (VLPs) can be used to manufacture a vaccine with the capacity to activate the immune system without the need to present an assembled viral particle. A vaccine against human papilloma virus (HPV) can be manufactured using the L1 and/or L2 major capsid proteins, which self-assemble into virus-like particles that are effective as antigens. These VLPs can be assembled in yeast, insect cells, mammalian cells, or bacteria. An example of this type of vaccine is GARDASIL[®] (Merck Sharp & Dohme Corp., Whitehouse Station, NJ). Multiple HPV L1 virus-like particles can be included in a single vaccine for the broadest spectrum of immunity.

[0118] VLPs of a combination of HPV types selected from types 6, 11, 16, 18, 31, 35, 45, 52, and 58 are selected. The VLPs can be of L1 or L1 + L2 protein. The nucleic acid sequences coding for these proteins are identified and separated into fragments as described above using an appropriate software with appropriate regulatory sequences for expression in the chosen cell type included. The nucleic acid sequence information is then transmitted by a transmitting unit to a receiving unit of a system of the invention located in a remote location. The receiving unit provides the sequence information to the assembly unit of the system, and the VLPs are synthesized. As described above, the DNA molecule can be assembled from overlapping oligonucleotide fragments into one or more whole dsDNA molecules.

[0119] After synthesis, as described above, the DNA is transferred to a sub-unit of the assembly unit (or to another zone of the reaction container) for *in vitro* translation in an established cell culture being maintained on the system. The vaccine is produced by the cells in that sub-unit. The translated proteins are provided to another sub-unit of the system where they are pooled and self-assemble into VLPs that are useful as a vaccine against HPV.

Example 9 - Synthesis of a Complete Viral Vaccine

[0120] In another embodiment a whole virus vaccine can be synthesized in cell culture using a reverse genetics technique. The influenza A genome is composed of eight viral gene segments, including the HA and NA segments, which are important in immune response to the virus. In one embodiment the assembly unit has a subunit maintaining, for example, MDCK cells, 293T cells, or Vero cells for virus production (these components can also be maintained in one or more additional zones of the reaction container). Plasmids containing multiple transcription cassettes can be constructed containing the eight viral genes and necessary regulatory sequences. The eight genes can be present on multiple plasmids or a single plasmid if desired. The HA and NA segments are based upon sequences derived from a virus that is presenting a local threat. Thus, in one embodiment the six non-varying genes are prepared ahead of time and only the HA and NA genes are synthesized by the system. The six non-varying genes can then be treated as reagents in the method. But in another embodiment a subunit of the assembly unit synthesizes the plasmids based upon information received from the receiving unit. In either embodiment, after synthesis the plasmids are transfected into the cell culture being maintained in another subunit of the assembly unit under conditions for high

transfection efficiency. The transfected cells produce whole virus particles which can be formulated as a live, attenuated vaccine, or as a killed or further inactivated vaccine, as desired.

[0121] All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains.

[0122] The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of" and "consisting of" may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible.

[0123] In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group. For example, if X is described as selected from the group consisting of bromine, chlorine, and iodine, claims for X being bromine and claims for X being bromine and chlorine are fully described.

[0124] Other embodiments are within the following claims.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **GEALL et al.** Nonviral delivery of self-amplifying RNA vaccines PNAS, 2012, vol. 109, 3614604-09 **[0081]**

- SMITH et al. PNAS, 2003, [0093]

PATENTKRAV

1. Automatiseret system til syntesen af et dsDNA-molekyle ifølge tilvejebragt sekvensinformation, omfattende:

5

en samlingsenhed til at samle dsDNA-molekylet ifølge tilvejebragt sekvensinformation,

10

hvor samlingsenheden omfatter eller er forbundet til kar indeholdende nukleotidderivater eller nukleosidphosphoramiditter og indeholdende komponenter til at transportere reagenser i systemet og til at udføre trin i en automatiseret fremgangsmåde til at samle dsDNA-molekylet gennem flere niveauer af biologisk samling, hvor niveauer af biologisk samling omfatter 1) syntesen af en flerhed af oligonukleotider ved at forbinde nukleotidderivater eller nukleosidphosphoramiditter, og 2) samlingen af et eller flere dsDNA-molekyler ved at forbinde flerheden af oligonukleotider; hvor der ikke sker nogen menneskelig indgriben, efter at fremgangsmåden er påbegyndt, før fremgangsmåden er afsluttet; og

15

hvor samlingsenheden yderligere omfatter softwareprogrammeringsinstruktioner, som dirigerer manipulationer af delene af samlingsenheden til at syntetisere flerheden af oligonukleotider fra nukleotidderivater eller nukleosidphosphoramiditter i overensstemmelse med sekvensinformationen og samle oligonukleotiderne til det automatiserede dsDNA-molekyle;

20

den automatiserede fremgangsmåde omfattende at udføre på flerheden af oligonukleotider et trin med PCR-amplifikation i en første reaktionszone i en reaktionsbeholder, et trin med fejlkorrektion på de PCR-amplificerede oligonukleotider i en anden reaktionszone i en reaktionsbeholder og et andet PCR trin i en tredje reaktionszone i en reaktionsbeholder;

25

hvor fejlkorrektionen udføres under anvendelse af en kombination af et medlem af CEL-nukleasefamilien af mismatch-specifikke nukleaser og exonuklease III;

30

hvor samlingsenheden anvender robotteknologi til at transportere prøver mellem reaktionszoner, således at den automatiserede fremgangsmåde kan udføres i én reaktionsbeholder;

hvor oligonukleotiderne er fragmenter af dsDNA-molekylet eller varianter deraf;
og yderligere hvor samlingsenheden udfører samling i en automatiseret fremgangsmåde, der omfatter syntesen af dsDNA-molekylet, der er større end 100
5 bp, ved at forbinde oligonukleotiderne.

2. Automatiseret system ifølge krav 1, hvor samlingsenheden yderligere omfatter komponenter til transport af reagenser i systemet og til udførelse af trin i en automatiseret fremgangsmåde til udførelse af et andet niveau af biologisk samling omfattende syntesen af et RNA-molekyle ifølge dsDNA'et molekyle.
10

3. Automatiseret system ifølge krav 1 eller 2, hvor oligonukleotiderne er tilvejebragt til en underenhed af samlingsenheden til amplifikation af oligonukleotiderne.

15 4. Automatiseret system ifølge et hvilket som helst af kravene 1-3, hvor oligonukleotiderne er samlet i det større dsDNA-molekyle ved anvendelse af isotermisk *in vitro*-rekombination i en enkelt, isoterm reaktion, eller

hvor oligonukleotiderne er fra 40-100 nukleotider lange, eller
20 hvor dsDNA-molekylet direkte er en DNA-vaccine eller komponent deraf.

5. Automatiseret system ifølge et hvilket som helst af kravene 1-4, hvor flerheden af oligonukleotider indeholder et sekvensoverlap til DNA-samling.

25 6. Automatiseret system ifølge krav 1 eller 2, hvor syntesen af flerheden af oligonukleotider ved at forbinde nukleotidderivater eller nukleosidphosphoramiditter udføres af en nukleinsyresyntese-underenhed, som syntetiserer oligonukleotiderne i den automatiserede fremgangsmåde.

30 7. Automatiseret system ifølge krav 1 eller 2, hvor dsDNA-molekylet er sammensat med en eller flere yderligere DNA-sekvenser syntetiseret af eller tilvejebragt som præ-lavede sekvenser til systemet, hvor den ene eller flere yderligere DNA-sekvenser er udvalgt blandt gruppen bestående af et plasmid, en vektor, en regulatorisk sekvens, en

promotorsekvens, et bindingselement for en trans-virkende faktor, en signalsekvens og en hvilken som helst kombination af to eller flere deraf.

- 5 8. Automatiseret system ifølge krav 7, hvor den yderligere DNA-sekvens er en vektor til anvendelse i en værtscelle, og det automatiserede system yderligere omfatter værtscellen, reagenser og komponenter til at transportere reagenser i systemet og til at udføre trin i en automatiseret fremgangsmåde fra softwareprogrammeringsinstruktioner til at transformere eller transficere værtscellen med dsDNA-molekylet og vektoren.
- 10 9. Automatiseret system ifølge krav 8, hvor den transformerende eller transficerede værtscelle danner et eller flere proteiner eller proteinunderenheder.
- 15 10. Automatiseret system ifølge krav 1 eller 2, hvor dsDNA- eller RNA-molekylet er en fag- eller viralt genom, og det automatiserede system yderligere omfatter værtscellen, reagenserne og komponenterne til at transportere reagenser i systemet og til at udføre trin i en automatiseret fremgangsmåde fra softwareprogrammeringsinstruktioner til at inficere eller transficere værtscellen med fagen eller det virale genom.
- 20 11. Automatiseret system ifølge krav 1 eller 2, hvor det automatiserede system samler flere dsDNA-molekyler.
- 25 12. Automatiseret system ifølge krav 2, hvor samlingsenheden yderligere omfatter reagenser og komponenter til transport af reagenser i systemet og til udførelse af trin i en automatiseret fremgangsmåde fra softwareprogrammeringsinstruktioner til udførelse af et andet niveau af biologisk samling omfattende syntesen af et protein- eller peptidmolekyle ved at oversætte RNA-molekylet.
- 30 13. Automatiseret system ifølge krav 7, hvor samlingsenheden udfører et andet niveau af biologisk samling omfattende samlingen af to eller flere proteiner til et protein med kvartær struktur.
14. Automatiseret system ifølge krav 12, hvor protein- eller peptidmolekylet er et produkt udvalgt blandt gruppen bestående af et proteindomæne, en bindingsunderenhed, et enzym, en enzymunderenhed, der har enzymatisk aktivitet, et protein eller pep-

tidevand, der har antigen aktivitet, et viralt protein eller en underenhed deraf, et viralt kappeprotein, og hvor det automatiserede system yderligere omfatter komponenter til udvinding af proteinet eller peptidmolekylet til anvendelse som en vaccine eller komponent deraf.

5

15. Automatiseret system ifølge krav 12, hvor det automatiserede system tilvejebringer multiple protein- eller peptidmolekyler.

10

16. Automatiseret system ifølge krav 1 eller 2, hvor det automatiserede system yderligere omfatter en modtageenhed til at modtage et signal, der koder for biologisk sekvensinformation transmitteret fra en sendeenhed.

DRAWINGS




Fig. 1




Fig. 2A




Fig. 2B

SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

