




(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/12/13
(87) Date publication PCT/PCT Publication Date: 2020/06/18
(85) Entrée phase nationale/National Entry: 2021/05/18
(86) N° demande PCT/PCT Application No.: US 2019/066376
(87) N° publication PCT/PCT Publication No.: 2020/124032
(30) Priorités/Priorities: 2018/12/14 (US62/779,885);
2019/04/19 (US62/836,520)

(51) Cl.Int./Int.Cl. *A61K 38/03* (2006.01),
A61K 49/00 (2006.01), *A61P 43/00* (2006.01),
C07K 7/08 (2006.01), *C07K 9/00* (2006.01)
(71) Demandeur/Applicant:
FRED HUTCHINSON CANCER RESEARCH CENTER,
US
(72) Inventeurs/Inventors:
CROOK, ZACHARY, US;
BRUSNIAK, MI-YOUN, US;
OLSON, JAMES, US;
MHYRE, ANDREW JAMES, US;
YIN, CHUNFENG, US;
HOPPING, GENE GREGORY, US;

(54) Titre : PEPTIDES CIBLANT UN RECEPTEUR DE LA TRANSFERRINE
(54) Title: TRANSFERRIN RECEPTOR TARGETING PEPTIDES



(57) Abrégé/Abstract:

Described herein are peptides and variants thereof as well as peptide constructs comprising peptides conjugated to, linked to, or fused to agents, wherein the peptides and peptide constructs are capable of binding TfR. Binding of a peptide or peptide construct as described herein to TfR can enable transcytosis across an endothelial layer, e.g., the blood brain barrier, or the crossing of a cell membrane. Pharmaceutical compositions and uses of peptides and peptide constructs, as well as methods of designing and manufacturing such peptides and peptide constructs, to treat a disease or condition are also described herein.

(72) Inventeurs(suite)/Inventors(continued): STRONG, ROLAND, US

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

18 June 2020 (18.06.2020)



(10) International Publication Number

WO 2020/124032 A1

(51) International Patent Classification:

A61K 38/03 (2006.01) *C07K 7/08* (2006.01)*A61P 43/00* (2006.01) *C07K 9/00* (2006.01)*A61K 49/00* (2006.01)

(21) International Application Number:

PCT/US2019/066376

(22) International Filing Date:

13 December 2019 (13.12.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/779,885 14 December 2018 (14.12.2018) US

62/836,520 19 April 2019 (19.04.2019) US

(71) Applicant: **FRED HUTCHINSON CANCER****RESEARCH CENTER** [US/US]; 1100 Fairview Avenue North, Seattle, Washington 98109 (US).

(72) **Inventors:** **CROOK, Zachary**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US). **BRUSNIAK, Mi-Young**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US). **OLSON, James**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US). **MHYRE, Andrew James**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US). **YIN, Chunfeng**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US). **HOPPING, Gene Gregory**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US). **STRONG, Roland**; 1100 Fairview Avenue North, Seattle, Washington 98109 (US).

(74) **Agent:** **LIPPERT, Lisa G.**; WILSON SONSINI GOODRICH & ROSATI, 650 Page Mill Road, Palo Alto, California 94304-1050 (US).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,

(54) Title: TRANSFERRIN RECEPTOR TARGETING PEPTIDES




FIG. 7

(57) **Abstract:** Described herein are peptides and variants thereof as well as peptide constructs comprising peptides conjugated to, linked to, or fused to agents, wherein the peptides and peptide constructs are capable of binding TfR. Binding of a peptide or peptide construct as described herein to TfR can enable transcytosis across an endothelial layer, e.g., the blood brain barrier, or the crossing of a cell membrane. Pharmaceutical compositions and uses of peptides and peptide constructs, as well as methods of designing and manufacturing such peptides and peptide constructs, to treat a disease or condition are also described herein.

[Continued on next page]



WO 2020/124032 A1

WO 2020/124032 A1

KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 239

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 239

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

TRANSFERRIN RECEPTOR TARGETING PEPTIDES

CROSS-REFERENCE

[0001] The present application claims priority to and benefit of U.S. Provisional Application Nos. 62/779,885 filed December 14, 2018, and 62/836,520 filed April 19, 2019, the entire contents of each of which are herein incorporated by reference.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on December 12, 2019, is named 44189-723_601_SL.txt and is 552,571 bytes in size.

BACKGROUND

[0003] Drug delivery to targets in the central nervous system (CNS) is hampered by the blood-brain barrier (BBB), a term for the particularly impassable vascular endothelial cells in CNS capillaries. The BBB exists to keep toxic metabolites and pathogens out of the brain, but also serves to render diseases of the CNS particularly difficult to treat using conventional medicines. For example, primary CNS tumors (e.g., gliomas) and neuroinflammatory and neurodegenerative diseases (e.g., Multiple Sclerosis and Alzheimer's Disease) respond particularly poorly to therapeutic modalities that would otherwise be more effective in similar diseases affecting peripheral tissues. Thus, approaches for an improved delivery of therapeutic and/or diagnostic molecules into the CNS and other organs and tissues are needed.

SUMMARY

[0004] In various aspects, the present disclosure provides a composition comprising an engineered transferrin receptor (TfR)-binding peptide, or a variant, homolog, fragment, or analog thereof.

[0005] In various aspects, the present disclosure provides a composition comprising a peptide construct, wherein the peptide construct comprises: a) a transferrin receptor (TfR)-binding peptide, or a variant, homolog, fragment, or analog thereof; and b) an active agent, wherein the peptide is conjugated to, linked to, or fused to the active agent.

[0006] In various aspects, the present disclosure provides a composition comprising an engineered peptide having at least 75%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a functional fragment thereof.

[0007] In some aspects, the peptide penetrates a cellular layer comprising a blood brain barrier (BBB). In some aspects, the peptide penetrates a membrane and wherein the membrane is the membrane of a cell. In some aspects, the cell expresses a TfR. In some aspects, the peptide penetrates the membrane of the cell via TfR-mediated endocytosis.

[0008] In some aspects, the peptide is delivered across the blood-brain barrier to the CNS by binding and then dissociating from TfR. In some aspects, the peptide is delivered across the blood-brain barrier by receptor-mediated transcytosis. In some aspects, the cell is a tumor cell. In some aspects, a cell or an organ in which the peptide localizes overexpresses a TfR. In some aspects, the peptide comprises SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344.

[0009] In some aspects, the peptide comprises a surface interface residue corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophilic surface-distal residues corresponding to any one of the positions 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophilic residues at a position corresponding to any one of 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises a hydrophilic residue corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32.

[0010] In some aspects, the peptide comprises hydrophobic residues at a position corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. In some aspects, the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32.

[0011] In some aspects, the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. In some aspects, the aliphatic residue is selected from the amino acids A, M, I, L, or V. In some aspects, the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32.

[0012] In some aspects, the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. In some aspects, the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. In some aspects, the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 1. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 2. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 4. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 30. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 32.

[0013] In some aspects, the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. In some aspects, the cell-penetrating moiety comprises polycations, polyorganic acids, endosomal releasing polymers, poly(2-propylacrylic acid), poly(2-ethylacrylic acid), Tat peptide, Arg patch, a knotted peptide, CysTAT, S19-TAT, R8 (SEQ ID NO: 73), pAntp, Pas-TAT, Pas-R8 (SEQ ID NO: 76), Pas-FHV, Pas-pAntP, F2R4 (SEQ ID NO: 79), B55, aurein, IMT-P8, BR2, OMOTAG1, OMOTAG2, pVEC, SynB3, DPV1047, C105Y, Transportan, MTS, hLF, PFVYLI (SEQ ID NO: 93), maurocalcine, imperatoxin, hadrucalin, hemicalcin, opicalcin-1, opicalcin-2, midkine (62-104), MCoTI-II, chlorotoxin, DRI-TAT, cF Φ R₄ (SEQ ID

NO: 96), R₆W₃ (SEQ ID NO: 421), myristate, yBBR, or a fragment or variant thereof, or any combination thereof. In some aspects, the cell-penetrating moiety is any one of SEQ ID NO: 71 – SEQ ID NO: 96, SEQ ID NO: 98 – SEQ ID NO: 101, SEQ ID NO: 116 – SEQ ID NO: 126, or SEQ ID NO: 403 – SEQ ID NO: 455.

[0014] In some aspects, the peptide sequence comprises at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36, at least 37, at least 38, at least 39, at least 40, at least 41, at least 42, at least 43, at least 44, at least 45, at least 46, at least 47, at least 48, at least 49, at least 50, at least 51, at least 52, at least 53, at least 54, at least 55, at least 56, at least 57, at least 58 residues, at least 59, at least 60, at least 61, at least 62, at least 63, at least 64, at least 65, at least 66, at least 67, at least 68, at least 69, at least 70, at least 71, at least 72, at least 73, at least 74, at least 75, at least 76, at least 77, at least 78, at least 79, at least 80, or at least 81 amino acid residues.

[0015] In some aspects, an active sequence is grafted onto the peptide or wherein the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct. In some aspects, the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400, or SEQ ID NO: 401.

[0016] In some aspects, the active agent is linked via a linker, wherein the linker is selected from **TABLE 7** or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. In some aspects, the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or

pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter. In some aspects, the Kv1.3 inhibitor is Vm24, ShK-170, ShK-186, or ShK-192. In some aspects, the Vm24 has a sequence of SEQ ID NO: 378 or SEQ ID NO: 391. In some aspects, the Shk-185 has a sequence of SEQ ID NO: 379, SEQ ID NO: 390, or SEQ ID NO: 402. In some aspects, the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. In some aspects, the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350 or SEQ ID NO: 365 – SEQ ID NO: 369.

[0017] In some aspects, the composition further comprises a half-life modifying agent coupled to the peptide. In some aspects, the half-life modifying agent comprises a polymer, a polyethylene glycol (PEG), a hydroxyethyl starch, polyvinyl alcohol, a water soluble polymer, a zwitterionic water soluble polymer, a water soluble poly(amino acid), a water soluble polymer of proline, alanine and serine, a water soluble polymer containing glycine, glutamic acid, and serine, an Fc region, a fatty acid, palmitic acid, or a molecule that binds to albumin. In some aspects, the half-life modifying agent comprises SA21. In some aspects, the SA21 has a sequence of SEQ ID NO: 376 or SEQ ID NO: 377.

[0018] In some aspects, the composition further comprises a detectable agent coupled to the peptide by a linker. In some aspects, the detectable agent is linked via a linker, wherein the linker is selected from **TABLE 7** or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. In some aspects, the detectable agent is a fluorophore, a near-infrared dye, a contrast agent, a nanoparticle, a metal-containing nanoparticle, a metal chelate, an X-ray contrast agent, a PET agent, a radionuclide, or a radionuclide chelator.

[0019] In various aspects, the present disclosure provides a pharmaceutical composition comprising any of the compositions described herein, or a salt thereof, and a pharmaceutically acceptable carrier.

[0020] In various aspects, the present disclosure provides a method for transporting a composition across a cell layer, the method comprising: a) contacting the cell layer with the composition comprising any of the peptides described herein or any of the pharmaceutical compositions described herein; and b) transporting the composition across the cell layer.

[0021] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising: a) administering to the subject a composition

comprising any of the peptides described herein or any of the pharmaceutical compositions described herein, and b) transporting the composition across a cell layer.

[0022] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising administering to the subject an amount of a composition comprising any of the peptides described herein or any of the pharmaceutical compositions described herein based on the expression of TfR in a tissue of the subject.

[0023] In some aspects, the peptide construct is capable of binding and activating a neurotensin receptor. In some aspects, the method further comprises reducing a level of pain upon administration of the peptide to the subject. In some aspects, the administered is inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof. In some aspects, the composition is administered intravenously as a bolus, infusion, or prolonged infusion.

[0024] In some aspects, the method further comprises administering the peptide to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, pain, epilepsy, a mood disorder, or an endocrine disorder. In some aspects, the subject has a cancer. In some aspects, the subject has a disease of the CNS. In some aspects, the subject has neuropathic pain, migraine, anxiety, or epilepsy.

[0025] In various aspects, the present disclosure provides a method for transporting a composition according to any one of claims 1-53 or the pharmaceutical composition of claim 54 across a cell layer, the method comprising contacting the cell layer with the composition, and transporting the composition across the cell layer.

[0026] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising administering to the subject the composition of any one of claims 1-53 or the pharmaceutical composition of claim 54.

[0027] In some aspects, the transporting comprises binding of the composition to TfR. In some aspects, the transporting further comprises transcytosis. In some aspects, the transcytosis is vesicular transcytosis. In some aspects, the transcytosis is TfR-mediated. In some aspects, the transporting further comprises dissociation of the composition from the TfR. In some aspects, the transporting further comprises releasing an active agent from the composition.

[0028] In various aspects, the present disclosure provides a method for designing a TfR-binding peptide, the method comprising: a) using a computer program to identify a peptide from a library of peptides with optimized stability and folding, wherein each peptide of the library of peptides

comprises at least three intramolecular disulfide bonds; b) using a computer program to superimpose amino acid residues of a binding patch from a peptide:TfR complex to determine a stable peptide that binds to TfR after grafting the binding patch onto the peptide; and c) grafting the binding patch onto the peptide to produce the stable peptide.

[0029] In some aspects, the method further comprises performing site saturation mutagenesis to obtain a mutated peptide with a higher binding affinity for TfR than the stable peptide. In some aspects, the peptide from a library of peptides is between 20 and 75 amino acid residues in length.

[0030] In various aspects, the present disclosure provides a composition, the composition comprising a non-naturally occurring peptide capable of binding to a transferrin receptor (TfR), or a variant, homolog, fragment, or analog thereof.

[0031] In various aspects, the present disclosure provides a composition, the composition comprising a peptide construct, wherein the peptide construct comprises: a peptide capable of binding to a transferrin receptor (TfR), or a variant, homolog, fragment, or analog thereof; and an active agent, wherein the peptide is conjugated to, linked to, or fused to the active agent.

[0032] In some aspects, the transferrin receptor (TfR) is a human transferrin receptor, or murine transferrin receptor, or a human transferrin receptor and murine transferrin receptor.

[0033] In various aspects, the present disclosure provides a composition, the composition comprising a peptide having at least 75%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a functional fragment thereof.

[0034] In some aspects, the peptide penetrates a cellular layer comprising endothelial cells. In some aspects, the peptide penetrates a cellular layer comprising a blood brain barrier (BBB). In some aspects, the peptide penetrates a cellular layer comprising endothelial cells via TfR-mediated transcytosis. In some aspects, the peptide penetrates a membrane and the membrane is the membrane of a cell. In some aspects, the cell expresses a TfR. In some aspects, the peptide penetrates the membrane of the cell via TfR-mediated endocytosis. In some aspects, the peptide is capable of localizing in or with any cell that expresses a TfR. In some aspects, the peptide remains bound to a cell surface membrane expressing TfR. In some aspects, the peptide is internalized into a cell by binding TfR. In some aspects, the peptide is delivered across the blood-

brain barrier to the CNS by binding and then dissociating from TfR. In some aspects, the peptide is delivered across the blood-brain barrier by receptor-mediated transcytosis.

[0035] In some aspects, the cell is a CNS cell, an erythrocyte, an erythrocyte precursor cell, an immune cell, a stem cell, a muscle cell, a brain cell, a thyroid cell, a parathyroid cell, an adrenal gland cell, a bone marrow cell, an appendix cell, a lymph node cell, a tonsil cell, a spleen cell, a muscle cell, a liver cell, a gallbladder cell, a pancreas cell, a cell of the gastrointestinal tract, a glandular cell, a kidney cell, a urinary bladder cell, an endothelial cell, an epithelial cell, a choroid plexus epithelial cell, a glial cell, an astrocyte, or a cell associated with a nervous system.

[0036] In some aspects, the cell is a tumor cell. In some aspects, the tumor cell is an ovarian cancer cell, a colon cancer cell, a lung cancer cell, a cancer cell located in the bone or bone marrow, a glioblastoma cell, an astrocytoma cell, a glioma cell, a medulloblastoma cell, an ependymoma cell, a choroid plexus carcinoma cell, a midline glioma cell, a diffuse intrinsic pontine glioma (DIPG) cell, a breast cancer cell, a liver cancer cell, a colon cancer cell, a brain cancer cell, a spleen cancer cell, a cancer cell of the salivary gland, a kidney cancer cell, a muscle cancer cell, a bone marrow cell cancer cell, a skin cancer cell, a genitourinary cancer cell, an osteosarcoma cell, a muscle-derived sarcoma cell, a melanoma cell, a head and neck cancer cell, a neuroblastoma cell, a prostate cancer cell, a bladder cancer cell, an acute lymphocytic leukemia cell, an acute myeloid leukemia cell, a chronic lymphocytic leukemia cell, a chronic myeloid leukemia cell, a Hodgkin lymphoma cell, a Non-Hodgkin lymphoma cell, or a CMYC-overexpressing cancer cell. In some aspects, the peptide localizes to an organ or tissue.

[0037] In some aspects, the organ or tissue is brain, thyroid, parathyroid, adrenal glands, bone marrow, appendix, lymph node, tonsil, spleen, muscle, liver, gallbladder, pancreas, gastrointestinal tract, kidney, urinary bladder, fallopian tube, breast, vagina, cervix, endometrium, ovary, placenta, and skin. In some aspects, the cell or organ in which the peptide localizes overexpresses a TfR. In some aspects, the peptide localizes to the parenchyma of the organ.

[0038] In some aspects, the peptide binds at least one amino acid residues selected from the group of residues 506-510, 523-531, or 611-662 corresponding to SEQ ID NO: 363 (MMDQARSAFSNLFGGEPLSYTRFSLARQVDGDNHVMKLA VDEEENADNNTKANVT KPKRCSGSICYGTIAVIVFFLIGFMIGYLG YCKGVEPKTECERLAGTESPVREEPGEDFPA ARRLYWDDLKRKLSEKLDSTDFGTIKLLNENS YVPREAGSQKDENLALYVENQFREFK LSKVWRDQHFVKIQVKDSAQNSVIIVDKNGRLVYLVENPGGYVAYSKAATVTGKLVHA

NFGTKKDFEDLYTPVNGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELS
 FFGHAHLGTGDPYTPGFPSFNHTQFPPSRSSGLPNIPVQTISRAAAELFGNMEGDCPSD
 WKTDSTCRMVTSSESKNVKLTVSNVLKEIKILNIFGVIKGFVEPDHYVVVGAQRDAWGPG
 AAKSGVGTALLLKLQMFSDMVLKDGFPQRSIIFASWSAGDFGSGVATEWLEGYLSSL
 HLKAFTYINLDKAVLGTSNFKVSASPLLYTLIEKTMQNVKHPVTGQFLYQDSNWASKVE
 KLTLDNAAFPFLAYSGIPAVSFCFCEDTDYPYLGTTMDTYKELIERIPELNKVARAAAEV
 AGQFVIKLTHDVELNLDYERYNSQLLSFVRDLNQYRADIKEMGLSLQWLYSARGDFFR
 ATSRLTTDFGNAEKTDRFVMKKLNDRVMRVEYHFLSPYVSPKESPFRHVFWSGSGSHTLP
 ALLENLKLKQNNGAFNETLFRNQLALATWTIQGAANALSGDVWDIDNEF) of human
 TfR1 (UniProtKB - P02786). In some aspects, the peptide comprises SEQ ID NO: 206 – SEQ ID
 NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344. In some aspects, peptide comprises a surface
 interface residue corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47,
 S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects,
 the peptide comprises hydrophilic surface-distal residues corresponding to any one of the
 positions 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any
 combination thereof, with reference to SEQ ID NO: 32.

[0039] In some aspects, the hydrophilic surface-distal residues are selected from the amino acids
 D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic
 surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29,
 T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference
 to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophilic residues at a position
 corresponding to any one of 15, 35, 39, or 49, or any combination thereof, with reference to SEQ
 ID NO: 32.

[0040] In some aspects, the hydrophilic residues are selected from the amino acids D, E, H, K, R,
 N, Q, S, or T. In some aspects, the peptide comprises a hydrophilic residue corresponding to
 D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. In some
 aspects, the peptide comprises hydrophobic residues at a position corresponding to 11, 25, or 27,
 or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophobic
 residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof.
 In some aspects, the peptide comprises any one of the hydrophobic residues corresponding to
 M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects,
 the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ

ID NO: 32. In some aspects, the aliphatic residue is selected from the amino acids A, M, I, L, or V.

[0041] In some aspects, the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. In some aspects, the peptide comprises three disulfide bonds. In some aspects, the peptide comprises at least two, at least four, or at least six cysteine residues. In some aspects, the peptide comprises six cysteine residues.

[0042] In some aspects, the peptide comprises cysteine residues at positions corresponding to C6, C10, C20, C34, C44, and C48, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, at least one cysteine residue of the peptide participates in a disulfide bond. In some aspects, the six cysteine residues participate in disulfide bonds. In some aspects, a disulfide bond increases the stability of the peptide.

[0043] In some aspects, the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. In some aspects, the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof.

[0044] In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 1. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 2. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 4. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 30. In some aspects, the peptide comprises the sequence set forth in SEQ ID NO: 32.

[0045] In some aspects, the peptide comprising three disulfide bonds is derived from a cystine-dense peptide (CDP). In some aspects, the peptide comprising three disulfide bonds is a CDP. In some aspects, the peptide competes with transferrin, or a homolog, variant, or fragment thereof, for binding to TfR. In some aspects, the peptide sequesters TfR or prevents another protein from binding to TfR. In some aspects, the peptide has equal or greater affinity or specificity for TfR as compared to that of transferrin, or an endogenous derivative thereof. In some aspects, the peptide

has equal or lower dissociation constant for TfR as compared to that of transferrin, or an endogenous derivative thereof. In some aspects, the peptide has a K_D of less than 40 nM, less than 30 nM, less than 20 nM, less than 10 nM, less than 1 nM, or less than 0.1 nM. In some aspects, the peptide penetrates or localizes into a target cell.

[0046] In some aspects, the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. In some aspects, the cell-penetrating moiety comprises polycations, polyorganic acids, endosomal releasing polymers, poly(2-propylacrylic acid), poly(2-ethylacrylic acid), Tat peptide, Arg patch, a knotted peptide, CysTAT, S19-TAT, R8 (SEQ ID NO: 73), pAntp, Pas-TAT, Pas-R8 (SEQ ID NO: 76), Pas-FHV, Pas-pAntP, F2R4 (SEQ ID NO: 79), B55, aurin, IMT-P8, BR2, OMOTAG1, OMOTAG2, pVEC, SynB3, DPV1047, C105Y, Transportan, MTS, hLF, PFVYLI (SEQ ID NO: 93), maurocalcine, imperatoxin, hadrucalin, hemicalcin, opicalcin-1, opicalcin-2, midkine (62-104), MCoTI-II, chlorotoxin, DRI-TAT, cF Φ R₄ (SEQ ID NO: 96), myristate, yBBR, or a fragment or variant thereof, or any combination thereof. In some aspects, the cell-penetrating moiety is maurocalcine, imperatoxin, hadrucalcine, hemicalcin, opicalcin-1, opicalcin-2, midkine (62-104), MCoTI-II, or chlorotoxin.

[0047] In some aspects, the peptide targets, homes, accumulates in, or selectively penetrates a tissue or cell. In some aspects, the peptide localizes into a target cell's nucleus. In some aspects, the peptide localizes into the target cell's nucleus by formulation with, fusion to, or conjugation to a nuclear localization signal peptide or moiety. In some aspects, the peptide is further conjugated to, linked to, or fused to a moiety or a second peptide that targets, homes, accumulates in, or selectively penetrates a cancerous tissue or cancer cell. In some aspects, the moiety or a second peptide is a knotted peptide or variant or fragment thereof. In some aspects, the target cell is a cancerous cell or a tumor cell. In some aspects, the target cell is a CNS cell, an erythrocyte, an erythrocyte precursor cell, an immune cell, a stem cell, a muscle cell, a brain cell, a thyroid cell, a parathyroid cell, an adrenal gland cell, a bone marrow cell, an appendix cell, a lymph node cell, a tonsil cell, a spleen cell, a muscle cell, a liver cell, a gallbladder cell, a pancreas cell, a cell of the gastrointestinal tract, a glandular cell, a kidney cell, a urinary bladder cell, an endothelial cell, an epithelial cell, a choroid plexus epithelial cell, a glial cell, an astrocyte, or a cell associated with a nervous system.

[0048] In some aspects, the peptide comprises a disulfide through disulfide knot. In some aspects, the peptide comprises a plurality of disulfide bridges formed between cysteine residues. In some aspects, the peptide comprises three or more disulfide bridges formed between cysteine

residues, wherein one of the disulfide bridges passes through a loop formed by two other disulfide bridges. In some aspects, the peptide comprises no disulfide bond or a disulfide knot. In some aspects, at least one amino acid residue of the peptide is in an L configuration or, wherein at least one amino acid residue is in a D configuration.

[0049] In some aspects, the peptide comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 positively charged residues. In some aspects, the peptide comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 consecutive positively charged residues. In some aspects, the positively charged residues are Arg, Lys, or any combination thereof. In some aspects, the peptide comprises an Arg patch. In some aspects, the peptide comprises two or more consecutive Arg residues at the N-terminus.

[0050] In some aspects, the peptide comprises a Tat peptide, or a fragment thereof. In some aspects, the Tat peptide comprises a sequence of YGRKKRRQRRR (SEQ ID NO: 99), GRKKRRQRRR (SEQ ID NO: 100), or a fragment or variant thereof. In some aspects, the Tat peptide is appended to the N-terminus or C-terminus of the peptide using a linker having a sequence of (GS)_x (SEQ ID NO: 105) or G_xS_y (SEQ ID NO: 104), wherein x and y is independently 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. In some aspects, the Tat peptide is appended to any residue of the peptide.

[0051] In some aspects, the peptide comprises at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, or at least 16 cysteine residues. In some aspects, the peptide comprises one or more regions of negatively charged residues. In some aspects, the peptide comprises one or more regions of positively charged residues.

[0052] In some aspects, the peptide sequence comprises at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36, at least 37, at least 38, at least 39, at least 40, at least 41, at least 42, at least 43, at least 44, at least 45, at least 46, at least 47, at least 48, at least 49, at least 50, at least 51, at least 52, at least 53, at least 54, at least 55, at least 56, at least 57, at least 58 residues, at least 59, at least 60, at least 61, at least 62, at least 63, at least 64, at least 65, at least 66, at least 67, at least 68, at least 69, at least 70, at least 71, at least

72, at least 73, at least 74, at least 75, at least 76, at least 77, at least 78, at least 79, at least 80, or at least 81 amino acid residues.

[0053] In some aspects, the peptide comprises or is derived from a human protein or peptide. In some aspects, the peptide is arranged in a multimeric structure with at least one other peptide. In some aspects, the multimeric structure comprises a dimer, trimer, tetramer, pentamer, hexamer, or heptamer. In some aspects, the peptide comprises an isoelectric point within a range from about 3.0 to about 10.0. In some aspects, the peptide comprises an isoelectric point within a range from about 4.5 to about 8.9. In some aspects, the peptide comprises a non-uniform charge distribution. In some aspects, the peptide is stable at physiological or intracellular pH values.

[0054] In some aspects, the peptide is stable at a pH of about 7 or 7.5. In some aspects, the peptide is stable at a pH of about 6.5 to 7.5. In some aspects, the peptide is stable at pH values less than or equal to about 5.0, less than or equal to about 3.0, or within a range from about 3.0 to about 5.0. In some aspects, the peptide is stable at pH values within a range from about 5.0 to about 7.0. In some aspects, the peptide comprises a hydrophobic core.

[0055] In some aspects, the peptide is resistant to protease degradation. In some aspects, the protease is any one of pepsin, trypsin, chymotrypsin, serum proteases, intracellular proteases, or any combination thereof. In some aspects, the peptide is resistant to reduction or is stable in a reducing environment. In some aspects, the reduction or reducing environment comprises DTT or GSH. In some aspects, the peptide is stable at an elevated temperature. In some aspects, the peptide exhibits an anti-cancer effect. In some aspects, the peptide comprises one or more chemical modifications. In some aspects, the chemical modification modifies one or more pharmacokinetic properties of the peptide. In some aspects, the chemical modification extends the plasma half-life and/or the biological half-life of the peptide.

[0056] In some aspects, the chemical modification is blocking the N-terminus of the peptide. In some aspects, the chemical modification is methylation, acetylation, or acylation. In some aspects, the chemical modification is: methylation of one or more lysine residues or analogue thereof; methylation of the N-terminus; or methylation of one or more lysine residue or analogue thereof and methylation of the N-terminus. In some aspects, the peptide is linked to an acyl adduct. In some aspects, an active sequence is grafted onto the peptide or the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct.

[0057] In some aspects, the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 359 – SEQ ID NO: 361. In some aspects, the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361. In some aspects, the active agent is conjugated to, linked to, or fused to the peptide at an N-terminus or a C-terminus of the peptide. In some aspects, the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid.

[0058] In some aspects, the nucleic acid is DNA. In some aspects, the nucleic acid is RNA. In some aspects, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 active agents are linked to the peptide by a linker. In some aspects, the peptide is linked to the active agent via a cleavable linker or a pH sensitive linker. In some aspects, the peptide is linked to the active agent at an N-terminus, at the epsilon amine of an internal lysine residue, at the carboxylic acid of an aspartic acid or glutamic acid residue, or a C-terminus of the peptide by a linker. In some aspects, the composition further comprises a non-natural amino acid, wherein the non-natural amino acid is an insertion, appendage, or substitution for another amino acid. In some aspects, the peptide is linked to the active agent at the non-natural amino acid by a linker. In some aspects, the linker comprises an amide bond, an ester bond, an ether bond, a triazole, a macrocycle, an oxime bond, a carbamate bond, a carbonate bond, a hydrazone bond, an azo bond, an oxime bond, a disulfide bond, a thioester bond, a thioether bond, a carbon-carbon single, double, or triple bond, a disulfide bond, a two carbon bridge between two cysteines, a three carbon bridge between two cysteines, or a carbon-nitrogen bond.

[0059] In some aspects, the cleavable linker comprises a cleavage site for a protease. In some aspects, the protease is a matrix metalloproteinases, thrombin, cathepsins, or beta-glucuronidase. In some aspects, the linker is a hydrolytically labile linker. In some aspects, the peptide is linked to the active agent via a noncleavable linker. In some aspects, the active agent is linked via a linker, wherein the linker is selected from **TABLE 4**. In some aspects, the active agent is an anti-cancer agent. In some aspects, the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFNgamma agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a

calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter.

[0060] In some aspects, the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. In some aspects, the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues.

[0061] In some aspects, the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. In some aspects, the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350. In some aspects, the peptide construct is capable of binding and activating a neurotensin receptor. In some aspects, the peptide is capable of reducing a level of pain in a subject. In some aspects, the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, or an endocrine disorder. In some aspects, the peptide interacts with dopamine signaling neurons. In some aspects, the peptide increases dopamine release in neurons. In some aspects, the composition further comprises a half-life modifying agent coupled to the peptide. In some aspects, the half-life modifying agent comprises a polymer, a polyethylene glycol (PEG), a hydroxyethyl starch, polyvinyl alcohol, a water soluble polymer, a zwitterionic water soluble polymer, a water soluble poly(amino acid), a water soluble polymer of proline, alanine and serine, a water soluble polymer containing glycine, glutamic acid, and serine, an Fc region, a fatty acid, palmitic acid, or a molecule that binds to albumin.

[0062] In some aspects, the composition further comprises a detectable agent coupled to the peptide by a linker. In some aspects, the detectable agent is conjugated to, linked to, or fused with the peptide at an N-terminus or a C-terminus of the peptide. In some aspects, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 detectable agents are linked to the peptide. In some aspects, the peptide is linked to the detectable agent via a cleavable linker. In some aspects, the peptide is linked to the detectable agent at an N-terminus, at the epsilon amine of an internal lysine residue, or a C-terminus of the peptide by a linker. In some aspects, the composition further comprises a non-natural amino acid,

wherein the non-natural amino acid is an insertion, appendage, or substitution for another amino acid. In some aspects, the peptide is linked to the detectable agent at the non-natural amino acid by a linker. In some aspects, the linker comprises an amide bond, an ester bond, an ether bond, a triazole, a macrocycle, an oxime bond, a carbamate bond, a carbonate bond, a hydrazone bond, an azo bond, an oxime bond, a disulfide bond, a thioester bond, a thioether bond, a carbon-carbon single, double, or triple bond, a disulfide bond, a two carbon bridge between two cysteines, a three carbon bridge between two cysteines, or a carbon-nitrogen bond. In some aspects, the cleavable linker comprises a cleavage site for matrix metalloproteinases, thrombin, cathepsins, or beta-glucuronidase. In some aspects, the peptide is linked to the detectable agent via a stable linker. In some aspects, the detectable agent is linked via a linker, wherein the linker is selected from **TABLE 4**.

[0063] In some aspects, the detectable agent is a fluorophore, a near-infrared dye, a contrast agent, a nanoparticle, a metal-containing nanoparticle, a metal chelate, an X-ray contrast agent, a PET agent, a radionuclide, or a radionuclide chelator. In some aspects, the detectable agent is a fluorescent dye.

[0064] In various aspects, the present disclosure provides a pharmaceutical composition comprising any of the compositions disclosed herein, or a salt thereof, and a pharmaceutically acceptable carrier.

[0065] In some aspects, the pharmaceutical composition is formulated for administration to a subject. In some aspects, the pharmaceutical composition is formulated for oral administration, intravenous administration, subcutaneous administration, intramuscular administration, or a combination thereof.

[0066] In various aspects, the present disclosure provides a method for transporting a composition across a cell layer, the method comprising: contacting the cell layer with the composition; and transporting the composition across the cell layer.

[0067] In some aspects, the composition comprises a peptide. In some aspects, the peptide comprises an amino acid sequence comprising any one of SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344, or a functional fragment thereof. In some aspects, the peptide is capable of binding a transferrin receptor (TfR). In some aspects, the peptide comprises a surface interface residues corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises a hydrophilic surface-distal residue at a position corresponding to

3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophilic residues at any one of the positions corresponding to 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32.

[0068] In some aspects, the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic residues corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophobic residues at any of the positions corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. In some aspects, the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. In some aspects, the aliphatic residue is selected from the amino acids A, M, I, L, or V. In some aspects, the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32.

[0069] In some aspects, the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. In some aspects, the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. In some aspects, the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof. In some aspects, the peptide penetrates or localizes into or on a target cell. In some aspects, an active sequence is grafted onto the peptide or the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct.

[0070] In some aspects, the peptide is conjugated to, linked to, or fused to the active agent. In some aspects, the active agent is linked via a linker, wherein the linker is selected from **TABLE 4**. In some aspects, the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. In some aspects, the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 359 – SEQ ID NO: 361. In some aspects, the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361.

[0071] In some aspects, the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter. In some aspects, the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof.

[0072] In some aspects, the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. In some aspects, the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. In some aspects, the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350.

[0073] In some aspects, the peptide construct is capable of binding and activating a neurotensin receptor. In some aspects, the peptide is capable of reducing a level of pain upon administration to a subject. In some aspects, the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, or an endocrine disorder. In some aspects, the peptide interacts with dopamine signaling neurons. In some aspects, the peptide increases dopamine release in neurons.

In some aspects, the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. In some aspects, the cell layer is an endothelial cell layer or an epithelial cell layer. In some aspects, the endothelial cell layer is the blood brain barrier. In some aspects, the endothelial cell layer is an intestinal endothelial cell layer. In some aspects, the epithelial cell layer is an intestinal epithelial cell layer. In some aspects, the transporting comprises binding of the composition to TfR. In some aspects, the transporting further comprises transcytosis. In some aspects, the transcytosis is vesicular transcytosis. In some aspects, the transcytosis is TfR-mediated. In some aspects, the transporting further comprises dissociation of the composition from the TfR.

[0074] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising: administering to the subject a composition, and transporting the composition across a cell layer.

[0075] In some aspects, the composition comprises a peptide. In some aspects, the peptide comprises an amino acid sequence of any one of SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344. In some aspects, the peptide is capable of binding a transferrin receptor (TfR). In some aspects, the peptide comprises a surface interface residues corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises a hydrophilic surface-distal residues at any one of the positions corresponding to 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32.

[0076] In some aspects, the peptide comprises hydrophilic residues at any one of the positions corresponding to 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic residues corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophobic residues at any of the positions corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. In

some aspects, the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. In some aspects, the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32.

[0077] In some aspects, the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. In some aspects, the aliphatic residue is selected from the amino acids A, M, I, L, or V. In some aspects, the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds.

[0078] In some aspects, the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. In some aspects, the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof.

[0079] In some aspects, the peptide penetrates or localizes into a target cell. In some aspects, an active sequence is grafted onto the peptide or the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct. In some aspects, the peptide is conjugated to, linked to, or fused to the active agent. In some aspects, the active agent is linked via a linker, wherein the linker is selected from **TABLE 4**.

[0080] In some aspects, the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. In some aspects, the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 359 – SEQ ID NO: 361. In some aspects, the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361.

[0081] In some aspects, the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFNgamma agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter.

[0082] In some aspects, the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. In some aspects, the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. In some aspects, the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. In some aspects, the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350.

[0083] In some aspects, the peptide construct is capable of binding and activating a neurotensin receptor. In some aspects, the peptide is capable of reducing a level of pain upon administration to the subject. In some aspects, the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, or an endocrine disorder. In some aspects, the peptide interacts with dopamine signaling neurons. In some aspects, the peptide increases dopamine release in neurons. In some aspects, the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. In some aspects, the cell layer is an endothelial cell layer or an epithelial cell layer. In some aspects, the endothelial cell layer is the blood brain barrier.

[0084] In some aspects, the endothelial cell layer is an intestinal endothelial cell layer. In some aspects, the epithelial cell layer is an intestinal epithelial cell layer. In some aspects, the transporting comprises binding of the composition to TfR. In some aspects, the transporting further comprises transcytosis. In some aspects, the transcytosis is vesicular transcytosis. In some aspects, the transcytosis is TfR-mediated. In some aspects, the transporting further comprises

dissociation of the composition from the TfR. In some aspects, the transporting further comprises releasing the active agent from the composition.

[0085] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising administering to the subject an amount of a composition based on the expression of TfR in the subject.

[0086] In some aspects, the composition comprises a peptide. In some aspects, the peptide comprises an amino acid sequence of any one of SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344. In some aspects, the peptide is capable of binding a transferrin receptor (TfR). In some aspects, the peptide comprises a surface interface residues corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises a hydrophilic surface-distal residues at any one of the positions corresponding to 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophilic residues at any one of the positions corresponding to 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. In some aspects, the peptide comprises any one of the hydrophilic residues corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises hydrophobic residues at any of the positions corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32.

[0087] In some aspects, the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. In some aspects, the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. In some aspects, the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. In some aspects, the aliphatic residue is selected from the amino acids A, M, I, L, or V. In some aspects, the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. In some aspects,

the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds.

[0088] In some aspects, the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. In some aspects, the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof.

[0089] In some aspects, the peptide penetrates or localizes into a target cell. In some aspects, an active sequence is grafted onto the peptide or the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct. In some aspects, the peptide is conjugated to, linked to, or fused to the active agent. In some aspects, the active agent is linked via a linker, wherein the linker is selected from **TABLE 4**.

[0090] In some aspects, the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. In some aspects, the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 359 – SEQ ID NO: 361. In some aspects, the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361.

[0091] In some aspects, the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter.

[0092] In some aspects, the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. In some aspects, the functional fragment is at least at least 5 amino

acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. In some aspects, the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. In some aspects, the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350.

[0093] In some aspects, the peptide construct is capable of binding and activating a neurotensin receptor. In some aspects, the peptide is capable of reducing a level of pain upon administration to the subject. In some aspects, the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, or an endocrine disorder. In some aspects, the peptide interacts with dopamine signaling neurons. In some aspects, the peptide increases dopamine release in neurons. In some aspects, the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. In some aspects, the composition is administered by inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof.

[0094] In some aspects, the composition is administered intravenously as a bolus, infusion, or prolonged infusion. In some aspects, the composition or the pharmaceutical composition is administered multiple times a day, twice a day, once a day, twice a week, three times a week, once a week, once every two weeks, or once a month or once every three months. In some aspects, the composition is administered subcutaneously once a day, once a week, or once a month. In some aspects, the condition is a cancer.

[0095] In some aspects, the cancer is ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin

lymphoma, or a CMYC-overexpressing cancer. In some aspects, the condition is a brain cancer. In some aspects, the brain cancer is glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, or diffuse intrinsic pontine glioma.

[0096] In some aspects, the condition is a disease of the CNS. In some aspects, the disease of the CNS is a neuroinflammatory or neurodegenerative disease. In some aspects, the neurodegenerative disease is Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufoor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. In some aspects, the condition is neuropathic pain, migraine, anxiety, or epilepsy. In some aspects, the condition is chronic pain, acute pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, or breakthrough pain.

[0097] In some aspects, the condition is chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. In some aspects, the condition is muscular dystrophy, or spinal bulbar muscular dystrophy (SBMA). In some aspects, the condition is an inflammatory bowel disease. In some aspects, the inflammatory bowel disease is a Crohn's disease.

[0098] In various aspects, the present disclosure provides a method for transporting any of the compositions disclosed herein across a cell layer, the method comprising contacting the cell layer with the composition, and transporting the composition across the cell layer.

[0099] In some aspects, the cell layer is an endothelial cell layer or an epithelial cell layer. In some aspects, the endothelial cell layer is the blood brain barrier. In some aspects, the endothelial cell layer is an intestinal endothelial cell layer. In some aspects, the epithelial cell layer is an intestinal epithelial cell layer. In some aspects, the transporting comprises binding of the

composition to TfR. In some aspects, the transporting further comprises transcytosis. In some aspects, the transcytosis is vesicular transcytosis. In some aspects, the transcytosis is TfR-mediated. In some aspects, the transporting further comprises dissociation of the composition from the TfR. In some aspects, the transporting further comprises releasing an active agent from the composition.

[0100] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising administering to the subject any of the compositions disclosed herein or any of the pharmaceutical compositions disclosed herein.

[0101] In some aspects, the composition is administered by inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof. In some aspects, the composition is administered intravenously as a bolus, infusion, or prolonged infusion. In some aspects, the composition or the pharmaceutical composition is administered multiple times a day, twice a day, once a day, twice a week, three times a week, once a week, once every two weeks, or once a month or once every three months. In some aspects, the composition is administered subcutaneously once a day, once a week, or once a month. In some aspects, the condition is a cancer. In some aspects, the cancer is ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer. In some aspects, the condition is a brain cancer.

[0102] In some aspects, the brain cancer is glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, or diffuse intrinsic pontine glioma. In some aspects, the condition is a disease of the CNS. In some aspects, the condition of the CNS is a neuroinflammatory or neurodegenerative disease. In some aspects, the neurodegenerative disease is Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease,

Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufoer–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. In some aspects, the condition is neuropathic pain, migraine, anxiety, or epilepsy.

[0103] In some aspects, the condition is chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. In some aspects, the condition is muscular dystrophy, or spinal bulbar muscular dystrophy (SBMA).

[0104] In some aspects, the condition is an inflammatory bowel disease. In some aspects, the inflammatory bowel disease is a Crohn's disease. In some aspects, the method further comprises transporting the composition across a cell layer. In some aspects, the cell layer is an endothelial cell layer or an epithelial cell layer. In some aspects, the endothelial cell layer is the blood brain barrier. In some aspects, the endothelial cell layer is an intestinal endothelial cell layer.

[0105] In some aspects, the epithelial cell layer is an intestinal epithelial cell layer. In some aspects, the transporting comprises binding of the composition to TfR. In some aspects, the transporting further comprises transcytosis. In some aspects, the transcytosis is vesicular transcytosis. In some aspects, the transcytosis is TfR-mediated. In some aspects, the transporting further comprises dissociation of the composition from the TfR. In some aspects, the transporting further comprises releasing an active agent from the composition.

[0106] In various aspects, the present disclosure provides a method of treating a condition in a subject in need thereof, the method comprising administering to the subject an amount of any of the compositions disclosed herein or any of the pharmaceutical compositions disclosed herein that is dependent on the expression of TfR in the subject.

[0107] In some aspects, the composition is administered by inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof. In some aspects, the composition is

administered intravenously as a bolus, infusion, or prolonged infusion. In some aspects, the composition or the pharmaceutical composition is administered multiple times a day, twice a day, once a day, twice a week, three times a week, once a week, once every two weeks, or once a month or once every three months. In some aspects, the composition is administered subcutaneously once a day, once a week, or once a month. In some aspects, the condition is a cancer.

[0108] In some aspects, the cancer is ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer. In some aspects, the condition is a brain cancer. In some aspects, the brain cancer is glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, or diffuse intrinsic pontine glioma.

[0109] In some aspects, the condition is a disease of the CNS. In some aspects, the disease of the CNS is a neuroinflammatory or neurodegenerative disease. In some aspects, the neurodegenerative disease is Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia.

[0110] In some aspects, the condition is neuropathic pain, migraine, anxiety, or epilepsy. In some aspects, the condition is chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia,

phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. In some aspects, the condition is muscular dystrophy, or spinal bulbar muscular dystrophy (SBMA).

[0111] In some aspects, the condition is an inflammatory bowel disease. In some aspects, the inflammatory bowel disease is a Crohn's disease. In some aspects, the method further comprises transporting the composition across a cell layer. In some aspects, the cell layer is an endothelial cell layer or an epithelial cell layer. In some aspects, the endothelial cell layer is the blood brain barrier. In some aspects, the endothelial cell layer is an intestinal endothelial cell layer. In some aspects, the epithelial cell layer is an intestinal epithelial cell layer.

[0112] In some aspects, the transporting comprises binding of the composition to TfR. In some aspects, the transporting further comprises transcytosis. In some aspects, the transcytosis is vesicular transcytosis. In some aspects, the transcytosis is TfR-mediated. In some aspects, the transporting further comprises dissociation of the composition from the TfR. In some aspects, the transporting further comprises releasing an active agent from the composition.

[0113] In various aspects, the present disclosure provides a method for designing a peptide capable of binding to TfR, the method comprising: using a computer program to identify a peptide from a library of peptides with optimized stability and folding, wherein each peptide of the library of peptides comprises at least three intramolecular disulfide bonds; using a computer program to superimpose amino acid residues of a binding patch from a peptide:TfR complex to determine a stable peptide that binds to TfR after grafting the binding patch onto the peptide; and grafting the binding patch onto the peptide to produce the stable peptide.

[0114] In some aspects, the method further comprises performing site saturation mutagenesis to obtain a mutated peptide with a higher binding affinity for TfR than the stable peptide. In some aspects, the peptide from a library of peptides is between 20 and 75 amino acid residues in length.

[0115] In various aspects, the present disclosure provides any of the compositions disclosed herein, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof.

[0116] In various aspects, the present disclosure provides any of the methods disclosed herein, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof.

[0117] In various aspects, the present disclosure provides any of the methods disclosed herein, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof.

[0118] In various aspects, the present disclosure provides any of the methods disclosed herein, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof.

INCORPORATION BY REFERENCE

[0119] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0120] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee. The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0121] **FIG. 1** illustrates a Coomassie stained gel of human soluble transferrin receptor (hTfR) ectodomain protein and flow cytometry plots showing successive enrichment of cells that bind to hTfR ectodomain from a pooled, highly diverse peptide library.

[0122] **FIG. 1A** illustrates a Coomassie stained gel of transferrin receptor (TfR) protein.

[0123] **FIG. 1B** illustrates a flow cytometry plot showing cells that bind to TfR after one flow sort.

[0124] **FIG. 1C** illustrates a flow cytometry plot showing cells of a control stain after one flow sort.

[0125] **FIG. 1D** illustrates a flow cytometry plot showing cells that bind to TfR after two flow sorts.

[0126] **FIG. 1E** illustrates a flow cytometry plot showing cells of a control stain after two flow sorts.

[0127] **FIG. 1F** illustrates a flow cytometry plot showing cells that bind to TfR after three flow sorts.

[0128] **FIG. 1G** illustrates a flow cytometry plot showing cells of a control stain after three flow sorts.

[0129] **FIG. 2** illustrates cells displaying TfR and a single clonal peptide. The control protein used in this experiment has an amino acid sequence set forth in SEQ ID NO: 333

(MRLAVGALLVCAVLGLCLADYKDEHHHHHHGLNDIFEAQKIEWHEGGGSKTVRWCAVSEHEA
TKCQSFDRDHMKSVIPSDGPSVACVKKASYLDCIRAIANEADAVTL DAGLVYDAYLAPNNLKPV
VAEFYGSKEDPQTFYYAVAVVKKDSGFQMNQLRGKKSCHTGLGRSAGWNIPIGLLYCDLPEPRK
PLEKAVANFFSGSCAPCADGTDFPQLCQLCPGCGCSTLNQYFGYSGAFKCLKDGAGDVA FVKHS
TIFENLANKADRDQYELLCLDNTRKPVDEYKDCHLAQVPSHTVVARSMGGKEDLIWELLNQAQ
EHFGKDKSKEFQLFSSPHGKDLLFKDSA HGF LKVP PRMDAKMYLGYEYVTAIRNLREGTCPEAP
TDECKPVKWCALSHHERLKCDEWSVNSVGKIECVSAETTEDCIAKIMNGEADAMSLDGGFVYIA
GKCGLVPVLAENYDKSDNCEDTPEAGYFAVAVVKKASDLTWDNLKGKKSCHTAVGRTAGWN
IPMGLLYNKINHCRFDEFFSEGCAPGSKKDSSLCKLCMGSGNLCEPNNKEGYGYTGAFRCLVE
KGDVAFVKHQ TVPQNTGGKNPDPWAKNLNEKDYELLCLDGTRKPV E EYANCHLARAPNHAVV
TRKDKEACVHKILRQQH LFGSDVTDCSGNFCLFRSETKDLLFRDDTVCLAKLHDRNTYEKYL
G E EYVKA VG NLRKCSTSSLLEACTFRRPGSS ELYENKPRRPYIL).

[0130] **FIG. 2A** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and a control protein (y-axis, stained with a fluorescent anti-His antibody).

[0131] **FIG. 2B** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and a control protein (y-axis, stained with a fluorescent streptavidin).

[0132] **FIG. 2C** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and TfR (y-axis, stained with a fluorescent anti-His antibody).

[0133] **FIG. 2D** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and a TfR (y-axis, stained with a fluorescent streptavidin).

[0134] **FIG. 3** illustrates TfR-binding for peptide variants arising from permuting enriched variants from site-saturation mutagenesis (SSM). Each graph represents a round of SSM completed and the colored bars within the graphs indicate the number of mutations in the variant peptides in comparison to the respective peptide sequence with which the round of SSM was started. The data represent the last step of SSM, after which the next generation molecule is settled upon.

[0135] **FIG. 3A** illustrates the amount of hTfR bound for variants comprising sequences of SEQ ID NO: 3 – SEQ ID NO: 23, respectively, derived from a site-saturation mutagenesis (SSM) for affinity maturation of the peptide having a sequence of SEQ ID NO: 1.

[0136] **FIG. 3B** illustrates the amount of hTfR bound for peptide variants having sequences of SEQ ID NO: 24 – SEQ ID NO: 31, respectively derived from a site-saturation mutagenesis (SSM) for affinity maturation of the starting peptide having a sequence of SEQ ID NO: 2.

[0137] **FIG. 4** illustrates sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) under reducing (DTT) or non-reducing (PBS) conditions and RP-HPLC plots of five soluble TfR-binding peptide variants having a sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 30, and SEQ ID NO: 32, respectively. The data show that the TfR-binding peptides exhibit high solubility. Comparison of the gels under reducing conditions (10 mM DTT) and non-reducing (PBS) conditions demonstrates that the peptides form disulfide bonds.

[0138] **FIG. 4A** illustrates SDS-PAGE and an RP-HPLC chromatogram of SEQ ID NO: 1.

[0139] **FIG. 4B** illustrates SDS-PAGE and an RP-HPLC chromatogram of SEQ ID NO: 2.

[0140] **FIG. 4C** illustrates SDS-PAGE and an RP-HPLC chromatogram of SEQ ID NO: 4.

[0141] **FIG. 4D** illustrates SDS-PAGE and an RP-HPLC chromatogram of SEQ ID NO: 30.

[0142] **FIG. 4E** illustrates SDS-PAGE and an RP-HPLC chromatogram of SEQ ID NO: 32.

[0143] **FIG. 5** illustrates that later generation TfR-binding peptides have improved resistance to glutathione (GSH) reduction. **FIG. 5** illustrates overlays (one per peptide tested) of reversed-phase HPLC traces of peptides subjected to either no treatment (red), 10 μ M GSH (blue), or 10 μ M DTT (green).

[0144] **FIG. 5A** illustrates resistance of a TfR-binding peptide having a sequence of SEQ ID NO: 1 to glutathione (GSH) reduction.

[0145] **FIG. 5B** illustrates improved TfR-binding peptide resistance to glutathione (GSH) reduction of the peptide having a sequence of SEQ ID NO: 2.

[0146] **FIG. 5C** illustrates improved TfR-binding peptide resistance to glutathione (GSH) and dithiothreitol (DTT) reduction of the peptide having a sequence of SEQ ID NO: 32.

[0147] **FIG. 5D** illustrates improved TfR-binding peptide resistance to glutathione (GSH) reduction of the peptide having a sequence of SEQ ID NO: 30.

[0148] **FIG. 6** illustrates improved resistance of affinity matured TfR-binding peptides to pepsin improves as peptides mature. **FIG. 6** illustrates overlays (one per peptide tested) of reversed-

phase HPLC traces of TfR-binding peptides subjected to either no treatment (red), trypsin (blue), or pepsin (green).

[0149] **FIG. 6A** illustrates treatment with trypsin (blue) and pepsin (green) of the peptide having a sequence of SEQ ID NO: 1.

[0150] **FIG. 6B** illustrates treatment with trypsin (blue) and pepsin (green) of the peptide having a sequence of SEQ ID NO: 2.

[0151] **FIG. 6C** illustrates treatment with trypsin (blue) and pepsin (green) of the peptide having a sequence of SEQ ID NO: 32.

[0152] **FIG. 6D** illustrates treatment with trypsin (blue) and pepsin (green) of the peptide having a sequence of SEQ ID NO: 30.

[0153] **FIG. 7** illustrates surface plasmon resonance (SPR) curves showing binding of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, and SEQ ID NO: 32 to hTfR.

[0154] **FIG. 8** illustrates a surface plasmon resonance (SPR) trace showing hTfR-binding for varying concentrations of the peptide having a sequence of SEQ ID NO: 2.

[0155] **FIG. 9** illustrates a surface plasmon resonance (SPR) trace showing hTfR-binding for varying concentrations of the peptide having a sequence of SEQ ID NO: 4.

[0156] **FIG. 10** illustrates binding and single cycle kinetics data of SEQ ID NO: 32 binding to captured biotinylated hTfR by surface plasmon resonance (SPR). 5 concentrations of a peptide having a sequence of SEQ ID NO: 32 (0.037 nM, 0.11 nM, 0.33 nM, 1 nM, 3 nM) were injected over 2 densities of captured biotinylated (Bt)-hTfR and analyzed globally.

[0157] **FIG. 11** illustrates binding and single cycle kinetics data of SEQ ID NO: 30 binding to captured biotinylated hTfR by SPR. 5 concentrations of a peptide having a sequence of SEQ ID NO: 30 (0.037 nM, 0.11 nM, 0.33 nM, 1 nM, 3 nM) were injected over 2 densities of captured Bt-hTfR and analyzed globally.

[0158] **FIG. 12** illustrates a bar graph quantitating flow cytometry data from a competitive binding assay in which peptides having a sequence of SEQ ID NO: 2 and SEQ ID NO: 32 were incubated with hTfR in the presence of 10 μ M of iron-loaded human holo-transferrin or empty human apo-transferrin.

[0159] **FIG. 13** illustrates co-crystallization of hTfR and the hTfR-binding peptide having a sequence of SEQ ID NO: 32.

[0160] **FIG. 13A** illustrates a first view of hTfR co-crystallized with the peptide having a sequence of SEQ ID NO: 32. The structures show that the binding site of the hTfR-binding

peptides overlaps with that of the endogenous binder transferrin. The linking sequence between the two helices, depicted as a solid, black coiled line, was not resolved in crystal structure and is based on structural modeling. The crystal structure of the transferrin receptor (TfR) is displayed as a space-filling electron density map. It appears as a large, globular protein with two prominent lobes. In between the two lobes the crystal structure of a short peptide, corresponding to SEQ ID NO: 32, is shown as interacting with the globular structure of the transferrin receptor. The peptide of SEQ ID NO: 32 is shown as a cartoon model with two anti-parallel α -helices oriented diagonally from upper left to lower right. Stick models corresponding to the amino acid side chains of SEQ ID NO: 32 are shown protruding from the α -helices, some of which are interacting with the globular protein of TfR.

[0161] FIG. 13B illustrates a zoomed in view of **FIG. 13A** and shows hTfR co-crystallized with a peptide having a sequence of SEQ ID NO: 32. Amino acid side chains either mutated from endogenous CDP or those that are part of the interface are shown as sticks. Interface-adjacent residues are shown in black, whereas interface-distal residues are shown in cyan. A portion of the TfR space-filling electron density map depicted in **FIG. 13A** that interacts with the peptide of SEQ ID NO: 32 is shown. The peptide of SEQ ID NO: 32, as shown in **FIG. 13A**, is displayed as a cartoon model with two anti-parallel α -helices and is positioned directly above the structure of TfR, with the α -helices positioned horizontally. The amino acid side chains of SEQ ID NO: 32 are shown as sticks. The side chains that protrude downward from the α -helices of SEQ ID NO: 32 toward the globular structure of TfR interact with TfR.

[0162] FIG. 13C illustrates the cartoon representations of a structural model of SEQ ID NO: 32 (top, N-terminus in blue, C-terminus in red, visible side chains colored by atom type), the crystal structure of free SEQ ID NO: 32 (middle, blue; 2.55 Å resolution, 4-fold superposition), and the crystal structure of SEQ ID NO: 32 when complexed with TfR (bottom, red; 1.85 Å resolution, 2-fold superposition). Side-chains with an atom within 4 Å of TfR in the complex structure are shown as sticks. N- and C-termini are labeled. Red arrows mark the most significant structural differences between the model, unbound, and bound structures of SEQ ID NO: 32 near the N-terminus. Gray crosses mark the location of the disordered inter-helical linker. The top structure, labeled “Model,” shows the *in silico* model of the peptide corresponding to SEQ ID NO: 32 displayed as a cartoon model. The anti-parallel α -helices, oriented horizontally, are connected on the right side by a short loop. The N-terminus of the peptide is located at the left side of the bottom helix, and the C-terminus is located at the left side of the upper helix. Amino acid side

chains are shown as sticks protruding from the α -helices. The middle structure, labeled “4x free,” shows the crystal structure of the peptide corresponding to SEQ ID NO: 32 crystallized in the absence of TfR displayed as a cartoon model. The loop seen in the *in silico* model is not visible in this structure, as denoted by an X to the right of the two helices. The N-terminus of the peptide is located at the left side of the bottom helix, and the C-terminus is located at the left side of the upper helix. Amino acid side chains are shown as sticks protruding from the α -helices. The bottom structure, labeled “2x TfR bound,” shows the crystal structure of the peptide corresponding to SEQ ID NO: 32 crystallized in the presence of TfR displayed as a cartoon model. The loop seen in the *in silico* model is not visible in this structure, as denoted by an X to the right of the two helices. The N-terminus of the peptide is located at the left side of the bottom helix, and the C-terminus is located at the left side of the upper helix. Amino acid side chains are shown as sticks protruding from the α -helices. Three triangular arrows pointing to the N-terminal region at the left end of the lower helix in each structure indicate the regions of the most significant structural differences between the three structures.

[0163] **FIG. 13D** illustrates a zoomed-in view of binding of SEQ ID NO: 32 bound to TfR, showing numerous polar and non-polar interactions driving binding. Non-polar interactions are primarily Leu and Met, while polar interactions are a mix of several charged and non-charged residues. A portion of the crystal structure of TfR is shown as a space-filling electron density map. The peptide corresponding to SEQ ID NO: 2 is shown as a ribbon diagram on top of the TfR structure. The α -helices are shown oriented vertically, and a loop connects the top ends of the two helices. Amino acid side chains are shown as sticks, and side chains facing the structure of TfR are interacting with TfR. Amino acid residues of SEQ ID NO: 32 are labeled as follows (from top to bottom, then from left to right): E16, M9, S6, N12, L43, L15, L36, and D44.

[0164] **FIG. 13E** illustrates the alignment of the TfR:SEQ ID NO: 32 co-crystal with that of the published TfR:transferrin complex (PDB 3S9L; TfR hidden) and is zoomed in on the binding interactions shown in **FIG. 14A**. The structural alignment (left) demonstrates an overlap of the binding sites for SEQ ID NO: 32 (blue) and transferrin

(MRLAVGALLVCAVLGLCLADYKDEHHHHHHGLNDIFEAQKIEWHEGGGSKTVRWCA
VSEHEATKCQSFRDHMKSVIPSDGPSVACVKKASYLDCIRAI AANEADAVTL DAGLVYD
AYLAPNNLKPVVAEFYGSKEDPQTFYYAVAVVKKDSGFQMNQLRGKKSCHTGLGRSA
GWNIPIGLLYCDLPEPRKPLEKAVANFFSGSCAPCADGTDFPQLCQLCPGCGCSTLNQYF
GYSGAFKCLKDGAGDVAFVKHSTIFENLANKADRDQYELLCLDNTRKPVDEYKDCHLA

QVPSHTVVARSMGGKEDLIWELLNQAQEHFGKDKSKEFQLFSSPHGKDLLFKDSAHGFL
 KVPPRMDAKMYLGYEYVTAIRNLREGTCPEAPTDECKPVKWCALSHHERLKCDEWSV
 NSVGKIECVSAETTEDCIAKIMNGEADAMSLDGGFVYIAGKCGLVPVLAENYDKSDNCE
 DTPEAGYFAVAVVKKASDLTWDNLKGKKSCHTAVGRTAGWNIPMGLLYNKINHCRF
 DEFFSEGCAPGSKKDSLLCKLCMGSGNLCEPNNKEGYGYTGAFRCLVEKGDVAFVK
 HQTVPQNTGGKNPDPWAKNLNEKDYELLCLDGTRKPVVEYANCHLARAPNHAVVTRK
 DKEACVHKILRQQQHLFGSDVTDCSGNFCLFRSETKDLLFRDDTVCLAKLHNRNTYEKY
 LGEEYVKA VGNLRKCSTSSLLEACTFRRP, SEQ ID NO: 353, green). The SEQ ID NO: 32

binding site, colored for human (SEQ ID NO: 353) and murine

(MRLAVGALLVCAVLGLCLADYKDEHHHHHHGLNDIFEAQKIEWHEGGGSVPDKTVK
 WCAVSEHENTKCISFRDHMKTVLPPDGPRLACVKKTSYPDCIKAISASEADAMTLDGG
 WVYDAGLTPNNLKPVAAEFYGSVEHPQTYYYAVAVVKKGTDFQLNQLEGKKSCHTGL
 GRSAGWVIPIGLLFCKLSEPRSPLEKAVSSFFSGSCVPCADPVAFPKLCQLCPGCGCSSTQ
 PFFGYVGAFKCLKDGGGDVAFVKHTTIFEVLPEKADRDQYELLCLDNTRKPVDQYEDC
 YLARIPSHAVVARKNNGKEDLIWEILKVAQEHFGKGKSKDFQLFSSPLGKDLLFKDSAF
 GLLRVPPRMDYRLYLGHNYVTAIRNQEGVCPEGSIDNSPVKWCALSHLERTKCDEWSI
 ISEGKIECESAETTEDCIEKIVNGEADAMTLDGGHAYIAGQCGLVPVMAEYYESSNCAIP
 SQQGIFPKGYAVAVVKKASDTSITWNNLKGKKSCHTGVDRTAGWNIPMGMLYNRINHC
 KFDEFFSQGCAPGYEKNSTLCDLCIGPLKCAPNNKEEYNGYTGAFRCLVEKGDVAFVKH
 QTVLDNTEGKNPAEWAKNLKQEDFELLCPDGTRKPVKDFASCHLAQAPNHVVVSRKE
 KAARVKAVLTSQETLFGGSDCTGNFCLFKSTTKDLLFRDDTKCFVKLPEGTTPEKYLGA
 EYMQSVGNMRKCSTSRLLLEACTFHKH, SEQ ID NO: 354) transferrin receptor homology,

shows that most of the binding surface on TfR shares identity or similarity between the two species (right). The left panel shows a portion of the structure of TfR as a space-filling electron density map. The ribbon diagram of transferrin from the 3S9L crystal structure (labeled "Tf from 3S9L") is positioned to the right of TfR and superimposed on the TfR structure. The Tf structure contains numerous α -helices which fit with the TfR structure. The structure also contains a β -sheet structure in the lower right region, as well as numerous loops protruding from the ends of the α -helices. The structure of SEQ ID NO: 32 crystallized with TfR is shown as a cartoon model interacting with TfR on the upper right side. It consists of two α -helices positioned vertically that overlap with the superimposed Tf structure. The right panel shows a rotated view of the TfR and SEQ ID NO: 32 structure shown in the left panel. TfR is shown as a space-filling electron density

map, and SEQ ID NO: 32 appears as a cartoon model containing two α -helices oriented diagonally from upper left to lower right. Shaded regions on the TfR structure denote regions where the HsTfR and MmTfR binding sites are either similar or dissimilar. Exemplary similar and dissimilar regions are indicated by arrows. The remaining regions of the structure not denoted as similar or dissimilar, making up the majority of the surface of the TfR structure, has identical binding sites. A small region directly above the right helix of SEQ ID NO: 32 and a small region directly beneath the upper end of the right helix are denoted as having a similar binding site. A small region directly above the right helix of SEQ ID NO: 32, a small region directly beneath the lower end of the right helix, and a small region below the lower end of the left helix are denoted as having a dissimilar binding site.

[0165] FIG. 14 illustrates a TfR:transferrin complex and a TfR:peptide complex showing superimposition of each of transferrin (data taken from published literature (RCSB PDB 1SUV)) and the peptide having a sequence of SEQ ID NO: 32 bound to hTfR, illustrating that transferrin and the peptide having a sequence of SEQ ID NO: 32 overlap in their binding sites on hTfR.

[0166] FIG. 14A illustrates the superimposition of each of transferrin (data taken from published literature (RCSB PDB 1SUV)) and the peptide having a sequence of SEQ ID NO: 32 bound to hTfR. The structure of TfR appears as a space-filling electron density map of a large, globular protein having two lobes. The structure of transferrin from the 1SUV crystal structure (labeled “Transferrin from RCSB 1SUV”) appears as a cartoon model with numerous α -helices and connecting loops. The transferrin structure is positioned to the lower right of TfR and fits with the TfR structure. An arrow labeled “SEQ ID NO: 32” points to a site to the right of TfR, directly above transferrin, and indicates the binding site of the peptide corresponding to SEQ ID NO: 32.

[0167] FIG. 14B illustrates a zoomed in view of the peptide having a sequence of SEQ ID NO: 32 bound to TfR, indicating that, when compared to the binding site of endogenous transferrin in **FIG. 14A**, both the peptide having a sequence of SEQ ID NO: 32 and transferrin have overlapping binding sites on hTfR. Amino acid side chains either mutated from endogenous CDP or that are part of the interface are shown as sticks. Interface-adjacent residues are shown in black, whereas interface-distal residues are shown in cyan. A portion of the TfR space-filling electron density map depicted in **FIG. 13A** that interacts with the peptide of SEQ ID NO: 32 is shown. The peptide of SEQ ID NO: 32, as shown in **FIG. 13A**, is displayed as a cartoon model with two anti-parallel α -helices and is positioned directly above the structure of TfR, with the α -helices positioned horizontally. The amino acid side chains of SEQ ID NO: 32 are shown as

sticks. The side chains that protrude downward from the α -helices of SEQ ID NO: 32 toward the globular structure of TfR interact with TfR.

[0168] FIG. 15 illustrates that TfR-binding peptides are cross-reactive with murine TfR (mTfR) in cell surface binding assays. 293F cells expressing either human or mouse TfR from their surface were stained with soluble TfR-binding peptides that were directly labeled with AlexaFluor 647 dye.

[0169] FIG. 15A illustrates the species specificity of the TfR used in these experiments, in this case human TfR. Data is displayed as two topographical density maps and indicates flow cytometry data of transferrin stained with Anti-hTfR (CD71) antibody. The upper density map, oriented diagonally from lower left to upper right, depicts 293ST+SDGF-hTfR. The lower density map, oriented horizontally, depicts 293ST+SDGF-mTfR. The y-axis shows hTfR + Streptavidin from 0 to 10^7 , in increments of 10 on a log scale. The x-axis shows GFP from 0 to 10^6 , in increments of 10 on a log scale.

[0170] FIG. 15B illustrates the species specificity of the TfR used in these experiments, in this case murine TfR. Data is displayed as two topographical density maps and indicates flow cytometry data of transferrin stained with Anti-mTfR (CD71) antibody. The upper density map, oriented diagonally from lower left to upper right, depicts 293ST+SDGF-mTfR. The lower density map, having three lobes, depicts 293ST+SDGF-hTfR. The y-axis shows hTfR + Streptavidin from 10^{-4} to 10^7 , in increments of 10 on a log scale. The x-axis shows GFP from 0 to 10^6 , in increments of 10 on a log scale.

[0171] FIG. 15C illustrates that the peptide having a sequence of SEQ ID NO: 1, the peptide having a sequence of SEQ ID NO: 2, the peptide having a sequence of SEQ ID NO: 30, and the peptide having a sequence of SEQ ID NO: 32 bind human TfR. Data is displayed as four topographical density maps and indicates flow cytometry data using 293ST cells + SDGF-hTfR. Three density maps appear nearly superimposed and are oriented above a fourth density map. The lower density map is oriented horizontally and depicts SEQ ID NO: 1 (1st gen). The upper three density maps are oriented diagonally from lower left to upper right. The density map slightly above the other two corresponds to SEQ ID NO: 32 (3rd gen). The density map slightly below the other two corresponds to SEQ ID NO: 2 (2nd gen). The third density map corresponds to SEQ ID NO: 30 (3rd gen). The y-axis shows hTfR + Streptavidin from 0 to 10^7 , in increments of 10 on a log scale. The x-axis shows GFP from 0 to 10^6 , in increments of 10 on a log scale.

[0172] **FIG. 15D** illustrates that the peptide having a sequence of SEQ ID NO: 1, the peptide having a sequence of SEQ ID NO: 2, the peptide having a sequence of SEQ ID NO: 30, and the peptide having a sequence of SEQ ID NO: 32 bind murine TfR. Data is displayed as four topographical density maps and indicates flow cytometry data using 293ST cells + SDGF-mTfR. Three density maps appear nearly superimposed and are oriented above a fourth density map. The lower density map is oriented horizontally and depicts SEQ ID NO: 1 (1st gen). The upper three density maps are oriented diagonally from lower left to upper right. The density map slightly above the other two corresponds to SEQ ID NO: 32 (3rd gen). The density map slightly below the other two corresponds to SEQ ID NO: 2 (2nd gen). The third density map corresponds to SEQ ID NO: 30 (3rd gen). The y-axis shows hTfR + Streptavidin from 0 to 10⁷, in increments of 10 on a log scale. The x-axis shows GFP from 0 to 10⁶, in increments of 10 on a log scale.

[0173] **FIG. 16** illustrates whole body autoradiography of mice injected intravenously with ¹⁴C-labeled TfR-binding peptides (SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 30, and SEQ ID NO: 32) to assess biodistribution. Images were taken 3 hours after a dose of 100 nmol peptide (20 mg/kg, assuming approximately 25 g body weight) was administered to mice bearing xenograft tumors (subcutaneous human astrocytoma U87 cells). The images demonstrate substantial accumulation in spleen, liver, kidney, and also high accumulation in muscle, bone marrow, and skin. CNS accumulation is less than other tissues but still substantial (>25% of serum) compared to historical values of compounds and biologics that do not penetrate the blood-brain barrier (BBB) - background blood level found in brain is 3% that of cardiac blood signal.

[0174] **FIG. 16A** illustrates whole body autoradiography of mice injected intravenously with ¹⁴C-labeled SEQ ID NO: 1.

[0175] **FIG. 16B** illustrates whole body autoradiography of mice injected intravenously with ¹⁴C-labeled SEQ ID NO: 2.

[0176] **FIG. 16C** illustrates whole body autoradiography of mice injected intravenously with ¹⁴C-labeled SEQ ID NO: 30.

[0177] **FIG. 16D** illustrates whole body autoradiography of mice injected intravenously with ¹⁴C-labeled SEQ ID NO: 32.

[0178] **FIG. 17** illustrates whole body autoradiography of mice injected intravenously with ¹⁴C-labeled peptides (SEQ ID NO: 1 and SEQ ID NO: 32) to assess biodistribution. Substantial accumulation in flank tumors was also seen. Accumulation in tumors is dispersed throughout,

rather than concentrated at sites rich in blood. This suggests extravasation and dispersal throughout the tumor parenchyma.

[0179] FIG. 17A illustrates whole body autoradiography of mice injected intravenously with ^{14}C -labeled SEQ ID NO: 1.

[0180] FIG. 17B illustrates whole body autoradiography of mice injected intravenously with ^{14}C -labeled SEQ ID NO: 1.

[0181] FIG. 17C illustrates whole body autoradiography of mice injected intravenously with ^{14}C -labeled SEQ ID NO: 32.

[0182] FIG. 17D illustrates whole body autoradiography of mice injected intravenously with ^{14}C -labeled SEQ ID NO: 32.

[0183] FIG. 18 illustrates the quantitation of peptide biodistribution and organ accumulation after a single 20 mg/kg IV dose of either a TfR-binding peptide having a sequence of SEQ ID NO: 1 or a peptide having a sequence of SEQ ID NO: 32. This dose achieved levels of >150 nM in the CNS, ~ 1 μM in tumor after 3 hours. The bar graph quantifies the levels of each ^{14}C -labeled peptides with known radioactivity in the same image and to peptide specific activity in various tissues from mice injected intravenously with peptides, 3 hours post-administration, as measured by whole body autoradiography.

[0184] FIG. 19 illustrates the quantitation of peptide biodistribution and organ accumulation for the peptides having a sequence of SEQ ID NO: 1 and SEQ ID NO: 32. The values are shown as levels in the tissue versus levels in the blood, where no brain penetration would give a level of 3% given that brain is 3% blood. The bar graph quantifies the levels of each ^{14}C -labeled peptides normalized to controls with known radioactivity in the same image and to peptide specific activity in various tissues from mice injected intravenously with peptides, 3 hours post-administration, as measured by whole body autoradiography.

[0185] FIG. 20 illustrates scintillation counting over time in various tissues after administration of 20 mg/Kg of TfR-binding peptide having a sequence of SEQ ID NO: 32 with a specific activity of 147 Ci/mol, validating whole body autoradiography (WBA) tissue accumulation.

[0186] FIG. 20A illustrates scintillation counting over time in the kidneys. The y-intercept at time 0 (Y_0) is 1900 pCi, the half-life ($t_{1/2}$) is 0.89 hrs, and the R^2 of the fit is 0.89.

[0187] FIG. 20B illustrates scintillation counting over time in the liver. The y-intercept at time 0 (Y_0) is 400 pCi, the half-life ($t_{1/2}$) is 7.22 hrs, and the R^2 of the fit is 0.97.

[0188] **FIG. 20C** illustrates scintillation counting over time in the brain. The y-intercept at time 0 (Y0) is 7.3 pCi, the half-life ($t_{1/2}$) is 0.72 hrs, and the R^2 of the fit is 0.45.

[0189] **FIG. 20D** illustrates scintillation counting over time in the spleen. The y-intercept at time 0 (Y0) is 168 pCi, the half-life ($t_{1/2}$) is 12.1 hrs, and the R^2 of the fit is 0.93.

[0190] **FIG. 20E** illustrates scintillation counting over time in the muscle. The y-intercept at time 0 (Y0) is 31.4 pCi, the half-life ($t_{1/2}$) is 2.54 hrs, and the R^2 of the fit is 0.85.

[0191] **FIG. 20F** illustrates scintillation counting over time in the skin. The y-intercept at time 0 (Y0) is 68.3 pCi, the half-life ($t_{1/2}$) is 0.33 hrs, and the R^2 of the fit is 0.31.

[0192] **FIG. 21** illustrates a serum elimination plot, including two-phase elimination regression analysis showing fast (95%, 15.6 min $t_{1/2}$) and slow (5%, 10.3 hr $t_{1/2}$) phase kinetics in mice intravenously administered 20 mg/kg of peptide having a sequence of SEQ ID NO: 32.

[0193] **FIG. 22** illustrates multiple time regression analysis and capillary depletion analysis of TfR-binding peptides having a sequence of SEQ ID NO: 1 and SEQ ID NO: 32.

[0194] **FIG. 22A** illustrates multiple regression analysis of SEQ ID NO: 1 and SEQ ID NO: 32 in a single plot, wherein the y-axis indicates the brain to serum ratio.

[0195] **FIG. 22B** illustrates parenchyma and capillary distribution of TfR-binding peptides having a sequence of SEQ ID NO: 1 and SEQ ID NO: 32, wherein the y-axis indicates the tissue-to-serum ratio ($\mu\text{L/g}$).

[0196] **FIG. 23** illustrates the purification and testing of soluble transferrin receptor (TfR) ectodomain.

[0197] **FIG. 23A** illustrates the surface plasmon resonance (SPR) of holo or apo transferrin (Tf) binding to the purified TfR. Holo Tf binds TfR, but apo Tf does not, as seen by the increase in response (RU) over time.

[0198] **FIG. 23B** illustrates a schematic of the vector display scaffold and target engagement used to screen for and optimize peptide binding properties. The surface display vector (SDGF) encoding a GFP-tagged construct of the binder (e.g., SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 32) is expressed on the cell surface. A target protein (e.g., TfR) labeled with a fluorescent dye ("Co-Stain") bind to the surface-expressed binder.

[0199] **FIG. 23C** illustrates flow cytometry to screen for specific TfR binding, as measured by the amount of Alexa Fluor 647-TfR (co-stain in **FIG. 23B**) bound. Cells transfected with Machupo virus glycoprotein (SDGF-MaCV) are tested with a combination of biotinylated TfR and Alexa Fluor 647-labeled streptavidin (Strep-647), SDGF-MaCV cells and Alexa Fluor 647-

labeled elastase, or SDGF-elafin cells and TfR + Strep-647. The elastase and elafin cells conjugates fail to bind cells.

[0200] **FIG. 24** illustrates whole body autoradiography (WBA) of ^{14}C peptide variant distribution. Histological sections of mice treated with ^{14}C peptide variant are shown 30 minutes and 180 minutes post-administration. Representative brain and heart (inset) WBA images of three peptide generations, plus that of a control, non-TfR binding CDP are shown. Full body images of the same sections are shown in **FIG. 25** and some of the 180 min. sections are also shown in **FIG. 16**.

[0201] **FIG. 25** illustrates the full body images of the whole body radiography shown in **FIG. 24**, at 30 minutes (left column) and 180 minutes (right column) post-administration. Some of the sections from 180 min. are also shown in **FIG. 16**.

[0202] **FIG. 26** illustrates the heat stability of SEQ ID NO: 32 under reduced and non-reduced conditions as measured using circular dichroism. Relative ellipticity at 210 nm (mdeg) is plotted as a function of temperature in degrees C. NR indicates non-reduced. DTT indicates reducing the peptide with DTT. Vertical lines represent calculated melting points ($\sim 74^\circ\text{C}$ for NR, $\sim 38^\circ\text{C}$ for 10 mM DTT treated) based on non-linear curve fitting (sigmoidal, variable slope, constrained bottom = 0; $R^2 > 0.98$ for both). Note that the NR sample failed to reach equilibrium up to 95°C , so it is likely that the protein is not fully “melted” at high temperature, but simply more disordered, particularly in the extended loop region.

[0203] **FIG. 27** illustrates the purification and characterization of murine transferrin, murine transferrin-NT fusion, and various CDP-neurotensin (NT)-complexes of the present disclosure. SDS-PAGE under reducing (DTT) and non-reducing (PBS) conditions and gels stained with Coomassie dye show the purity of the peptides following purification. As shown, are murine endogenous transferrin (denoted MmTf, SEQ ID NO: 354), murine endogenous transferrin conjugated to neurotensin (denoted MmTf-NT, SEQ ID NO: 355), and CDP-NT peptide constructs (SEQ ID NO: 33 – SEQ ID NO: 35). Gels are run under reducing (DTT) or non-reducing (PBS) conditions. Reduction with 10 mM DTT suggested that the peptides form disulfide bonds.

[0204] **FIG. 28** illustrates CDP-NT peptide constructs which induce an IP_1 response downstream of the neurotensin receptor (NTSR) both in CRE-Luciferase (CRE-Luc) mice and in mammalian cells.

[0205] **FIG. 28A** illustrates the relevant pathways influencing CRE-driven luciferase in the CRE-Luc mice. PLC denotes phospholipase C. AC denotes adenylyl cyclase. CaMK denotes calmodulin-dependent protein kinase. CREB denotes the cAMP response element binding protein. PKA denotes protein kinase A. PDE denotes cAMP phosphodiesterase. FS denotes forskolin. Rol denotes rolipram. GPCR denotes a G-protein-coupled receptor.

[0206] **FIG. 28B** illustrates *in vitro* neurotensin (NT) receptor engagement showing IP₁ accumulation only in response to NT or NT peptide constructs in HEK-293 cells expressing *NTSR1*. IP₁ is measured using an assay kit (CisBio 62IP APEB) with a readout of FRET ratio. N = 3 wells for all except vehicle, which had N = 36. Horizontal bar indicates sample mean. mTF = murine transferrin. Baseline HEK293 = mean assay value for HEK293 cells (N=36 wells) that do not express *NTSR1*, included as a reference.

[0207] **FIG. 29** illustrates immunohistochemistry stained tissue samples from mice administered NT peptide constructs. Tissue staining shows enhanced luciferase expression in the cortex, striatum, and thalamus of mice 4 hours after CDP-NT administration of SEQ ID NO: 33, SEQ ID NO: 34, and SEQ ID NO: 35. Mice administered Transferrin-NT (“Transferrin-NT,” SEQ ID NO: 355) or no peptide (“Unstimulated”) are shown as controls. Scale bar (top left panel) is 100 μ m; all panels are the same magnification.

[0208] **FIG. 30** illustrates quantitation of the effects of CDP-NT peptide constructs in CRE-luciferase (CRE-Luc) mice.

[0209] **FIG. 30A** illustrates luminescence (via intraperitoneal luciferin dosage) either before (unstimulated) or four hours after (denoted “NT fusion” or “Parent”) intravenous administration of parent TfR-binding peptides (“Parent”) (e.g., SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 32, or SEQ ID NO: 354 as denoted) or cystine dense peptide (CDP)-NT constructs or murine transferrin-NT constructs (e.g., comprising SEQ ID NO: 33 – SEQ ID NO: 35 or SEQ ID NO: 355 as denoted) (“NT fusion”), using matched cohorts. Horizontal bar indicates sample mean. Significance was determined using a T-test (unpaired, 2-tailed). *: P < 0.05. **: P < 0.01. #: P < 0.0001. For both the parent (SEQ ID NO: 354) and NT transferrin peptide constructs (SEQ ID NO: 355), murine transferrin was used.

[0210] **FIG. 30B** illustrates images from the SEQ ID NO: 32 cohort quantitated in **FIG. 30A** with pseudocolored luminescence intensity. All images were equally background-subtracted and contrast-enhanced for ease of viewing only.

[0211] **FIG. 31** illustrates *in vivo* CRE-luciferase luminescence four hours after forskolin and rolipram induction. Images show luciferase fluorescence images of mice and serve as positive controls for the experiments depicted in **FIG. 30**. Forskolin and rolipram activate CRE-dependent luciferase expression independent of the neurotensin receptor pathway. Forskolin and rolipram induce CRE expression via activation of adenylyl cyclase and inhibition of cAMP phosphodiesterase, respectively. N = 8 mice. Panels show replicate animals treated identically with rolipram and forskolin. One mouse was censored after the luciferin injection failed to hit the peritoneal cavity.

[0212] **FIG. 32** illustrates luciferase expression in CRE-Luc mice after intravenous administration of free neurotensin peptide. Intravenous free neurotensin fails to induce luciferase expression in CRE-Luc mice. *In vivo* luminescence of CRE-luciferase mice 4 hours after intravenous administration of high dose free neurotensin peptide (300 nmol) or vehicle (10% DMSO in PBS) was assessed with intraperitoneal luciferin injection and imaged in an IVIS imager. “ns” indicates that the results were not significant (P = 0.48), as calculated by T test (unpaired, 2 tailed).

[0213] **FIG. 33** illustrates the quantitation of peptide biodistribution and organ accumulation after a single 20 mg/kg IV dose of either a TfR-binding peptide having a sequence of SEQ ID NO: 1 or a peptide having a sequence of SEQ ID NO: 32. Peptide levels are measured at 30 min and 3 hours (3 hour data is also shown in **FIG. 18**). This dose achieved levels of >150 nM in the CNS, ~1 μ M in tumor after 3 hours. The bar graph quantifies the levels of each ^{14} C-labeled peptides with known radioactivity in the same image and to peptide specific activity in various tissues from mice injected intravenously with peptides, 3 hours post-administration, as measured by whole body autoradiography. Tissue data is plotted as seven clusters of four bars each. The 4-bar clusters correspond to different tissue types, from left to right, of Skin, Adipose, Muscle, Spleen, Kidney, Liver, Brain, and Blood, as shown on the x-axis. The four bars in each cluster correspond to, from left to right, SEQ ID NO: 1 30 min (N=10 sections), SEQ ID NO: 1 180 min (N=14 sections), SEQ ID NO: 32 30 min (N=12 sections), and SEQ ID NO: 32 180 min (N=21 sections). The y-axis shows nmol g⁻¹ tissue of peptide from 0.01 to 100 in increments of 10 on a log scale. The values of each bar, from left to right are, Skin: 1.33, 1.12, 1.70, 1.02, Adipose: 0.79, 0.52, 0.52, 0.27, Muscle: 0.63, 0.35, 0.44, 0.35, Spleen: 4.37, 20.03, 1.17, 3.60, Kidney: 29.05, 9.71, 16.91, 3.10, Liver: 4.79, 22.18, 1.80, 8.54, Brain: 0.23, 0.16, 0.21, 0.09, and Blood: 1.58, 0.62, 0.86, 0.34. The bars corresponding to Spleen, Kidney, and Liver are noticeably higher

than the other sets. The inset shows the same data from the brain normalized to blood. The four bars correspond to, from left to right, SEQ ID NO: 1 30 min (N=10 sections), SEQ ID NO: 1 180 min (N=14 sections), SEQ ID NO: 32 30 min (N=12 sections), and SEQ ID NO: 32 180 min (N=21 sections). The y-axis shows peptide normalized to blood from 1 to 100 in increments of 10 on a log scale. The values for each bar, from left to right, are: 14%, 26%, 25%, and 27% of blood peptide levels.

[0214] FIG. 34 illustrates the purification of SA21 fusion peptides.

[0215] FIG. 34A shows purification of a TfR-binding peptide fused to a serum albumin peptide (SA21) corresponding to SEQ ID NO: 373. Purity was verified by SDS-PAGE (left) and RP-HPLC (right) under DTT reducing (“R”) or non-reducing (“NR”) conditions. SDS-PAGE was also run on the uncleaved (“U”) siderocalin-CDP fusion peptide.

[0216] FIG. 34B shows purification of a peptide fused to SA21 corresponding to SEQ ID NO: 456

(GSRLIEDICLPRWGCLWEDDGGGGSGGGGSVRIPVSCKHSGQCLKPCKDAGMRFGKC MNGKCDCTPK). Purity was verified by SDS-PAGE (left) and RP-HPLC (right) under DTT reducing (“R”) or non-reducing (“NR”) conditions. SDS-PAGE was also run on the uncleaved (“U”) siderocalin-CDP fusion peptide.

DETAILED DESCRIPTION

[0217] Drug delivery to targets located in the central nervous system (CNS) can be hindered by the blood-brain barrier (BBB), a term for the vascular endothelial cells in CNS capillaries. This system exists to prevent toxic metabolites and pathogens from entering the CNS (e.g., the brain), but also serves to render diseases of the CNS particularly difficult to treat using conventional medicines. It can be for this reason that primary CNS tumors (e.g., gliomas) and neuroinflammatory and neurodegenerative diseases such as Multiple Sclerosis and Alzheimer’s disease respond particularly poorly to therapeutics that can otherwise be more effective in similar diseases affecting peripheral tissues where drug delivery to target cells can be less hindered. The myriad disorders of the CNS, from brain cancer to neurodegeneration to age-associated inflammatory processes, necessitate varied approaches to CNS drug delivery. While the BBB allows osmolytes and nutrients into the brain from serum, and CNS astrocytes provide many of the growth and survival signals required by neurons, several larger hormones and proteins such as transferrin (an iron chaperone), insulin, and leptin can cross the BBB. CNS transport of larger

molecules such as transferrin can be accomplished by receptor-mediated transcytosis or “vesicular transcytosis” (e.g., transport of cargo from the apical to the basal side, or vice versa, in intracellular vesicles). For example, variants of the endogenous receptors of insulin, leptin, and transferrin – InsR, ObR, and TfR, respectively – that contain the normal ectodomain can be employed by CNS vascular endothelial cells in order to facilitate transport of these molecules into the CNS. In this highly selective way, certain large molecules like transferrin (molecular weight is approximately 75 kDa) can access the brain parenchyma.

[0218] In various embodiments, the present disclosure provides compositions that enable transport of cargo molecules or active agents (e.g., small molecules, peptides, or proteins) across cell layers or barriers, including endothelial (e.g., the BBB) or epithelial cell layers and methods of using these compositions. In some embodiments, the present disclosure provides compositions and methods that enable delivery of various molecules into the CNS that would otherwise not be able to pass the BBB. In some cases, the present disclosure provides compositions and methods for delivery of therapeutic and/or diagnostic molecules into the CNS, e.g., the brain. Thus, in various embodiments, the present disclosure provides peptides capable of crossing the BBB. These peptides can have an affinity and selectively for a transferrin receptor (TfR). The TfR-binding peptides can be cystine-dense peptides (CDPs). In some cases, the peptides of the present disclosure can cross the BBB via TfR-mediated transcytosis.

[0219] In some embodiments, the presently described peptides can be peptide conjugates, peptide constructs, fusion peptides, or fusion molecules such as linked by chemical conjugation of any molecule type, such as small molecules, peptides, or proteins, or by recombinant fusions of peptides or proteins, respectively (e.g., a peptide construct). The terms “fusion peptide” and “peptide fusion” are used interchangeably herein. In some embodiments, the peptide constructs can be produced biologically or synthetically. Thus, in some cases, a TfR-binding peptide can comprise a TfR-binding peptide domain linked to another molecule or group of molecules such as small molecules, peptides, or proteins or other macromolecules such as nanoparticles.

[0220] In some embodiments, the present disclosure provides methods and compositions that enable transport across cellular or molecular barriers. In various embodiments, the present disclosure provides methods and compositions that enable TfR-mediated transport across cellular layers (e.g., endothelial cells or epithelial cells) or cell membranes. In some cases, the TfR-binding peptides of the present disclosure enable transport across the blood brain barrier (BBB). In various aspects, the peptides of the present disclosure can be used to target any cell expressing

TfR. In addition to the BBB, various other cells, tissues, and organs express TfR. Cells expressing TfR can include hepatocytes, erythrocytes and erythrocyte precursors in bone marrow, immune cells, stem cells, and rapidly dividing cells. Tissues and organs expressing TfR can include the brain (e.g., cerebral cortex, hippocampus, caudate, cerebellum), endocrine tissues (e.g., thyroid, parathyroid, and adrenal glands), bone marrow and immune system (e.g., appendix, lymph node, tonsil, spleen), muscle tissues (e.g., heart, skeletal, and smooth muscle), liver, gallbladder, pancreas, gastrointestinal tract (e.g., oral mucosa, esophagus, stomach, duodenum, small intestine, colon, rectum), kidney, urinary bladder, female tissues (e.g., fallopian tube, breast, vagina, cervix, endometrium, ovary, and placenta), adipose and soft tissue, and skin. Thus, the TfR-binding peptides of the present disclosure can be used to target these cells, tissues, and organs and deliver an active agent to these cells, tissues, and organs via, for example, TfR-mediated transcytosis (e.g., across cellular barrier such as the BBB) or TfR-mediated endocytosis (e.g., across cell membranes into cells).

[0221] In various embodiments, the present disclosure provides methods and compositions that enable TfR-mediated transport and delivery to cancer cells expressing TfR or CMYC-overexpressing cancers, as, in some cases, CMYC-overexpression can cause TfR-overexpression. Cancers overexpressing TfR can include ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer.

[0222] In some embodiments, the presently described peptides can be peptide constructs comprising one or more TfR-binding peptides as described herein conjugated to, linked to, or fused to one or more active agents. Peptide constructs as described herein can include chemical conjugates and recombinant fusion molecules. In some cases, a chemical conjugate can comprise a TfR-binding peptide as described herein that is chemically conjugated to or linked to another molecule. Molecules can include small molecules, peptides, polypeptides, proteins, or other macromolecules (e.g., nanoparticles) and polymers (e.g., nucleic acids, polylysine, or

polyethylene glycol). In some cases, a TfR-binding peptide of the present disclosure is conjugated to another molecule via a linker. Linker moieties can include cleavable (e.g., pH sensitive or enzyme-labile linkers) or stable linkers. In some embodiments, a peptide construct is a fusion molecule (e.g., a fusion peptide or fusion protein) that can be recombinantly expressed, and wherein the fusion molecule can comprise one or more TfR-binding peptides fused to one or more other molecules peptides, polypeptides, proteins, or other macromolecules that can be recombinantly expressed.

[0223] In some cases, the TfR-binding peptides of the present disclosure are conjugated to, linked to, or fused to a therapeutic (also referred to as an “active agent”) cargo molecule capable of exerting a certain biological effect or diagnostic (also referred to as a “detectable agent”) cargo molecule, resulting in peptide constructs. In some cases, the therapeutic or diagnostic cargo molecules used herein can be transported across the BBB via TfR-mediated transcytosis. Once inside the CNS, the cargo molecules can target a specific cell, cell population, or tissue. In some cases, the therapeutic or diagnostic cargo molecules are known compounds that, when used in combination with the compositions and methods disclosed herein, are able to exert a significantly higher potency inside the CNS due to effective transport across the BBB.

[0224] The peptides of the present disclosure, or derivatives, fragments, or variants thereof, can have an affinity and selectively for TfR, or a derivative or analog thereof. In some cases, the peptides of the present disclosure can be engineered using site-saturation mutagenesis (SSM) to exhibit improved TfR-binding properties or promote transcytosis more effectively. In some cases, the peptides of the present disclosure are cystine-dense peptides (CDPs), related to knotted peptides or hitchin-derived peptides or knottin-derived peptides. The TfR-binding peptides can be cystine-dense peptides (CDPs). The terms “peptides”, “CDPs”, “TfR-binding peptides,” “TfR-binding CDPs,” and “TfR-binding peptides” are used interchangeably herein. Hitchins can be a subclass of CDPs wherein six cysteine residues form disulfide bonds according to the connectivity [1-4], 2-5, 3-6 indicating that the first cysteine residue forms a disulfide bond with the fourth residue, the second with the fifth, and the third cysteine residue with the sixth. The brackets in this nomenclature indicate cysteine residues form the knotting disulfide bond. (See e.g., Correnti et al. Screening, large-scale production, and structure-based classification for cystine-dense peptides. *Nat Struct Mol Biol.* 2018 Mar; 25(3): 270–278). Knottins can be a subclass of CDPs wherein six cysteine residues form disulfide bonds according to the connectivity 1-4, 2-5, [3-6]. Knottins are a class of peptides, usually ranging from about 20 to

about 80 amino acids in length that are often folded into a compact structure. Knottins are typically assembled into a complex tertiary structure that is characterized by a number of intramolecular disulfide crosslinks and may contain beta strands and other secondary structures. The presence of the disulfide bonds gives knottins remarkable environmental stability, allowing them to withstand extremes of temperature and pH and to resist the proteolytic enzymes of the blood stream. In some cases, the peptides described herein can be derived from knotted peptides. The amino acid sequences of peptides as disclosed herein can comprise a plurality of cysteine residues. In some cases, at least cysteine residues of the plurality of cysteine residues present within the amino acid sequence of a peptide participate in the formation of disulfide bonds. In some cases, all cysteine residues of the plurality of cysteine residues present within the amino acid sequence of a peptide participate in the formation of disulfide bonds. As described herein, the term “knotted peptide” can be used interchangeably with the terms “cystine-dense peptide”, “CDP”, or “peptide”.

[0225] Provide herein are methods of identification, maturation, characterization, and utilization of CDPs that bind the transferrin receptor and allow accumulation of bioactive molecules at therapeutically relevant concentrations in a subject (e.g., a human or non-human animal). This disclosure demonstrates the utility of CDPs as a diverse scaffold family that can be screened for applicability to modern drug discovery strategies. CDPs comprise alternatives to existing biologics, primarily antibodies, which may bypass some of the liabilities of the immunoglobulin scaffold, including poor tissue permeability, immunogenicity, and long serum half-life that can become problematic if toxicities arise. Peptides of the present disclosure in the 20-80 amino acid range represent medically relevant therapeutics that are mid-sized, with many of the favorable binding specificity and affinity characteristics of antibodies but with improved stability, reduced immunogenicity, and simpler manufacturing methods. The intramolecular disulfide architecture of CDPs provides particularly high stability metrics, reducing fragmentation and immunogenicity, while their smaller size could improve tissue penetration or cell penetration and facilitate tunable serum half-life. Disclosed herein are peptides representing candidate peptides that can serve as CNS drug delivery vehicles. A peptide of the present disclosure may be a cell-penetrating peptide (CPP). A CPP may be cell-penetrating, tissue-penetrating, or both.

[0226] In some embodiments, a CPP may be a CDP. CPPs may comprise peptides that facilitate cellular intake or uptake of various moieties and agents, and themselves may translocate across a cell membrane. CPPs may contain protein transduction domains, which are a class of short

peptide sequences which can translocate across the cell membrane. A cell-penetrating peptide may directly or indirectly enter the cytosol of a cell, the nucleus of a cell, or other subcellular locations of a cell. A cell-penetrating peptide can be used as an appropriate carrier for various cargos including nucleic acid, peptides, proteins, siRNA, dsRNA, nucleic acids with alternative backbone chemistries (e.g., linked nucleic acids (LNAs) or peptide nucleic acids (PNAs)), radionuclides, imaging agents, fluorescent agents, therapeutic agents, nanoparticles, and the like. CPPs can be an artificial or engineered sequence (e.g., a synthetic sequence). CPPs may comprise multiple protein sequences, whether such protein sequences are native, synthetic, or a variant (e.g., chimeric). CPPs can be derived from a single protein sequence, whether such protein sequences are native synthetic or variant (e.g., a protein-derived sequence). CPPs can exhibit a variety of physiochemical properties such as cationic, amphipathic or hydrophobic. Some mechanisms for internalization of CPPs include direct cell penetration, use of the endocytosis pathway, and translocation through the formation of a transitory structure. It is understood that the description and use of a cell-penetrating peptide (CPP) herein is non-limiting. Non-limiting examples of CPPs can be found, for example, in Derakhshankhah and Jafari, “Cell penetrating peptides: A concise review with emphasis on biomedical applications” (Biomedicine & Pharmacotherapy; Volume 108, December 2018, Pages 1090-1096), which is incorporated by reference in its entirety.

[0227] Some therapeutic molecules that can be used in combination with the herein disclosed methods and compositions can be able to target one or more specific cells or tissues that are part of the CNS. For example, the neurotensin receptor (NTSR), a G-protein coupled receptor, is targeted with a fusion protein comprising a CDP linked to neurotensin (abbreviated herein as “NT”; ELYENKPRRPYIL (SEQ ID NO: 350)), or a derivative thereof, a neuropeptide that activates the NTSR. In some cases, a CDP-NT peptide construct is used to prevent or treat chronic pain or neuropathic pain in a subject (e.g., a human).

[0228] Also described herein are peptides that selectively home, target, are directed to, migrate to, are able to reach, are retained by, or accumulate in and/or bind to specific regions, tissues, structures or cells of the central nervous system (CNS) that are involved in sensing, modulating, managing, decreasing, ablating or reducing pain, including nociceptive pain, or other therapeutic indications as described herein. A peptide that homes, targets, migrates to, is directed to, is retained by, or accumulates in and/or binds to one or more specific regions, tissues, structures or cells of the affected region can have fewer off-target and potentially

negative effects, for example, side effects that often limit use and efficacy of pain drugs. In addition, such peptides can deliver active agents to regions, such as the CNS, where those active agents are otherwise unable to reach the CNS at therapeutic levels, such as due to the blood-brain barrier. Such peptides can also reduce need for other pain medication, including opioid medications. In addition, such peptides can reduce dosage and increase the efficacy of existing drugs by directly targeting them to a specific region, tissue, structure or cell of the affected region and helping to contact the affected region or increasing the local concentration of agent. The peptide itself can modulate pain or it can be conjugated to an agent that modulates pain. Such pain modulation may operate by various mechanisms such as modulating inflammation, autoimmune responses, direct or indirect action on pain receptors, cell killing, or programmed cell death (whether via an apoptotic and/or non-apoptotic pathway of diseased cells or tissues, and the like (Tait et al. J Cell Sci 127(Pt 10):2135-44 (2014))).

[0229] CDPs may be advantageous for delivery to the CNS, as compared to other molecules such as antibodies due to smaller size, greater tissue or cell penetration, and quicker clearance from serum, and as compared to smaller peptides due to resistance to proteases (both for stability and for immunogenicity reduction). In some embodiments, the TfR-binding peptides of the present disclosure (e.g., CDPs, knotted peptides, or hitchhins), TfR-binding peptide conjugates (e.g., comprising one or more TfR-binding peptides and one or more active agents), or engineered TfR-binding fusion peptides (e.g., comprising one or more TfR-binding peptides and one or more peptides) may have properties that are superior to TfR-binding antibodies. For example, the peptides and constructs described herein can provide superior, deeper, and/or faster tissue or cell penetration to cells and targeted tissues (e.g., brain parenchyma penetration, solid tumor penetration) and faster clearance from non-targeted tissues and serum. The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may have lower molecular weights than TfR-binding antibodies. The lower molecular weight may confer advantageous properties on the TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure as compared to TfR-binding antibodies. For example, the TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may penetrate a cell or tissue more readily than an anti-TfR antibody or may have lower molar dose toxicity than an anti-TfR antibody. The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may be advantageous for lacking the Fc function of an antibody. The TfR-binding peptides, TfR-binding

peptide conjugates, or TfR-binding fusion peptides of this disclosure may be advantageous for allowing higher concentrations, on a molar basis, of formulations.

[0230] The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may have a wider therapeutic window (e.g., the dosage above which a therapeutic pharmacodynamic response is observed but below which toxicity is observed) as compared to TfR-binding antibody-based therapeutics. The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may be used at higher molar dosage with less risk of toxicity as compared to TfR-binding antibody-based therapeutics. The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may have fewer epitopes to trigger an adaptive immune response, resulting in reduced immunogenicity as compared to TfR-binding antibody-based therapeutics. The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may exhibit more facile and less disruptive incorporation of active agents into protein fusion constructs as compared to TfR-binding antibody-based therapeutics. The TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may have a smaller surface area, resulting in lower risk for off-target binding, as compared to TfR-binding antibody-based therapeutics. An exemplary comparison is described in **EXAMPLE 54**.

[0231] In some embodiments, the TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure exhibit lower on-target toxicity than an anti-TfR antibody when administered to a subject at the same molar dose or at a similarly effective dose. In some embodiments, the TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides exhibit lower off-target toxicity than an antibody when administered to a subject at the same molar dose or a similarly effective dose. For example, the TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure may be administered to a subject at about 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 15-fold, 20-fold, 25-fold, 30-fold, 40-fold, 50-fold, 60-fold, 70-fold, 80-fold, 90-fold, 100-fold higher molar dose than an antibody while providing similar or lower observed toxicity. In some embodiments, the TfR-binding peptides, TfR-binding peptide conjugates, or TfR-binding fusion peptides of this disclosure exhibit higher efficacy than an anti-TfR antibody when administered to a subject at the same dose by weight as the anti-TfR antibody. The TfR-binding peptides of the present disclosure, when fused to a half-life extending moiety (e.g., Fc, SA21, PEG), can be delivered at

even lower doses while preserving activity and efficacy and, thus, is far superior to administering an anti-TfR antibody.

[0232] The presently disclosed TfR-binding peptides when fused to an active agent in a peptide construct can exhibit higher efficacy than an engineered anti-TfR bispecific antibody against the same target. For example, the engineered TfR-binding peptides disclosed herein can be further fused to a BACE inhibitor. These engineered TfR-binding peptide-BACE inhibitor constructs can be administered at a higher molar dose than a bispecific anti-TfR/BACE1 antibody while still maintaining the same or higher efficacy. For example, the engineered TfR-binding peptide-BACE inhibitor constructs can cause less reticulocytopenia and effectively inhibit amyloid beta formation, with low or no toxicity to the subject to which the engineered TfR-binding peptide-BACE inhibitor constructs is administered. In contrast, the bispecific anti-TfR/BACE1 antibody may cause more reticulocytopenia and/or have lower efficacy in preventing amyloid beta formation and can have higher toxicity to subjects, resulting in lethargy, distress, and signs of hemolysis.

[0233] In some embodiments, the present disclosure provides peptides (e.g., CDPs, knotted peptides, or hitchhins), chemical conjugates (e.g., comprising one or more TfR-binding peptides and one or more active agents), or recombinantly expressed fusion molecules (e.g., comprising one or more TfR-binding peptides and one or more active agents) that bind to TfR. The TfR-binding peptides can be cystine-dense peptides (CDPs). The terms “peptides”, “CDPs”, “TfR-binding peptides,” “TfR-binding CDPs,” “TfR-binding peptides,” and “engineered TfR-binding peptides” are used interchangeably herein. The binding of peptides described in the present disclosure to TfR can facilitate transcytosis of the peptide or peptide construct (e.g., fusion protein, or peptide conjugated to, linked to, or fused to an agent) across a cell barrier (e.g., the BBB). Also disclosed herein is the use of a mammalian surface display screening platform to screen a diverse library of CDPs and identify CDPs that specifically bind to human TfR. Further affinity maturation can be subsequently implemented to produce an allelic series of TfR-binding CDPs with varying affinities. In some embodiments, TfR-binding CDPs are identified and binding can be determined by crystallography. Peptides of the present disclosure can have cross-reactivity across species. For example, the peptides disclosed herein, in some cases, bind to human and murine TfR. Peptides disclosed herein can accumulate in the CNS and can penetrate the BBB via engagement of the TfR, following intravenous administration, including induction of a cAMP response element signaling cascade using neurotensin peptide constructs. Disclosed

herein are TfR-binding CDPs for use as therapeutic delivery agents in oncology, autoimmune disease, acute and chronic neurodegeneration, and pain management. Delivery of active or pharmaceutical agents via TfR-binding CDP can be advantageous over conventional anti-TfR antibodies due to simpler manufacturing (peptides can be made via biologic or synthetic means), improved stability, and smaller size (less potential for steric hindrance of cargo activity). Thus, the methods and compositions of the present disclosure can provide a solution to the problem of effectively transporting cargo molecules (e.g., therapeutic and/or diagnostic small molecules, peptides or proteins) into the CNS (e.g., the brain). For example, the peptides of the present disclosure aid in drug delivery to tumors located in the brain.

[0234] In some embodiments of the present disclosure, a diverse library of CDPs, knotted peptides, hitchins, or peptides derived from knotted peptides or hitchins can be used in combination with a mammalian surface display screening platform is used to identify peptides that specifically bind to human TfR. (See e.g., Crook et al. (2017) Mammalian display screening of diverse cysteine-dense peptides for difficult to drug targets. Nat Commun 8:2244). In some embodiments, a diverse library of CDPs, knotted peptides, hitchins, or peptides derived from knotted peptides or hitchins is mutagenized from endogenous peptide sequences to provide novel peptide sequences. Once TfR-binding peptides have been identified, affinity maturation (e.g., site-saturation mutagenesis) can be performed to produce an allelic series of binders with varying (e.g., improved) affinities for TfR. These techniques can be used in combination with various other analytical methods (e.g., crystallography or spectroscopy) in order to determine the nature of peptide-receptor interaction (e.g., critical amino acid residues for receptor binding etc.). In some cases, the peptides of the present disclosure are developed to bind human TfR; however, those peptides can show cross-reactivity with TfR derived from other species such as murine TfR.

[0235] The peptides of the present disclosure can have varying biodistribution patterns *in vivo*. In some cases, mouse biodistribution and pharmacokinetic studies are performed using, for example, radiolabeled peptides (e.g., ¹⁴C-labeled peptides), in order to determine organ distribution, the uptake and residence times of the peptides in target organs (e.g., brain), and their mode of clearance (e.g., renal or hepatic).

[0236] In various embodiments, the present disclosure provides peptides that accumulate in the CNS (e.g., the brain). In some cases, CNS accumulation of those peptides is due to penetration across the BBB. In some cases, engineered TfR-binding peptide variants as described herein

cross the BBB by TfR-mediated transcytosis. In some cases, the TfR-binding peptides of the present disclosure accumulate in tumor tissue that is located in the CNS. In some cases, the tumor that the peptide accumulates in is a brain tumor (e.g., glioma). As described herein, the terms “accumulates” and “accumulates in” can be used interchangeably and be used to refer to “accumulation in or accumulation on a cell and/or a tissue and can be understood as a gradual increase of concentration of the respective molecule (e.g., a TfR-binding peptide or peptide construct) over time.

[0237] In some embodiments, the engineered peptides of the present disclosure (e.g., histidine-containing or histidine-enriched TfR-binding peptides) can have a high TfR binding affinity at physiological pH but a significantly reduced binding affinity at lower pH levels such as endosomal pH of 5.4. In some cases, the TfR-binding peptides of the present disclosure can be optimized for improved intra-vesicular (e.g., intra-endosomal) and/or intracellular delivery function while retaining high TfR binding capabilities. In some cases, histidine scans and comparative binding experiments can be performed to develop and screen for such peptides. In addition, some peptides can comprise (e.g., conjugated to, linked to, or fused to) a motif that facilitates low-pH endosomal escape of the peptide or peptide construct for enhanced delivery functions (e.g., intracellular delivery of a therapeutic agent). In some embodiments, an amino acid residue in a peptide of the present disclosure is substituted with a different amino acid residue to alter a pH-dependent binding affinity to TfR. The amino acid substitution may increase a binding affinity at low pH, increase a binding affinity at high pH, decrease a binding affinity at low pH, decrease a binding affinity at high pH, or a combination thereof.

[0238] Exemplary peptides of the present disclosure are shown in **TABLE 1** with amino acid sequences set forth in SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, and SEQ ID NO: 171 – SEQ ID NO: 202.

[0239] In various embodiments, the peptides of the present disclosure are part of a peptide construct, such as a peptide construct or a fusion peptide (e.g., those having an amino acid sequence of any one of SEQ ID NO 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400, and SEQ ID NO: 401), designed to transport therapeutic and/or diagnostic molecules (e.g., small molecules, peptides, or

proteins) across cell layers or cell barriers such as the BBB. Exemplary peptide constructs are shown in **TABLE 2**, **TABLE 5**, and **TABLE 6**. A peptide construct of the present disclosure can comprise a TfR targeting peptide linked (e.g., chemically conjugated or fused) to a therapeutic and/or diagnostic molecule or compound. The TfR-binding peptide construct can cross the BBB via TfR-mediated transcytosis and hence deliver the therapeutic and/or diagnostic molecule or compound into the CNS. The methods and compositions of the present disclosure therefore can have a profound impact on the diagnosis, prevention, and treatment of various diseases or conditions. Disease areas for which the methods and compositions of the present disclosure can be used include, but are not limited to, oncology, autoimmune diseases, acute and chronic neurodegeneration, muscle disorders, neuropsychological disorders, epilepsy, schizophrenia, depression, anxiety, bipolar disorder, developmental brain disorders (e.g., autism spectrum), pain, epilepsy, mood disorder, and pain management.

[0240] Hence, a subject having a disorder of the CNS is otherwise unable to benefit from a drug or drug class whose limited by poor BBB penetration, can benefit from the compositions and methods described herein because therapeutically effective concentrations of the drug in the CNS can be achieved by conjugation to a TfR-binding peptide provided herein. In addition to pharmacological advances, the compositions and methods disclosed herein comprising TfR-binding peptides can be superior over conventional CNS- and TfR-targeting agents (e.g., anti-TfR antibodies) in terms of their production, quality control, and safety. For example, the peptide and peptide constructs of the present disclosure are manufactured in a simpler, less expensive, and more resource-effective manner (e.g., peptides can be synthesized via biologic or synthetic approaches); the peptide and peptide constructs of the present disclosure can show improved ex vivo and in vivo stability, are smaller in size (e.g., less potential for steric hindrance of cargo activity and the ability to penetrate dense tissue such as solid tumors as well as the potential for faster clearance from systemic circulation), exhibit a higher tissue or cell penetration, and have lower immunogenicity. Peptides of this disclosure can be engineered to have lower immunogenicity by combining crystallographic data with major histocompatibility complex (MHC) or human leukocyte antigen (HLA) peptide fragment binding experiments or computational predictions thereof, or a combination of the foregoing.

[0241] The identification of therapeutic or diagnostic agents that have the ability to cross cellular layers or barriers such as the BBB have been challenging, as such methods involve demand for high specificity or targeting, high affinity binders for receptors that can promote transcytosis

(e.g., TfR), and the ability to target cells and act on target proteins (low off-target adverse effects) after transcytosis. With few exceptions, high throughput screening campaigns with small molecule libraries failed to provide specific compounds capable of crossing endothelial or epithelial layers and/or transporting cargo across those layers.

[0242] Described herein are, in some embodiments, peptides and peptide constructs and methods of screening for peptides and peptide constructs that target a protein of interest, such as TfR. Compared to wildtype or endogenous molecules such as transferrin, the methods and compositions as described herein can provide peptides with improved TfR-binding capabilities, or peptides that exhibit improved transport capabilities across the BBB, or any combination thereof. In some cases, the presently described peptides efficiently transport cargo molecules (e.g., therapeutic or diagnostic small molecules or proteins) across endothelial cell layers (e.g., the BBB) or epithelial layers. In some embodiments, the TfR-binding peptides of the present disclosure bind to a TfR and promote vesicular transcytosis. In some cases, the TfR-binding peptides of the present disclosure bind to a cell that overexpress a TfR (e.g., a cancer cell) and promotes uptake of the peptide by the cell. In some aspects, a TfR binding peptide or peptide constructs as described herein promotes vesicular transcytosis and uptake by a TfR-overexpressing cell such as a cancer, or a combination thereof.

[0243] The TfR-binding peptides of the present disclosure can bind TfR of different species including human, monkey, mouse, and rat TfR. In some cases, variations or mutations in any of the amino acid residues of a TfR-binding peptide may influence cross-reactivity. In some cases, variations or mutations in any of the amino acid residues of a TfR-binding peptide that interact with the binding site of TfR may influence cross-reactivity.

[0244] Described herein are peptides, including, but not limited to, designed or engineered peptides, recombinant peptides, and cystine-dense peptides (CDPs)/small disulfide-knotted peptides (e.g., knotted peptides, hitchins, and peptides derived therefrom), that can be large enough to carry a cargo molecule while retaining the ability to bind a target protein with high affinity (e.g., TfR), but yet small enough to access cellular compartments, such as the cytosol or the nucleus, or tissues, such as the center of cell agglomerates (e.g., solid tumors). In some cases, the peptides as described herein carry cargo molecules across the BBB into the CNS (e.g., the parenchyma) via vascular transcytosis. In some cases, the transcytosis is TfR-mediated.

[0245] Further described herein are methods and compositions for determining the nature of peptide-receptor interactions (e.g., using X-ray crystallography) as well as their

pharmacodynamic and pharmacokinetic properties in vivo, including accumulation in the CNS (e.g., brain). Some of the peptides described herein have the ability to target and accumulate in tumor cells. In some cases, the tumor cells overexpress TfR. In some aspects, the peptides of the present disclosure have high in vivo stabilities, e.g., high protease stability, high tolerability of reducing agents such as glutathione (GSH), and tolerate elevated temperatures (e.g., up to 95 °C).

[0246] The present disclosure provides, in some embodiments, a peptide or protein design approach based on the 3D protein or receptor structure for identifying peptides or proteins capable of binding such receptor. In some cases, the receptor is a transferrin receptor.

[0247] Additional aspects and advantages of the present disclosure will become apparent to those skilled in this art from the following detailed description, wherein illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various respects, all without departing from the disclosure. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

[0248] As used herein, the abbreviations for the natural L-enantiomeric amino acids are conventional and are as follows: alanine (A, Ala); arginine (R, Arg); asparagine (N, Asn); aspartic acid (D, Asp); cysteine (C, Cys); glutamic acid (E, Glu); glutamine (Q, Gln); glycine (G, Gly); histidine (H, His); isoleucine (I, Ile); leucine (L, Leu); lysine (K, Lys); methionine (M, Met); phenylalanine (F, Phe); proline (P, Pro); serine (S, Ser); threonine (T, Thr); tryptophan (W, Trp); tyrosine (Y, Tyr); valine (V, Val). Typically, Xaa can indicate any amino acid. In some embodiments, X can be asparagine (N), glutamine (Q), histidine (H), lysine (K), or arginine (R).

[0249] Some embodiments of the disclosure contemplate D-amino acid residues of any standard or non-standard amino acid or analogue thereof. When an amino acid sequence is represented as a series of three-letter or one-letter amino acid abbreviations, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy terminal direction, in accordance with standard usage and convention.

[0250] The terms “peptide”, “polypeptide”, “protein”, “hitchin”, “cystine-dense peptide”, “knotted peptides” or “CDP” can be used interchangeably herein to refer to a polymer of amino acid residues. In various embodiments, “peptides”, “polypeptides”, and “proteins” can be chains of amino acids whose alpha carbons are linked through peptide bonds. The terminal amino acid at one end of the chain (e.g., amino terminal) therefore can have a free amino group, while the

terminal amino acid at the other end of the chain (e.g., carboxy terminal) can have a free carboxyl group. As used herein, the term “amino terminus” (e.g., abbreviated N-terminus) can refer to the free α -amino group on an amino acid at the amino terminal of a peptide or to the α -amino group (e.g., imino group when participating in a peptide bond) of an amino acid at any other location within the peptide. Similarly, the term “carboxy terminus” can refer to the free carboxyl group on the carboxy terminus of a peptide or the carboxyl group of an amino acid at any other location within the peptide. Peptides also include essentially any polyamino acid including, but not limited to, peptide mimetics such as amino acids joined by an ether or thioether as opposed to an amide bond.

[0251] As used herein, the term “peptide construct” can refer to a molecule comprising one or more peptides of the present disclosure that can be conjugated to, linked to, or fused to one or more cargo molecules. In some cases, cargo molecules are active agents. The term “active agent” can refer to any molecule, e.g., any molecule that is capable of eliciting a biological effect and/or a physical effect (e.g., emission of radiation) which can allow the localization, detection, or visualization of the respective peptide construct. In various embodiments, the term “active agent” refers to a therapeutic and/or diagnostic agent. A peptide construct of the present disclosure can comprise a TfR-binding peptide that is linked to one or more active agents via one or more linker moieties (e.g., cleavable or stable linker) as described herein.

[0252] As used herein, the terms “comprising” and “having” can be used interchangeably. For example, the terms “a peptide comprising an amino acid sequence of SEQ ID NO: 32” and “a peptide having an amino acid sequence of SEQ ID NO: 32” can be used interchangeably.

[0253] As used herein, and unless otherwise stated, the term “TfR” or “transferrin receptor” is a class of protein used herein and can refer to a transferrin receptor from any species (e.g., human or murine TfR or any human or non-human animal TfR). In some cases, and as used herein, the term “TfR” or “transferrin receptor” refers to human TfR (hTfR) and can include TfR or any of the known TfR homologs or orthologs, including TfR1, TfR2, soluble TfR, or any combination or fragment (e.g., ectodomain) thereof.

[0254] The term “engineered,” when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such engineered molecules are those that are separated from their natural

environment and include cDNA and genomic clones (i.e., a prokaryotic or eukaryotic cell with a vector containing a fragment of DNA from a different organism). Engineered DNA molecules of the present invention are free of other genes with which they are ordinarily associated but may include naturally occurring or non-naturally occurring 5' and 3' untranslated regions such as enhancers, promoters and terminators.

[0255] An “engineered” polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. In a preferred form, the engineered polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, e.g., greater than 95% pure, more preferably greater than 98% pure or greater than 99% pure. When used in this context, the term “engineered” does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers, heterodimers and multimers, or alternatively glycosylated, carboxylated, modified, or derivatized forms.

[0256] An “engineered” peptide or protein is a polypeptide that is distinct from a naturally occurring polypeptide structure, sequence, or composition. Engineered peptides include non-naturally occurring, artificial, isolated, synthetic, designed, modified, or recombinantly expressed peptides. Provided herein are engineered TfR-binding peptides, variants, or fragments thereof. These engineered TfR-binding peptides can be further linked to an active agent or a detectable agent. The active agent can be a half-life extending moiety.

[0257] Polypeptides of the disclosure include polypeptides that have been modified in any way, for example, to: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (5) confer or modify other physicochemical or functional properties. For example, single or multiple amino acid substitutions (e.g., conservative amino acid substitutions) are made in the naturally occurring sequence (e.g., in the portion of the polypeptide outside the domain(s) forming intermolecular contacts). A “conservative amino acid substitution” can refer to the substitution in a polypeptide of an amino acid with a functionally similar amino acid. The following six groups each contain amino acids that can be conservative substitutions for one another: i) Alanine (A), Serine (S), and Threonine (T); ii) Aspartic acid (D) and Glutamic acid (E); iii) Asparagine (N) and Glutamine (Q); iv) Arginine (R) and Lysine (K); v) Isoleucine (I), Leucine (L), Methionine (M), and Valine (V); vi) Phenylalanine (F), Tyrosine (Y), and Tryptophan (W).

[0258] The terms “polypeptide fragment” and “truncated polypeptide” as used herein can refer to

a polypeptide that has an amino-terminal and/or carboxy-terminal deletion as compared to a corresponding full-length peptide or protein. In various embodiments, fragments are at least 5, at least 10, at least 25, at least 50, at least 100, at least 150, at least 200, at least 250, at least 300, at least 350, at least 400, at least 450, at least 500, at least 600, at least 700, at least 800, at least 900 or at least 1000 amino acids in length. In various embodiments, fragments can also be, *e.g.*, at most 1000, at most 900, at most 800, at most 700, at most 600, at most 500, at most 450, at most 400, at most 350, at most 300, at most 250, at most 200, at most 150, at most 100, at most 50, at most 25, at most 10, or at most 5 amino acids in length. A fragment can further comprise, at either or both of its ends, one or more additional amino acids, for example, a sequence of amino acids from a different naturally-occurring protein (*e.g.*, an Fc or leucine zipper domain) or an artificial amino acid sequence (*e.g.*, an artificial linker sequence).

[0259] In various embodiments, the interface residues of the TfR-binding peptides of the present disclosure (*e.g.*, those amino acid residues that interact with TfR for receptor binding) can be divided between two largely helical domains of the peptide. In some cases, the interface residues can comprise residues corresponding to residues 5-25 (*e.g.*, and comprising corresponding residues G5, A7, S8, M11, N14, L17, E18, and E21), with reference to SEQ ID NO: 32, or corresponding to residues 35-51 (*e.g.*, and comprising corresponding residues L38, L41, L42, L45, D46, H47, H49, S50, and Q51), with reference to SEQ ID NO: 32, or both. For example, the interface residues can comprise residues corresponding to residues 5-25 (*e.g.*, and comprising corresponding residues G5, A7, S8, M11, N14, L17, E18, and E21), with reference to SEQ ID NO: 32, or corresponding to residues 35-51 (*e.g.*, and comprising corresponding residues L38, L41, L42, L45, D46, H47, H49, S50, and Q51), with reference to SEQ ID NO: 32. In some embodiments, a TfR-binding peptide can comprise a fragment of a peptide provided herein, wherein the fragment comprises the minimum interface residues for binding, for example residues corresponding to residues 5-25 (*e.g.*, and comprising corresponding residues G5, A7, S8, M11, N14, L17, E18, and E21), with reference to SEQ ID NO: 32, or corresponding to residues 35-51 (*e.g.*, and comprising corresponding residues L38, L41, L42, L45, D46, H47, H49, S50, and Q51), with reference to SEQ ID NO: 32. In some cases, the TfR-binding peptide is a peptide having the sequence set forth in SEQ ID NO: 32 comprising the TfR-binding residues corresponding to residues G5, A7, S8, M11, N14, L17, E18, and E21 of the domain and corresponding to residues L38, L41, L42, L45, D46, H47, H49, S50, and Q51 of the second domain, with reference to SEQ ID NO: 32.

[0260] As used herein, the terms “peptide” or “polypeptide” in conjunction with “variant” “mutant” or “enriched mutant” or “permuted enriched mutant” can refer to a peptide or polypeptide that can comprise an amino acid sequence wherein one or more amino acid residues are inserted into, deleted from and/or substituted into the amino acid sequence relative to another polypeptide sequence. In various embodiments, the number of amino acid residues to be inserted, deleted, or substituted is at least 1, at least 2, at least 3, at least 4, at least 5, at least 10, at least 25, at least 50, at least 75, at least 100, at least 125, at least 150, at least 175, at least 200, at least 225, at least 250, at least 275, at least 300, at least 350, at least 400, at least 450 or at least 500 amino acids in length. Variants of the present disclosure include peptide conjugates or fusion molecules (e.g., peptide constructs).

[0261] A “derivative” of a peptide or polypeptide can be a peptide or polypeptide that can have been chemically modified, e.g., conjugation to another chemical moiety such as, for example, polyethylene glycol, albumin (e.g., human serum albumin), phosphorylation, and glycosylation.

[0262] The term “% sequence identity” can be used interchangeably herein with the term “% identity” and can refer to the level of amino acid sequence identity between two or more peptide sequences or the level of nucleotide sequence identity between two or more nucleotide sequences, when aligned using a sequence alignment program. For example, as used herein, 80% identity means the same thing as 80% sequence identity determined by a defined algorithm, and means that a given sequence is at least 80% identical to another length of another sequence. In various embodiments, the % identity is selected from, e.g., at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% or more sequence identity to a given sequence. In various embodiments, the % identity is in the range of, e.g., about 60% to about 70%, about 70% to about 80%, about 80% to about 85%, about 85% to about 90%, about 90% to about 95%, or about 95% to about 99%.

[0263] The terms “% sequence homology” or “percent sequence homology” or “percent sequence identity” can be used interchangeably herein with the terms “% homology,” “% sequence identity,” or “% identity” and can refer to the level of amino acid sequence homology between two or more peptide sequences or the level of nucleotide sequence homology between two or more nucleotide sequences, when aligned using a sequence alignment program. For example, as used herein, 80% homology means the same thing as 80% sequence homology determined by a defined algorithm, and accordingly a homologue of a given sequence has greater than 80% sequence homology over a length of the given sequence. In various embodiments, the

% homology is selected from, *e.g.*, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% or more sequence homology to a given sequence. In various embodiments, the % homology is in the range of, *e.g.*, about 60% to about 70%, about 70% to about 80%, about 80% to about 85%, about 85% to about 90%, about 90% to about 95%, or about 95% to about 99%.

[0264] A protein or polypeptide can be “substantially pure,” “substantially homogeneous”, or “substantially purified” when at least about 60% to 75% of a sample exhibits a single species of polypeptide. The polypeptide or protein can be monomeric or multimeric. A substantially pure polypeptide or protein can typically comprise about 50%, 60%, 70%, 80% or 90% W/W of a protein sample, more usually about 95%, and *e.g.*, will be over 99% pure. Protein purity or homogeneity can be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band upon staining the gel with a stain well known in the art. For certain purposes, higher resolution is provided by using high-pressure liquid chromatography (*e.g.*, HPLC) or other high-resolution analytical techniques (*e.g.*, LC-mass spectrometry).

[0265] As used herein, the term “pharmaceutical composition” can generally refer to a composition suitable for pharmaceutical use in a subject such as an animal (*e.g.*, human or mouse). A pharmaceutical composition can comprise a pharmacologically effective amount of an active agent and a pharmaceutically acceptable carrier. The term “pharmacologically effective amount” can refer to that amount of an agent effective to produce the intended biological or pharmacological result.

[0266] As used herein, the term “pharmaceutically acceptable carrier” can refer to any of the standard pharmaceutical carriers, vehicles, buffers, and excipients, such as a phosphate buffered saline solution, 5% aqueous solution of dextrose, and emulsions, such as an oil/water or water/oil emulsion, and various types of wetting agents and/or adjuvants. Suitable pharmaceutical carriers and formulations are described in Remington's Pharmaceutical Sciences, 21st Ed. 2005, Mack Publishing Co, Easton. A “pharmaceutically acceptable salt” can be a salt that can be formulated into a compound for pharmaceutical use including, *e.g.*, metal salts (sodium, potassium, magnesium, calcium, etc.) and salts of ammonia or organic amines.

[0267] As used herein, the terms “treat”, “treating” and “treatment” can refer to a method of alleviating or abrogating a biological disorder and/or at least one of its attendant symptoms. As used herein, to “alleviate” a disease, disorder or condition, for example, means reducing the

severity and/or occurrence frequency of the symptoms of the disease, disorder, or condition. Further, references herein to “treatment” can include references to curative, palliative, and prophylactic or diagnostic treatment.

[0268] Generally, a cell of the present disclosure can be a eukaryotic cell or a prokaryotic cell. A cell can be an epithelial cell. A cell can be an animal cell or a plant cell. An animal cell can include a cell from a marine invertebrate, fish, insects, amphibian, reptile, or mammal. A mammalian cell can be obtained from a primate, ape, equine, bovine, porcine, canine, feline, or rodent. A mammal can be a primate, ape, dog, cat, rabbit, ferret, or the like. A rodent can be a mouse, rat, hamster, gerbil, hamster, chinchilla, or guinea pig. A bird cell can be from a canary, parakeet or parrots. A reptile cell can be from a turtles, lizard or snake. A fish cell can be from a tropical fish. For example, the fish cell can be from a zebrafish (e.g., *Danio rerio*). A worm cell can be from a nematode (e.g., *C. elegans*). An amphibian cell can be from a frog. An arthropod cell can be from a tarantula or hermit crab.

[0269] A mammalian cell can also include cells obtained from a primate (e.g., a human or a non-human primate). A mammalian cell can include a blood cell, a stem cell, an epithelial cell, connective tissue cell, hormone secreting cell, a nerve cell, a skeletal muscle cell, or an immune system cell. In preferred embodiments, the methods and compositions of the present disclosure are used in combination with one or more mammalian blood cells.

[0270] As used herein, the term “vector,” generally refers to a DNA molecule capable of replication in a host cell and/or to which another DNA segment can be operatively linked so as to bring about replication of the attached segment. A plasmid is an exemplary vector.

[0271] As used herein, the term “subject,” generally refers to a human or to another animal. A subject can be of any age, for example, a subject can be an infant, a toddler, a child, a pre-adolescent, an adolescent, an adult, or an elderly individual.

[0272] Ranges can be expressed herein as from “about” one particular value, and/or to “about” another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent “about,” it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are in relation to the other endpoint, and independently of the other endpoint. The term “about” as used herein refers to a range that is 15% plus or minus from a stated numerical value within the context of the particular usage. For example, about 10 can include a range from 8.5 to 11.5.

Peptides

[0273] Disclosed herein are peptide sequences, such as those listed in **TABLE 1**, capable of binding to TfR or any of the known TfR homologs, including TfR1, TfR2, soluble TfR, or any combination or fragment (e.g., ectodomain) thereof. A peptide capable of binding a transferrin receptor or a TfR homolog may be referred to herein as a transferrin receptor-binding peptide or a TfR-binding peptide. In some embodiments, peptides disclosed herein can penetrate, cross, or enter target cells in a TfR-mediated manner. These cell layers or cells can include TfR-expressing endothelial cells, epithelial cells, and TfR-expressing cells of various tissues or organs such as tumor cells, brain cells, cancerous or tumor cells, liver cells, pancreas cells, colon cells, ovarian cells, breast cells, and/or lung cells, or any combination thereof.

[0274] In various embodiments, the present disclosure provides methods and compositions that enable TfR-mediated transport across cellular layers (e.g., endothelial cells or epithelial cells) or cell membranes. In addition to the BBB, various other cells, tissues, and organs express TfR. Single cells expressing TfR can include hepatocytes, erythrocytes and erythrocyte precursors in bone marrow, immune cells, stem cells, and rapidly dividing cells. Tissues and organs expressing TfR can include the brain (e.g., cerebral cortex, hippocampus, caudate, cerebellum), endocrine tissues (e.g., thyroid, parathyroid, and adrenal glands), bone marrow and immune system (e.g., appendix, lymph node, tonsil, spleen), muscle tissues (e.g., heart, skeletal, and smooth muscle), liver, gallbladder, pancreas, gastrointestinal tract (e.g., oral mucosa, esophagus, stomach, duodenum, small intestine, colon, rectum), kidney, urinary bladder, female tissues (e.g., fallopian tube, breast, vagina, cervix, endometrium, ovary, and placenta), adipose and soft tissue, and skin. Thus, the TfR-binding peptides of the present disclosure can be used to target these cells, tissues, and organs and deliver an active agent to these cells, tissues, and organs via, for example, TfR-mediated transcytosis (e.g., across cellular barrier such as the BBB) or TfR-mediated endocytosis (e.g., across cell membranes into cells) or TfR-mediated accumulation in tissues to treat and/or prevent a disease or condition in one or more of these cells, tissues, or organs.

[0275] In various embodiments, the present disclosure provides methods and compositions that enable TfR-mediated transport and delivery to cancer cells expressing TfR. Cancers overexpressing TfR can include ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland,

kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer.

[0276] TfR-binding peptide constructs that can be used to prevent and/or treat a cancer are those comprising a TfR binding peptide and an active agent with anti-tumor activity such as a fused IL15/IL15Ra complex, IFNgamma, and anti-CD3 agents. Examples of such peptide constructs (e.g., fusion peptides) are those having an amino acid sequence set forth in any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332 or SEQ ID NO: 359 – SEQ ID NO: 361. In some embodiments, TfR-binding peptide constructs are fused to IL-15 to recruit and stimulate the immune T cells or NK cells. In some embodiments, TfR-binding peptide constructs are fused to IFNγ to activate macrophages and upregulated MHC in tumor cells. In some embodiments, TfR-binding peptide constructs are fused to CD3 to bind T cells in addition to binding TfR on cancer cells to create an immune synapse and activate T cells to target the cancer.

[0277] Generally, the TfR-binding peptides of the present disclosure can be used in combination with various classes of active agent. The TfR-binding peptides of the present disclosure can be conjugated to, linked to, or fused to one or more of those active agents. In various aspects, an active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL15/IL15Ra complex agent, an IFNgamma agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter such as neurotensin.

[0278] In some embodiments, the peptides as disclosed herein can cross cellular layers or barriers (e.g., BBB) or cell membranes via, for example, TfR-mediated vesicular transcytosis and TfR-mediated endocytosis, respectively. In addition to binding TfR and promote transcytosis and/or endocytosis, the peptides of the present disclosure can also bind to additional target proteins on cells such as cancer cells. In some cases, a peptide is a peptide or peptide construct comprising a

TfR-binding peptide conjugated to, linked to, or fused to a targeting moiety or an active agent (e.g., a therapeutic or diagnostic agent) such as a small molecule or a peptide that has an affinity for an additional target protein (e.g., receptor or enzyme). In some cases, the TfR-binding peptide is linked to a cargo molecule and enables or promotes TfR-mediated transcytosis of the cargo molecule across the BBB or TfR-mediated endocytosis into a cell. In some instances, and subsequent to transcytosis, a peptide construct comprising the TfR-binding peptide and a cargo moiety can target a specific cell or tissue in the CNS and exert a biological effect (e.g., binding a target protein) upon reaching said cell or tissue. In some cases, a peptide construct of the present disclosure exerts a biological effect that is mediated by the TfR-binding peptide, the cargo molecule or active agent, or a combination thereof. In some cases, a TfR-binding peptide construct of the present disclosure comprising one or more active agents (e.g., therapeutic agents) can transport and/or deliver the one or more active agents into cells that express TfR. In some cases, the TfR-binding peptide accumulates in tissues in the CNS. In some cases, off-target effects are reduced due to CNS-specific accumulation. In some cases, the TfR-binding peptide accumulates in tissue outside of the CNS (e.g., liver, kidney, spleen, or skin). In some cases, the cells expressing TfR are tumor cells and the TfR-binding peptide construct delivers anti-tumor agents to these tumor cells. In some cases, the anti-tumor agents alone show no or only very limited therapeutic efficacy against the tumor cells; however, when the anti-tumor agents are combined with the TfR-binding peptides of the present disclosure as, for example, a peptide construct, the therapeutic efficacy of these anti-tumor agents is significantly improved.

[0279] In some embodiments, the TfR-binding peptides of the present disclosure can induce a biologically relevant response. In some embodiments, the biologically relevant response can be induced after intravenous dose, and in some embodiments, after a single intravenous dose. In some embodiments, the TfR-binding peptides can be used in combination with various other classes of therapeutic compounds used to treat and/or prevent pain, neuropathic pain or other neurological disorders such as neurodegenerative disorders, infectious diseases, immunological disorders (e.g., autoimmune diseases). Binding of the herein described peptides and peptide constructs (e.g., peptide conjugates, fusion peptides, or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via TfR-mediated vesicular transcytosis) or a cell membrane (e.g., via TfR-mediated endocytosis) can have implications in a number of diseases, conditions, or disorders associated with chronic

pain (e.g., headaches or migraine), neuropathic pain, obesity, insulin resistance, opioid addiction, or other neurologic or psychiatric disorders in a subject (e.g., a human).

[0280] Binding of the herein described peptides and peptide constructs (e.g., peptide conjugates, fusion peptides, or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via vesicular transcytosis) or a cell membrane (e.g., via endocytosis) can have implications in a number of diseases, conditions, or disorders associated with neurodegeneration. Neurodegenerative diseases that can be treated, prevented, or diagnosed with the herein described TfR-binding peptides can include Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia.

[0281] Binding of the herein described peptides and peptide constructs (e.g., peptide conjugates, fusion peptides, or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via vesicular transcytosis) or a cell membrane (e.g., via endocytosis) can have implications in a number of autoimmune diseases to therapeutically address auto-immune or inflammatory disorders in the brain. In some cases, an autoimmune disease can be treated and/or prevented by conjugating, linking, or fusing the TfR-binding peptide to an active agent that can be an ion channel modulator such as Kv1.3 potassium channel inhibitor. Additional diseases that can be treated and/or prevented using ion channel modulator such as Kv1.3 potassium channel inhibitors can include psoriasis and other non-brain autoimmune diseases due to its effect on effector T cells, and for neuroinflammatory and neurodegenerative diseases, for example, multiple sclerosis, Alzheimer's, Parkinson's, traumatic brain injury, or radiation therapy toxicity. In some cases, Kv1.3 potassium channel inhibitors can be a small molecule (e.g., domatinostat tosylate), or a peptide (e.g., Vm24 or an ShK peptide, such as ShK-170, ShK-186, or ShK-192, or any fragments or derivatives thereof) or any combinations thereof. (See e.g., Bartok et al. An engineered scorpion toxin analogue with improved Kv1.3 selectivity displays reduced conformational flexibility, *Sci Rep.* 2015; 5: 18397).

[0282] In some embodiments, the TfR-binding peptides of the present disclosure can be used for the treatment and prevention of various neurological diseases including but not limited to epilepsy, schizophrenia, depression, anxiety, bipolar disorder, developmental brain disorders (e.g., autism spectrum), or mood disorder.

[0283] In some embodiments, the TfR-binding peptides of the present disclosure can be used for the treatment and prevention of Crohn's disease or, more generally, inflammatory bowel diseases. In some cases, the TfR-binding peptides of the present disclosure show high uptake and retention in glandular cells of the intestine, which can express high amounts of TfR.

[0284] In various embodiments, the therapeutic efficacy of these drugs can be significantly improved when used in combination with the TfR-binding peptides of the present disclosure compared to administration without conjugation to the TfR-binding peptides as described herein. In various embodiments, the efficacy can be improved due to higher delivery and achieved levels of drug in the tissue or cell of interest. In various embodiments, the efficacy can be improved by increasing the relative level of the drug in the tissue or cell of interest and reducing the level of the drug in other tissues or compartments, thereby improving the therapeutic window or reducing toxic side effects.

[0285] In other embodiments, peptides as described herein compete with endogenous molecules for binding to TfR. As described herein, "compete" or peptide competition for binding to target protein such as TfR encompasses, but is not limited to, steric hindrance, occupying binding sites of the target protein, non-covalent interactions, such as salt bridges or hydrophobic interactions, crosslinking, covalent interactions, sequestration, allosteric modulation, or any combination thereof.

[0286] In some embodiments, peptides of the present disclosure can bind to any of the known TfR homologs, including TfR1, TfR2, soluble TfR, or any combination or fragment (e.g., ectodomain) thereof. Thus, as used herein, "TfR" can refer to any known homolog, derivative, fragment, or member of the TfR family including TfR1, TfR2, and a soluble TfR. In other embodiments, peptides are capable of binding to one, one or more, or all TfR homologs. In some embodiments, peptides of the present disclosure can bind to a TfR and promote a particular biological effect such as vesicular transcytosis. In some embodiments, peptides of the present disclosure, including peptides and peptide constructs with amino acid sequences set forth in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 356 – SEQ ID NO: 361, and any derivatives or variant thereof,

prevent or decrease the binding of endogenous TfR binders (e.g., transferrin or any derivatives such as apo-transferrin or holo-transferrin) to TfR. In some embodiments, peptides or peptide constructs of the present disclosure comprise derivatives and variants with at least 70% homology, at least 75% homology, at least 80% homology, at least 85% homology, at least 90% homology, at least 95% homology, at least 96% homology, at least 97% homology, at least 98% homology, or at least 99% homology or at least 100% homology to amino acid sequences set forth in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 356 – SEQ ID NO: 361.

[0287] In some embodiments, peptides bind to TfR with equal, similar, or greater affinity (e.g., lower dissociation constant K_D) as compared to endogenous molecules (e.g., transferrin, holotransferrin (iron-bound transferrin), apotransferrin (transferrin not bound to iron), or any other endogenous TfR ligands) or other exogenous molecules. In some embodiments, the peptide can have a K_D of less than 50 μM , less than 5 μM , less than 500 nM, less than 100 nM, less than 40 nM, less than 30 nM, less than 20 nM, less than 10 nM, less than 1 nM, or less than 0.1 nM. In some embodiments, peptide transport by TfR is improved by having a lower affinity (e.g., a higher dissociation constant K_D) as compared to endogenous molecules. In some embodiments, peptide transport by TfR is improved by having a faster off rate or higher k_{off} than endogenous molecules. In some embodiments, the off rate or k_{off} is similar to that of transferrin. In some embodiments, peptide transport is improved by having a faster on rate or a higher k_{on} , optionally such as higher than that of transferrin. In other embodiments, one or more conserved residues at the transferrin (Tf)-TfR-binding interface are also present in the amino acid sequences of the peptides described herein.

[0288] In some embodiments, peptides that exhibit an improved TfR receptor binding show improved transcytosis function. In some embodiments, peptides that exhibit an improved TfR receptor binding show no or small changes in transcytosis function. In some embodiments, peptides that exhibit an improved TfR receptor binding show reduced transcytosis function. In some embodiments, the peptide binds at a site of high homology between human and murine TfR, including one or more, or all, of the amino acid domains corresponding to residues 506-510, 523-531, and 611-662 of the human TfR (SEQ ID NO: 363). In some embodiments, the regions of TfR to which the peptides disclosed herein or variants thereof bind all or in part to such TfR domains. In some embodiments, the peptides disclosed herein bind to any one, any two, or all three of the TfR regions of high homology including the amino acid domains corresponding to

residues 506-510, 523-531, and 611-662 of the human TfR (SEQ ID NO: 363). In some embodiments the peptides disclosed herein bind at least to the domain corresponding to residues 611-662 of the human TfR.

[0289] In some embodiments, the K_A and K_D values of a TfR-binding peptide can be modulated and optimized (e.g., via amino acid substitutions) to provide an optimal ratio of TfR-binding affinity and efficient transcytosis function.

[0290] In some embodiments, peptides disclosed herein or variants thereof bind to TfR at residues found in the binding interface (e.g., the binding domain or the binding pocket) of TfR with other exogenous or endogenous ligands (e.g., transferrin (Tf), Tf derivatives, or Tf-like peptides or proteins). In some embodiments, a peptide disclosed herein or a variant thereof, which binds to TfR, comprises at least 70% homology, at least 75% homology, at least 80% homology, at least 85% homology, at least 90% homology, at least 95% homology, at least 96% homology, at least 97% homology, at least 98% homology, or at least 99% homology or at least 100% homology to a sequence that binds residues of TfR, which make up the binding pocket. In some embodiments, a peptide disclosed herein or a variant thereof, which binds to TfR, comprises at least 70% homology, at least 75% homology, at least 80% homology, at least 85% homology, at least 90% homology, at least 95% homology, at least 96% homology, at least 97% homology, at least 98% homology, or at least 99% homology or at least 100% homology to an endogenous or exogenous polypeptide known to bind TfR, for example, endogenous Transferrin or any one of the peptides listed in **TABLE 1**. In other embodiments, a peptide described herein binds to a protein of interest, which comprises at least 70% homology, at least 75% homology, at least 80% homology, at least 85% homology, at least 90% homology, at least 95% homology, at least 96% homology, at least 97% homology, at least 98% homology, or at least 99% homology or at least 100% homology to TfR, a fragment, homolog, or a variant thereof.

[0291] In some embodiments, peptides disclosed herein or variants thereof bind regions of TfR that comprise the amino acid residues corresponding to residues 506-510, 523-531, and 611-662 (the numbering of these amino acid residues is based on the following Uniprot reference protein sequence of endogenous human TFRC UniProtKB - P02786 (SEQ ID NO: 363, TFR1_HUMAN)). In some embodiments, the regions of TfR to which the peptides disclosed herein or variants thereof bind overlap with those of Tf, a fragment, homolog, or a variant thereof.

[0292] In other embodiments, a nucleic acid, vector, plasmid, or donor DNA comprises a sequence that encodes a peptide, peptide construct, or variant or functional fragment thereof, as described in the present disclosure. In further embodiments, certain parts or fragments of TfR-binding motifs (e.g., conserved binding motifs) can be grafted onto a peptide or peptide construct with a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361.

[0293] In some embodiments, peptides inhibit binding between TfR and Tf, or between TfR and any other protein. In some embodiments, peptides prevent TfR from protein-protein interaction and/or prevent TfR localization to a cell's nucleus. In some cases, peptides deactivate TfR. In some embodiments, peptides can cause TfR to be degraded, or prevent TfR from localization to a cell's nucleus, or prevent TfR from interacting with Tf or Tf-like proteins.

[0294] In some embodiments, peptides competitively bind to TfR as compared to endogenous Tf or any other endogenous or exogenous TfR binder by binding to a certain amino acid residue or motif of amino acid residues in TfR. Furthermore, a peptide can be selected for further testing or use based upon its ability to bind to the certain amino acid residue or motif of amino acid residues. The certain amino acid residue or motif of amino acid residues in TfR can be identified an amino acid residue or sequence of amino acid residues that are involved in the binding of TfR to Tf. A certain amino acid residue or motif of amino acid residues can be identified from a crystal structure of the TfR:Tf complex. In some embodiments, peptides (e.g., CDPs) demonstrate the resistance to heat, protease (pepsin), and reduction.

[0295] The peptides and peptide constructs (e.g., peptide conjugates or fusion peptides) comprising one or more of the amino acid sequences set forth in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 356 – SEQ ID NO: 361 can bind to a protein of interest. In some embodiments, the protein of interest is a TfR. In some embodiments, the peptides and peptide constructs (e.g., peptide conjugates or fusion peptides) that bind to a TfR comprise at least one of the amino acid sequences set forth in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 356 – SEQ ID NO: 361. In some embodiments, peptides, peptide constructs (e.g., peptide conjugates and fusion molecules) of the present disclosure that bind to a TfR comprise peptide derivatives or variants having at least 70% homology, at least 75% homology, at least 80% homology, at least 85% homology, at least 90% homology, at least 95% homology, at least 96% homology, at least 97% homology, at least 98%

homology, or at least 99% homology or at least 100% homology to amino acid sequences set forth in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, and SEQ ID NO: 356 – SEQ ID NO: 361.

[0296] In some embodiments, a peptide or a library of peptides is designed *in silico* without derivation from a naturally occurring scaffold of a knotted peptide. In other embodiments, a peptide or a library of peptides is designed *in silico* by derivation, grafting relevant protein-binding residues, or conserved residues in the protein-binding interface a naturally occurring peptide or protein known to bind to a protein or receptor of interest. In some embodiments, the peptide (e.g., SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, or SEQ ID NO: 171 – SEQ ID NO: 202) is a simple helix-turn-helix. In some embodiments, the helix-turn-helix can be used for pharmacophore transfer onto other scaffolds, for example engraftment of the required TfR-engaging surface onto the helix-turn-helix scaffold using fusion tagging.

[0297] In some embodiments, a peptide comprising SEQ ID NO: 1 is used as a scaffold or base sequence for further modifications, including addition, deletion, or amino acid substitution. In some embodiments, short sequences of amino acid residues such as GS are added at the N-terminus of a peptide. In some embodiments, peptides lack GS at the N-terminus. In some instances, peptides undergo one or more post-translational modifications.

[0298] TABLE 1 lists exemplary peptide sequences according to the methods and compositions of the present disclosure.

TABLE 1 – Exemplary Peptide Sequences

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 1	GSREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 2	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 3	GSREGCASRCTKYNAELEKCEARVVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 4	GSREGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 5	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 6	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELFDLLHCLDHCHSQ
SEQ ID NO: 7	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 8	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 9	GSREGCASRCTKYNAELEKCEARVVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 10	GSREGCASRCTKYNAELEKCEARVVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 11	GSREGCASRCTKYNAELEKCEARVVMSMSNTEEDCEQELFDLLHCLDHCHSQ

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 12	GSREGCASRCTKYNAELEKCEARVVSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 13	GSREGCASRCTKYNAELEKCEARVVSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 14	GSREGCASRCTKYNAELEKCEARVMSMSNTEETCVQELEDLLHCLDHCHSQ
SEQ ID NO: 15	GSREGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 16	GSREGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCVDHCHSQ
SEQ ID NO: 17	GSREGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCLDHCVSQ
SEQ ID NO: 18	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 19	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCVDHCHSQ
SEQ ID NO: 20	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCLDHCVSQ
SEQ ID NO: 21	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 22	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 23	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCVDHCVSQ
SEQ ID NO: 24	GSREGCASRCMKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 25	GSREGCASRCTKYNDELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 26	GSREGCASRCTKYNAELEKCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 27	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 28	GSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 29	GSREGCASRCMKYNDELEKCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 30	GSREGCASRCMKYNAELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 31	GSREGCASRCTKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 32	GSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 36	REGCASRCTKYNAELEKCEARVSSMSNTEETCVQELEDLLHCVDHCVSQ
SEQ ID NO: 37	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 38	REGCASRCTKYNAELEKCEARVVSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 39	REGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 40	REGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 41	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 42	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 43	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 44	REGCASRCTKYNAELEKCEARVVSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 45	REGCASRCTKYNAELEKCEARVVSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 46	REGCASRCTKYNAELEKCEARVVSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 47	REGCASRCTKYNAELEKCEARVVSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 48	REGCASRCTKYNAELEKCEARVVSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 49	REGCASRCTKYNAELEKCEARVMSMSNTEETCVQELEDLLHCLDHCHSQ
SEQ ID NO: 50	REGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 51	REGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCVDHCHSQ
SEQ ID NO: 52	REGCASRCTKYNAELEKCEARVMSMSNTEETCEQELEDLLHCLDHCVSQ
SEQ ID NO: 53	REGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 54	REGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCVDHCHSQ
SEQ ID NO: 55	REGCASRCTKYNAELEKCEARVMSMSNTEEDCVQELEDLLHCLDHCVSQ
SEQ ID NO: 56	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELFDLLHCVDHCHSQ
SEQ ID NO: 57	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELFDLLHCLDHCVSQ
SEQ ID NO: 58	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCVDHCVSQ
SEQ ID NO: 59	REGCASRCMKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 60	REGCASRCTKYNDELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 61	REGCASRCTKYNAELEKCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 62	REGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 63	REGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 64	REGCASRCMKYNDELEKCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 65	REGCASRCMKYNAELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 66	REGCASRCTKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 67	REGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 136	GSREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 137	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 138	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 139	GSREGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 140	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 141	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCLDHCHSQ
SEQ ID NO: 142	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 143	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 144	GSREGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 145	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 146	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCLDHCHSQ
SEQ ID NO: 147	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 148	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 149	GSREGCASRCTRYNAELERCEARVMSMSNTEETCVQELEDLLHCLDHCHSQ
SEQ ID NO: 150	GSREGCASRCTRYNAELERCEARVMSMSNTEETCEQELFDLLHCLDHCHSQ
SEQ ID NO: 151	GSREGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCVDHCHSQ
SEQ ID NO: 152	GSREGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCLDHCVSQ
SEQ ID NO: 153	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCVQELFDLLHCLDHCHSQ
SEQ ID NO: 154	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCVDHCHSQ
SEQ ID NO: 155	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCLDHCVSQ
SEQ ID NO: 156	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCVDHCHSQ
SEQ ID NO: 157	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCLDHCVSQ
SEQ ID NO: 158	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCVDHCVSQ
SEQ ID NO: 159	GSREGCASRCMRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 160	GSREGCASRCTRYNDELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 161	GSREGCASRCTRYNAELERCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 162	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 163	GSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 164	GSREGCASRCMRYNDELERCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 165	GSREGCASRCMRYNAELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 166	GSREGCASRCTRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 167	GSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 171	REGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 172	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 173	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 174	REGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 175	REGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 176	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCLDHCHSQ
SEQ ID NO: 177	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 178	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 179	REGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCLDHCHSQ
SEQ ID NO: 180	REGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCLDHCHSQ
SEQ ID NO: 181	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCLDHCHSQ
SEQ ID NO: 182	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCVDHCHSQ
SEQ ID NO: 183	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCVSQ
SEQ ID NO: 184	REGCASRCTRYNAELERCEARVMSMSNTEETCVQELEDLLHCLDHCHSQ
SEQ ID NO: 185	REGCASRCTRYNAELERCEARVMSMSNTEETCEQELFDLLHCLDHCHSQ
SEQ ID NO: 186	REGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCVDHCHSQ
SEQ ID NO: 187	REGCASRCTRYNAELERCEARVMSMSNTEETCEQELEDLLHCLDHCVSQ
SEQ ID NO: 188	REGCASRCTRYNAELERCEARVMSMSNTEEDCVQELFDLLHCLDHCHSQ
SEQ ID NO: 189	REGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCVDHCHSQ
SEQ ID NO: 190	REGCASRCTRYNAELERCEARVMSMSNTEEDCVQELEDLLHCLDHCVSQ
SEQ ID NO: 191	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCVDHCHSQ
SEQ ID NO: 192	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELFDLLHCLDHCVSQ
SEQ ID NO: 193	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCVDHCVSQ
SEQ ID NO: 194	REGCASRCMRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 195	REGCASRCTRYNDELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 196	REGCASRCTRYNAELERCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 197	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 198	REGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 199	REGCASRCMRYNDELERCEARVMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 200	REGCASRCMRYNAELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 201	REGCASRCTRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 202	REGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ

[0299] In some embodiments, a TfR-binding peptide disclosed herein comprises GSREGCAX₁RCX₂KYX₄DEX₂X₃KCX₃ARMMSMSNTEEDCEQEX₂EDX₂X₂YCX₂X₃X₅CX₅X₁X₄ (SEQ ID NO: 206) or REGCAX₁RCX₂KYX₄DEX₂X₃KCX₃ARMMSMSNTEEDCEQEX₂EDX₂X₂YCX₂X₃X₅CX₅X₁X₄ (SEQ ID NO: 334), wherein X₁ can be independently selected from S, T, D, or N, X₂ can be independently selected from A, M, I, L, or V, X₃ can be independently selected from D, E, N, Q, S, or T, X₄ can be independently selected from D, E, H, K, R, N, Q, S, or T, and X₅ can be independently selected from H, K, R, N, Q, S, or T.

[0300] In some embodiments, a TfR-binding peptide disclosed herein comprises GSREX₁CX₂X₃RCX₄KYX₅DEX₆X₇KCX₈ARMMSMSNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 207) or REX₁CX₂X₃RCX₄KYX₅DEX₆X₇KCX₈ARMMSMSNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 335), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇ and X₈ are TfR binding interface residues and can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQEX₁EDX₂X₃YCX₄X₅X₆CX₇X₈X₉ (SEQ ID NO: 208) or REGCASRCMKYNDELEKCEARMMSMSNTEEDCEQEX₁EDX₂X₃YCX₄X₅X₆CX₇X₈X₉ (SEQ ID NO: 336), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, and X₉ are TfR binding interface residues and can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSREX₁CX₂X₃RCX₄KYX₅DEX₆X₇KCX₈ARMMSMSNTEEDCEQEX₉EDX₁₀X₁₁YCX₁₂X₁₃X₁₃CX₁₅X₁₆X₁₇ (SEQ ID NO: 209) or REX₁CX₂X₃RCX₄KYX₅DEX₆X₇KCX₈ARMMSMSNTEEDCEQEX₉EDX₁₀X₁₁YCX₁₂X₁₃X₁₃CX₁₅X₁₆X₁₇ (SEQ ID NO: 337), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆ and X₁₇ are TfR binding interface residues and can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises GSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 32).

[0301] In some embodiments, a TfR-binding peptide disclosed herein comprises X₁X₂X₃X₄GX₅ASX₆X₇MX₈X₉NX₁₀X₁₁LEX₁₂X₁₃EX₁₄X₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄X₂₅X₂₆

X₂₇X₂₈X₂₉X₃₀X₃₁X₃₂X₃₃X₃₄X₃₅X₃₆X₃₇X₃₈X₃₉X₄₀X₄₁X₄₂X₄₃ (SEQ ID NO 210), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, X₃₄, X₃₅, X₃₆, X₃₇, X₃₈, X₃₉, X₄₀, X₄₁, X₄₂, and X₄₃ can independently be any amino acid.

[0302] In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂X₃X₄X₅X₆X₇X₈X₉X₁₀X₁₁X₁₂X₁₃X₁₄X₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄X₂₅X₂₆X₂₇X₂₈X₂₉X₃₀X₃₁X₃₂X₃₃X₃₄X₃₅X₃₆X₃₇LX₃₈X₃₉LLX₄₀X₄₁LDHX₄₂HSQ (SEQ ID NO: 211), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, X₃₄, X₃₅, X₃₆, X₃₇, X₃₈, X₃₉, X₄₀, X₄₁, and X₄₂ can independently be any amino acid.

[0303] In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂X₃X₄GX₅ASX₆MX₇NX₈NX₉LEX₁₀X₁₁LEX₁₂X₁₃EX₁₄X₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄X₂₅X₂₆X₂₇X₂₈X₂₉LX₃₀X₃₁LLX₃₂X₃₃LDHX₃₄HSQ (SEQ ID NO: 212), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, and X₃₄ can independently be any amino acid.

[0304] In some embodiments, a TfR-binding peptide or peptide construct disclosed herein

comprises at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 100% sequence homology to any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361, or any variant, homolog, or functional fragment thereof. In some embodiments, a TfR-binding peptide or peptide construct disclosed herein comprises any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361, or any variant, homolog, or functional fragment thereof. In some embodiments, a peptide that binds to a TfR comprises the amino acid sequence set forth in SEQ ID NO: 32.

[0305] In some embodiments, a TfR-binding peptide comprises canonical amino acid residues as surface interface residues at any one of the corresponding positions 5, 7, 8, 14, 17, 18, 21, 38, 42, 45, 46, 47, 50, 51, with reference to SEQ ID NO: 32 or a combination thereof. In some embodiments, a TfR-binding peptide comprises canonical amino acid residues as surface interface residues at any one of the corresponding positions G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, Q51, with reference to SEQ ID NO: 32 or a combination thereof.

In some embodiments, the peptide or peptide construct of the present disclosure comprises at least one or more of these corresponding residues in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361. Such peptides can accordingly be engineered with enhanced binding to TfR. In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂X₃X₄GX₅ASX₆X₇X₈X₉X₁₀NX₁₁X₁₂LEX₁₃X₁₄EX₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄X₂₅X₂₆X₂₇X₂₈X₂₉X₃₀LX₃₁X₃₂X₃₃LX₃₄X₃₅LDHX₃₆X₃₇SQ (SEQ ID NO: 213), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, X₃₄, X₃₅, X₃₆, and X₃₇ can independently be any amino acid.

[0306] In some embodiments, surface-distal hydrophilic amino acid residues (e.g., D, E, H, K, R, N, Q, S, or T) present in the amino acid sequence of a peptide contribute to peptide solubility. In some embodiments, a peptide as disclosed herein comprises a hydrophilic amino acid residue at any one of the corresponding positions 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, 40, with reference to SEQ ID NO: 32, or any combination thereof. In some instances, a peptide of the present disclosure comprises hydrophilic amino acid residues at the following corresponding positions: R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, D40, with reference to SEQ ID NO: 32, or any combination thereof. In some embodiments, any one of or any combination of corresponding positions R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, D40 with reference to SEQ ID NO: 32, can be mutated to another hydrophilic residue without significantly impacting solubility or TfR-binding. In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂REX₃X₄X₅X₆RX₇X₈KX₉X₁₀DEX₁₁X₁₂KX₁₃X₁₄X₁₅RX₁₆X₁₇SX₁₈SNTEEDX₁₉EQEX₂₀EDX₂₁X₂₂X₂₃X₂₄X₂₅X₂₆X₂₇X₂₈X₂₉X₃₀X₃₁ (SEQ ID NO: 214), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, and X₃₁ can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSX₁X₂GCASX₃CMX₄YNX₅X₆LEX₇CEAX₈MMX₉MX₁₀X₁₁X₁₂X₁₃X₁₄X₁₅CX₁₆X₁₇X₁₈LX₁₉X₂₀LLYCLDHCHSQ (SEQ ID NO: 215) or

X₁X₂GCASX₃CMX₄YNX₅X₆LEX₇CEAX₈MMX₉MX₁₀X₁₁X₁₂X₁₃X₁₄X₁₅CX₁₆X₁₇X₁₈LX₁₉X₂₀LLYCLDHCHSQ (SEQ ID NO: 338), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃,

X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, and X₂₀ can be independently selected from D, E, H, K, R, N, Q, S, or T.

[0307] In some embodiments, a peptide of the present disclosure comprises hydrophilic residues (e.g., D, E, H, K, R, N, Q, S, or T) at corresponding positions 15, 35, 39, 49, with reference to SEQ ID NO: 32, or any combination thereof. In some instances, a peptide of the present disclosure comprises hydrophilic amino acid residues at the following corresponding positions: D15, E35, E39, H49, with reference to SEQ ID NO: 32, or any combination thereof. In some embodiments, any one of or any combination of corresponding positions D15, E35, E39, H49 with reference to SEQ ID NO: 32, can be mutated to another hydrophilic residue without significantly impacting solubility or TfR-binding. In some embodiments, a TfR-binding peptide disclosed herein comprises. In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂X₃X₄X₅X₆X₇X₈X₉X₁₀X₁₁X₁₂X₁₃X₁₄DX₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄X₂₅X₂₆X₂₇X₂₈X₂₉X₃₀X₃₁X₃₂X₃₃EX₃₄X₃₅X₃₆EX₃₇X₃₈X₃₉X₄₀X₄₁X₄₂X₄₃X₄₄X₄₅HX₄₆X₄₇ (SEQ ID NO: 216), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, X₃₄, X₃₅, X₃₆, X₃₇, X₃₈, X₃₉, X₄₀, X₄₁, X₄₂, X₄₃, X₄₄, X₄₅, X₄₆, and X₄₇ can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSREGCASRCMKYNX₁ELEKCEARMMSMSNTEEDCX₂QELX₃DLLYCLDHXCX₄SQ (SEQ ID NO: 217) or

REGCASRCMKYNX₁ELEKCEARMMSMSNTEEDCX₂QELX₃DLLYCLDHXCX₄SQ (SEQ ID NO: 339), wherein X₁, X₂, X₃, and X₄ can be independently selected from D, E, H, K, R, N, Q, S, or T.

[0308] In some embodiments, a peptide of the present disclosure comprises hydrophobic residues (e.g., A, M, I, L, V, F, W, or Y) at corresponding positions 15, 35, 39, 49, with reference to SEQ ID NO: 32, or any combination thereof. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSREGCASRCMKYNX₁ELEKCEARMMSMSNTEEDCX₂QELX₃DLLYCLDHXCX₄SQ (SEQ ID NO: 218) or

REGCASRCMKYNX₁ELEKCEARMMSMSNTEEDCX₂QELX₃DLLYCLDHXCX₄SQ (SEQ ID NO: 340), wherein X₁, X₂, X₃, and X₄ can be independently selected from A, M, I, L, V, F, W, or Y. In some embodiments, hydrophilic amino acid residues at any one of the corresponding

positions 15, 35, 39, and 49, with reference to SEQ ID NO: 32, are associated with higher binding affinity for TfR (e.g., target engagement) and higher solubility. In some embodiments, mutation of an amino acid residue at any one of the corresponding positions 15, 35, 39, and 49, with reference to SEQ ID NO: 32, from a hydrophobic to a hydrophilic residue can lead to higher binding affinity for TfR (e.g., target engagement) and higher solubility.

[0309] In some embodiments, a peptide of the present disclosure comprises hydrophobic residues (e.g., A, M, I, L, V, F, W, or Y) at corresponding positions 11, 25, 27, with reference to SEQ ID NO: 32, or any combination thereof. In some embodiments, a peptide of the present disclosure comprises hydrophilic residues (e.g., D, E, H, K, R, N, Q, S, or T) at corresponding positions 11, 25, 27, with reference to SEQ ID NO: 32, or any combination thereof. In some embodiments, hydrophobic amino acid residues at any one of the corresponding positions 11, 25, and 27, with reference to SEQ ID NO: 32, are associated with higher binding affinity for TfR (e.g., target engagement) and higher solubility. In some embodiments, mutation of an amino acid residue at any one of the corresponding positions 11, 25, and 27, with reference to SEQ ID NO: 32, from a hydrophilic residue to a hydrophobic residue can lead to higher binding affinity for TfR (e.g., target engagement) and higher solubility. In some embodiments, a peptide of the present disclosure comprises hydrophobic amino acid residues at the corresponding positions M11, M25, M27, with reference to SEQ ID NO: 32, or any combination thereof. In some instances, a peptide comprises the hydrophobic amino acid residues at the corresponding positions M11, M25, and M27, with reference to SEQ ID NO: 32. In some embodiments, any combination of the corresponding positions M11, M25, and M27, with reference to SEQ ID NO: 32, can be mutated to another hydrophobic residue without significantly impacting solubility or TfR-binding. In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂X₃X₄X₅X₆X₇X₈X₉X₁₀MX₁₁X₁₂X₁₃X₁₄X₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃MX₂₄MX₂₅X₂₆X₂₇X₂₈X₂₉X₃₀X₃₁X₃₂X₃₃X₃₄X₃₅X₃₆X₃₇X₃₈X₃₉X₄₀X₄₁X₄₂X₄₃X₄₄X₄₅X₄₆X₄₇X₄₈ (SEQ ID NO: 219),

wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, X₃₄, X₃₅, X₃₆, X₃₇, X₃₈, X₃₉, X₄₀, X₄₁, X₄₂, X₄₃, X₄₄, X₄₅, X₄₆, X₄₇, and X₄₈ can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSREGCASRCX₁KYNDELEKCEARMX₂SX₃SNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 220) or REGCASRCX₁KYNDELEKCEARMX₂SX₃SNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 341), wherein X₁, X₂, and X₃ can be independently selected from A, M, I, L, V, F,

W, or Y. In some embodiments, a TfR-binding peptide disclosed herein comprises GSREGCASRCX₁KYNDELEKCEARMX₂SX₃SNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 221) or REGCASRCX₁KYNDELEKCEARMX₂SX₃SNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 342), wherein X₁, X₂, and X₃ can be independently selected from D, E, H, K, R, N, Q, S, or T.

In some embodiments, a peptide of the present disclosure comprises an aliphatic amino acid residue (e.g., A, M, I, L, or V) at corresponding position 45, with reference to SEQ ID NO: 32. In some embodiments, a peptide of the present disclosure comprises an aromatic amino acid residue (e.g., F, W, or Y) at corresponding position 45. In some embodiments, an aliphatic amino acid residue at corresponding position 45 is associated with higher binding affinity to TfR. In some instances, a peptide comprises the aliphatic amino acid residue corresponding to L45, with reference to SEQ ID NO: 32. In some embodiments, mutation of an amino acid residue at corresponding position 45 from an aromatic residue to an aliphatic residue can lead to higher binding affinity for TfR (e.g., target engagement) and higher solubility. In some embodiments, mutating corresponding position L45 to another aliphatic residue may not significantly impact solubility or TfR-binding. In some embodiments, a TfR-binding peptide disclosed herein comprises

X₁X₂X₃X₄X₅X₆X₇X₈X₉X₁₀X₁₁X₁₂X₁₃X₁₄X₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄X₂₅X₂₆X₂₇X₂₈X₂₉X₃₀X₃₁X₃₂X₃₃X₃₄X₃₅X₃₆X₃₇X₃₈X₃₉X₄₀X₄₁X₄₂X₄₃X₄₄LX₄₅X₄₆X₄₇X₄₈X₄₉X₅₀ (SEQ ID NO: 222), wherein X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₄, X₂₅, X₂₆, X₂₇, X₂₈, X₂₉, X₃₀, X₃₁, X₃₂, X₃₃, X₃₄, X₃₅, X₃₆, X₃₇, X₃₈, X₃₉, X₄₀, X₄₁, X₄₂, X₄₃, X₄₄, X₄₅, X₄₆, X₄₇, X₄₈, X₄₉, and X₅₀ can independently be any amino acid. In some embodiments, a TfR-binding peptide disclosed herein comprises

GSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCX₁DHCHSQ (SEQ ID NO: 223) or REGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCX₁DHCHSQ (SEQ ID NO: 343), wherein X₁ can be independently selected from A, M, I, L, or V.

[0310] In some embodiments, a peptide of the present disclosure comprises

GSREGCASRCMX₁YNDELEX₂CEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 224) or REGCASRCMX₁YNDELEX₂CEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 344), wherein X₁ and X₂ can be independently selected from K or R. In some embodiments, these residues at corresponding position 12 and 19, with reference to SEQ ID NO: 32, can be used for chemical conjugation to another molecule (e.g., an active or a detectable

agent). In some embodiments, X₁ and X₂ are both R and chemical conjugation occurs at the N-terminus of the peptide.

[0311] In some embodiments, mutations in any one or more of the amino acid residues of a peptide of the present disclosure can improve binding affinity of the peptide to TfR. In some embodiments, mutations in 5-80% of amino acid residues of a peptide of the present disclosure improve the binding affinity of the peptide to TfR. In some embodiments, mutations in 1-100%, 5-100%, or 5-50% of amino acid residues of a peptide of the present disclosure improve binding affinity of the peptide to TfR. In some embodiments, mutations in 15-50% of amino acid residues of a peptide of the present disclosure improve binding affinity of the peptide to TfR. In some embodiments, mutations in 15-30% of amino acid residues of a peptide of the present disclosure improve binding affinity of the peptide to TfR. In some embodiments, mutations in 25-30% of amino acid residues of a peptide of the present disclosure improve binding affinity of the peptide to TfR. For example, mutations in 14 of the 51 amino acid residues (27.5%) of a peptide having a sequence of SEQ ID NO: 32 can improve binding affinity of the peptide to TfR.

[0312] In some embodiments, mutations in any one or more of the amino acid residues of a peptide of the present disclosure can lie at the binding interface of TfR. In some embodiments, a mutation to a peptide can improve binding affinity, which can be beneficial to binding and transcytosis of a peptide or peptide construct disclosed herein. In some embodiments, the peptides provided herein can have many mutations or few mutations to obtain optimal activity, wherein optimal activity is sufficient binding for engagement of the TfR, but not necessarily binding that is so strong as to preclude release of the peptide and/or peptide construct after transcytosis. Thus, peptides of the present disclosure can comprise a number of mutations (also referred to as % mutated amino acid residues) that tune binding affinity and off rate to obtain optimal binding, function (e.g., transcytosis, BBB-penetration, cell membrane penetration, transport across a biological barrier), and release of the peptide or peptide construct. Thus, mutations that result in the highest possible affinity may not necessarily correlate to a superior peptide having optimal binding and transcytosis.

[0313] In some embodiments, 1-100% or 5-100% of amino acid residues of a peptide of the present disclosure lie at the binding interface of TfR. In some embodiments, 10-90% of amino acid residues of a peptide of the present disclosure lie at the binding interface of TfR. In some embodiments, 20-80% of amino acid residues of a peptide of the present disclosure lie at the binding interface of TfR. In some embodiments, 30-70% of amino acid residues of a peptide of

the present disclosure lie at the binding interface of TfR. In some embodiments, 40-60% of amino acid residues of a peptide of the present disclosure lie at the binding interface of TfR. In some embodiments, 30-35% of amino acid residues of a peptide of the present disclosure lie at the binding interface of TfR. For example, 17 of the 51 amino acid residues (33%) of a peptide having a sequence of SEQ ID NO: 32 can lie at the binding interface of TfR.

[0314] In some embodiments, mutations in any one or more of the amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR can improve binding affinity of the peptide to TfR. In some embodiments, mutations in 1-100% or 5-100% of amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 5-80% of amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 10-70% of amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 15-60% of amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 20-50% of amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 25-30% of amino acid residues of a peptide of the present disclosure that lie at the binding interface of TfR improve binding affinity of the peptide to TfR. For example, mutations in 5 of the 17 amino acid residues (29%) of a peptide having a sequence of SEQ ID NO: 32 that lie at the binding interface of TfR and can improve binding affinity of the peptide to TfR.

[0315] In some embodiments, mutations in any one or more of the amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR. In some embodiments, 1-100% or 5-100% of amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR. In some embodiments, 10-90% of amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR. In some embodiments, 20-80% of amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR. In some embodiments, 30-70% of amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR. In some embodiments, 40-60% of amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR. In some embodiments, 65-70% of amino acid residues of a peptide of the present disclosure are distal to

the binding interface of TfR. For example, 34 of the 51 amino acid residues (66%) of a peptide having a sequence of SEQ ID NO: 32 can lie at the binding interface of TfR.

[0316] In some embodiments, mutations in any one or more of the amino acid residues of a peptide of the present disclosure are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 1-100% or 5-100% of amino acid residues of a peptide of the present disclosure that are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 5-80% of amino acid residues of a peptide of the present disclosure that are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 10-70% of amino acid residues of a peptide of the present disclosure that are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 15-60% of amino acid residues of a peptide of the present disclosure that are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 20-50% of amino acid residues of a peptide of the present disclosure that are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. In some embodiments, mutations in 25-30% of amino acid residues of a peptide of the present disclosure that are distal to the binding interface of TfR improve binding affinity of the peptide to TfR. For example, mutations in 5 of the 17 amino acid residues that are distal to the binding interface of TfR can improve binding affinity of the peptide to TfR. For example, mutations in 9 of the 34 amino acid residues (26.5%) of a peptide having a sequence of SEQ ID NO: 32 that are distal to the binding interface of TfR can improve binding affinity of the peptide to TfR. In some embodiments, and without being bound to any theory, one or more mutations in the amino acid residues of the peptide that are distal to the binding interface of TfR can improve protein folding, enhance protein solubility, and/or alter the backbone geometry that can improve binding through an optimized interface shape complementarity.

[0317] In some embodiments, a peptide of the present disclosure can comprise a sequence having cysteine residues at one or more of positions 11, 12, 13, 14, 19, 20, 21, 22, 36, 38, 39, 41. In some embodiments, a peptide comprises Cys at positions 11, 12, 19, 20, 36, 39, or any combination thereof. For example, in certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 11. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 12. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 13. In certain embodiments, a peptide

can comprise a sequence having a cysteine residue at position 14. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 19. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 20. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 21. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 22. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 36. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 38. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 39. In certain embodiments, a peptide can comprise a sequence having a cysteine residue at position 41. In some embodiments, the first cysteine residue in the sequence can be disulfide bonded with the 4th cysteine residue in the sequence, the 2nd cysteine residue in the sequence can be disulfide bonded to the 5th cysteine residue in the sequence, and the 3rd cysteine residue in the sequence can be disulfide bonded to the 6th cysteine residue in the sequence. Optionally, a peptide can comprise one disulfide bridge that passes through a ring formed by two other disulfide bridges, also known as a “two-and-through” structure system. In some embodiments, the peptides disclosed herein can have one or more cysteines mutated to serine.

[0318] In some embodiments, peptides of the present disclosure comprise at least one cysteine residue. In some embodiments, peptides of the present disclosure comprise at least two cysteine residues. In some embodiments, peptides of the present disclosure comprise at least three cysteine residues. In some embodiments, peptides of the present disclosure comprise at least four cysteine residues. In some embodiments, peptides of the present disclosure comprise at least five cysteine residues. In some embodiments, peptides of the present disclosure comprise at least six cysteine residues. In some embodiments, peptides of the present disclosure comprise at least ten cysteine residues. In some embodiments, a peptide of the present disclosure comprises six cysteine residues.

[0319] In some embodiments, a peptide of the present disclosure comprises an amino acid sequence having cysteine residues at one or more positions. In some embodiments, the one or more cysteine residues are located at any one of the amino acid positions 6, 10, 20, 34, 44, 48, or any combination thereof. In some aspects of the present disclosure, the one or more cysteine (C) residues participate in disulfide bonds with various pairing patterns (e.g., C10-C20). In some embodiments, the pairing patterns are C6-C48, C10-C44, and C20-C34. In some embodiments,

the peptides as described herein comprise at least one, at least two, or at least three disulfide bonds. In some embodiments, at least one, at least two, or at least three disulfide bonds are arranged according to the C6-C48, C10-C44, and C20-C34 pairing patterns, or a combination thereof. In some embodiments, peptides as described herein comprise three disulfide bonds with the pairing patterns C6-C48, C10-C44, and C20-C34.

[0320] In certain embodiments, a peptide comprises a sequence having a cysteine residue at position 6. In certain embodiments, a peptide comprises a sequence having a cysteine residue at position 10. In certain embodiments, a peptide comprises a sequence having a cysteine residue at position 20. In certain embodiments, a peptide comprises a sequence having a cysteine residue at position 34. In certain embodiments, a peptide comprises a sequence having a cysteine residue at position 44. In certain embodiments, a peptide comprises a sequence having a cysteine residue at position 50. In some embodiments, the first cysteine residue in the sequence is disulfide bonded with the last cysteine residue in the sequence. In some embodiments, the second cysteine residue in the sequence is disulfide bonded with the second to the last cysteine residue in the sequence. In some embodiments, the third cysteine residue in the sequence is disulfide bonded with the third to the last cysteine residue in the sequence and so forth.

[0321] In some embodiments, the first cysteine residue in the sequence is disulfide bonded with the 6th cysteine residue in the sequence, the 2nd cysteine residue in the sequence is disulfide bonded to the 5th cysteine residue in the sequence, and the 3rd cysteine residue in the sequence is disulfide bonded to the 4th cysteine residue in the sequence. Optionally, a peptide can comprise one disulfide bridge that passes through a ring formed by two other disulfide bridges, also known as a “two-and-through” structure system. In some embodiments, the peptides disclosed herein have one or more cysteines mutated to serine.

[0322] In some embodiments, a peptide comprises no cysteine or disulfides. In some embodiments, a peptide comprises 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 11 or more, 12 or more, 13 or more, 14 or more, or 15 or more cysteine or disulfides. In other embodiments, 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more cysteine residues have been replaced with serine residues. In some embodiments, 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more cysteine residues have been replaced with threonine residues.

[0323] In some embodiments, a peptide comprises no Cys or disulfides. In some embodiments, a peptide comprises 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 11 or more, 12 or more, 13 or more, 14 or more, or 15 or more Cys or disulfides. In other embodiments, 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more Cys residues have been replaced with Ser residues. In some embodiments, 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more Cys residues have been replaced with Thr residues.

[0324] In some instances, one or more or all of the methionine residues in the peptide are replaced by leucine or isoleucine. In some instances, one or more or all of the tryptophan residues in the peptide are replaced by phenylalanine or tyrosine. In some instances, one or more or all of the asparagine residues in the peptide are replaced by glutamine. In some embodiments, the N-terminus of the peptide is blocked, such as by an acetyl group. Alternatively or in combination, in some instances, the C-terminus of the peptide is blocked, such as by an amide group. In some embodiments, the peptide is modified by methylation on free amines.

[0325] For example, full methylation can be accomplished through the use of reductive methylation with formaldehyde and sodium cyanoborohydride.

[0326] In some embodiments, the peptides or peptide constructs as described herein target and/or penetrate a TfR-expressing cellular layer or barrier and/or the membrane of a TfR-expressing cell. In some embodiments, a peptide targets and/or penetrates a cell membrane of a cell, wherein said cell is located in the CNS such as the brain. For example, a peptide construct comprising a TfR-binding peptide and one or more active agents (e.g., a therapeutic or diagnostic compound) crosses a cellular barrier (e.g., BBB) via vesicular transcytosis, and subsequently targets and/or penetrates the cell membrane of a cell located within the CNS to deliver said one or more active agents to that cell.

[0327] In various embodiments, a peptide construct comprising a TfR-binding peptide as described herein and one or more active agents (e.g., a therapeutic or diagnostic compound) targets and/or penetrates the cell membrane of a TfR-expressing cell located in the gastrointestinal tract, spleen, liver, kidney, muscle, bone marrow, brain, or skin. In some cases, the TfR-expressing cell is a tumor cell, an immune cell, an erythrocyte, an erythrocyte precursor cell, a stem cell, a bone marrow cell, or stem cell. In some cases, the TfR-binding peptide is responsible for targeting the cell, e.g., in cases where the cell is overexpressing a TfR. In various

embodiments, a peptide construct as described herein comprising a TfR-binding peptide conjugated to, linked to, or fused to one or more cargo molecules (e.g., an active and/or detectable agent) targets and/or penetrates the cell membrane of a cell located within various organs such as the spleen, brain, liver, kidney, muscle, bone marrow, gastrointestinal tract, or skin.

[0328] In some cases, the cargo molecule promotes the targeting of a specific cell, cell population, or tissue. In some cases, it is a combination of TfR-mediated cell targeting and cargo molecule promoted cell targeting. In some aspects, a peptide or peptide construct (e.g., peptide conjugate or fusion peptide) of the present disclosure is used to target said cell, cell population, or tissue in order to exert a certain biological (e.g., therapeutic) effect. In some aspects, a peptide or peptide construct (e.g., peptide conjugate or fusion peptide) of the present disclosure is used to deliver the cargo molecule into said cell to exert a certain biological effect.

[0329] In some embodiments, peptides can comprise at least one or more tag peptide sequences for improved cell penetration. For example, peptides can comprise at least one or multiple Arg residues or residues from Tat protein for improved cell penetration property. Additional tag peptide sequences can include CysTat (CYRKKRRQRRR; SEQ ID NO: 71), S19-TAT (PFVIGAGVLGALGTGIGGIGRKKRRQRRR; SEQ ID NO: 72), R8 (RRRRRRRR; SEQ ID NO: 73), pAntp (RQIKIWFQNRRMKWKK; SEQ ID NO: 74), Pas-TAT (FFLIPKGGRKKRRQRRR; SEQ ID NO: 75), Pas-R8 (FFLIPKGRRRRRRRRR; SEQ ID NO: 76), PasFHV (FFLIPKGRRRRNRTRNRNRRVR; SEQ ID NO: 77), Pas-pAntp (FFLIPKGRQIKIWFQNRRMKWKK; SEQ ID NO: 78), F2R4 (FFRRRR; SEQ ID NO: 79), B55 (KAVLGATKIDLPVDINDPYDLGLLLRHLRHHSNLLANIGDPAVREQVLSAMQEEE; SEQ ID NO: 80), auzurin (LSTAADMQGVVTDGMASGLDKDYLPDD; SEQ ID NO: 81), IMT-P8 (RRWRRWNRFNRRRCR; SEQ ID NO: 82), BR2 (RAGLQFPVGRLLRLLR; SEQ ID NO: 83), OMOTAG1 (KRAHHNALERKRR; SEQ ID NO: 84), OMOTAG2 (RRMKANARERNRM; SEQ ID NO: 85), pVEC (LLIILRRRIRKQAAHAHSK; SEQ ID NO: 86), SynB3 (RRLSYSRRRF; SEQ ID NO: 87), DPV1047 (VKRGLKLRHVRPRVTRMDV; SEQ ID NO: 88), CY105Y (CSIPPEVKFNKPFVYLI; SEQ ID NO: 89), Transportan (GWTLSAGYLLGKINLKALAALAKKIL; SEQ ID NO: 90), MTS (KGEAAVLLPVLLAAPG; SEQ ID NO: 91), hLF (KCFQWQRNMRKVRGPPVSCIQR; SEQ ID NO: 92), PFVYLI (PFVYLI; SEQ ID NO: 93), yBBR (VLDSLEFIASKL, SEQ ID NO: 94), DRI-TAT31 (rrrqrrkrgy, wherein the lowercase notation indicates D-amino acids; SEQ ID NO:

95), cyclic heptapeptide cyclo (cFΦR₄) (FΦRRRRQ, where Φ is l-2-naphthylalanine, the entire moiety is cyclized, and Gln serves as a conjugation handle and can be substituted for other functional groups such as Lys (for amine coupling) or Cys (for sulfhydryl coupling); SEQ ID NO: 96), DPV3 (RKKRRRESRKKRRRES; SEQ ID NO: 403), C10H (RKGfYKRKQCKPSRGRKR; SEQ ID NO: 404), VP22 (NAATATRGRSAASRPTQRPRAPARSASRPRRPVQ; SEQ ID NO: 405), TP10 (AGYLLGKINLKALAALAKKIL; SEQ ID NO: 406), MAP (KLALKLALKALKAALKLA; SEQ ID NO: 407), BPrPp (MVKSKIGSWILVLFVAMWSDVGLCKKRP; SEQ ID NO: 408), ARF (MVRRLVTLRIRACGPPRVRV; SEQ ID NO: 409), GALA (WEAALAEALAEALAEHLAEALAEALAA; SEQ ID NO: 410), SAP (VRLPPPVRLLPPPVRLLPPP; SEQ ID NO: 411), MPG (GLAFLGFLGAAGSTMGAWSQPKKKRKV; SEQ ID NO: 412), Pep-1 (KETWWETWWTEWSQPKKKRKV; SEQ ID NO: 413), [WR]₄ (WRWRWRWR; SEQ ID NO: 414), Ig(v) (MGLGLHLLVLAAALQGAKKKKRKV; SEQ ID NO: 415), K-FGF (AAVALLPAVLLAHLAP; SEQ ID NO: 416), Melittin (GIGAVLKVLTTGLPALISWIKRKRQQ; SEQ ID NO: 417), gH625 (HGLASTLTRWAHYNALIRAF; SEQ ID NO: 418), HIV-1 TAT protein (48–60) (GRKKRRQRRRPPQ; SEQ ID NO: 419), MPG HIV-gp41/SV40 T-antigen (GALFLGFLGAAGSTMGAWSQPKKKRKV; SEQ ID NO: 420), R6W3 (RRWWRRWRR; SEQ ID NO: 421), NLS (CGYGPKKKRKYGG; SEQ ID NO: 422), 8-lysines (KKKKKKKK; SEQ ID NO: 423), HRSV (RRIPNRRPRR; SEQ ID NO: 424), AIP6 (RLRWR; SEQ ID NO: 425), Pep-1 (KETWWETWWTEWSQPKKKRKV; SEQ ID NO: 426), MAP17 (QLALQLALQALQAALQLA; SEQ ID NO: 427), VT5 (DPKGD PKGVTVTVTVTVTGKGD PKPD; SEQ ID NO: 428), Bac7 (RRIRPRPPRLPRPRPRPLPFPRPG; SEQ ID NO: 429), (PPR)_n ((PPRPPRPPR; SEQ ID NO: 430), (PPRPPRPPRPPR; SEQ ID NO: 431), (PPRPPRPPRPPRPPR; SEQ ID NO: 432), (PPRPPRPPRPPRPPRPPR; SEQ ID NO: 433)), INF7 (GLFEAIEGFIENGWEGMIDGWYGC; SEQ ID NO: 434), CADY (GLWRALWRLRLSLWRLWRA; SEQ ID NO: 435), Pep-7 (SDLWEMMMVSLACQY; SEQ ID NO: 436), TGN (TGNYKALHPHNG ; SEQ ID NO: 437), Ku-70 (VPMLK ; SEQ ID NO: 438), CPP (RRRRRGRRRRRG; SEQ ID NO: 452) (RRRRRGRRRRRG ; SEQ ID NO: 439), SVS-1 (KVKVKVKVDPPTKVKVKVK ; SEQ ID NO: 440), L-CPP (LAGRRRRRRRRRK; SEQ ID NO: 441), RLW (RLWMRWYSPRTRAYG;

SEQ ID NO: 442), K16ApoE (KKKKKKKKKKKKKKKKKKLRVRLASHLRKLRKRLLRDA ; SEQ ID NO: 443), Angiopep-2 (TFFYGGSRGKRNNFKTEEY; SEQ ID NO: 444), ACPP (EEEEEEEEPLGLAGRRRRRRRRN; SEQ ID NO: 445), KAFK (KAFKLAARLYRKALARQLGVAA; SEQ ID NO: 446), hCT (9-32) (LGTYTQDFNKFHTFPQTAIGVGAP; SEQ ID NO: 447), VP22(version2) (DAATATRGRSAASRPTQRPRAPARSASRRRPVE; SEQ ID NO: 448), MPG (GALFLGFLGAAGSTMGAWSQPKSKRKV ; SEQ ID NO: 449), hPP3 (KPKRKRRKKKGHWGR; SEQ ID NO: 450), PepNeg (SGTQEEY; SEQ ID NO: 451), CM18-TAT (KWKLFFKIGAVLKVLTTG; SEQ ID NO: 453), PTD4 (YARAAARQARA; SEQ ID NO: 454), or WaTx (MKYFTLALTLLFLLLINPCKDMNFAWAESSEKVERASPQQAQKCYEQCNVNVKVPFDQC YQMCSPLERS; SEQ ID NO: 455). For example, in some embodiments, the peptide can comprise an Arginine patch (Arg patch), for example, an RRRRRRRR (SEQ ID NO: 73), or a variant or fragment thereof, sequence can be appended to either the N-terminus or the C-terminus of a peptide. In some embodiments, the Arg patch comprises two or more Arg residues, or Arg_n wherein n is a whole number and can be 2, 3, 4, 5, 6, 7, 8, 9, or 10 (SEQ ID NO: 98). In other embodiments, the peptide can comprise a Tat peptide (Tat proteins are reviewed in Gump et al. TAT transduction: the molecular mechanism and therapeutic prospects. Trends Mol Med. 2007 Oct;13(10):443-8 and Harada et al. Antitumor protein therapy; application of the protein transduction domain to the development of a protein drug for cancer treatment. Breast Cancer. 2006;13(1):16-26). The Tat peptide can have a sequence of, for example, YGRKKRRQRRR (SEQ ID NO: 99), GRKKRRQRRR (SEQ ID NO: 100), or any modification, variant, or fragment thereof, can be appended to the N-terminus or C-terminus of any TfR-binding peptide of the present disclosure. In some embodiments, the Tat peptide sequence can be GRKKRRQRRRPQ (SEQ ID NO: 101), GRKKRRQRRR (SEQ ID NO: 100), or a fragment or variant thereof. In some embodiments, the Tat peptide can be appended to the N-terminus of any TfR-binding peptide of the present disclosure following an N-terminal GS dipeptide and preceding, for example, a GGGS (SEQ ID NO: 103) spacer. In some embodiments, a cell-penetrating tag peptides, such as any one of SEQ ID NO: 71 – SEQ ID NO: 96, SEQ ID NO: 98 – SEQ ID NO: 101, SEQ ID NO: 116 – SEQ ID NO: 124, SEQ ID NO: 126, or SEQ ID NO: 403 – SEQ ID NO: 455 can be appended to either the N-terminus or C-terminus of any peptide disclosed herein using a peptide linker such as G_xS_y (SEQ ID NO: 104) peptide linker, wherein x

and y can be any whole number, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. In other embodiments, a cell-penetrating peptide, such as a Tat peptide or an Arg patch, or any other moiety, can be appended to either the N-terminus or C-terminus of any peptide disclosed herein using a peptide linker such as G_xS_y (SEQ ID NO: 104) peptide linker, wherein x and y can be any whole number, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. In some embodiments, the peptide linker comprises $(GS)_x$ (SEQ ID NO: 105), wherein x can be any whole number, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. In some embodiments, the peptide linker comprises GGSSG (SEQ ID NO: 106), GGGGG (SEQ ID NO: 107), GSGSGSGS (SEQ ID NO: 108), GSGG (SEQ ID NO: 109), GGGGS (SEQ ID NO: 110), GGS (SEQ ID NO: 103), GGS (SEQ ID NO: 112), GGGSGGGSGGGS (SEQ ID NO: 113), or a variant or fragment thereof. Additionally, KKYKPYVPVTTN (SEQ ID NO: 114) from DkTx, and EPKSSDKTHT (SEQ ID NO: 115) from human IgG3 can be used as a peptide linker. In other embodiments, the tag peptide can be appended to the peptide at any amino acid residue. In further embodiments, the tag peptide can be appended to the peptide at any amino acid residue without interfering with TfR-binding activity. In some embodiments, the tag peptide is appended via conjugation, linking, or fusion techniques. In other embodiments, the Tat peptide can be appended to the peptide at any amino acid residue. In further embodiments, the Tat peptide can be appended to the peptide at any amino acid residue without interfering with TfR-binding activity. In some embodiments, the Tat peptide is appended via conjugation, linking, or fusion techniques to a TfR-binding peptide to obtain a TfR-binding Cell Penetrating Peptide fusion (CPP fusion).

[0330] Cell-penetrating peptides include, but are not limited to, short amphipathic or cationic short peptides with a positive net charge and are capable of penetrating cellular membrane and transferring a molecular or cargo either covalently or non-covalently attached to the peptides into a cell. Such cell-penetrating peptides can be synthesized or derived from known proteins, such as penetratin, Tat peptide, pVEC, or chimeric peptides, such as transportan, MPG, Pep-1, or synthetic peptides, such as polyarginines, MAP, and R_6W_3 .

[0331] In some embodiments, peptides can comprise at least one or more cell penetrating peptide sequences for improved cell penetration. For example, a cell penetrating peptide can include maurocaline (GDCLPHLKLCKENKDCCSKKCKRRGTNIEKRCR; SEQ ID NO: 116), imperatoxin (GDCLPHLKRCKADNDCCGKKCKRRGTNAEKRCR; SEQ ID NO: 117), hadrucalcin (SEKDCIKHLQRCRENKDCCSKKCSRRGTNPEKRCR; SEQ ID NO: 118), hemicalcin (GDCLPHLKLCKADKDCCSKKCKRRGTNPEKRCR; SEQ ID NO: 119),

opicalcin-1 (GDCLPHLKRCCKENNDCCSKKCKRRGTNPEKRCR; SEQ ID NO: 120), opicalcin-2 (GDCLPHLKRCCKENNDCCSKKCKRRGANPEKRCR; SEQ ID NO: 121, midkine (62-104) (CKYKFENWGACDGGTGTKVRQGTLLKKARYNAQCQETIRVTKPC; SEQ ID NO: 122), MCoTI-II (SGSDGGVCPKILKKCRDSDCPGACICRGNGYCG; SEQ ID NO: 123), or chlorotoxin (MCMPCFTTDHQMARCDDCCGGKGRGKCYGPQCLCR; SEQ ID NO: 124). In some embodiments, the cell penetrating peptide can have at least 80%, 90%, 95%, or 99% sequence identity with any sequence of SEQ ID NO: 71 – SEQ ID NO: 96, SEQ ID NO: 98 – SEQ ID NO: 101, SEQ ID NO: 116 – SEQ ID NO: 124, SEQ ID NO: 126, or SEQ ID NO: 403 – SEQ ID NO: 455.

[0332] In some embodiments, a peptide is conjugated to, linked to, or fused to one or more cell-penetrating peptides, such as arginine-rich, amphipathic and lysine-rich, and hydrophobic residues or peptides capable of penetrating plasma membrane or nucleus for in vivo delivery of a protein or macromolecular cargo. Conjugation or fusion can be direct or with a spacer in between (chemical or peptide-based). A spacer can be any peptide linker. For example, a spacer can be GGGSGGSGGGS (SEQ ID NO: 125), KKYKPYVPVTTN (SEQ ID NO: 114) from DkTx, EPKSSDKTHT (SEQ ID NO: 115) from human IgG3 or any variant or fragment thereof. In some embodiments, the cell penetrating peptide sequence can be appended to either the N-terminus or the C-terminus of a peptide. In some embodiments, the cell penetrating peptide can be appended to the N-terminus of any TfR-binding peptide of the present disclosure following an N-terminal GS dipeptide and preceding, for example, a GGS (SEQ ID NO: 103) spacer. In some embodiments, a cell-penetrating tag peptide, such as any one of SEQ ID NO: 71 – SEQ ID NO: 96, SEQ ID NO: 98 – SEQ ID NO: 101, SEQ ID NO: 116 – SEQ ID NO: 124, SEQ ID NO: 126, or SEQ ID NO: 403 – SEQ ID NO: 455, can be appended to either the N-terminus or C-terminus of any peptide disclosed herein using a peptide linker such as G_xS_y (SEQ ID NO: 104) peptide linker, wherein x and y can be any whole number, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. In some embodiments, the peptide linker comprises GGSSG (SEQ ID NO: 106), GGGGG (SEQ ID NO: 107), GSGSGSGS (SEQ ID NO: 108), GSGG (SEQ ID NO: 109), GGGGS (SEQ ID NO: 110), GGS (SEQ ID NO: 103), GGS (SEQ ID NO: 112), GGGSGGGSGGGS (SEQ ID NO: 113), GGGSGGGSGGGS (SEQ ID NO: 225), or a variant or fragment thereof. Additionally, KKYKPYVPVTTN (SEQ ID NO: 114) from DkTx, and EPKSSDKTHT (SEQ ID NO: 115) from human IgG3 can be used as a peptide linker. In other embodiments, the cell penetrating peptide can be appended to the peptide at any amino acid residue. In further embodiments, the

cell penetrating peptide can be appended to the peptide at any amino acid residue without interfering with TfR-binding activity. In some embodiments, the cell penetrating peptide is appended via conjugation, linking, or fusion techniques. In other embodiments, the cell penetrating peptide can be appended to the peptide at any amino acid residue.

[0333] In other embodiments, cell penetration can be increased by using high dosage of a peptide described here, such as up to 10 μ M, or 10 μ M or more of the peptide. In some cases, an Arg patch can be fused, conjugated to, linked to, or co-delivered with a peptide. Up 10 μ M, or 10 μ M or more Arg patch can be co-delivered with a peptide to facilitate cell penetration. Protein transfection agents can also be used to increase cell penetration of a peptide. In some embodiments, direct cytosolic expression of a peptide can be used. In other embodiments, physical disruption methods such as electroporation can be used to improve delivery of a peptide into a cell.

[0334] In some embodiments, cell penetrance of a peptide described herein can be improved. For example, the binding interface of a peptide described herein can be grafted on a scaffold that is known to be cell-penetrant, such as a calcine. Similarly, sequences that are known to be cell penetrant can be grafted onto peptides of this disclosure. Some non-limiting examples of calcines can be imperatoxin-A, maurocalcine, hemicalcin, opiclacine 1, opicalcin 2, and hadrucalcin. The scaffold can comprise at least 60%, 70%, 80%, 90%, 95%, or 98% with any one of SEQ ID NO: 116 – SEQ ID NO: 124. In some embodiments, the cell penetrance peptide can be calcines, modified calcines, derivatives of calcines, or fragments thereof, which can be used to increase cell penetration. Modified calcines, derivatives of calcines, or fragments can be screened for cell penetration activity such as activation of sarcoplasmic reticulum ryanodine receptors, activity on ryanodine-sensitive Ca^{2+} channels RyR1, Ryr2, or both, or as a selective agonist of the foregoing. Moreover, modified calcines can include substitution, addition or reduction of Lysine residues, or other charged residues, within a calcine in order to modify activity and optimize such calcine cell-penetration activity or activity on the RyR1 or RyR2 receptors. As an example, the six amino acid portion of helix 3 (MLICLF; SEQ ID NO: 126) can be transplanted onto a calcine or modified calcine scaffold to produce a bi-functional peptide that retains the cell penetration of the calcine with the novel TfR-binding function of the TfR-binding peptides. As another example, a peptide as described herein can have improved cell penetrating capabilities using cis-acting elements, including inclusion of K/R-rich sequences like TAT or octa-arginine, intra-helical

arginine patches, or fusion to larger fragments of proteins identified in cell penetration screening like penetratin or melittin.

[0335] In some embodiments, nuclear localization signals can be couple to, conjugated to, linked to, or fused to a peptide described herein to promote nuclear localization. In some embodiments, TfR-binding peptides are conjugated to, linked to, or fused to a nuclear localization signal, such as a four-residue sequence of K-K/R-X-K/R (SEQ ID NO: 129), wherein X can be any amino acid, or a variant thereof. In some embodiments, TfR-binding peptides are conjugated to, linked to, or fused to a nuclear localization signal as described in Lange et al, J Biol Chem. 2007 Feb 23;282(8):5101-5, such as PKKKRRV (SEQ ID NO: 130) or KRPAATKKAGQAKKKK (SEQ ID NO: 131). In some embodiments, a peptide described herein is conjugated to, linked to, or fused to a nuclear localization signal comprising KxRy (SEQ ID NO: 132), wherein x and y independently can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, such as KKRR (SEQ ID NO: 133), KKKRR (SEQ ID NO: 134), or KKKK (SEQ ID NO: 135). Other cell penetrating moieties can also be linked to, conjugated to, linked to, or fused to the peptides described herein, including, but not limited to, polycations, polyorganic acids, endosomal releasing polymers, poly(2-propylacrylic acid), poly(2-ethylacrylic acid), or any combination thereof.

[0336] In some embodiments, the peptides of the present disclosure (e.g., histidine-containing or histidine-enriched TfR-binding peptides) can have a high TfR binding affinity at physiological pH but a significantly reduced binding affinity at lower pH levels such as endosomal pH of 5.4. In some cases, the TfR-binding peptides of the present disclosure can be optimized for improved intra-vesicular (e.g., intra-endosomal) and/or intracellular delivery function while retaining high TfR binding capabilities. In some cases, histidine scans and comparative binding experiments can be performed to develop and screen for such peptides. In addition, some peptides can comprise (e.g., conjugated to, linked to, or fused to) a motif that facilitates low-pH endosomal escape of the peptide or peptide construct for enhanced delivery functions (e.g., intracellular delivery of a therapeutic agent). In some embodiments, an amino acid residue in a peptide of the present disclosure is substituted with a different amino acid residue to alter a pH-dependent binding affinity to TfR. The amino acid substitution may increase a binding affinity at low pH, increase a binding affinity at high pH, decrease a binding affinity at low pH, decrease a binding affinity at high pH, or a combination thereof.

[0337] In some embodiments, peptides of the present disclosure are capable of vesicular transcytosis across a cell layer or cell barrier such as the BBB. In some embodiments, the

peptides target and/or penetrate into a cell or a nucleus of a cell (e.g., a brain cell). Examples of cells or tissues that can be targeted include cells or tissues associated with a disease or condition such as cells or tissues of the CNS, brain cells, cancerous cells, and other cell types, wherein certain biological pathway can be dysregulated in said cells. Further examples of cells or tissues that can be targeted include cells or tissues of the spleen, liver, kidney, muscle, bone marrow, or skin. A cell that can be targeted with the peptides of the present disclosure can be a human cell, a mammalian cell, a human or mammalian cell line, a cancer cell line, a cell extracted from a subject, in vivo, or in vitro.

[0338] In some instances, a peptide as disclosed herein can contain only one lysine residue, or no lysine residues. In some instances, one or more or all of the lysine residues in the peptide are replaced with arginine residues. In some instances, one or more or all of the methionine residues in the peptide are replaced by leucine or isoleucine. One or more or all of the tryptophan residues in the peptide can be replaced by phenylalanine or tyrosine. In some instances, one or more or all of the asparagine residues in the peptide are replaced by glutamine. In some embodiments, one or more or all of the aspartic acid residues can be replaced by glutamic acid residues. In some instances, one or more or all of the lysine residues in the peptide are replaced by alanine or arginine. In some embodiments, the N-terminus of the peptide is blocked or protected, such as by an acetyl group or a *tert*-butyloxycarbonyl group. Alterly or in combination, the C-terminus of the peptide can be blocked or protected, such as by an amide group or by the formation of an ester (e.g., a butyl or a benzyl ester). In some embodiments, the peptide is modified by methylation on free amines. For example, full methylation is accomplished through the use of reductive methylation with formaldehyde and sodium cyanoborohydride.

[0339] In some embodiments, the dipeptide GS can be added as the first two N-terminal amino acids, as shown in SEQ ID NO: 1 – SEQ ID NO: 35, or SEQ ID NO: 136 – SEQ ID NO: 170, or such N-terminal dipeptide GS can be absent as shown in SEQ ID NO: 36 – SEQ ID NO: 70, or SEQ ID NO: 171 – SEQ ID NO: 205, or can be substituted by any other one or two amino acids. In some embodiments, the dipeptide GS is used as a linker or used to couple to a linker to form a peptide conjugate or fusion molecules such as a peptide construct. In some embodiments, the linker comprises a G_xS_y (SEQ ID NO: 104) peptide, wherein x and y independently are any whole number, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. In some embodiments, the peptide linker comprises $(GS)_x$ (SEQ ID NO: 105), wherein x can be any whole number, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10. In some embodiments, the peptide linker comprises GGSSG (SEQ ID NO: 106),

GGGGG (SEQ ID NO: 107), GSGSGSGS (SEQ ID NO: 108), GGGGS (SEQ ID NO: 110), GGGG (SEQ ID NO: 103), or a variant or fragment thereof.

[0340] In some embodiments of the present disclosure, a peptide or peptide construct as described herein comprises an amino acid sequence set forth in any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361. A peptide as disclosed herein can be a fragment comprising a contiguous fragment of any one of SEQ ID NO: 1– SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, or SEQ ID NO: 171 – SEQ ID NO: 202 that is at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36 at least 37, at least 38, at least 39, at least 40, at least 41, at least 42, at least 43, at least 44, at least 45, at least 46, at least 47, at least 48, at least 49, at least 50, at least 51, at least 52, at least 53, at least 54, at least 55, at least 56, at least 57, at least 58, at least 59, at least 60, at least 61, at least 62, at least 63, at least 64, at least 65 residues long, wherein the peptide fragment is selected from any portion of the peptide. In some embodiments, the peptide sequence is flanked by additional amino acids. One or more additional amino acids, for example, confer a particular *in vivo* charge, isoelectric point, chemical conjugation site, stability, or physiologic property to a peptide.

[0341] In some instances, the peptides as described herein that are capable of targeting and binding to a TfR comprise no more than 80 amino acids in length, or no more than 70, no more than 60, no more than 40, no more than 35, no more than 30, no more than 25, no more than 20, no more than 15, or no more than 10 amino acids in length.

[0342] In other embodiments, peptides can be conjugated to, linked to, or fused to a carrier or a molecule with targeting or homing function for a cell of interest or a target cell. In other embodiments, peptides can be conjugated to, linked to, or fused to a molecule that extends half-life or modifies the pharmacodynamic and/or pharmacokinetic properties of the peptides, or any combination thereof.

[0343] In some instances, a peptide comprises at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 positively charged residues, such as Arg or Lys, or any combination thereof. In some instances, one or more lysine

residues in the peptide are replaced with arginine residues. In some embodiments, peptides comprise one or more Arg patches. In some embodiments, an Arg patch is positioned in the N-terminus of a peptide. In other aspects, an Arg patch is positioned in the C-terminus of a peptide. In some embodiments, an Arg patch comprises 8 consecutive Arg residues (SEQ ID NO: 73). In some embodiments, 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, or 10 or more Arg or Lys residues are solvent exposed on a peptide. In some embodiments, an Arg patch can be two or more consecutive Arg residues. In some embodiments, an Arg patch comprises one or more Arg substituted with Lys.

[0344] The peptides of the present disclosure can further comprise neutral amino acid residues. In some embodiments, the peptide has 35 or fewer neutral amino acid residues. In other embodiments, the peptide has 81 or fewer neutral amino acid residues, 70 or fewer neutral amino acid residues, 60 or fewer neutral amino acid residues, 50 or fewer neutral amino acid residues, 40 or fewer neutral amino acid residues, 36 or fewer neutral amino acid residues, 33 or fewer neutral amino acid residues, 30 or fewer neutral amino acid residues, 25 or fewer neutral amino acid residues, or 10 or fewer neutral amino acid residues.

[0345] The peptides of the present disclosure can further comprise negative amino acid residues. In some embodiments the peptide has 6 or fewer negative amino acid residues, 5 or fewer negative amino acid residues, 4 or fewer negative amino acid residues, 3 or fewer negative amino acid residues, 2 or fewer negative amino acid residues, or 1 or fewer negative amino acid residues. While negative amino acid residues can be selected from any negatively charged amino acid residues, in some embodiments, the negative amino acid residues are either E, or D, or a combination of both E and D.

[0346] In some embodiments of the present disclosure, a three-dimensional or tertiary structure of a peptide is primarily comprised of beta-sheets and/or alpha-helix structures. In some embodiments, designed or engineered TfR-binding peptides of the present disclosure are small, compact peptides or polypeptides stabilized by intra-chain disulfide bonds (e.g., mediated by cysteines) to form cystine and a hydrophobic core. In some embodiments, engineered TfR-binding peptides have structures comprising helical bundles with at least one disulfide bridge between each of the alpha helices, thereby stabilizing the peptides. In other embodiments, the engineered TfR-binding peptides comprise structures with three alpha helices and three intra-chain disulfide bonds, one between each of the three alpha helices in the bundle of alpha helices.

[0347] In other embodiments, peptides can be conjugated to, linked to, or fused to a molecule (e.g., small molecule, peptide, or protein) with targeting or homing function for a cell of interest or a target protein located on the surface or inside said cell. In other embodiments, peptides can be conjugated to, linked to, or fused to a molecule that extends the plasma and/or biological half-life, or modifies the pharmacodynamic (e.g., enhanced binding to a target protein) and/or pharmacokinetic properties (e.g., rate and mode of clearance) of the peptides, or any combination thereof.

[0348] Generally, the nuclear magnetic resonance (NMR) solution structures or X-ray crystallography structures of related structural homologs can be used to inform mutational strategies that can improve the folding, stability, and manufacturability of the peptides as described herein, while maintaining a particular biological function (e.g., binding to TfR). These techniques can be used to predict the 3D pharmacophore of a group of structurally homologous scaffolds, as well as to predict possible graft regions of related proteins to create chimeras with improved properties (e.g., binding properties). For example, this strategy is used to identify critical amino acid positions and loops that are used to design peptides with improved TfR receptor binding and transcytosis properties, high expression, high stability *in vivo*, or any combination of these properties.

[0349] In some embodiments, a peptide capable of binding TfR and transcytosis across a cell membrane comprises a sequence with at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity with any one of the exemplary peptide sequences listed in **TABLE 1** (SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, and SEQ ID NO: 171 – SEQ ID NO: 202), or a functional fragment thereof. Two or more peptides can share a degree of sequence identity or homology and share similar properties *in vivo*. For instance, a peptide can share a degree of sequence identity or homology with any one of the peptides of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, or SEQ ID NO: 171 – SEQ ID NO: 202. In some embodiments, one or more peptides of the present disclosure have up to about 20% pairwise sequence identity or homology, up to about 25% pairwise sequence identity or homology, up to about 30% pairwise sequence identity or homology, up to about 35% pairwise sequence identity or homology, up to about 40% pairwise sequence identity or homology, up to about 45% pairwise sequence identity or homology, up to about 50% pairwise

sequence identity or homology, up to about 55% pairwise sequence identity or homology, up to about 60% pairwise sequence identity or homology, up to about 65% pairwise sequence identity or homology, up to about 70% pairwise sequence identity or homology, up to about 75% pairwise sequence identity or homology, up to about 80% pairwise sequence identity or homology, up to about 85% pairwise sequence identity or homology, up to about 90% pairwise sequence identity or homology, up to about 95% pairwise sequence identity or homology, up to about 96% pairwise sequence identity or homology, up to about 97% pairwise sequence identity or homology, up to about 98% pairwise sequence identity or homology, up to about 99% pairwise sequence identity or homology, up to about 99.5% pairwise sequence identity or homology, or up to about 99.9% pairwise sequence identity or homology. In some embodiments, one or more peptides of the disclosure have at least about 20% pairwise sequence identity or homology, at least about 25% pairwise sequence identity or homology, at least about 30% pairwise sequence identity or homology, at least about 35% pairwise sequence identity or homology, at least about 40% pairwise sequence identity or homology, at least about 45% pairwise sequence identity or homology, at least about 50% pairwise sequence identity or homology, at least about 55% pairwise sequence identity or homology, at least about 60% pairwise sequence identity or homology, at least about 65% pairwise sequence identity or homology, at least about 70% pairwise sequence identity or homology, at least about 75% pairwise sequence identity or homology, at least about 80% pairwise sequence identity or homology, at least about 85% pairwise sequence identity or homology, at least about 90% pairwise sequence identity or homology, at least about 95% pairwise sequence identity or homology, at least about 96% pairwise sequence identity or homology, at least about 97% pairwise sequence identity or homology, at least about 98% pairwise sequence identity or homology, at least about 99% pairwise sequence identity or homology, at least about 99.5% pairwise sequence identity or homology, at least about 99.9% pairwise sequence identity or homology with a second peptide.

[0350] In some embodiments, peptides that exhibit an improved TfR receptor binding show improved transcytosis function. In some cases, peptides that exhibit an improved TfR receptor binding show no or small changes in transcytosis function. In some cases, peptides that exhibit an improved TfR receptor binding show reduced transcytosis function. In some embodiments, the K_A and K_D values of a TfR-binding peptide can be modulated and optimized (e.g., via amino acid

substitutions) to provide an optimal ratio of TfR-binding affinity and efficient transcytosis function.

[0351] Various methods and software programs can be used to determine the homology between two or more peptides, such as NCBI BLAST, Clustal W, MAFFT, Clustal Omega, AlignMe, Praline, or another suitable method or algorithm. Pairwise sequence alignment can be used to identify regions of similarity that can indicate functional, structural and/or evolutionary relationships between two biological sequences (e.g., amino acid or nucleic acid sequences). In addition, multiple sequence alignment (MSA) is the alignment of three or more biological sequences. From the output of MSA applications, homology can be inferred and the evolutionary relationship between the sequences assessed. As used herein, “sequence homology” and “sequence identity” and “percent (%) sequence identity” and “percent (%) sequence homology” are used interchangeably to mean the sequence relatedness or variation, as appropriate, to a reference polynucleotide or amino acid sequence.

[0352] In some instances, the peptide or peptide construct is any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361, or a functional fragment thereof. In other embodiments, the peptide or peptide construct of the disclosure further comprises a peptide with 99%, 95%, 90%, 85%, or 80% sequence identity or homology to any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 or functional fragment thereof.

[0353] In other instances, the peptide or peptide construct can be a peptide that is homologous to any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361, or a functional fragment thereof. As further described herein, the term “homologous” can be used herein to denote peptides or peptide constructs having at least 70%, at least 80%, at least 90%, at least 95%, or greater than 95% sequence identity or homology to a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 or a functional fragment thereof.

[0354] In still other instances, the nucleic acid molecules that encode a peptide or peptide construct of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 can be identified by either a determination of the sequence identity or homology of the encoded peptide amino

acid sequence with the amino acid sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361, or by a nucleic acid hybridization assay. Such peptide variants or peptide construct variants of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 can be characterized as nucleic acid molecules (1) that remain hybridized with a nucleic acid molecule having the nucleotide sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 (or its complement) under highly stringent washing conditions, in which the wash stringency is equivalent to $0.1 \times - 0.2 \times \text{SSC}$ with 0.1% SDS at 50-65° C., and (2) that encode a peptide having at least 70%, at least 80%, at least 90%, at least 95% or greater than 95% sequence identity or homology to the amino acid sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361.

Knotted Peptides

[0355] In some embodiments, TfR-binding peptides of the present disclosure comprise one or more Cys, or one or more disulfide bond. In some embodiments, the TfR-binding peptides are derived from cystine-dense peptides (CDPs), knotted peptides, or hitchins. As used herein, the term “peptide” is considered to be interchangeable with the terms “knotted peptide”, “cystine-dense peptide”, “CDP”, and “hitchin”. (See e.g., Correnti et al. Screening, large-scale production, and structure-based classification for cystine-dense peptides. *Nat Struct Mol Biol.* 2018 Mar; 25(3): 270–278).

[0356] In some embodiments, the TfR-binding peptides of the present disclosure can bind TfR, thereby preventing TfR interactions such as interactions of TfR with other exogenous or endogenous ligands (e.g., Tf or homologs or fragments thereof). In some embodiments, the TfR-binding peptides of the present disclosure can bind TfR without impacting the binding of other exogenous or endogenous ligands (e.g., Tf or homologs or fragments thereof) with TfR. In some embodiments, TfR-binding peptides may be engineered peptides. An engineered peptide may be a peptide that is non-naturally occurring, artificial, isolated, synthetic, designed, or recombinantly expressed. In some embodiments, the TfR-binding peptides of the present disclosure comprise

one or more properties of CDPs, knotted peptides, or hitchhins, such as stability, resistance to proteolysis, resistance to reducing conditions, and/or ability to cross the blood brain barrier.

[0357] In some embodiments, CDPs or knotted peptides, including engineered, non-naturally occurring CDPs and those found in nature, can be conjugated to, linked to, or fused to the TfR-binding peptides of the present disclosure, such as those described in **TABLE 1**, to provide additional homing or targeting function to a target cell, such as a cancer cell, pancreatic cell, liver cell, colon cell, ovarian cell, breast cell, lung cell, or any combination thereof. In some embodiments, CDPs or knotted peptides, including engineered CDPs, isolated CDPs, and CDPs found in nature, can be conjugated to, linked to, or fused to the TfR-binding peptides of the present disclosure, such as those described in **TABLE 1**, to provide additional homing or targeting function to a target cell, such as a cancer cell, pancreatic cell, liver cell, colon cell, ovarian cell, breast cell, lung cell, or any combination thereof. An engineered peptide may be a peptide that is non-naturally occurring, artificial, synthetic, designed, or recombinantly expressed. In some embodiments, a TfR-binding peptide of the present disclosure enables TfR-mediated transcytosis and/or cellular endocytosis, and the additional CDP or knotted peptide that is conjugated to, linked to, or fused to TfR-binding peptide can selectively target an enzyme or other protein of interest in a cell associated with a disease or condition. In some cases, the cell is a cancer cell. Cancers can include breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, ovarian cancer, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma, lung cancer, bone marrow cell cancers, or skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, a neuroblastoma, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, and diffuse intrinsic pontine glioma (DIPG), or a CMYC-overexpressing cancer. In some cases, other CDP or knotted peptides (e.g., those found in nature) are conjugated to, linked to, or fused to TfR-binding peptides and are capable of localizing TfR-binding peptides across the blood brain barrier to deliver TfR-binding peptides to target cells in the central nervous system.

[0358] CDPs (e.g., knotted peptides) are a class of peptides, usually ranging from about 11 to about 81 amino acids in length that are often folded into a compact structure. Knotted peptides are typically assembled into a complex tertiary structure that is characterized by a number of intramolecular disulfide crosslinks and may contain beta strands, alpha helices, and other

secondary structures. The presence of the disulfide bonds gives knotted peptides remarkable environmental stability, allowing them to withstand extremes of temperature and pH and to resist the proteolytic enzymes of the blood stream. The presence of a disulfide knot may provide resistance to reduction by reducing agents. The rigidity of knotted peptides also allows them to bind to targets without paying the “entropic penalty” that a floppy peptide accrues upon binding a target. For example, binding is adversely affected by the loss of entropy that occurs when a peptide binds a target to form a complex. Therefore, “entropic penalty” is the adverse effect on binding, and the greater the entropic loss that occurs upon this binding, the greater the “entropic penalty.” Furthermore, unbound molecules that are flexible lose more entropy when forming a complex than molecules that are rigidly structured, because of the loss of flexibility when bound up in a complex. However, rigidity in the unbound molecule also generally increases specificity by limiting the number of complexes that molecule can form. The peptides can bind targets with antibody-like affinity, or with nanomolar or picomolar affinity. A wider examination of the sequence structure and sequence identity or homology of knotted peptides reveals that they have arisen by convergent evolution in all kinds of animals and plants. In animals, they are often found in venoms, for example, the venoms of spiders and scorpions and have been implicated in the modulation of ion channels. The knotted proteins of plants can inhibit the proteolytic enzymes of animals or have antimicrobial activity, suggesting that knotted peptides can function in molecular defense systems found in plants.

[0359] The peptides of the present disclosure comprise cysteine amino acid residues. In some embodiments, the peptide has at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 cysteine amino acid residues. In some embodiments, the peptide has at least 8 cysteine amino acid residues. In other embodiments, the peptide has at least 10 cysteine amino acid residues, at least 12 cysteine amino acid residues, at least 14 cysteine amino acid residues or at least 16 cysteine amino acid residues.

[0360] A knotted peptide can comprise disulfide bridges. A knotted peptide can be a peptide wherein 5% or more of the residues are cysteines forming intramolecular disulfide bonds. A disulfide-linked peptide can be a drug scaffold. In some embodiments, the disulfide bridges form a knot. A disulfide bridge can be formed between cysteine residues, for example, between cysteines 1 and 4, 2 and 5, or, 3 and 6. In some embodiments, one disulfide bridge passes through a loop formed by the other two disulfide bridges, for example, to form the knot. In other embodiments, the disulfide bridges can be formed between any two cysteine residues.

[0361] The present disclosure further includes peptide scaffolds that, e.g., can be used as a starting point for generating additional peptides. In some embodiments, these scaffolds can be derived from a variety of knotted peptides (such as CDPs or knotted peptides). In certain embodiments, CDPs (e.g., knotted peptides) are assembled into a complex tertiary structure that is characterized by a number of intramolecular disulfide crosslinks, and optionally contain beta strands and other secondary structures such as an alpha helix. For example, CDPs (e.g., knotted peptides) include, in some embodiments, small disulfide-rich proteins characterized by a disulfide through disulfide knot. This knot can be, e.g., obtained when one disulfide bridge crosses the macrocycle formed by two other disulfides and the interconnecting backbone. In some embodiments, the knotted peptides can include growth factor cysteine knots or inhibitor cysteine knots. Other possible peptide structures include peptide having two parallel helices linked by two disulfide bridges without β -sheets (e.g., hefutoxin).

[0362] Some peptides of the present disclosure can comprise at least one amino acid residue in an L configuration. A peptide can comprise at least one amino acid residue in D configuration. In some embodiments, a peptide is 15-75 amino acid residues long. In other embodiments, a peptide is 11-55 amino acid residues long. In still other embodiments, a peptide is 11-65 amino acid residues long. In further embodiments, a peptide is at least 20 amino acid residues long.

[0363] Some CDPs (e.g., knotted peptides) can be derived or isolated from a class of proteins known to be present or associated with toxins or venoms. In some cases, the peptide can be derived from toxins or venoms associated with scorpions or spiders. The peptide can be derived from venoms and toxins of spiders and scorpions of various genus and species. For example, the peptide can be derived from a venom or toxin of the *Leiurus quinquestriatus hebraeus*, *Buthus occitanus tunetanus*, *Hottentotta judaicus*, *Mesobuthus eupeus*, *Buthus occitanus israelis*, *Hadrurus gertschi*, *Androctonus australis*, *Centruroides noxius*, *Heterometrus laoticus*, *Opisthophthalmus carinatus*, *Haplopelma schmidtii*, *Isometrus maculatus*, *Haplopelma huwenum*, *Haplopelma hainanum*, *Haplopelma schmidtii*, *Agelenopsis aperta*, *Hadronyche versuta*, *Selenocosmia huwena*, *Heteropoda venatoria*, *Grammostola rosea*, *Ornithoctonus huwena*, *Hadronyche versuta*, *Atrax robustus*, *Agelenopsis aperta*, *Psalmopoeus cambridgei*, *Hadronyche infensa*, *Paracoelotes luctosus*, and *Chilobrachys jingzhaoor* another suitable genus or species of scorpion or spider. In some cases, a peptide can be derived from a *Buthus martensii* Karsh (scorpion) toxin.

Sequence Identity and Homology

[0364] Percent (%) sequence identity or homology is determined by conventional methods. (See e.g., Altschul et al. (1986), *Bull. Math. Bio.* 48:603 (1986), and Henikoff and Henikoff (1992), *Proc. Natl. Acad. Sci. USA* 89:10915). Briefly, two amino acid sequences can be aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the “BLOSUM62” scoring matrix of Henikoff and Henikoff (Id.). The sequence identity or homology is then calculated as: $(\text{[Total number of identical matches]} / \text{[length of the longer sequence plus the number of gaps introduced into the longer sequence in order to align the two sequences]}) \times 100$.

[0365] Additionally, there are several established algorithms available to align two amino acid sequences. For example, the “FASTA” similarity search algorithm of Pearson and Lipman can be a suitable protein alignment method for examining the level of sequence identity or homology shared by an amino acid sequence of a peptide disclosed herein and the amino acid sequence of a peptide variant. The FASTA algorithm is described, for example, by Pearson and Lipman, *Proc. Nat'l Acad. Sci. USA* 85:2444 (1988), and by Pearson, *Meth. Enzymol.* 183:63 (1990). Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO: 1) and a test sequence that has either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are “trimmed” to include only those residues that contribute to the highest score. If there are several regions with scores greater than the “cutoff” value (calculated by a predetermined formula based upon the length of the sequence and the ktup value), then the trimmed initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman and Wunsch, *J. Mol. Biol.* 48:444 (1970); Sellers, *Siam J. Appl. Math.* 26:787 (1974)), which allows for amino acid insertions and deletions. For example, illustrative parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file (“SMATRIX”), as explained in Appendix 2 of Pearson, *Meth. Enzymol.* 183:63 (1990).

[0366] FASTA can also be used to determine the sequence identity or homology of nucleic acid sequences or molecules using a ratio as disclosed above. For nucleic acid sequence comparisons, the ktup value can range between one to six, preferably from three to six, most preferably three, with other parameters set as described herein.

[0367] Some examples of common amino acids that are a “conservative amino acid substitution” are illustrated by a substitution among amino acids within each of the following groups: (1) glycine, alanine, valine, leucine, and isoleucine, (2) phenylalanine, tyrosine, and tryptophan, (3) serine and threonine, (4) aspartate and glutamate, (5) glutamine and asparagine, and (6) lysine, arginine and histidine. The BLOSUM62 table is an amino acid substitution matrix derived from about 2,000 local multiple alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff and Henikoff, Proc. Nat'l Acad. Sci. USA 89:10915 (1992)). Accordingly, the BLOSUM62 substitution frequencies can be used to define conservative amino acid substitutions that can be introduced into the amino acid sequences of the present invention. Although it is possible to design amino acid substitutions based solely upon chemical properties (as discussed above), the language “conservative amino acid substitution” preferably refers to a substitution represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. According to this system, preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 1 (e.g., 1, 2 or 3), while more preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 2 (e.g., 2 or 3).

[0368] Determination of amino acid residues that are within regions or domains that are critical to maintaining structural integrity can be determined. Within these regions one can determine specific residues that can be more or less tolerant of change and maintain the overall tertiary structure of the molecule. Methods for analyzing sequence structure include, but are not limited to, alignment of multiple sequences with high amino acid or nucleotide identity or homology and computer analysis using available software (e.g., the Insight II.RTM. viewer and homology modeling tools; MSI, San Diego, Calif.), secondary structure propensities, binary patterns, complementary packing and buried polar interactions (Barton, G.J., Current Opin. Struct. Biol. 5:372-6 (1995) and Cordes, M.H. et al., Current Opin. Struct. Biol. 6:3-10 (1996)). In general, when designing modifications to molecules or identifying specific fragments, determination of structure can typically be accompanied by evaluating activity of modified molecules.

Physicochemical Properties of Peptides

[0369] In some embodiments, the TfR-binding peptides of the present disclosure can comprise a wide range of physicochemical properties such as molecular size and structure, pH, isoelectric point, and overall molecular net charge. These parameters can have an effect on the peptides ability to bind TfR, promote transcytosis, and transport of cargo molecules across cell barrier such as the BBB.

[0370] A peptide of the present disclosure can comprise at least one amino acid residue in D configuration. In some embodiments, a peptide is about 5-100 amino acid residues long. In some embodiments, a peptide is about 10-90 amino acid residues long. In some embodiments, a peptide is about 15-80 amino acid residues long. In some embodiments, a peptide is about 15-75 amino acid residues long. In some embodiments, a peptide is about 15-70 amino acid residues long. In some embodiments, a peptide is about 20-65 amino acid residues long. In some embodiments, a peptide is about 20-60 amino acid residues long. In some embodiments, a peptide is about 25-55 amino acid residues long. In some embodiments, a peptide is about 25-50 amino acid residues long. In some embodiments, a peptide is about 25-40 amino acid residues long. In some embodiments, a peptide is about 11-35 amino acid residues long. In some embodiments, a peptide is about 10-25 amino acid residues long.

[0371] In some embodiments, a peptide is at least 5 amino acid residues long. In some embodiments, a peptide is at least 10 amino acid residues long. In some embodiments, a peptide is at least 15 amino acid residues long. In some embodiments, a peptide is at least 20 amino acid residues long. In some embodiments, a peptide is at least 25 amino acid residues long. In some embodiments, a peptide is at least 30 amino acid residues long. In some embodiments, a peptide is at least 35 amino acid residues long. In some embodiments, a peptide is at least 40 amino acid residues long. In some embodiments, a peptide is at least 45 amino acid residues long. In some embodiments, a peptide is at least 50 amino acid residues long. In some embodiments, a peptide is at least 55 amino acid residues long. In some embodiments, a peptide is at least 60 amino acid residues long. In some embodiments, a peptide is at least 65 amino acid residues long. In some embodiments, a peptide is at least 70 amino acid residues long. In some embodiments, a peptide is at least 75 amino acid residues long.

[0372] In some embodiments, an amino acid sequence of a peptide as described herein comprises at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least

27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36, at least 37, at least 38, at least 39, at least 40, at least 41, at least 42, at least 43, at least 44, at least 45, at least 46, at least 47, at least 48, at least 49, at least 50, at least 51, at least 52, at least 53, at least 54, at least 55, at least 56, at least 57, at least 58 residues, at least 59, at least 60, at least 61, at least 62, at least 63, at least 64, at least 65, at least 66, at least 67, at least 68, at least 69, at least 70, at least 71, at least 72, at least 73, at least 74, at least 75, at least 76, at least 77, at least 78, at least 79, at least 80, or at least 81 amino acid residues.

[0373] In some embodiments of the present disclosure, a three-dimensional or tertiary structure of a peptide is primarily comprised of beta-sheets and/or alpha-helix structures. In some embodiments, designed or engineered TfR-binding peptides of the present disclosure are small, compact peptides or polypeptides stabilized by intra-chain disulfide bonds (e.g., mediated by cysteines) and a hydrophobic core. In some embodiments, engineered TfR-binding peptides have structures comprising helical bundles with at least one disulfide bridge between each of the alpha helices, thereby stabilizing the peptides. In other embodiments, the engineered TfR-binding peptides comprise structures with three alpha helices and three intra-chain disulfide bonds, one between each of the three alpha helices in the bundle of alpha helices.

[0374] At physiological pH, peptides as described herein can have an overall molecular net charge, for example, of -5, -4, -3, -2, -1, 0, +1, +2, +3, +4, or +5. When the net charge is zero, the peptide can be uncharged or zwitterionic. In some embodiments, a peptide contains one or more disulfide bonds and has a positive net charge at physiological pH where the net charge can be +0.5 or less than +0.5, +1 or less than +1, +1.5 or less than +1.5, +2 or less than +2, +2.5 or less than +2.5, +3 or less than +3, +3.5 or less than +3.5, +4 or less than +4, +4.5 or less than +4.5, +5 or less than +5, +5.5 or less than +5.5, +6 or less than +6, +6.5 or less than +6.5, +7 or less than +7, +7.5 or less than +7.5, +8 or less than +8, +8.5 or less than +8.5, +9 or less than +9.5, +10 or less than +10. In some embodiments, a peptide has a negative net charge at physiological pH where the net charge can be -0.5 or less than -0.5, -1 or less than -1, -1.5 or less than -1.5, -2 or less than -2, -2.5 or less than -2.5, -3 or less than -3, -3.5 or less than -3.5, -4 or less than -4, -4.5 or less than -4.5, -5 or less than -5, -5.5 or less than -5.5, -6 or less than -6, -6.5 or less than -6.5, -7 or less than -7, -7.5 or less than -7.5, -8 or less than -8, -8.5 or less than -8.5, -9 or less than -9.5, -10 or less than -10.

[0375] In some embodiments, peptides of the present disclosure can have an isoelectric point (pI) value from 3 and 10. In other embodiments, peptides of the present disclosure can have a pI

value from 4.3 and 8.9. In some embodiments, peptides of the present disclosure can have a pI value from 3-4. In some embodiments, peptides of the present disclosure can have a pI value from 3-5. In some embodiments, peptides of the present disclosure can have a pI value from 3-6. In some embodiments, peptides of the present disclosure can have a pI value from 3-7. In some embodiments, peptides of the present disclosure can have a pI value from 3-8. In some embodiments, peptides of the present disclosure can have a pI value from 3-9. In some embodiments, peptides of the present disclosure can have a pI value from 4-5. In some embodiments, peptides of the present disclosure can have a pI value from 4-6. In some embodiments, peptides of the present disclosure can have a pI value from 4-7. In some embodiments, peptides of the present disclosure can have a pI value from 4-8. In some embodiments, peptides of the present disclosure can have a pI value from 4-9. In some embodiments, peptides of the present disclosure can have a pI value from 4-10. In some embodiments, peptides of the present disclosure can have a pI value from 5-6. In some embodiments, peptides of the present disclosure can have a pI value from 5-7. In some embodiments, peptides of the present disclosure can have a pI value from 5-8. In some embodiments, peptides of the present disclosure can have a pI value from 5-9. In some embodiments, peptides of the present disclosure can have a pI value from 5-10. In some embodiments, peptides of the present disclosure can have a pI value from 6-7. In some embodiments, peptides of the present disclosure can have a pI value from 6-8. In some embodiments, peptides of the present disclosure can have a pI value from 6-9. In some embodiments, peptides of the present disclosure can have a pI value from 6-10. In some embodiments, peptides of the present disclosure can have a pI value from 7-8. In some embodiments, peptides of the present disclosure can have a pI value from 7-9. In some embodiments, peptides of the present disclosure can have a pI value from 7-10. In some embodiments, peptides of the present disclosure can have a pI value from 8-9. In some embodiments, peptides of the present disclosure can have a pI value from 8-10. In some embodiments, peptides of the present disclosure can have a pI value from 9-10.

[0376] In some cases, the engineering of one or more mutations within a peptide of the present disclosure (e.g., a TfR-binding peptide) yields a peptide with an altered isoelectric point, charge, surface charge, or rheology at physiological pH. Such engineering of a mutation to a peptide that can be derived from a scorpion or spider complex can change the net charge of the peptide, for example, by decreasing the net charge by 1, 2, 3, 4, or 5, or by increasing the net charge by 1, 2,

3, 4, or 5. In such cases, the engineered mutation can facilitate the ability of the peptide to bind a target protein, promote transcytosis, and penetrate a cell, an endosome, or the nucleus. Suitable amino acid modifications for improving the rheology and potency of a peptide can include conservative or non-conservative mutations.

[0377] A peptide can comprise at most 1 amino acid mutation, at most 2 amino acid mutations, at most 3 amino acid mutations, at most 4 amino acid mutations, at most 5 amino acid mutations, at most 6 amino acid mutations, at most 7 amino acid mutations, at most 8 amino acid mutations, at most 9 amino acid mutations, at most 10 amino acid mutations, or another suitable number as compared to the sequence of the venom or toxin component that the peptide is derived from. In other embodiments, a peptide, or a functional fragment thereof, comprises at least 1 amino acid mutation, at least 2 amino acid mutations, at least 3 amino acid mutations, at least 4 amino acid mutations, at least 5 amino acid mutations, at least 6 amino acid mutations, at least 7 amino acid mutations, at least 8 amino acid mutations, at least 9 amino acid mutations, at least 10 amino acid mutations, or another suitable number as compared to the sequence of the venom or toxin component that the peptide is derived from. In some embodiments, mutations can be engineered within a peptide to provide a peptide that has a desired charge or stability at physiological pH.

[0378] Generally, the NMR solution structures, the X-ray crystal structures, as well as the primary structure sequence alignment of related structural peptide or protein homologs or *in silico* design can be used to generate mutational strategies that can improve the folding, stability, and/or manufacturability, while maintaining a particular biological function (e.g., TfR affinity/binding). A general strategy for producing homologs or *in silico* designed peptides or polypeptides can include identification of a charged surface patch or conserved residues of a protein, mutation of critical amino acid positions and loops, followed by *in vitro* and *in vivo* testing of the peptides. The overall peptide optimization process can be of iterative nature to the extent that, for example, information obtained during *in vitro* or *in vivo* testing is used for the design of the next generation of peptides. Hence, the herein disclosed methods can be used to design peptides with improved properties or to correct deleterious mutations that complicate folding and manufacturability. Key amino acid positions and loops can be retained while other residues in the peptide sequences can be mutated to improve, change, remove, or otherwise modify function, such as binding, transcytosis, or the ability to penetrate a cell, endosome, or nucleus in a cell, homing, or another activity of the peptide.

[0379] The present disclosure also encompasses multimers of the various peptides described herein. Examples of multimers include dimers, trimers, tetramers, pentamers, hexamers, heptamers, and so on. A multimer may be a homomer formed from a plurality of identical subunits or a heteromer formed from a plurality of different subunits. In some embodiments, a peptide of the present disclosure is arranged in a multimeric structure with at least one other peptide, or two, three, four, five, six, seven, eight, nine, ten, or more other peptides. In certain embodiments, the peptides of a multimeric structure each have the same sequence. In other embodiments, one or more or all of the peptides of a multimeric structure have different sequences.

[0380] In some embodiments, the present disclosure provides peptide scaffolds that can be used as a starting point for generating additional, next-generation peptides with more specific or improved properties. In some embodiments, these scaffolds are derived from a variety of CDPs or knotted peptides. Some suitable peptides for scaffolds can include, but are not limited to, chlorotoxin, brazzein, circulin, stecrisp, hanatoxin, midkine, hefutoxin, potato carboxypeptidase inhibitor, bubble protein, attractin, α -GI, α -GID, μ -PIIIA, ω -MVIIA, ω -CVID, χ -MrIA, ρ -TIA, conantokin G, contulakin G, GsMTx4, margatoxin, shK, toxin K, chymotrypsin inhibitor (CTI), and EGF epiregulin core. In some embodiments, the peptide sequence is flanked by additional amino acids. One or more additional amino acids can confer a desired *in vivo* charge, isoelectric point, chemical conjugation site, stability, or physiologic property to a peptide.

Chemical Modification of Peptides

[0381] A peptide can be chemically modified one or more of a variety of ways. In some embodiments, the peptide can be mutated to add function, delete function, or modify the *in vivo* behavior. For example, in some embodiments, peptides of the presenting disclosure may be chemically modified with a molecule that would lead to proteasomal degradation of TfR (e.g., ubiquitin ligase engaging conjugate or fusion or cereblon-binding molecule). One or more loops between the disulfide linkages can be modified or replaced to include active elements from other peptides (such as described in Moore and Cochran, *Methods in Enzymology*, 503, p. 223-251, 2012). Amino acids can also be mutated, such as to increase half-life, modify, add or delete binding behavior *in vivo*, add new targeting function, modify surface charge and hydrophobicity, or allow conjugation sites. N-methylation is one example of methylation that can occur in a peptide of the disclosure. In some embodiments, the peptide is modified by methylation on free

amines. For example, full methylation may be accomplished through the use of reductive methylation with formaldehyde and sodium cyanoborohydride.

[0382] The peptides can be modified to add function, such as to graft loops or sequences from other proteins or peptides onto peptides of this disclosure. Likewise, domains, loops, or sequences from this disclosure can be grafted onto other peptides or proteins such as antibodies that have additional function.

[0383] A chemical modification can, for instance, extend the half-life of a peptide or change the biodistribution or pharmacokinetic profile. A chemical modification can comprise a polymer, a polyether, polyethylene glycol, a biopolymer, a polyamino acid, a fatty acid, a dendrimer, an Fc region, a simple saturated carbon chain such as palmitate or myristolate, or albumin. A polyamino acid can include, for example, a poly amino acid sequence with repeated single amino acids (e.g., poly glycine), and a poly amino acid sequence with mixed poly amino acid sequences (e.g., gly-ala-gly-ala) that may or may not follow a pattern, or any combination of the foregoing.

[0384] The peptides of the present disclosure can be modified such that the modification increases the stability and/or the half-life of the peptides. The attachment of a hydrophobic moiety, such as to the N-terminus, the C-terminus, or an internal amino acid, can be used to extend half-life of a peptide of the present disclosure. The peptides can also be modified to increase or decrease the gut permeability or cellular permeability of the peptide. In some cases, the peptides of the present disclosure show high accumulation in glandular cells of the intestine, demonstrating applicability in the treatment and-or prevention of diseases or conditions of the intestines, such as Crohn's disease or more generally inflammatory bowel diseases. The peptide of the present disclosure can include post-translational modifications (e.g., methylation and/or amidation and/or glycosylation), which can affect, e.g., serum half-life. In some embodiments, simple carbon chains (e.g., by myristoylation and/or palmitylation) can be conjugated to, linked to, the fusion proteins or peptides. The simple carbon chains can render the fusion proteins or peptides easily separable from the unconjugated material. For example, methods that can be used to separate the fusion proteins or peptides from the unconjugated material include, but are not limited to, solvent extraction and reverse phase chromatography. Lipophilic moieties can extend half-life through reversible binding to serum albumin. Conjugated moieties can, e.g., be lipophilic moieties that extend half-life of the peptides through reversible binding to serum albumin. In some embodiments, the lipophilic moiety can be cholesterol or a cholesterol derivative including cholestenes, cholestanes, cholestadienes and oxysterols. In some

embodiments, the peptides can be conjugated to, linked to, myristic acid (tetradecanoic acid) or a derivative thereof. In other embodiments, the peptides of the present disclosure can be coupled (e.g., conjugated, linked, or fused) to a half-life modifying agent. Examples of half-life modifying agents can include, but is not limited to: a polymer, a polyethylene glycol (PEG), a hydroxyethyl starch, polyvinyl alcohol, a water soluble polymer, a zwitterionic water soluble polymer, a water soluble poly(amino acid), a water soluble polymer of proline, alanine and serine, a water soluble polymer containing glycine, glutamic acid, and serine, an Fc region, a fatty acid, palmitic acid, or a molecule that binds to albumin. In some embodiments, the half-life modifying agent may be a serum albumin binding peptide, for example SA21 (SEQ ID NO: 376, RLIEDICLPRWGCLWEDD). In some embodiments, a SA21 peptide may be conjugated or fused to the CDPs of the present disclosure (e.g., any of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358). A SA21 fusion peptide may include the SA21 TfR-binding peptide constructs disclosed herein (e.g., SEQ ID NO: 373 or SEQ ID NO: 375). The SA21 peptide may comprise a linker sequence for conjugation to, or fusion between, one or more peptides (e.g., SEQ ID NO: 377, GGGGSGGGGSRLEDICLPRWGCLWEDDGGGGSGGGGS). Exemplary SA21 peptides, fusion peptides, and linkers are provided in **TABLE 2**. A control SA21 fusion peptide may comprise a control peptide fused to SA21 (e.g., SEQ ID NO 372 (GSRLEDICLPRWGCLWEDDGGGGSGGGGSKCLPPGKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 374 (RLIEDICLPRWGCLWEDDGGGGSGGGGSKCLPPGKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 456 (GSRLEDICLPRWGCLWEDDGGGGSGGGGSVRIPVSCKHSGQCLKPCKDAGMRFGKC MNGKCDCTPK), or SEQ ID NO: 457 (RLIEDICLPRWGCLWEDDGGGGSGGGGSVRIPVSCKHSGQCLKPCKDAGMRFGKCMN GKCDCTPK)). Additionally, conjugation of the peptide to a near infrared dye, such as Cy5.5, or to an albumin binder such as Albu-tag can extend serum half-life of any peptide as described herein. In some embodiments, immunogenicity is reduced by using minimal non-human protein sequences to extend serum half-life of the peptide.

TABLE 2 – Exemplary TfR-Binding peptide constructs: TfR-binding peptides fused with Serum Albumin binding peptides for half-life extension

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 373	GSRLIEDICLPRWGCLWEDDGGGGSGGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 375	RLIEDICLPRWGCLWEDDGGGGSGGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ

[0385] In some embodiments, the first two N-terminal amino acids (GS) of SEQ ID NO: 1 – SEQ ID NO: 35 or SEQ ID NO: 136 – SEQ ID NO: 170 serve as a spacer or linker in order to facilitate conjugation or fusion to another molecule, as well as to facilitate cleavage of the peptide from such conjugated to, linked to, or fused molecules. In some embodiments, the fusion proteins or peptides of the present disclosure can be conjugated to, linked to, or fused to other moieties that, e.g., can modify or effect changes to the properties of the peptides.

Peptide-Active Agent Conjugates (also referred to as “Peptide Constructs”)

[0386] In some embodiments, the peptides of the present disclosure can themselves be used to bind TfR. In other embodiments, peptides of the present disclosure can also be used to deliver another active agent. Peptides according to the present disclosure can be conjugated to, linked to, or fused to an agent for use in the treatment of tumors and cancers. For example, in certain embodiments, the peptides described herein are fused to another molecule, such as an active agent that provides an additional functional capability. A peptide can be fused with an active agent through expression of a vector containing the sequence of the peptide with the sequence of the active agent. In various embodiments, the sequence of the peptide and the sequence of the active agent can be expressed from the same Open Reading Frame (ORF). In various embodiments, the sequence of the peptide and the sequence of the active agent can comprise a contiguous sequence. The peptide and the active agent can each retain similar functional capabilities in the peptide construct compared with their functional capabilities when expressed separately. In certain embodiments, examples of active agents can include other peptides.

[0387] In certain embodiments, examples of active agents include other peptides such as neurotensin peptide. Neurotensin is a 13 amino acid neuropeptide that can be involved in the regulation of luteinizing hormone and prolactin release, and can interact with the dopaminergic

system, but does not cross the blood brain barrier and is also rapidly metabolized by peptidases (Wang, et al., *Curr. Pharm. Des.* 21(7); 840-848 (2015)).

[0388] Neurotensin is a 13-amino acid peptide found in the central nervous system and the gastrointestinal tract and plays a role in a wide range of physiologic and pathologic processes. Various activities of neurotensin and potential therapeutic roles of neurotensin agonists are described in the following: Mustain, Rychahou, and Evers, *Curr Opin Endocrinol Diabetes Obes* (2011) and Boules, Li, Smith, Fredrickson, and Richelson, *Front Endocrinol (Lausanne)* (2013). NT plays a role in gut motility, modulation of the cardiovascular system, naloxone-independent antinociception, hypothermia, controls of anterior pituitary hormone secretion, muscle relaxation, central blood pressure, and inflammation. NT mediates its effects through three receptors NTS1, NTS2, and NTS3. NTS1 is a high affinity receptor and is expressed broadly throughout the CNS, including medial septal nucleus, nucleus basalis magnocellularis, suprachiasmatic nucleus, SN, and VTA, small dorsal root ganglion neurons of the spinal cord, in both neurons and glial cells. NTS2 is localized mainly in the olfactory system, the cerebral and cerebellar cortices, the hippocampal formation, and selective hypothalamic nuclei, VTA, and SN. NTS3 is also expressed at various locations throughout the brain. NT is involved in modulating dopamine neurotransmission, may play a role in the serotonergic system including antinociception, sleep-wake cycle, and stress, a role in glutamate release and may have antipsychotic effects. NT is involved in neuroendocrine regulation, including paracrine and autocrine roles, CRH, GnRH, GHRH, prolactin, ACTH, gonadotropic hormones, growth hormones, prolactin, thyroid hormone modulation, as well as gut motility, intestinal inflammation, lipid metabolism, appetite, and food intake. NT has been implicated in the pathophysiology of several CNS disorders such as schizophrenia, drug abuse, Parkinson's disease (PD), pain, central control of blood pressure, eating disorders, stroke, Alzheimer's, as well as, cancer and inflammation and other disorders of endocrine function. Disruption in the normal mechanisms can lead to diseases ranging from schizophrenia to colorectal cancer. By selective targeting or blockade of specific neurotensin receptors, investigators have identified potential drugs for use in the treatment of schizophrenia, alcoholism, chronic pain, or cancer. NT can provide opioid-independent analgesia and can treat thermal, visceral, and persistent inflammatory pain, somatic pain and visceral pain, and plays a role in stress-induced antinociception. NT exerts a mu-opioid-independent, anti-nociceptive effects. NT agonists can block effects of psychostimulants such as cocaine and amphetamine and nicotine. Plasma NT levels are higher in Parkinson's disease patients, and administration of NT

can reduce muscular rigidity and tremors. NT can also reduce body temperature and reduce blood pressure.

[0389] **TABLE 3** discloses exemplary neurotensin (NT) peptide variants (Boules, *et al.*, Diverse roles of neurotensin agonists in the central nervous system, *Frontiers in Endocrinology*, 2013). Any peptides of the present disclosure (e.g., any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358) can be fused or otherwise conjugated to any of the NT peptide variants disclosed herein (e.g., any of the NT peptide variants disclosed in **TABLE 3**), or fragments thereof. In some embodiments, the peptides of the disclosure are fused or otherwise conjugated a neurotensin (NT) peptide variant comprising a at least 53%, at least 60%, at least 69%, at least 75%, at least 77%, at least 85%, least 90%, at least 92% sequence identity with SEQ ID NO: 350. In other embodiments, the peptides of the disclosure are fused or otherwise conjugated a neurotensin (NT) peptide variant comprising the sequence ELYENKP (SEQ ID NO: 370) or ELYENKP-X₁-X₂-P-X₃-X₄-L (SEQ ID NO: 371), wherein X₁ and X₂ can be Lys or Arg, X₃ can be Trp or Tyr, and X₄ can be Ile or Leu, or wherein none, or one or more of X₁-X₄ and Leu comprises a modified amino acid residue or non-natural amino acid residue.

TABLE 3 – Neurotensin Peptide Variants

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 350	ELYENKPRRPYIL
SEQ ID NO: 365	ELYENKP-[N-methyl-Arg]-KP-[l-neo-Trp]-[<i>tert</i> -Leu]-L
SEQ ID NO: 366	ELYENKPR-[d-Lys]-P-[l-neo-Trp]-[<i>tert</i> -Ile]-L
SEQ ID NO: 367	ELYENKPR-[d-Orn]-P-[l-neo-Trp]-[<i>tert</i> -Leu]-L
SEQ ID NO: 368	ELYENKP-[N-methyl-Arg]-RP-[d-3,1-Nal]-[<i>tert</i> -Leu]-L
SEQ ID NO: 369	ELYENKP-[H-Lys-psi[CH ₂ NH]]-KPY-[Tle]-[Leu-Oet]
SEQ ID NO: 370	ELYENKP
SEQ ID NO: 371	ELYENKPX ₁ X ₂ PX ₃ X ₄ L

[0390] Therefore, the fusion of neurotensin peptide and one of the peptides described herein that can cross the blood brain barrier can produce a fusion peptide capable of crossing the blood barrier which can retain the functional capabilities of neurotensin peptide. For example, the DNA sequence of a peptide of the present disclosure is inserted into the gene of neurotensin to manufacture peptide-neurotensin fusions. In some embodiments, the peptides of this disclosure

are fused to neurotensin or a functional fragment thereof. A functional fragment of neurotensin can be at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues, or at least 13 amino acid residues long.

[0391] As another example, in certain embodiments, the peptides described herein are attached to another molecule, such as an active agent that provides a functional capability. In some embodiments, an active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFNgamma agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter such as neurotensin.

[0392] In some embodiments, TfR-binding peptides can direct the active agent into the cell. In further embodiments, TfR-binding peptides can direct the active agent into the nucleus. In some embodiments, the active agent has intrinsic tumor-homing properties or the active agent can be engineering to have tumor-homing properties. In some embodiments, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 active agents can be linked to a peptide. Multiple active agents can be attached by methods such as conjugating to multiple lysine residues and/or the N-terminus, or by linking the multiple active agents to a scaffold, such as a polymer or dendrimer and then attaching that agent-scaffold to the peptide (such as described in Yurkovetskiy, A. V., Cancer Res 75(16): 3365–72 (2015)).

Examples of active agents include but are not limited to: a peptide, an oligopeptide, a polypeptide, a peptidomimetic, a polynucleotide, a polyribonucleotide, a DNA, a cDNA, a ssDNA, a RNA, a dsRNA, a micro RNA, an oligonucleotide, an antibody, a single chain variable fragment (scFv), an antibody fragment, an aptamer, a cytokine, an interferon, a hormone, an enzyme, a growth factor, a checkpoint inhibitor, a PD-1 inhibitor, a PD-L1 inhibitor, a CD47 inhibitor, a CTLA4 inhibitor, a CD antigen, a chemokine, a neurotransmitter, an ion channel inhibitor, an ion channel activator, a G-protein coupled receptor inhibitor, a G-protein coupled receptor activator, a chemical agent, a radiosensitizer, a radioprotectant, a radionuclide, a therapeutic small molecule, a steroid, a corticosteroid, an anti-inflammatory agent, an immune modulator, a complement fixing peptide or protein, a tumor necrosis factor inhibitor, a tumor

necrosis factor activator, a tumor necrosis factor receptor family agonist, a tumor necrosis receptor antagonist, a Tim-3 inhibitor, a protease inhibitor, an amino sugar, a chemotherapeutic, a cytotoxic molecule, a toxin, a tyrosine kinase inhibitor, an anti-infective agent, an antibiotic, an anti-viral agent, an anti-fungal agent, an aminoglycoside, a nonsteroidal anti-inflammatory drug (NSAID), a statin, a nanoparticle, a liposome, a polymer, a biopolymer, a polysaccharide, a proteoglycan, a glycosaminoglycan, polyethylene glycol, a lipid, a dendrimer, a fatty acid, or an Fc region, or an active fragment or a modification thereof.

[0393] In some embodiments, the peptide is covalently or non-covalently linked to an active agent, e.g., directly or via a linker. For example, cytotoxic molecules that can be used include auristatins, MMAE, MMAF, dolostatin, auristatin F, monomethylauristatin D, DM1, DM4, maytansinoids, maytansine, calicheamicins, N-acetyl- γ -calicheamicin, pyrrolobenzodiazepines, PBD dimers, doxorubicin, vinca alkaloids (4-deacetylvinblastine), duocarmycins, cyclic octapeptide analogs of mushroom amatoxins, epothilones, and anthracyclines, CC-1065, taxanes, paclitaxel, cabazitaxel, docetaxel, SN-38, irinotecan, vincristine, vinblastine, platinum compounds, cisplatin, methotrexate, BACE (beta-secretase 1) inhibitors such as verubecestat, chlorambucil, mitomycin C. Additional examples of active agents are described in McCombs, J. R., AAPS J, 17(2): 339–51 (2015), Ducry, L., Antibody Drug Conjugates (2013), and Singh, S. K., Pharm Res. 32(11): 3541–3571 (2015). Additional examples of therapeutic payloads which therapeutic efficacy can be significantly improved when used in combination with the compositions and methods of the present disclosure include Carmustine, Cisplatin, Cyclophosphamide, Etoposide, Irinotecan, Lomustine, Procarbazine, Temozolomide, Vincristine, and Bevacizumab. Additional examples of therapeutic payloads are compounds that have therapeutic benefit in neurodegenerative diseases such as BACE inhibitors or auto-immunity diseases, and effective treatments for pain and migraine such as CGRP Receptor antagonists. Active agents used to treat pain are also consistent with the present disclosure. For example, in some embodiments, the active agent is a neurotransmitter, such as neurotensin. Thus, peptide constructs disclosed herein are peptide-neurotensin constructs. Exemplary linkers suitable for use with the embodiments herein are discussed in further detail below.

[0394] Only a small fraction of currently available drug molecules have applicability in CNS diseases due to their poor BBB penetration capabilities. About 98% of small molecule drugs do not or do only to a very limited degree cross the BBB. In addition, nearly 100% of macromolecular drug molecules (e.g., antibodies) do not exhibit significant BBB penetration

capabilities. (See e.g., Mikitsh et al. Pathways for Small Molecule Delivery to the Central Nervous System Across the Blood-Brain Barrier, *Perspect Medicin Chem.* 2014; 6: 11–24). Thus, drug molecules available or currently in preclinical or clinical development can be used in combination with the presently described methods and compositions to provide improved therapeutic activity of these drugs in the cells, tissues, or organs expressing TfR (e.g., cancer cells or immune cells) and thus provide superior clinical outcomes compared to conventional therapies. For example, a drug molecule that shows activity against a specific biological target (e.g., a protein or a nucleic acid) but is prevented from reaching said target in vivo due to cellular barriers or cell membranes, said drug molecule can be conjugated to, linked to, or fused to a TfR-binding peptide of the present disclosure for enhanced delivery via TfR-mediated transport across those cellular barriers (e.g., BBB) or cell membranes (e.g., a cell membrane of a cancer cell). The peptide-drug conjugate or fusion molecule can cross the BBB via TfR-mediated transcytosis and thus provide significantly higher drug concentrations (e.g., therapeutic concentrations in the CNS to treat or prevent a CNS disorder (e.g., a brain cancer). For example, palbociclib has been shown efficacy in the treatment of breast cancer, but activity against CNS tumors has been limited. Thus, conjugating the drug palbociclib to a TfR-binding peptide of the present disclosure can enable its use in the treatment and/or prevention of tumors that are not accessible for the drug molecule itself, for instance those located in the CNS (e.g., brain tumors). Brain tumors that can be treated or prevented using conjugates or fusion molecules comprising one or more TfR-binding peptides of the present disclosure can include glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, and diffuse intrinsic pontine glioma.

[0395] In some embodiments, the TfR-binding peptides as described herein can be used to treat and/or prevent diseases in non-CNS and/or peripheral cells, tissues, or organs. For example, the TfR-binding peptides as described herein can be used to treat and/or prevent muscle diseases. Drug or drug candidates that may be used in combination with the TfR-binding peptides to treat or prevent muscle diseases can include drugs or drug candidates for muscular dystrophy, spinal bulbar muscular dystrophy (SBMA), myotonic dystrophy, cachexia, or sarcopenia. In some embodiments, drugs can be senolytics or other anti-aging molecules (e.g., caspases), or gene therapies for diseases of the muscle.

[0396] In some embodiments, the TfR-binding peptides as described herein can be used to treat pain. The TfR binding peptides can be linked to neurotensin (NT) to generate CDP-NT peptide

constructs (e.g., SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, or SEQ ID NO: 359 – SEQ ID NO: 361). These peptide constructs can target neurotensin receptors in the CNS to suppress pain. In some embodiments, the peptide is capable of reducing a level of pain in a subject upon administration.

[0397] Kinds of pain include chronic pain, acute pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain. Neurotensin is a non-opioid but can provide analgesia. Opioid can cause addiction and neurotensin delivery to the CNS offers the possibility of non-addictive pain relief. Neurotensin peptide fusions can also be used to treat dopaminergic, serotonergic, GABAergic, glutamatergic, and cholinergic systems, schizophrenia, psychosis, drug abuse, Parkinson's disease (PD), pain, central control of blood pressure, eating disorders, as well as, cancer and inflammation. Neurotensin peptide fusions can be used to treat pain including chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. Neurotensin peptide fusions can be used in neuroendocrine regulation, such as of the ACTH axis, the HPA axis, hormone regulation, and ion channel regulation, such as in diseases and disorders including Bartter's syndrome, Andersen's syndrome, congenital hyperinsulinism, neonatal diabetes, dilated cardiomyopathy, episodic ataxia type 1, long QT syndrome, short QT syndrome, benign neonatal febrile convulsions, nonsyndromic deafness, long QT syndrome, short QT syndrome, polycystic kidney disease, familial episodic pain syndrome, focal segmental glomerulosclerosis, retinitis pigmentosa, epilepsy, severe myoclonic epilepsy, long QT syndrome, cerebellar ataxia, erythromelalgia, paroxysmal extreme pain disorder, congenital indifference to pain, benign familial neonatal seizures, timothy syndrome, episodic ataxia type 2, stiff baby syndrome, juvenile myoclonic epilepsy, and autosomal dominant nocturnal frontal lobe epilepsy. In some embodiments, peptide constructs (e.g., peptide-NT constructs) interact with dopamine-signaling neurons to reduce pain.

[0398] In some embodiments, the TfR-binding peptides of the present disclosure can be used for the treatment and prevention of various neurological diseases including but not limited to epilepsy, schizophrenia, depression, anxiety, bipolar disorder, developmental brain disorders (e.g., autism spectrum), or mood disorder.

[0399] Binding of the herein described engineered peptides and peptide constructs (e.g., peptide conjugates, fusion peptides, or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via vesicular transcytosis) or a cell membrane (e.g., via endocytosis) can have implications in a number of diseases, conditions, or disorders associated with neurodegeneration. Neurodegenerative diseases that can be treated, prevented, or diagnosed with the herein described TfR-binding peptides can include Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. In some embodiments, the TfR-binding peptide can be used in combination with BACE inhibitors, galantamine, amantadine, benztropine, biperiden, bromocriptin, carbidopa, donepezil, entacapone, levodopa, pergolide, pramipexole, procyclidine, rivastigmine, ropinirole, selegiline, tacrine, tolcapone, or trihexyphenidyl to treat and/or prevent a neurodegenerative disease.

[0400] In some embodiments, modulation (e.g., inhibition) of ion channels such as Kv1.3 potassium channels using TfR-binding peptides and ion channel modulator such as Kv1.3 potassium channel inhibitor conjugates or peptide constructs can be used for treating or preventing inflammation in the brain. Kv1.3 potassium channels, for example, can be highly expressed on microglia in the brain. Thus, neuroinflammatory and neurodegenerative diseases such as multiple sclerosis, Alzheimer's, Parkinson's, traumatic brain injury, radiation therapy toxicity and other neurodegenerative and neuroinflammatory diseases can be treated with TfR-binding peptides that are conjugated to, linked to, or fused to a Kv1.3 inhibitor. These diseases can be marked by upregulation of Kv1.3. In some cases, Kv1.3 inhibition can be used for treatment of psoriasis and other non-brain autoimmune diseases due to its effect on effector T cells. In some embodiments, a Kv1.3 inhibitor peptide may be Vm24 (SEQ ID NO: 378 or SEQ ID NO: 391, GSAAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCYYC) or ShK-186 (SEQ

ID NO: 379). A Kv1.3 inhibitor peptide may be conjugated to any of the peptides disclosed herein (e.g., any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358). Exemplary Kv1.3 inhibitor peptides are provided in **TABLE 4**.

TABLE 4 – Exemplary Kv1.3 Inhibitor Peptides

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 378	AAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYC
SEQ ID NO: 379	RSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTC
SEQ ID NO: 390	GSEWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTC
SEQ ID NO: 391	GSAAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYC
SEQ ID NO: 402	EWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTC

[0401] In some embodiments, a Kv1.3 inhibitor, for example a Kv1.3 peptide inhibitor, may be conjugated to the TfR-binding peptides of the present disclosure. Exemplary Kv1.3 inhibitor fusion peptides are provided in **TABLE 5**. A control for a Kv1.3 inhibitor fusion peptide of the present disclosure may comprise a control peptide fused to a Kv1.3 inhibitor (e.g., SEQ ID NO: 380

(GSAAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYCGGGGSGGGGSRLEDICLP RWGLWEDDGGGGSGGGGSKCLPPGKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 382

(GSEWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCGGGGSGGGGSGGGGSKCL PPGKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 383

(GSEWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCKKYKPYVPVTTNKCLPPG KPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 386

(GSAAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYCGGGGSGGGGSGGGGSKCLP PGKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 387

(GSAAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYCKKYKPYVPVTTNKCLPPGK PCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 392

(AAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYCGGGGSGGGGSRLEDICLP RWGLWEDDGGGGSGGGGSKCLPPGKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 394

(EWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCGGGGSGGGGSGGGGSKCLPP GKPCYGATQKIPCCGVCSHNNCT), SEQ ID NO: 395

(EWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCKKYKPYVPVTTNKCLPPGKPCY
GATQKIPCCGVCSHNNCT), SEQ ID NO: 398

(AAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCY YCGGGGSGGGGSGGGGSKCLPPG
KPCY GATQKIPCCGVCSHNNCT), and SEQ ID NO: 399

(AAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCY YCKKYKPYVPVTTNKCLPPGKPC
YGATQKIPCCGVCSHNNCT)). In some embodiments, a Kv1.3 peptide inhibitor may comprise
a linker for fusing or conjugating to a TfR-binding peptide. In some embodiments, a Kv1.3
peptide inhibitor fused to a TfR-binding peptide may comprise a peptide with a sequence of SEQ
ID NO: 390 or SEQ ID NO: 402. In some embodiments, a linker may have a sequence of any one
of SEQ ID NO: 103 – SEQ ID NO: 115 or SEQ ID NO: 125. A Kv1.3 binding peptide (e.g., SEQ
ID NO: 378, SEQ ID NO: 379, SEQ ID NO: 391, or SEQ ID NO: 402) may be conjugated to a
TfR-binding peptide with a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID
NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID
NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358.

TABLE 5 – Peptide-Kv1.3 Inhibitor Fusions

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 381	GSAAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCY YCGGGGSGGG GSRLIEDICLPRWGCLWEDDGGGGSGGGGSREGCASRCMKYNDELE KCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 384	GSEWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCCGGGGSGG GGSGGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELE DLLYCLDHCHSQ
SEQ ID NO: 385	GSEWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCKKYKPYV PVTNREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLY CLDHCHSQ
SEQ ID NO: 388	GSAAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCY YCGGGGSGGG GSGGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDL LYCLDHCHSQ
SEQ ID NO: 389	GSAAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCY YCKKYKPYVP VTNREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYC LDHCHSQ
SEQ ID NO: 393	AAAI SCVGSPECPPKCRAQGCKNGKCMNRKCKCY YCGGGGSGGGGS RLIEDICLPRWGCLWEDDGGGGSGGGGSREGCASRCMKYNDELEK EARMMSMSNTEEDCEQELEDLLYCLDHCHSQ

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 396	EWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCTGGGGSGGG GSGGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDL LYCLDHCHSQ
SEQ ID NO: 397	EWSSRSCIDTIPKSRCTAFQCKHSMKYRLSFCRKTCGTCKKYKPYVPV TTNREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCL DHCHSQ
SEQ ID NO: 400	AAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYCGGGGSGGGGS GGGGSGREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLL YCLDHCHSQ
SEQ ID NO: 401	AAAIscVGSPECPPKCRAQGCKNGKCMNRKCKCYCKKYKPYVPVT TNREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLD HCHSQ

[0402] In some embodiments, the TfR-binding peptides of the present disclosure can be used for the treatment and prevention of Crohn's disease or, more generally, inflammatory bowel diseases. In some cases, the TfR-binding peptides of the present disclosure show high uptake and retention in glandular cells of the intestine, which can express high amounts of TfR.

[0403] In some embodiments, the TfR-binding peptides as described herein can be used to treat and/or prevent cancers located both in and outside the CNS, including but not limited to ovarian cancer, colon cancer, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma, and lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, and diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, or skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, a neuroblastoma, or a CMYC-overexpressing cancer.

[0404] In some embodiments, TfR-binding peptide constructs that can be used to prevent and/or treat a cancer are those comprising a TfR binding peptide and an active agent with anti-tumor activity such as an IL15, a fused IL15/IL15Ra, IFN γ , and anti-CD3 agents. Examples of such peptide constructs (e.g., fusion peptides) are those having an amino acid sequence set forth in any one of SEQ ID NO: 225 – SEQ ID NO: 332. In some embodiments, a peptide construct of the present disclosure can have at least 80%, at least 90%, at least 92%, at least 93%, at least

94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 225 – SEQ ID NO: 332. Exemplary peptide constructs of the present disclosure are shown below in **TABLE 6**. These exemplary peptide constructs can comprise secretion murine IgKappa leader sequences, an anti-CD3 scFv, lysine to arginine mutants of the scFv, shorty-Flag sequences, His-tags, IL-15Ra, IL-15, IFN gamma, any linker disclosed herein, any peptides disclosed herein, and any fragments or derivatives thereof. Also shown in **TABLE 6** are peptide-neurotensin constructs (e.g., peptide constructs with a peptide of this disclosure complexed to neurotensin), for example those shown in SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, and SEQ ID NO: 359 – SEQ ID NO: 361. In some embodiments, a peptide construct of the present disclosure can have at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, and SEQ ID NO: 359 – SEQ ID NO: 361.

TABLE 6 – Exemplary Peptide Construct Sequences

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 225	METDTLLLWVLLLWVPGSTGDIKLQQSGAELARPGASVKMSCKTSGYTFTRY TMHWVKQRPGQGLEWIGYINPSRGYTNYNQKFKDKATLTDDKSSSTAYMQLS SLTSEDSAVYYCARYYDDHYCLDYWGQGTTLTVSSVEGGSGHHHHHHGSGG VDDIQLTQSPAIMSASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTS KVASGVPYRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPLTFGAGTKLE LKGGGSREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCVDHC VSQ
SEQ ID NO: 226	METDTLLLWVLLLWVPGSTGDIKLQQSGAELARPGASVKMSCKTSGYTFTRY TMHWVKQRPGQGLEWIGYINPSRGYTNYNQKFKDKATLTDDKSSSTAYMQLS SLTSEDSAVYYCARYYDDHYCLDYWGQGTTLTVSSVEGGSGHHHHHHHHGSGG VDDIQLTQSPAIMSASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTS KVASGVPYRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPLTFGAGTKLE LKGGGSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHC HSQ
SEQ ID NO: 227	METDTLLLWVLLLWVPGSTGDIKLQQSGAELARPGASVKMSCKTSGYTFTRY TMHWVKQRPGQGLEWIGYINPSRGYTNYNQKFKDKATLTDDKSSSTAYMQLS SLTSEDSAVYYCARYYDDHYCLDYWGQGTTLTVSSVEGGSGHHHHHHHHGSGG VDDIQLTQSPAIMSASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTS KVASGVPYRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPLTFGAGTKLE LKGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDH CHSQ
SEQ ID NO: 228	DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPGQGLEWIGYIN PSRGYTNYNQKFKDKATLTDDKSSSTAYMQLSSLTSEDSAVYYCARYYDDHY CLDYWGQGTTLTVSSVEGGSGHHHHHHHHGSGGVDDIQLTQSPAIMSASPGEKV

SEQ ID NO	Amino Acid Sequence
	TMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGSGTSYS LTISSMEAEDAATYYCQQWSSNPLTFGAGTKLELKGGGSREGCASRCTKYNA ELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 229	DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPGQGLEWIGYIN PSRGYTNYNQKFKDKATLTDDKSSSTAYMQLSSLTSEDSAVYYCARYYDDHY CLDYWGQGTTLTVSSVEGGSGHHHHHHGSGGVDDIQLTQSPAISASPGEKV TMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGSGTSYS LTISSMEAEDAATYYCQQWSSNPLTFGAGTKLELKGGGSREGCASRCTKYNA ELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 230	DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPGQGLEWIGYIN PSRGYTNYNQKFKDKATLTDDKSSSTAYMQLSSLTSEDSAVYYCARYYDDHY CLDYWGQGTTLTVSSVEGGSGHHHHHHGSGGVDDIQLTQSPAISASPGEKV TMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGSGTSYS LTISSMEAEDAATYYCQQWSSNPLTFGAGTKLELKGGGSREGCASRCMKYND ELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 231	METDTLLLWVLLLWVPGSTGDIRLQQSGAELARPGASVRMSCRTSGYTFTRY TMHWVRQRPGQGLEWIGYINPSRGYTNYNQRFDRATLTDDRSSSTAYMQLS SLTSEDSAVYYCARYYDDHYCLDYWGQGTTLTVSSVEGGSGHHHHHHGSGG VDDIQLTQSPAISASPGERVMTMTCRASSSVSYMNWYQQRSSTSPRRWIYDTS RVASGVPYRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPLTFGAGTRLE LRGGGSREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHC VSQ
SEQ ID NO: 232	METDTLLLWVLLLWVPGSTGDIRLQQSGAELARPGASVRMSCRTSGYTFTRY TMHWVRQRPGQGLEWIGYINPSRGYTNYNQRFDRATLTDDRSSSTAYMQLS SLTSEDSAVYYCARYYDDHYCLDYWGQGTTLTVSSVEGGSGHHHHHHGSGG VDDIQLTQSPAISASPGERVMTMTCRASSSVSYMNWYQQRSSTSPRRWIYDTS RVASGVPYRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPLTFGAGTRLE LRGGGSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHC HSQ
SEQ ID NO: 233	METDTLLLWVLLLWVPGSTGDIRLQQSGAELARPGASVRMSCRTSGYTFTRY TMHWVRQRPGQGLEWIGYINPSRGYTNYNQRFDRATLTDDRSSSTAYMQLS SLTSEDSAVYYCARYYDDHYCLDYWGQGTTLTVSSVEGGSGHHHHHHGSGG VDDIQLTQSPAISASPGERVMTMTCRASSSVSYMNWYQQRSSTSPRRWIYDTS RVASGVPYRFSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPLTFGAGTRLE LRGGGSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDH CHSQ
SEQ ID NO: 234	DIRLQQSGAELARPGASVRMSCRTSGYTFTRYTMHWVRQRPGQGLEWIGYIN PSRGYTNYNQRFDRATLTDDRSSSTAYMQLSSLTSEDSAVYYCARYYDDHY CLDYWGQGTTLTVSSVEGGSGHHHHHHGSGGVDDIQLTQSPAISASPGERV TMTCRASSSVSYMNWYQQRSSTSPRRWIYDTSRVASGVPYRFSGSGSGTSYSL TISSMEAEDAATYYCQQWSSNPLTFGAGTRLELRGGGSREGCASRCTRYNAEL ERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 235	DIRLQQSGAELARPGASVRMSCRTSGYTFTRYTMHWVRQRPGQGLEWIGYIN PSRGYTNYNQRFDRATLTDDRSSSTAYMQLSSLTSEDSAVYYCARYYDDHY CLDYWGQGTTLTVSSVEGGSGHHHHHHGSGGVDDIQLTQSPAISASPGERV TMTCRASSSVSYMNWYQQRSSTSPRRWIYDTSRVASGVPYRFSGSGSGTSYSL TISSMEAEDAATYYCQQWSSNPLTFGAGTRLELRGGGSREGCASRCTRYNAEL ERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 236	DIRLQQSGAELARPGASVRMSCRTSGYTFTRYTMHWVRQRPGQGLEWIGYIN PSRGYTNYNQRFDRATLTDDRSSSTAYMQLSSLTSEDSAVYYCARYYDDHY

SEQ ID NO	Amino Acid Sequence
	CLDYWGQGTTLTVSSVEGGSGHHHHHHGSGGVDDIQLTQSPAIMSASPGERV TMTCRASSSVSYMNWYQQRSGTSPRRWIYDTSRVASGVPPYRFSGSGSGTSYSL TISSMEAEDAATYYCQQWSSNPLTFGAGTRLELRGGGSREGCASRCMRYNDE LERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 237	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGSGGGGSGGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSC KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEEKTFTEFLQSFIRIVQMFINTSGGGSREGCASRCKYNAELEKCEARVSS MSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 238	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGSGGGGSGGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSC KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEEKTFTEFLQSFIRIVQMFINTSGGGSREGCASRCKYNAELEKCEARVM MSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 239	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGSGGGGSGGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSC KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEEKTFTEFLQSFIRIVQMFINTSGGGSREGCASRCMKYNDELEKCEARM MSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 240	DYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECEELEEKTFTEFLQSFIRIVQMFIN TSGGGSREGCASRCKYNAELEKCEARVSSMSNTEETCVQELFDLLHCVDHC VSQ
SEQ ID NO: 241	DYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECEELEEKTFTEFLQSFIRIVQMFIN TSGGGSREGCASRCKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHC HSQ
SEQ ID NO: 242	DYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECEELEEKTFTEFLQSFIRIVQMFIN TSGGGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDH CHSQ
SEQ ID NO: 243	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGSGTTCPPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSGGGSREGCASRCTRYNAELERCEARVSSMS NTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 244	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGSGTTCPPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE

SEQ ID NO	Amino Acid Sequence
	ELEERTFTEFLQSFIRIVQMFINTSGGGSREGCASRCTRYNAELERCEARVMSM SNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 245	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGTTCPPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNV AHWTTPSLRCIRDPSLAGG GGSGGGSGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSGGGSREGCASRCMRYNDELERCEARMMS MSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 246	DYRDEHHHHHHHSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNV AHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCV SQ
SEQ ID NO: 247	DYRDEHHHHHHHSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNV AHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHC HSQ
SEQ ID NO: 248	DYRDEHHHHHHHSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNV AHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHC HSQ
SEQ ID NO: 249	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGHGTVIESLES LN NYFNSSGI DVEEKSLFLDIWRNWQKD GDMKILQS QIISFYLR LFEVLKDNQAISNNISVIESH LITTFFSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRK RSRGGGSHGTVIESLES LN NYFNSSGIDVEEKSLFLDIWRNWQKD GDMKILQS QIISFYLR LFEVLKDNQAISNNISVIESHLITTFFSNSKAKKDAFMSIAKFEVNNP QVQRQAFNELIRVVHQLLPESSLRKRKR RSRGGGSREGCASRCTKYNAELEKCE ARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 250	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGHGTVIESLES LN NYFNSSGI DVEEKSLFLDIWRNWQKD GDMKILQS QIISFYLR LFEVLKDNQAISNNISVIESH LITTFFSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRK RSRGGGSHGTVIESLES LN NYFNSSGIDVEEKSLFLDIWRNWQKD GDMKILQS QIISFYLR LFEVLKDNQAISNNISVIESHLITTFFSNSKAKKDAFMSIAKFEVNNP QVQRQAFNELIRVVHQLLPESSLRKRKR RSRGGGSREGCASRCTKYNAELEKCE ARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 251	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGHGTVIESLES LN NYFNSSGI DVEEKSLFLDIWRNWQKD GDMKILQS QIISFYLR LFEVLKDNQAISNNISVIESH LITTFFSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRK RSRGGGSHGTVIESLES LN NYFNSSGIDVEEKSLFLDIWRNWQKD GDMKILQS QIISFYLR LFEVLKDNQAISNNISVIESHLITTFFSNSKAKKDAFMSIAKFEVNNP QVQRQAFNELIRVVHQLLPESSLRKRKR RSRGGGSREGCASRCMKYNDELEKC EARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 252	DYKDEHHHHHHHSGHGTVIESLES LN NYFNSSGIDVEEKSLFLDIWRNWQKD G DMKILQS QIISFYLR LFEVLKDNQAISNNISVIESHLITTFFSNSKAKKDAFMSIA KFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRKR RSRGGGSHGTVIESLES LN

SEQ ID NO	Amino Acid Sequence
	YFNSSGIDVEEKSLFLDIWRNWQKDGMKILQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRKRSRGGGSREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 253	DYKDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEEKSLFLDIWRNWQKDGMKILQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRKRSRGGGSHTVIESLESLESLNNYFNSSGIDVEEKSLFLDIWRNWQKDGMKILQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRKRSRGGGSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 254	DYKDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEEKSLFLDIWRNWQKDGMKILQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRKRSRGGGSHTVIESLESLESLNNYFNSSGIDVEEKSLFLDIWRNWQKDGMKILQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVHQLLPESSLRKRKRSRGGGSREGCASRCMKYNDLEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 255	METDTLLLWVLLLWVPGSTG DYRDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSHTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 256	METDTLLLWVLLLWVPGSTG DYRDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSHTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 257	METDTLLLWVLLLWVPGSTG DYRDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSHTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 258	DYRDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 259	DYRDEHHHHHHGSHGTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNISVIESHLITTFSSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESSLRRRRRSRGGGSHTVIESLESLESLNNYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRRLFVLRDNQAISNNIS

SEQ ID NO	Amino Acid Sequence
	VIESHLITTFFSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESLRRRRRSRGGGSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 260	DYRDEHHHHHHGSHGTVIESLES LN NYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRLEVL RDNQAISNNISVIESHLITTFFSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESLRRRRRSRGGGSHTVIESLES LN NYFNSSGIDVEERSLFLDIWRNWQRDGMRLQSQIISFYLRLEVL RDNQAISNNISVIESHLITTFFSNSRARRDAFMSIARFEVNNPQVQRQAFNELIRVVHQLLPESLRRRRRSRGGGSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 261	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTPSLKCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCV DHCVSQ
SEQ ID NO: 262	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTPSLKCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 263	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTPSLKCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 264	DYKDEHHHHHHHSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTPSLKCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCV DHCVSQ
SEQ ID NO: 265	DYKDEHHHHHHHSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTPSLKCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 266	DYKDEHHHHHHHSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTPSLKCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 267	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTPSLRCI RDPSLAGGGSGGGSGGGSGGGSGNWDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE

SEQ ID NO	Amino Acid Sequence
	ELEERTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCTRYNAELERCEARV SSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 268	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGGTTCTPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCTRYNAELERCEARV MSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 269	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGGTTCTPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSGGGSGGGSGREGCASRCMRYNDELERCEAR MMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 270	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGSGGGSGGGSGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYDSDFHPSRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCV DHCVSQ
SEQ ID NO: 271	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGSGGGSGGGSGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYDSDFHPSRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHC LDHCHSQ
SEQ ID NO: 272	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGSGGGSGGGSGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYDSDFHPSRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYC LDHCHSQ
SEQ ID NO: 273	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGGTTCTPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPS KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGGGSGREGCASRCTKYNAEL EKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 274	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGGTTCTPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPS KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGGGSGREGCASRCTKYNAEL EKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 275	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGGTTCTPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPS KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEKTFTEFLQSFIRIVQMFINTSGGGSGGGSGGGSGREGCASRCMKYNDEL EKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 276	DYKDEHHHHHHHSGGTTCTPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCKVTAMNCFLELQVILHEYNSM TLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDL LHCVDHCVSQ
SEQ ID NO: 277	DYKDEHHHHHHHSGGTTCTPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCKVTAMNCFLELQVILHEYNSM TLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELED LLHCLDHCHSQ
SEQ ID NO: 278	DYKDEHHHHHHHSGGTTCTPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCKVTAMNCFLELQVILHEYNSM TLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCMKYNDLEKCEARMMSMSNTEEDCEQELED DLLYCLDHCHSQ
SEQ ID NO: 279	METDTLLLWVLLLWVPGSTG DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGGSGGGGGSNWIDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCR VTAMNCFLELQVILHEYNSMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCTRYNAELERC EARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 280	METDTLLLWVLLLWVPGSTG DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGGSGGGGGSNWIDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCR VTAMNCFLELQVILHEYNSMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCTRYNAELERC EARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 281	METDTLLLWVLLLWVPGSTG DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGGSGGGGGSNWIDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCR VTAMNCFLELQVILHEYNSMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCMRYNDELER CEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 282	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCRVTAMNCFLELQVILHEYNSM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDL LHCVDHCVSQ
SEQ ID NO: 283	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCRVTAMNCFLELQVILHEYNSM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGGSGGGSGGGSGREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELED LLHCLDHCHSQ
SEQ ID NO: 284	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW

SEQ ID NO	Amino Acid Sequence
	IDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSGGSGGGSGGGSGGGSCASRCMRYNDELERCEARMMSMSNTEEDCEQELED LLYCLDHCHSQ
SEQ ID NO: 285	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGGSGGGSGGGSGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSC KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEKTFTEFLQSFIRIVQMFINTESEPRVPITQNREGCASRCTKYNAELEKCE ARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 286	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGGSGGGSGGGSGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSC KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEKTFTEFLQSFIRIVQMFINTESEPRVPITQNREGCASRCTKYNAELEKCE ARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 287	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGGSGGGSGGGSGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSC KVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEKTFTEFLQSFIRIVQMFINTESEPRVPITQNREGCASRCMKYNDELEKCE ARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 288	DYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGSGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCKVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECEELEKTFTEFLQSFIRIVQMFIN TSEPRVPITQNREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHC VDHCVSQ
SEQ ID NO: 289	DYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGSGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCKVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECEELEKTFTEFLQSFIRIVQMFIN TSEPRVPITQNREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHC LDHCHSQ
SEQ ID NO: 290	DYKDEHHHHHHHSGSGTTCPPPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGSGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLYTDSDFHPSCKVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECEELEKTFTEFLQSFIRIVQMFIN TSEPRVPITQNREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLY CLDHCHSQ
SEQ ID NO: 291	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGSGTTCPPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTESEPRVPITQNREGCASRCTRYNAELERCEAR VSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 292	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHSGSGTTCPPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGSGGGSGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLYTDSDFHPSCR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE

SEQ ID NO	Amino Acid Sequence
	ELEERTFTEFLQSFIRIVQMFINTSEPRVPITQNREGCASRCTRYNAELERCEAR VMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 293	METDTLLLWVLLLVWPGSTGDYRDEHHHHHHHSGGTTCTPPVSIHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGGSGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSR VTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSEPRVPITQNREGCASRCMRYNDELERCEAR MMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 294	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSGNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSEPRVPITQNREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCV DHCVSQ
SEQ ID NO: 295	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSGNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSEPRVPITQNREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHC LDHCHSQ
SEQ ID NO: 296	DYRDEHHHHHHHSGGTTCTPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSGNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFIN TSEPRVPITQNREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLY CLDHCHSQ
SEQ ID NO: 297	METDTLLLWVLLLVWPGSTGDYKDEHHHHHHHENLYFQGSREGCASRCTKYN AELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQQGGGSGTTCTPPVSI HADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKC IRDPSLAGGGGSGGGGSGGGGSGGGGSGNWIDVRYDLEKIESLIQSIHIDTTLTY DSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNV AESGCKECELEEKTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 298	METDTLLLWVLLLVWPGSTGDYKDEHHHHHHHENLYFQGSREGCASRCTKYN AELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQGGGSGTTCTPPVSI HADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKC IRDPSLAGGGGSGGGGSGGGGSGGGGSGNWIDVRYDLEKIESLIQSIHIDTTLTY DSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNV AESGCKECELEEKTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 299	METDTLLLWVLLLVWPGSTGDYKDEHHHHHHHENLYFQGSREGCASRCMKYN DELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQGGGSGTTCTPPVSI HADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKC IRDPSLAGGGGSGGGGSGGGGSGGGGSGNWIDVRYDLEKIESLIQSIHIDTTLTY DSDFHPSCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNV AESGCKECELEEKTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 300	DYKDEHHHHHHHENLYFQGSREGCASRCTKYNAELEKCEARVSSMSNTEETCV QELFDLLHCVDHCVSQQGGGSGTTCTPPVSIHADIRVKNYSVNSRERYVCNS GFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGG SGGGGSGNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMNCFLELQ VILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQS FIRIVQMFINTS

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 301	DYKDEHHHHHHENLYFQGSREGCASRCTKYNAELEKCEARVMSMSNTEEDC EQELEDLLHCLDHCHSQGGGGSGTTCTPPVSIHADIRVKNYSVNSRERYVCN SGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGG GSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSCKVTAMNCFLEL QVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFL QSFIRIVQMFINTS
SEQ ID NO: 302	DYKDEHHHHHHENLYFQGSREGCASRCMKYNDELEKCEARMMSMSNTEED CEQELEDLLYCLDHCHSQGGGGSGTTCTPPVSIHADIRVKNYSVNSRERYVC NSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGG GGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSCKVTAMNCFLE LQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFL QSFIRIVQMFINTS
SEQ ID NO: 303	METDTLLWVLLWVPGSTGDYRDEHHHHHHENLYFQGSREGCASRCTRYN AELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQGGGGSGTTCTPPVSI HADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLR CIRDPSLAGGGGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTL TYDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRN VAESGCRECEELEERTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 304	METDTLLWVLLWVPGSTGDYRDEHHHHHHENLYFQGSREGCASRCTRYN AELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQGGGGSGTTCTPPVSI HADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLR CIRDPSLAGGGGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTL TYDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRN VAESGCRECEELEERTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 305	METDTLLWVLLWVPGSTGDYRDEHHHHHHENLYFQGSREGCASRCMRYN DELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQGGGGSGTTCTPPVSI HADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLR CIRDPSLAGGGGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTL TYDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRN VAESGCRECEELEERTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 306	DYRDEHHHHHHENLYFQGSREGCASRCTRYNAELERCEARVSSMSNTEETCV QELFDLLHCVDHCVSQGGGGSGTTCTPPVSIHADIRVRNYSVNSRERYVCNS GFRRRAGTSTLIECVINRNTNVAHWTTPSLR CIRDPSLAGGGGSGGGGSGGG SGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSCRVTAMNCFLELQ VILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQ SFIRIVQMFINTS
SEQ ID NO: 307	DYRDEHHHHHHENLYFQGSREGCASRCTRYNAELERCEARVMSMSNTEEDC EQELEDLLHCLDHCHSQGGGGSGTTCTPPVSIHADIRVRNYSVNSRERYVCN SGFRRRAGTSTLIECVINRNTNVAHWTTPSLR CIRDPSLAGGGGSGGGGSGGG GSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSCRVTAMNCFLELQ VILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQ SFIRIVQMFINTS
SEQ ID NO: 308	DYRDEHHHHHHENLYFQGSREGCASRCMRYNDELERCEARMMSMSNTEEDC EQELEDLLYCLDHCHSQGGGGSGTTCTPPVSIHADIRVRNYSVNSRERYVCN SGFRRRAGTSTLIECVINRNTNVAHWTTPSLR CIRDPSLAGGGGSGGGGSGGG GSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSCRVTAMNCFLELQ VILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQ SFIRIVQMFINTS
SEQ ID NO: 309	METDTLLWVLLWVPGSTGDYKDEHHHHHHGSGTTCTPPVSIHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG

SEQ ID NO	Amino Acid Sequence
	GGGSGGGGSGGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSC KVTAMNCFLLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEEKTFTEFLQSFIRIVQMFINTSEPRIPKPSTPPGSSREGCASRCKYNAELE KCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 310	METDTLLLWVLLWVPGSTGDYKDEHHHHHHHSGGTTCTPPVVSIEHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGGSGGGGSGGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSC KVTAMNCFLLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEEKTFTEFLQSFIRIVQMFINTSEPRIPKPSTPPGSSREGCASRCKYNAELE KCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 311	METDTLLLWVLLWVPGSTGDYKDEHHHHHHHSGGTTCTPPVVSIEHADIRVKN YSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAG GGGSGGGGSGGGGSGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSC KVTAMNCFLLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKE CEELEEKTFTEFLQSFIRIVQMFINTSEPRIPKPSTPPGSSREGCASRCMKYNDEL EKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 312	DYKDEHHHHHHHSGGTTCTPPVVSIEHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSCKVTAMNCFLLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINT TSEPRIPKPSTPPGSSREGCASRCKYNAELEKCEARVSSMSNTEETCVQELFDL LHCVDHCVSQ
SEQ ID NO: 313	DYKDEHHHHHHHSGGTTCTPPVVSIEHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSCKVTAMNCFLLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINT TSEPRIPKPSTPPGSSREGCASRCKYNAELEKCEARVMSMSNTEEDCEQELED LLHCLDHCHSQ
SEQ ID NO: 314	DYKDEHHHHHHHSGGTTCTPPVVSIEHADIRVKNYSVNSRERYVCNSGFKRKAGT STLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGSNW IDVRYDLEKIESLIQSIHIDTTLTYDSDFHPSCKVTAMNCFLLELQVILHEYSNM TLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINT TSEPRIPKPSTPPGSSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELED LLYCLDHCHSQ
SEQ ID NO: 315	METDTLLLWVLLWVPGSTGDYRDEHHHHHHHSGGTTCTPPVVSIEHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSCR VTAMNCFLLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSEPRIPRPSTPPGSSREGCASRCTRYNAELERC EARVSSMSNTEETCVQELFDLLHCVDHCVSQ
SEQ ID NO: 316	METDTLLLWVLLWVPGSTGDYRDEHHHHHHHSGGTTCTPPVVSIEHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSCR VTAMNCFLLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE ELEERTFTEFLQSFIRIVQMFINTSEPRIPRPSTPPGSSREGCASRCTRYNAELERC EARVMSMSNTEEDCEQELEDLLHCLDHCHSQ
SEQ ID NO: 317	METDTLLLWVLLWVPGSTGDYRDEHHHHHHHSGGTTCTPPVVSIEHADIRVRN YSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGG GGSGGGGSGGGGSGGGGSNWIDVRYDLERIESLIQSIHIDTTLTYDSDFHPSCR VTAMNCFLLELOVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECE

SEQ ID NO	Amino Acid Sequence
	ELEERTFTEFLQSFIRIVQMFINTSEPRIPRPSTPPGSSREGCASRCMRYNDELER CEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ
SEQ ID NO: 318	DYRDEHHHHHHGSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFINT TSEPRIPRPSTPPGSSREGCASRCRYNAELERCEARVSSMSNTEETCVQELFDL LHCVDHCVSQ
SEQ ID NO: 319	DYRDEHHHHHHGSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFINT TSEPRIPRPSTPPGSSREGCASRCRYNAELERCEARVMSMSNTEEDCEQELED LLHCLDHCHSQ
SEQ ID NO: 320	DYRDEHHHHHHGSGTTCPPPVSIHADIRVRNYSVNSRERYVCNSGFRRRAGT STLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNW IDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNM TLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFTEFLQSFIRIVQMFINT TSEPRIPRPSTPPGSSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELED LLYCLDHCHSQ
SEQ ID NO: 321	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHENLYFQGSREGCASRCTKYN AELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQGGGGSGGGGSGTTCPP PPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTT PSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNWIDVRYDLEKIESLIQSIHIDT TLTYTDSDFHPCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSS NKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 322	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHENLYFQGSREGCASRCTKYN AELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQGGGGSGGGGSGTTCPP PPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTT PSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNWIDVRYDLEKIESLIQSIHIDT TLTYTDSDFHPCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSS NKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 323	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHENLYFQGSREGCASRCMKYN DELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQGGGGSGGGGSGTTCPP PPVSIHADIRVKNYSVNSRERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTT PSLKCIRDPSLAGGGGSGGGGSGGGGSGGGGGSNWIDVRYDLEKIESLIQSIHIDT TLTYTDSDFHPCKVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSS NKNVAESGCKECELEEKTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 324	DYKDEHHHHHHHENLYFQGSREGCASRCTKYNAELEKCEARVSSMSNTEETCV QELFDLLHCVDHCVSQGGGGSGGGGSGTTCPPPVSIHADIRVKNYSVNSRER YVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGGG SGGGGSGGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPCKVTAMNCF LLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTF TEFLQSFIRIVQMFINTS
SEQ ID NO: 325	DYKDEHHHHHHHENLYFQGSREGCASRCTKYNAELEKCEARVMSMSNTEEDC EQELEDLLHCLDHCHSQGGGGSGGGGSGTTCPPPVSIHADIRVKNYSVNSRE RYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGGG GGGGSGGGGGSNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPCKVTAMNC FLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEKTF TEFLQSFIRIVQMFINTS

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 326	DYKDEHHHHHHENLYFQGSREGCASRCMKYNDELEKCEARMMSMSNTEED CEQELEDLLYCLDHCHSQGGGGSGGGGSGTTCPPPVSIEHADIRVKNYSVNSR ERYVCNSGFKRKAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAGGGGSGG GGSGGGGSGGGGSGNWIDVRYDLEKIESLIQSIHIDTTLTYTDSDFHPSCKVTAMN CFLLELQVILHEYSNMTLNETVRNVLYLANSTLSSNKNVAESGCKECELEEK TFTEFLQSFIRIVQMFINTS
SEQ ID NO: 327	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHENLYFQGSREGCASRCTRYN AELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQGGGGSGGGGSGTTCP PPVSIEHADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTT SLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDT LYTDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSSN RNVAESGCRECEELEERTTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 328	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHENLYFQGSREGCASRCTRYN AELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQGGGGSGGGGSGTTCP PPVSIEHADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTT PSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDT TLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSS NRNVAESGCRECEELEERTTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 329	METDTLLLWVLLLWVPGSTGDYRDEHHHHHHHENLYFQGSREGCASRCMRYN DELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQGGGGSGGGGSGTTCP PPVSIEHADIRVRNYSVNSRERYVCNSGFRRRAGTSTLIECVINRNTNVAHWTT PSLRCIRDPSLAGGGGSGGGGSGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDT TLTYTDSDFHPSCRVTAMNCFLELQVILHEYSNMTLNETVRNVLYLANSTLSS NRNVAESGCRECEELEERTTFTEFLQSFIRIVQMFINTS
SEQ ID NO: 330	DYRDEHHHHHHHENLYFQGSREGCASRCTRYNAELERCEARVSSMSNTEETCV QELFDLLHCVDHCVSQGGGGSGGGGSGTTCPPPVSIEHADIRVRNYSVNSRER YVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGGG SGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCF LLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFT EFLQSFIRIVQMFINTS
SEQ ID NO: 331	DYRDEHHHHHHHENLYFQGSREGCASRCTRYNAELERCEARVMSMSNTEEDC EQELEDLLHCLDHCHSQGGGGSGGGGSGTTCPPPVSIEHADIRVRNYSVNSRE RYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGG GSGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCF LLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFT TEFLQSFIRIVQMFINTS
SEQ ID NO: 332	DYRDEHHHHHHHENLYFQGSREGCASRCMRYNDELERCEARMMSMSNTEEDC EQELEDLLYCLDHCHSQGGGGSGGGGSGTTCPPPVSIEHADIRVRNYSVNSRE RYVCNSGFRRRAGTSTLIECVINRNTNVAHWTTPSLRCIRDPSLAGGGGSGGG GSGGGGSGGGGSGNWIDVRYDLERIESLIQSIHIDTTLTYTDSDFHPSCRVTAMNCF LLELQVILHEYSNMTLNETVRNVLYLANSTLSSNRNVAESGCRECEELEERTFT TEFLQSFIRIVQMFINTS
SEQ ID NO: 33	GSREGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ SSELYENKPRRPYIL
SEQ ID NO: 34	GSREGCASRCTKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ SSELYENKPRRPYIL
SEQ ID NO: 35	GSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ GSSELYENKPRRPYIL
SEQ ID NO: 68	REGCASRCTKYNAELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQSSE LYENKPRRPYIL

SEQ ID NO	Amino Acid Sequence
SEQ ID NO: 69	REGCASRCKYNAELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQGSS ELYENKPRRPYIL
SEQ ID NO: 70	REGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQGS SELYENKPRRPYIL
SEQ ID NO: 168	GSREGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQGS SELYENRPRRPYIL
SEQ ID NO: 169	GSREGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQG SSELYENRPRRPYIL
SEQ ID NO: 170	GSREGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ GSSELYENRPRRPYIL
SEQ ID NO: 203	REGCASRCTRYNAELERCEARVSSMSNTEETCVQELFDLLHCVDHCVSQGSSE LYENRPRRPYIL
SEQ ID NO: 204	REGCASRCTRYNAELERCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQGSS ELYENRPRRPYIL
SEQ ID NO: 205	REGCASRCMRYNDELERCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQGSS ELYENRPRRPYIL
SEQ ID NO: 359	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTSDLPAPPLSKVPL QQNFQDNQFQGKWYVVGLAGNAILREDKDPQKMYATIYELKEDKSYNVT SVLFRKKKCDYWIRTFVPGSQPGEFTLGNIKSYPLTSYLVRVVSTNYNQHAMV FVKVSNREYFKITLYGRTELKELTSELKENFIRFSKSLGLPENHIVFPVPIDQ CIDGGSENLYFQGSREGCASRCKYNAELEKCEARVSSMSNTEETCVQELFDLL HCVHCVSQGSSELYENKPRRPYIL
SEQ ID NO: 360	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTSDLPAPPLSKVPL QQNFQDNQFQGKWYVVGLAGNAILREDKDPQKMYATIYELKEDKSYNVT SVLFRKKKCDYWIRTFVPGSQPGEFTLGNIKSYPLTSYLVRVVSTNYNQHAMV FVKVSNREYFKITLYGRTELKELTSELKENFIRFSKSLGLPENHIVFPVPIDQ CIDGGSENLYFQGSREGCASRCKYNAELEKCEARVMSMSNTEEDCEQELEDLL HCLDHCHSQGSSELYENKPRRPYIL
SEQ ID NO: 361	METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTSDLPAPPLSKVPL QQNFQDNQFQGKWYVVGLAGNAILREDKDPQKMYATIYELKEDKSYNVT SVLFRKKKCDYWIRTFVPGSQPGEFTLGNIKSYPLTSYLVRVVSTNYNQHAMV FVKVSNREYFKITLYGRTELKELTSELKENFIRFSKSLGLPENHIVFPVPIDQ CIDGGSENLYFQGSREGCASRCMKYNDELEKCEARMMSMSNTEEDCEQELEDLL YCLDHCHSQGSSELYENKPRRPYIL

[0405] Exemplary peptide-neurotensin constructs are set forth in SEQ ID NO 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, and SEQ ID NO: 359 – SEQ ID NO: 361. The peptides of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, and SEQ ID NO: 356 – SEQ ID NO: 358 could further be fused to neurotensin. Each of said peptides or peptide-NT fusions may include a peptide of SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 32 (or SEQ ID NO: 36, SEQ ID NO: 37, or SEQ ID NO: 67 with no N-terminal GS), and each of which can further include an expression tag or sequence to improve protein expression, secretion, and/or purification, such as the genetic fusion

expression tag siderocalin. In some embodiments, the TfR-binding peptides of the present disclosure are conjugated to, linked to, or fused to anti-cancer drugs that can include an EGFRvIII inhibitor, Copiktra (duvelisib), Erleada (apalutamide), Libtayo (cemiplimab-rwlc), Lorbrena (lorlatinib), Lumoxiti (moxetumomab pasudotox-tdf), Lutathera (lutetium Lu 177 dotatate), Talzenna (talazoparib), Vizimpro (dacomitinib), Aliqopa (copanlisib), Alunbrig (brigatinib), Bavencio (avelumab), Besponsa (inotuzumab ozogamicin), Calquence (acalabrutinib), IDHIFA (enasidenib), Imfinzi (durvalumab), Kiskali (ribociclib), Kymriah (tisagenlecleucel), Nerlynx (neratinib), Rydapt (midostaurin), Vyxeos (daunorubicin and cytarabine), Xermelo (telotristat ethyl), Yescarta (axicabtagene ciloleucel), Zejula (niraparib), Cabometyx (cabozantinib), Keytruda (pembrolizumab), Lartruvo (olaratumab), Lenvima (lenvatinib), Opdivo (nivolumab), Rubraca (rucaparib), Sustol (granisetron), Syndros (dronabinol oral solution), Tecentriq (atezolizumab), Venclexta (venetoclax), Alecensa (alectinib), Cotellic (cobimetinib), Darzalex (daratumumab), Empliciti (elotuzumab), Farydak (panobinostat), Ibrance (palbociclib), Imlygic (talimogene laherparepvec), Lenvima (lenvatinib), Lonsurf (trifluridine and tipiracil), Ninlaro (ixazomib), Odomzo (sonidegib), Onivyde (irinotecan liposome injection), Portrazza (necitumumab), Tagrisso (osimertinib), Unituxin (dinutuximab), Varubi (rolapitant), Vistogard (uridine triacetate), Yondelis (trabectedin), Akynzeo (netupitant and palonosetron), Beleodaq (belinostat), Blincyto (blinatumomab), Cyramza (ramucirumab), Imbruvica (ibrutinib), Lynparza (olaparib), Zydrelig (idelalisib), Zykadia (ceritinib), Gazyva (obinutuzumab), Gilotrif (afatinib), Imbruvica (ibrutinib), Kadcyla (ado-trastuzumab emtansine), Mekinist (trametinib), Pomalyst (pomalidomide), Revlimid (lenalidomide), Stivarga (regorafenib), Tafenlar (dabrafenib), Valchlor (mechlorethamine) gel, Xgeva (denosumab), Xofigo (radium Ra 223 dichloride), Abraxane (paclitaxel protein-bound particles for injectable suspension), Afinitor (everolimus), Afinitor (everolimus), Bosulif (bosutinib), Cometriq (cabozantinib), Erivedge (vismodegib), Iclusig (ponatinib), Inlyta (axitinib), Kyprolis (carfilzomib), Marqibo (vinCRISTine sulfate LIPOSOME injection), Neutroval (tbo-filgrastim), Perjeta (pertuzumab), Picato (ingenol mebutate) gel, Stivarga (regorafenib), Subsys (fentanyl sublingual spray), Synribo (omacetaxine mepesuccinate), Votrient (pazopanib), Xtandi (enzalutamide), Zaltrap (ziv-aflibercept), Abstral (fentanyl sublingual tablets), Adcetris (brentuximab vedotin), Afinitor (everolimus), Erwinaze (asparaginase Erwinia chrysanthemi), Lazanda (fentanyl citrate) nasal spray, Sutent (sunitinib malate), Sylatron (peginterferon alfa-2b), Vandetanib (vandetanib), Xalkori (crizotinib), Yervoy (ipilimumab), Zelboraf (vemurafenib), Zytiga (abiraterone acetate), Halaven (eribulin mesylate),

Herceptin (trastuzumab), Jevtana (cabazitaxel), Provenge (sipuleucel-T), Xgeva (denosumab), Zuplenz (ondansetron oral soluble film), Afinitor (everolimus), Arzerra (ofatumumab), Avastin (bevacizumab), Cervarix [Human Papillomavirus Bivalent (Types 16 and 18) Vaccine, Recombinant, Elitek (rasburicase), Folutyn (pralatrexate injection), Istodax (romidepsin), Onsolis (fentanyl buccal), Votrient (pazopanib), Degarelix (degarelix for injection), Fusilev (levoleucovorin); Mozobil (plerixafor injection), Sancuso (granisetron), Treanda (bendamustine hydrochloride), Evista (raloxifene hydrochloride), Hycamtin (topotecan hydrochloride), Ixempra (ixabepilone), Tassigna (nilotinib hydrochloride monohydrate), Torisel (temsirolimus), Tykerb (lapatinib), Gardasil (quadrivalent human papillomavirus (types 6, 11, 16, 18) recombinant vaccine), Sprycel (dasatinib), Sutent (sunitinib), Vectibix (panitumumab), Arranon (nelarabine), Nexavar (sorafenib), Alimta (pemetrexed for injection), Avastin (bevacizumab); Clolar (clofarabine), Erbitux (cetuximab), Sensipar (cinacalcet), Tarceva (erlotinib, OSI 774), Aloxi (palonosetron), Emend (aprepitant), Iressa (gefitinib); Plenaxis (abarelix for injectable suspension); Premarin (conjugated estrogens); UroXatral (alfuzosin HCl extended-release tablets); Velcade (bortezomib); Eligard (leuprolide acetate); Eloxatin (oxaliplatin/5-fluorouracil/leucovorin); Faslodex (fulvestrant); Gleevec (imatinib mesylate); Neulasta; SecreFlo (secretin); Zevalin (ibritumomab tiuxetan); Zometa (zoledronic acid); Campath; Femara (letrozole); Gleevec (imatinib mesylate); Kytril (granisetron) solution; Trelstar LA (triptorelin pamoate); Xeloda; Zometa (zoledronic acid); Mylotarg (gemtuzumab ozogamicin); Trelstar Depot (triptorelin pamoate); Trisenox (arsenic trioxide); Viadur (leuprolide acetate implant); Aromasin Tablets; Busulfex; Doxil (doxorubicin HCl liposome injection); Ellence; Ethyol (amifostine); Temodar; UVADEX Sterile Solution; Zofran; Actiq; Anzemet; Camptosar; Gemzar (gemcitabine HCL); Herceptin; Inform HER-2/neu breast cancer test; Neupogen; Nolvadex; Photofrin; Proleukin; Sclerosol Intrapleural Aerosol; Valstar; Xeloda; Zofran; Anzemet; Bromfenac; Femara (letrozole); Gliadel Wafer (polifeprosan 20 with carmustine implant); Intron A (interferon alfa-2b, recombinant); Kytril (granisetron) tablets; Lupron Depot (leuprolide acetate for depot suspension); Miraluma test; Neumega; Quadramet (Samarium Sm 153 Lexidronam Injection); Rituxan; Taxol; Anexsia; Aredia (pamidronate disodium for injection); Arimidex (anastrozole); Campostar; Elliotts B Solution (buffered intrathecal electrolyte/dextrose injection); Eulexin (flutamide); Gemzar (gemcitabine HCL); Hycamtin (topotecan hydrochloride); Kadian; Leukine (sargramostim); Lupron Depot (leuprolide acetate for depot suspension); Photodynamic Therapy; Taxotere (Docetaxel); Visipaque (iodixanol); Ethyol

(amifostine); Intron A (Interferon alfa-2b, recombinant), and Leukine (sargramostim). In some embodiments, the TfR-binding peptides of the present disclosure are conjugated to, linked to, or fused to Abemaciclib, Acalabrutinib, Afatinib, Alectinib, Axitinib, Baricitinib, Binimetinib, Bosutinib, Brigatinib, Cabozantinib, Ceritinib, Cobimetinib, Crizotinib, Dabrafenib, dacomitinib, Dasatinib, encorafenib, Erlotinib, Everolimus, Fostamatinib, Gefitinib, Ibrutinib, Imatinib, Lapatinib, Lenvatinib, Lorlatinib, Midostaurin, Neratinib, Nilotinib, Nintedanib, Osimertinib, Palbociclib, Pazopanib, Ponatinib, Regorafenib, Ribociclib, Ruxolitinib, Sirolimus, Sorafenib, Sunitinib, Temsirolimus, Tofacitinib, Trametinib, Vandetanib, Vemurafenib, an IL15, a fused IL15/IL15Ra, an IFN γ , and anti-CD3 agents, or fragment(s) thereof.

[0406] In some embodiments, the TfR-binding peptides as described herein can be used to treat and/or prevent an infection including infections of the central nervous system. In some embodiments, the TfR-binding peptides of the present disclosure are conjugated to, linked to, or fused to an anti-infective agent. Anti-infective agents can include antibiotic, antiviral, and antifungal agents. Anti-infective agent that can be used include artemisinin, Aemcolo (rifamycin), Biktarvy (bictegravir/emtricitabine/tenofovir alafenamide), Nuzyra (omadacycline), Trogarzo (ibalizumab-uiyk), Xofluza (baloxavir marboxil), Baxdela (delafloxacin) tablets and injection, Benznidazole, Giapreza (angiotensin II), Hephisav-B [Hepatitis B Vaccine (Recombinant), Adjuvanted], Juluca (dolutegravir and rilpivirine), KedRab [Rabies Immune Globulin (Human)], Mavyret (glecaprevir and pibrentasvir), Prevymis (letermovir), Shingrix (Zoster Vaccine Recombinant, Adjuvanted), Solosec (secnidazole), Vabomere (meropenem and vaborbactam), Xepi (ozenoxacin), Anthim (obiltoximab), Descovy (emtricitabine and tenofovir alafenamide), Epclusa (sofosbuvir and velpatasvir), Odefsey (emtricitabine, rilpivirine, and tenofovir alafenamide), Vaxchora (Cholera Vaccine, Live, Oral), Vemlidy (tenofovir alafenamide), Zepatier (elbasvir and grazoprevir), Zinplava (bezlotoxumab), Avycaz (ceftazidime-avibactam), Bexsero (Meningococcal Group B Vaccine), Cresemba (isavuconazonium sulfate), Daklinza (daclatasvir), Evotaz (atazanavir and cobicistat), Flud (trivalent influenza vaccine), Genvoya (elvitegravir, cobicistat, emtricitabine, and tenofovir alafenamide), Prezcoibix (darunavir and cobicistat), Technivie, (ombitasvir, paritaprevir and ritonavir), Dalvance (dalbavancin), Harvoni (ledipasvir and sofosbuvir), Impavido (miltefosine), Kerydin (tavaborole), Orbactiv (oritavancin), Rapivab (peramivir injection), Sivextro (tedizolid phosphate), Triumeq (abacavir, dolutegravir, and lamivudine), Zerbaxa (ceftolozane + tazobactam), Flublok (seasonal influenza vaccine), Luzu (luliconazole) Cream 1%, Olysio

(simeprevir), Sitavig (acyclovir) buccal tablets, Sovaldi (sofosbuvir), VariZIG, Varicella Zoster Immune Globulin (Human), Abthrax (raxibacumab), Afinitor (everolimus), Cystaran (cysteamine hydrochloride), Dymista (azelastine hydrochloride and fluticasone propionate), Flucelvax, Influenza Virus Vaccine, Jetrea (ocriplasmin), Linzess (linaclotide), Mytesi (crofelemer), Sirturo (bedaquiline), Sklice (ivermectin) lotion, Stribild (elvitegravir, cobicistat, emtricitabine, tenofovir disoproxil fumarate), Tudorza Pressair (aclidinium bromide inhalation powder), Afinitor (everolimus), Complera (emtricitabine/rilpivirine/tenofovir disoproxil fumarate), Difcicl (fidaxomicin), Edurant (rilpivirine), Eylea (aflibercept), Firazyr (icatibant), Gralise (gabapentin), Incivek (telaprevir), Victrelis (boceprevir), Egrifta (tesamorelin for injection), Menveo (meningitis vaccine), Zymaxid (gatifloxacin ophthalmic solution), Afinitor (everolimus), Bepreve (bepotastine besilate ophthalmic solution), Hiberix (Haemophilus b Conjugate Vaccine, Tetanus Toxoid Conjugate), Vibativ (telavancin), Aptivus (tipranavir), Astepro (azelastine hydrochloride nasal spray), Intelence (etravirine), Patanase (olopatadine hydrochloride), Viread (tenofovir disoproxil fumarate), Isentress (raltegravir), Selzentry (maraviroc), Veramyst (fluticasone furoate), Xyzal (levocetirizine dihydrochloride), Eraxis (anidulafungin), Noxafil (posaconazole), Prezista (darunavir), Tyzeka (telbivudine), Veregen (kunecatechins), Aptivus (tipranavir), Baraclude (entecavir), FluMist (Influenza Virus Vaccine), Fuzeon (enfuvirtide), Lexiva (fosamprenavir calcium), Reyataz (atazanavir sulfate), Botox Cosmetic (botulinum toxin type A), Clarinex, Hepsera (adefovir dipivoxil), Pediarix Vaccine, Pegasys (peginterferon alfa-2a), Sustiva, Vfend (voriconazole), Peg-Intron (peginterferon alfa-2b), Rebetol (ribavirin), Spectracef, Tavist (clemastine fumarate), Twinrix, Valcyte (valganciclovir HCl), Viread (tenofovir disoproxil fumarate), Xigris (drotrecogin alfa [activated]), ABREVA (docosanol), Cefazolin and Dextrose USP, Children's Motrin Cold, REBETRON (TM) Combination Therapy, Synagis, Viroptic, Aldara (imiquimod), Ceftin (cefuroxime axetil), Combivir, Famvir (famciclovir), Floxin otic, Fortovase, INFERGEN (interferon alfacon-1), Intron A (interferon alfa-2b, recombinant), Taxol, Timentin, Trovan, VIRACEPT (nelfinavir mesylate), Zerit (stavudine), AK-Con-A (naphazoline ophthalmic), Allegra (fexofenadine hydrochloride), Atrovent (ipratropium bromide), Augmentin (amoxicillin/clavulanate), Crixivan (Indinavir sulfate), Elmiron (pentosan polysulfate sodium), Havrix, IvyBlock, Leukine (sargramostim), Merrem I.V. (meropenem), Tavist (clemastine fumarate), Videx (didanosine), Viramune (nevirapine), Zithromax (azithromycin), Cedax (ceftibuten), Clarithromycin (Biaxin), Epivir

(lamivudine), Invirase (saquinavir), Valtrex (valacyclovir HCl), Zerit (stavudine), Zyrtec (cetirizine HCl), or a functional fragment thereof.

[0407] In some embodiments, the TfR-binding peptides as described herein can be used to treat and/or prevent an inflammation including inflammation of the central nervous system and the gastrointestinal (GI) tract. In some embodiments, the TfR-binding peptides of the present disclosure are conjugated to, linked to, or fused to an anti-inflammatory agent. Anti-inflammatory agents can include steroidal and nonsteroidal anti-inflammatory agents. Nonsteroidal anti-inflammatory agents can include aspirin, celecoxib, diclofenac, diflunisal, etodolac, ibuprofen, indomethacin, ketoprofen, ketorolac, nabumetone, naproxen, oxaprozin, piroxicam, salsalate, sulindac, tolmetin, piroxicam, flurbiprofen, and fenoprofen. Anti-inflammatory drugs that can be used in combination with the herein disclosed methods and compositions can include corticosteroids and aminosalicylates including but not limited to mesalamine, balsalazide, and olsalazine. In various cases, anti-inflammatory drugs include tumor necrosis factor (TNF)-alpha inhibitors such as infliximab, adalimumab, and golimumab, and biological drug (e.g., antibodies, fragments or derivatives thereof) targeting other biological targets including natalizumab, vedolizumab, and ustekinumab.

[0408] As compared to antibody-drug conjugates (e.g., Adcetris, Kadcyla, Mylotarg) or transferrin, in some aspects the peptide conjugated to, linked to, or fused to an active agent as described herein can exhibit better penetration of solid tumors or of the CNS due to its smaller size. In certain aspects, the peptide conjugated to, linked to, or fused to an active agent as described herein can carry different or higher doses of active agents as compared to antibody-drug conjugates. In some embodiments, the peptide as described conjugated to, linked to, or fused to an active agent as described herein can deliver higher doses of an active agent to skeletal muscle tissue compared to antibody-drug conjugates. Thus, treatment of muscular dystrophy, SBMA, or muscle-derived sarcomas can be of particular interest using the peptides of the present disclosure. In some aspects, the TfR-binding peptides as described herein can exhibit significantly lower plasma half-lives compared to antibodies or antibody-drug conjugates, and thus can allow for a more rapid peripheral clearance (can be advantageous e.g., when radiolabeled conjugates are used providing significantly lower organ dosimetries). In still other aspects, the peptide conjugated to, linked to, or fused to an active agent as described herein can have better site specific delivery of defined drug ratio as compared to antibody-drug conjugates. In other aspects, the peptide can be amenable to solvation in organic solvents (in addition to

water), which can allow more synthetic routes for solvation and conjugation of a drug (which often has low aqueous solubility) and higher conjugation yields, higher ratios of drug conjugated to, linked to, or fused to peptide (versus an antibody), and/or reduce aggregate/high molecular weight species formation during conjugation. Additionally, a unique amino acid residue(s) can be introduced into the peptide via a residue that is not otherwise present in the short sequence or via inclusion of a non-natural amino acid, allowing site specific conjugation to the peptide.

[0409] The peptides or peptide constructs of the present disclosure can also be conjugated to, linked to, or fused to other moieties that can serve other roles, such as providing an affinity handle (e.g., biotin) for retrieval of the peptides from tissues or fluids. For example, peptides or peptide constructs of the present disclosure can also be conjugated to, linked to, or fused to biotin. In addition to extension of half-life, biotin can also act as an affinity handle for retrieval of peptides or peptide constructs from tissues or other locations. In some embodiments, fluorescent biotin conjugates that can act both as a detectable label and an affinity handle can be used. Non limiting examples of commercially available fluorescent biotin conjugates can include Atto 425-Biotin, Atto 488-Biotin, Atto 520-Biotin, Atto-550 Biotin, Atto 565-Biotin, Atto 590-Biotin, Atto 610-Biotin, Atto 620-Biotin, Atto 655-Biotin, Atto 680-Biotin, Atto 700-Biotin, Atto 725-Biotin, Atto 740-Biotin, fluorescein biotin, biotin-4-fluorescein, biotin-(5-fluorescein) conjugate, and biotin-B-phycoerythrin, Alexa fluor 488 biocytin, Alexa fluor 546, Alexa Fluor 549, lucifer yellow cadaverine biotin-X, Lucifer yellow biocytin, Oregon green 488 biocytin, biotin-rhodamine and tetramethylrhodamine biocytin. In some other examples, the conjugates can include chemiluminescent compounds, colloidal metals, luminescent compounds, enzymes, radioisotopes, and paramagnetic labels. In some embodiments, the peptide described herein can also be attached to another molecule. For example, the peptide sequence also can be attached to another active agent (e.g., small molecule, peptide, polypeptide, polynucleotide, antibody, aptamer, cytokine, growth factor, neurotransmitter, an active fragment or modification of any of the preceding agents, fluorophore, radioisotope, radionuclide chelator, acyl adduct, chemical linker, or sugar). In some embodiments, the peptide can be conjugated to, linked to, or fused with, or covalently or non-covalently linked to an active agent.

[0410] In some embodiments, peptides or peptide constructs of the present disclosure can also be conjugated to, linked to, or fused to other affinity handles. Other affinity handles may include genetic fusion affinity handles. Genetic fusion affinity handles may include 6xHis (HHHHHH (SEQ ID NO: 345); immobilized metal affinity column purification possible), FLAG

(DYKDDDDK (SEQ ID NO: 346); anti-FLAG immunoprecipitation), “shorty” FLAG (DYKDE (SEQ ID NO: 347); anti-FLAG immunoprecipitation), hemagglutinin (YPYDVPDYA (SEQ ID NO: 348); anti-HA immunoprecipitation), and streptavidin binding peptide (DVEAWLGAR (SEQ ID NO: 349); streptavidin-mediated precipitation). In some embodiments, peptides or peptide constructs of the present disclosure can also be conjugated to, linked to, or fused to an expression tag or sequence to improve protein expression and/or purification. Such expression tags may include genetic fusion expression tags. Genetic fusion expression tags may include siderocalin (SEQ ID NO: 351,

METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTSDLPAPPLSKVPLQQNFQD
 NQFQGKQWYVVGLAGNAILREDKDPQKMYATIELKEDKSYNVTSVLFRKKKCDYWIR
 TFVPGSQPGFTLGNIKSYPLTSYLVRVVSTNYNQHAMVFFKKVSQNREYFKITLYGRT
 KELTSELKENFIRFSKSLGLPENHIVFPVPIDQCIDGGGSENLYFQ). Exemplary siderocalin
 peptide fusions include human soluble transferrin fused to siderocalin
 (METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTSDLPAPPLSKVPLQQNFQD
 NQFQGKQWYVVGLAGNAILREDKDPQKMYATIELKEDKSYNVTSVLFRKKKCDYWIR
 TFVPGSQPGFTLGNIKSYPLTSYLVRVVSTNYNQHAMVFFKKVSQNREYFKITLYGRT
 KELTSELKENFIRFSKSLGLPENHIVFPVPIDQCIDGGGSENLYFQGSRLYWDDLKRKLSE
 KLDSTDFSTIKLLNENSYVPREAGSQKDENLALYVENQFREFKLSKVWRDQHFVKIQV
 KDSAQNSVIIVDKNGRLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLYTPV
 NGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSFFGHAHLGTGDPYTP
 GFPSFNHTQFPSSRSGLPNIPVQTISRAAAEKLFGNMEGDCPSDWKTDSTCRMVTSSEK
 NVKLTVS NVLKEIKILNIFGVIKGFVEPDHYVVVGAQRDAWGPGAAGSGVGTALLLKL
 QMFSDMVLKDGFPQRSIIIFASWSAGDFGSGVATEWLEGYLSSLHLKAFTYINLDKAVL
 GTSNFKVSASPLLYTLIEKTMQNVKHPVTGQFLYQDSNWASKVEKLTLDNAAFPFLAYS
 GIPAVSFCFCEDTDYPYLGTTMDTYKELIERIPELNKVARAAAEVAGQFVIKLT HDVELN
 LDYERYNSQLLSFVRDLNQYRADIKEMGLSLQWLYSARGDFFRATSRLTTDFGNAEKT
 DRFVMKKLNDRVMRVEYHFLSPYVSPKESPF RHVFWGSGSHTLPALLENLKLKQNNG
 AFNETLFRNQLALATWTIQGAANALSGDVWDIDNEFGGGSHHHHHHGGGSLNDIFEAQ
 KIEWHE; SEQ ID NO: 352) and various peptides of the present disclosure fused to siderocalin,
 such as

METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTSDLPAPPLSKVPLQQNFQD
 NQFQGKQWYVVGLAGNAILREDKDPQKMYATIELKEDKSYNVTSVLFRKKKCDYWIR

TFVPGSQPGFTLGNIKSYPLTSYLVRVVSTNYNQHAMVFFKKVSQNREYFKITLYGRT
 KELTSELKENFIRFSKSLGLPENHIVFPVPIDQCIDGGGSENLYFQGSREGCASRCTKYNA
 ELEKCEARVSSMSNTEETCVQELFDLLHCVDHCVSQ) (SEQ ID NO: 356),
 METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTDLIPAPPLSKVPLQQNFQD
 NQFQGKQWYVVGLAGNAILREDKDPQKMYATIELKEDKSYNVTSVLFRKKKCDYWIR
 TFVPGSQPGFTLGNIKSYPLTSYLVRVVSTNYNQHAMVFFKKVSQNREYFKITLYGRT
 KELTSELKENFIRFSKSLGLPENHIVFPVPIDQCIDGGGSENLYFQGSREGCASRCTKYNA
 ELEKCEARVMSMSNTEEDCEQELEDLLHCLDHCHSQ (SEQ ID NO: 357), and
 (METDTLLLWVLLLWVPGSTGDYKDEHHHHHHHGGSQDSTDLIPAPPLSKVPLQQNFQD
 NQFQGKQWYVVGLAGNAILREDKDPQKMYATIELKEDKSYNVTSVLFRKKKCDYWIR
 TFVPGSQPGFTLGNIKSYPLTSYLVRVVSTNYNQHAMVFFKKVSQNREYFKITLYGRT
 KELTSELKENFIRFSKSLGLPENHIVFPVPIDQCIDGGGSENLYFQGSREGCASRCMKYND
 ELEKCEARMMSMSNTEEDCEQELEDLLYCLDHCHSQ (SEQ ID NO: 358).. Notably CDP-
 NT peptide constructs can be expressed with N-terminal siderocalin (e.g., SEQ ID NO: 359 –
 SEQ ID NO: 361).

[0411] In some embodiments, peptides or peptide constructs of the present disclosure can also be conjugated to, linked to, or fused to other moieties that can serve other roles, such as providing an affinity handle (e.g., biotin) for recruiting a non-conjugated therapeutic to or into the tissue that is targeted such as a tumor. In some cases, the therapeutic is a small molecule, peptide, antibody, or cellular therapeutic (e.g., a CAR T cell or another autologous or allogeneic engineered cell). For example, CAR T cells engineered with an affinity label (e.g., avidin) can be recruited to tissues that are targeted by the TfR-binding peptides that comprise a biotin moiety. In addition, biotin- (or other affinity labeled)-TfR-binding peptides as described herein can be administered (e.g., co-administered) with a variety of different avidin-tagged drugs, such as small molecules, peptides, or proteins that can provide pharmacologic activity in the brain, a tumor, or other tissues that the TfR-binding peptides target or accumulate in.

[0412] Additionally, more than one peptide sequence derived from a toxin or knotted venom protein can be present on, conjugated to, linked to, or fused with a particular peptide. A peptide can be incorporated into a biomolecule by various techniques. A peptide can be incorporated by a chemical transformation, such as the formation of a covalent bond, such as an amide bond. A peptide can be incorporated, for example, by solid phase or solution phase peptide synthesis. A peptide can be incorporated by preparing a nucleic acid sequence encoding the biomolecule,

wherein the nucleic acid sequence includes a subsequence that encodes the peptide. The subsequence can be in addition to the sequence that encodes the biomolecule, or can substitute for a subsequence of the sequence that encodes the biomolecule.

Detectable Agent Peptide Conjugates

[0413] A peptide can be conjugated to, linked to, or fused to an agent used in imaging, research, therapeutics, theranostics, pharmaceuticals, chemotherapy, chelation therapy, targeted drug delivery, and radiotherapy. In some embodiments, a peptide is conjugated to, linked to, or fused with detectable agents, such as a fluorophore, a near-infrared dye, a contrast agent, a nanoparticle, a metal-containing nanoparticle, a metal chelate, an X-ray contrast agent, a PET agent, a metal, a radioisotope, a dye, radionuclide chelator, or another suitable material that can be used in imaging.

[0414] In some embodiments, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 detectable agents can be linked to a peptide. Non-limiting examples of radioisotopes include alpha emitters, beta emitters, positron emitters, and gamma emitters. In some embodiments, the metal or radioisotope is selected from the group consisting of actinium, americium, bismuth, cadmium, cesium, cobalt, europium, gadolinium, iridium, lead, lutetium, manganese, palladium, polonium, radium, ruthenium, samarium, strontium, technetium, thallium, and yttrium. In some embodiments, the metal is actinium, bismuth, lead, radium, strontium, samarium, or yttrium. In some embodiments, the radioisotope is actinium-225 or lead-212. In some embodiments, the near-infrared dyes are not easily quenched by biological tissues and fluids. In some embodiments, the fluorophore is a fluorescent agent emitting electromagnetic radiation at a wavelength between 650 nm and 4000 nm, such emissions being used to detect such agent. Non-limiting examples of fluorescent dyes that could be used as a conjugating molecule in the present disclosure include DyLight-680, DyLight-750, VivoTag-750, DyLight-800, IRDye-800, VivoTag-680, Cy5.5, or indocyanine green (ICG). In some embodiments, near infrared dyes often include cyanine dyes (e.g., Cy7, Cy5.5, and Cy5). Additional non-limiting examples of fluorescent dyes for use as a conjugating molecule in the present disclosure include acradine orange or yellow, Alexa Fluors (e.g., Alexa Fluor 790, 750, 700, 680, 660, and 647) and any derivative thereof, 7-actinomycin D, 8-anilinonaphthalene-1-sulfonic acid, ATTO dye and any derivative thereof, auramine-rhodamine stain and any derivative thereof, bensanthrone, bimane, 9-10-bis(phenylethynyl)anthracene, 5,12-bis(phenylethynyl)naphthalene, bisbenzimidazole, brainbow, calcein, carbodan, fluorescein and any

derivative thereof, 1-chloro-9,10-bis(phenylethynyl)anthracene and any derivative thereof, DAPI, DiOC6, DyLight Fluors and any derivative thereof, epicocconone, ethidium bromide, FLAsH-EDT2, Fluo dye and any derivative thereof, FluoProbe and any derivative thereof, Fluorescein and any derivative thereof, Fura and any derivative thereof, GelGreen and any derivative thereof, GelRed and any derivative thereof, fluorescent proteins and any derivative thereof, m isoform proteins and any derivative thereof such as for example mCherry, hetamethine dye and any derivative thereof, hoeschst stain, iminocoumarin, indian yellow, indo-1 and any derivative thereof, laurdan, lucifer yellow and any derivative thereof, luciferin and any derivative thereof, luciferase and any derivative thereof, mercocyanine and any derivative thereof, Nile dyes and any derivative thereof, perylene, phloxine, phyco dye and any derivative thereof, propium iodide, pyranine, rhodamine and any derivative thereof, ribogreen, RoGFP, rubrene, stilbene and any derivative thereof, sulforhodamine and any derivative thereof, SYBR and any derivative thereof, synapto-pHluorin, tetraphenyl butadiene, tetrasodium tris, Texas Red, Titan Yellow, TSQ, umbelliferone, violanthrone, yellow fluorescent protein and YOYO-1. Other Suitable fluorescent dyes include, but are not limited to, fluorescein and fluorescein dyes (e.g., fluorescein isothiocyanine or FITC, naphthofluorescein, 4', 5'-dichloro-2',7' -dimethoxyfluorescein, 6-carboxyfluorescein or FAM, etc.), carbocyanine, merocyanine, styryl dyes, oxonol dyes, phycoerythrin, erythrosin, eosin, rhodamine dyes (e.g., carboxytetramethyl-rhodamine or TAMRA, carboxyrhodamine 6G, carboxy-X-rhodamine (ROX), lissamine rhodamine B, rhodamine 6G, rhodamine Green, rhodamine Red, tetramethylrhodamine (TMR), etc.), coumarin and coumarin dyes (e.g., methoxycoumarin, dialkylaminocoumarin, hydroxycoumarin, aminomethylcoumarin (AMCA), etc.), Oregon Green Dyes (e.g., Oregon Green 488, Oregon Green 500, Oregon Green 514, etc.), Texas Red, Texas Red-X, SPECTRUM RED, SPECTRUM GREEN, cyanine dyes (e.g., CY-3, Cy-5, CY-3.5, CY-5.5, etc.), ALEXA FLUOR dyes (e.g., ALEXA FLUOR 350, ALEXA FLUOR 488, ALEXA FLUOR 532, ALEXA FLUOR 546, ALEXA FLUOR 568, ALEXA FLUOR 594, ALEXA FLUOR 633, ALEXA FLUOR 660, ALEXA FLUOR 680, etc.), BODIPY dyes (e.g., BODIPY FL, BODIPY R6G, BODIPY TMR, BODIPY TR, BODIPY 530/550, BODIPY 558/568, BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY 630/650, BODIPY 650/665, etc.), IRDyes (e.g., IRD40, IRD 700, IRD 800, etc.), and the like. Additional suitable detectable agents are described in PCT/US14/56177. Non-limiting examples of radioisotopes include alpha emitters, beta emitters, positron emitters, and gamma emitters. In some embodiments, the metal or radioisotope is

selected from the group consisting of actinium, americium, bismuth, cadmium, cesium, cobalt, europium, gadolinium, iridium, lead, lutetium, manganese, palladium, polonium, radium, ruthenium, samarium, strontium, technetium, thallium, and yttrium. In some embodiments, the metal is actinium, bismuth, lead, radium, strontium, samarium, or yttrium. In some embodiments, the radioisotope is actinium-225 or lead-212. Additionally, the following radionuclides can be used for diagnosis and/or therapy: carbon (e.g., ^{11}C or ^{14}C), nitrogen (e.g., ^{13}N), fluorine (e.g., ^{18}F), gallium (e.g., ^{67}Ga or ^{68}Ga), copper (e.g., ^{64}Cu or ^{67}Cu), zirconium (e.g., ^{89}Zr), lutetium (e.g., ^{177}Lu).

[0415] Peptides can be conjugated to, linked to, or fused to a radiosensitizer or photosensitizer. Examples of radiosensitizers include but are not limited to: ABT-263, ABT-199, WEHI-539, paclitaxel, carboplatin, cisplatin, oxaliplatin, gemcitabine, etanidazole, misonidazole, tirapazamine, and nucleic acid base derivatives (e.g., halogenated purines or pyrimidines, such as 5-fluorodeoxyuridine). Examples of photosensitizers can include but are not limited to: fluorescent molecules or beads that generate heat when illuminated, nanoparticles, porphyrins and porphyrin derivatives (e.g., chlorins, bacteriochlorins, isobacteriochlorins, phthalocyanines, and naphthalocyanines), metalloporphyrins, metallophthalocyanines, angelicins, chalcogenapyrrillium dyes, chlorophylls, coumarins, flavins and related compounds such as alloxazine and riboflavin, fullerenes, pheophorbides, pyropheophorbides, cyanines (e.g., merocyanine 540), pheophytins, sapphyrins, texaphyrins, purpurins, porphycenes, phenothiaziniums, methylene blue derivatives, naphthalimides, Nile blue derivatives, quinones, perylenequinones (e.g., hypericins, hypocrellins, and cercosporins), psoralens, quinones, retinoids, rhodamines, thiophenes, verdins, xanthene dyes (e.g., eosins, erythrosins, rose bengals), dimeric and oligomeric forms of porphyrins, and prodrugs such as 5-aminolevulinic acid. Advantageously, this approach can allow for highly specific targeting of diseased cells (e.g., cancer cells) using both a therapeutic agent (e.g., drug) and electromagnetic energy (e.g., radiation or light) concurrently. In some embodiments, the peptide is conjugated to, linked to, fused with, or covalently or non-covalently linked to the agent, e.g., directly or via a linker. Exemplary linkers suitable for use with the embodiments herein are discussed in further detail below.

[0416] In some aspects of the present disclosure, the radionuclide is attached to a peptide or peptide construct as described herein using a chelator. Exemplary chelator moieties can include 2,2',2''-(3-(4-(3-(1-(4-(1,2,4,5-tetrazin-3-yl)phenyl)-1-oxo-5,8,11,14,17,20,23-hepta-oxa-2-

azapentacosan-25-yl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2',2''-(3-(4-(3-(1-(4-(1,2,4,5-tetrazin-3-yl)phenyl)-1-oxo-5,8,11,14,17,20,23,26,29,32,35-undecaoxa-2-azaheptatriacontan-37-yl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2'-(7-(4-(3-(1-(4-(1,2,4,5-tetrazin-3-yl)phenyl)-1-oxo-5,8,11,14,17,20,23,26,29,32,35-undecaoxa-2-azaheptatriacontan-37-yl)thioureido)benzyl)-1,4,7-triazonane-1,4-diyl)diacetic acid; 2,2',2''-(3-(4-(3-(1-(4-(1,2,4,5-tetrazin-3-yl)phenyl)-3,7-dioxo-11,14,17,20,23,26,29-hepta-2,8-diazahentriacontan-31-yl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2',2''-(3-(4-(3-(1-(4-(1,2,4,5-tetrazin-3-yl)phenyl)-3,7-dioxo-11,14,17,20,23,26,29,32,35,38,41-undecaoxa-2,8-diazatritetracontan-43-yl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2',2''-(3-(4-(3-(25,28-dioxo-28-((6-(6-(pyridin-2-yl)-1,2,4,5-tetrazin-3-yl)pyridin-3-yl)amino)-3,6,9,12,15,18,21-hepta-24-aza-octacosyl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2',2''-(3-(4-(3-(37,40-dioxo-40-((6-(6-(pyridin-2-yl)-1,2,4,5-tetrazin-3-yl)pyridin-3-yl)amino)-3,6,9,12,15,18,21,24,27,30,33-undecaoxa-36-azatetracontyl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2',2''-(3-(4-(1-(4-(6-methyl-1,2,4,5-tetrazin-3-yl)phenyl)-3-oxo-6,9,12,15,18,21,24-hepta-2-azaheptacosan-27-amido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2',2''-(2-(4-(1-(4-(6-methyl-1,2,4,5-tetrazin-3-yl)phenoxy)-3,6,9,12,15,18,21,24,27,30,33-undecaoxahexatriacontan-36-amido)benzyl)-1,4,7-triazonane-1,4,7-triyl)triacetic acid; 2,2',2''-(3-(4-(3-(5-amino-6-((4-(6-methyl-1,2,4,5-tetrazin-3-yl)benzyl)amino)-6-oxohexyl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; 2,2'-(7-(4-(3-(5-amino-6-((4-(6-methyl-1,2,4,5-tetrazin-3-yl)benzyl)amino)-6-oxohexyl)thioureido)benzyl)-1,4,7-triazonane-1,4-diyl)diacetic acid; 2,2',2''-(3-(4-(3-(5-amino-6-((5-amino-6-((4-(6-methyl-1,2,4,5-tetrazin-3-yl)benzyl)amino)-6-oxohexyl)amino)-6-oxohexyl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid; and 2,2',2''-(3-(4-(3-(5-amino-6-((5-amino-6-((5-amino-6-((4-(6-methyl-1,2,4,5-tetrazin-3-yl)benzyl)amino)-6-oxohexyl)amino)-6-oxohexyl)amino)-6-oxohexyl)thioureido)benzyl)-1,4,7-triazonane-2,5,8-triyl)triacetic acid.

Linkers

[0417] Peptides according to the present disclosure can be attached to a cargo molecule (e.g., a therapeutically active agent or a detectable agent), such as a small molecule, another peptide, a protein, an antibody, an antibody fragment, an aptamer, polypeptide, polynucleotide, a fluorophore, a radioisotope, a radionuclide chelator, a polymer, a biopolymer, a fatty acid, an

acyl adduct, a chemical linker, or sugar or other active agent or detectable agent described herein through a linker, or directly in the absence of a linker. In case a linker is absent, a cargo molecule is conjugated, linked, or fused directly to the N-terminus and/or the C-terminus of a peptide. In cases where a linker is present, the linker can be a cleavable or stable linker is used. Cleavable linkers can be used for *in vivo* delivery of a cargo molecule, as a linker can be cleaved upon entry in a tissue, cell, endosome, or a nucleus. Stable linkers can be used for delivery of a cargo molecule that is active when conjugated or is degraded such as by catabolism. As described herein, a linker can also be used to covalently attach a TfR-binding peptide to another a cargo moiety having a separate function, such a targeting, cytotoxic, therapeutic, homing, imaging, or diagnostic functions.

[0418] In some embodiments, a cargo molecule can be conjugated to, linked to, or fused to the N-terminus and/or the C-terminus of a peptide to create an active agent or detectable agent peptide construct. In some embodiments, the TfR-targeting peptide can be attached at the N-terminus, an internal lysine, glutamic acid, or aspartic acid residue, or the C-terminus of another functional peptide or protein via a linker. In some embodiments, the peptide can be attached to another functional peptide or protein via a side chain, such as the side chain of a lysine, serine, threonine, cysteine, tyrosine, aspartic acid, a non-natural amino acid residue, or glutamic acid residue. As used herein, molecules can be linked, attached, or fused to each other via an amide bond, an ester bond, an ether bond, a carbamate bond, a carbonate bond, a carbon-nitrogen bond, a triazole, a macrocycle, an oxime bond, a thioester bond, a thioether bond a hydrazone bond, an azo bond, a carbon-carbon single, double, or triple bond, a disulfide bond, a two carbon bridge between two cysteines, a three carbon bridge between two cysteines, or a thioether bond. In some embodiments, the peptide comprises one or more non-natural amino acid, wherein the non-natural amino acid can be an insertion, appendage, or substitution for another amino acid. In some embodiments, similar regions of the disclosed peptide(s) itself (such as a terminus of the amino acid sequence, an amino acid side chain, such as the side chain of a lysine, serine, threonine, cysteine, tyrosine, aspartic acid, a non-natural amino acid residue, or glutamic acid residue, via an amide bond, an ester bond, an ether bond, a carbamate bond, a carbon-nitrogen bond, a triazole, a macrocycle, an oxime bond, a hydrazone bond, a carbon-carbon single, double, or triple bond, a disulfide bond, a thioether bond, or other linker as described herein) can be used to link other molecules. In some embodiments, the peptide is attached to a terminus of the amino acid sequence of a larger polypeptide or peptide molecule, or is attached to a side

chain, such as the side chain of a lysine, serine, threonine, cysteine, tyrosine, aspartic acid, a non-natural amino acid residue, or glutamic acid residue.

[0419] Attachment via a linker can involve incorporation of a linker moiety between the other molecule (e.g., an active agent and/or a detectable agent) and the peptide. The peptide and the other molecule can both be covalently attached to the linker. The linker can be cleavable, stable, self-immolating, hydrophilic, or hydrophobic. The linker can have bulky side groups or chains that sterically limit access of enzymes, water, or other chemicals to the linking group. The linker can have at least two functional groups, one bonded to the other molecule, and one bonded to the peptide, and a linking portion between the two functional groups. Some example linkers are described in Jain, N., *Pharm Res.* 32(11): 3526–40 (2015), Doronina, S.O., *Bioconj Chem.* 19(10): 1960–3 (2008), Pillow, T.H., *J Med Chem.* 57(19): 7890–9 (2014), Dorywalksa, M., *Bioconj Chem.* 26(4): 650–9 (2015), Kellogg, B.A., *Bioconj Chem.* 22(4): 717–27 (2011), and Zhao, R.Y., *J Med Chem.* 54(10): 3606–23 (2011).

[0420] Non-limiting examples of the functional groups for attachment can include functional groups capable of forming, for example, an amide bond, an ester bond, an ether bond, a carbonate bond, a carbamate bond, a carbon-nitrogen bond, a triazole, a macrocycle, an oxime bond, a hydrazone bond, a carbon-carbon single, double, or triple bond, a disulfide bond or a thioether bond. Non-limiting examples of functional groups capable of forming such bonds can include amino groups; carboxyl groups; hydroxyl groups; aldehyde groups; azide groups; alkyne and alkene groups; ketones; hydrazides; hydrazines; acid halides such as acid fluorides, chlorides, bromides, and iodides; acid anhydrides, including symmetrical, mixed, and cyclic anhydrides; carbonates; carbonyl functionalities bonded to leaving groups such as cyano, succinimidyl, and *N*-hydroxysuccinimidyl; maleimides; linkers containing maleimide groups that are designed to hydrolyze; maleimidocaproyl; MCC ([*N*-maleimidomethyl]cyclohexane-1-carboxylate); *N*-ethylmaleimide; maleimide alkane; mc-vc-PABC; DUBA (DuocarmycinhydroxyBenzamide-Azaindole linker); SMCC Succinimidyl-4-(*N*-maleimidomethyl) cyclohexane-1-carboxylate; SPDP (*N*-succinimidyl-3-(2-pyridyldithio) propionate); SPDB *N*-succinimidyl-4-(2-pyridyldithio) butanoate; sulfo-SPDB *N*-succinimidyl-4-(2-pyridyldithio) -2-sulfo butanoate; SPP *N*-succinimidyl 4-(2-pyridyldithio)pentanoate; a dithiopyridylmaleimide (DTM); a hydroxylamine, a vinyl-halo group; haloacetamido groups; bromoacetamido; hydroxyl groups; sulfhydryl groups; and molecules possessing, for example,

alkyl, alkenyl, alkynyl, allylic, or benzylic leaving groups, such as halides, mesylates, tosylates, triflates, epoxides, phosphate esters, sulfate esters, and besylates.


[0421] Non-limiting examples of the linking portion can include alkylene, alkenylene, alkynylene, polyether, such as polyethylene glycol (PEG), hydroxy carboxylic acids, polyester, polyamide, polyamino acids, polypeptides, cleavable peptides, Val-Cit, Phe-Lys, Val-Lys, Val-Ala, other peptide linkers as given in Doronina et al., 2008, linkers cleavable by beta glucuronidase, linkers cleavable by a cathepsin or by cathepsin B, D, E, H, L, S, C, K, O, F, V, X, or W, Val-Cit-p-aminobenzyloxycarbonyl, glucuronide-MABC, valine-citrulline, aminobenzylcarbamates, D-amino acids, and polyamine, any of which being unsubstituted or substituted with any number of substituents, such as halogens, hydroxyl groups, sulfhydryl groups, amino groups, nitro groups, nitroso groups, cyano groups, azido groups, sulfoxide groups, sulfone groups, sulfonamide groups, carboxyl groups, carboxaldehyde groups, imine groups, alkyl groups, halo-alkyl groups, alkenyl groups, halo-alkenyl groups, alkynyl groups, halo-alkynyl groups, alkoxy groups, aryl groups, aryloxy groups, aralkyl groups, arylalkoxy groups, heterocyclyl groups, acyl groups, acyloxy groups, carbamate groups, amide groups, urethane groups, epoxides, charged groups, zwitterionic groups, and ester groups. Other non-limiting examples of reactions to link, fuse, or conjugate molecules together include click chemistry, copper-free click chemistry, HIPS ligation, Staudinger ligation, and hydrazine-iso-Pictet-Spengler.

[0422] In some embodiments, a linker for linking a repressor domain to a dCas9 can be about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, or 50 amino acid residues in length. A linker can be about 10 amino acid residues in length. A linker can be about 11 amino acid residues in length. A linker can be about 12 amino acid residues in length. A linker can be about 13 amino acid residues in length. A linker can be about 14 amino acid residues in length. A linker can be about 15 amino acid residues in length. A linker can be about 16 amino acid residues in length. A linker can be about 17 amino acid residues in length. A linker can be about 18 amino acid residues in length. A linker can be about 19 amino acid residues in length. A linker can be about 20 amino acid residues in length. A linker can be about 21 amino acid residues in length. A linker can be about 22 amino acid residues in length. A linker can be about 23 amino acid residues in length. A linker can be about 24 amino acid residues in length. A linker can be about 25 amino acid residues in length. A linker can be about 26 amino acid residues in length. A linker can be about 27 amino acid residues in length. A

linker can be about 28 amino acid residues in length. A linker can be about 29 amino acid residues in length. A linker can be about 30 amino acid residues in length.

[0423] In some embodiments, a linker of the present disclosure can comprise a cleavable or stable linker moiety. In some embodiments, cleavable linkers of the present disclosure can include, for example, protease cleavable peptide linkers, nuclease sensitive nucleic acid linkers, lipase sensitive lipid linkers, glycosidase sensitive carbohydrate linkers, pH sensitive linkers, hypoxia sensitive linkers, photo-cleavable linkers, heat-labile linkers, enzyme cleavable linkers (e.g., esterase cleavable linker), ultrasound-sensitive linkers, and X-ray cleavable linkers. In some embodiments, the linker is not a cleavable linker.



[0424] Non-limiting examples of linkers include:



is independently 0 to about 1,000; 1 to about 1,000; 0 to about 500; 1 to about 500; 0 to about 250; 1 to about 250; 0 to about 200; 1 to about 200; 0 to about 150; 1 to about 150; 0 to about 100; 1 to about 100; 0 to about 50; 1 to about 50; 0 to about 40; 1 to about 40; 0 to about 30; 1 to about 30; 0 to about 25; 1 to about 25; 0 to about 20; 1 to about 20; 0 to about 15; 1 to about 15; 0 to about 10; 1 to about 10; 0 to about 5; or 1 to about 5. In some embodiments, each n is independently 0, about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, or about 50. In some embodiments, m is 1 to about 1,000; 1 to about 500; 1 to about 250; 1 to about 200; 1 to about 150; 1 to about 100; 1 to about 50; 1 to about 40; 1 to about 30; 1 to about 25; 1 to about 20; 1 to about 15; 1 to about 10; or 1 to about 5. In some embodiments, m is 0, about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about


9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, or about 50, or any linker as disclosed in Jain, N., Pharm Res. 32(11): 3526-40 (2015) or Ducry, L., Antibody Drug Conjugates (2013).


[0425] In some cases, a linker can comprise a cyclic group, such as an organic nonaromatic or aromatic ring, optionally with 3-10 carbons in the ring, optionally built from a carboxylic acid,



or a substituted analog or a stereoisomer thereof. This linker can optionally be used to form a carbamate linkage. In some cases, a carbamate linkage can be more resistant to cleavage, such as by hydrolysis, enzymes such as esterases, or other chemical reactions, than an ester linkage.


[0426] In some cases, a linker can comprise a cyclic carboxylic acid, for example a cyclic dicarboxylic acid, for example one of the following groups: 1,4-cyclohexane dicarboxylic acid, 1,2-cyclohexane dicarboxylic acid, or 1,3-cyclohexane dicarboxylic acid, 1,1-cyclopentanediacetic acid,






. In some instances, the linker can optionally be used to form an ester linkage. In some cases, a cyclic ester linkage can be more sterically resistant to cleavage, such as by hydrolysis by water, enzymes such as esterases, or other chemical reactions, than a noncyclic or linear ester linkage.

[0427] In some cases, a linker can comprise an aromatic dicarboxylic acid, for example terephthalic acid, isophthalic acid, phthalic acid




or a substituted analog thereof.

[0428] In some cases, a linker can comprise a natural or non-natural amino acid, for example



cysteine, or a substituted analog or a stereoisomer thereof. In some instances, a linker can comprise alanine (A, Ala); arginine (R, Arg); asparagine (N, Asn); aspartic acid (D, Asp); glutamic acid (E, Glu); glutamine (Q, Gln); glycine (G, Gly); histidine (H, His); isoleucine (I, Ile); leucine (L, Leu); lysine (K, Lys); methionine (M, Met); phenylalanine (F, Phe); proline (P, Pro); serine (S, Ser); threonine (T, Thr); tryptophan (W, Trp); tyrosine (Y, Tyr); valine (V, Val); or any plurality or combination thereof. In some embodiments, the non-natural amino acid can comprise one or more functional groups, e.g., alkene or alkyne, that can be used as functional handles.


[0429] In some cases, a linker can comprise one of the following groups:



$n1 = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, \text{ or } 10$
 $n2 = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, \text{ or } 10$

or a substituted analog or a stereoisomer


thereof. In some instances, the linker is selected from one of the following groups:



or a substituted analog or a

stereoisomer thereof.


[0430] In some cases, a linker can comprise one of the following groups:



$n1 = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, \text{ or } 10$
 $n2 = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, \text{ or } 10$

or a substituted analog or a stereoisomer

thereof. In some instances, the linker is selected from one of the following groups:





or a substituted analog or a

stereoisomer thereof.


[0431] In some cases, a substituted analog or a stereoisomer is a structural analog of a compound disclosed herein, for which one or more hydrogen atoms of the compound can be substituted by one or more groups of halo (e.g., Cl, F, Br), alkyl (e.g., methyl, ethyl, propyl), alkenyl, alkynyl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroalkyl, heterocycloalkyl, or any combination thereof. In some cases, a stereoisomer can be an enantiomer, a diastereomer, a cis or trans stereoisomer, a E or Z stereoisomer, or a R or S stereoisomer.

[0432] Non-limiting examples of linear linkers include;





wherein each n1, or n2 or m is independently 0 to about 1,000; 1 to about 1,000; 0 to about 500; 1 to about 500; 0 to about 250; 1 to about 250; 0 to about 200; 1 to about 200; 0 to about 150; 1 to about 150; 0 to about 100; 1 to about 100; 0 to about 50; 1 to about 50; 0 to about 40; 1 to about 40; 0 to about 30; 1 to about 30; 0 to about 25; 1 to about 25; 0 to about 20; 1 to about 20; 0 to about 15; 1 to about 15; 0 to about 10; 1 to about 10; 0 to about 5; or 1 to about 5. In some embodiments, each n is independently 0, about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, or about 50. In some embodiments, m is 1 to about 1,000; 1 to about 500; 1 to about 250; 1 to about 200; 1 to about 150; 1 to about 100; 1 to about 50; 1 to about 40; 1 to about 30; 1 to about 25; 1 to about 20; 1 to about 15; 1 to about 10; or 1 to about 5. In some embodiments, m is 0, about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, or about 50. In some instances, the linker can comprise a linear dicarboxylic acid, e.g., one of the following groups: succinic acid, 2,3-dimethylsuccinic acid, glutaric acid, adipic acid, 2,5-dimethyladipic acid,







or a substituted analog or a stereoisomer thereof. In some cases, the linker can be used to form a










carbamate linkage. In some embodiments, the carbamate linkage can be more resistant to cleavage, such as by hydrolysis, enzymes such as esterases, or other chemical reactions, than an ester linkage. In some cases, the linker can be used to form a linear ester linkage. In some embodiments, the linear ester linkage can be more susceptible to cleavage, such as by hydrolysis, enzymes such as esterases, or other chemical reactions, than a cyclic ester or carbamate linkage. Side chains such as methyl groups on the linear ester linkage can optionally make the linkage less susceptible to cleavage than without the side chains.









[0433] In some cases a linker can be a succinic linker, and a drug can be attached to a peptide via an ester bond or an amide bond with two methylene carbons in between. In other cases, a linker can be any linker with both a hydroxyl group and a carboxylic acid, such as hydroxy hexanoic acid or lactic acid.

[0434] In some cases, a drug can be attached to a peptide using any one or more of the linkers shown below in TABLE 7.

TABLE 7 – Exemplary Linkers for Use in Peptide-Drug Conjugates

Compound number	Chemical structure	Linker is based on:
1		Carbonic acid
2		<i>trans</i> -1,4-cyclohexanedicarboxylic acid
3		1,4-cyclohexanedicarboxylic acid
4		<i>trans</i> -1-aminomethylamine-cyclohexane-4-carboxylic acid (carbamate)
5		1-aminomethylamine-cyclohexane-4-carboxylic acid

Compound number	Chemical structure	Linker is based on:
6		4-(carboxyoxymethyl)-cyclohexane-1-carboxylic acid
7		4-(carboxyamino)-cyclohexane-1-carboxylic acid
8		<i>trans</i> -4-(carboxyamino)-cyclohexane-1-carboxylic acid
9		4-(carboxyamino)benzoic acid
10		4-(carboxyoxymethyl)benzoic acid
11		2-(4-(carboxyoxymethyl)phenyl)acetic acid
12		4-(((carboxyoxymethyl)methyl)benzoic acid
13		2,2'-(cyclopentane-1,1-diyl)diacetic acid
14		<i>cis</i> -5-norbornene-endo-2,3-dicarboxylic acid

Compound number	Chemical structure	Linker is based on:
15		(3-amino-3-oxypropyl)carbamic acid
16		[¹⁴ C]-Cysteine
17		Cysteine
18		Succinic acid
19		Glutaric acid
20		Adipic acid
21		2,5-dimethyladipic acid (DMA)
22		<i>trans</i> -β-hydromuconic acid

[0435] In some cases, a drug molecule is attached to a linker wherein a nucleophilic functional group (e.g., a hydroxyl group) of the drug molecule acts as the nucleophile and replaces a leaving group on the linker moiety, thereby attaching the drug to the linker.

[0436] In other cases, a drug molecule is attached to a linker wherein a nucleophilic functional group (e.g., thiol group, amine group, etc) of the linker replaces a leaving group on the drug molecule, thereby attaching the drug to the linker. Such leaving group (or functional group that may be converted into a leaving group) may be a primary alcohol to form a thioether bond, thereby attaching the drug to the linker. A primary alcohol can be converted into a leaving group such as a mesylate, a tosylate, or a nosylate in order to accelerate the nucleophilic substitution reaction.

[0437] The peptide-drug conjugates of the present disclosure can comprise a drug, a linker, and/or a peptide of the present disclosure. A general connectivity between these three components can be drug-linker-peptide, such that the linker is attached to both the drug and the peptide. In many cases, the peptide is attached to a linker via an amide bond. Amide bonds can be relatively stable (e.g., *in vivo*) compared to other bonds described herein, such as esters, carbonates, etc. The amide bond between the peptide and the linker may thus provide advantageous properties due to its *in vivo* stability if the drug is sought to be cleaved from a peptide-drug-conjugate without the linker being attached to the drug after such *in vivo* cleavage. Thus, in various cases, a drug is attached to the linker-peptide moiety via linkages such as ester, carbonate, carbamate, etc., wherein the peptide is attached to the linker via an amide bond. This can allow for selective cleavage of the drug-linker bond (as opposed to the linker-peptide bond) allowing the drug to be released without a linker moiety attached to it after cleavage. The use of such different drug-linker bonds or linkages can allow the modulation of drug release *in vivo*, e.g., in order to achieve a therapeutic function while minimizing off-target effects (e.g., reduction in drug release during circulation).

[0438] The linker can be a cleavable or a stable linker. The use of a cleavable linker permits release of the conjugated moiety (e.g., a therapeutic agent) from the peptide, e.g., after targeting to the target tissue or cell or subcellular compartment. In some cases the linker is enzyme cleavable, e.g., a valine-citrulline linker that can be cleavable by cathepsin, or an ester linker that can be cleavable by esterase. In some embodiments, the linker contains a self-immolating portion. In other embodiments, the linker includes one or more cleavage sites for a specific protease, such as a cleavage site for matrix metalloproteases (MMPs), thrombin, urokinase-type plasminogen activator, or cathepsin (e.g., cathepsin K).

[0439] Thus, in some cases, a peptide-active agent conjugate of the present disclosure can comprise one or more, about two or more, about three or more, about five or more, about ten or more, or about 15 or more amino acids that can form an amino acid sequence cleavable by an enzyme. Such enzymes can include proteinases. A peptide-drug conjugate can comprise an amino acid sequence that can be cleaved by a Cathepsin, a Chymotrypsin, an Elastase, a Subtilisin, a Thrombin I, or a Urokinas, or any combination thereof.

[0440] Alternatively or in combination, the cleavable linker can be cleaved, dissociated, or broken by other mechanisms, such as via pH, reduction, or hydrolysis. Hydrolysis can occur directly due to water reaction, or be facilitated by an enzyme, or be facilitated by presence of

other chemical species. A hydrolytically labile linker, (amongst other cleavable linkers described herein) can be advantageous in terms of releasing active agents from the peptide. For example, an active agent in a conjugate form with the peptide may not be active, but upon release from the conjugate after targeting to the target tissue or cell or subcellular compartment, the active agent is active. The cleaved active agent may retain the chemical structure of the active agent before cleavage, or may be modified. In some embodiments, a stable linker may optionally not cleave in buffer over extended periods of time (e.g., hours, days, or weeks). In some embodiments, a stable linker may optionally not cleave in body fluids such as plasma or synovial fluid over extended periods of time (e.g., hours, days, or weeks). In some embodiments, a stable linker optionally may cleave, such as after exposure to enzymes, reactive oxygen species, other chemicals or enzymes that may be present in cells (such as macrophages), cellular compartments (such as endosomes and lysosomes), inflamed areas of the body (such as inflamed joints), or tissues or body compartments. In some embodiments, a stable linker may optionally not cleave in vivo but present an active agent that is still active when conjugated to, linked to, or fused to the peptide.

[0441] The rate of hydrolysis of the linker (e.g., a linker of a peptide conjugate) can be tuned. For example, the rate of hydrolysis of linkers with unhindered esters may be faster compared to the hydrolysis of linkers with bulky groups next an ester carbonyl. A bulky group can be a methyl group, an ethyl group, a phenyl group, a ring, or an isopropyl group, or any group that provides steric bulk. In another example, the rate of hydrolysis can be faster with hydrophilic groups, such as alcohols, acids, or ethers, or near an ester carbonyl. In another example, hydrophobic groups present as side chains or by having a longer hydrocarbon linker can slow cleavage of the ester. In some embodiments, cleavage of a carbamate group can also be tuned by hindrance, hydrophobicity, and the like. In another example, using a less labile linker, such as a carbamate rather than an ester, can slow the cleavage rate of the linker. In some cases, the steric bulk can be provided by the drug itself, such as by ketorolac when conjugated via its carboxylic acid. The rate of hydrolysis of the linker can be tuned according to the residency time of the conjugate in the target tissue or cell or subcellular compartment, according to how quickly the peptide accumulates in the target tissue or cell or subcellular compartment, or according to the desired time frame for exposure to the active agent in the target tissue or cell or subcellular compartment. For example, when a peptide is cleared from the target tissue or cell or subcellular compartment relatively quickly, the linker can be tuned to rapidly hydrolyze. In contrast, for example, when a peptide has a longer residence time in the target tissue or cell or subcellular

compartment, a slower hydrolysis rate can allow for extended delivery of an active agent. This can be important when the peptide is used to deliver a drug to the target tissue or cell or subcellular compartment (e.g., a tumor cell or a tumor tissue). “Programmed hydrolysis in designing paclitaxel prodrug for nanocarrier assembly” Sci Rep 2015, 5, 12023 Fu et al., provides an example of modified hydrolysis rates. In some embodiments, rates of cleavage can vary by species, body compartment, and disease state. For instance, cleavage by esterases may be more rapid in rat or mouse plasma than in human plasma, such as due to different levels of carboxyesterases. In some embodiments, a linker may be tuned for different cleavage rates for similar cleavage rates in different species.

[0442] In some cases a linker can be a succinic linker, and a drug can be attached to a peptide via an ester bond or an amide bond with two methylene carbons in between. In other cases, a linker can be any linker with both a hydroxyl group and a carboxylic acid, such as hydroxy hexanoic acid or lactic acid.

[0443] In some embodiments, the linker can release the active agent in an unmodified form. In other embodiments, the active agent can be released with chemical modification. In still other embodiments, catabolism can release the active agent still linked to parts of the linker and/or peptide.

[0444] The linker can be a stable linker or a cleavable linker. In some embodiments, the stable linker can slowly release the conjugated moiety by an exchange of the conjugated moiety onto the free thiols on serum albumin. In some embodiments, the use of a cleavable linker can permit release of the conjugated moiety (e.g., a therapeutic agent) from the peptide, e.g., after administration to a subject in need thereof. In other embodiments, the use of a cleavable linker can permit the release of the conjugated therapeutic from the peptide. In some cases the linker is enzyme cleavable, e.g., a valine-citrulline linker. In some embodiments, the linker contains a self-immolating portion. In other embodiments, the linker includes one or more cleavage sites for a specific protease, such as a cleavage site for matrix metalloproteases (MMPs), thrombin, cathepsins, peptidases, or beta-glucuronidase. Alternatively or in combination, the linker is cleavable by other mechanisms, such as via pH, reduction, or hydrolysis.

[0445] The rate of hydrolysis or reduction of the linker can be fine-tuned or modified depending on an application. For example, the rate of hydrolysis of linkers with unhindered esters can be faster compared to the hydrolysis of linkers with bulky groups next to an ester carbonyl. A bulky group can be a methyl group, an ethyl group, a phenyl group, a ring, or an isopropyl group, or

any group that provides steric bulk. In some cases, the steric bulk can be provided by the drug itself, such as by ketorolac when conjugated, linked, or fused via its carboxylic acid. The rate of hydrolysis of the linker can be tuned according to the residency time of the conjugate or fusion in the target location. For example, when a peptide is cleared from a tumor, or the brain, relatively quickly, the linker can be tuned to rapidly hydrolyze. When a peptide has a longer residence time in the target location, a slower hydrolysis rate would allow for extended delivery of an active agent.

[0446] The rate of hydrolysis of the linker (e.g., a linker of a peptide conjugate) can be measured. Such measurements can include determining free active agent in plasma, or synovial fluid, or other fluid or tissue of a subject *in vivo* and/or by incubating a linker or a peptide conjugate comprising a linker of the present disclosure with a buffer (e.g., PBS) or blood plasma from a subject (e.g., rat plasma, human plasma, etc.) or synovial fluid or other fluids or tissues *ex vivo*. The methods for measuring hydrolysis rates can include taking samples during incubation or after administration and determine free active agent, free peptide, or any other parameter indicate of hydrolysis, including also measuring total peptide, total active agent, or conjugated active agent-peptide. The results of such measurements can then be used to determine a hydrolysis half-life of a given linker or peptide conjugate comprising the linker. A hydrolysis half-life of a linker can differ depending on the plasma or fluid or species or other conditions used to determine such half-life. This can be due to certain enzymes or other compounds present in a certain plasma (e.g., rat plasma). For instance, different fluids (such as plasma or synovial fluid) can contain different amounts of enzymes such as esterases, and these levels of these compounds can also vary depending on species (such as rat versus human) as well as disease state (such as normal versus arthritic).

[0447] The conjugates of the present disclosure can be described as having a modular structure comprising various components, wherein each of the components (e.g., peptide, linker, active agent and/or detectable agent) can be selected dependently or independently of any other component. For example, a conjugate for use in the treatment of pain can comprise a TfR-binding peptide of the present disclosure (e.g., those having the amino acid sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358), a linker (e.g., any linker described in **TABLE 7**) and an active agent (e.g., an NSAID pain reliever, ibuprofen, or a molecule to treat disorders of the CNS or of cancer). The linker, for example, can

be selected and/or modified to achieve a certain active agent release (e.g., a certain release rate) via a certain mechanism (e.g., via hydrolysis, such as enzyme and/or pH-dependent hydrolysis) at the target site (e.g., in the brain) and/or to minimize systemic exposure to the active agent. During the testing of a conjugate any one or more of the components of the conjugate can be modified and/or altered to achieve certain *in vivo* properties of the conjugate, e.g., pharmacokinetic (e.g., clearance time, bioavailability, uptake and retention in various organs) and/or pharmacodynamic (e.g., target engagement) properties. Thus, the conjugates of the present disclosure can be modulated to prevent, treat, and/or diagnose a variety of diseases and conditions, while reducing side effects (e.g., side effects that occur if such active agents are administered alone (i.e., not conjugated to a peptide)).


[0448] In some embodiments, the non-natural amino acid can comprise one or more functional groups, e.g., alkene or alkyne, that can be used as functional handles. For example, a multiple bond of such functional groups can be used to add one or more molecules to the conjugate. The one or more molecules can be added using various synthetic strategies, some of which may include addition and/or substitution chemistries. For example, an addition reaction using a multiple bond can comprise the use of hydrobromic acid, wherein the bromine can act as a leaving group and thus be substituted with various moieties, e.g., active agents, detectable agents, agents that can modify or alter the pharmacokinetic (e.g., plasma half-life, retention and/or uptake in central nervous system (CNS) or elsewhere) and/or pharmacodynamic (e.g., hydrolysis rate such as an enzymatic hydrolysis rate) properties of the conjugate.

[0449] In some embodiments, a conjugate as described herein comprises one or more non-natural amino acid and/or one or more linkers. Such one or more non-natural amino acid and/or one or more linkers can comprise one or more functional groups, e.g., alkene or alkyne (e.g., non-terminal alkenes and alkynes), which can be used as functional handles. For example, a multiple bond of such functional groups can be used to add one or more molecules to the conjugate. The one or more molecules can be added using various synthetic strategies, some of which may include addition and/or substitution chemistries, cycloadditions, etc. For example, an addition reaction using a multiple bond can comprise the use of hydrogen bromide (e.g., via hydrohalogenation reactions), wherein the bromide substituent, once attached, can act as a leaving group and thus be substituted with various moieties comprising a nucleophilic functional groups, e.g., active agents, detectable agents, agents. As another example, a multiple bond can be used as a functional handle in a cycloaddition reaction. Cycloaddition reactions

can comprise 1,3-dipolar cycloadditions, [2+2]-cycloadditions (e.g., photocatalyzed), Diels-Alder reactions, Huisgen cycloadditions, nitron-olefin cycloadditions, etc. Such cycloaddition reactions can be used to attached various functional groups, functional moieties, active agents, detectable agents, and so forth to the conjugate. For example, a 1,3-dipolar cycloaddition reaction can be used to attach a molecule to a conjugate, wherein the molecule comprises a 1,3-dipole that can react with, e.g., an alkyne to form a 5-membered ring, thereby attaching said molecule to the conjugate.

[0450] The addition of such agents or molecules (e.g., via nucleophilic or electrophilic addition followed by nucleophilic substitution) can have various application. For example, attaching such molecule or agent can modify or alter the pharmacokinetic (e.g., plasma half-life, retention and/or uptake in CNS or biodistribution) and/or pharmacodynamic (e.g., hydrolysis rate such as an enzymatic hydrolysis rate) properties of the conjugate. Attaching such molecule or agent can also alter (e.g., increase) the depot effect of a conjugate, or provide functionality for in vivo tracking, e.g., using fluorescence or other types of detectable agents.

[0451] In some embodiments, a conjugate of the present disclosure can comprise a linker comprising one or more of the following groups:



or a substituted analog or a stereoisomer thereof,

wherein each n_1 and n_2 is independently a value from 1 to 10. Such a group can be used as a handle to attach one or more molecules to a conjugate, e.g., to alter the pharmacokinetic (e.g., plasma half-life, retention and/or uptake in central nervous system (CNS) or elsewhere) and/or pharmacodyna via nucleophilic or electrophilic addition followed by nucleophilic substitution mic properties of the conjugate. Functionalization of such a group can occur using one or more multiple bonds (e.g., double bonds, triple bonds, etc.) of the groups. Such functionalization can comprise addition and/or substitution chemistries. For example, a functional group of a linker, such as a double bond, can be converted into a single bond (e.g., via an addition reaction such as a nucleophilic addition reaction), wherein one or both of the carbon atoms of the newly formed single bond can have a leaving group (e.g., a bromine) attached to them. Such a leaving group can then be used (e.g., via nucleophilic substitution reaction) to

attach a specific molecule (e.g., an active agent, a detectable agent, etc.) to that carbon atom(s) of the linker.

[0452] As another example, a multiple bond can be used as a functional handle in a cycloaddition reaction. Cycloaddition reactions can comprise 1,3-dipolar cycloadditions, [2+2]-cycloadditions (e.g., photocatalyzed), Diels-Alder reactions, Huisgen cycloadditions, nitron-olefin cycloadditions, etc. Such cycloaddition reactions can be used to attached various functional groups, functional moieties, active agents, detectable agents, and so forth to the conjugate. For example, a 1,3-dipoalr cycloaddition reaction can be used to attach a molecule to a conjugate, wherein the molecule comprises a 1,3-dipole that can react with, e.g., an alkyne to form a 5-membered ring, thereby attaching said molecule (e.g., active agent, detectable agent, etc.) to the conjugate. In some cases, molecules may be attached to a conjugate to e.g., modulate the half-life, increase the depot effect, or provide new functionality of a conjugate, such as fluorescence for tracking.

Pharmacokinetics of Peptides

[0453] The pharmacokinetics of any of the peptides of the present disclosure can be determined after administration of the peptide via different routes of administration. For example, the pharmacokinetic parameters of a peptide of this disclosure can be quantified after intravenous, subcutaneous, intramuscular, rectal, aerosol, parenteral, ophthalmic, pulmonary, transdermal, vaginal, optic, nasal, oral, sublingual, inhalation, dermal, intrathecal, intranasal, peritoneal, buccal, synovial, intratumoral, or topical administration. Peptides of the present disclosure can be analyzed by using tracking agents such as radiolabels or fluorophores. For example, radiolabeled peptides of this disclosure can be administered via various routes of administration. Peptide concentration or dose recovery in various biological samples such as plasma, urine, feces, any organ, skin, muscle, and other tissues can be determined using a range of methods including HPLC, fluorescence detection techniques (TECAN quantification, flow cytometry, iVIS), or liquid scintillation counting.

[0454] The methods and compositions described herein relate to pharmacokinetics of peptide administration via any route to a subject. Pharmacokinetics can be described using methods and models, for example, compartmental models or noncompartmental methods. Compartmental models include but are not limited to monocompartmental model, the two compartmental model, the multicompartmental model or the like. Models are often divided into different compartments

and can be described by the corresponding scheme. For example, one scheme is the absorption, distribution, metabolism and excretion (ADME) scheme. For another example, another scheme is the liberation, absorption, distribution, metabolism and excretion (LADME) scheme. In some aspects, metabolism and excretion can be grouped into one compartment referred to as the elimination compartment. For example, liberation includes liberation of the active portion of the composition from the delivery system, absorption includes absorption of the active portion of the composition by the subject, distribution includes distribution of the composition through the blood plasma and to different tissues, metabolism, which includes metabolism or inactivation of the composition and finally excretion, which includes excretion or elimination of the composition or the products of metabolism of the composition. Compositions administered intravenously to a subject can be subject to multiphasic pharmacokinetic profiles, which can include but are not limited to aspects of tissue distribution and metabolism/excretion. As such, the decrease in plasma or serum concentration of the composition is often biphasic, including, for example an alpha phase and a beta phase, occasionally a gamma, delta or other phase is observed.

[0455] Pharmacokinetics includes determining at least one parameter associated with administration of a peptide to a subject. In some aspects, parameters include at least the dose (D), dosing interval (τ), area under curve (AUC), maximum concentration ($C_{\max}$), minimum concentration reached before a subsequent dose is administered ($C_{\min}$), minimum time ($T_{\min}$), maximum time to reach $C_{\max}$ ($T_{\max}$), volume of distribution (V_d), steady-state volume of distribution (V_{ss}), back-extrapolated concentration at time 0 (C_0), steady state concentration (C_{ss}), elimination rate constant (k_e), infusion rate (k_{in}), clearance (CL), bioavailability (f), fluctuation (%PTF) and elimination half-life ($t_{1/2}$).

[0456] In certain embodiments, the peptides or peptide constructs of any of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 exhibit optimal pharmacokinetic parameters after oral administration. In other embodiments, the peptides or peptide constructs of any of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 exhibit optimal pharmacokinetic parameters after any route of administration, such as oral administration, inhalation, intranasal administration, topical administration, intravenous administration, subcutaneous administration, intra-articular administration, intramuscular administration, intraperitoneal administration, intra-synovial, or any combination thereof.

[0457] In some embodiments, any peptide or peptide construct of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 exhibits an average $T_{\max}$ of 0.5 – 12 hours, or 1-48 hours at which the $C_{\max}$ is reached, an average bioavailability in serum of 0.1% - 10% in the subject after administering the peptide to the subject by an oral route, an average bioavailability in serum of less than 0.1% after oral administration to a subject for delivery to the GI tract, an average bioavailability in serum of 10-100% after parenteral administration, an average $t_{1/2}$ of 0.1 hours – 168 hours, or 0.25 hours – 48 hours in a subject after administering the peptide to the subject, an average clearance (CL) of 0.5-100 L/hour or 0.5 – 50 L/hour of the peptide after administering the peptide to a subject, an average volume of distribution (V_d) of 200 – 20,000 mL in the subject after systemically administering the peptide to the subject, or optionally no systemic uptake, any combination thereof.

Peptide Stability

[0458] A peptide of the present disclosure can be stable in various biological or physiological conditions, such as the pH or reducing environments inside a cell, in the cytosol, in a cell nucleus, or endosome or a tumor. For example, any peptide or peptide construct of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361 can exhibit resistance to reducing agents, proteases, oxidative conditions, or acidic conditions.

[0459] In some cases, biologic molecules (such as peptides and proteins) can provide therapeutic functions, but such therapeutic functions are decreased or impeded by instability caused by the in vivo environment. (Moroz et al. *Adv Drug Deliv Rev* 101:108-21 (2016), Mitragotri et al. *Nat Rev Drug Discov* 13(9):655-72 (2014), Bruno et al. *Ther Deliv* (11):1443-67 (2013), Sinha et al. *Crit Rev Ther Drug Carrier Syst.* 24(1):63-92 (2007), Hamman et al. *BioDrugs* 19(3):165-77 (2005)). For instance, the GI tract can contain a region of low pH (e.g. pH ~1), a reducing environment, or a protease-rich environment that can degrade peptides and proteins. Proteolytic activity in other areas of the body, such as the mouth, eye, lung, intranasal cavity, joint, skin, vaginal tract, mucous membranes, and serum, can also be an obstacle to the delivery of functionally active peptides and polypeptides. Additionally, the half-life of peptides in serum can be very short, in part due to proteases, such that the peptide can be degraded too quickly to have a lasting therapeutic effect when administering reasonable dosing regimens. Likewise, proteolytic

activity in cellular compartments such as lysosomes and reduction activity in lysosomes and the cytosol can degrade peptides and proteins such that they may be unable to provide a therapeutic function on intracellular targets. Therefore, peptides that are resistant to reducing agents, proteases, and low pH may be able to provide enhanced therapeutic effects or enhance the therapeutic efficacy of co-formulated or conjugated, linked, or fused active agents in vivo.

[0460] Additionally, oral delivery of drugs can be desirable in order to target certain areas of the body (e.g., disease in the GI tract such as colon cancer, irritable bowel disorder, infections, metabolic disorders, and constipation) despite the obstacles to the delivery of functionally active peptides and polypeptides presented by this method of administration. For example, oral delivery of drugs can increase compliance by providing a dosage form that is more convenient for patients to take as compared to parenteral delivery. Oral delivery can be useful in treatment regimens that have a large therapeutic window. Therefore, peptides that are resistant to reducing agents, proteases, and low pH can allow for oral delivery of peptides without nullifying their therapeutic function.

[0461] Peptide Resistance to Reducing Agents. Peptides of this disclosure can contain one or more cysteines, which can participate in disulfide bridges that can be integral to preserving the folded state of the peptide. Exposure of peptides to biological environments with reducing agents can result in unfolding of the peptide and loss of functionality and bioactivity. For example, glutathione (GSH) is a reducing agent that can be present in many areas of the body and in cells, and can reduce disulfide bonds. As another example, a peptide can become reduced during trafficking of a peptide across the gastrointestinal epithelium after oral administration. A peptide can become reduced upon exposure to various parts of the GI tract. The GI tract can be a reducing environment, which can inhibit the ability of therapeutic molecules with disulfide bonds to have optimal therapeutic efficacy, due to reduction of the disulfide bonds. A peptide can also be reduced upon entry into a cell, such as after internalization by endosomes or lysosomes or into the cytosol, or other cellular compartments. Reduction of the disulfide bonds and unfolding of the peptide can lead to loss of functionality or affect key pharmacokinetic parameters such as bioavailability, peak plasma concentration, bioactivity, and half-life. Reduction of the disulfide bonds can also lead to loss of functionality due to increased susceptibility of the peptide to subsequent degradation by proteases, resulting in rapid loss of intact peptide after administration. In some embodiments, a peptide that is resistant to reduction can remain intact and can impart a

functional activity for a longer period of time in various compartments of the body and in cells, as compared to a peptide that is more readily reduced.

[0462] In certain embodiments, the peptides of this disclosure can be analyzed for the characteristic of resistance to reducing agents to identify stable peptides. In some embodiments, the peptides of this disclosure can remain intact after being exposed to different molarities of reducing agents such as 0.00001 M – 0.0001 M, 0.0001 M – 0.001 M, 0.001 M – 0.01 M, 0.01 M – 0.05 M, 0.05 M – 0.1 M, for 15 minutes or more. In some embodiments, the reducing agent used to determine peptide stability can be dithiothreitol (DTT), Tris(2-carboxyethyl)phosphine HCl (TCEP), 2-Mercaptoethanol, (reduced) glutathione (GSH), or any combination thereof. In some embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to a reducing agent. In some embodiments, peptides are completely resistant to GSH reducing conditions and are partially resistant to degradation in DTT reducing conditions. In some embodiments, peptides described herein can withstand or are resistant to degradation in physiological reducing conditions.

[0463] Peptide Resistance to Proteases. The stability of peptides of this disclosure can be determined by resistance to degradation by proteases. Proteases, also referred to as peptidases or proteinases, are enzymes that can degrade peptides and proteins by breaking bonds between adjacent amino acids. Families of proteases with specificity for targeting specific amino acids can include serine proteases, cysteine proteases, threonine proteases, aspartic proteases, glutamic proteases, and asparagine proteases. Additionally, metalloproteases, matrix metalloproteases, elastase, carboxypeptidases, Cytochrome P450 enzymes, and cathepsins can also digest peptides and proteins. Proteases can be present at high concentration in blood, in mucous membranes, lungs, skin, the GI tract, the mouth, nose, eye, and in compartments of the cell. Misregulation of proteases can also be present in various diseases such as rheumatoid arthritis and other immune disorders. Degradation by proteases can reduce bioavailability, biodistribution, half-life, and bioactivity of therapeutic molecules such that they are unable to perform their therapeutic function. In some embodiments, peptides that are resistant to proteases can better provide therapeutic activity at reasonably tolerated concentrations in vivo.

[0464] In some embodiments, peptides of this disclosure can resist degradation by any class of protease. In certain embodiments, peptides of this disclosure resist degradation by pepsin (which can be found in the stomach), trypsin (which can be found in the duodenum), serum proteases, or

any combination thereof. In some embodiments, the proteases used to determine peptide stability can be pepsin, trypsin, chymotrypsin, or any combination thereof. In certain embodiments, peptides of this disclosure can resist degradation by lung proteases (e.g., serine, cysteinyl, and aspartyl proteases, metalloproteases, neutrophil elastase, alpha-1 antitrypsin, secretory leucoprotease inhibitor, and elafin), or any combination thereof. In some embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to a protease.

[0465] Peptide Stability in Acidic Conditions. Peptides of this disclosure can be administered in biological environments that are acidic. For example, after oral administration, peptides can experience acidic environmental conditions in the gastric fluids of the stomach and gastrointestinal (GI) tract. The pH of the stomach can range from about 1-4 and the pH of the GI tract ranges from acidic to normal physiological pH descending from the upper GI tract to the colon. In addition, the vagina, late endosomes, and lysosomes can also have acidic pH values, such as less than pH 7. These acidic conditions can lead to denaturation of peptides and proteins into unfolded states. Unfolding of peptides and proteins can lead to increased susceptibility to subsequent digestion by other enzymes as well as loss of biological activity of the peptide. In certain embodiments, the peptides of this disclosure can resist denaturation and degradation in acidic conditions and in buffers, which simulate acidic conditions. In certain embodiments, peptides of this disclosure can resist denaturation or degradation in buffer with a pH less than 1, a pH less than 2, a pH less than 3, a pH less than 4, a pH less than 5, a pH less than 6, a pH less than 7, or a pH less than 8. In some embodiments, peptides of this disclosure remain intact at a pH of 1-3. In certain embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to a buffer with a pH less than 1, a pH less than 2, a pH less than 3, a pH less than 4, a pH less than 5, a pH less than 6, a pH less than 7, or a pH less than 8. In other embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to a buffer with a pH of 1-3. In other embodiments, the peptides of this disclosure can be resistant to denaturation or degradation in simulated gastric fluid (pH 1-2). In some embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at

least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to simulated gastric fluid. In some embodiments, low pH solutions such as simulated gastric fluid can be used to determine peptide stability.

[0466] In some embodiments, the peptides described herein are resistant to degradation in vivo, in the serum of a subject, or inside a cell. In some embodiments, the peptides are stable at physiological pH ranges, such as about pH 7, about pH 7.5, between about pH 5 to 7.5, between about 6.5 to 7.5, between about pH 5 to 8, or between about pH 5 to 7. In some embodiments, the peptides described herein are stable in acidic conditions, such as less than or equal to about pH 5, less than or equal to about pH 3, or within a range from about 3 to about 5. In some embodiments, the peptides are stable in conditions of an endosome or lysosome, or inside a nucleus.

[0467] Peptide Stability at High Temperatures. Peptides of this disclosure can be administered in biological environments with high temperatures. For example, after oral administration, peptides can experience high temperatures in the body. Body temperature can range from 36°C to 40°C. High temperatures can lead to denaturation of peptides and proteins into unfolded states. Unfolding of peptides and proteins can lead to increased susceptibility to subsequent digestion by other enzymes as well as loss of biological activity of the peptide. In some embodiments, a peptide of this disclosure can remain intact at temperatures from 25°C to 100°C. High temperatures can lead to faster degradation of peptides. Stability at a higher temperature can allow for storage of the peptide in tropical environments or areas where access to refrigeration is limited. In certain embodiments, 5%-100% of the peptide can remain intact after exposure to 25°C for 6 months to 5 years. 5%-100% of a peptide can remain intact after exposure to 70°C for 15 minutes to 1 hour. 5%-100% of a peptide can remain intact after exposure to 100°C for 15 minutes to 1 hour. In other embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to 25°C for at least 6 months to 5 years. In other embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to 70°C for 15 minutes to 1 hour. In other embodiments, at least 5%-10%, at least 10%-20%, at least 20%-30%, at least 30%-40%, at least 40%-50%, at least 50%-60%, at least 60%-70%, at

least 70%-80%, at least 80%-90%, or at least 90%-100% of the peptide remains intact after exposure to 100°C for 15 minutes to 1 hour.

Methods

[0468] Peptide Production and Purification. In some embodiments, amino acid sequences that are part of a peptide library are synthesized using expression vectors or solid phase or solution phase peptide synthesis methods. For example, TfR-binding peptides can be cloned into a secreted, soluble protein production/expression vector and subsequently be purified. Peptide purification methods include, but are not limited to, affinity purification columns, ion exchange (cation and/or anion column chromatography), reversed-phase chromatography, hydrophobic interaction chromatography, and size exclusion column chromatography. SDS-PAGE followed by Coomassie staining and reverse phase high-pressure liquid chromatography (HPLC) can be used to analyze a sample of the purified protein. Protein concentrations were determined by UV spectral absorption (e.g., absorbance at 280 nm) and/or amino acid analysis. In some embodiments, the peptides may be produced and purified using the methods described in Crook ZR *et al.* Methods Mol Biol. 2020; 2070:363-396.

[0469] Surface Display Vector Construction. In various embodiments, surface display vectors, SDGF and SDPR (a variant of the SDGF vector, but with all basic and aromatic residues within the stalk removed to prevent trypsin/chymotrypsin cleavage, and with a 6xHis tag (SEQ ID NO: 345) added to the C-terminus of the peptide) based on the original Daedalus vector (Bandaranayake et al., 2011), can be used to clone and express recombinant peptides as described herein. Peptides can be cloned or expressed with a modified coding sequence and without an IRES-GFP. In some cases, the coding sequence of the vector comprises: GFP; the transmembrane domain of the single pass Type II transmembrane protein FASLG (human Fas ligand); a poly-GGGS spacer (SEQ ID NO: 103); a 9-residue bovine rhodopsin antigen; a poly-GGGS spacer (SEQ ID NO: 103); a TEV-cleavage site, including a GS-linker; the displayed peptide; and for SDPR only, a 6xHis tag (SEQ ID NO: 345). In some cases, the coding sequence includes a sequence encoding siderocalin at the N-terminal of the peptide (e.g., SEQ ID NO: 352, or SEQ ID NO: 356 – SEQ ID NO: 361). As with the original Daedalus vector, the entire construct, in some embodiments, is cloned with lentiviral LTRs, and introduced to cells by transient transfection or transduction. Peptide sequences can be inserted between unique BamHI and NotI cut sites for SDGF vector, or between unique BamHI and AgeI cut sites for SDPR. In

either SDGF or SDPR construct, a number of standard cloning strategies can be used to insert a desired coding sequence, including restriction digestion (BamHI and NotI, NEB; T4 DNA Ligase, Invitrogen), In-Fusion (Clontech), or Gibson Assembly (NEB), with all transformations using Stellar chemically competent cells (Clontech), or electrocompetent cells (NEB 10-beta cells), transformed by electroporation. DNA can be PCR-amplified using exemplary primers such as: SDGF Forward primer: TGTACTTCCAGGGAGGATCC (SEQ ID NO: 127); SDGF Reverse primer: AATGGTGATGAGCGGCC (SEQ ID NO: 128).

[0470] The use of both magnetic and flow sorting may allow for about 10^9 cells to be screened. One-eighth of the number of cells to be screened, or $>1.2 \times 10^8$ cells, may be transduced to allow for the 3 days for optimal construct expression and an approximately 24-hour doubling time. To test a limited number of peptide scaffolds, library diversity can be achieved through saturation mutagenesis (e.g., NNK). However, if a more diverse scaffold library is preferred, the library may be reduced to about 10,000 members, ordered as an oligonucleotide pool, and further diversified it by error-prone PCR mutagenesis. For libraries diversified by error-prone PCR mutagenesis, the total library size (the number of library variants) may be heavily dependent on the degree of mutagenesis imposed. Since stop codons and alteration of the cysteine topology can render a peptide nonfunctional. A lower mutagenesis rates (1–1.5 per peptide) may reduce complications from cysteine topology disruption in libraries comprising CDPs, but the mutagenesis rate may be adjusted for each library.

[0471] An exemplary method for library design for optimized cloning into SDGF may comprise the steps of: 1) Determining the desired peptide library. The peptide library may be identified from cysteine-rich sequences in existing databases (e.g., Uniprot) or may be derived from *in silico* docking simulations using protein design software (e.g., Rosetta); 2) Converting the peptide sequences to cloning-ready oligonucleotides. Converting the peptide sequences to cloning-ready oligonucleotides may comprise the steps of: (a) Reverse-translating the peptide sequences, using human codon-optimization; (b) Incorporating uniform flanking sequences to facilitate PCR amplification and cloning; (c) If NGS is to be employed for library characterization, verifying that each library member contains a unique sequence within the entirety of the NGS read. Silent mutations may be introduced to ensure unique mapping is possible; and (d) Ordering the finalized library.

[0472] Cloning Pooled Oligonucleotide Libraries into SDGF. An exemplary method for cloning pooled oligonucleotide libraries into SDGF may comprise the steps of: 1) resuspending

the pooled oligonucleotide library in 50 μ L molecular biology-grade water or Tris–EDTA buffer; 2) preparing four reactions to PCR amplify the library, running the four reactions at different cycle numbers (e.g., 14, 16, 18, and 20); 3) running all PCR products on an agarose gel and extracting the expected band from a lane lacking the inappropriate, high molecular weight amplicon smear above the desired product, which may be a result of PCR product cross-priming when primers are depleted, and purifying the bands using a gel extraction kit; 4) optionally, if further diversification is desired, performing mutagenesis by error-prone PCR using a Gene-Morph II Random Mutagenesis Kit and the SDGF Gibson primers using manufacturer-recommended reaction conditions, with the PCR product from the previous step as the template, and running on a gel and purify the desired band from a lane without a substantial cross-priming smear, as in step 3; 5) preparing 2 μ g per library or 100 ng per singleton to be cloned of BamHI/NotI-digested SDGF backbone; 6) cloning the amplicon into SDGF, for example by incubate a Gibson reaction for 2 h at 50°C instead of the recommended 1 hour; 7) de-salting the reaction using a PCR purification kit, eluting in a reduced volume of 20 μ L; 8) electroporating the library into competent *E. coli* and incubating at 37°C overnight, for example, using specialized electrocompetent cells, plating electroporated cells onto a large plate (25 x 25 cm) of LB agar + carbenicillin, taking care to spread completely and allowing to dry completely before inversion and overnight 37 °C incubation with control petri dishes containing serial dilutions of the library to establish library colony forming units (CFUs); 9) counting the control plates to establish CFUs and, if CFUs are sufficient, scraping the plate(s) by adding 25–50 mL LB media plus carbenicillin (100 μ g/mL) and scraping with a cell scraper, lifter, or spreader, repeating with more media and scraping until transfer is complete; 10) shaking bacteria in a final volume of 200 mL for 1 hour at 37°C, and then pelleting 50 mL aliquots (4000 x g for 20 min); and 11) freezing all but one pellet at -20°C, maxipreping the remaining pellet, and preferably sterile filtering the final DNA product prior to lentiviral production.

[0473] Mammalian Surface Display. Validation of peptides or single candidates for binding to TfR can be validated using a suspension of HEK-293 cells. In some embodiments, cells are transfected with 2.5 μ g SDGF vector comprising a putative or candidate peptide and 3.5 μ g polyethyleneimine to 2E6 cells in 1 mL media, in a 24-well suspension tissue culture dish. After incubation of cells, all pooled screening can be performed in 293T cells transduced with lentivirus (VSV-G coated, produced by standard methods in 293T cells), delivering the SDGF construct, at an MOI of approximately 1, or up to 5. Transduced or transfected suspension HEK-

293 cells can be pelleted (500xg, 5 min) and resuspended at up to 8E6 cells/mL in Flow Buffer (PBS with 0.5% BSA and 2 mM EDTA) containing DAPI and target protein, with or without Alexa Fluor 647-conjugated streptavidin (ThermoFisher) in equimolar quantity to the target protein. All staining incubations took place at 4 C for 30 mins with mild agitation. For the low stringency protocol (initial assay validation and pooled screening), 200 nM target protein can be used, and streptavidin can be pre-mixed with target protein. Cells can be incubated, diluted 10-fold with Flow Buffer, pelleted (500xg, 5 min), and resuspended in fresh Flow Buffer (up to 6.5E6 cells/mL) before flow cytometry. For the medium stringency protocol (SSM maturation), only 20 nM target protein can be used, and cells can be incubated with target protein alone, diluted 10-fold with Flow Buffer, pelleted, and then resuspended with 20 nM streptavidin. Cells were incubated again, diluted 10-fold with Flow Buffer, pelleted (500xg, 5 min), and resuspended in fresh Flow Buffer (up to 6.5E6 cells/mL) before flow cytometry. The high stringency protocol is identical to the medium stringency protocol, except that an additional Flow Buffer-only wash step is included immediately after target protein staining. Initial screen was conducted with 200 nM TfR concentration, using a one-step staining protocol for tetravalent target avidity, as opposed to a two-step staining protocol for weaker, monovalent avidity. Subsequent affinity maturation used 20 nM one-step staining (first affinity maturation screen) and 8 nM two-step staining (second affinity maturation screen). Singleton variant validation and permutation test staining took place under the same conditions as the screen from which the variants were enriched and identified. For the transferrin competition staining assay, TfR levels were 8 nM with two-step staining, and transferrin levels were 10 μ M. Transfected cells were incubated first with transferrin (10 μ M), then TfR solution was added for the first staining step. Cells were pelleted and then resuspended in streptavidin solution for the second staining step. After this second incubation, cells were pelleted and analyzed by flow cytometry for streptavidin staining (representing TfR-binding). For the murine transferrin vs. human transferrin cross-reactivity testing, soluble peptides were labeled with AlexaFluor 647 dye via NHS-ester conjugation, and then stained at 50 nM on cells expressing either human or mouse TfR on their cell surface. No co-stain was necessary, but staining protocol is otherwise identical. In some cases, flow sorting is performed on a Beckton-Dickinson Aria II, while analysis is performed on a combination of Beckton-Dickson LSR II and Acea NovoCyte flow cytometers. Data analysis can be performed on FlowJo (FlowJo, LLC) and Excel (Microsoft).

[0474] Site Saturation Mutagenesis (SSM) of Peptides. SSM can be employed for affinity maturation of peptides of the present disclosure, for example, for those peptides that are identified in a the first mammalian surface display experiment. During each round of maturation, a library of every possible non-cysteine point variant can be constructed, and screened against TfR at higher stringency than first screen. Variants with improved binding can be enriched and identified by Sanger sequencing. Such enriched variant mutations can be combined with one another in various permutations (shown in the data) to identify a composite, improved binder.

[0475] Next Generation Sequencing. In some embodiments, screens for enrichment or depletion of variants affecting TfR-binding can be assessed by Illumina sequencing. Such sequencing method involves collecting cell pellets (1.5E6, 3 technical replicates) resuspended in 50 µL Terra Direct PCR Mix (from Clontech) and amplified for 16 cycles using the original cloning primers. Up to four aliquots can be diluted 16-fold into 60 µL Phusion DNA Polymerase reactions and amplified using distinct Illumina primers, containing adaptor sequences for flow cell adherence. Forward primers can include a 6 bp barcode for multiplexing. Illumina HiSeq 2500 in rapid mode can be used to run samples, which Bowtie2 software can be used for mapping, and Excel (Microsoft) and MATLAB (MathWorks) used for data analysis.

[0476] His-Avi-TfR Production, Purification, and Biotinylation. In some embodiments, in vitro assays are used to analyze candidate peptides for ability to bind TfR. In some cases, biotinylated, and His-Avi-TfR can be expressed in mammalian cells. Optionally, proteins can be biotinylated following purification (e.g., using BirA to biotinylate an encoded AviTag). In some embodiments, the expressed peptide is fused to siderocalin (SEQ ID NO: 351) at the N-terminus for increased stability during translation or secretion (e.g., SEQ ID NO: 352, or SEQ ID NO: 356 – SEQ ID NO: 361). Peptides as disclosed herein can be generated in mammalian cell culture using a published methodology. (A.D. Bandaranayake, C. Correnti, B.Y. Ryu, M. Brault, R.K. Strong, D. Rawlings. 2011. Daedalus: a robust, turnkey platform for rapid production of decigram quantities of active recombinant proteins in human cell lines using novel lentiviral vectors. *Nucleic Acids Research*. (39)21, e143). The peptide sequence was reverse-translated into DNA, synthesized, and cloned in-frame with siderocalin using standard molecular biology techniques. (M.R. Green, Joseph Sambrook. *Molecular Cloning*. 2012 Cold Spring Harbor Press.). The resulting construct was packaged into a lentivirus, transduced into HEK-293 cells, expanded, isolated by immobilized metal affinity chromatography (IMAC), cleaved with tobacco etch virus protease, and purified (e.g., separation of peptide and siderocalin carrier (SEQ ID NO:

351)) to homogeneity by size-exclusion chromatography using a gradient of acetonitrile and 0.1% TFA. Peptides expressed as siderocalin fusions can be treated with tobacco etch virus (TEV) protease to cleave siderocalin and leave an N-terminal GS. Thus, the sequences of this disclosure can optionally include an N-terminal GS. Peptides can be separated from the siderocalin carrier using chromatography, e.g., size exclusion chromatography (SEC) or reverse phase high performance liquid chromatography (RP-HPLC). Following purification, each peptide was lyophilized and stored frozen. Alternatively, purified proteins can be flash frozen in aqueous solution (e.g., PBS) with a cryoprotectant (e.g., 5% glycerol) and stored frozen.

[0477] Fluorophore Conjugation. In some embodiments, screening peptides or a library of peptides involves labeling of a protein of interest or a protein partner with a fluorophore or any other detectable moiety, such that detection of a signal from the fluorophore or detectable moiety is indicative of binding to the peptide. An example of a fluorophore that can be used is Alexa Fluor 647 NHS Ester (Life Technologies), which can be used to label a protein of interest as per manufacturer's protocol. Saturation is approximately 1 fluorophore per molecule, as determined by mass spectrometry. Free dye can be removed following labeling (e.g., by spin column dialysis).

[0478] Surface Plasmon Resonance (SPR) Interaction Analyses. In some embodiments, TfR-binding can be assessed using surface plasmon resonance (SPR). The peptides of the present disclosure can be tobacco etch virus (TEV)-cleavable and thus can be analyzed as independent, soluble proteins.

[0479] Various peptides of the present disclosure can be analyzed for binding affinity to TfR. Binding affinity can be analyzed by SPR experiments using captured biotinylated TfR, which can be performed at 25 °C on a Biacore T100 instrument (GE Healthcare) with Series S SA chips. HBS-EP+ (10 mM HEPES, pH 7.4, 150 mM NaCl, 3 mM EDTA, 0.05% surfactant P20) can be used as a running buffer in the experiments with 0.1 mg/mL bovine serum albumin (BSA). Soluble TfR-binding peptides can be evaluated for binding by incubation of a dilution series, in which the concentration range can be varied depending on the TfR-binding peptide being tested with 2 ug/ml TfR, capturing ~300 resonance units (RUs) of protein for SPR experiments.

[0480] In some cases, SPR is performed by injecting the peptides over two densities of captured and biotinylated hTfR and the resulting data can then be analyzed globally. Binding constants (e.g., K_D values) and on- and off-rates can be calculated for the interactions between TfR and the TfR-binding peptide.

[0481] In some embodiments, SPR is performed with purified apo-transferrin and holo-transferrin. Holo-transferrin can be produced from apo-transferrin by loading apo-transferrin with an iron solution (e.g., 0.4 equivalents of $\text{Fe}(\text{NTA})_2$ added every 5 minutes until 3.2 equivalents have been added). Excess iron can be subsequently removed through buffer exchange (e.g., spin column dialysis into PBS).

[0482] **Circular Dichroism.** Protein secondary structures can be assessed using Circular Dichroism (CD). CD spectra can be measured with a Jasco J-720W spectropolarimeter using a 1.0mm path length cell. Protein samples in 10mM phosphate buffer (pH = 7.4) can be at 25-30 μM protein concentration. Samples can be analyzed at wavelength ranges of 260-190 nm. Data can be expressed in terms of relative ellipticity $[\theta]$; (mdeg). To determine thermal stability of proteins, samples can be subjected to incremental increase in temperature at a ramp of $2^\circ\text{C}/\text{min}$ from 20°C to 95°C . Stability and protein unfolding were monitored at 210 nm for α -helix secondary structure. Protein unfolding can also be monitored at 215 nm or 220 nm. Data can be expressed in terms of relative ellipticity $[\theta]$, reported in mdeg.

[0483] **Protease Resistance Testing.** Tumor environments, for instance, contain high concentrations of proteases. Furthermore, resistance to proteolysis (degradation or cleaving by proteases) reduces the likelihood of peptide degradation and subsequent immunogenicity, for example, in circulation. Resistance to proteolysis can further increase a peptide's serum half-life and the overall bioavailability of the peptide. Thus, it is advantageous for peptides of the present disclosure to be resistant to degradation by proteases.

[0484] To determine resistance to proteolytic digestion, various peptides of the present disclosure can be mixed with 50 U of porcine pepsin (Sigma-Aldrich P7012) in Simulated Gastric Fluid at pH 1.05, or 50 U of porcine trypsin (Sigma-Aldrich 6567) in PBS, incubated for 30 minutes at 37°C , and analyzed by reverse phase HPLC (RP-HPLC). Separately, the peptides can be mixed with 50 U of porcine pepsin (Sigma-Aldrich P7012) in Simulated Gastric Fluid⁴³ at pH 1.05, or 50 U of porcine trypsin (Sigma-Aldrich 6567) in PBS, incubated for 30 m at 37°C , and analyzed by RP-HPLC. Control peptides can be incubated in PBS. All peptides can be incubated in non-reducing conditions.

[0485] In some embodiments, SDPR protease resistance testing can be performed having a sequence of SEQuencing-grade enzymes, including Trypsin and Chymotrypsin. Trypsin and trypsin inhibitor can be used for HPLC analysis.

[0486] Peptide Stability testing under Reducing Conditions. Peptides that can be designed or engineered to interact with one or more target proteins in the cell, such as proteins in the nucleus, can be exposed to reducing conditions in the cytosolic compartment of cells. Thus, it can be advantageous for peptides of this disclosure to display stability in reducing conditions. Stability of peptides of this disclosure can be tested in 10 mM dithiothreitol (DTT) and/or 10 mM glutathione (GSH). GSH can be a more physiologically relevant reducing agent for testing peptide stability in intracellular conditions. An example of a peptide-target protein interaction is a peptide binding to a TfR protein. Stability to the tested peptides can be evaluated using HPLC (or radio-HPLC when radiolabeled peptides are used), for example.

[0487] Thermal Shift Assay. Protein melting temperature (T_m) determination can be performed by monitoring protein unfolding using SYPRO Orange dye (Molecular Probes). In brief, 0.1 mg/mL protein sample in 20 μ L total volume PBS buffer can be mixed with 2 μ L of 10X SYPRO Orange dye. Dye intercalation into the hydrophobic protein core following protein unfolding was assayed using the C1000 Touch Thermal Cycler with CFX96 Deep Well Real-Time System (BioRad). Samples were heated from 20 $^{\circ}$ C to 95 $^{\circ}$ C with stepwise increments of 0.5 $^{\circ}$ C per minute and a 5 sec hold step for every point, followed by fluorescence reading. T_m were calculated by analyzing the derivatives of Relative fluorescence Units (RFU).

[0488] Cross Reactivity of TfR-binding Peptides to Murine TfR. Cross reactivity to human and murine TfR of TfR-binding peptides as disclosed herein can be performed using cell surface binding assays. A reversal of the surface display paradigm is further described in **EXAMPLE 3**. 293F cells expressing either human or mouse TfR from their surface were stained with soluble TfR-binding peptides that were directly labeled with AlexaFluor 647 dye. Flow cytometry can subsequently be used for data analysis to assess human versus mouse reactive if TfR-binding peptides.

[0489] Co-crystalizing the TfR-binding peptides with TfR and solving the crystal structure. Co-crystalizing the TfR-binding peptides with TfR can be performed as follows. The peptide can be resuspended at a target concentration of approximately 5-15 mg/mL, in some cases 10 mg/mL. Higher concentrations of up to 80 mg/mL can be used for crystallization when, for example, a single peptide is present in the solution without any other components in it. Crystallization screening can be performed at room temperature by vapor diffusion, with 1:1 protein solution:reservoir solution sitting drops, set up using the Nextal JCSG+, PEGs, and $(\text{NH}_4)_2\text{SO}_4$ factorial suites (Qiagen) and sub-microliter robotics (TTP Labtech mosquito). Diffraction data

can be collected from single crystals using a Rigaku MicroMax-007 HF home source or beamline 5.0.1 at the Advanced Light Source (Lawrence Berkley National Laboratory, Berkeley, CA). Initial phases can be determined either by molecular replacement (MR), using PHASER (McCoy, A.J., *J Appl Crystallogr.*, 40(Pt 4): 658-674 (2007)) in the CCP4 program suite (Winn, M.D., *Acta Crystallogr D Biol Crystallogr.*, 67(Pt 4): 235-42 (2011)) using homologous structures from the RCSB PDB (Berman, H.M., *Nucleic Acids Res.*, 28(1): 235-42 (2000)) as search models, or sulfur single-wavelength anomalous diffraction (sSAD) (Liu, Q., *Acta Crystallogr D Biol Crystallogr.*, 69(Pt 7): 1314-32 (2013)), using CuKalpha radiation to maximize the anomalous signal, and determining sulfur substructures with SHELX (Sheldrick, G.M., *Acta Crystallogr D Biol Crystallogr.*, 66 (Pt 4): 479-85 (2010)). For sSAD phasing, Bijvoet pair measurement can be optimized by collecting data through 5° wedges with alternating phi rotations of 180°, in 1° oscillations. Data can be reduced and scaled with HKL2000 (Otwinowski, Z., *Methods Enzymol.*, 276: 307-326 (1997)). Iterative cycles of model building and refinement can be performed with COOT (Emsely, P., *Acta Crystallogr D Biol Crystallogr.*, 60(Pt 12 Pt 1): 2126-32 (2004)) and REFMAC (Murshudov, G.N., *Acta Crystallogr D Biol Crystallogr.*, 53(Pt 3): 240-55 (1997)). Structure validation can be performed with MolProbity (Davis, I.W., *Nucleic Acids Res.*, 35(Web Server issue): W375-83 (2007)).

[0490] Whole-body autoradiography to elucidate biodistribution of the TfR-binding

peptides in mice. Whole body autoradiography (WBA) sagittal sectioning can be performed using the following steps. Each peptide can be radiolabeled by methylating lysines at the N-terminus as described further in **EXAMPLE 14**. As such, the peptide can contain methyl or dimethyl lysines and a methylated or dimethylated amino terminus. A dose of 100 nmol radiolabeled peptide can be administered via tail vein injection in Female Harlan athymic nude mice with intact kidneys, weighing 20-25 g. Each radiolabeled peptide can be allowed to freely circulate within the animal for the described time period before the animals are euthanized and sectioned.

[0491] All mice of treatment cohorts can be dosed with 100 nmol radiolabeled peptide or peptide construct (~20 mg/kg). Control mice can be injected with saline or PBS. At the end of the dosing period, mice are frozen in a hexane/dry ice bath and then embedded in a frozen block of carboxymethylcellulose. Whole animal sagittal slices can be prepared that resulted in thin frozen sections for imaging. Thin frozen sections can be obtained using a microtome and allowed visualization of tissues such as brain, tumor, liver, kidney, lung, heart, spleen, pancreas, muscle,

adipose, gall bladder, upper gastrointestinal tract, lower gastrointestinal tract, bone, bone marrow, reproductive tract, eye, cartilage, stomach, skin, spinal cord, bladder, salivary gland, and more. Sections are allowed to dessicate in a freezer prior to imaging.

[0492] For the autoradiography imaging, tape mounted thin sections were freeze dried and radioactive samples can be exposed to phosphorimager plates for about 7 days. These plates are developed and the signal (densitometry) from each organ was normalized to the signal found in the cardiac blood of each animal. A signal in tissue darker than the signal expected from blood in that tissue indicates accumulation in a region, tissue, structure, or cell.

[0493] Reported quantitation of peptide in tissue (nmol/g) was accomplished by referencing an in-block standard curve and the peptide's specific ^{14}C activity.

[0494] Serum and tissue quantitation of peptide levels over a time course. Depending on the utilized radionuclide either ex vivo liquid scintillation counting (e.g., ^{14}C) or gamma counting (e.g., all positron emitting nuclides) can be used to determine peptide levels of various organs at specific time points as well as peptide behavior over time such as for blood half-life and organ uptake and clearance. All mice of treatment cohorts can be dosed with 100 nmol radiolabeled peptide or peptide construct (~20 mg/kg). Control mice can be injected with saline or PBS. Each radiolabeled peptide can be allowed to freely circulate within the animal for the described time period before the animals were euthanized and sectioned. Collected organs including kidney, liver, brain, spleen, muscle, and skin can be homogenized or weighed before or after counting. Internal reference standards and control samples can also be counted. The peptide uptake data can be visualized as amount of peptide per gram tissue, or as percent injected dose per gram (or per mole or per volume).

[0495] General determination of peptide uptake in cells or tissues. The uptake of a peptide or peptide construct as disclosed herein can be determined in a specific cell, cell population, tissue, or organ either *ex vivo* or *in vivo*. In the same way, the efficacy of cargo or payload delivery can be determined *ex vivo* or *in vivo*, for example, in cases where the cargo molecule comprises a detectable agent. *Ex vivo* analyses include organ harvest and fixation (e.g., using 4% formaldehyde) of harvested tissue prior to analyses. Tissue samples can be analyzed using a variety of analytical methods including microscopy, spectroscopy, flow cytometry, polymerase chain reaction (PCR), and via measurements of ultrasound, electromagnetic radiation (e.g., UV/VIS, X-ray) or radioactivity. For example, tissue uptake can be determined by measuring luminescence or bioluminescence of a cell, cell population, tissue, or organ sample, or by

measuring radioactivity of a cell, cell population, tissue, or organ sample and by calculating uptake values such as percent injected dose per gram (or per mole or per volume). In addition, the acquisition of nuclear images visualizing the biodistribution at a specific point in time (e.g., static imaging) or over a time course (e.g. dynamic imaging) can be performed using various techniques (e.g., PET or SPECT) and appropriately radiolabeled peptides.

[0496] MTR BBB transcytosis model, and capillary depletion quantitation of parenchyma/capillary distribution. Mice (strain CD-1, N=1 per time point) can be anesthetized, and the left jugular vein and right carotid arteries can be surgically exposed. Labeled peptides (e.g., 0.25-1 μ Ci in about 200 μ L volume; specific activities were 22-40 Ci/mol for SEQ ID NO: 1 and 127-150 Ci/mol for SEQ ID NO: 32) can be injected into the jugular vein. After a given period time within a time course (0-30 minutes), arterial blood can be collected, followed by decapitation and brain collection. Brains and serum can be tested via liquid scintillation counting for ^{14} C levels, and the data can be incorporated into a multiple time regression mathematic model to quantify CNS accumulation rate. To determine whether this CNS accumulation is limited to capillaries or actually represents vascular transcytosis, a separate set of animals (N=3/peptide) can be euthanized 5 minutes after dosage, and brain homogenates are subjected to centrifugation through a dextran density gradient. This can allow separation of capillaries from parenchyma, and the two tissues can be subjected separately to scintillation counting to quantitate ^{14} C signal. Normalized signal ratios can be an indicator of BBB penetration; for example, SEQ ID NO: 1 can have a substantially higher signal in the parenchyma than the capillaries, interpreted as evidence of BBB transcytosis.

[0497] Methods for NT fusions and activation of the neuronal CRE reporter mouse.

Transgenic mice expressing firefly luciferase under the control of a cyclic AMP response element (CRE) can be used. Under conditions of elevated cyclic AMP (cAMP) or other mechanisms that activate the transcription factor CRE Binding Protein 1 (CREB), cells in these mice can express luciferase, which can be measured in whole animal luminescence instrumentation (IVIS imager) after IV dosage with a luciferin formulated for animal use. Mice can be validated to express the transgene under proper response element control by dosing these mice with forskolin and rolipram, two drugs that stimulate neuronal cAMP production and inhibit its degradation (respectively). Animals dosed with forskolin and rolipram can then be tested for luciferase expression 4 hours later as described herein. High levels of luminescence from the animal's brain can be typical in these control experiments. Animals that validate in this assay can

be given about 1 week to return to steady state (low) luciferase expression, and can then be tested with neurotensin or TfR binding proteins fused with neurotensin. Neurotensin receptor expression can be limited to subsets of cells in the CNS, including the frontal cortex. Upon activation, a signal transduction pathway can be activated that culminates in CREB phosphorylation and CRE-mediated transcription. Its ligand, neurotensin, is a neuropeptide that cannot cross the BBB. Therefore, IV dosage of a fusion protein comprising neurotensin and a TfR binding peptide can demonstrate BBB penetration if CRE-driven luciferase is induced in the brains of these mice.

[0498] Peptide constructs, which are NT fusions include any sequence having at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO:70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, or SEQ ID NO: 359 – SEQ ID NO: 361.

Methods of Manufacture

[0499] Various expression vector/host systems can be utilized for the recombinant expression of peptides described herein. Non-limiting examples of such systems include microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing a nucleic acid sequence encoding peptides, peptide constructs, or peptide fusion proteins/chimeric proteins described herein, yeast transformed with recombinant yeast expression vectors containing the aforementioned nucleic acid sequence, insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the aforementioned nucleic acid sequence, plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus (CaMV), tobacco mosaic virus (TMV)), or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing the aforementioned nucleic acid sequence, or animal cell systems infected with recombinant virus expression vectors (e.g., adenovirus, vaccinia virus, lentivirus) including cell lines engineered to contain multiple copies of the aforementioned nucleic acid sequence, either stably amplified (e.g., CHO/dhfr, CHO/glutamine synthetase) or unstably amplified in double-minute chromosomes (e.g., murine cell lines). Disulfide bond formation and folding of the peptide could occur during expression or after expression or both.

[0500] A host cell can be adapted to express one or more peptides described herein. The host cells can be prokaryotic, eukaryotic, or insect cells. In some cases, host cells are capable of modulating the expression of the inserted sequences, or modifying and processing the gene or protein product in the specific fashion desired. For example, expression from certain promoters can be elevated in the presence of certain inducers (e.g., zinc and cadmium ions for metallothionine promoters). In some cases, modifications (e.g., phosphorylation) and processing (e.g., cleavage) of peptide products can be important for the function of the peptide. Host cells can have characteristic and specific mechanisms for the post-translational processing and modification of a peptide. In some cases, the host cells used to express the peptides secrete minimal amounts of proteolytic enzymes.

[0501] In the case of cell- or viral-based samples, organisms can be treated prior to purification to preserve and/or release a target polypeptide. In some embodiments, the cells are fixed using a fixing agent. In some embodiments, the cells are lysed. The cellular material can be treated in a manner that does not disrupt a significant proportion of cells, but which removes proteins from the surface of the cellular material, and/or from the interstices between cells. For example, cellular material can be soaked in a liquid buffer, or, in the case of plant material, can be subjected to a vacuum, in order to remove proteins located in the intercellular spaces and/or in the plant cell wall. If the cellular material is a microorganism, proteins can be extracted from the microorganism culture medium. Alternatively, the peptides can be packed in inclusion bodies. The inclusion bodies can further be separated from the cellular components in the medium. In some embodiments, the cells are not disrupted. A cellular or viral peptide that is presented by a cell or virus can be used for the attachment and/or purification of intact cells or viral particles. In addition to recombinant systems, peptides can also be synthesized in a cell-free system prior to extraction using a variety of known techniques employed in protein and peptide synthesis.

[0502] In some cases, a host cell produces a peptide that has an attachment point for a cargo molecule (e.g., a therapeutic agent). An attachment point could comprise a lysine residue, an N-terminus, a cysteine residue, a cysteine disulfide bond, a glutamic acid or aspartic acid residue, a C-terminus, or a non-natural amino acid. The peptide could also be produced synthetically, such as by solid-phase peptide synthesis, or solution-phase peptide synthesis. Peptide synthesis can be performed by fluorenylmethyloxycarbonyl (Fmoc) chemistry or by butyloxycarbonyl (Boc) chemistry. The peptide could be folded (formation of disulfide bonds) during synthesis or after

synthesis or both. Peptide fragments could be produced synthetically or recombinantly. Peptide fragments can be then be joined together enzymatically or synthetically.

[0503] In other aspects, the peptides of the present disclosure can be prepared by conventional solid phase chemical synthesis techniques, for example according to the Fmoc solid phase peptide synthesis method (“Fmoc solid phase peptide synthesis, a practical approach,” edited by W. C. Chan and P. D. White, Oxford University Press, 2000).

[0504] In some embodiments, the peptides of this disclosure can be more stable during manufacturing. For example, peptides of this disclosure can be more stable during recombinant expression and purification, resulting in lower rates of degradation by proteases that are present in the manufacturing process, a higher purity of peptide, a higher yield of peptide, or any combination thereof. In some embodiments, the peptides can also be more stable to degradation at high temperatures and low temperatures during manufacturing, storage, and distribution. For example, in some embodiments peptides of this disclosure can be stable at 25 °C. In other embodiments, peptides of this disclosure can be stable at 70 °C or higher than 70 °C. In some embodiments, peptides of this disclosure can be stable at 100 °C or higher than 100 °C.

Pharmaceutical Compositions

[0505] A pharmaceutical composition of the disclosure can be a combination of any peptide as described herein with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, antioxidants, solubilizers, buffers, osmolytes, salts, surfactants, amino acids, encapsulating agents, bulking agents, cryoprotectants, and/or excipients. The pharmaceutical composition facilitates administration of a peptide described herein to an organism. In some cases, the pharmaceutical composition comprises factors that extend half-life of the peptide and/or help the peptide to penetrate the target cells.

[0506] Pharmaceutical compositions can be administered in therapeutically-effective amounts as pharmaceutical compositions by various forms and routes including, for example, intravenous, subcutaneous, intramuscular, rectal, aerosol, parenteral, ophthalmic, pulmonary, transdermal, vaginal, optic, nasal, oral, sublingual, inhalation, dermal, intrathecal, intratumoral, intranasal, and topical administration. A pharmaceutical composition can be administered in a local or systemic manner, for example, via injection of the peptide described herein directly into an organ, optionally in a depot.

[0507] Parenteral injections can be formulated for bolus injection or continuous infusion. The pharmaceutical compositions can be in a form suitable for parenteral injection as a sterile suspension, solution or emulsion in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Pharmaceutical formulations for parenteral administration include aqueous solutions of a peptide described herein in water-soluble form. Suspensions of peptide-antibody complexes described herein can be prepared as oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. The suspension can also contain suitable stabilizers or agents which increase the solubility and/or reduce the aggregation of such peptide-antibody complexes described herein to allow for the preparation of highly concentrated solutions.

[0508] Formulations or approaches to increase cell penetration of a peptide as described herein can include, but not limited to, using high dosage of a peptide as described herein, such as up to 10 μM ; conjugating or fusing a Tat peptide or a cell-penetrating peptide, or a variant or derivative thereof to a peptide; co-delivering a Tat peptide or a cell-penetrating peptide, or a variant or derivative thereof with a peptide; conjugating or fusing an Arg patch to a peptide; or co-delivering a peptide with an Arg patch peptide, such as up to 10 μM . In some embodiments, protein transfection agents, direct cytosolic expression of the peptide, or electroporation of the peptide can be used to increase cell penetration. In some embodiments, other excipients can be formulated with a peptide in order to increase the cell penetration of the peptide, such as those approaches described in "Protein and Peptide Drug Delivery: Oral Approaches" Indian J. Pharm. Sci., Shaji and Patole, v70(3) 269-277, 2008. Any combination of these formulations or approaches can be used to increase cell penetration of a peptide as described herein.

[0509] Alternatively, the peptide described herein can be lyophilized or in powder form for reconstitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use. In some embodiments, a purified peptide is administered intravenously. A peptide described herein can be administered to a subject in order to home, target, migrate to, or be directed to a CNS cell, a brain cell, a cancerous cell, or a tumor. In some embodiments, a peptide can be conjugated to, linked to, or fused to another peptide that provides a targeting function to a specific target cell type in the central nervous system or across the blood brain barrier. Exemplary target cells include a

CNS cell, erythrocyte, an erythrocyte precursor cell, an immune cell, a stem cell, a muscle cell, a brain cell, a thyroid cell, a parathyroid cell, an adrenal gland cell, a bone marrow cell, an appendix cell, a lymph node cell, a tonsil cell, a spleen cell, a muscle cell, a liver cell, a gallbladder cell, a pancreas cell, a cell of the gastrointestinal tract, a glandular cell, a kidney cell, a urinary bladder cell, an endothelial cell, an epithelial cell, a choroid plexus epithelial cell, a neuron, a glial cell, an astrocyte, or a cell associated with a nervous system.

[0510] A peptide of the disclosure can be applied directly to an organ, or an organ tissue or cells, such as brain or brain tissue or cells, during a surgical procedure. The recombinant peptide described herein can be administered topically and can be formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, medicated sticks, balms, creams, and ointments. Such pharmaceutical compositions can contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

[0511] In practicing the methods of treatment or use provided herein, therapeutically-effective amounts of a peptide described herein can be administered in pharmaceutical compositions to a subject suffering from a condition that affects the immune system. In some embodiments, the subject is a mammal such as a human or a primate. A therapeutically-effective amount can vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compounds used, and other factors.

[0512] In some embodiments, a peptide is cloned into a viral or non-viral expression vector. Such expression vector can be packaged in a viral particle, a virion, or a non-viral carrier or delivery mechanism, which is administered to patients in the form of gene therapy. In other embodiments, patient cells are extracted and modified to express a peptide capable of binding TfR *ex vivo* before the modified cells are returned back to the patient in the form of a cell-based therapy, such that the modified cells will express the peptide once transplanted back in the patient.

[0513] Pharmaceutical compositions can be formulated using one or more physiologically-acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active compounds into preparations that can be used pharmaceutically. Formulation can be modified depending upon the route of administration chosen. Pharmaceutical compositions comprising a peptide described herein can be manufactured, for example, by expressing the peptide in a recombinant system, purifying the peptide, lyophilizing the peptide, mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or compression

processes. The pharmaceutical compositions can include at least one pharmaceutically acceptable carrier, diluent, or excipient and compounds described herein as free-base or pharmaceutically-acceptable salt form.

[0514] Methods for the preparation of peptide described herein comprising the compounds described herein include formulating peptide described herein with one or more inert, pharmaceutically-acceptable excipients or carriers to form a solid, semi-solid, or liquid composition. Solid compositions include, for example, powders, tablets, dispersible granules, capsules, cachets, and suppositories. These compositions can also contain minor amounts of nontoxic, auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, and other pharmaceutically-acceptable additives.

[0515] Non-limiting examples of pharmaceutically-acceptable excipients can be found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), each of which is incorporated by reference in its entirety.

[0516] Pharmaceutical compositions can also include permeation or absorption enhancers (Aungst et al. AAPS J. 14(1):10-8. (2012) and Moroz et al. Adv Drug Deliv Rev 101:108-21. (2016)). Permeation enhancers can facilitate uptake of molecules from the GI tract into systemic circulation. Permeation enhancers can include salts of medium chain fatty acids, sodium caprate, sodium caprylate, N-(8-[2-hydroxybenzoyl]amino)caprylic acid (SNAC), N-(5-chlorosalicyloyl)-8-aminocaprylic acid (5-CNAC), hydrophilic aromatic alcohols such as phenoxyethanol, benzyl alcohol, and phenyl alcohol, chitosan, alkyl glycosides, dodecyl-2-N,N-dimethylamino propionate (DDAIPP), chelators of divalent cations including EDTA, EGTA, and citric acid, sodium alkyl sulfate, sodium salicylate, lecithin-based, or bile salt-derived agents such as deoxycholates.

[0517] Compositions can also include protease inhibitors including soy bean trypsin inhibitor, aprotinin, sodium glycocholate, camostat mesilate, vacitracin, or cyclopentadecalactone.

Use of Peptides in Treatments

[0518] In some embodiments, the method includes administering an effective amount of a peptide as described herein to a subject in need thereof.

[0519] In one embodiment, the method includes administering an effective amount of a peptide as described herein to a subject in need thereof.

[0520] TfR can be expressed in various tissues such as the brain, the stomach, the liver, of the gall bladder. Hence, the TfR-binding peptides of the present disclosure can be used in the diagnosis and treatment of disease and conditions associated with various tissues and organs. For example, drug delivery to these tissues and organs can be improved by using the herein described peptides and peptide constructs carrying a diagnostic and/or therapeutic payload.

[0521] The term “effective amount,” as used herein, refers to a sufficient amount of an agent or a compound being administered which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system.

Compositions containing such agents or compounds can be administered for prophylactic, enhancing, and/or therapeutic treatments. An appropriate “effective” amount in any individual case can be determined using techniques, such as a dose escalation study.

[0522] The methods, compositions, and kits of this disclosure can comprise a method to prevent, treat, arrest, reverse, or ameliorate the symptoms of a condition. The treatment can comprise treating a subject (e.g., an individual, a domestic animal, a wild animal, or a lab animal afflicted with a disease or condition) with a peptide of the disclosure. The disease can be a cancer or tumor. In treating the disease, the peptide can contact the tumor or cancerous cells. The subject can be a human. Subjects can be humans; non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. A subject can be of any age. Subjects can be, for example, elderly adults, adults, adolescents, pre-adolescents, children, toddlers, infants, and fetuses in utero.

[0523] Treatment can be provided to the subject before clinical onset of disease. Treatment can be provided to the subject after clinical onset of disease. Treatment can be provided to the subject after 1 day, 1 week, 6 months, 12 months, or 2 years or more after clinical onset of the disease. Treatment can be provided to the subject for more than 1 day, 1 week, 1 month, 6 months, 12 months, 2 years or more after clinical onset of disease. Treatment can be provided to the subject

for less than 1 day, 1 week, 1 month, 6 months, 12 months, or 2 years after clinical onset of the disease. Treatment can also include treating a human in a clinical trial. A treatment can comprise administering to a subject a pharmaceutical composition, such as one or more of the pharmaceutical compositions described throughout the disclosure. A treatment can comprise a once daily dosing. A treatment can comprise delivering a peptide of the disclosure to a subject, either intravenously, subcutaneously, intramuscularly, by inhalation, dermally, topically, by intra-articular injection, orally, sublingually, intrathecally, transdermally, intranasally, via a peritoneal route, directly into a tumor e.g., injection directly into a tumor, directly into the brain, e.g., via and intracerebral ventricle route, or directly onto a joint, e.g. via topical, intra-articular injection route. A treatment can comprise administering a peptide-active agent complex to a subject, either intravenously, subcutaneously, intramuscularly, by inhalation, by intra-articular injection, dermally, topically, orally, intrathecally, transdermally, intranasally, parenterally, orally, via a peritoneal route, nasally, sublingually, or directly onto cancerous tissues.

[0524] Binding of the herein described peptides and peptide constructs (e.g., peptide conjugates or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via vesicular transcytosis) or a cell membrane (e.g., via endocytosis such as receptor-mediated endocytosis) or accumulation in a tissue can have implications in a number of diseases, conditions, or disorders such as those associated with cell growth, cell proliferation, angiogenesis, organogenesis, tumor progression, and/or metastasis, cell death (e.g., senescence), neurodegeneration, and pain in various cells, tissues and organs, particularly those cells tissues or organs that express and/or overexpress TfR. In some embodiments, peptide constructs (e.g., peptide-NT constructs) interact with dopamine-signaling neurons to reduce pain. Compositions comprising any one of the peptides disclosed herein that bind TfR, or a pharmaceutical composition thereof, can be used in a method of treating, preventing, or diagnosing a cancer, tumor progression, dysregulated cell growth in a variety of cancers including ovarian cancer, colon cancer, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma, and lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, and diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, or skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma,

melanoma, head and neck cancer, a neuroblastoma, or a CMYC-overexpressing cancer. In some embodiments, the TfR-binding peptides of the present disclosure enable delivery to tumor cells, bone marrow cells (e.g., erythroid precursor cells), spleen cells, immune cells, cancers of the salivary gland, or muscle cells. The TfR-binding peptides of the present disclosure can be used in combination with drug used for the treatment or prevention of neurological and CNS-related disorder (e.g., neurodegenerative diseases). Additional diseases, disorders, or conditions can include multiple myeloma, plastic anemia, myelodysplasia, and related bone marrow failure syndromes, myeloproliferative diseases, acute and chronic myeloid leukemia, malignancies of lymphoid cells, hematologic malignancies, plasma cell disorders, skeletal muscle disorder, myopathy, muscular dystrophy (e.g., Becker muscular dystrophy, Duchenne muscular dystrophy, Emery-Dreifuss muscular dystrophy, Facioscapulohumeral muscular dystrophy, Myotonia congenita, and myotonic dystrophy), and chronic obstructive pulmonary disorder. In some embodiments, compositions/peptides disclosed herein are used to treat dysregulated cell growth, cancer, tumor, and/or metastasis associated with any of the following cell, tissue, or organ types: brain, lung, liver, spleen, kidney, lymph node, small intestine, blood cell, bone marrow cell, hematopoietic stem cell, pancreatic, colon, stomach, cervix, breast, endometrial, muscular, connective, prostate, testicle, ovarian, skin, head and neck, esophageal, oral tissue, and bone marrow. In further embodiments, compositions/peptides disclosed herein are used to treat nay of the following: sarcoma, osteosarcoma, muscle-derived sarcoma, hepatocellular carcinoma, malignant mesothelioma, schwannoma, meningioma, renal carcinoma, cholangiocarcinoma, bile duct hamartoma, soft tissue carcinoma, myeloma, multiple myeloma, leukemia, lymphoma, ovarian carcinoma, colonic adenoma, T cell acute lymphoblastic leukemia, gastrointestinal hyperplasia, fibrosarcoma, pancreatic ductal metaplasia, squamous cell carcinoma, kaposi sarcoma, and HIV-induced non-Hodgkin's lymphoma.

[0525] In various embodiments, the present disclosure provides methods and compositions that enable TfR-mediated transport across cellular layers (e.g., endothelial cells or epithelial cells) or cell membranes. In addition to the BBB, various other cells, tissues, and organs express TfR. Single cells expressing TfR can include hepatocytes, erythrocytes and erythrocyte precursors in bone marrow, immune cells, stem cells, and rapidly dividing cells. Tissues and organs expressing TfR can include the brain (e.g., cerebral cortex, hippocampus, caudate, cerebellum), endocrine tissues (e.g., thyroid, parathyroid, and adrenal glands), bone marrow and immune system (e.g., appendix, lymph node, tonsil, spleen), muscle tissues (e.g., heart, skeletal, and smooth muscle),

liver, gallbladder, pancreas, gastrointestinal tract (e.g., oral mucosa, esophagus, stomach, duodenum, small intestine, colon, rectum), kidney, urinary bladder, female tissues (e.g., fallopian tube, breast, vagina, cervix, endometrium, ovary, and placenta), adipose and soft tissue, and skin. Thus, the TfR-binding peptides of the present disclosure can be used to target these cells, tissues, and organs and deliver an active agent to these cells, tissues, and organs via, for example, TfR-mediated transcytosis (e.g., across cellular barrier such as the BBB) or TfR-mediated endocytosis (e.g., across cell membranes into cells).

[0526] In various embodiments, the present disclosure provides methods and compositions that enable TfR-mediated transport and delivery to tumor cells expressing TfR. Cancers overexpressing TfR can include ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer. Brain cancers include but are not limited to glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, and diffuse intrinsic pontine glioma.

[0527] TfR-binding peptide constructs that can be used to prevent and/or treat a cancer are those comprising a TfR binding peptide and an active agent with anti-tumor activity such as an IL15, a fusion of IL15/IL15Ra, IFN γ , and anti-CD3 agents. Examples of TfR-binding peptide constructs comprising an active agent with anti-tumor activity are those having an amino acid sequence set forth in any one of SEQ ID NO: 225 – SEQ ID NO: 332). An active agent with active tumor activity (e.g., IL15, a fusion of IL15/IL15Ra, IFN γ , or anti-CD3 agents) may be fused to a TfR-binding peptide (e.g., any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358) to form a peptide-active agent construct.

[0528] Binding of the herein described peptides and peptide constructs (e.g., peptide conjugates or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via TfR-mediated vesicular transcytosis) or a cell

membrane (e.g., via TfR-mediated endocytosis) can have implications in a number of diseases, conditions, or disorders associated with chronic pain (e.g., headaches or migraine), neuropathic pain, obesity, insulin resistance, opioid addiction, or other neurologic or psychiatric disorders in a subject (e.g., a human). In some embodiments, peptide constructs (e.g., peptide-NT constructs) interact with dopamine-signaling neurons to reduce pain.

[0529] Binding of the herein described peptides and peptide constructs (e.g., peptide conjugates, fusion peptides, or recombinantly produced peptide constructs) to TfR and subsequent transport across a cell layer or barrier such as the BBB (e.g., via vesicular transcytosis) or a cell membrane (e.g., via endocytosis) can have implications in a number of diseases, conditions, or disorders associated with neurodegeneration. Neurodegenerative diseases that can be treated, prevented, or diagnosed with the herein described TfR-binding peptides can include Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. In some cases, the TfR-binding peptide can be used in combination with BACE inhibitors, galantamine, amantadine, benztropine, biperiden, bromocriptin, carbidopa, donepezil, entacapone, levodopa, pergolide, pramipexole, procyclidine, rivastigmine, ropinirole, selegiline, tacrine, tolcapone, or trihexyphenidyl to treat and/or prevent a neurodegenerative disease.

[0530] In some embodiments, modulation (e.g., inhibition) of ion channels such as Kv1.3 potassium channels using TfR-binding peptides and ion channel modulators such as Kv1.3 potassium channel inhibitor conjugates or peptide constructs can be used for treating or preventing inflammation in the brain. Kv1.3 potassium channels, for instance, can be highly expressed on microglia in the brain. Thus, neuroinflammatory and neurodegenerative diseases such as multiple sclerosis, Alzheimer's, Parkinson's, traumatic brain injury, radiation therapy toxicity and other neurodegenerative and neuroinflammatory diseases can be treated with TfR-binding peptides that are conjugated to, linked to, or fused to a Kv1.3 inhibitor. These diseases can be marked by upregulation of Kv1.3. In some cases, Kv1.3 inhibition can be used for treatment of psoriasis and other non-brain autoimmune diseases due to its effect on effector T cells.

[0531] In some embodiments, the TfR-binding peptides of the present disclosure can be used for the treatment and prevention of Crohn's disease or, more generally, inflammatory bowel diseases. In some embodiments, the TfR-binding peptides of the present disclosure show high uptake and retention in glandular cells of the intestine, which can express high amounts of TfR.

[0532] In some embodiments, the peptides or peptide constructs of the present disclosure (e.g., SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361) are used to access and treat these disorders due to their enhanced stability in various biological environments, including low pH, protease-rich environments, acidic environments, reducing environments, or environments with varying temperatures.

[0533] In some embodiments, compositions comprise any one of the TfR-binding peptides disclosed herein and are capable of transporting one or more cargo molecules or active agents per TfR-binding peptides across cell layers or cell barriers such as endothelial layers (e.g., the BBB) or epithelial layers, or cell membranes. In some embodiments, the cargo molecule or active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15, a fusion of IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter such as neurotensin.

Peptide Kits

[0534] In one aspect, peptides described herein can be provided as a kit. In another embodiment, peptide constructs described herein can be provided as a kit. In another embodiment, a kit comprises amino acids encoding a peptide described herein, a vector, a host organism, and an instruction manual. In some embodiments, a kit includes written instructions on the use or administration of the peptides.

Numbered Embodiments

[0535] The following embodiments recite non-limiting permutations of combinations of features disclosed herein. Other permutations of combinations of features are also contemplated. In

particular, each of these numbered embodiments is contemplated as depending from or relating to every previous or subsequent numbered embodiment, independent of their order as listed. 1. A composition comprising an engineered transferrin receptor (TfR)-binding peptide, or a variant, homolog, fragment, or analog thereof. 2. A composition comprising a peptide construct, wherein the peptide construct comprises: a) a transferrin receptor (TfR)-binding peptide, or a variant, homolog, fragment, or analog thereof; and b) an active agent, wherein the peptide is conjugated to, linked to, or fused to the active agent. 3. The composition of any one of embodiments 1-2, wherein the TfR is a human TfR, or murine TfR, or a human TfR and murine TfR. 4. A composition comprising an engineered peptide having at least 75%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a functional fragment thereof. 5. The composition of any one of embodiments 1-4, wherein the peptide penetrates a cellular layer comprising endothelial cells. 6. The composition of embodiment 5, wherein the endothelial cells are brain vascular endothelial cells. 7. The composition of any one of embodiments 1-6, wherein the the peptide penetrates a cellular layer comprising a blood brain barrier (BBB). 8. The composition of any one of embodiments 1-7, wherein the peptide penetrates a cellular layer comprising endothelial cells via TfR-mediated transcytosis. 9. The composition of any one of embodiments 1-8, wherein the peptide penetrates a membrane and wherein the membrane is the membrane of a cell. 10. The composition of embodiment 9, wherein the cell expresses a TfR. 11. The composition of any one of embodiments 9-10, wherein the peptide penetrates the membrane of the cell via TfR-mediated endocytosis. 12. The composition of any one of embodiments 1-11, wherein the peptide is capable of localizing in or with any cell that expresses a TfR. 13. The composition of any one of embodiments 1-12, wherein the peptide remains bound to a cell surface membrane expressing TfR. 14. The composition of any one of embodiments 1-13, wherein the peptide is internalized into a cell by binding TfR. 15. The composition of any one of embodiments 1-14, wherein the peptide is delivered across the blood-brain barrier to the CNS by binding and then dissociating from TfR. 16. The composition of any one of embodiments 1-15, wherein the peptide is delivered across the blood-brain barrier by receptor-mediated transcytosis. 17. The composition of any one of embodiments 9-16, wherein the cell is a CNS cell, an erythrocyte, an erythrocyte precursor cell, an immune cell, a stem cell, a muscle cell, a brain cell,

a thyroid cell, a parathyroid cell, an adrenal gland cell, a bone marrow cell, an appendix cell, a lymph node cell, a tonsil cell, a spleen cell, a muscle cell, a liver cell, a gallbladder cell, a pancreas cell, a cell of the gastrointestinal tract, a glandular cell, a kidney cell, a urinary bladder cell, an endothelial cell, an epithelial cell, a choroid plexus epithelial cell, a glial cell, an astrocyte, or a cell associated with a nervous system. 18. The composition of any one of embodiments 9-17, wherein the cell is a tumor cell. 19. The composition of embodiment 18, wherein the tumor cell is an ovarian cancer cell, a colon cancer cell, a lung cancer cell, a cancer cell located in the bone or bone marrow, a glioblastoma cell, an astrocytoma cell, a glioma cell, a medulloblastoma cell, an ependymoma cell, a choroid plexus carcinoma cell, a midline glioma cell, a diffuse intrinsic pontine glioma (DIPG) cell, a breast cancer cell, a liver cancer cell, a colon cancer cell, a brain cancer cell, a spleen cancer cell, a cancer cell of the salivary gland, a kidney cancer cell, a muscle cancer cell, a bone marrow cell cancer cell, a skin cancer cell, a genitourinary cancer cell, an osteosarcoma cell, a muscle-derived sarcoma cell, a melanoma cell, a head and neck cancer cell, a neuroblastoma cell, a prostate cancer cell, a bladder cancer cell, an acute lymphocytic leukemia cell, an acute myeloid leukemia cell, a chronic lymphocytic leukemia cell, a chronic myeloid leukemia cell, a Hodgkin lymphoma cell, a Non-Hodgkin lymphoma cell, or a CMYC-overexpressing cancer cell. 20. The composition of any one of embodiments 1-19, wherein the peptide localizes to an organ or tissue. 21. The composition of embodiment 20, wherein the organ or tissue is brain, thyroid, parathyroid, adrenal glands, bone marrow, appendix, lymph node, tonsil, spleen, muscle, liver, gallbladder, pancreas, gastrointestinal tract, kidney, urinary bladder, fallopian tube, breast, vagina, cervix, endometrium, ovary, placenta, and skin. 22. The composition of any one of embodiments 9-21, wherein the cell or organ in which the peptide localizes overexpresses a TfR. 23. The composition of embodiment 22, wherein the cell or organ overexpressing the TfR is a tumor cell or cancerous tissue. 24. The composition of any one of embodiments 22-23, wherein the peptide localizes to the parenchyma of the organ. 25. The composition of any one of embodiments 1-24, wherein the peptide binds at least one amino acid residues selected from the group of residues 506-510, 523-531, or 611-662 corresponding to SEQ ID NO: 363 of human TfR1 (UniProtKB - P02786). 26. The composition of any one of embodiments 1-25, wherein the peptide comprises SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344. 27. The composition of any one of embodiments 1-26, wherein the peptide comprises a surface interface residue corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and

Q51, or any combination thereof, with reference to SEQ ID NO: 32. 28. The composition of any one of embodiments 1-27, wherein the peptide comprises hydrophilic surface-distal residues corresponding to any one of the positions 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. 29. The composition of embodiment 28, wherein the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 30. The composition of any one of embodiments 26-29, wherein the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. 31. The composition of any one of embodiments 1-30, wherein the peptide comprises hydrophilic residues at a position corresponding to any one of 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. 32. The composition of embodiment 28, wherein the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 33. The composition of any one of embodiments 31-32, wherein the peptide comprises a hydrophilic residue corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. 34. The composition of any one of embodiments 1-33, wherein the peptide comprises hydrophobic residues at a position corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. 35. The composition of embodiment 34, wherein the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. 36. The composition of any one of embodiments 34-35, wherein the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. 37. The composition of any one of embodiments 1-36, wherein the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. 38. The composition of embodiment 37, wherein the aliphatic residue is selected from the amino acids A, M, I, L, or V. 39. The composition of any one of embodiments 37-38, wherein the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. 40. The composition of any one of embodiment 1-39, wherein the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. 41. The composition of any one of embodiment 1-40, wherein the peptide comprises three disulfide bonds. 42. The composition of any one of embodiment 1-41, wherein the peptide comprises at least two, at least four, or at least six cysteine residues. 43. The composition of any one of embodiment 1-42, wherein the peptide

comprises six cysteine residues. 44. The composition of any one of embodiments 1-43, wherein the peptide comprises cysteine residues at positions corresponding to C6, C10, C20, C34, C44, and C48, or any combination thereof, with reference to SEQ ID NO: 32. 45. The composition of any one of embodiments 1-44, wherein at least one cysteine residue of the peptide participates in a disulfide bond. 46. The composition of embodiment 43, wherein the six cysteine residues participate in disulfide bonds. 47. The composition of any one of embodiments 1-46, wherein a disulfide bond increases the stability of the peptide. 48. The composition of any one of embodiments 1-47, wherein the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. 49. The composition of any one of embodiments 1-48, wherein the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof. 50. The composition of any one of embodiments 1-49, wherein the peptide comprises the sequence set forth in SEQ ID NO: 1. 51. The composition of any one of embodiments 1-49, wherein the peptide comprises the sequence set forth in SEQ ID NO: 2. 52. The composition of any one of embodiments 1-49, wherein the peptide comprises the sequence set forth in SEQ ID NO: 4. 53. The composition of any one of embodiments 1-49, wherein the peptide comprises the sequence set forth in SEQ ID NO: 30. 54. The composition of any one of embodiments 1-49, wherein the peptide comprises the sequence set forth in SEQ ID NO: 32. 55. The composition of any one of embodiment 1-54, wherein the peptide comprising three disulfide bonds is derived from a cystine-dense peptide (CDP). 56. The composition of any one of embodiment 1-55, wherein the peptide comprising three disulfide bonds is a CDP. 57. The composition of any one of embodiments 1-56, wherein the peptide competes with transferrin, or a homolog, variant, or fragment thereof, for binding to TfR. 58. The composition of any one of embodiments 1-57, wherein the peptide sequesters TfR or prevents another protein from binding to TfR. 59. The composition of any one of embodiments 1-58, wherein the peptide has equal or greater affinity or specificity for TfR as compared to that of transferrin, or an endogenous derivative thereof. 60. The composition of any one of embodiments 1-59, wherein the peptide has equal or lower dissociation constant for TfR as compared to that of transferrin, or an endogenous

derivative thereof. 61. The composition of embodiment 60, wherein the peptide has a KD of less than 40 nM, less than 30 nM, less than 20 nM, less than 10 nM, less than 1 nM, or less than 0.1 nM. 62. The composition of any one of embodiments 1-61, wherein the peptide penetrates or localizes into a target cell. 63. The composition of any one of embodiments 1-62, wherein the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. 64. The composition of embodiment 63, wherein the cell-penetrating moiety comprises polycations, polyorganic acids, endosomal releasing polymers, poly(2-propylacrylic acid), poly(2-ethylacrylic acid), Tat peptide, Arg patch, a knotted peptide, CysTAT, S19-TAT, R8 (SEQ ID NO: 73), pAntp, Pas-TAT, Pas-R8 (SEQ ID NO: 76), Pas-FHV, Pas-pAntP, F2R4 (SEQ ID NO: 79), B55, aurein, IMT-P8, BR2, OMOTAG1, OMOTAG2, pVEC, SynB3, DPV1047, C105Y, Transportan, MTS, hLF, PFVYLI (SEQ ID NO: 93), maurocalcine, imperatoxin, hadrucalin, hemicalcin, opicalcin-1, opicalcin-2, midkine (62-104), MCoTI-II, chlorotoxin, DRI-TAT, cFΦR4 (SEQ ID NO: 96), R6W3 (SEQ ID NO: 421), myristate, yBBR, or a fragment or variant thereof, or any combination thereof. 65. The composition of embodiment 63, wherein the cell-penetrating moiety is maurocalcin, imperatoxin, hadrucalin, hemicalcin, opicalcin-1, opicalcin-2, midkine (62-104), MCoTI-II, or chlorotoxin. 66. The composition of embodiment 63, wherein the cell-penetrating moiety is any one of SEQ ID NO: 71 – SEQ ID NO: 96, SEQ ID NO: 98 – SEQ ID NO: 101, SEQ ID NO: 116 – SEQ ID NO: 126, or SEQ ID NO: 403 – SEQ ID NO: 455. 67. The composition of any one of embodiments 1-66, wherein the peptide targets, homes, accumulates in, or selectively penetrates a tissue or cell. 68. The composition of any one of embodiments 1-67, wherein the peptide localizes into a target cell's nucleus. 69. The composition of embodiment 68, wherein the peptide localizes into the target cell's nucleus by formulation with, fusion to, or conjugation to a nuclear localization signal peptide or moiety. 70. The composition of any one of embodiments 1-69, wherein the peptide is further conjugated to, linked to, or fused to a moiety or a second peptide that targets, homes, accumulates in, or selectively penetrates a cancerous tissue or cancer cell. 71. The composition of any one of embodiments 1-70, wherein the moiety or a second peptide is a knotted peptide or variant or fragment thereof. 72. The composition of any one of embodiments 68-71, wherein the target cell is a cancerous cell or a tumor cell. 73. The composition of any one of embodiments 68-72, wherein the target cell is a CNS cell, an erythrocyte, an erythrocyte precursor cell, an immune cell, a stem cell, a muscle cell, a brain cell, a thyroid cell, a parathyroid cell, an adrenal gland cell, a bone marrow cell, an appendix cell, a lymph node cell, a tonsil cell, a spleen cell, a muscle cell, a liver cell, a gallbladder cell, a

pancreas cell, a cell of the gastrointestinal tract, a glandular cell, a kidney cell, a urinary bladder cell, an endothelial cell, an epithelial cell, a choroid plexus epithelial cell, a glial cell, an astrocyte, or a cell associated with a nervous system. 74. The composition of any one of embodiments 1-73, wherein the peptide comprises a disulfide through disulfide knot. 75. The composition of any one of embodiments 1-74, wherein the peptide comprises a plurality of disulfide bridges formed between cysteine residues. 76. The composition of any one of embodiments 1-75, wherein the peptide comprises three or more disulfide bridges formed between cysteine residues, wherein one of the disulfide bridges passes through a loop formed by two other disulfide bridges. 77. The composition of any one of embodiments 1-76, wherein the peptide comprises no disulfide bond or a disulfide knot. 78. The composition of any one of embodiments 1-77, wherein at least one amino acid residue of the peptide is in an L configuration or, wherein at least one amino acid residue is in a D configuration. 79. The composition of any of embodiments 1-78, wherein the peptide comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 positively charged residues. 80. The composition of any of embodiments 1-79, wherein the peptide comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 consecutive positively charged residues. 81. The composition of any of embodiments 1-80, wherein the positively charged residues are Arg, Lys, or any combination thereof. 82. The composition of any of embodiments 1-81, wherein the peptide comprises an Arg patch. 83. The composition of any of embodiments 1-82, wherein the peptide comprises two or more consecutive Arg residues at the N-terminus. 84. The composition of any of embodiments 1-83, wherein the peptide comprises a Tat peptide, or a fragment thereof. 85. The composition of embodiment 84, wherein the Tat peptide comprises a sequence of YGRKKRRQRRR (SEQ ID NO: 99), GRKKRRQRRR (SEQ ID NO: 100), or a fragment or variant thereof. 86. The composition of any of embodiments 84-85, wherein the Tat peptide is appended to the N-terminus or C-terminus of the peptide using a linker having a sequence of (GS)_x (SEQ ID NO: 105) or G_xS_y (SEQ ID NO: 104), wherein x and y is independently 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. 87. The composition of any of embodiments 84-86, wherein the Tat peptide is appended to any residue of the peptide. 88. The composition of any one of embodiments 1-87, wherein the peptide comprises at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, or at least 16 cysteine residues. 89. The composition

of any one of embodiments 1-88, wherein the peptide comprises one or more regions of negatively charged residues. 90. The composition of any one of embodiments 1-89, wherein the peptide comprises one or more regions of positively charged residues. 91. The composition of any one of embodiments 1-90, wherein the peptide sequence comprises at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36, at least 37, at least 38, at least 39, at least 40, at least 41, at least 42, at least 43, at least 44, at least 45, at least 46, at least 47, at least 48, at least 49, at least 50, at least 51, at least 52, at least 53, at least 54, at least 55, at least 56, at least 57, at least 58 residues, at least 59, at least 60, at least 61, at least 62, at least 63, at least 64, at least 65, at least 66, at least 67, at least 68, at least 69, at least 70, at least 71, at least 72, at least 73, at least 74, at least 75, at least 76, at least 77, at least 78, at least 79, at least 80, or at least 81 amino acid residues. 92. The composition of any one of embodiments 1-91, wherein the peptide comprises or is derived from a human protein or peptide. 93. The composition of any one of embodiments 1-92, wherein the peptide is arranged in a multimeric structure with at least one other peptide. 94. The composition of embodiment 93, wherein the multimeric structure comprises a dimer, trimer, tetramer, pentamer, hexamer, or heptamer. 95. The composition of any one of embodiments 1-94, wherein the peptide comprises an isoelectric point within a range from about 3.0 to about 10.0. 96. The composition of any one of embodiments 1-95, wherein the peptide comprises an isoelectric point within a range from about 4.5 to about 8.9. 97. The composition of any one of embodiments 1-96, wherein the peptide comprises a non-uniform charge distribution. 98. The composition of any one of embodiments 1-97, wherein the peptide is stable at physiological or intracellular pH values. 99. The composition of any one of embodiments 1-98, wherein the peptide is stable at a pH of about 7 or 7.5. 100. The composition of any one of embodiments 1-99, wherein the peptide is stable at a pH of about 6.5 to 7.5. 101. The composition of any one of embodiments 1-100, wherein the peptide is stable at pH values less than or equal to about 5.0, less than or equal to about 3.0, or within a range from about 3.0 to about 5.0. 102. The composition of any one of embodiments 1-101, wherein the peptide is stable at pH values within a range from about 5.0 to about 7.0. 103. The composition of any one of embodiments 1-102, wherein the peptide comprises a hydrophobic core. 104. The composition of any one of embodiments 1-103, wherein the peptide is resistant to protease degradation. 105. The composition of embodiment 104, wherein the protease is any one of

pepsin, trypsin, chymotrypsin, serum proteases, intracellular proteases, or any combination thereof. 106. The composition of any one of embodiments 1-105, wherein the peptide is resistant to reduction or is stable in a reducing environment. 107. The composition of embodiment 106, wherein the reduction or reducing environment comprises DTT or GSH. 108. The composition of any one of embodiments 1-107, wherein the peptide is stable at an elevated temperature. 109. The composition of any one of embodiments 1-108, wherein the peptide exhibits an anti-cancer effect. 110. The composition of any one of embodiments 1-109, wherein the peptide comprises one or more chemical modifications. 111. The composition of embodiment 110, wherein the chemical modification modifies one or more pharmacokinetic properties of the peptide. 112. The composition of any one or embodiments 110-111, wherein the chemical modification extends the plasma half-life and/or the biological half-life of the peptide. 113. The composition of any of embodiments 110-112, wherein the chemical modification is blocking the N-terminus of the peptide. 114. The composition of any of embodiments 110-113, wherein the chemical modification is methylation, acetylation, or acylation. 115. The composition of embodiment 110-114, wherein the chemical modification is: methylation of one or more lysine residues or analogue thereof; methylation of the N-terminus; or methylation of one or more lysine residue or analogue thereof and methylation of the N-terminus. 116. The composition of any one of embodiments 1-115, wherein the peptide is linked to an acyl adduct. 117. The composition of any one of embodiments 1-116, wherein an active sequence is grafted onto the peptide or wherein the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct. 118. The compositions of embodiment 117, wherein the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO:70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400, or SEQ ID NO: 401. 119. The composition of any one of embodiments 117-118, wherein the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361. 120. The composition of any one of embodiment 117-119, wherein the active agent is conjugated to, linked to, or fused to the peptide at an N-terminus or a C-terminus of the peptide. 121. The composition of any one of embodiments 117-120, wherein the active

agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. 122. The composition of embodiment 121, wherein the nucleic acid is DNA. 123. The composition of embodiment 121, wherein the nucleic acid is RNA. 124. The composition of any one of embodiments 117-123, wherein 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 active agents are linked to the peptide by a linker. 125. The composition of any one of embodiments 117-124, wherein the peptide is linked to the active agent via a cleavable linker or a pH sensitive linker. 126. The composition of any one of embodiments 117-125, wherein the peptide is linked to the active agent at an N-terminus, at the epsilon amine of an internal lysine residue, at the carboxylic acid of an aspartic acid or glutamic acid residue, or a C-terminus of the peptide by a linker. 127. The composition of any one of embodiments 117-126, further comprising a non-natural amino acid, wherein the non-natural amino acid is an insertion, appendage, or substitution for another amino acid. 128. The composition of embodiment 127, wherein the peptide is linked to the active agent at the non-natural amino acid by a linker. 129. The composition of any one of embodiments 124-128, wherein the linker comprises an amide bond, an ester bond, an ether bond, a triazole, a macrocycle, an oxime bond, a carbamate bond, a carbonate bond, a hydrazone bond, an azo bond, an oxime bond, a disulfide bond, a thioester bond, a thioether bond, a carbon-carbon single, double, or triple bond, a disulfide bond, a two carbon bridge between two cysteines, a three carbon bridge between two cysteines, or a carbon-nitrogen bond. 130. The composition of any one of embodiments 125-129, wherein the cleavable linker comprises a cleavage site for a protease. 131. The composition of embodiment 130, wherein the protease is a matrix metalloproteinases, thrombin, cathepsins, or beta-glucuronidase. 132. The composition of any one of embodiments 124-131, wherein the linker is a hydrolytically labile linker. 133. The composition of any one of embodiments 1-132, wherein the peptide is linked to the active agent via a noncleavable linker. 134. The composition of any one of embodiments 1-133, wherein the active agent is linked via a linker, wherein the linker is selected from TABLE 7 or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. 135. The composition of any one of embodiments 117-134, wherein the active agent is an anti-cancer agent. 136. The composition of any one of embodiments 117-134, wherein the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFNgamma agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a

Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter. 137. The composition of embodiment 136, wherein the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. 138. The composition of embodiment 137, wherein the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. 139. The composition of any one of embodiments 137-138, wherein the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. 140. The composition of any one of embodiments 137-139, wherein the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350. 141. The composition of any one of embodiments 117-140, wherein the peptide construct is capable of binding and activating a neurotensin receptor. 142. The composition of any one of embodiments 117-141, wherein the peptide is capable of reducing a level of pain in a subject. 143. The composition of any one of embodiments 117-142, wherein the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, pain, epilepsy, a mood disorder, or an endocrine disorder. 144. The composition of any one of embodiments 117-143, wherein the peptide interacts with dopamine signaling neurons. 145. The composition of any one of embodiments 117-144, wherein the peptide increases dopamine release in neurons. 146. The composition of any one of embodiments 1-145, further comprising a half-life modifying agent coupled to the peptide. 147. The composition of embodiment 146, wherein the half-life modifying agent comprises a polymer, a polyethylene glycol (PEG), a hydroxyethyl starch, polyvinyl alcohol, a water soluble polymer, a zwitterionic water soluble polymer, a water soluble poly(amino acid), a water soluble polymer of proline, alanine and serine, a water soluble polymer containing glycine, glutamic acid, and serine, an Fc region, a fatty acid, palmitic acid, or a molecule that binds to albumin. 148. The composition of any one of embodiments 1-147, further comprising a detectable agent coupled to the peptide by a linker. 149. The composition of embodiment 148, wherein the detectable agent is conjugated to, linked to, or fused with the peptide at an N-terminus or a C-terminus of the peptide. 150. The composition any one of

embodiments 148-149, wherein 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 detectable agents are linked to the peptide. 151. The composition of any one of embodiments 148-150, wherein the peptide is linked to the detectable agent via a cleavable linker. 152. The composition of any one of embodiments 148-151, wherein the peptide is linked to the detectable agent at an N-terminus, at the epsilon amine of an internal lysine residue, or a C-terminus of the peptide by a linker. 153. The composition of any one of embodiments 117-152, further comprising a non-natural amino acid, wherein the non-natural amino acid is an insertion, appendage, or substitution for another amino acid. 154. The composition of any one of embodiments 148-153, wherein the peptide is linked to the detectable agent at the non-natural amino acid by a linker. 155. The composition of any one of embodiments 148-154, wherein the linker comprises an amide bond, an ester bond, an ether bond, a triazole, a macrocycle, an oxime bond, a carbamate bond, a carbonate bond, a hydrazone bond, an azo bond, an oxime bond, a disulfide bond, a thioester bond, a thioether bond, a carbon-carbon single, double, or triple bond, a disulfide bond, a two carbon bridge between two cysteines, a three carbon bridge between two cysteines, or a carbon-nitrogen bond. 156. The composition of embodiment 151-155, wherein the cleavable linker comprises a cleavage site for matrix metalloproteinases, thrombin, cathepsins, or beta-glucuronidase. 157. The composition of any one of embodiments 148-150, wherein the peptide is linked to the detectable agent via a stable linker. 158. The composition of any one of embodiments 148-157, wherein the detectable agent is linked via a linker, wherein the linker is selected from TABLE 7 or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. 159. The composition of any one of embodiments 148-158, wherein the detectable agent is a fluorophore, a near-infrared dye, a contrast agent, a nanoparticle, a metal-containing nanoparticle, a metal chelate, an X-ray contrast agent, a PET agent, a radionuclide, or a radionuclide chelator. 160. The composition of any one of embodiments 148-159, wherein the detectable agent is a fluorescent dye. 161. A pharmaceutical composition comprising the composition of any one of embodiments 1-160, or a salt thereof, and a pharmaceutically acceptable carrier. 162. The pharmaceutical composition of embodiment 161, wherein the pharmaceutical composition is formulated for administration to a subject. 163. The pharmaceutical composition of any one of embodiments 161-162, wherein the pharmaceutical composition is formulated for oral administration, intravenous administration, subcutaneous administration, intramuscular administration, or a combination thereof. 164. A method for transporting a composition across a cell layer, the method comprising: a) contacting the cell layer with the composition; and b) transporting the composition across the cell layer. 165.

The method of embodiment 164, wherein the composition comprises an engineered peptide. 166. The method of embodiment 165, wherein the peptide comprises an amino acid sequence comprising any one of SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344, or a functional fragment thereof. 167. The method of any one of embodiments 165-166, wherein the peptide is a transferrin receptor (TfR)-binding peptide. 168. The method of any one of embodiments 165-167, wherein the peptide comprises a surface interface residues corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. 169. The method of any one of embodiments 165-168, wherein the peptide comprises a hydrophilic surface-distal residue at a position corresponding to 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. 170. The method of embodiment 169, wherein the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 171. The method of any one of embodiments 169-170, wherein the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. 172. The method of any one of embodiments 165-171, wherein the peptide comprises hydrophilic residues at any one of the positions corresponding to 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. 173. The method of embodiment 172, wherein the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 174. The method of any one of embodiments 172-173, wherein the peptide comprises any one of the hydrophilic residues corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. 175. The method of any one of embodiments 165-174, wherein the peptide comprises hydrophobic residues at any of the positions corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. 176. The method of embodiment 175, wherein the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. 177. The method of any one of embodiments 175-176, wherein the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. 178. The method of any one of embodiments 165-177, wherein the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. 179. The method of embodiment 178, wherein the aliphatic residue is selected from the amino acids A, M, I, L, or V. 180. The method of any one

of embodiments 178-179, wherein the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. 181. The method of any one of embodiment 165-170, wherein the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. 182. The method of any one of embodiments 165-181, wherein the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. 183. The method of any one of embodiments 165-182, wherein the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof. 184. The method of any one of embodiments 165-183, wherein the peptide penetrates or localizes into or on a target cell. 185. The method of any one of embodiments 165-184, wherein an active sequence is grafted onto the peptide or wherein the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct. 186. The method of embodiment 185, wherein the peptide is conjugated to, linked to, or fused to the active agent. 187. The method of embodiment 185 or 186, wherein the active agent is linked via a linker, wherein the linker is selected from TABLE 7 or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. 188. The method of any one of embodiments 185-187, wherein the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. 189. The method of any one of embodiments 185-188, wherein the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400 or SEQ ID NO: 401. 190. The method of any one of embodiments 185-189, wherein the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361. 191. The method of any one of embodiments 185-190, wherein the active agent is an immunotherapeutic agent, a CTLA-

4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter. 192. The method of embodiment 191, wherein the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. 193. The method of embodiment 192, wherein the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. 194. The method of any one of embodiments 192-193, wherein the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. 195. The method of any one of embodiments 192-194, wherein the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350. 196. The method of any one of embodiments 185-195, wherein the peptide construct is capable of binding and activating a neurotensin receptor. 197. The method of any one of embodiments 165-196, wherein the peptide is capable of reducing a level of pain upon administration to a subject. 198. The method of any one of embodiments 165-197, wherein the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, pain, epilepsy, a mood disorder, or an endocrine disorder. 199. The method of any one of embodiments 165-198, wherein the peptide interacts with dopamine signaling neurons. 200. The method of any one of embodiments 165-199, wherein the peptide increases dopamine release in neurons. 201. The method of any one of embodiments 165-200, wherein the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. 202. The method of any one of embodiments 164-201, wherein the cell layer is an endothelial cell layer or an epithelial cell layer. 203. The method of embodiment 202, wherein the endothelial cell layer is the blood brain barrier. 204. The method of embodiment 202, wherein the endothelial cell layer is an intestinal endothelial cell layer. 205. The method of embodiment 202, wherein the epithelial cell layer is an intestinal epithelial cell layer. 206. The

method of any one of embodiments 164-205, wherein the transporting comprises binding of the composition to TfR. 207. The method of any one of embodiments 164-206, wherein the transporting further comprises transcytosis. 208. The method of embodiment 207, wherein the transcytosis is vesicular transcytosis. 209. The method of any one of embodiments 207-208, wherein the transcytosis is TfR-mediated. 210. The method of embodiment 209, wherein the transporting further comprises dissociation of the composition from the TfR. 211. A method of treating a condition in a subject in need thereof, the method comprising: a) administering to the subject a composition, and b) transporting the composition across a cell layer. 212. The method of embodiment 211, wherein the composition comprises an engineered peptide. 213. The method of embodiment 212, wherein the peptide comprises an amino acid sequence of any one of SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344. 214. The method of any one of embodiments 212-213, wherein the peptide is a transferrin receptor (TfR)-binding peptide. 215. The method of any one of embodiments 212-214, wherein the peptide comprises a surface interface residues corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. 216. The method of any one of embodiments 212-215, wherein the peptide comprises a hydrophilic surface-distal residues at any one of the positions corresponding to 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. 217. The method of embodiment 216, wherein the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 218. The method of any one of embodiments 216-217, wherein the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. 219. The method of any one of embodiments 212-218, wherein the peptide comprises hydrophilic residues at any one of the positions corresponding to 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. 220. The method of embodiment 219, wherein the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 221. The method of any one of embodiments 219-220, wherein the peptide comprises any one of the hydrophilic residues corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. 222. The method of any one of embodiments 212-221, wherein the peptide comprises hydrophobic residues at any of the positions corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. 223. The method of

embodiment 222, wherein the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. 224. The method of any one of embodiments 221-223, wherein the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. 225. The method of any one of embodiments 212-224, wherein the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. 226. The method of embodiment 225, wherein the aliphatic residue is selected from the amino acids A, M, I, L, or V. 227. The method of any one of embodiments 225-226, wherein the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. 228. The method of any one of embodiment 212-227, wherein the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. 229. The method of any one of embodiments 212-228, wherein the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. 230. The method of any one of embodiments 212-229, wherein the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof. 231. The method of any one of embodiments 212-230, wherein the peptide penetrates or localizes into a target cell. 232. The method of any one of embodiments 212-231, wherein an active sequence is grafted onto the peptide or wherein the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct. 233. The method of embodiment 232, wherein the peptide is conjugated to, linked to, or fused to the active agent. 234. The method of embodiment 232 or 233, wherein the active agent is linked via a linker, wherein the linker is selected from TABLE 7 or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. 235. The method of any one of embodiments 232-234, wherein the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. 236. The method of any one of embodiments 232-235, wherein the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO:

35, SEQ ID NO: 68 – SEQ ID NO:70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400 or SEQ ID NO: 401. 237. The method of any one of embodiments 232-236, wherein the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361. 238. The method of any one of embodiments 232-237, wherein the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFNgamma agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter. 239. The method of embodiment 238, wherein the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. 240. The method of embodiment 239, wherein the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. 241. The method of any one of embodiments 239-240, wherein the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. 242. The method of any one of embodiments 239-241, wherein the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350. 243. The method of any one of embodiments 232-242, wherein the peptide construct is capable of binding and activating a neurotensin receptor. 244. The method of any one of embodiments 232-243, wherein the peptide is capable of reducing a level of pain upon administration to the subject. 245. The method of any one of embodiments 212-244, wherein the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, pain, epilepsy, a mood disorder, or an endocrine disorder. 246. The method of any one of embodiments 212-245, wherein the peptide interacts with dopamine signaling neurons. 247. The method of any one of embodiments 212-246, wherein

the peptide increases dopamine release in neurons. 248. The method of any one of embodiments 212-247, wherein the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. 249. The method of any one of embodiments 212-248, wherein the cell layer is an endothelial cell layer or an epithelial cell layer. 250. The method of embodiment 249, wherein the endothelial cell layer is the blood brain barrier. 251. The method of embodiment 249, wherein the endothelial cell layer is an intestinal endothelial cell layer. 252. The method of embodiment 249, wherein the epithelial cell layer is an intestinal epithelial cell layer. 253. The method of any one of embodiments 211-252, wherein the transporting comprises binding of the composition to TfR. 254. The method of any one of embodiments 211-253, wherein the transporting further comprises transcytosis. 255. The method of embodiment 254, wherein the transcytosis is vesicular transcytosis. 256. The method of any one of embodiments 254-255, wherein the transcytosis is TfR-mediated. 257. The method of any one of embodiments 253-256, wherein the transporting further comprises dissociation of the composition from the TfR. 258. The method of any one of embodiments 253-257, wherein the transporting further comprises releasing the active agent from the composition. 259. A method of treating a condition in a subject in need thereof, the method comprising administering to the subject an amount of a composition based on the expression of TfR in the subject. 260. The method of embodiment 259, wherein the composition comprises an engineered peptide. 261. The method of embodiment 250, wherein the peptide comprises an amino acid sequence of any one of SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344. 262. The method of any one of embodiments 260-261, wherein the peptide is a transferrin receptor (TfR)-binding peptide. 263. The method of any one of embodiments 260-262, wherein the peptide comprises a surface interface residues corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32. 264. The method of any one of embodiments 260-263, wherein the peptide comprises a hydrophilic surface-distal residues at any one of the positions corresponding to 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32. 265. The method of embodiment 264, wherein the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 266. The method of any one of embodiments 264-265, wherein the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32. 267. The method

of any one of embodiments 260-266, wherein the peptide comprises hydrophilic residues at any one of the positions corresponding to 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32. 268. The method of embodiment 267, wherein the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T. 269. The method of any one of embodiments 260-268, wherein the peptide comprises any one of the hydrophilic residues corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32. 270. The method of any one of embodiments 260-269, wherein the peptide comprises hydrophobic residues at any of the positions corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32. 271. The method of embodiment 270, wherein the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof. 272. The method of any one of embodiments 270-271, wherein the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32. 273. The method of any one of embodiments 260-272, wherein the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32. 274. The method of embodiment 273, wherein the aliphatic residue is selected from the amino acids A, M, I, L, or V. 275. The method of any one of embodiments 273-274, wherein the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32. 276. The method of any one of embodiment 260-275, wherein the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds. 277. The method of any one of embodiments 260-276, wherein the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof. 278. The method of any one of embodiments 260-277, wherein the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof. 279. The method of any one of embodiments 260-278, wherein the peptide penetrates or localizes into a target cell. 280. The method of any one of embodiments 260-279, wherein an active sequence is grafted onto the peptide or wherein the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide

construct. 281. The method of embodiment 280, wherein the peptide is conjugated to, linked to, or fused to the active agent. 282. The method of embodiment 280 or 281, wherein the active agent is linked via a linker, wherein the linker is selected from TABLE 7 or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125. 283. The method of any one of embodiments 280-282, wherein the active agent is an antibody, antibody fragment, Fc, single chain Fv, polypeptide, peptide, small molecule, or nucleic acid. 284. The method of any one of embodiments 280-283, wherein the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400 or SEQ ID NO: 401. 285. The method of any one of embodiments 280-284, wherein the peptide construct is SEQ ID NO: 359, SEQ ID NO: 360, or SEQ ID NO: 361. 286. The method of any one of embodiments 280-285, wherein the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter. 287. The method of embodiment 286, wherein the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof. 288. The method of embodiment 287, wherein the functional fragment is at least at least 5 amino acid residues, at least 6 amino acid residues, at least 7 amino acid residues, at least 8 amino acid residues, at least 9 amino acid residues, at least 10 amino acid residues, at least 11 amino acid residues, at least 12 amino acid residues. 289. The method of any one of embodiments 287-288, wherein the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 365 – SEQ ID NO: 369. 290. The method of any one of embodiments 287-289, wherein the neurotensin has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at

least 99%, or 100% sequence identity to SEQ ID NO: 350. 291. The method of any one of embodiments 280-290, wherein the peptide construct is capable of binding and activating a neurotensin receptor. 292. The method of any one of embodiments 260-291, wherein the peptide is capable of reducing a level of pain upon administration to the subject. 293. The method of any one of embodiments 260-292, wherein the peptide is administered to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, pain, epilepsy, a mood disorder, or an endocrine disorder. 294. The method of any one of embodiments 260-293, wherein the peptide interacts with dopamine signaling neurons. 295. The method of any one of embodiments 260-294, wherein the peptide increases dopamine release in neurons. 296. The method of any one of embodiments 260-295, wherein the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety. 297. The method of any one of embodiments 259-296, wherein the composition is administered by inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof. 298. The method of any one of embodiments 259-297, wherein the composition is administered intravenously as a bolus, infusion, or prolonged infusion. 299. The method of any one of embodiments 259-298, wherein the composition or the pharmaceutical composition is administered multiple times a day, twice a day, once a day, twice a week, three times a week, once a week, once every two weeks, or once a month or once every three months. 300. The method of any one of embodiments 259-2969, wherein the composition is administered subcutaneously once a day, once a week, or once a month. 301. The method of any one of embodiments 259-300, wherein the condition is a cancer. 302. The method of embodiment 301, wherein the cancer is ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer. 303. The method of any one of embodiments 259-302, wherein the condition is a brain cancer. 304. The method of embodiment 303, wherein the brain cancer is

glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, or diffuse intrinsic pontine glioma. 305. The method of any one of embodiments 259-304, wherein the condition is a disease of the CNS. 306. The method of embodiment 305, wherein the disease of the CNS is a neuroinflammatory or neurodegenerative disease. 307. The method of embodiment 306, wherein the neurodegenerative disease is Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. 308. The method of any one of embodiments 259-302, wherein the condition is neuropathic pain, migraine, anxiety, or epilepsy. 309. The method of any one of embodiments 259-302 or 308, wherein the condition is chronic pain, acute pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, or breakthrough pain. 310. The method of any one of embodiments 259-302 or 308-309, wherein the condition is chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. 311. The method of any one of embodiments 259-310, wherein the condition is muscular dystrophy, or spinal bulbar muscular dystrophy (SBMA). 312. The method of any one of embodiments 259-311, wherein the condition is an inflammatory bowel disease. 313. The method of embodiment 312, wherein the inflammatory bowel disease is a Crohn's disease. 314. A method for transporting a composition according to any one of embodiments 1-313 across a cell layer, the method comprising contacting the cell layer with the composition, and transporting the composition across the cell layer. 315. The method of embodiment 314, wherein the cell layer is an endothelial cell layer or an epithelial cell layer. 316. The method of any one of embodiments 314-315, wherein the endothelial cell layer is the blood brain barrier. 317. The method of any one of embodiments 314-

315, wherein the endothelial cell layer is an intestinal endothelial cell layer. 318. The method of embodiment 317, wherein the epithelial cell layer is an intestinal epithelial cell layer. 319. The method of any one of embodiments 314-318, wherein the transporting comprises binding of the composition to TfR. 320. The method of any one of embodiments 314-319, wherein the transporting further comprises transcytosis. 321. The method of embodiment 320, wherein the transcytosis is vesicular transcytosis. 322. The method of any one of embodiments 320-321, wherein the transcytosis is TfR-mediated. 323. The method of any one of embodiments 319-319, wherein the transporting further comprises dissociation of the composition from the TfR. 324. The method of any one of embodiments 314-323, wherein the transporting further comprises releasing an active agent from the composition. 325. A method of treating a condition in a subject in need thereof, the method comprising administering to the subject the composition of any one of embodiments 1-163 or the pharmaceutical composition of any one of embodiments 164-324. 326. The method of embodiment 325, wherein the composition is administered by inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof. 327. The method of any one of embodiments 325-326, wherein the composition is administered intravenously as a bolus, infusion, or prolonged infusion. 328. The method of any one of embodiments 325-327, wherein the composition or the pharmaceutical composition is administered multiple times a day, twice a day, once a day, twice a week, three times a week, once a week, once every two weeks, or once a month or once every three months. 329. The method of any one of embodiments 325-328, wherein the composition is administered subcutaneously once a day, once a week, or once a month. 330. The method of any one of embodiments 325-329, wherein the condition is a cancer. 331. The method of embodiment 330, wherein the cancer is ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer. 332. The method of any one of embodiments 325-331, wherein the condition is a brain cancer. 333. The method of

embodiment 332, wherein the brain cancer is glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, or diffuse intrinsic pontine glioma. 334. The method of any one of embodiments 325-329, wherein the condition is a disease of the CNS. 335. The method of embodiment 334, wherein the condition of the CNS is a neuroinflammatory or neurodegenerative disease. 336. The method of embodiment 335, wherein the neurodegenerative disease is Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufor–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. 337. The method of any one of embodiments 325-329, wherein the condition is neuropathic pain, migraine, anxiety, or epilepsy. 338. The method of any one of embodiments 325-329, wherein the condition is chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. 339. The method of any one of embodiments 325-329, wherein the condition is muscular dystrophy, or spinal bulbar muscular dystrophy (SBMA). 340. The method of any one of embodiments 301-305, wherein the condition is an inflammatory bowel disease. 341. The method of embodiment 340, wherein the inflammatory bowel disease is a Crohn's disease. 342. The method of any one of embodiments 325-341, further comprising transporting the composition across a cell layer. 343. The method of embodiment 342, wherein the cell layer is an endothelial cell layer or an epithelial cell layer. 344. The method of any one of embodiments 342-343, wherein the endothelial cell layer is the blood brain barrier. 345. The method of any one of embodiments 342-344, wherein the endothelial cell layer is an intestinal endothelial cell layer. 346. The method of embodiment 340, wherein the epithelial cell layer is an intestinal epithelial cell layer. 347. The method of any one of embodiments 325-346, wherein the transporting comprises binding of the composition to TfR.

348. The method of any one of embodiments 325-347, wherein the transporting further comprises transcytosis. 349. The method of embodiment 348, wherein the transcytosis is vesicular transcytosis. 350. The method of any one of embodiments 348-349, wherein the transcytosis is TfR-mediated. 351. The method of any one of embodiments 347-350, wherein the transporting further comprises dissociation of the composition from the TfR. 352. The method of any one of embodiments 325-351, wherein the transporting further comprises releasing an active agent from the composition. 353. A method of treating a condition in a subject in need thereof, the method comprising administering to the subject an amount of the composition of any one of embodiments 164-253 or the pharmaceutical composition of any one of embodiments 161-163 that is dependent on the expression of TfR in a tissue of the subject. 354. The method of embodiment 353, wherein the composition is administered by inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof. 355. The method of any one of embodiments 353-354, wherein the composition is administered intravenously as a bolus, infusion, or prolonged infusion. 356. The method of any one of embodiments 353-355, wherein the composition or the pharmaceutical composition is administered multiple times a day, twice a day, once a day, twice a week, three times a week, once a week, once every two weeks, or once a month or once every three months. 357. The method of any one of embodiments 353-356, wherein the composition is administered subcutaneously once a day, once a week, or once a month. 358. The method of any one of embodiments 353-357, wherein the condition is a cancer. 359. The method of any one of embodiments 354-358, wherein the tissue of the subject is a cancerous tissue. 360. The method embodiment 358, wherein the cancer is ovarian cancer, colon cancer, lung cancer, cancer located in the bone or bone marrow, glioblastoma, astrocytoma, glioma, medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, diffuse intrinsic pontine glioma (DIPG), breast cancer, liver cancer, colon cancer, brain cancer, spleen cancer, cancers of the salivary gland, kidney cancer, muscle cancers, bone marrow cell cancers, skin cancer, genitourinary cancer, osteosarcoma, muscle-derived sarcoma, melanoma, head and neck cancer, neuroblastoma, prostate cancer, bladder cancer, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, or a CMYC-overexpressing cancer. 361. The method of any one of embodiments 353-360, wherein the condition is a brain cancer. 362. The method of embodiment 361, wherein the brain cancer is glioblastoma, astrocytoma, glioma,

medulloblastoma, ependymoma, choroid plexus carcinoma, midline glioma, or diffuse intrinsic pontine glioma. 363. The method of any one of embodiments 353-357, wherein the condition is a disease of the CNS. 364. The method of embodiment 363, wherein the disease of the CNS is a neuroinflammatory or neurodegenerative disease. 365. The method of embodiment 364, wherein the neurodegenerative disease is Alzheimer's disease, Amyotrophic lateral sclerosis, Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease, Spinal muscular atrophy, Motor neuron disease, Lyme disease, Ataxia-telangiectasia, Autosomal dominant cerebellar ataxia, Batten disease, Corticobasal syndrome, Creutzfeldt–Jakob disease, Fragile X-associated tremor/ataxia syndrome, Kufo–Rakeb syndrome, Machado–Joseph disease, multiple sclerosis, chronic traumatic encephalopathy, or frontotemporal dementia. 366. The method of any one of embodiments 353-365, wherein the condition is neuropathic pain, migraine, anxiety, or epilepsy. 367. The method of any one of embodiments 353-366, wherein the condition is chronic pain, acute pain, injury, mechanical pain, heat pain, cold pain, ischemic pain, and chemical-induced pain, inflammatory pain, migraine-related pain, headache-related pain, irritable bowel syndrome-related pain, fibromyalgia-related pain, arthritic pain, skeletal pain, joint pain, gastrointestinal pain, muscle pain, angina pain, facial pain, pelvic pain, claudication, postoperative pain, post traumatic pain, tension-type headache, obstetric pain, gynecological pain, or chemotherapy-induced pain, nociceptive pain, neuropathic pain, allodynia, phantom pain, visceral pain, breakthrough pain, familial episodic pain syndrome, paroxysmal extreme pain disorder, congenital indifference to pain, pain associated with opioid withdrawal. 368. The method of any one of embodiments 353-367, wherein the condition is muscular dystrophy, or spinal bulbar muscular dystrophy (SBMA). 369. The method of any one of embodiments 353-367, wherein the condition is an inflammatory bowel disease. 370. The method of embodiment 369, wherein the inflammatory bowel disease is a Crohn's disease. 371. The method of any one of embodiments 353-370, further comprising transporting the composition across a cell layer. 372. The method of embodiment 371, wherein the cell layer is an endothelial cell layer or an epithelial cell layer. 373. The method of embodiment 372, wherein the endothelial cell layer is the blood brain barrier. 374. The method of any one of embodiments 372-373, wherein the endothelial cell layer is an intestinal endothelial cell layer. 375. The method of any one of embodiments 372-373, wherein the epithelial cell layer is an intestinal epithelial cell layer. 376. The method of any one of embodiments 371-375, wherein the transporting comprises binding of the composition to TfR. 377. The method of any one of embodiments 371-376, wherein the transporting further

comprises transcytosis. 378. The method of embodiment 377, wherein the transcytosis is vesicular transcytosis. 379. The method of any one of embodiments 377-378, wherein the transcytosis is TfR-mediated. 380. The method of any one of embodiments 371-379, wherein the transporting further comprises dissociation of the composition from the TfR. 381. The method of any one of embodiments 371-380, wherein the transporting further comprises releasing an active agent from the composition. 382. A method for designing a TfR-binding peptide, the method comprising: a) using a computer program to identify a peptide from a library of peptides with optimized stability and folding, wherein each peptide of the library of peptides comprises at least three intramolecular disulfide bonds; b) using a computer program to superimpose amino acid residues of a binding patch from a peptide:TfR complex to determine a stable peptide that binds to TfR after grafting the binding patch onto the peptide; and c) grafting the binding patch onto the peptide to produce the stable peptide. 383. The method of embodiment 382, further comprising performing site saturation mutagenesis to obtain a mutated peptide with a higher binding affinity for TfR than the stable peptide. 384. The method of embodiment 382-383, wherein the peptide from a library of peptides is between 20 and 75 amino acid residues in length. 385. The composition of any one of embodiments 137-145, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof. 386. The method of any one of embodiments 192-200, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof. 387. The method of any one of embodiments 239-247, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof. 388. The method of any one of embodiments 287-295, wherein the neurotensin comprises any one of SEQ ID NO: 350, SEQ ID NO: 365 – SEQ ID NO: 371, or a functional fragment thereof. 389. The composition of any one of embodiments 136-163, wherein the Kv1.3 inhibitor is Vm24, ShK-170, ShK-186, or ShK-192. 390. The composition of embodiment 389, wherein Vm24 has a sequence of SEQ ID NO: 378 or SEQ ID NO: 391. 391. The composition of embodiment 389, wherein Shk-185 has a sequence of SEQ ID NO: 379, SEQ ID NO: 390, or SEQ ID NO: 402. 392. The composition of any one of embodiments 148-163, wherein the half-life modifying agent comprises SA21. 393. The composition of embodiment, 392, wherein the SA21 has a sequence of SEQ ID NO: 376 or SEQ ID NO: 377. 394. The method of any one of embodiments 191-210, 238-258, or 286-313, wherein the Kv1.3 inhibitor is Vm24, ShK-170, ShK-186, or ShK-192. 395. The method of embodiment 394, wherein Vm24

comprises SEQ ID NO: 378 or SEQ ID NO: 391. 396. The method of embodiment 394, wherein Shk-185 comprises SEQ ID NO: 379, SEQ ID NO: 390, or SEQ ID NO: 402. 397. The method of any one of embodiments 165-210, 213-258, or 261-381, further comprising a half-life modifying agent coupled to the peptide. 398. The method of embodiment 397, wherein the half-life modifying agent comprises a polymer, a polyethylene glycol (PEG), a hydroxyethyl starch, polyvinyl alcohol, a water soluble polymer, a zwitterionic water soluble polymer, a water soluble poly(amino acid), a water soluble polymer of proline, alanine and serine, a water soluble polymer containing glycine, glutamic acid, and serine, an Fc region, a fatty acid, palmitic acid, or a molecule that binds to albumin. 399. The method of embodiment 397, wherein the half-life modifying agent comprises SA21. 400. The method of embodiment, 399, wherein the SA21 comprises SEQ ID NO: 376 or SEQ ID NO: 377. 401. The composition of any one of embodiments 1-163, 385, or 389-393, wherein the peptide has a wider therapeutic window than a TfR-binding antibody-based therapeutic agent. 402. The composition of embodiment 401, wherein the wider therapeutic window is a dosage above which a therapeutic pharmacodynamic response is observed but below which toxicity is observed. 403. The composition of any one of embodiments 1-163, 385, 389-393, 401, or 402, wherein the peptide has a lower on-target toxicity, a lower off-target toxicity, or a combination thereof than a TfR-binding antibody-based therapeutic agent when administered to a subject at the same molar dose. 404. The method of any one of embodiments 165-210, 212-258, 260-384, 386-388, or 394-400, wherein the peptide has a wider therapeutic window than a TfR-binding antibody-based therapeutic agent. 405. The method of embodiment 404, wherein the wider therapeutic window is a dosage above which a therapeutic pharmacodynamic response is observed but below which toxicity is observed. 406. The method of any one of embodiments 165-210, 212-258, 260-384, 386-388, 394-400, 404, or 405, wherein the peptide has a lower on-target toxicity or a lower off-target toxicity than a TfR-binding antibody-based therapeutic agent when administered to a subject at the same molar dose.

EXAMPLES

[0536] The following examples are included to further describe some aspects of the present disclosure, and should not be used to limit the scope of the invention.

EXAMPLE 1

Manufacture of Peptides

[0537] This example describes the manufacture of the peptides and peptide constructs described herein (e.g., any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361). Peptides derived from proteins were generated in mammalian cell culture using a published methodology. (A.D. Bandaranayake, C. Correnti, B.Y. Ryu, M. Brault, R.K. Strong, D. Rawlings. 2011. Daedalus: a robust, turnkey platform for rapid production of decigram quantities of active recombinant proteins in human cell lines using novel lentiviral vectors. *Nucleic Acids Research*. (39)21, e143).

[0538] The peptide sequence was reverse-translated into DNA, synthesized, and cloned in-frame with siderocalin using standard molecular biology techniques (M.R. Green, Joseph Sambrook. *Molecular Cloning*. 2012 Cold Spring Harbor Press). The resulting construct was packaged into a lentivirus, transduced into HEK-293 cells, expanded, isolated by immobilized metal affinity chromatography (IMAC), cleaved with tobacco etch virus (TEV) protease, and purified to homogeneity by reverse-phase chromatography. Following purification, each peptide was lyophilized and stored frozen.

EXAMPLE 2

Peptide Expression Using a Mammalian Expression System

[0539] This example describes expression of the peptides and peptide constructs using a mammalian expression system. Peptides were expressed according to the methods described in in Bandaranayake et al., *Nucleic Acids Res*. 2011 Nov; 39(21): e143. Peptides (e.g., any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361) were cleaved from siderocalin using tobacco etch virus protease and were purified by reversed-phase HPLC (RP-HPLC) in a gradient of acetonitrile and water with 0.1% TFA, and then aliquoted and lyophilized for later use. Molecular weight was verified by mass spectrometry.

[0540] To optimize and validate the screening methodology and to identify TfR-binding peptides, transferrin receptor (TfR) ectodomain (“soluble TfR”, SEQ ID NO: 353) was cloned into the Daedalus soluble protein production lentivector, and protein was purified from the growth media (a gel of soluble TfR is shown in **FIG. 1A**). The same strategy was used to

produce and purify human apo-transferrin (residues 23-698, SEQ ID NO: 364, KTVRWCAVSEHEATKCQSF RDHMKSVIPSDGPSVACVKKASYLDCIRAI AANEADAVTL DAGLVYDAYLAPNNLKPVVAEFYGS KEDPQTFYYAVAVVKKDSGFQMNQLRGKKSCH TGLGRSAGWNIPIGLLYCDLPEPRKPLEKAVANFFSGSCAPCADGTDFPQLCQLCPGCGC STLNQYFGYSGAFKCLKDGAGDVAFVKHSTIFENLANKADRDQYELLCLDNTRKPVDE YKDCHLAQVPSHTVVARS MGGKEDLIWELLNQAQEHFGKDKSKEFQLFSSPHGKDLLF KDSA HGFLKVPPRMDAKMYLGYEYVTAIRNLREGTCPEAPTDECKPVKWCALSHHERL KCDEWSVNSVGKIECVSAETTEDCIAKIMNGEADAMSLDGGFVYIAGKCGLVPVLAEN YNKSDNCEDTPEAGYFAIAVVKKSASDLTWDNLKGKKSCHTAVGRTAGWNIPMGLLY NKINHCRFDEFFSEGCAPGSKKDSSLCKLCMGSGNLNCEPNNKEGYGYTGAFRCLVEK GDVAFVKHQTV PQNTGGKNPDPWAKNLNEKDYELLCLDGTRKPV E EYANCHLARAPN HAVVTRKDKEACVHKILRQQQHLFGSNVTDCSGNFCLFRSETKDLLFRDDTVCLAKLH DRNTYEKYLGE EYVKAVGNLRK CSTSSLLEACTFRRP), and some of the material was iron-loaded to produce holo-transferrin. Both apo-transferrin and holo-transferrin were tested for binding to the soluble TfR ectodomain via surface plasmon resonance (**FIG. 23A**), where only holo-transferrin demonstrated any interaction with the immobilized TfR.

[0541] As further validation that the soluble TfR used in the screen represents the human, endogenous protein structure, interaction with the Machupo virus glycoprotein was tested (referred to as MaCV), which uses TfR to determine tropism and mediate cell entry. For this, MaCV was cloned into a mammalian surface display vector SDGF (**FIG. 23B**), and transfected suspension 293 Freestyle (293F) cells with SDGF-MaCV or a control protein (SDGF-Elafin, an inhibitor of elastase known to bind some native CDPs). Transfected cells were stained with 200 nM each biotinylated TfR (all TfR used in cell binding assays is biotinylated) and Alexa Fluor 647-labeled streptavidin, and then analyzed by flow cytometry (**FIG. 23C**). Cells transfected with MaCV were successfully stained with TfR, while SDGF-Elafin cells were not. Meanwhile, SDGF-MaCV cells incubated with 200 nM fluorescent elastase were not stained. This demonstrates both the specificity of TfR binding to its endogenous ligand, and the utility of SDGF as a means to identify novel TfR binding partners.

EXAMPLE 3

Expression and Purification of Functional Transferrin Receptor

[0542] Transferrin receptor (TfR) ectodomain (SEQ ID NO: 352, residues 121-760 of HsTfR by UniProt numbering), apo-transferrin (SEQ ID NO: 353 (human) and SEQ ID NO: 354 (mouse)), and all soluble CDPs were produced and purified as **EXAMPLE 2** and **EXAMPLE 3**. In brief, relevant sequences were cloned into the Daedalus vector, a lentivector driving secretion of proteins of interest. Proteins either contained their endogenous signal peptide or were fused at their N-terminus to siderocalin for stabilization during translation / secretion; siderocalin fusions were cleaved using TEV protease, leaving an N-terminal “GS” behind. VSV-G pseudotyped lentivirus was produced through standard methods in 293 cells. This virus was used to transduce 1×10^7 suspension 293 Freestyle (293F, ThermoFisher R79007) cells at an approximate multiplicity of infection (MOI) of 10 in FreeStyle 293 Expression Medium (Invitrogen 12338018). The culture was then grown (suspension culture incubator at 125 RPM, 37°C, 80% humidity, 8% CO₂) and expanded over the course of 10-12 days (2 L final volume), after which media was collected and sterile filtered. Proteins were purified from media by batch Ni-NTA resin binding. CDPs were separated from the siderocalin carrier by TEV protease cleavage, and a final polish with either SEC (ÄKTA Pure, GE Healthcare) or RP-HPLC (Agilent model 1260, C18 column, 214 nm absorbance, with an in-line Agilent 6120 LC/MS) was performed (**FIG. 1A**). TfR and transferrin were frozen in PBS + 5% glycerol, while the CDP peptides, purified by RP-HPLC, were aliquoted and lyophilized in vials. Peptide quantitation was established by amino acid analysis before use *in vivo*.

[0543] Proper folding and function of purified TfR was verified using surface plasmon resonance. Binding of either apo-transferrin (ApoTf) or holo-transferrin (HoloTf) to the purified TfR was measured (**FIG. 23A**). Preferential binding of HoloTf over ApoTf is indicative of proper TfR function.

[0544] All proteins were analyzed by SDS-PAGE (NuPAGE 4-12% Bis-Tris, Thermo Fisher) and Coomassie staining (**FIG. 4 and FIG. 27**). CDPs were run both non-reduced and reduced with 10 mM DTT. Proteolysis and reduction resistance testing was performed on the same RP-HPLC instrumentation as used for purification, after treatment with either 50 U of porcine pepsin (Sigma-Aldrich P7012) in simulated gastric fluid or 50 U of porcine trypsin (Sigma-Aldrich 6567) in PBS followed by incubation for 30 min at 37°C. Similarly, glutathione or DTT reduction involved treatment of the peptides with 10 mM glutathione or DTT in PBS for > 5 min

prior to RP-HPLC analysis. CD spectra of a peptide of SEQ ID NO: 32 were measured with a Jasco J-720W spectropolarimeter (1 mm path length) in 20 mM phosphate buffer, pH 7.4, with or without 10 mM DTT at a concentration of 15-25 μ M. Data are expressed in terms of relative ellipticity $[\theta]$ change at 210 nm, reported in mdeg.

[0545] Holo-transferrin was produced according to the method of Hamilton et al. In brief, a 10 mL solution of 4.5 mM Fe(NTA)₂, containing 4.5 mM FeCl₃ (Sigma Aldrich) and 9 mM nitrolotriactic acid (NTA, Sigma Aldrich) in water, was titrated to pH 4.0 with HCl and NaOH. Apo-transferrin solution in PBS was loaded with Fe(NTA)₂, 0.4 equivalents at a time every 5 minutes, until 3.2 equivalents had been added. The solution was buffer exchanged into PBS using spin column dialysis (Amicon Ultra-4 centrifugal filter units, 10 kDa molecular weight cutoff) for >1,000-fold free Fe(NTA)₂ clearance.

[0546] Alexa Fluor 647 (Thermo Fisher A37573) CDP labeling took place in PBS buffer with a molar ratio of ~1.5 NHS-ester dye : 1 CDP for > 1 hour, after which free dye was removed by spin column dialysis (Amicon Ultra-4 centrifugal filter units, 3 kDa molecular weight cutoff) for >1,000-fold free dye clearance. Biotinylation of Avitagged TfR used the BirA-500 kit (Avidity) as per manufacturer's protocol, followed by a final buffer exchange into PBS containing 5% glycerol, and storage in small aliquots at -80°C.

EXAMPLE 4

Mammalian Surface Display of TfR-binding Peptides

[0547] This example describes mammalian surface display of TfR-binding peptides of the present disclosure including SEQ ID NO: 1 (SEQ ID NO: 1 is SEQ ID NO: 36 with an added N-terminal GS), SEQ ID NO: 2 (SEQ ID NO: 2 is SEQ ID NO: 37 with an added N-terminal GS), SEQ ID NO: 30 (SEQ ID NO: 30 is SEQ ID NO: 65 with an added N-terminal GS), and SEQ ID NO: 32 (SEQ ID NO: 32 is SEQ ID NO: 67 with an added N-terminal GS), and SEQ ID NO: 33 – SEQ ID NO: 35 (SEQ ID NO: 33 – SEQ ID NO: 35 are SEQ ID NO: 68 – SEQ ID NO: 70, respectively, with an added N-terminal GS). Screening for TfR-binding peptides was performed by transfecting or transducing mammalian cells to display candidate peptides (**FIG. 23B**) followed by screening against soluble human transferrin receptor ectodomain SEQ ID NO: 352 (200 nM, **FIG. 23C**, **FIG. 1**). Mammalian cells had improved fidelity in folding disulfide crosslinked proteins, making them a suitable cell type for display of the peptides of the present disclosure.

[0548] Mammalian cell surface display screening was carried out as follows. The screening strategy used a surface display GFP FasL (SDGF) vector. In the vector, FasL-TM is the transmembrane domain of the FasL protein. More specifically, the designed peptides were cloned as a pool into SDGF, which were then made into lentivirus. 293F cells were transduced with this library at a multiplicity of infection of ~1, and after three days of growth, the pool of transduced cells are incubated with Alexa647-labeled TfR. For these experiments, fluorescent labeling was accomplished by co-staining TfR with fluorescent streptavidin or fluorescent anti-His antibodies. Soluble TfR contained both His tag and biotin label. Either Alexa Fluor 647 or iFluor 647 was used for the antibody/streptavidin fluorescence. A percentage of the highest-staining TfR-positive cells, from GFP and TfR double-positive cells, were sorted and expanded. At every expansion, a portion of the cells were collected, and at the end, the enriched peptides were identified by sequencing. Flow cytometry was used to assess gating criteria to identify GFP+ 293F cells expressing proteins on their surface via the SDGF peptide construct. Gating progresses using FSC-H vs SSC-H to gate out debris; FSC-H vs FSC-A to gate out doublets; FSC-H vs Pacific Blue H to gate out DAPI+ dead cells; and an optional FITC-H histogram to identify GFP+ cells. Once gated as such, Alexa Fluor 647 (a co-stain for detecting target binding) was used for sorting and analysis.

[0549] Screening was performed using a combination of magnetic sorting and flow sorting. Magnetic sorting was performed as follows: 2×10^8 293F cells were transduced with the SDGF CDP library at an MOI of ~1, and expanded until 3 days post-transduction. For initial screening, magnetic cell sorting was performed. 1×10^9 transduced cells were resuspended in a binding buffer containing 200 nM biotinylated TfR, 2 mL anti-biotin MicroBeads (UltraPure, Miltenyi 130-105-637), and 21 mL Flow Buffer (PBS + 2 mM EDTA and 0.5% bovine serum albumin) in a final volume of 25 mL. Cells were incubated on ice with agitation (mild inversion every 2-5 min) for 30 mins, and were then diluted 10-fold to 250 mL with Flow Buffer, pelleted (500 x g, 5 mins), and resuspended to 40 mL with High BSA Flow Buffer (PBS containing 2 mM EDTA and 3% BSA). Cells were split into four 10 mL aliquots and run through a Miltenyi autoMACS® Pro Separator using the “posseld” protocol and “quick rinses” after each sort. The running and wash buffers were High BSA Flow Buffer and PBS + 2 mM EDTA, respectively. Eluted cells were pooled, pelleted, and their CDP sequences PCR amplified (Terra™ PCR Direct Polymerase Mix (Takara 639271) for 16 cycles followed by Phusion). This sub-library was cloned into SDGF as

above, made into lentivirus, and transduced into a new batch of 293F cells (1×10^7 cells, MOI ~ 1) for flow sorting.

[0550] Flow sorting was performed as follows: Flow sorting took place using 2.4×10^7 cells stained in 3 mL Flow Buffer with 200 nM TfR, 200 nM streptavidin Alexa Fluor 647 conjugate (ThermoFisher S21374), and $1 \mu\text{g mL}^{-1}$ DAPI. Cells were diluted 4-fold to 12 mL with Flow Buffer, pelleted ($500 \times g$, 5 mins), and resuspended in 3.6 mL Flow Buffer. Cells were sorted on a FACSAria II System (BD), gating based on FSC-A (medium), SSC-A (medium), DAPI-A (negative), GFP-A (positive), and APC-A (top 7% of GFP+) channels. After each flow sort, cells were cultured in FreeStyle Media starting at 0.5-1 mL in a suspension 24-well plate, shaking at 300 rpm, expanding to a final volume of 30 mL in a 125 mL baffled flask shaken at 125 RPM. At this point, cells were re-sorted as above. After the third flow sort, cells were expanded and frozen in 1.5×10^6 cell pellets. Pellets were PCR amplified as above (Terra Direct PCR followed by InFusion), CDP inserts were subcloned into SDGF, transformed (Stellar competent cells), and colonies picked for miniprepping and sequencing of the cloned CDPs. Enriched variants were those that appeared in the sequence analysis more than once in ~ 100 picked colonies. All site-saturation mutagenesis (SSM) affinity maturation screening was done as above, with the following changes. 1) Staining was sequential; first with TfR, and then with an equimolar amount of dye-labeled streptavidin. 2) TfR / streptavidin concentrations were reduced to 20 nM for the first SSM maturation and to 8 nM for the second SSM maturation screen. After each SSM screen, enriched variants were studied to assemble a compound mutant (peptide of SEQ ID NO: 2 or SEQ ID NO: 32) that showed higher TfR staining than any of the variants containing either 1 or 2 of the individual mutations.

[0551] Flow cytometry plots in **FIG. 1** illustrate successive enrichment of cells that bind to TfR from pooled, high diversity library. **FIG. 1A** illustrates a Coomassie stained gel of transferrin receptor (TfR) protein. **FIG. 1B** illustrates a flow cytometry plot showing cells that bind to TfR after one flow sort. **FIG. 1C** illustrates a flow cytometry plot showing cells of a control stain after one flow sort. **FIG. 1D** illustrates a flow cytometry plot showing cells that bind to TfR after two flow sorts. **FIG. 1E** illustrates a flow cytometry plot showing cells of a control stain after two flow sorts. **FIG. 1F** illustrates a flow cytometry plot showing cells that bind to TfR after three flow sorts. **FIG. 1G** illustrates a flow cytometry plot showing cells of a control stain after three flow sorts.

[0552] Each flow sort represents growing the library of cells to >30 million cells, staining for TfR-binding, and flow sorting the top binders. Sorted binders were allowed to grow before the next flow sort was performed.

EXAMPLE 5

Identification of TfR-binding Peptides

[0553] This example describes identification of TfR-binding peptides using a mammalian surface display system, as described in **EXAMPLE 4**.

[0554] Using the mammalian surface display system of **EXAMPLE 4**, a single clonal peptide was identified having a sequence of SEQ ID NO: 1 (SEQ ID NO: 1 is SEQ ID NO: 36 with an added N-terminal GS). A library of oligonucleotides encoding 10,000 CDPs was amplified and mutagenized. The CDPs were 17-50 amino acids in length, with 4, 6, 8, or 10 cysteines. While there was some weighting of the library towards annotated knottins or defensins, the library contained CDPs from every domain / kingdom of life. This library was cloned into SDGF, made into lentivirus, and transduced into suspension 293F cells. The transduced cells were subjected to staining with TfR (200 nM) and co-stain over the course of one round of magnetic cell sorting and three rounds of flow sorting, each round enriching for cells stained with TfR. Binding was validated to specifically bind TfR in the surface display assay using 200 nM of soluble AF647-TfR, which was either biotinylated or attached to a His tag. Staining was carried out using a one-step staining protocol for tetravalent target avidity. A single TfR-binding CDP, designated SEQ ID NO: 1, was identified by DNA sequencing of the final enriched cell population. It represents a randomly mutated variant of cytochrome BC1 complex subunit 6 from the marine choanoflagellate *Monosiga brevicolis* (Uniprot ID: A9V0D7, DOI: 10.1093/nar/gku989), is 49 amino acids in length (six cysteines), and has a predicted molecular mass of 5.6 kDa. SEQ ID NO: 36 was then subjected to affinity maturation using site saturation mutagenesis (SSM), wherein a library is created containing every possible non-cysteine single amino acid substitution (43 non-Cys amino acids x 18 possible non-Cys substitutions = 775 variants, including SEQ ID NO: 1).

[0555] Flow cytometry plots in **FIG. 2** illustrate cells displaying TfR and the single clonal peptide. **FIG. 2A** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and a control protein (y-axis, stained with a fluorescent anti-His antibody). **FIG. 2B** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP)

and a control protein (y-axis, stained with a fluorescent streptavidin). **FIG. 2C** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and TfR (y-axis, stained with a fluorescent anti-His antibody). **FIG. 2D** illustrates a flow cytometry plot of cells expressing a peptide of SEQ ID NO: 1 (x-axis, GFP) and a TfR (y-axis, stained with a fluorescent streptavidin).

[0556] TfR staining was observed with cells expressing the identified clone, with no staining seen with a control protein. This staining was observed when either fluorescent streptavidin or anti-His antibody was the co-stain, demonstrating that the nature of the binding is dependent only on TfR, not the co-stain. Double-positive cells (upper right hand quadrant) indicate peptide-expressing cells that are bound to TfR.

[0557] An alternative method was also used to identify and optimize (“mature”) TfR-binding peptides that are second and third generation binders using mammalian display screening. This library was screened with a modified staining protocol, using a lower concentration of target and co-stain (20 nM) and separate staining steps; the latter further increases stringency by eliminating the tetravalent avidity granted by streptavidin. Up to four rounds of flow sorting and enrichment were used to identify variants with improved TfR binding characteristics. Permutation of enriched variants identified an optimal mutant (SEQ ID NO: 2 (SEQ ID NO: 2 is SEQ ID NO: 37 with an added N-terminal GS)), and this process was repeated again (8 nM TfR and co-stain; otherwise, identical protocol) to generate SEQ ID NO: 32 (SEQ ID NO: 32 is SEQ ID NO: 67 with an added N-terminal GS). This twice-matured variant contains 14 point mutations from the original library member

(GSREGCASHCTKYKAELEKCEARVSSRSNTEETCVQELFDLHCVDHCVSQ, SEQ ID NO: 362). 4 point mutations from the parent sequence are found in SEQ ID NO: 1, while SEQ ID NO: 2 and SEQ ID NO: 32 contain 6 and 4 mutations, respectively, from the previous generation.

[0558] SEQ ID NO: 1 and its variants (e.g., SEQ ID NO: 2 and SEQ ID NO: 32) were produced as soluble peptides, and validated by reversed-phase HPLC (RP-HPLC), SDS-PAGE, and mass spectrometry (**FIG. 4A**, **FIG. 4C**, and **FIG. 4E**). Based on their masses of ~5-6 kDa, their slower-than-expected mobility in SDS-PAGE was a demonstration of the interesting electrophoretic mobility characteristics that some CDPs possess. All variants showed markedly different mobility upon DTT reduction (10 mM) in both SDS-PAGE and RP-HPLC, confirming disulfide bond stabilization. Their binding to TfR was verified by surface plasmon resonance

(**FIG. 7**), also confirming increased affinity of matured variants (SEQ ID NO: 32 [$K_D = 216 \pm 1$ pM] > SEQ ID NO: 2 [$K_D = 8.7 \pm 0.4$ nM] > SEQ ID NO: 1 [K_D not determined]). Peptides were further evaluated for stability under conditions of reduction, protease exposure, and heat (**FIG. 5**, **FIG. 6**, and **FIG. 26**). All variants demonstrated full or partial resistance to cellular reducing conditions (10 mM glutathione), while the affinity matured variants showed partial resistance to pepsin (though all were vulnerable to trypsin proteolysis). The intact, non-reduced peptide of SEQ ID NO: 32 protein demonstrates substantially improved heat tolerance to that of the DTT-reduced protein, with no substantial change in circular dichroism characteristics until well above common ambient temperatures (>50°C) and a failure to observe complete unfolding up to 95°C.

EXAMPLE 6

Site Saturation Mutagenesis of Peptides

[0559] This example illustrates site saturation mutagenesis (SSM) of peptides of this disclosure to identify beneficial or deleterious mutations. Site saturation mutagenesis was performed on a peptide of SEQ ID NO: 1 (SEQ ID NO: 1 is SEQ ID NO: 36 with an added N-terminal GS, results in **FIG. 3A**) and a peptide of SEQ ID NO: 2 (SEQ ID NO: 2 is SEQ ID NO: 37 with an added N-terminal GS, results in **FIG. 3B**).

[0560] **FIG. 3** shows the TfR-binding capabilities of TfR-binding peptide variants identified during peptide maturation. SSM was employed for affinity maturation of the peptide having a sequence of SEQ ID NO: 1 as identified in the first mammalian surface display experiments (see e.g., **FIG. 1** and **FIG. 2**). During each round of maturation, a library of every possible non-cysteine point variants was constructed, and screened against TfR at higher stringency than first screen. Variants with improved binding were enriched and identified by Sanger sequencing. Such enriched variant mutations were combined with one another in various permutations (shown in the data) to identify a composite, improved binder. Two rounds of SSM were completed, yielding matured peptides comprising SEQ ID NO: 2, and SEQ ID NO: 32, respectively (SEQ ID NO: 32 is SEQ ID NO: 67 with an added N-terminal GS). TfR concentration in the first round of SSM was 20 nM and SSM was carried out using one-step staining. TfR concentration in the second round of SSM was 8 nM and SSM was carried out using two-step staining.

[0561] TfR-binding of mammalian cells expressing peptides of the SSM library was performed as described in **EXAMPLE 4** and **EXAMPLE 5**, but with a higher stringency protocol. The higher stringency protocol included a lower concentration of TfR (e.g., 20 nM).

[0562] **FIG. 3A** illustrates the results of a first site-saturation mutagenesis screen in SEQ ID NO: 1 (SEQ ID NO: 1 is SEQ ID NO: 36 with an added N-terminal GS), with some variants exhibiting improved binding activity to TfR such as peptides having a sequence of SEQ ID NO: 4, SEQ ID NO: 5, and SEQ ID NO: 8 (SEQ ID NO: 4, SEQ ID NO: 5, and SEQ ID NO: 8 are SEQ ID NO: 39, SEQ ID NO: 40, and SEQ ID NO: 43, respectively, with an added N-terminal GS). **FIG. 3B** show TfR-binding of variants identified during the second variant mutation.

[0563] The x-axis shows SEQ ID NOs of all variants and the y-axis shows the amount TfR bound in relative fluorescence units (RFUs) extrapolated from flow cytometry experiments.

EXAMPLE 7

TfR-binding of SSM-Generated TfR-binding Peptide Variants

[0564] This example demonstrates TfR-binding of site saturation mutagenesis (SSM)-generated TfR-binding peptide variants, as identified during SSM as described in **EXAMPLE 6**

[0565] In vivo BBB penetration experiments revealed that the TfR-binding capability of a peptide does not necessarily correspond with the capability of promoting vesicular transcytosis.

[0566] The six cysteines corresponding to residues C6, C10, C20, C34, C44, and C48, with reference to SEQ ID NO: 32 (C4, C8, C18, C32, C42, and C46, with reference to SEQ ID NO: 67), participate in disulfides, and thus contribute to peptide stability.

[0567] The surface interface residues corresponding to residues G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, Q51, with reference to SEQ ID NO: 32 (G3, A5, S6, N12, L15, E16, E19, L36, L40, L43, D44, H45, S48, Q49, with reference to SEQ ID NO: 67), that are present in all three generations of TfR-binding peptides likely contribute to TfR-binding. In some embodiments, the peptide or peptide construct of the present disclosure comprises at least one or more of these corresponding residues in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361. Such peptides can accordingly be engineered with enhanced binding to TfR.

[0568] Hydrophilic surface-distal residues such as D, E, H, K, R, N, Q, S, or T likely contribute to peptide solubility corresponding to the following amino acid residues R3, E4, R9, K12, D14, E15, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, with reference to SEQ ID NO: 32 (R1, E3, R7, K10, D12, E13, K17, R21, S24, S26, N27, T28, E29, E30, D31, E33, Q34, E35, E37, and D38, with reference to SEQ ID NO: 67). In some embodiments, the peptide or peptide construct of the present disclosure comprises at least one or

more of these corresponding residues in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361. Such peptides can accordingly be engineered with enhanced solubility.

[0569] Higher binding affinity is associated with the presence of hydrophilic residues such as D, E, H, K, R, N, Q, S, or T as shown by improved binding from a mutation away from a nonpolar or hydrophobic residue such as A, M, I, L, V, F, W, or Y at the residues corresponding to D15, E35, E39, and H49, with reference to SEQ ID NO: 32 (D13, E33, E37, and H47, with reference to SEQ ID NO: 67). In some embodiments, the peptide or peptide construct of the present disclosure comprises at least one or more of these corresponding residues in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361. Such peptides can accordingly be engineered with modified binding affinity.

[0570] Higher binding affinity to TfR is associated with nonpolar or hydrophobic residues such as A, M, I, L, V, F, W, or Y as shown by improved binding from a mutation away from a hydrophilic residue such as D, E, H, K, R, N, Q, S, or T at the amino acid residues corresponding to M11, M25, and M27, with reference to SEQ ID NO: 32 (M9, M23, and M25, with reference to SEQ ID NO: 67). In some embodiments, the peptide or peptide construct of the present disclosure comprises at least one or more of these corresponding residues in SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361. Such peptides can accordingly be engineered with modified binding affinity.

[0571] A higher TfR-binding affinity is associated with aliphatic residues such as A, M, I, L, or V as shown by improved binding from a mutation away from a large, aromatic residues such as F, W, or Y at the amino acid residue corresponding to L45 with reference to SEQ ID NO: 32 (L43 with reference to SEQ ID NO: 67). Substitutions of any one or more F, W, or Y in a peptide of the present disclosure to an aliphatic residue comprising A, M, I, L, or V can be used to enhance the binding affinity of the peptide to TfR.

[0572] Any of peptides or peptide constructs of the present disclosure (e.g., any one of SEQ ID NO: 1 – SEQ ID NO: 70, SEQ ID NO: 136 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, or SEQ ID NO: 356 – SEQ ID NO: 361) can be modified at one or more of the corresponding residues described herein, to generate peptide variants with improved properties including enhanced stability and increased (or decreased) binding properties or modified TfR-

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 239

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 239

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

CLAIMS

WHAT IS CLAIMED IS:

1. A composition comprising an engineered transferrin receptor (TfR)-binding peptide, or a variant, homolog, fragment, or analog thereof.
2. A composition comprising a peptide construct, wherein the peptide construct comprises:
 - a) a transferrin receptor (TfR)-binding peptide, or a variant, homolog, fragment, or analog thereof; and
 - b) an active agent,wherein the peptide is conjugated to, linked to, or fused to the active agent.
3. A composition comprising an engineered peptide having at least 75%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, or SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a functional fragment thereof.
4. The composition of any one of claims 1-3, wherein the peptide penetrates a cellular layer comprising a blood brain barrier (BBB).
5. The composition of any one of claims 1-3, wherein the peptide penetrates a membrane and wherein the membrane is the membrane of a cell.
6. The composition of claim 5, wherein the cell expresses a TfR.
7. The composition of claim 5, wherein the peptide penetrates the membrane of the cell via TfR-mediated endocytosis.
8. The composition of any one of claims 1-3, wherein the peptide is delivered across the blood-brain barrier to the CNS by binding and then dissociating from TfR.
9. The composition of any one of claims 1-3, wherein the peptide is delivered across the blood-brain barrier by receptor-mediated transcytosis.
10. The composition of claim 5, wherein the cell is a tumor cell.
11. The composition of any one of claims 1-3, wherein a cell or an organ in which the peptide localizes overexpresses a TfR.
12. The composition of any one of claims 1-3, wherein the peptide comprises SEQ ID NO: 206 – SEQ ID NO: 224 and SEQ ID NO: 334 – SEQ ID NO: 344.

13. The composition of any one of claims 1-3, wherein the peptide comprises a surface interface residue corresponding to G5, A7, S8, N14, L17, E18, E21, L38, L42, L45, D46, H47, S50, and Q51, or any combination thereof, with reference to SEQ ID NO: 32.

14. The composition of any one of claims 1-3, wherein the peptide comprises hydrophilic surface-distal residues corresponding to any one of the positions 3, 4, 9, 11, 15, 16, 19, 23, 26, 28, 29, 30, 31, 32, 33, 35, 36, 37, 39, and 40, or any combination thereof, with reference to SEQ ID NO: 32.

15. The composition of claim 14, wherein the hydrophilic surface-distal residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T.

16. The composition of claim 12, wherein the peptide comprises any one of the hydrophilic surface-distal residues corresponding to R3, E4, R9, K12, D15, E16, K19, R23, S26, S28, N29, T30, E31, E32, D33, E35, Q36, E37, E39, and D40, or any combination thereof, with reference to SEQ ID NO: 32.

17. The composition of any one of claims 1-3, wherein the peptide comprises hydrophilic residues at a position corresponding to any one of 15, 35, 39, or 49, or any combination thereof, with reference to SEQ ID NO: 32.

18. The composition of claim 14, wherein the hydrophilic residues are selected from the amino acids D, E, H, K, R, N, Q, S, or T.

19. The composition of any one of claims 17-18, wherein the peptide comprises a hydrophilic residue corresponding to D15, E35, E39, and H49, or any combination thereof, with reference to SEQ ID NO: 32.

20. The composition of any one of claims 1-3, wherein the peptide comprises hydrophobic residues at a position corresponding to 11, 25, or 27, or any combination thereof, with reference to SEQ ID NO: 32.

21. The composition of claim 20, wherein the hydrophobic residues are selected from the amino acids A, M, I, L, V, F, W, or Y, or any combination thereof.

22. The composition of claim 20, wherein the peptide comprises any one of the hydrophobic residues corresponding to M11, M25, M27, or any combination thereof, with reference to SEQ ID NO: 32.

23. The composition of any one of claims 1-3, wherein the peptide comprises an aliphatic residue corresponding to position 45, with reference to SEQ ID NO: 32.

24. The composition of claim 23, wherein the aliphatic residue is selected from the amino acids A, M, I, L, or V.

25. The composition of claim 23, wherein the peptide comprises an aliphatic residue corresponding to L45, with reference to SEQ ID NO: 32.

26. The composition of any one of claims 1-3, wherein the peptide comprises at least one disulfide bond, at least two disulfide bonds, at least three disulfide bonds, or at least five disulfide bonds.

27. The composition of any one of claims 1-3, wherein the peptide comprises a sequence that has at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or a fragment thereof.

28. The composition of any one of claims 1-3, wherein the peptide comprises a sequence of any one of SEQ ID NO: 1 – SEQ ID NO: 32, SEQ ID NO: 36 – SEQ ID NO: 67, SEQ ID NO: 136 – SEQ ID NO: 167, SEQ ID NO: 171 – SEQ ID NO: 202, SEQ ID NO: 356 – SEQ ID NO: 358, or a variant or fragment thereof.

29. The composition of any one of claims 1-3, wherein the peptide comprises the sequence set forth in SEQ ID NO: 1.

30. The composition of any one of claims 1-3, wherein the peptide comprises the sequence set forth in SEQ ID NO: 2.

31. The composition of any one of claims 1-3, wherein the peptide comprises the sequence set forth in SEQ ID NO: 4.

32. The composition of any one of claims 1-3, wherein the peptide comprises the sequence set forth in SEQ ID NO: 30.

33. The composition of any one of claims 1-3, wherein the peptide comprises the sequence set forth in SEQ ID NO: 32.

34. The composition of any one of claims 1-3, wherein the peptide is further formulated with, fused to, or conjugated to a cell-penetrating moiety.

35. The composition of claim 34, wherein the cell-penetrating moiety comprises polycations, polyorganic acids, endosomal releasing polymers, poly(2-propylacrylic acid), poly(2-ethylacrylic acid), Tat peptide, Arg patch, a knotted peptide, CysTAT, S19-TAT, R8

(SEQ ID NO: 73), pAntp, Pas-TAT, Pas-R8 (SEQ ID NO: 76), Pas-FHV, Pas-pAntP, F2R4 (SEQ ID NO: 79), B55, aurein, IMT-P8, BR2, OMOTAG1, OMOTAG2, pVEC, SynB3, DPV1047, C105Y, Transportan, MTS, hLF, PFVYLI (SEQ ID NO: 93), maurocalcine, imperatoxin, hadrucalin, hemicalcin, opicalcin-1, opicalcin-2, midkine (62-104), MCoTI-II, chlorotoxin, DRI-TAT, cF Φ R₄ (SEQ ID NO: 96), R₆W₃ (SEQ ID NO: 421), myristate, yBBR, or a fragment or variant thereof, or any combination thereof.

36. The composition of claim 34, wherein the cell-penetrating moiety is any one of SEQ ID NO: 71 – SEQ ID NO: 96, SEQ ID NO: 98 – SEQ ID NO: 101, SEQ ID NO: 116 – SEQ ID NO: 126, or SEQ ID NO: 403 – SEQ ID NO: 455.

37. The composition of any one of claims 1-3, wherein the peptide sequence comprises at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, at least 30, at least 31, at least 32, at least 33, at least 34, at least 35, at least 36, at least 37, at least 38, at least 39, at least 40, at least 41, at least 42, at least 43, at least 44, at least 45, at least 46, at least 47, at least 48, at least 49, at least 50, at least 51, at least 52, at least 53, at least 54, at least 55, at least 56, at least 57, at least 58 residues, at least 59, at least 60, at least 61, at least 62, at least 63, at least 64, at least 65, at least 66, at least 67, at least 68, at least 69, at least 70, at least 71, at least 72, at least 73, at least 74, at least 75, at least 76, at least 77, at least 78, at least 79, at least 80, or at least 81 amino acid residues.

38. The composition of any one of claims 1-3, wherein an active sequence is grafted onto the peptide or wherein the peptide is conjugated to, linked to, or fused to or grafted onto an active agent resulting in a peptide construct.

39. The compositions of claim 38, wherein the peptide construct comprises at least 80%, at least 90%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity with any one of SEQ ID NO: 33 – SEQ ID NO: 35, SEQ ID NO: 68 – SEQ ID NO: 70, SEQ ID NO: 168 – SEQ ID NO: 170, SEQ ID NO: 203 – SEQ ID NO: 205, SEQ ID NO: 225 – SEQ ID NO: 332, SEQ ID NO: 359 – SEQ ID NO: 361, SEQ ID NO: 373, SEQ ID NO: 375, SEQ ID NO: 381, SEQ ID NO: 384, SEQ ID NO: 385, SEQ ID NO: 388, SEQ ID NO: 389, SEQ ID NO: 393, SEQ ID NO: 396, SEQ ID NO: 397, SEQ ID NO: 400, or SEQ ID NO: 401.

40. The composition of any one of claims 1-3, wherein the active agent is linked via a linker, wherein the linker is selected from **TABLE 7** or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125.

41. The composition of any one of claims 38-40, wherein the active agent is an immunotherapeutic agent, a CTLA-4 targeting agent, a PD-1 targeting agent, a PDL-1 targeting agent, an IL15 agent, a fused IL-15/IL-15Ra complex agent, an IFN γ agent, an anti-CD3 agent, an ion channel modulator, a Kv1.3 inhibitor, an auristatin, MMAE, a maytansinoid, DM1, DM4, doxorubicin, a calicheamicin, a platinum compound, cisplatin, a taxane, paclitaxel, SN-38, a BACE inhibitor, a Bcl-xL inhibitor, WEHI-539, venetoclax, ABT-199, navitoclax, AT-101, obatoclax, a pyrrolobenzodiazepine or pyrrolobenzodiazepine dimer, a dolastatin, or a neurotransmitter.

42. The composition of claim 41, wherein the Kv1.3 inhibitor is Vm24, ShK-170, ShK-186, or ShK-192.

43. The composition of claim 42, wherein the Vm24 has a sequence of SEQ ID NO: 378 or SEQ ID NO: 391.

44. The composition of claim 42, wherein the Shk-185 has a sequence of SEQ ID NO: 379, SEQ ID NO: 390, or SEQ ID NO: 402.

45. The composition of claim 41, wherein the neurotransmitter is neurotensin, a neurotensin peptide variant, or a functional fragment thereof.

46. The composition of claim 45, wherein the neurotensin peptide variant comprises at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 350 or SEQ ID NO: 365 – SEQ ID NO: 369.

47. The composition of any one of claims 1-3, further comprising a half-life modifying agent coupled to the peptide.

48. The composition of claim 48, wherein the half-life modifying agent comprises a polymer, a polyethylene glycol (PEG), a hydroxyethyl starch, polyvinyl alcohol, a water soluble polymer, a zwitterionic water soluble polymer, a water soluble poly(amino acid), a water soluble polymer of proline, alanine and serine, a water soluble polymer containing glycine, glutamic acid, and serine, an Fc region, a fatty acid, palmitic acid, or a molecule that binds to albumin.

49. The composition of claim 47, wherein the half-life modifying agent comprises SA21.

50. The composition of claim, 49, wherein the SA21 has a sequence of SEQ ID NO: 376 or SEQ ID NO: 377.

51. The composition of any one of claims 1-3, further comprising a detectable agent coupled to the peptide by a linker.

52. The composition of claim 51, wherein the detectable agent is linked via a linker, wherein the linker is selected from **TABLE 7** or any one of SEQ ID NO: 103 – SEQ ID NO: 115, or SEQ ID NO: 125.

53. The composition of claim 51, wherein the detectable agent is a fluorophore, a near-infrared dye, a contrast agent, a nanoparticle, a metal-containing nanoparticle, a metal chelate, an X-ray contrast agent, a PET agent, a radionuclide, or a radionuclide chelator.

54. A pharmaceutical composition comprising the composition of any one of claims 1-53, or a salt thereof, and a pharmaceutically acceptable carrier.

55. A method for transporting a composition across a cell layer, the method comprising:

- a) contacting the cell layer with the composition comprising the peptide of any one of claims 1-53 or the pharmaceutical composition of claim 54; and
- b) transporting the composition across the cell layer.

56. A method of treating a condition in a subject in need thereof, the method comprising:

- a) administering to the subject a composition comprising the peptide of any one of claims 1-53 or the pharmaceutical composition of claim 54, and
- b) transporting the composition across a cell layer.

57. A method of treating a condition in a subject in need thereof, the method comprising administering to the subject an amount of a composition comprising the peptide of any one of claims 1-53 or the pharmaceutical composition of claim 54 based on the expression of TfR in a tissue of the subject.

58. The method of any one of claims 55-57, wherein the peptide is capable of binding and activating a neurotensin receptor.

59. The method of any one of claims 56-58, further comprising reducing a level of pain upon administration of the peptide to the subject.

60. The method of claim 59, wherein the administered is inhalation, intranasally, orally, topically, intravenously, subcutaneously, intramuscularly administration, intraperitoneally, intratumoral, intrathecal, or a combination thereof.

61. The method of claim 60, wherein the composition is administered intravenously as a bolus, infusion, or prolonged infusion.

62. The method of any one of claims 56-58, further comprising administering the peptide to a patient with schizophrenia, drug abuse, Parkinson's disease, high blood pressure, an eating disorder, stroke, Alzheimer's disease, cancer, inflammation, pain, epilepsy, a mood disorder, or an endocrine disorder.

63. The method of any one of claims 56-58, wherein the subject has a cancer.

64. The method of any one of claims 56-58, wherein the subject has a disease of the CNS.

65. The method of any one of claims 56-58, wherein the subject has neuropathic pain, migraine, anxiety, or epilepsy.

66. A method for transporting a composition according to any one of claims 1-53 or the pharmaceutical composition of claim 54 across a cell layer, the method comprising contacting the cell layer with the composition, and transporting the composition across the cell layer.

67. A method of treating a condition in a subject in need thereof, the method comprising administering to the subject the composition of any one of claims 1-53 or the pharmaceutical composition of claim 54.

68. The method of any one of claims 55-56 or 60, wherein the transporting comprises binding of the composition to TfR.

69. The method of claim 68, wherein the transporting further comprises transcytosis.

70. The method of claim 69, wherein the transcytosis is vesicular transcytosis.

71. The method of claim 69, wherein the transcytosis is TfR-mediated.

72. The method of claim 69, wherein the transporting further comprises dissociation of the composition from the TfR.

73. The method of claim 72, wherein the transporting further comprises releasing an active agent from the composition.

74. A method for designing a TfR-binding peptide, the method comprising:

a) using a computer program to identify a peptide from a library of peptides with optimized stability and folding, wherein each peptide of the library of peptides comprises at least three intramolecular disulfide bonds;

b) using a computer program to superimpose amino acid residues of a binding patch from a peptide:TfR complex to determine a stable peptide that binds to TfR after grafting the binding patch onto the peptide; and

c) grafting the binding patch onto the peptide to produce the stable peptide.

75. The method of claim 74, further comprising performing site saturation mutagenesis to obtain a mutated peptide with a higher binding affinity for TfR than the stable peptide.

76. The method of claim 74, wherein the peptide from a library of peptides is between 20 and 75 amino acid residues in length.




FIG. 1D




FIG. 1G

FIG. 1C




FIG. 1F

FIG. 1B




FIG. 1E

FIG. 1

2 / 35




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D

FIG. 2

3 / 35




FIG. 3A




FIG. 3B

FIG. 3




FIG. 4




FIG. 5A




FIG. 5B




FIG. 5C








FIG. 5D

FIG. 5



7 / 35




FIG. 7

8 / 35




FIG. 8

9 / 35




FIG. 9

10 / 35

SEQ ID NO: 32 Binding to Captured Biotinylated hTfR by SPR

Single Cycle Kinetics Analysis


 $k_a = 8.55(1) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ (Note: k_a is approaching the instrument limit) $k_d = 1.85(1) \times 10^{-3} \text{ s}^{-1}$ $K_D = 216(1) \text{ pM}$

FIG. 10

SEQ ID NO: 30 Binding to Captured Biotinylated hTfR by SPR

Single Cycle Kinetics Analysis


 $k_a = 8.57(2) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ (Note: k_a is approaching the instrument limit) $k_d = 4.16(1) \times 10^{-3} \text{ s}^{-1}$ $K_D = 486(1) \text{ pM}$

FIG. 11

12 / 35




FIG. 12

13 / 35




FIG. 13A




FIG. 13B

FIG. 13

14 / 35




FIG. 13C




FIG. 13D




FIG. 13E




FIG. 13

15 / 35




FIG. 14A

SEQ ID NO: 32




FIG. 14B

FIG. 14




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 15D

FIG. 15

17 / 35

SEQ ID NO: 1




FIG. 16A

SEQ ID NO: 2



FIG. 16B

SEQ ID NO: 30




FIG. 16C

SEQ ID NO: 32



FIG. 16D

FIG. 16

18 / 35




FIG. 17A




FIG. 17B




FIG. 17C




FIG. 17D

FIG. 17

19 / 35




FIG. 18

20 / 35




FIG. 19

21 / 35




FIG. 20A




FIG. 20B




FIG. 20C




FIG. 20D




FIG. 20E





FIG. 20F

FIG. 20

22 / 35



23 / 35




FIG. 22A




FIG. 22B

FIG. 22

24 / 35




FIG. 23A




FIG. 23B




FIG. 23C

FIG. 23




FIG. 24

26 / 35




FIG. 25

27 / 35




FIG. 26

28 / 35





FIG. 27



Compound (100 nM) on HEK293-NTSR1 cells

FIG. 28B




FIG. 28A

FIG. 28





FIG. 29



336

[illegible]




FIG. 31

33 / 35




FIG. 32

34 / 35




FIG. 33





FIG. 34A





FIG. 34B

FIG. 34




FIG. 7