



US00PP30827P3

(12) **United States Plant Patent**
Wain

(10) **Patent No.:** **US PP30,827 P3**

(45) **Date of Patent:** **Aug. 20, 2019**

(54) **CHRYSANTHEMUM PLANT NAMED**
'FICHRYSWIROS'

CPC **A01H 6/1424** (2018.05); **A01H 5/02**
(2013.01)

(50) Latin Name: ***Chrysanthemum X morifolium***
Varietal Denomination: **Fichryswiros**

(58) **Field of Classification Search**
USPC Plt./263.1, 284, 286
See application file for complete search history.

(71) Applicant: **DUMMEN GROUP B.V.**, De Lier
(NL)

(56) **References Cited**

(72) Inventor: **Peter Wain**, Locks Heath (GB)

PUBLICATIONS

(73) Assignee: **Dümmen Group B.V.**, De Lier (NL)

Straathof Plants 2017 catalog (<http://www.straathofplants.nl/downloads/straathof%20brochure%202017.pdf>). Citation for 'Swiftly Rosso'. 5 pages. (Year: 2017).*

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

* cited by examiner

(21) Appl. No.: **15/732,523**

Primary Examiner — Susan McCormick Ewoldt
Assistant Examiner — Karen M Redden

(22) Filed: **Nov. 21, 2017**

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(65) **Prior Publication Data**

US 2018/0146590 P1 May 24, 2018

Related U.S. Application Data

(60) Provisional application No. 62/497,574, filed on Nov. 22, 2016.

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/14 (2018.01)

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named 'Fichryswiros', characterized by its relatively compact, upright to outwardly spreading and uniformly mounded plant habit; moderately vigorous growth habit; freely branching habit; dense and full plant form; uniform and freely flowering habit; medium semi-decorative type inflorescences with dark red purple-colored ray florets; early flowering habit, response time about 37 days under controlled photoperiodic treatments; and good postproduction longevity.

(52) **U.S. Cl.**
USPC **Plt./286**

1 Drawing Sheet

1

2

Botanical designation: *Chrysanthemum X morifolium*.
Cultivar denomination: 'FICHRYSWIROS'.

Lier, The Netherlands in February, 2015. Asexual reproduction by terminal vegetative cuttings has shown that the unique features of this new *Chrysanthemum* plant are stable and reproduced true to type in successive generations.

BACKGROUND OF THE INVENTION

SUMMARY OF THE INVENTION

The present invention relates to a new and distinct *Chrysanthemum* plant, botanically known as *Chrysanthemum X morifolium*, commercially grown as a garden *Chrysanthemum* plant, referred to as code number 65838 in U.S. Provisional Patent Application Ser. No. 62/497,574 and hereinafter referred to by the name 'Fichryswiros'.

The new *Chrysanthemum* plant is a product of a planned breeding program conducted by the Inventor in Fareham, Hampshire, United Kingdom and De Lier, The Netherlands. The objective of the breeding program is to create new potted *Chrysanthemum* plants with numerous attractive inflorescences.

The new *Chrysanthemum* plant is a naturally-occurring whole plant mutation of *Chrysanthemum X morifolium* 'Swiftly Purple', not patented. The new *Chrysanthemum* plant was discovered and selected by the Inventor as a single flowering plant from within a population of plants of 'Swiftly Purple' in a controlled greenhouse environment in De Lier, The Netherlands in January, 2015.

Asexual reproduction of the new *Chrysanthemum* plant by terminal vegetative cuttings was first conducted in De

5

10

15

20

25

Plants of the new *Chrysanthemum* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature, daylength and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Fichryswiros'. These characteristics in combination distinguish 'Fichryswiros' as a new and distinct *Chrysanthemum* plant:

1. Relatively compact, upright to outwardly spreading and uniformly mounded plant habit.
2. Moderately vigorous growth habit.
3. Freely branching habit; dense and full plant form.
4. Uniform and freely flowering habit.
5. Medium semi-decorative type inflorescences with dark red purple-colored ray florets.

6. Early flowering habit, response time about 37 days under controlled photoperiodic treatments.

7. Good postproduction longevity.

Plants of the new *Chrysanthemum* can be compared to plants of the mutation parent, 'Swiftly Purple'. Plants of the new *Chrysanthemum* differ primarily from plants of 'Swiftly Purple' in ray floret color as plants of 'Swiftly Purple' have inflorescences with purple-colored ray florets.

Plants of the new *Chrysanthemum* can be compared to plants of *Chrysanthemum* X *morifolium* 'Breeze Cardinal', not patented. In side-by-side comparisons, plants of the new *Chrysanthemum* differ primarily from plants of 'Breeze Cardinal' in the following characteristics:

1. Plants of the new *Chrysanthemum* are more compact than plants of 'Breeze Cardinal'.
2. Plants of the new *Chrysanthemum* flower about nine days earlier than plants of 'Breeze Cardinal'.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying photograph illustrates the overall appearance of the new *Chrysanthemum* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum* plant. The photograph is a side perspective view of a typical flowering plant of 'Fichryswiros' grown in a 14-cm container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations and measurements describe plants grown during the summer in 14-cm containers in a glass-covered greenhouse in Fareham, Hampshire, United Kingdom and under cultural practices typical of commercial garden *Chrysanthemum* production. During the production of the plants, day and night temperatures ranged from 17° C. to 21° C. and light levels averaged 6,000 lux. Plants were grown under long day/short night conditions for about two weeks and then grown under short day/long night conditions to induce inflorescence initiation and development. Plants were eight weeks old when the photograph and detailed description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, Fifth Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Chrysanthemum* X *morifolium* 'Fichryswiros'.

Parentage: Naturally-occurring whole plant mutation of *Chrysanthemum* X *morifolium* 'Swiftly Purple', not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots, summer.—About ten days at temperatures about 21° C.

Time to initiate roots, winter.—About twelve days at temperatures about 21° C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures about 21° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures about 21° C.

Root description.—Fine, fibrous; typically light brown in color, actual color of the roots is dependent on

substrate composition, water quality, fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; medium density.

Plant description:

Plant and growth habit.—Herbaceous semi-decorative type potted *Chrysanthemum*; relatively compact; stems upright to outwardly spreading giving a uniformly mounded appearance to the plant; numerous lateral branches and relatively short internodes, dense and full plant form; moderately vigorous growth habit.

Plant height.—About 10 cm.

Plant width.—About 23 cm.

Branching habit.—Freely branching habit; about five primary lateral branches develop after removal of terminal apex (pinching).

Lateral branches.—Length: About 7 cm. Diameter: About 3 mm. Internode length: About 5 mm. Strength: Strong. Aspect: About 60° from vertical and then bending upwardly. Texture: Fine pubescence. Color: Close to 147B.

Leaf description:

Arrangement.—Alternate, simple.

Length.—About 3 cm.

Width.—About 2 cm.

Shape.—Palmately-lobed; roughly ovate with three to five lobes.

Apex.—Broadly acuminate.

Base.—Attenuate.

Margin.—Slightly dentate and palmately lobed; sinuses between lateral lobes mostly divergent.

Texture, upper and lower surfaces.—Fine pubescence; slightly rough; veins prominent on lower surface.

Color.—Developing leaves, upper surface: Close to N137A. Developing leaves, lower surface: Close to 147B. Fully expanded leaves, upper surface: Close to N137D; venation, close to 145A. Fully expanded leaves, lower surface: Close to 147B; venation, close to 145A.

Petioles.—Length: About 8 mm. Diameter: About 2 mm. Texture, upper and lower surfaces: Fine pubescence; slightly rough. Color, upper and lower surfaces: Close to 145A.

Inflorescence description:

Form and flowering habit.—Semi-decorative type inflorescence form with ligulate-shaped ray florets; inflorescences borne on terminals above and beyond the foliar plane; disc and ray florets arranged acropetally on a capitulum; freely flowering habit with about 24 inflorescences developing per plant during the flowering season.

Fragrance.—Mildly fragrant; pungent, herbaceous.

Flowering response.—Early flowering habit, plants flower uniformly about 37 days after starting short day/long night photoperiodic treatments.

Inflorescence longevity.—Good postproduction longevity; inflorescences maintain good color and substance for about two to three weeks on the plant; inflorescences persistent.

Inflorescence buds.—Height: About 6 mm. Diameter: About 7 mm. Shape: Oblate. Color: Close to 137C.

Inflorescence diameter.—About 3.7 cm.

Inflorescence height.—About 1.2 cm.

Disc diameter.—About 9 mm.

Receptacles.—Height: About 2 mm. Diameter: About 3 mm. Shape: Conical. Color: Close to 144C.

Ray florets.—Number of ray florets per inflorescence: About 33 arranged in about three whorls. Orientation: Initially upright, then about 45° from vertical. Length: About 7 mm. Width: About 6 mm. Shape: Ligulate. Apex: Mammillate. Base: Fused into a short tube. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; double-keeled. Color: When opening and fully opened, upper surface: Close to 59A; with development, color becoming closer to 60A. When opening and fully opened, lower surface: Close to 59B; with development, color becoming closer to 185C.

Disc florets.—Number of disc florets per inflorescence: About 194 massed at the center of the receptacle. Length: About 5 mm. Diameter: About 1 mm. Shape: Tubular, elongated; apices, acute. Texture, inner and outer surfaces: Smooth, glabrous. Color, when opening: Apex: Close to 9B. Mid-section: Close to 145C. Base: Close to 157C. Color, fully opened: Apex: Close to 9A. Mid-section: Close to 145B. Base: Close to 157C.

Phyllaries.—Number of phyllaries per inflorescence: About 23 arranged in about three whorls. Length: About 5 mm. Width: About 2 mm. Shape: Lanceolate. Apex: Acute. Base: Obtuse. Margin: Entire.

Texture, upper surface: Smooth, glabrous; waxy. Texture, lower surface: Fine pubescence; waxy. Color, upper and lower surfaces: Close to 137C.

Peduncles.—Length, terminal peduncle: About 1.2 cm. Diameter, terminal peduncle: About 2 mm. Angle: Erect to about 10° from vertical. Strength: Moderately strong, flexible. Texture: Densely pubescent. Color: Close to 138B.

Reproductive organs.—Androecium: None observed. Gynoecium: Present only on ray florets. Pistil length: About 6 mm. Stigma shape: Bi-parted. Stigma color: Close to 5C. Style length: About 0.2 mm. Style color: Close to 1D. Ovary color: Close to 1D.

Seeds and fruits.—Seed and fruit production has not been observed on plants of the new *Chrysanthemum* to date.

Disease & pest resistance: Resistance to pathogens and pests common to *Chrysanthemum* plants has not been observed on plants of the new *Chrysanthemum* to date.

Garden performance: Plants of the new *Chrysanthemum* have demonstrated good garden performance and to tolerate temperatures from about 0° C. to about 35° C.

It is claimed:

1. A new and distinct *Chrysanthemum* plant named 'Fichryswiros' as illustrated and described.

* * * * *

