

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



(10) International Publication Number

WO 2015/191951 A2

(43) International Publication Date

17 December 2015 (17.12.2015)

(51) International Patent Classification:

A61K 38/10 (2006.01) A61K 47/48 (2006.01)

(21) International Application Number:

PCT/US2015/035473

(22) International Filing Date:

12 June 2015 (12.06.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/011,368	12 June 2014 (12.06.2014)	US
62/077,460	10 November 2014 (10.11.2014)	US
62/108,772	28 January 2015 (28.01.2015)	US

(71) Applicant: RA PHARMACEUTICALS, INC. [US/US]; One Kendall Square, Suite B14301, Cambridge, Massachusetts 02139 (US).

(72) Inventors: ARATA, Michelle Denise; 10 Overlook Ridge Drive, Unit 328, Malden, Massachusetts 02148 (US). DHAMNASKAR, Ketki Ashok; 101 Western Ave., Apartment #37, Cambridge, Massachusetts 02139 (US). ELBAUM, Daniel; 25 Sherrin Road, Newton, Massachusetts 02462 (US). JOSEPHSON, Kristopher; 7 York Road, Wayland, Massachusetts 01778 (US). LARSON, Kelley Cronin; 84 Crescent Street, Quincy, Massachusetts 02169 (US). MA, Zhong; 341 Marret Road, Unit 8, Lexington, Massachusetts 02421 (US). NIMS, Nathan Ezekiel; 55 Brookside Avenue, Winchester, Massachusetts 01890 (US). RICARDO, Alonso; 35 Pleasant Street, #1, Cambridge, Massachusetts 02139 (US). SEYB, Kathleen;

368 Dorchester St, Unit 1, South Boston, Massachusetts 02127 (US). TANG, Guo-Qing; 93 Drummer Road, Acton, Massachusetts 01720 (US). TRECO, Douglas A.; 87 Brantwood Road, Arlington, Massachusetts 02476 (US). WANG, Zhaolin; 88 Kingsbury Street, Wellesley, Massachusetts 02481 (US). YE, Ping; 27 Deering Avenue, Lexington, Massachusetts 02421 (US). ZHENG, Hong; 25 Lake Shore Ct., Apt. 4, Brighton, Massachusetts 02135 (US). PERLMUTTER, Sarah Jacqueline; 15 Renfrew Street, Medford, Massachusetts 02155 (US).


(74) Agents: WARD, Donna T. et al.; DT Ward, PC, 142A Main Street, Groton, Massachusetts 01450 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) Title: MODULATION OF COMPLEMENT ACTIVITY



(57) Abstract: The present invention provides modulators of complement activity. Also provided are methods of utilizing such modulators as therapeutics.



SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, — *with sequence listing part of description (Rule 5.2(a))*
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report (Rule 48.2(g))*

MODULATION OF COMPLEMENT ACTIVITY

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Patent Application No. 62/011,368 filed June 12, 2014, entitled Modulation of Complement Activity, U.S. Provisional Patent Application No. 62/077,460 filed November 10, 2014, entitled Modulation of Complement Activity, and U.S. Provisional Patent Application No. 62/108,772 filed January 28, 2015, entitled Modulation of Complement Activity, the contents of each of which are herein incorporated by reference in their entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on June 11, 2015, is named 20111005PCT_SL.txt and is 130,110 bytes in size.

FIELD OF THE INVENTION

[0003] The present invention relates to compounds, including polypeptides, which are useful as inhibitors and/or antagonists of complement activity. Also provided are methods of utilizing the inhibitors as therapeutics.

BACKGROUND OF THE INVENTION

[0004] The vertebrate immune response is comprised of adaptive and innate immunity components. While the adaptive immune response is selective for particular pathogens and is slow to respond, components of the innate immune response recognize a broad range of pathogens and respond rapidly upon infection. One such component of the innate immune response is the complement system.

[0005] The complement system includes about 20 circulating proteins, synthesized primarily by the liver. Components of this particular immune response were first termed “complement” due to the observation that they complemented the antibody response in the destruction of bacteria. These proteins remain in an inactive form prior to activation in response to infection. Activation occurs by way of a pathway of proteolytic cleavage initiated by pathogen recognition

and leading to pathogen destruction. Three such pathways are known in the complement system and are referred to as the classical pathway, the lectin pathway and the alternative pathway. The classical pathway is activated when an IgG or IgM molecule binds to the surface of a pathogen. The lectin pathway is initiated by the mannan-binding lectin protein recognizing the sugar residues of a bacterial cell wall. The alternative pathway remains active at low levels in the absence of any specific stimuli. While all three pathways differ with regard to initiating events, all three pathways converge with the cleavage of complement component C3. C3 is cleaved into two products termed C3a and C3b. Of these, C3b becomes covalently linked to the pathogen surface while C3a acts as a diffusible signal to promote inflammation and recruit circulating immune cells. Surface-associated C3b forms a complex with other components to initiate a cascade of reactions among the later components of the complement system. Due to the requirement for surface attachment, complement activity remains localized and minimizes destruction to non-target cells.

[0006] Pathogen-associated C3b facilitates pathogen destruction in two ways. In one pathway, C3b is recognized directly by phagocytic cells and leads to engulfment of the pathogen. In the second pathway, pathogen-associated C3b initiates the formation of the membrane attack complex (MAC). In the first step, C3b complexes with other complement components to form the C5-convertase complex. Depending on the initial complement activation pathway, the components of this complex may differ. C5-convertase formed as the result of the classical complement pathway comprises C4b and C2a in addition to C3b. When formed by the alternative pathway, C5-convertase comprises two subunits of C3b as well as one Bb component.

[0007] Complement component C5 is cleaved by either C5-convertase complex into C5a and C5b. C5a, much like C3a, diffuses into the circulation and promotes inflammation, acting as a chemoattractant for inflammatory cells. C5b remains attached to the cell surface where it triggers the formation of the MAC through interactions with C6, C7, C8 and C9. The MAC is a hydrophilic pore that spans the membrane and promotes the free flow of fluid into and out of the cell, thereby destroying it.

[0008] An important component of all immune activity is the ability of the immune system to distinguish between self and non-self cells. Pathology arises when the immune system is unable to make this distinction. In the case of the complement system, vertebrate cells express proteins that protect them from the effects of the complement cascade. This ensures that targets of the

complement system are limited to pathogenic cells. Many complement-related disorders and diseases are associated with abnormal destruction of self cells by the complement cascade. In one example, subjects suffering from paroxysmal nocturnal hemoglobinuria (PNH) are unable to synthesize functional versions of the complement regulatory proteins CD55 and CD59 on hematopoietic stem cells. This results in complement-mediated hemolysis and a variety of downstream complications. Other complement-related disorders and diseases include, but are not limited to autoimmune diseases and disorders, neurological diseases and disorders, blood diseases and disorders and infectious diseases and disorders. Experimental evidence suggests that many complement-related disorders are alleviated through inhibition of complement activity. Therefore, there is a need for the development of compounds capable of selectively blocking complement-mediated cell destruction.

SUMMARY OF THE INVENTION

[0009] The present invention provides compounds, including polypeptides (e.g., peptidomimetics, cyclic polypeptides and cyclic peptidomimetics,) small molecules, antibodies, antibody fragments, and aptamers, and methods of using said compounds for the modulation of complement. According to the present invention, polypeptides may have the sequences set forth in SEQ ID NOS 1-201 or 211. They may be linear or cyclic and may comprise a cyclic loop formed by a bridging moiety between two amino acids. The bridging moieties may comprise a feature selected from the group consisting of a disulfide bond, an amide bond (lactam), a thioether bond, an aromatic ring, an unsaturated aliphatic hydrocarbon chain, a saturated aliphatic hydrocarbon chain, a triazole ring or combinations thereof. Furthermore, the cyclic loop may vary in length and may be 1 amino acid, 2 amino acids, 3 amino acids, 4 amino acids, 5 amino acids, 6 amino acids, 7 amino acids, 8 amino acids, 9, amino acids, 10 amino acids, 11 amino acids, 12 amino acids, 13 amino acids, 14 amino acids, 15 amino acids, 16 amino acids, or more.

[0010] The bridging moiety may comprise a disulfide between two cysteine residues or an aromatic ring produced by reaction with a poly(bromomethyl)benzene and such poly(bromomethyl)benzene may be selected from the group consisting of 1,2-bis(bromomethyl)benzene, 1,3-bis(bromomethyl)benzene and 1,4-bis(bromomethyl)benzene.

[0011] The bridging moiety may also comprise an aromatic ring produced by reaction with a compound selected from the group consisting of 2,6-bis(bromomethyl)pyridine, (E)-1,4-dibromobut-2-ene and 1,2-bis(bromomethyl)-4-alkylbenzene.

[0012] The bridging moiety may also comprise an amide bond produced by the reaction of the side chains of the following pairs of amino acids: lysine and glutamate, lysine and aspartate, ornithine and glutamate, ornithine and aspartate, homolysine and glutamic acid, homolysine and aspartic acid, other combinations of amino acids, unnatural amino acids or non-amino acid residues comprising a primary amine and a carboxylic acid.

[0013] According to the present invention, the polypeptides are useful in that they may exhibit complement C5 inhibitory activity. In some cases, such activity may range from a half-maximal inhibitory concentration (IC_{50}) of about 50 nM to about 200 nM or be less than 50 nM.

[0014] In some embodiments, the present invention provides polypeptides having tryptophan or one or more tryptophan analogs selected from the group consisting of 5-fluorotryptophan, 5-methyl-O-tryptophan, 1-methyltryptophan, D-tryptophan and 7-azatryptophan present within the amino acid sequence. Such compounds include those of SEQ ID NOS: 3, 9, 11, 19, 45, 46, 50 and 59, among others.

[0015] In some embodiments, polypeptides of the present invention may bind to a region on C5, wherein said region is distal to the C5a-C5b cleavage site. In some cases, polypeptides of the present invention may inhibit the cleavage of C5 into C5a and C5b cleavage products.

[0016] In some cases, polypeptides of the present invention may be part of an antibody.

[0017] Polypeptides of the present invention may be comprised in compositions. Such compositions may comprise one or more acceptable carrier or excipient. Such polypeptides may further be conjugated to another non-polypeptide moiety such as for example a water-soluble polymer. These polymers may be hydrophilic or hydrophobic. In some embodiments the hydrophilic polymer is selected from the group consisting of polyalkylene oxide homopolymers, polypropylene glycols, polyoxyethylenated polyols and copolymers thereof. It may also comprise polyethylene glycol (PEG). Compositions may further comprise polypeptides conjugated to an albumin binding polypeptide selected from the group consisting of SEQ ID NOS: 202-204. In some cases compositions may comprise polypeptides conjugated to a cell penetrating polypeptide selected from the group consisting of SEQ ID NOS: 205-210.

Compositions may further comprise polypeptides conjugated to a lipidic moiety. Lipidic moieties

may include, but are not limited to fatty acids, phospholipids and sterols. In some embodiments, lipidic moieties may be directly conjugated to polypeptides of the invention or linked to a polypeptide-PEG conjugate. Some compositions may comprise a pharmaceutically acceptable excipient.

[0018] In some embodiments, the present invention provides a method of inhibiting C5 cleavage in a cellular system comprising the use of a composition according to the present invention, such as those described above. Some such methods may reduce C5a levels. Further methods may reduce membrane attack complex (MAC) formation.

[0019] Some methods of the invention may lead to C5 cleavage inhibition with an IC₅₀ of from about 50 nM to about 200 nM. In some cases, methods of the present invention may comprise C5 cleavage inhibition with an IC₅₀ of less than 50 nM.

[0020] Some methods of the present invention may be applied to a cellular systems that is selected from an *in vitro* system, an *in vivo* system and an *ex vivo* system. Some *in vivo* systems may include subjects, such as human subjects. In some cases, the polypeptides and polypeptide compositions may be used in the treatment or prevention of a complement-related disease, disorder and/or condition in a subject. Administration to a subject may be selected from the group consisting of buccal, intranasal, oral, intravenous, intramuscular, intraperitoneal, subcutaneous, transdermal and intravitreal. In some cases, compositions may be used to treat paroxysmal nocturnal hemoglobinuria and/or atypical hemolytic uremic syndrome. In some cases, compositions of the invention may be used to treat one or more of an inflammatory indication, a wound, an injury, an autoimmune disease, a vascular indication, a neurological indication, a kidney-related indication, and an ocular disease.

[0021] In some embodiments, the present invention provides methods of inhibiting thrombin-induced complement activation in cellular systems comprising contacting such systems with polypeptides of the invention. Further provided are methods of treating hemolysis in subjects, comprising administering a polypeptide described herein. According to some such methods, the hemolysis treated is caused by thrombin-induced complement activation.

[0022] In some embodiments are provided kits for the diagnosis, prognosis, prophylaxis or treatment of a disease, disorder and/or condition in a mammal, including a human. Such kits would contain one or more polypeptides or polypeptide compositions of the present invention and optionally reagents and/or instructions for use. In such kits, the polypeptides may comprise a

detectable label or the ability to bind to a detectable label to form a detectable complex.

According to one embodiment, the label comprises a BODIPY-TMR label.

DESCRIPTION OF THE FIGURES

[0023] The foregoing and other objects, features and advantages will be apparent from the following description of particular embodiments of the invention, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating the principles of various embodiments of the invention.

[0024] Figure 1 is a graph displaying the results of an enzyme immunoassay (EIA) for the detection of C5a in supernatant from a human red blood cell (RBC) hemolysis assay with increasing concentrations of inhibitors R3002 (SEQ ID NO: 3) and R3008 (SEQ ID NO: 9). Levels of C5a correlate with complement activity and are therefore an indicator of the ability of the compounds tested to inhibit complement activity. Supernatant from the hemolysis assay was diluted 1:50 and assayed for C5a levels. C5a levels decreased in human hemolysis assay supernatant samples with increasing levels of either inhibitor assayed. R3002 (SEQ ID NO: 3) had an IC₅₀ of 5.4 nM while R3008 (SEQ ID NO: 9) had an IC₅₀ of 54.5 nM. As used herein, the term “IC₅₀” refers to the half maximal inhibitory concentration, a value used to indicate the amount of the inhibitor needed to reduce a given reaction or process by half.

[0025] Figure 2 is a graph displaying the results of an EIA for the detection of the membrane attack complex (MAC) in supernatant from a human RBC hemolysis assay with increasing concentrations of R3008 (SEQ ID NO: 9). Levels of the MAC correlate with complement activity and are therefore an indicator of the ability of R3008 (SEQ ID NO: 9) to inhibit complement activity. Supernatant from the hemolysis assay was diluted 1:5 and assayed for MAC levels. MAC levels decreased in hemolysis assay supernatant samples with increasing levels of the inhibitor assayed with an IC₅₀ of 33 nM.

[0026] Figure 3 is a graph displaying competitive fluorescence polarization (FP) data for test articles R3003 (SEQ ID NO: 4), R3011 (SEQ ID NO: 31), R3014 (SEQ ID NO: 55), R3023 (SEQ ID NO: 104), R3043 (SEQ ID NO: 50) and R3050 (SEQ ID NO: 23). FP allows binding events to be measured in a homogenous solution. A competitive binding assay was conducted wherein a 25 nM solution of compound R3076 (SEQ ID NO: 40), which has a fluorescent tag,

was combined with increasing amounts of the test articles and measured for changes in FP (in milli-polarization units; mP). Decreasing mP levels correlate with successful competition for C5 by the test articles. The averages of two independent experiments conducted in triplicate (+/- standard deviation) are shown. Of the articles tested, R3003 (SEQ ID NO: 4) was the most potent while R3023 (SEQ ID NO: 104), a control polypeptide, showed no activity at the highest concentration tested.

[0027] Figure 4 depicts results from a study in cynomolgus monkey. Changes in R3152 (SEQ ID NO: 153) plasma concentration (circles) following a single 3 mg/kg IV dose in cynomolgus monkey are shown. Also shown are changes in hemolytic activity (squares) at the same time points.

[0028] Figure 5 depicts results of compound monitoring in plasma following intravenous (IV; squares) or subcutaneous (SC; circles) administration of 2 mg/kg of R3152 (SEQ ID NO: 153) in male Sprague-Dawley rats. Monitoring comprised determination of combined plasma concentrations of R3152 (SEQ ID NO: 153) as well as its equipotent C-terminally deamidated metabolite, R3201 (SEQ ID NO: 211.)

[0029] Figure 6 shows the pharmacokinetics of compounds of the present invention in rats. Left: Male Sprague-Dawley rats (n=3) were injected intravenously at a single 2 mg/kg dose. Blood samples were collected at indicated time points, processed into plasma, and analyzed for the indicated compound by LC-MS. Black circles: R3176 (SEQ ID NO: 177) (unlipidated compound); Open circles: R3183 (SEQ ID NO: 184) (C16 lipidated compound). Right: Male Sprague-Dawley rats (n=3) were injected subcutaneously at a single 15 mg/kg dose. Blood samples were collected at indicated time points, processed into plasma, and analyzed for the indicated compound by LC-MS. Black circles: R3176 (SEQ ID NO: 177) (unlipidated compound); Open circles: R3183 (SEQ ID NO: 184) (C16 lipidated compound).

[0030] Figure 7 is a graph presenting the effects of R3183 (SEQ ID NO: 184) (C16 lipidated compound) or an anti-C5 monoclonal antibody similar to ECULIZUMAB® on inhibition of hemolysis via the thrombin-induced complement pathway.

DETAILED DESCRIPTION


[0031] The present invention relates to the discovery of novel C5 modulatory compounds. Such compounds may include, but are not limited to polypeptides (e.g. cyclic polypeptides,

peptidomimetics and cyclic peptidomimetics, small molecules, antibodies, antibody fragments, and aptamers. In some cases, C5 modulatory compounds are polypeptides, such as cyclic polypeptides, peptidomimetics and cyclic peptidomimetics which are useful in the diagnosis and/or treatment of diseases in which the inhibition of complement activation is desirable. In some embodiments, polypeptides of the invention specifically bind complement component C5. In further embodiments, polypeptides of the invention reduce complement-mediated cell lysis by preventing cleavage of C5 into C5a and C5b fragments.

[0032] As used herein, a “mimetic” refers to a molecule which exhibits some of the properties or features of another molecule. A “peptidomimetic” or “polypeptide mimetic” is a mimetic in which the molecule contains structural elements that are not found in natural polypeptides (*i.e.* polypeptides comprised of only the 20 proteinogenic amino acids). In a preferred embodiment, peptidomimetics are capable of recapitulating or mimicking the biological action(s) of a natural peptide. A peptidomimetic may differ in many ways from natural polypeptides, including, but not limited to changes in backbone structure and the presence of amino acids that do not occur in nature. In some cases, peptidomimetics may include amino acids with side chains that are not found among the known 20 proteinogenic amino acids, non-polypeptide-based bridging moieties used to effect cyclization between the ends or internal portions of the molecule, substitutions of the amide bond hydrogen moiety by methyl groups (N-methylation) or other alkyl groups, replacement of a peptide bond with a chemical group or bond that is resistant to chemical or enzymatic treatments, N- and C-terminal modifications, and conjugation with a non-peptidic extension (such as polyethylene glycol, lipids, carbohydrates, nucleosides, nucleotides, nucleoside bases, various small molecules, or phosphate or sulfate groups).

[0033] Some polypeptides of the invention may be cyclic. Cyclic polypeptides include any polypeptides that have as part of their structure one or more cyclic features such as a loop, bridging moiety, and/or an internal linkage. As used herein, the term “bridging moiety” refers to one or more components of a bridge formed between two adjacent or non-adjacent amino acids, unnatural amino acids or non-amino acids in a polypeptide. Bridging moieties may be of any size or composition. In some embodiments, bridging moieties may comprise one or more chemical bonds between two adjacent or non-adjacent amino acids, unnatural amino acids, non-amino acid residues or combinations thereof. In some embodiments, such chemical bonds may be between one or more functional groups on adjacent or non-adjacent amino acids, unnatural amino acids,



non-amino acid residues or combinations thereof. Bridging moieties may comprise one or more features including, but not limited to an amide bond (lactam), disulfide bond, thioether bond, aromatic ring, triazole ring, and hydrocarbon chain. In some embodiments, bridging moieties comprise an amide bond between an amine functionality and a carboxylate functionality, each present in an amino acid, unnatural amino acid or non-amino acid residue side chain. In some embodiments, the amine or carboxylate functionalities are part of a non-amino acid residue or unnatural amino acid residue. In some cases, bridging moieties may comprise bonds formed between residues that may include, but are not limited to (S)-2-amino-5-azidopentanoic acid (also referred to herein as “X02”), (S)-2-aminohept-6-enoic acid (also referred to herein as “X30”), (S)-2-aminopent-4-ynoic acid (also referred to herein as “X31”) and (S)-2-aminopent-4-enoic acid (also referred to herein as “X12”.) Bridging moieties may be formed through cyclization reactions using olefin metathesis. In some cases, such bridging moieties may be formed between X12 and X30 residues. In some embodiments, the bridging moiety comprises a disulfide bond formed between two thiol containing residues. In some embodiments, the bridging moiety comprises one or more thioether bonds. Such thioether bonds, may include those found in cyclo-thioalkyl compounds. These bonds are formed during a chemical cyclization reaction between chloro acetic acid (also referred to herein as “X35”) N-terminal modified groups and cysteine residues. In some cases, bridging moieties comprise one or more triazole ring. Such triazole rings may include, but are not limited to those formed by cyclization reaction between X02 and X31. In some embodiments, bridging moieties comprise non-protein or non-polypeptide based moieties, including, but not limited to cyclic rings (including, but not limited to aromatic ring structures (e.g. xylyls)). Such bridging moieties may be introduced by reaction with reagents containing multiple reactive halides, including, but not limited to poly(bromomethyl)benzenes, poly(bromomethyl)pyridines, poly(bromomethyl)alkylbenzenes and/or (E)-1,4-dibromobut-2-ene. In some embodiments, bridging moieties of the present invention include, but are not limited to the following structures:




wherein each X is independently N or CH, such that no ring contains more than 2 N; each Z is independently absent or selected from a bond, NR, O, S, CH₂, C(O)NR, NRC(O), S(O)_vNR and NRS(O)_v; each m is independently selected from 0, 1, 2, and 3; each v is independently selected from 1 and 2; each R is independently selected from H and C₁-C₆; and each bridging moiety is connected to the polypeptide by an independently selected bond or C₁-C₆ spacer.

[0034] In certain embodiments of this invention, polypeptides are rendered macrocyclic by formation of covalent bonds between atoms present within the linear polypeptide and atoms of a bridging moiety. This bridging moiety serves the purpose of chemically tethering two reactive sites on the linear polypeptide so as to furnish a cyclic polypeptide product. Embodiments of the present invention include polypeptides cyclized in the aforementioned manner and comprising a bridging moiety containing an aromatic, 6-membered ring. In these embodiments, atoms of the

linear polypeptide that form explicit chemical bonds with the bridging moiety may be heteroatoms (including, but not limited to, nitrogen, oxygen and sulfur), or saturated or unsaturated carbon atoms. In each of these embodiments of this invention, the atoms of the polypeptide side chain may be bonded directly to a carbon atom within the aromatic ring of the bridging moiety. In alternative forms, the atoms of the polypeptide side chain may be bonded to a saturated -CH₂- group that is in turn directly bonded to a carbon atom within the aromatic ring of the bridging moiety. In certain cases, the aromatic, 6-membered ring within the bridging moiety is benzene, as in the following structures wherein Z may be selected from NH, S, O and (CH)₂:




[0035] In alternative forms of this invention, the aromatic, 6-membered ring that comprises the bridging moiety is heterocyclic and contains one or more nitrogen atoms. In these embodiments, the aromatic heterocycle may be pyridine, containing a single nitrogen atom in the aromatic ring [e.g. any of the structures below wherein Z may be selected from NH, S, O and (CH)₂]:



$Z = \text{NH, S, O, } (\text{CH}_2)_2$


[0036] Aromatic heterocycles may alternatively be pyridazine, containing two adjacent nitrogen atoms in a 1,2-orientation within the aromatic ring [e.g. any of the structures below wherein Z may be selected from NH, S, O and $(\text{CH}_2)_2$]:

XXXVIII




$Z = \text{NH, S, O, } (\text{CH}_2)_2$

[0037] In other embodiments, the aromatic heterocycle may be pyrimidine, containing two nitrogen atoms in a 1,3-orientation within the aromatic ring [e.g. any of the structures below wherein Z may be selected from NH, S, O and $(\text{CH}_2)_2$]:




$Z = \text{NH, S, O, } (\text{CH}_2)_2$


[0038] Alternatively, the aromatic heterocycle may be pyrazine, containing two nitrogen atoms in a 1,4-orientation within the aromatic ring [e.g. any of the structures below wherein Z may be selected from NH, S, O and $(\text{CH}_2)_2$]:



[0039] In alternative forms of this invention, polypeptides are rendered macrocyclic as a result of the formation of covalent bonds between atoms of the linear polypeptide and atoms of a bridging moiety consisting of a heterocyclic, aromatic, 5-membered ring. In these embodiments, atoms of the linear polypeptide that form explicit chemical bonds with the bridging moiety may be heteroatoms (including, but not limited to, nitrogen, oxygen and sulfur), or saturated or unsaturated carbon atoms. In each of these embodiments of this invention, the atoms of the polypeptide side chain may be bonded directly to a carbon atom or nitrogen atom within the aromatic ring of the bridging moiety. In alternative forms, the atoms of the polypeptide side chain may be bonded to a saturated -CH₂- group that is in turn directly bonded to a carbon atom or nitrogen atom within the aromatic ring of the bridging moiety. In certain cases, the heterocyclic, aromatic, 5-membered ring within the bridging moiety is 1,2,3-triazole. In these embodiments, the aromatic ring may be substituted at positions 1 and 4 with chemical functionality of the linear polypeptide that are being tethered. Alternatively, the 1,2,3-triazole scaffold may be substituted at positions 1 and 4 with -CH₂- groups that are directly bonded to the atoms of the linear polypeptide being tethered [e.g. either of the structures below wherein Z may be selected from NH, S, O and (CH₂)₂]:



[0040] In other embodiments of this invention, the heterocyclic, aromatic, 5-membered ring that comprises the bridging moiety is pyrazole. In these embodiments, the aromatic ring may be substituted either at positions 1 and 3 or at positions 1 and 4 with chemical functionality of the linear polypeptide that are being tethered. Alternatively, the pyrazole scaffold may be substituted either at positions 1 and 3 or at positions 1 and 4 with -CH₂- groups that are directly bonded to the atoms of the linear polypeptide being tethered [e.g. any of the structures below wherein Z may be selected from NH, S, O and (CH₂)₂]:



[0041] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the cyclic polypeptides and methods featured in the invention, suitable methods and materials are described below.

Polypeptides as Drugs

[0042] By virtue of their size and complexity, polypeptides are able to form numerous, highly specific contacts with their biological targets and can show a high level of selectivity for the correct or desired target as compared to a closely related target within the same family. Off-target effects (known also as side effects) often cause highly effective drugs to fail regulatory approval due to safety concerns.

[0043] Numerous polypeptides (including, but not limited to peptidomimetics, have been developed into effective drugs. These include, but are not limited to, insulin, glucagon-like peptide 1 (GLP-1), somatostatin, vasopressin, cyclosporine A, and the like. The therapeutic polypeptide can be identical to the naturally occurring molecule (i.e. that which circulates in humans and is considered “wild-type” in the human population). In many other cases, the polypeptide is not suitable or sub-optimal for therapeutic use due to a short circulating half-life that is often due to metabolic instability in the body. In these cases a modified or a variant form of the polypeptide (peptidomimetic) is used which results in improved pharmacokinetic and pharmacodynamic behavior. In other cases a polypeptide derived from a natural source has an equivalent mechanism of action and a preferred pharmaceutical profile and can be used as a therapy. For example, exenatide, a synthetic version of exendin-4, has biological properties similar to human glucagon-like peptide-1 (GLP-1) but improved pharmacokinetics, and has been approved by the FDA for the treatment of diabetes mellitus type 2. As another example, salmon calcitonin, calcitonin extracted from the Ultimobranchial glands of salmon, resembles human calcitonin but is more active than human calcitonin and may be used to treat postmenopausal osteoporosis, hypercalcaemia, Paget’s disease, bone metastases and phantom limb pain.

[0044] Polypeptides are typically limited to non-oral routes of administration. In nearly all cases, polypeptides must be delivered by injection, since even very short polypeptides (e.g., polypeptides with 4-10 amino acid residues) are incapable or poorly capable of passing through the cell membranes lining the intestinal tract. For efficient oral availability, drugs typically need to pass through both the luminal and basolateral membranes of gut epithelial cells in order to enter the systemic circulation. The poor membrane permeability and lack of oral bioavailability of polypeptides significantly limits their therapeutic use.

[0045] The effectiveness of a polypeptide as a drug may be influenced by its proteolytic stability. Within the body, polypeptides can be modified or degraded by enzymes, which can limit their effectiveness for interacting with an intended target.

[0046] Metabolic stability of polypeptides is important as it is related to their global flexibility, intramolecular fluctuations, various internal dynamic processes as well as many biologic functions. The metabolic stability of polypeptides may be critical in the development of pharmaceuticals, affecting parameters such as, but not limited to, clearance, half-life and bioavailability of the drugs.

[0047] Maintaining a given level of a therapeutic polypeptide within the body or the bloodstream may be difficult due to efflux. The rate of efflux of a polypeptide from the body may vary and should be monitored when considering the administration of therapeutic polypeptides.

[0048] There remains a significant medical need for complement activation inhibitors or inhibitors of complement activity and inhibitor formulations that are highly potent and highly specific.

Discovery of peptidomimetics

[0049] Peptidomimetics may be identified by a variety of means. In some cases a naturally occurring peptide or a sequence found in a natural protein is used as a starting point. In these instances, the starting peptide sequence has been chosen because it is known to physically interact with a desired target molecule. A natural peptide may be chosen because it is an agonist or antagonist for a receptor, inhibits an enzyme, or modulates a channel. A sequence found in a natural protein may be chosen because it comprises a domain that participates in an interaction with another protein or some other molecule in a human or animal. In many cases, structural data on interacting proteins can be obtained from public databases (e.g. the RCSB Protein Data Bank; H.M. Berman, J. Westbrook, Z. Feng, G. Gilliland, T.N. Bhat, H. Weissig, I.N. Shindyalov, P.E. Bourne (2000) The Protein Data Bank Nucleic Acids Research, 28: 235-242) and the specific region of a protein that interacts with the desired target can be identified from crystallographic data on the protein complex. In other cases, polypeptides corresponding to various portions of a protein can be prepared and tested for binding to a target of interest. Once identified, chemical modifications are introduced to improve its stability and potency, with the resulting peptidomimetic having improved pharmacokinetic or pharmacodynamic parameters.

[0050] In other cases, a polypeptide is isolated by one of several methods for isolating polypeptide sequences from libraries of polypeptides based on their affinities to specific target proteins, nucleic acids, carbohydrates, lipids, or whole cells. Such methods include phage display, mRNA display, ribosome display, DNA display, DNA-encoded assembly, and two-hybrid screening, as well as their modifications (See, e.g., Takashashi, T.T et al. (2003). Trends in Biochem. Sci. 28(3):159-165; Kay, B.K. et al. (2001). Methods. 24:240-246; He, M and Taussig, M (2002). Briefs in Functional Genomics and Proteomics. 1(2): 204-212; Rothe, A. et

al. (2006). The FASEB Journal. 20(10):1599-1610; all of which are included herein by reference in their entireties.)

[0051] Polypeptides can adopt three-dimensional structures that are capable of binding to other biological molecules with certain degrees of affinity and specificity. Some will bind with very high affinity and specificity. A library of random polypeptide sequences will be populated by molecules with a wide variety of three-dimensional structures. In order to isolate a polypeptide with a conformation that interacts with a specific target protein, individual sequences from the library can be prepared and tested or screened for their affinity to the target. However, for very large libraries ($>10^6$ members), the screening of individual sequences for binding affinity is not feasible. To overcome this limitation, a number of techniques have been developed to select novel polypeptides from extremely large, complex mixtures by virtue of their binding affinity to a target. Since high affinity binding polypeptides are predicted to be present at a very low frequency within the population, these selection methods rely on maintaining a physical link between the polypeptide and the genetic material (generally a nucleic acid such as DNA or RNA) encoding the polypeptide so that selection of the polypeptide automatically includes selection of a nucleic acid encoding it. The nucleic acid encoding the selected polypeptide can be amplified and sequenced to reveal the sequence of both the nucleic acid and the polypeptide. In one approach, phage display (see Cwirla, S.E. et al. (1990). Proc. Natl. Acad. Sci. U.S.A. 87:6378-6382; Dower, W.J. and Cwirla, S.E. U.S. Patent Nos. 5,427,908 and 5,580,717), each random polypeptide member of the library is displayed on the surface of a bacteriophage particle as part of a fusion protein between the polypeptide and one of the phage coat proteins. The phage particle provides the link between the polypeptide and the encoding DNA by co-localizing them within the same physical entity, and the encoding DNA can be subsequently amplified by infecting bacteria with the selected phage. In another approach, ribosome display (see Kawasaki, G.H. U.S. Patent Nos. 5,658,754 and 5,643,768), a mixture of messenger RNA (mRNA) molecules is translated *in vitro* in a manner that produces, for each mRNA in the mixture, a stabilized complex of ribosome, mRNA, and newly synthesized polypeptide protruding from the ribosome. Stabilizing the complex permits it to be held together while the polypeptides are screened for binding to a target of interest. The mRNAs encoding the selected polypeptides can be amplified using polymerase chain reaction (PCR), and then characterized, e.g., by sequencing.

[0052] In yet another approach, mRNA display (see Szostak, J.W. and Roberts, R.W., U.S. Patent No. 6,258,558, the contents of which are incorporated herein by reference in their entirety), each mRNA molecule in the library is modified by the covalent addition of a puromycin-like moiety at its 3' terminus. The puromycin-like moiety is an aminoacyl-tRNA acceptor stem analog that functions as a peptidyl acceptor, and can be added to a growing polypeptide chain by the peptidyl transferase activity of a ribosome translating the mRNA. During *in vitro* translation, the mRNA and the encoded polypeptide become covalently linked through the puromycin-like moiety, creating an RNA-peptide fusion. After selecting a fusion molecule by binding of its polypeptide component to a target, the RNA component of the selected fusion molecule can be amplified using PCR, and then characterized. Several other methods have been developed to produce a physical linkage between a polypeptide and its encoding nucleic acid to facilitate selection and amplification (see Yanagawa, H., Nemoto, N., Miyamoto, E., and Husimi, Y., U.S. Patent No. 6,361,943; Nemoto, H., Miyamoto-Sato, E., Husimi, H., and Yanagawa, H. (1997). FEBS Lett. 414:405-408; Gold, L., Tuerk, C., Pribnow, D., and Smith, J.D., U.S. Patent Nos. 5,843,701 and 6,194,550; Williams, R.B., U.S. Patent No. 6,962,781; Baskerville, S. and Bartel, D.P. (2002). Proc. Natl. Acad. Sci. USA 99:9154-9159; Baskerville, D.S. and Bartel, D.P., U.S. Patent No. 6,716,973; Sergeeva, A. et al. (2006). Adv. Drug Deliv. Rev. 58:1622-1654; the contents of each of which are incorporated herein by reference in their entirety).

[0053] mRNA display is a particularly useful method for creating large libraries of polypeptides. Accordingly, provided herein are methods of selecting for a polypeptide (or an mRNA encoding a polypeptide) that interacts with complement protein C5. A library will generally contain at least 10^2 members, more preferably at least 10^6 members, and more preferably at least 10^9 members (e.g., any of the mRNA-polypeptide complexes). In some embodiments, the library will include at least 10^{12} members or at least 10^{14} members. In general, the members will differ from each other; however, it is expected there will be some degree of redundancy in any library. The library can exist as a single mixture of all members, or can be divided into several pools held in separate containers or wells, each containing a subset of the library, or the library can be a collection of containers or wells on a plate, each container or well containing just one or a few members of the library.

[0054] Each mRNA in the library preferably comprises a translation initiation sequence, a start codon, and a variable polypeptide (e.g., protein or short peptide) coding region that is generated by, for example, a random or semi-random assembly of nucleotides, and varies from mRNA to mRNA in the library (though there will likely be some degree of redundancy within the library). The translation initiation sequence, start codon, and variable polypeptide coding region can be flanked by known, fixed sequences that can be used for PCR amplification of the mRNA, e.g., after selection. Other fixed sequences that can be present include those corresponding to sequences that encode amino acids that can participate in chemical or enzymatic cross-linking reactions, such that the polypeptide produced can be modified or derivatized after translation, or that encode a fixed C-terminal extension such as a polypeptide tag that can facilitate purification of the peptide-mRNA fusions.

[0055] Once a library of mRNA derivatized with puromycin is generated, the library can be translated. The resulting polypeptides (e.g., displayed polypeptides) will be linked to their corresponding mRNAs as described herein (e.g., as an mRNA-polypeptide complex).

[0056] Numerous *in vitro* translation systems have been described in the literature. The most common systems utilize rabbit reticulocyte lysates, wheat germ extracts, or *E. coli* extracts, which are available from a number of commercial sources in kit form (e.g., Ambion, Austin, TX; Promega, Madison, WI; Novagen/EMD Chemicals, Gibbstown, NJ; Qiagen, Valencia, CA).

[0057] Unlike phage display or other systems that rely on translation within cells, mRNA display can be adapted to directly produce libraries of peptidomimetics by performing *in vitro* translation with unnatural or non-standard amino acids. The 20 natural proteinogenic amino acids are identified and referred to herein by either the one-letter or three-letter designations as follows: aspartic acid (Asp:D), isoleucine (Ile:I), threonine (Thr:T), leucine (Leu:L), serine (Ser:S), tyrosine (Tyr:Y), glutamic acid (Glu:E), phenylalanine (Phe:F), proline (Pro:P), histidine (His:H), glycine (Gly:G), lysine (Lys:K), alanine (Ala:A), arginine (Arg:R), cysteine (Cys:C), tryptophan (Trp:W), valine (Val:V), glutamine (Gln:Q) methionine (Met:M), asparagine (Asn:N). Naturally occurring amino acids exist in their levorotatory (L) stereoisomeric forms. Amino acids referred to herein are L-stereoisomers except where otherwise indicated.

[0058] Unnatural amino acids have side chains or other features not present in the 20 naturally-occurring amino acids listed above and include, but are not limited to: N-methyl amino acids, N-alkyl amino acids, alpha, alpha substituted amino acids, beta-amino acids, alpha-

hydroxy amino acids, D-amino acids, and other unnatural amino acids known in the art (See, e.g., Josephson et al., (2005) *J. Am. Chem. Soc.* 127: 11727-11735; Forster, A.C. et al. (2003) *Proc. Natl. Acad. Sci. USA* 100: 6353-6357; Subtelny et al., (2008) *J. Am. Chem. Soc.* 130: 6131-6136; Hartman, M.C.T. et al. (2007) *PLoS ONE* 2:e972; and Hartman et al., (2006) *Proc. Natl. Acad. Sci. USA* 103:4356-4361).

[0059] Essentially any amino acid that, when attached to an appropriate tRNA, can be assembled into a polymer by natural or mutant ribosomes can be used (see Sando, S. et al., (2007) *J. Am. Chem. Soc.* 129:6180-6186; Dedkova, L. et al. (2003) *J. Am. Chem. Soc.* 125: 6616-6617; Josephson, K., Hartman, M.C.T., and Szostak, J.W. (2005) *J. Am. Chem. Soc.* 127:11727-11735; Forster, A.C. et al. (2003) *Proc. Natl. Acad. Sci. USA* 100:6353-6357; Subtelny, A.O., Hartman, M.C.T., and Szostak, J.W. (2008) *J. Am. Chem. Soc.* 130:6131-6136; and Hartman, M.C.T. et al. (2007) *PLoS ONE* 2:e972).

[0060] When unnatural amino acids are desired, it may be advantageous to use a purified translation system that lacks endogenous aminoacylated tRNAs (Shimizu, Y. et al. (2001) *Nat. Biotech.* 19:751-755; Josephson, K., Hartman, M.C.T., and Szostak, J.W. (2005) *J. Am. Chem. Soc.* 127: 11727-11735; Forster, A.C. et al. (2003) *Proc. Natl. Acad. Sci. USA* 100: 6353-6357). If unnatural amino acids are used with an *in vitro* translation system based on a lysate or extract, it may be desirable to deplete the extract of endogenous tRNAs, as previously described (see Jackson, R.J., Napthine, S., and Brierley, I. (2001) *RNA* 7:765-773). A system based on purified *E. coli* translation factors is commercially available (PUREEXPRESSTM; New England Biolabs, Ipswich, MA). These systems are particularly useful for translation with unnatural amino acids to produce peptidomimetics.

[0061] When using natural amino acids with an *in vitro* translation system based on a lysate or extract, translation is dependent on the enzymatic charging of amino acids onto tRNAs by tRNA synthetases, all of which are components of the extracts. Alternatively, *in vitro* translation systems that use purified translation factors and ribosomes, or tRNA-depleted extracts, require that aminoacylated tRNAs be provided. In these instances, purified or *in vitro* synthesized tRNAs can be charged with amino acids using chemical (see Frankel, A., Millward, S.W., and Roberts, R.W. (2003) *Chem. Biol.* 10:1043-1050) or enzymatic procedures (Josephson, K.,

Hartman, M.C.T., and Szostak, J.W. (2005) J. Am. Chem. Soc. 127: 11727-11735; Murakami, H. et al. (2006) Nat. Methods 3:357-359).

[0062] Numerous publications describe the recovery of mRNA-displayed polypeptides from translation complexes, and these are suitable for use with the methods described herein (Liu, R. et al. (2000). Methods Enzymol. 318:268-293; Baggio, R. et al. (2002). J. Mol. Recognit. 15:126-134; U.S. Pat. No. 6,261,804). The recovery of mRNA-displayed polypeptides may be facilitated by the use of various “tags” that are included in the polypeptide by translation of fixed sequences of the polypeptide coding sequence and which bind to specific substrates or molecules. Numerous reagents for capturing such tags are commercially available, including reagents for capturing the His-tag, FLAG-tag, glutathione-S-transferase (GST) tag, strep-tag, HSV-tag, T7-tag, S-tag, DsbA-tag, DsbC-tag, Nus-tag, myc-tag, hemagglutinin (HA)-tag, or Trx-tag (Novagen, Gibbstown, NJ; Pierce, Rockford, IL). mRNA-displayed polypeptides can also be isolated by binding of a polyA tail on the mRNA to polydT resin, or a combination of a polyA tail and a His-tag.

[0063] After the *in vitro* translation reaction has been performed, and prior to the selection step, the mRNA portion of the functionalized RNA is typically reversed-transcribed to produce a RNA-DNA hybrid molecule. This serves to protect the RNA from degradation, and also prevents the RNA from folding into a secondary structure that could bind to the selection target, which would lead to selection of inappropriate products (e.g., the selection of RNA aptamers rather than polypeptide aptamers).

[0064] After *in vitro* translation and isolation of polypeptide-mRNA fusions, the polypeptide moiety may be modified by intramolecular or intermolecular cross-linking, chemical conjugation, enzymatic cleavage, truncation, or extension with additional amino acid monomers. One way to accomplish this is by incorporating unnatural amino acids with reactive side chains into the polypeptides that make up the library. After translation, the newly formed polypeptides can be reacted with molecules that react specifically with the reactive side chain of the incorporated amino acid. For example, an amino acid with a terminal alkyne side chain can be incorporated into the polypeptide library and subsequently reacted with an azido sugar, creating a library of displayed polypeptides with sugars attached at the positions of the alkynyl side chains (Josephson, K., Hartman, M.C.T., and Szostak, J.W. (2005) J. Am. Chem. Soc. 127: 11727-

11735). A variety of reactive side chains can be used for such post-translational conjugation, including amines, carboxyl groups, azides, terminal alkynes, alkenes, and thiols.

[0065] One particularly useful modification is based on the cross-linking of amino acids to produce cyclic structures. Cyclic regions in a polypeptide contain a rigid domain, which reduces conformational flexibility and degrees of rotational freedom, leading to very high affinity binding to target proteins. A number of methods for cyclizing a polypeptide are available to those skilled in the art and are incorporated herein by reference. Typically, the chemical reactivity of specific amino acid side chains and/or the carboxyl or amino termini of the polypeptide are exploited to crosslink two sites of the polypeptide to produce a cyclic molecule. In one method, the thiol group of a cysteine residue is cross-linked with another cysteine residue to form a disulfide bond. In some embodiments, thiol groups of cysteine residues react with bromomethyl groups of poly(bromomethyl)benzene molecules to form stable linkages (see Timmerman, P. et al., (2005) *ChemBioChem* 6:821-824, the contents of which are herein incorporated by reference in their entirety). Poly(bromomethyl)benzene molecules of the present invention may include, but are not limited to 1,2-bis(bromomethyl)benzene, 1,3-bis(bromomethyl)benzene and 1,4-bis(bromomethyl)benzene. Bis-, tris- and tetrakis(bromomethyl)benzene molecules, for example, can be used to generate bridging moieties to produce polypeptides with one, two or three loops, respectively. Bromomethyl groups of a poly(bromomethyl)benzene molecule may be arranged on the benzene ring on adjacent ring carbons (ortho- or o-), with a ring carbon separating the two groups (meta- or m-) or on opposite ring carbons (para- or p-). In some embodiments, m-bis(bromomethyl)benzene (also referred to herein as m-dibromoxylene) is utilized in the formation of cyclic polypeptides. In some embodiments, o-bis(bromomethyl)benzene (also referred to herein as o-dibromoxylene) or p-bis(bromomethyl)benzene (also referred to herein as p-dibromoxylene) are utilized in the formation of cyclic polypeptides. In some embodiments, thiol groups of cysteine residues react with other reagents comprising one or more bromo functional groups to form stable linkages. Such reagents may include, but are not limited to poly(bromomethyl)pyridines (including, but not limited to 2,6-bis(bromomethyl)pyridine), poly(bromomethyl)alkylbenzenes (including, but not limited to 1,2-bis(bromomethyl)-4-alkylbenzene) and/or (E)-1,4-dibromobut-2-ene.

[0066] In another exemplary method, a side chain amino group and a terminal amino group are cross-linked with disuccinimidyl glutarate (see Millward, S.W. et al., *J. Am. Chem. Soc.*

127:14142-14143, 2005). In other approaches, cyclization is accomplished by forming a thioether bond between two sites on the polypeptide (see Timmerman, P. et al., (2005) *ChemBioChem* 6:821-824; incorporated by reference herein in its entirety). An enzymatic method relies on the reaction between (1) a cysteine and (2) a dehydroalanine or dehydrobutyryne group, catalyzed by a lantibiotic synthetase, to create the thioether bond (see Levengood, M.R. and Van der Donk, W.A., *Bioorg. and Med. Chem. Lett.* 18:3025-3028, 2008). The dehydro functional group can also be generated chemically by the oxidation of selenium containing amino acid side chains incorporated during translation (see Seebeck, F.P. and Szostak, J.W. *J. Am. Chem. Soc.* 2006).

[0067] A library of mRNA-polypeptide fusions (also referred to herein as an mRNA display library) generated using the methods described above, and which may or not have been subjected to a post-translational modification (such as cyclization of the polypeptide, as described above), can be subjected to a batch selection step to isolate those complexes displaying desirable polypeptides.

[0068] Typically, C5 is conjugated to a solid substrate, such as an agarose or synthetic polymer bead. Numerous methods are available for immobilizing C5 to a solid support. In one particularly useful method, C5 is conjugated to biotin and streptavidin beads are used to immobilize the protein. The beads comprising the immobilized C5 are mixed with the mRNA display library and incubated under conditions (e.g., temperature, ionic strength, divalent cations, and competing binding molecules) that permit specific members of the library to bind the target. Alternatively, the biotinylated enzyme can be free in solution and, after binding to an appropriate polypeptide, the mRNA-polypeptide fusions bound to C5 are captured by appropriately modified beads.

[0069] The binding conditions can be varied in order to change the stringency of the selection. For example, low concentrations of a competitive binding agent can be added to ensure that the selected polypeptides have a relatively higher affinity. Alternatively, the incubation period can be chosen to be very brief, such that only polypeptides with high k_{on} rates (rate of association) will be isolated. In this manner, the incubation conditions play an important role in determining the properties of the selected polypeptides. Negative selections can also be employed. In this case, a selection to remove polypeptides with affinity to the substrate to which the target is bound (e.g., Sepharose) is carried out by applying the displayed library to substrate

beads lacking the target protein. This step can remove mRNAs and their encoded polypeptides that are not specific for the target protein. Numerous references describing how to conduct selection experiments are available. (See, e.g., U.S. Patent No. 6,258,558, Smith, G.P. and Petrenko, V.A., (1997) *Chem. Rev.* 97:391-410; Keefe, A.D. and Szostak, J.W. (2001) *Nature* 15:715-718; Baggio, R. et al. (2002) *J. Mol. Recog.* 15:126-134 and Sergeeva, A. et al. (2006) *Adv. Drug Deliv. Rev.* 58:1622-1654; the contents of each of which are herein incorporated by reference in their entirety).

[0070] The frequency at which binding molecules are present in a library of random sequences is expected to be very low. Thus, in the initial selection step, very few polypeptides meeting the selection criteria (and their associated mRNAs) should be recovered. Typically, the selection is repeated with mRNAs selected from the first round of selection. This is accomplished by using PCR to amplify the mRNAs or corresponding cDNAs selected in the first round, followed by *in vitro* transcription to produce a new library of mRNAs. PCR primers corresponding to the 5' and 3' ends of the mRNAs in the library are used. Typically, the 5' primer will extend in the 5' direction beyond the end of the mRNA so that a bacterial promoter, such as a T7 promoter, is added to the 5' end of each amplified molecule. Once amplified, the double-stranded DNA can be used in an *in vitro* transcription reaction to generate the mRNA for a subsequent round of selection.

[0071] The selection process typically involves a number of rounds or cycles, in which the pool of selected molecules is incrementally enriched in a specific set of sequences at the end of each round. The selection conditions may be the same for each round, or the conditions may change, for example, in order to increase the stringency of selection in later rounds. The progress of selection may be monitored by the use of isotopically-labeled amino acids, such as ³⁵S methionine. The amount of radiolabeled polypeptide bound to the target at each round is measured, and a progressive increase in recovered radiolabel is indicative of a progressive enrichment in RNA molecules encoding polypeptides with binding affinity to the target. After any round, the PCR products may be cloned and sequenced. Generally, cloning and sequencing is performed after a round in which appreciable (e.g. >2% over background to beads lacking immobilized C5) amounts of radiolabeled polypeptide are recovered in the target-bound pool. Sequences that are found in multiple isolates are candidates for encoding polypeptides that bind specifically to the target. Alternatively, high throughput sequencing of thousands of clones can

be performed after the first or subsequent rounds. Sequences that increase in frequency between, for example, the third and fourth rounds are candidates for encoding polypeptides that bind specifically to the target. The polypeptide encoded by any sequence may be translated or synthesized and tested for binding affinity to the original target protein used in the selection.

[0072] The libraries and methods of the present invention may be used to optimize the function or properties of a polypeptide. In one approach, mutagenic PCR (Keefe, A.D. and Szostak, J.W. (2001). *Nature* 15:715-718) is used to introduce sequence variation in the library once the population is enriched in polypeptides with a certain level of binding affinity.

Alternatively, a single RNA sequence encoding a polypeptide with defined binding properties can be replicated but with a defined level of mutations, or mutagenic PCR can be performed to produce a pool of mutant molecules. Upon *in vitro* translation the resulting mixture of mRNA molecules produced from such a pool is expected to encode polypeptides with a range of improved, similar, or reduced affinities as compared to the starting sequence, and a selection performed on mRNAs from such a pool may be expected to identify polypeptides with improved affinity if an appropriate stringency regimen is used during the selection.

[0073] In a second approach, optimization is performed in a directed manner. A sequence encoding a polypeptide with established binding or functional properties is subjected to site-directed mutagenesis, whereby a series of sequences is produced, with each sequence having one codon replaced with, for example, an alanine codon. The number of sequences in the set is equal to the number of amino acid residues that are to be mutated. After *in vitro* translation, the polypeptide product of each “alanine scanning” mutant is tested for binding or functional properties. Sites at which an alanine substitution affects the binding or function of the polypeptide are considered critical residues. Similarly, an N-methyl scan may be performed, such that each residue is replaced with the N-methyl derivative, and positions in the polypeptide backbone that can tolerate N-methyl substitutions can be identified.

[0074] Alternatively, the sequences can be pooled, subjected to one or more rounds of a high stringency selection, and a pool of sequences representing high affinity binding polypeptides is isolated. Critical residues are identified after DNA sequencing of the recovered DNA as those that cannot be substituted by an alanine residue without loss of activity. Once the critical residues are identified, a pool of mRNA molecules encoding a wide variety of natural (or unnatural) amino acids at each critical position is produced. The resulting pool is subjected to one or more

rounds of a high stringency selection (with the appropriate mixture of tRNAs charged with natural or unnatural amino acids), and sequences representing high affinity binding polypeptides are isolated after *in vitro* translation. In this manner, an optimal polypeptide can be identified. Since the optimal sequence may not necessarily be identified by combining optimal residues at individual sites, it is useful to test mutations at multiple sites in combination.

[0075] Both alanine and N-methyl scanning can also be performed using chemical synthesis approaches, such as solid phase polypeptide synthesis (see e.g., Coin, I et al. (2007); *Nature Protocols* 2(12):3247-56, the contents of which are herein incorporated by reference in their entirety).

[0076] Once a pool, population or subset of polypeptides is identified, they may be evaluated for therapeutic or diagnostic applications, including improved pharmacokinetic and/or pharmacodynamic properties.

[0077] In one embodiment, polypeptides are evaluated for one or more of target binding affinity, activity in biochemical or cell based assays, protease resistance, *in vitro* or *in vivo* permeability, properties related to suitability for use as a pharmaceutical agent such as plasma protein binding, metabolism (in microsomes, hepatocytes, or plasma), P-glycoprotein (Pgp) inhibition and Cytochrome P450 inhibition. Polypeptides of the present invention may also undergo testing for oral bioavailability, toxicity, human ether-a-go-go related gene product (hERG) inhibition, circulating half-life, other pharmacokinetic and pharmacodynamic parameters, and efficacy in animal models of disease.

Polypeptides of the invention

[0078] According to the present invention, once a single polypeptide or a pool of candidate polypeptide molecules is identified, they may undergo one or more rounds of structure-activity relationship (SAR) optimization using standard chemical and polypeptide synthesis techniques. Such optimization may include considerations such as avoiding charged polar side chains (Asp, Glu, Arg, Lys) that may inhibit cell penetration, avoidance of side chains that pose metabolic liabilities (Tyr, Met, Trp, Cys), improving solubility, avoidance of unnecessary molecular weight, avoidance of rotatable bonds, and altering lipophilicity.

[0079] In one embodiment, it is a goal of the present invention to provide cyclic peptidomimetics designed to be metabolically stable, cell permeable, and/or orally available.

Amino acid variants

[0080] As used herein, the term “amino acid” includes the residues of the natural amino acids as well as unnatural amino acids. The term also includes amino acids bearing a conventional amino protecting group (e.g. acetyl or benzyloxycarbonyl), as well as natural and unnatural amino acids protected at the carboxy terminus (e.g., as a (C1-C6) alkyl, phenyl or benzyl ester or amide; or as an alpha-methylbenzyl amide). Other suitable amino and carboxy protecting groups are known to those skilled in the art (See for example, Greene, T. W.; Wutz, P. G. M., Protecting Groups In Organic Synthesis; second edition, 1991, New York, John Wiley & sons, Inc., and documents cited therein). Polypeptides and/or polypeptide compositions of the present invention may also include modified amino acids.

[0081] Unnatural amino acids useful for the optimization of polypeptides and/or polypeptide compositions of the present invention include, but are not limited to 1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid, 1-amino-2,3-hydro-1H-indene-1-carboxylic acid, homolysine, homoarginine, homoserine, 2-amino adipic acid, 3-amino adipic acid, beta-alanine, aminopropionic acid, 2-aminobutyric acid, 4-aminobutyric acid, 5-aminopentanoic acid, 5-aminohexanoic acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisobutyric acid, 2-aminopimelic acid, desmosine, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, homoproline, hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, allo-isoleucine, N-methylpentylglycine, naphthylalanine, ornithine, pentylglycine, thioproline, norvaline, tert-butylglycine, phenylglycine, azatryptophan, 5-azatryptophan, 7-azatryptophan, 4-fluorophenylalanine, penicillamine, sarcosine, homocysteine, 1-aminocyclopropanecarboxylic acid, 1-aminocyclobutanecarboxylic acid, 1-aminocyclopentanecarboxylic acid, 1-aminocyclohexanecarboxylic acid, 4-aminotetrahydro-2H-pyran-4-carboxylic acid, (S)-2-amino-3-(1H-tetrazol-5-yl)propanoic acid, cyclopentylglycine, cyclohexylglycine, cyclopropylglycine, η - ω -methyl-arginine, 4-chlorophenylalanine, 3-chlorotyrosine, 3-fluorotyrosine, 5-fluorotryptophan, 5-chlorotryptophan, citrulline, 4-chloro-homophenylalanine, homophenylalanine, 4-aminomethyl-phenylalanine, 3-aminomethyl-phenylalanine, octylglycine, norleucine, tranexamic acid, 2-amino pentanoic acid, 2-amino hexanoic acid, 2-amino heptanoic acid, 2-amino octanoic acid, 2-amino nonanoic acid, 2-amino decanoic acid, 2-amino undecanoic

acid, 2-amino dodecanoic acid, aminovaleric acid, and 2-(2-aminoethoxy)acetic acid, pipecolic acid, 2-carboxy azetidine, hexafluoroleucine, 3-Fluorovaline, 2-amino-4,4-difluoro-3-methylbutanoic acid, 3-fluoro-isoleucine, 4-fluoroisoleucine, 5-fluoroisoleucine, 4-methyl-phenylglycine, 4-ethyl-phenylglycine, 4-isopropyl-phenylglycine, (S)-2-amino-5-azidopentanoic acid (also referred to herein as “X02”), (S)-2-aminohept-6-enoic acid (also referred to herein as “X30”), (S)-2-aminopent-4-ynoic acid (also referred to herein as “X31”), (S)-2-aminopent-4-enoic acid (also referred to herein as “X12”), (S)-2-amino-5-(3-methylguanidino) pentanoic acid, (S)-2-amino-3-(4-(aminomethyl)phenyl)propanoic acid, (S)-2-amino-3-(3-(aminomethyl)phenyl)propanoic acid, (S)-2-amino-4-(2-aminobenzo[*d*]oxazol-5-yl)butanoic acid, (S)-leucinol, (S)-valinol, (S)-tert-leucinol, (*R*)-3-methylbutan-2-amine, (S)-2-methyl-1-phenylpropan-1-amine, and (S)-*N*,2-dimethyl-1-(pyridin-2-yl)propan-1-amine, (S)-2-amino-3-(oxazol-2-yl)propanoic acid, (S)-2-amino-3-(oxazol-5-yl)propanoic acid, (S)-2-amino-3-(1,3,4-oxadiazol-2-yl)propanoic acid, (S)-2-amino-3-(1,2,4-oxadiazol-3-yl)propanoic acid, (S)-2-amino-3-(5-fluoro-1*H*-indazol-3-yl)propanoic acid, and (S)-2-amino-3-(1*H*-indazol-3-yl)propanoic acid, (S)-2-amino-3-(oxazol-2-yl)butanoic acid, (S)-2-amino-3-(oxazol-5-yl) butanoic acid, (S)-2-amino-3-(1,3,4-oxadiazol-2-yl) butanoic acid, (S)-2-amino-3-(1,2,4-oxadiazol-3-yl) butanoic acid, (S)-2-amino-3-(5-fluoro-1*H*-indazol-3-yl) butanoic acid, and (S)-2-amino-3-(1*H*-indazol-3-yl) butanoic acid, 2-(2'MeOphenyl)-2-amino acetic acid, tetrahydro 3-isoquinolinecarboxylic acid and stereoisomers thereof (including, but not limited, to D and L isomers).

[0082] Additional unnatural amino acids that are useful in the optimization of polypeptides or polypeptide compositions of the invention include but are not limited to fluorinated amino acids wherein one or more carbon bound hydrogen atoms are replaced by fluorine. The number of fluorine atoms included can range from 1 up to and including all of the hydrogen atoms.

Examples of such amino acids include but are not limited to 3-fluoroproline, 3,3-difluoroproline, 4-fluoroproline, 4,4-difluoroproline, 3,4-difluoroproline, 3,3,4,4-tetrafluoroproline, 4-fluorotryptophan, 5-fluorotryptophan, 6-fluorotryptophan, 7-fluorotryptophan, and stereoisomers thereof.

[0083] Further unnatural amino acids that are useful in the optimization of polypeptides or polypeptide compositions of the invention include but are not limited to those that are disubstituted at the α -carbon. These include amino acids in which the two substituents on the α -carbon are the same, for example α -amino isobutyric acid, and 2-amino-2-ethyl butanoic acid, as

well as those where the substituents are different, for example α -methylphenylglycine and α -methylproline. Further the substituents on the α -carbon may be taken together to form a ring, for example 1-aminocyclopentanecarboxylic acid, 1- aminocyclobutanecarboxylic acid, 1-aminocyclohexanecarboxylic acid, 3-aminotetrahydrofuran-3-carboxylic acid, 3-aminotetrahydropyran-3-carboxylic acid, 4-aminotetrahydropyran-4-carboxylic acid, 3-aminopyrrolidine-3-carboxylic acid, 3-aminopiperidine-3-carboxylic acid, 4-aminopiperidin-4-carboxylic acid, and stereoisomers thereof.

[0084] Additional unnatural amino acids that are useful in the optimization of polypeptides or polypeptide compositions of the invention include but are not limited to analogs of tryptophan in which the indole ring system is replaced by another 9 or 10 membered bicyclic ring system comprising 0,1,2,3 or 4 heteroatoms independently selected from N,O, or S. Each ring system may be saturated, partially unsaturated or fully unsaturated. The ring system may be substituted by 0,1,2,3, or 4 substituents at any substitutable atom. Each substituent is independently selected from H, F, Cl, Br, CN, COOR, CONRR', oxo, OR, NRR'. Each R and R' is independently selected from H, C1-C20 alkyl, C1-C20 alkyl-O-C1-20 alkyl.

[0085] In some embodiments, analogs of tryptophan (also referred to herein as "tryptophan analogs") that are useful in the optimization of polypeptides or polypeptide compositions of the invention include, but are not limited to 5-fluorotryptophan [(5-F)W], 5-methyl-O-tryptophan [(5-MeO)W], 1-methyltryptophan [(1-Me-W) or (1-Me)W], D-tryptophan (D-Trp), azatryptophan (including, but not limited to 4-azatryptophan, 7-azatryptophan and 5-azatryptophan,) 5-chlorotryptophan, 4-fluorotryptophan, 6-fluorotryptophan, 7-fluorotryptophan, and stereoisomers thereof. Except where indicated to the contrary, the term "azatryptophan" and its abbreviation, "azaTrp," as used herein, refer to 7-azatryptophan.

[0086] Modified amino acid residues useful for the optimization of polypeptides and/or polypeptide compositions of the present invention include, but are not limited to those which are chemically blocked, reversibly or irreversibly, or chemically modified on their N-terminal amino group or their side chain groups, or chemically modified in the amide backbone, as for example, N-methylated, D (unnatural amino acids) and L (natural amino acids) stereoisomers or residues wherein the side chain functional groups are chemically modified to another functional group. For example, modified amino acids include without limitation, methionine sulfoxide; methionine sulfone; aspartic acid-(beta-methyl ester), a modified amino acid of aspartic acid; N-

ethylglycine, a modified amino acid of glycine; or alanine carboxamide, and a modified amino acid of alanine. Unnatural amino acids may be purchased from Sigma-Aldrich (St. Louis, MO), Bachem (Torrance, CA) or other suppliers. Unnatural amino acids may further include any of those listed in Table 2 of US patent publication US 2011/0172126, the contents of which are incorporated herein by reference in their entirety.

[0087] In some embodiments, the amino acid sequences of polypeptides and/or polypeptide compositions of the present invention may comprise only naturally occurring amino acids. While it is known in the art that the terms peptides, polypeptides, and/or fragments thereof imply relative size, these terms as used herein should not be considered limiting with respect to the size of the various polypeptide based molecules referred to herein and which are encompassed within this invention, unless otherwise noted. In some embodiments of the present invention, polypeptides may comprise both naturally and non-naturally occurring and/or modified amino acids or be exclusively comprised of non-naturally occurring amino acids.

Polypeptide variants

[0088] According to the present invention, any amino acid based molecule (natural or unnatural) may be termed a “polypeptide” and this term embraces “peptides”, “peptidomimetics” and “proteins.” Polypeptides are also a category of protein and are traditionally considered to range in size from about 4 to about 50 amino acids. Dipeptides, those having two amino acid residues, are a category of polypeptide as are tripeptides (polypeptides comprising three amino acids). Polypeptides larger than about 50 amino acids are generally termed “proteins.”

Polypeptide sequences may be linear or cyclic. For example, a cyclic polypeptide can be prepared or may result from the formation of disulfide bonds between two cysteine residues in a sequence. A polypeptide can be cyclized through the carboxy terminus, the amino terminus, or through any other convenient point of attachment, such as, for example, through the sulfur of a cysteine or any side-chain of an amino acid residue or other linkage including, but not limited to, a maleimide linkage, an amide linkage, an ester linkage, an ether linkage, a thiol ether linkage, a hydrazone linkage, or an acetamide linkage. In some embodiments, cyclic polypeptides are formed when a molecule acts as a bridging moiety to link two or more regions of the polypeptide.

[0089] The term “amino acid sequence variant” refers to polypeptides with some differences in their amino acid sequences as compared to a starting, reference, or native sequence. The amino acid sequence variants may possess substitutions, deletions, and/or insertions at certain positions within the amino acid sequence. Ordinarily, variants will possess at least about 70% homology to a native or starting sequence, and preferably, they will be at least about 80%, more preferably at least about 90% homologous to a native or starting sequence.

[0090] “Homology” as it applies to amino acid sequences is defined as the percentage of residues in the candidate amino acid sequence that are identical with the residues in the amino acid sequence of a second sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology. Methods and computer programs for the alignment are well known in the art. It is understood that homology depends on a calculation of percent identity but may differ in value due to gaps and penalties introduced in the calculation.

[0091] By “homologs” as it applies to amino acid sequences is meant the corresponding sequence of other species having substantial identity to a second sequence of a second species.

[0092] “Analogs” is meant to include amino acid sequence variants which differ by one or more amino acid alterations, e.g., substitutions, additions or deletions of amino acid residues that still maintain one or more of the properties of the parent or starting polypeptide.

[0093] The present invention contemplates several types of compositions that include polypeptides including variants and derivatives. These include substitutional, insertional, deletion and covalent variants and derivatives. The term “derivative” is used synonymously with the term “variant” and refers to a molecule that has been modified or changed in any way relative to a reference molecule or starting molecule.

[0094] As such, included within the scope of this invention are polypeptides containing substitutions, insertions and/or additions, deletions and covalent modifications. For example, sequence tags or amino acids, such as one or more lysines, can be added to the polypeptide sequences of the invention (e.g., at the N-terminal or C-terminal ends). Sequence tags can be used for polypeptide purification or localization. Lysines can be used to increase polypeptide solubility or to allow for site specific modifications, such as, but not limited to, biotinylation or PEGylation. In some cases, polypeptides may be desthiobiotinylated. As used herein, a polypeptide that is desthiobiotinylated may comprise a desthiobiotin (Dtb) moiety conjugated to the epsilon-amino group of a lysine residue. Such lysine residues may be C-terminal residues in

some instances. Alternatively, amino acid residues located at the carboxy and amino terminal regions of the amino acid sequence of a polypeptide may optionally be deleted, providing for truncated sequences. Certain amino acids (e.g., C-terminal or N-terminal residues) may alternatively be deleted depending on the use of the sequence, as for example, expression of the sequence as part of a larger sequence, which is soluble, or linked to a solid support.

[0095] “Substitutional variants” when referring to polypeptides are those that have at least one amino acid residue in a native or starting sequence removed and a different amino acid inserted in its place at the same position. The substitutions may be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more amino acids have been substituted in the same molecule.

[0096] As used herein the term “conservative amino acid substitution” refers to the substitution of an amino acid that is normally present in the sequence with a different amino acid of similar size, charge, or polarity. Examples of conservative substitutions include the substitution of a non-polar (hydrophobic) residue such as isoleucine, valine and leucine for another non-polar residue. Likewise, examples of conservative substitutions include the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, and between glycine and serine. Additionally, the substitution of a basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue such as aspartic acid or glutamic acid for another acidic residue are additional examples of conservative substitutions. Examples of non-conservative substitutions include the substitution of a non-polar (hydrophobic) amino acid residue such as isoleucine, valine, leucine, alanine, methionine for a polar (hydrophilic) residue such as cysteine, glutamine, glutamic acid or lysine and/or a polar residue for a non-polar residue.

[0097] “Isosteres” are one of two or more molecules that exhibit some similarity of biological properties as a result of having the same number of total or valence electrons in the same arrangement and that consist of different atoms, not necessarily the same number of atoms. There are two classes of isosteres, classical and non-classical. Classical isosteres have the same number of atoms and/or the same number of valence electrons whereas non-classical isosteres are molecules that produce a similar biological effect *in vivo* but do not have the same number of atoms and/or valence electrons.

[0098] According to the present invention, “peptide bond isosteres” are defined as isosteres having properties resembling peptide bonds. Peptide bond isosteres may be of a linear type comprising at least one peptide bond replacement or may be cyclic and comprise an amine and a carboxylic acid function. Such replacement may be with any moiety which improves the physicochemical, structural or functional properties of the molecule. Replacement of the peptide bond may increase the metabolic stability of the polypeptides and reduce or increase the flexibility. Peptide bond isosteres described herein may be mono-, di-, tri-, tetra-, penta-, sexta-, septa-, octa-, nona-, or deca-peptide bond isosteres, meaning that at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 peptidic bonds may be replaced. Non-limiting examples of linear dipeptide bond isosteres for amide (peptidic) bonds include thioamide, sulfonamide, sulfonate, phosphonamide, phosphonate phosphothioate, phosphinate, alkane, 1 or 2 hydroxyethylene, dihydroxylethylene, C-C single bond (alkane), C-C double bond (alkene), C-C triple bond (alkyne), C-O bond (methyleneoxy), O-N or N-O bond,(methylenemino), triazole, hydrazide, urea, ketone, urethane bond, (di)haloalkene, methylenemercapto, methyleneamino, trifluoroethylamino, hydrazide, amideoxy, and others known to those of skill in the art.

[0099] Peptide bond isosteres may also be cyclic molecules that are decorated with an amine and a carboxylic acid function. Non-limiting examples of cyclic peptide bond isosteres with varying ring sizes include carbacycles, azacycles and oxacycles. Azacycles may be based on an alkaloid core which forms a bicyclic structure isostere. An example of an azacyclic isostere includes an isostere based on a triazole ring formed by a copper catalyzed azide-alkyne cycloaddition. Cyclic peptide bond isosteres described herein may be bi-, tri-, tetra-, penta-, sexta-, septa-, octa- nona- deca-peptide cyclic isosteres

[00100] “Insertional variants” when referring to polypeptides are those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position in a native or starting sequence. “Immediately adjacent” to an amino acid means connected to either the alpha-carboxy or alpha-amino functional group of the amino acid.

[00101] “Deletional variants” when referring to polypeptides are those with one or more amino acids in the native or starting amino acid sequence removed. Ordinarily, deletional variants will have one or more amino acids deleted in a particular region of the molecule.

[00102] “Truncated variants” when referring to polypeptides are those with one or more amino acids in the native or starting amino acid sequence removed from either terminus of the polypeptide.

[00103] According to the present invention, polypeptides may be modified by the addition of one or more conjugate groups. In some embodiments, polypeptides may be administered in combination with one or more additional molecules.

[00104] As used herein, a “conjugate” refers to any molecule or moiety appended to another molecule. In the present invention, conjugates may be polypeptide (amino acid) based or not. Conjugates may comprise lipids, small molecules, RNA, DNA, polypeptides, polymers, or combinations thereof. Functionally, conjugates may serve as targeting molecules or may serve as payload to be delivered to a cell, organ or tissue. Conjugates are typically covalent modifications introduced by reacting targeted amino acid residues or the termini of the polypeptide with an organic derivatizing agent that is capable of reacting with selected side-chains or terminal residues. Such modifications are within the ordinary skill in the art and are performed without undue experimentation.

[00105] The conjugation process may involve PEGylation, lipidation, albumination, biotinylation, desthiobiotinylation, the addition of other polypeptide tails, or grafting onto antibody Fc domains, CDR regions of intact antibodies, or antibody domains produced by any number of means. The conjugate may include anchors including cholesterol oleate moiety, cholesteryl laurate moiety, an α -tocopherol moiety, a phytol moiety, an oleate moiety, or an unsaturated cholesterol-ester moiety or a lipophilic compound selected from acetanilides, anilides, aminoquinolines, benzhydryl compounds, benzodiazepines, benzofurans, cannabinoids, cyclic polypeptides, dibenzazepines, digitalis glycosides, ergot alkaloids, flavonoids, imidazoles, quinolines, macrolides, naphthalenes, opiates (such as, but not limited to, morphinans or other psychoactive drugs), oxazines, oxazoles, phenylalkylamines, piperidines, polycyclic aromatic hydrocarbons, pyrrolidines, pyrrolidinones, stilbenes, sulfonylureas, sulfones, triazoles, tropanes, and vinca alkaloids.

[00106] As used herein, the term “covalent derivative” when referring to a polypeptide includes modification of a native or starting polypeptide with an organic proteinaceous or non-proteinaceous derivatizing agent, and/or post-translational modification. Covalent modifications are traditionally introduced by reacting targeted amino acid residues of the polypeptide with an

organic derivatizing agent that is capable of reacting with selected side-chains or terminal residues, or by harnessing mechanisms of post-translational modifications that function in selected recombinant host cells. The resultant covalent derivatives are useful in programs directed at identifying residues important for biological activity, for immunoassays, or for the preparation of anti-protein antibodies for immunoaffinity purification of the recombinant protein. Such modifications are within the ordinary skill in the art and are performed without undue experimentation.

[00107] Certain post-translational modifications are the result of the action of recombinant host cells on an expressed polypeptide. Glutaminyl and asparaginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues may be present in the polypeptides produced in accordance with the present invention.

[00108] Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of tyrosinyl, seryl or threonyl residues, methylation of the alpha-amino groups of lysine, arginine, and histidine side chains (Creighton, T. E., *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, 1983, pp. 79-86).

[00109] Covalent modifications specifically include the bonding of non-proteinaceous polymers to polypeptides of the invention. Non-proteinaceous polymers may include a hydrophilic synthetic polymer, i.e., a polymer not otherwise found in nature. However, polymers that exist in nature and are produced by recombinant or *in vitro* methods are useful, as are polymers which are isolated from nature. Hydrophilic polyvinyl polymers fall within the scope of this invention, e.g. polyvinylalcohol and polyvinylpyrrolidone. The polypeptides may be linked to various non-proteinaceous polymers, such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in U.S. Pat. No. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

[00110] “Features” when referring to polypeptides are defined as distinct amino acid sequence-based components of a molecule. Features of the polypeptide of the present invention include surface manifestations, local conformational shape, folds, loops, half-loops, domains, half-domains, sites, termini or any combination thereof.

[00111] As used herein when referring to polypeptides the term “fold” refers to the resultant conformation of an amino acid sequence upon energy minimization. A fold may occur at the

secondary or tertiary level of the folding process. Examples of secondary level folds include beta sheets and alpha helices. Examples of tertiary folds include domains and regions formed due to aggregation or separation of physicochemically distinct regions. Regions formed in this way include hydrophobic and hydrophilic pockets, and the like.

[00112] As used herein the term “turn” as it relates to protein conformation means a bend which alters the direction of the backbone of a polypeptide and may involve one, two, three or more amino acid residues.

[00113] As used herein when referring to polypeptides the term “loop” refers to a structural feature of a polypeptide which may serve to reverse the direction of the backbone of a polypeptide. Where the loop is found in a polypeptide and only alters the direction of the backbone, it may comprise four or more amino acid residues. Oliva et al. have identified at least 5 classes of protein loops (Oliva, B. et al., J Mol Biol. 1997 Mar 7;266(4):814-30; the contents of which are herein incorporated by reference in their entirety). Loops may be open or closed.

Closed loops or “cyclic” loops may be formed when two amino acids are connected by a bridging moiety. The cyclic loop comprises the amino acids along the polypeptide present between the bridged amino acids. Cyclic loops may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acids.

[00114] As used herein when referring to polypeptides the term “half-loop” refers to a portion of an identified loop having at least half the number of amino acid residues as the loop from which it is derived. It is understood that loops may not always contain an even number of amino acid residues. Therefore, in those cases where a loop contains or is identified to comprise an odd number of amino acids, a half-loop of the odd-numbered loop will comprise the whole number portion or next whole number portion of the loop (number of amino acids of the loop/2+/-0.5 amino acids). For example, a loop identified as a 7 amino acid loop could produce half-loops of 3 amino acids or 4 amino acids (7/2=3.5+/-0.5 being 3 or 4).

[00115] As used herein when referring to proteins, the term “region” refers to a zone or general area. In some embodiments, when referring to a protein, a region may comprise a linear sequence of amino acids along the protein or may comprise a specific secondary or tertiary structure and/or one or more features or protein domains.

[00116] As used herein, the term “domain,” when referring to proteins, refers to a motif of a polypeptide having one or more identifiable structural (such as secondary or tertiary structures)

or functional characteristics or properties (e.g., binding capacity, serving as a site for protein-protein interactions.)

[00117] As used herein, the term “half-domain,” when referring to proteins, refers to a portion of an identified domain having at least half the number of amino acid residues as the domain from which it is derived. It is understood that domains may not always contain an even number of amino acid residues. Therefore, in those cases where a domain contains or is identified to comprise an odd number of amino acids, a half-domain of the odd-numbered domain will comprise the whole number portion or next whole number portion of the domain (number of amino acids of the domain/2+/-0.5 amino acids). For example, a domain identified as a 7 amino acid domain could produce half-domains of 3 amino acids or 4 amino acids (7/2=3.5+/-0.5 being 3 or 4). It is also understood that sub-domains may be identified within domains or half-domains, these subdomains possessing less than all of the structural or functional properties identified in the domains or half domains from which they were derived. It is also understood that the amino acids that comprise any of the domain types herein need not be contiguous along the backbone of the polypeptide (i.e., nonadjacent amino acids may fold structurally to produce a domain, half-domain or subdomain).

[00118] As used herein when referring to polypeptides the term “site” as it pertains to amino acid based embodiments is used synonymously with “amino acid residue” and “amino acid side chain.” A site represents a position within a polypeptide that may be modified, manipulated, altered, derivatized or varied within the polypeptide based molecules of the present invention.

[00119] As used herein the terms “termini” or “terminus” when referring to polypeptides refers to an extremity of a polypeptide. Such extremity is not limited only to the first or final site of the polypeptide but may include additional amino acids in the terminal regions. The polypeptide based molecules of the present invention may be characterized as having both an N-terminus and a C-terminus. Polypeptides and/or polypeptide compositions of the present invention are in some cases made up of multiple polypeptide chains brought together by disulfide bonds or by non-covalent forces (multimers, oligomers). These sorts of proteins will have multiple N- and C-termini. Alternatively, the termini of the polypeptides may be modified such that they begin or end, as the case may be, with a non-polypeptide based moiety such as an organic conjugate.

[00120] In one embodiment, polypeptides of the present invention may include a terminal region. As used herein, “terminal region” is a terminal region of amino acids that may include a

cysteine. The terminal region may be an N- and/or a C-terminal region. In some embodiments, terminal regions may be connected to the parent polypeptides using a bridging moiety. As used herein, “parent polypeptide” refers to the part of the polypeptide that does not include the terminal region. The terminal region may be separated from the parent polypeptide by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more residues. The residues added may be selected from, but are not limited to, any natural or unnatural amino acid, the N-methylated form of any natural or unnatural amino acid, the D-stereoisomer of any natural or unnatural amino acid, norvaline, tert-butylglycine, phenylglycine, azatryptophan, 7-azatryptophan, 4-fluorophenylalanine, penicillamine, sarcosine, homocysteine, 1-aminocyclopropanecarboxylic acid, 1-aminocyclobutanecarboxylic acid, 1-aminocyclopentanecarboxylic acid, 1-aminocyclohexanecarboxylic acid, 4-aminotetrahydro-2H-pyran-4-carboxylic acid, aminoisobutyric acid, (S)-2-amino-3-(1*H*-tetrazol-5-yl)propanoic acid, cyclopentylglycine, cyclohexylglycine, cyclopropylglycine, η - ω -methyl-arginine, 4-chlorophenylalanine, 3-chlorotyrosine, 3-fluorotyrosine, 5-fluorotryptophan, 5-chlorotryptophan, citrulline, 4-chloro-homophenylalanine, homophenylalanine, 4-aminomethyl-phenylalanine, 3-aminomethyl-phenylalanine, octylglycine, norleucine, tranexamic acid, 2-amino pentanoic acid, 2-amino hexanoic acid, 2-amino heptanoic acid, 2-amino octanoic acid, 2-amino nonanoic acid, 2-amino decanoic acid, 2-amino undecanoic acid, 2-amino dodecanoic acid, aminovaleric acid, and 2-(2-aminoethoxy)acetic acid, pipecolic acid, 2-carboxy azetidine, hexafluoroleucine, 3-Fluorovaline, 2-amino-4,4-difluoro-3-methylbutanoic acid, 3-fluoro-isoleucine, 4-fluoroisoleucine, 5-fluoroisoleucine, 4-methyl-phenylglycine, 4-ethyl-phenylglycine, 4-isopropyl-phenylglycine, (S)-2-amino-5-(3-methylguanidino) pentanoic acid, (S)-2-amino-3-(4-(aminomethyl)phenyl)propanoic acid, (S)-2-amino-3-(3-(aminomethyl)phenyl)propanoic acid, (S)-2-amino-4-(2-aminobenzo[*d*]oxazol-5-yl)butanoic acid, (S)-leucinol, (S)-valinol, (S)-tert-leucinol, (R)-3-methylbutan-2-amine, (S)-2-methyl-1-phenylpropan-1-amine, and (S)-*N*,2-dimethyl-1-(pyridin-2-yl)propan-1-amine, (S)-2-amino-3-(oxazol-2-yl)propanoic acid, (S)-2-amino-3-(oxazol-5-yl)propanoic acid, (S)-2-amino-3-(1,3,4-oxadiazol-2-yl)propanoic acid, (S)-2-amino-3-(1,2,4-oxadiazol-3-yl)propanoic acid, (S)-2-amino-3-(5-fluoro-1*H*-indazol-3-yl)propanoic acid, and (S)-2-amino-3-(1*H*-indazol-3-yl)propanoic acid.

[00121] Additional unnatural amino acids that are useful in the optimization of polypeptides of the invention include but are not limited to fluorinated amino acids wherein one or more carbon bound hydrogen atoms are replaced by fluorine. The number of fluorine atoms included can

range from 1 up to and including all of the hydrogen atoms. Examples of such amino acids include but are not limited to 3-fluoroproline, 3,3-difluoroproline, 4-fluoroproline, 4,4-difluoroproline, 3,4-difluoroproline, 3,3,4,4-tetrafluoroproline, 4-fluorotryptophan, 6-fluorotryptophan, 7-fluorotryptophan, and stereoisomers thereof.

[00122] Further unnatural amino acids that are useful in the optimization of polypeptides of the invention include but are not limited to those that are disubstituted at the α -carbon. These include amino acids in which the two substituents on the α -carbon are the same, for example α -amino isobutyric acid, and 2-amino-2-ethyl butanoic acid, as well as those where the substituents are different, for example α -methylphenylglycine and α -methylproline. Further the substituents on the α -carbon may be taken together to form a ring, for example 1-aminocyclopentanecarboxylic acid, 1-aminocyclobutanecarboxylic acid, 1-aminocyclohexanecarboxylic acid, 3-aminotetrahydrofuran-3-carboxylic acid, 3-aminotetrahydropyran-3-carboxylic acid, 4-aminotetrahydropyran-4-carboxylic acid, 3-aminopyrrolidine-3-carboxylic acid, 3-aminopiperidine-3-carboxylic acid, 4-aminopiperidin-4-carboxylic acid, and stereoisomers thereof.

[00123] Additional unnatural amino acids that are useful in the optimization of polypeptides of the invention include but are not limited to analogs of tryptophan in which the indole ring system is replaced by another 9 or 10 membered bicyclic ring system comprising 0,1,2,3, or 4 heteroatoms independently selected from N, O, or S. Each ring system may be saturated, partially unsaturated or fully unsaturated. The ring system may be substituted by 0,1,2,3, or 4 substituents at any substitutable atom. Each substituent is independently selected from H, F, Cl, Br, CN, oxo, COOR, CONRR', OR, NRR'. Each R and R' is independently selected from H, C1-C20 alkyl, C1-C20 alkyl-O-C1-20 alkyl.

[00124] In some embodiments, analogs of tryptophan (also referred to herein as "tryptophan analogs") that are useful in the optimization of polypeptides of the invention include, but are not limited to 5-fluorotryptophan [(5-F)W], 5-methyl-O-tryptophan [(5-MeO)W], 1-methyltryptophan [(1-Me-W) or (1-Me)W], D-tryptophan (D-Trp), 7-azatryptophan (including, but not limited to 4-azatryptophan, 7-azatryptophan and 5-azatryptophan,) 5-chlorotryptophan, 4-fluorotryptophan, 6-fluorotryptophan, 7-fluorotryptophan, and stereoisomers thereof. Except where indicated to the contrary, the term "azatryptophan" and its abbreviation, "azaTrp," as used herein, refer to 7-azatryptophan.

[00125] In one embodiment, polypeptides of the present invention may include a terminal modification at the N- or C-termini with the addition of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more residues and/or a cysteine in the terminal region. The residues added may be selected from, but are not limited to, any natural or unnatural amino acid, the N-methylated form of any natural or unnatural amino acid, the D-stereoisomer of any amino acid, norvaline, tert-butylglycine, phenylglycine, azatryptophan, 7-azatryptophan, 4-fluorophenylalanine, penicillamine, sarcosine, homocysteine, 1-aminocyclopropanecarboxylic acid, 1-aminocyclobutanecarboxylic acid, 1-aminocyclopentanecarboxylic acid, 1-aminocyclohexanecarboxylic acid, 4-aminotetrahydro-2H-pyran-4-carboxylic acid, aminoisobutyric acid, (S)-2-amino-3-(1*H*-tetrazol-5-yl)propanoic acid, cyclopentylglycine, cyclohexylglycine, cyclopropylglycine, η - ω -methyl-arginine, 4-chlorophenylalanine, 3-chlorotyrosine, 3-fluorotyrosine, 5-fluorotryptophan, 5-chlorotryptophan, citrulline, 4-chloro-homophenylalanine, homophenylalanine, 4-aminomethyl-phenylalanine, 3-aminomethyl-phenylalanine, octylglycine, norleucine, tranexamic acid, 2-amino pentanoic acid, 2-amino hexanoic acid, 2-amino heptanoic acid, 2-amino octanoic acid, 2-amino nonanoic acid, 2-amino decanoic acid, 2-amino undecanoic acid, 2-amino dodecanoic acid, aminovaleric acid, and 2-(2-aminoethoxy)acetic acid, pipecolic acid, 2-carboxy azetidine, hexafluoroleucine, 3-Fluorovaline, 2-amino-4,4-difluoro-3-methylbutanoic acid, 3-fluoro-isoleucine, 4-fluoroisoleucine, 5-fluoroisoleucine, 4-methyl-phenylglycine, 4-ethyl-phenylglycine, 4-isopropyl-phenylglycine, (S)-2-amino-5-(3-methylguanidino) pentanoic acid, (S)-2-amino-3-(4-(aminomethyl)phenyl)propanoic acid, (S)-2-amino-3-(3-(aminomethyl)phenyl)propanoic acid, (S)-2-amino-4-(2-aminobenzo[*d*]oxazol-5-yl)butanoic acid, (S)-leucinol, (S)-valinol, (S)-tert-leucinol, (R)-3-methylbutan-2-amine, (S)-2-methyl-1-phenylpropan-1-amine, and (S)-*N*,2-dimethyl-1-(pyridin-2-yl)propan-1-amine, (S)-2-amino-3-(oxazol-2-yl)propanoic acid, (S)-2-amino-3-(oxazol-5-yl)propanoic acid, (S)-2-amino-3-(1,3,4-oxadiazol-2-yl)propanoic acid, (S)-2-amino-3-(1,2,4-oxadiazol-3-yl)propanoic acid, (S)-2-amino-3-(5-fluoro-1*H*-indazol-3-yl)propanoic acid, and (S)-2-amino-3-(1*H*-indazol-3-yl)propanoic acid.

[00126] Polypeptides of the present invention may be conjugated to a polypeptide that increases or decreases plasma protein binding including but not limited to those described in Dennis, M.S. et al., Albumin binding as a general strategy for improving the pharmacokinetics of proteins. *J Biol Chem.* 2002 Sep 20; 277(38):35035-43; Nguyen, A. et al., The pharmacokinetics of an albumin-binding Fab (AB Fab) can be modulated as a function of affinity for albumin.

Protein Eng Des Sel. 2006 Jul;19(7):291-7 and Langerheim, J.F. et al., Improving the pharmacokinetics/pharmacodynamics of prolactin, GH, and their antagonists by fusion to a synthetic albumin-binding polypeptide. J Endocrinol. 2009 Dec;203(3):375-87. In some embodiments, such polypeptides bind serum albumin (referred to herein as “albumin-binding polypeptides”). In some embodiments, albumin-binding polypeptides are cyclized by disulfide bond formation between cysteine residues present in their polypeptide sequences. In some embodiments, albumin-binding polypeptides are conjugated by either their N or C-terminal ends. In some embodiments, conjugation to an albumin-binding polypeptide modulates the amount of time that a polypeptide of the present invention remains intact in a subject. In a preferred embodiment, conjugation to an albumin-binding polypeptide increases the amount of time that a polypeptide of the present invention remains in the blood of a subject. Polypeptides of the present invention may be conjugated to polypeptides that have cell penetrating properties (referred to herein as “cell penetrating polypeptides”) including but not limited to those disclosed in Milletti, F., Cell-penetrating peptides: classes, origin, and current landscape. Drug Discov Today. 2012 Aug;17(15-16):850-60. Additional cell penetrating polypeptides are known to those skilled in the art. Polypeptides of the present invention may be conjugated to any of the polypeptide conjugates taught, for example, in US patent publications US20110172126 or US20030040472 the contents of which are incorporated herein by reference in their entirety. Polypeptides of the present invention may be conjugated to a lipophilic molecule that increases plasma protein binding such as the lipophilic substituents taught, for example, in US Patent No. 6,268,343 or US Publication No. US2013/0053311, the contents of each of which are herein incorporated by reference in their entirety.

[00127] Once any of the features have been identified or defined as a desired component of a polypeptide, any of several manipulations and/or modifications of these features may be performed by moving, swapping, inverting, deleting, randomizing or duplicating. Furthermore, it is understood that manipulation of features may result in the same outcome as a modification to the molecules of the invention. For example, a manipulation which involved deleting a domain would result in the alteration of the length of a molecule just as modification of a nucleic acid to encode less than a full length molecule would.

[00128] Modifications and manipulations can be accomplished by methods known in the art such as, but not limited to, site directed mutagenesis. The resulting modified molecules may then

be tested for activity using *in vitro* or *in vivo* assays such as those described herein or any other suitable screening assay known in the art.

[00129] According to the present invention, the polypeptides may comprise a consensus sequence which is discovered through rounds of experimentation. As used herein a “consensus” sequence is a single sequence which represents a collective population of sequences allowing for variability at one or more sites.

[00130] The term “identity” as known in the art, refers to a relationship between the sequences of two or more polypeptides, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between polypeptides, as determined by the number of matches between strings of two or more amino acid residues. Identity measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., “algorithms”). Identity of related polypeptides can be readily calculated by known methods. Such methods include, but are not limited to, those described previously by others (Lesk, A. M., ed., Computational Molecular Biology, Oxford University Press, New York, 1988; Smith, D. W., ed., Biocomputing: Informatics and Genome Projects, Academic Press, New York, 1993; Griffin, A. M. et al., ed., Computer Analysis of Sequence Data, Part 1, Humana Press, New Jersey, 1994; von Heinje, G., Sequence Analysis in Molecular Biology, Academic Press, 1987; Gribskov, M. et al., ed., Sequence Analysis Primer, M. Stockton Press, New York, 1991; and Carillo et al., Applied Math, SIAM J, 1988, 48, 1073).

[00131] In some embodiments, a polypeptide variant may have the same or a similar activity as the reference polypeptide. Alternatively, a variant may have an altered activity (e.g., increased or decreased) relative to a reference polypeptide. Generally, variants of a particular polypeptide of the invention will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% but less than 100% sequence identity to that of a particular reference polypeptide as determined by sequence alignment programs and parameters described herein and known to those skilled in the art. Such tools for alignment include those of the BLAST suite (Altschul, S.F. et al., Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, Nucleic Acids Res. 1997, 25:3389-3402) Other tools are described herein, specifically in the definition of “identity.”


[00132] Default parameters in the BLAST algorithm include, for example, an expected threshold of 10, Word size of 28, Match/Mismatch Scores 1, -2, Gap costs Linear. Any filter can be applied as well as a selection for species specific repeats, e.g., *Homo sapiens*.

Abbreviations used in polypeptides

[00133] As used herein, abbreviations have the following meaning: “Ac” and “NH2” indicate acetyl and amidated termini, respectively; “Nvl” stands for norvaline; “Phg” stands for phenylglycine; “Tbg” stands for tert-butylglycine; “Chg” stands for cyclohexylglycine; “(N-Me)X” stands for the N-methylated form of the amino acid indicated by the letter or three letter amino acid code in place of variable “X” written as N-methyl-X [e.g. (N-Me)A or (N-Me)Ala stand for the N-methylated form of alanine or N-methyl-alanine]; “azaTrp” stands for azatryptophan; “(4-F)Phe” stands for 4-fluorophenylalanine; “Tyr(OMe)” stands for O-methyl tyrosine, “Aib” stands for amino isobutyric acid; “(homo)F” or “(homo)Phe” stands for homophenylalanine; “(2-OMe)Phg” refers to 2-O-methylphenylglycine; “(5-F)W” refers to 5-fluorotryptophan; “D-X” refers to the D-stereoisomer of the given amino acid “X” [e.g. (D-Chg) stands for D-cyclohexylglycine]; “(5-MeO)W” refers to 5-methyl-O-tryptophan; “homoC” refers to homocysteine; “(1-Me-W)” or “(1-Me)W” refers to 1-methyltryptophan; “Nle” refers to norleucine; “Tiq” refers to a tetrahydroisoquinoline residue; “Asp(T)” refers to (S)-2-amino-3-(1*H*-tetrazol-5-yl)propanoic acid; “(3-Cl-Phe)” refers to 3-chlorophenylalanine; “[N-Me-4-F)Phe]” or “[N-Me-4-F)Phe]” refers to N-methyl-4-fluorophenylalanine; “[m-Cl-homo)Phe]” refers to meta-chloro homophenylalanine; “[des-amino)C]” refers to 3-thiopropionic acid; “[alpha-methyl)D]” refers to alpha-methyl L-aspartic acid; “[2Nal]” refers to 2-naphthylalanine; “[3-aminomethyl)Phe]” refers to 3-aminomethyl-L-phenylalanine; “[Cle]” refers to cycloleucine; “[Ac-Pyran]” refers to 4-amino-tetrahydro-pyran-4-carboxylic acid; “[Lys-C16]” refers to N- ϵ -palmitoyl lysine; “[Lys-C12]” refers to N- ϵ -lauryl lysine; “[Lys-C10]” refers to N- ϵ -capryl lysine; “[Lys-C8]” refers to N- ϵ -caprylic lysine; “[xXyl(y, z)]” refers to the xyl bridging moiety between two thiol containing amino acids where x may be m, p or o to indicate the use of meta-, para- or ortho- dibromoxylene (respectively) to generate bridging moieties and the numerical identifiers, y and z, place the amino acid position within the polypeptide of the amino acids participating in the cyclization; “[cyclo(y, z)]” refers to the formation of a bond between two amino acid residues where the numerical identifiers, y and z, place the position of the


residues participating in the bond; “[cyclo-olefinyl(y,z)]” refers to the formation of a bond between two amino acid residues by olefin metathesis where the numerical identifiers, y and z , place the position of the residues participating in the bond; “[cyclo-thioalkyl(y,z)]” refers to the formation of a thioether bond between two amino acid residues where the numerical identifiers, y and z , place the position of the residues participating in the bond; “[cyclo-triazolyl(y,z)]” refers to the formation of a triazole ring between two amino acid residues where the numerical identifiers, y and z , place the position of the residues participating in the bond. “B20” refers to N- ϵ -(PEG2- γ -glutamic acid-N- α -octadecanedioic acid) lysine [also known as (1*S*,28*S*)-1-amino-7,16,25,30-tetraoxo-9,12,18,21-tetraoxa-6,15,24,29-tetraazahexatetracontane-1,28,46-tricarboxylic acid.]

B20



“B28” refers to N- ϵ -(PEG24- γ -glutamic acid-N- α -hexadecanoyl)lysine.

B28



“K14” refers to N- ϵ -1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)-3-methylbutyl-L-lysine. All other symbols refer to the standard one-letter amino acid code.

Antibodies

[00134] In some embodiments, compounds and/or compositions of the present invention may comprise antibodies or fragments thereof. As used herein, the term “antibody” is referred to in the broadest sense and specifically covers various embodiments including, but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies

formed from at least two intact antibodies), and antibody fragments such as diabodies so long as they exhibit a desired biological activity. Antibodies of the present invention may also comprise human antibodies or humanized antibodies. Antibodies are primarily amino-acid based molecules but may also comprise one or more modifications (including, but not limited to the addition of sugar moieties, fluorescent moieties, chemical tags, etc.)

[00135] As used herein the term, “antibody fragment” refers to any portion of an intact antibody. In some embodiments, antibody fragments comprise antigen binding regions from intact antibodies. Examples of antibody fragments may include, but are not limited to Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments. Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, each with a single antigen-binding site. Also produced is a residual “Fc” fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen. Compounds and/or compositions of the present invention may comprise one or more of these fragments. For the purposes herein, an “antibody” may comprise a heavy and light variable domain as well as an Fc region.

[00136] As used herein, the term “native antibody” refers to a usually heterotetrameric glycoprotein of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain.

[00137] As used herein, the term “variable domain” refers to specific antibody domains that differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen.

As used herein, the term “Fv” refers to antibody fragments comprising complete antigen-

recognition and antigen-binding sites. These regions consist of a dimer of one heavy chain and one light chain variable domain in tight, non-covalent association.

[00138] As used herein, the term “light chain” refers to a component of an antibody from any vertebrate species assigned to one of two clearly distinct types, called kappa and lambda based on amino acid sequences of constant domains.

[00139] Depending on the amino acid sequence of the constant domain of their heavy chains, antibodies can be assigned to different classes. There are five major classes of intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

[00140] As used herein, the term “Single-chain Fv” or “scFv” refers to a fusion protein of V_H and V_L antibody domains, wherein these domains are linked together into a single polypeptide chain. In some embodiments, the Fv polypeptide linker enables the scFv to form the desired structure for antigen binding.

[00141] As used herein, the term “diabody” refers to a small antibody fragment with two antigen-binding sites. Diabodies comprise a heavy chain variable domain V_H connected to a light chain variable domain V_L in the same polypeptide chain. By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al. (Hollinger, P. et al., “Diabodies”:Small bivalent and bispecific antibody fragments. PNAS. 1993. 90:6444-8) the contents of each of which are incorporated herein by reference in their entirety.

[00142] As used herein, the term “monoclonal antibody” refers to an antibody obtained from a population of substantially homogeneous cells (or clones), i.e., the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variants that may arise during production of the monoclonal antibodies, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen

[00143] The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. The monoclonal antibodies herein

include “chimeric” antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies.

[00144] As used herein, the term “humanized antibody” refers to a chimeric antibody comprising a minimal portion from one or more non-human (e.g., murine) antibody source with the remainder derived from one or more human immunoglobulin sources. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from the hypervariable region from an antibody of the recipient are replaced by residues from the hypervariable region from an antibody of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and/or capacity.

[00145] As used herein, the term “hypervariable region” refers to regions within the antigen binding domain of an antibody comprising amino acid residues responsible for antigen binding. The amino acids present within the hypervariable regions determine the structure of the complementarity determining region (CDR). As used herein, the term “CDR” refers to regions of antibodies comprising a structure that is complimentary to its target antigen or epitope.

[00146] In some embodiments, compounds and/or compositions of the present invention may be or comprise antibody mimetics. As used herein, the term “antibody mimetic” refers to any molecule which mimics the function or effect of an antibody and which binds specifically and with high affinity to their molecular targets. In some embodiments, antibody mimetics may be monobodies, designed to incorporate the fibronectin type III domain (Fn3) as a protein scaffold (US 6,673,901 and US 6,348,584, the contents of each of which are herein incorporated by reference in their entirety). In some embodiments, antibody mimetics may include those known in the art including, but not limited to affibody molecules, affilins, affitins, anticalins, avimers, Centyrins, DARPINSTM, Fynomers, Adnectins, and Kunitz domain peptides. In other embodiments, antibody mimetics may include one or more non-peptide region.

[00147] As used herein, the term “antibody variant” refers to a biomolecule resembling an antibody in structure and/or function comprising some differences in their amino acid sequence, composition or structure as compared to a native antibody.

[00148] The preparation of antibodies, whether monoclonal or polyclonal, is known in the art. Techniques for the production of antibodies are well known in the art and described, e.g. in Harlow and Lane “Antibodies, A Laboratory Manual”, Cold Spring Harbor Laboratory Press, 1988 and Harlow and Lane “Using Antibodies: A Laboratory Manual” Cold Spring Harbor Laboratory Press, 1999.

[00149] In some embodiments, polypeptide sequences provided herein may be utilized in the production of one or more antibodies. In some cases, such polypeptide sequences may be incorporated into antibody variable domains. Such variable domains may be incorporated into antibodies, antibody mimetics or antibody variants.

Small molecules

[00150] In some embodiments, compounds of the present invention may be small molecules. Such compounds may comprise a size from about 100 to about 2000 Daltons (e.g. from about 100 to about 200, to about 300, to about 400, to about 500, to about 600, to about 700, to about 800, to about 900, to about 1000, to about 1100, to about 1200, to about 1300, to about 1400, to about 1500, to about 1600, to about 1700, to about 1800, to about 1900 or to about 2000 Daltons.) Small molecules may be non-peptidic or share some or many characteristics of polypeptides and cyclic polypeptides, including amide bonds, cyclic structures, and amino acid-like substituents.

Aptamers

[00151] In some embodiments, compounds of the present invention may comprise aptamers (Keefe, A.D., Pai, S. and Ellington, A. (2010). *Nat. Rev. Drug Discovery* **9**:537-550). As used herein, the term “aptamer” refers to oligonucleic or polypeptide molecules that are capable of binding specific target molecules. Some aptamers may adopt a three-dimensional conformation capable of binding such target molecules with high affinity and specificity.

Isotopic variations

[00152] Polypeptides of the present invention may comprise one or more atoms that are isotopes. As used herein, the term “isotope” refers to a chemical element that has one or more additional neutrons. In one embodiment, polypeptides of the present invention may be deuterated. As used herein, the term “deuterated” refers to a substance that has had one or more

hydrogen atoms replaced by deuterium isotopes. Deuterium isotopes are isotopes of hydrogen. The nucleus of hydrogen contains one proton while deuterium nuclei contain both a proton and a neutron. Compounds and pharmaceutical compositions of the present invention may be deuterated in order to change a physical property, such as stability, or to allow them to be used in diagnostic and experimental applications.

Formulation and delivery

[00153] The term “pharmaceutical composition” refers to a composition comprising at least one active ingredient (e.g., such as a polypeptide) in a form and amount that permits the active ingredient to be therapeutically effective.

[00154] Polypeptide formulations of the present invention include controlled duodenal release formulations, time release formulations, osmotic-controlled release delivery systems, microemulsions, microspheres, liposomes, nanoparticles, patches, pumps, drug depots, and the like. Specifically included in the present invention are solid oral dosage forms, such as powders, softgels, gelcaps, capsules, pills, and tablets.

[00155] The pharmaceutical compositions of the present invention may be administered by any route that results in a therapeutically effective outcome. These include, but are not limited to enteral, gastroenteral, epidural, oral, peridural, intracerebral (into the cerebrum), intratracheal (into the airways for delivery to the lung), intracerebroventricular (into the cerebral ventricles), epicutaneous (application onto the skin), intradermal, (into the skin itself), subcutaneous (under the skin), nasal administration (through the nose), intravenous (into a vein), intraarterial (into an artery), intramuscular (into a muscle), intracardiac (into the heart), intraosseous infusion (into the bone marrow), intrathecal (into the spinal canal), intraperitoneal, (infusion or injection into the peritoneum), intravesical infusion, intravitreal, (into the posterior chamber of the eye), intracavernous injection, (into the base of the penis), intravaginal administration, intrauterine, extra-amniotic administration, transdermal (diffusion through the intact skin for systemic distribution), transmucosal (diffusion through a mucous membrane), insufflation (snorting), buccal, sublingual, sublabial, enema, eye drops (onto the conjunctiva), or in ear drops.

[00156] In some embodiments, polypeptides of the present invention are formulated in a sterile aqueous solution. In some embodiments, polypeptides of the present invention are formulated in a lipid or non-lipid formulation. In another embodiment, polypeptides of the present invention

are formulated in a cationic or non-cationic lipid formulation. In either embodiment, the sterile aqueous solution may contain additional active or inactive components. Inactive components, also referred to herein as “excipients,” can include, but are not limited to, physiologically compatible salts, sugars, bulking agents, surfactants, or buffers.

[00157] Polypeptides and/or polypeptide compositions of the present invention may comprise or be formulated or delivered in conjunction with one or more carrier agents. As used herein, the term “carrier” refers to a substance that aids in the delivery or improves the effectiveness of the polypeptides and/or polypeptide compositions of the present invention. The carrier agent can be a naturally occurring substance, such as a protein (e.g., human serum albumin (HSA), low-density lipoprotein (LDL), or globulin); carbohydrate (e.g., a dextran, pullulan, chitin, chitosan, inulin, cyclodextrin, or hyaluronic acid); or lipid. The carrier molecule can also be a recombinant or synthetic molecule, such as a synthetic polymer, e.g., a synthetic polyamino acid. Examples of polyamino acids include poly-L-lysine (PLL), poly-L-aspartic acid and , poly-L-glutamic acid, as well as polymers comprising the D-stereoisomers of these amino acids. Other carriers include poly(L-lactide-co-glycolide) copolymer, polyethylene glycol (PEG), polyvinyl alcohol (PVA), poly(2-ethylacrylic acid), and N-isopropylacrylamide polymers. Other useful carrier molecules can be identified by routine methods.

[00158] In some embodiments, compounds of the present invention may be combined with one or more pharmaceutically acceptable excipient to form a pharmaceutical composition. As used herein, the term “pharmaceutically acceptable” refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. The phrase “pharmaceutically acceptable excipient,” as used herein, refers any ingredient other than the inventive compounds described herein (for example, a vehicle capable of suspending or dissolving the active compound) and having the properties of being substantially nontoxic and non-inflammatory in a patient. Excipients may include, for example: antiadherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluents), film formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, and waters of hydration. Exemplary excipients include, but are

not limited to: butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E, vitamin C, and xylitol. In some embodiments, pharmaceutical compositions comprise one or more active polypeptide ingredients together with ethanol, corn oil-mono-di-triglycerides, hydrogenated castor oil, DL-tocopherol, propylene glycol, gelatin, glycerol, colorants, flavors and sweeteners.

[00159] In other embodiments, pharmaceutical compositions comprise one or more active polypeptide ingredients together with a delivery agent such as 4-(2-hydroxy-4-methoxybenzamido)butanoic acid (or any of the delivery agents described in US patent number 7,744,910B2, the contents of which are incorporated herein by reference in their entirety), a pharmaceutically acceptable buffer, a disintegrant, a detergent, hydroxypropylmethylcellulose, colorants, flavors and sweeteners.

[00160] In other embodiments, pharmaceutical compositions comprise one or more active polypeptide ingredients together with ethanol, soy phosphatidyl choline, glycerol diolate which is injected into an excess of saline solution as described in US patent application 2008/0146490A1, the contents of which are incorporated herein by reference in their entirety.

[00161] The delivery of one or more polypeptides to a subject in need thereof can be achieved in a number of different ways. *In vivo* delivery can be performed directly by administering a composition comprising one or more polypeptides, to a subject. Alternatively, delivery can be performed indirectly by administering one or more vectors that encode and direct the expression of the polypeptides.

[00162] Local delivery avoids gut permeability and systemic exposure. For example, polypeptides and/or polypeptide compositions of the present invention may be used in the eye as a drop or in the posterior section of the eye by direct injection. They may be applied in the gut to target enzymes. They may be used topically in dermatologic applications (e.g., creams, ointments, transdermal patches).

[00163] Polypeptides and/or polypeptide compositions of the present invention may comprise or be formulated with one or more fusogenic agents. As used herein, the term “fusogenic agent” refers to an agent that is responsive to changes, such as pH changes in the environment for example. Upon encountering the pH of an endosome, a fusogenic agent can cause a physical change, *e.g.*, a change in osmotic properties that disrupts or increases the permeability of the endosome membrane. Preferably, the fusogenic agent changes charge, *e.g.*, becomes protonated, at pH lower than physiological range. For example, the fusogenic agent can become protonated at pH 4.5-6.5. A fusogenic agent may serve to release a polypeptide into the cytoplasm of a cell after a composition is taken up, *e.g.*, via endocytosis, by the cell, thereby increasing the cellular concentration of the polypeptide in the cell.

[00164] In some embodiments, fusogenic agents may have a moiety, *e.g.*, an amino group, which, when exposed to a specified pH range, will undergo a change, *e.g.*, in charge, *e.g.*, protonation. Changes in charge of fusogenic agents can trigger changes, *e.g.*, osmotic changes, in vesicles, *e.g.*, endocytic vesicles, *e.g.*, endosomes. For example, the fusogenic agent, upon being exposed to the pH environment of an endosome, will cause a solubility or osmotic change substantial enough to increase the porosity of (preferably, to rupture) the endosomal membrane.

[00165] Fusogenic agents may be polymers, preferably polyamino chains, *e.g.*, polyethyleneimine (PEI). PEI may be linear, branched, synthetic or natural. PEI may be, *e.g.*, alkyl substituted PEI, or lipid substituted PEI.

[00166] In other embodiments, fusogenic agents may be polyhistidine, polyimidazole, polypyridine, polypropyleneimine, mellitin, or polyacetal substances, *e.g.*, cationic polyacetals. In some embodiments, fusogenic agents may have an alpha helical structure. Fusogenic agents may be membrane disruptive agents, *e.g.*, mellitin. Other suitable fusogenic agents can be tested and identified by a skilled artisan.

[00167] Polypeptides and/or polypeptide compositions of the present invention may comprise or be formulated with one or more condensing agents. Condensing agents of compositions described herein may interact with (*e.g.*, attract, hold, or bind to) polypeptides and act to (a) condense, *e.g.*, reduce the size or charge of polypeptides and/or (b) protect polypeptides, *e.g.*, protect polypeptides against degradation. Condensing agents may include a moiety, *e.g.*, a charged moiety, which can interact with polypeptides by ionic interactions. Condensing agents would preferably be charged polymers, *e.g.*, polycationic chains. Condensing agents can be

polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quarternary salt of a polyamine, or an alpha helical peptide.

[00168] In some embodiments, polypeptides of the present invention may be bicyclic polypeptides. As used herein, the term “bicyclic polypeptide” refers to a polypeptide with two loops. As a non-limiting example, bicyclic polypeptide inhibitors of C5 may be produced in combinatorial libraries. The bicyclic polypeptides may have 2, 3, 4, 5, 6 or more amino acids per loop.

[00169] In some embodiments, polypeptides and/or polypeptide compositions of the present invention may be provided as prodrugs. As used herein, the term “prodrug” refers to a drug that is provided in an inactive form that becomes active at some point after administration. In some embodiments wherein polypeptides are administered in the form of a prodrug, amino acids critical to polypeptide inhibitory activity are unavailable to interact with the target due to a reversible chemical bond, e.g., an ester bond. Upon administration, such prodrugs may be subject to cleavage of the reversible chemical bond, e.g., through enzymatic or acid hydrolysis in the stomach, blood and/or cells of a given target tissue.

C5 inhibitors

[00170] Some polypeptides and/or polypeptide compositions of the present invention inhibit complement activation at the level of complement component C5, referred to herein as “C5 inhibitors.” Some C5 inhibitors function by preventing the cleavage of C5 to the cleavage products C5a and C5b, such inhibitors are referred to herein as “C5 cleavage inhibitors.” In some embodiments, methods of the present invention may comprise inhibiting C5 cleavage in a system. As used herein, a “system” refers to a group of related parts that function together. Such systems include those comprising C5, referred to here as “C5 systems.” C5 systems may include, but are not limited to solutions, matrices, cells, tissues, organs, and bodily fluids (including, but not limited to blood.) In some cases, C5 systems may be cellular systems. As used herein the term “cellular system” refers to a system that comprises one or more cells or one or more components or products of a cell. In some cases, C5 systems may include *in vivo* systems, *in vitro* systems and *ex vivo* systems. *In vivo* C5 systems may comprise or be comprised in a subject. As used herein, the term “subject” refers to any organism to which a compound in

accordance with the invention may be administered, *e.g.*, for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include animals (*e.g.*, mammals such as mice, rats, rabbits, non-human primates, and humans.)

[00171] In some cases, C5 inhibitors of the invention may include any of the polypeptides listed in Table 1.

Table 1. Compounds of the invention

Compound Number	Sequence	SEQ ID NO.
R3000	Ac-Nvl-C-Y-K-N-Y-H-azaTrp-E-Y-P-Tbg-Y-NH2	1
R3001	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-(N-Me)G-Nvl-(N-Me)S-NH2	2
R3002	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	3
R3003	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-P-NH2	4
R3004	Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-Y-P-Phg-Tbg-NH2	5
R3005	Ac-Nvl-C-Y-azaTrp-(N-Me)G-Tbg-Nvl-azaTrp-E-Y-P-Phg-P-NH2	6
R3006	Ac-Y-E-N-Tbg-Y-azaTrp-E-Y-(N-Me)G-Nvl-(N-Me)S-NH2	7
R3007	[mXyl(2,7)]Ac-Nvl-C-K-E-Phg-Y-C-(N-Me)S-Tbg-K-azaTrp-E-Y-NH2	8
R3008	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	9
R3020	[mXyl(2,7)]M-C-S-E-R-Y-C-E-V-R-W-E-Y-NH2	10
R3021	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	11
R3079	Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	12
R3055	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	13
R3120	Ac-Nvl-Nvl-Y-E-(N-Me)N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	14
R3057	[mXyl(2,7)]M-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	15
R3056	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	16
R3054	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	17
R3029	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-NH2	18
R3048	[mXyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	19
R3072	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-K-NH2	20
R3024	Ac-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	21
R3114	Ac-Nvl-Nvl-(N-Me)Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	22
R3050	[pXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	23
R3025	Ac-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	24
R3061	Ac-Nvl-S-Y-E-A-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	25
R3041	Ac-Y-E-N-Tbg-Y-W-E-Y-P-Phg-Nvl-NH2	26
R3077	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-K-(PEG2000)NH2	27
R3030	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-NH2	28
R3062	Ac-Nvl-S-Y-E-N-A-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	29
R3066	Ac-Nvl-S-Y-E-N-Tbg-A-azaTrp-E-Y-P-Chg-Nvl-NH2	30
R3011	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-	31

	NH2	
R3070	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-Y-A-Chg-Nvl-NH2	32
R3071	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-Y-P-A-Nvl-NH2	33
R3033	[mXylyl(2,10)]Ac-Nvl-C-Phg-A-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	34
R3038	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	35
R3012	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	36
R3060	Ac-Nvl-S-Y-A-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	37
R3039	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-A-NH2	38
R3037	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)A-H-C-Nvl-NH2	39
R3076	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-K-(BODIPY-TMR-X) NH2	40
R3074	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Tyr(OMe)-(N-Me)S-H-C-Nvl-NH2	41
R3013	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-NH2	42
R3065	[pXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-P-H-C-Nvl-NH2	43
R3073	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Phe(4-F)-(N-Me)S-H-C-Nvl-NH2	44
R3116	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-(N-Me)W-E-Y-P-Chg-Nvl-NH2	45
R3091	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-W-E-Y-(N-Me)S-A-C-Nvl-NH2	46
R3078	PEG2000-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	47
R3100	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-F-(N-Me)S-A-C-Nvl-NH2	48
R3121	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-(N-Me)Phg-Nvl-NH2	49
R3043	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-NH2	50
R3102	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-P-H-C-Nvl-NH2	51
R3026	Ac-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	52
R3031	[mXylyl(2,10)]Ac-A-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	53
R3019	[mXylyl(2,14)]Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-C-NH2	54
R3014	[mXylyl(1,9)]Ac-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	55
R3104	[pXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	56
R3059	Ac-Nvl-S-A-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	57
R3115	Ac-Nvl-Nvl-Y-(N-Me)E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	58
R3110	Ac-Y-E-N-Tbg-Y-(1-Me)W-E-Y-P-Phg-Nvl-NH2	59
R3126	Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-C-P-Phg-Tbg-NH2	60
R3049	[oXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	61
R3069	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-A-P-Chg-Nvl-NH2	62
R3015	[mXylyl(1,9)]Ac-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-NH2	63
R3068	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-A-Y-P-Chg-Nvl-NH2	64
R3105	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	65
R3106	[pXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	66
R3111	[mXylyl(4,10)]Ac-Nvl-T-Phg-C-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	67
R3112	[mXylyl(2,10)]Ac-Nle-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	68
R3113	[mXylyl(3,11)]Ac-Y-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	69

R3134	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-(3-Cl-Phe)-(N-Me)S-A-C-Nvl-NH2	70
R3018	[mXylyl(2,10)]Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-C-P-Phg-Nvl-NH2	71
R3027	Ac-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	72
R3028	Ac-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	73
R3032	[mXylyl(2,10)]Ac-Nvl-C-A-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	74
R3058	[pXylyl(2,10)]Ac-Nvl-C-Chg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	75
R3067	Ac-Nvl-S-Y-E-N-Tbg-Y-A-E-Y-P-Chg-Nvl-NH2	76
R3117	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-(N-Me)Y-P-Chg-Nvl-NH2	77
R3022	Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	78
R3016	[mXylyl(1,9)]Ac-C-Tbg-Y-azaTrp-E-Y-(N-Me)S-H-C-NH2	79
R3089	[mXylyl(2,10)]Ac-Chg-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	80
R3083	[mXylyl(2,10)]Ac-V-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	81
R3087	[mXylyl(2,10)]Ac-Nvl-C-(2-OMe)Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	82
R3103	[mXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	83
R3135	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-(D-Ala)-C-Nvl-NH2	84
R3034	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-A-E-Y-(N-Me)S-H-C-Nvl-NH2	85
R3035	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-A-Y-(N-Me)S-H-C-Nvl-NH2	86
R3036	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-A-(N-Me)S-H-C-Nvl-NH2	87
R3044	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-NH2	88
R3080	[mXylyl(2,9)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-C-Nvl-NH2	89
R3085	[mXylyl(2,10)]heptanoyl-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	90
R3086	[mXylyl(5,13)]Ac-Nvl-S-Y-E-C-Tbg-Y-azaTrp-E-Y-P-Chg-C-Nvl-NH2	91
R3092	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-F-E-Y-(N-Me)S-A-C-Nvl-NH2	92
R3095	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-(homo)F-(N-Me)S-A-C-Nvl-NH2	93
R3096	[mXylyl(2,10)]Ac-Nvl-C-Phg-Aib-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	94
R3122	[mXylyl(2,10)]Ac-Nvl-C-Tiq-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	95
R3075	[mXylyl(2,11)]Nvl-C-Y-(N-Me)S-Phg-(N-Me-4-F)Phe-(N-Me)S-H-(N-Me-4-F)Phe-G-C-NH2	96
R3107	[mXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	97
R3108	[pXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	98
R3127	[mXylyl(2,10)]Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-C-P-Phg-Tbg-NH2	99
R3133	[mXylyl(2,10)]Ac-Nvl-C-Phg-(D-Ala)-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	100
R3009	[mXylyl(2,10)]Ac-Nvl-C-Y-E-(N-Me)G-Tbg-Y-azaTrp-E-C-Nvl-P-Nvl-NH2	101
R3010	[mXylyl(2,13)]Ac-Nvl-C-Y-E-(N-Me)G-Tbg-Y-azaTrp-E-Nvl-Nvl-P-C-NH2	102
R3017	[mXylyl(2,8)]Ac-Nvl-C-Y-E-N-Tbg-Y-C-E-Y-P-Phg-Nvl-NH2	103

R3023	Ac-Y-P-Y-C-Phg-azaTrp-Tbg-E-Nvl-N-Y-Nvl-E-NH2	104
R3040	[cyclo(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl	105
R3042	[cyclo(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	106
R3045	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-NH2	107
R3046	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-NH2	108
R3047	[mXyl(2,7)]M-C-V-E-R-F-C-NH2	109
R3051	[mXyl(2,11)]Nvl-C-Y-(N-Me)S-Phg-(N-Me-4-F)Phe-(N-Me)S-H-(N-Me-4-F)Phe-(N-Me)G-C-NH2	110
R3052	[mXyl(2,9)]Nvl-C-Y-Tbg-Phg-N-(N-Me)G-L-C-Phg-(N-Me)A-NH2	111
R3053	[mXyl-bicyclo]Nvl-C-C-N-Tbg-Phg-C-Tbg-(N-Me)S-C-Tbg-NH2	112
R3063	Ac-Tbg-Y-azaTrp-E-Y-NH2	113
R3064	Ac-Y-azaTrp-E-Y-P-NH2	114
R3081	Ac-Y-E-N-Tbg-Y-azaTrp-(N-Me)E-Y-P-Phg-Nvl-NH2	115
R3082	[mXyl(1,9)]heptanoyl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	116
R3084	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-S-A-C-Nvl-NH2	117
R3088	[mXyl(2,10)]Ac-Nvl-C-Phg-T-(5-F)W-E-Y-(N-Me)S-A-C-Nvl-NH2	118
R3090	[mXyl(2,10)]Ac-Nvl-C-F-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	119
R3093	[mXyl(2,10)]Ac-Nvl-C-(D-Chg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	120
R3094	Ac-Y-E-N-Tbg-Y-(5-MeO)W-E-Y-P-Phg-Nvl-NH2	121
R3097	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-D-Y-(N-Me)S-A-C-Nvl-NH2	122
R3098	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-Q-Y-(N-Me)S-A-C-Nvl-NH2	123
R3099	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-N-Y-(N-Me)S-A-C-Nvl-NH2	124
R3101	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-G-C-Nvl-NH2	125
R3109	[mXyl(2,10)]Ac-Nvl-C-Phg-T-(1-Me-W)-E-Y-(N-Me)S-A-C-Nvl-NH2	126
R3118	Ac-Y-E-N-Tbg-Y-(D-Trp)-E-Y-P-Phg-Nvl-NH2	127
R3119	Ac-Y-E-N-Y-(D-Trp)-E-Y-P-Phg-Nvl-NH2	128
R3123	Ac-Y-E-N-Tbg-Y-azaTrp-(D-Glu)-Y-P-Phg-Nvl-NH2	129
R3124	[mXyl(1,6)]Ac-C-V-E-R-F-C-V-Y-W-E-F-NH2	130
R3125	[mXyl(1,6)]Ac-C-V-E-R-F-C-W-E-F-NH2	131
R3128	Ac-Nvl-C-Y-N-N-Tbg-E-C-E-Y-P-Phg-Tbg-NH2	132
R3129	[mXyl(2,8)]Ac-Nvl-C-Y-N-N-Tbg-E-C-E-Y-P-Phg-Tbg-NH2	133
R3130	Ac-Nvl-Nvl-Y-E-N-Tbg-(N-Me)Y-azaTrp-E-Y-P-Chg-Nvl-NH2	134
R3131	[mXyl(2,10)]Ac-Nvl-C-Phg-T-W-Asp(T)-Y-(N-Me)S-H-C-Nvl-NH2	135
R3132	[mXyl(2,10)]Ac-Nvl-C-Phg-T-(D-Trp)-E-Y-(N-Me)S-H-C-Nvl-NH2	136
R3136	[mXyl(2,10)]heptanoyl-Nvl-C-(D-Phg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	137
R3137	[mXyl(1,9)]heptanoyl-C-(D-Phg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	138
R3138	[mXyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-W-E-F-NH2	139
R3139	[mXyl(1,6)]Ac-C-Tbg-E-R-F-C-D-Tbg-Y-W-E-F-NH2	140

R3140	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-Y-P-NH2	141
R3141	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-P-NH2	142
R3142	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-azaTrp-E-Y-P-NH2	143
R3143	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-W-E-Y-P-NH2	144
R3144	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	145
R3145	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	146
R3146	[mXylyl(1,6)]Ac-C-Tbg-E-R-F-C-D-V-Y-W-E-F-NH2	147
R3147	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-Propargyl-Gly-NH2	148
R3148	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-W-E-Y-P-Phg-Nvl-NH2	149
R3149	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-W-E-Y-P-(D-Phg)-Nvl-NH2	150
R3150	[mXylyl(1,6)]Ac-C-V-A-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	151
R3151	[mXylyl(1,6)]Ac-C-V-A-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	152
R3152	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	153
R3153	[mXylyl(1,6)]Ac-C-V-E-A-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	154
R3154	[mXylyl(1,6)]Ac-C-V-E-A-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	155
R3155	[mXylyl(1,6)]Ac-C-V-E-R-A-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	156
R3156	[mXylyl(1,6)]Ac-C-V-E-R-A-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	157
R3157	[mXylyl(1,6)]Ac-C-V-E-R-F-C-A-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	158
R3158	[mXylyl(1,6)]Ac-C-V-E-R-F-C-A-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	159
R3159	[mXylyl(1,6)](des-amino)C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	160
R3160	[mXylyl(1,6)](des-amino)C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	161
R3161	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-K-NH2	162
R3162	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-K-NH2	163
R3163	[cyclo(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	164
R3164	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C12)-NH2	165
R3165	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C10)-NH2	166
R3166	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C8)-NH2	167
R3167	[mXylyl(1,6)]Ac-C-V-E-R-F-C-(alpha-methyl)D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	168
R3168	[mXylyl(1,6)]Ac-C-V-E-R-F-C-Asp(T)-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	169

R3169	[cyclo(1,6)]Ac-K-V-E-R-F-D-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	170
R3170	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-K	171
R3171	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C12)	172
R3172	[mXylyl(1,6)]Ac-C-V-E-R-F-C-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	173
R3173	[cyclo(1,6)]Ac-K-V-E-R-F-D-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	174
R3174	[cyclo(1,6)]Ac-K-V-E-R-F-D-Asp(T)-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	175
R3175	[cyclo(1,6)]Ac-K-V-E-R-F-D-D-Tbg-Y-azaTrp-E-Y-P-Chg-B20	176
R3176	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	177
R3177	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-W-P-Chg-Nvl	178
R3178	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-(homo)Phe-P-Chg-Nvl	179
R3179	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-(m-Cl-homo)Phe-P-Chg-Nvl	180
R3180	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-2Nal-P-Chg-Nvl	181
R3181	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-(3-aminomethyl)Phe-E-Y-P-Chg-Nvl	182
R3182	[cyclo-triazolyl(1,6)]Ac-X02-V-E-R-F-X31-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	183
R3183	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)	184
R3184	[cyclo-thioalkyl(1,5)]V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	185
R3185	[mXylyl(1,6)]Ac-C-V-E-R-F-C-Cle-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	186
R3186	[mXylyl(1,6)]Ac-C-V-E-R-F-C-(Ac-Pyran)-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	187
R3187	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-(3-aminomethyl)Phe-P-Chg-Nvl	188
R3188	[cyclo-olefinyl(1,6)]Ac-X30-V-E-R-F-X12-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	189
R3189	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C16)	190
R3190	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-B20	191
R3191	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-K	192
R3192	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-K-NH2	193
R3193	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-B28	194
R3194	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)-NH2	195
R3195	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-W-E-Y-P-Chg-(Lys-C16)	196
R3196	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-W-E-Y-P-Chg-K	197
R3197	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-W-E-Y-P-Chg-K14	198
R3198	[cyclo(1,6)](desamino)C-V-E-R-F-C-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)	199

R3199	[cyclo(1,6)](desamino)C-(D-Ala)-E-R-F-C-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)	200
R3200	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Aib-(Lys-C16)	201
R3201	[mXyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	211

[00172] In C5 systems, C5 and other system components may be in solution or may be fixed, such as in an assay well. C5 systems may further comprise other components of complement, in some cases including all of the components necessary to form the membrane attack complex (MAC.) In some cases, polypeptides and/or polypeptide compositions of the invention may be used to inhibit C5 cleavage in a human subject. Such polypeptides and/or polypeptide compositions may find utility in treating various complement-related disorders and/or diseases as well as accompanying inflammatory conditions. Certain C5 inhibitors are known in the art and are taught in U.S. Pat Nos. 7,348,401 and 6,355,245; both of which are herein incorporated by reference in their entireties.

[00173] Cleavage of C5 yields the proteolytic products C5a and C5b. The cleavage site of C5 that is cleaved to yield these products is referred to herein as the C5a-C5b cleavage site. C5b contributes to the formation of the membrane attack complex (MAC) while C5a stimulates the immune system and the inflammatory response. In some embodiments, polypeptides and/or polypeptide compositions of the present invention prevent the cleavage of C5 and therefore may be useful in the treatment of inflammation through the inhibition of inflammatory events including, but not limited to chemotaxis and activation of inflammatory cells (e.g. macrophages, mast cells, neutrophils and platelets), proliferation of endothelial cells and edema.

[00174] Many of the components of the complement system, including but not limited to C3, C4, and C5, are functionally inert in their native state until targeted for cleavage into multiple active components. Cleavage of C3 or C4 causes a conformational change that exposes an internal thioester domain. Within the domain, an internal thioester linkage between cysteine and glutamine residue side chains is a chemically labile bond that confers the ability of C3 and C4 to bind cell surface and/or biological molecules. The cleavage of C3 and C4 also provides the components of the C5 convertase, either C3bC4bC2a or (C3b)2Bb. (Law, S.K., et al. (1997). Protein Science. 6:263-274; van den Elsen, J.M.H., (2002). J. Mol. Biol. 322:1103-1115; the contents of each of which are herein incorporated by reference in their entireties.)

[00175] The multiple domain structure of C5 is similar to C3 and C4. The C5 convertase cleaves C5 into the components C5a and C5b. The cleavage of C5 causes a conformational change that exposes the C5b thioester-like domain, which plays a role in C5 binding C6, followed by interactions with C7 and C8 to form the cytolytic MAC. The domain structures of C5 comprise regulatory features that are critical for the processing and downstream activity of complement. (Fredslund, F. et al. (2008). *Nature*. 9:753-760; Hadders, M.A. et al. (2012). *Cell Reports*. 1:200-207.)

[00176] In some embodiments, compounds of the present invention may bind C5 and prevent cleavage of C5 into C5a and C5b cleavage products.

[00177] Recently, a new paradigm for complement activation was proposed, based upon the discovery that thrombin generates previously unidentified C5 products that support the terminal complement activation pathway (Krisinger, et al., (2014). *Blood*. 120(8):1717-1725).

[00178] Thrombin acts in the coagulation cascade, a second circulation-based process by which organisms, in response to injury, are able to limit bleeding, restore vascular integrity, and promote healing. Subsequent to vessel damage, tissue factor is exposed to the circulation, setting off a cascade of proteolytic reactions that leads to the generation of the central coagulation enzyme thrombin, which converts fibrinogen into a fibrin clot.

[00179] Historically, the complement activation pathway has been viewed separately from the coagulation cascade; however, the interplay of these two systems is worthy of renewed consideration. Coagulation and complement are coordinately activated in an overlapping spatiotemporal manner in response to common pathophysiologic stimuli to maintain homeostasis, and disease emerges when there is unchecked activation of the innate immune and coagulation responses, as evidenced by, for example, atherosclerosis, stroke, coronary heart disease, diabetes, ischemia-reperfusion injury, trauma, paroxysmal nocturnal hemoglobinuria, age-related macular degeneration, and atypical hemolytic-uremic syndrome. In fact, introduction of complement inhibitors has been found to simultaneously treat the inflammatory and thrombotic disturbances associated with some of these disorders.

[00180] As noted above, the complement system is activated via three main pathways, all converging with proteolytic activation of the central complement component C3. Subsequently, the formation of C5 convertases results in cleavage of C5 at arginine 751 (R751) to liberate a chemotactic and anaphylatoxic C5a fragment and generate C5b. C5b is the initiating factor for

assembly of the C5b dependent lytic membrane attack complex (MAC; also known as C5b-9), responsible for destroying damaged cells and pathogens.

[00181] Several molecular links between complement and coagulation have been identified. Most notably in what was described as a new complement activation pathway, thrombin was found to be capable of directly promoting activation of complement by cleaving C5, presumably at R751, thereby releasing C5a in the absence of C3 (Huber-Lang, et al., 2006. *Nature Med.* 12(6):682-687). However, these studies did not compare thrombin with the bona fide C5 convertase, and only limited biochemical analyses were performed; thus, the physiologic relevance of the pathway was not evaluable.

[00182] Using purified and plasma-based systems, the effects of thrombin and C5 convertase on C5 were assessed by measuring release of the anaphylatoxin C5a and generation of the C5b, component of MAC. It was discovered that, while thrombin cleaved C5 poorly at R751, yielding minimal C5a and C5b, it efficiently cleaved C5 at a newly identified, highly conserved R947 site, generating previously undescribed intermediates C5_T and C5b_T. Tissue factor-induced clotting of plasma led to proteolysis of C5 at a thrombin-sensitive site corresponding to this new R947 site and not R751. Combined treatment of C5 with thrombin and C5 convertase yielded C5a and C5b_T, the latter forming a C5b_T-9 membrane attack complex with significantly more lytic activity than with C5b-9. Thus, a new paradigm has been proposed for complement activation, in which thrombin is an invariant and critical partner with C5 convertase in initiating formation of a more active MAC via formation of previously unidentified C5 products that are generated via cooperative proteolysis by the two enzymes. These discoveries provide new insights into the regulation of innate immunity in the context of coagulation activation occurring in many diseases. (Krisinger, et al., (2014). *Blood.* 120(8):1717-1725).

[00183] In some embodiments, polypeptides and/or polypeptide compositions of the invention may inhibit thrombin-induced complement activation. Such polypeptides and/or polypeptide compositions may therefore be used to treat hemolysis resulting from thrombin-induced complement activation.

[00184] Given the findings of molecular links between the complement and coagulation pathways, it is believed that complement may be activated by additional components of the coagulation and/or inflammation cascades. For example, other serine proteases with slightly different substrate specificity may act in a similar way. Huber-Lang et al. (2006) showed that

thrombin not only cleaved C5 but also *in vitro*-generated C3a when incubated with native C3 (Huber-Lang, et al., 2006. *Nature Med.* 12(6):682-687; the contents of which are herein incorporated by reference in their entirety). Similarly, other components of the coagulation pathway, such as FXa, FXIa and plasmin, have been found to cleave both C5 and C3.

[00185] Specifically, in a mechanism similar to the one observed via thrombin activation, it has been observed that plasmin, FXa, FIXa and FXIa are able to cleave C5 to generate C5a and C5b (Amara, et al., (2010). *J. Immunol.* 185:5628-5636; Amara, et al., (2008) "Interaction Between the Coagulation and Complement System" in *Current Topics in Complement II*, J.D. Lambris (ed.), pp. 71-79). The anaphylatoxins produced were found to be biologically active as shown by a dose-dependent chemotactic response of neutrophils and HMC-1 cells, respectively. Plasmin-induced cleavage activity could be dose-dependently blocked by the serine protease inhibitor aprotinin and leupeptine. These findings suggest that various serine proteases belonging to the coagulation system are able to activate the complement cascade independently of the established pathways. Moreover, functional C5a and C3a are generated (as detected by immunoblotting and ELISA), both of which are known to be crucially involved in the inflammatory response.

[00186] In some embodiments, polypeptides and/or polypeptides compositions of the invention may inhibit activation of C5 by plasmin, FXa, FIXa, FXIa and other proteases of the coagulation pathway.

[00187] Human leukocyte elastase (HLE), an enzyme secreted by neutrophils and macrophages during inflammatory processes, has long been known to also release from C5 a chemotactic, C5a-like fragment. However, this C5a-like fragment, is not identical with C5a, as HLE does not cleave peptide bonds at the cleavage site that ordinarily cleaves C5 into C5a and C5b after the exposure to the complement convertases. Rather, cleavage of complement C5 by HLE has also been found to generate a functionally active C5b-like molecule that is able to participate in MAC formation (Vogt, (1999). *Immunobiology*. 201:470-477).

[00188] In some embodiments, polypeptides and/or polypeptides compositions of the invention may inhibit activation of C5 by HLE and other proteases of the inflammation cascade.

[00189] In some embodiments, polypeptides and/or polypeptide compositions of the present invention may be useful in the treatment of diseases, disorders and/or conditions where C5 cleavage leads to progression of the disease, disorder and/or condition. Such diseases, disorders

and/or conditions may include, but are not limited to immune and autoimmune, neurological, cardiovascular, pulmonary, and ocular diseases, disorders and/or conditions. Immune and autoimmune diseases and/or disorders may include, but are not limited to Acute Disseminated Encephalomyelitis (ADEM), Acute necrotizing hemorrhagic leukoencephalitis, Addison's disease, Agammaglobulinemia, Alopecia areata, Amyloidosis, Ankylosing spondylitis, Acute antibody-mediated rejection following organ transplantation, Anti-GBM/Anti-TBM nephritis, Antiphospholipid syndrome (APS), Autoimmune angioedema, Autoimmune aplastic anemia, Autoimmune dysautonomia, Autoimmune hepatitis, Autoimmune hyperlipidemia, Autoimmune immunodeficiency, Autoimmune inner ear disease (AIED), Autoimmune myocarditis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune thrombocytopenic purpura (ATP), Autoimmune thyroid disease, Autoimmune urticaria, Axonal & neuronal neuropathies, Bacterial sepsis and septic shock, Balo disease, Behcet's disease, Bullous pemphigoid, Cardiomyopathy, Castleman disease, Celiac disease, Chagas disease, Chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome, Cicatricial pemphigoid/benign mucosal pemphigoid, Crohn's disease, Cogans syndrome, Cold agglutinin disease, Congenital heart block, Coxsackie myocarditis, CREST disease, Essential mixed cryoglobulinemia, Demyelinating neuropathies, Dermatitis herpetiformis, Dermatomyositis, Devic's disease (neuromyelitis optica), Diabetes Type I, Discoid lupus, Dressler's syndrome, Endometriosis, Eosinophilic esophagitis, Eosinophilic fasciitis, Erythema nodosum, Experimental allergic encephalomyelitis, Evans syndrome, Fibromyalgia, Fibrosing alveolitis, Giant cell arteritis (temporal arteritis), Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis (GPA) see Wegener's, Graves' disease, Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, Hemolytic anemia (including atypical hemolytic uremic syndrome and plasma therapy-resistant atypical hemolytic-uremic syndrome), Henoch-Schonlein purpura, Herpes gestationis, Hypogammaglobulinemia, Idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgG4-related sclerosing disease, Immunoregulatory lipoproteins, Inclusion body myositis, Insulin-dependent diabetes (type1), Interstitial cystitis, Juvenile arthritis, Juvenile diabetes, Kawasaki syndrome, Lambert-Eaton syndrome, Large vessel vasculopathy, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, Linear IgA disease (LAD), Lupus (SLE), Lyme disease, Meniere's disease,

Microscopic polyangiitis, Mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, Multiple endocrine neoplasia syndromes, Multiple sclerosis, Multifocal motor neuropathy, Myositis, Myasthenia gravis, Narcolepsy, Neuromyelitis optica (Devic's), Neutropenia, Ocular cicatricial pemphigoid, Optic neuritis, Osteoarthritis, Palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcus), Paraneoplastic cerebellar degeneration, Paroxysmal nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, Parsonnage-Turner syndrome, Pars planitis (peripheral uveitis), Pemphigus, Peripheral neuropathy, Perivenous encephalomyelitis, Pernicious anemia, POEMS syndrome, Polyarteritis nodosa, Type I, II, & III autoimmune polyglandular syndromes, Polyendocrinopathies, Polymyalgia rheumatica, Polymyositis, Postmyocardial infarction syndrome, Postpericardiotomy syndrome, Progesterone dermatitis, Primary biliary cirrhosis, Primary sclerosing cholangitis, Psoriasis, Psoriatic arthritis, Idiopathic Pulmonary fibrosis, Pyoderma gangrenosum, Pure red cell aplasia, Raynaud's phenomenon, Reactive arthritis, Reflex sympathetic dystrophy, Reiter's syndrome, Relapsing polychondritis, Restless legs syndrome, Retroperitoneal fibrosis, Rheumatic fever, Rheumatoid arthritis, Sarcoidosis, Schmidt syndrome, Scleritis, Scleroderma, Shiga-Toxin producing Escherichia Coli Hemolytic-Uremic Syndrome (STEC-HUS), Sjogren's syndrome, Small vessel vasculopathy, Sperm & testicular autoimmunity, Stiff person syndrome, Subacute bacterial endocarditis (SBE), Susac's syndrome, Sympathetic ophthalmia, Takayasu's arteritis, Temporal arteritis/Giant cell arteritis, Thrombocytopenic purpura (TTP), Tolosa-Hunt syndrome, Transverse myelitis, Tubular autoimmune disorder, Ulcerative colitis, Undifferentiated connective tissue disease (UCTD), Uveitis, Vesiculobullous dermatosis, Vasculitis, Vitiligo and Wegener's granulomatosis (also known as Granulomatosis with Polyangiitis (GPA)). Neurological diseases, disorders and/or conditions may include, but are not limited to Alzheimer's disease, Parkinson's disease, Lewy body dementia and Multiple sclerosis. Cardiovascular diseases, disorders and/or conditions may include, but are not limited to atherosclerosis, myocardial infarction, stroke, vasculitis, trauma and conditions arising from cardiovascular intervention (including, but not limited to cardiac bypass surgery, arterial grafting and angioplasty). Pulmonary diseases, disorders and/or conditions may include, but are not limited to asthma, pulmonary fibrosis, chronic obstructive pulmonary disease (COPD) and adult respiratory distress syndrome. Ocular related applications include, but are not limited to: Age-related macular degeneration, allergic and giant papillary

conjunctivitis, Behcet's disease, choroidal inflammation, complications related to intraocular surgery, corneal transplant rejection, corneal ulcers, cytomegalovirus retinitis, dry eye syndrome, endophthalmitis, Fuch's disease, Glaucoma, immune complex vasculitis, inflammatory conjunctivitis, ischemic retinal disease, keratitis, macular edema, ocular parasitic infestation/migration, retinitis pigmentosa, scleritis, Stargardt disease, subretinal fibrosis, uveitis, vitreo-retinal inflammation, and Vogt-Koyanagi-Harada disease.

[00190] Polypeptides and/or polypeptide compositions of the present invention may be particularly useful in the treatment of patients with PNH that show a poor response to monoclonal antibody therapies, such as ECULIZUMAB® therapy, due to mutations in the C5 gene that prevent binding of the antibody to C5 (Nishimura, J-I. (2012). 54th ASH Annual Meeting, Abstract 3197).

[00191] Polypeptides and/or polypeptide compositions of the present invention may be useful in the treatment of infectious diseases, disorders and/or conditions, for example, in a subject having an infection. In some preferred embodiments the subject has an infection and is at risk of developing sepsis or a septic syndrome. Polypeptides and/or polypeptide compositions of the present invention are particularly useful in the treatment of sepsis.

[00192] Polypeptides and/or polypeptide compositions of the present invention may also be administered to improve the outcome of clinical procedures wherein complement inhibition is desired. Such procedures may include, but are not limited to grafting, transplantation, implantation, catheterization, intubation and the like. In some embodiments, polypeptides and/or polypeptide compositions of the invention are used to coat devices, materials and/or biomaterials used in such procedures. In some embodiments, the inner surface of a tube may be coated with polypeptides and/or polypeptide compositions to prevent complement activation within a bodily fluid that passes through the tube, either *in vivo* or *ex vivo*, e.g., extracorporeal shunting, e.g., dialysis and cardiac bypass.

[00193]

Methods of Use

Therapeutic indications

[00194] The invention relates in particular to the use of polypeptide (e.g. peptidomimetics and cyclic polypeptides) and compositions containing at least one polypeptide, for the treatment of a

disorder, condition or disease. In some cases, compounds and compositions of the invention may be used to treat subjects suffering from paroxysmal nocturnal hemoglobinuria (PNH). Subjects with PNH are unable to synthesize functional versions of the complement regulatory proteins CD55 and CD59 on hematopoietic stem cells. This results in complement-mediated hemolysis and a variety of downstream complications. Other complement-related disorders and diseases include, but are not limited to autoimmune diseases and disorders, neurological diseases and disorders, blood diseases and disorders and infectious diseases and disorders. Experimental evidence suggests that many complement-related disorders are alleviated through inhibition of complement activity.

[00195] An acquired mutation in the phosphatidylinositol glycan anchor biosynthesis, class A (PIG-A) gene that originates from a multipotent hematopoietic stem cell results in a rare disease known as paroxysmal nocturnal hemoglobinuria (PNH) (Pu, J.J. et al., Paroxysmal nocturnal hemoglobinuria from bench to bedside. *Clin Transl Sci.* 2011 Jun;4(3):219-24). PNH is characterized by bone marrow disorder, hemolytic anemia and thrombosis. The PIG-A gene product is necessary for the production of a glycolipid anchor, glycosylphosphatidylinositol (GPI), utilized to tether proteins to the plasma membrane. Two complement-regulatory proteins, CD55 and CD59, become nonfunctional in the absence of GPI. This leads to complement-mediated destruction of these cells. Polypeptides and/or polypeptide compositions of the present invention are particularly useful in the treatment of PNH. As used herein the terms “treat,” “treatment,” and the like, refer to relief from or alleviation of pathological processes. In the context of the present invention insofar as it relates to any of the other conditions recited herein below, the terms “treat,” “treatment,” and the like mean to relieve or alleviate at least one symptom associated with such condition, or to slow or reverse the progression or anticipated progression of such condition, such as slowing the progression of a malignancy or cancer, or increasing the clearance of an infectious organism to alleviate/reduce the symptoms caused by the infection, e.g., hepatitis caused by infection with a hepatitis virus or reducing the destruction of red blood cells (as measured by reduced transfusion requirements or increased hematocrit or hemoglobin levels) resulting from paroxysmal nocturnal hemoglobinuria.

[00196] By “lower” or “reduce” in the context of a disease marker or symptom is meant a statistically significant decrease in such level. The decrease can be, for example, at least 10%, at

least 20%, at least 30%, at least 40% or more, and is preferably down to a level accepted as within the range of normal for an individual without such disorder.

[00197] By “increase” or “raise” in the context of a disease marker or symptom is meant a statistically significant rise in such level. The increase can be, for example, at least 10%, at least 20%, at least 30%, at least 40% or more, and is preferably up to a level accepted as within the range of normal for an individual without such disorder.

[00198] As used herein, the phrases “therapeutically effective amount” and “prophylactically effective amount” refer to an amount that provides a therapeutic benefit in the treatment, prevention, or management of pathological processes or an overt symptom of one or more pathological processes. The specific amount that is therapeutically effective can be readily determined by an ordinary medical practitioner, and may vary depending on factors known in the art, such as, for example, the type of pathological processes, patient history and age, the stage of pathological processes, and the administration of other agents that inhibit pathological processes.

[00199] As used herein, a “pharmaceutical composition” comprises a pharmacologically effective amount of a polypeptide and a pharmaceutically acceptable carrier. As used herein, “pharmacologically effective amount,” “therapeutically effective amount” or simply “effective amount” refers to that amount of a polypeptide effective to produce the intended pharmacological, therapeutic or preventive result. For example, if a given clinical treatment is considered effective when there is at least a 10% alteration (increase or decrease) in a measurable parameter associated with a disease or disorder, a therapeutically effective amount of a drug for the treatment of that disease or disorder is the amount necessary to effect at least a 10% alteration in that parameter. For example, a therapeutically effective amount of a polypeptide may be one that alters binding of a target to its natural binding partner by at least 10%.

[00200] The term “pharmaceutically acceptable carrier” refers to a carrier for administration of a therapeutic agent. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The term specifically excludes cell culture medium. For drugs administered orally, pharmaceutically acceptable carriers include, but are not limited to pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and

calcium phosphate, and lactose, while corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin, while the lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate, to delay absorption in the gastrointestinal tract. Agents included in drug formulations are described further herein below.

[00201] Efficacy of treatment or amelioration of disease can be assessed, for example by measuring disease progression, disease remission, symptom severity, reduction in pain, quality of life, dose of a medication required to sustain a treatment effect, level of a disease marker or any other measurable parameter appropriate for a given disease being treated or targeted for prevention. It is well within the ability of one skilled in the art to monitor efficacy of treatment or prevention by measuring any one of such parameters, or any combination of parameters. In connection with the administration of a polypeptide or pharmaceutical composition thereof, "effective against" a disease or disorder indicates that administration in a clinically appropriate manner results in a beneficial effect for at least a fraction of patients, such as an improvement of symptoms, a cure, a reduction in disease load, reduction in tumor mass or cell numbers, extension of life, improvement in quality of life, a reduction in the need for blood transfusions or other effect generally recognized as positive by medical doctors familiar with treating the particular type of disease or disorder.

[00202] A treatment or preventive effect is evident when there is a statistically significant improvement in one or more parameters of disease status, or by a failure to worsen or to develop symptoms where they would otherwise be anticipated. As an example, a favorable change of at least 10% in a measurable parameter of disease, and preferably at least 20%, 30%, 40%, 50% or more can be indicative of effective treatment. Efficacy for a given polypeptide drug or formulation of that drug can also be judged using an experimental animal model for the given disease as known in the art. When using an experimental animal model, efficacy of treatment is evidenced when a statistically significant modulation in a marker or symptom is observed.

[00203] The polypeptide and an additional therapeutic agent can be administered in combination in the same composition, *e.g.*, parenterally, or the additional therapeutic agent can be administered as part of a separate composition or by another method described herein.

Inflammatory Indications

[00204] In some embodiments, compounds and compositions of the invention may be used to treat subjects with diseases, disorders and/or conditions related to inflammation. Inflammation may be upregulated during the proteolytic cascade of the complement system. Although inflammation may have beneficial effects, excess inflammation may lead to a variety of pathologies (Markiewski et al. 2007. Am J Pathol. 17: 715-27). Accordingly, compounds and compositions of the present invention may be used to reduce or eliminate inflammation associated with complement activation.

Sterile inflammation

[00205] In some embodiments, compounds and compositions of the present invention may be used to treat, prevent or delay development of sterile inflammation. Sterile inflammation is inflammation that occurs in response to stimuli other than infection. Sterile inflammation may be a common response to stress such as genomic stress, hypoxic stress, nutrient stress or endoplasmic reticulum stress caused by a physical, chemical, or metabolic noxious stimuli. Sterile inflammation may contribute to pathogenesis of many diseases such as, but not limited to, ischemia-induced injuries, rheumatoid arthritis, acute lung injuries, drug-induced liver injuries, inflammatory bowel diseases and/or other diseases, disorders or conditions. Mechanism of sterile inflammation and methods and compositions for treatment, prevention and/or delaying of symptoms of sterile inflammation may include any of those taught by Rubartelli et al. in *Frontiers in Immunology*, 2013, 4:398-99, Rock et al. in *Annu Rev Immunol.* 2010, 28:321-342 or in United States Patent No. 8,101,586, the contents of each of which are herein incorporated by reference in their entirety.

Systemic inflammatory response (SIRS) and sepsis

[00206] In some embodiments, compounds and compositions of the invention may be used to treat and/or prevent systemic inflammatory response syndrome (SIRS). SIRS is inflammation affecting the whole body. Where SIRS is caused by an infection, it is referred to as sepsis. SIRS may also be caused by non-infectious events such as trauma, injury, burns, ischemia, hemorrhage and/or other conditions. During sepsis and SIRS, complement activation leads to excessive generation of complement activation products which may cause multi organ failure (MOF) in subjects. Compounds and compositions of the invention may be used to control and/or balance

complement activation for prevention and treatment of SIRS, sepsis and/or MOF. The methods of applying complement inhibitors to treat SIRS and sepsis may include those taught by Rittirsch et al. in *Clin Dev Immunol*, 2012, 962927, in U.S. publication No. US2013/0053302 or in United States Patent No. 8,329,169, the contents of each of which are herein incorporated by reference in their entirety.

Acute respiratory distress syndrome (ARDS)

[00207] In some embodiments, compounds and compositions of the invention may be used to treat and/or prevent development of acute respiratory distress syndrome (ARDS). ARDS is a widespread inflammation of the lungs and may be caused by trauma, infection (e.g., sepsis), severe pneumonia and/or inhalation of harmful substances. ARDS is typically a severe, life-threatening complication. Studies suggest that neutrophils may contribute to development of ARDS by affecting the accumulation of polymorphonuclear cells in the injured pulmonary alveoli and interstitial tissue of the lungs. Accordingly, compounds and compositions of the invention may be administered to reduce and/or prevent tissue factor production in alveolar neutrophils. Compounds and compositions of the invention may further be used for treatment, prevention and/or delaying of ARDS, in some cases according to any of the methods taught in International publication No. WO2009/014633, the contents of which are herein incorporated by reference in their entirety.

Periodontitis

[00208] In some embodiments, compounds and compositions of the invention may be used to treat or prevent development of periodontitis and/or associated conditions. Periodontitis is a widespread, chronic inflammation leading to the destruction of periodontal tissue which is the tissue supporting and surrounding the teeth. The condition also involves alveolar bone loss (bone that holds the teeth). Periodontitis may be caused by a lack of oral hygiene leading to accumulation of bacteria at the gum line, also known as dental plaque. Certain health conditions such as diabetes or malnutrition and/or habits such as smoking may increase the risk of periodontitis. Periodontitis may increase the risk of stroke, myocardial infarction, atherosclerosis, diabetes, osteoporosis, pre-term labor, as well as other health issues. Studies demonstrate a correlation between periodontitis and local complement activity. Periodontal

bacteria may either inhibit or activate certain components of the complement cascade. Accordingly, compounds and compositions of the invention may be used to prevent and/or treat periodontitis and associated diseases and conditions. Complement activation inhibitors and treatment methods may include any of those taught by Hajishengallis in Biochem Pharmacol. 2010, 15; 80(12): 1 and Lambris or in US publication No. US2013/0344082, the contents of each of which are herein incorporated by reference in their entirety.

Wounds and injuries

[00209] Compounds and compositions of the invention may be used to treat and/or promote healing of different types of wounds and/or injuries. As used herein, the term “injury” typically refers to physical trauma, but may include localized infection or disease processes. Injuries may be characterized by harm, damage or destruction caused by external events affecting body parts and/or organs. Wounds are associated with cuts, blows, burns and/or other impacts to the skin, leaving the skin broken or damaged. Wounds and injuries are often acute but if not healed properly they may lead to chronic complications and/or inflammation.

Wounds and burn wounds

[00210] In some embodiments, compounds and compositions of the invention may be used to treat and/or to promote healing of wounds. Healthy skin provides a waterproof, protective barrier against pathogens and other environmental affectors. The skin also controls body temperature and fluid evaporation. When skin is wounded these functions are disrupted making skin healing challenging. Wounding initiates a set of physiological processes related to the immune system that repair and regenerate tissue. Complement activation is one of these processes. Complement activation studies have identified several complement components involved with wound healing as taught by van de Goot et al. in J Burn Care Res 2009, 30:274–280 and Cazander et al. Clin Dev Immunol, 2012, 2012:534291, the contents of each of which are herein incorporated by reference in their entirety. In some cases, complement activation may be excessive, causing cell death and enhanced inflammation (leading to impaired wound healing and chronic wounds). In some cases, compounds and compositions of the present invention may be used to reduce or eliminate such complement activation to promote wound healing. Treatment with compounds and compositions of the invention may be carried out according to any of the methods for

treating wounds disclosed in International publication number WO2012/174055, the contents of which are herein incorporated by reference in their entirety.

Head trauma

[00211] In some embodiments, compounds and compositions of the invention may be used to treat and/or promote healing of head trauma. Head traumas include injuries to the scalp, the skull or the brain. Examples of head trauma include, but are not limited to concussions, contusions, skull fracture, traumatic brain injuries and/or other injuries. Head traumas may be minor or severe. In some cases, head trauma may lead to long term physical and/or mental complications or death. Studies indicate that head traumas may induce improper intracranial complement cascade activation, which may lead to local inflammatory responses contributing to secondary brain damage by development of brain edema and/or neuronal death (Stahel et al. in *Brain Research Reviews*, 1998, 27: 243–56, the contents of which are herein incorporated by reference in their entirety). Compounds and compositions of the invention may be used to treat head trauma and/or to reduce or prevent related secondary complications. Methods of using compounds and compositions of the invention to control complement cascade activation in head trauma may include any of those taught by Holers et al. in United States Patent No. 8,911,733, the contents of which are herein incorporated by reference in their entirety.

Crush injury

[00212] In some embodiments, compounds and compositions of the invention may be used to treat and/or promote healing of crush injuries. Crush injuries are injuries caused by a force or a pressure put on the body causing bleeding, bruising, fractures, nerve injuries, wounds and/or other damages to the body. Compounds and compositions of the invention may be used to reduce complement activation following crush injuries, thereby promoting healing after crush injuries (e.g. by promoting nerve regeneration, promoting fracture healing, preventing or treating inflammation, and/or other related complications). Compounds and compositions of the invention may be used to promote healing according to any of the methods taught in United States Patent No. 8,703,136; International Publication Nos. WO2012/162215; WO2012/174055; or US publication No. US2006/0270590, the contents of each of which are herein incorporated by reference in their entirety.

Autoimmune disease

[00213] The compounds and compositions of the invention may be used to treat subjects with autoimmune diseases and/or disorders. The immune system may be divided into innate and adaptive systems, referring to nonspecific immediate defense mechanisms and more complex antigen-specific systems, respectively. The complement system is part of the innate immune system, recognizing and eliminating pathogens. Additionally, complement proteins may modulate adaptive immunity, connecting innate and adaptive responses. Autoimmune diseases and disorders are immune abnormalities causing the system to target self tissues and substances. Autoimmune disease may involve certain tissues or organs of the body. Compounds and compositions of the invention may be used to modulate complement in the treatment and/or prevention of autoimmune diseases. In some cases, such compounds and compositions may be used according to the methods presented in Ballanti et al. *Immunol Res* (2013) 56:477–491, the contents of which are herein incorporated by reference in their entirety.

Anti-phospholipid syndrome (APS) and catastrophic anti-phospholipid syndrome (CAPS)

[00214] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat anti-phospholipid syndrome (APS) by complement activation control. APS is an autoimmune condition caused by anti-phospholipid antibodies that cause the blood to clot. APS may lead to recurrent venous or arterial thrombosis in organs, and complications in placental circulations causing pregnancy-related complications such as miscarriage, still birth, preeclampsia, premature birth and/or other complications. Catastrophic anti-phospholipid syndrome (CAPS) is an extreme and acute version of a similar condition leading to occlusion of veins in several organs simultaneously. Studies suggest that complement activation may contribute to APS-related complications including pregnancy-related complications, thrombotic (clotting) complications, and vascular complications. Compound and compositions of the invention may be used to treat APS-related conditions by reducing or eliminating complement activation. In some cases, compounds and compositions of the invention may be used to treat APS and/or APS-related complications according to the methods taught by Salmon et al. *Ann Rheum Dis* 2002;61(Suppl II):ii46–ii50 and Mackworth-Young in *Clin Exp Immunol* 2004, 136:393–401, the contents of which are herein incorporated by reference in their entirety.

Cold agglutinin disease

[00215] In some embodiments, compounds and compositions of the invention may be used to treat cold agglutinin disease (CAD), also referred to as cold agglutinin-mediated hemolysis. CAD is an autoimmune disease resulting from a high concentration of IgM antibodies interacting with red blood cells at low range body temperatures [Engelhardt et al. Blood, 2002, 100(5):1922-23]. CAD may lead to conditions such as anemia, fatigue, dyspnea, hemoglobinuria and/or acrocyanosis. CAD is related to robust complement activation and studies have shown that CAD may be treated with complement inhibitor therapies. Accordingly, the present invention provides methods of treating CAD using compounds and compositions of the invention. In some cases, compounds and compositions of the invention may be used to treat CAD according to the methods taught by Roth et al in Blood, 2009, 113:3885-86 or in International publication No. WO2012/139081, the contents of each of which are herein incorporated by reference in their entirety.

Vascular indications

[00216] In some embodiments, compounds and compositions of the invention may be used to treat vascular indications affecting blood vessels (e.g., arteries, veins, and capillaries). Such indications may affect blood circulation, blood pressure, blood flow, organ function and/or other bodily functions.

Thrombotic microangiopathy (TMA)

[00217] In some embodiments, compounds and compositions of the invention may be used to treat and/or prevent thrombotic microangiopathy (TMA) and associated diseases.

Microangiopathies affect small blood vessels (capillaries) of the body causing capillary walls to become thick, weak, and prone to bleeding and slow blood circulation. TMAs tend to lead to the development of vascular thrombi, endothelial cell damage, thrombocytopenia, and hemolysis. Organs such as the brain, kidney, muscles, gastrointestinal system, skin, and lungs may be affected. TMAs may arise from medical operations and/or conditions that include, but are not limited to, hematopoietic stem cell transplantation (HSCT), renal disorders, diabetes and/or other conditions. TMAs may be caused by underlying complement system dysfunction, as described by Meri et al. in European Journal of Internal Medicine, 2013, 24: 496-502, the contents of

which are herein incorporated by reference in their entirety. Generally, TMAs may result from increased levels of certain complement components leading to thrombosis. In some cases, this may be caused by mutations in complement proteins or related enzymes. Resulting complement dysfunction may lead to complement targeting of endothelial cells and platelets leading to increased thrombosis. In some embodiments, TMAs may be prevented and/or treated with compounds and compositions of the invention. In some cases, methods of treating TMAs with compounds and compositions of the invention may be carried out according to those described in US publication Nos. US2012/0225056 or US2013/0246083, the contents of each of which are herein incorporated by reference in their entirety.

Disseminated intravascular coagulation (DIC)

[00218] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat disseminated intravascular coagulation (DIC) by controlling complement activation. DIC is a pathological condition where the clotting cascade in blood is widely activated and results in formation of blood clots especially in the capillaries. DIC may lead to an obstructed blood flow of tissues and may eventually damage organs. Additionally, DIC affects the normal process of blood clotting that may lead to severe bleeding. Compounds and compositions of the invention may be used to treat, prevent or reduce the severity of DIC by modulating complement activity. In some cases compounds and compositions of the invention may be used according to any of the methods of DIC treatment taught in US Patent No. 8,652,477, the contents of which are herein incorporated by reference in their entirety.

Vasculitis

[00219] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat vasculitis. Generally, vasculitis is a disorder related to inflammation of blood vessels, including veins and arteries, characterized by white blood cells attacking tissues and causing swelling of the blood vessels. Vasculitis may be associated with an infection, such as in Rocky Mountain spotted fever, or autoimmunity. An example of autoimmunity associated vasculitis is Anti-Neutrophil Cytoplasmic Autoantibody (ANCA) vasculitis. ANCA vasculitis is caused by abnormal antibodies attacking the body's own cells and tissues. ANCAs attack the cytoplasm of certain white blood cells and neutrophils, causing them to attack the walls of the

vessels in certain organs and tissues of the body. ANCA vasculitis may affect skin, lungs, eyes and/or kidney. Studies suggest that ANCA disease activates an alternative complement pathway and generates certain complement components that create an inflammation amplification loop resulting in a vascular injury (Jennette et al. 2013, Semin Nephrol. 33(6): 557-64, the contents of which are herein incorporated by reference in their entirety). In some cases, compounds and compositions of the invention may be used to prevent and/or treat ANCA vasculitis by inhibiting complement activation.

Neurological indications

[00220] The compounds and compositions of the invention may be used to prevent, treat and/or ease the symptoms of neurological indications, including, but not limited to neurodegenerative diseases and related disorders. Neurodegeneration generally relates to a loss of structure or function of neurons, including death of neurons. These disorders may be treated by inhibiting the effect of complement on neuronal cells using compounds and compositions of the invention. Neurodegenerative related disorders include, but are not limited to, Amyotrophic Lateral Sclerosis (ALS), Multiple Sclerosis (MS), Parkinson's disease and Alzheimer's disease.

Amyotrophic lateral sclerosis (ALS)

[00221] In some embodiments, compounds and compositions of the invention may be used to prevent, treat and/or ease the symptoms of ALS. ALS is a fatal motor neuron disease characterized by the degeneration of spinal cord neurons, brainstems and motor cortex. ALS causes loss of muscle strength leading eventually to a respiratory failure. Complement dysfunction may contribute to ALS, and therefore ALS may be prevented, treated and/or the symptoms may be reduced by therapy with compounds and compositions of the invention targeting complement activity. In some cases, compounds and compositions of the invention may be used to promote nerve regeneration. In some cases, compounds and compositions of the invention may be used as complement inhibitors according to any of the methods taught in US publication No. US2014/0234275 or US2010/0143344, the contents of each of which are herein incorporated by reference in their entirety.

Alzheimer's disease

[00222] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat Alzheimer's disease by controlling complement activity. Alzheimer's disease is a chronic neurodegenerative disease with symptoms that may include disorientation, memory loss, mood swings, behavioral problems and eventually loss of bodily functions. Alzheimer's disease is thought to be caused by extracellular brain deposits of amyloid that are associated with inflammation-related proteins such as complement proteins (Sjoberg et al. 2009. Trends in Immunology. 30(2): 83-90, the contents of which are herein incorporated by reference in their entirety). In some cases, compounds and compositions of the invention may be used as complement inhibitors according to any of the Alzheimer's treatment methods taught in US publication No. US2014/0234275, the contents of which are herein incorporated by reference in their entirety.

Kidney-related indications

[00223] The compounds and compositions of the invention may be used to treat certain diseases, disorders and/or conditions related to kidneys, in some cases by inhibiting complement activity. Kidneys are organs responsible for removing metabolic waste products from the blood stream. Kidneys regulate blood pressure, the urinary system, and homeostatic functions and are therefore essential for a variety of bodily functions. Kidneys may be more seriously affected by inflammation (as compared to other organs) due to unique structural features and exposure to blood. Kidneys also produce their own complement proteins which may be activated upon infection, kidney disease, and renal transplantations. In some cases, compounds and compositions of the invention may be used as complement inhibitors in the treatment of certain diseases, conditions, and/or disorders of the kidney according to the methods taught by Quigg, J Immunol 2003; 171:3319-24, the contents of which are herein incorporated by reference in their entirety.

Lupus Nephritis

[00224] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat lupus nephritis by inhibiting complement activity. Lupus nephritis is a kidney inflammation caused by an autoimmune disease called systemic lupus erythematosus (SLE). Symptoms of lupus nephritis include high blood pressure; foamy urine; swelling of the

legs, the feet, the hands, or the face; joint pain; muscle pain; fever; and rash. Lupus nephritis may be treated by inhibitors that control complement activity, including compounds and compositions of the present invention. Methods and compositions for preventing and/or treating Lupus nephritis by complement inhibition may include any of those taught in US publication No. US2013/0345257 or United States Patent No. 8,377,437, the contents of each of which are herein incorporated by reference in their entirety.

Membranous glomerulonephritis (MGN)

[00225] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat membranous glomerulonephritis (MGN) disorder by inhibiting the activation of certain complement components. MGN is a disorder of the kidney that may lead to inflammation and structural changes. MGN is caused by antibodies binding to a soluble antigen in kidney capillaries (glomerulus). MGN may affect kidney functions, such as filtering fluids and may lead to kidney failure. Compounds and compositions of the invention may be used according to methods of preventing and/or treating MGN by complement inhibition taught in U.S. publication No. US2010/0015139 or in International publication No. WO2000/021559, the contents of each of which are herein incorporated by reference in their entirety.

Hemodialysis complications

[00226] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat complications associated with hemodialysis by inhibiting complement activation. Hemodialysis is a medical procedure used to maintain kidney function in subjects with kidney failure. In hemodialysis, the removal of waste products such as creatinine, urea, and free water from blood is performed externally. A common complication of hemodialysis treatment is chronic inflammation caused by contact between blood and the dialysis membrane. Another common complication is thrombosis referring to a formation of blood clots that obstructs the blood circulation. Studies have suggested that these complications are related to complement activation. Hemodialysis may be combined with complement inhibitor therapy to provide means of controlling inflammatory responses and pathologies and/or preventing or treating thrombosis in subjects going through hemodialysis due to kidney failure. Methods of using compounds and compositions of the invention for treatment of hemodialysis complications

may be carried out according to any of the methods taught by DeAngelis et al in Immunobiology, 2012, 217(11): 1097–1105 or by Kourtzelis et al. Blood, 2010, 116(4):631-639, the contents of each of which are herein incorporated by reference in their entirety.

Ocular diseases

[00227] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat certain ocular related diseases, disorders and/or conditions. In a healthy eye the complement system is activated at a low level and is continuously regulated by membrane-bound and soluble intraocular proteins that protect against pathogens. Therefore the activation of complement plays an important role in several complications related to the eye and controlling complement activation may be used to treat such diseases. Compounds and compositions of the invention may be used as complement inhibitors in the treatment of ocular disease according to any of the methods taught by Jha et al. in Mol Immunol. 2007; 44(16): 3901–3908 or in US Patent No. 8,753,625, the contents of each of which are herein incorporated by reference in their entirety.

Age-related macular degeneration (AMD)

[00228] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat age-related macular degeneration (AMD) by inhibiting ocular complement activation. AMD is a chronic ocular disease causing blurred central vision, blind spots in central vision, and/or eventual loss of central vision. Central vision affects ability to read, drive a vehicle and/or recognize faces. AMD is generally divided into two types, non-exudative (dry) and exudative (wet). Dry AMD refers to the deterioration of the macula which is the tissue in the center of the retina. Wet AMD refers to the failure of blood vessels under the retina leading to leaking of blood and fluid. Several human and animal studies have identified complement proteins that are related to AMD and novel therapeutic strategies included controlling complement activation pathways, as discussed by Jha et al. in Mol Immunol. 2007; 44(16): 3901–8. Methods of the invention involving the use of compounds and compositions of the invention for prevention and/or treatment of AMD may include any of those taught in US publication Nos. US2011/0269807 or US2008/0269318, the contents of each of which are herein incorporated by reference in their entirety.

Corneal disease

[00229] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat corneal diseases by inhibiting ocular complement activation. The complement system plays an important role in protection of the cornea from pathogenic particles and/or inflammatory antigens. The cornea is the outermost front part of the eye covering and protecting the iris, pupil and anterior chamber and is therefore exposed to external factors. Corneal diseases include, but are not limited to, keratoconus, keratitis, ocular herpes and/or other diseases. Corneal complications may cause pain, blurred vision, tearing, redness, light sensitivity and/or corneal scarring. The complement system is critical for corneal protection, but complement activation may cause damage to the corneal tissue after an infection is cleared as certain complement compounds are heavily expressed. Methods of the present invention for modulating complement activity in the treatment of corneal disease may include any of those taught by Jha et al. in Mol Immunol. 2007; 44(16): 3901–8, the contents of which are herein incorporated by reference in their entirety.

Autoimmune uveitis

[00230] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat uveitis, which is an inflammation of the uveal layer of the eye. Uvea is the pigmented area of the eye comprising the choroids, iris and ciliary body of the eye. Uveitis causes redness, blurred vision, pain, synechia and may eventually cause blindness. Studies have indicated that complement activation products are present in the eyes of patients with autoimmune uveitis and complement plays an important role in disease development. In some cases, compounds and compositions of the invention may be used to treat and/or prevent uveitis according to any of the methods identified in Jha et al. in Mol Immunol. 2007. 44(16): 3901–8, the contents of which are herein incorporated by reference in their entirety.

Diabetic retinopathy

[00231] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat diabetic retinopathy which is a disease caused by changes in retinal blood vessels in diabetic patients. Retinopathy may cause blood vessel swelling and fluid leaking and/or growth of abnormal blood vessels. Diabetic retinopathy affects vision and may eventually

lead to blindness. Studies have suggested that activation of complement has an important role in the development of diabetic retinopathy. In some cases, compounds and compositions of the invention may be used according to methods of diabetic retinopathy treatment described in Jha et al. Mol Immunol. 2007; 44(16): 3901–8, the contents of which are herein incorporated by reference in their entirety.

Pre-eclampsia and HELLP- syndrome

[00232] In some embodiments, compounds and compositions of the invention may be used to prevent and/or treat pre-eclampsia and/or HELLP (abbreviation standing for syndrome features of 1) hemolysis, 2) elevated liver enzymes and 3) low platelet count) syndrome by complement inhibitor therapy. Pre-eclampsia is a disorder of pregnancy with symptoms including elevated blood pressure, swelling, shortness of breath, kidney dysfunction, impaired liver function and/or low blood platelet count. Pre-eclampsia is typically diagnosed by a high urine protein level and high blood pressure. HELLP syndrome is a combination of hemolysis, elevated liver enzymes and low platelet conditions. Hemolysis is a disease involving rupturing of red blood cells leading to the release of hemoglobin from red blood cells. Elevated liver enzymes may indicate a pregnancy-induced liver condition. Low platelet levels lead to reduced clotting capability, causing danger of excessive bleeding. HELLP is associated with a pre-eclampsia and liver disorder. HELLP syndrome typically occurs during the later stages of pregnancy or after childbirth. It is typically diagnosed by blood tests indicating the presence of the three conditions it involves. Typically HELLP is treated by inducing delivery.

[00233] Studies suggest that complement activation occurs during HELLP syndrome and pre-eclampsia and that certain complement components are present at increased levels during HELLP and pre-eclampsia. Complement inhibitors may be used as therapeutic agents to prevent and/or treat these conditions. Compounds and compositions of the invention may be used according to methods of preventing and/or treating HELLP and pre-eclampsia taught by Heager et al. in *Obstetrics & Gynecology*, 1992, 79(1):19-26 or in International publication No. WO201/078622, the contents of each of which are herein incorporated by reference in their entirety.

Dosage and administration

[00234] For use as treatment of human subjects, polypeptides can be formulated as pharmaceutical compositions. Depending on the subject to be treated, the mode of administration, and the type of treatment desired (e.g., prevention, prophylaxis, or therapy) the polypeptides are formulated in ways consonant with these parameters. A summary of such techniques is found in Remington: The Science and Practice of Pharmacy, 21st Edition, Lippincott Williams & Wilkins, (2005); and Encyclopedia of Pharmaceutical Technology, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York, each of which is incorporated herein by reference.

[00235] Compositions of the present invention are preferably provided in a therapeutically effective amount, which may be, for example, a daily amount of from about 0.1 mg to about 100 mg, from about 0.5 mg to about 200 mg, from about 1 mg to about 300 mg, from about 5 mg to about 500 mg, from about 10 mg to about 750 mg, from about 50 mg to about 1000 mg or at least 1000 mg. In one embodiment, a pharmaceutical composition comprises a capsule, for example in unit dosage form.

Unit dosage forms

[00236] The polypeptides of the invention may be present in amounts totaling 0.1-95% by weight of the total weight of the composition. The composition may be provided in a dosage form that is suitable for oral administration. Thus, the pharmaceutical composition may be in the form of, e.g., hard capsules (e.g., hard gelatin capsules or hard hydroxypropyl methylcellulose capsules), soft gelatin capsules, tablets, caplets, enteric coated tablets, chewable tablets, enteric coated hard gelatin capsules, enteric coated soft gelatin capsules, minicapsules, lozenges, films, strips, gelcaps, dragees, solutions, emulsions, suspensions, syrups, or sprays.

[00237] Subjects may be administered a therapeutic amount of a polypeptide, such as 0.01 mg/kg, 1.0 mg/kg, or 15 mg/kg. For administration to human subjects, the dosage of polypeptides of the present invention is typically 0.01 to 15 mg/kg, more preferably 3 to 5 mg/kg. However, dosage levels can be highly dependent on the nature of the condition; drug efficacy; the condition of the patient; the judgment of the practitioner; and the frequency and mode of administration.

[00238] In other embodiments, the polypeptides are administered at a frequency of e.g., every 4 hr, every 6 hr, every 12 hr, every 18 hr, every 24 hr, every 36 hr, every 72 hr, every 84 hr,

every 96 hr, every 5 days, every 7 days, every 10 days, every 14 days, every 3 weeks, or more. The compositions can be administered once daily or the polypeptide can be administered as two, three, or more sub-doses at appropriate intervals throughout the day or delivery through a controlled release formulation. In that case, the polypeptide contained in each sub-dose must be correspondingly smaller in order to achieve the total daily dosage. The dosage unit can also be compounded for delivery over several days, e.g., using a conventional sustained release formulation, which provides sustained release of the polypeptide over a several-day-period.

[00239] Sustained release formulations are well known in the art and are particularly useful for delivery of agents to a particular site, such as could be used with the polypeptide compositions of the present invention. The effect of a single dose can be long-lasting, such that subsequent doses are administered at not more than 3-, 4-, or 5-day intervals, or at not more than 1, 2-, 3-, or 4-week intervals.

[00240] The polypeptide can be administered by intravenous infusion over a period of time, such as over a 5 minute, 10 minute, 15 minute, 20 minute, or 25 minute period. The administration may be repeated, for example, on a regular basis, such as biweekly (*i.e.*, every two weeks) for one month, two months, three months, four months or longer. After an initial treatment regimen, the treatments can be administered on a less frequent basis. For example, after administration biweekly for three months, administration can be repeated once per month, for six months or a year or longer. Administration of the polypeptide or composition can reduce, lower, increase or alter binding or any physiologically deleterious process, e.g., in a cell, tissue, blood, urine or other compartment of a patient by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80 % or at least 90% or more.

[00241] Before administration of a full dose of the polypeptide and/or polypeptide composition, patients can be administered a smaller dose, such as 5% of a full dose, and monitored for adverse effects, such as an allergic reaction or infusion reaction, or for elevated lipid levels or blood pressure. In another example, the patient can be monitored for unwanted immunostimulatory effects, such as increased cytokine (*e.g.*, TNF-alpha, IL-1, IL-6, or IL-10) levels.

[00242] Genetic predisposition plays a role in the development of some diseases or disorders. Therefore, a patient in need of a polypeptide and/or polypeptide composition may be identified

by taking a family history, or, for example, screening for one or more genetic markers or variants. A healthcare provider, such as a doctor, nurse, or family member, can take a family history before prescribing or administering a therapeutic composition of the present invention.

Kits

[00243] Any of the compositions described herein may be comprised in a kit. In a non-limiting example, polypeptides may be included in a kit for treating a disease. The kit may include a vial of sterile, dry polypeptide powder, sterile solution for dissolving the dried powder, and a syringe for infusion set for administering the polypeptide.

[00244] When polypeptides are provided as a dried powder it is contemplated that between 10 micrograms and 1000 milligrams of polypeptide, or at least or at most those amounts are provided in kits of the invention

[00245] The container means will generally include at least one vial, test tube, flask, bottle, syringe and/or other container means, into which the polypeptide formulations are placed, preferably, suitably allocated. The kits may also comprise a second container means for containing a sterile, pharmaceutically acceptable buffer and/or other diluent.

[00246] A kit can include instructions for employing the kit components as well the use of any other reagent not included in the kit. Instructions may include variations that can be implemented.

[00247] While various embodiments of the invention have been particularly shown and described, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended claims.

EQUIVALENTS AND SCOPE

[00248] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments in accordance with the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the appended claims.

[00249] In the claims, articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions

that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[00250] It is also noted that the term “comprising” is intended to be open and permits but does not require the inclusion of additional elements or steps. When the term “comprising” is used herein, the terms “consisting of” and “or including” are thus also encompassed and disclosed.

[00251] Where ranges are given, endpoints are included. Furthermore, it is to be understood that unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or subrange within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

[00252] In addition, it is to be understood that any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Since such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the compositions of the invention (e.g., any nucleic acid or protein encoded thereby; any method of production; any method of use; etc.) can be excluded from any one or more claims, for any reason, whether or not related to the existence of prior art.

[00253] All cited sources, for example, references, publications, databases, database entries, and art cited herein, are incorporated into this application by reference, even if not expressly stated in the citation. In case of conflicting statements of a cited source and the instant application, the statement in the instant application shall control.

[00254] Section and table headings are not intended to be limiting.

EXAMPLES

Example 1. Preparation of biotinylated C5

[00255] An effective final molar ratio of 1:4 C5 to biotin was used for large scale biotinylation. A 10 mM solution of EZ-Link Sulfo-NHS-LC Biotin (Thermo Scientific, Billerica, MA) was prepared as per the manufacturer's instructions. To 1 mg of 1 mg/ml C5 (Complement Tech, Tyler TX), 2.1 μ l of the 10mM biotin solution was added and incubated on ice for 2 hours. The reaction was quenched for 30 minutes at 4°C after the addition of 100 μ l of 1 M Tris HCl pH 7.5. The reaction was dialyzed overnight against cold PBST (phosphate buffered saline (PBS) + 0.1% Tween 80). The biotinylated-C5 was aliquotted and stored at -80°C. Biotinylated C5 was characterized by SDS-PAGE under reducing and non-reducing conditions and characterized for activity by a red blood cell hemolysis assay. Biotinylated C5 was also checked for recovery on streptavidin beads (Invitrogen, Grand Island, NY). Capture was performed using conditions recommended by the manufacturer. To capture 4 μ g of biotinylated-C5 from a 100 nM solution, 40 μ l of bead slurry was used and incubated at 4°C for 1 hour. The concentration of the captured biotinylated C5 was calculated by running a known amount of C5 on a NuPage 4-12% Bis-Tris gel (Invitrogen, Grand Island, NY).

Example 2. Human hemolysis assay for QC of biotinylated C5

[00256] A hemolysis assay was performed with C5-depleted sera and biotinylated C5 and non-biotinylated C5 to compare the lysis activities of C5 before and after biotinylation. Antibody-sensitized sheep erythrocytes (Complement Technology, Tyler TX) in solution at 5×10^8 cell/ml were centrifuged at 2,090 x gravity for 3 minutes and resuspended in GVB++ buffer (Complement Technology, Tyler TX). C5-depleted human sera (Complement Technology, Tyler TX) was rapidly thawed at 37°C and placed on ice until diluted in GVB++. Non-biotinylated C5 protein (Complement Technology, Tyler TX) and biotinylated C5 protein (in-house biotinylation) was rapidly thawed at 37°C and placed on a wet ice slurry until diluted in GVB++. 100 μ l of cells (at a final concentration of 2.5×10^7 cells/ml) was combined with C5-depleted human sera and 50 μ l biotinylated C5 or non-biotinylated C5 (with final concentrations of either 10 μ g/ml, 3 μ g/ml or 1 μ g/ml) in a tissue culture-treated clear 96-well microtitre plate (USA Scientific, Ocala, FL). The plate was incubated for 1 hour at 37°C. After incubation, plates were then centrifuged at 2,090 x gravity for 2 minutes before transferring 100 μ l of supernatant to a new microtitre plate. Absorbance was read at 412 nm and percent lysis activity of non-biotinylated C5 and biotinylated C5 was compared.

Example 3. Selection of polypeptides binding C5

[00257] C5 inhibitors were identified through several rounds of mRNA display and selection. mRNA display was performed generally as described (Roberts, R.W., and Szostak, J.W. (1997). Proc. Natl. Acad. Sci. USA 94, 12297–12302; WO2009067191; herein incorporated by reference in its entirety) with modifications as described herein. RNA pools, were generated by in vitro transcription from DNA synthesized with fixed N-terminal methionine and cysteine codons, followed by three positions of a sixteen codon phosphoramidite mixture, followed by eight positions of a second codon mixture also containing the cysteine codon. The resulting mRNA library has a fixed initiating methionine followed by a cysteine residue, followed by three positions lacking cysteine, followed by eight positions in which cysteine occurs with a frequency of 12.5%. To conduct the selection, the first round of enrichment comprised a first step in which RNA pools containing a 3' terminal UV cross-linked oligonucleotide containing puromycin were translated *in vitro* with the purified translation components listed in Table 2. Translation was carried out under two separate conditions to generate two unique libraries based on amino acid variation. The first condition utilized only the 20 natural amino acids while the second condition utilized natural amino acids (0.1 mM of histidine, threonine, proline, lysine, asparagine, tyrosine, glutamic acid and cysteine), unnatural amino acids (2 mM tertbutyl-glycine (Tbg), 0.8 mM 7-azatryptophan (abbreviated by “azaTrp” in this example) and 1 mM norvaline (Nvl), azaleucine and phenyl-glycine (Phg)) and N-methyl amino acids (450 µM mix of N-methylated serine [(N-Me)S], alanine [(N-Me)A], glycine [(N-Me)G] and 4-fluoro-N-methylphenylalanine [(N-Me-4-F)Phe]). ³⁵S-labeled cysteine residues were included in both conditions to enable monitoring of polypeptide enrichment per round.

Table 2. *In vitro* translation components

Component	Conc.
Creatine phosphate	20 mM
MeTHF, pH 7.6	15 µg/ml
HEPES-KOH, pH 7.6	51 mM
KCl	101 mM
Spermidine	2 mM
DTT	1 mM
Creatine kinase	4 mM
Myokinase	3 mM
Nucleotide diphosphate kinase	1 mM

Pyrophosphate	1 mM
ATP+GTP*	2 mM each
EF-Tu	50 μ M
Ribosomes	1 μ M
MTF	0.56 μ M
1F1	0.96 μ M
IF2	0.40 μ M
IF3	0.44 μ M
EF-G	0.64 μ M
EF-Ts	1.58 μ M
RF1	0.24 μ M
RF3	0.17 μ M
RRF	0.46 μ M
Mg	17.46 mM

[00258] The tRNAs were enzymatically charged with their respective amino acids using tRNA synthetases. The four N-methyl tRNAs were pre-charged, whereas all other tRNAs were enzymatically charged during the *in vitro* translation reaction. The tRNA synthetases were added on a volume basis irrespective of their concentrations. 0.1 μ l of each tRNA synthetase (except for methionine tRNA synthetase, which was added at 0.4 μ l per 25 μ l translation reaction) was added for in-situ charging during translation for a 25 μ l translation reaction. Cross-linked mRNA was added at a final concentration of 0.75 μ M. The translation reaction was kept at 37°C for 1 hour. After translation, the fusion of the translated polypeptides to their respective mRNAs was carried out by adding high salt to the translation mix and incubating at 37°C for 1.5 hours. A library for the selection of natural polypeptides was prepared from eight individual libraries with a fixed cysteine codon in positions 5-11. The random positions in these libraries, with all 20 amino acids possible, were made combinatorially with repeating codon units of NNS (N is A,G,C, or T; S is G or C) (Devlin, J.J., et. al., (1990). Science 249, 404-406.) The translation of these libraries into natural polypeptides was done using a rabbit reticulocyte *in vitro* translation kit rather than the reconstituted system described above.

[00259] Recovery of the mRNA-displayed polypeptides was done using both Oligo dT and Ni-NTA affinity, to isolate fusion molecules containing both polyA mRNA and His tagged polypeptides. Oligo dT bead-bound polypeptides were then cyclized with dibromoxylene as described by others (J. Am. Chem. Soc. 127:1 1727 (2005)).

[00260] Direct selection of the polypeptides by target affinity was then performed. mRNA-displayed polypeptides were allowed to bind for 1 hour at 4°C to biotinylated C5 in a 100 nM

solution of biotinylated C5 in PBST. The RNA corresponding to the affinity selected polypeptides was reverse transcribed and PCR amplified to create a double-stranded DNA pool. The DNA pool was *in vitro* transcribed to generate mRNA, and the mRNA produced was cross-linked as before at its 3' terminus with a puromycin-containing oligonucleotide. The mRNA-puromycin fusions were subjected to *in vitro* translation to generate the second round of the library, which is now enriched in polypeptides that bind complement component C5. The selection cycle was repeated for six rounds. After the sixth round, the DNA pool representing the selected polypeptides was cloned and sequenced, and the amino acid sequences of candidate C5 inhibitors were determined based on the DNA sequences. The polypeptide sequences identified are listed in Table 3.

[00261] As used in all of the following tables as well as in the sequence listing, abbreviations have the following meaning: “Ac” and “NH2” indicate acetyl and amidated termini, respectively; “Nvl” stands for norvaline; “Phg” stands for phenylglycine; “Sar” stands for sarcosine; “Tbg” stands for tert-butylglycine; “Trt” stands for trityl or triphenylmethyl; “Chg” stands for cyclohexylglycine; “(N-Me)X” stands for the N-methylated form of the amino acid indicated by the letter or three letter abbreviation for that amino acid in place of variable “X” written as N-methyl-X [e.g. (N-Me)A and (N-Me)Ala both stand for the N-methylated form of alanine or N-methyl-alanine]; 7-aza-tryptophan is incorporated where “azaTrp” is indicated; “(4-F)Phe” stands for 4-fluorophenylalanine; “Tyr(OMe)” stands for O-methyl tyrosine, “Aib” stands for amino isobutyric acid; “(homo)F” or “(homo)Phe” stands for homophenylalanine; “(2-OMe)Phg” refers to 2-O-methylphenylglycine; “PropargylGly” refers to propargyl-glycine; “(5-F)W” or “(5-F)Trp” refers to 5-fluorotryptophan; “D-X” refers to the D-stereoisomer of the given amino acid “X” wherein the amino acid may be abbreviated using single or three letter code [e.g. (D-Chg) stands for D-cyclohexylglycine and (D-W) stands for D-tryptophan]; “(5-MeO)W” or “(5-MeO)Trp” refers to 5-methyl-O-tryptophan; “homoC” refers to homocysteine; “(1-Me-W)” or “(1-Me)W” or “(1-Me-Trp)” or “(1-Me)Trp” refers to 1-methyltryptophan; “Nle” refers to norleucine; 1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid is incorporated where “Tiq” is indicated; “Asp(T)” refers to (S)-2-amino-3-(1*H*-tetrazol-5-yl)propanoic acid; “(3-Cl-Phe)” refers to 3-chlorophenylalanine; “[N-Me-4-F)Phe]” or “[N-Me-4-F)Phe” refers to N-methyl-4-fluorophenylalanine; “Boc” is a tert-Butyloxycarbonyl protecting group; “[xXyl(y, z)]” refers to the xylyl bridging moiety between two cysteines where x may be m, p or o to

indicate the use of meta-, para- or ortho- dibromoxylanes (respectively) to generate bridging moieties and the numerical identifiers, y and z , place the amino acid position within the polypeptide of the cysteines participating in the cyclization; “[cyclo(y,z)]” refers to the formation of a bond between two residues where the numerical identifiers, y and z , place the position of the residues participating in the bond; “[mXyl-bicyclo]” indicates that the polypeptide comprises two cyclic loops and that the bridging moiety is generated by reaction with a meta-dibromoxylene. All other symbols refer to the standard one-letter amino acid code. Additionally, polypeptides comprising PEG2000 or BODIPY-TMR-X sequence tags are indicated.

Table 3. Polypeptide Sequences

Compound Number	Sequence	SEQ ID NO
R3000	Ac-Nvl-C-Y-K-N-Y-H-azaTrp-E-Y-P-Tbg-Y-NH2	1
R3001	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-(N-Me)G-Nvl-(N-Me)S-NH2	2
R3002	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	3
R3003	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-P-NH2	4
R3004	Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-Y-P-Phg-Tbg-NH2	5
R3005	Ac-Nvl-C-Y-azaTrp-(N-Me)G-Tbg-Nvl-azaTrp-E-Y-P-Phg-P-NH2	6
R3006	Ac-Y-E-N-Tbg-Y-azaTrp-E-Y-(N-Me)G-Nvl-(N-Me)S-NH2	7
R3007	[mXyl(2,7)]Ac-Nvl-C-K-E-Phg-Y-C-(N-Me)S-Tbg-K-azaTrp-E-Y-NH2	8
R3008	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	9
R3020	[mXyl(2,7)]M-C-S-E-R-Y-C-E-V-R-W-E-Y-NH2	10
R3021	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	11

Example 4. Polypeptide synthesis

[00262] Polypeptides are synthesized using standard solid-phase Fmoc/tBu methods. The synthesis is typically performed on a Liberty automated microwave peptide synthesizer (CEM, Matthews NC) using standard protocols with Rink amide resin, although other automated synthesizers without microwave capability may also be used. All amino acids are obtained from commercial sources unless otherwise noted. The coupling reagent used is 2-(6-chloro-1-H-benzotriazole-1yl)-1,1,3,3,-tetramethylaminium hexafluorophosphate (HCTU) and the base is diisopropylethylamine (DIEA). Polypeptides are cleaved from resin with 95% TFA, 2.5% TIS

and 2.5% water for 3 hours and isolated by precipitation with ether. The crude polypeptides are purified on a reverse phase preparative HPLC using a C18 column, with an acetonitrile/water 0.1% TFA gradient from 20%-50% over 30 min. Fractions containing the pure polypeptide are collected and lyophilized and all polypeptides are analyzed by LC-MS.

Example 5. Formation of disulfide cyclized polypeptides

[00263] To produce disulfide cyclized polypeptides, the linear polypeptide is dissolved in a mixture of water and DMSO and the resulting solution is stirred vigorously under an air atmosphere for 12 hrs.

Example 6. Dibromoxylene polypeptide cyclization

[00264] A 100 mL flask is charged with acetonitrile (12 mL) and water (24 mL) and is degassed with argon for about 5 min. Linear polypeptide (0.01 mmole) and 200 mM ammonium bicarbonate (6mL) are added followed by 0.012 mmole or 1,3-bis(bromomethyl) benzene, 1,2-bis(bromomethyl)benzene, 1,4-bis(bromomethyl)benzene, 2,6-bis(bromomethyl)pyridine or (E)-1,4-dibromobut-2-ene. The reaction mixture is stirred under argon at room temperature for approximately 2 hours and monitored by LC-MS. After the reaction is complete, the reaction solution is frozen and lyophilized. HPLC purification of the crude lyophilized product followed by lyophilization of fractions containing pure polypeptide yield the final cyclized product as a white powder.

Example 7. Lactam polypeptide cyclization

[00265] Cyclization of polypeptides using a lactam moiety was performed in the solid phase. A polypeptide was first synthesized on a solid support Wang resin by standard Fmoc chemistry. Fmoc-ASP(allyl)-OH and Fmoc-LYS(allo)-OH were incorporated in the polypeptide at the indicated positions as the two precursor monomers for the lactam bridge formation. After full elongation the resin was washed with dry dichloromethane (3x) and purged with dry Nitrogen gas for 10 min. To remove the allyl and allo protecting groups, the resin was treated with a 5 fold molar excess of phenylsilane and purged with Nitrogen for 10 min. A catalytic amount of tetrakis Pd(0) was dissolved in dry dichloromethane and added to the suspension of resin. After one hour the resin was washed sequentially with dichloromethane (3x), dimethylformamide (3x), sodium diethyldithiocarbamate trihydrate (3x), dimethylformamide (3x) and dichloromethane

(3x). Lactam cyclization was achieved in dimethylformamide (DMF) by treating the deprotected polypeptide containing resin with PyAOP ((3-Hydroxy-3H-1,2,3-triazolo[4,5-b]pyridinato-O)tri-1-pyrrolidinyl- phosphorus hexafluorophosphate) and diisopropylethylamine and allowed to react overnight. The resin was rinsed with DMF and treated with fresh PyAOP and diisopropylethylamine for additional 60 minutes at 45 °C. The resin was rinsed and washed with dimethylformamide five times. The polypeptide was cleaved and purified as described in example 4.

Example 8. Triazole polypeptide cyclization

[00266] Cyclization of polypeptides containing an azide and an alkyne moiety was performed on the solid phase. Polypeptide containing resin (0.05 mmol) was treated with dichloromethane and allowed to swell for 10 min. The solvent was then exchanged to DMF (3-5 mL) and after 10 min, a solution of Cu-TBTA ligand was added (125 µL of a 20 mM solution). The suspension was purged with Argon gas and then ascorbic acid (5 µmoles) was added. The solution was shaken for 2h and the excess reagents removed, the resin was washed with a solution of EDTA in DMF to remove excess copper. The polypeptide was cleaved and purified as described in Example 4.

Example 9. Polypeptides of the current invention

[00267] Polypeptides of the current invention were synthesized. These include the compounds listed in Table 4.

Table 4. Compounds of the invention

Compound Number	Sequence	SEQ ID NO.
R3000	Ac-Nvl-C-Y-K-N-Y-H-azaTrp-E-Y-P-Tbg-Y-NH2	1
R3001	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-(N-Me)G-Nvl-(N-Me)S-NH2	2
R3002	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	3
R3003	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-P-NH2	4
R3004	Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-Y-P-Phg-Tbg-NH2	5
R3005	Ac-Nvl-C-Y-azaTrp-(N-Me)G-Tbg-Nvl-azaTrp-E-Y-P-Phg-P-NH2	6
R3006	Ac-Y-E-N-Tbg-Y-azaTrp-E-Y-(N-Me)G-Nvl-(N-Me)S-NH2	7
R3007	[mXyl(2,7)]Ac-Nvl-C-K-E-Phg-Y-C-(N-Me)S-Tbg-K-azaTrp-E-Y-NH2	8
R3008	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	9

R3020	[mXylyl(2,7)]M-C-S-E-R-Y-C-E-V-R-W-E-Y-NH2	10
R3021	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	11
R3079	Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	12
R3055	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	13
R3120	Ac-Nvl-Nvl-Y-E-(N-Me)N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	14
R3057	[mXylyl(2,7)]M-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	15
R3056	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	16
R3054	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	17
R3029	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-NH2	18
R3048	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	19
R3072	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-K-NH2	20
R3024	Ac-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	21
R3114	Ac-Nvl-Nvl-(N-Me)Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	22
R3050	[pXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	23
R3025	Ac-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	24
R3061	Ac-Nvl-S-Y-E-A-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	25
R3041	Ac-Y-E-N-Tbg-Y-W-E-Y-P-Phg-Nvl-NH2	26
R3077	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-K-(PEG2000) NH2	27
R3030	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-NH2	28
R3062	Ac-Nvl-S-Y-E-N-A-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	29
R3066	Ac-Nvl-S-Y-E-N-Tbg-A-azaTrp-E-Y-P-Chg-Nvl-NH2	30
R3011	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P- NH2	31
R3070	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-Y-A-Chg-Nvl-NH2	32
R3071	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-Y-P-A-Nvl-NH2	33
R3033	[mXylyl(2,10)]Ac-Nvl-C-Phg-A-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	34
R3038	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	35
R3012	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	36
R3060	Ac-Nvl-S-Y-A-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	37
R3039	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-A-NH2	38
R3037	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)A-H-C-Nvl-NH2	39
R3076	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-K-(BODIPY-TMR-X) NH2	40
R3074	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Tyr(OMe)-(N-Me)S-H-C-Nvl-NH2	41
R3013	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-NH2	42
R3065	[pXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-P-H-C-Nvl-NH2	43
R3073	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Phe(4-F)-(N-Me)S-H-C-Nvl-NH2	44
R3116	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-(N-Me)W-E-Y-P-Chg-Nvl-NH2	45
R3091	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-W-E-Y-(N-Me)S-A-C-Nvl-NH2	46
R3078	PEG2000-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	47
R3100	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-F-(N-Me)S-A-C-Nvl-NH2	48
R3121	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-Y-P-(N-Me)Phg-Nvl-NH2	49
R3043	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-NH2	50
R3102	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-P-H-C-Nvl-NH2	51

R3026	Ac-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	52
R3031	[mXylyl(2,10)]Ac-A-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	53
R3019	[mXylyl(2,14)]Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-C-NH2	54
R3014	[mXylyl(1,9)]Ac-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	55
R3104	[pXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	56
R3059	Ac-Nvl-S-A-E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	57
R3115	Ac-Nvl-Nvl-Y-(N-Me)E-N-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	58
R3110	Ac-Y-E-N-Tbg-Y-(1-Me)W-E-Y-P-Phg-Nvl-NH2	59
R3126	Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-C-P-Phg-Tbg-NH2	60
R3049	[oXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	61
R3069	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-E-A-P-Chg-Nvl-NH2	62
R3015	[mXylyl(1,9)]Ac-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-NH2	63
R3068	Ac-Nvl-S-Y-E-N-Tbg-Y-azaTrp-A-Y-P-Chg-Nvl-NH2	64
R3105	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	65
R3106	[pXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	66
R3111	[mXylyl(4,10)]Ac-Nvl-T-Phg-C-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	67
R3112	[mXylyl(2,10)]Ac-Nle-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	68
R3113	[mXylyl(3,11)]Ac-Y-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	69
R3134	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-(3-Cl-Phe)-(N-Me)S-A-C-Nvl-NH2	70
R3018	[mXylyl(2,10)]Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-C-P-Phg-Nvl-NH2	71
R3027	Ac-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	72
R3028	Ac-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	73
R3032	[mXylyl(2,10)]Ac-Nvl-C-A-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	74
R3058	[pXylyl(2,10)]Ac-Nvl-C-Chg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	75
R3067	Ac-Nvl-S-Y-E-N-Tbg-Y-A-E-Y-P-Chg-Nvl-NH2	76
R3117	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-azaTrp-E-(N-Me)Y-P-Chg-Nvl-NH2	77
R3022	Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	78
R3016	[mXylyl(1,9)]Ac-C-Tbg-Y-azaTrp-E-Y-(N-Me)S-H-C-NH2	79
R3089	[mXylyl(2,10)]Ac-Chg-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	80
R3083	[mXylyl(2,10)]Ac-V-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	81
R3087	[mXylyl(2,10)]Ac-Nvl-C-(2-OMe)Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	82
R3103	[mXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	83
R3135	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-(D-Ala)-C-Nvl-NH2	84
R3034	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-A-E-Y-(N-Me)S-H-C-Nvl-NH2	85
R3035	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-A-Y-(N-Me)S-H-C-Nvl-NH2	86
R3036	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-A-(N-Me)S-H-C-Nvl-NH2	87
R3044	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-NH2	88
R3080	[mXylyl(2,9)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-C-Nvl-NH2	89

R3085	[mXylyl(2,10)]heptanoyl-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	90
R3086	[mXylyl(5,13)]Ac-Nvl-S-Y-E-C-Tbg-Y-azaTrp-E-Y-P-Chg-C-Nvl-NH2	91
R3092	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-F-E-Y-(N-Me)S-A-C-Nvl-NH2	92
R3095	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-(homo)F-(N-Me)S-A-C-Nvl-NH2	93
R3096	[mXylyl(2,10)]Ac-Nvl-C-Phg-Aib-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	94
R3122	[mXylyl(2,10)]Ac-Nvl-C-Tiq-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	95
R3075	[mXylyl(2,11)]Nvl-C-Y-(N-Me)S-Phg-(N-Me-4-F)Phe-(N-Me)S-H-(N-Me-4-F)Phe-G-C-NH2	96
R3107	[mXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	97
R3108	[pXylyl(2,10)]Ac-Nvl-homoC-Phg-T-azaTrp-E-Y-(N-Me)S-H-homoC-Nvl-NH2	98
R3127	[mXylyl(2,10)]Ac-Nvl-C-Y-N-N-Tbg-E-azaTrp-E-C-P-Phg-Tbg-NH2	99
R3133	[mXylyl(2,10)]Ac-Nvl-C-Phg-(D-Ala)-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	100
R3009	[mXylyl(2,10)]Ac-Nvl-C-Y-E-(N-Me)G-Tbg-Y-azaTrp-E-C-Nvl-P-Nvl-NH2	101
R3010	[mXylyl(2,13)]Ac-Nvl-C-Y-E-(N-Me)G-Tbg-Y-azaTrp-E-Nvl-Nvl-P-C-NH2	102
R3017	[mXylyl(2,8)]Ac-Nvl-C-Y-E-N-Tbg-Y-C-E-Y-P-Phg-Nvl-NH2	103
R3023	Ac-Y-P-Y-C-Phg-azaTrp-Tbg-E-Nvl-N-Y-Nvl-E-NH2	104
R3040	[cyclo(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl	105
R3042	[cyclo(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-NH2	106
R3045	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-Y-NH2	107
R3046	[mXylyl(2,7)]M-C-V-E-R-F-C-D-V-NH2	108
R3047	[mXylyl(2,7)]M-C-V-E-R-F-C-NH2	109
R3051	[mXylyl(2,11)]Nvl-C-Y-(N-Me)S-Phg-(N-Me-4-F)Phe-(N-Me)S-H-(N-Me-4-F)Phe-(N-Me)G-C-NH2	110
R3052	[mXylyl(2,9)]Nvl-C-Y-Tbg-Phg-N-(N-Me)G-L-C-Phg-(N-Me)A-NH2	111
R3053	[mXylyl-bicyclo]Nvl-C-C-N-Tbg-Phg-C-Tbg-(N-Me)S-C-Tbg-NH2	112
R3063	Ac-Tbg-Y-azaTrp-E-Y-NH2	113
R3064	Ac-Y-azaTrp-E-Y-P-NH2	114
R3081	Ac-Y-E-N-Tbg-Y-azaTrp-(N-Me)E-Y-P-Phg-Nvl-NH2	115
R3082	[mXylyl(1,9)]heptanoyl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	116
R3084	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-S-A-C-Nvl-NH2	117
R3088	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-(5-F)W-E-Y-(N-Me)S-A-C-Nvl-NH2	118
R3090	[mXylyl(2,10)]Ac-Nvl-C-F-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	119
R3093	[mXylyl(2,10)]Ac-Nvl-C-(D-Chg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	120
R3094	Ac-Y-E-N-Tbg-Y-(5-MeO)W-E-Y-P-Phg-Nvl-NH2	121
R3097	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-D-Y-(N-Me)S-A-C-Nvl-NH2	122

R3098	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-Q-Y-(N-Me)S-A-C-Nvl-NH2	123
R3099	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-N-Y-(N-Me)S-A-C-Nvl-NH2	124
R3101	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-G-C-Nvl-NH2	125
R3109	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-(1-Me-W)-E-Y-(N-Me)S-A-C-Nvl-NH2	126
R3118	Ac-Y-E-N-Tbg-Y-(D-Trp)-E-Y-P-Phg-Nvl-NH2	127
R3119	Ac-Y-E-N-Y-(D-Trp)-E-Y-P-Phg-Nvl-NH2	128
R3123	Ac-Y-E-N-Tbg-Y-azaTrp-(D-Glu)-Y-P-Phg-Nvl-NH2	129
R3124	[mXylyl(1,6)]Ac-C-V-E-R-F-C-V-Y-W-E-F-NH2	130
R3125	[mXylyl(1,6)]Ac-C-V-E-R-F-C-W-E-F-NH2	131
R3128	Ac-Nvl-C-Y-N-N-Tbg-E-C-E-Y-P-Phg-Tbg-NH2	132
R3129	[mXylyl(2,8)]Ac-Nvl-C-Y-N-N-Tbg-E-C-E-Y-P-Phg-Tbg-NH2	133
R3130	Ac-Nvl-Nvl-Y-E-N-Tbg-(N-Me)Y-azaTrp-E-Y-P-Chg-Nvl-NH2	134
R3131	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-W-Asp(T)-Y-(N-Me)S-H-C-Nvl-NH2	135
R3132	[mXylyl(2,10)]Ac-Nvl-C-Phg-T-(D-Trp)-E-Y-(N-Me)S-H-C-Nvl-NH2	136
R3136	[mXylyl(2,10)]heptanoyl-Nvl-C-(D-Phg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	137
R3137	[mXylyl(1,9)]heptanoyl-C-(D-Phg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	138
R3138	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-W-E-F-NH2	139
R3139	[mXylyl(1,6)]Ac-C-Tbg-E-R-F-C-D-Tbg-Y-W-E-F-NH2	140
R3140	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-Y-P-NH2	141
R3141	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-P-NH2	142
R3142	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-azaTrp-E-Y-P-NH2	143
R3143	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-W-E-Y-P-NH2	144
R3144	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	145
R3145	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	146
R3146	[mXylyl(1,6)]Ac-C-Tbg-E-R-F-C-D-V-Y-W-E-F-NH2	147
R3147	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-Propargyl-Gly-NH2	148
R3148	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-W-E-Y-P-Phg-Nvl-NH2	149
R3149	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-W-E-Y-P-(D-Phg)-Nvl-NH2	150
R3150	[mXylyl(1,6)]Ac-C-V-A-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	151
R3151	[mXylyl(1,6)]Ac-C-V-A-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	152
R3152	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	153
R3153	[mXylyl(1,6)]Ac-C-V-E-A-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	154
R3154	[mXylyl(1,6)]Ac-C-V-E-A-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	155
R3155	[mXylyl(1,6)]Ac-C-V-E-R-A-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	156

	NH2	
R3156	[mXylyl(1,6)]Ac-C-V-E-R-A-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	157
R3157	[mXylyl(1,6)]Ac-C-V-E-R-F-C-A-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	158
R3158	[mXylyl(1,6)]Ac-C-V-E-R-F-C-A-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	159
R3159	[mXylyl(1,6)](des-amino)C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	160
R3160	[mXylyl(1,6)](des-amino)C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-Nvl-NH2	161
R3161	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-K-NH2	162
R3162	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-(D-Phg)-K-NH2	163
R3163	[cyclo(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	164
R3164	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C12)-NH2	165
R3165	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C10)-NH2	166
R3166	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C8)-NH2	167
R3167	[mXylyl(1,6)]Ac-C-V-E-R-F-C-(alpha-methyl)D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	168
R3168	[mXylyl(1,6)]Ac-C-V-E-R-F-C-Asp(T)-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	169
R3169	[cyclo(1,6)]Ac-K-V-E-R-F-D-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	170
R3170	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-K	171
R3171	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C12)	172
R3172	[mXylyl(1,6)]Ac-C-V-E-R-F-C-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	173
R3173	[cyclo(1,6)]Ac-K-V-E-R-F-D-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	174
R3174	[cyclo(1,6)]Ac-K-V-E-R-F-D-Asp(T)-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	175
R3175	[cyclo(1,6)]Ac-K-V-E-R-F-D-D-Tbg-Y-azaTrp-E-Y-P-Chg-B20	176
R3176	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	177
R3177	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-W-P-Chg-Nvl	178
R3178	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-(homo)Phe-P-Chg-Nvl	179
R3179	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-(m-Cl-homo)Phe-P-Chg-Nvl	180
R3180	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-2Nal-P-Chg-Nvl	181
R3181	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-(3-aminomethyl)Phe-E-Y-P-Chg-Nvl	182
R3182	[cyclo-triazolyl(1,6)]Ac-X02-V-E-R-F-X31-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	183
R3183	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)	184

R3184	[cyclo-thioalkyl(1,5)]V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	185
R3185	[mXylyl(1,6)]Ac-C-V-E-R-F-C-Cle-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	186
R3186	[mXylyl(1,6)]Ac-C-V-E-R-F-C-(Ac-Pyran)-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	187
R3187	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-(3-aminomethyl)Phe-P-Chg-Nvl	188
R3188	[cyclo-olefinyl(1,6)]Ac-X30-V-E-R-F-X12-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	189
R3189	[mXylyl(1,6)]Ac-C-A-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Phg-(Lys-C16)	190
R3190	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-B20	191
R3191	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-K	192
R3192	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-K-NH2	193
R3193	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-B28	194
R3194	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)-NH2	195
R3195	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-W-E-Y-P-Chg-(Lys-C16)	196
R3196	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-W-E-Y-P-Chg-K	197
R3197	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-W-E-Y-P-Chg-K14	198
R3198	[cyclo(1,6)](desamino)C-V-E-R-F-C-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)	199
R3199	[cyclo(1,6)](desamino)C-(D-Ala)-E-R-F-C-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Chg-(Lys-C16)	200
R3200	[cyclo(1,6)]Ac-K-V-E-R-F-D-(N-Me)D-Tbg-Y-azaTrp-E-Y-P-Aib-(Lys-C16)	201

[00268] Polypeptides R3183 (SEQ ID NO: 184) and R3193 (SEQ ID NO: 194) were synthesized according to the amino acid sequences of R3176 (SEQ ID NO: 177) with the exception of the replacement of Nvl-NH₂ with Lys. The side chain amine group of the Lys residue was modified with the different lipophilic moieties resulting in lipidated polypeptides.

Example 10. Optimization and testing of C5 inhibitors

[00269] Polypeptides selected according to Example 3 and listed in Table 3 were tested for their ability to inhibit complement-mediated cell lysis. Additionally, a variety of optimized polypeptides were synthesized according to the methods of Examples 4-8 and tested as well (see Table 4). Optimized polypeptide sequences include those obtained by making a variety of truncations, deletions, additions and/or substitutions to compounds R3002 (SEQ ID NO: 3),

R3008 (SEQ ID NO: 9) and R3021 (SEQ ID NO: 11) or through the formation of hybrid polypeptides comprising combinations of regions selected from any of the three.

Human hemolysis assay (RBC lysis assay using complete human sera)

[00270] Polypeptides listed in Table 3, as well as their optimized derivatives (see Table 4) were assessed for inhibitor activity using a red blood cell hemolysis assay. Antibody-sensitized sheep erythrocytes (Complement Technology, Tyler, TX) were plated at 2.5×10^7 cells/well with complete human sera (Complement Technology, Tyler TX) and polypeptides to determine the inhibitory effect of the polypeptides on the lysis of red blood cells. Cells were centrifuged for 3 minutes at 2,090 x gravity and resuspended in fresh GVB++ buffer (Complement Technology, Tyler TX). Human sera was rapidly thawed at 37°C and then stored on ice until diluted into GVB++. Ten 6-fold serial dilutions of polypeptides (10mM stock, DMSO) were performed in DMSO and then added to buffer. 50 μ l of each polypeptide dilution was combined with sera and 100 μ l of cells in individual wells of a 96-well tissue culture-treated clear microtitre plate (USA Scientific, Ocala, FL) and resuspended by pipetting. Samples were incubated at 37°C for one hour. Following incubation, plates were centrifuged at 2,090 x gravity for 2 minutes. 100 μ l of supernatant was transferred to a new plate and the absorbance was read at 412 nm. Data was fit with a log-logit formula producing a dose-response curve and IC₅₀. As used herein, the term “IC₅₀” refers to the half maximal inhibitory concentration, a value used to indicate the amount of the inhibitor needed to reduce a given reaction or process by half. Compounds tested are listed in Table 5.

Table 5. Compounds analyzed

Compound Number	Avg. IC ₅₀ (nM)	SEQ ID NO.
R3000	>10,000	1
R3001	67.2	2
R3002	11.9	3
R3003	13.9	4
R3004	53.5	5
R3005	66.7	6
R3006	267	7
R3007	314	8
R3008	97	9

R3009	>100,000	101
R3010	>100,000	102
R3011	112	31
R3012	148.5	36
R3013	344	42
R3014	1420	55
R3015	>3,960	63
R3016	>13,000	79
R3017	>100,000	103
R3018	>5,730	71
R3019	1320	54
R3020	24.6	10
R3021	27.5	11
R3022	>10,200	78
R3023	>100,000	104
R3024	32.5	21
R3025	47.5	24
R3026	1020	52
R3027	>10,000	72
R3028	>10,000	73
R3029	18.5	18
R3030	83.6	28
R3031	1090	53
R3032	>10,000	74
R3033	131	34
R3034	>50,000	85
R3035	>50,000	86
R3036	>50,000	87
R3037	276	39
R3038	140	35
R3039	240	38
R3040	>100,000	105
R3041	71.3	26
R3042	>100,000	106
R3043	934	50
R3044	>50,000	88
R3045	>100,000	107
R3046	>100,000	108
R3047	>100,000	109
R3048	19.3	19
R3049	>3,100	61
R3050	42.9	23
R3051	>100,000	110
R3052	>100,000	111
R3053	>100,000	112
R3054	14.1	17
R3055	10.4	13
R3056	13.8	16

R3057	12.4	15
R3058	>10,000	75
R3059	2160	57
R3060	161	37
R3061	53.9	25
R3062	89.9	29
R3063	>100,000	113
R3064	>100,000	114
R3065	394	43
R3066	104	30
R3067	>10,000	76
R3068	>4,500	64
R3069	>3,670	62
R3070	123	32
R3071	128	33
R3072	26.9	20
R3073	403	44
R3074	308	41
R3075	>75,000	96
R3076	297	40
R3077	81.7	27
R3078	568	47
R3079	7.3	12
R3080	>50,000	89
R3081	>100,000	115
R3083	>25,000	81
R3084	>100,000	117
R3086	>50,000	91
R3087	>25,000	82
R3088	>100,000	118
R3089	>15,000	80
R3090	>100,000	119
R3091	483	46
R3092	>50,000	92
R3093	>100,000	120
R3094	>100,000	121
R3095	>50,000	93
R3096	>50,000	94
R3097	>100,000	122
R3098	>100,000	123
R3099	>100,000	124
R3100	626	48
R3101	>100,000	125
R3102	978	51
R3103	>25,000	83
R3104	>2,000	56
R3105	>5,000	65
R3106	>5,000	66

R3107	>75,000	97
R3108	>75,000	98
R3109	>100,000	126
R3110	2940	59
R3111	>5,000	67
R3112	>5,000	68
R3113	>5,000	69
R3114	36.6	22
R3115	2780	58
R3116	441	45
R3117	>10,000	77
R3118	>100,000	127
R3119	>100,000	128
R3120	12.2	14
R3121	804	49
R3122	>50,000	95
R3123	>100,000	129
R3124	>100,000	130
R3125	>100,000	131
R3126	>3000	60
R3127	>75,000	99
R3128	>100,000	132
R3129	>100,000	133
R3130	>100,000	134
R3131	>100,000	135
R3132	>100,000	136
R3133	>75,000	100
R3134	>5,000	70
R3135	>25,000	84
R3136	>50,000	137
R3137	>100,000	138
R3138	87.2	139
R3139	97.2	140
R3140	17.9	141
R3141	24.5	142
R3142	44.6	143
R3143	18.6	144
R3144	6.7	145
R3145	39	146
R3146	107	147
R3147	138	148
R3148	8.5	149
R3149	13.6	150
R3150	32	151
R3151	165	152
R3152	11	153

R3153	175	154
R3154	592	155
R3155	1530	156
R3156	>10,000	157
R3157	84.5	158
R3158	327	159
R3159	7.6	160
R3160	37.1	161
R3161	7	162
R3162	16.5	163
R3163	17	164
R3164	36	165
R3165	18.5	166
R3166	17.5	167
R3167	11	168
R3168	7.5	169
R3169	5	170
R3170	4.5	171
R3172	12	173

Example 11. Alternative human hemolysis assay using C5 depleted sera

[00271] Polypeptides listed in Table 6 were tested for functional activity in a red blood cell hemolysis assay using human C5 depleted sera and purified human C5 rather than complete human sera. To assess activity, antibody-sensitized sheep erythrocytes (Complement Technology, Tyler, TX) were plated 2.5×10^7 cells/well with 1.5% human C5 depleted sera (Complement Technology, Tyler, TX) and 0.5nM purified human C5 (Complement Technology, Tyler, TX). The antibody-sensitized sheep erythrocytes were centrifuged at 2,090 x gravity for 3 minutes and then resuspended in fresh GVB ++ (Complement Technology, Tyler, TX). The human C5 depleted sera and purified human C5 were rapidly thawed at 37°C and then stored on ice or wet ice, respectively. The polypeptide stock (10mM, DMSO) was serially diluted in DMSO in order to obtain 10 6-fold dilutions and then GVB++ was added to them. 50 μ l of each polypeptide dilution was combined with 25 μ l C5 depleted sera, 25 μ l purified human C5 and 100 μ l cells in individual wells of a 96-well tissue culture-treated clear microtitre plate (USA Scientific, Ocala, FL) and resuspended by pipetting. The samples were incubated at 37°C for one hour. At the completion of the incubation, the plates were centrifuged at 2,090x gravity for 2

minutes. 100 μ l of supernatant was transferred to a new plate and the absorbance was read at 412 nm. Data was fit with a log-logit formula producing a dose-response curve and IC₅₀.

Table 6. Compounds analyzed

Compound Number	Avg. IC ₅₀ (nM)	SEQ ID NO.
R3171	5.67	172
R3173	2.5	174
R3174	2.3	175
R3176	1.1	177
R3177	12	178
R3179	83	180
R3180	29	181
R3181	1496	182
R3182	13	183
R3183	13.25	184
R3184	4	185
R3185	12.5	189
R3186	18	187
R3189	81.5	190
R3190	35.33	191
R3191	2.5	192
R3192	1.5	193
R3193	24	194
R3194	15.5	195
R3195	62.5	196
R3196	3	197
R3197	4	198
R3198	142	199
R3199	112	200
R3200	88.5	201

Example 12. Enzyme immunoassay to assess C5 inhibition

[00272] C5 inhibitory activity was assessed by enzyme immunoassay (EIA). Inhibition of the production of C5a and the membrane attack complex (MAC) were measured by MicroVue EIA kits (Quidel Corporation, San Diego, CA).

C5a EIA

[00273] Supernatant from a human RBC hemolysis assay of R3002 (SEQ ID NO: 3) and R3008 (SEQ ID NO: 9) was diluted 1:50 and assayed by C5a EIA (Figure 1). Both polypeptides inhibited the formation of C5a. R3002 (SEQ ID NO: 3) had an IC₅₀ of 5.4 nM, R3008 (SEQ ID NO: 9) had an IC₅₀ of 54.5 nM.

Membrane attack complex (MAC) EIA

[00274] The MAC EIA was performed on the diluted supernatant (1:5) of R3008 (SEQ ID NO: 9) from a human RBC hemolysis assay (Figure 2). This polypeptide was shown to inhibit the formation of the MAC with an IC₅₀ of 33 nM.

Example 13. Characterization of peptidomimetic binding by fluorescence polarization

[00275] Fluorescence polarization (FP) allows binding events to be measured in a homogenous solution (Banks, P. et al., Impact of a red-shifted dye label for high throughput fluorescence polarization assays of G protein-coupled receptors. *J Biomol Screen.* 2000 Oct;5(5):329-34 and Parker, G.J. et al., Development of high throughput screening assays using fluorescence polarization: nuclear receptor-ligand-binding and kinase/phosphatase assays. *J Biomol Screen.* 2000 Apr;5(2):77-88). The key concept of FP is that the degree by which a fluorophore polarizes light is inversely related to its molecular rotation (Lea, W.A. et al., Fluorescence polarization assays in small molecule screening. *Expert Opin Drug Discov.* 2011 Jan;6(1):17-32), and a fluorophore bound to a much larger target protein rotates more slowly than an unbound fluorophore, resulting in an increase in polarization that can be quantified. FP has been used increasingly in high throughput campaigns as a method to measure ligand-target binding (Parker, G.J. et al., Development of high throughput screening assays using fluorescence polarization: nuclear receptor-ligand-binding and kinase/phosphatase assays. *J Biomol Screen.* 2000 Apr;5(2):77-88), for equilibrium dissociation constant (K_D) determination (Prystay, L. et al., Determination of equilibrium dissociation constants in fluorescence polarization. *J Biomol Screen.* 2001 Jun;6(3):141-50), and lead discovery through competitive binding assays (Tian, W. et al., Development of novel fluorescence polarization-based assay for studying the β-catenin/Tcf4 interaction. *J Biomol Screen.* 2012 Apr;17(4):530-4).

Materials and methods

[00276] FP was used for screening competitive polypeptide inhibitors of the C5 protein. Probe R3076 (SEQ ID NO: 40) was generated by incubating the parent polypeptide, R3072 (SEQ ID NO: 20) with BODIPY-TMR-X, SE (Life Technologies, Grand Island, NY) in DMF (Sigma, Saint Louis, MO) for 4 hours. The BODIPY-TMR dye attached to the C-terminal lysine of the protein and the subsequent labeled probe was purified by HPLC.

[00277] The equilibrium dissociation constant (K_D) for binding of R3076 (SEQ ID NO: 40) to human C5 protein (Complement Technology, Tyler, TX) was determined by incubating a solution of 25 nM R3076 (SEQ ID NO: 40) with increasing concentrations of C5 protein. Polarization was measured over time, until binding reached equilibrium. K_D was determined using Graphpad Prism (using “Saturating Binding Curves, One Site – Specific Binding With Hill Slope” as the curve fit to determine K_D). Equilibrium was reached after 10 minutes, with values for K_D , hill slope and maximal binding remaining stable over 60 minutes. A final K_D value of 8.07 nM (0.53 standard deviation) was determined by averaging K_D values from 10 to 60 minutes. Based on this information, 25 nM and 50 nM concentrations for R3076 (SEQ ID NO: 40) and C5 protein, respectively, were chosen for use in the competition assay. These concentrations represented the approximate level of protein necessary for 95% probe binding to C5 protein. R3023 (SEQ ID NO: 104) is a scrambled polypeptide variant of R3002 (SEQ ID NO: 3) and was included in all assays as a negative control.

[00278] Human C5 protein was diluted to 200 nM in assay buffer, composed of TBS (EMD Millipore, Billerica, MA) + 0.005% Triton-X (Sigma, Saint Louis, MO). 10 μ l of assay buffer was added to all wells of a black, non-binding, 384-well assay plate (Greiner, Monroe, NC) and 10 μ l of diluted C5 protein stock was added to experimental and designated control wells.

[00279] Probe R3076 (SEQ ID NO: 40) was diluted 1 to 10 in DMSO (Life Technologies, Grand Island, NY) and 30 μ l of that stock was diluted in 3 ml of assay buffer to yield a 100 nM stock. 10 μ l of this working stock was then added to each well in the assay plate. The assay plate was incubated at room temperature, protected from light, for 20 minutes to allow binding to reach equilibrium.

[00280] Test articles listed in Table 7 were subsequently diluted in DMSO then assay buffer, comprising 10 2-fold dilutions and were then added to the assay plate in triplicate, rapidly. The assay plate was then incubated in the Paradigm (Molecular Devices, Sunnyvale, CA) plate reader for 60 minutes at 25°C.

[00281] After incubation for 60 minutes, the plate was read using the Paradigm FP protocol (Molecular Devices, Sunnyvale, CA) and raw polarization values were imported into Graphpad Prism. K_i (using the One Site K_i Curve fitting model, probe concentration = 25 nM, K_D = 8.07 nM, with baseline constrained to the average of the 0% binding control) and IC_{50} (log inhibitor vs response, 4 parameter curve fit) were determined in Graphpad.

Results

[00282] All test articles were able to compete with the labeled probe for binding to human C5 protein (Figure 3, Table 7). R3003 (SEQ ID NO: 4) was the most potent polypeptide tested, with a K_i value of 9.54 nM. R3023 (SEQ ID NO: 104) binding was not detected at the highest concentration tested.

Table 7. Competitive fluorescence polarization data

Compound Number	SEQ ID NO	K_i nM, Exp 1	K_i nM, Exp 2	IC_{50} nM, Exp 1	IC_{50} nM, Exp 2	Avg, nM K_i	Std. Dev. K_i	Avg, nM IC_{50}	Std. Dev. IC_{50}
R3003	4	10.39	8.69	72.65	64.06	9.54	1.20	68.36	6.07
R3011	31	73.17	86.91	294.8	261.3	80.04	9.72	278.1	23.69
R3014	55	1405	1585	6064	7579	1495	127.3	6822	1071
R3023	104	>1000	>1000	>1000	>1000	>1000	--	>1000	--
R3043	50	866.1	882.1	4332	4122	874.1	11.31	4227	148.5
R3050	23	71.08	57.6	266.9	292.7	64.34	9.53	279.8	18.24

[00283] Data shown in Table 7 were obtained from curve fitting analysis performed by Graphpad Prism software as described above. Triplicate values were averaged to yield the data points presented in each experiment. Of the polypeptides tested, R3003 (SEQ ID NO: 4) was identified originally by mRNA display selection. The affinity of R3003 (SEQ ID NO: 4) for C5 was verified by the results of FP analysis displaying low K_i as well as IC_{50} values. Inhibitors R3011 (SEQ ID NO: 31) and R3050 (SEQ ID NO: 23) also displayed relatively strong affinity for C5. A control polypeptide, R3023 (SEQ ID NO: 104), displayed no affinity for C5, while inhibitors R3014 (SEQ ID NO: 55) and R3043 (SEQ ID NO: 50) displayed weak affinity.

Example 14. Analysis of compound stability in plasma

[00284] Compounds were assayed for stability in human plasma under the following conditions. Human plasma was obtained from Bioreclamation (Westbury, NY) and collected in

sodium heparin. Plasma was adjusted to pH 7.4. DMSO stocks at 10 mM concentration were prepared for the test compounds. Aliquots of the DMSO solutions were dosed into 1 mL of plasma, which had been pre-warmed to 37°C, at a final test compound concentration of 10 µM. The vials were kept in a benchtop THERMOMIXER® (Eppendorf, Hauppauge, NY) for the duration of the experiment. Aliquots (100 µL) were taken at each timepoint and added to a 96-well plate that had been pre-filled with 300 µL of an acetonitrile solution containing mixture of the internal standards (metoprolol, propranolol and warfarin each at 500 ng/mL). Samples were stored at 4°C until the end of the experiment. After the final timepoint was sampled, the plate was mixed and then centrifuged at 3,000 rpm for 10 minutes. Aliquots of the supernatant were removed and analyzed by LC-HRAMS. Liquid chromatography settings are listed in Table 8 and mass spectrometry settings are listed in Table 9.

Table 8. Liquid chromatography settings

Column:	Luna C18 (Luna, Torrance, CA) 50 mm x 2.0 mm, 3 µm			
M.P. Buffer:	Aqueous Reservoir (A): 0.1% Acetic acid in water			
	Organic Reservoir (B): 0.1% Acetic acid in MeOH:MeCN=1:1			
Gradient Program:	Time (Min)	Flow rate (mL/min)	% A	% B
	0.0	0.3	100	0
	5	0.3	0	100
	7.5	0.3	0	100
	7.6	0.45	100	0
	10.5	0.3	100	0
Total Run Time:	10.5 minutes			
Autosampler:	Agilent 1100 Bin (Agilent, Santa Clara, CA)			
Injection loop volume:	20µL			
Injection volume:	10µL			
Autosampler Wash 1:	Methanol / water 1:1; with 0.2% formic acid			
Autosampler Wash 2:	Methanol / 2-propanol:1/1; with 0.2% formic acid			

Table 9. Mass spectrometry settings

Instrument:	LTQ Orbitrap XL (Thermo Scientific, St. Louis, MO)
Positive Mode:	Electrospray , positive mode (+5000V)
Interface:	High Resolution Mass Spectroscopy
Mode:	Capillary Temperature: 275°C
Ion Source Settings:	Capillary Voltage: 47

	<p>Sheath gas: 45 Auxiliary gas: 15 Sweep gas: 10</p>
Orbitrap Settings:	<p>Scan Range 200-2000, Resolution =30000 (Full width at half maximum)</p> <p>Setting for MS/MS Data Dependent Acquisition Isolation Width: 2 Normalized Collision Energy: 35</p>

[00285] The concentration of test compound R3050 (SEQ ID NO: 23) was determined by comparison to a previously determined calibration curve (Table 10).

Table 10. Stability profile

Percent remaining at time (hrs)					
0hrs	2hrs	4hrs	12hrs	24hrs	Half-Life (min.)
100	108.9	100.4	88	98.1	>1440

[00286] Under these conditions, R3050 (SEQ ID NO: 23) was shown to be highly stable.

Example 15. Polypeptide variants comprising tryptophan analogs

[00287] In some embodiments, polypeptides of the present invention comprise 7-azatryptophan. To determine the importance of this residue in C5 inhibition, amino acid substitution analysis was carried out wherein 7-azatryptophan was replaced by natural tryptophan as well as various other tryptophan analogs including 5-fluorotryptophan [(5-F)W], 1-methyl-tryptophan [(1-Me)W], D-tryptophan and 5-methyl-O-tryptophan [(5-MeO)W]. Similar polypeptides with non-tryptophan substitutions were analyzed as well.

[00288] Polypeptide variants of R3002 (SEQ ID NO: 3) and R3008 (SEQ ID NO: 9) were synthesized and tested for their ability to inhibit red blood cell lysis as described in Example 10 (see Tables 11 and 12). Of the variants tested, all with substitution of the 7-azatryptophan residue demonstrated a decreased ability to inhibit red blood cell lysis as indicated by increasing average IC₅₀ values (a measure of the half maximal inhibitory concentration, a value used to indicate the amount of the inhibitor needed to reduce a given reaction or process by half).

Table 11. 7-azatryptophan variant polypeptides of R3002 (SEQ ID NO: 3) analyzed by human hemolysis assay

Compound Number	Sequence	Avg. IC ₅₀ (nM)	SEQ ID NO
R3002	Ac-Nvl-C-Y-E-N-Tbg-Y-azaTrp-E-Y-P-Phg-Nvl-NH2	11.9	3
R3041	Ac-Y-E-N-Tbg-Y-W-E-Y-P-Phg-Nvl-NH2	71.3	26
R3094	Ac-Y-E-N-Tbg-Y-(5-MeO)W-E-Y-P-Phg-Nvl-NH2	>100,000	121
R3110	Ac-Y-E-N-Tbg-Y-(1-Me)W-E-Y-P-Phg-Nvl-NH2	2,940	59
R3118	Ac-Y-E-N-Tbg-Y-(D-Trp)-E-Y-P-Phg-Nvl-NH2	>100,000	127
R3119	Ac-Y-E-N-Y-(D-Trp)-E-Y-P-Phg-Nvl-NH2	>100,000	128
R3128	Ac-Nvl-C-Y-N-N-Tbg-E-C-E-Y-P-Phg-Tbg-NH2	>100,000	132
R3067	Ac-Nvl-S-Y-E-N-Tbg-Y-A-E-Y-P-Chg-Nvl-NH2	>10,000	76
R3116	Ac-Nvl-Nvl-Y-E-N-Tbg-Y-(N-Me)W-E-Y-P-Chg-Nvl-NH2	441	45
R3017	[mXyl(2,8)]Ac-Nvl-C-Y-E-N-Tbg-Y-C-E-Y-P-Phg-Nvl-NH2	>100,000	103
R3129	[mXyl(2,8)]Ac-Nvl-C-Y-N-N-Tbg-E-C-E-Y-P-Phg-Tbg-NH2	>100,000	133

Table 12. 7-azatryptophan variant polypeptides of R3008 (SEQ ID NO: 9) analyzed by human hemolysis assay

Compound Number	Sequence	Avg. IC ₅₀ (nM)	SEQ ID NO
R3008	[mXyl(2,10)]Ac-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-H-C-Nvl-P-Nvl-NH2	97	9
R3088	[mXyl(2,10)]Ac-Nvl-C-Phg-T-(5-F)W-E-Y-(N-Me)S-A-C-Nvl-NH2	>100,000	118
R3091	[mXyl(2,10)]Ac-Nvl-C-Phg-T-W-E-Y-(N-Me)S-A-C-Nvl-NH2	483	46
R3092	[mXyl(2,10)]Ac-Nvl-C-Phg-T-F-E-Y-(N-Me)S-A-C-Nvl-NH2	>50,000	92
R3109	[mXyl(2,10)]Ac-Nvl-C-Phg-T-(1-Me)W-E-Y-(N-Me)S-A-C-Nvl-NH2	>100,000	126
R3131	[mXyl(2,10)]Ac-Nvl-C-Phg-T-W-Asp(T)-Y-(N-Me)S-H-C-Nvl-NH2	>100,000	135
R3034	[mXyl(2,10)]Ac-Nvl-C-Phg-T-A-E-Y-(N-Me)S-H-C-Nvl-NH2	>50,000	85
R3132	[mXyl(2,10)]Ac-Nvl-C-Phg-T-(D-Trp)-E-Y-(N-Me)S-H-C-Nvl-NH2	>100,000	136

Example 16. Effect of polypeptide truncation and amino acid deletion analyzed by human hemolysis assay

[00289] C-terminally truncated polypeptide variants of R3021 (SEQ ID NO: 11) were synthesized and assayed by the human hemolysis assay described in Example 10 for their ability to inhibit C5-dependent red blood cell lysis. Average IC₅₀ values (a measure of the half maximal inhibitory concentration, a value used to indicate the amount of the inhibitor needed to reduce a given reaction or process by half) for each polypeptide tested are listed in Table 13. Truncated polypeptides demonstrated decreased ability (as indicated by an increase in the IC₅₀ value) to inhibit red blood cell lysis with those variants lacking tryptophan having the largest IC₅₀ values.

[00290] Additionally, polypeptide variants of R3021 (SEQ ID NO: 11) with internal amino acid deletions were synthesized and assayed (see Table 14) for their ability to inhibit C5-dependent red blood cell lysis according to the method described in Example 10. Additionally, the N-terminal methionine in these variants was replaced by an acetyl group. Average IC₅₀ values for each polypeptide tested are listed in Table 14. Interestingly, acetyl group replacement of the N-terminal methionine alone [R3048 (SEQ ID NO: 19)] increased the ability of the polypeptide to inhibit red blood cell lysis. Removal of internal residue D [R3124 (SEQ ID NO: 130)] or internal residues DVY [R3125 (SEQ ID NO: 131), corresponding to residues 8, 9 and 10 from R3021 (SEQ ID NO: 11)] led to a decrease in the ability of the polypeptide to inhibit red blood cell lysis.

Table 13. C-terminally truncated polypeptide variants of R3021 (SEQ ID NO: 11) analyzed by human hemolysis assay

Compound Number	Sequence	Avg. IC ₅₀ (nM)	SEQ ID NO
R3021	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	27.5	11
R3043	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-NH2	934	50
R3044	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-NH2	>50,000	88
R3045	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-NH2	>100,000	107
R3046	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-NH2	>100,000	108
R3047	[mXyl(2,7)]M-C-V-E-R-F-C-NH2	>100,000	109

Table 14. Polypeptide variants of R3021 (SEQ ID NO: 11) with internal amino acid deletions analyzed by human hemolysis assay

Compound Number	Sequence	Avg. IC ₅₀ (nM)	SEQ ID NO
R3021	[mXyl(2,7)]M-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	27.5	11

R3048	[mXylyl(1,6)]Ac-C-V-E-R-F-C-D-V-Y-W-E-F-NH2	19.3	19
R3124	[mXylyl(1,6)]Ac-C-V-E-R-F-C-V-Y-W-E-F-NH2	>100,000	130
R3125	[mXylyl(1,6)]Ac-C-V-E-R-F-C-W-E-F-NH2	>100,000	131

Example 17. Incorporation of albumin-binding polypeptides

[00291] Polypeptides are conjugated to one or more polypeptides that modulate plasma protein binding. These polypeptides, referred to herein as “albumin-binding polypeptides” are listed in Table 15.

Table 15. Albumin-binding polypeptides

Albumin-binding polypeptide sequence	SEQ ID NO
Ac-R-L-I-E-D-I-C-L-I-P-R-W-G-C-L-W-E-D-D-NH2	202
Q-R-L-M-E-D-I-C-L-P-R-W-G-C-L-W-E-D-D-F-NH2	203
Ac-Q-R-L-I-E-D-I-C-L-P-R-W-G-C-L-W-E-D-D-F-NH2	204

[00292] Albumin-binding polypeptides are cyclized by disulfide bond formation at cysteine residues. In some embodiments, albumin-binding polypeptides are conjugated by either their N or C-terminal ends, thus having slightly different structures (e.g. no acetyl group).

Example 18. Incorporation of cell penetrating polypeptides

[00293] Polypeptides are conjugated to a polypeptide that has cell penetrating properties. These polypeptides are listed in Table 16 and are described in Milletti, F., Cell-penetrating polypeptides: classes, origin, and current landscape. Drug Discov Today. 2012 Aug;17(15-16):850-60.

Table 16. Cell penetrating polypeptides

Cell penetrating polypeptide	SEQ ID NO
R-K-K-R-R-R-E-S-R-K-K-R-R-E-S	205
R-K-K-R-R-Q-R-R-R	206
R-Q-I-K-I-W-F-Q-N-R-R-M-K-W-K-K	207
A-A-V-L-L-P-V-L-L-A-A-P	208
V-P-T-L-K	209
P-L-I-L-L-R-L-L-R-G-Q-F	210

Example 19. Analysis of polypeptide mixtures comprising amino acid stereoisomers

[00294] Polypeptides R3136 (SEQ ID NO: 137) and R3137 (SEQ ID NO: 138) were synthesized according to the amino acid sequences of R3085 (SEQ ID NO: 90) and R3082 (SEQ

ID NO: 116), respectively with the exception of the replacement of Phg in each with D-Phg (see Table 17). Compositions comprising either R3136 (SEQ ID NO: 137) and R3085 (SEQ ID NO: 90) or R3137 (SEQ ID NO: 138) and R3082 (SEQ ID NO: 116) were analyzed for their ability to inhibit red blood cell lysis according to the human hemolysis assay described in Example 10. The composition comprising R3136 (SEQ ID NO: 137) and R3085 (SEQ ID NO: 90) yielded an average IC₅₀ (nM) of >50,000, while the composition comprising R3137 (SEQ ID NO: 138) and R3082 (SEQ ID NO: 116) yielded an average IC₅₀ (nM) of >100,000.

Table 17. Compounds used in amino acid stereoisomer polypeptide mixtures

Compound Number	Sequence	SEQ ID NO.
R3136	[mXyl(2,10)]heptanoyl-Nvl-C-(D-Phg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	137
R3137	[mXyl(1,9)]heptanoyl-C-(D-Phg)-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	138
R3085	[mXyl(2,10)]heptanoyl-Nvl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	90
R3082	[mXyl(1,9)]heptanoyl-C-Phg-T-azaTrp-E-Y-(N-Me)S-A-C-Nvl-NH2	116

Example 20. Pharmacokinetic studies in non-human primates

[00295] Pharmacokinetic studies were carried out in non-human primates using the compounds listed in Table 18. In the table, “Cmpd” refers to compound and “Avg” refers to average.

Table 18. Compounds tested in *in vivo* studies

Cmpd No.	Sequence	Avg IC ₅₀	SEQ ID NO.
R3152	[mXyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl-NH2	16.4	153
R3201	[mXyl(1,6)]Ac-C-V-E-R-F-C-D-Tbg-Y-azaTrp-E-Y-P-Chg-Nvl	7.7	211

[00296] The plasma concentration of polypeptide R3152 (SEQ ID NO: 153) was determined in cynomolgus monkeys following a single intravenous (IV) dose. Three male animals received 3 mg/kg polypeptide and plasma concentrations of polypeptide were determined using LC/MS-MS following acetonitrile precipitation and extraction on Sirocco Protein Precipitation plate (Waters Corporation, Milford, MA.) Pharmacokinetic (PK) parameters were calculated from the time course (see Figure 4) of the combined plasma concentrations of R3152 (SEQ ID NO: 153,) determined immediately post-dose to up to 48 h post-dose. Plasma drug levels fell rapidly

during an initial distribution phase (<1 hour) and then plateaued and were detectable for up to 48 hours. R3152 (SEQ ID NO: 153) had a mean terminal half-life of 10.9 ± 0.8 hours. The mean clearance rate was 0.129 ± 0.0122 L/hr/kg which is approximately 5% of the liver blood flow of a typical monkey (2.6 L/hr/kg). The mean volume of distribution was 1.49 ± 0.152 L/kg which is approximately double the total body water for a typical monkey (0.7 L/kg). The average AUC_{∞} was 23319 ± 2120 hr*ng/mL.

[00297] R3152 (SEQ ID NO: 153) binds with high affinity to primate C5 protein and blocks the complement pathway by preventing the generation of the C5a and C5b products and the production of a multimeric Membrane Attack Complex (MAC). The inhibition of complement-mediated MAC formation in plasma samples from the above PK study was examined using an established *ex vivo* assay (see the human hemolysis assay described in Example 10), in which plasma was diluted 1:100 and incubated with activated sheep red blood cells (Complement Technology, Tyler, TX). At each time-point, the hemolytic activity was determined as an indicator of active serum complement (see Figure 4) In plasma containing >200 ng/mL R3152 (SEQ ID NO: 153) there was a clear inhibition of complement-mediated hemolysis, indicating a blockade of MAC formation. Exogenous R3152 (SEQ ID NO: 153) added to normal cynomolgus plasma has an $IC_{50} = 2-20$ ng/mL. Hemolytic activity returned to normal levels 48 hours after dosing as plasma levels of R3152 (SEQ ID NO: 153) fell below 100 ng/ml.

Example 21. Pharmacokinetic studies in rat

[00298] R3152 (SEQ ID NO: 153) was delivered as an intravenous (IV) or subcutaneous (SC) dose to male rats at 2 and 30 mg/kg, respectively (Figure 5) Following IV dosing, R3152 (SEQ ID NO: 153) was monitored by using LC/MS-MS following acetonitrile precipitation and extraction on Sirocco Protein Precipitation plate (Waters Corporation, Milford, MA) as described above. Pharmacokinetic (PK) parameters were calculated from the time course of the combined plasma concentrations (see Figure 5) of R3152 (SEQ ID NO: 153) and its equipotent C-terminally deamidated metabolite, R3201 (SEQ ID NO: 211).

[00299] R3152 (SEQ ID NO: 153)/ R3201 (SEQ ID NO: 211) exhibited a fast distribution phase, followed by a slow elimination with a $t_{1/2} = 5.3$ hrs. A similar elimination rate was observed after SC dosing of 30 mg/kg, with approximately 65% bioavailability of dose, based on AUC. The T_{max} of 4 hrs and prolonged drug exposure seen in SC doses allowed for extended

coverage of the therapeutic concentration in plasma. As R3152 (SEQ ID NO: 153) and R3201 (SEQ ID NO: 211) do not bind to rat C5, very little inhibitory activity was observed in *ex vivo* hemolysis assays.

[00300] Lipidated and non-lipidated compounds R3183 (SEQ ID NO: 184) and R3176 (SEQ ID NO: 177), respectively, were evaluated for pharmacokinetic properties in Male Sprague-Dawley Rats following intravenous or subcutaneous administration. Figure 6 shows the results. The left panel of Figure 6 shows Male Sprague-Dawley rats (n=3) injected intravenously with a single 2 mg/kg dose. Blood samples were collected at indicated time points, processed into plasma, and analyzed for the indicated compound by LC-MS. Black circles: R3176 (SEQ ID NO: 177) (unlipidated compound); Open circles: R3183 (SEQ ID NO: 184) (C16 lipidated compound). The right panel of Figure 6 shows Male Sprague-Dawley rats (n=3) injected subcutaneously with a single 15 mg/kg dose. Blood samples were collected at indicated time points, processed into plasma, and analyzed for the indicated compound by LC-MS. Black circles: R3176 (SEQ ID NO: 177) (unlipidated compound); Open circles: R3183 (SEQ ID NO: 184) (C16 lipidated compound). Lipidation resulted in an increase in exposure as assessed by determination of the Area under the curve (AUC) by 2.1-fold by the intravenous route and 2.7-fold by the subcutaneous route.

Example 22. Inhibition of hemolysis in the Thrombin-Induced Complement Pathway

[00301] Thrombin can induce complement activity by cleaving C5 into C5_T which will then be cleaved into C5a and C5b_T. C5b_T, like C5b, will associate with C6 and the remaining terminal components of the complement pathway, C7, C8 and C9, which will lead to formation of the Membrane Attack Complex (MAC) causing lysis of red blood cells (Krisinger, et al., (2014). Blood. 120(8):1717-1725). Thus, R3183 and an anti-C5 monoclonal antibody similar to ECULIZUMAB® were tested for their ability to inhibit hemolysis through the thrombin-induced complement pathway.

[00302] To assess inhibitor activity, C5 (Complement Technology, Tyler, TX) was added to achieve a concentration of 400 nM, and the sample was incubated with C6 at a final concentration of 600 nM (Complement Technology, Tyler, TX) and thrombin at a concentration of 50 nM (Enzyme Research Laborites, South Bend, IN) at 37°C for 30 minutes, in the presence of either R3183 or the anti-C5 monoclonal antibody similar to ECULIZUMAB®, or no inhibitor.

The reaction was stopped with the addition of hirudin to 150 nM (Cell Sciences, Canton, MA) in the buffer GVB+EDTA (Complement Technology, Tyler, TX) and incubated 5 minute at room temperature. These diluted samples were mixed with antibody-sensitized sheep erythrocytes (Complement Technology, Tyler, TX) in a 96-well microtitre plate (USA Scientific, Ocala, FL) and incubated at 37°C for 5 minutes. C7 (Complement Technology, Tyler, TX) was then added to the wells to achieve a concentration of 15 nM and the plate was returned to 37°C for 15 minutes. A complex of C8 (10 nM; Complement Technology, Tyler, TX) and C9 (25 nM; Complement Technology, Tyler, TX) was then added to the assay mixture and the samples were incubated for 30 minutes at 37°C. Following incubation, the plate was centrifuged at 1000 x g and 100 µL of supernatant was transferred to a new microtitre plate and the absorbance was read at 412 nm. The resulting data are shown in Figure 7. R3183 was found to inhibit hemolysis by the thrombin-induced complement pathway at concentrations higher than 6 ng/mL, while the anti-C5 monoclonal antibody did not.

CLAIMS

What is claimed is:


1. A polypeptide of the formula R₁-Xaa0-Xaa1-Xaa2-Xaa3-Xaa4-Xaa5-Xaa6-Xaa7-Xaa8-Xaa9-Xaa10-Xaa11-Xaa12-Xaa13-Xaa14-Xaa15-Xaa16-Xaa17-Xaa18-R₂ (SEQ ID NO: 212), wherein:
 - a. R₁ is absent or selected from the group consisting of H, an acetyl group, an acyl group containing a linear or branched, saturated or unsaturated hydrocarbon chain from 1 to 20 carbon atoms, a heptanoyl group, an amide, a carbamate, urea, PEG, hydroxyalkyl starch and a polypeptide;
 - b. Xaa0 is absent, or an amino acid selected from the group consisting of Met and norvaline;
 - c. Xaa1 is absent or selected from the group consisting of (S)-2-amino-5-azidopentanoic acid, chloroacetic acid, (S)-2-aminohept-6-enoic acid, 4-aminobutyric acid, 5-aminopentanoic acid, 5-aminohexanoic acid, ornithine, Lys, homolysine, Glu, Asp, 3-thiopropionic acid and Cys;
 - d. Xaa2 is absent or selected from the group consisting of Ala, D-Ala, tertbutyl-glycine, Lys, Ser, Cys and Val;
 - e. Xaa3 is absent or selected from the group consisting of Ala, norvaline, Val and Glu;
 - f. Xaa4 is absent or selected from the group consisting of Ala, Cys, Arg, Ser, Glu, phenylglycine and norvaline;
 - g. Xaa5 is absent or selected from the group consisting of Tyr, Arg, Cys, Phe, N-methyl-tyrosine and Ala;

- h. Xaa6 is absent or selected from the group consisting of Glu, N-methyl-glutamic acid, cyclohexylglycine, Lys, Tyr, Pro, N-methyl-serine, tertbutyl-glycine, Val, norleucine, norvaline, 7-azatryptophan, Asn, Asp, (S)-2-aminopent-4-ynoic acid, (S)-2-aminopent-4-enoic acid, Cys and Ala;
- i. Xaa7 is absent or selected from the group consisting of Asn, N-methyl-asparagine, N-methyl-glycine, N-methyl-serine, homocysteine, Thr, Tyr, phenylglycine, tert-butylglycine, alpha-methyl L-aspartic acid, (S)-2-amino-3-(1*H*-tetrazol-5-yl)propanoic acid, N-methyl-aspartic acid, cycloleucine, 4-amino-tetrahydro-pyran-4-carboxylic acid, Arg, Glu, Asp, Cys and Ala;
- j. Xaa8 is absent or selected from the group consisting of tert-butylglycine, phenylglycine, D-phenylglycine, cyclohexylglycine, D-cyclohexylglycine, 1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid, Tyr, tert-butylglycine, Asn, Cys, N-methyl-4-fluorophenylalanine, 2-O-methylphenylglycine, Phe, Val and Ala;
- k. Xaa9 is selected from the group consisting of Tyr, N-methyl-tyrosine, Thr, Glu, norvaline, Lys, Ala, D-Ala, His, Cys, phenylglycine, N-methyl-serine, N-methyl-glycine, amino isobutyric acid and Arg;
- l. Xaa10 is selected from the group consisting of 3-aminomethyl-L-phenylalanine, 7-azatryptophan, N-methyl-tryptophan, 1-methyl-tryptophan, 5-fluorotryptophan, Phe, D-Trp, 5-methyl-O-tryptophan, Ala, His, Leu, tert-butylglycine, Cys and Trp;
- m. Xaa11 is selected from the group consisting of Glu, D-Glu, N-methyl-glutamic acid, Asn, Asp, Gln, tert-butylglycine, Cys, N-methyl-4-fluorophenylalanine, N-methyl-serine, (S)-2-amino-3-(1*H*-tetrazol-5-yl)propanoic acid and Ala;
- n. Xaa12 is absent or selected from the group consisting of Trp, homophenylalanine, meta-chloro homophenylalanine, 2-naphthylalanine, 3-aminomethyl-L-phenylalanine, Tyr, N-methyl-tyrosine, Cys, Phe, Ala, Glu, Gly, N-methyl-

glycine, phenylglycine, 4-fluorophenylalanine, O-methyl-tyrosine, homophenylalanine, 3-chlorophenylalanine and norvaline;

- o. Xaa13 is absent or selected from the group consisting of propargyl-glycine, Pro, Ala, N-methyl-glycine, Ser, N-methyl-serine, N-methyl-alanine, norvaline, Cys and Tbg;
- p. Xaa14 is absent or selected from the group consisting of amino isobutyric acid, tert-butylglycine, Cys, Pro, Asn, phenylglycine, D-phenylglycine, N-methyl-phenylglycine, norvaline, His, Ala, D-Ala, and cyclohexylglycine;
- q. Xaa15 is absent or selected from the group consisting of norvaline, Lys, N- ϵ -caprylic lysine, N- ϵ -capryl lysine, N- ϵ -lauryl lysine, N- ϵ -palmitoyl lysine, Pro, Cys, Tyr, Gly, propargyl-glycine, homoCys, N-methyl-serine and tert-butylglycine;
- r. Xaa16 is absent or selected from the group consisting of norvaline, Cys, Lys and Ala;
- s. Xaa17 is absent or selected from the group consisting of Pro, Glu and Nvl;
- t. Xaa18 is absent or is norvaline; and
- u. R₂ is absent or selected from the group consisting of B20, B28, K14, -NH₂ and -N(CH₃)₂.

2. The polypeptide of claim 1, further comprising a bridging moiety between two amino acids.
3. The polypeptide of claim 2, wherein said bridging moiety comprises a structure selected from the group consisting of structures I-XIX;




wherein each X is independently N or CH, such that no ring contains more than 2 N; each Z is independently a bond, NR, O, S, CH₂, C(O)NR, NRC(O), S(O)_vNR, NRS(O)_v; each m is independently selected from 0, 1, 2, and 3; each v is independently selected from 1 and 2; each R is independently selected from H and C₁-C₆; and each bridging moiety is connected to the polypeptide by independently selected C₀-C₆ spacers.

4. The polypeptide of claim 2, wherein the bridging moiety comprises a feature selected from the group consisting of a disulfide bond, an amide bond (lactam), a thioether bond an aromatic ring, an unsaturated aliphatic hydrocarbon chain, a saturated aliphatic hydrocarbon chain and a triazole ring.
5. The polypeptide of any of claims 2-4, wherein the cyclic loop is of a length selected from the group consisting of 1 amino acid, 2 amino acids, 3 amino acids, 4 amino acids, 5 amino acids, 6 amino acids, 7 amino acids, 8 amino acids, 9, amino acids, 10 amino

acids, 11 amino acids, 12 amino acids, 13 amino acids, 14 amino acids, 15 amino acids and 16 amino acids.

6. The polypeptide of claim 2, wherein residue Xaa1 is cysteine and wherein the bridging moiety joins residue Xaa1 and a residue selected from the group consisting of Xaa4, Xaa5, Xaa6, Xaa7, Xaa8, Xaa9, Xaa10, Xaa11, Xaa12, Xaa13, Xaa14, Xaa15 and Xaa16.
7. The polypeptide of claim 4, wherein said feature comprises an aromatic ring and wherein said bridging moiety is formed by reaction with a poly(bromomethyl)benzene.
8. The polypeptide of claim 7, wherein the poly(bromomethyl)benzene is selected from the group consisting of 1,2-bis(bromomethyl)benzene, 1,3-bis(bromomethyl)benzene and 1,4-bis(bromomethyl)benzene.
9. The polypeptide of claim 8 wherein the reagent is 1,3-bis(bromomethyl)benzene.
10. The polypeptide of claim 4, wherein said feature comprises a disulfide bond between two cysteine residues.
11. The polypeptide of claim 4, wherein said feature comprises an aromatic ring and wherein said bridging moiety is produced by reaction with a compound selected from the group consisting of 2,6-bis(bromomethyl)pyridine, (E)-1,4-dibromobut-2-ene and 1,2-bis(bromomethyl)-4-alkylbenzene.
12. The polypeptide of claim 1, wherein Xaa8 is tert-butylglycine, Xaa9 is Tyr, Xaa10 is 7-azatryptophan, Xaa11 is Glu and Xaa12 is Tyr.
13. The polypeptide of claim 12, comprising a bridging moiety between two amino acids.
14. The polypeptide of claim 13 wherein said bridging moiety comprises a structure selected from the group consisting of structures I-XIX;



wherein each X is independently N or CH, such that no ring contains more than 2 N; each Z is independently a bond, NR, O, S, CH₂, C(O)NR, NRC(O), S(O)_vNR, NRS(O)_v; each m is independently selected from 0, 1, 2, and 3; each v is independently selected from 1 and 2; each R is independently selected from H and C₁-C₆; and each bridging moiety is connected to the polypeptide by independently selected C₀-C₆ spacers.

15. The polypeptide of claim 13, wherein the bridging moiety comprises a feature selected from the group consisting of a disulfide bond, an amide bond (lactam), a thioether bond an aromatic ring, an unsaturated aliphatic hydrocarbon chain, a saturated aliphatic hydrocarbon chain and a triazole ring.
16. The polypeptide of claim 13, wherein said bridging moiety is formed by reaction with a poly(bromomethyl)benzene.


17. The polypeptide of claim 1, wherein said polypeptide is selected from the group consisting of SEQ ID NOs 1-201 and 211.
18. The polypeptide of claim 17, wherein said polypeptide inhibits the cleavage of C5 into C5a and C5b cleavage products.
19. The polypeptide of claim 1, wherein said polypeptide is part of an antibody.
20. A composition comprising the polypeptide of claim 1 and an acceptable carrier or excipient.
21. The composition of claim 20, wherein said polypeptide is conjugated to a hydrophilic polymer.
22. The composition of claim 21, wherein the hydrophilic polymer is selected from the group consisting of polyalkylene oxide homopolymers, polypropylene glycols, polyoxyethylenated polyols and copolymers thereof.
23. The composition of claim 22, wherein the water soluble polymer is polyethylene glycol (PEG).
24. The composition of any of claims 20-23, wherein said polypeptide is conjugated to an albumin-binding polypeptide selected from the group consisting of SEQ ID NOs: 202-204.
25. The composition of any of claims 20-24, wherein said polypeptide is conjugated to a cell penetrating polypeptide selected from the group consisting of SEQ ID NOs: 205-210.
26. The composition of any of claims 20-25, comprising an excipient, wherein said excipient comprises a pharmaceutically acceptable excipient.
27. A method of inhibiting C5 cleavage in a cellular system, comprising contacting said cellular system with the composition of any of claims 20-26 or a polypeptide of any of claims 1-19.

28. The method of claim 27, wherein C5a levels are reduced.
29. The method of claim 28, wherein formation of the membrane attack complex (MAC) is reduced.
30. The method of claim 27, wherein said polypeptide inhibits the cleavage of C5 with an IC₅₀ of from about 50 nM to about 200 nM.
31. The method of claim 27, wherein said polypeptide inhibits the cleavage of C5 with an IC₅₀ of less than 50 nM.
32. The method of any of claims 27-31, wherein said cellular system is selected from the group consisting of an *in vitro* system, an *in vivo* system and an *ex vivo* system.
33. The method of claim 32, comprising an *in vivo* system, wherein said *in vivo* system comprises a subject.
34. The method of claim 33, wherein said subject is a human subject.
35. The method of claim 34, wherein said human subject comprises a complement-related disease, disorder and/or condition.
36. The method of claim 35, wherein said contacting comprises a method of administration selected from the group consisting of buccal administration, intranasal administration, oral administration, intravenous administration, intramuscular administration, intraperitoneal administration, subcutaneous administration, transdermal administration and intravitreal administration.
37. The method of either of claims 35 or 36, wherein said complement-related disease, disorder and/or condition is selected from the group consisting of an inflammatory indication, a wound, an injury, an autoimmune disease, a vascular indication, a neurological indication, a kidney-related indication, an ocular disease, paroxysmal nocturnal hemoglobinuria and atypical hemolytic uremic syndrome.

38. A method of inhibiting thrombin-induced complement activation in a cellular system, comprising contacting said cellular system with the polypeptide of any of claims 1-19 or the composition of any of claims 20-26.
39. A method of treating hemolysis in a subject, comprising administering to said subject the polypeptide of any of claims 1-19 or the composition of any of claims 20-26.
40. The method of claim 39, wherein said hemolysis is caused by thrombin-induced complement activation.


1/7

Figure 1




2/7

Figure 2




3/7


Figure 3



4/7

Figure 4





6/7




Figure 6

7/7

