

[19] Patents Registry
The Hong Kong Special Administrative Region
香港特別行政區
專利註冊處

[11] 1237377 B
EP 3168309 B1

[12] **STANDARD PATENT (R) SPECIFICATION**
轉錄標準專利說明書

[21] Application no. 申請編號 17111501.0
[51] Int. Cl. C12Q 1/6851 (2018.01) C12Q 1/6883 (2018.01)
[22] Date of filing 提交日期 08.11.2017

[54] DETECTION OF FOETAL CHROMOSOMAL ANEUPLOIDIES USING DNA REGIONS THAT ARE DIFFERENTIALLY METHYLATED BETWEEN THE FOETUS AND THE PREGNANT FEMALE
使用胎兒和懷孕女性之間的差異甲基化的 DNA 區域檢測胎兒染色體非整倍體

[43] Date of publication of application 申請發表日期
13.04.2018
[45] Date of publication of grant of patent 批予專利的發表日期
26.03.2021
EP Application no. & date 歐洲專利申請編號及日期
EP 15193966.7 10.11.2015
EP Publication no. & date 歐洲專利申請發表編號及日期
EP 3168309 17.05.2017
Date of grant in designated patent office 指定專利當局批予專利日期
08.04.2020

[73] Proprietor 專利所有人
Eurofins LifeCodexx GmbH
Line-Eid-Strasse 3
78467 Konstanz
GERMANY
[72] Inventor 發明人
WERLER, Steffi
HOFMANN, Wera
SACHSE, Matthias
[74] Agent and / or address for service 代理人及/或送達地址
KANGXIN IP LIMITED
812 Silvercord, Tower 1, 30 Canton Road
Tsimshatsui, Kowloon
HONG KONG

(19)



(11)

EP 3 168 309 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
08.04.2020 Bulletin 2020/15

(51) Int Cl.:
C12Q 1/6851 (2018.01) C12Q 1/6883 (2018.01)

(21) Application number: **15193966.7**

(22) Date of filing: **10.11.2015**

(54) **DETECTION OF FOETAL CHROMOSOMAL ANEUPLOIDIES USING DNA REGIONS THAT ARE DIFFERENTIALLY METHYLATED BETWEEN THE FOETUS AND THE PREGNANT FEMALE**

NACHWEIS VON FÖTALEN CHROMOSOMALEN ANEUPLOIDIEN MITTELS DNA-REGIONEN MIT UNTERSCHIEDLICHER METHYLIERUNG ZWISCHEN DEM FÖTUS UND DER SCHWANGEREN FRAU

DÉTECTION D'ANEUPLOÏDIES CHROMOSOMIQUES FOETALES AU MOYEN DE RÉGIONS D'ADN QUI SONT MÉTHYLÉES DE MANIÈRE DIFFÉRENTIELLE ENTRE LE FOETUS ET LA FEMME ENCEINTE

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
Designated Extension States:
BA ME
Designated Validation States:
MA MD

(43) Date of publication of application:
17.05.2017 Bulletin 2017/20

(73) Proprietor: **Eurofins LifeCodexx GmbH**
78467 Konstanz (DE)

(72) Inventors:
• **WERLER, Steffi**
78467 Konstanz (DE)
• **HOFMANN, Wera**
78467 Konstanz (DE)
• **SACHSE, Matthias**
78467 Konstanz (DE)

(74) Representative: **Dörries, Hans Ulrich**
df-mp Dörries Frank-Molnia & Pohlman
Patentanwälte Rechtsanwälte PartG mbB
Theatinerstrasse 16
80333 München (DE)

(56) References cited:
WO-A1-03/062441 WO-A1-2012/007783
WO-A2-00/47764

- **SUZANNE SNELLENBERG ET AL: "Development of a multiplex methylation-specific PCR as candidate triage test for women with an HPV-positive cervical scrape", BMC CANCER, BIOMED CENTRAL, LONDON, GB, vol. 12, no. 1, 23 November 2012 (2012-11-23), page 551, XP021128495, ISSN: 1471-2407, DOI: 10.1186/1471-2407-12-551**
- **EKATERINA OLKHOV-MITSEL ET AL: "Novel Multiplex MethyLight Protocol for Detection of DNA Methylation in Patient Tissues and Bodily Fluids", SCIENTIFIC REPORTS, vol. 4, 21 March 2014 (2014-03-21), XP055143311, DOI: 10.1038/srep04432**
- **THERESA SWIFT-SCANLAN ET AL: "Two-color quantitative multiplex methylation-specific PCR", BIOTECHNIQUES, vol. 40, no. 2, 1 February 2006 (2006-02-01), pages 210-219, XP055143300, ISSN: 0736-6205, DOI: 10.2144/000112097**
- **DA EUN LEE ET AL: "Non-Invasive Prenatal Testing of Trisomy 18 by an Epigenetic Marker in First Trimester Maternal Plasma", PLOS ONE, vol. 8, no. 11, 1 November 2013 (2013-11-01), page e78136, XP055258783, DOI: 10.1371/journal.pone.0078136**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 168 309 B1

- TONG YU K ET AL: "Noninvasive prenatal detection of fetal trisomy 18 by epigenetic allelic ratio analysis in maternal plasma: Theoretical and empirical considerations", CLINICAL CHEMISTRY, AMERICAN ASSOCIATION FOR CLINICAL CHEMISTRY, WASHINGTON, DC, vol. 52, no. 12, 1 December 2006 (2006-12-01), pages 2194-2202, XP002603754, ISSN: 0009-9147, DOI: 10.1373/CLINCHEM.2006.076851 [retrieved on 2006-10-13]

Remarks:

The complete document including Reference Tables and the Sequence Listing can be downloaded from the EPO website

Description

[0001] The present invention relates to methods for detecting a chromosomal aneuploidy in a foetus carried by a pregnant female. Such methods are based on one or more of particular configurations and/or detections and/or analyses of two or more regions of DNA, including those that show differential methylation between DNA that originates from cells of a foetus (and/or the placenta of a foetus) and DNA of maternal origin. Such methods utilise a sample taken from said pregnant female, which sample comprises DNA that originates from cells of a foetus and/or the placenta of a foetus in admixture with differently methylated DNA of maternal origin. Such methods have diagnostic, prognostic and/or predictive utility; in particular for the detection/diagnosis of chromosomal aneuploidy, such as a trisomy, in a foetus, and/or for detecting an increased risk of an pregnant female suffering from or developing a pregnancy-associated medical condition.

[0002] In most developed countries, many pregnant women undergo prenatal testing for which the main objective is the identification of chromosomal aneuploidy in the developing foetus. Until recently, almost all reliable such prenatal tests were invasive, such as chorionic villus sampling (CVS) or amniocentesis, and carried a not insignificant risk of procedure-related miscarriage despite the diagnostic accuracy of such testing having been reported as between 97.% to 99.8% (for review, see Norwitz & Levy 2013, Rev Obs Gyn 6:48). Conventional non-invasive prenatal testing for foetal chromosomal aneuploidy includes screening by ultra-sound and/or analysis of various maternal serum biochemical markers. However, these tests are primarily targeted to the detection of Down syndrome (trisomy chromosome 21 (T21)) and, to a lesser extent, T18. However, with reported detection rates of only 75% to 96% (depending on the screening approach utilised) and with false-positive rates ranging from 5% to 10%, such ultrasound and maternal serum analyses are commonly considered only as screening procedures, with both requiring follow-up by CVS or amniocentesis in screen-positive cases for a definitive diagnosis of a chromosome abnormality in the foetus (Norwitz & Levy 2013).

[0003] The discovery of cell-free DNA (cfDNA) in maternal circulation that originates from the foetus (Lo et al 1997, Lancet 350:485) has provided an alternative approach for the development of assays for non-invasive prenatal testing (NIPT), including assays for foetal sex determination (Lo et al 1997) as well as foetal Rhesus D testing in pregnancies where the mother is Rhesus D negative (Lo 1998, N Eng J Med 229:1734). However, given such foetal cfDNA existing as a minor fraction of total cfDNA isolated from maternal plasma (Lo et al 1998, Am J Hum Genet 63:768), the detection of foetal chromosome aneuploidies using on the basis of foetal cfDNA from maternal plasma remains challenging. The application of high-throughput next-generation sequencing (NGS) technologies has enabled the sequencing of both foetal and maternal contributions to cfDNA in maternal plasma, and only in recent years has it been possible to offer NGS-based NIPT for chromosome 21 and certain other common foetal aneuploidies routinely to pregnant women, for example with the advent and general availability of commercial tests. Commercially available tests can use random NGS such as "MATERNIT21" (www.sequenom.com), "PRENATEST" (www.lifecodexx.com), "VERIFI" (www.illumina.com) or "IONA" (www.premaita.com), or may use targeted approaches, the aim of which is to enrich specific genomic regions of interest before sequencing to reduce the number of sequence tags needed to perform a reliable statistical analysis (eg "HARMONY" www.ariosadx.com or "PANORAMA" www.natera.com), polymorphism analysis or digital PCR (for review, see Go et al 2011, Human Reprod Update 17:372).

[0004] To date, such commercially available NIPTs that follow a sequencing approach require the use of complicated and very expensive equipment such as massively parallel sequencers, and have a turn around or several to up to ten days; largely driven by the need for substantial sample processing prior to analysis. For example, NIPT for foetal aneuploidy by NGS requires the time- and cost-consuming step of consisting a so-called "sequencing library" of multimillions of individual fragments of isolated cfDNA. Alternative NIPTs that follow digital PCR require the use of such equipment and technology that is not typically available except in very specialised and localised research laboratories, and at relatively low throughput (Hindson et al 2011, Anal Chem 83:8604).

[0005] Recognising this significant limitation on NIPT from cfDNA, a number of researches have sought to investigate and develop alternative, and in particular cheaper and faster, ways of conducting NIPT for foetal aneuploidy. Many of these alternatives for NIPT have focused on the use of foetal-specific characteristics of the foetal DNA found in maternal plasma (such as specific species of or polymorphisms in foetal cfDNA). In particular, the presence of regions of DNA that are specifically methylated in the foetal DNA compared to maternal DNA has led to substantial efforts into the use of such epigenetic markers for NIPT of foetal chromosomes of interest. WO 03/062441A1 discloses the use of a first primer pair on a potentially aneuploid chromosome and a second primer pair on a non-aneuploid chromosome to determine the ratio of the quantity of the two PCR products amplified from bisulphite-treated DNA that had been isolated from maternal serum, thereby detecting foetal aneuploidies. The primers were specific to only amplify the bisulphite-treated unmethylated DNA.

[0006] The first demonstration of the use of an epigenetic marker to specifically identify foetal DNA in maternal plasma was made as long ago as 2002 (Poon et al 2002, Clin Chem 48:35; WO 2003/020974), although the utility of the marker used was limited to certain foetal-maternal pairs as it was polymorphic-dependent. A more generally applicable foetal-specific epigenetic marker was the methylated form of the serpin peptidase inhibitor clade B member 5 (also known as maspin) gene ("U-maspin" or "U-SERPINB5") was identified in 2005 following a candidate gene approach (Chim et al

2005, PNAS 102:14753). The maspin gene is located on human chromosome 18, and the principle demonstrated for the first time that non-invasive identification of foetal trisomy 18 could be conducted using such an epigenetic marker (Tong et al 2006, Clin Chem 52:2194; WO 2005/118852).

[0007] With such an in-principle demonstration, a number of groups set out to identify further foetal epigenetic markers, in particular those on chromosomes associated with foetal aneuploidy such as chromosome 21 (for example, Old et al 2007, Reprod Biomed Online; Chim et al 2008, Clin Chem 54:500; Papageorgiou et al 2009, Am J Pathol 174:1609; WO 2007/132166; WO 2007/132167; WO 2011/034631; and WO 2011/092592), with the goal of exploiting such epigenetic markers for NIPT of foetal aneuploidy by quantifying eg a chromosome 21-derived sequence in epigenetically identified foetal cfDNA, relative to a reference sequence derived from another autosome or sex chromosome (eg Old et al 2007).

[0008] Certain of these epigenetic markers have been investigated as tools to NIPT. Indeed, the unmethylated form of the phosphodiesterase 9A gene (U-PDE9A) has been used in an NIPT test for foetal trisomy 21 (Lim et al 2011, PLoS One 6:e27709). In another study, the methylated form of the holocarboxylase synthetase gene (M-HLCS) has been used as a foetal epigenetic marker to compare relative chromosome 21 dosage normalised against the concentrations of a foetal genetic marker, zinc finger protein Y-Linked (ZFY) gene, on chromosome Y. This epigenetic-genetic ("EGG") approach was demonstrated to be useful for the non-invasive detection of foetal trisomy 21 (Tong et al 2010, Clin Chem 56:90). Using a different approach, Papageorgiou and colleagues described the use of an immunoprecipitation and real-time quantitative polymerase chain reaction (qPCR) approach to quantify the relative amount of a foetal-specific differentially methylated regions (DMRs) chromosome 21 compared to maternal DNA from whole blood, to argue that the slight excess in chromosome 21 amount in trisomy cases could be detected and diagnostic of foetal trisomy 21 (Papageorgiou et al 2011, Nat Med 17:510), and they hypothesised that a combination of DMRs and not a single DMR may be able to give an accurate NIPD of normal and trisomy 21 cases. In a later study, it was reported that in a larger blinded validation study such an approach demonstrated 100% sensitivity and 99.2 specificity (Tsaliki et al 2012, Prenat Diag 21:996). Not only does such an approach require a cumbersome and technically challenging methylated DNA immunoprecipitation (MeDiP) step, but significant technical concerns about using such an approach have been raised (Tong et al 2012, Nat Med 18:1327). In particular, that by using such an approach it is not possible to conclude that the observation of an elevated level of foetal-derived DNA methylation of chromosome 21 markers in maternal blood is due to the presence of a foetus with trisomy 21, as compared to a euploid foetus that happens to release a high amount of cell-free foetal DNA or a high number of foetal cells into the maternal blood. In their commentary of the MeDiP approach, Tong and colleagues suggest that a normalisation step could be included to control for variations in foetal-DNA concentrations using foetal epigenetic markers outside of chromosome 21, but that the previously published study using RASSF1, a gene on chromosome 3, has indicated that compared to the EGG approach, the effect of such normalisation is typically suboptimal, even for placental tissues, which contain primarily foetal DNA (Tong et al 2010).

[0009] However, in a recent study, starting with a step to enrich for methylated DNA from maternal primer, Lim and colleagues use conventional qPCR to measure the relative dosage of chromosome 21 (using M-HLCS) compared to the reference chromosome 3 (M-RASSF1A) to report a sensitivity of 90% and a specificity of 92.5% in the non-invasive prenatal detection of foetal trisomy 21 (Lim et al 2014, Clin Chem Lab Med 52:641). Despite this seeming improvement on the initial work reported by Tong and colleagues (Tong et al 2010), It has been reported that there is room for improvement for Lim's approach (Chim 2014, Clin Chem Lab Med 52:585). Indeed, to explore ways to improve the performance of using foetal epigenetic markers for NIPT for foetal trisomy 21, Chim tabulated the salient features of other studies similar to Lim (see TABLE A, adapted from Chim 2014).

TABLE A: Studies on non-invasive prenatal detection of trisomy 21 using foetal epigenetic markers (adapted from Chim 2014)

Study	Pretreatment*	Quant. Method	Foetal Chr21 marker	Foetal Ref marker (Chr)	Nature of ref marker	Sensitivity	Specificity
Tong et al 2010	MSRE	Digital PCR	M-HLCS	ZFY	Genetic	100.0%	95.8%
Lim et al 2011	MBD	qPCR	U-PDE9A	M-PDE9A + U-PDE9A	Epigenetic	83.3%	94.4%
Lim et al 2014	MBD	qPCR	M-HLCS	M-RASSF1A	Epigenetic	90.0%	92.5%

*MSRE: methylation-sensitive restriction enzyme digestion; MBD: methyl-CpG binding domain-based enrichment;

[0010] Chim summarise that: (a) digital PCR seems to give better sensitivity and specificity, compared with the con-

ventional quantitative and suggest that translating the exponential but analogue nature of qPCR into a '1' or '0' signal in digital PCR, digital counting platforms should facilitate more precise and accurate quantification; and (b) the use of foetal genetic markers seems to give better sensitivity and specificity, rather than foetal epigenetic markers, to quantify a reference chromosome for relative chromosome dosage analysis. Indeed, similarly presenting the results obtained by Tong and colleagues (2010) in their comparative study of the EGG approach compared to the analogous epigenetic-epigenetic approach, Chim describe that normalising against a foetal epigenetic marker, namely M-RASSF1A located on chromosome 3, Tong and colleagues found that the relative dosage of chromosome 21 of more than half of the trisomy 21 placentas studies overlapped with the euploid (normal) reference interval leading to an unworkable sensitivity of 25% (see TABLE B).

TABLE B: Data from Tong et al (2010) on the prenatal detection of trisomy 21 using foetal epigenetic or genetic markers as reference (adapted from Chim 2014)

Sample	Pre-treatment*	Quant. Method	Foetal Chr21 marker	Foetal Ref marker (Chr)	Nature of ref marker	Sensitivity	Specificity
Placenta	MSRE	Digital PCR	M-HLCS	M-RASSF1A	Epigenetic	25.0%	100.0%
Placenta	MSRE	Digital PCR	M-HLCS	ZFY	Genetic	100.0%	100.0%
Maternal plasma	MSRE	Digital PCR	M-HLCS	ZFY	Genetic	100.0%	95.8%

[0011] Therefore, despite there being a long-felt need for improved NIPT for foetal aneuploidies and, since 2002, the theoretical possibility to use foetal-specific epigenetic markers to do so, there has to date been no satisfactory, reliable, sensitive, specific, cost effective and/or technically straightforward approach to do so. Indeed, even in his recent review, Chim argues there is still a need for improvement, and concludes that only the use of digital PCR platform and normalisation with a foetal genetic marker will further improve the performance of such epigenetic-based tests. However, digital PCR approaches require the use of technically challenging instruments which are not typically available except in very specialised and localised research laboratories, and the use of a foetal genetic marker (ie, one that is genetically different to the maternal genome) limits the applicability of such tests only to certain foetal-maternal pairs where such a genetic difference exists (such as male foeti and use of ZFY as the genetic marker) and requires the additional sex or genetic testing for such a genetic difference before the NIPT for foetal aneuploidy can be interpreted.

[0012] Accordingly there is a need, from one or more of the above or perspectives, for improved methods, and related other aspects, to detect, indicate or diagnose the presence of a chromosomal aneuploidy in a foetus in particular such methods that are non-invasive and/or use more conventional processing, technical and/or detection steps/equipment, and hence may be conducted faster, more cost effectively and/or more widely in the community.

[0013] Accordingly, it is an object of the present invention to provide alternative, improved, simpler, cheaper and/or integrated means or methods that address one or more of these or other problems. Such an object underlying the present invention is solved by the subject matter as disclosed or defined anywhere herein, for example by the subject matter of the attached claims.

[0014] Generally, and by way of brief description, the main aspects of the present invention can be described as follows:

[0015] In a first aspect, and as may be further described, defined, claimed or otherwise disclosed herein, the invention relates to a method for detecting a chromosomal aneuploidy in a foetus carried by a pregnant female, said method comprising the steps:

(a) providing a sample taken from said pregnant female, which sample comprises DNA that originates from cells of a foetus and/or the placenta of a foetus in admixture with differently methylated DNA of maternal origin;

(b) treating the DNA present in said sample with a reagent that differentially modifies methylated and non-methylated DNA, wherein said reagent is an agent that selectively digests unmethylated over methylated DNA and said agent comprises a methylation sensitive restriction enzyme;

(c) determining an amount of a first target species of DNA, being the chromosome relevant to the chromosomal aneuploidy, in said sample by detecting in said sample the presence of methylation at two or more first target differentially methylated regions (DMRs) located on said chromosome, said first target DMRs differentially methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, wherein said DMRs are hypermethylated in foetal DNA and hypomethylated in maternal DNA, the modification of DNA of the first target DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of

methyated DNA at one or more of said first target DMRs indicates said amount of first target species of DNA in said sample, wherein an amount of the first target species of DNA is determined from the detected amount of methylation at one of the first target DMRs and an amount of the first target species of DNA is determined from the detected amount of methylation at another of the first target DMRs;

(d) determining an amount of reference species of DNA, being one or more reference chromosomes, in said sample by detecting in said sample the presence of methylation at two or more reference DMRs located on said reference chromosome(s), said reference DMRs differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, wherein said DMRs are hypermethylated in foetal DNA and hypomethylated in maternal DNA, the modification of DNA of such reference DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of methylated DNA at one or more of said reference DMRs indicates said amount of reference species of DNA in said sample; and

(e) determining relative amount(s), preferably ratio(s), of an amount determined from step (c) and an amount determined from step (d), wherein two or more such relative amounts are determined for two or more amounts of first target species of DNA, each in respect of one or more of said first target DMRs, and wherein two or more of said relative amounts indicate the presence or absence of the chromosomal aneuploidy in the foetus,

wherein, said detections in step (c) and step (d) are made using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said reference DMRs, and using: (x) the same detectable label(s) for at least two of said reference DMRs; and (y) a different detectable label(s) for at least two of said first target DMRs; and:

wherein in step (e) said two or more relative amounts or ratios are compared with threshold(s) and/or reference distribution(s), wherein said two or more relative amounts or ratios higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus.

[0016] The figures show:

FIGURE 1 depicts: (a) a schematic representation of the differentially methylated region(s) ("DMR") and other regions(s) ("OR") used in a first differential methylation-based DNA detection method; and (b) a schematic representation of the differentially methylated regions ("DMR") and other regions(s) ("OR") used in a second differential methylation-based DNA detection method.

FIGURE 2 depicts a schematic representation of the differentially methylated regions ("DMR") and other regions ("OR") used in Example 1.

FIGURE 3 depicts the correlation of the amount of male specific DNA (Y chromosomal-representation) to the foetal cfDNA fraction measured by a differential methylation-based DNA detection method (Example 1) for study twin cases with known foetal genders.

FIGURE 4 depicts the improved sensitivity of the detection of reference chromosomes using a multiplex differential methylation-based DNA detection method compared to detection of reference chromosomes detected using separate reactions of a single DMR. The number of PCR cycles (Cp) required for detection of foetal cfDNA derived from reference chromosomes (Example 2) in a sample using either RASSF1A or TBX3 alone as a single DMR, or as a multiplex (using the same labels) of RASSF1A and TBX3.

FIGURE 5 depicts a schematic representation of the operations conducted by a computer program product of EXAMPLE 3.

FIGURE 6 depicts a schematic representation of the differentially methylated regions ("DMR") (and optional other regions ("OR")) used in EXAMPLE 4 as a method of the present invention.

FIGURE 7 depicts a scatter plot, for each sample from a first run of 59 samples, of the ratio of the amount of chromosome 21 (as determined by one target DMR located on chromosome 21) to the amount of reference chromosomes (5 and 12) [X-axis] against the ratio of the amount of chromosome 21 (as determined by the other target DMR located on chromosome 21) to the amount of reference chromosomes [Y-axis], wherein such amounts are determined by a method of the invention as described in EXAMPLE 4. Points representing women carrying a foetus with trisomy 21 (T21) are marked by "+", and are distinguishable from the non-T21 samples.

FIGURE 8 depicts a scatter plot, for each sample from all three runs, totalling 168 samples, of the normalised ratio of the amount of chromosome 21 (as determined by one target DMR located on chromosome 21) to the amount of reference chromosomes (5 and 12) [X-axis] against the normalised ratio of the amount of chromosome 21 (as determined by the other target DMR located on chromosome 21) to the amount of reference chromosomes, wherein such amounts are determined by a method of the invention as described in EXAMPLE 4. Points representing women carrying a foetus with trisomy 21 (T21) are marked by "+", and are distinguishably clustered from the non-T21 samples, for example as demarcated by the dotted boundary shown. Note that the outlying non-T21 sample referred to in Example 4 lies outside the range of the X-axis (but not within the T21 cluster)

FIGURE 9 depicts a scatter plot of the distance of each point from FIGURE 8 from the centre point of the curved

demarcation line (0.5,0.55) against the run number. Clear separation is shown, across all runs, between the nine T21 samples (in this figure represented by "o") from the non-T21 samples (in this figure represented by "+").

FIGURE 10 depicts a scatter plot, for each sample from all three runs, totalling 168 samples, of the z-score for the ratio of the amount of chromosome 21 (as determined by one target DMR located on chromosome 21) to the amount of reference chromosomes (5 and 12) [X-axis] against the z-score for the ratio of the amount of chromosome 21 (as determined by the other target DMR located on chromosome 21) to the amount of reference chromosomes, wherein such amounts are determined by a method of the invention as described in EXAMPLE 4. Points representing women carrying a foetus with trisomy 21 (T21) are marked by "+", and are distinguishably clustered from the non-T21 samples. Note that the outlying non-T21 sample referred to in Example 4 lies outside the range of the X-axis (but not within the T21 cluster).

[0017] The present invention, and particular non-limiting aspects and/or embodiments thereof, can be described in more detail as follows:

[0018] In a first aspect, and as may be further described, defined, claimed or otherwise disclosed herein, the invention relates to **a method** for detecting a chromosomal aneuploidy in a foetus carried by a pregnant female, said method comprising the steps:

(a) providing a sample taken from said pregnant female, which sample comprises DNA that originates from cells of a foetus and/or the placenta of a foetus in admixture with differently methylated DNA of maternal origin;

(b) treating the DNA present in said sample with a reagent that differentially modifies methylated and non-methylated DNA, wherein said reagent is an agent that selectively digests unmethylated over methylated DNA and said agent comprises a methylation sensitive restriction enzyme;

(c) determining an amount of a first target species of DNA, being the chromosome relevant to the chromosomal aneuploidy, in said sample by detecting in said sample the presence of methylation at two or more first target differentially methylated regions (DMRs) located on said chromosome, said first target DMRs differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, wherein said DMRs are hypermethylated in foetal DNA and hypomethylated in maternal DNA, the modification of DNA of the first target DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of methylated DNA at one or more of said first target DMRs indicates said amount of first target species of DNA in said sample, wherein an amount of the first target species of DNA is determined from the detected amount of methylation at one of the first target DMRs and an amount of the first target species of DNA is determined from the detected amount of methylation at another of the first target DMRs;

(d) determining an amount of reference species of DNA, being one or more reference chromosomes, in said sample by detecting in said sample the presence of methylation at two or more reference DMRs located on said reference chromosome(s), said reference DMRs differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, wherein said DMRs are hypermethylated in foetal DNA and hypomethylated in maternal DNA, the modification of DNA of such reference DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of methylated DNA at one or more of said reference DMRs indicates said amount of reference species of DNA in said sample; and

(e) determining relative amount(s), preferably ratio(s), of an amount determined from step (c) and an amount determined from step (d), wherein two or more such relative amounts are determined for two or more amounts of first target species of DNA, each in respect of one or more of said first target DMRs, and wherein two or more of said relative amounts indicate the presence or absence of the chromosomal aneuploidy in the foetus,

wherein, said detections in step (c) and step (d) are made using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said reference DMRs, and using: (x) the same detectable label(s) for at least two of said reference DMRs; and (y) a different detectable label(s) for at least two of said first target DMRs; and:

wherein in step (e) said two or more relative amounts or ratios are compared with threshold(s) and/or reference distribution(s),

wherein said two or more relative amounts or ratios higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus.

[0019] Terms as set forth herein are generally to be understood by their common meaning unless indicated otherwise. Where the term "comprising" or "comprising of" is used herein, it does not exclude other elements. For the purposes of the present invention, the term "consisting of" is considered to be a particular embodiment of the term "comprising of". If hereinafter a group is defined to comprise at least a certain number of embodiments, this is also to be understood to disclose a group that consists of all and/or only of these embodiments. Where used herein, "and/or" is to be taken as specific disclosure of each of the two specified features or components with or without the other. For example "A and/or

B" is to be taken as specific disclosure of each of (i) A, (ii) B and (iii) A and B, just as if each is set out individually herein. In the context of the present invention, the terms "about" and "approximately" denote an interval of accuracy that the person skilled in the art will understand to still ensure the technical effect of the feature in question. The term typically indicates deviation from the indicated numerical value by $\pm 20\%$, $\pm 15\%$, $\pm 10\%$, and for example $\pm 5\%$. As will be appreciated by the person of ordinary skill, the specific such deviation for a numerical value for a given technical effect will depend on the nature of the technical effect. For example, a natural or biological technical effect may generally have a larger such deviation than one for a man-made or engineering technical effect. Where an indefinite or definite article is used when referring to a singular noun, e.g. "a", "an" or "the", this includes a plural of that noun unless something else is specifically stated.

[0020] In certain embodiments of the present invention, the pregnant female is a human or a non-human animal, where such non-human animal may, in particular embodiments, be selected from the group consisting of: horse, sheep, cow, pig, chicken, mouse and rat. In a more specific embodiment, the pregnant female may be suspected of being at increased risk of developing or suffering (or suffering from) a medical condition, such as one or more of the medical conditions disclosed herein. Such a method of the present invention is not intended to be practiced on the human or animal body; for example it is intended to be practiced in an in-vitro manner.

[0021] In certain embodiments of the present invention, said species of DNA and/or said differently methylated DNA is/are cell-free DNA, and in particular of such embodiments is/are circulating cell-free DNA. In one particular embodiment, said species of DNA and the differently methylated DNA that is admixed therewith are circulating cell-free DNA. The term "cell-free DNA" (or "cfDNA") is art recognised, and includes the meaning of DNA that is found outside of a cell, such as in a biological fluid (eg blood, or a blood fraction) of an individual. "Circulating" is also an art-recognised term, and includes the meaning that an entity or substance (eg cfDNA) is present in, detected or identified in, or isolated from, a circulatory system of the individual, such as the blood system or the lymphatic system. In particular, when cfDNA is "circulating" it is not located in a cell, and hence may be present in the plasma or serum of blood, or it may be present in the lymph of lymphatic fluid.

[0022] The term "differentially methylated region" or "DMR" will be recognised by the person of ordinary skill in the art, and is also intended to refer to a region in chromosomal DNA that is differentially methylated (eg at a CpG motif) between said (foetal) species of DNA and the other (maternal) DNA with which it is admixed in the sample. For example, the DMRs used in the present invention are differentially methylated between foetal and maternal DNA. In the present invention, the DMRs are hypermethylated in foetal DNA and hypo methylated in maternal DNA. That is, in such regions exhibit a greater degree (ie more) methylation in said (foetal) species of DNA as compared to the other (maternal) DNA, such as about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% of, or more of, the sites available for methylation at a given DMR are methylated in said (foetal) species of DNA as compared to the same sites in the other (maternal) DNA.

[0023] As will be understood by the person of ordinary skill, the term "an amount" can represent variables such as a concentration, a genomic equivalent, a molar number or a mass, as well as a value that is a surrogate for any of such variables. For example, a (quantitative) signal or measurement, such as one generated by analogue or digital (counting) quantification techniques, may also be considered an "amount". In particular, for those embodiments of the methods of the present invention that use quantitative PCR, the amount may be represented by a measurement collected by the qPCR apparatus, such as a Ct (Threshold cycle) [also called Cp, Cross point cycle, for LightCycler terminology]: the cycle number at which the (fluorescence) signal from qPCR crosses a threshold. As will be understood by the person of ordinary skill, such Ct (or Cp) numbers inversely correlate with initial concentrations (amounts) of the DNA template being quantitated (eg the species of DNA that is differentially methylated).

[0024] In step (e) of the method of the present invention, the relative amount(s) of an amount determined from step (c) and an amount determined from step (d) is indicative of the presence or absence of the chromosomal aneuploidy in the foetus. For example, an excess or increase of the amount of the (first) target species of DNA (being the chromosome relevant to the chromosomal aneuploidy) compared that which would be expected for a euploid sample (eg, by comparison to the amount of reference species of DNA, being one or more reference chromosomes) represents that an excess amount of the target species of DNA is present and indicates that a partial or complete extra copy of the chromosome relevant to the chromosomal aneuploidy is present in the foetus. In particular, such an excess of an amount of target species of DNA can indicate a partial or complete chromosomal trisomy, such as human trisomy 21 (T21). In a contrasting example, a deficient or a reduction of the amount of the (first) target species of DNA (being the chromosome relevant to the chromosomal aneuploidy) compared that which would be expected for a euploid sample (eg, by comparison to the amount of reference species of DNA, being one or more reference chromosomes) represents that a reduced amount of the target species of DNA is present and indicates that all or part of at least one copy of the chromosome relevant to the chromosomal aneuploidy is absent in the foetus. In particular, such a reduced amount of target species of DNA can indicate a partial or complete chromosomal loss (eg a monosomy), such as Turner syndrome (45,X).

[0025] The amounts determined from step (c) and an amount determined from step (d) can be compared from a theoretical perspective. For example, for a human trisomy (such as T21, T18 and/or T13), the amount of the (first) target species of DNA (being the chromosome relevant to the chromosomal aneuploidy) should theoretically have a ratio of

amounts equivalent to 3:2 compared to the amount of reference species of DNA (being one or more reference chromosomes). However, with differences in detection sensitivities and quantification for different regions of target and/or reference chromosomes (and/or DMRs), such theoretical ratios may not be achievable in practice. Accordingly, the presence of the chromosomal aneuploidy in the foetus may be indicated where an amount determined from step (c) is not equivalent to an amount determined from step (d), where equivalence of the amounts does not mean that they have the same value. Indeed, the presence of the chromosomal aneuploidy in the foetus may be indicated by a difference in, distortion of or otherwise a bias in the amount determined from step (c) compared to that expected for a euploid foetus, such as from the amount determined from step (d).

[0026] As will be appreciated by the person of ordinary skill, a definitive indication of the presence (or absence) of the chromosomal aneuploidy in the foetus may not be possible in all circumstances or samples. Accordingly, the present invention also envisions methods whereby in step (e) the indication of the presence (or absence) of the chromosomal aneuploidy in the foetus is represented as a likely or as a possible presence (or absence) rather than a definitive indication. Such envisioned methods include embodiments wherein said method includes an additional step of signalling or flagging that an additional diagnostic test is conducted to provide more certainty, confidence or a definitive diagnosis. Such a subsequent test may include a different NIPT for the detection of foetal aneuploidy (such as NGS-based NIPT as described in EXAMPLE 1) and/or may include CVS or amniocentesis.

[0027] In such method of the present invention, said detections in step (c) and step (d) are made using: (x) the same detectable label(s) for at least two of said reference DMRs; and (y) a different detectable label(s) for at least two of said first target DMRs; and in particular embodiments, said detections in step (c) and step (d) are made using: (x) the same detectable label(s) for each of said at least two reference DMRs; and (y) a different detectable label(s) for each of said at least two first target DMRs.

[0028] In alternative methods described herein, said detections in step (c) and step (d) are made using: (x) the same detectable label(s) for at least two of said first target DMRs; and/or (y) a different detectable label(s) for at least two of said reference DMRs; and in particular of such embodiments, said detections in step (c) and step (d) are made using: (x) the same detectable label(s) for each of said at least two first target DMRs; and (y) a different detectable label(s) for each of said at least two reference DMRs.

[0029] In a further alternative method described herein, said detections in step (c) and step (d) are made using: (x) the same detectable label(s) for at least two (preferably each) of said reference DMRs; and (y) the same detectable label(s) for at least two (preferably each) of said first target DMRs, wherein the detectable label(s) used for (x) is/are different to the detectable label(s) used for (y).

[0030] In alternative methods described herein, said detections in step (c) and step (d) are made using: (x) a different detectable label(s) for at least two (preferably each of said first target DMRs; and/or (y) a different detectable label(s) for at least two (preferably each of said reference DMRs.

[0031] In particular certain embodiments a different detectable label(s) is/are used for such first target DMRs and as is/are used for the reference DMRs and, if OR(s) are used in the method, as is/are used for such OR(s).

[0032] In certain embodiments of the present invention, in step (c) more than two first target DMRs may be used. For example, three, four, five, six, seven, eight, nine or more than nine first target DMRs may be used (such as about 10, 15, 20, 30, 50 or more than 50 of such first target DMRs). In the invention where a different detectable label(s) is used for at least two of said first target DMRs, and then one or more (preferable two, three, four, five or more than five) of the other first target DMRs may be detected using a further different detectable label(s). Alternatively, a different detectable label(s) used for at least two of said first target DMRs, and one or more (preferable two, three, four, five, six, seven, eight or more than eight) of the other first target DMRs may have the same detectable label(s) used for one of said two of said first target DMRs. For example, two first target DMRs may have the same detectable label(s) and a two further first target DMRs may have a different detectable label(s). Such embodiments may provide further advantages to the present invention, such as providing yet further increased sensitivity of specificity without having to resort to additional detectable labels; the number of which may be practically limited given expenses and technical demands of multi-channel detection equipment and eg for fluorescent labels the overlap of detection spectra and colour bleaching.

[0033] A reagent is used in the present invention that differentially (eg selectively) modifies methylated as compared to non-methylated DNA. Treatment of DNA with bisulphite (bisulfite) converts cytosine residues to uracil, but leaves 5-methylcytosine residues unaffected. Thus, bisulphite treatment introduces specific changes in the DNA sequence that depend on the methylation status of individual cytosine residues, yielding single- nucleotide resolution information about the methylation status of a segment of DNA. Various analyses can be performed on the altered sequence to retrieve this information, including the use of PCR primers and/or probes that can distinguish between such single-nucleotide changes.

[0034] Such a reagent comprises a restriction enzyme that is sensitive to the DNA methylation states. Cleavage of such a restriction enzyme's recognition sequence may be blocked, or impaired, when a particular base in the enzyme's recognition site is modified, eg methylated. In all aspects of the invention, the reagent comprises a methylation-sensitive restriction enzyme, such as a methylation-sensitive restriction enzyme disclosed herein; including such embodiments

that comprise two, three, four, five or more of such methylation-sensitive restriction enzymes.

[0035] Prior to step (b), the sample may be processed to isolate, enrich and/or purify, the DNA present therein. For example, a plasma sample may be processed using a cfDNA isolation process or kit to provide a (non-natural) subsequent solution that comprises an admixture of said species of DNA together with the differentially methylated DNA (eg of maternal origin) that does not originate from the foetus and/of the placenta of the foetus. The step of treating in (b) may comprise the step of adding a separate solution that comprises said reagent (eg a methylation sensitive restriction enzyme) to the admixed DNA of the sample (eg, to a non-natural solution comprising such admixed DNA); and/or may comprise maintaining (or changing to) certain conditions. In particular, when said reagent comprises one or more methylation sensitive restriction enzyme, the step of treating in (b) may comprise incubating the DNA and the enzyme(s) together at about 37°C for between about 5min and 300min, such as between about 30min and 90min or about 60min, and optionally may comprise a step of incubating such mixture at a higher temperature (for example, between about 50°C and 90°C, such as about 80°C) so as to deactivate the enzyme(s). In certain embodiments, the composition formed for a treating step of (b) may be non-naturally occurring. For example, particular salts of components of the solution (or buffer); and/or the mixture of (eg human) cfDNA together with one or more bacterial-derived restriction enzymes (or a non-natural mutant thereof) may be a non-natural composition or mixture. Furthermore, any of the methods of the present invention may produce (and hence a composition of the present disclosure may comprise) an in-vitro-produced nucleic acid molecule, such as a DNA product of a PCR reaction (eg a "PCR product"). One or more of such in-vitro-produced nucleic acid molecules may be non-natural because they comprise a nucleotide primer and/or probe that includes at least one detectable label, such a nucleic acid molecule having been generated by polymerase extension (or partial nuclease digestion) of such a labelled primer and/or probe, and hence providing at least a fraction of such nucleic acid molecules that include a detectable label, such that even though the nucleic acid sequence of the nucleic acid molecules may comprise a naturally occurring sequence (or fragment thereof), such an in-vitro-produced nucleic acid molecule is non-natural by virtue of (at least) the non-natural detectable label that it includes. In particular embodiments, prior to step (b), the sample taken from said pregnant female may be processed to extract DNA present in said sample; for example, said processes may extract total cell-free DNA from said sample, or said processes may extract and/or enrich for foetal cell-free DNA.

[0036] In the method of the present invention, the relative amount(s) or ratio(s) determined in step (e) are compared with threshold(s) and/or reference distribution(s), wherein two or more of said relative amount(s) or ratio(s) higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus. The threshold(s) and/or reference distribution(s) may come from prior studies, including those of control samples, or they may be generated from a plurality of samples analysed by a method of the invention. For example, if a number of different samples (such as about 10, 20, 30, 40, 50, 60, 80, 90, 10, 120, 150, 175, 200, 250, 500 or more than about 500 samples) are so analysed, and one or other individual samples show an amount from (c) or a relative amount from (e) that is an outlier to the population of analogous amounts from most other samples, then this indicates that the foetus carried by the pregnant female from which the individual sample was taken has a chromosomal aneuploidy. As described above, it will be appreciated that even if the relative amount(s) or ratio(s) determined in step (e) are compared with threshold(s) and/or reference distribution(s), a definitive indication of the presence (or absence) of the chromosomal aneuploidy in the foetus may not be possible in all circumstances or every sample. Accordingly in such embodiments of the present invention, also include those where a likely or possible presence (or absence), rather than a definitive indication, of the chromosomal aneuploidy is indicated, including those embodiments that include an additional step of signalling or flagging that an additional diagnostic test is conducted to provide more certainty or a definitive diagnosis. Exemplary additional diagnostic tests include any of those are described elsewhere herein, in particular such as NGS-based NIPT as described in EXAMPLE 1.

[0037] An indication that the chromosomal aneuploidy is present in the foetus may instead or additionally be assessed by considering: (i) the increase (or decrease) of a measure or signal representing an amount of the first target species of DNA (such as one determined from one or other of the first target DMRs) compared to that expected in relation to a measure or signal representing an amount of the reference species of DNA (such as one determined from the reference DMRs); and/or (ii) into which a measure or signal representing an amount of the first target species of DNA (such as one determined from one or other of the first target DMRs) determined from the sample falls, from consideration of a reference distribution.

[0038] In certain embodiments, the threshold amount(s) may be established by a standard control; for example, established experimentally from a known sample (or a plurality of known samples) once or separately, and/or a threshold amount(s) that is established (eg a from a sample or plurality of known samples) at about the same time as the test sample (or test samples), such as in the same run, in particularly by establishing the threshold amount(s) by practicing a method of the present invention on samples contained in wells of a microtiter plates where one or more known samples placed in one or more (separate) wells and one or more test samples placed in other wells. In other embodiments of the present invention, a comparison with a threshold amount and/or reference distribution of amounts is made from the relative amount (such as the ratio of) an amount of the first (or second) target species of DNA (ie, a chromosome relevant

to the chromosomal aneuploidy) to an amount of the reference species of DNA (ie, the reference chromosome). For example, from a theoretical perspective an amount of the first target species of DNA originating from a normal diploid set of human chromosomes 21 would be expected to show about a 2:2 (ie 1:1) ratio to the amount of the reference species of DNA originating from a reference (diploid) set of eg chromosome 2. However, in the event of trisomy 21, such a ratio would be expected to be about 3:2. As will now be understood by the person of ordinary skill, other chromosomal (or partial chromosomal) aneuploidies would be expected to show other theoretical ratios, for example 1:2 in the case of a loss of a complete chromosome, or a partial loss such as a partial deletion of the location of the first species of DNA, compared to the reference chromosome comprising the location of the second species of DNA. In certain embodiments, eg if not differentiated such as by methylation differences, the presence of other DNA (ie in a mixture with such other DNA) such as euploid maternal cfDNA in admixture with aneuploid foetal cfDNA could result in modified such ratios depending on the relative amounts of (euploid) maternal and (aneuploid) foetal cfDNAs. As will also be understood by the person of ordinary skill, modified such ratios may result from other factors such as the relative reaction (eg PCR reaction) efficiency of each amplicon. Accordingly, in certain of such embodiments, the threshold amount is a ratio that is (detectably and/or significantly) greater or smaller than 2:2 (100%) such as about 3:2 (150%), about 2:3 (66%), about 1:2 (50%) or about 2:1 (200%); including threshold amounts and/or ratios that are greater than about 200%, less than about 50%, or is one selected from the list consisting of about: 190%, 180%, 170%, 160%, 150%, 140%, 130%, 120%, 110%, 105%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, and 55%. Alternatively, in particular embodiments, the threshold amount may be determined merely by there being no detectable amount of the first (or second) species of DNA, such as in Turners syndrome (a human female with a "45, X" karyotype rather than the fully euploid "46,XX" karyotype). However, as described above, differences in detection sensitivities and quantification for different regions of target and/or reference chromosomes (and/or DMRs), such theoretical ratios may not be achievable in practice. Accordingly, the presence of the chromosomal aneuploidy in the foetus may be indicated where an amount determined from step (c) is not equivalent to an amount determined from step (d), where equivalence of the amounts does not mean that they have the same value. Indeed, the presence of the chromosomal aneuploidy in the foetus may be indicated by a difference in, distortion of or otherwise a bias in the amount determined from step (c) compared to that expected for a euploid foetus, such as from the amount determined from step (d).

[0039] In certain embodiments, a parameter (such as a mean, median, standard deviation, median absolute deviation or z-score) is calculated in respect of a set of samples within each run, plate or detection/analysis data-set. In certain of such embodiments, such a calculated parameter is used to identify outliers (such as trisomy samples) from those test samples detected/analysed in such run, plate or data-set (eg, a "run-specific" analysis). In particular embodiments, such a parameter is calculated from all test samples without knowledge of the identity of any outliers (eg a "masked" analysis). In other particular embodiments, such a parameter is calculated from a set of reference samples known to be (non-outlying) standards (such as samples known to contain cfDNA from euploid fetuses) or test samples that are presumed to be (or are unlikely to be) such standards.

[0040] Comparing and detecting differences between sample distributions and reference distributions, or sample outliers from reference distributions will be known to the person of ordinary skill, and include the use of parametric and non-parametric statistical testing such as the use of (one- or two-tailed) t-tests, Mann-Whitney Rank Sum test and others, including the use of a z-score, such as a Median Absolute Deviation based z-score (eg, such as used by Stumm et al 2014, Prenat Diagn 34:185). When comparing a distribution to (or outliers from) a reference distribution, then in certain embodiments of the invention, the comparison is distinguished (and/or identified as being significantly different) if the separation of the means, medians or individual samples are greater than about 1.5, 1.6, 1.7, 1.8, 1.9, 1.95, 1.97, 2.0, or greater than about 2.0 standard deviations ("SD") of the reference distribution; and/or if an individual sample separates from the reference distribution with a z-score of greater than about 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.5, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.75, 4.0, 4.5, 5.0 or greater than about 5.0.

[0041] In certain embodiments, in the context of a data set a z-score (or an equivalent statistic based on the distribution pattern of replicates of a parameter) may be calculated to identify an outlying data point(s) (for example, representing an excessive amount of the species of DNA such as in the context of a test seeking to identify a pregnant female predicted to have or having an increased risk of suffering or developing preeclampsia, or representing an excessive amount of one chromosome over a reference chromosome such as in the context of a test seeking to identify a foetus suffering from a chromosomal aneuploidy), the data representing such data point removed from the data set and a subsequent z-score analysis be conducted on the data set to seek to identify further outliers. Such an iterative z-score analysis may be particularly helpful in detection of foetal chromosomal aneuploidies using a method of the present invention, where sometimes two or more aneuploidy samples in one run may skew a single z-score analysis, and/or where follow-up tests are available to confirm false positives and hence avoiding false negatives is potentially more important than the (initial) identification of false positives.

[0042] The practice of the method of the present invention can enable the relative detection (or amount) of the first (or second) target species of DNA (eg, from a chromosome, or part thereof, related to a chromosomal aneuploidy such as human chromosome 21) and reference species of DNA (for example, from a reference chromosome, or part thereof,

such as chromosome 12, 5 or 2), and hence aid the rapid, simple and cost-effective detection, identification or diagnosis of a chromosomal aneuploidy in a foetus. Such an approach may be more easily established and practiced in many laboratories, requiring for example, a relatively simple, reliable and cost-effective quantitative PCR machine; and not requiring expensive and specialised high-throughput next-generation sequencing machines. Indeed, in the present invention, detection in step (c) of said first target DMRs and said detection in step (d) of said reference DMRs are made using the same aliquot of DNA of said sample, and in the same reaction/detection vessel, and effectively simultaneously for such DMRs, and using: (x) the same detectable labels(s) for at least two (preferably each) of said reference DMRs; and (y) a different detectable label(s) for at least two (preferably each) of said first target DMRs. The relative detection, identification or quantification of the first (or second) target and reference species of DNA (via the target DMRs and the reference DMRs) is conducted in the same reaction/detection vessel and effectively simultaneously, and advantageously may be made by the use of detectable labels that can distinguish at least two of the first (or second) target DMRs from the reference DMRs and, if used, their corresponding ORs detected in optional step (f). In a particular further such embodiment where detection (c) and (d) is conducted in the same reaction/detection vessel and effectively simultaneously includes where said detection in step (c) and said detection in step (d) are made by multiplex real-time quantitative probe-based PCR using at least one labelled probe specific for each of said DMRs and, if used, each OR.

[0043] In all aspects of the present invention, embodiments are included when the presence of the chromosomal aneuploidy is a total or partial such aneuploidy, for a chromosomal abnormality that is associated with a foetal abnormality and/or congenital disorder. For example, such a chromosomal abnormality may be selected from the group consisting of: a trisomy (such as trisomy 21, trisomy 18, or trisomy 13), a sex-chromosome abnormality (such as Turners syndrome, Klinefelter syndrome, [Noonan syndrome,] Triple X syndrome, XXY syndrome, or Fragile X syndrome or XYY syndrome or XYYY syndrome), a chromosomal deletion (such as Prader-Willi syndrome, Cris-du-chat syndrome, Wolf-Hirschhorn syndrome, or 22q11 deletion syndrome, Duchene muscular dystrophy), Beckwith-Wiedemann syndrome, Canvan syndrome, and neurofibromatosis. Of most relevance, in terms of prevalence and hence medical and social significance is where such chromosomal abnormality is a trisomy, such as one selected from the list consisting of trisomy 21, trisomy 18, or trisomy 13.

[0044] In those embodiments of the present invention in which the pregnant female is human, the chromosome relevant to the chromosomal aneuploidy may be a human chromosome selected from the list consisting of: chromosome 21, chromosome 18, chromosome 13, X-chromosome and Y-chromosome, *preferably* chromosome 21, chromosome 18, chromosome 13. In particular of such embodiments, the chromosome relevant to the chromosomal aneuploidy is human chromosome 21. In certain of any of such embodiment, said chromosomal aneuploidy may be an aneuploidy of said chromosome. In all aspects of the present invention, the chromosomal aneuploidy is a monosomy or a trisomy of said chromosome, which as will be known by the person of ordinary skill, may be a partial or complete aneuploidy such as a partial or complete monosomy or trisomy of said chromosome, in which a "partial" aneuploidy includes the meaning of an imbalance of genetic material caused by loss or gain of part of a chromosome. For example, a partial aneuploidy may include the situation of an unbalanced translocation, where an individual carries a derivative chromosome formed through the breakage and fusion of two different chromosomes. In this example, the individual would have three copies of part of one chromosome (two normal copies and the portion that exists on the derivative chromosome) and only one copy of part of the other chromosome involved in the derivative chromosome.

[0045] Correspondingly in those embodiments of the present invention in which the pregnant female is human, one or more of the reference chromosome(s) may be one selected from the list consisting of: human chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, human chromosome 14, 15, 16, 17, human chromosome 19, human chromosome 20 human chromosome 22 and human chromosome 23. In particular of such embodiments, the reference chromosome is human chromosome 5 or human chromosome 12.

[0046] In certain embodiments of the present invention, two or more of said reference DMRs are located on different reference chromosomes. For example, in certain embodiment, one of said reference DMR is located on human chromosome 5 and another of said reference DMR is located on human chromosome 12.

[0047] However, also encompassed within the scope of the present invention are embodiments where one (or more) of the chromosomes used as the reference chromosome is a chromosome which may be associated with a chromosome aneuploidy but a different chromosome to that relevant to the chromosomal aneuploidy for which the detection method is being applied. For example, if the method of the present invention is being practiced to detect trisomy 21 in a foetus carried by a pregnant human female, then for example human chromosome 18 or human chromosome 13, in particular chromosome 13, may be used as a "reference" chromosome. In such an example, given the negligible likelihood of the foetus carrying both a trisomy 21 and a trisomy 13, an excess of chromosome 21 compared to chromosome 13 would indicate a trisomy of chromosome 21. Correspondingly, by use of the same assay but now instead consideration of the chromosome 13 as the being the chromosome relevant to the chromosomal aneuploidy for which the detection method is being applied, then an excess of chromosome 13 compared to chromosome 21 (equivalent to a reduction in chromosome 21 compare to chromosome 13) would indicate a trisomy of chromosome 13. Such a selection of "reference" chromosomes and practice/consideration of the methods of the present invention would have particular advantages of

providing the possibility to detect the presence (or absence) of an aneuploidy at more than one chromosome relevant to a chromosomal aneuploidy in a single assay. Such an assay would not only save time and cost, but would may also enable the detection of the presence of such more than one chromosomal aneuploidy from a single sample; which given the relatively low level of foetal-derived DNA typically present in a maternal sample, would then require fewer sampling or less sample taken from the pregnant female.

[0048] In particular embodiments, the inventive method further comprises the step of:

(f) determining an amount of total DNA in said sample by detecting at least one other region (OR) that is not differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, the modification of which OR(s) by said reagent is insensitive to methylation of DNA,

wherein, said detections in step (c) and step (d) and step (f) are made using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said reference DMRs and said other region(s), and, in one embodiment, using: (x) the same detectable labels(s) for at least two (preferably each) of said reference DMRs; and/or (y) a different detectable label(s) for at least two (preferably each) of said first target DMRs and for at least one (preferably each) of said OR(s); optionally wherein said OR(s) is(are) located on one or more reference chromosome(s), preferably at least one of said ORs being located on the same reference chromosome(s) as at least one of said reference DMRs. In alternative disclosures, detections in step (c) and step (d) and step (f) may be made using: a different detectable label(s) for at least two (preferably each) of said reference DMRs, for at least two (preferably each) of said first target DMRs and for at least one (preferably each) of said.

[0049] In contrast to DMRs, an "other region" ("OR"), if used in the present invention, is not (significantly) differentially methylated between the species of DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin with which it is admixed in the sample. For example, under the conditions and nature of the reagent used, there is no detectable difference between modification by such reagent at the other region of said species of (foetal) DNA as compared to the other region of the admixed (maternal) DNA. Such a non-difference may be achieved if the other region comprises no sites for methylation, if there is no difference in the degree of methylation if such sites are present, or by the use of a reagent that does not recognise any sites of methylation present in the other region. Accordingly, in alternative embodiments of the present invention, the at least one OR used in optional step (f) is one for which no (significant) difference in methylation between said species of (foetal) DNA and the other (maternal) DNA is (or can be) recognised or detected (or recognisable or detectable) with said reagent.

[0050] In certain embodiments of the present inventive method, said OR(s) is(are) located on one or more reference chromosome(s). One or more of said reference chromosome(s) may, in the case of a pregnant human female, be one selected from the list consisting of: human chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, human chromosome 14, 15, 16, 17, human chromosome 19, human chromosome 20 human chromosome 22 and human chromosome 23. In particular of such embodiments, the reference chromosome is human chromosome 5 or human chromosome 12. In certain particular embodiments, at least one of said ORs may be located on the same reference chromosome(s) as at least one of said reference DMRs, for example one of said reference DMRs and one of said ORs may be both located on human chromosome 5 or human chromosome 12. In other embodiments, said OR(s) is(are) located on a chromosome(s) relevant to a chromosomal aneuploidy.

[0051] The OR(s) (that is/are not so differentially methylated), if used in the present invention, may be non-overlapping with one or more of the (eg reference) DMRs used in the present invention. For example, an OR can be located further than about 10bp, 20bp, 50bp, or more than 100bp, 500bp, 1kb or 10kb, away from one (or all) of the (eg reference) DMRs, such as is located between about 20bp and about 20kb upstream or downstream of (including embodiments being located within the same gene as) said (eg reference) DMR. In particular, the genomic location of the OR, if used in the present invention, may generally be located in the same portion of the genome, such as between about 20bp and about 20kb upstream or downstream of (including embodiments being located within the same gene as) the genomic location of at least one of the (eg reference) DMRs used herein. Particularly in the context of foetal fraction of cfDNA, detection (and particularly quantification) of such species of DNA is enhanced (eg, in terms of sensitivity, accuracy and/or precision) if the OR is so located in the same portion of the genome as one of the reference DMRs. Without being bound by theory, it is believed that with such similarly-located DMR(s) and OR, particularly when used in such context, the effect of variation in chromatin/nucleosome packing across the genome - and hence stability/degradation of different regions of genomic DNA - is mitigated, such that any difference in stability/degradation of a (eg reference) DMR (eg detecting the reference chromosome species of foetal DNA) as compared to the OR (is detecting total DNA) is less, and hence a relative (and absolute) quantification may be made without it being (significantly) confounded by quantitative differences brought about by (significantly) differential chromatin/nucleosome packing across the genome between such a DMR and an OR.

[0052] The present invention includes the optional use of one OR in optional step (f) to provide for the detection of an amount of total DNA in the admixture. However, the present invention also encompasses embodiments that use more than one OR. For example, the invention includes such embodiments wherein said detection in step (f) comprises using at least two of said ORs, such as two, three or four of said ORs. In particular embodiments of all aspects of the present

invention, the number of said ORs used in optional step (f) is the same as the number of reference DMRs used in step (d). For example, if two reference DMRs are used then two other regions are used in such an embodiment, and if three reference DMRs are used then three other regions are used (such as depicted in FIGURE 1).

[0053] As described elsewhere herein, the present invention includes embodiments where the optional OR is generally located in the same portion of the genome, such as between about 20bp and about 20kb upstream or downstream of (including embodiments within the same gene as) the genomic location of at least one of the DMRs (such as a reference DMR) used herein. Also as described elsewhere herein, certain embodiments of the present invention include where the OR is generally located in the same portion of the genome, such as between about 20bp and about 20kb upstream or downstream of (including embodiments within the same gene as) the genomic location of at least one of the DMRs used herein (such as a target DMR). In certain embodiments, the OR does not overlap with such a DMR. Accordingly, if multiple ORs are used in the present invention, then embodiments are included where two or more of such ORs are similarly located in the genome to the two or more DMRs (such as reference DMRs). For example, one of said ORs may be located between about 20bp and about 20kb upstream or downstream of (including embodiments within the same gene as) a reference DMR used in step (d) and each other of the said OR (eg, a second OR) is located between about 20bp and about 20kb upstream or downstream of (including embodiments within the same gene as) another of said (eg, non-overlapping) DMRs (eg, a second reference DMR). In certain embodiments an additional OR, may overlap with a DMR (such as a reference DMR).

[0054] An OR used in the present invention, when generally located in the same portion of the genome as a DMR (such as a reference DMR), may be located upstream or downstream of one of said DMRs within a distance selected from the group consisting of: between about 16kb to 20bp, 14kb to 20bp, 12kb to 20bp, 10kb to 20bp, 8kb to 20bp, 6kb to 20bp, 5kb to 20bp, 4kb to 20bp, 3kb to 2bp, 16kb to 20bp, 1kb to 20bp, 500bp to 20bp, 200bp to 20bp, 20kb to 15kb, 15kb to 10kb, 12kb to 8kb, 10kb to 8kb, 11kb to 7kb, 11kb to 10kb, 9kb to 8kb, 8kb to 6kb, 6kb to 4kb, 4kb to 2kb, 2kb to 500bp, 1kb to 100bp, 500bp to 50bp, 400bp to 200bp and 500bp to 100bp and 500bp to 300 bp. In particular embodiments, each OR used in the present invention is so generally located to a different of the DMRs used (such as reference DMRs).

[0055] If multiple ORs are used, then the present invention includes embodiments where the detection in optional step (f) is made using the same detectable label for each of said ORs and/or comprises multiplex real-time quantitative PCR using at least two labelled probes each of which is specific for one of said ORs.

[0056] In certain embodiments of the present inventive method, the detection in step (f), if conducted, comprises using at least two of said ORs. In particular of such embodiments, the number of said ORs may be the same as the number of reference DMRs used in step (d). In a further of such embodiments, one of said reference DMRs and one of said ORs may be both located on one reference chromosome (eg human chromosome 5), and a second reference DMRs and a second of said ORs may be both located on another reference chromosome (eg human chromosome 12). In certain such further embodiments, one of said ORs is located between about 20bp and about 20kb upstream or downstream of a reference DMR used in step (d) and each other of the said ORs is located between about 20bp and about 20kb upstream or downstream of another of said reference DMRs.

[0057] In an embodiment of the present invention, the detection of the various DNA regions, ie the DMR(s) (and the OR(s), if included) may occur in a simplified process. Correspondingly, one feature of the present invention is that the detection of the various DNA regions, ie the DMRs (and the OR(s), if included), may occur in a simplified process. For example, using a single aliquot of DNA from the sample, such DNA regions may be detected in a single vessel. This feature may simplify the method(s), and can provide for more efficient and accurate detection (especially in those embodiments when detection is quantitative). The term "vessel" will be art recognised, and includes embodiments of a vessel (such as a tube, well of a microtiter plate, nano-well, capillary reaction vessel etc) in which a process or procedure comprised in the method occurs, such as a reaction and/or detection process or a step of a method of the present invention. Other such vessels may include droplets in oil/water emulsions, nanoparticles or a hybridisation chamber; as appropriate to the detection technology used. The detectable labels used, in certain embodiments of the present invention may be different for each of the target DMRs, and/or may be the same for each reference DMR and/or may be the same for each OR (if included), provided that, when detected essentially simultaneously, the label(s) used for the target DMRs is/are different (ie, can be separately detected) to the label(s) used for the reference DMR(s) (and for the ORs, if used). Alternatively, the detectable labels used, in certain embodiments of the present invention may be different (eg, not the same) for each DMR and/or may be different (eg, not the same) for each OR (if included). The detectable labels used in the method of the invention may be the same for each reference DMR and, in certain embodiments, may be the same for each OR (if included), provided that the label(s) used for the target DMRs is different, (ie, can be separately detected) to the label(s) used for the reference DMRs. And different to the label(s) used for the ORs (if used). Detectable labels that are "the same", can also include labels while structurally different, are functionally (essentially) similar as they cannot be significantly differentiated by the detection technology employed. For example, structurally different fluorescent dyes may be considered "the same" if their excitation and emission spectra are (substantially or essentially) similar, or overlap to such a degree that they are able to be excited and detected simultaneously with the same wavelength(s). Suitable

labels (and detection modalities) are further described elsewhere herein. Preferably, the detection of the DMR(s) and OR(s) (if included) may be made effectively simultaneously. For example, within the same (reaction/detection) vessel, all such regions (and hence said species of DNA and total DNA) can be detected within less than about 5s, 1s, 0.5s, 100ms, 10ms, 1ms, 100us, 10us or 1us of each other, and for example without transferring the vessel, or the reaction/mixture, to any subsequent vessel, assay or equipment, or for example, without adapting or recalibrating the detection process for (any/either of) the DMR(s) or the OR(s) separately. The use of detectable label(s) - for example at least one for said first target DMR(s) and at least one for the reference DMR(s) - utilises components, process and/or steps that are non-natural. For example, a composition of two specific labels together with the specific DNA regions would (generally) not be found in nature. In particular, short probes used in quantitative probe-based PCR, while may comprise a DNA sequence that is a fragment of that found in a natural genome, when linked to a one or more labels (such as a fluorescent dye) form a specific labelled fragment that is non-natural.

[0058] By way of graphical description, a schematic representation of the general arrangement of the first target DMR(s), the reference DMRs the OR(s), if used, and the detectable label(s), as used for one embodiment of the present invention, is presented in FIGURE 6. (1a) The presence of methylation in DNA at one first target DMR (DMR1) located on the chromosome relevant to the chromosomal aneuploidy is detected using a detectable label(s); (1b) The presence of methylation in DNA at a second first target DMR (DMR2) located on the same chromosome relevant to the chromosomal aneuploidy is detected using a different detectable label(s); (1') The presence of methylation in DNA at two reference DMRs is detected with the same detectable label(s) (but different to those used to detect the first target DMRs); (3') The detection of the reference DMRs may be detected in the context of two optional other regions ("OR1" and "OR2'')) using the same detectable label(s) (but different to those used to detect the DMRs); (2) One or more of such ORs may be located in the same chromosome as a reference DMR; in particular (4) an OR may be in the same portion of the genome (eg, between about 20bp and about 20kb upstream or downstream of) a reference DMR.

[0059] In the context of the present invention, methylation at any particular DMR used in the inventive method (such as first target DMR or a reference DMR) may be detected in the context of at least one OR. By way of graphical description, a schematic representation of one possible general arrangement of such DMR(s), the OR(s) and the detectable label(s), as used for one embodiment of such DMR detection, is presented in FIGURE 1(a). (1) The presence of methylation in DNA at DMR1 is detected in the context of an other region ("OR1") located within the same portion of the genome (eg, between about 20bp and about 20kb upstream or downstream of) DMR1. (2) Optionally, additional DMRs and/or ORs (such as DMR2 and/or OR2, and up to DMRn and ORn) may be detected, and pairs of such additional DMRs and ORs may each be co-located in the same portion of the genome (eg, between about 20bp and about 20kb upstream or downstream of) as each other. Optionally, (3) the presence of methylation in DNA is detected at multiple DMRs - when located on the same target chromosome or located on one or more reference chromosomes - using the same detectable label(s) and/or (4) the amount of total DNA detected using at least one OR (OR1, and optionally, OR2 or up to ORn is detected using different detectable label(s) to those used to detect methylation at the DMR(s) (optionally, the detectable label(s) used is the same for all the ORs - when located on the same target chromosome or located on one or more reference chromosomes -).

[0060] Also in the context of the present invention, when methylation at two or more DMRs used (such as first target DMR or a reference DMR) - when located on the same target chromosome or located on one or more reference chromosomes - may be detected in using the same detectable label(s). By way of graphical description, a schematic representation of the general arrangement of such DMRs, the OR(s) (if used) and the detectable label(s), as used for one embodiment of such DMR detection, is presented in FIGURE 1(b). (1) The presence of methylation in DNA at two or more DMRs, DMR1 and DMR2 (and, optionally, up to DMRn) - when located on the same target chromosome or located on one or more reference chromosomes, is in each case detected using the same detectable label(s). (2) Optionally, an other region ("OR") may be located within the same portion of the genome (eg, between about 20bp and about 20kb upstream or downstream of) one of the DMRs). (3) The amount of total DNA detected using at least one OR (OR1, and optionally, OR2 or up to ORn) is detected using different detectable label(s) to those used to detect methylation at the DMRs (optionally, the detectable label(s) used is the same for all the ORs - when located on the same target chromosome or located on one or more reference chromosomes). (4) Optionally, methylation at more than two DMRs is so detected, and/or the amount of total DNA is detected at more than one OR.

[0061] The use of such a combination of features in the present invention provides opportunity for efficiency improvements and/or synergistic enhancement of outcome. For example, an improved sensitivity and/or accuracy and/or specificity and/or precision of detection (eg, detection of a quantitative amount) or said species of (foetal) DNA, in particular of the reference chromosome, can be obtained by the use of such a combination; the degree of improvement of which can be synergistic, as compared to the use of each feature alone; eg the enhancement obtained by use of the combined features being greater than the sum of each enhancement obtained by the use of each feature individually.

[0062] As will now be apparent to the person of ordinary skill following the disclosure of the present invention, an analogous and additional step to step (c) may be conducted in the method so as to detect methylation - and hence determine/quantitate/compare - the amount of a second target chromosome (eg relative to the reference chromosome(s))

where such second target chromosome is also a (eg different) chromosome relevant to a (eg different) chromosomal aneuploidy. Such a method that is able to detect the presence of two (or more) different chromosomal aneuploidies in a foetus carried by a pregnant female has particular advantages in terms of cost, speed and convenience.

[0063] Accordingly, in certain embodiments of the present invention, then method further comprises the steps of:

(c)' determining an amount of a second target species of DNA, being a chromosome relevant to a different chromosomal aneuploidy, in said sample by detecting in said sample the presence of methylation at two or more second target DMRs located on the chromosome relevant to said different chromosomal aneuploidy, said second target DMRs differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, the modification of DNA of the second target DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of methylated DNA at one or more of said second target DMRs indicates said amount of second target species of DNA in said sample; and

(e)' determining relative amount(s), preferable ratio(s), of an amount determined from step (c)' and an amount determined from step (d), wherein one or more of said relative amount(s) indicates the presence or absence of the different chromosomal aneuploidy in the foetus,

preferably wherein, said detections in step (c) and step (c)' and step (d) and optional step (f) are made using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said second target DMRs and said reference DMRs, and, optionally, using: (x) the same detectable label(s) for at least two (preferably each) of said reference DMRs; and/or (y) a different detectable label(s) for at least two (preferably each) of said first target DMRs and for at least two (preferably each) of said second target DMRs and for the optional OR(s).

[0064] In the context of the second target DMRs, the various possible arrangements of detectable label(s) used for such second target DMRs may be as described for the first target DMRs, provide however that in certain embodiments a different detectable label(s) is/are used for such second target DMRs that is/are used for such first target DMRs and as is used for the reference DMRs and, if OR(s) are used in the method, as is/are used for such OR(s).

[0065] In the context of such embodiment of the present invention when the method is practiced to seek to detect two or more chromosomal aneuploidies in a foetus carried by a pregnant human female, the chromosome relevant to the different chromosomal aneuploidy may be a human chromosome selected from the list consisting of: chromosome 21, chromosome 18, chromosome 13, X-chromosome and Y-chromosome, in particular where such chromosome is different to the first target chromosome. For example, the chromosome relevant to the different chromosomal aneuploidy may be human chromosome 18 or chromosome 13, in particular human chromosome 18: wherein said different chromosomal aneuploidy is an aneuploidy of said different chromosome, such as a trisomy of said different chromosome.

[0066] In particular of such embodiments when the method is practiced to seek to detect two or more chromosomal aneuploidies in a foetus carried by a pregnant human female, said chromosome relevant to the chromosomal aneuploidy and said different chromosome relevant to the different chromosomal aneuploidy may be the pairs of human chromosomes selected from the list consisting of: human chromosomes 21 and 18, human chromosomes 21 and 13, human chromosomes 18 and 13; wherein said chromosomal aneuploidy and said different chromosomal aneuploidy is an aneuploidy of the respective chromosome, *preferably* a trisomy of the respective chromosome.

[0067] In more particular of such embodiments when the method is practiced to seek to detect two or more chromosomal aneuploidies in a foetus carried by a pregnant human female, each of said first target DMRs may be located on human chromosome 21, on human chromosome 18 or on human chromosome 13, *preferably* on human chromosome 21; and optionally, if present each of said second target DMRs may be located on human chromosome 18 or on human chromosome 13.

[0068] The present invention also encompasses where further analogous additional steps to step (c) (and step (e)) are practiced; each further analogous additional step using further target DMRs to indicate if a different chromosome aneuploidy (such as two, three, four or more than four chromosome aneuploidies) is present in the foetus. For example, one step (c) (and step (e)) may be configured to detect an aneuploidy at human chromosome 21, a first additional step (c) (and step (e)) may be configured to detect an aneuploidy at human chromosome 18 and a second additional step (c) (and step (e)) may be configured to detect an aneuploidy at human chromosome 13.

[0069] Any possible embodiment described herein as being applicable to a step (c) (or a step (e)), may equally be applicable to any of such analogous additional step (c) (or a step (e)), such as to step (c)' (or step (e)'). For example, in particular of such embodiments of the present invention with a step (e)', the method may further include wherein in step (e)' said relative amount(s) or ratio(s) are compared with threshold(s) and/or reference distribution(s), wherein one or more of said relative amount(s) or ratio(s) higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the different chromosomal aneuploidy in the foetus.

[0070] In certain embodiments, prior to or as part of the detection that occurs as part of a step (c) and/or a step (d) and/or optional step (c)' and/or optional step (f) of any method of present invention, each DNA region comprising said DMR(s) and/or said optional OR(s), respectively, is(are) amplified. Amplification of DNA may be conducted using any

suitable replication process, and in particular of such embodiments, each of the DMR(s) and/or optional OR(s), is amplified by a polymerase chain reaction (PCR) using primers suitably designed for each DMR and/or OR. The person of ordinary skill will readily be able to design such PCR primers for use in the method of the invention, for example by use of primer design algorithms and programs such as Clone Manager Professional 9 (Sci-Ed Software), Vector NTI (Life Technologies), or web-based tools such as those found from www.ncbi.nlm.nih.gov/tools/primer-blast/ or molbiol-tools.ca/PCR.htm. Those embodiments of the present invention that comprise PCR amplification can further comprises specific steps that are related to the practice of PCR, such as any of those described herein, or in particular the steps of: (A) providing a reaction mixture comprising a double-stranded target DNA, a pair of primers (for example, a pair of primers disclosed herein) designed to amplify a region of such DNA (such as a DMR or an optional OR as described herein) wherein the first primer is complementary to a sequence on the first strand of the target DNA and the second primer is complementary to a sequence on the second strand of the target DNA, Taq polymerase, and a plurality of free nucleotides comprising adenine, thymine, cytosine and guanine; (B) heating the reaction mixture to a first predetermined temperature for a first predetermined time to separate the strands of the target DNA from each other; (C) cooling the reaction mixture to a second predetermined temperature for a second predetermined time under conditions to allow the first and second primers to hybridise with their complementary sequences on the first and second strands of the target DNA, and to allow the Taq polymerase to extend the primers; and (D) repeating steps (B) and (C) at least 20 times.

[0071] The term "probe-based" quantitative PCR is art recognised, and encompasses various embodiments described and marketed under different brand names (such as "TaqMan" PCR of Roche), and uses a (eg fluorescent) reporter probe that is specific for the detection of a given amplicon (eg a DMR or an other region). Probe-based quantitative PCR is distinct from quantitative PCR using double-stranded DNA-binding dyes (eg SYBR Green) as reporters, as such double-stranded DNA-binding dyes bind non-specifically to any double-stranded amplicon and eg cannot be used to distinguish between detection of the DMR(s) (ie said species of DNA) from detection of the other region(s) (ie detection of total DNA). As the person of ordinary skill will appreciate, a specific amplicon of PCR may be detected using a single probe or by using multiple probes (such as two or three probes) for an amplicon. In particular, probe-based quantitative PCR can include amplification reactions into which have been incorporated processes of detecting a target nucleic acid using labelled oligonucleotides that use the 5' to 3' nuclease activity of a nucleic acid polymerase to cleave annealed labelled oligonucleotide (eg the probe) from hybridised duplexes and release labelled oligonucleotide fragments for detection. Such approaches and processes are known in the art and are described in more general terms by Gelfand et al (US5804375, EP0543942 and related family members) and/or Livak et al (US6258569, EP0792374 and related family members), and include where the probe comprises a detectable label in combination with a quencher molecule that quenches the detectability of the label when bound, such that 5' to 3' nuclease (and hence amplification) is required to occur before the detectable label is released into the reaction mixture (away from the quencher) and hence may be detected. Furthermore, "probe-based" quantitative PCR approaches may be alternatively or additionally enhanced by the use of probes that comprise an oligonucleotide-fluorophore-quencher-minor groove binder conjugates, such as described by Reed et al (US6727356, EP1235938 and related family members).

[0072] Such probe-based quantitative PCR may be conducted in an analogue-approach, using a machine such as a LightCycler in which the intensity of signal (eg over time) is measured and used to quantitatively determine detection. Systems and approaches for such detection are described by Woudenberg et al (US6929907, EP0706649 and related family members) and/or Higuchi (US5994056, EP0512334 and related family members). Alternatively, digital PCR (dPCR), ie, PCR conducted in multiple events so as to determine the number of amplification events as method to quantitate an amount of detected DNA. For example, dPCR that is conducted in nano-wells or droplets (ddPCR). Commercially available suppliers of ddPCR technology include Bio-Rad and Rain Dance.

[0073] The person of ordinary skill will be able to design suitable primers and probes (and with suitable labels, eg dyes) for probe-based quantitative PCR detection of the DMRs and/or OR(s); for example by using primer/probe design software as described elsewhere herein. As will be known, the PCR primers may overlap methylation site(s) specific for the methylation-specific modifying reagent used in the methods, in particular when the reagent comprises one or more methylation sensitive restriction enzyme, such as one (or a combination thereof) as disclosed herein. In particular such embodiments, one or other (or when considered together, both) of the PCR primers for a given DMR may overlap two or three such methylation sites (such as two or three restriction sites for methylation-sensitive restriction enzymes, each of which may comprise, or comprises, a methylation site). Alternatively or in addition, the primers for a DMR may be designed to flank one, two, three or more such methylation sites, such as up to 10, 15, 20, 25 or 50 such methylation sites, in particular flanking restriction sites for one, two, three or more such methylation sites, such as up to 10, 15, 20, 25 or 50 methylation-sensitive restriction enzymes, each of which may comprise, or comprises, a methylation site.

[0074] In other embodiments, a detectable label used in step (c) and/or a step (d) and/or optional step (c)' and/or optional step (f) of a method of the invention is independently selected from the group consisting of: fluorescent, protein, small molecule or radioactive label. For example, fluorescent labels that are the same (including, by having similar or overlapping excitation and/or emission spectra) may be used for the DMR(s), and a fluorescent label that has an excitation and/or emission spectra (in particular, a different emission spectrum) may be used for detection of the optional OR(s).

The person of ordinary skill will be able to select appropriate such fluorescent label(s) for use in the present invention from, for example, the group consisting of: FAM, TET, JOE, VIC, HEX, NED, PET, ROX, TAMRA, Quasar and Texas Red and LCCyan500, 6FAM, Cy5, and LCRed610. In other embodiments, a detectable label may be a protein or small molecule tag that, for example, can be detected using a specific antibody and ELISA-type detection approaches. The use of the same protein or small molecule or a detectably different protein or small molecule as applicable for the same or different labels and appropriate detection of the various DMRs and/or optional OR(s), may also be utilised for the detectable label(s) used in the present invention. Different radioactive labels may be distinguished by their emission energy, penetration/excitation characteristics and particle-type (for example, by distinguishing between alpha and beta particles). Other detectable labels (such as nucleic-acid coded tag) may also be employed in the present invention.

[0075] In particular embodiments of the present invention, the detection in step (c) and/or step (c)' of a method of the example comprises real-time quantitative probe-based PCR, eg by using at least one labelled probe which is specific for one of the first target DMR(s) and/or optional second target DMR(s). In those embodiments where PCR amplification of multiple DMRs and/or optional OR(s) is made in the same reaction, such PCR can be considered as "multiplex" (or "duplex" if only two DMRs and/or OR(s) are so amplified). Likewise, the detection in step (d) in the methods of the invention may, in addition or alternatively, comprise real-time quantitative probe-based PCR, such as by using at least one labelled probe specific for one of said reference DMR(s). Also likewise, the detection in optional step (f) in the methods of the invention may, in addition or alternatively, comprise real-time quantitative probe-based PCR, such as by using at least one labelled probe specific for one of said OR(s).

[0076] The primers and/or probes used in the methods of the present disclosure may be configured or designed to detect methylation at one or more of the DMRs by sequence specific detection following eg bisulphite conversion of methylated CpGs to uracil. By way of non-limiting example, such detection may occur MSP or MethylLight as described elsewhere herein.

[0077] In particular embodiments of the present invention, the detection in step (c) of a method of the invention comprises real-time quantitative probe-based PCR, eg by using at least two labelled probes, each of which is specific for one of said two or more first target DMRs; and/or said detection in step (d) comprises multiplex real-time probe-based quantitative probe-based PCR using at least two labelled probes each of which specific for one of said reference DMRs.

[0078] In other embodiments of the present invention, said detection in step (f) comprises real-time quantitative PCR using at least one labelled probe specific for one of said OR(s).

[0079] When two or more of said OR(s) are used, the present invention includes embodiments where detection in optional step (f) is made using: (x) the same detectable label(s) for each of said ORs; and /or wherein said detection in step (f) comprises multiplex real-time quantitative probe-based PCR using at least two labelled probes each of which is specific for one of said ORs.

[0080] In particular embodiments, one or more (preferably all) detection steps (ie, those required for DMR(s) and OR(s)) are conducted in an efficient and effective manner using multiplex quantitative probe-based (eg TaqMan) PCR, in one process step or reaction. For example, in particular embodiments of the present invention, said detection step (c) and step (d), and optional step (c)' and optional step (f), are made using the same aliquot of DNA of said sample, and in the same reaction/detection vessel, and effectively simultaneously with each other, and by multiplex real-time quantitative probe-based PCR using at least one labelled probe specific for each of the said DMRs and optional OR(s). In particular of such embodiments, the reagent comprises one or more methylation sensitive restriction enzyme, such as one (or a combination thereof) as disclosed herein.

[0081] The present invention may also include further procedures, such as one or more control procedures. For example, the present invention can include one or more steps directed to the detection of a third class of DNA region that acts as a control for the modification step (eg, as a control for restriction enzyme digestion). Such embodiments may, for example, also be conducted using multiplex real-time quantitative probe-based PCR wherein such control region is amplified and detected by a further set of primer/probe(s) with a further detectable label used for such class of region.

[0082] In the present invention, said target and reference species of DNA originate from cells of a foetus and/or the placenta of a foetus and said sample is from a pregnant female. In certain embodiments, the sample may be obtained in a non-invasive manner. For example, said species of DNA is circulating cell-free DNA that has been detected from the sample being blood or a blood fraction (such as plasma or serum) that has been obtained from the pregnant female by conventional means such as a blood collection tube. In such embodiments, the sample can comprise DNA that has a maternal origin; that is it originates from cells (and hence the genome of) the pregnant female.

[0083] In all aspects of the present invention, there exist embodiments wherein the sample is a tissue sample or a sample of biological fluid. In particular, the sample is whole blood or a blood fraction (eg, such as plasma or serum). In alternative embodiments, the sample is biological fluid selected from the group consisting of: urine, saliva, sweat, tears, phlegm, breast milk, breast aspirate, vaginal secretion, vaginal wash and colonic wash. In more particular embodiments, the sample is a plasma or serum sample from the pregnant female, or is urine from the pregnant female. In other embodiments, the sample is largely (or essentially) free from cells, and/or is not a whole blood sample.

[0084] In particular embodiments of all aspects of the invention, said species of DNA that originate from cells of a foetus and/or the placenta of a foetus is circulating cell-free DNA and said sample is a blood fraction such as plasma or serum.

[0085] In the present invention the DMR(s) (such as the first or second target DMRs and the reference DMRs) are hypermethylated in foetal DNA and hypo methylated in maternal DNA. In certain embodiments, such a DMR may be located in a promoter, enhancer region or an exon of a gene, such as a gene disclosed herein, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. Alternatively, a DMR may be located in an intron of such a gene, or in a non-coding region of the genome. In particular embodiments of all aspects of the present invention, such genome and/or gene is a human genome or gene.

[0086] Specifically included in the present invention are embodiments wherein said DMR(s) comprises at least one, preferably at least two, methylation site(s) specific for said reagent. For example, comprising three, four, five, six, seven, eight, nine, ten or more than then, such as about 15, 20, 25, 50 or more than 50 methylation site(s) specific for said reagent.

[0087] In the context of the DMRs located on the chromosome relevant to the chromosomal aneuploidy, the present invention encompasses any suitable DMR located on such a chromosome, such as those described anywhere herein.

[0088] In particular embodiments of the present invention, for example when the method is configured to seek the detection of a chromosomal aneuploidy of human chromosome 21, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from one disclosed in WO 2011/092592, including one selected from the list consisting of: EP1, EP2, EP3, EP4, EP5, EP6, EP7, EP8, EP9, EP10, EP11 and EP12 of WO 2011/092592 (SEQ ID NOs: 33-44 of WO 2011/092592), as further investigated in Lim et al 2014, BMC Medical Genomics 7:1).

[0089] In other particular embodiments of the present invention, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from the list consisting of: AIRE (ENSG00000160224; human Autoimmune regulator gene; Chr21:44,285,838-44,298,648 forward strand, GRCh38:CM000683.2), SIM2 (ENSG00000159263; human Single-minded homolog 2; Chr21:36,699,133-36,749,917 forward strand, GRCh38:CM000683.2) and ERG (ENSG00000157554; human ETS-related gene; Chr21:38,380,027-38,661,780 reverse strand, GRCh38:CM000683.2), or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene.

[0090] In other particular embodiments of the present invention, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from the list consisting of: PDE9A (ENSG00000160191; human phosphodiesterase 9A; Chr21:42,653,636-42,775,509 forward strand, GRCh38:CM000683.2), PPP1R2P2 (ENSG00000234008; human protein phosphatase 1, regulatory (inhibitor) subunit 2 pseudogene 2 gene; Chr21:35,887,195-35,887,807 forward strand, GRCh38:CM000683.2), CBR1 (ENSG00000159228; human carbonyl reductase 1 gene; Chr 21:36,069,941-36,073,166 forward strand, GRCh38:CM000683.2), DSCAM (ENSG00000171587; human Down syndrome cell adhesion molecule gene; Chromosome 21: 40,010,999-40,847,139 reverse strand, GRCh38:CM000683.2), C21orf29 (ENSG00000175894; human thrombospondin-type laminin G domain and EAR repeats gene; Chromosome 21: 44,497,892-44,711,580 reverse strand, GRCh38:CM000683.2), HLCS (ENSG00000159267; human holocarboxylase synthetase gene; Chromosome 21: 36,750,888-36,990,236 reverse strand, GRCh38:CM000683.2), or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene, in particular each case a region disclosed in WO 2007/132176; or selected from the list consisting of: CG137, Similarity to FemIA (C.elegans), CG1009 and CG132, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene, in particular each case a region disclosed in WO 2007/132176. Or, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene being C21orf57 (ENSG00000182362; ybeY metalloproteinase; Chromosome 21: 46,286,337-46,297,751 forward strand, GRCh38:CM000683.2), or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. In particular of such embodiments, at least one of said first target DMRs (or at least one optional second target DMR) is located in in DSCAM or C21orf57, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; such as where one of said first (or optional second) target DMRs is located in DSCAM and another of said first (or optional second) target DMRs is located in C21orf57, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene.

[0091] In other particular embodiments of the present invention, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from the list consisting of: SEQ ID No NOs 33, 34, 35, 36, 37, 38, 39, 176, 179, 180, 184, 188, 189, 190, 191, 193, 195, 198, 199, 200, 201, 202, 203, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 221, 223, 225, 226, 231, 232, 233, 235, 239, 241, 257, 258, 259, and 261 of WO 2011/034631, such as independently selected from the list consisting of: SEQ ID No NOs 37, 257, 258 and 259, in particular any such region and/or gene disclosed in WO 2011/034631 as being hypermethylated in foetal DNA relative to maternal DNA. In certain of such particular embodiments of the present invention, one or said first target DMRs is

located in a region and/or gene being SEQ ID NO 37 of WO 2011/034631 (SEQ ID NO.: 51 of the present application) and another of said first target DMRs is located in a region and/or gene being SEQ ID NO.: 258 of WO 2011/034631 (SEQ ID NO.: 185 of the present application)

[0092] In alternative embodiments of the present invention, for example when the method is configured to seek the detection of a chromosomal aneuploidy of human chromosome 18, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from: VAPA-APCDDI (ENSG00000101558; VAMP (vesicle-associated membrane protein)-associated protein A; Chromosome 18: 9,914,002-9,960,021 forward strand, GRCh38:CM000680.2) and maspin (ENSG00000206075; also known as SERPINB5; serpin peptidase inhibitor, clade B (ovalbumin), member 5; Chromosome 18: 63,476,761-63,505,085 forward strand, GRCh38:CM000680.2), or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. In particular such embodiments, said first target DMRs, and/or said optional second target DMRs, are each located in a region of the maspin (aka "SERPINB5") gene that described in EP 1 751 307 as being differentially methylated between a foetus and its mother, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene.

[0093] In other alternative embodiments of the present invention, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from the list consisting of: SEQ ID No NOs 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32 of WO 2011/034631, in particular any such region and/or gene disclosed in WO 2011/034631 as being hypermethylated in foetal DNA relative to maternal DNA.

[0094] In further embodiments of the present invention, for example when the method is configured to seek the detection of a chromosomal aneuploidy of human chromosome 13, said first target DMRs, and/or said optional second target DMRs, are each located in a region and/or gene independently selected from: the list consisting of: SEQ ID No NOs 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 and 20 of WO 2011/034631, in particular any such region and/or gene disclosed in WO 2011/034631 as being hypermethylated in foetal DNA relative to maternal DNA.

[0095] In the context of the DMRs located on the reference chromosome, the present invention encompasses any suitable DMR located on such a chromosome, such as those described anywhere herein.

[0096] In particular embodiments of the present invention, for example when the method is configured to seek the detection of a reference chromosome, at least one of said reference DMRs is (preferably two or more of said reference DMRs each) located in a region and/or gene independently selected from the list consisting of: RASSF1A (ENSG00000068028; Ras association (RaGDS/AF-6) domain family member 1; Chromosome 3: 50,329,782-50,340,980 reverse strand, GRCh38:CM000665.2), TBX3 (ENSG00000135111; T-box 3; Chromosome 12: 114,670,254-114,684,164 reverse strand, GRCh38:CM000674.2), CDC42EP1 (ENSG00000128283; CDC42 effector protein (Rho GTPase binding) 1; Chromosome 22: 37,560,447-37,569,405 forward strand, GRCh38:CM000684.2), PCDHGA1 (ENSG00000204956; protocadherin gamma subfamily A, 1; Chromosome 5: 141,330,571-141,512,981 forward strand, GRCh38:CM000667.2), and SPN (ENSG00000197471; sialophorin; Chromosome 16: 29,662,979-29,670,876 forward strand, GRCh38:CM000678.2); or selected from SOX14 (ENSG00000168875; SRY (sex determining region Y)-box 14; Chromosome 3: 137,764,284-137,766,338 forward strand, GRCh38:CM000665.2) and ZFY (ENSG00000067646; zinc finger protein, Y-linked; Chromosome Y: 2,935,281-2,982,506 forward strand, GRCh38:CM000686.2), or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. Or located in MGC15523 (ENSG00000157637; also known as "SLC38A10", solute carrier family A 10; Chromosome 17: 81,245,000-81,295,547 reverse strand, GRCh38:CM000679.2) or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. In particular of such embodiments, at least one of said reference DMRs is located in TBX3 or PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; such as where one of said reference DMRs is located in TBX3 and another of said reference DMRs is located in PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene.

[0097] In other particular embodiments of the present invention, said reference DMRs are each located in a region and/or gene independently selected from the list consisting of: SEQ ID NOs: 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59 and 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162 and 163 of WO 2011/034631, such as independently selected from the list consisting of: SEQ ID No NOs 52, 118 and 142 of WO 2011/034631, in particular any such region and/or gene disclosed in WO 2011/034631 as being hypermethylated in foetal DNA relative to maternal DNA. In certain of such particular embodiments of the present invention, one or said reference DMRs is located in a region and/or gene being SEQ ID NO 52 of WO 2011/034631 (SEQ ID NO.: 66 of the present application) and another of said reference DMRs is located in a region and/or gene being SEQ ID NO.: 118 of WO 2011/034631 (SEQ ID NO.: 102 of the present application)

[0098] TABLE C shows the conversion of the sequence identifiers used in the WO 2011/034631 and WO 2011/092592 to the sequence identifiers used in the present invention.

TABLE C: Conversion table for sequence identifiers

	SEQ ID NO.:	SEQ ID NO.:	SEQ ID NO.:
	Present invention	WO 2011/034631	WO 2011/092592
5	SEQ ID NO.: 15	1	-
	SEQ ID NO.: 16	2	-
10	SEQ ID NO.: 17	3	-
	SEQ ID NO.: 18	4	-
	SEQ ID NO.: 19	5	-
	SEQ ID NO.: 20	6	-
	SEQ ID NO.: 21	7	-
15	SEQ ID NO.: 22	8	-
	SEQ ID NO.: 23	9	-
	SEQ ID NO.: 24	10	-
	SEQ ID NO.: 25	11	-
20	SEQ ID NO.: 26	12	-
	SEQ ID NO.: 27	13	-
	SEQ ID NO.: 28	14	-
	SEQ ID NO.: 29	15	-
	SEQ ID NO.: 30	16	-
25	SEQ ID NO.: 31	17	-
	SEQ ID NO.: 32	18	-
	SEQ ID NO.: 33	19	-
	SEQ ID NO.: 34	20	-
30	SEQ ID NO.: 35	21	-
	SEQ ID NO.: 36	22	-
	SEQ ID NO.: 37	23	-
	SEQ ID NO.: 38	24	-
	SEQ ID NO.: 39	25	-
35	SEQ ID NO.: 40	26	-
	SEQ ID NO.: 41	27	-
	SEQ ID NO.: 42	28	-
	SEQ ID NO.: 43	29	-
40	SEQ ID NO.: 44	30	-
	SEQ ID NO.: 45	31	-
	SEQ ID NO.: 46	32	-
	SEQ ID NO.: 47	33	-
	SEQ ID NO.: 48	34	-
45	SEQ ID NO.: 49	35	-
	SEQ ID NO.: 50	36	-
	SEQ ID NO.: 51	37	-
	SEQ ID NO.: 52	38	-
50	SEQ ID NO.: 53	39	-
	SEQ ID NO.: 54	40	-
	SEQ ID NO.: 55	41	-
	SEQ ID NO.: 56	42	-
	SEQ ID NO.: 57	43	-
55	SEQ ID NO.: 58	44	-
	SEQ ID NO.: 59	45	-
	SEQ ID NO.: 60	46	-

EP 3 168 309 B1

(continued)

	SEQ ID NO.: Present invention	SEQ ID NO.: WO 2011/034631	SEQ ID NO.: WO 2011/092592
5	SEQ ID NO.: 61	47	-
	SEQ ID NO.: 62	48	-
	SEQ ID NO.: 63	49	-
	SEQ ID NO.: 64	50	-
	SEQ ID NO.: 65	51	-
10	SEQ ID NO.: 66	52	-
	SEQ ID NO.: 67	53	-
	SEQ ID NO.: 68	54	-
	SEQ ID NO.: 69	55	-
15	SEQ ID NO.: 70	56	-
	SEQ ID NO.: 71	57	-
	SEQ ID NO.: 72	58	-
	SEQ ID NO.: 73	59	-
	SEQ ID NO.: 74	90	-
20	SEQ ID NO.: 75	91	-
	SEQ ID NO.: 76	92	-
	SEQ ID NO.: 77	93	-
	SEQ ID NO.: 78	94	-
25	SEQ ID NO.: 79	95	-
	SEQ ID NO.: 80	96	-
	SEQ ID NO.: 81	97	-
	SEQ ID NO.: 82	98	-
	SEQ ID NO.: 83	99	-
30	SEQ ID NO.: 84	100	-
	SEQ ID NO.: 85	101	-
	SEQ ID NO.: 86	102	-
	SEQ ID NO.: 87	103	-
35	SEQ ID NO.: 88	104	-
	SEQ ID NO.: 89	105	-
	SEQ ID NO.: 90	106	-
	SEQ ID NO.: 91	107	-
	SEQ ID NO.: 92	108	-
40	SEQ ID NO.: 93	109	-
	SEQ ID NO.: 94	110	-
	SEQ ID NO.: 95	111	-
	SEQ ID NO.: 96	112	-
45	SEQ ID NO.: 97	113	-
	SEQ ID NO.: 98	114	-
	SEQ ID NO.: 99	115	-
	SEQ ID NO.: 100	116	-
	SEQ ID NO.: 101	117	-
50	SEQ ID NO.: 102	118	-
	SEQ ID NO.: 103	119	-
	SEQ ID NO.: 104	120	-
	SEQ ID NO.: 105	121	-
55	SEQ ID NO.: 106	122	-
	SEQ ID NO.: 107	123	-
	SEQ ID NO.: 108	124	-
	SEQ ID NO.: 109	125	-

EP 3 168 309 B1

(continued)

	SEQ ID NO.: Present invention	SEQ ID NO.: WO 2011/034631	SEQ ID NO.: WO 2011/092592
5	SEQ ID NO.: 110	126	-
	SEQ ID NO.: 111	127	-
	SEQ ID NO.: 112	128	-
	SEQ ID NO.: 113	129	-
	SEQ ID NO.: 114	130	-
10	SEQ ID NO.: 115	131	-
	SEQ ID NO.: 116	132	-
	SEQ ID NO.: 117	133	-
	SEQ ID NO.: 118	134	-
15	SEQ ID NO.: 119	135	-
	SEQ ID NO.: 120	136	-
	SEQ ID NO.: 121	137	-
	SEQ ID NO.: 122	138	-
	SEQ ID NO.: 123	139	-
20	SEQ ID NO.: 124	140	-
	SEQ ID NO.: 125	141	-
	SEQ ID NO.: 126	142	-
	SEQ ID NO.: 127	143	-
25	SEQ ID NO.: 128	144	-
	SEQ ID NO.: 129	145	-
	SEQ ID NO.: 130	146	-
	SEQ ID NO.: 131	147	-
	SEQ ID NO.: 132	148	-
30	SEQ ID NO.: 133	149	-
	SEQ ID NO.: 134	150	-
	SEQ ID NO.: 135	151	-
	SEQ ID NO.: 136	152	-
35	SEQ ID NO.: 137	153	-
	SEQ ID NO.: 138	154	-
	SEQ ID NO.: 139	155	-
	SEQ ID NO.: 140	156	-
	SEQ ID NO.: 141	157	-
40	SEQ ID NO.: 142	158	-
	SEQ ID NO.: 143	159	-
	SEQ ID NO.: 144	160	-
	SEQ ID NO.: 145	161	-
45	SEQ ID NO.: 146	162	-
	SEQ ID NO.: 147	163	-
	SEQ ID NO.: 148	176	-
	SEQ ID NO.: 149	179	-
	SEQ ID NO.: 150	180	-
50	SEQ ID NO.: 151	184	-
	SEQ ID NO.: 152	188	-
	SEQ ID NO.: 153	189	-
	SEQ ID NO.: 154	190	-
55	SEQ ID NO.: 155	191	-
	SEQ ID NO.: 156	193	-
	SEQ ID NO.: 157	195	-
	SEQ ID NO.: 158	198	-

EP 3 168 309 B1

(continued)

	SEQ ID NO.: Present invention	SEQ ID NO.: WO 2011/034631	SEQ ID NO.: WO 2011/092592
5	SEQ ID NO.: 159	199	-
	SEQ ID NO.: 160	200	-
	SEQ ID NO.: 161	201	-
	SEQ ID NO.: 162	202	-
	SEQ ID NO.: 163	203	-
10	SEQ ID NO.: 164	205	-
	SEQ ID NO.: 165	206	-
	SEQ ID NO.: 166	207	-
	SEQ ID NO.: 167	208	-
15	SEQ ID NO.: 168	209	-
	SEQ ID NO.: 169	210	-
	SEQ ID NO.: 170	211	-
	SEQ ID NO.: 171	212	-
	SEQ ID NO.: 172	213	-
20	SEQ ID NO.: 173	214	-
	SEQ ID NO.: 174	221	-
	SEQ ID NO.: 175	223	-
	SEQ ID NO.: 176	225	-
25	SEQ ID NO.: 177	226	-
	SEQ ID NO.: 178	231	-
	SEQ ID NO.: 179	232	-
	SEQ ID NO.: 180	233	-
	SEQ ID NO.: 181	235	-
30	SEQ ID NO.: 182	239	-
	SEQ ID NO.: 183	241	-
	SEQ ID NO.: 184	257	-
	SEQ ID NO.: 185	258	-
35	SEQ ID NO.: 186	259	-
	SEQ ID NO.: 187	261	-
	SEQ ID NO.: 188	-	33
	SEQ ID NO.: 189	-	34
	SEQ ID NO.: 190	-	35
40	SEQ ID NO.: 191	-	36
	SEQ ID NO.: 192	-	37
	SEQ ID NO.: 193	-	38
	SEQ ID NO.: 194	-	39
45	SEQ ID NO.: 195	-	40
	SEQ ID NO.: 196	-	41
	SEQ ID NO.: 197	-	42
	SEQ ID NO.: 198	-	43
50	SEQ ID NO.: 199	-	44

[0099] If two DMRs are used that are located on the same chromosome, then in particular embodiments of all aspects of the present invention, they are not located in the same portion of the genomic and/or gene. For example, such DMRs may be separated by more than about 20kb, or more than about 15kb, 10kb, 8kb, 6kb, 4kb, 2kb, 1kb, 500bp or 200bp. Alternatively, it is envisioned, that when two (or more) DMRs are used in the present invention that are located on the same chromosome, they may, in certain embodiments, be located in the same region or gene (such as one described herein) and, further, may overlap with each other.

[0100] In particular embodiments of the present invention, when two of said reference DMRs are used (or more than

two reference DMRs are being used) at least one (preferable each) of said reference DMRs is located in a portion of the genome and/or gene (preferably that is human) that is RASSF1A and/or TBX3 and/or PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; and/or at least one (preferable each) of said reference DMRs is located between about positions 4,700bp and 5,600bp of RASSF1A (NCBI Reference Sequence: NG_023270.1: Homo sapiens Ras association (RalGDS/AF-6) domain family member 1 (RASSF1), RefSeqGene on chromosome 3; SEQ ID NO.: 13); and/or is located between about positions 1,660bp and 2,400bp of TBX3 (NCBI Reference Sequence: NG_008315.1: Homo sapiens T-box 3 (TBX3), RefSeqGene on chromosome 12; SEQ ID NO.: 14) and/or is located in PCDHGA1 between 141,330,571 to 141,512,981, of NCBI Reference Sequence Homo sapiens chromosome 5, GRCh38.p2 Primary Assembly: NC_000005.10 GI:568815593; SEQ ID No.: 217), or between about 141,492,450 to 141,492,750 or 141,492,550 to 141,492,700 or 141,492,580 to 141,492,7690 of such sequence. In more particular embodiments: (i) one reference DMR comprises one located between about positions 4,700bp and 5,600bp of RASSF1A and a second reference DMR comprises one located between about positions 1,660bp and 2,400bp of TBX3; or (ii) one reference DMR comprises one located between about positions about 141,492,450 to 141,492,750 or 141,492,550 to 141,492,700 or 141,492,580 to 141,492,7690 of PCDHGA1 (NCBI Reference Sequence Homo sapiens chromosome 5, GRCh38.p2 Primary Assembly: NC_000005.10 GI:568815593) and a second reference DMR comprises one located between about positions 1,660bp and 2,400bp of TBX3.

[0101] In particular embodiments, a reference DMR is located in RASSF1A between about positions 4,900bp and 5,500bp, 5,000bp and 5,400bp, or 5,100bp and 5,300bp of RASSF1A; and/or is located in TBX3 between about positions 1,800bp and 2,260bp, 1,920bp and 2,160bp or 1,920bp and 2,080bp of TBX3 (such as SEQ ID No.: 203); and/or is located in DSCAM between about positions 40,841,600bp and 40,841,900bp, 40,841,625bp and 40,841,840bp or 40,841,650bp and 40,841,790bp; and/or is located between about positions 141,492,450 to 141,492,750, 141,492,550 to 141,492,700 or 141,492,585 to 141,492,690 of PCDHGA1 (NCBI Reference Sequence Homo sapiens chromosome 5, GRCh38.p2 Primary Assembly: NC_000005.10 GI:568815593) such as SEQ ID No.: 221.

[0102] In particular embodiments of the present invention, when two of said first (or second) target DMRs are used (or more than two first or second target DMRs are being used) at least one (preferable each) of said first (or second) target DMRs is located in a portion of the genome and/or gene (preferably that is human) that is DSCAM and/or C21orf57, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; and/or at least one (preferable each) of said first (or second) target DMRs is located between about positions 40,841,584 and 40,842,020 of DSCAM; (Down Syndrome Cell Adhesion Molecule; NCBI Reference Sequence Homo sapiens chromosome 21, GRCh38.p2 Primary Assembly: NC_000021.9 GI:568815577, region 40,010,999 to 40,847,113; SEQ ID No.: 200); and/or is located within no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from C21orf57 (YBEY; Chromosome 21: 46,286,337-46,297,751 forward strand, GRCh38:CM000683.2; such gene, including 250bp upstream/downstream flanking regions, SEQ ID No.: 218).

[0103] In particular embodiments, a first (or second) target DMR is located in in DSCAM between about positions 40,841,600bp and 40,841,900bp, 40,841,625bp and 40,841,840bp or 40,841,650bp and 40,841,790bp of DSCAM (with reference to Homo sapiens chromosome 21, GRCh38.p2 Primary Assembly: NC_000021.9 GI:568815577, region), such as SEQ ID No.: 201; and/or is located within about 200bp, 150bp, or 100bp upstream or downstream of C21orf57, such as between about positions 46,297,700 to 46,297,940, 46,297,750 to 46,297,900 or 46,297,790 to 46,297,890 of Chromosome 21 forward strand, GRCh38:CM000683.2, such as SEQ ID No.: 219.

[0104] By way of one non-limiting example, in one embodiment of the present invention: (i) one of said first target DMRs is located in DSCAM and another of said first target DMRs is located in C21orf57 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; and (ii) one of said reference DMRs is located in TBX3 and another of said reference DMRs is located in PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. In particular of such embodiments: (i) one of said first target DMRs comprises SEQ ID No No.: 201 and another of said first target DMRs comprises SEQ ID No.: 219; and (ii) one of said reference DMRs comprises SEQ ID No.: 203 and another of said reference DMRs comprises SEQ ID No.: 220.

[0105] The general arrangement of the DMRs and other regions ("OR") of the differential methylation-based DNA detection used in EXAMPLE 1, is graphically represented by FIGURE 2: (1a) DMR1 is found in exon 2 of RASSF1A and OR1 is located within exon 4 of RASSF1A, with DMR1 located between positions 50,340,672bp and 50,340,784bp and OR1 located between positions 50,331,604bp and 50,331,702bp of the RASS1A genomic sequence (NCBI Reference Sequence: NC_000003.12 Homo sapiens chromosome 3, GRCh38 Primary Assembly), separating DMR1 and OR1 by a distance of 8,969bp. (1b) DMR2 is found in the promoter region of TBX3, with DMR2 located between positions 114,687,095bp and 114,687,189bp and OR2 is located between positions 114,676,384bp and 114,676,454bp of the TBX3 genomic sequence (NCBI Reference Sequence: NC_000012.12 Homo sapiens chromosome 12, GRCh38 Primary Assembly), separating DMR2 and OR2 by a distance of 10,640bp. (2) Methylation in DNA at the two DMRs is detected

using probe-based quantitative PCR using the respective forward (F) and reverse (R) PCR primers and region-specific probes, each probe labelled with the same labels (P*). (3) Total DNA is detected at two ORs using probe-based quantitative PCR using the respective forward (F) and reverse (R) PCR primers and region-specific probes, each probe labelled with the same labels for the ORs that is different to the labels used for the two DMRs (P**). Details of primer and probe sequences and probe labels are set out in TABLE 1.

[0106] The general arrangement of the DMRs and other regions ("OR") used in one non-limiting embodiment of the present invention, is graphically represented by FIGURE 6: (1a) one first target DMR (DMR1) is found in eg DSCAM (Down Syndrome Cell Adhesion Molecule; NCBI Reference Sequence Homo sapiens chromosome 21, GRCh38.p2 Primary Assembly: NC_000021.9 GI:568815577, region 40,010,999 to 40,847,113; SEQ ID No.: 200) such as one located in SEQ ID No.: 51, and (1b) another first target DMR (DMR2) is found in a region no more than 250pb upstream or downstream of eg C21orf57 (YBEY; Chromosome 21: 46,286,337-46,297,751 forward strand, GRCh38:CM000683.2; such gene, including 250bp upstream/downstream flanking regions, SEQ ID No.: 218) such as one located in SEQ ID No.: 185; (1') a first reference DMR (DMR1') is found in in the promoter region of TBX3 (such as one located in SEQ ID No.: 66) and OR1' is located in TBX3 (on human chromosome 12 (2a'), with DMR1' located between positions 114,687,093bp and 114,687,191bp and OR1' located between positions 114,676,384bp and 114,676,454 of the TBX3 genomic sequence (NCBI Reference Sequence: NC_000012.12 Homo sapiens chromosome 12, GRCh38 Primary Assembly), separating DMR1' and OR1' by a distance between about 10,600bp and 10,810bp (4a'); and a second reference DMR (DMR2') is found in PCDHGA1 (such as one located in SEQ ID No.: 102) and OR2' is located in PCDHGA1 (on human chromosome 5 (2b'), with DMR2' located between positions 141,492,593 - 141,492,687 and OR2' located between positions 141,492,918 - 141,493,009 of the PCDHGA1 genomic sequence (NCBI Reference Sequence Homo sapiens chromosome 5, GRCh38.p2 Primary Assembly: NC_000005.10 GI:568815593); separating DMR2' and OR2' by a distance between about 300pb (4b').

[0107] In the embodiment of the present invention graphically represented by FIGURE 6, each of the two first target DMRs are detected with the different detectable label(s) (1a) and (1b); the two reference DMRs (DMR1' and DMR 2') are detected with the same detectable label(s) (1'); and the optional two ORs (OR1' and OR2') are detected with the same detectable label(s) (3').

[0108] Certain embodiments of the present invention, in the context of the methods thereof, comprise or comprise the use of one or more of the foregoing DMRs, ORs, sequences of the primers and/or probes, in particular any of those set forth in TABLE 1, TABLE D or TABLE 5. In certain of such embodiments, a given probe comprises a sequence set forth in TABLE 1, TABLE D or TABLE 5 and any one of the label and quencher pairs (optionally, with a minor binding groove moiety) as set forth in TABLE 1, TABLE D or TABLE 5. In particular, the probe may comprise the combination of the sequence with the label and quencher pair (optionally, with the minor binding groove moiety) as set forth in TABLE 1, TABLE D or TABLE 5 for such probe. Other embodiments of the present invention, particularly in the context of the methods thereof, comprise or comprise the use of the specific combination of two or more (for example, of all) the foregoing DMRs, ORs, sequences of the primers and/or probes, in particular the combination of the primers/probes as set forth in TABLE 1 and/or the combination of the primers/probes as set forth in TABLE 5, including a primer or probe from TABLE D.

[0109] The term "methylation site(s)" will be art-recognised, and has a meaning that encompasses, for example, a CpG motif within a short nucleotide sequence (eg one that is 4, 6, 8, 10 or 12 bp in length) that is, preferably, recognised by a methylation-sensitive restriction enzyme, such as one disclosed elsewhere herein.

[0110] Analogously, and particularly in the context of those embodiments of the present invention that utilise one or more OR, the OR, when located in particular portions and/or genes of the genome, may be located in a promoter, enhancer region or an exon of a gene, or alternatively, located in an intron of such a gene, or in a non-coding region of the genome. In particular embodiments of all aspects of the present invention, such genome and/or gene is a human genome or gene. In particular embodiments, when an OR is used in the present invention and is located in the same portion of the genome and/or gene that features one or more DMRs (preferably, non-overlapping with a DMR used in the invention), then it is located in a portion of the genome and/or gene such as a gene (eg human, and/or in particular when said species of DNA is foetal cfDNA) that is RASSF1A and/or TBX3 and/or PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene, or is selected from the group consisting of: RASSF1A, TBX3, HLCS, ZFY, CDC42EP1, PCDHGA1, MGC15523, SOX14 and SPN and DSCAM and C21orf57 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. When not co-located with a DMR (for example, when a second or multiple other region is used), then such other region may, in certain embodiments, be located in a (eg human) housekeeping gene (such as GAPDH, beta-actin, ALB, APOE or RNASEP). Analogously, and particularly in the context of other embodiments of the present invention that use one or more OR, the OR may be located in particular portions and/or genes of the genome, and may be located in a promoter, enhancer region or an exon of a gene, or alternatively, located in an intron of such a gene, or in a non-coding region of the genome. In particular such embodiments of all aspects of the present invention, such genome and/or gene is a human genome or gene. In

particular embodiments, an OR used in the present invention is located in a (eg human) housekeeping gene (such as GAPDH, beta-actin, ALB, APOE or RNASEP) or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene. Alternatively said OR may be located in the same portion of the genome and/or gene that feature one or more DMRs (such as those located RASSF1A, TBX3, HLCS, ZFY, CDC42EP1, PCDHGA1, MGC15523, SOX14 or SPN or DSCAM or PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene), and preferably does not overlap with a DMR used in the invention.

[0111] In particular embodiments of all aspects of the invention that use one or more OR, said OR comprises a portion of the genome without a methylation site specific for said reagent, and said OR is located in the (eg human) genes RASSF1A or TBX3 (eg SEQ ID NOs: 13 and 14 respectively) or DSCAM (SEQ ID No.: 200) or PCDHGA1 (SEQ ID No.: 217) or C21orf57 (such gene including 250bp upstream/downstream flanking regions SEQ ID No.: 218) or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene, and includes more particular embodiments wherein two or more of said other regions are used in detection step (c) and the first other region is located between about positions 14,220bp and 13,350bp of such RASSF1A and the second other region is located between about positions 12,400bp and 13,000bp of such TBX3. In particular embodiments, an other region is located in RASSF1A between about positions 14,230bp and 14,340bp, 14,230bp and 14,330bp, 14,230bp and 14,320bp, or 14,230bp and 14,310bp of such RASSF1A; and/or is located in TBX3 between about positions 12,400bp and 12,940bp, 12,700bp and 12,850bp or 12,710bp and 12,790bp of such TBX3 (such as SEQ ID No.: 204); and/or is located in DSCAM between about positions 40,841,150bp and 40,841,525bp, 40,841,200bp and 40,841,475bp or 40,841,250bp and 40,841,425bp of DSCAM (with reference to Homo sapiens chromosome 21, GRCh38.p2 Primary Assembly: NC_000021.9 GI:568815577, region), such as SEQ ID No.: 202; and/or is located in PCDHGA1 between about positions 141,492,800 - 141,493,100, 141,492,850 - 141,493,050, 141,492,900 - 141,493,020, 141,492,918 - 141,493,009 of PCDHGA1 (NCBI Reference Sequence Homo sapiens chromosome 5, GRCh38.p2 Primary Assembly: NC_000005.10), such as SEQ ID No.: 221. Alternatively, an OR may be located in an exon such as between about positions 13,790bp and 13,880bp, or 14,490bp and 14,600bp of such RASSF1A, or between about positions 8,040bp and 8,180bp or 6,230bp and 6,350bp of such TBX3; or an OR may be located in an intron such as between about positions 10,500bp and 11,90bp of such RASSF1A, or between about positions 10,000bp and 11,000bp of such TBX3. Alternatively, an OR may be located in DSCAM, such as the sequence TCCGTGTGCTCCACCCTTTGAATTCAGAACGACATAGTGGA TACTCCGTGGGGCTGCT-GGAATCTTCCaTTCcCACT GCCTTATCTT (SEQ ID NO.: 202), which may be amplified and detected with the following probes and primers:

TABLE D: Primer and probes for a DSCAM OR

Chr.	Region	Component	Sequence (5'-3')**	SEQ ID No.*
21	DSCAM Other region	Chr21OR-For	TCCGTGTGCTCCACCCTTTG	208
		Chr21OR-Rev	AAGATAAGGCAGTGGAATGGAAG	209
		Chr21OR-Probe	[Cy5]-CCAGCAGCCCCACGGAGTATCC-[BHQ3]	210
*Only nucleotide sequence listed, without dyes/quenchers				
**The dyes/quenchers used for each probe are shown in "[]" parentheses				

[0112] There is now strong evidence that the level of foetal cfDNA (and/or total cfDNA) present in the circulatory system (eg in plasma) of a pregnant female is a marker of one or more forms of preeclampsia, such as early-onset preeclampsia, mild and/or severe preeclampsia (see Hahn et al 2011, Placenta 32(Supl):S17). The present invention shows particular utility in the efficient, effective, sensitive and/or low-variability detection/quantification of foetal cfDNA present in plasma of pregnant females, and the present invention has particular utility therein. Accordingly, in particular embodiments of the present invention, the individual is a pregnant (eg human) female who is susceptible to suffering or developing a pregnancy-associated medical condition; particularly where said pregnancy-associated medical condition is preeclampsia. As used herein, an individual "susceptible to" a medical condition may alternatively be described as "is suspected to" or to "be considered at risk of being susceptible to" suffering or developing a medical condition; and in certain embodiments, the present invention is used to screen and/or diagnose the individual for susceptibility to, risk of suffering or developing, or suffering from or developing, a medical condition.

[0113] In alternative embodiments, the individual is a pregnant (eg human) female who is susceptible to (or considered at risk of being susceptible to) suffering or developing a pregnancy-associated medical condition selected from the group consisting of: preterm labour, intrauterine growth retardation and vanishing twin. In particular, by comparison to

EXAMPLE1, the sensitivity of the present invention is such that discrepancies between cfDNA levels determined by the method of the invention and that determined by counts of Y-chromosome sequences as determined by massively parallel sequencing approaches, may be useful in identifying one or more cases of a vanishing twin in (mixed-sex) twin pregnancies that previously were believed to be singleton pregnancies, and/or to follow the relative development and health of one or other of such (mixed-sex) twin pregnancies. The present invention may also be utilised in gender determination of twin pregnancies, by consideration of the relative values for foetal cfDNA compared to counts of Y-chromosome sequences determined from cfDNA (eg by using parallel sequencing approaches). In these regards, it should be noted that approaches that use massively-parallel sequencing of random cfDNA in maternal blood typically always count a very low frequency of "Y-chromosome" sequences (such as between about 0.003% and 0.004% of all sequences, or between about 0.0015% and 0.01% or 0.002% and 0.005% of all sequences) in all female pregnancies due to homology of certain Y-chromosome short sequences to other chromosomes. A cut off of "Y-chromosome" sequence counts of about 0.005%, or between about 0.003%, 0.004%, 0.006% or 0.007%, may therefore be employed for female samples.

[0114] In all aspects of the present invention, the reagent that differentially modifies methylated and non-methylated DNA comprises an agent that selectively digests unmethylated over methylated DNA (for example, such agent may digest unmethylated DNA but not methylated DNA). The reagent/agent comprises: at least one methylation sensitive restriction enzyme; and/or an agent selected from the group consisting of: AatII, Acil, AclI, Afel, Agel, Agel-HF, Ascl, AsiSI, Aval, BceAI, BmgBI, BsaAI, BsaHI, BsiEI, BsiWI, BsmBI, BspDI, BsrFI, BssHII, BstBI, BstUI, ClaI, EagI, Faul, FseI, FspI, HaeII, HgaI, HhaI, HinfPII, HpaII, Hpy99I, HpyCH4IV, KasI, MluI, NaeI, NarI, NgoMIV, NotI, NotI-HF, NruI, Nt.BsmAI, Nt.CviPII, PaeR7I, PluTI, PmlI, PvuI, PvuI-HF, RsrII, SacII, Sall, Sall-HF, SfoI, SgrAI, SmaI, SnaBI, TspMI and ZraI. In particular embodiments, said reagent is one selected from the group consisting of: BstUI, HhaI and HpaII.

[0115] In related embodiments, the reagent may comprise two or more of any of the reagents disclosed herein. For example, it may comprise two, three, four, five or more (eg up to seven, eight or ten) methylation sensitive restriction enzymes, including a reagent comprising or essentially consisting of two or three of the methylation sensitive restriction enzymes selected from the group consisting of: BstUI, HhaI and HpaII

[0116] The use of bisulphite or methylation-sensitive restriction enzymes to study differential methylation will be well known to the person of ordinary skill, who may apply teachings of standard texts or adaptation of published methods such as Poon et al (2002), Nygren et al (2010) or Yegnashubramanian ; et al (2006, Nuc Acid Res 34:e19). By way of illustration, the inventors provide examples herein that employ the use of methylation-sensitive restriction enzymes as the reagent that differentially modifies methylated and non-methylated DNA. For further illustration using bisulphite as reagent, it will be apparent to the person of ordinary skill that bisulphite-modified DNA methylation sites may be detected using eg methylation-specific PCR (such as using primers and/or probes that selectively bind to the bisulphite-modified sequences) and/or by the subsequent use of restriction enzymes the recognition site of which is created upon such bisulphite-modification. Methylation-specific PCR ("MSP") is described by Herman et al (US6200756, EP0954608 and related family members); and a further development of MSP using probe-based PCR (known as "MethylLight") is described by Laird et al (US6331393, EP1185695 and related family members).

[0117] In particular embodiments of all aspects of the invention, a quantitative amount of the first or second target species of DNA and/or of the reference species if DNA (and/or or said total DNA) is to be detected and/or determined. Accordingly in such embodiments, one or more (eg each) of said detection and/or determination steps comprises quantitative detection and said detected amount of said species of DNA is expressed as a relative concentration of said species of DNA to the total DNA present in said sample. For example, in certain embodiments of the present invention, in each of said determination steps each of said determined amounts of said first target species of DNA, and of said optional second target species of DNA, is expressed as relative concentrations of said first target species of DNA, and of said optional second target species of DNA, in each case to the total DNA in said sample. In those embodiments where a third (or fourth) target species of DNA is detected, then the present invention also envisions that the amount of such third (or fourth) target species of DNA is expressed as relative concentration.

[0118] If an absolute amount of total DNA is known, then correspondingly an absolute amount (for example, as represented by a concentration such as $\mu\text{g/mL}$ or genome-equivalents such as Eg/mL) of any of species of DNA can be determined from such relative concentration. An absolute amount of total DNA for a sample may be determined, for certain embodiments, by including the further steps of: detecting and/or determining an amount of total DNA in a standard sample of DNA of known amount using the same other regions(s) as used in step (c); and comparing the signal detected from said standard sample of DNA to the signal detected in step (c). Such a standard sample of DNA (of known amount/concentration) is readily available from commercial sources, and especially if prepared and analysed using a dilution series, can readily and efficiently be used to determine (by interpolation/estimation from the standard curve) an absolute amount of total DNA present in the sample. Practically, such standard curve may be prepared and analysed essentially as described for an optional OR (but in a separate set of standard vessels/reactions), preferably in the same run as the detection of the DMRs/OR(s); and may even use the same reaction master-mix. Accordingly, while the "DMR(s)" of the DNA control may be detected for such standard DNA, such a signal is not required to generate a standard curve. Accordingly, if the signal from a such a standard DNA sample is used to compare, the in certain embodiments

where each of said detection and/or determination steps comprises quantitative detection, said detected amount of said species of DNA can be expressed as an absolute amount of said species of DNA in said sample.

[0119] Accordingly, in one embodiment of the method of the present invention, there further comprises the steps of:

- determining an amount of DNA in a standard sample of DNA of known amount in respect of the same DMRs as used in step (c) and/or step (d), and/or in optional step (c)' and/or optional step (f); and
- comparing each of the signals detected from said standard sample of DNA to the respective signals detected in step in step (c) and/or step (d), and/or in optional step (c)' and/or optional step (f).

[0120] In any of the embodiments of the present invention that utilise a standard sample of DNA, there includes certain of such embodiments wherein the standard sample of DNA is a sample of human genomic DNA of known concentration. For example, such human genomic DNA of known concentration may be derived from (such as isolated or purified from) *in-vitro* cultured human cells, such as those known to have either a euploid or an aneuploidy (such as a T21) complement of chromosomes. In alternative embodiments of the present invention, the standard sample of DNA is a synthetic or engineered sample of DNA of known concentration.

[0121] As will be now apparent to the person of ordinary skill, the standard sample of DNA may not be uniformly, appropriately, differentially or otherwise methylated (in particularly if isolated from a human cell-culture),. Accordingly, the present invention includes those embodiments where the standard sample of DNA is not treated with the reagent that differentially modifies methylated and non-methylated DNA (such as, a reagent described herein).

[0122] In one particular embodiment of the present invention, in each of said determination steps each of said determined amounts of said first target species of DNA, and of said optional second target species of DNA, and of the reference species of DNA, is expressed as an absolute amount of said species of DNA in said sample. However, as one aspect of the invention relates to the comparison of (or consideration of relative amounts or ratios) the determined amounts of one or more (such as two) first target species of DNA (ie the chromosome relevant to the chromosomal aneuploidy) to the determined amounts of the reference species of DNA (ie a reference chromosome), such a comparison can be conducted and does not require the consideration of the absolute amounts. By way of explanation, the ratio of "signal-Target-Chromosome"/"signal-Total-DNA" to "signal-Reference-Chromosome"/"signal-Total-DNA" is, because of mathematical cancelation of the "signal-Total-DNA" component, the same as the ratio of "signal-Target-Chromosome" to "signal-Reference-Chromosome". Accordingly, the present invention also includes embodiments where the identification of the presence (or absence) of the chromosomal aneuploidy present in the foetus is conducted in step (e) without knowledge or consideration of the amount of total DNA present in the sample. For example, the determination in step (e) may be made directly from the Ct (or Cp) numbers generated from qPCR as they inversely correlate with initial concentrations (amounts) of the species of DNA being quantitated.

[0123] In certain embodiments of the present invention, a plurality of sets of determinations are made for step (c) and/or step (d), and optional step (c)' and optional step (f), with each set of said determinations made using a different aliquot of DNA of said sample, for example in a different vessel and effectively simultaneously with each other member of the plurality of sets of determinations. Each member of such plurality can be considered a "replicate" of the assay for the same sample (eg, taken from the same pregnant female). Replicates may be efficiently incorporated into the method of the assay by, for example, configuring such a replicate assay to be practiced in a different vessel, such as a different well on the same (or different) microtiter plate such as one used in a qPCR embodiment of the present invention.

[0124] In those embodiments of the present invention that use replicates of the samples, said number of replicates for the sample (plurality of sets of determinations) may be between 2 and about 50 (sets), such as between 2 and about 20, between 2 and about 10 or between about 5 and about 15 (sets); preferably wherein said number of replicates (plurality of sets) of determinations is selected from the group consisting of: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12 (sets of said determinations). For example, for those embodiments of the assay using a microtitre plate, the number of replicates may be represented by the method of the present invention being practiced on aliquots of the same sample conducted in such number of (different) wells of such microtiter plate.

[0125] Inclusion of replicates in the method, and optionally, their analysis, can provide additional information on eg the robustness, the error or other variability in the method and hence one or more of the amounts of DNA determined therefrom. Indeed, in certain embodiments of the invention that use such replicates, the relative amount(s) or ratio(s) determined in step (e), and/or in optional step (e)', is determined from said plurality of sets of determinations made for step (c) and/or step (d), and/or optional step (c)' and/or optional step (f). Such amount may be determined, in such embodiments, by using an average amount of DNA determined for each set (of replicates), such as a mean, median or mode amount of DNA determined for each set (of replicates), preferably a mean amount of DNA determined for each set (of replicates).

[0126] Alternatively, or in addition, the method of the present invention may be practiced (with or without replicates) on a plurality of different samples, each taken from a different pregnant (eg human) female. In such embodiments, especially when in the absence of any molecular tagging methodology, said method is conducted on a plurality of samples

each taken from a different pregnant female, each method is conducted in a separate vessel, and effectively simultaneously with each other member of the plurality of samples. As will be now understood by the person of ordinary skill, the method may be practiced on each sample in a different well of eg a microtiter plate.

[0127] When practicing the method of the present invention on multiple samples taken from different pregnant females, it may be practiced on a number (plurality) of such different samples being between 2 and about 500 (or more than 500) samples, such as between 2 and about 200, between 2 and about 100 or between about 5 and about 150 samples; preferably wherein said plurality of samples is selected from the group consisting of about: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, and 140 samples (or more than 150 samples) each taken from a different pregnant female.

[0128] In particular, the practice of the method of multiple samples taken from multiple pregnant (eg human) females, following the disclosure of the present invention, will now be possible for the person of ordinary skill; for example by the use of microtiter-plate based qPCR system such as the LightCycler from Roche. Such multiple of samples may be located in different wells of the same microtiter plate or may be located on different microtiter plates. The inclusion and analysis of such a plurality of samples, especially if in combination with a plurality of replicates for each sample, may enable the identification of the presence (or absence) of the chromosomal aneuploidy to be made with more certainty or confidence. For example, the relative magnitude of the variation in a given sample's determined amounts (eg the standard deviation) for the various replicates compared to the relative magnitude of the variation on amount determined for all (different) samples, can provide additional information in step (e) (or optional step (e')) for the detection of the chromosomal aneuploidy. For example, a large difference in an amount with little relative variation between replicates will give more confidence that a difference from the other samples exists.

[0129] The inclusion of multiple samples (and/or replicate) in a single run (eg a microtiter plate) may also provide advantages in enabling the comparison of one run to another run; each run containing at least some different samples. Comparison between runs may also be assisted by the use of control or standard samples common to the different runs, which can enable normalisation of measured or calculated amounts between or across runs. However, with a number of different samples (eg about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 130, 140, 150 or more than 150) in each run, amounts determined from one run may be compared to other runs by normalisation methods such those known to the person of ordinary skill or as described elsewhere herein.

[0130] One feature the present invention is that an amount of the chromosome relevant to the chromosomal aneuploidy is estimated or determined using two (or more) first target DMRs located on such chromosome. Use and/or consideration of the amounts determined from each of such two (or more) DMRs can provide additional certainty or confidence to an indication of there being a chromosomal aneuploidy present (or absent) in the foetus. For example, if the determined amount in relation to only one target DMR indicates there being present a chromosomal aneuploidy, then this may be considered a less certain or confident indication than a result in which the amounts determined in relation to two (or more) target DMRs indicates there being present a chromosomal aneuploidy.

[0131] Accordingly, the present invention includes where an amount of the first target species of DNA is determined from the detected amount of methylation at one of the first target DMRs and an amount of the first target species of DNA is determined from the detected amount of methylation at another of the first target DMRs. As will now be apparent, an analogous embodiment may be used for amounts determined from two (or more) of the optional second target DMRs.

[0132] In particular, two or more relative amount(s), preferably ratio(s), are determined in step (e) (and optional step (e')) for two or more amounts of first target species of DNA, each in respect of one or more of said first target DMRs.

[0133] In certain embodiments, a combined analysis of these two (or more) amounts may be considered, such a mean or median of such two or more amounts. Alternatively, as described herein, in particular embodiments two or more (preferably each) of the relative amount(s) or ratios determined in step (e) (and/or optional step (e')) may be independently or separately used to consider, determine or indicate the presence or absence of the chromosomal aneuploidy in the foetus. Such a "two-dimensional" consideration or analysis may be represented or visualised using a scatter plot such as one shown in FIGURE 7; and/or following the use of a large number of different samples and replicates (optionally with normalisation between runs) as one shown in FIGURE 8. As will now be apparent, the use of more than two (such as a third) DMR as a target DMR would provide yet further certainty or confidence on the analysis, and the present invention specifically envisions the use of yet additional target DMRs to, eg, provide a "three-dimensional" analysis to further aid the identification of the chromosomal aneuploidy.

[0134] Each of the amounts of the target chromosome determined from a first (or second) target DMRs may be considered in relation to a threshold or reference distribution of amounts. In particular, certain embodiments of the present invention are when in step (e) two or more (preferably each) of said relative amount(s) or ratio(s) may be compared with threshold(s) and/or reference distribution(s), wherein two or more (preferably each) of said relative amount(s) or ratio(s) higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus.

[0135] As described herein, the threshold(s) and/or reference distribution(s) amount may be identified or provided from external records or information, or may be determined from the use, consideration and/or analysis of multiple samples analysed and amounts determined in accordance with a method of the present invention. Accordingly, in

particular embodiments a threshold and/or reference distribution may be determined from a plurality of samples, each sample taken from a different pregnant female, such as by practicing the method as described above on such samples; optionally wherein a replicates (plurality of sets) of determinations are made for step (c) and/or step (d), and optional step (c)' and optional step (f) for each sample such as by practicing the method as set forth herein.

[0136] When analysing a plurality of samples and/or replicates in a method of the invention, the samples (and/or replicates) may be arranged in different groups or runs (such as in different microtiter plates in a qPCR experiment). In such embodiments, two or more groups of such plurality of samples/replicates may be analysed on a group-by-group basis, such as each group analysed in a separate run, assay or microtiter plate, and the threshold and/or reference distribution is determined by normalisation of the amount(s) or ratio(s) determined from each group of samples. "Normalisation" can be considered as any suitable method that enables the results (eg the amounts) determined from the samples/replicates of one group to be compared/comparable to those of another group. For example, to be able to compare analyses conducted on one run (or microtiter plate) to those of another run (or microtiter plate). Various normalisation approaches will be available to the person of ordinary skill, including wherein said normalisation between groups of samples may be conducted by considering the difference between a sample-specific amount or ratio (such as a mean of sets of determinations made for such sample) and an average (such as a median) of the amount or ratio determined for all samples in the same group as said specific sample; in particular as described in EXAMPLE 4 and represented in FIGURE 8.

[0137] Given such numbers of sample-specific amounts, the presence of the chromosomal aneuploidy in the foetus may be indicated by the sample-specific amount or ratio in respect of such foetus that is an outlier compared with threshold(s) and/or reference distribution(s) of amount(s) or ratio(s) determined in respect of to one or more (preferably to two or to each) of the first target DMRs. In particular of such embodiments, the presence of the chromosomal aneuploidy in the foetus may be indicated by the sample-specific amount or ratio in respect of such foetus being located within a numerical cluster located: (a) outside a cluster of sample-specific amounts or ratios in respect of a plurality of other foeti that are (presumed) euploid; and/or (b) within a cluster of sample-specific amounts or ratios in respect of a plurality of other samples representing foeti having the chromosomal aneuploidy; preferably wherein said plurality of other samples representing foeti is selected from the group consisting of about: 2-10, 12-15, 16-25, 26-30, 32-40, 42-50, 52-75, 78-100, 105-125, 130-150, 155-175, 180-200, 205-250, 255-300, 305-350, 355-400, 405-450, 455-500 and more than 500 other such samples. In related embodiments, the cluster may be defined by a circular or ovular shape visualised in a scatter plot of amounts, wherein the coordinates of such circular or ovular shape may be predefined or calculated from the samples analysed in the method. The amount(s) of the target chromosome compared to the reference chromosome determined at two or more target DMRs may then be considered relative to such coordinates, where a sample having amounts that lie within such a boundary are indicative that the chromosomal aneuploidy is present in the foetus carried by the pregnant female from which that sample was taken. As will be apparent, different boundaries may be set so as to identify aneuploidy samples with various degrees of certainty or confidence. For example, those lying closer to the boundary (a so-called "grey-zone") may be subject to further analysis or testing.

[0138] In other embodiments, a combination of such cluster/circular/oval analysis with a Z-score analysis may be conducted. For example, such a cluster/circular/oval analysis may be conducted to identify euploid samples and a Z-score analysis (such as one described herein) may be conducted to identify aneuploid samples. Greater certainty on the identification of a chromosomal aneuploidy would then be attributed to a sample that shows concordance with such different analysis techniques.

[0139] Prior to or as part of such analysis, one or more of the samples, measures, signals and/or amounts determined from the method are subject to one or more quality control process or steps. For example, the concentration and/or fragmentation pattern of the DNA present in and/or extracted from the sample may be considered prior to step (b) and either the sample not included in the analysis or (in the case of a high concentration) the DNA diluted.

[0140] In particular of such embodiments of the present invention, prior to step (b) the concentration of the DNA may be measured eg with a DNF-474 High Sensitivity NGS Kit (Advances Analytical Technologies, Inc.; Ankeny, USA) using a fragment analyser (Advances Analytical Technologies, Inc.; Ankeny, USA). The DNA-concentration of the first significant peak, that may be located around 80 to 220 bp of size and peaks on average at 165bp, can be determined, which peak can represent the isolated cell-free DNA. In certain embodiments, samples that show a higher DNA-concentration than about 0.3 ng/uL of this peak are diluted to about 0.3 ng/uL. Such dilution may assist to avoid false-positive prediction of samples due to too high amounts of DNA-concentration in comparison to low concentrated samples. Another QC criterion that may applied with such a fragment analysis includes to detect (and eg exclude) those samples with fragmentation of cell-free DNA (represented for example by jagged peaks that occur at sizes less than about 100 bp) and/or contamination by genomic DNA.

[0141] Other possible quality control criteria that may be incorporated into any aspect of the present invention include those where the data from a sample are only included in the analysis if one or more of the following is true:

- The number of replicates present of generating data is greater than about four, five, six, seven, or eight; and/or

- The mean Cp at a qPCR channel for one or more of the DMRs or optional ORs used is less than about 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36; and/or
- The standard deviation of the mean of Cp values at a qPCR channel for one or more of the DMRs or optional ORs used is less than about 0.5, 0.6, 0.7, 0.9, 0.9 or 1.0; and/or
- All replicates have valid values in all of the qPCR channels the DMRs and optional ORs used; and/or
- In a fragment analysis, the sample does not show jagged peaks below about 100bp and/or contamination from genomic DNA; and/or
- The concentration of cfDNA is greater than about 0.05, 0.1, 0.15, 0.2 ng/uL and/or less than about 0.3, 0.4 or 0.5ng/uL

[0142] In those embodiments of the present invention in which total DNA (and hence eg, the foetal fraction of cfDNA) present in the sample is determined, as well as the identification of a chromosomal aneuploidy in the fetus, such method of the present invention may be applied for detecting an increased risk of an pregnant female suffering from or developing a pregnancy-associated medical condition.

[0143] Accordingly, another aspect described herein relates to **a method** for detecting an increased risk of an pregnant female suffering from or developing a pregnancy-associated medical condition; said method comprising the steps:

- (i) conducting a method of the invention as described above;
- (ii) determining at least one amount, such as an absolute or relative amount, of foetal DNA present in the sample; and
- (iii) comparing the amount of foetal DNA determined with a threshold and/or reference distribution,

wherein an increase in, or outlying of, the amount of said foetal of DNA from said threshold and/or reference distribution indicates an increased risk of the pregnant female suffering from or developing said pregnancy-associated medical condition.

[0144] Such a risk may instead or additional be assessed by considering: (i) the fold-increase (eg 1.5, 3, 3.5 or 4-fold increase) of foetal cfDNA (determined for such woman compared to a threshold amount), factoring into the determination that for later-term pregnancies a higher fold-increase in foetal cfDNA may be utilised (Zeybek et al 2013, J Obstet Gynaecol Res 39:632); and/or (ii) into which percentile the amount of cfDNA determined from the woman falls, from consideration of a reference distribution of amounts such as those determined from low-risk women or those which did not suffer from or develop preeclampsia, for example if the foetal cfDNA fraction falls within the 90th percentile of such a distribution, then the woman may be considered to have an increased risk of suffering mild or severe preeclampsia (Jakobsen et al 2013, Transfusion 53:1956). Other relevant factors may be considered in determining a suitable threshold amount. For example, a pregnant woman who is also suffering from breast cancer, may have a higher bias of methylation at RASSF1A present in her plasma due to both factors.

[0145] Analogously, certain embodiments of such aspect of described herein further include the step of: comparing the amount of said species of DNA detected with a threshold amount and/or reference distribution of amounts, wherein an amount of said species of DNA in excess to said threshold (or is not an outlier compared to said population) indicates that a diagnosis for an abnormality in the said species of DNA present in said sample may be performed on, preferably a separate aliquot of DNA of, said sample. For example, if foetal cfDNA fraction is greater than about 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or 0.5% of total cfDNA present in maternal plasma, then there would be sufficient fraction or foetal cfDNA to effectively conduct a subsequent test to investigate one or more characteristics of the foetal cfDNA, for example to investigate the chance or existence of a chromosomal anomaly of mutation comprised within such foetal cfDNA (such as using NIPT based on massively parallel sequencing). In the case of twin pregnancies, the inventors determine that a minimum foetal fraction of cfDNA for NIPT of a twin pregnancy could be considered to be 8%, or about 5%, 6%, 7%, 9% or 10%, and for monochorionic twin pregnancies with concordant genotypes (apart from rare exceptions, Chen et al, 2013, Am J Med Genet A, 161A:1817), a foetal cfDNA fraction of 4%, or about 2%, 3% or 5%, would be sufficient.

[0146] An additional aspect of the present disclosure relates to certain compositions (eg, one that is useful for, used in or generated as part of a method of the present invention). A composition may be a mixture or other combination of various components. Accordingly, the present disclosure also related to **a composition**, said composition comprising two (or more) labelled probes (eg nucleic acid probes, such as those for qPCR), each probe for detecting one or said two or more first target DMRs (such as, one probe for detecting a DMR located in SEQ ID No.: 51 and another probe for detecting a DMR located in SEQ ID No.: 185). In particular such disclosures, two or more or such probes present in the composition are labelled with different detectable label(s). The probes (and labels) may be for any of the first target DMRs disclosed herein, in particular those disclosed in TABLE 1 and/or TABLE 5. The composition may comprise the probes for detecting two, three, four, five or more than five first target DMRs, and labelled with detectable label(s) as set forth elsewhere herein.

[0147] In certain embodiments of the present disclosure, the composition may further comprise one or more of the following additional components:

- two or more pairs of PCR primers, each pair for amplifying one of said two or more first target DMRs, for example for amplifying the first target DMRs for which the probes in the composition are for detecting (such as, one primer pair for amplifying a DMR located in SEQ ID No.: 51 and another primer pair for amplifying a DMR located in SEQ ID No.: 185); and/or
- two or more labelled probes (eg nucleic acid probes, such as those for qPCR), each probe for detecting one of said two or more reference DMRs (such as, one probe for detecting a DMR located in SEQ ID No.: 66 and another probe for detecting a DMR located in SEQ ID No.: 102). In particular such embodiments, two or more or such probes present in the composition are labelled with the same detectable label(s). The probes (and labels) may be for any of the reference DMRs disclosed herein, in particular those disclosed in TABLE 1 and/or TABLE 5; and/or
- two or more pairs of PCR primers, each pair for amplifying one of said two or more reference DMRs, for example for amplifying the reference DMRs for which the probes in the composition are for detecting (such as, one primer pair for amplifying a DMR located in SEQ ID No.: 66 and another primer pair for amplifying a DMR located in SEQ ID No.: 102).

[0148] In alternative or additional embodiments of the present disclosure, the composition may further comprise one or more of the following additional components:

- at least one further pair of primers (preferably two further pairs of primers), each pair for amplifying one of said at least one OR and/or
- at least one further labelled probe (preferably two further labelled probes), each probe for detecting at least one of said ORs.. In particular such embodiments, two or more or such probes present in the composition are labelled with the same detectable label(s). The probes (and labels) may be for any of the ORs disclosed herein, in particular those disclosed in TABLE 1 and/or TABLE D and/or TABLE 5.

[0149] In further alternative or additional embodiments, the composition may further comprise one or more of the additional components for amplifying and/or detecting one, two, three or more second target DMRs, in particular where such second target DMRs are located on a different chromosome relevant to a chromosomal aneuploidy than the first target DMRs.

[0150] A yet further additional aspect disclosed herein relates to a **kit** (for example a kit of separate components; such as a kit of holders or vessels, each holding a different component of the kit), such kit comprising a combination of primers and/or probes wherein said combination is as described in the context of any of the compositions of the present disclosure. For example, a kit described herein may contain (packaged or contained separately or in admixture): (1) two (or more) labelled probes (eg nucleic acid probes, such as those for qPCR), each probe for detecting one of said two or more first target DMRs (one probe for detecting a DMR located in SEQ ID No.: 51 and another probe for detecting a DMR located in SEQ ID No.: 185), where two or more or such probes present in the kit are labelled with different detectable label(s); and (2) additional probes (one probe for detecting a DMR located in SEQ ID No.: 66 and another probe for detecting a DMR located in SEQ ID No.: 102), where such probes are labelled with the same detectable label(s). Additional probes and/or primers (such as those described in the context of the compositions disclosed herein optionally may be present in a kit disclosed herein (such as, one primer pair for amplifying a DMR located in SEQ ID No.: 51 and another primer pair for amplifying a DMR located in SEQ ID No.: 185; and/or one primer pair for amplifying a DMR located in SEQ ID No.: 66 and another primer pair for amplifying a DMR located in SEQ ID No.: 102).

[0151] In particular embodiments disclosed herein, the sequence of such probes and/or primers present in the kit or composition may be modified to detect bisulphite converted DNA in a sequence specific manner; for example such sequence modified primer or probes may be used or useful for MSP or MethylLight embodiments of the inventive method.

[0152] A kit disclosed herein may be used or useful for detecting a chromosomal aneuploidy in a foetus carried by a pregnant female and/or for detecting an increased risk of an pregnant female suffering from or developing a medical condition.

[0153] In further embodiments, a kit disclosed herein may comprise additional components. For example, the kit may additionally comprise: (i) a printed manual or computer readable memory comprising instructions to use said primers and/or probes to practice a method of the invention and/or to produce or use a composition disclosed herein; and/or (ii) one or more other item, component or reagent useful for the practice of a method of the invention and/or the production or use of a composition disclosed herein, including any such item, component or reagent disclosed herein useful for such practice or production. For example, the kit may further comprise the reagent that differently modifies methylated and non-methylated DNA as set above, Taq polymerase, and/or reactions buffers etc.

[0154] By way of further non-limiting example, a kit of the present invention may additionally comprise one or more of the following components:

- means to collect and/or store a tissue sample, such as blood, to be taken from said pregnant female, preferably

wherein said means is a blood collection tube; and/or

- means to extract DNA, preferably cell-free DNA, from the sample to be taken from said pregnant female, preferably wherein said means is a cell-free DNA extraction kit; and/or
- a printed manual or computer readable memory comprising instructions to identify, obtain and/or use one or both of said means in the context of a method of the present invention; and/or
- a computer program product of the present disclosure, or instructions on how to obtain access to or generate results from such a computer program product.

[0155] In particular embodiments of the composition or the kit of the present disclosure, one or more of the primers or probes comprised therein comprises (or consists of) a primer or probe sequence selected from one set forth in TABLE 1, TABLE D and/or TABLE 5 (optionally with the detectable label(s) described therein for such probe). In certain of such embodiments, the composition or the kit comprises the pair or primers and a probe as set forth in TABLE 5 for each of (x) two of the first target DMRs; and (y) two of the reference DMRs, and optionally with (z) one or two of the ORs; in particular the probes for the first target DMRs are differently labelled, the probes for the two reference DMRs are labelled with the same detectable label(s) and, if present, the probes for the two ORs are labelled with the same detectable label(s).

[0156] In particular embodiments, the probes may be labelled with the label/quencher (and optionally a minor groove binding moiety) as set forth in the respective table for such probe.

[0157] In another aspect, the present disclosure relates to a **nucleic acid**, preferably a synthetic, isolated and/or purified nucleic acid, that consists (or comprises) the sequence of any of the primers or probes defined herein, or of the DMRs amplified. In particular embodiments, the primer is covalently linked to a detectable label. In particular embodiments, the sequence of such probes and/or primers may be modified to detect bisulphite converted DNA in a sequence specific manner; for example such sequence modified primer or probes may be useful for MSP or MethylLight embodiments of the inventive method.

[0158] In alternative aspects, a **kit** of the present disclosure may comprise one or more of such nucleic acids together with: (i) a printed manual or computer readable memory comprising instructions to identify, obtain and/or use said nucleic acid in the context of a method; and/or (ii) a computer program product of the present invention, or instructions on how to obtain access to or generate results from such a computer program product. Such a kit may further comprise and additional components, such as one or more other item, component or reagent useful for the practice of a method of the present invention and/or the production or use of the composition of the present disclosure including any such item, component or reagent disclosed herein useful for such practice or production.

[0159] Another further aspect of the present disclosure relates to a **computer program product** comprising: a computer readable medium encoded with a plurality of instructions for controlling a computing system to perform and/or manage an operation for detecting a chromosomal aneuploidy in a foetus carried by a pregnant female from a sample taken from said pregnant female, which sample comprises DNA that originates from cells of a foetus and/or the placenta of a foetus in admixture with differently methylated DNA of maternal origin, the DNA present in said sample being treated with a reagent that differentially modifies methylated and non-methylated DNA such conducted in a method of the invention; said operation comprising the steps of:

- receiving: (i) two (or more) signals, each representing an amount of the first target species of DNA, being the chromosome relevant to the chromosomal aneuploidy, present in said sample such as as set forth in step (c) of the method of the present invention; and (ii) one signal representing an amount of reference species of DNA, being one or more reference chromosomes, present in said sample such as as set forth in step (d) of the method of the present invention;
- determining a classification of whether a chromosomal aneuploidy is likely to be present in the foetus carried by said pregnant female based on at least one (preferable at least two) relative amount(s) of the chromosome relevant to the chromosomal aneuploidy represented by at least one signal (preferably two signals) received in (i) compared to an amount of reference chromosomes represented by the signal received in (ii), wherein at least one (preferably at least two) indicates the presence or absence of the chromosomal aneuploidy in the foetus

[0160] In certain embodiments of the computer program product of the present disclosure, the relative amount can be calculated or otherwise generated from two (or more) of the signals received in (i). For example, a mean, median and/or a Z-Score.

[0161] In other certain embodiments of the computer program product of the present disclosure, the signals representing said amounts of DNA determined in step (c) and step (d) may be generated using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said reference DMRs; optionally using: (x) the same detectable labels(s) for at least two (preferably each) of said reference DMRs; and (y) a different detectable label(s) for at least two (preferably each) of said first target DMRs.

[0162] The operation controlled by the computer program product may conduct such classification by making com-

parison to threshold(s) and/or reference distribution(s), wherein one or more (preferably two or more) of said relative amount(s) or ratio(s) higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the fetus. In particular, said operation computes said threshold(s) and/or reference distribution(s) from a plurality of samples each taken from a different pregnant female by receiving a plurality of signals (i) and (ii) in respect of said plurality of samples; optionally wherein said plurality of samples is analysed in one or more groups of samples by a method the present invention and wherein said method is conducted on each sample in a given group in a separate vessel, and effectively simultaneously with each other member of the plurality of samples in such group.

[0163] In certain embodiments of the computer program product of the present disclosure, said operation further comprises one or more of the steps:

- determining a classification of whether the pregnant female has an increased risk of suffering from or developing a pregnancy-associated medical condition based on comparing said amount of foetal DNA present to a threshold and/or reference distribution, wherein an increase in, or outlying of, the amount of said foetal of DNA from said threshold and/or reference distribution indicates an increased risk of the pregnant female suffering from or developing said pregnancy-associated medical condition; and/or
- a quality-control calculation, process or step, such as one disclosed here.

[0164] In certain embodiments, of the present disclosure, a quality-control calculation, process or step may be practiced prior to any determination step. In this way, those samples/analyses that do not pass the required QC criteria are not included in any further step, process or analysis.

[0165] It is to be understood that application of the teachings of the present invention to a specific problem or environment, and the inclusion of variations of the present invention or additional features thereto (such as further aspects and embodiments), will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein.

[0166] Unless context dictates otherwise, the descriptions and definitions of the features set out above are not limited to any particular aspect or embodiment of the invention and apply equally to all aspects and embodiments which are described.

[0167] Certain aspects and embodiments of the invention will now be illustrated by way of example and with reference to the description, figures and tables set out herein. Such examples of the methods, uses and other aspects of the present invention are representative only, and should not be taken to limit the scope of the present invention to only such representative examples.

EXAMPLE 1 (Comparative): Use of a differential methylation-based DNA detection method in a quantification step prior to MGS-based NIPT in multiple pregnancies, including in cases of vanishing twins

Sample collection, processing and DNA extraction:

[0168] 36 blood samples from women pregnant with multiple gestations (mono-, di- and trichorionic twin and triplet pregnancies) were collected between November 6th 2012 and November 16th 2013, for research & development (R&D) purposes and as part of routine non-invasive prenatal testing (NIPT) laboratory procedure. One blood sample came from a woman pregnant with triplets, the remaining 35 samples came from twin pregnancies. From each pregnant woman carrying a multiple pregnancy two samples each with 7-10ml venous blood were collected using Streck cell-free DNA blood collection tubes (Streck). The blood samples were shipped to the diagnostic laboratory with a maximum delivery time of 4 days. Other blood samples from pregnant females analysed herein were similarly collected.

[0169] Plasma preparation was performed by centrifugation (1600g for 10min at 4°C) and plasma separation followed by a second centrifugation step (16000g for 10min at 4°C). Extraction of total cell-free DNA (cfDNA) was performed with QIAamp Circulating Nucleic Acid Kit (Qiagen) according to the manufacturer protocol using 3.0-4.0ml plasma with a final elution volume of 60ul AVE-buffer (Qiagen).

DNA quantification:

[0170] Foetal cell-free DNA (foetal cfDNA) was detected and quantified in relation to total cell-free DNA (total cfDNA) in order to determine the foetal cfDNA fraction as both a relative concentration and absolute amount using a differential methylation-based DNA detection. From the eluted cell-free DNA, 11ul were digested with the CpG-methylation sensitive enzymes *HhaI* (0.4U/ul), *HpaII* (0.3U/ul) and *BstUI* (0.3 U/ul) in a 22ul reaction using CutSmartTM Buffer (New England Biolabs). The reaction was incubated for 60min at 37°C and 60min at 60°C. 10ul from the digestion reaction was used as template DNA for quantitative probe-based PCR (reactions were conducted in duplicate), described briefly as follows.

[0171] A 25ul PCR reaction using a 2-fold concentrated PCR master mix (QuantiFast Multiplex PCR Kit, Qiagen) was conducted. Primers that span CpG methylation sensitive restriction enzyme sites of the respective region that is differentially methylated between foetal and maternal DNA (as a DMR) were used in combination with FAM-labelled probes for such DMRs, and primers that do not span any restriction enzyme sites, an other region that is not differentially methylated between foetal and maternal DNA (as an OR) are used in combination with VIC-labelled probes for such ORs. The sequences of the primers and labelled probes used in this example are described in TABLE 1, and the thermocycler profiles used for the quantitative probe-based (TaqMan) PCR (LightCycler 480 II Instrument; Roche) are described in TABLE 2. In this example, the probes used to detect the presence of the two DMRs, are each labelled with the same detectable fluorescein amidite (FAM) fluorescent moiety, and each with the same minor binding groove (MGB) non-fluorescent quencher (NFQ) moiety, and the probes used to detect the presence of the two ORs, are each labelled with the same detectable VIC (life Technologies) fluorescent moiety, and each with the same MGBNFQ moiety.

TABLE 1: Quantitative (probe-based) PCR components

Region	Component	Sequence (5'-3')**	SEQ ID No.*	Stock Conc	ul for 1x	Final uM Conc
	Master-mix	N/A		2x	12.5	1x
RASSF1A DMR	DMR1-For	ATT GAG CTG CGG GAG CTG GC	1	100uM	0.35	1.4
	DMR1-Rev	TGC CGT GTG GGG TTG CAC	2	100uM	0.35	1.4
	DMR1-Probe	[FAM]-ACC CGG CTG GAG CGT-[MGBNFQ]	3	100uM	0.035	0.14
RASSF1A Other region	OR1-For	GGT CAT CCA CCA CCA AGA AC	4	100uM	0.35	1.4
	OR1-Rev	TGC CCA AGG ATG CTG TCA AG	5	100uM	0.35	1.4
	OR1-Probe	[VIC]-GGG CCT CAA TGA CTT CAC GT-[MGBNFQ]	6	100uM	0.035	0.14
TBX3 DMR	DMR2-For	GGT GCG AAC TCC TCT TTG TC	7	100uM	0.35	1.4
	DMR2-Rev	TTA ATC ACC CAG CGC ATG GC	8	100uM	0.35	1.4
	DMR2-Probe	[FAM]-CCC TCC CGG TGG GTG ATA AA-[MGBNFQ]	9	100uM	0.035	0.14
TBX3 Other region	OR2-For	TGT TCA CTG GAG GAC TCA TC	10	100uM	0.35	1.4
	OR2-Rev	CAG TCC ATG AGG GTG TTT G	11	100uM	0.35	1.4
	OR2-Probe	[VIC]-GAG GTC CCA TTC TCC TTT-[MGBNFQ]	12	100uM	0.035	0.14
General reagents	DMSO	N/A		100%	0.025	0.625
	MgCl ₂	N/A		50mM	2	1
	DNA sample	N/A				10
	Water					-
	Total					25
*Only nucleotide sequence listed, without dyes/quenchers						
**The dyes/quenchers used for each probe are shown in "[]" parentheses						

TABLE 2: Thermocycler profiles

Step	Temperature	Time	Cycles	Analysis mode
Pre-incubation	95°C	5min	1	None

(continued)

Step	Temperature	Time	Cycles	Analysis mode
Denaturation	95°C	10sec	45	Quantification
Annealing	60°C	10sec		None
Elongation	72°C	8sec		Single
Cooling	40°C			None

[0172] The differential methylation-based DNA detection assay design used in this example is based on two marker DMRs which are described to be hypomethylated in maternal DNA and hypermethylated in foetal DNA (Nygren, et al, 2010: Clin Chem 56, 1627; Chan et al, 2006: Clin Chem 42, 2211; Chiu et al, 2007: Am J Pathol 170, 941), and two other regions (ORs) not differentially methylated between maternal and foetal DNA which are each located between about 20bp and 20kb of their DMR. In particular, the methylation insensitive locus located in RASSF1A is located between 8kb and 9kb (8.97kb) downstream of the methylation sensitive locus located in RASSF1A, and the methylation insensitive locus located in TBX3 is located between 10kb and 11kp (10.64kb) downstream of the methylation sensitive locus located in TBX3. FIGURE 2 depicts the respective arrangements and detection modalities of the two DMRs and the two other regions used in this example.

[0173] Parallel probe-based quantitative PCR reactions were performed (in separate reactions within the same PCR run) using for template a serial dilution of male genomic DNA (Promega) having known concentrations as a standard. The foetal cfDNA fraction was calculated by relative quantification of signals in the FAM channel (DMR; ie detecting foetal cfDNA) versus the VIC channel (ORs; ie detecting total cfDNA), and the absolute total cfDNA amount was calculated by absolute quantification of signals in the VIC channel obtained from the sample compared to the VIC channel obtained from the dilution series of standard DNA of known concentration. Such relative and absolute quantifications were conducted using LightCycler 480 Software release 1.5.0 (Roche).

Maternal plasma DNA sequencing and data analysis to identify foetal aneuploidy:

[0174] DNA sequencing libraries were prepared using NEBNext Ultra™ DNA Library Prep Kit from Illumina. Libraries were prepared according to the manufacturer protocol automated on a Hamilton STARplus robot. Library quality and quantity was measured using a Bioanalyzer instrument (Agilent) and a Qbit Fluorometer (Invitrogen). Based on the library quantification dilutions and equimolar pools of 12 samples per pool were prepared. The pooled samples were sequenced on one lane of an Illumina v3 flow cell on an Illumina HiSeq2000 sequencer. Clonal clusters were generated using TruSeq SR Cluster Kit v3-cBot-HS on a cBot Cluster generation System according to the manufacturer protocol. Bioinformatic analysis to identify foetal chromosomal aneuploidy was carried out as described previously, with z-scores ≥ 3 indicating the presence of a foetal trisomy 21 (Stumm et al 2014, Prenat Diag 34:185). In cases of a positive test result for foetal aneuploidy from this NGS-based method, the result was confirmed by invasive diagnostic methods.

Results:

[0175] Characteristics, % foetal fraction of cfDNA determined by the method of the present example and aneuploidy test results determined by such NGS-based method for the blood samples are given in TABLE 3. There were two positive NGS-based test results indicating foetal trisomy 21. Both were confirmed by karyotyping after amniocentesis; thus, the false positive rate in this NGS-based study was 0%. One blood sample represented monochorionic twins with concordant karyotypes [47,XY,+21] and the other one represented dichorionic twins with discordant karyotypes [47,XY,+21 and 46,XX]. In both samples the foetal fraction was as high as 18.0 and 24.8 %, respectively. All other NGS-based NIPT results were negative for trisomies 21, 18 and 13. There is no evidence for false-negative NIPT results so far in the pregnancies included in this study.

TABLE 3: Characteristics and NIPT results for the collected blood samples

Sample	Chr13 z-score	Chr18 z-score	Chr21 z-score	Foetal DNA fraction (%)	Gestation al age (p.m.)	No. of foetuses, chorionicity amnionicity	NIPT result
LCMPC05	1.3	-1.0	-0.8	16.7	11+5	3, trichorionic, triamniotic	negative

EP 3 168 309 B1

(continued)

5	Sample	Chr13 z-score	Chr18 z-score	Chr21 z-score	Foetal DNA fraction (%)	Gestation al age (p.m.)	No. of foetuses, chorinicity amnioncity	NIPT result
	LCMPC06	-0.4	1.1	8.5	18.0	13+2	2, monochorionic, n.a.	T21 positive
10	LCMPC07	-1.0	0.3	0.9	7.9	19+0	2, dichorionic, diamniotic	negative
	LCMPC08	0.7	1.2	0.0	16.5	18+1	2, dichorionic, diamniotic	negative
15	LCMPC09	0.6	-0.8	0.7	8.9	11+5	2, monochorionic, diamniotic	negative
	LCMPC10	0.3	0.7	-0.7	17.6	20+4	2, dichorionic, diamniotic	negative
20	LCMPC11	-0.9	-0.8	0.7	11.5	23+0	2, dichorionic, diamniotic	negative
	LCMPC12	-0.9	-0.7	-2.0	13.3	11+1	2, monochorionic, diamniotic	negative
25	LCMPC13	1.3	0.1	0.3	21.4	16+0	2, dichorionic, diamniotic	negative
	LCMPC14	0.2	-0.3	0.0	6.8	12+5	2, n.a., n.a.	negative
30	LCMPC15	2.2	0.1	14.7	24.8	16+0	2, dichorionic, diamniotic	T21 positive
	LCMPC16	1.1	1.7	0.5	5.4	12+5	2, n.a., n.a.	negative
	LCMPC17	0.7	1.4	0.5	16.5	14+2	2, n.a., n.a.	negative
	LCMPC18	0.3	2.6	0.0	18.5	18+3	2, n.a., n.a.	negative
35	LCMPC19	-0.2	0.8	0.3	16.6	14+0	2, dichorionic, diamniotic	negative
	LCMPC20	-0.7	-0.9	0.1	13.1	15+4	2, dichorionic, diamniotic	negative
40	LCMPC21	1.0	-0.7	1.2	8.4	9+3	2, dichorionic, diamniotic	negative
	LCMPC22	-1.1	-0.2	0.3	5.6	16+2	2, monochorionic, n.a.	negative
45	LCMPC23	-2.2	2.2	-0.8	20.6	19+5	2, monochorionic, n.a.	negative
	LCMPC24	-1.6	-0.4	-0.5	14.7	22+2	2, monochorionic, diamniotic	negative
50	LCMPC25	-0.8	-0.2	-1.5	12.1	11+5	2, n.a., n.a.	negative
	LCMPC26	-0.4	-0.6	-1.3	7.5	13+0	2, dichorionic, diamniotic	negative
	LCMPC27	0.5	-0.8	-0.4	16.3	12+6	2, n.a., n.a.	negative
55	LCMPC28	-1.2	-0.3	-0.7	19.4	10+1	2, dichorionic, diamniotic	negative

(continued)

Sample	Chr13 z-score	Chr18 z-score	Chr21 z-score	Foetal DNA fraction (%)	Gestation al age (p.m.)	No. of fetuses, chorinicity amnioncity	NIPT result
LCMPC29	-0.8	0.7	-0.4	14.2	13+2	2, monochorionic, n.a.	negative
LCMPC30	0.7	0.3	0.9	14.9	12+2	2, monochorionic, monoamniotic	negative
LCMPC31	-0.2	0.3	-0.9	19.3	19+1	2, dichorionic, diamniotic	negative
LCMPC32	-1.1	2.5	-2.2	11.6	20+0	2, dichorionic, diamniotic	negative
LCMPC33	0.2	2.2	-1.6	8.6	11+0	2, dichorionic, diamniotic	negative
LCMPC34	-1.0	1.2	0.0	15.1	15+4	2, dichorionic, diamniotic	negative
LCMPC35	-0.3	-0.8	-0.3	19.2	12+0	2, dichorionic, diamniotic	negative
LCMPC36	-1.4	-0.5	-0.8	13.9	12+0	2, dichorionic, diamniotic	negative
LCMPC37	1.8	-0.7	0.1	13.8	17+6	2, dichorionic, diamniotic	negative
LCMPC38	-0.1	1.1	-0.7	13.4	13+1	2, dichorionic, diamniotic	negative
LCMPC39	-1.9	0.2	-2.2	15.0	17+0	2, dichorionic, diamniotic	negative
LCMPC40	0.6	-0.4	0.8	16.2	18+3	2, dichorionic, diamniotic	negative

[0176] The reliable detection of foetal aneuploidy in twin pregnancies by prior art NGS-based NIPT is dependent on a sufficiently high amount of foetal cfDNA from each foetus in the maternal blood. Different data and considerations have been published on how the lower limit of foetal cfDNA fraction should be defined to ensure that each twin's contribution is above the detection threshold (Leung et al 2013, Prenat Diag 33:675; Qu et al 2007, Am J Pathol 170:941; Struble et al 2013, Fetal Diagn Ther Dec 7 Epub ahead of print). This is especially important for dichorionic twin pregnancies with discordant karyotypes. In the study described above, supporting information was used for the definition of the minimum foetal cfDNA fraction for twin pregnancies derived from the Y-chromosomal representation, if only one of the two fetuses is male. Using the method of the present example, the total foetal cfDNA fraction can be determined, which reflects the summary of foetal cfDNA derived from both fetuses. Using the Y-chromosomal representation from the next generation sequencing, the foetal cfDNA amount can be determined for male fetuses (as described in Stumm et al 2014). Thus, in the case of mixed foetal gender the contributing amount of each foetus can be determined by subtraction of the amount of foetal cfDNA determined by the Y-chromosomal representation from the foetal cfDNA fraction measured by method of the present example. The foetal cfDNA fractions determined by the method of the present example were compared with the values obtained from Y-chromosomal reads from next generation sequencing for cases with known gender (see FIGURE 3). There is a correlation of the amount of male specific cfDNA (y axis) to the foetal cfDNA fraction measured by method of the present example (x axis). Thus, for twin pregnancies with male/male gender approximately true is: $[y = x]$, for female/male genders it is: $[y = 0.5x]$ and for female/female: $[y = 1]$. The genders of cases with similar values are male/male and in case of differing values with low Y-chromosomal representation the genders are female/female. The intermediate cases, which show about half the percentage of foetal fraction as Y-chromosomal representation, are of mixed gender. The data presented in FIGURE 3 show that it is not only possible to determine the foetal genders using prior art NGS-based NIPT results for twin pregnancies, but also that the measurement

of the amount of foetal fraction of cfDNA determined by the method of the present example is accurate as compared to frequency counting of Y chromosome sequences. On the other hand, these data support the hypothesis that each foetus of a twin pregnancy contributes roughly about half of the total foetal cfDNA fraction. This leads to the conclusion that for twin pregnancies, twice the amount of foetal cfDNA would be required, and thus a recommended minimum foetal fraction of cfDNA for prior art NGS-based NIPT of a twin pregnancy could be considered to be 8%.

[0177] For monochorionic twin pregnancies with concordant genotypes (apart from rare exceptions, Chen et al 2013, Am J Med Genet A 161A:1817), a foetal cfDNA fraction of 4 % would be enough to detect a foetal aneuploidy, just as for single pregnancies. However, for routine laboratory NIPT service one major issue speaks against the implication of such different quality criteria for mono- and dichorionic pregnancies: the determination of chorionicity is dependent on the gestational age and the practical experience of the physician performing the ultrasound examination. The chorionicity is clearly detectable in the first trimester of a multiple pregnancy, but in later stages detection becomes more difficult (Sperling et al 2001, Acta Obstet Gynecol Scand 80:287). Therefore, it is a safer strategy to generally define a minimum foetal cfDNA fraction for twin pregnancies, which is applicable for monochorionic as well as for dichorionic multiple pregnancies.

Identification of vanishing twins:

[0178] In two cases of NIPT aneuploidy testing in which the foetal cfDNA fraction was measured using the method of the present example, identified a trisomy 21 (z-scores 13.5 and 3.4 respectively), but also a striking discrepancy between the total foetal cfDNA fraction measured by the method of the present example and the cf-Foetal-DNA amount measured by Y-chromosome representation were observed.

[0179] For case A, two analyses of blood samples (first and back-up samples) estimated the total foetal cfDNA fraction measured the method of the present example was 20.7% and 24.8%, respectively, whereas the foetal cfDNA according to the Y-chromosomal representation from next generation sequencing was 9.2 % and 9.3 %, respectively. It was speculated, and reported to the physician, that the pregnancy may be a mixed-sex twin pregnancy, who confirmed that a deceased twin had been observed during ultrasound scan at week 10. A further blood sample taken in the third trimester of the pregnancy (38+2) turned out to be negative for trisomy 21 and the foetal cfDNA amount measured by Y-chromosomal representation correlated with the foetal amount measured by QuantYfeX (21.7 % and 21.4), which matched the male gender determined by karyotyping of the living foetus. At birth a foetus papyraceus was found in the placental tissue from which a sufficient amount of cells could be isolated for cell culture and following GTG banding, a trisomy 21 positive, female karyotype was confirmed (47,XX,+21).

[0180] For case B, a slightly increased Y-chromosomal representation was monitored indicating male specific cf-Foetal-DNA of 3.0 % and 2.7 % respectively. As the foetal cfDNA fraction estimates measured by the method of the present example were far above that (13.4 % and 10.0 %) we hypothesized from this discrepancy in the foetal fraction measured, that two foetuses with discordant gender contribute to the foetal fraction and the male foetus being the one affected by trisomy 21. This suggestion was derived from the correlation of Y-chromosome specific foetal cfDNA amount of roughly 3 % with the elevated z-score around the cut-off value of 3.0. Since the examination was clearly requested for a singleton pregnancy, the male specific foetal cfDNA was suspected to stem from a vanishing twin - maybe the carrier of a trisomy 21 - that was either not recognized or not indicated on the consent form for NIPT. Thus, the result was reported to be indecisive for chromosome 21 and the conflicting data was reported to the responsible physician, including a notice regarding the potential vanishing twin, for further clarification via ultrasound. The responsible physician subsequently confirmed that the pregnancy had started as twin and later continued as a singleton pregnancy. The gender of the living and apparently healthy foetus was confirmed to be female and thus, the foetal cfDNA that caused the increased z-score for trisomy 21 can clearly be assigned to a deceased male foetus..

EXAMPLE 2 (Comparative): Improved detection sensitivity of reference chromosomes using two differentially methylated regions using the same detectable moiety/moieties for each differentially methylated region.

[0181] The inventors observe that a complex and multiplex reaction detecting two DMRs using the same detectable moiety/moieties for each of said DMR (as well as two other regions (OR) not differentially methylated) was more sensitive to detect foetal derived reference chromosomes than previous detection reactions that each detected - in separate PCR reactions - a single DMR (as well as a single OR) (FIGURE 4).

[0182] In another differential methylation-based DNA detection method, two DMRs (those found in RASSF1A and TBX3, as described in Example 1), and located on human chromosomes 3 and 12 respectively, were detected (over 4 dilutions) with the same aliquot of DNA and reaction - effectively simultaneously (using quantitative probe-based (TaqMan) PCR) - also with two ORs (those found in RASSF1A and TBX3, as described in Example 1), using the same detectable moiety/moieties for each of said DMR (and a detectable moiety/moieties for said at least one OR that is/are different to the detectable moiety/moieties used for said DMRs). In comparison, detection of foetal derived reference chromosomes

in cfDNA was less sensitive, as shown by detection at higher cycle numbers (Cp), if each DMR (and corresponding OR) was detected independently in separate reactions. The regions/markers, primers/probes and detection methodology was substantially as described in Example 1, except that for the single locus reactions, only the DMR (and corresponding OR) from a given gene (RASSF1A or TBX3) were detected simultaneously in a single reaction.

[0183] In contrast, detection of foetal derived reference chromosomes in cfDNA using a multiplex reaction of the two DMRs using different detectable moieties (eg FAM for the RASSF1A locus and VIC for the TBX3 locus) is determined to be even less sensitive (and further is difficult to detect simultaneously with any OR); without being bound by theory, believed due to the higher complexity of colour compensation, the limited number of separately detectable fluorescent markers and/or the "bleaching" effects from so many fluorescent markers being present in the same reaction.

[0184] Given the exponential nature of quantitative PCR detection, a higher sensitivity of detection (ie lower cycle numbers) would also equate to higher accuracy of quantification, as the correction to standard curves, and interpolation between data points, would be subject to less error than that arising with the amounts of DNA (ie of quantification of the reference chromosomes) correlating to detection at higher cycle numbers.

EXAMPLE 3 (Comparative): Use of a differential methylation-based DNA quantification method for the detection of an increased risk of a pregnant woman suffering from or developing preeclampsia (prophetic example).

[0185] Using a method of a comparative example, pregnant women are assessed for their risk of suffering from or developing preeclampsia as follows. Firstly, a blood sample is collected from the woman for whom such risk to be assessed and total cfDNA extracted from the plasma of such sample substantially in accordance with the procedures described in Example 1. Secondly, using a method substantially as described in Example 1, a relative and/or absolute amount of foetal cfDNA and total cfDNA present in the plasma is determined, where the absolute amount of foetal and/or total cfDNA can be expressed as the amount of genome equivalents ("Eq"). Thirdly, such determined amount of cfDNA and/or total cfDNA is compared to a threshold amount or a reference distribution of amounts, and the women is determined to be at increased risk of suffering from or developing preeclampsia if the amount of foetal cfDNA or total cfDNA exceeds such threshold value and/or is an outlier in such distribution.

[0186] For example, using published threshold values (Papantoniou et al 2013, Prenat Diag 33:682) if the total cfDNA exceeds an amount of about 7,500Eg/mL plasma or if the foetal cfDNA fraction exceeds an amount of about 500Eg/mL plasma, then the woman is determined to have such an increased risk. Such a risk may instead or additional be assessed by considering: (i) the fold-increase (eg 1.5, 3, 3.5 or 4-fold increase) of foetal cfDNA (determined for such woman compared to a threshold amount), factoring into the determination that for later-term pregnancies a higher fold-increase in foetal cfDNA may be utilised (Zeybek et al 2013, J Obstet Gynaecol Res 39:632); and/or (ii) into which percentile the amount of cfDNA determined from the woman falls, from consideration of a reference distribution of amounts determined from low-risk women or women who did not suffer from or develop preeclampsia, for example if the foetal cfDNA fraction falls within the 90th percentile of such a distribution, then the woman is considered to have an increased risk of suffering mild or severe preeclampsia (Jakobsen et al 2013, Transfusion 53:1956).

[0187] In this example, the detection of a risk is conducted using a computer program product that performs the operations represented by FIGURE 5. Operation (A) receives signals (1) and (2) representing, respectively, foetal and total cfDNA are used by the computer program product to determine a parameter (4) that represents the relative and/or absolute amount of foetal (or total) cfDNA present in the plasma of the woman. This operation may optional receive a signal (3) representing an absolute amount of standard DNA. A second operation (B) compares such determined parameter (4) against a threshold amount (5) and/or a reference population of amounts (6) so as to determine and report (7) whether or not - and based on such comparison - the woman is determined to be at increase risk of suffering or developing preeclampsia.

EXAMPLE 4: Use of a method of the invention as a direct and efficient method for NIPT to detect trisomy 21

[0188] The present invention was able to identify cfDNA samples obtained from pregnant females that were carrying a trisomy 21 foetus.

[0189] In a single multiplex PCR reaction of a method of the present invention, the amount of the foetal chromosomal 21 DNA species present in cfDNA samples obtained from 58 pregnant human females was determined at each of two target DMRs, and the amount of foetal chromosomes 5 and 12 (as reference chromosomes) reference DNA species present in the cfDNA sample was determined by use of two reference DMRs (see FIGURE 6), and in respect of each pregnant female. The relative amounts of the foetal chromosomal 21 species (as estimated at each of the two target DMRs) and such foetal chromosomal reference species (as estimated by both of the two reference DMRs) was calculated as a ratio, and each ratio used to plot a scatter plot displaying the sample-specific mean (n=6 replicates) of each of the two ratios for each pregnant human females (FIGURE 7). This scatter plot shows that the three "+" points, each representing a sample obtained from a pregnant female known to carry a foetus with trisomy 21 ("T21"), can be easily

recognised as outliers. This scatter plot also demonstrates one advantageous feature of the inventive method, in that one non-T21 sample was observed as an outlier for one, but not the other, target DMR on chromosome 21. Only by the use of an method and analysis that can analyse at least two of such markers, such as the present inventive method, would have correctly identified this sample as an outlier from the non-T21 population rather than as a T21 sample. For example, if only the target DMR represent by the X axis had been considered, or if the average of both target DMRs had be consider (ie, both target DMRS considered together), then such a sample would have been falsely identified as being a T21 sample.

[0190] The data shown in FIGURE 7 was generated from all 58 samples taken from 58 pregnant human females in a single qPCR run. However, an additional 110 samples taken from 110 pregnant human females were subsequently analysed by the same inventive method in two further qPCR runs, each run analysing samples from 54 and 56 pregnant human females, respectively. By normalisation of the data generated from the three separate runs, the total of all 168 samples could be visualised in a single scatter plot (FIGURE 8). Separation of the cluster of all nine T21 samples from the non-T21 samples is clearly seen; for example as demarcated by the curved boundary shown (dotted). Normalisation between runs was made by subtracting the median value of the applicable ratio across all replicates from a single run as follows from the mean of each ratio for each sample. The resulting normalised values were plotted using a logarithmic scale after conversion to positive values by adding two.

[0191] The robustness of the method of the invention to analysis numerous samples (in this example 168) across multiple runs (in this example 3) is demonstrated (FIGURE 9) by considering the run-independent separation, for each of runs 1 to 3, of the, in total, nine T21 samples (in this figure represented by "o") from the non-T21 samples (in this figure represented by "+"), of the distance of each point from FIGURE 8 (by run) from the centre point of the curved demarcation line (in this example, 0.5,0.55). All nine samples were successfully classified with no false negative or false positive result.

[0192] The amount of a foetal chromosomal 21 DNA species was determined by using two target DMRs: (i) a first such DMR being located in the DSCAM gene (Down Syndrome Cell Adhesion Molecule; NCBI Reference Sequence Homo sapiens chromosome 21, GRCh38.p2 Primary Assembly: NC_000021.9 GI:568815577, region 40010999 to 40847113; SEQ ID No.: 200) located on human chromosome 21q22.2; and (ii) a second DMR located within about 250bp upstream/downstream of C21orf57 (YBEY; Chromosome 21: 46,286,337-46,297,751 forward strand, GRCh38:CM000683.2; such gene including 250bp upstream/downstream flanking regions SEQ ID No.: 218 located on human chromosome 21q22.3.

[0193] The amount of a foetal reference chromosomal DNA species was determined by using two reference DMRs: (i) a first such DMR present in the TBX3 gene (as described above) located on human chromosome 12q24.21; and (ii) a second such DMR present in the PCDHGA1 gene (protocadherin gamma subfamily A, 1; NCBI Reference Sequence Homo sapiens chromosome 5, GRCh38.p2 Primary Assembly: NC_000005.10 GI:, region 141,330,571 to 141,512,981; SEQ ID No.: 217) located on chromosome 5q31.3. Also amplified in each multiple reaction was a first OR located about 10Kb from the DMR in TBX3 (see above) and a second OR located about 300bp from the DMR in PCDHGA1. The sequences of the respective DMRs (and ORs) used are described in TABLE 4.

TABLE 4: Chromosome 21 and chromosome 12 DMRs and ORs

Chr.	Chr. location	Gene	Type	Sequence (5'-3')	SEQ ID No.
21	40841691-40841781	DSCAM	DMR	ATTGGAAGGTCAgCCAAATCAGGCGGGAGCTGCTCCC GG (t) AGCTGCCACCTCCGAGGCGCGGCCACGCCGG GGTCCcTcGCGGCTTTGGA	201
21	46297794 - 46297886	C21orf57	DMR	CGAGCCGTGGCATCGAGAGGGCGTCTGGAGTTCAGGGAA CGCTGGCCCCCGCCCGGAGACCCGCGCAGCGCTCGCC TCTCGCCCTTCAAGG	219
12	114687093-114687191	TBX3	DMR	AAGGTGCGAACTCCTCTTTTGTCTCTGCGTGcCCGGCG CGCCCCCTCCCGGTGGGTGATAAAcCCACTCTGGCG CCGGcCATGCGcTGGgTGATTAATT	203
12	114676384-114676454	TBX3	OR	TGTTcACTGGAGGACTCATCAGAGGTCCCATTCTCCT TTTTGTGTCTTTTCATCAAAACACCCCTCATGGACTG	204
5	141492593 - 141492687	PCDHGA1	DMR	AGCGACTGCCGCTCTAAAGTGCCGGCGGGCAGGACTCTA CGATCCTTGGGCCAGAGGTCCGGATGGTCCCCGGGACTCC GTCTCAAGGGTCGGCGA	220
5	141492918 - 141493009	PCDHGA1	OR	GCGATCTAGGGTCAGAGATTTGGAGGTGACCAAACTATC TGACACTCTAAACAAGTCCCTGTCTCCTCTGGCAGATGGAA AGCTATAGGCTCTG	221
Methylation sensitive sites are underlined and locations of known SNPs are shown by non-capitalisation					

[0194] cfDNA samples from 168 pregnant human females (including 9 of which were known to carry a foetus with T21 were collected, prepared and digested with the CpG-methylation sensitive enzymes *HhaI*, *HpaI* and *BstUI* as described in EXAMPLE 1. A multiplex quantitative probe-based PCR reaction of the four separate loci described in TABLE 4 was conducted on replicates (n = 6) of each such sample as described in EXAMPLE 1, except that the PCR buffer used was

5 PerfeCTa MultiPlex qPCR ToughMix (Quanta BioSciences), using the PCR primers and labelled probes (with quenchers) as set forth in TABLE 5 and splitting the samples across three runs of 58, 54 and 56 samples respectively - each run consisting of such an inventive assay conducted in a single 384-well microtiter plate (Roche) with a LightCycler 480II (Roche).

10

15

20

25

30

35

40

45

50

55

TABLE 5: Primer and probes

Chr.	Region	Component	Sequence (5'-3')**	SEQ ID No.*
21	DSCAM DMR	Chr21DMR1-For	ATTGAAGGTCAGCCCAATCAGG	205
		Chr21DMR1-Rev	TCCAAAGCCGCGAGGGAAC	206
		Chr21DMR1-Probe	[LCyan500]-CGCCTCGGAGGTGGCAGCTC-[BHQ1]	207
21	C210f57 DMR	Chr21DMR2-For	CGAGCCGTGGCATCGA	222
		Chr21DMR2-Rev	CCTGAAGGCGGAGAGG	223
		Chr21DMR2-Probe	[Cy5]-CGTTCCTGAACTCCAGACGC-[BHQ1]	224
12	TBX3 DMR	Chr12DMR-For	AAGTGCGAACTCCTCTTTGTC	211
		Chr12DMR-Rev	AATTAATCACCCAGCGCATGGC	212
		Chr12DMR-Probe	[6FAM]-CCCCCTCCGGTGGTGATAAAACC-[Eclipse]	213
12	TBX3 OR	Chr120R-For	TGTTCACTGGAGGACTCATC	214
		Chr120R-Rev	CAGTCCATGAGGGTGTTTG	215
		Chr120R-Probe	[LCRed610]-AGGTCCCATTCTCCTTTTGTGTCTTTC-[BBQ650]	216
5	PCDHGA1 DMR	Chr5DMR-For	AGCGACTGCCGCTCTAA	225
		Chr5DMR-Rev	TCGCCGACCCCTTGAGAC	226
		Chr5DMR-Probe	[6FAM] - TCTGGCCCAAGGATCGTAGAGTCC - [Eclipse]	227
5	PCDHGA1 OR	Chr50R-For	GCGATCTAGGGTCAGAGATTG	228
		Chr50R-Rev	CAGAGCCTATAGCTTTCCATCTG	229
		Chr50R-Probe	[LCRed610]-ACACTCTAACAAAGTCCTGTCTCCTCTGG - [BBQ650]	230

*Only nucleotide sequence listed, without dyes/quenchers

**The dyes/quenchers used for each probe are shown in "[]" parentheses

[0195] The format of such assay is generally as depicted in FIGURE 6, where the DMR1 of such figure is located in the DSCAM gene of human chromosome 21, DMR2 of such figure is located within about 250bp upstream/downstream of the C21orf57 gene of human chromosome 21, the DMR1' of such figure is located in the TBX3 gene of human chromosome 12 and the DMR2' of such figure is located in the PCDHGA1 gene of human chromosome 5. For the purposes of calculating a relative amount of, or ratio between, the target DMRs on chromosome 21 and the reference DMRs on chromosomes 5 and 12 the ORs are not required as if used to calculate a foetal fraction of the respective chromosome, such value would be cancelled in the mathematical formulae. Hence, the amount of total DNA that may be estimated from use of the ORs is not required to analysis the T21-status of the foetus, but may still be utilised to calculate the fraction of foetal cfDNA from the total DNA; for example for the purposes as described in EXAMPLES 6 and/or 7, and/or as a possible or additional quality control measure (eg EXAMPLE 6), or for the purposes of other diagnosis such as described in EXAMPLE 7. As will be observed from TABLE 5, the probe for the chromosome 21 target DMR located in the DSCAM is labelled differently from the probe for the chromosome 21 target DMR located within about 250bp upstream/downstream of C21orf57; also the reference chromosome 12 reference DMR located in TBX3 and that on chromosome 5 located in PCDHGA1 are labelled with the same label (different from the target DMRs). Finally, if required for an estimation of total DNA, the two ORs are both labelled with the same label, which too is different from the other labels used in the multiplex assay. Overall, 4 labels are used, which enables the amount of chromosome 21 to be independently estimated for each target DMR and the amount of reference chromosomes to be more sensitively detected (see EXAMPLE 2), for example by reference to the respective signals for each DMR (or OR) from the sample compared to signals for the respective DMR (or OR) obtained from the dilution series of standard DNA of known concentration (as described in EXAMPLE 1) as may be provided in each qPCR plate (run) as the test samples. Such relative and absolute quantifications were conducted using LightCycler 480 Software release 1.5.0 (Roche). The mean [n=6 replicates] sample-specific ratio of each %foetal cfDNA of chromosome 21 target DNA + species and the %foetal cfDNA of both reference chromosome reference DNA species was calculated for each sample, and for each plate the overall median of all replicates was calculated. Analysis was as described above.

EXAMPLE 5: Iterative z-score analysis to detect trisomy 21 in NIPT without reference to known internal euploid standards

[0196] By the application of an iterative z-score approach, the inventors were able to identify all known Trisomy 21 samples from the test samples without reference to the known euploid samples in the estimation of mean and standard deviations in the z-score analysis.

[0197] The data from each run (qPCR plate) of samples analysed in EXAMPLE 4 was re-analysed as follows. Firstly, the mean ratio of each chromosome 21 target DMR to the reference chromosomes DMRs of each replicate of the samples (n=6) was calculated, and for all samples present in such run (plate) an overall mean and standard deviation for each ratio was calculated without reference to whether a sample was known to be euploid or trisomy. Secondly, based on such run-specific means and standard deviation, (run specific) separate z-scores for each ratio associated with each target DMR for each sample present in the run were calculated. Thirdly, for each DMR, such ratios were checked for those equal or exceeding 1.95, and any such samples represented by such outlying z-scores were, independently for each of the two target DMRs, excluded from the respective reference set samples used for any subsequent determination of z-score calculation factors (run specific mean ratio and standard deviation. A second mean and standard deviation for each ratio was calculated, independently for each target DMR, on the data in respect of the remaining samples in the reference data set and used to conduct a second z-score analysis of each ratio for all samples in such set. Fourthly, the resulting z-scores (ie, those calculated after the first iterative elimination) are checked for outliers, independently for each target DMR, as in the third step, as above and those samples associated with a z-score equal or greater to 1.95 are excluded from the reference data set used to calculate z-scores in respect of each DMR. A third mean and standard deviation was then calculated, independently for each target DMR, for each ratio on the data in respect of the remaining samples in the respective reference data set and used to conduct a third z-score analysis for each ratio of all samples in such set. This iterative process was repeated for 20 times and no sample which was once excluded from the respective DMR reference set was re-included in a reference data set, even if the respective sample's z-score would again change to below 1.95 after one iterative elimination. The resulting sample z-scores for each ratio and sample are shown in FIGURE 10, which displays complete separation of the euploid and T21 samples (represented by z-scores greater than about 3.0 for each of the two ratios; and/or by the mean of the two z-scores being greater than about 3.0; and/or by a cluster or distance approach analogous to that described in EXAMPLE 4/FIGURE 9). Diploid outliers which obtain values above mean z-scores of 3 were excluded by defining a rule: discordant results (Marker 1 and 2) are controlled for the extent of difference. If the difference was above 4.0 they were non positive/non reportable.

EXAMPLE 6: Use of a method of the invention to estimate the foetal cfDNA fraction

[0198] The % fraction of foetal DNA in the total cfDNA isolated from the pregnant female was estimated by calculating

the ratio of the amount of DNA measured using the two (commonly-labelled) reference DMRs present on the two reference chromosomes and the two (commonly-labelled) ORs; each within the same gene as their respective DMR. Information on the foetal fraction of cfDNA can be used for quality control purposes, including to determine if sufficient foetal DNA is present in the sample to permit subsequent analysis such as NGS-based analysis as described in EXAMPLE 1.

EXAMPLE 7: Use of a method of the invention to detect an increased risk of a pregnant woman suffering from or developing preeclampsia (prophetic example)

[0199] In addition to determining whether a fetus carried by a pregnant female is aneuploid, the format of the inventive assay described in EXAMPLE 4 (ie, one carrying at least one OR to measure total DNA) is used to also determine if the pregnant female is at risk of suffering from or developing preeclampsia as follows.

[0200] Using a method substantially as described in EXAMPLE 4, a relative and/or absolute amount of foetal cfDNA and total cfDNA present in the plasma is determined, where the absolute amount of foetal and/or total cfDNA can be expressed as the amount of genome equivalents ("Eq"). Thirdly, such determined amount of cfDNA and/or total cfDNA is compared to a threshold amount or a reference distribution of amounts, and the woman is determined to be at increased risk of suffering from or developing preeclampsia if the amount of foetal cfDNA or total cfDNA exceeds such threshold value and/or is an outlier in such distribution. Published values, and other ways of assessing such a risk include those described in EXAMPLE 3. Such a method of the present invention may be used to generate signals to utilise the computer program product described in EXAMPLE 3 (and represented by FIGURE 5) to detect such risk.

EXAMPLE 8: Preparation and distribution of a kit of the present disclosure (prophetic example)

[0201] A kit disclosed herein is prepared by forming a buffered solution for each of the primers and probes listed in TABLE 5. An aliquot of each buffered solution (eg, 200ul) is placed in separate eppendorf tubes, and such tubes packaged in a cardboard container.

[0202] In an alternative kit, a single buffered solution is prepared that comprises an admixture of each of the primers and probes listed in TABLE 5.

[0203] In a further alternative kit, the cardboard container further includes an aliquot of the probes used to detect the two first target DRMs set forth in TABLE 5.

[0204] The cardboard container of the kit further contains a printed instruction manual, instructing the user to use the components therein (together with other components) to practice the method of the present invention to seek to detect a chromosomal aneuploidy in a fetus.

EXAMPLE 9: Trial with collected samples (prophetic example)

[0205] A number of samples collected from numerous pregnant females known to carry an aneuploid or euploid fetus are analysed by a method of the present invention (such as by using an assay as described in EXAMPLE 4), but in a blinded fashion (ie, the investigators do not know which sample is from which class of pregnant female) to seek to (eg correctly) identify those samples believed to carry foeti having a chromosomal aneuploidy (eg trisomy 21). After the analysis has been conducted, the samples are un-blinded (ie, whether they did indeed carry a trisomy fetus, such as determined by NGS and/or other tests), and such information is compared to generate estimates or measures of sensitivity and specificity for the method of the present invention.

Claims

1. A method for detecting a chromosomal aneuploidy in a fetus carried by a pregnant female, said method comprising the steps:

- (a) providing a sample that has been taken from said pregnant female, which sample comprises DNA that originates from cells of a fetus and/or the placenta of a fetus in admixture with differently methylated DNA of maternal origin;
- (b) treating the DNA present in said sample with a reagent that differentially modifies methylated and non-methylated DNA, wherein said reagent is an agent that selectively digests unmethylated over methylated DNA and said agent comprises a methylation sensitive restriction enzyme;
- (c) determining an amount of a first target species of DNA, being the chromosome relevant to the chromosomal aneuploidy, in said sample by detecting in said sample the presence of methylation at two or more first target differentially methylated regions (DMRs) located on said chromosome, said first target DMRs differently meth-

ylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, wherein said DMRs are hypermethylated in foetal DNA and hypomethylated in maternal DNA, the modification of DNA of the first target DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of methylated DNA at one or more of said first target DMRs indicates said amount of first target species of DNA in said sample, wherein an amount of the first target species of DNA is determined from the detected amount of methylation at one of the first target DMRs and an amount of the first target species of DNA is determined from the detected amount of methylation at another of the first target DMRs;

(d) determining an amount of reference species of DNA, being one or more reference chromosomes, in said sample by detecting in said sample the presence of methylation at two or more reference DMRs located on said reference chromosome(s), said reference DMRs differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, wherein said DMRs are hypermethylated in foetal DNA and hypomethylated in maternal DNA, the modification of DNA of such reference DMRs by said reagent is sensitive to methylation of DNA, wherein a detected amount of methylated DNA at one or more of said reference DMRs indicates said amount of reference species of DNA in said sample; and

(e) determining relative amount(s), preferably ratio(s), of an amount determined from step (c) and an amount determined from step (d), wherein two or more such relative amounts are determined for two or more amounts of first target species of DNA, each in respect of one or more of said first target DMRs, and wherein two or more of said relative amounts indicate the presence or absence of the chromosomal aneuploidy in the foetus,

wherein, said detections in step (c) and step (d) are made using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said reference DMRs, and using: (x) the same detectable label(s) for at least two of said reference DMRs; and (y) a different detectable label(s) for at least two of said first target DMRs; and:

wherein in step (e) said two or more relative amounts or ratios are compared with threshold(s) and/or reference distribution(s), wherein said two or more relative amounts or ratios higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus.

2. The method of claim 1, wherein said detections in step (c) and step (d) are made using: (x) the same detectable label(s) for each of said at least two reference DMRs; and (y) a different detectable label(s) for each of said at least two first target DMRs.

3. The method of claims 1 or 2, wherein two or more of said reference DMRs are located on different reference chromosomes.

4. The method of any one of claims 1 to 3 further comprising the step:

(f) determining an amount of total DNA in said sample by detecting at least one other region (OR) that is not differently methylated between DNA that originates from cells of a foetus and/or the placenta of a foetus and DNA of maternal origin, the modification of which OR(s) by said reagent is insensitive to methylation of DNA,

wherein, said detections in step (c) and step (d) and step (f) are made using the same aliquot of DNA of said sample, and in the same vessel, and effectively simultaneously for said first target DMRs and said reference DMRs and said other region(s), and using: (x) the same detectable label(s) for at least two of said reference DMRs; and (y) a different detectable label(s) for at least two of said first target DMRs and for at least one of said OR(s); *optionally*:

wherein said OR(s) is(are) located on one or more reference chromosome(s), preferably at least one of said ORs being located on the same reference chromosome(s) as at least one of said reference DMRs.

5. The method of any one of claims 1 to 4, where in step (e) each of said relative amount(s) or ratio(s) is compared with threshold(s) and/or reference distribution(s), wherein each of said relative amount(s) or ratio(s) higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus.

6. The method of claims 4 or 5, wherein (A) said detection in step (f) comprises using at least two of said ORs; *preferably wherein*, the number of said ORs is the same as the number of reference DMRs used in step (d); *more preferably wherein*, one of said ORs is located on the same chromosome as, such as between about 20bp and about 20kb upstream or downstream of, a reference DMR used in step (d) and each other of the said ORs is located on the same chromosome as, such as between about 20bp and about 20kb upstream or downstream of, another of said reference DMRs; and

(B) said detection in step (f) is made using: (x) the same detectable label(s) for each of said ORs.

7. The method of any one of claims 1 to 6, wherein said detection step (c) and step (d), and optional step (f), are made using the same aliquot of DNA of said sample, and in the same reaction/detection vessel, and effectively simultaneously with each other, and by multiplex real-time quantitative probe-based PCR using at least one labelled probe specific for each of the said DMRs and optional OR(s).

8. The method of any one of claims 1 to 7, wherein the pregnant female is human and the chromosome relevant to the chromosomal aneuploidy is a human chromosome selected from the lists consisting of: chromosome 21, chromosome 18 and chromosome 13.

9. The method of any one of claims 1 to 8, wherein said species of DNA that originate from cells of a foetus and/or the placenta of a foetus is circulating cell-free DNA and said sample is a blood fraction such as plasma or serum.

10. The method of any one of items 1 to 9, wherein said reagent that differentially modifies methylated and non-methylated DNA comprises

an agent selected from the group consisting of: AatII, Acil, AclI, Afel, AgeI, AgeI-HF, AscI, AsiSI, Aval, BceAI, BmgBI, BsaAI, BsaHI, BsiEI, BsiWI, BsmBI, BspDI, BsrFI, BssHII, BstBI, BstUI, ClaI, EagI, FaeI, FseI, FspI, HaeII, HgaI, HhaI, HinfI, HpaII, Hpy99I, HpyCH4IV, KasI, MluI, NaeI, NarI, NgoMIV, NotI, NotI-HF, NruI, Nt.BsmAI, Nt.CviPII, PaeR7I, PfuTI, PmlI, PvuI, PvuI-HF, RsrII, SacII, SalI, SalI-HF, SfiI, SgrAI, SmaI, SnaBI, TspMI and ZraI.

11. The method of any one of claim 1 to 10, wherein one reference DMR is located on human chromosome 5 and another reference DMR is located on human chromosome 12.

12. The method of any one of claims 1 to 11, wherein said first target DMRs are each located in a region and/or gene independently selected from:

- one selected from the list consisting of: SEQ ID NOs: 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198 and 199; or
- the list consisting of: AIRE, SIM2, and ERG or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; or
- the list consisting of: PDE9A, PPP1 R2P2, CBR1, DSCAM, C21orf29 and HLCS or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; or
- C21orf57 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; or
- the list consisting of: SEQ ID NOs: 47, 48, 49, 50, 51, 52, 53, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186 and 187.

13. The method of claim 12, wherein said first target DMRs are each located in a region and/or gene independently selected from the list consisting of: SEQ ID NOs: 47, 48, 49, 50, 51, 52, 53, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186 and 187.

14. The method of claim 13, wherein said first target DMRs are each located in a region and/or gene independently selected from the list consisting of: SEQ ID NOs: 51, 184, 185 and 186.

15. The method of claim 14, wherein one of said first target DMRs is located in a region and/or gene being SEQ ID NO.: 51 and another of said first target DMRs is located in a region and/or gene being SEQ ID NO.: 185.

16. The method of any one of claims 1 to 15, wherein said first target DMRs are each located in a region and/or gene independently selected from:

- VAPA-APCDDI or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; or
- maspin or a DNA sequence of no more than 10kbp, 5kbp, 1 kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene
- the list consisting of: SEQ ID NOs: 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 and 46.

17. The method of any one of claims 1 to 16, wherein said first target DMRs are each located in a region and/or gene independently selected from:

- the list consisting of: SEQ ID NOs: 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33 and 34.

18. The method of any one of claims 1 to 17, wherein said reference DMRs are each located in a region and/or gene independently selected from:

- the list consisting of: RASSF1A, TBX3, ZFY, CDC42EP1, PCDHGA1, SOX14 and SPN or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; or
- the list consisting of: SEQ ID NOs: 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146 and 147.

19. The method of claim 18, wherein said reference DMRs are each located in a region and/or gene independently selected from the list consisting of: SEQ ID NOs: 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146 and 147.

20. The method of claim 19, wherein said reference DMRs are each located in a region and/or gene independently selected from the list consisting of SEQ ID NOs: 66, 102 and 126.

21. The method of claim 20, wherein one of said reference DMRs is located in a region and/or gene being SEQ ID NO.: 66 and another of said reference DMRs is located in a region and/or gene being SEQ ID NO.: 102.

22. The method of any one of claims 1 to 21, wherein at least one of said first target DMRs is located in a portion of the genome and/or a gene that is DSCAM and/or C21orf57, or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene.

23. The method of any one of claims 1 to 21, wherein:

- (A) one of said first target DMRs is located in DSCAM and another of said first target DMRs is located in C21orf57 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene; and
- (B) one of said reference DMRs is located in TBX3 and another of said reference DMRs is located in PCDHGA1 or a DNA sequence of no more than 10kbp, 5kbp, 1kbp, 500bp, 250bp, 150bp, 100bp or 50bp upstream and/or downstream from such region and/or gene.

24. The method of any one of claims 1 to 21, wherein: (i) one of said first target DMRs comprises SEQ ID NO.: 201 and another of said first target DMRs comprises SEQ ID NO.: 219; and (ii) one of said reference DMRs comprises SEQ ID NO.: 203 and another of said reference DMRs comprises SEQ ID NO.: 220.

25. The method of any one of claims 1 to 24,

- (A) wherein a plurality of sets of determinations are made for step (c) and step (d), and optional step (f), with each set of said determinations made using a different aliquot of DNA of said sample, in a different vessel and effectively simultaneously with each other member of the plurality of sets of determinations; and
- (B) wherein the relative amount(s) or ratio(s) determined in step (e) is determined from said plurality of sets of determinations made for step (c) and step (d), and optional step (f), preferably by using an average amount of DNA determined for each set, such as a mean, median or mode amount of DNA determined for each set, preferably a mean amount of DNA determined for each set.

26. The method of any one of claims 1 to 25, wherein in step (e) said relative amount(s) or ratio(s) are compared with

threshold(s) and/or reference distribution(s), wherein two or more of said relative amount(s) or ratio(s) higher or lower than said threshold(s) and/or reference distribution(s) indicates the presence of the chromosomal aneuploidy in the foetus; and wherein a threshold and/or reference distribution is determined from a plurality of samples, each sample taken from a different pregnant female, such as by practicing the method as set forth in any one of claims 1 to 22 on such samples, wherein said method is conducted on a plurality of samples each taken from a different pregnant female, each method conducted in a separate vessel, and effectively simultaneously with each other member of the plurality of samples

27. The method of any one of claims 1 to 26, wherein a given DMR is amplified by PCR primers that:

- overlap two or three restriction sites for the methylation sensitive restriction enzyme(s), each of which comprises a methylation site; and/or
- are designed to flank one, two, three or more restriction sites for the methylation sensitive restriction enzyme(s), each of which comprises a methylation site.

28. The method of any one of claims 1 to 27, wherein said methylation sensitive restriction enzyme(s):

- is one selected from the group consisting of: BstUI, Hha I and HpaII; or
- comprise two or three of the methylation sensitive restriction enzymes selected from the group consisting of: BstUI, Hha I and HpaII.

29. The method of any one of claims 1 to 27 or 28, wherein the detectable labels are fluorescent labels.

Patentansprüche

1. Verfahren zum Nachweis einer chromosomalen Aneuploidie in einem Fötus, der von einem schwangeren Weibchen ausgetragen wird, wobei das Verfahren die folgenden Schritte umfasst:

- (a) Bereitstellen einer Probe, die einem schwangeren Weibchen entnommen wurde, wobei die Probe DNA umfasst, die aus Zellen eines Fötus und/oder der Plazenta eines Fötus im Gemisch mit differentiell methylierter DNA mütterlicher Herkunft stammt;
- (b) Behandeln der in der Probe vorhandenen DNA mit einem Reagens, das methylierte und nicht-methylierte DNA differentiell modifiziert, wobei das Reagens ein Agens ist, das selektiv unmethylierte gegenüber methylierter DNA verdaut, und wobei das Agens ein methylierungsempfindliches Restriktionsenzym umfasst;
- (c) Bestimmen einer Menge einer ersten DNA-Zielspezies, die das für die chromosomale Aneuploidie relevante Chromosom darstellt, in der Probe, indem in dieser Probe das Vorhandensein einer Methylierung an zwei oder mehr ersten differentiell methylierten Regionen (DMRs) nachgewiesen wird, die sich auf dem Chromosom befinden, wobei die ersten Ziel-DMRs zwischen DNA, die aus Zellen eines Fötus und/oder der Plazenta eines Fötus stammt und DNA mütterlichen Ursprungs unterschiedlich methyliert sind, wobei die DMRs in fötaler DNA hypermethyliert und in mütterlicher DNA hypomethyliert sind, wobei die Modifikation der DNA der ersten Ziel-DMRs durch das Reagens empfindlich gegenüber Methylierung von DNA ist, wobei eine nachgewiesene Menge an methylierter DNA an einem oder mehreren der ersten Ziel-DMRs die Menge der ersten DNA-Zielspezies in der Probe angibt, wobei eine Menge der ersten DNA-Zielspezies aus der nachgewiesenen Methylierungsmenge an einer der ersten Ziel-DMRs bestimmt wird, und eine Menge der ersten DNA-Zielspezies aus der nachgewiesenen Methylierungsmenge an einer weiteren der ersten Ziel-DMRs bestimmt wird;
- (d) Bestimmen einer Menge von DNA-Referenzspezies, die ein oder mehrere Referenzchromosomen darstellt, in der Probe durch Nachweisen des Vorhandenseins einer Methylierung an zwei oder mehr Referenz-DMRs, die sich auf dem/den Referenzchromosom(en) befinden, wobei die Referenz-DMRs zwischen DNA, die aus Zellen eines Fötus und/oder der Plazenta eines Fötus stammt und DNA mütterlichen Ursprungs unterschiedlich methyliert sind, wobei die DMRs in fötaler DNA hypermethyliert und in mütterlicher DNA hypomethyliert sind, wobei die Modifikation der DNA solcher Referenz-DMRs durch das Reagenz empfindlich gegenüber Methylierung von DNA ist, wobei eine nachgewiesene Menge an methylierter DNA an einer oder mehreren der Referenz-DMRs die Menge an DNA-Referenzspezies in der Probe angibt; und
- (e) Bestimmen der relativen Menge(n), vorzugsweise Verhältnis(se) einer aus Schritt (c) bestimmten Menge und einer aus Schritt (d) bestimmten Menge, wobei zwei oder mehrere solcher relativer Mengen für zwei oder mehrere Mengen der ersten DNA-Zielspezies bestimmt werden, jeweils in Bezug auf eine oder mehrere der ersten Ziel-DMRs, und wobei zwei oder mehrere der relativen Mengen das Vorhandensein oder Nichtvorhandensein

densein der chromosomalen Aneuploidie beim Fötus anzeigen,

wobei die Nachweise in Schritt (c) und Schritt (d) unter Verwendung des gleichen Aliquots von DNA der Probe durchgeführt werden und in demselben Gefäß, und effektiv gleichzeitig für die ersten Ziel-DMRs und die Referenz-DMRs und unter Verwendung von: (x) derselben/denselben nachweisbaren Markierung(en) für mindestens zwei der Referenz-DMRs; und (y) (einer) unterschiedlichen nachweisbaren Markierung(en) für mindestens zwei der ersten Ziel-DMRs; und:

wobei in Schritt (e) die zwei oder mehreren relativen Mengen oder Verhältnisse mit (einem) Schwellenwert(en) und/oder (einer) Referenzverteilung(en) verglichen werden, wobei zwei oder mehrere relative Mengen oder Verhältnisse, die höher oder niedriger als der/die Schwellenwert(e) und/oder Referenzverteilung(en) sind, das Vorhandensein der chromosomalen Aneuploidie beim Fötus anzeigt.

2. Verfahren nach Anspruch 1, wobei die Nachweise in Schritt (c) und Schritt (d) durchgeführt werden unter Verwendung von: (x) derselben/denselben nachweisbaren Markierung(en) für jede der mindestens zwei Referenz-DMRs; und (y) (einer) unterschiedlichen nachweisbaren Markierung(en) für jede der mindestens zwei ersten Ziel-DMRs.

3. Verfahren nach Anspruch 1 oder 2, wobei sich zwei oder mehrere der Referenz-DMRs auf verschiedenen Referenzchromosomen befinden.

4. Verfahren nach einem der Ansprüche 1 bis 3, das ferner den folgenden Schritt umfasst:

(f) Bestimmen einer Menge an Gesamt-DNA in der Probe durch Nachweisen von mindestens einer anderen Region (OR), die zwischen DNA, die aus Zellen eines Fötus und/oder der Plazenta eines Fötus stammt, und DNA mütterlichen Ursprungs nicht differentiell methyliert ist, wobei die Modifikation dieser OR(s) durch das Reagens unempfindlich gegenüber Methylierung von DNA ist,

wobei die Nachweise in Schritt (c) und Schritt (d) und Schritt (f) unter Verwendung des gleichen DNA-Aliquots der Probe und in demselben Gefäß durchgeführt werden, und effektiv gleichzeitig für die ersten Ziel-DMRs und die Referenz-DMRs und die andere(n) Region(en), und unter Verwendung von: (x) derselben nachweisbaren Markierung(en) für mindestens zwei der Referenz-DMRs; und (y) (einer) unterschiedlichen nachweisbaren Markierung(en) für mindestens zwei der ersten Ziel-DMRs und für mindestens eine der OR(s); *wahlweise*:

wobei sich die OR(s) auf einem oder mehreren Referenzchromosom(en) befindet (befinden), wobei sich vorzugsweise mindestens eine der ORs auf demselben/denselben Referenzchromosom(en) befindet wie mindestens eine der Referenz-DMRs.

5. Verfahren nach einem der Ansprüche 1 bis 4, wobei in Schritt (e) jede der relativen Menge(n) oder Verhältnis(se) mit (einem) Schwellenwert(en) und/oder (einer) Referenzverteilung(en) verglichen wird, wobei jeder der relativen Mengen oder Verhältnisse, der höher oder niedriger als der/die Schwellenwert(e) und/oder die Referenzverteilung(en) ist, das Vorhandensein der chromosomalen Aneuploidie beim Fötus anzeigt.

6. Verfahren nach Anspruch 4 oder 5, wobei (A) der Nachweis in Schritt (f) das Verwenden von mindestens zwei der ORs umfasst; *wobei vorzugsweise* die Anzahl der ORs gleich der Anzahl der Referenz-DMRs ist, die in Schritt (d) verwendet werden; *bevorzugter*, *wobei* sich eine der ORs auf demselben Chromosom befindet, etwa zwischen ungefähr 20 bp und ungefähr 20 kb aufwärts oder abwärts davon, wie eine Referenz-DMR, die in Schritt (d) verwendet wird, und jede andere der ORs sich auf demselben Chromosom befindet, etwa zwischen ungefähr 20 bp und ungefähr 20 kb aufwärts oder abwärts davon, wie eine weitere der Referenz-DMRs; und

(B) der Nachweis in Schritt (f) erfolgt unter Verwendung von: (x) derselben (denselben) nachweisbaren Markierung(en) für jede der ORs.

7. Verfahren nach einem der Ansprüche 1 bis 6, wobei der Nachweisschritt (c) und der Schritt (d) und der optionale Schritt (f) unter Verwendung des gleichen Aliquots der DNA der Probe und in demselben Reaktions-/Nachweisgefäß durchgeführt werden, und effektiv gleichzeitig miteinander, und mittels quantitativer Multiplex-Echtzeit-PCR auf Sondenbasis, unter Verwendung mindestens einer markierten Sonde, die für jede der DMRs und optionalen OR(s) spezifisch ist.

8. Verfahren nach einem der Ansprüche 1 bis 7, wobei das schwangere Weibchen ein Mensch ist und das für die chromosomale Aneuploidie relevante Chromosom ein aus der Liste, bestehend aus Chromosom 21, Chromosom 18 und Chromosom 13 ausgewähltes menschliches Chromosom ist.

9. Verfahren nach einem der Ansprüche 1 bis 8, wobei die DNA-Spezies, die aus Zellen eines Fötus und/oder der

Plazenta eines Fötus stammt, zirkulierende zellfreie DNA und die Probe eine Blutfraktion, wie etwa Plasma oder Serum, ist.

10. Verfahren nach einem der Ansprüche 1 bis 9, wobei das Reagens, das methylierte und unmethylierte DNA differentiell modifiziert, Folgendes umfasst:

ein Agens, das ausgewählt wird aus der Gruppe, bestehend aus: AatII, AclI, AclI, AfeI, AgeI, AgeI-HF, AscI, AsiSI, Aval, BceAI, BmgBI, BsaAI, BsaHI, BsiEI, BsiWI, BsmBI, BspDI, BsrFI, BssHII, BstBI, BstUI, ClaI, EagI, FaeI, FseI, FspI, HaeIII, HgaI, HhaI, HinfI, HpaII, Hpy99I, HpyCH4IV, KasI, MluI, NaeI, NarI, NgoMIV, NotI, NotI-HF, NruI, Nt.BsmAI, Nt.CviPII, PaeR7I, PfuTI, PmlI, PvuI, PvuI-HF, RsrII, SacII, SalI, SalI-HF, SfiI, SgrAI, SmaI, SnaBI, TspMI und ZraI.

11. Verfahren nach einem der Ansprüche 1 bis 10, wobei sich eine Referenz-DMR auf dem menschlichen Chromosom 5 und eine weitere Referenz-DMR auf dem menschlichen Chromosom 12 befindet.

12. Verfahren nach einem der Ansprüche 1 bis 11, wobei die ersten Ziel-DMRs sich jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus:

- einem, das ausgewählt ist aus der Liste, bestehend aus: SEQ ID NOs: 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198 und 199; oder

- der Liste, bestehend aus: AIRE, SIM2 und ERG oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz; oder

- der Liste, bestehend aus: PDE9A, PPP1R2P2, CBR1, DSCAM, C21orf29 und HLCS oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz; oder

- c21 orf57 oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz; oder

- der Liste, bestehend aus: SEQ ID NOs: 47, 48, 49, 50, 51, 52, 53, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186 und 187.

13. Verfahren nach Anspruch 12, wobei sich die ersten Ziel-DMRs jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus der Liste, bestehend aus: SEQ ID NOs: 47, 48, 49, 50, 51, 52, 53, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 173, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186 und 187.

14. Verfahren nach Anspruch 13, wobei sich die ersten Ziel-DMRs jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus der Liste, bestehend aus: SEQ ID NO: 51, 184, 185 und 186.

15. Verfahren nach Anspruch 14, wobei sich eine der ersten Ziel-DMRs in einer Region und/oder einem Gen befindet, die/das durch SEQ ID NO: 51 dargestellt wird und sich eine weitere der ersten Ziel-DMRs in einer Region und/oder einem Gen befindet, die/das durch SEQ ID NO: 185 dargestellt wird.

16. Verfahren nach einem der Ansprüche 1 bis 15, wobei sich die ersten Ziel-DMRs in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus:

- VAPA-APCDDI oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz; oder

- Maspin oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz

- der Liste, bestehend aus: SEQ ID NOs: 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 und 46.

17. Verfahren nach einem der Ansprüche 1 bis 16, wobei die ersten Ziel-DMRs sich jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus:

- der Liste, bestehend aus: SEQ ID NOs: 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33 und 34.

18. Verfahren nach einem der Ansprüche 1 bis 17, wobei sich die Referenz-DMRs jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus:

- der Liste, bestehend aus: RASSF1A, TBX3, ZFY, CDC42EP1, PCDHGA1, SOX14 und SPN oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz; oder
- der Liste, bestehend aus: SEQ ID NOs: 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146 und 147.

19. Verfahren nach Anspruch 18, wobei sich die Referenz-DMRs jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus der Liste, bestehend aus: SEQ ID NOs: 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146 und 147.

20. Verfahren nach Anspruch 19, wobei sich die Referenz-DMRs jeweils in einer Region und/oder einem Gen befinden, die unabhängig ausgewählt sind aus der Liste, bestehend aus SEQ ID NOs: 66, 102 und 126.

21. Verfahren nach Anspruch 20, wobei sich eine der Referenz-DMRs in einer Region und/oder einem Gen befindet, die/das durch SEQ ID NO: 66 dargestellt wird und eine weitere der Referenz-DMRs sich in einer Region und/oder einem Gen befindet, die/das durch SEQ ID NO: 102 dargestellt wird.

22. Verfahren nach einem der Ansprüche 1 bis 21, wobei sich mindestens eine der ersten Ziel-DMRs in einem Teil des Genoms und/oder einem Gen befindet, das DSCAM und/oder C21orf57 darstellt, oder in einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz.

23. Verfahren nach einem der Ansprüche 1 bis 21, wobei:

- (A) sich eine der ersten Ziel-DMRs in DSCAM befindet und sich eine weitere der ersten Ziel-DMRs in C21orf57 oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz befindet; und
- (B) sich eine der Referenz-DMRs in TBX3 befindet und sich eine weitere der Referenz-DMRs in PCDHGA1 oder einer nicht mehr als 10 kbp, 5 kbp, 1 kbp, 500 bp, 250 bp, 150 bp, 100 bp oder 50 bp aufwärts und/oder abwärts von einer solchen Region und/oder einem solchen Gen gelegenen DNA-Sequenz befindet.

24. Verfahren nach einem der Ansprüche 1 bis 21, wobei: (i) eine der ersten Ziel-DMRs die SEQ ID NO: 201 umfasst und eine weitere der ersten Ziel-DMRs die SEQ ID NO: 219 umfasst; und (ii) eine der Referenz-DMRs die SEQ ID NO: 203 umfasst und eine weitere der Referenz-DMRs die SEQ ID NO: 220 umfasst.

25. Verfahren nach einem der Ansprüche 1 bis 24,

- (A) wobei eine Vielzahl von Bestimmungssätzen für Schritt (c) und Schritt (d) und den optionalen Schritt (f) durchgeführt werden, wobei jeder Satz der Bestimmungen unter Verwendung eines unterschiedlichen DNA-Aliquots der Probe in einem anderen Gefäß und effektiv gleichzeitig mit jedem anderen Mitglied der Vielzahl von Bestimmungssätzen erfolgt; und
- (B) wobei die in Schritt (e) bestimmte(n) relative(n) Menge(n) oder Verhältnis(se) aus der Vielzahl von für Schritt (c) und Schritt (d) und optional Schritt (f) durchgeführten Bestimmungen bestimmt wird, vorzugsweise unter Verwendung von einer durchschnittlichen Menge an DNA, die für jeden Satz bestimmt wird, wie etwa eine für jeden Satz ermittelte mittlere, mediane oder modale Menge an DNA, vorzugsweise eine für jeden Satz bestimmte mittlere Menge an DNA.

26. Verfahren nach einem der Ansprüche 1 bis 25, wobei in Schritt (e) der/die relative(n) Menge(n) oder Verhältnis(se) mit (einem) Schwellenwert(en) und/oder (einer) Referenzverteilung(en) verglichen werden, wobei zwei oder mehrere

der relativen Mengen oder Verhältnisse, die höher oder niedriger als der/die Schwellenwert(e) und/oder die Referenzverteilung(en) sind, das Vorhandensein der chromosomalen Aneuploidie beim Fötus anzeigt; und wobei ein Schwellenwert und/oder eine Referenzverteilung aus einer Vielzahl von Proben bestimmt wird, wobei jede Probe von einem anderen schwangeren Weibchen entnommen wurde, beispielsweise durch Anwenden des in einem der Ansprüche 1 bis 22 ausgeführten Verfahrens auf solche Proben, wobei das Verfahren an einer Vielzahl von Proben durchgeführt wird, die jeweils von einem anderen schwangeren Weibchen entnommen wurden, wobei jedes Verfahren in einem separaten Gefäß und effektiv gleichzeitig mit jedem anderen Mitglied der Vielzahl von Proben erfolgt.

27. Verfahren nach einem der Ansprüche 1 bis 26, wobei eine gegebene DMR durch PCR-Primer amplifiziert wird, die:

- zwei oder drei Restriktionsstellen für das/die methylierungsempfindliche(n) Restriktionsenzym(e) überlappen, von denen jede eine Methylierungsstelle umfasst; und/oder
- derart gestaltet sind, dass sie eine, zwei, drei oder mehr Restriktionsstellen für das/die methylierungsempfindliche(n) Restriktionsenzym(e) flankieren, von denen jede eine Methylierungsstelle umfasst.

28. Verfahren nach einem der Ansprüche 1 bis 27, wobei das/die methylierungsempfindliche(n) Restriktionsenzym(e):

- eines aus der Gruppe, bestehend aus: BstUI, HhaI und HpaII ist; oder
- zwei oder drei der methylierungsempfindlichen Restriktionsenzyme ausgewählt aus der Gruppe, bestehend aus: BstUI, HhaI und HpaII umfassen.

29. Verfahren nach einem der Ansprüche 1 bis 27 oder 28, wobei die nachweisbaren Markierungen Fluoreszenzmarkierungen sind.

Revendications

1. Procédé de détection d'une aneuploïdie chromosomique chez un fœtus porté par une femelle enceinte, ledit procédé comprenant les étapes suivantes :

(a) fourniture d'un échantillon qui a été prélevé sur ladite femelle enceinte, lequel échantillon comprend de l'ADN qui provient de cellules d'un fœtus et/ou du placenta d'un fœtus mélangé avec de l'ADN différemment méthylé d'origine maternelle ;

(b) traitement de l'ADN présent dans ledit échantillon avec un réactif qui modifie différemment l'ADN méthylé et non méthylé, dans lequel ledit réactif est un agent qui digère sélectivement l'ADN non méthylé par rapport à l'ADN méthylé et ledit agent comprend une enzyme de restriction sensible à la méthylation ;

(c) détermination d'une quantité d'une première espèce cible d'ADN, étant le chromosome en lien avec l'aneuploïdie chromosomique, dans ledit échantillon par détection dans ledit échantillon de la présence de méthylation au niveau de deux premières régions différemment méthylées (DMR) cibles ou plus situées sur ledit chromosome, lesdites premières DMR cibles différemment méthylées entre l'ADN qui provient de cellules d'un fœtus et/ou du placenta d'un fœtus et l'ADN d'origine maternelle, dans lequel lesdites DMR sont hyperméthylées dans l'ADN fœtal et hypométhylées dans l'ADN maternel, la modification de l'ADN des premières DMR cibles par ledit réactif est sensible à la méthylation de l'ADN, dans lequel une quantité détectée d'ADN méthylé au niveau d'une ou plusieurs desdites premières DMR cibles indique ladite quantité de première espèce cible d'ADN dans ledit échantillon, dans lequel une quantité de la première espèce cible d'ADN est déterminée à partir de la quantité détectée de méthylation au niveau de l'une des premières DMR cibles et une quantité de la première espèce cible d'ADN est déterminée à partir de la quantité détectée de méthylation au niveau d'une autre des premières DMR cibles ;

(d) détermination d'une quantité d'espèces de référence d'ADN, étant un ou plusieurs chromosomes de référence, dans ledit échantillon par détection dans ledit échantillon de la présence de méthylation au niveau de deux DMR de référence ou plus sur ledit (lesdits) chromosome(s) de référence, lesdites DMR de référence différemment méthylées entre l'ADN qui provient de cellules d'un fœtus et/ou du placenta d'un fœtus et l'ADN d'origine maternelle, dans lequel lesdites DMR sont hyperméthylées dans l'ADN fœtal et hypométhylées dans l'ADN maternel, la modification de l'ADN de ces DMR de référence par ledit réactif est sensible à la méthylation de l'ADN, dans lequel une quantité détectée d'ADN méthylé au niveau d'une ou plusieurs desdites DMR de référence indique ladite quantité d'espèces de référence d'ADN dans ledit échantillon ; et

(e) détermination d'une (de) quantité(s) relative(s), de préférence d'un (de) rapport(s), d'une quantité déterminée à partir de l'étape (c) et d'une quantité déterminée à partir de l'étape (d), dans lequel deux ou plus de ces

quantités relatives sont déterminées pour deux quantités de première espèce cible d'ADN ou plus, chacune relativement à une ou plusieurs desdites premières DMR cibles, et dans lequel deux ou plus desdites quantités relatives indiquent la présence ou l'absence de l'aneuploïdie chromosomique chez le fœtus,

- 5 dans lequel, lesdites détections à l'étape (c) et à l'étape (d) sont réalisées en utilisant la même aliquote d'ADN dudit échantillon, et dans le même récipient, et effectivement simultanément pour lesdites premières DMR cibles et lesdites DMR de référence, et en utilisant : (x) le(s) même(s) marqueur(s) détectable(s) pour au moins deux desdites DMR de référence ; et (y) un (des) marqueur(s) détectable(s) différent(s) pour au moins deux desdites premières DMR cibles ; et :
- 10 dans lequel à l'étape (e) lesdit(e)s deux quantités relatives ou rapports ou plus sont comparés ou à un(e) (des) seuil(s) et/ou une (des) distribution(s) de référence, dans lequel lesdit(e)s deux quantités relatives ou rapports ou plus supérieurs ou inférieurs au dit (auxdits) seuil(s) et/ou à ladite (auxdites) distribution(s) de référence indiquent la présence de l'aneuploïdie chromosomique chez le fœtus.
- 15 2. Procédé selon la revendication 1, dans lequel lesdites détections à l'étape (c) et à l'étape (d) sont réalisées en utilisant : (x) le(s) même(s) marqueur(s) détectable(s) pour chacune desdites au moins deux DMR de référence ; et (y) un (des) marqueur(s) détectable(s) différent(s) pour chacune desdites au moins deux premières DMR cibles.
- 20 3. Procédé selon la revendication 1 ou 2, dans lequel deux ou plus desdites DMR de référence se trouvent sur des chromosomes de référence différents.
4. Procédé selon l'une quelconque des revendications 1 à 3, comprenant en outre l'étape suivante :
 (f) détermination d'une quantité d'ADN total dans ledit échantillon par détection d'au moins une autre région (OR) qui n'est pas différemment méthylée entre l'ADN qui provient de cellules d'un fœtus et/ou du placenta d'un fœtus
 25 et l'ADN d'origine maternelle, la modification de cette(s) OR par ledit réactif n'est pas sensible à la méthylation de l'ADN,
 dans lequel, lesdites détections à l'étape (c) et à l'étape (d) et à l'étape (f) sont réalisées en utilisant la même aliquote d'ADN dudit échantillon, et dans le même récipient, et effectivement simultanément pour lesdites premières DMR cibles et lesdites DMR de référence et sur ladite (lesdites) autre(s) région(s), et en utilisant : (x) le(s) même(s)
 30 marqueur(s) détectable(s) pour au moins deux desdites DMR de référence ; et (y) un (des) marqueur(s) détectable(s) différent(s) pour au moins deux desdites premières DMR cibles et pour au moins l'une de ladite (desdites) OR ;
 facultativement :
 dans lequel ladite (lesdites) OR est (sont) située(s) sur un ou plusieurs chromosome(s) de référence, de préférence au moins l'une desdites OR étant située sur le(s) même(s) chromosome(s) de référence qu'au moins l'une desdites
 35 DMR de référence.
5. Procédé selon l'une quelconque des revendications 1 à 4, dans lequel à l'étape (e) chacune de ladite (desdites) quantité(s) relative(s) ou dudit (desdits) rapport(s) est comparé à un (des) seuil(s) et/ou une (des) distribution(s) de référence, dans lequel chacune de ladite (desdites) quantité(s) relative(s) ou dudit (desdits) rapport(s) supérieurs
 40 ou inférieurs au dit (auxdits) seuil(s) et/ou à ladite (auxdites) distribution(s) de référence indique la présence de l'aneuploïdie chromosomique chez le fœtus.
6. Procédé selon la revendication 4 ou 5, dans lequel (A) ladite détection à l'étape (f) comprend l'utilisation d'au moins deux desdites OR ; de préférence dans lequel, le nombre desdites OR est le même que le nombre de DMR de référence utilisées à l'étape (d) ; de manière davantage préférée dans lequel, l'une desdites OR se trouve sur le même chromosome qu'une, par exemple entre environ 20 pb et environ 20 kb en amont ou en aval d'une, DMR de référence utilisée à l'étape (d) et chaque autre desdites OR se trouve sur le même chromosome qu'une, par exemple entre environ 20 pb et environ 20 kb en amont ou en aval d'une, autre desdites DMR de référence ; et
 45 (B) ladite détection à l'étape (f) est faite en utilisant : (x) le(s) même(s) marqueur(s) détectable(s) pour chacune desdites OR.
 50
7. Procédé selon l'une quelconque des revendications 1 à 6, dans lequel ladite étape de détection (c) et ladite étape (d), et l'étape facultative (f), sont réalisées en utilisant la même aliquote d'ADN dudit échantillon, et dans le même récipient de réaction/détection, et effectivement simultanément les unes avec les autres, et par PCR quantitative
 55 en temps réel multiplexe basée sur des sondes utilisant au moins une sonde marquée spécifique de chacune desdites DMR et de ladite ou desdites OR facultative(s).
8. Procédé selon l'une quelconque des revendications 1 à 7, dans lequel la femelle enceinte est un être humain et le

chromosome en lien avec l'aneuploidie chromosomique est un chromosome humain sélectionné dans les listes constituées des : chromosome 21, chromosome 18 et chromosome 13.

9. Procédé selon l'une quelconque des revendications 1 à 8, dans lequel ladite espèce d'ADN qui provient de cellules d'un fœtus et/ou du placenta d'un fœtus est un ADN acellulaire circulant et ledit échantillon est une fraction du sang telle que le plasma ou le sérum.

10. Procédé selon l'une quelconque des revendications 1 à 9, dans lequel ledit réactif qui modifie différenciellement l'ADN méthylé et l'ADN non méthylé comprend

un agent sélectionné dans le groupe constitué des : AatII, Acil, AclI, Afel, Agel, Agel-HF, Ascl, AsiSI, Aval, BceAI, BmgBI, BsaAI, BsaHI, BsiEI, BsiWI, BsmBI, BspDI, BsrFI, BssHII, BstBI, BstUI, ClaI, EagI, Faul, FseI, FspI, HaeII, HgaI, HhaI, HinPII, HpaII, Hpy99I, HpyCH4IV, Kasi, MluI, NaeI, NarI, NgoMIV, NotI, NotI-HF, Nrul, Nt.BsmAI, Nt.CviPII, PaeR7I, PluTI, PmlI, PvuI, PvuI-HF, RsrII, SacII, Sall, Sall-HF, SfoI, SgrAI, SmaI, SnaBI, TspMI et ZraI.

11. Procédé selon l'une quelconque des revendications 1 à 10, dans lequel une DMR de référence se trouve sur le chromosome 5 humain et l'autre DMR de référence se trouve sur le chromosome 12 humain.

12. Procédé selon l'une quelconque des revendications 1 à 11, dans lequel lesdites premières DMR cibles sont chacune situées dans une région et/ou un gène indépendamment sélectionnés parmi :

- un sélectionné dans la liste constituée de : SEQ ID NO : 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198 et 199 ; ou
- la liste constituée de : AIRE, SIM2, et ERG ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ; ou
- la liste constituée de : PDE9A, PPP1 R2P2, CBR1, DSCAM, C21orf29 et HLCS ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ; ou
- C21 orf57 ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ; ou
- la liste constituée de : SEQ ID NO : 47, 48, 49, 50, 51, 52, 53, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186 et 187.

13. Procédé selon la revendication 12, dans lequel lesdites premières DMR cibles sont chacune situées dans une région et/ou un gène indépendamment sélectionnés dans la liste constituée de : SEQ ID NO : 47, 48, 49, 50, 51, 52, 53, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186 et 187.

14. Procédé selon la revendication 13, dans lequel lesdites premières DMR cibles sont chacune situées dans une région et/ou un gène indépendamment sélectionnés dans la liste constituée de : SEQ ID NO : 51, 184, 185 et 186.

15. Procédé selon la revendication 14, dans lequel l'une desdites premières DMR cibles se trouve dans une région et/ou un gène étant SEQ ID NO : 51 et une autre desdites premières DMR cibles se trouve dans une région et/ou un gène étant SEQ ID NO : 185.

16. Procédé selon l'une quelconque des revendications 1 à 15, dans lequel lesdites premières DMR cibles sont chacune situées dans une région et/ou un gène indépendamment sélectionnés parmi :

- VAPA-APCDDI ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ; ou
- maspin ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ;
- la liste constituée de : SEQ ID NO : 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 et 46.

17. Procédé selon l'une quelconque des revendications 1 à 16, dans lequel lesdites premières DMR cibles sont chacune situées dans une région et/ou un gène indépendamment sélectionnés parmi :

- la liste constituée de : SEQ ID NO : 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33 et 34.

18. Procédé selon l'une quelconque des revendications 1 à 17, dans lequel lesdites DMR de référence sont chacune situées dans une région et/ou un gène indépendamment sélectionnés parmi :

- la liste constituée de : RASSF1A, TBX3, ZFY, CDC42EP1, PCDHGA1, SOX14 et SPN ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ; ou
- la liste constituée de : SEQ ID NO : 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146 et 147.

19. Procédé selon la revendication 18, dans lequel lesdites DMR de référence sont chacune situées dans une région et/ou un gène indépendamment sélectionnés dans la liste constituée de : SEQ ID NO : 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146 et 147.

20. Procédé selon la revendication 19, dans lequel lesdites DMR de référence sont chacune situées dans une région et/ou un gène indépendamment sélectionnés dans la liste constituée des SEQ ID NO : 66, 102 et 126.

21. Procédé selon la revendication 20, dans lequel l'une desdites DMR de référence se trouve dans une région et/ou un gène étant SEQ ID NO : 66 et une autre desdites DMR de référence se trouve dans une région et/ou un gène étant SEQ ID NO : 102.

22. Procédé selon l'une quelconque des revendications 1 à 21, dans lequel au moins l'une desdites premières DMR cibles se trouve dans une partie du génome et/ou d'un gène qui est DSCAM et/ou C21orf57, ou une séquence d'ADN n'excédant pas 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène.

23. Procédé selon l'une quelconque des revendications 1 à 21, dans lequel :

- (A) l'une desdites premières DMR cibles se trouve dans DSCAM et une autre desdites premières DMR cibles se trouve dans C21orf57 ou une séquence d'ADN pas supérieure à 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène ; et
- (B) une desdites DMR de référence se trouve dans TBX3 et une autre desdites DMR de référence se trouve dans PCDHGA1 ou une séquence d'ADN pas supérieure à 10 kpb, 5 kpb, 1 kpb, 500 pb, 250 pb, 150 pb, 100 pb ou 50 pb en amont et/ou en aval d'une telle région et/ou d'un tel gène.

24. Procédé selon l'une quelconque des revendications 1 à 21, dans lequel : (i) l'une desdites premières DMR cibles comprend SEQ ID NO : 201 et l'autre desdites premières DMR cibles comprend SEQ ID NO : 219 ; et (ii) l'une desdites DMR de référence comprend SEQ ID NO : 203 et l'autre desdites DMR de référence comprend SEQ ID NO : 220.

25. Procédé selon l'une quelconque des revendications 1 à 24,

- (A) dans lequel une pluralité d'ensembles de déterminations sont faits pour l'étape (c) et l'étape (d), et l'étape facultative (f), avec chaque ensemble desdites déterminations fait en utilisant une aliquote différente d'ADN dudit échantillon, dans un récipient différent et efficacement simultanément avec chaque autre élément de la pluralité d'ensembles de déterminations ; et
- (B) dans lequel la (les) quantité(s) relative(s) ou le (les) rapport(s) déterminé(s) à l'étape (e) sont déterminés à partir de ladite pluralité d'ensembles de déterminations faits pour l'étape (c) et l'étape (d), et l'étape facultative (f), de préférence en utilisant une quantité moyenne d'ADN déterminée pour chaque ensemble, par exemple une quantité moyenne, médiane ou modale d'ADN déterminée pour chaque ensemble, de préférence une quantité moyenne d'ADN déterminée pour chaque ensemble.

26. Procédé selon l'une quelconque des revendications 1 à 25, dans lequel à l'étape (e) ladite (desdites) quantité(s)

relative(s) ou ledit (lesdits) rapport(s) sont comparés à un (des) seuil(s) et/ou une (des) distribution(s) de référence, dans lequel deux ou plus de ladite (desdites) quantité(s) relative(s) ou dudit (desdits) rapport(s) supérieurs ou inférieurs au dit (auxdits) seuil(s) et/ou distribution(s) de référence indique la présence de l'aneuploïdie chromosomique chez le fœtus ; et

dans lequel un seuil et/ou une distribution de référence sont déterminés à partir d'une pluralité d'échantillons, chaque échantillon prélevé chez une femelle enceinte différente, par exemple en pratiquant le procédé selon l'une quelconque des revendications 1 à 22 sur ces échantillons, dans lequel ledit procédé est réalisé sur une pluralité d'échantillons prélevés chacun chez une femelle enceinte différente, chaque procédé réalisé dans un récipient séparé, et efficacement simultanément avec chaque autre membre de la pluralité d'échantillons.

27. Procédé selon l'une quelconque des revendications 1 à 26, dans lequel une DMR donnée est amplifiée par des amorces de PCR qui :

- chevauchent deux ou trois sites de restriction pour l'enzyme (les enzymes) de restriction sensible(s) à la méthylation, chacune d'entre elles comprenant un site de méthylation ; et/ou
- sont conçues pour flanquer un, deux, trois sites de restriction ou plus pour l'enzyme (les enzymes) de restriction sensible(s) à la méthylation, chacune d'entre elles comprenant un site de méthylation.

28. Procédé selon l'une quelconque des revendications 1 à 27, dans lequel ladite enzyme (lesdites enzymes) de restriction sensible(s) à la méthylation :

- sont une sélectionnée dans le groupe constitué de : BstUI, Hha I et HpaII ; ou
- comprennent deux ou trois des enzymes de restriction sensibles à la méthylation sélectionnées dans le groupe constitué de : BstUI, Hha I et HpaII.

29. Procédé selon l'une quelconque des revendications 1 à 27 ou 28, dans lequel les marqueurs détectables sont des marqueurs fluorescents.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9





FIGURE 10



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 03062441 A1 [0005]
- WO 2003020974 A [0006]
- WO 2005118852 A [0006]
- WO 2007132166 A [0007]
- WO 2007132167 A [0007]
- WO 2011034631 A [0007] [0091] [0093] [0094] [0097] [0098]
- WO 2011092592 A [0007] [0088] [0098]
- US 5804375 A, Gelfand [0071]
- EP 0543942 A [0071]
- US 6258569 B, Livak [0071]
- EP 0792374 A [0071]
- US 6727356 B, Reed [0071]
- EP 1235938 A [0071]
- US 6929907 B, Woudenberg [0072]
- EP 0706649 A [0072]
- US 5994056 A [0072]
- EP 0512334 A [0072]
- WO 2007132176 A [0090]
- EP 1751307 A [0092]
- US 6200756 B, Herman [0116]
- EP 0954608 A [0116]
- US 6331393 B, Laird [0116]
- EP 1185695 A [0116]

Non-patent literature cited in the description

- NORWITZ ; LEVY. *Rev Obs Gyn*, 2013, vol. 6, 48 [0002]
- LO et al. *Lancet*, 1997, vol. 350, 485 [0003]
- LO. *N Eng J Med*, 1998, vol. 229, 1734 [0003]
- LO et al. *Am J Hum Genet*, 1998, vol. 63, 768 [0003]
- GO et al. *Human Reprod Update*, 2011, vol. 17, 372 [0003]
- HINDSON et al. *Anal Chem*, 2011, vol. 83, 8604 [0004]
- POON et al. *Clin Chem*, 2002, vol. 48, 35 [0006]
- CHIM et al. *PNAS*, 2005, vol. 102, 14753 [0006]
- TONG et al. *Clin Chem*, 2006, vol. 52, 2194 [0006]
- OLD et al. *Reprod Biomed Online*, 2007 [0007]
- CHIM. *Clin Chem*, 2008, vol. 54, 500 [0007]
- PAPAGEORGIOU et al. *Am J Pathol*, 2009, vol. 174, 1609 [0007]
- LIM et al. *PLoS One*, 2011, vol. 6, e27709 [0008]
- TONG et al. *Clin Chem*, 2010, vol. 56, 90 [0008]
- PAPAGEORGIOU et al. *Nat Med*, 2011, vol. 17, 510 [0008]
- TSALIKI et al. *Prenat Diag*, 2012, vol. 21, 996 [0008]
- TONG et al. *Nat Med*, 2012, vol. 18, 1327 [0008]
- LIM et al. *Clin Chem Lab Med*, 2014, vol. 52, 641 [0009]
- CHIM. *Clin Chem Lab Med*, 2014, vol. 52, 585 [0009]
- LIM et al. *BMC Medical Genomics*, 2014, vol. 7, 1 [0088]
- HAHN et al. *Placenta*, 2011, vol. 32, S17 [0112]
- YEGNASUBRAMANIAN et al. *Nuc Acid Res*, 2006, vol. 34, e19 [0116]
- ZEYBEK et al. *J Obstet Gynaecol Res*, 2013, vol. 39, 632 [0144] [0186]
- JAKOBSEN et al. *Transfusion*, 2013, vol. 53, 1956 [0144] [0186]
- CHEN et al. *Am J Med Genet A*, 2013, vol. 161A, 1817 [0145] [0177]
- NYGREN et al. *Clin Chem*, 2010, vol. 56, 1627 [0172]
- CHAN et al. *Clin Chem*, 2006, vol. 42, 2211 [0172]
- CHIU et al. *Am J Pathol*, 2007, vol. 170, 941 [0172]
- STUMM et al. *Prenat Diag*, 2014, vol. 34, 185 [0174]
- LEUNG et al. *Prenat Diag*, 2013, vol. 33, 675 [0176]
- QU et al. *Am J Pathol*, 2007, vol. 170, 941 [0176]
- SPERLING et al. *Acta Obstet Gynecol Scand*, 2001, vol. 80, 287 [0177]
- PAPANTONIOU et al. *Prenat Diag*, 2013, vol. 33, 682 [0186]