



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/04/21
(87) Date publication PCT/PCT Publication Date: 2017/10/26
(85) Entrée phase nationale/National Entry: 2018/10/09
(86) N° demande PCT/PCT Application No.: US 2017/028996
(87) N° publication PCT/PCT Publication No.: 2017/185061
(30) Priorité/Priority: 2016/04/21 (US62/325,860)

(51) Cl.Int./Int.Cl. *A61K 31/519* (2006.01),
A61P 25/00 (2006.01)
(71) Demandeurs/Applicants:
ASTROCYTE PHARMACEUTICALS, INC., US;
THE BOARD OF REGENTS OF THE UNIVERSITY OF
TEXAS SYSTEM, US;
THE UNITED STATES OF AMERICA, AS
REPRESENTED BY THE SECRETARY,
DEPARTMENT OF HEALTH AND HUMAN
SERVICES, US
(72) Inventeurs/Inventors:
KORINEK, WILLIAM S., US;
LECHLEITER, JAMES D., US; ...

(54) Titre : COMPOSES ET PROCEDES POUR TRAITER DES ETATS NEUROLOGIQUES ET CARDIOVASCULAIRES
(54) Title: COMPOUNDS AND METHODS FOR TREATING NEUROLOGICAL AND CARDIOVASCULAR CONDITIONS

(57) **Abrégé/Abstract:**

The present invention relates to compounds and methods of use thereof for treatment of certain disorders and conditions, for example brain injuries such as stroke or traumatic brain injuries.

(21) **3 020 490**

(13) **A1**

(72) **Inventeurs(suite)/Inventors(continued):** LISTON, THEODORE E., US; JACOBSON, KENNETH A., US

(74) **Agent:** GOWLING WLG (CANADA) LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
26 October 2017 (26.10.2017)



(10) International Publication Number
WO 2017/185061 A1

(51) International Patent Classification:

A61K 31/519 (2006.01) *A61P 25/00* (2006.01)

(21) International Application Number:

PCT/US2017/028996

(22) International Filing Date:

21 April 2017 (21.04.2017)

Published:

— with international search report (Art. 21(3))

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/325,860 21 April 2016 (21.04.2016) US

(71) Applicants: **ASTROCYTE PHARMACEUTICALS, INC.** [US/US]; 245 First Street, Suite 1800, Cambridge, Massachusetts 02142 (US). **THE BOARD OF REGENTS OF THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 8403 Floyd Curl Drive, Mail Code 7746, San Antonio, Texas 07829 (US). **THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES** [US/US]; Office of Technology Transfer, National Institutes of Health, 6011 Executive Boulevard, Suite 325, MSC 7660, Bethesda, Maryland 20892-7660 (US).

(72) Inventors: **KORINEK, William S.**; 163 Long Wharf Drive, Mystic, Connecticut 06355 (US). **LECHLEITER, James D.**; 112 Martha Glynn Court, San Antonio, Texas 78213 (US). **LISTON, Theodore E.**; 33 Kidds Way, Stonington, Connecticut 06378 (US). **JACOBSON, Kenneth A.**; 11218 Watermill Lane, Silver Spring, Maryland 20902 (US).

(74) Agent: **REID, Andrea L.C.** et al.; One International Place, 40th Floor, 100 Oliver Street, Boston, Massachusetts 02110-2605 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,

(54) Title: COMPOUNDS AND METHODS FOR TREATING NEUROLOGICAL AND CARDIOVASCULAR CONDITIONS

(57) Abstract: The present invention relates to compounds and methods of use thereof for treatment of certain disorders and conditions, for example brain injuries such as stroke or traumatic brain injuries.

WO 2017/185061 A1

COMPOUNDS AND METHODS FOR TREATING NEUROLOGICAL AND CARDIOVASCULAR CONDITIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/325,860, filed on April 21, 2016, the entirety of which is hereby incorporated by reference.

FIELD OF THE INVENTION

[0002] The present invention relates to compounds and methods of use thereof for treating, ameliorating, or promoting recovery from certain conditions of the brain, central nervous system (CNS), or cardiovascular system such as a brain injury, a neurodegenerative condition, or cardiac ischemia.

STATEMENT OF GOVERNMENT SUPPORT

[0003] This invention was made with government support under grant no. NS093756 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

[0004] Brain injuries are a distressingly common medical condition and one of the leading causes of morbidity and mortality worldwide. The brain is particularly susceptible to injury as neurons have a limited capacity to repair. When an individual is born, the brain already has essentially all the neurons it will have in life. Unlike other cells in the body, neurons stop reproducing shortly after birth. If these cells are injured or die, they are not replaced, often culminating in the disabling and largely irreversible degradation of a person's cognitive and sensorimotor capacity. Conditions that result in nerve cell death and damage range from ischemic episodes (e.g., stroke) and trauma, to degenerative disorders (e.g., Alzheimer's disease).

[0005] Injury to the Central Nervous System (CNS) is a substantial cause of death and disability worldwide. For example, according to the CDC approximately 1.7 million people sustain a Traumatic Brain Injury (TBI) annually, costing the U.S. economy in

excess of \$60 billion per year in terms of medical costs and lost productivity (Finkelstein, E; Corso, P; Miller, T, *The Incidence and Economic Burden of Injuries in the United States*, Oxford University Press: New York, 2006). Additionally, stroke is the third leading cause of death in the U.S. with an estimated incidence of 795,000 cases annually, a major cause of disability, and costing the U.S. economy over \$34 billion per year (NINDS, 2014; stroke.nih.gov; and Mozaffarian D, Benjamin EJ, Go AS, *et al.* “Heart disease and stroke statistics—2015 update: a report from the American Heart Association,” *Circulation*. 2015 ;e29-322).

[0006] In the acute setting, there is an opportunity to treat patients within 24 hours that can limit the extent of the damage. Immediately after an ischemic or hemorrhagic stroke, the site of insult in the brain typically contains a core of tissue that is irreversibly damaged, and then also an area of viable but at-risk tissue called the penumbra. During this period, the insufficient oxygen and glucose supply to brain cells results in further secondary injury to the penumbra. The lack of oxygen and glucose decreases energy production by cell mitochondria. An immediate effect of this energy depletion is failure of the ion pumps, which by elevating extracellular potassium (K^+) ions, results in waves of recurrent spreading depolarizations in brain tissue. At the same time, influx of sodium (Na^+) ions into cells, followed by chloride (Cl^-) ions, results in the swelling of cells due to osmotic pressure elevation, pressuring nearby neurons and their processes, ultimately leading to lysis (cell rupture) and inflammatory responses. In general, this disruption of ion homeostasis leads to excitotoxicity, cell swelling and cell death that extends damage to adjacent tissue and expands lesions by secondary mechanisms. There is a need for effective treatments during the initial 24 hours to protect the stressed brain cells. The propagation of brain damage in stroke is similar to that observed in other forms of brain injury such as trauma and concussions.

[0007] Beyond acute treatment, effective astrocyte function plays a key role in broader neurorestoration – in the period 24-96 hours following brain insult, in the period months-years in patients with neurodegeneration such as Alzheimer’s, or most generally in aged individuals. The inability of brain cells to regenerate requires the remaining intact brain tissue to reorganize in an attempt to recover any loss of function. This potential for neural reorganization is diminished in older individuals.


[0008] GPCR receptors have been suggested to mediate cardioprotective effects. Therefore, there is potential to treat heart and cardiovascular conditions by similar mechanisms of action via modulation of these receptors.

[0009] There is urgent and compelling unmet medical need for more effective treatments for brain injuries, CNS injuries, heart and cardiovascular diseases, and related conditions, as well as promoting neurorestoration in patients having a neurodegenerative condition such as Alzheimer's.

SUMMARY OF THE INVENTION


[0010] In one aspect, the present invention provides a method of treating an injury, disease, or condition selected from traumatic brain injury (TBI), stroke, a neurodegenerative condition, or a heart or cardiovascular disease, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R).

[0011] In one aspect, the present invention provides a method of treating an injury, disease, or condition selected from traumatic brain injury (TBI), stroke, a neurodegenerative condition, or a heart or cardiovascular disease, comprising administering to a patient in need thereof an effective amount of a biased agonist, partial agonist, or biased partial agonist of an A₃ adenosine receptor (A₃R) selected from




a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.


[0012] In another aspect, the present invention provides a method of treating a brain or central nervous system (CNS) injury or condition selected from traumatic brain injury (TBI) or stroke, comprising administering to a patient in need thereof an effective amount of a compound selected from:



a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.

[0013] In another aspect, the present invention provides a compound selected from the group consisting of:





[0014] In another aspect, the present invention provides a method of treating or ameliorating a traumatic brain injury (TBI), radiation damage, stroke, migraine headache, a heart or cardiovascular disease, or neurodegenerative disorder, comprising administering to a patient in need thereof an effective amount of a disclosed compound.

[0015] In another aspect, the present invention provides a method of treating or ameliorating a traumatic brain injury (TBI), radiation damage, stroke, migraine headache, a heart or cardiovascular disease, or neurodegenerative disorder, comprising administering to a patient in need thereof an effective amount of a compound selected from: adenosine, ADP, 2-methylthio-ADP trisodium salt, ATP, ATP disodium salt, α,β -methylene ATP, α,β -methyleneadenosine 5'-triphosphate trisodium salt, 2-methylthioadenosine triphosphate tetrasodium salt, 2-MeSATP, BzATP triethylammonium salt, inosine, cytidine, acylated cytidines, cytidine-monophosphate (CMP), cytidine diphosphate (CDP), cytidine triphosphate (CTP), CDP-choline, CMP-choline, denufosol, denufosol tetrasodium, GTP, ITP, MRS 541, MRS 542, MRS 1760, MRS 2179, MRS 2279, MRS 2341, MRS 2365, MRS 2500, MRS 2690, MRS 2698, MRS 3558, MRS 4322, MRS 5151, MRS 5676, MRS 5678, MRS 5697, MRS 5698, MRS 5923, MRS 5930, Benzyl-NECA, IB-MECA, Cl-IB-MECA, LJ529, DPMA, CCPA, DBXRM, HEMADO, PEMADO, HENECA, PENECA, CP608,039, CP532,903,

CGS21680, AR132, VT72, VT158, VT160, VT163, PSB 0474, uridine 5'-diphosphate (UDP), UDP-glucose, uridine β -thiodiphosphate (UDP β S), uridine 5'-triphosphate (UTP), uridine γ -thiophosphate (UTP γ S), 2-thioUTP tetrasodium salt, UTP γ S trisodium salt, uridine-5'-diphosphoglucose, diuridine triphosphate, 2-(hexylthio) (HT)-AMP, diadenosine pentaphosphate, 2'-deoxy-2'-amino-UTP, 2-thio-UTP, triacetyluridine, diacetyl/acyl uridine, uridine, suramin, dipyridamole analogs, diadenosine tetraphosphate Ap₄U, Ap₄A, INS365, INS37217, or INS48823; wherein each sugar may be replaced with a methanocarba sugar in the North or South conformation or each sugar may be replaced with a D-ribosugar; or a pharmaceutically acceptable salt thereof.

BRIEF DESCRIPTION OF THE FIGURES

[0016] **Figure 1A** shows plasma and brain concentration-time profiles of MRS4322 in mice following intraperitoneal administration. **Figure 1B** shows plasma and brain extracellular fluid concentration-time profiles of MRS4322 in neonatal pigs following intravenous administration.

[0017] **Figure 2** shows plasma and brain concentration-time profiles of MRS4322 in mice following intraperitoneal administration of MRS2365.

[0018] **Figure 3** shows *in vitro* stability of MRS2365 in EDTA-treated mouse plasma.

[0019] **Figure 4** shows *in vitro* stability of MRS2365 in EDTA-treated human plasma.

[0020] **Figure 5** shows *in vitro* stability of MRS2365 in EDTA-treated mouse whole blood.

[0021] **Figure 6** shows *in vitro* stability of MRS2365 in EDTA-treated human whole blood.

[0022] **Figure 7** shows *in vitro* stability of MRS2365 in EDTA- and Heparin-treated mouse plasma over a 90-second incubation period.

[0023] **Figure 8** shows *in vitro* stability of MRS2365 in EDTA- and Heparin-treated human plasma over a 90-second incubation period.

[0024] **Figure 9** shows positive-ion LC/MS-MS ion chromatograms and product ion spectra of MRS2365 metabolites in heparinized human plasma following a 30-minute incubation.

[0025] **Figure 10** shows positive-ion LC/MS-MS ion chromatograms and product ion spectra of MRS2365 metabolites in heparinized human whole blood following a 30-minute incubation.

[0026] **Figure 11** shows a scheme representing metabolites of MRS2365 detected by positive- and negative-ion LC/MS-MS in heparinized human plasma and whole blood. Two metabolites of MRS2365 were observed following incubation at 100 μ M in human whole blood and plasma (lithium heparin as anti-coagulant): M2, identified as the partially dephosphorylated metabolite of MRS2365 (i.e. with one phosphate group remaining, MRS2347); and M1, identified as the fully dephosphorylated metabolite of MRS2365 (i.e. MRS4322).

[0027] **Figure 12A-D** shows the results of MRS4322, Cl-IB-MECA and MRS2365 treatments on mice exposed to TBI or sham (control experiment). TBI normally induces increases in GFAP expression. Mice underwent sham or TBI (on **Ipsilateral** side of brain) and received treatments as labeled 30 min post-TBI. Plasma was obtained from mice at 7 days post-injury, then the mice were sacrificed to obtain brain homogenates from the ipsilateral (“Ipsi”) and contralateral (“Cntr”) hemispheres (middle third). Western blot analysis were normalized to actin. (**Figure 12A** and **12C**) Representative blots are shown for **Ipsilateral** brain homogenates and the plasma at day 7. Administration of MRS4322 or MRS2365 decreased brain levels of GFAP expression in mice exposed to TBI. (**Figure 12B** and **12D**) Data were pooled from 3 distinct experiments (N = number of mice per treatment) and plotted as bar histograms, shown as mean of control $\pm$ SEM. * $p < 0.05$ and ** $p < 0.01$ from TBI untreated (red bar).

[0028] **Figure 13A-D** shows the results of photothrombosis-induced stroke infarction experiments and the effects of administering MRS4322 and/or the specific adenosine type A₃ receptor antagonist MRS1523. MRS4322 reduced the effects of stroke infarction, but these effects were reversed by MRS1523. (**Figure 13A**) Coronal sections of mouse brains (each group of sections is from a single mouse) stroked with photothrombosis in vehicle (saline injected) and in treated mice, MRS4322, and MRS2365 as labelled. Stroked mice received either vehicle or treatments (IP injections) within 30 minutes of ischemia. Mice were then sacrificed at 24 hours post-stroke, their brains removed, sectioned and stained with TTC. (**Figure 13B**) Coronal sections from stroked mice pre-

injected with the A₃ receptor antagonist MRS1523. **(Figure 13C)** Average TTC-stained stroke volumes of MRS4322, MRS2365, MRS5698 and Cl-IBMECA as labeled. **(Figure 13D)** Stroke volumes in mice pretreated with the A₃ receptor antagonist MRS1523 as indicated. Data were pooled from 2 experiments (N = number of mice per treatment) and plotted as mean +/- SEM. ** p<0.01 and ***p<0.001.

[0029] **Figure 14A-B** shows the results of photothrombosis-induced stroke infarction experiments and the effects of administering MRS4322, MRS1873, or vehicle. Photothrombosis-induced stroke infarctions are reduced by MRS4322 and MRS1873. **(Figure 14A)** Coronal sections of brains stroked with photothrombosis in vehicle (saline injected) and in treated mice, MRS4322 and MRS1873 as labelled. Stroked mice received either vehicle or treatments (IP injections) within 30 minutes of ischemia. Mice were then sacrificed at 24 hours post-stroke, their brains removed, sectioned and stained with TTC. **(Figure 14B)** Average TTC-stained stroke volumes as labeled. Data were pooled from 3 experiments (N = number of mice per treatment) and plotted as mean +/- SEM. ** p<0.01.

[0030] **Figure 15** shows the structures of certain compounds suitable for use in the claimed invention.

[0031] **Figure 16** shows brain and cerebrospinal fluid concentration-time profiles of MRS4322 in neonatal pigs following intravenous administration. Following intravenous administration to neonatal pigs, MRS4322 concentrations were detectable in plasma, brain, brain extracellular fluid and cerebrospinal fluid samples.

[0032] **Figure 17** shows competition binding experiments of MRS4322 versus the A₃ agonist radioligand [³H]NECA (10 nM) at human A₃ receptors expressed in CHO cells. The calculated K_i-value for MRS4322 was **1490 ± 410 nM**.

[0033] **Figure 18** shows competition binding experiments of MRS4322 versus the A₃ agonist radioligand [³H]NECA (10 nM) at mouse A₃ receptors expressed in CHO cells. The calculated K_i-value for MRS4322 was **4940 ± 974 nM**.

[0034] **Figure 19** shows cAMP accumulation experiments of MRS4322 and NECA at human A₃ receptors expressed in CHO cells. The calculated EC₅₀-value for MRS4322 was **3630 ± 370 nM**; for NECA an EC₅₀ value of **41.8 ± 6.3 nM** was determined.

[0035] **Figure 20** shows cAMP accumulation experiments of MRS4322 and NECA at mouse A₃ receptors expressed in CHO cells. The calculated EC₅₀-value for MRS4322 was 759 ± 170 nM; for NECA an EC₅₀ value of 6.85 ± 0.88 nM was determined.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

1. Brain, CNS, Cardiovascular, and Other Injuries and Conditions

[0036] In some embodiments, the present invention provides a new approach to preventing and/or treating brain damage associated with acute brain trauma as well as longer term diseases of the brain and CNS and heart and cardiovascular diseases and conditions. In one aspect, the present invention provides methods of treating such injuries, diseases, and conditions by utilizing neuroprotective and neurorestorative effects mediated by astrocytes, which are now understood as the key natural caretaker cell of neurons, as well as the astrocyte mitochondria, which supply a significant portion of the brain's energy. In another aspect, the present invention provides methods of treating such injuries, diseases, and conditions by cardioprotective and regenerative effects mediated by A₃R receptors. Regarding neuroprotective and neurorestorative effects, without wishing to be bound by theory, it is believed that selective enhancement of astrocyte energy metabolism mediated by A₃R and/or P2Y₁ receptors promotes astrocyte caretaker functions, such as their neuroprotective and neurorestorative functions, in turn enhancing the resistance of neurons and other cells to both acute injury and long term stress. In some cases, it may be advantageous to achieve biased, i.e. selective or preferential, of one or more pathways mediated by A₃R and/or P2Y₁ receptors wherein one or more undesired pathways are not activated, or activated to a lesser degree. In addition to or as an alternative to astrocytes, neuroprotective or neurorestorative function of glia, microglia, neurons, endothelium cells and other brain and/or CNS cell types may be activated. Accordingly, in one aspect, the present invention provides compounds and methods of use thereof for treating, ameliorating, or promoting recovery from certain conditions of the brain or central nervous system (CNS) such as brain injuries, for example by increasing neuroprotection and/or neurorestorative effects mediated by

astrocytes, glia, microglia, neurons, endothelium cells or other cells of the brain and/or CNS, comprising administering to a patient in need thereof an effective amount of a disclosed compound.

[0037] Astrocytes play key roles in supporting and protecting neurons and they critically affect the outcome of brain injuries that cause brain damage, such as ischemic injuries. The central role astrocyte mitochondria themselves play in these brain functions is less well appreciated. For example, inhibition of astrocyte mitochondria increases swelling and leads to necrotic cell death. Neurons are permanently injured by recurrent spreading depolarizations only if astrocyte mitochondrial function fails, and astrocyte mitochondria are required for reduction of pathophysiological elevations of extracellular K^+ , which initiate spreading depolarizations. Activation of purinergic receptors on astrocytes results in increased mitochondrial Ca^{2+} that enhances mitochondrial citric acid cycle function and increases respiration and ATP production. Accordingly, in one aspect, the present invention relates to the discovery that activation of astrocyte purinergic receptors enhances brain cell survival signaling pathways, enabling both astrocyte and neuronal viability during oxidative stress. Furthermore, activated astrocytes generate and supply reduced glutathione, a key antioxidant that aids in the resistance of both astrocytes and neurons to oxidative stress. Thus, in one aspect, the present invention provides a method of modulating astrocyte purinergic receptors to promote survival and viability of one or more cell types in the brain of a patient after oxidative stress, such as oxidative stress caused by a brain injury, ischemia-reperfusion or a neurodegenerative condition, comprising administering to a patient in need thereof a disclosed compound.

[0038] In some embodiments, activation of astrocytes is achieved through contacting with a disclosed compound one or more purinergic receptors such as adenosine receptors (ARs), for example those associated with or expressed by astrocytes, thus modulating the activity of the one or more receptors. In some embodiments, through effects on adenosine receptors such as A_1 , A_{2A} , A_{2B} and A_3 on astrocytes, the compound activates astrocytes to treat one or more disclosed diseases or conditions. In some embodiments, after administration to a patient in need thereof, a disclosed compound influences one or more functions such as glutamate uptake, reactive gliosis, swelling, and release of neurotrophic and neurotoxic factors having an impact on metabolic stress and its

consequences, thus treating one or more diseases or conditions. In some embodiments, the compound is an AR agonist. In some embodiments, the purinergic receptor is an A₃ adenosine receptor (A₃R). In some embodiments, the compound is an A₃R agonist. In some embodiments, the compound is a partial agonist or biased agonist or biased partial agonist, at an A₃ receptor (A₃R), such as a human A₃ receptor (hA₃R). In some embodiments, the compound is a biased antagonist at an A₃ receptor. In some embodiments, the compound is MRS4322 or MRS1873 or a pharmaceutically acceptable salt thereof.

[0039] P2Y receptors are G-protein-coupled receptors and different subtypes of these receptors have important roles in processes such as synaptic communication, cellular differentiation, ion flux, vasodilation, blood brain barrier permeability, platelet aggregation and neuromodulation. Characterized members of the purinergic P2Y receptor family include the mammalian P2Y₁, P2Y₁₁, P2Y₁₂ and P2Y₁₃ receptors, which bind to adenine nucleotides; the P2Y₄, P2Y₆, and P2Y₁₄ receptors, that bind to uracil nucleotides; and the P2Y₂ and rodent P2Y₄ receptors, which have mixed selectivity. In some embodiments, activation of astrocytes is achieved through contacting with a disclosed compound one or more purinergic receptors such as P2Y receptors, for example those associated with or expressed by astrocytes, thus modulating the activity of the one or more receptors. In some embodiments, through effects on P2Y receptors such as P2Y₁, P2Y₁₁, P2Y₁₂ and P2Y₁₃ receptors associated with or expressed by astrocytes, the compound activates astrocytes to treat one or more disclosed diseases or conditions. In some embodiments, the P2Y receptor is a P2Y₁ receptor. In some embodiments, the P2Y₁ receptor is located on intracellular mitochondrial membranes. In some embodiments, the compound is a P2Y agonist. In some embodiments, the compound is a P2Y₁ agonist, e.g. at a human P2Y₁ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist at a P2Y₁ receptor, such as a human P2Y₁ receptor. In some embodiments, the compound is a biased antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or a pharmaceutically acceptable salt thereof.

[0040] In another aspect, the present invention provides a method of treating or ameliorating a brain injury, such as a brain injury resulting from a TBI or progressive

neurodegenerative disorder, in a patient in need thereof, comprising administering to the patient an effective amount of a disclosed compound. In some embodiments, the subject has suffered a TBI, concussion, stroke, partial or total spinal cord transection, or malnutrition. In other embodiments, the subject has suffered toxic neuropathies, meningoencephalopathies, neurodegeneration caused by a genetic disorder, age-related neurodegeneration, or a vascular disease; or another disease disclosed in US 8,691,775, which is hereby incorporated by reference. In some embodiments, the present invention provides a method of treating or ameliorating a brain injury, such as a brain injury resulting from a TBI or progressive neurodegenerative disorder, in a patient in need thereof, comprising administering to the patient an effective amount of an A₃R agonist. In other embodiments, the present invention provides a method of treating or ameliorating a brain injury, such as a brain injury resulting from a TBI or progressive neurodegenerative disorder, in a patient in need thereof, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

[0041] In another aspect, the present invention provides a method of promoting astrocyte-mediated neuroprotection or neurorestoration in a patient in need thereof, comprising administering to the patient an effective amount of a disclosed compound. In some embodiments, the present invention provides a method of promoting astrocyte-mediated neuroprotection or neurorestoration in a patient in need thereof, comprising administering to the patient an effective amount of an A₃R agonist. In other embodiments, the present invention provides a method of promoting astrocyte-mediated neuroprotection or neurorestoration in a patient in need thereof, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

[0042] In another aspect, the present invention provides a method of promoting survival of neurons, glial cells, endothelial cells or other brain cells, such as those in an ischemic penumbra in a patient in need thereof, comprising administering to the patient an effective amount of a compound disclosed herein. In some embodiments, the present invention provides a method of promoting survival of neurons, glial cells, or other brain cells, such as those in an ischemic penumbra in a patient in need thereof, comprising administering to the patient an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of promoting survival of neurons, glial cells, endothelial cells or other brain cells, such as those in an ischemic penumbra in a patient in need thereof, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

[0043] In further embodiments, the patient has or is at risk of acquiring a brain injury such as those below. Accordingly, methods of treating the conditions discussed below are also provided.

Traumatic Brain Injuries

[0044] Traumatic brain injuries (TBI) are a distressingly common medical condition and are predicted to become the third major cause of global morbidity and mortality by 2020. There are no approved treatments for TBI, and most TBI patients are discharged from the hospital with no pharmacological treatment (Witt 2006). Repetitive TBI such as concussions can trigger age-associated neurodegeneration that results in a range of symptoms and disabilities over decades (McKee 2013). TBIs can happen through sports-related injuries, motor vehicle accidents, falls, explosive impacts, physical assaults, etc. Injuries range widely in their complexity and severity, from “mild” concussions with brief alterations in mental status, cognitive difficulties, or loss of consciousness to “severe” with prolonged periods of unconsciousness and/or amnesia after the injury. In the U.S., approximately 1.7 million people have an injury resulting in a TBI annually and seek medical intervention (USCSF and CDC), and the CDC estimates that 1.6 to 3.8

million additional concussion incidents occur in sports and other recreational pursuits annually that do not present to hospital or emergency departments. (CDC; Langlois 2006) Approximately 5-10% of athletes will receive a concussion each sport season. (Sports Concussion Institute 2012) Football is the sport with the highest concussion risk for males (75% chance for concussion), while soccer has the highest concussion risk for females (50% chance for concussion). TBI is the leading cause of death and disability in children and young adults (CDC) and the most commonly received military-related injury; approximately 20% of U.S. Service Members deployed since 2003 have sustained at least one TBI. (Chronic Effects of Neurotrauma Consortium (CENC); Warden 2006; Scholten 2012; Taylor 2012; Gavett 2011; Guskiewicz 2005; Omalu 2005) Total TBI-related indirect and direct medical costs are estimated at \$77 billion annually (UCSF and CDC). At least 5 million Americans require ongoing daily support in performing activities as a result of TBI (CDC and Thurman 1999).

[0045] Activation of astrocytes according to the present invention represents a new treatment option for such conditions. Accordingly, provided herein in one aspect is a method of treating TBI or promoting recovery from TBI, comprising administering to a patient in need thereof an effective amount of a disclosed compound. In some embodiments, the TBI is selected from traumatic injuries to the brain (such as concussion, blast injury, combat-related injury) or spinal cord (such as partial or total spinal cord transection). In some embodiments, the TBI results from a mild, moderate, or severe blow to the head, comprises an open or closed head wound, or results from a penetrating or non-penetrating blow to the head. In some embodiments, the present invention provides a method of treating TBI or promoting recovery from TBI, comprising administering to a patient in need thereof an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating TBI or promoting recovery from TBI, comprising administering to a patient in need thereof an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

Stroke

[0046] A stroke occurs when a blood vessel that transports oxygen and nutrients to the brain is disrupted due to an ischemic blockage or from the hemorrhagic rupture of a blood vessel in the brain, causing neurons, glia and endothelial cells in the disrupted region of the brain to die. The outcome of the stroke depends upon the location and breadth of damage, and the impacts of that damage are observed in the body functions regulated by the damaged brain region. Strokes can cause unilateral or bilateral paralysis, speech and language disabilities, memory loss, behavioral changes, and even death. Stroke is the fourth leading cause of death in the United States and is a major cause of adult disability. Each year, ~800,000 people experience a new or recurrent stroke. Each day, over 2000 Americans will have a stroke, resulting in death in over 400 of these incidents. Stroke accounted for ~1 of every 19 deaths in the United States in 2010. An estimated 6.8 million Americans ≥ 20 years of age has had a stroke. (AHA and Go 2014) As of 2010, the annual direct and indirect cost of stroke was estimated at \$36.5 billion. Within minutes of a stroke, the lack of blood flow will permanently damage a core of brain tissue. Between this damaged core and normal brain tissue is a region of tissue known as the penumbra – tissue that is under gradated stress from lessened blood flow and some disruption of energy metabolism. Over the first 24-48 hours following a stroke incident, the stress on neuronal and glia cells in the penumbra resolves either with some recovery or further cell death.

[0047] In one aspect, the present invention provides a method of neuroprotective therapy in a stroke patient, comprising administering to a patient in need thereof an effective amount of a disclosed compound. In some embodiments, such therapy salvages as much of the penumbra as possible, and/or limits further acute tissue damage, and/or promotes neuron recovery. In another aspect is provided a method of treating stroke or promoting recovery from stroke, comprising administering to a patient in need thereof an effective amount of a disclosed compound. In another aspect is provided a method of treating stroke or promoting recovery from stroke, comprising administering to a patient in need thereof an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating stroke or promoting recovery from stroke, comprising administering to a patient in need thereof an effective amount of a P2Y₁

agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or a pharmaceutically acceptable salt thereof.

[0048] In some embodiments, the stroke is selected from selected from ischemic stroke, hemorrhagic stroke, subarachnoid hemorrhage, cerebral vasospasm, or transient ischemic attacks (TIA). In some embodiments, the stroke is ischemic. In some embodiments, the stroke is hemorrhagic. In some embodiments, the compound is administered within 48 hours of the stroke. In some embodiments, the compound is administered within 24 hours of the stroke. In some embodiments, the compound is administered within 16 hours of the stroke. In some embodiments, the compound is administered within 8, 4, 2, or 1 hours of the stroke. In some embodiments, the compound is administered for at least the first 1-72 hours following the stroke. In some embodiments, the compound is administered for at least the first 8-52 hours following the stroke. In some embodiments, the compound is administered for at least the first 8-48 hours following the stroke. In some embodiments, the compound is administered for at least the first 24-48 hours following the stroke. In some embodiments, the compound is administered chronically to treat the stroke as it occurs. In some embodiments, the compound is administered chronically to treat Transient Ischemic Attacks (TIA).

[0049] In some embodiments, the compound is administered chronically to treat ischemic stroke, hemorrhagic stroke, a subarachnoid hemorrhage, cerebral vasospasm, transient ischemic attacks (TIA), or treat a patient who is at an increased risk for a stroke, such as a patient who has had a stroke in the past and is at risk for a further stroke, such as a patient over the age of 40, 45, 50, 55, 60, 65, 70, 75, or 80 years of age.

[0050] In some embodiments, the compound treats an ischemia-reperfusion injury caused by the stroke.

Neurodegenerative Diseases

[0051] Neurodegenerative diseases are incurable, progressive, and ultimately debilitating syndromes resulting from the progressive degeneration and/or death of neurons in the brain and spinal cord. Neurodegeneration results in movement (ataxias)

and/or cognitive function (dementias) disorders, and includes a spectrum of diseases such as Alzheimer's Disease (AD), Parkinson's Disease (PD), Huntington's Disease (HD), Multiple Sclerosis (MS), amyotrophic lateral sclerosis (ALS), and chronic traumatic encephalopathy (CTE). While many neurodegenerative diseases are principally genetic in origin, other causes can include viruses, alcoholism, tumors or toxins, and as is now clear, repetitive brain injuries.

[0052] Neurons accumulate cellular damage over time due to the foregoing factors, which is generally considered the reason why many neurodegenerative diseases associated with prolonged cellular stress, such as Alzheimer's disease and Parkinson's disease, occur in aged individuals. Dementias represent the predominant outcome of neurodegenerative diseases with AD representing approximately 60-70% of cases. (Kandale 2013) As discussed above, activation of neuroprotective and neurorestorative mechanisms can ameliorate the progression of one or more neurodegenerative diseases. Accordingly, in one aspect the present invention provides a method of treating a neurodegenerative disease or promoting recovery from a neurodegenerative disease, comprising administering to a patient in need thereof an effective amount of a disclosed compound.

[0053] In one aspect, the present invention provides a method of promoting neuroprotection or neurorestoration in a patient suffering from a neurodegenerative disease, comprising administering to the patient an effective amount of a disclosed compound. In some embodiments is provided a method of promoting neuroprotection or neurorestoration in a patient suffering from a neurodegenerative disease, comprising administering to the patient an effective amount of an A₃R agonist. In other embodiments is provided a method of promoting neuroprotection or neurorestoration in a patient suffering from a neurodegenerative disease, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or a pharmaceutically acceptable salt thereof.

Alzheimer's Disease (AD)

[0054] An estimated 5.2 million Americans of all ages had AD in 2014; 11% of the population age 65 and older have AD. (Alzheimer's Association) By 2050, the number of people age 65 and older with AD is projected to nearly triple to a projected 13.8 million. In the U.S., the cost of providing care for AD patients is about \$214 billion per year; 70% of this cost is covered by Medicare and Medicaid . The current trends would project these costs to grow to \$1.2 trillion per year by 2050.

[0055] Activation of astrocytes and promoting neuroprotection and neurorestoration according to the present invention represents a new treatment option for AD. Accordingly, provided herein in one aspect is a method of treating AD or promoting neuroprotection or neurorestoration in a patient suffering from AD, comprising administering to the patient an effective amount of a compound disclosed herein. In some embodiments, the present invention provides a method of treating AD or promoting neuroprotection or neurorecovery in a patient suffering from AD, comprising administering to the patient an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating AD or promoting neuroprotection or neurorecovery in a patient suffering from AD, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or a pharmaceutically acceptable salt thereof.

Parkinson's Disease (PD)

[0056] As many as one million Americans live with PD, and each year approximately 60,000 Americans are newly diagnosed not including the thousands of cases that go undetected. (Parkinson's Disease Foundation) The total combined direct and indirect cost of PD, including medical treatment, social security payments and lost income, is estimated to be nearly \$25 billion per year in the United States. (Parkinson's Disease Foundation and Huse 2005)

[0057] Activation of neuroprotection and neurorestoration according to the present invention represents a new treatment option for PD. Accordingly, provided herein in one aspect is a method of treating PD or promoting neuroprotection or neurorestoration in a

patient suffering from PD, comprising administering to the patient an effective amount of a disclosed compound. In some embodiments, the present invention provides a method of treating PD or promoting neuroprotection or neurorecovery in a patient suffering from PD, comprising administering to the patient an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating PD or promoting neuroprotection or neurorecovery in a patient suffering from PD, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

Multiple Sclerosis (MS)

[0058] More than 400,000 people in the United States have MS. In young adults, MS represents the most prevalent disease of the central nervous system. (Multiple Sclerosis Foundation) There is potential for astrocytes to reverse the destruction of nerve cell myelin coatings that is caused by MS by their neurorestorative effects and promotion of healing in the damaged CNS of MS patients.

[0059] Activation of neuroprotection and neurorestoration in the CNS according to the present invention thus represents a new treatment option for MS. Accordingly, provided herein in one aspect is a method of treating MS or promoting neuroprotection or neurorestoration in a patient suffering from MS, comprising administering to the patient an effective amount of a disclosed compound. In some embodiments, the present invention provides a method of treating MS or promoting neuroprotection or neurorecovery in a patient suffering from MS, comprising administering to the patient an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating MS or promoting neuroprotection or neurorecovery in a patient suffering from MS, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a

P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

Amyotrophic Lateral Sclerosis (ALS) / Lou Gehrig's Disease

[0060] Approximately 5,600 people in the U.S. are diagnosed with ALS each year; as many as 30,000 Americans may have the disease concurrently. (ALS Association) Activation of astrocytes can provide stimulation of recovery and repair of the neurons and their connections in an ALS patient.

[0061] Accordingly, provided herein in one aspect is a method of treating ALS or promoting neuroprotection or neurorestoration in a patient suffering from ALS, comprising administering to the patient an effective amount of a disclosed compound. Also provided in other embodiments is a method of stimulating recovery and repair of the neurons and their connections in an ALS patient, comprising administering to the patient an effective amount of a compound disclosed herein. In some embodiments, the present invention provides a method of treating ALS or promoting neuroprotection or neurorecovery in a patient suffering from ALS, comprising administering to the patient an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating ALS or promoting neuroprotection or neurorecovery in a patient suffering from ALS, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or a pharmaceutically acceptable salt thereof.

Chronic Traumatic Encephalopathy (CTE)

[0062] CTE (a form of tauopathy) is a progressive neurodegenerative disease found in individuals who have suffered one or more (often multiple, or repeated over the course of time) severe blows to the head. CTE is most often diagnosed in professional athletes in American football, soccer, hockey, professional wrestling, stunt performing, bull riding and rodeo performing, motocross, and other contact sports who have experienced brain trauma and/or repeated concussions. A subset of CTE sufferers have chronic traumatic encephalomyopathy (CTEM), which is characterized by motor neuron disease

symptoms that mimic ALS. Progressive muscle weakness and motor and gait abnormalities are believed to be early signs of CTEM. First stage symptoms of CTE include progressive attention deficit, disorientation, dizziness, and headaches. Second stage symptoms comprise memory loss, social instability, erratic behavior, and poor judgment. In third and fourth stages, patients suffer progressive dementia, slowed movements, tremors, hypomimia, vertigo, speech impediments, hearing loss, and suicidality, and may further include dysarthria, dysphagia, and ocular abnormalities, e.g. ptosis.

[0063] Accordingly, provided herein in one aspect is a method of treating or preventing CTE or promoting neuroprotection or neurorestoration in a patient suffering from CTE, comprising administering to the patient an effective amount of a disclosed compound. Also provided in other embodiments is a method of stimulating recovery and repair of the neurons and their connections in a CTE patient, comprising administering to the patient an effective amount of a disclosed compound. In some embodiments, the compound treats one or more symptoms of first stage, second stage, third stage, or fourth stage CTE. In some embodiments, the present invention provides a method of treating CTE or promoting neuroprotection or neurorecovery in a patient suffering from CTE, comprising administering to the patient an effective amount of an A₃R agonist. In some embodiments, the present invention provides a method of treating CTE or promoting neuroprotection or neurorecovery in a patient suffering from CTE, comprising administering to the patient an effective amount of a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873 or a pharmaceutically acceptable salt thereof.

[0064] On a microscopic scale the pathology includes neuronal death, tau deposition, TAR DNA-binding Protein 43 (TDP 43) beta-amyloid deposition, white matter changes, and other abnormalities. Tau deposition includes the increasing presence of dense neurofibrillary tangles (NFT), neurites, and glial tangles, which are made up of astrocytes and other glial cells. Thus, in some embodiments, the method treats, enhances clearance

or prevents neuronal death, tau deposition, TAR DNA-binding Protein 43 (TDP 43) beta-amyloid deposition, white matter changes, and other abnormalities associated with CTE.

[0065] In some embodiments, the present invention provides long-term administration of a compound disclosed herein, such as a biased agonist, partial agonist, or biased partial agonist of A₃R, or a biased agonist, partial agonist, or biased partial agonist of P2Y₁, to treat a neurodegenerative disease, such as those discussed above and below.

Cardiovascular Diseases

[0066] Disclosed compounds are also useful in treating a variety of cardiovascular diseases and conditions. In some embodiments, the present invention provides a method of treating a heart or cardiovascular disease, such as cardiac ischemia, myocardial infarction, a cardiomyopathy, coronary artery disease, arrhythmia, myocarditis, pericarditis, angina, hypertensive heart disease, endocarditis, rheumatic heart disease, congenital heart disease, or atherosclerosis, comprising administering an effective amount of a disclosed compound to a patient in need thereof, such as MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof. In some embodiments, a disclosed compound provides for modulation of ATP-sensitive potassium channels, for example via biased agonism, partial agonism, or biased partial agonism at an A₃R receptor.

[0067] In some embodiments, the heart or cardiovascular disease is cardiac ischemia or myocardial infarction.

Other Diseases

[0068] Compounds that modulate beneficial effects such as neuroprotection, for example by increasing astrocyte mitochondrial activity, also have the potential to treat a variety of other diseases. For example, due to the role of astrocytes in neuroprotection disclosed in the present invention, activation of astrocytes, for example via modulation of A₃R and/or a P2Y₁ receptor, may be useful in treating various diseases and conditions discussed below. Accordingly, in some embodiments, the present invention provides a method of treating or promoting neuroprotection or neuroregeneration in a patient suffering from a disease or condition, comprising administering to the patient an effective amount of a disclosed compound, for example MRS4322 or a pharmaceutically

acceptable salt thereof. In some embodiments, the disease or condition is selected from autoimmune diseases, allergic diseases, and/or transplant rejection and graft-versus-host disease (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, WO 2007/20018, hereby incorporated by reference). In other embodiments, the disease or condition is selected from intraocular hypertension and/or glaucoma (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, WO 2011/77435, hereby incorporated by reference). In other embodiments, the disease or condition is selected from odor sensitivity and/or an olfactory disorder (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, EP1624753, hereby incorporated by reference). In other embodiments, the disease or condition is selected from type 2 diabetes and/or pain control (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, US 2010/0256086, hereby incorporated by reference).

[0069] In other embodiments, the disease or condition is selected from respiratory diseases and/or cardiovascular (CV) diseases (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, FASEB J. (2013) 27:1118.4 (abstract of meeting), hereby incorporated by reference). In other embodiments, the disease or condition is selected from deficits in CNS function, deficits in learning and/or deficits in cognition (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, Neuropsychopharmacology. 2015 Jan;40(2):305-14. doi: 10.1038/npp.2014.173. Epub 2014 Jul 15. "Impaired cognition after stimulation of a P2Y₁ receptor in the rat medial prefrontal cortex," Koch, H. *et al.* PMID: 25027332, hereby incorporated by reference). In other embodiments, the disease or condition is selected from a neurodegenerative disease such as Alzheimer's disease, Parkinson's disease, Huntington's disease, prion disease, and/or amyotrophic lateral sclerosis (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, US 8,691,775, hereby incorporated by reference). In other embodiments, the disease or condition is selected from otic disorders, Meniere's disease, endolymphatic hydrops, progressive hearing loss, dizziness, vertigo, tinnitus, collateral brain damage associated with radiation cancer therapy, and/or migraine treatment (for the use of certain nucleoside and nucleotide compounds in treating these

conditions, see, for example, US 2009/0306225; UY31779; and US 8,399,018, each of which is hereby incorporated by reference). In other embodiments, the disease or condition is selected from pathological sleep perturbations, depression, sleep disorders in the elderly, Parkinson's disease, Alzheimer's disease, epilepsy, schizophrenia, and/or symptoms experienced by recovering alcoholics (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, US 2014/0241990, hereby incorporated by reference). In other embodiments, the disease or condition is selected from damage to neurons or nerves of the peripheral nervous system during surgery (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, US 8,685,372, hereby incorporated by reference). In other embodiments, the disease or condition is a cancer such as prostate cancer (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, *Biochem Pharmacol.* 2011 August 15; 82(4): 418–425. doi:10.1016/j.bcp.2011.05.013. “Activation of the P2Y1 Receptor Induces Apoptosis and Inhibits Proliferation of Prostate Cancer Cells,” Qiang Wei et al., hereby incorporated by reference). In other embodiments, the disease or condition is selected from one or more gastrointestinal conditions such as constipation and/or diarrhea (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, *Acta Physiol (Oxf)*. 2014 Dec;212(4):293-305. doi: 10.1111/apha.12408. “Differential functional role of purinergic and nitrergic inhibitory cotransmitters in human colonic relaxation,” Mañé N1, Gil V, Martínez-Cutillas M, Clavé P, Gallego D, Jiménez M.; and *Neurogastroenterol. Motil.* 2014 Jan;26(1):115-23. doi: 10.1111/nmo.12240. Epub 2013 Oct 8. “Calcium responses in subserosal interstitial cells of the guinea-pig proximal colon,” Tamada H., Hashitani H. PMID: 24329947, hereby incorporated by reference). In other embodiments, the disease or condition is selected from pain mediated by the CNS, such as neuropathic pain, inflammatory pain, and/or acute pain (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, *Br J Pharmacol.* 2010 Mar;159(5):1106-17. doi: 10.1111/j.1476-5381.2009.00596.x. Epub 2010 Feb 5. “A comparative analysis of the activity of ligands acting at P2X and P2Y receptor subtypes in models of neuropathic,

acute and inflammatory pain.” Andó RD1, Méhész B, Gyires K, Illes P, Sperlágh B. PMID: 20136836), hereby incorporated by reference).

[0070] In other embodiments, the disease or condition is selected from cancer of the brain, such as glioblastoma (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, *Purinergic Signal*. 2015 Sep;11(3):331-46. doi: 10.1007/s11302-015-9454-7. Epub 2015 May 15. “Potentiation of temozolomide antitumor effect by purine receptor ligands able to restrain the *in vitro* growth of human glioblastoma stem cells.” D’Alimonte, I. *et al.* PMID: 25976165, hereby incorporated by reference). In other embodiments, the disease or condition is pain (for the use of certain nucleoside and nucleotide compounds in treating pain, see, for example, *Pharmacol Biochem Behav*. 2015 Jan;128:23-32. doi: 10.1016/j.pbb.2014.11.001. Epub 2014 Nov 6. “Participation of peripheral P2Y₁, P2Y₆ and P2Y₁₁ receptors in formalin-induced inflammatory pain in rats.” Barragán-Iglesias P. *et al.* PMID: 25449358; and *Neuropharmacology*. 2014 Apr;79:368-79. doi: 10.1016/j.neuropharm.2013.12.005. Epub 2013 Dec 12. “Blockade of peripheral P2Y₁ receptors prevents the induction of thermal hyperalgesia via modulation of TRPV1 expression in carrageenan-induced inflammatory pain rats: involvement of p38 MAPK phosphorylation in DRGs.” Kwon SG, Roh DH, Yoon SY, Moon JY, Choi SR, Choi HS, Kang SY, Han HJ, Beitz AJ, Lee JH. PMID: 24333674, each of which is hereby incorporated by reference). In other embodiments, the disease or condition is selected from a gastrointestinal disorder such as diarrhea (for the use of certain nucleoside and nucleotide compounds in treating these conditions, see, for example, *Acta Physiol (Oxf)*. 2014 Dec;212(4):293-305. doi: 10.1111/apha.12408. “Differential functional role of purinergic and nitrergic inhibitory cotransmitters in human colonic relaxation,” Mañé N., Gil V, Martínez-Cutillas M, Clavé P, Gallego D, Jiménez M., hereby incorporated by reference). In other embodiments, the disease or condition is impaired cognition (for the use of certain nucleoside and nucleotide compounds in treating this condition, see, for example, *Neuropsychopharmacology*. 2015 Jan;40(2):305-14. doi: 10.1038/npp.2014.173. Epub 2014 Jul 15. “Impaired cognition after stimulation of P2Y₁ receptors in the rat medial prefrontal cortex,” Koch H, Beshpalov A, Drescher K, Franke H, Krügel U. PMID: 25027332, hereby incorporated by reference).

[0071] In some embodiments, the present invention provides a method of treating a disease or condition associated with brain injury or a neurodegenerative condition, such as epilepsy, migraine, collateral brain damage associated with radiation cancer therapy, depression, mood or behavioral changes, dementia, erratic behavior, suicidality, tremors, Huntington's chorea, loss of coordination of movement, deafness, impaired speech, dry eyes, hypomimia, attention deficit, memory loss, cognitive difficulties, vertigo, dysarthria, dysphagia, ocular abnormalities, or disorientation, comprising administering to a patient in need thereof an effective amount of a disclosed compound. In some embodiments, the compound is an A₃R agonist. In some embodiments, the compound is a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

[0072] In further embodiments, the present invention provides a method of treating a neurodegenerative disease selected from the group consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple sclerosis, amyotrophic lateral sclerosis, and prion disease in a patient in need thereof, comprising administering an effective amount of a disclosed compound. In some embodiments, the compound is an A₃R agonist. In some embodiments, the compound is a P2Y₁ agonist. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist or antagonist at a P2Y₁ receptor. In some embodiments, the compound is MRS4322 or MRS1873, or a pharmaceutically acceptable salt thereof.

[0073] In some embodiments, the improvement in cognitive or neurological function is measured as a score increase between about 1% and 20% in the delayed verbal recall task of the revised Wechsler Memory Scale. For example, the improvement in cognitive function may be measured as a score increase between about 1% and 10%, or between about 1% and 5%.

2. Description of Certain Compounds of the Present Invention

[0074] In one aspect, the present invention provides compounds useful for treating, ameliorating, or promoting recovery from certain conditions of the brain or central nervous system (CNS) such as a brain injury or a neurodegenerative condition. In some embodiments, a disclosed compound increases neuroprotection and neuroregeneration mediated by astrocytes, thereby treating, ameliorating, or promoting recovery from the condition. In some embodiments, the compound is selective for an A₃ receptor, for example selective for an A₃ receptor by at least 10-fold relative to other adenosine receptors; or for example more than 25-fold, 50-fold, 100-fold, 500-fold, or 1000-fold relative to other adenosine receptors. In some embodiments, the compound selectively modulates an A₃ receptor. In some embodiments, the compound is a selective agonist at an A₃ receptor. In some embodiments, the compound is a selective partial agonist at an A₃ receptor. In some embodiments, the compound is a biased full or partial agonist. In some embodiments, the compound is a biased full or partial antagonist.

[0075] In further embodiments, the compound is selective for a P2Y₁ receptor, for example selective for P2Y₁ receptors by at least 10-fold relative to other P2Y receptors; or for example more than 25-fold, 50-fold, 100-fold, 500-fold, or 1000-fold relative to other P2Y receptors. In some embodiments, the compound selectively modulates a P2Y₁ receptor. In some embodiments, the compound is a selective agonist at a P2Y₁ receptor. In some embodiments, the compound is a selective partial agonist at a P2Y₁ receptor. In some embodiments, the compound is a biased full or partial agonist. In some embodiments, the compound is a biased full or partial antagonist.


[0076] The term “biased” refers to compounds that preferentially modulate, activate, agonize, or antagonize one or more, but not all, of the pathways associated with a receptor.


[0077] Without wishing to be bound by theory, it is believed that biased full or partial agonism or antagonism allows for selective modulation of one or more pathways linked to an A₃ or P2Y₁ receptor, which may lead to improved treatment of a disease or condition and avoidance of undesired pathway modulation (which would lead to side effects). Selective modulation may preferentially activate astrocytes as disclosed herein, for example to treat a brain injury or neurodegenerative disease or condition. Accordingly, in some embodiments, a disclosed compound is a biased full or partial


agonist or antagonist of one or more G-coupled or G-independent pathways linked to the adenosine A₃ receptor or P2Y₁ receptor. In some embodiments, the compound selectively modulates a pathway mediated by A₃ or a P2Y₁ receptor, such as beta-arrestin activation, intracellular calcium mobilization, cAMP modulation, ATP-dependent potassium channel activation, or ERK1/2 phosphorylation, or other downstream cellular activities associated with such pathways. In some embodiments, the pathway increases or is related to neuroprotection or neurorestoration, or cardioprotection or cardioregeneration. In some embodiments, the compound is selected from a (N)-methanocarba nucleoside such as MRS4322; or a pharmaceutically acceptable salt thereof.

[0078] The term “methanocarba nucleoside” as used herein refers to a nucleoside analog in which the oxygen present in the tetrahydrofuran ring of the ribose sugar is replaced with a methylene unit and the resulting carbocyclic ring is fused to a


cyclopropyl ring to form a bicyclo[3.1.0]hexane, such as the structures


or . Without being bound by theory, it is believed that methanocarba nucleosides mimic a sugar conformation or pseudoconformation believed to be favored by certain receptor subtypes. In some embodiments, North methanocarba nucleosides are those that mimic or prefer a C3'-endo/C2'-exo sugar conformation and South methanocarba nucleosides are those that mimic or prefer a C3'-exo/C2'-endo conformation. In some embodiments, a (N)-methanocarba (“North” methanocarba) sugar

has the following structure: . In some embodiments, a (N)-methanocarba

sugar has the following structure: , referred to herein as a “D-(N)-

methanocarba sugar.” In other embodiments, a methanocarba sugar is in the South, or (S)-methanocarba, configuration. In some embodiments, such methanocarba sugars are

represented by the structure: . In some embodiments, a (S)-

methanocarba sugar has the following structure: , referred to herein as a “D-(S)-methanocarba sugar.”

[0079] In some embodiments, the compound is functionally selective at the A₃ or P2Y₁ receptor, i.e., selectively discriminates among pathways mediated by A₃ or a P2Y₁ receptor, for example by modulating one or more pathways but not others, or by activating one or more pathways and deactivating one or more other pathways. In some embodiments, the compound is an antagonist as measured by cAMP signaling, but a partial agonist for β-arrestin recruitment. In other embodiments, the compound is an agonist of Gq/11-mediated Ca²⁺ mobilization and a partial agonist or antagonist of arrestin recruitment. In some embodiments, the present invention provides a method of treating a brain injury or neurodegenerative disease or condition via biased or functionally selective A₃ receptor modulation (e.g., by selective agonism or antagonism among pathways such as those mentioned above), comprising administering an effective amount of a disclosed compound to a patient in need thereof. In some embodiments, the compound is selected from DMPA, CCPA, MRS1760, or MRS542 (see Verzijl D, *et al.*, “Functional selectivity of adenosine receptor ligands,” *Purinergic Signaling* 7: 171-192 (2011)). In some embodiments, the compound is DBXRM. In some embodiments, the compound is selected from a (N)-methanocarba nucleoside such as MRS4322; or a pharmaceutically acceptable salt thereof.

[0080] It has been surprisingly found that certain purine nucleoside mono-, di-, and tri-phosphates, such as those described in detail herein, are rapidly dephosphorylated *in vivo*, possibly by ectonucleotidases, enzymes responsible for the dephosphorylation of nucleotides that are present both on the surface of cell membranes and circulating in blood and plasma (See Ziganshin et al. *Pflugers Arch.* (1995) 429:412-418). It is often

extremely difficult to predict which nucleotide analogs will be substrates for ectonucleotidases and will thus be expected to be dephosphorylated *in vivo*. In some embodiments, the dephosphorylated compound is responsible for the therapeutic efficacy. Thus, in some embodiments the corresponding, phosphorylated mono-, di-, or tri-phosphate, or a phosphate ester such as an alkyl or phenyl ester thereof, is a prodrug or precursor to the agent responsible for the therapeutic effect.

[0081] In some embodiments, compounds of the present invention are able to cross the blood-brain barrier (BBB). The term “blood-brain barrier” or “BBB”, as used herein, refers to the BBB proper as well as to the blood-spinal barrier. The blood-brain barrier, which consists of the endothelium of the brain vessels, the basal membrane and neuroglial cells, acts to limit penetration of substances into the brain. In some embodiments, the brain/plasma ratio of total drug is at least approximately 0.01 after administration (e.g. oral or intravenous administration) to a patient. In some embodiments, the brain/plasma ratio of total drug is at least approximately 0.03. In some embodiments, the brain/plasma ratio of total drug is at least approximately 0.06. In some embodiments, the brain/plasma ratio of total drug is at least approximately 0.1. In some embodiments, the brain/plasma ratio of total drug is at least approximately 0.2.

[0082] Prototypical adenosine A₃ agonists such as Cl-IB-MECA and MRS5698 are low-solubility, lipophilic compounds with cLogP values typically >2. This lipophilicity is a major factor contributing to these compounds’ high plasma protein binding, high brain binding and resulting low free fraction of drug available to interact with the A₃ receptor in the brain. In some embodiments, for example neurological and neurodegenerative conditions, the physicochemical properties of compounds of the present invention such as MRS4322 and MRS1873 are substantially different; these and related compounds are hydrophilic compounds with cLogP <0, resulting in high solubility, low plasma and brain binding and high unbound drug concentrations available to interact with the A₃ receptor.

[0083] Accordingly, in some embodiments the compound has a cLogP less than about 0.8, about 0.7, about 0.6, about 0.5, about 0.4, about 0.3, about 0.2, about 0.1, about 0.05, about 0.01, or about 0.005. In some embodiments, the compound has a cLogP less than about 0, such as less than about -0.1, -0.2, -0.3, -0.4, -0.5, -0.6, -0.7, -0.8, or -0.9 or

less. In some embodiments, the compound has an unbound fraction in plasma of about 0.5 to 0.9. In some embodiments, the compound has an unbound fraction in plasma of about 0.6 to 0.85, 0.7 to 0.8, or about 0.75. In some embodiments, the compound has an unbound fraction in brain of at least about 0.02, or at least about 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.12, 0.15, or 0.17 or greater. In some embodiments, the compound has an unbound fraction in plasma of about 0.6 to 0.85, 0.7 to 0.8, or about 0.75 and/or at least 0.08 unbound fraction in brain.

[0084] Compounds of the invention may be prepared using methods known in the art, using no more than routine experimentation. For example, certain compounds of the invention may be prepared following the procedures provided in U.S. Patent No. 7,087,589 (and references cited therein), which is hereby incorporated by reference.

[0085] In some embodiments, the compound is selected from adenosine, ADP, 2-methylthio-ADP trisodium salt, ATP, ATP disodium salt, α,β -methylene ATP, α,β -methyleneadenosine 5'-triphosphate trisodium salt, 2-methylthioadenosine triphosphate tetrasodium salt, 2-MeSATP, BzATP triethylammonium salt, inosine, cytidine, acylated cytidines, cytidine-monophosphate (CMP), cytidine diphosphate (CDP), cytidine triphosphate (CTP), CDP-choline, CMP-choline, denufosol, denufosol tetrasodium, GTP, ITP, MRS 541, MRS 542, MRS 1760, MRS 2179, MRS 2279, MRS 2341, MRS 2365, MRS 2500, MRS 2690, MRS 2698, MRS 3558, MRS 4322, MRS 5151, MRS 5676, MRS 5678, MRS 5697, MRS 5698, MRS 5923, MRS 5930, Benzyl-NECA, IB-MECA, Cl-IB-MECA, LJ529, DPMA, CCPA, DBXRM, HEMADO, PEMADO, HENECA, PENECA, CP608,039, CP532,903, CGS21680, AR132, VT72, VT158, VT160, VT163, PSB 0474, uridine 5'-diphosphate (UDP), UDP-glucose, uridine β -thiodiphosphate (UDP β S), uridine 5'-triphosphate (UTP), uridine γ -thiophosphate (UTP γ S), 2-thioUTP tetrasodium salt, UTP γ S trisodium salt, uridine-5'-diphosphoglucose, diuridine triphosphate, 2-(hexylthio) (HT)-AMP, diadenosine pentaphosphate, 2'-deoxy-2'-amino-UTP, 2-thio-UTP, triacetyluridine, diacetyl/acyl uridine, uridine, suramin, dipyrindamole analogs, diadenosine tetraphosphate Ap₄U, Ap₄A, INS365, INS37217, or INS48823; wherein each sugar may be replaced with a methanocarba sugar in the North or South conformation or each sugar may be replaced with a D-ribosugar; or a pharmaceutically acceptable salt thereof.










[0086] In some embodiments, 2-methylthio-ADP or a pharmaceutically acceptable salt thereof is useful in the methods of the present invention. Without wishing to be bound by theory, it is believed that 2-MeS ADP is rapidly hydrolyzed to 2-methylthioadenosine in vivo, where it is a biased agonist, partial agonist, or biased partial agonist of A₃R. 2-methylthioadenosine is believed to have receptor data very similar to that of MRS4322.





















[0087] In some embodiments, the compound is an A₃R agonist such as N⁶-benzyladenosine-5'-N-methyluronamides such as N⁶-(3-iodobenzyl)-adenosine-5'-N-methyluronamide, also known as IB-MECA or Can-Fite CF-101, or 2-Chloro-N⁶-(3-iodobenzyl)-adenosine-5'-N-methyluronamide (also known as 2-CI-IB-MECA or Can-Fite CF-102; (N)-methanocarba nucleosides such as (1R,2R,3S,4R)-4-(2-chloro-6-((3-chlorobenzyl)amino)-9H-purin-9-yl)-2,3-di-hydroxy-N-methylbicyclo[3.1.0]hexane-1-carboxamide (also known as CF502, Can-Fite Biopharma, MA); (2S,3S,4R,5R)-3-amino-5-[6-(2,5-dichlorobenzylamino)purin-9-yl]-4-hydroxy-tetrahydrofuran-2-carboxylic acid methylamide (also known as CP532,903); (1'S,2'R,3'S,4'R,5'S)-4-(2-chloro-6-(3-chlorobenzylamino)-9H-purin-9-yl)-2,3-dihydroxy-N-methylbicyclo[3.1.0]hexane-1-carboxamide (also known as MRS3558), 2-(1-hexynyl)-N-methyladenosine; (1S,2R,3S,4R)-2,3-dihydroxy-4-(6-((3-iodobenzyl)amino)-4H-purin-9(5H)-yl)-N-methylcyclopentanecarboxamide (also known as CF101, Can-Fite); (1S,2R,3S,4R)-4-(2-chloro-6-((3-iodobenzyl)amino)-4H-purin-9(5H)-yl)-2,3-dihydroxy-N-methylcyclopentanecarboxamide (also known as CF102, Can-Fite); (1'R,2'R,3'S,4'R,5'S)-4-{2-chloro-6-[(3-iodophenylmethyl)amino]purin-9-yl}-1-(methylaminocarbonyl)-bicyclo[3.1.0]hexane-2,3-diol (also known as MRS1898); or 2-dialkynyl derivatives of (N)-methanocarba nucleosides; or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is selected from IB-MECA (also known as CF101), or Cl-IB-MECA (also known as CF102); or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is selected from a (N)-methanocarba nucleoside such as those disclosed above; or a pharmaceutically acceptable salt thereof.


[0088] Also included are A₃R allosteric modulators which enhance the receptor activity in the presence of the native ligand, such as 2-cyclohexyl-N-(3,4-dichlorophenyl)-1H-imidazo[4,5-c]quinolin-4-amine (also known as CF602, Can-Fite).

However, the above-listed A₃R agonists are by no means exclusive and other such agonists may also be used. The administration of A₃R agonists covalently bound to polymers is also contemplated. For example, A₃R agonists may be administered in the form of conjugates where an agonist is bound to a polyamidoamine (PAMAM) dendrimer.

[0089] Without wishing to be bound by theory, it is believed that full or partial agonism, including biased agonism, by certain uridine analogs will allow for selective modulation of one or more pathways which may lead to improved treatment of a disclosed disease or condition and avoidance of undesired pathway modulation (which would lead to side effects). In some embodiments, selective modulation preferentially activates astrocytes or other glial cells such as microglia and oligodendrocytes to treat a disclosed brain injury or neurodegenerative disease or condition. Certain uridine analog compounds suitable for use in the present invention are disclosed in WO 2014/160502, which is hereby incorporated by reference in its entirety. In some embodiments, the compound is an A₃R agonist. In some embodiments, the compound is a P2Y₁ agonist. In some embodiments, the compound is a biased agonist at an adenosine receptor, such as an A₁, A_{2A}, A_{2B} or A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist at an A₃ receptor. In some embodiments, the compound is a biased agonist, partial agonist, or biased partial agonist at a P2Y₁ receptor. In some embodiments, the compound is selected from the group consisting of:









or a phosphorylated analog thereof; or a pharmaceutically acceptable salt thereof.

[0090] In some embodiments, the compound is selected from:




(see Beukers MW *et al.*, (2004) “New, non-adenosine, high-potency agonists for the human adenosine A2B receptor with an improved selectivity profile compared to the reference agonist N-ethylcarboxamidoadenosine,” *J. Med. Chem.*




Devine 2010 #7b


47(15):3707–3709);




Devine 2010



Devine 2010 #9a




Devine 2010 #9b



Devine 2010 #9c

(see Devine SM *et al.* “Synthesis and Evaluation of new A₃R agonists,” *Bioorg Med Chem* 18, 3078-3087. 2010; and Muller CE, Jacobson KA. “Recent Developments in adenosine


receptor ligands and their potential for novel drugs,” *Biochimica et Biophysica Acta*



1808, 1290-1308. 2011);

(see Ben DD *et al.* “Different


efficacy of adenosine and NECA derivatives at the human A3 receptor: Insight into the receptor activation switch,” *Biochem Pharm* 87, 321-331. 2014; and Camaioni E, Di Francesco E, Vittori S, Volpini R, Cristalli G. “Adenosine receptor agonists: synthesis and biological evaluation of the diastereoisomers of 2-(3-hydroxy-3-phenyl-1-propyn-1-



yl)NECA,” *Bioorg Med Chem* 1997;5:2267–75);

(see


Klotz, K.N. “2-Substituted N-ethylcarboxamidoadenosine derivatives as high-affinity agonists at human A3 adenosine receptors,” *Naunyn Schmiedebergs Arch Pharmacol.* 1999 Aug;360(2):103-8; and Cristalli G *et al.* (1995) “2-Aralkynyl and 2-heteroalkynyl derivatives of adenosine-5'-N-ethyluronamide as selective A2a adenosine receptor



agonists,” *J Med Chem* 38 :1462–1472);


(see Kim S *et al.*

“3D quantitative SAR at A3R,” *J Chem Inf. Model* 47, 1225-1233 2007);



(MRS1898);

(see Lee, K. *et al.* "Ring-Constrained (N)-Methanocarba Nucleosides as Adenosine Receptor Agonists," *Bioorg Med Chem Lett*




2001, 11, 1333-1337);


(see Kenneth A. Jacobson *et al.*

Chapter 6. A3 Adenosine Receptor Agonists: History and Future Perspectives pp 96-97.


Book – Springer: A3 Adenosine Receptors from Cell Biology to Pharmacology and




Therapeutics, 2009); (see Lee K *et al.* “Ring-Constrained (N)-Methanocarba Nucleosides as Adenosine Receptor Agonists,” *Bioorg Med Chem Lett* 2001, 11, 1333-1337; and Gao *et al.* “Structural Determinants of A3R Activation: Nucleoside Ligands at the Agonist/Antagonist Boundary,” *J. Med. Chem.*, 2002, 45,




4471-4484); $R = H$: IB-MECA (CF101)
 $R = Cl$: Cl-IB-MECA (CF102) (see Muller CE, Jacobson KA, “Recent Developments in adenosine receptor ligands and their potential for novel drugs,”




Biochimica et Biophysica Acta 2011, 1808, 1290-1308); (MRS5930; see Jacobson KA *et al.* “John Daly Lecture: Structure-guided Drug Design for Adenosine and P2Y Receptors,” *Comp. and Struct. Biotechnology Jour* 13. 286-298.



2015); (MRS5923; see Jacobson KA *et al.* “John Daly Lecture: Structure-guided Drug Design for Adenosine and P2Y Receptors,” *Comp. and*




Struct. Biotechnology Jour 13. 286-298. 2015); CP532,903 (see Tracey WR *et al.* "Novel n6-substituted adenosine 5'-N-methyluronamides with high selectivity for human A3R reduce ischemic myocardial injury," Am J Physiol Heart Circ Physiol 285. 2003; Muller CE, Jacobson KA, "Recent Developments in adenosine receptor ligands and their potential for novel drugs," Biochimica et Biophysica Acta 1808, 1290-1308. 2011; and Wan TC *et al.* "The A3R Agonist CP-532,903 Protects against Myocardial Ischemia/Reperfusion Injury," J. of Pharmacology and Exptl




R = $n\text{C}_4\text{H}_9$: HEMADO
R = Ph: PEMADO

Therapies 324,1. 2008); (see Volpini R *et al.* "HEMADO as Potent and Selective Agonists of hA3R," J Med Chem 45, 3271-3279. 2002; Muller CE *et al.* "Recent Developments in adenosine receptor ligands and their potential for novel drugs," Biochimica et Biophysica Acta 1808, 1290-1308. 2011; and Volpini R *et al.* "Synthesis and Evaluation of Potent and Highly Selective Agonists for hA3R," J of




MRS5151

Med Chem 52, 7897-7900. 2009);




LJ529

; (see Muller CE, Jacobson KA. "Recent Developments in adenosine receptor ligands and their potential for novel drugs,"




Biochimica et Biophysica Acta 1808, 1290-1308. 2011);



wherein R is H or cyclopentylmethyl;

wherein R is


H, butyl, or pyridin-2-yl (see Cosyn L. et al., "2-triazole-substituted adenosines," J Med



Chem 2006. 49. 7373-7383);

MRS3558 (CF502)

;




CP608,039

(see Muller CE, Jacobson KA. "Recent

Developments in adenosine receptor ligands and their potential for novel drugs,"


Biochimica et Biophysica Acta 1808, 1290-1308. 2011);



MRS5698

(see Jacobson KA et al. "John Daly Lecture:


Structure-guided Drug Design for Adenosine and P2Y Receptors," Comp. and Struct.





Biotechnology Jour 13. 286-298. 2015);

wherein Ar is selected from phenyl, *p*-CH₃CO-phenyl, *p*-fluorophenyl, or 2-pyridyl (see

Volpini R et al. "Synthesis and Evaluation of Potent and Highly Selective Agonists for




hA3R," J Med Chem 52, 7897-7900. 2009);




(see Pugliese AM *et al.*,

"Role of A3R on CA1 hippocampal neurotransmission during OGD," Biochem







Pharmacology 74. 2007);









; or




(see Klotz KN "Adenosine receptors and their ligands NS's," Arch Pharmacol. 362. 382-391. 2000); or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is selected from a (N)-methanocarba nucleoside such as those disclosed above; or a pharmaceutically acceptable salt thereof.

[0091] In some embodiments, the compound is selected from:









or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is selected from a (N)-methanocarba nucleoside such as those disclosed above; or a pharmaceutically acceptable salt thereof.

[0092] In some embodiments, the compound is selected from





pharmaceutically acceptable salt thereof. In some embodiments, the compound is selected from the group consisting of:





conformation or the methanocarba sugar may be replaced with a D-ribosugar; or a pharmaceutically acceptable salt thereof; or a mono-, di-, or triphosphate thereof or pharmaceutically acceptable salt of the mono-, di-, or triphosphate. In some embodiments, the methanocarba sugar is a D-(N)-methanocarba sugar. In some embodiments, the methanocarba sugar is a D-(S)-methanocarba sugar.

[0093] In some embodiments, the compound is selected from:








(MRS2367),




(MRS1873),



(MRS2371),






pharmaceutically acceptable salt thereof. In some embodiments, the compound is


selected from a (N)-methanocarba nucleoside such as those disclosed above; or a pharmaceutically acceptable salt thereof.

[0094] In some embodiments, the compound is selected from



(MRS2179)

or




•


or a

pharmaceutically acceptable salt thereof.


[0095] In some embodiments, the compound is selected from



2




•



, or


, or

, or




; or a pharmaceutically

acceptable salt thereof.




or a

[0096] In some embodiments, the compound is $\text{HO}-\text{C}_6\text{H}_4-\text{OH}$ or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is




or a pharmaceutically acceptable salt thereof. In some



embodiments, the compound is

or a pharmaceutically


acceptable salt thereof.




[0097] In some embodiments, the compound is

or a

pharmaceutically acceptable salt thereof. In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof. In some




embodiments, the compound is


or a pharmaceutically

acceptable salt thereof.

In some embodiments, the compound is




or a pharmaceutically acceptable salt thereof. In some




embodiments, the compound is

[0098] In one aspect, the present invention provides a pharmaceutical composition comprising a disclosed compound or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient. In some embodiments, the compound is




or




; or a pharmaceutically

acceptable salt thereof.

In some embodiments, the compound is




or a pharmaceutically acceptable salt thereof. In some




embodiments, the compound is

or a pharmaceutically


acceptable salt thereof. In some embodiments, the compound is



[0099] In some embodiments, the compound is selected from those in Figure 15, or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is




2




5


selected from




6, MRS1947




7




8






12



14




15



(MRS1898), or MRS1873; or a pharmaceutically


acceptable salt thereof.

[00100] In one aspect, the present invention provides a method of treating a brain or central nervous system (CNS) injury or condition selected from traumatic brain injury (TBI) or stroke, comprising administering to a patient in need thereof an effective amount of a compound selected from:




a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.

[00101] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00102] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00103] In some embodiments, the brain or central nervous system (CNS) injury or condition is TBI.

[00104] In some embodiments, the TBI is selected from concussion, blast injury, combat-related injury, or a mild, moderate or severe blow to the head. .

[00105] In some embodiments, the compound is administered within 24 hours of the TBI or stroke.

[00106] In some embodiments, the compound is administered within 8 hours of the TBI or stroke.

[00107] In some embodiments, the compound is administered at least during the first 8-48 hours following the TBI or stroke.

[00108] In some embodiments, the brain or central nervous system (CNS) injury or condition is stroke.

[00109] In some embodiments, the compound is administered chronically to treat the stroke during the time period after the stroke has occurred as it resolves.

[00110] In some embodiments, neuroprotection or neurorestoration is increased in the patient as compared with an untreated patient.


[00111] In some embodiments, the compound is a biased partial agonist at a human A₃ adenosine receptor (A₃R).

[00112] In some embodiments, the A₃R is partially agonized in a manner biased toward neuroprotective functions of the A₃R receptor.

[00113] In some embodiments, the compound is administered orally, intravenously, or parenterally.


[00114] In one aspect, the present invention provides a method of increasing neuroprotection or neurorestoration in a patient who has suffered a TBI or stroke,

comprising administering to a patient in need thereof an effective amount of a compound selected from




a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.

[00115] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00116] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00117] In some embodiments, the neuroprotection or neurorestoration decreases the recovery period after the TBI or stroke as compared with an untreated patient.


[00118] In some embodiments, the compound is a biased partial agonist at a human A₃ adenosine receptor (A₃R) and the A₃R is partially agonized in a manner biased toward neuroprotective functions of the A₃R receptor.

[00119] In some embodiments, the compound is administered orally, intravenously, or parenterally.


[00120] In some embodiments, the compound or pharmaceutically acceptable salt thereof has an unbound fraction in plasma of at least 0.7 or at least 0.08 unbound fraction in brain, or both.

[00121] In some embodiments, the compound or pharmaceutically acceptable salt thereof has an unbound fraction in plasma of at least 0.7 or at least 0.08 unbound fraction in brain, or both.

[00122] In one aspect, the present invention provides a method of treating an injury, disease, or condition selected from traumatic brain injury (TBI), stroke, a neurodegenerative condition, or a heart or cardiovascular disease, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R) selected from




(MRS4322) or



(MRS1873),


a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.

[00123] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00124] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00125] In some embodiments, the injury, disease, or condition is TBI.

[00126] In some embodiments, the TBI is selected from concussion, blast injury, combat-related injury, or a mild, moderate or severe blow to the head.

[00127] In some embodiments, the injury, disease, or condition is a stroke selected from ischemic stroke, hemorrhagic stroke, subarachnoid hemorrhage, cerebral vasospasm, or transient ischemic attacks (TIA).

[00128] In some embodiments, the neurodegenerative disease is selected from Alzheimer's Disease (AD), Parkinson's Disease (PD), Huntington's Disease (HD), Multiple Sclerosis (MS), amyotrophic lateral sclerosis (ALS), chronic traumatic encephalopathy (CTE), or a neurodegenerative condition caused by a virus, alcoholism, tumor, toxin, or repetitive brain injuries.

[00129] In some embodiments, the injury, disease, or condition is Parkinson's Disease.

[00130] In some embodiments, the injury, disease, or condition is Alzheimer's Disease, migraine, brain surgery, or a neurological side effect associated with cancer chemotherapy.

[00131] In some embodiments, the heart or cardiovascular disease is selected from cardiac ischemia, myocardial infarction, a cardiomyopathy, coronary artery disease, arrhythmia, myocarditis, pericarditis, angina, hypertensive heart disease, endocarditis, rheumatic heart disease, congenital heart disease, or atherosclerosis.

[00132] In some embodiments, the heart or cardiovascular disease is cardiac ischemia or myocardial infarction.

[00133] In some embodiments, the compound is administered chronically to treat the stroke, cardiac ischemia, or myocardial infarction during the time period after the injury has occurred as it resolves.

[00134] In some embodiments, neuroprotection or neurorestoration is increased in the patient as compared with an untreated patient.

[00135] In some embodiments, the A₃R is agonized in a biased manner toward neuroprotective functions of the A₃R receptor via preferential activation of intracellular calcium mobilization with less, or no, activation of other A₃R-mediated pathways, or via


preferential activation of Gq11-mediated intracellular calcium mobilization, Gi-mediated modulation of cAMP production, or Gi-mediated phosphorylation of ERK1/2 and Akt.

[00136] In some embodiments, the A₃R is partially agonized in a manner biased toward cardioprotective functions of the A₃R receptor via preferential activation of intracellular calcium mobilization with less, or no, activation of other A₃R-mediated pathways, or via preferential activation of Gq11-mediated intracellular calcium mobilization, Gi-mediated modulation of cAMP production, or Gi-mediated phosphorylation of ERK1/2 and Akt.

[00137] In some embodiments, the method increases neuroprotection or neurorestoration in a patient who is suffering from a neurological condition associated with or resulting from cancer chemotherapy or brain surgery.


[00138] In some embodiments, the compound is administered orally.

[00139] In one aspect, the present invention provides a method of increasing neuroprotection or neurorestoration in a patient who has suffered a TBI or stroke, thereby treating the TBI or stroke, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R) selected from




a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.

[00140] In one aspect, the present invention provides a method of increasing cardioprotection or regeneration of damaged heart tissue in a patient who has suffered a cardiac ischemia or myocardial infarction, thereby treating the cardiac ischemia or myocardial infarction, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R) selected from




(MRS4322) or



(MRS1873),


a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable composition comprising the same.

[00141] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00142] In some embodiments, the compound is



or a pharmaceutically acceptable salt thereof.

[00143] In some embodiments, the recovery period after the TBI, stroke, cardiac ischemia, or myocardial infarction is decreased as compared with an untreated patient.

[00144] In some embodiments, the A₃R is partially agonized in a manner biased toward neuroprotective functions of the A₃R receptor.

[00145] In some embodiments, the A₃R is partially agonized in a manner biased toward cardioprotective functions of the A₃R receptor.

[00146] In some embodiments, the compound is administered orally.

[00147] In some embodiments, the compound is a biased agonist of an A₃R with improved cardioprotection function relative to a full A₃R agonist.

[00148] In some embodiments, the compound is a biased agonist of an A₃R with improved cardioprotection function relative to a full A₃R agonist via preferential activation of one or more of the following A₃R-mediated pathways: activation of Gq11-mediated intracellular calcium mobilization, Gi-mediated modulation of cAMP production, Gi-mediated phosphorylation of ERK1/2 and Akt, or modulation of Beta-Arrestin activation.

[00149] In some embodiments, the compound is a biased agonist of an A₃R with improved cardioprotection function relative to a full A₃R agonist via preferential activation of intracellular calcium mobilization with less or no activation of the other A₃R-mediated pathways.

[00150] In some embodiments, the compound is a partial agonist of the A₃R with improved cardioprotection function relative to a full A₃R agonist.

[00151] The amount of a disclosed compound (i.e., active agent) that should be present in a composition for use a disclosed method or a disclosed pharmaceutical composition will generally be a therapeutically effective amount. A “therapeutically effective amount” or dose (or “effective amount”) refers to that amount of the active agent sufficient to result in a desired therapeutic result, such as activation of neuroprotection, neuroregeneration, and/or improvement in cognitive or neurological function. Toxicity and therapeutic efficacy of compositions of active agents can be determined by procedures known in the art in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the test group) and the ED₅₀ (the dose therapeutically effective in 50% of the test group). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compositions which exhibit large therapeutic indices are advantageous. Data obtained from cell culture assays and animal studies can be used in formulating a range of dosages for use in humans. In some embodiments, the dosage of such compositions lies within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized.

[00152] In some embodiments, the effective dose and/or desired therapeutic result are established by comparing the test subject or patient’s cognitive function or another

parameter for at least two measurements; however, more than two measurements may also be used. The initial cognitive function measurement establishes an initial baseline for the test subject or patient. The cognitive function can be measured using an established cognitive test such as the delayed verbal recall task of the revised Wechsler Memory Scale. A further test after treatment with the cognitive test will establish a second measurement. An effective amount is established when the comparison of the second measurement to the first measurement demonstrates an improvement of at least about 1%. In some embodiments, the improvement in cognitive function, as measured by the delayed verbal recall task of the revised Wechsler Memory Scale, is between about 1% and 20%. In some embodiments, the improvement is between about 1% and 10%. In some embodiments, the improvement is between about 1% and 5%. It is understood by one skilled in the art that other methods of determining cognitive function improvement are equally applicable as long as they do not measure stages of dementia.

[00153] Accordingly, the invention encompasses methods of improving cognitive or neurological function by administering an effective amount of a disclosed compound to a subject in need thereof, wherein the enhancement in neurological and cognitive function is measured as a score increase between 1% and 20% in the in the delayed verbal recall task score of the revised Wechsler Memory Scale.

[00154] Disclosed methods of treatment may encompass administration of a disclosed compound as needed to obtain the desired therapeutic effect. The composition can be administered as long as necessary to maintain the desired therapeutic effect. In some embodiments, the compound is administered between about one and 12 months. In some embodiments, the compound is administered between one and six months. In some embodiments, the compound is administered between one and three months.

[00155] In one aspect of the invention, a disclosed compound is administered in an amount between about 5 mg/day and 10 g/day. In some embodiments, each dose of the compound is in an amount between about 5 mg/dose and 10 g/dose. For example, satisfactory results are obtained by oral administration of a disclosed compound of the invention at dosages between about 0.05 and 10 mg/kg/day, between about 0.1 and 7.5 mg/kg/day, between about 0.1 and 2 mg/kg/day, or 0.5 mg/kg/day administered once or, in divided doses, 2 to 4 times per day. For parenteral administration, for example by i.v.

drip or infusion, dosages between about 0.01 and 5 mg/kg/day, between about 0.05 and 1.0 mg/kg/day and between about 0.1 and 1.0 mg/kg/day can be used. Suitable daily dosages for patients are thus between about 2.5 and 500 mg p.o., between about 5 and 250 mg p.o., between about 5 and 100 mg p.o., or between about 0.5 and 250 mg i.v., between about 2.5 and 125 mg i.v. and between about 2.5 and 50 mg i.v.

3. Uses, Formulation and Administration

Pharmaceutically acceptable compositions

[00156] According to another embodiment, the invention provides a composition comprising a disclosed compound and a pharmaceutically acceptable carrier, adjuvant, or vehicle. In certain embodiments, a composition of this invention is formulated for administration to a patient in need of such composition. In some embodiments, a composition of this invention is formulated for oral administration to a patient.

[00157] The term “patient,” as used herein, means an animal, preferably a mammal, and most preferably a human.

[00158] The term “pharmaceutically acceptable carrier, adjuvant, or vehicle” refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[00159] A “pharmaceutically acceptable derivative” means any non-toxic salt, ester, salt of an ester or other derivative of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

[00160] Compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term “parenteral” as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. In some embodiments, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer’s solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

[00161] For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[00162] Pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for

oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[00163] Alternatively, pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[00164] Pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[00165] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[00166] For topical applications, provided pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, provided pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[00167] For ophthalmic use, provided pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

[00168] Pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[00169] In some embodiments, pharmaceutically acceptable compositions of this invention are formulated for oral administration. Such formulations may be administered with or without food. In some embodiments, pharmaceutically acceptable compositions of this invention are administered without food. In other embodiments, pharmaceutically acceptable compositions of this invention are administered with food.

[00170] In other embodiments, pharmaceutically acceptable compositions of this invention are formulated for intravenous (IV) administration.

[00171] The amount of compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, provided compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

[00172] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

Uses of Compounds and Pharmaceutically Acceptable Compositions

[00173] Compounds and compositions described herein are generally useful for the treatment of various diseases and conditions, such as brain injuries and neurodegenerative conditions, and the various methods disclosed herein.

[00174] The activity of a compound utilized in the present invention may be assayed *in vitro*, *in vivo* or in a cell line. *In vitro* assays include assays that determine modulation or binding to a protein. Detailed conditions for assaying a compound are set forth in the Examples below.

[00175] As used herein, the terms “treatment,” “treat,” and “treating” refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

[00176] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of a disclosed disease or condition, or associated condition or symptom. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease or condition, the particular agent, its mode of administration, and the like. Compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression “dosage unit form” as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well

known in the medical arts. The term “patient”, as used herein, means an animal, in some embodiments a mammal, or in certain other embodiments a human.

[00177] Pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, sublingually, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), intraocularly (such as eye drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the disease or condition being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg or from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[00178] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00179] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be

employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00180] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00181] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[00182] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00183] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia,

c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[00184] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00185] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or in a certain

part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[00186] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[00187] Depending upon the particular condition, or disease, to be treated, additional therapeutic agents that are normally administered to treat that condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat a particular disease, or condition, are known as “appropriate for the disease, or condition, being treated.”

[00188] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared and used according to the following general procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds of the present invention, the following general methods, and other methods known to one of ordinary skill in the art, can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

[00189] The contents of each document cited in the specification are herein incorporated by reference in their entireties.

EXEMPLIFICATION

Example 1: Pharmacokinetics of MRS4322 Following Intraperitoneal Administration to Mice

Purpose

[00190] This study was designed to determine the plasma and brain concentrations of MRS4322 following intraperitoneal administration of a dose used in mouse photothrombosis and traumatic brain injury models.

Methods

[00191] **Chemicals:** MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD). Analytical-grade tolbutamide was obtained from commercial supplies at Seventh Wave Laboratories (Maryland Heights, MO). All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO).

[00192] **Animals:** Female C576BL/6J mice weighing approximately 0.02 kg were used for this study, supplied by the University of Texas Health Sciences Center (San Antonio, TX). All studies were conducted under approved University of Texas Health Sciences Center IACUC protocols.

[00193] **Drug Administration:** MRS4322 was solubilized in DMSO and then diluted in saline to prepare dosing solution. Final dosing solution concentration of MRS4322 was 100 μ M. A 100 μ L volume of dosing solution was administered intraperitoneally to each mouse per 20 gram body weight; MRS4322 was administered intraperitoneally at 0.16 mg/kg or 0.5 μ mol/kg. Three mice were administered MRS4322 for each sampling timepoint.

[00194] **Tissue Sampling:** Blood and brain samples were obtained at 0, 0.083, 0.25, 0.5, 1, 2 and 8 hours post-dose. At each timepoint, mice (3/timepoint) were euthanized in a carbon monoxide chamber. Whole blood was obtained by cardiac puncture into Microtainer tubes containing heparin and immediately centrifuged for preparation of plasma; plasma was stored at -80 °C. At each timepoint, whole brain samples were obtained by decapitation, rinsed in ice-cold phosphate-buffered saline and weighed. Brain samples were then immediately flash-frozen in liquid nitrogen and stored at -80 °C.

Bioanalysis

[00195] Plasma and brain concentrations of MRS4322 were determined by LC-MS/MS utilizing tolbutamide as an internal standard. The following table outlines the LC and MS/MS conditions employed:

Table 1: Bioanalytical Methods for MRS4322 Plasma and Brain Concentration Determinations

Bioanalytical Methods			
System Components			
Module	Manufacturer	Model	
HPLC	Shimadzu	Prominence LC-20AD Binary Pumps	
Autosampler	Leap	Prominence Sil-30AC	
Mass Spectrometer	AB Sciex	4000 Qtrap	
HPLC Method			
Column	Phenomenex Kinetex C18 (2.1 x 50 mm, 2.6 μm)		
Elution	Gradient, 0.4 mL/min		
	Mobile Phase A: 0.1% Formic Acid		
	Mobile Phase B: 0.1% Formic Acid in Acetonitrile		
MS Detection and Calibration for MRS4322 in Mouse Plasma			
Peak Name: Tolbutamide			
Use as Internal Standard			
Q1/Q3 Masses: 271.20/155.20 Da			
Peak Name: MRS4322			
Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.10/182.00 Da			
Fit	Linear	Weighting	1 / (x * x)
Intercept	0.000447		
Slope	0.00141		
Correlation coefficient	0.9996		
Use Area			
MS Detection and Calibration for MRS4322 in Mouse Brain Homogenate			
Peak Name: Tolbutamide			
Use as Internal Standard			
Q1/Q3 Masses: 271.20/155.20 Da			
Peak Name: MRS4322			

Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.10/182.00 Da			
Fit	Linear	Weighting	1 / (x * x)
Intercept	0.0000465		
Slope	0.000883		
Correlation coefficient	0.9996		
Use Area			

[00196] For each tissue matrix, standard curves were created and LLOQ/ULOQ concentrations determined. The calibration range for the MRS4322 plasma concentration standard curve was 2.42-242 ng/mL. The calibration range for MRS4322 brain concentrations was 2.41-233 ng/mL.

[00197] For bioanalysis of brain concentrations of MRS4322, brain samples were homogenized in ice-cold phosphate-buffered saline in a 4x dilution. Aliquots of the resulting diluted brain homogenate were treated with acetonitrile and analyzed by LC-MS/MS. Because of the 4x homogenate dilution, the calibration range of the MRS4322 brain standard curve was translated to 9.64-932 ng/gm. In several samples, MRS4322 was detectable but fell below the brain LLOQ of 9.64 ng/gm but were above background; in these cases the final reported brain concentration was extrapolated based on MS peak heights.

Results

[00198] Following intraperitoneal administration to mice, MRS4322 concentrations were detectable in both plasma and brain samples (Figure 1A and Table 2). Note that as described in Example 11 below and shown in Figures 1B and 16, MRS4322 concentrations were detectable in neonatal pig plasma and brain samples. Following intravenous administration to neonatal pigs, MRS4322 concentrations were detectable in plasma, brain, brain extracellular fluid and cerebrospinal fluid samples (Figure 1B and Figure 16, Table 13).

Table 2: Plasma and Brain Concentrations of MRS4322 in Mice Following Intraperitoneal Administration

Group	Matrix	Test Article	Dose (mg/kg)	Time (h)	Plasma Concentration (ng/mL) by Subject					
					A	B	C	Mean	SD	N
1	Plasma	MRS4322	0.16	0	BLQ	BLQ	BLQ	BLQ	---	0
				0.083	40.2	42.5	65.8	49.5	14.2	3
				0.25	37.5	41.6	46.3	41.8	4.40	3
				0.5	14.8	26.8	21.6	21.1	6.00	3
				1	7.35	4.23	85.0*	5.79	NA	2
				2	0.539 [†]	0.638 [†]	1.15 [†]	0.776	0.328	3
				8	BLQ	BLQ	BLQ	BLQ	---	0
Group	Matrix	Test Article	Dose (mg/kg)	Time (h)	Brain Concentration (ng/g) by Subject					
					A	B	C	Mean	SD	N
1	Brain	MRS4322	0.16	0	BLQ	BLQ	BLQ	BLQ	---	0
				0.083	2.04 [†]	2.18 [†]	2.55 [†]	2.11	0.100	3
				0.25	3.08 [†]	3.03 [†]	3.43 [†]	3.06	0.0400	3
				0.5	1.77 [†]	3.00 [†]	1.81 [†]	2.39	0.870	3
				1	BLQ	BLQ	BLQ	BLQ	---	0
				2	BLQ	BLQ	BLQ	BLQ	---	0
				8	BLQ	BLQ	BLQ	BLQ	---	0

BLQ = Below the Lower Limit of Quantitation (2.40 ng/mL for plasma, 9.64 ng/g for Brain due to 4x dilution)

* Animal F at 1 hour value of 85.0 was considered an outlier and was excluded from summary statistics

[†] Calculated concentrations were <LLOQ. Values reported are extrapolated

Plasma concentrations allowed initial estimates of Tmax, Cmax, half-life and AUC (Table 3).

Table 3: Plasma Pharmacokinetics of MRS4322 in Mice Following Intraperitoneal Administration

Group	Matrix	Analyte	Dose (mg/kg)	Animal ID	C _{max} (ng/mL)	T _{max} (hr)	AUC _{last} (hr*ng/mL)	AUC _{INF} (hr*ng/mL)	Half-life (hr)
1	Plasma	MRS4322	0.16	A	40.2	0.083	24.2	24.4	0.306
				B	42.5	0.083	27.5	27.8	0.288
				C	65.8	0.083	37.6	38.2	0.337
				N	3	3	3	3	3
				Mean	49.5	0.083	29.8	30.1	0.310

				SD	14.2	0.000	7.01	7.19	0.0247
				CV%	28.6	0.0	23.5	23.8	8.0

Note: BLQ values were converted to zero, extrapolated values were included, and Animal F at 1 hour was excluded for PK analyses

[00199] Although brain concentrations were detectable, data was insufficient for estimation of half-life or other pharmacokinetic parameters other than C_{max} and T_{max}. However, based on the available plasma and brain data, it was estimated that the brain/plasma ratio of total drug was approximately 0.06 based on mean C_{max} concentrations in plasma and brain.

[00200] These results confirm that circulating plasma concentrations of MRS4322 are detectable following intraperitoneal administration to mice under the dosing conditions used in models of photothrombosis and traumatic brain injury, and that MRS4322 distributes to the brain under these dosing conditions.

Example 2: Pharmacokinetics of MRS4322 Following Intraperitoneal Administration of MRS2365 to Mice

Purpose

[00201] This study was designed to determine the plasma and brain concentrations of MRS4322 following intraperitoneal administration of the P2Y₁ agonist MRS2365 at a dose used in mouse photothrombosis and traumatic brain injury models.

Methods

[00202] **Chemicals:** MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD). MRS2365 was obtained from Tocris Biosciences (Bristol, UK). Analytical-grade tolbutamide was obtained from commercial supplies at Seventh Wave Laboratories (Maryland Heights, MO). All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO).

[00203] **Animals:** Female C576BL/6J mice weighing approximately 0.02 Kg were used for this study, supplied by the University of Texas Health Sciences Center (San Antonio, TX). All studies were conducted under approved University of Texas Health Sciences Center IACUC protocols.

[00204] Drug Administration: MRS2365 was solubilized in phosphate-buffered saline and then diluted in phosphate-buffered saline to prepare dosing solution. Final dosing solution concentration of MRS2365 was 100 μ M. A 100 μ L volume of dosing solution was administered intraperitoneally to each mouse per 20 gram body weight; MRS2365 was administered intraperitoneally at 0.5 μ mol/kg or 0.24 mg/kg. Three mice were administered MRS2365 for each sampling timepoint.

[00205] Tissue Sampling: Blood and brain samples were obtained at 0, 0.083, 0.25, 0.5, 1, 2 and 8 hours post-dose. At each timepoint, mice (3/timepoint) were euthanized in a carbon monoxide chamber. Whole blood was obtained by cardiac puncture into Microtainer tubes containing heparin and immediately centrifuged for preparation of plasma; plasma was stored at -80 °C. At each timepoint, whole brain samples were obtained by decapitation, rinsed in ice-cold phosphate-buffered saline and weighed. Brain samples were then immediately flash-frozen in liquid nitrogen and stored at -80 °C.

[00206] Bioanalysis: Previous studies had demonstrated that no detectable circulating and brain concentrations of MRS2365 were observed following intraperitoneal or intravenous administration, so this study focused only on the detection and quantitation of its metabolite MRS4322. Plasma and brain concentrations of MRS4322 were determined by LC-MS/MS utilizing tolbutamide as an internal standard. The following table outlines the LC and MS/MS conditions employed:

Table 4: Bioanalytical Methods for MRS4322 Plasma and Brain Concentration Determinations

Bioanalytical Methods			
System Components			
Module	Manufacturer	Model	
HPLC	Shimadzu	Prominence LC-20AD Binary Pumps	
Autosampler	Leap	Prominence Sil-30AC	
Mass Spectrometer	AB Sciex	4000 Qtrap	
HPLC Method			
Column	Phenomenex Kinetex C18 (2.1 x 50 mm, 2.6 μm)		
Elution	Gradient, 0.4 mL/min		
	Mobile Phase A: 0.1% Formic Acid		
	Mobile Phase B: 0.1% Formic Acid in Acetonitrile		
MS Detection and Calibration for MRS4322 in Mouse Plasma			
Peak Name: Tolbutamide			

Use as Internal Standard			
Q1/Q3 Masses: 271.20/155.20 Da			
Peak Name: MRS4322			
Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.10/182.00 Da			
Fit	Linear	Weighting	1 / x
Intercept	0.000704		
Slope	0.00151		
Correlation coefficient	0.9998		
Use Area			
MS Detection and Calibration for MRS4322 in Mouse Brain Homogenate			
Peak Name: Tolbutamide			
Use as Internal Standard			
Q1/Q3 Masses: 271.20/155.20 Da			
Peak Name: MRS4322			
Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.10/182.00 Da			
Fit	Linear	Weighting	1 / x
Intercept	0.000209		
Slope	0.000864		
Correlation coefficient	1.00		
Use Area			

[00207] For each tissue matrix, standard curves were created and LLOQ/ULOQ concentrations determined. The calibration range for the MRS4322 plasma concentration standard curve was 2.26-241 ng/mL. The calibration range for MRS4322 brain concentrations was 2.35-242 ng/mL.

[00208] For bioanalysis of brain concentrations of MRS4322, brain samples were homogenized in ice-cold phosphate-buffered saline in a 4x dilution. Aliquots of the resulting diluted brain homogenate were treated with acetonitrile and analyzed by LC-MS/MS. Because of the 4x homogenate dilution, the calibration range of the MRS4322 brain standard curve was translated to 9.40-968 ng/gm. In several samples, MRS4322 were detectable but fell below the brain LLOQ of 9.40 ng/gm but were above background; in these cases the final reported brain concentration was extrapolated based on MS peak heights.

Results

[00209] Following intraperitoneal administration of MRS2365 to mice, MRS4322 concentrations were detectable in both plasma and brain samples (Figure 2 and Table 5).

Table 5: Plasma and Brain Concentrations of MRS4322 in Mice Following Intraperitoneal Administration of MRS2365

Group	Matrix	Test Article	Dose (mg/kg)	Time (h)	Plasma Concentration (ng/mL) by Subject					
					A	B	C	Mean	SD	N
1	Plasma	MRS4322	0.24 MRS2365	0	BLQ	BLQ	BLQ	BLQ	---	0
				0.083	24.5	64.1	82.8	57.1	29.8	3
				0.25	46.5	43.2	47.4	45.7	2.20	3
				0.5	22.3	25.1	31.0	26.1	4.40	3
				1	2.30	4.02	3.53	3.28	0.890	3
				2	BLQ	1.02*	0.680*	0.850	NA	2
				8	BLQ	BLQ	BLQ	BLQ	---	0
Group	Matrix	Test Article	Dose	Time (h)	Brain Concentration (ng/g) by Subject					
					A	B	C	Mean	SD	N
1	Brain ^a	MRS4322	0.24 MRS2365	0	2.76*	10.0	4.61*	5.79	3.76	3
				0.083	BLQ	BLQ	BLQ	BLQ	---	0
				0.25	3.92*	3.26*	4.16*	3.78	0.470	3
				0.5	3.19*	2.80*	2.70*	2.90	0.260	3
				1	BLQ	BLQ	BLQ	BLQ	---	0
				2	BLQ	BLQ	BLQ	BLQ	---	0
				8	BLQ	BLQ	BLQ	BLQ	---	0

BLQ = Below the Lower Limit of Quantitation (2.26 ng/mL for plasma; 9.40 ng/g for Brain due to 4-fold dilution)

* Projected values; these values were below the limit of quantitation

^a It appears possible that the 0 and 0.083 samples were mislabeled upon receipt

Plasma concentrations allowed initial estimates of T_{max}, C_{max}, half-life and AUC (Table 6).

Table 6: Plasma Pharmacokinetics of MRS4322 in Mice Following Intraperitoneal Administration of MRS2365

Group	Matrix	Analyte	Dose (mg/kg)	Animal ID	C _{max} (ng/mL)	T _{max} (hr)	AUC _{last} (hr*ng/mL)	AUC _{INF} (hr*ng/mL)	Half-life (hr)
1	Plasm	MRS432	0.24	A ^a	46.5	0.25	21.7	22.3	0.17

	a	2	MRS2 365	B	64.1	0.08	30.0	30.5	0.34
				C	82.8	0.08	34.8	35.1	0.28
				N	3	3	3	3	3
				Mean	64.5	0.14	28.8	29.3	0.27
				SD	18.2	0.10	6.65	6.51	0.09
				CV%	28.2	69.5	23.1	22.2	33.1

BLQ values were converted to zero; extrapolated values were included for PK analyses.

[00210] Although brain concentrations were detectable, data was insufficient for estimation of half-life or other pharmacokinetic parameters other than C_{max} and T_{max}. However, based on the available plasma and brain data, it was estimated that the brain/plasma ratio of total drug was approximately 0.10 based on mean C_{max} concentrations in plasma and brain.


[00211] These results confirm that circulating plasma concentrations of MRS4322 are detectable following intraperitoneal administration of the P2Y₁ agonist MRS2365 to mice under the dosing conditions used in models of photothrombosis and traumatic brain injury, and that MRS4322 distributes to the brain under these dosing conditions.

[00212] MRS4322 plasma and brain concentrations were determined in two different studies, following intraperitoneal administration of MRS4322 itself (see Example 1) and following intraperitoneal administration of the P2Y₁ agonist MRS2365. In both of these studies, MRS4322 or MRS2365 were administered at equimolar doses of 0.5 µmol/kg. Comparing the results of both studies, the observed plasma concentrations of MRS4322 were virtually identical and brain concentrations of MRS4322 were very similar (compare Figures 1 and 2). There was no statistically significant difference in the half-life and AUC values for MRS4322 following administration of equimolar doses of MRS4322 or MRS2365. These data indicate that following intraperitoneal administration to mice, MRS2365 is rapidly and completely metabolized to MRS4322, resulting in MRS4322 plasma and brain pharmacokinetics very similar to those following intraperitoneal administration of MRS4322 itself.

Example 3: Plasma and Brain Binding of MRS4322 in Mice

Purpose

[00213] This study was designed to determine the plasma and brain free fraction of MRS4322 in mice and compare free fractions to those of a prototypical adenosine A₃ receptor agonist, MRS5698. MRS5698 has the following chemical structure:



Methods

[00214] **Chemicals:** MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD). MRS5698 was acquired from Tocris Biosciences (Bristol, UK). Analytical-grade sulfamethoxazole and warfarin were obtained from commercial supplies at Seventh Wave Laboratories (Maryland Heights, MO). All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO).

[00215] **Animals and Tissue Preparation:** Plasma from male CD-1 mice was obtained from BioreclamationIVT (Westbury, NY) and stored at -80 °C until use. Male CD-1 mouse brains were obtained from BioreclamationIVT (Westbury, NY).

[00216] Plasma ultrafiltrate blank samples were prepared by thawing frozen plasma and then pre-warming plasma in a humidified 5% CO₂ chamber at 37 °C for 60 minutes. Aliquots of 800 µL were transferred to Centrifree Centrifugal Filters (Ultracel regenerated cellulose (NMWL 30,000 amu) Lot R5JA31736) and centrifuged at 2900 RPM at 37 °C for 10 minutes; plasma water filtrates were collected and used in preparation of standards, blanks and QC standards.

[00217] Brains were weighed and homogenized with 1:9 phosphate-buffered saline, pH 7.4 using an Omni tissue homogenizer. Brains from four mice were homogenized, pooled and mixed to form one sample.

[00218] Plasma Binding Determination: MRS4322, MRS5698, sulfamethaxazole and warfarin were solubilized in DMSO and then diluted in 1:1 acetonitrile:water to prepare 100 μ M dialysis stock solutions. Sulfamethaxazole and warfarin were utilized as study standards with known plasma binding values. Plasma samples were pre-warmed for 60 minutes in a humidified, 5% CO₂ incubator maintained at 37 °C. Three mL aliquots of pre-warmed plasma were each spiked with MRS4322, MRS5698, sulfamethaxazole or warfarin using 100 μ M stock solutions for each compound resulting in final test concentrations of 1 μ M. Spiked plasma samples were incubated on a rotary mixer in a humidified 5% CO₂ chamber at 37 °C for a minimum of 60 minutes. After 60 minutes, three 800 μ L aliquots of each sample were added to Centrifree centrifugal filters. The filters were subjected to centrifugation at 2900 rpm for 10 minutes at 37 °C. Three 100 μ L aliquots of residual plasma were collected along with ultrafiltrate for bioanalysis.

[00219] Brain Binding Determination: MRS4322, MRS5698, sulfamethoxazole and warfarin were solubilized in DMSO and diluted in 1:1 acetonitrile:water to prepare 100 μ M dialysis stock solutions. Pooled homogenized brains were pre-warmed for 60 minutes in a humidified, 5% CO₂ incubator maintained at 37 °C. Three mL aliquots of brain homogenate were each spiked with MRS4322, MRS5698, sulfamethaxazole or warfarin using the 100 μ M stock solutions for each compound resulting in final spiked concentrations of 1 μ M. Spiked pooled brain homogenates were placed on a Nutator mixer in a humidified, 5% CO₂ incubator at 37 °C for 60 minutes. After 60 minutes, three 800 μ L aliquots of each sample were added to Centrifree centrifugal filters. The filters were subjected to centrifugation at 2900 rpm for 10 minutes at 37 °C. Aliquots of residual brain homogenate and ultrafiltrate were collected for bioanalysis.

Bioanalysis

[00220] Plasma and brain concentrations of MRS4322 and MRS5698 in spiked plasma, brain homogenates and associated ultrafiltrates were determined by LC-MS/MS utilizing tolbutamide as an internal standard. Associated concentrations of sulfamethaxazole and warfarin were also determined by LC-MS/MS using standard conditions (data not shown). The following tables outline the LC and MS/MS conditions employed (Tables 7 and 8). Bioanalytical methods were identical for all matrices;

standard curve statistics (e.g. Fit, Intercept, Slope, Correlation Coefficient) were determined for each matrix but were not significantly different and thus are not shown for each matrix.

Table 7: Bioanalytical Methods for MRS4322 for Determination of Plasma, Brain Homogenate, Plasma Ultrafiltrate and Brain Homogenate Ultrafiltrate Concentrations

Bioanalysis of Plasma, Brain Homogenates, Plasma Ultrafiltrate and Brain Homogenate Ultrafiltrate Samples			
System Components			
Module	Manufacturer	Model	
HPLC	Shimadzu	Prominence LC20AD Binary Pumps	
Autosampler	CTC Leap	HTC PAL	
Mass Spectrometer	AB Sciex	API 4000	
HPLC Method			
Column	Phenomenex Kinetex C18 (2.1 x 50 mm, 2.6 μm)		
Elution	Gradient, 0.4 mL/min		
	Mobile Phase A: 0.1% Formic acid in Water		
	Mobile Phase B: 0.1% Formic acid in Acetonitrile		
Mass Spectrometer Conditions for MRS4322			
Peak Name: MRS4322			
Q1/Q3 Masses: 324.10/182.00 Da			
Peak Name: Tolbutamide			
Q1/Q3 Masses: 271.20/155.20 Da			
Fit	Linear	Weighting	1 / x
Intercept	0.0181		
Slope	0.00192		
Correlation coefficient	0.9988		

Table 8: Bioanalytical Methods for MRS5698 for Determination of Plasma, Brain Homogenate, Plasma Ultrafiltrate and Brain Homogenate Ultrafiltrate Concentrations

System Components		
Module	Manufacturer	Model
HPLC	Shimadzu	Prominence LC20AD Binary Pumps
Autosampler	CTC Leap	HTC PAL

Mass Spectrometer	AB Sciex	API 4000
HPLC Method		
Column	Phenomenex Kinetex C18 (2.1 x 50 mm, 2.6 µm)	
Elution	Gradient, 0.4 mL/min	
	Mobile Phase A: 0.1% Formic acid in Water	
	Mobile Phase B: 0.1% Formic acid in Acetonitrile	
Mass Spectrometer Conditions for MRS5698		
Peak Name: MRS5698		
Q1/Q3 Masses: 565.10/395.90 Da		
Peak Name: Tolbutamide		
Q1/Q3 Masses: 271.20/155.20 Da		
Fit	Linear	Weighting 1 / x
Intercept	0.107	
Slope	0.00786	
Correlation coefficient	0.9986	

[00221] For each tissue matrix, standard curves were created and LLOQ/ULOQ concentrations determined. The calibration range for the MRS4322 and MRS5698 plasma concentration standard curves was 400-1200 nM. The calibration range for MRS4322 and MRS5698 plasma ultrafiltrate standard curves was 100-1200 nM. The calibration range for MRS4322 brain homogenate and brain homogenate ultrafiltrate standard curves were 400-1200 nM and 100-1200 nM, respectively. The calibration range for MRS5698 brain homogenate and brain homogenate ultrafiltrate standard curves were 400-1200 nM and 1-500 nM, respectively.

Results

[00222] Plasma binding and free fraction were determined for MRS4322 and MRS5698 utilizing plasma ultrafiltration. Plasma binding was 25.8% for MRS4322; associated free fraction was 0.742 (Table 9).

[00223] MRS5698 concentrations were not detected in the plasma ultrafiltrate; MRS5698 was completely recovered in the residual spiked plasma samples obtained from the donor side of the Centrifree devices (data not shown). This indicates that the low MRS5698 ultrafiltrate concentrations were not due to low analytical recovery of the compound. Overall, these data are consistent with the high protein binding (99.88%) of MRS5698 in mouse plasma and tissue reported in the literature (Tosh, D.K. et al.

Purinergic Signalling (2015) 11:371-387). Binding of the study standards sulfamethaxazole and warfarin were consistent with literature values.

Table 9: Fraction Unbound and Binding of MRS4322 and MRS5698 in Mouse Plasma

Substrate	Fraction Unbound		% Bound		N
	Mean	SD	Mean	SD	
MRS4322	0.742	0.065	25.8	6.5	3
MRS5698	*	*	*	*	3
Sulfamethoxazole	0.345	0.027	65.5	2.7	3
Warfarin	0.0485	0.0031	95.1	0.3	3

* Plasma ultrafiltrate concentrations of MRS5698 were BLQ

[00224] Brain binding and free fraction were determined for MRS4322 and MRS5698 utilizing brain homogenate ultrafiltration. Brain binding was 87% for MRS4322; associated free fraction was 0.13 (Table 10).

[00225] MRS5698 concentrations were not detected in the brain homogenate ultrafiltrate. For estimation purposes, the MRS5698 brain homogenate ultrafiltrate LLOQ was utilized to calculate a brain binding value. The resulting brain binding value was 99.99%. Overall, these data are consistent with the high protein binding (99.88%) of MRS5698 in mouse plasma and tissue reported in the literature (Tosh, D.K. et al. Purinergic Signalling (2015) 11:371-387). Binding of the study standards sulfamethaxazole and warfarin were consistent with literature values.

Table 10: Fraction Unbound and Binding of MRS4322 and MRS5698 in Mouse Brain Homogenates

Assay	Substrate	Fraction Unbound		% Bound		N
		Mean	SD	Mean	SD	
1	MRS4322	0.130	0.010	87.0	1.0	3
	MRS5698	<0.000122*	---	>99.99*	---	3
	Sulfamethoxazole	0.431	0.049	56.9	4.9	3
	Warfarin	0.208	0.038	79.2	3.8	3

* MRS5698 was not detected in ultrafiltrate; assay LLOQ was utilized to calculate an estimated binding value

[00226] Overall, these data indicate that MRS4322 has a substantially higher free fraction and lower protein binding than the adenosine A₃ agonist MRS5698 in both plasma and brain. These data indicate that for a given total plasma or brain concentration, substantially higher concentrations of MRS4322 would be available to interact with effector sites than would be available for MRS5698.

Example 4: *In Vitro* Stability and Metabolism of MRS2365 in Mouse and Human Blood and Plasma

Purpose

[00227] This study was designed to determine the *in vitro* stability and metabolic fate of the P2Y₁ agonist MRS2365 in mouse and human blood and plasma.

Methods

[00228] **Chemicals:** MRS2365 was obtained from Tocris Biosciences (Bristol, UK). MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD). All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO). Enalapril and procaine were used as plasma and blood stability standards for mouse and human, respectively.

[00229] **Tissue Preparation:** Plasma from male CD-1 mice and humans was obtained from BioreclamationIVT (Westbury, NY) and stored at -80 °C until use. Whole blood was obtained from male CD-1 mice and human volunteers at Seventh Wave Laboratories (Maryland Heights, MO). Plasma and blood samples were prepared using either EDTA (1 mM) or lithium heparin as anticoagulants.

[00230] **Plasma Stability Determination:** MRS2365, enalapril and procaine were solubilized in phosphate-buffered saline, pH 7.4. Plasma samples (prepared from blood with either EDTA or lithium heparin as anticoagulant) were pre-warmed for 60 minutes in a humidified, 5% CO₂ incubator maintained at 37 °C. Stability incubations were initiated with the addition of MRS2365 (1 µM final concentration). Initial assessments of stability in EDTA-generated plasma utilized incubation timepoints of 0, 10, 30, 60, 120 and 240 minutes (Figures 3 and 4). Subsequent studies comparing EDTA- and heparin-generated plasma utilized incubation timepoints of 0, 1, 2.5, 5, 7.5, 10 and 30 minutes (Figure 7). Additional plasma stability incubations comparing EDTA- and heparin-

generated plasma were performed using timepoints of 0, 5, 10, 20, 30, 45, 60 and 90 seconds (Figure 8). For metabolite scouting analyses, MRS2365 was incubated in heparinized human plasma at a concentration of 100 μ M for 10 or 30 minutes (Figure 9). In all studies, plasma was immediately placed in microtainer tubes, frozen on dry ice and stored at -80 °C until analysis.

[00231] Blood Stability Determination: MRS2365, enalapril and procaine were solubilized in phosphate-buffered saline, pH 7.4. Blood samples (EDTA- or lithium heparin-treated) were pre-warmed for 60 minutes in a humidified, 5% CO₂ incubator maintained at 37 °C. Stability incubations were initiated with the addition of MRS2365 (1 μ M final concentration) (Figures 5 and 6). Blood sample aliquots were obtained at 0, 1, 2.5, 7.5, 10 and 30 minutes and placed in microtainer tubes. For metabolite scouting analyses, MRS2365 was incubated in heparinized human whole blood at a concentration of 100 μ M for 10 or 30 minutes (Figure 10). In all studies, plasma was immediately prepared by centrifugation at 4 °C, placed in microtainer tubes, frozen on dry ice and stored at -80 °C until analysis.

Bioanalysis

[00232] Plasma and blood concentrations of MRS2365 were determined by LC-MS/MS utilizing tolbutamide as an internal standard. The following table outlines the LC and MS/MS conditions employed (Table 11). Bioanalytical methods were identical for all matrices; standard curve statistics (e.g. Fit, Intercept, Slope, Correlation Coefficient) were determined for each matrix but were not significantly different and thus are not shown for each matrix.

Table 11: Bioanalytical Methods for MRS2365 for Determination of Plasma and Blood Concentrations in In Vitro Stability Determinations

Bioanalysis Methods – Plasma and Blood		
System Components		
Module	Manufacturer	Model
HPLC	Shimadzu	Prominence
Autosampler	Leap	CTC Pal
Mass Spectrometer	AB Sciex	API 4000
HPLC Method		
Column	Luna C18 (2.1 x 50 mm, 2.6 μ m)	

Elution	Gradient, 0.4 mL/min Mobile Phase A: 2 mM Ammonium acetate, pH 7.65 Mobile Phase B: 2 mM Ammonium acetate in Acetonitrile		
MS Detection and Calibration for MRS2365 in Plasma and Blood			
Peak Name: Tolbutamide Use as Internal Standard Q1/Q3 Masses: 269.00/170.00 Da			
Peak Name: MRS2365 Internal Standard: Tolbutamide Q1/Q3 Masses: 482.11/384.00 Da			
Fit	Linear	Weighting	1 / (x * x)
Intercept	0.00004		
Slope	0.00002		
Correlation coefficient	0.9854		
Use Area			

[00233] For each tissue matrix, standard curves were created and LLOQ/ULOQ concentrations determined. The calibration range for MRS2365 plasma and blood concentration standard curves was 5-5000 ng/mL. Quality control samples were utilized at concentrations of 32, 160, 800 and 4000 ng/mL.

[00234] Plasma and blood samples were also qualitatively analyzed for MRS2365 metabolites utilizing a LC-MS/MS-based “metabolite scouting” protocol. From review of the metabolic pathways of other nucleotides (e.g. ATP, ADP), it was hypothesized that the most likely metabolites of MRS2365 would be the dephosphorylated (i.e., monophosphate) metabolite (MRS2347, mw = 403.07) and/or the completely dephosphorylated riboside metabolite (MRS4322, mw = 323.10). Blood and plasma samples were analyzed by both positive- and negative-ion LC/MS-MS for parent compound and metabolites as it was anticipated that the phosphorylated compounds would preferentially generate negative ions while the riboside MRS4322 would preferentially generate positive ions. In positive-ion mode, extracted ion chromatograms monitored masses in the range of 323.7-324.7 and 403.7-404.7 to scout for MRS4322 and MRS2347, respectively. Any ion peaks in these ranges were further analyzed to generate product ion spectra for these ion chromatogram peaks. In negative-ion mode, extracted ion chromatograms monitored masses in the range of 321.70-322.70, 401.70-402.70 and 481.70-482.70 to scout for MRS4322, MRS2347 and MRS2365, respectively; peaks were

further analyzed to generate product ion spectra. Because authentic standards of MRS4322 and MRS2347 were unavailable, only qualitative identification of metabolites could be made as ion peak heights/areas could not be compared due to potential differences in ionization efficiency.

Results

[00235] Two time course protocols were utilized in assessing the plasma stability of MRS2365 in mouse and human plasma. Preliminary data were generated with EDTA-prepared plasma over a 240 minute time course. A subsequent study utilized timepoints from 1-30 minutes post-incubation, and the final study utilized a shorter time course of 5-90 seconds post-incubation. In all protocols, plasma stability was analyzed in plasma prepared from blood with either EDTA or lithium heparin as the anticoagulant. *In vitro* stability data for enalapril (mouse) and procaine (human) in plasma and whole blood were consistent with published values (data not shown).

[00236] In plasma stability studies utilizing EDTA-prepared plasma, MRS2365 was essentially stable over the course of the incubation out to 240 minutes post-incubation (Figures 3 and 4). Half-life values could not be calculated over the time course of the incubation, and thus were >240 minutes. In subsequent metabolite scouting analyzes, neither MRS2347 nor MRS4322 could be detected. These data suggested that MRS2365 was stable *in vitro* in mouse and human plasma.

[00237] Similar observations were made when the stability of MRS2365 was assessed in mouse and human whole-blood that had been treated with EDTA as the anticoagulant (Figures 5 and 6). In those studies, the half-life of MRS2365 in EDTA-treated mouse whole blood was 47 minutes, suggesting that EDTA did not completely inhibit clearance of MRS2365 in this matrix. However, MRS2365 was again completely stable in EDTA-treated human whole blood, with an estimated half-life of >240 minutes. No metabolites were detected in these whole blood studies.

[00238] These data were inconsistent with initial *in vivo* pharmacokinetic assessments of MRS2365 in mice that utilized heparin as a blood anticoagulant for plasma preparation, in which MRS2365 appeared to be rapidly cleared at a rate approximating cardiac output, suggesting extensive extrahepatic clearance of the compound. It is known that EDTA chelates divalent cations that are required for the enzymatic activity of

ectonucleotidases, enzymes responsible for the dephosphorylation of nucleotides, that are present both on the surface of cell membranes and circulating in blood and plasma (See Ziganshin et al. Pflugers Arch. (1995) 429:412-418). Thus, it was possible that the primary metabolic pathway of the P2Y₁ agonist nucleotide analog MRS2365 was completely inhibited by the EDTA used to prepare the plasma and whole blood for studies described above.

[00239] To investigate this possibility, additional stability studies were conducted to compare the stability of MRS2365 in plasma that had been generated by either EDTA or lithium heparin, the latter being an anticoagulant with no reported inhibitory effects on ectonucleotidases. Studies were first conducted with heparinized plasma and whole blood over a time course of 0-30 minutes. Quantitation of MRS2365 concentrations revealed extremely low concentrations of MRS2365 at all timepoints, with insufficient data to calculate an *in vitro* half-life (data not shown). Since low/undetectable MRS2365 concentrations were observed at even the shortest timepoints in this study (0 and 1 minutes), the study was repeated with a significantly shorter time course (0-90 seconds) to compare *in vitro* stability in both EDTA- and heparin-treated plasma and to attempt to calculate an *in vitro* half-life. In this shorter study, MRS2365 was relatively stable in EDTA-treated mouse and human plasma (Figures 5 and 6); variability in MRS2365 concentrations were likely due to incomplete mixing at the initiation of the short timecourse and variability in sampling multiple timepoints over the course of the 90 second incubation period. However, in heparin-treated plasma, MRS2365 concentrations were extremely low at even the shortest (0 and 5 seconds) timepoints. MRS2365 was completely stable when incubated in phosphate-buffered saline.

[00240] Together, these data suggest that although MRS2365 is inherently stable in solution, it is rapidly degraded in mouse and human plasma and blood by a process that is inhibited by EDTA. Considering that MRS2365 is a nucleotide analog and EDTA is a known inhibitor of ectonucleotidases that dephosphorylate nucleotides, these data strongly suggest that MRS2365 is susceptible to rapid dephosphorylation in plasma and whole blood, consistent with the *in vivo* pharmacokinetic data generated for MRS2365 in mouse.

[00241] To further investigate the possibility of dephosphorylation of MRS2365 as the cause of instability in mouse and human plasma and whole blood, metabolite scouting was conducted in heparinized human plasma and blood following incubation of MRS2365 at 100 μ M for 10 or 30 minutes; the higher substrate concentration was used to ensure detection of any metabolites formed. Representative ion chromatograms and product ion spectra under positive-ionization conditions are illustrated (Figures 7 and 8); negative-ionization chromatograms and product ion spectra yielded similar results (data not shown). Although the lack of chemical standards prevented the determination of absolute concentrations of metabolites, the relative abundances of metabolite M1 increased compared to metabolite M2 from 10 to 30 minutes post-incubation. Metabolites M2 and M1 parent ion chromatograms and resulting product ion spectra under both ionization conditions were consistent with the mass and structure of the dephosphorylated monophosphate metabolite MRS2347 and the fully dephosphorylated riboside metabolite MRS4322, respectively (Figure 9). The detection of both metabolites and the increase in abundance of metabolite M1 (MRS4322) relative to metabolite M2 (MRS2347) suggested a stepwise dephosphorylation of MRS2365 in human plasma and blood, progressing through the monophosphorylated intermediate MRS2347 that is then further dephosphorylated to the unphosphorylated riboside MRS4322.

[00242] Together, these data support the hypothesis that MRS2365 is rapidly metabolized in plasma and blood by circulating ectonucleotidases that dephosphorylate the parent compound, ultimately resulting in the unphosphorylated riboside MRS4322. This process is inhibited by EDTA, an agent known to chelate divalent cations that are required for ectonucleotidase activity.

Example 5: Neuroprotective Efficacy of MRS4322 after TBI in Mice

Purpose

[00243] This study was designed to determine the neuroprotective efficacy of MRS4322 in mice subjected to traumatic brain injury (TBI) and to compare free mice treated with MRS2365 and an adenosine A₃ receptor full agonist, Cl-IB-MECA.

Methods

[00244] Chemicals: MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD). Cl-IB-MECA is commercially available from Tocris Biosciences (Bristol, UK) and several other vendors. All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO).

[00245] Animals and traumatic brain injury (TBI): TBI was performed with a controlled closed skull injury model as described in Talley-Watts *et al.* 2012 (*J. Neurotrauma* 30, 55-66). Following the method described therein, a pneumatic impact device was used to generate a moderate TBI leaving the skull and dura matter intact. To achieve this, C57BL/6 mice were anesthetized with isoflurane (3% induction, 1% maintenance) in 100% oxygen. A body temperature of 37°C was maintained using a temperature-controlled heated surgical table. A small midline incision was made on the scalp using aseptic surgical techniques. A 5mm stainless steel disc was positioned on the skull and fixed using superglue on the right parietal bone between bregma and lamda over the somatosensory cortex. The mouse was then positioned on a stage directly under the pneumatic impact tip. A calibrated impact was delivered at 4.5m/s at a depth of 2mm which generates a moderate injury in the mouse. Scalp incisions were closed using 4-0 nylon braided suture and antibiotic ointment applied to the incision. Mice were placed in a Thermo-Intensive Care Unit (Braintree Scientific model FV-1; 37°C; 27% O₂) and monitored until fully awake and moving freely. Thirty minutes following injury or sham (uninjured), mice were treated with either vehicle (saline) or drugs (MRS4322, Cl-IB-MECA or MRS2365). Doses of MRS4322, Cl-IB-MECA and MRS2365 were 0.16, 0.24 and 0.2 mg/kg, respectively, each equivalent to equimolar doses of approximately 0.5 µmol/kg.

[00246] Western Blot Analysis for GFAP: At selected survival times, mice were anesthetized under isoflurane and sacrificed. The brain was removed and placed on ice for dissection into impacted and non-impacted brain hemispheres. The isolated tissue was rapidly homogenized in chilled homogenization buffer (0.32 M Sucrose, 1 mM EDTA, 1 M Tris-HCL pH = 7.8) on ice using a Wheaton glass dounce (20 strokes). The homogenate was transferred to a 2 mL tube and centrifuged at 1000 g for 10 minutes at 4 °C and the supernatant was collected and analyzed. Protein concentration was determined by the BCA assay using a 1:50 dilution. 100 µg of protein was removed as

an aliquot for each sample and Laemmli buffer containing β -mercaptoethanol added and the sample placed in a heat block for 3 minutes at 95 °C. Samples were loaded on a 12% gel and ran at 80 V for 20 minutes followed by 40 minutes at 130 V. Samples were transferred to nitrocellulose membrane at 100 V for 1 hour. The membrane was blocked with 5% milk in TBS-T for 30 minutes. GFAP (1:1000-Imgenex IMG-5083-A) was added and placed at 4 °C overnight. The membrane was washed with TBS-T three times for 10 minutes. Secondary antibody for GFAP (Donkey anti-rabbit HRP conjugated (ImmunoJackson Laboratories; 711-035-152; 1:20000) was applied at room temperature for 1 hour. The membranes were washed with TBS-T for 15 minutes (3 times) and developed using the Western Lightning Plus-ECL kit (PerkinElmer, Inc.) following manufacturer's directions.

Results

[00247] MRS4322 reduces GFAP expression in the mouse brains following TBI. Glial Fibrillary acidic protein (GFAP) expression was used as a biomarker for reactive gliosis after TBI (Talley-Watts *et al.* 2012; Sofroniew, 2005). We performed Western blot analysis for GFAP expression in Sham, TBI or TBI retreated (MRS4322 or MRS2365) mice sacrificed at 7 days post-injury. First, western blot analysis confirmed that TBI induced a significant increase in GFAP expression, both in the Ipsilateral (where the impact was centered) and contralateral sides of the brain at 7 days post-injury (Figure 12A). GFAP expression was significantly lower in blots from mice treated with MRS4322 or MRS2365, which were injected within 30 minutes of the initial trauma (Figure. 12A). For loading controls, beta-actin western blots were used and shown below the respective lanes. Western blots shown in Figure 12A were all obtained from a representative experiment and were run on the same gel. Data averaged from 3 separate experiments and showing the relative change in GFAP/actin ratios (band intensities measured in Image J software) are presented as mean +/- SEM in Figure 12B. Values have been normalized to TBI levels at day 7 (100%) in order to consolidate the data. The total number of mice for a given experimental treatment is indicated by N in Figure 12B.

[00248] MRS4322 and an Adenosine A3 Receptor agonist reduce GFAP expression in the mouse plasma following TBI. GFAP levels in the plasma have also been used as a biomarker for TBI, due to the breakdown of the blood brain barrier (BBB)

after a trauma. Consequently, we also collected plasma samples at day 7 from TBI mice. Similar to brain tissue, we found that GFAP levels were easily detected at day 7 by western blot analysis (Figure 12C). More importantly, we found that western blots obtained the plasma of TBI-mice treated with either MRS4322 or Cl-IB-MECA (Adenosine A₃ receptor agonist) exhibited significantly reduced levels of GFAP relative to actin (Figure 12C). Histogram plots of the average GFAP/actin ratios for each experimental condition are presented in Figure 12D. The total number of mice per experimental treatment is indicated by the value N.

[00249] MRS4322 is a low-affinity (4900 nM) agonist of the A₃ receptor in the mouse. Conversely, Cl-IB-MECA is a high-affinity (0.18 nM) agonist in the mouse - the differences in affinity of these two compounds is approximately 25,000-fold. However, in the mouse photothrombotic stroke and TBI models, MRS4322 demonstrates significant efficacy that is blocked by the A₃ antagonist MRS1523, whereas Cl-IB-MECA is either inactive (stroke) or weakly active (TBI, Figure 12). This is clearly a non-obvious result from the perspective of receptor affinity. Our current explanation for this finding is based on proprietary ADME/PK data we have generated for MRS4322 and Cl-IB-MECA. Cl-IB-MECA is a lipophilic compound (cLogP approx 2.5) that is highly bound to plasma proteins (free fraction 0.002) and highly bound nonspecifically to brain tissue (free fraction 0.002). MRS4322 is a very hydrophilic compound (cLogP <0) that has a very large unbound fraction in plasma (0.74) and brain (0.13). Only unbound drug is available for distribution across membranes and interaction with receptors. Thus, despite its lower receptor affinity, the fraction of MRS4322 available to interact with the A₃ receptor in these mouse models is at least 1000-fold higher than that of Cl-IB-MECA. We believe that these significant differences in compound physicochemical properties and ADME/PK characteristics contribute to the non-obvious efficacy of MRS4322 as compared to Cl-IB-MECA (and MRS5698, another lipophilic and highly-bound/high-affinity full A₃R agonist) in these mouse models.

[00250] Biased Agonism. The adenosine A₃ receptor is a G protein-coupled pleiotropic receptor, i.e. agonism of this receptor potentially activates multiple downstream pathways via multiple G proteins as well as Beta-arrestin. Pathways that are activated by A₃ receptor agonism have currently been identified, but may not be limited


to, Gq11-mediated intracellular calcium mobilization, Gi-mediated modulation of cAMP production, and Gi-mediated phosphorylation of ERK1/2 and Akt. One aspect of our discoveries is in the A₃-mediated mobilization of intracellular calcium resulting in promotion of mitochondrial ATP production in astrocytes.

[00251] An emerging concept in receptor pharmacology is biased agonism. This concept states that for pleiotropic receptors there are actually different classes of agonists, some of which may activate all downstream pathways while others demonstrate bias in activating a subset of the downstream pathways. In drug discovery and receptor pharmacology, biased agonism introduces the possibility of increased specificity in pathway activation with fewer off-target effects, i.e. fewer side-effects. There is evidence for biased agonism for the A₃ receptor. However, prototypical high-affinity agonists such as Cl-IB-MECA and MRS5698 are full agonists that do not demonstrate biased activation of the aforementioned downstream pathways. Accordingly, and without wishing to be bound by any particular theory, it is believed that MRS4322 is a biased agonist that preferentially activates intracellular calcium mobilization with less/no activation of the other A₃-mediated pathways. This finding explains the MRS4322 efficacy observed in mouse stroke and TBI models relative to the full/unbiased agonists Cl-IB-MECA and MRS5698.

Example 6: Neuroprotective Efficacy of MRS4322 After Stroke in Mice

Purpose

[00252] This study was designed to determine the neuroprotective efficacy of MRS4322 in mice subjected to stroke and compare mice treated with MRS2365, with and without the A₃ receptor antagonist MRS1523, and in comparison to full A₃R agonists MRS5698 and Cl-IB-MECA. MRS1523 has the following structure:



Methods

[00253] Chemicals: MRS4322 was obtained courtesy of Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD). Cl-IB-MECA, MRS5698 and MRS2365 are commercially available from Tocris Bioscience (Bristol, UK) and several other vendors. All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO).

[00254] Photothrombosis-induced Stroke: Photothrombosis was performed as described in Zheng et al 2010 (PloS One 5 (12): e14401). In brief, Rose Bengal is a fluorescent dye that when injected into the vasculature and excited, generates singlet oxygen that damages the endothelial wall and induces a local thrombosis (clot). Using this technique, mice were given a 0.1 mL tail-vein injection of sterilized Rose Bengal (RB, Sigma, U.S.A.) in artificial cerebral spinal fluid (aCSF). The RB concentration was 20 mg/mL. A cortical region was centered in the imaging field and illuminated with a green laser (543 nm, 5 mW) using a 0.8-NA 40x water-immersion objective (Nikon, Tokyo). The clot formation was monitored in real time until the targeted vessel or downstream capillaries were firmly occluded. Stable clots were subsequently identified by a non-fluorescent vessel segmentation ending with highly fluorescent regions. In control experiments, either laser illumination or Rose Bengal itself did not lead to clot formation. Treatments, MRS4322 (0.16 mg/kg; 0.5 μ mol/kg) or MRS2365 (0.24 mg/kg; 0.5 μ mol/kg) were introduced via intraperitoneal injections (i.p.). For experiments with A₃ receptor antagonist MRS1523, mice were administered intraperitoneal injections (2 mg/kg) at the 0 and 2 hour timepoints to ensure receptor antagonism throughout the course of the study.

[00255] Animals and Photothrombosis-induced Stroke: Stroke was performed as described in Zheng et al 2010 (PloS One 5 (12): e14401). Female C57Bl/6 mice (4-6 months) were used in this study. From the methods of this manuscript: Mice were anesthetized at 3% isoflurane with 100% oxygen and subsequently maintained at 1% isoflurane through a nosecone. Depth of anesthesia was monitored and regulated according to vital signs, pinch withdrawal and eye blinks. Body temperature was maintained at 37 °C by a feedback-controlled heating pad (Gaymar T/Pump). Vital signs including oxygen saturation, respiratory rate, and heart rate were continuously monitored by using the MouseOx system (STARR Life Sciences). The hair on each mouse's head

was trimmed and a small incision was made in the scalp to expose the skull. A custom-made stainless steel plate was glued to the skull with VetBond Tissue Adhesive (3M, St Paul, MN). A cranial thinned-skull imaging window was created over the right primary somatosensory cortex (~1.5 mm posterior to Bregma and 2 mm lateral from midline) depending on the experiment. In brief, a large region of the skull was first thinned with the electric drill and then further thinned with a surgical blade. The final thickness of the thinned skull was approximately 50 μm . After the cranial imaging window was created, mice were transferred to microscope stage and used for photothrombosis or imaging experiments. For the repeat imaging experiments, the plate was carefully detached from the skull and the scalp was sutured (Ethicon 6-0 sild suture). After each experiment, the mice were either returned to cages until the next timepoint or sacrificed. All procedures were approved by the Institutional Animal Care and Use Committee (IACUC) at University of Texas Health Science Center at San Antonio. Thirty minutes following stroke or sham (uninjured), mice were treated with either vehicle (saline) or drugs (MRS4322, MRS2365, MRS5698 or Cl-IBMECA).

[00256] Post photothrombotic infarction evaluation. The size of cerebral infarcts was evaluated using 2,3,5-Triphenyltetrazolium chloride (TTC) staining as described in Zheng et al 2010 (PloS One 5 (12): e14401). In brief, RB-induced lesions in brain slices were stained with TTC. TTC is a colorless dye that stains healthy brain tissue red when reduced by the mitochondrial enzyme succinyl dehydrogenase (Bederson JB et al., 1986). The absence of staining in necrotic tissue is then used to define the area of a brain infarction. Mice were sacrificed by cervical dislocation, their brains removed and then placed in ice cold HBSS for 3 minutes. The brain was subsequently transferred to a brain mold (KOPF), sliced into 1 mm sections and immersed in 2% TTC (5 min) at 37 °C. The sections were fixed in 10% buffered formaldehyde solution overnight at 4 °C. Slices were imaged on a flatbed scanner (HP scanjet 8300) for analysis of the lesion size at 1200 dpi.

Results

[00257] MRS4322 treatment reduces brain infarctions after stroke. Multivessel photothrombotic strokes were induced in mice using tail-vein injected in conjunction with RB as described above. Within 30 minutes of clot formation, mice were injected

intraperitoneally with either vehicle (saline control), MRS4322 (0.16 mg/kg; 0.5 μ mol/kg) or MRS2365 (0.24 mg/kg; 0.5 μ mol/kg). Twenty-four hours after the initial stroke, the brain infarction size was evaluated with TTC staining as described above. Representative TTC-stained brain slices are presented in Figure 13A. We found that both MRS4322 and MRS2365 significantly reduced the size of the brain infarction. A histogram plot of the average size of brain infarctions for vehicle, MRS4322, MRS2365, MRS5698 or Cl-IBMECA-treated mice is presented in Figure 13C. These data are pooled from 2 independent experiments. N refers to the total number of mice tested.

[00258] The A₃ Receptor Antagonist MRS1523 inhibits neuroprotection of MRS4322 and MRS2365 treatments after stroke. Multivessel photothrombotic strokes were induced in mice as described above. However, in this experiment, mice were treated with intraperitoneal injections of the A₃ receptor antagonist, MRS1523 (2 mg/kg) at the 0 and 2 hour timepoints to ensure receptor antagonism. Mice were then injected with either vehicle, MRS4322, MRS2365, MRS5698 or Cl-IBMECA within 30 minutes of clot formation at the concentrations described above. Twenty-four hours later, brain infarction sizes were evaluated with TTC staining. Representative TTC-stained brain slices are presented in Figure 13B. We found that the size of brain infarctions in mice pretreated with MRS1523 were not reduced by treatments with either MRS4322 or MRS2365, or MRS5698. A histogram plot of the average size of brain infarctions for these experiments is presented in Figure 13D. Data are pooled from 2 independent experiments. N refers to the total number of mice tested.


Example 7: Neuroprotective Efficacy of MRS1873 After Stroke in Mice

Purpose

[00259] This study was designed to determine the neuroprotective efficacy of MRS1873, which is the corresponding 2-chloro analog of MRS4322. Experiments were performed in mice subjected to stroke and compared mice treated with MRS4322 and vehicle treated.

Methods

[00260] Chemicals: MRS1873 and MRS4322 were obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD.). MRS1873 has the following structure:

**MRS1873**

[00261] Photothrombosis-induced Stroke: Photothrombosis was performed as described in Zheng et al. 2010 (PloS One 5 (12): e14401). In brief, Rose Bengal is a fluorescent dye that when injected into the vasculature and excited, generates singlet oxygen that damages the endothelial wall and induces a local thrombosis (clot). Using this technique, mice were given a 0.1ml tail-vein injection of sterilized Rose Bengal (RB, Sigma, U.S.A.) in artificial cerebral spinal fluid (aCSF). The RB concentration was 20 mg/ml. A cortical region was centered in the imaging field and illuminated with a green laser (543 nm, 5mW) using a 0.8-NA 40x water-immersion objective (Nikon, Tokyo). The clot formation was monitored in real time until the targeted vessel or downstream capillaries were firmly occluded. Stable clots were subsequently identified by a non-fluorescent vessel segmentation ending with highly fluorescent regions. In control experiments, either laser illumination or Rose Bengal itself did not lead to clot formation. Treatments, MRS1873 (100 ul of 100 uM) or MRS4322 (100 ul of 100 uM) were introduced via intraperitoneal injections (i.p.).

[00262] Animals and Photothrombosis-induced Stroke: Stroke was performed as described in Zheng et al 2010 (PloS One 5 (12): e14401). Female C57Bl/6 mice (4-6 months) were used in this study. From the methods of this manuscript: Mice were anesthetized at 3% isoflurane with 100% oxygen and subsequently maintained at 1% isoflurane through a nosecone. Depth of anesthesia was monitored and regulated according to vital signs, pinch withdrawal and eye blinks. Body temperature was maintained at 37 °C by a feedback-controlled heating pad (Gaymar T/Pump). Vital signs including oxygen saturation, respiratory rate, and heart rate were continuously monitored by using the MouseOx system (STARR Life Sciences). The hair on the mouse head was trimmed and a small incision was made in the scalp to expose the skull. A custom-made stainless steel plate was glued to the skull with VetBond Tissue Adhesive (3M, St

Paul, MN). A cranial thinned-skull imaging window was created over the right primary somatosensory cortex (~1.5 mm posterior to Bregma and 2 mm lateral from midline) depending on the experiment. In brief, a large region of the skull was first thinned with the electric drill and then further thinned with a surgical blade. The final thickness of the thinned skull was approximately 50 μm . After the cranial imaging window was created, mice were transferred to microscope stage and used for photothrombosis or imaging experiments. For the repeat imaging experiments, the plate was carefully detached from the skull and the scalp was sutured (Ethicon 6-0 sild suture). After each experiment, the mice were either returned to cages until the next timepoint or sacrificed. All procedures were approved by the Institutional Animal Care and Use Committee (IACUC) at University of Texas Health Science Center at San Antonio. Thirty minutes following stroke or sham (uninjured), mice were treated with either vehicle (saline) or drugs (MRS4322 or MRS2365).

[00263] Post photothrombotic infarction evaluation. The size of cerebral infarcts was evaluated using 2,3,5-Triphenyltetrazolium chloride (TTC) staining as described in Zheng et al 2010 (PloS One 5 (12): e14401). In brief, RB-induced lesions in brain slices were stained with TTC. TTC is a colorless dye that stains healthy brain tissue red when reduced by the mitochondrial enzyme succinyl dehydrogenase (Bederson JB et al., 1986). The absence of staining in necrotic tissue is then used to define the area of a brain infarction. Mice were sacrificed by cervical dislocation, their brains removed and then placed in ice cold HBSS for 3 minutes. The brain was subsequently transferred to a brain mold (KOPF), sliced into 1 mm sections and immersed in 2% TTC (5 min) at 37 °C. The sections were fixed in 10% buffered formaldehyde solution overnight at 4 °C. Slices were imaged on a flatbed scanner (HP scanjet 8300) for analysis of the lesion size at 1200 dpi.

Results

[00264] MRS1873 treatment reduces brain infarctions after stroke: Multivessel photothrombotic strokes were induced in mice using tail-vein injected in conjunction with RB as described above. Within 30 minutes of clot formation, mice were injected intraperitoneal with either vehicle (saline control), MRS1873 (100 μl of 100 μM) or MRS4322 (100 μl of 100 μM). Twenty-four hours after the initial stroke, the brain infarction size was evaluated with TTC staining as described above. Representative

TTC-stained brain slices are presented in Figure 14B. We found that both MRS1873 and MRS4322 significantly reduced the size of the brain infarction. A histogram plot of the average size of brain infarctions for vehicle, MRS1873 and MRS4322-treated mice is presented in Figure 14B. These data are pooled from 3 independent experiments. N refers to the total number of mice tested.

Pharmacokinetics and Rationale for Efficacy

[00265] MRS1873 is the 2-Cl analog of MRS4322 and is also an adenosine A₃ agonist. MRS1873 physicochemical properties are identical to those of MRS4322 in terms of low molecular weight, hydrophilicity (cLogP <0) and topological polar surface area. Since ADME/PK parameters such as plasma and brain binding, clearance and volume of distribution are driven by these physicochemical properties, we have demonstrated similar pharmacokinetics for MRS4322 and MRS1873. In addition, we have demonstrated similar efficacy for MRS4322 and MRS1873 in a murine photothrombotic model of stroke.

Example 8: Experimental Protocol for Determining Biased Agonism of Compounds at A₃ Adenosine Receptor (A₃R)

[00266] The following assay may be used to determine whether a disclosed compound, such as MRS4322 or MRS1873, exhibits biased agonism (also known as functional selectivity or agonist trafficking) at the A₃ receptor.

[00267] **Materials.** Fluo-4, Dulbecco's modified Eagle's medium (DMEM), and penicillin-streptomycin may be purchased from Invitrogen (Carlsbad, CA). Adenosine deaminase (ADA) and hygromycin-B may be purchased from Roche (Basel, Switzerland). Fetal bovine serum (FBS) may be purchased from ThermoTrace (Melbourne, Australia). AlphaScreen SureFire extracellular signal-regulated kinases 1 and 2 (ERK1/2), Akt 1/2/3, and cAMP kits may be obtained from PerkinElmer (Boston, MA). All compounds prefixed with MRS may be synthesized as described previously (Tosh et al., 2012a,b). All other reagents were purchased from Sigma-Aldrich (St. Louis, MO).

[00268] **Cell Culture.** The sequence of the human A₃R may be cloned into the Gateway entry vector, pDONR201, and then transferred in the Gateway destination

vector, pEF5/ FRT/V5-dest, using methods described previously (Stewart et al., 2009). A₃-FlpIn-CHO cells may be generated using methods described previously (May et al., 2007) and maintained at 37 °C in a humidified incubator containing 5% CO₂ in DMEM supplemented with 10% FBS and the selection antibiotic hygromycin-B (500 µg/ml). For cell survival, ERK1/2 phosphorylation, Akt 1/2/3 phosphorylation, and calcium mobilization assays, cells may be seeded into 96-well culture plates at a density of 4 x 10⁴ cells/ well. After 6 hours, cells are washed with serum-free DMEM and maintained in serum-free DMEM for 12–18 hours at 37 °C in 5% CO₂ before assaying. For cAMP assays, cells may be seeded into 96-well culture plates at a density of 2 x 10⁴ cells/well and incubated overnight at 37°C in 5% CO₂ prior to assay.

[00269] Cell Survival Assays. Media is removed and replaced with HEPES-buffered saline solution (10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), 146 mM NaCl, 10 mM D-glucose, 5 mM KCl, 1mM MgSO₄, 1.3 mM CaCl₂, and 1.5 mM NaHCO₃, pH 7.45) containing ADA (1 U/ml) and penicillin-streptomycin (0.05 U/ml) in the absence and presence of A₃R ligands. Plates are then maintained at 37 °C in a humidified incubator for 24 hours, after which 5 mg/ml propidium iodide is added to cells. Plates may be then read on an EnVision plate reader (PerkinElmer), with excitation and emission set to 320 nm and 615 nm, respectively. Data will be normalized to 100% cell survival and 0% cell survival, determined at t = 0 hours in HEPES buffer and t = 24 hours in Milli-Q water, respectively.

[00270] ERK1/2 and Akt 1/2/3 Phosphorylation Assays. A concentration-response curve of ERK1/2 and Akt 1/2/3 phosphorylation for each ligand may be performed in serum-free DMEM containing 1 U/ml ADA (5-minute exposure at 37°C). Agonist stimulation may be terminated by removal of media and the addition of 100 µl of SureFire lysis buffer to each well. Plates are then agitated for 5 minutes. Detection of pERK1/2 may involve an 80:20:120:1:1 v/v/v/v/v dilution of lysate: activation buffer: reaction buffer: AlphaScreen acceptor beads: AlphaScreen donor beads in a total volume of 11 µl in a 384-well ProxiPlate. Plates may be incubated in the dark at 37°C for 1 hour followed by measurement of fluorescence by an EnVision plate reader (PerkinElmer) with excitation and emission set to 630 nm and 520–620 nm, respectively. Detection of Akt 1/2/3 phosphorylation may employ a 40:9.8:39.2:1 v/v/v/v dilution of lysate:

activation buffer: reaction buffer: AlphaScreen acceptor beads in a total volume of 9 l in a 384-well Proxiplate. Plates may be incubated in the dark at room temperature for 2 hours, after which a 19:1 v/v dilution of dilution buffer: AlphaScreen donor beads may be added in a total volume of 11 µl. Plates may be incubated at room temperature for a further 2 hours, followed by measurement of fluorescence by an EnVision plate reader (PerkinElmer) with excitation and emission set to 630 nm and 520–620 nm, respectively. Agonist concentration-response curves are normalized to the phosphorylation mediated by 10% FBS (5-minute stimulation).

[00271] Calcium Mobilization Assays. Media may be removed from 96-well plates and replaced with HEPES-buffered saline solution containing 1 U/ml ADA, 2.5 mM probenecid, 0.5% bovine serum albumin (BSA), and 1 M Fluo4. Plates may be incubated in the dark for 1 hour at 37 °C in a humidified incubator. A FlexStation plate reader (Molecular Devices, Sunnyvale, CA) may perform the addition of HEPES-buffered saline solution in the absence and presence of agonist and measured fluorescence (excitation, 485 nm; emission, 520 nm) every 1.52 seconds for 75 seconds. The difference between the peak and baseline fluorescence may be measured as a marker for intracellular Ca²⁺ mobilization. A₃R agonist concentration-response curves may be normalized to the response mediated by 100 µM ATP to account for differences in cell number and loading efficiency.

[00272] Inhibition of cAMP Accumulation Assays. Media may be replaced with a stimulation buffer (140 mM NaCl, 5 mM KCl, 0.8 M MgSO₄, 0.2 mM Na₂HPO₄, 0.44 mM KH₂PO₄, 1.3 mM CaCl₂, 5.6 mM D-glucose, 5 mM HEPES, 0.1% BSA, 1 U/ml ADA, and 10 µM rolipram, pH 7.45) and incubated at 37 °C for 1 hour. Inhibition of cAMP accumulation may be assessed by preincubation of A₃-FlpIn-CHO cells with A₃R agonists for 10 minutes, after which 3 µM forskolin was added for a further 30 minutes. The reaction may be terminated by rapid removal of buffer and addition of 50 µl ice-cold 100% ethanol. Ethanol is allowed to evaporate before the addition of 50 µl detection buffer (0.1% BSA, 0.3% Tween-20, 5 mM HEPES, pH 7.45). Plates are agitated for 10 minutes, after which 10 µl lysate was transferred to a 384-well Optiplate. Detection may employ addition of a 5 µl 1:49 v/v dilution of AlphaScreen acceptor beads: stimulation buffer. Following this, a 15 µl 1:146:3 v/v/v dilution of AlphaScreen donor beads:

detection buffer: 3.3 U/ μ l biotinylated cAMP to form a total volume of 30 μ l. The donor bead/biotinylated cAMP mixture may be equilibrated for 30 minutes prior to addition. Plates may be incubated overnight in the dark at room temperature, followed by measurement of fluorescence by an EnVision plate reader (PerkinElmer) with excitation and emission set to 630 nm and 520–620 nm, respectively. Agonist concentration-response curves may be normalized to the response mediated by 3 μ M forskolin (0%) or buffer (100%) alone.

[00273] Molecular Modeling. Docking simulations can be performed for all the compounds investigated in this study using homology models of the human A₃R. In particular, three previously reported models can be used: a model entirely based on an agonist-bound hA_{2A}AR crystal structure (PDB ID: 3QAK), a model based on a hybrid A_{2A}AR- β 2 adrenergic receptor template, and a model based on a hybrid A_{2A}AR-opsin template (β 2 adrenoceptor X-ray structure PDB ID: 3SN6; opsin crystal X-ray crystal structure PDB ID: 3DQB) (Tosh et al., 2012a). Models based on hybrid templates will show an outward movement of TM2 compared with the A_{2A}AR-based model. Structures of A₃R ligands may be built and prepared for docking using the Builder and the LigPrep tools implemented in the Schrödinger suite (Schrödinger Release 2013-3, Schrödinger, LLC, New York, NY, 2013). Molecular docking of the ligands at the A₃R models may be performed by means of the Glide package part of the Schrödinger suite. In particular, a Glide Grid may be centered on the centroid of some key residues of the binding pocket of adenosine receptors, namely, Phe (EL2), Asn (6.55), Trp (6.48), and His (7.43). The Glide Grid may be built using an inner box (ligand diameter midpoint box) of 14 Å x 14 Å x 14 Å and an outer box (box within which all the ligand atoms must be contained) that extends 25 Å in each direction from the inner one. Docking of ligands may be performed in the rigid binding site using the XP (extra precision) procedure. The top scoring docking conformations for each ligand may be subjected to visual inspection and analysis of protein-ligand interactions to select the proposed binding conformations in agreement with the experimental data.


[00274] Data Analysis. Statistical analyses and curve fitting may be performed using Prism 6 (GraphPad Software, San Diego, CA). To quantify signaling bias, agonist concentration-response curves may be analyzed by nonlinear regression using a

derivation of the Black-Leff operational model of agonism, as described previously (Kenakin et al., 2012; Wootten et al., 2013; van der Westhuizen et al., 2014). The transduction coefficient, τ/K_A [expressed as a logarithm, $\text{Log}(\tau/K_A)$], may be used to quantify biased agonism. To account for cell-dependent effects on agonist response, the transduction ratio may be normalized to the values obtained for the reference agonist, IB-MECA, to generate $\text{ALog}(\tau/K_A)$. To determine the bias for each agonist at different signaling pathways, the $\text{ALog}(\tau/K_A)$ will be normalized to a reference pathway, pERK1/2, to generate $\text{AALog}(\tau/K_A)$. Bias may be defined as $10^{\text{AALog}(\tau/K_A)}$ where a lack of bias will result in values that are not statistically different from 1, or 0 when expressed as a logarithm. All results may be expressed as the mean $\pm$ S.E.M. Statistical analyses would involve an F test or a one-way analysis of variance with a Tukey or Dunnett's post hoc test, with statistical significance determined as P, 0.05.


Example 9: Synthetic Route for MRS4322

[00275] MRS4322 and similar compounds such as MRS1873 may be prepared according to methods known in the art. For example, MRS4322 may be prepared from D-ribose by following routes described in Choi, W. J. et al. J. Org. Chem. 2004, 69, 2634-2636, Tosh, D. K. et al. Purinergic Signalling 2015, 11, 371-387; and Chem. Eur. J., 2009, 15, 6244-6257. Schemes 1 and 2 below show the synthetic route.


Scheme 1: Synthesis of MRS4322



[00276] Zhan cat-1B has the following structure:



[00277] Scheme 2 shows the remainder of the synthesis.

Scheme 2: Synthesis of MRS4322 (continued)**Example 10: Rationale for Cardioprotection of Disclosed Compounds**

[00278] We have demonstrated the extremely rapid and quantitative dephosphorylation of P2Y1 agonists to compounds, such as, MRS4322 that are adenosine A₃ agonists. These adenosine A₃ agonists were surprisingly found to be efficacious in multiple murine models of stroke and traumatic brain injury, as described herein. Without wishing to be bound by any particular theory, it is believed that presently described A₃ agonists, such as MRS4322 and MRS1873, are effective as cardioprotective agents.

[00279] Our data showing rapid and quantitative dephosphorylation of MRS2365 and related phosphorylated nucleosides to compounds such as MRS4322 and MRS1873 supports our claim that the purported cardioprotective efficacy of these P2X₄ agonists is actually due to the adenosine A₃ agonism of their dephosphorylated metabolites. Indeed, we have demonstrated that MRS1873 is an adenosine A₃ agonist that is efficacious in stroke and traumatic brain injury models.

Example 11: Pharmacokinetics and Binding of MRS4322 Following Intravenous Administration to Neonatal Pigs

Purpose

[00280] This study was designed to determine the plasma, brain and CSF concentrations of MRS4322 following intravenous administration to neonatal pigs.

Methods

[00281] **Chemicals.** MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD.).

[00282] **Animals.** Four-week old female neonatal pigs weighing approximately 7.5 Kg were used for this study, supplied by the Department of Bioengineering, University of Pennsylvania (Philadelphia, PA.). Animals were equipped with brain microdialysis probes to obtain brain extracellular fluid samples for drug concentration determinations during the study. All studies were conducted under approved University of Pennsylvania IACUC protocols.

[00283] **Drug Administration:** MRS4322 was solubilized in DMSO and then diluted in saline to prepare dosing solution. A 10 mL volume of dosing solution was administered by intravenous bolus administration to each neonatal pig (n=3).

[00284] **Tissue Sampling:** Blood samples were obtained at 0.25, 0.5, 1, 2, 4 and 6 hours post-dose. Brain extracellular fluid samples were obtained from implanted microdialysis probes at 1, 4 and 6 hours post-dose. Whole blood (1 mL) was obtained at each timepoint and placed in vacutainer tubes containing heparin and immediately centrifuged for preparation of plasma; plasma was stored at -80 °C. Brain extracellular and cerebrospinal fluid samples were stored at -80 °C. At the time of euthanasia (6 hours post-dose), cerebrospinal fluid samples were obtained and frozen, while brain samples from the cortex and hippocampus were obtained by decapitation, rinsed in ice-cold phosphate-buffered saline and weighed. Brain samples were then immediately flash-frozen in liquid nitrogen and stored at -80 °C.

Bioanalysis

[00285] Plasma, brain, brain extracellular fluid and cerebrospinal fluid concentrations of MRS4322 were determined by LC-MS/MS utilizing tolbutamide as an internal standard. The following table outlines the LC and MS/MS conditions employed:

Table 12: Bioanalytical Methods for MRS4322 Plasma, Brain, Brain Extracellular and Cerebrospinal Fluid Concentration Determinations

System Components			
Module	Manufacturer	Model	
LC	Waters	Acquity I Class	
Autosampler	Waters	Acquity Sample Manager	
MS Detection	AB Sciex	API 5500 QQQ	
HPLC Method			
Column	Phenomenex Kinetex C18 (2.1 x 50 mm, 2.6 μm)		
Elution	Gradient, 0.6 mL/min		
	Mobile Phase A: 0.1% Formic Acid		
	Mobile Phase B: 0.1% Formic Acid in Acetonitrile		
MS Detection and Calibration for MRS4322 in Swine Plasma			
Peak Name: Tolbutamide			
Use as Internal Standard			
Q1/Q3 Masses: 271.00/155.00 Da			
Peak Name: MRS4322			
Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.08/182.10 Da			
Fit	Linear	Weighting	1 / x
Intercept	0.000570		
Slope	0.00699		
Correlation coefficient	0.9998		
Use Area			
MS Detection and Calibration for MRS4322 in Swine Brain Homogenate			
Peak Name: Tolbutamide			
Use as Internal Standard			
Q1/Q3 Masses: 271.00/155.00 Da			
Peak Name: MRS4322			
Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.08/182.10 Da			
Fit	Quadratic	Weighting	1 / x
a0	0.000244		
a1	0.00675		
a2	-0.00000134		
Correlation coefficient	0.9999		
Use Area			
MS Detection and Calibration for MRS4322 in Swine ECF and CSF			
Peak Name: Tolbutamide			
Use as Internal Standard			
Q1/Q3 Masses: 271.00/155.00 Da			
Peak Name: MRS4322			
Internal Standard: Tolbutamide			
Q1/Q3 Masses: 324.08/182.10 Da			
Fit	Quadratic	Weighting	1 / x
a0	0.000297		
a1	0.00783		
a2	0.000000130		
Correlation coefficient	0.9998		
Use Area			

[00286] For each tissue matrix, standard curves were created and LLOQ/ULOQ concentrations determined. The calibration ranges for all of the MRS4322 matrices' standard curves were 0.1-1000 ng/mL.

[00287] For bioanalysis of brain concentrations of MRS4322, brain samples were homogenized in ice-cold phosphate-buffered saline in a 4x dilution. Aliquots of the resulting diluted brain homogenate were treated with acetonitrile and analyzed by LC-MS/MS.

Results

[00288] Following intravenous administration to neonatal pigs, MRS4322 concentrations were detectable in plasma, brain, brain extracellular fluid and cerebrospinal fluid samples (Figure 1B and Figure 16, Table 13).

Table 13: Plasma, Brain, Brain Extracellular and Cerebrospinal Fluid Concentrations of MRS4322 in Neonatal Pigs Following Intravenous Administration

Test Article	Matrix	Route	Dose (mg/kg)	Time (h)	Plasma Concentration (ng/mL) by Subject					
					161128U	161128U-1	161128U-2	Mean	SD	n
MRS4322	Plasma	IV	0.2	0	BLQ	BLQ	BLQ	BLQ	---	0
				0.25	133	120	163	139	22	3
				0.5	85.3	64.1	98.7	79.4	22.9	3
				1	58.9	51.4	63.0	57.8	5.9	3
				2	27.2	18.7	46.8	28.9	14.4	3
				4	23.3	21.9	26.7	24.8	3.6	3
Test Article	Matrix	Route	Dose (mg/kg)	Time (h)	Brain Concentration (ng/g) by Subject					
					161128U	161128U-1	161128U-2	Mean	SD	n
					5.34	3.80	6.16	5.10	1.19	3
					6.47	4.56	5.39	5.47	0.96	3
					Extracellular Fluid Concentration (ng/mL) by Subject					
					161128U	161128U-1	161128U-2	Mean	SD	n
MRS4322	Matrix	Route	Dose (mg/kg)	Time (h)	1.55	0.641	1.70	1.30	0.57	3
					4	0.693	0.579	1.36	0.941	3
					6	0.602	0.459	1.08	0.714	3
					Cerebro-Spinal Fluid Concentration (ng/mL) by Subject					
					161128U	161128U-1	161128U-2	Mean	SD	n
					1.37	1.04	1.75	1.39	0.36	3
MRS4322	CSF	IV	0.2	6	1.37	1.04	1.75	1.39	0.36	3

BLQ = Below the Lower Limit of Quantitation (0.0975 ng/mL for Plasma)

Table 14: Plasma Pharmacokinetics of MRS4322 in Neonatal Pigs Following Intravenous Administration

Test Article	Route	Dose (mg/kg)	Animal ID	C0 (ng/mL)	Cmax (ng/mL)	Tmax (hr)	AUClast (hr*ng/mL)	AUCINF (hr*ng/mL)	Cl (mL/min/kg)	Vz (L/kg)	Half Life (hr)
MRS4322	IV	0.2	161128U	297	133	0.25	236	344	0.68	4.04	4.82
			161128U-1	286	120	0.25	203	233	14.3	2.86	2.31
			161128U-2	289	163	0.25	307	401	0.31	2.25	3.13
N				3	3	3	3	3	3	3	
Mean				248	139	0.25	249	326	10.8	3.05	3.42
SD				34.9	22.1	0.00	52.9	85.3	3.13	0.907	1.28
CV%				14.1	15.9	0.0	21.2	26.1	29.1	29.8	37.4

[00289] Plasma concentrations allowed initial estimates of T_{max}, C_{max}, plasma clearance, volume of distribution, half-life and AUC (Table 14).

[00290] Although MRS4322 concentrations in the brain, brain extracellular fluid and cerebrospinal fluid were detectable, data was insufficient for estimation of half-life or other pharmacokinetic parameters other than C_{max} and T_{max}. However, based on the available plasma and brain data at 6 hours post-dose when samples were obtained for all matrices, it was estimated that the brain/plasma ratio of total drug was approximately 0.3 based on mean concentrations in plasma and brain.

[00291] These results confirm that circulating plasma concentrations of MRS4322 are detectable following intravenous administration to neonatal pigs, and that MRS4322 is well distributed to the brain under these dosing conditions.

Example 12: Plasma and Brain Binding of MRS4322 in the Neonatal Pigs

Purpose

[00292] This study was designed to determine the plasma and brain free fraction of MRS4322 in neonatal pigs.

Methods

[00293] **Chemicals.** MRS4322 was obtained courtesy Dr. Ken Jacobson of the National Institute of Diabetes, Digestive and Kidney Diseases (Bethesda, MD.). Analytical-grade sulfamethoxazole and warfarin were obtained from commercial supplies at Seventh Wave Laboratories (Maryland Heights, MO.). All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO.).

[00294] **Animals and Tissue Preparation.** Plasma and brain samples from female neonatal pigs were obtained from the University of Pennsylvania and stored at -80 °C until use.

[00295] Plasma ultrafiltrate blank samples were prepared by thawing frozen plasma and then pre-warming plasma in a humidified 5% CO₂ chamber at 37°C for 60 minutes. Aliquots of 800 µl were transferred to Centrifree Centrifugal Filters (Ultracel regenerated cellulose (NMWL 30,000 amu) Lot R5JA31736) and centrifuged at 2900 RPM at 37°C for 10 minutes; plasma water filtrates were collected and used in preparation of standards, blanks and QC standards.

[00296] Brains were weighed and homogenized with 1:9 phosphate-buffered saline, pH 7.4 using an Omni tissue homogenizer. Brains from four mice were homogenized, pooled and mixed to form one sample.

[00297] **Plasma Binding Determination.** MRS4322, sulfamethaxazole and warfarin were solubilized in DMSO and then diluted in 1:1 acetonitrile:water to prepare 100 µM dialysis stock solutions. Sulfamethaxazole and warfarin were utilized as study standards with known plasma binding values. Plasma samples were pre-warmed for 60 minutes in a humidified, 5% CO₂ incubator maintained at 37 °C. Three ml aliquots of pre-warmed plasma were each spiked with MRS4322, sulfamethaxazole or warfarin using 100 µM stock solutions for each compound resulting in final test concentrations of 1 µM. Spiked plasma samples were incubated on a rotary mixer in a humidified 5% CO₂ chamber at 37 °C for a minimum of 60 minutes. After 60 minutes, three 800 µl aliquots of each sample were added to Centrifree centrifugal filters. The filters were subjected to centrifugation at 2900 rpm for 10 minutes at 37 °C. Three 100 µl aliquots of residual plasma were collected along with ultrafiltrate for bioanalysis.

[00298] **Brain Binding Determination:** MRS4322, sulfamethoxazole and warfarin were solubilized in DMSO and diluted in 1:1 acetonitrile:water to prepare 100 µM dialysis stock solutions. Pooled homogenized brains were pre-warmed for 60 minutes in a humidified, 5% CO₂ incubator maintained at 37 °C. Three ml aliquots of brain homogenate were each spiked with MRS4322, sulfamethaxazole or warfarin using the 100 µM stock solutions for each compound resulting in final spiked concentrations of 1 µM. Spiked pooled brain homogenates were placed on a Nutator mixer in a humidified, 5% CO₂ incubator at 37 °C for 60 minutes. After 60 minutes, three 800 µl aliquots of each sample were added to Centrifree centrifugal filters. The filters were subjected to

centrifugation at 2900 rpm for 10 minutes at 37C. Aliquots of residual brain homogenate and ultrafiltrate were collected for bioanalysis.

Bioanalysis

[00299] Plasma and brain concentrations of MRS4322 in spiked plasma, brain homogenates and associated ultrafiltrates were determined by LC-MS/MS utilizing tolbutamide as an internal standard. Associated concentrations of sulfamethaxazole and warfarin were also determined by LC-MS/MS using standard conditions (data not shown). The following tables outline the LC and MS/MS conditions employed (Tables 15 and 16). Bioanalytical methods were identical for all matrices; standard curve statistics (e.g. Fit, Intercept, Slope, Correlation Coefficient) were determined for each matrix but were not significantly different and thus are not shown for each matrix.

Table 15: Bioanalytical Methods for MRS4322 for Determination of Plasma, Brain Homogenate, Plasma Ultrafiltrate and Brain Homogenate Ultrafiltrate Concentrations

Bioanalysis of Plasma, Brain Homogenates, Plasma Ultrafiltrate and Brain Homogenate Ultrafiltrate Samples			
System Components			
Module	Manufacturer	Model	
HPLC	Shimadzu	Prominence LC20AD Binary Pumps	
Autosampler	CTC Leap	HTC PAL	
Mass Spectrometer	AB Sciex	API 4000	
HPLC Method			
Column	Phenomenex Kinetex C18 (2.1 x 50 mm, 2.6 μm)		
Elution	Gradient, 0.4 mL/min		
	Mobile Phase A: 0.1% Formic acid in Water		
	Mobile Phase B: 0.1% Formic acid in Acetonitrile		
Mass Spectrometer Conditions for MRS4322			
Peak Name: MRS4322			
Q1/Q3 Masses: 324.10/182.00 Da			
Peak Name: Tolbutamide			
Q1/Q3 Masses: 271.20/155.20 Da			
Fit	Linear	Weighting	1 / x
Intercept	0.0181		
Slope	0.00192		
Correlation coefficient	0.9988		

[00300] For each tissue matrix, standard curves were created and LLOQ/ULOQ concentrations determined. The calibration range for the MRS4322 plasma concentration

standard curves was 5-1000 nM. The calibration range for MRS4322 plasma ultrafiltrate standard curves was 5-1000 nM. The calibration range for MRS4322 brain homogenate and brain homogenate ultrafiltrate standard curves were 5-1000 nM and 5-1000 nM, respectively.

Results

[00301] Plasma binding and free fraction were determined for MRS4322 utilizing plasma ultrafiltration. Plasma binding was 21.6% for MRS4322; associated free fraction was 0.784 (Table 16). Binding of the study standards sulfamethaxazole and warfarin were consistent with literature values.

Table 16: Fraction Unbound and Binding of MRS4322 in Neonatal Pig Plasma

Substrate	Fraction Unbound		% Bound		N
	Mean	SD	Mean	SD	
MRS4322	0.784	0.061	21.6	6.1	3
Sulfamethoxazole	0.606	0.016	39.4	1.6	2
Warfarin	0.0159	0.0021	98.4	0.2	3

* Plasma ultrafiltrate concentrations of MRS5698 were BLQ.

[00302] Brain binding and free fraction were determined for MRS4322 utilizing brain homogenate ultrafiltration. Brain binding was 74.7% for MRS4322; associated free fraction was 0.253 (Table 17). Binding of the study standards sulfamethaxazole and warfarin were consistent with literature values.

Table 17: Fraction Unbound and Binding of MRS4322 in Neonatal Pig Brain Homogenates

Substrate	Fraction Unbound		% Bound		N
	Mean	SD	Mean	SD	
MRS4322	0.253	0.043	74.7	4.3	3
Sulfamethoxazole	0.309	0.029	69.1	2.9	3
Warfarin	0.189	0.006	81.1	0.6	3

[00303] These data indicate that for a given total plasma or brain concentration, substantial unbound drug concentrations of MRS4322 will be available in the brain for interaction with the adenosine A3 receptor. These findings are consistent with those also observed in the mouse.

Example 13: Pharmacological Characterization of MRS4322

[00304] The compound MRS4322 was investigated in competition binding studies at human and mouse A₃ adenosine receptors recombinantly expressed in Chinese hamster ovary (CHO) cells using cell membrane preparations. [³H]NECA was employed as an A₃ agonist radioligand. The non-selective agonist NECA could be used because CHO cells do not natively express adenosine receptors. Concentration-dependent displacement of the radioligand by MR4322 was determined.

[00305] Additionally cAMP experiments were conducted at CHO cells recombinantly expressing human A₃ or mouse A₃ adenosine receptors, respectively. The non-selective agonist NECA was used as a control.

Results

[00306] MRS4322 showed a K_i value of **1490 ± 410 nM** at human A₃ receptors and a K_i value of **4940 ± 974 nM** at mouse A₃ receptors in radioligand binding studies versus [³H]NECA.

[00307] In functional cAMP accumulation experiments in CHO cells expressing A₃ adenosine receptors, MRS4322 showed agonistic activity with an EC₅₀ value of **3630 ± 370 nM** at the human A₃ receptor, and an EC₅₀ value of **759 ± 170 nM** at the mouse A₃ receptor. EC₅₀ values of agonists at GPCRs are dependent on receptor expression levels. The human A₃ receptor cell line seems to have a lower expression level than the cell lines with the mouse A₃ receptors since higher EC₅₀ values are observed, which was also the case for the control agonist NECA.

Table 18: Affinities of agonists at adenosine receptors stably expressed in CHO cells determined in radioligand binding studies [K_i ± SEM (nM)]

Receptor	Human A ₃ R	Mouse A ₃ R
Agonist	[³ H]NECA	[³ H]NECA
MRS4322	1490 ± 410	4940 ± 974
NECA	6.18 (K_D)^a	15.1 (K_D)^a

^aAlnouri M.W. et al. Selectivity is species-dependent: Characterization of standard agonists and antagonists at human, rat, and mouse adenosine receptors. Purinergic Signal. **2015**, 11, 389-407. The same cell lines were used in the present and in the published study.

Table 19: Potencies of agonists at A₃ adenosine receptors stably expressed in CHO cells determined in cAMP accumulation assays

	Human A ₃ R	Mouse A ₃ R
	EC ₅₀ ± SEM (nM) (efficacy at 100 μM concentration in % relative to maximal effect of NECA)	
Agonist		
MRS4322	3630 ± 370 (70 ± 12)	759 ± 170 (72 ± 18)
NECA	41.8 ± 6.3	6.85 ± 0.88

[00308] **Figure 17** shows competition binding experiments of MRS4322 versus the A₃ agonist radioligand [³H]NECA (10 nM) at human A₃ receptors expressed in CHO cells. The calculated K_i-value for MRS4322 was **1490 ± 410 nM**.

[00309] **Figure 18** shows competition binding experiments of MRS4322 versus the A₃ agonist radioligand [³H]NECA (10 nM) at mouse A₃ receptors expressed in CHO cells. The calculated K_i-value for MRS4322 was **4940 ± 974 nM**. **Figure 19** shows cAMP accumulation experiments of MRS4322 and NECA at human A₃ receptors expressed in CHO cells. The calculated EC₅₀-value for MRS4322 was **3630 ± 370 nM**; for NECA an EC₅₀ value of **41.8 ± 6.3 nM** was determined. **Figure 20** shows cAMP accumulation experiments of MRS4322 and NECA at mouse A₃ receptors expressed in CHO cells. The calculated EC₅₀-value for MRS4322 was **759 ± 170 nM**; for NECA an EC₅₀ value of **6.85 ± 0.88 nM** was determined.

[00310] These results are lower than but related to the known binding data for MRS1873. Human A₃, K_i data: MRS1873 K_i = 353 nM; EC₅₀ = 803 nM, published in:

J. Med. Chem. 2002 45:4471-4484. "Structural Determinants of A₃ Adenosine Receptor Activation: Nucleoside Ligands at the Agonist/Antagonist Boundary," Gao, Z-G. et al.

[00311] An earlier paper lists the human A₃ K_i as 85 nM: Bioorganic and Medicinal Chemistry Letters 2001 11:1333-1337. "Ring-Constrained (N)-Methanocarba Nucleosides as Adenosine Receptor Agonists: Independent 5'-Uronamide and 2'-Deoxy Modifications," Lee, K. et al.


[00312] While a number of embodiments of this invention are described, it is understood that the particular examples described above may be altered using routine experimentation to provide other embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined only by the following claims rather than by the specific embodiments that have been provided.

CLAIMS

We claim:


1. A method of treating an injury, disease, or condition selected from traumatic brain injury (TBI), stroke, a neurodegenerative condition, or a heart or cardiovascular disease, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R).

2. The method of claim 1, wherein the compound is



or a pharmaceutically acceptable salt thereof.

3. The method of claim 1, wherein the compound is



or a pharmaceutically acceptable salt thereof.

4. The method of any one of claims 1-3, wherein the injury, disease, or condition is TBI.

5. The method of claim 4, wherein the TBI is selected from concussion, blast injury, combat-related injury, or a mild, moderate or severe blow to the head.

6. The method of any one of claims 1-3, wherein the injury, disease, or condition is a stroke selected from ischemic stroke, hemorrhagic stroke, subarachnoid hemorrhage, cerebral vasospasm, or transient ischemic attacks (TIA).
7. The method according to any one of claims 4-6, wherein neuroprotection or neurorestoration is increased in the patient as compared with an untreated patient.
8. The method of any one of claims 1-3, wherein the neurodegenerative disease is selected from Alzheimer's Disease (AD), Parkinson's Disease (PD), Huntington's Disease (HD), Multiple Sclerosis (MS), amyotrophic lateral sclerosis (ALS), chronic traumatic encephalopathy (CTE), or a neurodegenerative condition caused by a virus, alcoholism, tumor, toxin, or repetitive brain injuries.
9. The method of claim 8, wherein neurodegenerative disease is Parkinson's Disease.
10. The method of claim 8, wherein the injury, disease, or condition is Alzheimer's Disease, migraine, brain surgery, or a neurological side effect associated with cancer chemotherapy.
11. The method according to any one of claims 1-7, wherein the recovery period after the TBI, stroke, cardiac ischemia, or myocardial infarction is decreased as compared with an untreated patient.
12. The method of any one of claims 1-3, wherein the heart or cardiovascular disease is selected from cardiac ischemia, myocardial infarction, a cardiomyopathy, coronary artery disease, arrhythmia, myocarditis, pericarditis, angina, hypertensive heart disease, endocarditis, rheumatic heart disease, congenital heart disease, or atherosclerosis.
13. The method of claim 12, wherein the heart or cardiovascular disease is cardiac ischemia or myocardial infarction.

14. The method of any one of claims 1-3, wherein the compound is administered chronically to treat stroke, cardiac ischemia, or myocardial infarction during the time period after the injury has occurred as it resolves.

15. The method according to any one of claims 1-14, wherein the A₃R is partially agonized.

16. A method of treating an injury, disease, or condition selected from traumatic brain injury (TBI), stroke, or a neurodegenerative condition, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R), wherein the A₃R is agonized in a biased manner toward neuroprotective functions of the A₃R receptor via preferential activation of intracellular calcium mobilization with less, or no, activation of other A₃R-mediated pathways, or via preferential activation of Gq11-mediated intracellular calcium mobilization, Gi-mediated modulation of cAMP production, or Gi-mediated phosphorylation of ERK1/2 and Akt.

17. The method of claim 16, wherein the injury, disease, or condition is TBI.

18. The method of claim 17, wherein the TBI is selected from concussion, blast injury, combat-related injury, or a mild, moderate or severe blow to the head.

19. The method of claim 16, wherein the injury, disease, or condition is a stroke selected from ischemic stroke, hemorrhagic stroke, subarachnoid hemorrhage, cerebral vasospasm, or transient ischemic attacks (TIA).



20. The method of claim 16, wherein the neurodegenerative disease is selected from Alzheimer's Disease (AD), Parkinson's Disease (PD), Huntington's Disease (HD), Multiple Sclerosis (MS), amyotrophic lateral sclerosis (ALS), chronic traumatic encephalopathy (CTE), or a neurodegenerative condition caused by a virus, alcoholism, tumor, toxin, or repetitive brain injuries.

21. The method of claim 20, wherein the neurodegenerative disease is Parkinson's Disease.
22. The method of claim 20, wherein the injury, disease, or condition is Alzheimer's Disease, migraine, brain surgery, or a neurological side effect associated with cancer chemotherapy.
23. The method according to any one of claims 16-22, wherein neuroprotection or neurorestoration is increased in the patient as compared with an untreated patient.
24. The method according to any one of claims 16-23, wherein the A₃R is partially agonized with improved neuroprotective or neurorestoration function relative to a full A₃R agonist..
25. A method of treating a heart or cardiovascular disease, comprising administering to a patient in need thereof an effective amount of an agonist of an A₃ adenosine receptor (A₃R), wherein the agonist is a biased agonist of an A₃R with improved cardioprotection function relative to a full A₃R agonist via preferential activation of one or more of the following A₃R-mediated pathways: activation of Gq11-mediated intracellular calcium mobilization, Gi-mediated modulation of cAMP production, Gi-mediated phosphorylation of ERK1/2 and Akt, modulation of ATP- sensitive Potassium channels, or modulation of Beta-Arrestin activation.
26. The method according to claim 25, wherein the patient has suffered a cardiac ischemia or myocardial infarction.
27. The method according to claim 25, wherein said method increases cardioprotection or regeneration of damaged heart tissue in the patient.

28. The method according to any one of claims 25-27, wherein the recovery period after the TBI, stroke, cardiac ischemia, or myocardial infarction is decreased as compared with an untreated patient.


29. The method according to any one of claims 25-28, wherein the compound is a partial agonist of the A₃R with improved cardioprotection function relative to a full A₃R agonist.

1/20

Figure 1A**MRS4322 Plasma & Brain Concentration vs Time Profiles in Mice****Figure 1B****MRS4322 Plasma & Brain Concentration vs Time Profiles in Neonatal Pigs**

2/20


Figure 2



3/20


Figure 3

**MRS2365 and MRS2365 Monophosphate Percent Observed v Time
Following Incubation with Male CD-1 Mouse EDTA-Treated Plasma at 37°C**



4/20


Figure 4



5/20


Figure 5

**MRS2365 and MRS2365 Monophosphate Percent Observed v Time
Following Incubation with Male CD-1 Mouse EDTA-Treated Whole Blood at 37°C**




6/20

Figure 6




7/20

Figure 7




8/20

Figure 8




9/20

Figure 9




10/20

Figure 10




11/20


Figure 11

**MRS 2365**Chemical Formula: $C_{13}H_{19}N_5O_9P_2S$

Exact Mass: 483.0379

**M2**Chemical Formula: $C_{13}H_{18}N_5O_6PS$

Exact Mass: 403.0715


MRS2347**M1**Chemical Formula: $C_{13}H_{17}N_5O_3S$

Exact Mass: 323.1052

MRS4322


12/20

Figure 12




13/20

Figure 13




14/20

Figure 14



15/20


Figure 15



16/20


Figure 16

**Brain and Cerebrospinal Fluid Concentration-Time Profiles of MRS4322 in Neonatal Pigs
Following Intravenous Administration**




17/20

Figure 17

Competition binding experiments of MRS4322 at human A3 receptors


18/20

Figure 18

Competition binding experiments of MRS4322 at mouse A3 receptors

19/20

Figure 19

cAMP accumulation experiments of MRS4322 and NECA at human A3 receptors

20/20

Figure 20

cAMP accumulation experiments of MRS4322 and NECA at mouse A3 receptors

