

(19) AUSTRALIAN PATENT OFFICE

(54) Title	Process for the preparation of prostaglandins and analogues thereof		
(51) ⁶ International Patent Classification(s)	C07D 307/93 20060101ALI2005122 (2006.01) OBMJP C07C C07B 53/00 (2006.01) 405/00 C07B 61/00 (2006.01) 20060101ALI2005100 C07C 405/00 8BMEP C07D (2006.01) 307/935 C07D 307/935 20060101ALI2005100 (2006.01) 8BMEP C07F C07F 7/18 (2006.01) 7/18 C07F 9/38 (2006.01) 20060101ALI2005100 C25B 3/02 (2006.01) 8BMEP C07F C25B 3/02 (2006.01) 9/38 C07D 307/93 20060101ALI2005122 20060101AFI2005122 OBMJP C25B 53/00 C07B 3/02 20060101ALI2005122 20060101ALI2005122 OBMJP C07B OBMJP 61/00 PCT/GB02/02462		
(21) Application No:	2002321396	(22) Application Date:	2002.05.24
(87) WIPO No:	W002/096898		
(30) Priority Data			
(31) Number	(32) Date	(33) Country	
0112699.4	2001.05.24	GB	
(43) Publication Date :	2002.12.09		
(43) Publication Journal Date :	2003.05.08		
(71) Applicant(s)	Resolution Chemicals Limited		
(72) Inventor(s)	Thompson, David, George, Greenwood, Alan, Kenneth, Clissold, Derek, Wyndham, McHattie, Derek		
(74) Agent/Attorney	Shelston IP, Level 21 60 Margaret Street, Sydney, NSW, 2000		
(56) Related Art	Skuballa et al, J. Med. Chem. 1978, vol. 21(5), page 445 WO 93/00329 WO 98/21181 Nooy et al, Synthesis, 1995, pages 1153-1174 Liljebris et al, Tetrahedron, 1995, vol 51(33) pages 9139-9154 EP 0472338 US 4680415		

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number
WO 02/096898 A3

(51) International Patent Classification⁵: C07D 307/935, (74) Agents: RITTER, Stephen, David et al.; Mathys & C07C 405/00, C07F 7/18 Squire, 100 Gray's Inn Road, London WC1X 8AI. (GB).

(21) International Application Number: PCT/GB02/02462

(81) Designated States (national): AE, AG, AL, AM, AT, AU,

(22) International Filing Date: 24 May 2002 (24.05.2002)

AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

(25) Filing Language: English

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

(26) Publication Language: English

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

(30) Priority Data: 0112699.4 24 May 2001 (24.05.2001) GB

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,

(71) Applicants (for all designated States except US):

MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,

RESOLUTION CHEMICALS LIMITED [GB/GB]; Wedgwood Way, Stevenage, Hertfordshire SG1 4QT (GB). CASCADE BIOCHEM LIMITED [GB/GB]; Earley Gate, Whiteknights Road, Reading, Berkshire RG6 6BZ (GB).

SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(72) Inventors; and

(84) Designated States (regional): ARIPO patent (GH, GM,

(75) Inventors/Applicants (for US only): GREENWOOD, Alan, Kenneth [GB/GB]; Resolution Chemicals Limited, Wedgwood Way, Stevenage, Hertfordshire SG1 4QT (GB). MCHATTIE, Derek [GB/GB]; Resolution Chemicals Limited, Wedgwood Way, Stevenage, Hertfordshire SG1 4QT (GB). THOMPSON, David, George [GB/GB]; Resolution Chemicals Limited, Wedgwood Way, Stevenage, Hertfordshire SG1 4QT (GB). CLISSOLD, Derek, Wyndham [GB/GB]; Cascade Biochem Limited, Earley Gate, Whiteknights Road, Reading, Berkshire RG6 6BZ (GB).

KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

20 March 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/096898 A3

(54) Title: PROCESS FOR THE PREPARATION OF PROSTAGLANDINS AND ANALOGUES THEREOF


(57) Abstract: Disclosed are processes for the synthesis and purification of prostaglandins and analogues thereof, especially analogues of PGF_{2α}.

PROCESS FOR THE PREPARATION OF PROSTAGLANDINS AND
ANALOGUES THEREOF

5 The present invention relates to a novel process for the synthesis of prostaglandins and prostaglandin analogues. In particular, this invention relates to the synthesis of PGF_{2 α} and analogues thereof.

Prostaglandin F_{2 α} {PGF_{2 α} - 7-[3,5-dihydroxy-2-(3-hydroxy-1-octenyl)-cyclopentyl]-5-heptenoic acid]} has the structure:

10



This compound causes uterine contraction and is used clinically to induce and accelerate labour, and as an abortifacient.


15

Prostaglandins are generally characterised by the substituents on the cyclopentyl ring. The PGF_{2 α} prostaglandins and prostaglandin analogues generally have two hydroxyl groups in a *cis* configuration relative to the cyclopentane ring, and two side chains in a *trans* configuration relative to each other, each side chain having one double bond. Analogues of PGF_{2 α} can have a different number of double bonds in the side chains, and the substituents along the side chains may vary. Additionally, in some PGF_{2 α} analogues, the side chain carboxylic acid group may be esterified.

20


Examples of PGF_{2 α} analogues having therapeutic use are cloprostenol, which contains a chlorophenyl ether side chain substituent, fluprostenol, which contains a trifluoromethylphenyl ether side chain substituent, and travoprost:

25



These compounds have prostaglandin F agonist activity and are used in the clinic for treating glaucoma and ocular hypertension.


5 Latanoprost [13,14-dihydro-15(R)-17-phenyl-18,19,20-trinor-PGF_{2α}-isopropyl] is an example of a PGF_{2α} analogue having one saturated side chain and wherein the carboxylic acid group is esterified:




10 This compound is used in the clinic for the reduction of elevated intra-ocular pressure in patients with open angle glaucoma and ocular hypertension.

15 Prostaglandin analogues based on PGF_{2α} for use in the treatment of glaucoma and ocular hypertension are described in, for example, European patent number 0 364 417 B1. The procedures for the synthesis of PGF_{2α} analogues described therein start from an advanced-stage intermediate, 16-phenyl-17,18,19,20-trinor PGF_{2α}, or the tetranor homologue thereof.


European patent number EP 0 544 899 B1 describes a process for the synthesis of 13,14-dihydro-15(R)-17-phenyl-18,19,20-trinor-PGF_{2α} esters of the formula:



5 wherein the group R represents alkyl, phenyl or benzyl. The starting material for the process disclosed therein is the *para*-phenylbenzoyl (PPB)-protected Corey lactone:




In the disclosed process, the intermediate of formula:




is prepared by transforming the hydroxymethyl group of the protected Corey lactone, with an oxidising agent (dicyclohexylcarbodiimide) to form the corresponding aldehyde. Reaction of the aldehyde with a phenylphosphonium salt forms the above intermediate.

15

The intermediate is then reduced to form the corresponding hydroxy compound, which is subjected to a hydrogenation reaction to form the saturated side chain intermediate of the formula:



The lactone oxo group of the above intermediate is reduced to form the corresponding hydroxy analogue, which is subsequently deprotected to give the following intermediate:



A subsequent Wittig reaction followed by esterification forms the desired PGF_{2α} product.

The process disclosed in EP 0 544 899 B1 suffers from a major drawback in
 10 that only moderate or low yields are achieved in certain steps. In particular, according to the examples, the yield for the first step (reaction of the PPB-protected Corey lactone with triphenyl-(4-phenyl-3-oxobutyl)-phosphonium iodide on a small scale was only around 49%. Even on a large scale, the reported yield was only around 53%.

These low yields at the outset result in, at best, difficulties in purifying the intermediates, and at worse, a loss of valuable and expensive starting material. Additionally, the reported yield of 38% in the second step (reduction of the side chain oxo group), is also less than satisfactory.

Furthermore, only moderate yields are achieved in subsequent reaction steps, resulting in a dramatic reduction of the overall yield.


In an alternative synthesis described in B. Resul *et al.*, J. Med. Chem. (1993), 36, pp. 243-248, a similar route was employed, with the exception that the deprotection step was carried out before the reduction of the oxo group of the lactone. This process also suffers from low yields (about 58%) in the first step, and only moderate or low yields were reported for subsequent steps.

In view of the prior art procedures and the problems associated therewith, it is an object of the present invention to provide an alternative process for the synthesis of PGF2 α and analogues and salts thereof.

It is an object of the present invention to overcome or ameliorate at least one of the disadvantages of the prior art, or to provide a useful alternative.

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

Accordingly, the present invention provides a process for the preparation of prostaglandin derivatives having the Formulae (I-A) and (I-B):



wherein:


B represents a substituent selected from the group consisting of:

(i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo and CF₃; and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo or CF₃; and

R" represents C₁-C₂₀ alkyl (preferably a C₁ to C₆ alkyl group, e.g. methyl, ethyl, propyl and iso-propyl), C₃ to C₈ cycloalkyl (e.g. cyclohexyl, cyclopropyl, cyclobutyl) or C₆ to C₁₀ aryl (preferably phenyl). A preferred R" group is iso-propyl.

It will be appreciated by a person of skill in the art that the present invention can be applied to the synthesis of compounds of Formula (I-A) and (I-B) wherein R" is other than an alkyl, cycloalkyl or aryl group. In fact, the group R" can represent any group that can be introduced by reaction of an acid intermediate of the compound of Formula (I-A) or (I-B) with a compound R"-X (X = halo, and preferably iodo) in the presence of a base, such as DBU. Examples of other suitable R" groups include, but are not limited to, unsaturated C₁ to C₂₀ alkyl, unsaturated C₃ to C₈ cycloalkyl, wherein the saturated or unsaturated alkyl or cycloalkyl groups, or aryl groups can be substituted with one or more (typically 1 to 3) substituents such as CF₃, C₁ to C₆ alkoxy, CN. Other examples of suitable R" groups include C₆ to C₁₀ heterocycloalkyl (e.g. piperidinyl), C₆ to C₁₀ heteroaryl (such as pyridyl) and substituted C₆ to C₁₀ aryl (including substituents such as CF₃, C₁ to C₆ alkoxy, CN).

According to a first aspect of the invention, there is provided a process for the synthesis of a compound of Formula (I-A) or (I-B):



wherein:


20 B represents a substituent selected from the group consisting of:

(i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo and CF₃ and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo or CF₃; and

R" represents a C₁ to C₆ alkyl group or a C₃ to C₈ cycloalkyl group;

comprising the steps of:


subjecting a compound of Formula (X):



wherein


5 A represents C₆ to C₁₀ aryl which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C₁ to C₆ alkyl and (iii) unsubstituted C₆ to C₁₀ aryl;

to an oxidation reaction in the presence of a catalytic amount of a stable organic nitroxyl radical, to form a compound of formula (IX):



10


(2) subjecting a compound of Formula (IX) as defined above to reaction with a compound having the formula:




wherein B is as defined above and each R'' is the same or different and each 15 represents a C₁ to C₆ alkyl group,

- 6b -


the process being carried out in the presence of lithium chloride and an organic base to form the compound of Formula (VIII):




(3) reducing the side chain oxo group of the compound of Formula (VIII) with a
5 reducing agent to form a compound of Formula (VII):



(4) deprotecting the hydroxyl group of the compound of Formula (VII) to form
the corresponding hydroxy-substituted compound of Formula (VI):



10 (5) when the dashed line in the compound of Formula (I-A) or (I-B) is absent:
hydrogenating the double bond of a compound of Formula (VI), to form a compound of
Formula (VI-A):




- 6c -

(6) subjecting the compound of Formula (VI) or (VI-A) to reaction with a silylating agent of formula




wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl and X represents F , Cl , Br or I , to form a compound of Formula (V):



wherein R' represents:



(7) reducing the lactone oxo group of the compound of Formula (V) to form a compound of Formula (IV):




(8) subjecting the compound of Formula (IV) to a Wittig reaction with an ylide, the ylide being formed from the reaction of a compound of formula:




with a strong base, wherein R^w represents C_1 to C_6 alkyl or C_6 to C_{10} aryl and X represents F , Cl , Br or I ,

to form a compound of Formula (IIIa) or (IIIb) or a mixture thereof:



(9) for compounds of Formula (I-B): subjecting a compound of Formula (IIa) or Formula (IIb) or a mixture thereof, to reaction with an alkyl halide of formula $R''\text{-Hal}$,
5 wherein R'' is as defined above and "Hal" represents chloro, bromo or iodo, in the presence of DBU [1,8-diaza-bicyclo(5.4.0)undec-7-ene], to form a compound of Formula (IIa) or (IIb):




or a mixture thereof; and


10 (10) removing the protecting groups from: (a) the compound of Formula (IIIa) or
(IIIb) or a mixture thereof to form the compound of Formula (I-A), or (b) a compound of
Formula (IIa) or (IIb) or a mixture thereof to form the compound of Formula (I-B).

According to a second aspect of the invention, there is provided a process for the synthesis of a compound of Formula (I-B) as defined in the first aspect, wherein the dashed line is absent, comprising carrying out steps (1) and (2) as defined in the first aspect, then carrying out the following steps (3'), (4') and (5'):


(3') subjecting the double bond of the compound of Formula (VIII) to a hydrogenation reaction to form a compound of Formula (XIII):



20 (4') reducing the side chain oxo group of the compound of Formula (XIII) to form a compound of Formula (XIV);




(5') deprotecting the hydroxyl group of the compound Formula (XIV) to form a compound of Formula (VI-A):




5 subjecting the above compound of Formula (VI-A) to reaction steps (6), (7), (8),
(9) and (10) as defined in the first aspect.

According to a third aspect of the invention, there is provided a process for the synthesis of a compound of Formula (I-A) or (I-B) as defined in the first aspect comprising producing a compound of Formula (IIIa) or Formula (IIIb) or a mixture thereof as defined in either the first aspect or the second aspect:




and further carrying out the steps of:

(9a) subjecting the compound of Formula (IIa) or (IIb) or a mixture thereof, to reaction with at least one molar equivalent of silylating agent to form a compound of Formula (XI):




(10a) for compounds of Formula (I-B): subjecting the compound of Formula (XI) to a transesterification reaction with an alcohol having the formula $R''-OH$, wherein R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl optionally in the presence of a weak acid catalyst to form the compound of Formula (XII):



; and


(11a) removing the protecting groups R' from either (a) the compound of Formula (XI) to form the compound (I-A) or (b) the compound of Formula (XII) to form the compound (I-B).

5 According to a fourth aspect of the invention, there is provided a process for the synthesis of a compound of Formula (I-B) as defined in the first aspect comprising carrying out steps (1) to (9) as defined in either the first aspect or the second aspect to form a compound of formula (IIa) or (IIb) or a mixture thereof:



10 and further carrying out the steps of:


(10b) subjecting the compound of Formula (IIa) or (IIb) or a mixture thereof to reaction with at least one molar equivalent of silylating agent to form a compound of Formula (XII):



; and


15 (11b) removing the protecting groups R'.

According to a fifth aspect of the invention, there is provided a process for the synthesis of a compound of Formula (I-B) as defined in the first aspect comprising removing the silyl protecting groups, R', from a compound of Formula (XII), as prepared by the process of either the third aspect or the fourth aspect




According to a sixth aspect of the invention, there is provided a process for the synthesis of a compound of Formula (I-A) as defined in the first aspect comprising removing the silyl protecting groups, R', from a compound selected from the group consisting of


5 (i) a compound of Formula (IIIa) or Formula (IIIb), as prepared by the process of the first aspect, or a mixture thereof:



(ii) a compound of Formula (XI), as prepared by the process of the third aspect:




According to a seventh aspect of the invention, there is provided a process for the synthesis of a compound of Formula (IX):



wherein

A represents C₆ to C₁₀ aryl which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C₁ to C₆ alkyl and (iii) unsubstituted C₆ to C₁₀ aryl;


15 comprising subjecting a compound of Formula (X):



- 6h -

to an oxidation reaction in the presence of a catalytic amount of a stable organic nitroxyl radical.

According to an eighth aspect of the invention, there is provided a process for the synthesis of a compound of Formula (XI):



wherein the groups R' are the same and each represents the substituent:




wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;

10 comprising reacting a compound of Formula (IIIa) or Formula (IIIb), as defined in the first aspect, or a mixture thereof, to reaction with at least a molar equivalent of silylating agent having the formula:



wherein R^x, R^y and R^z are as defined above and X represents F, Cl, Br or I.

15 According to a ninth aspect of the invention, there is provided a process for the synthesis of a compound of Formula (XII):



wherein

each R' is the same and each represents the substituent:



20

wherein

R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl; and


R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl;

comprising subjecting a compound of Formula (IIa) or Formula (IIb), as defined in the first aspect, or a mixture thereof, to reaction with at least one molar equivalent of silylating agent having the formula:



wherein R^x , R^y and R^z are as defined above and the group X represents F, Cl, Br or I.

10 According to a tenth aspect of the invention, there is provided a process for the synthesis of a compound of Formula (XII):



wherein:


each R' is preferably the same and each represents the substituent:



15 wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl; and

R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl;

the process comprising subjecting a compound of Formula (XI), as defined in the third aspect:




wherein the groups R' are the same and each represents the substituent:



wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;


to a transesterification reaction with an alcohol having the formula R"-OH, wherein R" represents C₁ to C₆ alkyl or C₃ to C₈ cycloalkyl.

5 According to an eleventh aspect of the invention, there is provided a compound of Formula (XI)




or a single enantiomer thereof, wherein B and R' are as defined in the first aspect.

10 According to a twelfth aspect of the invention, there is provided a compound of Formula (XII)




or a single enantiomer thereof, wherein B and R'' are as defined in the first aspect.


15 According to a thirteenth aspect of the invention, there is provided a compound of Formula (VIII):



wherein A represents unsubstituted C₆ to C₁₀ aryl; and
B represents a substituent selected from the group consisting of:




According to a fourteenth aspect of the invention, there is provided a compound of Formula (VII):




5 or a single isomer thereof, wherein

A represents unsubstituted C₆ to C₁₀ aryl; and

B represents a substituent selected from the group consisting of:




According to a fifteenth aspect of the invention, there is provided a compound of
10 Formula (VI):




or a single isomer thereof, wherein

B represents a substituent selected from the group consisting of:




15 According to a sixteenth aspect of the invention, there is provided a compound of
Formula (VI-A):

- 61 -




or a single isomer thereof, wherein

B represents a substituent selected from the group consisting of:




5 According to a seventeenth aspect of the invention, there is provided a compound of Formula (V):




or a single isomer thereof, wherein


R' represents:



10 B represents a substituent selected from the group consisting of:



According to an eighteenth aspect of the invention, there is provided a compound of Formula (IV):




15


or a single isomer thereof, wherein

R' represents:


- 6m -



B represents a substituent selected from the group consisting of:




According to a nineteenth aspect of the invention, there is provided a compound of
5 Formula (IIIa) or (IIIb):




or a single isomer thereof, wherein


10 R' represents:



B represents a substituent selected from the group consisting of:




15 According to a twentieth aspect of the invention, there is provided a compound of
Formula (IIa) or (IIb):



or a single isomer thereof, wherein


R' represents:

- 6n -




, and

B represents a substituent selected from the group consisting of:



According to a twenty-first aspect of the invention, there is provided a compound of
5 formula (XIV):



or a single isomer thereof, wherein

A represents unsubstituted C₆ to C₁₀ aryl; and

B represents C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or
10 substituted with one to three substituents independently selected from the group
consisting of C₁ to C₆ alkyl, halo and CF₃.

According to a twenty-second aspect of the invention, there is provided a process for the
purification of latanoprost by HPLC comprising the use as an eluent, of a mixture
comprising one or more hydrocarbons, one or more alcohols and, optionally,
15 acetonitrile.


According to a twenty-third aspect of the invention, there is provided latanoprost
substantially free of the 15(S)-*cis* isomer, the 15(S)-*trans* isomer and the 15(R)-*trans*
isomer.


According to a twenty-fourth aspect of the invention, there is provided latanoprost
20 having a purity of greater than 98%.

Unless the context clearly requires otherwise, throughout the description and the
claims, the words "comprise", "comprising", "consisting" and the like are to be
construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to
say, in the sense of "including, but not limited to".

The embodiments of the process of the present invention are shown in Schemes 1-4:

Scheme 1 illustrates one route to the synthesis of compounds of Formula (I-A) and (I-B),
starting from a protected-Corey lactone compound of Formula (X):






(a) oxidation; (b) modified Horner-Wadsworth-Emmons reaction; (c) reduction; (d) deprotection
 (e) hydrogenation; (f) silylation; (g) silylation; (h) reduction; (i) Wittig reaction;
 (j) alkylation; (k) deprotection; (l) deprotection

SCHEME 1

In an alternative procedure according to the present invention, the
 5 intermediate (VI-A) in Scheme 1 can be made by carrying out steps (a) and

(b) as shown in Scheme 1, and substituting steps (c), (d), and (e) in Scheme 1 with the steps (e'), (c') and (d') as shown in the following Scheme 2:




(e') hydrogenation; (c') reduction; (d') deprotection

5

SCHEME 2


Scheme 3 illustrates an alternative procedure for the synthesis of compounds of Formula (I-A) and (I-B), starting from intermediates of structure (IIIa) and (IIIb):



(m) silylation; (n) deprotection; (o) transesterification; (p) deprotection

SCHEME 3


Scheme 4 shows an alternative procedure for the synthesis of compounds of
5 Formula (I-B) starting from the intermediates of Formula (IIa) and (IIb):



(q) silylation; (r) deprotection


SCHEME 4

In one aspect of the present invention, there is provided a process for the
5 production of a compound of Formula (IX):



wherein

A represents C₆ to C₁₀ aryl which may be substituted with one to three
10 substituents independently selected from the group consisting of
(i) halo, (ii) C₁ to C₃ alkyl and (iii) unsubstituted C₆ to C₁₀ aryl;
the process comprising subjecting a compound of Formula (X)



to an oxidation reaction in the presence of a catalytic amount of a stable organic nitroxyl radical.

5 The above reaction may be carried out by electrooxidation in the presence of the organic nitroxyl radical.

10 Alternatively, the oxidation reaction may be carried out in the presence of a nitroxyl radical and at least one molar equivalent of a co-oxidant selected from the group consisting of m-chloroperbenzoic acid, high-valent metal salts, sodium bromite, sodium or calcium hypochlorite, N-chlorosuccinimide or hypervalent iodine compounds such as [bis(acetoxyl)iodo]benzene. Preferably, the co-oxidant is sodium hypochlorite.


15 The stable organic radical preferably comprises a completely α -substituted piperidin-1-oxy radical, such as 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical (TEMPO, free radical).

20 Prior art oxidation procedures for oxidising the compound of Formula (X) to form the compound of Formula (IX) include the use of dimethylsulfoxide-dicyclohexylcarbodiimide. However, such a method requires isolation of the aldehyde (IX). Since the aldehyde (IX) is not particularly stable in solution, an amount of decomposition product is usually observed during work-up.

25 Advantageously, in the present method of oxidation of the compound of Formula (X) using a stable organic radical such as TEMPO free radical, the aldehyde (IX) solution obtained in this step can be employed in the subsequent step without isolation of the aldehyde, thus minimising any decomposition.

30 In a further aspect of the present invention, there is provided a process for the production of a compound of Formula (VIII):

12




wherein


A represents C_6 to C_{10} aryl which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C_1 to C_6 alkyl and (iii) unsubstituted C_6 to C_{10} aryl;

B represents a substituent selected from the group consisting of: (i) C_1 to C_6 alkyl, (ii) C_7 to C_{16} aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo and CF_3 and (iii) $-(CH_2)_nOR^a$, wherein n represents 1, 2 or 3 and R^a represents a C_6 to C_{10} aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo or CF_3 ,

15 the process comprising subjecting a compound of Formula (IX):




to reaction with a phosphonate compound having the structure:




wherein A and B are as defined as above, and each R^{'''} is the same or different (preferably the same) and each represents a C₁ to C₆ alkyl group (preferably methyl), the process being carried out in the presence of lithium chloride and an organic base (such as tertiary alkylamines, e.g. 5 di-*iso*-propylethylamine).


The compounds of Formula (IX) are commercially available or can be made from commercially available starting materials. For example, the compound of Formula (IX) may be prepared by the process described in US 3,778,450. 10 The phosphonate compound of formula:



may be produced by methods known in the art [see for example, B. Resul *et al.*, J. Med. Chem. (1993), 36, 243-248]. As an example, 15 dimethyl-(2-oxo-4-phenylbutyl)phosphonate may be produced from dimethyl-(2-oxo-propyl)phosphonate via the following reaction:




A preferred process for producing the phosphonate compound of formula:




comprises the reaction of a dialkylmethylphosphonate of structure:



with a strong base (e.g. n BuLi) to generate an anion, followed by reaction of the anion with a compound of formula $B-CO_2R^y$, wherein R^y can represent any group that can form a leaving group -OR y . Typical R^y groups include C₁ to C₆ alkyl, such as methyl, or ethyl (methyl is preferred). Thus the reaction of 5 dimethylmethyl phosphonate (methylphosphonic acid dimethyl ester) with a strong base such as n -butyllithium, to generate the corresponding anion, followed by reaction of the anion with a compound of Formula $B-CO_2R^y$ is shown below:



10 Dimethyl(2-oxo-4-phenylbutyl)phosphonate can be produced in good yield from the following reaction:



15 The product from this reaction can be typically purified by distillation. Advantageously, this reaction is usually free of side reactions compared with the prior art process using dimethyl(2-oxopropyl)phosphonate.

20 The process described above for producing the compound of formula VIII is a modified Horner-Wadsworth-Emmons reaction carried out in milder conditions, in which the usual base used to generate the anion of the phosphonate, sodium hydride in THF or potassium carbonate in toluene, is replaced by a base selected from the group consisting of tertiary alkylamines, such as triethylamine and diisopropylethylamine and DBU [1,8-diazabicyclo(5.4.0)undec-7-ene]. Advantageously, unlike the prior art methods which can lead to poor to moderate yields of the product (VIII), the 25 use of milder reagents, i.e. lithium cations from lithium chloride, in

combination with these tertiary alkylamines leads to cleaner reactions with high yield of product.


5 The reaction is preferably carried out at temperatures in the range of -20°C to 40°C, and preferably -10°C to 30°C. Suitable solvents for this reaction include those selected from the group consisting of benzene, toluene, acetonitrile, dichloromethane, diethylether, and mixtures thereof.

10 In the compound of Formula (VIII), the group A preferably represents an unsubstituted C₆ to C₁₀ aryl group (e.g. phenyl).

15 Other preferred substituents for the group A include those selected from C₆ to C₁₀ aryl group being substituted with one substituent selected from halo or phenyl. Further preferred substituents for the group A include unsubstituted or substituted phenyl wherein the substituent is selected from halo or phenyl. In a preferred process, the group A represents phenyl.

According to another aspect of the present invention, there is provided a process for the production of a compound of Formula (VII):

20



wherein


A represents unsubstituted C₆ to C₁₀ aryl;

B represents a substituent selected from the group consisting of:

25 (i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo

and CF_3 and (iii) $-(\text{CH}_2)_n\text{OR}^a$, wherein n represents 1, 2 or 3 and R^a represents a C_6 to C_{10} aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo or CF_3 ;

5 the process comprising reducing the oxo group in the side chain of a compound of Formula (VIII):



Suitable reducing agents for the reduction of the side chain oxo group include borane-dimethylsulfide complex, lithium tri-*sec*-butylborohydride, 10 $\{\text{LiB}[\text{CH}(\text{CH}_3)\text{CH}(\text{C}_2\text{H}_5)_3\text{H}\}$ (L-Selectride^{RTM}) and sodium borohydride. Where the final product is desired as a racemate (such as fluprostenol), a non-stereoselective reducing agent may be used (e.g. LiAlH_4 , NaBH_4 and other metallic hydrides).

15 Where the final product is desired as a single isomer (e.g. as in latanoprost), the reducing agent suitably comprises borane-dimethylsulfide complex in the presence of a chiral oxazaborolidine catalyst ("Corey catalyst") because of the greater selectivity towards the production of a major amount of the desired isomer. Although this reaction strongly favours the production of the desired 20 isomer, any undesired isomer which may be formed, may be separated by chromatographic techniques, such as flash column chromatography.


Thus, a preferred reagent for the reduction reaction is borane-dimethylsulfide complex in the presence of a chiral oxazaborolidine catalyst (Corey catalyst).

25 In this embodiment of the present invention, the group A in the compound of Formula (VII), in addition to being unsubstituted C_6 to C_{10} aryl, can also

represent C_6 to C_{10} aryl substituted with one to three substituents independently selected from the group consisting of (i) halo, i.e. fluoro, chloro, bromo or iodo, (ii) C_1 to C_6 alkyl and (iii) C_6 to C_{10} aryl, such as phenyl.

5 The use of borane-dimethylsulfide complex in combination with a Corey catalyst is especially preferred because the reaction takes place with excellent selectivity. In fact, a marked improvement in stereoselectivity is seen compared with the reaction using L-SelectrideTM. A further advantage is that the reduction reaction using borane-dimethylsulfide complex can be 10 carried out at a higher temperature (typically -15°C to -18°C) compared with L-SelectrideTM, which requires a reaction temperature of less than -70°C .


The Corey catalyst comprises a chiral oxazaborolidine compound [see J. Am. Chem. Soc., 109, 5551, (1987) and J. Am. Chem. Soc. 109, 7925, (1987) and references cited in Lancaster Catalogue 2000-2001, page 819] such as (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrole[1,2-c][1,3,2]oxazaborole, may be prepared by reaction of the appropriate chiral prolinol [such as the commercially available (R)-(+)- α,α -diphenylprolinol] with a trialkyl boroxine, e.g.:



The reaction is carried out in inert conditions in a solvent such as toluene, diethylether or tetrahydrofuran. The oxazaborolidine catalyst is employed as a solution in the reduction step.

25 According to another aspect of the present invention, there is provided a process for the production of a compound of Formula (V):

18



(V)

wherein

the dashed line forms an optional double bond;

B represents a substituent selected from the group consisting of:

5 (i) C_1 to C_6 alkyl, (ii) C_7 to C_{16} aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo and CF_3 and (iii) $-(CH_2)_nOR^a$, wherein n represents 1, 2 or 3 and R^a represents a C_6 to C_{10} aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo or CF_3 ; and


10 R' represents the substituent:



15 wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl;

the process comprising the steps of:

(a) deprotecting the hydroxyl group of a compound of Formula (VII):




(VII)

wherein A represents C_6 to C_{10} aryl which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C_1 to C_6 alkyl and (iii) unsubstituted C_6 to C_{10} aryl;


5

to form the corresponding hydroxy-substituted compound of Formula (VI):



10

(b) optionally, when the dashed line of the compound of Formula (V) represents a single bond, the double bond of the compound of Formula (VI) is hydrogenated to form the compound of Formula (VI-A):



15

(c) subjecting a compound of Formula (VI) or (VI-A) to a reaction with a silylating agent having the formula:



wherein R^x , R^y and R^z are as defined above and X represents F, Cl, Br or I.

The deprotection step (a) wherein the protecting group A on the hydroxyl group of the cyclopentane ring is removed, is preferably carried out in the presence of a base. Preferred bases for use in the deprotection reaction includes those selected from the group consisting of K_2CO_3 , Na_2CO_3 and Li_2CO_3 , with K_2CO_3 being particularly preferred.

Suitable solvents for the deprotection reaction include alcohols, such as methanol, ethanol and isopropanol.

10 The isolation of the deprotected product from step (a) may be carried out by standard chromatography procedures. However, it has been found that the deprotected product can advantageously be isolated by extraction with hexane fractions, thus avoiding the use of time consuming and expensive chromatographic procedures.

15

For the production of prostaglandin derivatives having one saturated side chain, such as latanoprost described above, it is convenient at this stage, i.e. after the deprotection step (a), to hydrogenate the double bond in the side chain to form the intermediate of Formula (VI-A).

20

The hydrogenation step can be carried out using any suitable hydrogenation catalyst such as palladium, platinum or rhodium, which may be supported on an inert support, such as carbon. An example of a suitable hydrogenation catalyst is 5% palladium on carbon.

25

In a preferred procedure, the hydrogenation reaction is carried out in the presence of sodium nitrite, preferably in aqueous solution. This procedure avoids the formation of elimination products and thus results in improved yields (typically greater than 95%) of the compounds of Formula (VI-A).

30

Suitable solvents for the hydrogenation reaction include alcohols such as methanol and ethanol. On completion of the hydrogenation reaction, the mixture is preferably stirred with dilute hydrochloric acid to remove the nitrite

(by conversion to nitrous acid, which decomposes at ambient temperature). This procedure ensures that nitrite is not carried through to the subsequent synthetic procedures.

5 Without wishing to be bound by theory, the role of the sodium nitrite in the hydrogenation reaction is to avoid the formation of elimination products, that is, the elimination of a water molecule from the side chain of the starting material and, as a consequence, formation of the fully saturated deoxygenated analogue of the desired product.

10 In order to produce the compound of Formula (V), the compound produced in step (a) having the Formula (VI) or the compound produced in step (b) having the Formula (VI-A) is reacted with a silylating agent (X)Si(R^x)(R^y)(R^z). The groups R^x, R^y, and R^z can be the same or different each represents a C₁ to C₆ alkyl group or a C₆ to C₁₀ aryl group. Preferably, each of the groups R^x, R^y, and R^z are independently selected from methyl, ethyl, butyl, isopropyl. Particularly preferred silylating agents for use in step (c) are selected from the group consisting of trimethylsilyl chloride, triethylsilyl chloride and *tert*-butyldimethylsilyl chloride. Triethylsilyl chloride is particularly preferred.

15

20 The silylation step is preferably carried out in the presence of a base, for example an organic base, such as imidazole or trialkylamines, such as triethylamine.

25 Suitable solvents for use in the silylation reaction include polar aprotic solvents such as tetrahydrofuran or dimethylsulfoxide, or chlorinated solvents such as dichloromethane. Preferably, however, the reaction is carried out in a solvent comprising dimethylformamide.


30 The use of silyl protecting groups in accordance with the present invention is advantageous because it generally results in cleaner reactions, with higher yields compared with reactions wherein the hydroxyl group is not protected.

The use of silyl protecting groups in the present process has particular advantages compared with the prior art process employing, e.g. benzoyl- and para-phenylbenzoyl (PPB)-protecting groups because silyl groups are stable to the subsequent reduction reaction with e.g. DIBAL-H (di-*iso*-butylaluminium).
5

In fact, the presence of benzoyl- and PPB-protecting groups prevents the use of many reducing agents, including DIBAL-H, in the subsequent reduction of the lactone ring to form the lactol, because such a reaction can result in the
10 reduction of the oxo moieties in the protecting groups. Thus, in prior art procedures wherein DIBAL is employed as a reducing agent and wherein the starting material comprises benzoyl- or PPB-protecting groups, starting materials must be deprotected prior to the reduction step. However, the presence of unprotected hydroxyl groups in the DIBAL-H reduction step is
15 undesirable because each free hydroxyl group or the starting material coordinates with DIBAL-H. As a result, the use of additional equivalents of DIBAL-H becomes necessary.


A second advantage of using silyl protecting groups in the subsequent Wittig reaction [step (i) in Scheme 1], is that the formation of the desired *cis* isomer
20 is favoured. Silyl protecting groups have the further advantage in that they generally increase the lipophilic character of the molecules, so that their derivatives are readily soluble in organic solvents. Thus, in the Wittig reaction, removal of the phosphine oxide by-product is facilitated because the silyl-protected Wittig reaction product [(IIIa)/(IIIb)] is soluble in hexane, whereas the triphenylphosphine oxide is insoluble, thus allowing separation by filtration. Subsequent purification of the product can be carried out by
25 silica gel filtration, rather than a full chromatographic purification.
A further advantage of employing silyl protecting groups is that these
30 protecting groups can be removed under mild conditions, as discussed below.

Alternatively, the intermediate compound of Formula (V) wherein the solid and dashed lines represent a single bond (i.e. compounds of formula (V-A)):




5 can be made from compounds of Formula (VIII) by a process comprising the steps of:

(a) hydrogenating the double bond of the compound of Formula (VIII):




to form a compound of Formula (XIII):




10

(b) reducing the side chain oxo group of the compound of Formula (XIII) to form a compound of formula (XIV):



15

(c) deprotecting the hydroxyl group in the compound of Formula (XIV) to form a compound of formula (VI-A):



; and


(d) subjecting the compound of Formula (VI-A) to a reaction with a silylating agent having the formula:



5 wherein R^x , R^y and R^z are as defined above and X represents F , Cl , Br or I .

10 Steps (a) to (c) of this process are depicted as steps (e'), (c') and (d') in Scheme 2. Step (d) corresponds to step (f) of Scheme 1, the product of which is a compound of Formula (V) wherein the dashed and solid line represents a single bond. The hydrogenation, reduction, deprotection and silylation steps in this alternative procedure are carried out as for the immediately preceding process to form the compounds of Formula (V).

15 According to another aspect of the present invention, there is provided a process for the preparation of a compound of Formula (IV):



wherein:

the dashed line forms an optional double bond;


20 B represents a substituent selected from the group consisting of:

(i) C_1 to C_6 alkyl, (ii) C_7 to C_{16} aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo and CF_3 and (iii) $-(CH_2)_nOR^a$, wherein n represents 1, 2 or 3 and R^a represents a C_6 to C_{10} aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo or CF_3 ; and
 5 R' represents the substituent:




10 wherein R^X , R^Y and R^Z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl;

the process comprising reducing the lactone oxo group of the compound of Formula (V):



A suitable reducing agent for this process is di-*iso*-butylaluminium hydride (DIBAL-H), and the reaction may be carried out in e.g. tetrahydrofuran.

According to a further aspect of the present invention there is provided a 20 process for the production of a compound of Formula (IIIa) or (IIIb), or a mixture thereof:



wherein

the dashed line forms an optional double bond;

B represents a substituent selected from the group consisting of:


5 (i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo and CF₃ and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo or CF₃; and

10 R' represents the substituent:




15 wherein R^X, R^Y and R^Z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;

the process comprising subjecting a compound of Formula (IV):



20 to a Wittig reaction with an ylide, the ylide being formed by reaction of a compound of formula:



with a strong base, wherein

R^w represents C_1 to C_6 alkyl or C_6 to C_{10} aryl; and


X represents fluoro, chloro, bromo or iodo.

5

Preferably, the group R^w represents phenyl. The group X preferably represents bromo.

Reagents of formula:

10




are commercially available, or can be prepared by reaction of a phosphine, $\text{P}(\text{R}^w)_3$, with $\text{HO}_2\text{C}(\text{CH}_2)_4\text{X}'$ (wherein X' represents halide, e.g. F, Cl, Br or I).

15

Suitable bases for forming the ylide include those selected from the group consisting of butyllithium, sodium amide, sodium hydride, and alkali metal alkoxides, including sodium methoxide, sodium ethoxide, potassium ethoxide and potassium *tert*-butoxide. Potassium *tert*-butoxide is a particularly preferred base. A suitable solvent for this reaction is tetrahydrofuran.

20

Thus, in a preferred embodiment, the ylide may be formed by the reaction of (4-carboxybutyl)-triphenylphosphonium bromide with potassium *tert*-butoxide:



25

The ylide can be generated using 3 equivalents of the phosphonium halide and 6 equivalents of base, i.e. a ratio of phosphonium halide and base of 1:2, but is preferably generated using 2.15 equivalents of the phosphonium halide and 4 equivalents of base.

During the Wittig reaction, the silyl protecting groups of the hydroxyl substituent on the cyclopentyl ring may migrate to the hydroxyl group formed by the opening of the lactol ring, to result in a mixture of 9- and 11-silylated isomers of Formula (IIIa) and (IIIb).


5

Although the mixture of compounds (IIIa) and (IIIb) can be separated e.g. by chromatographic procedures, separation is not necessary at this stage because the protecting groups are removed in subsequent reaction steps. Thus, in the process according to the invention, it is preferred that such a mixture of the 9- and 11-silylated isomers (IIIa) and (IIIb) is used in the subsequent reaction step without separation.

10

In a further aspect of the present invention there is provided a process for the preparation of a compound of Formula (XI):

15



wherein

the groups R' are the same and each represents the substituent:



20

wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;

the process comprising reacting a compound of Formula (IIIa) or Formula (IIIb), or a mixture thereof, to reaction with a silylating agent having the formula:




wherein R^x, R^y and R^z are as defined above and X represents F, Cl, Br or I.

5 This procedure is advantageously carried out where a mixture of the compounds of Formula (IIIa) and Formula (IIIb) are formed as the products of the Wittig reaction. The reaction of such a mixture with at least one molar equivalent of a silylating agent, preferably the same silylating agent as is used to protect the hydroxyl groups of the compounds of Formula (V), enables the mixture of compounds of Formula (IIIa) and (IIIb) to be "amalgamated" into a single product of Formula (XI) for subsequent reaction steps. Preferably, in 10 this step, at least a molar equivalent of silylating agent to starting material is employed. Typically, 1.1 to 2 molar equivalents are employed. The formation of a single product allows for better control of subsequent reaction steps and purification.

15

According to another aspect of the present invention, there is provided a process for the preparation of a compound of Formula (IIa) or (IIb) or a mixture thereof:



20 wherein

the dashed line represents an optional double bond;

B represents a substituent selected from the group consisting of:

(i) C_1 to C_6 alkyl, (ii) C_7 to C_{16} aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo

and CF_3 and (iii) $-(\text{CH}_2)_n\text{OR}^a$, wherein n represents 1, 2 or 3 and R^a represents a C_6 to C_{10} aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo or CF_3 ;

5 R' represents the substituent:




wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl; and

10 R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl;
the process comprising subjecting a compound of Formula (IIIa) or Formula (IIIb) or a mixture thereof, to reaction with an alkyl halide of formula $R''\text{-Hal}$, wherein R'' represents a C_1 to C_6 alkyl group (such as isopropyl) or a C_3 to C_8 cycloalkyl group, and "Hal" represents chloro, bromo, or iodo (preferably iodo), in the presence of DBU.

20 As indicated above, when the Wittig reaction results in migration of the silyl protecting group to form a mixture of 9- and 11-silylated isomers of Formula (IIIa) and (IIIb), the mixture may be alkylated by the process indicated above, to form a mixture of 9- and 11-silylated esters of Formula (IIa) and (IIb). Instead of carrying out a silylation of a mixture of compounds of Formula (IIIa) and (IIIb), it is also possible to carry out the silylation after the alkylation step, i.e. on the mixture of compounds of Formula (IIa) and (IIb).

25 Accordingly, a further aspect of the present invention provides a process for the production of a compound of Formula (XII):



wherein

R' represents the substituent:



5 wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl; and

10 R'' represents C₁ to C₆ alkyl or C₃ to C₈ cycloalkyl,
the process comprising subjecting a compound of Formula (IIa) or Formula (IIb) or a mixture thereof to reaction with at least one molar equivalent of silylating agent having the formula:




wherein R^x, R^y and R^z are as defined above and X represents F, Cl, Br or I.

15 As indicated above for the silylation of the mixture of compounds of Formula (IIIa) and (IIIb), this process is preferably carried out in the presence of at least a molar equivalent of silylating agent, and even more preferably 1.1 to 2 molar equivalents of silylating agent is employed. Again, this step leads to the "amalgamation" of the mixture of compounds of Formula (IIa) and (IIb) to form a single product [i.e. compounds of Formula (XII)] which facilitates control of subsequent reaction steps and purification of subsequent intermediates.

It has been found that in the steps wherein the mixtures of compound of Formula (IIa) and (IIb) or mixtures of compounds of Formula (IIIa) and (IIIb) are reacted with silylating agents to form single products of Formula (XII) and Formula (XI) respectively (i.e. precursors to the target compounds of Formula IA and IB), purification is facilitated. Advantageously, it has been found that the compounds of Formula (XII) and (XI) can be purified by simply by filtration through silica gel, thus obviating the need to perform a full chromatographic separation, which is particularly undesirable at the late stage of the synthetic process. One reason for this is that the presence of three silyl groups in the compounds of Formula (XII) and (XI) leads to highly lipophilic molecules, which are highly soluble in heptane and thus can be easily separated from the more polar impurities.

According to another aspect of the present invention, there is provided a process for the preparation of a compound of Formula (XII):



wherein:


R' is the same and each represents the substituent:



wherein R^X, R^Y and R^Z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl; and

R'' represents C₁ to C₆ alkyl or C₃ to C₆ cycloalkyl;

the process comprising subjecting a compound of Formula (XI):



wherein the groups R' are preferably the same and each represents the substituent:




5 wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;

10 to a reaction with an alcohol, having the formula R''-OH, wherein R'' represents a C₁ to C₆ alkyl group (e.g. isopropanol) or C₃ to C₈ cycloalkyl (e.g. cyclohexanol).

The above process may be carried out optionally in the presence of a weak acid catalyst, such as pyridinium p-toluenesulfonate. The reaction should be carried out in the absence of water, to avoid deprotection of the silyl groups.

15

According to a further aspect of the present invention, there is provided a process for the production of a compound of Formula (I-A):



wherein


20

B represents a substituent selected from the group consisting of:


(i) C₁ to C₈ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₈ alkyl, halo and CF₃ and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₈ alkyl, halo or CF₃;

the process comprising removing the silyl protecting groups, R', from a compound selected from the group consisting of

10 (i) a compound of Formula (IIIa), Formula (IIIb) or a mixture thereof;




(ii) a compound of Formula (XI):



15 wherein B and R' are as defined above.

According to another aspect of the present invention, there is provided a process for the production of a compound of Formula (I-B):



wherein


B represents a substituent selected from the group consisting of:

(i) C_1 to C_6 alkyl, (ii) C_7 to C_{16} aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo and CF_3 and (iii) $-(CH_2)_nOR^a$, wherein n represents 1, 2 or 3 and R^a represents a C_6 to C_{10} aryl group which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C_1 to C_6 alkyl, halo or CF_3 ; and

10 R" represents C₁ to C₆ alkyl or C₃ to C₈ cycloalkyl;

the process comprising removing the silyl protecting groups R' from a compound selected from the group consisting of:


(i) a compound of Formula (IIa) or (IIb) or a mixture thereof:



15

and

(ii) a compound of Formula (XII):



wherein B and R" are as defined as above.

20

The deprotection of the silyl protecting groups R' is advantageously carried out in mild reaction conditions. Suitable reagents for removal of the silyl

groups from the compounds of Formula (IIIa), (IIIb), (XI), (IIa), (IIb) and (XII) include weak acids such as acetic acid and citric acid. An especially preferred weak acid is pyridinium p-toluenesulfonate.


5 The reaction may be carried out in any suitable solvent or solvent mixtures. An especially preferred solvent for the deprotection reaction comprises acetone and water.

In accordance with preferred embodiments of the present invention, the 10 compounds of Formulae (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A), (I-B), (XI), (XII) and (XIV) are single enantiomers (i.e. the wavy line in the side chain represents — or —). As discussed above, such 15 compounds can be made by using a stereoselective reducing agent [e.g. borane-dimethylsulfide complex in the presence of a chiral oxazaborolidine (Corey) catalyst] in step (c) of Scheme 1, or step (c') in Scheme 2.

In accordance with preferred embodiments of the present invention, the group B in the compounds of Formula (XII), (XI), (VIII), (VII), (VI), (VI-A), (V), 20 (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) or (I-B) is selected from the group consisting of (i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl wherein the aryl group is unsubstituted and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which is substituted with a substituent selected from halo or CF₃.

25 Even more preferred are compounds of Formula (XII), (XI), (VIII), (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) or (I-B) wherein B is selected from the group consisting of (i) C₁ to C₆ straight chain alkyl, (ii) -(CH₂)₂Ph and (iii) -CH₂OR^a, wherein R^a represents a phenyl group substituted with a chloro or CF₃ group.

Especially preferred are compounds of Formula (XII), (XI), (VIII), (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) or (I-B) wherein B represents a substituent selected from the group consisting of:




5

In a further preferred embodiment of the present invention, the solid and dashed lines in each of Formulae (XII), (XI) (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) and (I-B) forms a single bond.


10 In yet another preferred embodiment, the solid and dashed lines in each of Formulae (XII), (XI) (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) and (I-B) forms a double bond.

15 For the compounds of Formulae (XII), (XI) (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) and (I-B) wherein the solid and dashed line represents a single bond, B preferably represents $-\text{CH}_2\text{CH}_2\text{Ph}$.

20 For the compounds of Formulae (XII), (XI) (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (I-A) and (I-B) wherein the dashed line represents a double bond, B preferably represents a substituent selected from the group consisting of:




25 As will be appreciated by a person of skill in the art, the process of the present invention is generally applicable for the synthesis of prostaglandins and prostaglandin analogues, particularly $\text{PGF}_{2\alpha}$ and analogues thereof. The process is particularly useful for the production of a compound selected from the group consisting of:




In a preferred embodiment, the present invention provides a process for the synthesis of latanoprost comprising the steps of:

5 (1) subjecting a compound of Formula (X):




wherein A represents a C₆ to C₁₀ aryl group, preferably phenyl, which may be substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C₁ to C₆ alkyl and (iii) 10 unsubstituted C₆ to C₁₀ aryl,
 to an oxidation reaction with sodium hypochlorite, the oxidation reaction being carried out in the presence of a catalytic amount of a

stable organic nitroxyl radical (preferably TEMPO free-radical), to form a compound of Formula (IX):




(2) subjecting the compound of Formula (IX) as defined above to reaction with a compound having the structure:




wherein each R''' is the same or different and each represents a C₁ to C₆ alkyl group (preferably methyl), in the presence of lithium chloride and an organic base, to form the compound of Formula (VIII) wherein B is -CH₂CH₂Ph:

10




15

(3) reducing the side chain oxo group of the compound of Formula (VIII) using borane-dimethylsulfide complex, the reduction being carried out in the presence of a chiral oxazaborolidine catalyst, such as (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrole[1,2c][1,3,2]oxaza-borole, to form a compound of Formula (VII) wherein B is -CH₂CH₂Ph:




(4) deprotecting the hydroxyl group of the compound of Formula (VII), preferably using K_2CO_3 in methanol, to form the compound of Formula (VI) wherein B is $-CH_2CH_2Ph$:



5

10

(5) hydrogenating the double bond of a compound of Formula (VI) in the presence of a hydrogenation catalyst (such as palladium, platinum or rhodium), the reaction optionally being carried out in the presence of sodium nitrite, to form a compound of Formula (VI-A) wherein B is $-CH_2CH_2Ph$:




(6) subjecting the compound of Formula (VI-A) to reaction with a silylating agent of formula



wherein R^x , R^y and R^z are as defined above (a particularly preferred silylating agent being triethylsilyl chloride) as defined above to form a compound of Formula (V) wherein B is $-\text{CH}_2\text{CH}_2\text{Ph}$:

5




wherein R' represents:




(7) reducing the lactone oxo group of the compound of Formula (V), using a reducing agent such as DIBAL-H, to form a compound of Formula (IV) wherein B is $-\text{CH}_2\text{CH}_2\text{Ph}$:

10




15

(8) subjecting the compound of Formula (IV) to a Wittig reaction with an ylide, the ylide being formed from the reaction of a compound of formula



wherein X represents F, Cl, Br or I, with a strong base (such as potassium *t*-butoxide), to form a compound of Formula (IIIa) or (IIIb) wherein B is -CH₂CH₂Ph:


5



or a mixture thereof;

(9) alkylating the carboxylic acid group of the compound of Formula (IIIa) or (IIIb), or a mixture thereof, with isopropyl iodide in the presence of DBU, to form the compound of Formula (IIa) or (IIb), wherein B is -CH₂CH₂Ph and R" is *iso*-propyl:


10



or a mixture thereof; and

(10) removing the protecting groups from the compound of Formula (IIa) or (IIb), or a mixture thereof, to form latanoprost:

15




The above steps (1) to (10) correspond to steps (a)-(b)-(c)-(d)-(e)-(f)-(h)-(i)-(j)-(k) in Scheme 1 above.

In an alternative process, latanoprost can be formed by a procedure involving carrying out steps (1) and (2) of the preceding process, replacing steps (3), (4) and (5) with the following steps (3'), (4') and (5'), and thereafter carrying out steps (6)-(10) as described in the preceding process. Steps (3'), (4') and (5') are as follows:


5 (3') subjecting the double bond of the compound of Formula (VIII) to a hydrogenation reaction in the presence of a hydrogenation catalyst (e.g. palladium, platinum or rhodium), the reaction being carried out optionally in the presence of sodium nitrite, to form a compound of Formula (XIII) wherein B is -CH₂CH₂Ph:

10



15 (4') reducing the side chain oxo group of the above compound of Formula (XIII) using borane-dimethylsulfide complex, the reduction being carried out in the presence of a chiral oxazaborolidine catalyst, such as (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrole[1,2c]-[1,3,2]oxazaborole, to form a compound of Formula (XIV) wherein B is -CH₂CH₂Ph:


15



20


(5') deprotecting the hydroxyl group of the compound of the above compound of Formula (XIV), preferably using K₂CO₃ in methanol, to

form a compound of Formula (VI), wherein the dashed and solid lines represent a single bond, and B is $-\text{CH}_2\text{CH}_2\text{Ph}$:




5 This procedure corresponds to steps (a)-(b)-(e')-(c')-(d')-(f)-(h)-(i)-(j)-(k) depicted in Scheme 1 and Scheme 2.

In a further preferred embodiment, there is provided a process for the synthesis of latanoprost comprising carrying out steps (1) to (8) according to either of the preceding procedures, to form a mixture comprising the compounds of Formula (IIIa) and Formula (IIIb) having the respective structures:




and further carrying out the steps of:

15 (9a) subjecting said mixture to reaction with at least one molar equivalent (preferably at least a molar equivalent, and more preferably 1.1 to 2 molar equivalents) of silylating agent to form a compound of Formula (XI) having the structure:



wherein each R' is as defined as above;


(10a) subjecting the compound of Formula (XI) to a transesterification reaction with isopropanol optionally in the presence of a weak acid catalyst such as pyridinium p-toluenesulfonate, to form the compound of Formula (XII) having the structure:



(11a) removing the protecting groups R' to form latanoprost.

These steps (9a) to (11a) correspond to steps (m)-(o)-(p) in Scheme 2 above.


10 According to a still further embodiment of the present process, there is provided a process for the production of latanoprost comprising carrying out steps (1) to (9) to form a mixture comprising the compounds of Formula (IIa) and Formula (IIb) having the respective structures:



15 and carrying out the further steps of:

(10b) subjecting said mixture to reaction with at least one molar equivalent (preferably at least 1.1 molar equivalents, and more preferably 1.15 to 2 molar equivalents) of silylating agent to form a compound of Formula (XII) having the structure:

46



; and


(11b) removing the protecting groups R' to form latanoprost.

The above steps (10b) and (11b) correspond to steps (q)-(r) shown in
5 Scheme 3 above.

According to a further aspect of the present invention, there are provided
novel intermediates for the synthesis of a compound of Formula (I-A) or (I-B)
as defined above. The novel intermediates include the following:

10

- compounds of the Formula (VIII):




wherein

A represents unsubstituted C₆ to C₁₀ aryl; and

15 B represents a substituent selected from the group consisting
of: (i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group
may be unsubstituted or substituted with one to three
substituents independently selected from the group consisting of
C₁ to C₆ alkyl, halo and CF₃; and (iii) -(CH₂)_nOR^a, wherein n
20 represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group
which may be unsubstituted or substituted with one to three


substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo or CF₃,

- compounds of the Formula (VII):




and single enantiomers thereof, wherein A and B are as defined herein;

- compounds of the Formula (VI):




and single enantiomers thereof wherein B is as defined herein;

- compounds of the Formula (VI-A):



15 and single enantiomers thereof wherein B is as defined herein;

- compounds of the Formula (V):




and single enantiomers thereof wherein B is as defined herein and R' represents the substituent:

5




wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;

10 - compounds of the Formula (IV):




and single enantiomers thereof wherein B and R' are as defined herein;

15 - compounds of the Formula (IIIa) or (IIIb):




and single enantiomers, and mixtures thereof wherein B and R' are as defined herein;

5 - compounds of the Formula (IIa) or (IIb):




and single enantiomers and mixtures thereof wherein B, R' and R'' are as defined herein;

10 - compounds of the Formula (XI):



and single enantiomers thereof wherein B and R' are as defined herein;


15 - compounds of the Formula (XII):



and single enantiomers thereof wherein B and R' are as defined herein;

- compounds of Formula XIII:

5



wherein A represents unsubstituted C₆ to C₁₀ aryl, and B is as defined herein; and

10 - compounds of Formula (XIV):


15

Preferably, in any of the structures herein, the group A represents phenyl.

20

In accordance with preferred embodiments of the present invention, the compounds of Formulae (VII), (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (XI), (XII), (XIII) and (XIV) are single enantiomers (i.e. the wavy line in the side chain represents — or ·····).


In the intermediates of Formula (VIII), (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (X), (XI), (XII) and (XIV) the group B is selected from:



Preferably, for the intermediates of Formula (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (XI) and (XII), the solid and dashed lines represent a single bond and B represents $-\text{CH}_2\text{CH}_2\text{Ph}$.

5

Further preferred are intermediates of Formula (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (XI) and (XII) wherein the solid and dashed lines represent a double bond and B represents a substituent selected from the group consisting of:



10

In the intermediates of Formula (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (XI) and (XII), the group R' is preferably:



In the compounds of Formula (XII), (IIa), (IIb) and (I-B), the group R" preferably represents *iso*-propyl.

15

The present invention further provides the use of any novel intermediate as defined as above in the manufacture of latanoprost and the use of any novel intermediate as defined as above in the manufacture of cloprosteno Δ l, fluprostenol, PGF $_{2\alpha}$, travoprost, or a PGF (preferably PGF $_{2\alpha}$) analogue.

20


The present invention also provides the use of a silylating reagent of formula



wherein the group X represents F, Cl, Br or I and R^x, R^y and R^z are as previously defined, for protecting a hydroxyl group of an intermediate in the synthesis of a prostaglandin or prostaglandin analogue, such as prostaglandin or prostaglandin analogues based on PG-A, PG-B, PG-C PG-D or PGF. The use of these silylating agents is particularly suitable in the synthesis of prostaglandin PGF_{2α} or prostaglandin analogues based on PGF_{2α}, including latanoprost, cloprosteno, fluprostenol and travoprost. Of these, latanoprost is particularly preferred.

Preferred R^x, R^y and R^z are methyl, ethyl and *tert*-butyl. A particularly preferred silylating reagent is triethylsilylchloride.

Whilst the above processes and steps refer to PGF_{2α} and analogues thereof, it will also be understood by a person of skill in the art that the process and steps could be successfully applied to the synthesis of other prostaglandin analogues, for example,



by the provision of one or more additional reaction steps in order to modify the functional groups on the cyclopentyl ring – for example, oxidation of the cyclopentyl ring hydroxyl group, in the PGF derivatives would result in either the PGD or PGE analogues.

As used herein hatched lines attached to the cyclopentane ring indicate bonds that are below the plane of the ring (i.e. bonds in an alpha configuration). Solid wedges attached to the cyclopentane ring indicate bonds that are above the plane of the ring (rings in the beta configuration). It is to be understood that a wavy line, i.e.



represents bonds in either the alpha or beta configuration, and includes single enantiomers, i.e.:




or




5 or mixtures thereof, including racemic mixtures. Thus:


Formula (XIV) includes:



Formula (XIII) includes:




Formula (XI) includes:




10


Formula (VII) includes:






Formula (VI) includes:



15





5


racemic

The single and dashed lines in each of the above formulae can represent optional double bonds. Thus, “—” represents either a single bond or a double bond.

10


It has been found that the major impurities in latanoprost, particular when produced by the methods of the present invention, include the following:

- the 15(S)-*cis* isomer of latanoprost:



15


- the 15(S)-*trans* isomer of latanoprost:



20

- and the 15(R)-*trans* isomer of latanoprost:

56



In particular, it has been found that the isomers of latanoprost are particularly difficult to remove because they have similar physical and chemical properties to latanoprost.

The present applicant has found that it is possible to obtain a separation of latanoprost from the 15(S)-*cis*-, 15(S)-*trans*- and 15(R)-*trans* isomers of latanoprost using an HPLC purification system with a chiral column. However chiral columns are expensive and are not practical for large scale separations.

These problems have been solved by the use of a non-chiral column and a particular eluent mixture in the HPLC purification of latanoprost. Accordingly, the present invention further provides a process for the purification of latanoprost by HPLC comprising the use as an eluent, of a mixture comprising a hydrocarbon, an alcohol and, optionally, acetonitrile.

The use of such an eluent combination is a particularly preferred method of purification of latanoprost when made by the processes of the present invention. It has been found that the above eluent mixtures can achieve an extremely high purity of the latanoprost.

Preferably, the eluent comprises a hydrocarbon, an alcohol and acetonitrile. Advantageously, it has been found that the use of acetonitrile as a component of the eluent in the HPLC purification of latanoprost results in an improved separation of the impurities. In particular, the use of acetonitrile as a component of the above eluent mixture results in a significantly improved separation of the hitherto difficult to separate 15(S)-*trans* isomer of

latanoprost. In particular, it has been found that the 15(S)-*trans* isomer does not co-elute with the latanoprost, i.e. a baseline separation of latanoprost of the 15(S)-*trans* isomer can be achieved. As a result, a higher yield of pure latanoprost may be obtained using the above eluent system compared with prior art procedures.

In particular the eluent systems in the present purification process comprises a hydrocarbon in an amount range of 80-99 volume percent and an alcohol in an amount range of 1-20 volume percent. Preferably, the eluent comprises a hydrocarbon in an amount range of 85-99 volume percent and an alcohol in an amount range of 1-15 volume percent. Even more preferred is an eluent comprising a hydrocarbon in an amount range of 88-98 volume percent and an alcohol in an amount range of 2-12 volume percent.

In the preferred acetonitrile-containing eluent systems of the present purification process, the eluent comprises a hydrocarbon in an amount range of 85-99 volume percent, an alcohol in an amount of 0.5-10 volume percent and acetonitrile in an amount of 0.5-5 volume percent. Preferred is an eluent comprises a hydrocarbon in an amount range of 86-98 volume percent, an alcohol in an amount of 1-8 volume percent and acetonitrile in an amount of 1-6 volume percent. Especially preferred is an eluent comprising a hydrocarbon in an amount range of 90-96 volume percent, an alcohol in an amount of 2-6 volume percent and acetonitrile in an amount of 2-4 volume percent.

In the above eluent systems it is preferred that the hydrocarbon is a C₅ to C₆ straight chain, branched or cyclic hydrocarbon, wherein the hydrocarbon is preferably an *n*-alkane. Hexane and heptane are especially preferred. The hydrocarbon employed in the eluent may comprise a mixture of e.g. alkanes, such as hexane fraction. It has been found that good results have been obtained with *n*-heptane.

Preferred alcohols in the above eluent systems are C₁ to C₈ straight chain, branched or cyclic alkanols, with C₁ to C₅ straight chain or branched alkanols being particularly preferred. Of these, methanol, ethanol, propan-1-ol, propan-2-ol, butan-1-ol or butan-2-ol are especially useful. Good results have been obtained with propan-2-ol (isopropanol) as the alcohol component. The use of ethanol as the alcohol component has also produced good separations. The alcohol component of the eluent can include mixtures of one or more alkanols, e.g. combinations of two or more of isopropanol, ethanol and methanol may be used, e.g. mixtures in ratios of 1:1 to 3:1 have been successfully employed.

A preferred eluent system comprises n-heptane : isopropanol : acetonitrile. Preferred volume percent ranges of n-heptane : isopropanol : acetonitrile are 90-96% : 2-7% : 2-5%. Preferably, the ratios of n-heptane : isopropanol : acetonitrile are in the ranges 92-94% : 3-5% : 2-4%. Good results have been obtained with ratios of 93% : 4% : 3%.

The above HPLC procedures are preferably carried out on a silica gel column. Examples of suitable columns include WatersTM Spherisorb, PhenomenexTM Luna Cyano and PhenomenexTM Luna Silica.

By using the above separation procedures, it has been found possible to produce latanoprost that is substantially free of the 15(S)-*cis* isomer, the 15(S)-*trans* isomer and the 15(R)-*trans* isomer. In particular, latanoprost containing less than 0.3% in total of any combination of: 15(S)-*cis* isomer, 15(S)-*trans* isomer and 15(R)-*trans* isomer may be produced. Using the process of the present invention, latanoprost containing less than 0.1% each of 15(S)-*cis*-, 15(S)-*trans*- and 15(R)-*trans* isomers can be produced.


The above processes thus enable latanoprost having an extremely high degree of purity can be obtained, e.g. greater than 98% pure, greater than

99% pure, greater than 99.5% pure. Indeed, it has been found possible to achieve latanoprost purities of greater than 99.8%.

The following abbreviations have been used:

- 5 HPLC = high pressure liquid chromatography
- MPLC = medium pressure liquid chromatography
- L-Selectride = lithium tri-sec-butylborohydride
- DMS = dimethylsulfide
- DIBAL-H = di-*iso*-butylaluminiumhydride
- 10 PPB = *para*-phenylbenzoyl
- DBU = [1,8-diazabicyclo(5.4.0)undec-7-ene]
- ⁿBuLi = *n*-butyllithium
- TEMPO, free radical = 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical
- EtOAc = ethylacetate
- 15 DMF = dimethylformamide
- TsOH = *para*-toluenesulphonic acid
- KO'Bu = potassium *tert*-butoxide

Benzoyl Corey lactone =



- 20 As used herein the term "alkyl" refers to C₁ to C₆ straight or branched carbon chains. Particularly preferred alkyl groups for the compounds and processes of the invention include methyl, ethyl, propyl, isopropyl, butyl, and tertiary butyl.
- 25 The term "aryl" represents a carbocyclic group containing from six to fifteen carbon atoms and having at least one aromatic ring. Particularly preferred aryl groups for any of the compounds and methods of the present invention include phenyl and naphthyl.

The term "aralkyl" refers to an alkyl group as defined above wherein one or more hydrogen atoms (preferably one) have been replaced by unsubstituted C₆ to C₁₀ aryl groups as defined above. A preferred aralkyl group for the compounds and methods of the invention is benzyl.

5

The term "halo" refers to fluoro, chloro, bromo or iodo.

The term "substantially free" of a particular impurity refers to less than 1%, preferably less than 0.5%, more preferably, less than 0.3% and even more 10 preferably, less than 0.1% of the impurity.

The following examples illustrates the preparation of latanoprost in accordance with the present invention:

EXAMPLES

Solvents (chromatography grade) were dried over 3Å molecular sieves prior to use. Purified water was obtained from Loveridge.

5

The products obtained in the reactions were sealed under argon and stored in a cold room at 4°C until required.


Column chromatography was carried out on Merck silica gel 60.

10

As a precaution, where indicated, reactions were carried out under argon. Reaction progress was monitored by thin layer chromatography.

Unless indicated otherwise, final products were dried under high vacuum at a pressure of about 0.01 kPa.

EXAMPLE 1: Preparation of (1*S*, 5*R*, 6*R*, 7*R*)-7-Benzoyloxy-6-formyl-2-oxabicyclo[3.3.0]octan-3-one (PGX-5)




A 0.5M solution of potassium bromide was prepared by dissolving KBr (11.9g) in purified water and then diluting with additional water to 200ml.

2.1M Sodium hypochlorite solution (476ml, 1M) was diluted to 1L by the addition of purified water to give a 1M solution. The pH was then adjusted to 8.83 by the addition, with stirring, of solid sodium bicarbonate.

5 (-)-Benzyl Corey Lactone (PGX-4) (200g, 0.724M, 1eq.) was dissolved in dichloromethane (2L) under an inert atmosphere and 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) free radical (1.13g, 7.24mmol, 0.01eq.) and the aqueous potassium bromide solution prepared above added. The mixture was cooled to approximately 0°C with stirring and a portion of the aqueous 10 sodium hypochlorite solution prepared above (100ml) added at this temperature. The reaction was allowed to warm to 10°C and an additional quantity of the aqueous sodium hypochlorite solution (896ml) added at 10-15°C over 40 minutes. After stirring for 20 minutes the reaction was shown to be complete by TLC. The layers were separated, the aqueous phase 15 extracted with dichloromethane (200ml) and the combined organic phases then washed with water (200ml) before drying over magnesium sulfate. Filtration and washing of the filtercake with dichloromethane (2 x 200ml) gave (1S, 5R, 6R, 7R)-7-benzoyloxy-6-formyl-2-oxabicyclo[3.3.0]octan-3-one (PGX-5) as a solution in dichloromethane.

20

EXAMPLE 2: Preparation of Dimethyl (2-oxo-4-phenylbutyl)-phosphonate (PGX-3)




25

Dimethyl methyl phosphonate (PGX-1) (346.2g, 2.79M, 1.95eq.) was dissolved in tetrahydrofuran (2.4L) and the resulting solution cooled to -70°C under an inert atmosphere. n-Butyllithium (2.5M in tetrahydrofuran, 1.0L,

2.5M, 1.75eq.) was added over 105 minutes whilst maintaining the temperature below -65°C. The resulting white suspension was stirred at approximately -70°C for 20 minutes. A solution of methyl 3-phenyl propionate (PGX-2) (234.8g, 1.43M, 1.0eq.) in tetrahydrofuran (400ml) was then added over 30 minutes whilst maintaining the temperature below -65°C. The mixture was left to stir at approximately -70°C for 2.5h until the reaction was complete by TLC. After allowing to warm to approximately 0°C the reaction was quenched by addition of water (800ml). tert-Butylmethyl ether (1.0L) was added and the mixture allowed to separate. The organic phase was extracted with water (5 x 500ml), and the combined aqueous phases acidified to pH 2 by the addition of 10% hydrochloric acid (ca. 110ml). The product was extracted into tert-butylmethyl ether (2 x 800ml, then 5 x 500ml) and the combined organics dried over magnesium sulfate. Filtration, washing the filtercake with tert-butylmethyl ether (400ml) and evaporation gave the crude product as an oil (240.4g). Distillation under reduced pressure (bp 152-156°C at 8 x 10⁻⁵ – 1 x 10⁻⁴ torr) gave dimethyl (2-oxo-4-phenylbutyl)phosphonate as an almost colourless oil (214.1g 58%).

EXAMPLE 3: Preparation of (1S, 5R,6R,7R)-7-Benzoyloxy-6-[3-oxo-5-phenyl-(E)-1-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-6)




The phosphonate ester (PGX-3) (176.2g, 0.69M, 0.95eq.) was dissolved in dry acetonitrile (1.8L) in an inert atmosphere and lithium chloride (153.4g, 3.62M, 5.0eq.) added. The mixture was cooled to approximately -15°C and dry diisopropyl ethylamine (178.1g, 1.38M, 1.9eq.) added over 10 minutes

whilst maintaining this temperature. After stirring for about 100 minutes a solution of the aldehyde (PGX-5) in dry dichloromethane (from Example 1) (assumed 0.72M, 1.0eq.) was added over 20 minutes whilst maintaining the temperature at ca. -15°C. The reaction was then allowed to warm to ambient temperature with stirring until TLC showed it was complete. Water (200ml) was added and the mixture evaporated *in vacuo* to remove the majority of the acetonitrile. The residual slurry was partitioned between water (2.0L) and ethyl acetate (2.0L). The organic phase was separated off and the aqueous extracted with ethyl acetate (2 x 500ml). The combined organics were washed with saturated brine (2 x 1L) and dried over magnesium sulfate. After filtration, the solvent was evaporated off to a residual weight of ca. 540g and hexane (1L) then added to the residue. After cooling to 4°C with stirring for 90 minutes the solid was filtered off, washed with hexane:ethyl acetate 4:1 (2 x 200ml) and dried *in vacuo* at ambient temperature to give (1S,5R,6R,7R)-7-benzyloxy-6-[3-oxo-5-phenyl-(E)-1-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one as a white solid (225.8g, 77%).

EXAMPLE 4: Preparation of (R)-Tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo[1,2-c][1,3,2]oxazaborole (Corey Catalyst)


20



(R)-(+)-1,1-Diphenylprolinol (15.20g, 60.0mMol) was dissolved in toluene (600ml) under an inert atmosphere and trimethylboroxine (7.57g, 60.3mMol, 1.0eq.) added. The mixture was heated to reflux and approximately half of the toluene was distilled off via a Dean-Stark trap. Further toluene (350ml) was added and this was also distilled off via the Dean-Stark trap until a total of 710ml of distillate had been collected. The reaction mixture was then allowed

to cool to ambient temperature to give (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo[1,2-c][1,3,2]oxazaborole as a 0.25M solution in toluene.

5 **EXAMPLE 5: Preparation of (1S,5R,6R,7R)-7-Benzoyloxy-6-[3(S)-3-hydroxy-5-phenyl-1 (E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-7)**




10 PGX-6 (225.1g, 0.556M) was dissolved in dry tetrahydrofuran (3.5L) in an inert atmosphere and 'Corey catalyst' prepared according to Example 4 (0.25M in toluene, 223ml, 0.1eq.) added. The mixture was cooled to approximately -15°C and a solution of borane-methyl sulfide complex (10M BH₃ concentration, 41.7ml, 417mMol, 0.75eq.) in dry tetrahydrofuran (450ml) was added whilst maintaining the temperature at *ca.* -15°C. The mixture was 15 then stirred at this temperature for 2h until the reaction was shown to be complete by TLC. Methanol (200ml) was added over 30 minutes and the mixture allowed to warm to approximately 0°C before acidification with 10% aqueous hydrochloric acid (1.2L). Ethyl acetate (2.4L) was added and the layers allowed to separate. The aqueous phase was extracted with ethyl 20 acetate (1 x 0.8L, 1 x 0.4L) and the combined organic phases washed with water (1 x 1.6L, 3 x 0.8L) and then brine (1L). Drying over magnesium sulfate, filtration and evaporation *in vacuo* gave the crude product (containing a mixture of the desired 3(S)-isomer, PGX-7, and the undesired 3(R)-isomer 25 PGX-8) as an oil (242.7g). This was dissolved in dichloromethane : ethyl acetate 2:1 (720ml) and chromatographed on silica gel 60 (12.8 Kg) eluting with dichloromethane : ethyl acetate 2:1. Impure fractions containing a mixture of PGX-7 and PGX-8 were evaporated and re-chromatographed

under the same conditions. Fractions free from PGX-8 (by TLC) were combined and evaporated. Drying *in vacuo* at ambient temperature gave the desired product (1S,5R,6R,7R)-7-benzoyloxy-6-[3(S)-3-hydroxy-5-phenyl-1(E)-pentenyl]-2-oxa-bicyclo[3.3.0]octan-3-one as a white solid (167.9g, 74.2%).

5

EXAMPLE 5a: Preparation of (1S,5R,6R,7R)-7-Benzoyloxy-6-[3(S)-3-hydroxy-5-phenyl-1 (E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX 7)



10

PGX 6 (315.2g, 0.779M) was dissolved in dry tetrahydrofuran (5.7L) in an inert atmosphere and 'Corey catalyst' (0.25M in toluene, 312ml, 0.1eq.) added. The mixture was cooled to approximately -15°C and a solution of borane-methyl sulfide complex (10M BH₃ concentration, 58.5ml, 0.585M, 0.75eq.) in dry tetrahydrofuran (630ml) added whilst maintaining the temperature at *ca.* -15°C. The mixture was then stirred at this temperature for 2h until the reaction was shown to be complete by TLC. Methanol (290ml) was added over 30 minutes and the mixture allowed to warm to approximately 0°C before acidification with 10% aqueous hydrochloric acid (1.7L). Ethyl acetate (3L) was added and the layers allowed to separate. The aqueous phase was extracted with ethyl acetate (1 x 1.1L, 1 x 0.55L) and the combined organic phases washed with water (1 x 2L, 3 x 1L) and then brine (1.5L). Drying over magnesium sulfate, filtration and evaporation *in vacuo* gave the crude product (containing a mixture of PGX 7 and the undesired isomer PGX 8) as an oil (349.3g). The oil was purified by the procedure given in Example 5b.

15

20

25

30

NOTE:

The chromatographic procedure described in Example 5 in which a 2:1 mixture of dichloromethane : ethylacetate is employed as an eluent for

medium pressure chromatography (MPLC) on a silica gel column, suffers from the disadvantage that the column must be re-packed after each separation before subsequent aliquots of the epimeric product mixture can be processed. Thus, such a procedure employs large quantities of both stationary phase and eluent, as well as being time consuming. An improved procedure is given below (Example 5b):

EXAMPLE 5b: Separation of (1S,5R,6R,7R)-7-Benzoyloxy-6-[3(S)-3-hydroxy-5-phenyl-1-(E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-7) from (1S,5R,6R,7R)-7-Benzoyloxy-6-[3(R)-3-hydroxy-5-phenyl-1-(E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-8)

According to an improved procedure, the crude mixture of epimers PGX-7/PGX-8 (686.5g) made according to the synthetic procedure described in Example 5a was crystallised from heptane fraction/ethyl acetate (7:3) to give a crystalline mixture of epimers PGX-7 and PGX-8 (480.4g) that is free of other impurities. The filtrates from the crystallisation were evaporated to give an oil (150.3g) comprising mainly impure PGX-7/PGX-8.

The crystalline mixture was dissolved in dichloromethane:ethyl acetate (2:1) mixture to give a stock solution (A). The evaporated impure PGX-7/PGX-8 mixture was dissolved in dichloromethane:ethyl acetate (2:1) mixture to give a second stock solution (B). The isomer PGX-7 was then separated from PGX-8 by medium pressure chromatography (MPLC) on a silica gel column by sequential injection of aliquots of the stock solutions according to the following protocol:

Injection 1: Injection of an aliquot of crystalline PGX-7/PGX-8 stock solution (A) and collection of appropriate fractions containing pure PGX-7 and impure PGX-7. The column was then flushed with eluent to elute any remaining PGX-8.

Injection 2: Injection of an aliquot of crystalline PGX-7/PGX-8 stock solution (A) and collection of appropriate fractions containing pure PGX-7 and impure PGX-7. Column flushed with methanol then equilibration of column with eluent.


5 Injection 3: Injection of an aliquot of impure PGX-7/PGX-8 stock solution (B) and collection of appropriate fractions containing pure PGX-7 and impure PGX-7 and impure PGX-8. Discarded silica gel and re-packed column with fresh silica gel and repeat cycle.

10 The above strategy increases the throughput of PGX-7 considerably and avoids the need to re-pack the column after every injection.

15 The column fractions containing the impure PGX-7 are recycled to give more pure PGX-7 utilising the above protocol, except that more than three injections per column can be performed before re-packing the column.

Fractions free from PGX-8 (by TLC) were combined and evaporated. Drying *in vacuo* at ambient temperature gave (1S,5R,6R,7R)-7-benzoyloxy-6-[3(S)-3-hydroxy-5-phenyl-1(E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-7) as a white solid (445.2g).


EXAMPLE 6: Preparation of (1S, 5R, 6R, 7R)-7-Hydroxy-6-[3(S)-3-hydroxy-5-phenyl-1 (E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-9)



PGX-7 (152.0g, 0.374M, 1.0eq.) was dissolved in dry methanol (2.28L) under an inert atmosphere and potassium carbonate (31.0g, 0.224M, 0.6eq.) added in one portion. The mixture was stirred at ambient temperature for 3h until TLC showed the reaction was complete. 5M Hydrochloric acid (65.5ml) was added to adjust the apparent pH of the solution to about 6.8-7.0 and the mixture then evaporated to dryness *in vacuo*. The sticky residue was treated with water (1.5L) and the pH adjusted to 6.8-7.0 by the addition of 1M hydrochloric acid (7ml). Heptane (0.45L) was added, the mixture agitated vigorously and the precipitated solid filtered off and washed with heptane (2 x 150ml) on the filter. The solid was then triturated with a further quantity of heptane (2 x 150ml). All the heptane washes were combined and added to the original filtrates. The aqueous phase was separated off, washed with heptane (2 x 150ml) and then extracted with ethyl acetate (1 x 450ml, 3 x 150ml). The previously isolated solid was added to the combined ethyl acetate extracts and the mixture shaken until a solution formed. This was washed with saturated brine (2 x 250ml) and dried over magnesium sulfate. Filtration and evaporation *in vacuo* gave (1S,5R,6R,7R)-7-hydroxy-6-[3(S)-3-hydroxy-5-phenyl-(E)-pentenyl]-2-oxabicyclo[3.3.0]octan-3-one as a pale yellow oil (112.1g, 99.1%).

20

EXAMPLE 7: Preparation of (1S,5R,6R,7R)-7-Hydroxy-6-[3(R)-(3-hydroxy-5-phenyl)pentyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX-10)



25

PGX-9 (111.5g 0.369M, 1.0eq.) was dissolved in ethanol (1.67L) and 5% palladium on carbon (5.58g) added followed by a solution of sodium nitrite


(8.90g, 0.129M, 0.35eq.) in water (100ml). The mixture was then hydrogenated for 5h until shown to be complete by TLC. 1M Hydrochloric acid (260ml) was added and the mixture stirred for 1h. The solids were removed by filtration through celite, the filtrates then evaporated to give an oily-solid residue which was partitioned between ethyl acetate (0.45L) and water (0.45L). The layers were separated and the aqueous extracted with ethyl acetate (2 x 0.33L). The combined organics were washed with brine (2 x 0.22L) and then dried over sodium sulfate containing 5% sodium hydrogen carbonate. Filtration and evaporation gave (1S, 5R, 6R, 7R)-7-Hydroxy-6-[3(R)-(3-hydroxy-5-phenyl)pentyl]-2-oxabicyclo[3.3.0]octan-3-one as a yellow oil (110.4g, 98.4%)

NOTE:

It has been found to be possible to reduce the time period during which the product PGX-10 is in the acid medium during the HCl work-up, and thus minimise the potential of an acid-catalysed epimerisation of the side-chain hydroxyl group of (1S,5R,6R,7R)-7-Hydroxy-6-[3(R)-(3-hydroxy-5-phenyl)pentyl]-2-oxabicyclo[3.3.0]octan-3-one

Thus, in the modified procedure, on completion of the reaction described in Example 7, 1M hydrochloric acid was added and the reaction mixture stirred for 60 minutes as described in Example 7. The pH of the mixture was adjusted to between 5 and 6 by addition of solid sodium hydrogen carbonate prior to the filtration of the used catalyst, and evaporation of the filtrate to dryness. The work-up is then completed as described in Example 7 by extraction of the product into ethyl acetate. The following Example 7a illustrates the improved procedure:

EXAMPLE 7a: Preparation of (1S, 5R, 6R, 7R)-7-Hydroxy-6-[3(R)-(3-hydroxy-5-phenyl)pentyl]-2-oxabicyclo[3.3.0]octan-3-one (PGX 10)




5

PGX 9 (326.5g, 1.08M, 1.0eq.) was dissolved in ethanol (6.5L) and 5% palladium on carbon (16.3g) added followed by a solution of sodium nitrite (26g, 0.377M, 0.35eq.) in water (200ml). The mixture was then hydrogenated for 1.5h until shown to be complete by TLC. 1M Hydrochloric acid (750ml) was added and the mixture stirred for 1h. The pH was adjusted to 5-6 by the addition of solid sodium hydrogen carbonate (55g). The solids were removed by filtration through celite, the filtrates then evaporated to give an oily-solid residue which was partitioned between ethyl acetate (1.3L) and water (1.3L). The layers were separated and the aqueous extracted with ethyl acetate (2 x 0.975L). The combined organics were washed with brine (2 x 0.65L) and then dried over sodium sulfate. Filtration and evaporation gave (1S, 5R, 6R, 7R)-7-Hydroxy-6-[3(R)-(3-hydroxy-5-phenyl)pentyl]-2-oxabicyclo[3.3.0]octan-3-one as a yellow oil (311.5g, 94.8%).

10


15

EXAMPLE 8: Preparation of (1S, 5R, 6R, 7R)-6-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-7-triethylsilyloxy-2-oxabicyclo[3.3.0]octan-3-one (PGX-11)




PGX-10 (109.7g, 0.360M, 1.0eq.) was dissolved in dry dimethyl formamide (720ml) under an inert atmosphere. Imidazole (29.4g, 0.432M, 1.2eq.) and triethylamine (102.9ml, 74.69g, 0.738M, 2.05eq.) were added and the mixture then cooled to approximately 0°C. Triethylchlorosilane (111.2g, 0.738M, 2.05eq.) was added over 15 minutes at less than 10°C. The mixture was allowed to warm to room temperature and stirred for 2h until TLC showed the reaction was complete. After re-cooling to 10°C, hexane (0.55L) and water (1L) were added and the layers separated. The aqueous phase was extracted with hexane (1 x 0.2L, 1 x 0.1L) and the combined organics then washed with water (2 x 0.5L) and brine (2 x 0.5L). Drying over magnesium sulfate, filtration and evaporation *in vacuo* gave (1S, 5R, 6R, 7R)-6-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-7-triethylsilyloxy-2-oxabicyclo[3.3.0] octan-3-one as a yellow oil (187.4g, 97.6%).

EXAMPLE 9: Preparation of (1*S*, 3*RS*, 5*R*, 6*R*, 7*R*)-6-[3(*R*)-5-phenyl-3-triethylsilyloxy]pentyl]-7-triethylsilyloxy-2-oxabicyclo[3.3.0]octan-3-ol (PGX-12)



PGX-11 (186.8g, 0.3505M, 1.0eq.) was dissolved in dry tetrahydrofuran (1.86L) under an inert atmosphere and the solution cooled to less than -70°C. Diisobutylaluminium hydride (1.1M in toluene solution, 701ml, 0.7711M, 2.2eq.) was added to the reaction whilst maintaining the temperature below -70°C. The mixture was then stirred at this temperature for 2h until the reaction was shown to be complete by TLC. Methanol (132ml) was added and the mixture allowed to warm to -5°C before addition of water (2L) followed by acidification to pH 3 with 2M aqueous sodium hydrogen sulphate solution (1.54L, 8.8eq.). Ethyl acetate (0.66L) was added and the layers allowed to separate. The aqueous phase was extracted with ethyl acetate (1 x 0.26L, 1 x 0.13L) and the combined organics then washed with water (2 x 1.3L) and brine (2 x 1.3L). Drying over magnesium sulfate and evaporation *in vacuo* gave (1S,3RS,5R,6R,7R)-6-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-7-triethylsilyloxy-2-oxabicyclo[3.3.0]octan-3-ol as an almost colourless oil (182.5g).

EXAMPLE 10: Preparation of (Z)-7-((1R, 2R, 3R, 5S)-5-Hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-3-(triethylsilyloxy)cyclopentyl)-5-heptenoic acid (PGX-13) and (Z)-7-((1 R, 2R, 3R, 5S)-3-Hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy) pentyl]-5-(triethylsilyloxy)cyclopentyl)-5-heptenoic acid (PGX-14)



(4-Carboxybutyl)-triphenylphosphonium bromide (459.0g, 1.0353M, 3.0eq.) was suspended in dry tetrahydrofuran (1.84L) under an inert atmosphere and the mixture cooled to 0°C. 1.07M Potassium tert-butoxide in tetrahydrofuran solution (1.806L, 1.93M, 5.6eq.) was then added over 60 minutes at 0°C. After the addition was complete the reaction was allowed to warm to approximately 20°C and stirred for 1 hour before re-cooling to 0°C. A solution of PGX-12 (184.6g, 0.3451M, 1.0eq.) in tetrahydrofuran (0.27L) was then added at this temperature. The mixture was allowed to warm to room temperature and stirred for 75 minutes until TLC showed the reaction was complete. After re-cooling to 0°C the mixture was quenched by the addition of water (3L) and then acidified to pH 5 with 5% aqueous citric acid solution (1.6L). The product was extracted into ethyl acetate (1 x 1.0L, 2 x 0.4L) and the combined organics then washed with brine (2 x 0.7L). Drying over magnesium sulfate, filtration and evaporation *in vacuo* gave a mixture of (Z)-7-((1R, 2R, 3R, 5S)-5-hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-3-(triethylsilyloxy)cyclopentyl)-5-heptenoic acid and (Z)-7-((1 R, 2R, 3R, 5S)-3-hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy) pentyl]-5-(triethylsilyloxy)cyclopentyl)-5-heptenoic acid.

hydroxy-2-[3(R)-(5-phenyl-3-triethyl-silyloxy) pentyl]-5-(triethylsilyloxy)cyclopentyl}-5-heptenoic acid as a yellow oil (461.7g). This crude product containing triphenylphosphine oxide by-product was used without purification in the next step of the synthesis.

5

NOTE:

The amount of (4-carboxybutyl)triphenylphosphonium bromide used in the above reaction is 3.0 equivalents with respect to the amount of starting material PGX-12 used. In order to generate the ylide and the carboxylate salt, it is treated with 5.6 equivalents of potassium *tert*-butoxide. The slight deficit in the amount of potassium *tert*-butoxide used with respect to (4-carboxybutyl)triphenylphosphonium bromide (2.0 equivalents required; 1.87 equivalents used) is deliberate to ensure that all of the potassium *tert*-butoxide is consumed and is not present during the reaction with the lactol.

10


However, the present applicant has found that 2.15 equivalents of (4-carboxybutyl)triphenylphosphonium bromide and 4.0 equivalents of potassium *tert*-butoxide with respect to the amount of the lactol starting material PGX-12 can be used. Advantageously, this leads to a better conversion of PGX-12 to PGX-13/PGX-14. This is illustrated in Example 10a:

15

20

**EXAMPLE 10a: Preparation of (Z)-7-((1R, 2R, 3R, 5S)-5-Hydroxy-2-[3(R)-
(5-phenyl-3-triethylsilyloxy)pentyl]-3-(triethylsilyloxy)cyclopentyl]-5-
hept-enoic acid (PGX-13) and (Z)-7-((1 R, 2R, 3R, 5S)-3-Hydroxy-2-[3(R)-
(5-phenyl-3-triethylsilyloxy)pentyl]-5-(triethylsilyloxy)cyclopentyl]-5-
hept-enoic acid (PGX-14)**

5



10 (4-Carboxybutyl)-triphenylphosphonium bromide (965.5g, 2.18M, 2.15eq.) was suspended in dry tetrahydrofuran (4.4L) under an inert atmosphere and the mixture cooled to 0°C. 0.98M Potassium tert-butoxide in tetrahydrofuran solution (4.135L, 4.05M, 4.0eq.) was then added over 60 minutes at 0°C. After the addition was complete the reaction was allowed to warm to approximately 20°C and stirred for 1 hour before re-cooling to 0°C. A solution of PGX-12 (542.3g, 1.013M, 1.0eq.) in tetrahydrofuran (1L) was then added at this temperature. The mixture was allowed to warm to room temperature and stirred for 60 minutes until TLC showed the reaction was complete. After re-cooling to 0°C the mixture was quenched by the addition of water (8.6L) and then acidified to pH 5 with 5% aqueous citric acid solution (4.5L). The product was extracted into ethyl acetate (1 x 3L, 2 x 1.2L) and the combined organics then washed with brine (2 x 2L). Drying over magnesium sulfate, filtration and evaporation *in vacuo* gave a mixture of (Z)-7-((1R, 2R, 3R, 5S)-5-hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-3-(triethylsilyloxy)cyclopentyl]-5-heptenoic acid and (Z)-7-((1 R, 2R, 3R, 5S)-3-hydroxy-2-[3(R)-

15


20

(5-phenyl-3-triethylsilyloxy) pentyl]-5-(triethylsilyloxy)cyclo-pentyl]-5-heptenoic acid as a yellow oil (1095.6g). This crude product containing triphenylphosphine oxide by-product was used without purification in the next step of the synthesis.

5

EXAMPLE 11: Preparation of Isopropyl (Z)-7-[(1R, 2R, 3R, 5S)-5-Hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-3-(triethylsilyloxy)-cyclo-pentyl]5-heptenoate (PGX-15) and Isopropyl (Z)-7-[(1R, 2R, 3R, 5S)-3-Hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-5-(triethylsilyloxy)cyclopentyl]-5-heptenoate (PGX-16)


10



The PGX-13/14 mixture (461.1g, assumed to contain 0.3446M, 1.0eq.) was dissolved in acetone (2.13L) under an inert atmosphere and dry 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (290.1ml, 1.94M, 5.63eq.) added at ambient temperature. After stirring for approximately 15 minutes, 2-iodopropane (329.7g, 1.94M, 5.63eq.) was added and the mixture then stirred for 16h at ambient temperature. TLC showed that the reaction was incomplete. An additional quantity of DBU (19.1ml, 0.1275M, 0.37eq.) was added followed, after 20 minutes, by 2-iodopropane (21.7g, 0.1275M, 0.37eq.). After stirring for 4.5h at ambient temperature the reaction was shown to be complete by TLC. The bulk of the acetone (ca. 1.72L) was evaporated *in vacuo* and the residue partitioned between ethyl acetate


(0.67L) and 3% aqueous citric acid solution (1.6L). The organic layer was separated off and the aqueous layer acidified to pH 6 with an additional quantity of 3% aqueous citric acid solution (0.72L) and then re-extracted with ethyl acetate (2 x 0.33L). The combined organic phases were washed with 5 3% aqueous citric acid solution (2 x 0.8L), 5% sodium hydrogen carbonate solution (2 x 0.8L) and saturated brine (2 x 1.6L). After drying over magnesium sulfate followed by filtration the solvent was evaporated off and heptane (1.0L) and ethyl acetate (80ml) added to the residue. The mixture was cooled to -20°C and agitated vigorously. After 30 minutes at -20°C the 10 precipitated solid was filtered off and washed on the filter with heptane : ethyl acetate 9:1 (6 x 200ml). Evaporation of the filtrates *in vacuo* gave a yellow oil. Heptane (600ml) was added and the solution cooled to -20°C. After 30 minutes the precipitate was filtered off and washed on the filter with heptane (3 x 100ml). The filtrate was evaporated *in vacuo* to give a mixture of 15 isopropyl (Z)-7-[(1R,2R,3R,5S)-5-hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-3-(triethylsilyloxy)cyclopentyl]-5-heptenoate and isopropyl (Z)-7-[(1R,2R,3R,5S)-3-hydroxy-2-[3(R)-(5-phenyl-3-triethylsilyloxy)pentyl]-5-(triethylsilyloxy)cyclo-pentyl]-5-heptenoate as a yellow oil (238.4g).

EXAMPLE 12: Preparation of Isopropyl (Z)-7-[(1R, 2R, 3R, 5S)-3,5-bis(triethylsiloxy)-2-[3(R)-(5-phenyl-3-triethylsiloxy)pentyl]-cyclopentyl]-5-heptenoate (PGX-17)



The PGX-15/16 mixture (237.79g, assumed to contain 0.344M) was dissolved in dry dimethylformamide (700ml) under an inert atmosphere. Imidazole (14.04g, 0.206M, 0.6eq.) and triethylamine (50.3ml, 0.361M, 1.05eq.) were added and the mixture cooled to approximately 0°C. Triethylchlorosilane (60.6ml, 54.4g, 0.361M, 1.05eq.) was then added and the mixture allowed to warm to ambient temperature. Stirring was continued for 2h until the reaction was shown to be complete by TLC. The reaction mixture was then partitioned between heptane (1.07L) and water (2.67L). After separation, the aqueous phase was further extracted with heptane (2 x 0.27L). The combined organics were washed with water (2 x 1.3L) and then saturated brine (2 x 1.3L). After drying over magnesium sulfate followed by filtration the solvent was evaporated *in vacuo* to give the crude product (235.0g) as a thick yellow oil. This was dissolved in heptane (90ml) and purified by column chromatography on silica gel (352.5g) eluting with heptane and then heptane : ethyl acetate 95:5. The relevant fractions were combined and the solvent evaporated *in vacuo* to give isopropyl (Z)-7-[(1R,2R,3R,5S)-3,5-bis(triethylsiloxy)-2-[3(R)-(5-phenyl-3-triethylsiloxy)pentyl]-cyclopentyl]-5-heptenoate as a pale yellow oil (210.5g).

EXAMPLE 13: Preparation of Isopropyl (Z)-7-[(1R, 2R, 3R, 5S)-3,5-dihydroxy-2-[3(R)-(3-hydroxy-5-phenyl)pentyl]cyclopentyl]-5-heptenoate (Latanoprost)



PGX-17 (0.504 g, 0.65 mmol) was weighed into a 50 ml round bottom flask equipped with a magnetic follower. Acetone (6.5 ml) was added and the resulting colourless solution stored at room temperature under a gentle stream of argon.


10 A solution of pyridinium-p-toluenesulfonate (Aldrich, 13 mg) in water (0.9 ml) was prepared and added to the reaction solution. The resulting white cloudy suspension was stored at room temperature under a stream of argon. The 15 mixture cleared after about 20 min to give a colourless solution. TLC after 3hr indicated complete conversion to the required product. The stirring was then stopped and the reaction mixture concentrated under reduced pressure to remove the acetone. The residue was partitioned between ethylacetate (10 ml) and saturated NaCl (15 ml). The organic phase was separated and the aqueous layer extracted with more ethylacetate (2 x 7.5 ml). The 20 combined organic extracts were washed with saturated NaCl (2 x 10 ml), dried over $MgSO_4$ (0.5 g) and evaporated to dryness under reduced pressure. The crude product was obtained as a colourless oil.

25 The crude product was purified by flash column chromatography. A flash column was prepared using silica gel 60 (6 g) and hexane fraction/EtOAc (1:1) as the eluent. The crude product (7 mg was removed as retention sample) was dissolved in the eluent (2 ml) and loaded onto the column. The column was then eluted with hexane fraction/EtOAc mixtures as follows:

Hexane fraction/EtOAc	1:1	50 ml
	1:2	150 ml
	1:3	80 ml

5 The fractions containing the pure product were combined and evaporated to dryness under reduced pressure. The pure product was obtained as a colourless viscous oil which was dried under high vacuum (0.01 kPa) to constant weight (0.251 g, yield 89.3%).

10 **EXAMPLE 13a: Preparation of Isopropyl (Z)-7-[(1R, 2R, 3R, 5S)-3,5-dihydroxy-2-[3(R)-(3-hydroxy-5-phenylpentyl)cyclopentyl]-5-heptenoate (Latanoprost, R23)**



15 PGX-17 (170.0g, 0.219M) was dissolved in acetone (1.9L) under an inert atmosphere and a solution of pyridinium-p-toluenesulphonate (4.52g, 18.0mMol, 0.08eq.) in water (0.3L) added. The resulting mixture was stirred at ambient temperature for 3h until TLC showed the reaction to be complete. After evaporation of the organic volatiles the residue was added to a mixture of ethyl acetate (2.1L) and brine (2L). The layers were separated and the aqueous phase further extracted with ethyl acetate (1L). The organics were combined and washed with brine (0.5L). Evaporation *in vacuo* gave a pale yellow oil (166.4g). This was dissolved in hexane : ethyl acetate 1:1 (0.25L) and purified by chromatography on silica gel (1.87Kg) eluting with hexane : ethyl acetate (1:1 then 1:2 then 1:3). The relevant fractions were combined and evaporated to give an almost colourless oil (78.0g). This was further purified by isocratic preparative HPLC on a silica column eluting with heptane (88-95%) : methanol/isopropanol (1/2) (5-12%). Evaporation of the relevant combined cuts gave isopropyl (Z)-7-[(1R,2R,3R,5S)-3,5-dihydroxy-2-[3(R)-(3-

20

25

hydroxy-5-phenyl)pentyl]cyclopentyl]-5-heptenoate (latanoprost) as an almost colourless oil (50.2g, 53.0%).

EXAMPLE 13b: HPLC Purification of latanoprost

5

HPLC purification of latanoprost was carried out using a WatersTM Spherisorb silica gel column. The isocratic eluent system comprised a hydrocarbon and an alcohol in volume percent ranges of 88-98% and 2-12% respectively. The hydrocarbons used were n-hexane, hexane fraction, n-heptane or heptane fraction. The alcohols used were isopropanol, ethanol or methanol, either singly or in combination in ratios of 1:1 to 3:1.

10

The results of typical HPLC runs using the above eluent system are shown below:

15

Product	Amount	
	Run 1	Run 2
Latanoprost	99.89%	99.83%
15(S)- <i>cis</i> isomer	0.08%	0.09%
15(S)- <i>trans</i> isomer	0.03%	0.07%
15(R)- <i>trans</i> isomer	not detected	not detected

Thus, it can be seen that the above solvent system achieves a latanoprost purity of greater than 99.8% and the amounts of the undesired isomers are all less than 0.1%.

20

EXAMPLE 13c: HPLC Separation of latanoprost and isomers

HPLC separations of latanoprost were carried out using a WatersTM Spherisorb silica gel column. The isocratic eluent system comprised hydrocarbon : alcohol : acetonitrile in volume percent ratios of 90-96% : 2-6% : 2-4%. The hydrocarbons used were *n*-hexane, hexane fraction, *n*-heptane or heptane fraction. The alcohols used were either isopropanol or ethanol.

10 The following results are shown for a HPLC separation of latanoprost using as eluent, a mixture of heptane : isopropanol : acetonitrile in the volume percent ratios of 93% : 4% : 3%. The relative retention times of latanoprost, and the 15(S)-*cis*, 15(S)-*trans* and 15(R)-*trans* isomers are shown below:


Product	Relative Retention Time
Latanoprost	1.00
15(S)- <i>cis</i> isomer	0.95
15(S)- <i>trans</i> isomer	1.13
15(R)- <i>trans</i> isomer	1.23

15 The above results indicate that an excellent degree of separation of latanoprost from the undesired isomers thereof can be achieved using an eluent system comprising acetonitrile. Similar results were achieved with PhenomenexTM Luna Cyano and PhenomenexTM Luna Silica columns.

20 Using this procedure, it is possible to achieve latanoprost that is substantially pure, or is completely free, of the hitherto difficult to remove isomers.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A process for the synthesis of a compound of Formula (I-A) or (I-B):



wherein:


5 B represents a substituent selected from the group consisting of:

(i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo and CF₃ and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which may be unsubstituted 10 or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo or CF₃; and

R["] represents a C₁ to C₆ alkyl group or a C₃ to C₈ cycloalkyl group;

comprising the steps of:

subjecting a compound of Formula (X):




15


wherein

A represents C₆ to C₁₀ aryl which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C₁ to C₆ alkyl and (iii) unsubstituted C₆ to C₁₀ aryl;

to an oxidation reaction in the presence of a catalytic amount of a stable organic nitroxyl radical, to form a compound of formula (IX):




(2) subjecting a compound of Formula (IX) as defined above to reaction with a
5 compound having the formula:




wherein B is as defined above and each R'' is the same or different and each
represents a C₁ to C₆ alkyl group,


the process being carried out in the presence of lithium chloride and an organic
10 base to form the compound of Formula (VIII):




(3) reducing the side chain oxo group of the compound of Formula (VIII) with a
reducing agent to form a compound of Formula (VII):



(4) deprotecting the hydroxyl group of the compound of Formula (VII) to form the corresponding hydroxy-substituted compound of Formula (VI):




5 (5) when the dashed line in the compound of Formula (I-A) or (I-B) is absent: hydrogenating the double bond of a compound of Formula (VI), to form a compound of Formula (VI-A):



10 (6) subjecting the compound of Formula (VI) or (VI-A) to reaction with a silylating agent of formula




wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl and X represents F , Cl , Br or I , to form a compound of Formula (V):



wherein R' represents:



(7) reducing the lactone oxo group of the compound of Formula (V) to form a compound of Formula (IV):




(8) subjecting the compound of Formula (IV) to a Wittig reaction with an ylide, the ylide being formed from the reaction of a compound of formula:




10 with a strong base, wherein R^w represents C₁ to C₆ alkyl or C₆ to C₁₀ aryl and X represents F, Cl, Br or I,

to form a compound of Formula (IIIa) or (IIIb) or a mixture thereof:



(9) for compounds of Formula (I-B): subjecting a compound of Formula (IIa) or
15 Formula (IIb) or a mixture thereof, to reaction with an alkyl halide of formula $R''\text{-Hal}$,
wherein R'' is as defined above and " Hal " represents chloro, bromo or iodo, in the

presence of DBU [1,8-diaza-bicyclo(5.4.0)undec-7-ene], to form a compound of Formula (IIa) or (IIb):




or a mixture thereof; and


5 (10) removing the protecting groups from: (a) the compound of Formula (IIIa) or (IIIb) or a mixture thereof to form the compound of Formula (I-A), or (b) a compound of Formula (IIa) or (IIb) or a mixture thereof to form the compound of Formula (I-B).

2. A process for the synthesis of a compound of Formula (I-B) as defined in claim 1, wherein the dashed line is absent, comprising carrying out steps (1) and (2) as defined 10 in claim 1, then carrying out the following steps (3'), (4') and (5'):


(3') subjecting the double bond of the compound of Formula (VIII) to a hydrogenation reaction to form a compound of Formula (XIII):



15 (4') reducing the side chain oxo group of the compound of Formula (XIII) to form a compound of Formula (XIV):




(5') deprotecting the hydroxyl group of the compound Formula (XIV) to form a compound of Formula (VI-A):




20 subjecting the above compound of Formula (VI-A) to reaction steps (6), (7), (8), (9) and (10) as defined in claim 1.

3. A process for the synthesis of a compound of Formula (I-A) or (I-B) as defined in claim 1 comprising producing a compound of Formula (IIIa) or Formula (IIIb) or a mixture thereof as defined in either claim 1 or claim 2:




and further carrying out the steps of:

(9a) subjecting the compound of Formula (IIa) or (IIb) or a mixture thereof, to reaction with at least one molar equivalent of silylating agent to form a compound of Formula (XI):




(10a) for compounds of Formula (I-B): subjecting the compound of Formula (XI) to a transesterification reaction with an alcohol having the formula $R''-OH$, wherein R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl optionally in the presence of a weak acid catalyst to form the compound of Formula (XII):




(11a) removing the protecting groups R' from either (a) the compound of Formula (XI) to form the compound (I-A) or (b) the compound of Formula (XII) to form the compound (I-B).

4. A process for the synthesis of a compound of Formula (I-B) as defined in claim 1
20 comprising carrying out steps (1) to (9) as defined in either claim 1 or claim 2 to form a compound of formula (IIa) or (IIb) or a mixture thereof:



and further carrying out the steps of:


(10b) subjecting the compound of Formula (IIa) or (IIb) or a mixture thereof to reaction with at least one molar equivalent of silylating agent to form a compound of Formula (XII):



; and


(11b) removing the protecting groups R'.

5. A process for the synthesis of a compound of Formula (I-B) as defined in claim 1 comprising removing the silyl protecting groups, R', from a compound of Formula (XII), as prepared by the process of either claim 3 or claim 4




6. A process for the synthesis of a compound of Formula (I-A) as defined in claim 1 comprising removing the silyl protecting groups, R', from a compound selected from the group consisting of

(i) a compound of Formula (IIIa) or Formula (IIIb), as prepared by the process of
15 claim 1, or a mixture thereof:





(ii) a compound of Formula (XI), as prepared by the process of claim 3;




7. A process according to any one of claims 1 to 6 wherein the oxidation reaction of
20 step (1) is carried out by electrooxidation.

8. A process according to any one of claims 1 to 6 wherein the oxidation reaction is carried out in the presence of at least one mole of co-oxidant selected from the group consisting of *m*-chloroperbenzoic acid, high valent metal salts, sodium bromite, sodium or calcium hypochlorite, N-chloro-succinimide or hypervalent iodine compounds.
- 5 9. A process according to claim 8 wherein the hypervalent iodine compound is [bis(acetoxy)iodo]benzene.
10. A process according to any one of claims 1 to 9 wherein the nitroxyl radical comprises a completely α -substituted piperidin-1-oxy radical.
11. A process according to claim 10 wherein the α -ubstituted piperidin-1-oxy radical is TEMPO free radical.
12. A process according to any one of claims 1 to 11 wherein the reducing agent for reducing the side chain oxo group is selected from the group consisting of borane-dimethylsulfide complex, lithium tri-*sec*-butylborohydride and sodium borohydride.
13. A process according to claim 12 wherein the reduction of the side chain oxo group is carried out in the presence of a chiral oxazaborolidine catalyst (Corey catalyst) to form a single enantiomer of the compound of Formula (VII).
14. A process according to claim 13 wherein the reduction of the side chain oxo group is carried out in the presence of (R)-tetrahydro-1-methyl-3,3-diphenyl-1H, 3H-pyrrole-[1,2-c][1,3,2]-oxazaborole.
- 20 15. A process according to any one of claims 1 to 14 wherein the hydroxyl deprotection step to produce the compound of formula (VI) or (VI-A) is carried out in the presence of a base selected from the group consisting of K_2CO_3 , Na_2CO_3 and Li_2CO_3 .
- 25 16. A process according to any one of claims 1 to 15 wherein the hydrogenation reaction is carried out by catalytic hydrogenation in the presence of a catalyst comprising palladium, platinum or rhodium optionally in the presence of sodium nitrite.


17. A process according to any one of claims 1 to 16 wherein the silylating agent is selected from the group consisting of trimethylsilyl chloride, triethylsilyl chloride and *tert*-butyldimethylsilyl chloride.
18. A process according to any one of claims 1 to 17 wherein the reducing agent in step (7) is di-*iso*-butylaluminium hydride.
19. A process according to any one of claims 1 to 18 wherein the removal of the protecting group R' is carried out using a weak acid selected from the group consisting of acetic acid, citric acid or pyridinium p-toluene sulfonate.
20. A process according to any one of claims 1 to 19 wherein A represents phenyl.
- 10 21. A process according to any one of claims 1 to 20 wherein R^w represents phenyl and X represents Br.
22. A process according to any one of claims 1 to 21 wherein the group R" represents isopropyl.
- 15 23. A process according to any one of claims 1 to 22 wherein B is selected from the group consisting of (i) C₁ to C₆ alkyl, (ii) C₇ to C₁₆ aralkyl, wherein the aryl group is unsubstituted and (iii) -(CH₂)_nOR^a, wherein n represents 1, 2 or 3 and R^a represents a C₆ to C₁₀ aryl group which is substituted with a substituent selected from halo or CF₃.
24. A process according to claim 23 wherein B represents a substituent selected from the group consisting of:
20 
25. A process according to any one of claims 1 to 24 wherein the wavy line in side chain of compounds of any of Formulae (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb), (XI), (XII) or (XIV) represents a single enantiomer.
26. A process according to any one of claims 1 to 25 for the synthesis of a compound selected from the group consisting of:



27. A process according to any one of claims 1 to 5 for the synthesis of latanoprost.



28. A process for the synthesis of a compound of Formula (IX):




5

wherein

A represents C_6 to C_{10} aryl which may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of (i) halo, (ii) C_1 to C_6 alkyl and (iii) unsubstituted C_6 to C_{10} aryl;


10 comprising subjecting a compound of Formula (X):



to an oxidation reaction in the presence of a catalytic amount of a stable organic nitroxyl radical.

29. A process for the synthesis of a compound of Formula (XI):

5



wherein the groups R' are the same and each represents the substituent:




wherein R^x, R^y and R^z are the same or different and each independently represents C₁ to C₆ alkyl, C₆ to C₁₀ aryl or C₇ to C₁₆ aralkyl;

10 comprising reacting a compound of Formula (IIIa) or Formula (IIIb), as defined in claim 1, or a mixture thereof, to reaction with at least a molar equivalent of silylating agent having the formula:



wherein R^x, R^y and R^z are as defined above and X represents F, Cl, Br or I.

15 30. A process for the synthesis of a compound of Formula (XII):



wherein

each R' is the same and each represents the substituent:

- 95 -



wherein

R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl; and


5 R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl;

comprising subjecting a compound of Formula (IIa) or Formula (IIb), as defined in claim 1, or a mixture thereof, to reaction with at least one molar equivalent of silylating agent having the formula:



10 wherein R^x , R^y and R^z are as defined above and the group X represents F, Cl, Br or I.

31. A process for the synthesis of a compound of Formula (XII):



wherein:


15 each R' is preferably the same and each represents the substituent:



wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl; and

R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl;

20 the process comprising subjecting a compound of Formula (XI), as defined in claim 3:




wherein the groups R' are the same and each represents the substituent



wherein R^x , R^y and R^z are the same or different and each independently represents C_1 to C_6 alkyl, C_6 to C_{10} aryl or C_7 to C_{16} aralkyl;


to a transesterification reaction with an alcohol having the formula $\text{R}''\text{-OH}$,
5 wherein R'' represents C_1 to C_6 alkyl or C_3 to C_8 cycloalkyl.

32. A compound of Formula (XI)




or a single enantiomer thereof, wherein B and R' are as defined in claim 1.

33. A compound of Formula (XII)




10 or a single enantiomer thereof, wherein B and R'' are as defined in claim 1.

34. A compound according to either claim 32 or claim 33 wherein B represents




35. A compound according to claim 34 having the structure:




15

36. A compound according to claim 34 having the structure:




37. A compound of Formula (VIII):




wherein A represents unsubstituted C₆ to C₁₀ aryl; and

B represents a substituent selected from the group consisting of:




38. A compound of Formula (VII):




or a single isomer thereof, wherein

A represents unsubstituted C₆ to C₁₀ aryl; and

B represents a substituent selected from the group consisting of:




39. A compound of Formula (VI):




or a single isomer thereof, wherein

B represents a substituent selected from the group consisting of:




40. A compound of Formula (VI-A):




or a single isomer thereof, wherein

B represents a substituent selected from the group consisting of:




41. A compound of Formula (V):




or a single isomer thereof, wherein


R' represents:



B represents a substituent selected from the group consisting of:




42. A compound of Formula (IV):




15 or a single isomer thereof, wherein

R' represents:




B represents a substituent selected from the group consisting of:




43. A compound of Formula (IIIa) or (IIIb):

5




or a single isomer thereof, wherein

R' represents:




10

B represents a substituent selected from the group consisting of:




44. A compound of Formula (IIa) or (IIb):

15




or a single isomer thereof, wherein


R' represents:



B represents a substituent selected from the group consisting of:



45. A compound of formula (XIV):



or a single isomer thereof, wherein

A represents unsubstituted C₆ to C₁₀ aryl; and

5 B represents C₇ to C₁₆ aralkyl, wherein the aryl group may be unsubstituted or substituted with one to three substituents independently selected from the group consisting of C₁ to C₆ alkyl, halo and CF₃.

46. A compound according to claim 45 wherein B represents -(CH₂)₂-phenyl.

47. A compound according to claim 45 or 46 wherein A represents phenyl.

10 48. A process for the purification of latanoprost by HPLC comprising the use as an eluent, of a mixture comprising one or more hydrocarbons, one or more alcohols and, optionally, acetonitrile.

49. A process according to claim 48 wherein the eluent comprises at least one hydrocarbon, at least one alcohol and acetonitrile.

15 50. A process according to claim 49 wherein the eluent comprises a hydrocarbon or a mixture of hydrocarbons in an amount range of 80 to 99 volume percent and an alcohol or a mixture of alcohols in an amount range of 1 to 20 volume percent.

51. A process according to claim 50 wherein the eluent comprises a hydrocarbon or a mixture of hydrocarbons in an amount range of 88 to 98 volume percent and an alcohol

20 or a mixture of alcohols in an amount range of 2 to 12 volume percent.

52. A process according to claim 51 wherein the eluent comprises a hydrocarbon or a mixture of hydrocarbons in an amount range of 85 to 99 volume percent, an alcohol or a mixture of alcohols in an amount of 0.5 to 10 volume percent and acetonitrile in an amount of 0.5 to 5 volume percent.

53. A process according to any one of claims 48 to 52 wherein the hydrocarbon is hexane or heptane.
54. A process according to any one of claims 48 to 53 wherein the alcohol component comprises at least one C₁ to C₈ straight chain, branched or cyclic alkanol.
55. A process according to any one of claims 48 to 54 wherein the alcohol comprises propan-2-ol.
56. Latanoprost substantially free of the 15(S)-*cis* isomer, the 15(S)-*trans* isomer and the 15(R)-*trans* isomer.
57. Latanoprost according to claim 56 containing less than 0.3% in total of any combination of: 15(S)-*cis* isomer, 15(S)-*trans* isomer and 15(R)-*trans* isomer.
58. Latanoprost according to claim 57 containing less than 0.1% each of 15(S)-*cis* isomer, 15(S)-*trans* isomer and 15(R)-*trans* isomer.
59. Latanoprost having a purity of greater than 98%.
60. Latanoprost according to claim 59 having a purity of greater than 99%.
61. Latanoprost according to claim 60 having a purity of greater than 99.5%.
62. Latanoprost according to claim 61 having a purity of greater than 99.8%.
63. A process for the synthesis of a compound of Formula (I-A), (I-B), (IX) or (XII), substantially as herein described with reference to any one or more of the examples but excluding comparative examples.
64. A compound of Formula (XI), (XII), (VIII), (VII), (VI), (VI-A), (V), (IV), (IIIa), (IIIb), (IIa), (IIb) or (XIV), substantially as herein described with reference to any one or more of the examples but excluding comparative examples.
65. A process for purification of latanoprost, substantially as herein described with reference to any one or more of the examples but excluding comparative examples.

66. Latanoprost substantially free to the 15(S)-*cis* isomer, the 15(S)-*trans* isomer and the 15(R)-*trans* isomer and/or having a purity of greater than 98%, substantially as herein described with reference to any one or more of the examples but excluding comparative examples.

2002321396 08 Oct 2007