

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
2 June 2005 (02.06.2005)

PCT

(10) International Publication Number
WO 2005/048810 A2

- (51) International Patent Classification⁷: **A61B**
- (21) International Application Number:
PCT/US2004/025304
- (22) International Filing Date: 30 July 2004 (30.07.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/491,582 31 July 2003 (31.07.2003) US
- (71) Applicant: **WASHINGTON UNIVERSITY IN ST. LOUIS** [US/US]; One Brookings Drive, St. Louis, MO 63130 (US).
- (72) Inventors: **MACH, Robert, H.**; 540 Orchard Lane, Eureka, MO 63025 (US). **WELCH, Michael, J.**; 312 N. Meramec #200, Clayton, MO 63105 (US). **ROWLAND, Douglas, J., Jr.**; 5024 Kain Dr., Shrewsbury, MO 63119 (US). **TU, Zhude**; 656 Legends View Drive, Eureka, MO 63025 (US).
- (74) Agents: **RASCHE, Patrick, W.** et al.; Armstrong Teasdale LLP, One Metropolitan Square, Suite 2600, St. Louis, MO 63102-2740 (US).
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SIGMA-2 RECEPTOR RADIOTRACERS FOR IMAGING THE PROLIFERATIVE STATUS OF SOLID TUMORS



(57) Abstract: Novel benzamide compounds of Formula (I), Formula (II) and Formula (III), salts, water soluble salts, analogs and radiolabeled counterparts thereof as sigma-2 receptor radiotracers for imaging the proliferative status of solid tumors. A method for diagnosing a mammal for the presence of a mammalian tumor therein comprises administering to the mammal a diagnostic imaging detectable effective amount of a benzamide compound having a structure illustrated in Formula (I), Formula (III) and Formula (III) and detecting binding of the compound to a tumor in the mammal. A method for diagnostic imaging of a mammalian tissue having cell surface sigma-2 receptors comprising administering to a mammal a diagnostic imaging amount of a compound having a structure illustrated in Formula (I) Formula (II) and Formula (III) and detecting an image of a tissue having an ample cells with sigma-2 receptors.

SIGMA-2 RECEPTOR RADIOTRACERS FOR IMAGING THE PROLIFERATIVE STATUS OF SOLID TUMORS

This application claims the benefit of U.S. Provisional Patent Application, Serial No. 60/491,582, filed July 31, 2003 which is incorporated herein in its entirety by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0001] This research was funded by grants DA 12647, CA86307 and CA102869 awarded by the National Institutes of Health and grant DAMD17-01-1-0446 awarded by the Department of Defense Breast Cancer Research Program of the US Army Medical Research and Material Command Office. The Government has certain rights in the invention.

FIELD OF THE INVENTION

[0002] This discovery relates to benzamide compositions comprising benzamide compounds, salts and analogs. This invention also relates to the use of benzamide compositions and benzamide compounds to prepare radiolabeled benzamide and to its uses in medicine and as a research tool.

BACKGROUND OF THE INVENTION

[0003] Cancer (malignant neoplasm) is the number two killer of people in the U.S. Each year in the U.S. more than a million people are diagnosed with cancer and half of those will ultimately die from the disease. Cancer also afflicts many other living mammals.

[0004] Cancer occurs when normal living mammalian cells undergo neoplastic (malignant) transformation. Cancer is tenacious in its ability to uncontrollably metastasize throughout the mammalian body thus giving rise to a high mortality rate in many situations, particularly breast cancer.

[0005] Breast cancer is characterized by a high proliferative potential that can vary considerably from patient to patient. No vaccine or other universally successful method for the prevention or treatment of breast cancer is currently available. The rate of tumor cell proliferation has been shown in breast tumors to predict the response to radiation therapy and chemotherapy. Presently, measures of tumor cell proliferation are obtained by histological or flow-cytometric analysis. Both methods are limited by sampling procedures and unfortunately only about 60% to about 70% of patient samples are suitable for flow cytometric analysis.

[0006] It has been demonstrated that sigma-2 (σ_2) receptors are expressed in high density in a number of human and rodent breast cancer cell lines (Cancer Research, 55, 408 (1995)). However, their expression in such cell lines is heterogenous, and their function is unknown.

[0007] Sigma (σ) receptors have also been identified as a distinct class of receptors that are expressed in liver, kidneys, endocrine glands, and in the central nervous system. Apart from the normal expression of sigma receptors in these tissues, several studies have reported their over expression in human and murine tumors (1-3). It has also been shown that there are two types of this receptor, σ_1 and σ_2 receptors. The σ_2 receptor has been demonstrated to be a reliable biomarker for the proliferative status of solid tumors (2-4). Up regulation of σ_2 receptors during proliferation was shown by selectively recruiting cells into quiescent and proliferative states and then measuring the receptor concentration (2-4). It was found that σ_2 receptor concentrations increased tenfold when cells were recruited to proliferative states. Therefore, radioligands are desired that have both high affinity and high selectivity for σ_2 receptors as tracers for the non-invasive assessment of the proliferative status of human solid tumors using noninvasive diagnostic imaging procedures such as PET and SPECT. To this end, a high affinity and highly selective σ_2 radioligand is needed for the assessment of tumor status.

[0008] One of the major problems in the clinical management of breast cancer is the early detection and identification of an appropriate treatment strategy. A complication that has limited successful treatment is the inability to assess

the proliferative status of breast tumors since breast cancer has a malignant potential that can vary considerably from patient to patient. The use of surrogate markers of proliferation such as the presence or absence of tumors in axillary lymph nodes suffers from a low sensitivity and specificity. Other methods such as determining the S-phase fraction of tumor biopsy or fine needle aspirates suffers from sampling problems associated with tumor heterogeneity that may not provide a true representation of the proliferative status of a solid tumor.

[0009] A recent strategy has focused on using noninvasive imaging procedures such as Positron Emission Tomography (PET) in order to assess the proliferative status of an entire tumor. This approach has relied primarily on the development of agents that target the increase in metabolic activity (i.e., [^{18}F]FDG) or increased DNA (DNA precursors such as [^{11}C]thymidine) or protein (i.e., [^{11}C]methionine) synthesis associated with tumor proliferation. However, the majority of these agents have proven to be inadequate for providing an accurate measure of the proliferative status of solid tumors for a variety of reasons.


[0010] Cancer cure rates have increased dramatically over the years. This trend as a result of the widespread use of improved screening procedures that often lead to the early diagnosis/detection of cancer. However, as more selective treatment strategies have been developed, it is necessary to develop new and improved diagnostic procedures that can be used earlier to determine a potential treatment strategy based on the biological properties and proliferation of the tumor. In addition, it is desired to develop and have available non-invasive procedures that can provide the means for determining either a positive or negative response to a treatment strategy as early as possible thus extending the mammalian host's viability.

[0011] Additionally a continuing need exists for enhanced non-invasive methods that can accurately assess the proliferative status of breast cancer, as such methods could have a significant positive impact on determining an optimal therapy for treating human breast cancer patients.


BRIEF DESCRIPTION OF THE INVENTION

[0012] In an aspect, the discovery comprises a benzamide composition comprising benzamide compound(s), salts, analogs and radiolabeled counterparts thereof, having at least one structure or substantially similar structure of at least one of the structures illustratively depicted hereinafter in one or more of Formula (I), Formula (II) and Formula (III), analogs and salts thereof and their respective radioactive labeled counterparts of such benzamide compounds, salts and their analogs having use as a radioimaging agent(s).


[0013] In an aspect benzamide compositions including benzamide compounds include those compounds having a structure shown in the structure in Formula (I):



[0014] where Ar symbolically represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H (hydrogen), halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ each symbolically represent chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of




wherein B and C each symbolically represent independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and compound 13 and compound 14.


[0015] In an aspect, compounds 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 and 23 of Formula (I) and compound 13 of Formula (II) and compound 14 of Formula (III) are depicted respectively structurally as:




[0016] Compound 1




[0017] Compound 2




[0018] Compound 3




[0019] Compound 4




[0020] Compound 5




[0021] Compound 6




[0022] Compound 7




[0023] Compound 8




[0024] Compound 9



[0025] Compound 10




[0026] Compound 11




Including compound 23 and

[0027] Compound 12




[0028] Compound 13




[0029] Compound 14

and (the) respective radiolabeled counterparts of compounds 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12, 13, 14 and 23.


In an aspect, a benzamide composition comprises a benzamide compound further comprises compound 6 labeled with at least one of ^{76}Br , ^{123}I , ^{124}I and ^{125}I . In an aspect, the compound is compound 10 labeled with (^{11}C).

[0030] In an aspect, the novel compound 6 comprises ^{76}Br .


[0031] In an aspect, the novel compound 6i comprises ^{123}I .




[0032] In an aspect, the novel compound 6i comprises ^{124}I .



[0033] In an aspect, the novel compound 6i comprises ^{125}I .





[0034] In an aspect, the novel compound 10 comprises ^{11}C .



[0035] In an aspect, a process for the preparation of a compound selected from one of compounds 1, 2, 3, 4, 5, 6 depicted structurally in Figure 4 comprises respectively reacting a compound


- (a) having a formula wherein R is hydrogen or methoxy



with a compound of the formula

wherein

R in this structural depiction is a substitutable moiety selected from hydrogen or methoxy and independently m is an integer from one to 3 with bromoacetonitrile or bromobutyronitrile to produce an N-alkylated product




wherein when R is hydrogen, n is an integer independently 2-4 and wherein R is independently methoxy then n is an integer 2-4 and

- (b) reducing the N-alkylated product with lithium aluminum hydride in THF or hydrogenating the N-alkylated compound over palladium on charcoal to provide an intermediate amine product depicted structurally as compound 19, 20, 21 and 22 in Figure 4, and

- (c) condensing the intermediate amine product with either 2-methoxy-5-bromonaphthol chloride or 5-bromo-2,3-dimethoxybenzoic acid to

produce a compound having as its structure a structure selected from one structures shown for compounds 1-6.


[0036] In an aspect, a process for the preparation of a compound 7, 8, 9 and 10 structurally depicted Figure 5 comprises reacting a compound of the formula




[0037] wherein X is a substitutable moiety independently halogen or alkyl C₁-C₈ with thionyl chloride and one of butyronitrile or bromobutyronitrile to produce compound of 7, 8, 9 and 10 of Fig 5. In an aspect X comprises bromide. In an aspect X comprises methyl.

[0038] In an aspect, a process for the preparation of a compound having a structure depicted as 11 and/or 12 comprises:


(a) reacting a compound having a structure of the formula



(b) with bromoacetonitrile to produce a compound having the structure shown in the formula




- (b) hydrogenating that compound over palladium on charcoal to prepare a compound of the structural formula



- (c) and condensing that compound with either of 2-methoxy-5-bromonaphthoyl chloride or 5-bromo-2,3-dimethoxybenzoic acid respectively to produce compound (11) and compound (12) respectively.

[0039] In an aspect, a process for the preparation of a compound having a structure depicted as compound 23, which comprises reacting the corresponding phenol with 1-bromo-2-fluoroethane.



[0040]

Compound 23

[0041] Compound 23 is depicted in Formula I. Compound 23 is useful in all aspects diagnostic, treatment, medicinal and research tool use aspects of this discovery.

[0042] In an aspect, a non-invasive method for diagnosing a mammal for the presence of a cancer comprises administering to the mammal a diagnostic imaging detectable effective amount of a benzamide composition comprising a benzamide detectably radio labeled compound having a structure selected from at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and detecting binding of the at least one benzamide compound in the mammal. In an aspect the method comprises determining that a mammalian tumor is present in the mammal upon detecting binding, thus diagnosing the mammal. In an aspect the method further comprises producing an acquisition of the detection.

[0043]—In an aspect, a marker for cancer comprises a detectably-labeled benzamide composition comprising a benzamide compound having a structure of at least one of the structures illustratively depicted of benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and having an explicit provocative binding efficacy to a tumor in a living mammal. In an aspect the detectably labeled benzamide compound is a highly selective σ_2 radioligand having the aforementioned structure effectively and functionally appended with a radioactive ligand.

[0044]—In an aspect, a novel pharmaceutical composition comprises a benzamide composition comprising a benzamide compound having as a structure at least one of structures shown for novel benzamide compounds illustratively depicted in Formula (I) compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and a pharmaceutically (pharmacologically) acceptable diluent or carrier. In an aspect a pharmaceutical composition comprises a detectably labeled benzamide compound having a structure illustratively depicted in one of the compounds 1-12 and 23 illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and a pharmaceutically acceptable diluent or carrier. In an aspect the detectably labeled benzamide compound is a highly selective σ_2 radioligand having the aforementioned structure appended with a radioactive ligand.

[0045]—In an aspect, a pharmaceutical composition effective for treating human or non-human neoplastic disorder comprises a detectably labeled pharmaceutically effective amount of at least one compound having a structure of at least one of the structures illustratively depicted in one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) in a composition including a pharmaceutically acceptable carrier. In an aspect the detectably labeled benzamide compound is a highly selective σ_2 radioligand having the aforementioned benzamide structure appended with a radioactive ligand.

[0046] In an aspect, a non-invasive method to determine the proliferative status of a cancer cell in a living mammal comprises administering to a living mammal afflicted with a solid malignant tumor, an effective amount of a detectably labeled benzamide composition comprising a benzamide compound having a structure of at least one of the benzamide compounds 12 illustratively depicted in Formula (I) compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and determining the extent to which the detectably labeled benzamide compound binds to cells of a tumor in the mammal, the extent providing a measure of the proliferative status of the cancer cells. In an aspect the living mammal is a human. In an aspect, determining the proliferative status includes assessing the proliferative status of a breast cancerous tumor.

[0047] In an aspect, a method for pharmacologically treating a mammalian tumor as a disorder in a mammal comprises administering to a mammal having a tumor a composition including a tumor-inhibiting amount of at least one detectably-labeled benzamide compound having a structure of the structure shown of at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) In an aspect the mammal is a living human.

[0048] In an aspect, a non-invasive method for diagnostic imaging of a mammalian tissue having ample cell surface sigma-2 receptors comprises administering to the tissue of the mammal a diagnostic imaging amount of at least one

compound having a structure of at least one of the detectably labeled benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and detecting an image of a tissue having an ample cells with sigma-2 receptors. In an aspect the mammal is a human. In an aspect, the image is used to diagnose living mammalian tissue.

[0049] In an aspect, a non-invasive method for in vitro detection of a cancer cell in living mammalian tissue sample comprises contacting a mammalian tissue sample comprising a cell with an in vitro diagnostic imaging amount of at least one detectably radiolabeled benzamide composition comprising a benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) for a time and under conditions sufficient and effective for binding of the compound to the cell and detecting such binding indicative of an association with the present of cancer. In an aspect the mammal is a living human. In an aspect the detecting is by image acquisition.

[0050] In an aspect, the cell(s) is in a previously obtained representative biological sample from a mammal. In an aspect, the novel compounds herein, including the radiolabeled counterparts are useful in the treatment of cancer.

[0051] In an aspect, a non-invasive method for in vitro detection of a cancer cell in a living mammalian tissue sample comprises contacting a mammalian tissue sample with an in vitro diagnostic imaging detectable and acquisitionable amount of at least one detectably radiolabeled benzamide composition/compound having as a structure a structure of at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) for a time and under cellular conditions functionally sufficient and effective for binding of the compound to the cancer cell and detecting such binding. In an aspect such binding is indicative of the presence of a cancer cell. In an aspect the mammal is a human.

[0052] In an aspect, a method for determining proliferation and./or progression of a cancer as a disorder in a living mammal comprises administering to a

living mammal a diagnostic imaging detectable amount of at least one detectably radiolabeled benzamide composition comprising a benzamide compound having as a structure a structure of at least one of the benzamide compounds illustratively structurally depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first selected time, detecting an image of a tissue having ample cells with sigma-2 receptors at a second selected (later) time respectively detecting an image of a tissue having ample cells with sigma-2 receptors at both times, comparing the images and determining if the detected image at the later time is smaller than the detected image at the first time. In an aspect the elapsed time between the first time and second time is selected to be a time duration significant amount. In an aspect the mammal is a living human. In an aspect the comparison is used to determine proliferation and/or progression of a cancer in a mammal.

[0053] In an aspect, a non-invasive method for identifying a modulating effect (and regression effect) of a cancer in a living mammal with a disorder, comprises administering to the mammal a diagnostic imaging detectable amount of at least one detectably radiolabeled benzamide composition comprising a benzamide compound and having a structure of at least one of the structures illustrated in Formula (I), Formula (II) and Formula (III) at a first time, detecting and acquisitioning an image of a tissue having ample available cells with sigma-2 receptors, administering a detectably-labeled benzamide compound having a structure of at least one of the compound structures illustrated in Formula (I) including compound 23, Formula (II) and Formula (III) to the mammal and at a second (later) time respectively detecting and acquisitioning an image of a tissue having an abundance of cells with sigma receptors, comparing the respective images and determining that there has been an prophylactic effect and/or regression and/or modulation. In an aspect the comparison shows the amount of regression over time. In an aspect the mammal is a living human. In an aspect the comparison shows the prophylactic effect of the compound and its toxicity to cancer. In an aspect the comparison shows the efficacy of the compound towards killing cancer in a living mammal. In an aspect the detectably-labeled benzamide compound is tagged with a radioactive ligand.

[0054] In an aspect, a non-invasive method for diagnosing and determining the response of a mammalian patient(s) with a disorder to tailored drug therapy comprises administering to a mammal a diagnostic imaging detectable amount of at least one detectably labeled benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I) including compound 23, compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first and second (later) times respectively, detecting an image of a tissue having an abundance of cells with sigma receptors at both times, comparing the images and determining/diagnosing if the image at a later time is larger than the image at the first time that there has been a proliferative effect or progression of cancer. In an aspect the dynamic comparison shows the proliferation and progression of the cancer over elapsed time. In an aspect the mammal is a living human. In an aspect a method for diagnosis and determining the response is a determination of a prophylaxis or management of a disorder associated with neoplastic cells.

[0055] In an aspect, a non-invasive method of screening candidate chemicals for toxicity/lethality to cancer comprises administering to a mammal a diagnostic imaging detectable amount of at least one detectably labeled benzamide compound selected from the benzamide compounds illustratively depicted in Formula (I) including compound 23, compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first or initial time, detecting and acquisitioning an image of a tissue having ample cells with sigma-2 receptors, administering to the mammal a candidate chemical, detecting and acquisitioning an image of tissue having ample cells with sigma-2 receptors at a secure time subsequent the first time, comparing the detected images and making a determination as to whether there has been a proliferative effect and progression of the cancer. In an aspect the mammal is a human. In an aspect the chemicals are organic or inorganic.

[0056] In another aspect, a non-invasive medical treatment for a mammal and a method of medically treating a mammal comprises administering to a mammal a diagnostic imaging detectable amount of at least one detectably labeled benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I) including compound 23, compound 13 depicted

in Formula (II) and compound 14 depicted in Formula (III) at a first and second (later) time respectively, detecting an image of a tissue having ample cells with sigma receptors at a first time and a second time, the second time subsequent the first time comparing the images taken respectively and determining if the image at the later time is larger than the image at the first time that there has been a proliferative effect and progression of the cancer. In an aspect the mammal is a human. In an aspect the treatment is a cancer treatment. In an aspect medical treatment includes a method of retarding, preventing, and ameliorating disease or a medical affliction in a mammal.

[0057] In an aspect, a non-invasive method of customizing drug therapy for a living mammalian subject comprises effectively and capably administering to the subject, the mammal undergoing drug therapy, a diagnostic imaging detectable amount of at least one radiolabeled benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I) including compound 23, compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III), detecting an image of a tumor having ample capably radionuclide detectable sigma-2 receptors and making a determination regarding a drug therapy based on the captured images. In an aspect, the determination comprises an adjuvant therapy to the drug therapy.

BRIEF DESCRIPTION OF THE FIGURES

[0058] Figure 1 illustratively shows structured depictions of novel conformationally flexible benzamide compounds 1, 2, 3, 4, 5 and 6 and compound 23 of Formula (I).

[0059] Figure 2 illustratively shows structured depictions of novel conformationally flexible benzamide compounds 7, 8, 9 and 10 of Formula (I).

[0060] Figure 3 illustratively shows structured depictions of novel conformationally flexible benzamide compounds 11 and 12 of Formula (I).

[0061] Figure 4 shows an illustrative process for preparing novel conformationally flexible benzamide compounds (1, 2, 3, 4, 5 and 6) in 1 of Formula (I).

[0062] Figure 5 shows an illustrative process for preparing novel conformationally flexible benzamide compounds (7, 8, 9, and 10) of Formula (I).

[0063] Figure 6 shows an illustrative process for preparing novel conformationally flexible benzamide compounds (11 and 12) of Formula (I).

[0064] Figure 7 shows structured depictions of conformationally flexible benzamide compound (13) of Formula (II) and conformationally flexible benzamide compound (14) of Formula (III) respectively labeled with a metal chelating group comprising an emitting radionuclide "(M)".

[0065] Table IA shows structures and in vitro binding data.

[0066] Figure 8 shows in vivo blocking studies of [^{125}I]6i (left), [^{76}Br]6 (center), and [^{11}C]10 (right) in tumor-bearing mice. The blocking agent was YUN 143, which has a high affinity for σ_1 and σ_2 receptors. Animals were sacrificed at the time point displaying the highest %I.D./gram tumor. The data are consistent with the labeling of σ_2 receptors in vivo by each radiotracer.

[0067] Figure 9 shows a comparison of [^{18}F]FLT ((18)F-3'-fluoro-3'-deoxy-L-thymidine PET (FLT PET)) and [^{11}C]10. Although [^{18}F]FLT has a high uptake in tumors (top graph), the high uptake of radioactivity in normal tissues results in a lower tumor:background ratio of [^{18}F]FLT relative to [^{11}C]10, particularly the tumor:muscle, tumor:blood, and tumor:heart ratios (bottom graph).

[0068] Figure 10 shows a comparison of [^{18}F]FLT with [^{76}Br]6 at 2 hrs post-i.v. injection. Notice the higher tumor:background ratios of [^{76}Br]6 versus that of [^{18}F]FLT.

[0069] Figure 11 shows a comparison of [^{18}F]FLT with [^{125}I]6i at 2 hrs post-i.v. injection. Notice the higher tumor:blood, tumor:muscle, and tumor:heart ratios of [^{125}I]6i versus that of [^{18}F]FLT.

[0070] Figure 12 shows MicroPET[®] imaging studies of [^{76}Br]6 in Balb-c mice bearing an EMT-6 breast tumor xenograft (EMT-6 BALB-C mice). The

no-carrier-added (NCA) study is on the left and the sigma receptor blocking study (1 mg/kg, i.v. YUN-143) is shown on the right.

[0071] Figure 13 shows graphs comparing the uptake of the benzamide analogs with % uptake in breast tumors at 5 min (top) and the tumor:fat ratio at the optimal imaging time (bottom). There was no correlation between tumor uptake or tumor:fat ratio and log P. Note that all compounds had a high tumor uptake at 5 min post-injection. The high tumor:fat ratio of [⁷⁶Br]6 and [¹²⁵I]6i was due to the later time point after injection used because of the longer half-life of Br-76 and I-125 versus that of C-11.

DETAILED DESCRIPTION OF THE INVENTION

[0072] The present discovery is understood more readily by reference to the following detailed description of the discovery and the Examples included therein.

[0073] As used herein the phrase “benzamide compound” illustratively depicted in Formula (I) including compound 23, compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III)” includes any moiety derived from or of the benzamide compound in any form including pharmaceutically effective and acceptable water soluble salt form, radical, analogs, ionic form, ion, conformational form, radiolabeled conformational form, radiolabeled ion and mammalian and metabolic derivatives thereof. Acceptable salts include but are not limited to tosylate, methanesulfonate, acetate, citrate, malonate, tartrate, succinate, benzoate, ascorbate, alpha-ketoglutarate, and alpha-glycerophosphate. Suitable inorganic salts include hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

[0074] This discovery relates to benzamide compounds having a specific affinity for sigma-2 receptors believed indicative of the presence of carcinoma(s) and melanoma(s) in a living mammal such as a human. Further this discovery relates to benzamide compositions as sigma-2 receptor radiotracers for imaging the proliferative status of solid tumors.

[0075] The invention relates to benzamide compounds and to their analogs and salts and to the use of benzamide benzamide compounds, salts and analogs thereof to prepare detectably radiolabeled benzamide compounds, including salts and analogs thereof which are useful in biological competitive assays to test for and detect the presence of sigma-2 receptors associated with cancer or pre-cancerous conditions. The discovery further relates to the use of benzamide compound(s) as radiolabeled benzamides including salts and analogs thereof which are useful in medicinal and diagnostic chemistry for detecting cancer in living mammals.

[0076] In an aspect, a benzamide compound is provided to a living mammal such as to a human by effective administration of an aqueous composition (such as a saline composition) comprising a salt of benzamide associated with an emitting radionuclide or a salt of an analog of benzamide with an associated (labeled) emitting radionuclide to the living mammal such as a human. In an aspect the salt is a water soluble salt. An adjuvant may be employed if desired in aqueous compositions of this invention.

[0077] As used herein, the term "radiolabeled counterparts" includes respective radiolabeled benzamide compounds which are also novel.

[0078] As used herein, the term "tumor:background ratio" (i.e., tumor:fat ratio, tumor:muscle ratio, tumor:lung ratio, tumor:blood ratio) refers to the % injected dose/g tissue of the radiotracer in the tumor divided by the % injected dose/g tumor in the background tissue (i.e., fat, muscle, lung, blood).

[0079] As used herein, the term "detectably labeled" includes the respective pharmacologically acceptable radiolabeled benzamide compounds having an effective amount of an emitting radiolabel therewith and suitably accepting an emitting radiolabel for use in administration to living mammals.

[0080] As used herein, the term "⁷⁶Br Compound 6" means a compound having a structure of that structure depicted in structure 6 radiolabeled with ⁷⁶Br similarly schematically for ¹²³I, ¹²⁴I, ¹²⁵I, and for ¹¹C Compound 10.

[0081] In an aspect the term, "C₁-C₄" comprises at least one of methyl, ethyl, propyl, isopropyl, butyl, isobutyl and isobutyl.

[0082] In an aspect the term, "alkoxy C₁-C₄" comprises at least one of methoxy, ethoxy, propoxy and butoxy.

[0083] As used herein the term "fluoroalkyl" includes fluoro C₁ to C₄" and includes fluormethyl, fluoroethyl, fluoropropyl and fluorobutyl.

[0084] As used herein the term "fluoroalkoxy" include fluoro C₁ to C₄ alkoxy and includes fluoromethoxy, fluoroethoxy, fluoropropoxy and fluorobutoxy.

[0085] As used herein, the term "salt" includes water soluble salt and includes alkali and alkaline earth metal salts including but not limited to sodium, potassium, magnesium and calcium.

[0086] As used herein, the term "alkyl" and "alkoxy" include substituted alkyl and substituted alkoxy.

[0087] As used herein, the symbol "I" means iodine, the symbol "Br" means bromine, the symbol "Cl" means chlorine, the symbol "F" means fluorine and the symbol "H" means hydrogen.

[0088] As used herein, the term "provocatively" means aggressively and actively and having a sufficient and effective reaction capability.

[0089] As used herein, the term "medicinal chemistry" means a chemistry-based medicinal practice involving one or more of the biological, medical and pharmaceutical sciences.

[0090] As used herein, the term "administration" includes the giving of a compound by any useful effective means to a living mammal and its successful introduction into the mammal such as in its gastrointestinal tract in an effective method which results in that compound, its salt, its ions, metabolites, analogs, radionuclides or derivatives being made biologically available to that mammal

receiving administration of benzamide for medicinal use. In an aspect the mammal is a living human. In an aspect the living mammal is a nonhuman such as a canine or feline. In an aspect the benzamide compound is made biologically available to the gastro intestinal tract of the mammal patient.

[0091] As used herein, the expression "pharmaceutically or pharmacologically acceptable" includes a benzamide composition comprising a benzamide compound and its radiolabeled counterpart which contains composition ingredients that are compatible with other ingredients of the composition as well as physiologically acceptable to the recipient, e.g. a mammal such as a living human, without the resulting production of excessive undesirable and unacceptable physiological effects or a deleterious impact on the mammal being administered the pharmaceutical composition. In an aspect, a composition for use comprises one or more carriers, useful excipients and/or diluents.

[0092] As used herein, the term "dosage" includes that amount of novel benzamide compound which when effectively administered to a living mammal provides an effective amount of biologically available benzamide compound to the living mammal.


[0093] In an aspect, as used herein the term "patient" includes a human subject and a human individual. In an aspect the patient includes a human, and a non-human such as feline, canine, horse and murine.

[0094] Terminology used herein is for the purpose of describing particular aspects only and is not intended to be limiting in any regard.


[0095] It must be noted that, as used in the specification the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise.

[0096] As used in the specification, claims and drawings, for convenience numbers 1-12 refer respectively to various structures of novel benzamide compounds of Formula (I) structurally depicted respectively in Figures 1, 2 and 3., Formula (II) and Formula (III) of Figure 7.


[0097] The invention provides benzamide compounds of Formula (I), Formula (II) Formula (III) and medicinal methods using such compounds administered to patients for detecting and treating cancer including but not limited to cancers such as blastomas, gliomas, pheochromocytomas, melanomas, colon, renal, prostate, lung and breast carcinomas in living mammals.



[0098] where Ar is a substitutable moiety comprising either a substitutable aromatic ring or substitutable heteroaromatic ring; X, Y, Z is a moiety selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer ranging from 2 to about 10; and NRR' is at least one of (a):



[0099] wherein A and A1 are each substitutable moieties selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or









[00101] wherein B, C are each substitutable moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂ and R





and R1 are two conjoined and linked cyclic aromatic organic moieties, n is an integer varying from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently individual multi-substitutable aromatic rings and their respective novel radiolabeled counterparts.




[00102] In an aspect, novel benzamide compounds comprise conformationally flexible benzamide compounds having a structure shown as one of structures 1, 2, 3, 4, 5 and 6 shown in Figure 1; a structure such as a structure shown structures 7, 8, 9 and 10 in Figure 2; a structure such one of structures 11 and 12 in Fig. 3 and structures 13 of Formula (II) and structure 14 of Formula (III) of Figure 7. In an aspect, novel detectably-labeled benzamide compounds include such aforerecited novel benzamide compounds having radiolabels attached.

[00103] In an aspect, compounds 1-12 and 23 of Formula (I) are depicted structurally as:











[00117] Compound 13 of Formula (II)



[00118] and Compound 14 of Formula (III).




[00119] In a chemical aspect compounds 1, 2, 3, 4, 5, 6, and 23 are benzamide analogs. Compounds 7, 8, 9, and 10 are benzamides and compounds 11 and 12 are 2 and 3-dichlorophenylpiperazine benzamide and naphthamide analogs.

[00120] In an aspect the inventive compounds comprise conformationally flexible detectably-labeled benzamide compounds of Formula (I) including compound 23, Formula (II) and Formula (III) which provocatively bind to a cell surface sigma receptor and exhibit exquisite cell specificity and affinity for cancerous cells such as at least one of the aforerecited blastomas, gliomas, pheochromocytomas, melanomas, colon, renal, prostate, lung and breast carcinomas in living mammals and for cells having sigma-2 receptors.


[00121] The invention also encompasses a process for the preparation of benzamide compounds and their radiolabeled counterparts, salts, analogs and pharmaceutical compositions thereof.

[00122] In an aspect a process for the preparation of a benzamide compound selected from compounds 1, 2, 3, 4, 5, 6 and 23 of Formula (I) comprises respectively:


- (c) reacting a compound having a formula wherein R is



hydrogen or methoxy with a compound of the formula



wherein R is an independently substitutable moiety selected from hydrogen or methoxy and m is an integer from one to 3 with bromoacetonitrile or bromobutyronitrile to produce an N-alkylated product




wherein when R is hydrogen, n is integer identity 2-4 and wherein R is methoxy and n is an integer ranging from 2-4 and

- (b) reducing said N-alkylated product with either lithium aluminum hydride in (THF) or hydrogenating said N-alkylated compound over palladium on charcoal to provide an intermediate amine product denoted as compound 19, 20, 21 and 22 on Figure 4, and

- (c) condensing said amine product with either 2-methoxy-5-bromonaphthol chloride or 5-bromo-2,3-dimethoxybenzoic acid to produce a compound selected from one of compounds 1, 2, 3, 4, 5 and 6.


[00123] In an aspect a process for the preparation of compound depicted as 7, 8, 9 and 10 of Formula (I) comprises reacting a compound of the formula




[00124] wherein X is a substitutable halogen or alkyl C₁-C₈ with thionyl chloride and one of butyronitrile or bromobutyronitrile to respectively produce compound 7, 8, 9 and 10 depicted structurally Fig 5. In an aspect X comprises bromide. In an aspect X comprises methyl.

[00125] In an aspect a process for the preparation of a compound(s) denoted as 11 and 12 respectively of Formula (III) comprises:


- (a) reacting a compound having a structure of the formula depicted as



with bromoacetonitrile to produce a compound of the formula



(b) hydrogenating that compound over palladium on charcoal to provide a compound of the formula



[00126] and condensing that compound with either of 2-methoxy-5-bromonaphthoyl chloride or 5-bromo-2,3-dimethoxybenzoic acid to provide compound 11 and compound 12.

[00127] In an aspect reaction conditions are employed which provide for the effective reaction of the reaction processes disclosed herein. In an aspect effective reaction time and reaction conditions are provided so as to allow preparation of the desired intermediates and final products of the reactions.

[00128] More in particular the invention relates to the use of one or more of a detectably-radiolabeled benzamide compound(s) of Formula (I), Formula (II) and Formula (III) and their pharmaceutically acceptable salts for non-invasive diagnostic imaging of living mammalian tissue having cells expressing sigma-2 receptors.

[00129] The present invention provides detectably radiolabeled benzamide compounds of Formula (I) Formula (II) and Formula (III) and their radiolabeled nuclides and their pharmaceutically acceptable salts thereof as agents for diagnostic imaging and for detecting and treating cancer cells in living mammals.

[00130] The present invention also relates to a method of determining the proliferative effect and progression of a cancer in a living mammal.

[00131] A noninvasive novel method using novel benzamide compounds of Formula (I) Formula (II) and Formula (III) and their radiolabeled nuclides is provided to detect cancer cells and to assess the proliferative status of cancer cells which express sigma-2 (sigma.2) receptors, such as cells of solid tumors, in vitro and in vivo.

[00132] In an aspect, Formula (I) Formula (II) and Formula (III) compounds are used to diagnose the presence of cancer in a living mammal or in a biological sample thereof. Formula (I) compounds include the compounds structurally shown of Figures 1, 2 and 3 and their pharmaceutically acceptable salts thereof, their radiolabeled counterparts and their mammalian metabolic derivatives. Formula (II) compounds and Formula (III) compounds include the compounds structurally shown of Figure 7 and their pharmaceutically acceptable salts thereof, radiolabeled counterparts and their mammalian metabolic derivatives.

[00133] In an aspect, a method of detecting cancer comprises administering to a human patient afflicted with a solid tumor, such as breast cancer, a functionally effective amount of a detectably labeled conformationally-flexible benzamide compound having a structure of at least one of the structures illustrated in Formula (I) Formula (II) and Formula (III) and determining the extent to which the compound binds to cells of the cancer, the extent providing a measure of the presence and/or proliferative status of the cells, which status correlates to the extent of sigma-2 receptor expression by said cells. The method is based on the ability of the compounds illustratively depicted in Figures 1, 2, 3, and 7 to selectively bind to sigma-2 (sigma.2) receptors versus sigma.1 receptors.

[00134] In an aspect, a benzamide compound having a structure of one of the structures illustrated in Formula (I), Formula (II) and Formula (III) is radiolabeled as a radioligand. In an aspect the radioligand is an emitting radioligand.

[00135] Typically the label is a fluorescent label or radionuclide (M) such as a radioisotope of halogen (^{125}I , ^{123}I , ^{124}I , ^{76}Br , ^{77}Br , ^{18}F) or (^{11}C). In an aspect a functional emitting effective amount of radiolabeled benzamide compound(s) is administered parenterally, i.e., by intravenous, i.p., intrathecal or enteral administration to a living mammal patient. In an aspect the radiolabeled emits a functional detectable amount of desired radioactivity. In a medical aspect the amount of emitted radioactivity is an amount which imparts a therapeutic benefit to the mammalian patient having cancer. In an aspect a therapeutic benefit is that benefit which is medicinally and therapeutically beneficial to the living mammalian afflicted with cancer. In an aspect a cytotoxic amount is an effective lethal amount of a therapeutic compound which beneficially kills or retards cancer cells.

[00136] In an aspect, unlabeled benzamide compounds having a structure of at least one of the structures depicted in Formula (I), Formula (II) and Formula (III) are used as precursor intermediates to make corresponding radiolabeled novel detectably labeled compounds or as sigma-2 specific ligands which can be used in competitive assays to assay for the presence of sigma-2 receptors as described hereinafter.

[00137] In an aspect a detectably labeled benzamide compound having a structure of at least one of the structures depicted in Formula (I), Formula (II) and Formula (III) is effectively administered to a living mammal or to a biological sample thereof or therefrom and the sample is analyzed and a diagnosis is made or obtained. In an aspect, a biological sample of the mammal comprises a representative sample taken of at least one of blood, vessels, atheroma, liver, and other body tissues as well as biopsies of body organs such as a liver biopsy or a muscle biopsy of a living mammal.

[00138] In an aspect the weight of a biological sample is a minimum of tissue about 1 mg, the amount of cells is about 2,000, and of blood about 2 μl or comparable functionally adequate amounts, quantities or volumes of other biological sample(s). In an aspect, the amount of biological sample is that amount or volume which is sufficient to provide for an analysis.

[00139] As used herein, the term “biological sample” or “biologic sample” includes a sample of a suitable size such as a sample of size and composition suitable to use in the methods disclosed herein.

[00140] Novel processes for preparing novel benzamide compounds illustratively depicted having a structure of at least one of the structures depicted in Formula (I), Formula (II) and Formula (III) are provided in illustrative process preparation schematics in Fig. 4, Fig. 5 and Fig. 6 as further aspects of the invention and are illustrated by these procedures in which the meanings of the generic radicals are as noted above unless otherwise qualified.

[00141] For example, the reaction may conveniently be carried out under conditions similar to those described in Example 1.

[00142] Starting materials employed in the synthetic processes described above are commercially available, are reported in the scientific literature, or can be prepared using methods analogous to those described in the literature. The reaction conditions will be such as to allow a successful carrying out of the desired reaction to the desired degree of completion.

[00143] In instances where the novel benzamide compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compounds as salts may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which produce a mammalian physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartrate, succinate, benzoate, ascorbate, alpha-ketoglutarate, and alpha-glycerophosphate. Suitable inorganic salts include hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

[00144] In an aspect pharmaceutically acceptable salts are prepared using procedures well known in the art such as by reacting a basic compound such as an amine with a suitable acid to prepare a mammalian physiologically acceptable anion. Alkali metal (sodium, potassium or lithium) and alkaline earth metal (calcium, magnesium) salts can be made. In an aspect an oxalate salt is a pharmaceutically

acceptable salt of a compound of Formula (I) Formula (II) and Formula (III). In an aspect an oxalate salt is prepared.

[00145] In an aspect, compounds having a structure illustrated in at least one of Formula (I), Formula (II) and Formula (III) are formulated as pharmaceutical compositions generally as their respective pharmaceutically acceptable salts and effectively administered to a mammalian host.

[00146] In an aspect a pharmaceutical composition is prepared comprising at least one compound having a structure depicted in at least one of the structures depicted in Formula (I), Formula (II) and Formula (III) as a radiolabeled benzamide compound.

[00147] In an aspect a living mammal host is selected from at least one of a human and non human animal such as canine, feline, equestrian, murine including dogs, cats, rabbits, guinea pigs, hamsters, mice, rats, horses, goats, sheep, pigs and cows. In an aspect a veterinarian treats a dog having cancer. In an aspect the mammal host is a patient.

[00148] In an aspect an inventive benzamide compound is systematically successfully administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle such as an inert diluent or an assimilable mammalian edible carrier to the patient. The novel benzamide compound may be enclosed in a gelatin capsule, may be compressed into a tablet, or may be a part of the patient's diet. The compound may be combined with one or more compatible functional excipients and administered as ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, crackers and pills. In an aspect the selected administration is, orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes of effective administration.

[00149] Pharmaceutical compositions may be used such as those formulated using readily available formulation ingredients with one or more benzamide radiolabeled compounds. In an aspect at least one novel benzamide compound having a structure depicted in Figures 1, 2, 3 and 7 is incorporated,

optionally together with other active substances, with one or more conventional pharmacological acceptable carriers, diluents and/or excipients, to produce a conventional preparation(s) such as tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, ointments, gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

[00150] In an aspect, such compositions and preparations contain at least a pharmaceutically effective amount of the benzamide compound generally at least about 0.1% of the compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be dosage form about 2 % to about 60% of the weight of a given unit dosage form. The amount of benzamide compound in such therapeutically useful compositions is such that an effective mammalian dosage level will be obtained for administration. In an aspect the amount is a threshold detectable amount. In an aspect the administration comprises two benzamide compounds. One or more of the benzamide compounds may be administered individually, serially, simultaneously, nearly simultaneous and sequentially. The compounds may be formulated as a time release composition if desired.

[00151] In an aspect, the preparations are consumed orally by a living mammal. In an aspect, the compositions are consumed orally by a human. In an aspect two compounds are administered to a living mammal.

[00152] In an aspect, depending on its form, the administered formulation may contain a binder, disintegrating agent, lubricant, sweetener, a liquid carrier. Pills may be coated or uncoated. A syrup may contain sucrose or fructose as a sweetener. Formulation ingredients used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed to a living mammal. In addition, if desired the compound may be incorporated into sustained-release preparations and administered using devices therefor. In an aspect an anti-microbial ingredient is included. In an aspect a preservative is included.

[00153] In an aspect, the benzamide compound is administered to a mammal as a pharmacologically acceptable composition such as in an aqueous

composition. Pharmacologically acceptable compositions such as solutions of a benzamide compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared. Any pharmacologically suitable humane method of administration may be employed so that an effective radiolabeled amount of benzamide compound is made available to the biochemical system of the mammal.

[00154] In an aspect, the pharmaceutical dosage forms suitable for injection or infusion include sterile aqueous solutions or dispersions or sterile powders comprising a radiolabeled or unlabeled compound of Formula (I), Formula (II) and Formula (III) adapted for preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. The administered dosage form should be sterile, fluid and stable. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising water, ethanol, a polyol, vegetable oils, and nontoxic glyceryl esters.

[00155] The fluidity of preparations can be maintained by the formation of liposomes and by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. Antibacterial and antifungal agents may be employed. An isotonic agent such as an effective amount of a sugar, buffer or sodium chloride may be employed. Delaying absorption agents such as aluminum monostearate and gelatin may be employed if desired.

[00156] In an aspect sterile injectable solutions are prepared by incorporating the novel benzamide compound(s) in the desired amount in the appropriate solvent with other ingredients as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the labeled or unlabeled compound of Formula (I), Formula (II) and Formula (III) along with any additional desired ingredient present in the previously sterile-filtered solutions.

[00157] For topical administration, a benzamide compound of Formula (I), Formula (II) and Formula (III) may be applied in pure form, i.e., when

they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

[00158] Useful effective dosages of the benzamide compounds of Formula (I), Formula (II) and Formula (III) can be determined by comparing their in vitro activity and in vivo activity using animal models. Methods for the extrapolation of effective dosages of some moieties in mice, and other animals, to humans are known for example, see U.S. Pat. No. 4,938,949.

[00159] In an aspect the amount of time elapsing between imagining is a time which provides for a useful and meaningful comparison of acquired images.

[00160] In an aspect the benzamide compound is applied as a radiolabeled benzamide compound.

[00161] Generally, the concentration of the benzamide compound(s) of Formula (I) Formula (II) and Formula (III) in a liquid composition, such as a lotion, will be from about 0.1- about 25 wt-%, preferably from about 0.5- about 10 wt-%. The concentration in a semi-solid or solid composition such as a gel or a powder will be about 0.1-about 5 wt-%, preferably about 0.5- about 2.5 wt-%. Single dosages for injection, infusion or ingestion will generally vary between 50-1500 mg, and may be administered, i.e., 1-3 times daily, to yield levels of about 0.5-50 mg/kg, for adult humans.

[00162] Accordingly, this discovery includes a pharmaceutical composition comprising a labeled or unlabeled compound of Formula (I) Formula (II) and Formula (III) as described hereinabove; or a pharmaceutically acceptable salt thereof; and as a pharmaceutically acceptable diluent or carrier.

[00163] In an aspect a pharmaceutically acceptable salt includes any water soluble salt which is pharmaceutically suitable to the mammalian recipient of the benzamide or radiolabeled compound. In an aspect a pharmaceutically acceptable diluent or carrier includes an aqueous diluent or any diluent or carrier which is innocuous to the mammal recipient of the benzamide compound and which provides

for facilitation of the administration of the benzamide compound(s) and their radionuclide counterparts.

[00164] The precise dosage of the detectably labeled benzamide compound to be administered and the length of time over which administration is carried out will depend on a number of factors including the age and weight of the mammal patient and the route of administration.

[00165] In an aspect a therapeutic rate titration is performed wherein the living mammalian afflicted with cancer or believing to be so afflicted with cancer is administered a series of dosages and respective effects therefrom or thereafter are determined by an inventive method herein at respective dosages and times. In this manner a therapeutic dosage curve or titration is obtained for determining dosage for that mammal patient.

[00166] In an aspect, the radionuclide is purified.

[00167] Administration may be performed by local or systemic application as appropriate. Administration of compositions may be done by inhalation, orally, rectally or parenterally, such as by intramuscular, subcutaneous, intraarticular, intracranial, intradermal, intraocular, intraperitoneal, intrathecal and intravenous injection. The injection may be by stereotaxic injection. Local administration may also be performed, e.g. at an affected site e.g. by use of a catheter or syringe. Treatment by topical application of a composition, e.g. an ointment, to the skin is appropriate. Administration may be performed at intervals of time, such as two or more applications, at some intervals, such as several times a day, or at periodic intervals of the daily or daily.

[00168] Compounds of Formula (I) Formula (II) and Formula (III) can be suitably radiolabeled using any of a number of techniques which are well known in the art. For example, a radioisotope can be incorporated into said compound or appended to said compound of Formula (I) using techniques well known in the art, for example, techniques analogous to those described in Arthur Murry III, D. Lloyd Williams; Organic Synthesis with Isotopes, vol. I and II, Interscience

Publishers Inc., N.Y. (1958) and Melvin Calvin et al. Isotopic Carbon John Wiley and Sons Inc., N.Y. (1949). Preferably, a compound of Formula (I) may be labeled effectively by appending a radioisotope of a halogen to the aromatic rings comprising Ar, Ar' and Ar". In an aspect compounds of Formula (I), Formula (II) and Formula (III) are radiolabeled prior to formulation and administration.

[00169] Additionally, a benzamide compound of Formula (I), Formula (II) and Formula (III) can be labeled with a metal chelating group optionally comprising a radionuclide, such as a metallic radioisotope. Such chelating groups are well known in the art and include polycarboxylic acids such as for example diethylenetriaminepentaacetic acid, ethylenediaminetetraacetic acid, and the like, or analogs or homologs thereof, as well as the chelating groups disclosed in Anderson and Welch (Chem Rev. 99: 2219-2234, 1999) and Jurisson and Lydon (Chem. Rev. 99: 2205-2218, 1999).

[00170] The chelating group or the radionuclide therein may be attached directly to a compound of Formula (I), Formula (II) and Formula (III) or may be attached to a compound of Formula (I), Formula (II) and Formula (III) by means of a divalent or bifunctional organic linker group. Such bifunctional organic linker groups are well known in the art and are preferably less than about 50 angstroms in length. Examples of suitable bifunctional linker groups include 2-carboxymethyl, 3-carboxypropyl, 4-carboxybutyl, and the like. Preferably, the bifunctional linker group is attached to a compound of Formula (I) at the amino nitrogen which is substituted by the group, NRR' in Formula (I). The linker group may also be attached at any synthetically feasible position. For example, Figure 7 shows two compounds of the invention (compounds II and III) which are compounds of Formula (I), labeled with a metal chelating group comprising a radionuclide (M).

[00171] Any metallic radioisotope capable of being detected in a PET or SPECT or microPET[®] diagnostic imaging procedure can be employed as a functional radionuclide.

Suitable nonlimiting examples of useful radionuclides include: Actinium-225, Astatine-211, Bismuth-212, Bismuth-213, Bromine-75, Bromine-76, Carbon-11, Cerium-

¹⁴¹, Chromium-⁵¹, Copper-⁶⁰, Copper-⁶¹, Copper-⁶², Copper-⁶⁴, Copper-⁶⁷, Dysprosium-¹⁶⁶, Fluorine-¹⁸, Gadolinium-¹⁵², Gadolinium-¹⁵³, Gold-^{195m}, Holmium-¹⁶⁶, Indium-¹¹¹, Indium-^{113m}, Iodine-¹²³, Iodine-¹²⁴, Iodine-¹²¹, Iron-⁵⁵, Iron-⁵⁹, Lutetium-¹⁷⁷, Nitrogen-¹³, Oxygen-¹⁵, Palladium-¹⁰³, Radium-²²³, Radium-²²⁴, Rhenium-¹⁸⁶, Rhenium-¹⁸⁸, Rubidium-⁸¹, Rubidium-⁸², Rubidium-⁸⁶, Ruthenium-¹⁰³, Ruthenium-¹⁰⁶, Samarium-¹⁵³, Scandium-⁴⁶, Tantalum-¹⁷⁸, Technetium-^{94m}, Technetium-^{99m}, Thallium-²⁰¹, Titanium-⁴⁵, Ytterbium-¹⁶⁹, Yttrium-⁸⁶, Yttrium-⁹⁰, and Zirconium-⁸⁹, In an aspect technetium-^{99m} is used for SPECT imaging studies, and rhenium-¹⁸⁸, rhenium-¹⁸⁶, copper-⁶⁴ and yttrium-⁹⁰ are useful for radiotherapy of breast tumors. In an aspect the compounds of Formula (I), Formula (II) and Formula (III) including compound 23 are radiolabeled with at least one of tritium and iodine ¹²⁵ radiolabels using standard radiolabeling conditions to produce at least one of a tritiated and iodine ¹²⁵ labeled analog that can be used in the invitro detection of sigma 2 receptors in tumors and normal tissue.

[00172] PET (or "Positron Emission Tomography"), and SPECT are non-invasive molecular diagnostic imaging (standard) medical procedures that produce (i.e. capture and optionally record) multiple acquisitions i.e. images of the body's biological functions and in an aspect are used to determine the extent of malignant disease. In an aspect, these imaging procedures show the presence and distribution of a radiolabeled detectable functionally emitting radiolabeled chemical i.e. a radionuclide acquisitioned at various selected times. Advantageously these two imaging procedures depict both metabolic characteristics of tissues and changes therein.

[00173] In an aspect, positive emission tomography (PET imaging) comprises detection of γ -rays emitted from radionuclides such as ⁷⁶Br Compound 6, ¹²³I Compound 6, ¹²⁴I Compound 6, ¹²⁵I Compound 6 and ¹¹C Compound 10, that decay by positron emission and are located within a mammalian patient's body. In an aspect this is possible by virtue of administration of a radiolabeled benzamide compound to the patient.

[00174] In an aspect, single photon emission computed tomography ("SPECT imaging") comprises a collimation of gamma rays emitted by a

radiopharmaceutical distribution such as detectable radioactivity emitting a benzamide compound of the invention or a pharmaceutical composition comprising a novel benzamide compound within the mammalian body undergoing treatment and analysis. Generally collimators for SPECT imaging are lead and comprise thousands of various shaped parallel channels through which - and only through which - gamma rays are allowed to pass. Generally such collimators are positioned over a single crystal of NaI contained in the Gamma camera in an arrangement referred to as an Anger camera, <http://www.amershamhealth.com/medcyclopaedia/Volume%20I/Anger%20camera>). The image or acquisition from the camera is the captured image which is presented to a human operator as part of a viewable image. This may be a screen shot or a captured digital image, which may be stored in a computer storage. In an aspect multi-acquisition is used. In an aspect a multi-acquisition is carried out over an elapsed time interval.

[00175] Single-photon emission computerized tomography (SPECT) scan creates three-dimensional images of internal organs revealing anatomy and function. A tiny amount of radioactive tracer is injected through a vein and a camera detects the radiation emitted by these tracers.

[00176] In an aspect an external measurement is made of the two high energy photons emitted in opposite directions when a positron-emitting radionuclide decays in a patient. A large number of scintillation detectors detect these photon pairs and measure the sum of radioactivity along many different paths through the patient undergoing measurement. Appropriate software associated with the operating instrument reconstructs a three-dimensional image of the patient and the concentrations of radionuclides can be expressed in quantitative units of radiotracer concentration per ml of tissue.

[00177] In an aspect images are acquisitioned (taken) over elapsed time in dynamic fashion to assemble a developing or developed scenario of developing or changing situations in the mammalian patient. It is believed that the tumorous or cancerous areas have a higher density of these receptors than surrounding normal tissue and thus that is why such areas show up on the image.

[00178] In an aspect a PET and/or a SPECT image is taken of a living mammal after administration of a compound of Formula (I) Formula (II) and Formula (III) to a living mammal. The image may be retained in computer storage if desired. A number of images may be acquired as a function of elapsed time to produce a profile over time of the images.

[00179] In an aspect a radioactive substance is produced in a process and is attached, or tagged, to a benzamide compound referred to as radiolabeling. After this radiolabeled benzamide as compound 6 labeled with one of ^{76}Br , ^{123}I , ^{124}I , ^{125}I or compound 10 labeled with ^{11}C is administered to a patient, radioactivity travels through gastro-intestinal tract through the vascular circulator (blood) system of the body and localizes in the appropriate areas of the body and is detected by PET or SPECT scanner.

[00180] Typically an adequate amount of time is allowed to lapse for the treated living mammal (i.e. having received the radiolabeled benzamide) to come to an equilibrium state following satisfactory administration of the benzamide radioligand to the mammal. Typically the mammal is placed in a position near the PET instrument or SPECT instrument or microPET[®] instrument allowing satisfactory operation of the PET instrument and/or SPECT instrument. The PET and SPECT instruments are equipped with all necessary operable software and operation requirements. These instruments are turned on by supplying 100 volts electric power to the instruments.

[00181] Generally after mammal has received its administration of the radiolabeled benzamide the mammal is placed in/on the PET scanner, which has a opening in the middle. In the PET scanner there are multiple rings of detectors that record the emission of energy from the radioactive substance now within in the mammal. In an aspect the mammal is comfortably moved into the hole of the machine. The images are displayed on the monitor of a computer, suitably equipped and operably coupled to the PET scanner instrument for acquiring.

[00182] Examples 1-5 following are illustrative of but are not meant to be limiting of this discovery.

EXAMPLES

Example 1

[00183] Synthesis of target tetrahydroisoquinoline analogs is shown in Figure 4 and Figure 5. Reaction of the secondary amine of compounds 13 and 14 with either bromoacetonitrile or bromobutyronitrile gave N-alkylated products, 15, 16, 17, and 18, in 75 – 85% yield. Reduction with either lithium aluminum hydride in THF or hydrogenation over palladium on charcoal gave the corresponding amines, 19, 20, 21, 22, in quantitative yields. Condensation of amines 19, 20, 21, 22 with either 2-methoxy-5-bromonaphthoyl chloride or 5-bromo-2,3-dimethoxybenzoic acid gave the corresponding amide analogs, 1, 2, 3, 4, 5, 6, in excess of 90% yield. Condensation of amines 19, 21 and 22 with either 5-bromo-2-methoxybenzoic acid or 5-methyl-2-methoxybenzoic acid gave amides 7, 8, 9, and 10 in excess of 90% yield. Synthesis of the 2,3-dichlorophenylpiperazine benzamide and naphthamide analogs, 11 and 12, was accomplished using a similar reaction sequence as outlined in Figure 5.

Example 2

Physical Data

[00184] The following physical data was obtained for and identifies and characterizes benzamide compounds and attests to their preparation denoted respectively by numbers (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12) herein. For example the immediately following paragraph presents the melting point of the respective oxalate salt, the results of a NMR (nuclear magnetic resonance analysis) and comparison calculated and observed molecular weights.

[00185] Physical Data for Compound 1. MP (Melting Point) 142 °C-144 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 2.99 (s, 2H), 3.10 (s, 2H), 3.61–3.63 (m, 3H), 3.73 (s, 3H), 3.85 (s, 3H), 4.15 (s, 3H), 7.13-7.23 (m, 4H), 7.34 (s, 1H), 7.36 (s, 1H), 8.59 (t, J = 5.0 Hz, 1H). Calculated: C: 51.88; H: 4.95; N: 5.50. Observed: C: 51.73; H: 4.95; N: 5.51.

[00186] Physical Data for Compound 2. MP 91°C-93 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 2.92 (s, 2H), 3.13 (s, 2H), 3.59-3.63 (m, 3H), 3.71 (s, 3H), 3.73 (s, 3H), 3.74 (s, 3H), 3.85 (s, 3H), 4.10 (s, 3H), 6.72 (s, 1H), 6.78 (s, 1H), 7.34 (s, 1H), 7.37 (s, 1H), 8.60 (s, 1H). Calculated: C: 50.63; H: 5.13; N: 4.92. Observed: C: 50.49; H: 5.30; N: 4.60.

[00187] Physical Data for Compound 3. MP 180 °C-182 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 3.02 (s, 2H), 3.19 9s, 2H), 3.72-3.74 (m, 3H), 3.94 (s, 3H), 4.21 (s, 3H), 7.16-7.25 (m, 4H), 7.75 (t, J = 8.1 Hz, 1H), 7.81 (t, J = 8.1 Hz, 1H), 8.05 (s, 1H), 8.15 (d, J = 8.3 Hz, 1H), 8.24 (d, J = 8.3 Hz, 1H), 8.77 (t, J = 5.4 Hz, 1H). Calculated: C: 56.72; H: 4.76; N: 5.29. Observed: C: 56.50; H: 4.91; N: 5.14.

[00188] Physical Data for Compound 4. MP 187°C-189 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 2.94 (s, 3H), 3.19 (s, 4H), 3.71 (s, 3H), 3.73 (s, 3H), 3.94 (s, 3H), 4.12 (s, 3H), 6.74 (s, 1H), 6.79 (s, 1H), 7.75 (t, J = 7.6 Hz, 1H), 7.82 (t, J = 7.6 Hz, 1H), 8.05 (s, 1H), 8.15 (d, J = 8.3 Hz, 1H), 8.25 (d, J = 8.3 Hz, 1H), 8.76 (s, 1H). Calculated: C: 55.02; H: 4.96; N: 4.75. Observed: C: 54.76; H: 5.04; N: 4.65.

[00189] Physical Data for Compound 5. MP 165°C-167 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 1.61-1.80 (m, 4H), 2.96 (s, 2H), 3.36-3.39 (m, 5H), 3.71 (s, 3H), 3.73 (s, 3H), 3.95 (s, 3H), 4.19 (s, 3H), 6.77 (s, 1H), 6.80 (s, 1H), 7.75 (t, J = 7.4 Hz, 1H), 7.81 (t, J = 7.4 Hz, 1H), 7.95 (s, 1H), 8.15 (d, J = 8.3 Hz, 1H), 8.25 (d, J = 8.3 Hz, 1H), 8.60 (t, J = 5.6 Hz, 1H). Calculated: C: 56.41; H: 5.39; N: 4.54. Observed: C: 56.32; H: 5.45; N: 4.45.

[00190] Physical Data for Compound 6. ¹H-NMR (CDCl₃) δ 1.70-1.79 (m, 4H), 2.59 (s, 2H), 2.73-2.76 (m, 2H), 2.81-2.84 (m, 2H), 3.51-3.52 (m, 2H), 3.58 (s, 2H), 3.85 (s, 3H), 3.86 (s, 3H), 3.88 (s, 3H), 3.90 (s, 3H), 6.52 (s, 1H), 6.60 (s, 1H), 7.12 (d, J = 2.7 Hz, 1H), 7.78 (d, J = 2.7 Hz, 1H), 8.05 (s, 1H).

[00191] Physical Data for Compound 7. MP 166°C-168 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 3.02 (s, 3H), 3.14 (s, 2H), 3.66 (s, 2H), 3.83 (s, 3H), 4.19 (s, 3H), 7.12-7.25 (m, 5H), 7.65-7.67 (m, 1H), 7.86 (s, 1H), 8.56 (t, J = 5.4 Hz, 1H). Calculated: C: 52.62; H: 4.84; N: 5.84. Observed: C: 52.38; H: 4.75; N: 5.69.

[00192] Physical Data for Compound 8. MP 158°C-160 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 2.93 (s, 2H), 3.16 (s, 2H), 3.60-3.66 (m, 3H), 3.71 (s, 3H), 3.73 (s, 3H), 3.84 (s, 3H), 4.14 (s, 3H), 6.73 (s, 1H), 6.78 (s, 1H), 7.13 (d, J = 8.9 Hz, 1H), 7.66 (d, J = 8.7 Hz, 1H), 7.86 (s, 1H), 8.57 (t, J = 5.4 Hz, 1H). Calculated: C: 51.22; H: 5.05; N: 5.19. Observed: C: 51.23; H: 5.07; N: 5.04.

[00193] Physical Data for Compound 9. MP 160°C-162 °C (oxalate salt); ¹H-NMR (d₆-DMSO) δ 2.27 (s, 3H), 2.94 (s, 2H), 3.16 (s, 2H), 3.65-3.66 (m, 3H), 3.72 (s, 3H), 3.73 (s, 3H), 3.82 (s, 3H), 4.14 (s, 3H), 6.73 (s, 1H), 6.79 (s, 1H), 7.03 (d, J = 8.4 Hz, 1H), 7.29 (d, J = 8.4 Hz, 1H), 7.63 (s, 1H), 8.51 (s, 1H). Calculated : C: 60.18; H: 6.42; N: 5.85. Observed: C: 60.32; H: 6.39; N: 5.56.

[00194] Physical Data for Compound 10. ¹H-NMR (CDCl₃) δ 1.70-1.79 (m, 4H), 2.32 (s, 3H), 2.59 (s, 2H), 2.73-2.76 (m, 2H), 2.81-2.84 (m, 2H), 3.51-3.52 (m, 2H), 3.58 (s, 2H), 3.81 (s, 3H), 3.83 (s, 3H), 3.89 (s, 3H), 6.50 (s, 1H), 6.58 (s, 1H), 6.86-6.96 (m, 2H), 7.11 (s, 1H), 8.00 (s, 1H).

[00195] Physical Data for Compound 11. ¹H-NMR (CDCl₃) δ 1.66 (s, 4H), 2.44-2.66 (m, 6H), 3.06 (s, 2H), 3.17-3.21 (m, 2H), 3.48-3.50 (m, 2H), 3.88 (s, 6H), 6.76-6.96 (m, 2H), 7.12-7.16 (m, 2H), 7.77-7.79 (m, 1H), 7.99 (s, 1H).

[00196] Physical Data for Compound 12. ¹H-NMR (CDCl₃) δ 1.72 (s, 4H), 2.47-2.67 (m, 6H), 3.03 (s, 2H), 3.15-3.20 (m, 2H), 3.54-3.63 (m, 2H), 3.99 (s, 3H), 6.73-6.93 (m, 2H), 7.11-7.17 (m, 1H), 7.60-7.72 (m, 2H), 8.01-8.26 (m, 3H), 8.38 (s, 1H).

[00197] Physical data for Compound 23. ¹H-NMR (CDCl₃) δ 1.60-1.70 (m, 4H), 2.33(s, 3H), 2.51-2.56 (t, 2H), 2.67-2.72 (t, 2H), 2.78-2.83 (t, 2H), 3.48-3.54 (m, 4H), 3.82-3.83 (s, 6H), 4.21-4.25 (t, 1H), 4.30-4.34 (t, 1H), 4.68-4.72 (t, 1H), 4.84-4.88 (t, 1H), 6.49 (s, 1H), 6.57 (s, 1H), 6.99-6.83 (d, 1H), 7.19-7.30 (d, 1H), 7.95 (s, 1H), 7.99 (d, 1H).

[00198] Example 3 σ receptor binding assays.

[00199] The σ_1 receptor binding assay was conducted using guinea pig brain membrane homogenates (100 μ g protein). Membrane homogenates were incubated with 3 nM [3 H](+)-pentazocine (31.6 Ci/mmol) in 50 mM Tris-HCl (pH 8.0) at 25 °C for either 120 or 240 min. Test compounds were dissolved in ethanol then diluted in buffer for a total incubation volume of 0.5 ml. Test compounds were added in concentrations ranging from 0.005 to 1000 nM. Assays were terminated by the addition of ice-cold 10 mM Tris-HCl (pH 8.0) followed by rapid filtration through Whatman GF/B glass fiber filters (presoaked in 0.5 % polyethylenimine) using a Brandel cell harvester (Gaithersburg, MD). Filters were washed twice with 5 ml of ice cold buffer. Nonspecific binding was determined in the presence of 10 μ M (+)-pentazocine. Liquid scintillation counting was carried out in EcoLite(+) (ICN Radiochemicals; Costa Mesa, CA now Valeant Pharmaceuticals International, Valeant Plaza, 3300 Hyland Avenue, Costa Mesa, CA 92626 USA), using a Beckman LS 6000IC spectrometer with a counting efficiency of 50%.

[00200] The σ_2 receptor binding assay was conducted using rat liver membrane homogenates (35 μ g of protein). Membrane homogenates were incubated with 3 nM [3 H]DTG (38.3 Ci/mmol) in the presence of 100 nM (+)-pentazocine to block σ_1 sites. Incubations were carried out in 50 mM Tris-HCl (pH 8.0) for 120 min at 25 °C in a total incubation volume of 0.5 ml. Test compounds were added in concentrations ranging from 0.005 to 1000 nM. Assays were terminated by the addition of ice-cold 10 mM Tris-HCl (pH 8.0) followed by rapid filtration through Whatman GF/B glass fiber filters (presoaked in 0.5% polyethylenimine) using a Brandel cell harvester (Brandel, 8561 Atlas Dr., Gaithersburg, MD 20877, USA). Filters were washed twice with 5 ml of ice cold buffer. Nonspecific binding was determined in the presence of 5 μ M DTG. Liquid scintillation counting was carried out in EcoLite(+) (ICN Radiochemicals; Costa Mesa, CA, now Valeant Pharmaceuticals International, Valeant Plaza, 3300 Hyland Avenue, Costa Mesa, CA 92626 USA) using a Beckman LS 6000IC spectrometer with a counting efficiency of 50%.

[00201] The IC₅₀ values at sigma sites were generally determined in triplicate from non-linear regression of binding data as analyzed by JMP (SAS

Institute Incv., JMP Software, SAS Campus Drive; Cary, NC 27513), using 8 concentrations of each compound. K_i values were calculated using the method of Cheng-Prusoff (Biochem. Pharmacol. 1973, 22, 3099-4022) and represent mean values \pm SEM. All curves were best fit to a one site fit and gave Hill coefficients of 0.8 – 1.0. The K_d value used for [3 H]DTG in rat liver was 17.9 nM and was 4.8 nM for [3 H](+)-pentazocine in guinea pig brain.^{11,12}

[00202] Generally compounds of the invention demonstrate high selectivity for sigma-2 versus sigma-1 receptors (Table).

Example 4

[00203] The σ_2 radioligand ([76 Br]6), was labeled with 76 Br via an electrophilic destannylation reaction. The reaction takes place under aqueous (200 μ L Milli-Q) conditions in the presence of 5% peracetic acid/acetic acid (40 μ L) for 40 min. (A) was purified via HPLC with yields typically between 50-60%.

[00204] This compound was studied in mature Balb/C mice that were implanted with EMT-6 mammary tumors. The mice were implanted in the nape of the neck 7 days prior to the study. The biodistribution study consisted of three groups: 1hr low dose (~ 6 μ Ci), 4hr low dose and a 4 hr high dose (~ 150 μ Ci). The %ID/g at 1 hr for tumor, brain, fat, blood and liver were 4.0 ± 0.4 , 0.25 ± 0.02 , 1.1 ± 0.4 , 2.1 ± 0.3 and 5.4 ± 0.4 , respectively. At 4 hr for the low dose animals the values decreased to 1.2 ± 0.2 , 0.15 ± 0.02 , 0.3 ± 0.2 , 0.82 ± 0.08 and 1.3 ± 0.2 , respectively. The %ID/g for the 4 hr high dose animals was not significantly different. The activity injected into the high dose animals was enough to perform microPET[®] imaging studies. At 1 and 2 hr the tumors were clearly identified in the three animals that were injected. This initial study has shown that this σ_2 receptor compound has a high uptake into EMT-6 mammary tumors and can be imaged non-invasively. (MicroPET[®] is a dedicated PET scanner designed for high resolution imaging of small laboratory animals. It is available from Concorde Microsystems, Inc. 10427 Cogdill Rd, Suite 500 Knoxville, TN 37932 USA.)

[00205] Without being bound by theory, it is believed that compounds of Formula (I), Formula (II) and Formula (III) when labeled with detectable radionuclides provide detectably labeled ligands that selectively bind to cancer cells and can be quantified by using one or more of functional imaging techniques such as positron emission tomography (PET), including MicroPET[®], single photon emission computed tomography (SPECT), MRI, computed tomography (CT) and functional magnetic resonance imaging (fMRI). Said components have the potential to noninvasively assess the proliferative status of known or suspected tumor cells or cells subject to hyperplasia, in bladder, colon, prostate, breast, lung, gut, pancreas, reproductive system, brain and the like. The labeled compounds having a structure illustrated in Formula (I) can also be used to treat cancer or abnormally dividing cells, by selectively inhibiting their proliferation.

[00206] The benzamides of Formula (I) are useful as radiotracers for imaging dopamine D₃ receptors with PET including compound 11 (Table IA and Table 2).

[00207] Since many of the compounds (1-12) depicted in Table IA have a high affinity for dopamine D₃ receptors, they are useful in the treatment of affective disorders such as schizophrenia, and depression as well as movement disorders such as Parkinson's Disease.

Example 5


[00208] BIOLOGICAL DATA

[00209] This section focuses on the use of the sigma-2 (σ_2) receptor as a biomarker of tumor proliferation. This is based on inventor data demonstrating that σ_2 receptors are expressed in about 10-fold higher density in proliferative mouse mammary adenocarcinoma cells versus the nonproliferative or quiescent cell population under both in vitro (cell culture) and in vivo (tumor xenographs) conditions. Therefore, a σ_2 receptor PET radiotracer has the potential to provide information regarding the proliferative status of breast cancer. An in vivo imaging procedure that can provide information about the proliferative status of primary breast

tumors would represent a significant improvement over current methods used in making this assessment. Our preliminary data also indicate that σ_2 -selective radiotracers are predicted to have a better tumor:non-tumor ratio than other agents, such as FDG and the DNA precursors, currently used to assess proliferation in PET oncology studies.

[00210] A. Carbon-11 labeled analogs. The inventors identified a number of lead compounds having a high affinity and selectivity for σ_2 versus σ_1 receptors (Table I). The presence of a 2-methoxy group in the lead compounds indicates that it is possible to prepare a ^{11}C -labeled version of the σ_2 ligands using standard radiochemistry procedures. This was accomplished via O-alkylation of the phenol precursor. The overall yield (15-75% from $[^{11}\text{C}]$ methyl iodide) and specific activity (1000 – 4000 mCi/ μmol) of each radiotracer was suitable for *in vivo* studies.

Table I.



| | X | R | n | σ_1 | σ_2 | $\sigma_1: \sigma_2$ | % Yield | Specific |
|----|------------------|-----------------|---|------------|------------|----------------------|---------|--------------------------------|
| # | | | | | | Ratio | | Activity (EOB) |
| 9 | H | CH ₃ | 2 | 10,412 | 13.3 | 783 | 60-75 | ~5,000 mCi/ μmol |
| 10 | H | CH ₃ | 4 | 3,078 | 10.3 | 300 | 60-75 | ~4,000 mCi/ μmol |
| 8 | H | Br | 2 | 5,484 | 12.2 | 442 | 10-15 | ~1,000 mCi/ μmol |
| 6 | OCH ₃ | Br | 4 | 12,900 | 8.2 | 1,573 | 30-40 | ~4,000 mCi/ μmol |

[00211] B. Biodistribution Studies of ^{11}C analogs in Tumor-Bearing Mice. Biodistribution studies were conducted in mature Balb/c mice that were implanted with EMT-6 mammary tumors identified herein as EMT-6 BALB/C mice to create laboratory tumor-bearing mice. The mice were implanted in the scapular region seven days prior to the study. Animals were injected with 100-150 μCi of the ^{11}C -labeled radiotracer and the animals were sacrificed at 5, 30, and 60 min post-i.v. injection of the radiotracer.

[00212] The results of the biodistribution studies are given respectively in Tables II-V following.

Table II.

| [^{11}C]9 Biodistribution in EMT-6 BALB/C mice | | | |
|---|------------------|-----------------|-----------------|
| %ID per gram | 5 min | 30 min | 1 hour |
| blood | 5.89 ± 0.29 | 2.62 ± 0.22 | 1.98 ± 0.35 |
| lung | 5.69 ± 0.70 | 1.42 ± 0.15 | 1.39 ± 0.67 |
| liver | 18.49 ± 2.87 | 3.87 ± 0.67 | 1.70 ± 0.23 |
| kidney | 44.07 ± 1.67 | 2.77 ± 0.42 | 1.01 ± 0.12 |
| muscle | 1.75 ± 0.21 | 0.56 ± 0.19 | 0.41 ± 0.22 |
| fat | 3.07 ± 0.40 | 0.38 ± 0.12 | 0.26 ± 0.09 |
| heart | 2.89 ± 0.36 | 0.76 ± 0.06 | 0.76 ± 0.46 |
| brain | 1.63 ± 0.30 | 0.11 ± 0.01 | 0.10 ± 0.04 |
| tumor | 3.10 ± 0.25 | 1.08 ± 0.08 | 0.85 ± 0.14 |
| ratio | | | |
| Tumor: blood | 0.53 ± 0.04 | 0.41 ± 0.02 | 0.44 ± 0.06 |
| Tumor: lung | 0.55 ± 0.07 | 0.77 ± 0.14 | 0.68 ± 0.19 |
| Tumor: muscle | 1.79 ± 0.20 | 2.08 ± 0.59 | 2.40 ± 0.84 |
| Tumor: fat | 1.03 ± 0.21 | 3.10 ± 1.21 | 3.46 ± 0.91 |
| Tumor: heart | 1.08 ± 0.10 | 1.43 ± 0.14 | 1.32 ± 0.45 |

Table III.

| [¹¹ C]10 Biodistribution in EMT-6 BALB/C mice | | | |
|---|--------------|-------------|-------------|
| %ID per gram | 5 min | 30 min | 1 hour |
| blood | 3.09 ± 0.33 | 1.31 ± 0.11 | 0.73 ± 0.05 |
| lung | 14.02 ± 1.40 | 2.27 ± 0.42 | 1.09 ± 0.26 |
| liver | 12.32 ± 1.73 | 9.65 ± 2.00 | 3.00 ± 0.21 |
| kidney | 20.50 ± 1.86 | 4.12 ± 0.51 | 2.26 ± 0.36 |
| muscle | 4.49 ± 0.45 | 0.75 ± 0.13 | 0.49 ± 0.11 |
| fat | 1.88 ± 0.50 | 0.68 ± 0.19 | 0.33 ± 0.24 |
| heart | 5.86 ± 0.47 | 0.95 ± 0.17 | 0.50 ± 0.11 |
| brain | 2.29 ± 0.28 | 0.28 ± 0.03 | 0.15 ± 0.01 |
| tumor | 4.22 ± 1.01 | 2.35 ± 0.27 | 1.32 ± 0.17 |
| ratio | | | |
| Tumor:blood | 1.37 ± 0.33 | 1.80 ± 0.26 | 1.81 ± 0.11 |
| Tumor:lung | 0.31 ± 0.11 | 1.06 ± 0.24 | 1.28 ± 0.41 |
| Tumor:muscle | 0.93 ± 0.13 | 3.18 ± 0.51 | 2.78 ± 0.62 |
| Tumor:fat | 2.28 ± 0.32 | 3.68 ± 1.14 | 5.36 ± 2.38 |
| Tumor:heart | 0.71 ± 0.12 | 2.53 ± 0.55 | 2.78 ± 0.79 |

Table IV.

| [¹¹ C]8 Biodistribution in EMT-6 BALB/C mice | | | |
|--|--------------|--------------|--------------|
| %ID per gram | 5 min | 30 min | 1 hour |
| blood | 5.25 ± 0.39 | 2.35 ± 0.16 | 1.88 ± 0.16 |
| lung | 5.72 ± 0.40 | 1.83 ± 0.13 | 1.32 ± 0.11 |
| liver | 19.88 ± 3.10 | 5.89 ± 0.82 | 2.65 ± 0.29 |
| kidney | 51.03 ± 7.14 | 34.19 ± 1.74 | 19.78 ± 1.99 |
| muscle | 1.73 ± 0.11 | 0.52 ± 0.23 | 0.36 ± 0.08 |
| fat | 2.05 ± 0.49 | 0.63 ± 0.19 | 0.37 ± 0.13 |
| heart | 3.18 ± 0.26 | 0.77 ± 0.08 | 0.56 ± 0.05 |
| brain | 2.52 ± 0.15 | 0.26 ± 0.10 | 0.14 ± 0.02 |
| tumor | 1.82 ± 0.39 | 1.06 ± 0.09 | 0.87 ± 0.09 |
| ratio | | | |
| Tumor:blood | 0.35 ± 0.07 | 0.45 ± 0.05 | 0.46 ± 0.02 |
| Tumor:lung | 0.32 ± 0.06 | 0.58 ± 0.07 | 0.66 ± 0.03 |
| Tumor:muscle | 1.05 ± 0.21 | 2.24 ± 0.63 | 2.52 ± 0.66 |
| Tumor:fat | 0.93 ± 0.27 | 1.76 ± 0.40 | 2.64 ± 1.12 |
| Tumor:heart | 0.57 ± 0.09 | 1.39 ± 0.17 | 1.56 ± 0.16 |


Table V.

| [¹¹ C]6 Biodistribution in EMT-6 BALB/C mice | | | |
|--|--------------|-------------|-------------|
| %ID per gram | 5 min | 30 min | 1 hour |
| blood | 7.12 ± 1.01 | 0.99 ± 0.15 | 0.45 ± 0.04 |
| lung | 6.01 ± 0.77 | 1.34 ± 0.23 | 0.70 ± 0.26 |
| liver(all) | 25.02 ± 3.70 | 2.48 ± 0.52 | 1.19 ± 0.17 |
| kidney | 19.48 ± 1.46 | 2.57 ± 0.78 | 1.34 ± 0.19 |
| muscle | 1.94 ± 0.13 | 1.67 ± 0.14 | 0.26 ± 0.07 |
| fat | 1.66 ± 0.41 | 0.46 ± 0.19 | 0.20 ± 0.08 |


| | | | |
|--------------|-------------|-------------|-------------|
| heart | 3.54 ± 0.31 | 0.68 ± 0.12 | 0.29 ± 0.10 |
| brain | 0.33 ± 0.09 | 0.10 ± 0.00 | 0.03 ± 0.00 |
| tumor | 2.82 ± 0.36 | 0.92 ± 0.10 | 0.50 ± 0.09 |
| ratio | | | |
| Tumor:blood | 0.40 ± 0.01 | 0.94 ± 0.05 | 1.10 ± 0.11 |
| Tumor:lung | 0.47 ± 0.04 | 0.69 ± 0.08 | 0.79 ± 0.30 |
| Tumor:muscle | 1.46 ± 0.19 | 0.59 ± 0.05 | 1.84 ± 0.25 |
| Tumor:fat | 1.82 ± 0.45 | 2.05 ± 0.57 | 2.77 ± 0.82 |
| Tumor:heart | 0.80 ± 0.07 | 1.36 ± 0.12 | 1.86 ± 0.64 |

[00213] C. Synthesis of Radiohalogenated σ_2 Receptor Ligands and in vivo studies. The presence of a bromine atom in compounds 8 and 6 (Table I) indicates that it is possible to prepare radiohalogenated probes of the σ_2 receptors by isotopic substitution with B-76, or by replacing the bromine atom with I-125. This was accomplished by preparing the corresponding tin precursor and conducting the oxidative radiohalogenation reactions outlined in Schemes I and II. The ^{125}I -labeled analogs, 6i and 8i, were obtained in an overall yield of 50% and a specific activity of 2200 mCi/ μmol . Similarly, the ^{76}Br -labeled analogs of 5 and 7 were obtained in a yield of 50-60% and a specific activity >1,000 mCi/ μmol .

Scheme I



Scheme II



A series of biodistribution studies in tumor-bearing mice were also conducted with [⁷⁶Br]8, [⁷⁶Br]6, [¹²⁵I]8i, and [¹²⁵I]6i. The results of these studies are presented in Tables VI-IX.

Table VI.

| [⁷⁶ Br]8 Biodistribution in EMT-6 BALB/C mice | | | | |
|---|--------------|--------------|--------------|-------------|
| %ID per gram | 5 min. | 30 min. | 1 hour | 2 hour |
| blood | 4.55 ± 0.59 | 2.55 ± 0.06 | 2.08 ± 0.69 | 3.46 ± 0.66 |
| lung | 5.52 ± 0.73 | 1.59 ± 0.10 | 1.20 ± 0.27 | 1.75 ± 0.20 |
| liver(all) | 17.85 ± 3.17 | 5.57 ± 1.31 | 3.09 ± 0.48 | 3.77 ± 0.80 |
| spleen | 3.22 ± 0.75 | 0.92 ± 0.26 | 0.59 ± 0.06 | 0.72 ± 0.05 |
| kidney | 50.08 ± 2.09 | 32.24 ± 4.64 | 15.60 ± 2.01 | 3.61 ± 0.52 |
| muscle | 1.42 ± 0.05 | 0.57 ± 0.07 | 0.56 ± 0.26 | 0.45 ± 0.02 |

| | | | | |
|--------------|-------------|-------------|-------------|-------------|
| fat | 2.61 ± 0.45 | 0.80 ± 0.05 | 0.66 ± 0.11 | 0.69 ± 0.18 |
| heart | 2.44 ± 0.43 | 0.99 ± 0.13 | 0.78 ± 0.14 | 1.23 ± 0.18 |
| brain | 1.69 ± 0.18 | 0.24 ± 0.05 | 0.15 ± 0.01 | 0.21 ± 0.01 |
| bone | 1.55 ± 0.20 | 0.52 ± 0.02 | 0.50 ± 0.17 | 0.49 ± 0.17 |
| tumor | 2.28 ± 0.14 | 1.30 ± 0.08 | 1.12 ± 0.20 | 1.19 ± 0.13 |
| ratio | | | | |
| Tumor:blood | 0.50 ± 0.04 | 0.51 ± 0.02 | 0.58 ± 0.22 | 0.35 ± 0.21 |
| Tumor:lung | 0.42 ± 0.05 | 0.82 ± 0.05 | 0.98 ± 0.34 | 0.68 ± 0.10 |
| Tumor:muscle | 1.61 ± 0.05 | 2.31 ± 0.40 | 2.47 ± 1.53 | 2.63 ± 2.47 |
| Tumor:fat | 0.90 ± 0.20 | 1.63 ± 0.02 | 1.71 ± 0.29 | 1.82 ± 2.22 |
| Tumor:heart | 0.95 ± 0.13 | 1.32 ± 0.19 | 1.50 ± 0.49 | 0.98 ± 0.13 |

Table VII.

| ⁷⁶ Br]6 Biodistribution in EMT-6 BALB/C mice | | | | | |
|---|--------------|--------------|--------------|-------------|-------------|
| %ID per gram | 5 min. | 30 min. | 1 hour | 2 hour | 4 hour |
| blood | 2.12 ± 0.20 | 2.20 ± 0.24 | 1.60 ± 0.22 | 0.46 ± 0.07 | 0.21 ± 0.03 |
| lung | 24.64 ± 2.74 | 5.81 ± 1.12 | 2.45 ± 0.17 | 0.74 ± 0.04 | 0.29 ± 0.03 |
| liver(all) | 10.99 ± 0.29 | 8.85 ± 0.52 | 4.58 ± 0.36 | 1.67 ± 0.10 | 0.71 ± 0.08 |
| spleen | 12.50 ± 1.46 | 6.91 ± 1.22 | 2.61 ± 0.62 | 0.60 ± 0.03 | 0.20 ± 0.03 |
| kidney | 31.20 ± 2.92 | 18.51 ± 2.66 | 10.81 ± 1.72 | 1.85 ± 0.54 | 0.57 ± 0.12 |
| muscle | 3.62 ± 0.27 | 1.54 ± 0.49 | 0.61 ± 0.11 | 0.20 ± 0.03 | 0.07 ± 0.01 |
| fat | 3.78 ± 0.97 | 2.27 ± 0.16 | 0.81 ± 0.16 | 0.22 ± 0.05 | 0.04 ± 0.02 |
| heart | 7.31 ± 0.70 | 2.15 ± 0.30 | 1.08 ± 0.07 | 0.30 ± 0.03 | 0.11 ± 0.02 |
| brain | 1.60 ± 0.15 | 0.41 ± 0.06 | 0.17 ± 0.02 | 0.05 ± 0.00 | 0.03 ± 0.00 |

| | | | | | |
|--------------|-----------------|-----------------|-----------------|-----------------|-------------------|
| bone | 3.10 ± 0.67 | 2.76 ± 0.58 | 1.38 ± 0.09 | 0.56 ± 0.20 | 0.12 ± 0.03 |
| tumor | 4.78 ± 0.78 | 5.31 ± 0.62 | 3.98 ± 0.58 | 1.71 ± 0.17 | 0.68 ± 0.15 |
| ratio | | | | | |
| Tumor:bone | 2.25 ± 0.28 | 2.41 ± 0.08 | 2.53 ± 0.54 | 3.79 ± 0.99 | 3.17 ± 0.58 |
| Tumor:liver | 0.19 ± 0.02 | 0.93 ± 0.09 | 1.64 ± 0.30 | 2.30 ± 0.13 | 2.36 ± 0.46 |
| Tumor:muscle | 1.32 ± 0.16 | 3.70 ± 1.06 | 6.76 ± 2.09 | 8.81 ± 0.91 | 9.48 ± 2.14 |
| Tumor:fat | 1.30 ± 0.24 | 2.35 ± 0.34 | 5.09 ± 1.39 | 7.97 ± 1.95 | 20.69 ± 10.77 |
| Tumor:heart | 0.65 ± 0.07 | 2.48 ± 0.17 | 3.72 ± 0.72 | 5.77 ± 0.60 | 6.24 ± 1.28 |

Table VIII.

| $[^{125}\text{I}]\text{8I}$ Biodistribution in EMT-6 BALB/C mice | | | | |
|--|------------------|------------------|------------------|-----------------|
| %ID per gram | 5 min. | 30 min. | 1 hour | 2 hour |
| blood | 6.90 ± 1.29 | 2.44 ± 0.65 | 1.17 ± 0.29 | 3.37 ± 1.23 |
| lung | 5.95 ± 1.30 | 1.39 ± 0.29 | 0.81 ± 0.20 | 1.27 ± 0.37 |
| liver(all) | 41.37 ± 7.23 | 8.20 ± 1.05 | 2.87 ± 0.34 | 3.47 ± 1.05 |
| kidney | 54.77 ± 8.42 | 38.59 ± 2.05 | 18.07 ± 2.07 | 8.46 ± 1.26 |
| muscle | 1.47 ± 0.20 | 0.70 ± 0.25 | 0.37 ± 0.23 | 0.38 ± 0.06 |
| fat | 3.46 ± 0.59 | 1.03 ± 0.44 | 0.77 ± 0.66 | 0.37 ± 0.13 |
| heart | 2.79 ± 0.35 | 0.94 ± 0.19 | 0.40 ± 0.02 | 0.86 ± 0.26 |
| brain | 1.42 ± 0.33 | 0.20 ± 0.06 | 0.06 ± 0.02 | 0.09 ± 0.03 |
| tumor | 2.91 ± 0.40 | 1.33 ± 0.13 | 0.71 ± 0.12 | 0.82 ± 0.09 |
| ratio | | | | |
| Tumor:bone | 0.42 ± 0.03 | 0.57 ± 0.12 | 0.62 ± 0.09 | 0.26 ± 0.08 |
| Tumor:liver | 0.50 ± 0.07 | 0.98 ± 0.16 | 0.91 ± 0.24 | 0.67 ± 0.14 |
| Tumor:muscle | 2.00 ± 0.37 | 2.06 ± 0.69 | 2.14 ± 0.99 | 2.17 ± 0.23 |

| | | | | |
|-------------|-------------|-------------|-------------|-------------|
| Tumor:fat | 0.84 ± 0.06 | 1.49 ± 0.65 | 1.93 ± 1.87 | 2.44 ± 0.77 |
| Tumor:heart | 1.04 ± 0.08 | 1.44 ± 0.18 | 1.81 ± 0.32 | 1.00 ± 0.27 |

Table IX.

| ¹²⁵ I]6i Biodistribution in EMT-6 BALB/C mice | | | | | |
|--|--------------|--------------|-------------|-------------|-------------|
| %ID per gram | 5 min. | 30 min. | 1 hour | 2 hour | 4 hour |
| blood | 2.37 ± 0.26 | 2.19 ± 0.16 | 1.52 ± 0.49 | 0.65 ± 0.07 | 0.29 ± 0.11 |
| lung | 27.13 ± 1.61 | 5.50 ± 0.62 | 2.12 ± 0.19 | 1.01 ± 0.18 | 0.29 ± 0.05 |
| liver(all) | 13.20 ± 2.04 | 8.77 ± 0.88 | 4.24 ± 0.60 | 1.91 ± 0.20 | 0.90 ± 0.21 |
| kidney | 29.51 ± 2.23 | 13.69 ± 0.35 | 5.94 ± 1.22 | 2.45 ± 0.25 | 0.63 ± 0.11 |
| muscle | 4.10 ± 0.33 | 1.21 ± 0.17 | 0.87 ± 0.22 | 0.31 ± 0.10 | 0.10 ± 0.04 |
| fat | 4.15 ± 0.91 | 1.73 ± 0.20 | 0.74 ± 0.17 | 0.33 ± 0.08 | 0.10 ± 0.05 |
| heart | 6.55 ± 0.54 | 1.94 ± 0.09 | 0.91 ± 0.10 | 0.40 ± 0.05 | 0.15 ± 0.04 |
| brain | 1.53 ± 0.14 | 0.39 ± 0.03 | 0.15 ± 0.03 | 0.06 ± 0.01 | 0.02 ± 0.01 |
| tumor | 4.02 ± 0.55 | 4.50 ± 0.43 | 3.53 ± 0.42 | 1.88 ± 0.76 | 0.82 ± 0.09 |
| ratio | | | | | |
| Tumor:blood | 1.73 ± 0.46 | 2.07 ± 0.32 | 2.44 ± 0.53 | 2.83 ± 1.62 | 3.11 ± 0.85 |
| Tumor:lung | 0.15 ± 0.02 | 0.82 ± 0.09 | 1.66 ± 0.17 | 1.99 ± 1.04 | 2.83 ± 0.19 |
| Tumor:muscle | | | | | |
| e | 0.98 ± 0.13 | 3.94 ± 0.55 | 4.26 ± 0.74 | 6.99 ± 4.44 | 8.98 ± 3.01 |
| Tumor:fat | 1.00 ± 0.20 | 2.62 ± 0.27 | 4.85 ± 0.67 | 5.86 ± 2.93 | 9.59 ± 4.55 |
| Tumor:heart | 0.61 ± 0.07 | 2.32 ± 0.18 | 3.89 ± 0.08 | 4.85 ± 2.31 | 5.91 ± 1.65 |

[00214] E. Blocking studies. The results of the above biodistribution studies in tumor bearing rodents (mice) showed the utility that [¹¹C]10, [⁷⁶Br]6, and [¹²⁵I]6i as potential candidates for further evaluation.

[00215] The inventors then conducted blocking studies in order to confirm that the radiotracer labeled σ_2 receptors in the breast tumors. These studies were conducted using the nonselective sigma ligand, YUN 143 (1 mg/kg, i.v.), which has a high affinity for both σ_1 and σ_2 receptors. We have previously reported that [¹⁸F]YUN 143 labels σ_1 and σ_2 receptors in breast tumor xenografts (Mach et al., 2001). The results of our blocking study are shown in Figure 2 and are consistent with the labeling of σ_2 receptors in vivo.

[00216] F. Comparison with [^{18}F]FLT. One of the specific aims of this project was to compare the σ_2 receptor imaging approach with the nucleoside analog, [^{18}F]FLT. These studies were conducted and the results are summarized in Figures 3-5. In this study, the uptake of [^{18}F]FLT at 1 hr was compared with the one hr data for [^{11}C]Compound 10. For studies comparing [^{18}F]FLT with [^{76}Br]6 and [^{125}I]6i, the two hr post-i.v. injection time point was used.

[00217] G. MicroPET[®] Imaging Studies -- The inventors conducted microPET[®] imaging studies with [^{76}Br]6 in Balb-c mice bearing EMT-6 breast tumor xenografts to model a living mammal afflicted with cancer. The results of the imaging studies are shown in Figure 6. Note the high uptake of the radiotracer in the NCA study (left image), which can be blocked with a known sigma receptor ligand (right image). These data show that [^{76}Br] Compound 6 is a potential radiotracer for imaging the σ_2 receptor status of breast tumors.

[00218] H. Log P Calculations. The lipophilicity of each compound was calculated using the program Clog P (Advanced Chemistry Development, Inc.; Toronto, Canada). The results of the log P calculations are shown in Table X. The log P values ranged from 2.31 to 3.43.

Table X.

| # | Log P |
|----|-------|
| 9 | 2.31 |
| 10 | 2.84 |
| 8 | 3.17 |
| 7 | 3.33 |
| 6 | 3.43 |
| 6i | 3.24 |

[00219] The relationship between the uptake of the radiolabeled benzamide analogs (%I.D. at 5 min post-i.v. injection) and log P is shown in Figure X. There was no clear trend with respect to log P and tumor uptake (top graph) and tumor:fat ratio (bottom graph). The higher tumor:fat ratio of [^{76}Br]6 and [^{125}I]6i was

due to the later time point that can be imaged because of the longer half-life of Br-76 and I-125 versus that of C-11.

[00220] The inventors identified a σ_2 receptor imaging agent that has in vivo properties equal to or greater than that of [^{18}F]FLT. Based on the data described above we have identified three σ_2 radiotracers:

1. [^{11}C]10, which has tumor:lung and tumor:fat ratios equal to [^{18}F]FLT and tumor:blood, tumor:muscle and tumor:heart ratios that are greater than that of [^{18}F]FLT;
2. [^{76}Br]6, which has higher tumor:background ratios greater than that observed with [^{18}F]FLT;
3. [^{125}I]6i, which has a similar tumor:fat ratio as [^{18}F]FLT and exceeds [^{18}F]FLT in all other tumor:background ratios.

[00221] We have specifically identified two potential PET radiotracers ([^{11}C]10 and [^{76}Br]6) and one SPECT radiotracer (^{123}I]6i) for imaging the σ_2 receptor status of breast tumors. In addition, 6i can also be labeled with ^{124}I , potentially yielding another PET radiotracer for imaging σ_2 receptors in breast tumors.

[00222] In conclusion, the inventors have identified and successfully prepared three radiotracers that are either equal to or greater than [^{18}F]FLT in a living rodent model of breast tumor.

[00223] References

[00224] Mach RH, Smith CR, Al-Nabulsi I, Whirrett BR, Childers SR, Wheeler KT. Sigma-2 receptors as potential biomarkers of proliferation in breast cancer. *Cancer Res* 1997; 57:156-161.

[00225] Al-Nabulsi I, Mach RH, Wang L-M, Wallen CA, Keng PC, Sten K, Childers SR, Wheeler KT. Effect of ploidy, recruitment, environmental

factors, and tamoxifen treatment on the expression of sigma-2 receptors in proliferating and quiescent tumor cells. *Br J Cancer* 1999; 81:925-933.


[00226] Wheeler KT, Wang L-M, Wallen CA, Childers SR, Cline JM, Keng PC, Mach RH. Sigma-2 receptors as a biomarker of proliferation in solid tumors. *Br J Cancer* 2000; 86:1223-1232.

[00227] Mach RH, Huang Y, Buchheimer N, Kuhner R, Wu L, Morton TE, Wang L-M, Ehrenkaufer RL, Wallen CA, Wheeler KT. [^{18}F]N-4'-fluorobenzyl-4-(3-bromophenyl)acetamide for imaging the sigma receptor status of shown tumors: comparison with [^{18}F]FDG and [^{125}I]IUDR. *Nucl Med Biol* 2001; 28:451-458.


[00228] While the invention has been described in terms of various specific embodiments, those skilled in the art will recognize that the invention can be practiced with modification within the spirit and scope of the claims.

WHAT IS CLAIMED IS:


1. A benzamide compound having a structure shown in the structure in Formula (I) :



wherein Ar is a substitutable moiety comprising either a substitutable aromatic ring or substitutable heteroaromatic ring; X, Y, Z is a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer ranging from 2 to about 10; and NRR' is at least one of (a):



wherein A and A1 are each moieties substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄), C₁-C₄ alkoxy, fluoroalkyl (C₁-C₄), fluoroalkoxy (C₁-C₄), or one of




wherein B, C are each substitutable moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R1 are two independently selected conjoined and/or linked cyclic aromatic organic


moieties, n is an integer independently varying from 1 to ten and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings, compound 13 and compound 14 and their analogs, salts and their radioactive labeled counterparts thereof..

2. A compound in accordance with Claim 1 having a structure of at least one of:


Compound 1




Compound 2



Compound 3




Compound 4




Compound 4

Compound 5




Compound 5

Compound 6




Compound 6

Compound 7




Compound 7


Compound 8




Compound 9





Compound 10




Compound 11




Compound 12



Compound 13



Compound 14



and its respective radiolabeled counterparts thereof.

3. A compound in accordance with Claim 2 wherein the compound has a structure depicted in the structure of compound 6.


4. A compound in accordance with Claim 3 wherein the compound is radiolabeled with ^{76}Br .


5. A compound in accordance with Claim 3 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.

6. A compound in accordance with Claim 3 wherein the compound has a structure depicted in the structure of compound 10.

7. A compound in accordance with Claim 6 wherein the compound is labeled with ^{11}C .

8. A process for the preparation of a compound selected from one of compounds 1, 2, 3, 4, 5 and 6 of Figure 4 which comprises respectively reacting a compound having a formula wherein R is hydrogen or methoxy






with a compound of the formula

wherein

R is a substitutable moiety selected from hydrogen or methoxy and independently m is an integer from one to 3 with bromoacetonitrile or bromobutyronitrile to produce an



N-alkylated product

wherein when R is hydrogen, n is an integer independently 2-4 and wherein R is independently methoxy then n is an integer independent 2-4 and

(b) reducing the N-alkylated product with lithium aluminum hydride in THF or hydrogenating the N-alkylated compound over palladium on charcoal to provide an intermediate amine product depicted as compound 19, 20, 21 and 22 on Figure 4, and

(c) condensing the intermediate amine product with either 2-methoxy-5-bromonaphthol chloride or 5-bromo-2,3-dimethoxybenzoic acid to produce a compound having as its structure a structure selected from one structures shown for compounds 1-6.

9. A process in accordance with Claim 8 wherein the compound is Compound 1.

10. A process in accordance with Claim 8 wherein the compound is Compound 2.


11. A process in accordance with Claim 8 wherein the compound is Compound 3.

12. A process in accordance with Claim 8 wherein the compound is Compound 4.

13. A process in accordance with Claim 8 wherein the compound is Compound 5.

14. A process in accordance with Claim 8 wherein the compound is Compound 6.

15. A process for the preparation of a compound 7, 8, 9 and 10 of Figure 5 which comprises reacting a compound of the formula



wherein X is a substitutable moiety independently halogen or alkyl C_1 - C_8 with thionyl chloride and one of butyronitrile or bromobutyronitrile to produce compound of 7, 8, 9 and 10 of Fig 5. In an aspect X comprises bromide.

16. In an aspect X comprises at least one of methyl and bromide.

17. A process in accordance with Claim 15 wherein the compound is Compound 7.


18. A process in accordance with Claim 15 wherein the compound is Compound 8.

19. A process in accordance with Claim 15 wherein the compound is Compound 9.


20. A process in accordance with Claim 15 wherein the compound is Compound 10.

21. A process for the preparation of a compound depicted as 11 and 12 which comprises:


(c) reacting a compound of the formula



(d) with bromoacetonitrile to produce a compound of the formula



(d) hydrogenating that compound over palladium on charcoal to provide a compound of the formula




and condensing that compound with either one of 2-methoxy-5-bromonaphthoyl chloride or 5-bromo-2,3-dimethoxybenzoic acid respectively to produce compound 11 and compound 12.


22. A process in accordance with Claim 15 wherein the compound is Compound 11.

23. A process in accordance with Claim 15 wherein the compound is Compound 12.


24. A method for diagnosing a mammal for the presence of a cancer comprises administering to the mammal a diagnostic imaging detectable effective amount of a novel benzamide detectably labeled compound having a structure selected from at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 structurally depicted in Formula (II) and compound 14 structurally depicted in Formula (III) and detecting binding of the at least one benzamide compound to a tumor in the mammal;





where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):



wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



25. A compound in accordance with Claim 24 wherein the compound has a structure depicted in the structure of compound 6.

26. A compound in accordance with Claim 25 wherein compound 6 is radiolabeled with ⁷⁶Br.

27. A compound in accordance with Claim 25 wherein compound 6 is radiolabeled with at least one of ^{123}I , ^{124}I and ^{125}I .


28. A compound in accordance with Claim 24 wherein the compound has a structure depicted in the structure of compound 10.

29. A compound in accordance with Claim 28 wherein compound 10 is labeled with ^{11}C .


30. A method in accordance with Claim 24 which comprises determining that a mammalian tumor is present upon detecting binding.

31. A method in accordance with Claim 24 which further comprises producing an acquisition of the detection.


32. A marker for cancer comprising a detectably-labeled benzamide compound having a structure of at least one of the structures illustratively depicted of benzamide compounds illustratively depicted in Formula (I) including compound 23, compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and having an explicit provocative binding efficacy to a tumor in a living mammal and in an aspect the detectably labeled benzamide compound is a highly selective σ_2 radioligand having the aforementioned structure appended with a radioactive ligand;




where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ alky, $\text{C}_1\text{-C}_4$ fluoroalkyl, $\text{C}_1\text{-C}_4$ fluoroalkoxy, CF_3 , OCF_3 , SCH_3 , SCF_3 , NH_2 ; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and





and Compound 14.

33. A compound in accordance with Claim 32 wherein the compound has a structure depicted in the structure of compound 6.

34. A compound in accordance with Claim 33 wherein compound 6 is radiolabeled with ^{76}Br .

35. A compound in accordance with Claim 32 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.

36. A compound in accordance with Claim 32 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and iodine 125 for invitro measurement of sigma-2 receptors in tumors and normal tissue.


37. A compound in accordance with Claim 32 wherein the compound is radiolabeled with at least one of tritium and iodine 125.

38. Radiolabel compounds of Formula (I), Formula (II) and Formula (III) including compound 23 having at least one of tritium and/or iodine 125 radiolabels using standard radiolabeling conditions to produce at least one of a tritiated and iodine 125 labeled analog that can be used in the invitro detection of sigma 2 receptors in tumors and normal tissue.


39. A compound in accordance with Claim 32 wherein the compound has a structure depicted in the structure of compound 10.

40. A compound in accordance with Claim 36 wherein compound 10 is labeled with ^{11}C .


41. A pharmaceutical composition comprising a benzamide compound having as a structure at least one of structures shown for novel benzamide compounds illustratively depicted in Formula (I) compound, 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and a pharmaceutically acceptable diluent or carrier; conformationally flexible benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :



where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):





wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl,

C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



42. A compound in accordance with Claim 41 wherein the compound has a structure depicted in the structure of compound 6.

43. A compound in accordance with Claim 42 wherein the compound is radiolabeled with ⁷⁶Br.

44. A compound in accordance with Claim 42 wherein the compound is radiolabeled with at least one of ¹²³I, ¹²⁴I, ¹²⁵I and/or tritium.


45. A compound in accordance with Claim 41 wherein the compound has a structure depicted in the structure of compound 10.

46. A compound in accordance with Claim 45 wherein the compound is labeled with ^{11}C .


47. A benzamide composition comprising a detectably labeled benzamide compound further comprising a highly selective σ_2 radioligand.

48. A pharmaceutical composition effective for treating human or non-human neoplastic disorder comprises a detectably labeled pharmaceutically effective amount of at least one compound having a structure of at least one of the structures illustratively depicted in one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) in a composition including a pharmaceutically acceptable carrier.


49. A composition in accordance with Claim 48 when the detectably labeled benzamide compound is a highly selective σ_2 radioligand having the aforementioned structure appended with a radioactive ligand; wherein the benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :




where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



Compound 13



and Compound 14.

50. A compound in accordance with Claim 49 wherein the compound has a structure depicted in the structure of compound 6.

51. A compound in accordance with Claim 50 wherein the compound is radiolabeled with ^{76}Br .


52. A compound in accordance with Claim 50 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I and/or ^{125}I .

53. A compound in accordance with Claim 49 wherein the compound has a structure depicted in the structure of compound 10.


54. A compound in accordance with Claim 53 wherein the compound is labeled with ^{11}C .

55. A method for determining the proliferative status of a cancer cell in a living mammal comprises administering to a living mammal afflicted with a solid malignant tumor, an effective amount of a detectably-labeled benzamide compound having a structure of at least one of the one of the benzamide compounds illustratively depicted in Formula (I) compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and determining the extent to which the detectably-labeled benzamide compound binds to cells of a tumor in the mammal, the extent providing a measure of the proliferative status of the cells.


56. A method in accordance with Claim 55 wherein the living mammal is a human and benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :





where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):



wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and compound 23 and



57. A method in accordance with Claim 56 wherein the benzamide compound has a structure depicted in the structure of compound 6.

58. A method in accordance with Claim 57 wherein the benzamide compound is radiolabeled with ^{76}Br .

59. A method in accordance with Claim 57 wherein the benzamide compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and/or tritium.

60. A method in accordance with Claim 55 wherein the benzamide compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and/or iodine 125 for invitro measurement of these receptors in tumors and normal tissue.

61. A method in accordance with Claim 60 wherein the benzamide compound is radiolabeled with at least one of tritium and/or iodine 125.

62. Radiolabel compounds and compositions of Formula (I), Formula (II) and Formula (III) including compound 23 wherein at least one benzamide compound(s) is radiolabeled with at least one of tritium and/or iodine 125 radiolabels


using standard radiolabeling conditions to produce at least one of a tritiated and iodine 125 labeled analog that can be used in the invitro detection of sigma 2 receptors in tumors and normal tissue.

63. A compound in accordance with Claim 55 wherein the compound has a structure depicted in the structure of compound 10.


64. A compound in accordance with Claim 63 wherein the compound is labeled with ^{11}C .

65. A method in accordance with Claim 64 when determining the proliferative status includes assessing the proliferative status of a breast cancerous tumor.


66. A method for pharmacologically treating a mammalian tumor as a disorder in a mammal comprises administering to a mammal having a tumor a composition including a tumor-inhibiting amount of at least one detectably-labeled benzamide compound having a structure of the structure shown of at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) In an aspect the mammal is a human; novel conformationally flexible benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :




where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and





and Compound 14.

67. A compound in accordance with Claim 66 wherein the compound has a structure depicted in the structure of compound 6.


68. A compound in accordance with Claim 67 wherein the compound is radiolabeled with ^{76}Br .

69. A compound in accordance with Claim 67 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.


70. A compound in accordance with Claim 66 wherein the compound has a structure depicted in the structure of compound 10.

71. A compound in accordance with Claim 70 wherein the compound is labeled with ^{11}C .


72. A method for diagnostic imaging of a mammalian tissue having ample cell surface sigma-2 receptors comprising administering to the tissue of the mammal a diagnostic imaging amount of at least one compound having a structure of at least one benzamide compound illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) and detecting an image of a tissue having an ample cells with sigma-2 receptors and including those compounds having a structure shown in the structure in Formula (I) :



where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):





wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein

Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



73. A compound in accordance with Claim 72 wherein the compound has a structure depicted in the structure of compound 6.

74. A compound in accordance with Claim 73 wherein the compound is radiolabeled with ^{76}Br .

75. A compound in accordance with Claim 73 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium

76. A compound in accordance with Claim 72 wherein the compound has a structure depicted in the structure of compound 10.

77. A compound in accordance with Claim 76 wherein the compound is labeled with ^{11}C .


78. A method for in vitro detection of a cancer cell in living mammalian tissue sample comprising contacting a mammalian tissue sample comprising a cell with an in vitro diagnostic imaging amount of at least one detectably

labeled benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) for a time and under conditions sufficient and effective for binding of the compound to the cell and detecting such binding indicative of an association with the present of cancer.


79. A method in accordance with Claim 78 wherein the mammal is a living human.

80. A method in accordance with Claim 79 wherein the detecting is by image acquisition.


81. A method in accordance with Claim 80 wherein the compound is selected from novel conformationally flexible benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :





where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):



wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and compound 23 and



82. A method for in vitro detection of a cancer cell in a living mammalian tissue sample comprising contacting a mammalian tissue sample with an


in vitro diagnostic imaging detectable and acquisitionable amount of at least one detectably labeled benzamide compound having as a structure a structure of at least one of the benzamide compounds including compound 23 illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) for a time and under cellular conditions functionally sufficient and effective for binding of the compound to the cancer cell and detecting such binding.

83. A compound in accordance with Claim 82 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and/or iodine 125 for in vitro measurement of these receptors in tumors and normal tissue.

84. A compound in accordance with Claim 82 wherein the compound is radiolabeled with at least one of tritium and/or iodine 125.


85. Radiolabel compounds of Formula (I), Formula (II) and Formula (III) including compound 23 are radiolabeled with at least one of tritium and iodine 125 radiolabels using standard radiolabeling conditions to produce at least one of a tritiated and iodine 125 labeled analog that can be used in the in vitro detection of sigma 2 receptors in tumors and normal tissue.

86. Benzamide compounds having a structure shown in the structure in Formula (I) :




where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an


integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and compound 23 and





and Compound 14.

87. A compound in accordance with Claim 86 wherein the compound has a structure depicted in the structure of compound 6.

88. A compound in accordance with Claim 87 wherein the compound is radiolabeled with ^{76}Br .

89. A compound in accordance with Claim 87 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.

90. A compound in accordance with Claim 86 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and iodine 125 for in vitro measurement of these receptors in tumors and normal tissue.

91. A compound in accordance with Claim 86 wherein the compound is radiolabeled with at least one of tritium and iodine 125.


92. A compound in accordance with Claim 86 wherein the compound has a structure depicted in the structure of compound 10.

93. A compound in accordance with Claim 92 wherein the compound is labeled with ^{11}C .


94. A method in accordance with Claim 93 wherein the mammal is a human.

95. A method for determining proliferation and progression of a cancer as a disorder in a living mammal comprising administering to a living mammal a diagnostic imaging detectable amount of at least one detectably-labeled benzamide


compound having as a structure a structure of at least one of the benzamide compounds illustratively structurally depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first selected time, detecting an image of a tissue having ample cells with sigma-2 receptors at a second selected (later) time respectively detecting an image of a tissue having ample cells with sigma-2 receptors at both times, comparing the images and determining if the detected image at the later time is smaller than the detected image at the first time; novel conformationally flexible benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :





where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):



wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



96. A compound in accordance with Claim 95 wherein the compound has a structure depicted in the structure of compound 6.

97. A compound in accordance with Claim 96 wherein the compound is radiolabeled with ⁷⁶Br.

98. A compound in accordance with Claim 96 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I and ^{125}I .

99. A compound in accordance with Claim 96 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and/or iodine 125 for in vitro measurement of these receptors in tumors and normal tissue.

100. A compound in accordance with Claim 96 wherein the compound is radiolabeled with at least one of tritium and/or iodine 125.

101. A compound in accordance with Claim 96 wherein the compound has a structure depicted in the structure of compound 10.


102. A compound in accordance with Claim 101 wherein the compound is labeled with ^{11}C .

103. A method in accordance with Claim 102 wherein the elapsed time between the first time and second time is selected to be a time duration significant amount.


104. A method in accordance with Claim 103 wherein the mammal is a living human.

105. A method for identifying a modulating effect (and regression effect) of a cancer in a living mammal with a disorder, comprising administering to the mammal a diagnostic imaging detectable amount of at least one detectably-labeled benzamide compound having a structure of at least one of the structures illustrated in Formula (I) Formula (II) and Formula (III) at a first time, detecting and acquisitioning an image of a tissue having ample available cells with sigma-2 receptors, administering a detectably-labeled benzamide compound having a structure of at least one of the compound structures illustrated in Formula (I), Formula (II) and Formula (III) to the mammal and at a second (later) time respectively detecting and acquisitioning an image of a tissue having an abundance of cells with sigma receptors, comparing the respective images and determining that there has been an prophylactic


effect and/or regression.; novel conformationally flexible benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :



where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):





wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein,

independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



106. A compound in accordance with Claim 105 wherein the compound has a structure depicted in the structure of compound 6.


107. A compound in accordance with Claim 106 wherein the compound is radiolabeled with ^{76}Br .

108. A compound in accordance with Claim 108 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.


109. A compound in accordance with Claim 106 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and iodine 125 for invitro measurement of these receptors in tumors and normal tissue.

110. A compound in accordance with Claim 106 wherein the compound is radiolabeled with at least one of tritium and/or iodine 125.


111. A compound in accordance with Claim 106 wherein the compound has a structure depicted in the structure of compound 10.
112. A compound in accordance with Claim 111 wherein the compound is labeled with ^{11}C .
113. A method in accordance with Claim 112 wherein the comparison shows the amount of regression over time.
114. A method in accordance with Claim 113 wherein the mammal is a living human.
115. A method in accordance with Claim 114 wherein the comparison shows the prophylactic effect of the compound and its toxicity to cancer.
116. A method in accordance with Claim 114 wherein the comparison shows the efficacy of the compound towards killing cancer in a living mammal and wherein the detectably-labeled benzamide compound is tagged with a radioactive ligand.
117. A method for diagnosing and determining the response of a mammalian patient(s) with a disorder to tailored drug therapy comprising administering to a mammal a diagnostic imaging detectable amount of at least one detectably labeled benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first and second (later) times respectively, detecting an image of a tissue having an abundance of cells with sigma receptors at both times, comparing the images and determining if the image at a later time is larger than the image at the first time that there has been a proliferative effect and progression of cancer; novel conformationally flexible benzamide compounds including those compounds having a structure shown in the structure in Formula (I) :



where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):





wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein

Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and compound 23 and



118. A compound in accordance with Claim 117 wherein the compound has a structure depicted in the structure of compound 6.

119. A compound in accordance with Claim 118 wherein the compound is radiolabeled with ^{76}Br .

120. A compound in accordance with Claim 118 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.

121. A compound in accordance with Claim 117 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and/or iodine 125 for in vitro measurement of these receptors in tumors and normal tissue.

122. A compound in accordance with Claim 117 wherein the compound is radiolabeled with at least one of tritium and/or iodine 125.


123. A compound in accordance with Claim 117 wherein the compound has a structure depicted in the structure shown for compound 10.

124. A compound in accordance with Claim 123 wherein the compound is labeled with ^{11}C .

125. A method in accordance with Claim 117 where the dynamic comparison shows the proliferation and progression of the cancer over elapsed time. In an aspect the mammal is a living human.


126. A method in accordance with Claim 125 where a method for diagnosis and determining the response is a determination of a prophylaxis or management of a disorder associated with neoplastic cells.

127. A method of screening candidate chemicals for toxicity/lethality to cancer comprising administering to a mammal a diagnostic imaging detectable amount of at least one detectably labeled benzamide compound selected from the benzamide compounds illustratively depicted in Formula (I), compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first time, detecting and acquisitioning an image of a tissue having ample cells with sigma-2 receptors, administering to the mammal a candidate chemical, detecting and acquisitioning an image of tissue having ample cells with sigma-2 receptors, comparing the detected images and making a determination as to whether there has been a proliferative effect and progression of the cancer; novel conformationally flexible benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :




where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar


and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



Compound 13



and Compound 14.

128. A compound in accordance with Claim 127 wherein the compound has a structure depicted in the structure of compound 6.

129. A compound in accordance with Claim 128 wherein the compound is radiolabeled with ^{76}Br .

130. A compound in accordance with Claim 128 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium.

131. A compound in accordance with Claim 128 wherein the compound has a structure depicted in the structure (depicted in the structure depicted for compound 10) and is radiolabeled with at least one of tritium and/or iodine 125 for invitro measurement of these receptors in tumors and normal tissue.

132. A compound in accordance with Claim 128 wherein the compound is radiolabeled with at least one of tritium and/or iodine 125.


133. A compound in accordance with Claim 128 wherein the compound has a structure depicted in the structure of compound 10.

134. A compound in accordance with Claim 133 wherein the compound is labeled with ^{11}C .


135. A method in accordance with Claim 134 wherein the mammal is a human.

136. A medical treatment for a mammal and a method of medically treating a mammal comprising administering to a mammal a diagnostic imaging detectable amount of at least one benzamide compound having a structure of at least


one of the benzamide compounds illustratively depicted in Formula (I) compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) at a first and second (later) time respectively, detecting an image of a tissue having ample cells with sigma receptors at the first and second times, comparing the images taken respectively and determining if the image at the later time is larger than the image at the first time that there has been a proliferative effect and progression of the cancer; wherein benzamide compounds include those compounds having a structure shown in the structure in Formula (I) :





where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, C₁-C₄ alky, C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃, NH₂; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):



wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



137. A compound in accordance with Claim 136 wherein the compound has a structure depicted in the structure of compound 6.

138. A compound in accordance with Claim 137 wherein the compound is radiolabeled with ⁷⁶Br.

139. A compound in accordance with Claim 137 wherein the compound is radiolabeled with at least one of ¹²³I, ¹²⁴I, ¹²⁵I and tritium.

140. A compound in accordance with Claim 136 wherein the compound has a structure depicted in the structure depicted for compound 10.


141. A compound in accordance with the compound of Claim 136 wherein the compound is labeled with ^{11}C .

142. A method in accordance with Claim 141 wherein the mammal is a human.


143. A method in accordance with Claim 142 wherein the treatment is a cancer treatment.

144. A method in accordance with Claim 143 wherein medical treatment includes a method of retarding, preventing, and ameliorating disease or a medical affliction in a mammal.


145. A method of customizing drug therapy for a living mammalian subject comprising administering to the subject to the mammal undergoing drug therapy a diagnostic imaging detectable amount of at least benzamide compound having a structure of at least one of the benzamide compounds illustratively depicted in Formula (I) compound 13 depicted in Formula (II) and compound 14 depicted in Formula (III) detecting an image of a tumor having ample sigma-2 receptors and making a determination regarding the drug therapy based on the captured images; having conformationally flexible benzamide compounds including those compounds having a structure shown in the structure in Formula (I) :




where Ar represents a substitutable chemical moiety comprising one of either a substitutable aromatic ring or a substitutable heteroaromatic ring; X, Y, Z represents a moiety substitutable on Ar and selected from the group consisting of H, halogen (I, Br, Cl, F), $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ alky, $\text{C}_1\text{-C}_4$ fluoroalkyl, $\text{C}_1\text{-C}_4$ fluoroalkoxy, CF_3 , OCF_3 , SCH_3 , SCF_3 , NH_2 ; n is an integer independently ranging from 2 to about 10 such as 2, 3, 4, 5, 6, 7, 8, 9, 10; and NRR' is at least one of (a):




wherein A and A₁ are each chemical moieties independently substitutable on Ar' selected from the group consisting of independently H, alkyl (C₁-C₄) or C₁-C₄ alkoxy or one of



wherein B and C are each independent substitutable chemical moieties independently selected from the group consisting of H, halogen (I, Br, Cl, F), C₁-C₄ alkoxy, alkyl (C₁-C₄), C₁-C₄ fluoroalkyl, C₁-C₄ fluoroalkoxy, CF₃, OCF₃, SCH₃, SCF₃ and NH₂, and R and R₁ are two independently selected conjoined and/or linked cyclic aromatic organic moieties, n is an integer the same or different as N in (a) above and N herein, independently varies from 1 to ten such as 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10 and wherein Ar, Ar', Ar'' are independently selected substitutable aromatic rings and their respective radioactive labeled counterparts and



Compound 13



and Compound 14.

146. A compound in accordance with Claim 145 wherein the compound has a structure depicted in the structure of compound 6.

147. A compound in accordance with Claim 146 wherein the compound is radiolabeled with ^{76}Br .

148. A compound in accordance with Claim 146 wherein the compound is radiolabeled with at least one of ^{123}I , ^{124}I , ^{125}I and tritium


149. A compound in accordance with Claim 146 wherein the compound has a structure depicted in the structure (depicted in compound 10) and is radiolabeled with at least one of tritium and/or iodine 125 for invitro measurement of these receptors in tumors and normal tissue.

150. A compound in accordance with Claim 149 wherein the compound is radiolabeled with at least one of tritium and/or iodine 125.

151. A compound in accordance with Claim 146 wherein the compound has a structure depicted in the structure of compound 10.

152. A compound in accordance with Claim 151 wherein the compound is labeled with ^{11}C .


153. A compound having a structure depicted as



Compound 23

154. A compound in accordance with Claim 153 wherein the compound is radiolabeled.

155. A process for the preparation of a compound having a structure depicted as compound 23, which comprises reacting the corresponding phenol with 1-bromo-2-fluoroethane.



Compound 23

156. A compound in accordance with Claim 1 having ¹²³iodine associated therewith as a radiolabel.

157. A compound in accordance with Claim 1 having ¹²⁴Iodine associated therewith as a radiolabel.

158. A compound in accordance with Claim 1 having ¹²⁵Iodine associated therewith as a radiolabel.

159. A compound in accordance with Claim 1 having Carbon 11 associated therewith as a radiolabel.

160. A compound in accordance with Claim 1 having tritium associated therewith as a radiolabel.










Figure 1

**Compound 7****Compound 8****Compound 9****Compound 10****Figure 2**

**Figure 3**




Figure 4. (a) BrCH₂CN or BrCH₂CH₂CH₂CN/Et₃N/CH₂Cl₂/RT; (b) LiAlH₄/THF or H₂/Pd(c)/ethanol; (c) Ref 10; (d) Ref 12.




Figure 5. (a) SOCl₂/benzene/reflux; (b) **19**/Et₃N/CH₂Cl₂/RT; (c) **21**/Et₃N/CH₂Cl₂/RT; (d) **22**/CH₂Cl₂/Et₃N/RT.




Figure 6. (a) $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CN}/\text{Et}_3\text{N}/\text{CH}_2\text{Cl}_2$; (b) H_2/Pd (c)/ethanol; (c) Figure 5.



**Figure 7.**

Table IA. Binding Affinities for Dopamine D₂/D₃ and Sigma σ_1 / σ_2 Receptors

| # | K _i (nM) ^a | | | |
|----|----------------------------------|-----------------------------|-------------------------|-------------------------|
| | D ₂ ^b | D ₃ ^c | σ_1 ^d | σ_2 ^e |
| 1 | 429.7 ± 76.1 | 17.8 ± 0.5 | 276.5 ± 35.7 | 716.5 ± 9.8 |
| 2 | 714.0 ± 133.7 | 21.4 ± 2.3 | 2,932 ± 28 | 16.4 ± 2.0 |
| 3 | 131.6 ± 24.6 | 81.6 ± 21.28 | 15.1 ± 1.7 | 47.7 ± 2.5 |
| 4 | 240.5 ± 19.4 | 126.5 ± 42.4 | 189.1 ± 2.6 | 21.2 ± 0.1 |
| 5 | 741.0 ± 287.3 | 106.5 ± 24.3 | 1,159 ± 7 | 17.6 ± 0.7 |
| 6 | 2,200 ± 390 | 627 ± 244 | 12,900 ± 111 | 8.2 ± 1.4 |
| 7 | 2,190 ± 351 | 310.7 ± 54.4 | 21.8 ± 5.6 | 89.4 ± 13.9 |
| 8 | 3,570 ± 796 | 488.0 ± 70.7 | 5,484 ± 266 | 12.4 ± 1.8 |
| 9 | 2,850 ± 316 | 3,760 ± 618 | 10,412 ± 462 | 13.3 ± 0.1 |
| 10 | 642.0 ± 141.0 | 313.0 ± 141.0 | 3,078 ± 87 | 10.3 ± 1.5 |
| 11 | 58.8 ± 13.7 | 2.1 ± 0.4 | 809 ± 66 | 75.0 ± 4.1 |
| 12 | 107.0 ± 19.0 | 10.2 ± 5.3 | 751 ± 6 | 26.4 ± 1.4 |
| 23 | N/A | N/A | 330.17±24.5 | 6.95±1.63 |

^A MEAN ± SEM, K_i VALUES WERE DETERMINED





Figure 8. In vivo blocking studies of [^{125}I]6i (left), [^{76}Br]6 (center), and [^{11}C]10(right) in tumor-bearing mice. The blocking agent was YUN 143, which has a high affinity for σ_1 and σ_2 receptors. Animals were sacrificed at the time point displaying the highest %I.D./gram tumor. The data are consistent with the labeling of σ_2 receptors in vivo by each radiotracer.

Uptake comparison for FLT vs. [C-11]10 in EMT-6 BALB/C mice



Tumor to background for FLT vs. [C-11] 10 in EMT-6 BALB/C mice




Figure 9. Comparison of [^{18}F]FLT and [^{11}C]10. Although [^{18}F]FLT has a high uptake in tumors (top graph), the high uptake of radioactivity in normal tissues results in a lower tumor:background ratio of [^{18}F]FLT relative to [^{11}C]10, particularly the tumor : muscle, tumor : blood, and tumor : heart ratios (bottom graph).




Figure 10 Comparison of [^{18}F]FLT with [^{76}Br]6 at 2 hrs post-i.v. injection. Notice the higher tumor:background ratios of [^{76}Br]6 versus that of [^{18}F]FLT.





Figure 11. Comparison of [18 F]FLT with [125 I]6i at 2 hrs post-i.v. injection. Notice the higher tumor:blood, tumor:muscle, and tumor:heart ratios of [125 I]6i versus that of [18 F]FLT




FIGURE 12. MICROPET[®] IMAGING STUDIES OF [⁷⁶BR]6 IN BALB-C MICE BEARING AN EMT-6 BREAST TUMOR XENOGRAFT. THE NO-CARRIER-ADDED (NCA) STUDY IS ON THE LEFT AND THE SIGMA RECEPTOR BLOCKING STUDY (1 MG/KG, I.V. YUN-143) IS SHOWN ON THE RIGHT.

14/14




Figure 13. Graphs comparing the uptake of the benzamide analogs with % uptake in breast tumors at 5 min (top) and the tumor:fat ratio at the optimal imaging time (bottom). There was no correlation between tumor uptake or tumor:fat ratio and log P. Note that all compounds had a high tumor uptake at 5 min post-injection. The high tumor:fat ratio of [⁷⁶Br]6 and [¹²⁵I]6i was due to the later time point after injection used because of the longer half-life of Br-76 and I-125 versus that of C-11.