



(51) International Patent Classification:

A01K 67/027 (2006.01) C12N 15/90 (2006.01)

(21) International Application Number:

PCT/US2018/044615

(22) International Filing Date:

31 July 2018 (31.07.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/539,275 31 July 2017 (31.07.2017) US

(71) Applicant: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).(72) Inventors: **GONG, Guochun**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **HUNT, Charleen**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **KAJIMURA, Daisuke**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **HARTFORD, Suzanne**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **ZAMBROWICZ, Brian**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).(74) Agent: **SCHILLING, Stephen H.** et al.; Alston & Bird LLP, Bank of America Plaza, 101 South Tryon Street, Suite 4000, Charlotte, North Carolina 28280-4000 (US).(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: CAS-TRANSGENIC MOUSE EMBRYONIC STEM CELLS AND MICE AND USES THEREOF




FIG. 1

(57) Abstract: Methods and compositions are provided herein for assessing CRISPR/Cas-mediated non-homologous end joining (NHEJ) activity and/or CRISPR/Cas-induced recombination of a target genomic locus with an exogenous donor nucleic acid in vivo and ex vivo. The methods and compositions employ cells and non-human animals comprising a Cas expression cassette such as a genomically integrated Cas expression cassette so that the Cas protein can be constitutively available or available in a tissue- specific or temporal- specific manner. Methods and compositions are also provided for making and using these non-human animals, including use of these non-human animals to assess CRISPR/Cas activity in vivo via adeno-associated virus (AAV)-mediated delivery of guide RNAs to the non-human animals.



CAS-TRANSGENIC MOUSE EMBRYONIC STEM CELLS AND MICE AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of US Application No. 62/539,275, filed July 31, 2017, which is herein incorporated by reference in its entirety for all purposes.

REFERENCE TO A SEQUENCE LISTING

SUBMITTED AS A TEXT FILE VIA EFS WEB

[0002] The Sequence Listing written in file 516569SEQLIST.txt is 178 kilobytes, was created on July 30, 2018, and is hereby incorporated by reference.

BACKGROUND

[0003] CRISPR/Cas technology is a promising new therapeutic modality. However, there is a need for better means of assessing the efficiency of mutation generation or targeted gene modification by an introduced CRISPR/Cas agent *in vivo*. One limitation of testing the system *in vivo* is the need to simultaneously introduce all components into a living organism. The typical method of intruding these components is to transiently transfect DNA constructs into cells that will generate the appropriate RNAs and protein. Though effective, this approach has an inherent disadvantage as the cells must rely on the plasmid DNA constructs to first undergo transcription and then translation before the Cas9 protein is available to interact with the sgRNA component. Better methods and tools are needed to more effectively assess the activity of introduced CRISPR/Cas agents and to assess different delivery methods and parameters for targeting specific tissues or cell types *in vivo*.

[0004] In addition, the delivery of biologically active agents such as CRISPR/Cas agents to subjects is often hindered by difficulties in the components reaching the target cell or tissue. These restrictions can result, for example, in the need to use much higher concentrations of the agents than is desirable to achieve a result, which increases the risk of toxic effects and side effects. Improved delivery methods and methods of assessing such delivery methods *in vivo* are needed.

SUMMARY

[0005] Cas9-ready non-human animals are provided, and methods and compositions are provided for assessing the ability of CRISPR/Cas nuclease agents to modify a target genomic locus *in vivo*. In one aspect, provided are methods of testing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*. Some such methods comprise: (a) introducing into a non-human animal a guide RNA designed to target a guide RNA target sequence at the target genomic locus, wherein the non-human animal comprises a genomically integrated Cas expression cassette comprising an NLS-Cas coding sequence, and wherein the guide RNA is introduced via adeno-associated virus (AAV)-mediated delivery; and (b) assessing the modification of the target genomic locus. Some such methods comprise: (a) introducing into a non-human animal a guide RNA designed to target a guide RNA target sequence at the target genomic locus, wherein the non-human animal comprises a genomically integrated Cas expression cassette comprising an NLS-Cas coding sequence, and wherein the guide RNA is introduced via lipid nanoparticle (LNP)-mediated delivery; and (b) assessing the modification of the target genomic locus.

[0006] In some such methods, the AAV is AAV7, AAV8, or AAV9, and step (b) comprises assessing modification of the target genomic locus in the liver. Optionally, the AAV is AAV8.

[0007] In some such methods, the route of administration of the AAV to the non-human animal is intravenous injection, intraparenchymal injection, intraperitoneal injection, nasal installation, or intravitreal injection.

[0008] In some such methods, an exogenous donor nucleic acid is introduced in step (a), wherein the exogenous donor nucleic acid is designed to recombine with the target genomic locus. Optionally, the exogenous donor nucleic acid is a single-stranded oligodeoxynucleotide (ssODN).

[0009] In some such methods, the non-human animal is a rat or mouse. Optionally, the non-human animal is a mouse.

[0010] In some such methods, the target genomic locus comprises a target gene, and step (b) comprises measuring expression of the target gene or activity of a protein encoded by the target gene.

[0011] In some such methods, step (b) comprises sequencing the target genomic locus in one or more cells isolated from the non-human animal.

[0012] In some such methods, step (b) comprises isolating a target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the target organ or tissue. Optionally, step (b) comprises assessing modification of the target genomic locus in two or more different cell types within the target organ or tissue.

[0013] In some such methods, step (b) comprises isolating a non-target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the non-target organ or tissue.

[0014] In some such methods, the NLS-Cas coding sequence is an NLS-Cas9 coding sequence.

[0015] In some such methods, the Cas expression cassette further comprises a polyadenylation signal upstream of the NLS-Cas coding sequence, wherein the polyadenylation signal is flanked by recombinase recognition sites, and wherein the polyadenylation signal in the Cas expression cassette has been excised in a tissue-specific manner. Optionally, the polyadenylation signal upstream of the NLS-coding sequence in the Cas expression cassette has been excised in the liver. Optionally, the recombinase that recognizes the recombinase recognition sites in the Cas expression cassette is a Cre recombinase. Optionally, the non-human animal further comprises a genomically integrated Cre recombinase expression cassette, wherein the Cre recombinase expression cassette comprises a Cre recombinase coding sequence operably linked to a tissue-specific promoter. Optionally, the Cre recombinase gene is operably linked to one of the promoters set forth in Table 2.

[0016] In some such methods, the Cas expression cassette further comprises a polyadenylation signal upstream of the NLS-Cas coding sequence, wherein the polyadenylation signal is flanked by recombinase recognition sites, and wherein the method further comprises introducing a recombinase into the non-human animal in a tissue-specific manner. Optionally, the recombinase is introduced via adeno-associated virus (AAV)-mediated delivery or lipid nanoparticle (LNP)-mediated delivery. Optionally, the recombinase is introduced via AAV8-mediated delivery. Optionally, the recombinase is introduced into the liver.

[0017] In some such methods, the Cas expression cassette further comprises a fluorescent protein coding sequence. Optionally, the Cas expression cassette comprises a multicistronic nucleic acid comprising the NLS-Cas coding sequence and the fluorescent protein coding sequence separated by an intervening internal ribosome entry site (IRES) or an intervening 2A

peptide coding sequence. Optionally, the multicistronic nucleic acid in the Cas expression cassette comprises the NLS-Cas coding sequence and a green fluorescent protein coding sequence separated by an intervening P2A peptide coding sequence. In some such methods, the Cas expression cassette does not further comprise a fluorescent protein coding sequence. In some such methods, the NLS-Cas coding sequence encodes a Cas protein comprising a protein tag.

[0018] In some such methods, the Cas expression cassette is operably linked to an endogenous promoter. In some such methods, the Cas expression cassette is operably linked to an exogenous, constitutive promoter.

[0019] In some such methods, the 5' end of the Cas expression cassette further comprises a 3' splicing sequence.

[0020] In some such methods, the Cas expression cassette encodes a protein comprising the sequence set forth in SEQ ID NO: 13, 16, 19, or 22. Optionally, the Cas expression cassette comprises the sequence set forth in SEQ ID NO: 28, 29, 30, or 31. Optionally, the Cas expression cassette comprises the sequence set forth in SEQ ID NO: 1, 12, 14, 15, 17, 18, 20, or 21.

[0021] In some such methods, the Cas expression cassette is integrated at a safe harbor locus. Optionally, the safe harbor locus is a *Rosa26* locus. Optionally, the Cas expression cassette is integrated into the first intron of the *Rosa26* locus.

[0022] In some such methods, the non-human animal is heterozygous for the Cas expression cassette. In some such methods, the non-human animal is homozygous for the Cas expression cassette.

[0023] In some such methods, the non-human animal is a mouse, the AAV is an AAV8, the Cas expression cassette is operably linked to the endogenous *Rosa26* promoter, is inserted into the first intron of the *Rosa26* locus, and comprises from 5' to 3': (i) a 3' splicing sequence; and (ii) an NLS-Cas9 coding sequence, and step (b) comprises assessing modification of the target genomic locus in the liver of the non-human animal. In some such methods, the non-human animal is a mouse, the AAV is an AAV8 delivered to the non-human animal by intravenous injection, the Cas expression cassette is operably linked to the endogenous *Rosa26* promoter, is inserted into the first intron of the *Rosa26* locus, and comprises from 5' to 3': (i) a 3' splicing sequence; and (ii) an NLS-Cas9 coding sequence, and step (b) comprises assessing modification

of the target genomic locus in the liver of the non-human animal.

[0024] In another aspect, provided are methods of optimizing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*. Some such methods comprises: (I) performing the any of the above methods of testing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo* a first time in a first non-human animal; (II) changing a variable and performing the method of step (I) a second time with the changed variable in a second non-human animal; and (III) comparing the modification of the target genomic locus in step (I) with the modification of the target genomic locus in step (II), and selecting the method resulting in the modification of the target genomic locus with one or more of higher efficacy, higher precision, higher consistency, or higher specificity.

[0025] In some such methods, the changed variable in step (II) is the AAV serotype. In some such methods, the changed variable in step (II) is the route of administration of introducing the guide RNA into the non-human animal. In some such methods, the changed variable in step (II) is the concentration or amount of the guide RNA introduced into the non-human animal. In some such methods, the changed variable in step (II) is the guide RNA (e.g., the form or sequence of the guide RNA) introduced into the non-human animal. In some such methods, the method comprises introducing an exogenous donor nucleic acid, and wherein the changed variable in step (II) is the delivery method of introducing the exogenous donor nucleic acid into the non-human animal. In some such methods, the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the route of administration of introducing the exogenous donor nucleic acid into the non-human animal. In some such methods, the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the concentration or amount of the exogenous donor nucleic acid introduced into the non-human animal. In some such methods, the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the concentration or amount of the guide RNA introduced into the non-human animal relative to the concentration or amount of exogenous donor nucleic acid introduced into the non-human animal. In some such methods, the changed variable in step (II) is the exogenous donor nucleic acid (e.g., the form of exogenous donor nucleic acid) introduced into the non-human animal.

BRIEF DESCRIPTION OF THE FIGURES

[0026] **Figure 1** shows a Cas9 allele (MAID2599; not to scale), comprising from 5' to 3': a 3' splicing sequence; a first loxP site, a neomycin resistance gene; a polyadenylation signal; a second loxP site; an NLS-Cas9 coding sequence; a P2A peptide coding sequence; and a GFP coding sequence.

[0027] **Figure 2A** shows NHEJ activity in wild type F1H4 mouse embryonic stem cells (mESCs) and Cas9-ready mESCs with and without the lox-stop-lox neomycin cassette (MAID2599 and MAID2600, respectively) following introduction of a two sgRNAs (in plasmid form or as RNAs) targeting the start and stop codon regions of a first target gene, optionally in combination with introduction of a Cas9 plasmid. 5' cutting efficiency was measured in the left panel, 3' cutting efficiency was measured in the middle panel, the rate in which the intervening DNA was deleted completely was measured in the right panel.

[0028] **Figure 2B** shows cutting efficiency (left panel) and HDR efficiency (right panel) following introduction of an sgRNA (in plasmid form or as RNA) targeting a second target gene along with a single-stranded oligodeoxynucleotide (ssODN) as a point mutation donor, optionally in combination with a Cas9 plasmid.

[0029] **Figures 3A-3F** show bright field images of liver (**Figure 3A**), kidney (**Figure 3B**), and brain (**Figure 3C**) tissues from wild type mice and heterozygous Cas9-ready mice (MAID2600), and GFP fluorescence images of liver (**Figure 3D**), kidney (**Figure 3E**), and brain (**Figure 3F**) tissues in wild type mice and Cas9-ready mice (MAID2600).

[0030] **Figure 4A** shows Cas9 mRNA expression levels in various tissues isolated from heterozygous Cas9-ready mice (MAID2600) as determined by RT-qPCR. The y-axis shows the delta Ct + 1 compared to the average Cas9 Ct from brain tissue.

[0031] **Figure 4B** shows Cas9 protein expression in various tissues isolated from wild-type mice and heterozygous Cas9-ready mice (MAID2600). Actin was used as a control.

[0032] **Figure 4C** shows Cas9 average Cas9 and beta-2-microglobulin (B2m) mRNA expression levels in various tissues isolated from heterozygous Cas9-ready mice (MAID2600) as determined by RT-qPCR. The number of samples tested from each type of tissue is indicated above the bars.

[0033] **Figures 5A-5B** show percent NHEJ activity (indel frequency) at a third target gene (target gene 3) in primary hepatocytes isolated from wild type mice (**Figure 5A**) and cassette-

deleted Cas9 mice (MAID2600; **Figure 5B**) following lipid nanoparticle (LNP) delivery of either GFP mRNA and a control (dead) sgRNA, GFP mRNA and a target gene 3 sgRNA, or Cas9 mRNA and a target gene 3 sgRNA. mRNA concentrations of 15.6, 62.5, 250, and 1000 ng/mL were tested.

[0034] **Figures 6A-6D** show serum levels of a protein that is secreted by the liver and found in serum and is encoded by the third target gene (target gene 3) following introduction of a target gene 3 sgRNA into wild type mice (msCas9⁻) or cassette-deleted Cas9-ready mice (msCas9⁺; MAID2600) via hydrodynamic DNA delivery (HDD), lipid nanoparticle (LNP) delivery, or adeno-associated virus (AAV) delivery by tail vein injection. In some cases, Cas9 was also introduced (in mRNA form for LNP delivery, and in DNA form for HDD (Cas9 plasmid) and AAV delivery). Untreated mice, LNP control mice, AAV control mice, and HDD control mice were used as negative controls. For LNP-mediated delivery, three groups of mice were tested: (1) Cas9-ready mice (3 male + 3 female; 2 mg/kg control guide RNA + GFP mRNA); (2) Cas9-ready mice (3 male + 3 female; 2 mg/kg guide RNA for target gene 3 + GFP mRNA); and (3) WT mice (3 male + 3 female; 2 mg/kg guide RNA for target gene 3 + Cas9 mRNA). For AAV-mediated delivery, two groups of mice were tested: (1) Cas9-ready mice (3 male + 3 female; AAV8-guide RNA for target gene 3); and (2) WT mice (3 male + 3 female; AAV8-guide RNA for target gene 3 + AAV8-Cas9). For HDD, two groups of mice were tested: (1) Cas9-ready mice (3 male + 3 female; guide RNA for target gene 3); and (2) WT mice (3 male + 3 female; guide RNA for target gene 3 + Cas9). Serum levels of the protein encoded by target gene 3 were measured in male mice (**Figures 6A** and **6B**) and female mice (**Figures 6C** and **6D**) and were measured at Day 7 (**Figures 6A** and **6C**) and Day 21 (**Figures 6B** and **6D**).

[0035] **Figure 7** shows percent NHEJ activity (indel frequency) at the target gene 3 locus in liver in wild type mice (msCas9⁻) and cassette-deleted Cas9 mice (msCas9⁺; MAID2600) one month after lipid nanoparticle (LNP) delivery of sgRNA alone or together with Cas9 mRNA, hydrodynamic delivery (HDD) of sgRNA plasmid alone or together with Cas9 plasmid, or AAV8-sgRNA alone or together with AAV8-Cas9.

[0036] **Figure 8A** shows percent NHEJ activity (indel frequency) at the target gene 4 locus in liver in cassette-deleted Cas9 mice (MAID2600) 3-4 weeks after AAV8 delivery of sgRNA by tail vein injection. UNT = untreated control.

[0037] **Figure 8B** shows relative levels of target gene 4 expression as determined by

TAQMAN analysis in liver tissue isolated from in cassette-deleted Cas9 mice (MAID2600) 3-4 weeks after AAV8 delivery of sgRNA by tail vein injection. WT mastermix refers to all five sgRNA viruses mixed together and injected into wild type mice as a negative control.

[0038] **Figure 9** shows a western blot of Cas9 expression in LSL-Cas9 mice (MAID2599) in liver, spleen, and kidney samples isolated one week after LNP-Cre was injected via tail vein injection. Mice without LNP-Cre injections were used as a negative control. Cassette-deleted Cas9 mice (MAID2600) were used as a positive control.

[0039] **Figure 10** shows serum levels of a protein that is secreted by the liver and found in serum and is encoded by the third target gene (target gene 3) 1 week and 3 weeks following injection of a target gene 3 sgRNA into LSL-Cas9 mice (MAID2599) via AAV8, either alone or together with LNP-Cre. Mice with neither LNP-Cre nor AAV8-gRNA were used as a negative control. All conditions were also tested in cassette-deleted mice (ROSA Cas9; MAID2600).

[0040] **Figure 11** shows percent NHEJ activity (indel frequency) at the target gene 3 locus in livers isolated 3 weeks following injection of a target gene 3 sgRNA into LSL-Cas9 mice (MAID2599) via AAV8, either alone or together with LNP-Cre. Mice with neither LNP-Cre nor AAV8-gRNA were used as a negative control. All conditions were also tested in cassette-deleted mice (ROSA Cas9; MAID2600).

[0041] **Figure 12A** shows a western blot for Cas9 in livers isolated from LSL-Cas9 mice (MAID2599) and LSL-Cas9/Alb-Cre mice. Actin was used as a loading control.

[0042] **Figure 12B** shows a western blot for Cas9 in brains isolated from LSL-Cas9 mice (MAID2599) and LSL-Cas9/Alb-Cre mice. Actin was used as a loading control.

[0043] **Figure 13** shows serum levels of a protein that is secreted by the liver and found in serum and is encoded by the third target gene (target gene 3) 1 week following injection of a target gene 3 sgRNA into WT mice, cassette-deleted Cas9 mice (ROSA Cas9; MAID2600), LSL-Cas9 mice (MAID2599), albumin-Cre mice, or LSL-Cas9/Alb-Cre mice via AAV8.

[0044] **Figure 14** shows four different Cas9 alleles (not to scale), including the MAID2599 allele (MAID2600 once the lox-stop-lox (LSL) cassette is deleted), the MAID2658 allele (MAID2659 once the LSL cassette is deleted), the MAID2660 allele (MAID2661 once the LSL cassette is deleted), and the MAID2672 allele (MAID2673 once the LSL cassette is deleted).

DEFINITIONS

[0045] The terms “protein,” “polypeptide,” and “peptide,” used interchangeably herein, include polymeric forms of amino acids of any length, including coded and non-coded amino acids and chemically or biochemically modified or derivatized amino acids. The terms also include polymers that have been modified, such as polypeptides having modified peptide backbones.

[0046] Proteins are said to have an “N-terminus” and a “C-terminus.” The term “N-terminus” relates to the start of a protein or polypeptide, terminated by an amino acid with a free amine group (-NH₂). The term “C-terminus” relates to the end of an amino acid chain (protein or polypeptide), terminated by a free carboxyl group (-COOH).

[0047] The terms “nucleic acid” and “polynucleotide,” used interchangeably herein, include polymeric forms of nucleotides of any length, including ribonucleotides, deoxyribonucleotides, or analogs or modified versions thereof. They include single-, double-, and multi-stranded DNA or RNA, genomic DNA, cDNA, DNA-RNA hybrids, and polymers comprising purine bases, pyrimidine bases, or other natural, chemically modified, biochemically modified, non-natural, or derivatized nucleotide bases.

[0048] Nucleic acids are said to have “5’ ends” and “3’ ends” because mononucleotides are reacted to make oligonucleotides in a manner such that the 5’ phosphate of one mononucleotide pentose ring is attached to the 3’ oxygen of its neighbor in one direction via a phosphodiester linkage. An end of an oligonucleotide is referred to as the “5’ end” if its 5’ phosphate is not linked to the 3’ oxygen of a mononucleotide pentose ring. An end of an oligonucleotide is referred to as the “3’ end” if its 3’ oxygen is not linked to a 5’ phosphate of another mononucleotide pentose ring. A nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5’ and 3’ ends. In either a linear or circular DNA molecule, discrete elements are referred to as being “upstream” or 5’ of the “downstream” or 3’ elements.

[0049] The term “genomically integrated” refers to a nucleic acid that has been introduced into a cell such that the nucleotide sequence integrates into the genome of the cell. Any protocol may be used for the stable incorporation of a nucleic acid into the genome of a cell.

[0050] The term “expression vector” or “expression construct” refers to a recombinant nucleic acid containing a desired coding sequence operably linked to appropriate nucleic acid

sequences necessary for the expression of the operably linked coding sequence in a particular host cell or organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, as well as other sequences. Eukaryotic cells are generally known to utilize promoters, enhancers, and termination and polyadenylation signals, although some elements may be deleted and other elements added without sacrificing the necessary expression.

[0051] The term “targeting vector” refers to a recombinant nucleic acid that can be introduced by homologous recombination, non-homologous-end-joining-mediated ligation, or any other means of recombination to a target position in the genome of a cell.

[0052] The term “viral vector” refers to a recombinant nucleic acid that includes at least one element of viral origin and includes elements sufficient for or permissive of packaging into a viral vector particle. The vector and/or particle can be utilized for the purpose of transferring DNA, RNA, or other nucleic acids into cells either *ex vivo* or *in vivo*. Numerous forms of viral vectors are known.

[0053] The term “isolated” with respect to proteins, nucleic acids, and cells includes proteins, nucleic acids, and cells that are relatively purified with respect to other cellular or organism components that may normally be present *in situ*, up to and including a substantially pure preparation of the protein, nucleic acid, or cell. The term “isolated” also includes proteins and nucleic acids that have no naturally occurring counterpart or proteins or nucleic acids that have been chemically synthesized and are thus substantially uncontaminated by other proteins or nucleic acids. The term “isolated” also includes proteins, nucleic acids, or cells that have been separated or purified from most other cellular components or organism components with which they are naturally accompanied (e.g., other cellular proteins, nucleic acids, or cellular or extracellular components).

[0054] The term “wild type” includes entities having a structure and/or activity as found in a normal (as contrasted with mutant, diseased, altered, or so forth) state or context. Wild type genes and polypeptides often exist in multiple different forms (e.g., alleles).

[0055] The term “endogenous sequence” refers to a nucleic acid sequence that occurs naturally within a cell or non-human animal. For example, an endogenous *Rosa26* sequence of a non-human animal refers to a native *Rosa26* sequence that naturally occurs at the *Rosa26* locus in the non-human animal.

[0056] “Exogenous” molecules or sequences include molecules or sequences that are not normally present in a cell in that form. Normal presence includes presence with respect to the particular developmental stage and environmental conditions of the cell. An exogenous molecule or sequence, for example, can include a mutated version of a corresponding endogenous sequence within the cell, such as a humanized version of the endogenous sequence, or can include a sequence corresponding to an endogenous sequence within the cell but in a different form (i.e., not within a chromosome). In contrast, endogenous molecules or sequences include molecules or sequences that are normally present in that form in a particular cell at a particular developmental stage under particular environmental conditions.

[0057] The term “heterologous” when used in the context of a nucleic acid or a protein indicates that the nucleic acid or protein comprises at least two segments that do not naturally occur together in the same molecule. For example, the term “heterologous,” when used with reference to segments of a nucleic acid or segments of a protein, indicates that the nucleic acid or protein comprises two or more sub-sequences that are not found in the same relationship to each other (e.g., joined together) in nature. As one example, a “heterologous” region of a nucleic acid vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid vector could include a coding sequence flanked by sequences not found in association with the coding sequence in nature. Likewise, a “heterologous” region of a protein is a segment of amino acids within or attached to another peptide molecule that is not found in association with the other peptide molecule in nature (e.g., a fusion protein, or a protein with a tag). Similarly, a nucleic acid or protein can comprise a heterologous label or a heterologous secretion or localization sequence.

[0058] “Codon optimization” takes advantage of the degeneracy of codons, as exhibited by the multiplicity of three-base pair codon combinations that specify an amino acid, and generally includes a process of modifying a nucleic acid sequence for enhanced expression in particular host cells by replacing at least one codon of the native sequence with a codon that is more frequently or most frequently used in the genes of the host cell while maintaining the native amino acid sequence. For example, a nucleic acid encoding a Cas9 protein can be modified to substitute codons having a higher frequency of usage in a given prokaryotic or eukaryotic cell, including a bacterial cell, a yeast cell, a human cell, a non-human cell, a mammalian cell, a

rodent cell, a mouse cell, a rat cell, a hamster cell, or any other host cell, as compared to the naturally occurring nucleic acid sequence. Codon usage tables are readily available, for example, at the “Codon Usage Database.” These tables can be adapted in a number of ways. *See Nakamura et al. (2000) Nucleic Acids Research 28:292*, herein incorporated by reference in its entirety for all purposes. Computer algorithms for codon optimization of a particular sequence for expression in a particular host are also available (*see, e.g., Gene Forge*).

[0059] A “promoter” is a regulatory region of DNA usually comprising a TATA box capable of directing RNA polymerase II to initiate RNA synthesis at the appropriate transcription initiation site for a particular polynucleotide sequence. A promoter may additionally comprise other regions which influence the transcription initiation rate. The promoter sequences disclosed herein modulate transcription of an operably linked polynucleotide. A promoter can be active in one or more of the cell types disclosed herein (e.g., a eukaryotic cell, a non-human mammalian cell, a human cell, a rodent cell, a pluripotent cell, a one-cell stage embryo, a differentiated cell, or a combination thereof). A promoter can be, for example, a constitutively active promoter, a conditional promoter, an inducible promoter, a temporally restricted promoter (e.g., a developmentally regulated promoter), or a spatially restricted promoter (e.g., a cell-specific or tissue-specific promoter). Examples of promoters can be found, for example, in WO 2013/176772, herein incorporated by reference in its entirety for all purposes.

[0060] A constitutive promoter is one that is active in all tissues or particular tissues at all developing stages. Examples of constitutive promoters include the human cytomegalovirus immediate early (hCMV), mouse cytomegalovirus immediate early (mCMV), human elongation factor 1 alpha (hEF1 α), mouse elongation factor 1 alpha (mEF1 α), mouse phosphoglycerate kinase (PGK), chicken beta actin hybrid (CAG or CBh), SV40 early, and beta 2 tubulin promoters.

[0061] Examples of inducible promoters include, for example, chemically regulated promoters and physically-regulated promoters. Chemically regulated promoters include, for example, alcohol-regulated promoters (e.g., an alcohol dehydrogenase (alcA) gene promoter), tetracycline-regulated promoters (e.g., a tetracycline-responsive promoter, a tetracycline operator sequence (tetO), a tet-On promoter, or a tet-Off promoter), steroid regulated promoters (e.g., a rat glucocorticoid receptor, a promoter of an estrogen receptor, or a promoter of an ecdysone receptor), or metal-regulated promoters (e.g., a metalloprotein promoter). Physically regulated

promoters include, for example temperature-regulated promoters (e.g., a heat shock promoter) and light-regulated promoters (e.g., a light-inducible promoter or a light-repressible promoter).

[0062] Tissue-specific promoters can be, for example, neuron-specific promoters, glia-specific promoters, muscle cell-specific promoters, heart cell-specific promoters, kidney cell-specific promoters, bone cell-specific promoters, endothelial cell-specific promoters, or immune cell-specific promoters (e.g., a B cell promoter or a T cell promoter).

[0063] Developmentally regulated promoters include, for example, promoters active only during an embryonic stage of development, or only in an adult cell.

[0064] “Operable linkage” or being “operably linked” includes juxtaposition of two or more components (e.g., a promoter and another sequence element) such that both components function normally and allow the possibility that at least one of the components can mediate a function that is exerted upon at least one of the other components. For example, a promoter can be operably linked to a coding sequence if the promoter controls the level of transcription of the coding sequence in response to the presence or absence of one or more transcriptional regulatory factors. Operable linkage can include such sequences being contiguous with each other or acting in trans (e.g., a regulatory sequence can act at a distance to control transcription of the coding sequence).

[0065] “Complementarity” of nucleic acids means that a nucleotide sequence in one strand of nucleic acid, due to orientation of its nucleobase groups, forms hydrogen bonds with another sequence on an opposing nucleic acid strand. The complementary bases in DNA are typically A with T and C with G. In RNA, they are typically C with G and U with A. Complementarity can be perfect or substantial/sufficient. Perfect complementarity between two nucleic acids means that the two nucleic acids can form a duplex in which every base in the duplex is bonded to a complementary base by Watson-Crick pairing. “Substantial” or “sufficient” complementary means that a sequence in one strand is not completely and/or perfectly complementary to a sequence in an opposing strand, but that sufficient bonding occurs between bases on the two strands to form a stable hybrid complex in set of hybridization conditions (e.g., salt concentration and temperature). Such conditions can be predicted by using the sequences and standard mathematical calculations to predict the T_m (melting temperature) of hybridized strands, or by empirical determination of T_m by using routine methods. T_m includes the temperature at which a population of hybridization complexes formed between two nucleic acid strands are 50% denatured (i.e., a population of double-stranded nucleic acid molecules becomes half dissociated

into single strands). At a temperature below the T_m , formation of a hybridization complex is favored, whereas at a temperature above the T_m , melting or separation of the strands in the hybridization complex is favored. T_m may be estimated for a nucleic acid having a known G+C content in an aqueous 1 M NaCl solution by using, e.g., $T_m = 81.5 + 0.41(\% \text{ G+C})$, although other known T_m computations take into account nucleic acid structural characteristics.

[0066] “Hybridization condition” includes the cumulative environment in which one nucleic acid strand bonds to a second nucleic acid strand by complementary strand interactions and hydrogen bonding to produce a hybridization complex. Such conditions include the chemical components and their concentrations (e.g., salts, chelating agents, formamide) of an aqueous or organic solution containing the nucleic acids, and the temperature of the mixture. Other factors, such as the length of incubation time or reaction chamber dimensions may contribute to the environment. *See, e.g.,* Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, 2nd ed., pp. 1.90-1.91, 9.47-9.51, 11.47-11.57 (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), herein incorporated by reference in its entirety for all purposes.

[0067] Hybridization requires that the two nucleic acids contain complementary sequences, although mismatches between bases are possible. The conditions appropriate for hybridization between two nucleic acids depend on the length of the nucleic acids and the degree of complementation, variables which are well known. The greater the degree of complementation between two nucleotide sequences, the greater the value of the melting temperature (T_m) for hybrids of nucleic acids having those sequences. For hybridizations between nucleic acids with short stretches of complementarity (e.g. complementarity over 35 or fewer, 30 or fewer, 25 or fewer, 22 or fewer, 20 or fewer, or 18 or fewer nucleotides) the position of mismatches becomes important (*see* Sambrook *et al.*, *supra*, 11.7-11.8). Typically, the length for a hybridizable nucleic acid is at least about 10 nucleotides. Illustrative minimum lengths for a hybridizable nucleic acid include at least about 15 nucleotides, at least about 20 nucleotides, at least about 22 nucleotides, at least about 25 nucleotides, and at least about 30 nucleotides. Furthermore, the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the region of complementation and the degree of complementation.

[0068] The sequence of polynucleotide need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, a polynucleotide may hybridize over one or more segments such that intervening or adjacent segments are not involved in the

hybridization event (e.g., a loop structure or hairpin structure). A polynucleotide (e.g., gRNA) can comprise at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, or 100% sequence complementarity to a target region within the target nucleic acid sequence to which they are targeted. For example, a gRNA in which 18 of 20 nucleotides are complementary to a target region, and would therefore specifically hybridize, would represent 90% complementarity. In this example, the remaining noncomplementary nucleotides may be clustered or interspersed with complementary nucleotides and need not be contiguous to each other or to complementary nucleotides.

[0069] Percent complementarity between particular stretches of nucleic acid sequences within nucleic acids can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs (Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410; Zhang and Madden (1997) *Genome Res.* 7:649-656) or by using the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482-489).

[0070] The methods and compositions provided herein employ a variety of different components. Some components throughout the description can have active variants and fragments. Such components include, for example, Cas proteins, CRISPR RNAs, tracrRNAs, and guide RNAs. Biological activity for each of these components is described elsewhere herein. The term “functional” refers to the innate ability of a protein or nucleic acid (or a fragment or variant thereof) to exhibit a biological activity or function. Such biological activities or functions can include, for example, the ability of a Cas protein to bind to a guide RNA and to a target DNA sequence. The biological functions of functional fragments or variants may be the same or may in fact be changed (e.g., with respect to their specificity or selectivity or efficacy) in comparison to the original, but with retention of the basic biological function.

[0071] The term “variant” refers to a nucleotide sequence differing from the sequence most prevalent in a population (e.g., by one nucleotide) or a protein sequence different from the sequence most prevalent in a population (e.g., by one amino acid).

[0072] The term “fragment” when referring to a protein means a protein that is shorter or has fewer amino acids than the full-length protein. The term “fragment” when referring to a nucleic acid means a nucleic acid that is shorter or has fewer nucleotides than the full-length nucleic

acid. A fragment can be, for example, an N-terminal fragment (i.e., removal of a portion of the C-terminal end of the protein), a C-terminal fragment (i.e., removal of a portion of the N-terminal end of the protein), or an internal fragment.

[0073] “Sequence identity” or “identity” in the context of two polynucleotides or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins, residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have “sequence similarity” or “similarity.” Means for making this adjustment are well known. Typically, this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).

[0074] “Percentage of sequence identity” includes the value determined by comparing two optimally aligned sequences (greatest number of perfectly matched residues) over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity. Unless otherwise specified (e.g., the shorter sequence includes a linked heterologous sequence), the comparison window is the full length of the shorter of the two sequences being compared.

[0075] Unless otherwise stated, sequence identity/similarity values include the value obtained using GAP Version 10 using the following parameters: % identity and % similarity for a nucleotide sequence using GAP Weight of 50 and Length Weight of 3, and the nwsgapdna.cmp scoring matrix; % identity and % similarity for an amino acid sequence using GAP Weight of 8 and Length Weight of 2, and the BLOSUM62 scoring matrix; or any equivalent program thereof. “Equivalent program” includes any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by GAP Version 10.

[0076] The term “conservative amino acid substitution” refers to the substitution of an amino acid that is normally present in the sequence with a different amino acid of similar size, charge, or polarity. Examples of conservative substitutions include the substitution of a non-polar (hydrophobic) residue such as isoleucine, valine, or leucine for another non-polar residue. Likewise, examples of conservative substitutions include the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, or between glycine and serine. Additionally, the substitution of a basic residue such as lysine, arginine, or histidine for another, or the substitution of one acidic residue such as aspartic acid or glutamic acid for another acidic residue are additional examples of conservative substitutions. Examples of non-conservative substitutions include the substitution of a non-polar (hydrophobic) amino acid residue such as isoleucine, valine, leucine, alanine, or methionine for a polar (hydrophilic) residue such as cysteine, glutamine, glutamic acid or lysine and/or a polar residue for a non-polar residue. Typical amino acid categorizations are summarized in **Table 1** below.

[0077] Table 1. Amino Acid Categorizations.

Alanine	Ala	A	Nonpolar	Neutral	1.8
Arginine	Arg	R	Polar	Positive	-4.5
Asparagine	Asn	N	Polar	Neutral	-3.5
Aspartic acid	Asp	D	Polar	Negative	-3.5
Cysteine	Cys	C	Nonpolar	Neutral	2.5
Glutamic acid	Glu	E	Polar	Negative	-3.5
Glutamine	Gln	Q	Polar	Neutral	-3.5
Glycine	Gly	G	Nonpolar	Neutral	-0.4
Histidine	His	H	Polar	Positive	-3.2
Isoleucine	Ile	I	Nonpolar	Neutral	4.5
Leucine	Leu	L	Nonpolar	Neutral	3.8
Lysine	Lys	K	Polar	Positive	-3.9
Methionine	Met	M	Nonpolar	Neutral	1.9
Phenylalanine	Phe	F	Nonpolar	Neutral	2.8
Proline	Pro	P	Nonpolar	Neutral	-1.6
Serine	Ser	S	Polar	Neutral	-0.8
Threonine	Thr	T	Polar	Neutral	-0.7
Tryptophan	Trp	W	Nonpolar	Neutral	-0.9
Tyrosine	Tyr	Y	Polar	Neutral	-1.3
Valine	Val	V	Nonpolar	Neutral	4.2

[0078] The term “*in vitro*” includes artificial environments and to processes or reactions that occur within an artificial environment (e.g., a test tube). The term “*in vivo*” includes natural environments (e.g., a cell or organism or body) and to processes or reactions that occur within a natural environment. The term “*ex vivo*” includes cells that have been removed from the body of an individual and to processes or reactions that occur within such cells.

[0079] The term “reporter gene” refers to a nucleic acid having a sequence encoding a gene product (typically an enzyme) that is easily and quantifiably assayed when a construct comprising the reporter gene sequence operably linked to an endogenous or heterologous promoter and/or enhancer element is introduced into cells containing (or which can be made to contain) the factors necessary for the activation of the promoter and/or enhancer elements. Examples of reporter genes include, but are not limited, to genes encoding beta-galactosidase (lacZ), the bacterial chloramphenicol acetyltransferase (cat) genes, firefly luciferase genes, genes encoding beta-glucuronidase (GUS), and genes encoding fluorescent proteins. A “reporter protein” refers to a protein encoded by a reporter gene.

[0080] The term “fluorescent reporter protein” as used herein means a reporter protein that is detectable based on fluorescence wherein the fluorescence may be either from the reporter protein directly, activity of the reporter protein on a fluorogenic substrate, or a protein with affinity for binding to a fluorescent tagged compound. Examples of fluorescent proteins include green fluorescent proteins (e.g., GFP, GFP-2, tagGFP, turboGFP, eGFP, Emerald, Azami Green, Monomeric Azami Green, CopGFP, AceGFP, and ZsGreen1), yellow fluorescent proteins (e.g., YFP, eYFP, Citrine, Venus, YPet, PhiYFP, and ZsYellow1), blue fluorescent proteins (e.g., BFP, eBFP, eBFP2, Azurite, mKalamal, GFPuv, Sapphire, and T-sapphire), cyan fluorescent proteins (e.g., CFP, eCFP, Cerulean, CyPet, AmCyan1, and Midoriishi-Cyan), red fluorescent proteins (e.g., RFP, mKate, mKate2, mPlum, DsRed monomer, mCherry, mRFP1, DsRed-Express, DsRed2, DsRed-Monomer, HcRed-Tandem, HcRed1, AsRed2, eqFP611, mRaspberry, mStrawberry, and Jred), orange fluorescent proteins (e.g., mOrange, mKO, Kusabira-Orange, Monomeric Kusabira-Orange, mTangerine, and tdTomato), and any other suitable fluorescent protein whose presence in cells can be detected by flow cytometry methods.

[0081] Repair in response to double-strand breaks (DSBs) occurs principally through two conserved DNA repair pathways: homologous recombination (HR) and non-homologous end joining (NHEJ). *See* Kasperek & Humphrey (2011) *Seminars in Cell & Dev. Biol.* 22:886-897, herein incorporated by reference in its entirety for all purposes. Likewise, repair of a target nucleic acid mediated by an exogenous donor nucleic acid can include any process of exchange of genetic information between the two polynucleotides.

[0082] The term “recombination” includes any process of exchange of genetic information between two polynucleotides and can occur by any mechanism. Recombination can occur via homology directed repair (HDR) or homologous recombination (HR). HDR or HR includes a form of nucleic acid repair that can require nucleotide sequence homology, uses a “donor” molecule as a template for repair of a “target” molecule (i.e., the one that experienced the double-strand break), and leads to transfer of genetic information from the donor to target. Without wishing to be bound by any particular theory, such transfer can involve mismatch correction of heteroduplex DNA that forms between the broken target and the donor, and/or synthesis-dependent strand annealing, in which the donor is used to resynthesize genetic information that will become part of the target, and/or related processes. In some cases, the donor polynucleotide, a portion of the donor polynucleotide, a copy of the donor polynucleotide,

or a portion of a copy of the donor polynucleotide integrates into the target DNA. *See Wang et al. (2013) Cell 153:910-918; Mandalos et al. (2012) PLOS ONE 7:e45768:1-9; and Wang et al. (2013) Nat Biotechnol. 31:530-532*, each of which is herein incorporated by reference in its entirety for all purposes.

[0083] NHEJ includes the repair of double-strand breaks in a nucleic acid by direct ligation of the break ends to one another or to an exogenous sequence without the need for a homologous template. Ligation of non-contiguous sequences by NHEJ can often result in deletions, insertions, or translocations near the site of the double-strand break. For example, NHEJ can also result in the targeted integration of an exogenous donor nucleic acid through direct ligation of the break ends with the ends of the exogenous donor nucleic acid (i.e., NHEJ-based capture). Such NHEJ-mediated targeted integration can be preferred for insertion of an exogenous donor nucleic acid when homology directed repair (HDR) pathways are not readily usable (e.g., in non-dividing cells, primary cells, and cells which perform homology-based DNA repair poorly). In addition, in contrast to homology-directed repair, knowledge concerning large regions of sequence identity flanking the cleavage site (beyond the overhangs created by Cas-mediated cleavage) is not needed, which can be beneficial when attempting targeted insertion into organisms that have genomes for which there is limited knowledge of the genomic sequence. The integration can proceed via ligation of blunt ends between the exogenous donor nucleic acid and the cleaved genomic sequence, or via ligation of sticky ends (i.e., having 5' or 3' overhangs) using an exogenous donor nucleic acid that is flanked by overhangs that are compatible with those generated by the Cas protein in the cleaved genomic sequence. *See, e.g., US 2011/020722, WO 2014/033644, WO 2014/089290, and Maresca et al. (2013) Genome Res. 23(3):539-546*, each of which is herein incorporated by reference in its entirety for all purposes. If blunt ends are ligated, target and/or donor resection may be needed to generation regions of microhomology needed for fragment joining, which may create unwanted alterations in the target sequence.

[0084] Compositions or methods “comprising” or “including” one or more recited elements may include other elements not specifically recited. For example, a composition that “comprises” or “includes” a protein may contain the protein alone or in combination with other ingredients. The transitional phrase “consisting essentially of” means that the scope of a claim is to be interpreted to encompass the specified elements recited in the claim and those that do not materially affect the basic and novel characteristic(s) of the claimed invention. Thus, the term

“consisting essentially of” when used in a claim of this invention is not intended to be interpreted to be equivalent to “comprising.”

[0085] “Optional” or “optionally” means that the subsequently described event or circumstance may or may not occur and that the description includes instances in which the event or circumstance occurs and instances in which it does not.

[0086] Designation of a range of values includes all integers within or defining the range, and all subranges defined by integers within the range.

[0087] Unless otherwise apparent from the context, the term “about” encompasses values within a standard margin of error of measurement (e.g., SEM) of a stated value.

[0088] The term “and/or” refers to and encompasses any and all possible combinations of one or more of the associated listed items, as well as the lack of combinations when interpreted in the alternative (“or”).

[0089] The term “or” refers to any one member of a particular list and also includes any combination of members of that list.

[0090] The singular forms of the articles “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. For example, the term “a Cas protein” or “at least one Cas protein” can include a plurality of Cas proteins, including mixtures thereof.

[0091] Statistically significant means $p \leq 0.05$.

DETAILED DESCRIPTION

I. Overview

[0092] The CRISPR/Cas9 system is a powerful tool for genome engineering. One limitation of the system *in vivo* is the need to simultaneously introduce all components into a living organism. The typical method of intruding these components is to transiently transfect DNA constructs into cells that will generate the appropriate RNAs and protein. Though effective, this approach has an inherent disadvantage as the cells must rely on the plasmid DNA constructs to first undergo transcription and then translation before the Cas9 protein is available to interact with the sgRNA component. Better methods and tools are needed to more effectively assess the activity of CRISPR/Cas agents and to assess different delivery methods and parameters for targeting specific tissues or cell types *in vivo*.

[0093] Methods and compositions are provided herein for assessing CRISPR/Cas-mediated non-homologous end joining (NHEJ) activity and/or CRISPR/Cas-induced recombination of a target genomic locus with an exogenous donor nucleic acid *in vivo* and *ex vivo*. The methods and compositions employ cells and non-human animals comprising a Cas expression cassette (e.g., a genomically integrated Cas expression cassette) so that the Cas protein can be constitutively available or, for example, available in a tissue-specific or temporal-specific manner.

[0094] Non-human animals comprising the Cas expression cassettes simplify the process for testing delivery and activity of CRISPR/Cas components *in vivo* because only the guide RNAs need to be introduced into the non-human animal. In addition, the Cas expression cassettes can optionally be conditional Cas expression cassettes that can be selectively expressed in particular tissues or developmental stages, thereby reducing the risk of Cas-mediated toxicity *in vivo*, or can be constitutively expressed to enable testing of activity in any and all types of cells, tissues, and organs.

[0095] Methods and compositions are also provided for making and using these non-human animals to test and measure the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*. In some such methods of testing and measuring the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*, a guide RNA can be delivered to the Cas-ready non-human animal via AAV-mediated delivery. As shown in Example 1, AAV-mediated delivery of guide RNAs to Cas9-ready mice, and particularly AAV8-mediated delivery to the liver, results in surprisingly higher levels of CRISPR/Cas targeting than delivery of guide RNAs via LNPs or HDD to Cas9-ready mice or delivery of both Cas9 and guide RNAs to wild type mice.

II. Non-Human Animals Comprising Cas Expression Cassettes

[0096] The methods and compositions disclosed herein utilize non-human animals or cells comprising Cas expression cassettes to assess the ability of Clustered Regularly Interspersed Short Palindromic Repeats (CRISPR)/CRISPR-associated (Cas) systems or components of such systems (e.g., guide RNAs introduced into the non-human animal or cell) to modify a target genomic locus *in vivo* or *ex vivo*.

[0097] CRISPR/Cas systems include transcripts and other elements involved in the expression of, or directing the activity of, Cas genes. A CRISPR/Cas system can be, for example, a type I, a type II, or a type III system. Alternatively, a CRISPR/Cas system can be a type V system (e.g., subtype V-A or subtype V-B). CRISPR/Cas systems used in the compositions and methods disclosed herein can be non-naturally occurring. A “non-naturally occurring” system includes anything indicating the involvement of the hand of man, such as one or more components of the system being altered or mutated from their naturally occurring state, being at least substantially free from at least one other component with which they are naturally associated in nature, or being associated with at least one other component with which they are not naturally associated. For example, non-naturally occurring CRISPR/Cas systems can employ CRISPR complexes comprising a gRNA and a Cas protein that do not naturally occur together, a Cas protein that does not occur naturally, or a gRNA that does not occur naturally.

[0098] The methods and compositions disclosed herein employ the CRISPR/Cas systems by testing the ability of CRISPR complexes (comprising a guide RNA (gRNA) complexed with a Cas protein) to induce site-directed cleavage events within a target genomic locus *in vivo* to modify the target genomic locus via non-homologous end joining (NHEJ), via homology-directed repair in the presence of an exogenous donor nucleic acid, or via any other means of repair or recombination.

A. Cas9-Ready Non-Human Animals

[0099] The cells and non-human animals disclosed herein comprise a Cas expression cassette. Cas proteins generally comprise at least one RNA recognition or binding domain that can interact with guide RNAs (gRNAs, described in more detail below), and nuclease domains. A nuclease domain possesses catalytic activity for nucleic acid cleavage, which includes the breakage of the covalent bonds of a nucleic acid molecule. Cleavage can produce blunt ends or staggered ends, and it can be single-stranded or double-stranded. A Cas protein can have full cleavage activity to create a double-strand break in the target nucleic acid (e.g., a double-strand break with blunt ends), or it can be a nickase that creates a single-strand break in the target nucleic acid.

[00100] Cells or non-human animals comprising a Cas expression cassette have the advantage of needing delivery only of guide RNAs in order to detect CRISPR/Cas-mediated modification of a target genomic locus.

(1) Cas Expression Cassettes

[00101] The cells and non-human animals described herein comprise a Cas expression cassette. The Cas expression cassette can be stably integrated into the genome (i.e., into a chromosome) of the cell or non-human animal or it can be located outside of a chromosome (e.g., extrachromosomally replicating DNA). Optionally, the Cas expression cassette is stably integrated into the genome. The stably integrated Cas expression cassette can be randomly integrated into the genome of the non-human animal (i.e., transgenic), or it can be integrated into a predetermined region of the genome of the non-human animal (i.e., knock in). Optionally, the Cas expression cassette is stably integrated into a safe harbor locus as described elsewhere herein. The target genomic locus at which the Cas expression cassette is stably integrated can be heterozygous for the Cas expression cassette or homozygous for the Cas expression cassette.

[00102] The Cas protein encoded by the Cas expression cassette can be any Cas protein (e.g., a Cas9 protein), examples of which are described below. The encoded Cas protein can further comprise one or more nuclear localization signals (NLSs) (e.g., an N-terminal NLS and a C-terminal NLS), and the sequence encoding the Cas protein can be codon-optimized for the cell or non-human animal as described below. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9 protein sequence set forth in SEQ ID NO: 19. The coding sequence can comprise, consist essentially of, or consist of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9 coding sequence set forth in SEQ ID NO: 30.

[00103] An example of a Cas expression cassette comprises a Cas coding sequence downstream of a polyadenylation signal or transcription terminator flanked by recombinase recognition sites recognized by a site-specific recombinase. The polyadenylation signal or transcription terminator prevents transcription and expression of the Cas protein. However, upon exposure to the site-specific recombinase, the polyadenylation signal or transcription terminator will be excised, and the Cas protein can be expressed.

[00104] Such a configuration for a Cas expression cassette can enable tissue-specific expression or developmental-stage-specific expression in non-human animals comprising the Cas expression cassette if the polyadenylation signal or transcription terminator is excised in a tissue-specific or developmental-stage-specific manner. This may reduce toxicity due to prolonged expression of the Cas protein in a cell or non-human animal or expression of the Cas protein at undesired developmental stages or in undesired cell or tissue types within a non-human animal. For example, toxicity could result from cleavage and disruption of off-target sites. *See, e.g., Parikh et al. (2015) PLoS One 10(1):e0116484.* Inducible expression may also be beneficial because the possibility of editing some genes in certain tissues (e.g., such as immune cells) may be detrimental, along with potentially causing an immune response. For example, in some cases, if a gene is mutated throughout the individual it may be lethal, but if it is mutated in a specific tissue or cell type, it would be beneficial. Excision of the polyadenylation signal or transcription terminator in a tissue-specific or developmental-stage-specific manner can be achieved if the non-human animal comprising the Cas expression cassette further comprises the site-specific recombinase operably linked to a tissue-specific or developmental-stage-specific promoter (e.g., albumin promoter for liver-specific expression or insulin 2 promoter for pancreas-specific expression). Similarly, LNP formulations specific for liver or other tissues can be used to deliver the recombinase, or AAV delivery methods or AAV serotypes specific for particular tissues (e.g., AAV8 for liver, or AAV direct injection for pancreas) can be used to deliver the recombinase. The polyadenylation signal or transcription terminator will then be excised only in those tissues or at those developmental stages, enabling tissue-specific expression or developmental-stage-specific expression of the Cas protein. In one example, the Cas protein can be expressed in a liver-specific manner. Examples of such promoters that have been used to develop such “recombinase deleter” strains of non-human animals are disclosed elsewhere herein.

[00105] Any transcription terminator or polyadenylation signal can be used. A “transcription terminator” as used herein refers to a DNA sequence that causes termination of transcription. In eukaryotes, transcription terminators are recognized by protein factors, and termination is followed by polyadenylation, a process of adding a poly(A) tail to the mRNA transcripts in presence of the poly(A) polymerase. The mammalian poly(A) signal typically consists of a core sequence, about 45 nucleotides long, that may be flanked by diverse auxiliary sequences that

serve to enhance cleavage and polyadenylation efficiency. The core sequence consists of a highly conserved upstream element (AATAAA or AAUAAA) in the mRNA, referred to as a poly A recognition motif or poly A recognition sequence), recognized by cleavage and polyadenylation-specificity factor (CPSF), and a poorly defined downstream region (rich in Us or Gs and Us), bound by cleavage stimulation factor (CstF). Examples of transcription terminators that can be used include, for example, the human growth hormone (HGH) polyadenylation signal, the simian virus 40 (SV40) late polyadenylation signal, the rabbit beta-globin polyadenylation signal, the bovine growth hormone (BGH) polyadenylation signal, the phosphoglycerate kinase (PGK) polyadenylation signal, an AOX1 transcription termination sequence, a CYC1 transcription termination sequence, or any transcription termination sequence known to be suitable for regulating gene expression in eukaryotic cells.

[00106] Site-specific recombinases include enzymes that can facilitate recombination between recombinase recognition sites, where the two recombination sites are physically separated within a single nucleic acid or on separate nucleic acids. Examples of recombinases include Cre, Flp, and Dre recombinases. One example of a Cre recombinase gene is Crei, in which two exons encoding the Cre recombinase are separated by an intron to prevent its expression in a prokaryotic cell. Such recombinases can further comprise a nuclear localization signal to facilitate localization to the nucleus (e.g., NLS-Crei). Recombinase recognition sites include nucleotide sequences that are recognized by a site-specific recombinase and can serve as a substrate for a recombination event. Examples of recombinase recognition sites include FRT, FRT11, FRT71, attp, att, rox, and lox sites such as loxP, lox511, lox2272, lox66, lox71, loxM2, and lox5171.

[00107] The Cas expression cassette can be operably linked to any suitable promoter for expression *in vivo* within a non-human animal. The non-human animal can be any suitable non-human animal as described elsewhere herein. As one example, the Cas expression cassette can be operably linked to an endogenous promoter at a target genomic locus, such as a *Rosa26* promoter. Alternatively, the Cas expression cassette can be operably linked to an exogenous promoter, such as a constitutively active promoter (e.g., a CAG promoter or a chicken beta actin promoter/enhancer coupled with the cytomegalovirus (CMV) immediate-early enhancer (CAGG)), a conditional promoter, an inducible promoter, a temporally restricted promoter (e.g., a developmentally regulated promoter), or a spatially restricted promoter (e.g., a cell-specific or

tissue-specific promoter). Such promoters are well-known and are discussed elsewhere herein. An exemplary CAGG promoter is set forth in SEQ ID NO: 38 or comprises, consists essentially of, or consists of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 38.

[00108] The Cas expression cassettes disclosed herein can comprise other components as well. A Cas expression cassette can further comprise a 3' splicing sequence at the 5' end of the Cas expression cassette and/or a second polyadenylation signal following the coding sequence for the Cas protein at the 3' end of the Cas expression cassette. A Cas expression cassette can further comprise a selection cassette comprising, for example, the coding sequence for a drug resistance protein. Examples of suitable selection markers include neomycin phosphotransferase (*neo_r*), hygromycin B phosphotransferase (*hyg_r*), puromycin-N-acetyltransferase (*puro_r*), blasticidin S deaminase (*bsr_r*), xanthine/guanine phosphoribosyl transferase (*gpt*), and herpes simplex virus thymidine kinase (*HSV-k*). Optionally, the selection cassette can be flanked by recombinase recognition sites for a site-specific recombinase. If the Cas expression cassette also comprises recombinase recognition sites flanking a polyadenylation signal upstream of the Cas coding sequence as described above, optionally a different set of recombinase recognition sites recognized by a different recombinase are used to flank the selection cassette. Alternatively, the same set of recombinase recognition sites can flank both the polyadenylation signal upstream of the Cas coding sequence and the selection cassette. An exemplary *neo_r*-polyadenylation sequence is set forth in SEQ ID NO: 37 or comprises, consists essentially of, or consists of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 37.

[00109] A Cas expression cassette can also comprise a nucleic acid encoding a protein tag, such as a 3xFLAG tag. An example of such a tag is set forth in SEQ ID NO: 23, which is optionally encoded by SEQ ID NO: 34. For example, the tag can be at the N-terminus of the Cas protein, at the C-terminus of the Cas protein, or internally within the Cas protein. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the 3xFLAG-Cas9 protein sequence set forth in SEQ ID NO: 22 or the 3xFLAG-Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 16. The coding sequence can comprise, consist essentially of, or consist of a sequence at least 90%, 91%, 92%, 93%, 94%,

95%, 96%, 97%, 98%, 99%, or 100% identical to the 3xFLAG-Cas9 coding sequence set forth in SEQ ID NO: 31 or the 3xFLAG-Cas9-P2A-eGFP coding sequence set forth in SEQ ID NO: 29, respectively.

[00110] A Cas expression cassette can also comprise a nucleic acid encoding one or more reporter proteins, such as a fluorescent protein (e.g., a green fluorescent protein). Any suitable reporter protein can be used. For example, a fluorescent reporter protein as defined elsewhere herein can be used, or a non-fluorescent reporter protein can be used. Examples of fluorescent reporter proteins are provided elsewhere herein. Non-fluorescent reporter proteins include, for example, reporter proteins that can be used in histochemical or bioluminescent assays, such as beta-galactosidase, luciferase (e.g., Renilla luciferase, firefly luciferase, and NanoLuc luciferase), and beta-glucuronidase. A Cas expression cassette can include a reporter protein that can be detected in a flow cytometry assay (e.g., a fluorescent reporter protein such as a green fluorescent protein) and/or a reporter protein that can be detected in a histochemical assay (e.g., beta-galactosidase protein). One example of such a histochemical assay is visualization of *in situ* beta-galactosidase expression histochemically through hydrolysis of X-Gal (5-bromo-4-chloro-3-indoyl-b-D-galactopyranoside), which yields a blue precipitate, or using fluorogenic substrates such as beta-methyl umbelliferyl galactoside (MUG) and fluorescein digalactoside (FDG).

[00111] The Cas expression cassette in such cases can comprise a multicistronic nucleic acid. For example, such nucleic acids can the Cas protein coding sequence and the reporter protein coding sequence (in either order) separated by an intervening internal ribosome entry site (IRES) or an intervening 2A peptide coding sequence. Multicistronic expression constructs simultaneously express two or more separate proteins from the same mRNA (i.e., a transcript produced from the same promoter). Suitable strategies for multicistronic expression of proteins include, for example, the use of a 2A peptide and the use of an internal ribosome entry site (IRES). For example, such nucleic acids can comprise coding sequences for two or more reporter proteins separated by an intervening internal ribosome entry site (IRES) or an intervening 2A peptide coding sequence. As one example, such multicistronic vectors can use one or more internal ribosome entry sites (IRES) to allow for initiation of translation from an internal region of an mRNA. As another example, such multicistronic vectors can use one or more 2A peptides. These peptides are small “self-cleaving” peptides, generally having a length of 18–22 amino acids and produce equimolar levels of multiple genes from the same mRNA.

Ribosomes skip the synthesis of a glycyl-prolyl peptide bond at the C-terminus of a 2A peptide, leading to the “cleavage” between a 2A peptide and its immediate downstream peptide. *See, e.g., Kim et al. (2011) PLoS One 6(4): e18556*, herein incorporated by reference in its entirety for all purposes. The “cleavage” occurs between the glycine and proline residues found on the C-terminus, meaning the upstream cistron will have a few additional residues added to the end, while the downstream cistron will start with the proline. As a result, the “cleaved-off” downstream peptide has proline at its N-terminus. 2A-mediated cleavage is a universal phenomenon in all eukaryotic cells. 2A peptides have been identified from picornaviruses, insect viruses and type C rotaviruses. *See, e.g., Szymczak et al. (2005) Expert Opin Biol Ther 5:627-638*, herein incorporated by reference in its entirety for all purposes. Examples of 2A peptides that can be used include *Thosea asigna* virus 2A (T2A); porcine teschovirus-1 2A (P2A); equine rhinitis A virus (ERAV) 2A (E2A); and FMDV 2A (F2A). Exemplary T2A, P2A, E2A, and F2A sequences include the following: T2A (EGRGSLTCTCGDVEENPGP; SEQ ID NO: 2); P2A (ATNFSLLKQAGDVEENPGP; SEQ ID NO: 3); E2A (QCTNYALLKLAGDVESNPGP; SEQ ID NO: 4); and F2A (VKQTLNFDLLKLAGDVESNPGP; SEQ ID NO: 5). GSG residues can be added to the 5' end of any of these peptides to improve cleavage efficiency. An exemplary coding sequence for P2A with GSG residues added at the 5' end is set forth in SEQ ID NO: 32 or comprises, consists essentially of, or consists of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 32. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 13 or the 3xFLAG-Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 16. The coding sequence can comprise, consist essentially of, or consist of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9-P2A-eGFP coding sequence set forth in SEQ ID NO: 28 or the 3xFLAG-Cas9-P2A-eGFP coding sequence set forth in SEQ ID NO: 29, respectively.

[00112] Cas expression cassettes can also comprise other elements, such as posttranscriptional regulatory elements or polyadenylation signals downstream of the Cas coding sequence. An exemplary posttranscriptional regulatory element is the woodchuck hepatitis virus

posttranscriptional regulatory element (WPRE) set forth in SEQ ID NO: 35 or comprises, consists essentially of, or consists of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 35. An exemplary polyadenylation signal is the bovine growth hormone polyadenylation signal set forth in SEQ ID NO: 36 or comprises, consists essentially of, or consists of a sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 36.

[00113] One exemplary Cas expression cassette comprises from 5' to 3': (a) a 3' splicing sequence; (b) a polyadenylation signal flanked by first and second recombinase recognition sites for a recombinase (e.g., loxP sites for a Cre recombinase); (c) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end); (d) a 2A protein coding sequence (e.g., a P2A coding sequence); and (e) a coding sequence for a reporter protein (e.g., a fluorescent reporter protein, such as a green fluorescent protein). *See, e.g., Figure 1 and MAID2599 in Figure 14.* Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 1. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 13.

[00114] Another exemplary Cas expression cassette comprises from 5' to 3': (a) a 3' splicing sequence; (b) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end); (c) a 2A protein coding sequence (e.g., a P2A coding sequence); and (d) a coding sequence for a reporter protein (e.g., a fluorescent reporter protein, such as a green fluorescent protein). *See, e.g., MAID2600 in Figure 14.* Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 12. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 13.

[00115] Another exemplary Cas expression cassette comprises from 5' to 3': (a) a 3' splicing sequence; (b) a polyadenylation signal flanked by first and second recombinase recognition sites for a recombinase (e.g., loxP sites for a Cre recombinase); and (c) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end). *See, e.g.,* MAID2660 in **Figure 14**. Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 17. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9 protein sequence set forth in SEQ ID NO: 19.

[00116] Another exemplary Cas expression cassette comprises from 5' to 3': (a) a 3' splicing sequence; and (b) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end). *See, e.g.,* MAID2661 in **Figure 14**. Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 18. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the Cas9 protein sequence set forth in SEQ ID NO: 19.

[00117] Another exemplary Cas expression cassette comprises from 5' to 3': (a) an exogenous promoter (e.g., a constitutive promoter, such as a CAGG promoter); (b) a polyadenylation signal flanked by first and second recombinase recognition sites for a recombinase (e.g., loxP sites for a Cre recombinase); (c) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end, optionally with a tag at the N-terminal or C-terminal end, such as a 3xFLAG tag at the N-terminal end); (d) a 2A protein coding sequence (e.g., a P2A coding sequence); and (e) a coding sequence for a reporter protein (e.g., a fluorescent reporter protein, such as a green fluorescent protein). *See, e.g.,* MAID2658 in **Figure 14**. Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 14. For example, such an expression cassette

can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the 3xFLAG-Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 16.

[00118] Another exemplary Cas expression cassette comprises from 5' to 3': (a) an exogenous promoter (e.g., a constitutive promoter, such as a CAGG promoter); (b) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end, optionally with a tag at the N-terminal or C-terminal end, such as a 3xFLAG tag at the N-terminal end); (c) a 2A protein coding sequence (e.g., a P2A coding sequence); and (d) a coding sequence for a reporter protein (e.g., a fluorescent reporter protein, such as a green fluorescent protein). *See, e.g.,* MAID2659 in **Figure 14**. Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 15. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the 3xFLAG-Cas9-P2A-eGFP protein sequence set forth in SEQ ID NO: 16.

[00119] Another exemplary Cas expression cassette comprises from 5' to 3': (a) an exogenous promoter (e.g., a constitutive promoter, such as a CAGG promoter); (b) a polyadenylation signal flanked by first and second recombinase recognition sites for a recombinase (e.g., loxP sites for a Cre recombinase); and (c) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an NLS at the C-terminal end, optionally with a tag at the N-terminal or C-terminal end, such as a 3xFLAG tag at the N-terminal end). *See, e.g.,* MAID2672 in **Figure 14**. Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 20. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the 3xFLAG-Cas9 protein sequence set forth in SEQ ID NO: 22.

[00120] Another exemplary Cas expression cassette comprises from 5' to 3': (a) an exogenous promoter (e.g., a constitutive promoter, such as a CAGG promoter); and (b) a Cas protein coding sequence (e.g., an NLS-Cas9 coding sequence, such as with an NLS at the N-terminal end and an

NLS at the C-terminal end, optionally with a tag at the N-terminal or C-terminal end, such as a 3xFLAG tag at the N-terminal end). *See, e.g.*, MAID2673 in **Figure 14**. Such an expression cassette can comprise, consist essentially of, or consist of a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequence set forth in SEQ ID NO: 21. For example, such an expression cassette can encode a protein comprising, consisting essentially of, or consisting of an amino acid sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the 3xFLAG-Cas9 protein sequence set forth in SEQ ID NO: 22.

[00121] The Cas expression cassettes described herein can be in any form. For example, a Cas expression cassette can be in a vector or plasmid, such as a viral vector. The Cas expression cassette can be operably linked to a promoter in an expression construct capable of directing expression of the Cas protein upon removal of the upstream polyadenylation signal.

Alternatively, a Cas expression cassette can be in a targeting vector as defined elsewhere herein. For example, the targeting vector can comprise homology arms flanking the Cas expression cassette, wherein the homology arms are suitable for directing recombination with a desired target genomic locus to facilitate genomic integration.

[00122] The Cas expression cassettes described herein can be *in vitro*, they can be within a cell (e.g., an embryonic stem cell) *ex vivo* (e.g., genomically integrated or extrachromosomal), or they can be in an organism (e.g., a non-human animal) *in vivo* (e.g., genomically integrated or extrachromosomal). If *ex vivo*, the Cas expression cassette can be in any type of cell from any organism, such as a totipotent cell such as an embryonic stem cell (e.g., a mouse or a rat embryonic stem cell) or an induced pluripotent stem cell (e.g., a human induced pluripotent stem cell). If *in vivo*, the Cas expression cassette can be in any type of organism (e.g., a non-human animal as described further elsewhere herein).

(2) Cas Proteins and Polynucleotides Encoding Cas Proteins

[00123] Cas proteins generally comprise at least one RNA recognition or binding domain that can interact with guide RNAs (gRNAs, described in more detail below). Cas proteins can also comprise nuclease domains (e.g., DNase or RNase domains), DNA-binding domains, helicase domains, protein-protein interaction domains, dimerization domains, and other domains. Some such domains (e.g., DNase domains) can be from a native Cas protein. Other such domains can

be added to make a modified Cas protein. A nuclease domain possesses catalytic activity for nucleic acid cleavage, which includes the breakage of the covalent bonds of a nucleic acid molecule. Cleavage can produce blunt ends or staggered ends, and it can be single-stranded or double-stranded. For example, a wild type Cas9 protein will typically create a blunt cleavage product. Alternatively, a wild type Cpf1 protein (e.g., FnCpf1) can result in a cleavage product with a 5-nucleotide 5' overhang, with the cleavage occurring after the 18th base pair from the PAM sequence on the non-targeted strand and after the 23rd base on the targeted strand. A Cas protein can have full cleavage activity to create a double-strand break at a target genomic locus (e.g., a double-strand break with blunt ends), or it can be a nickase that creates a single-strand break at a target genomic locus.

[00124] Examples of Cas proteins include Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas5e (CasD), Cas6, Cas6e, Cas6f, Cas7, Cas8a1, Cas8a2, Cas8b, Cas8c, Cas9 (Csn1 or Csx12), Cas10, Cas10d, CasF, CasG, CasH, Csy1, Csy2, Csy3, Cse1 (CasA), Cse2 (CasB), Cse3 (CasE), Cse4 (CasC), Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, and Cu1966, and homologs or modified versions thereof.

[00125] An exemplary Cas protein is a Cas9 protein or a protein derived from Cas9. Cas9 proteins are from a type II CRISPR/Cas system and typically share four key motifs with a conserved architecture. Motifs 1, 2, and 4 are RuvC-like motifs, and motif 3 is an HNH motif. Exemplary Cas9 proteins are from *Streptococcus pyogenes*, *Streptococcus thermophilus*, *Streptococcus sp.*, *Staphylococcus aureus*, *Nocardiopsis dassonvillei*, *Streptomyces pristinaespiralis*, *Streptomyces viridochromogenes*, *Streptomyces viridochromogenes*, *Streptosporangium roseum*, *Streptosporangium roseum*, *Alicyclobacillus acidocaldarius*, *Bacillus pseudomycooides*, *Bacillus selenitireducens*, *Exiguobacterium sibiricum*, *Lactobacillus delbrueckii*, *Lactobacillus salivarius*, *Microscilla marina*, *Burkholderiales bacterium*, *Polaromonas naphthalenivorans*, *Polaromonas sp.*, *Crocospaera watsonii*, *Cyanothece sp.*, *Microcystis aeruginosa*, *Synechococcus sp.*, *Acetohalobium arabaticum*, *Ammonifex degensii*, *Caldicelulosiruptor beccii*, *Candidatus Desulfurudis*, *Clostridium botulinum*, *Clostridium difficile*, *Finegoldia magna*, *Natronaerobius thermophilus*, *Pelotomaculum thermopropionicum*, *Acidithiobacillus caldus*, *Acidithiobacillus ferrooxidans*, *Allochroamatium vinosum*, *Marinobacter sp.*, *Nitrosococcus halophilus*, *Nitrosococcus watsoni*, *Pseudoalteromonas*

haloplanktis, *Ktedonobacter racemifer*, *Methanohalobium evestigatum*, *Anabaena variabilis*, *Nodularia spumigena*, *Nostoc sp.*, *Arthrospira maxima*, *Arthrospira platensis*, *Arthrospira sp.*, *Lyngbya sp.*, *Microcoleus chthonoplastes*, *Oscillatoria sp.*, *Petrotoga mobilis*, *Thermosiphon africanus*, *Acaryochloris marina*, *Neisseria meningitidis*, or *Campylobacter jejuni*. Additional examples of the Cas9 family members are described in WO 2014/131833, herein incorporated by reference in its entirety for all purposes. Cas9 from *S. pyogenes* (SpCas9) (assigned SwissProt accession number Q99ZW2) is an exemplary Cas9 protein. Cas9 from *S. aureus* (SaCas9) (assigned UniProt accession number J7RUA5) is another exemplary Cas9 protein. Cas9 from *Campylobacter jejuni* (CjCas9) (assigned UniProt accession number Q0P897) is another exemplary Cas9 protein. See, e.g., Kim *et al.* (2017) *Nat. Comm.* 8:14500, herein incorporated by reference in its entirety for all purposes. SaCas9 is smaller than SpCas9, and CjCas9 is smaller than both SaCas9 and SpCas9.

[00126] Another example of a Cas protein is a Cpf1 (CRISPR from *Prevotella* and *Francisella* 1) protein. Cpf1 is a large protein (about 1300 amino acids) that contains a RuvC-like nuclease domain homologous to the corresponding domain of Cas9 along with a counterpart to the characteristic arginine-rich cluster of Cas9. However, Cpf1 lacks the HNH nuclease domain that is present in Cas9 proteins, and the RuvC-like domain is contiguous in the Cpf1 sequence, in contrast to Cas9 where it contains long inserts including the HNH domain. See, e.g., Zetsche *et al.* (2015) *Cell* 163(3):759-771, herein incorporated by reference in its entirety for all purposes. Exemplary Cpf1 proteins are from *Francisella tularensis* 1, *Francisella tularensis* subsp. *novicida*, *Prevotella albensis*, *Lachnospiraceae* bacterium MC2017 1, *Butyrivibrio proteoclasticus*, *Peregrinibacteria bacterium* GW2011_GWA2_33_10, *Parcubacteria bacterium* GW2011_GWC2_44_17, *Smithella sp.* SCADC, *Acidaminococcus sp.* BV3L6, *Lachnospiraceae* bacterium MA2020, *Candidatus Methanoplasma termitum*, *Eubacterium eligens*, *Moraxella bovoculi* 237, *Leptospira inadai*, *Lachnospiraceae* bacterium ND2006, *Porphyromonas crevioricanis* 3, *Prevotella disiens*, and *Porphyromonas macacae*. Cpf1 from *Francisella novicida* U112 (FnCpf1; assigned UniProt accession number A0Q7Q2) is an exemplary Cpf1 protein.

[00127] Cas proteins can be wild type proteins (i.e., those that occur in nature), modified Cas proteins (i.e., Cas protein variants), or fragments of wild type or modified Cas proteins. Cas proteins can also be active variants or fragments with respect to catalytic activity of wild type or

modified Cas proteins. Active variants or fragments with respect to catalytic activity can comprise at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to the wild type or modified Cas protein or a portion thereof, wherein the active variants retain the ability to cut at a desired cleavage site and hence retain nick-inducing or double-strand-break-inducing activity. Assays for nick-inducing or double-strand-break-inducing activity are known and generally measure the overall activity and specificity of the Cas protein on DNA substrates containing the cleavage site.

[00128] Cas proteins can be modified to increase or decrease one or more of nucleic acid binding affinity, nucleic acid binding specificity, and enzymatic activity. Cas proteins can also be modified to change any other activity or property of the protein, such as stability. For example, one or more nuclease domains of the Cas protein can be modified, deleted, or inactivated, or a Cas protein can be truncated to remove domains that are not essential for the function of the protein or to optimize (e.g., enhance or reduce) the activity or a property of the Cas protein.

[00129] One example of a modified Cas protein is the modified SpCas9-HF1 protein, which is a high-fidelity variant of *Streptococcus pyogenes* Cas9 harboring alterations (N497A/R661A/Q695A/Q926A) designed to reduce non-specific DNA contacts. *See, e.g.,* Kleinstiver *et al.* (2016) *Nature* 529(7587):490-495, herein incorporated by reference in its entirety for all purposes. Another example of a modified Cas protein is the modified eSpCas9 variant (K848A/K1003A/R1060A) designed to reduce off-target effects. *See, e.g.,* Slaymaker *et al.* (2016) *Science* 351(6268):84-88, herein incorporated by reference in its entirety for all purposes. Other SpCas9 variants include K855A and K810A/K1003A/R1060A.

[00130] Cas proteins can comprise at least one nuclease domain, such as a DNase domain. For example, a wild type Cpf1 protein generally comprises a RuvC-like domain that cleaves both strands of target DNA, perhaps in a dimeric configuration. Cas proteins can also comprise at least two nuclease domains, such as DNase domains. For example, a wild type Cas9 protein generally comprises a RuvC-like nuclease domain and an HNH-like nuclease domain. The RuvC and HNH domains can each cut a different strand of double-stranded DNA to make a double-stranded break in the DNA. *See, e.g.,* Jinek *et al.* (2012) *Science* 337:816-821, herein incorporated by reference in its entirety for all purposes.

[00131] One or more of the nuclease domains can be deleted or mutated so that they are no longer functional or have reduced nuclease activity. For example, if one of the nuclease domains is deleted or mutated in a Cas9 protein, the resulting Cas9 protein can be referred to as a nickase and can generate a single-strand break at a guide RNA target sequence within a double-stranded DNA but not a double-strand break (i.e., it can cleave the complementary strand or the non-complementary strand, but not both). An example of a mutation that converts Cas9 into a nickase is a D10A (aspartate to alanine at position 10 of Cas9) mutation in the RuvC domain of Cas9 from *S. pyogenes*. Likewise, H939A (histidine to alanine at amino acid position 839), H840A (histidine to alanine at amino acid position 840), or N863A (asparagine to alanine at amino acid position N863) in the HNH domain of Cas9 from *S. pyogenes* can convert the Cas9 into a nickase. Other examples of mutations that convert Cas9 into a nickase include the corresponding mutations to Cas9 from *S. thermophilus*. See, e.g., Saprunauskas *et al.* (2011) *Nucleic Acids Research* 39:9275-9282 and WO 2013/141680, each of which is herein incorporated by reference in its entirety for all purposes. Such mutations can be generated using methods such as site-directed mutagenesis, PCR-mediated mutagenesis, or total gene synthesis. Examples of other mutations creating nickases can be found, for example, in WO 2013/176772 and WO 2013/142578, each of which is herein incorporated by reference in its entirety for all purposes.

[00132] Examples of inactivating mutations in the catalytic domains of *Staphylococcus aureus* Cas9 proteins are also known. For example, the *Staphylococcus aureus* Cas9 enzyme (SaCas9) may comprise a substitution at position N580 (e.g., N580A substitution) and a substitution at position D10 (e.g., D10A substitution) to generate a nuclease-inactive Cas protein. See, e.g., WO 2016/106236, herein incorporated by reference in its entirety for all purposes.

[00133] Examples of inactivating mutations in the catalytic domains of Cpf1 proteins are also known. With reference to Cpf1 proteins from *Francisella novicida* U112 (FnCpf1), *Acidaminococcus* sp. BV3L6 (AsCpf1), *Lachnospiraceae bacterium* ND2006 (LbCpf1), and *Moraxella bovoculi* 237 (MbCpf1 Cpf1), such mutations can include mutations at positions 908, 993, or 1263 of AsCpf1 or corresponding positions in Cpf1 orthologs, or positions 832, 925, 947, or 1180 of LbCpf1 or corresponding positions in Cpf1 orthologs. Such mutations can include, for example one or more of mutations D908A, E993A, and D1263A of AsCpf1 or corresponding mutations in Cpf1 orthologs, or D832A, E925A, D947A, and D1180A of LbCpf1 or

corresponding mutations in Cpf1 orthologs. *See, e.g.*, US 2016/0208243, herein incorporated by reference in its entirety for all purposes.

[00134] Cas proteins can also be operably linked to heterologous polypeptides as fusion proteins. For example, a Cas protein can be fused to a cleavage domain or an epigenetic modification domain. *See* WO 2014/089290, herein incorporated by reference in its entirety for all purposes. Cas proteins can also be fused to a heterologous polypeptide providing increased or decreased stability. The fused domain or heterologous polypeptide can be located at the N-terminus, the C-terminus, or internally within the Cas protein.

[00135] As one example, a Cas protein can be fused to one or more heterologous polypeptides that provide for subcellular localization. Such heterologous polypeptides can include, for example, one or more nuclear localization signals (NLS) such as the monopartite SV40 NLS and/or a bipartite alpha-importin NLS for targeting to the nucleus, a mitochondrial localization signal for targeting to the mitochondria, an ER retention signal, and the like. *See, e.g.*, Lange *et al.* (2007) *J. Biol. Chem.* 282:5101-5105, herein incorporated by reference in its entirety for all purposes. Such subcellular localization signals can be located at the N-terminus, the C-terminus, or anywhere within the Cas protein. An NLS can comprise a stretch of basic amino acids, and can be a monopartite sequence or a bipartite sequence. Optionally, a Cas protein can comprise two or more NLSs, including an NLS (e.g., an alpha-importin NLS or a monopartite NLS) at the N-terminus and an NLS (e.g., an SV40 NLS or a bipartite NLS) at the C-terminus. A Cas protein can also comprise two or more NLSs at the N-terminus and/or two or more NLSs at the C-terminus.

[00136] Cas proteins can also be operably linked to a cell-penetrating domain or protein transduction domain. For example, the cell-penetrating domain can be derived from the HIV-1 TAT protein, the TLM cell-penetrating motif from human hepatitis B virus, MPG, Pep-1, VP22, a cell penetrating peptide from Herpes simplex virus, or a polyarginine peptide sequence. *See, e.g.*, WO 2014/089290 and WO 2013/176772, each of which is herein incorporated by reference in its entirety for all purposes. The cell-penetrating domain can be located at the N-terminus, the C-terminus, or anywhere within the Cas protein.

[00137] Cas proteins can also be operably linked to a heterologous polypeptide for ease of tracking or purification, such as a fluorescent protein, a purification tag, or an epitope tag. Examples of fluorescent proteins include green fluorescent proteins (e.g., GFP, GFP-2, tagGFP,

turboGFP, eGFP, Emerald, Azami Green, Monomeric Azami Green, CopGFP, AceGFP, ZsGreen1), yellow fluorescent proteins (e.g., YFP, eYFP, Citrine, Venus, YPet, PhiYFP, ZsYellow1), blue fluorescent proteins (e.g., eBFP, eBFP2, Azurite, mKalamal, GFPuv, Sapphire, T-sapphire), cyan fluorescent proteins (e.g., eCFP, Cerulean, CyPet, AmCyan1, Midoriishi-Cyan), red fluorescent proteins (e.g., mKate, mKate2, mPlum, DsRed monomer, mCherry, mRFP1, DsRed-Express, DsRed2, DsRed-Monomer, HcRed-Tandem, HcRed1, AsRed2, eqFP611, mRaspberry, mStrawberry, Jred), orange fluorescent proteins (e.g., mOrange, mKO, Kusabira-Orange, Monomeric Kusabira-Orange, mTangerine, tdTomato), and any other suitable fluorescent protein. Examples of tags include glutathione-S-transferase (GST), chitin binding protein (CBP), maltose binding protein, thioredoxin (TRX), poly(NANP), tandem affinity purification (TAP) tag, myc, AcV5, AU1, AU5, E, ECS, E2, FLAG, hemagglutinin (HA), nus, Softag 1, Softag 3, Strep, SBP, Glu-Glu, HSV, KT3, S, S1, T7, V5, VSV-G, histidine (His), biotin carboxyl carrier protein (BCCP), and calmodulin.

[00138] A nucleic acid encoding a Cas protein can be codon optimized for efficient translation into protein in a particular cell or organism. For example, the nucleic acid encoding the Cas protein can be modified to substitute codons having a higher frequency of usage in a human cell, a non-human cell, a mammalian cell, a rodent cell, a mouse cell, a rat cell, or any other host cell of interest, as compared to the naturally occurring polynucleotide sequence.

(3) *Guide RNAs*

[00139] A “guide RNA” or “gRNA” is an RNA molecule that binds to a Cas protein (e.g., Cas9 protein) and targets the Cas protein to a specific location within a target DNA. Guide RNAs can comprise two segments: a “DNA-targeting segment” and a “protein-binding segment.” “Segment” includes a section or region of a molecule, such as a contiguous stretch of nucleotides in an RNA. Some gRNAs, such as those for Cas9, can comprise two separate RNA molecules: an “activator-RNA” (e.g., tracrRNA) and a “targeter-RNA” (e.g., CRISPR RNA or crRNA). Other gRNAs are a single RNA molecule (single RNA polynucleotide), which can also be called a “single-molecule gRNA,” a “single-guide RNA,” or an “sgRNA.” *See, e.g.,* WO 2013/176772, WO 2014/065596, WO 2014/089290, WO 2014/093622, WO 2014/099750, WO 2013/142578, and WO 2014/131833, each of which is herein incorporated by reference in its entirety for all purposes. For Cas9, for example, a single-guide RNA can comprise a crRNA

fused to a tracrRNA (e.g., via a linker). For Cpf1, for example, only a crRNA is needed to achieve binding to and/or cleavage of a target sequence. The terms “guide RNA” and “gRNA” include both double-molecule (i.e., modular) gRNAs and single-molecule gRNAs.

[00140] An exemplary two-molecule gRNA comprises a crRNA-like (“CRISPR RNA” or “targeter-RNA” or “crRNA” or “crRNA repeat”) molecule and a corresponding tracrRNA-like (“trans-acting CRISPR RNA” or “activator-RNA” or “tracrRNA”) molecule. A crRNA comprises both the DNA-targeting segment (single-stranded) of the gRNA and a stretch of nucleotides (i.e., the crRNA tail) that forms one half of the dsRNA duplex of the protein-binding segment of the gRNA. An example of a crRNA tail, located downstream (3’) of the DNA-targeting segment, comprises, consists essentially of, or consists of GUUUUAGAGCUAUGCU (SEQ ID NO: 25). Any of the DNA-targeting segments disclosed herein can be joined to the 5’ end of SEQ ID NO: 25 to form a crRNA.

[00141] A corresponding tracrRNA (activator-RNA) comprises a stretch of nucleotides that forms the other half of the dsRNA duplex of the protein-binding segment of the gRNA. A stretch of nucleotides of a crRNA are complementary to and hybridize with a stretch of nucleotides of a tracrRNA to form the dsRNA duplex of the protein-binding domain of the gRNA. As such, each crRNA can be said to have a corresponding tracrRNA. An example of a tracrRNA sequence comprises, consists essentially of, or consists of AGCAUAGCAAGUUA AAAUAAGGCUAGUCCGUUAUCAACUUGAAAAAGUGGCACC GAGUCGGUGCUUU (SEQ ID NO: 26).

[00142] In systems in which both a crRNA and a tracrRNA are needed, the crRNA and the corresponding tracrRNA hybridize to form a gRNA. In systems in which only a crRNA is needed, the crRNA can be the gRNA. The crRNA additionally provides the single-stranded DNA-targeting segment that targets a guide RNA target sequence by hybridizing to the opposite strand (i.e., the complementary strand). If used for modification within a cell, the exact sequence of a given crRNA or tracrRNA molecule can be designed to be specific to the species in which the RNA molecules will be used. *See, e.g., Mali et al. (2013) Science 339:823-826; Jinek et al. (2012) Science 337:816-821; Hwang et al. (2013) Nat. Biotechnol. 31:227-229; Jiang et al. (2013) Nat. Biotechnol. 31:233-239; and Cong et al. (2013) Science 339:819-823, each of which is herein incorporated by reference in its entirety for all purposes.*

[00143] The DNA-targeting segment (crRNA) of a given gRNA comprises a nucleotide sequence that is complementary to a sequence (i.e., the complementary strand of the guide RNA recognition sequence on the strand opposite of the guide RNA target sequence) in a target DNA. The DNA-targeting segment of a gRNA interacts with a target DNA in a sequence-specific manner via hybridization (i.e., base pairing). As such, the nucleotide sequence of the DNA-targeting segment may vary and determines the location within the target DNA with which the gRNA and the target DNA will interact. The DNA-targeting segment of a subject gRNA can be modified to hybridize to any desired sequence within a target DNA. Naturally occurring crRNAs differ depending on the CRISPR/Cas system and organism but often contain a targeting segment of between 21 to 72 nucleotides length, flanked by two direct repeats (DR) of a length of between 21 to 46 nucleotides (*see, e.g.*, WO 2014/131833, herein incorporated by reference in its entirety for all purposes). In the case of *S. pyogenes*, the DRs are 36 nucleotides long and the targeting segment is 30 nucleotides long. The 3' located DR is complementary to and hybridizes with the corresponding tracrRNA, which in turn binds to the Cas protein.

[00144] The DNA-targeting segment can have a length of at least about 12 nucleotides, at least about 15 nucleotides, at least about 17 nucleotides, at least about 18 nucleotides, at least about 19 nucleotides, at least about 20 nucleotides, at least about 25 nucleotides, at least about 30 nucleotides, at least about 35 nucleotides, or at least about 40 nucleotides. Such DNA-targeting segments can have a length from about 12 nucleotides to about 100 nucleotides, from about 12 nucleotides to about 80 nucleotides, from about 12 nucleotides to about 50 nucleotides, from about 12 nucleotides to about 40 nucleotides, from about 12 nucleotides to about 30 nucleotides, from about 12 nucleotides to about 25 nucleotides, or from about 12 nucleotides to about 20 nucleotides. For example, the DNA targeting segment can be from about 15 nucleotides to about 25 nucleotides (*e.g.*, from about 17 nucleotides to about 20 nucleotides, or about 17 nucleotides, about 18 nucleotides, about 19 nucleotides, or about 20 nucleotides). *See, e.g.*, US 2016/0024523, herein incorporated by reference in its entirety for all purposes. For Cas9 from *S. pyogenes*, a typical DNA-targeting segment is between 16 and 20 nucleotides in length or between 17 and 20 nucleotides in length. For Cas9 from *S. aureus*, a typical DNA-targeting segment is between 21 and 23 nucleotides in length. For Cpf1, a typical DNA-targeting segment is at least 16 nucleotides in length or at least 18 nucleotides in length.

[00145] TracrRNAs can be in any form (e.g., full-length tracrRNAs or active partial tracrRNAs) and of varying lengths. They can include primary transcripts or processed forms. For example, tracrRNAs (as part of a single-guide RNA or as a separate molecule as part of a two-molecule gRNA) may comprise, consist essentially of, or consist of all or a portion of a wild type tracrRNA sequence (e.g., about or more than about 20, 26, 32, 45, 48, 54, 63, 67, 85, or more nucleotides of a wild type tracrRNA sequence). Examples of wild type tracrRNA sequences from *S. pyogenes* include 171-nucleotide, 89-nucleotide, 75-nucleotide, and 65-nucleotide versions. See, e.g., Deltcheva *et al.* (2011) *Nature* 471:602-607; WO 2014/093661, each of which is herein incorporated by reference in its entirety for all purposes. Examples of tracrRNAs within single-guide RNAs (sgRNAs) include the tracrRNA segments found within +48, +54, +67, and +85 versions of sgRNAs, where “+n” indicates that up to the +n nucleotide of wild type tracrRNA is included in the sgRNA. See US 8,697,359, herein incorporated by reference in its entirety for all purposes.

[00146] The percent complementarity between the DNA-targeting sequence and the complementary strand of the guide RNA recognition sequence within the target DNA can be at least 60% (e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100%). The percent complementarity between the DNA-targeting sequence and the complementary strand of the guide RNA recognition sequence within the target DNA can be at least 60% over about 20 contiguous nucleotides. As an example, the percent complementarity between the DNA-targeting sequence and the complementary strand of the guide RNA recognition sequence within the target DNA is 100% over the 14 contiguous nucleotides at the 5' end of the complementary strand of the guide RNA recognition sequence within the complementary strand of the target DNA and as low as 0% over the remainder. In such a case, the DNA-targeting sequence can be considered to be 14 nucleotides in length. As another example, the percent complementarity between the DNA-targeting sequence and the complementary strand of the guide RNA recognition sequence within the target DNA is 100% over the seven contiguous nucleotides at the 5' end of the complementary strand of the guide RNA recognition sequence within the complementary strand of the target DNA and as low as 0% over the remainder. In such a case, the DNA-targeting sequence can be considered to be 7 nucleotides in length. In some guide RNAs, at least 17 nucleotides within the DNA-targeting sequence are complementary to the target DNA. For

example, the DNA-targeting sequence can be 20 nucleotides in length and can comprise 1, 2, or 3 mismatches with the complementary strand of the guide RNA recognition sequence.

Optionally, the mismatches are not adjacent to a protospacer adjacent motif (PAM) sequence (e.g., the mismatches are in the 5' end of the DNA-targeting sequence, or the mismatches are at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 base pairs away from the PAM sequence).

[00147] The protein-binding segment of a gRNA can comprise two stretches of nucleotides that are complementary to one another. The complementary nucleotides of the protein-binding segment hybridize to form a double-stranded RNA duplex (dsRNA). The protein-binding segment of a subject gRNA interacts with a Cas protein, and the gRNA directs the bound Cas protein to a specific nucleotide sequence within target DNA via the DNA-targeting segment.

[00148] Single-guide RNAs have the DNA-targeting segment and a scaffold sequence (i.e., the protein-binding or Cas-binding sequence of the guide RNA). For example, such guide RNAs have a 5' DNA-targeting segment and a 3' scaffold sequence. Exemplary scaffold sequences comprise, consist essentially of, or consist of:

GUUUUAGAGCUAGAAAUAGCAAGUUAAAAUAAGGCUAGUCCGUUAUCAACUUGA
AAAAGUGGCACCGAGUCGGUGCU (version 1; SEQ ID NO: 27);

GUUGGAACCAUUCAAAACAGCAUAGCAAGUUAAAAUAAGGCUAGUCCGUUAUCA
ACUUGAAAAAGUGGCACCGAGUCGGUGC (version 2; SEQ ID NO: 6);

GUUUUAGAGCUAGAAAUAGCAAGUUAAAAUAAGGCUAGUCCGUUAUCAACUUGA
AAAAGUGGCACCGAGUCGGUGC (version 3; SEQ ID NO: 7); and

GUUUUAGAGCUAUGCUGGAAACAGCAUAGCAAGUUUAAAUAAGGCUAGUCCGUU
AUCAACUUGAAAAAGUGGCACCGAGUCGGUGC (version 4; SEQ ID NO: 8). Guide

RNAs targeting any guide RNA target sequence can include, for example, a DNA-targeting segment on the 5' end of the guide RNA fused to any of the exemplary guide RNA scaffold sequences on the 3' end of the guide RNA. That is, any of the DNA-targeting segments disclosed herein can be joined to the 5' end of any one of SEQ ID NOS: 27, 6, 7, or 8 to form a single guide RNA (chimeric guide RNA). Guide RNA versions 1, 2, 3, and 4 as disclosed elsewhere herein refer to DNA-targeting segments joined with scaffold versions 1, 2, 3, and 4, respectively.

[00149] Guide RNAs can include modifications or sequences that provide for additional desirable features (e.g., modified or regulated stability; subcellular targeting; tracking with a fluorescent label; a binding site for a protein or protein complex; and the like). Examples of such modifications include, for example, a 5' cap (e.g., a 7-methylguanylate cap (m7G)); a 3' polyadenylated tail (i.e., a 3' poly(A) tail); a riboswitch sequence (e.g., to allow for regulated stability and/or regulated accessibility by proteins and/or protein complexes); a stability control sequence; a sequence that forms a dsRNA duplex (i.e., a hairpin); a modification or sequence that targets the RNA to a subcellular location (e.g., nucleus, mitochondria, chloroplasts, and the like); a modification or sequence that provides for tracking (e.g., direct conjugation to a fluorescent molecule, conjugation to a moiety that facilitates fluorescent detection, a sequence that allows for fluorescent detection, and so forth); a modification or sequence that provides a binding site for proteins (e.g., proteins that act on DNA, including DNA methyltransferases, DNA demethylases, histone acetyltransferases, histone deacetylases, and the like); and combinations thereof. Other examples of modifications include engineered stem loop duplex structures, engineered bulge regions, engineered hairpins 3' of the stem loop duplex structure, or any combination thereof. *See, e.g.*, US 2015/0376586, herein incorporated by reference in its entirety for all purposes. A bulge can be an unpaired region of nucleotides within the duplex made up of the crRNA-like region and the minimum tracrRNA-like region. A bulge can comprise, on one side of the duplex, an unpaired 5'-XXX-3' where X is any purine and Y can be a nucleotide that can form a wobble pair with a nucleotide on the opposite strand, and an unpaired nucleotide region on the other side of the duplex.

[00150] Unmodified nucleic acids can be prone to degradation. Exogenous nucleic acids can also induce an innate immune response. Modifications can help introduce stability and reduce immunogenicity. Guide RNAs can comprise modified nucleosides and modified nucleotides including, for example, one or more of the following: (1) alteration or replacement of one or both of the non-linking phosphate oxygens and/or of one or more of the linking phosphate oxygens in the phosphodiester backbone linkage; (2) alteration or replacement of a constituent of the ribose sugar such as alteration or replacement of the 2' hydroxyl on the ribose sugar; (3) replacement of the phosphate moiety with dephospho linkers; (4) modification or replacement of a naturally occurring nucleobase; (5) replacement or modification of the ribose-phosphate backbone; (6) modification of the 3' end or 5' end of the oligonucleotide (e.g., removal, modification or

replacement of a terminal phosphate group or conjugation of a moiety); and (7) modification of the sugar. Other possible guide RNA modifications include modifications of or replacement of uracils or poly-uracil tracts. *See, e.g.*, WO 2015/048577 and US 2016/0237455, each of which is herein incorporated by reference in its entirety for all purposes. Similar modifications can be made to Cas-encoding nucleic acids, such as Cas mRNAs.

[00151] As one example, nucleotides at the 5' or 3' end of a guide RNA can include phosphorothioate linkages (e.g., the bases can have a modified phosphate group that is a phosphorothioate group). For example, a guide RNA can include phosphorothioate linkages between the 2, 3, or 4 terminal nucleotides at the 5' or 3' end of the guide RNA. As another example, nucleotides at the 5' and/or 3' end of a guide RNA can have 2'-O-methyl modifications. For example, a guide RNA can include 2'-O-methyl modifications at the 2, 3, or 4 terminal nucleotides at the 5' and/or 3' end of the guide RNA (e.g., the 5' end). *See, e.g.*, WO 2017/173054 A1 and Finn et al. (2018) *Cell Reports* 22:1-9, each of which is herein incorporated by reference in its entirety for all purposes. In one specific example, the guide RNA comprises 2'-O-methyl analogs and 3' phosphorothioate internucleotide linkages at the first three 5' and 3' terminal RNA residues. In another specific example, the guide RNA is modified such that all 2'OH groups that do not interact with the Cas9 protein are replaced with 2'-O-methyl analogs, and the tail region of the guide RNA, which has minimal interaction with Cas9, is modified with 5' and 3' phosphorothioate internucleotide linkages. *See, e.g.*, Yin et al. (2017) *Nat. Biotech.* 35(12):1179-1187, herein incorporated by reference in its entirety for all purposes.

[00152] Guide RNAs can be provided in any form. For example, the gRNA can be provided in the form of RNA, either as two molecules (separate crRNA and tracrRNA) or as one molecule (sgRNA), and optionally in the form of a complex with a Cas protein. The gRNA can also be provided in the form of DNA encoding the gRNA. The DNA encoding the gRNA can encode a single RNA molecule (sgRNA) or separate RNA molecules (e.g., separate crRNA and tracrRNA). In the latter case, the DNA encoding the gRNA can be provided as one DNA molecule or as separate DNA molecules encoding the crRNA and tracrRNA, respectively.

[00153] When a gRNA is provided in the form of DNA, the gRNA can be transiently, conditionally, or constitutively expressed in the cell. DNAs encoding gRNAs can be stably integrated into the genome of the cell and operably linked to a promoter active in the cell. Alternatively, DNAs encoding gRNAs can be operably linked to a promoter in an expression

construct. For example, the DNA encoding the gRNA can be in a vector comprising a heterologous nucleic acid, such as a nucleic acid encoding a Cas protein. Alternatively, it can be in a vector or a plasmid that is separate from the vector comprising the nucleic acid encoding the Cas protein. Promoters that can be used in such expression constructs include promoters active, for example, in one or more of a eukaryotic cell, a human cell, a non-human cell, a mammalian cell, a non-human mammalian cell, a rodent cell, a mouse cell, a rat cell, a hamster cell, a rabbit cell, a pluripotent cell, an embryonic stem (ES) cell, an adult stem cell, a developmentally restricted progenitor cell, an induced pluripotent stem (iPS) cell, or a one-cell stage embryo. Such promoters can be, for example, conditional promoters, inducible promoters, constitutive promoters, or tissue-specific promoters. Such promoters can also be, for example, bidirectional promoters. Specific examples of suitable promoters include an RNA polymerase III promoter, such as a human U6 promoter, a rat U6 polymerase III promoter, or a mouse U6 polymerase III promoter.

[00154] Alternatively, gRNAs can be prepared by various other methods. For example, gRNAs can be prepared by *in vitro* transcription using, for example, T7 RNA polymerase (*see, e.g.*, WO 2014/089290 and WO 2014/065596, each of which is herein incorporated by reference in its entirety for all purposes). Guide RNAs can also be a synthetically produced molecule prepared by chemical synthesis.

(4) Guide RNA Recognition Sequences and Guide RNA Target Sequences

[00155] The term “guide RNA recognition sequence” includes nucleic acid sequences present in a target DNA to which a DNA-targeting segment of a gRNA will bind, provided sufficient conditions for binding exist. The term guide RNA recognition sequence as used herein encompasses both strands of the target double-stranded DNA (i.e., the sequence on the complementary strand to which the guide RNA hybridizes and the corresponding sequence on the non-complementary strand adjacent to the protospacer adjacent motif (PAM)). The term “guide RNA target sequence” as used herein refers specifically to the sequence on the non-complementary strand adjacent to the PAM (i.e., upstream or 5' of the PAM). That is, the guide RNA target sequence refers to the sequence on the non-complementary strand corresponding to the sequence to which the guide RNA hybridizes on the complementary strand. A guide RNA target sequence is equivalent to the DNA-targeting segment of a guide RNA, but with thymines

instead of uracils. As one example, a guide RNA target sequence for a Cas9 enzyme would refer to the sequence on the non-complementary strand adjacent to the 5'-NGG-3' PAM. Guide RNA recognition sequences include sequences to which a guide RNA is designed to have complementarity, where hybridization between the complementary strand of a guide RNA recognition sequence and a DNA-targeting sequence of a guide RNA promotes the formation of a CRISPR complex. Full complementarity is not necessarily required, provided that there is sufficient complementarity to cause hybridization and promote formation of a CRISPR complex. Guide RNA recognition sequences or guide RNA target sequences also include cleavage sites for Cas proteins, described in more detail below. A guide RNA recognition sequence or a guide RNA target sequence can comprise any polynucleotide, which can be located, for example, in the nucleus or cytoplasm of a cell or within an organelle of a cell, such as a mitochondrion or chloroplast.

[00156] The guide RNA recognition sequence within a target DNA can be targeted by (i.e., be bound by, or hybridize with, or be complementary to) a Cas protein or a gRNA. Suitable DNA/RNA binding conditions include physiological conditions normally present in a cell. Other suitable DNA/RNA binding conditions (e.g., conditions in a cell-free system) are known (*see, e.g., Molecular Cloning: A Laboratory Manual*, 3rd Ed. (Sambrook *et al.*, Harbor Laboratory Press 2001), herein incorporated by reference in its entirety for all purposes). The strand of the target DNA that is complementary to and hybridizes with the Cas protein or gRNA can be called the “complementary strand,” and the strand of the target DNA that is complementary to the “complementary strand” (and is therefore not complementary to the Cas protein or gRNA) can be called “non-complementary strand” or “template strand.”

[00157] The Cas protein can cleave the nucleic acid at a site within or outside of the nucleic acid sequence present in the target DNA to which the DNA-targeting segment of a gRNA will bind. The “cleavage site” includes the position of a nucleic acid at which a Cas protein produces a single-strand break or a double-strand break. For example, formation of a CRISPR complex (comprising a gRNA hybridized to the complementary strand of a guide RNA recognition sequence and complexed with a Cas protein) can result in cleavage of one or both strands in or near (e.g., within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from) the nucleic acid sequence present in a target DNA to which a DNA-targeting segment of a gRNA will bind. If the cleavage site is outside of the nucleic acid sequence to which the DNA-targeting segment of

the gRNA will bind, the cleavage site is still considered to be within the “guide RNA recognition sequence” or guide RNA target sequence. The cleavage site can be on only one strand or on both strands of a nucleic acid. Cleavage sites can be at the same position on both strands of the nucleic acid (producing blunt ends) or can be at different sites on each strand (producing staggered ends (i.e., overhangs)). Staggered ends can be produced, for example, by using two Cas proteins, each of which produces a single-strand break at a different cleavage site on a different strand, thereby producing a double-strand break. For example, a first nickase can create a single-strand break on the first strand of double-stranded DNA (dsDNA), and a second nickase can create a single-strand break on the second strand of dsDNA such that overhanging sequences are created. In some cases, the guide RNA recognition sequence or guide RNA target sequence of the nickase on the first strand is separated from the guide RNA recognition sequence or guide RNA target sequence of the nickase on the second strand by at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 75, 100, 250, 500, or 1,000 base pairs.

[00158] Site-specific binding and/or cleavage of target DNA by Cas proteins can occur at locations determined by both (i) base-pairing complementarity between the gRNA and the target DNA and (ii) a short motif, called the protospacer adjacent motif (PAM), in the target DNA. The PAM can flank the guide RNA target sequence on the non-complementary strand opposite of the strand to which the guide RNA hybridizes. Optionally, the guide RNA target sequence can be flanked on the 3' end by the PAM. Alternatively, the guide RNA target sequence can be flanked on the 5' end by the PAM. For example, the cleavage site of Cas proteins can be about 1 to about 10 or about 2 to about 5 base pairs (e.g., 3 base pairs) upstream or downstream of the PAM sequence. In some cases (e.g., when Cas9 from *S. pyogenes* or a closely related Cas9 is used), the PAM sequence of the non-complementary strand can be 5'-N₁GG-3', where N₁ is any DNA nucleotide and is immediately 3' of the guide RNA recognition sequence of the non-complementary strand of the target DNA (i.e., immediately 3' of the guide RNA target sequence). As such, the PAM sequence of the complementary strand would be 5'-CCN₂-3', where N₂ is any DNA nucleotide and is immediately 5' of the guide RNA recognition sequence of the complementary strand of the target DNA. In some such cases, N₁ and N₂ can be complementary and the N₁-N₂ base pair can be any base pair (e.g., N₁=C and N₂=G; N₁=G and N₂=C; N₁=A and N₂=T; or N₁=T, and N₂=A). In the case of Cas9 from *S. aureus*, the PAM can be NNGRRT or NNGRR, where N can A, G, C, or T, and R can be G or A. In the case of Cas9

from *C. jejuni*, the PAM can be, for example, NNNNACAC or NNNNRYAC, where N can be A, G, C, or T, and R can be G or A. In some cases (e.g., for FnCpf1), the PAM sequence can be upstream of the 5' end and have the sequence 5'-TTN-3'.

[00159] Examples of guide RNA target sequences or guide RNA target sequences in addition to a PAM sequence are provided below. For example, the guide RNA target sequence can be a 20-nucleotide DNA sequence immediately preceding an NGG motif recognized by a Cas9 protein. Examples of such guide RNA target sequences plus a PAM sequence are GN₁₉NGG (SEQ ID NO: 9) or N₂₀NGG (SEQ ID NO: 10). *See, e.g.*, WO 2014/165825, herein incorporated by reference in its entirety for all purposes. The guanine at the 5' end can facilitate transcription by RNA polymerase in cells. Other examples of guide RNA target sequences plus a PAM sequence can include two guanine nucleotides at the 5' end (e.g., GGN₂₀NGG; SEQ ID NO: 11) to facilitate efficient transcription by T7 polymerase *in vitro*. *See, e.g.*, WO 2014/065596, herein incorporated by reference in its entirety for all purposes. Other guide RNA target sequences plus a PAM sequence can have between 4-22 nucleotides in length of SEQ ID NOS: 9-11, including the 5' G or GG and the 3' GG or NGG. Yet other guide RNA target sequences can have between 14 and 20 nucleotides in length of SEQ ID NOS: 9-11.

[00160] The guide RNA recognition sequence or guide RNA target sequence can be any nucleic acid sequence endogenous or exogenous to a cell. The guide RNA recognition sequence or guide RNA target sequence can be a sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory sequence) or can include both.

B. Cells and Non-Human Animals Comprising Cas Expression Cassettes

[00161] Cells and non-human animals comprising the Cas expression cassettes described herein are also provided. The Cas expression cassette can be stably integrated into the genome (i.e., into a chromosome) of the cell or non-human animal or it can be located outside of a chromosome (e.g., extrachromosomally replicating DNA). Optionally, the Cas expression cassette is stably integrated into the genome. The stably integrated Cas expression cassette can be randomly integrated into the genome of the non-human animal (i.e., transgenic), or it can be integrated into a predetermined region of the genome of the non-human animal (i.e., knock in). Optionally, the Cas expression cassette is stably integrated into a predetermined region of the genome, such as a safe harbor locus. The target genomic locus at which the Cas expression

cassette is stably integrated can be heterozygous for the Cas expression cassette or homozygous for the Cas expression cassette. A diploid organism has two alleles at each genetic locus. Each pair of alleles represents the genotype of a specific genetic locus. Genotypes are described as homozygous if there are two identical alleles at a particular locus and as heterozygous if the two alleles differ.

[00162] The cells provided herein can be, for example, eukaryotic cells, which include, for example, fungal cells (e.g., yeast), plant cells, animal cells, mammalian cells, non-human mammalian cells, and human cells. The term “animal” includes mammals, fishes, and birds. A mammalian cell can be, for example, a non-human mammalian cell, a human cell, a rodent cell, a rat cell, a mouse cell, or a hamster cell. Other non-human mammals include, for example, non-human primates, monkeys, apes, cats, dogs, rabbits, horses, bulls, deer, bison, livestock (e.g., bovine species such as cows, steer, and so forth; ovine species such as sheep, goats, and so forth; and porcine species such as pigs and boars). Birds include, for example, chickens, turkeys, ostrich, geese, ducks, and so forth. Domesticated animals and agricultural animals are also included. The term “non-human” excludes humans.

[00163] The cells can also be any type of undifferentiated or differentiated state. For example, a cell can be a totipotent cell, a pluripotent cell (e.g., a human pluripotent cell or a non-human pluripotent cell such as a mouse embryonic stem (ES) cell or a rat ES cell), or a non-pluripotent cell. Totipotent cells include undifferentiated cells that can give rise to any cell type, and pluripotent cells include undifferentiated cells that possess the ability to develop into more than one differentiated cell types. Such pluripotent and/or totipotent cells can be, for example, ES cells or ES-like cells, such as an induced pluripotent stem (iPS) cells. ES cells include embryo-derived totipotent or pluripotent cells that are capable of contributing to any tissue of the developing embryo upon introduction into an embryo. ES cells can be derived from the inner cell mass of a blastocyst and are capable of differentiating into cells of any of the three vertebrate germ layers (endoderm, ectoderm, and mesoderm).

[00164] Examples of human pluripotent cells include human ES cells, human adult stem cells, developmentally restricted human progenitor cells, and human induced pluripotent stem (iPS) cells, such as primed human iPS cells and naïve human iPS cells. Induced pluripotent stem cells include pluripotent stem cells that can be derived directly from a differentiated adult cell. Human iPS cells can be generated by introducing specific sets of reprogramming factors into a

cell which can include, for example, Oct3/4, Sox family transcription factors (e.g., Sox1, Sox2, Sox3, Sox15), Myc family transcription factors (e.g., c-Myc, l-Myc, n-Myc), Krüppel-like family (KLF) transcription factors (e.g., KLF1, KLF2, KLF4, KLF5), and/or related transcription factors, such as NANOG, LIN28, and/or Glis1. Human iPS cells can also be generated, for example, by the use of miRNAs, small molecules that mimic the actions of transcription factors, or lineage specifiers. Human iPS cells are characterized by their ability to differentiate into any cell of the three vertebrate germ layers, e.g., the endoderm, the ectoderm, or the mesoderm. Human iPS cells are also characterized by their ability propagate indefinitely under suitable *in vitro* culture conditions. *See, e.g.,* Takahashi and Yamanaka (2006) *Cell* 126:663-676, herein incorporated by reference in its entirety for all purposes. Primed human ES cells and primed human iPS cells include cells that express characteristics similar to those of post-implantation epiblast cells and are committed for lineage specification and differentiation. Naïve human ES cells and naïve human iPS cells include cells that express characteristics similar to those of ES cells of the inner cell mass of a pre-implantation embryo and are not committed for lineage specification. *See, e.g.,* Nichols and Smith (2009) *Cell Stem Cell* 4:487-492, herein incorporated by reference in its entirety for all purposes.

[00165] The cells provided herein can also be germ cells (e.g., sperm or oocytes). The cells can be mitotically competent cells or mitotically-inactive cells, meiotically competent cells or meiotically-inactive cells. Similarly, the cells can also be primary somatic cells or cells that are not a primary somatic cell. Somatic cells include any cell that is not a gamete, germ cell, gametocyte, or undifferentiated stem cell. For example, the cells can be liver cells, kidney cells, hematopoietic cells, endothelial cells, epithelial cells, fibroblasts, mesenchymal cells, keratinocytes, blood cells, melanocytes, monocytes, mononuclear cells, monocytic precursors, B cells, erythroid-megakaryocytic cells, eosinophils, macrophages, T cells, islet beta cells, exocrine cells, pancreatic progenitors, endocrine progenitors, adipocytes, preadipocytes, neurons, glial cells, neural stem cells, neurons, hepatoblasts, hepatocytes, cardiomyocytes, skeletal myoblasts, smooth muscle cells, ductal cells, acinar cells, alpha cells, beta cells, delta cells, PP cells, cholangiocytes, white or brown adipocytes, or ocular cells (e.g., trabecular meshwork cells, retinal pigment epithelial cells, retinal microvascular endothelial cells, retinal pericyte cells, conjunctival epithelial cells, conjunctival fibroblasts, iris pigment epithelial cells, keratocytes, lens epithelial cells, non-pigment ciliary epithelial cells, ocular choroid fibroblasts, photoreceptor

cells, ganglion cells, bipolar cells, horizontal cells, or amacrine cells).

[00166] Suitable cells provided herein also include primary cells. Primary cells include cells or cultures of cells that have been isolated directly from an organism, organ, or tissue. Primary cells include cells that are neither transformed nor immortal. They include any cell obtained from an organism, organ, or tissue which was not previously passed in tissue culture or has been previously passed in tissue culture but is incapable of being indefinitely passed in tissue culture. Such cells can be isolated by conventional techniques and include, for example, somatic cells, hematopoietic cells, endothelial cells, epithelial cells, fibroblasts, mesenchymal cells, keratinocytes, melanocytes, monocytes, mononuclear cells, adipocytes, preadipocytes, neurons, glial cells, hepatocytes, skeletal myoblasts, and smooth muscle cells. For example, primary cells can be derived from connective tissues, muscle tissues, nervous system tissues, or epithelial tissues.

[00167] Other suitable cells provided herein include immortalized cells. Immortalized cells include cells from a multicellular organism that would normally not proliferate indefinitely but, due to mutation or alteration, have evaded normal cellular senescence and instead can keep undergoing division. Such mutations or alterations can occur naturally or be intentionally induced. Examples of immortalized cells include Chinese hamster ovary (CHO) cells, human embryonic kidney cells (e.g., HEK 293 cells or 293T cells), and mouse embryonic fibroblast cells (e.g., 3T3 cells). Numerous types of immortalized cells are well known. Immortalized or primary cells include cells that are typically used for culturing or for expressing recombinant genes or proteins.

[00168] The cells provided herein also include one-cell stage embryos (i.e., fertilized oocytes or zygotes). Such one-cell stage embryos can be from any genetic background (e.g., BALB/c, C57BL/6, 129, or a combination thereof for mice), can be fresh or frozen, and can be derived from natural breeding or *in vitro* fertilization.

[00169] The cells provided herein can be normal, healthy cells, or can be diseased or mutant-bearing cells.

[00170] Non-human animals comprising a Cas expression cassette as described herein can be made by the methods described elsewhere herein. The term “animal” includes mammals, fishes, and birds. Mammals include, for example, humans, non-human primates, monkeys, apes, cats, dogs, horses, bulls, deer, bison, sheep, rabbits, rodents (e.g., mice, rats, hamsters, and guinea

pigs), and livestock (e.g., bovine species such as cows and steer; ovine species such as sheep and goats; and porcine species such as pigs and boars). Birds include, for example, chickens, turkeys, ostrich, geese, and ducks. Domesticated animals and agricultural animals are also included. The term “non-human animal” excludes humans. Preferred non-human animals include, for example, rodents, such as mice and rats.

[00171] The non-human animals can be from any genetic background. For example, suitable mice can be from a 129 strain, a C57BL/6 strain, a mix of 129 and C57BL/6, a BALB/c strain, or a Swiss Webster strain. Examples of 129 strains include 129P1, 129P2, 129P3, 129X1, 129S1 (e.g., 129S1/SV, 129S1/SvIm), 129S2, 129S4, 129S5, 129S9/SvEvH, 129S6 (129/SvEvTac), 129S7, 129S8, 129T1, and 129T2. *See, e.g., Festing et al. (1999) Mammalian Genome 10:836, herein incorporated by reference in its entirety for all purposes.* Examples of C57BL strains include C57BL/A, C57BL/An, C57BL/GrFa, C57BL/KaL_wN, C57BL/6, C57BL/6J, C57BL/6ByJ, C57BL/6NJ, C57BL/10, C57BL/10ScSn, C57BL/10Cr, and C57BL/Ola. Suitable mice can also be from a mix of an aforementioned 129 strain and an aforementioned C57BL/6 strain (e.g., 50% 129 and 50% C57BL/6). Likewise, suitable mice can be from a mix of aforementioned 129 strains or a mix of aforementioned BL/6 strains (e.g., the 129S6 (129/SvEvTac) strain).

[00172] Similarly, rats can be from any rat strain, including, for example, an ACI rat strain, a Dark Agouti (DA) rat strain, a Wistar rat strain, a LEA rat strain, a Sprague Dawley (SD) rat strain, or a Fischer rat strain such as Fisher F344 or Fisher F6. Rats can also be obtained from a strain derived from a mix of two or more strains recited above. For example, a suitable rat can be from a DA strain or an ACI strain. The ACI rat strain is characterized as having black agouti, with white belly and feet and an *RT1^{avl}* haplotype. Such strains are available from a variety of sources including Harlan Laboratories. The Dark Agouti (DA) rat strain is characterized as having an agouti coat and an *RT1^{avl}* haplotype. Such rats are available from a variety of sources including Charles River and Harlan Laboratories. In some cases, suitable rats can be from an inbred rat strain. *See, e.g., US 2014/0235933, herein incorporated by reference in its entirety for all purposes.*

C. Target Genomic Loci

[00173] The Cas expression cassettes described herein can be genomically integrated at a

target genomic locus in a cell or a non-human animal. Any target genomic locus capable of expressing a gene can be used.

[00174] An example of a target genomic locus into which the Cas expression cassettes described herein can be stably integrated is a safe harbor locus in the genome of the non-human animal. Interactions between integrated exogenous DNA and a host genome can limit the reliability and safety of integration and can lead to overt phenotypic effects that are not due to the targeted genetic modification but are instead due to unintended effects of the integration on surrounding endogenous genes. For example, randomly inserted transgenes can be subject to position effects and silencing, making their expression unreliable and unpredictable. Likewise, integration of exogenous DNA into a chromosomal locus can affect surrounding endogenous genes and chromatin, thereby altering cell behavior and phenotypes. Safe harbor loci include chromosomal loci where transgenes or other exogenous nucleic acid inserts can be stably and reliably expressed in all tissues of interest without overtly altering cell behavior or phenotype (i.e., without any deleterious effects on the host cell). *See, e.g., Sadelain et al. (2012) Nat. Rev. Cancer* 12:51-58, herein incorporated by reference in its entirety for all purposes. Optionally, the safe harbor locus is one in which expression of the inserted gene sequence is not perturbed by any read-through expression from neighboring genes. For example, safe harbor loci can include chromosomal loci where exogenous DNA can integrate and function in a predictable manner without adversely affecting endogenous gene structure or expression. Safe harbor loci can include extragenic regions or intragenic regions such as, for example, loci within genes that are non-essential, dispensable, or able to be disrupted without overt phenotypic consequences.

[00175] For example, the *Rosa26* locus and its equivalent in humans offer an open chromatin configuration in all tissues and is ubiquitously expressed during embryonic development and in adults. *See, e.g., Zambrowicz et al. (1997) Proc. Natl. Acad. Sci. USA* 94:3789-3794, herein incorporated by reference in its entirety for all purposes. In addition, the *Rosa26* locus can be targeted with high efficiency, and disruption of the *Rosa26* gene produces no overt phenotype. Other examples of safe harbor loci include CCR5, HPRT, AAVS1, and albumin. *See, e.g., US Patent Nos. 7,888,121; 7,972,854; 7,914,796; 7,951,925; 8,110,379; 8,409,861; 8,586,526; and US Patent Publication Nos. 2003/0232410; 2005/0208489; 2005/0026157; 2006/0063231; 2008/0159996; 2010/00218264; 2012/0017290; 2011/0265198; 2013/0137104; 2013/0122591; 2013/0177983; 2013/0177960; and 2013/0122591, each of which is herein incorporated by*

reference in its entirety for all purposes. Biallelic targeting of safe harbor loci such as the *Rosa26* locus has no negative consequences, so different genes or reporters can be targeted to the two *Rosa26* alleles. In one example, a Cas expression cassette is integrated into an intron of the *Rosa26* locus, such as the first intron of the *Rosa26* locus.

D. Recombinase Deleter Non-Human Animals

[00176] Cells or non-human animals comprising a Cas expression cassette comprising a Cas coding sequence downstream of a polyadenylation signal or transcription terminator flanked by recombinase recognition sites recognized by a site-specific recombinase as disclosed herein can further comprise a recombinase expression cassette that drives expression of the site-specific recombinase. Site-specific recombinases include enzymes that can facilitate recombination between recombinase recognition sites, where the two recombination sites are physically separated within a single nucleic acid or on separate nucleic acids. Examples of recombinases include Cre, Flp, and Dre recombinases. One example of a Cre recombinase gene is *Crei*, in which two exons encoding the Cre recombinase are separated by an intron to prevent its expression in a prokaryotic cell. Such recombinases can further comprise a nuclear localization signal to facilitate localization to the nucleus (e.g., NLS-*Crei*). Recombinase recognition sites include nucleotide sequences that are recognized by a site-specific recombinase and can serve as a substrate for a recombination event. Examples of recombinase recognition sites include FRT, FRT11, FRT71, attP, att, rox, and lox sites such as loxP, lox511, lox2272, lox66, lox71, loxM2, and lox5171.

[00177] The Cas expression cassette and the recombinase expression cassette can be integrated at different target genomic loci, or they can be genomically integrated at the same target locus (e.g., a *Rosa26* locus, such as integrated in the first intron of the *Rosa26* locus). For example, the cell or non-human animal can be heterozygous for each of the Cas expression cassette and the recombinase expression cassette, with one allele of the target genomic locus comprising the Cas expression cassette, and a second allele of the target genomic locus comprising the recombinase expression cassette.

[00178] The recombinase gene in a recombinase expression cassette can be operably linked to any suitable promoter. Examples of promoters are disclosed elsewhere herein. For example, the promoter can be a tissue-specific promoter (e.g., albumin promoter for liver-specific expression

or insulin 2 promoter for pancreas-specific expression) or a developmental-stage-specific promoter. The advantage provided by such promoters is that Cas expression can be activated selectively in a desired tissue or only at a desired developmental stage, thereby reducing the possibility of Cas-mediated toxicity *in vivo*. A non-limiting list of exemplary promoters for mouse recombinase delete strains is provided in **Table 2**.

[00179] Table 2. Exemplary Promoters Used in Mouse Recombinase Deleter Strains.

Promoter (Species)	Site of Expression
<i>ACTA1</i> (human)	Adult striated muscle fibers and embryonic striated muscle cells of the somites and heart
<i>Adipoq</i> , adiponectin, C1Q and collagen domain containing (mouse)	White adipose tissue (WAT) and brown adipose tissue (BAT)
<i>Agrp</i> (mouse)	ArGP neurons in the hypothalamus
<i>Alb</i> , albumin (rat)	Liver
<i>Alb1</i> , albumin (mouse)	Liver
<i>Amh</i> (mouse)	Testis Sertoli cells
<i>Aqp2</i> (mouse)	Kidney cells (collecting duct, left) and testes (sperm, right).
<i>Calb2</i> , calbindin 2	Calretinin interneurons in the brain and cortex
<i>Camk2a</i> , calcium/calmodulin-dependent protein kinase II alpha (mouse)	Forebrain, specifically CA1 pyramidal cell layer in hippocampus
<i>Cck</i> , cholecystokinin (mouse)	Cholecystokinin positive neurons (interneurons) of the cortex and in adult spinal cord and embryonic day 15.5 spinal cord and heart
CD2, CD2 molecule (human)	T cells and B cells (all committed B cell and T cell progenitors)
<i>Cd19</i>	B cells
<i>Cdh5</i> , cadherin 5	Endothelium of developing and quiescent vessels, and a subset of hematopoietic cells
<i>Chd16</i> (mouse)	Renal tubules, especially collecting ducts, loops of Henle and distal tubules
<i>Chat</i> , choline acetyltransferase (mouse)	Cholinergic neurons
<i>Ckmm</i> (mouse)	Skeletal and cardiac muscle.
<i>Cort</i> , cortistatin	Cort-expressing cells (CST positive neurons)
<i>Crh</i> , corticotropin releasing hormone	CRH-positive neurons
<i>Cspg4</i> (mouse)	NG2-expressing glia (polydendrocytes, oligodendrocyte progenitor cells) in central nervous system and NG2-expressing cells in other organs; Corpus Callosum; CNS and other tissues such as testes and blood vessels
<i>Cyp39a1</i> , cytochrome P450, family 39, subfamily a, polypeptide 1 (mouse)	Cerebral cortex, hippocampus, striatum, olfactory bulb, and cerebellum
<i>dlx6a</i> , distal-less homeobox gene 6a	GABAergic forebrain neurons
Ella, adenovirus (adenovirus)	Wide range of tissues, including the germ cells that transmit the genetic alteration to progeny
<i>Emx1</i> , empty spiracles homolog 1 (<i>Drosophila</i>)	Neurons of neocortex and hippocampus, and in glial cells of pallium
<i>En1</i> , engrailed 1	Spinal cord V1 interneurons, the embryonic mesencephalon and rhombomere 1 by E9, as well as in the ventral ectoderm of the limbs, in a subset of somite cells, and some mesoderm-derived tissues
<i>Fabp4</i> , fatty acid binding protein 4	Brown and white adipose tissue.
<i>Foxd1</i> (mouse)	Kidney development in metanephric mesenchyme in cells fated to become stromal cells of kidney, and multiple organs throughout body

Promoter (Species)	Site of Expression
<i>Foxp3</i> (mouse)	<i>Cd4+Cd25<high>Cd127<low></i> T cells from the lymph nodes, spleen and thymus; ovary
<i>Gad2</i> , glutamic acid decarboxylase 2	Gad2-positive neurons
<i>GFAP</i> , glial fibrillary acidic protein (human)	Central nervous system, including astrocytes, oligodendroglia, ependyma and some neurons; also periportal cells of the liver
<i>Gfap</i> (mouse)	Astrocytes in the brain and spinal cord, as well as postnatal and adult GFAP-expressing neural stem cells and their progeny in the brain; cartilage primordium at e15.5; thymus, myocardium, eye lens, peripheral nerves embedded in bladder and intestinal muscle of adults
<i>Gfap</i> (mouse)	Most astrocytes throughout the healthy brain and spinal cord and to essentially all astrocytes after Central Nervous System (CNS) injury; subpopulation of the adult stems in the subventricular zone
<i>Grik4</i> , glutamate receptor, ionotropic, kainate 4 (mouse)	At 14 days old in area CA3 of the hippocampus, and at 8 weeks of age, recombination is observed in nearly 100% of pyramidal cells in area CA3; other brain areas
<i>Hspa2</i> , heat shock protein 2 (mouse)	Leptotene/zygotene spermatocytes
<i>Ins2</i> , insulin 2 (rat)	Pancreatic beta cells, as well as the hypothalamus
<i>Itgax</i> , integrin alpha X (mouse)	CD8-, CD8+dendritic cells, tissue derived dendritic cells from lymph nodes, lung and epidermis and plasmacytoid dendritic cells
<i>Kap</i> (mouse)	Proximal tubule cells of the renal cortex in male mice; uterus and liver
<i>KRT14</i> , keratin 14 (human)	Skin, the oral ectoderm including the dental lamina at 11.75 d.p.c., and dental epithelium by 14.5 d.p.c.
<i>Lck</i> , lymphocyte protein tyrosine kinase (mouse)	Thymocytes
<i>Lck</i> (mouse)	Thymus
<i>Lepr</i> (mouse)	Hypothalamus (arcuate, dorsomedial, lateral, and ventromedial nuclei), limbic and cortical brain regions (basolateral amygdaloid nucleus, piriform cortex, and lateral entorhinal cortex), and retrosplenial cortex
<i>Lyve1</i> (mouse)	Lymphatic endothelium
<i>Lyz2</i> , Lysozyme 2 (mouse)	Myeloid cells, including monocytes, mature macrophages and granulocytes
MMTV	Mammary gland, salivary gland, seminal vesicle, skin, erythrocytes, B cells and T cells; lower in lung, kidney, liver and brain tissues
<i>Mnx1</i> , motor neuron and pancreas homeobox 1 (mouse)	Motor neurons
<i>Myf5</i> , myogenic factor 5	Skeletal muscle and the dermis, and in several ectopic locations
<i>Myh6</i> (mouse)	Cardiac tissue
<i>Nes</i> , nestin (rat)	Central and peripheral nervous system; a few isolated kidney and heart cells
<i>Neurog3</i> , neurogenin 3, (rat)	Islets of the adult pancreas, small intestine enteroendocrine cells, endocrine portions of the stomach, all pancreatic endocrine cells, and some non-endocrine intestinal cells
<i>Nkx2-1</i>	Cre recombinase activity is directed to brain interneuron progenitors, developing lung, thyroid, and pituitary by the <i>Nkx2.1</i> promoter/enhancer regions
<i>NPHS2</i> (human)	Podocytes during late capillary loop stage of glomerular development and podocytes of mature glomeruli
<i>Nr5a1</i> , Nuclear receptor subfamily 5 group A member 1 (mouse)	Ventromedial Hypothalamus, Cortex, Adrenal Gland, Pituitary Gland and Gonads
<i>Omp</i> , Olfactory Marker Protein (mouse)	Mature olfactory sensory neurons
<i>Pax3</i> , paired box gene 3	Dorsal neural tube and somites of E9 to 11.5 embryos and cardiac neural crest cells and colonic epithelia of E11.5 embryos
<i>Pf4</i> , platelet factor 4 (mouse)	Megakaryocytes

Promoter (Species)	Site of Expression
<i>Pomc1</i> (mouse)	POMC neurons in the arcuate nucleus of the hypothalamus and scattered in the dentate gyrus of the hippocampus
<i>Prdm1</i> (mouse)	Primordial germ cells
<i>Prm</i> (mouse)	Male germ line
<i>Pvalb</i> , parvalbumin	Neurons that express parvalbumin, such as interneurons in the brain and proprioceptive afferent sensory neurons in the dorsal root ganglia
<i>Scnn1a</i> (mouse)	Cortex, thalamus, midbrain, and cerebellum
<i>Shh</i> , sonic hedgehog	Endogenous Shh expression patterns
<i>Sim1</i> , single-minded homolog 1 (Drosophila)(mouse)	Paraventricular hypothalamus and other parts of the brain
<i>Slc6a3</i> , solute carrier family 6 (neurotransmitter transporter, dopamine), member 3	Dopaminergic cell groups (substantia nigra (SN) and ventral tegmental area (VTA), as well as in the retrorubral field)
<i>Slc17a6</i> (mouse)	Excitatory glutamatergic neuron cell bodies
<i>Sst</i> , somatostatin	Somatostatin positive neurons (including dendritic inhibitory interneurons such as Martinotti cells and Oriens-Lacunosum-Moleculare cells)
<i>Stra8</i> (mouse)	Postnatal, premeiotic, male germ cells
<i>Syn1</i> (rat)	Neuronal cells, including brain, spinal cord and DRGs, as early as E12.5, as well as in neurons in adult
<i>Tagln</i> , transgelin (mouse)	Smooth muscle
<i>Tagln</i> (mouse)	Adult smooth muscle cells (such as arteries, veins, and visceral organs) and cardiac myocytes
<i>Tek</i> (mouse)	Endothelial cells during embryogenesis and adulthood
<i>Thy1</i> (mouse)	Neurons of the cortex and hippocampus
<i>Twist2</i> , twist basic helix-loop-helix transcription factor 2	Mesoderm as early as embryonic day 9.5, in mesodermal tissues such as branchial arches and somites, and in condensed mesenchyme-derived chondrocytes and osteoblasts
<i>Vav1</i> (mouse)	Variegated germline (testis and ovaries), and heart and gut
<i>Vill</i> , villin 1 (mouse)	Villi and crypts of the small and large intestine
<i>Vip</i> , vasoactive intestinal polypeptide	Some GABAergic interneurons
<i>Wnt1</i> , wingless-related MMTV integration site 1 (mouse)	Embryonic neural tube, midbrain, dorsal and ventral midlines of the midbrain and caudal diencephalon, the mid-hindbrain junction and dorsal spinal cord
<i>Wnt1</i> (mouse)	Developing neural crest and midbrain
<i>Krt17</i> , keratin 17 (mouse)	Endogenous keratin 17 expression patterns
<i>Osr2</i> , odd-skipped related 2 (Drosophila), mouse, laboratory	Developing palate and urogenital tract
<i>Trp63</i> , transformation related protein 63 (mouse)	Endogenous <i>Trp63</i> expression patterns
<i>Prrx1</i> , paired related homeobox 1 (rat)	Early limb bud mesenchyme and in a subset of craniofacial mesenchyme, along with limited female germline expression
<i>Tbx22</i> , T-box transcription factor 22 (mouse)	Endogenous <i>Tbx22</i> expression patterns
<i>Tgfb3</i> , transforming growth factor, beta 3 (mouse)	Heart, pharyngeal arches, otic vesicle, mid brain, limb buds, midline palatal epithelium, and whisker follicles during embryo and fetus development
<i>Wnt1</i> , wingless-related MMTV integration site 1 (mouse)	Embryonic neural tube, midbrain, caudal diencephalon, the mid-hindbrain junction, dorsal spinal cord, and neural crest cells
<i>ACTB</i> , actin, beta (chicken)	Most tissue types
<i>Col2a1</i> , collagen, type II, alpha 1 (mouse)	Cells of chondrogenic lineage (cartilage) during embryogenesis and postnatally.
<i>Dlx5</i> , distal-less homeobox 5	Cortex

Promoter (Species)	Site of Expression
<i>KRT14</i> , keratin 14 (human)	Keratinocytes
<i>Lgr5</i> , leucine rich repeat containing G protein coupled receptor 5	Crypt base columnar cells in small intestine (stem cells of the small intestine) and colon
<i>Myh6</i> , myosin, heavy polypeptide 6, (mouse)	Developing and adult heart
<i>Plp1</i> , proteolipid protein (myelin) 1 (mouse)	Oligodendrocytes and Schwann cells
<i>UBC</i> , ubiquitin C (human)	All tissue types
<i>Wfs1</i> , Wolfram syndrome 1 homolog (human)	Cortex, hippocampus, striatum, thalamus and cerebellum
<i>Gt(ROSA)26Sor</i> (mouse)	Most tissue types preimplantation onward, including cells of developing germline
Chicken beta-actin promoter and an hCMV immediate early enhancer	Ubiquitous

III. Methods of Assessing CRISPR/Cas Activity In Vivo

[00180] Various methods are provided for assessing CRISPR/Cas delivery to and for assessing CRISPR/Cas activity in tissues and organs of a live animal. Such methods make use of non-human animals comprising a Cas expression cassette as described elsewhere herein.

A. Methods of Testing Ability of CRISPR/Cas to Modify a Target Genomic Locus In Vivo or Ex Vivo

[00181] Various methods are provided for assessing the ability of a CRISPR/Cas nickase or nuclease to modify a target genomic locus *in vivo* using the non-human animals comprising a Cas expression cassette described herein. Such methods can comprise: (a) introducing into the non-human animal a guide RNA designed to target a guide RNA target sequence at the target genomic locus; and (b) assessing the modification of the target locus. Modification of a target genomic locus will be induced when the guide RNA forms a complex with the Cas protein and directs the Cas protein to the target genomic locus, and the Cas/guide RNA complex cleaves the guide RNA target sequence, triggering repair by the cell (e.g., via non-homologous end joining (NHEJ) if no donor sequence is present).

[00182] Optionally, two or more guide RNAs can be introduced, each designed to target a different guide RNA target sequence within the target genomic locus. For example, two guide RNAs can be designed to excise a genomic sequence between the two guide RNA target sequences. Modification of a target genomic locus will be induced when the first guide RNA forms a complex with the Cas protein and directs the Cas protein to the target genomic locus, the

second guide RNA forms a complex with the Cas protein and directs the Cas protein to the target genomic locus, the first Cas/guide RNA complex cleaves the first guide RNA target sequence, and the second Cas/guide RNA complex cleaves the second guide RNA target sequence, resulting in excision of the intervening sequence. Alternatively, two or more guide RNAs can be introduced, each designed to target to a different guide RNA target sequence at a different target genomic locus (i.e., multiplexing).

[00183] Optionally, an exogenous donor nucleic acid capable of recombining with and modifying the target genomic locus is also introduced into the non-human animal. Optionally, the Cas protein can be tethered to the exogenous donor nucleic acid as described elsewhere herein. Optionally, two or more exogenous donor nucleic acids are introduced, each capable of recombining with and modifying a different target genomic locus. Modification of the target genomic locus will be induced, for example, when the guide RNA forms a complex with the Cas protein and directs the Cas protein to the target genomic locus, the Cas/guide RNA complex cleaves the guide RNA target sequence, and the target genomic locus recombines with the exogenous donor nucleic acid to modify the target genomic locus. The exogenous donor nucleic acid can recombine with the target genomic locus, for example, via homology-directed repair (HDR) or via NHEJ-mediated insertion. Any type of exogenous donor nucleic acid can be used, examples of which are provided elsewhere herein.

[00184] Likewise, the various methods provided above for assessing CRISPR/Cas activity *in vivo* can also be used to assess CRISPR/Cas activity *ex vivo* using cells comprising a Cas expression cassette as described elsewhere herein.

[00185] Guide RNAs and optionally exogenous donor nucleic acids can be introduced into the cell or non-human animal via any delivery method (e.g., AAV, LNP, or HDD) and any route of administration as disclosed elsewhere herein. In particular methods, the guide RNA (or guide RNAs) is delivered via AAV-mediated delivery. For example, AAV8 can be used if the liver is being targeted.

[00186] Methods for assessing modification of the target genomic locus are provided elsewhere herein and are well known. Assessment of modification of the target genomic locus can be in any cell type, any tissue type, or any organ type as disclosed elsewhere herein. In some methods, modification of the target genomic locus is assessed in liver cells (e.g., assessing serum levels of a secreted protein expressed by the target genomic locus in liver cells). For example,

the target genomic locus comprises a target gene, and assessment can comprise measuring expression of the target gene or activity of a protein encoded by the target gene. Alternatively or additionally, assessment can comprise sequencing the target genomic locus in one or more cells isolated from the non-human animal. Assessment can comprise isolating a target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the target organ or tissue. Assessment can also comprise assessing modification of the target genomic locus in two or more different cell types within the target organ or tissue. Similarly, assessment can comprise isolating a non-target organ or tissue (e.g., two or more non-target organs or tissues) from the non-human animal and assessing modification of the target genomic locus in the non-target organ or tissue.

(1) Exogenous Donor Nucleic Acids

[00187] The methods and compositions disclosed herein utilize exogenous donor nucleic acids to modify a target genomic locus following cleavage of the target genomic locus with a Cas protein. In such methods, the Cas protein cleaves the target genomic locus to create a single-strand break (nick) or double-strand break, and the exogenous donor nucleic acid recombines the target genomic locus via non-homologous end joining (NHEJ)-mediated ligation or through a homology-directed repair event. Optionally, repair with the exogenous donor nucleic acid removes or disrupts the guide RNA target sequence or the Cas cleavage site so that alleles that have been targeted cannot be re-targeted by the Cas protein.

[00188] Exogenous donor nucleic acids can comprise deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), they can be single-stranded or double-stranded, and they can be in linear or circular form. For example, an exogenous donor nucleic acid can be a single-stranded oligodeoxynucleotide (ssODN). *See, e.g., Yoshimi et al. (2016) Nat. Commun. 7:10431*, herein incorporated by reference in its entirety for all purposes. An exemplary exogenous donor nucleic acid is between about 50 nucleotides to about 5 kb in length, is between about 50 nucleotides to about 3 kb in length, or is between about 50 to about 1,000 nucleotides in length. Other exemplary exogenous donor nucleic acids are between about 40 to about 200 nucleotides in length. For example, an exogenous donor nucleic acid can be between about 50-60, 60-70, 70-80, 80-90, 90-100, 100-110, 110-120, 120-130, 130-140, 140-150, 150-160, 160-170, 170-180, 180-190, or 190-200 nucleotides in length. Alternatively, an exogenous donor nucleic acid can

be between about 50-100, 100-200, 200-300, 300-400, 400-500, 500-600, 600-700, 700-800, 800-900, or 900-1000 nucleotides in length. Alternatively, an exogenous donor nucleic acid can be between about 1-1.5, 1.5-2, 2-2.5, 2.5-3, 3-3.5, 3.5-4, 4-4.5, or 4.5-5 kb in length.

Alternatively, an exogenous donor nucleic acid can be, for example, no more than 5 kb, 4.5 kb, 4 kb, 3.5 kb, 3 kb, 2.5 kb, 2 kb, 1.5 kb, 1 kb, 900 nucleotides, 800 nucleotides, 700 nucleotides, 600 nucleotides, 500 nucleotides, 400 nucleotides, 300 nucleotides, 200 nucleotides, 100 nucleotides, or 50 nucleotides in length. Exogenous donor nucleic acids (e.g., targeting vectors) can also be longer.

[00189] In one example, an exogenous donor nucleic acid is an ssODN that is between about 80 nucleotides and about 200 nucleotides in length. In another example, an exogenous donor nucleic acid is an ssODN that is between about 80 nucleotides and about 3 kb in length. Such an ssODN can have homology arms, for example, that are each between about 40 nucleotides and about 60 nucleotides in length. Such an ssODN can also have homology arms, for example, that are each between about 30 nucleotides and 100 nucleotides in length. The homology arms can be symmetrical (e.g., each 40 nucleotides or each 60 nucleotides in length), or they can be asymmetrical (e.g., one homology arm that is 36 nucleotides in length, and one homology arm that is 91 nucleotides in length).

[00190] Exogenous donor nucleic acids can include modifications or sequences that provide for additional desirable features (e.g., modified or regulated stability; tracking or detecting with a fluorescent label; a binding site for a protein or protein complex; and so forth). Exogenous donor nucleic acids can comprise one or more fluorescent labels, purification tags, epitope tags, or a combination thereof. For example, an exogenous donor nucleic acid can comprise one or more fluorescent labels (e.g., fluorescent proteins or other fluorophores or dyes), such as at least 1, at least 2, at least 3, at least 4, or at least 5 fluorescent labels. Exemplary fluorescent labels include fluorophores such as fluorescein (e.g., 6-carboxyfluorescein (6-FAM)), Texas Red, HEX, Cy3, Cy5, Cy5.5, Pacific Blue, 5-(and-6)-carboxytetramethylrhodamine (TAMRA), and Cy7. A wide range of fluorescent dyes are available commercially for labeling oligonucleotides (e.g., from Integrated DNA Technologies). Such fluorescent labels (e.g., internal fluorescent labels) can be used, for example, to detect an exogenous donor nucleic acid that has been directly integrated into a cleaved target nucleic acid having protruding ends compatible with the ends of the exogenous donor nucleic acid. The label or tag can be at the 5' end, the 3' end, or internally

within the exogenous donor nucleic acid. For example, an exogenous donor nucleic acid can be conjugated at 5' end with the IR700 fluorophore from Integrated DNA Technologies (5'IRDYE®700).

[00191] Exogenous donor nucleic acids can also comprise nucleic acid inserts including segments of DNA to be integrated at target genomic loci. Integration of a nucleic acid insert at a target genomic locus can result in addition of a nucleic acid sequence of interest to the target genomic locus, deletion of a nucleic acid sequence of interest at the target genomic locus, or replacement of a nucleic acid sequence of interest at the target genomic locus (i.e., deletion and insertion). Some exogenous donor nucleic acids are designed for insertion of a nucleic acid insert at a target genomic locus without any corresponding deletion at the target genomic locus. Other exogenous donor nucleic acids are designed to delete a nucleic acid sequence of interest at a target genomic locus without any corresponding insertion of a nucleic acid insert. Yet other exogenous donor nucleic acids are designed to delete a nucleic acid sequence of interest at a target genomic locus and replace it with a nucleic acid insert.

[00192] The nucleic acid insert or the corresponding nucleic acid at the target genomic locus being deleted and/or replaced can be various lengths. An exemplary nucleic acid insert or corresponding nucleic acid at the target genomic locus being deleted and/or replaced is between about 1 nucleotide to about 5 kb in length or is between about 1 nucleotide to about 1,000 nucleotides in length. For example, a nucleic acid insert or a corresponding nucleic acid at the target genomic locus being deleted and/or replaced can be between about 1-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, 90-100, 100-110, 110-120, 120-130, 130-140, 140-150, 150-160, 160-170, 170-180, 180-190, or 190-200 nucleotides in length. Likewise, a nucleic acid insert or a corresponding nucleic acid at the target genomic locus being deleted and/or replaced can be between 1-100, 100-200, 200-300, 300-400, 400-500, 500-600, 600-700, 700-800, 800-900, or 900-1000 nucleotides in length. Likewise, a nucleic acid insert or a corresponding nucleic acid at the target genomic locus being deleted and/or replaced can be between about 1-1.5, 1.5-2, 2-2.5, 2.5-3, 3-3.5, 3.5-4, 4-4.5, or 4.5-5 kb in length or longer.

[00193] The nucleic acid insert can comprise a sequence that is homologous or orthologous to all or part of sequence targeted for replacement. For example, the nucleic acid insert can comprise a sequence that comprises one or more point mutations (e.g., 1, 2, 3, 4, 5, or more) compared with a sequence targeted for replacement at the target genomic locus. Optionally, such

point mutations can result in a conservative amino acid substitution (e.g., substitution of aspartic acid [Asp, D] with glutamic acid [Glu, E]) in the encoded polypeptide.

(2) Donor Nucleic Acids for Non-Homologous-End-Joining-Mediated Insertion

[00194] Some exogenous donor nucleic acids have short single-stranded regions at the 5' end and/or the 3' end that are complementary to one or more overhangs created by Cas-protein-mediated cleavage at the target genomic locus. These overhangs can also be referred to as 5' and 3' homology arms. For example, some exogenous donor nucleic acids have short single-stranded regions at the 5' end and/or the 3' end that are complementary to one or more overhangs created by Cas-protein-mediated cleavage at 5' and/or 3' target sequences at the target genomic locus. Some such exogenous donor nucleic acids have a complementary region only at the 5' end or only at the 3' end. For example, some such exogenous donor nucleic acids have a complementary region only at the 5' end complementary to an overhang created at a 5' target sequence at the target genomic locus or only at the 3' end complementary to an overhang created at a 3' target sequence at the target genomic locus. Other such exogenous donor nucleic acids have complementary regions at both the 5' and 3' ends. For example, other such exogenous donor nucleic acids have complementary regions at both the 5' and 3' ends e.g., complementary to first and second overhangs, respectively, generated by Cas-mediated cleavage at the target genomic locus. For example, if the exogenous donor nucleic acid is double-stranded, the single-stranded complementary regions can extend from the 5' end of the top strand of the donor nucleic acid and the 5' end of the bottom strand of the donor nucleic acid, creating 5' overhangs on each end. Alternatively, the single-stranded complementary region can extend from the 3' end of the top strand of the donor nucleic acid and from the 3' end of the bottom strand of the template, creating 3' overhangs.

[00195] The complementary regions can be of any length sufficient to promote ligation between the exogenous donor nucleic acid and the target nucleic acid. Exemplary complementary regions are between about 1 to about 5 nucleotides in length, between about 1 to about 25 nucleotides in length, or between about 5 to about 150 nucleotides in length. For example, a complementary region can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides in length. Alternatively, the

complementary region can be about 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, 90-100, 100-110, 110-120, 120-130, 130-140, or 140-150 nucleotides in length, or longer.

[00196] Such complementary regions can be complementary to overhangs created by two pairs of nickases. Two double-strand breaks with staggered ends can be created by using first and second nickases that cleave opposite strands of DNA to create a first double-strand break, and third and fourth nickases that cleave opposite strands of DNA to create a second double-strand break. For example, a Cas protein can be used to nick first, second, third, and fourth guide RNA target sequences corresponding with first, second, third, and fourth guide RNAs. The first and second guide RNA target sequences can be positioned to create a first cleavage site such that the nicks created by the first and second nickases on the first and second strands of DNA create a double-strand break (i.e., the first cleavage site comprises the nicks within the first and second guide RNA target sequences). Likewise, the third and fourth guide RNA target sequences can be positioned to create a second cleavage site such that the nicks created by the third and fourth nickases on the first and second strands of DNA create a double-strand break (i.e., the second cleavage site comprises the nicks within the third and fourth guide RNA target sequences). Optionally, the nicks within the first and second guide RNA target sequences and/or the third and fourth guide RNA target sequences can be off-set nicks that create overhangs. The offset window can be, for example, at least about 5 bp, 10 bp, 20 bp, 30 bp, 40 bp, 50 bp, 60 bp, 70 bp, 80 bp, 90 bp, 100 bp or more. *See Ran et al. (2013) Cell 154:1380-1389; Mali et al. (2013) Nat. Biotech. 31:833-838; and Shen et al. (2014) Nat. Methods 11:399-404, each of which is herein incorporated by reference in its entirety for all purposes.* In such cases, a double-stranded exogenous donor nucleic acid can be designed with single-stranded complementary regions that are complementary to the overhangs created by the nicks within the first and second guide RNA target sequences and by the nicks within the third and fourth guide RNA target sequences. Such an exogenous donor nucleic acid can then be inserted by non-homologous-end-joining-mediated ligation.

(3) Donor Nucleic Acids for Insertion by Homology-Directed Repair

[00197] Some exogenous donor nucleic acids comprise homology arms. If the exogenous donor nucleic acid also comprises a nucleic acid insert, the homology arms can flank the nucleic acid insert. For ease of reference, the homology arms are referred to herein as 5' and 3' (i.e.,

upstream and downstream) homology arms. This terminology relates to the relative position of the homology arms to the nucleic acid insert within the exogenous donor nucleic acid. The 5' and 3' homology arms correspond to regions within the target genomic locus, which are referred to herein as "5' target sequence" and "3' target sequence," respectively.

[00198] A homology arm and a target sequence "correspond" or are "corresponding" to one another when the two regions share a sufficient level of sequence identity to one another to act as substrates for a homologous recombination reaction. The term "homology" includes DNA sequences that are either identical or share sequence identity to a corresponding sequence. The sequence identity between a given target sequence and the corresponding homology arm found in the exogenous donor nucleic acid can be any degree of sequence identity that allows for homologous recombination to occur. For example, the amount of sequence identity shared by the homology arm of the exogenous donor nucleic acid (or a fragment thereof) and the target sequence (or a fragment thereof) can be at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity, such that the sequences undergo homologous recombination. Moreover, a corresponding region of homology between the homology arm and the corresponding target sequence can be of any length that is sufficient to promote homologous recombination. Exemplary homology arms are between about 25 nucleotides to about 2.5 kb in length, are between about 25 nucleotides to about 1.5 kb in length, or are between about 25 to about 500 nucleotides in length. For example, a given homology arm (or each of the homology arms) and/or corresponding target sequence can comprise corresponding regions of homology that are between about 25-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, 90-100, 100-150, 150-200, 200-250, 250-300, 300-350, 350-400, 400-450, or 450-500 nucleotides in length, such that the homology arms have sufficient homology to undergo homologous recombination with the corresponding target sequences within the target nucleic acid. Alternatively, a given homology arm (or each homology arm) and/or corresponding target sequence can comprise corresponding regions of homology that are between about 0.5 kb to about 1 kb, about 1 kb to about 1.5 kb, about 1.5 kb to about 2 kb, or about 2 kb to about 2.5 kb in length. For example, the homology arms can each be about 750 nucleotides in length. The homology arms can be symmetrical (each about the same size in length), or they can be asymmetrical (one longer than the other).

[00199] When a CRISPR/Cas system is used in combination with an exogenous donor nucleic acid, the 5' and 3' target sequences are optionally located in sufficient proximity to the Cas cleavage site (e.g., within sufficient proximity to a the guide RNA target sequence) so as to promote the occurrence of a homologous recombination event between the target sequences and the homology arms upon a single-strand break (nick) or double-strand break at the Cas cleavage site. The term "Cas cleavage site" includes a DNA sequence at which a nick or double-strand break is created by a Cas enzyme (e.g., a Cas9 protein complexed with a guide RNA). The target sequences within the targeted locus that correspond to the 5' and 3' homology arms of the exogenous donor nucleic acid are "located in sufficient proximity" to a Cas cleavage site if the distance is such as to promote the occurrence of a homologous recombination event between the 5' and 3' target sequences and the homology arms upon a single-strand break or double-strand break at the Cas cleavage site. Thus, the target sequences corresponding to the 5' and/or 3' homology arms of the exogenous donor nucleic acid can be, for example, within at least 1 nucleotide of a given Cas cleavage site or within at least 10 nucleotides to about 1,000 nucleotides of a given Cas cleavage site. As an example, the Cas cleavage site can be immediately adjacent to at least one or both of the target sequences.

[00200] The spatial relationship of the target sequences that correspond to the homology arms of the exogenous donor nucleic acid and the Cas cleavage site can vary. For example, target sequences can be located 5' to the Cas cleavage site, target sequences can be located 3' to the Cas cleavage site, or the target sequences can flank the Cas cleavage site.

B. Methods of Optimizing Ability of CRISPR/Cas to Excise a Target Genomic Nucleic Acid *In Vivo* or *Ex Vivo*

[00201] Various methods are provided for optimizing delivery of CRISPR/Cas to a cell or non-human animal or optimizing CRISPR/Cas activity *in vivo*. Such methods can comprise, for example: (a) performing the method of testing the ability of CRISPR/Cas to modify a target genomic locus as described above a first time in a first non-human animal or first cell; (b) changing a variable and performing the method a second time in a second non-human animal (i.e., of the same species) or a second cell with the changed variable; and (c) comparing modification of the target genomic locus in step (a) with the modification of the target genomic locus in step (b), and selecting the method resulting in the more effective modification of the

target genomic locus.

[00202] More effective modification of the target genomic locus can mean different things depending on the desired effect within the non-human animal or cell. For example, more effective modification of the target genomic locus can mean one or more or all of higher efficacy, higher precision, higher consistency, or higher specificity. Higher efficacy refers to higher levels of modification of the target genomic locus (e.g., a higher percentage of cells is targeted within a particular target cell type, within a particular target tissue, or within a particular target organ). Higher precision refers to more precise modification of the target genomic locus (e.g., a higher percentage of targeted cells having the same modification or having the desired modification without extra unintended insertions and deletions (e.g., NHEJ indels)). Higher consistency refers to more consistent modification of the target genomic locus among different types of targeted cells, tissues, or organs if more than one type of cell, tissue, or organ is being targeted (e.g., modification of a greater number of cell types within a target organ). If a particular organ is being targeted, higher consistency can also refer to more consistent modification throughout all locations within the organ. Higher specificity can refer to higher specificity with respect to the genomic locus or loci targeted, higher specificity with respect to the cell type targeted, higher specificity with respect to the tissue type targeted, or higher specificity with respect to the organ targeted. For example, increased genomic locus specificity refers to less modification of off-target genomic loci (e.g., a lower percentage of targeted cells having modifications at unintended, off-target genomic loci instead of or in addition to modification of the target genomic locus). Likewise, increased cell type, tissue, or organ type specificity refers to less modification of off-target cell types, tissue types, or organ types if a particular cell type, tissue type, or organ type is being targeted (e.g., when a particular organ is targeted (e.g., the liver), there is less modification of cells in organs or tissues that are not intended targets).

[00203] The variable that is changed can be any parameter. As one example, the changed variable can be the packaging or the delivery method by which one or more or all of the guide RNA (or guide RNA packaged in AAV) and the exogenous donor nucleic acid are introduced into the cell or non-human animal. Examples of delivery methods, such as LNP, HDD, and AAV, are disclosed elsewhere herein. For example, the changed variable can be the AAV serotype. As another example, the changed variable can be the route of administration for

introduction of one or more or all of the guide RNA (e.g., packaged in AAV) and the exogenous donor nucleic acid into the cell or non-human animal. Examples of routes of administration, such as intravenous, intravitreal, intraparenchymal, and nasal instillation, are disclosed elsewhere herein.

[00204] As another example, the changed variable can be the concentration or amount of one or more or all of the guide RNA (e.g., packaged in AAV) introduced and the exogenous donor nucleic acid introduced. As another example, the changed variable can be the concentration or the amount of guide RNA (e.g., packaged in AAV) introduced relative to the concentration or the amount of exogenous donor nucleic acid introduced.

[00205] As another example, the changed variable can be the timing of introducing one or more or all of the guide RNA (e.g., packaged in AAV) and the exogenous donor nucleic acid relative to the timing of measuring expression or activity of the one or more reporter proteins. As another example, the changed variable can be the number of times or frequency with which one or more or all of the guide RNA (e.g., packaged in AAV) and the exogenous donor nucleic acid are introduced. As another example, the changed variable can be the timing of introduction of guide RNA relative to the timing of introduction of exogenous donor nucleic acid.

[00206] As another example, the changed variable can be the form in which one or more or all of the guide RNA and the exogenous donor nucleic acid are introduced. For example, the guide RNA can be introduced in the form of DNA or in the form of RNA. The exogenous donor nucleic acid can be DNA, RNA, single-stranded, double-stranded, linear, circular, and so forth. Similarly, each of the components can comprise various combinations of modifications for stability, to reduce off-target effects, to facilitate delivery, and so forth. As another example, the changed variable can be one or more or all of the guide RNA that is introduced (e.g., introducing a different guide RNA with a different sequence) and the exogenous donor nucleic acid that is introduced (e.g., introducing a different exogenous donor nucleic acid with a different sequence).

C. Introducing Guide RNAs and Other Components into Cells and Non-Human Animals

[00207] The methods disclosed herein comprise introducing into a cell or non-human animal one or more or all of guide RNAs and exogenous donor nucleic acids. “Introducing” includes presenting to the cell or non-human animal the nucleic acid or protein in such a manner that the

nucleic acid or protein gains access to the interior of the cell or to the interior of cells within the non-human animal. The introducing can be accomplished by any means, and two or more of the components (e.g., two of the components, or all of the components) can be introduced into the cell or non-human animal simultaneously or sequentially in any combination. For example, an exogenous donor nucleic acid can be introduced into a cell or non-human animal before introduction of a guide RNA, or it can be introduced following introduction of the guide RNA (e.g., the exogenous donor nucleic acid can be administered about 1, 2, 3, 4, 8, 12, 24, 36, 48, or 72 hours before or after introduction of the guide RNA). *See, e.g.*, US 2015/0240263 and US 2015/0110762, each of which is herein incorporated by reference in its entirety for all purposes. In addition, two or more of the components can be introduced into the cell or non-human animal by the same delivery method or different delivery methods. Similarly, two or more of the components can be introduced into a non-human animal by the same route of administration or different routes of administration.

[00208] A guide RNA can be introduced into the cell in the form of an RNA (e.g., *in vitro* transcribed RNA) or in the form of a DNA encoding the guide RNA. When introduced in the form of a DNA, the DNA encoding a guide RNA can be operably linked to a promoter active in the cell. For example, a guide RNA may be delivered via AAV and expressed *in vivo* under a U6 promoter. Such DNAs can be in one or more expression constructs. For example, such expression constructs can be components of a single nucleic acid molecule. Alternatively, they can be separated in any combination among two or more nucleic acid molecules (i.e., DNAs encoding one or more CRISPR RNAs and DNAs encoding one or more tracrRNAs can be components of a separate nucleic acid molecules).

[00209] Nucleic acids encoding guide RNAs can be operably linked to a promoter in an expression construct. Expression constructs include any nucleic acid constructs capable of directing expression of a gene or other nucleic acid sequence of interest and which can transfer such a nucleic acid sequence of interest to a target cell. Suitable promoters that can be used in an expression construct include promoters active, for example, in one or more of a eukaryotic cell, a human cell, a non-human cell, a mammalian cell, a non-human mammalian cell, a rodent cell, a mouse cell, a rat cell, a hamster cell, a rabbit cell, a pluripotent cell, an embryonic stem (ES) cell, an adult stem cell, a developmentally restricted progenitor cell, an induced pluripotent stem (iPS) cell, or a one-cell stage embryo. Such promoters can be, for example, conditional promoters,

inducible promoters, constitutive promoters, or tissue-specific promoters. Optionally, the promoter can be a bidirectional promoter driving expression of both a guide RNA in one direction and another component in the other direction. Such bidirectional promoters can consist of (1) a complete, conventional, unidirectional Pol III promoter that contains 3 external control elements: a distal sequence element (DSE), a proximal sequence element (PSE), and a TATA box; and (2) a second basic Pol III promoter that includes a PSE and a TATA box fused to the 5' terminus of the DSE in reverse orientation. For example, in the H1 promoter, the DSE is adjacent to the PSE and the TATA box, and the promoter can be rendered bidirectional by creating a hybrid promoter in which transcription in the reverse direction is controlled by appending a PSE and TATA box derived from the U6 promoter. *See, e.g.*, US 2016/0074535, herein incorporated by references in its entirety for all purposes. Use of a bidirectional promoter to express genes encoding a guide RNA and another component simultaneously allows for the generation of compact expression cassettes to facilitate delivery.

[00210] Exogenous donor nucleic acids and guide RNAs (or nucleic acids encoding guide RNAs) can be provided in compositions comprising a carrier increasing the stability of the exogenous donor nucleic acid or guide RNA (e.g., prolonging the period under given conditions of storage (e.g., -20°C, 4°C, or ambient temperature) for which degradation products remain below a threshold, such below 0.5% by weight of the starting nucleic acid or protein; or increasing the stability *in vivo*). Non-limiting examples of such carriers include poly(lactic acid) (PLA) microspheres, poly(D,L-lactic-coglycolic-acid) (PLGA) microspheres, liposomes, micelles, inverse micelles, lipid cochleates, and lipid microtubules.

[00211] Various methods and compositions are provided herein to allow for introduction of a nucleic acid or protein into a cell or non-human animal. Methods for introducing nucleic acids into various cell types are known in the art and include, for example, stable transfection methods, transient transfection methods, and virus-mediated methods.

[00212] Transfection protocols as well as protocols for introducing nucleic acid sequences into cells may vary. Non-limiting transfection methods include chemical-based transfection methods using liposomes; nanoparticles; calcium phosphate (Graham *et al.* (1973) *Virology* 52 (2): 456–67, Bacchetti *et al.* (1977) *Proc. Natl. Acad. Sci. USA* 74 (4): 1590–4, and Kriegler, M (1991). *Transfer and Expression: A Laboratory Manual*. New York: W. H. Freeman and Company. pp. 96–97); dendrimers; or cationic polymers such as DEAE-dextran or

polyethylenimine. Non-chemical methods include electroporation, Sono-poration, and optical transfection. Particle-based transfection includes the use of a gene gun, or magnet-assisted transfection (Bertram (2006) *Current Pharmaceutical Biotechnology* 7, 277–28). Viral methods can also be used for transfection.

[00213] Introduction of nucleic acids or proteins into a cell can also be mediated by electroporation, by intracytoplasmic injection, by viral infection, by adenovirus, by adeno-associated virus, by lentivirus, by retrovirus, by transfection, by lipid-mediated transfection, or by nucleofection. Nucleofection is an improved electroporation technology that enables nucleic acid substrates to be delivered not only to the cytoplasm but also through the nuclear membrane and into the nucleus. In addition, use of nucleofection in the methods disclosed herein typically requires much fewer cells than regular electroporation (e.g., only about 2 million compared with 7 million by regular electroporation). In one example, nucleofection is performed using the LONZA[®] NUCLEOFECTOR[™] system.

[00214] Introduction of nucleic acids or proteins into a cell (e.g., a zygote) can also be accomplished by microinjection. In zygotes (i.e., one-cell stage embryos), microinjection can be into the maternal and/or paternal pronucleus or into the cytoplasm. If the microinjection is into only one pronucleus, the paternal pronucleus is preferable due to its larger size. Alternatively, microinjection can be carried out by injection into both the nucleus/pronucleus and the cytoplasm: a needle can first be introduced into the nucleus/pronucleus and a first amount can be injected, and while removing the needle from the one-cell stage embryo a second amount can be injected into the cytoplasm. Methods for carrying out microinjection are well known. *See, e.g.,* Nagy *et al.* (Nagy A, Gertsenstein M, Vintersten K, Behringer R., 2003, *Manipulating the Mouse Embryo*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press); *see also* Meyer *et al.* (2010) *Proc. Natl. Acad. Sci. USA* 107:15022-15026 and Meyer *et al.* (2012) *Proc. Natl. Acad. Sci. USA* 109:9354-9359.

[00215] Other methods for introducing nucleic acid or proteins into a cell or non-human animal can include, for example, vector delivery, particle-mediated delivery, exosome-mediated delivery, lipid-nanoparticle-mediated delivery, cell-penetrating-peptide-mediated delivery, or implantable-device-mediated delivery. As specific examples, a nucleic acid or protein can be introduced into a cell or non-human animal in a carrier such as a poly(lactic acid) (PLA) microsphere, a poly(D,L-lactic-coglycolic-acid) (PLGA) microsphere, a liposome, a micelle, an

inverse micelle, a lipid cochleate, or a lipid microtubule. Some specific examples of delivery to a non-human animal include hydrodynamic delivery, virus-mediated delivery (e.g., adeno-associated virus (AAV)-mediated delivery), and lipid-nanoparticle-mediated delivery.

[00216] Introduction of nucleic acids and proteins into cells or non-human animals can be accomplished by hydrodynamic delivery (HDD). Hydrodynamic delivery has emerged as a method for intracellular DNA delivery *in vivo*. For gene delivery to parenchymal cells, only essential DNA sequences need to be injected via a selected blood vessel, eliminating safety concerns associated with current viral and synthetic vectors. When injected into the bloodstream, DNA is capable of reaching cells in the different tissues accessible to the blood. Hydrodynamic delivery employs the force generated by the rapid injection of a large volume of solution into the incompressible blood in the circulation to overcome the physical barriers of endothelium and cell membranes that prevent large and membrane-impermeable compounds from entering parenchymal cells. In addition to the delivery of DNA, this method is useful for the efficient intracellular delivery of RNA, proteins, and other small compounds *in vivo*. See, e.g., Bonamassa *et al.* (2011) *Pharm. Res.* 28(4):694-701, herein incorporated by reference in its entirety for all purposes.

[00217] Introduction of nucleic acids can also be accomplished by virus-mediated delivery, such as AAV-mediated delivery or lentivirus-mediated delivery. Other exemplary viruses/viral vectors include retroviruses, adenoviruses, vaccinia viruses, poxviruses, and herpes simplex viruses. The viruses can infect dividing cells, non-dividing cells, or both dividing and non-dividing cells. The viruses can integrate into the host genome or alternatively do not integrate into the host genome. Such viruses can also be engineered to have reduced immunity. The viruses can be replication-competent or can be replication-defective (e.g., defective in one or more genes necessary for additional rounds of virion replication and/or packaging). Viruses can cause transient expression, long-lasting expression (e.g., at least 1 week, 2 weeks, 1 month, 2 months, or 3 months), or permanent expression (e.g., of Cas9 and/or gRNA). Exemplary viral titers (e.g., AAV titers) include 10^{12} , 10^{13} , 10^{14} , 10^{15} , and 10^{16} vector genomes/mL.

[00218] The ssDNA AAV genome consists of two open reading frames, Rep and Cap, flanked by two inverted terminal repeats that allow for synthesis of the complementary DNA strand. When constructing an AAV transfer plasmid, the transgene is placed between the two ITRs, and Rep and Cap can be supplied *in trans*. In addition to Rep and Cap, AAV can require a helper

plasmid containing genes from adenovirus. These genes (E4, E2a, and VA) mediated AAV replication. For example, the transfer plasmid, Rep/Cap, and the helper plasmid can be transfected into HEK293 cells containing the adenovirus gene E1+ to produce infectious AAV particles. Alternatively, the Rep, Cap, and adenovirus helper genes may be combined into a single plasmid. Similar packaging cells and methods can be used for other viruses, such as retroviruses.

[00219] Multiple serotypes of AAV have been identified. These serotypes differ in the types of cells they infect (i.e., their tropism), allowing preferential transduction of specific cell types. Serotypes for CNS tissue include AAV1, AAV2, AAV4, AAV5, AAV8, and AAV9. Serotypes for heart tissue include AAV1, AAV8, and AAV9. Serotypes for kidney tissue include AAV2. Serotypes for lung tissue include AAV4, AAV5, AAV6, and AAV9. Serotypes for pancreas tissue include AAV8. Serotypes for photoreceptor cells include AAV2, AAV5, and AAV8. Serotypes for retinal pigment epithelium tissue include AAV1, AAV2, AAV4, AAV5, and AAV8. Serotypes for skeletal muscle tissue include AAV1, AAV6, AAV7, AAV8, and AAV9. Serotypes for liver tissue include AAV7, AAV8, and AAV9, and particularly AAV8.

[00220] Tropism can be further refined through pseudotyping, which is the mixing of a capsid and a genome from different viral serotypes. For example AAV2/5 indicates a virus containing the genome of serotype 2 packaged in the capsid from serotype 5. Use of pseudotyped viruses can improve transduction efficiency, as well as alter tropism. Hybrid capsids derived from different serotypes can also be used to alter viral tropism. For example, AAV-DJ contains a hybrid capsid from eight serotypes and displays high infectivity across a broad range of cell types *in vivo*. AAV-DJ8 is another example that displays the properties of AAV-DJ but with enhanced brain uptake. AAV serotypes can also be modified through mutations. Examples of mutational modifications of AAV2 include Y444F, Y500F, Y730F, and S662V. Examples of mutational modifications of AAV3 include Y705F, Y731F, and T492V. Examples of mutational modifications of AAV6 include S663V and T492V. Other pseudotyped/modified AAV variants include AAV2/1, AAV2/6, AAV2/7, AAV2/8, AAV2/9, AAV2.5, AAV8.2, and AAV/SASTG.

[00221] To accelerate transgene expression, self-complementary AAV (scAAV) variants can be used. Because AAV depends on the cell's DNA replication machinery to synthesize the complementary strand of the AAV's single-stranded DNA genome, transgene expression may be delayed. To address this delay, scAAV containing complementary sequences that are capable of

spontaneously annealing upon infection can be used, eliminating the requirement for host cell DNA synthesis. However, single-stranded AAV (ssAAV) vectors can also be used.

[00222] To increase packaging capacity, longer transgenes may be split between two AAV transfer plasmids, the first with a 3' splice donor and the second with a 5' splice acceptor. Upon co-infection of a cell, these viruses form concatemers, are spliced together, and the full-length transgene can be expressed. Although this allows for longer transgene expression, expression is less efficient. Similar methods for increasing capacity utilize homologous recombination. For example, a transgene can be divided between two transfer plasmids but with substantial sequence overlap such that co-expression induces homologous recombination and expression of the full-length transgene.

[00223] Introduction of nucleic acids and proteins can also be accomplished by lipid nanoparticle (LNP)-mediated delivery. For example, LNP-mediated delivery can be used to deliver a guide RNA in the form of RNA. Delivery through such methods results in transient presence of the guide RNA, and the biodegradable lipids improve clearance, improve tolerability, and decrease immunogenicity. Lipid formulations can protect biological molecules from degradation while improving their cellular uptake. Lipid nanoparticles are particles comprising a plurality of lipid molecules physically associated with each other by intermolecular forces. These include microspheres (including unilamellar and multilamellar vesicles, e.g., liposomes), a dispersed phase in an emulsion, micelles, or an internal phase in a suspension. Such lipid nanoparticles can be used to encapsulate one or more nucleic acids or proteins for delivery. Formulations which contain cationic lipids are useful for delivering polyanions such as nucleic acids. Other lipids that can be included are neutral lipids (i.e., uncharged or zwitterionic lipids), anionic lipids, helper lipids that enhance transfection, and stealth lipids that increase the length of time for which nanoparticles can exist *in vivo*. Examples of suitable cationic lipids, neutral lipids, anionic lipids, helper lipids, and stealth lipids can be found in WO 2016/010840 A1, herein incorporated by reference in its entirety for all purposes. An exemplary lipid nanoparticle can comprise a cationic lipid and one or more other components. In one example, the other component can comprise a helper lipid such as cholesterol. In another example, the other components can comprise a helper lipid such as cholesterol and a neutral lipid such as DSPC. In another example, the other components can comprise a helper lipid such as cholesterol, an optional neutral lipid such as DSPC, and a stealth lipid such as S010, S024, S027,

S031, or S033.

[00224] The LNP may contain one or more or all of the following: (i) a lipid for encapsulation and for endosomal escape; (ii) a neutral lipid for stabilization; (iii) a helper lipid for stabilization; and (iv) a stealth lipid. *See, e.g.,* Finn et al. (2018) *Cell Reports* 22:1-9 and WO 2017/173054 A1, each of which is herein incorporated by reference in its entirety for all purposes. In certain LNPs, the cargo can include a guide RNA or a nucleic acid encoding a guide RNA. In certain LNPs, the cargo can include a guide RNA or a nucleic acid encoding a guide RNA and an exogenous donor nucleic acid.

[00225] The lipid for encapsulation and endosomal escape can be a cationic lipid. The lipid can also be a biodegradable lipid, such as a biodegradable ionizable lipid. One example of a suitable lipid is Lipid A or LP01, which is (9Z,12Z)-3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl octadeca-9,12-dienoate, also called 3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl (9Z,12Z)-octadeca-9,12-dienoate. *See, e.g.,* Finn et al. (2018) *Cell Reports* 22:1-9 and WO 2017/173054 A1, each of which is herein incorporated by reference in its entirety for all purposes. Another example of a suitable lipid is Lipid B, which is ((5-((dimethylamino)methyl)-1,3-phenylene)bis(oxy))bis(octane-8,1-diyl)bis(decanoate), also called ((5-((dimethylamino)methyl)-1,3-phenylene)bis(oxy))bis(octane-8,1-diyl)bis(decanoate). Another example of a suitable lipid is Lipid C, which is 2-((4-(((3-(dimethylamino)propoxy)carbonyl)oxy)hexadecanoyl)oxy)propane-1,3-diyl(9Z,9'Z,12Z,12'Z)-bis(octadeca-9,12-dienoate). Another example of a suitable lipid is Lipid D, which is 3-(((3-(dimethylamino)propoxy)carbonyl)oxy)-13-(octanoyloxy)tridecyl 3-octylundecanoate. Other suitable lipids include heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (also known as Dlin-MC3-DMA (MC3))).

[00226] Some such lipids suitable for use in the LNPs described herein are biodegradable *in vivo*. For example, LNPs comprising such a lipid include those where at least 75% of the lipid is cleared from the plasma within 8, 10, 12, 24, or 48 hours, or 3, 4, 5, 6, 7, or 10 days. As another example, at least 50% of the LNP is cleared from the plasma within 8, 10, 12, 24, or 48 hours, or 3, 4, 5, 6, 7, or 10 days.

[00227] Such lipids may be ionizable depending upon the pH of the medium they are in. For example, in a slightly acidic medium, the lipids may be protonated and thus bear a positive

charge. Conversely, in a slightly basic medium, such as, for example, blood where pH is approximately 7.35, the lipids may not be protonated and thus bear no charge. In some embodiments, the lipids may be protonated at a pH of at least about 9, 9.5, or 10. The ability of such a lipid to bear a charge is related to its intrinsic pKa. For example, the lipid may, independently, have a pKa in the range of from about 5.8 to about 6.2.

[00228] Neutral lipids function to stabilize and improve processing of the LNPs. Examples of suitable neutral lipids include a variety of neutral, uncharged or zwitterionic lipids. Examples of neutral phospholipids suitable for use in the present disclosure include, but are not limited to, 5-heptadecylbenzene-1,3-diol (resorcinol), dipalmitoylphosphatidylcholine (DPPC), distearoylphosphatidylcholine (DSPC), phosphocholine (DOPC), dimyristoylphosphatidylcholine (DMPC), phosphatidylcholine (PLPC), 1,2-distearoyl-sn-glycero-3-phosphocholine (DAPC), phosphatidylethanolamine (PE), egg phosphatidylcholine (EPC), dilauryloylphosphatidylcholine (DLPC), dimyristoylphosphatidylcholine (DMPC), 1-myristoyl-2-palmitoyl phosphatidylcholine (MPPC), 1-palmitoyl-2-myristoyl phosphatidylcholine (PMPC), 1-palmitoyl-2-stearoyl phosphatidylcholine (PSPC), 1,2-diarachidoyl-sn-glycero-3-phosphocholine (DBPC), 1-stearoyl-2-palmitoyl phosphatidylcholine (SPPC), 1,2-dieicosenoyl-sn-glycero-3-phosphocholine (DEPC), palmitoyloleoyl phosphatidylcholine (POPC), lysophosphatidyl choline, dioleoyl phosphatidylethanolamine (DOPE), dilinoleoylphosphatidylcholine distearoylphosphatidylethanolamine (DSPE), dimyristoyl phosphatidylethanolamine (DMPE), dipalmitoyl phosphatidylethanolamine (DPPE), palmitoyloleoyl phosphatidylethanolamine (POPE), lysophosphatidylethanolamine, and combinations thereof. For example, the neutral phospholipid may be selected from the group consisting of distearoylphosphatidylcholine (DSPC) and dimyristoyl phosphatidyl ethanolamine (DMPE).

[00229] Helper lipids include lipids that enhance transfection. The mechanism by which the helper lipid enhances transfection can include enhancing particle stability. In certain cases, the helper lipid can enhance membrane fusogenicity. Helper lipids include steroids, sterols, and alkyl resorcinols. Examples of suitable helper lipids suitable include cholesterol, 5-heptadecylresorcinol, and cholesterol hemisuccinate. In one example, the helper lipid may be cholesterol or cholesterol hemisuccinate.

[00230] Stealth lipids include lipids that alter the length of time the nanoparticles can exist in

vivo. Stealth lipids may assist in the formulation process by, for example, reducing particle aggregation and controlling particle size. Stealth lipids may modulate pharmacokinetic properties of the LNP. Suitable stealth lipids include lipids having a hydrophilic head group linked to a lipid moiety.

[00231] The hydrophilic head group of stealth lipid can comprise, for example, a polymer moiety selected from polymers based on PEG (sometimes referred to as poly(ethylene oxide)), poly(oxazoline), poly(vinyl alcohol), poly(glycerol), poly(N-vinylpyrrolidone), polyaminoacids, and poly N-(2-hydroxypropyl)methacrylamide. The term PEG means any polyethylene glycol or other polyalkylene ether polymer. In certain LNP formulations, the PEG, is a PEG-2K, also termed PEG 2000, which has an average molecular weight of about 2,000 daltons. *See, e.g.,* WO 2017/173054 A1, herein incorporated by reference in its entirety for all purposes.

[00232] The lipid moiety of the stealth lipid may be derived, for example, from diacylglycerol or diacylglycamide, including those comprising a dialkylglycerol or dialkylglycamide group having alkyl chain length independently comprising from about C4 to about C40 saturated or unsaturated carbon atoms, wherein the chain may comprise one or more functional groups such as, for example, an amide or ester. The dialkylglycerol or dialkylglycamide group can further comprise one or more substituted alkyl groups.

[00233] As one example, the stealth lipid may be selected from PEG-dilauroylglycerol, PEG-dimyristoylglycerol (PEG-DMG), PEG-dipalmitoylglycerol, PEG-distearoylglycerol (PEG-DSPE), PEG-dilaurylglycamide, PEG-dimyristylglycamide, PEG-dipalmitoylglycamide, and PEG-distearoylglycamide, PEG-cholesterol (1-[8'-(Cholest-5-en-3[beta]-oxy)carboxamido-3',6'-dioxaoctanyl]carbamoyl-[omega]-methyl-poly(ethylene glycol)), PEG-DMB (3,4-ditetradecoxybenzyl-[omega]-methyl-poly(ethylene glycol)ether), 1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000] (PEG2k-DMG), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000] (PEG2k-DSPE), 1,2-distearoyl-sn-glycerol, methoxypolyethylene glycol (PEG2k-DSG), poly(ethylene glycol)-2000-dimethacrylate (PEG2k-DMA), and 1,2-distearoyloxypropyl-3-amine-N-[methoxy(polyethylene glycol)-2000] (PEG2k-DSA). In one particular example, the stealth lipid may be PEG2k-DMG.

[00234] The LNPs can comprise different respective molar ratios of the component lipids in the formulation. The mol-% of the CCD lipid may be, for example, from about 30 mol-% to

about 60 mol-%, from about 35 mol-% to about 55 mol-%, from about 40 mol-% to about 50 mol-%, from about 42 mol-% to about 47 mol-%, or about 45%. The mol-% of the helper lipid may be, for example, from about 30 mol-% to about 60 mol-%, from about 35 mol-% to about 55 mol-%, from about 40 mol-% to about 50 mol-%, from about 41 mol-% to about 46 mol-%, or about 44 mol-%. The mol-% of the neutral lipid may be, for example, from about 1 mol-% to about 20 mol-%, from about 5 mol-% to about 15 mol-%, from about 7 mol-% to about 12 mol-%, or about 9 mol-%. The mol-% of the stealth lipid may be, for example, from about 1 mol-% to about 10 mol-%, from about 1 mol-% to about 5 mol-%, from about 1 mol-% to about 3 mol-%, about 2 mol-%, or about 1 mol-%.

[00235] The LNPs can have different ratios between the positively charged amine groups of the biodegradable lipid (N) and the negatively charged phosphate groups (P) of the nucleic acid to be encapsulated. This may be mathematically represented by the equation N/P. For example, the N/P ratio may be from about 0.5 to about 100, from about 1 to about 50, from about 1 to about 25, from about 1 to about 10, from about 1 to about 7, from about 3 to about 5, from about 4 to about 5, about 4, about 4.5, or about 5.

[00236] In some LNPs, the cargo can comprise exogenous donor nucleic acid and gRNA. The exogenous donor nucleic acid and gRNAs can be in different ratios. For example, the LNP formulation can include a ratio of exogenous donor nucleic acid to gRNA nucleic acid ranging from about 25:1 to about 1:25, ranging from about 10:1 to about 1:10, ranging from about 5:1 to about 1:5, or about 1:1. Alternatively, the LNP formulation can include a ratio of exogenous donor nucleic acid to gRNA nucleic acid from about 1:1 to about 1:5, about 5:1 to about 1:1, about 10:1, or about 1:10. Alternatively, the LNP formulation can include a ratio of exogenous donor nucleic acid to gRNA nucleic acid of about 1:10, 25:1, 10:1, 5:1, 3:1, 1:1, 1:3, 1:5, 1:10, or 1:25.

[00237] A specific example of a suitable LNP has a nitrogen-to-phosphate (N/P) ratio of 4.5 and contains biodegradable cationic lipid, cholesterol, DSPC, and PEG2k-DMG in a 45:44:9:2 molar ratio. The biodegradable cationic lipid can be (9Z,12Z)-3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl octadeca-9,12-dienoate, also called 3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl (9Z,12Z)-octadeca-9,12-dienoate. *See, e.g.,* Finn et al. (2018) *Cell Reports* 22:1-9, herein incorporated by reference in its entirety for all

purposes. Another specific example of a suitable LNP contains Dlin-MC3-DMA (MC3), cholesterol, DSPC, and PEG-DMG in a 50:38.5:10:1.5 molar ratio.

[00238] The mode of delivery can be selected to decrease immunogenicity. For example, a gRNA and an exogenous donor nucleic acid may be delivered by different modes (e.g., bi-modal delivery). These different modes may confer different pharmacodynamics or pharmacokinetic properties on the subject delivered molecule (e.g., gRNA or nucleic acid encoding, or exogenous donor nucleic acid/repair template). For example, the different modes can result in different tissue distribution, different half-life, or different temporal distribution. Some modes of delivery (e.g., delivery of a nucleic acid vector that persists in a cell by autonomous replication or genomic integration) result in more persistent expression and presence of the molecule, whereas other modes of delivery are transient and less persistent (e.g., delivery of an RNA or a protein). Delivery of components in a more transient manner, for example as RNA or protein, can ensure that the Cas/gRNA complex is only present and active for a short period of time and can reduce immunogenicity. Such transient delivery can also reduce the possibility of off-target modifications.

[00239] Administration *in vivo* can be by any suitable route including, for example, parenteral, intravenous, oral, subcutaneous, intra-arterial, intracranial, intrathecal, intraperitoneal, topical, intranasal, or intramuscular. Systemic modes of administration include, for example, oral and parenteral routes. Examples of parenteral routes include intravenous, intraarterial, intraosseous, intramuscular, intradermal, subcutaneous, intranasal, and intraperitoneal routes. A specific example is intravenous infusion. Local modes of administration include, for example, intrathecal, intracerebroventricular, intraparenchymal (e.g., localized intraparenchymal delivery to the striatum (e.g., into the caudate or into the putamen), cerebral cortex, precentral gyrus, hippocampus (e.g., into the dentate gyrus or CA3 region), temporal cortex, amygdala, frontal cortex, thalamus, cerebellum, medulla, hypothalamus, tectum, tegmentum, or substantia nigra), intraocular, intraorbital, subconjunctival, intravitreal, subretinal, and transscleral routes. Significantly smaller amounts of the components (compared with systemic approaches) may exert an effect when administered locally (for example, intraparenchymal or intravitreal) compared to when administered systemically (for example, intravenously). Local modes of administration may also reduce or eliminate the incidence of potentially toxic side effects that may occur when therapeutically effective amounts of a component are administered

systemically.

[00240] Administration *in vivo* can be by any suitable route including, for example, parenteral, intravenous, oral, subcutaneous, intra-arterial, intracranial, intrathecal, intraperitoneal, topical, intranasal, or intramuscular. A specific example is intravenous infusion. Nasal instillation and intravitreal injection are other specific examples. Compositions comprising the guide RNAs (or nucleic acids encoding the guide RNAs) can be formulated using one or more physiologically and pharmaceutically acceptable carriers, diluents, excipients or auxiliaries. The formulation can depend on the route of administration chosen. The term “pharmaceutically acceptable” means that the carrier, diluent, excipient, or auxiliary is compatible with the other ingredients of the formulation and not substantially deleterious to the recipient thereof.

[00241] The frequency of administration and the number of dosages can be depend on the half-life of the exogenous donor nucleic acids or guide RNAs (or nucleic acids encoding the guide RNAs) and the route of administration among other factors. The introduction of nucleic acids or proteins into the cell or non-human animal can be performed one time or multiple times over a period of time. For example, the introduction can be performed at least two times over a period of time, at least three times over a period of time, at least four times over a period of time, at least five times over a period of time, at least six times over a period of time, at least seven times over a period of time, at least eight times over a period of time, at least nine times over a period of times, at least ten times over a period of time, at least eleven times, at least twelve times over a period of time, at least thirteen times over a period of time, at least fourteen times over a period of time, at least fifteen times over a period of time, at least sixteen times over a period of time, at least seventeen times over a period of time, at least eighteen times over a period of time, at least nineteen times over a period of time, or at least twenty times over a period of time.

D. Measuring CRISPR/Cas Activity *In Vivo* and Assessing Modification of a Target Genomic Locus

[00242] The methods disclosed herein can further comprise assessing modification of the target genomic locus. The methods for detecting or measuring expression or activity will depend on the target genomic locus being modified.

[00243] For example, if the target genomic locus comprises a gene encoding an RNA or

protein, and the intended modification is to change expression of the encoded RNA or protein, the method of assessing modification of the target genomic locus can comprise measuring expression or activity of the encoded RNA or protein. For example, if the encoded protein is a protein released into the serum, serum levels of the encoded protein can be measured. Assays for measuring levels and activity of RNA and proteins are well known.

[00244] Alternatively, the methods disclosed herein can further comprise identifying a cell having a modified target genomic locus in which the sequence has been modified by non-homologous end joining (e.g., presence of small insertions or deletions (indels)) following cleavage by CRISPR/Cas, in which a sequence at the target genomic locus between two guide RNA target sequences has been excised, or in which the target genomic locus has been modified by recombination with an exogenous donor nucleic acid. Various methods can be used to identify cells having a targeted genetic modification. The screening can comprise a quantitative assay for assessing modification of allele (MOA) of a parental chromosome. For example, the quantitative assay can be carried out via a quantitative PCR, such as a real-time PCR (qPCR). The real-time PCR can utilize a first primer set that recognizes the target locus and a second primer set that recognizes a non-targeted reference locus. The primer set can comprise a fluorescent probe that recognizes the amplified sequence. Other examples of suitable quantitative assays include fluorescence-mediated in situ hybridization (FISH), comparative genomic hybridization, isothermic DNA amplification, quantitative hybridization to an immobilized probe(s), INVADER[®] Probes, TAQMAN[®] Molecular Beacon probes, or ECLIPSE[™] probe technology (*see, e.g.*, US 2005/0144655, herein incorporated by reference in its entirety for all purposes).

[00245] Next-generation sequencing (NGS) can also be used for screening. Next-generation sequencing can also be referred to as “NGS” or “massively parallel sequencing” or “high throughput sequencing.” NGS can be used as a screening tool in addition to the MOA assays to define the exact nature of the targeted genetic modification and whether it is consistent across cell types or tissue types or organ types.

[00246] Assessing modification of the target genomic locus in a non-human animal can be in any cell type from any tissue or organ. For example, modification of the target genomic locus can be assessed in multiple cell types from the same tissue or organ or in cells from multiple locations within the tissue or organ. This can provide information about which cell types within

a target tissue or organ are being modified or which sections of a tissue or organ are being reached by the CRISPR/Cas and modified. As another example, modification of the target genomic locus can be assessed in multiple types of tissue or in multiple organs. In methods in which a particular tissue or organ is being targeted, this can provide information about how effectively that tissue or organ is being targeted and whether there are off-target effects in other tissues or organs.

[00247] In some specific examples, Cas9-ready non-human animals can be used to evaluate the editing rates of various guide RNAs. Guide RNAs may be introduced as either single guide RNA (modified and unmodified) or duplex RNA, or expressed under a U6 promoter (e.g., via AAV). Cas9-ready non-human animals can also be crossed to non-human animals comprising humanized alleles non-human animals expressing guide RNAs for evaluation in disease modeling.

IV. Methods of Making Non-Human Animals Comprising a Cas Expression Cassette and/or a Recombinase Expression Cassette

[00248] Various methods are provided for making a non-human animal comprising one or more or all of a Cas expression cassette and a recombinase expression cassette as disclosed elsewhere herein. Any convenient method or protocol for producing a genetically modified organism is suitable for producing such a genetically modified non-human animal. *See, e.g.,* Cho *et al.* (2009) *Current Protocols in Cell Biology* 42:19.11:19.11.1–19.11.22 and Gama Sosa *et al.* (2010) *Brain Struct. Funct.* 214(2-3):91-109, each of which is herein incorporated by reference in its entirety for all purposes. Such genetically modified non-human animals can be generated, for example, through gene knock-in at a targeted locus (e.g., a safe harbor locus such as *Rosa26*) or through use of a randomly integrating transgene. *See, e.g.,* WO 2014/093622 and WO 2013/176772, each of which is herein incorporated by reference in its entirety for all purposes. Methods of targeting a construct to the *Rosa26* locus are described, for example, in US 2012/0017290, US 2011/0265198, and US 2013/0236946, each of which is herein incorporated by reference in its entirety for all purposes.

[00249] For example, the method of producing a non-human animal comprising one or more or all of a Cas expression cassette and a recombinase expression cassette as disclosed elsewhere herein can comprise: (1) modifying the genome of a pluripotent cell to comprise one or more or

all of a Cas expression cassette and a recombinase expression cassette; (2) identifying or selecting the genetically modified pluripotent cell comprising the one or more or all of a Cas expression cassette and a recombinase expression cassette; (3) introducing the genetically modified pluripotent cell into a non-human animal host embryo; and (4) implanting and gestating the host embryo in a surrogate mother. Optionally, the host embryo comprising modified pluripotent cell (e.g., a non-human ES cell) can be incubated until the blastocyst stage before being implanted into and gestated in the surrogate mother to produce an F0 non-human animal. The surrogate mother can then produce an F0 generation non-human animal comprising one or more or all of a Cas expression cassette and a recombinase expression cassette.

[00250] The methods can further comprise identifying a cell or animal having a modified target genomic locus. Various methods can be used to identify cells and animals having a targeted genetic modification.

[00251] The screening step can comprise, for example, a quantitative assay for assessing modification of allele (MOA) of a parental chromosome. For example, the quantitative assay can be carried out via a quantitative PCR, such as a real-time PCR (qPCR). The real-time PCR can utilize a first primer set that recognizes the target locus and a second primer set that recognizes a non-targeted reference locus. The primer set can comprise a fluorescent probe that recognizes the amplified sequence.

[00252] Other examples of suitable quantitative assays include fluorescence-mediated in situ hybridization (FISH), comparative genomic hybridization, isothermic DNA amplification, quantitative hybridization to an immobilized probe(s), INVADER[®] Probes, TAQMAN[®] Molecular Beacon probes, or ECLIPSE[™] probe technology (*see, e.g.,* US 2005/0144655, incorporated herein by reference in its entirety for all purposes).

[00253] An example of a suitable pluripotent cell is an embryonic stem (ES) cell (e.g., a mouse ES cell or a rat ES cell). The modified pluripotent cell can be generated, for example, by (a) introducing into the cell one or more targeting vectors comprising an insert nucleic acid flanked by 5' and 3' homology arms corresponding to 5' and 3' target sites, wherein the insert nucleic acid comprises one or more or all of a Cas expression cassette and a recombinase expression cassette; and (b) identifying at least one cell comprising in its genome the insert nucleic acid integrated at the target genomic locus. Alternatively, the modified pluripotent cell can be generated by (a) introducing into the cell: (i) a nuclease agent, wherein the nuclease agent

induces a nick or double-strand break at a target sequence within the target genomic locus; and (ii) one or more targeting vectors comprising an insert nucleic acid flanked by 5' and 3' homology arms corresponding to 5' and 3' target sites located in sufficient proximity to the target sequence, wherein the insert nucleic acid comprises one or more or all of a Cas expression cassette and a recombinase expression cassette; and (c) identifying at least one cell comprising a modification (e.g., integration of the insert nucleic acid) at the target genomic locus. Any nuclease agent that induces a nick or double-strand break into a desired target sequence can be used. Examples of suitable nucleases include a Transcription Activator-Like Effector Nuclease (TALEN), a zinc-finger nuclease (ZFN), a meganuclease, and Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)/CRISPR-associated (Cas) systems or components of such systems (e.g., CRISPR/Cas9). *See, e.g.*, US 2013/0309670 and US 2015/0159175, each of which is herein incorporated by reference in its entirety for all purposes.

[00254] The donor cell can be introduced into a host embryo at any stage, such as the blastocyst stage or the pre-morula stage (i.e., the 4 cell stage or the 8 cell stage). Progeny that are capable of transmitting the genetic modification through the germline are generated. *See, e.g.*, US Patent No. 7,294,754, herein incorporated by reference in its entirety for all purposes.

[00255] Alternatively, the method of producing the non-human animals described elsewhere herein can comprise: (1) modifying the genome of a one-cell stage embryo to comprise one or more or all of a Cas expression cassette and a recombinase expression cassette using the methods described above for modifying pluripotent cells; (2) selecting the genetically modified embryo; and (3) implanting and gestating the genetically modified embryo into a surrogate mother. Progeny that are capable of transmitting the genetic modification through the germline are generated.

[00256] Nuclear transfer techniques can also be used to generate the non-human mammalian animals. Briefly, methods for nuclear transfer can include the steps of: (1) enucleating an oocyte or providing an enucleated oocyte; (2) isolating or providing a donor cell or nucleus to be combined with the enucleated oocyte; (3) inserting the cell or nucleus into the enucleated oocyte to form a reconstituted cell; (4) implanting the reconstituted cell into the womb of an animal to form an embryo; and (5) allowing the embryo to develop. In such methods, oocytes are generally retrieved from deceased animals, although they may be isolated also from either oviducts and/or ovaries of live animals. Oocytes can be matured in a variety of well-known

media prior to enucleation. Enucleation of the oocyte can be performed in a number of well-known manners. Insertion of the donor cell or nucleus into the enucleated oocyte to form a reconstituted cell can be by microinjection of a donor cell under the zona pellucida prior to fusion. Fusion may be induced by application of a DC electrical pulse across the contact/fusion plane (electrofusion), by exposure of the cells to fusion-promoting chemicals, such as polyethylene glycol, or by way of an inactivated virus, such as the Sendai virus. A reconstituted cell can be activated by electrical and/or non-electrical means before, during, and/or after fusion of the nuclear donor and recipient oocyte. Activation methods include electric pulses, chemically induced shock, penetration by sperm, increasing levels of divalent cations in the oocyte, and reducing phosphorylation of cellular proteins (as by way of kinase inhibitors) in the oocyte. The activated reconstituted cells, or embryos, can be cultured in well-known media and then transferred to the womb of an animal. *See, e.g.*, US 2008/0092249, WO 1999/005266, US 2004/0177390, WO 2008/017234, and US Patent No. 7,612,250, each of which is herein incorporated by reference in its entirety for all purposes.

[00257] The various methods provided herein allow for the generation of a genetically modified non-human F0 animal wherein the cells of the genetically modified F0 animal comprise the one or more or all of a Cas expression cassette and a recombinase expression cassette. It is recognized that depending on the method used to generate the F0 animal, the number of cells within the F0 animal that have the one or more or all of a Cas expression cassette and a recombinase expression cassette will vary. The introduction of the donor ES cells into a pre-morula stage embryo from a corresponding organism (e.g., an 8-cell stage mouse embryo) via for example, the VELOCIMOUSE[®] method allows for a greater percentage of the cell population of the F0 animal to comprise cells having the nucleotide sequence of interest comprising the targeted genetic modification. For example, at least 50%, 60%, 65%, 70%, 75%, 85%, 86%, 87%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the cellular contribution of the non-human F0 animal can comprise a cell population having the targeted modification.

[00258] The cells of the genetically modified F0 animal can be heterozygous for one or more or all of a Cas expression cassette and a recombinase expression cassette or can be homozygous for one or more or all of a Cas expression cassette and a recombinase expression cassette.

[00259] All patent filings, websites, other publications, accession numbers and the like cited

above or below are incorporated by reference in their entirety for all purposes to the same extent as if each individual item were specifically and individually indicated to be so incorporated by reference. If different versions of a sequence are associated with an accession number at different times, the version associated with the accession number at the effective filing date of this application is meant. The effective filing date means the earlier of the actual filing date or filing date of a priority application referring to the accession number if applicable. Likewise, if different versions of a publication, website or the like are published at different times, the version most recently published at the effective filing date of the application is meant unless otherwise indicated. Any feature, step, element, embodiment, or aspect of the invention can be used in combination with any other unless specifically indicated otherwise. Although the present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.

BRIEF DESCRIPTION OF THE SEQUENCES

[00260] The nucleotide and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three-letter code for amino acids. The nucleotide sequences follow the standard convention of beginning at the 5' end of the sequence and proceeding forward (i.e., from left to right in each line) to the 3' end. Only one strand of each nucleotide sequence is shown, but the complementary strand is understood to be included by any reference to the displayed strand. When a nucleotide sequence encoding an amino acid sequence is provided, it is understood that codon degenerate variants thereof that encode the same amino acid sequence are also provided. The amino acid sequences follow the standard convention of beginning at the amino terminus of the sequence and proceeding forward (i.e., from left to right in each line) to the carboxy terminus.

[00261] Table 3. Description of Sequences.

SEQ ID NO	Type	Description
1	DNA	MAID2599 Cas9 Allele
2	Protein	T2A
3	Protein	P2A
4	Protein	E2A
5	Protein	F2A
6	RNA	Generic Guide RNA Scaffold v.2
7	RNA	Generic Guide RNA Scaffold v.3

SEQ ID NO	Type	Description
8	RNA	Generic Guide RNA Scaffold v.4
9	DNA	Generic Guide RNA Target Sequence plus PAM v.1
10	DNA	Generic Guide RNA Target Sequence plus PAM v.2
11	DNA	Generic Guide RNA Target Sequence plus PAM v.3
12	DNA	MAID2600 Cas9 Allele
13	Protein	Cas9-P2A-eGFP Protein
14	DNA	MAID2658 Cas9 Allele
15	DNA	MAID2659 Cas9 Allele
16	Protein	3xFLAG-Cas9-P2A-eGFP Protein
17	DNA	MAID2660 Cas9 Allele
18	DNA	MAID2661 Cas9 Allele
19	Protein	Cas9 Protein
20	DNA	MAID2672 Cas9 Allele
21	DNA	MAID2673 Cas9 Allele
22	Protein	3xFLAG-Cas9 Protein
23	Protein	3xFLAG
24	Protein	eGFP
25	RNA	crRNA Tail
26	RNA	TracrRNA
27	RNA	Generic Guide RNA Scaffold v. 1
28	DNA	Cas9-P2A-eGFP Coding Sequence
29	DNA	3xFLAG-Cas9-P2A-eGFP Coding Sequence
30	DNA	Cas9 Coding Sequence
31	DNA	3xFLAG-Cas9 Coding Sequence
32	DNA	GSG-P2A Coding Sequence
33	DNA	eGFP Coding Sequence
34	DNA	3xFLAG Coding Sequence
35	DNA	Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element
36	DNA	Bovine Growth Hormone Polyadenylation Signal
37	DNA	Neo-PolyA
38	DNA	Chicken Beta Actin Promoter/Enhancer Coupled with the Cytomegalovirus (CMV) Immediate-Early Enhancer (CAGG)

EXAMPLES

Example 1. Validation of Cas9-Ready Mice

[00262] CRISPR/Cas9, an RNA-guided DNA endonuclease, catalyzes the creation of a double-strand break (DSB) of DNA at the binding site of its RNA guide. An exemplary RNA guide can consist of a 42 nucleotide CRISPR RNA (crRNA) that joins with an 87 nucleotide trans-activating RNA (tracrRNA). The tracrRNA is complementary to and base pairs with the crRNA, forming a functional crRNA/tracrRNA guide. This duplex RNA becomes bound to the Cas9 protein to form an active ribonucleoprotein (RNP) that can interrogate the genome for complementarity with the 20-nucleotide guide portion of the crRNA. A secondary requirement for strand breakage is that the Cas9 protein must recognize a protospacer adjacent motif (PAM) directly adjacent to the sequence complementary to the guide portion of crRNA (the crRNA

target sequence). Alternatively, an active RNP complex can also be formed by replacing the crRNA/tracrRNA duplex with a single guide RNA (sgRNA) formed by covalently joining the crRNA and the tracrRNA. Such a sgRNA can be formed, for example, by fusing the 20 nucleotide guide portion of the crRNA directly to the processed tracrRNA sequence. The sgRNA can interact with both the Cas9 protein and the DNA in the same way and with similar efficiency as the crRNA/tracrRNA duplex would. The CRISPR bacterial natural defense mechanism has been shown to function effectively in mammalian cells and to activate break induced endogenous repair pathways. When a double strand break occurs in the genome, repair pathways will attempt to fix the DNA by either the canonical or alternative non-homologous end joining (NHEJ) pathways or homologous recombination, also referred to as homology-directed repair (HDR), if an appropriate template is available. We can leverage these pathways to facilitate site specific deletion of genomic regions or insertion of exogenous DNA or HDR in mammalian cells.

[00263] The CRISPR/Cas9 system is a powerful tool for genome engineering. However, one limitation of the system for use *in vivo* is the need to simultaneously introduce all components into a living organism. The typical method of introducing these components into cells is to transiently transfect DNA constructs into cells that will generate the appropriate RNAs and protein. Though effective, this approach has an inherent disadvantage as the cells must rely on the plasmid DNA constructs to first undergo transcription and then translation before the Cas9 protein is available to interact with the sgRNA component. We believe that Cas9-induced mutation frequency and recombination frequency can be vastly improved by having the protein constitutively available.

[00264] The wild-type Cas9 coding sequence (CDS) was codon-optimized for expression in mice. An N-terminal monopartite nuclear localization (NLS) signal, a C-terminal bipartite NLS, and C-terminal P2A linked GFP fluorescent reporter were then incorporated. The Cas9 expression cassette (MAID2599) is depicted in **Figures 1** and **14** and SEQ ID NO: 1. The P2A-GFP can be used for better tracking of Cas9 expression *in vivo*. These components were engineered into the first intron of the *Rosa26* locus of the mouse genome along with a preceding floxed neomycin resistance cassette (neo cassette) with appropriate splicing signals and a strong polyadenylation (polyA) signal. The components of the Cas9 expression cassette from 5' to 3' are shown in **Table 4**, and the components of the Cas9 allele following removal of the floxed

neomycin cassette are shown in **Table 5**. The Cas9-P2A-eGFP protein sequence encoded by the allele is set forth in SEQ ID NO: 13.

[00265] Table 4. Components of MAID2599 Cas9 Allele.

Component	Nucleotide Region Within SEQ ID NO: 1
Mouse Rosa26 upstream sequence	1-170
First loxP site	300-333
Sequence encoding neomycin phosphotransferase for resistance to neomycin family antibiotics (e.g. G418), with a polyadenylation signal	424-2489
Second loxP site	2517-2550
Kozak sequence	2599-2608
Codon-optimized Cas9 coding sequence	2605-6777
N-terminal monopartite NLS	2614-2634
C-terminal bipartite NLS	6730-6777
P2A coding sequence	6778-6843
eGFP coding sequence	6844-7557
Woodchuck hepatitis virus posttranscriptional regulatory element (WPRE)	7607-8203
Bovine growth hormone polyadenylation signal (bGH polyA)	8204-8419
Mouse Rosa26 downstream sequence	8479-8628

[00266] Table 5. Components of MAID2600 Cas9 Allele.

Component	Nucleotide Region Within SEQ ID NO: 12
Mouse Rosa26 upstream sequence	1-170
LoxP site	300-333
Kozak sequence	382-391
Codon-optimized Cas9 coding sequence	388-4560
N-terminal monopartite NLS	397-417
C-terminal bipartite NLS	4513-4560
P2A coding sequence	4561-4626
eGFP coding sequence	4627-5340
Woodchuck hepatitis virus posttranscriptional regulatory element (WPRE)	5390-5986
Bovine growth hormone polyadenylation signal (bGH polyA)	5987-6202
Mouse Rosa26 downstream sequence	6262-6411

[00267] Prior to removal of the cassette by the action of Cre recombinase, the neomycin resistance gene will normally be efficiently transcribed and translated; however, the Cas9 CDS will not normally be expressed due to the presence of the strong poly(A) region, which can effectively block run-through transcription. Upon removal of the neo cassette by the action of Cre recombinase, the hybrid mRNA for the Cas9 and GFP proteins will normally be constitutively expressed by the *Rosa26* promoter. Targeted cells before and after neo cassette removal were first verified by loss-of-allele screening to detect the single, site-specific integration of the targeting vector at the *Rosa26* locus. Cas9 and GFP expression were validated

by extracting total RNA from targeted mESCs, followed by reverse transcription to generate cDNA and TAQMAN[®] qPCR to detect the reverse transcribed cDNA (RT-qPCR). Taken together, the system that was created is capable of expressing consistent levels of Cas9 protein continuously or conditionally (by requiring the removal of a neomycin resistance cassette) in mESCs and mice derived from them.

[00268] Another version was designed without the P2A-eGFP (MAID2660). *See Figure 14.* The components of the Cas9 expression cassette from 5' to 3' are shown in **Table 6**, and the components of the Cas9 allele following removal of the floxed neomycin cassette are shown in **Table 7**. The Cas9 protein sequence encoded by the allele is set forth in SEQ ID NO: 19.

[00269] **Table 6. Components of MAID2660 Cas9 Allele.**

Component	Nucleotide Region Within SEQ ID NO: 17
Mouse Rosa26 upstream sequence	1-170
First loxP site	300-333
Sequence encoding neomycin phosphotransferase for resistance to neomycin family antibiotics (e.g. G418), with a polyadenylation signal	424-2489
Second loxP site	2517-2550
Kozak sequence	2599-2608
Codon-optimized Cas9 coding sequence	2605-6777
N-terminal monopartite NLS	2614-2634
C-terminal bipartite NLS	6730-6777
Bovine growth hormone polyadenylation signal (bGH polyA)	6783-6998
Mouse Rosa26 downstream sequence	7058-7207

[00270] **Table 7. Components of MAID2661 Cas9 Allele.**

Component	Nucleotide Region Within SEQ ID NO: 18
Mouse Rosa26 upstream sequence	1-170
LoxP site	300-333
Kozak sequence	382-391
Codon-optimized Cas9 coding sequence	388-4560
N-terminal monopartite NLS	397-417
C-terminal bipartite NLS	4513-4560
Bovine growth hormone polyadenylation signal (bGH polyA)	4566-4781
Mouse Rosa26 downstream sequence	4841-4990

[00271] In addition, two versions with exogenous CAGG promoters and 3xFLAG tag sequences were designed. The first version included the P2A-eGFP (MAID2658), and the second version was designed without the P2A-eGFP (MAID2672). *See Figure 14.* The components of first version of the CAGG-Cas9 expression cassette from 5' to 3' are shown in **Table 8**, and the components of the first version of the CAGG-Cas9 allele following removal of

the floxed neomycin cassette are shown in **Table 9**. The 3xFLAG-Cas9-P2A-eGFP protein sequence encoded by this allele is set forth in SEQ ID NO: 16. The components of second version of the CAGG-Cas9 expression cassette from 5' to 3' are shown in **Table 10**, and the components of the second version of the CAGG-Cas9 allele following removal of the floxed neomycin cassette are shown in **Table 11**. The 3xFLAG-Cas9 protein sequence encoded by this allele is set forth in SEQ ID NO: 22.

[00272] Table 8. Components of MAID2658 Cas9 Allele.

Component	Nucleotide Region Within SEQ ID NO: 14
Mouse Rosa26 upstream sequence	1-170
Chicken beta actin promoter/enhancer coupled with the cytomegalovirus (CMV) immediate-early enhancer (CAGG)	195-1913
First loxP site	1996-2029
Sequence encoding neomycin phosphotransferase for resistance to neomycin family antibiotics (e.g. G418), with a polyadenylation signal	2120-4185
Second loxP site	4213-4246
Kozak sequence	4341-4350
3xFLAG	4350-4415
Codon-optimized Cas9 coding sequence	4416-8588
N-terminal monopartite NLS	4425-4445
C-terminal bipartite NLS	8541-8588
P2A coding sequence	8589-8654
eGFP coding sequence	8655-9368
Woodchuck hepatitis virus posttranscriptional regulatory element (WPRE)	9418-10014
Bovine growth hormone polyadenylation signal (bGH polyA)	10015-10230
Mouse Rosa26 downstream sequence	10290-10439

[00273] Table 9. Components of MAID2659 Cas9 Allele.

Component	Nucleotide Region Within SEQ ID NO: 15
Mouse Rosa26 upstream sequence	1-170
Chicken beta actin promoter/enhancer coupled with the cytomegalovirus (CMV) immediate-early enhancer (CAGG)	195-1913
LoxP site	1996-2029
Kozak sequence	2124-2133
3xFLAG	2133-2198
Codon-optimized Cas9 coding sequence	2199-6371
N-terminal monopartite NLS	2208-2228
C-terminal bipartite NLS	6324-6371
P2A coding sequence	6372-6437
eGFP coding sequence	6438-7151
Woodchuck hepatitis virus posttranscriptional regulatory element (WPRE)	7201-7797
Bovine growth hormone polyadenylation signal (bGH polyA)	7798-8013
Mouse Rosa26 downstream sequence	8073-8222

[00274] Table 10. Components of MAID2672 Cas9 Allele.

Component	Nucleotide Region Within SEQ ID NO: 20
Mouse Rosa26 upstream sequence	1-170
Chicken beta actin promoter/enhancer coupled with the cytomegalovirus (CMV) immediate-early enhancer (CAGG)	205-1923
First loxP site	2006-2039
Sequence encoding neomycin phosphotransferase for resistance to neomycin family antibiotics (e.g. G418), with a polyadenylation signal	2130-4195
Second loxP site	4223-4256
Kozak sequence	4351-4360
3xFLAG	4360-4425
Codon-optimized Cas9 coding sequence	4426-8598
N-terminal monopartite NLS	4435-4455
C-terminal bipartite NLS	8551-8598
Woodchuck hepatitis virus posttranscriptional regulatory element (WPRE)	8645-9241
Bovine growth hormone polyadenylation signal (bGH polyA)	9249-9464
Mouse Rosa26 downstream sequence	9524-9673

[00275] Table 11. Components of MAID2673 Cas9 Allele.

Component	Nucleotide Region Within SEQ ID NO: 21
Mouse Rosa26 upstream sequence	1-170
Chicken beta actin promoter/enhancer coupled with the cytomegalovirus (CMV) immediate-early enhancer (CAGG)	205-1923
LoxP site	2006-2039
Kozak sequence	2134-2143
3xFLAG	2143-2208
Codon-optimized Cas9 coding sequence	2209-6381
N-terminal monopartite NLS	2218-2238
C-terminal bipartite NLS	6334-6381
Woodchuck hepatitis virus posttranscriptional regulatory element (WPRE)	6428-7024
Bovine growth hormone polyadenylation signal (bGH polyA)	7032-7247
Mouse Rosa26 downstream sequence	7307-7456

[00276] To validate the MAID2599/MAID2600 system, Cas9 mESCs with and without the neomycin cassette (MAID2599 and MAID2600, respectively) were transfected with two sgRNAs targeting the start and stop codon regions of a first target gene. *See Figure 2A.* Cas9 cleavage efficiency was then assayed by loss-of-allele screening to assess the proportion of mESC clones having insertion-deletion mutations at the gRNA-targeted Cas9 cleavage sites. The proportion of mESC clones in which the DNA between the Cas9 cleavage sites was deleted on or both target alleles, causing a null mutation, was also determined. Cas9 was able to induce these genomic changes only when the neomycin cassette and poly(A) (stop sequence) had been removed (MAID2600). To better assess genome editing capabilities in regards to homology-

directed repair, an sgRNA targeting a second target gene was introduced along with a single stranded oligodeoxynucleotide (ssODN) as a point mutation donor. *See Figure 2B*. The constitutive Cas9 expression system described herein was compared to traditional methods of introducing Cas9 and sgRNA via plasmids along with an ssODN. The Cas9 expression system described herein, when combined with a plasmid expressing the sgRNA, was able to activate break-induced endogenous repair pathways to incorporate our desired point mutation at a frequency that was equal to that when both Cas9 and the sgRNA were expressed from exogenous plasmids. However, when the Cas9 expression system described herein was combined with directly delivered sgRNA, it induced homology-directed insertional mutagenesis at nearly double the efficiency of plasmid delivery methods.

[00277] To determine the effectiveness of endogenously expressed Cas9 in live mice, these targeted mESCs were microinjected into 8-cell mouse embryos using the VELOCIMOUSE[®] method. *See, e.g.,* US 7,576,259; US 7,659,442; US 7,294,754; US 2008/007800; and Poueymirou et al. (2007) *Nature Biotech.* 25(1):91-99, each of which is herein incorporated by reference in its entirety for all purposes. Specifically, a small hole was created in the zona pellucida to facilitate the injection of targeted mESC. These injected 8-cell embryos were transferred to surrogate mothers to produce live pups carrying the transgene. Upon gestation in a surrogate mother, the injected embryos produced F0 mice that carry no detectable host embryo contribution. The fully ES cell-derived mice were normal, healthy, and fertile (with germline transmission). Tissue was harvested from cassette-deleted F0 mice (MAID2600) for GFP visualization, Cas9 mRNA expression, and Cas9 protein expression. *See Figures 3A-3F* (bright field and GFP visualization) and **Figures 4A-4C** (Cas9 mRNA expression in **Figures 4A** and **4C**, and protein expression in **Figure 4B**). **Figure 3D** shows eGFP expression in heterozygous Rosa26Cas9 mice (MAID2600) but a lack of corresponding eGFP expression in wild type mice in liver, **Figure 3E** shows eGFP expression in heterozygous Rosa26Cas9 mice but a lack of corresponding eGFP expression in wild type mice in kidney, and **Figure 3F** shows eGFP expression in heterozygous Rosa26Cas9 mice but a lack of corresponding eGFP expression in wild type mice in brain. Likewise, Cas9 mRNA expression, assayed by RT-qPCR, was observed in heterozygous Rosa26Cas9 mice in brain, heart, kidney, liver, lung, quadriceps, spleen, and thymus, but no Cas9 mRNA expression was observed in the corresponding tissues from wild-type mice. *See Figures 4A* and **4C**. In the experiments, equal mass amounts of RNA from each

tissue were assayed by RT-qPCR. The data show that Cas9-ready mice express Cas9 mRNA at an easily detectable level in all tissues. Various tissues were harvested from Cas9-ready mice. Three tissues were harvested from fourteen mice and an additional five tissues were harvested from four mice to assess differences from mouse to mouse as well as from tissue to tissue within a mouse. Each of these tissues had the RNA extracted. The genomic DNA was degraded so that it would not count towards the qPCR reaction. The RNA was reverse transcribed and then an assay specific to Cas9 was used to detect Cas9 transcripts. As expected, the Cas9 mouse showed significant expression (ct values below 30) while WT mice showed ct values of 30 and higher indicating that there is no endogenous expression of Cas9 protein.

[00278] Similarly, Cas9 protein expression as determined by western blot using ThermoFisher Cas9 antibody MA5-23519 at a 1:250 dilution and using actin as a control showed Cas9 protein expression in heterozygous Rosa26Cas9 mice (MAID2600) in spleen, liver, and brain, whereas Cas9 protein was not observed in the same tissues in wild type mice. *See Figure 4B*. All three tests indicated a consistent level of expression in all assayed tissues.

[00279] An experiment to knock out target gene 3, which encodes a protein secreted by the liver and found in serum, was then performed by introducing an sgRNA into primary hepatocytes isolated from cassette-deleted Cas9 mice (MAID2600) via lipid nanoparticle (LNP) delivery. *See Figure 5B*. As a control, the same methods of sgRNA introduction were paired with exogenous Cas9 expression in primary hepatocytes isolated from wild type (WT) mice. *See Figure 5A*. Non-homologous end joining was then assessed by next-generation sequencing (NGS) to measure indel frequencies at the target gene 3 locus. In the experiment, there were three conditions: (1) LNP-mediated delivery of GFP mRNA and a control (i.e., dead) sgRNA; (2) LNP-mediated delivery of GFP mRNA and a target gene 3 sgRNA; and (3) LNP-mediated delivery of a Cas9 mRNA and a target gene 3 sgRNA. For each condition, four concentrations of mRNA were tested: 15.6 ng/mL, 62.5 ng/mL, 250 ng/mL, and 1000 ng/mL. In wild type primary mouse hepatocytes, a dose-dependent increase in insertion/deletion frequency was seen only when both Cas9 mRNA and the target gene 3 sgRNA were introduced. In contrast, in Cas9-ready primary mouse hepatocytes, a similar dose-dependent effect was seen when target gene 3 guide RNA was introduced with control GFP mRNA instead of Cas9 mRNA, and the level insertion/deletion frequency was essentially identical to the levels seen when Cas9 mRNA was also introduced.

[00280] An experiment to knock out target gene 3 *in vivo* was then performed by introducing an sgRNA into cassette-deleted Cas9 mice (MAID2600) via hydrodynamic DNA delivery (HDD), lipid nanoparticle (LNP) delivery, or introduction of an adeno-associated virus (AAV) carrying an sgRNA expression sequence by tail vein injection. *See Figures 6A-6D.* As a control, the same methods of sgRNA introduction were paired with exogenous Cas9 expression in wild type (WT) mice. For LNP-mediated delivery, three groups of mice were tested: (1) Cas9-ready mice (3 male + 3 female; 2 mg/kg control sgRNA + GFP mRNA); (2) Cas9-ready mice (3 male + 3 female; 2 mg/kg sgRNA for target gene 3 + GFP mRNA); and (3) WT mice (3 male + 3 female; 2 mg/kg sgRNA for target gene 3 + Cas9 mRNA). For AAV-mediated delivery, two groups of mice were tested: (1) Cas9-ready mice (3 male + 3 female; AAV8-sgRNA for target gene 3); and (2) WT mice (3 male + 3 female; AAV8-sgRNA for target gene 3 + AAV8-Cas9). For HDD, two groups of mice were tested: (1) Cas9-ready mice (3 male + 3 female; sgRNA for target gene 3); and (2) WT mice (3 male + 3 female; sgRNA for target gene 3 + Cas9). Cas9-ready mice had consistently and significantly more targeted gene inactivation than WT mice with exogenous Cas9 expression.

[00281] Surprisingly, AAV8-mediated delivery of target gene 3 sgRNA to Cas9-ready mice (MAID2600) was more effective than either LNP-mediated delivery or HDD at targeting liver target gene 3. *See Figures 6A-6D.* In addition, AAV8-mediated delivery of target gene 3 sgRNA to Cas9-ready mice was more effective than AAV8-mediated delivery of both target gene 3 sgRNA and Cas9 to WT mice, whereas not much difference was observed between both conditions using LNP-mediated delivery or HDD. *See Figures 6A-6D.* These results indicate that AAV-mediated delivery of guide RNAs to Cas9-ready mice can be a particularly effective means for testing gRNA activity *in vivo*. Serum levels of the protein encoded by target gene 3 (i.e., target protein 3) were measured in female and male mice on days 7 and 21 following introduction of the CRISPR/Cas components. The mice tested included Cas9-ready mice and wild type mice. Controls included Cas9-ready mice and WT mice in which neither Cas9 nor the target gene 3 sgRNA were introduced. Hydrodynamic delivery of the guide RNA or the combination of Cas9 and the guide RNA did not reduce serum levels of target protein 3 in a significant way over control WT mice in which neither Cas9 nor guide RNA were introduced. For LNP-mediated delivery, introduction of the sgRNA into Cas9-ready mice resulted in similar serum levels of target protein compared to WT mice in which both Cas9 and the sgRNA were

introduced, and each of these conditions resulted in reduced serum levels of target protein 3 by about 50% compared to Cas9-ready control mice in which neither Cas9 nor guide RNA was introduced. For AAV8-mediated delivery, however, delivery of the sgRNA to Cas9-ready mice resulted in a several-fold decrease in target protein 3 serum levels compared to WT mice in which both Cas9 and the sgRNA were introduced, and an even more dramatic decrease compared to control WT mice in which neither Cas9 nor the guide RNA were introduced. By day 21, serum levels of target protein 3 had dropped to near the limit of detection in Cas9-ready mice in which the sgRNA was introduced via AAV8.

[00282] **Figure 7** shows percent NHEJ activity (indel frequency) at the target gene 3 locus in liver in wild type mice and cassette-deleted Cas9 mice (MAID2600) one month after lipid nanoparticle (LNP) delivery of sgRNA alone or together with Cas9 mRNA, hydrodynamic delivery (HDD) of sgRNA plasmid alone or together with Cas9 plasmid, or AAV8-sgRNA alone or together with AAV8-Cas9. The percentage of liver cells with insertions/deletions (indels) at the locus was measured by NGS. Hydrodynamic delivery of the guide RNA or the combination of Cas9 and the guide RNA resulted in a low percentage of indels. For LNP-mediated delivery, introduction of the sgRNA into Cas9-ready mice resulted in a similar percentage of indels compared to WT mice in which both Cas9 and the sgRNA were introduced, and each of these conditions resulted in a percentage level of indels that was about 40%. For AAV8-mediated delivery, however, delivery of the sgRNA to Cas9-ready mice resulted in a much larger percentage of indels (~75%) compared to WT mice in which both Cas9 and the sgRNA were introduced (~35%), and an even more dramatic increase compared to control WT mice in which neither Cas9 nor the guide RNA were introduced.

[00283] Further next-generation sequencing (NGS) is also performed in harvested tissues. Amplicon sequencing is then used to assess the amount of editing in harvested tissues. Target specific primers are designed to produce a ~300 bp product that is slightly off center around the expected cut site of the guide. The primers then have “adapter” sequences added to them that will allow the individual samples to be barcoded in a secondary PCR reaction. Once the barcodes are added, the samples are all pooled together and loaded onto the MiSeq. Five thousand to ten thousand reads are expected over the region of interest. Informatic programs are then run to map the reads to determine the precise sequence of each edit. The program then counts the number of WT (unedited) reads and provides a breakdown of the type of edit done to

all edited reads (assessment of the number of base pairs added and/or deleted in the predicted region of editing).

[00284] An experiment to knock out target gene 4, which encodes a type II membrane-bound glycoprotein, was then performed by introducing Cas9 with a sgRNA targeting exon 2 of target gene 4 into primary hepatocytes isolated from wild type (WT) mice. Five different sgRNAs (guides 1-5) were tested. Cas9/sgRNA ribonucleoprotein (RNP) complexes were introduced into the cells via lipofectamine. Non-homologous end joining was then assessed by next-generation sequencing (NGS) to measure indel frequencies at the target gene 4 locus. Percent editing is a measure of total NHEJ events over total reads. NHEJ events are considered to be all edits (insertion, deletion, base change) that occur in the 20 bp before and after the cut site. The percent editing for guides 1 to 5 were as follows: 35.4%, 37.4%, 43.8%, 51.2%, and 55.8%, respectively (data not shown).

[00285] An experiment to knock out target gene 4 *in vivo* was then performed by introducing the same five sgRNAs into separate cassette-deleted Cas9 mice (MAID2600) via AAV8. Specifically, individual guides expressed by a U6 promoter were packaged in AAV8 and introduced into 6-12-week old cassette-deleted Cas9 mice by tail vein injection. The viral load introduced was between 1×10^{11} and 1×10^{12} , in an approximate volume of 50-100 μ L. Livers were harvested 3-4 weeks post-injection. Percent editing was calculated as it was in the primary hepatocytes and is shown in **Figure 8A**. Editing levels were consistent with, and in fact higher than, the editing levels observed in primary hepatocytes. Expression levels of mRNA transcribed from target gene 4 were also tested. As shown in **Figure 8B**, each gRNA reduced the relative levels of mRNA transcribed from target gene 4 in livers harvested 3-4 weeks post-injection.

[00286] Experiments to test percent editing in several other target genes in the liver were also performed. In each experiment, the age of the mice was about 6-12 weeks. For each target gene, five different guide RNAs were designed against critical exons. The guide RNAs were delivered via AAV8 by tail vein injection with viral loads between 1×10^{11} and 1×10^{12} in an approximate volume of 50-100 μ L. Livers were harvested 3-4 weeks post-injection. Percent editing was determined as explained above. The percent editing in the liver of cassette-deleted Cas9 mice (MAID2600) through delivery of AAV8-gRNA is shown in **Table 12**. The best gRNA for each gene resulted in 48%-70% editing in the liver *in vivo*.

[00287] Table 12. Percent Editing in Liver.

Target Gene	Guide#1	Guide#2	Guide#3	Guide#4	Guide#5
5	49.4%	37.1%	43.3%	21.3%	35.7%
6	25.6%	68.9%	44.8%	63.3%	42.1%
7	43.5%	36.1%	30.0%	48.2%	41.4%
8	24.5%	35.2%	66.1%	56.3%	45.5%
9	27.8%	32.7%	47.4%	65.0%	38.9%
4	52.3%	58.8%	63.6%	57.0%	61.5%

Example 2. Validation of Inducibility of Cas9-Ready Mice

[00288] The LSL-Cas9 allele described in Example 1 (MAID2599) includes a floxed strong poly(A) region (lox-stop-lox, or LSL) upstream of the Cas9 coding sequence. Prior to removal of the cassette by the action of Cre recombinase, the neomycin resistance gene will normally be efficiently transcribed and translated; however, the Cas9 CDS will not normally be expressed due to the presence of the strong poly(A) region, which can effectively block run-through transcription. Upon removal of the neo cassette by the action of Cre recombinase, the hybrid mRNA for the Cas9 and GFP proteins will normally be constitutively expressed by the *Rosa26* promoter. This makes the Cas9 allele inducible. This is beneficial for a number of reasons. The possibility of editing some genes in certain tissues (e.g., immune cells) may be detrimental, along with potentially causing an immune response. In addition, in certain circumstances, mutation of a gene throughout the targeted individual may be lethal, whereas mutation of the gene in a specific tissue or cell type would be beneficial. The inducible nature of the MAID2599 allele allows more specificity as to which tissue and cell type are being edited by only activating Cas9 in a tissue-specific or cell-specific manner.

[00289] To test the inducibility of Cas9 expression in the liver *in vivo*, lipid nanoparticles (LNPs) containing Cre recombinase mRNA were formulated on the Precision Nanosystems Benchtop NanoAssemblr. Cre mRNA from Trilink (cat#7211) was diluted in 10 mM sodium citrate and was combined through the NanoAssemblr cassette at 3:1 with the lipid combination of a cationic lipid, DSPC, cholesterol, and PEG-DMG at a molar ratio of 50:10:38.5:1.5. This formulation is readily absorbed by the liver. The resulting LNP-Cre was injected through the tail vein of LSL-Cas9 mice (MAID2599) at 1 mg/kg. In control mice, LNP-Cre was not injected. After 1 week, the mice were sacrificed, and organs were harvested for western analysis using anti-Cas9 (7A9) monoclonal antibody (Invitrogen Cat#MA5-23519) and anti-Actin (C4) monoclonal antibody (Millipore Sigma Cat#MAB1501). Organs from cassette-deleted Cas9

mice (MAID2600) were used as a positive control. In these mice, the LSL cassette had already been removed by Cre recombinase. The results are shown in **Figure 9**, which shows proof-of-concept for liver-specific Cas9 activation with LNP-Cre delivery for liver-specific gene editing.

[00290] The inducibility of Cas9-mediated gene editing was then tested *in vivo*. LNP-Cre was formulated as described above. Mice were dosed with LNP-Cre and AAV8-gRNA targeting target gene 3 (coinjection via tail vein injection) in the following groups: (1) 3 LSL-Cas9 mice treated with LNP-Cre and AAV8-gRNA; (2) 3 LSL-Cas9 mice treated with LNP-Cre and PBS; (3) 3 LSL-Cas9 mice treated with PBS and AAV8-gRNA; (4) 3 LSL-Cas9 mice treated with PBS alone; (5) 3 cassette-deleted Cas9 mice treated with LNP-Cre; (6) 3 cassette-deleted Cas9 mice treated with AAV8-gRNA; (7) 3 cassette-deleted Cas9 mice treated with PBS; and (8) 3 WT mice (untreated). In groups in which LNP-Cre was delivered, it was delivered at a concentration of 1 mg/kg. In groups in which AAV8-gRNA was delivered, it was delivered at a viral load of approximately 2×10^{11} . One and three weeks post-injection, mice were bled for serum chemistry and to measure circulating serum levels of target protein 3. At three weeks, tissues were also harvested for NGS and for western analysis. Serum levels of target protein 3 were measured by ELISA. The results are shown in **Figure 10**. Delivery of LNP-Cre to LSL-Cas9 mice together with AAV8-gRNA resulted in a decrease in serum levels of target protein 3 consistent with the decrease observed in cassette-deleted Cas9 mice in which AAV8-gRNA was delivered. These ELISA results were consistent with the NGS results for percent editing in target gene 3 in livers isolated from the mice 3 weeks post-injection. See **Figure 11**.

[00291] Next, LSL-Cas9 mice were crossed with albumin-Cre mice from Jax (3601-Tg(Alb-cre)21Mgn; MAID3601) in which the albumin promoter is operably linked to the Cre recombinase coding sequence and drives its expression in the liver. Following the cross, several tissues were harvested from the mice for western blot analysis. Corresponding tissues were harvested from LSL-Cas9 mice that were not crossed with the albumin-Cre mice. Western blots measuring Cas9 expression in the liver and brain were then performed. Actin was used as a loading control. The predicted size of Cas9 was 150.48 kD, and the predicted size of actin was 41.25 kD. 17.5 μ g of liver protein lysates and brain protein lysates were used. TruCut v2 Cas9 (17.5 μ g) was used as a positive control. As shown in **Figure 12A**, Cas9 expression was observed in the livers of LSL-Cas9/Alb-Cre mice but not in the livers of LSL-Cas9 mice. Cas9

expression was not observed in the brain tissues from any of the mice (*see Figure 12B*), confirming that Cas9 expression was induced specifically in the liver.

[00292] An experiment was performed to test Cas9-mediated gene editing *in vivo* in these mice. The experiment included five groups of 8-12-week old mice injected with AAV8-gRNA targeting target gene 3 via tail vein injection: (1) 3 mice LSL-Cas9:Alb-Cre (MAID2599 Het, MAID3601 Het); (2) 3 mice WT:Alb-Cre (MAID2599 WT, MAID3601 Het); (3) 3 mice LSL-Cas9:WT (MAID2599 Het, MAID3601 WT); (4) 3 Mice Cas9 (MAID2600 Het or Hom); and (5) 3 Mice 75/25 (50500 WT). Mice from each group also served as controls that were not injected with AAV8-gRNA. In groups in which AAV8-gRNA was delivered, it was delivered at a viral load of approximately 2×10^{11} . One week post-injection, the mice were bled for serum chemistry and ELISAs. Three weeks post-injection, the mice were bled and tissues were harvested. The results are shown in **Figure 13**. Crossing the LSL-Cas9 mice with the albumin-Cre mice and then injecting AAV8-gRNA resulted in a decrease in serum levels of target protein 3 consistent with the decrease observed in cassette-deleted Cas9 mice in which AAV8-gRNA was delivered.

[00293] The Cas9-ready mouse system described herein is able to induce more robust gene editing than other methods relying on exogenous introduction of Cas9. Further, the system can conditionally express Cas9 based on the deletion of a neomycin cassette. By combining this system with various Cre deleter mouse lines, the timing of Cas9-induced genome editing can be controlled and tissue-specific Cas9 expression can be provided *in vivo*.

We claim:

1. A method of testing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*, comprising:
 - (a) introducing into a non-human animal a guide RNA designed to target a guide RNA target sequence at the target genomic locus,
wherein the non-human animal comprises a genomically integrated Cas expression cassette comprising an NLS-Cas coding sequence, and
wherein the guide RNA is introduced via adeno-associated virus (AAV)-mediated delivery; and
 - (b) assessing the modification of the target genomic locus.
2. The method of claim 1, wherein the AAV is AAV7, AAV8, or AAV9, and step (b) comprises assessing modification of the target genomic locus in the liver.
3. The method of claim 2, wherein the AAV is AAV8.
4. The method of any preceding claim, wherein the route of administration of the AAV to the non-human animal is intravenous injection, intraparenchymal injection, intraperitoneal injection, nasal installation, or intravitreal injection.
5. The method of any preceding claim, wherein an exogenous donor nucleic acid is introduced in step (a), wherein the exogenous donor nucleic acid is designed to recombine with the target genomic locus.
6. The method of claim 5, wherein the exogenous donor nucleic acid is a single-stranded oligodeoxynucleotide (ssODN).
7. The method of any preceding claim, wherein the non-human animal is a rat or mouse.
8. The method of claim 7, wherein the non-human animal is a mouse.
9. The method of any preceding claim, wherein the target genomic locus comprises a target gene, and step (b) comprises measuring expression of the target gene or activity of a protein encoded by the target gene.

10. The method of any preceding claim, wherein step (b) comprises sequencing the target genomic locus in one or more cells isolated from the non-human animal.

11. The method of any preceding claim, wherein step (b) comprises isolating a target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the target organ or tissue.

12. The method of claim 11, wherein step (b) comprises assessing modification of the target genomic locus in two or more different cell types within the target organ or tissue.

13. The method of any preceding claim, wherein step (b) comprises isolating a non-target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the non-target organ or tissue.

14. The method of any preceding claim, wherein the NLS-Cas coding sequence is an NLS-Cas9 coding sequence.

15. The method of any preceding claim, wherein the Cas expression cassette further comprises a polyadenylation signal upstream of the NLS-Cas coding sequence, wherein the polyadenylation signal is flanked by recombinase recognition sites, and wherein the polyadenylation signal in the Cas expression cassette has been excised in a tissue-specific manner.

16. The method of claim 15, wherein the polyadenylation signal upstream of the NLS-Cas coding sequence has been excised in the liver.

17. The method of claim 15 or 16, wherein the recombinase that recognizes the recombinase recognition sites in the Cas expression cassette is a Cre recombinase.

18. The method of claim 17, wherein the non-human animal further comprises a genomically integrated Cre recombinase expression cassette comprising a Cre recombinase coding sequence operably linked to a tissue-specific promoter.

19. The method of claim 17 or 18, wherein the Cre recombinase gene is operably linked to one of the promoters set forth in Table 2.

20. The method of any one of claims 1-14, wherein the Cas expression cassette further comprises a polyadenylation signal upstream of the NLS-Cas coding sequence, wherein the polyadenylation signal is flanked by recombinase recognition sites, and wherein the method further comprises introducing a recombinase into the non-human animal in a tissue-specific manner.

21. The method of claim 20, wherein the recombinase is introduced via adeno-associated virus (AAV)-mediated delivery or lipid nanoparticle (LNP)-mediated delivery.

22. The method of any preceding claim, wherein the Cas expression cassette further comprises a fluorescent protein coding sequence.

23. The method of claim 22, wherein the Cas expression cassette comprises a multicistronic nucleic acid comprising the NLS-Cas coding sequence and the fluorescent protein coding sequence separated by an intervening internal ribosome entry site (IRES) or an intervening 2A peptide coding sequence.

24. The method of claim 23, wherein the multicistronic nucleic acid in the Cas expression cassette comprises the NLS-Cas coding sequence and a green fluorescent protein coding sequence separated by an intervening P2A peptide coding sequence.

25. The method of any one of claims 1-21, wherein the Cas expression cassette further does not comprise a fluorescent protein coding sequence.

26. The method of any preceding claim, wherein the NLS-Cas coding sequence encodes a Cas protein comprising a protein tag.

27. The method of any preceding claim, wherein the 5' end of the Cas expression cassette further comprises a 3' splicing sequence.

28. The method of any preceding claim, wherein the Cas expression cassette is operably linked to an endogenous promoter.

29. The method of any one of claims 1-27, wherein the Cas expression cassette is operably linked to an exogenous, constitutive promoter.

30. The method of any preceding claim, wherein the Cas expression cassette encodes a protein comprising the sequence set forth in SEQ ID NO: 13, 16, 19, or 22.

31. The method of claim 30, wherein the Cas expression cassette comprises the sequence set forth in SEQ ID NO: 28, 29, 30, or 31.

32. The method of claim 31, wherein the Cas expression cassette comprises the sequence set forth in SEQ ID NO: 1, 12, 14, 15, 17, 18, 20, or 21.

33. The method of any preceding claim, wherein the Cas expression cassette is integrated at a safe harbor locus.

34. The method of claim 33, wherein the safe harbor locus is a *Rosa26* locus.

35. The method of claim 34, wherein the Cas expression cassette is integrated into the first intron of the *Rosa26* locus.

36. The method of any preceding claim, wherein the non-human animal is heterozygous for the Cas expression cassette.

37. The method of any one of claims 1-35, wherein the non-human animal is homozygous for the Cas expression cassette.

38. The method of any preceding claim, wherein the non-human animal is a mouse,

wherein the AAV is an AAV8 delivered to the non-human animal by intravenous injection,

wherein the Cas expression cassette is operably linked to the endogenous *Rosa26* promoter, is inserted into the first intron of the *Rosa26* locus, and comprises from 5' to 3': (i) a 3' splicing sequence; and (ii) an NLS-Cas9 coding sequence, and

wherein step (b) comprises assessing modification of the target genomic locus in the liver of the non-human animal.

39. A method of optimizing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*, comprising:

(I) performing the method of any preceding claim a first time in a first non-human animal;

(II) changing a variable and performing the method of step (I) a second time with the changed variable in a second non-human animal; and

(III) comparing the modification of the target genomic locus in step (I) with the modification of the target genomic locus in step (II), and selecting the method resulting in the modification of the target genomic locus with one or more of higher efficacy, higher precision, higher consistency, or higher specificity.

40. The method of claim 39, wherein the changed variable in step (II) is the AAV serotype.

41. The method of claim 39, wherein the changed variable in step (II) is the route of administration of introducing the guide RNA into the non-human animal.

42. The method of claim 39, wherein the changed variable in step (II) is the concentration or amount of the guide RNA introduced into the non-human animal.

43. The method of claim 39, wherein the changed variable in step (II) is the guide RNA introduced into the non-human animal.

44. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and wherein the changed variable in step (II) is the delivery method of introducing the exogenous donor nucleic acid into the non-human animal.

45. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the route of administration of introducing the exogenous donor nucleic acid into the non-human animal.

46. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the concentration or amount of the exogenous donor nucleic acid introduced into the non-human animal.

47. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the concentration or amount of the guide RNA introduced into the non-human animal relative to the concentration or amount of exogenous donor nucleic acid introduced into the non-human animal.

48. The method of claim 39, wherein the changed variable in step (II) is the exogenous donor nucleic acid introduced into the non-human animal.

AMENDED CLAIMS

received by the International Bureau on 14 January 2019 (14.01.2019)

We claim:

1. A method of testing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*, comprising:
 - (a) introducing into a non-human animal a guide RNA designed to target a guide RNA target sequence at the target genomic locus,
wherein the non-human animal comprises a genomically integrated Cas expression cassette comprising a coding sequence for a Cas protein further comprising one or more nuclear localization signals, and
wherein the guide RNA is introduced via adeno-associated virus (AAV)-mediated delivery; and
 - (b) assessing the modification of the target genomic locus.
2. The method of claim 1, wherein the AAV is AAV7, AAV8, or AAV9, and step (b) comprises assessing modification of the target genomic locus in the liver.
3. The method of claim 2, wherein the AAV is AAV8.
4. The method of any preceding claim, wherein the route of administration of the AAV to the non-human animal is intravenous injection, intraparenchymal injection, intraperitoneal injection, nasal installation, or intravitreal injection.
5. The method of any preceding claim, wherein an exogenous donor nucleic acid is introduced in step (a), wherein the exogenous donor nucleic acid is designed to recombine with the target genomic locus.
6. The method of claim 5, wherein the exogenous donor nucleic acid is a single-stranded oligodeoxynucleotide (ssODN).
7. The method of any preceding claim, wherein the non-human animal is a rat or mouse.
8. The method of claim 7, wherein the non-human animal is a mouse.

9. The method of any preceding claim, wherein the target genomic locus comprises a target gene, and step (b) comprises measuring expression of the target gene or activity of a protein encoded by the target gene.

10. The method of any preceding claim, wherein step (b) comprises sequencing the target genomic locus in one or more cells isolated from the non-human animal.

11. The method of any preceding claim, wherein step (b) comprises isolating a target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the target organ or tissue.

12. The method of claim 11, wherein step (b) comprises assessing modification of the target genomic locus in two or more different cell types within the target organ or tissue.

13. The method of any preceding claim, wherein step (b) comprises isolating a non-target organ or tissue from the non-human animal and assessing modification of the target genomic locus in the non-target organ or tissue.

14. The method of any preceding claim, wherein the Cas protein is a Cas9 protein.

15. The method of any preceding claim, wherein the Cas expression cassette further comprises a polyadenylation signal upstream of the coding sequence for the Cas protein , wherein the polyadenylation signal is flanked by recombinase recognition sites, and wherein the polyadenylation signal in the Cas expression cassette has been excised in a tissue-specific manner.

16. The method of claim 15, wherein the polyadenylation signal upstream of the coding sequence for the Cas protein has been excised in the liver.

17. The method of claim 15 or 16, wherein the recombinase that recognizes the recombinase recognition sites in the Cas expression cassette is a Cre recombinase.

18. The method of claim 17, wherein the non-human animal further comprises a genomically integrated Cre recombinase expression cassette comprising a Cre recombinase coding sequence operably linked to a tissue-specific promoter.

19. The method of claim 17 or 18, wherein the Cre recombinase gene is operably linked to one of the promoters set forth in Table 2.

20. The method of any one of claims 1-14, wherein the Cas expression cassette further comprises a polyadenylation signal upstream of the coding sequence for the Cas protein, wherein the polyadenylation signal is flanked by recombinase recognition sites, and wherein the method further comprises introducing a recombinase into the non-human animal in a tissue-specific manner.

21. The method of claim 20, wherein the recombinase is introduced via adeno-associated virus (AAV)-mediated delivery or lipid nanoparticle (LNP)-mediated delivery.

22. The method of any preceding claim, wherein the Cas expression cassette further comprises a fluorescent protein coding sequence.

23. The method of claim 22, wherein the Cas expression cassette comprises a multicistronic nucleic acid comprising the coding sequence for the Cas protein and the fluorescent protein coding sequence separated by an intervening internal ribosome entry site (IRES) or an intervening 2A peptide coding sequence.

24. The method of claim 23, wherein the multicistronic nucleic acid in the Cas expression cassette comprises the coding sequence for the Cas protein and a green fluorescent protein coding sequence separated by an intervening P2A peptide coding sequence.

25. The method of any one of claims 1-21, wherein the Cas expression cassette further does not comprise a fluorescent protein coding sequence.

26. The method of any preceding claim, wherein the Cas protein comprises a protein tag.

27. The method of any preceding claim, wherein the 5' end of the Cas expression cassette further comprises a 3' splicing sequence.
28. The method of any preceding claim, wherein the Cas expression cassette is operably linked to an endogenous promoter.
29. The method of any one of claims 1-27, wherein the Cas expression cassette is operably linked to an exogenous, constitutive promoter.
30. The method of any preceding claim, wherein the Cas expression cassette encodes a protein comprising the sequence set forth in SEQ ID NO: 13, 16, 19, or 22.
31. The method of claim 30, wherein the Cas expression cassette comprises the sequence set forth in SEQ ID NO: 28, 29, 30, or 31.
32. The method of claim 31, wherein the Cas expression cassette comprises the sequence set forth in SEQ ID NO: 1, 12, 14, 15, 17, 18, 20, or 21.
33. The method of any preceding claim, wherein the Cas expression cassette is integrated at a safe harbor locus.
34. The method of claim 33, wherein the safe harbor locus is a *Rosa26* locus.
35. The method of claim 34, wherein the Cas expression cassette is integrated into the first intron of the *Rosa26* locus.
36. The method of any preceding claim, wherein the non-human animal is heterozygous for the Cas expression cassette.
37. The method of any one of claims 1-35, wherein the non-human animal is homozygous for the Cas expression cassette.
38. The method of any preceding claim, wherein the non-human animal is a mouse,
wherein the AAV is an AAV8 delivered to the non-human animal by intravenous injection,

wherein the Cas expression cassette is operably linked to the endogenous *Rosa26* promoter, is inserted into the first intron of the *Rosa26* locus, and comprises from 5' to 3': (i) a 3' splicing sequence; and (ii) a coding sequence for a Cas9 protein further comprising one or more nuclear localization signals, and

wherein step (b) comprises assessing modification of the target genomic locus in the liver of the non-human animal.

39. A method of optimizing the ability of a CRISPR/Cas nuclease to modify a target genomic locus *in vivo*, comprising:

(I) performing the method of any preceding claim a first time in a first non-human animal;

(II) changing a variable and performing the method of step (I) a second time with the changed variable in a second non-human animal; and

(III) comparing the modification of the target genomic locus in step (I) with the modification of the target genomic locus in step (II), and selecting the method resulting in the modification of the target genomic locus with one or more of higher efficacy, higher precision, higher consistency, or higher specificity.

40. The method of claim 39, wherein the changed variable in step (II) is the AAV serotype.

41. The method of claim 39, wherein the changed variable in step (II) is the route of administration of introducing the guide RNA into the non-human animal.

42. The method of claim 39, wherein the changed variable in step (II) is the concentration or amount of the guide RNA introduced into the non-human animal.

43. The method of claim 39, wherein the changed variable in step (II) is the guide RNA introduced into the non-human animal.

44. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and wherein the changed variable in step (II) is the delivery method of introducing the exogenous donor nucleic acid into the non-human animal.

45. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the route of administration of introducing the exogenous donor nucleic acid into the non-human animal.

46. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the concentration or amount of the exogenous donor nucleic acid introduced into the non-human animal.

47. The method of claim 39, wherein the method comprises introducing an exogenous donor nucleic acid, and the changed variable in step (II) is the concentration or amount of the guide RNA introduced into the non-human animal relative to the concentration or amount of exogenous donor nucleic acid introduced into the non-human animal.

48. The method of claim 39, wherein the changed variable in step (II) is the exogenous donor nucleic acid introduced into the non-human animal.

49. A non-human animal comprising a genomically integrated Cas expression cassette comprising a coding sequence for a Cas protein further comprising one or more nuclear localization signals.

50. A non-human animal cell comprising a genomically integrated Cas expression cassette comprising a coding sequence for a Cas protein further comprising one or more nuclear localization signals.

51. A non-human animal genome comprising a genomically integrated Cas expression cassette comprising a coding sequence for a Cas protein further comprising one or more nuclear localization signals.

52. A targeting vector comprising a Cas expression cassette flanked by homology arms, wherein the Cas expression cassette comprises a coding sequence for a Cas protein further comprising one or more nuclear localization signals, and wherein the homology arms are suitable for directing recombination with a desired target genomic locus to facilitate genomic integration.

53. A method for making the non-human animal of claim 49, comprising:

- (a) modifying the genome of a non-human animal pluripotent cell to comprise a Cas expression cassette comprising a coding sequence for a Cas protein further comprising one or more nuclear localization signals;
- (b) identifying or selecting the genetically modified pluripotent cell comprising the Cas expression cassette;
- (c) introducing the genetically modified pluripotent cell into a non-human animal host embryo; and
- (d) implanting and gestating the non-human animal host embryo in a surrogate mother.

54. A method for making the non-human animal of claim 49, comprising:

- (a) modifying the genome of a non-human animal one-cell stage embryo to comprise a Cas expression cassette comprising a coding sequence for a Cas protein further comprising one or more nuclear localization signals;
- (b) selecting the genetically modified one-cell stage embryo comprising the Cas expression cassette; and
- (c) implanting and gestating the genetically modified one-cell stage embryo in a surrogate mother.




FIG. 1






FIG. 2A

FIG. 2B




WT RosaCas9^{+/-}
FIG. 3A




WT RosaCas9^{+/-}
FIG. 3D




WT RosaCas9^{+/-}
FIG. 3B



WT RosaCas9^{+/-}
FIG. 3E



WT RosaCas9^{+/-}
FIG. 3C



WT RosaCas9^{+/-}
FIG. 3F




FIG. 4A





FIG. 4B



5/15




FIG. 5B








FIG. 5A






FIG. 7




FIG. 8B




FIG. 8A

9/15




FIG. 9




FIG. 10





FIG. 11

12/15






FIG. 12B




FIG. 13




FIG. 14

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2018/044615

A. CLASSIFICATION OF SUBJECT MATTER
INV. A01K67/027 C12N15/90
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A01K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PLATT RANDALL J ET AL: "CRISPR-Cas9 Knockin Mice for Genome Editing and Cancer Modeling", CELL, CELL PRESS, AMSTERDAM, NL, vol. 159, no. 2, 25 September 2014 (2014-09-25), pages 440-455, XP029073412, ISSN: 0092-8674, DOI: 10.1016/J.CELL.2014.09.014	1,2,4, 7-12,15, 17, 19-26, 29-37, 39-41, 43-45,48
Y	page 442, paragraph 2nd - last paragraph page 444, paragraph 3rd - page 451, paragraph 3rd; figures 1a, 3,4,7 -/--	1-48



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 November 2018

Date of mailing of the international search report

22/11/2018

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Chambonnet, F

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/044615

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	& RANDALL J. PLATT ET AL: "CRISPR-Cas9 Knockin Mice for Genome Editing and Cancer Modeling : Supplemental information", CELL, vol. 159, no. 2, 25 September 2014 (2014-09-25), XP055523070, AMSTERDAM, NL ISSN: 0092-8674, DOI: 10.1016/j.cell.2014.09.014 & Randall Platt: "CRISPR-Cas9 Knockin Mice for Genome Editing and Cancer Modeling : Supplementary material. Supplement 7:Figure S2", 25 September 2014 (2014-09-25), XP055523071, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-7.pdf [retrieved on 2018-11-12] & Randall Platt: "Supplementary material. Supplement 8:Figure S3", 25 September 2014 (2014-09-25), XP055523072, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-8.pdf [retrieved on 2018-11-12] & Randall J Platt: "Supplementary Material :Supplement 10", 25 September 2014 (2014-09-25), XP055523075, England DOI: 10.1038/nrc3950 Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-10.pdf [retrieved on 2018-11-12] & Randall Platt: "Supplementary Material :Supplement 4", 25 September 2014 (2014-09-25), XP055523080, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-4.xlsx [retrieved on 2018-11-12] -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/044615

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	& Randall Platt: "Supplementary Material :Supplement 3 Table S2", 25 September 2014 (2014-09-25), XP055523078, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-3.docx [retrieved on 2018-11-12] & Randall Platt: "Supplementary Material :Supplement 12", 25 September 2014 (2014-09-25), XP055523077, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-12.pdf [retrieved on 2018-11-12] & Randall Platt: "Supplementary material : supplement 11", 25 September 2014 (2014-09-25), XP055523076, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-11.pdf [retrieved on 2018-11-12] & Randall Platt: "Supplementary Material :Supplement 9", 25 September 2014 (2014-09-25), XP055523074, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-9.pdf [retrieved on 2018-11-12] & Randall Platt: "Supplementary material :Supplement 6 Figure S1", 25 September 2014 (2014-09-25), XP055523263, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-6.pdf [retrieved on 2018-11-13] -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/044615

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	& Randall Platt: "Table S1 - Electrophysiological characterization Parameter Wild-type Cas9 Rheobase current", 25 September 2014 (2014-09-25), pages 381-7283, XP055523259, Retrieved from the Internet: URL:https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265475/bin/NIHMS631163-supplement-2.docx [retrieved on 2018-11-13] -----	
X	US 2015/020223 A1 (ZHANG FENG [US] ET AL) 15 January 2015 (2015-01-15)	1-48
Y	paragraphs [0[5]], [[17]], [[22]], [[23]], [2[165]] - [[170]], [[203]], [[207]], [[210]] - paragraphs [[526]] - [[528]], [[538]], [[542]], [[549]], [[558]], [[561]], [[733]]; claims 42-51,53-61; figures 2B-D,4A,25A, 25B,26-30, 70; examples 14, 18, 19, 27,30, 31 -----	1-48
X	WO 2016/049024 A2 (BROAD INST INC [US]; MASSACHUSETTS INST TECHNOLOGY [US]) 31 March 2016 (2016-03-31)	1,2,7, 14,15, 17-26, 29,33-37
Y	paragraphs [0[9]], [[17]] - [[18]], [[54]] - [[56]], [[68]] - [[69]], [[72]], [[74]] - paragraphs [[103]] - [[106]], [[109]] - [[116]], [[141]] - [[232]]; claims 1-4,5-6,18-30; figures 1A-D, 2A-H, 5A-B, 7; examples 1-2 -----	1-48
X	WO 2016/049163 A2 (BROAD INST INC [US]; MASSACHUSETTS INST TECHNOLOGY [US]) 31 March 2016 (2016-03-31)	1,5, 7-24,26, 29,33-37
Y	paragraphs [0027] - [0038], [0139] - [0165]; claims 1-48,51-58, 74-76; examples 1,2,4 -----	1,5, 7-24,26, 29,33-37
Y	CHIANN-MUN CHEN ET AL: "A Comparison of Exogenous Promoter Activity at the ROSA26 Locus Using a PhiC31 Integrase Mediated Cassette Exchange Approach in Mouse ES Cells", PLOS ONE, vol. 6, no. 8, 11 August 2011 (2011-08-11) , page e23376, XP055523971, DOI: 10.1371/journal.pone.0023376 the whole document ----- -/--	1,28

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2018/044615

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ALBERTO CEBRIAN-SERRANO ET AL: "Maternal Supply of Cas9 to Zygotes Facilitates the Efficient Generation of Site-Specific Mutant Mouse Models", PLOS ONE, vol. 12, no. 1, 12 January 2017 (2017-01-12), page e0169887, XP055523935, DOI: 10.1371/journal.pone.0169887 figure 1; table 2 -----	1,28,29, 34-37
A	VAN TRUNG CHU ET AL: "Efficient CRISPR-mediated mutagenesis in primary immune cells using CrispRGold and a C57BL/6 Cas9 transgenic mouse line", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 113, no. 44, 11 October 2016 (2016-10-11), pages 12514-12519, XP055436418, US ISSN: 0027-8424, DOI: 10.1073/pnas.1613884113 page 12515, right-hand column, last paragraph - page 12516, left-hand column, last line; figure 1E -----	1,23,28, 33-37
A	TAKAYUKI SAKURAI ET AL: "A non-inheritable maternal Cas9-based multiple-gene editing system in mice", SCIENTIFIC REPORTS, vol. 6, no. 1, 28 January 2016 (2016-01-28), XP055523965, DOI: 10.1038/srep20011 page 2; figure 1 -----	1,22,29
A	ALLISON LANGE ET AL: "Classical Nuclear Localization Signals: Definition, Function, and Interaction with Importin [alpha]", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 282, no. 8, 23 February 2007 (2007-02-23), pages 5101-5105, XP055522010, US ISSN: 0021-9258, DOI: 10.1074/jbc.R600026200 cited in the application page 5103, paragraph 2nd -----	1,30-32

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2018/044615

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2015020223 A1	15-01-2015	AU 2013359199 A1	30-07-2015
		BR 112015013784 A2	11-07-2017
		CA 2894681 A1	19-06-2014
		CN 105164264 A	16-12-2015
		DK 2931897 T3	05-02-2018
		EP 2931897 A2	21-10-2015
		EP 3327127 A1	30-05-2018
		ES 2658401 T3	09-03-2018
		HK 1215048 A1	12-08-2016
		HK 1220725 A1	12-05-2017
		JP 2016501531 A	21-01-2016
		KR 20150105956 A	18-09-2015
		SG 10201707569Y A	30-10-2017
		SG 11201504523U A	30-07-2015
		US 2014179770 A1	26-06-2014
		US 2014242699 A1	28-08-2014
		US 2015020223 A1	15-01-2015
		US 2017107536 A1	20-04-2017
		US 2017152528 A1	01-06-2017
		WO 2014093622 A2	19-06-2014
WO 2016049024 A2	31-03-2016	US 2018010134 A1	11-01-2018
		WO 2016049024 A2	31-03-2016
WO 2016049163 A2	31-03-2016	US 2018044662 A1	15-02-2018
		WO 2016049163 A2	31-03-2016