



(22) Date de dépôt/Filing Date: 2008/02/07
(41) Mise à la disp. pub./Open to Public Insp.: 2008/08/09
(30) Priorités/Priorities: 2007/02/09 (EP07290164.8);
2007/02/09 (US11/704,313)

(51) Cl.Int./Int.Cl. *A61K 38/17* (2006.01),
A61K 9/70 (2006.01), *A61L 15/32* (2006.01),
A61L 15/44 (2006.01), *A61P 17/02* (2006.01)

(71) Demandeur/Applicant:
GENE SIGNAL INTERNATIONAL SA, CH

(72) Inventeurs/Inventors:
COLIN, SYLVIE, FR;
AL-MAHMOOD, SALMAN, FR

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : COMPOSITION ET AGENT DE TRAITEMENT D'UNE PLAIE
(54) Title: WOUND HEALING AGENT AND COMPOSITION

(57) **Abrégé/Abstract:**

The present invention relates to a wound-treating agent and to a composition for the treatment of wounds comprising a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2.

Abstract

The present invention relates to a wound-treating agent and to a composition for the treatment of wounds comprising a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2.

Wound healing agent and composition

The present invention relates to the healing of wounds. Particularly, the present invention relates to a
5 wound healing agent, compositions containing said wound-healing agent, and medical devices containing the same.

Wound healing in tissues is a complex reparative process. For example, the skin wound healing process involves the recruitment of a variety of specialised cells to the site
10 of the wound, extracellular matrix and basement membrane deposition, angiogenesis, selective protease activity and re-epithelialisation.

There is always a need to provide substances that promote the healing of wounds. It is often desirable to
15 increase the rate of healing in the case of acute wounds and chronic wounds, or for generally healing compromised individuals (for example the elderly). The wounds may severely influence quality of life or even result in death and therefore the rate of healing often needs to be increased
20 as much as is clinically possible.

Number of wound healing agents and compositions already exist on the market, but none are completely satisfying. Though, it is still an important issue to provide
25 alternative wound healing agents and compositions, resulting in effective, rapid and aesthetically acceptable wound repair.

WO 03/074073, in the name of the Applicant, describes a family of 54 genes involved in the regulation of angiogenesis. Amongst these genes, "gene 156" (SEQ ID N°1 in
30 this specification), which encodes "protein 156A" (SEQ ID N°2 in this specification), and also called Angiodensine in WO 03/074073, has been described as pro-angiogenic. Protein 156A comprises 217 amino-acids, and has a mitochondrial sequence signal, detected by *in silico* experiments. Protein 156A does

not contain any transmembrane domain nor conserved domain. WO 03/074073 describes that the expression of an antisense of the gene 156, i.e. the inhibition of gene 156, in human endothelial cells inhibits the formation of capillary tubes.
5 WO 03/074073 further foresees a potential pro-angiogenesis activity for gene 156 and protein 156a.

Going deeper in their researches, the inventors surprisingly found that protein 156A showed a strong *in vitro* and *in vivo* wound healing activity. WO 03/074073 mentioned
10 only potential *in vitro* pro-angiogenic activity for protein 156A, and the inventors had thus no idea about the possible behaviour of protein 156a on wound healing *in vivo*.

The inventors were thus really surprised to note the strong efficiency of protein 156A on wound repair.

15 In addition, inventors found that protein 156A was also particularly active on the final aesthetic aspect of the scar, which appeared more regular and less colored.

The present invention thus relates, in a first aspect, to a wound-treating agent comprising a polypeptide
20 having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2.

By the term "polypeptide having at least 50%,
25 preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2" it is meant any polypeptide having 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2 and presenting a wound healing activity. It also includes fragments of protein 156A
30 having a wound healing activity.

The term "wound" as used herein refers to, but is not limited to:

- injuries to epidermis and/or dermis of the skin,

- wounds resulting from damage, injury or trauma to an internal or external tissue or organ such as for example the eye, mucous, lung, kidney, heart, gut, tendons or liver,

- injuries or damages to vascular tissues, such as
5 for example veins, venules, arteries, and capillaries.

In this specification, the term "wound treating" is used to describe all the different steps involved in the healing of wounds. It therefore includes the steps of forming a clot that plugs the defect, invasion of the clot by
10 inflammatory cells and then of fibroblasts and capillaries to form a contractile granulation tissue that draws the wound margins together, and migration forward of the cut epidermal edges to cover the denuded wound surface. The term "Wound treating" is not limited to the healing of the skin, but also
15 includes the tissue repair of other types of wounds, as listed above.

In a second aspect, the present invention also relates to a method of treatment of a wound comprising administering to a subject in need thereof, a therapeutically
20 effective amount of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2.

By the term "therapeutically effective amount" it is meant an amount that allows the achievement of the
25 contemplated medical end, i.e. the healing of a wound, without producing unacceptable toxic symptoms. Said "therapeutically effective amount" will vary with the factors such as the particular condition being treated, the physical condition of the patients and the duration of the treatment.

30 The present invention further relates, in a third aspect, to a wound-treating composition comprising a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ

ID NO:2 in association with any suitable excipient for the treatment of wounds.

In a particular embodiment, the endotoxins are eliminated from the composition containing the polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2.

One of the principal advantages of the wound treating agent and of the wound treating composition according to the invention is the increase of the healing rate and the return of the injured tissue histologically very close to the native tissue.

Moreover, inventors pointed out that the wound treating agent and the wound treating composition according to the invention are particularly active on the stimulation of endothelial cell migration *in vitro*. Protein 156A also seems to play a global role on the healing of wounds, not limited to a single stimulation of angiogenesis but to a wide stimulation of the wound repair process, comprising the regeneration of the cells in the dermis and the epidermis.

The present invention also relates to a wound-treating composition comprising a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2 for use in a method of treatment of wounds of the human or animal body.

In a particular embodiment, the wound-treating compositions as described above, comprise at least two active substances, one of which being a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with

the amino acid sequence of SEQ ID NO:2. In another embodiment, a further active substance is a haemostatic active substance, a growth factor, an anti-infective substance, an analgesic substance, an anti-inflammatory substance, or a combination thereof.

In still another embodiment, the wound treating composition according to the invention is in a form suitable for topical, systemic, oral, subcutaneous, transderm, intramuscular or intra-peritoneal administration.

10 A suitable form for topical administration comprise liquid, ointment, cream, gel, hydrogel, cataplasm, pomade, liniment, milk, lotion, emulsion, spray, aerosol, collyrium, drops, powder.

A suitable form for systemic administration comprises injectable solution and suppository.

A suitable form for oral administration comprises drinkable suspension, syrup, tablets, capsules, pill.

20 According to the invention, in the wound treating composition as described above, said polypeptide is present in an amount from 0.01 to 90% in weight, preferably from 0.1% to 10% in weight, more preferably from 1% to 5% in weight.

In a particular embodiment, the composition of the invention is in a liquid form, and the active polypeptide is present in an amount from 0.01 mg/mL to 5 mg/mL, preferably from 0.1 mg/mL to 1 mg/mL, more preferably about 0.5 mg/mL.

25 A "composition" in the sense of the invention encompasses pharmaceutical, including dermatological, compositions and cosmetic compositions.

30 The term "suitable excipient", include well known raw materials such as animal and plant oils, mineral oils, synthetic oils, ester oils, waxes, linear higher alcohols, fatty acids, surfactants, phospholipids, gelling and/or thickening agents, alcohol, polyols (including glycerine and propylene glycol), fillers such as clay minerals, soft-focus

powders, preservatives, fragrances, pigments and purified water. This term also includes polysaccharides, such as for example mannans, gluco mannans, galactomannans, fucomannans, proteoglycans, glucosaminoglycans, chitins, chitomannans. It
5 is further mentioned by the Applicant that the present invention is not limited to the excipients listed above. The man in the art is able to choose the best excipients suitable to a particular administration form.

In a fourth aspect, the present invention further
10 relates to a wound-treating medical device comprising a wound treating agent or a wound treating composition as described above.

In a preferred embodiment, the medical device is in the form of a dressing, bandage, transdermic medical device,
15 controlled drug release medical device, or a drug-eluting stent.

Suitable dressings within the meaning of the invention are, without any limitation, hydrocolloid dressings, hydrocellular dressings, alginate dressings,
20 hydrogel dressings, chitosan based dressings, cellulose derivatives dressings and any other type of dressing dedicated to wound protection and repair.

By "transdermic medical device" it is meant a device for slow liberation via transdermic process of a substance,
25 such as for example adhesive patch.

By "drug-eluting stent", also called "coated" or "medicated" stent, it is meant a stent that has been coated with the active substance "protein 156A".

The invention also relates to the use of the wound-
30 treating agent, the wound-treating composition, or of the wound-treating medical device as described above, for the treatment of acute wounds and/or chronic wounds.

The invention also relates to the use of a polypeptide having the amino acid sequence of SEQ ID NO:2, or

of a polypeptide having at least 50%, preferably 70%, more preferably 90% identity with the amino acid sequence of SEQ ID NO:2, in the manufacture of a pharmaceutical composition, for the treatment of acute wounds and/or chronic wounds

5 Basically, acute wounds may be classified into different types, according to the object that caused the wound. For example, incisions or incised wounds, lacerations, abrasions and grazes, burns, puncture wounds caused by an object puncturing the skin, such as a nail or a needle,
10 penetration wounds caused by an object such a knife entering the body, gunshot wounds caused by a bullet or similar projectile driving into or through the body. Acute wounds may also be closed wounds, such as contusions or bruises, haematoma, crushing injuries caused by a great or extreme
15 amount of force applied over a long period of time. Other acute wounds are due to dermatologic diseases such as psoriasis, acne and eczema.

 Another type of wounds which may be treated by the invention is chronic wounds. Common chronic wounds are venous
20 ulcers, which usually occur in the legs and mostly affect the elderly, diabetic ulcers which is another major cause of chronic wounds, pressure ulcers, which usually occur in people with conditions such as paralysis that inhibit movement of body parts that are commonly subjected to
25 pressure such as the heels, shoulder blades and sacrum, corneal ulcers, most commonly caused by an infection with bacteria, viruses, fungi or amoebae, and digestive ulcers. Other types of chronic wounds may be due to causes such as ischemia and radiation poisoning.

30

 The present invention will now be further described with reference to the following non-limiting examples.

Fig. 1A and 1B: Pictures of HUVEC endothelial cell lawn after a wound *in vitro* assay. After wounding, cells are incubated with VEGF and FGF2 (1A) or with VEGF, FGF2 and protein 156A (1B).

5 Fig. 2A: Pictures of wounds performed on Nude mice.

Fig. 2B: Pictures of wounds performed on Nude mice 48 hours post the topical application of vehicle solution.

Fig. 2C: Pictures of wounds performed on Nude mice 48 hours post the topical application of protein 156A.

10 Fig. 3A: Pictures of 4 injured mice 96 hours post vehicle application.

Fig. 3B: Pictures of 4 injured mice 96 hours post protein 156A application.

15 Fig. 4: Picture of two trans-dermal parallel wounds performed on the back of a farm pig.

Fig. 5: Picture of the two trans-dermal parallel wounds few hours after application of vehicle solution ("C" for CONTROL) or protein 156A ("T" for TEST).

20 Fig. 6: Picture of the two trans-dermal parallel wounds 3 weeks after application of vehicle solution ("C" for CONTROL) or protein 156A ("T" for TEST).

Fig. 7A: Picture of a section of a wound harvested 5 days after application of a vehicle solution.

25 Fig. 7B: Picture of a section of a wound harvested 5 days after application of protein 156A.

Example 1: Production of the recombinant protein 156A

30 Protein 156A is a recombinant protein corresponding to the entire protein coded by the gene 156. Protein 156A was expressed in *Escherichia coli* B121(DE3)pLys, extracted from bacterial cell with 8M urea and purified onto metal ion (Ni) chelating columns using liquid chromatography. The purity of the purified recombinant protein 156A was controlled by SDS-PAGE and shown to be over 65%. Before use, endotoxins were

eliminated from the solution containing the purified protein 156A using Endo-Trap columns (Pharmacia) and tested. The purified endotoxin-free recombinant protein 156A was conserved in a Tris-HCl buffer solution at pH 7.5 containing
5 2M urea, 150mM NaCl and 0.1 mM CaCl₂.

Example 2: Stimulation of endothelial cell migration by the protein 156A *in vitro*

Cell migration was tested by the wound assay
10 described by Sato and Rifkin (J Cell Biol. 1988;107:1199) with few modifications. HUVEC grown in growth medium EGM-2MV (Cambrex) were seeded in 24-well plates at 80 000 cells per well in 500µL of growth medium and grown to confluence at 37°C in a humidified atmosphere containing 5% CO₂. Cells were
15 scrapped with a plastic tip on one line only. After wounding, the culture medium was changed for fresh medium supplemented with 500nM (15µg.mL⁻¹) of protein 156A (Test, **Fig 1B**) or not (Control, **Fig 1A**). After 18 hours of culture, cells were observed and photographed under the inverted microscope
20 (Analysis, Olympus, Rungis, France).

Figure 1A shows that, 18 hours post incubation, the wound is still present under the control condition (growth medium only).

Figure 1B shows that, 18 hours post incubation, the
25 wound is completely healed under the Test condition (growth medium + protein 156A).

Since migration of endothelial cells is one of the critical features of neovascularisation and wound repair, protein 156A is thus a potential therapeutic lead for wound
30 healing.

Example 3: Pilot experiment of the healing activity of protein 156A in Swiss Nude mice *in vivo*

Two animals were anesthetized by IP injection of Ketamine-Xylazine (80mg/kg - 12 mg/kg; Ref. K-113, Sigma, France) and then trans-dermal injured (1cm in length and about 1 to 2 mm in depth, see **Fig. 2A**) in the right flank of each mouse using 0.5/10 bladder. 200 μ L of either Vehicle solution (Tris-HCl pH 7.5, 2M Urea, 150 mM NaCl, 0.1 mM CaCl₂) or Test solution (Vehicle + 0.5mg.mL⁻¹ Protein 156A) was then topically applied on the wound of animals 10 min post injury.

The examination of wound healing at 48 hours post the topical application showed that the protein 156A treated wound (see **Fig. 2C**) was completely healed while the vehicle treated wound (see **Fig. 2B**) was not yet healed and keep widely opened.

15

Example 4: Test of the healing activity of protein 156A in Swiss Nude mice in vivo

Eight healthy female Swiss *Nude* mice were anesthetized as in example 3. Two trans-dermal wounds, each of 1 cm in length and about 1 to 2 mm in depth were realized in the right flank of each mouse using 0.5/10 bladder.

At D1, and 10 min post-injury of the animals, the 8 mice were randomized into 2 groups of 4 mice. All treatments were realized by topical application. The treatment schedule is summarized in Table 1.

Group	Animals n	Treatment	Treatment dose
1	4	Vehicle	0
2	4	Protein 156A	0.5mg.mL ⁻¹

Table 1

Mice of Group 1 were treated with 200 μ L of vehicle solution (Tris-HCl pH 7.5, 2M Urea, 150 mM NaCl, 0.1 mM

30

CaCl₂) topically applied on each wound, 10 minutes after wounding.

Mice of Group 2 were treated with 200µL of Test solution (Tris-HCl pH 7.5, 2M Urea, 150 mM NaCl, 0.1 mM CaCl₂, 0.5mg.mL⁻¹ Protein 156A) topically applied on each wound, 10 minutes after wounding.

Pictures of the wounds have been taken 96 hours after treatment with the vehicle or the test solution:

Fig.3A: Pictures of the mice of Group 1 96 hours post vehicle application.

Fig.3B: Pictures of the mice of Group 2 96 hours post Protein 156A application.

As illustrated by the pictures of Fig. 3, the examination of wound healing at 96 hours post the topical application showed that the Protein 156A treated wounds was completely healed (Group 2, Fig.3B) while the vehicle treated wounds (Group 1, Fig.3A) not yet healed and keep widely opened.

Example 5: Test of the healing activity of protein 156A in a porcine model in vivo

Skin wound healing activity

A farm pig was anaesthetised by IM injection of Ketamine/Azaperone (10 mg/kg - 2 mg/kg IM) and then two parallel longitudinal trans-dermal wounds all the way down to the hypodermis were performed (**Fig. 4**).

5 mL of either Vehicle solution (Tris-HCl pH 7.5, 2M Urea, 150 mM NaCl, 0.1 mM CaCl₂, CONTROL side "C") or Test solution (Vehicle + 0.5mg.mL⁻¹ Protein 156A, TEST side "T") was then topically applied on the wound of animals 10 min post injury.

Observation of both wounds showed a marked difference in the immediate postoperative period. A scab appeared in the few hours following surgery on the TEST side

"T" (**Fig. 5**). The latter scab stayed in place for two weeks and was then rubbed off. Healing under the scab seemed normal as judged on the macroscopic appearance of the wound. After 3 weeks, the wound was smooth/even and thin (**Fig. 6**). The scab
5 on the CONTROL side "C" developed only after two days. General aspects of the CONTROL wound was less even and ended up thicker and more contracted.

Cornea wound healing activity

10 A farm pig was anaesthetised by IM injection of Ketamine/Azaperone (10 mg/kg, 2 mg/kg IM) and then superficial corneal scarifications were performed.

15 1 mL of either Vehicle solution (Tris-HCl pH 7.5, 2M Urea, 150 mM NaCl, 0.1 mM CaCl₂, CONTROL side) or Test solution (Vehicle + 0.5mg.mL⁻¹ Protein 156A, TEST side) was then topically applied on the scarifications 10 min post injury.

At Day 2 the animal was sacrificed and the tissues were harvested.

20 Oedema of the ciliary processes was observed in both the treated and non treated zones. However, while the non-ulcerated corneal epithelium was detached laterally by intense oedema in the non treated zone, this oedema was virtually absent in the treated zone.

25

Example 6: Histological analysis of treated and non treated cutaneous wounds

30 A farm pig was anaesthetised by IM injection of Ketamine/Azaperone (10 mg/kg, 2 mg/kg IM) and then two parallel longitudinal trans-dermal wounds all the way down to the hypodermis were performed.

5 mL of either Vehicle solution (Tris-HCl pH 7.5, 2M Urea, 150 mM NaCl, 0.1 mM CaCl₂, CONTROL side) or Test

solution (Vehicule + 0.5mg.mL⁻¹ Protein 156A, TEST side) was then topically applied on the wound of animals 10 min post injury.

5 At day 5 the animal was sacrificed and areas of both CONTROL wound and TEST wound were harvested. Formalin-fixed, paraffin-embedded blocks were processed and serial five microns sections from the harvested areas were prepared for conventional HES staining.

10 In the non-treated animal (**Fig. 7A**), results showed that, at Day 5 post injury, the wound was still transfixing as far as the deep subdermis, while in the treated group (**fig. 7B**), the approximation of the edges concern all layers of the skin (dermis and epidermis).

15 These different tests showed that there was a strong difference between the treated wounds and the control wounds. Healing of the Test wounds (treated with protein 156A) was all times faster, thinner and more even and complete. Final aesthetic aspect of the wound was also greater. The molecule
20 further showed strong activity on the experimental model of corneal ulcer.

156A ST25.txt
SEQUENCE LISTING

<110> GENE SIGNAL INTERNATIONAL

<120> Wound healing agent and composition

<130> BEP

<160> 2

<170> PatentIn version 3.1

<210> 1

<211> 2153

<212> DNA

<213> Homo sapiens

<400> 1
 caaagagaca aacactgatg aagccgctaa tgacaatatg gcagttcgcc ccaaagagcg 60
 cagcagcctg agctctagac agcatccggt tgtaaggagc agtgtgatag tgcgctcaca 120
 gaccttttct ccaggagagc ggaaccagta catctgcagg ttaaatacggg gtgacagtga 180
 cagttcaacc ctggctaaaa aatcactggt tgtgagaaac tccaccgaac gccgcagttt 240
 gaggggtcaaa aggacgggtt gccagtcagt ccttagaaga acaacacagg aatgcccagt 300
 gcggacatct ctagacttag aactggacct tcaggcatct ctgacccggc agagccgcct 360
 caatgatgag ctgcaggcgc tgagggactt gcggcagaag ctggaggaac tgaaagctca 420
 gggagagact gaccttccac caggcgtgct ggaggatgag aggttccaga ggcttctgaa 480
 gcaagctgag aagcaggctg aacagtccaa agaagagcag aagcaaggctc tgaatgcaga 540
 gaagttgatg aggcaagtct ccaaggacgt gtgtcggctc cgggagcaga gccagaaggt 600
 gcctcggcag gtgcagtcct tcagggagaa gattgcctac ttcaccagag caaagataag 660
 catcccatcc ctgccagctg atgatgtgtg attacatgac ttaagaaatt attttttcat 720
 ctgttcactt tcttagggag ggtaaaagac tgaagatttg tgtttttggt ttggtgtttg 780
 gttttttttg gtaacgtaac tgtcaactct tgaagaactt ttatttcaca tcagattttc 840
 aacacattaa tttgtaaagt accttgagtg taatttttat atgcttaaaa aagaaaaaat 900
 catataagaa agatgggata acctgggata caaatgttgc ccagtatgtg cgcattgccca 960
 gtggacttca ttttgtaaaa atggaataga ggcagtaccc caccatgtgt actgttgagc 1020

156A ST25.txt

cggctggtga gaaacaactt ggttcagccg gtggttttgc ttctcctttg tgttcctgtg 1080
 agtaggctgt gccgatggca gcagcgccac gtggtatctg tgcgctggcg aggagaggag 1140
 cgctttccat ttggaggcag tgtgagcaca cgctggagtg cagcgttcca aagtccgccc 1200
 gaaatggatg ctgctgccac ggccatcacc tgtgcgggaa cacctggttt ccttaagaca 1260
 atttcctttt tgtcctttta ttttttcta aagaaaacta agatataaac taccaagtgc 1320
 tcttaagaat aaaaataaga ataagaatac aaaggagcac tactcttggc tacacgaaag 1380
 atcttgggat tcatgacact gagggcaggg agaagaaaga acaccagcca cgcagagata 1440
 caaataagct catcaaatga gcagctgggt tgtctcactc agcctgaatt tgcattaagt 1500
 gttcaaaagt gagtttctgg cttgcactag tgttttgccc taaatctaag gatggtgcct 1560
 agtgggtggc ctgtgttctg aggacagaga gagctggagg ggtggaggga aatgcatcgg 1620
 cactcatttg gtgtaaccaa gaagaggggg aaagatgtct tgatagaatc aaagttgcca 1680
 aaaacaaga agagtttatc aagcagattt ctaacattta gtagaaacta cttcatagct 1740
 ctcattttta gggggttcta tctataattt atggtaaata gttgggaggc aaaggaaga 1800
 atgtgtgtgg gggattatt aatttgaagt ataagctaaa ttgtttatta aaaagcctat 1860
 tttggatgaa caagtttaga gagataaact gactaatgtg tcctccagtt tagaatttca 1920
 ccttaccgga cccaggcact tcttaaaaac acagatcctc agtgcagctc tgccagcatc 1980
 aaggctttta aatatgtag tatttaggcc atttaaacca tacttattac tcaataaccc 2040
 aagaaacaat tttgggttag ccatctaata tagcaaataa aatagatttt ttaaagaaat 2100
 ccaaaattgt cttaaaggaa tggaccctt gaagaaaaaa aaaaaaaaaa aaa 2153

<210> 2

<211> 217

<212> PRT

<213> Homo sapiens

<400> 2

Met Ala Val Arg Pro Lys Glu Arg Ser Ser Leu Ser Ser Arg Gln His
 1 5 10 15

Pro Phe Val Arg Ser Ser Val Ile Val Arg Ser Gln Thr Phe Ser Pro
 20 25 30

Gly Glu Arg Asn Gln Tyr Ile Cys Arg Leu Asn Arg Ser Asp Ser Asp
 35 40 45

Ser Ser Thr Leu Ala Lys Lys Ser Leu Phe Val Arg Asn Ser Thr Glu
 50 55 60

156A ST25.txt

Arg Arg Ser Leu Arg Val Lys Arg Thr Val Cys Gln Ser Val Leu Arg
65 70 75 80

Arg Thr Thr Gln Glu Cys Pro Val Arg Thr Ser Leu Asp Leu Glu Leu
85 90 95

Asp Leu Gln Ala Ser Leu Thr Arg Gln Ser Arg Leu Asn Asp Glu Leu
100 105 110

Gln Ala Leu Arg Asp Leu Arg Gln Lys Leu Glu Glu Leu Lys Ala Gln
115 120 125

Gly Glu Thr Asp Leu Pro Pro Gly Val Leu Glu Asp Glu Arg Phe Gln
130 135 140

Arg Leu Leu Lys Gln Ala Glu Lys Gln Ala Glu Gln Ser Lys Glu Glu
145 150 155 160

Gln Lys Gln Gly Leu Asn Ala Glu Lys Leu Met Arg Gln Val Ser Lys
165 170 175

Asp Val Cys Arg Leu Arg Glu Gln Ser Gln Lys Val Pro Arg Gln Val
180 185 190

Gln Ser Phe Arg Glu Lys Ile Ala Tyr Phe Thr Arg Ala Lys Ile Ser
195 200 205

Ile Pro Ser Leu Pro Ala Asp Asp Val
210 215

CLAIMS

1. A wound-treating composition comprising a polypeptide having the amino acid sequence of SEQ ID NO:2, or
5 of a polypeptide having at least 70%, preferably 90% identity with the amino acid sequence of SEQ ID NO:2 in association with any suitable excipient for the treatment of wounds.

2. A wound-treating composition comprising a polypeptide having the amino acid sequence of SEQ ID NO:2, or
10 of a polypeptide having at least 70%, preferably 90% identity with the amino acid sequence of SEQ ID NO:2 for use in a method of treatment of wounds of the human or animal body.

3. A wound-treating composition according to claims 1-2, comprising at least two active substances, one of which
15 being a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 70%, preferably 90% identity with the amino acid sequence of SEQ ID NO:2.

4. A wound-treating composition according to claim 3, wherein a further active substance is an haemostatic active
20 substance, a growth factor, an anti-infective substance, an analgesic substance, an anti-inflammatory substance, or a combination thereof.

5. A wound treating composition according to anyone of claims 1-4, being in a form suitable for topical,
25 systemic, oral, subcutaneous, transderm, intramuscular or intra-peritoneal administration.

6. The wound-treating composition of claim 5, wherein the form suitable for topical administration comprise cream, gel, cataplasm, pomade, liniment, milk, lotion, emulsion,
30 spray, collyrium, drops, powder.

7. The wound-treating composition of claim 5, wherein the form suitable for systemic administration comprises injectable solution and suppository.

8. The wound-treating composition of claim 5, wherein the form suitable for oral administration comprises drinkable suspension, syrup, tablets, capsules, pill.

5 9. A wound treating composition according to claim 1-8, wherein said polypeptide is present in an amount from 0.01 to 90% in weight, preferably from 0.1% to 10% in weight, more preferably from 1% to 5% in weight.

10 10. A wound-treating medical device comprising a composition of claims 1-9.

11. A wound-treating medical device according to claim 10, being in the form of a dressing, bandage, transdermic medical device, controlled drug release medical device, or a drug-eluting stent.

15 12. Use of a polypeptide having the amino acid sequence of SEQ ID NO:2, or of a polypeptide having at least 70%, preferably 90% identity with the amino acid sequence of SEQ ID NO:2, in the manufacture of a pharmaceutical composition, for the treatment of acute wounds and/or chronic wounds.

20 13. The use according to claim 12, wherein the acute wounds comprise incisions, lacerations, abrasions, puncture wounds, penetration wounds, contusions, haematoma, crushing injuries.

25 14. The use according to claim 12, wherein the chronic wounds comprise wounds caused by venous ulcers, diabetic ulcers, pressure ulcers, corneal ulcers, digestive ulcers, ischemia, radiation poisoning, and dermatologic diseases such as psoriasis, acne and eczema.

Application number / numéro de demande: 2620029

Figures: 1A, 1B, 2A, 2B, 2C, 3A, 3B, 4, 5, 6, 7A, 7B

Pages: 1/7 to 7/7

Unscannable item(s)

received with this application

To inquire if you can order a copy of the unscannable items, please visit the
CIPO WebSite at [HTTP://CIPO.GC.CA](http://CIPO.GC.CA)

Item(s) ne pouvant être balayés

Documents reçus avec cette demande ne pouvant être balayés.

Pour vous renseigner si vous pouvez commander une copie des items ne
pouvant être balayés, veuillez visiter le site web de l'OPIC au [HTTP://CIPO.GC.CA](http://CIPO.GC.CA)