(12) STANDARD PATENT (11) Application No. **AU 2018368786 B2**
(19) AUSTRALIAN PATENT OFFICE

(54) Title
TIL expansion from fine needle aspirates and small biopsies

(51) International Patent Classification(s)
C12N 5/0783 (2010.01) ***A61K 39/00*** (2006.01)
A61K 35/17 (2015.01) ***A61P 35/00*** (2006.01)

(21) Application No: **2018368786** (22) Date of Filing: **2018.11.19**

(87) WIPO No: **WO19/100023**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/756,038	**2018.11.05**	**US**
62/621,515	**2018.01.24**	**US**
62/588,044	**2017.11.17**	**US**

(43) Publication Date: **2019.05.23**
(44) Accepted Journal Date: **2025.05.22**

(71) Applicant(s)
Iovance Biotherapeutics, Inc.

(72) Inventor(s)
SIMPSON-ABELSON, Michelle;CHARTIER-COURTAUD, Cecile

(74) Agent / Attorney
Pizzeys Patent and Trade Mark Attorneys Pty Ltd, PO Box 291, WODEN, ACT, 2606, AU

(56) Related Art
DONIA M ET AL: "Characterization and Comparison of 'Standard' and 'Young' Tumour-Infiltrating Lymphocytes for Adoptive Cell Therapy at a Danish Translational Research Institution", SCANDINAVIAN JOURNAL OF IMMUNOLOGY vol. 75, 2012 pg 157-167

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
23 May 2019 (23.05.2019)

WIPO | PCT

(10) International Publication Number
WO 2019/100023 A1

(51) International Patent Classification:
C12N 5/0783 (2010.01) *A61K 39/00* (2006.01)
A61K 35/17 (2015.01) *A61P 35/00* (2006.01)

(21) International Application Number:
PCT/US2018/061865

(22) International Filing Date:
19 November 2018 (19.11.2018)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

62/588,044	17 November 2017 (17.11.2017)	US
62/621,515	24 January 2018 (24.01.2018)	US
62/756,038	05 November 2018 (05.11.2018)	US

(71) Applicant: **IOVANCE BIOTHERAPEUTICS, INC.** [US/US]; 999 Skyway Raod, Suite 150, San Carlos, CA 94070 (US).

(72) Inventors: **SIMPSON-ABELSON, Michelle**; 15908 Oakleaf Run Drive, Lithia, FL 33547 (US). **CHARTIER-COURTAUD, Cecile**; 1357 Cowper Street, Palo Alto, CA 94301 (US).

(74) Agent: **MACDOUGALL, Christina, A.** et al.; Morgan, Lewis & Bockius LLP, One Market, Spear Tower, San Francisco, CA 94105 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:
— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:
— *with international search report (Art. 21(3))*
— *with sequence listing part of description (Rule 5.2(a))*

WO 2019/100023 A1

(54) Title: TIL EXPANSION FROM FINE NEEDLE ASPIRATES AND SMALL BIOPSIES

(57) Abstract: The present disclosure provides methods for expanding TIL populations from fine needle aspirates (FN As) or small biopsies which contain low numbers of TILs, using the methods disclosed herein including in a closed system that leads to improved phenotype and increased metabolic health of the TILs in a shorter time period.

TIL EXPANSION FROM FINE NEEDLE ASPIRATES AND SMALL BIOPSIES

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Patent Application No. 62/588,044, filed on November 17, 2017, U.S. Provisional Patent Application No. 62/621,515, filed on January 24, 2018, and U.S. Provisional Patent Application No. 62/756,038, filed on November 5, 2018, which are hereby incorporated by reference in their entireties.

BACKGROUND OF THE INVENTION

[0002] Treatment of bulky, refractory cancers using adoptive transfer of tumor infiltrating lymphocytes (TILs) represents a powerful approach to therapy for patients with poor prognoses. Gattinoni, *et al.*, *Nat. Rev. Immunol.* **2006,** *6,* 383-393. A large number of TILs are required for successful immunotherapy, and a robust and reliable process is needed for commercialization. This has been a challenge to achieve because of technical, logistical, and regulatory issues with cell expansion. IL-2-based TIL expansion followed by a "rapid expansion process" (REP) has become a preferred method for TIL expansion because of its speed and efficiency. Dudley, *et al.*, *Science* **2002,** *298,* 850-54; Dudley, *et al.*, *J. Clin. Oncol.* **2005,** *23,* 2346-57; Dudley, *et al.*, *J. Clin. Oncol.* **2008**, *26,* 5233-39; Riddell, *et al.*, *Science* **1992,** *257,* 238-41; Dudley, *et al.*, *J. Immunother.* **2003**, *26,* 332-42. REP can result in a 1,000-fold expansion of TILs over a 14-day period, although it requires a large excess (*e.g.*, 200-fold) of irradiated allogeneic peripheral blood mononuclear cells (PBMCs, also known as mononuclear cells (MNCs)), often from multiple donors, as feeder cells, as well as anti-CD3 antibody (OKT3) and high doses of IL-2. Dudley, *et al., J. Immunother.* **2003,** *26,* 332-42. TILs that have undergone an REP procedure have produced successful adoptive cell therapy following host immunosuppression in patients with melanoma. Current infusion acceptance parameters rely on readouts of the composition of TILs (*e.g.*, CD28, CD8, or CD4 positivity) and on fold expansion and viability of the REP product.

[0003] Current TIL manufacturing processes are limited by how the TILs are obtained from the patient. In some cases, where the tumor is sufficiently small or in a location in which tumor resection is not feasible, there remains a need for additional methods of obtaining TILs for expansion and treatment. The present invention meets this need by providing methods for

expanding TILs from a fine needle aspirate (FNA) or a core needle biopsy, which contain low numbers of TILs, and using these expanded TILs in treatment methods.

BRIEF SUMMARY OF THE INVENTION

[0004] The present invention provides improved and/or shortened methods for expanding TILs isolated from a fine needle aspirate or a core biopsy and producing therapeutic populations of TILs.

[0005] The present invention provides a method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(i) obtaining a first population of TILs from at least one fine needle aspirate (FNA) or at least one small biopsy from a tumor in a patient;

(ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is optionally supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for about 3 days to about 10 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 3 days to about 12 days when the first population of TILs is from a small biopsy; and

(iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 3 days to about 12 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs.

[0006] In some embodiments, the cell culture medium comprising IL-2 in step (ii) further comprises OKT-3 and is not optionally supplemented with OKT-3 at any one of days 1-3, and wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion.

[0007] The method according to claim 1 or 2, wherein after step (iii), the cells are removed from the cell culture and cryopreserved in a storage medium prior to performing step (iv).

[0008] In some embodiments, the cells are thawed prior to performing step (iv).

[0009] In some embodiments, step (iv) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

[0010] In some embodiments, steps (i) through (iii) or (iv) are performed within a period of about 17 days to about 24 days.

[0011] In some embodiments, steps (i) through (iii) or (iv) are performed within a period of about 18 days to about 22 days.

[0012] In some embodiments, steps (i) through (iii) or (iv) are performed within a period of about 20 days to about 22 days.

[0013] In some embodiments, steps (i) through (iii) or (iv) are performed within about 22 days.

[0014] In some embodiments, the cells from steps (iii) or (iv) express CD4, CD8, and TCR α β at levels similar to freshly harvested cells.

[0015] In some embodiments, the APCs are peripheral blood mononuclear cells (PBMCs).

[0016] In some embodiments, the PBMCs are added to the cell culture on any of days 3 through 12in step (ii) and/or any of days 11 through 14 in step (iii).

[0017] In some embodiments, the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs in step (iv) exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

[0018] In some embodiments, the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

[0019] In some embodiments, the APCs are artificial APCs (aAPCs) or autologous APCs.

[0020] In some embodiments, the therapeutic population of TILs are infused into a patient.

[0021] In some embodiments, the first expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0022] In some embodiments, the second expansion in step (iii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0023] In some embodiments, the FNA in step (i) comprises at least 400,000 TILs.

[0024] In some embodiments, the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian.

[0025] In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

[0026] In some embodiments, the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the patient has previously undergone surgical treatment.

[0027] In some embodiments, the TILs in step (i) are obtained from a FNA.

[0028] In some embodiments, the FNA is obtained using a 25-18 gauge needle.

[0029] In some embodiments, the TILs in step (i) are obtained from a small biopsy.

[0030] In some embodiments, the small biopsy is obtained using a 16-11 gauge needle.

[0031] In some embodiments, step (iii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

[0032] In some embodiments, the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

[0033] In some embodiments, the present invention provides a method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(i) performing a first expansion by culturing a first population of TILs from a fine needle aspirate (FNA) or a small biopsy from a tumor in a patient in a cell culture medium comprising IL-2 to obtain a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for about 3 days to about 12 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 3 days to about 12 days when the first population of TILs is from a small biopsy; and

(ii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs) to obtain a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 3 days to about 12 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs.

[0034] In some embodiments, the cell culture medium comprising IL-2 in step (ii) further comprises OKT-3 and is not optionally supplemented with OKT-3 at any one of days 1-3, and wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion step.

[0035] In some embodiments, the cells from the cell culture medium in step (ii) are removed and cryopreserved in a storage medium prior to step (iii).

[0036] In some embodiments, the cells are thawed prior to step (iii).

[0037] In some embodiments, the APCs are artificial APCs (aAPCs) or autologous APCs.

[0038] In some embodiments, the therapeutic population of TILs are infused into a patient.

[0039] In some embodiments, the first expansion in step (i) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0040] In some embodiments, the second expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0041] In some embodiments, the additional second expansion in step (iii) is performed by further supplementing the cell culture medium of the third population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0042] In some embodiments, the FNA in step (i) comprises at least 400,000 TILs.

[0043] In some embodiments, the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian.

[0044] In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

[0045] In some embodiments, the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the patient has previously undergone surgical treatment.

[0046] In some embodiments, the TILs in step (i) are obtained from a FNA.

[0047] In some embodiments, the FNA is obtained using a 25-18 gauge needle.

[0048] In some embodiments, the TILs in step (i) are obtained from a small biopsy.

[0049] In some embodiments, the small biopsy is obtained using a 16-11 gauge needle.

[0050] In some embodiments, step (ii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

[0051] In some embodiments, the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

[0052] In some embodiments, the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

[0053] In some embodiments, the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

[0054] The present invention also provides a method for treating a subject with cancer comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising:

(i) obtaining a first population of TILs from a fine needle aspirate (FNA) or a small biopsy obtained from a tumor in a patient;

(ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for about 3 days to about 12 days in order to obtain the second

population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 3 days to about 12 days when the first population of TILs is from a small biopsy;

(iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 3 days to about 12 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and

(iv) administering a therapeutically effective dosage of the third population of TILs to the patient.

[0055] In some embodiments, rein the cell culture medium comprising IL-2 in step (ii) further comprises OKT-3 and is not optionally supplemented with OKT-3 at any one of days 1-3, and wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion step.

[0056] In some embodiments, after step (ii) the cells are removed from the cell culture medium and cryopreserved in a storage medium prior to step (iv).

[0057] In some embodiments, the cells are thawed prior to step (iv).

[0058] In some embodiments, step (iii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

[0059] In some embodiments, the APCs are artificial APCs (aAPCs) or autologous APCs.

[0060] In some embodiments, the first expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0061] In some embodiments, the second expansion in step (iii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0062] In some embodiments, the FNA in step (i) comprises at least 400,000 TILs.

[0063] In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

[0064] In some embodiments, the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

[0065] In some embodiments, the TILs in step (i) are obtained from a FNA.

[0066] In some embodiments, the FNA is obtained using a 25-18 gauge needle.

[0067] In some embodiments, the TILs in step (i) are obtained from a small biopsy.

[0068] In some embodiments, the small biopsy is obtained using a 16-11 gauge needle.

[0069] In some embodiments, step (iii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

[0070] In some embodiments, the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

[0071] The present invention also provides a method according to any of claims 50 to 66, wherein the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

[0072] In some embodiments, the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

[0073] In some embodiments, the cancer is selected from the group consisting of melanoma, cervical cancer, head and neck cancer, glioblastoma, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma.

[0074] The present invention also provides a method for treating a subject with cancer comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising:

(i) performing a first expansion by culturing a first population of TILs from a fine needle

aspirate (FNA) or a small biopsy from a tumor in a patient in a cell culture medium comprising IL-2 to obtain a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for about 3 days to about 12 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 3 days to about 12 days when the first population of TILs is from a small biopsy;

(ii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs) to obtain a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 3 days to about 12 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and

(iii) administering a therapeutically effective dosage of the therapeutic population of TILs to the patient.

[0075] In some embodiments, the cell culture medium comprising IL-2 in step (ii) further comprises OKT-3 and is not optionally supplemented with OKT-3 at any one of days 1-3, and wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion step.

[0076] In some embodiments, the cells from the cell culture medium in step (ii) are removed and cryopreserved in a storage medium prior step (iii).

[0077] In some embodiments, the cells are thawed prior to step (iii).

[0078] In some embodiments, the APCs are artificial APCs (aAPCs) or autologous APCs.

[0079] In some embodiments, the APCs are peripheral blood mononuclear cells (PBMCs).

[0080] In some embodiments, the therapeutic population of TILs are infused into a patient.

[0081] In some embodiments, the first expansion in step (i) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0082] In some embodiments, the second expansion in step (iii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[0083] In some embodiments, the FNA in step (i) comprises at least 400,000 TILs.

[0084] In some embodiments, the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian.

[0085] In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

[0086] In some embodiments, the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

[0087] In some embodiments, the TILs in step (i) are obtained from a FNA.

[0088] In some embodiments, the FNA is obtained using a 25-18 gauge needle.

[0089] In some embodiments, the TILs in step (i) are obtained from a small biopsy.

[0090] In some embodiments, the small biopsy is obtained using a 16-11 gauge needle.

[0091] In some embodiments, step (ii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

[0092] In some embodiments, the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

[0093] In some embodiments, the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

[0094] In some embodiments, the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

[0095] The present invention also provides a method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(a) obtaining a first population of TILs from a fine needle aspirate (FNA) or a small biopsy obtained from a tumor in a patient;

(b) adding the first population into a closed system;

(c) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for about 3 days to about 12 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 3 days to about 12 days when the first population of TILs is from a small biopsy, wherein the first expansion is performed in a closed container providing a first gas-permeable surface area, wherein the first expansion is performed for about 3 days to about 12 days to obtain the second population of TILs, wherein the second population of TILs is at least 25-fold greater in number than the first population of TILs, and wherein the transition from step (b) to step (c) occurs without opening the system;

(d) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the second expansion is performed for about 3 days to about 12 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs, wherein the second expansion is performed in a closed container providing a second gas-permeable surface area, and wherein the transition from step (c) to step (d) occurs without opening the system;

(e) harvesting the therapeutic population of TILs obtained from step (d), wherein the transition from step (d) to step (e) occurs without opening the system; and

(f) transferring the harvested TIL population from step (e) to an infusion bag, wherein the transfer from step (e) to (f) occurs without opening the system.

[0096] In some embodiments, the cell culture medium comprising IL-2 in step (ii) further comprises OKT-3 and is not optionally supplemented with OKT-3 at any one of days 1-3, and wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion step.

[0097] In some embodiments, the method further comprises the step of cryopreserving the infusion bag comprising the harvested TIL population in step (f) using a cryopreservation process.

[0098] In some embodiments, the cryopreservation process is performed using a 1:1 ratio of harvested TIL population to CS10 media.

[0099] In some embodiments, the APCs are peripheral blood mononuclear cells (PBMCs).

[00100] In some embodiments, the PBMCs are irradiated and allogeneic.

[00101] In some embodiments, the PBMCs are added to the cell culture on any of days 3 through 12 in step (c) and/or any of days 3 through 12 in step (d).

[00102] In some embodiments, the antigen-presenting cells are artificial antigen-presenting cells (aAPCs) or autologous APCs.

[00103] In some embodiments, the therapeutic population of TILs are infused into a patient.

[00104] In some embodiments, the first expansion in step (c) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[00105] In some embodiments, the second expansion in step (d) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[00106] In some embodiments, the FNA in step (a) comprises at least 400,000 TILs.

[00107] In some embodiments, the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast and ovarian.

[00108] In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

[00109] In some embodiments, the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

[00110] In some embodiments, the TILs in step (a) are obtained from a FNA.

[00111] In some embodiments, the FNA is obtained using a 25-18 gauge needle.

[00112] In some embodiments, the TILs in step (a) are obtained from a small biopsy.

[00113] In some embodiments, the small biopsy is obtained using a 16-11 gauge needle.

[00114] In some embodiments, the harvesting in step (e) is performing using a LOVO cell processing system.

[00115] In some embodiments, the cell culture medium is provided in a container selected from the group consisting of a G-container and a Xuri cellbag.

[00116] In some embodiments, the infusion bag in step (f) is a hypothermosol infusion bag.

[00117] In some embodiments, steps (a) through (f) are performed within a period of about 17 days to about 24 days.

[00118] In some embodiments, steps (a) through (f) are performed within a period of about 18 days to about 22 days.

[00119] In some embodiments, steps (a) through (f) are performed within a period of about 20 days to about 22 days.

[00120] In some embodiments, steps (a) through (f) are performed in 22 days or less.

[00121] In some embodiments, steps (a) through (f) and cryopreservation are performed in 22 days or less.

[00122] In some embodiments, the therapeutic population of TILs harvested in step (e) comprises sufficient TILs for a therapeutically effective dosage of the TILs.

[00123] In some embodiments, the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

[00124] In some embodiments, steps (b) through (e) are performed in a single container, wherein performing steps (b) through (e) in a single container results in an increase in TIL yield per resected tumor as compared to performing steps (b) through (e) in more than one container.

[00125] In some embodiments, the antigen-presenting cells are added to the TILs during the second period in step (d) without opening the system.

[00126] In some embodiments, the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit one or more characteristics selected from the group consisting of expressing CD27+, expressing CD28+, longer telomeres, increased CD57 expression, and decreased CD56 expression relative to effector T cells, and/or central memory T cells obtained from the second population of cells.

[00127] In some embodiments, the effector T cells and/or central memory T cells obtained from the third population of TILs exhibit increased CD57 expression and decreased CD56 expression relative to effector T cells and/or central memory T cells obtained from the second population of cells.

[00128] In some embodiments, the risk of microbial contamination is reduced as compared to an open system.

[00129] In some embodiments, the TILs from step (g) are infused into a patient.

[00130] The present invention also provides a method for treating a subject with cancer, the method comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising:

(a) obtaining a first population of TILs from a fine needle aspirate (FNA) or a small biopsy from a tumor resected from a subject;

(b) adding the first population into a closed system;

(c) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for about 3 days to about 12 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 3 days to about 12 days when the first population of TILs is from a small biopsy, wherein the first expansion is performed in a closed container providing a first gas-permeable surface area, wherein the first expansion is performed for about 3-14 days to obtain the second population of TILs, wherein the second population of TILs is at least 25-fold greater in number than the first population of TILs, and wherein the transition from step (b) to step (c) occurs without opening the system;

(d) performing a second expansion by supplementing the cell culture medium of the

second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the second expansion is performed for about 3 days to about 12 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs, wherein the second expansion is performed in a closed container providing a second gas-permeable surface area, and wherein the transition from step (c) to step (d) occurs without opening the system;

(e) harvesting the therapeutic population of TILs obtained from step (d), wherein the transition from step (d) to step (e) occurs without opening the system; and

(f) transferring the harvested TIL population from step (e) to an infusion bag, wherein the transfer from step (e) to (f) occurs without opening the system;

(g) optionally cryopreserving the infusion bag comprising the harvested TIL population from step (f) using a cryopreservation process; and

(h) administering a therapeutically effective dosage of the third population of TILs from the infusion bag in step (g) to the patient.

[00131] In some embodiments, the therapeutic population of TILs harvested in step (e) comprises sufficient TILs for administering a therapeutically effective dosage of the TILs in step (h).

[00132] In some embodiments, the cell culture medium comprising IL-2 in step (ii) further comprises OKT-3 and is not optionally supplemented with OKT-3 at any one of days 1-3, and wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion step.

[00133] In some embodiments, the APCs are artificial APCs (aAPCs) or autologous APCs.

[00134] In some embodiments, the therapeutic population of TILs are infused into a patient.

[00135] In some embodiments, the first expansion in step (c) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[00136] In some embodiments, the second expansion in step (d) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

[00137] In some embodiments, the FNA in step (a) comprises at least 400,000 TILs.

[00138] In some embodiments, the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast and ovarian.

[00139] In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

[00140] In some embodiments, the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

[00141] In some embodiments, the TILs in step (i) are obtained from a FNA.

[00142] In some embodiments, the FNA is obtained using a 25-18 gauge needle.

[00143] In some embodiments, the TILs in step (i) are obtained from a small biopsy.

[00144] In some embodiments, the small biopsy is obtained using a 16-11 gauge needle.

[00145] In some embodiments, the number of TILs sufficient for administering a therapeutically effective dosage in step (h) is from about 2.3×10^{10} to about 13.7×10^{10}.

[00146] In some embodiments, the antigen presenting cells (APCs) are PBMCs.

[00147] In some embodiments, the PBMCs are added to the cell culture on any of days 3 through 12 in step (c) and/or any of days 3 through 12 in step (d).

[00148] In some embodiments, prior to administering a therapeutically effective dosage of TIL cells in step (h), a non-myeloablative lymphodepletion regimen has been administered to the patient.

[00149] In some embodiments, the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

[00150] In some embodiments, the method further comprises the step of treating the patient with a high-dose IL-2 regimen starting on the day after administration of the TIL cells to the patient in step (h).

[00151] In some embodiments, the high-dose IL-2 regimen comprises 600,000 or 720,000 IU/kg administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

[00152] In some embodiments, the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit one or more characteristics selected from the group consisting of expressing CD27+, expressing CD28+, longer telomeres, increased CD57 expression, and decreased CD56 expression relative to effector T cells, and/or central memory T cells obtained from the second population of cells.

[00153] In some embodiments, the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit increased CD57 expression and decreased CD56 expression relative to effector T cells and/or central memory T cells obtained from the second population of cells.

[00154] The present invention also provides a method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(a) obtaining a first population of TILs from a fine needle aspirate (FNA), a small biopsy, a core biopsy, or a small biopsy from a tumor in a patient;

(b) performing a priming first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2, OKT-3, and antigen presenting cells (APCs) to produce a second population of TILs, wherein the priming first expansion is performed for a first period of about 1 to 14 days in a container comprising a first gas-permeable surface area to obtain the second population of TILs, wherein the second population of TILs is greater in number than the first population of TILs;

(c) performing a rapid second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and APCs, to produce a third population of TILs, wherein the rapid second expansion is performed for a second period of about 1 to about 14 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and

(d) harvesting the therapeutic population of TILs from step (c).

[00155] In some embodiments, the first period is about 6 to 12 days.

[00156] In some embodiments, the second period is about 6 to 12 days.

[00157] In some embodiments, the first period is selected from the group consisting of 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, or 12 days.

[00158] In some embodiments, the second period is selected from the group consisting of 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, or 12 days.

[00159] In some embodiments, the APCs are peripheral blood mononuclear cells (PBMCs).

[00160] In some embodiments, the ratio of APCs used in Step (b) to the APCs used in Step (c) is about 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, 2:1, 2.1:1, 2.2:1, 2.3:1, 2.4:1, 2.5:1, 2.6:1, 2.7:1, 2.8:1, 2.9:1, 3:1, 3.1:1, 3.2:1, 3.3:1, 3.4:1, 3.5:1, 3.6:1, 3.7:1, 3.8:1, 3.9:1, 4:1, 4.1:1, 4.2:1, 4.3:1, 4.4:1, 4.5:1, 4.6:1, 4.7:1, 4.8:1, 4.9:1, or 5:1, or preferably about 1 to 2.

[00161] In some embodiments, the first population of TILs is obtained from a core biopsy.

[00162] In some embodiments, the core biopsy is obtained from a tumor selected from the group consisting of a melanoma tumor, an ovarian cancer tumor, a cervical cancer tumor, a non-small-cell lung cancer (NSCLC) tumor, a lung cancer tumor, a bladder cancer tumor, a breast cancer tumor, a tumor from a cancer caused by human papilloma virus, a head and neck cancer (including head and neck squamous cell carcinoma (HNSCC)) tumor, a glioblastoma tumor, a gastrointestinal cancer tumor, and a renal cancer tumor.

[00163] The present invention also provides compositions comprising TILs expanded using the methods as described herein.

[00164] In some embodiments, the composition further comprises a cyropreservant.

[00165] In some embodiments, the cryopreservant comprises dimethylsulfoxide.

[00166] In some embodiments, the composition further comprises a cyropreservant and an isotonic agent.

[00167] In some embodiments, the composition further comprises a cyropreservant comprising dimethylsulfoxide and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate.

[00168] In some embodiments, the composition further comprises a cyropreservant comprising dimethylsulfoxide and dextran 40 and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate.

[00169] In some emboidments, the composition comprising the TILs is provided in a sterile infusion bag.

BRIEF DESCRIPTION OF THE DRAWINGS

[00170] Figure 1: Exemplary Process 2A chart providing an overview of Steps A through F.

[00171] Figure 2: Process Flow Chart of Process 2A.

[00172] Figure 3: Shows a diagram of an embodiment of a cryopreserved TIL exemplary manufacturing process (~22 days).

[00173] Figure 4: Shows a diagram of an embodiment of process 2A, a 22-day process for TIL manufacturing.

[00174] Figure 5: Comparison table of Steps A through F from exemplary embodiments of process 1C and process 2A.

[00175] Figure 6: Detailed comparison of an embodiment of process 1C and an embodiment of process 2A.

[00176] Figure 7: Exemplary Small Biopsy chart providing an overview of Steps A through F of the Small Biopsy Expansion process.

[00177] Figure 8: Image of viable cells expanded from core biopsies taken from a patient with pancreatic cancer (P7015). A) Core biopsies treated with IL-2, and B) core biopsies treated with a combination of IL-2, IL-15, and IL-21.

[00178] Figure 9: FACS analysis of TILs expanded from pancreatic core biopsies. A) T cells expanded from core biopsies treated with IL-2 are CD4+ and CD8+. B) CD8+ T cells expanded from core biopsies treated with IL-2 are CD107a+. C) T cells expanded from core biopsies treated with a combination of IL-2, IL-15, and IL-21 are CD4+ and CD8+. D) CD8+ T cells expanded from core biopsies treated with a combination of IL-2, IL-15, and IL-21 are CD107a+.

[00179] Figure 10: FACS analysis of TILs expanded from fine needle aspirates from a patient with cervical carcinoma (C2005). A) Fine needle aspirates treated with IL-2, and B) fine needle aspirates treated with a combination of IL-2, IL-15, and IL-21.

[00180] Figure 11: FACS analysis of TILs expanded from fine needle aspirates from a patient with lung carcinoma (L4032). A) Fine needle aspirates treated with IL-2, and B) fine needle aspirates treated with a combination of IL-2, IL-15, and IL-21. C) As a control, lung tumor fragments were cultured with IL-2 and expanded to produce CD4+ and CD8+ T cells.

[00181] Figure 12: TILs from the FNA sample from a patient with lung carcinoma (L4032) were further expanding using a Rapid Expansion Protocol. A) Total cell counts after rapid expansion of TILs expanded from aspirates that were initially treated with either IL-2 alone, a combination of IL-2, IL-15, and IL-21, or initially a combination of IL-2, IL-15, and IL-21 and then IL-2 alone. B) CD4+ and CD8+ T cells were present in the expanded TIL population treated with Il-2.

[00182] Figure 13: FACS analysis of TILs expanded from fine needle aspirates from a patient with lung carcinoma (L4033). A) T cells from fine needle aspirates treated with IL-2, and B) as a control, T cells from lung tumor fragments cultured with IL-2.

[00183] Figure 14: TILs expanded from fine needle aspirates from a patient with lung carcinoma (L4033) were further expanded using a Rapid Expansion Protocol. A) Total cell counts after rapid expansion of TILs obtained from fine needle aspirates treated with IL-2.

[00184] Figure 15: TILs from fine needle aspirates from ovarian tumors taken from patients with ovarian carcinoma (OV8011, OV8012, and OV8013). Total cell counts after culturing the fine needle aspirates with IL-2 or a combination of IL-2, IL-15, and IL-21.

[00185] Figure 16: FACS analysis of TILs expanded from fine needle aspirates from a patient with melanoma (M1101). A) T cells from fine needle aspirates treated with IL-2, and B) as a control, T cells from lung tumor fragments cultured with IL-2.

[00186] Figure 17: Phenotypic and functional status of TILs derived from pancreatic tumor sample obtained by core biopsies. A) Results from pancreatic tumors cultured in culture medium with IL-2 or culture medium with IL-2 and cell culture additives. B and C) Results of CD107a expression in CD4+ cells and CD8+ cells in different culture mediums (IL-2 supplemented culture medium versus additive supplemented culture medium) after stimulation with phorbol 12-myristate 13-acetate (PMA).

[00187] Figure 18: Summary table of results from core biopsies and fine needle aspirates. A) Results from pancreatic cancer and lung cancer samples. B) Results from colorectal cancer, ovarian cancer, cervical cancer, and melanoma samples.

[00188] Figure 19: Phenotypic and functional status of TILs derived from melanoma samples obtained by fine needle aspirate biopsies. A) Total cell counts of cells cultures after 0 days, 11 days, 18 days, and 21 days. B) Phenotypic analysis of cells derived from fragments and aspirates. B) Functional analysis (CD107a expression) of cells derived from fragments and aspirates after stimulation with PMA.

[00189] Figure 20: Summary of data showing TIL can be expanded from the Pancreatic Cancer Core Biopsies obtained from UPMC (University of Pittsburgh Medical Center).

[00190] Figure 21: Data showing general phenotyping of TIL derived from pancreatic tumors.

[00191] Figure 22: TIL isolated from the pre-REP and REP of Pancreatic Cores are Functional as determined by IFNγ and CD107a.

[00192] Figure 23: An increase in CD3+/NK cell population was observed in low expanded Pre-REP lung TIL.

[00193] Figure 24: CD4/8 ratio is higher than 1 in majority of Pre-REP lung TIL.

[00194] Figure 25: Comparable fold expansion during REP was observed in both high and low yield.

[00195] Figure 26: Data regarding Sarcoma (UPMC 58).

[00196] Figure 27: Data regarding Cervical cancer (UPMC45). Pre-REP TIL were expanded from 4 fragments in G-REX 10 for 21 days.

[00197] Figure 28: Data regarding ES19001 (UPMC46). Pre-REP TIL were expanded from 4 fragments in G-REX 10 for 21 days.

[00198] Figure 29: Data regarding Hard Palate (UPMC 67) and Oral Cancer (UPMC 68). Pre-REP TIL were expanded from 4 fragments in G-REX 10 for 21 days.

[00199] Figure 30: Summary of data showing TIL can be expanded from the Pancreatic Cancer Core Biopsies provided by UPMC.

[00200] Figure 31: Pancreatic Core Biopsies: TIL can be expanded from Core Biopsies from Pancreatic Cancer. P7015 Experimental Summary and Results: One core was placed

into one well of a 24 well plate for a total of 5 wells. Two cores were placed into two wells, however there was little to no growth in these wells. Three wells were treated with 1) IL-2 and two wells with 2) IL-2/IL-15/IL-21.

[00201] Figure 32: Data showing that the majority of cells expanded from pancreatic cores were CD4 and CD8+ T cells, and were functional as determined by CD107a expression (P7015). P7015 Experimental Results: The majority of cells expanded from the pancreatic core biopsies are T cells (approximately 80% were CD4+ and CD8+). Stimulation with PMA/I demonstrated the ability of the CD8+ to degranulate as indicated by expression of CD107a+, thereby suggesting that these cells are functional. The % of CD8+ expressing CD107a+ was higher in the IL-2/IL-15/IL-21 triple cocktail condition, compared to IL-2 alone.

[00202] Figure 33: Pancreatic Core Biopsies: TIL can be expanded from Core Biopsies from Pancreatic Cancer. P7028 and P7031 pre-REP Experimental Summary and Results: 11 core biopsies were place into a G-Rex 10 with the triple cocktail (IL-2/IL-15/IL-21). P7028 TIL were assessed at Day 12 and kept in culture until Day 19. Cell counts: Day 12= 1.61e6 viable cells; Day 21 3.49e7 viable cells. P7028 TIL were assessed at Day 12 and kept in culture until Day 27. Cell counts: Day 12= 1.52e5 viable cells; Day 27: 1.14e7 viable cells.

[00203] Figure 34: Pancreatic Core Biopsies: TIL can be expanded from Core Biopsies and fragments treated with IL-2/IL-15/IL-21 from Pancreatic Cancer Day of harvest for TIL derived from fragments ranged from Day 11-21. Day of harvest for TIL derived from cores ranged from Day 21-28.

[00204] Figure 35: Phenotypic Characterization of TIL derived from cores and tumors biopsies. Day of harvest for TIL derived from fragments ranged from Day 11-21. Day of harvest for TIL derived from cores ranged from Day 21-28.

[00205] Figure 36: Phenotypic Characterization of TIL derived from cores and tumors biopsies.

[00206] Figure 37: TIL from core biopsies can express CD107a and secrete IFNγ similarly to TIL isolated from fragments.

[00207] Figure 38: A) Exemplary process for core biopsy expansion procedure. 1-2 core biopsies (3 pancreases, 4 melanoma, 1 breast and 1 ovarian) were subjected to an exemplary process that included the addition of OKT3 + feeders at Day 3. Pancreatic cores were expanded +/- IL-15 and IL-21 (REP1 & REP2; *i.e.*, first expansion and second expansion).

Ovarian cores were treated +/- OX40 (BMS) (n=1). All TIL were subjected to a second REP (second expansion). B) Exemplary process for core biopsy expansion procedure. 1-2 core biopsies are subjected to a process that includes the addition of OKT3 + feeders at Day 3

[00208] Figure 39: Summary table of pancreatic core biopsy expansion procedure.

[00209] Figure 40: Data regarding cell number for pancreatic core biopsy expansions.

[00210] Figure 41: Summary table of melanoma core biopsy expansion procedure.

[00211] Figure 42: Data regarding cell number for melanoma core biopsy expansions.

[00212] Figure 43: Phenotypic expression of the exhaustion and activation markers are similar between REP1 and REP2 (*i.e.*, first expansion and second expansion).

[00213] Figure 44: The addition of OKT3 + feeders to cores at Day 3 does not significantly alter the phenotype of TIL compared to Gen2 (*i.e.*, process 2A) derived TIL from fragments.

[00214] Figure 45: Summary table of breast and ovarian core biopsy expansion procedure.

[00215] Figure 46: Data regarding cell number for breast (left) and ovarian (right panel) core biopsy expansions.

[00216] Figure 47: Experiment showing that TILs can be expanded from Core Biopsies from Pancreatic Cancer. P7028 and P7031 pre-REP Experimental Summary and Results. Core biopsies were place into a G-Rex 10 with the triple cocktail (IL-2/IL-15/IL-21). P7028 TIL were assessed at Day 12 and kept in culture until Day 19. Cell counts: Day 12= 1.61×10^6 viable cells; Day 21 3.49×10^7 viable cells. P7030 TIL were assessed at Day 12 and kept in culture until Day 27. Cell counts: Day 12= 1.52×10^5 viable cells; Day 27: 1.14×10^7 viable cells.

[00217] Figure 48: TILs can be expanded from Core Biopsies and fragments treated with IL-2/IL-15/IL-21 from Pancreatic Cancer. Graphs showing cell expansion numbers for fragments and cores. Day of harvest for TIL derived from fragments ranged from Day 11-21. Day of harvest for TIL derived from cores ranged from Day 21-28.

[00218] Figure 49: TIL derived from cores and fragments are phenotypically similar. Day of harvest for TIL derived from fragments ranged from Day 11-21. Day of harvest for TIL derived from cores ranged from Day 21-28.

[00219] Figure 50: The "exhaustion markers" are differentially expressed when comparing TIL from cores and fragments.

[00220] Figure 51: TIL from core biopsies can express CD107a and secrete IFNγ similarly to TIL isolated from fragments.

[00221] Figure 52: The TC (triple cocktail; IL-2/IL-15/IL-21) does not significantly alter the total cell count in pancreatic cores in REP2 (second expansion), regardless of the initial culture conditions (REP1; first expansion).

[00222] Figure 53: The TC (triple cocktail; IL-2/IL-15/IL-21) does not significantly alter the CD4/CD8 ratio or the Memory subsets in REP2 (second expansion) in pancreatic cores.

[00223] Figure 54: CD107a is similar in the CD4 and CD8 populations from REP2 (second expansion) in all treatment conditions in pancreatic cores.

[00224] Figure 55: IFNγ secretion is similar regardless of REP2 (second expansion) treatment condition in pancreatic cores.

[00225] Figure 56: The addition of OKT3 & feeders at Day 3 (REP1; first expansion) does not significantly alter phenotypic expression, in comparison to pre-REP conditions in pancreatic cores.

[00226] Figure 57: The addition of OKT3 & feeders at Day 3 (REP1; first expansion) does not significantly alter phenotypic expression of the "exhaustion markers" in pancreatic cores.

[00227] Figure 58: The addition of OKT3 & feeders at Day 3 (REP1; first expansion) does not significantly alter phenotypic expression of the activation markers in pancreatic cores.

[00228] Figure 59: CD107a is similar in the CD4 and CD8 populations in the pre-REP, REP1 (first expansion) and REP2 (second expansion) in pancreatic cores.

[00229] Figure 60: IFNγ secretion is reduced in REP2 (second expansion), compared to REP1 (first expansion) in pancreatic cores.

[00230] Figure 61: TCM (central memory T-cells) population is increased and the TEM (effector memory T-cells) decreased in REP2 (second expansion), but CD27 and CD28 are unaltered in melanoma. TEMRA: terminally differentiated effector memory cells re-expressing CD45RA. TSCM: Stem memory T cells.

[00231] Figure 62: IFNγ and CD107a are similar in melanoma core REP1 (first expansion) and REP2 (second expansion).

[00232] Figure 63: Data showing CD4 and CD8 phenotypic markers in melanoma fragments versus melanoma cores.

[00233] Figure 64: Data showing CD4 and CD8 phenotypic markers in melanoma fragments versus melanoma cores.

[00234] Figure 65: Data showing CD107a level in CD4 and CD8 in melanoma fragments versus melanoma cores.

[00235] Figure 66: Treatment with OX40 enhanced the percentage of CD8+ cells in the ovarian cores.

[00236] Figure 67: Similar CD107a expression and IFNγ secretion in ovarian cores treated with OX40.

[00237] Figure 68: Enhanced TCM and differential expression of exhaustion and activation markers in REP1 (first expansion) and REP2 (second expansion) breast cores.

[00238] Figure 69: CD107a and IFNγ is similar in REP1 (first expansion) and REP2 (second expansion) of breast cores.

[00239] Figure 70: Schematic of an exemplary tumor core biopsy TIL expansion procedure.

[00240] Figure 71: TIL derived from pancreatic cores treated with OKT3 + feeders secrete IFNγ with restimulation.

[00241] Figure 72: TIL derived from melanoma cores express CD107a and secrete IFNγ upon restimulation.

[00242] Figure 73: Expanding TIL from Breast and Ovarian Core Biopsies.

[00243] Figure 74: The final product phenotype of TIL derived from breast cores treated with OKT3+feeders during REP1 were similar to cores treated with IL-2 alone during REP1.

[00244] Figure 75: The final product phenotype of TIL derived from breast cores treated with OKT3+feeders during REP1 were similar to cores treated with IL-2 alone during REP1.

[00245] Figure 76: Enhanced IFNγ secretion expression in breast cores treated with OKT3+feeders during REP1 compared to cores treated with IL-2 alone during REP1.

[00246] Figure 77: Treatment with OX40 enhanced the percentage of CD8+ cells in the ovarian cores in REP2.

[00247] Figure 78: Treatment with OX40 during both REPs altered the expression of PD-1, KLRG1 and CD25 in REP2.

[00248] Figure 79: IFNγ secretion in ovarian cores treated with OX40 is slightly higher in the final product (REP2).

[00249] Figure 80: CD107a expression in ovarian cores treated with OX40.

[00250] Figure 81: Treatment with OX40 through 2 REPs altered the expression of markers associated with T cell "youth".

[00251] Figure 82: Treatment with OX40 during both REPs slightly altered the expression altered the expression of PD-1, KLRG1 and CD25 in REP2.

[00252] Figure 83: Differential expression of exhaustion and activation markers in REP1 and REP 2 breast cores.

[00253] Figure 84: CD107a and IFNγ is similar in REP1 and REP2 of breast cores.

[00254] Figure 85A-85B: A) Shows a comparison between the 2A process (approximately 22-day process) and an embodiment of the Small Biopsy Gen 3 process for TIL manufacturing (approximately 17-days to 24-days process). **B)** Exemplary Process Gen3 chart providing an overview of Steps A through F (approximately 17-days to 24-days process).

[00255] Figure 86: Provides an experimental flow chart for comparability between GEN 2 (process 2A) versus GEN 3.

[00256] Figure 84A-87C: A) L4054 - Phenotypic characterization on TIL product on Gen 2 and Gen 3 process. **B)** L4055-Phenotypic characterization on TIL product on Gen 2 and Gen 3 process. **C)** M1085T-Phenotypic characterization on TIL product on Gen 2 and Gen 3 process.

[00257] Figure 88A-88C: **A)** L4054 – Memory markers analysis on TIL product from the Gen 2 and Gen 3 processes. **B)** L4055 – Memory markers analysis on TIL product from the Gen 2 and Gen 3 processes. **C)** M1085T- Memory markers analysis on TIL product from the Gen 2 and Gen 3 processes.

[00258] Figure 89: L4054 Activation and exhaustion markers **(A)** Gated on CD4+, **(B)** Gated on CD8+.

[00259] Figure 90: L4055 Activation and exhaustion markers **(A)** Gated on CD4+, **(B)** Gated on CD8+.

[00260] Figure 91: IFNγ production (pg/mL): (A) L4054, (B) L4055, and (C) M1085T for the Gen 2 and Gen 3 processes: Each bar represented here is mean + SEM for IFNγ levels of stimulated, unstimulated, and media control. Optical density measured at 450 nm.

[00261] Figure 92: ELISA analysis of IL-2 concentration in cell culture supernatant: **(A) L4054** and **(B) L4055**. Each bar represented here is mean + SEM for IL-2 levels on spent media. Optical density measured at 450 nm.

[00262] Figure 93: Quantification of glucose and lactate (g/L) in spent media: (A) Glucose and (B) Lactate: In the two tumor lines, and in both processes, a decrease in glucose was observed throughout the REP expansion. Conversely, as expected, an increase in lactate was observed. Both the decrease in glucose and the increase in lactate were comparable between the Gen 2 and Gen 3 processes.

[00263] Figure 94: A) Quantification of L-glutamine in spent media for L4054 and L4055. **B)** Quantification of Glutamax in spent media for L4054 and L4055. C) Quantification of ammonia in spent media for L4054 and L4055.

[00264] Figure 95: Telomere length analysis: The relative telomere length (RTL) value indicates that the average telomere fluorescence per chromosome/genome in Gen 2 and Gen 3 process of the telomere fluorescence per chromosome/genome in the control cells line (1301 Leukemia cell line) using DAKO kit.

[00265] Figure 96: Unique CDR3 sequence analysis for TIL final product on L4054 and L4055 under Gen 2 and Gen 3 process. Columns show the number of unique TCR B clonotypes identified from 1×10^6 cells collected on Harvest Day Gen 2 (*e.g.*, day 22) and Gen 3 process (*e.g.*, day 14-16). Gen 3 shows higher clonal diversity compared to Gen 2 based on the number of unique peptide CDRs within the sample.

[00266] Figure 97: Frequency of unique CDR3 sequences on L4054 IL harvested final cell product (Gen 2 (*e.g.*, day 22) and Gen 3 process (*e.g.*, day 14-16)).

[00267] Figure 98: Frequency of unique CDR3 sequences on L4055 TIL harvested final cell product (Gen 2 (*e.g.*, day 22) and Gen 3 process (*e.g.*, day 14-16)).

[00268] Figure 99: Diversity Index for TIL final product on L4054 and L4055 under Gen 2 and Gen 3 process. Shanon entropy diversity index is a more reliable and common metric for comparison. Gen 3 L4054 and L4055 showed a slightly higher diversity than Gen 2.

[00269] Figure 100: Raw data for cell counts Day 7-Gen 3 REP initiation presented in Table 37 (see Example 14 below).

[00270] Figure 101: Raw data for cell counts Day 11-Gen 2 REP initiation and Gen 3 Scale Up presented in Table 37 (see Example 14 below).

[00271] Figure 102: Raw data for cell counts Day 16-Gen 2 Scale Up and Gen 3 Harvest (*e.g.*, day 16) presented in Table 38 (see Example 14 below).

[00272] Figure 103: Raw data for cell counts Day 22-Gen 2 Harvest (*e.g.*, day 22) presented in Table 38 (see Example 14 below). For L4054 Gen 2, post LOVO count was extrapolated to 4 flasks, because was the total number of the study. 1 flask was contaminated, and the extrapolation was done for total = 6.67×10^{10}.

[00273] Figure 104: Raw data for flow cytometry results depicted in Figs. 87A, 87A, and 87B.

[00274] Figure 105: Raw data for flow cytometry results depicted in Figs. 87C and 87C.

[00275] Figure 106: Raw data for flow cytometry results depicted in Figs. 89 and 90.

[00276] Figure 107: Raw data for IFNγ production assay results for L4054 samples depicted in Fig. 91.

[00277] Figure 108: Raw data for IFNγ production assay results for L4055 samples depicted in Fig. 91.

[00278] Figure 109: Raw data for IFNγ production assay results for M1085T samples depicted in Fig. 91.

[00279] Figure 110: Raw data for IL-2 ELISA assay results depicted in Fig. 92.

[00280] Figure 111: Raw data for the metabolic substrate and metabolic analysis results presented in Figs. 93 and 94.

[00281] Figure 112: Raw data for the relative telomere length anaylsis results presented in Fig. 95.

[00282] Figure 113: Raw data for the unique CD3 sequence and clonal diversity analyses results presented in Figs. 96 and 99.

[00283] Figure 114: Shows a comparison between various Gen 2 (2A process) and the Gen 3.1 process embodiment.

[00284] Figure 115: Table describing various features of embodiments of the Gen 2, Gen 2.1 and Gen 3.0 process.

[00285] Figure 116: Overview of the media conditions for an embodiment of the Gen 3 process, referred to as Gen 3.1

[00286] Figure 117: Table describing various features of embodiments of the Gen 2, Gen 2.1 and Gen 3.0 process.

[00287] Figure 118: Table comparing various features of embodiments of the Gen 2 and Gen 3.0 processes.

[00288] Figure 119: Table providing media uses in the various embodiments of the described expansion processes.

[00289] Figure 120: Phenotype comparison: Gen 3.0 and Gen 3.1 embodiments of the process showed comparable CD28, CD27 and CD57 expression.

[00290] Figure 121: Higher production of IFNγ on Gen 3 final product. IFNγ analysis (by ELISA) was assessed in the culture frozen supernatant to compared both processes. For each tumor overnight stimulation with coated anti -CD3 plate, using fresh TIL product on each Gen 2 (*e.g.*, day 22) and Gen 3 process (*e.g.*, day 16). Each bar represents here are IFNγlevels of stimulated, unstimulated and media control.

[00291] Figure 122: Top: Unique CDR3 sequence analysis for TIL final product: Columns show the number of unique TCR B clonotypes identified from 1×10^6 cells collected on Gen 2 (*e.g.*, day 22) and Gen 3 process (*e.g.*, day 14-16). Gen 3 shows higher clonal diversity compared to Gen 2 based on the number of unique peptide CDRs within the sample. Bottom: Diversity Index for TIL final product: Shanon entropy diversity index is a more reliable a common metric for comparison. Gen 3 showed a slightly higher diversity than Gen 2.

[00292] Figure 123: 199 sequences are shared between Gen 3 and Gen 2 final product, corresponding to 97.07% of top 80% of unique CDR3 sequences from Gen 2 shared with Gen 3 final product.

[00293] Figure 124: 1833 sequences are shared between Gen 3 and Gen 2 final product, corresponding to 99.45% of top 80% of unique CDR3 sequences from Gen 2 shared with Gen 3 final product.

[00294] Figure 125: Schematic of an exemplary embodiment of the Gen 3 process (a 16-day process).

[00295] Figure 126: Schematic of an exemplary embodiment of a method for expanding TILS from hematopoietic malignancies using Gen 3 expansion platform.

[00296] Figure 127: Provides data showing the addition of OKT3 + feeders to cores at Day 3 did not significantly alter the phenotype of TIL compared to Gen2, indicating the TILs were healthy TILs.

[00297] Figure 128: Provides data showing expression of exhaustion and activation markers by melanoma core-derived TIL followed similar trends as that by pancreatic core-derived TIL. There were no differences observed in the phenotypic expression between IL-2 and the triple cocktail, in REP1 and REP2.

[00298] Figure 129: Provides data regarding expanding TIL from lung core biopsies.

[00299] Figure 130: Provides data showing that similar phenotypic profiles are observed in Day 0 versus Day 3 (OKT3 + feeders) treated lung cores.

[00300] Figure 131: Provides data showing enhanced phenotypic expression of activation and resident T cell markers in cores treated with OKT3 and feeders at Day 3.

[00301] Figure 132: TIL from lung cores are functional as determined by IFNγ secretion and CD107a mobilization in response to non-specific stimulation.

[00302] Figure 133: The CD8+ TIL from Day 3 treated lung cores are oligoclonal compared to the Day 0 treated TIL.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

[00303] SEQ ID NO:1 is the amino acid sequence of the heavy chain of muromonab.

[00304] SEQ ID NO:2 is the amino acid sequence of the light chain of muromonab.

[00305] SEQ ID NO:3 is the amino acid sequence of a recombinant human IL-2 protein.

[00306] SEQ ID NO:4 is the amino acid sequence of aldesleukin.

[00307] SEQ ID NO:5 is the amino acid sequence of a recombinant human IL-4 protein.

[00308] SEQ ID NO:6 is the amino acid sequence of a recombinant human IL-7 protein.

[00309] SEQ ID NO:7 is the amino acid sequence of a recombinant human IL-15 protein.

[00310] SEQ ID NO:8 is the amino acid sequence of a recombinant human IL-21 protein.

[00311] SEQ ID NO:9 is the amino acid sequence of human 4-1BB.

[00312] SEQ ID NO:10 is the amino acid sequence of murine 4-1BB.

[00313] SEQ ID NO:11 is the heavy chain for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00314] SEQ ID NO:12 is the light chain for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00315] SEQ ID NO:13 is the heavy chain variable region (VH) for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00316] SEQ ID NO:14 is the light chain variable region (VL) for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00317] SEQ ID NO:15 is the heavy chain CDRl for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00318] SEQ ID NO:16 is the heavy chain CDR2 for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00319] SEQ ID NO:17 is the heavy chain CDR3 for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00320] SEQ ID NO:18 is the light chain CDR1 for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00321] SEQ ID NO:19 is the light chain CDR2 for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00322] SEQ ID NO:20 is the light chain CDR3 for the 4-1BB agonist monoclonal antibody utomilumab (PF-05082566).

[00323] SEQ ID NO:21 is the heavy chain for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00324] SEQ ID NO:22 is the light chain for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00325] SEQ ID NO:23 is the heavy chain variable region (VH) for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00326] SEQ ID NO:24 is the light chain variable region (VL) for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00327] SEQ ID NO:25 is the heavy chain CDR1 for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00328] SEQ ID NO:26 is the heavy chain CDR2 for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00329] SEQ ID NO:27 is the heavy chain CDR3 for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00330] SEQ ID NO:28 is the light chain CDR1 for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00331] SEQ ID NO:29 is the light chain CDR2 for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00332] SEQ ID NO:30 is the light chain CDR3 for the 4-1BB agonist monoclonal antibody urelumab (BMS-663513).

[00333] SEQ ID NO:31 is an Fc domain for a TNFRSF agonist fusion protein.

[00334] SEQ ID NO:32 is a linker for a TNFRSF agonist fusion protein.

[00335] SEQ ID NO:33 is a linker for a TNFRSF agonist fusion protein.

[00336] SEQ ID NO:34 is a linker for a TNFRSF agonist fusion protein.

[00337] SEQ ID NO:35 is a linker for a TNFRSF agonist fusion protein.

[00338] SEQ ID NO:36 is a linker for a TNFRSF agonist fusion protein.

[00339] SEQ ID NO:37 is a linker for a TNFRSF agonist fusion protein.

[00340] SEQ ID NO:38 is a linker for a TNFRSF agonist fusion protein.

[00341] SEQ ID NO:39 is a linker for a TNFRSF agonist fusion protein.

[00342] SEQ ID NO:40 is a linker for a TNFRSF agonist fusion protein.

[00343] SEQ ID NO:41 is a linker for a TNFRSF agonist fusion protein.

[00344] SEQ ID NO:42 is an Fc domain for a TNFRSF agonist fusion protein.

[00345] SEQ ID NO:43 is a linker for a TNFRSF agonist fusion protein.

[00346] SEQ ID NO:44 is a linker for a TNFRSF agonist fusion protein.

[00347] SEQ ID NO:45 is a linker for a TNFRSF agonist fusion protein.

[00348] SEQ ID NO:46 is a 4-1BB ligand (4-1BBL) amino acid sequence.

[00349] SEQ ID NO:47 is a soluble portion of 4-1BBL polypeptide.

[00350] SEQ ID NO:48 is a heavy chain variable region (VH) for the 4-1BB agonist antibody 4B4-1-1 version 1.

[00351] SEQ ID NO:49 is a light chain variable region (VL) for the 4-1BB agonist antibody 4B4-1-1 version 1.

[00352] SEQ ID NO:50 is a heavy chain variable region (VH) for the 4-1BB agonist antibody 4B4-1-1 version 2.

[00353] SEQ ID NO:51 is a light chain variable region (VL) for the 4-1BB agonist antibody 4B4-1-1 version 2.

[00354] SEQ ID NO:52 is a heavy chain variable region (VH) for the 4-1BB agonist antibody H39E3-2.

[00355] SEQ ID NO:53 is a light chain variable region (VL) for the 4-1BB agonist antibody H39E3-2.

[00356] SEQ ID NO:54 is the amino acid sequence of human OX40.

[00357] SEQ ID NO:55 is the amino acid sequence of murine OX40.

[00358] SEQ ID NO:56 is the heavy chain for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00359] SEQ ID NO:57 is the light chain for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00360] SEQ ID NO:58 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00361] SEQ ID NO:59 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00362] SEQ ID NO:60 is the heavy chain CDRl for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00363] SEQ ID NO:61 is the heavy chain CDR2 for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00364] SEQ ID NO:62 is the heavy chain CDR3 for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00365] SEQ ID NO:63 is the light chain CDR1 for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00366] SEQ ID NO:64 is the light chain CDR2 for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00367] SEQ ID NO:65 is the light chain CDR3 for the OX40 agonist monoclonal antibody tavolixizumab (MEDI-0562).

[00368] SEQ ID NO:66 is the heavy chain for the OX40 agonist monoclonal antibody 11D4.

[00369] SEQ ID NO:67 is the light chain for the OX40 agonist monoclonal antibody 11D4.

[00370] SEQ ID NO:68 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody 11D4.

[00371] SEQ ID NO:69 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody 11D4.

[00372] SEQ ID NO:70 is the heavy chain CDRl for the OX40 agonist monoclonal antibody 11D4.

[00373] SEQ ID NO:71 is the heavy chain CDR2 for the OX40 agonist monoclonal antibody 11D4.

[00374] SEQ ID NO:72 is the heavy chain CDR3 for the OX40 agonist monoclonal antibody 11D4.

[00375] SEQ ID NO:73 is the light chain CDR1 for the OX40 agonist monoclonal antibody 11D4.

[00376] SEQ ID NO:74 is the light chain CDR2 for the OX40 agonist monoclonal antibody 11D4.

[00377] SEQ ID NO:75 is the light chain CDR3 for the OX40 agonist monoclonal antibody 11D4.

[00378] SEQ ID NO:76 is the heavy chain for the OX40 agonist monoclonal antibody 18D8.

[00379] SEQ ID NO:77 is the light chain for the OX40 agonist monoclonal antibody 18D8.

[00380] SEQ ID NO:78 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody 18D8.

[00381] SEQ ID NO:79 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody 18D8.

[00382] SEQ ID NO:80 is the heavy chain CDRl for the OX40 agonist monoclonal antibody 18D8.

[00383] SEQ ID NO:81 is the heavy chain CDR2 for the OX40 agonist monoclonal antibody 18D8.

[00384] SEQ ID NO:82 is the heavy chain CDR3 for the OX40 agonist monoclonal antibody 18D8.

[00385] SEQ ID NO:83 is the light chain CDR1 for the OX40 agonist monoclonal antibody 18D8.

[00386] SEQ ID NO:84 is the light chain CDR2 for the OX40 agonist monoclonal antibody 18D8.

[00387] SEQ ID NO:85 is the light chain CDR3 for the OX40 agonist monoclonal antibody 18D8.

[00388] SEQ ID NO:86 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody Hu119-122.

[00389] SEQ ID NO:87 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody Hu119-122.

[00390] SEQ ID NO:88 is the heavy chain CDRl for the OX40 agonist monoclonal antibody Hu119-122.

[00391] SEQ ID NO:89 is the heavy chain CDR2 for the OX40 agonist monoclonal antibody Hu119-122.

[00392] SEQ ID NO:90 is the heavy chain CDR3 for the OX40 agonist monoclonal antibody Hu119-122.

[00393] SEQ ID NO:91 is the light chain CDR1 for the OX40 agonist monoclonal antibody Hu119-122.

[00394] SEQ ID NO:92 is the light chain CDR2 for the OX40 agonist monoclonal antibody Hu119-122.

[00395] SEQ ID NO:93 is the light chain CDR3 for the OX40 agonist monoclonal antibody Hu119-122.

[00396] SEQ ID NO:94 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody Hu106-222.

[00397] SEQ ID NO:95 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody Hu106-222.

[00398] SEQ ID NO:96 is the heavy chain CDRl for the OX40 agonist monoclonal antibody Hu106-222.

[00399] SEQ ID NO:97 is the heavy chain CDR2 for the OX40 agonist monoclonal antibody Hu106-222.

[00400] SEQ ID NO:98 is the heavy chain CDR3 for the OX40 agonist monoclonal antibody Hu106-222.

[00401] SEQ ID NO:99 is the light chain CDR1 for the OX40 agonist monoclonal antibody Hu106-222.

[00402] SEQ ID NO:100 is the light chain CDR2 for the OX40 agonist monoclonal antibody Hu106-222.

[00403] SEQ ID NO:101 is the light chain CDR3 for the OX40 agonist monoclonal antibody Hu106-222.

[00404] SEQ ID NO:102 is an OX40 ligand (OX40L) amino acid sequence.

[00405] SEQ ID NO:103 is a soluble portion of OX40L polypeptide.

[00406] SEQ ID NO:104 is an alternative soluble portion of OX40L polypeptide.

[00407] SEQ ID NO:105 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody 008.

[00408] SEQ ID NO:106 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody 008.

[00409] SEQ ID NO:107 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody 011.

[00410] SEQ ID NO:108 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody 011.

[00411] SEQ ID NO:109 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody 021.

[00412] SEQ ID NO:110 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody 021.

[00413] SEQ ID NO:111 is the heavy chain variable region (VH) for the OX40 agonist monoclonal antibody 023.

[00414] SEQ ID NO:112 is the light chain variable region (VL) for the OX40 agonist monoclonal antibody 023.

[00415] SEQ ID NO:113 is the heavy chain variable region (VH) for an OX40 agonist monoclonal antibody.

[00416] SEQ ID NO:114 is the light chain variable region (VL) for an OX40 agonist monoclonal antibody.

[00417] SEQ ID NO:115 is the heavy chain variable region (VH) for an OX40 agonist monoclonal antibody.

[00418] SEQ ID NO:116 is the light chain variable region (VL) for an OX40 agonist monoclonal antibody.

[00419] SEQ ID NO:117 is the heavy chain variable region (VH) for a humanized OX40 agonist monoclonal antibody.

[00420] SEQ ID NO:118 is the heavy chain variable region (VH) for a humanized OX40 agonist monoclonal antibody.

[00421] SEQ ID NO:119 is the light chain variable region (VL) for a humanized OX40 agonist monoclonal antibody.

[00422] SEQ ID NO:120 is the light chain variable region (VL) for a humanized OX40 agonist monoclonal antibody.

[00423] SEQ ID NO:121 is the heavy chain variable region (VH) for a humanized OX40 agonist monoclonal antibody.

[00424] SEQ ID NO:122 is the heavy chain variable region (VH) for a humanized OX40 agonist monoclonal antibody.

[00425] SEQ ID NO:123 is the light chain variable region (VL) for a humanized OX40 agonist monoclonal antibody.

[00426] SEQ ID NO:124 is the light chain variable region (VL) for a humanized OX40 agonist monoclonal antibody.

[00427] SEQ ID NO:125 is the heavy chain variable region (VH) for an OX40 agonist monoclonal antibody.

[00428] SEQ ID NO:126 is the light chain variable region (VL) for an OX40 agonist monoclonal antibody.

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

[00429] Adoptive cell therapy utilizing TILs cultured *ex vivo* by the Rapid Expansion Protocol (REP) has produced successful adoptive cell therapy following host immunosuppression in patients with melanoma. Current infusion acceptance parameters rely on readouts of the composition of TILs (*e.g.*, CD28, CD8, or CD4 positivity) and on the numerical folds of expansion and viability of the REP product.

[00430] Current REP protocols give little insight into the health of the TIL that will be infused into the patient. T cells undergo a profound metabolic shift during the course of their maturation from naïve to effector T cells (see Chang, *et al.*, *Nat. Immunol.* **2016,** *17,* 364, hereby expressly incorporated in its entirety, and in particular for the discussion and markers of anaerobic and aerobic metabolism). For example, naïve T cells rely on mitochondrial respiration to produce ATP, while mature, healthy effector T cells such as TIL are highly glycolytic, relying on aerobic glycolysis to provide the bioenergetics substrates they require for proliferation, migration, activation, and anti-tumor efficacy.

[00431] Previous papers report that limiting glycolysis and promoting mitochondrial metabolism in TILs prior to transfer is desirable as cells that are relying heavily on glycolysis will suffer nutrient deprivation upon adoptive transfer which results in a majority of the transferred cells dying. Thus, the art teaches that promoting mitochondrial metabolism might

promote *in vivo* longevity and in fact suggests using inhibitors of glycolysis before induction of the immune response. See Chang et al. (Chang, *et al.*, *Nat. Immunol.* **2016,** 17(364).

[00432] The present invention is further directed in some embodiments to methods for evaluating and quantifying this increase in metabolic health. Thus, the present invention provides methods of assaying the relative health of a TIL population using one or more general evaluations of metabolism, including, but not limited to, rates and amounts of glycolysis, oxidative phosphorylation, spare respiratory capacity (SRC), and glycolytic reserve.

[00433] Furthermore, the present invention is further directed in some embodiments to methods for evaluating and quantifying this increase in metabolic health. Thus, the present invention provides methods of assaying the relative health of a TIL population using one or more general evaluations of metabolism, including, but not limited to, rates and amounts of glycolysis, oxidative phosphorylation, spare respiratory capacity (SRC), and glycolytic reserve.

[00434] In addition, optional additional evaluations include, but are not limited to, ATP production, mitochondrial mass and glucose uptake.

II. Definitions

[00435] The term "*in vivo*" refers to an event that takes place in a subject's body.

[00436] The term "*in vitro*" refers to an event that takes places outside of a subject's body. *In vitro* assays encompass cell-based assays in which cells alive or dead are employed and may also encompass a cell-free assay in which no intact cells are employed.

[00437] The term "rapid expansion" means an increase in the number of antigen-specific TILs of at least about 3-fold (or 4-, 5-, 6-, 7-, 8-, or 9-fold) over a period of a week, more preferably at least about 10-fold (or 20-, 30-, 40-, 50-, 60-, 70-, 80-, or 90-fold) over a period of a week, or most preferably at least about 100-fold over a period of a week. A number of rapid expansion protocols are outlined below.

[00438] By "tumor infiltrating lymphocytes" or "TILs" herein is meant a population of cells originally obtained as white blood cells that have left the bloodstream of a subject and migrated into a tumor. TILs include, but are not limited to, $CD8^+$ cytotoxic T cells (lymphocytes), Th1 and Th17 $CD4^+$ T cells, natural killer cells, dendritic cells and M1 macrophages. TILs include both primary and secondary TILs. "Primary TILs" are those that are obtained from patient tissue samples as outlined herein (sometimes referred to as "freshly

harvested"), and "secondary TILs" are any TIL cell populations that have been expanded or proliferated as discussed herein, including, but not limited to bulk TILs and expanded TILs ("REP TILs" or "post-REP TILs").

[00439] By "population of cells" (including TILs) herein is meant a number of cells that share common traits. In general, populations generally range from 1×10^6 to 1×10^{10} in number, with different TIL populations comprising different numbers. For example, initial growth of primary TILs in the presence of IL-2 results in a population of bulk TILs of roughly 1×10^8 cells. REP expansion is generally done to provide populations of 1.5×10^9 to 1.5×10^{10} cells for infusion.

[00440] By "small biopsy" herein is meant a biopsy that effects the removal of a volume of tissue from a tumor that is typically less, and in some embodiments substantially less, than the volume of the tumor. Small biopsy includes core biopsies, punch biopsies and the like. Any small biopsy is included, for example any biopsy with a volume less than about 1 mm^3, 2 mm^3, about 5 mm^3, about 10 mm^3, about 25 mm^3, about 50 mm^3, or about 100 mm^3 or with a cross-sectional area of less than about 1 mm^2, 2 mm^2, about 5 mm^2, about 10 mm^2, about 25 mm^2, about 50 mm^2, or about 100 mm^2. A small biopsy can also include multiple biopsies taken from the same or multiple locations in the body, including from different cancerous lesions in metastatic disease.

[00441] By "cryopreserved TILs" herein is meant that TILs, either primary, bulk, or expanded (REP TILs), are treated and stored in the range of about -150°C to -60°C. General methods for cryopreservation are also described elsewhere herein, including in the Examples. For clarity, "cryopreserved TILs" are distinguishable from frozen tissue samples which may be used as a source of primary TILs.

[00442] By "thawed cryopreserved TILs" herein is meant a population of TILs that was previously cryopreserved and then treated to return to room temperature or higher, including but not limited to cell culture temperatures or temperatures wherein TILs may be administered to a patient.

[00443] TILs can generally be defined either biochemically, using cell surface markers, or functionally, by their ability to infiltrate tumors and effect treatment. TILs can be generally categorized by expressing one or more of the following biomarkers: CD4, CD8, TCR αβ, CD27, CD28, CD56, CCR7, CD45Ra, CD95, PD-1, and CD25. Additionally and

alternatively, TILs can be functionally defined by their ability to infiltrate solid tumors upon reintroduction into a patient.

[00444] The term "cryopreservation media" or "cryopreservation medium" refers to any medium that can be used for cryopreservation of cells. Such media can include media comprising 7% to 10% DMSO. Exemplary media include CryoStor CS10, Hyperthermasol, as well as combinations thereof. The term "CS10" refers to a cryopreservation medium which is obtained from Stemcell Technologies or from Biolife Solutions. The CS10 medium may be referred to by the trade name "CryoStor® CS10". The CS10 medium is a serum-free, animal component-free medium which comprises DMSO.

[00445] By "population of cells" (including TILs) herein is meant a number of cells that share common traits. In general, populations generally range from 1×10^6 to 1×10^{10} in number, with different TIL populations comprising different numbers. For example, initial growth of primary TILs in the presence of IL-2 results in a population of bulk TILs of roughly 1×10^8 cells. REP expansion is generally done to provide populations of 1.5×10^9 to 1.5×10^{10} cells for infusion.

[00446] The term "central memory T cell" refers to a subset of T cells that in the human are CD45R0+ and constitutively express CCR7 ($CCR7^{hi}$) and CD62L ($CD62^{hi}$). The surface phenotype of central memory T cells also includes TCR, CD3, CD127 (IL-7R), and IL-15R. Transcription factors for central memory T cells include BCL-6, BCL-6B, MBD2, and BMI1. Central memory T cells primarily secret IL-2 and CD40L as effector molecules after TCR triggering. Central memory T cells are predominant in the CD4 compartment in blood, and in the human are proportionally enriched in lymph nodes and tonsils.

[00447] The term "effector memory T cell" refers to a subset of human or mammalian T cells that, like central memory T cells, are CD45R0+, but have lost the constitutive expression of CCR7 ($CCR7^{lo}$) and are heterogeneous or low for CD62L expression ($CD62L^{lo}$). The surface phenotype of central memory T cells also includes TCR, CD3, CD127 (IL-7R), and IL-15R. Transcription factors for central memory T cells include BLIMP1. Effector memory T cells rapidly secret high levels of inflammatory cytokines following antigenic stimulation, including interferon-γ, IL-4, and IL-5. Effector memory T cells are predominant in the CD8 compartment in blood, and in the human are proportionally enriched in the lung, liver, and gut. CD8+ effector memory T cells carry large amounts of perforin. The term "closed system" refers to a system that is closed to the outside

environment. Any closed system appropriate for cell culture methods can be employed with the methods of the present invention. Closed systems include, for example, but are not limited to closed G-containers. Once a tumor segment is added to the closed system, the system is no opened to the outside environment until the TILs are ready to be administered to the patient.

[00448] The terms "fragmenting," "fragment," and "fragmented," as used herein to describe processes for disrupting a tumor, includes mechanical fragmentation methods such as crushing, slicing, dividing, and morcellating tumor tissue as well as any other method for disrupting the physical structure of tumor tissue.

[00449] The terms "peripheral blood mononuclear cells" and "PBMCs" refers to a peripheral blood cell having a round nucleus, including lymphocytes (T cells, B cells, NK cells) and monocytes. When used as an antigen-presenting cell (PBMCs are a type of antigen-presenting cell), the peripheral blood mononuclear cells are irradiated allogeneic peripheral blood mononuclear cells.

[00450] The terms "peripheral blood lymphocytes" and "PBLs" refers to T-cells expanded from peripheral blood. In some embodiments, PBLs are separated from whole blood or apheresis product from a donor. In some embodiments, PBLs are separated from whole blood or apheresis product from a donor by positive or negative selection of a T cell phenotype, such as the T cell phenotype of CD3+ CD45+.

[00451] The term "anti-CD3 antibody" refers to an antibody or variant thereof, *e.g.*, a monoclonal antibody and including human, humanized, chimeric or murine antibodies which are directed against the CD3 receptor in the T cell antigen receptor of mature T cells. Anti-CD3 antibodies include OKT-3, also known as muromonab. Other anti-CD3 antibodies include, for example, otelixizumab, teplizumab, and visilizumab.

[00452] The term "OKT-3" (also referred to herein as "OKT3") refers to a monoclonal antibody or biosimilar or variant thereof, including human, humanized, chimeric, or murine antibodies, directed against the CD3 receptor in the T cell antigen receptor of mature T cells, and includes commercially-available forms such as OKT-3 (30 ng/mL, MACS GMP CD3 pure, Miltenyi Biotech, Inc., San Diego, CA, USA) and muromonab or variants, conservative amino acid substitutions, glycoforms, or biosimilars thereof. The amino acid sequences of the heavy and light chains of muromonab are given in Table 1 (SEQ ID NO:1 and SEQ ID NO:2). A hybridoma capable of producing OKT-3 is deposited with the American Type

Culture Collection and assigned the ATCC accession number CRL 8001. A hybridoma capable of producing OKT-3 is also deposited with European Collection of Authenticated Cell Cultures (ECACC) and assigned Catalogue No. 86022706.

TABLE 1. Amino acid sequences of muromonab.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:1 Muromonab heavy chain	QVQLQQSGAE LARPGASVKM SCKASGYTFT RYTMHWVKQR PGQGLEWIGY INPSRGYTNY NQKFKDKATL TTDKSSSTAY MQLSSLTSED SAVYYCARYY DDHYCLDYWG QGTTLTVSSA KTTAPSVYPL APVCGGTTGS SVTLGCLVKG YFPEPVTLTW NSGSLSSGVH TFPAVLQSDL YTLSSSVTVT SSTWPSQSIT CNVAHPASST KVDKKIEPRP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK	60 120 180 240 300 360 420 450
SEQ ID NO:2 Muromonab light chain	QIVLTQSPAI MSASPGEKVT MTCSASSSVS YMNWYQQKSG TSPKRWIYDT SKLASGVPAH FRGSGSGTSY SLTISGMEAE DAATYYCQQW SSNPFTFGSG TKLEINRADT APTVSIFPPS SEQLTSGGAS VVCFLNNFYP KDINVKWKID GSERQNGVLN SWTDQDSKDS TYSMSSTLTL TKDEYERHNS YTCEATHKTS TSPIVKSFNR NEC	60 120 180 213

[00453] The term "IL-2" (also referred to herein as "IL2") refers to the T cell growth factor known as interleukin-2, and includes all forms of IL-2 including human and mammalian forms, conservative amino acid substitutions, glycoforms, biosimilars, and variants thereof. IL-2 is described, *e.g.*, in Nelson, *J. Immunol.* **2004,** *172,* 3983-88 and Malek, *Annu. Rev. Immunol.* **2008,** *26,* 453-79, the disclosures of which are incorporated by reference herein. The amino acid sequence of recombinant human IL-2 suitable for use in the invention is given in Table 2 (SEQ ID NO:3). For example, the term IL-2 encompasses human, recombinant forms of IL-2 such as aldesleukin (PROLEUKIN, available commercially from multiple suppliers in 22 million IU per single use vials), as well as the form of recombinant IL-2 commercially supplied by CellGenix, Inc., Portsmouth, NH, USA (CELLGRO GMP) or ProSpec-Tany TechnoGene Ltd., East Brunswick, NJ, USA (Cat. No. CYT-209-b) and other commercial equivalents from other vendors. Aldesleukin (des-alanyl-1, serine-125 human IL-2) is a nonglycosylated human recombinant form of IL-2 with a molecular weight of approximately 15 kDa. The amino acid sequence of aldesleukin suitable for use in the invention is given in Table 2 (SEQ ID NO:4). The term IL-2 also encompasses pegylated forms of IL-2, as described herein, including the pegylated IL2 prodrug NKTR-214, available from Nektar Therapeutics, South San Francisco, CA, USA. NKTR-214 and pegylated IL-2 suitable for use in the invention is described in U.S. Patent Application Publication No. US 2014/0328791 A1 and International Patent Application Publication No. WO 2012/065086 Al, the disclosures of which are incorporated by reference herein. Alternative forms of conjugated IL-2 suitable for use in the invention are described in U.S. Patent Nos. 4,766,106, 5,206,344, 5,089,261 and 4902,502, the disclosures of which are incorporated by reference

herein. Formulations of IL-2 suitable for use in the invention are described in U.S. Patent No. 6,706,289, the disclosure of which is incorporated by reference herein.

TABLE 2. Amino acid sequences of interleukins.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:3 recombinant human IL-2 (rhIL-2)	MAPTSSSTKK TQLQLEHLLL DLQMILNGIN NYKNPKLTRM LTFKFYMPKK ATELKHLQCL EEELKPLEEV LNLAQSKNFH LRPRDLISNI NVIVLELKGS ETTFMCEYAD ETATIVEFLN RWITFCQSII STLT	60 120 134
SEQ ID NO:4 Aldesleukin	PTSSSTKKTQ LQLEHLLLDL QMILNGINNY KNPKLTRMLT FKFYMPKKAT ELKHLQCLEE ELKPLEEVLN LAQSKNFHLR PRDLISNINV IVLELKGSET TFMCEYADET ATIVEFLNRW ITFSQSIIST LT	60 120 132
SEQ ID NO:5 recombinant human IL-4 (rhIL-4)	MHKCDITLQE IIKTLNSLTE QKTLCTELTV TDIFAASKNT TEKETFCRAA TVLRQFYSHH EKDTRCLGAT AQQFHRHKQL IRFLKRLDRN LWGLAGLNSC PVKEANQSTL ENFLERLKTI MREKYSKCSS	60 120 130
SEQ ID NO:6 recombinant human IL-7 (rhIL-7)	MDCDIEGKDG KQYESVLMVS IDQLLDSMKE IGSNCLNNEF NFFKRHICDA NKEGMFLFRA ARKLRQFLKM NSTGDFDLHL LKVSEGTTIL LNCTGQVKGR KPAALGEAQP TKSLEENKSL KEQKKLNDLC FLKRLLQEIK TCWNKILMGT KEH	60 120 153
SEQ ID NO:7 recombinant human IL-15 (rhIL-15)	MNWVNVISDL KKIEDLIQSM HIDATLYTES DVHPSCKVTA MKCFLLELQV ISLESGDASI HDTVENLIIL ANNSLSSNGN VTESGCKECE ELEEKNIKEF LQSFVHIVQM FINTS	60 115
SEQ ID NO:8 recombinant human IL-21 (rhIL-21)	MQDRHMIRMR QLIDIVDQLK NYVNDLVPEF LPAPEDVETN CEWSAFSCFQ KAQLKSANTG NNERIINVSI KKLKRKPPST NAGRRQKHRL TCPSCDSYEK KPPKEFLERF KSLLQKMIHQ HLSSRTHGSE DS	60 120 132

[00454] The term "IL-4" (also referred to herein as "IL4") refers to the cytokine known as interleukin 4, which is produced by Th2 T cells and by eosinophils, basophils, and mast cells. IL-4 regulates the differentiation of naïve helper T cells (Th0 cells) to Th2 T cells. Steinke and Borish, *Respir. Res.* **2001,** *2,* 66-70. Upon activation by IL-4, Th2 T cells subsequently produce additional IL-4 in a positive feedback loop. IL-4 also stimulates B cell proliferation and class II MHC expression, and induces class switching to IgE and IgG_1 expression from B cells. Recombinant human IL-4 suitable for use in the invention is commercially available from multiple suppliers, including ProSpec-Tany TechnoGene Ltd., East Brunswick, NJ, USA (Cat. No. CYT-211) and ThermoFisher Scientific, Inc., Waltham, MA, USA (human IL-15 recombinant protein, Cat. No. Gibco CTP0043). The amino acid sequence of recombinant human IL-4 suitable for use in the invention is given in Table 2 (SEQ ID NO:5).

[00455] The term "IL-7" (also referred to herein as "IL7") refers to a glycosylated tissue-derived cytokine known as interleukin 7, which may be obtained from stromal and epithelial cells, as well as from dendritic cells. Fry and Mackall, *Blood* **2002,** *99,* 3892-904. IL-7 can stimulate the development of T cells. IL-7 binds to the IL-7 receptor, a heterodimer consisting of IL-7 receptor alpha and common gamma chain receptor, which in a series of signals important for T cell development within the thymus and survival within the periphery. Recombinant human IL-4 suitable for use in the invention is commercially available from

multiple suppliers, including ProSpec-Tany TechnoGene Ltd., East Brunswick, NJ, USA (Cat. No. CYT-254) and ThermoFisher Scientific, Inc., Waltham, MA, USA (human IL-15 recombinant protein, Cat. No. Gibco PHC0071). The amino acid sequence of recombinant human IL-7 suitable for use in the invention is given in Table 2 (SEQ ID NO:6).

[00456] The term "IL-15" (also referred to herein as "IL15") refers to the T cell growth factor known as interleukin-15, and includes all forms of IL-2 including human and mammalian forms, conservative amino acid substitutions, glycoforms, biosimilars, and variants thereof. IL-15 is described, *e.g.*, in Fehniger and Caligiuri, *Blood* **2001,** *97,* 14-32, the disclosure of which is incorporated by reference herein. IL-15 shares β and γ signaling receptor subunits with IL-2. Recombinant human IL-15 is a single, non-glycosylated polypeptide chain containing 114 amino acids (and an N-terminal methionine) with a molecular mass of 12.8 kDa. Recombinant human IL-15 is commercially available from multiple suppliers, including ProSpec-Tany TechnoGene Ltd., East Brunswick, NJ, USA (Cat. No. CYT-230-b) and ThermoFisher Scientific, Inc., Waltham, MA, USA (human IL-15 recombinant protein, Cat. No. 34-8159-82). The amino acid sequence of recombinant human IL-15 suitable for use in the invention is given in Table 2 (SEQ ID NO:7).

[00457] The term "IL-21" (also referred to herein as "IL21") refers to the pleiotropic cytokine protein known as interleukin-21, and includes all forms of IL-21 including human and mammalian forms, conservative amino acid substitutions, glycoforms, biosimilars, and variants thereof. IL-21 is described, *e.g.*, in Spolski and Leonard, *Nat. Rev. Drug. Disc.* **2014,** *13,* 379-95, the disclosure of which is incorporated by reference herein. IL-21 is primarily produced by natural killer T cells and activated human $CD4^+$ T cells. Recombinant human IL-21 is a single, non-glycosylated polypeptide chain containing 132 amino acids with a molecular mass of 15.4 kDa. Recombinant human IL-21 is commercially available from multiple suppliers, including ProSpec-Tany TechnoGene Ltd., East Brunswick, NJ, USA (Cat. No. CYT-408-b) and ThermoFisher Scientific, Inc., Waltham, MA, USA (human IL-21 recombinant protein, Cat. No. 14-8219-80). The amino acid sequence of recombinant human IL-21 suitable for use in the invention is given in Table 2 (SEQ ID NO:8).

[00458] When "an anti-tumor effective amount", "a tumor-inhibiting effective amount", or "therapeutic amount" is indicated, the precise amount of the compositions of the present invention to be administered can be determined by a physician with consideration of individual differences in age, weight, tumor size, extent of infection or metastasis, and condition of the patient (subject). It can generally be stated that a pharmaceutical composition

comprising the genetically modified cytotoxic lymphocytes described herein may be administered at a dosage of 10^4 to 10^{11} cells/kg body weight (*e.g.*, 10^5 to 10^6, 10^5 to 10^{10}, 10^5 to 10^{11}, 10^6 to 10^{10}, 10^6 to 10^{11},10^7 to 10^{11}, 10^7 to 10^{10}, 10^8 to 10^{11}, 10^8 to 10^{10}, 10^9 to 10^{11}, or 10^9 to 10^{10} cells/kg body weight), including all integer values within those ranges. Genetically modified cytotoxic lymphocytes compositions may also be administered multiple times at these dosages. The genetically modified cytotoxic lymphocytes can be administered by using infusion techniques that are commonly known in immunotherapy (see, *e.g.*, Rosenberg *et al.*, *New Eng. J. of Med.* **319**: 1676, 1988). The optimal dosage and treatment regime for a particular patient can readily be determined by one skilled in the art of medicine by monitoring the patient for signs of disease and adjusting the treatment accordingly.

[00459] The term "hematological malignancy," "hematologic malignancy" or terms of correlative meaning refer to mammalian cancers and tumors of the hematopoietic and lymphoid tissues, including but not limited to tissues of the blood, bone marrow, lymph nodes, and lymphatic system. Hematological malignancies are also referred to as "liquid tumors." Hematological malignancies include, but are not limited to, acute lymphoblastic leukemia (ALL), chronic lymphocytic lymphoma (CLL), small lymphocytic lymphoma (SLL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), acute monocytic leukemia (AMoL), Hodgkin's lymphoma, and non-Hodgkin's lymphomas. The term "B cell hematological malignancy" refers to hematological malignancies that affect B cells.

[00460] The term "solid tumor" refers to an abnormal mass of tissue that usually does not contain cysts or liquid areas. Solid tumors may be benign or malignant. The term "solid tumor cancer refers to malignant, neoplastic, or cancerous solid tumors. Solid tumor cancers include, but are not limited to, sarcomas, carcinomas, and lymphomas, such as cancers of the lung, breast, prostate, colon, rectum, and bladder. The tissue structure of solid tumors includes interdependent tissue compartments including the parenchyma (cancer cells) and the supporting stromal cells in which the cancer cells are dispersed and which may provide a supporting microenvironment.

[00461] The term "fine needle aspirate" or FNA refers to a type of biopsy procedure that can be employed for sampling or diagnostic procedures, including tumor sampling, in which a sample is taken but the tumor is not removed or resected. In fine needle aspiration, a hollow needle, for example 25-18 gauge, is inserted into the tumor or into an area containing the tumor and fluid and cells (including tissue) are obtained for further analysis or expansion, as

described herein. With an FNA, the cells are removed without preserving the histological architecture of the tissue cells. An FNA can comprise TILs. In some instances, a fine needle aspiration biopsy is performed using an ultrasound-guided fine needle aspiration biopsy needle. FNA needles are commercially available from Becton Dickinson, Covidien, and the like.

[00462] The term "core biopsy" or "core needle biopsy" refers to a type of biopsy procedure that can be employed for sampling or diagnostic procedures, including tumor sampling, in which a sample is taken but the tumor is not removed or resected. In a core biopsy, a hollow needle, example 16-11 gauge, is inserted into the tumor or into an area containing the tumor and fluid and cells (including tissue) are obtained for further analysis or expansion, as described herein. With a core biopsy, the cells can be removed with some preservation of the histological architecture of the tissue cells, given the larger needle size as compared to a FNA. The core biopsy needle is generally of a gauge size that is able to preserve at least some portion of the histological architecture of the tumor. A core biopsy can comprise TILs. In some instances, a core needle biopsy is performed using a biopsy instrument, a vacuum-assisted core-needle biopsy instrument, a steretactically guided core-needle biopsy instrument, an ultrasound-guided core-needle biopsy instrument, an MRI-guided core-needle biopsy instrument commercially available from Bard Medical, Becton Dickinson, and the like.

[00463] The term "liquid tumor" refers to an abnormal mass of cells that is fluid in nature. Liquid tumor cancers include, but are not limited to, leukemias, myelomas, and lymphomas, as well as other hematological malignancies. TILs obtained from liquid tumors may also be referred to herein as marrow infiltrating lymphocytes (MILs). TILs obtained from liquid tumors, including liquid tumors circulating in peripheral blood, may also be referred to herein as PBLs. The terms MIL, TIL, and PBL are used interchangeably herein and differ only based on the tissue type from which the cells are derived.

[00464] The term "microenvironment," as used herein, may refer to the solid or hematological tumor microenvironment as a whole or to an individual subset of cells within the microenvironment. The tumor microenvironment, as used herein, refers to a complex mixture of "cells, soluble factors, signaling molecules, extracellular matrices, and mechanical cues that promote neoplastic transformation, support tumor growth and invasion, protect the tumor from host immunity, foster therapeutic resistance, and provide niches for dominant metastases to thrive," as described in Swartz, *et al., Cancer Res.*, **2012,** *72,* 2473. Although

tumors express antigens that should be recognized by T cells, tumor clearance by the immune system is rare because of immune suppression by the microenvironment.

[00465] In an embodiment, the invention includes a method of treating a cancer with a population of remnant TILs (rTILs), wherein a patient is pre-treated with non-myeloablative chemotherapy prior to an infusion of rTILs according to the invention. In some embodiments, the population of rTILs may be provided with a population of normal emigrant TILs (eTILs), wherein a patient is pre-treated with nonmyeloablative chemotherapy prior to an infusion of rTILs and eTILs according to the invention. In an embodiment, the non-myeloablative chemotherapy is cyclophosphamide 60 mg/kg/d for 2 days (days 27 and 26 prior to rTIL infusion) and fludarabine 25 $mg/m^2/d$ for 5 days (days 27 to 23 prior to rTIL infusion). In an embodiment, after non-myeloablative chemotherapy and rTIL infusion (at day 0) according to the invention, the patient receives an intravenous infusion of IL-2 intravenously at 720,000 IU/kg every 8 hours to physiologic tolerance.

[00466] Experimental findings indicate that lymphodepletion prior to adoptive transfer of tumor-specific T lymphocytes plays a key role in enhancing treatment efficacy by eliminating regulatory T cells and competing elements of the immune system ("cytokine sinks"). Accordingly, some embodiments of the invention utilize a lymphodepletion step (sometimes also referred to as "immunosuppressive conditioning") on the patient prior to the introduction of the rTILs of the invention.

[00467] The terms "co-administration," "co-administering," "administered in combination with," "administering in combination with," "simultaneous," and "concurrent," as used herein, encompass administration of two or more active pharmaceutical ingredients (in a preferred embodiment of the present invention, for example, at least one potassium channel agonist in combination with a plurality of TILs) to a subject so that both active pharmaceutical ingredients and/or their metabolites are present in the subject at the same time. Co-administration includes simultaneous administration in separate compositions, administration at different times in separate compositions, or administration in a composition in which two or more active pharmaceutical ingredients are present. Simultaneous administration in separate compositions and administration in a composition in which both agents are present are preferred.

[00468] The term "effective amount" or "therapeutically effective amount" refers to that amount of a compound or combination of compounds as described herein that is sufficient to

effect the intended application including, but not limited to, disease treatment. A therapeutically effective amount may vary depending upon the intended application (*in vitro* or *in vivo*), or the subject and disease condition being treated (*e.g.*, the weight, age and gender of the subject), the severity of the disease condition, or the manner of administration. The term also applies to a dose that will induce a particular response in target cells (*e.g.*, the reduction of platelet adhesion and/or cell migration). The specific dose will vary depending on the particular compounds chosen, the dosing regimen to be followed, whether the compound is administered in combination with other compounds, timing of administration, the tissue to which it is administered, and the physical delivery system in which the compound is carried.

[00469] The terms "treatment", "treating", "treat", and the like, refer to obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease. "Treatment", as used herein, covers any treatment of a disease in a mammal, particularly in a human, and includes: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, *i.e.*, arresting its development or progression; and (c) relieving the disease, *i.e.*, causing regression of the disease and/or relieving one or more disease symptoms. "Treatment" is also meant to encompass delivery of an agent in order to provide for a pharmacologic effect, even in the absence of a disease or condition. For example, "treatment" encompasses delivery of a composition that can elicit an immune response or confer immunity in the absence of a disease condition, *e.g.*, in the case of a vaccine.

[00470] The term "heterologous" when used with reference to portions of a nucleic acid or protein indicates that the nucleic acid or protein comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, *e.g.*, a promoter from one source and a coding region from another source, or coding regions from different sources. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (*e.g.*, a fusion protein).

[00471] The terms "sequence identity," "percent identity," and "sequence percent identity" (or synonyms thereof, *e.g.*, "99% identical") in the context of two or more nucleic acids or

polypeptides, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned (introducing gaps, if necessary) for maximum correspondence, not considering any conservative amino acid substitutions as part of the sequence identity. The percent identity can be measured using sequence comparison software or algorithms or by visual inspection. Various algorithms and software are known in the art that can be used to obtain alignments of amino acid or nucleotide sequences. Suitable programs to determine percent sequence identity include for example the BLAST suite of programs available from the U.S. Government's National Center for Biotechnology Information BLAST web site. Comparisons between two sequences can be carried using either the BLASTN or BLASTP algorithm. BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. ALIGN, ALIGN-2 (Genentech, South San Francisco, California) or MegAlign, available from DNASTAR, are additional publicly available software programs that can be used to align sequences. One skilled in the art can determine appropriate parameters for maximal alignment by particular alignment software. In certain embodiments, the default parameters of the alignment software are used.

[00472] As used herein, the term "variant" encompasses but is not limited to antibodies or fusion proteins which comprise an amino acid sequence which differs from the amino acid sequence of a reference antibody by way of one or more substitutions, deletions and/or additions at certain positions within or adjacent to the amino acid sequence of the reference antibody. The variant may comprise one or more conservative substitutions in its amino acid sequence as compared to the amino acid sequence of a reference antibody. Conservative substitutions may involve, *e.g.*, the substitution of similarly charged or uncharged amino acids. The variant retains the ability to specifically bind to the antigen of the reference antibody. The term variant also includes pegylated antibodies or proteins.

[00473] The term "*in vivo*" refers to an event that takes place in a subject's body.

[00474] The term "*in vitro*" refers to an event that takes places outside of a subject's body. *In vitro* assays encompass cell-based assays in which cells alive or dead are employed and may also encompass a cell-free assay in which no intact cells are employed.

[00475] The term "rapid expansion" means an increase in the number of antigen-specific TILs of at least about 3-fold (or 4-, 5-, 6-, 7-, 8-, or 9-fold) over a period of a week, more preferably at least about 10-fold (or 20-, 30-, 40-, 50-, 60-, 70-, 80-, or 90-fold) over a period

of a week, or most preferably at least about 100-fold over a period of a week. A number of rapid expansion protocols are outlined below.

[00476] By "tumor infiltrating lymphocytes" or "TILs" herein is meant a population of cells originally obtained as white blood cells that have left the bloodstream of a subject and migrated into a tumor. TILs include, but are not limited to, $CD8^+$ cytotoxic T cells (lymphocytes), Th1 and Th17 $CD4^+$ T cells, natural killer cells, dendritic cells and M1 macrophages. TILs include both primary and secondary TILs. "Primary TILs" are those that are obtained from patient tissue samples as outlined herein (sometimes referred to as "freshly harvested"), and "secondary TILs" are any TIL cell populations that have been expanded or proliferated as discussed herein, including, but not limited to bulk TILs, expanded TILs ("REP TILs") as well as "reREP TILs" as discussed herein. reREP TILs can include for example second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs).

[00477] TILs can generally be defined either biochemically, using cell surface markers, or functionally, by their ability to infiltrate tumors and effect treatment. TILs can be generally categorized by expressing one or more of the following biomarkers: CD4, CD8, TCR αβ, CD27, CD28, CD56, CCR7, CD45Ra, CD95, PD-1, and CD25. Additionally, and alternatively, TILs can be functionally defined by their ability to infiltrate solid tumors upon reintroduction into a patient. TILS may further be characterized by potency – for example, TILS may be considered potent if, for example, interferon (IFN) release is greater than about 50 pg/mL, greater than about 100 pg/mL, greater than about 150 pg/mL, or greater than about 200 pg/mL.

[00478] By "population of cells" (including TILs) herein is meant a number of cells that share common traits. In general, populations generally range from 1×10^6 to 1×10^{10} in number, with different TIL populations comprising different numbers. For example, initial growth of primary TILs in the presence of IL-2 results in a population of bulk TILs of roughly 1×10^8 cells. REP expansion is generally done to provide populations of 1.5×10^9 to 1.5×10^{10} cells for infusion.

[00479] The term "pharmaceutically acceptable salt" refers to salts derived from a variety of organic and inorganic counter ions known in the art. Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Preferred inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid and phosphoric acid. Preferred organic acids from which salts

can be derived include, for example, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid and salicylic acid. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese and aluminum. Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins. Specific examples include isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine. In some embodiments, the pharmaceutically acceptable base addition salt is chosen from ammonium, potassium, sodium, calcium, and magnesium salts. The term "cocrystal" refers to a molecular complex derived from a number of cocrystal formers known in the art. Unlike a salt, a cocrystal typically does not involve hydrogen transfer between the cocrystal and the drug, and instead involves intermolecular interactions, such as hydrogen bonding, aromatic ring stacking, or dispersive forces, between the cocrystal former and the drug in the crystal structure.

[00480] The terms "pharmaceutically acceptable carrier" or "pharmaceutically acceptable excipient" are intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and inert ingredients. The use of such pharmaceutically acceptable carriers or pharmaceutically acceptable excipients for active pharmaceutical ingredients is well known in the art. Except insofar as any conventional pharmaceutically acceptable carrier or pharmaceutically acceptable excipient is incompatible with the active pharmaceutical ingredient, its use in the therapeutic compositions of the invention is contemplated. Additional active pharmaceutical ingredients, such as other drugs, can also be incorporated into the described compositions and methods.

[00481] The terms "about" and "approximately" mean within a statistically meaningful range of a value. Such a range can be within an order of magnitude, preferably within 50%, more preferably within 20%, more preferably still within 10%, and even more preferably within 5% of a given value or range. The allowable variation encompassed by the terms "about" or "approximately" depends on the particular system under study, and can be readily appreciated by one of ordinary skill in the art. Moreover, as used herein, the terms "about"

and "approximately" mean that dimensions, sizes, formulations, parameters, shapes and other quantities and characteristics are not and need not be exact, but may be approximate and/or larger or smaller, as desired, reflecting tolerances, conversion factors, rounding off, measurement error and the like, and other factors known to those of skill in the art. In general, a dimension, size, formulation, parameter, shape or other quantity or characteristic is "about" or "approximate" whether or not expressly stated to be such. It is noted that embodiments of very different sizes, shapes and dimensions may employ the described arrangements.

[00482] The transitional terms "comprising," "consisting essentially of," and "consisting of," when used in the appended claims, in original and amended form, define the claim scope with respect to what unrecited additional claim elements or steps, if any, are excluded from the scope of the claim(s). The term "comprising" is intended to be inclusive or open-ended and does not exclude any additional, unrecited element, method, step or material. The term "consisting of" excludes any element, step or material other than those specified in the claim and, in the latter instance, impurities ordinary associated with the specified material(s). The term "consisting essentially of" limits the scope of a claim to the specified elements, steps or material(s) and those that do not materially affect the basic and novel characteristic(s) of the claimed invention. All compositions, methods, and kits described herein that embody the present invention can, in alternate embodiments, be more specifically defined by any of the transitional terms "comprising," "consisting essentially of," and "consisting of."

III. TIL Manufacturing Processes – Process 2A (Gen 2 Processes)

[00483] In some embodiments, the invention provides the method of any of GEN 2 processes (*e.g.* process 2A processes) modified to use a small biopsy, core biopsy or fine needle aspirate as the source of T cells for expansion in the first expansion of any such GEN 2 process, wherein (1) the duration of the first expansion is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the second expansion of such GEN 2 process or (2) the duration of the second expansion of such GEN 2 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the second expansion or (3) the duration of the first expansion is lengthened and the duration of the

second expansion of such GEN 2 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the second expansion.

[00484] In some embodiments, the invention provides a population of TILs made by the method of any of GEN 2 processes (*e.g.* process 2A processes) modified to use a small biopsy, core biopsy or fine needle aspirate as the source of T cells for expansion in the first expansion of any such GEN 2 process, wherein (1) the duration of the first expansion is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the second expansion of such GEN 2 process or (2) the duration of the second expansion of such GEN 2 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the second expansion or (3) the duration of the first expansion is lengthened and the duration of the the second expansion of such GEN 2 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the second expansion.

[00485] Exemplary TIL processes known as process 2A and the small biopsy process containing some of these features is depicted in Figure 7, and some of the advantages of this embodiment of the present invention over process 1C are described in Figures 5 and 6. An embodiment of process 2A is shown Figure 1. Moreover, an overview as well as a comparison to process 2A with regard to the exemplary core biopsy process is provide in Figure 7. Process 2A (Gen 2), methods of treatment using TILs from process 2A, and compositions of TILs prepared by process 2A can be used in conjunction with small biopsy/core biopsy with durations of expansion periods adjusted as needed to achieve necessary cell counts, as provided herein below and throughout the present application.

[00486] As discussed herein, the present invention can include a step relating to the restimulation of cryopreserved TILs to increase their metabolic activity and thus relative health prior to transplant into a patient, and methods of testing said metabolic health. As generally outlined herein, TILs are generally taken from a patient sample and manipulated to expand their number prior to transplant into a patient. In some embodiments, the TILs may be optionally genetically manipulated as discussed below.

[00487] In some embodiments, the TILs may be cryopreserved. Once thawed, they may also be restimulated to increase their metabolism prior to infusion into a patient.

[00488] In some embodiments, the first expansion (including processes referred to as the pre-REP as well as processes shown in Figure 7 as Step A) is 1-19 days and the second expansion (including processes referred to as the REP as well as processes shown in Figure 7

as Step B) is shorted to 11-14 days, as discussed in detail below as well as in the examples and figures. In some embodiments, the first expansion (for example, an expansion described as Step B in Figure 7) is split into two periods of 1-2 days and 3-19, or in some cases 1-2 days and 3-10 days and the second expansion (for example, an expansion as described in Step D in Figure 7) is 11-14 days, as discussed in the Examples and shown in Figures 4, 5, and 7. In some embodiments, the combination of the first expansion and second expansion (for example, expansions described as Step B and Step D in Figure 7) is shortened to 22 days, as discussed in detail below and in the examples and figures.

[00489] The "Step" Designations A, B, C, *etc.*, below are in reference to Figure 7. The ordering of the Steps below and in Figure 7 is exemplary and any combination or order of steps, as well as additional steps, repetition of steps, and/or omission of steps is contemplated by the present application and the methods disclosed herein.

A. STEP A: Obtain Patient tumor sample

[00490] In general, TILs are initially obtained from a patient tumor sample ("primary TILs") obtained by a core biopsy or similar procedure and then expanded into a larger population for further manipulation as described herein, optionally cryopreserved, and optionally evaluated for phenotype and metabolic parameters.

[00491] A patient tumor sample may be obtained using methods known in the art, generally via small biopsy, core biopsy, needle biopsy or other means for obtaining a sample that contains a mixture of tumor and TIL cells. In general, the tumor sample may be from any solid tumor, including primary tumors, invasive tumors or metastatic tumors. The tumor sample may also be a liquid tumor, such as a tumor obtained from a hematological malignancy. In some embodiments, the sample can be from multiple small tumor samples or biopsies. In some embodiments, the sample can comprise multiple tumor samples from a single tumor from the same patient. In some embodiments, the sample can comprise multiple tumor samples from one, two, three, or four tumors from the same patient. In some embodiments, the sample can comprise multiple tumor samples from multiple tumors from the same patient. The solid tumor may be of any cancer type, including, but not limited to, breast, pancreatic, prostate, colorectal, lung, brain, renal, stomach, and skin (including but not limited to squamous cell carcinoma, basal cell carcinoma, and melanoma). In some embodiments, the cancer is selected from cervical cancer, head and neck cancer (including,

for example, head and neck squamous cell carcinoma (HNSCC)), glioblastoma (GBM), gastrointestinal cancer, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma (NSCLC). In some embodiments, useful TILs are obtained from malignant melanoma tumors, as these have been reported to have particularly high levels of TILs.

[00492] In general, the cell suspension obtained from the tumor core or fragment is called a "primary cell population" or a "freshly obtained" or a "freshly isolated" cell population. In certain embodiments, the freshly obtained cell population of TILs is exposed to a cell culture medium comprising antigen presenting cells, IL-2 and OKT-3.

[00493] In some embodiments, if the tumor is metastatic and the primary lesion has been efficiently treated/removed in the past, removal of one of the metastatic lesions may be needed. In some embodiments, the least invasive approach is to remove a skin lesion, or a lymph node on the neck or axillary area when available. In some embodiments, a skin lesion is removed or small biopsy thereof is removed. In some embodiments, a lymph node or small biopsy thereof is removed. In some embodiments, a lung or liver metastatic lesion, or an intra-abdominal or thoracic lymph node or small biopsy can thereof can be employed.

[00494] In some embodiments, the tumor is a melanoma. In some embodiments, the small biopsy for a melanoma comprises a mole or portion thereof.

[00495] In some embodiments, the small biopsy is a punch biopsy. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin. around a suspicious mole. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin, and a round piece of skin is removed. In some embodiments, the small biopsy is a punch biopsy and round portion of the tumor is removed.

[00496] In some embodiments, the small biopsy is an excisional biopsy. In some embodiments, the small biopsy is an excisional biopsy and the entire mole or growth is removed. In some embodiments, the small biopsy is an excisional biopsy and the entire mole or growth is removed along with a small border of normal-appearing skin.

[00497] In some embodiments, the small biopsy is an incisional biopsy. In some embodiments, the small biopsy is an incisional biopsy and only the most irregular part of a mole or growth is taken. In some embodiments, the small biopsy is an incisional biopsy and

the incisional biopsy is used when other techniques can't be completed, such as if a suspicious mole is very large.

[00498] In some embodiments, the small biopsy is a lung biopsy. In some embodiments, the small biopsy is obtained by bronchoscopy. Generally, bronchoscopy, the patient is put under anesthesia, and a small tool goes through the nose or mouth, down the throat, and into the bronchial passages, where small tools are used to remove some tissue. In some embodiments, where the tumor or growth cannot be reached via bronchoscopy, a transthoracic needle biopsy can be employed. Generally, for a transthoracic needle biopsy, the patient is also under anesthesia and a needle is inserted through the skin directly into the suspicious spot to remove a small sample of tissue. In some embodiments, a transthoracic needle biopsy may require interventional radiology (for example, the use of x-rays or CT scan to guide the needle). In some embodiments, the small biopsy is obtained by needle biopsy. In some embodiments, the small biopsy is obtained endoscopic ultrasound (for example, an endoscope with a light and is placed through the mouth into the esophagus). In some embodiments, the small biopsy is obtained surgically.

[00499] In some embodiments, the small biopsy is a head and neck biopsy. In some embodiments, the small biopsy is an incisional biopsy. In some embodiments, the small biopsy is an incisional biopsy, wherein a small piece of tissue is cut from an abnormal-looking area. In some embodiments, if the abnormal region is easily accessed, the sample may be taken without hospitalization. In some embodiments, if the tumor is deeper inside the mouth or throat, the biopsy may need to be done in an operating room, with general anesthesia. In some embodiments, the small biopsy is an excisional biopsy. In some embodiments, the small biopsy is an excisional biopsy, wherein the whole area is removed. In some embodiments, the small biopsy is a fine needle aspiration (FNA). In some embodiments, the small biopsy is a fine needle aspiration (FNA), wherein a very thin needle attached to a syringe is used to extract (aspirate) cells from a tumor or lump. In some embodiments, the small biopsy is a punch biopsy. In some embodiments, the small biopsy is a punch biopsy, wherein punch forceps are used to remove a piece of the suspicious area.

[00500] In some embodiments, the small biopsy is a cervical biopsy. In some embodiments, the small biopsy is obtained via colposcopy. Generally, colposcopy methods employ the use of a lighted magnifying instrument attached to magnifying binoculars (a colposcope) which is then used to biopsy a small section of the surface of the cervix. In some embodiments, the small biopsy is a conization/cone biopsy. In some embodiments, the small biopsy is a

conization/cone biopsy, wherein an outpatient surgery may be needed to remove a larger piece of tissue from the cervix. In some embodiments, the cone biopsy, in addition to helping to confirm a diagnosis, a cone biopsy can serve as an initial treatment.

[00501] The term "solid tumor" refers to an abnormal mass of tissue that usually does not contain cysts or liquid areas. Solid tumors may be benign or malignant. The term "solid tumor cancer refers to malignant, neoplastic, or cancerous solid tumors. Solid tumor cancers include, but are not limited to, sarcomas, carcinomas, and lymphomas, such as cancers of the lung, breast, triple negative breast cancer, prostate, colon, rectum, and bladder. In some embodiments, the cancer is selected from cervical cancer, head and neck cancer, glioblastoma, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma. The tissue structure of solid tumors includes interdependent tissue compartments including the parenchyma (cancer cells) and the supporting stromal cells in which the cancer cells are dispersed and which may provide a supporting microenvironment.

[00502] In some embodiments, the sample from the tumor is obtained as a fine needle aspirate (FNA), a core biopsy, a small biopsy (including, for example, a punch biopsy). In some embodiments, sample is placed first into a G-Rex 10. In some embodiments, sample is placed first into a G-Rex 10 when there are 1 or 2 core biopsy and/or small biopsy samples. In some embodiments, sample is placed first into a G-Rex 100 when there are 3, 4, 5, 6, 8, 9, or 10 or more core biopsy and/or small biopsy samples. In some embodiments, sample is placed first into a G-Rex 500 when there are 3, 4, 5, 6, 8, 9, or 10 or more core biopsy and/or small biopsy samples.

[00503] The FNA can be obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma. In some embodiments, the FNA is obtained from a lung tumor, such as a lung tumor from a patient with non-small cell lung cancer (NSCLC). In some cases, the patient with NSCLC has previously undergone a surgical treatment.

[00504] TILs described herein can be obtained from an FNA sample. In some cases, the FNA sample is obtained or isolated from the patient using a fine gauge needle ranging from an 18 gauge needle to a 25 gauge needle. The fine gauge needle can be 18 gauge, 19 gauge, 20 gauge, 21 gauge, 22 gauge, 23 gauge, 24 gauge, or 25 gauge. In some embodiments, the FNA sample from the patient can contain at least 400,000 TILs, *e.g.*,

400,000 TILs, 450,000 TILs, 500,000 TILs, 550,000 TILs, 600,000 TILs, 650,000 TILs, 700,000 TILs, 750,000 TILs, 800,000 TILs, 850,000 TILs, 900,000 TILs, 950,000 TILs, or more.

[00505] In some cases, the TILs described herein are obtained from a core biopsy sample. In some cases, the core biopsy sample is obtained or isolated from the patient using a surgical or medical needle ranging from an 11 gauge needle to a 16 gauge needle. The needle can be 11 gauge, 12 gauge, 13 gauge, 14 gauge, 15 gauge, or 16 gauge. In some embodiments, the core biopsy sample from the patient can contain at least 400,000 TILs, *e.g.*, 400,000 TILs, 450,000 TILs, 500,000 TILs, 550,000 TILs, 600,000 TILs, 650,000 TILs, 700,000 TILs, 750,000 TILs, 800,000 TILs, 850,000 TILs, 900,000 TILs, 950,000 TILs, or more.

[00506] In general, the harvested cell suspension is called a "primary cell population" or a "freshly harvested" cell population.

[00507] In some embodiments, the TILs are not obtained from tumor digests. In some embodiments, the solid tumor cores are not fragmented.

[00508] In some embodiments, the TILs are obtained from tumor digests. In some embodiments, tumor digests were generated by incubation in enzyme media, for example but not limited to RPMI 1640, 2mM GlutaMAX, 10 mg/mL gentamicin, 30 U/mL DNase, and 1.0 mg/mL collagenase, followed by mechanical dissociation (GentleMACS, Miltenyi Biotec, Auburn, CA). After placing the tumor in enzyme media, the tumor can be mechanically dissociated for approximately 1 minute. The solution can then be incubated for 30 minutes at 37 °C in 5% CO_2 and it then mechanically disrupted again for approximately 1 minute. After being incubated again for 30 minutes at 37 °C in 5% CO_2, the tumor can be mechanically disrupted a third time for approximately 1 minute. In some embodiments, after the third mechanical disruption if large pieces of tissue were present, 1 or 2 additional mechanical dissociations were applied to the sample, with or without 30 additional minutes of incubation at 37 °C in 5% CO_2. In some embodiments, at the end of the final incubation if the cell suspension contained a large number of red blood cells or dead cells, a density gradient separation using Ficoll can be performed to remove these cells.

[00509] In some embodiments, the harvested cell suspension prior to the first expansion step is called a "primary cell population" or a "freshly harvested" cell population.

[00510] In some embodiments, cells can be optionally frozen after sample harvest and stored frozen prior to entry into the expansion described in Step B, which is described in further detail below, as well as exemplified in Figure 7.

B. STEP B: First Expansion

[00511] In some embodiments, the present methods provide for obtaining young TILs, which are capable of increased replication cycles upon administration to a subject/patient and as such may provide additional therapeutic benefits over older TILs (*i.e.*, TILs which have further undergone more rounds of replication prior to administration to a subject/patient). Features of young TILs have been described in the literature, for example Donia, *et al.*, *Scandinavian Journal of Immunology*, 75:157–167 (2012); Dudley *et al.*, *Clin Cancer Res*, 16:6122-6131 (2010); Huang *et al.*, *J Immunother*, 28(3):258–267 (2005); Besser *et al.*, *Clin Cancer Res*, 19(17):OF1-OF9 (2013); Besser et al., *J Immunother* 32:415–423 (2009); Robbins, *et al.*, *J Immunol* 2004; 173:7125-7130; Shen *et al.*, J Immunother, 30:123–129 (2007); Zhou, *et al.*, *J Immunother*, 28:53–62 (2005); and Tran, *et al.*, J Immunother, 31:742–751 (2008), all of which are incorporated herein by reference in their entireties.

[00512] The diverse antigen receptors of T and B lymphocytes are produced by somatic recombination of a limited, but large number of gene segments. These gene segments: V (variable), D (diversity), J (joining), and C (constant), determine the binding specificity and downstream applications of immunoglobulins and T-cell receptors (TCRs). The present invention provides a method for generating TILs which exhibit and increase the T-cell repertoire diversity. In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity. In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity as compared to freshly harvested TILs and/or TILs prepared using other methods than those provide herein including for example, methods other than those embodied in Figure 1. In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity as compared to freshly harvested TILs and/or TILs prepared using methods referred to as process 1C, as exemplified in Figure 3 and/or Figure 4. In some embodiments, the TILs obtained in the first expansion exhibit an increase in the T-cell repertoire diversity. In some embodiments, the increase in diversity is an increase in the immunoglobulin diversity and/or the T-cell receptor diversity. In some embodiments, the diversity is in the immunoglobulin is in the immunoglobulin heavy chain. In some

embodiments, the diversity is in the immunoglobulin is in the immunoglobulin light chain. In some embodiments, the diversity is in the T-cell receptor. In some embodiments, the diversity is in one of the T-cell receptors selected from the group consisting of alpha, beta, gamma, and delta receptors. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) alpha and/or beta. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) alpha. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) beta. In some embodiments, there is an increase in the expression of TCRab (*i.e.*, TCRα/β).

[00513] After obtaining the small biopsy, core biopsy or fine needle aspirate tumor fragment or fragments (which may be referred to as "cores" or "fragments"), for example such as described in Step A of Figure 7, the resulting cells are cultured in serum containing IL-2 under conditions that favor the growth of TILs over tumor and other cells. In some embodiments, the tumor digests are incubated in 2 mL wells in media comprising inactivated human AB serum with 6000 IU/mL of IL-2. This primary cell population is cultured for a period of days, generally from 3 to 14 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, this primary cell population is cultured for a period of 7 to 14 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, this primary cell population is cultured for a period of 1 to 149 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, this primary cell population is cultured for a period of about 3-19 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, IL-2 is included during days 1-2 of the first expansion and OKT3 and feeders are added at day 3 of the first expansion. In some embodiments, the OKT3 and feeders are included for days 3-19. In some embodiments, the OKT3 and feeders are included for days 3-18. In some embodiments, the OKT3 and feeders are included for days 3-17. In some embodiments, the OKT3 and feeders are included for days 3-16. In some embodiments, the OKT3 and feeders are included for days 3-15. In some embodiments, the OKT3 and feeders are included for days 3-14. In some embodiments, the OKT3 and feeders are included for days 3-13. In some embodiments, the OKT3 and feeders are included for days 3-12. In some embodiments, the OKT3 and feeders are included for days 3-11. In some embodiments, the OKT3 and feeders are included for days 3-10. In some embodiments, the OKT3 and feeders are included for days 3-9. In a preferred embodiment, expansion of TILs may be performed using an initial bulk TIL expansion step (for example such as those described in Step B of

Figure 7, which can include processes referred to as pre-REP) as described below and herein, followed by a second expansions (Step D, including processes referred to as rapid expansion protocol (REP) steps) as described below under Step D and herein, followed by optional cryopreservation, and followed by a second Step D (including processes referred to as restimulation REP steps) as described below and herein. The TILs obtained from this process may be optionally characterized for phenotypic characteristics and metabolic parameters as described herein.

[00514] In embodiments where TIL cultures are initiated in 24-well plates, for example, using Costar 24-well cell culture cluster, flat bottom (Corning Incorporated, Corning, NY, each well can be seeded with 1×10^6 small biopsy, core biopsy, or fine needle aspirate tumor cells or one small biopsy, core biopsy, or fine needle aspirate tumor fragment in 2 mL of complete medium (CM) with IL-2 (6000 IU/mL; Chiron Corp., Emeryville, CA).

[00515] In some embodiments, the first expansion culture medium is referred to as "CM", an abbreviation for culture media. In some embodiments, CM for Step B consists of RPMI 1640 with GlutaMAX, supplemented with 10% human AB serum, 25 mM Hepes, and 10 mg/mL gentamicin. In embodiments where cultures are initiated in gas-permeable flasks with a 40 mL capacity and a 10 cm^2 gas-permeable silicon bottom (for example, G-Rex10; Wilson Wolf Manufacturing, New Brighton, MN) (Fig. 1), each flask was loaded with $10–40 \times 10^6$ viable small biopsy, core biopsy, or fine needle aspirate tumor cells or 5–30 small biopsy, core biopsy or fine needle aspirate tumor fragments in 10–40 mL of CM with IL-2. Both the G-Rex10 and 24-well plates were incubated in a humidified incubator at 37°C in 5% CO_2 and 5 days after culture initiation, half the media was removed and replaced with fresh CM and IL-2 and after day 5, half the media was changed every 2–3 days.

[00516] After harvest of the small biopsy, core biopsy or fine needle aspirate tumor fragments, the resulting cells (*i.e.*, fragments) are cultured in serum containing IL-2 under conditions that favor the growth of TILs over tumor and other cells. In some embodiments, the tumor fragments are incubated in 2 mL wells in media comprising inactivated human AB serum (or, in some cases, as outlined herein, in the presence of aAPC cell population) with 6000 IU/mL of IL-2. This primary cell population is cultured for a period of days, generally from 10 to 14 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, the growth media during the first expansion comprises IL-2 or a variant thereof. In some embodiments, the IL is recombinant human IL-2 (rhIL-2). In some embodiments the IL-2 stock solution has a specific activity of $20\text{-}30 \times 10^6$ IU/mg for a 1 mg

vial. In some embodiments the IL-2 stock solution has a specific activity of 20-30 × 10^6 IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 25 × 10^6 IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 30 × 10^6 IU/mg for a 1 mg vial. In some embodiments, the IL- 2 stock solution has a final concentration of 4-8 × 10^6 IU/mg of IL-2. In some embodiments, the IL- 2 stock solution has a final concentration of 5-7 × 10^6 IU/mg of IL-2. In some embodiments, the IL-2 stock solution has a final concentration of 6 × 10^6 IU/mg of IL-2. In some embodiments, the IL-2 stock solution is prepare as described in Example 4. In some embodiments, first expansion culture media comprises about 10,000 IU/mL of IL-2, about 9,000 IU/mL of IL-2, about 8,000 IU/mL of IL-2, about 7,000 IU/mL of IL-2, about 6000 IU/mL of IL-2 or about 5,000 IU/mL of IL-2. In some embodiments, first expansion culture media comprises about 9,000 IU/mL of IL-2, to about 5,000 IU/mL of IL-2. In some embodiments, first expansion culture media comprises about 8,000 IU/mL of IL-2, to about 6,000 IU/mL of IL-2. In some embodiments, first expansion culture media comprises about 7,000 IU/mL of IL-2, to about 6,000 IU/mL of IL-2. In some embodiments, first expansion culture media comprises about 6,000 IU/mL of IL-2. In an embodiment, the cell culture medium further comprises IL-2. In some embodiments, the cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the cell culture medium further comprises IL-2. In a preferred embodiment, the cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises about 1000 IU/mL, about 1500 IU/mL, about 2000 IU/mL, about 2500 IU/mL, about 3000 IU/mL, about 3500 IU/mL, about 4000 IU/mL, about 4500 IU/mL, about 5000 IU/mL, about 5500 IU/mL, about 6000 IU/mL, about 6500 IU/mL, about 7000 IU/mL, about 7500 IU/mL, or about 8000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises between 1000 and 2000 IU/mL, between 2000 and 3000 IU/mL, between 3000 and 4000 IU/mL, between 4000 and 5000 IU/mL, between 5000 and 6000 IU/mL, between 6000 and 7000 IU/mL, between 7000 and 8000 IU/mL, or about 8000 IU/mL of IL-2.

[00517] In some embodiments, the first expansion is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody. For instance, the cell culture medium of the second population of TILs is supplemented with OKT3, IL-15, an OX40 agonistic antibody, a 4-1BB agonistic antibody, a combination of OKT3 and IL-15, a combination of OKT3 and an OX40 agonistic antibody, a combination of OKT3 and a 4-1BB agonistic antibody, a combination of IL-15 and an OX40 agonistic antibody, a combination

of IL-15, and a 4-1BB agonistic antibody, a combination of an OX40 agonistic antibody and a 4-1BB agonistic antibody, a combination of OKT3, IL-15, and an OX40 agonistic antibody, a combination of OKT3, IL-15, and a 4-1BB agonistic antibody, a combination of OKT3, an OX40 agonistic antibody, and a 4-1BB agonistic antibody, a combination of IL-15, an OX40 agonistic antibody, and a 4-1BB agonistic antibody, a combination of OKT3, IL-15, an OX40 agonistic antibody, and a 4-1BB agonistic antibody, and any combination thereof. In some embodiments, IL-15 is not included.

[00518] After obtaining the small biopsy, core biopsy or fine needle aspirate tumor fragments, for example such as described in Step A of Figure 7A, the resulting cells are cultured in serum containing IL-2 under conditions that favor the growth of TILs over tumor and other cells. In some embodiments, the tumor fragments are incubated in 2 mL wells in media comprising inactivated human AB serum with 6000 IU/mL of IL-2. This primary cell population is cultured for a period of days, generally from 1 to 19 days, resulting in a TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 11 to day 19. In some embodiments, this primary cell population is cultured for a period of 1 to 18 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 18. In some embodiments, this primary cell population is cultured for a period of 1 to 17 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 17. In some embodiments, this primary cell population is cultured for a period of 1 to 16 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 16. In some embodiments, this primary cell population is cultured for a period of 1 to 15 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 15. In some embodiments, this primary cell population is cultured for a period of 1 to 14 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 14. In some embodiments, this primary cell population is cultured for a period of 1 to 13 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 13. In some embodiments, this primary cell population is cultured for a period of 1 to 12 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 12. In some embodiments, this primary cell population is cultured for a period of 1 to 11 days, resulting in a bulk TIL population of at least 5×10^7 (*i.e.*, 50 million) TIL cells by at least day 11. In some embodiments, TIL population is at least about 5×10^7 (*i.e.*, 50 million) TIL cells by the end of the first expansion, step (for example such as those described in Step B of Figure 7, which can include processes referred to as pre-REP).

[00519] In a preferred embodiment, expansion of TILs may be performed using an initial bulk TIL expansion step (for example such as those described in Step B of Figure7, which can include processes referred to as pre-REP) as described below and herein, followed by a second expansion (Step D, including processes referred to as rapid expansion protocol (REP) steps) as described below under Step D and herein, followed by optional cryopreservation, and followed by a second Step D (including processes referred to as restimulation REP steps) as described below and herein. The TILs obtained from this process may be optionally characterized for phenotypic characteristics and metabolic parameters as described herein.

[00520] In some embodiments, the first expansion culture medium is referred to as "CM", an abbreviation for culture media. In some embodiments, CM for Step B consists of RPMI 1640 with GlutaMAX, supplemented with 10% human AB serum, 25 mM Hepes, and 10 mg/mL gentamicin. In embodiments where cultures are initiated in gas-permeable flasks with a 40 mL capacity and a 10 cm^2 gas-permeable silicon bottom (for example, G-Rex10; Wilson Wolf Manufacturing, New Brighton, MN), each flask was loaded with 1-2 biopsy cores in 10–40 mL of CM with IL-2. Both the G-Rex10 and 24-well plates were incubated in a humidified incubator at 37°C in 5% CO_2 and 5 days after culture initiation, half the media was removed and replaced with fresh CM and IL-2 and after day 2, half the media was changed every 2–3 days. In some embodiments, when the core biopsy is from an ovarian tumor (*e.g.*, of ovarian origin), cores are incubated in CM and IL-2 for 1-2 days. In some embodiments, when the core biopsy is from a pancreatic tumor cells (*e.g.*, of pancreatic origin), cores are incubated in CM and IL-2/IL-15/IL-21 for 1-2 days. In some embodiments, at day 3, OKT3 and feeder cells are added to the cores. In some embodiments, at day 3, 30 ng OKT3 and 10^6 feeder cells are added to the cores.

[00521] The small biopsy fragments or cores are cultured in serum containing IL-2 under conditions that favor the growth of TILs over tumor and other cells. In some embodiments, the small biopsy fragments or cores are incubated in 2 mL wells in media comprising inactivated human AB serum (or, in some cases, as outlined herein, in the presence of aAPC cell population) with 6000 IU/mL of IL-2. This primary cell population is cultured for a period of days, generally from 1 to 19 days, resulting in a bulk TIL population, of at least about 5×10^7 TIL cells at day 11, and in some embodiments, 1×10^8 TILs at days 19 through 28. In some embodiments, the growth media during the first expansion comprises IL-2 or a variant thereof. In some embodiments, the IL is recombinant human IL-2 (rhIL-2). In some embodiments the IL-2 stock solution has a specific activity of 20-30$\times10^6$ IU/mg for a 1 mg

vial. In some embodiments the IL-2 stock solution has a specific activity of 20×10^6 IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 25×10^6 IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 30×10^6 IU/mg for a 1 mg vial. In some embodiments, the IL- 2 stock solution has a final concentration of 4-8×10^6 IU/mg of IL-2. In some embodiments, the IL- 2 stock solution has a final concentration of 5-7×10^6 IU/mg of IL-2. In some embodiments, the IL- 2 stock solution has a final concentration of 6×10^6 IU/mg of IL-2. In some embodiments, the IL-2 stock solution is prepare as described in Example E. In some embodiments, the first expansion culture media comprises about 10,000 IU/mL of IL-2, about 9,000 IU/mL of IL-2, about 8,000 IU/mL of IL-2, about 7,000 IU/mL of IL-2, about 6000 IU/mL of IL-2 or about 5,000 IU/mL of IL-2. In some embodiments, the first expansion culture media comprises about 9,000 IU/mL of IL-2 to about 5,000 IU/mL of IL-2. In some embodiments, the first expansion culture media comprises about 8,000 IU/mL of IL-2 to about 6,000 IU/mL of IL-2. In some embodiments, the first expansion culture media comprises about 7,000 IU/mL of IL-2 to about 6,000 IU/mL of IL-2. In some embodiments, the first expansion culture media comprises about 6,000 IU/mL of IL-2. In an embodiment, the cell culture medium further comprises IL-2. In some embodiments, the cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the cell culture medium further comprises IL-2. In a preferred embodiment, the cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises about 1000 IU/mL, about 1500 IU/mL, about 2000 IU/mL, about 2500 IU/mL, about 3000 IU/mL, about 3500 IU/mL, about 4000 IU/mL, about 4500 IU/mL, about 5000 IU/mL, about 5500 IU/mL, about 6000 IU/mL, about 6500 IU/mL, about 7000 IU/mL, about 7500 IU/mL, or about 8000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises between 1000 and 2000 IU/mL, between 2000 and 3000 IU/mL, between 3000 and 4000 IU/mL, between 4000 and 5000 IU/mL, between 5000 and 6000 IU/mL, between 6000 and 7000 IU/mL, between 7000 and 8000 IU/mL, or about 8000 IU/mL of IL-2.

[00522] In some embodiments, first expansion culture media comprises about 500 IU/mL of IL-15, about 400 IU/mL of IL-15, about 300 IU/mL of IL-15, about 200 IU/mL of IL-15, about 180 IU/mL of IL-15, about 160 IU/mL of IL-15, about 140 IU/mL of IL-15, about 120 IU/mL of IL-15, or about 100 IU/mL of IL-15. In some embodiments, the first expansion culture media comprises about 500 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the first expansion culture media comprises about 400 IU/mL of IL-15 to about

100 IU/mL of IL-15. In some embodiments, the first expansion culture media comprises about 300 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the first expansion culture media comprises about 200 IU/mL of IL-15. In some embodiments, the cell culture medium comprises about 180 IU/mL of IL-15. In an embodiment, the cell culture medium further comprises IL-15. In a preferred embodiment, the cell culture medium comprises about 180 IU/mL of IL-15. In some embodiments, IL-15 is included when the cores are from a pancreatic tumor (*e.g.*, of pancreatic origin).

[00523] In some embodiments, first expansion culture media comprises about 20 IU/mL of IL-21, about 15 IU/mL of IL-21, about 12 IU/mL of IL-21, about 10 IU/mL of IL-21, about 5 IU/mL of IL-21, about 4 IU/mL of IL-21, about 3 IU/mL of IL-21, about 2 IU/mL of IL-21, about 1 IU/mL of IL-21, or about 0.5 IU/mL of IL-21. In some embodiments, the first expansion culture media comprises about 20 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the first expansion culture media comprises about 15 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the first expansion culture media comprises about 12 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the first expansion culture media comprises about 10 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the first expansion culture media comprises about 5 IU/mL of IL-21 to about 1 IU/mL of IL-21. In some embodiments, the first expansion culture media comprises about 2 IU/mL of IL-21. In some embodiments, the cell culture medium comprises about 1 IU/mL of IL-21. In some embodiments, the cell culture medium comprises about 0.5 IU/mL of IL-21. In an embodiment, the cell culture medium further comprises IL-21. In a preferred embodiment, the cell culture medium comprises about 1 IU/mL of IL-21. In some embodiments, IL-21 is included when the cores are from a pancreatic tumor (*e.g.*, of pancreatic origin).

[00524] In an embodiment, the cell culture medium comprises OKT-3 antibody. In some embodiments, the cell culture medium comprises about 30 ng/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises about 0.1 ng/mL, about 0.5 ng/mL, about 1 ng/mL, about 2.5 ng/mL, about 5 ng/mL, about 7.5 ng/mL, about 10 ng/mL, about 15 ng/mL, about 20 ng/mL, about 25 ng/mL, about 30 ng/mL, about 35 ng/mL, about 40 ng/mL, about 50 ng/mL, about 60 ng/mL, about 70 ng/mL, about 80 ng/mL, about 90 ng/mL, about 100 ng/mL, about 200 ng/mL, about 500 ng/mL, and about 1 μg/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises between 0.1 ng/mL and 1 ng/mL, between 1 ng/mL and 5 ng/mL, between 5 ng/mL and 10 ng/mL, between 10 ng/mL and 20 ng/mL,

between 20 ng/mL and 30 ng/mL, between 30 ng/mL and 40 ng/mL, between 40 ng/mL and 50 ng/mL, and between 50 ng/mL and 100 ng/mL of OKT-3 antibody. In some embodiments, the cell culture medium does not comprise OKT-3 antibody. In some embodiments, the OKT-3 antibody is muromonab.

TABLE 3: Amino acid sequences of muromonab (exemplary OKT-3 antibody)

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:1 Muromonab heavy chain	QVQLQQSGAE LARPGASVKM SCKASGYTFT RYTMHWVKQR PGQGLEWIGY INPSRGYTNY NQKFKDKATL TTDKSSSTAY MQLSSLTSED SAVYYCARYY DDHYCLDYWG QGTTLTVSSA KTTAPSVYPL APVCGGTTGS SVTLGCLVKG YFPEPVTLTW NSGSLSSGVH TFPAVLQSDL YTLSSSVTVT SSTWPSQSIT CNVAHPASST KVDKKIEPRP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK	60 120 180 240 300 360 420 450
SEQ ID NO:2 Muromonab light chain	QIVLTQSPAI MSASPGEKVT MTCSASSSVS YMNWYQQKSG TSPKRWIYDT SKLASGVPAH FRGSGSGTSY SLTISGMEAE DAATYYCQQW SSNPFTFGSG TKLEINRADT APTVSIFPPS SEQLTSGGAS VVCFLNNFYP KDINVKWKID GSERQNGVLN SWTDQDSKDS TYSMSSTLTL TKDEYERHNS YTCEATHKTS TSPIVKSFNR NEC	60 120 180 213

[00525] In some embodiments, the cell culture medium comprises one or more TNFRSF agonists in a cell culture medium. In some embodiments, the TNFRSF agonist comprises a 4-1BB agonist. In some embodiments, the TNFRSF agonist is a 4-1BB agonist, and the 4-1BB agonist is selected from the group consisting of urelumab, utomilumab, EU-101, a fusion protein, and fragments, derivatives, variants, biosimilars, and combinations thereof. In some embodiments, the TNFRSF agonist is added at a concentration sufficient to achieve a concentration in the cell culture medium of between 0.1 μg/mL and 100 μg/mL. In some embodiments, the TNFRSF agonist is added at a concentration sufficient to achieve a concentration in the cell culture medium of between 20 μg/mL and 40 μg/mL.

[00526] In some embodiments, in addition to one or more TNFRSF agonists, the cell culture medium further comprises IL-2 at an initial concentration of about 3000 IU/mL and OKT-3 antibody at an initial concentration of about 30 ng/mL, and wherein the one or more TNFRSF agonists comprises a 4-1BB agonist.

[00527] In some embodiments, the first expansion culture medium is referred to as "CM", an abbreviation for culture media. In some embodiments, it is referred to as CM1 (culture medium 1). In some embodiments, CM consists of RPMI 1640 with GlutaMAX, supplemented with 10% human AB serum, 25 mM Hepes, and 10 mg/mL gentamicin. In embodiments where cultures are initiated in gas-permeable flasks with a 40 mL capacity and a 10 cm^2 gas-permeable silicon bottom (for example, G-Rex10; Wilson Wolf Manufacturing,

New Brighton, MN). In some embodiments, the CM is the CM1 described in the Examples, *see*, Example 1 In some embodiments, the first expansion occurs in an initial cell culture medium or a first cell culture medium. In some embodiments, the initial cell culture medium or the first cell culture medium comprises IL-2.

[00528] In some embodiments, the first expansion (including processes such as for example those described in Step B of Figure 7, which can include those sometimes referred to as the pre-REP) process is about 1-19 days, a time period sufficient to obtain 5×10^7 (*i.e.*, 50 million) TILs, as discussed in the examples and figures. In some embodiments, the first expansion (including processes such as for example those described in Step B of Figure 7, which can include those sometimes referred to as the pre-REP) is divided into two phases, 1-2 days and 3 to 19 days, as provided in Figure 7 and 38. In some embodiments, the first expansion of Step B is divided into two phases, 1-2 days (cultured with IL-2 or IL2/IL-15/IL-21) and 3 to 19 days (cultured with OKT3 and feeders), as provided in Figure 7 and 38.

[00529] In some embodiments, the first TIL expansion can proceed for 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 14 days, 14 days, 14 days, 14 days, 14 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 1 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 2 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 3 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 6 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 7 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 8 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 9 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 10 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 11 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 12 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 13 days to 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first

TIL expansion can proceed for 16 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 17 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 18 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the first TIL expansion can proceed for 19 days, in order to obtain at least 5×10^7 (*i.e.*, 50 million) TILs. In some embodiments, the small biopsy (*e.g.*, the core biopsy) is removed from the culture at about day 1, 2, or 3. In some embodiments, the small biopsy (*e.g.*, the core biopsy) is removed from the culture at day 3.

[00530] In some embodiments, a combination of IL-2, IL-7, IL-15, and/or IL-21 are employed as a combination during the first expansion. In some embodiments, IL-2, IL-7, IL-15, and/or IL-21 as well as any combinations thereof can be included during the first expansion, including for example during a Step B processes according to Figure 7, as well as described herein. In some embodiments, a combination of IL-2, IL-15, and IL-21 are employed as a combination during the first expansion. In some embodiments, IL-2, IL-15, and IL-21 as well as any combinations thereof can be included during Step B processes according to Figure 7 and as described herein. In some embodiments, IL-15 and/or IL-21 is added at day 1. In some embodiments, IL-15 and/or IL-21 is added at day 1 if the core biopsy is from a pancreatic tumor (*e.g.*, of pancreatic origin). In some embodiments, IL-15 is not included. In some embodiments, IL-21 is not included. In some embodiments, neither IL-15 nor IL-21 is not included.

[00531] In some embodiments, the first expansion, for example, Step B according to Figure 7, is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a single bioreactor is employed. In some embodiments, the single bioreactor employed is for example a GREX-10 or a GREX-100. In some embodiments, the closed system bioreactor is a single bioreactor.

[00532] In some embodiments, the first TIL expansion generates at least 50×10^6 TILs, *e.g.*, 50×10^6 (*i.e.*, 5×10^7), 55×10^6, 60×10^6, 65×10^6, 70×10^6, 75×10^6, 80×10^6, 85×10^6, 90×10^6, 95×10^6, 100×10^6, 105×10^6, 110×10^6, 115×10^6, 120×10^6, 125×10^6, 150×10^6, 200×10^6, 300×10^6, 400×10^6, or more.

C. STEP C: First Expansion to Second Expansion Transition

[00533] In some cases, the bulk TIL population obtained from the first expansion, including for example the TIL population obtained from for example, Step B as indicated in Figure 7, can be cryopreserved immediately, using the protocols discussed herein below. Alternatively, the TIL population obtained from the first expansion, referred to as the second TIL population, can be subjected to a second expansion (which can include expansions sometimes referred to as REP) and then cryopreserved as discussed below. Similarly, in the case where genetically modified TILs will be used in therapy, the first TIL population (sometimes referred to as the bulk TIL population) or the second TIL population (which can in some embodiments include populations referred to as the REP TIL populations) can be subjected to genetic modifications for suitable treatments prior to expansion or after the first expansion and prior to the second expansion.

[00534] In some embodiments, the TILs obtained from the first expansion (for example, from Step B as indicated in Figure 7) are stored until phenotyped for selection. In some embodiments, the TILs obtained from the first expansion (for example, from Step B as indicated in Figure 7) are not stored and proceed directly to the second expansion. In some embodiments, the TILs obtained from the first expansion are not cryopreserved after the first expansion and prior to the second expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 3 days, 4, days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days, 17 days, 18 days, or 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 3 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 4 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 4 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 7 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 10 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about

11 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 12 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 13 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 14 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 15 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 16 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 17 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 18 days to 19 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 10 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 11 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 12 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 13 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 14 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 15 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 16 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 17 days from when the cores are added to the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 18 days from when the cores are added to

the culture medium for the first expansion. In some embodiments, the transition from the first expansion to the second expansion occurs at about 19 days from when the cores are added to the culture medium for the first expansion.

[00535] In some embodiments, the TILs are not stored after the first expansion and prior to the second expansion, and the TILs proceed directly to the second expansion (for example, in some embodiments, there is no storage during the transition from Step B to Step D as shown in Figure 7). In some embodiments, the transition occurs in closed system, as described herein. In some embodiments, the TILs from the first expansion, the second population of TILs, proceeds directly into the second expansion with no transition period.

[00536] In some embodiments, the transition from the first expansion to the second expansion, for example, Step C according to Figure 7, is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a single bioreactor is employed. In some embodiments, the single bioreactor employed is for example a GREX-10 or a GREX-100. In some embodiments, the closed system bioreactor is a single bioreactor.

D. STEP D: Second Expansion

[00537] In some embodiments, the TIL cell population is expanded in number after harvest and initial bulk processing for example, after Step A and Step B, and the transition referred to as Step C, as indicated in Figure 7). This further expansion is referred to herein as the second expansion, which can include expansion processes generally referred to in the art as a rapid expansion process (REP; as well as processes as indicated in Step D of Figure 7). The second expansion is generally accomplished using a culture media comprising a number of components, including feeder cells, a cytokine source, and an anti-CD3 antibody, in a gas-permeable container.

[00538] In some embodiments, the second expansion or second TIL expansion (which can include expansions sometimes referred to as REP; as well as processes as indicated in Step D of Figure 7) of TIL can be performed using any TIL flasks or containers known by those of skill in the art. In some embodiments, the second TIL expansion can proceed for 10 days, 11 days, 12 days, 13 days, or 14 days. In some embodiments, the second TIL expansion can proceed for about 10 days to about 14 days. In some embodiments, the second TIL expansion can proceed for about 11 days to about 14 days. In some embodiments, the second TIL

expansion can proceed for about 12 days to about 14 days. In some embodiments, the second TIL expansion can proceed for about 13 days to about 14 days. In some embodiments, the second TIL expansion can proceed for about 14 days.

[00539] In an embodiment, the second expansion can be performed in a gas permeable container using the methods of the present disclosure (including for example, expansions referred to as REP; as well as processes as indicated in Step D of Figure 7). For example, TILs can be rapidly expanded using non-specific T-cell receptor stimulation in the presence of interleukin-2 (IL-2) or interleukin-15 (IL-15). The non-specific T-cell receptor stimulus can include, for example, about 30 ng/mL of OKT3, a mouse monoclonal anti-CD3 antibody (commercially available from Ortho-McNeil, Raritan, NJ or Miltenyi Biotech, Auburn, CA). TILs can be expanded to induce further stimulation of the TILs *in vitro* by including one or more antigens during the second expansion, including antigenic portions thereof, such as epitope(s), of the cancer, which can be optionally expressed from a vector, such as a human leukocyte antigen A2 (HLA-A2) binding peptide, *e.g.*, 0.3 μM MART-1 :26-35 (27 L) or gpl 00:209-217 (210M), optionally in the presence of a T-cell growth factor, such as 300 IU/mL IL-2 or IL-15. Other suitable antigens may include, *e.g.*, NY-ESO-1, TRP-1, TRP-2, tyrosinase cancer antigen, MAGE-A3, SSX-2, and VEGFR2, or antigenic portions thereof. TIL may also be rapidly expanded by re-stimulation with the same antigen(s) of the cancer pulsed onto HLA-A2-expressing antigen-presenting cells. Alternatively, the TILs can be further re-stimulated with, *e.g.*, example, irradiated, autologous lymphocytes or with irradiated HLA-A2+ allogeneic lymphocytes and IL-2. In some embodiments, the re-stimulation occurs as part of the second expansion. In some embodiments, the second expansion occurs in the presence of irradiated, autologous lymphocytes or with irradiated HLA-A2+ allogeneic lymphocytes and IL-2. In some embodiments, 13×10^6 feeder cells are added during the second expansion. In some embodiments, 13×10^6 feeder cells are added at the start of second expansion.

[00540] In an embodiment, the cell culture medium further comprises IL-2. In some embodiments, the cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises about 1000 IU/mL, about 1500 IU/mL, about 2000 IU/mL, about 2500 IU/mL, about 3000 IU/mL, about 3500 IU/mL, about 4000 IU/mL, about 4500 IU/mL, about 5000 IU/mL, about 5500 IU/mL, about 6000 IU/mL, about 6500 IU/mL, about 7000 IU/mL, about 7500 IU/mL, or about 8000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises between 1000 and 2000 IU/mL, between

2000 and 3000 IU/mL, between 3000 and 4000 IU/mL, between 4000 and 5000 IU/mL, between 5000 and 6000 IU/mL, between 6000 and 7000 IU/mL, between 7000 and 8000 IU/mL, or between 8000 IU/mL of IL-2.

[00541] In an embodiment, the cell culture medium comprises OKT3 antibody. In some embodiments, the cell culture medium comprises about 30 ng/mL of OKT3 antibody. In an embodiment, the cell culture medium comprises about 0.1 ng/mL, about 0.5 ng/mL, about 1 ng/mL, about 2.5 ng/mL, about 5 ng/mL, about 7.5 ng/mL, about 10 ng/mL, about 15 ng/mL, about 20 ng/mL, about 25 ng/mL, about 30 ng/mL, about 35 ng/mL, about 40 ng/mL, about 50 ng/mL, about 60 ng/mL, about 70 ng/mL, about 80 ng/mL, about 90 ng/mL, about 100 ng/mL, about 200 ng/mL, about 500 ng/mL, and about 1 μg/mL of OKT3 antibody. In an embodiment, the cell culture medium comprises between 0.1 ng/mL and 1 ng/mL, between 1 ng/mL and 5 ng/mL, between 5 ng/mL and 10 ng/mL, between 10 ng/mL and 20 ng/mL, between 20 ng/mL and 30 ng/mL, between 30 ng/mL and 40 ng/mL, between 40 ng/mL and 50 ng/mL, and between 50 ng/mL and 100 ng/mL of OKT3 antibody.

[00542] In some embodiments, a combination of IL-2, IL-7, IL-15, and/or IL-21 are employed as a combination during the second expansion. In some embodiments, IL-2, IL-7, IL-15, and/or IL-21 as well as any combinations thereof can be included during the second expansion, including for example during a Step D processes according to Figure 7, as well as described herein. In some embodiments, a combination of IL-2, IL-15, and IL-21 are employed as a combination during the second expansion. In some embodiments, IL-2, IL-15, and IL-21 as well as any combinations thereof can be included during Step D processes according to Figure 7 and as described herein.

[00543] In some embodiments, the second expansion can be conducted in a supplemented cell culture medium comprising IL-2, OKT-3, and antigen-presenting feeder cells. In some embodiments, the second expansion occurs in a supplemented cell culture medium. In some embodiments, the supplemented cell culture medium comprises IL-2, OKT-3, and antigen-presenting feeder cells. In some embodiments, the second cell culture medium comprises IL-2, OKT-3, and antigen-presenting cells (APCs; also referred to herein as antigen-presenting feeder cells, feeder cells or feeders). In some embodiments, the second expansion occurs in a cell culture medium comprising IL-2, OKT-3, and antigen-presenting feeder cells (*i.e.*, antigen presenting cells).

[00544] In some embodiments, the second expansion is performed by further supplementing the cell culture medium of the population of TILs in the second expansion with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody (sometimes referred to as the second population of TILs). For instance, the cell culture medium of the population of TILs in the second expansion is supplemented with IL-15, an OX40 agonistic antibody, a 4-1BB agonistic antibody, a combination of IL-15 and an OX40 agonistic antibody, a combination of IL-15, and a 4-1BB agonistic antibody, a combination of an OX40 agonistic antibody and a 4-1BB agonistic antibody, a combination of IL-15, an OX40 agonistic antibody, and a 4-1BB agonistic antibody, and any combinations thereof. In some embodiments, OX40 is included when the cores are from an ovarian tumor (*e.g.*, of ovarian origin). In some embodiments, OKT-3 is added during the first and/or second expansion. In some embodiuments, OKT-3 is added at day-3, along with the antigen-presenting feeder cells (APCs). In some embodiuments, when the sample is from an FNA, OKT-3 is added at day-1, along with the antigen-presenting feeder cells (APCs).

[00545] In some embodiments the antigen-presenting feeder cells (APCs) are PBMCs. In an embodiment, the ratio of TILs to PBMCs and/or antigen-presenting cells in the rapid expansion and/or the second expansion is about 1 to 25, about 1 to 50, about 1 to 100, about 1 to 125, about 1 to 150, about 1 to 175, about 1 to 200, about 1 to 225, about 1 to 250, about 1 to 275, about 1 to 300, about 1 to 325, about 1 to 350, about 1 to 375, about 1 to 400, or about 1 to 500. In an embodiment, the ratio of TILs to PBMCs in the rapid expansion and/or the second expansion is between 1 to 50 and 1 to 300. In an embodiment, the ratio of TILs to PBMCs in the rapid expansion and/or the second expansion is between 1 to 100 and 1 to 200.

[00546] In an embodiment, REP and/or the second expansion is performed in flasks with the bulk TILs being mixed with a 100- or 200-fold excess of inactivated feeder cells, 30 mg/mL OKT3 anti-CD3 antibody and 3000 IU/mL IL-2 in 150 ml media. Media replacement is done (generally 2/3 media replacement via respiration with fresh media) until the cells are transferred to an alternative growth chamber. Alternative growth chambers include GREX flasks and gas permeable containers as more fully discussed below.

[00547] In some embodiments, the second expansion (which can include processes referred to as the REP process) is 11-14 days, as discussed in the examples and figures. In some embodiments, the second expansion is 11 days. In some embodiments, the second expansion is 12 days. In some embodiments, the second expansion is 13 days. In some embodiments, the second expansion is 14 days.

[00548] In an embodiment, REP and/or the second expansion may be performed using T-175 flasks and gas permeable bags as previously described (Tran, *et al.*, *J. Immunother.* **2008,** *31,* 742-51; Dudley, *et al.*, *J. Immunother.* **2003,** *26,* 332-42) or gas permeable cultureware (G-Rex flasks). In some embodiments, the second expansion (including expansions referred to as rapid expansions) is performed in T-175 flasks, and about 1×10^6 TILs suspended in 150 mL of media may be added to each T-175 flask. The TILs may be cultured in a 1 to 1 mixture of CM and AIM-V medium, supplemented with 3000 IU per mL of IL-2 and 30 ng per ml of anti-CD3. The T-175 flasks may be incubated at 37° C in 5% CO_2. Half the media may be exchanged on day 5 using 50/50 medium with 3000 IU per mL of IL-2. In some embodiments, on day 7 cells from two T-175 flasks may be combined in a 3 L bag and 300 mL of AIM V with 5% human AB serum and 3000 IU per mL of IL-2 was added to the 300 ml of TIL suspension. The number of cells in each bag was counted every day or two and fresh media was added to keep the cell count between 0.5 and 2.0×10^6 cells/mL.

[00549] In an embodiment, the second expansion (which can include expansions referred to as REP, as well as those referred to in Step D of Figure 7) may be performed in 500 mL capacity gas permeable flasks with 100 cm gas-permeable silicon bottoms (G-Rex 100, commercially available from Wilson Wolf Manufacturing Corporation, New Brighton, MN, USA), 5×10^6 or 10×10^6 TIL may be cultured with PBMCs in 400 mL of 50/50 medium, supplemented with 5% human AB serum, 3000 IU per mL of IL-2 and 30 ng per ml of anti-CD3 (OKT3). The G-Rex 100 flasks may be incubated at 37°C in 5% CO_2. On day 5, 250 mL of supernatant may be removed and placed into centrifuge bottles and centrifuged at 1500 rpm ($491 \times g$) for 10 minutes. The TIL pellets may be re-suspended with 150 mL of fresh medium with 5% human AB serum, 3000 IU per mL of IL-2, and added back to the original G-Rex 100 flasks. When TIL are expanded serially in G-Rex 100 flasks, on day 7 the TIL in each G-Rex 100 may be suspended in the 300 mL of media present in each flask and the cell suspension may be divided into 3-100 mL aliquots that may be used to seed 3 G-Rex 100 flasks. Then 150 mL of AIM-V with 5% human AB serum and 3000 IU per mL of IL-2 may be added to each flask. The G-Rex 100 flasks may be incubated at 37° C in 5% CO_2 and after 4 days 150 mL of AIM-V with 3000 IU per mL of IL-2 may be added to each G-Rex 100 flask. The cells may be harvested on day 11 of culture. The cells may be harvested on day 12 of culture. The cells may be harvested on day 13 of culture. The cells may be harvested on day 14 of culture. In some embodiments, cultures are expanded into a G-Rex 500.

[00550] In an embodiment, the second expansion (including expansions referred to as REP) is performed in flasks with the bulk TILs being mixed with a 100- or 200-fold excess of inactivated feeder cells, 30 mg/mL OKT3 anti-CD3 antibody and 3000 IU/mL IL-2 in 150 ml media. In some embodiments, media replacement is done until the cells are transferred to an alternative growth chamber. In some embodiments, 2/3 of the media is replaced by aspiration of spent media and replacement with an equivalent volumne of fresh media. In some embodiments, alternative growth chambers include GRex flasks and gas permeable containers as more fully discussed below.

[00551] In an embodiment, the second expansion (including expansions referred to as REP) is performed and further comprises a step wherein TILs are selected for superior tumor reactivity. Any selection method known in the art may be used. For example, the methods described in U.S. Patent Application Publication No. 2016/0010058 A1, the disclosures of which are incorporated herein by reference, may be used for selection of TILs for superior tumor reactivity.

[00552] Optionally, a cell viability assay can be performed after the second expansion (including expansions referred to as the REP expansion), using standard assays known in the art. For example, a trypan blue exclusion assay can be done on a sample of the bulk TILs, which selectively labels dead cells and allows a viability assessment. In some embodiments, TIL samples can be counted and viability determined using a Cellometer K2 automated cell counter (Nexcelom Bioscience, Lawrence, MA). In some embodiments, viability is determined according to the Cellometer K2 Image Cytometer Automatic Cell Counter protocol.

[00553] In some embodiments, the second expansion (including expansions referred to as REP) of TIL can be performed using T-175 flasks and gas-permeable bags as previously described (Tran KQ, Zhou J, Durflinger KH, *et al.*, **2008**, *J Immunother.*, 31:742–751, and Dudley ME, Wunderlich JR, Shelton TE, *et al.* 2003, *J Immunother.*, 26:332–342) or gas-permeable G-Rex flasks. In some embodiments, the second expansion is performed using flasks. In some embodiments, the second expansion is performed using gas-permeable G-Rex flasks. In some embodiments, the second expansion is performed in T-175 flasks, and about 1×10^6 TIL are suspended in about 150 mL of media and this is added to each T-175 flask. The TIL are cultured with irradiated (50 Gy) allogeneic PBMC as "feeder" cells at a ratio of 1 to 100 and the cells were cultured in a 1 to 1 mixture of CM and AIM-V medium (50/50 medium), supplemented with 3000 IU/mL of IL-2 and 30 ng/mL of anti-CD3. The T-175

flasks are incubated at 37°C in 5% CO_2. In some embodiments, half the media is changed on day 5 using 50/50 medium with 3000 IU/mL of IL-2. In some embodiments, on day 7, cells from 2 T-175 flasks are combined in a 3 L bag and 300 mL of AIM-V with 5% human AB serum and 3000 IU/mL of IL-2 is added to the 300 mL of TIL suspension. The number of cells in each bag can be counted every day or two and fresh media can be added to keep the cell count between about 0.5 and about 2.0×10^6 cells/mL. In some embodiments, cultures are expanded into a G-Rex 500.

[00554] In some embodiments, the second expansion (including expansions referred to as REP) are performed in 500 mL capacity flasks with 100 cm^2 gas-permeable silicon bottoms (G-Rex100, Wilson Wolf) (Fig. 1), about 5×10^6 or 10×10^6 TIL are cultured with irradiated allogeneic PBMC at a ratio of 1 to 100 in 400 mL of 50/50 medium, supplemented with 3000 IU/mL of IL-2 and 30 ng/mL of anti-CD3. The G-Rex100 flasks are incubated at 37°C in 5% CO_2. In some embodiments, on day 5, 250 mL of supernatant is removed and placed into centrifuge bottles and centrifuged at 1500 rpm (491 × g) for 10 minutes. The TIL pellets can then be resuspended with 150 mL of fresh 50/50 medium with 3000 IU/ mL of IL-2 and added back to the original G-Rex100 flasks. In embodiments where TILs are expanded serially in G-Rex100 flasks, on day 7 the TIL in each G-Rex100 are suspended in the 300 mL of media present in each flask and the cell suspension was divided into three 100 mL aliquots that are used to seed 3 G-Rex100 flasks. Then 150 mL of AIM-V with 5% human AB serum and 3000 IU/mL of IL-2 is added to each flask. The G-Rex100 flasks are incubated at 37°C in 5% CO_2 and after 4 days 150 mL of AIM-V with 3000 IU/mL of IL-2 is added to each G-Rex100 flask. The cells are harvested on day 14 of culture. In some embodiments, cultures are expanded into a G-Rex 500.

[00555] In some embodiments, the second expansion culture medium (*e.g.*, sometimes referred to as CM2 or the second cell culture medium), comprises IL-2, OKT-3, as well as the antigen-presenting feeder cells (APCs), as discussed in more detail below.

[00556] In some embodiments, the second expansion, for example, Step D according to Figure 7, is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a single bioreactor is employed. In some embodiments, the single bioreactor employed is for example a GREX-10 or a GREX-100. In some embodiments, the closed system bioreactor is a single bioreactor.

1. Feeder Cells and Antigen Presenting Cells

[00557] In an embodiment, the second expansion procedures described herein (for example including expansion such as those described in Step D from Figure 7, as well as those referred to as REP) require an excess of feeder cells during REP TIL expansion and/or during the second expansion. In many embodiments, the feeder cells are peripheral blood mononuclear cells (PBMCs) obtained from standard whole blood units from healthy blood donors. The PBMCs are obtained using standard methods such as Ficoll-Paque gradient separation.

[00558] In general, the allogenic PBMCs are inactivated, either via irradiation or heat treatment, and used in the REP procedures, as described in the examples, in particular Example 4, which provides an exemplary protocol for evaluating the replication incompetence of irradiate allogeneic PBMCs.

[00559] In some embodiments, PBMCs are considered replication incompetent and accepted for use in the TIL expansion procedures described herein if the total number of viable cells on day 14 is less than the initial viable cell number put into culture on day 0 of the REP and/or day 0 of the second expansion (*i.e.*, the start day of the second expansion). See, for example, Examples 3 and/or 4.

[00560] In some embodiments, PBMCs are considered replication incompetent and accepted for use in the TIL expansion procedures described herein if the total number of viable cells, cultured in the presence of OKT3 and IL-2, on day 7 and day 14 has not increased from the initial viable cell number put into culture on day 0 of the REP and/or day 0 of the second expansion (*i.e.*, the start day of the second expansion). In some embodiments, the PBMCs are cultured in the presence of 30 ng/mL OKT3 antibody and 3000 IU/mL IL-2. See, for example, Examples 3 and/or 4.

[00561] In some embodiments, PBMCs are considered replication incompetent and accepted for use in the TIL expansion procedures described herein if the total number of viable cells, cultured in the presence of OKT3 and IL-2, on day 7 and day 14 has not increased from the initial viable cell number put into culture on day 0 of the REP and/or day 0 of the second expansion (*i.e.*, the start day of the second expansion). In some embodiments, the PBMCs are cultured in the presence of 5-60 ng/mL OKT3 antibody and 1000-6000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 10-50 ng/mL OKT3 antibody and 2000-5000 IU/mL IL-2. In some embodiments, the

PBMCs are cultured in the presence of 20-40 ng/mL OKT3 antibody and 2000-4000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 25-35 ng/ml OKT3 antibody and 2500-3500 IU/mL IL-2.

[00562] In some embodiments, the antigen-presenting feeder cells are PBMCs. In some embodiments, the antigen-presenting feeder cells are artificial antigen-presenting feeder cells. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is about 1 to 25, about 1 to 50, about 1 to 100, about 1 to 125, about 1 to 150, about 1 to 175, about 1 to 200, about 1 to 225, about 1 to 250, about 1 to 275, about 1 to 300, about 1 to 325, about 1 to 350, about 1 to 375, about 1 to 400, or about 1 to 500. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is between 1 to 50 and 1 to 300. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is between 1 to 100 and 1 to 200.

[00563] In an embodiment, the second expansion procedures described herein require an excess of feeder cells during the second expansion. In many embodiments, the feeder cells are peripheral blood mononuclear cells (PBMCs) obtained from standard whole blood units from healthy blood donors. The PBMCs are obtained using standard methods such as Ficoll-Paque gradient separation. In an embodiment, artificial antigen-presenting (aAPC) cells are used in place of PBMCs.

[00564] In general, the allogenic PBMCs are inactivated, either via irradiation or heat treatment, and used in the TIL expansion procedures described herein, including the exemplary procedures described in, for example, Figure 7.

[00565] In an embodiment, artificial antigen presenting cells are used in the second expansion as a replacement for, or in combination with, PBMCs.

[00566] The expansion methods described herein generally use culture media with high doses of a cytokine, in particular IL-2, as is known in the art.

[00567] Alternatively, using combinations of cytokines for the rapid expansion and or second expansion of TILS is additionally possible, with combinations of two or more of IL-2, IL-15 and IL-21 as is generally outlined in International Publication No. WO 2015/189356 and W International Publication No. WO 2015/189357, hereby expressly incorporated by reference in their entirety. Thus, possible combinations include IL-2 and IL-15, IL-2 and IL-21, IL-15 and IL-21 and IL-2, IL-15 and IL-21, with the latter finding particular use in many

embodiments. The use of combinations of cytokines specifically favors the generation of lymphocytes, and in particular T-cells as described therein.

E. STEP E: Harvest TILs

[00568] After the second expansion step, cells can be harvested. In some embodiments the TILs are harvested after one, two, three, four or more expansion steps, for example as provided in Figure 7. In some embodiments the TILs are harvested after two expansion steps, for example as provided in Figure 7.

[00569] TILs can be harvested in any appropriate and sterile manner, including for example by centrifugation. Methods for TIL harvesting are well known in the art and any such know methods can be employed with the present process. In some embodiments, TILS are harvest using an automated system.

[00570] In some embodiments, the harvest, for example, Step E according to Figure 7, is performed from a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a single bioreactor is employed. In some embodiments, the single bioreactor employed is for example a GREX-10 or a GREX-100. In some embodiments, the closed system bioreactor is a single bioreactor.

F. STEP F: Final Formulation/Transfer to Infusion Bag

[00571] After Steps A through E as provided in an exemplary order in Figure 7 and as outlined in detailed above and herein are complete, cells are transferred to a container for use in administration to a patient. In some embodiments, once a therapeutically sufficient number of TILs are obtained using the expansion methods described above, they are transferred to a container for use in administration to a patient.

[00572] In an embodiment, TILs expanded using processes of the present disclosure are administered to a patient as a pharmaceutical composition. In an embodiment, the pharmaceutical composition is a suspension of TILs in a sterile buffer. TILs expanded using processes of the present disclosure may be administered by any suitable route as known in the art. In some embodiments, the T-cells are administered as a single intra-arterial or

intravenous infusion, which preferably lasts approximately 30 to 60 minutes. Other suitable routes of administration include intraperitoneal, intrathecal, and intralymphatic.

[00573] In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant and an isotonic agent. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant comprising dimethylsulfoxide and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant comprising dimethylsulfoxide and dextran 40 and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions delivered in a sterile infusion bag, such compositions further comprising a cyropreservant comprising dimethylsulfoxide and dextran 40 and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate.

1. Pharmaceutical Compositions, Dosages, and Dosing Regimens

[00574] In an embodiment, TILs expanded using the methods of the present disclosure are administered to a patient as a pharmaceutical composition. In an embodiment, the pharmaceutical composition is a suspension of TILs in a sterile buffer. TILs expanded using processes of the present disclosure may be administered by any suitable route as known in the art. In some embodiments, the T-cells are administered as a single intra-arterial or intravenous infusion, which preferably lasts approximately 30 to 60 minutes. Other suitable routes of administration include intraperitoneal, intrathecal, and intralymphatic administration.

[00575] Any suitable dose of TILs can be administered. In some embodiments, a therapeutically sufficient number of TILs are needed for a suitable dosage. In some embodiments, from about 2.3×10^{10} to about 13.7×10^{10} TILs are administered, with an average of around 7.8×10^{10} TILs, particularly if the cancer is melanoma. In an embodiment, about 1.2×10^{10} to about 4.3×10^{10} of TILs are administered. In some embodiments, about 3×10^{10} to about 12×10^{10} TILs are administered. In some embodiments, about 4×10^{10} to about 10×10^{10} TILs are administered. In some embodiments, about 5×10^{10} to about 8×10^{10} TILs are administered. In some embodiments, about 6×10^{10} to about 8×10^{10} TILs are administered. In

some embodiments, about 7×10^{10} to about 8×10^{10} TILs are administered. In some embodiments, the therapeutically effective dosage is about 2.3×10^{10} to about 13.7×10^{10}. In some embodiments, the therapeutically effective dosage is about 7.8×10^{10} TILs, particularly of the cancer is melanoma. In some embodiments, the therapeutically effective dosage is about 1.2×10^{10} to about 4.3×10^{10} of TILs. In some embodiments, the therapeutically effective dosage is about 3×10^{10} to about 12×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 4×10^{10} to about 10×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 5×10^{10} to about 8×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 6×10^{10} to about 8×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 7×10^{10} to about 8×10^{10} TILs.

[00576] In some embodiments, the number of the TILs provided in the pharmaceutical compositions of the invention is about 1×10^{6}, 2×10^{6}, 3×10^{6}, 4×10^{6}, 5×10^{6}, 6×10^{6}, 7×10^{6}, 8×10^{6}, 9×10^{6}, 1×10^{7}, 2×10^{7}, 3×10^{7}, 4×10^{7}, 5×10^{7}, 6×10^{7}, 7×10^{7}, 8×10^{7}, 9×10^{7}, 1×10^{8}, 2×10^{8}, 3×10^{8}, 4×10^{8}, 5×10^{8}, 6×10^{8}, 7×10^{8}, 8×10^{8}, 9×10^{8}, 1×10^{9}, 2×10^{9}, 3×10^{9}, 4×10^{9}, 5×10^{9}, 6×10^{9}, 7×10^{9}, 8×10^{9}, 9×10^{9}, 1×10^{10}, 2×10^{10}, 3×10^{10}, 4×10^{10}, 5×10^{10}, 6×10^{10}, 7×10^{10}, 8×10^{10}, 9×10^{10}, 1×10^{11}, 2×10^{11}, 3×10^{11}, 4×10^{11}, 5×10^{11}, 6×10^{11}, 7×10^{11}, 8×10^{11}, 9×10^{11}, 1×10^{12}, 2×10^{12}, 3×10^{12}, 4×10^{12}, 5×10^{12}, 6×10^{12}, 7×10^{12}, 8×10^{12}, 9×10^{12}, 1×10^{13}, 2×10^{13}, 3×10^{13}, 4×10^{13}, 5×10^{13}, 6×10^{13}, 7×10^{13}, 8×10^{13}, and 9×10^{13}. In an embodiment, the number of the TILs provided in the pharmaceutical compositions of the invention is in the range of 1×10^{6} to 5×10^{6}, 5×10^{6} to 1×10^{7}, 1×10^{7} to 5×10^{7}, 5×10^{7} to 1×10^{8}, 1×10^{8} to 5×10^{8}, 5×10^{8} to 1×10^{9}, 1×10^{9} to 5×10^{9}, 5×10^{9} to 1×10^{10}, 1×10^{10} to 5×10^{10}, 5×10^{10} to 1×10^{11}, 5×10^{11} to 1×10^{12}, 1×10^{12} to 5×10^{12}, and 5×10^{12} to 1×10^{13}. In some embodiments, the therapeutically effective dosage is about 1×10^{6}, 2×10^{6}, 3×10^{6}, 4×10^{6}, 5×10^{6}, 6×10^{6}, 7×10^{6}, 8×10^{6}, 9×10^{6}, 1×10^{7}, 2×10^{7}, 3×10^{7}, 4×10^{7}, 5×10^{7}, 6×10^{7}, 7×10^{7}, 8×10^{7}, 9×10^{7}, 1×10^{8}, 2×10^{8}, 3×10^{8}, 4×10^{8}, 5×10^{8}, 6×10^{8}, 7×10^{8}, 8×10^{8}, 9×10^{8}, 1×10^{9}, 2×10^{9}, 3×10^{9}, 4×10^{9}, 5×10^{9}, 6×10^{9}, 7×10^{9}, 8×10^{9}, 9×10^{9}, 1×10^{10}, 2×10^{10}, 3×10^{10}, 4×10^{10}, 5×10^{10}, 6×10^{10}, 7×10^{10}, 8×10^{10}, 9×10^{10}, 1×10^{11}, 2×10^{11}, 3×10^{11}, 4×10^{11}, 5×10^{11}, 6×10^{11}, 7×10^{11}, 8×10^{11}, 9×10^{11}, 1×10^{12}, 2×10^{12}, 3×10^{12}, 4×10^{12}, 5×10^{12}, 6×10^{12}, 7×10^{12}, 8×10^{12}, 9×10^{12}, 1×10^{13}, 2×10^{13}, 3×10^{13}, 4×10^{13}, 5×10^{13}, 6×10^{13}, 7×10^{13}, 8×10^{13}, and 9×10^{13}.

[00577] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is less than, for example, 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%,

0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002% or 0.0001% w/w, w/v or v/v of the pharmaceutical composition.

[00578] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is greater than 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19.75%, 19.50%, 19.25% 19%, 18.75%, 18.50%, 18.25% 18%, 17.75%, 17.50%, 17.25% 17%, 16.75%, 16.50%, 16.25% 16%, 15.75%, 15.50%, 15.25% 15%, 14.75%, 14.50%, 14.25% 14%, 13.75%, 13.50%, 13.25% 13%, 12.75%, 12.50%, 12.25% 12%, 11.75%, 11.50%, 11.25% 11%, 10.75%, 10.50%, 10.25% 10%, 9.75%, 9.50%, 9.25% 9%, 8.75%, 8.50%, 8.25% 8%, 7.75%, 7.50%, 7.25% 7%, 6.75%, 6.50%, 6.25% 6%, 5.75%, 5.50%, 5.25% 5%, 4.75%, 4.50%, 4.25%, 4%, 3.75%, 3.50%, 3.25%, 3%, 2.75%, 2.50%, 2.25%, 2%, 1.75%, 1.50%, 125%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002% or 0.0001% w/w, w/v, or v/v of the pharmaceutical composition.

[00579] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is in the range from about 0.0001% to about 50%, about 0.001% to about 40%, about 0.01% to about 30%, about 0.02% to about 29%, about 0.03% to about 28%, about 0.04% to about 27%, about 0.05% to about 26%, about 0.06% to about 25%, about 0.07% to about 24%, about 0.08% to about 23%, about 0.09% to about 22%, about 0.1% to about 21%, about 0.2% to about 20%, about 0.3% to about 19%, about 0.4% to about 18%, about 0.5% to about 17%, about 0.6% to about 16%, about 0.7% to about 15%, about 0.8% to about 14%, about 0.9% to about 12% or about 1% to about 10% w/w, w/v or v/v of the pharmaceutical composition.

[00580] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is in the range from about 0.001% to about 10%, about 0.01% to about 5%, about 0.02% to about 4.5%, about 0.03% to about 4%, about 0.04% to about 3.5%, about 0.05% to about 3%, about 0.06% to about 2.5%, about 0.07% to about 2%, about 0.08% to about 1.5%, about 0.09% to about 1%, about 0.1% to about 0.9% w/w, w/v or v/v of the pharmaceutical composition.

[00581] In some embodiments, the amount of the TILs provided in the pharmaceutical compositions of the invention is equal to or less than 10 g, 9.5 g, 9.0 g, 8.5 g, 8.0 g, 7.5 g, 7.0 g, 6.5 g, 6.0 g, 5.5 g, 5.0 g, 4.5 g, 4.0 g, 3.5 g, 3.0 g, 2.5 g, 2.0 g, 1.5 g, 1.0 g, 0.95 g, 0.9 g, 0.85 g, 0.8 g, 0.75 g, 0.7 g, 0.65 g, 0.6 g, 0.55 g, 0.5 g, 0.45 g, 0.4 g, 0.35 g, 0.3 g, 0.25 g, 0.2 g, 0.15 g, 0.1 g, 0.09 g, 0.08 g, 0.07 g, 0.06 g, 0.05 g, 0.04 g, 0.03 g, 0.02 g, 0.01 g, 0.009 g, 0.008 g, 0.007 g, 0.006 g, 0.005 g, 0.004 g, 0.003 g, 0.002 g, 0.001 g, 0.0009 g, 0.0008 g, 0.0007 g, 0.0006 g, 0.0005 g, 0.0004 g, 0.0003 g, 0.0002 g, or 0.0001 g.

[00582] In some embodiments, the amount of the TILs provided in the pharmaceutical compositions of the invention is more than 0.0001 g, 0.0002 g, 0.0003 g, 0.0004 g, 0.0005 g, 0.0006 g, 0.0007 g, 0.0008 g, 0.0009 g, 0.001 g, 0.0015 g, 0.002 g, 0.0025 g, 0.003 g, 0.0035 g, 0.004 g, 0.0045 g, 0.005 g, 0.0055 g, 0.006 g, 0.0065 g, 0.007 g, 0.0075 g, 0.008 g, 0.0085 g, 0.009 g, 0.0095 g, 0.01 g, 0.015 g, 0.02 g, 0.025 g, 0.03 g, 0.035 g, 0.04 g, 0.045 g, 0.05 g, 0.055 g, 0.06 g, 0.065 g, 0.07 g, 0.075 g, 0.08 g, 0.085 g, 0.09 g, 0.095 g, 0.1 g, 0.15 g, 0.2 g, 0.25 g, 0.3 g, 0.35 g, 0.4 g, 0.45 g, 0.5 g, 0.55 g, 0.6 g, 0.65 g, 0.7 g, 0.75 g, 0.8 g, 0.85 g, 0.9 g, 0.95 g, 1 g, 1.5 g, 2 g, 2.5, 3 g, 3.5, 4 g, 4.5 g, 5 g, 5.5 g, 6 g, 6.5 g, 7 g, 7.5 g, 8 g, 8.5 g, 9 g, 9.5 g, or 10 g.

[00583] The TILs provided in the pharmaceutical compositions of the invention are effective over a wide dosage range. The exact dosage will depend upon the route of administration, the form in which the compound is administered, the gender and age of the subject to be treated, the body weight of the subject to be treated, and the preference and experience of the attending physician. The clinically-established dosages of the TILs may also be used if appropriate. The amounts of the pharmaceutical compositions administered using the methods herein, such as the dosages of TILs, will be dependent on the human or mammal being treated, the severity of the disorder or condition, the rate of administration, the disposition of the active pharmaceutical ingredients and the discretion of the prescribing physician.

[00584] In some embodiments, TILs may be administered in a single dose. Such administration may be by injection, *e.g.*, intravenous injection. In some embodiments, TILs may be administered in multiple doses. Dosing may be once, twice, three times, four times, five times, six times, or more than six times per year. Dosing may be once a month, once every two weeks, once a week, or once every other day. Administration of TILs may continue as long as necessary.

[00585] In some embodiments, an effective dosage of TILs is about 1×10^6, 2×10^6, 3×10^6, 4×10^6, 5×10^6, 6×10^6, 7×10^6, 8×10^6, 9×10^6, 1×10^7, 2×10^7, 3×10^7, 4×10^7, 5×10^7, 6×10^7, 7×10^7, 8×10^7, 9×10^7, 1×10^8, 2×10^8, 3×10^8, 4×10^8, 5×10^8, 6×10^8, 7×10^8, 8×10^8, 9×10^8, 1×10^9, 2×10^9, 3×10^9, 4×10^9, 5×10^9, 6×10^9, 7×10^9, 8×10^9, 9×10^9, 1×10^{10}, 2×10^{10}, 3×10^{10}, 4×10^{10}, 5×10^{10}, 6×10^{10}, 7×10^{10}, 8×10^{10}, 9×10^{10}, 1×10^{11}, 2×10^{11}, 3×10^{11}, 4×10^{11}, 5×10^{11}, 6×10^{11}, 7×10^{11}, 8×10^{11}, 9×10^{11}, 1×10^{12}, 2×10^{12}, 3×10^{12}, 4×10^{12}, 5×10^{12}, 6×10^{12}, 7×10^{12}, 8×10^{12}, 9×10^{12}, 1×10^{13}, 2×10^{13}, 3×10^{13}, 4×10^{13}, 5×10^{13}, 6×10^{13}, 7×10^{13}, 8×10^{13}, and 9×10^{13}. In some embodiments, an effective dosage of TILs is in the range of 1×10^6 to 5×10^6, 5×10^6 to 1×10^7, 1×10^7 to 5×10^7, 5×10^7 to 1×10^8, 1×10^8 to 5×10^8, 5×10^8 to 1×10^9, 1×10^9 to 5×10^9, 5×10^9 to 1×10^{10}, 1×10^{10} to 5×10^{10}, 5×10^{10} to 1×10^{11}, 5×10^{11} to 1×10^{12}, 1×10^{12} to 5×10^{12}, and 5×10^{12} to 1×10^{13}.

[00586] In some embodiments, an effective dosage of TILs is in the range of about 0.01 mg/kg to about 4.3 mg/kg, about 0.15 mg/kg to about 3.6 mg/kg, about 0.3 mg/kg to about 3.2 mg/kg, about 0.35 mg/kg to about 2.85 mg/kg, about 0.15 mg/kg to about 2.85 mg/kg, about 0.3 mg to about 2.15 mg/kg, about 0.45 mg/kg to about 1.7 mg/kg, about 0.15 mg/kg to about 1.3 mg/kg, about 0.3 mg/kg to about 1.15 mg/kg, about 0.45 mg/kg to about 1 mg/kg, about 0.55 mg/kg to about 0.85 mg/kg, about 0.65 mg/kg to about 0.8 mg/kg, about 0.7 mg/kg to about 0.75 mg/kg, about 0.7 mg/kg to about 2.15 mg/kg, about 0.85 mg/kg to about 2 mg/kg, about 1 mg/kg to about 1.85 mg/kg, about 1.15 mg/kg to about 1.7 mg/kg, about 1.3 mg/kg mg to about 1.6 mg/kg, about 1.35 mg/kg to about 1.5 mg/kg, about 2.15 mg/kg to about 3.6 mg/kg, about 2.3 mg/kg to about 3.4 mg/kg, about 2.4 mg/kg to about 3.3 mg/kg, about 2.6 mg/kg to about 3.15 mg/kg, about 2.7 mg/kg to about 3 mg/kg, about 2.8 mg/kg to about 3 mg/kg, or about 2.85 mg/kg to about 2.95 mg/kg.

[00587] In some embodiments, an effective dosage of TILs is in the range of about 1 mg to about 500 mg, about 10 mg to about 300 mg, about 20 mg to about 250 mg, about 25 mg to about 200 mg, about 1 mg to about 50 mg, about 5 mg to about 45 mg, about 10 mg to about 40 mg, about 15 mg to about 35 mg, about 20 mg to about 30 mg, about 23 mg to about 28 mg, about 50 mg to about 150 mg, about 60 mg to about 140 mg, about 70 mg to about 130 mg, about 80 mg to about 120 mg, about 90 mg to about 110 mg, or about 95 mg to about 105 mg, about 98 mg to about 102 mg, about 150 mg to about 250 mg, about 160 mg to about 240 mg, about 170 mg to about 230 mg, about 180 mg to about 220 mg, about 190 mg to about 210 mg, about 195 mg to about 205 mg, or about 198 to about 207 mg.

[00588] An effective amount of the TILs may be administered in either single or multiple doses by any of the accepted modes of administration of agents having similar utilities, including intranasal and transdermal routes, by intra-arterial injection, intravenously, intraperitoneally, parenterally, intramuscularly, subcutaneously, topically, by transplantation, or by inhalation.

IV. TIL Manufacturing Processes – Gen 3 Processes

[00589] In some embodiments, the invention provides a population of TILs made by the method of any of GEN 3 processes modified to use a small biopsy, core biopsy or fine needle aspirate as theIn some embodiments, the invention provides the method of any of GEN 3 processes modified to use a small biopsy, core biopsy or fine needle aspirate as the source of T cells for expansion in the priming first expansion of any such GEN 3 process, wherein (1) the duration of the priming first expansion is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the rapid second expansion of such GEN 3 process or (2) the duration of the rapid second expansion of such GEN 3 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the rapid second expansion or (3) the duration of the priming first expansion is lengthened and the duration of the rapid second expansion of such GEN 3 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the rapid second expansion.

[00590] In some embodiments, the invention provides a population of TILs made by the method of any of GEN 3 processes modified to use a small biopsy, core biopsy or fine needle aspirate as the source of T cells for expansion in the priming first expansion of any such GEN 3 process, wherein (1) the duration of the priming first expansion is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the rapid second expansion of such GEN 3 process or (2) the duration of the rapid second expansion of such GEN 3 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the rapid second expansion or (3) the duration of the priming first expansion is lengthened and the duration of the the rapid second expansion of such GEN 3 process is lengthened to achieve the desired TIL cell count in the population of TILs harvested after the rapid second expansion.

[00591] Without being limited to any particular theory, it is believed that the priming first expansion that primes an activation of T cells obtained from a tumor core or fragment obtained from a donor followed by the rapid second expansion that boosts the activation of T

cells as described in some methods of the invention allows the preparation of expanded T cells that retain a "younger" phenotype, and as such the expanded T cells of the invention are expected to exhibit greater cytotoxicity against cancer cells than T cells expanded by other methods. In particular, it is believed that an activation of T cells that is primed by exposure to an anti-CD3 antibody (*e.g.* OKT-3), IL-2 and optionally antigen-presenting cells (APCs) and then boosted by subsequent exposure to additional anti-CD-3 antibody (*e.g.* OKT-3), IL-2 and APCs as taught by the methods of the invention limits or avoids the maturation of T cells in culture, yielding a population of T cells with a less mature phenotype, which T cells are less exhausted by expansion in culture and exhibit greater cytotoxicity against cancer cells. Thus, in some embodiments, the invention provides a method of expanding TILs comprising: (a) performing a priming first expansion of a first population of T cells obtained from a tumor core or fragment obtained from a donor by culturing the first population of T cells to effect growth and to prime an activation of the first population of T cells; (b) after the activation of the first population of T cells primed in step (a) begins to decay, performing a rapid second expansion of the first population of T cells by culturing the first population of T cells to effect growth and to boost the activation of the first population of T cells to obtain a second population of T cells; and (c) harvesting the second population of T cells. The Gen 3 process, methods of treatment using TILs from the Gen 3 process, and compositions of TILs prepared by the Gen 3 can be used in conjunction with small biopsy/core biopsy with durations of expansion periods adjusted as needed to achieve necessary cell counts, as provided herein below and throughout the present application.

[00592] In some embodiments, the step of rapid second expansion is split into a plurality of steps to achieve a scaling up of the culture by: (a) performing the rapid second expansion by culturing T cells in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (b) effecting the transfer of the T cells in the small scale culture to a second container larger than the first container, *e.g.*, a G-REX 500MCS container, and culturing the T cells from the small scale culture in a larger scale culture in the second container for a period of about 4 to 7 days. In some embodiments, the step of rapid expansion is split into a plurality of steps to achieve a scaling out of the culture by: (a) performing the rapid second expansion by culturing T cells in a first small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (b) effecting the transfer and apportioning of the T cells from the first small scale culture into and amongst at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18,

19, or 20 second containers that are equal in size to the first container, wherein in each second container the portion of the T cells from first small scale culture transferred to such second container is cultured in a second small scale culture for a period of about 4 to 7 days. In some embodiments, the step of rapid expansion is split into a plurality of steps to achieve a scaling out and scaling up of the culture by: (a) performing the rapid second expansion by culturing T cells in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (b) effecting the transfer and apportioning of the T cells from the small scale culture into and amongst at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 second containers that are larger in size than the first container, *e.g.*, G-REX 500MCS containers, wherein in each second container the portion of the T cells from the small scale culture transferred to such second container is cultured in a larger scale culture for a period of about 4 to 7 days. In some embodiments, the step of rapid expansion is split into a plurality of steps to achieve a scaling out and scaling up of the culture by: (a) performing the rapid second expansion by culturing T cells in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 4 days, and then (b) effecting the transfer and apportioning of the T cells from the small scale culture into and amongst 2, 3 or 4 second containers that are larger in size than the first container, *e.g.*, G-REX 500MCS containers, wherein in each second container the portion of the T cells from the small scale culture transferred to such second container is cultured in a larger scale culture for a period of about 5 days.

[00593] In some embodiments, the rapid second expansion is performed after the activation of T cells effected by the priming first expansion begins to decrease, abate, decay or subside.

[00594] In some embodiments, the rapid second expansion is performed after the activation of T cells effected by the priming first expansion has decreased by at or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%.

[00595] In some embodiments, the rapid second expansion is performed after the activation of T cells effected by the priming first expansion has decreased by a percentage in the range of at or about 1% to 100%.

[00596] In some embodiments, the rapid second expansion is performed after the activation of T cells effected by the priming first expansion has decreased by a percentage in the range of at or about 1% to 10%, 10% to 20%, 20% to 30%, 30% to 40%, 40% to 50%, 50% to 60%, 60% to 70%, 70% to 80%, 80% to 90%, or 90% to 100%.

[00597] In some embodiments, the rapid second expansion is performed after the activation of T cells effected by the priming first expansion has decreased by at least at or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99%.

[00598] In some embodiments, the rapid second expansion is performed after the activation of T cells effected by the priming first expansion has decreased by up to at or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100%.

[00599] In some embodiments, the decrease in the activation of T cells effected by the priming first expansion is determined by a reduction in the amount of interferon gamma released by the T cells in response to stimulation with antigen.

[00600] In some embodiments, the priming first expansion of T cells is performed during a period of up to at or about 7 days.

[00601] In some embodiments, the priming first expansion of T cells is performed during a period of up to at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, or 7 days.

[00602] In some embodiments, the priming first expansion of T cells is performed during a period of 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, or 7 days.

[00603] In some embodiments, the rapid second expansion of T cells is performed during a period of up to at or about 11 days.

[00604] In some embodiments, the rapid second expansion of T cells is performed during a period of up to at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days or 11 days.

[00605] In some embodiments, the rapid second expansion of T cells is performed during a period of 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days or 11 days.

[00606] In some embodiments, the priming first expansion of T cells is performed during a period of from at or about 1 day to at or about 7 days and the rapid second expansion of T cells is performed during a period of from at or about 1 day to at or about 11 days.

[00607] In some embodiments, the priming first expansion of T cells is performed during a period of up to at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, or 7 days and the rapid second expansion of T cells is performed during a period of up to at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days or 11 days.

[00608] In some embodiments, the priming first expansion of T cells is performed during a period of from at or about 1 day to at or about 7 days and the rapid second expansion of T cells is performed during a period of from at or about 1 day to at or about 9 days.

[00609] In some embodiments, the priming first expansion of T cells is performed during a period of 7 days and the rapid second expansion of T cells is performed during a period of 9 days.

[00610] In some embodiments, the T cells are tumor infiltrating lymphocytes (TILs).

[00611] In some embodiments, the T cells are marrow infiltrating lymphocytes (MILs).

[00612] In some embodiments, the T cells are obtained from a donor suffering from a cancer.

[00613] In some embodiments, the T cells are TILs obtained from a core biopsy or fine needle aspirate obtained from a tumor in a patient suffering from a cancer.

[00614] In some embodiments, the T cells are MILs obtained from bone marrow of a patient suffering from a hematologic malignancy.

[00615] In some embodiments, the donor is suffering from a cancer. In some embodiments, the cancer is the cancer is selected from the group consisting of melanoma, ovarian cancer, cervical cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer (including head and neck squamous cell carcinoma (HNSCC)), glioblastoma (including GBM), gastrointestinal cancer, renal cancer, and renal cell carcinoma. In some embodiments, the cancer is selected from the group consisting of melanoma, ovarian cancer, cervical

cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer (including head and neck squamous cell carcinoma (HNSCC)), glioblastoma (including GBM), gastrointestinal cancer, renal cancer, and renal cell carcinoma. In some embodments, the donor is suffering from a tumor. In some embodiments, the tumor is a liquid tumor. In some embodiments, the tumor is a solid tumor. In some embodiments, the donor is suffering from a hematologic malignancy.

[00616] An exemplary TIL process known as process 3 (also referred to herein as GEN3) containing some of these features is depicted in Figure 85 (in particular, *e.g.*, Figure 85B), and some of the advantages of this embodiment of the present invention over process 2A are described in Figures 1, 2, 30, and 31 (in particular, *e.g.*, Figure 85B). Two embodiments of process 3 are shown in Figures 1 and 30 (in particular, *e.g.*, Figure 85B). Process 2A or Gen 2 is also described in U.S. Patent Publication No. 2018/0280436, incorporated by reference herein in its entirety. The Gen 3 process is also described in USSN 62/755,954 filed on November 5, 2018 (116983-5045-PR).

[00617] As discussed and generally outlined herein, TILs are taken from a patient sample and manipulated to expand their number prior to transplant into a patient using the TIL expansion process described herein and referred to as Gen 3. In some embodiments, the TILs may be optionally genetically manipulated as discussed below. In some embodiments, the TILs may be cryopreserved prior to or after expansion. Once thawed, they may also be restimulated to increase their metabolism prior to infusion into a patient.

[00618] In some embodiments, the priming first expansion (including processes referred herein as the pre-Rapid Expansion (Pre-REP), as well as processes shown in Figure 85 (in particular, *e.g.*, Figure 85B) as Step B) is shortened to 1 to 7 days and the rapid second expansion (including processes referred to herein as Rapid Expansion Protocol (REP) as well as processes shown in Figure 85 (in particular, *e.g.*, Figure 85B) as Step D) is shortened to 1 to 9 days, as discussed in detail below as well as in the examples and figures. In some embodiments, the priming first expansion (for example, an expansion described as Step B in Figure 85 (in particular, *e.g.*, Figure 85B)) is shortened to 7 days and the rapid second expansion (for example, an expansion as described in Step D in Figure 85 (in particular, *e.g.*, Figure 85B)) is 7 to 9 days. In some embodiments, the combination of the priming first expansion and rapid second expansion (for example, expansions described as Step B and Step D in Figure 85 (in particular, *e.g.*, Figure 85B)) is 14-16 days, as discussed in detail below

and in the examples and figures. Particularly, it is considered that certain embodiments of the present invention comprise a priming first expansion step in which TILs are activated by exposure to an anti-CD3 antibody, *e.g.*, OKT-3 in the presence of IL-2 or exposure to an antigen in the presence of at least IL-2 and an anti-CD3 antibody *e.g.* OKT-3. In certain embodiments, the TILs which are activated in the priming first expansion step as described above are a first population of TILs *i.e.* which are a primary cell population.

[00619] The "Step" Designations A, B, C, *etc.*, below are in reference to the non-limiting example in Figure 85 (in particular, *e.g.*, Figure 85B) and in reference to certain non-limiting embodiments described herein. The ordering of the Steps below and in Figure 85 (in particular, *e.g.*, Figure 85B) is exemplary and any combination or order of steps, as well as additional steps, repetition of steps, and/or omission of steps is contemplated by the present application and the methods disclosed herein.

[00620] In some embodiments, if the tumor is metastatic and the primary lesion has been efficiently treated/removed in the past, removal of one of the metastatic lesions may be needed. In some embodiments, the least invasive approach is to remove a skin lesion, or a lymph node on the neck or axillary area when available. In some embodiments, a skin lesion is removed or small biopsy thereof is removed. In some embodiments, a lymph node or small biopsy thereof is removed. In some embodiments, a lung or liver metastatic lesion, or an intra-abdominal or thoracic lymph node or small biopsy can thereof can be employed.

[00621] In some embodiments, the tumor is a melanoma. In some embodiments, the small biopsy for a melanoma comprises a mole or portion thereof.

[00622] In some embodiments, the small biopsy is a punch biopsy. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin. around a suspicious mole. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin, and a round piece of skin is removed. In some embodiments, the small biopsy is a punch biopsy and round portion of the tumor is removed.

[00623] In some embodiments, the small biopsy is an excisional biopsy. In some embodiments, the small biopsy is an excisional biopsy and the entire mole or growth is removed. In some embodiments, the small biopsy is an excisional biopsy and the entire mole or growth is removed along with a small border of normal-appearing skin.

[00624] In some embodiments, the small biopsy is an incisional biopsy. In some embodiments, the small biopsy is an incisional biopsy and only the most irregular part of a mole or growth is taken. In some embodiments, the small biopsy is an incisional biopsy and the incisional biopsy is used when other techniques can't be completed, such as if a suspicious mole is very large.

[00625] In some embodiments, the small biopsy is a lung biopsy. In some embodiments, the small biopsy is obtained by bronchoscopy. Generally, bronchoscopy, the patient is put under anesthesia, and a small goes through the nose or mouth, down the throat, and into the bronchial passages, where small tools are used to remove some tissue. In some embodiments, where the tumor or growth cannot be reached via bronchoscopy, a transthoracic needle biopsy can be employed. Generally, for a transthoracic needle biopsy, the patient is also under anesthesia and a needle is inserted through the skin directly into the suspicious spot to remove a small sample of tissue. In some embodiments, a transthoracic needle biopsy may require interventional radiology (for example, the use of x-rays or CT scan to guide the needle). In some embodiments, the small biopsy is obtained by needle biopsy. In some embodiments, the small biopsy is obtained endoscopic ultrasound (for example, an endoscope with a light and is placed through the mouth into the esophagus). In some embodiments, the small biopsy is obtained surgery.

[00626] In some embodiments, the small biopsy is a head and neck biopsy. In some embodiments, the small biopsy is an incisional biopsy. In some embodiments, the small biopsy is an incisional biopsy, wherein a small piece of tissue is cut from an abnormal-looking area. In some embodiments, if the abnormal region is easily accessed, the sample may be taken without hospitalization. In some embodiments, if the tumor is deeper inside the mouth or throat, the biopsy may need to be done in an operating room, with general anesthesia. In some embodiments, the small biopsy is an excisional biopsy. In some embodiments, the small biopsy is an excisional biopsy, wherein the whole area is removed. In some embodiments, the small biopsy is a fine needle aspiration (FNA). In some embodiments, the small biopsy is a fine needle aspiration (FNA), wherein a very thin needle attached to a syringe is used to extract (aspirate) cells from a tumor or lump. In some embodiments, the small biopsy is a punch biopsy. In some embodiments, the small biopsy is a punch biopsy, wherein punch forceps are used to remove a piece of the suspicious area.

[00627] In some embodiments, the small biopsy is a cervical biopsy. In some embodiments, the small biopsy is obtained via colposcopy. Generally, colposcopy methods employ the use

of a lighted magnifying instrument attached to magnifying binoculars (a colposcope) which is then used to biopsy a small section of the surface of the cervix. In some embodiments, the small biopsy is a conization/cone biopsy. In some embodiments, the small biopsy is a conization/cone biopsy, wherein an outpatient surgery may be needed to remove a larger piece of tissue from the cervix. In some embodiments, the cone biopsy, in addition to helping to confirm a diagnosis, a cone biopsy can serve as an initial treatment.

[00628] The term "solid tumor" refers to an abnormal mass of tissue that usually does not contain cysts or liquid areas. Solid tumors may be benign or malignant. The term "solid tumor cancer refers to malignant, neoplastic, or cancerous solid tumors. Solid tumor cancers include, but are not limited to, sarcomas, carcinomas, and lymphomas, such as cancers of the lung, breast, triple negative breast cancer, prostate, colon, rectum, and bladder. In some embodiments, the cancer is selected from cervical cancer, head and neck cancer, glioblastoma, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma. The tissue structure of solid tumors includes interdependent tissue compartments including the parenchyma (cancer cells) and the supporting stromal cells in which the cancer cells are dispersed and which may provide a supporting microenvironment.

[00629] In some embodiments, the sample from the tumor is obtained as a fine needle aspirate (FNA), a core biopsy, or a small biopsy (including, for example, a punch biopsy). In some embodiments, sample is placed first into a G-Rex 10. In some embodiments, sample is placed first into a G-Rex 10 when there are 1 or 2 core biopsy and/or small biopsy samples. In some embodiments, sample is placed first into a G-Rex 100 when there are 3, 4, 5, 6, 8, 9, or 10 or more core biopsy and/or small biopsy samples. In some embodiments, sample is placed first into a G-Rex 500 when there are 3, 4, 5, 6, 8, 9, or 10 or more core biopsy and/or small biopsy samples.

[00630] The FNA can be obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma. In some embodiments, the FNA is obtained from a lung tumor, such as a lung tumor from a patient with non-small cell lung cancer (NSCLC). In some cases, the patient with NSCLC has previously undergone a surgical treatment.

[00631] TILs described herein can be obtained from an FNA sample. In some cases, the FNA sample is obtained or isolated from the patient using a fine gauge needle ranging

from a 18 gauge needle to a 25 gauge needle. The fine gauge needle can be 18 gauge, 19 gauge, 20 gauge, 21 gauge, 22 gauge, 23 gauge, 24 gauge, or 25 gauge. In some embodiments, the FNA sample from the patient can contain at least 400,000 TILs, *e.g.*, 400,000 TILs, 450,000 TILs, 500,000 TILs, 550,000 TILs, 600,000 TILs, 650,000 TILs, 700,000 TILs, 750,000 TILs, 800,000 TILs, 850,000 TILs, 900,000 TILs, 950,000 TILs, or more.

[00632] In some cases, the TILs described herein are obtained from a core biopsy sample. In some cases, the small biopsy or core biopsy sample is obtained or isolated from the patient using a surgical or medical needle ranging from a 11 gauge needle to a 16 gauge needle. The needle can be 11 gauge, 12 gauge, 13 gauge, 14 gauge, 15 gauge, or 16 gauge. In some embodiments, the core biopsy sample from the patient can contain at least 400,000 TILs, *e.g.*, 400,000 TILs, 450,000 TILs, 500,000 TILs, 550,000 TILs, 600,000 TILs, 650,000 TILs, 700,000 TILs, 750,000 TILs, 800,000 TILs, 850,000 TILs, 900,000 TILs, 950,000 TILs, or more.

[00633] In some embodiments, the TILs are obtained from tumor digests. In some embodiments, tumor digests were generated by incubation in enzyme media, for example but not limited to RPMI 1640, 2mM GlutaMAX, 10 μg/mL gentamicin, 30 U/mL DNase, and 1.0 mg/mL collagenase, followed by mechanical dissociation (GentleMACS, Miltenyi Biotec, Auburn, CA). After placing the tumor in enzyme media, the tumor can be mechanically dissociated for approximately 1 minute. The solution can then be incubated for 30 minutes at 37 °C in 5% CO_2 and it then mechanically disrupted again for approximately 1 minute. After being incubated again for 30 minutes at 37 °C in 5% CO_2, the tumor can be mechanically disrupted a third time for approximately 1 minute. In some embodiments, after the third mechanical disruption if large pieces of tissue were present, 1 or 2 additional mechanical dissociations were applied to the sample, with or without 30 additional minutes of incubation at 37 °C in 5% CO_2. In some embodiments, at the end of the final incubation if the cell suspension contained a large number of red blood cells or dead cells, a density gradient separation using Ficoll can be performed to remove these cells.

[00634] In some embodiments, the harvested cell suspension prior to the first expansion step is called a "primary cell population" or a "freshly harvested" cell population.

[00635] In some embodiments, cells can be optionally frozen after sample harvest and stored frozen prior to entry into the expansion described in Step B, which is described in further detail below, as well as exemplified in Figure 85.

A. STEP A: Obtain Patient tumor sample

[00636] In general, TILs are initially obtained from a patient tumor sample ("primary TILs") obtained by a core biopsy or similar procedure and are then expanded into a larger population for further manipulation as described herein, optionally cryopreserved, and optionally evaluated for phenotype and metabolic parameters.

[00637] A patient tumor sample may be obtained using methods known in the art, generally via surgical resection, core biopsy, needle biopsy or other means for obtaining a sample that contains a mixture of tumor and TIL cells. In general, the tumor sample may be from any solid tumor, including primary tumors, invasive tumors or metastatic tumors. In some embodiments, the sample can be from multiple small tumor samples or biopsies. In some embodiments, the sample can comprise multiple tumor samples (such as multiple cores) from a single tumor from the same patient. In some embodiments, the sample can comprise multiple tumor samples from one, two, three, or four tumors from the same patient (such as core biopsies obtained from multiple lesions in metastatic disease). In some embodiments, the sample can comprise multiple tumor samples from multiple tumors from the same patient. The tumor sample may also be a liquid tumor, such as a tumor obtained from a hematological malignancy. The solid tumor may be of any cancer type, including, but not limited to, breast, pancreatic, prostate, colorectal, lung, brain, renal, stomach, and skin (including but not limited to squamous cell carcinoma, basal cell carcinoma, and melanoma). In some embodiments, the cancer is selected from cervical cancer, head and neck cancer (including, for example, head and neck squamous cell carcinoma (HNSCC)), glioblastoma (GBM), gastrointestinal cancer, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma. In some embodiments, useful TILs are obtained from malignant melanoma tumors, as these have been reported to have particularly high levels of TILs.

[00638] As indicated above, in some embodiments, the TILs are derived from solid tumor cores or fragments. In some embodiments, the solid tumor cores or fragments are subjected to enzymatic digestion. In some embodiments, the tumor cores or fragments are digested in in an enzyme mixture comprising collagenase, DNase, and hyaluronidase. In some

embodiments, the tumor cores or fragments are digested in in an enzyme mixture comprising collagenase, DNase, and hyaluronidase for 1-2 hours. In some embodiments, the tumors are digested in in an enzyme mixture comprising collagenase, DNase, and hyaluronidase for 1-2 hours at 37°C, 5% CO_2. In some embodiments, the tumor cores or fragments are digested in in an enzyme mixture comprising collagenase, DNase, and hyaluronidase for 1-2 hours at 37°C, 5% CO_2 with rotation. In some embodiments, the tumor cores or fragments are digested overnight with constant rotation. In some embodiments, the tumor cores or fragments are digested overnight at 37°C, 5% CO_2 with constant rotation. In some embodiments, the tumor cores or fragments are combined with with the enzymes to form a tumor digest reaction mixture.

[00639] In some embodiments, the TILs are not obtained from tumor digests. In some embodiments, the solid tumor cores are not fragmented.

[00640] In some embodiments, the tumor cores or fragments are reconstituted with the lyophilized enzymes in a sterile buffer. In some embodiments, the buffer is sterile HBSS.

[00641] In some embodiments, the enxyme mixture comprises collagenase. In some embodiments, the collagenase is collagenase IV. In some embodiments, the working stock for the collagenase is a 100 mg/mL 10X working stock.

[00642] In some embodiments, the enzyme mixture comprises DNAse. In some embodiments, the working stock for the DNAse is a 10,000IU/mL 10X working stock.

[00643] In some embodiments, the enzyme mixture comprises hyaluronidase. In some embodiments, the working stock for the hyaluronidase is a 10 mg/mL 10X working stock.

[00644] In some embodiments, the enzyme mixture comprises 10 mg/mL collagenase, 1000 IU/mL DNAse, and 1 mg/mL hyaluronidase.

[00645] In some embodiments, the enzyme mixture comprises 10 mg/mL collagenase, 500 IU/mL DNAse, and 1 mg/mL hyaluronidase.

[00646] In general, the cell suspension obtained from the tumor core or fragment is called a "primary cell population" or a "freshly obtained" or a "freshly isolated" cell population. In certain embodiments, the freshly obtained cell population of TILs is exposed to a cell culture medium comprising antigen presenting cells, IL-12 and OKT-3.

[00647] In some embodiments, TILs can be initially cultured from enzymatic tumor core or fragment digests and tumor cores or fragments obtained from patients. In an embodiment,

TILs can be initially cultured from enzymatic tumor core or fragment digests and tumor cores or fragments obtained from patients.

[00648] In some embodiments, the TILs are obtained from tumor fragment or core digests. In some embodiments, tumor fragment or core digests are generated by incubation in enzyme media, for example but not limited to RPMI 1640, 2 mM GlutaMAX, 10 mg/mL gentamicin, 30 U/mL DNase, and 1.0 mg/mL collagenase, followed by mechanical dissociation (GentleMACS, Miltenyi Biotec, Auburn, CA). After placing the tumor fragment or core in enzyme media, the tumor fragment or core can be mechanically dissociated for approximately 1 minute. The solution can then be incubated for 30 minutes at 37 °C in 5% CO_2 and it then mechanically disrupted again for approximately 1 minute. After being incubated again for 30 minutes at 37 °C in 5% CO_2, the tumor fragment or core can be mechanically disrupted a third time for approximately 1 minute. In some embodiments, after the third mechanical disruption if large pieces of tissue were present, 1 or 2 additional mechanical dissociations were applied to the sample, with or without 30 additional minutes of incubation at 37 °C in 5% CO_2. In some embodiments, at the end of the final incubation if the cell suspension contained a large number of red blood cells or dead cells, a density gradient separation using Ficoll can be performed to remove these cells.

[00649] In some embodiments, if the tumor is metastatic and the primary lesion has been efficiently treated/removed in the past, removal of one of the metastatic lesions may be needed. In some embodiments, the least invasive approach is to remove a skin lesion, or a lymph node on the neck or axillary area when available. In some embodiments, a skin lesion is removed or small biopsy thereof is removed. In some embodiments, a lymph node or small biopsy thereof is removed. In some embodiments, a lung or liver metastatic lesion, or an intra-abdominal or thoracic lymph node or small biopsy can thereof can be employed.

[00650] In some embodiments, the tumor is a melanoma. In some embodiments, the small biopsy for a melanoma comprises a mole or portion thereof.

[00651] In some embodiments, the small biopsy is a punch biopsy. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin. around a suspicious mole. In some embodiments, the punch biopsy is obtained with a circular blade pressed into the skin, and a round piece of skin is removed. In some embodiments, the small biopsy is a punch biopsy and round portion of the tumor is removed.

[00652] In some embodiments, the small biopsy is an excisional biopsy. In some embodiments, the small biopsy is an excisional biopsy and the entire mole or growth is removed. In some embodiments, the small biopsy is an excisional biopsy and the entire mole or growth is removed along with a small border of normal-appearing skin.

[00653] In some embodiments, the small biopsy is an incisional biopsy. In some embodiments, the small biopsy is an incisional biopsy and only the most irregular part of a mole or growth is taken. In some embodiments, the small biopsy is an incisional biopsy and the incisional biopsy is used when other techniques can't be completed, such as if a suspicious mole is very large.

[00654] In some embodiments, the small biopsy is a lung biopsy. In some embodiments, the small biopsy is obtained by bronchoscopy. Generally, bronchoscopy, the patient is put under anesthesia, and a small goes through the nose or mouth, down the throat, and into the bronchial passages, where small tools are used to remove some tissue. In some embodiments, where the tumor or growth cannot be reached via bronchoscopy, a transthoracic needle biopsy can be employed. Generally, for a transthoracic needle biopsy, the patient is also under anesthesia and a needle is inserted through the skin directly into the suspicious spot to remove a small sample of tissue. In some embodiments, a transthoracic needle biopsy may require interventional radiology (for example, the use of x-rays or CT scan to guide the needle). In some embodiments, the small biopsy is obtained by needle biopsy. In some embodiments, the small biopsy is obtained endoscopic ultrasound (for example, an endoscope with a light and is placed through the mouth into the esophagus). In some embodiments, the small biopsy is obtained surgery.

[00655] In some embodiments, the small biopsy is a head and neck biopsy. In some embodiments, the small biopsy is an incisional biopsy. In some embodiments, the small biopsy is an incisional biopsy, wherein a small piece of tissue is cut from an abnormal-looking area. In some embodiments, if the abnormal region is easily accessed, the sample may be taken without hospitalization. In some embodiments, if the tumor is deeper inside the mouth or throat, the biopsy may need to be done in an operating room, with general anesthesia. In some embodiments, the small biopsy is an excisional biopsy. In some embodiments, the small biopsy is an excisional biopsy, wherein the whole area is removed. In some embodiments, the small biopsy is a fine needle aspiration (FNA). In some embodiments, the small biopsy is a fine needle aspiration (FNA), wherein a very thin needle attached to a syringe is used to extract (aspirate) cells from a tumor or lump. In some

embodiments, the small biopsy is a punch biopsy. In some embodiments, the small biopsy is a punch biopsy, wherein punch forceps are used to remove a piece of the suspicious area.

[00656] In some embodiments, the small biopsy is a cervical biopsy. In some embodiments, the small biopsy is obtained via colposcopy. Generally, colposcopy methods employ the use of a lighted magnifying instrument attached to magnifying binoculars (a colposcope) which is then used to biopsy a small section of the surface of the cervix. In some embodiments, the small biopsy is a conization/cone biopsy. In some embodiments, the small biopsy is a conization/cone biopsy, wherein an outpatient surgery may be needed to remove a larger piece of tissue from the cervix. In some embodiments, the cone biopsy, in addition to helping to confirm a diagnosis, a cone biopsy can serve as an initial treatment.

[00657] The term "solid tumor" refers to an abnormal mass of tissue that usually does not contain cysts or liquid areas. Solid tumors may be benign or malignant. The term "solid tumor cancer refers to malignant, neoplastic, or cancerous solid tumors. Solid tumor cancers include, but are not limited to, sarcomas, carcinomas, and lymphomas, such as cancers of the lung, breast, triple negative breast cancer, prostate, colon, rectum, and bladder. In some embodiments, the cancer is selected from cervical cancer, head and neck cancer, glioblastoma, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma. The tissue structure of solid tumors includes interdependent tissue compartments including the parenchyma (cancer cells) and the supporting stromal cells in which the cancer cells are dispersed and which may provide a supporting microenvironment.

[00658] In some embodiments, the sample from the tumor is obtained as a fine needle aspirate (FNA), a core biopsy, a small biopsy (including, for example, a punch biopsy). In some embodiments, sample is placed first into a G-Rex 10. In some embodiments, sample is placed first into a G-Rex 10 when there are 1 or 2 core biopsy and/or small biopsy samples. In some embodiments, sample is placed first into a G-Rex 100 when there are 3, 4, 5, 6, 8, 9, or 10 or more core biopsy and/or small biopsy samples. In some embodiments, sample is placed first into a G-Rex 500 when there are 3, 4, 5, 6, 8, 9, or 10 or more core biopsy and/or small biopsy samples.

[00659] The FNA can be obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma. In some embodiments, the FNA is obtained from a lung tumor, such as a lung

tumor from a patient with non-small cell lung cancer (NSCLC). In some cases, the patient with NSCLC has previously undergone a surgical treatment.

[00660] TILs described herein can be obtained from an FNA sample. In some cases, the FNA sample is obtained or isolated from the patient using a fine gauge needle ranging from an 18 gauge needle to a 25 gauge needle. The fine gauge needle can be 18 gauge, 19 gauge, 20 gauge, 21 gauge, 22 gauge, 23 gauge, 24 gauge, or 25 gauge. In some embodiments, the FNA sample from the patient can contain at least 400,000 TILs, *e.g.*, 400,000 TILs, 450,000 TILs, 500,000 TILs, 550,000 TILs, 600,000 TILs, 650,000 TILs, 700,000 TILs, 750,000 TILs, 800,000 TILs, 850,000 TILs, 900,000 TILs, 950,000 TILs, or more.

[00661] In some cases, the TILs described herein are obtained from a core biopsy sample. In some cases, the core biopsy or small biopsy sample is obtained or isolated from the patient using a surgical or medical needle ranging from an 11 gauge needle to a 16 gauge needle. The needle can be 11 gauge, 12 gauge, 13 gauge, 14 gauge, 15 gauge, or 16 gauge. In some embodiments, the core biopsy sample from the patient can contain at least 400,000 TILs, *e.g.*, 400,000 TILs, 450,000 TILs, 500,000 TILs, 550,000 TILs, 600,000 TILs, 650,000 TILs, 700,000 TILs, 750,000 TILs, 800,000 TILs, 850,000 TILs, 900,000 TILs, 950,000 TILs, or more.

[00662] In some embodiments, the TILs are obtained from tumor core or fragment digests. In some embodiments, tumor core or fragment digests were generated by incubation in enzyme media, for example but not limited to RPMI 1640, 2mM GlutaMAX, 10 µg/mL gentamicin, 30 U/mL DNase, and 1.0 mg/mL collagenase, followed by mechanical dissociation (GentleMACS, Miltenyi Biotec, Auburn, CA). After placing the tumor core or fragment in enzyme media, the tumor can be mechanically dissociated for approximately 1 minute. The solution can then be incubated for 30 minutes at 37 °C in 5% CO_2 and it then mechanically disrupted again for approximately 1 minute. After being incubated again for 30 minutes at 37 °C in 5% CO_2, the tumor core or fragment can be mechanically disrupted a third time for approximately 1 minute. In some embodiments, after the third mechanical disruption if large pieces of tissue were present, 1 or 2 additional mechanical dissociations were applied to the sample, with or without 30 additional minutes of incubation at 37 °C in 5% CO_2. In some embodiments, at the end of the final incubation if the cell suspension contained a large number of red blood cells or dead cells, a density gradient separation using Ficoll can be performed to remove these cells.

[00663] In some embodiments, the cell suspension prior to the priming first expansion step is called a "primary cell population" or a "freshly obtained" or "freshly isolated" cell population.

[00664] In some embodiments, cells can be optionally frozen after sample isolation (*e.g.*, after obtaining the tumor sample and/or after obtaining the cell suspension from the tumor sample) and stored frozen prior to entry into the expansion described in Step B, which is described in further detail below, as well as exemplified in Figure 85 (in particular, *e.g.*, Figure 85B).

B. STEP B: Priming First Expansion

[00665] In some embodiments, the present methods provide for younger TILs, which may provide additional therapeutic benefits over older TILs (*i.e.*, TILs which have further undergone more rounds of replication prior to administration to a subject/patient). Features of young TILs have been described in the literature, for example Donia, *et al.*, *Scandinavian Journal of Immunology,* 75:157–167 (2012); Dudley *et al.*, *Clin Cancer Res,* 16:6122-6131 (2010); Huang *et al.*, *J Immunother*, 28(3):258–267 (2005); Besser *et al.*, *Clin Cancer Res*, 19(17):OF1-OF9 (2013); Besser *et al.*, *J Immunother* 32:415–423 (2009); Robbins, *et al.*, *J Immunol* 2004; 173:7125-7130; Shen et al., J Immunother, 30:123–129 (2007); Zhou, et al., *J Immunother*, 28:53–62 (2005); and Tran, *et al.*, J Immunother, 31:742–751 (2008), all of which are incorporated herein by reference in their entireties.

[00666] After biopsy or digestion of tumor fragments and/or tumor cores, for example such as described in Step A of Figure 85 (in particular, *e.g.*, Figure 85B), the resulting cells are cultured in serum containing IL-2, OKT-3, and feeder cells (*e.g.*, antigen-presenting feeder cells), under conditions that favor the growth of TILs over tumor and other cells. In some embodiments, the IL-2, OKT-3, and feeder cells are added at culture initiation along with the tumor fragment or core digest and/or tumor fragments or cores (*e.g.*, at Day 0). In some embodiments, the tumor fragments or core digests and/or tumor fragments or cores are incubated in a container with up to 60 fragments or cores per container and with 6000 IU/mL of IL-2. This primary cell population is cultured for a period of days, generally from 1 to 7 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, priming first expansion occurs for a period of 1 to 7 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, this priming first expansion occurs for a period of 5 to 7 days, resulting in a bulk TIL population,

generally about 1×10^8 bulk TIL cells. In some embodiments, this priming first expansion occurs for a period of about 6 to 7 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, this priming first expansion occurs for a period of about 7 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells.

[00667] In a preferred embodiment, expansion of TILs may be performed using a priming first expansion step (for example such as those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include processes referred to as pre-REP or priming REP and which contains feeder cells from Day 0 and/or from culture initiation) as described below and herein, followed by a rapid second expansion (Step D, including processes referred to as rapid expansion protocol (REP) steps) as described below under Step D and herein, followed by optional cryopreservation, and followed by a second Step D (including processes referred to as restimulation REP steps) as described below and herein. The TILs obtained from this process may be optionally characterized for phenotypic characteristics and metabolic parameters as described herein. In some embodiments, the tumor fragment is between about 1 mm^3 and 10 mm^3.

[00668] In some embodiments, the first expansion culture medium is referred to as "CM", an abbreviation for culture media. In some embodiments, CM for Step B consists of RPMI 1640 with GlutaMAX, supplemented with 10% human AB serum, 25 mM Hepes, and 10 µg/mL gentamicin.

[00669] In some embodiments, there are less than or equal to 240 tumor fragments or cores. In some embodiments, there are less than or equal to 240 tumor fragments or cores placed in less than or equal to 4 containers. In some embodiments, the containers are GREX100 MCS flasks. In some embodiments, less than or equal to 60 tumor fragments or cores are placed in 1 container. In some embodiments, each container comprises less than or equal to 500 mL of media per container. In some embodiments, the media comprises IL-2. In some embodiments, the media comprises 6000 IU/mL of IL-2. In some embodiments, the media comprises antigen-presenting feeder cells (also referred to herein as "antigen-presenting cells"). In some embodiments, the media comprises 2.5×10^8 antigen-presenting feeder cells per container. In some embodiments, the media comprises OKT-3. In some embodiments, the media comprises 30 ng/mL of OKT-3 per container. In some embodiments, the container is a GREX100 MCS flask. In some embodiments, the media comprises 6000 IU/mL of IL-2, 30 ng/mL of OKT-3, and 2.5×10^8 antigen-presenting feeder cells. In some embodiments, the

media comprises 6000 IU/mL of IL-2, 30 ng/mL of OKT-3, and 2.5×10^8 antigen-presenting feeder cells per container.

[00670] After preparation of the tumor fragments or cores, the resulting cells (*i.e.*, which are from the fragments or cores, and which constitute a primary cell population) are cultured in media containing IL-2, antigen-presenting feeder cells and OKT-3 under conditions that favor the growth of TILs over tumor and other cells and which allow for TIL priming and accelerated growth from initiation of the culture on Day 0. In some embodiments, the tumor fragment or core digests and/or tumor fragments or cores are incubated in with 6000 IU/mL of IL-2, as well as antigen-presenting feeder cells and OKT-3. This primary cell population is cultured for a period of days, generally from 1 to 7 days, resulting in a bulk TIL population, generally about 1×10^8 bulk TIL cells. In some embodiments, the growth media during the priming first expansion comprises IL-2 or a variant thereof, as well as antigen-presenting feeder cells and OKT-3. In some embodiments, the IL-2 is recombinant human IL-2 (rhIL-2). In some embodiments the IL-2 stock solution has a specific activity of $20\text{-}30\times10^6$ IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 20×10^6 IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 25×10^6 IU/mg for a 1 mg vial. In some embodiments the IL-2 stock solution has a specific activity of 30×10^6 IU/mg for a 1 mg vial. In some embodiments, the IL- 2 stock solution has a final concentration of $4\text{-}8\times10^6$ IU/mg of IL-2. In some embodiments, the IL- 2 stock solution has a final concentration of $5\text{-}7\times10^6$ IU/mg of IL-2. In some embodiments, the IL- 2 stock solution has a final concentration of 6×10^6 IU/mg of IL-2. In some embodiments, the IL-2 stock solution is prepared as described in Example 5. In some embodiments, the priming first expansion culture media comprises about 10,000 IU/mL of IL-2, about 9,000 IU/mL of IL-2, about 8,000 IU/mL of IL-2, about 7,000 IU/mL of IL-2, about 6000 IU/mL of IL-2 or about 5,000 IU/mL of IL-2. In some embodiments, the priming first expansion culture media comprises about 9,000 IU/mL of IL-2 to about 5,000 IU/mL of IL-2. In some embodiments, the priming first expansion culture media comprises about 8,000 IU/mL of IL-2 to about 6,000 IU/mL of IL-2. In some embodiments, the priming first expansion culture media comprises about 7,000 IU/mL of IL-2 to about 6,000 IU/mL of IL-2. In some embodiments, the priming first expansion culture media comprises about 6,000 IU/mL of IL-2. In an embodiment, the cell culture medium further comprises IL-2. In some embodiments, the priming first expansion cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the priming first expansion cell culture medium further comprises IL-2. In a

preferred embodiment, the priming first expansion cell culture medium comprises about 3000 IU/mL of IL-2. In an embodiment, the priming first expansion cell culture medium comprises about 1000 IU/mL, about 1500 IU/mL, about 2000 IU/mL, about 2500 IU/mL, about 3000 IU/mL, about 3500 IU/mL, about 4000 IU/mL, about 4500 IU/mL, about 5000 IU/mL, about 5500 IU/mL, about 6000 IU/mL, about 6500 IU/mL, about 7000 IU/mL, about 7500 IU/mL, or about 8000 IU/mL of IL-2. In an embodiment, the priming first expansion cell culture medium comprises between 1000 and 2000 IU/mL, between 2000 and 3000 IU/mL, between 3000 and 4000 IU/mL, between 4000 and 5000 IU/mL, between 5000 and 6000 IU/mL, between 6000 and 7000 IU/mL, between 7000 and 8000 IU/mL, or about 8000 IU/mL of IL-2.

[00671] In some embodiments, priming first expansion culture media comprises about 500 IU/mL of IL-15, about 400 IU/mL of IL-15, about 300 IU/mL of IL-15, about 200 IU/mL of IL-15, about 180 IU/mL of IL-15, about 160 IU/mL of IL-15, about 140 IU/mL of IL-15, about 120 IU/mL of IL-15, or about 100 IU/mL of IL-15. In some embodiments, the priming first expansion culture media comprises about 500 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the priming first expansion culture media comprises about 400 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the priming first expansion culture media comprises about 300 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the priming first expansion culture media comprises about 200 IU/mL of IL-15. In some embodiments, the priming first expansion cell culture medium comprises about 180 IU/mL of IL-15. In an embodiment, the priming first expansion cell culture medium further comprises IL-15. In a preferred embodiment, the priming first expansion cell culture medium comprises about 180 IU/mL of IL-15.

[00672] In some embodiments, priming first expansion culture media comprises about 20 IU/mL of IL-21, about 15 IU/mL of IL-21, about 12 IU/mL of IL-21, about 10 IU/mL of IL-21, about 5 IU/mL of IL-21, about 4 IU/mL of IL-21, about 3 IU/mL of IL-21, about 2 IU/mL of IL-21, about 1 IU/mL of IL-21, or about 0.5 IU/mL of IL-21. In some embodiments, the priming first expansion culture media comprises about 20 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the priming first expansion culture media comprises about 15 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the priming first expansion culture media comprises about 12 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the priming first expansion culture media comprises about 10 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the priming first

expansion culture media comprises about 5 IU/mL of IL-21 to about 1 IU/mL of IL-21. In some embodiments, the priming first expansion culture media comprises about 2 IU/mL of IL-21. In some embodiments, the priming first expansion cell culture medium comprises about 1 IU/mL of IL-21. In some embodiments, the priming first expansion cell culture medium comprises about 0.5 IU/mL of IL-21. In an embodiment, the cell culture medium further comprises IL-21. In a preferred embodiment, the priming first expansion cell culture medium comprises about 1 IU/mL of IL-21.

[00673] In an embodiment, the priming first expansion cell culture medium comprises OKT-3 antibody. In some embodiments, the priming first expansion cell culture medium comprises about 30 ng/mL of OKT-3 antibody. In an embodiment, the priming first expansion cell culture medium comprises about 0.1 ng/mL, about 0.5 ng/mL, about 1 ng/mL, about 2.5 ng/mL, about 5 ng/mL, about 7.5 ng/mL, about 10 ng/mL, about 15 ng/mL, about 20 ng/mL, about 25 ng/mL, about 30 ng/mL, about 35 ng/mL, about 40 ng/mL, about 50 ng/mL, about 60 ng/mL, about 70 ng/mL, about 80 ng/mL, about 90 ng/mL, about 100 ng/mL, about 200 ng/mL, about 500 ng/mL, and about 1 µg/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises between 0.1 ng/mL and 1 ng/mL, between 1 ng/mL and 5 ng/mL, between 5 ng/mL and 10 ng/mL, between 10 ng/mL and 20 ng/mL, between 20 ng/mL and 30 ng/mL, between 30 ng/mL and 40 ng/mL, between 40 ng/mL and 50 ng/mL, and between 50 ng/mL and 100 ng/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises between 15 ng/ml and 30 ng/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises 30 ng/mL of OKT-3 antibody. In some embodiments, the OKT-3 antibody is muromonab.

TABLE 18: Amino acid sequences of muromonab (exemplary OKT-3 antibody)

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:1 Muromonab heavy chain	QVQLQQSGAE LARPGASVKM SCKASGYTFT RYTMHWVKQR PGQGLEWIGY INPSRGYTNY NQKFKDKATL TTDKSSSTAY MQLSSLTSED SAVYYCARYY DDHYCLDYWG QGTTLTVSSA KTTAPSVYPL APVCGGTTGS SVTLGCLVKG YFPEPVTLTW NSGSLSSGVH TFPAVLQSDL YTLSSSVTVT SSTWPSQSIT CNVAHPASST KVDKKIEPRP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK	60 120 180 240 300 360 420 450
SEQ ID NO:2 Muromonab light chain	QIVLTQSPAI MSASPGEKVT MTCSASSSVS YMNWYQQKSG TSPKRWIYDT SKLASGVPAH FRGSGSGTSY SLTISGMEAE DAATYYCQQW SSNPFTFGSG TKLEINRADT APTVSIFPPS SEQLTSGGAS VVCFLNNFYP KDINVKWKID GSERQNGVLN SWTDQDSKDS TYSMSSTLTL TKDEYERHNS YTCEATHKTS TSPIVKSFNR NEC	60 120 180 213

[00674] In some embodiments, the priming first expansion cell culture medium comprises one or more TNFRSF agonists in a cell culture medium. In some embodiments, the TNFRSF agonist comprises a 4-1BB agonist. In some embodiments, the TNFRSF agonist is a 4-1BB agonist, and the 4-1BB agonist is selected from the group consisting of urelumab, utomilumab, EU-101, a fusion protein, and fragments, derivatives, variants, biosimilars, and combinations thereof. In some embodiments, the TNFRSF agonist is added at a concentration sufficient to achieve a concentration in the cell culture medium of between 0.1 μg/mL and 100 μg/mL. In some embodiments, the TNFRSF agonist is added at a concentration sufficient to achieve a concentration in the cell culture medium of between 20 μg/mL and 40 μg/mL.

[00675] In some embodiments, in addition to one or more TNFRSF agonists, the priming first expansion cell culture medium further comprises IL-2 at an initial concentration of about 3000 IU/mL and OKT-3 antibody at an initial concentration of about 30 ng/mL, and wherein the one or more TNFRSF agonists comprises a 4-1BB agonist. In some embodiments, in addition to one or more TNFRSF agonists, the priming first expansion cell culture medium further comprises IL-2 at an initial concentration of about 6000 IU/mL and OKT-3 antibody at an initial concentration of about 30 ng/mL, and wherein the one or more TNFRSF agonists comprises a 4-1BB agonist.

[00676] In some embodiments, the priming first expansion culture medium is referred to as "CM", an abbreviation for culture media. In some embodiments, it is referred to as CM1 (culture medium 1). In some embodiments, CM consists of RPMI 1640 with GlutaMAX, supplemented with 10% human AB serum, 25 mM Hepes, and 10 μg/mL gentamicin. In some embodiments, the CM is the CM1 described in the Examples, *see*, Examples 1 and 14. In some embodiments, the priming first expansion occurs in an initial cell culture medium or a first cell culture medium. In some embodiments, the priming first expansion culture medium or the initial cell culture medium or the first cell culture medium comprises IL-2, OKT-3 and antigen-presenting feeder cells (also referred to herein as feeder cells).

[00677] In some embodiments, the priming first expansion (including processes such as for example those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 1 to 7 days, as discussed in the examples and figures. In some embodiments, the priming first expansion (including processes such as for example those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 2 to 7 days. In some embodiments, the priming first expansion

(including processes such as for example those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 3 to 7 days. In some embodiments, the priming first expansion (including processes such as for example those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 4 to 7 days. In some embodiments, the priming first expansion (including processes such as for example those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 5 to 7 days. In some embodiments, the priming first expansion (including processes such as for example those described in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 6 to 7 days. In some embodiments, the priming first expansion (including processes such as for example those provided in Step B of Figure 85 (in particular, *e.g.*, Figure 85B), which can include those sometimes referred to as the pre-REP or priming REP) process is 7 days.

[00678] In some embodiments, the priming first TIL expansion can proceed for 1 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 2 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 3 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 4 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 5 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 6 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 1 days to 10 days from when

tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 2 days to 10 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 3 days to 10 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 4 days to 7 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 5 days to 10 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 6 days to 10 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 8 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 9 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 10 days from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the small biopsy (*e.g.*, the core biopsy) is removed from the culture at about day 1, 2, or 3. In some embodiments, the small biopsy (*e.g.*, the core biopsy) is removed from the culture at day 3.

[00679] In some embodiments, the priming first TIL expansion can proceed for 8 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 9 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 10 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 11 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL

expansion can proceed for 12 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 13 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 14 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 15 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated. In some embodiments, the priming first TIL expansion can proceed for 16 days to 17 days from when tumor cores or fragments are added to the cell culture and/or when the first priming expansion step is initiated.

[00680] In some embodiments, the priming first expansion of the TILs can proceed for 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9, days, or 10 days. In some embodiments, the first TIL expansion can proceed for 1 day to 7 days. In some embodiments, the first TIL expansion can proceed for 2 days to 7 days. In some embodiments, the first TIL expansion can proceed for 3 days to 7 days. In some embodiments, the first TIL expansion can proceed for 4 days to 7 days. In some embodiments, the first TIL expansion can proceed for 5 days to 7 days. In some embodiments, the first TIL expansion can proceed for 6 days to 7 days. In some embodiments, the first TIL expansion can proceed for 1 day to 10 days. In some embodiments, the first TIL expansion can proceed for 2 days to 10 days. In some embodiments, the first TIL expansion can proceed for 3 days to 10 days. In some embodiments, the first TIL expansion can proceed for 4 days to 10 days. In some embodiments, the first TIL expansion can proceed for 5 days to 10 days. In some embodiments, the first TIL expansion can proceed for 6 days to 10 days. In some embodiments, the first TIL expansion can proceed for 7 days to 10 days. In some embodiments, the first TIL expansion can proceed for 8 days to 10 days. In some embodiments, the first TIL expansion can proceed for 9 days to 10 days. In some embodiments, the first TIL expansion can proceed for 7 days. In some embodiments, the first TIL expansion can proceed for 8 days. In some embodiments, the first TIL expansion can proceed for 9 days. In some embodiments, the first TIL expansion can proceed for 10 days.

[00681] In some embodiments, the priming first expansion of the TILs can proceed for 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days or 17 days. In

some embodiments, the first TIL expansion can proceed for 8 days to 17 days. In some embodiments, the first TIL expansion can proceed for 9 days to 17 days. In some embodiments, the first TIL expansion can proceed for 10 days to 17 days. In some embodiments, the first TIL expansion can proceed for 11 days to 17 days. In some embodiments, the first TIL expansion can proceed for 12 days to 17 days. In some embodiments, the first TIL expansion can proceed for 13 days to 17 days. In some embodiments, the first TIL expansion can proceed for 14 days to 17 days. In some embodiments, the first TIL expansion can proceed for 15 days to 17 days. In some embodiments, the first TIL expansion can proceed for 16 days to 17 days. In some embodiments, the first TIL expansion can proceed for 17 days.

[00682] In some embodiments, a combination of IL-2, IL-7, IL-15, and/or IL-21 are employed as a combination during the priming first expansion. In some embodiments, IL-2, IL-7, IL-15, and/or IL-21 as well as any combinations thereof can be included during the priming first expansion, including, for example during Step B processes according to Figure 85 (in particular, *e.g.*, Figure 85B), as well as described herein. In some embodiments, a combination of IL-2, IL-15, and IL-21 are employed as a combination during the priming first expansion. In some embodiments, IL-2, IL-15, and IL-21 as well as any combinations thereof can be included during Step B processes according to Figure 85 (in particular, *e.g.*, Figure 85B) and as described herein.

[00683] In some embodiments, the priming first expansion, for example, Step B according to Figure 85 (in particular, *e.g.*, Figure 85B), is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a bioreactor is employed. In some embodiments, a bioreactor is employed as the container. In some embodiments, the bioreactor employed is for example a G-REX-10 or a G-REX-100. In some embodiments, the bioreactor employed is a G-REX-100. In some embodiments, the bioreactor employed is a G-REX-10.

1. Feeder Cells and Antigen Presenting Cells

[00684] In an embodiment, the priming first expansion procedures described herein (for example including expansion such as those described in Step B from Figure 85 (in particular, *e.g.*, Figure 85B), as well as those referred to as pre-REP or priming REP) require feeder cells (also referred to herein as "antigen-presenting cells") at the initiation of the TIL expansion and during the priming first expansion. In many embodiments, the feeder cells are peripheral

blood mononuclear cells (PBMCs) obtained from standard whole blood units from allogeneic healthy blood donors. The PBMCs are obtained using standard methods such as Ficoll-Paque gradient separation. In some embodiments, 2.5×10^8 feeder cells are used during the priming first expansion. In some embodiments, 2.5×10^8 feeder cells per container are used during the priming first expansion. In some embodiments, 2.5×10^8 feeder cells per GREX-10 are used during the priming first expansion. In some embodiments, 2.5×10^8 feeder cells per GREX-100 are used during the priming first expansion.

[00685] In general, the allogenic PBMCs are inactivated, either via irradiation or heat treatment, and used in the REP procedures, as described in the examples, which provides an exemplary protocol for evaluating the replication incompetence of irradiate allogeneic PBMCs.

[00686] In some embodiments, PBMCs are considered replication incompetent and acceptable for use in the TIL expansion procedures described herein if the total number of viable cells on day 14 is less than the initial viable cell number put into culture on day 0 of the priming first expansion.

[00687] In some embodiments, PBMCs are considered replication incompetent and acceptable for use in the TIL expansion procedures described herein if the total number of viable cells, cultured in the presence of OKT3 and IL-2, on day 7 have not increased from the initial viable cell number put into culture on day 0 of the priming first expansion. In some embodiments, the PBMCs are cultured in the presence of 30 ng/mL OKT3 antibody and 3000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 30 ng/mL OKT3 antibody and 6000 IU/mL IL-2.

[00688] In some embodiments, PBMCs are considered replication incompetent and acceptable for use in the TIL expansion procedures described herein if the total number of viable cells, cultured in the presence of OKT3 and IL-2, on day 7 have not increased from the initial viable cell number put into culture on day 0 of the priming first expansion. In some embodiments, the PBMCs are cultured in the presence of 5-60 ng/mL OKT3 antibody and 1000-6000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 10-50 ng/mL OKT3 antibody and 2000-5000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 20-40 ng/mL OKT3 antibody and 2000-4000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 25-35 ng/mL OKT3 antibody and 2500-3500 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence

of 30 ng/mL OKT3 antibody and 6000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 15 ng/mL OKT3 antibody and 3000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 15 ng/mL OKT3 antibody and 6000 IU/mL IL-2.

[00689] In some embodiments, the antigen-presenting feeder cells are PBMCs. In some embodiments, the antigen-presenting feeder cells are artificial antigen-presenting feeder cells. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is about 1 to 25, about 1 to 50, about 1 to 100, about 1 to 125, about 1 to 150, about 1 to 175, about 1 to 200, about 1 to 225, about 1 to 250, about 1 to 275, about 1 to 300, about 1 to 325, about 1 to 350, about 1 to 375, about 1 to 400, or about 1 to 500. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is between 1 to 50 and 1 to 300. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is between 1 to 100 and 1 to 200.

[00690] In an embodiment, the priming first expansion procedures described herein require a ratio of about 2.5×10^8 feeder cells to about 100×10^6 TILs. In another embodiment, the priming first expansion procedures described herein require a ratio of about 2.5×10^8 feeder cells to about 50×10^6 TILs. In yet another embodiment, the priming first expansion described herein require about 2.5×10^8 feeder cells to about 25×10^6 TILs. In yet another embodiment, the priming first expansion described herein require about 2.5×10^8 feeder cells. In yet another embodiment, the priming first expansion requires one-fourth, one-third, five-twelfths, or one-half of the number of feeder cells used in the rapid second expansion.

[00691] In some embodiments, the media in the priming first expansion comprises IL-2. In some embodiments, the media in the priming first expansion comprises 6000 IU/mL of IL-2. In some embodiments, the media in the priming first expansion comprises antigen-presenting feeder cells. In some embodiments, the media in the priming first expansion comprises 2.5×10^8 antigen-presenting feeder cells per container. In some embodiments, the media in the priming first expansion comprises OKT-3. In some embodiments, the media comprises 30 ng of OKT-3 per container. In some embodiments, the container is a GREX100 MCS flask. In some embodiments, the media comprises 6000 IU/mL of IL-2, 30 ng/mL of OKT-3, and 2.5×10^8 antigen-presenting feeder cells. In some embodiments, the media comprises 6000 IU/mL of IL-2, 30 ng/mL of OKT-3, and 2.5×10^8 antigen-presenting feeder cells per container. In some embodiments, the media comprises 500 mL of culture medium and 15 ug of OKT-3 per 2.5×10^8 antigen-presenting feeder cells per container. In some embodiments,

the media comprises 500 mL of culture medium and 15 ug of OKT-3 per container. In some embodiments, the container is a GREX100 MCS flask. In some embodiments, the media comprises 500 mL of culture medium and 6000 IU/mL of IL-2, 30 ng/mL of OKT-3, and 2.5 $\times 10^8$ antigen-presenting feeder cells. In some embodiments, the media comprises 500 mL of culture medium and 6000 IU/mL of IL-2, 15 ug of OKT-3, and 2.5 $\times 10^8$ antigen-presenting feeder cells per container. In some embodiments, the media comprises 500 mL of culture medium and 15 ug of OKT-3 per 2.5 $\times 10^8$ antigen-presenting feeder cells per container.

[00692] In an embodiment, the priming first expansion procedures described herein require an excess of feeder cells over TILs during the second expansion. In many embodiments, the feeder cells are peripheral blood mononuclear cells (PBMCs) obtained from standard whole blood units from allogeneic healthy blood donors. The PBMCs are obtained using standard methods such as Ficoll-Paque gradient separation. In an embodiment, artificial antigen-presenting (aAPC) cells are used in place of PBMCs.

[00693] In general, the allogenic PBMCs are inactivated, either via irradiation or heat treatment, and used in the TIL expansion procedures described herein, including the exemplary procedures described in the figures and examples.

[00694] In an embodiment, artificial antigen presenting cells are used in the priming first expansion as a replacement for, or in combination with, PBMCs.

2. Cytokines

[00695] The expansion methods described herein generally use culture media with high doses of a cytokine, in particular IL-2, as is known in the art.

[00696] Alternatively, using combinations of cytokines for the priming first expansion of TILs is additionally possible, with combinations of two or more of IL-2, IL-15 and IL-21 as is generally outlined in International Publication No. WO 2015/189356 and WO 2015/189357, hereby expressly incorporated by reference in their entirety. Thus, possible combinations include IL-2 and IL-15, IL-2 and IL-21, IL-15 and IL-21, and IL-2, IL-15 and IL-21, with the latter finding particular use in many embodiments. The use of combinations of cytokines specifically favors the generation of lymphocytes, and in particular T-cells as described therein.

TABLE 19: Amino acid sequences of interleukins.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:3 recombinant human IL-2 (rhIL-2)	MAPTSSSTKK TQLQLEHLLL DLQMILNGIN NYKNPKLTRM LTFKFYMPKK ATELKHLQCL EEELKPLEEV LNLAQSKNFH LRPRDLISNI NVIVLELKGS ETTFMCEYAD ETATIVEFLN RWITFCQSII STLT	60 120 134
SEQ ID NO:4 Aldesleukin	PTSSSTKKTQ LQLEHLLLDL QMILNGINNY KNPKLTRMLT FKFYMPKKAT ELKHLQCLEE ELKPLEEVLN LAQSKNFHLR PRDLISNINV IVLELKGSET TFMCEYADET ATIVEFLNRW ITFSQSIIST LT	60 120 132
SEQ ID NO:5 recombinant human IL-4 (rhIL-4)	MHKCDITLQE IIKTLNSLTE QKTLCTELTV TDIFAASKNT TEKETFCRAA TVLRQFYSHH EKDTRCLGAT AQQFHRHKQL IRFLKRLDRN LWGLAGLNSC PVKEANQSTL ENFLERLKTI MREKYSKCSS	60 120 130
SEQ ID NO:6 recombinant human IL-7 (rhIL-7)	MDCDIEGKDG KQYESVLMVS IDQLLDSMKE IGSNCLNNEF NFFKRHICDA NKEGMFLFRA ARKLRQFLKM NSTGDFDLHL LKVSEGTTIL LNCTGQVKGR KPAALGEAQP TKSLEENKSL KEQKKLNDLC FLKRLLQEIK TCWNKILMGT KEH	60 120 153
SEQ ID NO:7 recombinant human IL-15 (rhIL-15)	MNWVNVISDL KKIEDLIQSM HIDATLYTES DVHPSCKVTA MKCFLLELQV ISLESGDASI HDTVENLIIL ANNSLSSNGN VTESGCKECE ELEEKNIKEF LQSFVHIVQM FINTS	60 115
SEQ ID NO:8 recombinant human IL-21 (rhIL-21)	MQDRHMIRMR QLIDIVDQLK NYVNDLVPEF LPAPEDVETN CEWSAFSCFQ KAQLKSANTG NNERIINVSI KKLKRKPPST NAGRRQKHRL TCPSCDSYEK KPPKEFLERF KSLLQKMIHQ HLSSRTHGSE DS	60 120 132

C. STEP C: Priming First Expansion to Rapid Second Expansion Transition

[00697] In some cases, the bulk TIL population obtained from the priming first expansion (which can include expansions sometimes referred to as pre-REP), including, for example the TIL population obtained from for example, Step B as indicated in Figure 85 (in particular, *e.g.*, Figure 85B), can be subjected to a rapid second expansion (which can include expansions sometimes referred to as Rapid Expansion Protocol (REP)) and then cryopreserved as discussed below. Similarly, in the case where genetically modified TILs will be used in therapy, the expanded TIL population from the priming first expansion or the expanded TIL population from the rapid second expansion can be subjected to genetic modifications for suitable treatments prior to the expansion step or after the priming first expansion and prior to the rapid second expansion.

[00698] In some embodiments, the TILs obtained from the priming first expansion (for example, from Step B as indicated in Figure 85 (in particular, *e.g.*, Figure 85B)) are stored until phenotyped for selection. In some embodiments, the TILs obtained from the priming first expansion (for example, from Step B as indicated in Figure 85 (in particular, *e.g.*, Figure 85B)) are not stored and proceed directly to the rapid second expansion. In some embodiments, the TILs obtained from the priming first expansion are not cryopreserved after the priming first expansion and prior to the rapid second expansion. In some embodiments, the transition from the priming first expansion to the second expansion occurs at about 2

days, 3 days, 4, days, 5 days, 6 days, or 7 days, from when tumor cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs at about 3 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the second expansion occurs at about 4 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the second expansion occurs at about 5 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the second expansion occurs at about 6 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the second expansion occurs at about 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated.

[00699] In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs at 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, or 7 days, up to 10 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs 1 day to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the second expansion occurs 2 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the second expansion occurs 3 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs 4 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs 5 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step

is initiated. In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs 6 days to 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated. In some embodiments, the transition from the priming first expansion to the rapid second expansion occurs 7 days from when cores or fragments are added to the cell culture medium and/or when the first priming expansion step is initiated.

[00700] In some embodiments, the TILs are not stored after the primary first expansion and prior to the rapid second expansion, and the TILs proceed directly to the rapid second expansion (for example, in some embodiments, there is no storage during the transition from Step B to Step D as shown in Figure 85 (in particular, *e.g.*, Figure 85B)). In some embodiments, the transition occurs in closed system, as described herein. In some embodiments, the TILs from the priming first expansion, the second population of TILs, proceeds directly into the rapid second expansion with no transition period.

[00701] In some embodiments, the transition from the priming first expansion to the rapid second expansion, for example, Step C according to Figure 85 (in particular, *e.g.*, Figure 85B), is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a single bioreactor is employed. In some embodiments, the single bioreactor employed is for example a GREX-10 or a GREX-100. In some embodiments, the closed system bioreactor is a single bioreactor. In some embodiments, the transition from the priming first expansion to the rapid second expansion involves a scale-up in container size. In some embodiments, the priming first expansion is performed in a smaller container than the rapid second expansion. In some embodiments, the priming first expansion is performed in a GREX-100 and the rapid second expansion is performed in a GREX-500.

D. STEP D: Rapid Second Expansion

[00702] In some embodiments, the TIL cell population is further expanded in number after harvest and the priming first expansion, after Step A and Step B, and the transition referred to as Step C, as indicated in Figure 85 (in particular, *e.g.*, Figure 85B)). This further expansion is referred to herein as the rapid second expansion, which can include expansion processes generally referred to in the art as a rapid expansion process (Rapid Expansion Protocol or REP; as well as processes as indicated in Step D of Figure 85 (in particular, *e.g.*, Figure 85B)). The rapid second expansion is generally accomplished using a culture media

comprising a number of components, including feeder cells, a cytokine source, and an anti-CD3 antibody, in a gas-permeable container. In some embodiments, 1 day, 2 days, 3 days, or 4 days after initiation of the rapid second expansion (*i.e.*, at days 8, 9, 10, or 11 of the overall Gen 3 process), the TILs are transferred to a larger volume container.

[00703] In some embodiments, the rapid second expansion (which can include expansions sometimes referred to as REP; as well as processes as indicated in Step D of Figure 85 (in particular, *e.g.*, Figure 85B)) of TIL can be performed using any TIL flasks or containers known by those of skill in the art. In some embodiments, the second TIL expansion can proceed for 1 day, 2 days, 3 days, 4, days, 5 days, 6 days, 7 days, 8 days, or 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 1 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 2 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 3 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 4 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 5 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 6 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 7 days to about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 7 days to about 10 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 1 day after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 2 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 3 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 4 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 5 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 6 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 7 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 8 days after initiation of the rapid second expansion. In some embodiments, the second

TIL expansion can proceed for about 9 days after initiation of the rapid second expansion. In some embodiments, the second TIL expansion can proceed for about 10 days after initiation of the rapid second expansion.

[00704] In an embodiment, the rapid second expansion can be performed in a gas permeable container using the methods of the present disclosure (including, for example, expansions referred to as REP; as well as processes as indicated in Step D of Figure 85 (in particular, *e.g.*, Figure 85B)). In some embodiments, the TILs are expanded in the rapid second expansion in the presence of IL-2, OKT-3, and feeder cells (also referred herein as "antigen-presenting cells"). In some embodiments, the TILs are expanded in the rapid second expansion in the presence of IL-2, OKT-3, and feeder cells, wherein the feeder cells are added to a final concentration that is twice, 2.4 times, 2.5 times, 3 times, 3.5 times or 4 times the concentration of feeder cells present in the priming first expansion. For example, TILs can be rapidly expanded using non-specific T-cell receptor stimulation in the presence of interleukin-2 (IL-2) or interleukin-15 (IL-15). The non-specific T-cell receptor stimulus can include, for example, an anti-CD3 antibody, such as about 30 ng/mL of OKT3, a mouse monoclonal anti-CD3 antibody (commercially available from Ortho-McNeil, Raritan, NJ or Miltenyi Biotech, Auburn, CA) or UHCT-1 (commercially available from BioLegend, San Diego, CA, USA). TILs can be expanded to induce further stimulation of the TILs *in vitro* by including one or more antigens during the second expansion, including antigenic portions thereof, such as epitope(s), of the cancer, which can be optionally expressed from a vector, such as a human leukocyte antigen A2 (HLA-A2) binding peptide, *e.g.*, 0.3 μM MART-1 :26-35 (27 L) or gpl 00:209-217 (210M), optionally in the presence of a T-cell growth factor, such as 300 IU/mL IL-2 or IL-15. Other suitable antigens may include, *e.g.*, NY-ESO-1, TRP-1, TRP-2, tyrosinase cancer antigen, MAGE-A3, SSX-2, and VEGFR2, or antigenic portions thereof. TIL may also be rapidly expanded by re-stimulation with the same antigen(s) of the cancer pulsed onto HLA-A2-expressing antigen-presenting cells. Alternatively, the TILs can be further re-stimulated with, *e.g.*, example, irradiated, autologous lymphocytes or with irradiated HLA-A2+ allogeneic lymphocytes and IL-2. In some embodiments, the re-stimulation occurs as part of the second expansion. In some embodiments, the second expansion occurs in the presence of irradiated, autologous lymphocytes or with irradiated HLA-A2+ allogeneic lymphocytes and IL-2.

[00705] In an embodiment, the cell culture medium further comprises IL-2. In some embodiments, the cell culture medium comprises about 3000 IU/mL of IL-2. In an

embodiment, the cell culture medium comprises about 1000 IU/mL, about 1500 IU/mL, about 2000 IU/mL, about 2500 IU/mL, about 3000 IU/mL, about 3500 IU/mL, about 4000 IU/mL, about 4500 IU/mL, about 5000 IU/mL, about 5500 IU/mL, about 6000 IU/mL, about 6500 IU/mL, about 7000 IU/mL, about 7500 IU/mL, or about 8000 IU/mL of IL-2. In an embodiment, the cell culture medium comprises between 1000 and 2000 IU/mL, between 2000 and 3000 IU/mL, between 3000 and 4000 IU/mL, between 4000 and 5000 IU/mL, between 5000 and 6000 IU/mL, between 6000 and 7000 IU/mL, between 7000 and 8000 IU/mL, or between 8000 IU/mL of IL-2.

[00706] In an embodiment, the cell culture medium comprises OKT-3 antibody. In some embodiments, the cell culture medium comprises about 30 ng/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises about 0.1 ng/mL, about 0.5 ng/mL, about 1 ng/mL, about 2.5 ng/mL, about 5 ng/mL, about 7.5 ng/mL, about 10 ng/mL, about 15 ng/mL, about 20 ng/mL, about 25 ng/mL, about 30 ng/mL, about 35 ng/mL, about 40 ng/mL, about 50 ng/mL, about 60 ng/mL, about 70 ng/mL, about 80 ng/mL, about 90 ng/mL, about 100 ng/mL, about 200 ng/mL, about 500 ng/mL, and about 1 μg/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises between 0.1 ng/mL and 1 ng/mL, between 1 ng/mL and 5 ng/mL, between 5 ng/mL and 10 ng/mL, between 10 ng/mL and 20 ng/mL, between 20 ng/mL and 30 ng/mL, between 30 ng/mL and 40 ng/mL, between 40 ng/mL and 50 ng/mL, and between 50 ng/mL and 100 ng/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises between 30 ng/ml and 60 ng/mL of OKT-3 antibody. In an embodiment, the cell culture medium comprises about 30 ng/mL OKT-3. In an embodiment, the cell culture medium comprises about 60 ng/mL OKT-3. In some embodiments, the OKT-3 antibody is muromonab.

[00707] In some embodiments, the media in the rapid second expansion comprises IL-2. In some embodiments, the media comprises 6000 IU/mL of IL-2. In some embodiments, the media in the rapid second expansion comprises antigen-presenting feeder cells. In some embodiments, the media in the rapid second expansion comprises 7.5×10^8 antigen-presenting feeder cells per container. In some embodiments, the media in the rapid second expansion comprises OKT-3. In some embodiments, the in the rapid second expansion media comprises 500 mL of culture medium and 30 ug of OKT-3 per container. In some embodiments, the container is a GREX100 MCS flask. In some embodiments, the in the rapid second expansion media comprises 6000 IU/mL of IL-2, 60 ng/mL of OKT-3, and 7.5×10^8 antigen-presenting feeder cells. In some embodiments, the media comprises 500 mL of

culture medium and 6000 IU/mL of IL-2, 30 ug of OKT-3, and 7.5×10^8 antigen-presenting feeder cells per container.

[00708] In some embodiments, the media in the rapid second expansion comprises IL-2. In some embodiments, the media comprises 6000 IU/mL of IL-2. In some embodiments, the media in the rapid second expansion comprises antigen-presenting feeder cells. In some embodiments, the media comprises between 5×10^8 and 7.5×10^8antigen-presenting feeder cells per container. In some embodiments, the media in the rapid second expansion comprises OKT-3. In some embodiments, the media in the rapid second expansion comprises 500 mL of culture medium and 30 μg of OKT-3 per container. In some embodiments, the container is a GREX100 MCS flask. In some embodiments, the media in the rapid second expansion comprises 6000 IU/mL of IL-2, 60 ng/mL of OKT-3, and between 5×10^8 and 7.5×10^8 antigen-presenting feeder cells. In some embodiments, the media in the rapid second expansion comprises 500 mL of culture medium and 6000 IU/mL of IL-2, 30 μg of OKT-3, and between 5×10^8 and 7.5×10^8 antigen-presenting feeder cells per container.

[00709] In some embodiments, the cell culture medium comprises one or more TNFRSF agonists in a cell culture medium. In some embodiments, the TNFRSF agonist comprises a 4-1BB agonist. In some embodiments, the TNFRSF agonist is a 4-1BB agonist, and the 4-1BB agonist is selected from the group consisting of urelumab, utomilumab, EU-101, a fusion protein, and fragments, derivatives, variants, biosimilars, and combinations thereof. In some embodiments, the TNFRSF agonist is added at a concentration sufficient to achieve a concentration in the cell culture medium of between 0.1 μg/mL and 100 μg/mL. In some embodiments, the TNFRSF agonist is added at a concentration sufficient to achieve a concentration in the cell culture medium of between 20 μg/mL and 40 μg/mL.

[00710] In some embodiments, in addition to one or more TNFRSF agonists, the cell culture medium further comprises IL-2 at an initial concentration of about 3000 IU/mL and OKT-3 antibody at an initial concentration of about 30 ng/mL, and wherein the one or more TNFRSF agonists comprises a 4-1BB agonist.

[00711] In some embodiments, a combination of IL-2, IL-7, IL-15, and/or IL-21 are employed as a combination during the second expansion. In some embodiments, IL-2, IL-7, IL-15, and/or IL-21 as well as any combinations thereof can be included during the second expansion, including, for example during a Step D processes according to Figure 85 (in particular, *e.g.*, Figure 85B), as well as described herein. In some embodiments, a

combination of IL-2, IL-15, and IL-21 are employed as a combination during the second expansion. In some embodiments, IL-2, IL-15, and IL-21 as well as any combinations thereof can be included during Step D processes according to Figure 85 (in particular, *e.g.*, Figure 85B) and as described herein.

[00712] In some embodiments, the second expansion can be conducted in a supplemented cell culture medium comprising IL-2, OKT-3, antigen-presenting feeder cells, and optionally a TNFRSF agonist. In some embodiments, the second expansion occurs in a supplemented cell culture medium. In some embodiments, the supplemented cell culture medium comprises IL-2, OKT-3, and antigen-presenting feeder cells. In some embodiments, the second cell culture medium comprises IL-2, OKT-3, and antigen-presenting cells (APCs; also referred to as antigen-presenting feeder cells). In some embodiments, the second expansion occurs in a cell culture medium comprising IL-2, OKT-3, and antigen-presenting feeder cells (*i.e.*, antigen presenting cells).

[00713] In some embodiments, the second expansion culture media comprises about 500 IU/mL of IL-15, about 400 IU/mL of IL-15, about 300 IU/mL of IL-15, about 200 IU/mL of IL-15, about 180 IU/mL of IL-15, about 160 IU/mL of IL-15, about 140 IU/mL of IL-15, about 120 IU/mL of IL-15, or about 100 IU/mL of IL-15. In some embodiments, the second expansion culture media comprises about 500 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the second expansion culture media comprises about 400 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the second expansion culture media comprises about 300 IU/mL of IL-15 to about 100 IU/mL of IL-15. In some embodiments, the second expansion culture media comprises about 200 IU/mL of IL-15. In some embodiments, the cell culture medium comprises about 180 IU/mL of IL-15. In an embodiment, the cell culture medium further comprises IL-15. In a preferred embodiment, the cell culture medium comprises about 180 IU/mL of IL-15.

[00714] In some embodiments, the second expansion culture media comprises about 20 IU/mL of IL-21, about 15 IU/mL of IL-21, about 12 IU/mL of IL-21, about 10 IU/mL of IL-21, about 5 IU/mL of IL-21, about 4 IU/mL of IL-21, about 3 IU/mL of IL-21, about 2 IU/mL of IL-21, about 1 IU/mL of IL-21, or about 0.5 IU/mL of IL-21. In some embodiments, the second expansion culture media comprises about 20 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the second expansion culture media comprises about 15 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the second expansion culture media comprises about 12 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some

embodiments, the second expansion culture media comprises about 10 IU/mL of IL-21 to about 0.5 IU/mL of IL-21. In some embodiments, the second expansion culture media comprises about 5 IU/mL of IL-21 to about 1 IU/mL of IL-21. In some embodiments, the second expansion culture media comprises about 2 IU/mL of IL-21. In some embodiments, the cell culture medium comprises about 1 IU/mL of IL-21. In some embodiments, the cell culture medium comprises about 0.5 IU/mL of IL-21. In an embodiment, the cell culture medium further comprises IL-21. In a preferred embodiment, the cell culture medium comprises about 1 IU/mL of IL-21.

[00715] In some embodiments the antigen-presenting feeder cells (APCs) are PBMCs. In an embodiment, the ratio of TILs to PBMCs and/or antigen-presenting cells in the rapid expansion and/or the second expansion is about 1 to 10, about 1 to 15, about 1 to 20, about 1 to 25, about 1 to 30, about 1 to 35, about 1 to 40, about 1 to 45, about 1 to 50, about 1 to 75, about 1 to 100, about 1 to 125, about 1 to 150, about 1 to 175, about 1 to 200, about 1 to 225, about 1 to 250, about 1 to 275, about 1 to 300, about 1 to 325, about 1 to 350, about 1 to 375, about 1 to 400, or about 1 to 500. In an embodiment, the ratio of TILs to PBMCs in the rapid expansion and/or the second expansion is between 1 to 50 and 1 to 300. In an embodiment, the ratio of TILs to PBMCs in the rapid expansion and/or the second expansion is between 1 to 100 and 1 to 200.

[00716] In an embodiment, REP and/or the rapid second expansion is performed in flasks with the bulk TILs being mixed with a 100- or 200-fold excess of inactivated feeder cells, wherein the feeder cell concentration is at least 1.1 times (1.1X), 1.2X, 1.3X, 1.4X, 1.5X, 1.6X, 1.7X, 1.8X, 1.8X, 2X, 2.1X2.2X, 2.3X, 2.4X, 2.5X, 2.6X, 2.7X, 2.8X, 2.9X, 3.0X, 3.1X, 3.2X, 3.3X, 3.4X, 3.5X, 3.6X, 3.7X, 3.8X, 3.9X or 4.0X the feeder cell concentration in the priming first expansion, 30 ng/mL OKT3 anti-CD3 antibody and 6000 IU/mL IL-2 in 150 ml media. Media replacement is done (generally 2/3 media replacement via aspiration of 2/3 of spent media and replacement with an equal volume of fresh media) until the cells are transferred to an alternative growth chamber. Alternative growth chambers include G-REX flasks and gas permeable containers as more fully discussed below.

[00717] In some embodiments, the rapid second expansion (which can include processes referred to as the REP process) is 7 to 10 days, as discussed in the examples and figures. In some embodiments, the rapid second expansion (which can include processes referred to as the REP process) is 7 to 9 days, as discussed in the examples and figures. In some embodiments, the second expansion is 7 days. In some embodiments, the second expansion is

8 days. In some embodiments, the second expansion is 9 days. In some embodiments, the second expansion is 10 days.

[00718] In an embodiment, the second expansion (which can include expansions referred to as REP, as well as those referred to in Step D of Figure 85 (in particular, *e.g.*, Figure 85B)) may be performed in 500 mL capacity gas permeable flasks with 100 cm gas-permeable silicon bottoms (G-Rex 100, commercially available from Wilson Wolf Manufacturing Corporation, New Brighton, MN, USA), 5×10^6 or 10×10^6 TIL may be cultured with PBMCs in 400 mL of 50/50 medium, supplemented with 5% human AB serum, 3000 IU per mL of IL-2 and 30 ng per ml of anti-CD3 (OKT3). The G-Rex 100 flasks may be incubated at 37°C in 5% CO_2. On day 5, 250 mL of supernatant may be removed and placed into centrifuge bottles and centrifuged at 1500 rpm (491 × g) for 10 minutes. The TIL pellets may be re-suspended with 150 mL of fresh medium with 5% human AB serum, 6000 IU per mL of IL-2, and added back to the original GREX-100 flasks. When TIL are expanded serially in GREX-100 flasks, on day 10 or 11 the TILs can be moved to a larger flask, such as a GREX-500. The cells may be harvested on day 14 of culture. The cells may be harvested on day 15 of culture. The cells may be harvested on day 16 of culture. In some embodiments, media replacement is done until the cells are transferred to an alternative growth chamber. In some embodiments, 2/3 of the media is replaced by aspiration of spent media and replacement with an equal volume of fresh media. In some embodiments, alternative growth chambers include GREX flasks and gas permeable containers as more fully discussed below.

[00719] In an embodiment, the rapid second expansion (including expansions referred to as REP) is performed and further comprises a step wherein TILs are selected for superior tumor reactivity. Any selection method known in the art may be used. For example, the methods described in U.S. Patent Application Publication No. 2016/0010058 A1, the disclosures of which are incorporated herein by reference, may be used for selection of TILs for superior tumor reactivity.

[00720] Optionally, a cell viability assay can be performed after the rapid second expansion (including expansions referred to as the REP expansion), using standard assays known in the art. For example, a trypan blue exclusion assay can be done on a sample of the bulk TILs, which selectively labels dead cells and allows a viability assessment. In some embodiments, TIL samples can be counted and viability determined using a Cellometer K2 automated cell counter (Nexcelom Bioscience, Lawrence, MA). In some embodiments, viability is

determined according to the standard Cellometer K2 Image Cytometer Automatic Cell Counter protocol.

[00721] The diverse antigen receptors of T and B lymphocytes are produced by somatic recombination of a limited, but large number of gene segments. These gene segments: V (variable), D (diversity), J (joining), and C (constant), determine the binding specificity and downstream applications of immunoglobulins and T-cell receptors (TCRs). The present invention provides a method for generating TILs which exhibit and increase the T-cell repertoire diversity. In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity. In some embodiments, the TILs obtained in the second expansion exhibit an increase in the T-cell repertoire diversity. In some embodiments, the increase in diversity is an increase in the immunoglobulin diversity and/or the T-cell receptor diversity. In some embodiments, the diversity is in the immunoglobulin is in the immunoglobulin heavy chain. In some embodiments, the diversity is in the immunoglobulin is in the immunoglobulin light chain. In some embodiments, the diversity is in the T-cell receptor. In some embodiments, the diversity is in one of the T-cell receptors selected from the group consisting of alpha, beta, gamma, and delta receptors. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) alpha and/or beta. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) alpha. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) beta. In some embodiments, there is an increase in the expression of TCRab (*i.e.*, TCRα/β).

[00722] In some embodiments, the rapid second expansion culture medium (*e.g.*, sometimes referred to as CM2 or the second cell culture medium), comprises IL-2, OKT-3, as well as the antigen-presenting feeder cells (APCs), as discussed in more detail below. In some embodiments, the rapid second expansion culture medium (*e.g.*, sometimes referred to as CM2 or the second cell culture medium), comprises 6000 IU/mL IL-2, 30 μg/flask OKT-3, as well as 7.5×10^8 antigen-presenting feeder cells (APCs), as discussed in more detail below. In some embodiments, the rapid second expansion culture medium (*e.g.*, sometimes referred to as CM2 or the second cell culture medium), comprises IL-2, OKT-3, as well as the antigen-presenting feeder cells (APCs), as discussed in more detail below. In some embodiments, the rapid second expansion culture medium (*e.g.*, sometimes referred to as CM2 or the second cell culture medium), comprises 6000 IU/mL IL-2, 30 μg/flask OKT-3, as well as 5×10^8 antigen-presenting feeder cells (APCs), as discussed in more detail below.

[00723] In some embodiments, the rapid second expansion, for example, Step D according to Figure 85 (in particular, *e.g.*, Figure 85B), is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a bioreactor is employed. In some embodiments, a bioreactor is employed as the container. In some embodiments, the bioreactor employed is for example a G-REX-100 or a G-REX-500. In some embodiments, the bioreactor employed is a G-REX-100. In some embodiments, the bioreactor employed is a G-REX-500.

1. Feeder Cells and Antigen Presenting Cells

[00724] In an embodiment, the rapid second expansion procedures described herein (for example including expansion such as those described in Step D from Figure 85 (in particular, *e.g.*, Figure 85B), as well as those referred to as REP) require an excess of feeder cells during REP TIL expansion and/or during the rapid second expansion. In many embodiments, the feeder cells are peripheral blood mononuclear cells (PBMCs) obtained from standard whole blood units from healthy blood donors. The PBMCs are obtained using standard methods such as Ficoll-Paque gradient separation.

[00725] In general, the allogenic PBMCs are inactivated, either via irradiation or heat treatment, and used in the REP procedures, as described in the examples, which provides an exemplary protocol for evaluating the replication incompetence of irradiate allogeneic PBMCs.

[00726] In some embodiments, PBMCs are considered replication incompetent and acceptable for use in the TIL expansion procedures described herein if the total number of viable cells on day 7 or 14 is less than the initial viable cell number put into culture on day 0 of the REP and/or day 0 of the second expansion (*i.e.*, the start day of the second expansion).

[00727] In some embodiments, PBMCs are considered replication incompetent and acceptable for use in the TIL expansion procedures described herein if the total number of viable cells, cultured in the presence of OKT3 and IL-2, on day 7 and day 14 has not increased from the initial viable cell number put into culture on day 0 of the REP and/or day 0 of the second expansion (*i.e.*, the start day of the second expansion). In some embodiments, the PBMCs are cultured in the presence of 30 ng/mL OKT3 antibody and 3000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 30 ng/mL OKT3 antibody and 6000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 15 ng/mL OKT3 antibody and 3000 IU/mL IL-2. In some embodiments, the PBMCs are

cultured in the presence of 15 ng/mL OKT3 antibody and 6000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 60 ng/mL OKT3 antibody and 3000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 60 ng/mL OKT3 antibody and 6000 IU/mL IL-2.

[00728] In some embodiments, PBMCs are considered replication incompetent and acceptable for use in the TIL expansion procedures described herein if the total number of viable cells, cultured in the presence of OKT3 and IL-2, on day 7 and day 14 has not increased from the initial viable cell number put into culture on day 0 of the REP and/or day 0 of the second expansion (*i.e.*, the start day of the second expansion). In some embodiments, the PBMCs are cultured in the presence of 30-60 ng/mL OKT3 antibody and 1000-6000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 30-60 ng/mL OKT3 antibody and 2000-5000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 30-60 ng/mL OKT3 antibody and 2000-4000 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 30-60 ng/mL OKT3 antibody and 2500-3500 IU/mL IL-2. In some embodiments, the PBMCs are cultured in the presence of 30-60 ng/mL OKT3 antibody and 6000 IU/mL IL-2.

[00729] In some embodiments, the antigen-presenting feeder cells are PBMCs. In some embodiments, the antigen-presenting feeder cells are artificial antigen-presenting feeder cells. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is about 1 to 10, about 1 to 25, about 1 to 50, about 1 to 100, about 1 to 125, about 1 to 150, about 1 to 175, about 1 to 200, about 1 to 225, about 1 to 250, about 1 to 275, about 1 to 300, about 1 to 325, about 1 to 350, about 1 to 375, about 1 to 400, or about 1 to 500. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is between 1 to 50 and 1 to 300. In an embodiment, the ratio of TILs to antigen-presenting feeder cells in the second expansion is between 1 to 100 and 1 to 200.

[00730] In an embodiment, the second expansion procedures described herein require a ratio of about 5×10^8 feeder cells to about 100×10^6 TILs. In an embodiment, the second expansion procedures described herein require a ratio of about 7.5×10^8 feeder cells to about 100×10^6 TILs. In another embodiment, the second expansion procedures described herein require a ratio of about 5×10^8 feeder cells to about 50×10^6 TILs. In another embodiment, the second expansion procedures described herein require a ratio of about 7.5×10^8 feeder cells to about 50×10^6 TILs. In yet another embodiment, the second expansion procedures described herein require about 5×10^8 feeder cells to about 25×10^6 TILs. In yet another

embodiment, the second expansion procedures described herein require about 7.5×10^8 feeder cells to about 25×10^6 TILs. In yet another embodiment, the rapid second expansion requires twice the number of feeder cells as the rapid second expansion. In yet another embodiment, when the priming first expansion described herein requires about 2.5×10^8 feeder cells, the rapid second expansion requires about 5×10^8 feeder cells. In yet another embodiment, when the priming first expansion described herein requires about 2.5×10^8 feeder cells, the rapid second expansion requires about 7.5×10^8 feeder cells. In yet another embodiment, the rapid second expansion requires two times (2.0X), 2.5X, 3.0X, 3.5X or 4.0X the number of feeder cells as the priming first expansion.

[00731] In an embodiment, the rapid second expansion procedures described herein require an excess of feeder cells during the rapid second expansion. In many embodiments, the feeder cells are peripheral blood mononuclear cells (PBMCs) obtained from standard whole blood units from allogeneic healthy blood donors. The PBMCs are obtained using standard methods such as Ficoll-Paque gradient separation. In an embodiment, artificial antigen-presenting (aAPC) cells are used in place of PBMCs. In some embodiments, the PBMCs are added to the rapid second expansion at twice the concentration of PBMCs that were added to the priming first expansion.

[00732] In general, the allogenic PBMCs are inactivated, either via irradiation or heat treatment, and used in the TIL expansion procedures described herein, including the exemplary procedures described in the figures and examples.

[00733] In an embodiment, artificial antigen presenting cells are used in the rapid second expansion as a replacement for, or in combination with, PBMCs.

2. Cytokines

[00734] The rapid second expansion methods described herein generally use culture media with high doses of a cytokine, in particular IL-2, as is known in the art.

[00735] Alternatively, using combinations of cytokines for the rapid second expansion of TILs is additionally possible, with combinations of two or more of IL-2, IL-15 and IL-21 as is generally outlined in WO 2015/189356 and WO 2015/189357, hereby expressly incorporated by reference in their entirety. Thus, possible combinations include IL-2 and IL-15, IL-2 and IL-21, IL-15 and IL-21, and IL-2, IL-15 and IL-21, with the latter finding particular use in many embodiments. The use of combinations of cytokines specifically favors the generation of lymphocytes, and in particular T-cells as described therein.

E. STEP E: Harvest TILS

[00736] After the rapid second expansion step, cells can be harvested. In some embodiments the TILs are harvested after one, two, three, four or more expansion steps, for example as provided in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments the TILs are harvested after two expansion steps, for example as provided in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments the TILs are harvested after two expansion steps, one priming first expansion and one rapid second expansion, for example as provided in Figure 85 (in particular, *e.g.*, Figure 85B).

[00737] TILs can be harvested in any appropriate and sterile manner, including, for example by centrifugation. Methods for TIL harvesting are well known in the art and any such known methods can be employed with the present process. In some embodiments, TILS are harvested using an automated system.

[00738] Cell harvesters and/or cell processing systems are commercially available from a variety of sources, including, for example, Fresenius Kabi, Tomtec Life Science, Perkin Elmer, and Inotech Biosystems International, Inc. Any cell based harvester can be employed with the present methods. In some embodiments, the cell harvester and/or cell processing system is a membrane-based cell harvester. In some embodiments, cell harvesting is via a cell processing system, such as the LOVO system (manufactured by Fresenius Kabi). The term "LOVO cell processing system" also refers to any instrument or device manufactured by any vendor that can pump a solution comprising cells through a membrane or filter such as a spinning membrane or spinning filter in a sterile and/or closed system environment, allowing for continuous flow and cell processing to remove supernatant or cell culture media without pelletization. In some embodiments, the cell harvester and/or cell processing system can perform cell separation, washing, fluid-exchange, concentration, and/or other cell processing steps in a closed, sterile system.

[00739] In some embodiments, the rapid second expansion, for example, Step D according to Figure 85 (in particular, *e.g.*, Figure 85B), is performed in a closed system bioreactor. In some embodiments, a closed system is employed for the TIL expansion, as described herein. In some embodiments, a bioreactor is employed. In some embodiments, a bioreactor is employed as the container. In some embodiments, the bioreactor employed is for example a G-REX-100 or a G-REX-500. In some embodiments, the bioreactor employed is a G-REX-100. In some embodiments, the bioreactor employed is a G-REX-500.

[00740] In some embodiments, Step E according to Figure 85 (in particular, *e.g.*, Figure 85B), is performed according to the processes described herein. In some embodiments, the closed system is accessed via syringes under sterile conditions in order to maintain the sterility and closed nature of the system. In some embodiments, a closed system as described herein is employed.

[00741] In some embodiments, TILs are harvested according to the methods described in herein. In some embodiments, TILs between days 14 and 16 are harvested using the methods as described herein. In some embodiments, TILs are harvested at 14 days using the methods as described herein. In some embodiments, TILs are harvested at 15 days using the methods as described herein. In some embodiments, TILs are harvested at 16 days using the methods as described herein.

F. STEP F: Final Formulation/ Transfer to Infusion Bag

[00742] After Steps A through E as provided in an exemplary order in Figure 85 (in particular, *e.g.*, Figure 85B) and as outlined in detailed above and herein are complete, cells are transferred to a container for use in administration to a patient. In some embodiments, once a therapeutically sufficient number of TILs are obtained using the expansion methods described above, they are transferred to a container for use in administration to a patient.

[00743] In an embodiment, TILs expanded using the methods of the present disclosure are administered to a patient as a pharmaceutical composition. In an embodiment, the pharmaceutical composition is a suspension of TILs in a sterile buffer. TILs expanded as disclosed herein may be administered by any suitable route as known in the art. In some embodiments, the TILs are administered as a single intra-arterial or intravenous infusion, which preferably lasts approximately 30 to 60 minutes. Other suitable routes of administration include intraperitoneal, intrathecal, and intralymphatic.

[00744] In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant and an isotonic agent. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions further comprising a cyropreservant comprising dimethylsulfoxide and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered

as compositions further comprising a cyropreservant comprising dimethylsulfoxide and dextran 40 and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate. In an embodiment, TILs expanded using processes of the foregoing disclosure may be administered as compositions delivered in a sterile infusion bag, such compositions further comprising a cyropreservant comprising dimethylsulfoxide and dextran 40 and an isotonic agent comprising sodium chloride, sodium gluconate, and sodium acetate.

G. PBMC Feeder Cell Ratios

[00745] In some embodiments, the culture media used in expansion methods described herein (see for example, Figure 85 (in particular, *e.g.*, Figure 85B)) include an anti-CD3 antibody *e.g.* OKT-3. An anti-CD3 antibody in combination with IL-2 induces T cell activation and cell division in the TIL population. This effect can be seen with full length antibodies as well as Fab and F(ab')2 fragments, with the former being generally preferred; see, *e.g.,* Tsoukas *et al.*, *J. Immunol.* **1985,** *135,* 1719, hereby incorporated by reference in its entirety.

[00746] In an embodiment, the number of PBMC feeder layers is calculated as follows:

A. Volume of a T-cell (10 μm diameter): $V = (4/3)\ \pi r^3 = 523.6\ \mu m^3$

B. Columne of G-Rex 100 (M) with a 40 μm (4 cells) height: $V = (4/3)\ \pi r^3 = 4\times10^{12}\ \mu m^3$

C. Number cell required to fill column B: $4\times10^{12}\ \mu m^3\ /\ 523.6\ \mu m^3 = 7.6x10^8\ \mu m^3\ * 0.64 = 4.86\times10^8$

D. Number cells that can be optimally activated in 4D space: $4.86\times10^8 / 24 = 20.25\times10^6$

E. Number of feeders and TIL extrapolated to G-Rex 500: TIL: 100×10^6 and Feeder: 2.5×10^9

In this calculation, an approximation of the number of mononuclear cells required to provide an icosahedral geometry for activation of TIL in a cylinder with a 100 cm^2 base is used. The calculation derives the experimental result of ~5×10^8 for threshold activation of T-cells which closely mirrors NCI experimental data.[(1)] (C) The multiplier (0.64) is the random packing density for equivalent spheres as calculated by Jaeger and Nagel in 1992 [(2)]. (D) The divisor 24 is the number of equivalent spheres that could contact a similar object in 4 dimensional space "the Newton number"[(3)].

(1) Jin, Jianjian, et.al., Simplified Method of the Growth of Human Tumor Infiltrating Lymphocytes (TIL) in Gas-Permeable Flasks to Numbers Needed for Patient Treatment. J Immunother. 2012 Apr; 35(3): 283–292.

(2) Jaeger HM, Nagel SR. Physics of the granular state. Science. 1992 Mar 20;255(5051):1523-31.

(3) O. R. Musin (2003). "The problem of the twenty-five spheres". Russ. Math. Surv. 58 (4): 794–795.

[00747] In an embodiment, the number of antigen-presenting feeder cells exogenously supplied during the priming first expansion is approximately one-half the number of antigen-presenting feeder cells exogenously supplied during the rapid second expansion. In certain embodiments, the method comprises performing the priming first expansion in a cell culture medium which comprises approximately 50% fewer antigen presenting cells as compared to the cell culture medium of the rapid second expansion.

[00748] In another embodiment, the number of antigen-presenting feeder cells (APCs) exogenously supplied during the rapid second expansion is greater than the number of APCs exogenously supplied during the priming first expansion.

[00749] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 20:1.

[00750] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 10:1.

[00751] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 9:1.

[00752] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 8:1.

[00753] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 7:1.

[00754] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 6:1.

[00755] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 5:1.

[00756] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 4:1.

[00757] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion) is selected from a range of from at or about 1.1:1 to at or about 3:1.

[00758] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.9:1.

[00759] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.8:1.

[00760] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.7:1.

[00761] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.6:1.

[00762] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.5:1.

[00763] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.4:1.

[00764] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.3:1.

[00765] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.2:1.

[00766] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.1:1.

[00767] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2:1.

[00768] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 10:1.

[00769] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 5:1.

[00770] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 4:1.

[00771] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 3:1.

[00772] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.9:1.

[00773] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.8:1.

[00774] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.7:1.

[00775] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.6:1.

[00776] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.5:1.

[00777] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.4:1.

[00778] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.3:1.

[00779] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about about 2:1 to at or about 2.2:1.

[00780] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.1:1.

[00781] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is at or about 2:1.

[00782] In another embodiment, the ratio of the number of APCs exogenously supplied during the rapid second expansion to the number of APCs exogenously supplied during the priming first expansion is at or about 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, 2:1, 2.1:1, 2.2:1, 2.3:1, 2.4:1, 2.5:1, 2.6:1, 2.7:1, 2.8:1, 2.9:1, 3:1, 3.1:1, 3.2:1, 3.3:1, 3.4:1, 3.5:1, 3.6:1, 3.7:1, 3.8:1, 3.9:1, 4:1, 4.1:1, 4.2:1, 4.3:1, 4.4:1, 4.5:1, 4.6:1, 4.7:1, 4.8:1, 4.9:1, or 5:1.

[00783] In another embodiment, the number of APCs exogenously supplied during the priming first expansion is at or about 1×10^8, 1.1×10^8, 1.2×10^8, 1.3×10^8, 1.4×10^8, 1.5×10^8, 1.6×10^8, 1.7×10^8, 1.8×10^8, 1.9×10^8, 2×10^8, 2.1×10^8, 2.2×10^8, 2.3×10^8, 2.4×10^8, 2.5×10^8, 2.6×10^8, 2.7×10^8, 2.8×10^8, 2.9×10^8, 3×10^8, 3.1×10^8, 3.2×10^8, 3.3×10^8, 3.4×10^8 or 3.5×10^8 APCs, and the number of APCs exogenously supplied during the rapid second expansion is at or about 3.5×10^8, 3.6×10^8, 3.7×10^8, 3.8×10^8, 3.9×10^8, 4×10^8, 4.1×10^8, 4.2×10^8, 4.3×10^8, 4.4×10^8, 4.5×10^8, 4.6×10^8, 4.7×10^8, 4.8×10^8, 4.9×10^8, 5×10^8, 5.1×10^8, 5.2×10^8, 5.3×10^8, 5.4×10^8, 5.5×10^8, 5.6×10^8, 5.7×10^8, 5.8×10^8, 5.9×10^8, 6×10^8, 6.1×10^8, 6.2×10^8, 6.3×10^8, 6.4×10^8, 6.5×10^8, 6.6×10^8, 6.7×10^8, 6.8×10^8, 6.9×10^8, 7×10^8, 7.1×10^8, 7.2×10^8, 7.3×10^8, 7.4×10^8, 7.5×10^8, 7.6×10^8, 7.7×10^8, 7.8×10^8, 7.9×10^8, 8×10^8, 8.1×10^8, 8.2×10^8, 8.3×10^8, 8.4×10^8, 8.5×10^8, 8.6×10^8, 8.7×10^8, 8.8×10^8, 8.9×10^8, 9×10^8, 9.1×10^8, 9.2×10^8, 9.3×10^8, 9.4×10^8, 9.5×10^8, 9.6×10^8, 9.7×10^8, 9.8×10^8, 9.9×10^8 or 1×10^9 APCs.

[00784] In another embodiment, the number of APCs exogenously supplied during the priming first expansion is selected from the range of at or about 1.5×10^8 APCs to at or about 3×10^8 APCs, and the number of APCs exogenously supplied during the rapid second expansion is selected from the range of at or about 4×10^8 APCs to at or about 7.5×10^8 APCs.

[00785] In another embodiment, the number of APCs exogenously supplied during the priming first expansion is selected from the range of at or about 2×10^8 APCs to at or about 2.5×10^8 APCs, and the number of APCs exogenously supplied during the rapid second expansion is selected from the range of at or about 4.5×10^8 APCs to at or about 5.5×10^8 APCs.

[00786] In another embodiment, the number of APCs exogenously supplied during the priming first expansion is at or about 2.5×10^8 APCs, and the number of APCs exogenously supplied during the rapid second expansion is at or about 5×10^8 APCs.

[00787] In an embodiment, the number of APCs (including, for example, PBMCs) added at day 0 of the priming first expansion is approximately one-half of the number of PBMCs added at day 7 of the priming first expansion (*e.g.*, day 7 of the method). In certain embodiments, the method comprises adding antigen presenting cells at day 0 of the priming first expansion to the first population of TILs and adding antigen presenting cells at day 7 to the second population of TILs, wherein the number of antigen presenting cells added at day 0

is approximately 50% of the number of antigen presenting cells added at day 7 of the priming first expansion (*e.g.*, day 7 of the method).

[00788] In another embodiment, the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is greater than the number of PBMCs exogenously supplied at day 0 of the priming first expansion.

[00789] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density selected from a range of at or about 1.0×10^6 APCs/cm^2 to at or about 4.5×10^6 APCs/cm^2.

[00790] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density selected from a range of at or about 1.5×10^6 APCs/cm^2 to at or about 3.5×10^6 APCs/cm^2.

[00791] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density selected from a range of at or about 2×10^6 APCs/cm^2 to at or about 3×10^6 APCs/cm^2.

[00792] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density of at or about 2×10^6 APCs/cm^2.

[00793] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density of at or about 1.0×10^6, 1.1×10^6, 1.2×10^6, 1.3×10^6, 1.4×10^6, 1.5×10^6, 1.6×10^6, 1.7×10^6, 1.8×10^6, 1.9×10^6, 2×10^6, 2.1×10^6, 2.2×10^6, 2.3×10^6, 2.4×10^6, 2.5×10^6, 2.6×10^6, 2.7×10^6, 2.8×10^6, 2.9×10^6, 3×10^6, 3.1×10^6, 3.2×10^6, 3.3×10^6, 3.4×10^6, 3.5×10^6, 3.6×10^6, 3.7×10^6, 3.8×10^6, 3.9×10^6, 4×10^6, 4.1×10^6, 4.2×10^6, 4.3×10^6, 4.4×10^6 or 4.5×10^6 APCs/cm^2.

[00794] In another embodiment, the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 2.5×10^6 APCs/cm^2 to at or about 7.5×10^6 APCs/cm^2.

[00795] In another embodiment, the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 3.5×10^6 APCs/cm^2 to about 6.0×10^6 APCs/cm^2.

[00796] In another embodiment, the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 4.0×10^6 APCs/cm^2 to about 5.5×10^6 APCs/cm^2.

[00797] In another embodiment, the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 4.0 × 10^6 APCs/cm^2.

[00798] In another embodiment, the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density of at or about 2.5 × 10^6 APCs/cm^2, 2.6 × 10^6 APCs/cm^2, 2.7 × 10^6 APCs/cm^2, 2.8 × 10^6, 2.9 × 10^6, 3 × 10^6, 3.1 × 10^6, 3.2 × 10^6, 3.3 × 10^6, 3.4 × 10^6, 3.5 × 10^6, 3.6 × 10^6, 3.7 × 10^6, 3.8 × 10^6, 3.9 × 10^6, 4 × 10^6, 4.1 × 10^6, 4.2 × 10^6, 4.3 × 10^6, 4.4 × 10^6, 4.5 × 10^6, 4.6 × 10^6, 4.7 × 10^6, 4.8 × 10^6, 4.9 × 10^6, 5 × 10^6, 5.1 × 10^6, 5.2 × 10^6, 5.3 × 10^6, 5.4 × 10^6, 5.5 × 10^6, 5.6 × 10^6, 5.7 × 10^6, 5.8 × 10^6, 5.9 × 10^6, 6 × 10^6, 6.1 × 10^6, 6.2 × 10^6, 6.3 × 10^6, 6.4 × 10^6, 6.5 × 10^6, 6.6 × 10^6, 6.7 × 10^6, 6.8 × 10^6, 6.9 × 10^6, 7 × 10^6, 7.1 × 10^6, 7.2 × 10^6, 7.3 × 10^6, 7.4 × 10^6, or 7.5 × 10^6 APCs/cm^2.

[00799] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density of at or about 1.0 × 10^6, 1.1 × 10^6, 1.2 × 10^6, 1.3 × 10^6, 1.4 × 10^6, 1.5 × 10^6, 1.6 × 10^6, 1.7 × 10^6, 1.8 × 10^6, 1.9 × 10^6, 2 × 10^6, 2.1 × 10^6, 2.2 × 10^6, 2.3 × 10^6, 2.4 × 10^6, 2.5 × 10^6, 2.6 × 10^6, 2.7 × 10^6, 2.8 × 10^6, 2.9 × 10^6, 3 × 10^6, 3.1 × 10^6, 3.2 × 10^6, 3.3 × 10^6, 3.4 × 10^6, 3.5 × 10^6, 3.6 × 10^6, 3.7 × 10^6, 3.8 × 10^6, 3.9 × 10^6, 4 × 10^6, 4.1 × 10^6, 4.2 × 10^6, 4.3 × 10^6, 4.4 × 10^6, or 4.5 × 10^6 APCs/cm^2 and the the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density of at or about 2.5 × 10^6 APCs/cm^2, 2.6 × 10^6 APCs/cm^2, 2.7 × 10^6 APCs/cm^2, 2.8 × 10^6, 2.9 × 10^6, 3 × 10^6, 3.1 × 10^6, 3.2 × 10^6, 3.3 × 10^6, 3.4 × 10^6, 3.5 × 10^6, 3.6 × 10^6, 3.7 × 10^6, 3.8 × 10^6, 3.9 × 10^6, 4 × 10^6, 4.1 × 10^6, 4.2 × 10^6, 4.3 × 10^6, 4.4 × 10^6, 4.5 × 10^6, 4.6 × 10^6, 4.7 × 10^6, 4.8 × 10^6, 4.9 × 10^6, 5 × 10^6, 5.1 × 10^6, 5.2 × 10^6, 5.3 × 10^6, 5.4 × 10^6, 5.5 × 10^6, 5.6 × 10^6, 5.7 × 10^6, 5.8 × 10^6, 5.9 × 10^6, 6 × 10^6, 6.1 × 10^6, 6.2 × 10^6, 6.3 × 10^6, 6.4 × 10^6, 6.5 × 10^6, 6.6 × 10^6, 6.7 × 10^6, 6.8 × 10^6, 6.9 × 10^6, 7 × 10^6, 7.1 × 10^6, 7.2 × 10^6, 7.3 × 10^6, 7.4 × 10^6, or 7.5 × 10^6 APCs/cm^2.

[00800] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density selected from a range of at or about 1.0 × 10^6 APCs/cm^2 to at or about 4.5 × 10^6 APCs/cm^2, and the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 2.5 × 10^6 APCs/cm^2 to at or about 7.5 × 10^6 APCs/cm^2.

[00801] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density selected from a range of at or about 1.5

$\times 10^6$ APCs/cm^2 to at or about 3.5 $\times 10^6$ APCs/cm^2, and the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 3.5 $\times 10^6$ APCs/cm^2 to at or about 6 $\times 10^6$ APCs/cm^2.

[00802] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density selected from a range of at or about 2 $\times 10^6$ APCs/cm^2 to at or about 3 $\times 10^6$ APCs/cm^2, and the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density selected from a range of at or about 4 $\times 10^6$ APCs/cm^2 to at or about 5.5 $\times 10^6$ APCs/cm^2.

[00803] In another embodiment, the APCs exogenously supplied in the priming first expansion are seeded in the culture flask at a density at or about 2 $\times 10^6$ APCs/cm^2 and the APCs exogenously supplied in the rapid second expansion are seeded in the culture flask at a density of at or about 4 $\times 10^6$ APCs/cm^2.

[00804] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of PBMCs exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 20:1.

[00805] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of PBMCs exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 10:1.

[00806] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of PBMCs exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 9:1.

[00807] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 8:1.

[00808] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 7:1.

[00809] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 6:1.

[00810] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 5:1.

[00811] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 4:1.

[00812] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 3:1.

[00813] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.9:1.

[00814] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.8:1.

[00815] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.7:1.

[00816] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.6:1.

[00817] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.5:1.

[00818] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.4:1.

[00819] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.3:1.

[00820] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.2:1.

[00821] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2.1:1.

[00822] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 1.1:1 to at or about 2:1.

[00823] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 10:1.

[00824] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 5:1.

[00825] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 4:1.

[00826] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 3:1.

[00827] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.9:1.

[00828] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.8:1.

[00829] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.7:1.

[00830] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.6:1.

[00831] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.5:1.

[00832] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.4:1.

[00833] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.3:1.

[00834] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about about 2:1 to at or about 2.2:1.

[00835] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from a range of from at or about 2:1 to at or about 2.1:1.

[00836] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is at or about 2:1.

[00837] In another embodiment, the ratio of the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion to the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is at or about 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, 2:1, 2.1:1, 2.2:1, 2.3:1, 2.4:1, 2.5:1, 2.6:1, 2.7:1, 2.8:1, 2.9:1, 3:1, 3.1:1, 3.2:1, 3.3:1, 3.4:1, 3.5:1, 3.6:1, 3.7:1, 3.8:1, 3.9:1, 4:1, 4.1:1, 4.2:1, 4.3:1, 4.4:1, 4.5:1, 4.6:1, 4.7:1, 4.8:1, 4.9:1, or 5:1.

[00838] In another embodiment, the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is at or about 1×10^8, 1.1×10^8, 1.2×10^8, 1.3×10^8, 1.4×10^8, 1.5×10^8, 1.6×10^8, 1.7×10^8, 1.8×10^8, 1.9×10^8, 2×10^8, 2.1×10^8, 2.2×10^8, 2.3×10^8, 2.4×10^8, 2.5×10^8, 2.6×10^8, 2.7×10^8, 2.8×10^8, 2.9×10^8, 3×10^8, 3.1×10^8, 3.2×10^8, 3.3×10^8, 3.4×10^8 or 3.5×10^8 APCs (including, for example, PBMCs), and the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is at or about 3.5×10^8, 3.6×10^8, 3.7×10^8, 3.8×10^8, 3.9×10^8, 4×10^8, 4.1×10^8, 4.2×10^8, 4.3×10^8, 4.4×10^8, 4.5×10^8, 4.6×10^8, 4.7×10^8, 4.8×10^8, 4.9×10^8, 5×10^8, 5.1×10^8, 5.2×10^8, 5.3×10^8, 5.4×10^8, 5.5×10^8, 5.6×10^8, 5.7×10^8, 5.8×10^8, 5.9×10^8, 6×10^8, 6.1×10^8, 6.2×10^8, 6.3×10^8, 6.4×10^8, 6.5×10^8,

6.6×10^8, 6.7×10^8, 6.8×10^8, 6.9×10^8, 7×10^8, 7.1×10^8, 7.2×10^8, 7.3×10^8, 7.4×10^8, 7.5×10^8, 7.6×10^8, 7.7×10^8, 7.8×10^8, 7.9×10^8, 8×10^8, 8.1×10^8, 8.2×10^8, 8.3×10^8, 8.4×10^8, 8.5×10^8, 8.6×10^8, 8.7×10^8, 8.8×10^8, 8.9×10^8, 9×10^8, 9.1×10^8, 9.2×10^8, 9.3×10^8, 9.4×10^8, 9.5×10^8, 9.6×10^8, 9.7×10^8, 9.8×10^8, 9.9×10^8 or 1×10^9 APCs (including, for example, PBMCs).

[00839] In another embodiment, the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from the range of at or about 1×10^8 APCs (including, for example, PBMCs) to at or about 3.5×10^8 APCs (including, for example, PBMCs), and the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is selected from the range of at or about 3.5×10^8 APCs (including, for example, PBMCs) to at or about 1×10^9 APCs (including, for example, PBMCs).

[00840] In another embodiment, the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from the range of at or about 1.5×10^8 APCs to at or about 3×10^8 APCs (including, for example, PBMCs), and the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is selected from the range of at or about 4×10^8 APCs (including, for example, PBMCs) to at or about 7.5×10^8 APCs (including, for example, PBMCs).

[00841] In another embodiment, the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is selected from the range of at or about 2×10^8 APCs (including, for example, PBMCs) to at or about 2.5×10^8 APCs (including, for example, PBMCs), and the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is selected from the range of at or about 4.5×10^8 APCs (including, for example, PBMCs) to at or about 5.5×10^8 APCs (including, for example, PBMCs).

[00842] In another embodiment, the number of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion is at or about 2.5×10^8 APCs (including, for example, PBMCs) and the number of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is at or about 5×10^8 APCs (including, for example, PBMCs).

[00843] In an embodiment, the number of layers of APCs (including, for example, PBMCs) added at day 0 of the priming first expansion is approximately one-half of the number of

layers of APCs (including, for example, PBMCs) added at day 7 of the rapid second expansion. In certain embodiments, the method comprises adding antigen presenting cell layers at day 0 of the priming first expansion to the first population of TILs and adding antigen presenting cell layers at day 7 to the second population of TILs, wherein the number of antigen presenting cell layer added at day 0 is approximately 50% of the number of antigen presenting cell layers added at day 7.

[00844] In another embodiment, the number of layers of APCs (including, for example, PBMCs) exogenously supplied at day 7 of the rapid second expansion is greater than the number of layers of APCs (including, for example, PBMCs) exogenously supplied at day 0 of the priming first expansion.

[00845] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 2 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 4 cell layers.

[00846] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about one cell layer and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 3 cell layers.

[00847] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 1.5 cell layers to at or about 2.5 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 3 cell layers.

[00848] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about one cell layer and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 2 cell layers.

[00849] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of of at or about 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 3.1, 3.2, 3.3, 3.4,

3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9 or 8 cell layers.

[00850] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 1 cell layer to at or about 2 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 3 cell layers to at or about 10 cell layers.

[00851] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 2 cell layers to at or about 3 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 4 cell layers to at or about 8 cell layers.

[00852] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 2 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 4 cell layers to at or about 8 cell layers.

[00853] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 1, 2 or 3 cell layers and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with an average thickness of at or about 3, 4, 5, 6, 7, 8, 9 or 10 cell layers.

[00854] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:10.

[00855] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:8.

[00856] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:7.

[00857] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:6.

[00858] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:5.

[00859] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:4.

[00860] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:3.

[00861] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.1 to at or about 1:2.

[00862] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.2 to at or about 1:8.

[00863] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.3 to at or about 1:7.

[00864] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.4 to at or about 1:6.

[00865] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.5 to at or about 1:5.

[00866] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.6 to at or about 1:4.

[00867] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.7 to at or about 1:3.5.

[00868] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.8 to at or about 1:3.

[00869] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from the range of at or about 1:1.9 to at or about 1:2.5.

[00870] In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is at or about 1: 2.

In another embodiment, day 0 of the priming first expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a first average thickness equal to a first number of layers of APCs (including, for example, PBMCs) and day 7 of the rapid second expansion occurs in the presence of layered APCs (including, for example, PBMCs) with a second average thickness equal to a second number of layers of APCs (including, for example, PBMCs), wherein the ratio of the first number of layers of APCs (including, for example, PBMCs) to the second number of layers of APCs (including, for example, PBMCs) is selected from at or about 1:1.1, 1:1.2, 1:1.3, 1:1.4, 1:1.5, 1:1.6, 1:1.7, 1:1.8, 1:1.9, 1:2, 1:2.1, 1:2.2, 1:2.3, 1:2.4, 1:2.5, 1:2.6, 1:2.7, 1:2.8, 1:2.9, 1:3, 1:3.1, 1:3.2, 1:3.3, 1:3.4, 1:3.5, 1:3.6, 1:3.7, 1:3.8, 1:3.9, 1:4, 1:4.1, 1:4.2, 1:4.3, 1:4.4, 1:4.5, 1:4.6, 1:4.7, 1:4.8, 1:4.9, 1:5, 1:5.1, 1:5.2, 1:5.3, 1:5.4, 1:5.5, 1:5.6, 1:5.7, 1:5.8, 1:5.9, 1:6, 1:6.1, 1:6.2, 1:6.3, 1:6.4, 1:6.5, 1:6.6, 1:6.7, 1:6.8, 1:6.9, 1:7, 1:7.1, 1:7.2, 1:7.3, 1:7.4, 1:7.5, 1:7.6, 1:7.7, 1:7.8, 1:7.9, 1:8, 1:8.1, 1:8.2, 1:8.3, 1:8.4, 1:8.5, 1:8.6, 1:8.7, 1:8.8, 1:8.9, 1:9, 1:9.1, 1:9.2, 1:9.3, 1:9.4, 1:9.5, 1:9.6, 1:9.7, 1:9.8, 1:9.9 or 1:10.

V. Optional Cell Medium Components

1. Anti-CD3 Antibodies

[00871] In some embodiments of Process 2A, the culture media used in expansion methods described herein (including those referred to as REP, see for example, Figure 1) also includes an anti-CD3 antibody. An anti-CD3 antibody in combination with IL-2 induces T cell activation and cell division in the TIL population. This effect can be seen with full length antibodies as well as Fab and F(ab')2 fragments, with the former being generally preferred; see, *e.g.,* Tsoukas *et al.*, *J. Immunol.* **1985,** *135,* 1719, hereby incorporated by reference in its entirety.

[00872] As will be appreciated by those in the art, there are a number of suitable anti-human CD3 antibodies that find use in the invention, including anti-human CD3 polyclonal and monoclonal antibodies from various mammals, including, but not limited to, murine, human, primate, rat, and canine antibodies. In particular embodiments, the OKT3 anti-CD3 antibody is used (commercially available from Ortho-McNeil, Raritan, NJ or Miltenyi Biotech, Auburn, CA).

TABLE 4: Amino acid sequences of muromonab (exemplary OKT-3 antibody)

Identifier	Sequence (One-Letter Amino Acid Symbols)
SEQ ID NO:1	QVQLQQSGAE LARPGASVKM SCKASGYTFT RYTMHWVKQR PGQGLEWIGY INPSRGYTNY 60 NQKFKDKATL TTDKSSSTAY MQLSSLTSED SAVYYCARYY DDHYCLDYWG QGTTLTVSSA 120

Muromonab heavy chain	KTTAPSVYPL APVCGGTTGS SVTLGCLVKG YFPEPVTLTW NSGSLSSGVH TFPAVLQSDL YTLSSSVTVT SSTWPSQSIT CNVAHPASST KVDKKIEPRP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK	180 240 300 360 420 450
SEQ ID NO:2 Muromonab light chain	QIVLTQSPAI MSASPGEKVT MTCSASSSVS YMNWYQQKSG TSPKRWIYDT SKLASGVPAH FRGSGSGTSY SLTISGMEAE DAATYYCQQW SSNPFTFGSG TKLEINRADT APTVSIFPPS SEQLTSGGAS VVCFLNNFYP KDINVKWKID GSERQNGVLN SWTDQDSKDS TYSMSSTLTL TKDEYERHNS YTCEATHKTS TSPIVKSFNR NEC	60 120 180 213

2. 4-1BB (CD137) AGONISTS

[00873] In an embodiment, the TNFRSF agonist is a 4-1BB (CD137) agonist. The 4-1BB agonist may be any 4-1BB binding molecule known in the art. The 4-1BB binding molecule may be a monoclonal antibody or fusion protein capable of binding to human or mammalian 4-1BB. The 4-1BB agonists or 4-1BB binding molecules may comprise an immunoglobulin heavy chain of any isotype (*e.g.*, IgG, IgE, IgM, IgD, IgA, and IgY), class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. The 4-1BB agonist or 4-1BB binding molecule may have both a heavy and a light chain. As used herein, the term binding molecule also includes antibodies (including full length antibodies), monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific antibodies), human, humanized or chimeric antibodies, and antibody fragments, *e.g.*, Fab fragments, F(ab′) fragments, fragments produced by a Fab expression library, epitope-binding fragments of any of the above, and engineered forms of antibodies, *e.g.*, scFv molecules, that bind to 4-1BB. In an embodiment, the 4-1BB agonist is an antigen binding protein that is a fully human antibody. In an embodiment, the 4-1BB agonist is an antigen binding protein that is a humanized antibody. In some embodiments, 4-1BB agonists for use in the presently disclosed methods and compositions include anti-4-1BB antibodies, human anti-4-1BB antibodies, mouse anti-4-1BB antibodies, mammalian anti-4-1BB antibodies, monoclonal anti-4-1BB antibodies, polyclonal anti-4-1BB antibodies, chimeric anti-4-1BB antibodies, anti-4-1BB adnectins, anti-4-1BB domain antibodies, single chain anti-4-1BB fragments, heavy chain anti-4-1BB fragments, light chain anti-4-1BB fragments, anti-4-1BB fusion proteins, and fragments, derivatives, conjugates, variants, or biosimilars thereof. Agonistic anti-4-1BB antibodies are known to induce strong immune responses. Lee, *et al.*, *PLOS One* **2013**, *8,* e69677. In a preferred embodiment, the 4-1BB agonist is an agonistic, anti-4-1BB humanized or fully human monoclonal antibody (*i.e.*, an antibody derived from a single cell line). In an embodiment, the 4-1BB agonist is EU-101 (Eutilex Co. Ltd.), utomilumab, or urelumab, or a fragment, derivative, conjugate, variant, or

biosimilar thereof. In a preferred embodiment, the 4-1BB agonist is utomilumab or urelumab, or a fragment, derivative, conjugate, variant, or biosimilar thereof.

[00874] In a preferred embodiment, the 4-1BB agonist or 4-1BB binding molecule may also be a fusion protein. In a preferred embodiment, a multimeric 4-1BB agonist, such as a trimeric or hexameric 4-1BB agonist (with three or six ligand binding domains), may induce superior receptor (4-1BBL) clustering and internal cellular signaling complex formation compared to an agonistic monoclonal antibody, which typically possesses two ligand binding domains. Trimeric (trivalent) or hexameric (or hexavalent) or greater fusion proteins comprising three TNFRSF binding domains and IgG1-Fc and optionally further linking two or more of these fusion proteins are described, *e.g.*, in Gieffers, *et al.*, *Mol. Cancer Therapeutics* **2013,** *12,* 2735-47.

[00875] Agonistic 4-1BB antibodies and fusion proteins are known to induce strong immune responses. In a preferred embodiment, the 4-1BB agonist is a monoclonal antibody or fusion protein that binds specifically to 4-1BB antigen in a manner sufficient to reduce toxicity. In some embodiments, the 4-1BB agonist is an agonistic 4-1BB monoclonal antibody or fusion protein that abrogates antibody-dependent cellular toxicity (ADCC), for example NK cell cytotoxicity. In some embodiments, the 4-1BB agonist is an agonistic 4-1BB monoclonal antibody or fusion protein that abrogates antibody-dependent cell phagocytosis (ADCP). In some embodiments, the 4-1BB agonist is an agonistic 4-1BB monoclonal antibody or fusion protein that abrogates complement-dependent cytotoxicity (CDC). In some embodiments, the 4-1BB agonist is an agonistic 4-1BB monoclonal antibody or fusion protein which abrogates Fc region functionality.

[00876] In some embodiments, the 4-1BB agonists are characterized by binding to human 4-1BB (SEQ ID NO:9) with high affinity and agonistic activity. In an embodiment, the 4-1BB agonist is a binding molecule that binds to human 4-1BB (SEQ ID NO:9). In an embodiment, the 4-1BB agonist is a binding molecule that binds to murine 4-1BB (SEQ ID NO:10). The amino acid sequences of 4-1BB antigen to which a 4-1BB agonist or binding molecule binds are summarized in Table DD.

TABLE 5. Amino acid sequences of 4-1BB antigens.

Identifier	Sequence (One-Letter Amino Acid Symbols)
SEQ ID NO:9 human 4-1BB, Tumor necrosis factor receptor	MGNSCYNIVA TLLLVLNFER TRSLQDPCSN CPAGTFCDNN RNQICSPCPP NSFSSAGGQR 60 TCDICRQCKG VFRTRKECSS TSNAECDCTP GFHCLGAGCS MCEQDCKQGQ ELTKKGCKDC 120 CFGTFNDQKR GICRPWTNCS LDGKSVLVNG TKERDVVCGP SPADLSPGAS SVTPPAPARE 180 PGHSPQIISF FLALTSTALL FLLFFLTLRF SVVKRGRKKL LYIFKQPFMR PVQTTQEEDG 240

superfamily, member 9 (Homo sapiens)	CSCRFPEEEE GGCEL 255
SEQ ID NO:10 murine 4-1BB, Tumor necrosis factor receptor superfamily, member 9 (Mus musculus)	MGNNCYNVVV IVLLLVGCEK VGAVQNSCDN CQPGTFCRKY NPVCKSCPPS TFSSIGGQPN 60 CNICRVCAGY FRFKKFCSST HNAECECIEG FHCLGPQCTR CEKDCRPGQE LTKQGCKTCS 120 LGTFNDQNGT GVCRPWTNCS LDGRSVLKTG TTEKDVVCGP PVVSFSPSTT ISVTPEGGPG 180 GHSLQVLTLF LALTSALLLA LIFITLLFSV LKWIRKKFPH IFKQPFKKTT GAAQEEDACS 240 CRCPQEEEGG GGGYEL 256

[00877] In some embodiments, the compositions, processes and methods described include a 4-1BB agonist that binds human or murine 4-1BB with a K_D of about 100 pM or lower, binds human or murine 4-1BB with a K_D of about 90 pM or lower, binds human or murine 4-1BB with a K_D of about 80 pM or lower, binds human or murine 4-1BB with a K_D of about 70 pM or lower, binds human or murine 4-1BB with a K_D of about 60 pM or lower, binds human or murine 4-1BB with a K_D of about 50 pM or lower, binds human or murine 4-1BB with a K_D of about 40 pM or lower, or binds human or murine 4-1BB with a K_D of about 30 pM or lower.

[00878] In some embodiments, the compositions, processes and methods described include a 4-1BB agonist that binds to human or murine 4-1BB with a k_{assoc} of about 7.5×10^5 1/M·s or faster, binds to human or murine 4-1BB with a k_{assoc} of about 7.5×10^5 1/M·s or faster, binds to human or murine 4-1BB with a k_{assoc} of about 8×10^5 l/M·s or faster, binds to human or murine 4-1BB with a k_{assoc} of about 8.5×10^5 1/M·s or faster, binds to human or murine 4-1BB with a k_{assoc} of about 9×10^5 1/M·s or faster, binds to human or murine 4-1BB with a k_{assoc} of about 9.5×10^5 1/M·s or faster, or binds to human or murine 4-1BB with a k_{assoc} of about 1×10^6 1/M·s or faster.

[00879] In some embodiments, the compositions, processes and methods described include a 4-1BB agonist that binds to human or murine 4-1BB with a k_{dissoc} of about 2×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.1×10^{-5} 1/s or slower , binds to human or murine 4-1BB with a k_{dissoc} of about 2.2×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.3×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.4×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.5×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.6×10^{-5} 1/s or slower or binds to human or murine 4-1BB with a k_{dissoc} of about 2.7×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.8×10^{-5} 1/s or slower, binds to human or murine 4-1BB with a k_{dissoc} of about 2.9×10^{-5} 1/s or slower, or binds to human or murine 4-1BB with a k_{dissoc} of about 3×10^{-5} 1/s or slower.

[00880] In some embodiments, the compositions, processes and methods described include a 4-1BB agonist that binds to human or murine 4-1BB with an IC_{50} of about 10 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 9 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 8 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 7 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 6 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 5 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 4 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 3 nM or lower, binds to human or murine 4-1BB with an IC_{50} of about 2 nM or lower, or binds to human or murine 4-1BB with an IC_{50} of about 1 nM or lower.

[00881] In a preferred embodiment, the 4-1BB agonist is utomilumab, also known as PF-05082566 or MOR-7480, or a fragment, derivative, variant, or biosimilar thereof. Utomilumab is available from Pfizer, Inc. Utomilumab is an immunoglobulin G2-lambda, anti-[*Homo sapiens* TNFRSF9 (tumor necrosis factor receptor (TNFR) superfamily member 9, 4-1BB, T cell antigen ILA, CD137)], *Homo sapiens* (fully human) monoclonal antibody. The amino acid sequences of utomilumab are set forth in Table EE. Utomilumab comprises glycosylation sites at Asn59 and Asn292; heavy chain intrachain disulfide bridges at positions 22-96 (V_H-V_L), 143-199 (C_H1-C_L), 256-316 (C_H2) and 362-420 (C_H3); light chain intrachain disulfide bridges at positions 22'-87' (V_H-V_L) and 136'-195' (C_H1-C_L); interchain heavy chain-heavy chain disulfide bridges at IgG2A isoform positions 218-218, 219-219, 222-222, and 225-225, at IgG2A/B isoform positions 218-130, 219-219, 222-222, and 225-225, and at IgG2B isoform positions 219-130 (2), 222-222, and 225-225; and interchain heavy chain-light chain disulfide bridges at IgG2A isoform positions 130-213' (2), IgG2A/B isoform positions 218-213' and 130-213', and at IgG2B isoform positions 218-213' (2). The preparation and properties of utomilumab and its variants and fragments are described in U.S. Patent Nos. 8,821,867; 8,337,850; and 9,468,678, and International Patent Application Publication No. WO 2012/032433 A1, the disclosures of each of which are incorporated by reference herein. Preclinical characteristics of utomilumab are described in Fisher, *et al.*, *Cancer Immunolog. & Immunother.* **2012,** *61,* 1721-33. Current clinical trials of utomilumab in a variety of hematological and solid tumor indications include U.S. National Institutes of Health clinicaltrials.gov identifiers NCT02444793, NCT01307267, NCT02315066, and NCT02554812.

[00882] In an embodiment, a 4-1BB agonist comprises a heavy chain given by SEQ ID NO:11 and a light chain given by SEQ ID NO:12. In an embodiment, a 4-1BB agonist

comprises heavy and light chains having the sequences shown in SEQ ID NO:11 and SEQ ID NO:12, respectively, or antigen binding fragments, Fab fragments, single-chain variable fragments (scFv), variants, or conjugates thereof. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 99% identical to the sequences shown in SEQ ID NO:11 and SEQ ID NO:12, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 98% identical to the sequences shown in SEQ ID NO:11 and SEQ ID NO:12, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 97% identical to the sequences shown in SEQ ID NO:11 and SEQ ID NO:12, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 96% identical to the sequences shown in SEQ ID NO:11 and SEQ ID NO:12, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 95% identical to the sequences shown in SEQ ID NO:11 and SEQ ID NO:12, respectively.

[00883] In an embodiment, the 4-1BB agonist comprises the heavy and light chain CDRs or variable regions (VRs) of utomilumab. In an embodiment, the 4-1BB agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:13, and the 4-1BB agonist light chain variable region (V_L) comprises the sequence shown in SEQ ID NO:14, and conservative amino acid substitutions thereof. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14, respectively. In an embodiment, a 4-1BB agonist comprises an scFv antibody comprising V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14.

[00884] In an embodiment, a 4-1BB agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:15, SEQ ID NO:16, and SEQ ID NO:17, respectively, and conservative amino acid substitutions thereof, and light chain

CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:18, SEQ ID NO:19, and SEQ ID NO:20, respectively, and conservative amino acid substitutions thereof.

[00885] In an embodiment, the 4-1BB agonist is a 4-1BB agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to utomilumab. In an embodiment, the biosimilar monoclonal antibody comprises an 4-1BB antibody comprising an amino acid sequence which has at least 97% sequence identity, *e.g.,* 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is utomilumab. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a 4-1BB agonist antibody authorized or submitted for authorization, wherein the 4-1BB agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is utomilumab. The 4-1BB agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is utomilumab. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is utomilumab.

TABLE 6. Amino acid sequences for 4-1BB agonist antibodies related to utomilumab.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:11 heavy chain for utomilumab	EVQLVQSGAE VKKPGESLRI SCKGSGYSFS TYWISWVRQM PGKGLEWMGK IYPGDSYTNY SPSFQGQVTI SADKSISTAY LQWSSLKASD TAMYYCARGY GIFDYWGQGT LVTVSSASTK GPSVFPLAPC SRSTSESTAA LGCLVKDYFP EPVTVSWNSG ALTSGVHTFP AVLQSSGLYS LSSVVTVPSS NFGTQTYTCN VDHKPSNTKV DKTVERKCCV ECPPCPAPPV AGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVQF NWYVDGVEVH NAKTKPREEQ FNSTFRVVSV LTVVHQDWLN GKEYKCKVSN KGLPAPIEKT ISKTKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS DIAVEWESNG QPENNYKTTP PMLDSDGSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNH YTQKSLSLSP G	60 120 180 240 300 360 420 441
SEQ ID NO:12 light chain for utomilumab	SYELTQPPSV SVSPGQTASI TCSGDNIGDQ YAHWYQQKPG QSPVLVIYQD KNRPSGIPER FSGSNSGNTA TLTISGTQAM DEADYYCATY TGFGSLAVFG GGTKLTVLGQ PKAAPSVTLF PPSSEELQAN KATLVCLISD FYPGAVTVAW KADSSPVKAG VETTTPSKQS NNKYAASSYL	60 120 180

	SLTPEQWKSH RSYSCQVTHE GSTVEKTVAP TECS	214
SEQ ID NO:13 heavy chain variable region for utomilumab	EVQLVQSGAE VKKPGESLRI SCKGSGYSFS TYWISWVRQM PGKGLEWMG KIYPGDSYTN YSPSFQGQVT ISADKSISTA YLQWSSLKAS DTAMYYCARG YGIFDYWGQ GTLVTVSS	60 118
SEQ ID NO:14 light chain variable region for utomilumab	SYELTQPPSV SVSPGQTASI TCSGDNIGDQ YAHWYQQKPG QSPVLVIYQD KNRPSGIPER FSGSNSGNTA TLTISGTQAM DEADYYCATY TGFGSLAVFG GGTKLTVL	60 108
SEQ ID NO:15 heavy chain CDR1 for utomilumab	STYWIS	6
SEQ ID NO:16 heavy chain CDR2 for utomilumab	KIYPGDSYTN YSPSFQG	17
SEQ ID NO:17 heavy chain CDR3 for utomilumab	RGYGIFDY	8
SEQ ID NO:18 light chain CDR1 for utomilumab	SGDNIGDQYA H	11
SEQ ID NO:19 light chain CDR2 for utomilumab	QDKNRPS	7
SEQ ID NO:20 light chain CDR3 for utomilumab	ATYTGFGSLA V	11

[00886] In a preferred embodiment, the 4-1BB agonist is the monoclonal antibody urelumab, also known as BMS-663513 and 20H4.9.h4a, or a fragment, derivative, variant, or biosimilar thereof. Urelumab is available from Bristol-Myers Squibb, Inc., and Creative Biolabs, Inc. Urelumab is an immunoglobulin G4-kappa, anti-[*Homo sapiens* TNFRSF9 (tumor necrosis factor receptor superfamily member 9, 4-1BB, T cell antigen ILA, CD137)], *Homo sapiens* (fully human) monoclonal antibody. The amino acid sequences of urelumab are set forth in Table EE. Urelumab comprises N-glycosylation sites at positions 298 (and 298'') ; heavy chain intrachain disulfide bridges at positions 22-95 (V_H-V_L), 148-204 (C_H1-C_L), 262-322 (C_H2) and 368-426 (C_H3) (and at positions 22''-95'', 148''-204'', 262''-322'', and 368''-426'') ; light chain intrachain disulfide bridges at positions 23'-88' (V_H-V_L) and 136'-196' (C_H1-C_L) (and at positions 23'''-88''' and 136'''-196''') ; interchain heavy chain-heavy chain disulfide bridges at positions 227-227'' and 230-230'' ; and interchain heavy chain-light chain disulfide bridges at 135-216' and 135''-216'''. The preparation and properties of urelumab and its variants and fragments are described in U.S. Patent Nos. 7,288,638 and 8,962,804, the disclosures of which are incorporated by reference herein. The preclinical and clinical characteristics of urelumab are described in Segal, *et al.*, *Clin. Cancer Res.* **2016,** *available at* http:/dx.doi.org/ 10.1158/1078-0432.CCR-16-1272. Current clinical trials of urelumab in a variety of hematological and solid tumor indications include U.S. National Institutes of Health clinicaltrials.gov identifiers NCT01775631, NCT02110082, NCT02253992, and NCT01471210.

[00887] In an embodiment, a 4-1BB agonist comprises a heavy chain given by SEQ ID NO:21 and a light chain given by SEQ ID NO:22. In an embodiment, a 4-1BB agonist comprises heavy and light chains having the sequences shown in SEQ ID NO:21 and SEQ ID NO:22, respectively, or antigen binding fragments, Fab fragments, single-chain variable fragments (scFv), variants, or conjugates thereof. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 99% identical to the sequences shown in SEQ ID NO:21 and SEQ ID NO:22, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 98% identical to the sequences shown in SEQ ID NO:21 and SEQ ID NO:22, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 97% identical to the sequences shown in SEQ ID NO:21 and SEQ ID NO:22, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 96% identical to the sequences shown in SEQ ID NO:21 and SEQ ID NO:22, respectively. In an embodiment, a 4-1BB agonist comprises heavy and light chains that are each at least 95% identical to the sequences shown in SEQ ID NO:21 and SEQ ID NO:22, respectively.

[00888] In an embodiment, the 4-1BB agonist comprises the heavy and light chain CDRs or variable regions (VRs) of urelumab. In an embodiment, the 4-1BB agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:23, and the 4-1BB agonist light chain variable region (V_L) comprises the sequence shown in SEQ ID NO:24, and conservative amino acid substitutions thereof. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24, respectively. In an embodiment, a 4-1BB agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24, respectively. In an embodiment, a 4-1BB agonist comprises an scFv antibody comprising V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24.

[00889] In an embodiment, a 4-1BB agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27, respectively, and conservative amino acid substitutions thereof, and light chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:28, SEQ ID NO:29, and SEQ ID NO:30, respectively, and conservative amino acid substitutions thereof.

[00890] In an embodiment, the 4-1BB agonist is a 4-1BB agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to urelumab. In an embodiment, the biosimilar monoclonal antibody comprises an 4-1BB antibody comprising an amino acid sequence which has at least 97% sequence identity, *e.g.,* 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is urelumab. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a 4-1BB agonist antibody authorized or submitted for authorization, wherein the 4-1BB agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is urelumab. The 4-1BB agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is urelumab. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is urelumab.

TABLE 7: Amino acid sequences for 4-1BB agonist antibodies related to urelumab.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:21 heavy chain for urelumab	QVQLQQWGAG LLKPSETLSL TCAVYGGSFS GYYWSWIRQS PEKGLEWIGE INHGGYVTYN PSLESRVTIS VDTSKNQFSL KLSSVTAADT AVYYCARDYG PGNYDWYFDL WGRGTLVTVS SASTKGPSVF PLAPCSRSTS ESTAALGCLV KDYFPEPVTV SWNSGALTSG VHTFPAVLQS SGLYSLSSVV TVPSSSLGTK TYTCNVDHKP SNTKVDKRVE SKYGPPCPPC PAPEFLGGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSQE DPEVQFNWYV DGVEVHNAKT KPREEQFNST YRVVSVLTVL HQDWLNGKEY KCKVSNKGLP SSIEKTISKA KGQPREPQVY TLPPSQEEMT	60 120 180 240 300 360

	KNQVSLTCLV KGFYPSDIAV EWESNGQPEN NYKTTPPVLD SDGSFFLYSR LTVDKSRWQE GNVFSCSVMH EALHNHYTQK SLSLSLGK	420 448
SEQ ID NO:22 light chain for urelumab	EIVLTQSPAT LSLSPGERAT LSCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA RFSGSGSGTD FTLTISSLEP EDFAVYYCQQ RSNWPPALTF CGGTKVEIKR TVAAPSVFIF PPSDEQLKSG TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYSLSST LTLSKADYEK HKVYACEVTH QGLSSPVTKS FNRGEC	60 120 180 216
SEQ ID NO:23 variable heavy chain for urelumab	MKHLWFFLLL VAAPRWVLSQ VQLQQWGAGL LKPSETLSLT CAVYGGSFSG YYWSWIRQSP EKGLEWIGEI NHGGYVTYNP SLESRVTISV DTSKNQFSLK LSSVTAADTA VYYCARDYGP	60 120
SEQ ID NO:24 variable light chain for urelumab	MEAPAQLLFL LLLWLPDTTG EIVLTQSPAT LSLSPGERAT LSCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA RFSGSGSGTD FTLTISSLEP EDFAVYYCQQ	60 110
SEQ ID NO:25 heavy chain CDR1 for urelumab	GYYWS	5
SEQ ID NO:26 heavy chain CDR2 for urelumab	EINHGGYVTY NPSLES	16
SEQ ID NO:27 heavy chain CDR3 for urelumab	DYGPGNYDWY FDL	13
SEQ ID NO:28 light chain CDR1 for urelumab	RASQSVSSYL A	11
SEQ ID NO:29 light chain CDR2 for urelumab	DASNRAT	7
SEQ ID NO:30 light chain CDR3 for urelumab	QQRSDWPPAL T	11

[00891] In an embodiment, the 4-1BB agonist is selected from the group consisting of 1D8, 3Elor, 4B4 (BioLegend 309809), H4-1BB-M127 (BD Pharmingen 552532), BBK2 (Thermo Fisher MS621PABX), 145501 (Leinco Technologies B591), the antibody produced by cell line deposited as ATCC No. HB-11248 and disclosed in U.S. Patent No. 6,974,863, 5F4 (BioLegend 31 1503), C65-485 (BD Pharmingen 559446), antibodies disclosed in U.S. Patent Application Publication No. US 2005/0095244, antibodies disclosed in U.S. Patent No. 7,288,638 (such as 20H4.9-IgGl (BMS-663031)), antibodies disclosed in U.S. Patent No. 6,887,673 (such as 4E9 or BMS-554271), antibodies disclosed in U.S. Patent No. 7,214,493, antibodies disclosed in U.S. Patent No. 6,303,121, antibodies disclosed in U.S. Patent No. 6,569,997, antibodies disclosed in U.S. Patent No. 6,905,685 (such as 4E9 or BMS-554271), antibodies disclosed in U.S. Patent No. 6,362,325 (such as 1D8 or BMS-469492; 3H3 or BMS-469497; or 3El), antibodies disclosed in U.S. Patent No. 6,974,863 (such as 53A2); antibodies disclosed in U.S. Patent No. 6,210,669 (such as 1D8, 3B8, or 3El), antibodies described in U.S. Patent No. 5,928,893, antibodies disclosed in U.S. Patent No. 6,303,121, antibodies disclosed in U.S. Patent No. 6,569,997, antibodies disclosed in International Patent Application Publication Nos. WO 2012/177788, WO 2015/119923, and WO 2010/042433, and fragments, derivatives, conjugates, variants, or biosimilars thereof, wherein the disclosure of each of the foregoing patents or patent application publications is incorporated by reference here.

[00892] In an embodiment, the 4-1BB agonist is a 4-1BB agonistic fusion protein described in International Patent Application Publication Nos. WO 2008/025516 A1, WO 2009/007120 A1, WO 2010/003766 A1, WO 2010/010051 A1, and WO 2010/078966 A1; U.S. Patent Application Publication Nos. US 2011/0027218 A1, US 2015/0126709 A1, US 2011/0111494 A1, US 2015/0110734 A1, and US 2015/0126710 A1; and U.S. Patent Nos. 9,359,420, 9,340,599, 8,921,519, and 8,450,460, the disclosures of which are incorporated by reference herein.

[00893] In an embodiment, the 4-1BB agonist is a 4-1BB agonistic fusion protein as depicted in Structure I-A (C-terminal Fc-antibody fragment fusion protein) or Structure I-B (N-terminal Fc-antibody fragment fusion protein), or a fragment, derivative, conjugate, variant, or biosimilar thereof:

(I-A) (I-B)



In structures I-A and I-B, the cylinders refer to individual polypeptide binding domains. Structures I-A and I-B comprise three linearly-linked TNFRSF binding domains derived from *e.g.*, 4-1BBL or an antibody that binds 4-1BB, which fold to form a trivalent protein, which is then linked to a second triavelent protein through IgG1-Fc (including C_H3 and C_H2 domains) is then used to link two of the trivalent proteins together through disulfide bonds (small elongated ovals), stabilizing the structure and providing an agonists capable of bringing together the intracellular signaling domains of the six receptors and signaling proteins to form a signaling complex. The TNFRSF binding domains denoted as cylinders may be scFv domains comprising, *e.g.*, a V_H and a V_L chain connected by a linker that may comprise

hydrophilic residues and Gly and Ser sequences for flexibility, as well as Glu and Lys for solubility. Any scFv domain design may be used, such as those described in de Marco, *Microbial Cell Factories*, **2011,** *10*, 44; Ahmad, *et al.*, *Clin. & Dev. Immunol.* **2012,** 980250; Monnier, *et al.*, *Antibodies,* **2013,** *2,* 193-208; or in references incorporated elsewhere herein. Fusion protein structures of this form are described in U.S. Patent Nos. 9,359,420, 9,340,599, 8,921,519, and 8,450,460, the disclosures of which are incorporated by reference herein.

[00894] Amino acid sequences for the other polypeptide domains of structure I-A are given in Table GG. The Fc domain preferably comprises a complete constant domain (amino acids 17-230 of SEQ ID NO:31) the complete hinge domain (amino acids 1-16 of SEQ ID NO:31) or a portion of the hinge domain (*e.g.*, amino acids 4-16 of SEQ ID NO:31). Preferred linkers for connecting a C-terminal Fc-antibody may be selected from the embodiments given in SEQ ID NO:32 to SEQ ID NO:41, including linkers suitable for fusion of additional polypeptides.

TABLE 8: Amino acid sequences for TNFRSF fusion proteins, including 4-1BB fusion proteins, with C-terminal Fc-antibody fragment fusion protein design (structure I-A).

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:31 Fc domain	KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK	60 120 180 230
SEQ ID NO:32 linker	GGPGSSKSCD KTHTCPPCPA PE	22
SEQ ID NO:33 linker	GGSGSSKSCD KTHTCPPCPA PE	22
SEQ ID NO:34 linker	GGPGSSSSSS SKSCDKTHTC PPCPAPE	27
SEQ ID NO:35 linker	GGSGSSSSSS SKSCDKTHTC PPCPAPE	27
SEQ ID NO:36 linker	GGPGSSSSSS SSSKSCDKTH TCPPCPAPE	29
SEQ ID NO:37 linker	GGSGSSSSSS SSSKSCDKTH TCPPCPAPE	29
SEQ ID NO:38 linker	GGPGSSGSGS SDKTHTCPPC PAPE	24
SEQ ID NO:39 linker	GGPGSSGSGS DKTHTCPPCP APE	23
SEQ ID NO:40 linker	GGPSSSGSDK THTCPPCPAP E	21
SEQ ID NO:41 linker	GGSSSSSSSS GSDKTHTCPP CPAPE	25

[00895] Amino acid sequences for the other polypeptide domains of structure I-B are given in Table 9. If an Fc antibody fragment is fused to the N-terminus of an TNRFSF fusion protein as in structure I-B, the sequence of the Fc module is preferably that shown in SEQ ID NO:42, and the linker sequences are preferably selected from those embodiments set forth in SED ID NO:43 to SEQ ID NO:45.

TABLE 9: Amino acid sequences for TNFRSF fusion proteins, including 4-1BB fusion proteins, with N-terminal Fc-antibody fragment fusion protein design (structure I-B).

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:42 Fc domain	METDTLLLWV LLLWVPAGNG DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTPPVLDS DGSFFLYSKL TVDKSRWQQG NVFSCSVMHE ALHNHYTQKS LSLSPG	60 120 180 240 246
SEQ ID NO:43 linker	SGSGSGSGSG S	11
SEQ ID NO:44 linker	SSSSSSGSGS GS	12
SEQ ID NO:45 linker	SSSSSSGSGS GSGSGS	16

[00896] In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains selected from the group consisting of a variable heavy chain and variable light chain of utomilumab, a variable heavy chain and variable light chain of urelumab, a variable heavy chain and variable light chain of utomilumab, a variable heavy chain and variable light chain selected from the variable heavy chains and variable light chains described in Table GG, any combination of a variable heavy chain and variable light chain of the foregoing, and fragments, derivatives, conjugates, variants, and biosimilars thereof.

[00897] In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains comprising a 4-1BBL sequence. In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains comprising a sequence according to SEQ ID NO:46. In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains comprising a soluble 4-1BBL sequence. In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains comprising a sequence according to SEQ ID NO:47.

[00898] In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:13 and SEQ ID NO:14, respectively, wherein the V_H and V_L domains are connected by a linker. In an embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:23 and SEQ ID NO:24, respectively, wherein the V_H and V_L domains are connected by a linker. In an

embodiment, a 4-1BB agonist fusion protein according to structures I-A or I-B comprises one or more 4-1BB binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the V_H and V_L sequences given in Table II, wherein the V_H and V_L domains are connected by a linker.

TABLE 10: Additional polypeptide domains useful as 4-1BB binding domains in fusion proteins or as scFv 4-1BB agonist antibodies.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:46 4-1BBL	MEYASDASLD PEAPWPPAPR ARACRVLPWA LVAGLLLLLL LAAACAVFLA CPWAVSGARA SPGSAASPRL REGPELSPDD PAGLLDLRQG MFAQLVAQNV LLIDGPLSWY SDPGLAGVSL TGGLSYKEDT KELVVAKAGV YYVFFQLELR RVVAGEGSGS VSLALHLQPL RSAAGAAALA LTVDLPPASS EARNSAFGFQ GRLLHLSAGQ RLGVHLHTEA RARHAWQLTQ GATVLGLFRV TPEIPAGLPS PRSE	60 120 180 240 254
SEQ ID NO:47 4-1BBL soluble domain	LRQGMFAQLV AQNVLLIDGP LSWYSDPGLA GVSLTGGLSY KEDTKELVVA KAGVYYVFFQ LELRRVVAGE GSGSVSLALH LQPLRSAAGA AALALTVDLP PASSEARNSA FGFQGRLLHL SAGQRLGVHL HTEARARHAW QLTQGATVLG LFRVTPEIPA GLPSPRSE	60 120 168
SEQ ID NO:48 variable heavy chain for 4B4-1-1 version 1	QVQLQQPGAE LVKPGASVKL SCKASGYTFS SYWMHWVKQR PGQVLEWIGE INPGNGHTNY NEKFKSKATL TVDKSSSTAY MQLSSLTSED SAVYYCARSF TTARGFAYWG QGTLVTVS	60 118
SEQ ID NO:49 variable light chain for 4B4-1-1 version 1	DIVMTQSPAT QSVTPGDRVS LSCRASQTIS DYLHWYQQKS HESPRLLIKY ASQSISGIPS RFSGSGSGSD FTLSINSVEP EDVGVYYCQD GHSFPPTFGG GTKLEIK	60 107
SEQ ID NO:50 variable heavy chain for 4B4-1-1 version 2	QVQLQQPGAE LVKPGASVKL SCKASGYTFS SYWMHWVKQR PGQVLEWIGE INPGNGHTNY NEKFKSKATL TVDKSSSTAY MQLSSLTSED SAVYYCARSF TTARGFAYWG QGTLVTVSA	60 119
SEQ ID NO:51 variable light chain for 4B4-1-1 version 2	DIVMTQSPAT QSVTPGDRVS LSCRASQTIS DYLHWYQQKS HESPRLLIKY ASQSISGIPS RFSGSGSGSD FTLSINSVEP EDVGVYYCQD GHSFPPTFGG GTKLEIKR	60 108
SEQ ID NO:52 variable heavy chain for H39E3-2	MDWTWRILFL VAAATGAHSE VQLVESGGGL VQPGGSLRLS CAASGFTFSD YWMSWVRQAP GKGLEWVADI KNDGSYTNYA PSLTNRFTIS RDNAKNSLYL QMNSLRAEDT AVYYCARELT	60 120
SEQ ID NO:53 variable light chain for H39E3-2	MEAPAQLLFL LLLWLPDTTG DIVMTQSPDS LAVSLGERAT INCKSSQSLL SSGNQKNYL WYQQKPGQPP KLLIYYASTR QSGVPDRFSG SGSGTDFTLT ISSLQAEDVA	60 110

[00899] In an embodiment, the 4-1BB agonist is a 4-1BB agonistic single-chain fusion polypeptide comprising (i) a first soluble 4-1BB binding domain, (ii) a first peptide linker, (iii) a second soluble 4-1BB binding domain, (iv) a second peptide linker, and (v) a third soluble 4-1BB binding domain, further comprising an additional domain at the N-terminal and/or C-terminal end, and wherein the additional domain is a Fab or Fc fragment domain. In an embodiment, the 4-1BB agonist is a 4-1BB agonistic single-chain fusion polypeptide comprising (i) a first soluble 4-1BB binding domain, (ii) a first peptide linker, (iii) a second soluble 4-1BB binding domain, (iv) a second peptide linker, and (v) a third soluble 4-1BB binding domain, further comprising an additional domain at the N-terminal and/or C-terminal end, wherein the additional domain is a Fab or Fc fragment domain, wherein each of the soluble 4-1BB domains lacks a stalk region (which contributes to trimerisation and provides a

certain distance to the cell membrane, but is not part of the 4-1BB binding domain) and the first and the second peptide linkers independently have a length of 3-8 amino acids.

[00900] In an embodiment, the 4-1BB agonist is a 4-1BB agonistic single-chain fusion polypeptide comprising (i) a first soluble tumor necrosis factor (TNF) superfamily cytokine domain, (ii) a first peptide linker, (iii) a second soluble TNF superfamily cytokine domain, (iv) a second peptide linker, and (v) a third soluble TNF superfamily cytokine domain, wherein each of the soluble TNF superfamily cytokine domains lacks a stalk region and the first and the second peptide linkers independently have a length of 3-8 amino acids, and wherein each TNF superfamily cytokine domain is a 4-1BB binding domain.

[00901] In an embodiment, the 4-1BB agonist is a 4-1BB agonistic scFv antibody comprising any of the foregoing V_H domains linked to any of the foregoing V_L domains.

[00902] In an embodiment, the 4-1BB agonist is BPS Bioscience 4-1BB agonist antibody catalog no. 79097-2, commercially available from BPS Bioscience, San Diego, CA, USA. In an embodiment, the 4-1BB agonist is Creative Biolabs 4-1BB agonist antibody catalog no. MOM-18179, commercially available from Creative Biolabs, Shirley, NY, USA.

3. OX40 (CD134) AGONISTS

[00903] In an embodiment, the TNFRSF agonist is an OX40 (CD134) agonist. The OX40 agonist may be any OX40 binding molecule known in the art. The OX40 binding molecule may be a monoclonal antibody or fusion protein capable of binding to human or mammalian OX40. The OX40 agonists or OX40 binding molecules may comprise an immunoglobulin heavy chain of any isotype (*e.g.*, IgG, IgE, IgM, IgD, IgA, and IgY), class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. The OX40 agonist or OX40 binding molecule may have both a heavy and a light chain. As used herein, the term binding molecule also includes antibodies (including full length antibodies), monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific antibodies), human, humanized or chimeric antibodies, and antibody fragments, *e.g.*, Fab fragments, F(ab′) fragments, fragments produced by a Fab expression library, epitope-binding fragments of any of the above, and engineered forms of antibodies, *e.g.*, scFv molecules, that bind to OX40. In an embodiment, the OX40 agonist is an antigen binding protein that is a fully human antibody. In an embodiment, the OX40 agonist is an antigen binding protein that is a humanized antibody. In some embodiments, OX40 agonists for use in the presently disclosed methods and compositions include anti-

OX40 antibodies, human anti-OX40 antibodies, mouse anti-OX40 antibodies, mammalian anti-OX40 antibodies, monoclonal anti-OX40 antibodies, polyclonal anti-OX40 antibodies, chimeric anti-OX40 antibodies, anti-OX40 adnectins, anti-OX40 domain antibodies, single chain anti-OX40 fragments, heavy chain anti-OX40 fragments, light chain anti-OX40 fragments, anti-OX40 fusion proteins, and fragments, derivatives, conjugates, variants, or biosimilars thereof. In a preferred embodiment, the OX40 agonist is an agonistic, anti-OX40 humanized or fully human monoclonal antibody (*i.e.*, an antibody derived from a single cell line).

[00904] In a preferred embodiment, the OX40 agonist or OX40 binding molecule may also be a fusion protein. OX40 fusion proteins comprising an Fc domain fused to OX40L are described, for example, in Sadun, *et al.*, *J. Immunother.* **2009,** *182,* 1481-89. In a preferred embodiment, a multimeric OX40 agonist, such as a trimeric or hexameric OX40 agonist (with three or six ligand binding domains), may induce superior receptor (OX40L) clustering and internal cellular signaling complex formation compared to an agonistic monoclonal antibody, which typically possesses two ligand binding domains. Trimeric (trivalent) or hexameric (or hexavalent) or greater fusion proteins comprising three TNFRSF binding domains and IgG1-Fc and optionally further linking two or more of these fusion proteins are described, *e.g.*, in Gieffers, *et al.*, *Mol. Cancer Therapeutics* **2013,** *12,* 2735-47.

[00905] Agonistic OX40 antibodies and fusion proteins are known to induce strong immune responses. Curti, *et al.*, *Cancer Res.* **2013**, *73,* 7189-98. In a preferred embodiment, the OX40 agonist is a monoclonal antibody or fusion protein that binds specifically to OX40 antigen in a manner sufficient to reduce toxicity. In some embodiments, the OX40 agonist is an agonistic OX40 monoclonal antibody or fusion protein that abrogates antibody-dependent cellular toxicity (ADCC), for example NK cell cytotoxicity. In some embodiments, the OX40 agonist is an agonistic OX40 monoclonal antibody or fusion protein that abrogates antibody-dependent cell phagocytosis (ADCP). In some embodiments, the OX40 agonist is an agonistic OX40 monoclonal antibody or fusion protein that abrogates complement-dependent cytotoxicity (CDC). In some embodiments, the OX40 agonist is an agonistic OX40 monoclonal antibody or fusion protein which abrogates Fc region functionality.

[00906] In some embodiments, the OX40 agonists are characterized by binding to human OX40 (SEQ ID NO:54) with high affinity and agonistic activity. In an embodiment, the OX40 agonist is a binding molecule that binds to human OX40 (SEQ ID NO:54). In an embodiment, the OX40 agonist is a binding molecule that binds to murine OX40 (SEQ ID

NO:55). The amino acid sequences of OX40 antigen to which an OX40 agonist or binding molecule binds are summarized in Table 11.

TABLE 11: Amino acid sequences of OX40 antigens.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:54 human OX40 (Homo sapiens)	MCVGARRLGR GPCAALLLLG LGLSTVTGLH CVGDTYPSND RCCHECRPGN GMVSRCSRSQ NTVCRPCGPG FYNDVVSSKP CKPCTWCNLR SGSERKQLCT ATQDTVCRCR AGTQPLDSYK PGVDCAPCPP GHFSPGDNQA CKPWTNCTLA GKHTLQPASN SSDAICEDRD PPATQPQETQ GPPARPITVQ PTEAWPRTSQ GPSTRPVEVP GGRAVAAILG LGLVLGLLGP LAILLALYLL RRDQRLPPDA HKPPGGGSFR TPIQEEQADA HSTLAKI	60 120 180 240 277
SEQ ID NO:55 murine OX40 (Mus musculus)	MYVWVQQPTA LLLLGLTLGV TARRLNCVKH TYPSGHKCCR ECQPGHGMVS RCDHTRDTLC HPCETGFYNE AVNYDTCKQC TQCNHRSGSE LKQNCTPTQD TVCRCRPGTQ PRQDSGYKLG VDCVPCPPGH FSPGNNQACK PWTNCTLSGK QTRHPASDSL DAVCEDRSLL ATLLWETQRP TFRPTTVQST TVWPRTSELP SPPTLVTPEG PAFAVLLGLG LGLLAPLTVL LALYLLRKAW RLPNTPKPCW GNSFRTPIQE EHTDAHFTLA KI	60 120 180 240 272

[00907] In some embodiments, the compositions, processes and methods described include a OX40 agonist that binds human or murine OX40 with a K_D of about 100 pM or lower, binds human or murine OX40 with a K_D of about 90 pM or lower, binds human or murine OX40 with a K_D of about 80 pM or lower, binds human or murine OX40 with a K_D of about 70 pM or lower, binds human or murine OX40 with a K_D of about 60 pM or lower, binds human or murine OX40 with a K_D of about 50 pM or lower, binds human or murine OX40 with a K_D of about 40 pM or lower, or binds human or murine OX40 with a K_D of about 30 pM or lower.

[00908] In some embodiments, the compositions, processes and methods described include a OX40 agonist that binds to human or murine OX40 with a k_{assoc} of about 7.5×10^5 1/M·s or faster, binds to human or murine OX40 with a k_{assoc} of about 7.5×10^5 1/M·s or faster, binds to human or murine OX40 with a k_{assoc} of about 8×10^5 1/M·s or faster, binds to human or murine OX40 with a k_{assoc} of about 8.5×10^5 1/M·s or faster, binds to human or murine OX40 with a k_{assoc} of about 9×10^5 1/M·s or faster, binds to human or murine OX40 with a k_{assoc} of about 9.5×10^5 1/M·s or faster, or binds to human or murine OX40 with a k_{assoc} of about 1×10^6 1/M·s or faster.

[00909] In some embodiments, the compositions, processes and methods described include a OX40 agonist that binds to human or murine OX40 with a k_{dissoc} of about 2×10^{-5} 1/s or slower, binds to human or murine OX40 with a k_{dissoc} of about 2.1×10^{-5} 1/s or slower , binds to human or murine OX40 with a k_{dissoc} of about 2.2×10^{-5} 1/s or slower, binds to human or murine OX40 with a k_{dissoc} of about 2.3×10^{-5} 1/s or slower, binds to human or murine OX40 with a k_{dissoc} of about 2.4×10^{-5} 1/s or slower, binds to human or murine OX40 with a k_{dissoc} of about 2.5×10^{-5} 1/s or slower, binds to human or murine OX40 with a k_{dissoc} of about 2.6×10^{-5} 1/s or slower or binds to human or murine OX40 with a k_{dissoc} of about 2.7×10^{-5} 1/s or slower, binds to human or murine OX40 with a k_{dissoc} of about 2.8×10^{-5} 1/s or slower, binds

to human or murine OX40 with a k_{dissoc} of about 2.9×10^{-5} 1/s or slower, or binds to human or murine OX40 with a k_{dissoc} of about 3×10^{-5} 1/s or slower.

[00910] In some embodiments, the compositions, processes and methods described include OX40 agonist that binds to human or murine OX40 with an IC_{50} of about 10 nM or lower, binds to human or murine OX40 with an IC_{50} of about 9 nM or lower, binds to human or murine OX40 with an IC_{50} of about 8 nM or lower, binds to human or murine OX40 with an IC_{50} of about 7 nM or lower, binds to human or murine OX40 with an IC_{50} of about 6 nM or lower, binds to human or murine OX40 with an IC_{50} of about 5 nM or lower, binds to human or murine OX40 with an IC_{50} of about 4 nM or lower, binds to human or murine OX40 with an IC_{50} of about 3 nM or lower, binds to human or murine OX40 with an IC_{50} of about 2 nM or lower, or binds to human or murine OX40 with an IC_{50} of about 1 nM or lower.

[00911] In some embodiments, the OX40 agonist is tavolixizumab, also known as MEDI0562 or MEDI-0562. Tavolixizumab is available from the MedImmune subsidiary of AstraZeneca, Inc. Tavolixizumab is immunoglobulin G1-kappa, anti-[*Homo sapiens* TNFRSF4 (tumor necrosis factor receptor (TNFR) superfamily member 4, OX40, CD134)], humanized and chimeric monoclonal antibody. The amino acid sequences of tavolixizumab are set forth in Table KK. Tavolixizumab comprises N-glycosylation sites at positions 301 and 301'', with fucosylated complex bi-antennary CHO-type glycans; heavy chain intrachain disulfide bridges at positions 22-95 (V_H-V_L), 148-204 (C_H1-C_L), 265-325 (C_H2) and 371-429 (C_H3) (and at positions 22''-95'', 148''-204'', 265''-325'', and 371''-429''); light chain intrachain disulfide bridges at positions 23'-88' (V_H-V_L) and 134'-194' (C_H1-C_L) (and at positions 23'''-88''' and 134'''-194'''); interchain heavy chain-heavy chain disulfide bridges at positions 230-230'' and 233-233''; and interchain heavy chain-light chain disulfide bridges at 224-214' and 224''-214'''. Current clinical trials of tavolixizumab in a variety of solid tumor indications include U.S. National Institutes of Health clinicaltrials.gov identifiers NCT02318394 and NCT02705482.

[00912] In an embodiment, a OX40 agonist comprises a heavy chain given by SEQ ID NO:56 and a light chain given by SEQ ID NO:57. In an embodiment, a OX40 agonist comprises heavy and light chains having the sequences shown in SEQ ID NO:56 and SEQ ID NO:57, respectively, or antigen binding fragments, Fab fragments, single-chain variable fragments (scFv), variants, or conjugates thereof. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 99% identical to the sequences shown in SEQ ID NO:56 and SEQ ID NO:57, respectively. In an embodiment, a OX40 agonist

comprises heavy and light chains that are each at least 98% identical to the sequences shown in SEQ ID NO:56 and SEQ ID NO:57, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 97% identical to the sequences shown in SEQ ID NO:56 and SEQ ID NO:57, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 96% identical to the sequences shown in SEQ ID NO:56 and SEQ ID NO:57, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 95% identical to the sequences shown in SEQ ID NO:56 and SEQ ID NO:57, respectively.

[00913] In an embodiment, the OX40 agonist comprises the heavy and light chain CDRs or variable regions (VRs) of tavolixizumab. In an embodiment, the OX40 agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:58, and the OX40 agonist light chain variable region (V_L) comprises the sequence shown in SEQ ID NO:59, and conservative amino acid substitutions thereof. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59, respectively. In an embodiment, an OX40 agonist comprises an scFv antibody comprising V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59.

[00914] In an embodiment, a OX40 agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:60, SEQ ID NO:61, and SEQ ID NO:62, respectively, and conservative amino acid substitutions thereof, and light chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:63, SEQ ID NO:64, and SEQ ID NO:65, respectively, and conservative amino acid substitutions thereof.

[00915] In an embodiment, the OX40 agonist is a OX40 agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to tavolixizumab. In an embodiment, the biosimilar monoclonal antibody comprises an OX40 antibody comprising

an amino acid sequence which has at least 97% sequence identity, *e.g.*, 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is tavolixizumab. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a OX40 agonist antibody authorized or submitted for authorization, wherein the OX40 agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is tavolixizumab. The OX40 agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is tavolixizumab. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is tavolixizumab.

TABLE 12: Amino acid sequences for OX40 agonist antibodies related to tavolixizumab.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:56 heavy chain for tavolixizumab	QVQLQESGPG LVKPSQTLSL TCAVYGGSFS SGYWNWIRKH PGKGLEYIGY ISYNGITYHN PSLKSRITIN RDTSKNQYSL QLNSVTPEDT AVYYCARYKY DYDGGHAMDY WGQGTLVTVS SASTKGPSVF PLAPSSKSTS GGTAALGCLV KDYFPEPVTV SWNSGALTSG VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ TYICNVNHKP SNTKVDKRVE PKSCDKTHTC PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK ALPAPIEKTI SKAKGQPREP QVYTLPPSRE EMTKNQVSLT CLVKGFYPSD IAVEWESNGQ PENNYKTTPP VLDSDGSFFL YSKLTVDKSR WQQGNVFSCS VMHEALHNHY TQKSLSLSPG K	60 120 180 240 300 360 420 451
SEQ ID NO:57 light chain for tavolixizumab	DIQMTQSPSS LSASVGDRVT ITCRASQDIS NYLNWYQQKP GKAPKLLIYY TSKLHSGVPS RFSGSGSGTD YTLTISSLQP EDFATYYCQQ GSALPWTFGQ GTKVEIKRTV AAPSVFIFPP SDEQLKSGTA SVVCLLNNFY PREAKVQWKV DNALQSGNSQ ESVTEQDSKD STYSLSSTLT LSKADYEKHK VYACEVTHQG LSSPVTKSFN RGEC	60 120 180 214
SEQ ID NO:58 heavy chain variable region for tavolixizumab	QVQLQESGPG LVKPSQTLSL TCAVYGGSFS SGYWNWIRKH PGKGLEYIGY ISYNGITYHN PSLKSRITIN RDTSKNQYSL QLNSVTPEDT AVYYCARYKY DYDGGHAMDY WGQGTLVT	60 118
SEQ ID NO:59 light chain variable region for tavolixizumab	DIQMTQSPSS LSASVGDRVT ITCRASQDIS NYLNWYQQKP GKAPKLLIYY TSKLHSGVPS RFSGSGSGTD YTLTISSLQP EDFATYYCQQ GSALPWTFGQ GTKVEIKR	60 108

SEQ ID NO:60 heavy chain CDR1 for tavolixizumab	GSFSSGYWN	9
SEQ ID NO:61 heavy chain CDR2 for tavolixizumab	YIGYISYNGI TYH	13
SEQ ID NO:62 heavy chain CDR3 for tavolixizumab	RYKYDYDGGH AMDY	14
SEQ ID NO:63 light chain CDR1 for tavolixizumab	QDISNYLN	8
SEQ ID NO:64 light chain CDR2 for tavolixizumab	LLIYYTSKLH S	11
SEQ ID NO:65 light chain CDR3 for tavolixizumab	QQGSALPW	8

[00916] In some embodiments, the OX40 agonist is 11D4, which is a fully human antibody available from Pfizer, Inc. The preparation and properties of 11D4 are described in U.S. Patent Nos. 7,960,515; 8,236,930; and 9,028,824, the disclosures of which are incorporated by reference herein. The amino acid sequences of 11D4 are set forth in Table 13.

[00917] In an embodiment, a OX40 agonist comprises a heavy chain given by SEQ ID NO:66 and a light chain given by SEQ ID NO:67. In an embodiment, a OX40 agonist comprises heavy and light chains having the sequences shown in SEQ ID NO:66 and SEQ ID NO:67, respectively, or antigen binding fragments, Fab fragments, single-chain variable fragments (scFv), variants, or conjugates thereof. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 99% identical to the sequences shown in SEQ ID NO:66 and SEQ ID NO:67, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 98% identical to the sequences shown in SEQ ID NO:66 and SEQ ID NO:67, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 97% identical to the sequences shown in SEQ ID NO:66 and SEQ ID NO:67, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 96% identical to the sequences shown in SEQ ID NO:66 and SEQ ID NO:67, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 95% identical to the sequences shown in SEQ ID NO:66 and SEQ ID NO:67, respectively.

[00918] In an embodiment, the OX40 agonist comprises the heavy and light chain CDRs or variable regions (VRs) of 11D4. In an embodiment, the OX40 agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:68, and the OX40 agonist light

chain variable region (V_L) comprises the sequence shown in SEQ ID NO:69, and conservative amino acid substitutions thereof. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:68 and SEQ ID NO:69, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:68 and SEQ ID NO:69, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:68 and SEQ ID NO:69, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:68 and SEQ ID NO:69, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:68 and SEQ ID NO:69, respectively.

[00919] In an embodiment, a OX40 agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:70, SEQ ID NO:71, and SEQ ID NO:72, respectively, and conservative amino acid substitutions thereof, and light chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:73, SEQ ID NO:74, and SEQ ID NO:75, respectively, and conservative amino acid substitutions thereof.

[00920] In an embodiment, the OX40 agonist is a OX40 agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to 11D4. In an embodiment, the biosimilar monoclonal antibody comprises an OX40 antibody comprising an amino acid sequence which has at least 97% sequence identity, *e.g.*, 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 11D4. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a OX40 agonist antibody authorized or submitted for authorization, wherein the OX40 agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 11D4. The OX40 agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or

more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 11D4. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 11D4.

TABLE 13: Amino acid sequences for OX40 agonist antibodies related to 11D4.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:66 heavy chain for 11D4	EVQLVESGGG LVQPGGSLRL SCAASGFTFS SYSMNWVRQA PGKGLEWVSY ISSSSSTIDY ADSVKGRFTI SRDNAKNSLY LQMNSLRDED TAVYYCARES GWYLFDYWGQ GTLVTVSSAS TKGPSVFPLA PCSRSTSEST AALGCLVKDY FPEPVTVSWN SGALTSGVHT FPAVLQSSGL YSLSSVVTVP SSNFGTQTYT CNVDHKPSNT KVDKTVERKC CVECPPCPAP PVAGPSVFLF PPKPKDTLMI SRTPEVTCVV VDVSHEDPEV QFNWYVDGVE VHNAKTKPRE EQFNSTFRVV SVLTVVHQDW LNGKEYKCKV SNKGLPAPIE KTISKTKGQP REPQVYTLPP SREEMTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT TPPMLDSDGS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL SPGK	60 120 180 240 300 360 420 444
SEQ ID NO:67 light chain for 11D4	DIQMTQSPSS LSASVGDRVT ITCRASQGIS SWLAWYQQKP EKAPKSLIYA ASSLQSGVPS RFSGSGSGTD FTLTISSLQP EDFATYYCQQ YNSYPPTFGG GTKVEIKRTV AAPSVFIFPP SDEQLKSGTA SVVCLLNNFY PREAKVQWKV DNALQSGNSQ ESVTEQDSKD STYSLSSTLT LSKADYEKHK VYACEVTHQG LSSPVTKSFN RGEC	60 120 180 214
SEQ ID NO:68 heavy chain variable region for 11D4	EVQLVESGGG LVQPGGSLRL SCAASGFTFS SYSMNWVRQA PGKGLEWVSY ISSSSSTIDY ADSVKGRFTI SRDNAKNSLY LQMNSLRDED TAVYYCARES GWYLFDYWGQ GTLVTVSS	60 118
SEQ ID NO:69 light chain variable region for 11D4	DIQMTQSPSS LSASVGDRVT ITCRASQGIS SWLAWYQQKP EKAPKSLIYA ASSLQSGVPS RFSGSGSGTD FTLTISSLQP EDFATYYCQQ YNSYPPTFGG GTKVEIK	60 107
SEQ ID NO:70 heavy chain CDR1 for 11D4	SYSMN	5
SEQ ID NO:71 heavy chain CDR2 for 11D4	YISSSSSTID YADSVKG	17
SEQ ID NO:72 heavy chain CDR3 for 11D4	ESGWYLFDY	9
SEQ ID NO:73 light chain CDR1 for 11D4	RASQGISSWL A	11
SEQ ID NO:74 light chain CDR2 for 11D4	AASSLQS	7
SEQ ID NO:75 light chain CDR3 for 11D4	QQYNSYPPT	9

[00921] In some embodiments, the OX40 agonist is 18D8, which is a fully human antibody available from Pfizer, Inc. The preparation and properties of 18D8 are described in U.S. Patent Nos. 7,960,515; 8,236,930; and 9,028,824, the disclosures of which are incorporated by reference herein. The amino acid sequences of 18D8 are set forth in Table 14.

[00922] In an embodiment, a OX40 agonist comprises a heavy chain given by SEQ ID NO:76 and a light chain given by SEQ ID NO:77. In an embodiment, a OX40 agonist

comprises heavy and light chains having the sequences shown in SEQ ID NO:76 and SEQ ID NO:77, respectively, or antigen binding fragments, Fab fragments, single-chain variable fragments (scFv), variants, or conjugates thereof. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 99% identical to the sequences shown in SEQ ID NO:76 and SEQ ID NO:77, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 98% identical to the sequences shown in SEQ ID NO:76 and SEQ ID NO:77, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 97% identical to the sequences shown in SEQ ID NO:76 and SEQ ID NO:77, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 96% identical to the sequences shown in SEQ ID NO:76 and SEQ ID NO:77, respectively. In an embodiment, a OX40 agonist comprises heavy and light chains that are each at least 95% identical to the sequences shown in SEQ ID NO:76 and SEQ ID NO:77, respectively.

[00923] In an embodiment, the OX40 agonist comprises the heavy and light chain CDRs or variable regions (VRs) of 18D8. In an embodiment, the OX40 agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:78, and the OX40 agonist light chain variable region (V_L) comprises the sequence shown in SEQ ID NO:79, and conservative amino acid substitutions thereof. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:78 and SEQ ID NO:79, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:78 and SEQ ID NO:79, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:78 and SEQ ID NO:79, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:78 and SEQ ID NO:79, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:78 and SEQ ID NO:79, respectively.

[00924] In an embodiment, a OX40 agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:80, SEQ ID NO:81, and SEQ ID NO:82, respectively, and conservative amino acid substitutions thereof, and light chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:83, SEQ ID NO:84, and SEQ ID NO:85, respectively, and conservative amino acid substitutions thereof.

[00925] In an embodiment, the OX40 agonist is a OX40 agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to 18D8. In an embodiment, the biosimilar monoclonal antibody comprises an OX40 antibody comprising an amino acid sequence which has at least 97% sequence identity, *e.g.,* 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 18D8. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a OX40 agonist antibody authorized or submitted for authorization, wherein the OX40 agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 18D8. The OX40 agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 18D8. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is 18D8.

TABLE 14: Amino acid sequences for OX40 agonist antibodies related to 18D8.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:76 heavy chain for 18D8	EVQLVESGGG LVQPGRSLRL SCAASGFTFD DYAMHWVRQA PGKGLEWVSG ISWNSGSIGY ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TALYYCAKDQ STADYYFYYG MDVWGQGTTV TVSSASTKGP SVFPLAPCSR STSESTAALG CLVKDYFPEP VTVSWNSGAL TSGVHTFPAV LQSSGLYSLS SVVTVPSSNF GTQTYTCNVD HKPSNTKVDK TVERKCCVEC PPCPAPPVAG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVQFNW YVDGVEVHNA KTKPREEQFN STFRVVSVLT VVHQDWLNGK EYKCKVSNKG LPAPIEKTIS KTKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPM LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK	60 120 180 240 300 360 420 450
SEQ ID NO:77 light chain for 18D8	EIVVTQSPAT LSLSPGERAT LSCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA RFSGSGSGTD FTLTISSLEP EDFAVYYCQQ RSNWPTFGQG TKVEIKRTVA APSVFIFPPS DEQLKSGTAS VVCLLNNFYP REAKVQWKVD NALQSGNSQE SVTEQDSKDS TYSLSSTLTL SKADYEKHKV YACEVTHQGL SSPVTKSFNR GEC	60 120 180 213
SEQ ID NO:78 heavy chain variable region for 18D8	EVQLVESGGG LVQPGRSLRL SCAASGFTFD DYAMHWVRQA PGKGLEWVSG ISWNSGSIGY ADSVKGRFTI SRDNAKNSLY LQMNSLRAED TALYYCAKDQ STADYYFYYG MDVWGQGTTV TVSS	60 120 124
SEQ ID NO:79 light chain	EIVVTQSPAT LSLSPGERAT LSCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA RFSGSGSGTD FTLTISSLEP EDFAVYYCQQ RSNWPTFGQG TKVEIK	60 106

variable region for 18D8		
SEQ ID NO:80 heavy chain CDR1 for 18D8	DYAMH	5
SEQ ID NO:81 heavy chain CDR2 for 18D8	GISWNSGSIG YADSVKG	17
SEQ ID NO:82 heavy chain CDR3 for 18D8	DQSTADYYFY YGMDV	15
SEQ ID NO:83 light chain CDR1 for 18D8	RASQSVSSYL A	11
SEQ ID NO:84 light chain CDR2 for 18D8	DASNRAT	7
SEQ ID NO:85 light chain CDR3 for 18D8	QQRSNWPT	8

[00926] In some embodiments, the OX40 agonist is Hu119-122, which is a humanized antibody available from GlaxoSmithKline plc. The preparation and properties of Hu119-122 are described in U.S. Patent Nos. 9,006,399 and 9,163,085, and in International Patent Publication No. WO 2012/027328, the disclosures of which are incorporated by reference herein. The amino acid sequences of Hu119-122 are set forth in Table 15.

[00927] In an embodiment, the OX40 agonist comprises the heavy and light chain CDRs or variable regions (VRs) of Hu119-122. In an embodiment, the OX40 agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:86, and the OX40 agonist light chain variable region (V_L) comprises the sequence shown in SEQ ID NO:87, and conservative amino acid substitutions thereof. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:86 and SEQ ID NO:87, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:86 and SEQ ID NO:87, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:86 and SEQ ID NO:87, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:86 and SEQ ID NO:87, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:86 and SEQ ID NO:87, respectively.

[00928] In an embodiment, a OX40 agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:88, SEQ ID NO:89, and SEQ ID NO:90, respectively, and conservative amino acid substitutions thereof, and light chain

CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:91, SEQ ID NO:92, and SEQ ID NO:93, respectively, and conservative amino acid substitutions thereof.

[00929] In an embodiment, the OX40 agonist is a OX40 agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to Hu119-122. In an embodiment, the biosimilar monoclonal antibody comprises an OX40 antibody comprising an amino acid sequence which has at least 97% sequence identity, *e.g.,* 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu119-122. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a OX40 agonist antibody authorized or submitted for authorization, wherein the OX40 agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu119-122. The OX40 agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu119-122. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu119-122.

TABLE 15: Amino acid sequences for OX40 agonist antibodies related to Hu119-122.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:86 heavy chain variable region for Hu119-122	EVQLVESGGG LVQPGGSLRL SCAASEYEFP SHDMSWVRQA PGKGLELVAA INSDGGSTYY PDTMERRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARHY DDYYAWFAYW GQGTMVTVSS	60 120
SEQ ID NO:87 light chain variable region for Hu119-122	EIVLTQSPAT LSLSPGERAT LSCRASKSVS TSGYSYMHWY QQKPGQAPRL LIYLASNLES GVPARFSGSG SGTDFTLTIS SLEPEDFAVY YCQHSRELPL TFGGGTKVEI K	60 111
SEQ ID NO:88 heavy chain CDR1 for Hu119-122	SHDMS	5

SEQ ID NO:89 heavy chain CDR2 for Hu119-122	AINSDGGSTY YPDTMER	17
SEQ ID NO:90 heavy chain CDR3 for Hu119-122	HYDDYYAWFA Y	11
SEQ ID NO:91 light chain CDR1 for Hu119-122	RASKSVSTSG YSYMH	15
SEQ ID NO:92 light chain CDR2 for Hu119-122	LASNLES	7
SEQ ID NO:93 light chain CDR3 for Hu119-122	QHSRELPLT	9

[00930] In some embodiments, the OX40 agonist is Hu106-222, which is a humanized antibody available from GlaxoSmithKline plc. The preparation and properties of Hu106-222 are described in U.S. Patent Nos. 9,006,399 and 9,163,085, and in International Patent Publication No. WO 2012/027328, the disclosures of which are incorporated by reference herein. The amino acid sequences of Hu106-222 are set forth in Table 16.

[00931] In an embodiment, the OX40 agonist comprises the heavy and light chain CDRs or variable regions (VRs) of Hu106-222. In an embodiment, the OX40 agonist heavy chain variable region (V_H) comprises the sequence shown in SEQ ID NO:94, and the OX40 agonist light chain variable region (V_L) comprises the sequence shown in SEQ ID NO:95, and conservative amino acid substitutions thereof. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 99% identical to the sequences shown in SEQ ID NO:94 and SEQ ID NO:95, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 98% identical to the sequences shown in SEQ ID NO:94 and SEQ ID NO:95, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 97% identical to the sequences shown in SEQ ID NO:94 and SEQ ID NO:95, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 96% identical to the sequences shown in SEQ ID NO:94 and SEQ ID NO:95, respectively. In an embodiment, a OX40 agonist comprises V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:94 and SEQ ID NO:95, respectively.

[00932] In an embodiment, a OX40 agonist comprises heavy chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:96, SEQ ID NO:97, and SEQ ID NO:98, respectively, and conservative amino acid substitutions thereof, and light chain CDR1, CDR2 and CDR3 domains having the sequences set forth in SEQ ID NO:99, SEQ ID NO:100, and SEQ ID NO:101, respectively, and conservative amino acid substitutions thereof.

[00933] In an embodiment, the OX40 agonist is a OX40 agonist biosimilar monoclonal antibody approved by drug regulatory authorities with reference to Hu106-222. In an embodiment, the biosimilar monoclonal antibody comprises an OX40 antibody comprising an amino acid sequence which has at least 97% sequence identity, *e.g.*, 97%, 98%, 99% or 100% sequence identity, to the amino acid sequence of a reference medicinal product or reference biological product and which comprises one or more post-translational modifications as compared to the reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu106-222. In some embodiments, the one or more post-translational modifications are selected from one or more of: glycosylation, oxidation, deamidation, and truncation. In some embodiments, the biosimilar is a OX40 agonist antibody authorized or submitted for authorization, wherein the OX40 agonist antibody is provided in a formulation which differs from the formulations of a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu106-222. The OX40 agonist antibody may be authorized by a drug regulatory authority such as the U.S. FDA and/or the European Union's EMA. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu106-222. In some embodiments, the biosimilar is provided as a composition which further comprises one or more excipients, wherein the one or more excipients are the same or different to the excipients comprised in a reference medicinal product or reference biological product, wherein the reference medicinal product or reference biological product is Hu106-222.

TABLE 16: Amino acid sequences for OX40 agonist antibodies related to Hu106-222.

Identifier	Sequence (One-Letter Amino Acid Symbols)	
SEQ ID NO:94 heavy chain variable region for Hu106-222	QVQLVQSGSE LKKPGASVKV SCKASGYTFT DYSMHWVRQA PGQGLKWMGW INTETGEPTY ADDFKGRFVF SLDTSVSTAY LQISSLKAED TAVYYCANPY YDYVSYYAMD YWGQGTTVTV SS	60 120 122
SEQ ID NO:95 light chain variable region for Hu106-222	DIQMTQSPSS LSASVGDRVT ITCKASQDVS TAVAWYQQKP GKAPKLLIYS ASYLYTGVPS RFSGSGSGTD FTFTISSLQP EDIATYYCQQ HYSTPRTFGQ GTKLEIK	60 107
SEQ ID NO:96 heavy chain CDR1 for Hu106-222	DYSMH	5
SEQ ID NO:97 heavy chain CDR2 for Hu106-222	WINTETGEPT YADDFKG	17

SEQ ID NO:98 heavy chain CDR3 for Hu106-222	PYYDYVSYYA MDY	13
SEQ ID NO:99 light chain CDR1 for Hu106-222	KASQDVSTAV A	11
SEQ ID NO:100 light chain CDR2 for Hu106-222	SASYLYT	7
SEQ ID NO:101 light chain CDR3 for Hu106-222	QQHYSTPRT	9

[00934] In some embodiments, the OX40 agonist antibody is MEDI6469 (also referred to as 9B12). MEDI6469 is a murine monoclonal antibody. Weinberg, *et al.*, *J. Immunother*. **2006**, *29*, 575-585. In some embodiments the OX40 agonist is an antibody produced by the 9B12 hybridoma, deposited with Biovest Inc. (Malvern, MA, USA), as described in Weinberg, *et al.*, *J. Immunother*. **2006**, *29*, 575-585, the disclosure of which is hereby incorporated by reference in its entirety. In some embodiments, the antibody comprises the CDR sequences of MEDI6469. In some embodiments, the antibody comprises a heavy chain variable region sequence and/or a light chain variable region sequence of MEDI6469.

[00935] In an embodiment, the OX40 agonist is L106 BD (Pharmingen Product #340420). In some embodiments, the OX40 agonist comprises the CDRs of antibody L106 (BD Pharmingen Product #340420). In some embodiments, the OX40 agonist comprises a heavy chain variable region sequence and/or a light chain variable region sequence of antibody L106 (BD Pharmingen Product #340420). In an embodiment, the OX40 agonist is ACT35 (Santa Cruz Biotechnology, Catalog #20073). In some embodiments, the OX40 agonist comprises the CDRs of antibody ACT35 (Santa Cruz Biotechnology, Catalog #20073). In some embodiments, the OX40 agonist comprises a heavy chain variable region sequence and/or a light chain variable region sequence of antibody ACT35 (Santa Cruz Biotechnology, Catalog #20073). In an embodiment, the OX40 agonist is the murine monoclonal antibody anti-mCD134/mOX40 (clone OX86), commercially available from InVivoMAb, BioXcell Inc, West Lebanon, NH.

[00936] In an embodiment, the OX40 agonist is selected from the OX40 agonists described in International Patent Application Publication Nos. WO 95/12673, WO 95/21925, WO 2006/121810, WO 2012/027328, WO 2013/028231, WO 2013/038191, and WO 2014/148895; European Patent Application EP 0672141; U.S. Patent Application Publication Nos. US 2010/136030, US 2014/377284, US 2015/190506, and US 2015/132288 (including clones 20E5 and 12H3); and U.S. Patent Nos. 7,504,101, 7,550,140, 7,622,444, 7,696,175,

7,960,515, 7,961,515, 8,133,983, 9,006,399, and 9,163,085, the disclosure of each of which is incorporated herein by reference in its entirety.

[00937] In an embodiment, the OX40 agonist is an OX40 agonistic fusion protein as depicted in Structure I-A (C-terminal Fc-antibody fragment fusion protein) or Structure I-B (N-terminal Fc-antibody fragment fusion protein), or a fragment, derivative, conjugate, variant, or biosimilar thereof. The properties of structures I-A and I-B are described above and in U.S. Patent Nos. 9,359,420, 9,340,599, 8,921,519, and 8,450,460, the disclosures of which are incorporated by reference herein. Amino acid sequences for the polypeptide domains of structure I-A are given in Table GG. The Fc domain preferably comprises a complete constant domain (amino acids 17-230 of SEQ ID NO:31) the complete hinge domain (amino acids 1-16 of SEQ ID NO:31) or a portion of the hinge domain (*e.g.*, amino acids 4-16 of SEQ ID NO:31). Preferred linkers for connecting a C-terminal Fc-antibody may be selected from the embodiments given in SEQ ID NO:32 to SEQ ID NO:41, including linkers suitable for fusion of additional polypeptides. Likewise, amino acid sequences for the polypeptide domains of structure I-B are given in Table 9. If an Fc antibody fragment is fused to the N-terminus of an TNRFSF fusion protein as in structure I-B, the sequence of the Fc module is preferably that shown in SEQ ID NO:42, and the linker sequences are preferably selected from those embodiments set forth in SED ID NO:43 to SEQ ID NO:45.

[00938] In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains selected from the group consisting of a variable heavy chain and variable light chain of tavolixizumab, a variable heavy chain and variable light chain of 11D4, a variable heavy chain and variable light chain of 18D8, a variable heavy chain and variable light chain of Hu119-122, a variable heavy chain and variable light chain of Hu106-222, a variable heavy chain and variable light chain selected from the variable heavy chains and variable light chains described in Table OO, any combination of a variable heavy chain and variable light chain of the foregoing, and fragments, derivatives, conjugates, variants, and biosimilars thereof.

[00939] In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains comprising an OX40L sequence. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains comprising a sequence according to SEQ ID NO:102. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains comprising a soluble OX40L sequence. In an

embodiment, a OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains comprising a sequence according to SEQ ID NO:103. In an embodiment, a OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains comprising a sequence according to SEQ ID NO:104.

[00940] In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:58 and SEQ ID NO:59, respectively, wherein the V_H and V_L domains are connected by a linker. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:68 and SEQ ID NO:69, respectively, wherein the V_H and V_L domains are connected by a linker. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:78 and SEQ ID NO:79, respectively, wherein the V_H and V_L domains are connected by a linker. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:86 and SEQ ID NO:87, respectively, wherein the V_H and V_L domains are connected by a linker. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the sequences shown in SEQ ID NO:94 and SEQ ID NO:95, respectively, wherein the V_H and V_L domains are connected by a linker. In an embodiment, an OX40 agonist fusion protein according to structures I-A or I-B comprises one or more OX40 binding domains that is a scFv domain comprising V_H and V_L regions that are each at least 95% identical to the V_H and V_L sequences given in Table 17, wherein the V_H and V_L domains are connected by a linker.

TABLE 17: Additional polypeptide domains useful as OX40 binding domains in fusion proteins (*e.g.*, structures I-A and I-B) or as scFv OX40 agonist antibodies.

Identifier	Sequence (One-Letter Amino Acid Symbols)
SEQ ID NO:102 OX40L	MERVQPLEEN VGNAARPRFE RNKLLLVASV IQGLGLLLCF TYICLHFSAL QVSHRYPRIQ 60 SIKVQFTEYK KEKGFILTSQ KEDEIMKVQN NSVIINCDGF YLISLKGYFS QEVNISLHYQ 120 KDEEPLFQLK KVRSVNSLMV ASLTYKDKVY LNVTTDNTSL DDFHVNGGEL ILIHQNPGEF 180 CVL 183

SEQ ID NO:103 OX40L soluble domain	SHRYPRIQSI KVQFTEYKKE KGFILTSQKE DEIMKVQNNS VIINCDGFYL ISLKGYFSQE VNISLHYQKD EEPLFQLKKV RSVNSLMVAS LTYKDKVYLN VTTDNTSLDD FHVNGGELIL IHQNPGEFCV L	60 120 131
SEQ ID NO:104 OX40L soluble domain (alternative)	YPRIQSIKVQ FTEYKKEKGF ILTSQKEDEI MKVQNNSVII NCDGFYLISL KGYFSQEVNI SLHYQKDEEP LFQLKKVRSV NSLMVASLTY KDKVYLNVTT DNTSLDDFHV NGGELILIHQ NPGEFCVL	60 120 128
SEQ ID NO:105 variable heavy chain for 008	EVQLVESGGG LVQPGGSLRL SCAASGFTFS NYTMNWVRQA PGKGLEWVSA ISGSGGSTYY ADSVKGRFTI SRDNSKNTLY LQMNSLRAED TAVYYCAKDR YSQVHYALDY WGQGTLVTVS	60 120
SEQ ID NO:106 variable light chain for 008	DIVMTQSPDS LPVTPGEPAS ISCRSSQSLL HSNGYNYLDW YLQKAGQSPQ LLIYLGSNRA SGVPDRFSGS GSGTDFTLKI SRVEAEDVGV YYCQQYYNHP TTFGQGTK	60 108
SEQ ID NO:107 variable heavy chain for 011	EVQLVESGGG VVQPGRSLRL SCAASGFTFS DYTMNWVRQA PGKGLEWVSS ISGGSTYYAD SRKGRFTISR DNSKNTLYLQ MNNLRAEDTA VYYCARDRYF RQQNAFDYWG QGTLVTVSSA	60 120
SEQ ID NO:108 variable light chain for 011	DIVMTQSPDS LPVTPGEPAS ISCRSSQSLL HSNGYNYLDW YLQKAGQSPQ LLIYLGSNRA SGVPDRFSGS GSGTDFTLKI SRVEAEDVGV YYCQQYYNHP TTFGQGTK	60 108
SEQ ID NO:109 variable heavy chain for 021	EVQLVESGGG LVQPRGSLRL SCAASGFTFS SYAMNWVRQA PGKGLEWVAV ISYDGSNKYY ADSVKGRFTI SRDNSKNTLY LQMNSLRAED TAVYYCAKDR YITLPNALDY WGQGTLVTVS	60 120
SEQ ID NO:110 variable light chain for 021	DIQMTQSPVS LPVTPGEPAS ISCRSSQSLL HSNGYNYLDW YLQKPGQSPQ LLIYLGSNRA SGVPDRFSGS GSGTDFTLKI SRVEAEDVGV YYCQQYKSNP PTFGQGTK	60 108
SEQ ID NO:111 variable heavy chain for 023	EVQLVESGGG LVHPGGSLRL SCAGSGFTFS SYAMHWVRQA PGKGLEWVSA IGTGGGTYYA DSVMGRFTIS RDNSKNTLYL QMNSLRAEDT AVYYCARYDN VMGLYWFDYW GQGTLVTVSS	60 120
SEQ ID NO:112 variable light chain for 023	EIVLTQSPAT LSLSPGERAT LSCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA RFSGSGSGTD FTLTISSLEP EDFAVYYCQQ RSNWPPAFGG GTKVEIKR	60 108
SEQ ID NO:113 heavy chain variable region	EVQLQQSGPE LVKPGASVKM SCKASGYTFT SYVMHWVKQK PGQGLEWIGY INPYNDGTKY NEKFKGKATL TSDKSSSTAY MELSSLTSED SAVYYCANYY GSSLSMDYWG QGTSVTVSS	60 119
SEQ ID NO:114 light chain variable region	DIQMTQTTSS LSASLGDRVT ISCRASQDIS NYLNWYQQKP DGTVKLLIYY TSRLHSGVPS RFSGSGSGTD YSLTISNLEQ EDIATYFCQQ GNTLPWTFGG GTKLEIKR	60 108
SEQ ID NO:115 heavy chain variable region	EVQLQQSGPE LVKPGASVKI SCKTSGYTFK DYTMHWVKQS HGKSLEWIGG IYPNNGGSTY NQNFKDKATL TVDKSSSTAY MEFRSLTSED SAVYYCARMG YHGPHLDFDV WGAGTTVTVS P	60 120 121
SEQ ID NO:116 light chain variable region	DIVMTQSHKF MSTSLGDRVS ITCKASQDVG AAVAWYQQKP GQSPKLLIYW ASTRHTGVPD RFTGGGSGTD FTLTISNVQS EDLTDYFCQQ YINYPLTFGG GTKLEIKR	60 108
SEQ ID NO:117 heavy chain variable region of humanized antibody	QIQLVQSGPE LKKPGETVKI SCKASGYTFT DYSMHWVKQA PGKGLKWMGW INTETGEPTY ADDFKGRFAF SLETSASTAY LQINNLKNED TATYFCANPY YDYVSYYAMD YWGHGTSVTV SS	60 120 122
SEQ ID NO:118 heavy chain variable region of humanized antibody	QVQLVQSGSE LKKPGASVKV SCKASGYTFT DYSMHWVRQA PGQGLKWMGW INTETGEPTY ADDFKGRFVF SLDTSVSTAY LQISSLKAED TAVYYCANPY YDYVSYYAMD YWGQGTTVTV SS	60 120 122
SEQ ID NO:119 light chain variable region of humanized antibody	DIVMTQSHKF MSTSVRDRVS ITCKASQDVS TAVAWYQQKP GQSPKLLIYS ASYLYTGVPD RFTGSGSGTD FTFTISSVQA EDLAVYYCQQ HYSTPRTFGG GTKLEIK	60 107
SEQ ID NO:120 light chain variable region of humanized antibody	DIVMTQSHKF MSTSVRDRVS ITCKASQDVS TAVAWYQQKP GQSPKLLIYS ASYLYTGVPD RFTGSGSGTD FTFTISSVQA EDLAVYYCQQ HYSTPRTFGG GTKLEIK	60 107
SEQ ID NO:121 heavy chain variable region of humanized antibody	EVQLVESGGG LVQPGESLKL SCESNEYEFP SHDMSWVRKT PEKRLELVAA INSDGGSTYY PDTMERRFII SRDNTKKTLY LQMSSLRSED TALYYCARHY DDYYAWFAYW GQGTLVTVSA	60 120
SEQ ID NO:122 heavy chain variable region of humanized antibody	EVQLVESGGG LVQPGGSLRL SCAASEYEFP SHDMSWVRQA PGKGLELVAA INSDGGSTYY PDTMERRFTI SRDNAKNSLY LQMNSLRAED TAVYYCARHY DDYYAWFAYW GQGTMVTVSS	60 120
SEQ ID NO:123 light chain	DIVLTQSPAS LAVSLGQRAT ISCRASKSVS TSGYSYMHWY QQKPGQPPKL LIYLASNLES GVPARFSGSG SGTDFTLNIH PVEEEDAATY YCQHSRELPL TFGAGTKLEL K	60 111

variable region of humanized antibody		
SEQ ID NO:124 light chain variable region of humanized antibody	EIVLTQSPAT LSLSPGERAT LSCRASKSVS TSGYSYMHWY QQKPGQAPRL LIYLASNLES GVPARFSGSG SGTDFTLTIS SLEPEDFAVY YCQHSRELPL TFGGGTKVEI K	60 111
SEQ ID NO:125 heavy chain variable region	MYLGLNYVFI VFLLNGVQSE VKLEESGGGL VQPGGSMKLS CAASGFTFSD AWMDWVRQSP EKGLEWVAEI RSKANNHATY YAESVNGRFT ISRDDSKSSV YLQMNSLRAE DTGIYYCTWG EVFYFDYWGQ GTTLTVSS	60 120 138
SEQ ID NO:126 light chain variable region	MRPSIQFLGL LLFWLHGAQC DIQMTQSPSS LSASLGGKVT ITCKSSQDIN KYIAWYQHKP GKGPRLLIHY TSTLQPGIPS RFSGSGSGRD YSFSISNLEP EDIATYYCLQ YDNLLTFGAG TKLELK	60 120 126

[00941] In an embodiment, the OX40 agonist is a OX40 agonistic single-chain fusion polypeptide comprising (i) a first soluble OX40 binding domain, (ii) a first peptide linker, (iii) a second soluble OX40 binding domain, (iv) a second peptide linker, and (v) a third soluble OX40 binding domain, further comprising an additional domain at the N-terminal and/or C-terminal end, and wherein the additional domain is a Fab or Fc fragment domain. In an embodiment, the OX40 agonist is a OX40 agonistic single-chain fusion polypeptide comprising (i) a first soluble OX40 binding domain, (ii) a first peptide linker, (iii) a second soluble OX40 binding domain, (iv) a second peptide linker, and (v) a third soluble OX40 binding domain, further comprising an additional domain at the N-terminal and/or C-terminal end, wherein the additional domain is a Fab or Fc fragment domain wherein each of the soluble OX40 binding domains lacks a stalk region (which contributes to trimerisation and provides a certain distance to the cell membrane, but is not part of the OX40 binding domain) and the first and the second peptide linkers independently have a length of 3-8 amino acids.

[00942] In an embodiment, the OX40 agonist is an OX40 agonistic single-chain fusion polypeptide comprising (i) a first soluble tumor necrosis factor (TNF) superfamily cytokine domain, (ii) a first peptide linker, (iii) a second soluble TNF superfamily cytokine domain, (iv) a second peptide linker, and (v) a third soluble TNF superfamily cytokine domain, wherein each of the soluble TNF superfamily cytokine domains lacks a stalk region and the first and the second peptide linkers independently have a length of 3-8 amino acids, and wherein the TNF superfamily cytokine domain is an OX40 binding domain.

[00943] In some embodiments, the OX40 agonist is MEDI6383. MEDI6383 is an OX40 agonistic fusion protein and can be prepared as described in U.S. Patent No. 6,312,700, the disclosure of which is incorporated by reference herein.

[00944] In an embodiment, the OX40 agonist is an OX40 agonistic scFv antibody comprising any of the foregoing V_H domains linked to any of the foregoing V_L domains.

[00945] In an embodiment, the OX40 agonist is Creative Biolabs OX40 agonist monoclonal antibody MOM-18455, commercially available from Creative Biolabs, Inc., Shirley, NY, USA.

[00946] In an embodiment, the OX40 agonist is OX40 agonistic antibody clone Ber-ACT35 commercially available from BioLegend, Inc., San Diego, CA, USA.

VI. Optional Cell Viability Analyses

[00947] Optionally, a cell viability assay can be performed after the Step B first expansion of Process 2A (Gen 2) or Gen 3 processes, using standard assays known in the art. For example, a trypan blue exclusion assay can be done on a sample of the bulk TILs, which selectively labels dead cells and allows a viability assessment. Other assays for use in testing viability can include but are not limited to the Alamar blue assay; and the MTT assay.

1. Cell Counts, Viability, Flow Cytometry

[00948] In some embodiments of Process 2A (Gen 2) or Gen 3, cell counts and/or viability are measured. The expression of markers such as but not limited CD3, CD4, CD8, and CD56, as well as any other disclosed or described herein, can be measured by flow cytometry with antibodies, for example but not limited to those commercially available from BD Biosciences (BD Biosciences, San Jose, CA) using a FACSCanto™ flow cytometer (BD Biosciences). The cells can be counted manually using a disposable c-chip hemocytometer (VWR, Batavia, IL) and viability can be assessed using any method known in the art, including but not limited to trypan blue staining. In some embodiments of Process 2A (Gen 2) and/or Gen 3, the TILs exhibit an increase in CD8+ cells after either the first expansion, the second expansion, the priming first expansion, or the rapid second expansion. In some embodiments of Process 2A (Gen 2), the TILs exhibit an increase in CD8+ cells after either the first expansion and/or the second expansion. In some embodiments of Gen 3, the TILs exhibit an increase in CD8+ cells after the priming first expansion, or the rapid second expansion.

[00949] In some cases, the bulk TIL population can be cryopreserved immediately, using the protocols discussed below. Alternatively, the bulk TIL population can be subjected to REP and then cryopreserved as discussed below. Similarly, in the case where genetically modified

TILs will be used in therapy, the bulk or REP TIL populations can be subjected to genetic modifications for suitable treatments.

2. Cell Cultures

[00950] In an embodiment, a method for expanding TILs may include using about 5,000 mL to about 25,000 mL of cell medium, about 5,000 mL to about 10,000 mL of cell medium, or about 5,800 mL to about 8,700 mL of cell medium. In an embodiment, expanding the number of TILs uses no more than one type of cell culture medium. Any suitable cell culture medium may be used, *e.g.*, AIM-V cell medium (L-glutamine, 50 μM streptomycin sulfate, and 10 μM gentamicin sulfate) cell culture medium (Invitrogen, Carlsbad CA). In this regard, the inventive methods advantageously reduce the amount of medium and the number of types of medium required to expand the number of TIL. In an embodiment, expanding the number of TIL may comprise adding fresh cell culture media to the cells (also referred to as feeding the cells) no more frequently than every third or fourth day. Expanding the number of cells in a gas permeable container simplifies the procedures necessary to expand the number of cells by reducing the feeding frequency necessary to expand the cells.

[00951] In an embodiment, the cell medium in the first and/or second gas permeable container is unfiltered. The use of unfiltered cell medium may simplify the procedures necessary to expand the number of cells. In an embodiment, the cell medium in the first and/or second gas permeable container lacks beta-mercaptoethanol (BME).

[00952] In an embodiment, TILs are expanded in gas-permeable containers. Gas-permeable containers have been used to expand TILs using PBMCs using methods, compositions, and devices known in the art, including those described in U.S. Patent Application Publication No. 2005/0106717 A1, the disclosures of which are incorporated herein by reference. In an embodiment, TILs are expanded in gas-permeable bags. In an embodiment, TILs are expanded using a cell expansion system that expands TILs in gas permeable bags, such as the Xuri Cell Expansion System W25 (GE Healthcare). In an embodiment, TILs are expanded using a cell expansion system that expands TILs in gas permeable bags, such as the WAVE Bioreactor System, also known as the Xuri Cell Expansion System W5 (GE Healthcare). In an embodiment, the cell expansion system includes a gas permeable cell bag with a volume selected from the group consisting of about 100 mL, about 200 mL, about 300 mL, about 400 mL, about 500 mL, about 600 mL, about 700 mL, about 800 mL, about 900 mL, about 1 L, about 2 L, about 3 L, about 4 L, about 5 L, about 6 L, about 7 L, about 8 L, about 9 L, and

about 10 L. In an embodiment, TILs can be expanded in G-Rex flasks (commercially available from Wilson Wolf Manufacturing). Such embodiments allow for cell populations to expand from about 5×10^5 cells/cm^2 to between 10×10^6 and 30×10^6 cells/cm^2. In an embodiment this expansion is conducted without adding fresh cell culture media to the cells (also referred to as feeding the cells). In an embodiment, this is without feeding so long as medium resides at a height of about 10 cm in the GRex flask. In an embodiment this is without feeding but with the addition of one or more cytokines. In an embodiment, the cytokine can be added as a bolus without any need to mix the cytokine with the medium. Such containers, devices, and methods are known in the art and have been used to expand TILs, and include those described in U.S. Patent Application Publication No. US 2014/0377739A1, International Publication No. WO 2014/210036 A1, U.S. Patent Application Publication No. us 2013/0115617 A1, International Publication No. WO 2013/188427 A1, U.S. Patent Application Publication No. US 2011/0136228 A1, U.S. Patent No. US 8,809,050 B2, International publication No. WO 2011/072088 A2, U.S. Patent Application Publication No. US 2016/0208216 A1, U.S. Patent Application Publication No. US 2012/0244133 A1, International Publication No. WO 2012/129201 A1, U.S. Patent Application Publication No. US 2013/0102075 A1, U.S. Patent No. US 8,956,860 B2, International Publication No. WO 2013/173835 A1, U.S. Patent Application Publication No. US 2015/0175966 A1, the disclosures of which are incorporated herein by reference. Such processes are also described in Jin *et al.*, *J. Immunotherapy,* **2012**, 35:283-292.

VII. Metabolic Health of Expanded TILs

[00953] Spare respiratory capacity (SRC) and glycolytic reserve can be evaluated for TILs expanded with the methods disclosed herein. The Seahorse XF Cell Mito Stress Test measures mitochondrial function by directly measuring the oxygen consumption rate (OCR) of cells, using modulators of respiration that target components of the electron transport chain in the mitochondria. The test compounds (oligomycin, FCCP, and a mix of rotenone and antimycin A, described below) are serially injected to measure ATP production, maximal respiration, and non-mitochondrial respiration, respectively. Proton leak and spare respiratory capacity are then calculated using these parameters and basal respiration. Each modulator targets a specific component of the electron transport chain. Oligomycin inhibits ATP synthase (complex V) and the decrease in OCR following injection of oligomycin correlates to the mitochondrial respiration associated with cellular ATP production.

Carbonyl cyanide-4 (trifluoromethoxy) phenylhydrazone (FCCP) is an uncoupling agent that collapses the proton gradient and disrupts the mitochondrial membrane potential. As a result, electron flow through the electron transport chain is uninhibited and oxygen is maximally consumed by complex IV. The FCCP-stimulated OCR can then be used to calculate spare respiratory capacity, defined as the difference between maximal respiration and basal respiration. Spare respiratory capacity (SRC) is a measure of the ability of the cell to respond to increased energy demand. The third injection is a mix of rotenone, a complex I inhibitor, and antimycin A, a complex III inhibitor. This combination shuts down mitochondrial respiration and enables the calculation of nonmitochondrial respiration driven by processes outside the mitochondria

[00954] In some embodiments, the metabolic assay is basal respiration.

[00955] In general, TILs have a basal respiration rate that is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the basal respiration rate is from about 50% to about 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the basal respiration rate is from about 60% to about 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the basal respiration rate is from about 70% to about 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the basal respiration rate is from about 80% to about 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the basal respiration rate is from about 90% to about 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the basal respiration rate is from about 95% to about 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the TILs have a basal respiration rate that is not statistically significantly different than the basal respiration rate of freshly harvested TILs.

[00956] In some embodiments, the metabolic assay is spare respiratory capacity.

[00957] In some embodiments, the metabolic assay is glycolytic reserve. In some embodiments, the metabolic assay is spare respiratory capacity. To measure cellular (respiratory) metabolism cells were treated with inhibitors of mitochondrial respiration and glycolysis to determine a metabolic profile for the TIL consisting of the following measures: baseline oxidative phosphorylation (as measured by OCR), spare respiratory capacity, baseline glycolytic activity (as measured by ECAR), and glycolytic reserve. Metabolic

profiles were performed using the Seahorse Combination Mitochondrial/Glycolysis Stress Test Assay (including the kit commercially available from Agilent®), which allows for determining a cells' capacity to perform glycolysis upon blockage of mitochondrial ATP production. In some embodiments, cells are starved of glucose, then glucose is injected, followed by a stress agent. In some embodiments, the stress agent is selected from the group consisting of oligomycin, FCCP, rotenone, antimycin A and/or 2-deoxyglucose (2-DG), as well as combinations thereof. In some embodiments, oligomycin is added at 10 mM. In some embodiments, FCCP is added at 10 mM. In some embodiments, rotenone is added at 2.5 mM. In some embodiments, antimycin A is added at 2.5 mM. In some embodiments, 2-deoxyglucose (2-DG) is added at 500 mM. In some embodiments, glycolytic capacity, glycolytic reserve, and/or non-glycolytic acidification are measured. In general, TILs have a glycolytic reserve that is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 50% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 60% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 70% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 80% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 90% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 95% to about 99% of the *basal respiration rate* of freshly harvested TILs.

[00958] In some embodiments, the metabolic assay is basal glycolysis. In some the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of at least two-fold, at least three-fold, at least four-fold, at least five-fold, at least six-fold, at least 7-fold, at least eight-fold, at least nine-fold, or at least ten-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about ten-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs)

have an increase in basal glycolysis of about two-fold to about eight-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about three-fold to about seven-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about four-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about three-fold.

[00959] In general, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a spare respiratory capacity that is at least is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 50% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 50% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 60% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 70% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 80% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 90% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 95% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a spare respiratory capacity that is not statistically significantly different than the *basal respiration rate* of freshly harvested TILs.

VIII. FUNCTIONAL CHARACTERIZATION OF TILs

[00960] Granzyme B Production: Granzyme B is another measure of the ability of TIL to kill target cells. Media supernatants restimulated as described above using antibodies to CD3, CD28, and CD137/4-1BB were also evaluated for their levels of Granzyme B using the Human Granzyme B DuoSet ELISA Kit (R & D Systems, Minneapolis, MN) according to the manufacturer's instructions. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have increased Granzyme B production. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have increased cytotoxic activity.

[00961] In some embodiments, telomere length can be used as a measure of cell viability and/or cellular function. Telomere length measurement: Diverse methods have been used to measure the length of telomeres in genomic DNA and cytological preparations. The telomere restriction fragment (TRF) analysis is the gold standard to measure telomere length (de Lange *et al.*, 1990). However, the major limitation of TRF is the requirement of a large amount of DNA (1.5 μg). Two widely used techniques for the measurement of telomere lengths namely, fluorescence in situ hybridization (FISH; Agilent Technologies, Santa Clara, CA) and quantitative PCR can be employed with the present invention. In some embodiments, there is no change in telomere length between the initially harvest TILs in Step A and the expanded TILs from for example Step D as provided in Figure 7.

[00962] In some embodiments, TIL health is measured by IFN-gamma secretion. In some embodiments, a potency assay for IFN-γ production is employed. IFN-γ production is another measure of cytotoxic potential. IFN-γ production can be measured by determining the levels of the cytokine IFN-γ in the media of TIL stimulated with antibodies to CD3, CD28, and CD137/4-1BB. IFN-γ levels in media from these stimulated TIL can be determined using by measuring IFN-γ release. In some embodiments, an increase in IFN-γ production in for example Step D as provided in Figure 7 TILs as compared to initially harvested TILs in for example Step A as provided in Figure 7 is indicative of an increase in cytotoxic potential of the Step D TILs.

[00963] In some embodiments, the Bioluminescent Redirected Lysis Assay: The cytotoxic potential of TIL to lyse target cells was assessed using a co-culture assay of TIL with the bioluminescent cell line, P815 (Clone G6), according to a Bioluminescent redirected lysis assay (potency assay) for TIL assay which measures TIL cytotoxicity in a highly sensitive dose dependent manner.

[00964] In some embodiments, the present methods include an assay for assessing TIL viability, using the methods as described above and herein, including, for example, as described in the Examples and Figures. In some embodiments, the TILs are expanded as discussed above, including for example as provided in Figure 7. In some embodiments, the TILs are cryopreserved prior to being assessed for viability. In some embodiments, the viability assessment includes thawing the TILs prior to performing a first expansion, a second expansion, and an additional second expansion. In some embodiments, the present methods provide an assay for assessing cell proliferation, cell toxicity, cell death, and/or other terms related to viability of the TIL population. Viability can be measured by any of the TIL metabolic assays described above as well as any methods know for assessing cell viability that are known in the art. In some embodiments, the present methods provide as assay for assessment of cell proliferation, cell toxicity, cell death, and/or other terms related to viability of the TILs expanded using the methods described herein, including those exemplified in Figure 7.

[00965] In some embodiments, the metabolic assay is basal respiration. In general, second expansion TILs have a basal respiration rate that is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the basal respiration rate is from about 50% to about 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the basal respiration rate is from about 60% to about 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the basal respiration rate is from about 70% to about 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the basal respiration rate is from about 80% to about 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the basal respiration rate is from about 90% to about 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the basal respiration rate is from about 95% to about 99% of the basal

respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a basal respiration rate that is not statistically significantly different than the basal respiration rate of 2A process TILs (see, for example Figure 7).

[00966] In general, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a spare respiratory capacity that is at least is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the spare respiratory capacity is from about 50% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the spare respiratory capacity is from about 50% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the spare respiratory capacity is from about 60% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the spare respiratory capacity is from about 70% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the spare respiratory capacity is from about 80% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the spare respiratory capacity is from about 90% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the spare respiratory capacity is from about 95% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a spare respiratory capacity that is not statistically significantly different than the *basal respiration rate* of 2A process TILs (see, for example Figure 7).

[00967] In general, second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a spare respiratory capacity that is at least is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the metabolic assay measured is

glycolytic reserve. In some embodiments, the metabolic assay is spare respiratory capacity. To measure cellular (respiratory) metabolism cells were treated with inhibitors of mitochondrial respiration and glycolysis to determine a metabolic profile for the TIL consisting of the following measures: baseline oxidative phosphorylation (as measured by OCR), spare respiratory capacity, baseline glycolytic activity (as measured by ECAR), and glycolytic reserve. Metabolic profiles were performed using the Seahorse Combination Mitochondrial/Glycolysis Stress Test Assay (including the kit commercially available from Agilent®), which allows for determining a cells' capacity to perform glycolysis upon blockage of mitochondrial ATP production. In some embodiments, cells are starved of glucose, then glucose is injected, followed by a stress agent. In some embodiments, the stress agent is selected from the group consisting of oligomycin, FCCP, rotenone, antimycin A and/or 2-deoxyglucose (2-DG), as well as combinations thereof. In some embodiments, oligomycin is added at 10 mM. In some embodiments, FCCP is added at 10 mM. In some embodiments, rotenone is added at 2.5 mM. In some embodiments, antimycin A is added at 2.5 mM. In some embodiments, 2-deoxyglucose (2-DG) is added at 500 mM. In some embodiments, glycolytic capacity, glycolytic reserve, and/or non-glycolytic acidification are measured. In general, TILs have a glycolytic reserve that is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 50% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 60% to about 99% of the *basal respiration rate* of freshly harvested TILs. In some embodiments, the glycolytic reserve is from about 70% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 80% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 90% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 95% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7).

[00968] In some embodiments, the metabolic assay is basal glycolysis. In some embodiments, second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs)

have an increase in basal glycolysis of at least two-fold, at least three-fold, at least four-fold, at least five-fold, at least six-fold, at least 7-fold, at least eight-fold, at least nine-fold, or at least ten-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about ten-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about eight-fold. In some embodiments, second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about three-fold to about seven-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about four-fold. In some embodiments, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have an increase in basal glycolysis of about two-fold to about three-fold.

[00969] In general, the second expansion TILs or second additional expansion TILs (such as, for example, those described in Step D of Figure 7, including TILs referred to as reREP TILs) have a glycolytic reserve that is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% of the basal respiration rate of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 50% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 60% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 70% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 80% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 90% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7). In some embodiments, the glycolytic reserve is from about 95% to about 99% of the *basal respiration rate* of 2A process TILs (see, for example Figure 7).

[00970] Granzyme B Production: Granzyme B is another measure of the ability of TIL to kill target cells. Media supernatants restimulated as described above using antibodies to CD3, CD28, and CD137/4-1BB were also evaluated for their levels of Granzyme B using the Human Granzyme B DuoSet ELISA Kit (R & D Systems, Minneapolis, MN) according to the manufacturer's instructions.

[00971] In some embodiments, the present methods provide an assay for assessing TIL viability, using the methods as described above. In some embodiments, the TILs are expanded as discussed above, including for example as provided in Figure 7. In some embodiments, the TILs are cryopreserved prior to being assessed for viability. In some embodiments, the viability assessment includes thawing the TILs prior to performing a first expansion, a second expansion, and an additional second expansion. In some embodiments, the present methods provide an assay for assessing cell proliferation, cell toxicity, cell death, and/or other terms related to viability of the TIL population. Viability can be measured by any of the TIL metabolic assays described above as well as any methods know for assessing cell viability that are known in the art. In some embodiments, the present methods provide as assay for assessment of cell proliferation, cell toxicity, cell death, and/or other terms related to viability of the TILs expanded using the methods described herein, including those exemplified in Figure 7.

[00972] The present invention also provides assay methods for determining TIL viability. The present disclosure provides methods for assaying TILs for viability by expanding tumor infiltrating lymphocytes (TILs) into a larger population of TILs comprising:

(i) obtaining a first population of TILs which has been previously expanded;

(ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy; and

[00973] (iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least

100-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, and wherein the third population is further assayed for viability. In some embodiments, the method further comprises:

(iv) performing an additional second expansion by supplementing the cell culture medium of the third population of TILs with additional IL-2, additional OKT-3, and additional APCs, wherein the additional second expansion is performed for at least 14 days to obtain a larger population of TILs than obtained in step (iii), wherein the larger population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the third population of TILs, and wherein the third population is further assayed for viability.

[00974] In some embodiments, prior to step (i), the cells are cryopreserved.

[00975] In some embodiments, the cells are thawed prior to performing step (i).

[00976] In some embodiments, step (iv) is repeated one to four times in order to obtain sufficient TILs for analysis.

[00977] In some embodiments, steps (i) through (iii) or (iv) are performed within a period of about 21 days to about 33 days.

[00978] In some embodiments, steps (i) through (iii) or (iv) are performed within a period of about 21 days to about 30 days.

[00979] In some embodiments, steps (i) through (iii) or (iv) are performed within a period of about 21 days to about 28 days.

[00980] In some embodiments, steps (i) through (iii) or (iv) are performed within about 24 days.

[00981] In some embodiments, the cells from steps (iii) or (iv) express CD4, CD8, and TCR $\alpha\beta$ at levels similar to freshly harvested cells.

[00982] In some embodiments, the antigen presenting cells are peripheral blood mononuclear cells (PBMCs).

[00983] In some embodiments, the PBMCs are added to the cell culture on any of days 9 through 17 in step (iii).

[00984] In some embodiments, the effector T cells and/or central memory T cells in the larger population of TILs in step (iv) exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells, and/or central memory T cells in the third population of cells.

[00985] In some embodiments, the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

[00986] In some embodiments, the APCs are artificial APCs (aAPCs).

[00987] In some embodiments, the method further comprises the step of transducing the first population of TILs with an expression vector comprising a nucleic acid encoding a high-affinity T cell receptor.

[00988] In some embodiments, the step of transducing occurs before step (i).

[00989] In some embodiments, the method further comprises the step of transducing the first population of TILs with an expression vector comprising a nucleic acid encoding a chimeric antigen receptor (CAR) comprising a single chain variable fragment antibody fused with at least one endodomain of a T-cell signaling molecule.

[00990] In some embodiments, the step of transducing occurs before step (i).

[00991] In some embodiments, the TILs are assayed for viability.

[00992] In some embodiments, the TILs are assayed for viability after cryopreservation.

[00993] In some embodiments, the TILs are assayed for viability after cryopreservation and after step (iv).

[00994] In some embodiments, the present disclosure provides a method of treating a subject with cancer comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising: (i) obtaining a first population of TILs from a fine needle aspirate (FNA) or a core biopsy obtained from a tumor in a patient; (ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to

obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy; (iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and (iv) administering a therapeutically effective dosage of the third population of TILs to the patient.

[00995] According to the present disclosure, a method for assaying TILs for viability and/or further use in administration to a subject. In some embodiments, the method for assay tumor infiltrating lymphocytes (TILs) comprises:

(i) obtaining a first population of TILs;

(ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy; and

(iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 50-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs;

(iv) harvesting, washing, and cryopreserving the third population of TILs;

(v) storing the cryopreserved TILs at a cryogenic temperature;

(vi) thawing the third population of TILs to provide a thawed third population of TILs; and

(vii) performing an additional second expansion of a portion of the thawed third

population of TILs by supplementing the cell culture medium of the third population with IL-2, OKT-3, and APCs for a reREP period of at least 3 days, wherein the third expansion is performed to obtain a fourth population of TILs, wherein the number of TILs in the fourth population of TILs is compared to the number of TILs in the third population of TILs to obtain a ratio;

(viii) determining based on the ratio in step (vii) whether the thawed population of TILs is suitable for administration to a patient;

(ix) administering a therapeutically effective dosage of the thawed population of TILs to the patient when the ratio of the number of TILs in the fourth population of TILs to the number of TILs in the third population of TILs is determined to be greater than 5:1 in step (viii).

[00996] In some embodiments, the reREP period is performed until the ratio of the number of TILs in the fourth population of TILs to the number of TILs in the third population of TILs is greater than 50:1.

[00997] In some embodiments, the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

[00998] In some embodiments, steps (i) through (vii) are performed within a period of about 21 days to about 33 days. In some embodiments, steps (i) through (vii) are performed within a period of about 21 days to about 30 days. In some embodiments, steps (i) through (vii) are performed within a period of about 21 days to about 28 days. In some embodiments, steps (i) through (vii) are performed within about 24 days. In some embodiments, the cells from steps (iii) or (vii) express CD4, CD8, and TCR α β at levels similar to freshly harvested cells. In some embodiments the cells are TILs.

[00999] In some embodiments, the antigen presenting cells are peripheral blood mononuclear cells (PBMCs). In some embodiments, the PBMCs are added to the cell culture on any of days 3 through 12 in step (ii) and/or any of days 11 through 14 in step (iii).

[001000] In some embodiments, the effector T cells and/or central memory T cells in the larger population of TILs in steps (iii) or (vii) exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells, and/or central memory T cells in the third population of cells.

[001001] In some embodiments, the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

[001002] In some embodiments, the APCs are artificial APCs (aAPCs).

[001003] In some embodiments, the step of transducing the first population of TILs with an expression vector comprising a nucleic acid encoding a high-affinity T cell receptor.

[001004] In some embodiments, the step of transducing occurs before step (i).

[001005] In some embodiments, the step of transducing the first population of TILs with an expression vector comprising a nucleic acid encoding a chimeric antigen receptor (CAR) comprising a single chain variable fragment antibody fused with at least one endodomain of a T-cell signaling molecule.

[001006] In some embodiments, the step of transducing occurs before step (i).

[001007] In some embodiments, the TILs are assayed for viability after step (vii).

[001008] The present disclosure also provides further methods for assaying TILs. In some embodiments, the disclosure provides a method for assaying TILs comprising:

(i) obtaining a portion of a first population of cryopreserved TILs obtained from a core biopsy sample;

(ii) thawing the portion of the first population of cryopreserved TILs;

(iii) performing a first expansion by culturing the portion of the first population of TILs in a cell culture medium comprising IL-2, OKT-3, and antigen presenting cells (APCs) for a reREP period of at least 3 days, to produce a second population of TILs, wherein the portion from the first population of TILs is compared to the second population of TILs to obtain a ratio of the number of TILs, wherein the ratio of the number of TILs in the second population of TILs to the number of TILs in the portion of the first population of TILs is greater than 5:1, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days;

(iv) determining based on the ratio in step (iii) and the number of TILs in step (iii) whether the first population of TILs is suitable for use in therapeutic administration to a patient;

(v) determining the first population of TILs is suitable for use in therapeutic administration when the ratio of the number of TILs in the second population of TILs

to the number of TILs in the first population of TILs is determined to be greater than 5:1 in step (iv).

[001009] In some embodiments, the ratio of the number of TILs in the second population of TILs to the number of TILs in the portion of the first population of TILs is greater than 50:1.

[001010] In some embodiments, the method further comprises performing expansion of the entire first population of cryopreserved TILs from step (i) according to the methods as described in any of the embodiments provided herein.

[001011] In some embodiments, the method further comprises administering the entire first population of cryopreserved TILs from step (i) to the patient.

IX. Closed Systems for TIL Manufacturing

[001012] The present invention provides for the use of closed systems during the TIL culturing process using Gen 2 processes or Gen 3 processes. Such closed systems allow for preventing and/or reducing microbial contamination, allow for the use of fewer flasks, and allow for cost reductions. In some embodiments, the closed system uses two containers.

[001013] Such closed systems are well-known in the art and can be found, for example, at http://www.fda.gov/cber/guidelines.htm and https://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/ucm076779.htm.

[001014] Sterile connecting devices (STCDs) produce sterile welds between two pieces of compatible tubing. This procedure permits sterile connection of a variety of containers and tube diameters. In some embodiments, the closed systems include luer lock and heat sealed systems. In some embodiments, the closed system is accessed via syringes under sterile conditions in order to maintain the sterility and closed nature of the system. In some embodiments, a closed system is employed. In some embodiments, the TILs are formulated into a final product formulation container.

[001015] In some embodiments, the closed system uses one container from the time the tumor fragments are obtained until the TILs are ready for administration to the patient or cryopreserving. In some embodiments when two containers are used, the first container is a closed G-container and the population of TILs is centrifuged and transferred to an infusion bag without opening the first closed G-container. In some embodiments, when two containers

are used, the infusion bag is a HypoThermosol-containing infusion bag. A closed system or closed TIL cell culture system is characterized in that once the tumor sample and/or tumor fragments have been added, the system is tightly sealed from the outside to form a closed environment free from the invasion of bacteria, fungi, and/or any other microbial contamination.

[001016] In some embodiments, the reduction in microbial contamination is between about 5% and about 100%. In some embodiments, the reduction in microbial contamination is between about 5% and about 95%. In some embodiments, the reduction in microbial contamination is between about 5% and about 90%. In some embodiments, the reduction in microbial contamination is between about 10% and about 90%. In some embodiments, the reduction in microbial contamination is between about 15% and about 85%. In some embodiments, the reduction in microbial contamination is about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 97%, about 98%, about 99%, or about 100%.

[001017] The closed system allows for TIL growth in the absence and/or with a significant reduction in microbial contamination.

[001018] Moreover, pH, carbon dioxide partial pressure and oxygen partial pressure of the TIL cell culture environment each vary as the cells are cultured. Consequently, even though a medium appropriate for cell culture is circulated, the closed environment still needs to be constantly maintained as an optimal environment for TIL proliferation. To this end, it is desirable that the physical factors of pH, carbon dioxide partial pressure and oxygen partial pressure within the culture liquid of the closed environment be monitored by means of a sensor, the signal whereof is used to control a gas exchanger installed at the inlet of the culture environment, and the that gas partial pressure of the closed environment be adjusted in real time according to changes in the culture liquid so as to optimize the cell culture environment. In some embodiments, the present invention provides a closed cell culture system which incorporates at the inlet to the closed environment a gas exchanger equipped with a monitoring device which measures the pH, carbon dioxide partial pressure and oxygen partial pressure of the closed environment, and optimizes the cell culture environment by automatically adjusting gas concentrations based on signals from the monitoring device.

[001019] In some embodiments, the pressure within the closed environment is continuously or intermittently controlled. That is, the pressure in the closed environment can be varied by means of a pressure maintenance device for example, thus ensuring that the space is suitable for growth of TILs in a positive pressure state, or promoting exudation of fluid in a negative pressure state and thus promoting cell proliferation. By applying negative pressure intermittently, moreover, it is possible to uniformly and efficiently replace the circulating liquid in the closed environment by means of a temporary shrinkage in the volume of the closed environment.

[001020] In some embodiments, optimal culture components for proliferation of the TILs can be substituted or added, and including factors such as IL-2 and/or OKT3, as well as combination, can be added.

X. Optional Genetic Engineering of TILs

[001021] In some embodiments of Gen 2 or Gen 3 processes, the expanded TILs of the present invention are further manipulated before, during, or after an expansion step, including during closed, sterile manufacturing processes, each as provided herein, in order to alter protein expression in a transient manner. In some embodiments, the transiently altered protein expression is due to transient gene editing. In some embodiments, the expanded TILs of the present invention are treated with transcription factors (TFs) and/or other molecules capable of transiently altering protein expression in the TILs. In some embodiments, the TFs and/or other molecules that are capable of transiently altering protein expression provide for altered expression of tumor antigens and/or an alteration in the number of tumor antigen-specific T cells in a population of TILs.

[001022] In certain embodiments, the method comprises genetically editing a population of TILs. In certain embodiments, the method comprises genetically editing the first population of TILs, the second population of TILs and/or the third population of TILs.

[001023] In some embodiments, the present invention includes genetic editing through nucleotide insertion, such as through ribonucleic acid (RNA) insertion, including insertion of messenger RNA (mRNA) or small (or short) interfering RNA (siRNA), into a population of TILs for promotion of the expression of one or more proteins or inhibition of the expression of one or more proteins, as well as simultaneous combinations of both promotion of one set of proteins with inhibition of another set of proteins.

[001024] In some embodiments, the expanded TILs of the present invention undergo transient alteration of protein expression. In some embodiments, the transient alteration of protein expression occurs in the bulk TIL population prior to first expansion, including, for example in the TIL population obtained from for example, Step A as indicated in Figure 85 (particularly Figure 85B). In some embodiments, the transient alteration of protein expression occurs during the first expansion, including, for example in the TIL population expanded in for example, Step B as indicated in Figure 85 (for example Figure 85B). In some embodiments, the transient alteration of protein expression occurs after the first expansion, including, for example in the TIL population in transition between the first and second expansion (*e.g.* the second population of TILs as described herein), the TIL population obtained from for example, Step B and included in Step C as indicated in Figure 85. In some embodiments, the transient alteration of protein expression occurs in the bulk TIL population prior to second expansion, including, for example in the TIL population obtained from for example, Step C and prior to its expansion in Step D as indicated in Figure 85. In some embodiments, the transient alteration of protein expression occurs during the second expansion, including, for example in the TIL population expanded in for example, Step D as indicated in Figure 85 (*e.g.* the third population of TILs). In some embodiments, the transient alteration of protein expression occurs after the second expansion, including, for example in the TIL population obtained from the expansion in for example, Step D as indicated in Figure 85.

[001025] In an embodiment, a method of transiently altering protein expression in a population of TILs includes the step of electroporation. Electroporation methods are known in the art and are described, *e.g.*, in Tsong, *Biophys. J.* **1991,** *60,* 297-306, and U.S. Patent Application Publication No. 2014/0227237 A1, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of transiently altering protein expression in population of TILs includes the step of calcium phosphate transfection. Calcium phosphate transfection methods (calcium phosphate DNA precipitation, cell surface coating, and endocytosis) are known in the art and are described in Graham and van der Eb, *Virology* **1973,** *52,* 456-467; Wigler, *et al.*, *Proc. Natl. Acad. Sci.* **1979,** *76,* 1373-1376; and Chen and Okayarea, *Mol. Cell. Biol.* **1987,** *7,* 2745-2752; and in U.S. Patent No. 5,593,875, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of transiently altering protein expression in a population of TILs includes the step of liposomal transfection. Liposomal transfection methods, such as methods that employ a 1:1

(w/w) liposome formulation of the cationic lipid *N*-[1-(2,3-dioleyloxy)propyl]-*n,n,n*-trimethylammonium chloride (DOTMA) and dioleoyl phophotidylethanolamine (DOPE) in filtered water, are known in the art and are described in Rose, *et al.*, *Biotechniques* **1991,** *10,* 520-525 and Felgner, *et al.*, *Proc. Natl. Acad. Sci. USA*, **1987,** *84,* 7413-7417 and in U.S. Patent Nos. 5,279,833; 5,908,635; 6,056,938; 6,110,490; 6,534,484; and 7,687,070, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of transiently altering protein expression in a population of TILs includes the step of transfection using methods described in U.S. Patent Nos. 5,766,902; 6,025,337; 6,410,517; 6,475,994; and 7,189,705; the disclosures of each of which are incorporated by reference herein.

[001026] In some embodiments, transient alteration of protein expression results in an increase in Stem Memory T cells (TSCMs). TSCMs are early progenitors of antigen-experienced central memory T cells. TSCMs generally display the long-term survival, self-renewal, and multipotency abilities that define stem cells, and are generally desirable for the generation of effective TIL products. TSCM have shown enhanced anti-tumor activity compared with other T cell subsets in mouse models of adoptive cell transfer (Gattinoni *et al.* Nat Med 2009, 2011; Gattinoni, Nature Rev. Cancer, 2012; Cieri *et al.* Blood 2013). In some embodiments, transient alteration of protein expression results in a TIL population with a composition comprising a high proportion of TSCM. In some embodiments, transient alteration of protein expression results in an at least 5%, at least 10%, at least 10%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% increase in TSCM percentage. In some embodiments, transient alteration of protein expression results in an at least a 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, or 10-fold increase in TSCMs in the TIL population. In some embodiments, transient alteration of protein expression results in a TIL population with at least at least 5%, at least 10%, at least 10%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% TSCMs. In some embodiments, transient alteration of protein expression results in a therapeutic TIL population with at least at least 5%, at least 10%, at least 10%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% TSCMs.

[001027] In some embodiments, transient alteration of protein expression results in rejuvenation of antigen-experienced T-cells. In some embodiments, rejuvenation includes, for example, increased proliferation, increased T-cell activation, and/or increased antigen recognition.

[001028] In some embodiments, transient alteration of protein expression alters the expression in a large fraction of the T-cells in order to preserve the tumor-derived TCR repertoire. In some embodiments, transient alteration of protein expression does not alter the tumor-derived TCR repertoire. In some embodiments, transient alteration of protein expression maintains the tumor-derived TCR repertoire.

[001029] In some embodiments, transient alteration of protein results in altered expression of a particular gene. In some embodiments, the transient alteration of protein expression targets a gene including but not limited to PD-1 (also referred to as PDCD1 or CC279), TGFBR2, CCR4/5, CBLB (CBL-B), CISH, CCRs (chimeric co-stimulatory receptors), IL-2, IL-12, IL-15, IL-21, NOTCH 1/2 ICD, TIM3, LAG3, TIGIT, TGFβ, CCR2, CCR4, CCR5, CXCR1, CXCR2, CSCR3, CCL2 (MCP-1), CCL3 (MIP-1α), CCL4 (MIP1-β), CCL5 (RANTES), CXCL1/CXCL8, CCL22, CCL17, CXCL1/CXCL8, VHL, CD44, PIK3CD, SOCS1, and/or cAMP protein kinase A (PKA). In some embodiments, the transient alteration of protein expression targets a gene selected from the group consisting of PD-1, TGFBR2, CCR4/5, CBLB (CBL-B), CISH, CCRs (chimeric co-stimulatory receptors), IL-2, IL-12, IL-15, IL-21, NOTCH 1/2 ICD, TIM3, LAG3, TIGIT, TGFβ, CCR2, CCR4, CCR5, CXCR1, CXCR2, CSCR3, CCL2 (MCP-1), CCL3 (MIP-1α), CCL4 (MIP1-β), CCL5 (RANTES), CXCL1/CXCL8, CCL22, CCL17, CXCL1/CXCL8, VHL, CD44, PIK3CD, SOCS1, and/or cAMP protein kinase A (PKA). In some embodiments, the transient alteration of protein expression targets PD-1. In some embodiments, the transient alteration of protein expression targets TGFBR2. In some embodiments, the transient alteration of protein expression targets CCR4/5. In some embodiments, the transient alteration of protein expression targets CBLB. In some embodiments, the transient alteration of protein expression targets CISH. In some embodiments, the transient alteration of protein expression targets CCRs (chimeric co-stimulatory receptors). In some embodiments, the transient alteration of protein expression targets IL-2. In some embodiments, the transient alteration of protein expression targets IL-12. In some embodiments, the transient alteration of protein expression targets IL-15. In some embodiments, the transient alteration of protein expression targets IL-21. In some embodiments, the transient alteration of protein expression targets NOTCH 1/2 ICD. In some

embodiments, the transient alteration of protein expression targets TIM3. In some embodiments, the transient alteration of protein expression targets LAG3. In some embodiments, the transient alteration of protein expression targets TIGIT. In some embodiments, the transient alteration of protein expression targets TGFβ. In some embodiments, the transient alteration of protein expression targets CCR1. In some embodiments, the transient alteration of protein expression targets CCR2. In some embodiments, the transient alteration of protein expression targets CCR4. In some embodiments, the transient alteration of protein expression targets CCR5. In some embodiments, the transient alteration of protein expression targets CXCR1. In some embodiments, the transient alteration of protein expression targets CXCR2. In some embodiments, the transient alteration of protein expression targets CSCR3. In some embodiments, the transient alteration of protein expression targets CCL2 (MCP-1). In some embodiments, the transient alteration of protein expression targets CCL3 (MIP-1α). In some embodiments, the transient alteration of protein expression targets CCL4 (MIP1-β). In some embodiments, the transient alteration of protein expression targets CCL5 (RANTES). In some embodiments, the transient alteration of protein expression targets CXCL1. In some embodiments, the transient alteration of protein expression targets CXCL8. In some embodiments, the transient alteration of protein expression targets CCL22. In some embodiments, the transient alteration of protein expression targets CCL17. In some embodiments, the transient alteration of protein expression targets VHL. In some embodiments, the transient alteration of protein expression targets CD44. In some embodiments, the transient alteration of protein expression targets PIK3CD. In some embodiments, the transient alteration of protein expression targets SOCS1. In some embodiments, the transient alteration of protein expression targets cAMP protein kinase A (PKA).

[001030] In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of a chemokine receptor. In some embodiments, the chemokine receptor that is overexpressed by transient protein expression includes a receptor with a ligand that includes but is not limited to CCL2 (MCP-1), CCL3 (MIP-1α), CCL4 (MIP1-β), CCL5 (RANTES), CXCL1, CXCL8, CCL22, and/or CCL17.

[001031] In some embodiments, the transient alteration of protein expression results in a decrease and/or reduced expression of PD-1, CTLA-4, TIM-3, LAG-3, TIGIT, TGFβR2, and/or TGFβ (including resulting in, for example, TGFβ pathway blockade). In some

embodiments, the transient alteration of protein expression results in a decrease and/or reduced expression of CBLB (CBL-B). In some embodiments, the transient alteration of protein expression results in a decrease and/or reduced expression of CISH.

[001032] In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of chemokine receptors in order to, for example, improve TIL trafficking or movement to the tumor site. In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of a CCR (chimeric co-stimulatory receptor). In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of a chemokine receptor selected from the group consisting of CCR1, CCR2, CCR4, CCR5, CXCR1, CXCR2, and/or CSCR3.

[001033] In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of an interleukin. In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of an interleukin selected from the group consisting of IL-2, IL-12, IL-15, and/or IL-21.

[001034] In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of NOTCH 1/2 ICD. In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of VHL. In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of CD44. In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of PIK3CD. In some embodiments, the transient alteration of protein expression results in increased and/or overexpression of SOCS1,

[001035] In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of cAMP protein kinase A (PKA).

[001036] In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of a molecule selected from the group consisting of PD-1, LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB, BAFF (BR3), and combinations thereof. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of two molecules selected from the group consisting of PD-1, LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB, BAFF (BR3), and combinations thereof. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of PD-1 and one molecule selected from the group consisting of LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB,

BAFF (BR3), and combinations thereof. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of PD-1, LAG-3, CISH, CBLB, TIM3, and combinations thereof. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of PD-1 and one of LAG3, CISH, CBLB, TIM3, and combinations thereof. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of PD-1 and LAG3. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of PD-1 and CISH. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of PD-1 and CBLB. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of LAG3 and CISH. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of LAG3 and CBLB. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of CISH and CBLB. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of TIM3 and PD-1. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of TIM3 and LAG3. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of TIM3 and CISH. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of TIM3 and CBLB.

[001037] In some embodiments, an adhesion molecule selected from the group consisting of CCR2, CCR4, CCR5, CXCR2, CXCR3, CX3CR1, and combinations thereof, is inserted by a gammaretroviral or lentiviral method into the first population of TILs, second population of TILs, or harvested population of TILs (*e.g.*, the expression of the adhesion molecule is increased).

[001038] In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of a molecule selected from the group consisting of PD-1, LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB, BAFF (BR3), and combinations thereof, and increased and/or enhanced expression of CCR2, CCR4, CCR5, CXCR2, CXCR3, CX3CR1, and combinations thereof. In some embodiments, the transient alteration of protein expression results in decreased and/or reduced expression of a molecule selected from the group consisting of PD-1, LAG3, TIM3, CISH, CBLB, and combinations

thereof, and increased and/or enhanced expression of CCR2, CCR4, CCR5, CXCR2, CXCR3, CX3CR1, and combinations thereof.

[001039] In some embodiments, there is a reduction in expression of about 5%, about 10%, about 10%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 80%. In some embodiments, there is a reduction in expression of at least about 85%, In some embodiments, there is a reduction in expression of at least about 90%. In some embodiments, there is a reduction in expression of at least about 95%. In some embodiments, there is a reduction in expression of at least about 99%.

[001040] In some embodiments, there is an increase in expression of about 5%, about 10%, about 10%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is an increase in expression of at least about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is an increase in expression of at least about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is an increase in expression of at least about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is an increase in expression of at least about 85%, about 90%, or about 95%. In some embodiments, there is an increase in expression of at least about 80%. In some embodiments, there is an increase in expression of at least about 85%, In some embodiments, there is an increase in expression of at least about 90%. In some embodiments, there is an increase in expression of at least about 95%. In some embodiments, there is an increase in expression of at least about 99%.

[001041] In some embodiments, transient alteration of protein expression is induced by treatment of the TILs with transcription factors (TFs) and/or other molecules capable of transiently altering protein expression in the TILs. In some embodiments, the SQZ vector-

free microfluidic platform is employed for intracellular delivery of the transcription factors (TFs) and/or other molecules capable of transiently altering protein expression. Such methods demonstrating the ability to deliver proteins, including transcription factors, to a variety of primary human cells, including T cells (Sharei *et al.* PNAS 2013, as well as Sharei *et al.,* PLOS ONE 2015 and Greisbeck *et al.,* J. Immunology vol. 195, 2015) have been described; see, for example, International Patent Publications WO 2013/059343A1, WO 2017/008063A1, and WO 2017/123663A1, all of which are incorporated by reference herein in their entireties. Such methods as described in International Patent Publications WO 2013/059343A1, WO 2017/008063A1, and WO 2017/123663A1 can be employed with the present invention in order to expose a population of TILs to transcription factors (TFs) and/or other molecules capable of inducing transient protein expression, wherein said TFs and/or other molecules capable of inducing transient protein expression provide for increased expression of tumor antigens and/or an increase in the number of tumor antigen-specific T cells in the population of TILs, thus resulting in reprogramming of the TIL population and an increase in therapeutic efficacy of the reprogrammed TIL population as compared to a non-reprogrammed TIL population. In some embodiments, the reprogramming results in an increased subpopulation of effector T cells and/or central memory T cells relative to the starting or prior population (*i.e.*, prior to reprogramming) population of TILs, as described herein.

[001042] In some embodiments, the transcription factor (TF) includes but is not limited to TCF-1, NOTCH 1/2 ICD, and/or MYB. In some embodiments, the transcription factor (TF) is TCF-1. In some embodiments, the transcription factor (TF) is NOTCH 1/2 ICD. In some embodiments, the transcription factor (TF) is MYB. In some embodiments, the transcription factor (TF) is administered with induced pluripotent stem cell culture (iPSC), such as the commercially available KNOCKOUT Serum Replacement (Gibco/ThermoFisher), to induce additional TIL reprogramming. In some embodiments, the transcription factor (TF) is administered with an iPSC cocktail to induce additional TIL reprogramming. In some embodiments, the transcription factor (TF) is administered without an iPSC cocktail. In some embodiments, reprogramming results in an increase in the percentage of TSCMs. In some embodiments, reprogramming results in an increase in the percentage of TSCMs by about 5%, about 10%, about 10%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% TSCMs.

[001043] In some embodiments, a method of transient altering protein expression, as described above, may be combined with a method of genetically modifying a population of TILs includes the step of stable incorporation of genes for production of one or more proteins. In certain embodiments, the method comprises a step of genetically modifying a population of TILs. In certain embodiments, the method comprises genetically modifying the first population of TILs, the second population of TILs and/or the third population of TILs. In an embodiment, a method of genetically modifying a population of TILs includes the step of retroviral transduction. In an embodiment, a method of genetically modifying a population of TILs includes the step of lentiviral transduction. Lentiviral transduction systems are known in the art and are described, *e.g.*, in Levine, *et al.*, *Proc. Nat'l Acad. Sci.* **2006,** *103,* 17372-77; Zufferey, *et al., Nat. Biotechnol.* **1997,** *15,* 871-75; Dull, *et al.*, *J. Virology* **1998,** *72,* 8463-71, and U.S. Patent No. 6,627,442, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of genetically modifying a population of TILs includes the step of gamma-retroviral transduction. Gamma-retroviral transduction systems are known in the art and are described, *e.g.*, Cepko and Pear, *Cur. Prot. Mol. Biol.* **1996,** 9.9.1-9.9.16, the disclosure of which is incorporated by reference herein. In an embodiment, a method of genetically modifying a population of TILs includes the step of transposon-mediated gene transfer. Transposon-mediated gene transfer systems are known in the art and include systems wherein the transposase is provided as DNA expression vector or as an expressible RNA or a protein such that long-term expression of the transposase does not occur in the transgenic cells, for example, a transposase provided as an mRNA (*e.g.*, an mRNA comprising a cap and poly-A tail). Suitable transposon-mediated gene transfer systems, including the salmonid-type Tel-like transposase (SB or Sleeping Beauty transposase), such as SB10, SB11, and SB100x, and engineered enzymes with increased enzymatic activity, are described in, *e.g.*, Hackett, *et al.*, *Mol. Therapy* **2010,** *18,* 674-83 and U.S. Patent No. 6,489,458, the disclosures of each of which are incorporated by reference herein.

[001044] In some embodiments, transient alteration of protein expression is a reduction in expression induced by self-delivering RNA interference (sdRNA), which is a chemically-synthesized asymmetric siRNA duplex with a high percentage of 2′-OH substitutions (typically fluorine or -OCH_3) which comprises a 20-nucleotide antisense (guide) strand and a 13 to 15 base sense (passenger) strand conjugated to cholesterol at its 3′ end using a tetraethylenglycol (TEG) linker. In some embodiments, the method comprises transient

alteration of protein expression in a population of TILs, comprising the use of self-delivering RNA interference (sdRNA), which is a chemically-synthesized asymmetric siRNA duplex with a high percentage of 2′-OH substitutions (typically fluorine or -OCH_3) which comprises a 20-nucleotide antisense (guide) strand and a 13 to 15 base sense (passenger) strand conjugated to cholesterol at its 3' end using a tetraethylenglycol (TEG) linker. Methods of using sdRNA have been described in Khvorova and Watts, *Nat. Biotechnol.* **2017,** *35,* 238–248; Byrne, *et al.*, *J. Ocul. Pharmacol. Ther.* **2013,** *29,* 855-864; and Ligtenberg, *et al.*, *Mol. Therapy,* **2018,** in press, the disclosures of which are incorporated by reference herein. In an embodiment, delivery of sdRNA to a TIL population is accomplished without use of electroporation, SQZ, or other methods, instead using a 1 to 3 day period in which a TIL population is exposed to sdRNA at a concentration of 1 μM/10,000 TILs in medium. In certain embodiments, the method comprises delivery sdRNA to a TILs population comprising exposing the TILs population to sdRNA at a concentration of 1 μM/10,000 TILs in medium for a period of between 1 to 3 days. In an embodiment, delivery of sdRNA to a TIL population is accomplished using a 1 to 3 day period in which a TIL population is exposed to sdRNA at a concentration of 10 μM/10,000 TILs in medium. In an embodiment, delivery of sdRNA to a TIL population is accomplished using a 1 to 3 day period in which a TIL population is exposed to sdRNA at a concentration of 50 μM/10,000 TILs in medium. In an embodiment, delivery of sdRNA to a TIL population is accomplished using a 1 to 3 day period in which a TIL population is exposed to sdRNA at a concentration of between 0.1 μM/10,000 TILs and 50 μM/10,000 TILs in medium. In an embodiment, delivery of sdRNA to a TIL population is accomplished using a 1 to 3 day period in which a TIL population is exposed to sdRNA at a concentration of between 0.1 μM/10,000 TILs and 50 μM/10,000 TILs in medium, wherein the exposure to sdRNA is performed two, three, four, or five times by addition of fresh sdRNA to the media. Other suitable processes are described, for example, in U.S. Patent Application Publication No. US 2011/0039914 A1, US 2013/0131141 A1, and US 2013/0131142 A1, and U.S. Patent No. 9,080,171, the disclosures of which are incorporated by reference herein.

[001045] In some embodiments, sdRNA is inserted into a population of TILs during manufacturing. In some embodiments, the sdRNA encodes RNA that interferes with NOTCH 1/2 ICD, PD-1, CTLA-4 TIM-3, LAG-3, TIGIT, TGFβ, TGFBR2, cAMP protein kinase A (PKA), BAFF BR3, CISH, and/or CBLB. In some embodiments, the reduction in expression is determined based on a percentage of gene silencing, for example, as assessed by

flow cytometry and/or qPCR. In some embodiments, there is a reduction in expression of about 5%, about 10%, about 10%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 75%, about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 80%, about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 85%, about 90%, or about 95%. In some embodiments, there is a reduction in expression of at least about 80%. In some embodiments, there is a reduction in expression of at least about 85%, In some embodiments, there is a reduction in expression of at least about 90%. In some embodiments, there is a reduction in expression of at least about 95%. In some embodiments, there is a reduction in expression of at least about 99%.

[001046] The self-deliverable RNAi technology based on the chemical modification of siRNAs can be employed with the methods of the present invention to successfully deliver the sdRNAs to the TILs as described herein. The combination of backbone modifications with asymmetric siRNA structure and a hydrophobic ligand (*see*, for eample, Ligtenberg, *et al.*, *Mol. Therapy,* **2018** and US20160304873) allow sdRNAs to penetrate cultured mammalian cells without additional formulations and methods by simple addition to the culture media, capitalizing on the nuclease stability of sdRNAs. This stability allows the support of constant levels of RNAi-mediated reduction of target gene activity simply by maintaining the active concentration of sdRNA in the media. While not being bound by theory, the backbone stabilization of sdRNA provides for extended reduction in gene expression effects which can last for months in non-dividing cells.

[001047] In some embodiments, over 95% transfection efficiency of TILs and a reduction in expression of the target by various specific sdRNA occurs. In some embodiments, sdRNAs containing several unmodified ribose residues were replaced with fully modified sequences to increase potency and/or the longevity of RNAi effect. In some embodiments, a reduction in expression effect is maintained for 12 hours, 24 hours, 36 hours, 48 hours, 5 days, 6 days, 7 days, or 8 days or more. In some embodiments, the reduction in expression effect decreases at 10 days or more post sdRNA treatment of the TILs. In some embodiments, more than 70% reduction in expression of the target expression is maintained. In some embodiments, more

than 70% reduction in expression of the target expression is maintained TILs. In some embodiments, a reduction in expression in the PD-1/PD-L1 pathway allows for the TILs to exhibit a more potent *in vivo* effect, which is in some embodiments, due to the avoidance of the suppressive effects of the PD-1/PD-L1 pathway. In some embodiments, a reduction in expression of PD-1 by sdRNA results in an increase TIL proliferation.

[001048] Small interfering RNA (siRNA), sometimes known as short interfering RNA or silencing RNA, is a double stranded RNA molecule, generally 19-25 base pairs in length. siRNA is used in RNA interference (RNAi), where it interferes with expression of specific genes with complementary nucleotide sequences.

[001049] Double stranded DNA (dsRNA) can be generally used to define any molecule comprising a pair of complementary strands of RNA, generally a sense (passenger) and antisense (guide) strands, and may include single-stranded overhang regions. The term dsRNA, contrasted with siRNA, generally refers to a precursor molecule that includes the sequence of an siRNA molecule which is released from the larger dsRNA molecule by the action of cleavage enzyme systems, including Dicer.

[001050] sdRNA (self-deliverable RNA) are a new class of covalently modified RNAi compounds that do not require a delivery vehicle to enter cells and have improved pharmacology compared to traditional siRNAs. "Self-deliverable RNA" or "sdRNA" is a hydrophobically modified RNA interfering-antisense hybrid, demonstrated to be highly efficacious *in vitro* in primary cells and *in vivo* upon local administration. Robust uptake and/or silencing without toxicity has been demonstrated. sdRNAs are generally asymmetric chemically modified nucleic acid molecules with minimal double stranded regions. sdRNA molecules typically contain single stranded regions and double stranded regions, and can contain a variety of chemical modifications within both the single stranded and double stranded regions of the molecule. Additionally, the sdRNA molecules can be attached to a hydrophobic conjugate such as a conventional and advanced sterol-type molecule, as described herein. sdRNAs and associated methods for making such sdRNAs have also been described extensively in, for example, US20160304873, WO2010033246, WO2017070151, WO2009102427, WO2011119887, WO2010033247A2, WO2009045457, WO2011119852, all of which are incorporated by reference herein in their entireties for all purposes. To optimize sdRNA structure, chemistry, targeting position, sequence preferences, and the like, a proprietary algorithm has been developed and utilized for sdRNA potency prediction (*see*, for example, US 20160304873). Based on these analyses, functional sdRNA sequences have

been generally defined as having over 70% reduction in expression at 1 μM concentration, with a probability over 40%.

[001051] In some embodiments, the sdRNA sequences used in the invention exhibit a 70% reduction in expression of the target gene. In some embodiments, the sdRNA sequences used in the invention exhibit a 75% reduction in expression of the target gene.
In some embodiments, the sdRNA sequences used in the invention exhibit an 80% reduction in expression of the target gene. In some embodiments, the sdRNA sequences used in the invention exhibit an 85% reduction in expression of the target gene. In some embodiments, the sdRNA sequences used in the invention exhibit a 90% reduction in expression of the target gene. In some embodiments, the sdRNA sequences used in the invention exhibit a 95% reduction in expression of the target gene. In some embodiments, the sdRNA sequences used in the invention exhibit a 99% reduction in expression of the target gene. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 0.25 μM to about 4 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 0.25 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 0.5 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 0.75 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 1.0 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 1.25 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 1.5 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 1.75 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 2.0 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 2.25 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a

concentration of about 2.5 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 2.75 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 3.0 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 3.25 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 3.5 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 3.75 μM. In some embodiments, the sdRNA sequences used in the invention exhibit a reduction in expression of the target gene when delivered at a concentration of about 4.0 μM.

[001052] In some emodiments, the oligonucleotide agents comprise one or more modification to increase stability and/or effectiveness of the therapeutic agent, and to effect efficient delivery of the oligonucleotide to the cells or tissue to be treated. Such modifications can include a 2'-O-methyl modification, a 2'-O-Fluro modification, a diphosphorothioate modification, 2' F modified nucleotide, a2'-O-methyl modified and/or a 2'deoxy nucleotide. In some embodiments, the oligonucleotide is modified to include one or more hydrophobic modifications including, for example, sterol, cholesterol, vitamin D, naphtyl, isobutyl, benzyl, indol, tryptophane, and/or phenyl. In an additional particular embodiment, chemically modified nucleotides are combination of phosphorothioates, 2'-O-methyl, 2'deoxy, hydrophobic modifications and phosphorothioates. In some embodiments, the sugars can be modified and modified sugars can include but are not limited to D-ribose, 2'-O-alkyl (including 2'-O-methyl and 2'-0-ethyl), i.e., 2'-alkoxy, 2'-amino, 2'-S-alkyl, 2'-halo (including 2'-fluoro), T- methoxyethoxy, 2'-allyloxy (-$OCH_2CH{=}CH_2$), 2'-propargyl, 2'-propyl, ethynyl, ethenyl, propenyl, and cyano and the like. In one embodiment, the sugar moiety can be a hexose and incorporated into an oligonucleotide as described (Augustyns, K., *et al.*, Nucl. Acids. Res. 18:4711 (1992)).

[001053] In some embodiments, the double-stranded oligonucleotide of the invention is double-stranded over its entire length, *i.e.*, with no overhanging single-stranded sequence at either end of the molecule, *i.e.*, is blunt-ended. In some embodiments, the individual nucleic acid molecules can be of different lengths. In other words, a double-stranded oligonucleotide

of the invention is not double-stranded over its entire length. For instance, when two separate nucleic acid molecules are used, one of the molecules, *e.g.*, the first molecule comprising an antisense sequence, can be longer than the second molecule hybridizing thereto (leaving a portion of the molecule single-stranded). In some embodiments, when a single nucleic acid molecule is used a portion of the molecule at either end can remain single-stranded.

[001054] In some embodiments, a double-stranded oligonucleotide of the invention contains mismatches and/or loops or bulges, but is double-stranded over at least about 70% of the length of the oligonucleotide. In some embodiments, a double-stranded oligonucleotide of the invention is double-stranded over at least about 80% of the length of the oligonucleotide. In another embodiment, a double-stranded oligonucleotide of the invention is double-stranded over at least about 90%-95% of the length of the oligonucleotide. In some embodiments, a double-stranded oligonucleotide of the invention is double-stranded over at least about 96%-98% of the length of the oligonucleotide. In some embodiments, the double-stranded oligonucleotide of the invention contains at least or up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 mismatches.

[001055] In some embodiments, the oligonucleotide can be substantially protected from nucleases *e.g.*, by modifying the 3' or 5' linkages (*e.g.*, U.S. Pat. No. 5,849,902 and WO 98/13526). For example, oligonucleotides can be made resistant by the inclusion of a "blocking group." The term "blocking group" as used herein refers to substituents (*e.g.*, other than OH groups) that can be attached to oligonucleotides or nucleomonomers, either as protecting groups or coupling groups for synthesis (*e.g.*, FITC, propyl (CH_2-CH_2-CH_3), glycol (-0-CH_2-CH_2-O-) phosphate (PO_3^{2+}), hydrogen phosphonate, or phosphoramidite). "Blocking groups" can also include "end blocking groups" or "exonuclease blocking groups" which protect the 5' and 3' termini of the oligonucleotide, including modified nucleotides and non-nucleotide exonuclease resistant structures.

[001056] In some embodiments, at least a portion of the contiguous polynucleotides within the sdRNA are linked by a substitute linkage, *e.g.*, a phosphorothioate linkage.

[001057] In some embodiments, chemical modification can lead to at least a 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 enhancements in cellular uptake. In some embodiments, at least one of the C or U residues includes a hydrophobic

modification. In some embodiments, a plurality of Cs and Us contain a hydrophobic modification. In some embodiments, at least 10%, 15%, 20%, 30%, 40%, 50%, 55%, 60% 65%, 70%, 75%, 80%, 85%, 90% or at least 95% of the Cs and Us can contain a hydrophobic modification. In some embodiments, all of the Cs and Us contain a hydrophobic modification.

[001058] In some embodiments, the sdRNA or sd-rxRNAs exhibit enhanced endosomal release of sd-rxRNA molecules through the incorporation of protonatable amines. In some embodiments, protonatable amines are incorporated in the sense strand (in the part of the molecule which is discarded after RISC loading). In some embodiments, the sdRNA compounds of the invention comprise an asymmetric compound comprising a duplex region (required for efficient RISC entry of 10-15 bases long) and single stranded region of 4-12 nucleotides long; with a 13 nucleotide duplex. In some embodiments, a 6 nucleotide single stranded region is employed. In some embodiments, the single stranded region of the sdRNA comprises 2-12 phosphorothioate intemucleotide linkages (referred to as phosphorothioate modifications). In some embodiments, 6-8 phosphorothioate intemucleotide linkages are employed. In some embodiments, the sdRNA compounds of the invention also include a unique chemical modification pattern, which provides stability and is compatible with RISC entry.

[001059] The guide strand, for example, may also be modified by any chemical modification which confirms stability without interfering with RISC entry. In some embodiments, the chemical modification pattern in the guide strand includes the majority of C and U nucleotides being 2' F modified and the 5' end being phosphorylated.

[001060] In some embodiments, at least 30% of the nucleotides in the sdRNA or sd-rxRNA are modified. In some embodiments, at least 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% of the nucleotides in the sdRNA or sd-rxRNA are modified. In some embodiments, 100% of the nucleotides in the sdRNA or sd-rxRNA are modified.

[001061] In some embodiments, the sdRNA molecules have minimal double stranded regions. In some embodiments the region of the molecule that is double stranded ranges from

8-15 nucleotides long. In some embodiments, the region of the molecule that is double stranded is 8, 9, 10, 11, 12, 13, 14 or 15 nucleotides long. In some embodiments the double stranded region is 13 nucleotides long. There can be 100% complementarity between the guide and passenger strands, or there may be one or more mismatches between the guide and passenger strands. In some embodiments, on one end of the double stranded molecule, the molecule is either blunt-ended or has a one-nucleotide overhang. The single stranded region of the molecule is in some embodiments between 4-12 nucleotides long. In some embodiments, the single stranded region can be 4, 5, 6, 7, 8, 9, 10, 11 or 12 nucleotides long. In some embodiments, the single stranded region can also be less than 4 or greater than 12 nucleotides long. In certain embodiments, the single stranded region is 6 or 7 nucleotides long.

[001062] In some embodiments, the sdRNA molecules have increased stability. In some instances, a chemically modified sdRNA or sd-rxRNA molecule has a half-life in media that is longer than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or more than 24 hours, including any intermediate values. In some embodiments, the sd-rxRNA has a half-life in media that is longer than 12 hours.

[001063] In some embodiments, the sdRNA is optimized for increased potency and/or reduced toxicity. In some embodiments, nucleotide length of the guide and/or passenger strand, and/or the number of phosphorothioate modifications in the guide and/or passenger strand, can in some aspects influence potency of the RNA molecule, while replacing 2'-fluoro (2'F) modifications with 2'-0-methyl (2'OMe) modifications can in some aspects influence toxicity of the molecule. In some embodiments, reduction in 2'F content of a molecule is predicted to reduce toxicity of the molecule. In some embodiments, the number of phosphorothioate modifications in an RNA molecule can influence the uptake of the molecule into a cell, for example the efficiency of passive uptake of the molecule into a cell. In some embodiments, the sdRNA has no 2'F modification and yet are characterized by equal efficacy in cellular uptake and tissue penetration.

[001064] In some embodiments, a guide strand is approximately 18-19 nucleotides in length and has approximately 2-14 phosphate modifications. For example, a guide strand can contain 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or more than 14 nucleotides that are phosphate-modified. The guide strand may contain one or more modifications that confer increased stability without interfering with RISC entry. The phosphate modified nucleotides, such as phosphorothioate modified nucleotides, can be at the 3' end, 5' end or spread throughout the

guide strand. In some embodiments, the 3' terminal 10 nucleotides of the guide strand contain 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 phosphorothioate modified nucleotides. The guide strand can also contain 2'F and/or 2'OMe modifications, which can be located throughout the molecule. In some embodiments, the nucleotide in position one of the guide strand (the nucleotide in the most 5' position of the guide strand) is 2'OMe modified and/or phosphorylated. C and U nucleotides within the guide strand can be 2'F modified. For example, C and U nucleotides in positions 2-10 of a 19 nt guide strand (or corresponding positions in a guide strand of a different length) can be 2'F modified. C and U nucleotides within the guide strand can also be 2'OMe modified. For example, C and U nucleotides in positions 11-18 of a l9 nt guide strand (or corresponding positions in a guide strand of a different length) can be 2'OMe modified. In some embodiments, the nucleotide at the most 3' end of the guide strand is unmodified. In certain embodiments, the majority of Cs and Us within the guide strand are 2'F modified and the 5' end of the guide strand is phosphorylated. In other embodiments, position 1 and the Cs or Us in positions 11-18 are 2'OMe modified and the 5' end of the guide strand is phosphorylated. In other embodiments, position 1 and the Cs or Us in positions 11-18 are 2'OMe modified, the 5' end of the guide strand is phosphorylated, and the Cs or Us in position 2-10 are 2'F modified.

[001065] The self-deliverable RNAi technology provides a method of directly transfecting cells with the RNAi agent, without the need for additional formulations or techniques. The ability to transfect hard-to-transfect cell lines, high *in vivo* activity, and simplicity of use, are characteristics of the compositions and methods that present significant functional advantages over traditional siRNA-based techniques, and as such, the sdRNA methods are employed in several embodiments related to the methods of reduction in expression of the target gene in the TILs of the present invention. The sdRNAi methods allows direct delivery of chemically synthesized compounds to a wide range of primary cells and tissues, both *ex-vivo* and *in vivo*. The sdRNAs described in some embodiments of the invention herein are commercially available from Advirna LLC, Worcester, MA, USA.

[001066] The sdRNA are formed as hydrophobically-modified siRNA-antisense oligonucleotide hybrid structures, and are disclosed, for example in Byrne *et al.*, December 2013, *J. Ocular Pharmacology and Therapeutics*, 29(10): 855-864, incorporated by reference herein in its entirety.

[001067] In some embodiments, the sdRNA oligonucleotides can be delivered to the TILs described herein using sterile electroporation. In certain embodiments, the method comprises sterile electroporation of a population of TILs to deliver sdRNA oligonucleotides.

[001068] In some embodiments, the oligonucleotides can be delivered to the cells in combination with a transmembrane delivery system. In some embodimets, this transmembrane delivery system comprises lipids, viral vectors, and the like. In some embodiments, the oligonucleotide agent is a self-delivery RNAi agent, that does not require any delivery agents. In certain embodiments, the method comprises use of a transmembrane delivery system to deliver sdRNA oligonucleotides to a population of TILs.

[001069] Oligonucleotides and oligonucleotide compositions are contacted with (*e.g.*, brought into contact with, also referred to herein as administered or delivered to) and taken up by TILs described herein, including through passive uptake by TILs. The sdRNA can be added to the TILs as described herein during the first expansion, for example Figure 7 and/or Figure 85, Step B, after the first expansion, for example, during Step C, before or during the second expansion, for example before or during Step D, after Step D and before harvest in Step E, during or after harvest in Step F, before or during final formulation and/or transfer to infusion Bag in Step F, as well as before any optional cryopreservation step in Step F. Mroeover, sdRNA can be added after thawing from any cryopreservation step in Step F. In an embodiment, one or more sdRNAs targeting genes as described herein, including PD-1, LAG-3, TIM-3, CISH, and CBLB, may be added to cell culture media comprising TILs and other agents at concentrations selected from the group consisting of 100 nM to 20 mM, 200 nM to 10 mM, 500 nm to 1 mM, 1 μM to 100 μM, and 1 μM to 100 μM. In an embodiment, one or more sdRNAs targeting genes as described herein, including PD-1, LAG-3, TIM-3, CISH, and CBLB, may be added to cell culture media comprising TILs and other agents at amounts selected from the group consisting of 0.1 μM sdRNA/10,000 TILs/100 μL media, 0.5 μM sdRNA/10,000 TILs /100 μL media, 0.75 μM sdRNA/10,000 TILs /100 μL media, 1 μM sdRNA/10,000 TILs /100 μL media, 1.25 μM sdRNA/10,000 TILs /100 μL media, 1.5 μM sdRNA/10,000 TILs /100 μL media, 2 μM sdRNA/10,000 TILs /100 μL media, 5 μM sdRNA/10,000 TILs /100 μL media, or 10 μM sdRNA/10,000 TILs /100 μL media. In an embodiment, one or more sdRNAs targeting genes as described herein, including PD-1, LAG-3, TIM-3, CISH, and CBLB, may be added to TIL cultures during the pre-REP or REP stages twice a day, once a day, every two days, every three days, every four days, every five days, every six days, or every seven days.

[001070] Oligonucleotide compositions of the invention, including sdRNA, can be contacted with TILs as described herein during the expansion process, for example by dissolving sdRNA at high concentrations in cell culture media and allowing sufficient time for passive uptake to occur. In certain embodiments, the method of the present invention comprises contacting a population of TILs with an oligonucleotide composition as described herein. In certain embodiments, the method comprises dissolving an oligonucleotide *e.g.* sdRNA in a cell culture media and contacting the cell culture media with a population of TILs. The TILs may be a first population, a second population and/or a third population as described herein.

[001071] In some embodiments, delivery of oligonucleotides into cells can be enhanced by suitable art recognized methods including calcium phosphate, DMSO, glycerol or dextran, electroporation, or by transfection, *e.g.*, using cationic, anionic, or neutral lipid compositions or liposomes using methods known in the art (*see*, *e.g.*, WO 90/14074; WO 91/16024; WO 91/17424; U.S. Pat. No. 4,897,355; Bergan et a 1993. Nucleic Acids Research. 21:3567).

[001072] In some embodiments, more than one sdRNA is used to reduce expression of a target gene. In some embodiments, one or more of PD-1, TIM-3, CBLB, LAG3 and/or CISH targeting sdRNAs are used together. In some embodiments, a PD-1 sdRNA is used with one or more of TIM-3, CBLB, LAG3 and/or CISH in order to reduce expression of more than one gene target. In some embodiments, a LAG3 sdRNA is used in combination with a CISH targeting sdRNA to reduce gene expression of both targets. In some embodiments, the sdRNAs targeting one or more of PD-1, TIM-3, CBLB, LAG3 and/or CISH herein are commercially available from Advirna LLC, Worcester, MA, USA.

[001073] In some embodiments, the sdRNA targets a gene selected from the group consisting of PD-1, LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB, BAFF (BR3), and combinations thereof. In some embodiments, the sdRNA targets a gene selected from the group consisting of PD-1, LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB, BAFF (BR3), and combinations thereof. In some embodiments, one sdRNA targets PD-1 and another sdRNA targets a gene selected from the group consisting of LAG3, TIM3, CTLA-4, TIGIT, CISH, TGFβR2, PKA, CBLB, BAFF (BR3), and combinations thereof. In some embodiments, the sdRNA targets a gene selected from PD-1, LAG-3, CISH, CBLB, TIM3, and combinations thereof. In some embodiments, the sdRNA targets a gene selected from PD-1 and one of LAG3, CISH, CBLB, TIM3, and combinations thereof. In some embodiments, one sdRNA targets PD-1 and one sdRNA targets LAG3. In some embodiments, one sdRNA targets PD-1 and one sdRNA targets CISH. In some

embodiments, one sdRNA targets PD-1 and one sdRNA targets CBLB. In some embodiments, one sdRNA targets LAG3 and one sdRNA targets CISH. In some embodiments, one sdRNA targets LAG3 and one sdRNA targets CBLB. In some embodiments, one sdRNA targets CISH and one sdRNA targets CBLB. In some embodiments, one sdRNA targets TIM3 and one sdRNA targets PD-1. In some embodiments, one sdRNA targets TIM3 and one sdRNA targets LAG3. In some embodiments, one sdRNA targets TIM3 and one sdRNA targets CISH. In some embodiments, one sdRNA targets TIM3 and one sdRNA targets CBLB.

[001074] As discussed above, embodiments of the present invention provide tumor infiltrating lymphocytes (TILs) that have been genetically modified via gene-editing to enhance their therapeutic effect. Embodiments of the present invention embrace genetic editing through nucleotide insertion (RNA or DNA) into a population of TILs for both promotion of the expression of one or more proteins and inhibition of the expression of one or more proteins, as well as combinations thereof. Embodiments of the present invention also provide methods for expanding TILs into a therapeutic population, wherein the methods comprise gene-editing the TILs. There are several gene-editing technologies that may be used to genetically modify a population of TILs, which are suitable for use in accordance with the present invention.

[001075] In some embodiments, the method comprises a method of genetically modifying a population of TILs which include the step of stable incorporation of genes for production of one or more proteins. In an embodiment, a method of genetically modifying a population of TILs includes the step of retroviral transduction. In an embodiment, a method of genetically modifying a population of TILs includes the step of lentiviral transduction. Lentiviral transduction systems are known in the art and are described, *e.g.*, in Levine, *et al.*, *Proc. Nat'l Acad. Sci.* **2006,** *103,* 17372-77; Zufferey, *et al., Nat. Biotechnol.* **1997,** *15,* 871-75; Dull, *et al.*, *J. Virology* **1998,** *72*, 8463-71, and U.S. Patent No. 6,627,442, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of genetically modifying a population of TILs includes the step of gamma-retroviral transduction. Gamma-retroviral transduction systems are known in the art and are described, *e.g.*, Cepko and Pear, *Cur. Prot. Mol. Biol.* **1996,** 9.9.1-9.9.16, the disclosure of which is incorporated by reference herein. In an embodiment, a method of genetically modifying a population of TILs includes the step of transposon-mediated gene transfer. Transposon-mediated gene transfer systems are known in the art and include systems wherein the transposase is provided as DNA

expression vector or as an expressible RNA or a protein such that long-term expression of the transposase does not occur in the transgenic cells, for example, a transposase provided as an mRNA (*e.g.*, an mRNA comprising a cap and poly-A tail). Suitable transposon-mediated gene transfer systems, including the salmonid-type Tel-like transposase (SB or Sleeping Beauty transposase), such as SB10, SB11, and SB100x, and engineered enzymes with increased enzymatic activity, are described in, *e.g.*, Hackett, *et al.*, *Mol. Therapy* **2010,** *18,* 674-83 and U.S. Patent No. 6,489,458, the disclosures of each of which are incorporated by reference herein.

[001076] In an embodiment, the method comprises a method of genetically modifying a population of TILs *e.g.* a first population, a second population and/or a third population as described herein. In an embodiment, a method of genetically modifying a population of TILs includes the step of stable incorporation of genes for production or inhibition (*e.g.*, silencing) of one ore more proteins. In an embodiment, a method of genetically modifying a population of TILs includes the step of electroporation. Electroporation methods are known in the art and are described, *e.g.*, in Tsong, *Biophys. J.* **1991,** *60,* 297-306, and U.S. Patent Application Publication No. 2014/0227237 A1, the disclosures of each of which are incorporated by reference herein. Other electroporation methods known in the art, such as those described in U.S. Patent Nos. 5,019,034; 5,128,257; 5,137,817; 5,173,158; 5,232,856; 5,273,525; 5,304,120; 5,318,514; 6,010,613 and 6,078,490, the disclosures of which are incorporated by reference herein, may be used. In an embodiment, the electroporation method is a sterile electroporation method. In an embodiment, the electroporation method is a pulsed electroporation method. In an embodiment, the electroporation method is a pulsed electroporation method comprising the steps of treating TILs with pulsed electrical fields to alter, manipulate, or cause defined and controlled, permanent or temporary changes in the TILs, comprising the step of applying a sequence of at least three single, operator-controlled, independently programmed, DC electrical pulses, having field strengths equal to or greater than 100 V/cm, to the TILs, wherein the sequence of at least three DC electrical pulses has one, two, or three of the following characteristics: (1) at least two of the at least three pulses differ from each other in pulse amplitude; (2) at least two of the at least three pulses differ from each other in pulse width; and (3) a first pulse interval for a first set of two of the at least three pulses is different from a second pulse interval for a second set of two of the at least three pulses. In an embodiment, the electroporation method is a pulsed electroporation method comprising the steps of treating TILs with pulsed electrical fields to alter, manipulate,

or cause defined and controlled, permanent or temporary changes in the TILs, comprising the step of applying a sequence of at least three single, operator-controlled, independently programmed, DC electrical pulses, having field strengths equal to or greater than 100 V/cm, to the TILs, wherein at least two of the at least three pulses differ from each other in pulse amplitude. In an embodiment, the electroporation method is a pulsed electroporation method comprising the steps of treating TILs with pulsed electrical fields to alter, manipulate, or cause defined and controlled, permanent or temporary changes in the TILs, comprising the step of applying a sequence of at least three single, operator-controlled, independently programmed, DC electrical pulses, having field strengths equal to or greater than 100 V/cm, to the TILs, wherein at least two of the at least three pulses differ from each other in pulse width. In an embodiment, the electroporation method is a pulsed electroporation method comprising the steps of treating TILs with pulsed electrical fields to alter, manipulate, or cause defined and controlled, permanent or temporary changes in the TILs, comprising the step of applying a sequence of at least three single, operator-controlled, independently programmed, DC electrical pulses, having field strengths equal to or greater than 100 V/cm, to the TILs, wherein a first pulse interval for a first set of two of the at least three pulses is different from a second pulse interval for a second set of two of the at least three pulses. In an embodiment, the electroporation method is a pulsed electroporation method comprising the steps of treating TILs with pulsed electrical fields to induce pore formation in the TILs, comprising the step of applying a sequence of at least three DC electrical pulses, having field strengths equal to or greater than 100 V/cm, to TILs, wherein the sequence of at least three DC electrical pulses has one, two, or three of the following characteristics: (1) at least two of the at least three pulses differ from each other in pulse amplitude; (2) at least two of the at least three pulses differ from each other in pulse width; and (3) a first pulse interval for a first set of two of the at least three pulses is different from a second pulse interval for a second set of two of the at least three pulses, such that induced pores are sustained for a relatively long period of time, and such that viability of the TILs is maintained. In an embodiment, a method of genetically modifying a population of TILs includes the step of calcium phosphate transfection. Calcium phosphate transfection methods (calcium phosphate DNA precipitation, cell surface coating, and endocytosis) are known in the art and are described in Graham and van der Eb, *Virology* **1973,** *52,* 456-467; Wigler, *et al.*, *Proc. Natl. Acad. Sci.* **1979,** *76,* 1373-1376; and Chen and Okayarea, *Mol. Cell. Biol.* **1987,** *7,* 2745-2752; and in U.S. Patent No. 5,593,875, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of genetically modifying a population of TILs includes

the step of liposomal transfection. Liposomal transfection methods, such as methods that employ a 1:1 (w/w) liposome formulation of the cationic lipid *N*-[1-(2,3-dioleyloxy)propyl]-*n,n,n*-trimethylammonium chloride (DOTMA) and dioleoyl phophotidylethanolamine (DOPE) in filtered water, are known in the art and are described in Rose, *et al.*, *Biotechniques* **1991,** *10,* 520-525 and Felgner, *et al.*, *Proc. Natl. Acad. Sci. USA*, **1987,** *84,* 7413-7417 and in U.S. Patent Nos. 5,279,833; 5,908,635; 6,056,938; 6,110,490; 6,534,484; and 7,687,070, the disclosures of each of which are incorporated by reference herein. In an embodiment, a method of genetically modifying a population of TILs includes the step of transfection using methods described in U.S. Patent Nos. 5,766,902; 6,025,337; 6,410,517; 6,475,994; and 7,189,705; the disclosures of each of which are incorporated by reference herein. The TILs may be a first population, a second population and/or a third population of TILs as described herein.

[001077] According to an embodiment, the gene-editing process may comprise the use of a programmable nuclease that mediates the generation of a double-strand or single-strand break at one or more immune checkpoint genes. Such programmable nucleases enable precise genome editing by introducing breaks at specific genomic loci, *i.e.,* they rely on the recognition of a specific DNA sequence within the genome to target a nuclease domain to this location and mediate the generation of a double-strand break at the target sequence. A double-strand break in the DNA subsequently recruits endogenous repair machinery to the break site to mediate genome editing by either non-homologous end-joining (NHEJ) or homology-directed repair (HDR). Thus, the repair of the break can result in the introduction of insertion/deletion mutations that disrupt (*e.g.*, silence, repress, or enhance) the target gene product.

[001078] Major classes of nucleases that have been developed to enable site-specific genomic editing include zinc finger nucleases (ZFNs), transcription activator-like nucleases (TALENs), and CRISPR-associated nucleases (*e.g.*, CRISPR/Cas9). These nuclease systems can be broadly classified into two categories based on their mode of DNA recognition: ZFNs and TALENs achieve specific DNA binding via protein-DNA interactions, whereas CRISPR systems, such as Cas9, are targeted to specific DNA sequences by a short RNA guide molecule that base-pairs directly with the target DNA and by protein-DNA interactions. *See, e.g.,* Cox *et al.*, *Nature Medicine,* 2015, Vol. 21, No. 2.

[001079] Non-limiting examples of gene-editing methods that may be used in accordance with TIL expansion methods of the present invention include CRISPR methods, TALE

methods, and ZFN methods, which are described in more detail below. According to an embodiment, a method for expanding TILs into a therapeutic population may be carried out in accordance with any embodiment of the methods described herein (*e.g.*, GEN 3 process) or as described in PCT/US2017/058610, PCT/US2018/012605, or PCT/US2018/012633, wherein the method further comprises gene-editing at least a portion of the TILs by one or more of a CRISPR method, a TALE method or a ZFN method, in order to generate TILs that can provide an enhanced therapeutic effect. According to an embodiment, gene-edited TILs can be evaluated for an improved therapeutic effect by comparing them to non-modified TILs *in vitro, e.g.,* by evaluating *in vitro* effector function, cytokine profiles, etc. compared to unmodified TILs. In certain embodiments, the method comprises gene editing a population of TILs using CRISPR, TALE and/ or ZFN methods.

[001080] In some embodiments of the present invention, electroporation is used for delivery of a gene editing system, such as CRISPR, TALEN, and ZFN systems. In some embodiments of the present invention, the electroporation system is a flow electroporation system. An example of a suitable flow electroporation system suitable for use with some embodiments of the present invention is the commercially-available MaxCyte STX system. There are several alternative commercially-available electroporation instruments which may be suitable for use with the present invention, such as the AgilePulse system or ECM 830 available from BTX-Harvard Apparatus, Cellaxess Elektra (Cellectricon), Nucleofector (Lonza/Amaxa), GenePulser MXcell (BIORAD), iPorator-96 (Primax) or siPORTer96 (Ambion). In some embodiments of the present invention, the electroporation system forms a closed, sterile system with the remainder of the TIL expansion method. In some embodiments of the present invention, the electroporation system is a pulsed electroporation system as described herein, and forms a closed, sterile system with the remainder of the TIL expansion method.

[001081] A method for expanding TILs into a therapeutic population may be carried out in accordance with any embodiment of the methods described herein (*e.g.*, process GEN 3) or as described in PCT/US2017/058610, PCT/US2018/012605, or PCT/US2018/012633, wherein the method further comprises gene-editing at least a portion of the TILs by a CRISPR method (*e.g.,* CRISPR/Cas9 or CRISPR/Cpf1). According to particular embodiments, the use of a CRISPR method during the TIL expansion process causes expression of one or more immune checkpoint genes to be silenced or reduced in at least a portion of the therapeutic population of TILs. Alternatively, the use of a CRISPR method during the TIL expansion process

causes expression of one or more immune checkpoint genes to be enhanced in at least a portion of the therapeutic population of TILs.

[001082] CRISPR stands for "Clustered Regularly Interspaced Short Palindromic Repeats." A method of using a CRISPR system for gene editing is also referred to herein as a CRISPR method. There are three types of CRISPR systems which incorporate RNAs and Cas proteins, and which may be used in accordance with the present invention: Types I, II, and III. The Type II CRISPR (exemplified by Cas9) is one of the most well-characterized systems.

[001083] CRISPR technology was adapted from the natural defense mechanisms of bacteria and archaea (the domain of single-celled microorganisms). These organisms use CRISPR-derived RNA and various Cas proteins, including Cas9, to foil attacks by viruses and other foreign bodies by chopping up and destroying the DNA of a foreign invader. A CRISPR is a specialized region of DNA with two distinct characteristics: the presence of nucleotide repeats and spacers. Repeated sequences of nucleotides are distributed throughout a CRISPR region with short segments of foreign DNA (spacers) interspersed among the repeated sequences. In the type II CRISPR/Cas system, spacers are integrated within the CRISPR genomic loci and transcribed and processed into short CRISPR RNA (crRNA). These crRNAs anneal to trans-activating crRNAs (tracrRNAs) and direct sequence-specific cleavage and silencing of pathogenic DNA by Cas proteins. Target recognition by the Cas9 protein requires a "seed" sequence within the crRNA and a conserved dinucleotide-containing protospacer adjacent motif (PAM) sequence upstream of the crRNA-binding region. The CRISPR/Cas system can thereby be retargeted to cleave virtually any DNA sequence by redesigning the crRNA. The crRNA and tracrRNA in the native system can be simplified into a single guide RNA (sgRNA) of approximately 100 nucleotides for use in genetic engineering. The CRISPR/Cas system is directly portable to human cells by co-delivery of plasmids expressing the Cas9 endo-nuclease and the necessary crRNA components. Different variants of Cas proteins may be used to reduce targeting limitations (*e.g.*, orthologs of Cas9, such as Cpf1).

[001084] Non-limiting examples of genes that may be silenced or inhibited by permanently gene-editing TILs via a CRISPR method include PD-1, CTLA-4, LAG-3, HAVCR2 (TIM-3), Cish, TGFβ, PKA, CBL-B, PPP2CA, PPP2CB, PTPN6, PTPN22, PDCD1, BTLA, CD160, TIGIT, CD96, CRTAM, LAIR1, SIGLEC7, SIGLEC9, CD244, TNFRSF10B, TNFRSF10A, CASP8, CASP10, CASP3, CASP6, CASP7, FADD, FAS, SMAD2, SMAD3, SMAD4,

SMAD10, SKI, SKIL, TGIF1, IL10RA, IL10RB, HMOX2, IL6R, IL6ST, EIF2AK4, CSK, PAG1, SIT1, FOXP3, PRDM1, BATF, GUCY1A2, GUCY1A3, GUCY1B2, and GUCY1B3.

[001085] Non-limiting examples of genes that may be enhanced by permanently gene-editing TILs via a CRISPR method include CCR2, CCR4, CCR5, CXCR2, CXCR3, CX3CR1, IL-2, IL12, IL-15, and IL-21.

[001086] Examples of systems, methods, and compositions for altering the expression of a target gene sequence by a CRISPR method, and which may be used in accordance with embodiments of the present invention, are described in U.S. Patent Nos. 8,697,359; 8,993,233; 8,795,965; 8,771,945; 8,889,356; 8,865,406; 8,999,641; 8,945,839; 8,932,814; 8,871,445; 8,906,616; and 8,895,308, which are incorporated by reference herein. Resources for carrying out CRISPR methods, such as plasmids for expressing CRISPR/Cas9 and CRISPR/Cpf1, are commercially available from companies such as GenScript.

[001087] In an embodiment, genetic modifications of populations of TILs, as described herein, may be performed using the CRISPR/Cpf1 system as described in U.S. Patent No. US 9790490, the disclosure of which is incorporated by reference herein.

[001088] A method for expanding TILs into a therapeutic population may be carried out in accordance with any embodiment of the methods described herein (*e.g.*, process 2A) or as described in PCT/US2017/058610, PCT/US2018/012605, or PCT/US2018/012633, wherein the method further comprises gene-editing at least a portion of the TILs by a TALE method. According to particular embodiments, the use of a TALE method during the TIL expansion process causes expression of one or more immune checkpoint genes to be silenced or reduced in at least a portion of the therapeutic population of TILs. Alternatively, the use of a TALE method during the TIL expansion process causes expression of one or more immune checkpoint genes to be enhanced in at least a portion of the therapeutic population of TILs.

[001089] TALE stands for "Transcription Activator-Like Effector" proteins, which include TALENs ("Transcription Activator-Like Effector Nucleases"). A method of using a TALE system for gene editing may also be referred to herein as a TALE method. TALEs are naturally occurring proteins from the plant pathogenic bacteria genus *Xanthomonas*, and contain DNA-binding domains composed of a series of 33–35-amino-acid repeat domains that each recognizes a single base pair. TALE specificity is determined by two hypervariable amino acids that are known as the repeat-variable di-residues (RVDs). Modular TALE repeats are linked together to recognize contiguous DNA sequences. A specific RVD in the

DNA-binding domain recognizes a base in the target locus, providing a structural feature to assemble predictable DNA-binding domains. The DNA binding domains of a TALE are fused to the catalytic domain of a type IIS FokI endonuclease to make a targetable TALE nuclease. To induce site-specific mutation, two individual TALEN arms, separated by a 14-20 base pair spacer region, bring FokI monomers in close proximity to dimerize and produce a targeted double-strand break.

[001090] Several large, systematic studies utilizing various assembly methods have indicated that TALE repeats can be combined to recognize virtually any user-defined sequence. Custom-designed TALE arrays are also commercially available through Cellectis Bioresearch (Paris, France), Transposagen Biopharmaceuticals (Lexington, KY, USA), and Life Technologies (Grand Island, NY, USA). TALE and TALEN methods suitable for use in the present invention are described in U.S. Patent Application Publication Nos. US 2011/0201118 A1; US 2013/0117869 A1; US 2013/0315884 A1; US 2015/0203871 A1 and US 2016/0120906 A1, the disclosures of which are incorporated by reference herein.

[001091] Non-limiting examples of genes that may be silenced or inhibited by permanently gene-editing TILs via a TALE method include PD-1, CTLA-4, LAG-3, HAVCR2 (TIM-3), Cish, TGFβ, PKA, CBL-B, PPP2CA, PPP2CB, PTPN6, PTPN22, PDCD1, BTLA, CD160, TIGIT, CD96, CRTAM, LAIR1, SIGLEC7, SIGLEC9, CD244, TNFRSF10B, TNFRSF10A, CASP8, CASP10, CASP3, CASP6, CASP7, FADD, FAS, SMAD2, SMAD3, SMAD4, SMAD10, SKI, SKIL, TGIF1, IL10RA, IL10RB, HMOX2, IL6R, IL6ST, EIF2AK4, CSK, PAG1, SIT1, FOXP3, PRDM1, BATF, GUCY1A2, GUCY1A3, GUCY1B2, and GUCY1B3.

[001092] Non-limiting examples of genes that may be enhanced by permanently gene-editing TILs via a TALE method include CCR2, CCR4, CCR5, CXCR2, CXCR3, CX3CR1, IL-2, IL12, IL-15, and IL-21.

[001093] Examples of systems, methods, and compositions for altering the expression of a target gene sequence by a TALE method, and which may be used in accordance with embodiments of the present invention, are described in U.S. Patent No. 8,586,526, which is incorporated by reference herein.

[001094] A method for expanding TILs into a therapeutic population may be carried out in accordance with any embodiment of the methods described herein (*e.g.*, process GEN 3) or as described in PCT/US2017/058610, PCT/US2018/012605, or PCT/US2018/012633, wherein the method further comprises gene-editing at least a portion of the TILs by a zinc finger or

zinc finger nuclease method. According to particular embodiments, the use of a zinc finger method during the TIL expansion process causes expression of one or more immune checkpoint genes to be silenced or reduced in at least a portion of the therapeutic population of TILs. Alternatively, the use of a zinc finger method during the TIL expansion process causes expression of one or more immune checkpoint genes to be enhanced in at least a portion of the therapeutic population of TILs.

[001095] An individual zinc finger contains approximately 30 amino acids in a conserved ββα configuration. Several amino acids on the surface of the α-helix typically contact 3 bp in the major groove of DNA, with varying levels of selectivity. Zinc fingers have two protein domains. The first domain is the DNA binding domain, which includes eukaryotic transcription factors and contain the zinc finger. The second domain is the nuclease domain, which includes the FokI restriction enzyme and is responsible for the catalytic cleavage of DNA.

[001096] The DNA-binding domains of individual ZFNs typically contain between three and six individual zinc finger repeats and can each recognize between 9 and 18 base pairs. If the zinc finger domains are specific for their intended target site then even a pair of 3-finger ZFNs that recognize a total of 18 base pairs can, in theory, target a single locus in a mammalian genome. One method to generate new zinc-finger arrays is to combine smaller zinc-finger "modules" of known specificity. The most common modular assembly process involves combining three separate zinc fingers that can each recognize a 3 base pair DNA sequence to generate a 3-finger array that can recognize a 9 base pair target site. Alternatively, selection-based approaches, such as oligomerized pool engineering (OPEN) can be used to select for new zinc-finger arrays from randomized libraries that take into consideration context-dependent interactions between neighboring fingers. Engineered zinc fingers are available commercially; Sangamo Biosciences (Richmond, CA, USA) has developed a propriety platform (CompoZr®) for zinc-finger construction in partnership with Sigma–Aldrich (St. Louis, MO, USA).

[001097] Non-limiting examples of genes that may be silenced or inhibited by permanently gene-editing TILs via a zinc finger method include PD-1, CTLA-4, LAG-3, HAVCR2 (TIM-3), Cish, TGFβ, PKA, CBL-B, PPP2CA, PPP2CB, PTPN6, PTPN22, PDCD1, BTLA, CD160, TIGIT, CD96, CRTAM, LAIR1, SIGLEC7, SIGLEC9, CD244, TNFRSF10B, TNFRSF10A, CASP8, CASP10, CASP3, CASP6, CASP7, FADD, FAS, SMAD2, SMAD3, SMAD4, SMAD10, SKI, SKIL, TGIF1, IL10RA, IL10RB, HMOX2, IL6R, IL6ST,

EIF2AK4, CSK, PAG1, SIT1, FOXP3, PRDM1, BATF, GUCY1A2, GUCY1A3, GUCY1B2, and GUCY1B3.

[001098] Non-limiting examples of genes that may be enhanced by permanently gene-editing TILs via a zinc finger method include CCR2, CCR4, CCR5, CXCR2, CXCR3, CX3CR1, IL-2, IL12, IL-15, and IL-21.

[001099] Examples of systems, methods, and compositions for altering the expression of a target gene sequence by a zinc finger method, which may be used in accordance with embodiments of the present invention, are described in U.S. Patent Nos. 6,534,261, 6,607,882, 6,746,838, 6,794,136, 6,824,978, 6,866,997, 6,933,113, 6,979,539, 7,013,219, 7,030,215, 7,220,719, 7,241,573, 7,241,574, 7,585,849, 7,595,376, 6,903,185, and 6,479,626, which are incorporated by reference herein.

[001100] Other examples of systems, methods, and compositions for altering the expression of a target gene sequence by a zinc finger method, which may be used in accordance with embodiments of the present invention, are described in Beane, *et al.*, *Mol. Therapy*, **2015,** 23 1380-1390, the disclosure of which is incorporated by reference herein.

[001101] In some embodiments, the TILs are optionally genetically engineered to include additional functionalities, including, but not limited to, a high-affinity T cell receptor (TCR), *e.g.*, a TCR targeted at a tumor-associated antigen such as MAGE-1, HER2, or NY-ESO-1, or a chimeric antigen receptor (CAR) which binds to a tumor-associated cell surface molecule (*e.g.*, mesothelin) or lineage-restricted cell surface molecule (*e.g.*, CD19). In certain embodiments, the method comprises genetically engineering a population of TILs to include a high-affinity T cell receptor (TCR), *e.g.*, a TCR targeted at a tumor-associated antigen such as MAGE-1, HER2, or NY-ESO-1, or a chimeric antigen receptor (CAR) which binds to a tumor-associated cell surface molecule (*e.g.*, mesothelin) or lineage-restricted cell surface molecule (*e.g.*, CD19). Aptly, the population of TILs may be a first population, a second population and/or a third population as described herein.

XI. Closed Systems for TIL Manufacturing

[001102] The present invention provides for the use of closed systems during the TIL culturing process, for example, in conjunction with Gen 2 or Gen 3 processes. Such closed systems allow for preventing and/or reducing microbial contamination, allow for the use of

fewer flasks, and allow for cost reductions. In some embodiments, the closed system uses two containers.

[001103] Such closed systems are well-known in the art and can be found, for example, at http://www.fda.gov/cber/guidelines.htm and https://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/ucm076779.htm.

[001104] Sterile connecting devices (STCDs) produce sterile welds between two pieces of compatible tubing. This procedure permits sterile connection of a variety of containers and tube diameters. In some embodiments, the closed systems include luer lock and heat sealed systems as described. In some embodiments, the closed system is accessed via syringes under sterile conditions in order to maintain the sterility and closed nature of the system. In some embodiments, a closed system as described hereinis employed. In some embodiments, the TILs are formulated into a final product formulation container according to the method described herein.

[001105] In some embodiments, the closed system uses one container from the time the tumor fragments are obtained until the TILs are ready for administration to the patient or cryopreserving. In some embodiments when two containers are used, the first container is a closed G-container and the population of TILs is centrifuged and transferred to an infusion bag without opening the first closed G-container. In some embodiments, when two containers are used, the infusion bag is a HypoThermosol-containing infusion bag. A closed system or closed TIL cell culture system is characterized in that once the tumor sample and/or tumor fragments have been added, the system is tightly sealed from the outside to form a closed environment free from the invasion of bacteria, fungi, and/or any other microbial contamination.

[001106] In some embodiments, the reduction in microbial contamination is between about 5% and about 100%. In some embodiments, the reduction in microbial contamination is between about 5% and about 95%. In some embodiments, the reduction in microbial contamination is between about 5% and about 90%. In some embodiments, the reduction in microbial contamination is between about 10% and about 90%. In some embodiments, the reduction in microbial contamination is between about 15% and about 85%. In some embodiments, the reduction in microbial contamination is about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about

55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 97%, about 98%, about 99%, or about 100%.

[001107] The closed system allows for TIL growth in the absence and/or with a significant reduction in microbial contamination.

[001108] Moreover, pH, carbon dioxide partial pressure and oxygen partial pressure of the TIL cell culture environment each vary as the cells are cultured. Consequently, even though a medium appropriate for cell culture is circulated, the closed environment still needs to be constantly maintained as an optimal environment for TIL proliferation. To this end, it is desirable that the physical factors of pH, carbon dioxide partial pressure and oxygen partial pressure within the culture liquid of the closed environment be monitored by means of a sensor, the signal whereof is used to control a gas exchanger installed at the inlet of the culture environment, and the that gas partial pressure of the closed environment be adjusted in real time according to changes in the culture liquid so as to optimize the cell culture environment. In some embodiments, the present invention provides a closed cell culture system which incorporates at the inlet to the closed environment a gas exchanger equipped with a monitoring device which measures the pH, carbon dioxide partial pressure and oxygen partial pressure of the closed environment, and optimizes the cell culture environment by automatically adjusting gas concentrations based on signals from the monitoring device.

[001109] In some embodiments, the pressure within the closed environment is continuously or intermittently controlled. That is, the pressure in the closed environment can be varied by means of a pressure maintenance device for example, thus ensuring that the space is suitable for growth of TILs in a positive pressure state, or promoting exudation of fluid in a negative pressure state and thus promoting cell proliferation. By applying negative pressure intermittently, moreover, it is possible to uniformly and efficiently replace the circulating liquid in the closed environment by means of a temporary shrinkage in the volume of the closed environment.

[001110] In some embodiments, optimal culture components for proliferation of the TILs can be substituted or added, and including factors such as IL-2 and/or OKT3, as well as combination, can be added.

XII. Optional Cryopreservation of TILs

[001111] Either the bulk TIL population (for example the second population of TILs) or the expanded population of TILs (for example the third population of TILs) prepared by Gen 2 or Gen 3 processes can be optionally cryopreserved. In some embodiments, cryopreservation occurs on the therapeutic TIL population. In some embodiments, cryopreservation occurs on the TILs harvested after the second expansion. In some embodiments, cryopreservation occurs on the TILs in exemplary Step F of Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, the TILs are cryopreserved in the infusion bag. In some embodiments, the TILs are cryopreserved prior to placement in an infusion bag. In some embodiments, the TILs are cryopreserved and not placed in an infusion bag. In some embodiments, cryopreservation is performed using a cryopreservation medium. In some embodiments, the cryopreservation media contains dimethylsulfoxide (DMSO). This is generally accomplished by putting the TIL population into a freezing solution, *e.g.* 85% complement inactivated AB serum and 15% dimethyl sulfoxide (DMSO). The cells in solution are placed into cryogenic vials and stored for 24 hours at -80 °C, with optional transfer to gaseous nitrogen freezers for cryopreservation. See, Sadeghi, *et al., Acta Oncologica* **2013,** *52,* 978-986.

[001112] When appropriate, the cells are removed from the freezer and thawed in a 37 °C water bath until approximately 4/5 of the solution is thawed. The cells are generally resuspended in complete media and optionally washed one or more times. In some embodiments, the thawed TILs can be counted and assessed for viability as is known in the art.

[001113] In a preferred embodiment, a population of TILs is cryopreserved using CS10 cryopreservation media (CryoStor 10, BioLife Solutions). In a preferred embodiment, a population of TILs is cryopreserved using a cryopreservation media containing dimethylsulfoxide (DMSO). In a preferred embodiment, a population of TILs is cryopreserved using a 1:1 (vol:vol) ratio of CS10 and cell culture media. In a preferred embodiment, a population of TILs is cryopreserved using about a 1:1 (vol:vol) ratio of CS10 and cell culture media, further comprising additional IL-2.

[001114] As discussed above, and exemplified in Steps A through E as provided in Figure 85 (in particular, *e.g.*, Figure 85B), cryopreservation can occur at numerous points throughout the TIL expansion process. In some embodiments, the expanded population of TILs after the second expansion (as provided for example, according to Step D of Figure 85 (in particular,

e.g., Figure 85B)) can be cryopreserved. Cryopreservation can be generally accomplished by placing the TIL population into a freezing solution, *e.g.*, 85% complement inactivated AB serum and 15% dimethyl sulfoxide (DMSO). The cells in solution are placed into cryogenic vials and stored for 24 hours at -80 °C, with optional transfer to gaseous nitrogen freezers for cryopreservation. See Sadeghi, *et al., Acta Oncologica* **2013,** *52,* 978-986. In some embodiments, the TILs are cryopreserved in 5% DMSO. In some embodiments, the TILs are cryopreserved in cell culture media plus 5% DMSO. In some embodiments, the TILs are cryopreserved according to the methods provided herein.

[001115] When appropriate, the cells are removed from the freezer and thawed in a 37 °C water bath until approximately 4/5 of the solution is thawed. The cells are generally resuspended in complete media and optionally washed one or more times. In some embodiments, the thawed TILs can be counted and assessed for viability as is known in the art.

[001116] In some cases, the Step B TIL population can be cryopreserved immediately, using the protocols discussed below. Alternatively, the bulk TIL population can be subjected to Step C and Step D and then cryopreserved after Step D. Similarly, in the case where genetically modified TILs will be used in therapy, the Step B or Step D TIL populations can be subjected to genetic modifications for suitable treatments.

XIII. Phenotypic Characteristics of Expanded TILs

[001117] In some embodiments, the TILs are analyzed for expression of numerous phenotype markers after expansion, including those described herein and in the Examples. In an embodiment, expression of one or more phenotypic markers is examined. In some embodiments, the phenotypic characteristics of the TILs are analyzed after the first expansion in Step B. In some embodiments, the phenotypic characteristics of the TILs are analyzed during the transition in Step C. In some embodiments, the phenotypic characteristics of the TILs are analyzed during the transition according to Step C and after cryopreservation. In some embodiments, the phenotypic characteristics of the TILs are analyzed after the second expansion according to Step D. In some embodiments, the phenotypic characteristics of the TILs are analyzed after two or more expansions according to Step D.

[001118] In some embodiments, the marker is selected from the group consisting of CD8 and CD28. In some embodiments, expression of CD8 is examined. In some embodiments, expression of CD28 is examined. In some embodiments, the expression of CD8 and/or CD28

is higher on TILs produced according the current invention process, as compared to other processes (*e.g.*, the Gen 3 process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85B), as compared to the 2A process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85B)). In some embodiments, the expression of CD8 is higher on TILs produced according the current invention process, as compared to other processes (*e.g.*, the Gen 3 process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85B), as compared to the 2A process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85B)). In some embodiments, the expression of CD28 is higher on TILs produced according the current invention process, as compared to other processes (*e.g.*, the Gen 3 process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85B), as compared to the 2A process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85A)). In an embodiment, expression of one or more regulatory markers is measured.

[001119] In an embodiment, no selection of the first population of TILs, second population of TILs, third population of TILs, or harvested TIL population based on CD8 and/or CD28 expression is performed during any of the steps for the method for expanding tumor infiltrating lymphocytes (TILs) described herein.

[001120] In some embodiments, the percentage of central memory cells is higher on TILs produced according the current invention process, as compared to other processes (*e.g.*, the Gen 3 process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85B), as compared to the 2A process as provided for example in Figure 85 (in particular, *e.g.*, Figure 85A)). In some embodiments the memory marker for central memory cells is selected from the group consisting of CCR7 and CD62L.

[001121] In an embodiment, restimulated TILs can also be evaluated for cytokine release, using cytokine release assays. In some embodiments, TILs can be evaluated for interferon-γ (IFN-γ) secretion. In some embodiments, the IFN-γ secretion is measured by an ELISA assay. In some embodiments, the IFN-γ secretion is measured by an ELISA assay after the rapid second expansion step, after Step D as provided in for example, Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, TIL health is measured by IFN-gamma (IFN-γ) secretion. In some embodiments, IFN-γ secretion is indicative of active TILs. In some embodiments, a potency assay for IFN-γ production is employed. IFN-γ production is another measure of cytotoxic potential. IFN-γ production can be measured by determining the levels of the cytokine IFN-γ in the media of TIL stimulated with antibodies to CD3, CD28, and CD137/4-1BB. IFN-γ levels in media from these stimulated TIL can be determined using by

measuring IFN-γ release. In some embodiments, an increase in IFN-γ production in for example Step D in the Gen 3 process as provided in Figure 85 (in particular, *e.g.*, Figure 85B) TILs as compared to for example Step D in the 2A process as provided in Figure 85 (in particular, *e.g.*, Figure 85A) is indicative of an increase in cytotoxic potential of the Step D TILs. In some embodiments, IFN-γ secretion is increased one-fold, two-fold, three-fold, four-fold, or five-fold or more. In some embodiments, IFN-γ secretion is increased one-fold. In some embodiments, IFN-γ secretion is increased two-fold. In some embodiments, IFN-γ secretion is increased three-fold. In some embodiments, IFN-γ secretion is increased four-fold. In some embodiments, IFN-γ secretion is increased five-fold. In some embodiments, IFN-γ is measured using a Quantikine ELISA kit. In some embodiments, IFN-γ is measured in TILs *ex vivo*. In some embodiments, IFN-γ is measured in TILs *ex vivo*, including TILs produced by the methods of the present invention, including, for example Figure 85B methods.

[001122] The diverse antigen receptors of T and B lymphocytes are produced by somatic recombination of a limited, but large number of gene segments. These gene segments: V (variable), D (diversity), J (joining), and C (constant), determine the binding specificity and downstream applications of immunoglobulins and T-cell receptors (TCRs). The present invention provides a method for generating TILs which exhibit and increase the T-cell repertoire diversity. In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity. In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity as compared to freshly harvested TILs and/or TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, the TILs obtained by the present method exhibit an increase in the T-cell repertoire diversity as compared to freshly harvested TILs and/or TILs prepared using methods referred to as process 2A, as exemplified in Figure 85 (in particular, *e.g.*, Figure 85A). In some embodiments, the TILs obtained in the first expansion exhibit an increase in the T-cell repertoire diversity. In some embodiments, the increase in diversity is an increase in the immunoglobulin diversity and/or the T-cell receptor diversity. In some embodiments, the diversity is in the immunoglobulin is in the immunoglobulin heavy chain. In some embodiments, the diversity is in the immunoglobulin is in the immunoglobulin light chain. In some embodiments, the diversity is in the T-cell receptor. In some embodiments, the diversity is in one of the T-cell receptors selected from the group consisting of alpha, beta,

gamma, and delta receptors. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) alpha and/or beta. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) alpha. In some embodiments, there is an increase in the expression of T-cell receptor (TCR) beta. In some embodiments, there is an increase in the expression of TCRab (*i.e.*, TCRα/β). In some embodiments, the process as described herein (*e.g.*, the Gen 3 process) shows higher clonal diversity as compared to other processes, for example the process referred to as the Gen 2 based on the number of unique peptide CDRs within the sample (*see*, for example Figures 12-14).

[001123] In some embodiments, the phenotypic characterization is examined after cryopreservation.

XIV. Additional Process Embodiments

[001124] In some embodiments, the invention provides a method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising: (a) obtaining a first population of TILs from one or more tumor fragments or cores from a subject; (b) performing a priming first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 and OKT-3, wherein the priming first expansion is performed for about 1 to 17 days to obtain the second population of TILs, wherein the second population of TILs is greater in number than the first population of TILs; (c) performing a rapid second expansion by contacting the second population of TILs with a cell culture medium comprising IL-2, OKT-3 and exogenous antigen presenting cells (APCs) to produce a third population of TILs, wherein the rapid second expansion is performed for about 1 to 11 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and (d) harvesting the therapeutic population of TILs obtained from step (c). In some embodiments, the step of rapid second expansion is split into a plurality of steps to achieve a scaling up of the culture by: (1) performing the rapid second expansion by culturing the second population of TILs in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (2) effecting the transfer of the second population of TILs from the small scale culture to a second container larger than the first container, *e.g.*, a G-REX 500MCS container, wherein in the second container the second population of TILs from the small scale culture is cultured in a larger scale culture for a period of about 4 to 7 days. In some embodiments, the step of rapid expansion is split into a plurality of steps to achieve a scaling out of the culture by: (1)

performing the rapid second expansion by culturing the second population of TILs in a first small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (2) effecting the transfer and apportioning of the second population of TILs from the first small scale culture into and amongst at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 second containers that are equal in size to the first container, wherein in each second container the portion of the second population of TILs from the first small scale culture transferred to such second container is cultured in a second small scale culture for a period of about 4 to 7 days. In some embodiments, the step of rapid expansion is split into a plurality of steps to achieve a scaling out and scaling up of the culture by: (1) performing the rapid second expansion by culturing the second population of TILs in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (2) effecting the transfer and apportioning of the second population of TILs from the first small scale culture into and amongst at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 second containers that are larger in size than the first container, *e.g.*, G-REX 500MCS containers, wherein in each second container the portion of the second population of TILs transferred from the small scale culture to such second container is cultured in a larger scale culture for a period of about 4 to 7 days. In some embodiments, the step of rapid expansion is split into a plurality of steps to achieve a scaling out and scaling up of the culture by: (1) performing the rapid second expansion by culturing the second population of TILs in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 4 days, and then (2) effecting the transfer and apportioning of the second population of TILs from the first small scale culture into and amongst 2, 3 or 4 second containers that are larger in size than the first container, *e.g.*, G-REX 500MCS containers, wherein in each second container the portion of the second population of TILs transferred from the small scale culture to such second container is cultured in a larger scale culture for a period of about 5 days.

[001125] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by contacting the first population of TILs with a culture medium which further comprises exogneous antigen-presenting cells (APCs), wherein the number of APCs in the culture medium in step (c) is greater than the number of APCs in the culture medium in step (b).

[001126] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the culture medium is supplemented with additional exogenous APCs.

[001127] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 20:1.

[001128] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 10:1.

[001129] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 9:1.

[001130] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 8:1.

[001131] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 7:1.

[001132] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 6:1.

[001133] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 5:1.

[001134] In another embodiment, the invention provides the method described in any one of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 4:1.

[001135] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 3:1.

[001136] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.9:1.

[001137] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.8:1.

[001138] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.7:1.

[001139] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.6:1.

[001140] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.5:1.

[001141] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.4:1.

[001142] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.3:1.

[001143] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.2:1.

[001144] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2.1:1.

[001145] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 1.1:1 to at or about 2:1.

[001146] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 10:1.

[001147] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 5:1.

[001148] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 4:1.

[001149] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 3:1.

[001150] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.9:1.

[001151] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.8:1.

[001152] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.7:1.

[001153] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.6:1.

[001154] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.5:1.

[001155] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.4:1.

[001156] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.3:1.

[001157] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about about 2:1 to at or about 2.2:1.

[001158] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is selected from a range of from at or about 2:1 to at or about 2.1:1.

[001159] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is at or about 2:1.

[001160] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of number of APCs added in the rapid second expansion to the number of APCs added in step (b) is at or about 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1, 2:1, 2.1:1, 2.2:1, 2.3:1, 2.4:1, 2.5:1, 2.6:1, 2.7:1, 2.8:1, 2.9:1, 3:1, 3.1:1, 3.2:1, 3.3:1, 3.4:1, 3.5:1, 3.6:1, 3.7:1, 3.8:1, 3.9:1, 4:1, 4.1:1, 4.2:1, 4.3:1, 4.4:1, 4.5:1, 4.6:1, 4.7:1, 4.8:1, 4.9:1, or 5:1.

[001161] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the number of APCs added in the primary first expansion is at or about 1×10^8, 1.1×10^8, 1.2×10^8, 1.3×10^8, 1.4×10^8, 1.5×10^8, 1.6×10^8, 1.7×10^8, 1.8×10^8, 1.9×10^8, 2×10^8, 2.1×10^8, 2.2×10^8, 2.3×10^8, 2.4×10^8, 2.5×10^8, 2.6×10^8, 2.7×10^8, 2.8×10^8, 2.9×10^8, 3×10^8, 3.1×10^8, 3.2×10^8, 3.3×10^8, 3.4×10^8 or 3.5×10^8 APCs, and such that the number of APCs added in the rapid second expansion is at or about 3.5×10^8, 3.6×10^8, 3.7×10^8, 3.8×10^8, 3.9×10^8, 4×10^8, 4.1×10^8, 4.2×10^8, 4.3×10^8, 4.4×10^8, 4.5×10^8, 4.6×10^8, 4.7×10^8, 4.8×10^8, 4.9×10^8, 5×10^8, 5.1×10^8, 5.2×10^8, 5.3×10^8, 5.4×10^8, 5.5×10^8, 5.6×10^8, 5.7×10^8, 5.8×10^8, 5.9×10^8, 6×10^8, 6.1×10^8, 6.2×10^8, 6.3×10^8, 6.4×10^8, 6.5×10^8, 6.6×10^8, 6.7×10^8, 6.8×10^8, 6.9×10^8, 7×10^8, 7.1×10^8, 7.2×10^8, 7.3×10^8, 7.4×10^8, 7.5×10^8, 7.6×10^8, 7.7×10^8, 7.8×10^8, 7.9×10^8, 8×10^8, 8.1×10^8, 8.2×10^8, 8.3×10^8, 8.4×10^8, 8.5×10^8, 8.6×10^8, 8.7×10^8, 8.8×10^8, 8.9×10^8, 9×10^8, 9.1×10^8, 9.2×10^8, 9.3×10^8, 9.4×10^8, 9.5×10^8, 9.6×10^8, 9.7×10^8, 9.8×10^8, 9.9×10^8 or 1×10^9 APCs.

[001162] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the number of APCs added in the primary first expansion is selected from the range of at or about 1×10^8 APCs to at or

about 3.5×10^8 APCs, and wherein the number of APCs added in the rapid second expansion is selected from the range of at or about 3.5×10^8 APCs to at or about 1×10^9 APCs.

[001163] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the number of APCs added in the primary first expansion is selected from the range of at or about 1.5×10^8 APCs to at or about 3×10^8 APCs, and wherein the number of APCs added in the rapid second expansion is selected from the range of at or about 4×10^8 APCs to at or about 7.5×10^8 APCs.

[001164] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the number of APCs added in the primary first expansion is selected from the range of at or about 2×10^8 APCs to at or about 2.5×10^8 APCs, and wherein the number of APCs added in the rapid second expansion is selected from the range of at or about 4.5×10^8 APCs to at or about 5.5×10^8 APCs.

[001165] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that at or about 2.5×10^8 APCs are added to the primary first expansion and at or about 5×10^8 APCs are added to the rapid second expansion.

[001166] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the antigen-presenting cells are peripheral blood mononuclear cells (PBMCs).

[001167] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple tumor fragments are distributed into a plurality of separate containers, in each of which separate containers the first population of TILs is obtained in step (a), the second population of TILs is obtained in step (b), and the third population of TILs is obtained in step (c), and the therapeutic populations of TILs from the plurality of containers in step (c) are combined to yield the harvested TIL population from step (d).

[001168] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the first population of TILs is obtained from multiple tumor fragments or cores from the subject in step (a).

[001169] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple tumors are evenly distributed into the plurality of separate containers.

[001170] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the plurality of separate containers comprises at least two separate containers.

[001171] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the plurality of separate containers comprises from two to twenty separate containers.

[001172] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the plurality of separate containers comprises from two to fifteen separate containers.

[001173] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the plurality of separate containers comprises from two to ten separate containers.

[001174] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the plurality of separate containers comprises from two to five separate containers.

[001175] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the plurality of separate containers comprises 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 separate containers.

[001176] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that for each container in which the priming first expansion is performed on a first population of TILs in step (b) the rapid second expansion in step (c) is performed in the same container on the second population of TILs produced from such first population of TILs.

[001177] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each of the separate containers comprises a first gas-permeable surface area.

[001178] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple tumor fragments are distributed in a single container.

[001179] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the single container comprises a first gas-permeable surface area.

[001180] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about one cell layer to at or about three cell layers.

[001181] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 1.5 cell layers to at or about 2.5 cell layers.

[001182] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 2 cell layers.

[001183] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3 cell layers.

[001184] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 3 cell layers to at or about 10 cell layers.

[001185] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 4 cell layers to at or about 8 cell layers.

[001186] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered

onto the first gas-permeable surface area at an average thickness of at or about 3, 4, 5, 6, 7, 8, 9 or 10 cell layers.

[001187] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 4, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9 or 8 cell layers.

[001188] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the priming first expansion is performed in a first container comprising a first gas-permeable surface area and in step (c) the rapid second expansion is performed in a second container comprising a second gas-permeable surface area.

[001189] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the second container is larger than the first container.

[001190] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the priming first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about one cell layer to at or about three cell layers.

[001191] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 1.5 cell layers to at or about 2.5 cell layers.

[001192] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 2 cell layers.

[001193] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable modified such that in step (b) the APCs are layered onto

the first gas-permeable surface area at an average thickness of at or about 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3 cell layers.

[001194] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the second gas-permeable surface area at an average thickness of at or about 3 cell layers to at or about 10 cell layers.

[001195] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the second gas-permeable surface area at an average thickness of at or about 4 cell layers to at or about 8 cell layers.

[001196] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the second gas-permeable surface area at an average thickness of at or about 3, 4, 5, 6, 7, 8, 9 or 10 cell layers.

[001197] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable modified such that in step (c) the APCs are layered onto the second gas-permeable surface area at an average thickness of at or about 4, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9 or 8 cell layers.

[001198] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the priming first expansion is performed in a first container comprising a first gas-permeable surface area and in step (c) the rapid second expansion is performed in the first container.

[001199] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about one cell layer to at or about three cell layers.

[001200] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered

onto the first gas-permeable surface area at an average thickness of at or about 1.5 cell layers to at or about 2.5 cell layers.

[001201] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 2 cell layers.

[001202] In another embodiment, the invention provides the method described any of the preceding paragraphs as applicable above modified such that in step (b) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3 cell layers.

[001203] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 3 cell layers to at or about 10 cell layers.

[001204] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 4 cell layers to at or about 8 cell layers.

[001205] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 3, 4, 5, 6, 7, 8, 9 or 10 cell layers.

[001206] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (c) the APCs are layered onto the first gas-permeable surface area at an average thickness of at or about 4, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9 or 8 cell layers.

[001207] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the

average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:10.

[001208] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:9.

[001209] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:8.

[001210] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:7.

[001211] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:6.

[001212] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first

expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:5.

[001213] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:4.

[001214] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:3.

[001215] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.1 to at or about 1:2.

[001216] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the

average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.2 to at or about 1:8.

[001217] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.3 to at or about 1:7.

[001218] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.4 to at or about 1:6.

[001219] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.5 to at or about 1:5.

[001220] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.6 to at or about 1:4.

[001221] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first

expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.7 to at or about 1:3.5.

[001222] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.8 to at or about 1:3.

[001223] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:1.9 to at or about 1:2.5.

[001224] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of APCs layered in step (c) is selected from the range of at or about 1:2.

[001225] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the primary first expansion is performed by supplementing the cell culture medium of the first population of TILs with additional antigen-presenting cells (APCs), wherein the number of APCs added in step (c) is greater than the number of APCs added in step (b), and wherein the ratio of the average number of layers of APCs layered in step (b) to the average number of layers of

APCs layered in step (c) is selected from at or about 1:1.1, 1:1.2, 1:1.3, 1:1.4, 1:1.5, 1:1.6, 1:1.7, 1:1.8, 1:1.9, 1:2, 1:2.1, 1:2.2, 1:2.3, 1:2.4, 1:2.5, 1:2.6, 1:2.7, 1:2.8, 1:2.9, 1:3, 1:3.1, 1:3.2, 1:3.3, 1:3.4, 1:3.5, 1:3.6, 1:3.7, 1:3.8, 1:3.9, 1:4, 1:4.1, 1:4.2, 1:4.3, 1:4.4, 1:4.5, 1:4.6, 1:4.7, 1:4.8, 1:4.9, 1:5, 1:5.1, 1:5.2, 1:5.3, 1:5.4, 1:5.5, 1:5.6, 1:5.7, 1:5.8, 1:5.9, 1:6, 1:6.1, 1:6.2, 1:6.3, 1:6.4, 1:6.5, 1:6.6, 1:6.7, 1:6.8, 1:6.9, 1:7, 1:7.1, 1:7.2, 1:7.3, 1:7.4, 1:7.5, 1:7.6, 1:7.7, 1:7.8, 1:7.9, 1:8, 1:8.1, 1:8.2, 1:8.3, 1:8.4, 1:8.5, 1:8.6, 1:8.7, 1:8.8, 1:8.9, 1:9, 1:9.1, 1:9.2, 1:9.3, 1:9.4, 1:9.5, 1:9.6, 1:9.7, 1:9.8, 1:9.9 or 1:10.

[001226] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the number of TILs in the second population of TILs to the number of TILs in the first population of TILs is at or about 1.5:1 to at or about 100:1.

[001227] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the number of TILs in the second population of TILs to the number of TILs in the first population of TILs is at or about 50:1.

[001228] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the number of TILs in the second population of TILs to the number of TILs in the first population of TILs is at or about 25:1.

[001229] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the number of TILs in the second population of TILs to the number of TILs in the first population of TILs is at or about 20:1.

[001230] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the number of TILs in the second population of TILs to the number of TILs in the first population of TILs is at or about 10:1.

[001231] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the second population of TILs is at least at or about 50-fold greater in number than the first population of TILs.

[001232] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the second population of TILs is

at least at or about 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, 12-, 13-, 14-, 15-, 16-, 17-, 18-, 19-, 20-, 21-, 22-, 23-, 24-, 25-, 26-, 27-, 28-, 29-, 30-, 31-, 32-, 33-, 34-, 35-, 36-, 37-, 38-, 39-, 40-, 41-, 42-, 43-, 44-, 45-, 46-, 47-, 48-, 49- or 50-fold greater in number than the first population of TILs.

[001233] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that at or about 2 days or at or about 3 days after the commencement of the second period in step (c), the cell culture medium is supplemented with additional IL-2.

[001234] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified to further comprise the step of cryopreserving the harvested TIL population in step (d) using a cryopreservation process.

[001235] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified to comprise performing after step (d) the additional step of (e) transferring the harvested TIL population from step (d) to an infusion bag that optionally contains HypoThermosol.

[001236] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified to comprise the step of cryopreserving the infusion bag comprising the harvested TIL population in step (e) using a cryopreservation process.

[001237] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the cryopreservation process is performed using a 1:1 ratio of harvested TIL population to cryopreservation media.

[001238] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the antigen-presenting cells are peripheral blood mononuclear cells (PBMCs).

[001239] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the PBMCs are irradiated and allogeneic.

[001240] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the total number of APCs added to the cell culture in step (b) is 2.5×10^8.

[001241] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the total number of APCs added to the cell culture in step (c) is 5×10^8.

[001242] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the APCs are PBMCs.

[001243] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the PBMCs are irradiated and allogeneic.

[001244] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the antigen-presenting cells are artificial antigen-presenting cells.

[001245] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the harvesting in step (d) is performed using a membrane-based cell processing system.

[001246] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the harvesting in step (d) is performed using a LOVO cell processing system.

[001247] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 5 to at or about 60 cores or fragments per container in step (b).

[001248] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 10 to at or about 60 cores or fragments per container in step (b).

[001249] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 15 to at or about 60 cores or fragments per container in step (b).

[001250] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 20 to at or about 60 cores or fragments per container in step (b).

[001251] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 25 to at or about 60 cores or fragments per container in step (b).

[001252] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 30 to at or about 60 cores or fragments per container in step (b).

[001253] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 35 to at or about 60 cores or fragments per container in step (b).

[001254] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 40 to at or about 60 cores or fragments per container in step (b).

[001255] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 45 to at or about 60 cores or fragments per container in step (b).

[001256] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple fragments comprise at or about 50 to at or about 60 cores or fragments per container in step (b).

[001257] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the multiple cores or fragments comprise at or about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59 or 60 cores or fragment(s) per container in step (b).

[001258] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each cores or fragment has a volume of at or about 1 mm^3.

[001259] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each core or fragment has a volume of at or about 1 mm^3 to at or about 10 mm^3.

[001260] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each core fragment has a volume of at or about 1 mm^3 to at or about 9 mm^3.

[001261] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 8 mm^3.

[001262] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 7 mm^3.

[001263] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 6 mm^3.

[001264] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 5 mm^3.

[001265] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 4 mm^3.

[001266] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 3 mm^3.

[001267] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1 mm^3 to at or about 2 mm^3.

[001268] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each fragment has a volume of at or about 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 mm^3.

[001269] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the cell culture medium is provided in a container that is a G-container or a Xuri cellbag.

[001270] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the IL-2 concentration in the cell culture medium is about 10,000 IU/mL to about 5,000 IU/mL.

[001271] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the IL-2 concentration in the cell culture medium is about 6,000 IU/mL.

[001272] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the cryopreservation media comprises dimethlysulfoxide (DMSO).

[001273] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the cryopreservation media comprises 7% to 10% DMSO.

[001274] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the first period in step (b) is performed within a period of at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days or 17 days.

[001275] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the second period in step (c) is performed within a period of at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days or 11 days.

[001276] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the first period in step (b) and the second period in step (c) are each individually performed within a period of at or about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days,11 days or 12 days.

[001277] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the first period in step (b) and the second period in step (c) are each individually performed within a period of at or about 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, or 12 days.

[001278] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the first period in step (b) and

the second period in step (c) are each individually performed within a period of at or about 7 days.

[001279] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 14 days to at or about 28 days.

[001280] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 15 days to at or about 28 days.

[001281] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 16 days to at or about 28 days.

[001282] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 17 days to at or about 28 days.

[001283] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 18 days to at or about 28 days.

[001284] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 19 days to at or about 28 days.

[001285] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 20 days to at or about 28 days.

[001286] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 21 days to at or about 28 days.

[001287] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 22 days to at or about 28 days.

[001288] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 23 days to at or about 28 days.

[001289] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 24 days to at or about 28 days.

[001290] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 25 days to at or about 28 days.

[001291] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 26 days to at or about 28 days.

[001292] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 27 days to at or about 28 days.

[001293] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 14 days.

[001294] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 15 days.

[001295] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 16 days.

[001296] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 17 days.

[001297] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 18 days.

[001298] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 19 days.

[001299] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 20 days.

[001300] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 21 days.

[001301] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 22 days.

[001302] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 23 days.

[001303] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 24 days.

[001304] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 25 days.

[001305] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 26 days.

[001306] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 27 days.

[001307] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 28 days.

[001308] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 16 days or less.

[001309] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 20 days or less.

[001310] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 24 days or less.

[001311] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that steps (a) through (d) are performed in a total of at or about 28 days or less.

[001312] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the therapeutic population of TILs harvested in step (d) comprises sufficient TILs for a therapeutically effective dosage of the TILs.

[001313] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the number of TILs sufficient for a therapeutically effective dosage is from at or about 2.3×10^{10} to at or about 13.7×10^{10}.

[001314] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the third population of TILs in step (c) provides for increased efficacy, increased interferon-gamma production, and/or increased polyclonality.

[001315] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the third population of TILs in step (c) provides for at least a one-fold to five-fold or more interferon-gamma production as compared to TILs prepared by a process longer than 16 days.

[001316] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the effector T cells and/or central memory T cells obtained from the third population of TILs step (c) exhibit increased

CD8 and CD28 expression relative to effector T cells and/or central memory T cells obtained from the second population of cells step (b).

[001317] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each container recited in the method is a closed container.

[001318] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each container recited in the method is a G-container.

[001319] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each container recited in the method is a GREX-10.

[001320] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each container recited in the method is a GREX-100.

[001321] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that each container recited in the method is a GREX-500.

[001322] In another embodiment, the invention provides the therapeutic population of tumor infiltrating lymphocytes (TILs) made by the method described in any of the preceding paragraphs as applicable above.

[001323] In another embodiment, the invention provides a therapeutic population of tumor infiltrating lymphocytes (TILs) prepared from tumor tissue of a patient, wherein the therapeutic population of TILs provides for increased efficacy, increased interferon-gamma production, and/or increased polyclonality compared to TILs prepared by a process in which the first expansion of TILs is performed without any added antigen-presenting cells (APCs) or OKT3.

[001324] In another embodiment, the invention provides a therapeutic population of tumor infiltrating lymphocytes (TILs) prepared from tumor tissue of a patient, wherein the therapeutic population of TILs provides for increased efficacy, increased interferon-gamma production, and/or increased polyclonality compared to TILs prepared by a process in which the first expansion of TILs is performed without any added antigen-presenting cells (APCs).

[001325] In another embodiment, the invention provides a therapeutic population of tumor infiltrating lymphocytes (TILs) prepared from tumor tissue of a patient, wherein the therapeutic population of TILs provides for increased efficacy, increased interferon-gamma production, and/or increased polyclonality compared to TILs prepared by a process in which the first expansion of TILs is performed without any added OKT3.

[001326] In another embodiment, the invention provides a therapeutic population of tumor infiltrating lymphocytes (TILs) prepared from tumor tissue of a patient, wherein the therapeutic population of TILs provides for increased efficacy, increased interferon-gamma production, and/or increased polyclonality compared to TILs prepared by a process in which the first expansion of TILs is performed with no added antigen-presenting cells (APCs) and no added OKT3.

[001327] In another embodiment, the invention provides a therapeutic population of tumor infiltrating lymphocytes (TILs) prepared from tumor tissue of a patient, wherein the therapeutic population of TILs provides for increased efficacy, increased interferon-gamma production, and/or increased polyclonality compared to TILs prepared by a process by a process longer than 16 days.

[001328] In another embodiment, the invention provides for the therapeutic population of TILs described in any of the preceding paragraphs as applicable above that provides for increased interferon-gamma production.

[001329] In another embodiment, the invention provides for the therapeutic population of TILs described in any of the preceding paragraphs as applicable above that provides for increased polyclonality.

[001330] In another embodiment, the invention provides for the therapeutic population of TILs described in any of the preceding paragraphs as applicable above that provides for increased efficacy.

[001331] In another embodiment, the invention provides for the therapeutic population of TILs described in any of the preceding paragraphs as applicable above modified such that the therapeutic population of TILs is capable of at least one-fold more interferon-gamma production as compared to TILs prepared by a process longer than 16 days.

[001332] In another embodiment, the invention provides for the therapeutic population of TILs described in any of the preceding paragraphs as applicable above modified such that the

therapeutic population of TILs is capable of at least two-fold more interferon-gamma production as compared to TILs prepared by a process longer than 16 days.

[001333] In another embodiment, the invention provides for the therapeutic population of TILs described in any of the preceding paragraphs as applicable above modified such that the therapeutic population of TILs is capable of at least three-fold more interferon-gamma production as compared to TILs prepared by a process longer than 16 days.

[001334] In another embodiment, the invention provides for a therapeutic population of tumor infiltrating lymphocytes (TILs) that is capable of at least one-fold more interferon-gamma production as compared to TILs prepared by a process in which the first expansion of TILs is performed without any added antigen-presenting cells (APCs).

[001335] In another embodiment, the invention provides for a therapeutic population of tumor infiltrating lymphocytes (TILs) that is capable of at least one-fold more interferon-gamma production as compared to TILs prepared by a process in which the first expansion of TILs is performed without any added OKT3.

[001336] In another embodiment, the invention provides for a therapeutic population of TILs that is capable of at least two-fold more interferon-gamma production as compared to TILs prepared by a process in which the first expansion of TILs is performed without any added APCs.

[001337] In another embodiment, the invention provides for a therapeutic population of TILs that is capable of at least two-fold more interferon-gamma production as compared to TILs prepared by a process in which the first expansion of TILs is performed without any added OKT3.

[001338] In another embodiment, the invention provides for a therapeutic population of TILs that is capable of at least three-fold more interferon-gamma production as compared to TILs prepared by a process in which the first expansion of TILs is performed without any added APCs.

[001339] In another embodiment, the invention provides for a therapeutic population of TILs that is capable of at least three-fold more interferon-gamma production as compared to TILs prepared by a process in which the first expansion of TILs is performed without any added OKT3.

[001340] In another embodiment, the invention provides a method of expanding T cells comprising: (a) performing a priming first expansion of a first population of T cells obtained from a tumor fragment or core obtained from a donor by culturing the first population of T cells to effect growth and to prime an activation of the first population of T cells; (b) after the activation of the first population of T cells primed in step (a) begins to decay, performing a rapid second expansion of the first population of T cells by culturing the first population of T cells to effect growth and to boost the activation of the first population of T cells to obtain a second population of T cells; and (c) harvesting the second population of T cells. In another embodiment, the step of rapid second expansion is split into a plurality of steps to achieve a scaling up of the culture by: (a) performing the rapid second expansion by culturing the first population of T cells in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (b) effecting the transfer of the first population of T cells from the small scale culture to a second container larger than the first container, *e.g.*, a G-REX 500MCS container, and culturing the first population of T cells from the small scale culture in a larger scale culture in the second container for a period of about 4 to 7 days. In another embodiment, the step of rapid expansion is split into a plurality of steps to achieve a scaling out of the culture by: (a) performing the rapid second expansion by culturing the first population of T cells in a first small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (b) effecting the transfer and apportioning of the first population of T cells from the first small scale culture into and amongst at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 second containers that are equal in size to the first container, wherein in each second container the portion of the first population of T cells from first small scale culture transferred to such second container is cultured in a second small scale culture for a period of about 4 to 7 days. In another embodiment, the step of rapid expansion is split into a plurality of steps to achieve a scaling out and scaling up of the culture by: (a) performing the rapid second expansion by culturing the first population of T cells in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 3 to 4 days, and then (b) effecting the transfer and apportioning of the first population of T cells from the small scale culture into and amongst at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 second containers that are larger in size than the first container, *e.g.*, G-REX 500MCS containers, wherein in each second container the portion of the first population of T cells from the small scale culture transferred to such second container is cultured in a larger scale culture for a period of about 4 to 7 days. In another embodiment, the step of rapid expansion

is split into a plurality of steps to achieve a scaling out and scaling up of the culture by: (a) performing the rapid second expansion by culturing the first population of T cells in a small scale culture in a first container, *e.g.*, a G-REX 100MCS container, for a period of about 4 days, and then (b) effecting the transfer and apportioning of the first population of T cells from the small scale culture into and amongst 2, 3 or 4 second containers that are larger in size than the first container, *e.g.*, G-REX 500MCS containers, wherein in each second container the portion of the first population of T cells from the small scale culture transferred to such second container is cultured in a larger scale culture for a period of about 5 days.

[001341] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the priming first expansion of step (a) is performed during a period of up to 17 days.

[001342] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the rapid second expansion of step (b) is performed during a period of up to 11 days.

[001343] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the priming first expansion in step (a) is performed during a period of 17 days and the rapid second expansion of step (b) is performed during a period of up to 9 days.

[001344] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of T cells is cultured in a first culture medium comprising OKT-3 and IL-2.

[001345] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the first culture medium comprises OKT-3, IL-2 and antigen-presenting cells (APCs).

[001346] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the first population of T cells is cultured in a second culture medium comprising OKT-3, IL-2 and antigen-presenting cells (APCs).

[001347] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of T cells is cultured in a first culture medium in a container comprising a first gas-permeable surface, wherein the first culture medium comprises OKT-3, IL-2 and a first population of

antigen-presenting cells (APCs), wherein the first population of APCs is exogenous to the donor of the first population of T cells and the first population of APCs is layered onto the first gas-permeable surface, wherein in step (b) the first population of T cells is cultured in a second culture medium in the container, wherein the second culture medium comprises OKT-3, IL-2 and a second population of APCs, wherein the second population of APCs is exogenous to the donor of the first population of T cells and the second population of APCs is layered onto the first gas-permeable surface, and wherein the second population of APCs is greater than the first population of APCs.

[001348] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the number of APCs in the second population of APCs to the number of APCs in the first population of APCs is about 2:1.

[001349] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the number of APCs in the first population of APCs is about 2.5×10^8 and the number of APCs in the second population of APCs is about 5×10^8.

[001350] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is layered onto the first gas-permeable surface at an average thickness of 2 layers of APCs.

[001351] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the second population of APCs is layered onto the first gas-permeable surface at an average thickness selected from the range of 4 to 8 layers of APCs.

[001352] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the ratio of the average number of layers of APCs layered onto the first gas-permeable surface in step (b) to the average number of layers of APCs layered onto the first gas-permeable surface in step (a) is 2:1.

[001353] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 1.0×10^6 APCs/cm^2 to at or about 4.5×10^6 APCs/cm^2.

[001354] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 1.5×10^6 APCs/cm^2 to at or about 3.5×10^6 APCs/cm^2.

[001355] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 2.0×10^6 APCs/cm^2 to at or about 3.0×10^6 APCs/cm^2.

[001356] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density of at or about 2.0×10^6 APCs/cm^2.

[001357] In another embodiment, the invention provides the method described any of the preceding paragraphs as applicable above modified such that in step (b) the second population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 2.5×10^6 APCs/cm^2 to at or about 7.5×10^6 APCs/cm^2.

[001358] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the second population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 3.5×10^6 APCs/cm^2 to at or about 6.0×10^6 APCs/cm^2.

[001359] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (b) the second population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 4.0×10^6 APCs/cm^2 to at or about 5.5×10^6 APCs/cm^2.

[001360] In another embodiment, the invention provides the method described any of the preceding paragraphs as applicable above modified such that in step (b) the second population of APCs is seeded on the first gas permeable surface at a density of at or about 4.0×10^6 APCs/cm^2.

[001361] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 1.0×10^6 APCs/cm^2 to at or about 4.5×10^6 APCs/cm^2 and in step (b) the second

population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 2.5×10^6 APCs/cm^2 to at or about 7.5×10^6 APCs/cm^2.

[001362] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 1.5×10^6 APCs/cm^2 to at or about 3.5×10^6 APCs/cm^2 and in step (b) the second population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 3.5×10^6 APCs/cm^2 to at or about 6.0×10^6 APCs/cm^2.

[001363] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 2.0×10^6 APCs/cm^2 to at or about 3.0×10^6 APCs/cm^2 and in step (b) the second population of APCs is seeded on the first gas permeable surface at a density selected from the range of at or about 4.0×10^6 APCs/cm^2 to at or about 5.5×10^6 APCs/cm^2.

[001364] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that in step (a) the first population of APCs is seeded on the first gas permeable surface at a density of at or about 2.0×10^6 APCs/cm^2 and in step (b) the second population of APCs is seeded on the first gas permeable surface at a density of at or about 4.0×10^6 APCs/cm^2.

[001365] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the APCs are peripheral blood mononuclear cells (PBMCs).

[001366] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the PBMCs are irradiated and exogenous to the donor of the first population of T cells.

[001367] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the T cells are tumor infiltrating lymphocytes (TILs), the first population of T cells is a first population of TILs in step (a), the second population of T cells is a second population of TILs in step (b), the second population of TILs is a therapeutic population of TILs, and the therapeutic population of TILs is harvested in step (c).

[001368] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the T cells are marrow infiltrating lymphocytes (MILs).

[001369] In another embodiment, the invention provides the method described in any of the preceding paragraphs as applicable above modified such that the T cell phenotype is CD3+ and CD45+.

XV. Pharmaceutical Compositions, Dosages, and Dosing Regimens

[001370] In an embodiment, TILs expanded using the methods of the present disclosure are administered to a patient as a pharmaceutical composition. In an embodiment, the pharmaceutical composition is a suspension of TILs in a sterile buffer. TILs expanded using PBMCs of the present disclosure may be administered by any suitable route as known in the art. In some embodiments, the T-cells are administered as a single intra-arterial or intravenous infusion, which preferably lasts approximately 30 to 60 minutes. Other suitable routes of administration include intraperitoneal, intrathecal, and intralymphatic administration.

[001371] Any suitable dose of TILs can be administered. In some embodiments, from about 2.3×10^{10} to about 13.7×10^{10} TILs are administered, with an average of around 7.8×10^{10} TILs, particularly if the cancer is melanoma. In an embodiment, about 1.2×10^{10} to about 4.3×10^{10} of TILs are administered. In some embodiments, about 3×10^{10} to about 12×10^{10} TILs are administered. In some embodiments, about 4×10^{10} to about 10×10^{10} TILs are administered. In some embodiments, about 5×10^{10} to about 8×10^{10} TILs are administered. In some embodiments, about 6×10^{10} to about 8×10^{10} TILs are administered. In some embodiments, about 7×10^{10} to about 8×10^{10} TILs are administered. In some embodiments, the therapeutically effective dosage is about 2.3×10^{10} to about 13.7×10^{10}. In some embodiments, the therapeutically effective dosage is about 7.8×10^{10} TILs, particularly of the cancer is melanoma. In some embodiments, the therapeutically effective dosage is about 1.2×10^{10} to about 4.3×10^{10} of TILs. In some embodiments, the therapeutically effective dosage is about 3×10^{10} to about 12×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 4×10^{10} to about 10×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 5×10^{10} to about 8×10^{10} TILs. In some embodiments, the therapeutically

effective dosage is about 6×10^{10} to about 8×10^{10} TILs. In some embodiments, the therapeutically effective dosage is about 7×10^{10} to about 8×10^{10} TILs.

[001372] In some embodiments, the number of the TILs provided in the pharmaceutical compositions of the invention is about 1×10^{6}, 2×10^{6}, 3×10^{6}, 4×10^{6}, 5×10^{6}, 6×10^{6}, 7×10^{6}, 8×10^{6}, 9×10^{6}, 1×10^{7}, 2×10^{7}, 3×10^{7}, 4×10^{7}, 5×10^{7}, 6×10^{7}, 7×10^{7}, 8×10^{7}, 9×10^{7}, 1×10^{8}, 2×10^{8}, 3×10^{8}, 4×10^{8}, 5×10^{8}, 6×10^{8}, 7×10^{8}, 8×10^{8}, 9×10^{8}, 1×10^{9}, 2×10^{9}, 3×10^{9}, 4×10^{9}, 5×10^{9}, 6×10^{9}, 7×10^{9}, 8×10^{9}, 9×10^{9}, 1×10^{10}, 2×10^{10}, 3×10^{10}, 4×10^{10}, 5×10^{10}, 6×10^{10}, 7×10^{10}, 8×10^{10}, 9×10^{10}, 1×10^{11}, 2×10^{11}, 3×10^{11}, 4×10^{11}, 5×10^{11}, 6×10^{11}, 7×10^{11}, 8×10^{11}, 9×10^{11}, 1×10^{12}, 2×10^{12}, 3×10^{12}, 4×10^{12}, 5×10^{12}, 6×10^{12}, 7×10^{12}, 8×10^{12}, 9×10^{12}, 1×10^{13}, 2×10^{13}, 3×10^{13}, 4×10^{13}, 5×10^{13}, 6×10^{13}, 7×10^{13}, 8×10^{13}, and 9×10^{13}. In an embodiment, the number of the TILs provided in the pharmaceutical compositions of the invention is in the range of 1×10^{6} to 5×10^{6}, 5×10^{6} to 1×10^{7}, 1×10^{7} to 5×10^{7}, 5×10^{7} to 1×10^{8}, 1×10^{8} to 5×10^{8}, 5×10^{8} to 1×10^{9}, 1×10^{9} to 5×10^{9}, 5×10^{9} to 1×10^{10}, 1×10^{10} to 5×10^{10}, 5×10^{10} to 1×10^{11}, 5×10^{11} to 1×10^{12}, 1×10^{12} to 5×10^{12}, and 5×10^{12} to 1×10^{13}.

[001373] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is less than, for example, 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002% or 0.0001% w/w, w/v or v/v of the pharmaceutical composition.

[001374] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is greater than 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19.75%, 19.50%, 19.25% 19%, 18.75%, 18.50%, 18.25% 18%, 17.75%, 17.50%, 17.25% 17%, 16.75%, 16.50%, 16.25% 16%, 15.75%, 15.50%, 15.25% 15%, 14.75%, 14.50%, 14.25% 14%, 13.75%, 13.50%, 13.25% 13%, 12.75%, 12.50%, 12.25% 12%, 11.75%, 11.50%, 11.25% 11%, 10.75%, 10.50%, 10.25% 10%, 9.75%, 9.50%, 9.25% 9%, 8.75%, 8.50%, 8.25% 8%, 7.75%, 7.50%, 7.25% 7%, 6.75%, 6.50%, 6.25% 6%, 5.75%, 5.50%, 5.25% 5%, 4.75%, 4.50%, 4.25%, 4%, 3.75%, 3.50%, 3.25%, 3%, 2.75%, 2.50%, 2.25%, 2%, 1.75%, 1.50%, 125%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%,

0.0004%, 0.0003%, 0.0002% or 0.0001% w/w, w/v, or v/v of the pharmaceutical composition.

[001375] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is in the range from about 0.0001% to about 50%, about 0.001% to about 40%, about 0.01% to about 30%, about 0.02% to about 29%, about 0.03% to about 28%, about 0.04% to about 27%, about 0.05% to about 26%, about 0.06% to about 25%, about 0.07% to about 24%, about 0.08% to about 23%, about 0.09% to about 22%, about 0.1% to about 21%, about 0.2% to about 20%, about 0.3% to about 19%, about 0.4% to about 18%, about 0.5% to about 17%, about 0.6% to about 16%, about 0.7% to about 15%, about 0.8% to about 14%, about 0.9% to about 12% or about 1% to about 10% w/w, w/v or v/v of the pharmaceutical composition.

[001376] In some embodiments, the concentration of the TILs provided in the pharmaceutical compositions of the invention is in the range from about 0.001% to about 10%, about 0.01% to about 5%, about 0.02% to about 4.5%, about 0.03% to about 4%, about 0.04% to about 3.5%, about 0.05% to about 3%, about 0.06% to about 2.5%, about 0.07% to about 2%, about 0.08% to about 1.5%, about 0.09% to about 1%, about 0.1% to about 0.9% w/w, w/v or v/v of the pharmaceutical composition.

[001377] In some embodiments, the amount of the TILs provided in the pharmaceutical compositions of the invention is equal to or less than 10 g, 9.5 g, 9.0 g, 8.5 g, 8.0 g, 7.5 g, 7.0 g, 6.5 g, 6.0 g, 5.5 g, 5.0 g, 4.5 g, 4.0 g, 3.5 g, 3.0 g, 2.5 g, 2.0 g, 1.5 g, 1.0 g, 0.95 g, 0.9 g, 0.85 g, 0.8 g, 0.75 g, 0.7 g, 0.65 g, 0.6 g, 0.55 g, 0.5 g, 0.45 g, 0.4 g, 0.35 g, 0.3 g, 0.25 g, 0.2 g, 0.15 g, 0.1 g, 0.09 g, 0.08 g, 0.07 g, 0.06 g, 0.05 g, 0.04 g, 0.03 g, 0.02 g, 0.01 g, 0.009 g, 0.008 g, 0.007 g, 0.006 g, 0.005 g, 0.004 g, 0.003 g, 0.002 g, 0.001 g, 0.0009 g, 0.0008 g, 0.0007 g, 0.0006 g, 0.0005 g, 0.0004 g, 0.0003 g, 0.0002 g, or 0.0001 g.

[001378] In some embodiments, the amount of the TILs provided in the pharmaceutical compositions of the invention is more than 0.0001 g, 0.0002 g, 0.0003 g, 0.0004 g, 0.0005 g, 0.0006 g, 0.0007 g, 0.0008 g, 0.0009 g, 0.001 g, 0.0015 g, 0.002 g, 0.0025 g, 0.003 g, 0.0035 g, 0.004 g, 0.0045 g, 0.005 g, 0.0055 g, 0.006 g, 0.0065 g, 0.007 g, 0.0075 g, 0.008 g, 0.0085 g, 0.009 g, 0.0095 g, 0.01 g, 0.015 g, 0.02 g, 0.025 g, 0.03 g, 0.035 g, 0.04 g, 0.045 g, 0.05 g, 0.055 g, 0.06 g, 0.065 g, 0.07 g, 0.075 g, 0.08 g, 0.085 g, 0.09 g, 0.095 g, 0.1 g, 0.15 g, 0.2 g, 0.25 g, 0.3 g, 0.35 g, 0.4 g, 0.45 g, 0.5 g, 0.55 g, 0.6 g, 0.65 g, 0.7 g, 0.75 g, 0.8 g, 0.85 g, 0.9

g, 0.95 g, 1 g, 1.5 g, 2 g, 2.5, 3 g, 3.5, 4 g, 4.5 g, 5 g, 5.5 g, 6 g, 6.5 g, 7 g, 7.5 g, 8 g, 8.5 g, 9 g, 9.5 g, or 10 g.

[001379] The TILs provided in the pharmaceutical compositions of the invention are effective over a wide dosage range. The exact dosage will depend upon the route of administration, the form in which the compound is administered, the gender and age of the subject to be treated, the body weight of the subject to be treated, and the preference and experience of the attending physician. The clinically-established dosages of the TILs may also be used if appropriate. The amounts of the pharmaceutical compositions administered using the methods herein, such as the dosages of TILs, will be dependent on the human or mammal being treated, the severity of the disorder or condition, the rate of administration, the disposition of the active pharmaceutical ingredients and the discretion of the prescribing physician.

[001380] In some embodiments, TILs may be administered in a single dose. Such administration may be by injection, *e.g.*, intravenous injection. In some embodiments, TILs may be administered in multiple doses. Dosing may be once, twice, three times, four times, five times, six times, or more than six times per year. Dosing may be once a month, once every two weeks, once a week, or once every other day. Administration of TILs may continue as long as necessary.

[001381] In some embodiments, an effective dosage of TILs is about 1×10^6, 2×10^6, 3×10^6, 4×10^6, 5×10^6, 6×10^6, 7×10^6, 8×10^6, 9×10^6, 1×10^7, 2×10^7, 3×10^7, 4×10^7, 5×10^7, 6×10^7, 7×10^7, 8×10^7, 9×10^7, 1×10^8, 2×10^8, 3×10^8, 4×10^8, 5×10^8, 6×10^8, 7×10^8, 8×10^8, 9×10^8, 1×10^9, 2×10^9, 3×10^9, 4×10^9, 5×10^9, 6×10^9, 7×10^9, 8×10^9, 9×10^9, 1×10^{10}, 2×10^{10}, 3×10^{10}, 4×10^{10}, 5×10^{10}, 6×10^{10}, 7×10^{10}, 8×10^{10}, 9×10^{10}, 1×10^{11}, 2×10^{11}, 3×10^{11}, 4×10^{11}, 5×10^{11}, 6×10^{11}, 7×10^{11}, 8×10^{11}, 9×10^{11}, 1×10^{12}, 2×10^{12}, 3×10^{12}, 4×10^{12}, 5×10^{12}, 6×10^{12}, 7×10^{12}, 8×10^{12}, 9×10^{12}, 1×10^{13}, 2×10^{13}, 3×10^{13}, 4×10^{13}, 5×10^{13}, 6×10^{13}, 7×10^{13}, 8×10^{13}, and 9×10^{13}. In some embodiments, an effective dosage of TILs is in the range of 1×10^6 to 5×10^6, 5×10^6 to 1×10^7, 1×10^7 to 5×10^7, 5×10^7 to 1×10^8, 1×10^8 to 5×10^8, 5×10^8 to 1×10^9, 1×10^9 to 5×10^9, 5×10^9 to 1×10^{10}, 1×10^{10} to 5×10^{10}, 5×10^{10} to 1×10^{11}, 5×10^{11} to 1×10^{12}, 1×10^{12} to 5×10^{12}, and 5×10^{12} to 1×10^{13}.

[001382] In some embodiments, an effective dosage of TILs is in the range of about 0.01 mg/kg to about 4.3 mg/kg, about 0.15 mg/kg to about 3.6 mg/kg, about 0.3 mg/kg to about 3.2 mg/kg, about 0.35 mg/kg to about 2.85 mg/kg, about 0.15 mg/kg to about 2.85 mg/kg,

about 0.3 mg to about 2.15 mg/kg, about 0.45 mg/kg to about 1.7 mg/kg, about 0.15 mg/kg to about 1.3 mg/kg, about 0.3 mg/kg to about 1.15 mg/kg, about 0.45 mg/kg to about 1 mg/kg, about 0.55 mg/kg to about 0.85 mg/kg, about 0.65 mg/kg to about 0.8 mg/kg, about 0.7 mg/kg to about 0.75 mg/kg, about 0.7 mg/kg to about 2.15 mg/kg, about 0.85 mg/kg to about 2 mg/kg, about 1 mg/kg to about 1.85 mg/kg, about 1.15 mg/kg to about 1.7 mg/kg, about 1.3 mg/kg mg to about 1.6 mg/kg, about 1.35 mg/kg to about 1.5 mg/kg, about 2.15 mg/kg to about 3.6 mg/kg, about 2.3 mg/kg to about 3.4 mg/kg, about 2.4 mg/kg to about 3.3 mg/kg, about 2.6 mg/kg to about 3.15 mg/kg, about 2.7 mg/kg to about 3 mg/kg, about 2.8 mg/kg to about 3 mg/kg, or about 2.85 mg/kg to about 2.95 mg/kg.

[001383] In some embodiments, an effective dosage of TILs is in the range of about 1 mg to about 500 mg, about 10 mg to about 300 mg, about 20 mg to about 250 mg, about 25 mg to about 200 mg, about 1 mg to about 50 mg, about 5 mg to about 45 mg, about 10 mg to about 40 mg, about 15 mg to about 35 mg, about 20 mg to about 30 mg, about 23 mg to about 28 mg, about 50 mg to about 150 mg, about 60 mg to about 140 mg, about 70 mg to about 130 mg, about 80 mg to about 120 mg, about 90 mg to about 110 mg, or about 95 mg to about 105 mg, about 98 mg to about 102 mg, about 150 mg to about 250 mg, about 160 mg to about 240 mg, about 170 mg to about 230 mg, about 180 mg to about 220 mg, about 190 mg to about 210 mg, about 195 mg to about 205 mg, or about 198 to about 207 mg.

[001384] An effective amount of the TILs may be administered in either single or multiple doses by any of the accepted modes of administration of agents having similar utilities, including intranasal and transdermal routes, by intra-arterial injection, intravenously, intraperitoneally, parenterally, intramuscularly, subcutaneously, topically, by transplantation, or by inhalation.

[001385] In another embodiment, the invention provides an infusion bag comprising the therapeutic population of TILs described in any of the preceding paragraphs as applicable above.

[001386] In another embodiment, the invention provides a tumor infiltrating lymphocyte (TIL) composition comprising the therapeutic population of TILs described in any of the preceding paragraphs as applicable above and a pharmaceutically acceptable carrier.

[001387] In another embodiment, the invention provides an infusion bag comprising the TIL composition described in any of the preceding paragraphs as applicable above.

[001388] In another embodiment, the invention provides a cryopreserved preparation of the therapeutic population of TILs described in in any of the preceding paragraphs as applicable above.

[001389] In another embodiment, the invention provides a tumor infiltrating lymphocyte (TIL) composition comprising the therapeutic population of TILs described in any of the preceding paragraphs as applicable above and a cryopreservation media.

[001390] In another embodiment, the invention provides the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cryopreservation media contains DMSO.

[001391] In another embodiment, the invention provides the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cryopreservation media contains 7-10% DMSO.

[001392] In another embodiment, the invention provides a cryopreserved preparation of the TIL composition described in any of the preceding paragraphs as applicable above.

XVI. Methods of Treating Patients

[001393] Methods of treatment begin with the initial TIL collection and culture of TILs. Such methods have been both described in the art by, for example, Jin *et al., J. Immunotherapy*, **2012**, 35(3):283-292, incorporated by reference herein in its entirety. Embodiments of methods of treatment are described throughout the sections below, including the Examples.

[001394] The expanded TILs produced according the methods described herein, including, for example as described in Steps A through F above or according to Steps A through F above (also as shown, for example, in Figure 85 (in particular, *e.g.*, Figure 85B)) find particular use in the treatment of patients with cancer (for example, as described in Goff, *et al.*, *J. Clinical Oncology*, **2016**, 34(20):2389-239, as well as the supplemental content; incorporated by reference herein in its entirety. In some embodiments, TIL were grown from resected deposits of metastatic melanoma as previously described (see, Dudley, *et al.*, *J Immunother.*, **2003**, 26:332-342; incorporated by reference herein in its entirety). Fresh tumor can be dissected under sterile conditions. A representative sample can be collected for formal pathologic analysis. Single fragments of 2 mm^3 to 3 mm^3 may be used. In some embodiments, 5, 10, 15, 20, 25 or 30 samples per patient are obtained. In some embodiments, 20, 25, or 30 samples per patient are obtained. In some embodiments, 20, 22, 24, 26, or 28

samples per patient are obtained. In some embodiments, 24 samples per patient are obtained. Samples can be placed in individual wells of a 24-well plate, maintained in growth media with high-dose IL-2 (6,000 IU/mL), and monitored for destruction of tumor and/or proliferation of TIL. Any tumor with viable cells remaining after processing can be enzymatically digested into a single cell suspension and cryopreserved, as described herein.

[001395] In some embodiments, successfully grown TIL can be sampled for phenotype analysis (CD3, CD4, CD8, and CD56) and tested against autologous tumor when available. TIL can be considered reactive if overnight coculture yielded interferon-gamma (IFN-γ) levels > 200 pg/mL and twice background. (Goff, *et al.*, *J Immunother.*, **2010**, 33:840-847; incorporated by reference herein in its entirety). In some embodiments, cultures with evidence of autologous reactivity or sufficient growth patterns can be selected for a second expansion (for example, a second expansion as provided in according to Step D of Figure 85 (in particular, *e.g.*, Figure 85B)), including second expansions that are sometimes referred to as rapid expansion (REP). In some embodiments, expanded TILs with high autologous reactivity (for example, high proliferation during a second expansion), are selected for an additional second expansion. In some embodiments, TILs with high autologous reactivity (for example, high proliferation during second expansion as provided in Step D of Figure 85 (in particular, *e.g.*, Figure 85B)), are selected for an additional second expansion according to Step D of Figure 85 (in particular, *e.g.*, Figure 85B).

[001396] In some embodiments, the patient is not moved directly to ACT (adoptive cell transfer), for example, in some embodiments, after tumor harvesting and/or a first expansion, the cells are not utilized immediately. In some embodiments, TILs can be cryopreserved and thawed 2 days before administration to a patient. In some embodiments, TILs can be cryopreserved and thawed 1 day before administration to a patient. In some embodiments, the TILs can be cryopreserved and thawed immediately before the administration to a patient.

[001397] Cell phenotypes of cryopreserved samples of infusion bag TIL can be analyzed by flow cytometry (*e.g.*, FlowJo™) for surface markers CD3, CD4, CD8, CCR7, and CD45RA (BD BioSciences), as well as by any of the methods described herein. Serum cytokines were measured by using standard enzyme-linked immunosorbent assay techniques. A rise in serum IFN-g was defined as >100 pg/mL and greater than 4 3 baseline levels.

[001398] In some embodiments, the TILs produced by the methods provided herein, for example those exemplified in Figure 85 (in particular, *e.g.*, Figure 85B), provide for a

surprising improvement in clinical efficacy of the TILs. In some embodiments, the TILs produced by the methods provided herein, for example those exemplified in Figure 85 (in particular, *e.g.*, Figure 85B), exhibit increased clinical efficacy as compared to TILs produced by methods other than those described herein, including, for example, methods other than those exemplified in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, the methods other than those described herein include methods referred to as process 1C and/or Generation 1 (Gen 1). In some embodiments, the increased efficacy is measured by DCR, ORR, and/or other clinical responses. In some embodiments, the TILS produced by the methods provided herein, for example those exemplified in Figure 85 (in particular, *e.g.*, Figure 85B), exhibit a similar time to response and safety profile compared to TILs produced by methods other than those described herein, including, for example, methods other than those exemplified in Figure 85 (in particular, *e.g.*, Figure 85B), for example the Gen 1 process.

[001399] In some embodiments, IFN-gamma (IFN-γ) is indicative of treatment efficacy and/or increased clinical efficacy. In some embodiments, IFN-γ in the blood of subjects treated with TILs is indicative of active TILs. In some embodiments, a potency assay for IFN-γ production is employed. IFN-γ production is another measure of cytotoxic potential. IFN-γ production can be measured by determining the levels of the cytokine IFN-γ in the blood, serum, or TILs *ex vivo* of a subject treated with TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, an increase in IFN-γ is indicative of treatment efficacy in a patient treated with the TILs produced by the methods of the present invention. In some embodiments, IFN-γ is increased one-fold, two-fold, three-fold, four-fold, or five-fold or more as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased one-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased two-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ

secretion is increased three-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased four-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased five-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ is measured using a Quantikine ELISA kit. In some embodiments, IFN-γ is measured in TILs *ex vivo* of a subject treated with TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ is measured in blood of a subject treated with TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ is measured in TILs serum of a subject treated with TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B).

[001400] In some embodiments, the TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B), exhibit increased polyclonality as compared to TILs produced by other methods, including those not exemplified in Figure 85 (in particular, *e.g.*, Figure 85B), such as for example, methods referred to as process 1C methods. In some embodiments, significantly improved polyclonality and/or increased polyclonality is indicative of treatment efficacy and/or increased clinical efficacy. In some embodiments, polyclonality refers to the T-cell repertoire diversity. In some embodiments, an increase in polyclonality can be indicative of treatment efficacy with regard to administration of the TILs produced by the methods of the present invention. In some embodiments, polyclonality is increased one-fold, two-fold, ten-fold, 100-fold, 500-fold, or 1000-fold as compared to TILs prepared using methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased one-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other

methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased two-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased ten-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased 100-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased 500-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased 1000-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B).

[001401] Measures of efficacy can include the disease control rate (DCR) as well as overall response rate (ORR), as known in the art as well as described herein.

1. Methods of Treating Cancers and Other Diseases

[001402] The compositions and methods described herein can be used in a method for treating diseases. In an embodiment, they are for use in treating hyperproliferative disorders. They may also be used in treating other disorders as described herein and in the following paragraphs.

[001403] In some embodiments, the hyperproliferative disorder is cancer. In some embodiments, the hyperproliferative disorder is a solid tumor cancer. In some embodiments, the solid tumor cancer is selected from the group consisting of glioblastoma (GBM), gastrointestinal cancer, melanoma, ovarian cancer, cervical cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer (including head and neck squamous cell carcinoma

(HNSCC)), renal cancer, and renal cell carcinoma. In some embodiments, the hyperproliferative disorder is a hematological malignancy. In some embodiments, the solid tumor cancer is selected from the group consisting of chronic lymphocytic leukemia, acute lymphoblastic leukemia, diffuse large B cell lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, follicular lymphoma, and mantle cell lymphoma.

[001404] In some embodiments, the cancer is a hypermutated cancer phenotype. Hypermutated cancers are extensively described in Campbell, *et al.* (Cell, 171:1042-1056 (2017); incorporated by reference herein in its entirety for all purposes). In some embodiments, a hypermutated tumors comprise between 9 and 10 mutations per megabase (Mb). In some embodiments, pediatric hypermutated tumors comprise 9.91 mutations per megabase (Mb). In some embodiments, adult hypermutated tumors comprise 9 mutations per megabase (Mb). In some embodiments, enhanced hypermutated tumors comprise between 10 and 100 mutations per megabase (Mb). In some embodiments, enhanced pediatric hypermutated tumors comprise between 10 and 100 mutations per megabase (Mb). In some embodiments, enhanced adult hypermutated tumors comprise between 10 and 100 mutations per megabase (Mb). In some embodiments, an ultra-hypermutated tumors comprise greater than 100 mutations per megabase (Mb). In some embodiments, pediatric ultra-hypermutated tumors comprise greater than 100 mutations per megabase (Mb). In some embodiments, adult ultra-hypermutated tumors comprise greater than 100 mutations per megabase (Mb).

[001405] In some embodiments, the hypermutated tumors have mutations in replication repair pathways. In some embodiments, the hypermutated tumors have mutations in replication repair associated DNA polymerases. In some embodiments, the hypermutated tumors have microsatellite instability. In some embodiments, the ultra-hypermutated tumors have mutations in replication repair associated DNA polymerases and have microsatellite instability. In some embodiments, hypermutation in the tumor is correlated with response to immune checkpoint inhibitors. In some embodiments, hypermutated tumors are resistant to treatment with immune checkpoint inhibitors. In some embodiments, hypermutated tumors can be treated using the TILs of the present invention. In some embodiments, hypermutation in the tumor is caused by environmental factors (extrinsic exposures). For example, UV light can be the primary cause of the high numbers of mutations in malignant melanoma (*see*, for example, Pfeifer, G.P., You, Y.H., and Besaratinia, A. **(2005)**. *Mutat. Res.* 571, 19–31.; Sage, E. **(1993)**. *Photochem. Photobiol.* 57, 163–174.). In some embodiments, hypermutation in the tumor can be caused by the greater than 60 carcinogens in tobacco smoke for tumors of the

lung and larynx, as well as other tumors, due to direct mutagen exposure (*see*, for example, Pleasance, E.D., Stephens, P.J., O'Meara, S., McBride, D.J., Meynert, A., Jones, D., Lin, M.L., Beare, D., Lau, K.W., Greenman, C., *et al.* **(2010)**. *Nature* 463, 184–190). In some embodiments, hypermutation in the tumor is caused by dysregulation of apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like (APOBEC) family members, which has been shown to result in increased levels of C to T transitions in a wide range of cancers (see, for example, Roberts, S.A., Lawrence, M.S., Klimczak, L.J., Grimm, S.A., Fargo, D., Stojanov, P., Kiezun, A., Kryukov, G.V., Carter, S.L., Saksena, G., *et al.* **(2013)**. *Nat. Genet.* 45, 970–976). In some embodiments, hypermutation in the tumor is caused by defective DNA replication repair by mutations that compromise proofreading, which is performed by the major replicative enzymes Pol3 and Pold1. In some embodiments, hypermutation in the tumor is caused by defects in DNA mismatch repair, which are associated with hypermutation in colorectal, endometrial, and other cancers (see, for example, Kandoth, C., Schultz, N., Cherniack, A.D., Akbani, R., Liu, Y., Shen, H., Robertson, A.G., Pashtan, I., Shen, R., Benz, C.C., *et al.*; **(2013)**. *Nature* 497, 67–73.; Muzny, D.M., Bainbridge, M.N., Chang, K., Dinh, H.H., Drummond, J.A., Fowler, G., Kovar, C.L., Lewis, L.R., Morgan, M.B., Newsham, I.F., *et al.*; **(2012)**. *Nature* 487, 330–337). In some embodiments, DNA replication repair mutations are also found in cancer predisposition syndromes, such as constitutional or biallelic mismatch repair deficiency (CMMRD), Lynch syndrome, and polymerase proofreading-associated polyposis (PPAP).

[001406] In an embodiment, the invention includes a method of treating a cancer with a population of TILs, wherein the cancer is a hypermutated cancer. In an embodiment, the invention includes a method of treating a cancer with a population of TILs, wherein the cancer is an enhanced hypermutated cancer. In an embodiment, the invention includes a method of treating a cancer with a population of TILs, wherein the cancer is an ultra-hypermutated cancer.

[001407] In an embodiment, the invention includes a method of treating a cancer with a population of TILs, wherein a patient is pre-treated with non-myeloablative chemotherapy prior to an infusion of TILs according to the present disclosure. In an embodiment, the non-myeloablative chemotherapy is cyclophosphamide 60 mg/kg/d for 2 days (days 27 and 26 prior to TIL infusion) and fludarabine 25 mg/m^2/d for 5 days (days 27 to 23 prior to TIL infusion). In an embodiment, after non-myeloablative chemotherapy and TIL infusion (at day

0) according to the present disclosure, the patient receives an intravenous infusion of IL-2 intravenously at 720,000 IU/kg every 8 hours to physiologic tolerance.

[001408] Efficacy of the compounds and combinations of compounds described herein in treating, preventing and/or managing the indicated diseases or disorders can be tested using various models known in the art, which provide guidance for treatment of human disease. For example, models for determining efficacy of treatments for ovarian cancer are described, *e.g.*, in Mullany, *et al.*, *Endocrinology* **2012,** *153,* 1585-92; and Fong, *et al.*, *J. Ovarian Res.* **2009,** *2,* 12. Models for determining efficacy of treatments for pancreatic cancer are described in Herreros-Villanueva, *et al.*, *World J. Gastroenterol.* **2012,** *18,* 1286-1294. Models for determining efficacy of treatments for breast cancer are described, *e.g.*, in Fantozzi, *Breast Cancer Res.* **2006,** *8,* 212. Models for determining efficacy of treatments for melanoma are described, *e.g.*, in Damsky, *et al.*, *Pigment Cell & Melanoma Res.* **2010,** *23,* 853–859. Models for determining efficacy of treatments for lung cancer are described, *e.g.*, in Meuwissen, *et al.*, *Genes & Development*, **2005,** *19,* 643-664. Models for determining efficacy of treatments for lung cancer are described, *e.g.*, in Kim, *Clin. Exp. Otorhinolaryngol.* **2009,** *2,* 55-60; and Sano, *Head Neck Oncol.* **2009,** *1,* 32.

[001409] In some embodiments, IFN-gamma (IFN-γ) is indicative of treatment efficacy for hyperproliferative disorder treatment. In some embodiments, IFN-γ in the blood of subjects treated with TILs is indicative of active TILs. In some embodiments, a potency assay for IFN-γ production is employed. IFN-γ production is another measure of cytotoxic potential. IFN-γ production can be measured by determining the levels of the cytokine IFN-γ in the blood of a subject treated with TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, the TILs obtained by the present method provide for increased IFN-γ in the blood of subjects treated with the TILs of the present method as compared subjects treated with TILs prepared using methods referred to as the Gen 3 process, as exemplified Figure 85 (in particular, *e.g.*, Figure 85B) and throughout this application. In some embodiments, an increase in IFN-γ is indicative of treatment efficacy in a patient treated with the TILs produced by the methods of the present invention. In some embodiments, IFN-γ is increased one-fold, two-fold, three-fold, four-fold, or five-fold or more as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ

secretion is increased one-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased two-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased three-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased four-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ secretion is increased five-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, IFN-γ is measured using a Quantikine ELISA kit. In some embodiments, IFN-γ is measured using a Quantikine ELISA kit. In some embodiments, IFN-γ is measured in TILs *ex vivo* from a patient treated with the TILs produced by the methods of the present invention. In some embodiments, IFN-γ is measured in blood in a patient treated with the TILs produced by the methods of the present invention. In some embodiments, IFN-γ is measured in serum in a patient treated with the TILs produced by the methods of the present invention.

[001410] In some embodiments, the TILs prepared by the methods of the present invention, including those as described for example in Figure 85 (in particular, *e.g.*, Figure 85B), exhibit increased polyclonality as compared to TILs produced by other methods, including those not exemplified in Figure 85 (in particular, *e.g.*, Figure 85B), such as for example, methods referred to as process 1C methods. In some embodiments, significantly improved polyclonality and/or increased polyclonality is indicative of treatment efficacy and/or increased clinical efficacy for cancer treatment. In some embodiments, polyclonality refers to the T-cell repertoire diversity. In some embodiments, an increase in polyclonality can be indicative of treatment efficacy with regard to administration of the TILs produced by the

methods of the present invention. In some embodiments, polyclonality is increased one-fold, two-fold, ten-fold, 100-fold, 500-fold, or 1000-fold as compared to TILs prepared using methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased one-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased two-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased ten-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased 100-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased 500-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B). In some embodiments, polyclonality is increased 1000-fold as compared to an untreated patient and/or as compared to a patient treated with TILs prepared using other methods than those provide herein including, for example, methods other than those embodied in Figure 85 (in particular, *e.g.*, Figure 85B).

2. **Methods of co-administration**

[001411] In some embodiments, the TILs produced as described herein, including, for example TILs derived from a method described in Steps A through F of Figure 85 (in particular, *e.g.*, Figure 85B), can be administered in combination with one or more immune checkpoint regulators, such as the antibodies described below. For example, antibodies that target PD-1 and which can be co-administered with the TILs of the present invention include, *e.g.*, but are not limited to nivolumab (BMS-936558, Bristol-Myers Squibb; Opdivo®), pembrolizumab (lambrolizumab, MK03475 or MK-3475, Merck; Keytruda®), humanized anti-PD-1 antibody JS001 (ShangHai JunShi), monoclonal anti-PD-1 antibody TSR-042

(Tesaro, Inc.), Pidilizumab (anti-PD-1 mAb CT-011, Medivation), anti-PD-1 monoclonal Antibody BGB-A317 (BeiGene), and/or anti-PD-1 antibody SHR-1210 (ShangHai HengRui), human monoclonal antibody REGN2810 (Regeneron), human monoclonal antibody MDX-1106 (Bristol-Myers Squibb), and/or humanized anti-PD-1 IgG4 antibody PDR001 (Novartis). In some embodiments, the PD-1 antibody is from clone: RMP1-14 (rat IgG) - BioXcell cat# BP0146. Other suitable antibodies suitable for use in co-administration methods with TILs produced according to Steps A through F as described herein are anti-PD-1 antibodies disclosed in U.S. Patent No. 8,008,449, herein incorporated by reference. In some embodiments, the antibody or antigen-binding portion thereof binds specifically to PD-L1 and inhibits its interaction with PD-1, thereby increasing immune activity. Any antibodies known in the art which bind to PD-L1 and disrupt the interaction between the PD-1 and PD-L1, and stimulates an anti- tumor immune response, are suitable for use in co-administration methods with TILs produced according to Steps A through F as described herein. For example, antibodies that target PD-L1 and are in clinical trials, include BMS-936559 (Bristol-Myers Squibb) and MPDL3280A (Genentech). Other suitable antibodies that target PD-L1 are disclosed in U.S. Patent No. 7,943,743, herein incorporated by reference. It will be understood by one of ordinary skill that any antibody which binds to PD-1 or PD-L1, disrupts the PD-1/PD-L1 interaction, and stimulates an anti-tumor immune response, are suitable for use in co-administration methods with TILs produced according to Steps A through F as described herein. In some embodiments, the subject administered the combination of TILs produced according to Steps A through F is co administered with a and anti-PD-1 antibody when the patient has a cancer type that is refractory to administration of the anti-PD-1 antibody alone. In some embodiments, the patient is administered TILs in combination with and anti-PD-1 when the patient has refractory melanoma. In some embodiments, the patient is administered TILs in combination with and anti-PD-1 when the patient has non-small-cell lung carcinoma (NSCLC).

3. Optional Lymphodepletion Preconditioning of Patients

[001412] In an embodiment, the invention includes a method of treating a cancer with a population of TILs, wherein a patient is pre-treated with non-myeloablative chemotherapy prior to an infusion of TILs according to the present disclosure. In an embodiment, the invention includes a population of TILs for use in the treatment of cancer in a patient which has been pre-treated with non-myeloablative chemotherapy. In an embodiment, the population of TILs is for administration by infusion. In an embodiment, the non-

myeloablative chemotherapy is cyclophosphamide 60 mg/kg/d for 2 days (days 27 and 26 prior to TIL infusion) and fludarabine 25 mg/m^2/d for 5 days (days 27 to 23 prior to TIL infusion). In an embodiment, after non-myeloablative chemotherapy and TIL infusion (at day 0) according to the present disclosure, the patient receives an intravenous infusion of IL-2 (aldesleukin, commercially available as PROLEUKIN) intravenously at 720,000 IU/kg every 8 hours to physiologic tolerance. In certain embodiments, the population of TILs is for use in treating cancer in combination with IL-2, wherein the IL-2 is administered after the population of TILs.

[001413] Experimental findings indicate that lymphodepletion prior to adoptive transfer of tumor-specific T lymphocytes plays a key role in enhancing treatment efficacy by eliminating regulatory T cells and competing elements of the immune system ('cytokine sinks'). Accordingly, some embodiments of the invention utilize a lymphodepletion step (sometimes also referred to as "immunosuppressive conditioning") on the patient prior to the introduction of the TILs of the invention.

[001414] In general, lymphodepletion is achieved using administration of fludarabine or cyclophosphamide (the active form being referred to as mafosfamide) and combinations thereof. Such methods are described in Gassner, *et al., Cancer Immunol. Immunother.* **2011**, *60,* 75–85, Muranski, *et al.*, *Nat. Clin. Pract. Oncol.,* **2006,** *3,* 668–681, Dudley, *et al.*, *J. Clin. Oncol.* **2008**, *26,* 5233-5239, and Dudley, *et al.*, *J. Clin. Oncol.* **2005**, *23,* 2346–2357, all of which are incorporated by reference herein in their entireties.

[001415] In some embodiments, the fludarabine is administered at a concentration of 0.5 μg/mL -10 μg/mL fludarabine. In some embodiments, the fludarabine is administered at a concentration of 1 μg/mL fludarabine. In some embodiments, the fludarabine treatment is administered for 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, or 7 days or more. In some embodiments, the fludarabine is administered at a dosage of 10 mg/kg/day, 15 mg/kg/day, 20 mg/kg/day, 25 mg/kg/day, 30 mg/kg/day, 35 mg/kg/day, 40 mg/kg/day, or 45 mg/kg/day. In some embodiments, the fludarabine treatment is administered for 2-7 days at 35 mg/kg/day. In some embodiments, the fludarabine treatment is administered for 4-5 days at 35 mg/kg/day. In some embodiments, the fludarabine treatment is administered for 4-5 days at 25 mg/kg/day.

[001416] In some embodiments, the mafosfamide, the active form of cyclophosphamide, is obtained at a concentration of 0.5 μg/mL -10 μg/mL by administration of cyclophosphamide.

In some embodiments, mafosfamide, the active form of cyclophosphamide, is obtained at a concentration of 1 μg/mL by administration of cyclophosphamide. In some embodiments, the cyclophosphamide treatment is administered for 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, or 7 days or more. In some embodiments, the cyclophosphamide is administered at a dosage of 100 mg/m^2/day, 150 mg/m^2/day, 175 mg/m^2/day, 200 mg/m^2/day, 225 mg/m^2/day, 250 mg/m^2/day, 275 mg/m^2/day, or 300 mg/m^2/day. In some embodiments, the cyclophosphamide is administered intravenously *(i.e.*, i.v.) In some embodiments, the cyclophosphamide treatment is administered for 2-7 days at 35 mg/kg/day. In some embodiments, the cyclophosphamide treatment is administered for 4-5 days at 250 mg/m^2/day i.v. In some embodiments, the cyclophosphamide treatment is administered for 4 days at 250 mg/m^2/day i.v.

[001417] In some embodiments, lymphodepletion is performed by administering the fludarabine and the cyclophosphamide together to a patient. In some embodiments, fludarabine is administered at 25 mg/m^2/day i.v. and cyclophosphamide is administered at 250 mg/m^2/day i.v. over 4 days.

[001418] In an embodiment, the lymphodepletion is performed by administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

4. **IL-2 Regimens**

[001419] In an embodiment, the IL-2 regimen comprises a high-dose IL-2 regimen, wherein the high-dose IL-2 regimen comprises aldesleukin, or a biosimilar or variant thereof, administered intravenously starting on the day after administering a therapeutically effective portion of the therapeutic population of TILs, wherein the aldesleukin or a biosimilar or variant thereof is administered at a dose of 0.037 mg/kg or 0.044 mg/kg IU/kg (patient body mass) using 15-minute bolus intravenous infusions every eight hours until tolerance, for a maximum of 14 doses. Following 9 days of rest, this schedule may be repeated for another 14 doses, for a maximum of 28 doses in total.

[001420] In an embodiment, the IL-2 regimen comprises a decrescendo IL-2 regimen. Decrescendo IL-2 regimens have been described in O'Day, *et al.*, *J. Clin. Oncol.* **1999,** *17,* 2752-61 and Eton, *et al.*, *Cancer* **2000,** *88,* 1703-9, the disclosures of which are incorporated herein by reference. In an embodiment, a decrescendo IL-2 regimen comprises 18×10^6 IU/m^2 administered intravenously over 6 hours, followed by 18×10^6 IU/m^2 administered

intravenously over 12 hours, followed by 18×10^6 IU/m^2 administered intravenously over 24 hrs, followed by 4.5×10^6 IU/m^2 administered intravenously over 72 hours. This treatment cycle may be repeated every 28 days for a maximum of four cycles. In an embodiment, a decrescendo IL-2 regimen comprises 18,000,000 IU/m^2 on day 1, 9,000,000 IU/m^2 on day 2, and 4,500,000 IU/m^2 on days 3 and 4.

[001421] In an embodiment, the IL-2 regimen comprises administration of pegylated IL-2 every 1, 2, 4, 6, 7, 14 or 21 days at a dose of 0.10 mg/day to 50 mg/day.

5. **Adoptive Cell Transfer**

[001422] Adoptive cell transfer (ACT) is an effective form of immunotherapy and involves the transfer of immune cells with antitumor activity into cancer patients. ACT is a treatment approach that involves the identification, *in vitro*, of lymphocytes with antitumor activity, the *in vitro* expansion of these cells to large numbers and their infusion into the cancer-bearing host. Lymphocytes used for adoptive transfer can be derived from the stroma of resected tumors (tumor infiltrating lymphocytes or TILs). TILs for ACT can be prepared as described herein. In some embodiments, the TILs are prepared, for example, according to a method as described in Figure 85 (in particular, *e.g.*, Figure 85B). They can also be derived or from blood if they are genetically engineered to express antitumor T-cell receptors (TCRs) or chimeric antigen receptors (CARs), enriched with mixed lymphocyte tumor cell cultures (MLTCs), or cloned using autologous antigen presenting cells and tumor derived peptides. ACT in which the lymphocytes originate from the cancer-bearing host to be infused is termed autologous ACT. U.S. Publication No. 2011/0052530 relates to a method for performing adoptive cell therapy to promote cancer regression, primarily for treatment of patients suffering from metastatic melanoma, which is incorporated by reference in its entirety for these methods. In some embodiments, TILs can be administered as described herein. In some embodiments, TILs can be administered in a single dose. Such administration may be by injection, *e.g.*, intravenous injection. In some embodiments, TILs and/or cytotoxic lymphocytes may be administered in multiple doses. Dosing may be once, twice, three times, four times, five times, six times, or more than six times per year. Dosing may be once a month, once every two weeks, once a week, or once every other day. Administration of TILs and/or cytotoxic lymphocytes may continue as long as necessary.

6. Additional Methods of Treatment

[001423] In another embodiment, the invention provides a method for treating a subject with cancer comprising administering to the subject a therapeutically effective dosage of the therapeutic TIL population described in any of the preceding paragraphs as applicable above.

[001424] In another embodiment, the invention provides a method for treating a subject with cancer comprising administering to the subject a therapeutically effective dosage of the TIL composition described in any of the preceding paragraphs as applicable above

[001425] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that prior to administering the therapeutically effective dosage of the therapeutic TIL population and the TIL composition described in any of the preceding paragraphs as applicable above, respectively, a non-myeloablative lymphodepletion regimen has been administered to the subject.

[001426] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

[001427] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified to further comprise the step of treating the subject with a high-dose IL-2 regimen starting on the day after administration of the TIL cells to the subject.

[001428] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the the high-dose IL-2 regimen comprises 600,000 or 720,000 IU/kg administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

[001429] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is a solid tumor.

[001430] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such

that the cancer is melanoma, ovarian cancer, cervical cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer (including head and neck squamous cell carcinoma (HNSCC)), glioblastoma (including GBM), gastrointestinal cancer, renal cancer, or renal cell carcinoma.

[001431] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma, HNSCC, cervical cancers, NSCLC, glioblastoma (including GBM), and gastrointestinal cancer.

[001432] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma.

[001433] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is HNSCC.

[001434] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is a cervical cancer.

[001435] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is NSCLC.

[001436] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is glioblastoma (including GBM).

[001437] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is gastrointestinal cancer.

[001438] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is a hypermutated cancer.

[001439] In another embodiment, the invention provides the method for treating a subject with cancer described in any of the preceding paragraphs as applicable above modified such that the cancer is a pediatric hypermutated cancer.

[001440] In another embodiment, the invention provides the therapeutic TIL population described in any of the preceding paragraphs as applicable above for use in a method for treating a subject with cancer comprising administering to the subject a therapeutically effective dosage of the therapeutic TIL population.

[001441] In another embodiment, the invention provides the TIL composition described in any of the preceding paragraphs as applicable above for use in a method for treating a subject with cancer comprising administering to the subject a therapeutically effective dosage of the TIL composition.

[001442] In another embodiment, the invention provides the therapeutic TIL population described in any of the preceding paragraphs as applicable above or the TIL composition described in any of the preceding paragraphs as applicable above modified such that prior to administering to the subject the therapeutically effective dosage of the therapeutic TIL population described in any of the preceding paragraphs as applicable above or the TIL composition described in any of the preceding paragraphs as applicable above, a non-myeloablative lymphodepletion regimen has been administered to the subject.

[001443] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

[001444] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified to further comprise the step of treating patient with a high-dose IL-2 regimen starting on the day after administration of the TIL cells to the patient.

[001445] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the high-dose IL-2 regimen comprises 600,000 or 720,000 IU/kg administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

[001446] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a solid tumor.

[001447] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma, ovarian cancer, cervical cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer (including head and neck squamous cell carcinoma (HNSCC)), glioblastoma (including GBM), gastrointestinal cancer, renal cancer, or renal cell carcinoma.

[001448] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma, HNSCC, cervical cancers, NSCLC, glioblastoma (including GBM), and gastrointestinal cancer.

[001449] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma.

[001450] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is HNSCC.

[001451] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a cervical cancer.

[001452] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is NSCLC.

[001453] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is glioblastoma (including GBM).

[001454] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is gastrointestinal cancer.

[001455] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a hypermutated cancer.

[001456] In another embodiment, the invention provides the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a pediatric hypermutated cancer.

[001457] In another embodiment, the invention provides the use of the therapeutic TIL population described in any of the preceding paragraphs as applicable above in a method of treating cancer in a subject comprising administering to the subject a therapeutically effective dosage of the therapeutic TIL population.

[001458] In another embodiment, the invention provides the use of the TIL composition described in any of the preceding paragraphs as applicable above in a method of treating cancer in a subject comprising administering to the subject a therapeutically effective dosage of the TIL composition.

[001459] In another embodiment, the invention provides the use of the therapeutic TIL population described in any of the preceding paragraphs as applicable above or the TIL composition described in any of the preceding paragraphs as applicable above in a method of treating cancer in a subject comprising administering to the subject a non-myeloablative lymphodepletion regimen and then administering to the subject a therapeutically effective dosage of the therapeutic TIL population described in any of the preceding paragraphs as applicable above or the therapeutically effective dosage of the TIL composition described in any of the preceding paragraphs as applicable above.

[001460] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

[001461] In another embodiment, the invention provides the use of the therapeutic TIL population described in any of the preceding paragraphs as applicable above or the use of the TIL composition described in any of the preceding paragraphs as applicable above modified to further comprise the step of treating patient with a high-dose IL-2 regimen starting on the day after administration of the TIL cells to the patient.

[001462] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the high-dose IL-2 regimen comprises 600,000 or 720,000 IU/kg administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

[001463] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a solid tumor.

[001464] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma, ovarian cancer, cervical cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer (including head and neck squamous cell carcinoma (HNSCC)), glioblastoma (including GBM), gastrointestinal cancer, renal cancer, or renal cell carcinoma.

[001465] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma, HNSCC, cervical cancers, NSCLC, glioblastoma (including GBM), and gastrointestinal cancer.

[001466] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is melanoma.

[001467] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is HNSCC.

[001468] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a cervical cancer.

[001469] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is NSCLC.

[001470] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is glioblastoma (including GBM).

[001471] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is gastrointestinal cancer.

[001472] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a hypermutated cancer.

[001473] In another embodiment, the invention provides the use of the therapeutic TIL population or the TIL composition described in any of the preceding paragraphs as applicable above modified such that the cancer is a pediatric hypermutated cancer.

7. Exemplary Treatment Embodiments

[001474] In some embodiments, the present disclosure provides a method of treating a cancer with a population of tumor infiltrating lymphocytes (TILs) comprising the steps of (a) obtaining a first population of TILs from a tumor fragment or core obtained from a patient; (b) performing an initial expansion of the first population of TILs in a first cell culture medium to obtain a second population of TILs, wherein the second population of TILs is at least 5-fold greater in number than the first population of TILs, and wherein the first cell culture medium comprises IL-2; (c) performing a rapid expansion of the second population of TILs using a population of myeloid artificial antigen presenting cells (myeloid aAPCs) in a second cell culture medium to obtain a third population of TILs, wherein the third population of TILs is at least 50-fold greater in number than the second population of TILs after 7 days from the start of the rapid expansion; and wherein the second cell culture medium comprises IL-2 and OKT-3; (d) administering a therapeutically effective portion of the third population of TILs to a patient with the cancer. In some embodiments, the IL-2 is present at an initial concentration of about 3000 IU/mL and OKT-3 antibody is present at an initial concentration of about 30 ng/mL in the second cell culture medium. In some embodiments, first expansion is performed over a period not greater than 14 days. In some embodiments, the first expansion is performed using a gas permeable container. In some embodiments, the second expansion is performed using a gas permeable container. In some embodiments, the ratio of the second population of TILs to the population of aAPCs in the rapid expansion is between 1

to 80 and 1 to 400. In some embodiments, the ratio of the second population of TILs to the population of aAPCs in the rapid expansion is about 1 to 300. In some embodiments, the cancer for treatment is selected from the group consisting of melanoma, ovarian cancer, cervical cancer, non-small-cell lung cancer (NSCLC), lung cancer, bladder cancer, breast cancer, cancer caused by human papilloma virus, head and neck cancer, renal cancer, and renal cell carcinoma. In some embodiments, the cancer for treatment is selected from the group consisting of melanoma, ovarian cancer, and cervical cancer. In some embodiments, the cancer for treatment is melanoma. In some embodiments, the cancer for treatment is ovarian cancer. In some embodiments, the cancer for treatment is cervical cancer. In some embodiments, the method of treating cancer further comprises the step of treating the patient with a non-myeloablative lymphodepletion regimen prior to administering the third population of TILs to the patient. In some embodiments, the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days. In some embodiments, the high dose IL-2 regimen comprises 600,000 or 720,000 IU/kg of aldesleukin, or a biosimilar or variant thereof, administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

[001475] In some embodiments, the present disclosure provides a method of treating a subject with cancer comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising: (i) obtaining a first population of TILs from a fine needle aspirate (FNA) or a small biopsy obtained from a tumor in a patient; (ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a small biopsy; (iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and (iv) administering a therapeutically effective dosage of the third population of TILs

to the patient. In some embodiments, the IL-2 is present at an initial concentration of about 3000 IU/mL and OKT-3 antibody is present at an initial concentration of about 30 ng/mL in the second cell culture medium. In some embodiments, the first expansion is performed using a gas permeable container. In some embodiments, the second expansion is performed using a gas permeable container. In some embodiments, the ratio of the second population of TILs to the population of APCs in the rapid expansion is between 1 to 80 and 1 to 400. In some embodiments, the ratio of the second population of TILs to the population of APCs in the rapid expansion is about 1 to 300. In some embodiments, the APCs are PMBCs. In some embodiments, the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian. In some embodiments, the cancer to be treated is selected from the group consisting of pancreatic, melanoma, breast, and ovarian. In some embodiments, the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma. In some embodiments, the cancer to be treated is selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma. In some embodiments, the method of treating cancer further comprises the step of treating the patient with a non-myeloablative lymphodepletion regimen prior to administering the third population of TILs to the patient. In some embodiments, the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days. In some embodiments, the high dose IL-2 regimen comprises 600,000 or 720,000 IU/kg of aldesleukin, or a biosimilar or variant thereof, administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

XVII. Exemplary Embodiments

1. A method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(i) obtaining a first population of TILs from a fine needle aspirate (FNA) or a small biopsy from a tumor in a patient;

(ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is

performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy; and

(iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs.

2. The method according to claim 1, wherein the method further comprises:

(iv) performing an additional second expansion by supplementing the cell culture medium of the third population of TILs with additional IL-2, additional OKT-3, and additional APCs, wherein the additional second expansion is performed for at least 14 days to obtain a larger therapeutic population of TILs than obtained in step (iii), wherein the larger therapeutic population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the third population of TILs.

3. The method according to claim 2, wherein after step (iii), the cells are removed from the cell culture and cryopreserved in a storage medium prior to performing step (iv).

4. The method according to claim 3, wherein the cells are thawed prior to performing step (iv).

5. The method according to any of the preceding claims, wherein step (iv) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

6. The method of according to any of the preceding claims, wherein steps (i) through (iii) or (iv) are performed within a period of about 21 days to about 33 days.

7. The method according to any of the preceding claims, wherein steps (i) through (iii) or (iv) are performed within a period of about 21 days to about 30 days.

8. The method according to any of the preceding claims, wherein steps (i) through (iii) or (iv) are performed within a period of about 21 days to about 28 days.

9. The method according to any of the preceding claims, wherein steps (i) through (iii) or (iv) are performed within about 28 days.

10. The method according to any of the preceding claims, wherein the cells from steps (iii) or (iv) express CD4, CD8, and TCR α β at levels similar to freshly harvested cells.

11. The method according to claim 1, wherein the APCs are peripheral blood mononuclear cells (PBMCs).

12. The method according to claim 11, wherein the PBMCs are added to the cell culture on any of days 3 through 19 in step (ii) and/or any of days 11 through 14 in step (iii).

13. The method according to claims 2 to 12, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs in step (iv) exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

14. The method according to claim 13, wherein the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

15. The method according to any of the preceding claims, wherein the APCs are artificial APCs (aAPCs) or autologous APCs.

16. The method according to any of the preceding claims, wherein the therapeutic population of TILs are infused into a patient.

17. The method according to claim 1, wherein the first expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

18. The method according to claim 1, wherein the second expansion in step (iii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

19. The method according to claim 2, wherein the additional second expansion in step (iv) is performed by further supplementing the cell culture medium of the third population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

20. The method according to claim 1, wherein the FNA in step (i) comprises at least 400,000 TILs.

21. The method according to claim 1, wherein the core biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian.

22. The method according to claim 1, wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

23. The method according to claim 22, wherein the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the patient has previously undergone surgical treatment.

24. The method according to claim 1, wherein the TILs in step (i) are obtained from a FNA.

25. The method according to claim 1, wherein the FNA is obtained using a 25-18 gauge needle.

26. The method according to claim 1, wherein the TILs in step (i) are obtained from a core biopsy.

27. The method according to claim 1, wherein the core biopsy is obtained using a 16-11 gauge needle.

28. The method according to claim 1, wherein step (iii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

29. The method according to claim 28, where the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

30. A method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(i) performing a first expansion by culturing a first population of TILs from a fine needle aspirate (FNA) or a core biopsy from a tumor in a patient in a cell culture medium comprising IL-2 to obtain a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by

about 10 days to about 19 days when the first population of TILs is from a core biopsy; and

(ii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs) to obtain a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs.

31. The method of claim 30, wherein the method further comprises:

(iii) performing an additional second expansion of the third population of TILs by supplementing the cell culture medium of the third population of TILs with additional IL-2, additional OKT-3, and additional APCs, wherein the additional second expansion is performed for at least 14 days to obtain a larger therapeutic population of TILs than obtained in step (ii), wherein the larger therapeutic population of TILs exhibits an increased subpopulation of effector T cells and/or central memory T cells relative to the third population of TILs.

32. The method according to claim 31, wherein the cells from the cell culture medium in step (ii) are removed and cryopreserved in a storage medium prior to step (iii).

33. The method according to claim 32, wherein the cells are thawed prior to step (iii).

34. The method according to any of claims 30 to 33, wherein the APCs are artificial APCs (aAPCs) or autologous APCs.

35. The method according to any of claims 30 to 34, wherein the therapeutic population of TILs are infused into a patient.

36. The method according to claim 30, wherein the first expansion in step (i) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

37. The method according to claim 30, wherein the second expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

38. The method according to claim 30, wherein the additional second expansion in step (iii) is performed by further supplementing the cell culture medium of the third population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

39. The method according to claim 30, wherein the FNA in step (i) comprises at least 400,000 TILs.

40. The method according to claim 30, wherein the core biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian.

41. The method according to claim 30, wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

42. The method according to claim 41, wherein the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the patient has previously undergone surgical treatment.

43. The method according to claim 30, wherein the TILs in step (i) are obtained from a FNA.

44. The method according to claim 30, wherein the FNA is obtained using a 25-18 gauge needle.

45. The method according to claim 30, wherein the TILs in step (i) are obtained from a core biopsy.

46. The method according to claim 30, wherein the core biopsy is obtained using a 16-11 gauge needle.

47. The method according to claim 30, wherein step (ii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

48. The method according to claim 47, where the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

49. The method according to any of claims 30 to 48, wherein the effector T cells and/or central memory T cells exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased

CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

50. The method according to claim 49, wherein the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

51. A method for treating a subject with cancer comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising:

(i) obtaining a first population of TILs from a fine needle aspirate (FNA) or a core biopsy obtained from a tumor in a patient;

(ii) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy;

(iii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and

(iv) administering a therapeutically effective dosage of the third population of TILs to the patient.

52. The method of claim 51, wherein the method further comprises prior to step (iv) a step of performing an additional second expansion by supplementing the cell culture medium of the third population of TILs with additional IL-2, additional OKT-3, and additional APCs, wherein the additional second expansion is performed for at least 14 days to obtain a larger therapeutic population of TILs than obtained in step (iii), wherein the larger therapeutic population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the third population of TILs.

53. The method according to claim 51, wherein after step (ii) the cells are removed from the cell culture medium and cryopreserved in a storage medium prior to the additional second expansion of claim 52.

54. The method according to claim 51, wherein the cells are thawed prior to the additional second expansion of claim 52.

55. The method according to claim 51, wherein step (iii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

56. The method according to claims 51 to 55, wherein the APCs are artificial APCs (aAPCs) or autologous APCs.

57. The method according to claim 51, wherein the first expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

58. The method according to claim 51, wherein the second expansion in step (iii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

59. The method according to claim 52, wherein the additional second expansion is performed by further supplementing the cell culture medium of the third population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

60. The method according to claim 51, wherein the FNA in step (i) comprises at least 400,000 TILs.

61. The method according to claim 51, wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

62. The method according to claim 61, wherein the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

63. The method according to claim 51, wherein the TILs in step (i) are obtained from a FNA.

64. The method according to claim 51, wherein the FNA is obtained using a 25-18 gauge needle.

65. The method according to claim 51, wherein the TILs in step (i) are obtained from a core biopsy.

66. The method according to claim 51, wherein the core biopsy is obtained using a 16-11 gauge needle.

67. The method according to claim 51, wherein step (iii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

68. The method according to claim 67, where the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

69. A method according to any of claims 51 to 68, wherein the effector T cells and/or central memory T cells exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

70. The method according to claim 69, wherein the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

71. The method according to any of claims 51 to 70, wherein the cancer is selected from the group consisting of melanoma, cervical cancer, head and neck cancer, glioblastoma, ovarian cancer, sarcoma, pancreatic cancer, bladder cancer, breast cancer, triple negative breast cancer, and non-small cell lung carcinoma.

72. A method for treating a subject with cancer comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising:

(i) performing a first expansion by culturing a first population of TILs from a fine needle aspirate (FNA) or a core biopsy from a tumor in a patient in a cell culture medium comprising IL-2 to obtain a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core

biopsy;

(ii) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs) to obtain a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for about 11 days to about 14 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and

(iii) administering a therapeutically effective dosage of the therapeutic population of TILs to the patient.

73. The method of claim 72, wherein the method further comprises prior to step (iii) a step of performing an additional second expansion by supplementing the cell culture medium of the third population of TILs with additional IL-2, additional OKT-3, and additional APCs, wherein the additional second expansion is performed for at least 14 days to obtain a larger therapeutic population of TILs than obtained in step (ii), wherein the larger therapeutic population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the third population of TILs.

74. A method according to claim 72, wherein the cells from the cell culture medium in step (ii) are removed and cryopreserved in a storage medium prior to the additional second expansion in claim 73.

75. The method according to claim 74, wherein the cells are thawed prior to the additional second expansion in claim 73.

76. The method according to claim 74, wherein the cells are thawed prior to step (iii).

77. The method according to any of claims 72 to 76, wherein the APCs are artificial APCs (aAPCs) or autologous APCs.

78. The method according to any of claims 72 to 77, wherein the APCs are peripheral blood mononuclear cells (PBMCs).

79. The method according to any of claims 72 to 78, wherein the therapeutic population of TILs are infused into a patient.

80. The method according to claim 72, wherein the first expansion in step (i) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

81. The method according to claim 72, wherein the second expansion in step (iii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

82. The method according to claim 81, wherein the additional second expansion is performed by further supplementing the cell culture medium of the third population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

83. The method according to claim 72, wherein the FNA in step (i) comprises at least 400,000 TILs.

84. The method according to claim 72, wherein the core biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian.

85. The method according to claim 72, wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

86. The method according to claim 85, wherein the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

87. The method according to claim 72, wherein the TILs in step (i) are obtained from a FNA.

88. The method according to claim 72, wherein the FNA is obtained using a 25-18 gauge needle.

89. The method according to claim 72, wherein the TILs in step (i) are obtained from a core biopsy.

90. The method according to claim 72, wherein the core biopsy is obtained using a 16-11 gauge needle.

91. The method according to claim 72, wherein step (ii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs.

92. The method according to claim 91, where the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

93. The method according to any of claims 72 to 92, wherein the effector T cells and/or central memory T cells exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the third population of cells.

94. The method according to claim 93, wherein the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

95. A method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(a) obtaining a first population of TILs from a fine needle aspirate (FNA) or a core biopsy obtained from a tumor in a patient;

(b) adding the first population into a closed system;

(c) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy, wherein the first expansion is performed in a closed container providing a first gas-permeable surface area, wherein the first expansion is performed for about 3-14 days to obtain the second population of TILs, wherein the second population of TILs is at least 25-fold greater in number than the first population of TILs, and wherein the transition from step (b) to step (c) occurs without opening the system;

(d) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the second expansion is performed for about 11 days to about-14 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs which comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the second expansion is performed in a closed container providing a second gas-permeable surface area, and

wherein the transition from step (c) to step (d) occurs without opening the system;

(e) harvesting the therapeutic population of TILs obtained from step (d), wherein the transition from step (d) to step (e) occurs without opening the system; and

(f) transferring the harvested TIL population from step (e) to an infusion bag, wherein the transfer from step (e) to (f) occurs without opening the system.

96. The method according to claim 95, further comprising the step of cryopreserving the infusion bag comprising the harvested TIL population in step (f) using a cryopreservation process.

97. The method according to claim 96, wherein the cryopreservation process is performed using a 1:1 ratio of harvested TIL population to CS10 media.

98. The method according to claim 96, wherein the antigen-presenting cells are peripheral blood mononuclear cells (PBMCs).

99. The method according to claim 98, wherein the PBMCs are irradiated and allogeneic.

100. The method according to claim 98, wherein the PBMCs are added to the cell culture on any of days 3 through 19 in step (c) and/or any of days 11 through 14 in step (d).

101. The method according to claim 95, wherein the antigen-presenting cells are artificial antigen-presenting cells (aAPCs) or autologous APCs.

102. The method according to any of claims 95 to 101, wherein the therapeutic population of TILs are infused into a patient.

103. The method according to claim 95, wherein the first expansion in step (c) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

104. The method according to claim 95, wherein the second expansion in step (d) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

105. The method according to claim 95, wherein the FNA in step (a) comprises at least 400,000 TILs.

106. The method according to claim 95, wherein the core biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast and ovarian.

107. The method according to claim 95, wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

108. The method according to claim 107, wherein the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

109. The method according to claim 95, wherein the TILs in step (a) are obtained from a FNA.

110. The method according to claim 95, wherein the FNA is obtained using a 25-18 gauge needle.

111. The method according to claim 95, wherein the TILs in step (a) are obtained from a core biopsy.

112. The method according to claim 95, wherein the core biopsy is obtained using a 16-11 gauge needle.

113. The method according to claim 95, wherein the harvesting in step (e) is performing using a LOVO cell processing system.

114. The method according to claim 95, wherein the cell culture medium is provided in a container selected from the group consisting of a G-container and a Xuri cellbag.

115. The method according to claim 95, wherein the infusion bag in step (f) is a hypothermosol infusion bag.

116. The method according to claim 95, wherein steps (a) through (f) are performed within a period of about 21 days to about 33 days.

117. The method according to claim 95, wherein steps (a) through (f) are performed within a period of about 21 days to about 30 days.

118. The method according to claim 95, wherein steps (a) through (f) are performed within a period of about 21 days to about 28 days.

119. The method according to claim 95, wherein steps (a) through (f) are performed in 22 days or less.

120. The method according to claim 96, wherein steps (a) through (f) and cryopreservation are performed in 22 days or less.

121. The method according to any one of claims 95 to 120, wherein the therapeutic population of TILs harvested in step (e) comprises sufficient TILs for a therapeutically effective dosage of the TILs.

122. The method according to claim 121, wherein the number of TILs sufficient for a therapeutically effective dosage is from about 2.3×10^{10} to about 13.7×10^{10}.

123. The method according to any one of claims 95 to 122, wherein steps (b) through (e) are performed in a single container, wherein performing steps (b) through (e) in a single container results in an increase in TIL yield per resected tumor as compared to performing steps (b) through (e) in more than one container.

124. The method according to any one of claims 95 to 123, wherein the antigen-presenting cells are added to the TILs during the second period in step (d) without opening the system.

125. The method according to any one of claims 95 to 124, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit one or more characteristics selected from the group consisting of expressing CD27+, expressing CD28+, longer telomeres, increased CD57 expression, and decreased CD56 expression relative to effector T cells, and/or central memory T cells obtained from the second population of cells.

126. The method according to any one of claims 95 to 125, wherein the effector T cells and/or central memory T cells obtained from the third population of TILs exhibit increased CD57 expression and decreased CD56 expression relative to effector T cells and/or central memory T cells obtained from the second population of cells.

127. The method according to any one of claims 95 to 126, wherein the risk of microbial contamination is reduced as compared to an open system.

128. The method according to any one of claims 95 to 128, wherein the TILs from step (g) are infused into a patient.

129. A method for treating a subject with cancer, the method comprising administering expanded tumor infiltrating lymphocytes (TILs) comprising:

(a) obtaining a first population of TILs from a fine needle aspirate (FNA) or a core biopsy from a tumor resected from a subject;

(b) adding the first population into a closed system;

(c) performing a first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, , wherein the cell culture medium is supplemented with OKT-3 at day 3, wherein the first expansion is performed for about 3 days to about 19 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5×10^7 TILs by about 10 days to about 19 days when the first population of TILs is from a core biopsy, wherein the first expansion is performed in a closed container providing a first gas-permeable surface area, wherein the first expansion is performed for about 3-14 days to obtain the second population of TILs, wherein the second population of TILs is at least 25-fold greater in number than the first population of TILs, and wherein the transition from step (b) to step (c) occurs without opening the system;

(d) performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the second expansion is performed for about 7-14 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs which comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the second expansion is performed in a closed container providing a second gas-permeable surface area, and wherein the transition from step (c) to step (d) occurs without opening the system;

(e) harvesting the therapeutic population of TILs obtained from step (d), wherein the transition from step (d) to step (e) occurs without opening the system; and

(f) transferring the harvested TIL population from step (e) to an infusion bag, wherein the transfer from step (e) to (f) occurs without opening the system;

(g) optionally cryopreserving the infusion bag comprising the harvested TIL population from step (f) using a cryopreservation process; and

(h) administering a therapeutically effective dosage of the third population of TILs from the infusion bag in step (g) to the patient.

130. The method according to claim 129, wherein the therapeutic population of TILs harvested in step (e) comprises sufficient TILs for administering a therapeutically effective dosage of the TILs in step (h).

131. The method according to claim 129 or 130, wherein the APCs are artificial APCs (aAPCs) or autologous APCs.

132. The method according to any of claims 129 to 131, wherein the therapeutic population of TILs are infused into a patient.

133. The method according to claim 129, wherein the first expansion in step (c) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

134. The method according to claim 129, wherein the second expansion in step (d) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

135. The method according to claim 129, wherein the FNA in step (a) comprises at least 400,000 TILs.

136. The method according to claim 129, wherein the core biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast and ovarian.

137. The method according to claim 129, wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma.

138. The method according to claim 137, wherein the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

139. The method according to claim 129, wherein the TILs in step (i) are obtained from a FNA.

140. The method according to claim 139, wherein the FNA is obtained using a 25-18 gauge needle.

141. The method according to claim 129, wherein the TILs in step (i) are obtained from a core biopsy.

142. The method according to claim 141, wherein the core biopsy is obtained using a 16-11 gauge needle.

143. The method according to claim 129, wherein the number of TILs sufficient for administering a therapeutically effective dosage in step (h) is from about 2.3×10^{10} to about 13.7×10^{10}.

144. The method according to claim 129, wherein the antigen presenting cells (APCs) are PBMCs.

145. The method according to claim 144, wherein the PBMCs are added to the cell culture on any of days 3 through 19 in step (c) and/or any of days 9 through 14 in step (d).

146. The method according to any of claims 129 to 145, wherein prior to administering a therapeutically effective dosage of TIL cells in step (h), a non-myeloablative lymphodepletion regimen has been administered to the patient.

147. The method according to claim 146, where the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

148. The method according to any of claims 129 to 147, further comprising the step of treating the patient with a high-dose IL-2 regimen starting on the day after administration of the TIL cells to the patient in step (h).

149. The method according to claim 148, wherein the high-dose IL-2 regimen comprises 600,000 or 720,000 IU/kg administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

150. The method according to any of the claims 129 to 149, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit one or more characteristics selected from the group consisting of expressing CD27+, expressing CD28+, longer telomeres, increased CD57 expression, and decreased CD56 expression relative to effector T cells, and/or central memory T cells obtained from the second population of cells.

151. The method according to any of claims 129 to 150, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit increased CD57 expression and decreased CD56 expression relative to effector T cells and/or central memory T cells obtained from the second population of cells.

152. A method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

(a) obtaining a first population of TILs from a fine needle aspirate (FNA), a small biopsy, or a core biopsy from a tumor in a patient;

(b) performing a priming first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2, OKT-3, and antigen presenting cells (APCs) to produce a second population of TILs, wherein the priming first expansion is performed for a first period of about 1 to 7 days in a container comprising a first gas-permeable surface area to obtain the second population of TILs, wherein the second population of TILs is greater in number than the first population of TILs;

(c) performing a rapid second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and APCs, to produce a third population of TILs, wherein the rapid second expansion is performed for a second period of about 1 to about 11 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs; and

(d) harvesting the therapeutic population of TILs from step (c).

EXAMPLES

[001476] The embodiments encompassed herein are now described with reference to the following examples. These examples are provided for the purpose of illustration only and the disclosure encompassed herein should in no way be construed as being limited to these examples, but rather should be construed to encompass any and all variations which become evident as a result of the teachings provided herein.

EXAMPLE 1: PREPARATION OF MEDIA FOR PRE-REP AND REP PROCESSES

[001477] This Example describes the procedure for the preparation of tissue culture media for use in protocols involving the culture of tumor infiltrating lymphocytes (TIL) derived from various tumor types including, but not limited to, metastatic melanoma, head and neck squamous cell carcinoma (HNSCC), ovarian carcinoma, triple-negative breast carcinoma, and lung adenocarcinoma. This media can be used for preparation of any of the TILs described in the present application and Examples.

Preparation of CM1

[001478] Removed the following reagents from cold storage and warmed them in a 37°C water bath: (RPMI1640, Human AB serum, 200mM L-glutamine). Prepared CM1 medium according to Table 32 below by adding each of the ingredients into the top section of a 0.2 µm filter unit appropriate to the volume to be filtered. Store at 4°C.

TABLE 32: Preparation of CM1

Ingredient	Final concentration	Final Volume 500 ml	Final Volume IL
RPMI1640	NA	450 mL	900 mL
Human AB serum, heat-inactivated 10%	50 mL	100 mL	
200 mM L-glutamine	2 mM	5 mL	10 mL
55 mM BME	55 µM	0.5 mL	1 mL
50 mg/mL gentamicin sulfate	50 µg/mL	0.5 mL	1 mL

[001479] On the day of use, prewarmed required amount of CM1 in 37°C water bath and add 6000 IU/mL IL-2.

[001480] Additional supplementation - as needed according to Table 33.

TABLE 33: Additional supplementation of CM1, as needed.

Supplement	**Stock concentration**	**Dilution**	**Final concentration**
GlutaMAX™	200mM	1:100	2mM
Penicillin/streptomycin	10,000 U/mL penicillin	1:100	100 U/mL penicillin 100 µg/mL streptomycin

	10,000 µg/mL streptomycin		
Amphotericin B	250 µg/mL	1:100	2.5 µg/mL

Preparation of CM2

[001481] Removed prepared CM1 from refrigerator or prepare fresh CM1 as per Section 7.3 above. Removed AIM-V® from refrigerator and prepared the amount of CM2 needed by mixing prepared CM1 with an equal volume of AIM-V® in a sterile media bottle. Added 3000 IU/mL IL-2 to CM2 medium on the day of usage. Made sufficient amount of CM2 with 3000 IU/mL IL-2 on the day of usage. Labeled the CM2 media bottle with its name, the initials of the preparer, the date it was filtered/prepared, the two-week expiration date and store at 4°C until needed for tissue culture.

Preparation of CM3

[001482] Prepared CM3 on the day it was required for use. CM3 was the same as AIM-V® medium, supplemented with 3000 IU/ml IL-2 on the day of use. Prepared an amount of CM3 sufficient to experimental needs by adding IL-2 stock solution directly to the bottle or bag of AIM-V. Mixed well by gentle shaking. Label bottle with "3000 IU/mL IL-2" immediately after adding to the AIM-V. If there was excess CM3, stored it in bottles at 4°C labeled with the media name, the initials of the preparer, the date the media was prepared, and its expiration date (7 days after preparation). Discarded media supplemented with IL-2 after 7 days storage at 4°C.

Preparation of CM4

[001483] CM4 was the same as CM3, with the additional supplement of 2mM GlutaMAX™ (final concentration). For every 1 L of CM3, added 10 ml of 200 mM GlutaMAX™. Prepared an amount of CM4 sufficient to experimental needs by adding IL-2 stock solution and GlutaMAX™ stock solution directly to the bottle or bag of AIM-V. Mixed well by gentle shaking. Labeled bottle with "3000 IL/ml IL-2 and GlutaMAX" immediately after adding to the AIM-V. If there was excess CM4, stored it in bottles at 4°C labeled with the media name, "GlutaMAX", and its expiration date (7 days after preparation). Discarded media supplemented with IL-2 after 7-days storage at 4°C.

EXAMPLE 2: USE OF IL-2, IL-15, AND IL-21 CYTOKINE COCKTAIL

[001484] This example describes the use of IL-2, IL-15, and IL-21 cytokines, which serve as additional T cell growth factors, in combination with the TIL process of the Examples.

[001485] Using the processes described herein, TILs were grown from colorectal, melanoma, cervical, triple negative breast, lung and renal tumors in presence of IL-2 in one arm of the experiment and, in place of IL-2, a combination of IL-2, IL-15, and IL-21 in another arm at the initiation of culture. At the completion of the pre-REP, cultures were assessed for expansion, phenotype, function (CD107a+ and IFNγ) and TCR Vβ repertoire. IL-15 and IL-21 are described elsewhere herein and in Gruijl, et al., IL-21 promotes the expansion of CD27+CD28+ tumor infiltrating lymphocytes with high cytotoxic potential and low collateral expansion of regulatory T cells, *Santegoets, S. J.,* J Transl Med., **2013,** *11:*37 (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3626797/).

[001486] The results showed that enhanced TIL expansion (>20%), in both $CD4^+$ and $CD8^+$ cells in the IL-2, IL-15, and IL-21 treated conditions were observed in multiple histologies relative to the IL-2 only conditions. There was a skewing towards a predominantly $CD8^+$ population with a skewed TCR Vβ repertoire in the TILs obtained from the IL-2, IL-15, and IL-21 treated cultures relative to the IL-2 only cultures. IFNγ and CD107a were elevated in the IL-2, IL-15, and IL-21 treated TILs, in comparison to TILs treated only IL-2.

EXAMPLE 3: QUALIFYING INDIVIDUAL LOTS OF GAMMA-IRRADIATED PERIPHERAL MONONUCLEAR CELLS

[001487] This Example describes a novel abbreviated procedure for qualifying individual lots of gamma-irradiated peripheral mononuclear cells (PBMCs, also known as MNC) for use as allogeneic feeder cells in the exemplary methods described herein.

[001488] Each irradiated MNC feeder lot was prepared from an individual donor. Each lot or donor was screened individually for its ability to expand TIL in the REP in the presence of purified anti-CD3 (clone OKT3) antibody and interleukin-2 (IL-2). In addition, each lot of feeder cells was tested without the addition of TIL.to verify that the received dose of gamma radiation was sufficient to render them replication incompetent.

Background

[001489] Gamma-irradiated, growth-arrested MNC feeder cells were required for REP of TIL. Membrane receptors on the feeder MNCs bind to anti-CD3 (clone OKT3) antibody and crosslink to TIL in the REP flask, stimulating the TIL to expand. Feeder lots were prepared from the leukapheresis of whole blood taken from individual donors. The leukapheresis product was subjected to centrifugation over Ficoll-Hypaque, washed, irradiated, and cryopreserved under GMP conditions.

[001490] It is important that patients who received TIL therapy not be infused with viable feeder cells as this can result in Graft-Versus-Host Disease (GVHD). Feeder cells are therefore growth-arrested by dosing the cells with gamma-irradiation, resulting in double strand DNA breaks and the loss of cell viability of the MNC cells upon reculture.

Evaluation Criteria and Experimental Set-Up

[001491] Feeder lots were evaluated on two criteria: 1) their ability to expand TIL in co-culture >100-fold and 2) their replication incompetency.

[001492] Feeder lots were tested in mini-REP format utilizing two primary pre-REP TIL lines grown in upright T25 tissue culture flasks. Feeder lots were tested against two distinct TIL lines, as each TIL line is unique in its ability to proliferate in response to activation in a REP. As a control, a lot of irradiated MNC feeder cells which has historically been shown to meet the criteria above was run alongside the test lots.

[001493] To ensure that all lots tested in a single experiment receive equivalent testing, sufficient stocks of the same pre-REP TIL lines were available to test all conditions and all feeder lots.

[001494] For each lot of feeder cells tested, there was a total of six T25 flasks: Pre-REP TIL line #1 (2 flasks); Pre-REP TIL line #2 (2 flasks); and Feeder control (2 flasks). Flasks containing TIL lines #1 and #2 evaluated the ability of the feeder lot to expand TIL. The feeder control flasks evaluated the replication incompetence of the feeder lot.

EXPERIMENTAL PROTOCOL

Day -2/3, Thaw of TIL lines

Prepared CM2 medium. Warmed CM2 in 37°C water bath. Prepared 40 mL of CM2 supplemented with 3000 IU/ml IL-2. Keep warm until use. Placed 20 mL of pre-warmed CM2 without IL-2 into each of two 50 mL conical tubes labeled with names of the TIL lines

used. Removed the two designated pre-REP TIL lines from LN_2 storage and transferred the vials to the tissue culture room. Thawed vials by placing them inside a sealed zipper storage bag in a 37ºC water bath until a small amount of ice remains.

[001495] Using a sterile transfer pipet, immediately transferred the contents of vial into the 20 mL of CM2 in the prepared, labeled 50 mL conical tube. QS to 40 mL using CM2 without IL-2 to wash cells. Centrifuged at 400 x CF for 5 minutes. Aspirated the supernatant and resuspend in 5 mL warm CM2 supplemented with 3000 IU/mL IL-2.

[001496] Removed small aliquot (20 µL) in duplicate for cell counting using an automated cell counter. Record the counts. While counting, placed the 50 mL conical tube with TIL cells into a humidified 37ºC, 5% CO_2 incubator, with the cap loosened to allow for gas exchange. Determined cell concentration and diluted TIL to 1×10^6 cells/mL in CM2 supplemented with IL-2 at 3000 IU/mL.

[001497] Cultured in 2 mL/well of a 24-well tissue culture plate in as many wells as needed in a humidified 37ºC incubator until Day 0 of the mini-REP. Cultured the different TIL lines in separate 24-well tissue culture plates to avoid confusion and potential cross-contamination.

Day 0, initiate Mini-REP

[001498] Prepared enough CM2 medium for the number of feeder lots to be tested. (*e.g.*, for testing 4 feeder lots at one time, prepared 800 mL of CM2 medium). Aliquoted a portion of the CM2 prepared above and supplemented it with 3000 IU/mL IL-2 for the culturing of the cells. (*e.g.*, for testing 4 feeder lots at one time, prepare 500 mL of CM2 medium with 3000 IU/mL IL-2).

[001499] Working with each TIL line separately to prevent cross-contamination, removed the 24-well plate with TIL culture from the incubator and transferred to the BSC.

[001500] Using a sterile transfer pipet or 100-1000 µL Pipettor and tip, removed about 1 mL of medium from each well of TIL to be used and place in an unused well of the 24-well tissue culture plate.

[001501] Using a fresh sterile transfer pipet or 100-1000 µL Pipettor and tip, mixed remaining medium with TIL in wells to resuspend the cells and then transferred the cell suspension to a 50 mL conical tube labeled with the TIL name and recorded the volume.

[001502] Washed the wells with the reserved media and transferred that volume to the same 50 mL conical tube. Spun the cells at 400 x CF to collect the cell pellet. Aspirated off the

media supernatant and resuspend the cell pellet in 2-5 mL of CM2 medium containing 3000 IU/mL IL-2, volume to be used based on the number of wells harvested and the size of the pellet – volume should be sufficient to ensure a concentration of $>1.3 \times 10^6$ cells/mL.

[001503] Using a serological pipet, mixed the cell suspension thoroughly and recorded the volume. Removed 200 µL for a cell count using an automated cell counter. While counting, placed the 50 mL conical tube with TIL cells into a humidified, 5% CO_2, 37°C incubator, with the cap loosened to allow gas exchange. Recorded the counts.

[001504] Removed the 50 ml conical tube containing the TIL cells from the incubator and resuspend them cells at a concentration of 1.3×10^6 cells/mL in warm CM2 supplemented with 3000 IU/mL IL-2. Returned the 50 mL conical tube to the incubator with a loosened cap.

[001505] Repeated steps above for the second TIL line.

[001506] Just prior to plating the TIL into the T25 flasks for the experiment, TIL were diluted 1:10 for a final concentration of 1.3×10^5 cells/mL as per below.

Prepare MACS GMP CD3 pure (OKT3) working solution

[001507] Took out stock solution of OKT3 (1 mg/mL) from 4°C refrigerator and placed in BSC. A final concentration of 30 ng/mL OKT3 was used in the media of the mini-REP.

[001508] 600 ng of OKT3 were needed for 20 ml in each T25 flask of the experiment; this was the equivalent of 60 µL of a 10 µg/mL solution for each 20 mL, or 360 µL for all 6 flasks tested for each feeder lot.

[001509] For each feeder lot tested, made 400 µL of a 1:100 dilution of 1 mg/mL OKT3 for a working concentration of 10 µg/mL (*e.g.*, for testing 4 feeder lots at one time, make 1600 µL of a 1:100 dilution of 1 mg/mL OKT3: 16 µL of 1 mg/mL OKT3 + 1.584 mL of CM2 medium with 3000 IU/mL IL-2.)

Prepare T25 flasks

[001510] Labeled each flask and filled flask with the CM2 medium prior to preparing the feeder cells. Placed flasks into 37°C humidified 5% CO_2 incubator to keep media warm while waiting to add the remaining components. Once feeder cells were prepared, the components will be added to the CM2 in each flask.

TABLE 34: Solutions

Component	Volume in co-culture flasks	Volume in control (feeder only) flasks
CM2 + 3000 IU/ml IL-2	18ml	19ml
MNC: 1.3 x 10^7/ml in CM2 + 3000U IL-2 (final concentration 1.3 x 10^7/flask)	1ml	1ml
OKT3: 10µg/ml in CM2 + 3000U IL-2	60µl	60µl
TIL: 1.3 x 10^5/ml in CM2 with 3000U of IL-2 (final concentration 1.3 x 10^5/flask)	1ml	0

Prepare Feeder Cells

[001511] A minimum of 78 × 10^6 feeder cells were needed per lot tested for this protocol. Each 1 ml vial frozen by SDBB had 100 × 10^6 viable cells upon freezing. Assuming a 50% recovery upon thaw from LN_2 storage, it was recommended to thaw at least two 1 mL vials of feeder cells per lot giving an estimated 100 × 10^6 viable cells for each REP. Alternately, if supplied in 1.8 ml vials, only one vial provided enough feeder cells.

[001512] Before thawing feeder cells, pre-warmed approximately 50 mL of CM2 without IL-2 for each feeder lot to be tested. Removed the designated feeder lot vials from LN_2 storage, placed in zipper storage bag, and place on ice. Thawed vials inside closed zipper storage bag by immersing in a 37°C water bath. Removed vials from zipper bag, spray or wipe with 70% EtOH and transferred vials to BSC.

[001513] Using a transfer pipet immediately transferred the contents of feeder vials into 30 mL of warm CM2 in a 50 mL conical tube. Washed vial with a small volume of CM2 to remove any residual cells in the vial. Centrifuged at 400 x CF for 5 minutes. Aspirated the supernatant and resuspended in 4 mL warm CM2 plus 3000 IU/mL IL-2. Removed 200 µL for cell counting using the Automated Cell Counter. Recorded the counts.

[001514] Resuspended cells at 1.3 × 10^7 cells/mL in warm CM2 plus 3000 IU/mL IL-2. Diluted TIL cells from 1.3 × 10^6 cells/mL to 1.3 × 10^5 cells/mL.

Setup Co-Culture

[001515] Diluted TIL cells from 1.3 × 10^6 cells/mL to 1.3 × 10^5 cells/mL. Added 4.5 ml of CM2 medium to a 15 mL conical tube. Removed TIL cells from incubator and resuspended well using a 10 mL serological pipet. Removed 0.5 mL of cells from the 1.3 × 10^6 cells/mL TIL suspension and added to the 4.5 mL of medium in the 15 mL conical tube. Returned TIL stock vial to incubator. Mixed well. Repeated for the second TIL line.

Transferred flasks with pre-warmed media for a single feeder lot from the incubator to the BSC. Mixed feeder cells by pipetting up and down several times with a 1 mL pipet tip and transferred 1 mL (1.3×10^7 cells) to each flask for that feeder lot. Added 60 µL of OKT3 working stock (10 µg/mL) to each flask. Returned the two control flasks to the incubator.

[001516] Transferred 1 mL (1.3×10^5) of each TIL lot to the correspondingly labeled T25 flask. Returned flasks to the incubator and incubate upright. Did not disturb until Day 5.

[001517] Repeated for all feeder lots tested.

Day 5, Media change

Prepared CM2 with 3000 IU/mL IL-2. 10 ml is needed for each flask. With a 10 mL pipette, transferred 10 mL warm CM2 with 3000 IU/mL IL-2 to each flask. Returned flasks to the incubator and incubated upright until Day 7. Repeated for all feeder lots tested.

Day 7, Harvest

[001518] Removed flasks from the incubator and transfer to the BSC, care as taken not to disturb the cell layer on the bottom of the flask. Without disturbing the cells growing on the bottom of the flasks, removed 10 ml of medium from each test flask and 15 ml of medium from each of the control flasks.

[001519] Using a 10 mL serological pipet, resuspended the cells in the remaining medium and mix well to break up any clumps of cells. After thoroughly mixing cell suspension by pipetting, removed 200 µL for cell counting. Counted the TIL using the appropriate standard operating procedure in conjunction with the automatic cell counter equipment. Recorded counts in Day 7.

[001520] Repeated for all feeder lots tested.

[001521] Feeder control flasks were evaluated for replication incompetence and flasks containing TIL were evaluated for fold expansion from Day 0 according to Table TT below.

Day 7, Continuation of Feeder Control Flasks to Day 14

[001522] After completing the Day 7 counts of the feeder control flasks, added 15 mL of fresh CM2 medium containing 3000 IU/mL IL-2 to each of the control flasks. Returned the control flasks to the incubator and incubated in an upright position until Day 14.

Day 14, Extended Non-proliferation of Feeder Control Flasks

[001523] Removed flasks from the incubator and transfer to the BSC, care was taken not to disturb the cell layer on the bottom of the flask. Without disturbing the cells growing on the bottom of the flasks, removed approximately 17 ml of medium from each control flasks. Using a 5 mL serological pipet, resuspended the cells in the remaining medium and mixed well to break up any clumps of cells. Recorded the volumes for each flask.

[001524] After thoroughly mixing cell suspension by pipetting, removed 200 µl for cell counting. Counted the TIL using the appropriate standard operating procedure in conjunction with the automatic cell counter equipment. Recorded counts.

[001525] Repeated for all feeder lots tested.

RESULTS AND ACCEPTANCE CRITERIA

Results

[001526] The dose of gamma irradiation was sufficient to render the feeder cells replication incompetent. All lots were expected to meet the evaluation criteria and also demonstrated a reduction in the total viable number of feeder cells remaining on Day 7 of the REP culture compared to Day 0.

[001527] All feeder lots were expected to meet the evaluation criteria of 100-fold expansion of TIL growth by Day 7 of the REP culture.

[001528] Day 14 counts of Feeder Control flasks were expected to continue the non-proliferative trend seen on Day 7.

Acceptance Criteria

[001529] The following acceptance criteria were met for each replicate TIL line tested for each lot of feeder cells

[001530] Acceptance was two-fold, as follows (outlined in Table 35 below).

TABLE 35: Acceptance Criteria

Test	Acceptance criteria
Irradiation of MNC / Replication incompetence	No growth observed at 7 and 14 days
TIL expansion	At least a 100-fold expansion of each TIL (minimum of 1.3×10^7 viable cells)

[001531] Evaluated whether the dose of radiation was sufficient to render the MNC feeder cells replication incompetent when cultured in the presence of 30 ng/mL OKT3 antibody and

3000 IU/mL IL-2. Replication incompetence was evaluated by total viable cell count (TVC) as determined by automated cell counting on Day 7 and Day 14 of the REP.

[001532] Acceptance criteria was "No Growth," meaning the total viable cell number has not increased on Day 7 and Day 14 from the initial viable cell number put into culture on Day 0 of the REP.

[001533] Evaluated the ability of the feeder cells to support TIL expansion. TIL growth was measured in terms of fold expansion of viable cells from the onset of culture on Day 0 of the REP to Day 7 of the REP. On Day 7, TIL cultures achieved a minimum of 100-fold expansion, (*i.e.*, greater than 100 times the number of total viable TIL cells put into culture on REP Day 0), as evaluated by automated cell counting.

Contingency Testing of MNC Feeder Lots that do not meet acceptance criteria

[001534] In the event that an MNC feeder lot did not meet the either of the acceptance criteria outlined above, the following steps will be taken to retest the lot to rule out simple experimenter error as its cause.

[001535] If there are two or more remaining satellite testing vials of the lot, then the lot was retested. If there were one or no remaining satellite testing vials of the lot, then the lot was failed according to the acceptance criteria listed above.

[001536] In order to be qualified, the lot in question and the control lot had to achieve the acceptance criteria above. Upon meeting these criteria, the lot was then released for use.

EXAMPLE 4: QUALIFYING INDIVIDUAL LOTS OF GAMMA-IRRADIATED PERIPHERAL BLOOD MONONUCLEAR CELLS

[001537] This Example describes a novel abbreviated procedure for qualifying individual lots of gamma-irradiated peripheral blood mononuclear cells (PBMC) for use as allogeneic feeder cells in the exemplary methods described herein. This example provides a protocol for the evaluation of irradiated PBMC cell lots for use in the production of clinical lots of TIL. Each irradiated PBMC lot was prepared from an individual donor. Over the course of more than 100 qualification protocols, it has been shown that, in all cases, irradiated PBMC lots from SDBB (San Diego Blood Bank) expand TIL >100-fold on Day 7 of a REP. This modified qualification protocol was intended to apply to irradiated donor PBMC lots from SDBB which were then further tested to verify that the received dose of gamma radiation was

sufficient to render them replication incompetent. Once demonstrated that they maintained replication incompetence over the course of 14 days, donor PBMC lots were considered "qualified" for usage to produce clinical lots of TIL.

Background

[001538] Gamma-irradiated, growth-arrested PBMC were required for current standard REP of TIL. Membrane receptors on the PBMCs bind to anti-CD3 (clone OKT3) antibody and crosslink to TIL in culture, stimulating the TIL to expand. PBMC lots were prepared from the leukapheresis of whole blood taken from individual donors. The leukapheresis product was subjected to centrifugation over Ficoll-Hypaque, washed, irradiated, and cryopreserved under GMP conditions.

[001539] It is important that patients who received TIL therapy not be infused with viable PBMCs as this could result in Graft-Versus-Host Disease (GVHD). Donor PBMCs are therefore growth-arrested by dosing the cells with gamma-irradiation, resulting in double strand DNA breaks and the loss of cell viability of the PBMCs upon reculture.

Evaluation Criteria

[001540] 7.2.1 Evaluation criterion for irradiated PBMC lots was their replication incompetency.

Experimental Set-up

[001541] Feeder lots were tested in mini-REP format as if they were to be co-cultured with TIL, using upright T25 tissue culture flasks. Control lot: One lot of irradiated PBMCs, which had historically been shown to meet the criterion of 7.2.1, was run alongside the experimental lots as a control. For each lot of irradiated donor PBMC tested, duplicate flasks were run.

Experimental Protocol

Day 0

[001542] Prepared ~90 ml of CM2 medium for each lot of donor PBMC to be tested. Kept CM2 warm in 37°C water bath. Thawed an aliquot of 6×10^6 IU/mL IL-2. Returned the CM2 medium to the BSC, wiping with 70% EtOH prior to placing in hood. For each lot of PBMC tested, removed about 60 mL of CM2 to a separate sterile bottle. Added IL-2 from the thawed 6×10^6 IU/mL stock solution to this medium for a final concentration of 3000 IU/mL. Labeled this bottle as "CM2/IL2" (or similar) to distinguish it from the unsupplemented CM2.

Prepare OKT3

[001543] Took out the stock solution of anti-CD3 (OKT3) from the 4°C refrigerator and placed in the BSC. A final concentration of 30 ng/mL OKT3 was used in the media of the mini-REP. Prepared a 10 µg/mL working solution of anti-CD3 (OKT3) from the 1 mg/mL stock solution. Placed in refrigerator until needed.

[001544] For each PBMC lot tested, prepare 150 µL of a 1:100 dilution of the anti-CD3 (OKT3) stock. For example, for testing 4 PBMC lots at one time, prepare 600 µL of 10 µg/mL anti-CD3 (OKT3) by adding 6 µL of the 1 mg/mL stock solution to 594 µL of CM2 supplemented with 3000 IU/mL IL-2.

Prepare Flasks

[001545] Added 19 mL per flask of CM2/IL-2 to the labeled T25 flasks and placed flasks into 37°C, humidified, 5% CO_2 incubator while preparing cells.

Prepare Irradiate PBMC

[001546] Retrieved vials of PBMC lots to be tested from LN_2 storage. These were placed at -80°C or kept on dry ice prior to thawing. Placed 30 mL of CM2 (without IL-2 supplement) into 50 mL conical tubes for each lot to be thawed. Labeled each tube with the different lot numbers of the PBMC to be thawed. Capped tubes tightly and place in 37°C water bath prior to use. As needed, returned 50 mL conical tubes to the BSC, wiping with 70% EtOH prior to placing in the hood.

[001547] Removed a vial PBMC from cold storage and place in a floating tube rack in a 37°C water bath to thaw. Allowed thaw to proceed until a small amount of ice remains in the vial. Using a sterile transfer pipet, immediately transferred the contents of the vial into the 30 mL of CM2 in the 50 ml conical tube. Removed about 1 mL of medium from the tube to rinse the vial; returned rinse to the 50 mL conical tube. Capped tightly and swirl gently to wash cells.

[001548] Centrifuged at 400 x g for 5 min at room temperature. Aspirated the supernatant and resuspend the cell pellet in 1 mL of warm CM2/IL-2 using a 1000 µL pipet tip. Alternately, prior to adding medium, resuspended cell pellet by dragging capped tube along an empty tube rack. After resuspending the cell pellet, brought volume to 4 mL using CM2/IL-2 medium. Recorded volume.

[001549] Removed a small aliquot (*e.g.*, 100 µL) for cell counting using an automated cell counter. Performed counts in duplicate according to the particular automated cell counter

SOP. It most likely was necessary to perform a dilution of the PBMC prior to performing the cell counts. A recommended starting dilution was 1:10, but this varied depending on the type of cell counter used. Recorded the counts.

[001550] Adjusted concentration of PBMC to 1.3×10^7 cells/mL using CM2/IL-2 medium. Mixed well by gentle swirling or by gently aspirating up-and-down using a serological pipet.

Set Up Culture Flasks

[001551] Returned two labeled T25 flasks to the BSC from the tissue culture incubator. Returned the 10 µg/mL vial of anti-CD3/OKT3 to the BSC. Added 1 mL of the 1.3×10^7 PBMC cell suspension to each flask. Added 60 µL of the 10 µg/mL anti-CD3/OKT3 to each flask. Returned capped flasks to the tissue culture incubators for 14 days of growth without disturbance. Placed anti-CD3/OKT3 vial back into the refrigerator until needed for the next lot. Repeated for each lot of PBMC to be evaluated.

Day 14, Measurement of Non-proliferation of PBMC

[001552] Returned the duplicate T25 flasks to the BSC. For each flask, using a fresh 10 mL serological pipet, removed ~17 mL from each of the flasks, then carefully pulled up the remaining media to measure the volume remaining in the flasks. Recorded volume.

[001553] Mixed sample well by pipetting up and down using the same serological pipet.

[001554] Removed a 200 µL sample from each flask for counting. Counted cells using an automated cell counter. Repeated steps for each lot of PBMC being evaluated.

RESULTS AND ACCEPTANCE CRITERION

Results

[001555] The dose of gamma irradiation was expected to be sufficient to render the feeder cells replication incompetent. All lots were expected to meet the evaluation criterion, demonstrating a reduction in the total viable number of feeder cells remaining on Day 14 of the REP culture compared to Day 0.

Acceptance Criterion

[001556] The following acceptance criterion were met for each irradiated donor PBMC lot tested: "No growth" – meant that the total number of viable cells on Day 14 was less than the initial viable cell number put into culture on Day 0 of the REP.

Contingency Testing of PBMC lots which do not meet acceptance criterion.

[001557] In the event than an irradiated donor PBMC lot did not meet the acceptance criterion above, the following steps were taken to retest the lot to rule out simple experimenter error as the cause of its failure. If there were two or more remaining satellite vials of the lot, then the lot was retested. If there are one or no remaining satellite vials of the lot, then the lot was failed according to the acceptance criterion above.

[001558] To be qualified, a PBMC lot going through contingency testing had both the control lot and both replicates of the lot in question achieve the acceptance criterion. Upon meeting this criterion, the lot was then released for use.

EXAMPLE 5: PREPARATION OF IL-2 STOCK SOLUTION (CELLGENIX)

[001559] This Example describes the process of dissolving purified, lyophilized recombinant human interleukin-2 into stock samples suitable for use in further tissue culture protocols, including all of those described in the present application and Examples, including those that involve using rhIL-2.

Procedure

[001560] Prepared 0.2% Acetic Acid solution (HAc). Transferred 29 mL sterile water to a 50 mL conical tube. Added 1 mL 1 N acetic acid to the 50 mL conical tube. Mixed well by inverting tube 2-3 times. Sterilized the HAc solution by filtration using a Steriflip filter

[001561] Prepare 1% HSA in PBS. Added 4 mL of 25% HSA stock solution to 96 mL PBS in a 150 mL sterile filter unit. Filtered solution. Stored at 4°C. For each vial of rhIL-2 prepared, fill out forms.

[001562] Prepared rhIL-2 stock solution (6×10^6 IU/mL final concentration). Each lot of rhIL-2 was different and required information found in the manufacturer's Certificate of Analysis (COA), such as: 1) Mass of rhIL-2 per vial (mg), 2) Specific activity of rhIL-2 (IU/mg) and 3) Recommended 0.2% HAc reconstitution volume (mL).

[001563] Calculated the volume of 1% HSA required for rhIL-2 lot by using the equation below:

$$\left(\frac{Vial\ Mass\ (mg)\ x\ Biological\ Activity\ \left(\frac{IU}{mg}\right)}{6x10^6\ \frac{IU}{mL}}\right) - HAc\ vol\ (mL) = 1\%\ HSA\ vol\ (mL)$$

[001564] For example, according to CellGenix's rhIL-2 lot 10200121 COA, the specific activity for the 1 mg vial is 25×10^6 IU/mg. It recommends reconstituting the rhIL-2 in 2 mL 0.2% HAc.

$$\left(\frac{1mg\ x\ 25x10^6\ \frac{IU}{mg}}{6x10^6\ \frac{IU}{mL}}\right) - 2mL = 2.167mL\ HSA$$

[001565] Wiped rubber stopper of IL-2 vial with alcohol wipe. Using a 16G needle attached to a 3 mL syringe, injected recommended volume of 0.2% HAc into vial. Took care to not dislodge the stopper as the needle is withdrawn. Inverted vial 3 times and swirled until all powder is dissolved. Carefully removed the stopper and set aside on an alcohol wipe. Added the calculated volume of 1% HSA to the vial.

[001566] Storage of rhIL-2 solution. For short-term storage (<72 hrs), stored vial at 4°C. For long-term storage (>72 hrs), aliquoted vial into smaller volumes and stored in cryovials at -20°C until ready to use. Avoided freeze/thaw cycles. Expired 6 months after date of preparation. Rh-IL-2 labels included vendor and catalog number, lot number, expiration date, operator initials, concentration and volume of aliquot.

EXAMPLE 6: CRYOPRESERVATION PROCESS

[001567] This example describes the cryopreservation process method for TILs prepared with the abbreviated, closed procedure described in Example G using the CryoMed Controlled Rate Freezer, Model 7454 (Thermo Scientific).

[001568] The equipment used was as follows: aluminum cassette holder rack (compatible with CS750 freezer bags), cryostorage cassettes for 750 mL bags, low pressure (22 psi) liquid nitrogen tank, refrigerator, thermocouple sensor (ribbon type for bags), and CryoStore CS750 Freezing bags (OriGen Scientific).

[001569] The freezing process provides for a 0.5 °C rate from nucleation to -20 °C and 1 °C per minute cooling rate to -80 °C end temperature. The program parameters are as

follows: Step 1 - wait at 4 °C; Step 2: 1.0 °C/min (sample temperature) to -4 °C; Step 3: 20.0 °C/min (chamber temperature) to -45 °C; Step 4: 10.0 °C/min (chamber temperature) to -10.0 °C; Step 5: 0.5 °C/min (chamber temperature) to -20 °C; and Step 6: 1.0 °C/min (sample temperature) to -80 °C.

EXAMPLE 7: USE OF IL-2, IL-15, AND IL-21 CYTOKINE COCKTAIL

[001570] This example describes the use of IL-2, IL-15, and IL-21 cytokines, which serve as additional T cell growth factors, in combination with the TIL process of Examples 1 to 6.

[001571] Using the process of Examples 1 to 9, TILs were grown from colorectal, melanoma, cervical, triple negative breast, lung and renal tumors in presence of IL-2 in one arm of the experiment and, in place of IL-2, a combination of IL-2, IL-15, and IL-21 in another arm at the initiation of culture. At the completion of the pre-REP, cultures were assessed for expansion, phenotype, function (CD107a+ and IFNγ) and TCR Vβ repertoire. IL-15 and IL-21 are described elsewhere herein and in Gruijl, et al., IL-21 promotes the expansion of CD27+CD28+ tumor infiltrating lymphocytes with high cytotoxic potential and low collateral expansion of regulatory T cells, *Santegoets, S. J.,* J Transl Med., **2013,** *11:*37 (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3626797/).

[001572] The results showed that enhanced TIL expansion (>20%), in both $CD4^+$ and $CD8^+$ cells in the IL-2, IL-15, and IL-21 treated conditions were observed in multiple histologies relative to the IL-2 only conditions. There was a skewing towards a predominantly $CD8^+$ population with a skewed TCR Vβ repertoire in the TILs obtained from the IL-2, IL-15, and IL-21 treated cultures relative to the IL-2 only cultures. IFNγ and CD107a were elevated in the IL-2, IL-15, and IL-21 treated TILs, in comparison to TILs treated only IL-2.

EXAMPLE 8: TUMOR INFILTRATING LYMPHOCYTE (TIL) EXPANSION FROM FINE NEEDLE ASPIRATES (FNAS) AND CORE NEEDLE BIOPSIES

I. Background

[001573] This example provides data related to expanding tumor infiltrating lymphocytes (TILs) from tumor samples obtained by fine needle aspirations or core needle biopsies.

[001574] The goal of the study was to develop a method of expanding TILs from either core needle biopsies or fine needle aspirates. In some cases, 3 to 5 tumor samples were obtained from patients with cancer such as lung cancer (n=3), melanoma (n=1), head and neck cancer (n=1), cervical cancer (n=1), ovarian cancer (n=3), pancreatic cancer (n=2), glioblastoma (GBM), and colorectal cancer (n=1).

II. **Expanded TILs from core biopsies from a patient with pancreatic cancer (P7015).**

[001575] Pancreatic core biopsies were received for the study. One core was placed into one well of a 24-well plate for a total of 5 wells. When two cores were place into two wells, little to no growth was observed in these wells. Three wells were treated with culture medium supplemented with IL-2 and two wells were treated with culture medium supplemented with a cytokine cocktail containing IL-2, IL-15, and IL-21. Viable cells were observed in the wells treated with IL-2 alone (Figure 8A) and in the wells treated with the cytokine cocktail (Figure 8B). About 1.3×10^6 viable cells/well (1.2×10^6 viable cells/well on average) were obtained with the IL-2 treatment. About 2.3×10^6 viable cells/well (2.0×10^6 viable cells/well on average) were obtained with the cytokine cocktail treatment.

[001576] Figures 17A, 17B, and 17C show phenotyping and functional status of TILs derived from a core biopsy of pancreatic tumors. The TILs were expanded from pancreatic tumors and culture in medium containing IL-2 or medium supplemented with cell culture additives. TILs were expanded from pancreatic tumors in IL-2 supplemented culture medium and in culture medium containing cell culture additives (Figure 17D). Figure 17A shows the percentage of cells that were CD3+, CD4+, CD8+, CD19+, CD3+/CD56+, or CD3-/CD56+. Figure 17B shows that the percentage of CD107a cells in the CD4+ subpopulation after stimulation with phorbol 12-myristate 13-acetate (PMA) compared to an unstimulated condition. Figure 17C shows functional analysis of the expanded TILs from the pancreatic core biopsy. The graph shows the percentage of CD107a cells in the CD8+ subpopulation after stimulation with phorbol 12-myristate 13-acetate (PMA) compared to an unstimulated condition.

[001577] The majority of cells expanded from the pancreatic core biopsies are T cells (CD4+ cells and CD8+ cells). Stimulation with PMA demonstrated the ability of the CD8+ to degranulate as indicated by expression of CD107a+, thereby suggesting that these cells are

functional. The % of CD8+ expressing CD107a+ was higher in the wells treated with the culture additive wells compared to IL-2 alone (Figure 17C).

III. **Expanded TILs from fine needle aspirates from a patient with cervical carcinoma (C2005).**

[001578] A cervical tumor was received for the study. Five fine needle aspirates were isolated from the tumor using a 22 gauge needle and syringe. Each FNA sample was placed into at least three wells of a 24-well plate. Two wells of each sample were treated with IL-2 and one well of each sample was treated with the cytokine cocktail (IL-2, IL-15, and IL-21).

[001579] Figure 10A shows that CD4+ and CD8+ T cells were present in the FNAs treated with IL-2. On average 3.45×10^5 viable cells/well were present in the IL-2 treated wells. Figure 10B shows that CD4+ and CD8+ T cells were present in the FNAs treated with the cytokine cocktail. On average 8.69×10^5 viable cells/well were present in these wells. The IL-2. IL-15, and IL-21 treated aspirates demonstrated more cells/well compared to the IL-2 treated aspirates. The percentage of CD8+ cells was greater in the cocktail treated wells (20.1%) compared to the IL-2 treated wells (8.58%).

IV. **Expanded TILs from fine needle aspirates from a patient with lung carcinoma (L4032).**

[001580] Three lung tumors were received for the study. Fine needle aspirates were isolated from the tumors using a 22 gauge needle and syringe. One FNA sample included 1.6×10^6 cells with 28% viable cells. Two wells (each condition) were treated with culture medium supplemented with (1) IL-2 alone or (2) the cytokine cocktail. In addition, lung tumor fragments were cultured in a 24-well plate (two wells each) as a control. Cells were harvested at day 13 and evaluated by FACS analysis.

[001581] Figure 11A shows that T cells expanded from an FNA sample treated with IL-2. There were on average about 4.64×10^5 viable cells/well. Figure 11B shows that T cells expanded from an FNA sample treated with the cocktail containing IL-2, IL-15, and IL-21. There were on average about 7.75×10^5 viable cells/well. Figure 11C shows that T cells expanded from a lung tumor fragment treated with IL-2. There were on average about 4.28×10^5 viable cells/well.

[001582] Next, the TILs cultured from the FNA samples were expanded using a rapid expansion protocol (*e.g.*, second expansion protocol or step D). The cells were harvested at day 14 and evaluated by FACS analysis. Figure 12A shows the total cell count of the

expanded TILs from the FNA samples treated with (i) IL-2, (ii) IL-2, IL-15, and IL-21, and (iii) IL-2, IL-15, and IL-21 into IL-2. Figure 12A shows that after the second expansion, the TILs included CD4+ and CD8+ T cells (84.1% and 10.3%, respectively).

V. Expanded TILs from fine needle aspirates from a patient with lung carcinoma (L4033).

[001583] A fine needle aspirate was isolated from a lung tumor using a 22 gauge needle and syringe. The FNA sample included 5.5×10^6 cells with 76% viable cells. One G-Rex 10 flask (each condition) was treated with culture media containing (1) IL-2 alone or (2) the cytokine cocktail. In addition, four lung tumor fragments were cultured in a G-Rex 10 flask as a control. Cells were harvested at day 12 and evaluated by FACS analysis. Figure 13A shows that CD4+ T cells (4.25%) and CD8+ T cells (55.1%) were obtained from the FNA sample treated with IL-2. Figure 13B shows that CD4+ T cells (7.76%) and CD8+ T cells (54.5%) were obtained from the tumor fragments treated with IL-2.

[001584] Next, the TILs cultured from the FNA samples were expanded using a rapid expansion protocol (*e.g.*, second expansion protocol or step D). The aspirates were incubated for the first 6 days in a conical tube. The cells were harvested at day 14 and evaluated by FACS analysis. Figure 14 shows the total cell count of the expanded TILs from the FNA samples and the tumor fragments treated with IL-2. The graph shows that TILs from the FNA samples expanded similarly to the TILs from the tumor fragments.

[001585] In a similar experiment, FNA samples were isolated from three lung tumor samples from a patient with lung carcinoma (L4034). About 1.0×10^6 cells from an FNA sample were cultured in a G-Rex 10 flask for 15 days in culture medium supplemented with IL-2 alone. As a control, four tumor fragments were cultured in a G-Rex 10 flask culture medium supplemented with IL-2 alone. Few to no detectable cells were harvested from the FNA culture.

VI. Expanded TILs from fine needle aspirates from patients with ovarian carcinoma (OV8011, OV8012, and OV8013).

[001586] Three ovarian tumors (OV8011, OV8012, and OV8013) were received for the study. Fine needle aspirates were isolated from the tumors using a 22 gauge needle and syringe.

[001587] The FNA sample from OV8011 included 3.19×10^5 cells with 28% viable cells. This FNA sample was divided into four wells; two wells were cultured in culture

medium containing IL-2 and two wells were cultured in culture medium containing IL-2, IL-15, and IL-21. The FNA sample from OV8012 included 8.3×10^5 cells with 11% viable cells. This FNA sample was divided into two wells; one well was cultured in culture medium containing IL-2 and the other well was cultured in culture medium containing IL-2, IL-15, and IL-21. The FNA sample from OV8013 included 1.5×10^5 cells with 7% viable cells. This FNA sample was divided into two wells; one well was cultured in culture medium containing IL-2 and the other well was cultured in culture medium containing IL-2, IL-15, and IL-21. No expansion was observed from the FNA samples.

[001588] As a control, four tumor fragments from each tumor were cultured in a G-Rex 10 flask. Some flasks received culture medium containing IL-2 alone and other flasks received culture medium containing IL-2, Il-15, and IL-21. Figure 15 shows the total cell counts of the cells derived from the ovarian tumor fragments cultured in culture medium containing (1) IL-2 alone or (2) the cytokine cocktail. The results show that the ovarian tumors expanded poorly in the first expansion.

VII. Expanded TILs from fine needle aspirates from a patient with melanoma (M1101).

[001589] A melanoma tumor (M1101) was received for the study. An FNA sample was obtained from the tumor. The sample included 1.96×10^6 cells with 65% viable cells. The sample was place in a G-Rex 10 flask and treated with culture medium supplemented with IL-2. As a control, four melanoma tumor fragments were placed in a G-Rex 10 flask and treated with culture medium supplemented with IL-2. The cells were harvested at Day 12. Figure 16A shows that CD4+ T cells (21.4%) and CD8+ T cells (52.8%) expanded from the FNA sample treated with IL-2. Figure 16B shows that CD4+ T cells (18.0%) and CD8+ T cells (71.5%) expanded from the tumor fragments treated with IL-2.

[001590] Figures 18A and 18B provides a summary of the results obtained from the core biopsies and fine needle aspirates.

VIII. Expanded TILs from fine needle aspirates from a patient with melanoma (M1101).

[001591] On Day 11 of the pre-REP, cell counts were performed on the flasks containing the aspirate and fragments (n=2). The pre-REP cultures containing the fragments were harvested at Day 11 (Gen2). The pre-REP culture containing the aspirate was continued due to low cell count at Day 11. The aspirate was recounted at Day 18 and then again at Day

21 (Figure 19A). At Day 21, the culture had reached almost 60×10^6 cells, the cells were harvested at this time and assessed via flow cytometry. The phenotype (Figure 19B) and function (CD107a expression, Figure 19C) of the pre-REP TIL from the fragments and aspirates were shown to be comparable.

[001592] Figures 19A, 19B and 19C show the expansion of TILs from an FNA sample from another patient with melanoma (M1107). Figure 19A shows the total cell count at different time during the expansion. Figure 19B shows the phenotype of the expanded TILs. Figure 19C shows functional analysis of the expanded TILs from fragments and aspirates. The graph shows the percentage of CD107a cells in the CD4+ and CD8+ subpopulations after stimulation with PMA compared to an unstimulated condition.

EXAMPLE 9: TUMOR INFILTRATING LYMPHOCYTE (TIL) EXPANSION FROM FINE NEEDLE ASPIRATES (FNAS) AND CORE NEEDLE BIOPSIES

I. **Pancreatic cores: Data shows in Figures 20-29.**

[001593] Fifteen tumors (Pancreatic, Lung, Cervical, Mesothelioma, Colorectal) was reported. To date, 37 tumor samples from different histology including pancreatic, cervical, lung, oral and esophageal, sarcoma cancers have been received.

[001594] Upon arrival the tumors were fragmented and placed in flasks/plate with IL-2 for a pre-Rapid Expansion Protocol (REP). pre-REP TIL were further propagated in a REP protocol in the presence of irradiated PBMCs, anti-CD3 antibody (30 ng/mL), and IL-2 (3000 IU/mL). Lung and Sarcoma pre-REP TIL were expanded from 4 tumor fragments in 24-well-plate format for up 22 days. Other cancers were expanded from 4 tumor fragments in G-REX TIL for up to 22 days.

II. **TIL can be expanded from Core Biopsies from Pancreatic Cancer: data shown in Figures 30-37.**

General conclusions

[001595] The kinetics of growth demonstrated very few cells at Day 11-12 in both P7028 and P7031, however significant expansion was demonstrated between Day 21-28.

[001596] The majority of cells expanded from the pancreatic core biopsies are T cells and phenotypically CD4+. TIL isolated from pancreatic tumors are typically CD4+.

[001597] Stimulation with PMA demonstrated the ability of the CD4+ to degranulate as indicated by expression of CD107a+, thereby suggesting that these cells are functional, as was shown with the earlier studies with P7015. The expression of CD107a was similar in TIL derived from fragments.

[001598] The TIL derived from cores secreted significant levels of IFNγ in a restimulation assay with plate-bound anti-CD3, similar to TIL derived from fragments.

[001599] Extensive phenotyping demonstrated similar expression of the majority of markers assessed in TIL derived from tumor fragments, suggesting that the outcome of growing TIL from cores is similar to fragments.

[001600] The exhaustion/activation panel demonstrated significant differences in PD-1, Tim3 and KLRG1 in the CD4+ cells and LAG3, Tim3 and KLRG1 in the CD8+ cells. Additional pancreatic core studies are required to validate these findings.

General Proposals:

[001601] These results suggest that the process for growing TIL successfully from pancreatic cores (and other histologies) will likely require additional time in culture (beyond the pre-REP requirements for Gen2), unless additional "tweaking strategies" are utilized to enhance growth (see below).

[001602] Based upon the preliminary growth data in pancreatic cores, it is recommended that the pre-REP process be between Day 21-28, in order to obtain sufficient cells .

[001603] For this process the pre-REP should be initiated in G-Rex 10 or G-Rex 10M's will CM1 media and IL-2/IL-15/IL-21.

[001604] The REP should be initiated either in a G-Rex 100M or G-Rex 500M, depending on the pre-REP cell count.

[001605] G-Rex 500M: if the cell count is 5×10^7.

[001606] G-Rex 100M: if cell count is between $1\text{-}5 \times 10^7$.

[001607] Future experiments will be used to determine the minimum number of cells required for sufficient cell growth in each vessel .

[001608] Future experiments will compare the growth, phenotype and function of pancreatic cores expanded in IL-2/IL-15/IL-21 alone or with IL-2/IL-15/IL-21 with OKT3 and feeders.

EXAMPLE 10: EXPANDING TIL USING CORE BIOPSIES

[001609] Rationale:

[001610] Tumor removal/resection is not appropriate or feasible for all patients. In this cohort of patients, core biopsies may be a source of TIL for ACT (adoptive cellular therapy). Such core biopsies allow to sample multiple lesions and potentially obtain a broader TIL repertoire.

[001611] **Strategy:**

[001612] The present example provides a method to expand TIL from core biopsies, from multiple histologies and develop a defined process that is feasible for TIL production from core biopsies. See, for example Figure 38 which provides a diagram of two exemplary processes.

[001613] Data

[001614] The ability to expand TIL form biopsies has been demonstrated for pancreatic cores using Gen2, which also showed that TIL obtained from biopsies are phenotypically and functionally comparable to TIL obtained from fragments.

[001615] Seven sets of pancreatic cores (between 7-10 cores) were treated with IL-2/IL-15/IL-21 and pre-REP'ed and assessed for expansion. Three successfully expanded, however the growth was extremely variable. Pre-REP cultures required at least 21-28 days to acquire sufficient numbers of cells (*e.g.*, 21-28 days for Steps A through E of Figure 7). Extensive phenotyping demonstrated similar expression of the majority of markers assessed in TIL derived from pancreatic tumor fragment. The expression of CD107a and IFNγ was similar in TIL derived from fragments

Current protocol:

[001616] Addition of OKT3 & feeders at Day 3 of culture enhanced TIL yield. Cultured fewer cores/flask and assess growth (for example, cultured 1-2 cores).

[001617] 1-2 core biopsies (3 pancreas, 4 melanoma, 1 breast, and 1 ovarian) were subjected to a modified research-scale process that included the addition of OKT3 + feeders at Day 3 (See, Figures 39-40).

- TIL were subjected to a second REP (*e.g.*, a second expansion).
- Pancreatic cores were expanded +/- IL15 & 21 (the triple cocktail; REP1 & REP2; *e.g.*, a first expansion and a second expansion)
- Ovarian cores were treated +/- OX40 (BMS) (n=1, preliminary data).
- All TIL were subjected to a second REP.

PANCREATIC

[001618] TIL from pancreatic cores were successfully expanded in the presence of OKT3 + feeders. >50e6 cells were acquired between Day 11-17 of the first expansion.

[001619] Two cores were generally required to acquire the appropriate cell numbers.

[001620] The addition of the triple cocktail (TC: IL-2, IL-15/IL-21) to the 2nd REP (either previously cultured in IL-2 or the TC) did not significantly alter the expansion of TIL in comparison to IL-2 alone.

[001621] The addition of OKT3 + feeders at Day 3 to core biopsies does not significantly alter the phenotype (memory, CD27/CD28, exhaustion, activation) of TIL in comparison to cores grown using Gen2 like conditions. Phenotype (memory, CD27/CD28, exhaustion, activation) of TIL expanded through 2 REPs did not significantly differ from that of TIL grown using Gen2 conditions.

[001622] CD107a expression did not vary significantly with the addition of the OKT3 and feeders. TIL expanded through 2 REPs maintained the ability to respond to non-specific stimulus, as assessed by CD107a induction.

[001623] TIL expanded through 2 REPs maintained the ability to respond to non-specific stimulus, as assessed by CD107a induction.

[001624] IFNγ secretion was evaluated. TIL produce IFNγ upon stimulation with OKT3.

[001625] Pancreatic TILs can be expanded from Core Biopsies from Pancreatic Cancer.

P7028 and P7031 pre-REP Experimental Summary and Results

[001626] Core biopsies were place into a G-Rex 10 with the triple cocktail (IL-2/IL-15/IL-21)

[001627] P7028 TIL were assessed at Day 12 and kept in culture until Day 19

[001628] Cell counts: Day 12= 1.61×10^6 viable cells; Day 21 3.49×10^7 viable cells

[001629] P7030 TIL were assessed at Day 12 and kept in culture until Day 27

[001630] Cell counts: Day 12= 1.52×10^5 viable cells; Day 27: 1.14×10^7 viable cells

MELANOMA

[001631] TIL derived from melanoma cores were successfully expanded in the presence of OKT3 + feeders (n=4). $>50 \times 10^6$ cells were acquired by Day 12 of the first expansion.

[001632] One to two cores were generally required to acquire the appropriate cell numbers. The addition of OKT3 + feeders at Day 3 to core biopsies, does not significantly alter the phenotype (memory, CD27/CD28, exhaustion, activation) of TIL in comparison to fragments using Gen2 like conditions.

[001633] TIL expanded through 2 REPs responded to non-specific stimuli, as assessed by CD107a mobilization and IFNγ secretion.

LUNG

[001634] TILs from 1 core from lung were successfully expanded in the presence of IL-2+ OKT3 + feeders at Day 0. $>50 \times 10^6$ cells were acquired at Day 11.

[001635] Phenotype (memory, CD27/CD28, exhaustion, activation) of TIL expanded through 2-REPs varied slightly from TIL expanded from cores treated with OKT3 + feeders at Day 3 of culture initiation. There was an increase in the ratio of CD8+/CD4+ in the Day 3 treated cores.

[001636] There were slight differences in CD28, CD69, CD39, CD49a, Tim3 and CD101 expression were observed between the Day 0 and Day 3 culture conditions.

[001637] The TILs produced IFNγ upon stimulation with anti-CD3. The Day 0 OKT3 + feeders treated lung cores had greater levels of IFNγ in comparison to the Day 3 OKT3 + feeders treated lung cores.

[001638] The TILs expanded through 2 REPs maintained the ability to respond to non-specific stimulus, as assessed by CD107a induction.

[001639] The CD8+ TILs from Day 3 OKT3 + feeders treated lung cores were oligoclonal compared to the Day 0 OKT3 + feeders treated lung core TILs, indicating a different TCR repertoire depending on the timing of the addition of OKT3 + feeders.

Conclusions

[001640] The addition of OKT3 + feeders to cores at Day 3 of culture significantly improved TIL yield. $>50 \times 10^6$ cells (threshold for REP2 initiation) was acquired within 11-15 days of the first expansion in all histologies assessed (pancreatic, melanoma, breast and ovarian). Two cores (in the histologies assayed) were sufficient to acquire the number of cells required.

[001641] TILs from at least 2 cores from breast were successfully expanded in the presence of IL-2+ OKT3 + feeders, but not IL-2 alone. $>50 \times 10^6$ cells were acquired at Day 11. One to two cores are sufficient to acquire the number of cells required.

[001642] Data suggests that the phenotype and function of TIL expanded from cores (with early OKT3+ feeders) is similar to cores/fragments expanded under Gen2 conditions (IL-2 alone). Addition of OX40 in ovarian cores improved the expansion of TILs in comparison to IL-2 alone.

[001643] 1-2 pancreatic cores failed to expand in IL-2 alone.

[001644] Side to side experiments will be performed with fragments +/- OKT3+ feeders at Day 3. TILs produced IFNγ upon stimulation with OKT3 and further experiments are ongoing. Differences in KLRG1, CD25 and PD1 were observed in ovarian cores treated +/- OX40 (therapeutic TIL population).

[001645] Phenotype, function (potency) and TCR repertoire were and continue to be assessed (currently in progress). The addition of OKT3 + feeders at Day 3 to core biopsies, did not significantly alter the phenotype (memory, CD27/CD28, exhaustion, activation) of TIL in comparison to fragments using Gen2 like conditions. TIL expanded through 2 REPs responded to non-specific stimuli, as assessed by CD107a mobilization and IFNγ secretion. Phenotype (memory, CD27/CD28, exhaustion, activation) of TILs expanded through 2-REPs did not significantly differ from that of TILs expanded using a 1-REP process (ie. no OKT3+feeders during 1st expansion) in breast cores.

[001646] Slight differences in KLRG1 and PD1 expression were observed.

[001647] TILs expanded through 2 REPs maintained the ability to respond to non-specific stimulus, as assessed by CD107a induction.

[001648] Further assessment will include +/- removal of cores at Day 3 as well as further assessment of the effect of anti-OX40 on expanded TILs will be performed.

EXAMPLE 11: EXPANDING TIL FROM BREAST AND OVARIAN CORE BIOPSIES USING OKT3+ FEEDERS: SUMMARY OF EXAMPLES 9-10

[001649] TIL from at least 2 cores from breast were successfully expanded in the presence of IL-2+ OKT3 + feeders, but not IL-2 alone. $>50 \times 10^6$ cells were acquired at Day 11.

[001650] Addition of OX40 in ovarian cores improved the expansion of TIL in comparison to IL-2 alone.

[001651] Phenotype (memory, CD27/CD28, exhaustion, activation) of TIL expanded through 2-REPs did not significantly differ from that of TIL expanded using a 1-REP process (*i.e.* no OKT3+feeders during 1st expansion) in breast cores.

[001652] Slight differences in KLRG1 and PD1 expression were observed.

[001653] Differences in KLRG1, CD25 and PD1 were observed in ovarian cores treated +/-OX40 (final product).

[001654] TIL produce IFNγ upon stimulation with OKT3.

[001655] TIL expanded through 2 REPs maintained the ability to respond to non-specific stimulus, as assessed by CD107a induction.

[001656] The addition of OKT3 + feeders to cores at Day 3 significantly improved TIL yield. $>50 \times 10^6$ cells (threshold for REP initiation under Gen2 conditions) was acquired within 11-15 days in all histologies assessed (pancreatic, melanoma, breast, and ovarian). Two cores (in the histologies assayed) are sufficient to acquire the number of cells required. Preliminary data suggest that the phenotype and function of TIL expanded from cores (with early OKT3+ feeders) is similar to cores/fragments expanded under control conditions.

EXAMPLE 12: EXPANDING TIL FROM NEEDLE ASPIRATES

Strategy:

[001657] To expand TIL from needle aspirates, from multiple histologies and develop a defined process that is feasible for TIL production. Preliminary studies demonstrated that TIL could be expanded from "some" aspirates performed under optimal conditions (*i.e.*, taken directly from a tumor biopsies).

[001658] 22 aspirates were isolated from tumor biopsies of various histologies at Iovance using Gen2 conditions (*i.e.*, no OKT3 and feeders at Day 3). Growth was observed in 6 tumors (1 cervical, 2 lung and 3 melanoma). The number of cells isolated during the pre-REP was well below the 50×10^6 cell threshold.

[001659] Additionally, 10 needle aspirates have been received, including three from melanoma (n-3), and seven from breast (n-7). The aspirates were treated with OKT3 and feeders.

[001660] Due to small initial cell numbers, most of the experiments were initiated in a 96 well plate. $<50 \times 10^6$ cells were obtained at Day 11 in all aspirates. Several sub-cultures would be required to acquire the necessary number of cells.

EXAMPLE 13: EXPANDING TIL FROM TUMOR CORE BIOPSIES

INTRODUCTION

[001661] Adoptive T cell therapy (ACT) with autologous tumor infiltrating lymphocytes (TIL) has demonstrated clinical efficacy in patients with metastatic melanoma and cervical carcinoma. TIL are currently expanded from fragmented tumor that is removed surgically. Unfortunately, patients with unresectable tumors do not currently qualify for TIL therapy. If TIL expansion were feasible from the type of small tumor samples used by physicians for diagnostics and staging purposes, correlative studies, and biomarker analyses, TIL therapy would become available to a greater number of cancer patients.

[001662] A core needle biopsy is an example of such small tumor samples. It is an inexpensive, safe, simple and less invasive procedure than surgical resection. A core biopsy

removes a small cylinder of tumor tissue using a small incision and needle. Typically, a needle can be inserted into the incision up to 5 times, to excise an adequate amount of tissue. Core biopsies are small samplings of the intact tumor microenvironment, and therefore contain tumor cells, stroma and immune cells, including TIL. To date, there is one clinical study, in melanoma, that used tissue derived from core needle biopsies to expand TIL (Ullenhag, G.J., *et al.*, *Cancer Immunol Immunother*, 61(5):725-732 (2014)). The absolute cell numbers and fold expansion were similar between TIL derived from surgical resection versus TIL derived from core biopsies. Interestingly, most of the objective responses (4/5) in this study were treated with cells derived from core biopsies. No further studies have corroborated these findings.

[001663] In some embodimetns, a 2-REP process is employed to expand TIL derived from core biopsies, as described above and in the present example.

PURPOSE

[001664] The present example provides a protocol to expand TIL to clinically appropriate numbers from small tumor samples for ACT (adoptive cell transfer). This protocol was developed using tumor core biopsies from pancreatic, melanoma, breast, and ovarian tumors. The expanded TIL were assessed for growth, viability, phenotype, function (IFNγ secretion, CD107a mobilization), and TCR Vβ repertoire (RNA-sequencing).

Materials

[001665] Tumor Core Tissue. Tumor cores from pancreas, melanoma, breast, and ovarian are received from UPMC, Biotheme and MT group.

PROCEDURE

Core Processing and Culture Initiation

[001666] Freshly resected tumor core biopsies are received from research alliances (UPMC) and tissue procurement vendors (Biotheme and MTG group). The tumor core biopsies are shipped overnight in HypoThermosol (Biolife Solutions, Washington, Cat # 101104) (with antibiotic) or RPMI 1640 (Fisher Scientific, Pennsylvania, Cat # 11875-085) + male human AB serum (Access Biologicals, California, A13012).

[001667] Using a 0.22-μM filter on top of 50-mL conical tube, carefully and slowly pour the contents of the shipping container comprising the tumor core biopsies through the

filter. The cores are retained in the filter. Wash the cores by gently adding 10-20 mL of HBSS over the cores.

[001668] Gently remove each core from the filter membrane, while attempting to keep the core intact. Using either sterile plastic forceps or gently pipetting (with approximately 200 µL of HBSS in a 1-mL pipette), transfer the core to a G-Rex 10 with 40 mL of CM2 and 6000 IU/mL of IL-2.

REP1 initiation (*e.g.*, priming first expansion)

[001669] On Day 3 post-culture, OKT3 (30 ng/mL) (Miltenyi Biotec, Germany, Cat # 170-076-116) and Feeders (1:100) are added to the G-Rex 10 without disturbing the culture upon addition.

[001670] Assess the culture daily for growth, by microscopy. REP1 is conducted for 8 days after the addition of OKT3 + Feeders and ends at Day 11. The cells remain untouched for the entire length of the culture.

[001671] To count, remove approximately 30 mL of media and resuspend the cells in the remaining 10 mL of media, by pipetting up and down. Place cells in a 50-mL conical tube and centrifuge at 1500 rpm at RT for 5 min in a Sorvell Legend XFR centrifuge and swinging bucket (ThermoScientific, CA, Cat #75003180).

[001672] Determine cell and viability counts, using the Nexcelom Cellometer K2 (Nexcelom, MA).

[001673] If after centrifugation there is no pellet, or the pellet is < 200 µL, resuspend the pellet in 2 mL of media for counting. If the pellet is > 200 µL resuspend in 5 mL of CM2 media for counting.

[001674] Experiments are harvested at Day 11 post-culture initiation (8 days post-OKT3 + Feeders addition). REP1 product is assessed for cell count, viability, phenotype (TIL1, TIL2 and TIL3 (TIL2 and TIL3 panel for Surface Antigen Staining of TIL), TIL3), and function (CD107a (Assessing TIL function by CD107a mobilization), IFNγ (WRK LAB-059)). The remaining cells are frozen and stored in liquid nitrogen.

REP2 Initiation (*e.g.*, second rapid expansion)

[001675] For mini-REP2 initiation, 1×10^5 cells are placed into a G-Rex 10 with 40 mL of CM2 media and 3000 IU/mL of IL-2. Anti-CD3 (clone: OKT3, 30 ng/mL) and Feeders

(1:100 ratio, TIL: Feeders) are added at culture initation. All flasks are treated with OKT3 + Feeders in REP2, even if OKT3 + feeders were not added during REP1.

[001676] A media change is performed at Day 5-7 of REP2 (Day 16-18 of process). The media in the G-Rex 10 flask is volume reduced to approximately 10 mL and supplemented to 40 mL with either CM2 or AimV + 3000 IU/mL IL-2.

[001677] At Day 11 of REP2, volume is reduced as indicated above and cells centrifuged at 1500 rpm at RT for 5 min.

[001678] The final product is assessed for cell count, viability, phenotype (TIL1, TIL2 (TIL2 panel for Surface Antigen Staining of TIL, v1 and v2), TIL3 and function (CD107a (Assessing TIL function by CD107a mobilization, IFNγ). Additional cells ($1 \times 10^6 - 5 \times 10^6$ cells) are pelleted and frozen for RNA sequencing and analysis. The remaining cells are frozen and stored in liquid nitrogen.

RESULTS AND ACCEPTANCE CRITERIA

[001679] The REP1 yield is expected to be > 50 x 10^6 by D11 in flasks that were treated with OKT3 + Feeders. The fold expansion during REP2 is expected to be unaffected by the addition of OKT3 and feeders during REP1 and to be consistent with Gen2 REP fold expansion numbers. The phenotype of TIL derived from tumor core biopsies and exposed early to OKT3 + Feeders is expected to be similar to that of TIL derived from surgical resections (or small biopsies) and expanded according to the Gen 2 process.

EXAMPLE 14: GEN 3 EXEMPLARY PROCESS

[001680] The example provides a comparison between the Gen 2 and Gen 3 processes. This example describes the development of a robust TIL expansion platform. The modifications to the Gen 2 process reduce risk and streamline the manufacturing process by reducing the number of operator interventions, reduce the overall time of manufacturing, optimize the use of reagents, and facilitate a flexible semi-closed and semi-automated cell production process amenable to high-throughput manufacturing on a commercial scale.

[001681] Process Gen 2 and Gen 3 TILs are composed of autologous TIL derived from an individual patient through surgical resection of a tumor and then expanded *ex vivo*. The Priming First Expansion step of the Gen 3 process was a cell culture in the presence of

interleukin-2 (IL-2) and the monoclonal antibody OKT3, which targets the T-cell co-receptor CD3 on a scaffold of irradiated peripheral blood mononuclear cells (PBMCs).

[001682] The manufacture of Gen 2 TIL products consisted of two phases: 1) pre-Rapid Expansion (Pre-REP) and 2) Rapid Expansion Protocol (REP). During the Pre-REP resected tumors were cut up into ≤ 50 fragments 2-3 mm in each dimension which were cultured with serum-containing culture medium (RPMI 1640 media containing 10% HuSAB supplemented) and 6,000 IU/mL of Interleukin-2 (IL-2) for a period of 11 days. On day 11 TIL were harvested and introduced into the large-scale secondary REP expansion. The REP consists of activation of $\leq 200 \times 10^6$ of the viable cells from pre-REP in a co-culture of 5×10^9 irradiated allogeneic PBMCs feeder cells loaded with 150 μg of monoclonal anti-CD3 antibody (OKT3) in a 5L volume of CM2 supplemented with 3000 IU/mL of rhIL-2 for 5 days. On day 16 the culture is volume reduced 90% and the cell fraction is split into multiple G-REX-500 flasks at $\geq 1 \times 10^9$ viable lymphocytes/flask and QS to 5L with CM4. TIL are incubated an additional 6 days. The REP is harvested on day 22, washed, formulated, and cryo-preserved prior to shipping at -150°C to the clinical site for infusion.

[001683] The manufacture of Gen 3 TIL products consisted of three phases: 1) Priming First Expansion Protocol 2) Rapid Second Expansion Protocol (also referred to as rapid expansion phase or REP) and 3) Subculture Split. To effect the Priming First Expansion TIL propagation, resected tumor was cut up into ≤ 120 fragments 2-3 mm in each dimension. On day 0 of the Priming First Expansion, a feeder layer of approximately 2.5×10^8 allogeneic irradiated PBMCs feeder cells loaded with OKT-3 was established on a surface area of approximately 100 cm^2 in each of 3 100 MCS vessels. The tumor fragments were distributed among and cultured in the 3 100 MCS vessels each with 500 mL serum-containing CM1 culture medium and 6,000 IU/mL of Interleukin-2 (IL-2) and 15 ug OKT-3 for a period of 7 days. On day 7, REP was initiated by incorporating an additional feeder cell layer of approximately 5×10^8 allogeneic irradiated PBMCs feeder cells loaded with OKT-3 into the tumor fragmented culture phase in each of the 3 100 MCS vessels and culturing with 500 mL CM2 culture medium and 6,000 IU/mL IL-2 and 30 ug OKT-3. The REP initiation was enhanced by activating the entire Priming First Expansion culture in the same vessel using closed system fluid transfer of OKT3 loaded feeder cells into the 100MCS vessel. For Gen 3, the TIL scale up or split involved process steps where the whole cell culture was scaled to a larger vessel through closed system fluid transfer and was transferred (from 100 M flask to a 500 M flask) and additional 4L of CM4 media was added. The REP cells were

harvested on day 16, washed, formulated, and cryo-preserved prior to shipping at -150ºC to the clinical site for infusion.

[001684] Overall, the Gen 3 process is a shorter, more scalable, and easily modifiable expansion platform that will accommodate to fit robust manufacturing and process comparability.

Table 36: Comparison of Exemplary Gen 2 and Exemplary Gen 3 manufacturing process.

Step	Process (Gen 2)	Process (Gen 3)
Pre REP-day 0	Up to 50 fragments/ 1-G-Rex 100MCS - 11 days In 1 L of CM1 media + IL-2 (6000 IU/mL)	Whole tumor up to 120 fragments divided evenly among up to 3 flasks. 1 flask: 1-60 fragments 2 flasks: 61-89 fragments 3 flasks 90-120 fragments 7 days in 500 mL of CM1 media + IL-2 (6000 IU/mL) 2.5×10^8 feeder cells/flask 15ug OKT-3/flask
REP Initiation	Direct to REP- Day 11- $<200 \times 10^6$ TIL (1)G-Rex 500MCS in 5L CM2 media IL-2 (3000 IU/mL) 5×10^9 feeder cells 150ug OKT-3	Direct to REP- Day 7-all cells TIL- same G-Rex 100MCS Add 500 CM2 media IL-2 (6000 IU/mL) 5×10^8 feeder cells/flask 30ug OKT-3/flask
TIL propagation or Scale up	Volume reduce and split cell fraction in up to 5 G-REX 500MCS 4.5L CM4 media + IL-2 (3000 IU/mL) $\geq 1 \times 10^9$ TVC / flask Split day 16	Each G-REX 100MCS(1L) transfers to 1 G-REX 500MCS Add 4L CM4 media +IL-2 (3000 IU/mL) Scale up on day 9 to 11
Harvest	Harvest day 22, LOVO-automated cell washer	Harvest day 16 LOVO- automated cell washer
Final formulation	Cryopreserved Product 300IU/mL IL2- CS10 in LN_2, multiple aliquots	Cryopreserved product 300IU/mL IL-2-CS10 in LN_2, multiple aliquots
Process time	**22 days**	**16 days**

[001685] On day 0, for both processes, the tumor was washed 3 times and the fragments were randomized and divided into two pools; one pool per process. For the Gen 2 Process, the fragments were transferred to one GREX 100MCS flask with 1 L of CM1 media containing 6,000IU/mL rhIL-2. For the Gen 3 Process, fragments were transferred to one GRE × 100MCS flask with 500 mL of CM1 containing 6,000IU/mL rhIL-2, 15 μg OKT-3 and 2.5×10^8 feeder cells. Seeding of TIL for Rep initiation day occurred on different days according to each process. For the Gen 2 Process, in which the G-REX 100MCS flask was 90% volume reduced, collected cell suspension was transferred to a new G-REX 500MCS to start REP initiation on day 11 in CM2 media containing IL-2 (3000 IU/mL), plus 5×10^9 feeder cells and OKT-3 (30 ng/mL). Cells were expanded and split on day 16 into multiple GREX 500 MCS flasks with CM4 media with IL-2 (3000 IU/mL) per protocol. The culture was then harvested and cryopreserved on day 22 per protocol. For the Gen 3 process, the REP initiation occurred on day 7, in which the same G-REX 100MCS used for REP initiation. Briefly, 500 mL of CM2 media containing IL-2 (6000 IU/mL) and 5×10^8 feeder cells with 30 μg OKT-3 was added to each flask. On day 9-11 the culture was scaled up. The entire volume of the G-Re × 100M (1L) was transferred to a G-REX 500MCS and 4L of CM4 containing IL-2 (3000 IU/mL) was added. Flasks were incubated 5 days. Cultures were harvested and cryopreserved on Day 16.

[001686] Three different tumors were included in the comparison, two lung tumors (L4054 and L4055) and one melanoma tumor (M1085T).

[001687] CM1 (culture media 1), CM2 (culture media 2), and CM4 (culture media 4) media were prepared in advance and held at 4°C for L4054 and L4055. CM1 and CM2 media were prepared without filtration to compare cell growth with and without filtration of media.

[001688] Media was warmed at 37°C up to 24 hours in advance for L4055 tumor on REP initiation and scale-up.

Results Summary

[001689] Gen 3 fell within 30% of Gen 2 for total viable cells achieved. Gen 3 final product exhibited higher production of INF-γ after restimulation. Gen 3 final product exhibited increased clonal diversity as measured by total unique CDR3 sequences present. Gen 3 final product exhibited longer mean telomere length.

Results achieved

Cell count and % viability:

[001690] Pre REP and REP expansion on Gen 2 and Gen 3 processes followed details described above.

[001691] **Table 37: Pre-REP cell counts.** For each tumor, the two pools contained equal number of fragments. Due to the small size of tumors, the maximum number of fragments per flask was not achieved. Total pre-REP cells (TVC) were harvested and counted at day 11 for the Gen 2 process and at day 7 for the Gen 3 process. To compare the two pre-REP arms, the cell count was divided over the number of fragments provided in the culture in order to calculate an average of viable cells per fragment. As indicated in the table below, the Gen 2 process consistently grew more cells per fragment compared to the Gen 3 Process. An extrapolated calculation of the number of TVC expected for Gen 3 process at day 11, which was calculated dividing the pre-REP TVC by 7 and then multiply by 11.

Table 37: pre-REP cell counts

Tumor ID	L4054		L4055*		M1085T	
Process	Gen 2	Gen 3	Gen 2	Gen 3	Gen 2	Gen 3
pre-REP TVC	1.42E+08	4.32E+07	2.68E+07	1.38E+07	1.23E+07	3.50E+06
Number of fragments	21	21	24	24	16	16
Average TVC per fragment at pre-REP	6.65E+06	2.06E+06	1.12E+06	5.75E+05	7.66E+05	2.18E+05
Gen 3 extrapolated value at pre REP day 11	N/A	6.79E+07	N/A	2.17E+07	N/A	5.49E+06

*** L4055, unfiltered media.**

[001692] **Table 38: Total viable cell count and fold expansion on TIL final product:** For the Gen 2 and Gen 3 processes, TVC was counted per process condition and percent viable cells was generated for each day of the process. On harvest, day 22 (Gen 2) and day 16 (Gen 3) cells were collected and the TVC count was established. The TVC was then divided by the number of fragments provided on day 0, to calculate an average of viable cells per fragment. Fold expansion was calculated by dividing harvest TVC by over the REP initiation TVC. As exhibited in the table, comparing Gen 2 and the Gen 3, fold expansions were similar for L4054; in the case of L4055, the fold expansion was higher for the Gen 2 process. Specifically, in this case, the media was warmed up 24 in advance of REP initiation day. A higher fold expansion was also observed in Gen 3 for M1085T. An extrapolated calculation

of the number of TVC expected for Gen 3 process at day 22, which was calculated dividing the REP TVC by 16 and then multiply by 22.

Table 38: Total viable cell count and fold expansion on TIL final product

Tumor ID	L4054		L4055		M1085T	
Process	Gen 2	Gen 3	Gen 2	Gen 3	Gen 2	Gen 3
# Fragments	**21**	**21**	**24**	**24**	**16**	**16**
TVC /fragment (at Harvest)	3.18E+09	8.77E+08	2.30E+09	3.65E+08	7.09E+08	4.80E+08
REP initiation	1.42E+08	4.32E+07	2.68E+07	1.38E+07	1.23E+07	3.50E+06
Scale up	3.36E+09	9.35E+08	3.49E+09	8.44E+08	1.99E+09	3.25E+08
Harvest	6.67E+10	1.84E+10	5.52E+10	8.76E+09	1.13E+10	7.68E+09
Fold Expansion Harvest/ REP initiation	**468.4**	**425.9**	**2056.8**	**634.6**	**925.0**	**2197.2**
Gen 3 extrapolated value at REP harvest day 22	**N/A**	**2.53E+10**	**N/A**	**1.20E+10**	**N/A**	**1.06E+10**

*** L4055, unfiltered media.**

[001693] **Table 39: %Viability of TIL final product:** Upon harvest, the final TIL REP products were compared against release criteria for % viability. All of the conditions for the Gen 2 and Gen 3 processes surpassed the 70% viability criterion and were comparable across processes and tumors.

Table 39: % Viability of REP

Tumor ID	L4054		L4055		M1085T	
Process	Gen 2	Gen 3	Gen 2	Gen 3	Gen 2	Gen 3
REP initiation	98.23%	97.97%	97.43%	92.03%	81.85%	68.27%
Scale up	94.00%	93.57%	90.50%	95.93%	78.55%	71.15%
Harvest	87.95%	89.85%	87.50%	86.70%	86.10%	87.45%

[001694] **Table 40: Estimate cell count per additional flask for Gen 3 process.** Due to the number of fragments per flask below the maximum required number, an estimated cell

count at harvest day was calculated for each tumor. The estimation was based on the expectation that clinical tumors were large enough to seed 2 or 3 flasks on day 0.

Table 40: Extrapolated estimate cell count calculation to full scale 2 and 3 flask on Gen 3 Process

Tumor ID	L4054		L4055		M1085T	
Gen 3 Process	2 flasks	3 Flasks	2 flasks	3 Flasks	2 flasks	3 Flasks
Estimate Harvest	3.68E+10	5.52E+10	1.75E+10	2.63E+10	1.54E+10	2.30E+10

IMMUNOPHENOTYPING:

Phenotypic markers comparison on TIL final product:

[001695] Three tumors L4054, L4055, and M1085T underwent TIL expansion in both the Gen 2 and Gen 3 processes. Upon harvest, the REP TIL final products were subjected to flow cytometry analysis to test purity, differentiation, and memory markers. For all the conditions the percentage of TCR a/b+ cells was over 90%.

[001696] TIL harvested from the Gen 3 process showed a higher expression of CD8 and CD28 compared to TIL harvested from the Gen 2 process. The Gen 2 process showed a higher percentage of CD4+. **See, Figure 3 (A, B, C)**.

Memory markers comparison on TIL final product:

[001697] TIL harvested from the Gen 3 process showed a higher expression on central memory compartments compared to TIL from the Gen 2 process. **See, Figure 4 (A, B, C)**.

Activation and exhaustion markers comparison on TIL final product:

[001698] Activation and exhaustion marker were analyzed in TIL from two, tumors L4054 and L4055 to compare the final TIL product by from the Gen 2 and Gen 3 TIL expansion processes. Activation and exhaustion markers were comparable between the Gen 2 and Gen 3 processes**. See, Figure 5 (A, B); Figure 6 (A, B).**

INTERFERON GAMMA SECRETION UPON RESTIMULATION:

[001699] On harvest day, day 22 for Gen 2 and day 16 for Gen 3, TIL underwent an overnight restimulation with coated anti -CD3 plates for L4054 and L4055. The restimulation on M1085T was performed using anti-CD3, CD28, and CD137 beads. Supernatant was

collected after 24 hours of the restimulation in all conditions and the supernatant was frozen. IFNγ analysis by ELISA was assessed on the supernatant from both processes at the same time using the same ELISA plate. Higher production of IFNγ from the Gen 3 process was observed in the three tumors analyzed. **See, Figure 7 (A, B, C).**

MEASUREMENT OF IL-2 LEVELS IN CULTURE MEDIA:

[001700] To compare the IL-2 consumption between Gen 2 and Gen 3 process, cell supernatant was collected on REP initiation, scale up, and harvest day, on tumor L4054 and L4055. The quantity of IL-2 in cell culture supernatant was measured by Quantitate ELISA Kit from R&D. The general trend indicates that the IL-2 concentration remains higher in the Gen 3 process when compared to the Gen 2 process. This is likely due to the higher concentration of IL-2 on REP initiation (6000 IU/mL) for Gen 3 coupled with the carryover of the media throughout the process. **See, Figure 8 (A, B).**

METABOLIC SUBSTRATE AND METABOLITE ANALYSIS

[001701] The levels of metabolic substrates such as D-glucose and L-glutamine were measured as surrogates of overall media consumption. Their reciprocal metabolites, such lactic acid and ammonia, were measured. Glucose is a simple sugar in media that is utilized by mitochondria to produce energy in the form of ATP. When glucose is oxidized, lactic acid is produced (lactate is an ester of lactic acid). Lactate is strongly produced during the cells exponential growth phase. High levels of lactate have a negative impact on cell culture processes. **See, Figure 9 (A, B).**

[001702] Spent media for L4054 and L4055 was collected at REP initiation, scale up, and harvest days for both process Gen 2 and Gen 3. The spent media collection was for Gen 2 on Day 11, day 16 and day 22; for Gen 3 was on day 7, day 11 and day 16. Supernatant was analyzed on a CEDEX Bio-analyzer for concentrations of glucose, lactic acid, glutamine, glutamax, and ammonia.

[001703] L- glutamine is an unstable essential amino acid required in cell culture media formulations. Glutamine contains an amine, and this amide structural group can transport and deliver nitrogen to cells. When L-glutamine oxidizes, a toxic ammonia by-product is produced by the cell. To counteract the degradation of L-glutamine the media for the Gen 2 and Gen 3 processes was supplemented with Glutamax, which is more stable in aqueous

solutions and does not spontaneously degrade. In the two tumor lines, the Gen 3 arm showed a decrease in L-glutamine and Glutamax during the process and an increase in ammonia throughout the REP. In the Gen 2 arm a constant concentration of L-glutamine and Glutamax, and a slight increase in the ammonia production was observed. The Gen 2 and Gen 3 processes were comparable at harvest day for ammonia and showed a slight difference in L-glutamine degradation. **See, Figure 10 (A, B, C).**

TELOMERE REPEATS BY FLOW – FISH:

[001704] Flow-FISH technology was used to measure the average length of the telomere repeat on L4054 and L4055 under Gen 2 and Gen 3 process. The determination of a relative telomere length (RTL) was calculated using Telomere PNA kit/FITC for flow cytometry analysis from DAKO. Gen 3 showed comparable telomere length to Gen 2.

CD3 Analysis

[001705] To determine the clonal diversity of the cell products generated in each process, TIL final product harvested for L4054 and L4055, were sampled and assayed for clonal diversity analysis through sequencing of the CDR3 portion of the T-cell receptors.

[001706] Table 41: Comparison of Gen 2 and Gen3 of percentage shared unique CDR3 sequences on L4054 on TIL harvested cell product. 199 sequences are shared between Gen 3 and Gen 2 final product, corresponding to 97.07% of top 80% of unique CDR3 sequences from Gen 2 shared with Gen 3 final product.

Table 41: Comparison of shared uCDR3 sequences between Gen 2 and Gen 3 processes on L4054.

# uCDR3 (% Overlap)	All uCDR3's		Top 80% uCDR3's	
	Gen 2	Gen 3	Gen 2	Gen 3
Gen 2-L4054	8915	4355 **(48.85%)**	205	199 **(97.07%)**
Gen 3-L4054	-	18130	-	223

[001707] **Table 42: Comparison of Gen 2 and Gen3 of percentage shared unique CDR3 sequences on L4055 on TIL harvested cell product.** 1833 sequences are shared between Gen 3 and Gen 2 final product, corresponding to 99.45% of top 80% of unique CDR3 sequences from Gen 2 shared with Gen 3 final product.

Table 42: Comparison of shared uCDR3 sequences between Gen 2 and Gen 3 processes on L4055.

# uCDR3 (% Overlap)	All uCDR3's		Top 80% uCDR3's	
	Gen 2	Gen 3	Gen 2	Gen 3
Gen 2-L4055	12996	6599 **(50.77%)**	1843	1833 **(99.45%)**
Gen 3-L4055	-	27246	-	2616

[001708] CM1 and CM2 media was prepared in advanced without filtration and held at 4°C until use for tumor L4055 to use on Gen 2 and Gen 3 process.

[001709] Media was warmed up at 37°C for 24 hours in advance for tumor L4055 on REP initiation day for Gen 2 and Gen 3 process.

[001710] LDH was not measured in the supernatants collected on the processes.

[001711] M1085T TIL cell count was executed with K2 cellometer cell counter.

[001712] On tumor M1085T, samples were not available such as supernatant for metabolic analysis, TIL product for activation and exhaustion markers analysis, telomere length and CD3 - TCR vb Analysis.

CONCLUSIONS

[001713] This example compares 3 independent donor tumors tissue in terms of functional quality attributes, plus extended phenotypic characterization and media consumption among Gen 2 and Gen 3 processes.

[001714] Gen 2 and Gen 3 pre-REP and REP expansion comparison were evaluated in terms of total viable cells generated and viability of the total nucleated cell population. TVC cell doses at harvest day was not comparable between Gen 2 (22 days) and Gen 3 (16 days). Gen 3 cell doses were lower than Gen 2 at around 40% of total viable cells collected at harvest.

[001715] An extrapolated cell number was calculated for Gen 3 process assuming the pre-REP harvest occurred at day 11 instead day 7 and REP Harvest at Day 22 instead day 16. In both cases shows closer number on TVC compared to Gen 2 process, indicating that the early activation could allow an overall better performance on TIL growth. Table 4 and 5 bottom row.

[001716] In the case of extrapolated value for extra flasks (2 or 3) on Gen 3 process assuming a bigger size of tumor processed, and reaching the maximum number of fragments required per process as described. It was observed that a similar dose can be reachable on TVC at Day 16 Harvest for Gen 3 process compared to Gen 2 process at Day 22. This observation is important and indicates an early activation of the culture can allow better performance of TIL in less processing time.

[001717] Gen 2 and Gen 3 pre-REP and REP expansion comparison were evaluated in terms of total viable cells generated and viability of the total nucleated cell population. TVC cell doses at harvest day was not comparable between Gen 2 (22 days) and Gen 3 (16 days). Gen 3 cell doses were lower than Gen 2 at around 40% of total viable cells collected at harvest.

[001718] In terms of phenotypic characterization, a higher CD8+ and CD28+ expression was observed on three tumors on Gen 3 process compared to Gen 2 process. This data indicates the Gen 3 process has improved attributes of final TIL product compared to Gen 2.

[001719] Gen 3 process showed slightly higher central memory compartments compared to Gen 2 process.

[001720] Gen 2 and Gen 3 process showed comparable activation and exhaustion markers, despite the shorter duration of the Gen 3 process.

[001721] IFN gamma (IFNγ) production was 3 times higher on Gen 3 final product compared to Gen 2 in the three tumors analyzed. This data indicates the Gen 3 process generated a highly functional and more potent TIL product as compared to the Gen 2 process, possibly due to the higher expression of CD8 and CD28 expression on Gen 3. Phenotypic characterization suggested positive trends in Gen 3 toward CD8+, CD28+ expression on three tumors compared to Gen 2 process.

[001722] Telomere length on TIL final product between Gen 2 and Gen 3 were comparable.

[001723] Glucose and Lactate levels were comparable between Gen 2 and Gen 3 final product, suggesting the levels of nutrients on the media of Gen 3 process were not affected due to the non-execution of volume reduction removal in each day of the process and less volume media overall in the process, compared to Gen 2.

[001724] Overall Gen 3 process showed a reduction almost two times of the processing time compared to Gen 2 process, which would yield a substantial reduction on the cost of goods (COGs) for TIL product expanded by the Gen 3 process.

[001725] IL-2 consumption indicates a general trend of IL-2 consumption on Gen 2 process, and in Gen 3 process IL-2 was higher due to the non-removal of the old media.

[001726] The Gen 3 process showed a higher clonal diversity measured by CDR3 TCRab sequence analysis.

[001727] The addition of feeders and OKT-3 on day 0 of the pre-REP allowed an early activation of TIL and overall a better growth TIL performance using the Gen 3 process.

[001728] Table 43 describes various embodiments and outcomes for the Gen 3 process as compared to the current Gen 2 process:

Table 43: Exemplary Gen 3 process.

Step	Process Gen 2	Process Gen 3
Pre REP-day 0	≤50 fragments 1X G-Rex 100MCS 1L media IL-2 (6000IU/mL) 11 days	≤240 fragments ≤60 fragments/flask ≤4 flasks ≤2L media (500mL/flask) IL-2 (6000IU/mL) 2.5×10^8 feeder cells/flask 15µg OKT3/flask
REP Initiation	Fresh TIL direct to REP Day 11 $\leq 200 \times 10^6$ viable cells 5×10^9 feeder cells G-Rex 500MCS 5L CM2 media + IL-2 (3000 IU/mL) 150µg OKT3	Fresh TIL direct to REP Day 7 Activate entire culture 5×10^8 feeder cells 30 µg OKT3/flask G-Rex 100MCS 500mL media+ IL-2(6000IU/mL)
TIL Sub-culture or Scale up	≤5 G-REX 500MCS $\leq 1 \times 10$ viable cells/ flask 5L/flask Day 16	≤4 G-REX 500MCS Scale up entire culture 4L/flask Day 10-11
Harvest	Harvest Day 22, LOVO-automated cell washer 2 wash cycles	Harvest Day 16 LOVO- automated cell washer 5 wash cycles

Final formulation	Cryopreserved Product 300IU/mL IL2- CS10 in LN_2, multiple aliquots	Cryopreserved product 300IU/mL IL-2-CS10 in LN_2, multiple aliquots
Process time	**22 days**	**16 days**

[001729] The examples set forth above are provided to give those of ordinary skill in the art a complete disclosure and description of how to make and use the embodiments of the compositions, systems and methods of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Modifications of the above-described modes for carrying out the invention that are obvious to persons of skill in the art are intended to be within the scope of the following claims. All patents and publications mentioned in the specification are indicative of the levels of skill of those skilled in the art to which the invention pertains. All references cited in this disclosure are incorporated by reference to the same extent as if each reference had been incorporated by reference in its entirety individually.

TILs have been successfully expanded from core biopsies isolated from pancreatic tumors. TILs were also expanded from FNA's from certain tumor types. In the study no TIL growth was observed from ovarian tumors (n=3) and a colorectal tumor (n=1). TIL growth was observed from FNA's obtained from a lung tumor, melanoma, and a cervical tumor. Phenotyping and functional assays of the population of TILs from the first expansion and from the second expansion are currently in progress. The data from the study shows that a significant population of CD4+ and CD8+ T cells are present in the pre-REP aspirate.

[001730] All headings and section designations are used for clarity and reference purposes only and are not to be considered limiting in any way. For example, those of skill in the art will appreciate the usefulness of combining various aspects from different headings and sections as appropriate according to the spirit and scope of the invention described herein.

[001731] All references cited herein are hereby incorporated by reference herein in their entireties and for all purposes to the same extent as if each individual publication or patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

[001732] Many modifications and variations of this application can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments and examples described herein are offered by way of example only, and the application is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which the claims are entitled.

2018368786 05 May 2025

WHAT IS CLAIMED IS:

1. A method for expanding tumor infiltrating lymphocytes (TILs) into a therapeutic population of TILs comprising:

    i. performing a first expansion by culturing a first population of TILs from at least one fine needle aspirate (FNA) or at least one small biopsy from a tumor in a patient in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for 3 days to 12 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5 x 107 TILs by 3 days to 12 days when the first population of TILs is from a small biopsy; and

    ii. performing a second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the third population of TILs is at least 25-fold greater in number than the second population of TILs, and wherein the second expansion is performed for 3 days to 12 days in order to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs.

2. The method according to claim 1, wherein step (i) is a priming first expansion step and step (ii) is a rapid second expansion, wherein optionally after step (ii), the cells are removed from the cell culture and cryopreserved in a storage medium.

3. The method of according to any of the preceding claims, wherein steps (i) through (ii) are performed within a period of 17 days to 24 days, optionally within a period of 18 days to 22 days, further optionally within a period of 20 days to 22 days, and further optionally within a period of 22 days.

4. The method according to any of the preceding claims, wherein the cells in the therapeutic population of TILs from step (ii) express CD4, CD8, and TCR a 13 at levels similar to freshly harvested cells.

2018368786 05 May 2025

5. The method according to claim 1, wherein the APCs are peripheral blood mononuclear cells (PBMCs), wherein optionally the PBMCs are added to the cell culture in step (i) on any of days 3 through 12 after initiation of the culture in step (i) and/or in step (ii) on any of days 11 through 14 after initiation of the culture in step (i), further optionally wherein the APCs are artificial APCs (aAPCs) or autologous APCs.

6. The method according to claims 2 to 5, wherein the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs in step (ii) exhibit one or more characteristics selected from the group consisting of expression of CD27, expression of CD28, longer telomeres, increased CD57 expression, and decreased CD56 expression, relative to effector T cells and/or central memory T cells in the second population of cells, wherein optionally the effector T cells and/or central memory T cells exhibit increased CD57 expression and decreased CD56 expression.

7. The method according to any of the preceding claims, wherein the therapeutic population of TILs are for infusion into a patient.

8. The method according to claim 1, wherein the first expansion in step (i) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody and/or the second expansion in step (ii) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

9. The method according to claim 1, wherein the FNA in step (i) comprises at least 400,000 TILs.

10. The method according to claim 1, wherein the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast, and ovarian and optionally wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and

2018368786 05 May 2025

sarcoma, wherein optionally the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the patient has previously undergone surgical treatment.

11. The method according to claim 1, wherein the first population of TILs in step (i) are obtained from a FNA, wherein optionally the FNA is obtained using a 25-18 gauge needle.

12. The method according to claim 1, wherein the first population of TILs in step (i) are obtained from a small biopsy, and are optionally obtained from a core biopsy, and further optionally wherein the small biopsy is obtained using a 16-11 gauge needle.

13. The method according to claim 1, wherein step (ii) is repeated one to four times in order to obtain sufficient TILs in the therapeutic population of TILs for a therapeutically effective dosage of the TILs, optionally where the number of TILs sufficient for a therapeutically effective dosage is from $2.3x10^{10}$ to $13.7x10^{10}$, from about 7.5×10^{9} to about 75×10^{9}, or from about 1×10^{9} to about 1×10^{11}.

14. The method according to any preceding claim, wherein the first expansion is performed for 3 days to 10 days.

15. The method according to claim 14, which further comprises an additional second expansion in step (ii) which is performed by further supplementing the cell culture medium of the third population of TILs with IL-15, OX40 agonistic antibody and/or 4-IBB agonistic antibody.

16. The method according to any of claims 1 to 15, which comprises:

(a) adding the first population of TILs into a closed system;

(b) performing the first expansion by culturing the first population of TILs in a cell culture medium comprising IL-2 to produce a second population of TILs, wherein the cell culture medium is supplemented with OKT-3 at any one of days 1-3, wherein the first expansion is performed for 3 days to 12 days in order to obtain the second population of TILs, wherein the second population of TILs comprises at least 5 x 107 TILs by 3 days to 12 days when the first population of TILs is from a small biopsy, wherein the first expansion is performed in a closed container providing a first gas-permeable surface area, wherein the second population of TILs is at least 25-fold

2018368786 05 May 2025

greater in number than the first population of TILs, and wherein the transition from step (a) to step (b) occurs without opening the system;

(c) performing the second expansion by supplementing the cell culture medium of the second population of TILs with additional IL-2, OKT-3, and antigen presenting cells (APCs), to produce a third population of TILs, wherein the second expansion is performed for 3 days to 12 days to obtain the third population of TILs, wherein the third population of TILs is a therapeutic population of TILs, wherein the second expansion is performed in a closed container providing a second gas-permeable surface area, and wherein the transition from step (b) to step (c) occurs without opening the system;

(d) harvesting the therapeutic population of TILs obtained from step (c), wherein the transition from step (c) to step (d) occurs without opening the system; and

(e) transferring the harvested TIL population from step (d) to an infusion bag, wherein the transfer from step (d) to (e) occurs without opening the system.

17. The method according to claim 16, wherein step (b) is a priming first expansion step and step (c) is a rapid second expansion step, wherein optionally the method further comprises the step of cryopreserving the infusion bag comprising the harvested TIL population in step (e) using a cryopreservation process.

18. The method according to any one of claims 16 or 17, wherein the APCs are peripheral blood mononuclear cells (PBMCs) wherein optionally the PBMCs are irradiated and allogeneic and optionally, wherein the PBMCs are added to the cell culture on any of days 3 through 12 of the first period in step (b) and/or any of days 3 through 12 of the second period in step (c).

19. The method according to claim 16, wherein the first expansion in step (b) is performed by further supplementing the cell culture medium of the second population of TILs with OKT-3, IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody and/or wherein the second expansion in step (c) is performed by further supplementing the cell culture medium of the second population of TILs with IL-15, OX40 agonistic antibody and/or 4-1BB agonistic antibody.

2018368786 05 May 2025

20. The method according to claim 16, wherein the TILs in step (a) are obtained from a FNA, and optionally wherein the FNA in step (a) comprises at least 400,000 TILs further optionally wherein the FNA is obtained using a 25-18 gauge needle, and further optionally wherein the FNA is obtained from a tumor selected from the group consisting of lung, melanoma, head and neck, cervical, ovarian, pancreatic, glioblastoma, colorectal, and sarcoma, wherein optionally the lung tumor is a non-small cell lung carcinoma (NSCLC), and optionally wherein the subject has previously undergone surgical treatment.

21. The method according to claim 16, wherein the first population of TILs in step (a) are obtained from a small biopsy and optionally wherein the small biopsy is obtained from a tumor selected from the group consisting of pancreatic, melanoma, breast and ovarian, and further optionally wherein the small biopsy is obtained using a 16-11 gauge needle.

22. The method according to claim 16, wherein the harvesting in step (d) is performing using a cell processing system, wherein optionally the cell culture medium is provided in a container or cellbag and further optionally wherein the infusion bag in step (e) is a hypothermosol infusion bag.

23. The method according to claim 16, wherein steps (a) through (e) are performed within a period of 17 days to 24 days, optionally wherein steps (a) through (e) are performed within a period of 18 days to 22 days, further optionally wherein steps (a) through (e) are performed within a period of 20 days to 22 days.

24. The method according to claim 16, wherein steps (a) through (e) and optionally cryopreservation are performed in 22 days or less.

25. The method according to any one of claims 16 to 24, wherein the therapeutic population of TILs harvested in step (d) comprises sufficient TILs for a therapeutically effective dosage of the TILs, optionally wherein the number of TILs sufficient for a therapeutically effective dosage is from $2.3x10^{10}$ to $13.7x10^{10}$, from about 7.5×10^{9} to about 75×10^{9}, or from about 1×10^{9} to about 1×10^{11}.

26. The method according to any one of claims 16 to 25, wherein steps (a) through (d) are performed in a single container, wherein performing steps (a) through (d) in a single

2018368786 05 May 2025

container results in an increase in TIL yield per resected tumor as compared to performing steps (a) through (d) in more than one container, optionally wherein the antigen-presenting cells are added to the TILs during the second expansion in step (c) without opening the system, wherein optionally the third population of TILs comprises an increased subpopulation of effector T cells and/or central memory T cells relative to the second population of TILs, wherein the effector T cells and/or central memory T cells in the therapeutic population of TILs exhibit one or more characteristics selected from the group consisting of expressing CD27+, expressing CD28+, longer telomeres, increased CD57 expression, and decreased CD56 expression relative to effector T cells, and/or central memory T cells obtained from the second population of cells, and further optionally wherein the effector T cells and/or central memory T cells obtained from the third population of TILs exhibit increased CD57 expression and decreased CD56 expression relative to effector T cells and/or central memory T cells obtained from the second population of cells.

Figure 1

Process 2A: about 22 days from Steps A - E

1. **STEP A**

Obtain Patient Tumor Sample

2. **STEP B**

Fragmentation and First Expansion

3 days to 14 days

3. **STEP C**

First Expansion to Second Expansion Transition

No Storage and Closed System

4. **STEP D**

Second Expansion

IL-2, OKT-3, and antigen-presenting feeder cells

Closed System

5. **STEP E**

Harvest TILS from Step D

Closed System

6. **STEP F**

Final Formulation and/or Transfer to Infusion Bag

(optionally cryopreserve)

Figure 2A



Figure 2B

Figure 2B

Figure 2A



Figure 2C

Figure 2C


Figure 2B
Spent Medium removal ≤ 5x G-Rex 500MCS
supernate
Pool supernate 10L Bioprocess bag(s)
Mycoplasma PCR Sample Diluent
3L Cell culture Bag (Origen EV3000 or equivalent)
Cell Count & Viability NC-200 4x 1mL sample
Wash Buffer PL-A + 1% HSA
Volume Reduction LOVO 2 Cycles, 10:1 concentration 4000 RPM, 75mL/min
Formulate 1:1 with cold CS10*
IL-2 300 IU/mL
Add IL-2
*Formulate bulk Product Bag stored at 2-8°C in between processing steps
Satelllite Vial Prep
Cryopreserved Satellite Vial Sample 10 x 0.5mL/vial
Remove QC Release Test Samples & Air (Mycoplasma, Endotoxin, Sterility, Gram Stain, Cell Count, viability, flow, INF-g, Retain) (20mL)
Aliquot into CS750 cryobags 90-120mL/ bag
Manual Fill 0.5 mL / vial
Controlled Rate Freeze Preset-6 Profile
Visually Inspect
Store in vapor phase LN
Ship*


≤4.0 cm diameter
Excise tumor
Express courier from Clinical Site to CMO
Cut into fragments
≤50 Fragments
ex vivo culture
+ IL-2
Bulk TILs (~10^8)
Direct To REP
Tumor Fragment Culture (Pre-REP) 11 days
Rapid Expansion (REP) 11 days
GMP MANUFACTURING FACILITY
NMA-LD preconditioning therapy
Thaw, Infuse + IL-2 administration
Express courier in cryoshipper from CMO to Clinical Site
-190°C
LN2 cryopreserved LN-144 infusion product controlled rate freeze
Harvest
Expansion ex vivo scale up
IL-2 + OKT3
Co-Culture TIL And Feeder Cells

Figure 3


Excise Tumor
≤4.0 cm Diameter
≤96hrs
8-27 mm3 Fragments
≤50 Fragments
Closed single use bioreactor
+ IL-2
Bulk TILs (≤200x10)
Direct To Rapid Expansion
Tumor Fragment Culture (Pre-REP) 11 days
Rapid Expansion (REP) 11 days
GMP MANUFACTURING FACILITY
Thaw and infuse + IL-2
Scheduled cryoshipper
LN2-freeze in aliquots
Semi-automated Harvest/Wash/ Formulate/Aliquot
Scale Up
IL-2 + OKT3
Co-Culture TIL And Feeder Cells

Figure 4

Figure 5

Process 1C: 43-55 Days for Steps A - E	**Process 2A: about 22 days from Steps A - E**
1. STEP A Obtain Patient Tumor Sample	**1. STEP A** Obtain Patient Tumor Sample
2. STEP B Fragmentation and First Expansion 11 days to 21 days	**2. STEP B** Fragmentation and First Expansion 3 days to 14 days
3. STEP C First Expansion to Second Expansion Transition Optional Storage until Selection	**3. STEP C** First Expansion to Second Expansion Transition No Storage and Closed System
4. STEP D Second Expansion IL-2, OKT-3, antigen-presenting feeder cells Optionally repeat one or more times	**4. STEP D** Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells Closed System
5. STEP E Harvest TILS from Step D	**5. STEP E** Harvest TILS from Step D Closed System
6. STEP F Final Formulation and/or Transfer to Infusion Bag	**6. STEP F** Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)

Figure 6

Process Step	Process 1C Embodiment	Process 2A Embodiment	Advantages
Pre-REP	• 4 fragments per 10 GREX-10 flasks • 11-21 day duration	• 40 fragments per 1 GREX-100M flask • 11 day duration	• Increased tumor fragments per flask • Shortened culture time • Reduced number of steps • Amenable to closed system
Pre-REP to REP Transition	• Pre-REP TIL are frozen until phenotyped for selection then thawed to proceed to the REP (~day 30) • REP requires >40×10^6 TIL	• Pre-REP TIL directly move to REP on day 11 • REP requires 25-200×10^6 TIL	• Shortened pre-REP-to-REP process • Reduced number of steps • Eliminated phenotyping selection • Amenable to closed system
REP	• 6 GREX-100M flasks on REP day 0 • 5×10^6 TIL and 5×10^8 PBMC feeders per flask on REP day 0 • Split to 18-36 flasks on REP day 7 • 14 day duration	• 1 GREX-500M flask on day 11 • 25-200×10^6 TIL and 5x10^9 PBMC feeders on day 11 • Split to ≤ 6 GREX-500M flasks on day 16 • 11 day duration	• Reduced number of steps • Shorter REP duration • Closed system transfer of TIL between flasks • Closed system media exchanges
Harvest	• TIL harvested via centrifugation	• TIL harvested via LOVO automated cell washing system'	• Reduced number of steps • Automated cell washing • Closed system • Reduced loss of product during wash
Final Formulation	• Fresh product in Hypothermosol • Single infusion bag • Limited shipping stability	• Cyropreserved product in PlasmaLyte-A + 1% HSA and CS10 stored in LN_2 • Multiple aliquots • Longer shipping stability	• Shipping flexibility • Flexible patient scheduling • More timely release testing
Overall Estimated Process Time	• 43-55 days	• 22 days	• Faster turnaround to patient

Figure 7A

Small Biopsy Process: Version 1 **About 21-33 days from Steps A - E**
1. STEP A Obtain Patient Tumor Sample
2. STEP B First Expansion 1 days to 19 days (with OKT-3 and antigen-presenting feeder cells at day 3)
3. STEP C First Expansion to Second Expansion Transition No Storage and Closed System
4. STEP D Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells 11 to 14 days Closed System
5. STEP E Harvest TILS from Step D Closed System
6. STEP F Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)

Figure 7B

Small Biopsy Process: Version 2 **About 17-24 days from Steps A - E**
1. STEP A Obtain Patient Tumor Sample
2. STEP B First Expansion 3 days to 12 days (with OKT-3 and antigen-presenting feeder cells at any one of days 1-3)
3. STEP C First Expansion to Second Expansion Transition No Storage and Closed System
4. STEP D Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells 3 to 12 days Closed System
5. STEP E Harvest TILS from Step D Closed System
6. STEP F Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)

Figure 7C

Small Biopsy Process: Version 1 About 21-33 days from Steps A - E	Small Biopsy Process: Version 2 (GEN3 based process) About 17-24 days from Steps A - E	Process 2A: About 22 days from Steps A - E
1. **STEP A** Obtain Patient Tumor Sample	1. **STEP A** 1. Obtain Patient Tumor Sample	1. **STEP A** Obtain Patient Tumor Sample
2. **STEP B** First Expansion 1 days to 19 days (with OKT-3 and antigen-presenting feeder cells at day 3)	2. **STEP B** First Expansion 3 days to 12 days (with OKT-3 and antigen-presenting feeder cells at any one of days 1-3)	2. **STEP B** Fragmentation and First Expansion 3 days to 14 days
3. **STEP C** First Expansion to Second Expansion Transition No Storage and Optionally Closed System	3. **STEP C** First Expansion to Second Expansion Transition No Storage and optionally Closed System	3. **STEP C** First Expansion to Second Expansion Transition No Storage and Closed System
4. **STEP D** Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells 11 to 14 days Optionally Closed System	4. **STEP D** Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells 3 to 12 days Optionally Closed System	4. **STEP D** Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells Closed System
5. **STEP E** Harvest TILS from Step D Optionally Closed System	5. **STEP E** Harvest TILS from Step D Closed System	5. **STEP E** Harvest TILS from Step D Closed System
6. **STEP F** Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)	6. **STEP F** Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)	6. **STEP F** Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)

Figures 8A and 8B

Figure 8A



Figure 8B



Figures 9A and 9B

Figure 9A



Figure 9B



Figures 9C and 9D

Figure 9C



Figure 9D



Figures 10A and 10B

Figure 10A



Figure 10B



Figures 11A and 11B

Figure 11A



Figure 11B



Figure 11C

Figure 11C



Figures 12A and 12B

Figure 12A



Figure 12B



Figures 13A and 13B

Figure 13A



Figure 13B



Figure 14


Total Cell Counts
2.5×10^07
2.0×10^07
1.5×10^07
1.0×10^07
5.0×10^06
0
Fragments
Aspirates
Fragments
Aspirates

Figure 15


Ovarian pre-REP
(4 frags/G-Rex 10)
Total Cell Count
6×10^06
4×10^06
2×10^06
0
*
IL-2
IL-2/IL-15/IL-21
OV8011
OV8012
OV8013
* 30% viable

Figures 16A and 16B

Figure 16A



Figure 16B



Figures 17A and 17B

Figure 17A



Figure 17B



Figures 17C and 17D

Figure 17C



Figure 17D

Tumor ID	Cytokines	Days to reach Stage #1	Acceptance Criteria for Stage #2	Assayed Cells for Stage #2	Acceptance Criteria for Stage #3	Assayed Cells for Stage #3
Pan-17-004	IL-2 #1	27	20x10⁶	116x10⁶	100x10⁶	Media Failure
	IL-2 #2	27		130x10⁶		Media Failure
	Cell Culture Additives #1	14		Media Failure		N/A
	Cell Culture Additives #2	32		132x10⁶		148x10⁶
Pan-17-007	IL-2	16		Did not meet criteria to progress to REP		N/A
	Cell Culture Additives	25		118x10⁶		244x10⁶
Pan-17-008	IL-2	Did not meet criteria for G-Rex 10		N/A		N/A
	Cell Culture Additives	21		105x10⁶		144x10⁶

Figure 18A

Histology	FNA/Biopsy	Initial Cell Count (FNA's)	Conditions	Days in Culture	Final Cell Counts	Assays
Pancreatic (P7015)	Cores 24 well plate	N/A	**ONE CORE** 1) IL-2 (3 wells) 2) IL-2/IL-15/IL-21 (2 wells) **TWO CORES** 1) IL-2 (1 well) 2) IL-2/IL-15/IL-21 (1 wells)	28 days	**ONE CORE** IL-2 1.2e6 viable cells/well average IL-2/IL-15/IL-21 2e6 viable cells/well average **TWO CORES** No growth	**Phenotypic** FACS **Functional** CD107a
Pancreatic (P7017)	Cores 24 well plate	N/A	**ONE CORE**	N/A	No TIL growth	
Pancreatic (P7019)	Cores G-Rex 10	N/A	**6 CORES**	In progress		
Lung (L4032)	FNA 24 well plate	1.6e6 cells (28% viable)	1) IL-2 (2 wells) 2) IL-2/IL-15/IL-21 (2 wells) + Fragments	13 days	IL-2 4.64e5cells/well average, 76 % viable IL-2/IL-15/IL-21 7.75e5 cells/well average, 94% viable	**Phenotypic** FACS **Functional** CD107a (N/A) **REP** (recently harvested)
Lung (L4033)	**FNA G-Rex 10**	5.5e6 cells (76% viable)	1) IL-2 2) IL-2/IL-15/IL-21 + Fragments	12 days	IL-2 6.e6, 73 % viable IL-2/IL-15/IL-21 No detectable cells (no assay were run on this sample)	**Phenotypic** FACS **Functional** CD107a **REP** (recently harvested)
Lung (L4034)	FNA G-Rex 10	N/A			No TIL growth	No assays

Figure 18B

Histology	FNA/Biopsy	Initial Cell Count (FNA's)	Conditions	Days in Culture	Final Cell Counts	Assays
Colorectal (CC10024)	FNA 24 well plate	8.0e5 viable cells, 42% viable	1) IL-2 (2 wells) 2) IL-2/IL-15/IL-21 (2 wells) Fragments as a control	27 days	No TIL growth	No assays
Ovarian (OV8012)	FNA 24 well plate	3.19e5 viable cells, 28% viable	1) IL-2 (2 wells) 2) IL-2/IL-15/IL-21 (2 wells)		No TIL growth	No assays
Ovarian (OV8013)	FNA 24 well plate	8.3e5 viable cells, 11% viable	1) IL-2 (2 wells) 2) IL-2/IL-15/IL-21 (2 wells)		No TIL growth	No assays
Ovarian (OV8014)	FNA 24 well plate	1.0e5 viable cells, 7% viable	1) IL-2 (1 well) 2) IL-2/IL-15/IL-21 (1 well)		No TIL growth	No assays
Cervical (C2005)	FNA 24 well plate	6.84e5 (13% viable)	1) IL-2 (2 wells) 2) IL-2/IL-15/IL-21 (2 wells) + Fragments	21 days	IL-2 3.45e5 viable cells/well average, 39% viable IL-2/IL-15/IL-21 8.69e5 viable cells/well average, 40% viable	**Phenotypic** FACS **Functional** CD107a
Melanoma (M1101)	FNA G-Rex 10	1.96e6 (65% viable)	IL-2 (1 G-Rex 10) + Fragments	12 days	IL-2 3.0e6, 78 % viable	**Phenotypic** FACS **Functional** CD107a **REP** (recently harvested)
Melanoma (M1107)	FNA G-Rex 10	2.25e6 (69% viable)	IL-2 (2 G-Rex 10) IL-2/IL-15/IL-21 (2 G-Rex) + Fragments			Pre-REP in progress

Figures 19A and 19B

Figure 19A



Figure 19B



Figure 19C


80
60
40
20
0
% CD107a+
Fragments
Aspirates
Unstim
PMA
Unstim
PMA
CD4+
CD8+

Figure 20

Tumor Name	Cell Growth	Analysis
UPMC-3 Reported	24 well plate: 1) IL-2 versus 2) Cytokine Additives	Cell growth, viability, phenotyping, CD107a
UPMC-9 Reported	Expanded poorly	N/A
UPMC-17 Reported	Contaminated culture	N/A
UPMC-26	Expanded poorly	N/A
UPMC-30	pre-REP Day 11: 1.6e6, 40% viable pre-REP Day 18: 3.49e7, 89% viable	Cell growth, viability, phenotyping, CD107a, IFNγ, REP initiation and analysis
UPMC-61	pre-REP Day 11: 1.52e5, 66% viable pre-REP Day 18: 1.14e7, 1.14e7, 40% viable	Cell growth, viability, phenotyping, CD107a, IFNγ, REP initiation and analysis
UPMC-59	Expanded poorly:	N/A
UPMC-61	pre-REP: In progress	


A)
% of Live Cells
150
100
50
0
pre-REP
REP
CD4+
CD4+
CD27+
CD28+
TCM
TSCM
TEMRA
TEM
B)
% of Live Cells
150
100
50
0
pre-REP
REP
CD8+
CD8+
CD27+
CD28+
TCM
TSCM
TEMRA
TEM

Figure 21


A)
% of Live Cells
100
80
60
40
20
0
CD4
CD8
pre-REP
REP
B)
IFNγ (pg/ml)
40000
30000
20000
10000
0
pre-REP
REP
pre-REP
REP

Figure 22

Figure 23


A)
Total cell count
1.5×10^8
1×10^8
5×10^7
2×10^7
1.5×10^7
1×10^7
5×10^6
0
IL-2 alone
IL-2 + additives
n=6
B)
% of CD3+/CD3-CD56+/CD3-
100
90
80
70
30
20
10
0
CD3+
CD3-CD56+/CD3-
Low yield
High yield
IL-2 alone

Figure 24

A)


pre-REP CD4/8 ratio
0.7
CD8/CD4 ratio
3
2
1
0
Low yield
High yield

B)


Pre-REP vs Post-REP CD4/8 ratio
CD8/CD4 ratio
6
5
4
3
2
1
0
Low yield
High yield
PreREP
REP

Figure 25



Figure 26


A)
S13006 (UPMC 58)
Total cell count
3×10^7
2.5×10^7
2×10^7
1.5×10^7
1×10^7
1000000
800000
600000
400000
200000
0
PreREP S13006 NT
PreREP S13006 Culture additive
B)
% of T-cell subsets
100
80
60
40
20
0
CD3+
CD4+
CD8+
PreREP S13006 NT
PreREP S13006 additive

Figure 27


A)
Pre-REP cell count
Total cell count
2×10^6
1.5×10^6
1×10^6
5×10^5
0
C2007
B)
TCRab+
TCRab+ CD8+
TCRab+ CD4+
CD8+CD27+
CD4+CD27+
CD8+CD28+
CD4+CD28+
% of T-cell subset
100
80
60
40
20
0
TCRab+
TCRab+ CD8+
TCRab+ CD4+
CD8+CD27+
CD4+CD27+
CD8+CD28+
CD4+CD28+

Figure 28


A)
ES19001
Total cell count
1.5×10^7
1×10^7
5×10^6
0
ES19001
B)
ES19001 phenotype
% of T-cell subsets
100
80
60
40
20
0
CD3+CD16-
CD4+
CD8+
NK
CD3+CD16-
CD4+
CD8+
NK


A)
Oral Cancer
Total cell count
5×10^7
4×10^7
3×10^7
2×10^7
1×10^7
0
B)
% of T-cell subsets
100
80
60
40
20
0
CD3+CD16-
CD3+CD4+
CD3+CD8+

Figure 29

Figure 30

Tumor Name	Cell Growth	Analysis
UPMC-3 (P7011) Reported	24 well plate: 1) IL-2 versus 2) Cytokine Additives	Cell growth, viability, phenotyping, CD107a
UPMC-9 (P7017) Reported	Expanded poorly	N/A
UPMC-17 (P7019) Reported	Contaminated culture. Discarded.	N/A
UPMC-26 (P7022)	Expanded poorly	N/A
UPMC-30 (P7028)	pre-REP Day 12: 1.6e6, 40% viable pre-REP Day 21: 3.49e7, 89% viable	Cell growth, viability, phenotyping, CD107a, IFNγ, REP initiation and analysis
UPMC-61 (P7031)	pre-REP Day 12: 1.52e5, 66% viable pre-REP Day 28: 1.14e7, 1.14e7, 40% viable	Cell growth, viability, phenotyping, CD107a, IFNγ, REP initiation and analysis
UPMC-59 (P7033)	Expanded poorly	N/A
UPMC-61 (P7034)	pre-REP: In progress	

Figure 31

A)
IL-2 treated well
Pancreatic Core Biopsies
1.3e6 viable cells/well
1.2e6 viable cells/well average
B)
IL-2/15/21 treated well
2.3e6 viable cells/well
2.0e6 viable cells/well average

Figure 32


A)
IL-2 treated well
14.5
CD107a+ (CD8+)
B)
IL-2/15/21 treated well
44.1
CD107a+ (CD8+)

Figure 33


A)
P7028 Core Biopsies
CD4
CD8
CD45RA
CCR7
B)
P7031 Core Biopsies
CD4
CD8
CD45RA
CCR7

Figure 34


Total Cell Expansion
5×10^07
4×10^07
3×10^07
2×10^07
1×10^07
0
N=7
N=2
Fragments
Cores

Figure 35


A)
% of Live Lymphocytes
150
100
50
0
Fragments
Cores
CD28+
CD27+
CD28+
CD27+
B)
% of Live Lymphocytes
100
75
50
25
0
Fragments
Cores
TCM
TSCM
TEMRA
TEM
CD4+
TCM
TSCM
TEMRA
TEM
CD8+

Figure 36



Figure 37


A)
% CD107a
100
80
60
40
20
0
Unstimulated
PMA
Fragments
Cores
Fragments
Cores
CD4+
Fragments
Cores
Fragments
Cores
CD8+
B)
IFNγ (pg/ml)
50000
40000
30000
20000
10000
0
Fragments
Cores

Figure 38A


1-2 cores
+IL-2
+/- 30 ng OKT3
& 10e6 feeders
2-day
ex vivo culture
+/- IL15&21
(pancreatic cores)
10- to 19 day
co-culture
REP1 TIL
Cell count
Phenotype
Function
+30 ng OKT3
+13E6 feeders
+IL2
11- to 14-day
co-culture
REP2 TIL
Cell count
Phenotype
Potency

Figure 38B


1-2 cores
+IL-2
+OKT3
+feeders
Day 3
11-day
co-culture
REP1 TIL
Cell count
Phenotype
Function
+ OKT3
+ feeders
+IL2
11-day
co-culture
REP2 TIL
Cell count
Phenotype
Potency

Figure 39

	Culture conditions	Cores / cond. (#)	REP1	REP2
Pancreas (P7034)	REP1: + TC +OKT3 & feeders vs NO OKT3 & feeders REP2: +/- triple cocktail	2 (2 conditions)	**>50E6 cells** at Day 17 ONLY in the presence of OKT3 & feeders	**207- & 162-fold** expansion +/- triple cocktail **~3000 pg/ml IFNγ** +/- triple cocktail
Pancreas (P7037)	REP1: IL2 versus TC REP2: +/- triple cocktail	4 (2 conditions)	**>50E6 cells** at Day 11 only in the presence of OKT3 & feeders +/- TC	**207- & 231-fold** expansion +/- triple cocktail **~1000 pg/ml IFNγ**+/- triple cocktail
Pancreas (P7039)	REP1: TC + OKT3& feeders, TC NO OKT3 & feeders IL-2 + OKT3& feeders, IL-2 NO OKT3& feeders REP2: +/- TC	2 (4 conditions)	**>50E6 cells** at Day 12 (+/- TC) ONLY in the presence of OKT3 & feeders	**662- & 582-fold** expansion +/- triple cocktail **~10K & 3K pg/ml IFNγ**+/- triple cocktail

Figure 40


P7034
P7037
P7039
Total Cell Number
Days in culture
P7037+ IL-2
P7037 + TC
P7039 + IL-2
P7039 +TC
Total Cell Number
Days in culture

Figure 41

	Culture conditions	Cores / cond. (#)	REP1	REP2
Melanoma (M1114)	REP1: IL2 REP2: IL-2	2	**>50E6 cells** at Day 11	**215-fold** expansion **~1000 pg/ml IFNγ**
Melanoma (M1115)	REP1: IL2 REP2: IL-2	2	**>50E6 cells** at Day 12	**662-fold** expansion **~3000 pg/ml IFNγ**
Melanoma (M1118)	REP1: IL-2 REP2: IL-2	1	**>50E6 cells** at Day 11	**215-fold** expansion **~10,000 pg/ml IFNγ**
** Melanoma (M1121)	REP1: IL2 REP2: IL-2	1	**50E6 cells** at Day 9	**155-fold** expansion **~2000 pg/ml IFNγ**

Figure 42


M1114
M1115
M1118
M1121
Total Cell Number
2.5×10^08
2.0×10^08
1.5×10^08
1.0×10^08
5.0×10^07
0
6
10
11
12
13
14
15
16
17
18
19
20
21
Days in culture

Figure 43


REP1
REP2
Same trend observed in the pancreatic cores
% of CD3+
100
50
0
Lag3
PD1
Tigit
TIM3
CD4
CD8
CD69 and CD103 are reduced in REP2, also observed with the pancreatic cores
2B4
BTLA
CD25
CD69
CD103
KLRG1
CD4
CD8
© 2017, Iovance Biotherapeutics

Figure 44


Melanoma Fragments
pre-REP
REP
% of CD3+
Melanoma Cores
REP1
REP2
% of CD3+
Melanoma Fragments
N=6
pre-REP
REP
% of Live Cells
CD4+
CD8+
Melanoma Cores
N=4
REP1
REP2
% of Live Cells
CD4+
CD8+

Figure 45

	Culture conditions	Cores / cond. (#)	REP1	REP2
Breast (EP11032)	REP1: IL2 REP2: IL-2	1 vs 2	>50E6 cells at Day 11	173 and 158-fold (1/2 cores) expansion ~ pg/ml IFNg
Ovarian (OV8016)	REP1: IL2 +/- OX40 REP2: IL-2 +/- OX40 (in progress)	2	>50E6 cells at Day 11 (OX40 condition)	In progress

Figure 46


OV8016
OKT3+ Feeders
OKT3+ Feeders + OX40
Total Cell Number
8×10^07
6×10^07
4×10^07
2×10^07
0
REP1 (D11)
EP11032
D11
D14
Total Cell Number
2×10^08
2×10^08
1×10^08
5×10^07
0
1 core
2 cores
REP1

Figure 47


P7028 Core Biopsies
P7031 Core Biopsies
CD4
CD8
CD45RA
CCR7
CD4+
CD8+
CD4- CD8-


Total Cell Expansion
5×10^07
4×10^07
3×10^07
2×10^07
1×10^07
0
N=7
N=2
Fragments
Cores
Pre-REP

Figure 48

Figure 49


% of Live Lymphocytes
150
100
50
0
Fragments
Cores
CD28+
CD27+
CD28+
CD27+
% of Live Lymphocytes
100
75
50
25
0
Fragments
Cores
TCM
TSCM
TEMRA
TEM
CD4+
TCM
TSCM
TEMRA
TEM
CD8+

Figure 50


% of Live Lymphocytes
150
100
50
0
Fragments
Cores
PD-1
LAG3
TIM3
TIGIT
KLRG1
CD4+
PD-1
LAG3
TIM3
TIGIT
KLRG1
CD8+

Figure 51


IFNγ (pg/ml)
50000
40000
30000
20000
10000
0
Fragments
Cores
% CD107a
100
80
60
40
20
0
Unstimulate
PMA
Fragments
Cores
Fragments
Cores
CD4
CD8

Figure 52


IL-2 REP1 - IL2 REP2
TC REP1 - TC REP2
TC REP1 - IL-2 REP2
IL2 REP1 - TC REP2
Total Cell Expansion
1.5×10^08
1.0×10^08
5.0×10^07
0
REP2

Figure 53


IL-2 REP1 - IL2 REP2
TC REP1 - TC REP2
IL-REP1 - TC REP2
TC REP1 - IL-2 REP2
% of CD3
150
100
50
0
CD4+
CD27+ CD28+
CD8+
CD27+ CD28+
CD4
CD8
% of CD3+
100
80
60
40
20
0
TCM
TSCM
TEMRA
TEM
TCM
TSCM
TEMRA
TEM
CD4
CD8

Figure 54


IL-2 REP1 - IL2 REP2
TC REP 1 - TC REP 2
IL-2 REP1 - TC REP2
TC REP1- IL2 REP2
% of CD107a
100
80
60
40
20
0
Unstim
Stimulated
Unstim
Stimulated
REP2

Figure 55


REP1
IL-2 REP1 - IL-REP2
TC REP 1 - TC REP 2
IL-2 REP1 - TC REP2
TC REP 1 - IL-2 REP2
IFNγ (pg/ml)
40000
30000
20000
10000
0
REP1
REP2
Treated with the TC

Figure 56


"Gen2" like
pre-REP
REP
Cores cultured using Gen2 like conditions (ie. NO OKT3& feeders)
N=2
% Live Cells
150
100
50
0
CD4+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
CD8+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
All cores were cultured in the TC
REP1-REP2
REP1
REP2
Cores cultured with OKT3& feeders at Day 3.
N=3
% Live Cells
150
100
50
0
CD4+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
CD8+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM

Figure 57


% of CD3+
100
80
60
40
20
0
pre-REP
REP
N=2
Lag3
PD1
Tigit
TIM3
CD4
CD8
% of CD3
REP1
REP2
N=3

Figure 58


pre-REP
REP
N=2
% of CD3+
150
100
50
0
2B4
BTLA
CD25
CD69
CD103
KLRG1
CD4
CD8


REP1
REP2
N=3
% of CD3+
150
100
50
0
2B4
BTLA
CD25
CD69
CD103
KLRG1
CD4
CD8

Figure 59


% of CD107a
100
80
60
40
20
0
Unstimulated
PMA
N=2
CD4
CD8
pre-REP
REP
N=3
REP1
REP2
Treated with the TC

Figure 60


Unstimulated
PMA
N=3
IFNγ (pg/ml)
40000
30000
20000
10000
0
REP1
REP2

Figure 61


REP1 (IL-2)
REP2 (IL-2 REP1-IL-2 REP2)
% of Live Cells
100
50
0
CD3+
CD4+
CD27+
CD28+
CD8+
CD27+
CD28+
CD4
CD8
% of CD3+
150
100
50
0
TCM
TSCM
TEMRA
TEM
TCM
TSCM
TEMRA
TEM
CD4
CD8

Figure 62


% of CD107a
80
60
40
20
0
Unstimulated
Stimulated
CD4+
CD8+
CD4+
CD8+
REP1
REP2
IL-2 culture conditions
IFNγ (pg/ml)
25000
20000
15000
10000
5000
0
NT
OKT3
NT
OKT3
REP1
REP2

Figure 63


Melanoma Fragments
pre-REP
REP
% of CD3+
100
80
60
40
20
0
Lag3
PD-1
TIGIT
TIM3
CD4
LAG3+
PD-1
TIGIT
TIM3
CD8
Melanoma Cores
REP1
REP2
% of CD3+
100
80
60
40
20
0
Lag3
PD1
TIGIT
TIM3
CD4
Lag3
PD1
TIGIT
TIM3
CD8

Figure 64


Melanoma Cores
REP1
REP2
% of CD3+
100
80
60
40
20
0
CD25
CD69
CD103
KLRG1
CD4
CD8
Melanoma Fragments
pre-REP
REP
% of CD3+
100
80
60
40
20
0
CD25
CD69
CD103
KLRG1
CD4
CD8

Figure 65


Melanoma cores
Unstimulated
Stimulated
% of CD107a
0
20
40
60
80
CD4+
CD8+
CD4+
CD8+
REP1
REP2
Melanoma fragments
Unstimulated
Stimulated
% of CD107a
0
20
40
60
80
CD4+
CD8+
CD4+
CD8+
pre-REP
REP

Figure 66


% of Live Cells
150
100
50
0
OKT3 + Feeders
OKT3 + Feeders +OX40
CD3+
CD4+
CD27+ CD28+
TCM TSCM TEMRA TEM
CD8+
CD27+ CD28+
TCM TSCM TEMRA TEM
CD4
CD8
% of CD3+
100
80
60
40
20
0
2B4 BTLA CD25 CD69 CD103 KLRG1 Lag3 PD1 Tigit TIM3
2B4 BTLA CD25 CD69 CD103 KLRG1 Lag3 PD1 Tigit TIM3
CD4
CD8

Figure 67


Ovarian Cores
NT
OKT3
IFNγ (pg/ml)
40000
30000
20000
10000
0
OKT3 + F
OKT3 + F + OX40
REP1
Ovarian Cores
Unstimulated
Stimulated
% of CD107a
40
30
20
10
0
CD4+
CD8+
CD4+
CD8+
NT
REP1
OX40

Figure 68


% of CD3
150
100
50
0
REP1
REP2
CD4+
CD27+CD28+
TCM TSCM TEMRA EM
CD8+
CD27+CD28+
TCM TSCM TEMRA EM
CD4
CD8
% of CD3
100
80
60
40
20
0
REP1
REP2
2B4
BTLA
CD25
CD69
CD103
KLRG1
Lag3
PD1
Tigit
TIM3
CD4
CD8

Figure 69 A & B


Breast Cores
Unstimulated
Stimulated
% CD107a
0
10
20
30
40
50
1 core
2 core
1 core
2 core
CD4
REP1
CD8
Breast Cores
Unstimulated
Stimulated
% CD107a
0
10
20
30
40
50
1 core
2 core
1 core
2 core
CD4
REP2
CD8

Figure 69C


Breast Cores
NT
OKT3
IFNγ (pg/ml)
40000
30000
20000
10000
0
1 core
2 core
1 core
2 core
REP1
REP2

Figure 70


Day 0
•1-2 cores are placed in a G-Rex 10 in CM2 with IL-2
Day 3
•Add OKT3 + feeders
Day 11
•Volume reduce the culture media, count cells, and harvest REP1
•The following assays are run
•Cell count, viability
•Phenotype (TIL1, TIL2, TIL3)
•CD107a
•IFNγ (restimulation assay)
•REP2 initation
•Add OKT3 + feeders
D22
•Volume reduce the culture media, count cells, and harvest REP2
•The following assays are run on the harvested REP2 product
•Cell count, viability
•Phenotype (TIL1, TIL2, TIL3)
•CD107a
•IFNγ (restimulation assay)
•TCRvβ (freeze 1e6 for RNA-sequencing by iRepetoire)

Figure 71


IFNγ (pg/ml)
40000
30000
20000
10000
0
NT
OKT3
P7034
P7037
P7039

Figure 72


% of CD107a
80
60
40
20
0
Unstimulated
Stimulated
CD4+
CD8+
IFNγ (pg/ml)
25000
20000
15000
10000
5000
0
NT
OKT3
M1114
M1115
M1118
M1121

Figure 73A

	Culture conditions	Cores / cond. (#)	REP1	REP2
Breast (EP11026)	REP1: IL2 REP2: IL-2	1 vs 2	>50E6 cells at Day 11 (2 cores)	173 and 158-fold (1/2 cores) expansion ~ 34,0000 and 23,000 pg/ml IFNγ
Breast (EP11045)	REP1: IL-2 (+/- OKT3+feeders) REP2: IL-2 (OKT3+feeders)	2	>50e6 cells at Day 11 (OKT3+ feeders)	1178 and 1920-fold (+/- OKT3+feeders REP1) ~ 99,000 and 45,000 pg/ml IFNγ
Ovarian (OV8016)	REP1: IL2 +/- OX40 REP2: IL-2 +/- OX40	2	>50E6 cells at Day 11 (OX40 condition)	1860 and 1540-fold (NO OX40/OX40) expansion ~ 10,000 and 20,000 pg/ml IFNγ
Ovarian (OV8031)	REP1: IL2 REP2: IL-2	1(small core)	1.3e6 cells at Day 11	400-fold expansion ~ ELISA is in progress

Figure 73B


EP11045
IL-2
IL-2 + OKT3 + Feeders
1.5×10^8
1.0×10^8
5.0×10^7
0
OV8031
D11 (IL-2)
6×10^7
4×10^7
2×10^7
0
EP11028
D11
D14
2×10^8
1.5×10^8
1×10^8
5×10^7
0
1 core
2 cores
OV8016
OKT3+ Feeders
OKT3+ Feeders + OX40
8×10^07
6×10^07
4×10^07
2×10^07
0
Total Cell Number

Figure 74


EP11045
IL2
IL-2+OKT3 +Feeders
% Live Cells
100
80
60
40
20
0
CD4
CD8
REP1
REP2
CD4
CD27
CD28
CD127
TEMRA
TSCM
TCM
TEM
CD8
CD27
CD28
CD127
TEMRA
TSCM
TCM
TEM

Figure 75


EP11045
IL-2
IL-2+OKT3 +Feeders
% Live cells
100
80
60
40
20
0
CD25
CD69
KLRG1
Lag3
PD1
Tigit
TIM3
CD4
CD8
REP1
REP2
IOVANCE
© 2018, Iovance Biotherapeutics

Figure 76


EP11045
NT
OKT3
IFNγ (pg/ml)
10000
8000
6000
4000
2000
0
IL-2
IL-2 + OKT3+Feeders
REP1
NT
OKT3
IFNγ (pg/ml)
150000
100000
50000
0
IL-2
IL-2 + OKT3+Feeders
REP2

Figure 77


OKT3+Feeders
OKT3+Feeders +OX40
% Live Cells
100
80
60
40
20
0
CD4
CD27
CD28
CD123
TEMRA
TSCM
TCM
TEM
CD8
CD27
CD28
CD123
TEMRA
TSCM
TCM
TEM
REP2

Figure 78


OKT3+Feeders
OKT3+Feeders +OX40
% Live Cells
100
80
60
40
20
0
CD25
KLRG1
Lag3
PD1
Tigit
TIM3
CD4
REP2
CD25
KLRG1
Lag3
PD1
Tigit
TIM3
CD8

Figure 79


IFNγ (pg/ml)
40000
30000
20000
10000
0
NT
OKT3
Unstimulated
OKT3
OKT3+Feeders
OKT3+Feeders+ OX40
REP1
REP2

Figure 80


OKT3+Feeders
OKT3+Feeders +OX40
CD107a (Fold Change)
30
20
10
0
CD4+
CD8+
CD4+
CD8+
REP1
REP2

Figure 81


OKT3+Feeders
OKT3+Feeders +OX40
% Live cells
100
80
60
40
20
0
CD4
CD8
CD27+CD28+
CD27+CD28+CD127+
CD27+CD28+CD127+TEMRA
CD27+CD28+CD127+TSCM
CD27+CD28+CD127+TCM
CD27+CD28+CD127+TEM
REP1
REP2

Figure 82


%Live Cells
100
80
60
40
20
0
OKT3+Feeders
OKT3+Feeders +OX40
REP1
REP2
7BA
BTLA
CD25
CD69
CD103
KLRG1
LAG3
PD1
TIGIT
TIM3

Figure 83


EP1028
% of CD3
REP1
REP2
150
100
50
0
CD4+
CD27+ CD28+
TCM TSCM TEMRA TEM
CD8+
CD27+ CD28+
TCM TSCM TEMRA TEM
CD4
CD8
% of CD3
REP1
REP2
100
80
60
40
20
0
2B4 BTLA CD25 CD69 CD103 KLRG1 Lag3 PD1 Tigit TIM3
2B4 BTLA CD25 CD69 CD103 KLRG1 Lag3 PD1 Tigit TIM3
CD4
CD8

Figure 84


EP11028
Breast Cores
NT
OKT3
IFNγ (pg/ml)
40000
30000
20000
10000
0
1 core
2 core
1 core
2 core
REP1
REP2
Breast Cores
Unstimulated
Stimulated
% CD107a
50
40
30
20
10
0
1 core
2 core
1 core
2 core
CD4
REP2
CD8
Breast Cores
Unstimulated
Stimulated
% CD107a
50
40
30
20
10
0
1 core
2 core
1 core
2 core
CD4
REP1
CD8

Figure 85A

Process 2A: about 22 days from Steps A - E	Process GEN 3: about 14-16 days from Steps A - E
STEP A Obtain Patient Tumor Sample (optionally can be frozen before Step B)	**STEP A** Obtain Patient Tumor Sample (optionally can be frozen before Step B)
STEP B First Expansion (physical fragmentation to at least 40 fragments per container grown for about 3 days to 14 days with media comprising IL-2)	**STEP B** Priming First Expansion (physical fragmentation of up to 60 fragments per container grown for about 1 days to 7 days with media comprising IL-2, OKT-3, and antigen-presenting feeder cells)
STEP C First Expansion to Second Expansion Transition (Step B TILs directly move to Step D, optionally on Step B day 11)	**STEP C** Priming First Expansion to Rapid Second Expansion Transition (Step B TILs directly move to Step D on day 7)
STEP D Second Expansion (TILs grown in growth media medium comprising IL-2, OKT-3, and antigen-presenting feeder cells in a closed container)	**STEP D** Rapid Second Expansion (TILs grown in growth media medium comprising IL-2, OKT-3, and 2X antigen-presenting feeder cells; Days 10-11 scale up and add additional IL-2)
STEP E Harvest TILS from Step D (TILs harvested via closed system)	**STEP E** Harvest TILS from Step D
STEP F Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)	**STEP F** Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)

Figure 85B

Small Biopsy GEN 3 Process: about 17-24 days from Steps A - E

STEP A

Obtain Patient Tumor Sample

(optionally can be frozen before Step B)

STEP B

Priming First Expansion

(physical fragmentation of up to 60 fragments per container grown for about 3 days to 12 days with media comprising IL-2, OKT-3, and antigen-presenting feeder cells)

STEP C

Priming First Expansion to Rapid Second Expansion Transition

(Step B TILs directly move to Step D on day 7 to 12)

STEP D

Rapid Second Expansion

(TILs grown in growth media medium comprising IL-2, OKT-3, and 2X antigen-presenting feeder cells; Days 10-12 scale up and add additional IL-2)

STEP E

Harvest TILS from Step D

STEP F

Final Formulation and/or Transfer to Infusion Bag

(optionally cryopreserve)

Figure 85C

Small Biopsy Process: Version 2 (GEN3 based process) **About 17-24 days from Steps A - E**
STEP A Obtain Patient Tumor Sample
STEP B First Expansion 3 days to 12 days (with OKT-3 and antigen-presenting feeder cells at any one of days 1-3)
STEP C First Expansion to Second Expansion Transition No Storage and optionally Closed System
STEP D Second Expansion IL-2, OKT-3, and antigen-presenting feeder cells 3 to 12 days Optionally Closed System
STEP E Harvest TILS from Step D Closed System
STEP F Final Formulation and/or Transfer to Infusion Bag (optionally cryopreserve)

Figure 86


Tumor Dissection
Day 0
Up to 50 fragments in GREX 100MCS + IL-2 (6000 IU/mL) in 1 L CM1 media
Day 0
Whole tumor
1st flask Up to 60 fragments
2nd flask (61-89 fragments)
3rd flask (90-120 fragments)
in GREX 100MCS + IL-2 (6000 IU/mL) in 200mL CM1 media
Day 7
Direct to REP all cells in the same GREX 100MCS in CM2 media +IL-2 (6000 IU/mL)
TVC and % viability
Supernatant collection
Day 11
Direct to REP <200E+06 TIL in GREX 500MCS in CM2 media +IL-2 (3000 IU/mL)
Fragment filtration
Day 9 to 11
Transfer 1L cell solution into 1 GREX 500MCS with 4L CM4 media + IL-2 (3000 IU/mL)
Fragment Filtration
TVC and % viability
Supernatant collection
Day 16
90 % Volume reduce, split up to 5 GREX 500MCS in CM4 media + IL-2 (3000 IU/mL)
Day 16
Harvest
LOVO automated cell washer
Day 22
Harvest
LOVO automated cell washer
TVC and % Viability
INF-γ secretion by ELISA
CDR3 analysis
TIL:Tumor co- culture
Telomere (FlowFISH)
Metabolic substrate analysis
IL-2 levels by ELISA
Immunophenotyping

Figure 87



Tumor ID	L4054		L4055		M1085T	
Process	Gen 2	Gen 3	Gen 2	Gen 3	Gen 2	Gen 3
# Fragments	21	21	24	24	16	16

Figure 88


A
B
C
Memory L4054
Memory L4055
Memory M1085T
Gen 2 Process
Gen 3 Process
% Frequency of viable cells
Phenotype
CD3/CD4
TEMRA
NAIVE
CM
EM
CD3/CD8
CD3 / CD4
CD3 / CD8
Tumor ID
Process
Fragments
L4054
L4055
M1085T
Gen 2
Gen 3
21
21
24
24
16
16

Figure 89


A
L4054
% Gated on CD4+
100
80
60
40
20
0
2Bd
BTLA+
CD25+
CD69+
CD9+
CD11+
CXCP
KLRG1
LAG3+
PD1
TIGIT+
TIM3+
Gen 2 Process
Gen 3 Process
B
L4054
% Gated on CD8+
100
80
60
40
20
0
2Bd
BTLA+
CD25+
CD69+
CD9+
CD11+
CXCP
KLRG1
LAG3+
PD1
TIGIT+
TIM3+
Gen 2 Process
Gen 3 Process

Figure 90


A
L4055
% Gated on CD4+
100
80
60
40
20
0
2B4
BTLA
CD25
CD69
CD9
CD1F
CXCP
KLRG1
LAG3+
PD1
TIGIT
TIM3+
Gen 2 Process
Gen 3 Process
B
L4055
% Gated on CD8+
100
80
60
40
20
0
2B4
BTLA
CD25
CD69
CD9
CD1F
CXCP
KLRG1
LAG3+
PD1
TIGIT
TIM3+
Gen 2 Process
Gen 3 Process

Figure 91


A
L4054
IFNγ (pg/mL)
20000
15000
10000
5000
0
Stimulated
Unstimulated
MC Stimulated
B
L4055
IFNγ (pg/mL)
20000
15000
10000
5000
0
Stimulated
Unstimulated
MC Stimulated
C
M1085T
IFNγ (pg/mL)
20000
15000
10000
5000
0
Stimulated
Unstimulated
MC Stimulated
GEN 2 Process
GEN 3 Process

Figure 92


A
L4054
IL-2 (pg/mL)
40000
30000
20000
10000
0
REP Initiation
Scale Up
Harvest
GEN 2 Process
GEN 3 Process
B
L4055
IL-2 (pg/mL)
30000
20000
10000
0
REP Initiation
Scale Up
Harvest
GEN 2 Process
GEN 3 Process

Figure 93


A
Glucose Gen 2 & Gen 3
Glucose (g/L)
2.5
2.0
1.5
1.0
0.5
0.0
REP initiation
Scale Up
Harvest
Day of REP
Gen 2- L4054
Gen 3- L4054
Gen 2- L4055
Gen 3-L4055
B
Lactate Gen 2 & Gen 3
Lactate (g/L)
2.0
1.5
1.0
0.5
0.0
REP initiation
Scale Up
Harvest
Day of REP
Gen 2- L4054
Gen 3- L4054
Gen 2- L4055
Gen 3-L4055

Figure 94

A


Glutamine Gen 2 & Gen 3
Glutamine (mmol/L)
2.4
2.2
2.0
1.8
1.6
1.4
1.2
REP initiation
Scale Up
Harvest
Day of REP
Gen 2- L4054
Gen 3- L4054
Gen 2- L4055
Gen 3-L4055


Glutamax Gen 2 & Gen 3
Glutamax (mmol/L)
2.6
2.4
2.2
2.0
1.8
1.6
1.4
REP initiation
Scale Up
Harvest
Day of REP
Gen 2- L4054
Gen 3- L4054
Gen 2- L4055
Gen 3-L4055

B

Figure 94

C


Ammonia Gen 2 & Gen 3
Ammonia (mmol/L)
3.0
2.5
2.0
1.5
1.0
REP initiation
Scale Up
Harvest
Day of REP
Gen 2- L4054
Gen 3- L4054
Gen 2- L4055
Gen 3-L4055

Figure 95


Relative Telomere lenght (1301)
10
8
6
4
2
0
L4054
L4055
Tumor ID#
Gen 2 Process
Gen 3 Process

Figure 96


Unique CDR3-Gen2 vs Gen3
Unique CDR3
30000
20000
10000
0
Gen 2-L4054
Gen 3-L4054
Gen 2-L4055
Gen 3- L4055

Figure 97

# uCDR3 (% Overlap)	All uCDR3's		Top 80% uCDR3's	
	Gen 2	Gen 3	Gen 2	Gen 3
Gen 2-L4054	8915	4355 (48.85%)	205	199 (97.07%)
Gen 3-L4054	-	18130	-	223



- 199 sequences are shared between Gen 3 and Gen 2 final product, corresponding to 97.07% of top 80% of unique CDR3 sequences from Gen 2 shared with Gen 3 final product.

Figure 98

# uCDR3 (% Overlap)	All uCDR3's		Top 80% uCDR3's	
	Gen 2	Gen 3	Gen 2	Gen 3
Gen 2-L4055	12996	6599 (50.77%)	1843	1833 (99.45%)
Gen 3-L4055	-	27246	-	2616



- 1833 sequences are shared between Gen 3 and Gen 2 final product, corresponding to 99.45% of top 80% of unique CDR3 sequences from Gen 2 shared with Gen 3 final product.

23 © 2018. Iovance Biotherapeutics

Figure 99


Gen2 vs Gen3
Diversity Index
10
8
6
4
2
0
Gen 2-L4054
Gen 3-L4054
Gen 2-L4055
Gen 3- L4055

Figure 100

Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
188	Total	2.46E+05	2.28E+05	2.31E+05	2.35E+05	4.42E+07
	Live	2.37E+05	2.27E+05	2.26E+05	2.30E+05	4.32E+07
	Dead	9.26E+03	7.03E+02	4.63E+03	4.86E+03	9.14E+05
	% Viability	96.20%	99.70%	98.00%	97.97%	
188	Total	8.26E+04	8.00E+04	7.66E+04	7.97E+04	1.50E+07
	Live	7.60E+04	7.54E+04	6.89E+04	7.34E+04	1.38E+07
	Dead	6.61E+03	4.63E+03	7.73E+03	6.32E+03	1.19E+06
	% Viability	92.00%	94.20%	89.90%	92.03%	
200	Total	2.44E+04	4.20E+04	1.05E+04	2.56E+04	5.13E+06
	Live	1.74E+04	2.80E+04	7.03E+03	1.75E+04	3.50E+06
	Dead	7.03E+03	1.40E+04	3.48E+03	8.17E+03	1.63E+06
	% Viability	71.20%	66.70%	66.90%	68.27%	

Figure 101

ID	Volume (mL)		Count 1 (cells/mL)	Count 3 (cells/mL)	Count 2 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 3 #L4054 First Round	1000	Total	9.11E+05	1.01E+06	1.08E+06	1.00E+06	1.00E+09
		Live	8.60E+05	9.34E+05	1.01E+06	9.35E+05	9.35E+08
		Dead	5.09E+04	7.49E+04	6.74E+04	6.44E+04	6.44E+07
		% Viability	94.40%	92.60%	93.70%	93.57%	
Gen 3 #L4055 Second Round	1000	Total	8.56E+05	9.04E+05	8.81E+05	8.80E+05	8.80E+08
		Live	8.24E+05	8.67E+05	8.42E+05	8.44E+05	8.44E+08
		Dead	3.17E+04	3.70E+04	3.87E+04	3.58E+04	3.58E+07
		% Viability	96.30%	95.90%	95.60%	95.93%	
Gen 3 #M1085T	200	Total	2.32E+06	2.25E+06		2.29E+06	4.57E+08
		Live	1.68E+06	1.57E+06		1.63E+06	3.25E+08
		Dead	6.41E+05	6.79E+05		6.60E+05	1.32E+08
		% Viability	72.40%	69.90%		71.15%	
Gen 2 #L4054 First Round	142	Total	1.08E+06	9.84E+05	1.00E+06	1.02E+06	1.45E+08
		Live	1.06E+06	9.71E+05	9.79E+05	1.00E+06	1.42E+08
		Dead	2.05E+04	1.26E+04	2.12E+04	1.81E+04	2.57E+06
		% Viability	98.10%	98.70%	97.90%	98.23%	
Gen 2 #L4055 Second Round	96	Total	2.93E+05	3.05E+05	2.64E+05	2.87E+05	2.76E+07
		Live	2.88E+05	2.96E+05	2.55E+05	2.80E+05	2.68E+07
		Dead	4.72E+03	9.14E+03	8.26E+03	7.37E+03	7.08E+05
		% Viability	98.40%	97.00%	96.90%	97.43%	
Gen 2 #M1085T	200	Total	9.10E+04	5.60E+04		7.35E+04	1.47E+07
		Live	8.05E+04	4.21E+04		6.13E+04	1.23E+07
		Dead	1.04E+04	1.39E+04		1.22E+04	2.43E+06
		% Viability	88.50%	75.20%		81.85%	

Figure 102

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 3 #L4054 First Round -pre LOVO	1243.1	Total	1.40E+07	1.46E+07		1.43E+07	1.78E+10
		Live	1.32E+07	1.35E+07		1.34E+07	1.66E+10
		Dead	8.47E+05	1.12E+06		9.84E+05	1.22E+09
		% Viability	94.00%	92.40%		93.20%	
Gen 3 #L4054 First Round -post LOVO	330	Total	6.35E+07	6.07E+07		6.21E+07	2.05E+10
		Live	5.72E+07	5.44E+07		5.58E+07	1.84E+10
		Dead	6.35E+06	6.27E+06		6.31E+06	2.08E+09
		% Viability	90.00%	89.70%		89.85%	

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 3 #L4055 Second Round -pre LOVO	1032.6	Total	9.49E+06	1.05E+07		1.00E+07	1.03E+10
		Live	8.90E+06	9.79E+06		9.35E+06	9.65E+09
		Dead	5.85E+05	6.75E+05		6.30E+05	6.51E+08
		% Viability	93.80%	93.60%		93.70%	
Gen 3 #L4055 Second Round -post LOVO	330	Total	3.31E+07	2.82E+07	3.18E+07	3.07E+07	1.01E+10
		Live	2.85E+07	2.46E+07	2.77E+07	2.66E+07	8.76E+09
		Dead	4.56E+06	3.61E+06	4.12E+06	4.09E+06	1.35E+09
		% Viability	86.20%	87.20%	87.10%	86.70%	

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 3 #M1085T pre-LOVO	5000	Total	2.26E+06	2.64E+06		2.45E+06	1.23E+10
		Live	1.97E+06	2.31E+06		2.14E+06	1.07E+10
		Dead	2.85E+05	3.30E+05		3.08E+05	1.54E+09
		% Viability	87.40%	87.50%		87.45%	
Gen 3 #M1085T - post LOVO pre CS10 addition	150	Total	6.27E+07	5.44E+07		5.86E+07	8.78E+09
		Live	5.50E+07	4.74E+07		5.12E+07	7.68E+09
		Dead	7.70E+06	6.96E+06		7.33E+06	1.10E+09
		% Viability	87.70%	87.20%		87.45%	

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 2 #L4054 First Round	913	Total	3.53E+06	4.30E+06		3.92E+06	3.57E+09
		Live	3.35E+06	4.00E+06		3.68E+06	3.36E+09
		Dead	1.75E+05	3.03E+05		2.39E+05	2.18E+08
		% Viability	95.00%	93.00%		94.00%	
Gen 2 #L4055 Second Round	292	Total	1.29E+07	1.36E+07		1.33E+07	3.87E+09
		Live	1.16E+07	1.23E+07		1.20E+07	3.49E+09
		Dead	1.27E+06	1.23E+06		1.25E+06	3.65E+08
		% Viability	90.10%	90.90%		90.50%	
Gen 2 #M1085T	369	Total	6.75E+06	6.98E+06		6.87E+06	2.53E+09
		Live	5.22E+06	5.58E+06		5.40E+06	1.99E+09
		Dead	1.54E+06	1.40E+06		1.47E+06	5.42E+08
		% Viability	77.20%	79.90%		78.55%	

Figure 103

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 2 #L4054 First Round -pre LOVO	1385.7	Total	3.55E+07	4.12E+07	4.03E+07	3.90E+07	5.40E+10
		Live	3.26E+07	3.73E+07	3.67E+07	3.55E+07	4.92E+10
		Dead	2.93E+06	3.87E+06	3.59E+06	3.46E+06	4.80E+09
		% Viability	91.70%	90.60%	91.10%	91.13%	
Gen 2 #L4054 First Round -post LOVO	330	Total	1.70E+08	1.79E+08	1.68E+08	1.72E+08	5.69E+10
		Live	1.49E+08	1.58E+08	1.48E+08	1.52E+08	5.01E+10
		Dead	2.16E+07	2.04E+07	2.00E+07	2.07E+07	6.82E+09
		% Viability	87.30%	88.60%	88.10%	87.95%	

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 2 #L4055 First Round -pre LOVO	1968.2	Total	3.15E+07	2.51E+07	2.97E+07	2.88E+07	5.66E+10
		Live	2.89E+07	2.25E+07	2.72E+07	2.62E+07	5.16E+10
		Dead	2.57E+06	2.61E+06	2.52E+06	2.57E+06	5.05E+09
		% Viability	91.80%	89.60%	91.50%	90.97%	
Gen 2 #L4055 First Round -post LOVO	330	Total	2.33E+08	1.89E+08	1.53E+08	1.92E+08	6.33E+10
		Live	2.03E+08	1.66E+08	1.33E+08	1.67E+08	5.52E+10
		Dead	3.00E+07	2.24E+07	1.94E+07	2.39E+07	7.90E+09
		% Viability	87.10%	88.10%	87.30%	87.50%	

ID	Volume (mL)		Count 1 (cells/mL)	Count 2 (cells/mL)	Count 3 (cells/mL)	Average (cells/mL)	Total Cells Average*Volume
Gen 2 #M1085T-pre LOVO	N/A	Total				#DIV/0!	#DIV/0!
		Live				#DIV/0!	#DIV/0!
		Dead				#DIV/0!	#DIV/0!
		% Viability				#DIV/0!	
Gen 2 #M1085T - post LOVO pre CS10 addition	150	Total	8.51E+07	9.05E+07		8.78E+07	1.32E+10
		Live	7.33E+07	7.79E+07		7.56E+07	1.13E+10
		Dead	1.18E+07	1.26E+07		1.22E+07	1.83E+09
		% Viability	86.10%	86.10%		86.10%	

For L4054 Gen 2, post LOVO count was extrapolated to 4 flasks, because was the total number of the study. 1 flask was contaminated, and the extrapolation was done for total = 6.67E+10

Figure 104

	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD8/	S1/S2/All/Live/CD14-/TCRab/CD4/Q1: CCR7-PE-A-, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q2: CCR7-PE-A+, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q3: CCR7-PE-A+, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q4: CCR7-PE-A-, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q5: CD62L-BV421-A-, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q6: CD62L-BV421-A+, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q7: CD62L-BV421-A+, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD4/Q8: CD62L-BV421-A-, CD45RA-AF700-A- | Freq. of Parent (%)
	Freq. of Parent (%)	CD14+ Freq. of Parent (%)	CD33-/% Freq. of null (%)	CD19- (%)	CD4- Freq. of null (%)	/TCRab- | Freq. of null (%)	/TCRab/CD Freq. of null (%)	CD27- | Freq. of null (%)	CD28- Freq. of null (%)	CD56- Freq. of null (%)	CD57- | Freq. of null (%)								
												TEMRA	NAÏVE	CM	EM	TEMRA	NAÏVE	CM	EM
A1 gen2 1st.fcs	97.2	1.77E-03	0.06	0	96.9	85.8	3.35	89.1	0.06	5.24	0.25	0.073	3.37	96.3	0.056	0.032	21.1	78.8
A2 gen2 2nd.fcs	73.5	0	0.12	0	99.7	93.1	1.94	92.6	0.14	6.18	0.16	0.047	2.39	97.4	0.036	0.014	8.3	91.6
A3 gen3 1st.fcs	90.4	0	0.07	1.22E-03	86.8	49.1	4.9	48.9	0.18	6.02	0.2	0.059	2.29	97.5	0.059	0.049	30.7	69.2
A4 gen3 2nd.fcs	72.1	2.38E-04	0.44	0.018	99.3	64.4	6.27	68	0.26	4.3	0.25	0.18	7.12	92.5	0.12	0.072	20.1	79.7
A5 PBMC.fcs	93.7	6.07	11.6	6.12	78.9	41.1	45.6	49.4	0.04	5.01	0.91	36.1	16.2	46.9	11.4	23.1	16.4	49.1
I area was include to make the graph																		
TCRab+ are = CD3+																		

S1/S2/All/Live/CD14-/TCRab/CD8/ Freq. of null (%)	S1/S2/All/Live/CD14-/TCRab/CD8/CD27- | Freq. of null (%)	S1/S2/All/Live/CD14-/TCRab/CD8/CD28- Freq. of null (%)	S1/S2/All/Live/CD14-/TCRab/CD8/CD56- | Freq. of null (%)	S1/S2/All/Live/CD14-/TCRab/CD8/CD57- | Freq. of null (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q1: CCR7-PE-A-, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q2: CCR7-PE-A+, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q3: CCR7-PE-A+, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q4: CCR7-PE-A-, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q5: CD62L-BV421-A-, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q6: CD62L-BV421-A+, CD45RA-AF700-A+ | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q7: CD62L-BV421-A+, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRab/CD8/Q8: CD62L-BV421-A-, CD45RA-AF700-A- | Freq. of Parent (%)	S1/S2/All/Live/CD14-/TCRgd | Freq. of null (%)
					TEMRA	NAÏVE	CM	EM	TEMRA	NAÏVE	CM	EM	
37.8	4.25	37.3	0.41	2.11	0.028	0	4.24	95.7	0.033	9.37E-03	14.1	85.9	0
18.1	1.08	15.8	0.67	0.17	0.042	0.014	4.76	95.2	0.071	0.014	5.24	94.7	0
39.3	4.33	40.8	0.77	1.34	0.056	6.97E-03	3.11	96.8	0.077	0.017	25.5	74.4	0
34.6	2.46	34	1.57	0.5	0.18	0.13	10.2	89.5	0.29	0.044	10.6	89.1	0
29.6	25.8	37.4	0.29	2.81	8.05	40.9	3.02	48	22.3	27.1	6.23	44.3	0

Figure 105

	CD3/CD4 CD38	HLA-DR	CCR7- CD45RA+	CCR7+CD45RA+	CCR7+ CD45RA-	CCR7-CD45RA-
GEN 3-Day 16-Harvest	42.4	95.2	4.96	0.7	1.22	93.1
GEN 2-D22-Harvest	50.4	98	0.61	0.13	0.3	99

CD3	CD3/CD8 CD38	HLA-DR	CCR7-CD45RA+	CCR7+CD45RA+	CCR7+CD45RA-	CCR7-CD45RA-
CD8						
77.6	83.2	90.3	20.5	0.013	0.06	79.5
54.3	88	83.2	4.24	0.026	1.05	94.7

Figure 106

			CD4												
	CD3+	CD4+	2B4+	BTLA+	CD103+	CD25+	CD69+	CD95+	CXCR3+	KLRG1+	LAG3+	PD1+	TIGIT+	TIM3+	
Gen 2-L4054			0.61	99.50	1.05	14.80	43.20	99.40	52.90	2.65	5.36	81.20	88.40	78.60	
Gen 3-L4054			1.11	99.50	2.28	21.70	45.40	99.70	56.90	8.06	5.07	76.70	74.10	79.30	
Gen 2-L4055			1.61	98.60	1.56	23.10	51.10	99.80	63.30	3.86	15.30	77.60	86.00	79.50	
Gen 3- L4055			3.57	99.30	2.61	54.40	57.00	99.90	48.00	8.02	10.80	61.40	47.50	84.60	
			CD8												
Gen 2-L4054		CD8+	2B4+	BTLA4+	CD103+	CD25+	CD95+	CD69+	CXCR3+	KLRG1+	LAG3+	PD1+	TIGIT+	TIM3+	
Gen 3-L4054			35.20	99.50	74.90	11.80	99.80	58.30	99.10	8.03	33.10	80.60	99.10	88.30	
Gen 2-L4055			45.50	99.60	62.40	12.10	99.90	54.00	99.30	13.40	22.10	64.90	98.50	87.50	
Gen 3- L4055			40.60	98.20	19.50	32.00	99.70	75.50	96.80	22.80	27.40	28.10	92.30	75.10	
			62.80	99.20	29.20	45.50	99.70	58.80	99.60	13.10	16.60	34.10	71.70	63.60	

Figure 107



	1	2	3	4	5	6	7	8	9	10	11	12	
A	BTD1	BTD2	BTD3	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	MC-Stimulated xG2	MC-Stimulated xG2	MC-Stimulated xG2	Well ID
	1000	1000	1000	3	3	3	3	3	3	3	3	3	Conc/Dil
B	BTD4	BTD5	BTD6	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	MC-Stimulated xG2	MC-Stimulated xG2	MC-Stimulated xG2	Well ID
	500	500	500	9	9	9	9	9	9	9	9	9	Conc/Dil
C	BTD7	BTD8	BTD9	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	MC-Unstimulated xG2	MC-Unstimulated xG2	MC-Unstimulated xG2	Well ID
	250	250	250	27	27	27	27	27	27	3	3	3	Conc/Dil
D	BTD10	BTD11	BTD12	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 stimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	TIL Gen 2 Unstimulated	MC-Unstimulated xG2	MC-Unstimulated xG2	MC-Unstimulated xG2	Well ID
	125	125	125	81	81	81	81	81	81	9	9	9	Conc/Dil
E	BTD13	BTD14	BTD15	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	MC-Stimulated xG3	MC-Stimulated xG3	MC-Stimulated xG3	Well ID
	62.5	62.5	62.5	3	3	3	3	3	3	3	3	3	Conc/Dil
F	BTD16	BTD17	BTD18	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	MC-Stimulated xG3	MC-Stimulated xG3	MC-Stimulated xG3	Well ID
	31.3	31.3	31.3	9	9	9	9	9	9	9	9	9	Conc/Dil
G	BTD19	BTD20	BTD21	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	MC-Unstimulated xG3	MC-Unstimulated xG3	MC-Unstimulated xG3	Well ID
	15.6	15.6	15.6	27	27	27	27	27	27	3	3	3	Conc/Dil
H	BTD22	BTD23	BTD24	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-stimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	TIL Gen 3-Unstimulated	MC-Unstimulated xG3	MC-Unstimulated xG3	MC-Unstimulated xG3	Well ID
	8	8	8	81	81	81	81	81	81	9	9	9	Conc/Dil

22.6
22.6

	1	2	3	4	5	6	7	8	9	10	11	12	
A	0.956	0.962	0.966	2.797	2.104	3.118	0.117	0.116	0.117	0.105	0.108	0.102	450
	0.05	0.05	0.05	0.05	0.048	0.051	0.041	0.043	0.041	0.041	0.043	0.041	570
	0.906	0.912	0.916	2.747	2.056	3.067	0.076	0.074	0.076	0.064	0.065	0.06	Delta
	994.773	1002.343	1004.079	963.405	694.982	1018.375	1.825	1.421	1.952	<0.000	<0.000	<0.000	[Concentration]
				2890.216	2084.946	3055.126	5.474	4.262	5.855	<0.000	<0.000	<0.000	x-dil factor
B	0.626	0.654	0.65	1.181	0.9	1.315	0.107	0.107	0.108	0.106	0.105	0.102	450
	0.046	0.046	0.046	0.045	0.047	0.045	0.042	0.041	0.042	0.043	0.042	0.042	570
	1.613	1.608	1.604	1.136	0.853	1.27	0.066	0.066	0.067	0.062	0.063	0.061	Delta
	499.866	497.826	496.528	329.765	235.273	341.438	<0.000	<0.000	<0.000	<0.000	<0.000	<0.000	[Concentration]
				2967.887	2117.46	3072.942	<0.000	<0.000	<0.000	<0.000	<0.000	<0.000	x-dil factor
C	0.952	0.966	0.94	0.494	0.391	0.505	0.106	0.107	0.107	0.107	0.105	0.103	450
	0.044	0.048	0.043	0.042	0.042	0.043	0.041	0.042	0.042	0.043	0.042	0.042	570
	0.908	0.918	0.896	0.452	0.349	0.462	0.065	0.065	0.065	0.063	0.063	0.062	Delta
	253.244	256.8	249.399	109.334	78.83	112.411	<0.000	<0.000	<0.000	<0.000	<0.000	<0.000	[Concentration]
				2952.028	2128.412	3035.11	<0.000	<0.000	<0.000	<0.000	<0.000	<0.000	x-dil factor
D	0.549	0.541	0.542	0.258	0.218	0.259	0.102	0.101	0.103	0.111	0.102	0.101	450
	0.044	0.044	0.043	0.042	0.041	0.047	0.041	0.042	0.042	0.05	0.042	0.043	570
	0.505	0.498	0.499	0.216	0.176	0.212	0.06	0.059	0.061	0.061	0.06	0.057	Delta
	125.458	123.174	123.508	40.262	29.15	39.101	<0.000	<0.000	<0.000	<0.000	<0.000	<0.000	[Concentration]
				3261.251	2361.138	3167.212	<0.000	<0.000	<0.000	<0.000	<0.000	<0.000	x-dil factor
E	0.334	0.332	0.332	OVRFLW	OVRFLW	OVRFLW	0.114	0.11	0.113	0.103	0.102	0.104	450
	0.043	0.042	0.042	#####	#####	#####	0.041	0.041	0.042	0.042	0.042	0.043	570
	0.291	0.29	0.29	#####	#####	#####	0.073	0.069	0.072	0.062	0.061	0.061	Delta
	61.826	61.537	61.594	#####	#####	#####	1.145	0.285	0.822	<0.000	<0.000	<0.000	[Concentration]
				#####	#####	#####	3.435	0.854	2.465	<0.000	<0.000	<0.000	x-dil factor
F	0.222	0.222	0.224	0.09	0.085	0.093	0.114	0.104	0.107	0.101	0.103	0.101	450
	0.042	0.043	0.042	0.043	0.039	0.057	0.041	0.041	0.042	0.042	0.041	0.042	570
	0.18	0.179	0.181	0.047	0.046	0.036	0.073	0.063	0.066	0.059	0.062	0.059	Delta
	30.212	29.82	30.519	1474.838	>1500.000	1482.263	1.145	<0.000	<0.000	<0.000	<0.000	<0.000	[Concentration]
				13273.55	>13500.000	13340.37	10.304	<0.000	<0.000	<0.000	<0.000	<0.000	x-dil factor
G	0.174	0.17	0.172	0.09	0.085	0.093	0.114	0.109	0.114	0.106	0.107	0.104	450
	0.041	0.041	0.041	0.048	0.047	0.049	0.042	0.042	0.041	0.042	0.041	0.042	570
	0.132	0.128	0.131	0.042	0.038	0.044	0.073	0.067	0.073	0.063	0.065	0.062	Delta
	16.911	15.874	16.447	627.703	628.253	635.61	1.12	<0.000	1.17	<0.000	<0.000	<0.000	[Concentration]
				16947.99	16962.84	17161.47	30.239	<0.000	31.588	<0.000	<0.000	<0.000	x-dil factor
H	0.117	0.111	0.114	0.84	0.867	0.857	0.11	0.11	0.115	0.109	0.11	0.113	450
	0.045	0.043	0.044	0.044	0.044	0.044	0.041	0.042	0.043	0.042	0.043	0.042	570
	0.072	0.068	0.07	0.797	0.822	0.813	0.069	0.068	0.073	0.067	0.067	0.071	Delta
	0.822	0.073	0.405	216.931	225.269	222.13	0.119	0.05	1.145	<0.000	<0.000	0.748	[Concentration]
				17571.43	18246.77	17992.49	9.644	4.039	92.74	<0.000	<0.000	60.557	x-dil factor

				Average	Average
Gen 2-Stimulated+TIL 1.3	2890.216	2084.946	3055.126	2676.76267	2757.81
Gen 2-Stimulated+TIL 1.9	2967.887	2117.46	3072.942	2719.42967	
Gen 2-Stimulated+TIL 1.27	2952.028	2128.412	3035.11	2705.18333	
Gen 2-Stimulated+TIL 1.81	3261.251	2361.138	3167.212	2929.867	
Gen 3-Stimulated+TIL 1.3	#####	#####	#####	#DIV/0!	#DIV/0!
Gen 3-Stimulated+TIL 1.9	13273.546	>13500.00	13340.37	13306.9575	16089.32
Gen 3-Stimulated+TIL 1.27	16947.994	16962.84	17161.47	17024.1	
Gen 3-Stimulated+TIL 1.81	17571.427	18246.77	17992.49	17936.8973	
Gen 2-Unstimulated 1.3	5.474	4.262	5.855	5.197	5.20
Gen 2-Unstimulated 1.9	<0.000	<0.000	<0.000	#DIV/0!	
Gen 2-Unstimulated 1.27	<0.000	<0.000	<0.000	#DIV/0!	
Gen 2-Unstimulated 1.81	<0.000	<0.000	<0.000	#DIV/0!	
Gen 3-Unstimulated 1.3	3.435	0.854	2.465	2.25133333	19.74
Gen 3-Unstimulated 1.9	10.304	<0.000	<0.000	10.304	
Gen 3-Unstimulated 1.27	30.239	<0.000	31.588	30.9135	
Gen 3-Unstimulated 1.81	9.644	4.039	92.74	35.4743333	
Gen 2-Media Control Stimulated 1.3	<0.000	<0.000	<0.000	#DIV/0!	0.00
Gen 2-Media Control Stimulated 1.9	<0.000	<0.000	<0.000	#DIV/0!	
Gen 3-Media Control Stimulated 1.3	<0.000	<0.000	<0.000	#DIV/0!	0.00
Gen 3-Media Control Stimulated 1.9	<0.000	<0.000	<0.000	#DIV/0!	
Gen 2-Media Control Unstimulated 1.3	<0.000	<0.000	<0.000	#DIV/0!	0.00
Gen 2-Media Control Unstimulated 1.9	<0.000	<0.000	<0.000	#DIV/0!	
Gen 3-Media Control Unstimulated 1.3	<0.000	<0.000	<0.000	0	30.28
Gen 3-Media Control Unstimulated 1.9	<0.000	<0.000	60.557	60.557	

Figure 108



Figure 109



Figure 110



	1	2	3	4	5	6	7	8	9	10	11	12
A	5'Dt1 3'Dt1	5'Dt1 3'Dt2	5'Dt1 3'Dt3	Gen 3-D7 1ST RUN	Gen 3-D7 1ST RUN	Gen 3-D7 1ST RUN	Gen 3-D7 2ND RUN	Gen 3-D7 2ND RUN	Gen 3-D7 2ND RUN			
B	5'Dt2 3'Dt1	5'Dt2 3'Dt2	5'Dt2 3'Dt3	Gen 3-D11 1ST RUN	Gen 3-D11 1ST RUN	Gen 3-D11 1ST RUN	Gen 3-D11 2ND RUN	Gen 3-D11 2ND RUN	Gen 3-D11 2ND RUN			
C	5'Dt3 3'Dt1	5'Dt3 3'Dt2	5'Dt3 3'Dt3	Gen 3-D16 1ST RUN	Gen 3-D16 1ST RUN	Gen 3-D16 1ST RUN	Gen 3-D16 2ND RUN	Gen 3-D16 2ND RUN	Gen 3-D16 2ND RUN			
D	5'Dt4 3'Dt1	5'Dt4 3'Dt2	5'Dt4 3'Dt3	Gen 2-D11 1ST RUN	Gen 2-D11 1ST RUN	Gen 2-D11 1ST RUN	Gen 2-D11 2ND RUN	Gen 2-D11 2ND RUN	Gen 2-D11 2ND RUN			
E	5'Dt5 3'Dt1	5'Dt5 3'Dt2	5'Dt5 3'Dt3	Gen 2-D16 1ST RUN	Gen 2-D16 1ST RUN	Gen 2-D16 1ST RUN	Gen 2-D16 2ND RUN	Gen 2-D16 2ND RUN	Gen 2-D16 2ND RUN			
F	5'Dt6 3'Dt1	5'Dt6 3'Dt2	5'Dt6 3'Dt3	Gen 2-D22 1ST RUN	Gen 2-D22 1ST RUN	Gen 2-D22 1ST RUN	Gen 2-D22 2ND RUN	Gen 2-D22 2ND RUN	Gen 2-D22 2ND RUN			
G	5'Dt7 3'Dt1	5'Dt7 3'Dt2	5'Dt7 3'Dt3	Empty	Empty	Empty	Empty	Empty	Empty			
H	5'Dt8 3'Dt1	5'Dt8 3'Dt2	5'Dt8 3'Dt3	Empty	Empty	Empty	Empty	Empty	Empty			

	1	2	3	4	5	6	7	8	9	10	11	12
A	0.216 0.043 0.173 1970.541	0.219 0.043 0.176 2046.905	0.221 0.043 0.178 1987.758	0.218 0.043 0.175 124.53	0.22 0.042 0.177 126.082	0.215 0.043 0.172 121.662	0.187 0.044 0.143 99.302	0.187 0.043 0.145 100.904	0.184 0.051 0.133 92.083			
B	1.045 0.047 1.048 877.734	1.251 0.046 1.234 1051.728	1.276 0.047 1.229 1047.027	0.415 0.043 0.372 282.4	0.412 0.043 0.369 280.328	0.435 0.043 0.393 299.854	0.394 0.043 0.351 264.954	0.392 0.044 0.348 262.893	0.356 0.044 0.312 233.453			
C	0.682 0.045 0.636 510.008	0.681 0.044 0.637 510.619	0.68 0.044 0.637 508.261	0.376 0.043 0.333 252.774	0.372 0.043 0.33 247.768	0.376 0.043 0.333 250.557	0.356 0.043 0.313 233.943	0.338 0.043 0.295 219.211	0.33 0.043 0.287 212.888			
D	0.37 0.043 0.328 246.046	0.38 0.045 0.335 252.035	0.382 0.043 0.338 254.91	0.162 0.046 0.116 79.199	0.156 0.042 0.114 77.702	0.177 0.042 0.135 93.523	0.155 0.043 0.112 76.431	0.15 0.044 0.106 71.588	0.149 0.043 0.106 71.737			
E	0.216 0.043 0.173 122.514	0.219 0.043 0.176 124.995	0.22 0.043 0.176 123.909	0.235 0.044 0.191 136.754	0.227 0.043 0.185 131.526	0.228 0.043 0.185 131.682	0.253 0.046 0.205 147.807	0.231 0.045 0.186 132.695	0.222 0.045 0.177 125.539			
F	0.137 0.048 0.089 59.482	0.132 0.043 0.089 58.969	0.133 0.044 0.089 59.116	0.21 0.043 0.166 117.564	0.227 0.044 0.183 130.125	0.216 0.044 0.172 121.895	0.238 0.043 0.195 139.491	0.233 0.043 0.19 135.738	0.253 0.045 0.207 149.459			
G	0.088 0.042 0.046 28.375	0.088 0.043 0.046 28.237	0.089 0.043 0.046 28.514	0.2 0.05 0.15 2014.9	0.2 0.05 0.15 2081.651	0.2 0.05 0.15 1970.439	0.2 0.05 0.15 1973.799	0.2 0.05 0.15 1920.845	0.2 0.05 0.15 1986.127			
H	0.05 0.043 0.007 3.1	0.048 0.042 0.006 2.399	0.048 0.041 0.007 2.573	0.2 0.05 0.15 2023.073	0.2 0.05 0.15 2016.943	0.2 0.05 0.15 1960.16	0.2 0.05 0.15 1986.84	0.2 0.05 0.15 2002.036	0.2 0.05 0.15 1988.165			

Dilution Factor=100

	Reading A	Reading B	Reading C	Reading A *100	Reading B*100	Reading C*100	Average
Gen 3 Rep initiation 1st run	124.530	126.082	121.662	12453.0	12608.2	12166.2	12409.1
Gen 3 Scale up 1st run	282.400	280.328	299.854	28240.0	28032.8	29985.4	28752.7
Gen 3 Harvest 1st run	252.774	247.768	250.557	25277.4	24776.8	25055.7	25036.6
Gen 2 Rep initiation 1st run	79.199	77.702	93.523	7919.9	7770.2	9352.3	8347.5
Gen 2 Scale up 1st run	136.754	131.526	131.682	13675.4	13152.6	13168.2	13332.1
Gen 2 Harvest 1st run	117.564	130.125	121.895	11756.4	13012.5	12189.5	12319.5
Gen 3 Rep initiation 2nd run	99.302	100.904	92.083	9930.2	10090.4	9208.3	9743.0
Gen 3 Scale up 2nd run	264.954	262.893	233.453	26495.4	26289.3	23345.3	25376.7
Gen 3 Harvest 2nd run	233.943	219.211	212.888	23394.3	21921.1	21288.8	22201.4
Gen 2 Rep initiation 2nd run	76.431	71.588	71.737	7643.1	7158.8	7173.7	7325.2
Gen 2 Scale up 2nd run	147.807	132.695	125.539	14780.7	13269.5	12553.9	13534.7
Gen 2 Harvest 2nd run	139.491	135.738	149.459	13949.1	13573.8	14945.9	14156.3

Figure 111

Sample number #	Description	Glucose (g/L)	Lactate (g/L)	Ammonia (mmol/L)	Glutamine (mmol/L)	Glutamax (mmol/L)	Glutamax-Glutamine (mmol/L)
1	Gen 2-Rep initiation First Round #L4054	1.78	0.14	2.16	1.56	1.71	0.16
2	Gen 2-Scale Up First Round #L4054	1.36	0.89	1.40	1.76	1.90	0.15
3	Gen 2-Harvest First Round #L4054	0.83	1.68	2.46	1.69	1.85	0.17
4	Gen 3-REP initiation First Round #L4054	1.68	0.29	1.76	2.11	2.29	0.18
5	Gen 3-Scale Up First Round #L4054	1.28	0.94	1.64	1.70	1.87	0.16
6	Gen 3-Harvest First Round #L4054	1.07	1.53	2.23	1.39	1.54	0.14
7	Gen 2-Rep initiation Second Round #L4055	1.91	0.08	2.29	1.60	1.74	0.14
8	Gen 2-Scale up Second Round #L4055	1.61	0.77	1.40	1.78	1.93	0.15
9	Gen 2-Harvest Second Round #L4055	0.85	1.86	2.46	1.39	1.55	0.16
10	Gen 3-REP initiation Second Round #L4055	1.77	0.18	1.74	2.06	2.23	0.17
11	Gen 3-Scale up Second Round #L4055	1.60	0.72	1.54	1.68	1.84	0.15
12	Gen 3-Harvest Second Round #L4055	1.44	1.32	2.07	1.79	1.95	0.16

Figure 112

	$FIL	S/L/1301 | Geometr	S/L/Pt | Geometric Mean (FL03-A)
A1 gen2L4054 ne	A1 gen2L4054 neg.fcs	34.5	3.24
A2 gen2L4054 ne	A2 gen2L4054 neg.fcs	34	3.19
A3 gen2L4054 po	A3 gen2L4054 pos.fcs	1834	70.6
A4 gen2L4054 po	A4 gen2L4054 pos.fcs	1872	70.3
A5 gen2L4055 ne	A5 gen2L4055 neg.fcs	31.9	4.23
A6 gen2L4055 ne	A6 gen2L4055 neg.fcs	32.4	4.87
A7 gen2L4055 po	A7 gen2L4055 pos.fcs	1941	88.7
A8 gen2L4055 po	A8 gen2L4055 pos.fcs	1933	91.1
B1 gen3L4054 ne	B1 gen3L4054 neg.fcs	31.2	3.45
B2 gen3L4054 ne	B2 gen3L4054 neg.fcs	31.9	3.39
B3 gen3L4054 po	B3 gen3L4054 pos.fcs	2016	61.3
B4 gen3L4054 po	B4 gen3L4054 pos.fcs	2014	76.2
B5 gen3L4055 ne	B5 gen3L4055 neg.fcs	33.7	6.76
B6 gen3L4055 ne	B6 gen3L4055 neg.fcs	35.4	6.7
B7 gen3L4055 po	B7 gen3L4055 pos.fcs	2158	99.4
B8 gen3L4055 po	B8 gen3L4055 pos.fcs	1938	99.5
Mean	-	998	43.3
SD	-	999	41.3

Figure 113

Study	Sample	Sample id	Species	Chain	Reads	CDR3	Unique CDR3	D50
20180511_Study1_FCA	Gen2-L4054	59990	TRB	h	1181732	1181732	8915	0
20180511_Study1_FCA	Gen3-L4054	59991	TRB	h	1145697	1145697	18130	0
20180511_Study1_FCA	Gen2-L4055	59987	TRB	h	1166465	1166465	12996	0.1
20180511_Study1_FCA	Gen3-L4055	59982	TRB	h	1059985	1059985	27246	0.9

Figure 114


Gen 2 Process
Gen 2.1 Process
Gen 3.1 Process
Day 0:Pre-REP
Day 11: REP-Initiation
Day 16: TIL-Scale Up
Day 22: Harvest
Harvest-LOVO automated cell washer
Harvest-LOVO automated cell washer
Day 0:Pre-REP
Day 7: REP-initiation
Day 10 to 11: TIL-Scale Up
Day 14-16: Harvest
Harvest-LOVO automated cell washer
ADVANCE

Figure 115

STEP	Gen 2	Gen 2.1	Gen 3.0
Pre REP- day 0	≤ 50 fragments/ 1 G-Rex 100MCS - 11 days	≤ 180 fragments/ 3 G-Rex Pre-formulated CM1 warmed media 100MCS - 11 days	*Fresh or Frozen Tumor* Whole tumor with ≤ 30 fragments up to 60 fragments per 1 G-Rex 100MCS (up to 4 G-Rex), preformulated warmed media - 7 days, Pre REP: Feeders 2.5 E8 cells + OKT-3 (30ng/mL)
REP Initiation	Direct to REP- Day 11- <200 E6 TIL 1 G-Rex 500MCS	Direct to REP- Day 11- <200 E6 TIL Pre-formulated CM2 warmed media in one G-Rex 500MCS	Direct to REP- Day 7-all cells TIL- same G-Rex 100MCS (100MCS up to 4 GREX), Standard media or Defined Media (Serum free), Addition Feeders 5 E8 cells +OKT-3 (30ng/mL)
TIL propagation or Scale up	1 to 5 G-REX 500MCS Split day 16	2 to 5 G-REX 500MCS Pre-formulated CM4 warmed media Split day 16	From G-REX 100MCS transfer TIL suspension to G-REX 500MCS , up to 4 GREX 500 MCS- Standard media or Defined Media (Serum Free) Scale up on day 10 or 11
Harvest	Harvest day 22, LOVO-automated cell washer	Harvest day 22, LOVO-automated cell washer (5 wash cycle)	Harvest day 14 or 16 LOVO- automated cell washer (5 wash cycle)
Final formulation	Cryopreserved Product 300IU/ml IL2- CS10 in LN_2 multiple aliquots	Cryopreserved Product 300IU/ml IL2- CS10 in LN_2 multiple aliquots	Cryopreserved product 3001U/ml IL-2-CS 10 in LN_2 multiple aliquots
Process time	22 days	22 days	16 days

Figure 116

Process Day	Conditions	Gen 3.1
Day 0- pre REP initiation	Media CM1	500 mL
	IL-2 (6000 IU/mL)	+
	OKT-3 (30ng/mL)	+
	Feeders (250 E+06)	+

Process Day	Conditions	Gen 3.1
Day 7- REP initiation	Media CM2	500 mL
	IL-2 (6000 IU/mL)	+
	OKT-3 (30ng/mL) added on Day 7	+
	Feeders Added on Day 7	500 E06
	Total Feeders at Day	750 E+06

Process Day	Conditions	Gen 3.1
Day 9-11 -Scale Up	From G-REX 100MCS transfer TIL suspension to 1 G-REX 500MCS (up to 3 GREX 500MCS)	Yes
Day 16- Harvest	LOVO- automated cell washer	Yes

Figure 117

Process Comparison	Key Process Changes	Benefit
Gen 2 : Gen 2.1	• Initiate process with two flasks instead of one flask • Divide REP initiation feeder layer between 2 G-Rex500MCS Flasks • Pre-formulate media and warm prior to use	• Potential doubling of final cell count (dose) with increased TIL repertoire. • Process redundancy throughout process
Gen 2.1 : Gen 3.1	• Fresh or Frozen tumor • 14-16 day process (from 22 day) • Reduce total feeder layer on process • Feeder layer and OKT3 present at Day 0 • REP initiated with fragments • 100MCS scales to 500MCS • Scales to multiple pre-REP flasks • Standard Media and Defined Media (Serum Free)	• Increased potency • Improved phenotype • Decreased process time • Reduced reagent testing • Decreased process variability • Defined reagents • Increased repertoire • Reduce impurities (feeder) • Comparable or Higher Dose.

Figure 118

Process Comparison	Key Process Changes	Desired Improvement	Criteria for Success	Outcome
Gen 2 : Gen 3.0	• 14-16 days • Initiate REP with fragments up to 4 flask. • 100MCS scales to 500MCS	• Increased potency • Improved phenotype • Decreased process time	• Increase potency as measured by INF-g ✓ • Comparable phenotype ✓ • Comparable Dose ✓ • Comparable purity ✓ (feeder cell) • Maintain clonal diversity ✓	• Potency increased over Gen2 • Improved expression of CD28 on CD8 cells • Maximum capacity of flask reached by day 16 on Gen 3.1 • Reduced feeder cell usage • Increased diversity

Figure 119

Process	Gen 2	Gen 3
L4054	Standard Media	Standard Media
L4055	Standard Media	Standard Media
M1085T	Standard Media	Standard Media

Process	Gen 3	Gen 3.1 control	Gen 3.1
L4063	Standard Media	Standard Media	Standard Media
L4064	Defined Media	Defined Media	Defined Media

Standard Media:
Pre REP: CM1
REP initiation : CM2
Split or Scale up : CM4

Defined Media:
CTS Optimizer (Serum Free Media) in each day of the process

Figure 120



Tumor ID	L4063 Standard media			L4064 Defined Media		
Process	Gen 3.0	Gen 3.1 Control	Gen 3.1	Gen 3.0	Gen 3.1 Control	Gen 3.1
# Fragments	53	53	53	21	21	21

Figure 121





L4063- Standard Media:

L4064- Defined Media

Process Day	Conditions	Gen 3.0	Gen 3.1 CT (Control)	Gen 3.1 F
Day 0- pre REP initiation	Media CM1	200 mL	500 mL	500 mL
	IL-2 (6000 IU/mL)	+	+	+
	OKT-3 (30ng/mL)	-	+	+
	Feeders (250 E+06)	-	-	+

Figure 122


Unique CDR3
30000
20000
10000
0
L4054 Gen2
L4054 Gen3
L4055 Gen2
L4055 Gen3
L4063 Gen3.0
L4063 Gen3.1CT
L4063 Gen3.1
L4064 Gen3.0
L4064 Gen3.1CT
L4064 Gen3.1
Diversity Index
15
10
5
0
L4054 Gen2
L4054 Gen3
L4055 Gen2
L4055 Gen3
L4063 Gen3.0
L4063 Gen3.1CT
L4063 Gen3.1
L4064 Gen3.0
L4064 Gen3.1CT
L4064 Gen3.1

Figure 123

Conditions on Day 0	Gen 3.0	Gen 3.1 (control)	Gen 3.1
Media	500 mL	500 mL	500 mL
IL-2 (6E+06)	+	+	+
OKT-3 (30ng/mL)	-	+	+
Feeders (250 E+06)	-	-	+

Number of uCDR3 (% Overlap)	All uCDR3's			Top 80% uCDR3's		
	Gen 3.0	Gen 3.1 Control	Gen 3.1	Gen 3.1	Gen 3.1 control	Gen 3.1
Gen 3.0	14005	3240 (23)	2959 (21)	1105	1000 (90)	975 (88)
Gen 3.1 control	-	18149	-	-	2438	-
Gen 3.1	-	-	17181	-	-	2332



Figure 124

Conditions on Day 0	Gen 3.0	Gen 3.1 (control)	Gen 3.1
Media	500 mL	500 mL	500 mL
IL-2 (6E+06)	+	+	+
OKT-3 (30ng/mL)	-	+	+
Feeders (250 E+06)	-	-	+

Number of uCDR3 (% Overlap)	All uCDR3's			Top 80% uCDR3's		
	Gen 3.0	Gen 3.1 Control	Gen 3.1	Gen 3.0	Gen 3.1 Control	Gen 3.1
Gen 3.0	15107	5486 (36)	5541 (37)	2479	2186 (88)	2163 (87)
Gen 3.1 Control	-	21460	-	-	3247	-
Gen 3.1	-	-	21503	-	-	3264



Figure 125


Tumor Dissection
Day 0: Pre-REP
≤240 fragments
≤60 fragments/flask , ≤4 flasks
≤2L media (500mL/flask) + IL-2 (6000IU/mL)
2.5E8 feeder cells/flask
30ng OKT3/flask
Day 7: REP- Initiation
Fresh TIL direct to REP Day 7. Activate entire culture 5E8 feeder cells + 30 ng OKT3/flask in G-Rex 100MCS
Add 500mL CM1 media+ IL-2(6000IU/mL) / flask
Day 11: Scale Up
Scale up entire culture 24 G-REX 500MCS.
Add 4L of CM4 media /flask +IL-2(3000IU/mL)
Day 10-11
Day 16: Harvest
Harvest Day 16
LOVO- automated cell washer
5 wash cycles*

Figure 126


Day 0
•Isolate T cell fraction (CD3+,CD45+) from an apheresis product enriched for lymphocytes, whole blood, or tumor digest (fresh or thawed) using positve or negative selection methods,i.e removing the T-cells using a T-cell marker (CD2,CD3,etc, or removing other cells leaving T-cells), or gradient centrifugation.
–
•Enter Gen 3.1 process by seeding ~1x10^7 cells/ flask according to Gen3 process
Day 7
•Reactivate per Gen3
Day 9-11
•Scale up per Gen 3
Day 14-16
•Harvest per Gen 3

Figure 127


Melanoma Fragments
pre-REP
REP
N=6
% of Live Cells
150
100
50
0
CD4+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
CD8+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
CD4+
CD8+
Melanoma Cores
REP1
REP2
N=4
% of Live Cells
150
100
50
0
CD4+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
CD8+
CD27+ CD28+
TCM
TSCM
TEMRA
TEM
CD4+
CD8+

Figure 128


REP1
REP2
Same trend observed in the pancreatic cores
% of CD3+
100
50
0
Lag3
PD1
Tigit
TIM3
CD4
CD8
REP1
REP2
CD69 and CD103 are reduced in REP2, also observed with the pancreatic cores
% of CD3+
100
80
60
40
20
0
2B4
BTLA
CD25
CD69
CD103
KLRG1
CD4
CD8

Figure 129

	Culture conditions	Cores / cond. (#)	REP1	REP2
Lung (L4083)	Day 0: Addition of OKT3-feeders Day 3: Addition of OKT3-feeders	1	>50ES cells at Day 11 in Day 0 condition	1157-fold (Day 0) and 925-fold (1/2 cores) expansion ~ 10,350 and 7,663 pg/ml IFN$_\gamma$ (respectively)



Figure 130


100
75
50
25
0
%CD3
TCM
TSCM
TEMRA
TEM
REP2
Day 0
Day 3
100
75
50
25
0
%CD3
CD3
CD4
CD8
CD27
CD28
KLRG1
CD57
REP2
Day 0
Day 3

Figure 131


100
75
50
25
0
% CD3
Day 0
Day 3
CD25
CD69
CD134
CD137
CD39
CD49a
CD103
LAG3
PD-1+
TIM3
CD101
REP2

Figure 132


15000
10000
5000
0
IFNγ (pg/ml)
Day 0
Day 3
REP2
Unstimulated
Stimulated
40
30
20
10
0
%CD3
Day 0
Day 3
Day 0
Day 3
CD4
CD8
REP2
Unstimulated
PMA

Figure 133


% TCRvb (of CD4)
20
15
10
5
0
Day 0
Day 3
Vb1
Vb2
Vb3
Vb4
Vb5.1
Vb5.2
Vb5.3
Vb7
Vb7.1
Vb7.2
Vb8
Vb9
Vb11
Vb12
Vb13.1
Vb13.2
Vb13.6
Vb14
Vb16
Vb18
Vb20
Vb21.3
Vb22
Vb23
% TCRvb (of CD8)
60
40
20
0
Day 0
Day 3
Vb1
Vb2
Vb3
Vb4
Vb5.1
Vb5.2
Vb5.3
Vb7
Vb7.1
Vb7.2
Vb8
Vb9
Vb11
Vb12
Vb13.1
Vb13.2
Vb13.6
Vb14
Vb16
Vb18
Vb20
Vb21.3
Vb22
Vb23

SEQUENCE LISTING

<110> Iovance Biotherapeutics, Inc.

<120> TIL EXPANSION FROM FINE NEEDLE ASPIRATES AND SMALL BIOPSIES

<130> 116983-5033

<150> US 62/588,044
<151> 2017-11-17

<150> US 62/621,515
<151> 2018-01-24

<150> US 62/756,038
<151> 2018-11-05

<160> 126

<170> PatentIn version 3.5

<210> 1
<211> 450
<212> PRT
<213> Artificial Sequence

<220>
<223> Muromonab heavy chain

<400> 1

Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Arg Tyr
20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys

85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Thr Leu Thr Val Ser Ser Ala Lys Thr Thr Ala Pro Ser Val Tyr
115 120 125

Pro Leu Ala Pro Val Cys Gly Gly Thr Thr Gly Ser Ser Val Thr Leu
130 135 140

Gly Cys Leu Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Leu Thr Trp
145 150 155 160

Asn Ser Gly Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu
165 170 175

Gln Ser Asp Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Thr Ser Ser
180 185 190

Thr Trp Pro Ser Gln Ser Ile Thr Cys Asn Val Ala His Pro Ala Ser
195 200 205

Ser Thr Lys Val Asp Lys Lys Ile Glu Pro Arg Pro Lys Ser Cys Asp
210 215 220

Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
225 230 235 240

Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
245 250 255

Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
260 265 270

Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
275 280 285

Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg

```
    290                 295                 300

Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
305                 310                 315                 320

Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
                325                 330                 335

Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
            340                 345                 350

Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
        355                 360                 365

Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
    370                 375                 380

Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
385                 390                 395                 400

Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                405                 410                 415

Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
            420                 425                 430

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
        435                 440                 445

Gly Lys
    450


<210>  2
<211>  213
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  Muromonab light chain

<400>  2
```

```
Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1               5                  10                  15

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
            20                  25                  30

Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
        35                  40                  45

Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala His Phe Arg Gly Ser
    50                  55                  60

Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Gly Met Glu Ala Glu
65                  70                  75                  80

Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Phe Thr
                85                  90                  95

Phe Gly Ser Gly Thr Lys Leu Glu Ile Asn Arg Ala Asp Thr Ala Pro
            100                 105                 110

Thr Val Ser Ile Phe Pro Pro Ser Ser Glu Gln Leu Thr Ser Gly Gly
        115                 120                 125

Ala Ser Val Val Cys Phe Leu Asn Asn Phe Tyr Pro Lys Asp Ile Asn
    130                 135                 140

Val Lys Trp Lys Ile Asp Gly Ser Glu Arg Gln Asn Gly Val Leu Asn
145                 150                 155                 160

Ser Trp Thr Asp Gln Asp Ser Lys Asp Ser Thr Tyr Ser Met Ser Ser
                165                 170                 175

Thr Leu Thr Leu Thr Lys Asp Glu Tyr Glu Arg His Asn Ser Tyr Thr
            180                 185                 190

Cys Glu Ala Thr His Lys Thr Ser Thr Ser Pro Ile Val Lys Ser Phe
        195                 200                 205
```

Asn Arg Asn Glu Cys
210

<210> 3
<211> 134
<212> PRT
<213> Artificial Sequence

<220>
<223> recombinant human IL-2 (rhIL-2)

<400> 3

Met Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu
1 5 10 15

His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr
20 25 30

Lys Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro
35 40 45

Lys Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu
50 55 60

Lys Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His
65 70 75 80

Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu
85 90 95

Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr
100 105 110

Ala Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser
115 120 125

Ile Ile Ser Thr Leu Thr
130

<210> 4

<211> 132
<212> PRT
<213> Artificial Sequence

<220>
<223> Aldesleukin

<400> 4

```
Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His Leu
1               5                  10                  15


Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys Asn
            20                  25                  30


Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro Lys Lys
        35                  40                  45


Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys Pro
    50                  55                  60


Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His Leu Arg
65                  70                  75                  80


Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu Lys
                85                  90                  95


Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala Thr
            100                 105                 110


Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ser Gln Ser Ile Ile
        115                 120                 125


Ser Thr Leu Thr
    130
```

<210> 5
<211> 130
<212> PRT
<213> Artificial Sequence

<220>
<223> recombinant human IL-4

<400> 5

Met His Lys Cys Asp Ile Thr Leu Gln Glu Ile Ile Lys Thr Leu Asn
1 5 10 15

Ser Leu Thr Glu Gln Lys Thr Leu Cys Thr Glu Leu Thr Val Thr Asp
20 25 30

Ile Phe Ala Ala Ser Lys Asn Thr Thr Glu Lys Glu Thr Phe Cys Arg
35 40 45

Ala Ala Thr Val Leu Arg Gln Phe Tyr Ser His His Glu Lys Asp Thr
50 55 60

Arg Cys Leu Gly Ala Thr Ala Gln Gln Phe His Arg His Lys Gln Leu
65 70 75 80

Ile Arg Phe Leu Lys Arg Leu Asp Arg Asn Leu Trp Gly Leu Ala Gly
85 90 95

Leu Asn Ser Cys Pro Val Lys Glu Ala Asn Gln Ser Thr Leu Glu Asn
100 105 110

Phe Leu Glu Arg Leu Lys Thr Ile Met Arg Glu Lys Tyr Ser Lys Cys
115 120 125

Ser Ser
130

<210> 6
<211> 153
<212> PRT
<213> Artificial Sequence

<220>
<223> recombinant human IL-7 (rhIL-7)

<400> 6

Met Asp Cys Asp Ile Glu Gly Lys Asp Gly Lys Gln Tyr Glu Ser Val
1 5 10 15

```
Leu Met Val Ser Ile Asp Gln Leu Leu Asp Ser Met Lys Glu Ile Gly
            20                  25                  30

Ser Asn Cys Leu Asn Asn Glu Phe Asn Phe Phe Lys Arg His Ile Cys
        35                  40                  45

Asp Ala Asn Lys Glu Gly Met Phe Leu Phe Arg Ala Ala Arg Lys Leu
    50                  55                  60

Arg Gln Phe Leu Lys Met Asn Ser Thr Gly Asp Phe Asp Leu His Leu
65                  70                  75                  80

Leu Lys Val Ser Glu Gly Thr Thr Ile Leu Leu Asn Cys Thr Gly Gln
                85                  90                  95

Val Lys Gly Arg Lys Pro Ala Ala Leu Gly Glu Ala Gln Pro Thr Lys
            100                 105                 110

Ser Leu Glu Glu Asn Lys Ser Leu Lys Glu Gln Lys Lys Leu Asn Asp
        115                 120                 125

Leu Cys Phe Leu Lys Arg Leu Leu Gln Glu Ile Lys Thr Cys Trp Asn
    130                 135                 140

Lys Ile Leu Met Gly Thr Lys Glu His
145                 150
```

<210> 7
<211> 115
<212> PRT
<213> Artificial Sequence

<220>
<223> recombinant human IL-15 (rhIL-15)

<400> 7

```
Met Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu Asp Leu
1               5                   10                  15

Ile Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser Asp Val
```

20 25 30

His Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu Glu Leu
35 40 45

Gln Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp Thr Val
50 55 60

Glu Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn Gly Asn
65 70 75 80

Val Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu Lys Asn
85 90 95

Ile Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met Phe Ile
100 105 110

Asn Thr Ser
115

<210> 8
<211> 132
<212> PRT
<213> Artificial Sequence

<220>
<223> recombinant human IL-21 (rhIL-21)

<400> 8

Met Gln Asp Arg His Met Ile Arg Met Arg Gln Leu Ile Asp Ile Val
1 5 10 15

Asp Gln Leu Lys Asn Tyr Val Asn Asp Leu Val Pro Glu Phe Leu Pro
20 25 30

Ala Pro Glu Asp Val Glu Thr Asn Cys Glu Trp Ser Ala Phe Ser Cys
35 40 45

Phe Gln Lys Ala Gln Leu Lys Ser Ala Asn Thr Gly Asn Asn Glu Arg
50 55 60

```
Ile Ile Asn Val Ser Ile Lys Lys Leu Lys Arg Lys Pro Pro Ser Thr
65                  70                  75                  80


Asn Ala Gly Arg Arg Gln Lys His Arg Leu Thr Cys Pro Ser Cys Asp
                85                  90                  95


Ser Tyr Glu Lys Lys Pro Pro Lys Glu Phe Leu Glu Arg Phe Lys Ser
            100                 105                 110


Leu Leu Gln Lys Met Ile His Gln His Leu Ser Ser Arg Thr His Gly
        115                 120                 125


Ser Glu Asp Ser
    130


<210>  9
<211>  255
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  human 4-1BB, Tumor necrosis factor receptor superfamily, member 9
       (Homo sapiens)

<400>  9

Met Gly Asn Ser Cys Tyr Asn Ile Val Ala Thr Leu Leu Leu Val Leu
1               5                   10                  15


Asn Phe Glu Arg Thr Arg Ser Leu Gln Asp Pro Cys Ser Asn Cys Pro
            20                  25                  30


Ala Gly Thr Phe Cys Asp Asn Asn Arg Asn Gln Ile Cys Ser Pro Cys
        35                  40                  45


Pro Pro Asn Ser Phe Ser Ser Ala Gly Gly Gln Arg Thr Cys Asp Ile
    50                  55                  60


Cys Arg Gln Cys Lys Gly Val Phe Arg Thr Arg Lys Glu Cys Ser Ser
65                  70                  75                  80
```

```
Thr Ser Asn Ala Glu Cys Asp Cys Thr Pro Gly Phe His Cys Leu Gly
                85                  90                  95

Ala Gly Cys Ser Met Cys Glu Gln Asp Cys Lys Gln Gly Gln Glu Leu
            100                 105                 110

Thr Lys Lys Gly Cys Lys Asp Cys Cys Phe Gly Thr Phe Asn Asp Gln
        115                 120                 125

Lys Arg Gly Ile Cys Arg Pro Trp Thr Asn Cys Ser Leu Asp Gly Lys
    130                 135                 140

Ser Val Leu Val Asn Gly Thr Lys Glu Arg Asp Val Val Cys Gly Pro
145                 150                 155                 160

Ser Pro Ala Asp Leu Ser Pro Gly Ala Ser Ser Val Thr Pro Pro Ala
                165                 170                 175

Pro Ala Arg Glu Pro Gly His Ser Pro Gln Ile Ile Ser Phe Phe Leu
            180                 185                 190

Ala Leu Thr Ser Thr Ala Leu Leu Phe Leu Leu Phe Phe Leu Thr Leu
        195                 200                 205

Arg Phe Ser Val Val Lys Arg Gly Arg Lys Lys Leu Leu Tyr Ile Phe
    210                 215                 220

Lys Gln Pro Phe Met Arg Pro Val Gln Thr Thr Gln Glu Glu Asp Gly
225                 230                 235                 240

Cys Ser Cys Arg Phe Pro Glu Glu Glu Glu Gly Gly Cys Glu Leu
                245                 250                 255
```

<210> 10
<211> 256
<212> PRT
<213> Artificial Sequence

<220>
<223> murine 4-1BB, Tumor necrosis factor receptor superfamily, member 9 (Mus musculus)

<400> 10

```
Met Gly Asn Asn Cys Tyr Asn Val Val Val Ile Val Leu Leu Leu Val
1               5                  10                  15

Gly Cys Glu Lys Val Gly Ala Val Gln Asn Ser Cys Asp Asn Cys Gln
            20                  25                  30

Pro Gly Thr Phe Cys Arg Lys Tyr Asn Pro Val Cys Lys Ser Cys Pro
        35                  40                  45

Pro Ser Thr Phe Ser Ser Ile Gly Gly Gln Pro Asn Cys Asn Ile Cys
    50                  55                  60

Arg Val Cys Ala Gly Tyr Phe Arg Phe Lys Lys Phe Cys Ser Ser Thr
65                  70                  75                  80

His Asn Ala Glu Cys Glu Cys Ile Glu Gly Phe His Cys Leu Gly Pro
                85                  90                  95

Gln Cys Thr Arg Cys Glu Lys Asp Cys Arg Pro Gly Gln Glu Leu Thr
            100                 105                 110

Lys Gln Gly Cys Lys Thr Cys Ser Leu Gly Thr Phe Asn Asp Gln Asn
        115                 120                 125

Gly Thr Gly Val Cys Arg Pro Trp Thr Asn Cys Ser Leu Asp Gly Arg
    130                 135                 140

Ser Val Leu Lys Thr Gly Thr Thr Glu Lys Asp Val Val Cys Gly Pro
145                 150                 155                 160

Pro Val Val Ser Phe Ser Pro Ser Thr Thr Ile Ser Val Thr Pro Glu
                165                 170                 175

Gly Gly Pro Gly Gly His Ser Leu Gln Val Leu Thr Leu Phe Leu Ala
            180                 185                 190

Leu Thr Ser Ala Leu Leu Leu Ala Leu Ile Phe Ile Thr Leu Leu Phe
```

195 200 205

Ser Val Leu Lys Trp Ile Arg Lys Lys Phe Pro His Ile Phe Lys Gln
210 215 220

Pro Phe Lys Lys Thr Thr Gly Ala Ala Gln Glu Glu Asp Ala Cys Ser
225 230 235 240

Cys Arg Cys Pro Gln Glu Glu Glu Gly Gly Gly Gly Tyr Glu Leu
245 250 255

<210> 11
<211> 441
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain for utomilumab

<400> 11

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

Ser Leu Arg Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Ser Thr Tyr
20 25 30

Trp Ile Ser Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
35 40 45

Gly Lys Ile Tyr Pro Gly Asp Ser Tyr Thr Asn Tyr Ser Pro Ser Phe
50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ala Arg Gly Tyr Gly Ile Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val
100 105 110

```
Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala
        115                 120                 125

Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu
    130                 135                 140

Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly
145                 150                 155                 160

Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser
                165                 170                 175

Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe
            180                 185                 190

Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr
        195                 200                 205

Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro
    210                 215                 220

Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro
225                 230                 235                 240

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
                245                 250                 255

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp
            260                 265                 270

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
        275                 280                 285

Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val
    290                 295                 300

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
305                 310                 315                 320
```

```
Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly
                325                 330                 335


Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
            340                 345                 350


Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
        355                 360                 365


Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
    370                 375                 380


Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe
385                 390                 395                 400


Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
                405                 410                 415


Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
            420                 425                 430


Gln Lys Ser Leu Ser Leu Ser Pro Gly
        435                 440


<210>  12
<211>  214
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  light chain for utomilumab

<400>  12

Ser Tyr Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln
1               5                   10                  15


Thr Ala Ser Ile Thr Cys Ser Gly Asp Asn Ile Gly Asp Gln Tyr Ala
            20                  25                  30


His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Val Leu Val Ile Tyr
```

```
        35                  40                  45

Gln Asp Lys Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
    50                  55                  60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Met
65                  70                  75                  80

Asp Glu Ala Asp Tyr Tyr Cys Ala Thr Tyr Thr Gly Phe Gly Ser Leu
                85                  90                  95

Ala Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln Pro Lys
            100                 105                 110

Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln
        115                 120                 125

Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly
    130                 135                 140

Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys Ala Gly
145                 150                 155                 160

Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala
                165                 170                 175

Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Arg Ser
            180                 185                 190

Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr Val
        195                 200                 205

Ala Pro Thr Glu Cys Ser
    210
```

<210> 13
<211> 116
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region for utomilumab

<400> 13

```
Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1               5                  10                  15


Ser Leu Arg Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Ser Thr Tyr
            20                  25                  30


Trp Ile Ser Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
        35                  40                  45


Gly Lys Ile Tyr Pro Gly Asp Ser Tyr Thr Asn Tyr Ser Pro Ser Phe
    50                  55                  60


Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65                  70                  75                  80


Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
                85                  90                  95


Ala Arg Gly Tyr Gly Ile Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val
            100                 105                 110


Thr Val Ser Ser
        115
```

<210> 14
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region for utomilumab

<400> 14

```
Ser Tyr Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln
1               5                  10                  15


Thr Ala Ser Ile Thr Cys Ser Gly Asp Asn Ile Gly Asp Gln Tyr Ala
```

20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Val Leu Val Ile Tyr
35 40 45

Gln Asp Lys Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
50 55 60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Met
65 70 75 80

Asp Glu Ala Asp Tyr Tyr Cys Ala Thr Tyr Thr Gly Phe Gly Ser Leu
85 90 95

Ala Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu
100 105

<210> 15
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR1 for utomilumab

<400> 15

Ser Thr Tyr Trp Ile Ser
1 5

<210> 16
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR2 for utomilumab

<400> 16

Lys Ile Tyr Pro Gly Asp Ser Tyr Thr Asn Tyr Ser Pro Ser Phe Gln
1 5 10 15

Gly

<210> 17
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for utomilumab

<400> 17

Arg Gly Tyr Gly Ile Phe Asp Tyr
1 5

<210> 18
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for utomilumab

<400> 18

Ser Gly Asp Asn Ile Gly Asp Gln Tyr Ala His
1 5 10

<210> 19
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR2 for utomilumab

<400> 19

Gln Asp Lys Asn Arg Pro Ser
1 5

<210> 20
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for utomilumab

<400> 20

Ala Thr Tyr Thr Gly Phe Gly Ser Leu Ala Val
1 5 10

<210> 21
<211> 448
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain for urelumab

<400> 21

Gln Val Gln Leu Gln Gln Trp Gly Ala Gly Leu Leu Lys Pro Ser Glu
1 5 10 15

Thr Leu Ser Leu Thr Cys Ala Val Tyr Gly Gly Ser Phe Ser Gly Tyr
20 25 30

Tyr Trp Ser Trp Ile Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Ile
35 40 45

Gly Glu Ile Asn His Gly Gly Tyr Val Thr Tyr Asn Pro Ser Leu Glu
50 55 60

Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

Arg Asp Tyr Gly Pro Gly Asn Tyr Asp Trp Tyr Phe Asp Leu Trp Gly
100 105 110

Arg Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
115 120 125

Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala
130 135 140

```
Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
145                 150                 155                 160

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
                165                 170                 175

Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
            180                 185                 190

Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His
        195                 200                 205

Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly
    210                 215                 220

Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser
225                 230                 235                 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
                245                 250                 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro
            260                 265                 270

Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
        275                 280                 285

Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val
    290                 295                 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
305                 310                 315                 320

Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr
                325                 330                 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
            340                 345                 350
```

```
Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
        355                 360                 365


Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
    370                 375                 380


Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
385                 390                 395                 400


Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser
                405                 410                 415


Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
            420                 425                 430


Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
        435                 440                 445


<210>  22
<211>  216
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  light chain for urelumab

<400>  22

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1               5                   10                  15


Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
            20                  25                  30


Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
        35                  40                  45


Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
    50                  55                  60


Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
```

65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
85 90 95

Ala Leu Thr Phe Cys Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val
100 105 110

Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
115 120 125

Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
130 135 140

Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
145 150 155 160

Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser
165 170 175

Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
180 185 190

Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
195 200 205

Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> 23
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain for urelumab

<400> 23

Met Lys His Leu Trp Phe Phe Leu Leu Leu Val Ala Ala Pro Arg Trp
1 5 10 15

```
Val Leu Ser Gln Val Gln Leu Gln Gln Trp Gly Ala Gly Leu Leu Lys
            20                  25                  30


Pro Ser Glu Thr Leu Ser Leu Thr Cys Ala Val Tyr Gly Gly Ser Phe
        35                  40                  45


Ser Gly Tyr Tyr Trp Ser Trp Ile Arg Gln Ser Pro Glu Lys Gly Leu
    50                  55                  60


Glu Trp Ile Gly Glu Ile Asn His Gly Gly Tyr Val Thr Tyr Asn Pro
65                  70                  75                  80


Ser Leu Glu Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln
                85                  90                  95


Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr
            100                 105                 110


Tyr Cys Ala Arg Asp Tyr Gly Pro
        115                 120


<210>  24
<211>  110
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  variable light chain for urelumab

<400>  24

Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
1               5                   10                  15


Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser
            20                  25                  30


Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
        35                  40                  45


Val Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
```

50 55 60

Arg Leu Leu Ile Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala
65 70 75 80

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
85 90 95

Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln
100 105 110

<210> 25
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR1 for urelumab

<400> 25

Gly Tyr Tyr Trp Ser
1 5

<210> 26
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR2 for urelumab

<400> 26

Glu Ile Asn His Gly Gly Tyr Val Thr Tyr Asn Pro Ser Leu Glu Ser
1 5 10 15

<210> 27
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for urelumab

<400> 27

Asp Tyr Gly Pro Gly Asn Tyr Asp Trp Tyr Phe Asp Leu
1 5 10

<210> 28
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for urelumab

<400> 28

Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
1 5 10

<210> 29
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR2 for urelumab

<400> 29

Asp Ala Ser Asn Arg Ala Thr
1 5

<210> 30
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for urelumab

<400> 30

Gln Gln Arg Ser Asp Trp Pro Pro Ala Leu Thr
1 5 10

<210> 31
<211> 230
<212> PRT
<213> Artificial Sequence

<220>
<223> Fc domain

<400> 31

```
Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
1               5                  10                  15


Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
            20                  25                  30


Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
        35                  40                  45


Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
    50                  55                  60


Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
65                  70                  75                  80


Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
                85                  90                  95


Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro
            100                 105                 110


Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
        115                 120                 125


Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
    130                 135                 140


Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
145                 150                 155                 160


Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
                165                 170                 175


Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
            180                 185                 190
```

```
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
        195                 200                 205


Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
    210                 215                 220


Ser Leu Ser Pro Gly Lys
225                 230


<210>  32
<211>  22
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  linker

<400>  32

Gly Gly Pro Gly Ser Ser Lys Ser Cys Asp Lys Thr His Thr Cys Pro
1               5                   10                  15


Pro Cys Pro Ala Pro Glu
            20


<210>  33
<211>  22
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  linker

<400>  33

Gly Gly Ser Gly Ser Ser Lys Ser Cys Asp Lys Thr His Thr Cys Pro
1               5                   10                  15


Pro Cys Pro Ala Pro Glu
            20


<210>  34
<211>  27
<212>  PRT
```

<213> Artificial Sequence

<220>
<223> linker

<400> 34

Gly Gly Pro Gly Ser Ser Ser Ser Ser Ser Ser Lys Ser Cys Asp Lys
1 5 10 15

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
20 25

<210> 35
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 35

Gly Gly Ser Gly Ser Ser Ser Ser Ser Ser Ser Lys Ser Cys Asp Lys
1 5 10 15

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
20 25

<210> 36
<211> 29
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 36

Gly Gly Pro Gly Ser Ser Ser Ser Ser Ser Ser Ser Ser Lys Ser Cys
1 5 10 15

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
20 25

<210> 37

<211> 29
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 37

Gly Gly Ser Gly Ser Ser Ser Ser Ser Ser Ser Ser Ser Lys Ser Cys
1 5 10 15

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
20 25

<210> 38
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 38

Gly Gly Pro Gly Ser Ser Gly Ser Gly Ser Ser Asp Lys Thr His Thr
1 5 10 15

Cys Pro Pro Cys Pro Ala Pro Glu
20

<210> 39
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 39

Gly Gly Pro Gly Ser Ser Gly Ser Gly Ser Asp Lys Thr His Thr Cys
1 5 10 15

Pro Pro Cys Pro Ala Pro Glu
20

<210> 40
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 40

Gly Gly Pro Ser Ser Ser Gly Ser Asp Lys Thr His Thr Cys Pro Pro
1 5 10 15

Cys Pro Ala Pro Glu
20

<210> 41
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 41

Gly Gly Ser Ser Ser Ser Ser Ser Ser Ser Gly Ser Asp Lys Thr His
1 5 10 15

Thr Cys Pro Pro Cys Pro Ala Pro Glu
20 25

<210> 42
<211> 246
<212> PRT
<213> Artificial Sequence

<220>
<223> Fc domain

<400> 42

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
1 5 10 15

Ala Gly Asn Gly Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro

```
            20                  25                  30

Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
        35                  40                  45

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
    50                  55                  60

Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
65                  70                  75                  80

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
                85                  90                  95

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
            100                 105                 110

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu
        115                 120                 125

Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
    130                 135                 140

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys
145                 150                 155                 160

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
                165                 170                 175

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
            180                 185                 190

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
        195                 200                 205

Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
    210                 215                 220

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
```

225 230 235 240

Leu Ser Leu Ser Pro Gly
245

<210> 43
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 43

Ser Gly Ser Gly Ser Gly Ser Gly Ser Gly Ser
1 5 10

<210> 44
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 44

Ser Ser Ser Ser Ser Ser Gly Ser Gly Ser Gly Ser
1 5 10

<210> 45
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> linker

<400> 45

Ser Ser Ser Ser Ser Ser Gly Ser Gly Ser Gly Ser Gly Ser Gly Ser
1 5 10 15

<210> 46
<211> 254
<212> PRT

<213> Artificial Sequence

<220>
<223> 4-1BBL

<400> 46

```
Met Glu Tyr Ala Ser Asp Ala Ser Leu Asp Pro Glu Ala Pro Trp Pro
1               5                  10                  15


Pro Ala Pro Arg Ala Arg Ala Cys Arg Val Leu Pro Trp Ala Leu Val
            20                  25                  30


Ala Gly Leu Leu Leu Leu Leu Leu Leu Ala Ala Ala Cys Ala Val Phe
        35                  40                  45


Leu Ala Cys Pro Trp Ala Val Ser Gly Ala Arg Ala Ser Pro Gly Ser
    50                  55                  60


Ala Ala Ser Pro Arg Leu Arg Glu Gly Pro Glu Leu Ser Pro Asp Asp
65                  70                  75                  80


Pro Ala Gly Leu Leu Asp Leu Arg Gln Gly Met Phe Ala Gln Leu Val
                85                  90                  95


Ala Gln Asn Val Leu Leu Ile Asp Gly Pro Leu Ser Trp Tyr Ser Asp
            100                 105                 110


Pro Gly Leu Ala Gly Val Ser Leu Thr Gly Gly Leu Ser Tyr Lys Glu
        115                 120                 125


Asp Thr Lys Glu Leu Val Val Ala Lys Ala Gly Val Tyr Tyr Val Phe
    130                 135                 140


Phe Gln Leu Glu Leu Arg Arg Val Val Ala Gly Glu Gly Ser Gly Ser
145                 150                 155                 160


Val Ser Leu Ala Leu His Leu Gln Pro Leu Arg Ser Ala Ala Gly Ala
                165                 170                 175


Ala Ala Leu Ala Leu Thr Val Asp Leu Pro Pro Ala Ser Ser Glu Ala
```

180 185 190

Arg Asn Ser Ala Phe Gly Phe Gln Gly Arg Leu Leu His Leu Ser Ala
195 200 205

Gly Gln Arg Leu Gly Val His Leu His Thr Glu Ala Arg Ala Arg His
210 215 220

Ala Trp Gln Leu Thr Gln Gly Ala Thr Val Leu Gly Leu Phe Arg Val
225 230 235 240

Thr Pro Glu Ile Pro Ala Gly Leu Pro Ser Pro Arg Ser Glu
245 250

<210> 47
<211> 168
<212> PRT
<213> Artificial Sequence

<220>
<223> 4-1BBL soluble domain

<400> 47

Leu Arg Gln Gly Met Phe Ala Gln Leu Val Ala Gln Asn Val Leu Leu
1 5 10 15

Ile Asp Gly Pro Leu Ser Trp Tyr Ser Asp Pro Gly Leu Ala Gly Val
20 25 30

Ser Leu Thr Gly Gly Leu Ser Tyr Lys Glu Asp Thr Lys Glu Leu Val
35 40 45

Val Ala Lys Ala Gly Val Tyr Tyr Val Phe Phe Gln Leu Glu Leu Arg
50 55 60

Arg Val Val Ala Gly Glu Gly Ser Gly Ser Val Ser Leu Ala Leu His
65 70 75 80

Leu Gln Pro Leu Arg Ser Ala Ala Gly Ala Ala Ala Leu Ala Leu Thr
85 90 95

```
Val Asp Leu Pro Pro Ala Ser Ser Glu Ala Arg Asn Ser Ala Phe Gly
            100                 105                 110


Phe Gln Gly Arg Leu Leu His Leu Ser Ala Gly Gln Arg Leu Gly Val
        115                 120                 125


His Leu His Thr Glu Ala Arg Ala Arg His Ala Trp Gln Leu Thr Gln
    130                 135                 140


Gly Ala Thr Val Leu Gly Leu Phe Arg Val Thr Pro Glu Ile Pro Ala
145                 150                 155                 160


Gly Leu Pro Ser Pro Arg Ser Glu
                165


<210>  48
<211>  118
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  variable heavy chain for 4B4-1-1 version 1

<400>  48

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1               5                   10                  15


Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Ser Tyr
            20                  25                  30


Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Val Leu Glu Trp Ile
        35                  40                  45


Gly Glu Ile Asn Pro Gly Asn Gly His Thr Asn Tyr Asn Glu Lys Phe
    50                  55                  60


Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65                  70                  75                  80


Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
```

85 90 95

Ala Arg Ser Phe Thr Thr Ala Arg Gly Phe Ala Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser
115

<210> 49
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> variable light chain for 4B4-1-1 version 1

<400> 49

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Gln Ser Val Thr Pro Gly
1 5 10 15

Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Thr Ile Ser Asp Tyr
20 25 30

Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
35 40 45

Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Pro
65 70 75 80

Glu Asp Val Gly Val Tyr Tyr Cys Gln Asp Gly His Ser Phe Pro Pro
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 50
<211> 119
<212> PRT

<213> Artificial Sequence

<220>
<223> variable heavy chain for 4B4-1-1 version 2

<400> 50

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Ser Tyr
20 25 30

Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Val Leu Glu Trp Ile
35 40 45

Gly Glu Ile Asn Pro Gly Asn Gly His Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ser Phe Thr Thr Ala Arg Gly Phe Ala Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ala
115

<210> 51
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> variable light chain for 4B4-1-1 version 2

<400> 51

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Gln Ser Val Thr Pro Gly
1 5 10 15

```
Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Thr Ile Ser Asp Tyr
            20                  25                  30


Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
        35                  40                  45


Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
    50                  55                  60


Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Pro
65                  70                  75                  80


Glu Asp Val Gly Val Tyr Tyr Cys Gln Asp Gly His Ser Phe Pro Pro
                85                  90                  95


Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
            100                 105


<210>  52
<211>  120
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  variable heavy chain for H39E3-2

<400>  52

Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Ala Thr Gly
1               5                   10                  15


Ala His Ser Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln
            20                  25                  30


Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
        35                  40                  45


Ser Asp Tyr Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
    50                  55                  60


Glu Trp Val Ala Asp Ile Lys Asn Asp Gly Ser Tyr Thr Asn Tyr Ala
```

65 70 75 80

Pro Ser Leu Thr Asn Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100 105 110

Tyr Tyr Cys Ala Arg Glu Leu Thr
115 120

<210> 53
<211> 109
<212> PRT
<213> Artificial Sequence

<220>
<223> variable light chain for H39E3-2

<400> 53

Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
1 5 10 15

Asp Thr Thr Gly Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala
20 25 30

Val Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser
35 40 45

Leu Leu Ser Ser Gly Asn Gln Lys Asn Tyr Leu Trp Tyr Gln Gln Lys
50 55 60

Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr Tyr Ala Ser Thr Arg Gln
65 70 75 80

Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
85 90 95

Thr Leu Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala
100 105

<210> 54
<211> 277
<212> PRT
<213> Artificial Sequence

<220>
<223> human OX40 (Homo sapiens)

<400> 54

```
Met Cys Val Gly Ala Arg Arg Leu Gly Arg Gly Pro Cys Ala Ala Leu
1               5                   10                  15


Leu Leu Leu Gly Leu Gly Leu Ser Thr Val Thr Gly Leu His Cys Val
            20                  25                  30


Gly Asp Thr Tyr Pro Ser Asn Asp Arg Cys Cys His Glu Cys Arg Pro
        35                  40                  45


Gly Asn Gly Met Val Ser Arg Cys Ser Arg Ser Gln Asn Thr Val Cys
    50                  55                  60


Arg Pro Cys Gly Pro Gly Phe Tyr Asn Asp Val Val Ser Ser Lys Pro
65                  70                  75                  80


Cys Lys Pro Cys Thr Trp Cys Asn Leu Arg Ser Gly Ser Glu Arg Lys
                85                  90                  95


Gln Leu Cys Thr Ala Thr Gln Asp Thr Val Cys Arg Cys Arg Ala Gly
            100                 105                 110


Thr Gln Pro Leu Asp Ser Tyr Lys Pro Gly Val Asp Cys Ala Pro Cys
        115                 120                 125


Pro Pro Gly His Phe Ser Pro Gly Asp Asn Gln Ala Cys Lys Pro Trp
    130                 135                 140


Thr Asn Cys Thr Leu Ala Gly Lys His Thr Leu Gln Pro Ala Ser Asn
145                 150                 155                 160


Ser Ser Asp Ala Ile Cys Glu Asp Arg Asp Pro Pro Ala Thr Gln Pro
```

```
                165                 170                 175

Gln Glu Thr Gln Gly Pro Pro Ala Arg Pro Ile Thr Val Gln Pro Thr
            180                 185                 190

Glu Ala Trp Pro Arg Thr Ser Gln Gly Pro Ser Thr Arg Pro Val Glu
        195                 200                 205

Val Pro Gly Gly Arg Ala Val Ala Ala Ile Leu Gly Leu Gly Leu Val
    210                 215                 220

Leu Gly Leu Leu Gly Pro Leu Ala Ile Leu Leu Ala Leu Tyr Leu Leu
225                 230                 235                 240

Arg Arg Asp Gln Arg Leu Pro Pro Asp Ala His Lys Pro Pro Gly Gly
                245                 250                 255

Gly Ser Phe Arg Thr Pro Ile Gln Glu Glu Gln Ala Asp Ala His Ser
            260                 265                 270

Thr Leu Ala Lys Ile
        275
```

<210> 55
<211> 272
<212> PRT
<213> Artificial Sequence

<220>
<223> murine OX40 (Mus musculus)

<400> 55

```
Met Tyr Val Trp Val Gln Gln Pro Thr Ala Leu Leu Leu Leu Gly Leu
1               5                   10                  15

Thr Leu Gly Val Thr Ala Arg Arg Leu Asn Cys Val Lys His Thr Tyr
            20                  25                  30

Pro Ser Gly His Lys Cys Cys Arg Glu Cys Gln Pro Gly His Gly Met
        35                  40                  45
```

```
Val Ser Arg Cys Asp His Thr Arg Asp Thr Leu Cys His Pro Cys Glu
    50                  55                  60

Thr Gly Phe Tyr Asn Glu Ala Val Asn Tyr Asp Thr Cys Lys Gln Cys
65                  70                  75                  80

Thr Gln Cys Asn His Arg Ser Gly Ser Glu Leu Lys Gln Asn Cys Thr
                85                  90                  95

Pro Thr Gln Asp Thr Val Cys Arg Cys Arg Pro Gly Thr Gln Pro Arg
            100                 105                 110

Gln Asp Ser Gly Tyr Lys Leu Gly Val Asp Cys Val Pro Cys Pro Pro
        115                 120                 125

Gly His Phe Ser Pro Gly Asn Asn Gln Ala Cys Lys Pro Trp Thr Asn
    130                 135                 140

Cys Thr Leu Ser Gly Lys Gln Thr Arg His Pro Ala Ser Asp Ser Leu
145                 150                 155                 160

Asp Ala Val Cys Glu Asp Arg Ser Leu Leu Ala Thr Leu Leu Trp Glu
                165                 170                 175

Thr Gln Arg Pro Thr Phe Arg Pro Thr Thr Val Gln Ser Thr Thr Val
            180                 185                 190

Trp Pro Arg Thr Ser Glu Leu Pro Ser Pro Pro Thr Leu Val Thr Pro
        195                 200                 205

Glu Gly Pro Ala Phe Ala Val Leu Leu Gly Leu Gly Leu Gly Leu Leu
    210                 215                 220

Ala Pro Leu Thr Val Leu Leu Ala Leu Tyr Leu Leu Arg Lys Ala Trp
225                 230                 235                 240

Arg Leu Pro Asn Thr Pro Lys Pro Cys Trp Gly Asn Ser Phe Arg Thr
                245                 250                 255
```

```
Pro Ile Gln Glu Glu His Thr Asp Ala His Phe Thr Leu Ala Lys Ile
            260                 265                 270


<210>  56
<211>  451
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  heavy chain for tavolixizumab

<400>  56

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1               5                   10                  15


Thr Leu Ser Leu Thr Cys Ala Val Tyr Gly Gly Ser Phe Ser Ser Gly
            20                  25                  30


Tyr Trp Asn Trp Ile Arg Lys His Pro Gly Lys Gly Leu Glu Tyr Ile
        35                  40                  45


Gly Tyr Ile Ser Tyr Asn Gly Ile Thr Tyr His Asn Pro Ser Leu Lys
    50                  55                  60


Ser Arg Ile Thr Ile Asn Arg Asp Thr Ser Lys Asn Gln Tyr Ser Leu
65                  70                  75                  80


Gln Leu Asn Ser Val Thr Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
                85                  90                  95


Arg Tyr Lys Tyr Asp Tyr Asp Gly Gly His Ala Met Asp Tyr Trp Gly
            100                 105                 110


Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
        115                 120                 125


Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
    130                 135                 140


Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
```

```
145                 150                 155                 160

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
                165                 170                 175

Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
            180                 185                 190

Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
        195                 200                 205

Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys
    210                 215                 220

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
225                 230                 235                 240

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
                245                 250                 255

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
            260                 265                 270

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
        275                 280                 285

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
    290                 295                 300

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
305                 310                 315                 320

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
                325                 330                 335

Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
            340                 345                 350

Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser
```

355 360 365

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
370 375 380

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
385 390 395 400

Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
405 410 415

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
420 425 430

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
435 440 445

Pro Gly Lys
450

<210> 57
<211> 214
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain for tavolixizumab

<400> 57

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Tyr Thr Ser Lys Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

```
Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65                  70                  75                  80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Gly Ser Ala Leu Pro Trp
                85                  90                  95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
            100                 105                 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
        115                 120                 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
    130                 135                 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145                 150                 155                 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
                165                 170                 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
            180                 185                 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
        195                 200                 205

Phe Asn Arg Gly Glu Cys
    210


<210>  58
<211>  118
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  heavy chain variable region for tavolixizumab

<400>  58

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
```

1 5 10 15

Thr Leu Ser Leu Thr Cys Ala Val Tyr Gly Gly Ser Phe Ser Ser Gly
20 25 30

Tyr Trp Asn Trp Ile Arg Lys His Pro Gly Lys Gly Leu Glu Tyr Ile
35 40 45

Gly Tyr Ile Ser Tyr Asn Gly Ile Thr Tyr His Asn Pro Ser Leu Lys
50 55 60

Ser Arg Ile Thr Ile Asn Arg Asp Thr Ser Lys Asn Gln Tyr Ser Leu
65 70 75 80

Gln Leu Asn Ser Val Thr Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

Arg Tyr Lys Tyr Asp Tyr Asp Gly Gly His Ala Met Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr
115

<210> 59
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region for tavolixizumab

<400> 59

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Tyr Thr Ser Lys Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
    50   55   60

Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65   70   75   80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Gly Ser Ala Leu Pro Trp
                85   90   95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
            100   105

<210> 60
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR1 for tavolixizumab

<400> 60

Gly Ser Phe Ser Ser Gly Tyr Trp Asn
1   5

<210> 61
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR2 for tavolixizumab

<400> 61

Tyr Ile Gly Tyr Ile Ser Tyr Asn Gly Ile Thr Tyr His
1   5   10

<210> 62
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for tavolixizumab

<400> 62

Arg Tyr Lys Tyr Asp Tyr Asp Gly Gly His Ala Met Asp Tyr
1 5 10

<210> 63
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for tavolixizumab

<400> 63

Gln Asp Ile Ser Asn Tyr Leu Asn
1 5

<210> 64
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> LLIYYTSKLH S

<400> 64

Leu Leu Ile Tyr Tyr Thr Ser Lys Leu His Ser
1 5 10

<210> 65
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for tavolixizumab

<400> 65

Gln Gln Gly Ser Ala Leu Pro Trp
1 5

<210> 66
<211> 444
<212> PRT

<213> Artificial Sequence

<220>
<223> heavy chain for 11D4

<400> 66

```
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1               5                  10                  15


Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
            20                  25                  30


Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
        35                  40                  45


Ser Tyr Ile Ser Ser Ser Ser Ser Thr Ile Asp Tyr Ala Asp Ser Val
    50                  55                  60


Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65                  70                  75                  80


Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys
                85                  90                  95


Ala Arg Glu Ser Gly Trp Tyr Leu Phe Asp Tyr Trp Gly Gln Gly Thr
            100                 105                 110


Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
        115                 120                 125


Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
    130                 135                 140


Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
145                 150                 155                 160


Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
                165                 170                 175


Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
```

```
            180                 185                 190

Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
        195                 200                 205

Asn Thr Lys Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys
    210                 215                 220

Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe
225                 230                 235                 240

Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
                245                 250                 255

Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe
            260                 265                 270

Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
        275                 280                 285

Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr
    290                 295                 300

Val Val His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
305                 310                 315                 320

Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr
                325                 330                 335

Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
            340                 345                 350

Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
        355                 360                 365

Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
    370                 375                 380

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser
```

385 390 395 400

Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
405 410 415

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
420 425 430

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
435 440

<210> 67
<211> 214
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain for 11D4

<400> 67

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
100 105 110

```
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
        115                 120                 125


Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
    130                 135                 140


Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145                 150                 155                 160


Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
                165                 170                 175


Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
            180                 185                 190


Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
        195                 200                 205


Phe Asn Arg Gly Glu Cys
    210


<210>  68
<211>  118
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  heavy chain variable region for 11D4

<400>  68

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1               5                   10                  15


Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
            20                  25                  30


Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
        35                  40                  45


Ser Tyr Ile Ser Ser Ser Ser Ser Thr Ile Asp Tyr Ala Asp Ser Val
```

```
    50                  55                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65                  70                  75                  80

Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys
                85                  90                  95

Ala Arg Glu Ser Gly Trp Tyr Leu Phe Asp Tyr Trp Gly Gln Gly Thr
            100                 105                 110

Leu Val Thr Val Ser Ser
        115


<210>  69
<211>  107
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  light chain variable region for 11D4

<400>  69

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1               5                   10                  15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
            20                  25                  30

Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
        35                  40                  45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
    50                  55                  60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65                  70                  75                  80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
                85                  90                  95
```

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 70
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR1 for 11D4

<400> 70

Ser Tyr Ser Met Asn
1 5

<210> 71
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR2 for 11D4

<400> 71

Tyr Ile Ser Ser Ser Ser Ser Thr Ile Asp Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 72
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for 11D4

<400> 72

Glu Ser Gly Trp Tyr Leu Phe Asp Tyr
1 5

<210> 73

<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for 11D4

<400> 73

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
1 5 10

<210> 74
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR2 for 11D4

<400> 74

Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 75
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for 11D4

<400> 75

Gln Gln Tyr Asn Ser Tyr Pro Pro Thr
1 5

<210> 76
<211> 450
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain for 18D8

<400> 76

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg

```
1               5                   10                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
            20                  25                  30

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
        35                  40                  45

Ser Gly Ile Ser Trp Asn Ser Gly Ser Ile Gly Tyr Ala Asp Ser Val
    50                  55                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65                  70                  75                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr Tyr Cys
                85                  90                  95

Ala Lys Asp Gln Ser Thr Ala Asp Tyr Tyr Phe Tyr Tyr Gly Met Asp
            100                 105                 110

Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys
        115                 120                 125

Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
    130                 135                 140

Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
145                 150                 155                 160

Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
                165                 170                 175

Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
            180                 185                 190

Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
        195                 200                 205

Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg
```

```
    210                 215                 220

Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
225                 230                 235                 240

Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
                245                 250                 255

Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
            260                 265                 270

Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
        275                 280                 285

Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
    290                 295                 300

Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
305                 310                 315                 320

Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
                325                 330                 335

Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
            340                 345                 350

Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
        355                 360                 365

Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
    370                 375                 380

Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met
385                 390                 395                 400

Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
                405                 410                 415

Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
```

420 425 430

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
435 440 445

Gly Lys
450

<210> 77
<211> 213
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain for 18D8

<400> 77

Glu Ile Val Val Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Thr
85 90 95

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

```
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
    130                 135                 140


Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145                 150                 155                 160


Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
                165                 170                 175


Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
            180                 185                 190


Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
        195                 200                 205


Asn Arg Gly Glu Cys
    210


<210>  78
<211>  124
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  heavy chain variable region for 18D8

<400>  78

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1               5                   10                  15


Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
            20                  25                  30


Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
        35                  40                  45


Ser Gly Ile Ser Trp Asn Ser Gly Ser Ile Gly Tyr Ala Asp Ser Val
    50                  55                  60


Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
```

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr Tyr Cys
85 90 95

Ala Lys Asp Gln Ser Thr Ala Asp Tyr Tyr Phe Tyr Tyr Gly Met Asp
100 105 110

Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 79
<211> 106
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region for 18D8

<400> 79

Glu Ile Val Val Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Thr
85 90 95

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105

<210> 80
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR1 for 18D8

<400> 80

Asp Tyr Ala Met His
1 5

<210> 81
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR2 for 18D8

<400> 81

Gly Ile Ser Trp Asn Ser Gly Ser Ile Gly Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 82
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for 18D8

<400> 82

Asp Gln Ser Thr Ala Asp Tyr Tyr Phe Tyr Tyr Gly Met Asp Val
1 5 10 15

<210> 83
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for 18D8

<400> 83

Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
1   5   10

<210> 84
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR2 for 18D8

<400> 84

Asp Ala Ser Asn Arg Ala Thr
1   5

<210> 85
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for 18D8

<400> 85

Gln Gln Arg Ser Asn Trp Pro Thr
1   5

<210> 86
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region for Hu119-122

<400> 86

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1   5   10   15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Glu Tyr Glu Phe Pro Ser His

20 25 30

Asp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
35 40 45

Ala Ala Ile Asn Ser Asp Gly Gly Ser Thr Tyr Tyr Pro Asp Thr Met
50 55 60

Glu Arg Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg His Tyr Asp Asp Tyr Tyr Ala Trp Phe Ala Tyr Trp Gly Gln
100 105 110

Gly Thr Met Val Thr Val Ser Ser
115 120

<210> 87
<211> 111
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region for Hu119-122

<400> 87

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Lys Ser Val Ser Thr Ser
20 25 30

Gly Tyr Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
35 40 45

Arg Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80

Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln His Ser Arg
85 90 95

Glu Leu Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

<210> 88
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDRl for Hu119-122

<400> 88

Ser His Asp Met Ser
1 5

<210> 89
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR2 for Hu119-122

<400> 89

Ala Ile Asn Ser Asp Gly Gly Ser Thr Tyr Tyr Pro Asp Thr Met Glu
1 5 10 15

Arg

<210> 90
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for Hu119-122

<400> 90

His Tyr Asp Asp Tyr Tyr Ala Trp Phe Ala Tyr
1 5 10

<210> 91
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for Hu119-122

<400> 91

Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser Tyr Met His
1 5 10 15

<210> 92
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR2 for Hu119-122

<400> 92

Leu Ala Ser Asn Leu Glu Ser
1 5

<210> 93
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for Hu119-122

<400> 93

Gln His Ser Arg Glu Leu Pro Leu Thr
1 5

<210> 94
<211> 122
<212> PRT

<213> Artificial Sequence

<220>
<223> heavy chain variable region for Hu106-222

<400> 94

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
1   5   10   15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
            20   25   30

Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
        35   40   45

Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe
    50   55   60

Lys Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Val Ser Thr Ala Tyr
65   70   75   80

Leu Gln Ile Ser Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Tyr Cys
                85   90   95

Ala Asn Pro Tyr Tyr Asp Tyr Val Ser Tyr Tyr Ala Met Asp Tyr Trp
            100   105   110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser
        115   120

<210> 95
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region for Hu106-222

<400> 95

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1   5   10   15

```
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Thr Ala
            20                  25                  30


Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
        35                  40                  45


Tyr Ser Ala Ser Tyr Leu Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
    50                  55                  60


Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65                  70                  75                  80


Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser Thr Pro Arg
                85                  90                  95


Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
            100                 105


<210>  96
<211>  5
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  heavy chain CDR1 for Hu106-222

<400>  96

Asp Tyr Ser Met His
1               5


<210>  97
<211>  17
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  heavy chain CDR2 for Hu106-222

<400>  97

Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe Lys
1               5                   10                  15
```

Gly

<210> 98
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain CDR3 for Hu106-222

<400> 98

Pro Tyr Tyr Asp Tyr Val Ser Tyr Tyr Ala Met Asp Tyr
1 5 10

<210> 99
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR1 for Hu106-222

<400> 99

Lys Ala Ser Gln Asp Val Ser Thr Ala Val Ala
1 5 10

<210> 100
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR2 for Hu106-222

<400> 100

Ser Ala Ser Tyr Leu Tyr Thr
1 5

<210> 101
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain CDR3 for Hu106-222

<400> 101

Gln Gln His Tyr Ser Thr Pro Arg Thr
1   5

<210> 102
<211> 183
<212> PRT
<213> Artificial Sequence

<220>
<223> OX40L

<400> 102

Met Glu Arg Val Gln Pro Leu Glu Glu Asn Val Gly Asn Ala Ala Arg
1   5   10   15

Pro Arg Phe Glu Arg Asn Lys Leu Leu Leu Val Ala Ser Val Ile Gln
            20   25   30

Gly Leu Gly Leu Leu Leu Cys Phe Thr Tyr Ile Cys Leu His Phe Ser
        35   40   45

Ala Leu Gln Val Ser His Arg Tyr Pro Arg Ile Gln Ser Ile Lys Val
    50   55   60

Gln Phe Thr Glu Tyr Lys Lys Glu Lys Gly Phe Ile Leu Thr Ser Gln
65   70   75   80

Lys Glu Asp Glu Ile Met Lys Val Gln Asn Asn Ser Val Ile Ile Asn
                85   90   95

Cys Asp Gly Phe Tyr Leu Ile Ser Leu Lys Gly Tyr Phe Ser Gln Glu
            100   105   110

Val Asn Ile Ser Leu His Tyr Gln Lys Asp Glu Glu Pro Leu Phe Gln
        115   120   125

Leu Lys Lys Val Arg Ser Val Asn Ser Leu Met Val Ala Ser Leu Thr

130 135 140

Tyr Lys Asp Lys Val Tyr Leu Asn Val Thr Thr Asp Asn Thr Ser Leu
145 150 155 160

Asp Asp Phe His Val Asn Gly Gly Glu Leu Ile Leu Ile His Gln Asn
165 170 175

Pro Gly Glu Phe Cys Val Leu
180

<210> 103
<211> 131
<212> PRT
<213> Artificial Sequence

<220>
<223> OX40L soluble domain

<400> 103

Ser His Arg Tyr Pro Arg Ile Gln Ser Ile Lys Val Gln Phe Thr Glu
1 5 10 15

Tyr Lys Lys Glu Lys Gly Phe Ile Leu Thr Ser Gln Lys Glu Asp Glu
20 25 30

Ile Met Lys Val Gln Asn Asn Ser Val Ile Ile Asn Cys Asp Gly Phe
35 40 45

Tyr Leu Ile Ser Leu Lys Gly Tyr Phe Ser Gln Glu Val Asn Ile Ser
50 55 60

Leu His Tyr Gln Lys Asp Glu Glu Pro Leu Phe Gln Leu Lys Lys Val
65 70 75 80

Arg Ser Val Asn Ser Leu Met Val Ala Ser Leu Thr Tyr Lys Asp Lys
85 90 95

Val Tyr Leu Asn Val Thr Thr Asp Asn Thr Ser Leu Asp Asp Phe His
100 105 110

```
Val Asn Gly Gly Glu Leu Ile Leu Ile His Gln Asn Pro Gly Glu Phe
        115                 120                 125


Cys Val Leu
    130


<210>  104
<211>  128
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  OX40L soluble domain (alternative)

<400>  104

Tyr Pro Arg Ile Gln Ser Ile Lys Val Gln Phe Thr Glu Tyr Lys Lys
1               5                   10                  15


Glu Lys Gly Phe Ile Leu Thr Ser Gln Lys Glu Asp Glu Ile Met Lys
            20                  25                  30


Val Gln Asn Asn Ser Val Ile Ile Asn Cys Asp Gly Phe Tyr Leu Ile
        35                  40                  45


Ser Leu Lys Gly Tyr Phe Ser Gln Glu Val Asn Ile Ser Leu His Tyr
    50                  55                  60


Gln Lys Asp Glu Glu Pro Leu Phe Gln Leu Lys Lys Val Arg Ser Val
65                  70                  75                  80


Asn Ser Leu Met Val Ala Ser Leu Thr Tyr Lys Asp Lys Val Tyr Leu
                85                  90                  95


Asn Val Thr Thr Asp Asn Thr Ser Leu Asp Asp Phe His Val Asn Gly
            100                 105                 110


Gly Glu Leu Ile Leu Ile His Gln Asn Pro Gly Glu Phe Cys Val Leu
        115                 120                 125


<210>  105
```

<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain for 008

<400> 105

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30

Thr Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Asp Arg Tyr Ser Gln Val His Tyr Ala Leu Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser
115 120

<210> 106
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> variable light chain for 008

<400> 106

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Pro Val Thr Pro Gly

1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
20 25 30

Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Ala Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Tyr
85 90 95

Tyr Asn His Pro Thr Thr Phe Gly Gln Gly Thr Lys
100 105

<210> 107
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain for 011

<400> 107

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Thr Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ser Ile Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Arg Lys Gly
50 55 60

```
Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu Gln
65                  70                  75                  80


Met Asn Asn Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
                85                  90                  95


Asp Arg Tyr Phe Arg Gln Gln Asn Ala Phe Asp Tyr Trp Gly Gln Gly
            100                 105                 110


Thr Leu Val Thr Val Ser Ser Ala
        115                 120


<210>  108
<211>  108
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  variable light chain for 011

<400>  108

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Pro Val Thr Pro Gly
1               5                   10                  15


Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
            20                  25                  30


Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Ala Gly Gln Ser
        35                  40                  45


Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
    50                  55                  60


Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65                  70                  75                  80


Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Tyr
                85                  90                  95


Tyr Asn His Pro Thr Thr Phe Gly Gln Gly Thr Lys
```

100 105

<210> 109
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain for 021

<400> 109

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Arg Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Val Ile Ser Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Asp Arg Tyr Ile Thr Leu Pro Asn Ala Leu Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser
115 120

<210> 110
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> variable light chain for 021

<400> 110

```
Asp Ile Gln Met Thr Gln Ser Pro Val Ser Leu Pro Val Thr Pro Gly
1               5                  10                  15


Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
            20                  25                  30


Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
        35                  40                  45


Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
    50                  55                  60


Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65                  70                  75                  80


Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Tyr
                85                  90                  95


Lys Ser Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys
            100                 105
```

<210> 111
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain for 023

<400> 111

```
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val His Pro Gly Gly
1               5                  10                  15


Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe Ser Ser Tyr
            20                  25                  30


Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
        35                  40                  45
```

```
Ser Ala Ile Gly Thr Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Met
    50                  55                  60


Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu
65                  70                  75                  80


Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
                85                  90                  95


Arg Tyr Asp Asn Val Met Gly Leu Tyr Trp Phe Asp Tyr Trp Gly Gln
            100                 105                 110


Gly Thr Leu Val Thr Val Ser Ser
        115                 120


<210>  112
<211>  108
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  variable light chain for 023

<400>  112

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1               5                   10                  15


Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
            20                  25                  30


Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
        35                  40                  45


Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
    50                  55                  60


Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65                  70                  75                  80


Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Pro
```

85 90 95

Ala Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 113
<211> 119
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region

<400> 113

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Val Met His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Asn Tyr Tyr Gly Ser Ser Leu Ser Met Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Ser Val Thr Val Ser Ser
115

<210> 114
<211> 108
<212> PRT

<213> Artificial Sequence

<220>
<223> light chain variable region

<400> 114

Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly
1   5   10   15

Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr
            20   25   30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
        35   40   45

Tyr Tyr Thr Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
    50   55   60

Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Gln
65   70   75   80

Glu Asp Ile Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro Trp
                85   90   95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
            100   105

<210> 115
<211> 121
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region

<400> 115

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1   5   10   15

Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Tyr Thr Phe Lys Asp Tyr
            20   25   30

```
Thr Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
        35                  40                  45


Gly Gly Ile Tyr Pro Asn Asn Gly Gly Ser Thr Tyr Asn Gln Asn Phe
    50                  55                  60


Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65                  70                  75                  80


Met Glu Phe Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
                85                  90                  95


Ala Arg Met Gly Tyr His Gly Pro His Leu Asp Phe Asp Val Trp Gly
            100                 105                 110


Ala Gly Thr Thr Val Thr Val Ser Pro
        115                 120


<210>  116
<211>  108
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  light chain variable region

<400>  116

Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Leu Gly
1               5                   10                  15


Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Gly Ala Ala
            20                  25                  30


Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
        35                  40                  45


Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Thr Gly
    50                  55                  60


Gly Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Asn Val Gln Ser
```

65 70 75 80

Glu Asp Leu Thr Asp Tyr Phe Cys Gln Gln Tyr Ile Asn Tyr Pro Leu
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> 117
<211> 122
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region of humanized antibody

<400> 117

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Ser Met His Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Met
35 40 45

Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe
50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80

Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys
85 90 95

Ala Asn Pro Tyr Tyr Asp Tyr Val Ser Tyr Tyr Ala Met Asp Tyr Trp
100 105 110

Gly His Gly Thr Ser Val Thr Val Ser Ser
115 120

<210> 118
<211> 122
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region of humanized antibody

<400> 118

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
35 40 45

Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe
50 55 60

Lys Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Val Ser Thr Ala Tyr
65 70 75 80

Leu Gln Ile Ser Ser Leu Lys Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Asn Pro Tyr Tyr Asp Tyr Val Ser Tyr Tyr Ala Met Asp Tyr Trp
100 105 110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 119
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region of humanized antibody

<400> 119

```
Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Arg
1               5                  10                  15


Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Thr Ala
            20                  25                  30


Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
        35                  40                  45


Tyr Ser Ala Ser Tyr Leu Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
    50                  55                  60


Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala
65                  70                  75                  80


Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Ser Thr Pro Arg
                85                  90                  95


Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
            100                 105


<210>  120
<211>  107
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  light chain variable region of humanized antibody

<400>  120

Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Arg
1               5                  10                  15


Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Thr Ala
            20                  25                  30


Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
        35                  40                  45


Tyr Ser Ala Ser Tyr Leu Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
```

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala
65 70 75 80

Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Ser Thr Pro Arg
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 121
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region of humanized antibody

<400> 121

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Glu
1 5 10 15

Ser Leu Lys Leu Ser Cys Glu Ser Asn Glu Tyr Glu Phe Pro Ser His
20 25 30

Asp Met Ser Trp Val Arg Lys Thr Pro Glu Lys Arg Leu Glu Leu Val
35 40 45

Ala Ala Ile Asn Ser Asp Gly Gly Ser Thr Tyr Tyr Pro Asp Thr Met
50 55 60

Glu Arg Arg Phe Ile Ile Ser Arg Asp Asn Thr Lys Lys Thr Leu Tyr
65 70 75 80

Leu Gln Met Ser Ser Leu Arg Ser Glu Asp Thr Ala Leu Tyr Tyr Cys
85 90 95

Ala Arg His Tyr Asp Asp Tyr Tyr Ala Trp Phe Ala Tyr Trp Gly Gln
100 105 110

Gly Thr Leu Val Thr Val Ser Ala
        115   120

<210> 122
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region of humanized antibody

<400> 122

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1   5   10   15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Glu Tyr Glu Phe Pro Ser His
            20   25   30

Asp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
        35   40   45

Ala Ala Ile Asn Ser Asp Gly Gly Ser Thr Tyr Tyr Pro Asp Thr Met
    50   55   60

Glu Arg Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65   70   75   80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
                85   90   95

Ala Arg His Tyr Asp Asp Tyr Tyr Ala Trp Phe Ala Tyr Trp Gly Gln
            100   105   110

Gly Thr Met Val Thr Val Ser Ser
        115   120

<210> 123
<211> 111
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region of humanized antibody

<400> 123

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
1   5   10   15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Lys Ser Val Ser Thr Ser
            20   25   30

Gly Tyr Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
        35   40   45

Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
    50   55   60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
65   70   75   80

Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ser Arg
                85   90   95

Glu Leu Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
            100   105   110

<210> 124
<211> 111
<212> PRT
<213> Artificial Sequence

<220>
<223> light chain variable region of humanized antibody

<400> 124

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1   5   10   15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Lys Ser Val Ser Thr Ser
            20   25   30

Gly Tyr Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro

```
        35                  40                  45

Arg Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
    50                  55                  60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65                  70                  75                  80

Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln His Ser Arg
                85                  90                  95

Glu Leu Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
            100                 105                 110
```

<210> 125
<211> 138
<212> PRT
<213> Artificial Sequence

<220>
<223> heavy chain variable region

<400> 125

```
Met Tyr Leu Gly Leu Asn Tyr Val Phe Ile Val Phe Leu Leu Asn Gly
1               5                   10                  15

Val Gln Ser Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln
            20                  25                  30

Pro Gly Gly Ser Met Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
        35                  40                  45

Ser Asp Ala Trp Met Asp Trp Val Arg Gln Ser Pro Glu Lys Gly Leu
    50                  55                  60

Glu Trp Val Ala Glu Ile Arg Ser Lys Ala Asn Asn His Ala Thr Tyr
65                  70                  75                  80

Tyr Ala Glu Ser Val Asn Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser
                85                  90                  95
```

```
Lys Ser Ser Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr
            100                 105                 110


Gly Ile Tyr Tyr Cys Thr Trp Gly Glu Val Phe Tyr Phe Asp Tyr Trp
        115                 120                 125


Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
    130                 135


<210>  126
<211>  126
<212>  PRT
<213>  Artificial Sequence

<220>
<223>  light chain variable region

<400>  126

Met Arg Pro Ser Ile Gln Phe Leu Gly Leu Leu Leu Phe Trp Leu His
1               5                   10                  15


Gly Ala Gln Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
            20                  25                  30


Ala Ser Leu Gly Gly Lys Val Thr Ile Thr Cys Lys Ser Ser Gln Asp
        35                  40                  45


Ile Asn Lys Tyr Ile Ala Trp Tyr Gln His Lys Pro Gly Lys Gly Pro
    50                  55                  60


Arg Leu Leu Ile His Tyr Thr Ser Thr Leu Gln Pro Gly Ile Pro Ser
65                  70                  75                  80


Arg Phe Ser Gly Ser Gly Ser Gly Arg Asp Tyr Ser Phe Ser Ile Ser
                85                  90                  95


Asn Leu Glu Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp
            100                 105                 110


Asn Leu Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
```

115 120 125