



(51) International Patent Classification:

C12Q 1/34 (2006.01) G01N 33/573 (2006.01)
G01N 33/52 (2006.01) G01N 33/574 (2006.01)

(21) International Application Number:

PCT/US2011/024658

(22) International Filing Date:

11 February 2011 (11.02.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/303,509 11 February 2010 (11.02.2010) US
61/412,364 10 November 2010 (10.11.2010) US

(71) Applicant (for all designated States except US): **INDIANA UNIVERSITY RESEARCH AND TECHNOLOGY CORPORATION** [US/US]; 351 West 10th Street, Indianapolis, IN 46202 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **XU, Yan** [US/US];
11580 Winding Wood Drive, Indianapolis, IN 46235 (US).

(74) Agent: **EMANUELE, John, J.**; Baker & Daniels LLP,
300 North Meridian Street, Suite 2700, Indianapolis, IN
46204 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,

[Continued on next page]



(54) Title: **PLA₂ ACTIVITY AS A MARKER FOR OVARIAN AND OTHER GYNECOLOGIC CANCERS**

FIG. 1 E

(57) Abstract: Materials and methods are provided for diagnosis, monitoring, and personalized treatments of gynecological cancers. The methods comprise determining levels of PLA₂ activity with a patient sample and using elevated levels of this activity which correlate with diseases such as epithelial ovarian cancer (EOC) as a marker for pathology. These methods include assaying for PLA₂ activity within tissue, ascites, blood, and other tissue forms by exposing the patient sample to a fluorogenic compound such as DBPC. The methods disclosed herein further include correlating the fluorogenic detection with a disease state in the patient, including diseases such as gynecological cancers, such as EOC. The methods comprise determining levels of total PLA₂ activity, and of specific isoforms of PLA₂ such as iPLA₂, iPLA₂β, cPLA₂, among other isoforms.



ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

**PLA₂ ACTIVITY AS A MARKER FOR OVARIAN AND OTHER
GYNECOLOGIC CANCERS**

PRIORITY CLAIM

[0001] This application claims the benefit of United States Provisional Patent application Numbers 61/303,509 filed on February 11, 2010, and 61/412,364 filed on November 10, 2010, each of which is incorporated herein by reference in its entirety as if each were incorporated by reference individually.

FIELD OF THE DISCLOSURE

[0002] The present disclosure relates generally to cancer. More particularly, the present disclosure relates to Phospholipase A₂ (PLA₂) as a novel marker and as a target for diagnosing, evaluating and treating ovarian and other gynecologic cancers.

BACKGROUND

[0003] Gynecological cancers, and specifically ovarian, uterine, and cervical cancers, cause more than 26,000 deaths annually in the United States. A majority of epithelial ovarian cancer (EOC) patients present with late-stage disease. Curative treatment for late-stage EOC, and in particular for refractory and drug-resistant EOC, is often ineffective resulting in high mortality rates from EOC.

[0004] In general, there is not an acceptable method for screening patients for EOC. Current screening methods, such as ultrasound, are inadequate for early stage detection and in some cases for staging of EOC patient. Further, current markers for EOC, in general, are poor distinguishers for distinguishing between benign and malignant tumor development. Hence, development of markers for monitoring disease progress which also function as better targets for personalized treatment is urgently needed for a majority of EOC patients.

[0005] Therefore, a method and marker for better diagnosis, monitoring of gynecological cancer progression, and application in personalized gynecological cancer treatment are desirable. Some aspects of the present disclosure, disclosed herein, address these needs.

SUMMARY OF THE DISCLOSURE

[0006] Some embodiments of the disclosure include methods for evaluating a disease such as EOC including detecting and/or following the course of a disease including for example diagnosing, prognosing and evaluating the efficacy of treating a given disease, these methods may include the steps of measuring PLA₂ activity in a patient sample and providing information

associated with correlating the PLA₂ activity measured in said step of measuring, with a PLA₂ activity value indicative of a disease. Some embodiments further include the step of comparing the values measured in a given sample with values indicative of various diseases and concluding that a given sample includes evidence of disease. Some embodiments of the methods disclosed herein include measuring the activity of PLA₂ comprising, for example, iPLA₂. Some further embodiments include methods of correlating a PLA₂ activity value with a PLA₂ activity value indicative of a form of gynecological cancer. In further embodiments the form of gynecological cancer is EOC.

[0007] Some embodiments further include the step of obtaining the patient sample from a patient. In some embodiments of the disclosure, the patient sample may be selected from the group of sample consisting of tissue samples, cells, blood, bodily fluids, cellular fluids, a cellular fluids, discharges or fluids within or otherwise associated or produced by tumors, cysts or other growths and the like.

[0008] In some further embodiments of the methods disclosed herein, the step of measuring PLA₂ activity in a patient sample involves using a fluorogenic compound, including but not limited to the fluorogenic compound is DBPC. Still other embodiments include the use of still other compound that produces a detectable signal and can be used to measure or at least estimate PLA₂ activity in a given sample. In still other embodiments of the disclosed methods, the measuring step involves the use of radioactive compounds. Some embodiments may include a further separation steps and/or sample preparations to increase the sensitivity and or reproducibility of the activity assay. Still another embodiment includes normalizing the level of PLA₂ activity measured in a given sample to the number of cells associated with the cells or to the level of protein in the sample.

[0009] In even further embodiments of the methods disclosed herein, the step of providing information comprises providing information correlating PLA₂ activity in range of activity that is demonstratively higher in sample for a patient with a form of GYN cancer such as EOC. In other embodiments of the methods disclosed herein, the step of providing information comprises providing information correlating PLA₂ activity in the range of about 1.5 higher than that detected in similar but healthy tissue of the same type. In still other embodiments the level of activity that correlates with disease may be about 2x or higher than normal activity measured in sample from non-cancerous sources include healthy samples or at least benign samples.

[0010] Still other embodiments of the methods provided herein, include the step of introducing a compound to the patient which inhibits activity of at least one isoform of PLA₂. In some embodiments the isoforms of PLA₂ in the sample includes at least one of the following sPLA₂, cPLA₂ and iPLA₂. Inhibitors that can be used include, but are not limited to, BEL. Some embodiments may include the steps of conducting a given PLA₂ assay with and without the presence of a given PLA₂ inhibitor and using any difference observed in the assays to assign at least a portion of the reduced activity to the forms of PLA₂ thought to be sensitive to the specific inhibitor used in the assay.

[0011] Further embodiments of the methods described herein includes a method of detecting PLA₂ activity within a sample comprising the steps of introducing a fluorogenic compound to a sample, and detecting a signal produced by the fluorogenic compound in the presence of the sample. In some embodiments of the method described herein the sample is a tissue and the method further includes the step of fixing the tissue.

[0012] In yet further embodiments of the methods disclosed herein, the tissue sample is homogenized and the fluorogenic compound is introduced to the supernant resulting from the homogenization of the tissue sample.

[0013] Still further embodiments of the methods include the step of correlating the signal value, detected in the step of detecting, with a disease. Some embodiments of these methods include the disease being a form of cancer. In some embodiments, the form of cancer is EOC.

[0014] Some embodiments include identifying a using a marker for diseases such, but not necessarily limited to EOC. These markers include phospholipase A₂ (PLA₂) activity that is demonstratively higher in samples such as fluids, tissues, blood, cells and like associated with disease such as EOC than it is in samples of healthy sources or samples from benign tumors.

[0015] In some embodiments demonstratively higher activity indicative of a disease state is at least 1.5 or in some embodiments at least 2 fold higher in a sample from a patient with a disease then it is in a patient that is without disease or that presents with a benign tumor.

[0016] In some embodiments the identification of the maker is indicative of cancer especially GYN cancers such as EOC. In some embodiments the marker can be used to diagnose a disease, prognosticate on the course of the disease, following the course of a disease or to evaluate the effectiveness of a treatment of a disease. In some embodiment the maker is activity

that due primarily to elevated iPLA₂β activity. In some embodiments values of PLA₂ activity used as markers for disease are used to personalize the treatment of a patient.

[0017] These and other features, aspects, and advantages of the present invention will become better understood with reference to the following drawings, descriptions and claims.

BRIEF DESCRIPTION OF THE FIGURES

[0018] The above-mentioned aspects of the present disclosure and the manner of obtaining them will become more apparent, and aspects thereof will be better understood by reference to the following description of the embodiments of the disclosure, taken in conjunction with the accompanying drawings, figures, schemes, graphs, charts, and the like, wherein:

[0019] **FIG. 1A.** Photomicrographs of para-nuclear PLA₂ activity in cells from cancerous tissue.

[0020] **FIG. 1B.** Photomicrographs of para-nuclear PLA₂ activity only in the surface cells in benign tumors.

[0021] **FIG. 1C.** Photomicrographs of PLA₂ activity in the stromal portions of from normal tissue (non-cellular).

[0022] **FIG. 1D.** Photomicrographs of cells treated with an inhibitor of cPLA₂ and/or iPLA₂ (panels b, d) and an inhibitor of sPLA₂ (panel c).

[0023] **FIG. 1E.** Dot plots (panel a) and a histogram (panel b) of fluorescent intensity normalized to cell count.

[0024] **FIG. 2A.** Arachidonic acid (AA) concentrations in non-malignant (N; n=9) and EOC (n=8) ascites samples.

[0025] **FIG. 2B.** Histograms of fluorescent intensity (PLA₂ activity) measured in the blood sample from subjects: healthy controls, patients with benign gynecologic diseases, and patients with ovarian cancer Total PLA₂ activities in blood samples from healthy (n=7) subjects, benign (n=8) subjects, and subjects diagnosed with EOC (n=5).

[0026] **FIG. 2C.** PLA₂ activities in cell-free ascites samples mentioned in **FIG. 2A**.

[0027] **FIG. 2D.** Histograms showing PLA₂ activities in the exosomes and the supernatant (super) portions of ascites samples. The dark bars: 1, background fluorescence; 2. sPLA₂ positive control; 3. TA-PC treated sPLA₂. **: p<0.01. * p<0.05.

[0028] **FIG. 3A.** PLA₂ activities in mouse tissues measured in mice that were injected with A2780 human EOC cells and PLA₂ activities were analyzed in tumors: a. no DBPC; b. with DBPC; c. DAPI; d. merged.

[0029] **FIG. 3B.** The PLA₂ activities from tumors obtained from control (a) and BEL-treated (b) mice. The tumors in BEL-treated mice were significantly smaller and the PLA₂ activities were reduced.

[0030] **FIG. 3C.** PLA₂ activities detected in tumors derived from human SKOV3 (C) and mouse ID-8 cells (D) EOC cells.

[0031] **FIG. 3D.** PLA₂ activities detected in tumors derived from human SKOV3 (C) and mouse ID-8 cells (D) EOC cells.

[0032] **FIG. 3E.** Normal mouse kidney tissues mainly had sPLA₂-like activity, which was not associated with cell nucleus and minimally affected by BEL. The numbers in all figures are the quantified PLA₂ activity/cell. Representative results are shown

[0033] **FIG. 4A.** Shows a time dependent curve demonstrating fluorescent intensity having a linear relation with time

[0034] **FIG. 4B.** Shows another time dependent curve demonstrating fluorescent intensity having a linear relation with time

[0035] **FIG. 4C.** Is a standard curve demonstrating fluorescent intensity having a linear relation with DBPC substrate concentration.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0036] For the purposes of promoting an understanding of the principles of the novel technology, reference will now be made to the preferred embodiments thereof, and specific language will be used to describe the same. It will nevertheless be understood that no limitation of the scope of the novel technology is thereby intended, such alterations, modifications, and further applications of the principles of the novel technology being contemplated as would normally occur to one skilled in the art to which the novel technology relates are within the scope of this disclosure and what is claimed.

[0037] Unless explicitly or implicitly stated otherwise the term, 'about' as used herein means plus or minus 10 percent. For example about 1.1 encompasses the values between 0.9 and 1.1.

[0038] Unless explicitly or implicitly stated otherwise the term, 'demonstratively' as used to refer to a property that is integral to given sample and that can be reproduced with sufficient precision and accuracy to show or illustrate the similarity or difference between at least two different portions, samples and the like.

[0039] Phospholipase A₂s (PLA₂s) are enzymes commonly found in mammalian tissue. PLA₂s include several protein families, including secreted phospholipase A₂ (sPLA₂), cytosolic phospholipase A₂ (cPLA₂), and Ca²⁺ independent PLA₂ (iPLA₂). All PLA₂s catalyze hydrolysis of phospholipid substrates to generate a lysophospholipid (**LPL**) and a free fatty acid (**FA**), *e.g.*, arachidonic acid (**AA**). **AA** can be oxygenated to a variety of bioactive eicosanoids. Phosphatidylcholine (**PC**) is an abundant lipid substrate in cell membranes and lysophosphatidylcholin (**LPC**) is a more common product of PLA₂s. LPC is convertible to Lysophosphatidic acid (**LPA**) by autotaxin (**ATX**). Additionally, certain PLA₂s, including sPLA₂ and calcium independent iPLA₂ isoforms can utilize phosphatidic acid (**PA**) as a substrate for directly producing LPA. The compounds, LPA and AA are shown to be important lipid mediators in chemotaxis and/or chemokinesis.

[0040] Aberrant expression of some sPLA₂s and cPLA₂ has been shown in some cancer types (not inclusive of EOC). To date, only the sPLA₂ isoform, sPLA₂-XII, has shown potential as a possible predictive indicator of patient outcomes. However, currently there exists no comparison data between malignant and benign/normal tissues.

[0041] In regard to EOC, specifically, mRNA expression levels of 19 different PLA₂ isoforms (sPLA₂ and cPLA₂s) have demonstrated decreased or unchanged transcriptional levels in EOC versus non-tumoral immortalized (EONT) cells.

[0042] iPLA₂s, calcium-independent phospholipase A₂ beta, the Group VIA phospholipase A₂, are intracellular enzymes that do not require Ca²⁺ for their catalytic activity. iPLA₂s are, in general, known to be active in phospholipid remodeling, signal transduction, and cell proliferation and apoptosis.

[0043] Various iPLA₂ isoforms use **PA** as a substrate and produce LPA. LPA and AA are shown to be important lipid mediators in chemotaxis and/or chemokinesis. Further, iPLA₂ activity is known to regulate **LPA** and other lipid concentrations in tumor microenvironment.

[0044] Lysophosphatidic acid (**LPA**) is a bioactive lipid with multiple functions. Epidemiological, animal, and cell culture studies indicate that LPA plays a role in EOC

development. Over-expression or down-regulation of LPA receptors 1, 2, and 3 in several human EOC cell lines and *in vivo* studies, indicate that LPA receptors are involved in EOC development.

[0045] LPA activity is suppressible in various manners including inhibiting its production, increasing its degradation or conversion, preventing receptor occupancy, or and interfering with distal signaling pathways.

[0046] LPA is producible by the action of lysophospholipase(s) D (lyso-PLD), *e.g.*, autotaxin (ATX), or via phospholipase(s) A₂ (PLA₂), as discussed above.

[0047] The activation of iPLA₂β in non-apoptotic EOC cells occurs via a laminin-β₁-integrin-caspase 3 pathway resulting in LPA and AA increase from EOC cells. Also, iPLA₂ activity increases (in association with EOC) as disclosed herein, are not reflective of increased iPLA₂ expression levels in EOC and/or other GYN cancer tissues.

[0048] As detailed above, resulting LPA acts as an “oncolipid” in EOC. Unexpectedly, as disclosed herein the activity of PLA₂ in EOC tissues is increased compared to its activity in benign disease and/or normal tissues without a correlated increase in PLA₂ expression levels. The activity of the PLA₂ enzymes (there are over 20 isoforms of phospholipase A₂ enzymes) taught and described herein, are disclosed as utilizable in the assay and methods disclosed herein as useful markers for diagnosing, prognosis, and monitoring EOC progression, as well as a target for personalized therapeutics in treating EOC.

[0049] As disclosed in the Examples section provided herein, PLA₂ activities have been assessed in cancer vs. normal tissues and the differences in activity between these types of tissues are disclosed herein as usable and valuable in providing an assay for cancer diagnosis and monitoring. Similarly, PLA₂ activities have been assessed in EOC vs. tissue from benign tumors and usable differences between PLA₂ activities measured between these different tissues have been demonstrated to exist.

[0050] Further, the disclosure provided herein discloses PLA₂ activity as a robust and useful marker that can be used to distinguishing malignant EOC, and other GYN cancer tissues including, from benign and/or normal tissues. With reference to **FIG. 1E**, data is presented demonstrating PLA₂ activity in malignant and benign tissues provides distinguishability. Current markers utilized for EOC (including Ca¹²⁵) lack sufficient differentiation capabilities for distinguishing between benign and malignant tumor development. This common issue suggests

that the current markers' biological functions are similar, resulting in difficulty in distinguishing between benign and malignant tumor development. The methods and assays disclosed and described herein provide a further advantage over current markers by providing an assay for distinguishing malignant EOC, and other GYN cancer tissues, from benign and/or normal tissues based on PLA₂ activity.

[0051] As disclosed and described herein, PLA₂ activity provides a valuable marker with utility in personalized chemotherapy. As is noted herein, this disclosure provides a method corresponding to greater than 30 percent of all EOC patients (no other genetic changes, other than p53 mutations, is known to occur with an incidence of greater than 30% in EOC patients). Thus, personalized and targeted therapy is the promising goal for EOC treatment.

[0052] Further, PLA₂s are involved in cellular processes, sometimes associated in other forms of cancers. The major effects of iPLA₂, for example, in EOC cells are cell migration and invasion, two key steps of tumor metastasis, with a relatively less prominent effect on cell proliferation. Thus, the present embodiments further provides a clinical value (in utilizing of PLA₂ activity as disclosed herein) in cancer types of than EOC. For example, the instant disclosure for analysis of PLA₂ activity as a marker for metastases has a prognostic value in both EOC and other forms of GYN cancers. Hence, it is within the present disclosure to utilize the assay involving PLA₂ activity, disclosed herein, for distinguishing between metastases and primary EOC.

[0053] Metastasis is the major cause for deaths in solid cancer patients. As such, the assay disclosed herein provides a valuable marker and prognosis tool. Further, the methods associated with PLA₂ activity, disclosed and described herein, provide a valuable and necessary marker adapted for early detection in the diagnosis of GYN diseases.

[0054] One example demonstrating the value and novelty of the disclosed embodiments herein is demonstrated in early detection of Type II EOC. Recent data indicates that there exists two major types of EOC, Type I and Type II. Type I tumors are, characteristically speaking, slow growing and include low-grade micropapillary serous carcinoma, mucinous, and endometrioid carcinomas. Type I tumors are also genetically stable and are characterized by mutations in a number of different genes including KRAS, BRAF, pTEN, and beta-catenin. Type II tumors, however, are rapidly growing and highly aggressive neoplasms for which well-defined precursor lesions have not been described. Type II tumors include high-grade serous

carcinoma, malignant mixed mesodermal tumors (carcinosarcomas), and undifferentiated carcinomas and have a high level of genetic instability and are characterized by a mutation of TP53. This method and assay disclosed herein, helps to explain why current screening techniques, aimed at detecting stage I disease, have not been effective given that Type II tumors are rarely confined to the ovary and are not derived from Type I EOC. Current screening approaches for early EOC detection have utilized stage I cancers as model in early markers development. However, specifically in regard to Type II EOC, such screening approaches and models are flawed for the reasons discussed above. The diagnostic methods disclosed herein, however, provide an improved, novel, and unique method allowing for earlier diagnosis of Type II EOC.

[0055] Ascitic fluids, as disclosed herein, represents EOC tumor microenvironment and/or peritoneal washings (for those patients who do not develop ascites or are still at a relatively early stage of disease development) which can be obtained through minimally invasive ultrasound guided procedures. Accordingly, the PLA₂ activity based methods provided herein disclose a ready and less invasive means for diagnosing and monitoring the progress of these diseases.

[0056] Additionally, the single blood marker disclosed herein may be used as part of an array of diagnostic tools and has important value to existing EOC markers such as the markers marketed by Quest Diagnostics (OVA1, Ca¹²⁵ and HE4).

[0057] Further, tissue based PLA₂ activity assays, as disclosed herein, offer information related to tumor and host cell histology, and potentially better separation for subjects in different groups. DBPC is used in assays, disclosed and described herein, for *in vitro* diagnostic applications. Additionally, development of infrared fluorescence (NIRF) PLA₂ substrates suitable for *in vivo* tumor imaging provide for *in vivo* EOC imaging. Utilization of the PLA₂ activity assays, disclosed herein, for monitoring PLA₂ activity provides not only available diagnostic tool, but also an efficient means of monitoring the efficacy of various treatments including drug treatments.

[0058] The data presented herein provides and supports a method of using PLA₂ activity as a new marker for studying a disease, such as EOC, including methods of diagnosing the diseases, predicting the course of the diseases, evaluating the effectiveness of treatments for the diseases and formulating personalized treatment regimes for this disease. As discussed herein,

PLA₂ activity provides a useful marker for diagnosis, prognosis, monitoring, and/or predicting EOC and/or other GYN cancers. These methods are especially useful when PLA₂ activity is normalized to an inherent standard such as the number of cells in a sample of tissue. Still other means of normalizing PLA₂ activity include normalizing data based on the protein content of a given sample and the like. The data indicated herein further provides that PLA₂, including iPLA₂ alone or in combination with other PLA₂s provides a useful target for the treatment of EOC. Further, PLA₂-related activities, disclosed herein, are highly significant in EOC management and treatment decision making, as well as in designing and testing new therapeutics. Further, it should be understood that although genetic, epigenetic, and proteomic studies regarding EOC have been conducted, those studies have focused on expression at either the RNA or protein levels, not a direct focus on PLA₂ activity. Thus, a focus directly on PLA₂ as a marker for ovarian and other gynecological forms of cancer, as disclosed herein, represents a new and highly relevant and useful method for the diagnosis, monitoring, and targeted treatment of gynecological cancers such as EOC.

EXPERIMENTAL

Example 1. Detection Assays for PLA₂ activity.

[0059] Classical methods of assaying PLA₂ activity, in general, involve the use of a radioactive-labeled phosphatidylcholine (PC) substrate. Cell or tissue lysates are incubated in a buffer with or without calcium (assays specific for a form of iPLA₂ are performed without calcium and include 1 mM EDTA) and the substrates [dipalmitoyl phosphatidylcholine (DPPC), and 1-palmitoyl-2-[1-¹⁴C] palmitoyl-*sn*-glycero-3-phosphocholine (300,000 cpm/assay)]. Post incubation, the cleaved and ¹⁴C-labeled fatty acid (FA) is resolved using thin-layer-chromatography (TLC). The radioactivity associated with the cleaved FA is counted and calculated. However, this standard assay is not convenient or practical for high throughput PLA₂ activity assays.

[0060] According to the instant disclosure, a PLA₂ activity assay practical for high throughput utilizing DBPC (available from Echelon Biosciences Incorporated, Salt Lake City, UT 84108 Echelon Biosciences Incorporated, Salt Lake City, UT 84108) is disclosed. DBPC is a fluorogenic Dabcyl- and BODIPY-containing PC which functions as a PLA₂ substrate producing fluorescence. Usages of DBPC has the advantage of *not* involving radioactivity or a time-consuming and labor intensive TLC separation step. Further, the assay disclosed herein has

been adapted for use in frozen tissue sections with cell nuclei co-stained to examine the types of cells, having significant PLA₂ activity, and tissue histology, simultaneously.

Example 2: PLA₂ Activity Assay Using Tissue.

[0061] PLA₂ activity from human gynecologic (GYN) cancers (including both EOC and endometrial), benign GYN disease, and normal tissue samples were tested. Six samples from each of these groups were assayed in accordance with the instant disclosure.

[0062] Tissue samples were obtained and frozen in order to preserve enzymatic activity. The tissue samples (of average size 6-10 μ m) were fixed using acetone and methanol at -20°C according to typical fixation methods known within the art. As explained above, the substrate DBPC is a fluorogenic Dabcyl- and BODIPY – containing PC which represents PLA₂ activity with fluorescence.

[0063] The PLA₂ activity in the frozen tissue sections was quantified. Quantification of the PLA₂ signals in frozen tissue sections used the software (MetaMorph). The total integrated signaling intensity, with the background subtracted (tissues processed the same way in the absence of the PLA₂ substrate), were captured and measured, and then normalized based on cell number or per cell ratio (the cell numbers were counted by DAPI-stained nuclei).

[0064] Referring now to **FIG. 1**, the activity of PLA₂ is shown illustrating PLA₂ activity being elevated in cancer (**FIG. 1A**) vs. benign (**FIG. 1B**) and normal (**FIG. 1C**) GYN tissue samples. Representative tissue PLA₂ activities are shown. The first two rows in **FIGS. 1A to C**, (each comprising four panels labeled as lower case letters a to d) present: a.) no DBPC substrate; b.) addition of DBPC; c.) DAPI staining of cell nuclei; and d.) an overlay the DAPI and PLA₂ activity staining. Referring now to **FIG. 1D**, para-nuclear PLA₂ activities are demonstrated as being sensitive to inhibitors of iPLA₂ and cPLA₂ (e.g., AACOCF₃ and BEL) but not to a inhibitor of sPLA₂ (e.g., TA-PC) (the big bright spots were present in the absence of DBPC and thus were non-specific and were not a PLA₂ activity). Remaining with **FIG. 1D**, the numbers in each panel (a to d) are the activities expressed as fluorescent intensity/cell under each condition (as described herein).

[0065] Referring now to **FIG. 1E**. **FIG. 1E** presents PLA₂ activities measured in six samples from each of cancer (EOC and endometrial), benign GYN, and normal groups. Panel a of **FIG. 1E** presents individual values; whereas panel b of **FIG. 1E** presents mean and SD values (**, P< 0.01; *** P<0.001).

[0066] As shown herein, PLA₂ activities in EOC samples were significantly increased compared to those in either benign or normal tissues. The PLA₂'s activities/cell in the three groups of samples were 881+/-395; 276 +/- 58; and 55+/-16 for human cancer, benign, and normal tissues, respectively (**FIG. 1**). The results presented herein illustrate that high levels of para-nuclear PLA₂ activities in the cancer tissues were only observed in the ovarian and endometrial cancer tissues (**FIG. 1A**). For benign tumors, these activities were limited to the superficial cell layers of the benign tumors (**FIG. 1B**). For normal tissues, very little or no para-nuclear PLA₂ activity was observed (**FIG. 1C**). The long-thin bright lines detected in some normal tissues (**FIG. 1C**) are unlikely to be cellular PLA₂ activities, since they were insensitive to any of the PLA₂ inhibitors tested (data not shown). In contrast, cancer PLA₂ activities were sensitive to inhibitors for both cPLA₂ and iPLA₂ (e.g., AACOCF₃ and BEL) but not to TA-PC (an inhibitor of sPLA₂) (**FIG. 1D**). Thus the data presented herein illustrates that a majority (if not the total PLA₂ activity tested in tumor tissues) is derived from cPLA₂ and/or iPLA₂. It is worth noting that the individual values of PLA₂ activity from different groups were not overlapping, indicating this assays excellent separation and high specificity (**FIG. 1E**, panel a). As disclosed and described herein, a cut-off or determination value for identifying tumor tissues from both benign and normal tissues is approximately 400 (normalized fluorescent intensity values / cell). A determination or cut-off value of 400 (normalized fluorescent intensity values / cell), as disclosed herein, provides a 100% specificity in separating the various tissue states tested and described above.

Example 3: PLA₂ Activity Assay Using Tissue Homogenates.

[0067] Further, a tissue homogenization method, in which tissue lysate is utilized for measuring PLA₂ activity, is disclosed herein. The homogenization method described herein was compared against the current standard quantitative methods of tissue using a radioactive assay (as discussed above), for assessing quantification using tissue homogenates for measuring PLA₂ activity. As described herein, a tissue homogenization method suitable for PLA₂ activity analysis was developed.

[0068] According to the instant disclosure, frozen samples were pulverized and transferred to a microcentrifuge tube and mixed with 500μL lysis buffer. It is within the instant disclosure that samples may be pulverized in any known method (including using pulverizing instrumentation, bead ablation, and manual grinding). The lysis buffer comprised 10mM hepes,

pH 7.5, and 0.34 M sucrose with an additional 5 μ L of mammalian protease inhibitor cocktail obtained from Sigma-Aldrich, Inc. The tissue samples were next homogenized using tissue homogenization instrumentation (such as the Omni Tissue Homogenizer, Model No.: LR60902). Following lysis and homogenization, the samples were centrifuged at 16,000 x g, at a temperature of 4°C, for about 40 minutes. Following centrifugation the supernatants were transferred to a new eppendorf tube for assaying the PLA₂ activity.

[0069] PLA₂ activities measured using the standard radioactive methods (discussed above) were 1.76, 1.30, 0.52, and 0.57 [Fatty acid/(Fatty acid+PC)]% (normalized to the *same amount of protein (1 mg)* in each sample) for the two cancer samples and two normal samples, respectively. Taken together, these data show that EOC and possibly other forms of GYN cancers have elevated PLA₂ activities when compared to those of benign GYN and normal tissues.

Example 4: PLA₂ Activity in Ascites and LPA Levels in Plasma.

[0070] The method described herein, is also adaptable for use in identifying PLA₂ activity as a marker in body fluid. For reasons discussed in detail above, a body fluid PLA₂ activity marker provides a more convenient and clinically useful marker, than a genetic number marker or a marker based on RNA.

[0071] The PLA₂ activity detection method employed herein is simple and can be easily conducted in almost any clinical laboratory. In contrast, although LPA has been shown to be a potential EOC marker, because of obstacles such as low concentration in the blood, variable samples processing procedures, as well as the rather sophisticated (less clinic-friendly) and different electrospray ionization tandem mass spectrometry (ESI-MS/MS) methods used in it, has proven slow in moving to wide spread and accepted clinical use.

[0072] Referring now to **FIG. 2**, as reported herein, PLA₂ activity is increased, with a concurrent increase in PLA₂ direct and/or indirect products (such as LPC, LPA, and arachidonic acid (AA)) in EOC ascites and blood samples. Data measured with LPC and LPA have been published, and increased LPA in EOC blood samples has been confirmed in repeated and independent studies.

[0073] According to the present disclosure, PLA₂ activity in human ascites samples were collected and assayed for detection of DBPC derived fluorescence. According to the disclosed method, ascites samples (25 μ L per sample) were used in assessing PLA₂ activities. The

fluorescence detected (shown as total intensity in the accompanying figures) were approximately 2400 and 4,000 (wherein the numbers in the figures should be divided by 25, respectively, based on sample volume). The background fluorescence (intensities/ μL) in the human samples assayed, according to the method disclosed herein was observed to be relatively low (~ 40) and did not change in a considerable amount during a 24 hour incubation time.

[0074] According to the methods disclosed and described herein, human ascite samples were centrifuged at 3,000 g for 20 minutes at 4°C. The samples were then aliquoted and stored at -80°C. Ascite samples (25 μL of each) were suspended in 50 μL buffer [comprising 80 mM hepes (pH 7.4), 150 mM NaCl, 10 mM CaCl_2 , 4 mM Triton X-100, 60% glycerol, and 1 mg/mL BSA], respectively, and then mixed with 0.20 μg DBPC [dissolved in 50 μL DMSO/assay buffer (1:100)]. The total volume of each assay was approximately 200 μL . After incubation for 4 hrs, fluorescent intensities were measured at 485 nm/535 nm in a plate reader (Perkin Elmer Victor³V 1420 Multilabel Counter).

[0075] Referring now to **FIGS. 4A and B**, Two time-dependent curves are presented demonstrating fluorescence observed having a linear relationship with time over a 4 hour assay time period for different samples. The data in **FIG 4 A** fit to a line with the following values: $Y=230$, intercept of 7468 and a R squared value of 0.9559. The data in **FIG 4 B** fit to a line with the following values: $Y=352$, intercept of 25,328 and a R squared value of 0.7818.

[0076] With reference now to **FIG. 4C**, the method and assay disclosed herein is demonstrated as comprising good inter-day and intra-day precision with all assays performed. **FIG. 4C** presents a standard curve which correlates intensity value of assayed EOC ascite samples with DBPC concentrations used.

[0077] As shown by the methods described and disclosed herein, PLA_2 is associated with human EOC ascites and exosomes (from ascites and blood). Both cPLA_2 s and iPLA_2 s are cytosolic enzymes, only sPLA_2 s are secreted, data presented herein shows that elevated PLA_2 activity (which may be due to the activity of any of these three classes of PLA_2 s) can be detected in the cell-free blood and/or ascites samples from patients with EOC and/or other GYN cancers.

[0078] Exosomes are 40-100-nm diameter membrane vesicles released from multivesicular bodies (MVB) by intact cells and are thought to participate in intercellular signaling. Further, cell membrane lipids, including PC (a PLA_2 substrate) and sphingolipids (in particular ceramides) are prominent lipids within exosomes.

[0079] According to methods of the instant disclosure, the ExoQuick kit from SBI System Biosciences (which has been validated to be more effective in exosome isolation than the more traditional ultra-centrifugation method, available at <http://www.systembio.com/exoquick-exosomes/>) was utilized in isolation of exosomes from human ascite samples. As demonstrated by the instant disclosure, significant increases in PLA₂ activities were detected in EOC blood, cell-free ascites, and exosomes vs. similar samples collected from non-malignant liver diseases or healthy controls. With reference to **FIG. 2**, approximately 80% of cell-free ascitic PLA₂ activity is shown as associated with the exosome portion in ascites. In addition, the observed exosome-associated PLA₂ activity proved sensitive to methyl arachidonyl fluorophosphonate (MAFP) (an irreversible dual cPLA₂ and iPLA₂ activity inhibitor), BEL (an irreversible and selective iPLA₂ activity inhibitor), and/or thioether amide-PC (TAPC) (a selective sPLA₂ activity inhibitor). Referring now to **FIG. 2B**, the inhibitor's effectiveness is confirmed by its ability in blocking sPLA₂ activity when a positive control (from bee venom; Cayman) was used (**Fig. 2D**). As such, the method and disclosure herein, provides analysis of PLA₂ activity, which is demonstrated herein to be associated with human EOC ascites and exosome (from ascites and blood), presents a novel and highly useful method for diagnostic and personalized treatment purposes in GYN cancers.

Example 5: PLA₂ Activity in Mouse Tissue.

[0080] Referring now to **FIGs. 3A, C and D**, PLA₂ activities measured from several mouse models for GYN diseases are presented. These assays demonstrate high PLA₂ activities in tumors from human A2780 and SKOV3 xenografts, as well as from tumors derived from mouse EOC ID-8 cells. Further, and as shown herein, PLA₂ activity is greatly reduced in tumors from bromoenol lactone (BEL) -treated mice compared to those from controls. Further, as shown herein, BEL targets PLA₂ enzymes (see, **FIG. 3B**, panels a - b) and reduces tumorigenesis, correlating to a reduction in PLA₂ activity as reducing tumorigenesis.

[0081] Referring now to **FIG. 3E**, PLA₂ activity was measured in several other mouse tissues, including the kidney, lung, and liver. Tumors did not grow in or on these organs, and tumor cell injections did not significantly change the PLA₂ activities in these organs (data not shown). Further, it was determined that at least some (and possibly most) of the activities detected in these tissues appeared to be sPLA₂ activity of normal tissues (based on the activity's extracellular localization and insensitivity to BEL and AACACF3). In all experiments, controls

were run which did not include added substrate. Only the merged pictures and only kidney results among different tissues are shown in **FIGs. 3B to 3E** (data not shown). These data demonstrate the use of PLA₂ activity for following progression of these diseases and/or to monitor the effective of various treatments for the disease.

REFERENCES

1. Sengupta S, Wang Z, Tipps R, Xu Y. Biology of LPA in health and disease. *Semin Cell Dev Biol.* 2004 Oct;15(5):503-12.
2. Xu Y, Sengupta S, Singh S, Steinmetz R. Novel lipid signaling pathways in ovarian cancer cells. *Cell Science Reviews* 2006;3:168-97.
3. Mills GB, Moolenaar WH. The emerging role of lysophosphatidic acid in cancer. *Nat Rev Cancer.* 2003 Aug;3(8):582-91.
4. Xu Y, Casey G, Mills GB. Effect of lysophospholipids on signaling in the human Jurkat T cell line. *J Cell Physiol.* 1995 Jun;163(3):441-50.
5. Xu Y, Fang XJ, Casey G, Mills GB. Lysophospholipids activate ovarian and breast cancer cells. *Biochem J.* 1995 Aug 1;309 (Pt 3):933-40.
6. Shen Z, Belinson J, Morton RE, Xu Y, Xu Y. Phorbol 12-myristate 13-acetate stimulates lysophosphatidic acid secretion from ovarian and cervical cancer cells but not from breast or leukemia cells. *Gynecol Oncol.* 1998 Dec;71(3):364-8.
7. Xu Y, Shen Z, Wiper DW, Wu M, Morton RE, Elson P, et al. Lysophosphatidic acid as a potential biomarker for ovarian and other gynecologic cancers. *Jama.* 1998 Aug 26;280(8):719-23.
8. Xiao Y, Chen Y, Kennedy AW, Belinson J, Xu Y. Evaluation of plasma lysophospholipids for diagnostic significance using electrospray ionization mass spectrometry (ESI-MS) analyses. *Ann N Y Acad Sci.* 2000 Apr;905:242-59.
9. Schwartz BM, Hong G, Morrison BH, Wu W, Baudhuin LM, Xiao YJ, et al. Lysophospholipids increase interleukin-8 expression in ovarian cancer cells. *Gynecol Oncol.* 2001 May;81(2):291-300.
10. Shen Z, Wu M, Elson P, Kennedy AW, Belinson J, Casey G, et al. Fatty acid composition of lysophosphatidic acid and lysophosphatidylinositol in plasma from patients with ovarian cancer and other gynecological diseases. *Gynecol Oncol.* 2001 Oct;83(1):25-30.

11. Xiao YJ, Schwartz B, Washington M, Kennedy A, Webster K, Belinson J, et al. Electrospray ionization mass spectrometry analysis of lysophospholipids in human ascitic fluids: comparison of the lysophospholipid contents in malignant vs nonmalignant ascitic fluids. *Anal Biochem.* 2001 Mar;290(2):302-13.
12. Xu Y, Xiao YJ, Baudhuin LM, Schwartz BM. The role and clinical applications of bioactive lysolipids in ovarian cancer. *J Soc Gynecol Investig.* 2001 Jan-Feb;8(1):1-13.
13. Baudhuin LM, Cristina KL, Lu J, Xu Y. Akt activation induced by lysophosphatidic acid and sphingosine-1-phosphate requires both mitogen-activated protein kinase kinase and p38 mitogen-activated protein kinase and is cell-line specific. *Mol Pharmacol.* 2002 Sep;62(3):660-71.
14. Sengupta S, Xiao YJ, Xu Y. A novel laminin-induced LPA autocrine loop in the migration of ovarian cancer cells. *Faseb J.* 2003 Aug;17(11):1570-2.
15. Xu Y, Xiao YJ, Zhu K, Baudhuin LM, Lu J, Hong G, et al. Unfolding the pathophysiological role of bioactive lysophospholipids. *Curr Drug Targets Immune Endocr Metabol Disord.* 2003 Mar;3(1):23-32.
16. Baudhuin LM, Jiang Y, Zaslavsky A, Ishii I, Chun J, Xu Y. S1P3-mediated Akt activation and cross-talk with platelet-derived growth factor receptor (PDGFR). *Faseb J.* 2004 Feb;18(2):341-3.
17. Sutphen R, Xu Y, Wilbanks GD, Fiorica J, Grendys EC, Jr., LaPolla JP, et al. Lysophospholipids are potential biomarkers of ovarian cancer. *Cancer Epidemiol Biomarkers Prev.* 2004 Jul;13(7):1185-91.
18. Kim KS, Ren J, Jiang Y, Ebrahim Q, Tipps R, Cristina K, et al. GPR4 plays a critical role in endothelial cell function and mediates the effects of sphingosylphosphorylcholine. *Faseb J.* 2005 May;19(7):819-21.
19. Wang P, Wu XH, Chen WX, Shan BE, Guo Q. [Expression of lysophosphatidic acid receptor in human ovarian cancer cell lines 3AO, SKOV3, OVCAR3 and its significance]. *Di Yi Jun Yi Da Xue Xue Bao.* 2005 Nov;25(11):1422-4, 31.
20. Kim KS, Sengupta S, Berk M, Kwak YG, Escobar PF, Belinson J, et al. Hypoxia enhances lysophosphatidic acid responsiveness in ovarian cancer cells and lysophosphatidic acid induces ovarian tumor metastasis in vivo. *Cancer Res.* 2006 Aug 15;66(16):7983-90.

21. Ren J, Xiao YJ, Singh LS, Zhao X, Zhao Z, Feng L, et al. Lysophosphatidic acid is constitutively produced by human peritoneal mesothelial cells and enhances adhesion, migration, and invasion of ovarian cancer cells. *Cancer Res.* 2006 Mar 15;66(6):3006-14.
22. Zhao X, Wang D, Zhao Z, Xiao Y, Sengupta S, Xiao Y, et al. Caspase-3-dependent activation of calcium-independent phospholipase A2 enhances cell migration in non-apoptotic ovarian cancer cells. *J Biol Chem.* 2006 Sep 29;281(39):29357-68.
23. Sengupta S, Kim KS, Berk MP, Oates R, Escobar P, Belinson J, et al. Lysophosphatidic acid downregulates tissue inhibitor of metalloproteinases, which are negatively involved in lysophosphatidic acid-induced cell invasion. *Oncogene.* 2007 May 3;26(20):2894-901.
24. Wang H, Wu XH, Shi L. [Lysophosphatidic acid promotes transplanted tumor growth of human epithelium ovarian cancer in nude mice]. *Nan Fang Yi Ke Da Xue Xue Bao.* 2007 Feb;27(2):228-9, 31.
25. Li H, Wang D, Zhang H, Kirmani K, Zhao Z, Steinmetz R, et al. Lysophosphatidic acid stimulates cell migration, invasion, and colony formation as well as tumorigenesis/metastasis of mouse ovarian cancer in immunocompetent mice. *Mol Cancer Ther.* 2009 Jun;8(6):1692-701.
26. Xu Y, Wang D, Wang Z. Lipid generation and signaling in ovarian cancer. In: Stack MS, Fishman DA, editors. *Cancer Treatment and Research-Ovarian cancer.* 2 ed. New York, Dordrecht, Heidelberg, London: Springer; 2009. p. 241-68.
27. Murph MM, Liu W, Yu S, Lu Y, Hall H, Hennessy BT, et al. Lysophosphatidic acid-induced transcriptional profile represents serous epithelial ovarian carcinoma and worsened prognosis. *PLoS ONE.* 2009;4(5):e5583.
28. Aoki J, Inoue A, Okudaira S. Two pathways for lysophosphatidic acid production. *Biochim Biophys Acta.* 2008 Sep;1781(9):513-8.
29. Balsinde J, Balboa MA, Insel PA, Dennis EA. Regulation and inhibition of phospholipase A2. *Annu Rev Pharmacol Toxicol.* 1999;39:175-89.
30. Kudo I, Murakami M. [Phospholipase A2]. *Tanpakushitsu Kakusan Koso.* 1999 Jun;44(8 Suppl):1013-24.

31. Kudo I, Murakami M. Phospholipase A2 enzymes. Prostaglandins Other Lipid Mediat. 2002 Aug;68-69:3-58.
32. Aoki J. Mechanisms of lysophosphatidic acid production. Semin Cell Dev Biol. 2004 Oct;15(5):477-89.
33. Tang J, Kriz RW, Wolfman N, Shaffer M, Seehra J, Jones SS. A novel cytosolic calcium-independent phospholipase A2 contains eight ankyrin motifs. J Biol Chem. 1997 Mar 28;272(13):8567-75.
34. Ramanadham S, Yarasheski KE, Silva MJ, Wohltmann M, Novack DV, Christiansen B, et al. Age-related changes in bone morphology are accelerated in group VIA phospholipase A2 (iPLA₂beta)-null mice. Am J Pathol. 2008 Apr;172(4):868-81.
35. Song Y, Wilkins P, Hu W, Murthy KS, Chen J, Lee Z, et al. Inhibition of calcium-independent phospholipase A2 suppresses proliferation and tumorigenicity of ovarian carcinoma cells. Biochem J. 2007 Sep 15;406(3):427-36.
36. Sedlakova I, Vavrova J, Tosner J, Hanousek L. Lysophosphatidic acid: an ovarian cancer marker. Eur J Gynaecol Oncol. 2008;29(5):511-4.
37. Sedlakova I, Vavrova J, Tosner J, Hanousek L. [Lysophosphatidic acid in ovarian cancer patients]. Ceska Gynekol. 2006 Jul;71(4):312-7.
38. Seballos L, Zhang JZ, Sutphen R. Surface-enhanced Raman scattering detection of lysophosphatidic acid. Anal Bioanal Chem. 2005 Nov;383(5):763-7.
39. Meleh M, Pozlep B, Mlakar A, Meden-Vrtovec H, Zupancic-Kralj L. Determination of serum lysophosphatidic acid as a potential biomarker for ovarian cancer. J Chromatogr B Analyt Technol Biomed Life Sci. 2007 Oct 15;858(1-2):287-91.
40. Jeon ES, Moon HJ, Lee MJ, Song HY, Kim YM, Cho M, et al. Cancer-derived lysophosphatidic acid stimulates differentiation of human mesenchymal stem cells to myofibroblast-like cells. Stem Cells. 2008 Mar;26(3):789-97.
41. Kosaka N, Iguchi H, Ochiya T. Circulating microRNA in body fluid: a new potential biomarker for cancer diagnosis and prognosis. Cancer Sci. 2010 Jun 18.
42. Taylor DD, Gercel-Taylor C. MicroRNA signatures of tumor-derived exosomes as diagnostic biomarkers of ovarian cancer. Gynecol Oncol. 2008 Jul;110(1):13-21.

43. Subra C, Grand D, Laulagnier K, Stella A, Lambeau G, Paillasse M, et al. Exosomes account for vesicle-mediated transcellular transport of activatable phospholipases and prostaglandins. *J Lipid Res.* 2010 Aug;51(8):2105-20.
44. Burke JE, Dennis EA. Phospholipase A2 structure/function, mechanism, and signaling. *J Lipid Res.* 2009 Apr;50 Suppl:S237-42.
45. Hazen SL, Zupan LA, Weiss RH, Getman DP, Gross RW. Suicide inhibition of canine myocardial cytosolic calcium-independent phospholipase A2. Mechanism-based discrimination between calcium-dependent and -independent phospholipases A2. *J Biol Chem.* 1991 Apr 15;266(11):7227-32.
46. Burke JE, Dennis EA. Phospholipase A2 biochemistry. *Cardiovasc Drugs Ther.* 2009 Feb;23(1):49-59.
47. Baskakis C, Magrioti V, Cotton N, Stephens D, Constantinou-Kokotou V, Dennis EA, et al. Synthesis of polyfluoro ketones for selective inhibition of human phospholipase A2 enzymes. *J Med Chem.* 2008 Dec 25;51(24):8027-37.
48. Lopez-Vales R, Navarro X, Shimizu T, Baskakis C, Kokotos G, Constantinou-Kokotou V, et al. Intracellular phospholipase A(2) group IVA and group VIA play important roles in Wallerian degeneration and axon regeneration after peripheral nerve injury. *Brain.* 2008 Oct;131(Pt 10):2620-31.
49. Cummings BS. Phospholipase A2 as targets for anti-cancer drugs. *Biochem Pharmacol.* 2007 Oct 1;74(7):949-59.
50. Gorovetz M, Backelandt M, Berner A, Trope CG, Davidson B, Reich R. The clinical role of phospholipase A2 isoforms in advanced-stage ovarian carcinoma. *Gynecol Oncol.* 2006 Dec;103(3):831-40.
51. Iorio E, Ricci A, Bagnoli M, Pisanu ME, Castellano G, Di Vito M, et al. Activation of phosphatidylcholine cycle enzymes in human epithelial ovarian cancer cells. *Cancer Res.* 2010 Mar 1;70(5):2126-35.
52. Li H, Zhao Z, Wei G, Yan L, Wang D, Zhang H, et al. Group VIA phospholipase A2 in both host and tumor cells is involved in ovarian cancer development. *FASEB J.* 2010 Jun 8.

53. van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature*. 2002 Jan 31;415(6871):530-6.
54. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med*. 2002 Dec 19;347(25):1999-2009.
55. Zhao Z, Xiao Y, Elson P, Tan H, Plummer SJ, Berk M, et al. Plasma lysophosphatidylcholine levels: potential biomarkers for colorectal cancer. *J Clin Oncol*. 2007 Jul 1;25(19):2696-701.
56. Zhao Z, Xu Y. Measurement of endogenous lysophosphatidic acid by ESI-MS/MS in plasma samples requires pre-separation of lysophosphatidylcholine. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2009 Nov 1;877(29):3739-42.
57. Zhao Z, Xu Y. An extremely simple method for extraction of lysophospholipids and phospholipids from blood samples. *J Lipid Res*. 2009 Sep 25.
58. Li H, Wang D, Singh LS, Berk M, Tan H, Zhao Z, et al. Abnormalities in osteoclastogenesis and decreased tumorigenesis in mice deficient for ovarian cancer G protein-coupled receptor 1. *PLoS One*. 2009;4(5):e5705.
59. Vang R, Shih Ie M, Kurman RJ. Ovarian low-grade and high-grade serous carcinoma: pathogenesis, clinicopathologic and molecular biologic features, and diagnostic problems. *Adv Anat Pathol*. 2009 Sep;16(5):267-82.
60. Kurman RJ, Shih Ie M. Pathogenesis of ovarian cancer: lessons from morphology and molecular biology and their clinical implications. *Int J Gynecol Pathol*. 2008 Apr;27(2):151-60.
61. Bast RC, Jr., Hennessy B, Mills GB. The biology of ovarian cancer: new opportunities for translation. *Nat Rev Cancer*. 2009 Jun;9(6):415-28.
62. Schlaerth AC, Abu-Rustum NR. Role of minimally invasive surgery in gynecologic cancers. *Oncologist*. 2006 Sep;11(8):895-901.
63. Shao M, Cao L, Shen C, Satpathy M, Chelladurai B, Bigsby RM, et al. Epithelial-to-mesenchymal transition and ovarian tumor progression induced by tissue transglutaminase. *Cancer Res*. 2009 Dec 15;69(24):9192-201.

64. Jiang Y, Berk M, Singh LS, Tan H, Yin L, Powell CT, et al. KiSS1 suppresses metastasis in human ovarian cancer via inhibition of protein kinase C alpha. *Clin Exp Metastasis*. 2005;22(5):369-76.
65. Yang G, Rosen DG, Mercado-Urbe I, Colacino JA, Mills GB, Bast RC, Jr., et al. Knockdown of p53 combined with expression of the catalytic subunit of telomerase is sufficient to immortalize primary human ovarian surface epithelial cells. *Carcinogenesis*. 2007 Jan;28(1):174-82.
66. Wang D, Zhao Z, Caperell-Grant A, Yang G, Mok SC, Liu J, et al. S1P differentially regulates migration of human ovarian cancer and human ovarian surface epithelial cells. *Mol Cancer Ther*. 2008 Jul;7(7):1993-2002.
67. Murakami M, Kudo I. Phospholipase A2. *J Biochem*. 2002 Mar;131(3):285-92.
68. Lauber K, Bohn E, Krober SM, Xiao YJ, Blumenthal SG, Lindemann RK, et al. Apoptotic cells induce migration of phagocytes via caspase-3-mediated release of a lipid attraction signal. *Cell*. 2003 Jun 13;113(6):717-30.
69. Flesken-Nikitin A, Choi KC, Eng JP, Shmidt EN, Nikitin AY. Induction of carcinogenesis by concurrent inactivation of p53 and Rb1 in the mouse ovarian surface epithelium. *Cancer Res*. 2003 Jul 1;63(13):3459-63.
70. Dinulescu DM, Ince TA, Quade BJ, Shafer SA, Crowley D, Jacks T. Role of K-ras and Pten in the development of mouse models of endometriosis and endometrioid ovarian cancer. *Nat Med*. 2005 Jan;11(1):63-70.
71. Vanderhyden BC, Shaw TJ, Ethier JF. Animal models of ovarian cancer. *Reprod Biol Endocrinol*. 2003 Oct 7;1:67.
72. Fong MY, Kakar SS. Ovarian cancer mouse models: a summary of current models and their limitations. *J Ovarian Res*. 2009;2(1):12.
73. Ackermann EJ, Kempner ES, Dennis EA. Ca(2+)-independent cytosolic phospholipase A2 from macrophage-like P388D1 cells. Isolation and characterization. *J Biol Chem*. 1994;269(12):9227-33.
74. Deems RA, Eaton BR, Dennis EA. Kinetic analysis of phospholipase A2 activity toward mixed micelles and its implications for the study of lipolytic enzymes. *J Biol Chem*. 1975;250(23):9013-20.

75. Lucas KK, Dennis EA. Distinguishing phospholipase A2 types in biological samples by employing group-specific assays in the presence of inhibitors. Prostaglandins Other Lipid Mediat. 2005;77(1-4):235-48.

[0082] While the novel technology has been illustrated and described in detail in the figures and foregoing description, the same is to be considered as illustrative and not restrictive in character, it being understood that only the preferred embodiments have been shown and described and that all changes and modifications that come within the spirit of the novel technology are desired to be protected. As well, while the novel technology was illustrated using specific examples, theoretical arguments, accounts, and illustrations, these illustrations and the accompanying discussion should by no means be interpreted as limiting the technology. All patents, patent applications, and references to texts, scientific treatises, publications, and the like referenced in this application are incorporated herein by reference in their entirety.

CLAIMS

WHAT IS CLAIMED IS:

1. A method for evaluating a disease, comprising the steps of:
measuring phospholipase A₂ (PLA₂) activity in a patient sample; and
providing information associated with correlating the PLA₂ activity, measured in said step of measuring, with a PLA₂ activity value indicative of a disease, wherein said PLA₂ activity is elevated in a diseased state.
2. The method according to claim 1, further including the step of:
obtaining said patient sample from a patient.
3. The method according to claim 1, wherein said patient sample is a tissue sample.
4. The method according to claim 1, wherein said patient sample includes at least one cell.
5. The method according to claim 1, wherein said sample comprises a cellular fluid.
6. The method according to claim 1, wherein the patient sample is from tumorous tissue.
7. The method according to claim 1, wherein the disease is a gynecological disease.
8. The method according to claim 1, wherein the disease is EOC.
9. The method according to claim 1, wherein the measuring step includes the use of a fluorogenic compound.
10. The method according to claim 9 wherein the fluorogenic compound is DBPC.
11. The method according to claim 1, wherein the measuring step includes the use of a radioactive labeled substrate for PLA₂.
12. The method according to claim 1, further including the step of:
indicating the sample as being from a patient likely to have a form of cancer.
13. The method according to claim 10, wherein said information, provided in said step of providing, correlates PLA₂ activity of demonstratively higher activity in a diseased state than in a non-diseased state.
14. The method according to claim 10, wherein said information, provided in said step of providing, correlates PLA₂ activity that is about 1.5 fold higher activity in a diseased state than in a non-diseased state.
15. The method according to claim 13, wherein the form of cancer affects a mammalian reproductive tract.
16. The method according to claim 15, wherein the form of cancer is EOC.

17. The method according to claim 1 further including the step of introducing a compound to the patient which inhibits activity of an isoform of PLA₂
18. The method according to claim 17, wherein the isoform of PLA₂ is a sPLA₂.
19. The method according to claim 17, wherein the isoform of PLA₂ is an iPLA₂.
20. The method according to claim 17, wherein the isoform of PLA₂ is a cPLA₂.
21. The method according to claim 17, wherein the compound inhibits both a cPLA₂ and an iPLA₂s.
22. The method according to claim 1, wherein the PLA₂ includes at least one of iPLA₂ and cPLA₂.
23. The method according to claim 9, wherein the measuring step comprises a treatment step and a detection step.
24. The method according to claim 23, wherein the treatment step is performed in vivo.
25. A method of using PLA₂ activity to assess the health of a patient, comprising the steps of:
 - obtaining a sample from an individual;
 - contacting the sample with a fluorogenic compound;
 - detecting a signal produced by the fluorogenic compound in the presence of the sample; and
 - correlating the size of the signal with the health of the patient.
26. The method of claim 25, wherein the sample is a tissue.
27. The method of claim 26, further comprising the step of fixing the tissue
28. The method of claim 26, further comprising the step of:
 - homogenizing the sample;
 - collecting a supernatant of the homogenized sample, wherein the fluorogenic compound is introduced, in the step of introducing, to the supernatant.
29. The method of claim 25, further comprising the step of correlating a value of the signal, detected in the step of detecting, with a disease.
30. The method of claim 29, wherein the disease is a form of cancer.
31. The method of claim 25, further including the step of introducing a compound to the sample which inhibits activity of an isoform of PLA₂.
32. The method of claim 31, wherein the isoform of PLA₂ is iPLA₂.

33. A marker for disease, comprising:
a marker wherein said marker is phospholipase A₂ (PLA₂) activity that is demonstratively higher in a diseased state than it is in a healthy state.
34. The marker according to claim 33, wherein the activity is at least about 1.5fold higher in a sample from a patient with a disease than it is in a patient that is without disease.
35. The marker according to claim 33, wherein the marker is elevated in at least one sample for a portion of a patient's body selected from the group consisting of: cells, tissues, and fluids.
36. The marker according, wherein said marker is indicative of cancer.
37. The marker according to claim 36, wherein the marker is EOC.
38. The marker according to claim 33, wherein said marker can be used to diagnose a disease.
39. The marker according to claim 33, wherein said marker can be used to evaluate the effectiveness of a treatment for a disease.
40. The marker according to claim 33, wherein the marker can be used to determine the prognosis of a disease.
41. The marker according to claim 33, wherein the marker can be used to determine the personalized treatment of a patient.
42. The marker according to claim 33, wherein the activity is due primarily to elevated iPLA₂β activity.

1/17




FIG. 1 A

2/17




FIG. 1 B

3/17




FIG. 1 C

4/17




FIG. 1 D

5/17





FIG. 1 E

6/17




FIG. 2 A

7/17




FIG. 2 B

8/17




FIG. 2 C

9/17




FIG. 2 D

10/17




FIG. 3 A

11/17




FIG. 3 B

12/17




FIG. 3 C

13/17




FIG. 3 D

14/17





FIG. 3 E

15/17




FIG. 4 A

16/17




FIG. 4 B

17/17




FIG. 4 C