

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199738661 B2**  
**(10) Patent No. 718458**

(54) Title  
**Novel DNAs and process for producing proteins by using the same**

(51)<sup>7</sup> International Patent Classification(s)  
**C12N 015/00 C12P 021/00**

(21) Application No: **199738661**

(22) Application Date: **1997.08.19**

(87) WIPO No: **WO98/07840**

(30) Priority Data

(31) Number	(32) Date	(33) Country
<b>8-235928</b>	<b>1996.08.19</b>	<b>JP</b>

(43) Publication Date : **1998.05.11**

(43) Publication Journal Date : **1998.07.02**

(44) Accepted Journal Date : **2000.04.13**

(71) Applicant(s)  
**Snow Brand Milk Products Co., Ltd.**

(72) Inventor(s)  
**Nobuaki Nakagawa; Hisataka Yasuda; Tomonori Morinaga**

(74) Agent/Attorney  
**DAVIES COLLISON CAVE,GPO Box 3876,SYDNEY NSW 2001**

OPI DATE 11/05/98 APPLN. ID 38661/97  
PC' AOJP DATE 02/07/98 PCT NUMBER PCT/JP97/02859



AU9738661



(51) 国際特許分類6 C12N 15/00, C12P 21/00	A1	(11) 国際公開番号 WO98/07840
		(43) 国際公開日 1998年2月26日(26.02.98)


(21) 国際出願番号 PCT/JP97/02859	(74) 代理人 弁理士 藤野清也(FUJINO, Seiya) 〒160 東京都新宿区四谷1丁目2番1号 三浜ビル8階 Tokyo, (JP)
(22) 国際出願日 1997年8月19日(19.08.97)	
(30) 優先権データ 特願平8/235928 1996年8月19日(19.08.96) JP	(81) 指定国 AU, CA, CN, FI, HU, IL, KR, MX, NO, NZ, RU, US, 欧州特許 (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(71) 出願人 (米国を除くすべての指定国について) 雪印乳業株式会社 (SNOW BRAND MILK PRODUCTS CO., LTD.)(JP/JP) 〒065 北海道札幌市東区苗穂町6丁目1番1号 Hokkaido, (JP)	添付公開書類 国際調査報告書
(72) 発明者 ; および (75) 発明者 / 出願人 (米国についてののみ) 中川信明(NAKAGAWA, Nobuaki)(JP/JP) 〒329-05 栃木県下都賀郡石橋町石橋578-15 西蒲ハイツ2-4 Tochigi, (JP) 保田尚孝(YASUDA, Hisataka)(JP/JP) 〒329-04 栃木県河内郡南河内町緑2-3293-46 Tochigi, (JP) 森永伴法(MORINAGA, Tomonori)(JP/JP) 〒321-02 栃木県下都賀郡壬生町幸町3-11-12 Tochigi, (JP)	

(54)Title: NOVEL DNAs AND PROCESS FOR PRODUCING PROTEINS BY USING THE SAME

(54)発明の名称 新規DNA及びそれを用いた蛋白質の製造方法


(57) Abstract

~~DNAs represented by Sequence Listings (1 and 2) and a process for producing proteins which comprises inserting these DNAs into expression vectors to thereby produce proteins having molecular weights of about 60 kD (under reductive conditions) and about 60 kD and about 120 kD (under nonreductive conditions) and being capable of inhibiting the formation of osteoclasts. These proteins are useful in the treatment of osteoporosis and rheumatism.~~



### ABSTRACT OF THE DISCLOSURE

DNAs having the nucleotide sequences of the Sequences No. 1 and No. 2 in the Sequence Table and a process for producing a protein which comprises inserting these DNAs into expression vectors to thereby produce a protein having molecular weights of about 60 kD (under reducing conditions) and about 60 kD and 120 kD (under non-reducing conditions) and being capable of inhibiting formation of osteoclast. These proteins are useful in the treatment of osteoporosis and rheumatism.



## DESCRIPTION

### NOVEL DNA AND PROCESS FOR PREPARING PROTEIN USING THE DNA


## FIELD OF TECHNOLOGY

The present invention relates to a novel DNA and a process for preparing a protein which possesses an activity to inhibit osteoclast differentiation and/or maturation (hereinafter called osteoclastogenesis-inhibitory activity) by a genetic engineering technique using the DNA. More particularly, the present invention relates to a genomic DNA encoding a protein OCIF which possesses an osteoclastogenesis-inhibitory activity and a process for preparing said protein by a genetic engineering technique using the genomic DNA.

## BACKGROUND OF THE INVENTION

Human bones are constantly repeating a process of resorption and formation. Osteoblasts controlling formation of bones and osteoclasts controlling resorption of bones take major roles in this process. Osteoporosis is a typical disease caused by abnormal metabolism of bones. This disease is caused when bone resorption by osteoclasts exceeds bone formation by osteoblasts. Although the mechanism of this disease is still to be elucidated completely, the disease causes the bones to ache, makes the bones fragile, and may result in fracturing of the bones. As the population of the aged increases, this disease results in an increase in the number of bedridden aged people which becomes a social problem. Urgent development of a therapeutic agent for this disease is strongly desired.

Disease due to a




decrease in bone mass is expected to be treated by controlling bone resorption, accelerating bone formation, or improving balance between bone resorption and formation.

Osteogenesis is expected to be increased by accelerating proliferation, differentiation, or activation of the cells controlling the bone formation, or by controlling proliferation, differentiation, or activation of the cells involved in bone

resorption. In recent years, strong interest has been directed to physiologically active proteins (cytokines) exhibiting such activities as described above, and energetic research is ongoing on this subject. The cytokines which have been reported to accelerate proliferation or differentiation of osteoblasts include the proteins of fibroblast growth factor family (FGF: Rodan S. B. et al., Endocrinology vol. 121, p 1917, 1987), insulin-like growth factor I (IGF-I: Hock J. M. et al., Endocrinology vol. 122, p 254, 1988), insulin growth factor II (IGF-II: McCarthy T. et al., Endocrinology vol. 124, p 301, 1989), Activin A (Centrella M. et al., Mol. Cell. Biol., vol. 11, p 250, 1991), transforming growth factor- $\beta$ , (Noda M., The Bone, vol. 2, p 29, 1988), Vasculotropin (Varonique M. et al., Biochem. Biophys. Res. Commun., vol. 199, p 380, 1994), and the protein of heterotopic bone formation factor family (bone morphogenic protein; BMP: BMP-2; Yanaguchi A. et al., J. Cell Biol. vol. 113, p 682, 1991, OP-1; Sampath T. K. et al., J. Biol. Chem. vol. 267, p 20532, 1992, and Knutsen R. et al., Biochem. Biophys. Res. Commun. vol. 194, P 1352, 1993).

On the other hand, as the cytokines which suppress



differentiation and/or maturation of osteoclasts, transforming growth factor- $\beta$  (Chenu C, et. al., Proc. Natl. Acad. Sci. USA, vol. 85, p 5683, 1988), interleukin-4 (Kasano K. et al., Bone-Miner., vol. 21, p 179, 1993), and the like have been reported. Further, as the cytokines which suppress bone resorption by osteoclast, calcitonin (Bone-Miner., vol. 17, p 347, 1992), macrophage colony stimulating factor (Hattersley G. et al., J. Cell. Physiol. vol. 137, p 199. 1988), interleukin-4 (Watanabe, K. et al., Biochem. Biophys. Res. Commun. vol. 172. P 1035, 1990), and interferon- $\gamma$  (Gowen M. et al., J. Bone Miner. Res., vol. 1, p 46.9, 1986) have been reported.

These cytokines are expected to be used as agents for treating diseases accompanying bone loss by accelerating bone formation or suppressing bone resorption. Clinical tests are being undertaken to verify the effect of improving bone metabolism of some cytokines such as insulin-like growth

factor-I and the heterotopic bone formation factor family. In addition, calcitonin is already commercially available as a therapeutic agent for osteoporosis and a pain relief agent. At present, drugs for clinically treating bone diseases or shortening the period of treatment of bone diseases include activated vitamin D<sub>3</sub>, calcitonin and its derivatives, and hormone preparations such as estradiol agent, ipriflavon or calcium preparations. These agents are not necessarily satisfactory in terms of the efficacy and therapeutic results. Development of a novel therapeutic agent which can be used in




place of these agents is strongly desired.

In view of this situation, the present inventors have undertaken extensive studies. As a result, the present inventors had found protein OCIF exhibiting an osteoclastogenesis-inhibitory activity in a culture broth of human embryonic lung fibroblast IMR-90 (ATCC Deposition No. CCL186), and filed a patent application (PCT/JP96/00374). The present inventors have conducted further studies relating to the origin of this protein OCIF exhibiting the osteoclastogenesis-inhibitory activity. The studies have matured into determination of the sequence of a genomic DNA encoding the human origin OCIF. Accordingly, an object of the present invention is to provide a genomic DNA encoding protein OCIF exhibiting osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA.

#### DISCLOSURE OF THE INVENTION

Specifically, the present invention relates to a genomic DNA encoding protein OCIF exhibiting osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA. The DNA of the present invention includes the nucleotide sequences No. 1 and No. 2 in the Sequence Table attached hereto.

Moreover, the present invention relates to a process for preparing a protein, comprising inserting a DNA including the nucleotide sequences of the sequences No. 1 and No. 2 in the Sequence Table into an expression vector, producing a vector



capable of expressing a protein having the following physicochemical characteristics and exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts, and producing this protein by a genetic engineering technique,

(a) molecular weight (SDS-PAGE):

(i) Under reducing conditions: about 60 kD,

(ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 of the Sequence Table,

(c) affinity:


exhibits affinity to a cation exchanger and heparin, and

(d) thermal stability:

(i) the osteoclast differentiation and/or maturation inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,

(ii) the osteoclast differentiation and/or maturation inhibitory activity is lost when treated with heat at 90°C for 10 minutes.

The protein obtained by expressing the gene of the present invention exhibits an osteoclastogenesis-inhibitory activity. This protein is effective as an agent for the treatment and improvement of diseases involving decrease in the amount of bone such as osteoporosis, diseases relating to bone metabolism abnormality such as rheumatism, degenerative joint disease, or multiple myeloma, and is useful as an antigen to establish an





immunological diagnosis of such diseases.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a result of Western Blotting analysis of the protein obtained by causing genomic DNA of the present invention to express a protein in Example 4 (iii), wherein lane 1 indicates a size marker, lane 2 indicates the culture broth of COS7 cells in which a vector pWESR $\alpha$ OCIF (Example 4 (iii)) has been transfected, and lane 3 is the culture broth of COS7 cell in which a vector pWESR $\alpha$ (control) has been transfected.

#### BEST MODE FOR CARRYING OUT THE INVENTION

The genomic DNA encoding the protein OCIF which exhibits osteoclastogenesis-inhibitory activity in the present invention can be obtained by preparing a cosmid library using a human placenta genomic DNA and a cosmid vector and by screening this library using DNA fragments which are prepared based on the OCIF cDNA as a probe. The thus-obtained genomic DNA is inserted into a suitable expression vector to prepare an OCIF expression cosmid. A recombinant type OCIF can be obtained by transfecting the genomic DNA into a host organism such as various types of cells or microorganism strains and causing the DNA to express a protein by a conventional method. The resultant protein exhibiting osteoclastogenesis-inhibitory activity (an osteoclastogenesis-inhibitory factor) is useful as an agent for the treatment and improvement of diseases involving a decrease in bone mass such as osteoporosis and other diseases relating to bone metabolism abnormality and also as an antigen to prepare antibodies for establishing immunological diagnosis of such



diseases. The protein of the present invention can be prepared as a drug composition for oral or non-oral administration. Specifically, the drug composition of the present invention containing the protein which is an osteoclastogenesis-inhibitory factor as an active ingredient can be safely administered to humans and animals. As the form of drug composition, a composition for injection, composition for intravenous drip, suppository, nasal agent, sublingual agent, percutaneous absorption agent, and the like are given. In the case of the composition for injection, such a composition is a mixture of a pharmacologically effective amount of the

osteoclastogenesis-inhibitory factor of the present invention and a pharmaceutically acceptable carrier. The composition may further comprise amino acids, saccharides, cellulose derivatives, and other excipients and/or activation agents, including other organic compounds and inorganic compounds which are commonly added to a composition for injection. When an injection preparation is prepared using the osteoclastogenesis-inhibitory factor of the present invention and these excipients and activation agents, a pH adjuster, buffering agent, stabilizer, solubilizing agent, and the like may be added if necessary to prepare various types of injection agents.

The present invention will now be described in more detail by way of examples which are given for the purpose of illustration and not intended to be limiting of the present invention.



## Example 1

### < Preparation of a cosmid library >

A cosmid library was prepared using human placenta genomic DNA (Clontech; Cat. No. 6550-2) and pWE15 cosmid vector (Stratagene). The experiment was carried out following principally the protocol attached to the pWE15 cosmid vector kit of Stratagene Company, provided Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory (1989)) was referred to for common procedures for handling DNA, E. coli, and phage.

#### (i) Preparation of restriction enzyme digest of human-genomic DNA

Human placenta genomic DNA dissolved in 750  $\mu$ l of a solution containing 10 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, and 100 mM NaCl was added to four 1.5 ml Eppendorf tubes (tube A, B, C, and D) in the amount of 100  $\mu$ g each. Restriction enzyme MboI was added to these tubes in the amounts of 0.2 unit for tube A, 0.4 unit for tube B, 0.6 unit for tube C, and 0.8 unit for tube D, and DNA was digested for 1 hour. Then, EDTA in the amount to make a 20 mM concentration was added to each tube to terminate the reaction, followed by extraction with phenol/chloroform (1:1). A two-fold amount of ethanol was added to the aqueous layer to precipitate DNA. DNA was collected by centrifugation, washed with 70% ethanol, and DNA in each tube was dissolved in 100  $\mu$ l of TE (10mM Tris-HCl (pH 8.0) + 1mM EDTA) buffer solution, hereinafter called TE). DNA in four tubes was combined in one tube and incubated for 10 minutes at 68°C. After cooling to room



temperature, the mixture was overlayed onto a 10%-40% linear sucrose gradient which was prepared in a buffer containing 20 mM Tris-HCl (pH 8.0), 5mM EDTA, and 1M NaCl in a centrifuge

tube (38 ml). The tube was centrifuged at 26,000 rpm for 24 hours at 20°C using a rotor SRP28SA manufactured by Hitachi, Ltd. and 0.4 ml fractions of the sucrose gradient was collected using a fraction collector. A portion of each fraction was subjected to 0.4% agarose electrophoresis to confirm the size of DNA. Fractions containing DNA with a length of 30 kb (kilo base pair) to 40 kb were thus combined. The DNA solution was diluted with TE to make a sucrose concentration to 10% or less and 2.5-fold volumes of ethanol was added to precipitate DNA. DNA was dissolved in TE and stored at 4°C.

(ii) Preparation of cosmid vector

The pWE15 cosmid vector obtained from Stratagene Company was completely digested with restriction enzyme BamHI according to the protocol attached to the cosmid vector kit. DNA collected by ethanol precipitation was dissolved in TE to a concentration of 1 mg/ml. Phosphoric acid at the 5'-end of this DNA was removed using calf small intestine alkaline phosphatase, and DNA was collected by phenol extraction and ethanol precipitation. The DNA was dissolved in TE to a concentration of 1 mg/ml.

(iii) Ligation of genomic DNA to vector and in vitro packaging

1.5 micrograms of genomic DNA fractionated according to size and 3 µg of pWE15 cosmid vector which was digested with



restriction enzyme BamHI were ligated in 20  $\mu$ l of a reaction solution using Ready-To-Go T4DNA ligase of Pharmacia Company. The ligated DNA was packaged in vitro using Gigapack™ II packaging extract (Stratagene) according to the protocol. After the packaging reaction, a portion of the reaction mixture was diluted stepwise with an SM buffer solution and mixed with E. coli XL1-Blue MR (Stratagene) which was suspended in 10 mM  $MgCl_2$  to cause phage to infect, and plated onto LB agar plates containing 50  $\mu$ g/ml of ampicillin. The number of colonies produced was counted. The number of colonies per 1  $\mu$ l of packaging reaction was calculated based on this result.

(iv) Preparation of a cosmid library

The packaging reaction solution thus prepared was mixed with E. coli XL1-Blue MR and the mixture was plated onto agarose plates containing ampicillin so as to produce 50,000 colonies per agarose plate having a 15 cm of diameter. After incubating the plate overnight at 37°C, an LB culture medium was added in the amount of 3 ml per plate to suspend and collect E. coli cells. Each agarose plate was again washed with 3 ml of LB culture medium and the washing was combined with the original suspension of E. coli. The E. coli collected from all agarose plates was placed in a centrifuge tube, glycerol was added to a concentration of 20%, and ampicillin was further added to make a final concentration of 50 $\mu$ g/ml. A portion of the E. Coli suspension was removed and the remainder was stored at -80°C. The removed E. Coli was diluted stepwise and plated onto agar plates to estimate the number of colonies per 1 ml of




suspension.

## Example 2

### <Screening of cosmid library and purification of colony>

A nitrocellulose filter (Millipore) with a diameter of 14.2 cm was placed on each LB agarose plate with a diameter of 15 cm which contained 50 µg/ml of ampicillin. The cosmid library was plated onto the plates so as to produce 50,000 colonies of *E. coli* per plate, followed by incubation overnight at 37°C. *E. coli* on the nitrocellulose filter was transferred to another nitrocellulose filter according to a conventional method to obtain two replica filters. According to the protocol attached to the cosmid vector kit, cosmid DNA in the *E. coli* on the replica filters was denatured with an alkali, neutralized, and immobilized on the nitrocellulose filter using a Stratalinker (Stratagene). The filters were heated for two hours at 80°C in a vacuum oven. The nitrocellulose filters thus obtained were hybridized using two kinds of DNA produced, respectively, from 5'-end and 3'-end of human OCIF cDNA as probes. Namely, a plasmid was purified from *E. coli* pKB/OIF10 (deposited at The Ministry of International Trade and Industry, the Agency of Industrial Science and Technology, Biotechnology Laboratory, Deposition No. FERM BP-5267) containing OCIF cDNA. The plasmid containing OCIF cDNA was digested with restriction enzymes KpnI and EcoRI. Fragments thus obtained were separated using agarose gel electrophoresis. KpnI/EcoRI fragment with a length of 0.2 kb was purified using a Q1AEX II gel extraction kit (Qiagen). This DNA was labelled with <sup>32</sup>p using the Megaprime DNA labelling




System (Amersham) (5' -DNA probe). Apart from this, a BamHI/EcoRV fragment with a length of 0.2 kb which was produced from the above plasmid by digestion with restriction enzymes BamHI and EcoRV was purified and labeled with  $^{32}\text{p}$  (3' -DNA probe). One set of the replica filters described above was hybridized with the 5' -DNA probe and the other with the 3' -DNA probe. Hybridization and washing of the filters were carried out according to the protocol attached to the cosmid vector kit. Autoradiography detected several positive signals with each probe. One colony which gave positive signals with both probes was identified. The colony on the agar plate, which corresponding to the signal on the autoradiogram was isolated and purified. A cosmid was prepared from the purified colony by a conventional method. This cosmid was named pWEOCIF. The size of the human genomic DNA fragment contained in this cosmid was about 38 kb.

### Example 3

#### < Determination of the nucleotide sequence of the human OCIF genomic DNA fragment >

##### (i) Subcloning of the OCIF genomic DNA fragment

Cosmid pWEOCIF was digested with restriction enzyme EcoRI. After the separation of the DNA fragments thus produced by electrophoresis using a 0.7% agarose gel, the DNA fragments were transferred to a nylon membrane (Hybond -N, Amasham) by the Southern blot technique and immobilized on the nylon membrane using Stratalinker (Stratagene). On the other hand, plasmid pBKOCIF was digested with restriction enzyme EcoRI and a 1.6



kb fragment containing human OCIF cDNA was isolated by agarose gel electrophoresis. The fragment was labeled with  $^{32}\text{P}$  using the Megaprime DNA labeling system (Amasham).

Hybridization of the nylon membrane described above with the  $^{32}\text{P}$ -Labelled 1.6=kb OCIF cDNA, performed according to a conventional method, detected DNA fragments with a size of 6 kb, 4 kb, 3.6 kb, and 2.6 kb. These fragments hybridized with the human OCIF cDNA were isolated using agarose gel electrophoresis and individually subcloned into an EcoRI site of pBluescript II SK + vector (Strategene) by a conventional method. The resulting plasmids were named respectively, pBSE 6, pBSE 4, pBSE 3.6, and PBSE 2.6.

(ii) Determination of the nucleotide sequence

The nucleotide sequences of the human OCIF genomic DNA fragments which were subcloned into the plasmid were determined using the ABI Dideoxy Terminator Cycle Sequencing Ready Reaction kit (Perkin Elmer) and the 373 Sequencing System (Applied Biosystems). The primers used for the determination of the nucleotide sequences were synthesized based on the nucleotide sequence of human OCIF cDNA (Sequence ID No. 4 in the Sequence Table). The nucleotide sequences thus determined are given as the Sequences No. 1 and No. 2 in the Sequence Table. The Sequence ID No. 1 includes the first exon of the OCIF gene and the Sequence ID No. 2 includes the second, third, fourth, and fifth exons. A stretch of about 17 kb is present between the first and second exons.

Example 4

<Production of recombinant OCIF using COS-7 cells>





(i) Preparation of OCIF genomic DNA expression cosmid

To express OCIF genomic DNA in animal cells, an expression unit of expression plasmid pcDL-SR $\alpha$ 296 (Molecular and Cellular Biology, vol. 8, P466-472, 1988) was inserted into cosmid vector pWE15 (Stratagene). First of all, the expression plasmid pcDL-SR $\alpha$ 296 was digested with a restriction enzyme Sal I to cut out expression unit with a length of about 1.7 kb which includes an SR $\alpha$ promotor, SV40 later splice signal, poly (A) addition signal, and so on. The digestion products were separated by agarose gel electrophoresis and the 1.7-kb fragment was purified using the QIAEX II gel extraction kit (Qiagen). On the other hand, cosmid vector pWE15 was digested with a restriction enzyme EcoRI and fragments were separated using agarose gel electrophoresis. pWE15 DNA of 8.2 kb long was purified using the QIAEX II gel extraction kit (Qiagen). The ends of these two DNA fragments were bluntled using a DNA blunting kit (Takara Shuzo), ligated using a DNA ligation kit (Takara Shuzo), and transfected into E. coli DH5 $\alpha$  (Gibco BRL). The resultant transformant was grown and the expression cosmid pWESR $\alpha$  containing the expression unit was purified using a Qiagen column (Qiagen).

The cosmid pWE OCIF containing the OCIF genomic DNA fragment with a length of about 38 kb obtained in (i) above was digested with a restriction enzyme NotI to cut out the OCIF genomic DNA fragment of about 38 kb. After separation by agarose gel electrophoresis, the DNA fragment was purified using the QIAEX II gel extraction kit (Qiagen). On the other hand, the expression cosmid pWESR $\alpha$  was



digested with a restriction enzyme EcoRI and the digestion product was extracted with phenol and chloroform, ethanol-precipitated, and dissolved in TE.

pWESR $\alpha$  digested with a restriction enzyme EcoRI and an EcoRI-XmnI-NotI adapter (#1105, #1156 New England Biolaboratory Co.) were ligated using T4 DNA ligase (Takara Shuzo Co., Ltd.). After removal of the free adapter by agarose gel electrophoresis, the product was purified using QIAEX gel extraction kit (Qiagen). The OCIF genomic DNA fragment with a length of about 37 kb, which was derived from the digestion with

restriction enzyme NotI and the pWESR $\alpha$  to which the adapter was attached were ligated using T4 DNA ligase (Takara Shuzo). The DNA was packaged in vitro using the Gigapack packaging extract (Stratagene) and transfected to E. coli XL1-Blue MR (Stratagene). The resultant transformant was grown and the expression cosmid pWESR $\alpha$  OCIF which contained OCIF genomic DNA fragment was purified using a Qiagen column (Qiagen). The OCIF expression cosmid pWESR $\alpha$  OCIF was ethanol-precipitated, dissolved in sterile distilled water and used in the following analysis.

(ii) Transient expression of OCIF genomic DNA and measurement of OCIF activity

A recombinant OCIF was expressed as described below using the OCIF expression cosmid pWESR $\alpha$  OCIF obtained in (i) above and its activity was measured. COS-7 ( $8 \times 10^5$  cells/well) cells (Riken Cell Bank, RCB0539) were plated in a 6-well plate using DMEM culture medium (Gibco BRL) containing 10% fetal bovine serum (Gibco BRL). On the following day, the culture



medium was removed and cells were washed with serum-free DMEM culture medium. The OCIF expression cosmid pWESR $\alpha$ OCIF which had been diluted with OPTI-MEM culture medium (Gibco BRL) was mixed with lipopfectamine and the mixture was added to the cells in each well according to the attached protocol. The expression cosmid pWESR $\alpha$  was added to the cells in the same manner as a control. The amount of the cosmid DNA and Lipopfectamine was respectively 3  $\mu$ g and 12  $\mu$ l. After 24 hours, the culture medium was removed and 1.5 ml of fresh EX-CELL 301 culture medium (JRH Bioscience) was added to each well. The culture medium was recovered after 48 hours and used as a sample for the measurement of OCIF activity. The measurement of OCIF activity was carried out according to the method described by Kumegawa, M. et al. (Protein, Nucleic Acid, and Enzyme, Vol. 34, p 999 (1989)) and the method of TAKAHASHI, N. et al. (Endocrihology vol. 122, p 1373 (1988)). The osteoclast formation from bone marrow cells isolated from mice aged about 17 days in the presence of activated vitamin D<sub>3</sub> was evaluated by the induction of tartaric acid resistant acidic phosphatase activity. The reduction of the acid phosphatase was measured and used as the activity of the protein which possesses osteoclastogenesis-inhibitory activity (OCIF). Namely, 100  $\mu$ l of a OCIF sample which was diluted with  $\alpha$ -MEM culture medium (Gibco BRL) containing 2x10<sup>-8</sup> M activated vitamin D<sub>3</sub> and 10% fetal bovine serum was added to each well of a 96 well micro plate. Then, 3x10<sup>5</sup> bone marrow cells isolated from mice (about 17-days old) suspended in 100  $\mu$ l of  $\alpha$ -MEM culture medium containing 10% fetal bovine serum



were added to each well of the 96 well micro plate and cultured for a week at 37°C and 100% humidity under 5% CO<sub>2</sub> atmosphere. On days 3 and 5, 160 µl of the conditioned medium was removed from each well, and 160 µl of a sample which was diluted with  $\alpha$ -MEM culture medium containing  $1 \times 10^{-8}$  M activated vitamin D<sub>3</sub> and 10% fetal bovine serum was added. On day 7, the cells were washed with a phosphate buffered saline and fixed with a ethanol/acetone (1:1) solution for one

minute at room temperature. The osteoclast formation was detected by staining the cells using an acidic phosphatase activity measurement kit (Acid Phosphatase, Leucocyte, Cat. No. 387-A, Sigma Company). A decrease in the number of cells positive to acidic phosphatase activity in the presence of tartaric acid was taken as the OCIF activity. The results are shown in Table 1, which indicates that the conditioned medium exhibits the similar activity to natural type OCIF obtained from the IMR-90 culture medium and recombinant OCIF produced by CHO cells.



TABLE 1

Activity of OCIF in the conditioned medium of COS-7 cells.

Dilution	1/10	1/20	1/40	1/80	1/160	1/320
OCIF genomic DNA introduced	++	++	++	++	+	-
Vector introduced	-	-	-	-	-	-
Untreated	-	-	-	-	-	-

"++" indicates an activity inhibiting 80% or more of osteoclast formation, "+" indicates an activity inhibiting 30-80% of osteoclast formation, and "-" indicates that no inhibition of osteoclast formation is observed.

(iii) Identification of the product by Western Blotting

A buffer solution (10  $\mu$ l) for SDS-PAGE (0.5 M Tris-HCl, 20% glycerol, 4% SDS, 20  $\mu$ g/ml bromophenol blue, pH 6.8) was added to 10  $\mu$ l of the sample for the measurement of OCIF activity prepared in (ii) above. After boiling for 3 minutes at 100°C, the mixture was subjected to 10% SDS polyacrylamide electrophoresis under non-reducing conditions. The proteins were transferred from the gel to a PVDF membrane (ProBlott, Perkin Elmer) using semi-dry blotting apparatus (Biorad). The membrane was blocked and incubated for 2 hours at 37°C together with a horseradish peroxidase-labeled anti-OCIF antibody obtained by labelling the previously obtained anti-OCIF antibody with horseradish peroxidase according to a conventional method.

After washing, the protein which bound the anti-OCIF antibody was detected using the ECL system (Amasham). As shown in Figure 1, two bands, one with a molecular weight of about 120 kilo dalton and the other 60 kilo dalton, were detected in



the supernatant obtained from the culture broth of COS-7 cells in which pWESR $\alpha$ OCIF was transfected. On the other hand, these two bands with a molecular weight of about 120 kilo dalton and 60 kilo dalton were not detected in the supernatant obtained from the culture broth of COS-7 cells in which pWESR $\alpha$ vector was transfected, confirming that the protein obtained was OCIF.

#### INDUSTRIAL APPLICABILITY

The present invention provides a genomic DNA encoding a protein OCIF which possesses an osteoclastogenesis-inhibitory activity and a process for preparing this protein by a genetic engineering technique using the genomic DNA. The protein obtained by expressing the gene of the present invention exhibits an osteoclastogenesis-inhibitory activity and is useful as an agent for the treatment and improvement of diseases involving a decrease in the amount of bone such as osteoporosis, other diseases resulting from bone metabolism abnormality such as rheumatism, degenerative joint disease, and multiple myeloma. The protein is further useful as an antigen to establish antibodies useful for an immunological diagnosis of such diseases.

#### NOTE ON MICROORGANISM

Depositing Organization:

The Ministry of International Trade and Industry, National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology

Address: 1-3, Higashi-1-Chome, Tsukuba-shi, Ibaraki-ken, Japan



Date of Deposition: June 21, 1995 (originally deposited on  
June 21, 1995 and transferred to the international  
deposition according to the Budapest Treaty on  
October 25, 1995)

Accession No. FERM BP-5267




TABLE OF SEQUENCES

Sequence number: 1

Length of sequence: 1316

Sequence Type: nucleic acid


Strandedness: double

Topology: linear

Molecular type: genomic DNA (human OCIF genomic DNA-1)

Sequence:

```
CTGGAGACAT ATAACTTGAA CACTTGGCCC TGATGGGGAA GCAGCTCTGC AGGGACTTTT 60
TCAGCCATCT GTAAACAATT TCAGTGGCAA CCCGCCAACT GTAATCCATG AATGGGACCA 120
CACTTTACAA GTCATCAAGT CTAAC TTCTA GACCAGGGAA TTAATCGGGG AGACAGCGAA 180
CCCTAGAGCA AAGTGCCAAA CTTCTGTCGA TAGCTTGAGG CTAGTGAAA GACCTCGAGG 240
AGGCTACTCC AGAAGTTCAG CGCGTAGGAA GCTCCGATAC CAATAGCCCT TTGATGATGG 300
TGGGGTTGGT GAAGGGAACA GTGCTCCGCA AGGTTATCCC TGCCCCAGGC AGTCCAATTT 360
TCACTCTGCA GATTCTCTCT GGCTCTAACT ACCCCAGATA ACAAGGAGTG AATGCAGAAT 420
AGCACGGGCT TTAGGGCCAA TCAGACATTA GTTAGAAAAA TTCCTACTAC ATGGTTTATG 480
TAAACTTGAA GATGAATGAT TCGGAACTCC CCGAAAAGGG CTCAGACAAT GCCATGCATA 540
AAGAGGGGCC CTGTAATTTG AGGTTTCAGA ACCCGAAGTG AAGGGGTCAG GCAGCCGGGT 600
ACGGCGGAAA CTCACAGCTT TCGCCCAGCG AGAGGACAAA GGTCTGGGAC ACACTCCAAC 660
TGCGTCCGGA TCTTGGCTGG ATCGGACTCT CAGGGTGGAG GAGACACAAG CACAGCAGCT 720
GCCCAGCGTG TGCCCAGCCC TCCCACCGCT GGTCCCGGCT GCCAGGAGGC TGGCCGCTGG 780
CGGGAAGGGG CCGGGAACC TCAGAGCCCC GCGGAGACAG CAGCCGCCCT GTTCCTCAGC 840
CCGGTGGCTT TTTTTCCTCC TGCTCTCCCA GGGGACAGAC ACCACCGCCC CACCCCTCAC 900
GCCCCACCTC CCTGGGGGAT CTTTTCGGCC CCAGCCCTGA AAGCGTTAAT CCTGGAGCTT 960
TCTGCACACC CCCCAGCCGC TCCCGCCCAA GCTTCCTAAA AAAGAAAGGT GCAAAGTTTG 1020
GTCCAGGATA GAAAAATGAC TGATCAAAGG CAGGCGATAC TTCCTGTTGC CGGGACGCTA 1080
TATATAACGT GATGAGCGCA CGGGCTGCGG AGACGCACCG GAGCGCTCGC CCAGCCGCCG 1140
```





CCTCCAAGCC CCTGAGGTTT CCGGGGACCA CA ATG AAC AAG TTG CTG TGC TGC 1193

Met Asn Lys Leu Leu Cys Cys

-20

-15

GCG CTC GTG GTAAGTCCCT GGGCCAGCCG ACGGGTGCCC GCGCCTGGG 1242

Ala Leu Val

GAGGCTGCTG CCACCTGGTC TCCCAACCTC CCAGCGGACC GGCAGGGAGA AGGCTCCACT 1302

CGCTCCCTCC CAGG 1316

Sequence number: 2

Length of sequence: 9898

Sequence Type: nucleic acid

Strandedness: double

Topology: linear

Molecular type: genomic DNA (human OCIF genomic DNA-2)

Sequence:

GCTTACTTTG TGCCAAATCT CATTAGGCTT AAGGTAATAC AGGACTTTGA GTCAAATGAT 60

ACTGTTGCAC ATAAGAACAA ACCTATTTTC ATGCTAAGAT GATGCCACTG TGTTCCTTTC 120

TCCTTCTAG TTT CTG GAC ATC TCC ATT AAG TGG ACC ACC CAG GAA ACG TTT 171

Phe Leu Asp Ile Ser Ile Lys Trp Thr Thr Gln Glu Thr Phe

-10

-5

1

CCT CCA AAG TAC CTT CAT TAT GAC GAA GAA ACC TCT CAT CAG CTG TTG 219

Pro Pro Lys Tyr Leu His Tyr Asp Glu Glu Thr Ser His Gln Leu Leu

5

10

15



TGT GAC AAA TGT CCT CCT GGT ACC TAC CTA AAA CAA CAC TGT ACA GCA 267  
 Cys Asp Lys Cys Pro Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala  
 20 25 30 35

AAG TGG AAG ACC GTG TGC GCC CCT TGC CCT GAC CAC TAC TAC ACA GAC 315  
 Lys Trp Lys Thr Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp  
 40 45 50

AGC TGG CAC ACC AGT GAC GAG TGT CTA TAC TGC AGC CCC GTG TGC AAG 363  
 Ser Trp His Thr Ser Asp Glu Cys Leu Tyr Cys Ser Pro Val Cys Lys  
 55 60 65

GAG CTG CAG TAC GTC AAG CAG GAG TGC AAT CGC ACC CAC AAC CGC GTG 411  
 Glu Leu Gln Tyr Val Lys Gln Glu Cys Asn Arg Thr His Asn Arg Val  
 70 75 80

TGC GAA TGC AAG GAA GGG CGC TAC CTT GAG ATA GAG TTC TGC TTG AAA 459  
 Cys Glu Cys Lys Glu Gly Arg Tyr Leu Glu Ile Glu Phe Cys Leu Lys  
 85 90 95

CAT AGG AGC TGC CCT CCT GGA TTT GGA GTG GTG CAA GCT G GTACGTGTCA 509  
 His Arg Ser Cys Pro Pro Gly Phe Gly Val Val Gln Ala  
 100 105 110

ATGTGCAGCA AAATTAATTA GGATCATGCA AAGTCAGATA GTTGTGACAG TTTAGGAGAA 569



CACTTTTGTT CTGATGACAT TATAGGATAG CAAATTGCAA AGGTAATGAA ACCTGCCAGG 629  
 TAGGTACTAT GTGTCTGGAG TGCTTCCAAA GGACCATTGC TCAGAGGAAT ACTTTGCCAC 689  
 TACAGGGCAA TTTAATGACA AATCTCAAAT GCAGCAAATT ATTCTCTCAT GAGATGCATG 749  
 ATGGTTTTTT TTTTTTTTTT TAAAGAAACA AACTCAAGTT GCACTATTGA TAGTTGATCT 809  
 ATACCTCTAT ATTTCACTTC AGCATGGACA CCTTCAAACCT GCAGCACTTT TTGACAAACA 869  
 TCAGAAATGT TAATTTATAC CAAGAGAGTA ATTATGCTCA TATTAATCAG ACTCTGGAGT 929  
 GCTAACAATA AGCAGTTATA ATTAATTATG TAAAAAATGA GAATGGTCAG GCGAATTGCA 989  
 TTTCAATTATT AAAACAAGG CTAGTTCTTC CTTTAGCATG GGAGCTGAGT GTTTGGCAGG 1049  
 GTAAGGACTA TAGCAGAATC TCTTCAATGA GCTTATTCTT TATCTTAGAC AAAACAGATT 1109  
 GTCAAGCCAA GAGCAAGCAC TTGCCTATAA ACCAAGTGCT TTCTCTTTTG CATTTTGAAC 1169  
 AGCATTGGTC AGGGCTCATG TGTATTGAAT CTTTAAACC AGTAACCCAC GTTTTTTTTC 1229  
 TGCCACATTT GCGAAGCTTC AGTGCAGCCT ATAACCTTTC ATAGCTTGAG AAAATTAAGA 1289  
 GTATCCACTT ACTTAGATGG AAGAAGTAAT CAGTATAGAT TCTGATGACT CAGTTTGAAG 1349  
 CAGTGTTTCT CAACTGAAGC CCTGCTGATA TTTAAGAAA TATCTCGATT CCTAGGCTCG 1409  
 ACTCCTTTTT GTGGGCAGCT GTCCTGCGCA TTGTAGAATT TTGGCAGCAC CCCTGGACTC 1469  
 TAGCCACTAG ATACCAATAG CAGTCCTTCC CCCATGTGAC AGCCAAAAT GTCTTCAGAC 1529  
 ACTGTCAAAT GTCGCCAGGT GGCAAAATCA CTCCTGGTTG AGAACAGGGT CATCAATGCT 1589  
 AAGTATCTGT AACTATTTTA ACTCTCAAAA CTTGTGATAT ACAAAGTCTA AATTATTAGA 1649  
 CGACCAATAC TTTAGGTTTA AAGGCATACA AATGAAACAT TCAAAAATCA AAATCTATTC 1709  
 TGTCTCTCAA ATAGTGAATC TTATAAAATT AATCACAGAA GATGCAAATT GCATCAGAGT 1769  
 CCCTTAAAAT TCCTCTTCGT ATGAGTATTT GAGGGAGGAA TTGGTGATAG TTCCTACTTT 1829  
 CTATTGGATG GTACTTTGAG ACTCAAAGC TAAGCTAAGT TGTGTGTGTG TCAGGGTGCG 1889  
 GGGTGTGGA TCCCATCAGA TAAAAGCAAA TCCATGTAAT TCATTAGTA AGTTGTATAT 1949  
 GTAGAAAAAT GAAAAGTGGG CTATGCAGCT TCGAAACTAG AGAATTTTGA AAAATAATGG 2009  
 AAATCACAAG GATCTTTCTT AAATAAGTAA GAAAATCTGT TTGTAGAATG AAGCAAGCAG 2069  
 GCAGCCAGAA GACTCAGAAC AAAAGTACAC ATTTTACTCT GTGTACTG GCAGCACAGT 2129



GGGATTTATT TACCTCTCCC TCCCTAAAAA CCCACACAGC GGTTCCTCTT GGGAAATAAG 2189  
 AGGTTTCCAG CCCAAAGAGA AGGAAAGACT ATGTGGTGT ACTCTAAAAA GTATTTAATA 2249  
 ACCGTTTTGT TGTGCTGTT GCTGTTTTGA AATCAGATTG TCTCCTCTCC ATATTTTATT 2309  
 TACTTCATTC TGTTAATTCC TGTGGAATTA CTTAGAGCAA GCATGGTGAA TTCTCAACTG 2369  
 TAAAGCCAAA TTTCTCCATC ATTATAATTT CACATTTTGC CTGGCAGGTT ATAATTTTTA 2429  
 TATTTCCACT GATAGTAATA AGGTAAATC ATTACTTAGA TGGATAGATC TTTTTCATAA 2489  
 AAAGTACCAT CAGTTATAGA GGAAGTCAT GTTCATGTTT AGGAAGGTCA TTAGATAAAG 2549  
 CTTCTGAATA TATTATGAAA CATTAGTTCT GTCATTCTTA GATTCTTTTT GTTAAATAAC 2609  
 TTTAAAAGCT AACTTACCTA AAAGAAATAT CTGACACATA TGAACCTCTC ATTAGGATGC 2669  
 AGGAGAAGAC CCAAGCCACA GATATGTATC TGAAGAATGA ACAAGATTCT TAGGCCCGGC 2729  
 ACGGTGGCTC ACATCTGTAA TCTCAAGAGT TTGAGAGGTC AAGCGGGCA GATCACCTGA 2789  
 GGTCAAGGAGT TCAAGACCAG CCTGGCCAAC ATGATGAAAC CCTGCCTCTA CTAAAAATAC 2849  
 AAAAATTAGC AGGGCATGGT GGTGCATGCC TGCAACCCTA GCTACTCAGG AGGCTGAGAC 2909  
 AGGAGAATCT CTTGAACCCT CGAGGCGGAG GTTGTGCTGA GCTGAGATCC CTCTACTGCA 2969  
 CTCCAGCCTG GGTGACAGAG ATGAGACTCC GTCCCTGCCG CCGCCCCCGC CTTCCCCCCC 3029  
 AAAAAGATTC TTCTTCATGC AGAACATACG GCAGTCAACA AAGGGAGACC TGGGTCCAGG 3089  
 TGTCCAAGTC ACTTATTTTC AGTAAATTAG CAATGAAAGA ATGCCATGGA ATCCCTGCCC 3149  
 AAATACCTCT GCTTATGATA TTGTAGAATT TGATATAGAG TTGTATCCCA TTTAAGGAGT 3209  
 AGGATGTAGT AGGAAAGTAC TAAAAACAAA CACACAAACA GAAAACCCTC TTTGCTTTGT 3269  
 AAGGTGGTTC CTAAGATAAT GTCAGTGCAA TGCTGGAAT AATATTTAAT ATGTGAAGGT 3329  
 TTTAGGCTGT GTTTTCCCTT CCTGTTCTTT TTTTCTGCCA GCCCTTTGTC ATTTTGCAG 3389  
 GTCAATGAAT CATGTAGAAA GAGACAGGAG ATGAACTAG AACCAGTCCA TTTGCCCTT 3449  
 TTTTTTATTT TCTGGTTTTG GTAAAAGATA CAATGAGGTA GGAGGTTGAG ATTTATAAAT 3509  
 GAAGTTTAAT AAGTTTCTGT AGCTTTGATT TTTCTCTTC ATATTTGTTA TCTTCATAA 3569  
 GCCAGAATTG GCCTGTAAAA TCTACATATG GATATTGAAG TCTAAATCTG TTCAACTAGC 3629  
 TTACACTAGA TGGAGATATT TTCATATTCA GATACACTGG AATGTATGAT CTAGCCATGC 3689



GTAATATAGT CAAGTGTGTTG AAGGTATTTA TTTTAAATAG CGTCTTTAGT TGTGGACTGG 3749  
 TTCAAGTTTT TCTGCCAATG ATTTCTTCAA ATTTATCAAA TATTTTTCCA TCATGAAGTA 3809  
 AAATGCCCTT GCAGTCACCC TTCCTGAAGT TTGAACGACT CTGCTGTTTT AAACAGTTTA 3869  
 AGCAAATGGT ATATCATCTT CCGTTTACTA TGTAGCTTAA CTGCAGGCTT ACGCTTTTGA 3929  
 GTCAGCGGCC AACTTTATTG CCACCTTCAA AAGTTTATTA TAATGTTGTA AATTTTTACT 3989  
 TCTCAAGGTT AGCATACTTA GGAGTTGCTT CACAATTAGG ATTCAGGAAA GAAAGAACTT 4049  
 CAGTAGGAAC TGATTGGAAT TTAATGATGC AGCATTCAAT GGGTACTAAT TTCAAAGAAT 4109  
 GATATTACAG CAGACACACA GCAGTTATCT TGATTTTCTA GGAATAATTG TATGAAGAAT 4169  
 ATGGCTGACA ACACGGCCTT ACTGCCACTC AGCGGAGGCT GGACTAATGA ACACCCTACC 4229  
 CTTCTTTCCT TTCCTCTCAC ATTTTCATGAG CGTTTTGTAG GTAACGAGAA AATTGACTTG 4289  
 CATTTCATT ACAAGGAGGA GAAACTGGCA AAGGGGATGA TGGTGGAAGT TTTGTTCTGT 4349  
 CTAATGAAGT GAAAAATGAA AATGCTAGAG TTTTGTGCAA CATAATAGTA GCAGTAAAAA 4409  
 CCAAGTGAAA AGTCTTTCCA AAAGTGTGTT AAGAGGGCAT CTGCTGGGAA ACGATTTGAG 4469  
 GAGAAGGTAC TAAATTGCTT GGTATTTTCC GTAG GA ACC CCA GAG CGA AAT ACA 4523

Gly Thr Pro Glu Arg Asn Thr

115

GTT TGC AAA AGA TGT CCA GAT GGG TTC TTC TCA AAT GAG ACG TCA TCT 4571  
 Val Cys Lys Arg Cys Pro Asp Gly Phe Phe Ser Asn Glu Thr Ser Ser  
 120 125 130 135

AAA GCA CCC TGT AGA AAA CAC ACA AAT TGC AGT GTC TTT GGT CTC CTG 4619  
 Lys Ala Pro Cys Arg Lys His Thr Asn Cys Ser Val Phe Gly Leu Leu  
 140 145 150

CTA ACT CAG AAA GGA AAT GCA ACA CAC GAC AAC ATA TGT TCC GGA AAC 4667



Leu Thr Gln Lys Gly Asn Ala Thr His Asp Asn Ile Cys Ser Gly Asn

155

160

165


AGT GAA TCA ACT CAA AAA TGT GGA ATA G GTAATTACAT TCCAAAATAC 4715

Ser Glu Ser Thr Gln Lys Cys Gly Ile

170

175

GTCTTTGTAC GATTTTGTAG TATCATCTCT CTCTCTGAGT TGAACACAAG GCCTCCAGCC 4775  
ACATTCTTGG TCAAACCTTAC ATTTTCCCTT TCTTGAATCT TAACCAGCTA AGGCTACTCT 4835  
CGATGCATTA CTGCTAAAGC TACCACTCAG AATCTCTCAA AAATCATCT TCTCACAGAT 4895  
AACACCTCAA AGCTTGATTT TCTCTCCTTT CACACTGAAA TCAAATCTTG CCCATAGGCA 4955  
AAGGGCAGTG TCAAGTTTGC CACTGAGATG AAATTAGGAG AGTCCAAACT GTAGAATTCA 5015  
CGTTGTGTGT TATTACTTTC ACGAATGTCT GTATTATTAA CTAAAGTATA TATTGGCAAC 5075  
TAAGAAGCAA AGTGATATAA ACATGATGAC AAATTAGGCC AGGCATGGTG GCTTACTCCT 5135  
ATAATCCCAA CATTTTGGGG GGCCAAGGTA GGCAGATCAC TTGAGGTCAG GATTTCAAGA 5195  
CCAGCCTGAC CAACATGGTG AAACCTTGTC TCTACTAAAA ATACAAAAAT TAGCTGGGCA 5255  
TGGTAGCAGG CACTTCTAGT ACCAGCTACT CAGGGCTGAG GCAGGAGAAT CGTTGAACC 5315  
CAGGAGATGG AGGTTGCAGT GAGCTGAGAT TGTACCACTG CACTCCAGTC TGGGCAACAG 5375  
AGCAAGATTT CATCACACAC ACACACACAC ACACACACAC ACACATTAGA AATGTGTACT 5435  
TGGCTTTGTT ACCTATCGTA TTAGTGCATC TATTGCATGG AACTTCCAAG CTACTCTGGT 5495  
TGTGTTAAGC TCTTCATTGG GTACAGGTCA CTAGTATTAA GTTCAGGTTA TTCGGATGCA 5555  
TTCCACGGTA GTGATGACAA TTCATCAGGC TAGTGTGTGT GTTCACCTTG TCACTCCCAC 5615  
CACTAGACTA ATCTCAGACC TCACTCAAA GACACATTAC ACTAAAGATG ATTTGCTTTT 5675  
TTGTGTTTAA TCAAGCAATG GTATAAACCA GCTTGACTCT CCCCAAACAG TTTTTCGTAC 5735  
TACAAAGAAG TTTATGAAGC AGAGAAATGT GAATTGATAT ATATATGAGA TTCTAACCCA 5795  
GTTCCAGCAT TGTTTCATTG TGTAATTGAA ATCATAGACA AGCCATTTTA GCCTTTGCTT 5855



TCTTATCTAA AAAAAAAAAA AAAAAAATGA AGGAAGGGGT ATTAAAAGGA GTGATCAAAT 5915  
TTTAACATTC TCTTTAATTA ATTCATTTTT AATTTTACTT TTTTTCATTT ATTGTGCACT 5975  
TACTATGTGG TACTGTGCTA TAGAGGCTTT AACATTTATA AAAACACTGT GAAAGTTGCT 6035  
TCAGATGAAT ATAGGTAGTA GAACGGCAGA ACTAGTATTC AAAGCCAGGT CTGATGAATC 6095  
CAAAAACAAA CACCCATTAC TCCCATTTTC TGGGACATAC TTACTCTACC CAGATGCTCT 6155  
GGGCTTTGTA ATGCCTATGT AAATAACATA GTTTTATGTT TGGTTATTTT CCTATGTAAT 6215  
GTCTACTTAT ATATCTGTAT CTATCTCTTG CTTTGTTCCT AAAGGTAAAC TATGTGTCTA 6275  
AATGTGGGCA AAAAATAACA CACTATTCCA AATTACTGTT CAAATTCCTT TAAGTCAGTG 6335  
ATAATTATTT GTTTTGACAT TAATCATGAA GTTCCCTGTG GGTACTAGGT AAACCTTTAA 6395  
TAGAATGTTA ATGTTTGTAT TCATTATAAG AATTTTTGGC TGTTACTTAT TTACAACAAT 6455  
ATTTCACTCT AATTAGACAT TTACTAACT TTCTCTTGAA AACAATGCCC AAAAAAGAAC 6515  
ATTAGAAGAC ACGTAAGCTC AGTTGGTCTC TGCCACTAAG ACCAGCCAAC AGAAGCTTGA 6575  
TTTTATTCAA ACTTTGCATT TTAGCATATT TTATCTTGGA AAATTCAATT GTGTTGGTTT 6635  
TTTGTTTTTG TTTGTATTGA ATAGACTCTC AGAAATCCAA TTGTTGAGTA AATCTTCTGG 6695  
GTTTTCTAAC CTTTCTTTAG AT GTT ACC CTG TGT GAG GAG GCA TTC TTC AGG 6747

Asp Val Thr Leu Cys Glu Glu Ala Phe Phe Arg

180

185

TTT GCT GTT CCT ACA AAG TTT ACG CCT AAC TGG CTT AGT GTC TTG GTA 6795  
Phe Ala Val Pro Thr Lys Phe Thr Pro Asn Trp Leu Ser Val Leu Val

190

195


200

GAC AAT TTG CCT GGC ACC AAA GTA AAC GCA GAG AGT GTA GAG AGG ATA 6843  
Asp Asn Leu Pro Gly Thr Lys Val Asn Ala Glu Ser Val Glu Arg Ile

205

210

215



AAA CGG CAA CAC AGC TCA CAA GAA CAG ACT TTC CAG CTG CTG AAG TTA 6891  
 Lys Arg Gln His Ser Ser Gln Glu Gln Thr Phe Gln Leu Leu Lys Leu  
 220 225 230 235

TGG AAA CAT CAA AAC AAA GAC CAA GAT ATA GTC AAG AAG ATC ATC CAA G 6940  
 Trp Lys His Gln Asn Lys Asp Gln Asp Ile Val Lys Lys Ile Ile Gln  
 240 245 250

GTATGATAAT CTAAAATAAA AAGATCAATC AGAAATCAAA GACACCTATT TATCATAAAC 7000  
 CAGGAACAAG ACTGCATGTA TGTTTAGTTG TGTGGATCTT GTTCCCTGT TGGAA TCATT 7060  
 GTTGGACTGA AAAAGTTTCC ACCTGATAAT GTAGATGTGA TTCCACAAAC AGTTATACAA 7120  
 GGTTTTGTTT TCACCCCTGC TCCCAGTTT CCTTGTAAG TATGTTGAAC ACTCTAAGAG 7180  
 AAGAGAAATG CATTTGAAGG CAGGGCTGTA TCTCAGGGAG TCGCTTCCAG ATCCCTTAAC 7240  
 GCTTCTGTAA GCAGCCCTC TAGACCACCA AGGAGAAGCT CTATAACCAC TTTGTATCTT 7300  
 ACATTGCACC TCTACCAAGA AGCTCTGTTG TATTTACTTG GTAATTCTCT CCAGGTAGGC 7360  
 TTTTCGTAGC TTACAAATAT GTTCTTATTA ATCCTCATGA TATGGCCTGC ATTAAAATTA 7420  
 TTTTAATGGC ATATGTTATG AGAATTAATG AGATAAAATC TGAAAAGTGT TTGAGCCTCT 7480  
 TGTAGGAAAA AGCTAGTTAC AGCAAAATGT TCTCACATCT TATAAGTTTA TATAAAGATT 7540  
 CTCCTTTAGA AATGGTGTGA GAGAGAAACA GAGAGAGATA GGGAGAGAAG TGTGAAAGAA 7600  
 TCTGAAGAAA AGGAGTTTCA TCCAGTGTGG ACTGTAAGCT TTACGACACA TGATGGAAAG 7660  
 AGTTCTGACT TCAGTAAGCA TTGGGAGGAC ATGCTAGAAG AAAAAGGAAG AAGAGTTTCC 7720  
 ATAATGCAGA CAGGGTCAGT GAGAAATTCA TTCAGGTCCT CACCAGTAGT TAAATGACTG 7780  
 TATAGTCTTG CACTACCCTA AAAAATTCA AGTATCTGAA ACCGGGGCAA CAGATTTTAG 7840  
 GAGACCAACG TCTTTGAGAG CTGATTGCTT TTGCTTATGC AAAGAGTAAA CTTTATGTT 7900  
 TTGAGCAAAC CAAAAGTATT CTTTGAACGT ATAATTAGCC CTGAAGCCGA AAGAAAAGAG 7960  
 AAAATCAGAG ACCGTTAGAA TTGGAAGCAA CCAAATCCC TATTTTATAA ATGAGGACAT 8020





TTTAACCCAG AAAGATGAAC CGATTTGGCT TAGGGCTCAC AGATACTAAG TGA CT CATGT 8080  
 CATTAATAGA AATGTTAGTT CCTCCCTCTT AGGTTTGTAC CCTAGCTTAT TACTGAAATA 8140  
 TTCTCTAGGC TGTGTGTCTC CTTTAGTTCC TCGACCTCAT GTCTTTGAGT TTTCAGATAT 8200  
 CCTCCTCATG GAGGTAGTCC TCTGGTGCTA TGTGTATTCT TTAAAGGCTA GTTACGGCAA 8260  
 TTAAC TTATC AACTAGCGCC TACTAATGAA ACTTTGTATT ACAAAGTAGC TAACTTGAAT 8320  
 ACTTTCC TTT TTTTCTGAAA TGTTATGGTG GTAATTTCTC AAAC TTTTTC TTAGAAA ACT 8380  
 GAGAGTGATG TGTCTTATTT TCTACTGTTA ATTTTCAAAA TTAGGAGCTT CTTC CAAAGT 8440  
 TTTGTTGGAT GCCAAAAATA TATAGCATAT TATCTTATTA TAACAAAAAA TATTTATCTC 8500  
 AGTTCTTAGA AATAAATGGT GTCAC TTAAC TCCCTCTCAA AAGAAAAGGT TATCATTGAA 8560  
 ATATAATTAT GAAATTCTGC AAGAACCTTT TGCCTCACGC TTGTTTTATG ATGGCATTGG 8620  
 ATGAATATAA ATGATGTGAA CACTTATCTG GGCTTTTGCT TTATGCAG AT ATT GAC 8676

Asp Ile Asp

CTC TGT GAA AAC AGC GTG CAG CGG CAC ATT GGA CAT GCT AAC CTC ACC 8724  
 Leu Cys Glu Asn Ser Val Gln Arg His Ile Gly His Ala Asn Leu Thr  
 255 260 265 270

TTC GAG CAG CTT CGT AGC TTG ATG GAA AGC TTA CCG GGA AAG AAA GTG 8772  
 Phe Glu Gln Leu Arg Ser Leu Met Glu Ser Leu Pro Gly Lys Lys Val  
 275 280 285

GGA GCA GAA GAC ATT GAA AAA ACA ATA AAG GCA TGC AAA CCC AGT GAC 8820  
 Gly Ala Glu Asp Ile Glu Lys Thr Ile Lys Ala Cys Lys Pro Ser Asp  
 290 295 300

CAG ATC CTG AAG CTG CTC AGT TTG TGG CGA ATA AAA AAT GGC GAC CAA 8868



Gln Ile Leu Lys Leu Leu Ser Leu Trp Arg Ile Lys Asn Gly Asp Gln

305

310

315

GAC ACC TTG AAG GGC CTA ATG CAC GCA CTA AAG CAC TCA AAG ACG TAC 8916

Asp Thr Leu Lys Gly Leu Met His Ala Leu Lys His Ser Lys Thr Tyr

320

325

330

CAC TTT CCC AAA ACT GTC ACT CAG AGT CTA AAG AAG ACC ATC AGG TTC 8964

His Phe Pro Lys Thr Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe

335

340

345

350

CTT CAC AGC TTC ACA ATG TAC AAA TTG TAT CAG AAG TTA TTT TTA GAA 9012

Leu His Ser Phe Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu

355

360

365

ATG ATA GGT AAC CAG GTC CAA TCA GTA AAA ATA AGC TGC TTA 9054

Met Ile Gly Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu

370

375

380

TAACTGGAAA TGGCCATTGA GCTGTTTCCT CACAATTGGC GAGATCCCAT GGATGAGTAA 9114

ACTGTTTCTC AGGCACTTGA GGCTTTCAGT GATATCTTTC TCATTACCAG TGACTAATTT 9174

TGCCACAGGG TACTAAAAGA AACTATGATG TGGAGAAAGG ACTAACATCT CCTCCAATAA 9234

ACCCCAAATG GTTAATCCAA CTGTCAGATC TGGATCGTTA TCTACTGACT ATATTTTCCC 9294

TTATTACTGC TTGCAGTAAT TCAACTGGAA ATTAAAAAAA AAAAATACTAGA CTCCACTGGG 9354

CCTTACTAAA TATGGGAATG TCTAACTTAA ATAGCTTTGG GATTCCAGCT ATGCTAGAGG 9414

CTTTTATTAG AAAGCCATAT TTTTCTGT AAAAGTTACT AATATATCTG TAACACTATT 9474



ACAGTATTGC TATTTATATT CATTGAGATA TAAGATTGG ACATATTATC ATCCTATAAA 9534  
 GAAACGGTAT GACTTAATTT TAGAAAGAAA ATTATATTCT GTTTATTATG ACAAATGAAA 9594  
 GAGAAAATAT ATATTTTAA TGGAAGTTT GTAGCATTTT TCTAATAGGT ACTGCCATAT 9654  
 TTTTCTGTGT GCAGTATTTT TATAATTTTA TCTGTATAAG CTGTAATATC ATTTTATAGA 9714  
 AAATGCATTA TTTAGTCAAT TGTTTAATGT TGGAAAACAT ATGAAATATA AATTATCTGA 9774  
 ATATTAGATG CTCTGAGAAA TTGAATGTAC CTTATTTAAA AGATTTTATG GTTTTATAAC 9834  
 TATATAATG ACATTATTAA AGTTTTCAAA TTATTTTAA TTGCTTTCTC TGTGCTTTT 9894  
 ATTT 9898

Sequence number: 3

Length of sequence: 401

Sequence Type: amino acid

Strandedness: single stranded

Topology: linear

Molecular type: protein

Sequence:

Met Asn Asn Leu Leu Cys Cys Ala Leu Val Phe Leu Asp Ile Ser  
 -20 -15 -10  
 Ile Lys Trp Thr Thr Gln Glu Thr Phe Pro Pro Lys Tyr Leu His  
 -5 1 5  
 Tyr Asp Glu Glu Thr Ser His Gln Leu Leu Cys Asp Lys Cys Pro  
 10 15 20  
 Pro Gly Thr Tyr Leu Lys Gln His Cys Thr Ala Lys Trp Lys Thr  
 25 30 35  
 Val Cys Ala Pro Cys Pro Asp His Tyr Tyr Thr Asp Ser Trp His  
 40 45 50



Ile Gln Asp Ile Asp Leu Cys Glu Asn Ser Val Gln Arg His Ile		
250	255	260
Gly His Ala Asn Leu Thr Phe Glu Gln Leu Arg Ser Leu Met Glu		
265	270	275
Ser Leu Pro Gly Lys Lys Val Gly Ala Glu Asp Ile Glu Lys Thr		
280	285	290
Ile Lys Ala Cys Lys Pro Ser Asp Gln Ile Leu Lys Leu Leu Ser		
295	300	305
Leu Trp Arg Ile Lys Asn Gly Asp Gln Asp Thr Leu Lys Gly Leu		
310	315	320
Met His Ala Leu Lys His Ser Lys Thr Tyr His Phe Pro Lys Thr		
325	330	335
Val Thr Gln Ser Leu Lys Lys Thr Ile Arg Phe Leu His Ser Phe		
340	345	350
Thr Met Tyr Lys Leu Tyr Gln Lys Leu Phe Leu Glu Met Ile Gly		
355	360	365
Asn Gln Val Gln Ser Val Lys Ile Ser Cys Leu		
370	375	380

Sequence number: 4

Length of sequence: 1206

Sequence Type: nucleic acid

Strandedness: single stranded

Topology: linear

Molecular type: cDNA



Sequence:

ATGAACAAC	TGCTGTGCTG	CGCGCTCGTG	TTTCTGGACA	TCTCCATTAA	GTGGACCACC	60
CAGGAAACG	TTCCTCCAAA	GTACCTTCAT	TATGACGAAG	AAACCTCTCA	TCAGCTGTTG	120
TGTGACAAAT	GTCCTCCTGG	TACCTACCTA	AAACAACACT	GTACAGCAAA	GTGGAAGACC	180
GTGTGCGCCC	CTTGCCCTGA	CCACTACTAC	ACAGACAGCT	GGCACACCAG	TGACGAGTGT	240
CTATACTGCA	GCCCCGTGTG	CAAGGAGCTG	CAGTACGTCA	AGCAGGAGTG	CAATCGCACC	300
CACAACCGCG	TGTGCGAATG	CAAGGAAGGG	CGCTACCTTG	AGATAGAGTT	CTGCTTGAAA	360
CATAGGAGCT	GCCCTCCTGG	ATTTGGAGTG	GTGCAAGCTG	GAACCCGAGA	GCGAAATACA	420
GTTTGCAAAA	GATGTCCAGA	TGGGTTCTTC	TCAAATGAGA	CGTCATCTAA	AGCACCTGT	480
AGAAAACACA	CAAATTGCAG	TGTCTTTGGT	CTCCTGCTAA	CTCAGAAAGG	AAATGCAACA	540
CACGACAACA	TATGTTCCGG	AAACAGTGAA	TCAACTCAAA	AATGTGGAAT	AGATGTTACC	600
CTGTGTGAGG	AGGCATTCTT	CAGGTTTGCT	GTTCTACAA	AGTTTACGCC	TAAGTGGCTT	660
AGTGTCTTGG	TAGACAATTT	GCCTGGCACC	AAAGTAAACG	CAGAGAGTGT	AGAGAGGATA	720
AAACGGCAAC	ACAGCTCACA	AGAACAGACT	TTCCAGCTGC	TGAAGTTATG	GAAACATCAA	780
AACAAAGACC	AAGATATAGT	CAAGAAGATC	ATCCAAGATA	TTGACCTCTG	TGAAAACAGC	840
GTGCAGCGGC	ACATTGGACA	TGCTAACCTC	ACCTTCGAGC	AGCTTCGTAG	CTTGATGGAA	900
AGCTTACCGG	GAAAGAAAGT	GGGAGCAGAA	GACATTGAAA	AAACAATAAA	GGCATGCAAA	960
CCCAGTGACC	AGATCCTGAA	GCTGCTCAGT	TTGTGGCGAA	TAAAAAATGG	CGACCAAGAC	1020
ACCTTGAAGG	GCCTAATGCA	CGCACTAAAG	CACTCAAAGA	CGTACCACTT	TCCCAAAACT	1080
GTCACCTCAGA	GTCTAAAGAA	GACCATCAGG	TTCCTTCACA	GCTTCACAAT	GTACAAATTG	1140
TATCAGAAGT	TATTTTTAGA	AATGATAGGT	AACCAGGTCC	AATCAGTAAA	AATAAGCTGC	1200
TTATAA						1206



CLAIMS:

1. A DNA comprising the nucleotide sequences of the Sequences No. 1 and No. 2 in the Sequence Table.

2. The DNA according to claim 1, wherein the Sequence ID No. 1 includes the first exon of the OCIF gene and the Sequence ID No. 2 includes the second, third, fourth, and fifth exons.

3. A protein exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts and having the following physicochemical characteristics,

(a) molecular weight (SDS-PAGE):

(i) Under reducing conditions: about 60 kD,

(ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 in the Sequence Table,

(c) affinity:


exhibits affinity to a cation exchanger and heparin, and

(d) heat stability:

(i) the osteoclastogenesis-inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,

(ii) the osteoclastogenesis-inhibitory activity is lost when treated with heat at 90°C for 10 minutes.

4. A process for producing a protein exhibiting an



activity of inhibiting differentiation and/or maturation of osteoclasts and having the following physicochemical characteristics,

(a) molecular weight (SDS-PAGE):

(i) Under reducing conditions: about 60 kD,

(ii) Under non-reducing conditions: about 60 kD and about 120 kD;

(b) amino acid sequence:

includes an amino acid sequence of the Sequence ID No. 3 of the Sequence Table,

(c) affinity:

exhibits affinity to a cation exchanger and heparin, and

(d) heat stability:

(i) the osteoclastogenesis-inhibitory activity is reduced when treated with heat at 70°C for 10 minutes or at 56°C for 30 minutes,

(ii) the osteoclastogenesis-inhibitory activity is lost when treated with heat at 90°C for 10 minutes,

the process comprising inserting a DNA including the nucleotide sequences of the sequences No. 1 and No. 2 in the Sequence Table into an expression vector, producing a vector capable of expressing a protein having the above-mentioned physicochemical characteristics and exhibiting the activity of inhibiting differentiation and/or maturation of osteoclasts, and producing this protein by a genetic engineering technique.



Figure 1

