

(51) International Patent Classification:
C12N 15/82 (2006.01)(21) International Application Number:
PCT/US2009/002547(22) International Filing Date:
24 April 2009 (24.04.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/125,908 29 April 2008 (29.04.2008) US(71) Applicant (for all designated States except US): **MON-SANTO TECHNOLOGY, LLC** [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **COFFIN, Marie** [US/US]; Monsanto Technology, LLC, 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).(74) Agent: **HUELSKAMP, Nan**; Cy-mail Zone E2NA, Monsanto Technology, LLC, 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: GENES AND USES FOR PLANT ENHANCEMENT




FIG. #1

(57) Abstract: Transgenic seed for crops with enhanced agronomic traits are provided by trait-improving recombinant DNA in the nucleus of cells of the seed where plants grown from such transgenic seed exhibit one or more enhanced traits as compared to a control plant. Of particular interest are transgenic plants that have increased yield. The present invention also provides recombinant DNA molecules for expression of a protein, and recombinant DNA molecules for suppression of a protein.

GENES AND USES FOR PLANT ENHANCEMENT

INCORPORATION OF SEQUENCE LISTING

Two copies of the sequence listing (Copy 1 and Copy 2) and a computer readable
5 form (CRF) of the sequence listing, all on CD-Rs, each containing the file named 38-
21_54976_PCT_seqlisting.txt, which is 319,329,870 bytes (measured in MS-WINDOWS)
and was created on April 24, 2009, are incorporated herein by reference in their entirety.

INCORPORATION OF COMPUTER PROGRAM LISTING

A Computer Program Listing folder named 38-21(54976)-PCT_programListing
10 containing folders "hmmmer-2.3.2" and "495pfamDir" is contained on a CD-R and is
incorporated herein by reference in their entirety. Folder hmmmer-2.3.2 contains the source
code and other associated file for implementing the HMMer software for Pfam analysis.
Folder 495pfamDir contains 495 profile Hidden Markov Models. Both folders were created
on the disk on April 24, 2009 having a total size of 39,755,623 bytes when measured in MS-
15 WINDOWS ® operating system.

INCORPORATION OF TABLES

Two copies of Table 19 (Copy 1 and Copy 2) and a computer readable form (CRF),
all on CD-Rs, each containing the file named 38-21_54976_PCT_table19.txt, which is 864
KB (measured in MS-WINDOWS), were created on April 24, 2009, and comprise 417 pages
20 when viewed in MS Word, are herein incorporated by reference.

FIELD OF THE INVENTION

Disclosed herein are recombinant DNA for providing enhanced traits to
transgenic plants, seeds, pollen, plant cells and plant nuclei of such transgenic plants,
methods of making and using such recombinant DNA, plants, seeds, pollen, plant cells and
25 plant nuclei. Also disclosed are methods of producing hybrid seed comprising such
recombinant DNA.

SUMMARY OF THE INVENTION

This invention provides recombinant DNA comprising polynucleotides characterized
by SEQ ID NO: 1-803 and the cognate amino acid sequences of SEQ ID NO: 804-1606.
30 The recombinant DNA is used for providing enhanced traits when stably integrated into the
chromosomes and expressed in the nuclei of transgenic plants cells. In some aspects the

recombinant DNA encodes a protein; in other aspects the recombinant DNA is transcribed to RNA that suppresses the expression of a native gene.

Such recombinant DNA in a plant cell nucleus of this invention is provided in as a construct comprising a promoter that is functional in plant cells and that is operably linked to DNA that encodes a protein or to DNA that results in gene suppression. Such DNA in the construct is sometimes defined by protein domains of an encoded protein targeted for production or suppression *e.g.* a “Pfam domain module” (as defined herein below) from the group of Pfam domain modules identified in Table 20. Alternatively, *e.g.* where a Pfam domain module is not available, such DNA in the construct is defined a consensus amino acid sequence of an encoded protein that is targeted for production *e.g.* a protein having amino acid sequence with at least 90% identity to a consensus amino acid sequence in the group of SEQ ID NO: 94617 through SEQ ID NO: 94734. In a particular aspect of the invention the recombinant DNA is characterized by its cognate amino acid sequence that has at least 70% identity to any of SEQ ID NO: 804-1606.

This invention also provides transgenic plant cell nuclei comprising the recombinant DNA of the invention, transgenic plant cells comprising such nuclei, transgenic plants comprising a plurality of such transgenic plant cells, and transgenic seeds and transgenic pollen of such plants. Such transgenic plants are selected from a population of transgenic plants regenerated from plant cells transformed with recombinant DNA by screening transgenic plants for an enhanced trait as compared to control plants. The enhanced trait is one or more of enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced heat tolerance, enhanced shade tolerance, enhanced high salinity tolerance, enhanced seed protein and enhanced seed oil. Such recombinant DNA in a plant cell nucleus of this invention is provided in as a construct comprising a promoter that is functional in plant cells and that is operably linked to DNA that encodes a protein or to DNA that results in gene suppression. Such DNA in the construct is sometimes defined by protein domains of an encoded protein targeted for production or suppression, *e.g.* a “Pfam domain module” (as defined herein below) from the group of Pfam domain modules identified in Table 20. Alternatively, *e.g.* where a Pfam domain module is not available, such DNA in the construct is defined a consensus amino acid sequence of an encoded protein that is targeted for production *e.g.* a protein having amino acid sequence with at least 90% identity to a consensus amino acid sequence in the group of SEQ ID NO: 94617 through SEQ ID NO: 94734.

In another aspect of the invention the plant cell nuclei, cells, plants, seeds, and pollen further comprise DNA expressing a protein that provides tolerance from exposure to an herbicide applied at levels that are lethal to a wild type plant cell.

This invention also provides methods for manufacturing non-natural, transgenic seed
5 that can be used to produce a crop of transgenic plants with an enhanced trait resulting from expression of stably-integrated, recombinant DNA in the nucleus of the plant cells. More specifically the method comprises (a) screening a population of plants for an enhanced trait and recombinant DNA, where individual plants in the population can exhibit the trait at a level less than, essentially the same as or greater than the level that the trait is exhibited in
10 control plants which do not express the recombinant DNA; (b) selecting from the population one or more plants that exhibit the trait at a level greater than the level that said trait is exhibited in control plants and (c) collecting seed from a selected plant. Such method further comprises steps (a) verifying that the recombinant DNA is stably integrated in said selected plants; and (b) analyzing tissue of a selected plant to determine the production of a
15 protein having the function of a protein encoded by a recombinant DNA with a sequence of one of SEQ ID NO: 1-803; in one aspect of the invention the plants in the population further comprise DNA expressing a protein that provides tolerance to exposure to an herbicide applied at levels that are lethal to wild type plant cells and where the selecting is effected by treating the population with the herbicide, *e.g.* a glyphosate, dicamba, or glufosinate
20 compound. In another aspect of the invention the plants are selected by identifying plants with the enhanced trait. The methods are used for manufacturing corn, soybean, cotton, canola, alfalfa, wheat, rice seed or any combinations thereof selected as having one of the enhanced traits described above.

Another aspect of the invention provides a method of producing hybrid corn seed
25 comprising acquiring hybrid corn seed from a herbicide tolerant corn plant which also has a nucleus of this invention with stably-integrated, recombinant DNA. The method further comprises producing corn plants from said hybrid corn seed, where a fraction of the plants produced from said hybrid corn seed is homozygous for said recombinant DNA, a fraction of the plants produced from said hybrid corn seed is hemizygous for said recombinant DNA,
30 and a fraction of the plants produced from said hybrid corn seed has none of said recombinant DNA; selecting corn plants which are homozygous and hemizygous for said recombinant DNA by treating with an herbicide; collecting seed from herbicide-treated-surviving corn plants and planting said seed to produce further progeny corn plants; repeating the selecting

and collecting steps at least once to produce an inbred corn line; and crossing the inbred corn line with a second corn line to produce hybrid seed.

Still other aspects of this invention relate to transgenic plants with enhanced water use efficiency or enhanced nitrogen use efficiency. For instance, this invention provides methods of growing a corn, cotton, soybean, or canola crop without irrigation water comprising planting seed having plant cells of the invention which are selected for enhanced water use efficiency. Alternatively methods comprise applying reduced irrigation water, *e.g.* providing up to 300 millimeters of ground water during the production of a corn crop. This invention also provides methods of growing a corn, cotton, soybean or canola crop without added nitrogen fertilizer comprising planting seed having plant cells of the invention which are selected for enhanced nitrogen use efficiency.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1, 2 and 3 illustrate plasmid maps.

Figure 4 illustrates a consensus amino acid sequence of SEQ ID NO: 1325 and its homologs.

DETAILED DESCRIPTION OF THE INVENTION

In the attached sequence listing:

SEQ ID NO: 1-803 are nucleotide sequences of the coding strand of DNA for “genes” used in the recombinant DNA imparting an enhanced trait in plant cells, where each represents a coding sequence for a protein;

SEQ ID NO: 804-1606 are amino acid sequences of the cognate protein of the “genes” with nucleotide coding sequence 1-803;

SEQ ID NO: 1607-94613 are amino acid sequences of homologous proteins;

SEQ ID NO: 94614 is a nucleotide sequence of a plasmid base vector for corn transformation; and

SEQ ID NO: 94615 is a DNA sequence of a plasmid base vector for soybean transformation.

SEQ ID NO: 94616 is a DNA sequence of a plasmid base vector for cotton transformation.

SEQ ID NO: 94617-94734 are consensus sequences.

Table 1 lists the protein SEQ ID NOs and their corresponding consensus SEQ ID NOs.

Table 1.

NUC SEQ ID	PEP SEQ ID NO	GENE ID	CONSENSUS SEQ
------------	---------------	---------	---------------

NO			ID NO
5	808	CGPG1067	94617
6	809	CGPG1095	94618
7	810	CGPG1101	94619
8	811	CGPG1109	94620
10	813	CGPG1155	94621
12	815	CGPG1171	94622
13	816	CGPG1177	94623
19	822	CGPG1279	94624
20	823	CGPG1304	94625
22	825	CGPG1327	94626
57	860	CGPG1723	94627
78	881	CGPG1972	94628
88	891	CGPG2041	94629
90	893	CGPG2070	94630
98	901	CGPG2126	94631
99	902	CGPG213	94632
106	909	CGPG2193	94633
108	911	CGPG2224	94634
118	921	CGPG2319	94635
121	924	CGPG2359	94636
130	933	CGPG2408	94637
133	936	CGPG2414	94638
156	959	CGPG2863	94639
172	975	CGPG3183	94640
180	983	CGPG3280	94641
191	994	CGPG3405	94642
200	1003	CGPG3598	94643
202	1005	CGPG3606	94644
203	1006	CGPG3610	94645
204	1007	CGPG3620	94646
222	1025	CGPG3820	94647
226	1029	CGPG3958	94648
227	1030	CGPG3981	94649
236	1039	CGPG4083	94650
256	1059	CGPG4347	94651
257	1060	CGPG4354	94652
273	1076	CGPG4559	94653
294	1097	CGPG4802	94654
295	1098	CGPG4822	94655
296	1099	CGPG4826	94656
297	1100	CGPG4833	94657
298	1101	CGPG4841	94658
332	1135	CGPG5208	94659
393	1196	CGPG5750	94660
394	1197	CGPG5751	94661
413	1216	CGPG5922	94662
415	1218	CGPG5964	94663

432	1235	CGPG6137	94664
456	1259	CGPG6358	94665
501	1304	CGPG6803	94666
504	1307	CGPG6812	94667
506	1309	CGPG6827	94668
509	1312	CGPG6859	94669
514	1317	CGPG6886	94670
518	1321	CGPG6903	94671
520	1323	CGPG6933	94672
522	1325	CGPG6944	94673
523	1326	CGPG6949	94674
525	1328	CGPG6989	94675
527	1330	CGPG7003	94676
528	1331	CGPG7009	94677
531	1334	CGPG7062	94678
532	1335	CGPG7064	94679
534	1337	CGPG7109	94680
535	1338	CGPG7123	94681
538	1341	CGPG7159	94682
541	1344	CGPG7224	94683
569	1372	CGPG7505	94684
576	1379	CGPG7540	94685
582	1385	CGPG7568	94686
583	1386	CGPG7571	94687
584	1387	CGPG7587	94688
585	1388	CGPG7591	94689
588	1391	CGPG7633	94690
601	1404	CGPG7774	94691
602	1405	CGPG7777	94692
606	1409	CGPG7792	94693
613	1416	CGPG7851	94694
616	1419	CGPG7865	94695
622	1425	CGPG7934	94696
623	1426	CGPG7949	94697
624	1427	CGPG7954	94698
627	1430	CGPG7969	94699
641	1444	CGPG8072	94700
642	1445	CGPG8075	94701
644	1447	CGPG8100	94702
645	1448	CGPG8101	94703
646	1449	CGPG8104	94704
647	1450	CGPG8108	94705
648	1451	CGPG8116	94706
651	1454	CGPG8138	94707
654	1457	CGPG8163	94708
659	1462	CGPG8204	94709
662	1465	CGPG8234	94710
671	1474	CGPG8345	94711

686	1489	CGPG8515	94712
688	1491	CGPG8518	94713
690	1493	CGPG8523	94714
691	1494	CGPG8528	94715
693	1496	CGPG8553	94716
694	1497	CGPG8562	94717
697	1500	CGPG8589	94718
699	1502	CGPG8592	94719
703	1506	CGPG8648	94720
706	1509	CGPG8678	94721
707	1510	CGPG8694	94722
709	1512	CGPG8737	94723
710	1513	CGPG8745	94724
712	1515	CGPG8749	94725
715	1518	CGPG8781	94726
760	1563	CGPG9075	94727
762	1565	CGPG9094	94728
769	1572	CGPG9136	94729
770	1573	CGPG9156	94730
773	1576	CGPG9173	94731
781	1584	CGPG921	94732
788	1591	CGPG9281	94733
792	1595	CGPG9304	94734

The nuclei of this invention are identified by screening transgenic plants for one or more traits including enhanced drought stress tolerance, enhanced heat stress tolerance, enhanced cold stress tolerance, enhanced high salinity stress tolerance, enhanced low nitrogen availability stress tolerance, enhanced shade stress tolerance, enhanced plant growth and development at the stages of seed imbibition through early vegetative phase, and enhanced plant growth and development at the stages of leaf development, flower production and seed maturity.

“As used herein a “plant cell” means a plant cell that is transformed with stably-integrated, non-natural, recombinant DNA, *e.g.* by *Agrobacterium*-mediated transformation or by bombardment using microparticles coated with recombinant DNA or other means. A plant cell of this invention can be an originally-transformed plant cell that exists as a microorganism or as a progeny plant cell that is regenerated into differentiated tissue, *e.g.* into a transgenic plant with stably-integrated, non-natural recombinant DNA, or seed or pollen derived from a progeny transgenic plant.

As used herein a “transgenic plant” means a plant whose genome has been altered by the stable integration of recombinant DNA. A transgenic plant includes a plant regenerated

from an originally-transformed plant cell and progeny transgenic plants from later generations or crosses of a transformed plant.

As used herein “recombinant DNA” means DNA which has been a genetically engineered and constructed outside of a cell including DNA containing naturally occurring
5 DNA or cDNA or synthetic DNA.

As used herein a “homolog” means a protein in a group of proteins that perform the same biological function, *e.g.* proteins that belong to the same Pfam protein family and that provide a common enhanced trait in transgenic plants of this invention. Homologs are expressed by homologous genes. Homologous genes include naturally occurring alleles and
10 artificially-created variants. Degeneracy of the genetic code provides the possibility to substitute at least one base of the protein encoding sequence of a gene with a different base without causing the amino acid sequence of the polypeptide produced from the gene to be changed. Hence, a polynucleotide in the present invention can have any base sequence that has been changed from SEQ ID NO:1 through SEQ ID NO: 803 through substitution in
15 accordance with degeneracy of the genetic code. Homologs are proteins that, when optimally aligned, have at least about 60% identity, about 70% or higher, at least about 80% and at least about 90% identity over the full length of a protein identified as being associated with imparting an enhanced trait when expressed in plant cells. Homologs include proteins with an amino acid sequence that has at least about 90% identity to a consensus amino acid
20 sequence of proteins and homologs disclosed herein.

Homologs are identified by comparison of amino acid sequence, *e.g.* manually or by use of a computer-based tool using known homology-based search algorithms such as those commonly known and referred to as BLAST, FASTA, and Smith-Waterman. A local sequence alignment program, *e.g.* BLAST, can be used to search a database of sequences to
25 find similar sequences, and the summary Expectation value (E-value) used to measure the sequence base similarity. As a protein hit with the best E-value for a particular organism could be an ortholog or the only ortholog, a reciprocal query is used in the present invention to filter hit sequences with significant E-values for ortholog identification. The reciprocal query entails search of the significant hits against a database of amino acid sequences from
30 the base organism that are similar to the sequence of the query protein. A hit can be identified as an ortholog, when the reciprocal query’s best hit is the query protein itself or a protein encoded by a duplicated gene after speciation. A further aspect of the homologs encoded by DNA in the transgenic plants of the invention are those proteins that differ from a disclosed protein as the result of deletion or insertion of one or more amino acids in a native sequence.

Homologous genes are genes which encode proteins with the same or similar biological function to the protein encoded by the second gene. Homologous genes can be generated by the event of speciation (see ortholog) or by the event of genetic duplication (see paralog). "Orthologs" refer to a set of homologous genes in different species that evolved from a common ancestral gene by specification. Normally, orthologs retain the same function in the course of evolution; and "paralogs" refer to a set of homologous genes in the same species that have diverged from each other as a consequence of genetic duplication. Thus, homologous genes can be from the same or a different organism. As used herein, "homolog" means a protein that performs the same biological function as a second protein including those identified by sequence identity search.

As used herein, "percent identity" means the extent to which two optimally aligned DNA or protein segments are invariant throughout a window of alignment of components, for example nucleotide sequence or amino acid sequence. An "identity fraction" for aligned segments of a test sequence and a reference sequence is the number of identical components that are shared by sequences of the two aligned segments divided by the total number of sequence components in the reference segment over a window of alignment which is the smaller of the full test sequence or the full reference sequence. "Percent identity" ("percent identity") is the identity fraction times 100. Such optimal alignment is understood to be deemed as local alignment of DNA sequences. For protein alignment, a local alignment of protein sequences should allow introduction of gaps to achieve optimal alignment. Percent identity is calculated over the aligned length not including the gaps introduced by the alignment *per se*.

As used herein, "promoter" means regulatory DNA for initializing transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells whether or not its origin is a plant cell, *e.g.* is it well known that *Agrobacterium* promoters are functional in plant cells. Thus, plant promoters include promoter DNA obtained from plants, plant viruses and bacteria such as *Agrobacterium* and *Bradyrhizobium* bacteria. Examples of promoters under developmental control include promoters that preferentially initiate transcription in certain tissues, such as leaves, roots, or seeds. Such promoters are referred to as "tissue preferred". Promoters that initiate transcription only in certain tissues are referred to as "tissue specific". A "cell type" specific promoter primarily drives expression in certain cell types in one or more organs, for example, vascular cells in roots or leaves. An "inducible" or "repressible" promoter is a promoter which is under environmental control. Examples of

environmental conditions that can effect transcription by inducible promoters include anaerobic conditions, or certain chemicals, or the presence of light. Tissue specific, tissue preferred, cell type specific, and inducible promoters constitute the class of "non-constitutive" promoters. A "constitutive" promoter is a promoter which is active under most conditions.

5 As used herein, "operably linked" refers to the association of two or more nucleic acid elements in a recombinant DNA construct, *e.g.* as when a promoter is operably linked with DNA that is transcribed to RNA whether for expressing or suppressing a protein. Recombinant DNA constructs can be designed to express a protein which can be an endogenous protein, an exogenous homologue of an endogenous protein or an exogenous
10 protein with no native homologue. Alternatively, recombinant DNA constructs can be designed to suppress the level of an endogenous protein, *e.g.* by suppression of the native gene. Such gene suppression can be effectively employed through a native RNA interference (RNAi) mechanism in which recombinant DNA comprises both sense and anti-sense oriented DNA matched to the gene targeted for suppression where the recombinant
15 DNA is transcribed into RNA that can form a double-strand to initiate an RNAi mechanism. Gene suppression can also be effected by recombinant DNA that comprises anti-sense oriented DNA matched to the gene targeted for suppression. Gene suppression can also be effected by recombinant DNA that comprises DNA that is transcribed to a microRNA matched to the gene targeted for suppression. In the examples illustrating the invention
20 recombinant DNA for effecting gene suppression that imparts is identified by the term "antisense". It will be understood by a person of ordinary skill in the art that any of the ways of effecting gene suppression are contemplated and enabled by a showing of one approach to gene suppression.

 As used herein "expressed" means produced, *e.g.* a protein is expressed in a plant cell
25 when its cognate DNA is transcribed to mRNA that is translated to the protein.

 As used herein "suppressed" means decreased, *e.g.* a protein is suppressed in a plant cell when there is a decrease in the amount and/or activity of the protein in the plant cell. The presence or activity of the protein can be decreased by any amount up to and including a total loss of protein expression and/or activity.

30 As used herein a "control plant" means a plant that does not contain the recombinant DNA that expressed a protein that imparts an enhanced trait. A control plant is to identify and select a transgenic plant that has an enhance trait. A suitable control plant can be a non-transgenic plant of the parental line used to generate a transgenic plant, *e.g.* devoid of recombinant DNA. A suitable control plant can in some cases be a progeny of a

hemizygous transgenic plant line that does not contain the recombinant DNA, known as a negative segregant.

“Trait enhancement” means a detectable and desirable difference in a characteristic in a transgenic plant relative to a control plant or a reference. In some cases, the trait enhancement can be measured quantitatively. For example, the trait enhancement can entail at least about 2% difference in an observed trait, at least about 5% desirable difference, at least about about 10% difference, at least about about 20% difference, at least about 30% difference, at least about 40%, at least about 50% difference, at least about 60%, at least about 70% difference, at least about 80% difference, at least about 90% difference, or at least about a 100% difference, or an even greater difference. In other cases, the trait enhancement is only measured qualitatively. It is known that there can be a natural variation in a trait. Therefore, the trait enhancement observed entails a change of the normal distribution of the trait in the transgenic plant compared with the trait distribution observed in a control plant or a reference, which is evaluated by statistical methods provided herein. Trait enhancement includes, but is not limited to, yield increase, including increased yield under non-stress conditions and increased yield under environmental stress conditions. Stress conditions can include, for example, drought, shade, fungal disease, viral disease, bacterial disease, insect infestation, nematode infestation, cold temperature exposure, heat exposure, osmotic stress, reduced nitrogen nutrient availability, reduced phosphorus nutrient availability, high plant density, or any combinations thereof.

“Yield” can be affected by many properties including without limitation, plant height, pod number, pod position on the plant, number of internodes, incidence of pod shatter, grain size, efficiency of nodulation and nitrogen fixation, efficiency of nutrient assimilation, resistance to biotic and abiotic stress, carbon assimilation, plant architecture, resistance to lodging, percent seed germination, seedling vigor, and juvenile traits. Yield can also be affected by efficiency of germination (including germination in stressed conditions), growth rate (including growth rate in stressed conditions), ear number, seed number per ear, seed size, composition of seed (starch, oil, protein) and characteristics of seed fill.

Increased yield of a transgenic plant of the present invention can be measured in a number of ways, including test weight, seed number per plant, seed weight, seed number per unit area (*e.g.* seeds, or weight of seeds, per acre), bushels per acre, tonnes per acre, tons per acre, kilo per hectare. For example, maize yield can be measured as production of shelled corn kernels per unit of production area, for example in bushels per acre or metric tons per

hectare, often reported on a moisture adjusted basis, for example at 15.5 percent moisture. Increased yield can result from improved utilization of key biochemical compounds, such as nitrogen, phosphorous and carbohydrate, or from improved responses to environmental stresses, such as cold, heat, drought, salt, and attack by pests or pathogens. Recombinant DNA used in this invention can also be used to provide plants having improved growth and development, and ultimately increased yield, as the result of modified expression of plant growth regulators or modification of cell cycle or photosynthesis pathways. Also of interest is the generation of transgenic plants that demonstrate enhanced yield with respect to a seed component that could correspond to an increase in overall plant yield. Such properties include enhancements in seed oil, seed molecules such as tocopherol, protein and starch, or oil particular oil components as can be manifest by alterations in the ratios of seed components.

“Stress condition” means a condition unfavorable for a plant, which adversely affect plant metabolism, growth and/or development. A plant under the stress condition typically shows reduced germination rate, retarded growth and development, reduced photosynthesis rate, and eventually leading to reduction in yield. Specifically, “water deficit stress” used herein refers to the sub-optimal conditions for water and humidity needed for normal growth of natural plants. Relative water content (RWC) can be used as a physiological measure of plant water deficit. It measures the effect of osmotic adjustment in plant water status, when a plant is under stressed conditions. Conditions which can result in water deficit stress include, but are not limited to, heat, drought, high salinity and PEG induced osmotic stress.

“Cold stress” means the exposure of a plant to a temperatures below (two or more degrees Celsius below) those normal for a particular species or particular strain of plant.

“Nitrogen nutrient” means any one or any mix of the nitrate salts commonly used as plant nitrogen fertilizer, including, but not limited to, potassium nitrate, calcium nitrate, sodium nitrate, ammonium nitrate. The term ammonium as used herein means any one or any mix of the ammonium salts commonly used as plant nitrogen fertilizer, *e.g.*, ammonium nitrate, ammonium chloride, ammonium sulfate, etc.

“Low nitrogen availability stress” means a plant growth condition that does not contain sufficient nitrogen nutrient to maintain a healthy plant growth and/or for a plant to reach its typical yield under a sufficient nitrogen growth condition. For example, a limiting nitrogen condition can refers to a growth condition with 50% or less of the conventional nitrogen inputs. “Sufficient nitrogen growth condition” means a growth condition where the

soil or growth medium contains or receives optimal amounts of nitrogen nutrient to sustain a healthy plant growth and/or for a plant to reach its typical yield for a particular plant species or a particular strain. One skilled in the art would recognize what constitute such soil, media and fertilizer inputs for most plant species.

5 “Shade stress” means a growth condition that has limited light availability that triggers the shade avoidance response in plant. Plants are subject to shade stress when localized at lower part of the canopy, or in close proximity of neighboring vegetation. Shade stress can become exacerbated when the planting density exceeds the average prevailing density for a particular plant species.

10 In some embodiments of the invention a constitutively active mutant, is constructed to achieve the desired effect. In other embodiments of the invention, a dominant negative gene is constructed to adversely affect the normal, wild-type gene product within the same cell.

DNA constructs are assembled using methods well known to persons of ordinary skill in the art and typically comprise a promoter operably linked to DNA, the expression of which provides the enhanced agronomic trait. Other construct components can include additional regulatory elements, such as 5' leaders and introns for enhancing transcription, 3' untranslated regions (such as polyadenylation signals and sites), DNA for transit or signal peptides.

20 Numerous promoters that are active in plant cells have been described in the literature. These include promoters present in plant genomes as well as promoters from other sources, including nopaline synthase (NOS) promoter and octopine synthase (OCS) promoters carried on tumor-inducing plasmids of *Agrobacterium tumefaciens* and the CaMV35S promoters from the cauliflower mosaic virus as disclosed in US Patents No. 5,164,316 and 5,322,938. Promoters derived from plant genes are found in U.S. Patent 25 5,641,876, which discloses a rice actin promoter, U.S. Patent No. 7,151,204, which discloses a maize chloroplast aldolase promoter and a maize aldolase (FDA) promoter, and U.S. Patent Application Publication 2003/0131377 A1, which discloses a maize nicotianamine synthase promoter, all of which are incorporated herein by reference. These and numerous other promoters that function in plant cells are known to those skilled in the art and available for 30 use in recombinant polynucleotides of the present invention to provide for expression of desired genes in transgenic plant cells.

In other aspects of the invention, preferential expression in plant green tissues is desired. Promoters of interest for such uses include those from genes such as Arabidopsis

thaliana ribulose-1,5-bisphosphate carboxylase (Rubisco) small subunit (Fischhoff *et al.* (1992) *Plant Mol Biol.* 20:81-93), aldolase and pyruvate orthophosphate dikinase (PPDK) (Taniguchi *et al.* (2000) *Plant Cell Physiol.* 41(1):42-48).

Furthermore, the promoters can be altered to contain multiple “enhancer sequences” to assist in elevating gene expression. Such enhancers are known in the art. By including an enhancer sequence with such constructs, the expression of the selected protein can be enhanced. These enhancers often are found 5' to the start of transcription in a promoter that functions in eukaryotic cells, but can often be inserted upstream (5') or downstream (3') to the coding sequence. In some instances, these 5' enhancing elements are introns. In one embodiment, enhancers are the 5' introns of the rice actin 1 (see US Patent 5,641,876) and rice actin 2 genes, the maize alcohol dehydrogenase gene intron, the maize heat shock protein 70 gene intron (U.S. Patent 5,593,874) and the maize shrunken 1 gene.

In other aspects of the invention, sufficient expression in plant seed tissues is desired to affect improvements in seed composition. Exemplary promoters for use for seed composition modification include promoters from seed genes such as napin (U.S. 5,420,034), maize L3 oleosin (U.S. 6,433,252), zein Z27 (Russell *et al.* (1997) *Transgenic Res.* 6(2):157-166), globulin 1 (Belanger *et al.* (1991) *Genetics* 129:863-872), glutelin 1 (Russell (1997) *supra*), and peroxiredoxin antioxidant (Per1) (Stacy *et al.* (1996) *Plant Mol Biol.* 31(6):1205-1216).

Recombinant DNA constructs prepared in accordance with the invention will also generally include a 3' element that typically contains a polyadenylation signal and site. Well-known 3' elements include those from *Agrobacterium tumefaciens* genes such as *nos* 3', *tml* 3', *tmr* 3', *tms* 3', *ocs* 3', *tr7* 3', for example disclosed in U.S. 6,090,627, incorporated herein by reference; 3' elements from plant genes such as wheat (*Triticum aestivum*) heat shock protein 17 (*Hsp17* 3'), a wheat ubiquitin gene, a wheat fructose-1,6-bisphosphatase gene, a rice glutelin gene, a rice lactate dehydrogenase gene and a rice beta-tubulin gene, all of which are disclosed in U.S. published patent application 2002/0192813 A1, incorporated herein by reference; and the pea (*Pisum sativum*) ribulose biphosphate carboxylase gene (*rbs* 3'), and 3' elements from the genes within the host plant.

Constructs and vectors can also include a transit peptide for targeting of a gene to a plant organelle, particularly to a chloroplast, leucoplast or other plastid organelle. For descriptions of the use of chloroplast transit peptides see U.S. Patent 5,188,642 and U.S. Patent No. 5,728,925, incorporated herein by reference. For description of the transit peptide

region of an Arabidopsis EPSPS gene can be in the present invention, see Klee, H.J. *et al* (MGG (1987) 210:437-442).

Gene suppression includes any of the well-known methods for suppressing transcription of a gene or the accumulation of the mRNA corresponding to that gene thereby preventing translation of the transcript into protein. Posttranscriptional gene suppression is mediated by transcription of RNA that forms double-stranded RNA (dsRNA) having homology to a gene targeted for suppression. Suppression can also be achieved by insertion mutations created by transposable elements can also prevent gene function. For example, in many dicot plants, transformation with the T-DNA of *Agrobacterium* can be readily achieved and large numbers of transformants can be rapidly obtained. Also, some species have lines with active transposable elements that can efficiently be used for the generation of large numbers of insertion mutations, while some other species lack such options. Mutant plants produced by *Agrobacterium* or transposon mutagenesis and having altered expression of a polypeptide of interest can be identified using the polynucleotides of the present invention. For example, a large population of mutated plants can be screened with polynucleotides encoding the polypeptide of interest to detect mutated plants having an insertion in the gene encoding the polypeptide of interest.

Transgenic plants comprising or derived from plant cells of this invention transformed with recombinant DNA can be further enhanced with stacked traits, *e.g.* a crop plant having an enhanced trait resulting from expression of DNA disclosed herein in combination with herbicide and/or pest resistance traits. For example, genes of the current invention can be stacked with other traits of agronomic interest, such as a trait providing herbicide resistance, or insect resistance, such as using a gene from *Bacillus thuringensis* to provide resistance against lepidopteran, coliopteran, homopteran, hemiopteran, and other insects. Herbicides for which transgenic plant tolerance has been demonstrated and the method of the present invention can be applied include, but are not limited to, glyphosate, dicamba, glufosinate, sulfonylurea, bromoxynil and norflurazon herbicides. Polynucleotide molecules encoding proteins involved in herbicide tolerance are well-known in the art and include, but are not limited to, a polynucleotide molecule encoding 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) disclosed in U.S. Patent 5,094,945; 5,627,061; 5,633,435 and 6,040,497 for imparting glyphosate tolerance; polynucleotide molecules encoding a glyphosate oxidoreductase (GOX) disclosed in U.S. Patent 5,463,175 and a glyphosate-N-acetyl transferase (GAT) disclosed in U.S. Patent Application publication 2003/0083480 A1 also for imparting glyphosate tolerance; dicamba monooxygenase disclosed in U.S. Patent

Application publication 2003/0135879 A1 for imparting dicamba tolerance; a polynucleotide molecule encoding bromoxynil nitrilase (*Bxn*) disclosed in U.S. Patent 4,810,648 for imparting bromoxynil tolerance; a polynucleotide molecule encoding phytoene desaturase (*crtI*) described in Misawa et al, (1993) *Plant J.* 4:833-840 and in Misawa et al, (1994) *Plant J.* 6:481-489 for norflurazon tolerance; a polynucleotide molecule encoding acetohydroxyacid synthase (AHAS, *aka* ALS) described in Sathasiivan *et al.* (1990) *Nucl. Acids Res.* 18:2188-2193 for imparting tolerance to sulfonylurea herbicides; polynucleotide molecules known as *bar* genes disclosed in DeBlock, *et al.* (1987) *EMBO J.* 6:2513-2519 for imparting glufosinate and bialaphos tolerance; polynucleotide molecules disclosed in U.S. Patent Application Publication 2003/010609 A1 for imparting N-amino methyl phosphonic acid tolerance; polynucleotide molecules disclosed in U.S. Patent 6,107,549 for imparting pyridine herbicide resistance; molecules and methods for imparting tolerance to multiple herbicides such as glyphosate, atrazine, ALS inhibitors, isoxoflutole and glufosinate herbicides are disclosed in U.S. Patent 6,376,754 and U.S. Patent Application Publication 2002/0112260, all of said U.S. Patents and Patent Application Publications are incorporated herein by reference. Molecules and methods for imparting insect/nematode/virus resistance are disclosed in U.S. Patents 5,250,515; 5,880,275; 6,506,599; 5,986,175 and U.S. Patent Application Publication 2003/0150017 A1, all of which are incorporated herein by reference.

A “consensus amino acid sequence” means an artificial, amino acid sequence indicating conserved amino acids in the sequence of homologous proteins as determined by statistical analysis of an optimal alignment, *e.g.* CLUSTALW, of amino acid sequence of homolog proteins. The consensus sequences listed in the sequence listing were created by identifying the most frequent amino acid at each position in a set of aligned protein sequences. When there was 100% identity in an alignment the amino acid is indicated by a capital letter. When the occurrence of an amino acid is at least about 70% in an alignment, the amino acid is indicated by a lower case letter. When there is no amino acid occurrence of at least about 70%, *e.g.* due to diversity or gaps, the amino acid is indicated by an “x”. When used to defined embodiments of the invention, a consensus amino acid sequence will be aligned with a query protein amino acid sequence in an optimal alignment, *e.g.* CLUSTALW. An embodiment of the invention will have identity to the conserved amino acids indicated in the consensus amino acid sequence.

“*Arabidopsis*” means plants of *Arabidopsis thaliana*.

“Pfam” database is a large collection of multiple sequence alignments and hidden

Markov models covering many common protein families, *e.g.* Pfam version 19.0 (December 2005) contains alignments and models for 8183 protein families and is based on the Swissprot 47.0 and SP-TrEMBL 30.0 protein sequence databases. See S.R. Eddy, "Profile Hidden Markov Models", *Bioinformatics* 14:755-763, 1998. The Pfam database is currently
5 maintained and updated by the Pfam Consortium. The alignments represent some evolutionary conserved structure that has implications for the protein's function. Profile hidden Markov models (profile HMMs) built from the protein family alignments can be used for automatically recognizing that a new protein belongs to an existing protein family even if the homology by alignment appears to be low.

10 A "Pfam domain module" is a representation of Pfam domains in a protein, in order from N terminus to C terminus. In a Pfam domain module individual Pfam domains are separated by double colons "::". The order and copy number of the Pfam domains from N to C terminus are attributes of a Pfam domain module. Although the copy number of repetitive domains is important, varying copy number often enables a similar function. Thus,
15 a Pfam domain module with multiple copies of a domain should define an equivalent Pfam domain module with variance in the number of multiple copies. A Pfam domain module is not specific for distance between adjacent domains, but contemplates natural distances and variations in distance that provide equivalent function. The Pfam database contains both narrowly- and broadly-defined domains, leading to identification of overlapping domains on
20 some proteins. A Pfam domain module is characterized by non-overlapping domains. Where there is overlap, the domain having a function that is more closely associated with the function of the protein (based on the E value of the Pfam match) is selected.

Once one DNA is identified as encoding a protein which imparts an enhanced trait when expressed in transgenic plants, other DNA encoding proteins with the same Pfam
25 domain module are identified by querying the amino acid sequence of protein encoded by candidate DNA against the Hidden Markov Models which characterizes the Pfam domains using HMMER software, a current version of which is provided in the appended computer listing. Candidate proteins meeting the same Pfam domain module are in the protein family and have cognate DNA that is used in constructing recombinant DNA for the use in the plant
30 cells of this invention. Hidden Markov Model databases for use with HMMER software in identifying DNA expressing protein with a common Pfam domain module for recombinant DNA in the plant cells of this invention are also included in the appended computer listing.

Version 19.0 of the HMMER software and Pfam databases were used to identify known domains in the proteins corresponding to amino acid sequence of SEQ ID NO: 804

through SEQ ID NO:1606. All DNA encoding proteins that have scores higher than the gathering cutoff disclosed in Table 19 by Pfam analysis disclosed herein can be used in recombinant DNA of the plant cells of this invention, *e.g.* for selecting transgenic plants having enhanced agronomic traits. The relevant Pfam modules for use in this invention, as

5 more specifically disclosed below, are Syntaxin::SNARE, Pro_isomerase, Pkinase, ATP-synt_G, Carboxyl_trans, CDC50, GATase::GMP_synt_C, F-box::WD40::WD40::WD40, dsrm::dsrm, Pyr_redox_2::Pyr_redox_dim, WAK::Pkinase, Pkinase_Tyr, PTPA, Biotin_lipoyl::E3_binding::2-oxoacid_dh, AAA, LRRNT_2::LRR_1::LRR_1::LRR_1, PRA1, TIM, YTH, ThiF, Hep_59, Pkinase, PALP::Thr_dehydrat_C::Thr_dehydrat_C, zf-

10 Tim10_DDP, CAF1, Pkinase, Pyr_redox_2::Pyr_redox_dim, Response_reg, YIF1, NPH3, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::Pkinase_Tyr, NOI, Fer2, AA_permease, Caleosin, IF4E, Pkinase, Ribul_P_3_epim, F-box::WD40::WD40::WD40::WD40::WD40::WD40, MIP, p450, Ribosomal_S5::Ribosomal_S5_C, Hpt, TBC, Acyltransferase, Epimerase, Pkinase, Dus,

15 SelR, PH, Tyr-DNA_phospho, RRM_1, Pkinase, ECH, SRPRB, DUF599, Pkinase::efhand::efhand::efhand::efhand, ORC2, PME1, p450, GLTP, Suc_Fer-like, CBS::CBS, ADH_N::ADH_zinc_N, LRRNT_2::LRR_1::LRR_1::LRR_1::Pkinase, F-box, adh_short, polyprenyl_synt, GHMP_kinases_N::GHMP_kinases_C, PsbW, Na_H_Exchanger::TrkA_C, Ribosomal_L12, CDC50, DCP1, Pyr_redox_2::Pyr_redox_dim,

20 Steroid_dh, tRNA-synt_1c::tRNA-synt_1c_C, Cupin_1::Cupin_1, peroxidase, Sugar_tr, Pro_isomerase, Iso_dh, PTS_2-RNA, Aminotran_5, ENOD93, KH_1::KH_1::KH_1::KH_1::KH_1, Mito_carr::Mito_carr::Mito_carr, DUF569::DUF569, C1_2::C1_2::C1_3::C1_2::C1_3::C1_2, DnaJ::zf-C2H2, Aldedh, AAA::Rep_fac_C, DUF641, Aminotran_1_2, TB2_DP1_HVA22, zf-C3HC4, RRM_1::RRM_2, Pkinase,

25 PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR, Pyr_redox_2, IQ::IQ, SMP::SMP::SMP, AWPM-19, CLP_protease, Pkinase::NAF, PRMT5, PPR::PPR::PPR::PPR::PPR::PPR, Aldedh, ATP-synt_C::ATP-synt_C, adh_short, RCC1::RCC1::RCC1::RCC1::FYVE::DZC, p450, PfkB, PB1, Hexapep::Hexapep::Hexapep::Hexapep, Aminotran_1_2, DUF862, Aldedh,

30 ADH_N::ADH_zinc_N, CDC48_N::AAA::AAA, AA_permease, UQ_con, Mem_trans, GFA, OPT, DUF887, Di19, U-box::Arm::Arm, DAO, G-alpha, Aminotran_3, 2OG-FelI_Oxy, Lectin_legB::Pkinase, NAC::UBA, efhand_like::PI-PLC-X::PI-PLC-Y::C2, TLC, ATP-grasp_2::Ligase_CoA, Pkinase, PRC::PRC, MFS_1::Sugar_tr, Copine::zf-C3HC4, RRM_1::RRM_1::RRM_1, Pribosyltran, AA_kinase, PhzC-PhzF, Gp_dh_N::Gp_dh_C,

Sugar_tr, Pre-SET::SET, Alpha-amylase::Alpha-amyl_C2, Cyclin_N,
 FAE1_CUT1_RppA::ACP_syn_III_C, RPE65, efhand_like::PI-PLC-X::PI-PLC-Y::C2,
 Mito_carr::Mito_carr::Mito_carr, Invertase_neut, DSPc, LANC_like, Aminotran_5,
 Glutaminase, PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR, PGI, SSB, p450,
 5 La::RRM_1::RRM_3, Methyltransf_11, SBF,
 NTP_transferase::Hexapep::Hexapep::Hexapep, Mov34, Hydrolase, AAA, RRM_1::RRM_1,
 Pkinase::NAF, F-box::FBA_1, p450, F-box::LRR_2, zf-DHHC, TBP::TBP, DUF607, FMO-
 like, adh_short, ATP-synt_ab_N::ATP-synt_ab, RNase_PH, PP2C, UQ_con, Aminotran_1_2,
 p450, FMO-like, Gp_dh_N::Gp_dh_C, Mito_carr::Mito_carr::Mito_carr, zf-CCCH::zf-
 10 CCCH::zf-CCCH::zf-CCCH::zf-CCCH, GST_C, GH3, PAP_fibrillin, ThiF,
 Ribul_P_3_epim, DUF260, p450, MACPF, Steroid_dh, Response_reg, Carb_anhydrase,
 Aldedh, Pkinase, Methyltransf_11, PK::PK_C, Pkinase, DEK_C::SWIB::SWIB, TATA_RF,
 Tryp_alpha_amyl, Y_phosphatase2, TATA_RF, NifU_N, ENT, Pkinase, F-
 box::Kelch_2::Kelch_2, WD40::WD40, MtN3_slv::MtN3_slv, TB2_DP1_HVA22, Pkinase,
 15 Aldedh, AAA, Pyridoxal_deC, zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC, Peptidase_M24,
 LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::Pkinase, Arf, DUF1279, UPF0185,
 Rhodanese, adh_short, Tryp_alpha_amyl, Pkinase, UPF0139, CBS, Glyco_transf_29,
 iPGM_N::Metalloenzyme, Proteasome, Pkinase, UBX,
 ELFV_dehydrog_N::ELFV_dehydrog, p450, ADH_N::ADH_zinc_N, zf-C3HC4, NDK,
 20 NAD_Gly3P_dh_N::NAD_Gly3P_dh_C, GlutR_N::Shikimate_DH::GlutR_dimer, p450,
 SNF5, p450, Pribosyltran, AIG1, Response_reg::CCT, mTERF, DUF220, Pkinase::NAF, zf-
 CCCH::zf-CCCH, Pkinase, Pkinase, MATH::MATH, Asparaginase, Pkinase,
 Gp_dh_N::Gp_dh_C, Pkinase, KH_1, Fcf1, PK::PK_C::PEP-utilizers,
 CRAL_TRIO_N::CRAL_TRIO, Raffinose_syn, DUF584, BT1, Aminotran_1_2,
 25 C1_2::C1_3::C1_3::C1_2::C1_3::C1_3::C1_2, FMO-like, PLAC8, F-
 box::Kelch_2::Kelch_1::Kelch_2, XS::XH, STT3, ABC_tran::ABC_tran, Aldo_ket_red,
 NTF2::RRM_1, DUF594, Biotin_lipoyl::2-oxoacid_dh, RRM_1, FHA, Mov34, MT-A70,
 Brix, X8, Auxin_inducible, Peptidase_M18, Sec61_beta, F-box::Kelch_1::Kelch_1, Brix,
 TIM, CTP_transf_1, Tetraspannin, PALP, DUF822, Pkinase, B3::B3, MFS_1, SFT2, F-
 30 box::Kelch_1::Kelch_1, DUF588, Thioredoxin, Alba, Ion_trans_2::Ion_trans_2,
 Peptidase_C48, YDG_SRA::Pre-SET::SET, Aldedh, Abhydrolase_3, p450, U-
 box::Arm::Arm::Arm::Arm, AA_permease, LAG1, peroxidase, PAD_porph, Pkinase, F-box,
 AIG2, p450, AARP2CN::DUF663, UPF0153, NAD_binding_2::6PGD, Pkinase,
 B_lectin::S_locus_glycop::PAN_1::Pkinase, EMP24_GP25L, DUF6::DUF6, DUF163, Sina,

adh_short, Rho_GDI, zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC, VHS::GAT, Pyrophosphatase, DMRL_synthase, Cyclin_N::Cyclin_C, NDK, TPP_enzyme_N::TPP_enzyme_M::TPP_enzyme_C, SH3_1, UDPG_MGDP_dh_N::UDPG_MGDP_dh::UDPG_MGDP_dh_C, UIM::efhand, DUF579,

5 Pkinase, DHBP_synthase::GTP_cyclohydro2, Peptidase_C14, Glycolytic, Pkinase, Lipase_3, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1, Snf7, BT1, PPK_N::PEP-utilizers::PEP-utilizers_C, V-ATPase_G, RNA_pol_I_A49, Zip, WD40::WD40::WD40, Branch, RCC1::RCC1::RCC1::RCC1, TFIID-18kDa, Pkinase, Cullin, TFIID-31kDa, Asp, Pkinase, Ubie_methyltran, PPR::PPR, PPR::PPR::PPR::PPR::PPR::PPR, Fructosamin_kin,

10 PPR::PPR::PPR::PPR::PPR::PPR, Pkinase_Tyr, p450, AA_permease, Glycos_transf_1, ECH::3HCDH_N::3HCDH, MIP, ADH_N::ADH_zinc_N, Aa_trans, DUF1517, Redoxin, RRS1, Pribosyltran, Sulfate_transp::STAS, DUF167, MMR_HSR1::DUF933, Aminotran_1_2, AA_permease, DUF620, Aldedh, Aminotran_1_2, LRR_1, CDP-OH_P_transf, DHBP_synthase::GTP_cyclohydro2, NHL::NHL, WD40::U3_snoRNA_C,

15 Na_sulph_symp, adh_short, Thioredoxin, Ribosomal_L7Ae, OTCace_N::OTCace, Mito_carr::Mito_carr::Mito_carr, Pescadillo_N::BRCT, PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR, mTERF, Chalcone, Per1, PPR::PPR, Na_H_Exchange::TrkA_N, DEAD::Helicase_C, Cyclin_N::Cyclin_C, PP-binding, SRF-TF, SATase_N::Hexapep::Hexapep::Hexapep, PK::PK_C, adh_short,

20 DUF6::DUF6, Pyr_redox_2, G-patch, WD40::WD40::WD40::WD40, G-alpha, VQ, WD40::WD40::WD40::WD40::WD40, Aminotran_3, Pribosyltran, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1, eIF3_subunit, H_PPase, Nodulin-like, Whi5, Tryp_alpha_amyl, NIR_SIR_ferr::NIR_SIR::NIR_SIR_ferr, RRM_1::RRM_1, Band_7, Asp, MFS_1, F420_oxidored, Pkinase, PRK, Na_Ca_ex::Na_Ca_ex, Bombesin,

25 Pkinase, Epimerase, Pkinase, Inositol_P, Arm::Arm::Arm::Arm, PB1, Hin1, p450, RNA_pol_Rpb5_N::RNA_pol_Rpb5_C, Adap_comp_sub, Peptidase_M16::Peptidase_M16_C, Pkinase, PGK, Biotin_lipoyl::E3_binding::2-oxoacid_dh, Alg6_Alg8, F-box, MMR_HSR1, G6PD_N::G6PD_C, DAD, malic::Malic_M::PTA_PTB, ADH_N::ADH_zinc_N,

30 LRR_1::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1, PfkB, PA, Pyr_redox_2::Pyr_redox_dim, SIP1, PP2C, BT1, AhpC-TSA, GASA, Aminotran_3, Sad1_UNC, Pkinase, zf-MYND::PDCD2_C, Pkinase, PGK, ArfGap, GATase::GMP_synt_C, p450, DUF1637, 2OG-FeII_Oxy, Aldedh, DUF231, Glyco_transf_8, Sec1, DUF1475, Cellulose_synt, RRM_1::RRM_1::RRM_1, Usp, AA_permease,

Acetyltransf_1::Bromodomain, Glyco_tran_28_C, RNA_pol_N, NTP_transferase,
malic::Malic_M, Histone, Epimerase, UPF0061, ClpS, LEA_5, Auxin_inducible,
PAP_fibrillin, Aldo_ket_red, Gln-synt_C, DREPP, p450, SFT2, eIF2A, Cenp-O,
CPSase_sm_chain::GATase, WD40::WD40::WD40, Pkinase, Porin_3, Pkinase,
5 Aminotran_1_2, GST_N::GST_C, F-box::Kelch_1::Kelch_1, IU_nuc_hydro, SYF2, PDT,
Subtilisin_N::PA, Sugar_tr, WD40::WD40, p450, DUF829, Pyr_redox_2::Pyr_redox_dim,
Aminotran_1_2, Cyclin_N::Cyclin_C, YgbB, Sugar_tr, SRF-TF, DUF231, DUF584, BT1, zf-
C3HC4, Metallophos, FA_hydroxylase, p450, ACBP::Ank::Ank, LSM, SAM_1,
LRR_1::LRR_1::Pkinase, Cullin, F-box::FBA_1, peroxidase, Rib_5-P_isom_A, zf-A20::zf-
10 AN1, Dimerisation::Methyltransf_2, PGAM, PTR2, Copine, PALP, Exo_endo_phos,
G6PD_N::G6PD_C, PLAC8, Aldo_ket_red, Ank::Ank, FBPase, zf-C3HC4, ACT::ACT,
Suc_Fer-like, Pkinase_Tyr, G-alpha, RAMP4, Glutaredoxin, RRM_1::RRM_1, MT-A70,
Response_reg::CCT, Arf, AP2, DUF177, 2OG-FeII_Oxy,
LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1, LRR_1, Aldedh, DUF1350,
15 FAE1_CUT1_RppA::ACP_syn_III_C, Sybindin, zf-C3HC4, Nfu_N::NifU, Rho_GDI,
PK::PK_C, NOSIC::Nop, PK::PK_C, PRK::Pribosyltran, PfkB, p450, DUF231,
TFIID_30kDa, Aldedh, zf-AN1, GHMP_kinases_N::GHMP_kinases_C, DUF423, Pkinase,
Cu-oxidase_3::Cu-oxidase::Cu-oxidase_2, PCI, RNase_PH::RNase_PH_C, DUF59,
NTP_transferase, CH::EB1, Pkinase::Pkinase_C, DUF866, SMP::SMP::SMP, Aminotran_3,
20 Transket_pyr::Transketolase_C, Copine, His_biosynth, Tbf5, DUF543, p450, C2, DUF616,
Gp_dh_N::Gp_dh_C, Smg4_UPF3, DUF231, DUF89, WD40::WD40, Aldedh,
ACT::ACT::ACT::ACT, Pkinase, Pyridoxal_deC, Skp1_POZ::Skp1, NAP, FolB, p450,
Pentapeptide::Pentapeptide, NTP_transferase::MannoseP_isomer, SQS_PSY, Cyclase,
GASA, Rick_17kDa_Anti, FMO-like, B_lectin::S_locus_glycop::PAN_2, Peptidase_C12,
25 Ribosomal_S6e, Glyoxalase, Response_reg::CCT, PHD::SET, Redoxin, G_glu_transpept,
Synaptobrevin, p450, RALF, PALP, Branch, DUF579, Aminotran_3, Nramp,
Enolase_N::Enolase_C, Str_synth, FAD_binding_3, MOSC_N::MOSC, Spc97_Spc98,
Glycolytic, F-box::WD40::WD40::WD40::WD40, AA_kinase::ACT::ACT, RALF, and
ATP_bind_1.

30 **Plant Cell Transformation Methods**

Numerous methods for transforming chromosomes in a plant cell nucleus with recombinant DNA are known in the art and are used in methods of preparing a transgenic plant cell nucleus cell, and plant. Two effective methods for such transformation are

Agrobacterium-mediated transformation and microprojectile bombardment. Microprojectile bombardment methods are illustrated in US Patents 5,015,580 (soybean); 5,550,318 (corn); 5,538,880 (corn); 5,914,451 (soybean); 6,160,208 (corn); 6,399,861 (corn); 6,153,812 (wheat) and 6,365,807 (rice) and *Agrobacterium*-mediated transformation is described in US Patents 5,159,135 (cotton); 5,824,877 (soybean); 5,463,174 (canola, also known as rapeseed); 5,591,616 (corn); 6,384,301 (soybean), 7,026,528 (wheat) and 6,329,571 (rice), all of which are incorporated herein by reference for enabling the production of transgenic plants. Transformation of plant material is practiced in tissue culture on a nutrient media, *e.g.* a mixture of nutrients that will allow cells to grow *in vitro*. Recipient cell targets include, but are not limited to, meristem cells, hypocotyls, calli, immature embryos and gametic cells such as microspores, pollen, sperm and egg cells. Callus can be initiated from tissue sources including, but not limited to, immature embryos, hypocotyls, seedling apical meristems, microspores and the like. Cells containing a transgenic nucleus are grown into transgenic plants.

In addition to direct transformation of a plant material with a recombinant DNA, a transgenic plant cell nucleus can be prepared by crossing a first plant having cells with a transgenic nucleus with recombinant DNA with a second plant lacking the transgenic nucleus. For example, recombinant DNA can be introduced into a nucleus from a first plant line that is amenable to transformation to transgenic nucleus in cells that are grown into a transgenic plant which can be crossed with a second plant line to introgress the recombinant DNA into the second plant line. A transgenic plant with recombinant DNA providing an enhanced trait, *e.g.* enhanced yield, can be crossed with transgenic plant line having other recombinant DNA that confers another trait, for example herbicide resistance or pest resistance, to produce progeny plants having recombinant DNA that confers both traits. Typically, in such breeding for combining traits the transgenic plant donating the additional trait is a male line and the transgenic plant carrying the base traits is the female line. The progeny of this cross will segregate such that some of the plants will carry the DNA for both parental traits and some will carry DNA for one parental trait; such plants can be identified by markers associated with parental recombinant DNA, *e.g.* marker identification by analysis for recombinant DNA or, in the case where a selectable marker is linked to the recombinant, by application of the selecting agent such as a herbicide for use with a herbicide tolerance marker, or by selection for the enhanced trait. Progeny plants carrying DNA for both parental traits can be crossed back into the female parent line multiple times, for example usually 6 to 8 generations, to

produce a progeny plant with substantially the same genotype as one original transgenic parental line but for the recombinant DNA of the other transgenic parental line

In practice, DNA is introduced into only a small percentage of target cell nuclei. Marker genes are used to provide an efficient system for identification of those cells with nuclei that are stably transformed by receiving and integrating a recombinant DNA molecule into their genomes. Some marker genes provide selective markers that confer resistance to a selective agent, such as an antibiotic or herbicide. Potentially transformed cells with a nucleus of the invention are exposed to the selective agent. In the population of surviving cells will be those cells where, generally, the resistance-conferring gene has been integrated and expressed at sufficient levels to permit cell survival. Cells can be tested further to confirm stable integration of the exogenous DNA in the nucleus. Exemplary selective marker genes include those conferring resistance to antibiotics such as kanamycin (*nptII*), hygromycin B (*aph IV*), spectinomycin (*aadA*) and gentamycin (*aac3* and *aacC4*) or resistance to herbicides such as glufosinate (*bar* or *pat*), dicamba (DMO) and glyphosate (EPSPS). Examples of such selectable markers are illustrated in U.S. Patents 5,550,318; 5,633,435; 5,780,708 and 6,118,047, all of which are incorporated herein by reference. Screenable markers which provide an ability to visually identify transformants can also be employed, *e.g.*, a gene expressing a colored or fluorescent protein such as a luciferase or green fluorescent protein (GFP) or a gene expressing a *beta*-glucuronidase or *uidA* gene (GUS) for which various chromogenic substrates are known. It is also contemplated that combinations of screenable and selectable markers can be used for identification of transformed cells. See PCT publication WO 99/61129 (herein incorporated by reference) which discloses use of a gene fusion between a selectable marker gene and a screenable marker gene, *e.g.*, an NPTII gene and a GFP gene.

Plant cells that survive exposure to the selective agent, or plant cells that have been scored positive in a screening assay, can be cultured in regeneration media and allowed to mature into plants. Developing plantlets regenerated from transformed plant cells can be transferred to plant growth mix, and hardened off, for example, in an environmentally controlled chamber at about 85% relative humidity, 600 ppm CO₂, and 25-250 microeinsteins m⁻² s⁻¹ of light, prior to transfer to a greenhouse or growth chamber for maturation. Plants are regenerated from about 6 weeks to 10 months after a transformant is identified, depending on the initial tissue, and plant species. Plants can be pollinated using conventional plant breeding methods known to those of skill in the art and seed produced, for example self-pollination is commonly used with transgenic corn. The regenerated transformed plant or its

progeny seed or plants can be tested for expression of the recombinant DNA and selected for the presence of enhanced agronomic trait.

Transgenic Plants and Seeds

Transgenic plants derived from transgenic plant cells having a transgenic nucleus of this invention are grown to generate transgenic plants having an enhanced trait as compared to a control plant and produce transgenic seed and haploid pollen of this invention. Such plants with enhanced traits are identified by selection of transformed plants or progeny seed for the enhanced trait. For efficiency a selection method is designed to evaluate multiple transgenic plants (events) comprising the recombinant DNA, for example multiple plants from 2 to 20 or more transgenic events. Transgenic plants grown from transgenic seed provided herein demonstrate improved agronomic traits that contribute to increased yield or other trait that provides increased plant value, including, for example, improved seed quality. Of particular interest are plants having enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

To identify nuclei with recombinant DNA that confer enhanced traits to plants, *Arabidopsis thaliana* was transformed with a candidate recombinant DNA construct and screened for an enhanced trait.

Arabidopsis thaliana is used a model for genetics and metabolism in plants. A two-step screening process was employed which included two passes of trait characterization to ensure that the trait modification was dependent on expression of the recombinant DNA, but not due to the chromosomal location of the integration of the transgene. Twelve independent transgenic lines for each recombinant DNA construct were established and assayed for the transgene expression levels. Five transgenic lines with high transgene expression levels were used in the first pass screen to evaluate the transgene's function in T2 transgenic plants. Subsequently, three transgenic events, which had been shown to have one or more enhanced traits, were further evaluated in the second pass screen to confirm the transgene's ability to impart an enhanced trait. The following Table 3 summarizes the enhanced traits that have been confirmed as provided by a recombinant DNA construct.

Table 2 provides a list of protein encoding DNA ("genes") as recombinant DNA for production of transgenic plants with enhanced agronomic trait, the elements of Table 2 are described by reference to:

PEP SEQ ID NO" identifies an amino acid sequence from SEQ ID NO: 804 to 1606.

“NUC SEQ ID NO” identifies a DNA sequence from SEQ ID NO:1 to 803

“construct_id” refers to an arbitrary number used to identify a particular recombinant DNA construct comprising the particular DNA.

“Gene ID” refers to an arbitrary name used to identify the particular DNA.

- 5 “orientation” refers to the orientation of the particular DNA in a recombinant DNA construct relative to the promoter.

Table 2.

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
1	804	CGPG10	70244	SENSE	399	1202	CGPG5796	73036	SENSE
2	805	CGPG1008	71108	SENSE	400	1203	CGPG5800	72958	SENSE
3	806	CGPG1023	12816	SENSE	401	1204	CGPG5812	77006	SENSE
4	807	CGPG1038	12650	ANTI- SENSE	402	1205	CGPG5815	74732	SENSE
5	808	CGPG1067	12031	SENSE	403	1206	CGPG5838	74333	SENSE
6	809	CGPG1095	12217	SENSE	404	1207	CGPG5844	74735	SENSE
7	810	CGPG1101	12043	SENSE	405	1208	CGPG5846	74736	SENSE
8	811	CGPG1109	12147	ANTI- SENSE	406	1209	CGPG5851	74737	SENSE
9	812	CGPG1113	13638	ANTI- SENSE	407	1210	CGPG5857	76657	SENSE
10	813	CGPG1155	11860	ANTI- SENSE	408	1211	CGPG5862	76113	SENSE
11	814	CGPG1163	72361	SENSE	409	1212	CGPG5863	74747	SENSE
12	815	CGPG1171	12230	SENSE	410	1213	CGPG5867	77309	SENSE
13	816	CGPG1177	71626	SENSE	411	1214	CGPG5872	74322	SENSE
14	817	CGPG1205	13454	ANTI- SENSE	412	1215	CGPG5913	76212	SENSE
15	818	CGPG1207	12173	ANTI- SENSE	413	1216	CGPG5922	76514	SENSE
16	819	CGPG1215	17806	ANTI- SENSE	414	1217	CGPG5961	75242	SENSE
17	820	CGPG1223	12011	ANTI- SENSE	415	1218	CGPG5964	77324	SENSE
18	821	CGPG1250	72915	SENSE	416	1219	CGPG5968	74350	SENSE
19	822	CGPG1279	13231	SENSE	417	1220	CGPG5984	77606	SENSE
20	823	CGPG1304	13913	SENSE	418	1221	CGPG5991	76530	SENSE
21	824	CGPG1319	12749	ANTI- SENSE	419	1222	CGPG6006	76716	SENSE
22	825	CGPG1327	78901	SENSE	420	1223	CGPG6015	77328	SENSE
23	826	CGPG1329	75969	SENSE	421	1224	CGPG603	12204	SENSE
24	827	CGPG133	15606	SENSE	422	1225	CGPG6046	77011	SENSE
25	828	CGPG1332	75946	SENSE	423	1226	CGPG6063	77607	SENSE
26	829	CGPG1343	13050	ANTI- SENSE	424	1227	CGPG6083	74612	SENSE
27	830	CGPG1348	75958	SENSE	425	1228	CGPG6092	76126	SENSE
28	831	CGPG1349	17333	SENSE	426	1229	CGPG6104	74376	SENSE
29	832	CGPG1373	76002	SENSE	427	1230	CGPG6106	74377	SENSE
30	833	CGPG1377	78735	SENSE	428	1231	CGPG6111	74381	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
31	834	CGPG1412	13914	SENSE	429	1232	CGPG6113	78364	SENSE
32	835	CGPG1421	14405	SENSE	430	1233	CGPG6125	74623	SENSE
33	836	CGPG1426	75983	SENSE	431	1234	CGPG6132	74668	SENSE
34	837	CGPG1433	70721	ANTI- SENSE	432	1235	CGPG6137	74629	SENSE
35	838	CGPG1453	13811	SENSE	433	1236	CGPG6147	74635	SENSE
35	838	CGPG1453	13828	ANTI- SENSE	434	1237	CGPG6152	74638	SENSE
36	839	CGPG1454	13614	SENSE	435	1238	CGPG6154	76721	SENSE
37	840	CGPG1463	76050	SENSE	436	1239	CGPG6170	78365	SENSE
38	841	CGPG1464	13038	ANTI- SENSE	437	1240	CGPG6171	74655	SENSE
39	842	CGPG1471	13040	ANTI- SENSE	438	1241	CGPG6177	74660	SENSE
40	843	CGPG1481	70716	ANTI- SENSE	439	1242	CGPG6181	74661	SENSE
40	843	CGPG1481	76039	SENSE	440	1243	CGPG6188	75259	SENSE
41	844	CGPG1499	71611	SENSE	441	1244	CGPG6202	75282	SENSE
42	845	CGPG150	70253	SENSE	442	1245	CGPG6217	76724	SENSE
43	846	CGPG1536	14415	SENSE	443	1246	CGPG623	12350	ANTI- SENSE
44	847	CGPG1539	13718	ANTI- SENSE	444	1247	CGPG6239	76522	SENSE
45	848	CGPG155	10706	ANTI- SENSE	445	1248	CGPG6244	77903	SENSE
45	848	CGPG155	10709	ANTI- SENSE	446	1249	CGPG6254	76653	SENSE
46	849	CGPG1583	14340	ANTI- SENSE	447	1250	CGPG6266	76524	SENSE
47	850	CGPG1588	13471	ANTI- SENSE	448	1251	CGPG627	11356	SENSE
48	851	CGPG16	18304	SENSE	449	1252	CGPG6271	75275	SENSE
49	852	CGPG1609	13623	SENSE	450	1253	CGPG6278	76225	SENSE
50	853	CGPG1629	18302	SENSE	451	1254	CGPG6288	75281	SENSE
51	854	CGPG1637	15053	ANTI- SENSE	452	1255	CGPG6309	74384	SENSE
52	855	CGPG1653	15624	SENSE	453	1256	CGPG633	71238	SENSE
53	856	CGPG1658	70412	SENSE	454	1257	CGPG635	12354	ANTI- SENSE
54	857	CGPG1663	17902	SENSE	455	1258	CGPG6350	76236	SENSE
55	858	CGPG1682	14272	ANTI- SENSE	456	1259	CGPG6358	74682	SENSE
56	859	CGPG1701	15122	SENSE	457	1260	CGPG6365	73425	SENSE
56	859	CGPG1701	70723	ANTI- SENSE	458	1261	CGPG6374	73438	SENSE
57	860	CGPG1723	14248	SENSE	459	1262	CGPG6377	73474	SENSE
58	861	CGPG1724	13965	ANTI- SENSE	460	1263	CGPG638	70850	SENSE
59	862	CGPG1726	75912	SENSE	461	1264	CGPG6397	73429	SENSE
60	863	CGPG1736	14344	ANTI- SENSE	462	1265	CGPG6398	73441	SENSE
61	864	CGPG1741	14807	ANTI- SENSE	463	1266	CGPG6408	73466	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
62	865	CGPG1783	17307	SENSE	464	1267	CGPG6415	73455	SENSE
63	866	CGPG1790	16431	ANTI- SENSE	465	1268	CGPG6421	73432	SENSE
64	867	CGPG1797	73075	SENSE	466	1269	CGPG6442	73435	SENSE
65	868	CGPG1845	16323	SENSE	467	1270	CGPG6443	77726	SENSE
66	869	CGPG1855	14833	ANTI- SENSE	468	1271	CGPG6446	73483	SENSE
67	870	CGPG1870	17030	ANTI- SENSE	469	1272	CGPG6453	73472	SENSE
68	871	CGPG1879	76414	SENSE	470	1273	CGPG6466	73526	SENSE
69	872	CGPG1886	16021	SENSE	471	1274	CGPG6467	73538	SENSE
70	873	CGPG1903	78701	SENSE	472	1275	CGPG6470	73574	SENSE
71	874	CGPG1905	78702	SENSE	473	1276	CGPG6475	73539	SENSE
72	875	CGPG1914	15220	ANTI- SENSE	474	1277	CGPG6477	73563	SENSE
73	876	CGPG193	15623	SENSE	475	1278	CGPG6478	73575	SENSE
74	877	CGPG1939	19402	SENSE	476	1279	CGPG6493	73565	SENSE
75	878	CGPG1949	70559	SENSE	477	1280	CGPG6506	73531	SENSE
76	879	CGPG1959	15111	ANTI- SENSE	478	1281	CGPG6518	73592	SENSE
77	880	CGPG197	70832	SENSE	479	1282	CGPG6546	74149	SENSE
78	881	CGPG1972	15129	SENSE	480	1283	CGPG6556	74174	SENSE
79	882	CGPG1981	76086	SENSE	481	1284	CGPG6562	74151	SENSE
80	883	CGPG1999	15203	ANTI- SENSE	482	1285	CGPG6566	74104	SENSE
81	884	CGPG2	70216	SENSE	483	1286	CGPG6571	74164	SENSE
82	885	CGPG2006	14910	ANTI- SENSE	484	1287	CGPG6576	74129	SENSE
83	886	CGPG2010	70318	SENSE	485	1288	CGPG659	72340	SENSE
84	887	CGPG2011	15207	ANTI- SENSE	486	1289	CGPG6594	74155	SENSE
85	888	CGPG2014	14915	ANTI- SENSE	487	1290	CGPG6603	74168	SENSE
86	889	CGPG2023	15132	SENSE	488	1291	CGPG6608	74133	SENSE
87	890	CGPG2026	16229	SENSE	489	1292	CGPG6634	74160	SENSE
88	891	CGPG2041	14932	ANTI- SENSE	490	1293	CGPG6662	74485	SENSE
89	892	CGPG2064	76017	SENSE	491	1294	CGPG6667	74450	SENSE
90	893	CGPG2070	15964	ANTI- SENSE	492	1295	CGPG6678	74487	SENSE
91	894	CGPG2077	17227	SENSE	493	1296	CGPG6695	74406	SENSE
92	895	CGPG2095	72952	SENSE	494	1297	CGPG6699	74454	SENSE
93	896	CGPG2105	73706	SENSE	495	1298	CGPG670	11361	SENSE
94	897	CGPG2108	18203	SENSE	496	1299	CGPG6705	74431	SENSE
95	898	CGPG2111	16309	SENSE	497	1300	CGPG6735	74411	SENSE
96	899	CGPG2124	15986	ANTI- SENSE	498	1301	CGPG6744	74424	SENSE
97	900	CGPG2125	15987	ANTI- SENSE	499	1302	CGPG6753	74525	SENSE
98	901	CGPG2126	16204	SENSE	500	1303	CGPG6757	74573	SENSE
99	902	CGPG213	11133	SENSE	501	1304	CGPG6803	75827	SENSE
100	903	CGPG2134	73977	SENSE	502	1305	CGPG6806	77329	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
101	904	CGPG2139	15995	ANTI- SENSE	503	1306	CGPG6811	77330	SENSE
101	904	CGPG2139	18201	SENSE	504	1307	CGPG6812	75830	SENSE
102	905	CGPG2140	16887	SENSE	505	1308	CGPG6815	75832	SENSE
103	906	CGPG2153	16213	SENSE	506	1309	CGPG6827	75836	SENSE
104	907	CGPG2163	15507	ANTI- SENSE	507	1310	CGPG6830	77034	SENSE
105	908	CGPG2165	15508	ANTI- SENSE	508	1311	CGPG6839	76255	SENSE
106	909	CGPG2193	15959	ANTI- SENSE	509	1312	CGPG6859	76542	SENSE
107	910	CGPG2218	72427	ANTI- SENSE	510	1313	CGPG6864	77511	SENSE
108	911	CGPG2224	19041	ANTI- SENSE	511	1314	CGPG6877	76261	SENSE
109	912	CGPG2225	17802	SENSE	512	1315	CGPG6878	78456	SENSE
110	913	CGPG2229	19042	ANTI- SENSE	513	1316	CGPG6880	77041	SENSE
111	914	CGPG2243	76053	SENSE	514	1317	CGPG6886	76546	SENSE
112	915	CGPG2254	16120	SENSE	515	1318	CGPG6887	75848	SENSE
113	916	CGPG2268	15424	ANTI- SENSE	516	1319	CGPG6895	77515	SENSE
114	917	CGPG227	16401	SENSE	517	1320	CGPG6896	76266	SENSE
115	918	CGPG2312	72417	SENSE	518	1321	CGPG6903	75852	SENSE
116	919	CGPG2315	73702	SENSE	519	1322	CGPG6912	76556	SENSE
117	920	CGPG2316	70706	ANTI- SENSE	520	1323	CGPG6933	77523	SENSE
118	921	CGPG2319	71145	SENSE	521	1324	CGPG6942	75863	SENSE
119	922	CGPG235	11609	ANTI- SENSE	522	1325	CGPG6944	77052	SENSE
120	923	CGPG2358	71706	SENSE	523	1326	CGPG6949	75865	SENSE
121	924	CGPG2359	17325	SENSE	524	1327	CGPG6969	76551	SENSE
122	925	CGPG2361	70102	SENSE	525	1328	CGPG6989	76280	SENSE
123	926	CGPG2365	70106	SENSE	526	1329	CGPG6991	75874	SENSE
124	927	CGPG2372	70113	SENSE	527	1330	CGPG7003	75879	SENSE
125	928	CGPG2374	70115	SENSE	528	1331	CGPG7009	75880	SENSE
126	929	CGPG2377	70118	SENSE	529	1332	CGPG7037	77519	SENSE
127	930	CGPG2387	70128	SENSE	530	1333	CGPG7051	76745	SENSE
128	931	CGPG2389	70130	SENSE	531	1334	CGPG7062	76554	SENSE
129	932	CGPG2395	17103	ANTI- SENSE	532	1335	CGPG7064	76746	SENSE
130	933	CGPG2408	72748	SENSE	533	1336	CGPG7086	76750	SENSE
131	934	CGPG2409	17502	SENSE	534	1337	CGPG7109	76621	SENSE
132	935	CGPG2410	17503	SENSE	535	1338	CGPG7123	76623	SENSE
133	936	CGPG2414	78708	SENSE	536	1339	CGPG7136	76461	SENSE
134	937	CGPG2416	17504	SENSE	537	1340	CGPG7147	76462	SENSE
135	938	CGPG2441	17132	ANTI- SENSE	538	1341	CGPG7159	78987	SENSE
136	939	CGPG2450	72965	SENSE	539	1342	CGPG7186	76568	SENSE
137	940	CGPG2451	72796	SENSE	540	1343	CGPG7222	76176	SENSE
138	941	CGPG2492	16617	SENSE	541	1344	CGPG7224	76177	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
139	942	CGPG2495	16640	SENSE	542	1345	CGPG7233	78380	SENSE
140	943	CGPG2506	16603	SENSE	543	1346	CGPG7238	76629	SENSE
141	944	CGPG2515	16612	SENSE	544	1347	CGPG7253	76189	SENSE
142	945	CGPG2531	16614	SENSE	545	1348	CGPG7255	76761	SENSE
143	946	CGPG2581	78354	SENSE	546	1349	CGPG7286	78608	SENSE
144	947	CGPG2584	17818	SENSE	547	1350	CGPG7292	78115	SENSE
145	948	CGPG2592	78370	SENSE	548	1351	CGPG7301	74873	SENSE
146	949	CGPG2612	70411	SENSE	549	1352	CGPG7305	74826	SENSE
147	950	CGPG2660	19154	SENSE	550	1353	CGPG7307	74850	SENSE
148	951	CGPG2663	19155	SENSE	551	1354	CGPG7318	74887	SENSE
149	952	CGPG2679	18307	SENSE	552	1355	CGPG7319	74804	SENSE
150	953	CGPG2696	72030	SENSE	553	1356	CGPG7345	74831	SENSE
151	954	CGPG2772	17916	SENSE	554	1357	CGPG7368	74822	SENSE
152	955	CGPG2773	17917	SENSE	555	1358	CGPG7385	74836	SENSE
153	956	CGPG281	17334	SENSE	556	1359	CGPG7391	74901	SENSE
154	957	CGPG2846	71531	SENSE	557	1360	CGPG7405	74974	SENSE
155	958	CGPG2852	18441	SENSE	558	1361	CGPG7415	74904	SENSE
156	959	CGPG2863	19162	SENSE	559	1362	CGPG7416	74916	SENSE
157	960	CGPG2870	73205	SENSE	560	1363	CGPG7420	74964	SENSE
158	961	CGPG2877	18306	SENSE	561	1364	CGPG7432	74918	SENSE
159	962	CGPG289	70808	SENSE	562	1365	CGPG7436	74966	SENSE
160	963	CGPG2924	17653	SENSE	563	1366	CGPG7445	74979	SENSE
161	964	CGPG2947	19535	SENSE	564	1367	CGPG7465	74934	SENSE
162	965	CGPG2963	71719	SENSE	565	1368	CGPG7473	77814	SENSE
163	966	CGPG2987	18546	SENSE	566	1369	CGPG7489	75387	SENSE
164	967	CGPG3033	19534	SENSE	567	1370	CGPG7492	75328	SENSE
165	968	CGPG3045	18421	SENSE	568	1371	CGPG7499	77808	SENSE
166	969	CGPG3046	18422	SENSE	569	1372	CGPG7505	75389	SENSE
167	970	CGPG3060	19539	SENSE	570	1373	CGPG7507	75318	SENSE
168	971	CGPG3075	19244	SENSE	571	1374	CGPG7508	75330	SENSE
169	972	CGPG310	15803	SENSE	572	1375	CGPG7509	75342	SENSE
170	973	CGPG3103	19545	SENSE	573	1376	CGPG7511	75366	SENSE
171	974	CGPG315	10222	ANTI- SENSE	574	1377	CGPG7515	75319	SENSE
171	974	CGPG315	72356	SENSE	575	1378	CGPG7521	75391	SENSE
172	975	CGPG3183	70425	SENSE	576	1379	CGPG7540	75334	SENSE
173	976	CGPG3189	19237	SENSE	577	1380	CGPG7547	75323	SENSE
174	977	CGPG3204	71303	SENSE	578	1381	CGPG7561	75396	SENSE
175	978	CGPG3208	19240	SENSE	579	1382	CGPG7562	75401	SENSE
176	979	CGPG3219	18535	SENSE	580	1383	CGPG7563	75413	SENSE
177	980	CGPG3233	19450	SENSE	581	1384	CGPG7567	75461	SENSE
178	981	CGPG3263	18221	SENSE	582	1385	CGPG7568	75473	SENSE
179	982	CGPG3276	18233	SENSE	583	1386	CGPG7571	77817	SENSE
180	983	CGPG3280	18236	SENSE	584	1387	CGPG7587	75416	SENSE
181	984	CGPG3282	18238	SENSE	585	1388	CGPG7591	75464	SENSE
182	985	CGPG3300	18309	SENSE	586	1389	CGPG7597	75441	SENSE
183	986	CGPG3318	70337	SENSE	587	1390	CGPG7606	75454	SENSE
184	987	CGPG3319	18319	SENSE	588	1391	CGPG7633	75493	SENSE
185	988	CGPG3326	70338	SENSE	589	1392	CGPG7637	75446	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
186	989	CGPG333	10471	SENSE	590	1393	CGPG7649	75495	SENSE
187	990	CGPG3338	18331	SENSE	591	1394	CGPG7658	75705	SENSE
188	991	CGPG334	10228	ANTI- SENSE	592	1395	CGPG7664	77818	SENSE
188	991	CGPG334	10473	SENSE	593	1396	CGPG7666	75525	SENSE
189	992	CGPG3374	18261	SENSE	594	1397	CGPG7668	75549	SENSE
190	993	CGPG3402	18848	SENSE	595	1398	CGPG7742	75582	SENSE
191	994	CGPG3405	18268	SENSE	596	1399	CGPG7746	75535	SENSE
192	995	CGPG3413	18345	SENSE	597	1400	CGPG7747	75547	SENSE
193	996	CGPG3422	73607	SENSE	598	1401	CGPG7752	75512	SENSE
194	997	CGPG3436	18352	SENSE	599	1402	CGPG7766	75673	SENSE
195	998	CGPG3539	18357	SENSE	600	1403	CGPG7770	75626	SENSE
196	999	CGPG3550	19423	SENSE	601	1404	CGPG7774	75674	SENSE
197	1000	CGPG3551	78373	SENSE	602	1405	CGPG7777	75615	SENSE
198	1001	CGPG3552	70603	SENSE	603	1406	CGPG7778	75627	SENSE
199	1002	CGPG3572	19616	SENSE	604	1407	CGPG7786	75628	SENSE
200	1003	CGPG3598	19425	SENSE	605	1408	CGPG7788	75652	SENSE
201	1004	CGPG3599	19426	SENSE	606	1409	CGPG7792	75605	SENSE
202	1005	CGPG3606	18382	SENSE	607	1410	CGPG78	70228	SENSE
203	1006	CGPG3610	71536	SENSE	608	1411	CGPG783	13302	ANTI- SENSE
204	1007	CGPG3620	70419	SENSE	609	1412	CGPG7832	75610	SENSE
205	1008	CGPG364	70827	SENSE	610	1413	CGPG7841	75623	SENSE
206	1009	CGPG3676	19311	SENSE	611	1414	CGPG7845	75671	SENSE
207	1010	CGPG3679	19323	SENSE	612	1415	CGPG7847	75695	SENSE
208	1011	CGPG3686	19320	SENSE	613	1416	CGPG7851	75648	SENSE
209	1012	CGPG3694	73613	SENSE	614	1417	CGPG7853	75672	SENSE
210	1013	CGPG3696	70432	SENSE	615	1418	CGPG7857	75713	SENSE
211	1014	CGPG3698	70433	SENSE	616	1419	CGPG7865	75714	SENSE
212	1015	CGPG3699	70434	SENSE	617	1420	CGPG7869	75762	SENSE
213	1016	CGPG3702	72614	SENSE	618	1421	CGPG7891	77954	SENSE
214	1017	CGPG3703	70436	SENSE	619	1422	CGPG7892	77955	SENSE
215	1018	CGPG3707	74226	SENSE	620	1423	CGPG7906	77541	SENSE
216	1019	CGPG3710	70439	SENSE	621	1424	CGPG7924	78131	SENSE
217	1020	CGPG3730	70446	SENSE	622	1425	CGPG7934	78134	SENSE
218	1021	CGPG3731	70613	SENSE	623	1426	CGPG7949	77554	SENSE
219	1022	CGPG3734	70447	SENSE	624	1427	CGPG7954	78992	SENSE
220	1023	CGPG3745	71725	SENSE	625	1428	CGPG7964	77557	SENSE
221	1024	CGPG3764	70462	SENSE	626	1429	CGPG7968	78128	SENSE
222	1025	CGPG3820	70545	SENSE	627	1430	CGPG7969	77560	SENSE
223	1026	CGPG3851	70741	SENSE	628	1431	CGPG7972	77561	SENSE
224	1027	CGPG3911	19702	SENSE	629	1432	CGPG7982	77958	SENSE
225	1028	CGPG3948	19985	SENSE	630	1433	CGPG7985	77563	SENSE
226	1029	CGPG3958	19775	SENSE	631	1434	CGPG7993	78994	SENSE
227	1030	CGPG3981	19829	SENSE	632	1435	CGPG8	74508	SENSE
228	1031	CGPG3996	19982	SENSE	633	1436	CGPG80	70229	SENSE
229	1032	CGPG4006	19755	SENSE	634	1437	CGPG8001	77959	SENSE
230	1033	CGPG4025	19732	SENSE	635	1438	CGPG8009	77567	SENSE
231	1034	CGPG4028	70365	SENSE	636	1439	CGPG802	11749	ANTI-

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
									SENSE
232	1035	CGPG403	10476	SENSE	637	1440	CGPG8049	77578	SENSE
233	1036	CGPG4041	19987	SENSE	638	1441	CGPG806	11751	ANTI- SENSE
234	1037	CGPG4078	19893	SENSE	639	1442	CGPG8060	77963	SENSE
235	1038	CGPG4079	19983	SENSE	640	1443	CGPG8068	77925	SENSE
236	1039	CGPG4083	19825	SENSE	641	1444	CGPG8072	77927	SENSE
237	1040	CGPG4092	19919	SENSE	642	1445	CGPG8075	77345	SENSE
238	1041	CGPG4104	19979	SENSE	643	1446	CGPG81	70230	SENSE
239	1042	CGPG4127	19786	SENSE	644	1447	CGPG8100	77353	SENSE
240	1043	CGPG4135	70903	SENSE	645	1448	CGPG8101	77354	SENSE
241	1044	CGPG4161	70970	SENSE	646	1449	CGPG8104	77355	SENSE
242	1045	CGPG4180	70980	SENSE	647	1450	CGPG8108	77588	SENSE
243	1046	CGPG4191	70936	SENSE	648	1451	CGPG8116	77591	SENSE
244	1047	CGPG4199	71425	SENSE	649	1452	CGPG8125	77594	SENSE
245	1048	CGPG4241	78666	SENSE	650	1453	CGPG8134	77595	SENSE
246	1049	CGPG427	71202	SENSE	651	1454	CGPG8138	77364	SENSE
247	1050	CGPG4273	78973	SENSE	652	1455	CGPG8156	78904	SENSE
248	1051	CGPG4301	76411	SENSE	653	1456	CGPG8159	77937	SENSE
249	1052	CGPG4303	70635	SENSE	654	1457	CGPG8163	77369	SENSE
250	1053	CGPG4313	73715	SENSE	655	1458	CGPG8179	77940	SENSE
251	1054	CGPG4314	70638	SENSE	656	1459	CGPG8193	78746	SENSE
252	1055	CGPG4329	70642	SENSE	657	1460	CGPG8197	77372	SENSE
253	1056	CGPG4330	70643	SENSE	658	1461	CGPG8198	78126	SENSE
254	1057	CGPG4338	71564	SENSE	659	1462	CGPG8204	77951	SENSE
255	1058	CGPG4341	75039	SENSE	660	1463	CGPG8211	75901	SENSE
256	1059	CGPG4347	70653	SENSE	661	1464	CGPG823	12123	ANTI- SENSE
257	1060	CGPG4354	70655	SENSE	662	1465	CGPG8234	75987	SENSE
258	1061	CGPG4386	78321	SENSE	663	1466	CGPG8238	75940	SENSE
259	1062	CGPG4394	78307	SENSE	664	1467	CGPG8260	75919	SENSE
260	1063	CGPG4396	71808	SENSE	665	1468	CGPG8267	75908	SENSE
261	1064	CGPG4400	71810	SENSE	666	1469	CGPG8268	75920	SENSE
262	1065	CGPG4401	71313	SENSE	667	1470	CGPG8274	75992	SENSE
263	1066	CGPG441	72349	SENSE	668	1471	CGPG8277	75933	SENSE
264	1067	CGPG4413	71318	SENSE	669	1472	CGPG8279	77965	SENSE
265	1068	CGPG4427	71812	SENSE	670	1473	CGPG8284	77966	SENSE
266	1069	CGPG4446	74060	SENSE	671	1474	CGPG8345	77972	SENSE
267	1070	CGPG4448	71816	SENSE	672	1475	CGPG8408	78522	SENSE
268	1071	CGPG4474	71569	SENSE	673	1476	CGPG842	12194	ANTI- SENSE
269	1072	CGPG4482	70672	SENSE	674	1477	CGPG8421	78525	SENSE
270	1073	CGPG4511	71336	SENSE	675	1478	CGPG8435	78527	SENSE
271	1074	CGPG4517	71337	SENSE	676	1479	CGPG8440	78917	SENSE
272	1075	CGPG4551	71339	SENSE	677	1480	CGPG8453	78920	SENSE
273	1076	CGPG4559	71825	SENSE	678	1481	CGPG8470	78535	SENSE
274	1077	CGPG4567	73696	SENSE	679	1482	CGPG8479	78538	SENSE
275	1078	CGPG4586	70682	SENSE	680	1483	CGPG8489	78922	SENSE
276	1079	CGPG4600	70686	SENSE	681	1484	CGPG8498	77984	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
277	1080	CGPG4631	78316	SENSE	682	1485	CGPG85	73938	SENSE
278	1081	CGPG4642	71667	SENSE	683	1486	CGPG8501	78621	SENSE
279	1082	CGPG4645	71695	SENSE	684	1487	CGPG8507	77990	SENSE
280	1083	CGPG4646	71696	SENSE	685	1488	CGPG8509	78386	SENSE
281	1084	CGPG4649	72475	SENSE	686	1489	CGPG8515	78001	SENSE
282	1085	CGPG4653	71677	SENSE	687	1490	CGPG8517	78543	SENSE
283	1086	CGPG4656	71691	SENSE	688	1491	CGPG8518	78002	SENSE
284	1087	CGPG4668	72476	SENSE	689	1492	CGPG852	12259	SENSE
285	1088	CGPG469	70803	SENSE	690	1493	CGPG8523	78005	SENSE
286	1089	CGPG4708	71612	SENSE	691	1494	CGPG8528	78008	SENSE
287	1090	CGPG4712	71637	SENSE	692	1495	CGPG8552	78019	SENSE
288	1091	CGPG4714	71629	SENSE	693	1496	CGPG8553	78020	SENSE
289	1092	CGPG4719	71622	SENSE	694	1497	CGPG8562	78023	SENSE
290	1093	CGPG473	70810	SENSE	695	1498	CGPG8567	78548	SENSE
291	1094	CGPG4734	72452	SENSE	696	1499	CGPG8580	78161	SENSE
292	1095	CGPG4736	74071	SENSE	697	1500	CGPG8589	78162	SENSE
293	1096	CGPG474	11042	SENSE	698	1501	CGPG8590	78033	SENSE
294	1097	CGPG4802	72510	SENSE	699	1502	CGPG8592	78622	SENSE
295	1098	CGPG4822	72537	SENSE	700	1503	CGPG8594	78036	SENSE
296	1099	CGPG4826	72541	SENSE	701	1504	CGPG8606	78043	SENSE
297	1100	CGPG4833	72617	SENSE	702	1505	CGPG8628	78165	SENSE
298	1101	CGPG4841	73755	SENSE	703	1506	CGPG8648	78066	SENSE
299	1102	CGPG4850	72630	SENSE	704	1507	CGPG8654	78069	SENSE
300	1103	CGPG4868	72645	SENSE	705	1508	CGPG8668	78183	SENSE
301	1104	CGPG4871	72647	SENSE	706	1509	CGPG8678	78170	SENSE
302	1105	CGPG488	14316	SENSE	707	1510	CGPG8694	78953	SENSE
303	1106	CGPG4908	73689	SENSE	708	1511	CGPG8702	78929	SENSE
304	1107	CGPG4913	73345	SENSE	709	1512	CGPG8737	78564	SENSE
305	1108	CGPG4921	73347	SENSE	710	1513	CGPG8745	78566	SENSE
306	1109	CGPG4954	73231	SENSE	711	1514	CGPG8748	78591	SENSE
307	1110	CGPG4956	78325	SENSE	712	1515	CGPG8749	78181	SENSE
308	1111	CGPG4959	72658	SENSE	713	1516	CGPG8770	78190	SENSE
309	1112	CGPG4965	72660	SENSE	714	1517	CGPG8777	78571	SENSE
310	1113	CGPG4970	72662	SENSE	715	1518	CGPG8781	78933	SENSE
311	1114	CGPG4980	73352	SENSE	716	1519	CGPG8786	78573	SENSE
312	1115	CGPG4982	72814	SENSE	717	1520	CGPG8792	78632	SENSE
313	1116	CGPG4985	72816	SENSE	718	1521	CGPG8801	78959	SENSE
314	1117	CGPG4990	72820	SENSE	719	1522	CGPG8809	78191	SENSE
315	1118	CGPG4991	73353	SENSE	720	1523	CGPG8840	78595	SENSE
316	1119	CGPG5007	75066	SENSE	721	1524	CGPG8850	78646	SENSE
317	1120	CGPG5015	78337	SENSE	722	1525	CGPG8870	76325	SENSE
318	1121	CGPG5026	73317	SENSE	723	1526	CGPG8871	76337	SENSE
319	1122	CGPG5029	73318	SENSE	724	1527	CGPG8872	76349	SENSE
320	1123	CGPG5046	73749	SENSE	725	1528	CGPG8876	76302	SENSE
321	1124	CGPG508	18704	SENSE	726	1529	CGPG8902	76329	SENSE
322	1125	CGPG5103	73221	SENSE	727	1530	CGPG8904	76353	SENSE
323	1126	CGPG511	71125	SENSE	728	1531	CGPG8907	76389	SENSE
324	1127	CGPG5121	76203	SENSE	729	1532	CGPG8914	76378	SENSE
325	1128	CGPG5126	75067	SENSE	730	1533	CGPG8931	76392	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
326	1129	CGPG5136	73729	SENSE	731	1534	CGPG8933	76321	SENSE
327	1130	CGPG5146	75206	SENSE	732	1535	CGPG8935	76345	SENSE
328	1131	CGPG5149	73248	SENSE	733	1536	CGPG8938	76381	SENSE
329	1132	CGPG5181	73738	SENSE	734	1537	CGPG8944	76358	SENSE
330	1133	CGPG52	73332	SENSE	735	1538	CGPG895	12629	SENSE
331	1134	CGPG5206	78335	SENSE	736	1539	CGPG8950	76335	SENSE
332	1135	CGPG5208	75821	SENSE	737	1540	CGPG8961	76372	SENSE
333	1136	CGPG5232	72038	SENSE	738	1541	CGPG8963	76396	SENSE
334	1137	CGPG5239	72027	SENSE	739	1542	CGPG898	72314	SENSE
335	1138	CGPG5246	72016	SENSE	740	1543	CGPG8993	77840	SENSE
336	1139	CGPG525	70835	SENSE	741	1544	CGPG8994	76886	SENSE
337	1140	CGPG5268	72044	SENSE	742	1545	CGPG8995	76803	SENSE
338	1141	CGPG5272	72092	SENSE	743	1546	CGPG90	10302	ANTI- SENSE
339	1142	CGPG5333	72124	SENSE	744	1547	CGPG900	71107	SENSE
340	1143	CGPG5338	72110	SENSE	745	1548	CGPG9002	76887	SENSE
341	1144	CGPG5341	73916	SENSE	746	1549	CGPG9009	77842	SENSE
342	1145	CGPG5369	74268	SENSE	747	1550	CGPG9011	76805	SENSE
343	1146	CGPG5372	74269	SENSE	748	1551	CGPG9012	76817	SENSE
344	1147	CGPG5380	77902	SENSE	749	1552	CGPG9017	77843	SENSE
345	1148	CGPG5386	74273	SENSE	750	1553	CGPG9025	77844	SENSE
346	1149	CGPG5396	74280	SENSE	751	1554	CGPG9026	76890	SENSE
347	1150	CGPG5397	74281	SENSE	752	1555	CGPG9032	77839	SENSE
348	1151	CGPG5421	74287	SENSE	753	1556	CGPG9040	76868	SENSE
349	1152	CGPG5433	73767	SENSE	754	1557	CGPG9044	76821	SENSE
350	1153	CGPG5439	77304	SENSE	755	1558	CGPG9048	76869	SENSE
351	1154	CGPG5453	73761	SENSE	756	1559	CGPG9049	76881	SENSE
352	1155	CGPG5456	74721	SENSE	757	1560	CGPG9058	76894	SENSE
353	1156	CGPG5483	74719	SENSE	758	1561	CGPG906	11930	ANTI- SENSE
354	1157	CGPG5492	74257	SENSE	759	1562	CGPG9070	76848	SENSE
355	1158	CGPG5508	72749	SENSE	760	1563	CGPG9075	76901	SENSE
356	1159	CGPG5520	72703	SENSE	761	1564	CGPG9084	76914	SENSE
357	1160	CGPG5525	72763	SENSE	762	1565	CGPG9094	76939	SENSE
358	1161	CGPG5526	72775	SENSE	763	1566	CGPG9098	76987	SENSE
359	1162	CGPG5530	72728	SENSE	764	1567	CGPG9099	76904	SENSE
360	1163	CGPG5534	72776	SENSE	765	1568	CGPG9110	76941	SENSE
361	1164	CGPG5537	72717	SENSE	766	1569	CGPG9119	77113	SENSE
362	1165	CGPG5545	72718	SENSE	767	1570	CGPG9125	77185	SENSE
363	1166	CGPG555	11715	SENSE	768	1571	CGPG913	12911	SENSE
364	1167	CGPG5558	72779	SENSE	769	1572	CGPG9136	77127	SENSE
365	1168	CGPG5559	72791	SENSE	770	1573	CGPG9156	77177	SENSE
366	1169	CGPG5562	72732	SENSE	771	1574	CGPG9164	77178	SENSE
367	1170	CGPG5592	73027	SENSE	772	1575	CGPG9172	77179	SENSE
368	1171	CGPG5625	73058	SENSE	773	1576	CGPG9173	77191	SENSE
369	1172	CGPG5631	73126	SENSE	774	1577	CGPG9185	77145	SENSE
370	1173	CGPG5632	73138	SENSE	775	1578	CGPG9187	77169	SENSE
371	1174	CGPG5636	73174	SENSE	776	1579	CGPG9190	77110	SENSE
372	1175	CGPG5637	73186	SENSE	777	1580	CGPG9193	77146	SENSE

NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation	NUC SEQ ID NO	PEP SEQ ID NO	Gene ID	Construct ID	Orientation
373	1176	CGPG564	11141	SENSE	778	1581	CGPG9195	77170	SENSE
374	1177	CGPG5641	73139	SENSE	779	1582	CGPG9203	77171	SENSE
375	1178	CGPG5650	73128	SENSE	780	1583	CGPG9209	77148	SENSE
376	1179	CGPG5653	73164	SENSE	781	1584	CGPG921	11933	ANTI- SENSE
377	1180	CGPG5657	73117	SENSE	782	1585	CGPG9210	77160	SENSE
378	1181	CGPG5671	73142	SENSE	783	1586	CGPG9211	77172	SENSE
379	1182	CGPG5672	73990	SENSE	784	1587	CGPG9212	77184	SENSE
380	1183	CGPG5677	73166	SENSE	785	1588	CGPG9216	77225	SENSE
381	1184	CGPG5678	73178	SENSE	786	1589	CGPG9226	77250	SENSE
382	1185	CGPG5679	73190	SENSE	787	1590	CGPG9252	77277	SENSE
383	1186	CGPG5682	73131	SENSE	788	1591	CGPG9281	77245	SENSE
384	1187	CGPG5691	73108	SENSE	789	1592	CGPG9289	77246	SENSE
385	1188	CGPG5699	73074	SENSE	790	1593	CGPG9298	77462	SENSE
386	1189	CGPG5705	73132	SENSE	791	1594	CGPG9299	77414	SENSE
387	1190	CGPG5722	73146	SENSE	792	1595	CGPG9304	77439	SENSE
388	1191	CGPG5734	73147	SENSE	793	1596	CGPG9310	77463	SENSE
389	1192	CGPG5736	73080	SENSE	794	1597	CGPG9316	77488	SENSE
390	1193	CGPG5744	73077	SENSE	795	1598	CGPG9317	77441	SENSE
391	1194	CGPG5747	73183	SENSE	796	1599	CGPG9318	77464	SENSE
392	1195	CGPG5748	73018	SENSE	797	1600	CGPG9324	77442	SENSE
393	1196	CGPG5750	73112	SENSE	798	1601	CGPG9357	77419	SENSE
394	1197	CGPG5751	73988	SENSE	799	1602	CGPG965	12444	SENSE
395	1198	CGPG577	10908	ANTI- SENSE	800	1603	CGPG970	11791	ANTI- SENSE
395	1198	CGPG577	11147	SENSE	800	1603	CGPG970	12445	SENSE
396	1199	CGPG5780	72909	SENSE	801	1604	CGPG982	71249	SENSE
397	1200	CGPG5793	73987	SENSE	802	1605	CGPG994	71237	SENSE
398	1201	CGPG5795	72922	SENSE	803	1606	CGPG996	12315	ANTI- SENSE
/	/	/	/	/	803	1606	CGPG996	12366	SENSE

Recombinant DNA

DNA for use in the present invention to improve traits in plants have a nucleotide sequence of SEQ ID NO:1 through SEQ ID NO:803, as well as the homologs of such DNA molecules. A subset of the DNA for gene suppression aspects of the invention includes fragments of the disclosed full polynucleotides consisting of oligonucleotides of 21 or more consecutive nucleotides. In some embodiments, the nucleotides for gene suppression can be 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more consecutive nucleotides. Oligonucleotides having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 803 can be as probes and primers for detection of the polynucleotides used in the invention. Other embodiments this invention are variants of the DNA. Such variants can be naturally occurring, including DNA from homologous genes from the same or

a different species, or can be non-natural variants, for example DNA synthesized using chemical synthesis methods, or generated using recombinant DNA techniques. Degeneracy of the genetic code provides the possibility to substitute at least one base of the protein encoding sequence of a gene with a different base without causing the amino acid sequence of the polypeptide produced from the gene to be changed. Hence, a DNA in the present invention can have any base sequence that has been changed from the sequences provided herein by substitution in accordance with degeneracy of the genetic code.

Homologs of the genes providing DNA demonstrated uses in improving traits in model plants disclosed herein will generally have significant identity with the DNA disclosed herein. DNA is substantially identical to a reference DNA if, when the sequences of the polynucleotides are optimally aligned there is about 60% nucleotide equivalence; about 70% equivalence; about 80% equivalence; about 85% equivalence; about 90%; about 95%; about 98% , about 99% equivalence or about 99.5 equivalence over a comparison window. A comparison window can be at least 50-100 nucleotides or the entire length of the polynucleotide provided herein. Optimal alignment of sequences for aligning a comparison window can be conducted by algorithms; for example by computerized implementations of these algorithms (such as, the Wisconsin Genetics Software Package Release 7.0-10.0, Genetics Computer Group, 575 Science Dr., Madison, WI). The reference polynucleotide can be a full-length molecule or a portion of a longer molecule. In one embodiment, the window of comparison for determining polynucleotide identity of protein encoding sequences is the entire coding region.

Proteins used for imparting enhanced traits are entire proteins or at least a sufficient portion of the entire protein to impart the relevant biological activity of the protein. Proteins used for generation of transgenic plants having enhanced traits include the proteins with an amino acid sequence provided herein as SEQ ID NO: 804 through SEQ ID NO: 1606, as well as homologs of such proteins.

Homologs of the proteins in the invention are identified by comparison of the amino acid sequence of the protein to amino acid sequences of proteins from the same or different plant sources, *e.g.*, manually or by using known homology-based search algorithms such as those commonly known and referred to as BLAST, FASTA, and Smith-Waterman. As used herein, a homolog is a protein from the same or a different organism that performs the same biological function as the polypeptide to which it is compared. An orthologous relation between two organisms is not necessarily manifest as a one-to-one correspondence between two genes, because a gene can be duplicated or deleted after organism phylogenetic

separation, such as speciation. For a given protein, there can be no ortholog or more than one ortholog. Other complicating factors include alternatively spliced transcripts from the same gene, limited gene identification, redundant copies of the same gene with different sequence lengths or corrected sequence. A local sequence alignment program, *e.g.*, BLAST, can be used to search a database of sequences to find similar sequences, and the summary Expectation value (E-value) used to measure the sequence base similarity. As a protein hit with the best E-value for a particular organism could be an ortholog or the only ortholog, a reciprocal BLAST search is used in the present invention to filter hit sequences with significant E-values for ortholog identification. The reciprocal BLAST entails search of the significant hits against a database of amino acid sequences from the base organism that are similar to the sequence of the query protein. A hit is a likely ortholog, when the reciprocal BLAST's best hit is the query protein itself or a protein encoded by a duplicated gene after speciation. Thus, homolog is used herein to describe proteins that are assumed to have functional similarity by inference from sequence base similarity. The relationship of homologs with amino acid sequences of SEQ ID NO: 1607 to SEQ ID NO: 94613 to the proteins with amino acid sequences of SEQ ID NO: 804 to SEQ ID NO: 1606 are found in the listing of Table 19.

Other functional homolog proteins differ in one or more amino acids from those of a trait-improving protein disclosed herein as the result of one or more of the well-known conservative amino acid substitutions, *e.g.*, valine is a conservative substitute for alanine and threonine is a conservative substitute for serine. Conservative substitutions for an amino acid within the native sequence can be selected from other members of a class to which the naturally occurring amino acid belongs. Representative amino acids within these various classes include, but are not limited to: (1) acidic (negatively charged) amino acids such as aspartic acid and glutamic acid; (2) basic (positively charged) amino acids such as arginine, histidine, and lysine; (3) neutral polar amino acids such as glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; and (4) neutral nonpolar (hydrophobic) amino acids such as alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine. Conserved substitutes for an amino acid within a native amino acid sequence can be selected from other members of the group to which the naturally occurring amino acid belongs. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is

phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Naturally conservative amino acids substitution groups are: valine-leucine, valine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine. A further aspect of the invention comprises proteins that differ in one or more amino acids from those of a described protein sequence as the result of deletion or insertion of one or more amino acids in a native sequence.

Homologs of the trait-improving proteins provided herein will generally demonstrate significant sequence identity. Of particular interest are proteins having at least about 50% sequence identity, at least about 70% sequence identity or higher, *e.g.*, at least about 80% sequence identity with an amino acid sequence of SEQ ID NO: 804 to SEQ ID NO: 1606. Additional embodiments also include those with higher identity, *e.g.*, about 90%, about 92.5%, about 95%, about 98%, about 99%, or to about 99.5% identity. Identity of protein homologs is determined by optimally aligning the amino acid sequence of a putative protein homolog with a defined amino acid sequence and by calculating the percentage of identical and conservatively substituted amino acids over the window of comparison. The window of comparison for determining identity can be the entire amino acid sequence disclosed herein, *e.g.*, the full sequence of any of SEQ ID NO: 804 to SEQ ID NO: 1606.

Genes that are homologous to each other can be grouped into families and included in multiple sequence alignments. Then a consensus sequence for each group can be derived. This analysis enables the derivation of conserved and class- (family) specific residues or motifs that are functionally important. These conserved residues and motifs can be further validated with 3D protein structure if available. The consensus sequence can be used to define the full scope of the invention, *e.g.*, to identify proteins with a homolog relationship. Thus, the present invention contemplates that protein homologs include proteins with an amino acid sequence that has at least 90% identity to such a consensus amino acid sequence sequences.

Discovery of Trait-improving Recombinant DNA

To identify nuclei with recombinant DNA that confer enhanced traits to plants, *Arabidopsis thaliana* was transformed with a candidate recombinant DNA construct and screened for an enhanced trait.

Arabidopsis thaliana is used a model for genetics and metabolism in plants. *Arabidopsis* has a small genome, and well-documented studies are available. It is easy to grow in large numbers and mutants defining important genetically controlled mechanisms are either available, or can readily be obtained. Various methods to introduce and express isolated homologous genes are available (see Koncz, *et al.*, Methods in *Arabidopsis* Research *et al.*, (1992), World Scientific, New Jersey, New Jersey, in “Preface”).

A two-step screening process was employed which comprised two passes of trait characterization to ensure that the trait modification was dependent on expression of the recombinant DNA, but not due to the chromosomal location of the integration of the transgene. Twelve independent transgenic lines for each recombinant DNA construct were established and assayed for the transgene expression levels. Five transgenic lines with high transgene expression levels were used in the first pass screen to evaluate the transgene’s function in T2 transgenic plants. Subsequently, three transgenic events, which had been shown to have one or more enhanced traits, were further evaluated in the second pass screen to confirm the transgene’s ability to impart an enhanced trait. The following Table 3 summarizes the enhanced traits that have been confirmed as provided by a recombinant DNA construct.

In particular, Table 3 reports:

“PEP SEQ ID” which is the amino acid sequence of the protein cognate to the DNA in the recombinant DNA construct corresponding to a protein sequence of a SEQ ID NO. in the Sequence Listing.

“construct_id” is an arbitrary name for the recombinant DNA describe more particularly in Table 1.

“annotation” refers to a description of the top hit protein obtained from an amino acid sequence query of each PEP SEQ ID NO to GenBank database of the National Center for Biotechnology Information (ncbi). More particularly, “gi” is the GenBank ID number for the top BLAST hit.

“description” refers to the description of the top BLAST hit.

“e-value” provides the expectation value for the BLAST hit.

“ % id” refers to the percentage of identically matched amino acid residues along the length of the portion of the sequences which is aligned by BLAST between the sequence of interest provided herein and the hit sequence in GenBank.

“traits” identify by two letter codes the confirmed enhancement in a transgenic plant provided by the recombinant DNA . The codes for enhanced traits are:

“CK” which indicates cold tolerance enhancement identified under a cold shock tolerance screen;

“CS” which indicates cold tolerance enhancement identified by a cold germination tolerance screen;

5 “DS” which indicates drought tolerance enhancement identified by a soil drought stress tolerance screen;

“PEG” which indicates osmotic stress tolerance enhancement identified by a PEG induced osmotic stress tolerance screen;

10 “HS” which indicates heat stress tolerance enhancement identified by a heat stress tolerance screen;

“SS” which indicates high salinity stress tolerance enhancement identified by a salt stress tolerance screen;

“LN” which indicates nitrogen use efficiency enhancement identified by a limited nitrogen tolerance screen;

15 “LL” which indicates attenuated shade avoidance response identified by a shade tolerance screen under a low light condition;

“PP” which indicates enhanced growth and development at early stages identified by an early plant growth and development screen;

20 “SP” which indicates enhanced growth and development at late stages identified by a late plant growth and development screen provided herein.

Table 3.

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
804	70244	0	95	ref NP_193113.1 CYP83A1 (CYTOCHROME P450 83A1); oxygen binding [Arabidopsis thaliana]	HS			
805	71108	3.00E -54	100	gb AAB61093.1 F20P5.4 gene product [Arabidopsis thaliana]	LL			
806	12816	0	79	ref NP_190754.2 CAX3 (cation exchanger 3); cation:cation antiporter [Arabidopsis thaliana]	LN			
807	12650	1.00E -159	89	gb AAC25513.1 AAC25513Strong similarity to hypothetical protein gb Y09823 from A. thaliana.	LN			
808	12031	1.00E -56	88	ref NP_179413.1 unknown protein [Arabidopsis thaliana]	LN			
809	12217	1.00E -168	88	ref NP_194233.1 unknown protein [Arabidopsis thaliana]	LN			
810	12043	2.00E -46	100	ref NP_194730.1 unknown protein [Arabidopsis thaliana]	PP			
811	12147	2.00E -33	68	ref NP_190718.1 unknown protein [Arabidopsis thaliana]	LN			
812	13638	0	100	ref NP_189940.1 unknown protein [Arabidopsis thaliana]	DS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
813	11860	2.00E -81	93	ref NP_973419.1 unknown protein [Arabidopsis thaliana]	LN			
814	72361	1.00E -139	93	ref NP_178534.2 IBR5 (INDOLE-3-BUTYRIC ACID RESPONSE 5); protein tyrosine/serine/threonine phosphatase [Arabidopsis thaliana]	CK			
815	12230	1.00E -23	100	ref NP_564545.1 TOM6 (TRANSLOCASE OF THE OUTER MITOCHONDRIAL MEMBRANE 6) [Arabidopsis thaliana] sp Q9XIA7 TOM6_ARATH Mitochondrial import receptor subunit TOM6 homolog (Translocase of outer membrane 6 kDa subunit homolog)	LN			
816	71626	1.00E -47	93	ref NP_177455.1 unknown protein [Arabidopsis thaliana]	CK			
817	13454	0	96	ref NP_566136.1 armadillo/beta-catenin repeat family protein [Arabidopsis thaliana]	LN			
818	12173	4.00E -49	100	ref NP_186751.1 ATISU2/ISU2 (IscU-like 2); structural molecule [Arabidopsis thaliana]	CK			
819	17806	1.00E -35	54	ref NP_188341.2 unknown protein [Arabidopsis thaliana]	LN			
820	12011	1.00E -144	99	ref NP_567393.1 ATP-binding family protein [Arabidopsis thaliana]	LN			
821	72915	1.00E -135	77	sp Q94A43 BEH2_ARATHBES1/BZR1 homolog protein 2	SS			
822	13231	1.00E -161	96	ref NP_178361.2 nucleic acid binding [Arabidopsis thaliana]	HS			
823	13913	1.00E -147	88	ref NP_028242.1 PDV2 (PLASTID DIVISION2) [Arabidopsis thaliana] gb AAD26950.1 expressed protein [Arabidopsis thaliana]	HS			
824	12749	1.00E -140	100	ref NP_196620.1 unknown protein [Arabidopsis thaliana]	LN			
825	78901	0	97	ref NP_565919.1 COI1 (CORONATINE INSENSITIVE 1); ubiquitin-protein ligase [Arabidopsis thaliana]	HS	PP		
826	75969	0	100	ref NP_174005.1 ATCUL3/ATCUL3A/CUL3/CUL3A (Cullin 3A); protein binding / ubiquitin-protein ligase [Arabidopsis thaliana]	CS	HS		
827	15606	0	96	gb AAA32781.1 cyclin emb CAA44169.1 cyclin [Arabidopsis thaliana]	HS			
828	75946	0	96	ref NP_171797.2 CUL2 (cullin 2) [Arabidopsis thaliana]	LL			
829	13050	0	97	ref NP_175955.1 F-box family protein [Arabidopsis thaliana]	LN			
830	75958	0	94	ref NP_177170.1 lectin protein kinase, putative [Arabidopsis thaliana]	HS	LN		
831	17333	0	93	gb AAF02796.1 AF195115_16Similar to receptor-like protein kinase precursor; F5110.19 [Arabidopsis thaliana]	PP			
832	76002	0	100	ref NP_193038.1 MHK; kinase [Arabidopsis thaliana] sp P43294 MHK_ARATH Serine/threonine-protein kinase MHK	HS	PEG		
833	78735	0	94	ref NP_187132.1 protein kinase, putative [Arabidopsis thaliana]	LL	PEG		
834	13914	0	100	ref NP_195033.1 pyruvate decarboxylase, putative [Arabidopsis thaliana]	HS	SS		
835	14405	0	92	ref NP_187341.1 DIN3/LTA1 (DARK INDUCIBLE 3); alpha-ketoacid dehydrogenase [Arabidopsis thaliana]	LN	SP		
836	75983	0	100	ref NP_171655.1 ACS2 (1-Amino-cyclopropane-1-carboxylate synthase 2) [Arabidopsis thaliana]	CK			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
837	70721	0	94	ref NP_851005.1 LPD2 (LIPOAMIDE DEHYDROGENASE 2); FAD binding / dihydrolipoyl dehydrogenase/ disulfide oxidoreductase/ oxidoreductase [Arabidopsis thaliana]	HS	LN		
838	13811	0	92	ref NP_178093.1 ATNADP-ME4 (NADP-MALIC ENZYME 4); malate dehydrogenase (oxaloacetate-decarboxylating) (NADP+)/ malic enzyme/ oxidoreductase, acting on NADH or NADPH, NAD or NADP as acceptor [Arabidopsis thaliana]	DS			
838	13828	0	92	ref NP_178093.1 ATNADP-ME4 (NADP-MALIC ENZYME 4); malate dehydrogenase (oxaloacetate-decarboxylating) (NADP+)/ malic enzyme/ oxidoreductase, acting on NADH or NADPH, NAD or NADP as acceptor [Arabidopsis thaliana]	LN			
839	13614	1.00E -122	96	ref NP_176518.1 ribulose-phosphate 3-epimerase, cytosolic, putative / pentose-5-phosphate 3-epimerase, putative [Arabidopsis thaliana]	LN			
840	76050	0	97	ref NP_567714.1 AMY1 (ALPHA-AMYLASE-LIKE); alpha-amylase [Arabidopsis thaliana]	CS			
841	13038	0	100	ref NP_194382.1 fructose-bisphosphate aldolase, cytoplasmic [Arabidopsis thaliana]	LN			
842	13040	1.00E -104	92	gb AAB66906.1 eukaryotic initiation factor (iso)-4F p28 subunit [Arabidopsis thaliana]	LN			
843	70716	1.00E -172	100	ref NP_974170.1 serine/threonine protein kinase, putative [Arabidopsis thaliana]	HS			
843	76039	0	99	ref NP_001078396.1 PPDK (PYRUVATE ORTHOPHOSPHATE DIKINASE) [Arabidopsis thaliana]	HS			
844	71611	0	99	ref NP_001078396.1 PPDK (PYRUVATE ORTHOPHOSPHATE DIKINASE) [Arabidopsis thaliana]	CK	PP		
845	70253	0	100	ref NP_200694.1 CYP86A1 (cytochrome P450, family 86, subfamily A, polypeptide 1); oxygen binding [Arabidopsis thaliana]	HS	CK		
846	14415	0	96	ref NP_568054.1 SEC14 cytosolic factor, putative / phosphoglyceride transfer protein, putative [Arabidopsis thaliana]	LN			
847	13718	4.00E -56	100	ref NP_196413.1 unknown protein [Arabidopsis thaliana] sp Q9SD88 U139_ARATH UPF0139 protein At5g07960	LN			
848	10706	0	88	gb AAA19118.1 glutamyl-tRNA reductase	CK	HS		
848	10709	0	88	gb AAA19118.1 glutamyl-tRNA reductase	LN			
849	14340	1.00E -125	93	ref NP_175332.1 unknown protein [Arabidopsis thaliana]	LN			
850	13471	1.00E -136	95	ref NP_565858.1 unknown protein [Arabidopsis thaliana]	LN			
851	18304	1.00E -133	94	ref NP_189283.1 TIP2 (TONOPLAST INTRINSIC PROTEIN 2); water channel [Arabidopsis thaliana]	HS			
852	13623	0	96	ref NP_176001.2 GTP-binding protein-related [Arabidopsis thaliana]	DS			
853	18302	0	100	ref NP_194702.2 NFC5 (NUCLEOSOME/CHROMATIN ASSEMBLY FACTOR GROUP C 5) [Arabidopsis thaliana] sp Q9SU78 MSI5_ARATH WD-40 repeat-containing protein MSI5	HS			
854	15053	0	97	ref NP_001031261.1 unknown protein [Arabidopsis thaliana]	LN			
855	15624	0	93	ref NP_171975.2 RabGAP/TBC domain-containing protein [Arabidopsis thaliana]	LL	PP		
856	70412	0	81	ref NP_179257.1 nodulin family protein [Arabidopsis thaliana]	HS			
857	17902	0	91	ref NP_564843.1 integral membrane transporter family protein [Arabidopsis thaliana]	PP			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
858	14272	0	90	ref NP_566584.1 YSL5 (YELLOW STRIPE LIKE 5); oligopeptide transporter [Arabidopsis thaliana]	LN			
859	15122	0	95	ref NP_565817.2 glycogenin glucosyltransferase (glycogenin)-related [Arabidopsis thaliana]	LN			
859	70723	0	95	ref NP_565817.2 glycogenin glucosyltransferase (glycogenin)-related [Arabidopsis thaliana]	HS			
860	14248	1.00E -154	79	ref NP_176081.1 unknown protein [Arabidopsis thaliana]	DS			
861	13965	0	100	ref NP_173104.1 Per1-like family protein [Arabidopsis thaliana]	CK			
862	75912	1.00E -169	100	ref NP_199075.1 unknown protein [Arabidopsis thaliana]	CS	LL		
863	14344	1.00E -81	100	ref NP_566785.1 universal stress protein (USP) family protein [Arabidopsis thaliana]	CS			
864	14807	0	89	gb AAD20391.1 hypothetical protein [Arabidopsis thaliana]	CS			
865	17307	0	89	db BAE98505.1 hypothetical protein [Arabidopsis thaliana]	PP			
866	16431	0	73	ref NP_201248.1 octicosapeptide/Phox/Bem1p (PB1) domain-containing protein [Arabidopsis thaliana]	LN			
867	73075	0	96	ref NP_187459.2 protein binding / zinc ion binding [Arabidopsis thaliana]	CS			
868	16323	9.00E -57	89	ref NP_171768.2 transcription initiation factor IID (TFIID) 18 kDa subunit (TAFII-18) family protein [Arabidopsis thaliana]	LN			
869	14833	1.00E -141	90	ref NP_176342.1 CCR4-NOT transcription complex protein, putative [Arabidopsis thaliana]	SP			
870	17030	0	95	gb AAG28899.1 AC008113_15F12A21.29 [Arabidopsis thaliana]	LL			
871	76414	0	99	ref NP_180595.1 CIPK11 (SNF1-RELATED PROTEIN KINASE 3.22, SOS3-INTERACTING PROTEIN 4); kinase [Arabidopsis thaliana]	PEG			
872	16021	0	82	ref NP_201301.1 CDKC;2 (CYCLIN-DEPENDENT KINASE C;2); kinase [Arabidopsis thaliana]	LN			
873	78701	0	93	ref NP_564203.1 F-box family protein [Arabidopsis thaliana]	HS			
874	78702	0	93	ref NP_565142.1 F-box family protein (FBX3) [Arabidopsis thaliana]	PEG			
875	15220	0	89	ref NP_173623.1 kelch repeat-containing F-box family protein [Arabidopsis thaliana] sp Q9LM55 FBK8_ARATH F-box/Kelch-repeat protein	LN			
876	15623	0	88	ref NP_179623.1 transducin family protein / WD-40 repeat family protein [Arabidopsis thaliana]	HS			
877	19402	0	96	ref NP_568408.1 F-box family protein / WD-40 repeat family protein [Arabidopsis thaliana]	PP			
878	70559	0	98	ref NP_181061.1 SUVH5 (SU(VAR)3-9 HOMOLOG 5) [Arabidopsis thaliana]	LN			
879	15111	0	90	ref NP_197821.1 ATXR6 (Arabidopsis thaliana Trithorax-related protein 6); DNA binding sp Q9FNE9 ATXR6_ARATH Histone-lysine N-methyltransferase ATXR6 (Trithorax-related protein 6) (TRX-related protein 6)	DS			
880	70832	0	100	ref NP_172633.1 CYP51G1 (CYTOCHROME P450 51); oxygen binding [Arabidopsis thaliana]	HS			
881	15129	4.00E -82	77	ref NP_188372.1 PHD finger family protein [Arabidopsis thaliana]	LN			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
882	76086	0	91	ref NP_192814.1 EMB1706 (EMBRYO DEFECTIVE 1706); S-adenosylmethionine-dependent methyltransferase [Arabidopsis thaliana]	PP			
883	15203	1.00E -158	94	gb AAD30650.1 AC006085_23Similar to human CGI-33 protein [Arabidopsis thaliana]	CK			
884	70216	0	95	ref NP_567665.2 CYP706A1 (cytochrome P450, family 706, subfamily A, polypeptide 1); oxygen binding [Arabidopsis thaliana]	HS	PP		
885	14910	0	100	ref NP_201191.1 UVR8 (UVB-RESISTANCE 8) [Arabidopsis thaliana]	DS			
886	70318	0	92	ref NP_172325.2 ATCPNIFS/ATNFS2/ATSUFS/CPNIFS/S UFS (CHLOROPLASTIC NIFS-LIKE CYSTEINE DESULFURASE); cysteine desulfurase/ selenocysteine lyase/ transaminase [Arabidopsis thaliana]	HS			
887	15207	1.00E -44	99	gb AAG51737.1 AC068667_16beta-1,3 glucanase, putative; 26636-27432 [Arabidopsis thaliana]	SP			
888	14915	0	97	ref NP_199393.1 CIPK19 (CIPK19); kinase [Arabidopsis thaliana]	LN			
889	15132	1.00E -152	80	ref NP_187881.1 ZIP1 (ZINC TRANSPORTER 1 PRECURSOR); zinc ion transporter [Arabidopsis thaliana]	LN			
890	16229	0	100	gb ABG54330.1 double HA-tagged mitogen activated protein kinase 3 [synthetic construct]	SS			
891	14932	0	100	ref NP_568748.1 nucleotide-binding family protein [Arabidopsis thaliana]	DS			
892	76017	1.00E -180	100	ref NP_568235.1 unknown protein [Arabidopsis thaliana]	HS	PP		
893	15964	1.00E -44	100	ref NP_566894.1 unknown protein [Arabidopsis thaliana]	DS			
894	17227	8.00E -83	73	gb AAD03429.1 similar to nascent polypeptide associated complex alpha chain [Arabidopsis thaliana]	CS			
895	72952	0	96	ref NP_564480.1 unknown protein [Arabidopsis thaliana]	CS			
896	73706	0	96	ref NP_568221.1 protein kinase, putative [Arabidopsis thaliana]	HS	SS		
897	18203	0	92	ref NP_568919.1 APRR3 (PSEUDO-RESPONSE REGULATOR 3); transcription regulator [Arabidopsis thaliana]	HS			
898	16309	1.00E -163	88	ref NP_195558.1 GGR (GERANYLGERANYL REDUCTASE); farnesyltransferase [Arabidopsis thaliana]	LN			
899	15986	3.00E -99	84	dbj BAF02216.1 hypothetical protein [Arabidopsis thaliana]	DS			
900	15987	2.00E -54	79	ref NP_174170.1 glutaredoxin family protein [Arabidopsis thaliana]	DS			
901	16204	2.00E -59	100	ref NP_566191.1 NADH-ubiquinone oxidoreductase-related [Arabidopsis thaliana]	LN			
902	11133	1.00E -114	100	ref NP_563897.1 alanine racemase family protein [Arabidopsis thaliana]	LN			
903	73977	9.00E -67	100	ref NP_194677.1 mitochondrial ATP synthase g subunit family protein [Arabidopsis thaliana]	HS			
904	15995	1.00E -176	84	ref NP_564660.1 unknown protein [Arabidopsis thaliana]	DS			
904	18201	1.00E -176	84	ref NP_564660.1 unknown protein [Arabidopsis thaliana]	PP			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
905	16887	1.00E -164	100	ref NP_171820.1 phenazine biosynthesis PhzC/PhzF family protein [Arabidopsis thaliana]	PP			
906	16213	1.00E -140	82	ref NP_563991.1 unknown protein [Arabidopsis thaliana]	HS			
907	15507	0	96	ref NP_172255.1 cupin family protein [Arabidopsis thaliana]	LN			
908	15508	3.00E -86	100	ref NP_564937.1 unknown protein [Arabidopsis thaliana]	LN			
909	15959	3.00E -56	74	ref NP_180097.1 unknown protein [Arabidopsis thaliana]	LN			
910	72427	0	93	ref NP_568346.1 tubulin family protein [Arabidopsis thaliana]	HS			
911	19041	2.00E -85	82	ref NP_565723.1 unknown protein [Arabidopsis thaliana]	HS	LN		
912	17802	1.00E -118	100	ref NP_187425.1 emp24/gp25L/p24 family protein [Arabidopsis thaliana]	HS			
913	19042	0	94	ref NP_569048.1 ARA12; subtilase [Arabidopsis thaliana]	HS			
914	76053	0	94	ref NP_172544.1 unknown protein [Arabidopsis thaliana]	HS			
915	16120	1.00E -56	90	ref NP_192997.1 ribosomal protein L7Ae/L30e/S12e/Gadd45 family protein [Arabidopsis thaliana]	CK			
916	15424	1.00E -103	94	ref NP_196647.1 CBS domain-containing protein [Arabidopsis thaliana]	HS			
917	16401	0	90	ref NP_179995.1 CYP71B6 (CYTOCHROME P450 71B6); oxygen binding [Arabidopsis thaliana]	PP			
918	72417	0	100	gb AAK25868.1 AF360158_1 unknown protein [Arabidopsis thaliana]	CK			
919	73702	0	90	ref NP_564657.1 spliceosome protein-related [Arabidopsis thaliana]	HS	SS		
920	70706	0	100	gb AAD17366.1 similar to human phosphotyrosyl phosphatase activator PTPA (GB:X73478) [Arabidopsis thaliana]	HS			
921	71145	4.00E -57	86	gb AAG40386.1 AF325034_1AT5g02160 [Arabidopsis thaliana]	HS			
922	11609	0	97	gb AAC34228.1 putative cytochrome P450 [Arabidopsis thaliana]	LN			
923	71706	0	88	gb AAF03464.1 AC009327_3 hypothetical protein [Arabidopsis thaliana]	CK			
924	17325	1.00E -144	86	ref NP_563896.1 unknown protein [Arabidopsis thaliana]	PP			
925	70102	1.00E -166	100	ref NP_353899.2 hypothetical protein Atu0877 [Agrobacterium tumefaciens str. C58]	HS			
926	70106	1.00E -151	92	ref NP_353321.1 hypothetical protein Atu0291 [Agrobacterium tumefaciens str. C58]	HS			
927	70113	3.00E -94	94	ref NP_013690.1 Adenine phosphoribosyltransferase, catalyzes the formation of AMP from adenine and 5-phosphoribosylpyrophosphate; involved in the salvage pathway of purine nucleotide biosynthesis; Apt1p [Saccharomyces cerevisiae]	HS			
928	70115	0	100	ref NP_116623.1 Alanine:glyoxylate aminotransferase (AGT), catalyzes the synthesis of glycine from glyoxylate, which is one of three pathways for glycine biosynthesis in yeast; has similarity to mammalian and plant alanine:glyoxylate aminotransferases; Agx1p [Saccharomyces cerevisiae]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
929	70118	0	94	ref NP_012111.1 Mitochondrial glycerol-3-phosphate dehydrogenase; expression is repressed by both glucose and cAMP and derepressed by non-fermentable carbon sources in a Snf1p, Rsf1p, Hap2/3/4/5 complex dependent manner; Gut2p [<i>Saccharomyces cerevisiae</i>]	SP			
930	70128	9.00E -85	70	ref NP_638658.1 membrane-bound proton-translocating pyrophosphatase [<i>Xanthomonas campestris</i> pv. <i>campestris</i> str. ATCC 33913]	HS			
931	70130	1.00E -161	99	ref NP_639330.1 a-type carbonic anhydrase [<i>Xanthomonas campestris</i> pv. <i>campestris</i> str. ATCC 33913]	HS	PEG		
932	17103	0	88	ref NP_173155.1 CAT8 (CATIONIC AMINO ACID TRANSPORTER 8); cationic amino acid transporter [<i>Arabidopsis thaliana</i>]	SS			
933	72748	1.00E -33	74	ref NP_568635.1 unknown protein [<i>Arabidopsis thaliana</i>]	HS			
934	17502	3.00E -94	74	ref NP_565588.1 unknown protein [<i>Arabidopsis thaliana</i>]	HS			
935	17503	1.00E -130	100	ref NP_176178.1 ATGSTU16 (<i>Arabidopsis thaliana</i> Glutathione S-transferase (class tau) 16); glutathione transferase	HS			
936	78708	1.00E -168	95	gb AAK43902.1 AF370583_1Unknown protein [<i>Arabidopsis thaliana</i>]	LL			
937	17504	1.00E -139	95	ref NP_565634.1 harpin-induced protein-related / HIN1-related / harpin-responsive protein-related [<i>Arabidopsis thaliana</i>]	HS	LL	PP	
938	17132	1.00E -127	100	ref NP_187978.1 seven in absentia (SINA) family protein [<i>Arabidopsis thaliana</i>]	LN			
939	72965	0	91	gb AAN60250.1 unknown [<i>Arabidopsis thaliana</i>]	LL			
940	72796	3.00E -61	100	ref NP_201508.1 RALFL34 (RALF-LIKE 34) [<i>Arabidopsis thaliana</i>]	CS			
941	16617	5.00E -96	100	ref NP_355193.2 adenine phosphoribosyltransferase [<i>Agrobacterium tumefaciens</i> str. C58]	CS	HS		
942	16640	0	100	ref NP_355113.2 alcohol dehydrogenase [<i>Agrobacterium tumefaciens</i> str. C58]	HS			
943	16603	1.00E -132	85	ref NP_357530.1 ribokinase [<i>Agrobacterium tumefaciens</i> str. C58]	HS			
944	16612	1.00E -137	100	ref NP_010335.1 Triose phosphate isomerase, abundant glycolytic enzyme; mRNA half-life is regulated by iron availability; transcription is controlled by activators Reb1p, Gcr1p, and Rap1p through binding sites in the 5' non-coding region; Tpi1p [<i>Saccharomyces cerevisiae</i>]	LN	SP		
945	16614	1.00E -109	100	ref NP_010741.1 Stress inducible cytoplasmic thioredoxin peroxidase; cooperates with Tsa1p in the removal of reactive oxygen, nitrogen and sulfur species using thioredoxin as hydrogen donor; deletion enhances the mutator phenotype of tsa1 mutants; Tsa2p [<i>Saccharomyces cerevisiae</i>]	DS			
946	78354	7.00E -80	100	ref NP_192392.1 methionine sulfoxide reductase domain-containing protein / SelR domain-containing protein [<i>Arabidopsis thaliana</i>]	SS			
947	17818	6.00E -19	60	ref NP_564780.1 TIM13 (TIM13); protein translocase [<i>Arabidopsis thaliana</i>]	PP			
948	78370	3.00E -64	91	ref NP_194900.1 TAFII15 (SALT TOLERANCE DURING GERMINATION 1); transcription factor [<i>Arabidopsis thaliana</i>]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
949	70411	0	90	ref NP_849806.1 UBP1A; mRNA 3'-UTR binding [Arabidopsis thaliana]	SP			
950	19154	1.00E -151	89	ref NP_197310.1 transcriptional factor B3 family protein [Arabidopsis thaliana]	HS	LN	PP	
951	19155	0	82	ref NP_200906.2 nuclear transport factor 2 (NTF2) family protein / RNA recognition motif (RRM)-containing protein [Arabidopsis thaliana]	HS			
952	18307	0	94	ref NP_194507.1 ACBP2 (ACYL-COA BINDING PROTEIN ACBP 2) [Arabidopsis thaliana]	HS			
953	72030	0	86	ref NP_177367.1 zinc finger (C3HC4-type RING finger) family protein [Arabidopsis thaliana]	LN			
954	17916	1.00E -173	95	ref NP_568839.1 CSN6A (COP9 signalosome subunit 6A) [Arabidopsis thaliana]	PP			
955	17917	0	96	ref NP_188478.1 aspartyl protease family protein [Arabidopsis thaliana]	SS			
956	17334	0	94	ref NP_178146.1 chloroplast ADP, ATP carrier protein 1 / ADP, ATP translocase 1 / adenine nucleotide translocase 1 (AATP1) [Arabidopsis thaliana]	PP			
957	71531	0	100	ref NP_565656.1 aldo/keto reductase family protein [Arabidopsis thaliana]	HS			
958	18441	7.00E -75	100	ref NP_973779.1 lactoylglutathione lyase family protein / glyoxalase I family protein [Arabidopsis thaliana]	HS			
959	19162	2.00E -73	85	ref NP_563687.1 photosystem II family protein [Arabidopsis thaliana]	DS			
960	73205	0	98	ref NP_190414.1 zinc finger (CCCH-type) family protein [Arabidopsis thaliana]	PEG			
961	18306	1.00E -172	100	ref NP_566769.1 LAG1 (Longevity assurance gene 1) [Arabidopsis thaliana]	SS			
962	70808	0	97	ref NP_196136.1 CESA3 (CELLULOSE SYNTHASE 3); cellulose synthase/ transferase, transferring glycosyl groups [Arabidopsis thaliana]	CS	CK		
963	17653	1.00E -178	95	ref NP_199155.1 zinc finger (C3HC4-type RING finger) family protein [Arabidopsis thaliana]	LN			
964	19535	0	81	ref NP_567002.1 GCN5 (Histon acetyltransferase HAT1) [Arabidopsis thaliana]	HS			
965	71719	0	99	ref NP_192044.1 DC1 domain-containing protein [Arabidopsis thaliana]	HS	LL	PP	
966	18546	1.00E -102	92	ref NP_176285.1 AGL56 (AGAMOUS LIKE-56); DNA binding / transcription factor [Arabidopsis thaliana]	PP			
967	19534	0	78	ref NP_566721.1 binding [Arabidopsis thaliana]	HS			
968	18421	0	96	gb ABK06451.1 flag-tagged protein kinase domain of putative mitogen-activated protein kinase kinase kinase [synthetic construct]	HS			
969	18422	0	100	ref NP_194498.1 MSP1 protein, putative / intramitochondrial sorting protein, putative [Arabidopsis thaliana]	HS			
970	19539	0	95	ref NP_191747.1 CYP78A9 (CYTOCHROME P450 78A9); oxygen binding [Arabidopsis thaliana]	HS	PP		
971	19244	3.00E -74	88	ref NP_565113.1 auxin-responsive family protein [Arabidopsis thaliana]	HS			
972	15803	1.00E -142	100	ref NP_199675.2 cyclin family protein [Arabidopsis thaliana]	HS			
973	19545	0	100	ref NP_568665.1 pentatricopeptide (PPR) repeat-containing protein [Arabidopsis thaliana]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
974	10222	0	94	ref NP_190896.1 CYP71B5 (CYTOCHROME P450 71B5); oxygen binding [Arabidopsis thaliana]	LN			
974	72356	0	94	ref NP_190896.1 CYP71B5 (CYTOCHROME P450 71B5); oxygen binding [Arabidopsis thaliana]	CK	PEG		
975	70425	0	98	ref NP_200201.3 unknown protein [Arabidopsis thaliana]	HS			
976	19237	1.00E -150	95	ref NP_199836.1 31 kDa ribonucleoprotein, chloroplast, putative / RNA-binding protein RNP-T, putative / RNA-binding protein 1/2/3, putative / RNA-binding protein cp31, putative [Arabidopsis thaliana]	HS			
977	71303	0	82	ref NP_199431.1 KH domain-containing protein [Arabidopsis thaliana]	HS			
978	19240	1.00E -180	100	ref NP_566576.1 meprin and TRAF homology domain-containing protein / MATH domain-containing protein [Arabidopsis thaliana]	LN			
979	18535	0	100	ref NP_196179.1 oxidoreductase, 2OG-Fe(II) oxygenase family protein [Arabidopsis thaliana]	HS			
980	19450	0	94	ref NP_565422.1 pentatricopeptide (PPR) repeat-containing protein [Arabidopsis thaliana]	LN			
981	18221	1.00E -66	100	ref NP_193992.1 protease inhibitor/seed storage/lipid transfer protein (LTP) family protein [Arabidopsis thaliana]	PP			
982	18233	1.00E -110	87	ref NP_193179.1 enoyl-CoA hydratase/isomerase family protein [Arabidopsis thaliana]	PP			
983	18236	1.00E -31	70	ref NP_565410.1 unknown protein [Arabidopsis thaliana]	PP			
984	18238	1.00E -129	88	ref NP_197105.1 3-oxo-5-alpha-steroid 4-dehydrogenase family protein / steroid 5-alpha-reductase family protein [Arabidopsis thaliana]	HS			
985	18309	1.00E -128	93	gb AAM61596.1 DNA-binding protein [Arabidopsis thaliana]	CS	PEG		
986	70337	0	85	gb AAP84710.2 metacaspase 7 [Arabidopsis thaliana]	HS			
987	18319	1.00E -105	89	ref NP_172721.1 DDF1 (DWARF AND DELAYED FLOWERING 1); DNA binding / transcription factor [Arabidopsis thaliana]	LN			
988	70338	1.00E -172	85	ref NP_200445.1 zinc finger (C3HC4-type RING finger) family protein [Arabidopsis thaliana]	LL			
989	10471	1.00E -114	84	dbj BAF00210.1 hypothetical protein [Arabidopsis thaliana]	DS			
990	18331	1.00E -114	63	ref NP_565124.1 nucleic acid binding [Arabidopsis thaliana]	CS			
991	10228	0	97	ref NP_563880.1 TIF3H1 (eukaryotic translation initiation factor 3 subunit H1); translation initiation factor [Arabidopsis thaliana]	LN			
991	10473	0	97	ref NP_563880.1 TIF3H1 (eukaryotic translation initiation factor 3 subunit H1); translation initiation factor [Arabidopsis thaliana]	LN			
992	18261	1.00E -167	91	ref NP_563732.1 peroxidase, putative [Arabidopsis thaliana]	PP			
993	18848	0	94	ref NP_177805.1 O-methyltransferase family 2 protein [Arabidopsis thaliana]	LN			
994	18268	1.00E -130	92	gb AAM62813.1 unknown [Arabidopsis thaliana]	PP			
995	18345	0	92	ref NP_182320.1 SWA1 (SLOW WALKER1); nucleotide binding [Arabidopsis thaliana]	HS			
996	73607	0	92	ref NP_851068.1 WD-40 repeat family protein [Arabidopsis thaliana]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
997	18352	1.00E -166	83	ref NP_201275.1 nodulin MtN21 family protein [Arabidopsis thaliana]	HS	SS	PEG	
998	18357	1.00E -171	100	ref NP_172866.1 mitochondrial substrate carrier family protein [Arabidopsis thaliana]	PP			
999	19423	0	100	ref NP_177020.1 CUT1 (CUTICULAR 1); acyltransferase [Arabidopsis thaliana]	DS			
1000	78373	0	94	ref NP_179537.1 CKL5 (Casein Kinase I-like 5); casein kinase I/kinase [Arabidopsis thaliana]	CK	HS		
1001	70603	0	100	ref NP_175081.1 ALDH3H1 (ALDEHYDE DEHYDROGENASE 4); 3-chloroallyl aldehyde dehydrogenase/ aldehyde dehydrogenase (NAD) [Arabidopsis thaliana]	HS	LN		
1002	19616	1.00E -159	100	ref NP_563789.1 SMO2-2 (sterol 4-alpha-methyl-oxidase 2); C-4 methylsterol oxidase [Arabidopsis thaliana]	HS			
1003	19425	1.00E -164	91	ref NP_564407.1 electron carrier/ iron ion binding [Arabidopsis thaliana]	DS			
1004	19426	1.00E -152	89	ref NP_564428.1 unknown protein [Arabidopsis thaliana]	PP			
1005	18382	2.00E -51	87	ref NP_564651.1 unknown protein [Arabidopsis thaliana]	PP			
1006	71536	4.00E -86	86	ref NP_564747.1 ZCF37 [Arabidopsis thaliana]	HS			
1007	70419	6.00E -88	66	ref NP_564888.1 unknown protein [Arabidopsis thaliana]	LL	CK	DS	
1008	70827	6.00E -85	100	ref NP_189667.1 zinc finger (C3HC4-type RING finger) family protein [Arabidopsis thaliana]	HS			
1009	19311	0	94	ref NP_010849.1 Ferrioxamine B transporter, member of the ARN family of transporters that specifically recognize siderophore-iron chelates; transcription is induced during iron deprivation and diauxic shift; potentially phosphorylated by Cdc28p; Sit1p [Saccharomyces cerevisiae]	LN			
1010	19323	0	92	ref NP_354642.2 malic enzyme [Agrobacterium tumefaciens str. C58]	LL			
1011	19320	0	100	ref NP_012542.1 Glyceraldehyde-3-phosphate dehydrogenase, isozyme 2, involved in glycolysis and gluconeogenesis; tetramer that catalyzes the reaction of glyceraldehyde-3-phosphate to 1,3 bis-phosphoglycerate; detected in the cytoplasm and cell-wall; Tdh2p [Saccharomyces cerevisiae]	HS			
1012	73613	0	95	ref NP_197104.1 NIK1 (NSP-INTERACTING KINASE 1); kinase [Arabidopsis thaliana]	CK			
1013	70432	1.00E -162	91	ref NP_196709.1 GTP binding [Arabidopsis thaliana]	SS			
1014	70433	0	100	ref NP_196957.1 transducin family protein / WD-40 repeat family protein [Arabidopsis thaliana]	HS			
1015	70434	1.00E -142	94	ref NP_180497.1 tropinone reductase, putative / tropine dehydrogenase, putative [Arabidopsis thaliana]	HS			
1016	72614	1.00E -128	79	ref NP_197757.1 sterile alpha motif (SAM) domain-containing protein [Arabidopsis thaliana]	LL			
1017	70436	7.00E -35	70	ref NP_197966.1 early nodulin-related [Arabidopsis thaliana]	CS	HS		
1018	74226	0	92	dbj BAB02016.1 MAP kinase [Arabidopsis thaliana]	PP			
1019	70439	1.00E -123	91	ref NP_191247.1 VQ motif-containing protein [Arabidopsis thaliana]	HS	PEG		

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1020	70446	1.00E -128	100	ref NP_568972.2 unknown protein [Arabidopsis thaliana]	CS			
1021	70613	0	93	ref NP_174145.1 unknown protein [Arabidopsis thaliana]	PEG			
1022	70447	0	97	ref NP_175977.1 uracil phosphoribosyltransferase, putative / UMP pyrophosphorylase, putative / UPRTase, putative [Arabidopsis thaliana] ref NP_974037.1 uracil phosphoribosyltransferase, putative / UMP pyrophosphorylase, putative / UPRTase, putative [Arabidopsis thaliana]	CK			
1023	71725	0	67	gb ABE65946.1 nucleolin [Arabidopsis thaliana]	CK			
1024	70462	9.00E -73	92	ref NP_567037.1 ARR17 (response regulator 17); transcription regulator/ two-component response regulator [Arabidopsis thaliana]	PEG			
1025	70545	0	95	ref NP_176108.3 bZIP family transcription factor [Arabidopsis thaliana]	SP			
1026	70741	0	100	gb AAM78097.1 AT4g18060/F15J5_30 [Arabidopsis thaliana]	HS			
1027	19702	0	97	ref NP_011904.1 Protein of unknown function, green fluorescent protein (GFP)-fusion protein localizes to the endoplasmic reticulum; msc7 mutants are defective in directing meiotic recombination events to homologous chromatids; Msc7p [Saccharomyces cerevisiae]	HS			
1028	19985	1.00E -140	86	gb AAA88792.1 nucleosome assembly protein 1	HS			
1029	19775	1.00E -80	54	ref NP_175779.1 unknown protein [Arabidopsis thaliana]	CK	HS	PP	LN
1030	19829	3.00E -56	42	ref NP_565892.1 unknown protein [Arabidopsis thaliana]	CS			
1031	19982	1.00E -123	79	gb AAV49506.1 L-galactose-1-phosphate phosphatase [Actinidia deliciosa]	HS	PEG		
1032	19755	1.00E -110	63	emb CAO23381.1 unnamed protein product [Vitis vinifera]	HS			
1033	19732	0	96	dbj BAA22559.1 squalene synthase [Glycine max]	PP			
1034	70365	0	91	gb AAQ84169.1 1-deoxy-D-xylulose 5-phosphate synthase [Pueraria montana var. lobata]	CS			
1035	10476	1.00E -168	86	ref NP_194096.1 CDPK6 (CALCIUM-DEPENDENT PROTEIN KINASE 6); anion channel/ calcium- and calmodulin-dependent protein kinase/ kinase [Arabidopsis thaliana]	LN			
1036	19987	0	83	emb CAO67062.1 unnamed protein product [Vitis vinifera]	HS			
1037	19893	1.00E -166	85	gb ABC75353.2 Intracellular chloride channel [Medicago truncatula]	HS			
1038	19983	6.00E -92	60	emb CAO39739.1 unnamed protein product [Vitis vinifera]	HS	LL		
1039	19825	1.00E -40	39	emb CAO18125.1 unnamed protein product [Vitis vinifera]	HS			
1040	19919	1.00E -121	77	emb CAO61472.1 unnamed protein product [Vitis vinifera]	HS			
1041	19979	0	67	emb CAO68238.1 unnamed protein product [Vitis vinifera]	HS			
1042	19786	2.00E -80	75	emb CAO48404.1 unnamed protein product [Vitis vinifera]	HS	LN		
1043	70903	0	76	emb CAO66035.1 unnamed protein product [Vitis vinifera]	HS	PP	PEG	

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1044	70970	1.00E -102	67	ref NP_200538.1 unknown protein [Arabidopsis thaliana]	HS			
1045	70980	0	89	dbj BAE71243.1 putative galactose kinase [Trifolium pratense]	CK			
1046	70936	1.00E -158	49	emb CAO71465.1 unnamed protein product [Vitis vinifera]	HS			
1047	71425	1.00E -160	74	emb CAO14679.1 unnamed protein product [Vitis vinifera]	DS			
1048	78666	3.00E -45	83	ref NP_192698.1 GASA3 (GAST1 PROTEIN HOMOLOG 3) [Arabidopsis thaliana]	LN			
1049	71202	1.00E -163	95	ref NP_194179.1 serine/threonine protein kinase, putative [Arabidopsis thaliana]	LN			
1050	78973	0	90	ref NP_565672.1 DRB2 (DSRNA-BINDING PROTEIN 2); double-stranded RNA binding [Arabidopsis thaliana]	LL			
1051	76411	1.00E -179	99	ref NP_172864.1 2-oxoglutarate-dependent dioxygenase, putative [Arabidopsis thaliana]	PP			
1052	70635	4.00E -64	100	ref NP_851278.1 senescence-associated family protein [Arabidopsis thaliana]	HS			
1053	73715	0	92	ref NP_198428.1 G6PD1 (GLUCOSE-6-PHOSPHATE DEHYDROGENASE 1); glucose-6-phosphate 1-dehydrogenase [Arabidopsis thaliana]	PEG			
1054	70638	6.00E -79	89	ref NP_197398.1 AWPM-19-like membrane family protein [Arabidopsis thaliana]	HS	LN		
1055	70642	1.00E -176	92	ref NP_191315.1 aspartate/glutamate/uridylate kinase family protein [Arabidopsis thaliana]	HS			
1056	70643	0	96	ref NP_191703.1 ATCYSC1 (BETA-SUBSTITUTED ALA SYNTHASE 3;1); L-3-cyanoalanine synthase/ cysteine synthase [Arabidopsis thaliana]	HS			
1057	71564	1.00E -121	95	emb CAA47753.1 proteosome subunit [Arabidopsis thaliana]	HS	SP		
1058	75039	1.00E -175	90	ref NP_188659.1 mitochondrial substrate carrier family protein [Arabidopsis thaliana]	LN			
1059	70653	8.00E -82	100	ref NP_566365.1 unknown protein [Arabidopsis thaliana]	PEG			
1060	70655	4.00E -87	82	ref NP_566619.1 unknown protein [Arabidopsis thaliana]	LL	PEG		
1061	78321	0	97	ref NP_181404.1 MVD1 (mevalonate diphosphate decarboxylase 1) [Arabidopsis thaliana]	DS	HS		
1062	78307	0	100	ref NP_182079.1 CYP76C4 (cytochrome P450, family 76, subfamily C, polypeptide 4); oxygen binding [Arabidopsis thaliana]	CK			
1063	71808	0	91	ref NP_182082.2 CYP76C3 (cytochrome P450, family 76, subfamily C, polypeptide 3); oxygen binding [Arabidopsis thaliana]	HS			
1064	71810	0	97	gb AAF88087.1 AC025417_15T12C24.27 [Arabidopsis thaliana]	LN			
1065	71313	0	100	ref NP_187667.1 CYP77A7 (cytochrome P450, family 77, subfamily A, polypeptide 7, unfertilized embryo sac 9); oxygen binding [Arabidopsis thaliana]	CS			
1066	72349	1.00E -134	100	ref NP_191072.1 TT5 (TRANSPARENT TESTA 5); chalcone isomerase [Arabidopsis thaliana]	SP			
1067	71318	0	90	ref NP_189261.1 CYP71B34 (cytochrome P450, family 71, subfamily B, polypeptide 34); oxygen binding [Arabidopsis thaliana]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1068	71812	0	92	ref NP_195705.1 CYP79B2 (cytochrome P450, family 79, subfamily B, polypeptide 2); oxygen binding [Arabidopsis thaliana]	LN			
1069	74060	0	90	ref NP_188731.1 CYP705A32 (cytochrome P450, family 705, subfamily A, polypeptide 32); oxygen binding [Arabidopsis thaliana]	DS			
1070	71816	0	93	ref NP_568025.1 CYP81F3 (cytochrome P450, family 81, subfamily F, polypeptide 3); oxygen binding [Arabidopsis thaliana]	LL			
1071	71569	1.00E -142	95	ref NP_189364.1 ATPHB4 (PROHIBITIN 4) [Arabidopsis thaliana]	CS			
1072	70672	0	100	ref NP_567641.1 PRXR1 (peroxidase 42); peroxidase [Arabidopsis thaliana]	PP			
1073	71336	0	96	ref NP_563762.1 VHS domain-containing protein / GAT domain-containing protein [Arabidopsis thaliana]	DS			
1074	71337	1.00E -179	100	ref NP_179721.1 mannose 6-phosphate reductase (NADPH-dependent), putative [Arabidopsis thaliana]	HS			
1075	71339	0	93	ref NP_200747.1 XH/XS domain-containing protein [Arabidopsis thaliana]	CS			
1076	71825	0	87	ref NP_195998.2 TRFL10 (TRF-LIKE 10); DNA binding [Arabidopsis thaliana]	LN			
1077	73696	1.00E -166	86	ref NP_192864.1 MADS-box family protein [Arabidopsis thaliana]	SP			
1078	70682	1.00E -129	94	ref NP_001078595.1 RNA recognition motif (RRM)-containing protein [Arabidopsis thaliana]	HS			
1079	70686	1.00E -117	75	ref NP_199096.1 ATU2AF35B; RNA binding [Arabidopsis thaliana]	SP	LL		
1080	78316	0	100	ref NP_192085.1 DC1 domain-containing protein [Arabidopsis thaliana]	SS			
1081	71667	0	100	ref NP_013976.1 Glutamate decarboxylase, converts glutamate into gamma-aminobutyric acid (GABA) during glutamate catabolism; involved in response to oxidative stress; Gad1p [Saccharomyces cerevisiae]	DS			
1082	71695	0	77	emb CAA75577.1 L-ascorbate oxidase [Medicago truncatula]	CK			
1083	71696	0	96	gb AAB03258.1 phosphoinositide-specific phospholipase C P13	HS	LN		
1084	72475	1.00E -111	83	gb AAF21310.1 seed maturation protein PM24 [Glycine max]	HS			
1085	71677	1.00E -125	77	gb ABD28522.1 Cupin, RmIC-type [Medicago truncatula]	CK			
1086	71691	7.00E -34	35	emb CAO44124.1 unnamed protein product [Vitis vinifera]	CS			
1087	72476	1.00E -122	70	emb CAO47395.1 unnamed protein product [Vitis vinifera]	HS			
1088	70803	0	97	ref NP_850874.1 glutamate-tRNA ligase, putative / glutamyl-tRNA synthetase, putative / GluRS, putative [Arabidopsis thaliana]	SP	LN		
1089	71612	0	94	ref NP_192179.1 SULTR3;2 (SULFATE TRANSPORTER 3;2); sulfate transporter [Arabidopsis thaliana]	LL			
1090	71637	0	88	gb AAD52696.1 AF087819_1 auxin transport protein [Arabidopsis thaliana]	LN			
1091	71629	0	97	ref NP_565924.1 unknown protein [Arabidopsis thaliana]	HS	PEG		

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1092	71622	0	97	ref NP_566085.1 APRR9 (PSEUDO-RESPONSE REGULATOR 9); transcription regulator [Arabidopsis thaliana]	LN			
1093	70810	0	91	ref NP_175610.1 mitochondrial processing peptidase alpha subunit, putative [Arabidopsis thaliana]	HS	SS		
1094	72452	6.00E-76	77	ref NP_199889.1 auxin-responsive family protein [Arabidopsis thaliana]	LN			
1095	74071	0	93	ref NP_174516.1 ribose-phosphate pyrophosphokinase 2 / phosphoribosyl diphosphate synthetase 2 (PRS2) [Arabidopsis thaliana]	LL			
1096	11042	2.00E-89	93	ref NP_179709.1 peptidyl-prolyl cis-trans isomerase / cyclophilin (CYP2) / rotamase [Arabidopsis thaliana]	LN			
1097	72510	0	89	ref NP_196865.1 unknown protein [Arabidopsis thaliana]	CK			
1098	72537	1.00E-124	82	ref NP_851118.1 unknown protein [Arabidopsis thaliana]	HS	LN		
1099	72541	9.00E-57	88	ref NP_568684.1 unknown protein [Arabidopsis thaliana]	PP			
1100	72617	4.00E-88	100	ref NP_568761.1 unknown protein [Arabidopsis thaliana]	LL			
1101	73755	1.00E-132	100	ref NP_200501.1 cysteine protease inhibitor [Arabidopsis thaliana]	PP	SS		
1102	72630	2.00E-86	93	ref NP_201267.1 C/VIF2 (CELL WALL / VACUOLAR INHIBITOR OF FRUCTOSIDASE 2); pectinesterase inhibitor [Arabidopsis thaliana]	SP			
1103	72645	5.00E-37	75	ref NP_180615.1 photosystem II reaction center W (PsbW) protein-related [Arabidopsis thaliana]	LN			
1104	72647	5.00E-43	100	ref NP_566036.1 protease inhibitor/seed storage/lipid transfer protein (LTP) family protein [Arabidopsis thaliana]	LN			
1105	14316	1.00E-55	86	ref NP_188703.1 histone H2A, putative [Arabidopsis thaliana]	DS			
1106	73689	6.00E-79	100	ref NP_173037.1 transducin family protein / WD-40 repeat family protein [Arabidopsis thaliana]	DS	SS		
1107	73345	0	92	ref NP_174226.1 transducin family protein / WD-40 repeat family protein [Arabidopsis thaliana]	HS	LN		
1108	73347	0	100	ref NP_177513.2 transducin family protein / WD-40 repeat family protein [Arabidopsis thaliana]	SP			
1109	73231	0	84	ref NP_196332.1 leucine-rich repeat family protein [Arabidopsis thaliana]	PP			
1110	78325	0	84	ref NP_171637.1 aspartyl protease family protein [Arabidopsis thaliana]	HS			
1111	72658	1.00E-138	93	ref NP_172224.1 tropinone reductase, putative / tropine dehydrogenase, putative [Arabidopsis thaliana]	PEG			
1112	72660	3.00E-78	82	ref NP_564585.1 zinc finger (AN1-like) family protein [Arabidopsis thaliana]	LN			
1113	72662	5.00E-49	100	ref NP_179096.1 gibberellin-regulated family protein [Arabidopsis thaliana]	LN			
1114	73352	0	97	ref NP_566304.1 armadillo/beta-catenin repeat family protein / U-box domain-containing protein [Arabidopsis thaliana]	LN			
1115	72814	0	100	ref NP_566558.1 F-box family protein [Arabidopsis thaliana]	PP	CS	PEG	
1116	72816	0	100	ref NP_566824.1 CARA (CARBAMOYL PHOSPHATE SYNTHETASE A); carbamoyl-phosphate synthase (glutamine-hydrolyzing) [Arabidopsis thaliana]	PP			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1117	72820	0	89	ref NP_194653.1 leucine-rich repeat family protein / extensin family protein [Arabidopsis thaliana]	LL			
1118	73353	0	97	ref NP_194713.1 MTO2 (METHIONINE OVER- ACCUMULATOR); threonine synthase [Arabidopsis thaliana]	HS	PEG		
1119	75066	1.00E -134	91	ref NP_175644.1 ABA2 (ABA DEFICIENT 2); oxidoreductase [Arabidopsis thaliana]	PEG			
1120	78337	2.00E -40	78	dbj BAD43663.1 unknown protein [Arabidopsis thaliana]	PEG			
1121	73317	0	96	ref NP_200121.4 oxidoreductase [Arabidopsis thaliana]	LN			
1122	73318	0	94	ref NP_568145.1 short-chain dehydrogenase/reductase (SDR) family protein [Arabidopsis thaliana]	CS			
1123	73749	0	97	gb AAG51266.1 AC027135_7protein kinase, putative [Arabidopsis thaliana]	CS			
1124	18704	0	88	ref NP_201234.1 DCT/DIT2.1 (DICARBOXYLATE TRANSPORT); oxoglutarate:malate antiporter [Arabidopsis thaliana]	PEG			
1125	73221	1.00E -162	82	ref NP_563785.1 C2 domain-containing protein [Arabidopsis thaliana]	CK	LL		
1126	71125	1.00E -117	94	ref NP_172671.1 Rho GDP-dissociation inhibitor family protein [Arabidopsis thaliana]	HS			
1127	76203	0	97	ref NP_200678.2 phosphoinositide-specific phospholipase C family protein [Arabidopsis thaliana]	PEG			
1128	75067	0	100	ref NP_566565.1 peroxidase, putative [Arabidopsis thaliana]	PEG			
1129	73729	0	100	ref NP_194944.1 CYP96A2 (cytochrome P450, family 96, subfamily A, polypeptide 2); oxygen binding [Arabidopsis thaliana]	LL			
1130	75206	0	95	ref NP_197962.1 GA3 (GA REQUIRING 3); oxygen binding [Arabidopsis thaliana]	HS			
1131	73248	0	100	ref NP_200532.1 CYP81F2 (cytochrome P450, family 81, subfamily F, polypeptide 2); oxygen binding [Arabidopsis thaliana]	HS			
1132	73738	0	93	ref NP_198641.1 serine/threonine protein kinase, putative [Arabidopsis thaliana]	DS			
1133	73332	0	97	ref NP_173685.1 SUC2 (SUCROSE-PROTON SYMPORTER 2); carbohydrate transporter/ sucrose:hydrogen symporter/ sugar porter [Arabidopsis thaliana]	PP			
1134	78335	0	100	ref NP_189582.1 UDP-glucose 6-dehydrogenase, putative [Arabidopsis thaliana]	HS	LL	PP	
1135	75821	6.00E -76	90	ref NP_190424.1 unknown protein [Arabidopsis thaliana]	LL			
1136	72038	0	97	emb CAA18501.1 Calcium-dependent serine/threonine protein kinase [Arabidopsis thaliana]	HS			
1137	72027	0	91	ref NP_849857.1 copine-related [Arabidopsis thaliana]	DS			
1138	72016	0	97	ref NP_200887.1 ATGCN1 (Arabidopsis thaliana general control non-repressible 1)	CS			
1139	70835	1.00E -128	82	ref NP_181461.1 PGPS1 (PHOSPHATIDYLGLYCEROLPHOSPHATE+SYNTHASE +1); CDP-alcohol phosphatidyltransferase [Arabidopsis thaliana]	SS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1140	72044	0	99	ref NP_014446.1 Protein proposed to interact with phospholipid translocases, shares similarity to Cdc50p; Ynr048wp [Saccharomyces cerevisiae]	CS			
1141	72092	0	92	ref NP_010796.1 High-affinity glutamine permease, also transports Leu, Ser, Thr, Cys, Met and Asn; expression is fully dependent on Grr1p and modulated by the Ssy1p-Ptr3p-Ssy5p (SPS) sensor of extracellular amino acids; Gnp1p [Saccharomyces cerevisiae]	CK			
1142	72124	1.00E-173	55	emb CAO64952.1 unnamed protein product [Vitis vinifera]	HS			
1143	72110	1.00E-172	63	emb CAN75218.1 hypothetical protein [Vitis vinifera]	HS			
1144	73916	1.00E-162	57	ref NP_569047.1 SKIP2 (SKP1 INTERACTING PARTNER 2); ubiquitin-protein ligase [Arabidopsis thaliana]	HS			
1145	74268	0	100	ref NP_186851.1 aspartate kinase, lysine-sensitive, putative [Arabidopsis thaliana]	HS	LL		
1146	74269	0	94	ref NP_566764.1 bile acid:sodium symporter family protein [Arabidopsis thaliana]	DS			
1147	77902	0	96	ref NP_197021.2 tyrosyl-DNA phosphodiesterase-related [Arabidopsis thaliana]	SS			
1148	74273	1.00E-180	100	ref NP_199285.1 molybdenum cofactor sulfurase family protein [Arabidopsis thaliana]	HS			
1149	74280	1.00E-165	94	emb CAA58893.1 cysteine synthase [Arabidopsis thaliana]	HS			
1150	74281	0	94	ref NP_197655.1 prephenate dehydratase family protein [Arabidopsis thaliana]	LL			
1151	74287	0	100	ref NP_974217.1 SUVR4 [Arabidopsis thaliana]	CK	HS		
1152	73767	1.00E-156	100	ref NP_194231.1 nodulin MtN3 family protein [Arabidopsis thaliana]	LN			
1153	77304	0	100	ref NP_175700.2 catalytic [Arabidopsis thaliana]	CK	HS	PEG	
1154	73761	0	95	ref NP_566683.1 mitochondrial substrate carrier family protein [Arabidopsis thaliana]	LL			
1155	74721	0	93	ref NP_194568.1 AAC3 (ADP/ATP CARRIER 3); ATP:ADP antiporter/ binding [Arabidopsis thaliana]	LL			
1156	74719	0	91	ref NP_198474.1 protein phosphatase 2C, putative / PP2C, putative [Arabidopsis thaliana]	CK			
1157	74257	0	98	ref NP_179113.2 fatty acid elongase, putative [Arabidopsis thaliana]	DS	HS		
1158	72749	0	95	ref NP_176767.1 regulator of chromosome condensation (RCC1) family protein / zinc finger protein-related [Arabidopsis thaliana]	HS	LL		
1159	72703	2.00E-77	92	ref NP_013963.1 Subunit (17 kDa) of TFIID and SAGA complexes, involved in RNA polymerase II transcription initiation and in chromatin modification, similar to histone H3; Taf9p [Saccharomyces cerevisiae]	PEG			
1160	72763	0	98	ref NP_009691.1 Protein arginine N-methyltransferase that exhibits septin and Hsl1p-dependent bud neck localization and periodic Hsl1p-dependent phosphorylation; required along with Hsl1p for bud neck recruitment, phosphorylation, and degradation of Swe1p; Hsl7p [Saccharomyces cerevisiae]	SS			
1161	72775	1.00E-172	100	ref NP_009718.1 Catalytic subunit of the main cell cycle cyclin-dependent kinase (CDK); alternately associates with G1 cyclins (CLNs) and G2/M cyclins (CLBs) which direct the CDK to specific substrates; Cdc28p [Saccharomyces cerevisiae]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1162	72728	0	96	ref NP_013106.1 Cytoplasmic response regulator, part of a two-component signal transducer that mediates osmosensing via a phosphorelay mechanism; dephosphorylated form is degraded by the ubiquitin-proteasome system; potential Cdc28p substrate; Ssk1p [Saccharomyces cerevisiae]	CS			
1163	72776	1.00E-101	92	gb EDN62754.1 oxidant-induced cell cycle arrest [Saccharomyces cerevisiae YJM789]	HS			
1164	72717	1.00E-131	100	ref NP_011075.1 TATA-binding protein, general transcription factor that interacts with other factors to form the preinitiation complex at promoters, essential for viability; Spt15p [Saccharomyces cerevisiae]	DS			
1165	72718	1.00E-159	90	ref NP_011400.1 Activating gamma subunit of the AMP-activated Snf1p kinase complex (contains Snf1p and a Sip1p/Sip2p/Gal83p family member); activates glucose-repressed genes, represses glucose-induced genes; role in sporulation, and peroxisome biogenesis; Snf4p [Saccharomyces cerevisiae]	DS			
1166	11715	0	93	ref NP_187616.1 OMR1 (L-O-METHYLTHREONINE RESISTANT 1); threonine ammonia-lyase [Arabidopsis thaliana]	LN			
1167	72779	0	93	ref NP_011622.1 B-type cyclin involved in cell cycle progression; activates Cdc28p to promote the transition from G2 to M phase; accumulates during G2 and M, then targeted via a destruction box motif for ubiquitin-mediated degradation by the proteasome; Clb1p [Saccharomyces cerevisiae]	DS			
1168	72791	0	100	ref NP_009709.1 Protein of unknown function, required for normal localization of actin patches and for normal tolerance of sodium ions and hydrogen peroxide; localizes to both cytoplasm and nucleus; Apd1p [Saccharomyces cerevisiae]	DS	HS		
1169	72732	0	100	ref NP_013662.1 Putative protein of unknown function; non-essential gene; null mutant displays increased frequency of mitochondrial genome loss (petite formation); Aim32p [Saccharomyces cerevisiae]	HS			
1170	73027	1.00E-125	93	ref NP_187445.1 Rho GDP-dissociation inhibitor family protein [Arabidopsis thaliana]	CS			
1171	73058	0	96	ref NP_175591.1 leucine-rich repeat protein kinase, putative [Arabidopsis thaliana]	CK			
1172	73126	0	95	ref NP_173076.1 protein kinase family protein [Arabidopsis thaliana]	PEG			
1173	73138	0	91	ref NP_356682.2 aldehyde dehydrogenase [Agrobacterium tumefaciens str. C58]	HS			
1174	73174	0	94	ref NP_241069.1 glycine betaine aldehyde dehydrogenase [Bacillus halodurans C-125]	HS			
1175	73186	0	99	ref NP_241405.1 NADP-dependent aldehyde dehydrogenase [Bacillus halodurans C-125]	HS	LN		
1176	11141	0	100	ref NP_568064.1 AGT2 (ALANINE:GLYOXYLATE AMINOTRANSFERASE 2); alanine-glyoxylate transaminase [Arabidopsis thaliana]	LN			
1177	73139	0	89	ref NP_242876.1 aldehyde dehydrogenase [Bacillus halodurans C-125]	CS			
1178	73128	0	97	ref NP_388272.1 4-aminobutyrate aminotransferase [Bacillus subtilis subsp. subtilis str. 168]	HS			
1179	73164	0	93	ref NP_389241.1 transaminase [Bacillus subtilis subsp. subtilis str. 168]	PEG			
1180	73117	0	97	ref NP_391762.1 aldehyde dehydrogenase [Bacillus subtilis subsp. subtilis str. 168]	CS	HS		

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1181	73142	0	99	ref YP_585121.1 glyceraldehyde-3-phosphate dehydrogenase, type I [Ralstonia metallidurans CH34]	PP	SS		
1182	73990	1.00E-154	90	ref NP_949547.1 glutaminase [Rhodopseudomonas palustris CGA009]	HS			
1183	73166	1.00E-117	100	ref NP_384268.1 phosphoglyceromutase [Sinorhizobium meliloti 1021]	SS			
1184	73178	0	91	ref NP_385430.1 PUTATIVE AMINOTRANSFERASE PROTEIN [Sinorhizobium meliloti 1021]	LL			
1185	73190	1.00E-142	99	ref NP_385544.1 PROBABLE TRIOSEPHOSPHATE ISOMERASE PROTEIN [Sinorhizobium meliloti 1021]	PEG			
1186	73131	0	99	ref NP_386510.1 beta amino acid--pyruvate transaminase [Sinorhizobium meliloti 1021]	CK			
1187	73108	0	92	ref NP_929378.1 pyruvate kinase [Photobacterium luminescens subsp. laumondii TTO1]	CK			
1188	73074	1.00E-171	87	gb AAO17215.1 Epd [Photobacterium luminescens]	HS			
1189	73132	1.00E-80	100	ref NP_390154.1 nucleoside diphosphate kinase [Bacillus subtilis subsp. subtilis str. 168]	CS			
1190	73146	2.00E-94	93	ref NP_931715.1 inorganic pyrophosphatase [Photobacterium luminescens subsp. laumondii TTO1]	DS			
1191	73147	0	97	ref NP_015444.1 B-type cyclin involved in cell cycle progression; activates Cdc28p to promote the transition from G2 to M phase; accumulates during G2 and M, then targeted via a destruction box motif for ubiquitin-mediated degradation by the proteasome; Clb2p [Saccharomyces cerevisiae]	HS			
1192	73080	0	92	gb AAB17351.1 protein phosphatase type 2C [Saccharomyces cerevisiae]	LN			
1193	73077	0	97	ref NP_010018.1 Endosomal protein that regulates cell polarity, controls polarized growth; similar to Ynr048wp and Lem3p; Cdc50p [Saccharomyces cerevisiae]	HS			
1194	73183	0	93	ref NP_011895.1 Serine/threonine MAP kinase involved in regulating the maintenance of cell wall integrity and progression through the cell cycle; regulated by the PKC1-mediated signaling pathway; Sit2p [Saccharomyces cerevisiae]	CS	HS		
1195	73018	0	100	ref NP_014582.1 NAD-dependent glycerol 3-phosphate dehydrogenase, homolog of Gpd1p, expression is controlled by an oxygen-independent signaling pathway required to regulate metabolism under anoxic conditions; located in cytosol and mitochondria; Gpd2p [Saccharomyces cerevisiae]	HS			
1196	73112	0	97	ref NP_014271.1 Homolog of human tumor suppressor gene PTEN/MMAC1/TEP1 that has lipid phosphatase activity and is linked to the phosphatidylinositol signaling pathway; plays a role in normal sporulation; Tep1p [Saccharomyces cerevisiae]	HS			
1197	73988	0	83	ref NP_013664.1 ER localized integral membrane protein that may promote secretion of certain hexose transporters, including Gal2p; involved in glucose-dependent repression; Gsf2p [Saccharomyces cerevisiae]	SS	PEG		
1198	10908	2.00E-61	100	ref NP_199112.1 ATTRX3 (thioredoxin H-type 3); thiol-disulfide exchange intermediate [Arabidopsis thaliana]	CK			
1198	11147	2.00E-61	100	ref NP_199112.1 ATTRX3 (thioredoxin H-type 3); thiol-disulfide exchange intermediate [Arabidopsis thaliana]	LN			
1199	72909	0	95	ref NP_011569.1 High affinity methionine permease, integral membrane protein with 13 putative membrane-spanning regions; also involved in cysteine uptake; Mup1p [Saccharomyces cerevisiae]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1200	73987	0	94	ref NP_010444.1 Component of the SPS plasma membrane amino acid sensor system (Ssy1p-Ptr3p-Ssy5p), which senses external amino acid concentration and transmits intracellular signals that result in regulation of expression of amino acid permease genes; Ssy1p [Saccharomyces cerevisiae]	PP	SS		
1201	72922	0	91	ref NP_010082.1 Putative transporter, member of the sugar porter family; Ydl199cp [Saccharomyces cerevisiae]	LL			
1202	73036	0	95	ref NP_010022.1 Plasma membrane permease, mediates uptake of glycerophosphoinositol and glycerophosphocholine as sources of the nutrients inositol and phosphate; expression and transport rate are regulated by phosphate and inositol availability; Git1p [Saccharomyces cerevisiae]	CS	HS		
1203	72958	0	94	ref NP_014622.1 High affinity tryptophan and tyrosine permease, overexpression confers FK506 and FTY720 resistance; Tat2p [Saccharomyces cerevisiae]	HS	PP		
1204	77006	1.00E -167	100	ref NP_182073.1 ATAUR3 (ATAURORA3); ATP binding / histone serine kinase(H3-S10 specific) / protein kinase [Arabidopsis thaliana]	LL	LN		
1205	74732	0	90	ref NP_177415.1 casein kinase, putative [Arabidopsis thaliana]	CK			
1206	74333	1.00E -158	100	ref NP_180976.1 protein kinase family protein [Arabidopsis thaliana]	HS			
1207	74735	0	100	ref NP_173489.2 protein kinase [Arabidopsis thaliana]	LN			
1208	74736	0	100	ref NP_176353.1 protein kinase, putative [Arabidopsis thaliana]	LL			
1209	74737	0	100	gb AAF16665.1 AC012394_14putative protein kinase; 59396-62219 [Arabidopsis thaliana]	HS			
1210	76657	0	95	ref NP_187827.1 protein kinase family protein [Arabidopsis thaliana]	SP	LL		
1211	76113	0	94	ref NP_187165.2 protein kinase family protein [Arabidopsis thaliana]	CS			
1212	74747	0	92	ref NP_174161.1 protein kinase family protein [Arabidopsis thaliana]	HS			
1213	77309	0	87	ref NP_175879.2 protein kinase family protein [Arabidopsis thaliana]	PEG			
1214	74322	0	94	gb AAG52342.1 AC011663_21putative protein kinase; 29119-30743 [Arabidopsis thaliana]	LL			
1215	76212	1.00E -170	93	ref NP_200959.2 NADP-dependent oxidoreductase, putative [Arabidopsis thaliana]	HS	LN		
1216	76514	1.00E -159	85	ref NP_568357.1 AAA-type ATPase family protein [Arabidopsis thaliana]	LL	DS		
1217	75242	0	84	ref NP_200318.1 2-oxoacid dehydrogenase family protein [Arabidopsis thaliana]	LL			
1218	77324	0	97	gb AAF79717.1 AC020889_25T1N15.3 [Arabidopsis thaliana]	PEG	LN		
1219	74350	3.00E -95	93	ref NP_178122.1 APT2 (ADENINE PHOSPHORIBOSYL TRANSFERASE 2); adenine phosphoribosyltransferase [Arabidopsis thaliana]	LN			
1220	77606	0	96	ref NP_190925.1 AFC1 (ARABIDOPSIS FUS3-COMPLEMENTING GENE 1); kinase [Arabidopsis thaliana]	HS	SS		
1221	76530	0	92	ref NP_567069.1 F-box family protein-related [Arabidopsis thaliana]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1222	76716	1.00E -86	61	ref NP_565175.1 RNA recognition motif (RRM)-containing protein [Arabidopsis thaliana]	HS			
1223	77328	0	89	ref NP_567904.1 RNA recognition motif (RRM)-containing protein [Arabidopsis thaliana]	HS			
1224	12204	0	100	ref NP_189366.1 G6PD5 (GLUCOSE-6-PHOSPHATE DEHYDROGENASE 5); glucose-6-phosphate 1-dehydrogenase [Arabidopsis thaliana]	DS			
1225	77011	0	96	ref NP_563818.1 strictosidine synthase family protein [Arabidopsis thaliana]	LL			
1226	77607	0	94	ref NP_175157.1 NRAMP2 (NRAMP metal ion transporter 2); metal ion transporter [Arabidopsis thaliana]	LL	PEG		
1227	74612	1.00E -68	100	ref NP_177750.1 unknown protein [Arabidopsis thaliana]	HS			
1228	76126	1.00E -144	100	ref NP_179467.1 signal recognition particle binding [Arabidopsis thaliana]	HS			
1229	74376	1.00E -150	95	ref NP_181340.1 DET2 (DE-ETIOLATED 2) [Arabidopsis thaliana]	DS			
1230	74377	1.00E -46	100	ref NP_181546.1 ATEM6 (ARABIDOPSIS EARLY METHIONINE-LABELLED 6) [Arabidopsis thaliana]	DS	LL	PEG	
1231	74381	6.00E -63	100	ref NP_181990.1 mtACP-1 (MITOCHONDRIAL ACYL CARRIER PROTEIN 1); acyl carrier [Arabidopsis thaliana]	CS			
1232	78364	6.00E -88	87	ref NP_566062.1 UBC6 (UBIQUITIN-CONJUGATING ENZYME 6); ubiquitin-protein ligase [Arabidopsis thaliana]	HS			
1233	74623	1.00E -115	75	ref NP_188379.1 late embryogenesis abundant domain-containing protein / LEA domain-containing protein [Arabidopsis thaliana]	HS			
1234	74668	1.00E -122	88	ref NP_188888.1 late embryogenesis abundant protein, putative / LEA protein, putative [Arabidopsis thaliana]	DS			
1235	74629	5.00E -83	90	ref NP_566782.1 unknown protein [Arabidopsis thaliana]	LL			
1236	74635	1.00E -113	92	ref NP_190864.1 peroxiredoxin type 2, putative [Arabidopsis thaliana]	CS			
1237	74638	1.00E -100	100	ref NP_191788.1 ATARFA1E (ADP-ribosylation factor A1E); GTP binding / phospholipase activator/ protein binding [Arabidopsis thaliana]	HS	SS		
1238	76721	1.00E -148	77	ref NP_192093.1 KCO5 (Ca2+ activated outward rectifying K+ channel 5); outward rectifier potassium channel [Arabidopsis thaliana]	PEG			
1239	78365	2.00E -83	90	ref NP_568004.1 UBC17 (UBIQUITIN-CONJUGATING ENZYME 17); ubiquitin-protein ligase [Arabidopsis thaliana]	CK	HS		
1240	74655	1.00E -132	91	ref NP_195437.1 HCF164 (High chlorophyll fluorescence 164); thiol-disulfide exchange intermediate [Arabidopsis thaliana]	LN			
1241	74660	1.00E -160	92	ref NP_196119.1 ATTOC34/OEP34 (Translocase of chloroplast 34) [Arabidopsis thaliana]	CS	PEG		
1242	74661	1.00E -180	96	ref NP_197135.1 ubiquitin carboxyl-terminal hydrolase family 1 protein [Arabidopsis thaliana]	HS			
1243	75259	1.00E -101	85	ref NP_198051.1 drought-responsive family protein [Arabidopsis thaliana]	HS			
1244	75282	3.00E -93	100	ref NP_201055.1 ATHVA22B (Arabidopsis thaliana HVA22 homologue B)	PEG			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1245	76724	0	93	ref NP_173089.1 organic cation transporter-related [Arabidopsis thaliana]	DS			
1246	12350	1.00E -131	78	gb AAM61424.1 serine O-acetyltransferase (EC 2.3.1.30) Sat-52 [Arabidopsis thaliana]	LN			
1247	76522	0	93	ref NP_001077495.1 unknown protein [Arabidopsis thaliana]	PP			
1248	77903	0	96	ref NP_194952.2 protein kinase family protein [Arabidopsis thaliana]	HS	LL	SS	
1249	76653	0	100	ref NP_568860.1 CIPK21 (CBL-INTERACTING PROTEIN KINASE 21); kinase [Arabidopsis thaliana]	CS			
1250	76524	0	100	ref NP_199469.1 protein kinase family protein [Arabidopsis thaliana]	CS			
1251	11356	1.00E -172	100	ref NP_181165.1 APG10 (ALBINO AND PALE GREEN 10); 1-(5-phosphoribosyl)-5-[(5-phosphoribosylamino)methylideneamino]imidazole-4-carboxamide isomerase [Arabidopsis thaliana]	LN			
1252	75275	0	94	ref NP_199586.1 protein kinase, putative [Arabidopsis thaliana] dbj BAA01715.1 serine/threonine protein kinase [Arabidopsis thaliana]	LL			
1253	76225	0	94	ref NP_195285.3 CONNEXIN 32; kinase [Arabidopsis thaliana]	PEG			
1254	75281	0	94	ref NP_568893.1 protein kinase family protein [Arabidopsis thaliana]	HS			
1255	74384	1.00E -102	100	ref NP_175960.1 prenylated rab acceptor (PRA1) family protein [Arabidopsis thaliana]	LL			
1256	71238	1.00E -168	84	ref NP_563814.1 DCP1 (DECAPPING 1) [Arabidopsis thaliana]	LN			
1257	12354	0	100	ref NP_172738.1 UGE1 (UDP-D-glucose/UDP-D-galactose 4-epimerase 1); UDP-glucose 4-epimerase/ protein dimerization [Arabidopsis thaliana]	DS			
1258	76236	1.00E -157	100	ref NP_201384.1 leucine-rich repeat family protein [Arabidopsis thaliana]	HS			
1259	74682	4.00E -83	100	ref NP_565581.1 unknown protein [Arabidopsis thaliana]	CS			
1260	73425	0	95	ref NP_391649.1 transaminase [Bacillus subtilis subsp. subtilis str. 168]	LL			
1261	73438	0	99	ref YP_351136.1 Glutamate--ammonia ligase [Pseudomonas fluorescens PfO-1]	SP			
1262	73474	0	92	ref NP_790236.1 phosphoglycerate kinase [Pseudomonas syringae pv. tomato str. DC3000]	HS	PP		
1263	70850	0	97	ref NP_190853.1 ATELP1 (VACUOLAR SORTING RECEPTOR HOMOLOG) [Arabidopsis thaliana]	HS			
1264	73429	1.00E -144	100	ref ZP_01370194.1 pyrroline-5-carboxylate reductase [Desulfotobacterium hafniense DCB-2]	HS	SS	CS	
1265	73441	0	94	ref ZP_01370604.1 enolase [Desulfotobacterium hafniense DCB-2]	PEG			
1266	73466	0	97	ref ZP_00108577.1 COG1063: Threonine dehydrogenase and related Zn-dependent dehydrogenases [Nostoc punctiforme PCC 73102]	LN			
1267	73455	1.00E -159	100	ref NP_792905.1 UTP-glucose-1-phosphate uridylyltransferase [Pseudomonas syringae pv. tomato str. DC3000]	HS			
1268	73432	0	94	ref YP_582700.1 Alcohol dehydrogenase, zinc-binding [Ralstonia metallidurans CH34]	CS			
1269	73435	0	97	ref NP_391271.1 phosphoglyceromutase [Bacillus subtilis subsp. subtilis str. 168]	LN			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1270	77726	0	99	ref NP_391273.1 phosphoglycerate kinase [Bacillus subtilis subsp. subtilis str. 168]	HS			
1271	73483	0	96	ref NP_009362.1 Pyruvate kinase, functions as a homotetramer in glycolysis to convert phosphoenolpyruvate to pyruvate, the input for aerobic (TCA cycle) or anaerobic (glucose fermentation) respiration; Cdc19p [Saccharomyces cerevisiae]	LL			
1272	73472	0	99	ref NP_387397.1 ASPARTATE AMINOTRANSFERASE B PROTEIN [Sinorhizobium meliloti 1021]	DS			
1273	73526	0	94	ref NP_242877.1 benzyl alcohol dehydrogenase [Bacillus halodurans C-125]	DS			
1274	73538	0	100	ref NP_243126.1 hypothetical protein BH2260 [Bacillus halodurans C-125]	HS			
1275	73574	1.00E-177	100	ref NP_244591.1 2-keto-3-deoxygluconate kinase [Bacillus halodurans C-125]	HS			
1276	73539	1.00E-170	100	ref NP_287481.1 UTP--glucose-1-phosphate uridylyltransferase subunit GalU [Escherichia coli O157:H7 EDL933]	CS			
1277	73563	3.00E-62	90	ref NP_743010.1 nucleoside diphosphate kinase [Pseudomonas putida KT2440]	CK			
1278	73575	1.00E-175	100	ref NP_792508.1 fructokinase [Pseudomonas syringae pv. tomato str. DC3000]	LL			
1279	73565	0	95	ref NP_011707.1 High-affinity histidine permease, also involved in the transport of manganese ions; Hip1p [Saccharomyces cerevisiae]	HS			
1280	73531	0	93	ref NP_388151.1 hypothetical protein BSU02690 [Bacillus subtilis subsp. subtilis str. 168]	DS			
1281	73592	0	96	ref NP_391888.1 6-phosphogluconate dehydrogenase [Bacillus subtilis subsp. subtilis str. 168]	LN			
1282	74149	0	100	ref NP_794093.1 pyruvate kinase [Pseudomonas syringae pv. tomato str. DC3000]	LL			
1283	74174	0	99	ref NP_416880.1 predicted aminotransferase, PLP-dependent [Escherichia coli K12]	LL	LN		
1284	74151	0	100	ref NP_417148.1 4-aminobutyrate aminotransferase, PLP-dependent [Escherichia coli K12]	PP			
1285	74104	0	100	ref ZP_01369074.1 glucose-1-phosphate adenylyltransferase [Desulfotobacterium hafniense DCB-2]	CS	HS		
1286	74164	0	89	ref NP_242876.1 aldehyde dehydrogenase [Bacillus halodurans C-125]	DS			
1287	74129	0	77	ref YP_001185919.1 methylmalonate-semialdehyde dehydrogenase [Pseudomonas mendocina ymp]	CK			
1288	72340	0	97	ref NP_195674.2 GGT1; gamma-glutamyltransferase/ glutathione gamma-glutamylcysteinyltransferase [Arabidopsis thaliana]	HS			
1289	74155	0	100	ref NP_385556.1 dihydrolipoamide dehydrogenase [Sinorhizobium meliloti 1021]	HS	LL		
1290	74168	0	87	ref NP_927730.1 glutathione reductase [Photorhabdus luminescens subsp. laumondii TTO1]	LN			
1291	74133	0	96	ref ZP_00109554.2 COG1249: Pyruvate/2-oxoglutarate dehydrogenase complex, dihydrolipoamide dehydrogenase (E3) component, and related enzymes [Nostoc punctiforme PCC 73102]	HS			
1292	74160	4.00E-95	99	ref NP_441445.1 hypothetical protein slr0816 [Synechocystis sp. PCC 6803] dbj BAA18125.1 slr0816 [Synechocystis sp. PCC 6803]	LL			
1293	74485	0	88	ref NP_439939.1 hypothetical protein slr1495 [Synechocystis sp. PCC 6803]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1294	74450	0	100	ref NP_442868.1 LIM17 protein [Synechocystis sp. PCC 6803]	CK			
1295	74487	0	90	ref NP_357138.2 branched-chain alpha-keto acid dehydrogenase subunit E2 [Agrobacterium tumefaciens str. C58]	PP			
1296	74406	0	96	gb AAN78639.1 AE016755_139Dihydrolipoamide dehydrogenase [Escherichia coli CFT073]	LN			
1297	74454	1.00E -112	91	ref NP_927456.1 ribulose-phosphate 3-epimerase (pentose-5-phosphate 3-epimerase) (PPE) (R5P3E) [Photorhabdus luminescens subsp. laumondii TTO1]	LL			
1298	11361	6.00E -61	83	ref NP_176291.1 FED A (FERREDOXIN 2); electron carrier/ iron ion binding [Arabidopsis thaliana]	LN			
1299	74431	0	99	ref NP_390796.1 pyruvate kinase [Bacillus subtilis subsp. subtilis str. 168]	HS			
1300	74411	0	94	ref NP_949979.1 fructose-1,6-bisphosphatase [Rhodospseudomonas palustris CGA009]	SS			
1301	74424	1.00E -172	92	ref NP_386873.1 PROBABLE FRUCTOSE-BISPHOSPHATE ALDOLASE CLASS I PROTEIN [Sinorhizobium meliloti 1021]	PP			
1302	74525	0	83	ref NP_931210.1 glutamate synthase subunit beta [Photorhabdus luminescens subsp. laumondii TTO1]	HS			
1303	74573	1.00E -75	38	ref YP_720246.1 aspartate aminotransferase [Trichodesmium erythraeum IMS101]	LN			
1304	75827	2.00E -44	82	ref NP_565988.1 unknown protein [Arabidopsis thaliana]	LL			
1305	77329	0	98	ref NP_566005.1 mitochondrial transcription termination factor-related / mTERF-related [Arabidopsis thaliana]	CK			
1306	77330	0	97	ref NP_182296.1 auxin-responsive GH3 family protein [Arabidopsis thaliana]	PEG			
1307	75830	6.00E -61	100	ref NP_186754.1 MUB1 (MEMBRANE-ANCHORED UBIQUITIN-FOLD PROTEIN 1 PRECURSOR) [Arabidopsis thaliana]	CK			
1308	75832	2.00E -42	85	ref NP_186788.1 VMA10 (VACUOLAR MEMBRANE ATPASE 10) [Arabidopsis thaliana]	HS			
1309	75836	7.00E -41	73	ref NP_187208.1 unknown protein [Arabidopsis thaliana]	CS			
1310	77034	4.00E -28	95	ref NP_172702.1 unknown protein [Arabidopsis thaliana]	HS	PP		
1311	76255	0	92	ref NP_566390.1 unknown protein [Arabidopsis thaliana]	LL			
1312	76542	1.00E -77	69	ref NP_188337.1 unknown protein [Arabidopsis thaliana]	CS	LL		
1313	77511	0	90	ref NP_566642.1 unknown protein [Arabidopsis thaliana]	SS	CK		
1314	76261	1.00E -127	86	ref NP_563649.1 unknown protein [Arabidopsis thaliana]	HS			
1315	78456	0	97	ref NP_189202.1 MAP1B (METHIONINE AMINOPEPTIDASE 1C); metalloexopeptidase [Arabidopsis thaliana]	CK			
1316	77041	1.00E -125	95	ref NP_189236.3 plastid-lipid associated protein PAP / fibrillin family protein [Arabidopsis thaliana]	LN			
1317	76546	0	90	gb AAF79247.1 AC006917_32F10B6.11 [Arabidopsis thaliana]	LL			
1318	75848	9.00E -96	100	ref NP_189535.1 AIG2 (AVRRPT2-INDUCED GENE 2) [Arabidopsis thaliana]	CK	HS		

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1319	77515	0	80	ref NP_189945.1 zinc knuckle (CCHC-type) family protein [Arabidopsis thaliana]	HS			
1320	76266	0	82	emb CAB83070.1 putative protein [Arabidopsis thaliana]	PEG			
1321	75852	1.00E -26	100	ref NP_190227.1 unknown protein [Arabidopsis thaliana]	HS			
1322	76556	0	86	ref NP_190445.2 zinc finger (DHHC type) family protein [Arabidopsis thaliana]	PEG			
1323	77523	1.00E -174	96	ref NP_001078291.1 unknown protein [Arabidopsis thaliana]	HS			
1324	75863	1.00E -120	100	ref NP_191267.1 eukaryotic rpb5 RNA polymerase subunit family protein [Arabidopsis thaliana]	CS			
1325	77052	1.00E -171	96	ref NP_564000.2 unknown protein [Arabidopsis thaliana]	PP			
1326	75865	1.00E -56	80	ref NP_173123.1 ribosomal protein-related [Arabidopsis thaliana]	CS			
1327	76551	0	92	ref NP_173144.1 oxidoreductase, 2OG-Fe(II) oxygenase family protein [Arabidopsis thaliana]	PEG			
1328	76280	0	86	ref NP_564064.1 phytochrome kinase substrate-related [Arabidopsis thaliana]	LL			
1329	75874	7.00E -96	100	ref NP_193404.1 glycosyltransferase family protein 28 [Arabidopsis thaliana]	PP			
1330	75879	9.00E -62	100	ref NP_194229.1 ATGP4 (Arabidopsis thaliana geranylgeranylated protein)	HS			
1331	75880	7.00E -59	65	ref NP_173435.1 unknown protein [Arabidopsis thaliana]	HS			
1332	77519	1.00E -176	90	ref NP_196118.1 sad1/unc-84 protein-related [Arabidopsis thaliana]	HS	CK		
1333	76745	1.00E -111	88	dbj BAB09403.1 unnamed protein product [Arabidopsis thaliana]	HS			
1334	76554	1.00E -173	91	ref NP_197092.1 unknown protein [Arabidopsis thaliana]	DS			
1335	76746	3.00E -93	75	ref NP_568333.1 unknown protein [Arabidopsis thaliana]	CK			
1336	76750	1.00E -85	93	ref NP_197746.1 unknown protein [Arabidopsis thaliana]	LL			
1337	76621	3.00E -20	100	ref NP_198912.1 unknown protein [Arabidopsis thaliana]	HS			
1338	76623	1.00E -23	80	ref NP_568739.1 unknown protein [Arabidopsis thaliana]	LL			
1339	76461	1.00E -29	78	gb AAB86938.1 NOI protein [Arabidopsis thaliana]	CS			
1340	76462	5.00E -31	100	ref NP_564277.1 unknown protein [Arabidopsis thaliana]	HS	PP		
1341	78987	1.00E -147	81	ref NP_201096.1 protein binding / zinc ion binding [Arabidopsis thaliana]	HS			
1342	76568	0	96	ref NP_564484.1 unknown protein [Arabidopsis thaliana]	HS			
1343	76176	0	95	ref NP_565009.1 glycosyltransferase family 14 protein / core-2/I-branching enzyme family protein [Arabidopsis thaliana]	CK			
1344	76177	1.00E -97	100	ref NP_177287.1 unknown protein [Arabidopsis thaliana]	LN			
1345	78380	0	100	ref NP_565058.1 eukaryotic translation initiation factor-related [Arabidopsis thaliana]	LN			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1346	76629	2.00E -46	100	ref NP_177894.1 unknown protein [Arabidopsis thaliana]	HS			
1347	76189	2.00E -58	100	gb AAD15467.2 unknown protein [Arabidopsis thaliana]	CS			
1348	76761	1.00E -106	94	ref NP_179133.1 ATARFB1A (ADP-ribosylation factor B1A); GTP binding [Arabidopsis thaliana]	LL			
1349	78608	0	100	ref NP_973624.1 ATORC2/ORC2 (ORIGIN RECOGNITION COMPLEX SECOND LARGEST SUBUNIT) [Arabidopsis thaliana]	PP			
1350	78115	1.00E -153	100	ref NP_181669.1 embryo-abundant protein-related [Arabidopsis thaliana]	CK	DS		
1351	74873	0	100	ref NP_416275.1 glutamate dehydrogenase, NADP-specific [Escherichia coli K12]	DS			
1352	74826	0	99	ref NP_416651.1 predicted oxidoreductase [Escherichia coli K12]	HS			
1353	74850	0	99	gb AAN81953.1 AE016766_41D-erythrose 4-phosphate dehydrogenase [Escherichia coli CFT073]	HS			
1354	74887	4.00E -76	90	gb AAD25168.1 aquaporin [Saccharomyces cerevisiae]	PEG			
1355	74804	0	95	gb EDN59233.1 chromatin remodeling complex member, RSC [Saccharomyces cerevisiae YJM789]	HS	SS	PEG	
1356	74831	1.00E -167	81	gb EAY97794.1 hypothetical protein OsI_019027 [Oryza sativa (indica cultivar-group)]	LN			
1357	74822	0	69	emb CAO62934.1 unnamed protein product [Vitis vinifera]	PEG			
1358	74836	1.00E -121	70	ref NP_187378.1 transcriptional activator, putative [Arabidopsis thaliana]	LN			
1359	74901	3.00E -31	61	ref NP_001064928.1 Os10g0491900 [Oryza sativa (japonica cultivar-group)]	HS			
1360	74974	1.00E -112	87	ref NP_001063296.1 Os09g0443500 [Oryza sativa (japonica cultivar-group)]	HS			
1361	74904	1.00E -149	82	ref NP_001063966.1 Os09g0567900 [Oryza sativa (japonica cultivar-group)]	HS			
1362	74916	8.00E -92	76	gb EAZ28333.1 hypothetical protein OsJ_011816 [Oryza sativa (japonica cultivar-group)]	PEG			
1363	74964	1.00E -158	95	ref NP_417814.1 predicted phosphoribulokinase [Escherichia coli K12]	LL			
1364	74918	6.00E -82	82	ref NP_443015.1 hypothetical protein sl1109 [Synechocystis sp. PCC 6803]	LN			
1365	74966	0	99	ref NP_442146.1 hypothetical protein sl10209 [Synechocystis sp. PCC 6803]	HS			
1366	74979	1.00E -142	82	ref NP_440823.1 hypothetical protein slr1220 [Synechocystis sp. PCC 6803]	CK	LL		
1367	74934	0	96	ref YP_350971.1 betaine aldehyde dehydrogenase [Pseudomonas fluorescens PfO-1]	PEG			
1368	77814	0	92	ref NP_116585.1 F-box protein required for G1/S and G2/M transition, associates with Skp1p and Cdc53p to form a complex, SCFCdc4, which acts as ubiquitin-protein ligase directing ubiquitination of the phosphorylated CDK inhibitor Sic1p; Cdc4p [Saccharomyces cerevisiae]	HS			
1369	75387	1.00E -148	53	ref NP_001067713.1 Os11g0293900 [Oryza sativa (japonica cultivar-group)]	CK	LL		
1370	75328	0	93	ref NP_001050986.1 Os03g0699200 [Oryza sativa (japonica cultivar-group)]	PP			
1371	77808	0	74	emb CAO63310.1 unnamed protein product [Vitis vinifera]	CS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1372	75389	0	69	ref NP_001045085.1 Os01g0897100 [Oryza sativa (japonica cultivar-group)]	HS			
1373	75318	1.00E -126	49	emb CAO40140.1 unnamed protein product [Vitis vinifera]	CK			
1374	75330	0	70	emb CAH66847.1 H0525C06.10 [Oryza sativa (indica cultivar-group)]	CS	HS		
1375	75342	1.00E -75	69	ref NP_001056004.1 Os05g0509700 [Oryza sativa (japonica cultivar-group)]	CK			
1376	75366	4.00E -59	74	ref NP_001063106.1 Os09g0397700 [Oryza sativa (japonica cultivar-group)]	HS			
1377	75319	1.00E -121	73	ref NP_001050079.1 Os03g0343700 [Oryza sativa (japonica cultivar-group)]	HS			
1378	75391	0	96	gb ABU48662.1 G protein alpha subunit [Sorghum bicolor]	HS			
1379	75334	4.00E -45	92	ref NP_001049605.1 Os03g0257900 [Oryza sativa (japonica cultivar-group)]	HS			
1380	75323	0	82	ref NP_001053032.1 Os04g0466600 [Oryza sativa (japonica cultivar-group)]	HS			
1381	75396	5.00E -84	77	ref NP_001068537.1 Os11g0703400 [Oryza sativa (japonica cultivar-group)]	PEG			
1382	75401	2.00E -72	93	ref NP_001059557.1 Os07g0454700 [Oryza sativa (japonica cultivar-group)]	HS			
1383	75413	1.00E -139	90	ref NP_001050848.1 Os03g0666700 [Oryza sativa (japonica cultivar-group)]	LL			
1384	75461	0	92	ref NP_001066463.1 Os12g0236500 [Oryza sativa (japonica cultivar-group)]	HS	SS	PP	
1385	75473	6.00E -50	71	ref NP_001043914.1 Os01g0687600 [Oryza sativa (japonica cultivar-group)]	DS			
1386	77817	8.00E -26	93	ref NP_001045143.1 Os01g0908400 [Oryza sativa (japonica cultivar-group)]	LL			
1387	75416	1.00E -165	81	gb EAY92384.1 hypothetical protein Osl_013617 [Oryza sativa (indica cultivar-group)]	CK			
1388	75464	0	68	gb EAZ22667.1 hypothetical protein OsJ_006150 [Oryza sativa (japonica cultivar-group)]	HS			
1389	75441	5.00E -46	68	gb AAA33773.1 PVPR3	LN			
1390	75454	2.00E -82	73	ref NP_001064989.1 Os10g0502000 [Oryza sativa (japonica cultivar-group)]	HS			
1391	75493	4.00E -31	51	emb CAO48835.1 unnamed protein product [Vitis vinifera]	HS	PP	PEG	
1392	75446	2.00E -29	55	gb ABU54835.1 defender against apoptotic death [Penaeus monodon]	CS	HS		
1393	75495	2.00E -86	82	gb ABG23395.1 2C-methyl-D-erythritol 2,4-cyclodiphosphate synthase [Stevia rebaudiana]	SS			
1394	75705	1.00E -102	71	ref NP_001046126.1 Os02g0187100 [Oryza sativa (japonica cultivar-group)]	SP			
1395	77818	0	100	ref NP_241861.1 succinate-semialdehyde dehydrogenase [Bacillus halodurans C-125]	HS	PEG		
1396	75525	0	99	ref NP_415448.1 aspartate aminotransferase, PLP-dependent [Escherichia coli K12]	LN			
1397	75549	1.00E -115	37	dbj BAC10067.1 hypothetical protein [Oryza sativa Japonica Group]	SS			
1398	75582	0	68	emb CAO16731.1 unnamed protein product [Vitis vinifera]	CS	PEG		
1399	75535	6.00E -53	62	emb CAO18074.1 unnamed protein product [Vitis vinifera]	PEG			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1400	75547	1.00E -164	74	emb CAO42486.1 unnamed protein product [Vitis vinifera]	LL	LN		
1401	75512	1.00E -141	63	emb CAO40060.1 unnamed protein product [Vitis vinifera]	CS	HS		
1402	75673	1.00E -158	55	ref NP_001050262.1 Os03g0387900 [Oryza sativa (japonica cultivar-group)]	CK			
1403	75626	0	78	gb EAZ07965.1 hypothetical protein Osl_029197 [Oryza sativa (indica cultivar-group)]	HS			
1404	75674	4.00E -53	66	gb EAY98259.1 hypothetical protein Osl_019492 [Oryza sativa (indica cultivar-group)]	HS	LN		
1405	75615	2.00E -46	59	gb EAY75762.1 hypothetical protein Osl_003609 [Oryza sativa (indica cultivar-group)]	CK			
1406	75627	1.00E -135	92	gb EAZ15023.1 hypothetical protein OsJ_004848 [Oryza sativa (japonica cultivar-group)]	SS			
1407	75628	1.00E -103	88	gb AAQ24835.1 partner of Nob1 [Sorghum bicolor]	PEG			
1408	75652	1.00E -139	85	gb EAY93006.1 hypothetical protein Osl_014239 [Oryza sativa (indica cultivar-group)]	CK			
1409	75605	1.00E -121	55	ref NP_001047861.1 Os02g0704600 [Oryza sativa (japonica cultivar-group)]	CK	HS	LN	
1410	70228	1.00E -83	100	ref NP_189581.1 AHP2 (HISTIDINE-CONTAINING PHOSPHOTRANSFER 2); histidine phosphotransfer kinase/ signal transducer [Arabidopsis thaliana]	CK			
1411	13302	0	95	gb AAF21885.1 AF101056_1MEI2 [Arabidopsis thaliana]	LN			
1412	75610	0	86	ref NP_001054248.1 Os04g0675500 [Oryza sativa (japonica cultivar-group)]	CK			
1413	75623	0	95	ref NP_194630.1 AIM1 (ABNORMAL INFLORESCENCE MERISTEM); enoyl-CoA hydratase [Arabidopsis thaliana]	HS			
1414	75671	0	88	ref NP_176918.1 leucine-rich repeat family protein [Arabidopsis thaliana]	PP			
1415	75695	4.00E -78	63	emb CAO21783.1 unnamed protein product [Vitis vinifera]	HS			
1416	75648	2.00E -32	46	ref NP_001054395.1 Os05g0103500 [Oryza sativa (japonica cultivar-group)] dbj BAF16309.1 Os05g0103500 [Oryza sativa (japonica cultivar-group)]	SS			
1417	75672	0	84	ref NP_196925.1 leucine-rich repeat transmembrane protein kinase, putative [Arabidopsis thaliana] dbj BAB08300.1 receptor protein kinase-like protein [Arabidopsis thaliana]	PP			
1418	75713	1.00E -141	43	emb CAO21763.1 unnamed protein product [Vitis vinifera]	CK			
1419	75714	7.00E -35	36	gb ABK93511.1 unknown [Populus trichocarpa]	HS	PP		
1420	75762	1.00E -171	65	emb CAN68895.1 hypothetical protein [Vitis vinifera]	CK			
1421	77954	0	94	ref NP_172684.1 flavin-containing monooxygenase family protein / FMO family protein [Arabidopsis thaliana]	HS			
1422	77955	0	95	ref NP_172680.1 flavin-containing monooxygenase family protein / FMO family protein [Arabidopsis thaliana]	HS			
1423	77541	7.00E -85	100	gb ABK28567.1 unknown [Arabidopsis thaliana]	LN			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1424	78131	0	93	emb CAB42923.1 putative mitochondrial protein [Arabidopsis thaliana]	HS			
1425	78134	6.00E -95	91	ref NP_192051.2 CBL5 (CALCINEURIN B-LIKE PROTEIN 5) [Arabidopsis thaliana]	LN	SS		
1426	77554	1.00E -138	75	tpg DAA00869.1 TPA_exp: PDR2 ABC transporter [Arabidopsis thaliana]	HS			
1427	78992	1.00E -170	90	ref NP_194508.1 calcium-binding EF hand family protein [Arabidopsis thaliana]	CK			
1428	77557	0	84	ref NP_196016.1 IQD12 (IQ-domain 12); calmodulin binding [Arabidopsis thaliana]	LL	LN		
1429	78128	0	96	ref NP_196397.1 flavin-containing monooxygenase family protein / FMO family protein [Arabidopsis thaliana]	HS			
1430	77560	0	94	ref NP_568202.1 calcium-binding EF hand family protein [Arabidopsis thaliana]	CK	HS		
1431	77561	0	100	ref NP_196694.1 monooxygenase family protein [Arabidopsis thaliana]	CS	CK		
1432	77958	1.00E -108	74	ref NP_197694.1 TET12 (TETRASPANIN12) [Arabidopsis thaliana]	PP			
1433	77563	0	92	ref NP_197887.1 integral membrane transporter family protein [Arabidopsis thaliana]	CS	CK		
1434	78994	0	91	ref NP_001032056.1 F-box family protein [Arabidopsis thaliana]	HS			
1435	74508	0	90	ref NP_850439.1 CYP76C1 (cytochrome P450, family 76, subfamily C, polypeptide 1); heme binding / iron ion binding / monooxygenase [Arabidopsis thaliana]	CS	HS		
1436	70229	0	100	ref NP_194337.1 MEK1 (mitogen-activated protein kinase kinase 1); MAP kinase kinase/ kinase [Arabidopsis thaliana]	HS			
1437	77959	0	90	ref NP_200937.1 flavin-containing monooxygenase family protein / FMO family protein [Arabidopsis thaliana]	HS			
1438	77567	0	88	ref NP_171951.2 integral membrane transporter family protein [Arabidopsis thaliana]	HS	LN		
1439	11749	2.00E -52	74	ref NP_181812.1 PDF1 (PROTODERMAL FACTOR 1) [Arabidopsis thaliana]	CK			
1440	77578	0	90	ref NP_180886.1 integral membrane transporter family protein [Arabidopsis thaliana]	HS			
1441	11751	1.00E -110	90	ref NP_180871.1 SAR1 (SYNAPTOBREVIN-RELATED PROTEIN 1) [Arabidopsis thaliana]	LN			
1442	77963	0	86	ref NP_566652.1 unknown protein [Arabidopsis thaliana]	LN			
1443	77925	1.00E -149	94	ref NP_190985.1 ATFIP37 (ARABIDOPSIS THALIANA FKBP12 INTERACTING PROTEIN 37) [Arabidopsis thaliana]	HS	PP		
1444	77927	0	90	ref NP_564083.1 unknown protein [Arabidopsis thaliana]	HS			
1445	77345	2.00E -57	100	ref NP_564113.1 unknown protein [Arabidopsis thaliana]	HS			
1446	70230	0	96	gb ABF55662.1 double MYC-tagged mitogen activated protein kinase kinase 2 [synthetic construct]	HS			
1447	77353	1.00E -65	100	ref NP_564301.1 unknown protein [Arabidopsis thaliana]	HS			
1448	77354	6.00E -42	81	ref NP_564312.1 unknown protein [Arabidopsis thaliana]	LL			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1449	77355	1.00E -41	90	ref NP_564515.1 unknown protein [Arabidopsis thaliana]	PP			
1450	77588	1.00E -152	100	ref NP_176415.1 unknown protein [Arabidopsis thaliana]	CS			
1451	77591	1.00E -164	92	ref NP_565628.1 unknown protein [Arabidopsis thaliana]	LL			
1452	77594	1.00E -143	72	emb CAB46003.1 RING-H2 finger protein RHF1a [Arabidopsis thaliana]	DS			
1453	77595	1.00E -144	71	ref NP_567661.1 SWIB complex BAF60b domain-containing protein [Arabidopsis thaliana]	LL			
1454	77364	5.00E -76	84	ref NP_194770.1 transcription factor [Arabidopsis thaliana]	LN			
1455	78904	0	100	emb CAB87764.1 putative protein [Arabidopsis thaliana]	HS			
1456	77937	0	100	ref NP_197133.1 acetolactate synthase small subunit, putative [Arabidopsis thaliana]	HS			
1457	77369	1.00E -103	86	ref NP_197494.1 plastid-lipid associated protein PAP-related / fibrillin-related [Arabidopsis thaliana]	DS			
1458	77940	0	89	ref NP_568648.1 transducin family protein / WD-40 repeat family protein [Arabidopsis thaliana]	CK	HS		
1459	78746	0	100	ref NP_201387.2 ankyrin repeat family protein [Arabidopsis thaliana]	HS			
1460	77372	1.00E -119	94	ref NP_201515.1 F-box family protein [Arabidopsis thaliana]	DS			
1461	78126	0	86	gb AAF24593.1 AC007654_9T19E23.14 [Arabidopsis thaliana]	CK			
1462	77951	0	100	ref NP_565803.1 ATEYA (ARABIDOPSIS THALIANA EYES ABSENT HOMOLOG); protein tyrosine phosphatase, metal-dependent [Arabidopsis thaliana]	LL	PEG		
1463	75901	0	97	gb EDN61448.1 F-box protein [Saccharomyces cerevisiae YJM789]	CS			
1464	12123	1.00E -115	86	ref NP_196598.1 EMB3010 (EMBRYO DEFECTIVE 3010); structural constituent of ribosome [Arabidopsis thaliana]	LN			
1465	75987	9.00E -84	80	ref NP_001105929.1 high affinity nitrate transporter [Zea mays]	CS			
1466	75940	0	100	ref NP_842265.1 Phosphoglucose isomerase (PGI) [Nitrosomonas europaea ATCC 19718]	HS	PP		
1467	75919	1.00E -175	95	ref NP_780153.1 isocitrate dehydrogenase [Xylella fastidiosa Temecula1]	HS	LL	SS	
1468	75908	0	99	ref NP_440384.1 glutamate decarboxylase [Synechocystis sp. PCC 6803]	HS			
1469	75920	0	100	ref NP_440189.1 ferredoxin-sulfite reductase [Synechocystis sp. PCC 6803]	HS			
1470	75992	0	94	ref YP_001636715.1 pyruvate kinase [Chloroflexus aurantiacus J-10-fl]	HS			
1471	75933	1.00E -116	94	ref NP_442343.1 ribose-5-phosphate isomerase A [Synechocystis sp. PCC 6803]	HS	PP		
1472	77965	0	92	ref NP_181991.1 kelch repeat-containing F-box family protein [Arabidopsis thaliana]	DS			
1473	77966	0	89	ref NP_186818.1 unknown protein [Arabidopsis thaliana]	LN			
1474	77972	0	90	ref NP_189657.1 unknown protein [Arabidopsis thaliana]	HS			
1475	78522	0	96	ref NP_175744.1 unknown protein [Arabidopsis thaliana]	SP			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1476	12194	3.00E -78	91	ref NP_176772.1 TPX2 (THIOREDOXIN-DEPENDENT PEROXIDASE 2); antioxidant [Arabidopsis thaliana]	CK			
1477	78525	0	87	gb AAC24050.1 Similar to S. cerevisiae SIK1P protein	LN			
1478	78527	0	87	gb AAC18814.1 Contains homology to serine/threonine protein kinase gb X99618 from Mycobacterium tuberculosis.	PEG			
1479	78917	0	95	ref NP_177258.3 armadillo/beta-catenin repeat family protein / U-box domain-containing protein [Arabidopsis thaliana]	LN			
1480	78920	0	100	ref NP_565146.1 ACR3 (ACT Domain Repeat 3) [Arabidopsis thaliana]	PP			
1481	78535	0	96	ref NP_179324.2 pantothenate kinase-related [Arabidopsis thaliana]	CK			
1482	78538	0	98	ref NP_180538.1 kelch repeat-containing F-box family protein [Arabidopsis thaliana]	HS			
1483	78922	0	95	ref NP_181393.1 F-box family protein [Arabidopsis thaliana]	HS	PP	CS	
1484	77984	1.00E -109	100	ref NP_566068.1 unknown protein [Arabidopsis thaliana]	LN	CK		
1485	73938	0	100	ref NP_201190.1 AAP4 (amino acid permease 4); amino acid permease [Arabidopsis thaliana]	DS	CS		
1486	78621	0	100	gb AAD13704.1 hypothetical protein [Arabidopsis thaliana]	HS			
1487	77990	1.00E -74	85	ref NP_187290.1 integral membrane family protein [Arabidopsis thaliana]	DS			
1488	78386	1.00E -132	100	ref NP_187319.1 peptidyl-prolyl cis-trans isomerase cyclophilin-type family protein [Arabidopsis thaliana]	SP			
1489	78001	1.00E -70	100	gb AAF23196.1 AC016795_9unknown protein [Arabidopsis thaliana]	HS			
1490	78543	0	90	ref NP_187807.1 S-locus related protein SLR1, putative (S1) [Arabidopsis thaliana]	HS			
1491	78002	1.00E -136	100	ref NP_187847.1 unknown protein [Arabidopsis thaliana]	LL			
1492	12259	1.00E -134	100	gb AAM67061.1 ribosomal protein S2, putative [Arabidopsis thaliana]	LN			
1493	78005	3.00E -93	96	ref NP_172796.1 unknown protein [Arabidopsis thaliana]	LL			
1494	78008	1.00E -135	88	ref NP_188459.1 unknown protein [Arabidopsis thaliana]	PEG			
1495	78019	1.00E -176	94	ref NP_001031574.1 unknown protein [Arabidopsis thaliana]	CK	HS		
1496	78020	2.00E -74	89	gb ABK28619.1 unknown [Arabidopsis thaliana]	SS			
1497	78023	1.00E -121	94	ref NP_173373.2 unknown protein [Arabidopsis thaliana]	DS			
1498	78548	1.00E -146	86	ref NP_564100.1 abscisic acid-responsive HVA22 family protein [Arabidopsis thaliana]	LN	PP		
1499	78161	1.00E -103	94	ref NP_173738.2 caleosin-related family protein [Arabidopsis thaliana]	HS	PEG		
1500	78162	1.00E -179	100	ref NP_564270.1 unknown protein [Arabidopsis thaliana]	CK	LN	PP	PEG
1501	78033	1.00E -148	90	ref NP_564297.1 unknown protein [Arabidopsis thaliana]	LL			
1502	78622	0	95	ref NP_564315.1 unknown protein [Arabidopsis thaliana]	CK			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1503	78036	1.00E -67	100	ref NP_174257.1 unknown protein [Arabidopsis thaliana]	DS			
1504	78043	0	96	ref NP_175182.2 endonuclease/exonuclease/phosphatase family protein [Arabidopsis thaliana]	CS			
1505	78165	2.00E -92	84	ref NP_564788.1 unknown protein [Arabidopsis thaliana]	LL			
1506	78066	1.00E -116	86	gb AAF68106.1 AC010793_1F20B17.2 [Arabidopsis thaliana]	PP			
1507	78069	1.00E -124	64	ref NP_178248.2 pentatricopeptide (PPR) repeat-containing protein [Arabidopsis thaliana]	PP			
1508	78183	1.00E -115	100	ref NP_180390.2 unknown protein [Arabidopsis thaliana]	PP			
1509	78170	2.00E -37	100	gb AAS47599.1 At2g33690 [Arabidopsis thaliana]	PP			
1510	78953	1.00E -125	88	ref NP_565921.1 unknown protein [Arabidopsis thaliana]	PP			
1511	78929	0	97	ref NP_182026.1 CCD7 (more axillary growth 3) [Arabidopsis thaliana]	LN			
1512	78564	1.00E -63	100	ref NP_188669.1 unknown protein [Arabidopsis thaliana]	PEG			
1513	78566	0	100	ref NP_566813.1 unknown protein [Arabidopsis thaliana]	HS			
1514	78591	0	96	gb AAG12668.1 AC027033_3hypothetical protein; 23726-25026 [Arabidopsis thaliana]	SS			
1515	78181	3.00E -99	100	ref NP_563968.1 unknown protein [Arabidopsis thaliana]	LN			
1516	78190	2.00E -30	71	ref NP_564173.1 unknown protein [Arabidopsis thaliana]	LL			
1517	78571	0	93	gb AAF79874.1 AC000348_27T7N9.16 [Arabidopsis thaliana]	CK			
1518	78933	1.00E -107	84	gb AAG51731.1 AC068667_10hypothetical protein; 69078-67980 [Arabidopsis thaliana]	LN			
1519	78573	8.00E -73	80	ref NP_174417.1 LOB domain protein 4 / lateral organ boundaries domain protein 4 (LBD4) [Arabidopsis thaliana]	CK			
1520	78632	0	100	ref NP_564507.1 unknown protein [Arabidopsis thaliana]	HS	LL	PP	
1521	78959	0	100	ref NP_175985.1 pentatricopeptide (PPR) repeat-containing protein [Arabidopsis thaliana]	LN			
1522	78191	3.00E -51	63	ref NP_564986.1 ribosomal protein L12 family protein [Arabidopsis thaliana]	HS	LN		
1523	78595	1.00E -150	92	gb AAB70396.1 Similar to ATP-dependent Clp protease (gb D90915).	HS	PEG		
1524	78646	0	100	ref NP_172453.1 pentatricopeptide (PPR) repeat-containing protein [Arabidopsis thaliana]	HS			
1525	76325	1.00E -143	82	gb EAY88291.1 hypothetical protein Osl_009524 [Oryza sativa (indica cultivar-group)]	CS			
1526	76337	6.00E -76	87	ref NP_001053732.1 Os04g0595200 [Oryza sativa (japonica cultivar-group)]	HS			
1527	76349	1.00E -139	56	ref NP_001066102.1 Os12g0136200 [Oryza sativa (japonica cultivar-group)]	PP			
1528	76302	4.00E -75	82	ref NP_001047161.1 Os02g0564500 [Oryza sativa (japonica cultivar-group)]	LL			
1529	76329	0	92	ref NP_014151.1 RNA polymerase I subunit A49; Rpa49p [Saccharomyces cerevisiae]	PEG	PP		

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1530	76353	0	97	ref NP_010937.1 Nucleotide binding alpha subunit of the heterotrimeric G protein that interacts with the receptor Gpr1p, has signaling role in response to nutrients; green fluorescent protein (GFP)-fusion protein localizes to the cell periphery; Gpa2p [Saccharomyces cerevisiae]	SS			
1531	76389	0	83	ref NP_014172.1 Co-chaperone that stimulates the ATPase activity of Ssa1p, required for a late step of ribosome biogenesis; associated with the cytosolic large ribosomal subunit; contains a J-domain; mutation causes defects in fluid-phase endocytosis; Jij1p [Saccharomyces cerevisiae]	CK			
1532	76378	1.00E-119	79	ref NP_014726.1 Repressor of G1 transcription that binds to SCB binding factor (SBF) at SCB target promoters in early G1; phosphorylation of Whi5p by the CDK, Cln3p/Cdc28p relieves repression and promoter binding by Whi5; periodically expressed in G1; Whi5p [Saccharomyces cerevisiae]	HS	PEG		
1533	76392	1.00E-59	55	ref NP_001045725.1 Os02g0122300 [Oryza sativa (japonica cultivar-group)]	LL			
1534	76321	0	76	ref NP_001058970.1 Os07g0166100 [Oryza sativa (japonica cultivar-group)]	LN			
1535	76345	0	86	ref NP_001043794.1 Os01g0665200 [Oryza sativa (japonica cultivar-group)]	DS			
1536	76381	0	88	ref NP_001058289.1 Os06g0663200 [Oryza sativa (japonica cultivar-group)]	HS			
1537	76358	0	76	ref NP_001050845.1 Os03g0666200 [Oryza sativa (japonica cultivar-group)]	LN			
1538	12629	0	97	ref NP_194110.2 unknown protein [Arabidopsis thaliana]	HS			
1539	76335	1.00E-101	66	ref NP_001065630.1 Os11g0127700 [Oryza sativa (japonica cultivar-group)]	LL			
1540	76372	0	82	gb EAY93387.1 hypothetical protein Osl_014620 [Oryza sativa (indica cultivar-group)]	PP			
1541	76396	0	57	ref NP_001047355.1 Os02g0602100 [Oryza sativa (japonica cultivar-group)]	DS			
1542	72314	0	92	ref NP_187714.1 steroid hormone receptor/ transcription factor [Arabidopsis thaliana]	HS			
1543	77840	1.00E-166	87	ref NP_001066296.1 Os12g0176500 [Oryza sativa (japonica cultivar-group)]	CS	CK		
1544	76886	1.00E-143	85	dbj BAC99390.1 putative U3 snoRNP protein IMP4 [Oryza sativa Japonica Group]	PP	PEG		
1545	76803	1.00E-163	90	gb EAY89818.1 hypothetical protein Osl_011051 [Oryza sativa (indica cultivar-group)]	LL			
1546	10302	0	97	ref NP_172288.1 ATNRT2:1 (Arabidopsis thaliana high affinity nitrate transporter 2.1); nitrate transporter	HS			
1547	71107	0	96	ref NP_030560.1 unknown protein [Arabidopsis thaliana]	HS			
1548	76887	0	86	ref NP_001044302.1 Os01g0758400 [Oryza sativa (japonica cultivar-group)]	HS	PP	PEG	
1549	77842	0	76	ref NP_001047779.1 Os02g0688500 [Oryza sativa (japonica cultivar-group)]	LN			
1550	76805	1.00E-165	85	dbj BAD08085.1 putative ornithine carbamoyltransferase [Oryza sativa Japonica Group]	HS			
1551	76817	2.00E-51	78	ref NP_001063927.1 Os09g0560500 [Oryza sativa (japonica cultivar-group)]	HS			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1552	77843	1.00E -68	66	ref NP_001053206.1 Os04g0497400 [Oryza sativa (japonica cultivar-group)]	CS			
1553	77844	0	94	ref NP_001105843.1 sialyltransferase like protein [Zea mays]	PP			
1554	76890	0	98	ref NP_001044001.1 Os01g0703600 [Oryza sativa (japonica cultivar-group)]	HS			
1555	77839	1.00E -155	74	ref NP_001047096.1 Os02g0550700 [Oryza sativa (japonica cultivar-group)]	DS	PP		
1556	76868	0	83	ref NP_001052270.1 Os04g0223000 [Oryza sativa (japonica cultivar-group)]	HS	PP	PEG	CS
1557	76821	1.00E -144	80	ref NP_001042197.1 Os01g0179200 [Oryza sativa (japonica cultivar-group)]	LN			
1558	76869	3.00E -98	73	gb EAY29350.1 hypothetical protein OsJ_012833 [Oryza sativa (japonica cultivar-group)]	HS	PP		
1559	76881	0	88	ref NP_001049165.1 Os03g0180700 [Oryza sativa (japonica cultivar-group)]	PP			
1560	76894	4.00E -33	94	ref NP_001066312.1 Os12g0180400 [Oryza sativa (japonica cultivar-group)]	LN			
1561	11930	0	100	dbj BAF01049.1 hypothetical protein [Arabidopsis thaliana]	SS			
1562	76848	2.00E -80	90	ref NP_001043542.1 Os01g0610100 [Oryza sativa (japonica cultivar-group)]	HS			
1563	76901	2.00E -44	85	gb EAY92619.1 hypothetical protein Osl_013852 [Oryza sativa (indica cultivar-group)]	LN			
1564	76914	1.00E -139	78	gb EAY79090.1 hypothetical protein Osl_033049 [Oryza sativa (indica cultivar-group)]	PP			
1565	76939	1.00E -137	71	gb EAY98958.1 hypothetical protein Osl_020191 [Oryza sativa (indica cultivar-group)]	CK			
1566	76987	1.00E -150	72	gb ABF93996.1 Integral membrane protein DUF6 containing protein, expressed [Oryza sativa (japonica cultivar-group)]	CK			
1567	76904	4.00E -58	76	gb EAY05371.1 hypothetical protein Osl_026603 [Oryza sativa (indica cultivar-group)]	HS	LN		
1568	76941	4.00E -26	69	gb EAY83408.1 hypothetical protein Osl_037367 [Oryza sativa (indica cultivar-group)]	HS			
1569	77113	0	58	ref NP_001064033.1 Os10g0110600 [Oryza sativa (japonica cultivar-group)]	PEG			
1570	77185	0	80	dbj BAD45738.1 thiF family protein-like [Oryza sativa Japonica Group]	DS			
1571	12911	0	100	ref NP_001078414.1 binding [Arabidopsis thaliana]	LN	HS		
1572	77127	0	89	ref NP_001062585.1 Os09g0115600 [Oryza sativa (japonica cultivar-group)]	HS	PEG		
1573	77177	0	85	gb EAY84671.1 hypothetical protein Osl_005904 [Oryza sativa (indica cultivar-group)]	PEG			
1574	77178	0	100	ref NP_441510.1 bifunctional 3,4-dihydroxy-2-butanone 4-phosphate synthase/GTP cyclohydrolase II protein [Synechocystis sp. PCC 6803]	HS			
1575	77179	2.00E -15	60	gb EAY74575.1 hypothetical protein Osl_002422 [Oryza sativa (indica cultivar-group)]	HS			
1576	77191	8.00E -61	81	ref NP_001046564.1 Os02g0282100 [Oryza sativa (japonica cultivar-group)]	PP			
1577	77145	2.00E -48	81	ref NP_001051289.1 Os03g0751000 [Oryza sativa (japonica cultivar-group)]	CK	PP		
1578	77169	1.00E -137	87	gb AAP50960.1 putative aurora-related kinase [Oryza sativa (japonica cultivar-group)]	PP			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1579	77110	5.00E -30	56	ref NP_001042508.1 Os01g0233000 [Oryza sativa (japonica cultivar-group)]	LL	LN		
1580	77146	0	100	ref NP_353311.1 bifunctional GMP synthase/glutamine amidotransferase protein [Agrobacterium tumefaciens str. C58]	PP			
1581	77170	7.00E -59	82	emb CAH67592.1 OSIGBa0092M08.4 [Oryza sativa (indica cultivar-group)]	DS			
1582	77171	0	95	ref NP_244906.1 potassium/proton antiporter [Bacillus halodurans C-125]	HS			
1583	77148	0	100	ref NP_661081.1 bifunctional GMP synthase/glutamine amidotransferase protein [Chlorobium tepidum TLS]	CK			
1584	11933	1.00E -149	94	ref NP_849952.1 GONST1 (GOLGI NUCLEOTIDE SUGAR TRANSPORTER 1) [Arabidopsis thaliana]	LN			
1585	77160	0	87	ref NP_414589.1 potassium:proton antiporter [Escherichia coli K12]	DS			
1586	77172	0	92	gb AAC77846.1 mannose-1-P guanosyltransferase [Escherichia coli]	LL			
1587	77184	0	92	ref NP_243321.1 flagellum-specific ATP synthase [Bacillus halodurans C-125]	HS			
1588	77225	0	94	ref NP_662474.1 3,4-dihydroxy-2-butanone 4-phosphate synthase/GTP cyclohydrolase II [Chlorobium tepidum TLS]	LL			
1589	77250	0	97	ref NP_599940.1 detergent sensitivity rescuer dtsR1 [Corynebacterium glutamicum ATCC 13032]	LN			
1590	77277	0	95	ref NP_001048987.1 Os03g0151800 [Oryza sativa (japonica cultivar-group)] ica cultivar-group]]	DS			
1591	77245	7.00E -45	64	gb EAY86480.1 hypothetical protein OsI_007713 [Oryza sativa (indica cultivar-group)]	SS			
1592	77246	4.00E -49	98	gb EAZ21400.1 hypothetical protein OsJ_004883 [Oryza sativa (japonica cultivar-group)]	HS			
1593	77462	0	75	ref NP_001065111.1 Os10g0525000 [Oryza sativa (japonica cultivar-group)]	CK			
1594	77414	0	82	ref NP_001046919.1 Os02g0506500 [Oryza sativa (japonica cultivar-group)]	HS			
1595	77439	0	82	ref NP_001045101.1 Os01g0899500 [Oryza sativa (japonica cultivar-group)]	PEG			
1596	77463	1.00E -106	89	ref NP_001048776.1 Os03g0119000 [Oryza sativa (japonica cultivar-group)]	HS			
1597	77488	0	80	gb EAT84267.1 hypothetical protein SNOG_07991 [Phaeosphaeria nodorum SN15]	DS			
1598	77441	5.00E -87	82	ref NP_001053804.1 Os04g0606900 [Oryza sativa (japonica cultivar-group)]	SS	PEG		
1599	77464	1.00E -164	90	gb EAT77502.1 hypothetical protein SNOG_14959 [Phaeosphaeria nodorum SN15]	PEG			
1600	77442	0	83	ref NP_013943.1 SR protein kinase (SRPK) involved in regulating proteins involved in mRNA metabolism and cation homeostasis; similar to human SRPK1; Sky1p [Saccharomyces cerevisiae]	HS			
1601	77419	0	100	ref NP_191311.1 ATSIP2 (ARABIDOPSIS THALIANA SEED IMBIBITION 2); hydrolase, hydrolyzing O-glycosyl compounds [Arabidopsis thaliana]	CK	HS		
1602	12444	1.00E -146	100	ref NP_569036.1 GAMMA CA3 (GAMMA CARBONIC ANHYDRASE 3); carbonate dehydratase [Arabidopsis thaliana]	LN			

PEP SEQ ID NO	Construct	Annotation			traits			
		E- value	% Id	Description				
1603	11791	0	87	ref NP_188619.2 phototropic-responsive NPH3 family protein [Arabidopsis thaliana]	LN			
1603	12445	0	87	ref NP_188619.2 phototropic-responsive NPH3 family protein [Arabidopsis thaliana]	HS			
1604	71249	0	96	ref NP_189251.1 CYP71B22 (cytochrome P450, family 71, subfamily B, polypeptide 22); oxygen binding [Arabidopsis thaliana]	HS			
1605	71237	0	95	ref NP_192740.1 short-chain dehydrogenase/reductase (SDR) family protein [Arabidopsis thaliana]	CK			
1606	12315	1.00E-97	100	ref NP_567911.1 unknown protein [Arabidopsis thaliana] gb AAM66960.1 unknown [Arabidopsis thaliana]	DS			
1606	12366	1.00E-97	100	ref NP_567911.1 unknown protein [Arabidopsis thaliana] gb AAM66960.1 unknown [Arabidopsis thaliana]	DS			

Trait Enhancement Screens

DS- Enhancement of drought tolerance identified by a soil drought stress tolerance

Drought or water deficit conditions impose mainly osmotic stress on plants. Plants are particularly vulnerable to drought during the flowering stage. The drought condition in the screening process disclosed in Example 1B started from the flowering time and was sustained to the end of harvesting. The present invention provides recombinant DNA that can improve the plant survival rate under such sustained drought condition. Exemplary recombinant DNA for conferring such drought tolerance are identified as such in Table 3. Such recombinant DNA can find particular use in generating transgenic plants that are tolerant to the drought condition imposed during flowering time and in other stages of the plant life cycle. As demonstrated from the model plant screen, in some embodiments of transgenic plants with trait-improving recombinant DNA grown under such sustained drought condition can also have increased total seed weight per plant in addition to the increased survival rate within a transgenic population, providing a higher yield potential as compared to control plants.

PEG-Enhancement of drought tolerance identified by PEG induced osmotic stress

tolerance screen: Various drought levels can be artificially induced by using various concentrations of polyethylene glycol (PEG) to produce different osmotic potentials (Pilon-Smits *et al.*, (1995) Plant Physiol. 107:125-130). Several physiological characteristics have been reported as being reliable indications for selection of plants possessing drought tolerance. These characteristics include the rate of seed germination and seedling growth. The traits can be assayed relatively easily by measuring the growth rate of seedling in PEG solution. Thus, a PEG-induced osmotic stress tolerance screen can be a surrogate for drought tolerance screen. As demonstrated from the model plant screen, embodiments of transgenic

plants with trait-improving recombinant DNA identified in the PEG-induced osmotic stress tolerance screen can survive better drought conditions providing a higher yield potential as compared to control plants.

SS-Enhancement of drought tolerance identified by high salinity stress tolerance

5 screen: Three different factors are responsible for salt damages: (1) osmotic effects, (2) disturbances in the mineralization process, and (3) toxic effects caused by the salt ions, *e.g.*, inactivation of enzymes. While the first factor of salt stress results in the wilting of the plants that is similar to drought effect, the ionic aspect of salt stress is clearly distinct from drought. The present invention provides genes that help plants maintain biomass, root growth, and/or
10 plant development in high salinity conditions, which are identified as such in Table 3. Since osmotic effect is one of the major components of salt stress, which is common to the drought stress, trait-improving recombinant DNA identified in a high salinity stress tolerance screen can also provide transgenic crops with enhanced drought tolerance. As demonstrated from the model plant screen, embodiments of transgenic plants with trait-improving recombinant DNA
15 identified in a high salinity stress tolerance screen can survive better drought conditions and/or high salinity conditions providing a higher yield potential as compared to control plants.

HS-Enhancement of drought tolerance identified by heat stress tolerance screen:

Heat and drought stress often occur simultaneously, limiting plant growth. Heat stress can cause
20 the reduction in photosynthesis rate, inhibition of leaf growth and osmotic potential in plants. Thus, genes identified by the present invention as heat stress tolerance conferring genes can also impart enhanced drought tolerance to plants. As demonstrated from the model plant screen, embodiments of transgenic plants with trait-improving recombinant DNA identified in a heat stress tolerance screen can survive better heat stress conditions and/or drought
25 conditions providing a higher yield potential as compared to control plants.

CK and CS-Enhancement of tolerance to cold stress:

Low temperature can immediately result in mechanical constraints, changes in activities of macromolecules, and reduced osmotic potential. In the present invention, two screening conditions, *e.g.*, cold shock tolerance screen (CK) and cold germination tolerance screen (CS), were set up to look for
30 transgenic plants that display visual growth advantage at lower temperature. In cold germination tolerance screen, the transgenic *Arabidopsis* plants were exposed to a constant temperature of 8°C from planting until day 28 post plating. The trait-improving recombinant DNA identified by such screen can be used for the production of transgenic plant that can germinate more robustly in a cold temperature as compared to the wild type plants. In cold

shock tolerance screen, the transgenic plants were first grown under the normal growth temperature of 22°C until day 8 post plating, and subsequently were placed under 8°C until day 28 post plating. As demonstrated from the model plant screen, embodiments of transgenic plants with trait-improving recombinant DNA identified in a cold shock stress tolerance screen and/or a cold germination stress tolerance screen can survive better cold conditions providing a higher yield potential as compared to control plants.

Enhancement of tolerance to multiple stresses: Different kinds of stresses often lead to identical or similar reaction in the plants. Genes that are activated or inactivated as a reaction to stress can either act directly in a way the genetic product reduces a specific stress, or they can act indirectly by activating other specific stress genes. By manipulating the activity of such regulatory genes, *e.g.*, multiple stress tolerance genes, the plant can be enabled to react to different kinds of stresses. For examples, PEP SEQ ID NO: 1000 can be used to enhance both salt stress tolerance and heat stress tolerance in plants. Of particular interest, plants transformed with PEP SEQ ID NO: 1495 can resist salt and heat stress. Plants transformed with PEP SEQ ID NO: 1483 can also improve growth in early stage and under heat stress. In addition to these multiple stress tolerance genes, the stress tolerance conferring genes provided by the present invention can be used in combinations to generate transgenic plants that can resist multiple stress conditions.

PP-Enhancement of early plant growth and development: It has been known in the art that to minimize the impact of disease on crop profitability, it is important to start the season with healthy and vigorous plants. This means avoiding seed and seedling diseases, leading to increased nutrient uptake and increased yield potential. Traditionally early planting and applying fertilizer are the methods used for promoting early seedling vigor. In early development stage, plant embryos establish only the basic root-shoot axis, a cotyledon storage organ(s), and stem cell populations, called the root and shoot apical meristems that continuously generate new organs throughout post-embryonic development. "Early growth and development" used herein encompasses the stages of seed imbibition through the early vegetative phase. The present invention provides genes that can be used to produce transgenic plants that have advantages in one or more processes including, but not limited to, germination, seedling vigor, root growth and root morphology under non-stressed conditions. The transgenic plants starting from a more robust seedling are less susceptible to the fungal and bacterial pathogens that attach germinating seeds and seedling. Furthermore, seedlings with advantage in root growth are more resistant to drought stress due to extensive and deeper root architecture. Therefore, it can be recognized by those skilled in the art that genes

conferring the growth advantage in early stages to plants can also be used to generate transgenic plants that are more resistant to various stress conditions due to enhanced early plant development. The present invention provides such exemplary recombinant DNA that confer both the stress tolerance and growth advantages to plants, identified as such in Table 3, *e.g.*, PEP SEQ ID NO: 1043 which can improve the plant early growth and development, and impart heat stress tolerance to plants. As demonstrated from the model plant screen, embodiments of transgenic plants with trait-improving recombinant DNA identified in the early plant development screen can grow better under non-stress conditions and/or stress conditions providing a higher yield potential as compared to control plants.

SP-Enhancement of late plant growth and development: "Late growth and development" used herein encompasses the stages of leaf development, flower production, and seed maturity. In certain embodiments, transgenic plants produced using genes that confer growth advantages to plants provided by the present invention, identified as such in Table 3, exhibit at least one phenotypic characteristics including, but not limited to, increased rosette radius, increased rosette dry weight, seed dry weight, silique dry weight, and silique length. On one hand, the rosette radius and rosette dry weight are used as the indexes of photosynthesis capacity, and thereby plant source strength and yield potential of a plant. On the other hand, the seed dry weight, silique dry weight and silique length are used as the indexes for plant sink strength, which are considered as the direct determinants of yield. As demonstrated from the model plant screen, embodiments of transgenic plants with trait-improving recombinant DNA identified in the late development screen can grow better and/or have enhanced development during leaf development and seed maturation providing a higher yield potential as compared to control plants.

LL-Enhancement of tolerance to shade stress identified in a low light screen: The effects of light on plant development are especially prominent at the seedling stage. Under normal light conditions with unobstructed direct light, a plant seedling develops according to a characteristic photomorphogenic pattern, in which plants have open and expanded cotyledons and short hypocotyls. Then the plant's energy is devoted to cotyledon and leaf development while longitudinal extension growth is minimized. Under low light condition where light quality and intensity are reduced by shading, obstruction or high population density, a seedling displays a shade-avoidance pattern, in which the seedling displays a reduced cotyledon expansion, and hypocotyls extension is greatly increased. As the result, a plant under low light condition increases significantly its stem length at the expense of leaf, seed or fruit and storage organ development, thereby adversely affecting of yield. The present

invention provides recombinant DNA that enable plants to have an attenuated shade avoidance response so that the source of plant can be contributed to reproductive growth efficiently, resulting higher yield as compared to the wild type plants. As demonstrated from the model plant screen, embodiments of transgenic plants with trait-improving recombinant DNA identified in a shade stress tolerance screen can have attenuated shade response under shade conditions providing a higher yield potential as compared to control plants. The transgenic plants generated by the present invention can be suitable for a higher density planting, thereby resulting increased yield per unit area.

LN-Enhancement of tolerance to low nitrogen availability stress

Nitrogen is a key factor in plant growth and crop yield. The metabolism, growth and development of plants are profoundly affected by their nitrogen supply. Restricted nitrogen supply alters shoot to root ratio, root development, activity of enzymes of primary metabolism and the rate of senescence (death) of older leaves. All field crops have a fundamental dependence on inorganic nitrogenous fertilizer. Since fertilizer is rapidly depleted from most soil types, it must be supplied to growing crops two or three times during the growing season. Enhanced nitrogen use efficiency by plants should enable crops cultivated under low nitrogen availability stress condition resulted from low fertilizer input or poor soil quality.

According to the present invention, transgenic plants generated using the recombinant nucleotides, which confer enhanced nitrogen use efficiency, identified as such in Table 3, exhibit one or more desirable traits including, but not limited to, increased seedling weight, greener leaves, increased number of rosette leaves, increased or decreased root length. One skilled in the art can recognize that the transgenic plants provided by the present invention with enhanced nitrogen use efficiency can also have altered amino acid or protein compositions, increased yield and/or better seed quality. The transgenic plants of the present invention can be productively cultivated under low nitrogen growth conditions, *e.g.*, nitrogen-poor soils and low nitrogen fertilizer inputs, which would cause the growth of wild type plants to cease or to be so diminished as to make the wild type plants practically useless. The transgenic plants also can be advantageously used to achieve earlier maturing, faster growing, and/or higher yielding crops and/or produce more nutritious foods and animal feedstocks when cultivated using nitrogen non-limiting growth conditions.

Stacked Traits: The present invention also encompasses transgenic plants with stacked engineered traits, *e.g.*, a crop having an enhanced phenotype resulting from expression of a trait-improving recombinant DNA, in combination with herbicide and/or pest resistance

traits. For example, genes of the current invention can be stacked with other traits of agronomic interest, such as a trait providing herbicide resistance, for example a RoundUp Ready® trait, or insect resistance, such as using a gene from *Bacillus thuringiensis* to provide resistance against lepidopteran, coliopteran, homopteran, hemiopteran, and other insects.

5 Explementary herbicides include glyphosate herbicides, phosphinothricin herbicides, oxynil herbicides, imidazolinone herbicides, dinitroaniline herbicides, pyridine herbicides, sulfonylurea herbicides, bialaphos herbicides, sulfonamide herbicides and gluphosinate herbicides. To illustrate that the production of transgenic plants with herbicide resistance is a capability of those of ordinary skill in the art, reference is made to U.S. patent application
10 publications 2003/0106096A1 and 2002/0112260A1 and U.S. Patents 5,034,322; 5,776,760, 6,107,549 and 6,376,754, all of which are incorporated herein by reference. To illustrate that the production of transgenic plants with pest resistance is a capability of those of ordinary skill in the art reference is made to U.S. Patents 5,250,515 and 5,880,275 which disclose plants expressing an endotoxin of *Bacillus thuringiensis* bacteria, to U.S. Patent
15 6,506,599 which discloses control of invertebrates which feed on transgenic plants which express dsRNA for suppressing a target gene in the invertebrate, to U.S. Patent 5,986,175 which discloses the control of viral pests by transgenic plants which express viral replicase, and to U.S. Patent Application Publication 2003/0150017 A1 which discloses control of pests by a transgenic plant which express a dsRNA targeted to suppressing a gene in the pest, all of
20 which are incorporated herein by reference.

Once one recombinant DNA has been identified as conferring an enhanced trait of interest in transgenic *Arabidopsis* plants, several methods are available for using the sequence of that recombinant DNA and knowledge about the protein it encodes to identify homologs of that sequence from the same plant or different plant species or other organisms, *e.g.*, bacteria
25 and yeast. Thus, in one aspect, the invention provides methods for identifying a homologous gene with a DNA sequence homologous to any of SEQ ID NO: 1 through SEQ ID NO: 803, or a homologous protein with an amino acid sequence homologous to any of SEQ ID NO: 804 to SEQ ID NO: 1606. In another aspect, the present invention provides the protein sequences of identified homologs for a sequence listed as SEQ ID NO: 1607 through SEQ ID
30 NO: 94613. In yet another aspect, the present invention also includes linking or associating one or more desired traits, or gene function with a homolog sequence provided herein.

The trait-improving recombinant DNA and methods of using such trait-improving recombinant DNA for generating transgenic plants with enhanced traits provided by the present invention are not limited to any particular plant species. Indeed, the plants according

to the present invention can be of any plant species, *e.g.*, can be monocotyledonous or dicotyledonous. In one embodiment, they are agricultural plants, *e.g.*, plants cultivated by man for purposes of food production or technical, particularly industrial applications. Of particular interest in the present invention are corn and soybean plants. The recombinant DNA constructs optimized for soybean transformation and recombinant DNA constructs optimized for corn transformation are provided by the present invention. Other plants of interest in the present invention for production of transgenic plants having enhanced traits include, without limitation, cotton, canola, wheat, sunflower, sorghum, alfalfa, barley, millet, rice, tobacco, fruit and vegetable crops, and turfgrass.

In certain embodiments, the present invention contemplates to use an orthologous gene in generating the transgenic plants with similarly enhanced traits as the transgenic *Arabidopsis* counterpart. Enhanced physiological properties in transgenic plants of the present invention can be confirmed in responses to stress conditions, for example in assays using imposed stress conditions to detect enhanced responses to drought stress, nitrogen deficiency, cold growing conditions, or alternatively, under naturally present stress conditions, for example under field conditions. Biomass measures can be made on greenhouse or field grown plants and can include such measurements as plant height, stem diameter, root and shoot dry weights, and, for corn plants, ear length and diameter.

Trait data on morphological changes can be collected by visual observation during the process of plant regeneration as well as in regenerated plants transferred to soil. Such trait data includes characteristics such as normal plants, bushy plants, taller plants, thicker stalks, narrow leaves, striped leaves, knotted phenotype, chlorosis, albino, anthocyanin production, or altered tassels, ears or roots. Other enhanced traits can be identified by measurements taken under field conditions, such as days to pollen shed, days to silking, leaf extension rate, chlorophyll content, leaf temperature, stand, seedling vigor, internode length, plant height, leaf number, leaf area, tillering, brace roots, stay green, stalk lodging, root lodging, plant health, barrenness/prolificacy, green snap, and pest resistance. In addition, trait characteristics of harvested grain can be confirmed, including number of kernels per row on the ear, number of rows of kernels on the ear, kernel abortion, kernel weight, kernel size, kernel density and physical grain quality.

To confirm hybrid yield in transgenic corn plants expressing genes of the present invention, it can be desirable to test hybrids over multiple years at multiple locations in a geographical location where maize is conventionally grown, *e.g.*, in Iowa, Illinois or other

locations in the midwestern United States, under “normal” field conditions as well as under stress conditions, *e.g.*, under drought or population density stress.

Transgenic plants can be used to provide plant parts according to the invention for regeneration or tissue culture of cells or tissues containing the constructs described herein.

5 Plant parts for these purposes can include leaves, stems, roots, flowers, tissues, epicotyl, meristems, hypocotyls, cotyledons, pollen, ovaries, cells and protoplasts, or any other portion of the plant which can be used to regenerate additional transgenic plants, cells, protoplasts or tissue culture. Seeds of transgenic plants are provided by this invention can be used to propagate more plants containing the trait-improving recombinant DNA constructs of this invention. These descendants are intended to be included in the scope of this invention if they
10 contain a trait-improving recombinant DNA construct of this invention, whether or not these plants are selfed or crossed with different varieties of plants.

The various aspects of the invention are illustrated by means of the following examples which are in no way intended to limit the full breath and scope of claims.

15

EXAMPLES

Example 1. Identification of recombinant DNA that confers enhanced trait(s) to plants

A. Plant expression constructs for *Arabidopsis* transformation

Each gene of interest was amplified from a genomic or cDNA library using primers
20 specific to sequences upstream and downstream of the coding region. Transformation vectors were prepared to constitutively transcribe DNA in either sense orientation (for enhanced protein expression) or anti-sense orientation (for endogenous gene suppression) under the control of an enhanced *Cauliflower Mosaic Virus* 35S promoter (U.S. patent 5,359,142) directly or indirectly (Moore, *et al.*, PNAS 95:376-381, 1998; Guyer, *e.g.*, Genetics 149: 633-
25 639, 1998; International patent application NO. PCT/EP98/07577). The transformation vectors also contain a *bar* gene as a selectable marker for resistance to glufosinate herbicide. The transformation of *Arabidopsis* plants was carried out using the vacuum infiltration method known in the art (Bethtold, *et al.*, Methods Mol. Biol. 82:259-66, 1998). Seeds
30 harvested from the plants, named as T1 seeds, were subsequently grown in a glufosinate-containing selective medium to select for plants which were actually transformed and which produced T2 transgenic seed.

B. Soil Drought tolerance screen

This example describes a soil drought tolerance screen to identify *Arabidopsis* plants transformed with recombinant DNA that wilt less rapidly and/or produce higher seed yield when grown in soil under drought conditions

5 T2 seeds were sown in flats filled with Metro/Mix® 200 (The Scotts® Company, USA). Humidity domes were added to each flat and flats were assigned locations and placed in climate-controlled growth chambers. Plants were grown under a temperature regime of 22°C at day and 20°C at night, with a photoperiod of 16 hours and average light intensity of 170 $\mu\text{mol}/\text{m}^2/\text{s}$. After the first true leaves appeared, humidity domes were
10 removed. The plants were sprayed with glufosinate herbicide and put back in the growth chamber for 3 additional days. Flats were watered for 1 hour the week following the herbicide treatment. Watering was continued every seven days until the flower bud primordia became apparent, at which time plants were watered for the last time.

To identify drought tolerant plants, plants were evaluated for wilting response and
15 seed yield. Beginning ten days after the last watering, plants were examined daily until 4 plants/line had wilted. In the next six days, plants were monitored for wilting response. Five drought scores were assigned according to the visual inspection of the phenotypes: 1 for healthy, 2 for dark green, 3 for wilting, 4 severe wilting, and 5 for dead. A score of 3 or higher was considered as wilted.

20 At the end of this assay, seed yield measured as seed weight per plant under the drought condition was characterized for the transgenic plants and their controls and analyzed as a quantitative response according to example 1M.

Two approaches were used for statistical analysis on the wilting response. First, the risk score was analyzed for wilting phenotype and treated as a qualitative response according to the
25 example 1L. Alternatively, the survival analysis was carried out in which the proportions of wilted and non-wilted transgenic and control plants were compared over each of the six days under scoring and an overall log rank test was performed to compare the two survival curves using S-PLUS statistical software (S-PLUS 6, Guide to statistics, Insightful, Seattle, WA, USA). A list of recombinant DNA constructs which improve drought tolerance in transgenic
30 plants is illustrated in Table 4

Table 4.

PEP SEQ ID	Construct ID	Orientation	Drought score		Seed yield	
			Delta mean	P- value	Delta mean	P- value

PEP SEQ ID	Construct ID	Orientation	Drought score		Seed yield	
			Delta mean	P- value	Delta mean	P- value
989	10471	SENSE	0.069	0.285	0.368	0.021
1224	12204	SENSE	0.152	0.018	-0.519	0.041
1606	12315	ANTI-SENSE	0.067	0.029	-0.126	0.210
1257	12354	ANTI-SENSE	0.118	0.334	-0.663	0.001
1606	12366	SENSE	-0.105	0.735	-0.072	0.403
852	13623	SENSE	0.309	0.005	-0.202	0.403
812	13638	ANTI-SENSE	0.195	0.047	-0.930	0.045
838	13811	SENSE	-0.037	0.729	-0.578	0.032
860	14248	SENSE	0.205	0.013	-0.484	0.066
885	14910	ANTI-SENSE	-0.087	0.305	-0.261	0.220
891	14932	ANTI-SENSE	0.222	0.011	-0.019	0.957
879	15111	ANTI-SENSE	0.082	0.241	0.624	0.019
893	15964	ANTI-SENSE	0.003	0.023	-0.396	0.028
899	15986	ANTI-SENSE	0.199	0.128	-0.022	0.773
900	15987	ANTI-SENSE	0.223	0.416	-0.156	0.576
904	15995	ANTI-SENSE	0.281	0.196	-0.362	0.157
945	16614	SENSE	0.136	0.001	0.059	0.440
959	19162	SENSE	0.007	0.887	0.348	0.019
999	19423	SENSE	0.222	0.024	-0.460	0.069
1003	19425	SENSE	0.002	0.556	0.099	0.019
1081	71667	SENSE	0.064	0.790	-0.314	0.101
1164	72717	SENSE	0.043	0.031	-0.173	0.221
1165	72718	SENSE	0.119	0.037	-0.272	0.359
1167	72779	SENSE	-0.026	0.491	0.395	0.006
1168	72791	SENSE	0.284	0.046	-1.867	0.009
1272	73472	SENSE	0.159	0.036	-0.223	0.511
1273	73526	SENSE	-0.219	0.141	-0.250	0.145
1280	73531	SENSE	-0.277	0.022	-0.097	0.740
1132	73738	SENSE	0.386	0.022	-1.025	0.018
1485	73938	SENSE	0.117	0.050	0.309	0.052
1069	74060	SENSE	0.116	0.016	0.195	0.200
1286	74164	SENSE	0.602	0.031	0.287	0.137
1157	74257	SENSE	-0.139	0.622	-0.366	0.059
1146	74269	SENSE	0.176	0.057	-0.101	0.273
1229	74376	SENSE	-0.023	0.583	-0.747	0.053
1234	74668	SENSE	0.112	0.018	-0.931	0.063
1351	74873	SENSE	0.256	0.006	0.110	0.627
1385	75473	SENSE	0.108	0.031	-1.469	0.015
1535	76345	SENSE	-0.024	0.476	0.318	0.017
1541	76396	SENSE	-0.146	0.218	-0.137	0.518
1245	76724	SENSE	0.514	0.029	-1.467	0.021
1581	77170	SENSE	0.258	0.026	-0.436	0.026
1570	77185	SENSE	0.180	0.037	-0.602	0.086
1457	77369	SENSE	-0.117	0.194	-0.324	0.096

PEP SEQ ID	Construct ID	Orientation	Drought score		Seed yield	
			Delta mean	P- value	Delta mean	P- value
1460	77372	SENSE	0.623	0.017	-0.108	0.028
1597	77488	SENSE	0.001	0.973	-0.252	0.075
1452	77594	SENSE	0.200	0.032	-0.334	0.096
1555	77839	SENSE	0.082	0.021	-0.242	0.298
1472	77965	SENSE	0.643	0.003	-0.157	0.456
1487	77990	SENSE	0.246	0.022	-0.303	0.458
1497	78023	SENSE	0.364	0.004	-3.072	0.012
1503	78036	SENSE	0.651	0.010	-0.959	0.095
1350	78115	SENSE	0.412	0.027	-0.024	0.880
1061	78321	SENSE	0.254	0.044	-0.941	0.014
1230	74377	SENSE	0.327	0.028	/	/
1007	70419	SENSE	0.007	0.047	-0.599	0.012

Transgenic plants comprising recombinant DNA expressing protein as set forth in SEQ ID NO: 1047, 1073, 1105, 1106, 1137, 1190, 1216, 1334, 1585, or 1590 showed enhanced drought tolerance by the second criteria as illustrated in Example 1L.

C. Heat stress tolerance screen

5 Under high temperatures, *Arabidopsis* seedlings become chlorotic and root growth is inhibited. This example sets forth the heat stress tolerance screen to identify *Arabidopsis* plants transformed with the gene of interest that are more resistant to heat stress based on primarily their seedling weight and root growth under high temperature.

10 T2 seeds were plated on 1/2 X MS salts, 1/% phytagel, with 10 µg/ml BASTA (7 per plate with 2 control seeds; 9 seeds total per plate). Plates were placed at 4°C for 3 days to stratify seeds. Plates were then incubated at room temperature for 3 hours and then held vertically for 11 additional days at temperature of 34°C at day and 20°C at night. Photoperiod was 16 h. Average light intensity was ~140 µmol/m²/s. After 14 days of growth, plants were scored for glufosinate resistance, root length, final growth stage, visual color, and seedling fresh weight.

15 A photograph of the whole plate was taken on day 14.

The seedling weight and root length were analyzed as quantitative responses according to example 1M. The final grow stage at day 14 was scored as success if 50% of the plants had reached 3 rosette leaves and size of leaves are greater than 1mm (Boyes, *et al.*, (2001) The Plant Cell 13, 1499-1510). The growth stage data was analyzed as a qualitative response according to example 1L. A list of recombinant DNA constructs that improve heat tolerance in transgenic plants illustrated in Table 5.

Table 5.

PEP SEQ ID	Construct ID	Orientation	Root length day 14		Growth stage at day 14		Seedling weight	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
848	10706	ANTI-SENSE	-0.080	0.577	-0.056	0.371	0.423	0.023
822	13231	SENSE	0.267	0.037	1.633	0.010	1.492	0.000
834	13914	SENSE	0.283	0.096	0.466	0.230	1.497	0.016
827	15606	SENSE	-0.107	0.459	0.032	0.889	0.658	0.049
876	15623	SENSE	0.265	0.088	0.150	0.371	1.064	0.012
906	16213	SENSE	0.020	0.917	0.048	0.579	0.681	0.013
943	16603	SENSE	0.171	0.300	0.783	0.192	0.994	0.035
941	16617	SENSE	0.327	0.055	0.179	0.312	1.362	0.007
934	17502	SENSE	-0.082	0.580	0.989	0.402	0.519	0.026
935	17503	SENSE	0.149	0.052	0.015	0.828	0.697	0.003
897	18203	SENSE	-0.078	0.640	-0.221	0.300	0.728	0.041
984	18238	SENSE	0.031	0.845	-0.077	0.457	0.820	0.047
853	18302	SENSE	0.315	0.074	1.298	0.269	1.271	0.011
851	18304	SENSE	0.346	0.222	0.374	0.381	1.111	0.043
952	18307	SENSE	0.394	0.183	1.676	0.287	1.088	0.013
997	18352	SENSE	0.239	0.055	0.036	0.894	0.924	0.019
968	18421	SENSE	0.069	0.684	0.941	0.213	0.918	0.012
969	18422	SENSE	0.236	0.243	/	/	1.127	0.033
958	18441	SENSE	0.175	0.289	0.832	0.497	0.907	0.043
979	18535	SENSE	-0.007	0.946	-0.100	0.496	0.853	0.015
911	19041	ANTI-SENSE	0.160	0.012	-0.063	0.102	0.932	0.003
913	19042	ANTI-SENSE	0.068	0.618	0.024	0.725	0.767	0.030
950	19154	SENSE	0.109	0.609	1.642	0.111	0.877	0.041
951	19155	SENSE	0.244	0.237	0.085	0.434	0.905	0.018
976	19237	SENSE	0.094	0.296	0.492	0.362	0.663	0.047
971	19244	SENSE	0.473	0.104	1.137	0.259	1.324	0.020
1011	19320	SENSE	-0.265	0.212	/	/	0.329	0.043
967	19534	SENSE	0.064	0.756	0.920	0.315	0.630	0.039
964	19535	SENSE	0.190	0.158	-0.025	0.750	0.961	0.013
970	19539	SENSE	0.083	0.373	0.521	0.256	0.466	0.031
973	19545	SENSE	-0.083	0.150	/	/	0.609	0.038
1002	19616	SENSE	0.288	0.094	-0.068	0.565	0.689	0.003
1027	19702	SENSE	-0.017	0.915	-0.033	0.798	0.783	0.043
1032	19755	SENSE	0.312	0.129	1.611	0.029	1.285	0.008
1029	19775	SENSE	0.126	0.362	2.093	0.172	0.775	0.004
1042	19786	SENSE	0.226	0.265	0.580	0.372	1.083	0.040
1039	19825	SENSE	0.053	0.662	-0.055	0.759	0.872	0.022
1037	19893	SENSE	-0.089	0.168	-0.130	0.249	0.787	0.023
1040	19919	SENSE	-0.214	0.277	0.198	0.435	0.814	0.050
1041	19979	SENSE	0.202	0.080	0.738	0.222	1.048	0.047
1031	19982	SENSE	0.241	0.111	0.933	0.263	1.127	0.025
1038	19983	SENSE	0.238	0.218	0.834	0.367	0.876	0.002
1028	19985	SENSE	0.216	0.057	0.414	0.306	0.860	0.001
1036	19987	SENSE	0.052	0.568	0.584	0.442	0.834	0.028
925	70102	SENSE	-0.013	0.877	-0.115	0.461	0.779	0.012

PEP SEQ ID	Construct ID	Orientation	Root length day 14		Growth stage at day 14		Seedling weight	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
926	70106	SENSE	0.038	0.667	-0.016	0.692	0.869	0.028
927	70113	SENSE	-0.163	0.289	1.098	0.222	1.005	0.012
930	70128	SENSE	0.025	0.785	0.347	0.302	0.920	0.016
931	70130	SENSE	0.238	0.111	-0.041	0.447	1.275	0.000
884	70216	SENSE	0.012	0.932	0.165	0.689	0.708	0.007
1436	70229	SENSE	-0.074	0.761	0.228	0.259	0.494	0.034
1446	70230	SENSE	0.068	0.053	-0.005	0.908	0.554	0.008
804	70244	SENSE	0.019	0.894	/	/	0.468	0.036
845	70253	SENSE	-0.080	0.196	-0.189	0.436	1.085	0.004
856	70412	SENSE	0.151	0.185	1.151	0.374	0.757	0.023
975	70425	SENSE	-0.056	0.712	/	/	0.815	0.011
1014	70433	SENSE	0.196	0.242	/	/	0.737	0.021
1015	70434	SENSE	0.136	0.106	/	/	0.814	0.000
1017	70436	SENSE	0.199	0.196	0.008	0.918	0.969	0.017
1019	70439	SENSE	0.188	0.127	0.846	0.311	1.250	0.012
1001	70603	SENSE	0.430	0.028	0.301	0.281	1.573	0.004
1052	70635	SENSE	-0.199	0.020	-0.185	0.234	0.381	0.045
1054	70638	SENSE	0.004	0.943	-0.069	0.197	0.788	0.036
1055	70642	SENSE	-0.100	0.336	-0.278	0.049	0.385	0.003
1056	70643	SENSE	0.160	0.445	-0.001	0.423	0.947	0.021
1078	70682	SENSE	-0.203	0.156	-0.278	0.057	0.404	0.024
920	70706	ANTI- SENSE	-0.354	0.066	-0.162	0.208	0.517	0.010
843	70716	ANTI- SENSE	0.098	0.240	0.013	0.970	0.858	0.001
837	70721	ANTI- SENSE	0.254	0.047	0.346	0.068	0.859	0.008
859	70723	ANTI- SENSE	0.180	0.341	1.110	0.253	0.990	0.016
1026	70741	SENSE	0.046	0.693	0.469	0.265	0.927	0.008
1093	70810	SENSE	0.168	0.053	-0.072	0.275	0.897	0.013
1008	70827	SENSE	-0.106	0.519	0.878	0.158	0.646	0.045
880	70832	SENSE	0.224	0.152	1.398	0.318	0.912	0.008
1263	70850	SENSE	-0.002	0.986	0.312	0.296	0.887	0.035
1043	70903	SENSE	0.065	0.734	0.623	0.454	0.776	0.002
1046	70936	SENSE	0.393	0.063	0.889	0.020	1.352	0.008
1044	70970	SENSE	0.335	0.231	0.520	0.444	1.429	0.031
1547	71107	SENSE	-0.257	0.177	0.027	0.685	0.528	0.045
1126	71125	SENSE	0.236	0.091	1.197	0.112	1.308	0.029
921	71145	SENSE	0.093	0.534	/	/	0.535	0.030
1604	71249	SENSE	-0.150	0.372	0.056	0.768	0.866	0.035
977	71303	SENSE	0.103	0.377	/	/	0.858	0.025
1067	71318	SENSE	0.100	0.212	0.498	0.324	1.190	0.004
957	71531	SENSE	-0.087	0.696	-0.085	0.224	0.718	0.048
1006	71536	SENSE	0.021	0.831	-0.134	0.725	0.922	0.047
1057	71564	SENSE	0.305	0.153	0.033	0.795	0.793	0.013
1091	71629	SENSE	0.084	0.202	1.387	0.044	1.001	0.019
1083	71696	SENSE	0.222	0.304	0.850	0.296	0.885	0.043
965	71719	SENSE	0.113	0.605	0.088	0.689	0.989	0.046

PEP SEQ ID	Construct ID	Orientation	Root length day 14		Growth stage at day 14		Seedling weight	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
1063	71808	SENSE	0.158	0.043	0.589	0.172	0.956	0.011
1143	72110	SENSE	-0.031	0.607	-0.114	0.565	0.329	0.041
1142	72124	SENSE	-0.211	0.255	-0.100	0.411	0.568	0.049
1542	72314	SENSE	0.127	0.271	1.270	0.224	0.943	0.002
1288	72340	SENSE	0.089	0.574	0.893	0.258	0.864	0.032
910	72427	ANTI- SENSE	-0.024	0.813	-0.019	0.423	0.658	0.006
1084	72475	SENSE	-0.129	0.192	0.024	0.896	0.697	0.013
1087	72476	SENSE	-0.136	0.228	-0.168	0.134	0.729	0.007
1098	72537	SENSE	-0.124	0.341	-0.254	0.011	0.352	0.025
1169	72732	SENSE	0.191	0.165	0.090	0.641	0.700	0.020
933	72748	SENSE	0.125	0.369	1.403	0.138	1.010	0.000
1158	72749	SENSE	0.053	0.642	0.625	0.250	0.881	0.027
1161	72775	SENSE	-0.050	0.707	-0.157	0.333	0.861	0.049
1163	72776	SENSE	0.091	0.358	1.128	0.239	1.244	0.006
1168	72791	SENSE	-0.095	0.672	/	/	0.763	0.034
1199	72909	SENSE	0.254	0.065	-0.033	0.348	0.818	0.048
1203	72958	SENSE	-0.030	0.774	-0.052	0.358	0.840	0.034
1195	73018	SENSE	0.053	0.638	0.057	0.895	0.856	0.042
1202	73036	SENSE	-0.005	0.957	0.282	0.386	0.783	0.007
1188	73074	SENSE	-0.146	0.202	-0.081	0.319	0.667	0.028
1193	73077	SENSE	0.028	0.770	0.495	0.459	0.944	0.001
1196	73112	SENSE	0.021	0.846	/	/	0.886	0.019
1180	73117	SENSE	-0.060	0.537	0.124	0.759	0.996	0.018
1178	73128	SENSE	0.149	0.075	0.248	0.598	0.934	0.038
1191	73147	SENSE	0.689	0.024	0.822	0.293	1.572	0.010
1174	73174	SENSE	0.305	0.027	0.532	0.480	0.779	0.046
1194	73183	SENSE	0.113	0.617	0.180	0.573	0.969	0.004
1175	73186	SENSE	0.101	0.628	1.230	0.130	0.714	0.014
1131	73248	SENSE	0.289	0.103	0.455	0.104	1.034	0.018
1118	73353	SENSE	0.108	0.504	0.947	0.218	0.864	0.024
1264	73429	SENSE	0.305	0.067	0.867	0.057	1.190	0.003
1267	73455	SENSE	0.133	0.064	0.299	0.033	0.559	0.018
1262	73474	SENSE	-0.433	0.105	-0.164	0.092	0.433	0.024
1274	73538	SENSE	-0.003	0.962	0.241	0.339	0.741	0.003
1279	73565	SENSE	0.182	0.325	0.650	0.525	1.279	0.021
1275	73574	SENSE	0.067	0.721	0.363	0.636	0.758	0.043
996	73607	SENSE	-0.070	0.388	0.401	0.673	0.636	0.024
896	73706	SENSE	0.064	0.744	0.294	0.124	1.082	0.012
1144	73916	SENSE	-0.004	0.980	0.495	0.265	1.060	0.024
903	73977	SENSE	0.230	0.065	0.096	0.370	0.596	0.014
1182	73990	SENSE	0.133	0.346	/	/	1.140	0.025
1291	74133	SENSE	-0.170	0.121	-0.027	0.729	0.635	0.013
1289	74155	SENSE	0.211	0.210	0.458	0.428	0.945	0.017
1148	74273	SENSE	0.050	0.756	0.694	0.467	0.837	0.026
1149	74280	SENSE	0.117	0.251	-0.103	0.271	0.785	0.002
1151	74287	SENSE	0.044	0.644	-0.148	0.027	0.928	0.000
1206	74333	SENSE	0.144	0.114	0.144	0.521	0.748	0.023

PEP SEQ ID	Construct ID	Orientation	Root length day 14		Growth stage at day 14		Seedling weight	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
1299	74431	SENSE	0.002	0.987	-0.190	0.282	0.674	0.012
1293	74485	SENSE	-0.007	0.920	-0.098	0.273	0.923	0.007
1435	74508	SENSE	0.409	0.104	0.221	0.394	0.825	0.001
1302	74525	SENSE	0.110	0.261	0.080	0.467	0.967	0.004
1227	74612	SENSE	-0.319	0.220	-0.591	0.063	0.563	0.017
1233	74623	SENSE	-0.167	0.140	/	/	0.521	0.000
1237	74638	SENSE	0.036	0.848	0.418	0.399	0.828	0.049
1209	74737	SENSE	0.004	0.959	0.877	0.414	0.972	0.008
1212	74747	SENSE	0.254	0.074	1.728	0.157	0.961	0.027
1355	74804	SENSE	0.615	0.095	1.179	0.314	1.685	0.019
1352	74826	SENSE	-0.062	0.460	/	/	0.655	0.009
1353	74850	SENSE	0.167	0.271	0.910	0.352	0.950	0.040
1359	74901	SENSE	0.117	0.664	0.258	0.353	0.897	0.041
1361	74904	SENSE	-0.030	0.814	-0.030	0.791	0.866	0.016
1365	74966	SENSE	-0.041	0.658	0.114	0.770	0.671	0.015
1360	74974	SENSE	0.142	0.382	-0.117	0.116	0.793	0.047
1130	75206	SENSE	0.044	0.484	-0.217	0.093	0.671	0.010
1254	75281	SENSE	0.207	0.020	0.046	0.795	0.827	0.009
1380	75323	SENSE	0.041	0.684	-0.202	0.074	0.473	0.006
1374	75330	SENSE	-0.005	0.712	-0.001	0.998	0.753	0.001
1379	75334	SENSE	0.427	0.107	0.050	0.599	1.040	0.004
1376	75366	SENSE	-0.013	0.883	/	/	0.832	0.008
1372	75389	SENSE	0.304	0.038	0.181	0.377	0.923	0.005
1378	75391	SENSE	0.076	0.434	0.109	0.434	0.918	0.020
1382	75401	SENSE	0.020	0.910	/	/	0.854	0.005
1392	75446	SENSE	0.431	0.174	0.231	0.269	0.967	0.021
1390	75454	SENSE	0.055	0.585	0.288	0.378	0.770	0.022
1384	75461	SENSE	0.032	0.434	0.020	0.936	0.931	0.034
1388	75464	SENSE	-0.117	0.288	0.085	0.651	0.717	0.014
1391	75493	SENSE	0.007	0.891	-0.021	0.738	1.080	0.013
1401	75512	SENSE	0.089	0.652	0.860	0.279	0.806	0.032
1409	75605	SENSE	0.153	0.136	-0.016	0.721	1.065	0.014
1413	75623	SENSE	0.151	0.321	-0.030	0.721	0.751	0.000
1403	75626	SENSE	0.364	0.001	0.619	0.475	1.127	0.014
1404	75674	SENSE	-0.028	0.530	/	/	0.893	0.021
1415	75695	SENSE	0.132	0.300	0.072	0.640	1.108	0.010
1419	75714	SENSE	-0.049	0.701	0.169	0.588	0.853	0.025
1308	75832	SENSE	0.043	0.229	-0.006	0.944	0.691	0.013
1318	75848	SENSE	0.117	0.485	0.324	0.491	0.953	0.035
1321	75852	SENSE	-0.218	0.196	-0.017	0.895	0.468	0.039
1330	75879	SENSE	0.126	0.570	-0.007	0.937	0.863	0.042
1331	75880	SENSE	0.198	0.119	0.331	0.249	0.842	0.011
1468	75908	SENSE	0.057	0.686	-0.064	0.759	0.916	0.015
1467	75919	SENSE	-0.063	0.562	-0.056	0.586	0.640	0.036
1471	75933	SENSE	0.126	0.099	0.431	0.238	0.905	0.004
1466	75940	SENSE	0.147	0.429	1.316	0.343	0.872	0.049
830	75958	SENSE	0.036	0.515	-0.144	0.102	1.048	0.001
826	75969	SENSE	0.164	0.316	0.002	0.990	0.696	0.025

PEP SEQ ID	Construct ID	Orientation	Root length day 14		Growth stage at day 14		Seedling weight	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
1470	75992	SENSE	0.103	0.525	0.101	0.605	0.853	0.005
832	76002	SENSE	-0.016	0.860	0.032	0.815	0.868	0.023
892	76017	SENSE	0.107	0.424	0.359	0.280	0.978	0.022
843	76039	SENSE	0.226	0.299	0.039	0.656	1.072	0.008
914	76053	SENSE	0.293	0.058	0.940	0.022	1.142	0.004
1215	76212	SENSE	0.006	0.732	0.105	0.829	0.808	0.012
1258	76236	SENSE	-0.062	0.693	0.941	0.453	0.871	0.031
1314	76261	SENSE	0.065	0.490	0.699	0.344	0.901	0.039
1526	76337	SENSE	0.121	0.439	0.454	0.199	0.924	0.010
1532	76378	SENSE	-0.077	0.500	0.220	0.481	0.829	0.034
1536	76381	SENSE	-0.011	0.954	0.712	0.324	0.806	0.004
1340	76462	SENSE	-0.019	0.861	-0.004	0.959	0.716	0.038
1221	76530	SENSE	0.260	0.305	-0.053	0.801	1.088	0.004
1342	76568	SENSE	-0.008	0.748	0.059	0.677	0.610	0.016
1337	76621	SENSE	-0.098	0.270	0.001	0.995	0.622	0.006
1346	76629	SENSE	0.154	0.603	0.550	0.509	0.982	0.049
1222	76716	SENSE	-0.007	0.945	/	/	0.792	0.024
1333	76745	SENSE	0.044	0.738	-0.010	0.927	0.695	0.021
1550	76805	SENSE	0.369	0.113	0.041	0.519	0.950	0.018
1551	76817	SENSE	0.126	0.645	0.007	0.790	0.861	0.024
1562	76848	SENSE	-0.080	0.699	-0.112	0.185	0.804	0.006
1558	76869	SENSE	0.345	0.160	1.378	0.130	1.205	0.030
1548	76887	SENSE	0.319	0.292	/	/	0.812	0.036
1554	76890	SENSE	0.226	0.007	-0.032	0.300	0.659	0.027
1567	76904	SENSE	0.178	0.268	/	/	0.917	0.004
1310	77034	SENSE	-0.033	0.843	0.479	0.109	0.953	0.001
1572	77127	SENSE	0.061	0.705	-0.026	0.704	0.769	0.043
1582	77171	SENSE	0.054	0.519	-0.133	0.026	0.698	0.000
1575	77179	SENSE	0.146	0.392	0.039	0.517	1.072	0.010
1587	77184	SENSE	0.025	0.782	/	/	0.979	0.012
1153	77304	SENSE	0.174	0.297	1.437	0.321	1.010	0.047
1223	77328	SENSE	0.116	0.253	/	/	1.016	0.024
1445	77345	SENSE	0.222	0.005	0.215	0.529	0.833	0.010
1447	77353	SENSE	0.124	0.184	0.043	0.702	0.830	0.007
1594	77414	SENSE	0.115	0.229	/	/	0.965	0.002
1601	77419	SENSE	0.309	0.205	0.762	0.199	1.166	0.013
1600	77442	SENSE	0.112	0.214	0.779	0.067	0.825	0.019
1596	77463	SENSE	-0.027	0.865	-0.143	0.007	0.674	0.045
1319	77515	SENSE	0.242	0.195	0.088	0.276	1.411	0.014
1332	77519	SENSE	0.667	0.043	1.824	0.173	1.142	0.021
1323	77523	SENSE	0.190	0.379	1.044	0.059	1.312	0.014
1426	77554	SENSE	0.358	0.257	1.126	0.108	0.949	0.034
1430	77560	SENSE	-0.047	0.095	0.786	0.380	0.583	0.012
1438	77567	SENSE	-0.178	0.220	/	/	0.437	0.020
1440	77578	SENSE	0.213	0.119	1.089	0.387	0.931	0.013
1270	77726	SENSE	-0.060	0.338	/	/	0.990	0.007
1368	77814	SENSE	-0.032	0.843	-0.117	0.556	0.885	0.025
1395	77818	SENSE	0.094	0.220	0.336	0.311	0.879	0.004

PEP SEQ ID	Construct ID	Orientation	Root length day 14		Growth stage at day 14		Seedling weight	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
1248	77903	SENSE	0.014	0.798	0.015	0.874	0.675	0.017
1443	77925	SENSE	0.005	0.960	0.085	0.472	0.911	0.001
1444	77927	SENSE	-0.167	0.181	-0.114	0.590	0.638	0.002
1456	77937	SENSE	0.231	0.151	0.815	0.062	1.368	0.001
1458	77940	SENSE	-0.017	0.859	/	/	0.706	0.004
1421	77954	SENSE	0.326	0.111	1.981	0.208	1.139	0.014
1422	77955	SENSE	-0.078	0.652	-0.062	0.104	0.529	0.029
1437	77959	SENSE	0.345	0.055	0.505	0.429	1.016	0.038
1474	77972	SENSE	0.101	0.481	0.125	0.606	1.030	0.008
1495	78019	SENSE	0.030	0.321	0.320	0.481	0.931	0.034
1429	78128	SENSE	0.341	0.052	0.016	0.764	1.007	0.021
1424	78131	SENSE	0.042	0.138	0.018	0.829	0.687	0.005
1499	78161	SENSE	0.156	0.202	0.170	0.418	0.898	0.000
1522	78191	SENSE	0.177	0.313	0.964	0.512	0.854	0.020
1061	78321	SENSE	0.196	0.245	-0.007	0.773	1.013	0.048
1110	78325	SENSE	0.046	0.345	0.123	0.609	0.643	0.007
1232	78364	SENSE	0.054	0.615	0.837	0.243	0.475	0.038
1239	78365	SENSE	0.083	0.384	/	/	1.153	0.030
948	78370	SENSE	0.016	0.854	0.310	0.456	0.826	0.036
1000	78373	SENSE	0.018	0.774	0.595	0.263	1.046	0.009
1482	78538	SENSE	0.123	0.332	1.152	0.342	0.921	0.024
1490	78543	SENSE	-0.003	0.979	/	/	0.934	0.008
1513	78566	SENSE	0.131	0.429	0.479	0.532	0.820	0.017
1523	78595	SENSE	0.182	0.295	0.005	0.938	0.766	0.014
1520	78632	SENSE	0.394	0.058	0.997	0.314	1.143	0.017
1459	78746	SENSE	0.192	0.256	1.115	0.346	1.025	0.013
1455	78904	SENSE	0.150	0.222	0.970	0.084	0.946	0.030
1483	78922	SENSE	0.162	0.307	2.327	0.024	1.156	0.007
1341	78987	SENSE	-0.034	0.626	0.689	0.246	0.828	0.005
1434	78994	SENSE	0.282	0.000	2.384	0.009	0.940	0.002

Transgenic plants comprising recombinant DNA expressing protein as set forth in SEQ ID NO: 823, 825, 873, 886, 912, 916, 919, 928, 937, 942, 972, 986, 995, 1074, 1107, 1134, 1136, 1145, 1157, 1173, , 1220, 1228, 1242, 1243, 1285, 1377, 1469, 1486, 1489, 1524, 1538, 1546, 1556, 1568, 1571, 1574, 1592, or 1603 showed enhanced heat stress tolerance by the second criteria as illustrated in Example 1L and 1M.

D. Salt stress tolerance screen

This example sets forth the high salinity stress screen to identify *Arabidopsis* plants transformed with the gene of interest that are tolerant to high levels of salt based on their rate of development, root growth and chlorophyll accumulation under high salt conditions.

T2 seeds were plated on glufosinate selection plates containing 90 mM NaCl and grown under standard light and temperature conditions. All seedlings used in the experiment

were grown at a temperature of 22°C at day and 20°C at night, a 16-hour photoperiod, an average light intensity of approximately 120 $\mu\text{mol}/\text{m}^2$. On day 11, plants were measured for primary root length. After 3 more days of growth (day 14), plants were scored for transgenic status, primary root length, growth stage, visual color, and the seedlings were pooled for fresh weight measurement. A photograph of the whole plate was also taken on day 14.

The seedling weight and root length were analyzed as quantitative responses according to example 1M. The final growth stage at day 14 was scored as success if 50% of the plants reached 3 rosette leaves and size of leaves are greater than 1mm (Boyce, D.C., *et al.*, (2001), *The Plant Cell* 13, 1499/1510). The growth stage data was analyzed as a qualitative response according to example 1L. A list of recombinant DNA constructs that improve high salinity tolerance in transgenic plants illustrated in Table 6.

Table 6.

PEP SEQ ID	Orientation	Root length at day 11		Root length at day 14		Growth stage at day 14		Seedling weight at day 14	
		Delta mean	P-value	Delta mean	P-value	Delta mean	P-value	Delta mean	P-value
932	ANTI- SENSE	0.305	0.019	0.360	0.093	0.583	0.428	0.835	0.050
955	SENSE	0.028	0.336	0.085	0.047	-0.038	0.866	0.060	0.036
1093	SENSE	0.097	0.010	0.046	0.488	0.605	0.172	0.166	0.016
1139	SENSE	0.187	0.011	0.092	0.015	-0.038	0.923	-0.002	0.987
1183	SENSE	0.251	0.118	0.328	0.016	/	/	0.404	0.148
1106	SENSE	0.093	0.383	0.169	0.030	0.370	0.357	0.215	0.053
896	SENSE	0.349	0.000	0.270	0.069	0.057	0.797	0.421	0.047
1416	SENSE	0.062	0.516	0.081	0.035	-0.159	0.513	-0.076	0.670
1591	SENSE	0.034	0.592	0.051	0.003	0.255	0.623	0.194	0.052
1598	SENSE	0.227	0.004	0.224	0.002	1.203	0.316	0.281	0.258
1425	SENSE	-0.107	0.327	-0.028	0.724	-0.215	0.425	0.384	0.029
1384	SENSE	-0.088	0.195	-0.0557	0.531	-0.009	0.980	0.101	0.025

Transgenic plants comprising recombinant DNA expressing protein as set forth in SEQ ID NO: 821, 834, 890, 919, 946, 961, 997, 1013, 1080, 1101, 1147, 1160, 1181, 1197, 1200, 1220, 1237, 1248, 1264, 1300, 1313, 1355, 1393, 1397, 1406, 1467, 1496, 1514, 1530 or 1561 showed enhanced salt stress tolerance by the second criteria as illustrated in Example 1L and 1M.

E. Polyethylene Glycol (PEG) induced osmotic stress tolerance screen

There are numerous factors, which can influence seed germination and subsequent seedling growth, one being the availability of water. Genes, which can directly affect the

success rate of germination and early seedling growth, are usable agronomic traits for improving the germination and growth of crop plants under drought stress. In this assay, PEG was used to induce osmotic stress on germinating transgenic lines of *Arabidopsis thaliana* seeds in order to screen for osmotically resistant seed lines.

5 T2 seeds were plated on BASTA selection plates containing 3% PEG and grown under standard light and temperature conditions. Seeds were plated on each plate containing 3% PEG, 1/2 X MS salts, 1% phytigel, and 10 µg/ml glufosinate. Plates were placed at 4°C for 3 days to stratify seeds. On day 11, plants were measured for primary root length. After 3 more days of growth, *e.g.*, at day 14, plants were scored for transgenic status, primary root
10 length, growth stage, visual color, and the seedlings were pooled for fresh weight measurement. A photograph of the whole plate was taken on day 14.

Seedling weight and root length were analyzed as quantitative responses according to example 1M. The final growth stage at day 14 was scored as success or failure based on whether the plants reached 3 rosette leaves and size of leaves are greater than 1mm. The
15 growth stage data was analyzed as a qualitative response according to example 1L. A list of recombinant DNA constructs that improve osmotic stress tolerance in transgenic plants illustrated in Table 7.

Table 7.

PEP Seq ID	Orientation	Root length at day 11		Root length at day 14		Growth stage at day 14		Seedling weight at day 14	
		Delta mean	P- value	Delta mean	P-value	Delta mean	P- value	Delta mean	P- value
1019	SENSE	0.003	0.883	0.068	0.022	0.113	0.715	-0.139	0.215
960	SENSE	0.139	0.278	0.065	0.010	2.264	0.322	0.197	0.256
1241	SENSE	0.061	0.690	0.096	0.044	1.011	0.569	-0.095	0.666
1119	SENSE	0.002	0.978	-0.107	0.415	2.760	0.156	0.156	0.000
1391	SENSE	0.602	0.012	0.533	0.044	/	/	0.866	0.005
1322	SENSE	0.062	0.041	0.150	0.020	/	/	-0.132	0.218
1238	SENSE	0.099	0.387	0.118	0.004	2.693	0.175	-0.187	0.005
1556	SENSE	0.153	0.116	0.271	0.081	/	/	0.331	0.019
1153	SENSE	0.063	0.067	0.081	0.134	2.964	0.104	0.280	0.020
1218	SENSE	0.040	0.067	-0.089	0.170	0.492	0.806	0.198	0.030
1599	SENSE	0.099	0.168	0.103	0.237	/	/	0.292	0.031
1478	SENSE	0.216	0.115	0.197	0.041	1.353	0.414	0.234	0.388
1512	SENSE	-0.167	0.071	-0.230	0.141	-1.066	0.090	0.085	0.012
1572	SENSE	0.447	0.005	0.352	0.025	4	/	0.656	0.046

20 Transgenic plants comprising recombinant DNA expressing protein as set forth in SEQ ID NO: 832, 833, 871, 874, 931, 974, 985, 997, 1021, 1024, 1031, 1043, 1053, 1059, 1060, 1091, 1111, 1115, 1118, 1120, 1124, 1127, 1128, 1159, 1172, 1179, 1185, 1197, 1213,

1226, 1230, 1244, 1253, 1265, 1306, 1320, 1327, 1354, 1355, 1357, 1362, 1367, 1381, 1395, 1398, 1399, 1407, 1462, 1494, 1499, 1500, 1523, 1529, 1532, 1544, 1548, 1569, 1572, 1573, 1595, or 1598 showed enhanced PEG osmotic stress tolerance by the second criteria as illustrated in Example 1L and 1M.

5 F. Cold shock tolerance screen

This example set forth a screen to identify *Arabidopsis* plants transformed with the genes of interest that are more tolerant to cold stress subjected during day 8 to day 28 after seed planting. During these crucial early stages, seedling growth and leaf area increase were measured to assess tolerance when *Arabidopsis* seedlings were exposed to low temperatures.

10 Using this screen, genetic alterations can be found that enable plants to germinate and grow better than wild type plants under sudden exposure to low temperatures.

Eleven seedlings from T2 seeds of each transgenic line plus one control line were plated together on a plate containing ½ X Gamborg Salts with 0.8 Phytigel™, 1% Phytigel, and 0.3% Sucrose. Plates were then oriented horizontally and stratified for three days at 4°C. At day three, plates were removed from stratification and exposed to standard conditions (16 hr photoperiod, 22°C at day and 20°C at night) until day 8. At day eight, plates were removed from standard conditions and exposed to cold shock conditions (24 hr photoperiod, 8°C at both day and night) until the final day of the assay, *e.g.*, day 28. Rosette areas were measured at day 8 and day 28, which were analyzed as quantitative responses according to
20 example 1M. A list of recombinant nucleotides that improve cold shock stress tolerance in plants illustrated in Table 8.

Table 8.

PEP SEQ ID	Construct ID	Orientation	Rosette area at day 8		Rosette area at day 28		Rosette area difference	
			Delta mean	P-value	Risk score mean	P-value	Delta mean	P-value
814	72361	SENSE	0.172	0.712	0.657	0.000	0.757	0.001
816	71626	SENSE	0.634	0.013	0.456	0.032	0.513	0.003
861	13965	ANTI- SENSE	0.414	0.077	0.248	0.117	0.274	0.012
915	16120	SENSE	0.494	0.247	0.758	0.042	0.722	0.077
918	72417	SENSE	0.984	0.041	0.125	0.023	0.114	0.051
923	71706	SENSE	0.010	0.943	0.231	0.064	0.459	0.019
962	70808	SENSE	0.394	0.285	0.290	0.014	0.362	0.072
974	72356	SENSE	0.570	0.057	0.823	0.044	0.888	0.096
1000	78373	SENSE	-0.020	0.948	0.846	0.006	0.923	0.005
1012	73613	SENSE	0.574	0.024	0.579	0.010	0.570	0.012
1022	70447	SENSE	0.294	0.369	0.195	0.015	0.106	0.089

1023	71725	SENSE	0.201	0.492	0.199	0.026	0.225	0.055
1085	71677	SENSE	0.596	0.003	0.741	0.021	0.425	0.226
1097	72510	SENSE	0.813	0.250	0.891	0.009	1.105	0.015
1125	73221	SENSE	0.582	0.016	0.412	0.027	0.425	0.117
1141	72092	SENSE	0.092	0.750	0.568	0.006	0.480	0.027
1151	74287	SENSE	0.692	0.065	0.795	0.017	0.774	0.015
1153	77304	SENSE	0.664	0.194	1.164	0.014	1.479	0.014
1156	74719	SENSE	0.350	0.133	0.393	0.083	0.449	0.013
1171	73058	SENSE	0.500	0.180	0.322	0.027	0.329	0.049
1186	73131	SENSE	0.369	0.286	0.725	0.031	0.907	0.052
1187	73108	SENSE	0.579	0.065	0.725	0.018	0.644	0.025
1198	10908	ANTI-SENSE	-0.549	0.086	0.112	0.095	0.136	0.050
1205	74732	SENSE	0.348	0.082	0.457	0.059	0.513	0.019
1239	78365	SENSE	0.430	0.141	0.621	0.035	0.600	0.043
1307	75830	SENSE	0.865	0.113	1.430	0.003	1.558	0.002
1315	78456	SENSE	-0.352	0.118	0.534	0.077	0.650	0.049
1335	76746	SENSE	0.552	0.112	0.508	0.071	0.734	0.041
1343	76176	SENSE	0.435	0.095	1.337	0.018	1.472	0.018
1350	78115	SENSE	0.136	0.464	0.413	0.016	0.421	0.018
1375	75342	SENSE	1.019	0.029	1.486	0.039	1.752	0.046
1408	75652	SENSE	-0.425	0.454	0.388	0.005	0.415	0.009
1409	75605	SENSE	0.424	0.100	0.703	0.041	0.808	0.010
1420	75762	SENSE	1.078	0.039	0.558	0.037	0.480	0.011
1430	77560	SENSE	0.378	0.128	0.261	0.028	0.288	0.041
1431	77561	SENSE	-0.191	0.296	0.446	0.054	0.511	0.047
1439	11749	ANTI-SENSE	-0.458	0.494	0.622	0.028	0.710	0.008
1458	77940	SENSE	0.498	0.200	0.305	0.167	0.573	0.009
1461	78126	SENSE	-0.708	0.080	0.426	0.046	0.485	0.071
1500	78162	SENSE	1.502	0.000	1.496	0.025	1.778	0.024
1502	78622	SENSE	0.550	0.221	0.722	0.018	0.752	0.016
1517	78571	SENSE	1.529	0.033	1.204	0.001	1.303	0.001
1519	78573	SENSE	0.292	0.457	0.603	0.007	0.542	0.038
1543	77840	SENSE	-0.182	0.173	0.232	0.030	0.299	0.020
1565	76939	SENSE	0.598	0.154	0.520	0.037	0.597	0.035
1566	76987	SENSE	0.953	0.052	0.845	0.010	0.936	0.007
1577	77145	SENSE	0.408	0.180	0.403	0.041	0.368	0.140
1593	77462	SENSE	1.157	0.052	1.077	0.012	0.969	0.003
1601	77419	SENSE	0.156	0.513	0.763	0.010	0.844	0.031
1605	71237	SENSE	0.836	0.009	0.825	0.005	1.041	0.007
845	70253	SENSE	0.578	0.007	0.571	0.064	0.459	0.007
1007	70419	SENSE	-0.079	0.646	0.197	0.126	0.266	0.012
1313	77511	SENSE	0.822	0.011	0.131	0.013	0.063	0.189
1332	77519	SENSE	0.275	0.286	0.911	0.034	1.020	0.031
1484	77984	SENSE	-0.104	0.724	0.255	0.055	0.274	0.048

Transgenic plants comprising recombinant DNA expressing protein as set forth in SEQ ID NO: 818, 836, 844, 848, 883, 1029, 1045, 1062, 1082, 1277, 1287, 1294, 1305, 1318, 1366, 1369, 1373, 1387, 1402, 1405, 1410, 1412, 1418, 1427, 1433, 1476, 1481, 1495, 1531, or

1583 showed enhanced cold stress tolerance by the second criterial as illustrated in Example 1L and 1M.

G. Cold germination tolerance screen

This example sets forth a screen to identify *Arabidopsis* plants transformed with the genes of interests are resistant to cold stress based on their rate of development, root growth and chlorophyll accumulation under low temperature conditions.

T2 seeds were plated and all seedlings used in the experiment were grown at 8°C. Seeds were first surface disinfested using chlorine gas and then seeded on assay plates containing an aqueous solution of 1/2 X Gamborg's B/5 Basal Salt Mixture (Sigma/Aldrich Corp., St. Louis, MO, USA G/5788), 1% Phytigel™ (Sigma-Aldrich, P-8169), and 10 ug/ml glufosinate with the final pH adjusted to 5.8 using KOH. Test plates were held vertically for 28 days at a constant temperature of 8°C, a photoperiod of 16 hr, and average light intensity of approximately 100 umol/m²/s. At 28 days post plating, root length was measured, growth stage was observed, the visual color was assessed, and a whole plate photograph was taken.

The root length at day 28 was analyzed as a quantitative response according to example 1M. The growth stage at day 7 was analyzed as a qualitative response according to example 1L. A list of recombinant DNA constructs that improve cold stress tolerance in transgenic plants illustrated in Table 9.

Table 9.

NUC Seq ID NO	PEP SEQ ID NO	Construct ID	Nomination ID	Orientation	Root length at day 28	P-value	Growth stage at day 28	
					Delta mean		Delta mean	P- value
187	990	18331	CGPG3338	SENSE	/	/	/	/
227	1030	19829	CGPG3981	SENSE	-0.021	0.389	0.597	0.031
231	1034	70365	CGPG4028	SENSE	-0.088	0.312	0.725	0.013
214	1017	70436	CGPG3703	SENSE	0.051	0.747	1.952	0.005
217	1020	70446	CGPG3730	SENSE	-0.064	0.784	1.572	0.009
159	962	70808	CGPG289	SENSE	0.090	0.637	2.318	0.003
262	1065	71313	CGPG4401	SENSE	0.160	0.542	1.654	0.001
272	1075	71339	CGPG4551	SENSE	/	/	/	/
283	1086	71691	CGPG4656	SENSE	-0.048	0.305	1.121	0.003
335	1138	72016	CGPG5246	SENSE	0.385	0.012	1.855	0.161
337	1140	72044	CGPG5268	SENSE	0.442	0.044	1.908	0.003
359	1162	72728	CGPG5530	SENSE	0.243	0.300	2.142	0.002
137	940	72796	CGPG2451	SENSE	0.187	0.063	0.775	0.014
92	895	72952	CGPG2095	SENSE	/	/	/	/
399	1202	73036	CGPG5796	SENSE	0.246	0.127	2.235	0.001
64	867	73075	CGPG1797	SENSE	0.046	0.788	1.242	0.007
377	1180	73117	CGPG5657	SENSE	0.384	0.033	1.764	0.001
386	1189	73132	CGPG5705	SENSE	0.208	0.283	2.646	0.003

NUC Seq ID NO	PEP SEQ ID NO	Construct ID	Nomination ID	Orientation	Root length at day 28	P-value	Growth stage at day 28	
					Delta mean		Delta mean	P- value
374	1177	73139	CGPG5641	SENSE	0.040	0.536	1.183	0.002
473	1276	73539	CGPG6475	SENSE	0.004	0.967	1.052	0.002
482	1285	74104	CGPG6566	SENSE	0.107	0.490	1.060	0.003
428	1231	74381	CGPG6111	SENSE	0.060	0.631	0.457	0.049
632	1435	74508	CGPG8	SENSE	0.323	0.009	1.197	0.387
433	1236	74635	CGPG6147	SENSE	0.165	0.172	1.596	0.006
438	1241	74660	CGPG6177	SENSE	0.358	0.007	1.518	0.001
571	1374	75330	CGPG7508	SENSE	0.019	0.906	0.570	0.026
589	1392	75446	CGPG7637	SENSE	/	/	/	/
598	1401	75512	CGPG7752	SENSE	0.057	0.576	1.652	0.001
595	1398	75582	CGPG7742	SENSE	0.213	0.374	1.883	0.003
506	1309	75836	CGPG6827	SENSE	0.228	0.076	1.314	0.041
521	1324	75863	CGPG6942	SENSE	0.162	0.132	1.468	0.002
523	1326	75865	CGPG6949	SENSE	0.046	0.615	1.742	0.006
660	1463	75901	CGPG8211	SENSE	0.437	0.015	1.733	0.070
59	862	75912	CGPG1726	SENSE	-0.035	0.550	0.265	0.000
37	840	76050	CGPG1463	SENSE	-0.146	0.404	0.213	0.043
408	1211	76113	CGPG5862	SENSE	0.135	0.014	0.615	0.453
544	1347	76189	CGPG7253	SENSE	0.015	0.018	0.333	0.645
722	1525	76325	CGPG8870	SENSE	-0.141	0.553	0.557	0.026
447	1250	76524	CGPG6266	SENSE	0.099	0.300	1.123	0.015
446	1249	76653	CGPG6254	SENSE	0.507	0.101	3.257	0.004
628	1431	77561	CGPG7972	SENSE	0.278	0.061	1.265	0.009
647	1450	77588	CGPG8108	SENSE	0.063	0.677	1.028	0.032
749	1552	77843	CGPG9017	SENSE	0.382	0.011	1.325	0.006
701	1504	78043	CGPG8606	SENSE	0.237	0.204	0.859	0.003
91	894	17227	CGPG2077	SENSE	0.071	0.621	0.820	0.006
319	1122	73318	CGPG5029	SENSE	0.004	0.989	0.532	0.011
465	1268	73432	CGPG6421	SENSE	-0.038	0.743	0.532	0.011
568	1371	77808	CGPG7499	SENSE	0.170	0.037	1.138	0.023
312	1115	72814	CGPG4982	SENSE	0.080	0.592	1.467	0.013
461	1264	73429	CGPG6397	SENSE	0.320	0.206	1.589	0.017
682	1485	73938	CGPG85	SENSE	0.029	0.927	1.140	0.005
680	1483	78922	CGPG8489	SENSE	0.260	0.123	1.326	0.004

Transgenic plants comprising recombinant DNA expressing protein as set forth in 826, 863, 864, 941, 985, 1071, 1123, 1170, 1194, 1259, 1312, 1339, 1433, 1465, 1543, 1556 showed enhanced cold stress tolerance by the second criterial as illustrated in Example 1L and 1M.

H. shade tolerance screen

Plants undergo a characteristic morphological response in shade that includes the elongation of the petiole, a change in the leaf angle, and a reduction in chlorophyll content. While these changes can confer a competitive advantage to individuals, in a monoculture the

shade avoidance response is thought to reduce the overall biomass of the population. Thus, genetic alterations that prevent the shade avoidance response can be associated with higher yields. Genes that favor growth under low light conditions can also promote yield, as inadequate light levels frequently limit yield. This protocol describes a screen to look for *Arabidopsis* plants that show an attenuated shade avoidance response and/or grow better than control plants under low light intensity. Of particular interest, we were looking for plants that didn't extend their petiole length, had an increase in seedling weight relative to the reference and had leaves that were more close to parallel with the plate surface.

T2 seeds were plated on glufosinate selection plates with ½ MS medium. Seeds were sown on 1/2 X MS salts, 1% Phytigel, 10 ug/ml BASTA. Plants were grown on vertical plates at a temperature of 22°C at day, 20°C at night and under low light (approximately 30 uE/m²/s, far/red ratio (655/665/725/735) ~0.35 using PLAQ lights with GAM color filter #680). Twenty-three days after seedlings were sown, measurements were recorded including seedling status, number of rosette leaves, status of flower bud, petiole leaf angle, petiole length, and pooled fresh weights. A digital image of the whole plate was taken on the measurement day. Seedling weight and petiole length were analyzed as quantitative responses according to example 1M. The number of rosette leaves, flowering bud formation and leaf angel were analyzed as qualitative responses according to example 1L.

A list of recombinant DNA constructs that improve shade tolerance in plants illustrated in Table 10.

Table 10.

NUC Seq ID	PEP SEQ ID	Construct ID	Orientation	Seedling weight at day 23		Petiole length at day 23	
				Delta mean	P-value	Delta mean	P-value
52	855	15624	SENSE	0.327	0.041	0.141	0.189
398	1201	72922	SENSE	0.421	0.049	0.339	0.070
427	1230	74377	SENSE	0.411	0.039	0.374	0.067
405	1208	74736	SENSE	0.268	0.024	0.147	0.141
566	1369	75387	SENSE	0.329	0.008	0.342	0.010
59	862	75912	SENSE	0.470	0.035	0.572	0.058
664	1467	75919	SENSE	0.453	0.007	0.114	0.063
736	1539	76335	SENSE	-1.012	0.081	-0.594	0.007
783	1586	77172	SENSE	0.426	0.044	0.218	0.164
698	1501	78033	SENSE	0.060	0.033	-0.129	0.612

For "seedling weight", if p<0.05 and delta or risk score mean >0, the transgenic plants showed statistically significant trait enhancement as compared to the reference. If

$p < 0.2$ and delta or risk score mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference with $p < 0.2$.

For “petiole length”, if $p < 0.05$ and delta < 0 , the transgenic plants showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$ and delta < 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference.

Transgenic plants comprising recombinant DNA expressing protein as set forth in SEQ ID NO: 805, 828, 833, 870, 936, 937, 939, 965, 988, 1007, 1010, 1016, 1038, 1050, 1060, 1070, 1079, 1089, 1095, 1100, 1117, 1125, 1129, 1134, 1135, 1145, 1150, 1154, 1155, 1158, 1184, 1204, 1210, 1214, 1216, 1217, 1225, 1226, 1235, 1248, 1252, 1255, 1260, 1271, 1278, 1282, 1283, 1289, 1292, 1297, 1304, 1311, 1312, 1317, 1328, 1336, 1338, 1348, 1363, 1366, 1383, 1386, 1400, 1428, 1448, 1451, 1453, 1462, 1491, 1493, 1505, 1516, 1520, 1528, 1533, 1545, 1579, or 1588 showed enhanced tolerance to shade or low light condition by the second criteria as illustrated in Example 1L and 1M.

I. early plant growth and development screen

This example sets forth a plate based phenotypic analysis platform for the rapid detection of phenotypes that are evident during the first two weeks of growth. In this screen, we were looking for genes that confer advantages in the processes of germination, seedling vigor, root growth and root morphology under non-stressed growth conditions to plants. The transgenic plants with advantages in seedling growth and development were determined by the seedling weight and root length at day 14 after seed planting.

T2 seeds were plated on glufosinate selection plates and grown under standard conditions ($\sim 100 \text{ uE/m}^2/\text{s}$, 16 h photoperiod, 22°C at day, 20°C at night). Seeds were stratified for 3 days at 4°C . Seedlings were grown vertically (at a temperature of 22°C at day 20°C at night). Observations were taken on day 10 and day 14. Both seedling weight and root length at day 14 were analyzed as quantitative responses according to example 1M.

A list recombinant DNA constructs that improve early plant growth and development illustrated in Table 11.

Table 11.

NUC SEQ ID	PEP SEQ ID	Orientation	Root length at day 10		Root length at day 14		Seedling weight at day 14	
			Delta mean	P-value	Delta mean	P-value	Delta mean	P-value
52	855	SENSE	0.091	0.027	0.071	0.018	0.037	0.744
102	905	SENSE	0.188	0.210	0.104	0.247	0.308	0.021
121	924	SENSE	0.242	0.015	0.128	0.063	0.203	0.078
153	956	SENSE	0.153	0.027	0.122	0.197	0.256	0.010

NUC SEQ ID	PEP SEQ ID	Orientation	Root length at day 10		Root length at day 14		Seedling weight at day 14	
			Delta mean	P-value	Delta mean	P-value	Delta mean	P-value
134	937	SENSE	0.329	0.048	0.250	0.085	0.712	0.020
144	947	SENSE	0.127	0.146	0.045	0.566	0.371	0.023
101	904	SENSE	0.537	0.026	0.356	0.004	0.634	0.021
178	981	SENSE	0.372	0.136	0.273	0.033	0.346	0.082
179	982	SENSE	0.404	0.001	0.222	0.051	0.288	0.111
191	994	SENSE	0.282	0.008	0.285	0.009	0.445	0.012
163	966	SENSE	0.132	0.219	0.121	0.035	0.135	0.439
74	877	SENSE	0.282	0.033	0.166	0.015	0.143	0.363
162	965	SENSE	0.320	0.119	0.253	0.035	0.458	0.109
296	1099	SENSE	0.047	0.653	0.057	0.369	0.141	0.013
313	1116	SENSE	0.222	0.124	0.166	0.039	0.187	0.202
400	1203	SENSE	0.103	0.336	0.171	0.067	0.263	0.049
459	1262	SENSE	0.278	0.081	0.273	0.027	0.308	0.121
215	1018	SENSE	0.138	0.023	0.088	0.068	0.167	0.092
498	1301	SENSE	0.248	0.245	0.203	0.100	0.435	0.002
492	1295	SENSE	0.066	0.018	-0.052	0.334	0.215	0.171
567	1370	SENSE	-0.134	0.218	0.030	0.770	-0.132	0.305
611	1414	SENSE	0.232	0.220	0.168	0.035	0.165	0.319
614	1417	SENSE	0.142	0.277	0.147	0.174	0.167	0.562
616	1419	SENSE	0.038	0.549	0.046	0.130	-0.028	0.873
526	1329	SENSE	-0.044	0.135	0.070	0.044	-0.026	0.085
663	1466	SENSE	0.075	0.043	-0.065	0.418	0.047	0.639
79	882	SENSE	0.127	0.094	0.117	0.041	0.111	0.266
724	1527	SENSE	0.512	0.024	0.437	0.010	0.634	0.001
737	1540	SENSE	-0.011	0.931	0.120	0.151	-0.090	0.487
248	1051	SENSE	0.197	0.003	0.145	0.007	0.020	0.783
444	1247	SENSE	0.369	0.009	0.279	0.012	0.255	0.004
741	1544	SENSE	0.205	0.011	0.129	0.350	0.168	0.158
522	1325	SENSE	0.134	0.354	0.160	0.069	0.085	0.653
777	1580	SENSE	0.218	0.022	0.064	0.601	0.067	0.770
775	1578	SENSE	-0.192	0.322	0.019	0.780	-0.333	0.091
629	1432	SENSE	-0.042	0.151	0.075	0.016	-0.184	0.176
704	1507	SENSE	-0.074	0.546	0.123	0.058	0.113	0.433
697	1500	SENSE	0.173	0.049	0.078	0.546	0.109	0.054
706	1509	SENSE	0.074	0.018	0.076	0.301	0.059	0.664
331	1134	SENSE	0.107	0.013	0.050	0.569	0.202	0.178
546	1349	SENSE	-0.182	0.037	-0.050	0.104	-0.150	0.613
717	1520	SENSE	0.297	0.085	0.223	0.029	0.292	0.213
22	825	SENSE	/	/	/	/	0.444	0.025
581	1384	SENSE	0.116	0.126	0.193	0.013	0.181	0.175
726	1529	SENSE	0.375	0.012	0.088	0.188	0.353	0.015
755	1558	SENSE	0.409	0.031	0.203	0.091	0.346	0.198

Transgenic plants comprising recombinant DNA expressing a protein as set forth in 810, 831, 844, 857, 865, 884, 892, 917, 950, 954, 970, 983, 992, 998, 1004, 1005, 1029, 1033, 1043, 1072, 1101, 1109, 1115, 1133, 1181, 1200, 1284, 1310, 1340, 1384, 1391, 1443,

1449, 1471, 1480, 1483, 1498, 1506, 1508, 1510, 1529, 1548, 1553, 1555, 1556, 1558, 1559, 1564, 1576, or 1577 showed enhanced tolerance to shade or low light condition by the second criterion as illustrated in Example 1L and 1M.

J. late plant growth and development screen

5 This example sets forth a soil based phenotypic platform to identify genes that confer advantages in the processes of leaf development, flowering production and seed maturity to plants.

Arabidopsis plants were grown on a commercial potting mixture (Metro Mix 360, Scotts Co., Marysville, OH) consisting of 30-40% medium grade horticultural vermiculite, 35-55% sphagnum peat moss, 10-20% processed bark ash, 1-15% pine bark and a starter nutrient charge. Soil was supplemented with Osmocote time-release fertilizer at a rate of 30 mg/ft³. T2 seeds were imbibed in 1% agarose solution for 3 days at 4 °C and then sown at a density of ~5 per 2 ½" pot. Thirty-two pots were ordered in a 4 by 8 grid in standard greenhouse flat. Plants were grown in environmentally controlled rooms under a 16 h day length with an average light intensity of ~200 μmoles/m²/s. Day and night temperature set points were 22 °C and 20 °C, respectively. Humidity was maintained at 65%. Plants were watered by sub-irrigation every two days on average until mid-flowering, at which point the plants were watered daily until flowering was complete.

Application of the herbicide glufosinate was performed to select T2 individuals containing the target transgene. A single application of glufosinate was applied when the first true leaves were visible. Each pot was thinned to leave a single glufosinate-resistant seedling ~3 days after the selection was applied.

The rosette radius was measured at day 25. The silique length was measured at day 40. The plant parts were harvested at day 49 for dry weight measurements if flowering production was stopped. Otherwise, the dry weights of rosette and silique were carried out at day 53. The seeds were harvested at day 58. All measurements were analyzed as quantitative responses according to example 1M. A list of recombinant DNA constructs that improve late plant growth and development illustrated in Table 12.

Table 12

PEP SEQ ID	Construct ID	Rosette dry weight at day 53		Rosette radius at day 25		Seed net dry weight at day 62		Silique dry weight at day 53		Silique length at day 40	
		Delta mean	P- value	Delta mean	P- value	Delta mean	P- value	Delta mean	P-value	Delta mean	P-value
835	14405	0.110	0.430	/	/	0.557	0.001	0.200	0.214	0.043	0.252
869	14833	0.247	0.382	/	/	1.126	0.021	0.587	0.044	0.075	0.340

887	15207	0.238	0.023	-0.407	0.087	0.047	0.785	0.343	0.046	-0.015	0.540
949	70411	0.181	0.008	/	/	-0.268	0.192	0.061	0.025	-0.039	0.240
1057	71564	0.397	0.049	0.190	0.205	0.440	0.005	0.342	0.035	0.070	0.043
1066	72349	0.174	0.011	/	/	-0.526	0.049	-0.292	0.131	0.041	0.265
1102	72630	0.200	0.022	/	/	0.537	0.030	-0.257	0.034	-0.297	0.015
1108	73347	0.063	0.559	-0.046	0.475	0.848	0.035	0.243	0.188	-0.005	0.706
944	16612	-0.438	0.116	-0.280	0.026	0.493	0.023	-0.746	0.034	-0.023	0.289
929	70118	-0.186	0.237	-0.004	0.944	1.043	0.042	/	/	0.040	0.279
1025	70545	0.057	0.434	0.111	0.355	1.228	0.021	/	/	-0.060	0.418
1079	70686	-0.071	0.644	0.136	0.109	0.700	0.004	/	/	0.014	0.571
1088	70803	0.173	0.197	-0.207	0.053	0.691	0.008	/	/	0.067	0.094
1261	73438	0.397	0.057	-0.251	0.005	0.733	0.014	-0.056	0.542	-0.012	0.809
1077	73696	-0.384	0.425	-0.284	0.129	0.666	0.020	/	/	-0.097	0.154
1394	75705	0.570	0.049	0.019	0.797	0.477	0.165	0.213	0.337	-0.164	0.018
1210	76657	0.091	0.481	-0.370	0.031	0.744	0.019	-0.016	0.927	-0.038	0.395
1488	78386	0.092	0.010	-0.118	0.227	-0.578	0.034	/	/	0.017	0.502
1475	78522	0.386	0.003	0.728	0.001	1.532	0.007	/	/	0.170	0.038

If $p < 0.05$ and delta or risk score mean > 0 , the transgenic plants showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$ and delta or risk score mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference.

K. Limited nitrogen tolerance screen

Under low nitrogen conditions, *Arabidopsis* seedlings become chlorotic and have less biomass. This example sets forth the limited nitrogen tolerance screen to identify *Arabidopsis* plants transformed with the gene of interest that are altered in their ability to accumulate biomass and/or retain chlorophyll under low nitrogen condition.

T2 seeds were plated on glufosinate selection plates containing 0.5x N-Free Hoagland's T 0.1 mM NH_4NO_3 T 0.1% sucrose T 1% phytigel media and grown under standard light and temperature conditions. At 12 days of growth, plants were scored for seedling status (e.g., viable or non-viable) and root length. After 21 days of growth, plants were scored for BASTA resistance, visual color, seedling weight, number of green leaves, number of rosette leaves, root length and formation of flowering buds. A photograph of each plant was also taken at this time point.

The seedling weight and root length were analyzed as quantitative responses according to example 1M. The number green leaves, the number of rosette leaves and the flowerbud formation were analyzed as qualitative responses according to example 1L. The leaf color raw data were collected on each plant as the percentages of five color elements (Green, DarkGreen, LightGreen, RedPurple, YellowChlorotic) using a computer imaging

system. A statistical logistic regression model was developed to predict an overall value based on five colors for each plant.

A list of recombinant DNA constructs that improve low nitrogen availability tolerance in plants illustrated in Table 13.

5

Table 13

NUC SEQ ID	PEP SEQ ID	Construct ID	Orientation	Leaf color at day 21		Rosette weight at day 21	
				Risk score mean	P-value	Delta mean	P-value
188	991	10473	SENSE	4.078	0.017	-0.181	0.023
373	1176	11141	SENSE	4.800	0.027	-0.059	0.513
395	1198	11147	SENSE	3.499	0.012	-0.157	0.323
10	813	11860	ANTI-SENSE	-3.108	0.157	0.345	0.017
781	1584	11933	ANTI-SENSE	0.412	0.024	0.004	0.902
661	1464	12123	ANTI-SENSE	4.867	0.025	0.036	0.750
8	811	12147	ANTI-SENSE	2.164	0.134	0.074	0.007
6	809	12217	SENSE	3.494	0.047	0.086	0.245
12	815	12230	SENSE	1.878	0.077	0.111	0.015
799	1602	12444	SENSE	0.144	0.926	0.014	0.010
21	824	12749	ANTI-SENSE	0.315	0.022	0.109	0.325
3	806	12816	SENSE	-0.989	0.012	0.098	0.002
38	841	13038	ANTI-SENSE	2.207	0.450	0.074	0.040
39	842	13040	ANTI-SENSE	3.223	0.039	0.137	0.241
608	1411	13302	ANTI-SENSE	2.956	0.013	0.040	0.100
35	838	13828	ANTI-SENSE	3.733	0.002	-0.043	0.188
46	849	14340	ANTI-SENSE	3.417	0.023	-0.113	0.049
32	835	14405	SENSE	2.350	0.158	0.097	0.031
43	846	14415	SENSE	3.195	0.006	-0.131	0.066
85	888	14915	ANTI-SENSE	3.923	0.005	-0.081	0.079
78	881	15129	SENSE	3.751	0.002	0.096	0.164
104	907	15507	ANTI-SENSE	8.130	0.013	-0.086	0.312
106	909	15959	ANTI-SENSE	5.694	0.032	-0.165	0.121
98	901	16204	SENSE	7.476	0.013	-0.012	0.416
95	898	16309	SENSE	0.678	0.046	0.072	0.436
65	868	16323	SENSE	2.085	0.019	-0.008	0.779
16	819	17806	ANTI-SENSE	3.639	0.036	-0.087	0.208
184	987	18319	SENSE	1.043	0.023	-0.100	0.061
147	950	19154	SENSE	-1.213	0.201	0.146	0.033
175	978	19240	SENSE	0.175	0.828	0.098	0.002
289	1092	71622	SENSE	0.222	0.357	0.082	0.003
287	1090	71637	SENSE	0.074	0.772	0.061	0.001
261	1064	71810	SENSE	2.561	0.010	-0.038	0.111
416	1219	74350	SENSE	8.315	0.006	0.020	0.504
500	1303	74573	SENSE	4.232	0.015	0.093	0.207
555	1358	74836	SENSE	1.278	0.002	-0.133	0.018
561	1364	74918	SENSE	5.122	0.022	0.414	0.346
27	830	75958	SENSE	1.827	0.046	-0.078	0.101
731	1534	76321	SENSE	5.099	0.004	-0.120	0.313
754	1557	76821	SENSE	4.293	0.011	-0.157	0.044

NUC SEQ ID	PEP SEQ ID	Construct ID	Orientation	Leaf color at day 21		Rosette weight at day 21	
				Risk score mean	P-value	Delta mean	P-value
757	1560	76894	SENSE	3.806	0.048	0.123	0.102
513	1316	77041	SENSE	1.772	0.042	-0.186	0.098
786	1589	77250	SENSE	7.658	0.015	0.035	0.664
620	1423	77541	SENSE	1.896	0.045	-0.189	0.081
746	1549	77842	SENSE	4.863	0.048	-0.184	0.037
670	1473	77966	SENSE	1.716	0.120	0.218	0.022
719	1522	78191	SENSE	2.686	0.037	0.121	0.113
695	1498	78548	SENSE	1.766	0.191	0.116	0.027
226	1029	19775	SENSE	0.037	0.858	0.240	0.042
34	837	70721	ANTI-SENSE	1.858	0.012	0.052	0.047
280	1083	71696	SENSE	-0.133	0.002	0.085	0.049

For rosette weight, if $p < 0.05$ and delta or risk score mean > 0 , the transgenic plants showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$ and delta or risk score mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference with $p < 0.2$. For root length, if $p < 0.05$, the transgenic plants showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$, the transgenic plants showed a trend of trait enhancement as compared to the reference.

Transgenic plants comprising recombinant DNA expressing a protein as set forth in 807, 808, 817, 820, 829, 839, 847, 848, 850, 854, 858, 859, 866, 872, 875, 878, 889, 902, 908, 911, 922, 938, 944, 953, 963, 974, 980, 993, 1001, 1009, 1035, 1042, 1048, 1049, 1054, 1058, 1068, 1076, 1088, 1094, 1096, 1098, 1103, 1104, 1107, 1112, 1113, 1114, 1121, 1152, 1166, 1175, 1192, 1204, 1207, 1215, 1218, 1240, 1246, 1251, 1256, 1266, 1269, 1281, 1283, 1290, 1296, 1298, 1344, 1345, 1356, 1389, 1396, 1400, 1404, 1409, 1425, 1428, 1438, 1441, 1442, 1454, 1477, 1479, 1484, 1492, 1500, 1511, 1515, 1518, 1521, 1537, 1563, 1567, 1571, 1579, or 1603 showed enhanced tolerance to shade or low light condition by the second criterial as illustrated in Example 1L and 1M.

L. Statistic analysis for qualitative responses

A list of responses that were analyzed as qualitative responses illustrated in Table 14.

Table 14

response	Screen	categories (success vs. failure)
Wilting response Risk Score	Soil drought tolerance screen	non-wilted vs. wilted
growth stage at day 14	heat stress tolerance screen	50% of plants reach stage 1.03 vs. not
growth stage at day 14	salt stress tolerance screen	50% of plants reach stage 1.03 vs. not

response	Screen	categories (success vs. failure)
growth stage at day 14	PEG induced osmotic stress tolerance screen	50% of plants reach stage 1.03 vs. not
growth stage at day 7	cold germination tolerance screen	50% of plants reach stage 0.5 vs. not
number of rosette leaves at day 23	Shade tolerance screen	5 leaves appeared vs. not
Flower bud formation at day 23	Shade tolerance screen	flower buds appear vs. not
leaf angle at day 23	Shade tolerance screen	>60 degree vs. <60 degree
number of green leaves at day 21	limited nitrogen tolerance screen	6 or 7 leaves appeared vs. not
number of rosette leaves at day 21	limited nitrogen tolerance screen	6 or 7 leaves appeared vs. not
Flower bud formation at day 21	limited nitrogen tolerance screen	flower buds appear vs. not

Plants were grouped into transgenic and reference groups and were scored as success or failure according to Table 14. First, the risk (R) was calculated, which is the proportion of plants that were scored as of failure plants within the group. Then the relative risk (RR) was calculated as the ratio of R (transgenic) to R (reference). Risk score (RS) was calculated as - $\log_2^{RR}$. Two criteria were used to determine a transgenic with enhanced trait(s). Transgenic plants comprising recombinant DNA disclosed herein showed trait enhancement according to either or both of the two criteria.

For the first criteria, the risk scores from multiple events of the transgene of interest were evaluated for statistical significance by t-test using SAS statistical software (SAS 9, SAS/STAT User's Guide, SAS Institute Inc, Cary, NC, USA). RS with a value greater than 0 indicates that the transgenic plants perform better than the reference. RS with a value less than 0 indicates that the transgenic plants perform worse than the reference. The RS with a value equal to 0 indicates that the performance of the transgenic plants and the reference don't show any difference. If $p < 0.05$ and risk score mean > 0 , the transgenic plants showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$ and risk score mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference.

For the second criteria, the RS from each event was evaluated for statistical significance by t-test using SAS statistical software (SAS 9, SAS/STAT User's Guide, SAS Institute Inc, Cary, NC, USA). The RS with a value greater than 0 indicates that the transgenic plants from this event \ perform better than the reference. The RS with a value less than 0 indicates that the transgenic plants from this event perform worse than the reference.

The RS with a value equal to 0 indicates that the performance of the transgenic plants from this event and the reference don't show any difference. If $p < 0.05$ and risk score mean > 0 , the transgenic plants from this event showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$ and risk score mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference. If two or more events of the transgene of interest showed improvement in the same response, the transgene was deemed to show trait enhancement.

M. Statistic analysis for quantitative responses

A list of responses that were analyzed as quantitative responses illustrated in Table 15.

Table 15

response	screen
seed yield	Soil drought stress tolerance screen
seedling weight at day 14	heat stress tolerance screen
root length at day 14	heat stress tolerance screen
seedling weight at day 14	salt stress tolerance screen
root length at day 14	salt stress tolerance screen
root length at day 11	salt stress tolerance screen
seedling weight at day 14	PEG induced osmotic stress tolerance screen
root length at day 11	PEG induced osmotic stress tolerance screen
root length at day 14	PEG induced osmotic stress tolerance screen
rosette area at day 8	cold shock tolerance screen
rosette area at day 28	cold shock tolerance screen
difference in rosette area from day 8 to day 28	cold shock tolerance screen
root length at day 28	cold germination tolerance screen
seedling weight at day 23	Shade tolerance screen
petiole length at day 23	Shade tolerance screen
root length at day 14	Early plant growth and development screen
Seedling weight at day 14	Early plant growth and development screen
Rosette dry weight at day 53	Late plant growth and development screen
rosette radius at day 25	Late plant growth and development screen
seed dry weight at day 58	Late plant growth and development screen
silique dry weight at day 53	Late plant growth and development screen
silique length at day 40	Late plant growth and development screen
Seedling weight at day 21	Limited nitrogen tolerance screen
Root length at day 21	Limited nitrogen tolerance screen

The measurements (M) of each plant were transformed by $\log_2$ calculation. The Delta was calculated as $\log_2 M(\text{transgenic}) - \log_2 M(\text{reference})$. Two criteria were used to determine

trait enhancement. A transgene of interest could show trait enhancement according to either or both of the two criteria. The measurements (M) of each plant were transformed by $\log_2$ calculation. The Delta was calculated as $\log_2 M(\text{transgenic}) - \log_2 M(\text{reference})$. If the measured response was Petiole Length for the Low Light assay, Delta was subsequently multiplied by -1, to account for the fact that a shorter petiole length is considered an indication of trait enhancement.

For the first criteria, the Deltas from multiple events of the transgene of interest were evaluated for statistical significance by t-test using SAS statistical software (SAS 9, SAS/STAT User's Guide, SAS Institute Inc, Cary, NC, USA). Delta with a value greater than 0 indicates that the transgenic plants perform better than the reference. Delta with a value less than 0 indicates that the transgenic plants perform worse than the reference. The Delta with a value equal to 0 indicates that the performance of the transgenic plants and the reference don't show any difference. If $p < 0.05$ and risk score mean > 0 , the transgenic plants showed statistically significant trait enhancement as compared to the reference. If $p < 0.2$ and risk score mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference.

For the second criteria, the delta from each event was evaluated for statistical significance by t-test using SAS statistical software (SAS 9, SAS/STAT User's Guide, SAS Institute Inc, Cary, NC, USA). The Delta with a value greater than 0 indicates that the transgenic plants from this event perform better than the reference. The Delta with a value less than 0 indicates that the transgenic plants from this event perform worse than the reference. The Delta with a value equal to 0 indicates that the performance of the transgenic plants from this event and the reference don't show any difference. If $p < 0.05$ and delta mean > 0 , the transgenic plants from this event showed statistically significant trait improvement as compared to the reference. If $p < 0.2$ and delta mean > 0 , the transgenic plants showed a trend of trait enhancement as compared to the reference. If two or more events of the transgene of interest showed enhancement in the same response, the transgene was deemed to show trait improvement.

Example 2. Plant Expression Constructs

This example illustrates the construction of plasmids for transferring recombinant DNA into a plant cell nucleus that can be regenerated into transgenic plants.

Plant expression constructs for corn transformation

A base corn transformation vector pMON93039, as set forth in SEQ ID NO: 94614, illustrated in Table 16 and Figure 1, is made for use in preparing recombinant DNA for *Agrobacterium*-mediated transformation into corn tissue.

Table 16

Function	Name	Annotation	Coordinates of SEQ ID NO: 94614
Agrobacterium T-DNA transfer	B-AGRtu.right border	Agro right border sequence, essential for transfer of T- DNA.	11364-11720
Gene of interest expression cassette	E-Os.Act1	Upstream promoter region of the rice actin 1 gene	19-775
	E- CaMV.35S.2xA1- B3	Duplicated35S A1-B3 domain without TATA box	788-1120
	P-Os.Act1	Promoter region of the rice actin 1 gene	1125-1204
	L-Ta.Lhcb1	5' untranslated leader of wheat major chlorophyll a/b binding protein	1210-1270
	I-Os.Act1	First intron and flanking UTR exon sequences from the rice actin 1 gene	1287-1766
	T-St.Pis4	3' non-translated region of the potato proteinase inhibitor II gene which functions to direct polyadenylation of the mRNA	1838-2780
Plant selectable marker expression cassette	P-Os.Act1	Promoter from the rice actin 1 gene	2830-3670
	L-Os.Act1	First exon of the rice actin 1 gene	3671-3750
	I-Os.Act1	First intron and flanking UTR exon sequences from the rice actin 1 gene	3751-4228
	TS-At.ShkG-CTP2	Transit peptide region of Arabidopsis EPSPS	4238-4465
	CR-AGRtu.aroA- CP4.nat	Coding region for bacterial strain CP4 native aroA gene.	4466-5833
	T-AGRtu.nos	A 3' non-translated region of the nopaline synthase gene of <i>Agrobacterium</i> <i>tumefaciens</i> Ti plasmid which functions to direct polyadenylation of the mRNA.	5849-6101

Function	Name	Annotation	Coordinates of SEQ ID NO: 94614
Agrobacterium T-DNA transfer	B-AGRtu.left border	Agro left border sequence, essential for transfer of T- DNA.	6168-6609
Maintenance in E. coli	OR-Ec.oriV-RK2	The vegetative origin of replication from plasmid RK2.	6696-7092
	CR-Ec.rop	Coding region for repressor of primer from the ColE1 plasmid. Expression of this gene product interferes with primer binding at the origin of replication, keeping plasmid copy number low.	8601-8792
	OR-Ec.ori-ColE1	The minimal origin of replication from the E. coli plasmid ColE1.	9220-9808
	P-Ec.aadA- SPC/STR	Promoter for Tn7 adenylyltransferase (AAD(3"))	10339-10380
	CR-Ec.aadA- SPC/STR	Coding region for Tn7 adenylyltransferase (AAD(3")) conferring spectinomycin and streptomycin resistance.	10381-11169
	T-Ec.aadA- SPC/STR	3' UTR from the Tn7 adenylyltransferase (AAD(3")) gene of E. coli.	11170-11227

Primers for PCR amplification of protein coding nucleotides of recombinant DNA are designed at or near the start and stop codons of the coding sequence in order to eliminate most of the 5' and 3' untranslated regions. Each recombinant DNA coding for or suppressing a protein identified in Table 2 is amplified by PCR prior to insertion into the insertion site within the gene of interest expression cassette of the base vector.

B. Plant expression constructs for soy and canola transformation

Vectors for use in transformation of soybean and canola can also be prepared. Elements of an exemplary common expression vector pMON82053 (SEQ ID NO: 94615) are shown in Table 17 below and Figure 2.

Table 17.

Function	Name	Annotation	Coordinates of SEQ ID NO: 94615
----------	------	------------	---------------------------------------

Function	Name	Annotation	Coordinates of SEQ ID NO: 94615
Agrobacterium T-DNA transfer	B-AGRtu.left border	Agro left border sequence, essential for transfer of T-DNA.	6144-6585
Plant selectable marker expression cassette	P-At.Act7	Promoter from the Arabidopsis actin 7 gene	6624-7861
	L-At.Act7	5'UTR of Arabidopsis Act7 gene	
	I-At.Act7	Intron from the Arabidopsis actin7 gene	
	TS-At.ShkG- CTP2	Transit peptide region of Arabidopsis EPSPS	7864-8091
	CR-AGRtu.aroA- CP4.nno_At	Synthetic CP4 coding region with dicot preferred codon usage.	8092-9459
	T-AGRtu.nos	A 3' non-translated region of the nopaline synthase gene of Agrobacterium tumefaciens Ti plasmid which functions to direct polyadenylation of the mRNA.	9466-9718
Gene of interest expression cassette	P-CaMV.35S-enh	Promoter for 35S RNA from CaMV containing a duplication of the -90 to -350 region.	1-613
	T-Gb.E6-3b	3' untranslated region from the fiber protein E6 gene of sea-island cotton.	688-1002
Agrobacterium T-DNA transfer	B-AGRtu.right border	Agro right border sequence, essential for transfer of T-DNA.	1033-1389
Maintenance in E. coli	OR-Ec.oriV-RK2	The vegetative origin of replication from plasmid RK2.	5661-6057
	CR-Ec.rop	Coding region for repressor of primer from the ColE1 plasmid. Expression of this gene product interferes with primer binding at the origin of replication, keeping plasmid copy number low.	3961-4152
	OR-Ec.ori-ColE1	The minimal origin of replication from the E. coli plasmid ColE1.	2945-3533
	P-Ec.aadA- SPC/STR	Promoter for Tn7 adenylyltransferase (AAD(3"))	2373-2414
	CR-Ec.aadA- SPC/STR	Coding region for Tn7 adenylyltransferase (AAD(3")) conferring spectinomycin and streptomycin resistance.	1584-2372
	T-Ec.aadA- SPC/STR	3' UTR from the Tn7 adenylyltransferase (AAD(3")) gene of E. coli.	1526-1583

Primers for PCR amplification of protein coding nucleotides of recombinant DNA are designed at or near the start and stop codons of the coding sequence in order to eliminate

most of the 5' and 3' untranslated regions. Each recombinant DNA coding for or suppressing a protein identified in Table 2 is amplified by PCR prior to insertion into the insertion site within the gene of interest expression cassette of the base vector.

C. Cotton transformation vector

- 5 Plasmids for use in transformation of cotton tissue are prepared with elements of expression vector pMON99053 (SEQ ID NO: 94616) are shown in Table 18 below and Figure 3. Primers for PCR amplification of protein coding nucleotides of recombinant DNA are designed at or near the start and stop codons of the coding sequence in order to eliminate most of the 5' and 3' untranslated regions. Each recombinant DNA coding for or suppressing
- 10 a protein identified in Table 2 is amplified by PCR prior to insertion into the insertion site within the gene of interest expression cassette of the base vector.

Table 18.

Function	Name	Annotation	Coordinates of SEQ ID NO: 94616
Agrobacterium T-DNA transfer	B-AGRtu.right border	Agro right border sequence, essential for transfer of T-DNA.	1-357
Gene of interest expression cassette	Exp-CaMV.35S-enh+Ph.DnaK	Enhanced version of the 35S RNA promoter from CaMV plus the petunia hsp70 5' untranslated region	388-1091
	T-Ps.RbcS2-E9	The 3' non-translated region of the pea RbcS2 gene which functions to direct polyadenylation of the mRNA.	1165-1797
Plant selectable marker expression cassette	Exp-CaMV.35S	Promoter and 5' untranslated region from the 35S RNA of CaMV	1828-2151
	CR-Ec.nptII-Tn5	Coding region for neomycin phosphotransferase gene from transposon Tn5 which confers resistance to neomycin and kanamycin.	2185-2979
	T-AGRtu.nos	A 3' non-translated region of the nopaline synthase gene of Agrobacterium tumefaciens Ti plasmid which functions to direct polyadenylation of the mRNA.	3011-3263
Agrobacterium T-DNA transfer	B-AGRtu.left border	Agro left border sequence, essential for transfer of T-DNA.	3309-3750

Function	Name	Annotation	Coordinates of SEQ ID NO: 94616
Maintenance in E. coli	OR-Ec.oriV-RK2	The vegetative origin of replication from plasmid RK2.	3837-4233
	CR-Ec.rop	Coding region for repressor of primer from the ColE1 plasmid. Expression of this gene product interferes with primer binding at the origin of replication, keeping plasmid copy number low.	5742-5933
	OR-Ec.ori-ColE1	The minimal origin of replication from the E. coli plasmid ColE1.	6363-6949
	P-Ec.aadA-SPC/STR	Promoter for Tn7 adenylyltransferase (AAD(3"))	7480-7521
	CR-Ec.aadA-SPC/STR	Coding region for Tn7 adenylyltransferase (AAD(3")) conferring spectinomycin and streptomycin resistance.	7522-8310
	T-Ec.aadA-SPC/STR	3' UTR from the Tn7 adenylyltransferase (AAD(3")) gene of E. coli.	8311-8368

Example 3. Corn Transformation

This example illustrates transformation methods in producing a transgenic nucleus in a corn plant cell and the plants, seeds and pollen produced from a transgenic cell with such a nucleus having an enhanced trait, *e.g.* enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil. Plasmid vectors are prepared by cloning DNA identified in Table 2 in the base vector for use in corn transformation of corn tissue.

For *Agrobacterium*-mediated transformation of corn embryo cells corn plants of a readily transformable line are grown in the greenhouse and ears are harvested when the embryos are 1.5 to 2.0 mm in length. Ears are surface sterilized by spraying or soaking the ears in 80% ethanol, followed by air drying. Immature embryos are isolated from individual kernels on surface sterilized ears. Prior to inoculation of maize cells, *Agrobacterium* cells are grown overnight at room temperature. Immature maize embryo cells are inoculated with *Agrobacterium* shortly after excision, and incubated at room temperature with *Agrobacterium* for 5-20 minutes. Immature embryo plant cells are then co-cultured with *Agrobacterium* for 1 to 3 days at 23°C in the dark. Co-cultured embryos are transferred to selection media and cultured for approximately two weeks to allow embryogenic callus to develop.

Embryogenic callus is transferred to culture medium containing 100 mg/L paromomycin and subcultured at about two week intervals. Transformed plant cells are recovered 6 to 8 weeks after initiation of selection.

For *Agrobacterium*-mediated transformation of maize callus immature embryos are
5 cultured for approximately 8-21 days after excision to allow callus to develop. Callus is then incubated for about 30 minutes at room temperature with the *Agrobacterium* suspension, followed by removal of the liquid by aspiration. The callus and *Agrobacterium* are co-cultured without selection for 3-6 days followed by selection on paromomycin for approximately 6 weeks, with biweekly transfers to fresh media. Paromomycin resistant calli
10 are identified about 6-8 weeks after initiation of selection.

For transformation by microprojectile bombardment maize immature embryos are isolated and cultured 3-4 days prior to bombardment. Prior to microprojectile bombardment, a suspension of gold particles is prepared onto which the desired recombinant DNA expression cassettes are precipitated. DNA is introduced into maize cells as described
15 in U.S. Patents 5,550,318 and 6,399,861 using the electric discharge particle acceleration gene delivery device. Following microprojectile bombardment, tissue is cultured in the dark at 27°C. Additional transformation methods and materials for making transgenic plants of this invention, for example, various media and recipient target cells, transformation of immature embryos and subsequence regeneration of fertile transgenic plants are disclosed in U.S.
20 Patents 6,194,636, 6,232,526 and 7,151,204, which are incorporated herein by reference.

To regenerate transgenic corn plants a callus of transgenic plant cells resulting from transformation and selection is placed on media to initiate shoot development into plantlets which are transferred to potting soil for initial growth in a growth chamber at 26°C followed by a mist bench before transplanting to 5 inch pots where plants are grown to maturity. The
25 regenerated plants are self-fertilized and seed is harvested for use in one or more methods to select seeds, seedlings or progeny second generation transgenic plants (R2 plants) or hybrids, *e.g.* by selecting transgenic plants exhibiting an enhanced trait as compared to a control plant.

Transgenic corn plant cells are transformed with recombinant DNA from each of the genes identified in Table 2. Progeny transgenic plants and seed of the transformed plant cells
30 are screened for enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil as obtained in Example 7.

Example 4. Soybean transformation

This example illustrates plant transformation in producing the transgenic soybean plants of this invention and the production and identification of transgenic seed for transgenic soybean having enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

5 For *Agrobacterium* mediated transformation, soybean seeds are imbibed overnight and the meristem explants excised. The explants are placed in a wounding vessel. Soybean explants and induced *Agrobacterium* cells from a strain containing plasmid DNA with the gene of interest cassette and a plant selectable marker cassette are mixed no later than 14 hours from the time of initiation of seed imbibition, and wounded using sonication.
10 Following wounding, explants are placed in co-culture for 2-5 days at which point they are transferred to selection media for 6-8 weeks to allow selection and growth of transgenic shoots. Resistant shoots are harvested approximately 6-8 weeks and placed into selective rooting media for 2-3 weeks. Shoots producing roots are transferred to the greenhouse and potted in soil. Shoots that remain healthy on selection, but do not produce roots are
15 transferred to non-selective rooting media for an additional two weeks. Roots from any shoots that produce roots off selection are tested for expression of the plant selectable marker before they are transferred to the greenhouse and potted in soil. Additionally, a DNA construct can be transferred into the genome of a soybean cell by particle bombardment and the cell regenerated into a fertile soybean plant as described in U.S. Patent 5,015,580, herein
20 incorporated by reference.

Transgenic soybean plant cells are transformed with recombinant DNA from each of the genes identified in Table 2. Transgenic progeny plants and seed of the transformed plant cells are screened for enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil as
25 obtained in Example 10.

Example 5. Cotton transgenic plants with enhanced agronomic traits

Cotton transformation is performed as generally described in WO0036911 and in U.S. Pat. No. 5,846,797. Transgenic cotton plants containing each recombinant DNA having a sequence of SEQ ID NO: 1 through SEQ ID NO: 803 are obtained by transforming with
30 recombinant DNA from each of the genes identified in Table 2. Progeny transgenic plants are selected from a population of transgenic cotton events under specified growing conditions and are compared with control cotton plants. Control cotton plants are substantially the same cotton genotype but without the recombinant DNA, for example, either a parental cotton plant of the same genotype that was not transformed with the identical recombinant DNA or a

negative isoline of the transformed plant. Additionally, a commercial cotton cultivar adapted to the geographical region and cultivation conditions, *e.g.* cotton variety ST474, cotton variety FM 958, and cotton variety Siokra L-23, are used to compare the relative performance of the transgenic cotton plants containing the recombinant DNA. The specified culture conditions are growing a first set of transgenic and control plants under “wet” conditions, *e.g.* irrigated in the range of 85 to 100 percent of evapotranspiration to provide leaf water potential of -14 to -18 bars, and growing a second set of transgenic and control plants under “dry” conditions, *e.g.* irrigated in the range of 40 to 60 percent of evapotranspiration to provide a leaf water potential of -21 to -25 bars. Pest control, such as weed and insect control is applied equally to both wet and dry treatments as needed. Data gathered during the trial includes weather records throughout the growing season including detailed records of rainfall; soil characterization information; any herbicide or insecticide applications; any gross agronomic differences observed such as leaf morphology, branching habit, leaf color, time to flowering, and fruiting pattern; plant height at various points during the trial; stand density; node and fruit number including node above white flower and node above crack boll measurements; and visual wilt scoring. Cotton boll samples are taken and analyzed for lint fraction and fiber quality. The cotton is harvested at the normal harvest timeframe for the trial area. Enhanced water use efficiency is indicated by increased yield, improved relative water content, enhanced leaf water potential, increased biomass, enhanced leaf extension rates, and improved fiber parameters.

The transgenic cotton plants of this invention are identified from among the transgenic cotton plants by agronomic trait screening as having increased yield and enhanced water use efficiency.

Example 6. Canola transformation

This example illustrates plant transformation in producing the transgenic canola plants of this invention and the production and identification of transgenic seed for transgenic canola having enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

Tissues from *in vitro* grown canola seedlings are prepared and inoculated with overnight-grown *Agrobacterium* cells containing plasmid DNA with the gene of interest cassette and a plant selectable marker cassette. Following co-cultivation with *Agrobacterium*, the infected tissues are allowed to grow on selection to promote growth of transgenic shoots, followed by growth of roots from the transgenic shoots. The selected plantlets are then transferred to the greenhouse and potted in soil. Molecular characterization is performed to

confirm the presence of the gene of interest, and its expression in transgenic plants and progenies. Progeny transgenic plants are selected from a population of transgenic canola events under specified growing conditions and are compared with control canola plants. Control canola plants are substantially the same canola genotype but without the recombinant DNA, for example, either a parental canola plant of the same genotype that is not transformed with the identical recombinant DNA or a negative isolate of the transformed plant

Transgenic canola plant cells are transformed with recombinant DNA from each of the genes identified in Table 1. Transgenic progeny plants and seed of the transformed plant cells are screened for enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil as obtained in Example 7.

Example 7. Identification of Homologs

This example illustrates the identification of homologs of proteins encoded by the DNA identified in Table 1 which is used to provide transgenic seed and plants having enhanced agronomic traits. From the sequence of the homologs, homologous DNA sequence can be identified for preparing additional transgenic seeds and plants of this invention with enhanced agronomic traits.

An "All Protein Database" is constructed of known protein sequences using a proprietary sequence database and the National Center for Biotechnology Information (NCBI) non-redundant amino acid database (nr.aa). For each organism from which a polynucleotide sequence provided herein was obtained, an "Organism Protein Database" is constructed of known protein sequences of the organism; it is a subset of the All Protein Database based on the NCBI taxonomy ID for the organism.

The All Protein Database is queried using amino acid sequences provided herein as SEQ ID NO: 804 through SEQ ID NO: 1606 using NCBI "blastp" program with E-value cutoff of $1e^{-8}$. Up to 1000 top hits are kept, and separated by organism names. For each organism other than that of the query sequence, a list is kept for hits from the query organism itself with a more significant E-value than the best hit of the organism. The list contains likely duplicated genes of the polynucleotides provided herein, and is referred to as the Core List. Another list is kept for all the hits from each organism, sorted by E-value, and referred to as the Hit List.

The Organism Protein Database is queried using polypeptide sequences provided herein as SEQ ID NO: 804 through SEQ ID NO: 1606 using NCBI "blastp" program with E-value cutoff of $1e^{-4}$. Up to 1000 top hits are kept. A BLAST searchable database is

constructed based on these hits, and is referred to as “SubDB”. SubDB is queried with each sequence in the Hit List using NCBI “blastp” program with E-value cutoff of $1e^{-8}$. The hit with the best E-value is compared with the Core List from the corresponding organism. The hit is deemed a likely ortholog if it belongs to the Core List, otherwise it is deemed not a likely ortholog and there is no further search of sequences in the Hit List for the same organism. Likely orthologs from a large number of distinct organisms were identified and are reported by amino acid sequences of SEQ ID NO: 1607 to SEQ ID NO: 94613. These orthologs are reported in Tables 19 as homologs to the proteins cognate to genes used in trait-improving recombinant DNA

10 **Example 8. Consensus sequence build**

ClustalW program is selected for multiple sequence alignments of an amino acid sequence of SEQ ID NO: 804 and its homologs, through SEQ ID NO: 1606 and its homologs. Three major factors affecting the sequence alignments dramatically are (1) protein weight matrices; (2) gap open penalty; (3) gap extension penalty. Protein weight matrices available for ClustalW program include Blosom, Pam and Gonnet series. Those parameters with gap open penalty and gap extension penalty were extensively tested. On the basis of the test results, Blosom weight matrix, gap open penalty of 10 and gap extension penalty of 1 were chosen for multiple sequence alignment. The consensus sequence of SEQ ID NO: 1325 and its 30 homologs were derived according to the procedure described above and is displayed in Figure 4.

20 **Example 9. Pfam module annotation**

This example illustrates the identification of domain and domain module by Pfam analysis.

The amino acid sequence of the expressed proteins that were shown to be associated with an enhanced trait were analyzed for Pfam protein family against the current Pfam collection of multiple sequence alignments and hidden Markov models using the HMMER software in the appended computer listing. The Pfam domain modules and individual protein domain for the proteins of SEQ ID NO: 804 through 1606 are shown in Table 20 and Table 21 respectively. The Hidden Markov model databases for the identified patent families are also in the appended computer listing allowing identification of other homologous proteins and their cognate encoding DNA to enable the full breadth of the invention for a person of ordinary skill in the art. Certain proteins are identified by a single Pfam domain and others by multiple Pfam domains. For instance, the protein with amino acids of SEQ ID NO: 830 is characterized by two Pfam domains, *e.g.* “Lectin_legB” and “Pkinase”. See also the protein

with amino acids of SEQ ID NO: 817 which is characterized by four copies of the Pfam domain "Arm". In Table 21 "score" is the gathering score for the Hidden Markov Model of the domain which exceeds the gathering cutoff reported in Table 22.

Table 20.

PEP Seq ID No.	Construct ID	Pfam module	Position
804	CGPG10.pep	p450	31-495
805	CGPG1008.pep	Tryp_alpha_amyl	23-95
806	CGPG1023.pep	Na_Ca_ex::Na_Ca_ex	104-255::299-432
807	CGPG1038.pep	DUF1475	1-256
812	CGPG1113.pep	DUF1350	49-370
814	CGPG1163.pep	DSPc	49-182
817	CGPG1205.pep	Arm::Arm::Arm::Arm	94-134::135-175::176-216::258-298
818	CGPG1207.pep	NifU_N	26-105
820	CGPG1223.pep	ATP_bind_1	7-250
821	CGPG1250.pep	DUF822	12-160
824	CGPG1319.pep	DUF599	10-233
826	CGPG1329.pep	Cullin	29-631
827	CGPG133.pep	Cyclin_N::Cyclin_C	168-293::295-422
828	CGPG1332.pep	Cullin	12-646
829	CGPG1343.pep	LRR_1	389-412
830	CGPG1348.pep	Lectin_legB::Pkinase	27-248::334-604
831	CGPG1349.pep	B_lectin::S_locus_glycop::PAN_1::Pkinase	65-171::180-312::327-404::484-731
832	CGPG1373.pep	Pkinase	12-291
833	CGPG1377.pep	Pkinase	4-258
834	CGPG1412.pep	TPP_enzyme_N::TPP_enzyme_M::TPP_enzyme_C	45-221::243-376::431-578
835	CGPG1421.pep	Biotin_lipoyl::E3_binding::2-oxoacid_dh	76-149::180-218::249-480
836	CGPG1426.pep	Aminotran_1_2	48-432
837	CGPG1433.pep	Pyr_redox_2::Pyr_redox_dim	45-359::388-497
838	CGPG1453.pep	malic::Malic_M	171-360::362-615
839	CGPG1454.pep	Ribul_P_3_epim	7-207
840	CGPG1463.pep	Alpha-amylase::Alpha-amyl_C2	26-361::362-422
841	CGPG1464.pep	Glycolytic	11-358
842	CGPG1471.pep	IF4E	1-198
843	CGPG1481.pep	Pkinase	4-260
844	CGPG1499.pep	PPDK_N::PEP-utilizers::PEP-utilizers_C	85-445::496-586::598-955
845	CGPG150.pep	p450	30-507
846	CGPG1536.pep	CRAL_TRIO_N::CRAL_TRIO	73-142::159-345
847	CGPG1539.pep	UPF0139	8-107

PEP Seq ID No.	Construct ID	Pfam module	Position
848	CGPG155.pep	GlutR_N::Shikimate_DH::GlutR_dimer	100-251::255-407::420-526
849	CGPG1583.pep	PLAC8	88-187
850	CGPG1588.pep	PLAC8	75-214
851	CGPG16.pep	MIP	14-235
852	CGPG1609.pep	MMR_HSR1::DUF933	54-190::337-420
853	CGPG1629.pep	WD40::WD40::WD40	262-301::307-346::356-395
854	CGPG1637.pep	DUF220	163-262
855	CGPG1653.pep	TBC	200-415
856	CGPG1658.pep	Nodulin-like	14-260
857	CGPG1663.pep	BT1	30-441
858	CGPG1682.pep	OPT	62-685
859	CGPG1701.pep	Glyco_transf_8	76-299
861	CGPG1724.pep	Per1	63-334
862	CGPG1726.pep	DUF607	111-291
863	CGPG1736.pep	Usp	1-148
864	CGPG1741.pep	mTERF	279-627
865	CGPG1783.pep	PB1	42-132
866	CGPG1790.pep	PB1	59-153
868	CGPG1845.pep	TFIID-18kDa	29-119
869	CGPG1855.pep	CAF1	3-227
870	CGPG1870.pep	Pkinase	406-572
871	CGPG1879.pep	Pkinase::NAF	21-277::303-364
872	CGPG1886.pep	Pkinase	26-325
873	CGPG1903.pep	F-box	320-367
874	CGPG1905.pep	F-box::Kelch_2::Kelch_2	4-51::104-150::255-306
875	CGPG1914.pep	F-box::Kelch_1::Kelch_1	42-89::170-215::217-266
876	CGPG193.pep	WD40::WD40::WD40::WD40	167-205::268-307::318-355::363-401
877	CGPG1939.pep	F-box::WD40::WD40::WD40	66-113::152-190::247-283::325-364
878	CGPG1949.pep	YDG_SRA::Pre-SET::SET	360-519::543-639::641-771
879	CGPG1959.pep	PHD::SET	34-82::208-344
880	CGPG197.pep	p450	40-481
882	CGPG1981.pep	MT-A70	476-636
883	CGPG1999.pep	Nfu_N::NifU	78-193::221-291
884	CGPG2.pep	p450	71-531
885	CGPG2006.pep	RCC1::RCC1::RCC1::RCC1	32-82::138-187::190-239::294-343
886	CGPG2010.pep	Aminotran_5	74-448
887	CGPG2011.pep	X8	128-209
888	CGPG2014.pep	Pkinase::NAF	28-282::341-398
889	CGPG2023.pep	Zip	48-352
890	CGPG2026.pep	Pkinase	38-324

PEP Seq ID No.	Construct ID	Pfam module	Position
892	CGPG2064.pep	Cenp-O	123-201
894	CGPG2077.pep	NAC::UBA	88-147::195-232
895	CGPG2095.pep	AARP2CN::DUF663	228-309::483-780
896	CGPG2105.pep	Pkinase::Pkinase_C	102-404::422-471
897	CGPG2108.pep	Response_reg::CCT	64-180::442-484
898	CGPG2111.pep	polyprenyl_synt	74-324
899	CGPG2124.pep	UIM::efhand	139-156::220-248
900	CGPG2125.pep	Glutaredoxin	44-110
903	CGPG2134.pep	ATP-synt_G	8-122
904	CGPG2139.pep	DUF1517	94-391
905	CGPG2140.pep	PhzC-PhzF	6-282
907	CGPG2163.pep	Cupin_1::Cupin_1	5-157::190-339
908	CGPG2165.pep	ClpS	76-152
910	CGPG2218.pep	Spc97_Spc98	67-555
912	CGPG2225.pep	EMP24_GP25L	52-116
913	CGPG2229.pep	Subtilisin_N::PA	30-106::362-461
915	CGPG2254.pep	Ribosomal_L7Ae	20-114
916	CGPG2268.pep	CBS	66-188
917	CGPG227.pep	p450	37-494
918	CGPG2312.pep	UPF0061	75-571
919	CGPG2315.pep	SIP1	286-514
920	CGPG2316.pep	PTPA	98-396
922	CGPG235.pep	p450	74-510
923	CGPG2358.pep	Branch	54-278
925	CGPG2361.pep	Metallophos	9-231
926	CGPG2365.pep	Epimerase	3-211
927	CGPG2372.pep	Pribosyltran	29-169
928	CGPG2374.pep	Aminotran_5	8-370
929	CGPG2377.pep	DAO	69-466
930	CGPG2387.pep	H_PPase	1-239
931	CGPG2389.pep	Carb_anhydrase	52-275
932	CGPG2395.pep	AA_permease	89-526
934	CGPG2409.pep	DUF862	17-154
935	CGPG2410.pep	GST_N::GST_C	5-79::101-204
937	CGPG2416.pep	HinI	97-235
938	CGPG2441.pep	Sina	5-205
939	CGPG2450.pep	Epimerase	114-371
940	CGPG2451.pep	RALF	57-129
941	CGPG2492.pep	Pribosyltran	80-216
942	CGPG2495.pep	ADH_N::ADH_zinc_N	25-134::165-307
943	CGPG2506.pep	PfkB	5-289
944	CGPG2515.pep	TIM	5-244
945	CGPG2531.pep	AhpC-TSA	5-138
946	CGPG2581.pep	SelR	12-133
947	CGPG2584.pep	zf-Tim10_DDP	22-86
948	CGPG2592.pep	TFIID_30kDa	30-80
949	CGPG2612.pep	RRM_1::RRM_1::RRM_1	65-132::150-225::275-343
950	CGPG2660.pep	B3::B3	17-115::211-301

PEP Seq ID No.	Construct ID	Pfam module	Position
951	CGPG2663.pep	NTF2::RRM_1	15-131::295-365
952	CGPG2679.pep	ACBP::Ank::Ank	104-190::265-297::298-330
953	CGPG2696.pep	zf-C3HC4	177-218
954	CGPG2772.pep	Mov34	6-136
955	CGPG2773.pep	Asp	161-498
956	CGPG281.pep	TLC	91-574
957	CGPG2846.pep	Aldo_ket_red	49-365
958	CGPG2852.pep	Glyoxalase	9-132
960	CGPG2870.pep	zf-CCCH::zf-CCCH::zf-CCCH::zf-CCCH::zf-CCCH	111-137::159-185::205-231::347-373::393-419
961	CGPG2877.pep	LAG1	96-307
962	CGPG289.pep	Cellulose_synt	243-1060
963	CGPG2924.pep	zf-C3HC4	138-179
964	CGPG2947.pep	Acetyltransf_1::Bromodomain	265-343::460-548
965	CGPG2963.pep	C1_2::C1_3::C1_3::C1_2::C1_3::C1_3::C1_2	81-108::136-164::194-223::305-335::391-420::494-522::552-581
966	CGPG2987.pep	SRF-TF	Nov-66
968	CGPG3045.pep	Pkinase_Tyr	82-356
969	CGPG3046.pep	AAA	121-305
970	CGPG3060.pep	p450	67-528
971	CGPG3075.pep	Auxin_inducible	7-106
972	CGPG310.pep	Cyclin_N	18-143
973	CGPG3103.pep	PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR	14-48::49-83::84-118::119-153::155-188::189-223::224-258::259-293::295-329::330-364::365-398
974	CGPG315.pep	p450	29-491
976	CGPG3189.pep	RRM_1::RRM_1	115-186::209-280
977	CGPG3204.pep	KH_1::KH_1::KH_1::KH_1::KH_1	45-104::151-221::318-381::400-467::575-637
978	CGPG3208.pep	MATH::MATH	26-153::180-302
979	CGPG3219.pep	2OG-FeII_Oxy	220-320
980	CGPG3233.pep	PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR	160-194::195-229::231-265::266-300::301-335::336-370::371-405::406-440
981	CGPG3263.pep	Tryp_alpha_amyl	28-104
982	CGPG3276.pep	ECH	11-186
984	CGPG3282.pep	Steroid_dh	117-268
985	CGPG3300.pep	zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC	55-72::74-91::93-110::112-129::138-155::157-174::229-246

PEP Seq ID No.	Construct ID	Pfam module	Position
986	CGPG3318.pep	Peptidase_C14	3-416
987	CGPG3319.pep	AP2	28-79
988	CGPG3326.pep	zf-C3HC4	259-299
989	CGPG3333.pep	PTS_2-RNA	48-239
990	CGPG3338.pep	Alba	17-87
991	CGPG334.pep	Mov34	20-128
992	CGPG3374.pep	peroxidase	42-286
993	CGPG3402.pep	Dimerisation::Methyltransf_2	30-89::99-342
995	CGPG3413.pep	WD40::U3_snoRNA_C	168-207::377-523
996	CGPG3422.pep	WD40::WD40::WD40	241-279::387-425::519-559
997	CGPG3436.pep	DUF6::DUF6	17-144::198-327
998	CGPG3539.pep	Mito_carr::Mito_carr::Mito_carr	11-105::113-209::214-303
999	CGPG3550.pep	FAE1_CUT1_RppA::ACP_syn_III_C	81-370::384-468
1000	CGPG3551.pep	Pkinase	9-273
1001	CGPG3552.pep	Aldedh	23-441
1002	CGPG3572.pep	FA_hydroxylase	112-225
1004	CGPG3599.pep	DUF579	51-289
1008	CGPG364.pep	zf-C3HC4	97-137
1010	CGPG3679.pep	malic::Malic_M::PTA_PTB	36-179::181-418::446-768
1011	CGPG3686.pep	Gp_dh_N::Gp_dh_C	2-150::155-312
1012	CGPG3694.pep	LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::Pkinase_Tyr	38-78::106-128::130-152::154-176::178-201::312-583
1013	CGPG3696.pep	MMR_HSR1	136-257
1014	CGPG3698.pep	WD40::WD40	104-142::234-272
1015	CGPG3699.pep	adh_short	19-187
1016	CGPG3702.pep	SAM_1	228-291
1017	CGPG3703.pep	ENOD93	25-103
1018	CGPG3707.pep	Pkinase	42-333
1019	CGPG3710.pep	VQ	118-148
1020	CGPG3730.pep	DUF167	140-216
1021	CGPG3731.pep	DUF616	194-508
1022	CGPG3734.pep	PRK::Pribosyltran	46-233::280-425
1023	CGPG3745.pep	RRM_1::RRM_1	386-455::481-553
1024	CGPG3764.pep	Response_reg	20-145
1026	CGPG3851.pep	SH3_1	284-338
1027	CGPG3911.pep	Aldedh	104-587
1028	CGPG3948.pep	NAP	50-295
1031	CGPG3996.pep	Inositol_P	6-267
1032	CGPG4006.pep	ENT	1-74
1033	CGPG4025.pep	SQS_PSY	44-331
1034	CGPG4028.pep	Transket_pyr::Transketolase_C	398-565::579-702
1035	CGPG403.pep	Pkinase	78-336

PEP Seq ID No.	Construct ID	Pfam module	Position
1036	CGPG4041.pep	DEAD::Helicase_C	141-315::386-462
1037	CGPG4078.pep	GST_C	180-283
1038	CGPG4079.pep	RRS1	1-178
1041	CGPG4104.pep	LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::Pkinase	30-67::95-117::119-141::143-165::167-188::367-633
1042	CGPG4127.pep	Hydrolase	43-239
1043	CGPG4135.pep	Pkinase::efhand::efhand::efhand::efhand	118-376::423-451::459-487::495-523::529-557
1044	CGPG4161.pep	Methyltransf_11	56-159
1045	CGPG4180.pep	GHMP_kinases_N::GHMP_kinases_C	152-219::382-466
1046	CGPG4191.pep	Bombesin	163-176
1047	CGPG4199.pep	Copine::zf-C3HC4	117-265::385-417
1048	CGPG4241.pep	GASA	3-99
1049	CGPG427.pep	Pkinase_Tyr	8-262
1050	CGPG4273.pep	dsrm::dsrm	2-68::88-153
1051	CGPG4301.pep	2OG-FeII_Oxy	151-253
1052	CGPG4303.pep	Rhodanese	13-114
1053	CGPG4313.pep	G6PD_N::G6PD_C	94-273::276-573
1054	CGPG4314.pep	AWPM-19	15-156
1055	CGPG4329.pep	AA_kinase	86-324
1056	CGPG4330.pep	PALP	53-341
1057	CGPG4338.pep	Proteasome	21-217
1058	CGPG4341.pep	Mito_carr::Mito_carr::Mito_carr	51-134::158-246::252-343
1061	CGPG4386.pep	GHMP_kinases_N::GHMP_kinases_C	113-171::245-338
1062	CGPG4394.pep	p450	39-503
1063	CGPG4396.pep	p450	39-504
1064	CGPG4400.pep	p450	32-474
1065	CGPG4401.pep	p450	47-511
1066	CGPG441.pep	Chalcone	14-225
1067	CGPG4413.pep	p450	32-547
1068	CGPG4427.pep	p450	57-530
1069	CGPG4446.pep	p450	43-504
1070	CGPG4448.pep	p450	32-486
1071	CGPG4474.pep	Band_7	33-212
1072	CGPG4482.pep	peroxidase	47-291
1073	CGPG4511.pep	VHS::GAT	36-166::227-315
1074	CGPG4517.pep	Aldo_ket_red	5-292
1075	CGPG4551.pep	XS::XH	30-148::418-557
1077	CGPG4567.pep	SRF-TF	3-53
1078	CGPG4586.pep	RRM_1	8-73
1079	CGPG4600.pep	zf-CCCH::zf-CCCH	13-39::149-174
1080	CGPG4631.pep	Cl_2::Cl_2::Cl_3::Cl_2::Cl_3::Cl_2	88-115::209-237::264-292::319-349::404-433::566-

PEP Seq ID No.	Construct ID	Pfam module	Position
			595
1081	CGPG4642.pep	Pyridoxal_deC	63-412
1082	CGPG4645.pep	Cu-oxidase_3::Cu-oxidase::Cu-oxidase_2	31-148::157-325::414-551
1083	CGPG4646.pep	efhand_like::PI-PLC-X::PI-PLC-Y::C2	27-105::108-251::332-450::471-563
1084	CGPG4649.pep	SMP::SMP::SMP	14-72::130-191::195-256
1085	CGPG4653.pep	DUF1637	47-280
1087	CGPG4668.pep	SYF2	140-306
1088	CGPG469.pep	tRNA-synt_1c::tRNA-synt_1c_C	213-518::520-697
1089	CGPG4708.pep	Sulfate_transp::STAS	172-482::505-623
1090	CGPG4712.pep	Mem_trans	9-565
1091	CGPG4714.pep	DUF231	248-423
1092	CGPG4719.pep	Response_reg::CCT	37-153::417-461
1093	CGPG473.pep	Peptidase_M16::Peptidase_M16_C	88-234::239-423
1094	CGPG4734.pep	Auxin_inducible	19-119
1095	CGPG4736.pep	Pribosyltran	225-358
1096	CGPG474.pep	Pro_isomerase	6-172
1102	CGPG4850.pep	PMEI	25-174
1103	CGPG4868.pep	PsbW	1-133
1104	CGPG4871.pep	Tryp_alpha_amyl	52-133
1105	CGPG488.pep	Histone	19-92
1106	CGPG4908.pep	WD40::WD40	16-54::63-101
1108	CGPG4921.pep	WD40::WD40::WD40::WD40::WD40	202-241::253-291::295-333::337-375::472-510
1109	CGPG4954.pep	LRRNT_2::LRR_1::LRR_1::LRR_1	33-72::76-98::100-122::124-145
1110	CGPG4956.pep	Asp	141-482
1111	CGPG4959.pep	adh_short	15-183
1112	CGPG4965.pep	zf-A20::zf-AN1	15-39::114-154
1113	CGPG4970.pep	GASA	1-108
1114	CGPG4980.pep	U-box::Arm::Arm	73-147::246-287::288-328
1115	CGPG4982.pep	F-box::FBA_1	2-49::209-387
1116	CGPG4985.pep	CPSase_sm_chain::GATase	56-203::245-422
1117	CGPG4990.pep	LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::LR R_1	73-112::139-161::163-183::187-206::210-232::306-328
1118	CGPG4991.pep	PALP	164-473
1119	CGPG5007.pep	adh_short	21-192
1120	CGPG5015.pep	adh_short	13-126
1121	CGPG5026.pep	adh_short	46-221
1122	CGPG5029.pep	adh_short	80-252

PEP Seq ID No.	Construct ID	Pfam module	Position
1123	CGPG5046.pep	LRRNT_2::LRR_1::LRR_1::LRR_1::Pkinase	30-70::99-121::123-145::147-169::306-573
1124	CGPG508.pep	Na_sulph_symp	93-563
1125	CGPG5103.pep	C2	9-95
1126	CGPG511.pep	Rho_GDI	22-222
1127	CGPG5121.pep	efhand_like::PI-PLC-X::PI-PLC-Y::C2	30-113::116-258::331-449::470-562
1128	CGPG5126.pep	peroxidase	54-303
1129	CGPG5136.pep	p450	35-500
1130	CGPG5146.pep	p450	45-501
1131	CGPG5149.pep	p450	31-478
1132	CGPG5181.pep	Pkinase	278-540
1133	CGPG52.pep	Sugar_tr	32-472
1134	CGPG5206.pep	UDPG_MGDP_dh_N::UDPG_MGDP_dh::UDPG_MGDP_dh_C	2-200::209-306::328-452
1136	CGPG5232.pep	Pkinase	71-328
1137	CGPG5239.pep	Copine	136-284
1138	CGPG5246.pep	ABC_tran::ABC_tran	94-285::406-571
1139	CGPG525.pep	CDP-OH_P_transf	138-277
1140	CGPG5268.pep	CDC50	65-368
1141	CGPG5272.pep	AA_permease	153-621
1142	CGPG5333.pep	MFS_1	55-468
1143	CGPG5338.pep	RRM_1::RRM_1	8-77::108-178
1144	CGPG5341.pep	F-box	33-79
1145	CGPG5369.pep	AA_kinase::ACT::ACT	83-366::400-469::477-539
1146	CGPG5372.pep	SBF	142-322
1147	CGPG5380.pep	Tyr-DNA_phospho	137-581
1148	CGPG5386.pep	MOSC_N::MOSC	4-132::138-296
1149	CGPG5396.pep	PALP	9-297
1150	CGPG5397.pep	PDT	128-307
1151	CGPG5421.pep	Pre-SET::SET	119-267::269-414
1152	CGPG5433.pep	MtN3_slv::MtN3_slv	10-98::132-218
1153	CGPG5439.pep	LANC_like	69-410
1154	CGPG5453.pep	Mito_carr::Mito_carr::Mito_carr	14-116::125-215::233-334
1155	CGPG5456.pep	Mito_carr::Mito_carr::Mito_carr	78-175::183-279::283-373
1156	CGPG5483.pep	PP2C	66-369
1157	CGPG5492.pep	FAE1_CUT1_RppA::ACP_syn_III_C	61-358::372-456
1158	CGPG5508.pep	RCC1::RCC1::RCC1::RCC1::FYVE::DZC	310-359::413-462::478-526::582-630::633-701::967-1002
1159	CGPG5520.pep	TFIID-31kDa	19-152
1160	CGPG5525.pep	PRMT5	187-679

PEP Seq ID No.	Construct ID	Pfam module	Position
1161	CGPG5526.pep	Pkinase	8-295
1162	CGPG5530.pep	Response_reg	504-644
1163	CGPG5534.pep	Y_phosphatase2	3-169
1164	CGPG5537.pep	TBP::TBP	62-147::152-238
1165	CGPG5545.pep	CBS::CBS	35-175::193-318
1166	CGPG5555.pep	PALP::Thr_dehydrat_C::Thr_dehydrat_C	104-396::409-498::504-591
1167	CGPG5558.pep	Cyclin_N::Cyclin_C	212-337::339-458
1168	CGPG5559.pep	Suc_Fer-like	69-311
1169	CGPG5562.pep	Suc_Fer-like	59-308
1170	CGPG5592.pep	Rho_GDI	35-239
1171	CGPG5625.pep	LRR_1::LRR_1::Pkinase	433-455::457-476::586-854
1172	CGPG5631.pep	WAK::Pkinase	171-270::390-658
1173	CGPG5632.pep	Aldedh	29-491
1174	CGPG5636.pep	Aldedh	15-476
1175	CGPG5637.pep	Aldedh	32-493
1176	CGPG564.pep	Aminotran_3	80-413
1177	CGPG5641.pep	Aldedh	14-472
1178	CGPG5650.pep	Aminotran_3	32-374
1179	CGPG5653.pep	Aminotran_1_2	32-383
1180	CGPG5657.pep	Aldedh	15-474
1181	CGPG5671.pep	Gp_dh_N::Gp_dh_C	3-151::156-313
1182	CGPG5672.pep	Glutaminase	24-309
1183	CGPG5677.pep	PGAM	4-165
1184	CGPG5678.pep	Aminotran_1_2	35-385
1185	CGPG5679.pep	TIM	7-248
1186	CGPG5682.pep	Aminotran_3	31-384
1187	CGPG5691.pep	PK::PK_C	5-350::362-478
1188	CGPG5699.pep	Gp_dh_N::Gp_dh_C	3-155::160-316
1189	CGPG5705.pep	NDK	3-137
1190	CGPG5722.pep	Pyrophosphatase	17-175
1191	CGPG5734.pep	Cyclin_N::Cyclin_C	232-357::359-478
1192	CGPG5736.pep	PP2C	22-280
1193	CGPG5744.pep	CDC50	63-365
1194	CGPG5747.pep	Pkinase	23-318
1195	CGPG5748.pep	NAD_Gly3P_dh_N::NAD_Gly3P_dh_C	84-259::282-432
1198	CGPG577.pep	Thioredoxin	7-111
1199	CGPG5780.pep	AA_permease	68-517
1200	CGPG5793.pep	AA_permease	285-815
1201	CGPG5795.pep	Sugar_tr	123-565
1202	CGPG5796.pep	Sugar_tr	43-466
1203	CGPG5800.pep	AA_permease	86-546
1204	CGPG5812.pep	Pkinase	22-273
1205	CGPG5815.pep	Pkinase	9-272
1206	CGPG5838.pep	Pkinase	1-264
1207	CGPG5844.pep	Pkinase	78-351
1208	CGPG5846.pep	Pkinase	99-376

PEP Seq ID No.	Construct ID	Pfam module	Position
1209	CGPG5851.pep	Pkinase	121-401
1210	CGPG5857.pep	Pkinase	19-273
1211	CGPG5862.pep	Pkinase	181-447
1212	CGPG5863.pep	Pkinase	52-330
1213	CGPG5867.pep	Pkinase	142-423
1214	CGPG5872.pep	Pkinase	18-288
1215	CGPG5913.pep	ADH_N::ADH_zinc_N	28-109::140-284
1217	CGPG5961.pep	Biotin_lipoyl::2-oxoacid_dh	94-167::232-462
1219	CGPG5968.pep	Pribosyltran	36-172
1220	CGPG5984.pep	Pkinase	115-443
1221	CGPG5991.pep	LRR_1	266-293
1222	CGPG6006.pep	RRM_1	19-89
1223	CGPG6015.pep	La::RRM_1::RRM_3	14-85::118-188::302-404
1224	CGPG603.pep	G6PD_N::G6PD_C	35-222::224-508
1225	CGPG6046.pep	Str_synth	179-266
1226	CGPG6063.pep	Nramp	88-451
1228	CGPG6092.pep	SRPRB	53-235
1229	CGPG6104.pep	Steroid_dh	111-262
1230	CGPG6106.pep	LEA_5	1-92
1231	CGPG6111.pep	PP-binding	48-115
1232	CGPG6113.pep	UQ_con	7-143
1234	CGPG6132.pep	SMP::SMP::SMP	13-71::135-196::200-262
1236	CGPG6147.pep	Redoxin	74-234
1237	CGPG6152.pep	Arf	5-177
1238	CGPG6154.pep	Ion_trans_2::Ion_trans_2	122-204::253-328
1239	CGPG6170.pep	UQ_con	19-158
1240	CGPG6171.pep	Thioredoxin	120-227
1241	CGPG6177.pep	AIG1	39-236
1242	CGPG6181.pep	Peptidase_C12	2-214
1243	CGPG6188.pep	Di19	10-217
1244	CGPG6202.pep	TB2_DP1_HVA22	9-106
1245	CGPG6217.pep	MFS_1::Sugar_tr	40-460::90-507
1246	CGPG623.pep	SATase_N::Hexapep::Hexapep::Hexapep	45-149::203-220::229-246::247-264
1247	CGPG6239.pep	Sugar_tr	27-458
1248	CGPG6244.pep	Pkinase	39-303
1249	CGPG6254.pep	Pkinase::NAF	12-264::292-352
1251	CGPG627.pep	His_biosynth	48-289
1252	CGPG6271.pep	Pkinase	191-527
1253	CGPG6278.pep	Pkinase	86-368
1254	CGPG6288.pep	Pkinase_Tyr	207-463
1255	CGPG6309.pep	PRA1	29-181
1256	CGPG633.pep	DCP1	12-134
1257	CGPG635.pep	Epimerase	9-273
1258	CGPG6350.pep	LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1	32-78::105-127::129-151::153-175::177-196

PEP Seq ID No.	Construct ID	Pfam module	Position
1260	CGPG6365.pep	Aminotran_1_2	32-384
1261	CGPG6374.pep	Gln-synt_C	120-380
1262	CGPG6377.pep	PGK	2-383
1263	CGPG638.pep	PA	51-163
1264	CGPG6397.pep	F420_oxidored	3-246
1265	CGPG6398.pep	Enolase_N::Enolase_C	5-135::140-430
1266	CGPG6408.pep	ADH_N::ADH_zinc_N	25-155::185-350
1267	CGPG6415.pep	NTP_transferase	4-271
1268	CGPG6421.pep	ADH_N::ADH_zinc_N	25-148::179-318
1269	CGPG6442.pep	iPGM_N::Metalloenzyme	2-363::373-488
1270	CGPG6443.pep	PGK	1-391
1271	CGPG6446.pep	PK::PK_C	106-453::467-587
1272	CGPG6453.pep	Aminotran_1_2	41-402
1273	CGPG6466.pep	ADH_N::ADH_zinc_N	27-153::184-322
1274	CGPG6467.pep	Aminotran_3	19-366
1275	CGPG6470.pep	PfkB	1-302
1276	CGPG6475.pep	NTP_transferase	10-282
1277	CGPG6477.pep	NDK	4-138
1278	CGPG6478.pep	PfkB	1-308
1279	CGPG6493.pep	AA_permease	94-561
1280	CGPG6506.pep	Asparaginase	140-457
1281	CGPG6518.pep	NAD_binding_2::6PGD	2-173::177-467
1282	CGPG6546.pep	PK::PK_C	3-348::360-477
1283	CGPG6556.pep	Aminotran_1_2	124-475
1284	CGPG6562.pep	Aminotran_3	116-451
1285	CGPG6566.pep	NTP_transferase::Hexapep::Hexapep::Hexapep	95-350::398-415::432-449::460-477
1286	CGPG6571.pep	Aldedh	102-560
1287	CGPG6576.pep	Aldedh	99-565
1288	CGPG659.pep	G_glu_transpept	48-567
1289	CGPG6594.pep	Pyr_redox_2::Pyr_redox_dim	94-419::450-559
1290	CGPG6603.pep	Pyr_redox_2::Pyr_redox_dim	94-397::427-538
1291	CGPG6608.pep	Pyr_redox_2::Pyr_redox_dim	93-394::423-532
1292	CGPG6634.pep	DUF177	100-252
1294	CGPG6667.pep	Invertase_neut	89-577
1295	CGPG6678.pep	Biotin_lipoyl::E3_binding::2-oxoacid_dh	92-165::225-263::281-512
1296	CGPG6695.pep	Pyr_redox_2::Pyr_redox_dim	97-408::436-545
1297	CGPG6699.pep	Ribul_P_3_epim	94-295
1298	CGPG670.pep	Fer2	60-135
1299	CGPG6705.pep	PK::PK_C::PEP-utilizers	89-433::445-559::584-663
1300	CGPG6735.pep	FBPase	93-413
1301	CGPG6744.pep	Glycolytic	92-429

PEP Seq ID No.	Construct ID	Pfam module	Position
1302	CGPG6753.pep	Pyr_redox_2	236-538
1303	CGPG6757.pep	Aminotran_1_2	127-473
1305	CGPG6806.pep	mTERF	110-457
1306	CGPG6811.pep	GH3	10-567
1308	CGPG6815.pep	V-ATPase_G	5-109
1310	CGPG6830.pep	Tbf5	13-76
1313	CGPG6864.pep	DUF620	190-438
1314	CGPG6877.pep	Hep_59	101-194
1315	CGPG6878.pep	Peptidase_M24	107-343
1316	CGPG6880.pep	PAP_fibrillin	72-232
1318	CGPG6887.pep	AIG2	12-111
1319	CGPG6895.pep	zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC	166-183::208-225::325-342::356-373
1322	CGPG6912.pep	zf-DHHC	149-213
1324	CGPG6942.pep	RNA_pol_Rpb5_N::RNA_pol_Rpb5_C	18-108::149-222
1327	CGPG6969.pep	2OG-FeII_Oxy	223-323
1329	CGPG6991.pep	Glyco_tran_28_C	11-170
1332	CGPG7037.pep	Sad1_UNC	313-416
1333	CGPG7051.pep	PAP_fibrillin	84-236
1336	CGPG7086.pep	SFT2	44-163
1339	CGPG7136.pep	NOI	1-72
1340	CGPG7147.pep	RAMP4	1-64
1343	CGPG7222.pep	Branch	49-274
1345	CGPG7233.pep	eIF2A	214-407
1346	CGPG7238.pep	UPF0185	5-80
1347	CGPG7253.pep	ArfGap	15-117
1348	CGPG7255.pep	Arf	5-177
1349	CGPG7286.pep	ORC2	20-345
1350	CGPG7292.pep	Methyltransf_11	38-135
1351	CGPG7301.pep	ELFV_dehydrog_N::ELFV_dehydrog	57-187::202-445
1352	CGPG7305.pep	Pyr_redox_2	126-386
1353	CGPG7307.pep	Gp_dh_N::Gp_dh_C	11-163::168-324
1354	CGPG7318.pep	MIP	1-142
1355	CGPG7319.pep	SNF5	172-400
1356	CGPG7345.pep	Lipase_3	97-237
1357	CGPG7368.pep	MT-A70	516-676
1358	CGPG7385.pep	FHA	32-107
1359	CGPG7391.pep	G-patch	6-50
1360	CGPG7405.pep	TATA_RF	1-216
1361	CGPG7415.pep	IU_nuc_hydro	1-312
1362	CGPG7416.pep	GLTP	52-233
1363	CGPG7420.pep	PRK	95-302
1366	CGPG7445.pep	PRC::PRC	92-169::171-252
1367	CGPG7465.pep	Aldedh	103-567

PEP Seq ID No.	Construct ID	Pfam module	Position
1368	CGPG7473.pep	F-box::WD40::WD40::WD40::WD40	273-320::412-449::453-493::520-556::560-598
1369	CGPG7489.pep	UBX	295-376
1370	CGPG7492.pep	Pescadillo_N::BRCT	8-291::337-414
1371	CGPG7499.pep	G-alpha	423-839
1374	CGPG7508.pep	Smg4_UPF3	1-170
1375	CGPG7509.pep	SSB	76-187
1376	CGPG7511.pep	GFA	31-125
1377	CGPG7515.pep	Brix	45-297
1378	CGPG7521.pep	G-alpha	23-386
1380	CGPG7547.pep	PAD_porph	14-368
1381	CGPG7561.pep	Snf7	17-187
1382	CGPG7562.pep	Sybindin	6-136
1383	CGPG7563.pep	DUF887	40-279
1384	CGPG7567.pep	Peptidase_M18	14-461
1389	CGPG7597.pep	zf-AN1	76-116
1390	CGPG7606.pep	Pentapeptide::Pentapeptide	127-166::172-211
1392	CGPG7637.pep	DAD	2-112
1393	CGPG7649.pep	YgbB	71-227
1394	CGPG7658.pep	Cyclase	56-256
1395	CGPG7664.pep	Aldedh	9-463
1396	CGPG7666.pep	Aminotran_1_2	26-392
1397	CGPG7668.pep	DUF594	660-719
1399	CGPG7746.pep	UPF0153	49-139
1400	CGPG7747.pep	Copine	113-261
1401	CGPG7752.pep	zf-MYND::PDCD2_C	176-214::241-403
1403	CGPG7770.pep	YTH	263-353
1406	CGPG7778.pep	Ubic_methyltran	51-304
1407	CGPG7786.pep	KH_1	141-193
1408	CGPG7788.pep	YIF1	34-263
1410	CGPG78.pep	Hpt	46-132
1411	CGPG783.pep	RRM_1::RRM_2	288-354::682-778
1412	CGPG7832.pep	STT3	26-670
1413	CGPG7841.pep	ECH::3HCDH_N::3HCDH	17-186::311-490::492-585
1414	CGPG7845.pep	LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::LR R_1	23-64::97-119::121-143::145-167::169-192::194-216
1415	CGPG7847.pep	Peptidase_C48	30-225
1417	CGPG7853.pep	LRR_1::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1	142-164::166-188::190-212::214-235::258-280::282-304::306-325
1418	CGPG7857.pep	Response_reg::CCT	79-195::689-736
1420	CGPG7869.pep	Acyltransferase	117-264
1421	CGPG7891.pep	FMO-like	11-427
1422	CGPG7892.pep	FMO-like	9-416
1423	CGPG7906.pep	Skp1_POZ::Skp1	4-65::76-153

PEP Seq ID No.	Construct ID	Pfam module	Position
1424	CGPG7924.pep	AAA	249-462
1428	CGPG7964.pep	IQ::IQ	108-128::130-150
1429	CGPG7968.pep	FMO-like	13-427
1431	CGPG7972.pep	FAD_binding_3	5-372
1432	CGPG7982.pep	Tetraspannin	4-241
1433	CGPG7985.pep	BT1	57-469
1434	CGPG7993.pep	F-box::LRR_2	27-73::278-304
1435	CGPG8.pep	p450	39-503
1436	CGPG80.pep	Pkinase	68-328
1437	CGPG8001.pep	FMO-like	13-433
1438	CGPG8009.pep	BT1	41-511
1440	CGPG8049.pep	BT1	1-436
1441	CGPG806.pep	Synaptobrevin	127-215
1442	CGPG8060.pep	DUF829	161-425
1446	CGPG81.pep	Pkinase	70-330
1452	CGPG8125.pep	zf-C3HC4	46-86
1453	CGPG8134.pep	DEK_C::SWIB::SWIB	3-57::197-272::305-382
1455	CGPG8156.pep	PCI	258-362
1456	CGPG8159.pep	ACT::ACT	77-141::308-374
1458	CGPG8179.pep	WD40::WD40	138-177::264-305
1459	CGPG8193.pep	Ank::Ank	22-54::55-87
1460	CGPG8197.pep	F-box	5-53
1461	CGPG8198.pep	F-box::Kelch_2::Kelch_1::Kelch_2	74-119::174-221::223-278::329-369
1463	CGPG8211.pep	F-box::WD40::WD40::WD40::WD40::WD40::WD40	182-229::292-328::332-368::372-408::411-449::453-538::542-578
1464	CGPG823.pep	Ribosomal_S6e	1-129
1466	CGPG8238.pep	PGI	49-537
1467	CGPG8260.pep	Iso_dh	4-331
1468	CGPG8267.pep	Pyridoxal_deC	34-383
1469	CGPG8268.pep	NIR_SIR_ferr::NIR_SIR::NIR_SIR_ferr	62-131::164-345::360-432
1470	CGPG8274.pep	PK::PK_C	5-347::363-476
1471	CGPG8277.pep	Rib_5-P_isom_A	55-229
1472	CGPG8279.pep	F-box::Kelch_1::Kelch_1	22-69::124-168::170-213
1475	CGPG8408.pep	DUF641	60-192
1476	CGPG842.pep	Redoxin	5-162
1477	CGPG8421.pep	NOSIC::Nop	121-173::213-361
1478	CGPG8435.pep	NHL::NHL	55-82::115-142
1479	CGPG8440.pep	U-box::Arm::Arm::Arm::Arm	242-316::372-413::455-495::496-537::538-578
1480	CGPG8453.pep	ACT::ACT::ACT::ACT	36-99::129-201::265-332::343-406

PEP Seq ID No.	Construct ID	Pfam module	Position
1481	CGPG8470.pep	DUF89	18-357
1482	CGPG8479.pep	F-box::Kelch_1::Kelch_1	31-78::124-170::172-216
1483	CGPG8489.pep	F-box::FBA_1	3-50::219-384
1484	CGPG8498.pep	Fcf1	87-185
1485	CGPG85.pep	Aa_trans	19-454
1486	CGPG8501.pep	PPR::PPR::PPR::PPR::PPR::PPR	149-183::189-223::224-258::259-292::330-364::401-435
1487	CGPG8507.pep	DUF588	31-185
1488	CGPG8509.pep	Pro_isomerase	80-233
1490	CGPG8517.pep	B_lectin::S_locus_glycop::PAN_2	84-197::211-339::356-422
1492	CGPG852.pep	Ribosomal_S5::Ribosomal_S5_C	94-160::177-250
1498	CGPG8567.pep	TB2_DP1_HVA22	3-98
1499	CGPG8580.pep	Caleosin	22-195
1501	CGPG8590.pep	DUF579	42-289
1503	CGPG8594.pep	DUF584	1-139
1504	CGPG8606.pep	Exo_endo_phos	56-327
1505	CGPG8628.pep	DUF584	32-203
1507	CGPG8654.pep	PPR::PPR	73-107::141-175
1508	CGPG8668.pep	TATA_RF	1-200
1511	CGPG8702.pep	RPE65	50-622
1514	CGPG8748.pep	AAA	201-357
1516	CGPG8770.pep	DUF543	1-73
1517	CGPG8777.pep	DUF569::DUF569	1-144::227-368
1519	CGPG8786.pep	DUF260	13-113
1520	CGPG8792.pep	Abhydrolase_3	75-290
1521	CGPG8801.pep	PPR::PPR::PPR::PPR::PPR::PPR::PPR	110-144::145-179::181-215::216-250::251-285::286-320::321-355
1522	CGPG8809.pep	Ribosomal_L12	141-208
1523	CGPG8840.pep	CLP_protease	115-289
1524	CGPG8850.pep	PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR	189-223::224-258::297-331::332-366::367-401::402-436::437-470::471-505::506-540::542-576
1525	CGPG8870.pep	Fructosamin_kin	54-339
1526	CGPG8871.pep	DUF59	36-115
1527	CGPG8872.pep	RRM_1::RRM_1::RRM_1	138-205::214-288::308-378
1528	CGPG8876.pep	SFT2	43-162
1529	CGPG8902.pep	RNA_pol_I_A49	31-415
1530	CGPG8904.pep	G-alpha	103-447
1531	CGPG8907.pep	DnaJ::zf-C2H2	4-67::338-362
1532	CGPG8914.pep	Whi5	181-205

PEP Seq ID No.	Construct ID	Pfam module	Position
1533	CGPG8931.pep	eIF3_subunit	1-231
1534	CGPG8933.pep	MACPF	126-322
1535	CGPG8935.pep	Pkinase	33-324
1536	CGPG8938.pep	Pkinase	71-348
1537	CGPG8944.pep	PH	31-134
1538	CGPG895.pep	DUF231	251-427
1540	CGPG8961.pep	Dus	59-392
1541	CGPG8963.pep	Pkinase	319-612
1542	CGPG898.pep	DUF231	276-448
1543	CGPG8993.pep	AAA::Rep_fac_C	44-226::237-326
1544	CGPG8994.pep	Brix	86-262
1545	CGPG8995.pep	Porin_3	54-326
1546	CGPG90.pep	MFS_1	68-427
1547	CGPG900.pep	DUF231	241-420
1548	CGPG9002.pep	CTP_transf_1	51-382
1549	CGPG9009.pep	Alg6_Algl8	22-514
1550	CGPG9011.pep	OTCace_N::OTCace	70-212::216-369
1551	CGPG9012.pep	FolB	16-129
1552	CGPG9017.pep	DMRL_synthase	70-213
1553	CGPG9025.pep	Glyco_transf_29	105-355
1554	CGPG9026.pep	Adap_comp_sub	160-428
1555	CGPG9032.pep	RNase_PH::RNase_PH_C	33-165::194-262
1556	CGPG9040.pep	SecI	35-592
1557	CGPG9044.pep	Syntaxin::SNARE	57-163::256-318
1558	CGPG9048.pep	RNase_PH	15-135
1559	CGPG9049.pep	Glycos_transf_1	244-442
1560	CGPG9058.pep	RNA_pol_N	1-60
1561	CGPG906.pep	PPR::PPR	210-244::245-279
1562	CGPG9070.pep	ATP-synt_C::ATP-synt_C	20-85::104-169
1564	CGPG9084.pep	CH::EB1	15-116::209-255
1566	CGPG9098.pep	DUF6::DUF6	32-165::208-337
1567	CGPG9099.pep	Rick_17kDa_Anti	66-110
1568	CGPG9110.pep	RALF	53-118
1569	CGPG9119.pep	PTR2	115-504
1570	CGPG9125.pep	ThiF	98-233
1571	CGPG913.pep	PPR::PPR::PPR::PPR::PPR::PPR	86-120::188-222::223-257::289-323::325-358::361-395
1574	CGPG9164.pep	DHBP_synthase::GTP_cyclohydro2	5-202::207-377
1575	CGPG9172.pep	Sec61_beta	32-77
1577	CGPG9185.pep	DUF423	10-115
1578	CGPG9187.pep	Pkinase	12-263
1579	CGPG9190.pep	DREPP	2-203
1580	CGPG9193.pep	GATase::GMP_synt_C	11-196::432-524
1581	CGPG9195.pep	DUF1279	91-193
1582	CGPG9203.pep	Na_H_Exchanger::TrkA_C	13-391::418-486
1583	CGPG9209.pep	GATase::GMP_synt_C	9-197::423-515

PEP Seq ID No.	Construct ID	Pfam module	Position
1585	CGPG9210.pep	Na_H_Exchange::TrkA_N	10-376::402-517
1586	CGPG9211.pep	NTP_transferase::MannoseP_isomer	7-296::307-473
1587	CGPG9212.pep	ATP-synt_ab_N::ATP-synt_ab	22-87::143-353
1588	CGPG9216.pep	DHBP_synthase::GTP_cyclohydro2	10-206::211-379
1589	CGPG9226.pep	Carboxyl_trans	40-541
1590	CGPG9252.pep	CDC48_N::AAA::AAA	28-114::245-429::518-705
1592	CGPG9289.pep	LSM	10-75
1593	CGPG9298.pep	p450	41-516
1594	CGPG9299.pep	ThiF	82-227
1597	CGPG9316.pep	ATP-grasp_2::Ligase_CoA	34-242::285-421
1598	CGPG9317.pep	DUF163	43-196
1599	CGPG9318.pep	Aldo_ket_red	20-285
1600	CGPG9324.pep	Pkinase	158-706
1601	CGPG9357.pep	Raffinose_syn	7-757
1602	CGPG965.pep	Hexapep::Hexapep::Hexapep::Hexapep	70-87::119-136::142-159::160-177
1603	CGPG970.pep	NPH3	206-441
1604	CGPG982.pep	p450	29-495
1605	CGPG994.pep	adh_short	50-218
1606	CGPG996.pep	DUF866	1-167

Table 21.

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
806	CGPG1023.pep	Na_Ca_ex	104	255	94.2	4.10E-25
806	CGPG1023.pep	Na_Ca_ex	299	432	128.4	2.10E-35
807	CGPG1038.pep	DUF1475	1	256	709.5	2.40E-210
812	CGPG1113.pep	DUF1350	49	370	64.5	3.50E-16
814	CGPG1163.pep	DSPc	49	182	103	9.00E-28
817	CGPG1205.pep	Arm	94	134	37.2	5.90E-08
817	CGPG1205.pep	Arm	135	175	32.9	1.10E-06
817	CGPG1205.pep	Arm	176	216	31.7	2.60E-06
817	CGPG1205.pep	Arm	258	298	18.8	0.02
818	CGPG1207.pep	NifU_N	26	105	78.3	2.60E-20
820	CGPG1223.pep	ATP_bind_1	7	250	425.9	5.70E-125
821	CGPG1250.pep	DUF822	12	160	323.6	3.70E-94
824	CGPG1319.pep	DUF599	10	233	464.2	1.70E-136
826	CGPG1329.pep	Cullin	29	631	966.3	1.20E-287
827	CGPG133.pep	Cyclin_N	168	293	232.6	9.20E-67
827	CGPG133.pep	Cyclin_C	295	422	170.5	4.50E-48
828	CGPG1332.pep	Cullin	12	646	709.1	3.20E-210
829	CGPG1343.pep	LRR_1	389	412	9.2	6

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
830	CGPG1348.pep	Lectin_legB	27	248	295	1.40E-85
830	CGPG1348.pep	Pkinase	334	604	170.2	5.30E-48
830	CGPG1348.pep	Pkinase_Tyr	334	604	134.8	2.50E-37
831	CGPG1349.pep	B_lectin	65	171	95.8	1.30E-25
831	CGPG1349.pep	S_locus_glycop	180	312	105	2.40E-28
831	CGPG1349.pep	PAN_1	327	404	38.5	2.40E-08
831	CGPG1349.pep	Pkinase	484	731	167.1	4.70E-47
831	CGPG1349.pep	Pkinase_Tyr	484	731	89.6	1.00E-23
832	CGPG1373.pep	Pkinase	12	291	356.7	3.80E-104
833	CGPG1377.pep	Pkinase	4	258	247.3	3.30E-71
834	CGPG1412.pep	TPP_enzyme_N	45	221	286.5	5.40E-83
834	CGPG1412.pep	TPP_enzyme_M	243	376	82.9	1.00E-21
834	CGPG1412.pep	TPP_enzyme_C	431	578	35	3.50E-09
835	CGPG1421.pep	Biotin_lipoyl	76	149	85.1	2.30E-22
835	CGPG1421.pep	E3_binding	180	218	58.5	2.20E-14
835	CGPG1421.pep	2-oxoacid_dh	249	480	358.5	1.10E-104
836	CGPG1426.pep	Aminotran_1_2	48	432	525	8.50E-155
837	CGPG1433.pep	HI0933_like	44	365	-247.2	0.002
837	CGPG1433.pep	GIDA	45	369	-206	0.00022
837	CGPG1433.pep	Pyr_redox_2	45	359	241.5	1.90E-69
837	CGPG1433.pep	Pyr_redox	216	313	115.7	1.40E-31
837	CGPG1433.pep	Pyr_redox_dim	388	497	213.1	6.50E-61
838	CGPG1453.pep	malic	171	360	412.3	7.10E-121
838	CGPG1453.pep	Malic_M	362	615	468.8	7.10E-138
839	CGPG1454.pep	Ribul_P_3_epim	7	207	331.5	1.50E-96
840	CGPG1463.pep	Alpha-amylase	26	361	224.3	2.80E-64
840	CGPG1463.pep	Alpha-amyl_C2	362	422	132.9	9.40E-37
841	CGPG1464.pep	Glycolytic	11	358	852.5	2.20E-253
842	CGPG1471.pep	IF4E	1	198	287.7	2.20E-83
843	CGPG1481.pep	Pkinase	4	260	292.7	7.40E-85
844	CGPG1499.pep	PPDK_N	85	445	654.8	6.90E-194
844	CGPG1499.pep	PEP-utilizers	496	586	165.9	1.10E-46
844	CGPG1499.pep	PEP-utilizers_C	598	955	688.7	4.30E-204
845	CGPG150.pep	p450	30	507	128	2.80E-35
846	CGPG1536.pep	CRAL_TRIO_N	73	142	46.4	1.00E-10
846	CGPG1536.pep	CRAL_TRIO	159	345	145.1	2.00E-40
847	CGPG1539.pep	UPF0139	8	107	215.9	9.60E-62
848	CGPG155.pep	GlutR_N	100	251	288.2	1.70E-83
848	CGPG155.pep	Shikimate_DH	255	407	208.3	1.80E-59

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
848	CGPG155.pep	GlutR_dimer	420	526	180.6	4.10E-51
849	CGPG1583.pep	PLAC8	88	187	176.8	5.40E-50
850	CGPG1588.pep	PLAC8	75	214	156.3	8.30E-44
851	CGPG16.pep	MIP	14	235	396.7	3.60E-116
852	CGPG1609.pep	MMR_HSR1	54	190	118.9	1.50E-32
852	CGPG1609.pep	DUF933	337	420	205.4	1.40E-58
853	CGPG1629.pep	WD40	262	301	31.1	4.20E-06
853	CGPG1629.pep	WD40	307	346	36	1.30E-07
853	CGPG1629.pep	WD40	356	395	41.3	3.50E-09
854	CGPG1637.pep	DUF220	163	262	148.5	1.80E-41
855	CGPG1653.pep	TBC	200	415	-15	1.60E-06
856	CGPG1658.pep	Nodulin-like	14	260	516.4	3.30E-152
857	CGPG1663.pep	MFS_1	22	387	26.3	0.00011
857	CGPG1663.pep	BT1	30	441	633.5	1.90E-187
858	CGPG1682.pep	OPT	62	685	688.8	4.20E-204
859	CGPG1701.pep	Glyco_transf_8	76	299	13.5	1.90E-07
861	CGPG1724.pep	Per1	63	334	669.6	2.50E-198
862	CGPG1726.pep	DUF607	111	291	388.6	9.80E-114
863	CGPG1736.pep	Usp	1	148	64.9	2.70E-16
864	CGPG1741.pep	mTERF	279	627	84.1	4.40E-22
865	CGPG1783.pep	PB1	42	132	85	2.40E-22
866	CGPG1790.pep	PB1	59	153	109.6	9.20E-30
868	CGPG1845.pep	TFIID-18kDa	29	119	145.5	1.50E-40
869	CGPG1855.pep	CAF1	3	227	330.9	2.30E-96
870	CGPG1870.pep	Pkinase	406	572	74.8	2.90E-19
871	CGPG1879.pep	Pkinase	21	277	357.5	2.30E-104
871	CGPG1879.pep	Pkinase_Tyr	21	275	88.2	2.60E-23
871	CGPG1879.pep	NAF	303	364	98.7	1.80E-26
872	CGPG1886.pep	Pkinase	26	325	337.3	2.70E-98
873	CGPG1903.pep	F-box	320	367	41	4.30E-09
874	CGPG1905.pep	F-box	4	51	41.7	2.60E-09
874	CGPG1905.pep	Kelch_2	104	150	27.4	5.30E-05
874	CGPG1905.pep	Kelch_2	255	306	41.6	2.80E-09
875	CGPG1914.pep	F-box	42	89	39.8	9.50E-09
875	CGPG1914.pep	Kelch_2	170	215	26.3	0.00011
875	CGPG1914.pep	Kelch_1	170	215	52.2	1.80E-12
875	CGPG1914.pep	Kelch_2	217	266	31.3	3.60E-06
875	CGPG1914.pep	Kelch_1	217	266	59.1	1.50E-14
876	CGPG193.pep	WD40	167	205	28.5	2.40E-05
876	CGPG193.pep	WD40	268	307	24.4	0.00041
876	CGPG193.pep	WD40	318	355	25.1	0.00026
876	CGPG193.pep	WD40	363	401	39.3	1.40E-08
877	CGPG1939.pep	F-box	66	113	36.2	1.20E-07
877	CGPG1939.pep	WD40	152	190	26.2	0.00012
877	CGPG1939.pep	WD40	247	283	33.6	7.10E-07
877	CGPG1939.pep	WD40	325	364	22.9	0.0012
878	CGPG1949.pep	YDG_SRA	360	519	403.5	3.10E-118
878	CGPG1949.pep	Pre-SET	543	639	156.7	6.30E-44
878	CGPG1949.pep	SET	641	771	178.4	1.80E-50

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
879	CGPG1959.pep	PHD	34	82	54.3	4.30E-13
879	CGPG1959.pep	SET	208	344	105.1	2.20E-28
880	CGPG197.pep	p450	40	481	169.2	1.10E-47
882	CGPG1981.pep	MT-A70	476	636	321.4	1.70E-93
883	CGPG1999.pep	Nfu_N	78	193	145	2.10E-40
883	CGPG1999.pep	NifU	221	291	121.1	3.30E-33
884	CGPG2.pep	p450	71	531	324.3	2.20E-94
885	CGPG2006.pep	RCC1	32	82	31.3	3.60E-06
885	CGPG2006.pep	RCC1	138	187	40	8.30E-09
885	CGPG2006.pep	RCC1	190	239	33	1.10E-06
885	CGPG2006.pep	RCC1	294	343	42.8	1.20E-09
886	CGPG2010.pep	Aminotran_5	74	448	681.1	8.90E-202
886	CGPG2010.pep	Beta_elim_lyase	77	389	-107.3	0.0021
887	CGPG2011.pep	X8	128	209	153.5	5.60E-43
888	CGPG2014.pep	Pkinase	28	282	339.2	7.00E-99
888	CGPG2014.pep	NAF	341	398	86.9	6.70E-23
889	CGPG2023.pep	Zip	48	352	371	1.90E-108
890	CGPG2026.pep	Pkinase	38	324	328.9	9.40E-96
892	CGPG2064.pep	Cenp-O	123	201	107.6	3.80E-29
894	CGPG2077.pep	NAC	88	147	81.6	2.50E-21
894	CGPG2077.pep	UBA	195	232	29.8	1.00E-05
895	CGPG2095.pep	AARP2CN	228	309	121.6	2.30E-33
895	CGPG2095.pep	DUF663	483	780	615.9	3.60E-182
896	CGPG2105.pep	Pkinase	102	404	274.6	2.00E-79
896	CGPG2105.pep	Pkinase_C	422	471	39.6	1.10E-08
897	CGPG2108.pep	Response_reg	64	180	86.8	6.80E-23
897	CGPG2108.pep	CCT	442	484	65	2.50E-16
898	CGPG2111.pep	polyprenyl_synt	74	324	156.4	7.70E-44
899	CGPG2124.pep	UIM	139	156	20.6	0.0057
899	CGPG2124.pep	efhand	220	248	35.7	1.70E-07
900	CGPG2125.pep	Glutaredoxin	44	110	40.7	5.10E-09
903	CGPG2134.pep	ATP-synt_G	8	122	170.2	5.30E-48
904	CGPG2139.pep	DUF1517	94	391	709.7	2.10E-210
905	CGPG2140.pep	PhzC-PhzF	6	282	506.2	3.90E-149
907	CGPG2163.pep	Cupin_1	5	157	105.9	1.20E-28
907	CGPG2163.pep	Cupin_1	190	339	83.5	6.80E-22
908	CGPG2165.pep	ClpS	76	152	131.5	2.40E-36
910	CGPG2218.pep	Spc97_Spc98	67	555	750.6	1.10E-222
912	CGPG2225.pep	EMP24_GP25L	52	116	55.6	1.80E-13
913	CGPG2229.pep	Subtilisin_N	30	106	89.6	9.70E-24
913	CGPG2229.pep	Peptidase_S8	119	600	52.4	1.60E-12
913	CGPG2229.pep	PA	362	461	112	1.80E-30
915	CGPG2254.pep	Ribosomal_L7Ae	20	114	110.3	5.80E-30
916	CGPG2268.pep	CBS	66	188	107.9	3.00E-29
917	CGPG227.pep	p450	37	494	364.2	2.20E-106
918	CGPG2312.pep	UPF0061	75	571	361.8	1.20E-105

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
919	CGPG2315.pep	SIP1	286	514	516.5	3.10E-152
920	CGPG2316.pep	PTPA	98	396	696.6	1.80E-206
922	CGPG235.pep	p450	74	510	203.2	6.10E-58
923	CGPG2358.pep	Branch	54	278	425.3	8.70E-125
925	CGPG2361.pep	Metallophos	9	231	49.7	1.00E-11
926	CGPG2365.pep	Epimerase	3	211	-25.8	3.00E-05
927	CGPG2372.pep	Pribosyltran	29	169	147.9	2.70E-41
928	CGPG2374.pep	Aminotran_5	8	370	-92.6	4.20E-07
929	CGPG2377.pep	DAO	69	466	243.4	5.00E-70
929	CGPG2377.pep	FAD_binding_2	69	457	-124	0.0024
930	CGPG2387.pep	H_PPase	1	239	-188.8	6.80E-17
931	CGPG2389.pep	Carb_anhydrase	52	275	117.4	4.30E-32
932	CGPG2395.pep	AA_permease	89	526	-4.2	1.30E-05
934	CGPG2409.pep	DUF862	17	154	252.5	8.90E-73
935	CGPG2410.pep	GST_N	5	79	68.9	1.70E-17
935	CGPG2410.pep	GST_C	101	204	30.1	8.30E-06
937	CGPG2416.pep	Hin1	97	235	198.2	2.10E-56
938	CGPG2441.pep	Sina	5	205	188	2.30E-53
939	CGPG2450.pep	adh_short	112	282	-38.2	0.004
939	CGPG2450.pep	Epimerase	114	371	191.5	2.10E-54
939	CGPG2450.pep	3Beta_HSD	115	402	-72.8	1.60E-05
939	CGPG2450.pep	NAD_binding_4	116	342	-62.5	9.90E-05
940	CGPG2451.pep	RALF	57	129	123.9	4.80E-34
941	CGPG2492.pep	Pribosyltran	80	216	144.1	3.80E-40
942	CGPG2495.pep	ADH_N	25	134	143.2	7.40E-40
942	CGPG2495.pep	ADH_zinc_N	165	307	138	2.60E-38
943	CGPG2506.pep	PfkB	5	289	140.3	5.30E-39
944	CGPG2515.pep	TIM	5	244	446	5.10E-131
945	CGPG2531.pep	Redoxin	4	160	43.7	6.60E-10
945	CGPG2531.pep	AhpC-TSA	5	138	178.9	1.30E-50
946	CGPG2581.pep	SelR	12	133	289.3	7.50E-84
947	CGPG2584.pep	zf-Tim10_DDP	22	86	115.1	2.00E-31
948	CGPG2592.pep	TFIID_30kDa	30	80	135.9	1.10E-37
949	CGPG2612.pep	RRM_1	65	132	43	1.10E-09
949	CGPG2612.pep	RRM_1	150	225	83.9	5.30E-22
949	CGPG2612.pep	RRM_1	275	343	53.1	9.50E-13
950	CGPG2660.pep	B3	17	115	77.8	3.60E-20
950	CGPG2660.pep	B3	211	301	61.4	3.10E-15
951	CGPG2663.pep	NTF2	15	131	168.8	1.50E-47
951	CGPG2663.pep	RRM_1	295	365	45.3	2.10E-10
952	CGPG2679.pep	ACBP	104	190	54.4	3.80E-13
952	CGPG2679.pep	Ank	265	297	49.7	1.00E-11
952	CGPG2679.pep	Ank	298	330	35.5	2.00E-07
953	CGPG2696.pep	zf-C3HC4	177	218	43.9	5.70E-10
954	CGPG2772.pep	Mov34	6	136	111.9	1.90E-30
955	CGPG2773.pep	Asp	161	498	-73.6	3.00E-09
956	CGPG281.pep	TLC	91	574	1046.6	0

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
957	CGPG2846.pep	Aldo_ket_red	49	365	248	2.00E-71
958	CGPG2852.pep	Glyoxalase	9	132	37.4	5.20E-08
960	CGPG2870.pep	zf-CCCH	111	137	49.3	1.30E-11
960	CGPG2870.pep	zf-CCCH	159	185	34.6	3.60E-07
960	CGPG2870.pep	zf-CCCH	205	231	49.5	1.20E-11
960	CGPG2870.pep	zf-CCCH	347	373	29.9	9.60E-06
960	CGPG2870.pep	zf-CCCH	393	419	42.5	1.50E-09
961	CGPG2877.pep	LAG1	96	307	430.8	1.90E-126
962	CGPG289.pep	Cellulose_synt	243	1060	2161.1	0
963	CGPG2924.pep	zf-C3HC4	138	179	33.3	9.10E-07
964	CGPG2947.pep	Acetyltransf_1	265	343	50.6	5.60E-12
964	CGPG2947.pep	Bromodomain	460	548	122.2	1.50E-33
965	CGPG2963.pep	C1_3	80	108	35.7	1.60E-07
965	CGPG2963.pep	C1_2	81	108	37.1	6.10E-08
965	CGPG2963.pep	C1_1	122	172	14.3	0.00058
965	CGPG2963.pep	C1_3	136	164	36.3	1.10E-07
965	CGPG2963.pep	C1_3	194	223	27.2	6.10E-05
965	CGPG2963.pep	C1_2	305	335	40.5	5.90E-09
965	CGPG2963.pep	C1_3	391	420	40.1	8.20E-09
965	CGPG2963.pep	C1_3	494	522	25.6	0.00018
965	CGPG2963.pep	C1_3	551	581	27.5	5.00E-05
965	CGPG2963.pep	C1_2	552	581	47.9	3.50E-11
966	CGPG2987.pep	SRF-TF	11	66	24.4	0.00041
968	CGPG3045.pep	Pkinase	82	356	192	1.50E-54
968	CGPG3045.pep	Pkinase_Tyr	82	356	254	3.20E-73
969	CGPG3046.pep	AAA	121	305	245.7	1.00E-70
970	CGPG3060.pep	p450	67	528	268	1.90E-77
971	CGPG3075.pep	Auxin_inducible	7	106	121	3.50E-33
972	CGPG310.pep	Cyclin_N	18	143	31.8	1.80E-07
973	CGPG3103.pep	PPR	14	48	10.8	0.4
973	CGPG3103.pep	PPR	49	83	37.5	4.80E-08
973	CGPG3103.pep	PPR	84	118	38.8	1.90E-08
973	CGPG3103.pep	PPR	119	153	11.8	0.31
973	CGPG3103.pep	PPR	155	188	27.6	4.50E-05
973	CGPG3103.pep	PPR	189	223	34.8	3.20E-07
973	CGPG3103.pep	PPR	224	258	36.8	7.80E-08
973	CGPG3103.pep	PPR	259	293	7.7	0.95
973	CGPG3103.pep	PPR	295	329	39.7	1.10E-08
973	CGPG3103.pep	PPR	330	364	12.6	0.25
973	CGPG3103.pep	PPR	365	398	30.4	6.70E-06
974	CGPG315.pep	p450	29	491	346.8	3.80E-101
976	CGPG3189.pep	RRM_1	115	186	101.9	2.00E-27
976	CGPG3189.pep	RRM_1	209	280	104.9	2.40E-28
977	CGPG3204.pep	KH_1	45	104	53.4	7.60E-13
977	CGPG3204.pep	KH_1	151	221	58.6	2.20E-14
977	CGPG3204.pep	KH_1	318	381	44.6	3.60E-10
977	CGPG3204.pep	KH_2	318	365	11.3	0.019
977	CGPG3204.pep	KH_1	400	467	63.4	7.50E-16

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
977	CGPG3204.pep	KH_1	575	637	53.3	8.70E-13
978	CGPG3208.pep	MATH	26	153	47.1	6.00E-11
978	CGPG3208.pep	MATH	180	302	18.3	0.0015
979	CGPG3219.pep	2OG-Fell_Oxy	220	320	161.1	2.90E-45
980	CGPG3233.pep	PPR	160	194	21.8	0.0026
980	CGPG3233.pep	PPR	195	229	9.6	0.56
980	CGPG3233.pep	PPR	231	265	24.1	0.00052
980	CGPG3233.pep	PPR	266	300	40.7	5.10E-09
980	CGPG3233.pep	PPR	301	335	44	5.30E-10
980	CGPG3233.pep	PPR	336	370	45.3	2.20E-10
980	CGPG3233.pep	PPR	371	405	44.7	3.30E-10
980	CGPG3233.pep	PPR	406	440	7.4	1
981	CGPG3263.pep	Tryp_alpha_amyl	28	104	36.6	8.90E-08
982	CGPG3276.pep	ECH	11	186	48.1	3.10E-11
984	CGPG3282.pep	Steroid_dh	117	268	65.5	1.80E-16
985	CGPG3300.pep	zf-CCHC	55	72	30.4	6.40E-06
985	CGPG3300.pep	zf-CCHC	74	91	26.8	7.80E-05
985	CGPG3300.pep	zf-CCHC	93	110	29.8	1.00E-05
985	CGPG3300.pep	zf-CCHC	112	129	27.3	5.70E-05
985	CGPG3300.pep	zf-CCHC	138	155	23.8	0.00038
985	CGPG3300.pep	zf-CCHC	157	174	24.9	0.00026
985	CGPG3300.pep	zf-CCHC	229	246	26.5	9.70E-05
986	CGPG3318.pep	Peptidase_C14	3	416	273.4	4.50E-79
987	CGPG3319.pep	AP2	28	79	61.2	3.40E-15
988	CGPG3326.pep	zf-C3HC4	259	299	36.8	7.90E-08
989	CGPG3333.pep	PTS_2-RNA	48	239	409.9	3.80E-120
990	CGPG3338.pep	Alba	17	87	114.8	2.50E-31
991	CGPG3334.pep	Mov34	20	128	65.2	2.20E-16
992	CGPG3374.pep	peroxidase	42	286	426.2	4.60E-125
993	CGPG3402.pep	Dimerisation	30	89	93.1	8.60E-25
993	CGPG3402.pep	Methyltransf_2	99	342	287.6	2.40E-83
995	CGPG3413.pep	WD40	168	207	41.5	3.10E-09
995	CGPG3413.pep	U3_snoRNA_C	377	523	212.3	1.10E-60
996	CGPG3422.pep	WD40	241	279	41.2	3.70E-09
996	CGPG3422.pep	WD40	387	425	35.3	2.20E-07
996	CGPG3422.pep	WD40	519	559	21.9	0.0024
997	CGPG3436.pep	DUF6	17	144	33.8	6.40E-07
997	CGPG3436.pep	DUF6	198	327	50.4	6.10E-12
998	CGPG3539.pep	Mito_carr	11	105	84.6	3.10E-22
998	CGPG3539.pep	Mito_carr	113	209	115.7	1.40E-31
998	CGPG3539.pep	Mito_carr	214	303	102.2	1.60E-27
999	CGPG3550.pep	FAE1_CUT1_Rpp A	81	370	700.9	9.60E-208
999	CGPG3550.pep	Chal_sti_synt_C	327	470	2.6	0.0011

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
999	CGPG3550.pep	ACP_syn_III_C	384	468	21.5	8.90E-08
1000	CGPG3551.pep	Pkinase	9	273	145.9	1.10E-40
1001	CGPG3552.pep	Aldedh	23	441	176.5	6.90E-50
1002	CGPG3572.pep	FA_hydroxylase	112	225	133.2	7.50E-37
1004	CGPG3599.pep	DUF579	51	289	588.5	6.70E-174
1008	CGPG364.pep	zf-C3HC4	97	137	51	4.30E-12
1010	CGPG3679.pep	malic	36	179	249.1	9.50E-72
1010	CGPG3679.pep	Malic_M	181	418	426.3	4.30E-125
1010	CGPG3679.pep	PTA_PTB	446	768	457.7	1.60E-134
1011	CGPG3686.pep	Gp_dh_N	2	150	334.5	1.80E-97
1011	CGPG3686.pep	Gp_dh_C	155	312	388.4	1.10E-113
1012	CGPG3694.pep	LRRNT_2	38	78	64.2	4.40E-16
1012	CGPG3694.pep	LRR_1	106	128	11.7	2.1
1012	CGPG3694.pep	LRR_1	130	152	14.2	0.48
1012	CGPG3694.pep	LRR_1	154	176	16.7	0.089
1012	CGPG3694.pep	LRR_1	178	201	11.7	2.1
1012	CGPG3694.pep	Pkinase	312	583	104.7	2.80E-28
1012	CGPG3694.pep	Pkinase_Tyr	312	583	110.8	4.20E-30
1013	CGPG3696.pep	MMR_HSR1	136	257	120.2	5.90E-33
1014	CGPG3698.pep	WD40	104	142	31.6	2.80E-06
1014	CGPG3698.pep	WD40	234	272	22.5	0.0016
1015	CGPG3699.pep	adh_short	19	187	111.6	2.30E-30
1015	CGPG3699.pep	KR	19	203	-54.2	0.00034
1016	CGPG3702.pep	SAM_2	227	293	48.9	1.80E-11
1016	CGPG3702.pep	SAM_1	228	291	52.7	1.30E-12
1017	CGPG3703.pep	ENOD93	25	103	190.1	5.50E-54
1018	CGPG3707.pep	Pkinase	42	333	296.2	6.30E-86
1019	CGPG3710.pep	VQ	118	148	45.7	1.60E-10
1020	CGPG3730.pep	DUF167	140	216	99.7	9.10E-27
1021	CGPG3731.pep	DUF616	194	508	769.9	1.60E-228
1022	CGPG3734.pep	PRK	46	233	177.1	4.40E-50
1022	CGPG3734.pep	Pribosyltran	280	425	5.8	0.00083
1023	CGPG3745.pep	RRM_1	386	455	66.7	7.80E-17
1023	CGPG3745.pep	RRM_1	481	553	49.6	1.10E-11
1024	CGPG3764.pep	Response_reg	20	145	67.1	5.90E-17
1026	CGPG3851.pep	SH3_1	284	338	42.4	1.60E-09
1026	CGPG3851.pep	SH3_2	285	338	30.1	8.10E-06
1027	CGPG3911.pep	Aldedh	104	587	526.9	2.20E-155
1028	CGPG3948.pep	NAP	50	295	467.9	1.30E-137
1031	CGPG3996.pep	Inositol_P	6	267	363.1	4.60E-106
1032	CGPG4006.pep	ENT	1	74	140.3	5.60E-39
1033	CGPG4025.pep	SQS_PSY	44	331	455.8	5.80E-134
1034	CGPG4028.pep	Transket_pyr	398	565	233	6.90E-67
1034	CGPG4028.pep	Transketolase_C	579	702	152.2	1.40E-42
1035	CGPG403.pep	Pkinase	78	336	354.1	2.40E-103
1036	CGPG4041.pep	DEAD	141	315	219	1.10E-62
1036	CGPG4041.pep	Helicase_C	386	462	120.8	4.10E-33

PEP Seq ID No.	Construct ID	Pfam domain-name	Begin	Stop	score	E-value
1037	CGPG4078.pep	GST_C	180	283	23.1	0.00038
1038	CGPG4079.pep	RRS1	1	178	271.1	2.30E-78
1041	CGPG4104.pep	LRRNT_2	30	67	33.1	1.00E-06
1041	CGPG4104.pep	LRR_1	95	117	11.4	2.3
1041	CGPG4104.pep	LRR_1	119	141	15.5	0.21
1041	CGPG4104.pep	LRR_1	143	165	12.1	1.8
1041	CGPG4104.pep	LRR_1	167	188	16.1	0.13
1041	CGPG4104.pep	Pkinase	367	633	99.2	1.30E-26
1042	CGPG4127.pep	Hydrolase	43	239	93.3	7.50E-25
1043	CGPG4135.pep	Pkinase	118	376	357.9	1.70E-104
1043	CGPG4135.pep	efhand	423	451	41.5	3.10E-09
1043	CGPG4135.pep	efhand	459	487	24.8	0.00033
1043	CGPG4135.pep	efhand	495	523	30.6	5.70E-06
1043	CGPG4135.pep	efhand	529	557	40.3	7.00E-09
1044	CGPG4161.pep	Methyltransf_11	56	159	48	3.30E-11
1044	CGPG4161.pep	Methyltransf_12	56	157	31.8	2.40E-06
1045	CGPG4180.pep	GHMP_kinases_N	152	219	64.2	4.40E-16
1045	CGPG4180.pep	GHMP_kinases_C	382	466	58.4	2.50E-14
1046	CGPG4191.pep	Bombesin	163	176	14.8	0.16
1047	CGPG4199.pep	Copine	117	265	295.3	1.20E-85
1047	CGPG4199.pep	zf-C3HC4	385	417	16.2	0.0013
1048	CGPG4241.pep	GASA	3	99	179.9	6.40E-51
1049	CGPG427.pep	Pkinase	8	262	210	5.80E-60
1049	CGPG427.pep	Pkinase_Tyr	8	262	248	2.10E-71
1050	CGPG4273.pep	dsrm	2	68	70	7.70E-18
1050	CGPG4273.pep	dsrm	88	153	66.9	6.90E-17
1051	CGPG4301.pep	2OG-Fell_Oxy	151	253	97	6.00E-26
1052	CGPG4303.pep	Rhodanese	13	114	64.9	2.70E-16
1053	CGPG4313.pep	G6PD_N	94	273	322.4	8.50E-94
1053	CGPG4313.pep	G6PD_C	276	573	522.2	6.00E-154
1054	CGPG4314.pep	AWPM-19	15	156	335.5	9.50E-98
1055	CGPG4329.pep	AA_kinase	86	324	201.9	1.60E-57
1056	CGPG4330.pep	PALP	53	341	418.6	9.40E-123
1057	CGPG4338.pep	Proteasome	21	217	157.3	4.00E-44
1058	CGPG4341.pep	Mito_carr	51	134	78	3.10E-20
1058	CGPG4341.pep	Mito_carr	158	246	81.5	2.80E-21
1058	CGPG4341.pep	Mito_carr	252	343	82.8	1.10E-21
1061	CGPG4386.pep	GHMP_kinases_N	113	171	55.4	1.90E-13
1061	CGPG4386.pep	GHMP_kinases_C	245	338	29.5	1.20E-05
1062	CGPG4394.pep	p450	39	503	389.1	7.00E-114
1063	CGPG4396.pep	p450	39	504	355.6	8.20E-104
1064	CGPG4400.pep	p450	32	474	146.7	6.70E-41
1065	CGPG4401.pep	p450	47	511	383.1	4.60E-112
1066	CGPG441.pep	Chalcone	14	225	498.4	8.60E-147

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1067	CGPG4413.pep	p450	32	547	286.5	5.20E-83
1068	CGPG4427.pep	p450	57	530	225.1	1.60E-64
1069	CGPG4446.pep	p450	43	504	275.6	1.00E-79
1070	CGPG4448.pep	p450	32	486	332.3	8.70E-97
1071	CGPG4474.pep	Band_7	33	212	136.8	5.90E-38
1072	CGPG4482.pep	peroxidase	47	291	314.4	2.10E-91
1073	CGPG4511.pep	VHS	36	166	58.2	2.80E-14
1073	CGPG4511.pep	GAT	227	315	18.1	2.80E-05
1074	CGPG4517.pep	Aldo_ket_red	5	292	460.6	2.10E-135
1075	CGPG4551.pep	XS	30	148	250.2	4.60E-72
1075	CGPG4551.pep	XH	418	557	251.6	1.70E-72
1077	CGPG4567.pep	SRF-TF	3	53	85.4	1.80E-22
1078	CGPG4586.pep	RRM_1	8	73	41.5	2.90E-09
1079	CGPG4600.pep	zf-CCCH	13	39	35.4	2.00E-07
1079	CGPG4600.pep	zf-CCCH	149	174	24.1	0.00053
1080	CGPG4631.pep	C1_3	87	115	40.8	4.80E-09
1080	CGPG4631.pep	C1_2	88	115	41.2	3.80E-09
1080	CGPG4631.pep	C1_3	208	237	39.4	1.30E-08
1080	CGPG4631.pep	C1_2	209	237	45.5	1.80E-10
1080	CGPG4631.pep	C1_3	264	292	48.3	2.70E-11
1080	CGPG4631.pep	C1_2	319	349	41.5	2.90E-09
1080	CGPG4631.pep	C1_3	404	433	55.4	1.90E-13
1080	CGPG4631.pep	C1_2	566	595	40.8	4.70E-09
1081	CGPG4642.pep	Pyridoxal_deC	63	412	126.5	7.90E-35
1082	CGPG4645.pep	Cu-oxidase_3	31	148	198.9	1.30E-56
1082	CGPG4645.pep	Cu-oxidase	157	325	222.3	1.10E-63
1082	CGPG4645.pep	Cu-oxidase_2	414	551	155.1	2.00E-43
1083	CGPG4646.pep	efhand_like	27	105	109.6	9.60E-30
1083	CGPG4646.pep	PI-PLC-X	108	251	143.6	5.60E-40
1083	CGPG4646.pep	PI-PLC-Y	332	450	93.9	5.20E-25
1083	CGPG4646.pep	C2	471	563	82.3	1.50E-21
1084	CGPG4649.pep	SMP	14	72	102	1.90E-27
1084	CGPG4649.pep	SMP	130	191	127.6	3.60E-35
1084	CGPG4649.pep	SMP	195	256	62.7	1.30E-15
1085	CGPG4653.pep	DUF1637	47	280	366.2	5.30E-107
1087	CGPG4668.pep	SYF2	140	306	301.7	1.50E-87
1088	CGPG469.pep	tRNA-synt_1c	213	518	428.8	7.70E-126
1088	CGPG469.pep	tRNA-synt_1c_C	520	697	229.4	8.00E-66
1089	CGPG4708.pep	Sulfate_transp	172	482	504.9	9.60E-149
1089	CGPG4708.pep	STAS	505	623	142.8	9.80E-40
1090	CGPG4712.pep	Mem_trans	9	565	715.3	4.30E-212
1091	CGPG4714.pep	DUF231	248	423	336.8	3.80E-98
1092	CGPG4719.pep	Response_reg	37	153	91.6	2.50E-24
1092	CGPG4719.pep	CCT	417	461	85	2.40E-22
1093	CGPG473.pep	Peptidase_M16	88	234	184	3.90E-52
1093	CGPG473.pep	Peptidase_M16_C	239	423	161.3	2.70E-45

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1094	CGPG4734.pep	Auxin_inducible	19	119	55.4	1.90E-13
1095	CGPG4736.pep	Pribosyltran	225	358	139.9	7.20E-39
1096	CGPG474.pep	Pro_isomerase	6	172	426	5.20E-125
1102	CGPG4850.pep	PMEI	25	174	138.8	1.60E-38
1103	CGPG4868.pep	PsbW	1	133	333	5.40E-97
1104	CGPG4871.pep	Tryp_alpha_amyl	52	133	98.3	2.30E-26
1105	CGPG488.pep	Histone	19	92	110.5	5.00E-30
1105	CGPG488.pep	CBFD_NFYB_HM F	25	89	20.6	0.002
1106	CGPG4908.pep	WD40	16	54	22.1	0.002
1106	CGPG4908.pep	WD40	63	101	22.2	0.002
1108	CGPG4921.pep	WD40	202	241	26.8	7.90E-05
1108	CGPG4921.pep	WD40	253	291	41.3	3.40E-09
1108	CGPG4921.pep	WD40	295	333	33.5	7.40E-07
1108	CGPG4921.pep	WD40	337	375	47.3	5.20E-11
1108	CGPG4921.pep	WD40	472	510	25.7	0.00017
1109	CGPG4954.pep	LRRNT_2	33	72	38.6	2.20E-08
1109	CGPG4954.pep	LRR_1	76	98	9	6.5
1109	CGPG4954.pep	LRR_1	100	122	12.7	1.4
1109	CGPG4954.pep	LRR_1	124	145	8	10
1110	CGPG4956.pep	Asp	141	482	-61.6	5.90E-10
1111	CGPG4959.pep	adh_short	15	183	94.8	2.80E-25
1111	CGPG4959.pep	KR	16	200	-64.6	0.0014
1112	CGPG4965.pep	zf-A20	15	39	32.9	1.20E-06
1112	CGPG4965.pep	zf-AN1	114	154	64.5	3.50E-16
1113	CGPG4970.pep	GASA	1	108	225.3	1.40E-64
1114	CGPG4980.pep	U-box	73	147	95.4	1.80E-25
1114	CGPG4980.pep	Arm	246	287	20.1	0.0085
1114	CGPG4980.pep	Arm	288	328	22.4	0.0017
1115	CGPG4982.pep	F-box	2	49	42.8	1.20E-09
1115	CGPG4982.pep	FBA_1	209	387	313.8	3.20E-91
1116	CGPG4985.pep	CPSase_sm_chai n	56	203	298.5	1.30E-86
1116	CGPG4985.pep	GATase	245	422	238.5	1.50E-68
1117	CGPG4990.pep	LRRNT_2	73	112	28.9	1.90E-05
1117	CGPG4990.pep	LRR_1	139	161	13.9	0.59
1117	CGPG4990.pep	LRR_1	163	183	13.6	0.73
1117	CGPG4990.pep	LRR_1	187	206	13.9	0.62
1117	CGPG4990.pep	LRR_1	210	232	8.2	9.1
1117	CGPG4990.pep	LRR_1	306	328	9.8	4.6
1118	CGPG4991.pep	PALP	164	473	254.7	2.00E-73
1119	CGPG5007.pep	adh_short	21	192	65.9	1.30E-16
1119	CGPG5007.pep	Epimerase	23	271	-42.4	0.00053
1120	CGPG5015.pep	adh_short	13	126	-22.3	0.00036
1121	CGPG5026.pep	adh_short	46	221	22.7	4.00E-07
1122	CGPG5029.pep	adh_short	80	252	70.7	4.90E-18
1123	CGPG5046.pep	LRRNT_2	30	70	51.2	3.60E-12

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1123	CGPG5046.pep	LRR_1	99	121	9.2	6
1123	CGPG5046.pep	LRR_1	123	145	10.5	3.4
1123	CGPG5046.pep	LRR_1	147	169	15.8	0.16
1123	CGPG5046.pep	Pkinase_Tyr	306	573	109.4	1.10E-29
1123	CGPG5046.pep	Pkinase	306	573	145.6	1.40E-40
1124	CGPG508.pep	Na_sulph_symp	93	563	619	4.30E-183
1125	CGPG5103.pep	C2	9	95	64.8	3.00E-16
1126	CGPG511.pep	Rho_GDI	22	222	43.8	6.70E-12
1127	CGPG5121.pep	efhand_like	30	113	41.4	3.20E-09
1127	CGPG5121.pep	PI-PLC-X	116	258	140.7	4.20E-39
1127	CGPG5121.pep	PI-PLC-Y	331	449	88.6	2.00E-23
1127	CGPG5121.pep	C2	470	562	81.3	3.00E-21
1128	CGPG5126.pep	peroxidase	54	303	312	1.10E-90
1129	CGPG5136.pep	p450	35	500	88.4	2.30E-23
1130	CGPG5146.pep	p450	45	501	209	1.10E-59
1131	CGPG5149.pep	p450	31	478	306.7	4.40E-89
1132	CGPG5181.pep	Pkinase	278	540	112.7	1.10E-30
1133	CGPG52.pep	Sugar_tr	32	472	-75.6	5.00E-06
1134	CGPG5206.pep	UDPG_MGDP_dh_N	2	200	331.6	1.40E-96
1134	CGPG5206.pep	UDPG_MGDP_dh	209	306	189.1	1.10E-53
1134	CGPG5206.pep	UDPG_MGDP_dh_C	328	452	183.7	4.70E-52
1136	CGPG5232.pep	Pkinase	71	328	295.9	8.10E-86
1137	CGPG5239.pep	Copine	136	284	331.7	1.30E-96
1138	CGPG5246.pep	ABC_tran	94	285	132.8	9.70E-37
1138	CGPG5246.pep	ABC_tran	406	571	146.2	9.10E-41
1139	CGPG525.pep	CDP-OH_P_transf	138	277	132.1	1.60E-36
1140	CGPG5268.pep	CDC50	65	368	681.3	7.50E-202
1141	CGPG5272.pep	AA_permease	153	621	569.2	4.10E-168
1142	CGPG5333.pep	MFS_1	55	468	76	1.30E-19
1142	CGPG5333.pep	Sugar_tr	89	515	21.1	8.20E-10
1143	CGPG5338.pep	RRM_1	8	77	90.6	5.10E-24
1143	CGPG5338.pep	RRM_1	108	178	82.6	1.30E-21
1144	CGPG5341.pep	F-box	33	79	31.7	2.60E-06
1145	CGPG5369.pep	AA_kinase	83	366	213.4	5.30E-61
1145	CGPG5369.pep	ACT	400	469	20.9	0.0047
1145	CGPG5369.pep	ACT	477	539	9.7	0.71
1146	CGPG5372.pep	SBF	142	322	125.7	1.30E-34
1147	CGPG5380.pep	Tyr-DNA_phospho	137	581	1043.5	0
1148	CGPG5386.pep	MOSC_N	4	132	245	1.70E-70
1148	CGPG5386.pep	MOSC	138	296	66.5	8.80E-17
1149	CGPG5396.pep	PALP	9	297	469.3	5.00E-138
1150	CGPG5397.pep	PDT	128	307	347.9	1.70E-101
1151	CGPG5421.pep	Pre-SET	119	267	14.7	2.20E-05
1151	CGPG5421.pep	SET	269	414	151.3	2.60E-42

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1152	CGPG5433.pep	MtN3_slv	10	98	144.3	3.30E-40
1152	CGPG5433.pep	MtN3_slv	132	218	145.4	1.50E-40
1153	CGPG5439.pep	LANC_like	69	410	459.8	3.60E-135
1154	CGPG5453.pep	Mito_carr	14	116	102	1.80E-27
1154	CGPG5453.pep	Mito_carr	125	215	98.8	1.60E-26
1154	CGPG5453.pep	Mito_carr	233	334	104	4.60E-28
1155	CGPG5456.pep	Mito_carr	78	175	128.9	1.50E-35
1155	CGPG5456.pep	Mito_carr	183	279	100.4	5.50E-27
1155	CGPG5456.pep	Mito_carr	283	373	91.5	2.70E-24
1156	CGPG5483.pep	PP2C	66	369	92.8	1.10E-24
1157	CGPG5492.pep	FAE1_CUT1_RppA	61	358	683.7	1.40E-202
1157	CGPG5492.pep	Chal_sti_synt_C	312	458	-1.1	0.0023
1157	CGPG5492.pep	ACP_syn_III_C	372	456	4.3	5.30E-06
1158	CGPG5508.pep	RCC1	310	359	23.9	0.00059
1158	CGPG5508.pep	RCC1	413	462	39	1.70E-08
1158	CGPG5508.pep	RCC1	478	526	32.5	1.50E-06
1158	CGPG5508.pep	RCC1	582	630	28.9	1.80E-05
1158	CGPG5508.pep	FYVE	633	701	44.7	3.30E-10
1158	CGPG5508.pep	DZC	967	1002	74	4.80E-19
1159	CGPG5520.pep	TFIID-31kDa	19	152	319.5	6.10E-93
1160	CGPG5525.pep	PRMT5	187	679	1145.5	0
1161	CGPG5526.pep	Pkinase	8	295	335.7	8.20E-98
1162	CGPG5530.pep	Response_reg	504	644	85.4	1.80E-22
1163	CGPG5534.pep	Y_phosphatase2	3	169	331.1	2.00E-96
1164	CGPG5537.pep	TBP	62	147	136.9	5.90E-38
1164	CGPG5537.pep	TBP	152	238	157.1	4.70E-44
1165	CGPG5545.pep	CBS	35	175	26.9	7.50E-05
1165	CGPG5545.pep	CBS	193	318	80.7	4.60E-21
1166	CGPG555.pep	PALP	104	396	325.8	7.90E-95
1166	CGPG555.pep	Thr_dehydrat_C	409	498	105.4	1.70E-28
1166	CGPG555.pep	Thr_dehydrat_C	504	591	126.4	8.20E-35
1167	CGPG5558.pep	Cyclin_N	212	337	208.5	1.60E-59
1167	CGPG5558.pep	Cyclin_C	339	458	141	3.30E-39
1168	CGPG5559.pep	Suc_Fer-like	69	311	370.5	2.70E-108
1169	CGPG5562.pep	Suc_Fer-like	59	308	60.7	4.90E-15
1170	CGPG5592.pep	Rho_GDI	35	239	71.2	3.40E-18
1171	CGPG5625.pep	LRR_1	433	455	20.5	0.0061
1171	CGPG5625.pep	LRR_1	457	476	11.7	2.1
1171	CGPG5625.pep	Pkinase	586	854	138.6	1.80E-38
1171	CGPG5625.pep	Pkinase_Tyr	587	854	123.3	7.00E-34
1172	CGPG5631.pep	WAK	171	270	186.4	7.00E-53
1172	CGPG5631.pep	Pkinase_Tyr	390	660	146.8	5.90E-41
1172	CGPG5631.pep	Pkinase	390	658	159.5	9.00E-45
1173	CGPG5632.pep	Aldedh	29	491	831.6	4.30E-247

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1174	CGPG5636.pep	Aldedh	15	476	794.5	6.10E-236
1175	CGPG5637.pep	Aldedh	32	493	925.9	1.70E-275
1176	CGPG564.pep	Aminotran_3	80	413	293.8	3.30E-85
1177	CGPG5641.pep	Aldedh	14	472	720.7	1.10E-213
1178	CGPG5650.pep	Aminotran_3	32	374	476.5	3.50E-140
1179	CGPG5653.pep	Aminotran_1_2	32	383	268.2	1.70E-77
1180	CGPG5657.pep	Aldedh	15	474	701.6	5.70E-208
1181	CGPG5671.pep	Gp_dh_N	3	151	356.5	4.40E-104
1181	CGPG5671.pep	Gp_dh_C	156	313	350.1	3.90E-102
1182	CGPG5672.pep	Glutaminase	24	309	615.7	4.20E-182
1183	CGPG5677.pep	PGAM	4	165	177.7	3.10E-50
1184	CGPG5678.pep	Aminotran_1_2	35	385	204	3.60E-58
1185	CGPG5679.pep	TIM	7	248	398.9	8.00E-117
1186	CGPG5682.pep	Aminotran_3	31	384	432.3	7.00E-127
1187	CGPG5691.pep	PK	5	350	699.4	2.60E-207
1187	CGPG5691.pep	PK_C	362	478	198.7	1.40E-56
1188	CGPG5699.pep	Gp_dh_N	3	155	274	3.00E-79
1188	CGPG5699.pep	Gp_dh_C	160	316	231.1	2.60E-66
1189	CGPG5705.pep	NDK	3	137	354.1	2.40E-103
1190	CGPG5722.pep	Pyrophosphatase	17	175	197.9	2.50E-56
1191	CGPG5734.pep	Cyclin_N	232	357	208.7	1.40E-59
1191	CGPG5734.pep	Cyclin_C	359	478	150	6.70E-42
1192	CGPG5736.pep	PP2C	22	280	398.6	9.20E-117
1193	CGPG5744.pep	CDC50	63	365	697.3	1.20E-206
1194	CGPG5747.pep	Pkinase	23	318	294	2.90E-85
1195	CGPG5748.pep	NAD_Gly3P_dh_N	84	259	271	2.50E-78
1195	CGPG5748.pep	NAD_Gly3P_dh_C	282	432	282.8	6.90E-82
1198	CGPG577.pep	Thioredoxin	7	111	138.6	1.70E-38
1199	CGPG5780.pep	AA_permease	68	517	-83.3	0.0017
1200	CGPG5793.pep	AA_permease	285	815	226.5	5.90E-65
1201	CGPG5795.pep	Sugar_tr	123	565	30.1	3.70E-10
1202	CGPG5796.pep	Sugar_tr	43	466	340.9	2.30E-99
1202	CGPG5796.pep	MFS_1	47	417	74.4	3.70E-19
1203	CGPG5800.pep	AA_permease	86	546	489	5.90E-144
1204	CGPG5812.pep	Pkinase	22	273	334.2	2.30E-97
1204	CGPG5812.pep	Pkinase_Tyr	22	271	85.2	2.10E-22
1205	CGPG5815.pep	Pkinase	9	272	136.6	6.90E-38
1206	CGPG5838.pep	Pkinase	1	264	137.8	3.10E-38
1207	CGPG5844.pep	Pkinase	78	351	159.7	7.80E-45
1207	CGPG5844.pep	Pkinase_Tyr	78	351	122.7	1.10E-33
1208	CGPG5846.pep	Pkinase_Tyr	92	376	128.5	2.00E-35
1208	CGPG5846.pep	Pkinase	99	376	136.1	9.90E-38
1209	CGPG5851.pep	Pkinase_Tyr	121	401	138.2	2.30E-38
1209	CGPG5851.pep	Pkinase	121	401	139.5	9.70E-39
1210	CGPG5857.pep	Pkinase	19	273	159.1	1.20E-44

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1211	CGPG5862.pep	Pkinase	181	447	99.8	8.50E-27
1211	CGPG5862.pep	Pkinase_Tyr	181	447	95.5	1.70E-25
1212	CGPG5863.pep	Pkinase	52	330	86.5	8.60E-23
1213	CGPG5867.pep	Pkinase_Tyr	142	423	94.1	4.50E-25
1213	CGPG5867.pep	Pkinase	142	423	104	4.70E-28
1214	CGPG5872.pep	Pkinase_Tyr	18	288	125.8	1.20E-34
1214	CGPG5872.pep	Pkinase	18	288	153	7.90E-43
1215	CGPG5913.pep	ADH_N	28	109	95.4	1.80E-25
1215	CGPG5913.pep	ADH_zinc_N	140	284	151	3.30E-42
1217	CGPG5961.pep	Biotin_lipoyl	94	167	79.7	9.50E-21
1217	CGPG5961.pep	2-oxoacid_dh	232	462	478	1.20E-140
1219	CGPG5968.pep	Pribosyltran	36	172	151.2	2.90E-42
1220	CGPG5984.pep	Pkinase	115	443	246.8	4.70E-71
1221	CGPG5991.pep	LRR_1	266	293	9.3	5.7
1222	CGPG6006.pep	RRM_1	19	89	81.3	3.10E-21
1223	CGPG6015.pep	La	14	85	101.3	3.00E-27
1223	CGPG6015.pep	RRM_1	118	188	37.1	6.40E-08
1223	CGPG6015.pep	RRM_3	302	404	142.2	1.40E-39
1224	CGPG603.pep	G6PD_N	35	222	342	1.00E-99
1224	CGPG603.pep	G6PD_C	224	508	655.5	4.50E-194
1225	CGPG6046.pep	Str_synth	179	266	137.2	4.80E-38
1226	CGPG6063.pep	Nramp	88	451	733.6	1.40E-217
1228	CGPG6092.pep	SRPRB	53	235	117.6	3.70E-32
1229	CGPG6104.pep	Steroid_dh	111	262	337.7	2.00E-98
1230	CGPG6106.pep	LEA_5	1	92	209.3	9.10E-60
1231	CGPG6111.pep	PP-binding	48	115	75.2	2.10E-19
1232	CGPG6113.pep	UQ_con	7	143	205.4	1.40E-58
1234	CGPG6132.pep	SMP	13	71	121.1	3.30E-33
1234	CGPG6132.pep	SMP	135	196	113.9	4.90E-31
1234	CGPG6132.pep	SMP	200	262	105	2.30E-28
1236	CGPG6147.pep	Redoxin	74	234	181.5	2.20E-51
1236	CGPG6147.pep	AhpC-TSA	75	214	65.2	2.30E-16
1237	CGPG6152.pep	Arf	5	177	434.6	1.40E-127
1237	CGPG6152.pep	SRPRB	15	177	-23.5	7.20E-06
1237	CGPG6152.pep	Miro	19	129	20.4	0.00012
1237	CGPG6152.pep	Ras	19	179	-47.2	8.20E-06
1238	CGPG6154.pep	lon_trans_2	122	204	70	7.90E-18
1238	CGPG6154.pep	lon_trans_2	253	328	48.9	1.80E-11
1239	CGPG6170.pep	UQ_con	19	158	187.1	4.50E-53
1240	CGPG6171.pep	Thioredoxin	120	227	55.8	1.50E-13
1241	CGPG6177.pep	AlG1	39	236	212.9	7.60E-61
1241	CGPG6177.pep	MMR_HSR1	39	154	31.5	1.40E-06
1242	CGPG6181.pep	Peptidase_C12	2	214	311.3	1.80E-90
1243	CGPG6188.pep	Di19	10	217	479	6.00E-141
1244	CGPG6202.pep	TB2_DP1_HVA22	9	106	191.2	2.60E-54
1245	CGPG6217.pep	MFS_1	40	460	86.7	7.20E-23
1245	CGPG6217.pep	Sugar_tr	90	507	-7.3	1.10E-08
1246	CGPG623.pep	SATase_N	45	149	210	5.60E-60

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1246	CGPG623.pep	Hexapep	203	220	19.7	0.011
1246	CGPG623.pep	Hexapep	229	246	15.6	0.18
1246	CGPG623.pep	Hexapep	247	264	8	16
1247	CGPG6239.pep	Sugar_tr	27	458	289.4	7.10E-84
1247	CGPG6239.pep	MFS_1	32	413	90.4	5.50E-24
1248	CGPG6244.pep	Pkinase	39	303	81.8	2.10E-21
1249	CGPG6254.pep	Pkinase	12	264	310.9	2.40E-90
1249	CGPG6254.pep	Pkinase_Tyr	12	261	75.1	2.20E-19
1249	CGPG6254.pep	NAF	292	352	103	9.50E-28
1251	CGPG627.pep	His_biosynth	48	289	33.3	4.00E-12
1252	CGPG6271.pep	Pkinase	191	527	227.5	3.00E-65
1253	CGPG6278.pep	Pkinase_Tyr	86	365	146.9	5.60E-41
1253	CGPG6278.pep	Pkinase	86	368	158.7	1.60E-44
1254	CGPG6288.pep	Pkinase	207	463	183.7	4.70E-52
1254	CGPG6288.pep	Pkinase_Tyr	207	463	207.1	4.10E-59
1255	CGPG6309.pep	PRA1	29	181	233.1	6.50E-67
1256	CGPG633.pep	DCP1	12	134	258.3	1.70E-74
1257	CGPG635.pep	RmID_sub_bind	7	338	-145.7	0.00034
1257	CGPG635.pep	adh_short	7	177	-35.3	0.0026
1257	CGPG635.pep	Epimerase	9	273	226.4	6.40E-65
1257	CGPG635.pep	Polysacc_synt_2	9	319	-162.6	0.0011
1257	CGPG635.pep	3Beta_HSD	10	299	-79	3.60E-05
1257	CGPG635.pep	NAD_binding_4	11	240	-73	0.00047
1258	CGPG6350.pep	LRRNT_2	32	78	22.6	0.0015
1258	CGPG6350.pep	LRR_1	105	127	20.8	0.0051
1258	CGPG6350.pep	LRR_1	129	151	13.6	0.74
1258	CGPG6350.pep	LRR_1	153	175	13.4	0.84
1258	CGPG6350.pep	LRR_1	177	196	10.9	2.9
1260	CGPG6365.pep	Aminotran_1_2	32	384	403.5	3.20E-118
1261	CGPG6374.pep	Gln-synt_C	120	380	370.8	2.30E-108
1262	CGPG6377.pep	PGK	2	383	466.4	3.70E-137
1263	CGPG638.pep	PA	51	163	106	1.10E-28
1264	CGPG6397.pep	NAD_binding_2	1	157	-57	0.00027
1264	CGPG6397.pep	NAD_Gly3P_dh_N	2	121	-33.4	0.0009
1264	CGPG6397.pep	F420_oxidored	3	246	249.4	7.60E-72
1265	CGPG6398.pep	Enolase_N	5	135	212.4	1.10E-60
1265	CGPG6398.pep	Enolase_C	140	430	405	1.10E-118
1266	CGPG6408.pep	ADH_N	25	155	153.5	6.00E-43
1266	CGPG6408.pep	ADH_zinc_N	185	350	101.5	2.50E-27
1267	CGPG6415.pep	NTP_transferase	4	271	71.9	2.10E-18
1268	CGPG6421.pep	ADH_N	25	148	131.4	2.60E-36
1268	CGPG6421.pep	ADH_zinc_N	179	318	163.9	4.30E-46

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1269	CGPG6442.pep	iPGM_N	2	363	870.1	1.10E-258
1269	CGPG6442.pep	Metalloenzyme	373	488	191.4	2.20E-54
1270	CGPG6443.pep	PGK	1	391	727.8	7.60E-216
1271	CGPG6446.pep	PK	106	453	901.2	4.70E-268
1271	CGPG6446.pep	PK_C	467	587	259.4	7.60E-75
1272	CGPG6453.pep	Aminotran_1_2	41	402	404.3	1.80E-118
1273	CGPG6466.pep	ADH_N	27	153	129.2	1.20E-35
1273	CGPG6466.pep	ADH_zinc_N	184	322	135.3	1.80E-37
1274	CGPG6467.pep	Aminotran_3	19	366	361.7	1.20E-105
1275	CGPG6470.pep	PfkB	1	302	230	5.40E-66
1276	CGPG6475.pep	NTP_transferase	10	282	35.5	1.90E-09
1277	CGPG6477.pep	NDK	4	138	268.7	1.20E-77
1278	CGPG6478.pep	PfkB	1	308	70.6	5.30E-18
1279	CGPG6493.pep	AA_permease	94	561	594.7	9.00E-176
1280	CGPG6506.pep	Asparaginase	140	457	525.4	6.50E-155
1281	CGPG6518.pep	NAD_binding_2	2	173	303	5.60E-88
1281	CGPG6518.pep	6PGD	177	467	700.7	1.10E-207
1282	CGPG6546.pep	PK	3	348	560.5	1.80E-165
1282	CGPG6546.pep	PK_C	360	477	86.8	6.80E-23
1283	CGPG6556.pep	Aminotran_1_2	124	475	223.7	4.30E-64
1284	CGPG6562.pep	Aminotran_3	116	451	628.8	4.90E-186
1285	CGPG6566.pep	NTP_transferase	95	350	301.4	1.80E-87
1285	CGPG6566.pep	Hexapep	398	415	6.4	25
1285	CGPG6566.pep	Hexapep	432	449	16.6	0.097
1285	CGPG6566.pep	Hexapep	460	477	5.5	31
1286	CGPG6571.pep	Aldedh	102	560	720.7	1.10E-213
1287	CGPG6576.pep	Aldedh	99	565	569.2	4.30E-168
1288	CGPG659.pep	G_glu_transpept	48	567	839.1	2.30E-249
1289	CGPG6594.pep	DAO	94	524	-28.6	0.00082
1289	CGPG6594.pep	GIDA	94	429	-226.6	0.0045
1289	CGPG6594.pep	Pyr_redox_2	94	419	229.8	6.20E-66
1289	CGPG6594.pep	Pyr_redox	278	374	119.5	9.90E-33
1289	CGPG6594.pep	Pyr_redox_dim	450	559	194.7	2.20E-55
1290	CGPG6603.pep	Pyr_redox_2	94	397	239.1	9.60E-69
1290	CGPG6603.pep	Pyr_redox	257	350	101.2	3.10E-27
1290	CGPG6603.pep	Pyr_redox_dim	427	538	189.7	7.10E-54
1291	CGPG6608.pep	DAO	93	497	-33.9	0.0018
1291	CGPG6608.pep	Pyr_redox_2	93	394	145.4	1.60E-40
1291	CGPG6608.pep	Pyr_redox	256	348	92.2	1.70E-24
1291	CGPG6608.pep	Pyr_redox_dim	423	532	89.3	1.20E-23

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1292	CGPG6634.pep	DUF177	100	252	152.3	1.30E-42
1294	CGPG6667.pep	Invertase_neut	89	577	1313.4	0
1295	CGPG6678.pep	Biotin_lipoyl	92	165	75.9	1.30E-19
1295	CGPG6678.pep	E3_binding	225	263	51.7	2.60E-12
1295	CGPG6678.pep	2-oxoacid_dh	281	512	374.6	1.60E-109
1296	CGPG6695.pep	GIDA	97	428	-211.9	0.00053
1296	CGPG6695.pep	Pyr_redox_2	97	408	266.1	7.40E-77
1296	CGPG6695.pep	Pyr_redox	266	361	90.1	7.20E-24
1296	CGPG6695.pep	Pyr_redox_dim	436	545	184.3	3.20E-52
1297	CGPG6699.pep	Ribul_P_3_epim	94	295	491.3	1.20E-144
1298	CGPG670.pep	Fer2	60	135	120.2	6.20E-33
1299	CGPG6705.pep	PK	89	433	788.8	3.30E-234
1299	CGPG6705.pep	PK_C	445	559	164.3	3.30E-46
1299	CGPG6705.pep	PEP-utilizers	584	663	122.9	9.40E-34
1300	CGPG6735.pep	FBPase	93	413	525	8.60E-155
1301	CGPG6744.pep	Glycolytic	92	429	576.6	2.40E-170
1302	CGPG6753.pep	Pyr_redox	236	328	10.7	0.00051
1302	CGPG6753.pep	Pyr_redox_2	236	538	72.9	1.10E-18
1303	CGPG6757.pep	Aminotran_1_2	127	473	234	3.30E-67
1305	CGPG6806.pep	mTERF	110	457	550.4	2.00E-162
1306	CGPG6811.pep	GH3	10	567	1341	0
1308	CGPG6815.pep	V-ATPase_G	5	109	183.9	4.00E-52
1310	CGPG6830.pep	Tbf5	13	76	113.7	5.40E-31
1313	CGPG6864.pep	DUF620	190	438	698.7	4.50E-207
1314	CGPG6877.pep	Hep_59	101	194	198.3	1.90E-56
1315	CGPG6878.pep	Peptidase_M24	107	343	229.5	7.80E-66
1316	CGPG6880.pep	PAP_fibrillin	72	232	178.2	2.20E-50
1318	CGPG6887.pep	AIG2	12	111	89.1	1.40E-23
1319	CGPG6895.pep	zf-CCHC	166	183	26	0.00014
1319	CGPG6895.pep	zf-CCHC	208	225	24.9	0.00026
1319	CGPG6895.pep	zf-CCHC	325	342	38.1	3.30E-08
1319	CGPG6895.pep	zf-CCHC	356	373	32	2.20E-06
1322	CGPG6912.pep	zf-DHHC	149	213	119.1	1.30E-32
1324	CGPG6942.pep	RNA_pol_Rpb5_N	18	108	174.3	3.20E-49
1324	CGPG6942.pep	RNA_pol_Rpb5_C	149	222	68.2	2.80E-17
1327	CGPG6969.pep	2OG-Fell_Oxy	223	323	152.3	1.30E-42
1329	CGPG6991.pep	Glyco_tran_28_C	11	170	171.7	1.90E-48
1332	CGPG7037.pep	Sad1_UNC	313	416	-3.4	0.0011
1333	CGPG7051.pep	PAP_fibrillin	84	236	218.7	1.40E-62
1336	CGPG7086.pep	SFT2	44	163	182.6	9.70E-52
1339	CGPG7136.pep	NOI	1	72	159.7	8.10E-45
1340	CGPG7147.pep	RAMP4	1	64	152.5	1.20E-42
1343	CGPG7222.pep	Branch	49	274	399.8	4.00E-117

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1345	CGPG7233.pep	eIF2A	214	407	415.3	8.90E-122
1346	CGPG7238.pep	UPF0185	5	80	214.1	3.30E-61
1347	CGPG7253.pep	ArfGap	15	117	58.4	2.40E-14
1348	CGPG7255.pep	Arf	5	177	365.3	1.00E-106
1348	CGPG7255.pep	SRPRB	15	200	-3.7	2.60E-07
1348	CGPG7255.pep	Miro	19	129	40.3	7.10E-09
1348	CGPG7255.pep	Ras	19	179	-43.7	4.50E-06
1349	CGPG7286.pep	ORC2	20	345	558.7	6.00E-165
1350	CGPG7292.pep	Methyltransf_11	38	135	44.1	5.00E-10
1350	CGPG7292.pep	Methyltransf_12	38	133	38.4	2.60E-08
1351	CGPG7301.pep	ELFV_dehydrog_N	57	187	294.9	1.50E-85
1351	CGPG7301.pep	ELFV_dehydrog	202	445	484.3	1.50E-142
1352	CGPG7305.pep	DAO	126	401	-31.6	0.0013
1352	CGPG7305.pep	Pyr_redox_2	126	386	66.3	1.00E-16
1352	CGPG7305.pep	Pyr_redox	255	345	55.7	1.60E-13
1353	CGPG7307.pep	Gp_dh_N	11	163	300.6	3.00E-87
1353	CGPG7307.pep	Gp_dh_C	168	324	258.5	1.50E-74
1354	CGPG7318.pep	MIP	1	142	-31.2	9.10E-08
1355	CGPG7319.pep	SNF5	172	400	471.8	9.00E-139
1356	CGPG7345.pep	Lipase_3	97	237	131.1	3.30E-36
1357	CGPG7368.pep	MT-A70	516	676	316.4	5.20E-92
1358	CGPG7385.pep	FHA	32	107	51	4.10E-12
1359	CGPG7391.pep	G-patch	6	50	71.2	3.40E-18
1360	CGPG7405.pep	TATA_RF	1	216	160.8	3.70E-45
1361	CGPG7415.pep	IU_nuc_hydro	1	312	255.3	1.30E-73
1362	CGPG7416.pep	GLTP	52	233	208	2.30E-59
1363	CGPG7420.pep	PRK	95	302	426.6	3.60E-125
1366	CGPG7445.pep	PRC	92	169	37.2	5.70E-08
1366	CGPG7445.pep	PRC	171	252	48.6	2.20E-11
1367	CGPG7465.pep	Aldedh	103	567	834.4	6.30E-248
1368	CGPG7473.pep	F-box	273	320	35.1	2.60E-07
1368	CGPG7473.pep	WD40	412	449	41.5	2.90E-09
1368	CGPG7473.pep	WD40	453	493	49.7	1.00E-11
1368	CGPG7473.pep	WD40	520	556	30.6	5.70E-06
1368	CGPG7473.pep	WD40	560	598	41.7	2.70E-09
1369	CGPG7489.pep	UBX	295	376	70.8	4.50E-18
1370	CGPG7492.pep	Pescadillo_N	8	291	519	5.30E-153
1370	CGPG7492.pep	BRCT	337	414	50.7	5.00E-12
1371	CGPG7499.pep	G-alpha	423	839	50.2	3.20E-22
1374	CGPG7508.pep	Smg4_UPF3	1	170	236.2	7.10E-68
1375	CGPG7509.pep	SSB	76	187	95.3	1.90E-25
1376	CGPG7511.pep	GFA	31	125	74.2	4.20E-19
1377	CGPG7515.pep	Brix	45	297	232.5	9.60E-67
1378	CGPG7521.pep	G-alpha	23	386	560.5	1.80E-165
1380	CGPG7547.pep	PAD_porph	14	368	695.7	3.40E-206
1381	CGPG7561.pep	Snf7	17	187	219.6	7.40E-63

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1382	CGPG7562.pep	Sybindin	6	136	253.6	4.30E-73
1382	CGPG7562.pep	Sedlin_N	11	135	-21.6	0.00015
1383	CGPG7563.pep	DUF887	40	279	394.2	2.10E-115
1384	CGPG7567.pep	Peptidase_M18	14	461	680	1.90E-201
1389	CGPG7597.pep	zf-AN1	76	116	78.4	2.40E-20
1390	CGPG7606.pep	Pentapeptide	127	166	29.2	1.50E-05
1390	CGPG7606.pep	Pentapeptide	172	211	42.7	1.30E-09
1392	CGPG7637.pep	DAD	2	112	168.4	1.90E-47
1393	CGPG7649.pep	YgbB	71	227	244.8	2.00E-70
1394	CGPG7658.pep	Cyclase	56	256	109.9	7.70E-30
1395	CGPG7664.pep	Aldedh	9	463	801	7.20E-238
1396	CGPG7666.pep	Aminotran_1_2	26	392	491	1.50E-144
1397	CGPG7668.pep	DUF594	660	719	97.6	3.90E-26
1399	CGPG7746.pep	UPF0153	49	139	59.7	9.90E-15
1400	CGPG7747.pep	Copine	113	261	308.5	1.30E-89
1401	CGPG7752.pep	zf-MYND	176	214	48.9	1.80E-11
1401	CGPG7752.pep	PDCD2_C	241	403	38.2	1.00E-10
1403	CGPG7770.pep	YTH	263	353	218.6	1.40E-62
1406	CGPG7778.pep	Ubie_methyltran	51	304	393.3	3.70E-115
1406	CGPG7778.pep	Methyltransf_11	105	222	75	2.50E-19
1406	CGPG7778.pep	Methyltransf_12	105	220	43.1	1.00E-09
1407	CGPG7786.pep	KH_1	141	193	27.6	4.60E-05
1408	CGPG7788.pep	YIF1	34	263	341.8	1.20E-99
1410	CGPG78.pep	Hpt	46	132	55.5	1.80E-13
1411	CGPG783.pep	RRM_1	288	354	40	8.40E-09
1411	CGPG783.pep	RRM_2	682	778	232.4	1.00E-66
1412	CGPG7832.pep	STT3	26	670	626.4	2.50E-185
1413	CGPG7841.pep	ECH	17	186	192.8	8.40E-55
1413	CGPG7841.pep	3HCDH_N	311	490	295.4	1.10E-85
1413	CGPG7841.pep	3HCDH	492	585	116.2	9.50E-32
1414	CGPG7845.pep	LRRNT_2	23	64	49.2	1.50E-11
1414	CGPG7845.pep	LRR_1	97	119	10.5	3.5
1414	CGPG7845.pep	LRR_1	121	143	14.5	0.39
1414	CGPG7845.pep	LRR_1	145	167	12.1	1.7
1414	CGPG7845.pep	LRR_1	169	192	13.1	1.1
1414	CGPG7845.pep	LRR_1	194	216	14	0.55
1415	CGPG7847.pep	Peptidase_C48	30	225	-1.9	0.00023
1417	CGPG7853.pep	LRR_1	142	164	17.4	0.052
1417	CGPG7853.pep	LRR_1	166	188	9.9	4.4
1417	CGPG7853.pep	LRR_1	190	212	10	4.3
1417	CGPG7853.pep	LRR_1	214	235	13.8	0.65
1417	CGPG7853.pep	LRR_1	258	280	10.5	3.5
1417	CGPG7853.pep	LRR_1	282	304	14	0.56
1417	CGPG7853.pep	LRR_1	306	325	11.7	2.1

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1418	CGPG7857.pep	Response_reg	79	195	89.7	9.40E-24
1418	CGPG7857.pep	CCT	689	736	62.1	1.80E-15
1420	CGPG7869.pep	Acyltransferase	117	264	81.9	2.00E-21
1421	CGPG7891.pep	FMO-like	11	427	-207	1.20E-16
1421	CGPG7891.pep	DAO	13	268	-9.4	4.70E-05
1422	CGPG7892.pep	FMO-like	9	416	-157.4	2.80E-19
1423	CGPG7906.pep	Skp1_POZ	4	65	96.7	7.50E-26
1423	CGPG7906.pep	Skp1	76	153	154.3	3.40E-43
1424	CGPG7924.pep	AAA	249	462	54.2	4.60E-13
1428	CGPG7964.pep	IQ	108	128	25.7	0.00017
1428	CGPG7964.pep	IQ	130	150	12.2	1.9
1429	CGPG7968.pep	FMO-like	13	427	-229.9	2.10E-15
1429	CGPG7968.pep	DAO	15	291	-34.2	0.0019
1429	CGPG7968.pep	Pyr_redox_2	15	321	-18	0.00054
1431	CGPG7972.pep	FAD_binding_3	5	372	-97.4	5.20E-06
1432	CGPG7982.pep	Tetraspannin	4	241	241.9	1.40E-69
1433	CGPG7985.pep	BT1	57	469	419.2	6.00E-123
1434	CGPG7993.pep	F-box	27	73	15.1	0.26
1434	CGPG7993.pep	LRR_2	278	304	16.8	0.084
1435	CGPG8.pep	p450	39	503	399	7.30E-117
1436	CGPG80.pep	Pkinase	68	328	277.6	2.60E-80
1437	CGPG8001.pep	FMO-like	13	433	-243.5	1.10E-14
1437	CGPG8001.pep	Pyr_redox	215	295	5.3	0.0016
1438	CGPG8009.pep	BT1	41	511	517.8	1.30E-152
1440	CGPG8049.pep	BT1	1	436	331.8	1.20E-96
1441	CGPG806.pep	Synaptobrevin	127	215	146.7	6.20E-41
1442	CGPG8060.pep	DUF829	161	425	426	5.30E-125
1446	CGPG81.pep	Pkinase	70	330	287.7	2.30E-83
1452	CGPG8125.pep	zf-C3HC4	46	86	31.1	4.00E-06
1453	CGPG8134.pep	DEK_C	3	57	60.4	6.10E-15
1453	CGPG8134.pep	SWIB	197	272	140.9	3.60E-39
1453	CGPG8134.pep	SWIB	305	382	112.9	9.40E-31
1455	CGPG8156.pep	PCI	258	362	69.4	1.20E-17
1456	CGPG8159.pep	ACT	77	141	51.9	2.10E-12
1456	CGPG8159.pep	ACT	308	374	51.5	3.00E-12
1458	CGPG8179.pep	WD40	138	177	23.7	0.00069
1458	CGPG8179.pep	WD40	264	305	42.5	1.50E-09
1459	CGPG8193.pep	Ank	22	54	5.5	5.2
1459	CGPG8193.pep	Ank	55	87	36.4	1.00E-07
1460	CGPG8197.pep	F-box	5	53	22	0.0022
1461	CGPG8198.pep	F-box	74	119	15	0.28
1461	CGPG8198.pep	Kelch_2	174	221	19.7	0.011
1461	CGPG8198.pep	Kelch_1	223	278	15.6	0.041
1461	CGPG8198.pep	Kelch_2	329	369	14.2	0.51
1463	CGPG8211.pep	F-box	182	229	48.8	2.00E-11
1463	CGPG8211.pep	WD40	292	328	30.3	6.90E-06
1463	CGPG8211.pep	WD40	332	368	37.7	4.30E-08
1463	CGPG8211.pep	WD40	372	408	43.4	8.30E-10

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1463	CGPG8211.pep	WD40	411	449	40.9	4.60E-09
1463	CGPG8211.pep	WD40	453	538	32.2	1.90E-06
1463	CGPG8211.pep	WD40	542	578	38.9	1.80E-08
1464	CGPG823.pep	Ribosomal_S6e	1	129	258.3	1.70E-74
1466	CGPG8238.pep	PGI	49	537	857.4	7.40E-255
1467	CGPG8260.pep	Iso_dh	4	331	460.7	1.90E-135
1468	CGPG8267.pep	Pyridoxal_deC	34	383	352.2	8.70E-103
1469	CGPG8268.pep	NIR_SIR_ferr	62	131	43.9	5.60E-10
1469	CGPG8268.pep	NIR_SIR	164	345	193.4	5.50E-55
1469	CGPG8268.pep	NIR_SIR_ferr	360	432	76.9	6.80E-20
1470	CGPG8274.pep	PK	5	347	589.9	2.50E-174
1470	CGPG8274.pep	PK_C	363	476	81.9	2.10E-21
1471	CGPG8277.pep	Rib_5-P_isom_A	55	229	345	1.30E-100
1472	CGPG8279.pep	F-box	22	69	43	1.10E-09
1472	CGPG8279.pep	Kelch_1	124	168	35.2	2.40E-07
1472	CGPG8279.pep	Kelch_1	170	213	29.4	1.40E-05
1472	CGPG8279.pep	Kelch_2	170	213	24.3	0.00045
1475	CGPG8408.pep	DUF641	60	192	237.3	3.40E-68
1476	CGPG842.pep	Redoxin	5	162	182.6	9.80E-52
1476	CGPG842.pep	AhpC-TSA	6	142	38.6	2.20E-08
1477	CGPG8421.pep	NOSIC	121	173	90.1	6.90E-24
1477	CGPG8421.pep	Nop	213	361	257.3	3.20E-74
1478	CGPG8435.pep	NHL	55	82	5.9	3.4
1478	CGPG8435.pep	NHL	115	142	37.2	5.80E-08
1479	CGPG8440.pep	U-box	242	316	70.2	6.80E-18
1479	CGPG8440.pep	Arm	372	413	47.5	4.70E-11
1479	CGPG8440.pep	Arm	455	495	32.1	2.00E-06
1479	CGPG8440.pep	Arm	496	537	22.3	0.0018
1479	CGPG8440.pep	Arm	538	578	25.5	0.0002
1480	CGPG8453.pep	ACT	36	99	33.5	7.90E-07
1480	CGPG8453.pep	ACT	129	201	54.1	4.90E-13
1480	CGPG8453.pep	ACT	265	332	28.6	2.40E-05
1480	CGPG8453.pep	ACT	343	406	43.5	7.50E-10
1481	CGPG8470.pep	DUF89	18	357	-103.6	0.0022
1482	CGPG8479.pep	F-box	31	78	22.6	0.0015
1482	CGPG8479.pep	Kelch_1	124	170	35.4	2.00E-07
1482	CGPG8479.pep	Kelch_1	172	216	44.6	3.40E-10
1482	CGPG8479.pep	Kelch_2	172	216	21.7	0.0027
1483	CGPG8489.pep	F-box	3	50	44.4	3.90E-10
1483	CGPG8489.pep	FBA_1	219	384	291.3	2.00E-84
1484	CGPG8498.pep	Fcf1	87	185	245.9	9.10E-71
1485	CGPG85.pep	Aa_trans	19	454	556.5	2.80E-164
1486	CGPG8501.pep	PPR	149	183	11.6	0.32
1486	CGPG8501.pep	PPR	189	223	29	1.70E-05
1486	CGPG8501.pep	PPR	224	258	25.6	0.00019
1486	CGPG8501.pep	PPR	259	292	6.5	1.3
1486	CGPG8501.pep	PPR	330	364	4.2	2.5

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1486	CGPG8501.pep	PPR	401	435	12.3	0.27
1487	CGPG8507.pep	DUF588	31	185	202.1	1.30E-57
1488	CGPG8509.pep	Pro_isomerase	80	233	18.3	3.90E-10
1490	CGPG8517.pep	B_lectin	84	197	191	2.90E-54
1490	CGPG8517.pep	S_locus_glycop	211	339	250	5.20E-72
1490	CGPG8517.pep	PAN_2	356	422	105.1	2.10E-28
1492	CGPG852.pep	Ribosomal_S5	94	160	140.9	3.60E-39
1492	CGPG852.pep	Ribosomal_S5_C	177	250	134.4	3.30E-37
1498	CGPG8567.pep	TB2_DP1_HVA22	3	98	59.4	1.20E-14
1499	CGPG8580.pep	Caleosin	22	195	434.3	1.70E-127
1501	CGPG8590.pep	DUF579	42	289	571.9	6.60E-169
1503	CGPG8594.pep	DUF584	1	139	248.7	1.30E-71
1504	CGPG8606.pep	Exo_endo_phos	56	327	125	2.30E-34
1505	CGPG8628.pep	DUF584	32	203	125.8	1.30E-34
1507	CGPG8654.pep	PPR	73	107	13.2	0.21
1507	CGPG8654.pep	PPR	141	175	38.6	2.30E-08
1508	CGPG8668.pep	TATA_RF	1	200	306.9	3.90E-89
1511	CGPG8702.pep	RPE65	50	622	19.9	1.70E-17
1514	CGPG8748.pep	AAA	201	357	13.1	1.30E-06
1516	CGPG8770.pep	DUF543	1	73	107.6	3.90E-29
1517	CGPG8777.pep	DUF569	1	144	377.9	1.70E-110
1517	CGPG8777.pep	DUF569	227	368	391.4	1.40E-114
1519	CGPG8786.pep	DUF260	13	113	247.6	2.70E-71
1520	CGPG8792.pep	Abhydrolase_3	75	290	228.4	1.60E-65
1521	CGPG8801.pep	PPR	110	144	18.7	0.022
1521	CGPG8801.pep	PPR	145	179	9.1	0.64
1521	CGPG8801.pep	PPR	181	215	42.6	1.40E-09
1521	CGPG8801.pep	PPR	216	250	19.5	0.012
1521	CGPG8801.pep	PPR	251	285	23.6	0.00076
1521	CGPG8801.pep	PPR	286	320	37.2	6.10E-08
1521	CGPG8801.pep	PPR	321	355	12	0.3
1522	CGPG8809.pep	Ribosomal_L12	141	208	61.2	3.50E-15
1523	CGPG8840.pep	CLP_protease	115	289	45.1	1.30E-11
1524	CGPG8850.pep	PPR	189	223	17.3	0.057
1524	CGPG8850.pep	PPR	224	258	48.5	2.40E-11
1524	CGPG8850.pep	PPR	297	331	34.6	3.50E-07
1524	CGPG8850.pep	PPR	332	366	46.2	1.20E-10
1524	CGPG8850.pep	PPR	367	401	40.7	5.40E-09
1524	CGPG8850.pep	PPR	402	436	50	8.00E-12
1524	CGPG8850.pep	PPR	437	470	36.5	9.50E-08
1524	CGPG8850.pep	PPR	471	505	39.8	9.60E-09
1524	CGPG8850.pep	PPR	506	540	23.2	0.00094

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1524	CGPG8850.pep	PPR	542	576	46.6	8.80E-11
1525	CGPG8870.pep	Fructosamin_kin	54	339	208	2.30E-59
1525	CGPG8870.pep	APH	69	308	80.1	7.30E-21
1526	CGPG8871.pep	DUF59	36	115	57.6	4.20E-14
1527	CGPG8872.pep	RRM_1	138	205	78	3.00E-20
1527	CGPG8872.pep	RRM_1	214	288	32.4	1.70E-06
1527	CGPG8872.pep	RRM_1	308	378	60	8.10E-15
1528	CGPG8876.pep	SFT2	43	162	194.7	2.40E-55
1529	CGPG8902.pep	RNA_pol_I_A49	31	415	702.7	2.70E-208
1530	CGPG8904.pep	G-alpha	103	447	513.2	3.10E-151
1531	CGPG8907.pep	DnaJ	4	67	128.7	1.70E-35
1531	CGPG8907.pep	zf-C2H2	338	362	28	3.40E-05
1532	CGPG8914.pep	Whi5	181	205	50.1	7.80E-12
1533	CGPG8931.pep	eIF3_subunit	1	231	272.1	1.20E-78
1534	CGPG8933.pep	MACPF	126	322	60.6	5.20E-15
1535	CGPG8935.pep	Pkinase	33	324	308.5	1.30E-89
1536	CGPG8938.pep	Pkinase	71	348	187.5	3.40E-53
1536	CGPG8938.pep	Pkinase_Tyr	71	348	128.4	2.10E-35
1537	CGPG8944.pep	PH	31	134	48.5	2.40E-11
1538	CGPG895.pep	DUF231	251	427	223.7	4.30E-64
1540	CGPG8961.pep	Dus	59	392	236.5	5.90E-68
1541	CGPG8963.pep	Pkinase	319	612	332	1.10E-96
1542	CGPG898.pep	DUF231	276	448	289.8	5.30E-84
1543	CGPG8993.pep	AAA	44	226	54.9	2.80E-13
1543	CGPG8993.pep	Rep_fac_C	237	326	113.5	6.50E-31
1544	CGPG8994.pep	Brix	86	262	264.3	2.50E-76
1545	CGPG8995.pep	Porin_3	54	326	280.1	4.30E-81
1546	CGPG90.pep	MFS_1	68	427	55	2.50E-13
1547	CGPG900.pep	DUF231	241	420	229.6	6.90E-66
1548	CGPG9002.pep	CTP_transf_1	51	382	443.1	3.90E-130
1549	CGPG9009.pep	Alg6_Alg8	22	514	439.6	4.30E-129
1550	CGPG9011.pep	OTCace_N	70	212	188.6	1.60E-53
1550	CGPG9011.pep	OTCace	216	369	198.1	2.20E-56
1551	CGPG9012.pep	FolB	16	129	125	2.10E-34
1552	CGPG9017.pep	DMRL_synthase	70	213	195.5	1.30E-55
1553	CGPG9025.pep	Glyco_transf_29	105	355	-11	1.90E-06
1554	CGPG9026.pep	Adap_comp_sub	160	428	469.4	4.70E-138
1555	CGPG9032.pep	RNase_PH	33	165	135.6	1.40E-37
1555	CGPG9032.pep	RNase_PH_C	194	262	44.9	2.70E-10
1556	CGPG9040.pep	Sec1	35	592	217.7	2.70E-62
1557	CGPG9044.pep	Syntaxin	57	163	93.9	5.10E-25
1557	CGPG9044.pep	SNARE	256	318	79.5	1.10E-20
1558	CGPG9048.pep	RNase_PH	15	135	53.3	8.30E-13
1559	CGPG9049.pep	Glycos_transf_1	244	442	16	3.70E-05

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1560	CGPG9058.pep	RNA_pol_N	1	60	125.6	1.40E-34
1561	CGPG906.pep	PPR	210	244	28.2	3.10E-05
1561	CGPG906.pep	PPR	245	279	13.7	0.19
1562	CGPG9070.pep	ATP-synt_C	20	85	42.4	1.60E-09
1562	CGPG9070.pep	ATP-synt_C	104	169	21.7	0.00014
1564	CGPG9084.pep	CH	15	116	47.4	5.20E-11
1564	CGPG9084.pep	EB1	209	255	79.7	9.20E-21
1566	CGPG9098.pep	DUF6	32	165	22.6	0.0015
1566	CGPG9098.pep	TPT	199	337	-10.3	0.0023
1566	CGPG9098.pep	DUF6	208	337	66.1	1.20E-16
1567	CGPG9099.pep	Rick_17kDa_Anti	66	110	24.9	0.00029
1568	CGPG9110.pep	RALF	53	118	122.9	9.30E-34
1569	CGPG9119.pep	PTR2	115	504	350.4	3.10E-102
1570	CGPG9125.pep	ThiF	98	233	171.7	1.90E-48
1571	CGPG913.pep	PPR	86	120	32.9	1.20E-06
1571	CGPG913.pep	PPR	188	222	51.1	4.00E-12
1571	CGPG913.pep	PPR	223	257	7.4	1
1571	CGPG913.pep	PPR	289	323	26	0.00014
1571	CGPG913.pep	PPR	325	358	12	0.29
1571	CGPG913.pep	PPR	361	395	12.8	0.24
1574	CGPG9164.pep	DHBP_synthase	5	202	475.9	5.10E-140
1574	CGPG9164.pep	GTP_cyclohydro2	207	377	431.4	1.30E-126
1575	CGPG9172.pep	Sec61_beta	32	77	83	9.60E-22
1577	CGPG9185.pep	DUF423	10	115	84.7	2.90E-22
1578	CGPG9187.pep	Pkinase	12	263	298.2	1.60E-86
1579	CGPG9190.pep	DREPP	2	203	280.3	3.80E-81
1580	CGPG9193.pep	GATase	11	196	175.1	1.90E-49
1580	CGPG9193.pep	GMP_synt_C	432	524	172.7	9.30E-49
1581	CGPG9195.pep	DUF1279	91	193	127	5.60E-35
1582	CGPG9203.pep	Na_H_Exchanger	13	391	242.3	1.00E-69
1582	CGPG9203.pep	TrkA_C	418	486	58.4	2.50E-14
1583	CGPG9209.pep	GATase	9	197	199.6	7.80E-57
1583	CGPG9209.pep	GMP_synt_C	423	515	201.4	2.20E-57
1585	CGPG9210.pep	Na_H_Exchanger	10	376	229	1.10E-65
1585	CGPG9210.pep	TrkA_N	402	517	131	3.50E-36
1586	CGPG9211.pep	NTP_transferase	7	296	508.1	1.00E-149
1586	CGPG9211.pep	MannoseP_isomer	307	473	454	2.00E-133
1586	CGPG9211.pep	Cupin_2	388	458	48.1	3.10E-11
1587	CGPG9212.pep	ATP-synt_ab_N	22	87	53.4	7.60E-13
1587	CGPG9212.pep	ATP-synt_ab	143	353	429.3	5.40E-126
1588	CGPG9216.pep	DHBP_synthase	10	206	377.8	1.70E-110

PEP Seq ID No.	Construct ID	Pfam domain name	Begin	Stop	score	E-value
1588	CGPG9216.pep	GTP_cyclohydro2	211	379	395.3	9.20E-116
1589	CGPG9226.pep	Carboxyl_trans	40	541	967.5	5.20E-288
1590	CGPG9252.pep	CDC48_N	28	114	132.1	1.60E-36
1590	CGPG9252.pep	AAA	245	429	330.3	3.40E-96
1590	CGPG9252.pep	AAA_5	245	377	9.8	0.00052
1590	CGPG9252.pep	AAA	518	705	344.1	2.40E-100
1592	CGPG9289.pep	LSM	10	75	64.8	3.00E-16
1593	CGPG9298.pep	p450	41	516	138.8	1.60E-38
1594	CGPG9299.pep	ThiF	82	227	145.2	1.80E-40
1597	CGPG9316.pep	ATP-grasp_2	34	242	405.7	7.10E-119
1597	CGPG9316.pep	Ligase_CoA	285	421	229	1.10E-65
1598	CGPG9317.pep	DUF163	43	196	205.3	1.50E-58
1599	CGPG9318.pep	Aldo_ket_red	20	285	408	1.40E-119
1600	CGPG9324.pep	Pkinase	158	706	150.9	3.40E-42
1601	CGPG9357.pep	Raffinose_syn	7	757	1669.7	0
1602	CGPG965.pep	Hexapep	70	87	10.5	6.4
1602	CGPG965.pep	Hexapep	119	136	16.4	0.11
1602	CGPG965.pep	Hexapep	142	159	5.9	28
1602	CGPG965.pep	Hexapep	160	177	2.6	71
1603	CGPG970.pep	NPH3	206	441	272	1.20E-78
1604	CGPG982.pep	p450	29	495	389.8	4.30E-114
1605	CGPG994.pep	adh_short	50	218	46.7	8.30E-11
1606	CGPG996.pep	DUF866	1	167	456.5	3.60E-134

Table 22.

Pfam domain name	Accession #	gathering cutoff	domain description
2OG-FeII_Oxy	PF03171.11	11.5	2OG-Fe(II) oxygenase superfamily
3Beta_HSD	PF01073.10	-135.9	3-beta hydroxysteroid dehydrogenase/isomerase family
3HCDH	PF00725.13	-8.9	3-hydroxyacyl-CoA dehydrogenase, C-terminal domain
3HCDH_N	PF02737.9	-82.1	3-hydroxyacyl-CoA dehydrogenase, NAD binding domain
6PGD	PF00393.10	-232.3	6-phosphogluconate dehydrogenase, C-terminal domain
AAA	PF00004.20	12.3	ATPase family associated with various cellular activities (AAA)
AAA_5	PF07728.5	4	ATPase family associated with various cellular activities (AAA)
AARP2CN	PF08142.3	25	AARP2CN (NUC121) domain
AA_kinase	PF00696.19	-40	Amino acid kinase family
AA_permease	PF00324.12	-120.8	Amino acid permease
ABC_tran	PF00005.18	9.5	ABC transporter
ACBP	PF00887.10	-14	Acyl CoA binding protein
ACP_syn_III_C	PF08541.1	-21.6	3-Oxoacyl-[acyl-carrier-protein (ACP)] synthase III C terminal
ACT	PF01842.16	0.7	ACT domain
ADH_N	PF08240.3	-14.5	Alcohol dehydrogenase GroES-like domain
ADH_zinc_N	PF00107.17	23.8	Zinc-binding dehydrogenase
AIG1	PF04548.7	-40	AIG1 family
AIG2	PF06094.3	12.7	AIG2-like family
AP2	PF00847.11	21.7	AP2 domain
APH	PF01636.14	32.5	Phosphotransferase enzyme family

Pfam domain name	Accession #	gathering cutoff	domain description
ATP-grasp_2	PF08442.1	-118.8	ATP-grasp domain
ATP-synt_C	PF00137.12	5	ATP synthase subunit C
ATP-synt_G	PF04718.6	-7.9	Mitochondrial ATP synthase g subunit
ATP-synt_ab	PF00006.16	-37.5	ATP synthase alpha/beta family, nucleotide-binding domain
ATP-synt_ab_N	PF02874.14	17	ATP synthase alpha/beta family, beta-barrel domain
ATP_bind_1	PF03029.8	-19	Conserved hypothetical ATP binding protein
AWPM-19	PF05512.2	25	AWPM-19-like family
Aa_trans	PF01490.9	-128.4	Transmembrane amino acid transporter protein
Abhydrolase_3	PF07859.4	25.8	alpha/beta hydrolase fold
Acetyltransf_1	PF00583.15	18.6	Acetyltransferase (GNAT) family
Acyltransferase	PF01553.12	2.2	Acyltransferase
Adap_comp_sub	PF00928.12	-5.4	Adaptor complexes medium subunit family
AhpC-TSA	PF00578.12	4.8	AhpC/TSA family
Alba	PF01918.12	20	Alba
Aldedh	PF00171.13	-208.2	Aldehyde dehydrogenase family
Aldo_ket_red	PF00248.12	-97	Aldo/keto reductase family
Alg6_Alg8	PF03155.6	-235.5	ALG6, ALG8 glycosyltransferase family
Alpha-amyl_C2	PF07821.3	25	Alpha-amylase C-terminal beta-sheet domain
Alpha-amylase	PF00128.15	-92.6	Alpha amylase, catalytic domain
Aminotran_1_2	PF00155.12	-57.5	Aminotransferase class I and II
Aminotran_3	PF00202.12	-206.1	Aminotransferase class-III
Aminotran_5	PF00266.10	-164.4	Aminotransferase class-V
Ank	PF00023.21	0	Ankyrin repeat
Arf	PF00025.12	40	ADP-ribosylation factor family
ArfGap	PF01412.9	-17	Putative GTPase activating protein for Arf
Arm	PF00514.14	17	Armadillo/beta-catenin-like repeat
Asp	PF00026.14	-153.8	Eukaryotic aspartyl protease
Asparaginase	PF00710.11	25	Asparaginase
Auxin_inducible	PF02519.5	-15	Auxin responsive protein
B3	PF02362.12	26.5	B3 DNA binding domain
BRCT	PF00533.17	27.8	BRCA1 C Terminus (BRCT) domain
BT1	PF03092.7	-148.6	BT1 family
B_lectin	PF01453.15	28.2	D-mannose binding lectin
Band_7	PF01145.16	10.3	SPFH domain / Band 7 family
Beta_elim_lyase	PF01212.12	-110.3	Beta-eliminating lyase
Biotin_lipoyl	PF00364.13	-2.3	Biotin-requiring enzyme
Bombesin	PF02044.8	11.4	Bombesin-like peptide
Branch	PF02485.12	-83.5	Core-2/I-Branching enzyme
Brix	PF04427.9	11.4	Brix domain
Bromodomain	PF00439.16	8.9	Bromodomain
C1_1	PF00130.13	11.4	Phorbol esters/diacylglycerol binding domain (C1 domain)
C1_2	PF03107.7	26.4	C1 domain
C1_3	PF07649.3	25	C1-like domain
C2	PF00168.21	3.7	C2 domain
CAF1	PF04857.11	-100.5	CAF1 family ribonuclease
CBFD_NFYB_HMF	PF00808.14	18.4	Histone-like transcription factor (CBF/NF-Y) and archaeal histone
CBS	PF00571.19	17.5	CBS domain pair
CCT	PF06203.5	25	CCT motif
CDC48_N	PF02359.9	-2	Cell division protein 48 (CDC48), N-terminal domain

Pfam domain name	Accession #	gathering cutoff	domain description
CDC50	PF03381.6	25	LEM3 (ligand-effect modulator 3) family / CDC50 family
CDP-OH_P_transf	PF01066.12	0	CDP-alcohol phosphatidyltransferase
CH	PF00307.22	22.5	Calponin homology (CH) domain
CLP_protease	PF00574.14	-77.2	Clp protease
CPSase_sm_chain	PF00988.13	25	Carbamoyl-phosphate synthase small chain, CPSase domain
CRAL_TRIO	PF00650.11	-26	CRAL/TRIO domain
CRAL_TRIO_N	PF03765.6	16	CRAL/TRIO, N-terminus
CTP_transf_1	PF01148.11	-35.9	Cytidyltransferase family
Caleosin	PF05042.4	25	Caleosin related protein
Carb_anhydrase	PF00194.12	-105	Eukaryotic-type carbonic anhydrase
Carboxyl_trans	PF01039.13	-262.3	Carboxyl transferase domain
Cellulose_synt	PF03552.5	-257.9	Cellulose synthase
Cenp-O	PF09496.1	25	Cenp-O kinetochore centromere component
Chal_sti_synt_C	PF02797.6	-6.1	Chalcone and stilbene synthases, C-terminal domain
Chalcone	PF02431.6	25	Chalcone-flavanone isomerase
ClpS	PF02617.8	-15	ATP-dependent Clp protease adaptor protein ClpS
Copine	PF07002.7	-36.5	Copine
Cu-oxidase	PF00394.13	-18.9	Multicopper oxidase
Cu-oxidase_2	PF07731.5	-5.8	Multicopper oxidase
Cu-oxidase_3	PF07732.6	10	Multicopper oxidase
Cullin	PF00888.13	-33.3	Cullin family
Cupin_1	PF00190.13	-13	Cupin
Cupin_2	PF07883.2	19.4	Cupin domain
Cyclase	PF04199.4	-31.7	Putative cyclase
Cyclin_C	PF02984.10	-13	Cyclin, C-terminal domain
Cyclin_N	PF00134.14	-14.7	Cyclin, N-terminal domain
DAD	PF02109.7	25	DAD family
DAO	PF01266.15	-34.9	FAD dependent oxidoreductase
DCP1	PF06058.4	-16.8	Dcp1-like decapping family
DEAD	PF00270.20	7.2	DEAD/DEAH box helicase
DEK_C	PF08766.2	25	DEK C terminal domain
DHBP_synthase	PF00926.10	-95.1	3,4-dihydroxy-2-butanone 4-phosphate synthase
DMRL_synthase	PF00885.10	-43	6,7-dimethyl-8-ribityllumazine synthase
DREPP	PF05558.3	25	DREPP plasma membrane polypeptide
DSPc	PF00782.11	-21.8	Dual specificity phosphatase, catalytic domain
DUF1279	PF06916.4	25	Protein of unknown function (DUF1279)
DUF1350	PF07082.2	-145.7	Protein of unknown function (DUF1350)
DUF1475	PF07343.2	25	Protein of unknown function (DUF1475)
DUF1517	PF07466.2	25	Protein of unknown function (DUF1517)
DUF163	PF02590.8	-48.6	Uncharacterized ACR, COG1576
DUF1637	PF07847.3	25	Protein of unknown function (DUF1637)
DUF167	PF02594.7	-3.4	Uncharacterised ACR, YggU family COG1872
DUF177	PF02620.8	25	Uncharacterized ACR, COG1399
DUF220	PF02713.5	25	Domain of unknown function DUF220
DUF231	PF03005.6	-58	Arabidopsis proteins of unknown function
DUF260	PF03195.5	0.8	Protein of unknown function DUF260
DUF423	PF04241.6	-17.6	Protein of unknown function (DUF423)
DUF543	PF04418.3	25	Domain of unknown function (DUF543)
DUF569	PF04601.4	-47.6	Protein of unknown function (DUF569)

Pfam domain name	Accession #	gathering cutoff	domain description
DUF579	PF04669.4	25	Protein of unknown function (DUF579)
DUF584	PF04520.4	25	Protein of unknown function, DUF584
DUF588	PF04535.3	25	Domain of unknown function (DUF588)
DUF59	PF01883.10	-7.9	Domain of unknown function DUF59
DUF594	PF04578.4	25	Protein of unknown function, DUF594
DUF599	PF04654.3	25	Protein of unknown function, DUF599
DUF6	PF00892.11	21.7	Integral membrane protein DUF6
DUF607	PF04678.4	25	Protein of unknown function, DUF607
DUF616	PF04765.4	25	Protein of unknown function (DUF616)
DUF620	PF04788.3	25	Protein of unknown function (DUF620)
DUF641	PF04859.3	25	Plant protein of unknown function (DUF641)
DUF663	PF04950.3	25	Protein of unknown function (DUF663)
DUF822	PF05687.4	25	Plant protein of unknown function (DUF822)
DUF829	PF05705.5	25	Eukaryotic protein of unknown function (DUF829)
DUF862	PF05903.5	-21.4	PPPDE putative peptidase domain
DUF866	PF05907.4	30.1	Eukaryotic protein of unknown function (DUF866)
DUF887	PF05967.2	-72.3	Eukaryotic protein of unknown function (DUF887)
DUF89	PF01937.10	-113.8	Protein of unknown function DUF89
DUF933	PF06071.4	20	Protein of unknown function (DUF933)
DZC	PF08381.2	15.3	Disease resistance/zinc finger/chromosome condensation-like region
Di19	PF05605.3	25	Drought induced 19 protein (Di19)
Dimerisation	PF08100.2	25	Dimerisation domain
DnaJ	PF00226.22	-8	DnaJ domain
Dus	PF01207.8	-82.1	Dihydrouridine synthase (Dus)
E3_binding	PF02817.8	10	e3 binding domain
EB1	PF03271.8	25	EB1-like C-terminal motif
ECH	PF00378.11	-57.6	Enoyl-CoA hydratase/isomerase family
ELFV_dehydrog	PF00208.12	-26.4	Glutamate/Leucine/Phenylalanine/Valine dehydrogenase
ELFV_dehydrog_N	PF02812.9	31.8	Glu/Leu/Phe/Val dehydrogenase, dimerisation domain
EMP24_GP25L	PF01105.15	10.4	emp24/gp25L/p24 family/GOLD
ENOD93	PF03386.5	25	Early nodulin 93 ENOD93 protein
ENT	PF03735.5	25	ENT domain
Enolase_C	PF00113.13	-71.2	Enolase, C-terminal TIM barrel domain
Enolase_N	PF03952.7	11.3	Enolase, N-terminal domain
Epimerase	PF01370.12	-46.3	NAD dependent epimerase/dehydratase family
Exo_endo_phos	PF03372.14	11	Endonuclease/Exonuclease/phosphatase family
F-box	PF00646.24	14.4	F-box domain
F420_oxidored	PF03807.8	-34.5	NADP oxidoreductase coenzyme F420-dependent
FAD_binding_2	PF00890.15	-124.8	FAD binding domain
FAD_binding_3	PF01494.10	-136.6	FAD binding domain
FAE1_CUT1_RppA	PF08392.3	-192.7	FAE1/Type III polyketide synthase-like protein
FA_hydroxylase	PF04116.4	-15.3	Fatty acid hydroxylase superfamily
FBA_1	PF07734.4	-39.4	F-box associated
FBPase	PF00316.11	-170.3	Fructose-1-6-bisphosphatase
FHA	PF00498.17	25	FHA domain
FMO-like	PF00743.10	-381.6	Flavin-binding monooxygenase-like
FYVE	PF01363.12	28.5	FYVE zinc finger
Fcf1	PF04900.3	-7.6	Fcf1
Fer2	PF00111.18	7	2Fe-2S iron-sulfur cluster binding domain

Pfam domain name	Accession #	gathering cutoff	domain description
FolB	PF02152.9	25	Dihydroneopterin aldolase
Fructosamin_kin	PF03881.5	-125.6	Fructosamine kinase
G-alpha	PF00503.11	-230	G-protein alpha subunit
G-patch	PF01585.14	18.4	G-patch domain
G6PD_C	PF02781.7	-175	Glucose-6-phosphate dehydrogenase, C-terminal domain
G6PD_N	PF00479.13	-99	Glucose-6-phosphate dehydrogenase, NAD binding domain
GASA	PF02704.5	25	Gibberellin regulated protein
GAT	PF03127.5	-7	GAT domain
GATase	PF00117.19	-38.1	Glutamine amidotransferase class-I
GFA	PF04828.5	12.4	Glutathione-dependent formaldehyde-activating enzyme
GH3	PF03321.4	-336	GH3 auxin-responsive promoter
GHMP_kinases_C	PF08544.4	20.8	GHMP kinases C terminal
GHMP_kinases_N	PF00288.17	17.1	GHMP kinases N terminal domain
GIDA	PF01134.13	-226.7	Glucose inhibited division protein A
GLTP	PF08718.2	-71.1	Glycolipid transfer protein (GLTP)
GMP_synt_C	PF00958.13	25	GMP synthase C terminal domain
GST_C	PF00043.16	22.3	Glutathione S-transferase, C-terminal domain
GST_N	PF02798.11	14.6	Glutathione S-transferase, N-terminal domain
GTP_cyclohydro2	PF00925.11	-49	GTP cyclohydrolase II
G_glu_transpept	PF01019.12	-236.1	Gamma-glutamyltranspeptidase
Gln-synt_C	PF00120.15	-124	Glutamine synthetase, catalytic domain
GlutR_N	PF05201.6	-37.8	Glutamyl-tRNA ^{Glu} reductase, N-terminal domain
GlutR_dimer	PF00745.11	25	Glutamyl-tRNA ^{Glu} reductase, dimerisation domain
Glutaminase	PF04960.6	-143.6	Glutaminase
Glutaredoxin	PF00462.15	17.2	Glutaredoxin
Glyco_tran_28_C	PF04101.7	-10.4	Glycosyltransferase family 28 C-terminal domain
Glyco_transf_29	PF00777.9	-74	Glycosyltransferase family 29 (sialyltransferase)
Glyco_transf_8	PF01501.11	-43.2	Glycosyl transferase family 8
Glycolytic	PF00274.10	-174.5	Fructose-bisphosphate aldolase class-I
Glycos_transf_1	PF00534.11	-7.3	Glycosyl transferases group 1
Glyoxalase	PF00903.16	12.1	Glyoxalase/Bleomycin resistance protein/Dioxygenase superfamily
Gp_dh_C	PF02800.11	-64.1	Glyceraldehyde 3-phosphate dehydrogenase, C-terminal domain
Gp_dh_N	PF00044.15	-74.2	Glyceraldehyde 3-phosphate dehydrogenase, NAD binding domain
HI0933_like	PF03486.5	-255.8	HI0933-like protein
H_PPase	PF03030.7	-377	Inorganic H ⁺ pyrophosphatase
Helicase_C	PF00271.22	2.4	Helicase conserved C-terminal domain
Hep_59	PF07052.2	25	Hepatocellular carcinoma-associated antigen 59
Hexapep	PF00132.15	0.3	Bacterial transferase hexapeptide (three repeats)
Hin1	PF07320.4	25	Harpin-induced protein 1 (Hin1)
His_biosynth	PF00977.12	-102.4	Histidine biosynthesis protein
Histone	PF00125.15	17.4	Core histone H2A/H2B/H3/H4
Hpt	PF01627.14	25	Hpt domain
Hydrolase	PF00702.17	13.6	haloacid dehalogenase-like hydrolase
IF4E	PF01652.9	-35	Eukaryotic initiation factor 4E
IQ	PF00612.18	11.9	IQ calmodulin-binding motif
IU_nuc_hydro	PF01156.10	-154	Inosine-uridine preferring nucleoside hydrolase
Inositol_P	PF00459.16	-55.6	Inositol monophosphatase family
Invertase_neut	PF04853.3	-233.6	Plant neutral invertase
Ion_trans_2	PF07885.7	24.9	Ion channel

Pfam domain name	Accession #	gathering cutoff	domain description
Iso_dh	PF00180.11	-97	Isocitrate/isopropylmalate dehydrogenase
KH_1	PF00013.20	12.5	KH domain
KH_2	PF07650.8	5	KH domain
KR	PF08659.1	-74.3	KR domain
Kelch_1	PF01344.16	11.7	Kelch motif
Kelch_2	PF07646.6	14	Kelch motif
LAG1	PF03798.7	-57.5	Longevity-assurance protein (LAG1)
LANC_like	PF05147.4	-76.9	Lanthionine synthetase C-like protein
LEA_5	PF00477.8	25	Small hydrophilic plant seed protein
LRRNT_2	PF08263.3	18.6	Leucine rich repeat N-terminal domain
LRR_1	PF00560.24	7.7	Leucine Rich Repeat
LRR_2	PF07723.4	6	Leucine Rich Repeat
LSM	PF01423.13	13.7	LSM domain
La	PF05383.8	25	La domain
Lectin_legB	PF00139.10	-110.1	Legume lectin domain
Ligase_CoA	PF00549.10	-43.3	CoA-ligase
Lipase_3	PF01764.16	-8	Lipase (class 3)
MACPF	PF01823.10	-24.6	MAC/Perforin domain
MATH	PF00917.17	0.5	MATH domain
MFS_1	PF07690.7	23.5	Major Facilitator Superfamily
MIP	PF00230.11	-62	Major intrinsic protein
MMR_HSR1	PF01926.14	31.2	GTPase of unknown function
MOSC	PF03473.8	25	MOSC domain
MOSC_N	PF03476.7	25	MOSC N-terminal beta barrel domain
MT-A70	PF05063.5	-40.8	MT-A70
Malic_M	PF03949.6	-143.9	Malic enzyme, NAD binding domain
MannoseP_isomer	PF01050.9	-70	Mannose-6-phosphate isomerase
Mem_trans	PF03547.9	-73.4	Membrane transport protein
Metalloenzyme	PF01676.9	-14.4	Metalloenzyme superfamily
Metallophos	PF00149.19	22	Calcineurin-like phosphoesterase
Methyltransf_11	PF08241.3	21	Methyltransferase domain
Methyltransf_12	PF08242.3	23	Methyltransferase domain
Methyltransf_2	PF00891.9	-103.8	O-methyltransferase
Miro	PF08477.4	10.8	Miro-like protein
Mito_carr	PF00153.18	0	Mitochondrial carrier protein
Mov34	PF01398.12	0.9	Mov34/MPN/PAD-1 family
MtN3_slv	PF03083.7	9.7	MtN3/saliva family
NAC	PF01849.9	0	NAC domain
NAD_Gly3P_dh_C	PF07479.5	-50.8	NAD-dependent glycerol-3-phosphate dehydrogenase C-terminus
NAD_Gly3P_dh_N	PF01210.14	-44	NAD-dependent glycerol-3-phosphate dehydrogenase N-terminus
NAD_binding_2	PF03446.6	-63.5	NAD binding domain of 6-phosphogluconate dehydrogenase
NAD_binding_4	PF07993.3	-87.7	Male sterility protein
NAF	PF03822.5	4.5	NAF domain
NAP	PF00956.9	-122	Nucleosome assembly protein (NAP)
NDK	PF00334.10	-59.9	Nucleoside diphosphate kinase
NHL	PF01436.12	3.8	NHL repeat
NIR_SIR	PF01077.13	-19.6	Nitrite and sulphite reductase 4Fe-4S domain
NIR_SIR_ferr	PF03460.8	2.4	Nitrite/Sulfite reductase ferredoxin-like half domain
NOI	PF05627.2	0.3	Nitrate-induced NOI protein

Pfam domain name	Accession #	gathering cutoff	domain description
NOSIC	PF08060.4	25	NOSIC (NUC001) domain
NPH3	PF03000.5	25	NPH3 family
NTF2	PF02136.11	6	Nuclear transport factor 2 (NTF2) domain
NTP_transferase	PF00483.14	-39.9	Nucleotidyl transferase
Na_Ca_ex	PF01699.15	25	Sodium/calcium exchanger protein
Na_H_Exchanger	PF00999.12	-67.9	Sodium/hydrogen exchanger family
Na_sulph_symp	PF00939.10	-259	Sodium:sulfate symporter transmembrane region
Nfu_N	PF08712.2	20	Scaffold protein Nfu/NifU N terminal
NifU	PF01106.8	-5.6	NifU-like domain
NifU_N	PF01592.7	-27.4	NifU-like N terminal domain
Nodulin-like	PF06813.4	-57.8	Nodulin-like
Nop	PF01798.9	25	Putative snoRNA binding domain
Nramp	PF01566.9	-171.4	Natural resistance-associated macrophage protein
OPT	PF03169.6	-238.6	OPT oligopeptide transporter protein
ORC2	PF04084.5	-181.2	Origin recognition complex subunit 2
OTCace	PF00185.15	-42	Aspartate/ornithine carbamoyltransferase, Asp/Orn binding domain
OTCace_N	PF02729.12	-55	Aspartate/ornithine carbamoyltransferase, carbamoyl-P binding domain
PA	PF02225.13	13	PA domain
PAD_porph	PF04371.6	-180.8	Porphyromonas-type peptidyl-arginine deiminase
PALP	PF00291.16	-70	Pyridoxal-phosphate dependent enzyme
PAN_1	PF00024.17	1.4	PAN domain
PAN_2	PF08276.2	-4.9	PAN-like domain
PAP_fibrillin	PF04755.3	25	PAP_fibrillin
PB1	PF00564.15	12.3	PB1 domain
PCI	PF01399.18	25	PCI domain
PDCD2_C	PF04194.4	-40.9	Programmed cell death protein 2, C-terminal putative domain
PDT	PF00800.9	25	Prephenate dehydratase
PEP-utilizers	PF00391.14	0.6	PEP-utilising enzyme, mobile domain
PEP-utilizers_C	PF02896.9	-173	PEP-utilising enzyme, TIM barrel domain
PGAM	PF00300.13	-3	Phosphoglycerate mutase family
PGI	PF00342.10	-168.9	Phosphoglucose isomerase
PGK	PF00162.10	-39.3	Phosphoglycerate kinase
PH	PF00169.20	25.1	PH domain
PHD	PF00628.20	26.2	PHD-finger
PI-PLC-X	PF00388.10	18.8	Phosphatidylinositol-specific phospholipase C, X domain
PI-PLC-Y	PF00387.10	-11	Phosphatidylinositol-specific phospholipase C, Y domain
PK	PF00224.12	-244	Pyruvate kinase, barrel domain
PK_C	PF02887.7	-44	Pyruvate kinase, alpha/beta domain
PLAC8	PF04749.8	-1.1	PLAC8 family
PMEI	PF04043.6	25	Plant invertase/pectin methylesterase inhibitor
PP-binding	PF00550.16	25.3	Phosphopantetheine attachment site
PP2C	PF00481.12	-36.3	Protein phosphatase 2C
PPDK_N	PF01326.10	-150.2	Pyruvate phosphate dikinase, PEP/pyruvate binding domain
PPR	PF01535.11	0	PPR repeat
PRA1	PF03208.10	25	PRA1 family protein
PRC	PF05239.7	25	PRC-barrel domain
PRK	PF00485.9	-28.5	Phosphoribulokinase / Uridine kinase family
PRMT5	PF05185.7	25	PRMT5 arginine-N-methyltransferase
PTA_PTB	PF01515.10	-165.5	Phosphate acetyl/butaryl transferase

Pfam domain name	Accession #	gathering cutoff	domain description
PTPA	PF03095.6	-106	Phosphotyrosyl phosphate activator (PTPA) protein
PTR2	PF00854.12	-50	POT family
PTS_2-RNA	PF01885.7	-48	RNA 2'-phosphotransferase, Tpt1 / KptA family
Pentapeptide	PF00805.13	0	Pentapeptide repeats (8 copies)
Peptidase_C12	PF01088.12	10	Ubiquitin carboxyl-terminal hydrolase, family 1
Peptidase_C14	PF00656.13	-22.5	Caspase domain
Peptidase_C48	PF02902.10	-13.1	Ulp1 protease family, C-terminal catalytic domain
Peptidase_M16	PF00675.11	-38	Insulinase (Peptidase family M16)
Peptidase_M16_C	PF05193.12	7.9	Peptidase M16 inactive domain
Peptidase_M18	PF02127.6	-250.8	Aminopeptidase I zinc metalloprotease (M18)
Peptidase_M24	PF00557.15	-70.7	metallopeptidase family M24
Peptidase_S8	PF00082.13	-51	Subtilase family
Per1	PF04080.4	25	Per1-like
Pescadillo_N	PF06732.2	-167.1	Pescadillo N-terminus
PfkB	PF00294.15	-67.8	pfkB family carbohydrate kinase
PhzC-PhzF	PF02567.7	-66	Phenazine biosynthesis-like protein
Pkinase	PF00069.16	70.3	Protein kinase domain
Pkinase_C	PF00433.15	14.1	Protein kinase C terminal domain
Pkinase_Tyr	PF07714.8	65	Protein tyrosine kinase
Polysacc_synt_2	PF02719.6	-176	Polysaccharide biosynthesis protein
Porin_3	PF01459.13	-53	Eukaryotic porin
Pre-SET	PF05033.7	3.9	Pre-SET motif
Pribosyltran	PF00156.18	2	Phosphoribosyl transferase domain
Pro_isomerase	PF00160.12	-37	Cyclophilin type peptidyl-prolyl cis-trans isomerase/CLD
Proteasome	PF00227.17	-36.7	Proteasome A-type and B-type
PsbW	PF07123.3	6.7	Photosystem II reaction centre W protein (PsbW)
Pyr_redox	PF00070.18	5	Pyridine nucleotide-disulphide oxidoreductase
Pyr_redox_2	PF07992.5	-20	Pyridine nucleotide-disulphide oxidoreductase
Pyr_redox_dim	PF02852.13	-13	Pyridine nucleotide-disulphide oxidoreductase, dimerisation domain
Pyridoxal_deC	PF00282.10	-158.6	Pyridoxal-dependent decarboxylase conserved domain
Pyrophosphatase	PF00719.10	-41	Inorganic pyrophosphatase
RALF	PF05498.2	25	Rapid Alkalinization Factor (RALF)
RAMP4	PF06624.3	25	Ribosome associated membrane protein RAMP4
RCC1	PF00415.9	21.2	Regulator of chromosome condensation (RCC1)
RNA_pol_I_A49	PF06870.3	25	A49-like RNA polymerase I associated factor
RNA_pol_N	PF01194.8	25	RNA polymerases N / 8 kDa subunit
RNA_pol_Rpb5_C	PF01191.10	25	RNA polymerase Rpb5, C-terminal domain
RNA_pol_Rpb5_N	PF03871.5	-8.9	RNA polymerase Rpb5, N-terminal domain
RNase_PH	PF01138.12	4	3' exoribonuclease family, domain 1
RNase_PH_C	PF03725.6	20	3' exoribonuclease family, domain 2
RPE65	PF03055.6	-156.5	Retinal pigment epithelial membrane protein
RRM_1	PF00076.13	17.7	RNA recognition motif. (a.k.a. RRM, RBD, or RNP domain)
RRM_2	PF04059.3	25	RNA recognition motif 2
RRM_3	PF08777.2	0.5	RNA binding motif
RRS1	PF04939.3	-60.5	Ribosome biogenesis regulatory protein (RRS1)
Raffinose_syn	PF05691.3	-395.7	Raffinose synthase or seed imbibition protein Sip1
Ras	PF00071.13	-69.9	Ras family
Redoxin	PF08534.1	-1	Redoxin
Rep_fac_C	PF08542.2	-0.4	Replication factor C

Pfam domain name	Accession #	gathering cutoff	domain description
Response_reg	PF00072.15	4	Response regulator receiver domain
Rho_GDI	PF02115.8	-55	RHO protein GDP dissociation inhibitor
Rhodanese	PF00581.11	25	Rhodanese-like domain
Rib_5-P_isom_A	PF06026.5	-84.5	Ribose 5-phosphate isomerase A (phosphoriboisomerase A)
Ribosomal_L12	PF00542.10	25	Ribosomal protein L7/L12 C-terminal domain
Ribosomal_L7Ae	PF01248.17	6	Ribosomal protein L7Ae/L30e/S12e/Gadd45 family
Ribosomal_S5	PF00333.11	25	Ribosomal protein S5, N-terminal domain
Ribosomal_S5_C	PF03719.6	0	Ribosomal protein S5, C-terminal domain
Ribosomal_S6e	PF01092.10	25	Ribosomal protein S6e
Ribul_P_3_epim	PF00834.10	-97.3	Ribulose-phosphate 3 epimerase family
Rick_17kDa_Anti	PF05433.6	23.2	Rickettsia 17 kDa surface antigen
RmID_sub_bind	PF04321.8	-171.8	RmID substrate binding domain
SAM_1	PF00536.21	11.3	SAM domain (Sterile alpha motif)
SAM_2	PF07647.8	0	SAM domain (Sterile alpha motif)
SATase_N	PF06426.5	25	Serine acetyltransferase, N-terminal
SBF	PF01758.7	-27.8	Sodium Bile acid symporter family
SET	PF00856.19	23.5	SET domain
SFT2	PF07770.3	-14.4	SFT2-like protein
SH3_1	PF00018.19	-8	SH3 domain
SH3_2	PF07653.8	0	Variant SH3 domain
SIP1	PF04938.3	25	Survival motor neuron (SMN) interacting protein 1 (SIP1)
SMP	PF04927.3	25	Seed maturation protein
SNARE	PF05739.10	23.8	SNARE domain
SNF5	PF04855.3	25	SNF5 / SMARCB1 / INI1
SQS_PSY	PF00494.10	-78	Squalene/phytoene synthase
SRF-TF	PF00319.9	11	SRF-type transcription factor (DNA-binding and dimerisation domain)
SRPRB	PF09439.1	-45.6	Signal recognition particle receptor beta subunit
SSB	PF00436.16	0.4	Single-strand binding protein family
STAS	PF01740.12	0	STAS domain
STT3	PF02516.5	-175	Oligosaccharyl transferase STT3 subunit
SWIB	PF02201.9	-5.7	SWIB/MDM2 domain
SYF2	PF08231.3	25	SYF2 splicing factor
S_locus_glycop	PF00954.11	-12.7	S-locus glycoprotein family
Sad1_UNC	PF07738.4	-20.4	Sad1 / UNC-like C-terminal
Sec1	PF00995.14	-272	Sec1 family
Sec61_beta	PF03911.7	25	Sec61beta family
Sedlin_N	PF04628.4	-22.1	Sedlin, N-terminal conserved region
SeIR	PF01641.9	-66.5	SeIR domain
Shikimate_DH	PF01488.11	-4.4	Shikimate / quinate 5-dehydrogenase
Sina	PF03145.7	-48.4	Seven in absentia protein family
Skp1	PF01466.10	-2	Skp1 family, dimerisation domain
Skp1_POZ	PF03931.6	14.9	Skp1 family, tetramerisation domain
Smg4_UPF3	PF03467.6	-27.5	Smg-4/UPF3 family
Snf7	PF03357.12	-22.9	Snf7
Spc97_Spc98	PF04130.4	-136.8	Spc97 / Spc98 family
Steroid_dh	PF02544.7	-44.7	3-oxo-5-alpha-steroid 4-dehydrogenase
Str_synth	PF03088.7	4.7	Strictosidine synthase
Subtilisin_N	PF05922.7	26.1	Subtilisin N-terminal Region
Suc_Fer-like	PF06999.3	-42.4	Sucrase/ferredoxin-like

Pfam domain name	Accession #	gathering cutoff	domain description
Sugar_tr	PF00083.15	-85	Sugar (and other) transporter
Sulfate_transp	PF00916.11	-131.5	Sulfate transporter family
Sybindin	PF04099.3	-44.4	Sybindin-like family
Synaptobrevin	PF00957.12	25	Synaptobrevin
Syntaxin	PF00804.16	11.4	Syntaxin
TATA_RF	PF08612.2	25	TATA-binding related factor (TRF)
TB2_DP1_HVA22	PF03134.10	-6.9	TB2/DP1, HVA22 family
TBC	PF00566.9	-58	TBC domain
TBP	PF00352.12	-8	Transcription factor TFIID (or TATA-binding protein, TBP)
TFIID-18kDa	PF02269.7	-7.3	Transcription initiation factor IID, 18kD subunit
TFIID-31kDa	PF02291.6	25	Transcription initiation factor IID, 31kD subunit
TFIID_30kDa	PF03540.4	25	Transcription initiation factor TFIID 23-30kDa subunit
TIM	PF00121.9	-97	Triosephosphate isomerase
TLC	PF03219.5	25	TLC ATP/ADP transporter
TPP_enzyme_C	PF02775.12	19.7	Thiamine pyrophosphate enzyme, C-terminal TPP binding domain
TPP_enzyme_M	PF00205.13	-8.1	Thiamine pyrophosphate enzyme, central domain
TPP_enzyme_N	PF02776.9	-70	Thiamine pyrophosphate enzyme, N-terminal TPP binding domain
TPT	PF03151.7	-15.3	Triose-phosphate Transporter family
Tbf5	PF06331.3	25	Transcription factor TFIH complex subunit Tbf5
Tetraspanin	PF00335.11	-15.4	Tetraspanin family
ThiF	PF00899.12	-38.4	ThiF family
Thioredoxin	PF00085.11	-25.7	Thioredoxin
Thr_dehydrat_C	PF00585.9	-7.9	C-terminal regulatory domain of Threonine dehydratase
Transket_pyr	PF02779.15	-49	Transketolase, pyrimidine binding domain
Transketolase_C	PF02780.11	-15.5	Transketolase, C-terminal domain
TrkA_C	PF02080.12	25	TrkA-C domain
TrkA_N	PF02254.9	4.7	TrkA-N domain
Tryp_alpha_amyl	PF00234.13	-0.2	Protease inhibitor/seed storage/LTP family
Tyr-DNA_phospho	PF06087.3	-152.8	Tyrosyl-DNA phosphodiesterase
U-box	PF04564.6	-7.6	U-box domain
U3_snoRNA_C	PF09384.1	25	U3 small nucleolar RNA C terminal
UBA	PF00627.22	20.7	UBA/TS-N domain
UBX	PF00789.11	10	UBX domain
UDPG_MGDP_dh	PF00984.10	-6.9	UDP-glucose/GDP-mannose dehydrogenase family, central domain
UDPG_MGDP_dh_C	PF03720.6	0.8	UDP-glucose/GDP-mannose dehydrogenase family, UDP binding domain
UDPG_MGDP_dh_N	PF03721.5	-74.8	UDP-glucose/GDP-mannose dehydrogenase family, NAD binding domain
UIM	PF02809.11	16.6	Ubiquitin interaction motif
UPF0061	PF02696.5	-200	Uncharacterized ACR, YdiU/UPF0061 family
UPF0139	PF03669.4	25	Uncharacterised protein family (UPF0139)
UPF0153	PF03692.6	11.7	Uncharacterised protein family (UPF0153)
UPF0185	PF03671.5	11.6	Uncharacterised protein family (UPF0185)
UQ_con	PF00179.17	-30	Ubiquitin-conjugating enzyme
Ubie_methyltran	PF01209.9	-117	ubiE/COQ5 methyltransferase family
Usp	PF00582.17	21.6	Universal stress protein family
V-ATPase_G	PF03179.6	-19.8	Vacuolar (H ⁺)-ATPase G subunit
VHS	PF00790.10	-13.2	VHS domain
VQ	PF05678.5	25	VQ motif
WAK	PF08488.2	25	Wall-associated kinase

Pfam domain name	Accession #	gathering cutoff	domain description
WD40	PF00400.23	21.5	WD domain, G-beta repeat
Whi5	PF08528.2	25	Whi5 like
X8	PF07983.4	-28.8	X8 domain
XH	PF03469.5	25	XH domain
XS	PF03468.5	25	XS domain
YDG_SRA	PF02182.8	25	YDG/SRA domain
YIF1	PF03878.6	-81.2	YIF1
YTH	PF04146.6	25	YT521-B-like family
Y_phosphatase2	PF03162.4	-47.6	Tyrosine phosphatase family
YgbB	PF02542.7	25	YgbB family
Zip	PF02535.13	-25.6	ZIP Zinc transporter
adh_short	PF00106.16	-40.2	short chain dehydrogenase
dsrm	PF00035.16	24.6	Double-stranded RNA binding motif
eIF2A	PF08662.2	0	Eukaryotic translation initiation factor eIF2A
eIF3_subunit	PF08597.1	11.9	Translation initiation factor eIF3 subunit
efhand	PF00036.23	21.7	EF hand
efhand_like	PF09279.2	8.8	Phosphoinositide-specific phospholipase C, efhand-like
iPGM_N	PF06415.4	-263.4	BPG-independent PGAM N-terminus (iPGM_N)
mTERF	PF02536.5	-60	mTERF
malic	PF00390.10	25	Malic enzyme, N-terminal domain
p450	PF00067.13	-105	Cytochrome P450
peroxidase	PF00141.14	-10	Peroxidase
polyprenyl_synt	PF00348.8	-43	Polyprenyl synthetase
tRNA-synt_1c	PF00749.12	-130.2	tRNA synthetases class I (E and Q), catalytic domain
tRNA-synt_1c_C	PF03950.9	25	tRNA synthetases class I (E and Q), anti-codon binding domain
zf-A20	PF01754.7	25	A20-like zinc finger
zf-AN1	PF01428.7	0	AN1-like Zinc finger
zf-C2H2	PF00096.17	17.7	Zinc finger, C2H2 type
zf-C3HC4	PF00097.16	16	Zinc finger, C3HC4 type (RING finger)
zf-CCCH	PF00642.15	0	Zinc finger C-x8-C-x5-C-x3-H type (and similar)
zf-CCHC	PF00098.14	17.9	Zinc knuckle
zf-DHHC	PF01529.11	-11	DHHC zinc finger domain
zf-MYND	PF01753.9	11	MYND finger
zf-Tim10_DDP	PF02953.6	-5	Tim10/DDP family zinc finger

Example 10. Selection of transgenic plants with enhanced agronomic trait(s)

This example illustrates identification of plant cells of the invention by screening derived plants and seeds for enhanced trait. Transgenic seed and plants in corn, soybean, cotton or canola with recombinant DNA identified in Table 2 are prepared by plant cells transformed with DNA that is stably integrated into the genome of the corn cell. Transgenic corn plant cells are transformed with recombinant DNA from each of the genes identified in Table 2. Progeny transgenic plants and seed of the transformed plant cells are screened for enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil as compared to control plants

A. Selection for enhanced Nitrogen Use Efficiency

Transgenic corn seeds provided by the present invention are planted in fields with three levels of nitrogen (N) fertilizer being applied, *e.g.* low level (0 N), medium level (80 lb/ac) and high level (180 lb/ac). A variety of physiological traits are monitored. Plants with enhanced NUE provide higher yield as compared to control plants.

5 **B. Selection for increased yield**

Effective selection of enhanced yielding transgenic plants uses hybrid progeny of the transgenic plants for corn, cotton, and canola, or inbred progeny of transgenic plants for soybeanplants plant such as corn, cotton, canola, or inbred plant such as soy, canola and cottoncotton over multiple locations with plants grown under optimal production
10 management practices, and maximum pest control. A target for improved yield is about a 5% to 10% increase or more in yield as compared to yield produced by plants grown from seed for a control plant. Selection methods can be applied in multiple and diverse geographic locations, for example up to 16 or more locations, over one or more planting seasons, for example at least two planting seasons, to statistically distinguish yield improvement from
15 natural environmental effects.

C. Selection for enhanced water use efficiency (WUE)

The selection process imposes a water withholding period to induce stressdrought followed by watering. For for example, for corn, a selection process imposes 3 drought/re-water cycles on plants over a total period of 15 days after an initial stress free growth period
20 of 11 days. Each cycle consists of 5 days, with no water being applied for the first four days and a water quenching on the 5th day of the cycle. The primary phenotypes analyzed by the selection method are the changes in plant growth rate as determined by height and biomass during a vegetative drought treatment.

D. Selection for Growth Under Cold Stress

25 (1) Cold germination assay - Trays of transgenic and control seeds are placed in a growth chamber at 9.7°C for 24 days (no light). Seeds having higher germination rates as compared to the control are identified.

(2) Cold field efficacy trial - A cold field efficacy trial is used to identify gene constructs that confer enhanced cold vigor at germination and early seedling growth under
30 early spring planting field conditions in conventional-till and simulated no-till environments. Seeds are planted into the ground around two weeks before local farmers begin to plant corn so that a significant cold stress is exerted onto the crop, named as cold treatment. Seeds also are planted under local optimal planting conditions such that the crop has little or no exposure to cold condition, named as normal treatment. At each location, seeds are planted under both

cold and normal conditions with 3 repetitions per treatment. Two temperature monitors are set up at each location to monitor both air and soil temperature daily.

Seed emergence is defined as the point when the growing shoot breaks the soil surface. The number of emerged seedlings in each plot is counted everyday from the day the earliest plot begins to emerge until no significant changes in emergence occur. In addition, for each planting date, the latest date when emergence is 0 in all plots is also recorded. Seedling vigor is also rated at V3-V4 stage before the average of corn plant height reaches 10 inches, with 1=excellent early growth, 5=Average growth and 9=poor growth. Days to 50% emergence, maximum percent emergence and seedling vigor are used to determine plants with enhanced cold tolerance.

E. Screens for transgenic plant seeds with increased protein and/or oil levels

This example sets forth a high-throughput selection for identifying plant seeds with improvement in seed composition using the Infratec 1200 series Grain Analyzer, which is a near-infrared transmittance spectrometer used to determine the composition of a bulk seed sample (Table 26). Near infrared analysis is a non-destructive, high-throughput method that can analyze multiple traits in a single sample scan. An NIR calibration for the analytes of interest is used to predict the values of an unknown sample. The NIR spectrum is obtained for the sample and compared to the calibration using a complex chemometric software package that provides predicted values as well as information on how well the sample fits in the calibration.

Infratec Model 1221, 1225, or 1227 with transport module by Foss North America is used with cuvette, item # 1000-4033, Foss North America or for small samples with small cell cuvette, Foss standard cuvette modified by Leon Girard Co. Corn and soy check samples of varying composition maintained in check cell cuvettes are supplied by Leon Girard Co. NIT collection software is provided by Maximum Consulting Inc. Software. Calculations are performed automatically by the software. Seed samples are received in packets or containers with barcode labels from the customer. The seed is poured into the cuvettes and analyzed as received.

Table 23

Typical sample(s):	Whole grain corn and soybean seeds
Analytical time to run method:	Less than 0.75 min per sample
Total elapsed time per run:	1.5 minute per sample

Typical and minimum sample size:	Corn typical: 50cc; minimum 30cc Soybean typical: 50cc; minimum 5cc
Typical analytical range:	Determined in part by the specific calibration. Corn – moisture 5-15%, oil 5-20%, protein 5-30%, starch 50-75%, and density 1.0-1.3%. Soybean – moisture 5-15%, oil 15-25%, and protein 35-50%.

Example 11. Cotton transgenic plants with enhanced agronomic traits

Cotton transformation is performed as generally described in WO0036911 and in U.S. Pat. No. 5,846,797. Transgenic cotton plants containing each of the recombinant DNA having a sequence of SEQ ID NO: 1 through SEQ ID NO: 803 are obtained by transforming with recombinant DNA from each of the genes identified in Table 2. Progeny transgenic plants are selected from a population of transgenic cotton events under specified growing conditions and are compared with control cotton plants. Control cotton plants are substantially the same cotton genotype but without the recombinant DNA, for example, either a parental cotton plant of the same genotype that was not transformed with the identical recombinant DNA or a negative isoline of the transformed plant. Additionally, a commercial cotton cultivar adapted to the geographical region and cultivation conditions, *e.g.* cotton variety ST474, cotton variety FM 958, and cotton variety Siokra L-23, are used to compare the relative performance of the transgenic cotton plants containing the recombinant DNA. The specified culture conditions are growing a first set of transgenic and control plants under “wet” conditions, *e.g.* irrigated in the range of 85 to 100 percent of evapotranspiration to provide leaf water potential of -14 to -18 bars, and growing a second set of transgenic and control plants under “dry” conditions, *e.g.* irrigated in the range of 40 to 60 percent of evapotranspiration to provide a leaf water potential of -21 to -25 bars. Pest control, such as weed and insect control is applied equally to both wet and dry treatments as needed. Data gathered during the trial includes weather records throughout the growing season including detailed records of rainfall; soil characterization information; any herbicide or insecticide applications; any gross agronomic differences observed such as leaf morphology, branching habit, leaf color, time to flowering, and fruiting pattern; plant height at various points during the trial; stand density; node and fruit number including node above white flower and node above crack boll

measurements; and visual wilt scoring. Cotton boll samples are taken and analyzed for lint fraction and fiber quality. The cotton is harvested at the normal harvest timeframe for the trial area. Enhanced water use efficiency is indicated by increased yield, improved relative water content, enhanced leaf water potential, increased biomass, enhanced leaf extension rates, and improved fiber parameters.

The transgenic cotton plants of this invention are identified from among the transgenic cotton plants by agronomic trait screening as having increased yield and enhanced water use efficiency.

Example 12. Canola plants with enhanced agrominic traits

This example illustrates plant transformation in producing the transgenic canola plants of this invention and the production and identification of transgenic seed for transgenic canola having enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

Tissues from in vitro grown canola seedlings are prepared and inoculated with a suspension of overnight grown *Agrobacterium* containing plasmid DNA with the gene of interest cassette and a plant selectable marker cassette. Following co-cultivation with *Agrobacterium*, the infected tissues are allowed to grow on selection to promote growth of transgenic shoots, followed by growth of roots from the transgenic shoots. The selected plantlets are then transferred to the greenhouse and potted in soil. Molecular characterizations are performed to confirm the presence of the gene of interest, and its expression in transgenic plants and progenies. Progeny transgenic plants are selected from a population of transgenic canola events under specified growing conditions and are compared with control canola plants. Control canola plants are substantially the same canola genotype but without the recombinant DNA, for example, either a parental canola plant of the same genotype that is not transformed with the identical recombinant DNA or a negative isoline of the transformed plant

Transgenic canola plant cells are transformed with recombinant DNA from each of the genes identified in Table 2. Transgenic progeny plants and seed of the transformed plant cells are screened for enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

Example 13. Selection of transgenic plants with enhanced agronomic trait(s)

This example illustrates the preparation and identification by selection of transgenic seeds and plants derived from transgenic plant cells of this invention where the plants and seed are identified by screening for a transgenic plant having an enhanced

agronomic trait imparted by expression or suppression of a protein selected from the group including the homologous proteins identified in Example 7. Transgenic plant cells of corn, soybean, cotton, canola, alfalfa, wheat and rice are transformed with recombinant DNA for expressing or suppressing each of the homologs identified in Example 7. Plants are regenerated from the transformed plant cells and used to produce progeny plants and seed that are screened for enhanced water use efficiency, enhanced cold tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil. Plants are identified exhibiting enhanced traits imparted by expression or suppression of the homologous proteins.

Example 14. Monocot and dicot plant transformation for the suppression of endogenous protein

This example illustrates monocot and dicot plant transformation to produce nuclei of this invention in cells of a transgenic plant by transformation where the recombinant DNA suppresses the expression of an endogenous protein identified in Table 24.

Corn, soybean, cotton, or canola tissue are transformed as described in Examples 2-5 using recombinant DNA in the nucleus with DNA that is transcribed into RNA that forms double-stranded RNA targeted to an endogenous gene with DNA encoding the protein. The genes for which the double-stranded RNAs are targeted are the native gene in corn, soybean, cotton or canola that are homologs of the genes encoding the protein that has the function of the protein of *Arabidopsis thaliana* as identified in Table 24.

Populations of transgenic plants prepared in Examples 3, 4, 5, 6, or 13 with DNA for suppressing a gene identified in Table 2 as providing an enhanced trait by gene suppression are screened to identify an event from those plants with a nucleus of the invention by selecting the trait identified in this specification.

Table 24.

PEP SEQ ID	Pfam module	Construct ID	Traits	
115	AP2	10177	LN	
116	zf-C2H2	12155	CS	SS
118	F-box::Tub	11873	LN	PEG
154	Pex2_Pex12::zf-C3HC4	11113	CS	
188	Myb_DNA-binding::Myb_DNA-binding	12325	LN	
200	bZIP_1	71228	CK	

38-21(54976)-PCT

What is claimed is:

1. A plant cell nucleus with stably-integrated, recombinant DNA comprising a promoter that is functional in plant cells and that is operably linked to protein coding DNA encoding a protein having an amino acid sequence comprising a Pfam domain module

5 NTP_transferase::Hexapep::Hexapep::Hexapep;

wherein said plant cell nucleus is selected by screening a population of transgenic plants that have said recombinant DNA in its nuclei and express said protein for an enhanced trait as compared to control plants that do not have said recombinant DNA in their nuclei; and

10 wherein said enhanced trait is selected from group of enhanced traits consisting of enhanced water use efficiency, enhanced cold tolerance, enhanced heat tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

2. A plant cell nucleus with stably integrated, recombinant DNA, wherein

(a) said recombinant DNA comprises a promoter that is functional in said plant cell and that is operably linked to a protein coding DNA encoding a protein having an amino acid sequence
 15 comprising a Pfam domain module selected from the group consisting of Syntaxin::SNARE, Pro_isomerase, Pkinase, ATP-synt_G, Carboxyl_trans, CDC50, GATase::GMP_synt_C, F-box::WD40::WD40::WD40, dsrm::dsrm, Pyr_redox_2::Pyr_redox_dim, WAK::Pkinase, Pkinase_Tyr, PTPA, Biotin_lipoyl::E3_binding::2-oxoacid_dh, AAA, LRRNT_2::LRR_1::LRR_1::LRR_1, PRA1, TIM, YTH, ThiF, Hep_59, Pkinase,
 20 PALP::Thr_dehydrat_C::Thr_dehydrat_C, zf-Tim10_DDP, CAF1, Pkinase, Pyr_redox_2::Pyr_redox_dim, Response_reg, YIF1, NPH3, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::Pkinase_Tyr, NOI, Fer2, AA_permease, Caleosin, IF4E, Pkinase, Ribul_P_3_epim, F-box::WD40::WD40::WD40::WD40::WD40::WD40, MIP, p450, Ribosomal_S5::Ribosomal_S5_C, Hpt, TBC, Acyltransferase, Epimerase, Pkinase, Dus,
 25 SelR, PH, Tyr-DNA_phospho, RRM_1, Pkinase, ECH, SRPRB, DUF599, Pkinase::efhand::efhand::efhand::efhand, ORC2, PME1, p450, GLTP, Suc_Fer-like, CBS::CBS, ADH_N::ADH_zinc_N, LRRNT_2::LRR_1::LRR_1::LRR_1::Pkinase, F-box, adh_short, polyprenyl_synt, GHMP_kinases_N::GHMP_kinases_C, PsbW, Na_H_Exchange::TrkA_C, Ribosomal_L12, CDC50, DCP1, Pyr_redox_2::Pyr_redox_dim, Steroid_dh, tRNA-
 30 synt_1c::tRNA-synt_1c_C, Cupin_1::Cupin_1, peroxidase, Sugar_tr, Pro_isomerase, Iso_dh,

38-21(54976)-PCT

- PTS_2-RNA, Aminotran_5, ENOD93, KH_1::KH_1::KH_1::KH_1::KH_1,
Mito_carr::Mito_carr::Mito_carr, DUF569::DUF569, C1_2::C1_2::C1_3::C1_2::C1_3::C1_2,
DnaJ::zf-C2H2, Aldedh, AAA::Rep_fac_C, DUF641, Aminotran_1_2, TB2_DP1_HVA22, zf-
C3HC4, RRM_1::RRM_2, Pkinase,
- 5 PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR, Pyr_redox_2, IQ::IQ,
SMP::SMP::SMP, AWPM-19, CLP_protease, Pkinase::NAF, PRMT5,
PPR::PPR::PPR::PPR::PPR::PPR, Aldedh, ATP-synt_C::ATP-synt_C, adh_short,
RCC1::RCC1::RCC1::RCC1::FYVE::DZC, p450, PfkB, PB1,
Hexapep::Hexapep::Hexapep::Hexapep, Aminotran_1_2, DUF862, Aldedh,
- 10 ADH_N::ADH_zinc_N, CDC48_N::AAA::AAA, AA_permease, UQ_con, Mem_trans, GFA,
OPT, DUF887, Di19, U-box::Arm::Arm, DAO, G-alpha, Aminotran_3, 2OG-FeII_Oxy,
Lectin_legB::Pkinase, NAC::UBA, efhand_like::PI-PLC-X::PI-PLC-Y::C2, TLC, ATP-
grasp_2::Ligase_CoA, Pkinase, PRC::PRC, MFS_1::Sugar_tr, Copine::zf-C3HC4,
RRM_1::RRM_1::RRM_1, Pribosyltran, AA_kinase, PhzC-PhzF, Gp_dh_N::Gp_dh_C,
- 15 Sugar_tr, Pre-SET::SET, Alpha-amylase::Alpha-amyl_C2, Cyclin_N,
FAE1_CUT1_RppA::ACP_syn_III_C, RPE65, efhand_like::PI-PLC-X::PI-PLC-Y::C2,
Mito_carr::Mito_carr::Mito_carr, Invertase_neut, DSPc, LANC_like, Aminotran_5,
Glutaminase, PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR, PGI, SSB, p450,
La::RRM_1::RRM_3, Methyltransf_11, SBF, NTP_transferase::Hexapep::Hexapep::Hexapep,
- 20 Mov34, Hydrolase, AAA, RRM_1::RRM_1, Pkinase::NAF, F-box::FBA_1, p450, F-
box::LRR_2, zf-DHHC, TBP::TBP, DUF607, FMO-like, adh_short, ATP-synt_ab_N::ATP-
synt_ab, RNase_PH, PP2C, UQ_con, Aminotran_1_2, p450, FMO-like, Gp_dh_N::Gp_dh_C,
Mito_carr::Mito_carr::Mito_carr, zf-CCCH::zf-CCCH::zf-CCCH::zf-CCCH::zf-CCCH, GST_C,
GH3, PAP_fibrillin, ThiF, Ribul_P_3_epim, DUF260, p450, MACPF, Steroid_dh,
- 25 Response_reg, Carb_anhydrase, Aldedh, Pkinase, Methyltransf_11, PK::PK_C, Pkinase,
DEK_C::SWIB::SWIB, TATA_RF, Tryp_alpha_amyl, Y_phosphatase2, TATA_RF, NifU_N,
ENT, Pkinase, F-box::Kelch_2::Kelch_2, WD40::WD40, MtN3_slv::MtN3_slv,
TB2_DP1_HVA22, Pkinase, Aldedh, AAA, Pyridoxal_deC, zf-CCHC::zf-CCHC::zf-CCHC::zf-
CCHC, Peptidase_M24, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::Pkinase, Arf, DUF1279,
- 30 UPF0185, Rhodanese, adh_short, Tryp_alpha_amyl, Pkinase, UPF0139, CBS, Glyco_transf_29,
iPGM_N::Metalloenzyme, Proteasome, Pkinase, UBX, ELFV_dehydrog_N::ELFV_dehydrog,

38-21(54976)-PCT

- p450, ADH_N::ADH_zinc_N, zf-C3HC4, NDK, NAD_Gly3P_dh_N::NAD_Gly3P_dh_C, GlutR_N::Shikimate_DH::GlutR_dimer, p450, SNF5, p450, Pribosyltran, AIG1, Response_reg::CCT, mTERF, DUF220, Pkinase::NAF, zf-CCCH::zf-CCCH, Pkinase, Pkinase, MATH::MATH, Asparaginase, Pkinase, Gp_dh_N::Gp_dh_C, Pkinase, KH_1, Fcf1,
- 5 PK::PK_C::PEP-utilizers, CRAL_TRIO_N::CRAL_TRIO, Raffinose_syn, DUF584, BT1, Aminotran_1_2, C1_2::C1_3::C1_3::C1_2::C1_3::C1_3::C1_2, FMO-like, PLAC8, F-box::Kelch_2::Kelch_1::Kelch_2, XS::XH, STT3, ABC_tran::ABC_tran, Aldo_ket_red, NTF2::RRM_1, DUF594, Biotin_lipoyl::2-oxoacid_dh, RRM_1, FHA, Mov34, MT-A70, Brix, X8, Auxin_inducible, Peptidase_M18, Sec61_beta, F-box::Kelch_1::Kelch_1, Brix, TIM,
- 10 CTP_transf_1, Tetraspannin, PALP, DUF822, Pkinase, B3::B3, MFS_1, SFT2, F-box::Kelch_1::Kelch_1, DUF588, Thioredoxin, Alba, Ion_trans_2::Ion_trans_2, Peptidase_C48, YDG_SRA::Pre-SET::SET, Aldedh, Abhydrolase_3, p450, U-box::Arm::Arm::Arm::Arm, AA_permease, LAG1, peroxidase, PAD_porph, Pkinase, F-box, AIG2, p450, AARP2CN::DUF663, UPF0153, NAD_binding_2::6PGD, Pkinase,
- 15 B_lectin::S_locus_glycop::PAN_1::Pkinase, EMP24_GP25L, DUF6::DUF6, DUF163, Sina, adh_short, Rho_GDI, zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC::zf-CCHC, VHS::GAT, Pyrophosphatase, DMRL_synthase, Cyclin_N::Cyclin_C, NDK, TPP_enzyme_N::TPP_enzyme_M::TPP_enzyme_C, SH3_1, UDPG_MGDP_dh_N::UDPG_MGDP_dh::UDPG_MGDP_dh_C, UIM::efhand, DUF579,
- 20 Pkinase, DHBP_synthase::GTP_cyclohydro2, Peptidase_C14, Glycolytic, Pkinase, Lipase_3, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1, Snf7, BT1, PPK_N::PEP-utilizers::PEP-utilizers_C, V-ATPase_G, RNA_pol_I_A49, Zip, WD40::WD40::WD40, Branch, RCC1::RCC1::RCC1::RCC1, TFIID-18kDa, Pkinase, Cullin, TFIID-31kDa, Asp, Pkinase, Ubie_methyltran, PPR::PPR, PPR::PPR::PPR::PPR::PPR::PPR::PPR, Fructosamin_kin,
- 25 PPR::PPR::PPR::PPR::PPR::PPR, Pkinase_Tyr, p450, AA_permease, Glycos_transf_1, ECH::3HCDH_N::3HCDH, MIP, ADH_N::ADH_zinc_N, Aa_trans, DUF1517, Redoxin, RRS1, Pribosyltran, Sulfate_transp::STAS, DUF167, MMR_HSR1::DUF933, Aminotran_1_2, AA_permease, DUF620, Aldedh, Aminotran_1_2, LRR_1, CDP-OH_P_transf, DHBP_synthase::GTP_cyclohydro2, NHL::NHL, WD40::U3_snoRNA_C, Na_sulph_symp,
- 30 adh_short, Thioredoxin, Ribosomal_L7Ae, OTCace_N::OTCace, Mito_carr::Mito_carr::Mito_carr, Pescadillo_N::BRCT,

38-21(54976)-PCT

PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR::PPR, mTERF, Chalcone, Per1, PPR::PPR,
 Na_H_Exchanger::TrkA_N, DEAD::Helicase_C, Cyclin_N::Cyclin_C, PP-binding, SRF-TF,
 SATase_N::Hexapep::Hexapep::Hexapep, PK::PK_C, adh_short, DUF6::DUF6, Pyr_redox_2,
 G-patch, WD40::WD40::WD40::WD40, G-alpha, VQ, WD40::WD40::WD40::WD40::WD40,
 5 Aminotran_3, Pribosyltran, LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1,
 eIF3_subunit, H_PPase, Nodulin-like, Whi5, Tryp_alpha_amyl,
 NIR_SIR_ferr::NIR_SIR::NIR_SIR_ferr, RRM_1::RRM_1, Band_7, Asp, MFS_1,
 F420_oxidored, Pkinase, PRK, Na_Ca_ex::Na_Ca_ex, Bombesin, Pkinase, Epimerase, Pkinase,
 Inositol_P, Arm::Arm::Arm::Arm, PB1, Hin1, p450, RNA_pol_Rpb5_N::RNA_pol_Rpb5_C,
 10 Adap_comp_sub, Peptidase_M16::Peptidase_M16_C, Pkinase, PGK,
 Biotin_lipoyl::E3_binding::2-oxoacid_dh, Alg6_Alg8, F-box, MMR_HSR1,
 G6PD_N::G6PD_C, DAD, malic::Malic_M::PTA_PTB, ADH_N::ADH_zinc_N,
 LRR_1::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1, PfkB, PA,
 Pyr_redox_2::Pyr_redox_dim, SIP1, PP2C, BT1, AhpC-TSA, GASA, Aminotran_3, Sad1_UNC,
 15 Pkinase, zf-MYND::PDCD2_C, Pkinase, PGK, ArfGap, GATase::GMP_synt_C, p450,
 DUF1637, 2OG-FeII_Oxy, Aldedh, DUF231, Glyco_transf_8, Sec1, DUF1475, Cellulose_synt,
 RRM_1::RRM_1::RRM_1, Usp, AA_permease, Acetyltransf_1::Bromodomain,
 Glyco_tran_28_C, RNA_pol_N, NTP_transferase, malic::Malic_M, Histone, Epimerase,
 UPF0061, ClpS, LEA_5, Auxin_inducible, PAP_fibrillin, Aldo_ket_red, Gln-synt_C, DREPP,
 20 p450, SFT2, eIF2A, Cenp-O, CPSase_sm_chain::GATase, WD40::WD40::WD40, Pkinase,
 Porin_3, Pkinase, Aminotran_1_2, GST_N::GST_C, F-box::Kelch_1::Kelch_1, IU_nuc_hydro,
 SYF2, PDT, Subtilisin_N::PA, Sugar_tr, WD40::WD40, p450, DUF829,
 Pyr_redox_2::Pyr_redox_dim, Aminotran_1_2, Cyclin_N::Cyclin_C, YgbB, Sugar_tr, SRF-TF,
 DUF231, DUF584, BT1, zf-C3HC4, Metallophos, FA_hydroxylase, p450, ACBP::Ank::Ank,
 25 LSM, SAM_1, LRR_1::LRR_1::Pkinase, Cullin, F-box::FBA_1, peroxidase, Rib_5-P_isom_A,
 zf-A20::zf-AN1, Dimerisation::Methyltransf_2, PGAM, PTR2, Copine, PALP, Exo_endo_phos,
 G6PD_N::G6PD_C, PLAC8, Aldo_ket_red, Ank::Ank, FBpase, zf-C3HC4, ACT::ACT,
 Suc_Fer-like, Pkinase_Tyr, G-alpha, RAMP4, Glutaredoxin, RRM_1::RRM_1, MT-A70,
 Response_reg::CCT, Arf, AP2, DUF177, 2OG-FeII_Oxy,
 30 LRRNT_2::LRR_1::LRR_1::LRR_1::LRR_1::LRR_1, LRR_1, Aldedh, DUF1350,
 FAE1_CUT1_RppA::ACP_syn_III_C, Sybindin, zf-C3HC4, Nfu_N::NifU, Rho_GDI,

38-21(54976)-PCT

PK::PK_C, NOSIC::Nop, PK::PK_C, PRK::Pribosyltran, PfkB, p450, DUF231, TFIID_30kDa, Aldedh, zf-AN1, GHMP_kinases_N::GHMP_kinases_C, DUF423, Pkinase, Cu-oxidase_3::Cu-oxidase::Cu-oxidase_2, PCI, RNase_PH::RNase_PH_C, DUF59, NTP_transferase, CH::EB1, Pkinase::Pkinase_C, DUF866, SMP::SMP::SMP, Aminotran_3, Transket_pyr::Transketolase_C, Copine, His_biosynth, Tbf5, DUF543, p450, C2, DUF616, Gp_dh_N::Gp_dh_C, Smg4_UPF3, DUF231, DUF89, WD40::WD40, Aldedh, ACT::ACT::ACT::ACT, Pkinase, Pyridoxal_deC, Skp1_POZ::Skp1, NAP, FolB, p450, Pentapeptide::Pentapeptide, NTP_transferase::MannoseP_isomer, SQS_PSY, Cyclase, GASA, Rick_17kDa_Anti, FMO-like, B_lectin::S_locus_glycop::PAN_2, Peptidase_C12, Ribosomal_S6e, Glyoxalase, Response_reg::CCT, PHD::SET, Redoxin, G_glu_transpept, Synaptobrevin, p450, RALF, PALP, Branch, DUF579, Aminotran_3, Nramp, Enolase_N::Enolase_C, Str_synth, FAD_binding_3, MOSC_N::MOSC, Spc97_Spc98, Glycolytic, F-box::WD40::WD40::WD40::WD40, AA_kinase::ACT::ACT, RALF, and ATP_bind_1;

(b) said recombinant DNA comprises a promoter that is functional in said plant cell and that is operably linked to a protein coding DNA encoding a protein comprising an amino acid sequence with at least 90% identity to a consensus amino acid sequence selected from the group consisting of SEQ ID NO: 94617 through 94734;

(c) said recombinant DNA comprises a promoter that is functional in plant cells and that is operably linked to a protein coding DNA encoding a protein comprising an amino acid sequence selected from the group consisting of 819, 867, 906, 914, 967, 1009, 1040, 1086, 1107, 1227, 1233, 1250, 1293, 1311, 1320, 1342, 1364, 1365, 1373, 1398, 1402, 1439, 1443, 1473, 1495, 1539, 1596, and homologs thereof listed in table 19;

(d) said recombinant DNA comprises a promoter that is functional in said plant cell and that is operably linked to a protein coding recombinant DNA encoding a protein having an amino acid sequence having at least 70% identity to an amino acid sequence of SEQ ID NO 1227;

(e) said recombinant DNA suppresses comprises a promoter that is functional in said plant cell and operably linked to DNA that transcribe into RNA that suppresses the level of an endogenous protein wherein said endogenous protein has an amino acid sequence comprising a pfam domain module selected from the group consisting of DUF1475, DUF1350, Arm::Arm::Arm::Arm, NifU_N, ATP_bind_1, DUF599, LRR_1, Pyr_redox_2::Pyr_redox_dim, malic::Malic_M, Glycolytic, IF4E, Pkinase, UPF0139, GlutR_N::Shikimate_DH::GlutR_dimer, PLAC8, PLAC8,

38-21(54976)-PCT

DUF220, OPT, Glyco_transf_8, Per1, Usp, mTERF, PB1, CAF1, Pkinase, F-box::Kelch_1::Kelch_1, PHD::SET, Nfu_N::NifU, RCC1::RCC1::RCC1::RCC1, X8, Pkinase::NAF, UIM::efhand, Glutaredoxin, DUF1517, Cupin_1::Cupin_1, ClpS, Spc97_Spc98, Subtilisin_N::PA, CBS, PTPA, p450, AA_permease, Sina, p450, Mov34, Thioredoxin, SATase_N::Hexapep::Hexapep::Hexapep, Epimerase, RRM_1::RRM_2, Synaptobrevin, Ribosomal_S6e, Redoxin, MFS_1, PPR::PPR, NPH3, and DUF866; or

(f) said recombinant DNA comprises a promoter that is functional in said plant cell and operably linked to DNA that transcribe into RNA that suppresses the level of an endogenous protein wherein said endogenous protein has an amino acid sequence with at least 70% identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 811, 813, 819, 891, 893, 909, 911, 1439, 1584, and homologs thereof listed in table 16;

and wherein said plant cell nucleus is selected by screening a population of transgenic plants that have said recombinant DNA and an enhanced trait as compared to control plants that do not have said recombinant DNA in their nuclei; and wherein said enhanced trait is selected from group of enhanced traits consisting of enhanced water use efficiency, enhanced cold tolerance, enhanced heat tolerance, enhanced resistance to salt exposure, enhanced shade tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil.

3. The plant cell nucleus of claim 2 wherein said protein coding DNA encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NO: 804 through SEQ ID NO: 94613.

4. The plant cell nucleus of claim 2 further comprising DNA expressing a protein that provides tolerance from exposure to an herbicide applied at levels that are lethal to a wild type of said plant cell.

5. The plant cell nucleus of claim 4 wherein said herbicide is a glyphosate, dicamba, or glufosinate compound.

6. A transgenic plant cell or plant comprising a plurality of plant cells with the plant cell nucleus of claim 2.

38-21(54976)-PCT

7. The transgenic plant cell or plant of claim 6 which is homozygous for said recombinant DNA.
8. A transgenic seed comprising a plurality of plant cells with a plant cell nucleus of claim 2.
- 5 9. The transgenic seed of claim 8 from a corn, soybean, cotton, canola, alfalfa, wheat or rice plant.
10. A transgenic pollen grain comprising a haploid derivative of a plant cell nucleus of claim 2.
11. A method for manufacturing non-natural, transgenic seed that can be used to produce a
10 crop of transgenic plants with an enhanced trait resulting from expression of recombinant DNA in a nucleus of claim 2, wherein said method for manufacturing said transgenic seed comprising:
(a) screening a population of plants for said enhanced trait and said recombinant DNA, wherein individual plants in said population can exhibit said trait at a level less than, essentially the same as or greater than the level that said trait is exhibited in control plants which do not contain the
15 recombinant DNA, wherein said enhanced trait is selected from the group of enhanced traits consisting of enhanced water use efficiency, enhanced cold tolerance, enhanced heat tolerance, enhanced high salinity tolerance, enhanced shade tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein and enhanced seed oil,
(b) selecting from said population one or more plants that exhibit said trait at a level greater than
20 the level that said trait is exhibited in control plants, and
(c) collecting seeds from selected plant selected from step b.
12. The method of claim 11 wherein said method for manufacturing said transgenic seed further comprising
(a) verifying that said recombinant DNA is stably integrated in said selected plants, and
25 (b) analyzing tissue of said selected plant to determine the expression or suppression of a protein having the function of a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:804-1606.

38-21(54976)-PCT

13. The method of claim 11 wherein said seed is corn, soybean, cotton, alfalfa, canola wheat or rice seed.

14. A method of producing hybrid corn seed comprising:

- (a) acquiring hybrid corn seed from a herbicide tolerant corn plant which also has stably-integrated, recombinant DNA in a nucleus of claim 2;
- (b) producing corn plants from said hybrid corn seed, wherein a fraction of the plants produced from said hybrid corn seed is homozygous for said recombinant DNA, a fraction of the plants produced from said hybrid corn seed is hemizygous for said recombinant DNA, and a fraction of the plants produced from said hybrid corn seed has none of said recombinant DNA;
- (c) selecting corn plants which are homozygous and hemizygous for said recombinant DNA by treating with an herbicide;
- (d) collecting seed from herbicide-treated-surviving corn plants and planting said seed to produce further progeny corn plants;
- (e) repeating steps (c) and (d) at least once to produce an inbred corn line; and
- (f) crossing said inbred corn line with a second corn line to produce hybrid seed.

FIG. #1

1/13




FIG. #2

2/13




FIG. #3

3/13




FIG. #4

4/13

SEQ ID No 35795-----MS--
SEQ ID No 59830-----MSKN
SEQ ID No 23175-----MSKN
SEQ ID No 93133-----MSKN
5 SEQ ID No 19860-----
SEQ ID No 57392-----
SEQ ID No 91490-----
SEQ ID No 20139MLSVKPKKWLNRFNWSTGCARPIEDRLREAKKFLSLPVASSRSRLNLRSSQDLRS
SEQ ID No 88784-----
10 SEQ ID No 17591-----
SEQ ID No 43370-----
SEQ ID No 60203-----
SEQ ID No 1325-----
SEQ ID No 21321-----
15 SEQ ID No 45211-----MAVTTLRKKSASQGMLRTPSCPYPYISSAASSSSASMHGSSP
SEQ ID No 50671-----MTQTPTP-----
SEQ ID No 78019-----MALSTPT-----
SEQ ID No 45047-----
SEQ ID No 69645-----
20 SEQ ID No 13488-----
SEQ ID No 83296-----
SEQ ID No 58057-----
SEQ ID No 25554-----
SEQ ID No 4744-----
25 SEQ ID No 80927-----
SEQ ID No 68166-----
SEQ ID No 31956-----
SEQ ID No 29566-----
SEQ ID No 67810-----
30 SEQ ID No 36150-----
SEQ ID No 94673-----

35

40

45

50

FIG. #4

5/13

```

5  -----NGA-----GEFELPQEMLAVLPSPDY-----EQLDVARRITAM
   ESSYAGV-VGNNG-----GEFELPQEVFSVLPSDPY-----EQLDVARRITAM
   GSLYAGS-MGNG-----GEFELPQEVLSVLPSDPY-----EQLDVARRITAM
   RGFNAGSSMGNG-----GEFEISQELLSVLPSDPY-----EQLEVARRITGM
   -----MGNG-----GEFELSQEVLAFLPSDPY-----EQLDVARRITAM
10  -----MGNG-----GEFELSQEVSVLPSPDY-----EQIDIASRITAM
   -----MGSG-----GQGELPQDLVSVLPSPDY-----EQLDVARRITAM
   GCRLRPNGMSQNSN-YNSGSGPDFNITDDILAVIPTDPY-----DQLDLARKITSM
   -----MSQLT-----TG-GTDFSLPDEILAVIPMDPY-----DQLDLARKITSM
   -----MSSQG-----TGSVVDLFDLPDEILSVIPTDPY-----QQLDLARKITSM
15  -----MAHGG-----GGAPDFDLPEILSVMPDPY-----DQLDLARKITSM
   -----MGDDQ-----LDFELPEEVLSPVIMDPF-----EQDLARKITSM
   -----MGDDQ-----LDFELPEEVLSPVIMDPF-----EQDLARKITSM
   -----MGDDR-----LDLELSDEVLSVIMDPF-----EQDLARKITSM
   TGTRNLVAASPPPP---PLFPTSTGTYPAFISATTPDEC PAAALVMLPGKLEVARKITAV
20  -----AAAAPAP---AVAVSEAGLPDAIAAALPPDPY-----EQLEVARKITAV
   ----TAVAAPPPEPGPEPAPSAELPLDAIAAALPPDPY-----EQLEVARKITAV
   -----MPLAEG-----EFALPDEVLAVMRPDPY-----EQDLARRITAL
   -----MPLAEG-----EFALPDEVLAVMRPDPY-----EQDLARRITAL
   -----MPLAEG-----EFALPDEVLAVMRPDPY-----EQDLARRITAL
25  -----MPPAEGAAAGLVSEFALPDGVLAVLPRDPY-----EQDLARRITAL
   -----MPLGD-----DAAADFALPDELLAALPRDPY-----EQDLARRITAL
   -----MPLGD-----DAAADFALPDELLAALPRDPY-----EQDLARRITAL
   -----MPLGDG-----AEAADFVLPDELLAALPRDPY-----EQDLARRITAL
   -----MARHEAA---AGVVDLFHLPDEILAVIPTDPY-----EQLDVARKITSM
30  -----MARHEAA---AGVVDLFHLPDEILAVIPTDPY-----EQLDVARKITSM
   -----MARQEAVG-AATGVDFHLPDEILAVIPTDPY-----EQLDVARKITSM
   -----MSKGANYG-PDNASDFELPEELGILPTDPY-----EQLELARRITSI
   -----MSKAG-----EMQLADGILMNLPEHPD-----QQLEIAQQITRF
   -----
35  -----mxxxxx-----xxxxxfxlpexlxvxpdpdy-----eqldxarxitxx

```

40

45

50

6/13

10

50

7/13

10

20

25

30

35

40

45

50

FIG. #4

8/13

5

-----AESTVSRAQ--LLDDDGQSAK-SFSTSSSGTTEERGYREGAALKPPTRYQS
10 SSNLAIIRASQAESSLSR--ALDDDGQSTK-SFSTSSSVNTEERGFREGGASKPPTS YQT
NSSLAIIRASQAESTVSRSPILDDDHQPGKLSYESDSSVITEERGYRQGAVLKPPTSHQG
NPSLAIKRASQAESTVSRTPVSDDDYQSEKSSYD TDSSLTSEERGYQH--ASRPPTS YQN
PQSLALYKATQAETLSKAQI-FEDDGRYARASFSANPPQYHDDADSKS-----ATPTSQSA
PANSAMWKASRGTTT-----PKSQSA
15 APNSAIWKASQAESAVVKAQNFEDHGRYAKTSSRPDP PKYQGDLDVSDGRTPPKSRNG
TVDIGTCDQS---IPKAYPD---IDDG TNGYV V NQPLSGFTEYRNSTDEASRLAVQRF SV
TVDIGTCDQS---VPRAYSE---KDEGMNVYVAHHNFNGSTDMGN-TDEAPRHAGQRYSI
TIDIGTCDQS---VPKAYPD---KDDDGSGYMVHHSYNGPADVGKTND EASRYSGQRFSL
TVDIGTYDQT---VPKAYYT---GDE-PNGYTKHHSYSGSTESASLNDDVSKQTGQKH SI
20 PVDIRTC DQP-----GSYPG---KDGRINAHSIKQAYSGSTD TNNPVVEASKYTGNKFSM
PVDIRTC DQP-----GSYPG---KDGRINAHSIKQAYSGSTD TNNPVVEASKYTGNKFSM
PVDIKACDQSSIGSTGSYTG---KDERTTVHSTHRSYSGSSDMNPIVEASKYAGNKFSM
TVDIRTC EQSVA-KASSWKDGV AHSRHHHPVSSLADGSTEIES-VNQEVA-RPFEQKLSV
TVDIRTC EQSVA-KASSWKDGV AHSRHHHPVSSLADGSTEIES-VNQEVA-RPFEQKLSV
25 TVDIRTYEQSVA-KANSWKDGLEKN--SHPVSSLSD ESVEADS-VNQEVTTRPFEQKL TI
TVDITTC DQSISSKASSCGD---GDSITH TTTNLLSTSLDVGSTVQE--VSKPPIQKYAL
TVDITTC DQSISSKASSCGD---GDSITH TTTNLLSTSLDVGSTVQEGTVXKPPIQKYAL
--KMSGTNLSLIFSLN-IAD---GDSITH TTTNLLSTSLDVGSTVQEGTVSKPPIQKYAL
TVDITTC DQSVSSKTSSCGD---G-SISHTAANLLNGSGDLGSTTRE--AARTPVQKYAL
30 TVDITTC DQPVASKASSAD---GGSVSNPATNIFTESLDVGSTNRDGAAR PPIQKYSL
TVDITTC DQPVASKASSSAGYRYGGSVSNPATNIFTESLDVGSTNRDGT AAR PPIQKYSL
TVDITTC DQSVATKASSSGD---GGSGSHPTTNIFSES LDAGSTNRDGTASRPPIQKYAP
IGDDRDFDAN-----NNLTAR-----VPSWEDEVSSSRTSADSSSRSTMTESAQEHQFSV
IGDDRDFDAN-----NNLTAR-----VPSWK-----EHQFSV
35 TGEDRDVDAE-----NNGTAR-----IPSWKDEVSSSHTSSNTSSRSTITESAQGYQFSI
TQNTATRRVTSAKLSFSLSS-RDDSNVPIGETNQVSSAVSETSNSTLDGDNQASRHGKSK
NPGDATYAYAPEASASTYCS-----SEKSLQLPLIMSKSTEPPTPTKRKQQGQYSC
-----LEPSVAAADVAGERLVSEVLSSISKPPPPQMGV
XX

40

45

FIG. #4

9/13

5

10

15

-----SAHNSARG--TPHLSPKLTTP--SGSP-RGSGRTSPRRSMSLTDNSHRI SLPS
 -----SAHNSAQG--TPHLSPKLTSS--SG-----SGRASPRRSMSLTDNSHRI SLPS
 20 -----SAHNSARG--TPHLSPRLTTP--SGSPKQSQARTSPRRSMSSSENSHRI SLPS
 -----SAQNSARG--TPRLSPRLTTP--SVSPKQSQSARGSPQRSMSLTENSHRI SLPS
 RNSPKQIIRGTSSARG--TPSLTPRLTTP--TGSPKPQPKKGSPLPSS--YTLDSQHIQIPT
 HTSPKQIMRGIDSARGSMTPRLTPLHTS--MTSPQPKPKK-EALPR-----ASVTQIPS
 RTVPKQIVGGSTTLRP-----SSQLIP--IGSRLPKNVD-SPLNS-----MIQLPS
 25 TPYITPR-FTPSGTPKIFSTGGSPRGYS-AVASPNKTSGATSPTKSQYDGRGALSSWYPS
 TPYITPR-LTPSGTPKIISTASPKGYS-AAGSPQKTSGATSPTKPQYDGRASLSSWYPS
 TPYITPR-LTPTGTPKVIISTAGSPREYS-AAGSPKKTSGATSPTKLPYDGRSLSSWYSS
 TPYISPR-LTPTGTPKVIISTSVSPRRYS-AAGSPQKTSGITSPTK--HEGRGSLSSWYPS
 TSYISPR-LTPTATPKIISTSVSPRGYS-AAGSPKRTSGAVSPTK-----ATLWYPS
 30 TSYISPR-LTPTATPKIISTSVSPRGYS-AAGSPKRTSGAVSPTK-----ATLWYPS
 TPY-----
 T-HISPR-LTSDPAAKTRTAATSPRRYS-TAVSPKLAASATSP---R-----
 T-HISPR-LTSDPAAKTRTAATSPRRYS-TAVSPKLAASATSP---RLEGHMAMQPWLPS
 T-HITPR-LTSDPAKLRTAATSPRRYS-TAVSPKLASGATSP---RLEGHMAMSPWLPS
 35 SSHITPR-LTPEATPKIMSTSASPRRMS-TTATPKLMSGTTSPSKTRIEGYMSMTPWYPS
 SSHITPR-LTPEATPKIMSTSASPRRMS-TTATPKLMSGTTSPSKTRIEGYMSMTPWYPS
 SSHITPR-LTPEATPKIMSTSASPRRMS-TTATPKLMSGTTSPSKTRIEGYMSMTPWYPS
 SPHINQR-LTPEATPNIMSTSASPRGMS-TTATPKLVSGATSPSRTRIEGHMSMTPWY-S
 TTHITPR-LTPEATPKVMSTSASPRRMS-TTGTPKLMSGSTSPSKTRIESHMSMTPWYPS
 40 TTHITPR-LTPEATPKIMSTSASPRRMS-TTGTPKLMSGSTSPSKTRIEAHMSMTPWYPS
 SSHITPW-LTPKATPKIMSTSASPRQIS-TTATPKLMSGATSPTKSRMEAHMSMTPWYPS
 TPYTAPK-LTPG-----STPKFLSGPTSPTKSLSEVHSTFSSWHGS
 TPYTAPK-LTPG-----STPKFLSGPTSPTKSLSEVHSTFSSWHGS
 TPYVAPQ-ITPGSTPIISSSSGSPPLAYSTGPSTPKFYSGPTSPTRSRSEDQSAFSSWNGS
 45 GIPLTPHNNTPELTPKLTPNGSPKRLS-PPQSPRRHSASMSPTRHQFEGRLSSYSSLPA
 MMLTSQD-----SPRDSAKGEESGKAMEDDKWTDNYKSAGAISEKSQWEC
 GGAYGLRGMATAAATPAFPSTSGRPLYGASTPQAGQPPAASGPPYAAMTGMGMPAQAPPP
 xxxxxxx-xxxxxxxxxxxxxxxxspxxxx-xxxxpxxxxsxxxspxxxxxxxxxxxxxxxxxps

50

FIG. #4

10/13

5

10

15

SKPTSQSTTAPNSPPSHGSDPS-----RTPRVDGKDFFR----Q
SKPTSQSTTAPNSPPSHGSAPS-----RTPRVDGKDFFR----Q
SKPTSQSTTAPNSPPSSGSMPS-----RTPRVDGKDFFR----Q
20 SKATSQSTTAPNSPPSSGSMPS-----RTPRVDGKDFFR----Q
SQPTSRSS----SPPSSGSGQS-----RSTRLDGKEFFR----Q
SLPTSGSS----SPNSGHETLQGFHDNRSAFMNARVSVFIAARTARLDGKEFFR----Q
SQPTSGAS----SPPSHGSGQS-----RSTRLDGKEFFR----Q
S----QQSSAANSPPRARPLPG-----S
25 S----QQSSAANSPPRGHPIPG-----RAPRVDGKEFFR----Q
S----QQSSAANSPPWGRSLPV-----RTPKIDGKEFFR----Q
S----QQSSAANSPPRARPLPA-----RAPGIDGKEFFR----Q
S----QQSSAANSPPRNRTLPA-----RTPRMDGKEFFR----Q
S----QQSSAANSPPRNRTLPA-----RTPRMDGKEFFR----Q
30 -----RTPRMDGKEFFR----Q
-----RSTRLDGKEFFR----Q
S----KMSSAANSPPRAHSISG-----RSTRLDGKEFFR----Q
S----KMSSAANSPPRGQISG-----RSTRLDGKEFFR----Q
S----KQSSAANSPPRGRPNPG-----RTPRIDGKEFFR----Q
35 S----KQSSAANSPPRGRPNPG-----RTPRIDGKEFFR----Q
S----KQSSAANSPPRGRPNPG-----RTPRIDGKEFFR----Q
S----KQSSAANSPPRGRPNPG-----RTPRIDGKEFFR----Q
S----KQSSAANSPPRGRSNPG-----RTPRIDGKEFFR----Q
S----KQSSAANSPPRGRSNPG-----RTPRIDGKEFFR----Q
40 S----KQSSAANSPPRGRSNPG-----RTPRIDGKEFFR----Q
SS---HQYSAPTSPQHRSFAG-----R-PRIDGKEFFR----Q
SS---HQYSAPTSPQHRSFAG-----R-PRIDGKEFFR----Q
S----HQYSAPVSP-QRRSFAG-----R-PRIDGKEFFR----Q
S----HQATAPTSPPHSR--AQ-----AHVRIDGKEFFR----Q
45 P-----LTPRLPSDTGPAKS-----RRIDGKEVFRRASQIQ
S----TYATPHPHAQYGAAPG-----SPPRVDGKEFFR----Q
s----xxxxaxxspxxxxxxx-----rxxrxdgkeffr----q

50

FIG. #4

5

11/13

ARNRLSYEQFSAFLANIKELNAHRQTREETLRKAEEITFGPENKDLYAAFDGLLSRHLPS-
10 ARNRLSYEQFSAFLANIKELNAHRQTREETLRKAEEITFGPDNKDLYAAFDGLLSRHLPS-
ARNRLSYEQFSAFLANIKELNAHRQTREETLQKAQETFGPDNKDLYSAFEGLLSRHLPS-
ARNRLSYEQFSAFLANIKELNAHRQTREETLRKAQETFGPDNKDLYSAFEGLLSRHLPS-
ARARLSYEQFSSFLANIKELNAHRQTREETLANAEDIFGPENRDLYAAFEGLSRHLPS-
ARSRLSYEQFSSFLANIKELNAHRQTREETLAKAEDIFGPEHMDLHSAFEEVLSRHLSA-
15 ARSRLSYEQFSSFLANIKELNAHRQTREDTLGQAERIFGPENRDLSIAFEAILSRHLPS-
EKSRLSYEQFSAFLANIKELNAQKQSREETLRKAEEIFGTDNKDLYLSFQGLLRNRP--
ARSRLSYEQFSAFLANIKKLNQEQTREETLRKAEEIFGTDNKDLYFSFRGLLRNRIH--
ARSRLSYEQFSAFLANIKELNAQKQTREETLRKADEIFGSDNKDLYLSFQGLLRNAR--
ARSRLSLEQFSGFLANVKELNAQKQSREETLRKAEEIFGMDNKDLYISFQGLLRNRIH--
20 ARSRLSYEQFSSFLANIKELNAQKQTREETLRKADEIFGEENKDLYLSFQGLLRNMR--
ARSRLSYEQFSSFLANIKELNAQKQTREETLRKADEIFGEENKDLYLSFQGLLRNMR--

ARNRLSYEQFAAFLANIKELNAHRQSREETLQKADEIFGSENKDLFMSFQSLLSRSLSSG
ARNRLSYEQFAAFLANIKELNAHRQSREETLQKADEIFGSENKDLFMSFQSLLSRSLSS--
25 ARHRLSYEQFAAFLANIKELNAHRQSREETLRKADEIFGAENMDLFRSFQGLLRSLSS--
ARSRLSYEQFGAFLANIKELNAHKQSREDTLKKAAEEIFGPDNKDLYLSFQGLLRNRLP--
ARSRLSYEQFGAFLANIKELNAHKQSREDTLKKAAEEIFGPDNKDLYLSFQGLLRNRLP--
ARSRLSYEQFGAFLANIKELNAHKQSREDTLKKAAEEIFGPDNKDLYLSFQGLLRNRLP--
ARSRLSYEQFGAFLANIKELNAHKQSREETLKKAAEEIFGPDNKDLYLSFQGLLRNRLP--
30 ARSRLSYEQFGAFLANIKELNAHKQSREETLKKAAEEIFGPDNKDLYLSFQGLLRNRMVA-
ARSRLSYEQFGAFLANIKELNAHKQSREETLKKAAEEIFGPDNKDLYLSFQGLLRNRMVA-
ARSRLSYEQFGAFLANIKELNAHKQSREETLKKAAEEIFGPENKDLYLSFRGLLRNRSMP--
ARTRLSYEQFGAFLANIKEFNAQKQSREDTLSKAAEEIFGTEHKDLYISFQNLNRNHS--
ARTRLSYEQFGAFLANIKEFNAQKQSREDTLSKAAEEIFGTEHKDLYISFQNLNRNHS--
35 ARTRLSYEQFGAFLANIKEFNAQKQSREDTLSKAAEEIFGTEHKDLYISFQNLNRNQS--
ARNRLSFEQFSAFLANIKELNSHRQTREETLHKADEIFGSDNKDLYVIFDSLHRHLS--
SRARWTCEQFSDFLNNIREVNALRRAEEFLVCASSRAAYAYNVVPQCQLYGLNPPNWNCP
-----QARYLSVYKENLH-----
40 arxrlysyeqfxaflanikelnaxxqxrextlxkaxeifgxxnkdyxxfxllxxxxx-

45

50

FIG. #4

5

12/13

10

15

20

VKVYQMMDLSTTSVVAAKAYKYRAESLVKDYLLADCYVSYTAVLGGILMCKMVYDITHLI

25

30

35

PANCRVYTR

40

45

50

FIG. #4

5

13/13

10

15

20

25

30

35

40

45

50

SSLYYKGYGSLTKIQKLEWNNRGMSTVHAMFITLMSVYLVFFSNLFSDELDGPVTVRSSN