

(12) STANDARD PATENT

(11) Application No. AU 2005309820 B2

(19) AUSTRALIAN PATENT OFFICE

(54) Title
2-methylene-19,21-dinor-1alpha-hydroxy-bishomopregnacalciferol

(51) International Patent Classification(s)
C07C 401/00 (2006.01) **A61K 31/59** (2006.01)

(21) Application No: **2005309820** (22) Date of Filing: **2005.11.18**

(87) WIPO No: **WO06/057915**

(30) Priority Data

(31) Number
60/629,958 (32) Date
2004.11.22 (33) Country
US

(43) Publication Date: **2006.06.01**

(44) Accepted Journal Date: **2011.06.23**

(71) Applicant(s)
Wisconsin Alumni Research Foundation

(72) Inventor(s)
Deluca, Hector F.;Plum, Lori A.;Clagett-Dame, Margaret

(74) Agent / Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(56) Related Art

WO 2003/075932 A

WO 1998/041501 A

US 5843928 A

US 6566352 B1

WO 2005/051323 A = US 2005/0119242 A1

EP 0503923 B1

WO 2002/020021 A

PLUM LORI A et al. Proceedings of the National Academy of Sciences of USA, (2004), vol. 101 (18), pp. 6900-6904.

WO 2003/051828 A

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 June 2006 (01.06.2006)

PCT

(10) International Publication Number
WO 2006/057915 A3

(51) International Patent Classification:
A61K 31/59 (2006.01) *C07C 401/00* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CI, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2005/041888

(22) International Filing Date:
18 November 2005 (18.11.2005)

(25) Filing Language:
English

(26) Publication Language:
English

(30) Priority Data:
60/629,958 22 November 2004 (22.11.2004) US

(71) Applicant (for all designated States except US): **WISCONSIN ALUMNI RESEARCH FOUNDATION** [US/US]; Post Office Box 7365, Madison, Wisconsin 53707-7365 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments


(75) Inventors/Applicants (for US only): **DELUCA, Hector F.** [US/US]; 1809 Highway BB, Deerfield, Wisconsin 53531 (US). **PLUM, Lori A.** [US/US]; 6139 Highway H, Arena, Wisconsin 53503 (US). **CLAGETT-DAME, Margaret** [US/US]; 1809 Highway BB, Deerfield, Wisconsin 53531 (US).

(88) Date of publication of the international search report:
13 July 2006

(74) Agents: **MEARA, Joseph P.** et al.; **FOLEY & LARDNER LLP**, 150 East Gilman Street, Post Office Box 1497, Madison, Wisconsin 53701-1497 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 2-METHYLENE-19,21-DINOR-1ALPHA-HYDROXY-BISHOMOPREGNACALCIFEROL



(57) Abstract: Compounds of formula (I) are provided where X¹ and X² are independently selected from H or hydroxy protecting groups. Such compounds may be used in preparing pharmaceutical compositions and are useful in treating a variety of biological conditions.

WO 2006/057915 A3

2-METHYLENE-19,21-DINOR-1 α -HYDROXY-BISHOMOPREGNACALCIFEROL


FIELD OF THE INVENTION

[0001] This invention relates to vitamin D compounds, and more particularly to 2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol and to pharmaceutical formulations that include this compound. The invention also relates to the use of 2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol or salts thereof in the preparation of medicaments for use in treating various diseases.

BACKGROUND OF THE INVENTION

[0002] The natural hormone, 1 α ,25-dihydroxyvitamin D₃ (also referred to as 1 α ,25-dihydroxycholecalciferol and calcitriol) and its analog in the ergosterol series, i.e. 1 α ,25-dihydroxyvitamin D₂, are known to be highly potent regulators of calcium homeostasis in animals and humans, and their activity in cellular differentiation has also been established, Ostrem *et al.*, *Proc. Natl. Acad. Sci. USA*, 84, 2610 (1987). Many structural analogs of these metabolites have been prepared and tested, including 1 α -hydroxyvitamin D₃, 1 α -hydroxyvitamin D₂, various side chain homologated vitamins, and fluorinated analogs. Some of these compounds exhibit an interesting separation of activities in cell differentiation and calcium regulation. This difference in activity may be useful in the treatment of a variety of diseases as renal osteodystrophy, vitamin D-resistant rickets, osteoporosis, psoriasis, and certain malignancies. The structure of 1 α ,25-dihydroxyvitamin D₃ and the numbering system used to denote the carbon atoms in this compound are shown below.

Atty. Docket No. 032026-0906



$1\alpha,25$ -Dihydroxyvitamin D₃ = $1\alpha,25$ -Dihydroxycholecalciferol = Calcitriol

[0003] Another class of vitamin D analogs, i.e. the so called 19-nor-vitamin D compounds, is characterized by the replacement of the A-ring exocyclic methylene group (carbon 19), typical of the vitamin D system, by two hydrogen atoms. Biological testing of such 19-nor-analogs (e.g., $1\alpha,25$ -dihydroxy-19-nor-vitamin D₃) revealed a selective activity profile with high potency in inducing cellular differentiation, and very low calcium mobilizing activity. Thus, these compounds are potentially useful as therapeutic agents for the treatment of malignancies, or the treatment of various skin disorders. Two different methods of synthesis of such 19-nor-vitamin D analogs have been described (Perlman *et al.*, *Tetrahedron Lett.* 31, 1823 (1990); Perlman *et al.*, *Tetrahedron Lett.* 32, 7663 (1991), and DeLuca *et al.*, U.S. Pat. No. 5,086,191).

[0004] In U.S. Pat. No. 4,666,634, 2β -hydroxy and alkoxy (e.g., ED-71) analogs of $1\alpha,25$ -dihydroxyvitamin D₃ have been described and examined by the Chugai group as potential drugs for osteoporosis and as antitumor agents. See also Okano *et al.*, *Biochem. Biophys. Res. Commun.* 163, 1444 (1989). Other 2-substituted (with hydroxyalkyl, e.g., ED-120, and fluoroalkyl groups) A-ring analogs of $1\alpha,25$ -dihydroxyvitamin D₃ have also been prepared and tested (Miyamoto *et al.*, *Chem. Pharm. Bull.* 41, 1111 (1993); Nishii *et al.*, *Osteoporosis*

Atty. Docket No. 032026-0906

Int. Suppl. 1, 190 (1993); Posner *et al.*, *J. Org. Chem.* 59, 7855 (1994), and *J. Org. Chem.* 60, 4617 (1995)).

[0005] Various 2-substituted analogs of 1α ,25-dihydroxy-19-nor-vitamin D₃ have also been synthesized, i.e. compounds substituted at the 2-position with hydroxy or alkoxy groups (DeLuca *et al.*, U.S. Pat. No. 5,536,713), with 2-alkyl groups (DeLuca *et al.*, U.S. Patent No. 5,945,410), and with 2-alkylidene groups (DeLuca *et al.*, U.S. Patent No. 5,843,928), which exhibit interesting and selective activity profiles. All these studies indicate that binding sites in vitamin D receptors can accommodate different substituents at C-2 in the synthesized vitamin D analogs.

[0006] In a continuing effort to explore the 19-nor class of pharmacologically important vitamin D compounds, analogs which are characterized by the presence of a methylene substituent at carbon 2 (C-2), a hydroxyl group at carbon 1 (C-1), and a shortened side chain attached to carbon 20 (C-20) have also been synthesized and tested. 1α -Hydroxy-2-methylene-19-nor-pregnacalciferol is described in U.S. Patent No. 6,566,352 while 1α -hydroxy-2-methylene-19-nor-(20S)-homopregnacalciferol is described in U.S. Patent No. 6,579,861 and 1α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol is described in U.S. Patent No. 6,627,622. All three of these compounds have relatively high binding activity to vitamin D receptors and relatively high cell differentiation activity, but little if any calcemic activity as compared to 1α ,25-dihydroxyvitamin D₃. Their biological activities make these compounds excellent candidates for a variety of pharmaceutical uses, as set forth in the '352, '861 and '622 patents.


SUMMARY OF THE INVENTION

[0007] The invention provides 2-methylene-19,21-dinor- 1α -hydroxy-bishomopregnacalciferol and related compounds, pharmaceutical formulations that include 2-methylene-19,21-dinor- 1α -hydroxy-bishomopregnacalciferol, and

Atty. Docket No. 032026-0906


the use of this compound in the preparation of medicaments for use in treating various disease states.

[0008] Therefore, in one aspect, the invention provides a compound having the formula I shown below




where X^1 and X^2 may be the same or different and are independently selected from H or hydroxy-protecting groups. In some embodiments, X^1 and X^2 are both hydroxy protecting groups such as silyl groups. In some such embodiments, X^1 and X^2 are both t-butyldimethylsilyl groups. In other embodiments, X^1 and X^2 are both H such that the compound is 2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol having the formula IA as shown below:

Atty. Docket No. 032026-0906



IA

In some such embodiments, the compound of formula IA is a compound of formula IB and has the structure shown below:



IB

[0009] The above compound exhibits a desired, and highly advantageous, pattern of biological activity. This compound is characterized by relatively high binding to vitamin D receptors, but very low intestinal calcium transport activity, as compared to that of 1α ,25-dihydroxyvitamin D₃, and has very low ability to

Atty. Docket No. 032026-0906

mobilize calcium from bone, as compared to $1\alpha,25$ -dihydroxyvitamin D₃. Hence, this compound can be characterized as having little, if any, calcemic activity. Thus, it may be useful as a therapy for suppression of secondary hyperparathyroidism of renal osteodystrophy.

[0010] The compound of the invention is also especially suited for treatment and prophylaxis of human disorders which are characterized by an imbalance in the immune system, e.g. in autoimmune diseases, including multiple sclerosis, lupus, diabetes mellitus, host versus graft reaction, and rejection of organ transplants; and additionally for the treatment of inflammatory diseases, such as rheumatoid arthritis, asthma, and inflammatory bowel diseases such as celiac disease, ulcerative colitis and Crohn's disease. Acne, alopecia and hypertension are other conditions which may be treated with the compound of the invention.

[0011] The above compound is also characterized by relatively high cell differentiation activity. Thus, this compound also provides a therapeutic agent for the treatment of psoriasis, or as an anti-cancer agent, especially against leukemia, colon cancer, breast cancer and prostate cancer. In addition, due to its relatively high cell differentiation activity, this compound provides a therapeutic agent for the treatment of various skin conditions including wrinkles, lack of adequate dermal hydration, i.e. dry skin, lack of adequate skin firmness, i.e. slack skin, and insufficient sebum secretion. Use of this compound thus not only results in moisturizing of skin but also improves the barrier function of skin.

[0012] The compounds of the invention may be used to prepare pharmaceutical formulations or medicaments that include a compound of the invention in combination with a pharmaceutically acceptable carrier. Such pharmaceutical formulations and medicaments may be used to treat various biological disorders such as those described herein. Methods for treating such disorders typically include administering an effective amount of the compound or an appropriate amount of a pharmaceutical formulation or a medicament that includes the compound to a subject suffering from the biological disorder. In

Atty. Docket No. 032026-0906

some embodiments, the subject is a mammal. In some such embodiments, the mammal is selected from a rodent, a primate, a bovine, an equine, a canine, a feline, an ursine, a porcine, a rabbit, or a guinea pig. In some such embodiments, the mammal is a rat or is a mouse. In some embodiments, the subject is a primate such as, in some embodiments, a human.

[0013] The compound may be present in a composition to treat the above-noted diseases and disorders in an amount from about 0.01 μ g/gm to about 1 mg/gm of the composition, preferably from about 0.1 μ g/gm to about 500 μ g/gm of the composition, and may be administered topically, transdermally, orally, or parenterally in dosages of from about 0.01 μ g/day to about 1 mg/day, preferably from about 0.1 μ g/day to about 500 μ g/day.

[0014] Further objects, features and advantages of the invention will be apparent from the following detailed description and drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] Figures 1-7 illustrate various biological activities of 2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol (referred to as "19,21-dinor" in the Figures) compared with those of the native hormone 1 α ,25-dihydroxyvitamin D₃ (referred to as "1,25(OH)₂D₃" in the Figures).

[0016] Figure 1 is a graph comparing the relative activity of 19,21-dinor and 1,25(OH)₂D₃ to compete for binding with [³H]-1,25-(OH)₂-D₃ to the full-length recombinant rat vitamin D receptor.

[0017] Figure 2 is a graph comparing the percent HL-60 cell differentiation as a function of the concentration of 19,21-dinor with that of 1,25(OH)₂D₃.

[0018] Figure 3 is a graph comparing the *in vitro* transcription activity of 19,21-dinor with that of 1,25(OH)₂D₃.

[0019] Figure 4 is a bar graph comparing the bone calcium mobilization activity of 19,21-dinor with that of 1,25(OH)₂D₃.

Atty. Docket No. 032026-0906


[0020] Figure 5 is a bar graph comparing the intestinal calcium transport activity of 19,21-dinor with that of 1,25(OH)₂D₃.

[0021] Figure 6 is a bar graph comparing the serum calcium levels in adult rats after administration of 19,21-dinor and 1,25(OH)₂D₃.

[0022] Figure 7 is a bar graph comparing the percent parathyroid hormone (PTH) suppression in adult rats of 19,21-dinor with 1,25(OH)₂D₃.

DETAILED DESCRIPTION OF THE INVENTION


[0023] 2-Methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol was synthesized, and tested, and found to be useful in treating a variety of biological conditions as described herein. Structurally, this compound has the formula IA as shown below:



IA

[0024] Preparation of 2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol can be accomplished by condensing an appropriate bicyclic Windaus-Grundmann type ketone (II) with the allylic phosphine oxide III followed by deprotection (removal of the Y₁ and Y₂ groups).

Atty. Docket No. 032026-0906




[0025] In phosphine oxide III, Y₁ and Y₂ are preferably hydroxy-protecting groups such as silyl protecting groups. The t-butyldimethylsilyl (TBDMS) group is an example of a particularly useful hydroxy-protecting group. The process described above represents an application of the convergent synthesis concept, which has been applied effectively to the preparation of numerous vitamin D compounds (see Lythgoe *et al.*, *J. Chem. Soc. Perkin Trans. I*, 590 (1978); Lythgoe, *Chem. Soc. Rev.* 9, 449 (1983); Toh *et al.*, *J. Org. Chem.* 48, 1414 (1983); Baggioini *et al.*, *J. Org. Chem.* 51, 3098 (1986); Sardina *et al.*, *J. Org. Chem.* 51, 1264 (1986); *J. Org. Chem.* 51, 1269 (1986); DeLuca *et al.*, U.S. Patent No. 5,086,191; DeLuca *et al.*, U.S. Patent No. 5,536,713; and DeLuca *et al.*, U.S. Patent No. 5,843,928 all of which are hereby incorporated by reference in their entirety and for all purposes as if fully set forth herein).

[0026] Phosphine oxide III is a convenient reagent that can be used to prepare a large number of 19-nor vitamin D compounds and may be prepared according to the procedures described by Sicinski *et al.*, *J. Med. Chem.*, 41, 4662 (1998), DeLuca *et al.*, U.S. Patent No. 5,843,928; Perlman *et al.*, *Tetrahedron Lett.* 32, 7663 (1991); and DeLuca *et al.*, U.S. Patent No. 5,086,191. Scheme I shows the general procedure for synthesizing phosphine oxide III as outlined in U.S. Patent No. 5,843,928 which is hereby incorporated by reference in its entirety as if fully set forth herein. Modification of the method shown in Scheme I may be used to produce a large number of vitamin D analogs as will be apparent to those skilled in the art. For example, a wide variety of

Atty. Docket No. 032026-0906

phosphonium compounds may be used in place of the $\text{MePh}_3\text{P}^+ \text{Br}^-$ used to convert ketone B to alkene C. Examples of such compounds include EtPh_3P^+ Br^- , $\text{PrPh}_3\text{P}^+ \text{Br}^-$, and compounds generally prepared by reaction of triphenylphosphine with an alkyl halide, an alkenyl halide, a protected-hydroxyalkyl halide, and a protected hydroxyalkenyl halide. Alkenes prepared using this procedure may then be carried through to prepare a phosphine oxide in an analogous manner to that used to prepare phosphine oxide H in Scheme I. Alternatively, an alkene analogous to compound C of Scheme I may be reduced with $(\text{Ph}_3\text{P})_3\text{RhCl}$ and H_2 to provide other vitamin D analogs. See U.S. Patent No. 5,945,410 and Sicinski, R. R. *et al.*, *J. Med. Chem.*, 41, 4662-4674 (1998) both of which are hereby incorporated by reference in their entireties and for all purposes. Therefore, the procedure for forming the phosphine oxide shown in Scheme I may be used to prepare a wide variety of vitamin D analogs in addition to the compound of the present invention.

Atty. Docket No. 032026-0906

Scheme I

[0027] Hydriindanones of structure II can be prepared by known methods or adapted methods as will be readily apparent to one of skill in the art and described herein. Specific examples of some important bicyclic ketones used to synthesize vitamin D analogs are those described in Mincione *et al.*, *Synth. Commun.* 19, 723, (1989); and Peterson *et al.*, *J. Org. Chem.* 51, 1948, (1986).

[0028] An overall process for synthesizing 2-alkylidene-19-nor-vitamin D compounds is illustrated and described in U.S. Patent No. 5,843,928 which is

Atty. Docket No. 032026-0906

hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein.

[0029] As used herein, the term "hydroxy-protecting group" signifies any group commonly used for the temporary protection of the hydroxy (-OH) functional group, such as, but not limited to, alkoxy carbonyl, acyl, alkylsilyl or alkylarylsilyl groups (hereinafter referred to simply as "silyl" groups), and alkoxyalkyl groups. Alkoxy carbonyl protecting groups are alkyl-O-CO- groups such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, tert-butoxycarbonyl, benzyloxycarbonyl or allyloxycarbonyl. The term "acyl" signifies an alkanoyl group of 1 to 6 carbons, in all of its isomeric forms, or a carboxyalkanoyl group of 1 to 6 carbons, such as an oxanyl, malonyl, succinyl, glutaryl group, or an aromatic acyl group such as benzoyl, or a halo, nitro or alkyl substituted benzoyl group. Alkoxyalkyl protecting groups are groups such as methoxymethyl, ethoxymethyl, methoxyethoxymethyl, or tetrahydrofuranyl and tetrahydropyranyl. Preferred silyl-protecting groups are trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, dibutylmethylsilyl, diphenylmethylsilyl, phenyldimethylsilyl, diphenyl-t-butyldimethylsilyl and analogous alkylated silyl radicals. The term "aryl" specifies a phenyl-, or an alkyl-, nitro- or halo-substituted phenyl group. An extensive list of protecting groups for the hydroxy functionality may be found in Protective Groups in Organic Synthesis, Greene, T.W.; Wuts, P. G. M., John Wiley & Sons, New York, NY, (3rd Edition, 1999) which can be added or removed using the procedures set forth therein and which is hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein.

[0030] A "protected hydroxy" group is a hydroxy group derivatized or protected by any of the above groups commonly used for the temporary or permanent protection of hydroxy functional groups, e.g., the silyl, alkoxyalkyl, acyl or alkoxy carbonyl groups, as previously defined.

Atty. Docket No. 032026-0906

EXAMPLES

Synthesis of 2-Methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol


[0031] The synthesis and characteristics of various 19-nor vitamin D analogs is described in numerous United States patents including U.S. Patent No. 5,843,928, U.S. Patent No. 6,627,622, U.S. Patent No. 6,579,861, U.S. Patent No. 5,086,191, U.S. Patent No. 5,585,369, and U.S. Patent No. 6,537,981. Each of the above-described references is hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein.

[0032] Compounds of formula I, formula IA, and formula IB were prepared using the methods shown in Schemes I and II. The starting material, compound 1, was prepared using the procedure set forth by Andrzej R. Daniewski and Wen Liu (*J. Org. Chem.* 66, 626-628 (2001) using $\text{Ph}_3\text{P}^+\text{Pr Br}^-$ in place of $\text{Ph}_3\text{P}^+\text{Et Br}^-$ in the step for converting compound 7 to compound 8 in Scheme I of the article which is hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein. Starting material 1 was then hydrogenated using palladium on carbon as a catalyst to yield the saturated compound 2 having a molecular weight of 196. The reaction was followed using thin layer chromatography (TLC) using a 20% ethyl acetate in a hexane solvent system. Compound 2 was then oxidized with pyridinium chlorochromate as described in U.S. Patent Publication No. 2004/0220418, published on November 4, 2004 (U.S. Patent Application No. 10/847,040), hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein, to yield 3 as followed by TLC using the same solvent system. The Ring-A phosphine oxide compound 4 was synthesized as shown in Scheme I. The condensation was carried out again as described in the above-referenced patent documents with n-butyl lithium and was followed by TLC using a solvent of 5% ethyl acetate in hexane to yield compound 5. The t-butyldimethylsilyl protecting groups were removed using tetrabutylammonium fluoride as described in U.S. Patent Publication No. 2004/0220418 to yield compound 6 (2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol).

Atty. Docket No. 032026-0906

TLC was used to follow completion of the reaction using the solvent system described above. The final product was fully characterized as described below.

Scheme II



2-Methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol

[0033] UV (in EtOH) λ_{max} 244, 252, 262 nm; ^1H NMR (CDCl_3), 0.451 (3H, s, 18- H_3), 0.898 (3H, t, J = 6.8 Hz, 23- H_3), 1.8-2.05 (2H, m), 2.29 (1H, dd, J =

Atty. Docket No. 032026-0906

13.2, 8.7 Hz, 10 α -H), 2.33 (1H, dd, J = 13.5, 5.7 Hz, 4 β -H), 2.58 (1H, dd, J = 13.5, 3.8 Hz, 4 α -H), ca. 2.84 (1H, overlapped with 10 β -H, 9 β -H), 2.86 (1H, dd, J = 13.2, 4.5 Hz, 10 β -H), 4.49 (2H, m, 1 β - and 3 α -H), 5.10 and 5.11 (1H and 1H, each s, =CH₂), 5.89 and 6.37 (1H and 1H, each d, J = 11.4 Hz, 7- and 6-H); MS (APCI) m/z (relative intensity) 331 ([M + H]⁺, 7), 313 ([M + H]⁺ - H₂O, 100), 295 ([M + H]⁺ - 2•H₂O, 92).

BIOLOGICAL ACTIVITY

Vitamin D Receptor Binding

Test Material

Protein Source

[0034] Full-length recombinant rat receptor was expressed in *E. coli* BL21(DE3) Codon Plus RIL cells and purified to homogeneity using two different column chromatography systems. The first system was a nickel affinity resin that utilizes the C-terminal histidine tag on this protein. The protein that was eluted from this resin was further purified using ion exchange chromatography (S-Sepharose Fast Flow). Aliquots of the purified protein were quick frozen in liquid nitrogen and stored at -80°C until use. For use in binding assays, the protein was diluted in TEDK₅₀ (50 mM Tris, 1.5 mM EDTA, pH 7.4, 5 mM DTT, 150 mM KCl) with 0.1% Chaps detergent. The receptor protein and ligand concentration was optimized such that no more than 20% of the added radiolabeled ligand was bound to the receptor.

Study Drugs

[0035] Unlabeled ligands were dissolved in ethanol and the concentrations determined using UV spectrophotometry (1,25(OH)₂D₃: molar extinction coefficient = 18,200 and λ_{max} = 265 nm; Analogs: molar extinction coefficient = 42,000 and λ_{max} = 252 nm). Radiolabeled ligand (³H-1,25(OH)₂D₃, ~159 Ci/mmole) was added in ethanol at a final concentration of 1 nM.

Atty. Docket No. 032026-0906

Assay Conditions

[0036] Radiolabeled and unlabeled ligands were added to 100 mcL of the diluted protein at a final ethanol concentration of $\leq 10\%$, mixed and incubated overnight on ice to reach binding equilibrium. The following day, 100 mcL of hydroxylapatite slurry (50%) was added to each tube and mixed at 10-minute intervals for 30 minutes. The hydroxylapatite was collected by centrifugation and then washed three times with Tris-EDTA buffer (50 mM Tris, 1.5 mM EDTA, pH 7.4) containing 0.5% Triton X-100. After the final wash, the pellets were transferred to scintillation vials containing 4 mL of Biosafe II scintillation cocktail, mixed and placed in a scintillation counter. Total binding was determined from the tubes containing only radiolabeled ligand.

HL-60 Differentiation

Test Material

Study Drugs

[0037] The study drugs were dissolved in ethanol and the concentrations determined using UV spectrophotometry. Serial dilutions were prepared so that a range of drug concentrations could be tested without changing the final concentration of ethanol ($\leq 0.2\%$) present in the cell cultures.

Cells

[0038] Human promyelocytic leukemia (HL60) cells were grown in RPMI-1640 medium containing 10% fetal bovine serum. The cells were incubated at 37°C in the presence of 5% CO₂.

Assay Conditions

[0039] HL60 cells were plated at 1.2×10^5 cells/mL. Eighteen hours after plating, cells in duplicate were treated with drug. Four days later, the cells were harvested and a nitro blue tetrazolium reduction assay was performed (Collins *et al.*, 1979; *J. Exp. Med.* 149:969-974). The percentage of differentiated cells was determined by counting a total of 200 cells and recording the number that

Atty. Docket No. 032026-0906

contained intracellular black-blue formazan deposits. Verification of differentiation to monocytic cells was determined by measuring phagocytic activity (data not shown).

In Vitro Transcription Assay

[0040] Transcription activity was measured in ROS 17/2.8 (bone) cells that were stably transfected with a 24-hydroxylase (24Ohase) gene promoter upstream of a luciferase reporter gene (Arbour *et al.*, 1998). Cells were given a range of doses. Sixteen hours after dosing the cells were harvested and luciferase activities were measured using a luminometer. RLU = relative luciferase units.

Intestinal Calcium Transport and Bone Calcium Mobilization

[0041] Male, weanling Sprague-Dawley rats were placed on Diet 11 (0.47% Ca) diet +AEK for one week followed by Diet 11 (0.02% Ca) +AEK for 3 weeks. The rats were then switched to a diet containing 0.47% Ca for one week followed by two weeks on a diet containing 0.02% Ca. Dose administration began during the last week on 0.02% calcium diet. Four consecutive ip doses were given approximately 24 hours apart. Twenty-four hours after the last dose, blood was collected from the severed neck and the concentration of serum calcium determined as a measure of bone calcium mobilization. The first 10 cm of the intestine was also collected for intestinal calcium transport analysis using the everted gut sac method.

PTH Suppression and Hypercalcemia

Species

[0042] Adult, female Sprague-Dawley rats were obtained from Harlan (Madison, WI).

Animal Husbandry

Atty. Docket No. 032026-0906

[0043] Upon receipt, the animals were identified by individual tail marks. Animals were housed in suspended, stainless steel, wire-bottom cages. Each cage contained one animal. The animal rooms were maintained at a temperature of 68 to 72°F and a relative humidity of 25 to 75%. The holding rooms were set to provide 12 hours of light per day. Water and a purified rodent diet (Suda *et al.*, Purified Rodent Diet-Diet 11) containing 0.47% and 0.3% phosphorus and fat soluble vitamins A, D, E and K were provided *ad libitum*.

Treatment Groups

[0044] Animals were randomly assigned to treatment groups (5 animals/group). All doses were administered intraperitoneally in 100 microliters of propylene glycol. Four to seven consecutive doses were given approximately 24 hours apart. Dosing was initiated after the animals had been allowed to acclimate for at least one week.

Dose Preparation

Control Material

A. Negative Control Material

[0045] The negative control material was prepared by volumetrically measuring ethanol (< 5%) and propylene glycol, mixing, and then placing in storage at 2 to 8°C.

B. Positive Control Material

[0046] 1,25(OH)₂D₃ was prepared by determining the concentration of an ethanol stock solution using UV spectrophotometry (extinction coefficient = 18,200; λ_{max} = 265 nm). The required amount of 1,25(OH)₂D₃ was volumetrically measured into propylene glycol so that there was less than 5% ethanol in the final solution. The solution was mixed and then stored at 2 to 8°C.

Test Material

[0047] The analogs were prepared by first determining the concentration of an ethanol stock solution using UV spectrophotometry (extinction coefficient =

Atty. Docket No. 032026-0906

42,000; $\lambda_{\text{max}} = 252$ nm). The analog solutions were then volumetrically added to propylene glycol so that there was less than 5% ethanol in the final solution. The solution was mixed and stored at 2 to 8°C.

Dose Administration Method

[0048] Both control and test articles were administered by intraperitoneal injection in 100 microliters for 4-7 consecutive days spaced approximately 24 hours apart. $1,25(\text{OH})_2\text{D}_3$ was given for 4 consecutive days, whereas, the test drugs were given for 7 consecutive days.

Serum PTH Levels

[0049] Twenty-four hours after the final dose, blood was collected from the tail artery and the concentration of bioactive serum PTH was measured using the rat BioActive Intact PTH ELISA Kit from Immutopics, Inc. (San Clemente, CA).

Serum Calcium Analysis

[0050] Twenty-four hours after the final dose, approximately 1 ml of blood was collected from the tail artery of each experimental animal. The blood was allowed to coagulate at room temperature and then centrifuged at 3000 x g for 15 minutes. The serum was transferred to a polypropylene tube and stored frozen at -20°C. The level of calcium was determined by diluting the serum into 0.1% lanthum chloride and measuring the absorbance on an atomic absorption spectrophotometer (Perkin Elmer Model 3110, Shelton, CT).

[0051] 2-Methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol binds to the recombinant vitamin D receptor, but is about 5 times less active than is 1 α ,25-dihydroxyvitamin D₃ in this respect (see Figure 1). Furthermore, it is active in stimulating transcription of a reporter gene stably transfected in Ros17/2.8 (bone) cells, indicating significant biological activity (see Figure 3). However, it is about 15 times less active than 1 α ,25-dihydroxyvitamin D₃ in inducing differentiation of HL-60 cells (see Figure 2). It has no calcemic activity when measured either by intestinal calcium transport or bone calcium mobilization even when given at 100 times the dose of 1 α ,25-dihydroxyvitamin

Atty. Docket No. 032026-0906

D₃ (see Figures 4 and 5). However, it does possess significant activity in suppressing parathyroid hormone levels in normal rats (see Figure 7). This compound should find its uses in the treatment of autoimmune diseases such as multiple sclerosis, type I diabetes, rheumatoid arthritis, lupus, and other similar degenerative diseases. It should also have significant activity in treating malignant growth such as colorectal, breast and prostate cancers. All of these activities should be evident in the absence of raising serum calcium concentrations (see Figure 6). This compound should also be useful in treating secondary hyperparathyroidism found in patients who have lost kidney function such as those on hemodialysis or peritoneal dialysis.

[0052] The compounds of the invention are also useful in preventing or treating obesity, inhibiting adipocyte differentiations, inhibiting SCD-1 gene transcription, and/or reducing body fat in animal subjects. Therefore, in some embodiments, a method of preventing or treating obesity, inhibiting adipocyte differentiations, inhibiting SCD-1 gene transcription, and or reducing body fat in animal subject includes administering to the animal subject, an effective amount of the compound or a pharmaceutical composition that includes the compound. Administration of the compound or the pharmaceutical composition to the subject inhibits adipocyte differentiation, inhibits gene transcription, and/or reduces body fat in the animal subject.

[0053] For treatment purposes, the compounds defined by formula I, formula IA, and formula IB may be formulated for pharmaceutical applications as a solution in innocuous solvents, or as an emulsion, suspension or dispersion in suitable solvents or carriers, or as pills, tablets or capsules, together with solid carriers, according to conventional methods known in the art. Any such formulations may also contain other pharmaceutically acceptable and non-toxic excipients such as stabilizers, anti-oxidants, binders, coloring agents or emulsifying or taste-modifying agents. Pharmaceutically acceptable excipients and carriers are generally known to those skilled in the art and are thus included in the instant invention. Such excipients and carriers are described, for example,

Atty. Docket No. 032026-0906

in "Remingtons Pharmaceutical Sciences" Mack Pub. Co., New Jersey (1991), which is hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein.

[0054] The compounds may be administered orally, topically, parenterally, or transdermally. The compounds are advantageously administered by injection or by intravenous infusion or suitable sterile solutions, or in the form of liquid or solid doses via the alimentary canal, or in the form of creams, ointments, patches, or similar vehicles suitable for transdermal applications. In some embodiments, doses of from 0.001 μ g to about 1 mg per day of the compound are appropriate for treatment purposes. In some such embodiments an appropriate and effective dose may range from 0.01 μ g to 1 mg per day of the compound. In other such embodiments an appropriate and effective dose may range from 0.1 μ g to 500 μ g per day of the compound. Such doses will be adjusted according to the type of disease or condition to be treated, the severity of the disease or condition, and the response of the subject as is well understood in the art. The compound may be suitably administered alone, or together with another active vitamin D compound.

[0055] Compositions for use in the invention include an effective amount of 2-methylene-19,21-dinor-1 α -hydroxy-bishomopregnacalciferol as the active ingredient, and a suitable carrier. An effective amount of the compound for use in accordance with some embodiments of the invention will generally be a dosage amount such as those described herein, and may be administered topically, transdermally, orally, nasally, rectally, or parenterally.

[0056] The compounds of formula IA and IB, may be advantageously administered in amounts sufficient to effect the differentiation of promyelocytes to normal macrophages. Dosages as described above are suitable, it being understood that the amounts given are to be adjusted in accordance with the severity of the disease, and the condition and response of the subject as is well understood in the art.

Atty. Docket No. 032026-0906

[0057] The compound may be formulated as creams, lotions, ointments, aerosols, suppositories, topical patches, pills, capsules or tablets, or in liquid form as solutions, emulsions, dispersions, or suspensions in pharmaceutically innocuous and acceptable solvent or oils, and such preparations may contain, in addition, other pharmaceutically innocuous or beneficial components, such as stabilizers, antioxidants, emulsifiers, coloring agents, binders or taste-modifying agents.

[0058] The formulations of the present invention comprise an active ingredient in association with a pharmaceutically acceptable carrier therefore and optionally other therapeutic ingredients. The carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulations and not deleterious to the recipient thereof.

[0059] Formulations of the present invention suitable for oral administration may be in the form of discrete units as capsules, sachets, tablets or lozenges, each containing a predetermined amount of the active ingredient; in the form of a powder or granules; in the form of a solution or a suspension in an aqueous liquid or non-aqueous liquid; or in the form of an oil-in-water emulsion or a water-in-oil emulsion.

[0060] Formulations for rectal administration may be in the form of a suppository incorporating the active ingredient and carrier such as cocoa butter, or in the form of an enema.

[0061] Formulations suitable for parenteral administration conveniently comprise a sterile oily or aqueous preparation of the active ingredient which is preferably isotonic with the blood of the recipient.

[0062] Formulations suitable for topical administration include liquid or semi-liquid preparations such as liniments, lotions, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops; or as sprays.

Atty. Docket No. 032026-0906

[0063] For nasal administration, inhalation of powder, self-propelling or spray formulations, dispensed with a spray can, a nebulizer or an atomizer can be used. The formulations, when dispensed, preferably have a particle size in the range of 10 to 100 microns.

[0064] The formulations may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. By the term "dosage unit" is meant a unitary, i.e., a single dose which is capable of being administered to a patient as a physically and chemically stable unit dose comprising either the active ingredient as such or a mixture of it with solid or liquid pharmaceutical diluents or carriers.


[0065] All references cited herein are specifically incorporated by reference in their entireties and for all purposes as if fully set forth herein.

[0066] It is understood that the invention is not limited to the embodiments set forth herein for illustration, but embraces all such forms thereof as come within the scope of the following claims.

CLAIMS

What is claimed is:

1. A compound have the formula I




wherein X^1 and X^2 are independently selected from H and hydroxy protecting groups.

2. The compound of claim 1, wherein X^1 and X^2 are both hydroxy protecting groups.

3. The compound of claim 2, wherein X^1 and X^2 are both t-butyldimethylsilyl groups.

Atty. Docket No. 032026-0906

4. The compound of claim 1, wherein X^1 and X^2 are both H and the compound has the formula IA



IA

5. A pharmaceutical composition, comprising an effective amount of the compound of claim 4 and a pharmaceutically acceptable carrier.

6. The pharmaceutical composition of claim 5 wherein the effective amount comprises from about 0.01 μ g to about 1 mg of the compound per gram of the composition.

7. The pharmaceutical composition of claim 5 wherein the effective amount comprises from about 0.1 μ g to about 500 μ g of the compound per gram of the composition.

8. A method of treating a subject suffering from a biological condition, comprising administering an effective amount of the compound of claim 4 to the subject, wherein the biological condition is selected from psoriasis; leukemia; colon cancer; breast cancer; prostate cancer; multiple sclerosis; lupus; diabetes mellitus; host versus graft reaction; rejection of organ transplants; an inflammatory disease selected from rheumatoid arthritis, asthma, or

inflammatory bowel diseases; a skin condition selected from wrinkles, lack of adequate skin firmness, lack of adequate dermal hydration, or insufficient sebum secretion; renal osteodystrophy; or osteoporosis.

9. The method of claim 8, wherein the biological condition is psoriasis.

5 10. The method of claim 8, wherein the biological condition is selected from leukemia, colon cancer, breast cancer, or prostate cancer.

11. The method of claim 8, wherein the biological condition is selected from multiple sclerosis, lupus, diabetes mellitus host versus graft reaction, or rejection of organ transplants.

10 12. The method of claim 8, wherein the biological condition is selected from rheumatoid arthritis, asthma, or inflammatory bowel diseases selected from celiac disease, ulcerative colitis and Crohn's disease.

13. The method of claim 8, wherein the biological condition is selected from wrinkles, lack of adequate skin firmness, lack of adequate dermal hydration, or insufficient sebum secretion.


14. The method of claim 8, wherein the compound is administered orally to the subject.

15. The method of claim 8, wherein administration of the compound to the subject is selected from parenterally, transdermally or topically.

20 16. The method of claim 8, wherein the compound is administered in a dosage of from 0.01 μ g per day to 1 mg per day.

17. The use of the compound of claim 4 in the preparation of a medicament for the treatment of a biological condition selected from psoriasis; leukemia; colon cancer; breast cancer; prostate cancer; multiple sclerosis; lupus; diabetes mellitus; host versus graft reaction; rejection of organ transplants; an inflammatory disease selected from rheumatoid arthritis, asthma, or inflammatory bowel diseases; a skin condition selected from wrinkles, lack of adequate skin firmness, lack of adequate dermal hydration, or insufficient sebum secretion; renal osteodystrophy; or osteoporosis.

25 18. The compound of claim 1, wherein X^1 and X^2 are both H and the compound has the formula IB



19. The compound of claim 1 substantially as hereinbefore described with reference to any one of the examples.

20. A process for preparing the compound of claim 1 substantially as hereinbefore described with reference to any one of the examples.

Dated 30 May 2011
Wisconsin Alumni Research Foundation
Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

FIGURE 1




FIGURE 2

HL-60 Cell Differentiation





FIGURE 3

24-OHase Transcription



EC₅₀: 1,25(OH)₂D₃ = 1.7×10^{-10} M
19,21-dinor = 1.0×10^{-8} M

FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7
PTH Suppression in Adult Rats

