



(86) Date de dépôt PCT/PCT Filing Date: 2002/11/07
(87) Date publication PCT/PCT Publication Date: 2003/05/15
(85) Entrée phase nationale/National Entry: 2004/04/01
(86) N° demande PCT/PCT Application No.: US 2002/036002
(87) N° publication PCT/PCT Publication No.: 2003/040340
(30) Priorités/Priorities: 2001/11/07 (10/005,480) US;
2002/01/31 (10/062,109) US

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/12, A61K 31/7088, A61K 47/48,
A61K 39/395, A61K 38/17, A61K 49/16, A61K 35/14,
A01K 67/027, A61K 48/00, C07K 16/30, C07K 14/47,
C12Q 1/68, G01N 33/574

(71) Demandeurs/Applicants:
AGENSYS INC., US;
AGENSYS, INC., US

(72) Inventeurs/Inventors:
JAKOBOVITS, AYA, US;
RAITANO, ARTHUR B., US;
FARIS, MARY, US;
...

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : ACIDE NUCLEIQUE ET PROTEINE CORRESPONDANTE 161P2F10B UTILES DANS LE TRAITEMENT ET
LE DEPISTAGE DU CANCER

(54) Title: NUCLEIC ACID AND CORRESPONDING PROTEIN ENTITLED 161P2F10B USEFUL IN TREATMENT AND
DETECTION OF CANCER

1 GATCACACAT TAGGTTATNG ACTTCAATAT TTTCAAATGG TTCAACTTCA GTCTTCTCTT
61 TAAAACTGGG TCCATGTGCC AAGAAAGATA GCCTCCATGC TCCTAAACTC ATTGTTATAA
121 CCATGGTTGC CTCCTCCACA ATTTGTATTT GATTTACTCC TAACAGCCAG CCACTGTTGA
181 TC

(57) Abrégé/Abstract:

A novel gene 0161P2F10B (also designated 161P2F10B) and its encoded protein, and variants thereof, are described wherein 161P2F10B exhibits tissue specific expression in normal adult tissue, and is aberrantly expressed in the cancers listed in Table I. Consequently, 161P2F10B provides a diagnostic, prognostic, prophylactic and/or therapeutic target for cancer. The 161P2F10B gene of fragment thereof, or its encoded protein, or variants thereof, or a fragment thereof, can be used to elicit a humoral or cellular immune response; antibodies or T cells reactive with 161P2F10B can be used in active or passive immunization.



(72) **Inventeurs(suite)/Inventors(continued)**: HUBERT, RENE S., US; GE, WANGMAO, US;
MORRISON, KAREN JANE MEYRICK, US; MORRISON, ROBERT KENDALL, US; CHALLITA-EID, PIA M., US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2003 (15.05.2003)

PCT

(10) International Publication Number
WO 03/040340 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number: PCT/US02/36002

(22) International Filing Date:
7 November 2002 (07.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/005,480 7 November 2001 (07.11.2001) US
10/062,109 31 January 2002 (31.01.2002) US

(71) Applicant (*for all designated States except US*): **AGENSYS, INC.** [US/US]; 1545 17th Street, Santa Monica, CA 90404 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **JAKOBOVITS, Aya** [US/US]; 3135 Hutton Drive, Beverly Hills, CA 90210 (US). **RAITANO, Arthur, B.** [US/US]; 12685 Rose Avenue, Los Angeles, CA 90066 (US). **FARIS, Mary** [US/US]; 2538 Almaden Court, Los Angeles, CA 90077 (US). **HUBERT, Rene, S.** [CA/US]; 1644 No. Occidental Boulevard, Los Angeles, CA 90026 (US). **GE, Wangmao** [CN/US]; 4838 Hollow Corner Road, Apt. #314, Culver City, CA 90230 (US). **MORRISON, Karen, Jane, Meyrick** [GB/US]; 1044 Yale Street, Santa Monica, CA 90403 (US). **MORRISON, Robert, Kendall** [GB/US]; 1044 Yale Street, Santa Monica, CA 90403 (US). **CHALLITA-EID, Pia, M.** [LB/US]; 15745 Morrison Street, Encino, CA 91436 (US).

(74) Agents: **DEVERNOE, David, L.** et al.; Morrisson & Forster LLP, Suite 500, 3811 Valley Centre Drive, San Diego, CA 92130-2332 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(48) Date of publication of this corrected version:

7 August 2003

(15) Information about Correction:


see PCT Gazette No. 32/2003 of 7 August 2003, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID AND CORRESPONDING PROTEIN ENTITLED 161P2F10B USEFUL IN TREATMENT AND DETECTION OF CANCER

1 GATCACACAT TAGGTTATNG ACTTCAATAT TTTCAAATGG TTCAACTTCA GTCTTCTCTT
61 TAAAACTGGG TCCATGTGCC AAGAAAGATA GCCTCCATGC TCCTAAACTC ATTGTTATAA
121 CCATGGTTGC CTCCTCCACA ATTTGTATTT GATTTACTCC TAACAGCCAG CCACTGTTGA
181 TC

(57) Abstract: A novel gene 0161P2F10B (also designated 161P2F10B) and its encoded protein, and variants thereof, are described wherein 161P2F10B exhibits tissue specific expression in normal adult tissue, and is aberrantly expressed in the cancers listed in Table I. Consequently, 161P2F10B provides a diagnostic, prognostic, prophylactic and/or therapeutic target for cancer. The 161P2F10B gene of fragment thereof, or its encoded protein, or variants thereof, or a fragment thereof, can be used to elicit a humoral or cellular immune response; antibodies or T cells reactive with 161P2F10B can be used in active or passive immunization.



WO 03/040340 A2

**NUCLEIC ACID AND CORRESPONDING PROTEIN ENTITLED 161P2F10B
USEFUL IN TREATMENT AND DETECTION OF CANCER**

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority from pending United States patent application USSN 10/005,480, filed November 7, 2001; pending United States patent application number 10/062,109, filed January 31, 2002. The contents of each application listed in this paragraph are fully incorporated by reference herein.

STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

Not applicable.

FIELD OF THE INVENTION

The invention described herein relates to a gene and its encoded protein, termed 161P2F10B, expressed in certain cancers, and to diagnostic and therapeutic methods and compositions useful in the management of cancers that express 161P2F10B.

BACKGROUND OF THE INVENTION

Cancer is the second leading cause of human death next to coronary disease. Worldwide, millions of people die from cancer every year. In the United States alone, as reported by the American Cancer Society, cancer causes the death of well over a half-million people annually, with over 1.2 million new cases diagnosed per year. While deaths from heart disease have been declining significantly, those resulting from cancer generally are on the rise. In the early part of the next century, cancer is predicted to become the leading cause of death.

Worldwide, several cancers stand out as the leading killers. In particular, carcinomas of the lung, prostate, breast, colon, pancreas, and ovary represent the primary causes of cancer death. These and virtually all other carcinomas share a common lethal feature. With very few exceptions, metastatic disease from a carcinoma is fatal. Moreover, even for those cancer patients who initially survive their primary cancers, common experience has shown that their lives are dramatically altered. Many cancer patients experience strong anxieties driven by the awareness of the potential for recurrence or treatment failure. Many cancer patients experience physical debilitations following treatment. Furthermore, many cancer patients experience a recurrence.

Worldwide, prostate cancer is the fourth most prevalent cancer in men. In North America and Northern Europe, it is by far the most common cancer in males and is the second leading cause of cancer death in men. In the United States alone, well over 30,000 men die annually of this disease - second only to lung cancer. Despite the magnitude of these figures, there is still no effective treatment for metastatic prostate cancer. Surgical prostatectomy, radiation therapy, hormone ablation therapy, surgical castration and chemotherapy continue to be the main treatment modalities. Unfortunately, these treatments are ineffective for many and are often associated with undesirable consequences.

On the diagnostic front, the lack of a prostate tumor marker that can accurately detect early-stage, localized tumors remains a significant limitation in the diagnosis and management of this disease. Although the serum prostate specific antigen (PSA) assay has been a very useful tool, however its specificity and general utility is widely regarded as lacking in several important respects.

Progress in identifying additional specific markers for prostate cancer has been improved by the generation of prostate cancer xenografts that can recapitulate different stages of the disease in mice. The LAPC (Los Angelos Prostate Cancer) xenografts are prostate cancer xenografts that have survived passage in severe combined immune deficient (SCID) mice and have exhibited the capacity to mimic the transition from androgen dependence to androgen independence (Klein et

mice and have exhibited the capacity to mimic the transition from androgen dependence to androgen independence (Klein *et al.*, 1997, Nat. Med. 3:402). More recently identified prostate cancer markers include PCTA-1 (Su *et al.*, 1996, Proc. Natl. Acad. Sci. USA 93: 7252), prostate-specific membrane (PSM) antigen (Pinto *et al.*, Clin Cancer Res 1996 Sep 2 (9): 1445-51), STEAP (Hubert, *et al.*, Proc Natl Acad Sci U S A. 1999 Dec 7; 96(25): 14523-8) and prostate stem cell antigen (PSCA) (Reiter *et al.*, 1998, Proc. Natl. Acad. Sci. USA 95: 1735).

While previously identified markers such as PSA, PSM, PCTA and PSCA have facilitated efforts to diagnose and treat prostate cancer, there is need for the identification of additional markers and therapeutic targets for prostate and related cancers in order to further improve diagnosis and therapy.

Renal cell carcinoma (RCC) accounts for approximately 3 percent of adult malignancies. Once adenomas reach a diameter of 2 to 3 cm, malignant potential exists. In the adult, the two principal malignant renal tumors are renal cell adenocarcinoma and transitional cell carcinoma of the renal pelvis or ureter. The incidence of renal cell adenocarcinoma is estimated at more than 29,000 cases in the United States, and more than 11,600 patients died of this disease in 1998. Transitional cell carcinoma is less frequent, with an incidence of approximately 500 cases per year in the United States.

Surgery has been the primary therapy for renal cell adenocarcinoma for many decades. Until recently, metastatic disease has been refractory to any systemic therapy. With recent developments in systemic therapies, particularly immunotherapies, metastatic renal cell carcinoma may be approached aggressively in appropriate patients with a possibility of durable responses. Nevertheless, there is a remaining need for effective therapies for these patients.

Of all new cases of cancer in the United States, bladder cancer represents approximately 5 percent in men (fifth most common neoplasm) and 3 percent in women (eighth most common neoplasm). The incidence is increasing slowly, concurrent with an increasing older population. In 1998, there was an estimated 54,500 cases, including 39,500 in men and 15,000 in women. The age-adjusted incidence in the United States is 32 per 100,000 for men and eight per 100,000 in women. The historic male/female ratio of 3:1 may be decreasing related to smoking patterns in women. There were an estimated 11,000 deaths from bladder cancer in 1998 (7,800 in men and 3,900 in women). Bladder cancer incidence and mortality strongly increase with age and will be an increasing problem as the population becomes more elderly.

Most bladder cancers recur in the bladder. Bladder cancer is managed with a combination of transurethral resection of the bladder (TUR) and intravesical chemotherapy or immunotherapy. The multifocal and recurrent nature of bladder cancer points out the limitations of TUR. Most muscle-invasive cancers are not cured by TUR alone. Radical cystectomy and urinary diversion is the most effective means to eliminate the cancer but carry an undeniable impact on urinary and sexual function. There continues to be a significant need for treatment modalities that are beneficial for bladder cancer patients.

An estimated 130,200 cases of colorectal cancer occurred in 2000 in the United States, including 93,800 cases of colon cancer and 36,400 of rectal cancer. Colorectal cancers are the third most common cancers in men and women. Incidence rates declined significantly during 1992-1996 (-2.1% per year). Research suggests that these declines have been due to increased screening and polyp removal, preventing progression of polyps to invasive cancers. There were an estimated 56,300 deaths (47,700 from colon cancer, 8,600 from rectal cancer) in 2000, accounting for about 11% of all U.S. cancer deaths.

At present, surgery is the most common form of therapy for colorectal cancer, and for cancers that have not spread, it is frequently curative. Chemotherapy, or chemotherapy plus radiation, is given before or after surgery to most patients whose cancer has deeply perforated the bowel wall or has spread to the lymph nodes. A permanent colostomy (creation of an abdominal opening for elimination of body wastes) is occasionally needed for colon cancer and is infrequently required for rectal cancer. There continues to be a need for effective diagnostic and treatment modalities for colorectal cancer.

There were an estimated 164,100 new cases of lung and bronchial cancer in 2000, accounting for 14% of all U.S. cancer diagnoses. The incidence rate of lung and bronchial cancer is declining significantly in men, from a high of 86.5 per 100,000 in 1984 to 70.0 in 1996. In the 1990s, the rate of increase among women began to slow. In 1996, the incidence rate in women was 42.3 per 100,000.

Lung and bronchial cancer caused an estimated 156,900 deaths in 2000, accounting for 28% of all cancer deaths. During 1992–1996, mortality from lung cancer declined significantly among men (-1.7% per year) while rates for women were still significantly increasing (0.9% per year). Since 1987, more women have died each year of lung cancer than breast cancer, which, for over 40 years, was the major cause of cancer death in women. Decreasing lung cancer incidence and mortality rates most likely resulted from decreased smoking rates over the previous 30 years; however, decreasing smoking patterns among women lag behind those of men. Of concern, although the declines in adult tobacco use have slowed, tobacco use in youth is increasing again.

Treatment options for lung and bronchial cancer are determined by the type and stage of the cancer and include surgery, radiation therapy, and chemotherapy. For many localized cancers, surgery is usually the treatment of choice. Because the disease has usually spread by the time it is discovered, radiation therapy and chemotherapy are often needed in combination with surgery. Chemotherapy alone or combined with radiation is the treatment of choice for small cell lung cancer; on this regimen, a large percentage of patients experience remission, which in some cases is long lasting. There is however, an ongoing need for effective treatment and diagnostic approaches for lung and bronchial cancers.

An estimated 182,800 new invasive cases of breast cancer were expected to occur among women in the United States during 2000. Additionally, about 1,400 new cases of breast cancer were expected to be diagnosed in men in 2000. After increasing about 4% per year in the 1980s, breast cancer incidence rates in women have leveled off in the 1990s to about 110.6 cases per 100,000.

In the U.S. alone, there were an estimated 41,200 deaths (40,800 women, 400 men) in 2000 due to breast cancer. Breast cancer ranks second among cancer deaths in women. According to the most recent data, mortality rates declined significantly during 1992–1996 with the largest decreases in younger women, both white and black. These decreases were probably the result of earlier detection and improved treatment.

Taking into account the medical circumstances and the patient's preferences, treatment of breast cancer may involve lumpectomy (local removal of the tumor) and removal of the lymph nodes under the arm; mastectomy (surgical removal of the breast) and removal of the lymph nodes under the arm; radiation therapy; chemotherapy; or hormone therapy. Often, two or more methods are used in combination. Numerous studies have shown that, for early stage disease, long-term survival rates after lumpectomy plus radiotherapy are similar to survival rates after modified radical mastectomy. Significant advances in reconstruction techniques provide several options for breast reconstruction after mastectomy. Recently, such reconstruction has been done at the same time as the mastectomy.

Local excision of ductal carcinoma *in situ* (DCIS) with adequate amounts of surrounding normal breast tissue may prevent the local recurrence of the DCIS. Radiation to the breast and/or tamoxifen may reduce the chance of DCIS occurring in the remaining breast tissue. This is important because DCIS, if left untreated, may develop into invasive breast cancer. Nevertheless, there are serious side effects or sequelae to these treatments. There is, therefore, a need for efficacious breast cancer treatments.

There were an estimated 23,100 new cases of ovarian cancer in the United States in 2000. It accounts for 4% of all cancers among women and ranks second among gynecologic cancers. During 1992–1996, ovarian cancer incidence rates were significantly declining. Consequent to ovarian cancer, there were an estimated 14,000 deaths in 2000. Ovarian cancer causes more deaths than any other cancer of the female reproductive system.

Surgery, radiation therapy, and chemotherapy are treatment options for ovarian cancer. Surgery usually includes the removal of one or both ovaries, the fallopian tubes (salpingo-oophorectomy), and the uterus (hysterectomy). In some very early tumors, only the involved ovary will be removed, especially in young women who wish to have children. In advanced disease, an attempt is made to remove all intra-abdominal disease to enhance the effect of chemotherapy. There continues to be an important need for effective treatment options for ovarian cancer.

There were an estimated 28,300 new cases of pancreatic cancer in the United States in 2000. Over the past 20 years, rates of pancreatic cancer have declined in men. Rates among women have remained approximately constant but may be beginning to decline. Pancreatic cancer caused an estimated 28,200 deaths in 2000 in the United States. Over the past 20 years, there has been a slight but significant decrease in mortality rates among men (about -0.9% per year) while rates have increased slightly among women.

Surgery, radiation therapy, and chemotherapy are treatment options for pancreatic cancer. These treatment options can extend survival and/or relieve symptoms in many patients but are not likely to produce a cure for most. There is a significant need for additional therapeutic and diagnostic options for pancreatic cancer.

SUMMARY OF THE INVENTION

The present invention relates to a gene, designated 161P2F10B, that has now been found to be over-expressed in the cancer(s) listed in Table I. Northern blot expression analysis of 161P2F10B gene expression in normal tissues shows a restricted expression pattern in adult tissues. The nucleotide (Figure 2) and amino acid (Figure 2, and Figure 3) sequences of 161P2F10B are provided. The tissue-related profile of 161P2F10B in normal adult tissues, combined with the over-expression observed in the tissues listed in Table I, shows that 161P2F10B is aberrantly over-expressed in at least some cancers, and thus serves as a useful diagnostic, prophylactic, prognostic, and/or therapeutic target for cancers of the tissue(s) such as those listed in Table I.

The invention provides polynucleotides corresponding or complementary to all or part of the 161P2F10B genes, mRNAs, and/or coding sequences, preferably in isolated form, including polynucleotides encoding 161P2F10B-related proteins and fragments of 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more than 25 contiguous amino acids; at least 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 85, 90, 95, 100 or more than 100 contiguous amino acids of a 161P2F10B-related protein, as well as the peptides/proteins themselves; DNA, RNA, DNA/RNA hybrids, and related molecules, polynucleotides or oligonucleotides complementary or having at least a 90% homology to the 161P2F10B genes or mRNA sequences or parts thereof, and polynucleotides or oligonucleotides that hybridize to the 161P2F10B genes, mRNAs, or to 161P2F10B-encoding polynucleotides. Also provided are means for isolating cDNAs and the genes encoding 161P2F10B. Recombinant DNA molecules containing 161P2F10B polynucleotides, cells transformed or transduced with such molecules, and host-vector systems for the expression of 161P2F10B gene products are also provided. The invention further provides antibodies that bind to 161P2F10B proteins and polypeptide fragments thereof, including polyclonal and monoclonal antibodies, murine and other mammalian antibodies, chimeric antibodies, humanized and fully human antibodies, and antibodies labeled with a detectable marker or therapeutic agent. In certain embodiments, there is a *proviso* that the entire nucleic acid sequence of Figure 2 is not encoded and/or the entire amino acid sequence of Figure 2 is not prepared. In certain embodiments, the entire nucleic acid sequence of Figure 2 is encoded and/or the entire amino acid sequence of Figure 2 is prepared, either of which are in respective human unit dose forms.

The invention further provides methods for detecting the presence and status of 161P2F10B polynucleotides and proteins in various biological samples, as well as methods for identifying cells that express 161P2F10B. A typical embodiment of this invention provides methods for monitoring 161P2F10B gene products in a tissue or hematology sample having or suspected of having some form of growth dysregulation such as cancer.

The invention further provides various immunogenic or therapeutic compositions and strategies for treating cancers that express 161P2F10B such as cancers of tissues listed in Table I, including therapies aimed at inhibiting the transcription, translation, processing or function of 161P2F10B as well as cancer vaccines. In one aspect, the invention provides compositions, and methods comprising them, for treating a cancer that expresses 161P2F10B in a human subject wherein the composition comprises a carrier suitable for human use and a human unit dose of one or more than one agent that inhibits the production or function of 161P2F10B. Preferably, the carrier is a uniquely human carrier. In another aspect of the invention, the agent is a moiety that is immunoreactive with 161P2F10B protein. Non-limiting examples of such moieties include, but are not limited to, antibodies (such as single chain, monoclonal, polyclonal, humanized, chimeric, or human antibodies), functional equivalents thereof (whether naturally occurring or synthetic), and combinations thereof. The antibodies can be conjugated to a diagnostic or therapeutic moiety. In another aspect, the agent is a small molecule as defined herein.

In another aspect, the agent comprises one or more than one peptide which comprises a cytotoxic T lymphocyte (CTL) epitope that binds an HLA class I molecule in a human to elicit a CTL response to 161P2F10B and/or one or more than one peptide which comprises a helper T lymphocyte (HTL) epitope which binds an HLA class II molecule in a human to elicit an HTL response. The peptides of the invention may be on the same or on one or more separate polypeptide molecules. In a further aspect of the invention, the agent comprises one or more than one nucleic acid molecule that expresses one or more than one of the CTL or HTL response stimulating peptides as described above. In yet another aspect of the invention, the one or more than one nucleic acid molecule may express a moiety that is immunologically reactive with 161P2F10B as described above. The one or more than one nucleic acid molecule may also be, or encodes, a molecule that inhibits production of 161P2F10B. Non-limiting examples of such molecules include, but are not limited to, those complementary to a nucleotide sequence essential for production of 161P2F10B (e.g. antisense sequences or molecules that form a triple helix with a nucleotide double helix essential for 161P2F10B production) or a ribozyme effective to lyse 161P2F10B mRNA.

Note that to determine the starting position of any peptide set forth in Tables VIII-XXI and XXII to XLIX (collectively HLA Peptide Tables) respective to its parental protein, e.g., variant 1, variant 2, etc., reference is made to three factors: the particular variant, the length of the peptide in an HLA Peptide Table, and the Search Peptides in Table VII. Generally, a unique Search Peptide is used to obtain HLA peptides of a particular for a particular variant. The position of each Search Peptide relative to its respective parent molecule is listed in Table VII. Accordingly, if a Search Peptide begins at position "X", one must add the value "X - 1" to each position in Tables VIII-XXI and XXII to XLIX to obtain the actual position of the HLA peptides in their parental molecule. For example, if a particular Search Peptide begins at position 150 of its parental molecule, one must add 150 - 1, i.e., 149 to each HLA peptide amino acid position to calculate the position of that amino acid in the parent molecule.

One embodiment of the invention comprises an HLA peptide, that occurs at least twice in Tables VIII-XXI and XXII to XLIX collectively, or an oligonucleotide that encodes the HLA peptide. Another embodiment of the invention comprises an HLA peptide that occurs at least once in Tables VIII-XXI and at least once in tables XXII to XLIX, or an oligonucleotide that encodes the HLA peptide.

Another embodiment of the invention is antibody epitopes, which comprise a peptide regions, or an oligonucleotide encoding the peptide region, that has one two, three, four, or five of the following characteristics:

- i) a peptide region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Hydrophilicity profile of Figure 5;

- ii) a peptide region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or less than 0.5, 0.4, 0.3, 0.2, 0.1, or having a value equal to 0.0, in the Hydropathicity profile of Figure 6;
- iii) a peptide region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Percent Accessible Residues profile of Figure 7;
- iv) a peptide region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Average Flexibility profile of Figure 8; or
- v) a peptide region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Beta-turn profile of Figure 9.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. The 161P2F10B SSH sequence of 182 nucleotides.

Figure 2. A) The cDNA and amino acid sequence of 161P2F10B variant 1 (also called "161P2F10B v.1" or "161P2F10B variant 1") is shown in Figure 2A. The start methionine is underlined. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

B) The cDNA and amino acid sequence of 161P2F10B variant 2 (also called "161P2F10B v.2") is shown in Figure 2B. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

C) The cDNA and amino acid sequence of 161P2F10B variant 3 (also called "161P2F10B v.3") is shown in Figure 2C. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 44-2671 including the stop codon. The cDNA and amino acid sequence of 161P2F10B variant 4 (also called "161P2F10B v.4") is shown in

D) Figure 2D. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

E) The cDNA and amino acid sequence of 161P2F10B variant 5 (also called "161P2F10B v.5") is shown in Figure 2E. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

F) The cDNA and amino acid sequence of 161P2F10B variant 6 (also called "161P2F10B v.6") is shown in Figure 2F. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 84-2711 including the stop codon.

G) The cDNA and amino acid sequence of 161P2F10B variant 7 (also called "161P2F10B v.7") is shown in Figure 2G. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 276-2801 including the stop codon.

Figure 3.

A) Amino acid sequence of 161P2F10B v.1 is shown in Figure 3A; it has 875 amino acids.

B) The amino acid sequence of 161P2F10B v.2 is shown in Figure 3B; it has 875 amino acids.

C) The amino acid sequence of 161P2F10B v.3 is shown in Figure 3C; it has 875 amino acids.

D) The amino acid sequence of 161P2F10B v.4 is shown in Figure 3D; it has 875 amino acids.

E) The amino acid sequence of 161P2F10B v.7 is shown in Figure 3E; it has 841 amino acids. As used herein, a reference to 161P2F10B includes all variants thereof, including those shown in Figures 2, 3, 10, and 11, unless the context clearly indicates otherwise.

Figure 4. Figure 4A: Alignment of 161P2F10 with variant 1 carrying a K to R mutation. Figure 4B: Alignment of 161P2F10B and SNP variant carrying a T to P mutation.

Figure 5. Hydrophilicity amino acid profile of 161P2F10B determined by computer algorithm sequence analysis using the method of Hopp and Woods (Hopp T.P., Woods K.R., 1981. Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828) accessed on the Protscale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

Figure 6. Hydropathicity amino acid profile of 161P2F10B determined by computer algorithm sequence analysis using the method of Kyte and Doolittle (Kyte J., Doolittle R.F., 1982. J. Mol. Biol. 157:105-132) accessed on the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

Figure 7. Percent accessible residues amino acid profile of 161P2F10B determined by computer algorithm sequence analysis using the method of Janin (Janin J., 1979 Nature 277:491-492) accessed on the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

Figure 8.. Average flexibility amino acid profile of 161P2F10B determined by computer algorithm sequence analysis using the method of Bhaskaran and Ponnuswamy (Bhaskaran R., and Ponnuswamy P.K., 1988. Int. J. Pept. Protein Res. 32:242-255) accessed on the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

Figure 9. Beta-turn amino acid profile of 161P2F10B determined by computer algorithm sequence analysis using the method of Deleage and Roux (Deleage, G., Roux B. 1987 Protein Engineering 1:289-294) accessed on the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

Figure 10. Variants 161P2F10B v.2 through v.5 are variants with single nucleotide differences. Though these SNP variants are shown separately, they could also occur in any combinations and in any transcript variants that contains the base pairs. Variants 161P2F10B v.6 and v.7 are transcript variants. Variant 161P2F10B v.6 has extra 40 bases at the 5' end and a different 3' end portion, while variant 161P2F10B v.7 has an insertion of 130 bases in between positions 121 and 122 of 161P2F10B v.1. Numbers in "()" correspond to those of 161P2F10B v.1. Black box shows the same sequence as 161P2F10B v.1. SNPs are indicated above the box.

Figure 11. Protein variants correspond to nucleotide variants. Nucleotide variants 161P2F10B v.5 and v.6 in Figure 10 code for the same protein as 161P2F10B v.1. Nucleotide variants 161P2F10B v.6 and v.7 are splice variants of v.1, as shown in Figure 12. Single amino acid differences were indicated above the boxes. Black boxes represent the same sequence as 161P2F10B v.1. Numbers underneath the box correspond to 161P2F10B v.1.

Figure 12. Intentionally Omitted

Figure 13. The secondary structure of 161P2F10B (SEQ ID NO: 103), namely the predicted presence and location of alpha helices, extended strands, and random coils, is predicted from the primary amino acid sequence using the HNN - Hierarchical Neural Network method (Guerneur, 1997, on the World Wide Web at [://pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_nn.html](http://pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_nn.html)), accessed from the ExPasy molecular biology server (on the World Wide Web at expasy.ch/tools/). The analysis indicates that 161P2F10B is composed 31.31% alpha helix, 11.31% extended strand, and 57.37% random coil (Figure 13A). Shown graphically in Figure 13 panels B and C are the results of analysis using the TMpred (Figure 13B) and TMHMM (Figure 13C) prediction programs depicting the location of the transmembrane domain.

Figure 14. First strand cDNA was generated from normal stomach, normal brain, normal heart, normal liver, normal skeletal muscle, normal testis, normal prostate, normal bladder, normal kidney, normal colon, normal lung, normal

pancreas, and a pool of cancer specimens from prostate cancer patients, bladder cancer patients, kidney cancer patients, colon cancer patients, lung cancer patients, pancreas cancer patients, a pool of prostate cancer xenografts (LAPC-4AD, LAPC-4AI, LAPC-9AD and LAPC-9AI), and a pool of 2 patient prostate metastasis to lymph node. Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 161P2F10B, was performed at 26 and 30 cycles of amplification. Samples were run on an agarose gel, and PCR products were quantitated using the Alphamager software. Results show strong expression in prostate cancer, bladder cancer, kidney cancer, colon cancer, lung cancer, pancreas cancer, bone cancer, lymphoma cancer, uterus cancer, compared to all normal tissues tested. Strong expression was also detected in the xenograft pool as well as the prostate cancer metastasis to lymph node specimens.

Figure 15. First strand cDNA was prepared from a panel of kidney cancer clear cell carcinoma (A), kidney cancer papillary carcinoma (B), and in uterus patient cancer specimens (C). Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 161P2F10B, was performed at 26 and 30 cycles of amplification. Samples were run on an agarose gel, and PCR products were quantitated using the Alphamager software. Expression was recorded as absent, low, medium or strong. Results show expression of 161P2F10B in 94.7% of clear cell renal carcinoma, 62.5% of papillary renal cell carcinoma, and in 61.5% of uterus cancer.

Figure 16. Shows Phosphodiesterase Activity of 3T3-161P2F10B Stable Cells. Cell surface phosphodiesterase activity is assayed on 3T3 and 3T3-161P2F10B using the substrate p-nitrophenyl thymidine -5'-L-monophosphate.

Figure 17. Shows Protection from Apoptosis by 161P2F10B.

Figure 18. Shows that 161P2F10B Protects from Apoptotic Signals.

Figure 19. Shows that 161P2F10B Protects from Staurosporine and UV-Induced Apoptosis.

Figure 20. Shows that 161P2F10B Expression Protects Cells from Drug and UV-Induced Apoptosis. NIH 3T3 cells were treated with the staurosporine or UV, stained with Annexin V-FITC and propidium iodide, and analyzed by FACS.

Figure 21. Shows that 161P2F10B Protects from Apoptosis by Chemotherapeutic Agents.

Figure 22 Shows the effect of 161P2F10B on *In Vitro* Invasion. Invasion was determined by measuring the fluorescence of cells in the lower chamber relative to the fluorescence of the entire cell population.

Figure 23. Shows that 161P2F10B MAb Attenuates the Growth of Human Kidney Cancer Xenograft in SCID Mice.

Figure 24. Detection of 161P2F10B protein by immunohistochemistry in kidney cancer patient specimens. Two renal clear cell carcinoma tissue specimens and one renal papillary cell carcinoma were obtained from three different kidney cancer patients. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours. The slides were washed three times in buffer, and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed strong expression of 161P2F10B in all three renal carcinoma patient tissues (Figure 24 panels A-C). The expression was detected mostly around the cell membrane in the renal clear cell carcinoma specimens, indicating that 161P2F10B is membrane associated in this kidney cancer, and throughout the cells in the papillary cell carcinoma with an apparent predisposition towards the cell periphery.

Figure 25. Detection of 161P2F10B protein by immunohistochemistry in a prostate cancer patient specimen. Tissue specimens of prostate adenocarcinoma were obtained from eight different prostate cancer patients. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours. The slides were washed three times in buffer, and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA

Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed expression of 161P2F10B in six of the eight prostate cancer patient tissues, one of which is illustrated in this Figure 25. 161P2F10B was expressed on the tumor cells with an apparent proclivity towards the luminal cell surface.

Figure 26. Detection of 161P2F10B protein by immunohistochemistry in a colon cancer patient specimen. Tissue specimens of colon adenocarcinoma were obtained from nine different colon cancer patients. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours. The slides were washed three times in buffer, and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed strong expression of 161P2F10B in two of the nine colon cancer patient tissues, one of which is illustrated in this Figure 26. 161P2F10B was most strongly expressed on the tumor cells with a luminal cell surface but was also expressed throughout all the tumor tissue.

Figure 27. Detection by immunohistochemistry of 161P2F10B protein expression in kidney clear cell cancer patient specimens by specific binding of mouse monoclonal antibodies. Renal clear cell carcinoma tissue and its matched normal adjacent were obtained from a kidney cancer patient. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated either mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours (Figure 27 panels A, D), or mouse monoclonal antibody X41(3)50 (Figure 27 panels B, E), or mouse monoclonal antibody X41(3)37 (Figure 27 panels C, F). The slides were washed three times in buffer and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy (Figure 27 panels A-F). The results showed strong expression of 161P2F10B in the renal clear cell carcinoma patient tissue (Figure 27 panels A-C), but weakly in normal kidney (Figure 27 panels D-F). The expression was predominantly around the cell periphery indicating that 161P2F10B is membrane associated in kidney cancer tissues. The weak expression detected in normal kidney was localized to the kidney proximal tubules.

Figure 28. Expression of 161P2F10b in recombinant cell lines.

A.) Rat1, NIH3T3, NSO, and 300.19 cells stably expressing either 16P2F10b or a control vector (neo) were stained with PE-conjugated anti-CD203c MAb and examined by flow cytometry. (Light dotted line: control neo cells. Dark line: 161P2F10 cells)

B.) Rat1, NIH3T3, NSO, 300.19, and UT7 cells were stained with either PE-conjugated anti-CD203c MAb or control IgG1-PE Ab and examined by flow cytometry. (Light dotted line: control MAb. Dark line: 97A6 (CD203c) MAb.) Shown is the mean fluorescence of the staining of the control and 161P2F10b cells and the ratio of the values. This was used to rank the cells for relative expression levels of 161P2F10b.

C.) The relative cell surface phosphodiesterase enzymatic activity of the recombinant cells was measured by the addition of p-nitrophenyl thymidine-5'-L-monophosphate (p-nTMP) phosphodiesterase substrate. There is a correlation between expression levels determined by flow cytometry and surface enzyme activity.

Figure 29. Surface expression and phosphodiesterase activity of 161P2F10b.

A. 161P2F10b transfected 293T cells were stained with the commercially available (Coulter Immunotech) PE-conjugated anti-CD-203c MAb, a commercially available anti-ENPP3 (161P2F10b) MAb and examined by fluorescent microscopy.

B. 161P2F10b and vector transfected 293T cells were incubated in assay buffer containing the phosphodiesterase-1 colorimetric substrate p-nitrophenyl thymidine-5'-L-monophosphate (p-nTMP) and optical densities (O.D.) were obtained at 405nm.

Figure 30. Relative expression and enzymatic activity of 161P2F10b mutants in recombinant Caki kidney cancer cells. Caki kidney cancer cells were infected with retrovirus containing either wildtype 161P2F120b cDNA, or point mutant cDNAs encoding either a threonine to serine mutation (T/S) at amino acid 205, a threonine to alanine mutation (T/A) at amino acid 205, or a aspartic acid to glutamic acid mutation (D/E) at amino acid 80. Stably expressing cell lines were analyzed for 161P2F10b expression by flow cytometry with 97A6 (CD203c) MAb (A) and for enzymatic activity with p-nTMP substrate (B). Mutation of threonine 205 to aspartic acid or alanine abolishes the ability to cleave the substrate, demonstrating that threonine 205 is critical to the enzymatic activity of 161P2F10b.

Figure 31. Purification of a recombinant protein encoding the extracellular domain (ECD) of 161P2F10b. 293T cells were transfected with a Tag5 secretion expression vector encoding the ECD of 161P2F10b (amino acids 46-875). The recombinant protein was purified from the conditioned media using either metal chelate affinity chromatography (not shown) or with an immunoaffinity column comprised of anti-161P2F10b MAb X41.6 (shown). 2 ul of 2 separate purified lots were analyzed by SDS-PAGE and Coomassie staining. BSA protein was also analyzed as a quantitative standard.

Figure 32. 161P2F10b enzymatic assays utilizing P-nitrophenyl-thymidine monophosphate (p-nTMP).

A. Schematic of the colorimetric enzyme assay showing enzymatic cleavage of the p-nTMP substrate generating a soluble yellow product.

B. Kinetics and dose response of the enzymatic action of purified Tag5-ECD 161P2F10b protein on p-nTMP (2.5 mM). Optical densities (OD) of reactions were determined at 405nm.

C. Cell surface enzymatic assay of 161P2F10b-expressing Rat1 cells. The indicated number of Rat1-161P2F10b cells were incubated with p-nTMP substrate and the OD's of the wells were determined.

D. ATP and NAD (not shown) serve as competitive inhibitors 161P2F10b cleavage of p-nTMP. Purified Tag5-ECD protein (20 ng) was incubated with p-nTMP substrate in the absence or presence of the indicated amounts of ATP. The OD's of reactions were obtained at 405 nm.

Figure 33. Analysis of the internalization of anti-161P2F10b MAb X41.6.

Panel A. Schematic of the protocol. Rat1-161P2F10b cells are incubated with anti-161P2F10b MAb at 4C, washed, and then either kept at 4C and stained with anti-mouse IgG secondary-PE conjugated Ab at 4C (B, total surface staining) or moved to 37C for various times and then stained with secondary Ab at 4C (C, residual surface staining). Panels B and C demonstrate that MAb X41.6 engagement of surface 161P2F10b causes internalization at 37C of the complex indicated by the progressive decrease in mean fluorescence intensity (MFI).

Figure 34. Internalization of selected anti-161P2F10b murine MAbs. Internalization of selected anti-161P2F10b MAbs are by flow cytometry are shown. Internalization is indicated by a decrease in the mean fluorescence intensity (MFI) of cells moved to 37C versus cells stained at 4C.

Figure 35. Antibody engagement of 161P2F10b results in its internalization. Internalization of the commercially available MAb 97A6, anti-CD203c, is shown by fluorescence microscopy following staining of Rat1-161P2F10b cells. The cells were incubated with CD203c-PE conjugated MAb at 4C, washed, and then moved to 37C for the indicated times and then examined by fluorescence microscopy. At 4C, the staining of the cells is cell surface (bright halo of fluorescence around individual cells). Upon moving to 37C, there is a gradual loss of the surface fluorescence, concomitant with capping of the MAb to punctate regions on the surface, followed by the appearance of punctate and diffuse intracellular fluorescence and a total loss of surface fluorescence.

Figure 36. Effects of X41.50 MAb-saporin toxin conjugate on Caki-161P2F10b cells. Shown is the morphology of Caki-161P2F10b cells that were treated with and without the indicated concentrations of the internalizing anti-161P2F10b MAb and an anti-mouse IgG-saporin toxin secondary Ab (2 ug/ml). Saporin is unable to enter cells efficiently on its own and must be internalized for its toxic mechanism (protein synthesis inhibition) to work. Cells were first incubated at 4C with X41.50 MAb to allow surface binding, then either media or the saporin-conjugated secondary Ab was added and the cells were incubated for 72 hours at 37C. Cells incubated with either media alone, X41.50 alone, or the secondary-saporin Ab alone had no effect on Caki-161P2F10b growth and morphology, exemplified by a viable confluent monolayer. However, cells incubated with X41.50 MAb (2 and 0.5 ug/ml) and the secondary saporin-conjugate exhibited signs of growth inhibition (did not reach confluency) and apoptosis (small round floating apoptotic cells above the attached cell layer). This demonstrates the utility of anti-161P2F10b MAbs drug/toxin conjugates as a therapeutic approach for 161P2F10b-expressing cancers and diseased tissues.

Figure 37. Internalization-mediated downregulation of 161P2F10 protein by MAb X41.50. Rat1-161P2F10b cells were incubated with and without 10 ug/ml of MAb X41.50 for 72 hours. Cells were washed, fixed, permeabilized, and stained with PE-conjugated CD203c MAb to monitor total 161P2F10b protein expression. The data shows a marked decrease in staining following treatment of the cells with X41.50, demonstrating downregulation of 161P2F10b protein.

Figure 38. Anti-161P2F10b MAbs downregulate surface 161P2F10b enzymatic activity. Rat1-161P2F10b cells were treated with and without various concentrations of the indicated MAbs for 48 hours and then assayed for surface enzymatic activity using p-n-TMB substrate. The data demonstrates that engagement and internalization of surface 161P2F10b by MAbs results in the concomitant loss of surface 161P2F10b enzymatic activity.

Figure 39. Characteristics of mouse 161P2F10b MAbs. Shown is a summary of various characteristics of MAbs that recognize 161P2F10b.

The relative affinity of the MAbs was determined by saturation binding ELISA using the recombinant Tag5-ECD protein. The K_d of the binding reaction was determined using a one-site binding non-linear regression analysis of the data using GraphPad Prism software version 3.02 (Graphpad Software, San Diego, CA).

Relative surface staining was determined using 10 ug/ml of each MAb on RAT1-161P2F10b cells.

Relative ability to internalize was also carried out on Rat1-161P2F10b cells comparing staining with 10 ug/ml of MAb at 4C versus residual staining following incubation at 37C for 30 minutes.

The ability of the MAbs to downregulate surface enzyme activity was determined by incubation of Rat1-161P2F10b cells with 10 ug/ml of each MAb for 72 hours then assaying surface enzyme activity with p-nTMP substrate.

Relative specific immunohistochemical staining (IHC) was determined using 10 ug/ml of each MAb on 161P2F10b-expressing frozen section kidney clear cell carcinoma samples.

The epitope family was determined by competition binding ELISA using the Tag5-ECD protein as target. Tag5-ECD ELISA coated wells were first incubated with or without 10 ug/ml of competitor MAb, washed, and then incubated with 1 ug/ml of HRP-labeled test MAb. MAb that compete for binding (reduction of the signal of the test MAb with prior incubation with competitor) must share the same or an overlapping epitope and are thus assigned to an epitope family. Of the MAbs listed, at least 2 epitope families are defined.

Figure 40. Surface staining of selected anti-161P2F10b MAbs. Specific binding of cell surface 161P2F10b was determined by incubation of Rat1-161P2F10 (dark line) and Rat1-neo control cells (light dotted line) with 10 ug/ml of each MAb for 1.5 hours at 4C. Cells were washed, incubated with goat-anti-mouse-PE conjugated secondary Ab, washed again, and analyzed by flow cytometry. Shown are examples of MAb derived from DNA-based immunization of mice with an FC-fusion of the ECD (X41.6, X41.15, X41.17, X41.29, X41.37, X41.50), also DNA-based immunization with Tag5-ECD, and

with Rat1-161P2F10b cells (the last data was generated using the respective hybridoma supernatant at a 1:50 dilution) was performed.

Figure 41. Anti-161P2F10b MAbs X41.6 and 97A6 (CD203c) do not cross-react with ENPP1. Conditioned media from 293T cells transfected with either Tag5-161P2F10b or ENPP1 His-tagged vectors was subjected to immunoprecipitation analysis using 5 ug of MAb X41.6, MAb 97A6 (CD203c), or anti-His pAb. Following washing of the immune complexes, phosphodiesterase activity was determined by the addition of p-nTMP substrate. Enzymatic activity is seen in anti-His immune complexes from both Tag5 161P2F10b and Tag5 ENPP1 media due to the presence of the His epitope in both proteins. However, enzymatic activity is seen only in the immune complexes of X41.6 and 97A6 from Tag5 161P2F10 conditioned media and not with Tag5 ENPP1 media. These data demonstrate that MAbs X41.6 and 97A6 (CD203c) do not crossreact with the homologous ectonucleotide pyrophosphatase/phosphodiesterase family member ENPP1.

Figure 42. Detection of 161P2F10b in the conditioned media of 161P2F10b-expressing cells. Supernatants of the indicated 161P2F10b-expressing and non-expressing cell lines were analyzed for shedding/secretion of 161P2F10b protein by a capture ELISA. The capture ELISA was made using a 161P2F10b-specific MAb as the bottom capture MAb (1 ug/well), and X41.29 as the top detection MAb (2 ug/ml), and an anti-mouse IgG2a-HRP secondary and tetramethylbenzidine as substrate for development. Recombinant 161P2F10b Tag5 ECD protein was used as a standard. 161P2F10b protein was detected in the media from 769 and Caki kidney cancer cells engineered to express 161P2F10b but not in the parental lines, indicating that 161P2F10b protein is shed or secreted. Shed/secreted 161P2F10b may exert its activity on cells in an autocrine/paracrine manner. In addition, shed/secreted 161P2F10b is useful as a diagnostic marker for 161P2F10b-expressing cancer and/or other 161P2F10b-expressing diseased tissues.

Figure 43. Detection of secreted 161P2F10B in the serum of mice bearing UGK3 human kidney cancer xenografts. SCID mice inoculated subcutaneously with UGK3 kidney cancer cells were monitored for tumor growth (1 dimensional tumor measurements) and 161P2F10b serum levels (by capture ELISA) over the indicated times. The data demonstrates that 161P2F10b serum levels increase as the tumor size increases. This demonstrates that 161P2F10b is shed/secreted from 161P2F10b-expressing tissues in vivo and further demonstrates the utility of an ELISA to monitor 161P2F10b as a diagnostic marker.

Figure 44: Detection of 161P2F10B protein by immunohistochemistry in kidney cancer patient specimens. Renal clear cell carcinoma tissue and its matched normal adjacent tissue as well as its metastatic cancer to lymph node were obtained from a kidney cancer patient. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with PE-labeled mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours (Figure 44 panels A-F), or isotype control antibody (Figure 44 panels G-I). The slides were washed three times in buffer, and either analyzed by fluorescence microscopy (Figure 44 panels A, B and C), or further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour (Figure 44 panels D, E, and F). The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy (Figure 44 panels D, E and F). The results showed strong expression of 161P2F10B in the renal carcinoma patient tissue (Figure 44 panels A and D) and the kidney cancer metastasis to lymph node tissue (Figure 44 panels C and F), but weakly in normal kidney (Figure 44 B and E). The expression was detected mostly around the cell membrane indicating that 161P2F10B is membrane associated in kidney cancer tissues. The weak expression detected in normal kidney was localized to the kidney tubules. The sections stained with the isotype control antibody were negative showing the specificity of the anti-ENPP3 antibody (Figure 44 panels G-I).

Figure 45: Expression of 161P2F10B in Human Patient Cancers by Western Blot. Cell lysates from kidney cancer tissues (KiCa), kidney cancer metastasis to lymph node (KiCa Met), as well as normal kidney (NK) were subjected to

Western analysis using an anti-161P2F10B mouse monoclonal antibody. Briefly, tissues (~25 µg total protein) were solubilized in SDS-PAGE sample buffer and separated on a 10-20% SDS-PAGE gel and transferred to nitrocellulose. Blots were blocked in Tris-buffered saline (TBS) + 3% non-fat milk and then probed with purified anti-161P2F10B antibody in TBS + 0.15% Tween-20 + 1% milk. Blots were then washed and incubated with a 1:4,000 dilution of anti-mouse IgG-HRP conjugated secondary antibody. Following washing, anti-161P2F10B immunoreactive bands were developed and visualized by enhanced chemiluminescence and exposure to autoradiographic film. The specific anti-161P2F10B immunoreactive bands represent a monomeric form of the 161P2F10B protein, which runs at approximately 130kDa. These results demonstrate that 161P2F10B is useful as a diagnostic and therapeutic target for kidney cancers, metastatic cancers and other such as those as listed in Table I and other human cancers that express 161P2F10B.

Figure 46: Expression of 161P2F10B in Human Xenograft Tissues by Western Blot. Cell lysates from kidney cancer xenograft (KiCa Xeno), kidney cancer metastasis to lymph node xenograft (Met Xeno), as well as normal kidney (NK) were subjected to Western analysis using an anti-161P2F10B mouse monoclonal antibody. Briefly, tissues (~25 µg total protein) were solubilized in SDS-PAGE sample buffer and separated on a 10-20% SDS-PAGE gel and transferred to nitrocellulose. Blots were blocked in Tris-buffered saline (TBS) + 3% non-fat milk and then probed with purified anti-161P2F10B antibody in TBS + 0.15% Tween-20 + 1% milk. Blots were then washed and incubated with a 1:4,000 dilution of anti-mouse IgG-HRP conjugated secondary antibody. Following washing, anti-161P2F10B immunoreactive bands were developed and visualized by enhanced chemiluminescence and exposure to autoradiographic film. The specific anti-161P2F10B immunoreactive bands represent a monomeric form of the 161P2F10B protein, which runs at approximately 130kDa, and a multimer of approximately 260kDa. These results demonstrate that the human cancer xenograft mouse models can be used to study the diagnostic and therapeutic effects of 161P2F10B.

DETAILED DESCRIPTION OF THE INVENTION

Outline of Sections

- I.) Definitions**
- II.) 161P2F10B Polynucleotides**
 - II.A.) Uses of 161P2F10B Polynucleotides**
 - II.A.1.) Monitoring of Genetic Abnormalities**
 - II.A.2.) Antisense Embodiments**
 - II.A.3.) Primers and Primer Pairs**
 - II.A.4.) Isolation of 161P2F10B-Encoding Nucleic Acid Molecules**
 - II.A.5.) Recombinant Nucleic Acid Molecules and Host-Vector Systems**
 - III.) 161P2F10B-related Proteins**
 - III.A.) Motif-bearing Protein Embodiments**
 - III.B.) Expression of 161P2F10B-related Proteins**
 - III.C.) Modifications of 161P2F10B-related Proteins**
 - III.D.) Uses of 161P2F10B-related Proteins**
- IV.) 161P2F10B Antibodies**
- V.) 161P2F10B Cellular Immune Responses**
- VI.) 161P2F10B Transgenic Animals**
- VII.) Methods for the Detection of 161P2F10B**
- VIII.) Methods for Monitoring the Status of 161P2F10B-related Genes and Their Products**
- IX.) Identification of Molecules That Interact With 161P2F10B**

- X.) Therapeutic Methods and Compositions
 - X.A.) Anti-Cancer Vaccines
 - X.B.) 161P2F10B as a Target for Antibody-Based Therapy
 - X.C.) 161P2F10B as a Target for Cellular Immune Responses
 - X.C.1. Minigene Vaccines
 - X.C.2. Combinations of CTL Peptides with Helper Peptides
 - X.C.3. Combinations of CTL Peptides with T Cell Priming Agents
 - X.C.4. Vaccine Compositions Comprising DC Pulsed with CTL and/or HTL Peptides
 - X.D.) Adoptive Immunotherapy
 - X.E.) Administration of Vaccines for Therapeutic or Prophylactic Purposes
- XI.) Diagnostic and Prognostic Embodiments of 161P2F10B.
- XII.) Inhibition of 161P2F10B Protein Function
 - XII.A.) Inhibition of 161P2F10B With Intracellular Antibodies
 - XII.B.) Inhibition of 161P2F10B with Recombinant Proteins
 - XII.C.) Inhibition of 161P2F10B Transcription or Translation
 - XII.D.) General Considerations for Therapeutic Strategies
- XIII.) Identification, Characterization and Use of Modulators of 161P2F10b
- XIV.) KITS/Articles of Manufacture

I.) Definitions:

Unless otherwise defined, all terms of art, notations and other scientific terms or terminology used herein are intended to have the meanings commonly understood by those of skill in the art to which this invention pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a substantial difference over what is generally understood in the art. Many of the techniques and procedures described or referenced herein are well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized molecular cloning methodologies described in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2nd. edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. As appropriate, procedures involving the use of commercially available kits and reagents are generally carried out in accordance with manufacturer defined protocols and/or parameters unless otherwise noted.

The terms "advanced prostate cancer", "locally advanced prostate cancer", "advanced disease" and "locally advanced disease" mean prostate cancers that have extended through the prostate capsule, and are meant to include stage C disease under the American Urological Association (AUA) system, stage C1 - C2 disease under the Whitmore-Jewett system, and stage T3 - T4 and N+ disease under the TNM (tumor, node, metastasis) system. In general, surgery is not recommended for patients with locally advanced disease, and these patients have substantially less favorable outcomes compared to patients having clinically localized (organ-confined) prostate cancer. Locally advanced disease is clinically identified by palpable evidence of induration beyond the lateral border of the prostate, or asymmetry or induration above the prostate base. Locally advanced prostate cancer is presently diagnosed pathologically following radical prostatectomy if the tumor invades or penetrates the prostatic capsule, extends into the surgical margin, or invades the seminal vesicles.

"Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence 161P2F10B (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not

present in the native sequence 161P2F10B. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

The term "analog" refers to a molecule which is structurally similar or shares similar or corresponding attributes with another molecule (e.g. a 161P2F10B-related protein). For example, an analog of a 161P2F10B protein can be specifically bound by an antibody or T cell that specifically binds to 161P2F10B.

The term "antibody" is used in the broadest sense. Therefore, an "antibody" can be naturally occurring or man-made such as monoclonal antibodies produced by conventional hybridoma technology. Anti-161P2F10B antibodies comprise monoclonal and polyclonal antibodies as well as fragments containing the antigen-binding domain and/or one or more complementarity determining regions of these antibodies.

An "antibody fragment" is defined as at least a portion of the variable region of the immunoglobulin molecule that binds to its target, i.e., the antigen-binding region. In one embodiment it specifically covers single anti-161P2F10B antibodies and clones thereof (including agonist, antagonist and neutralizing antibodies) and anti-161P2F10B antibody compositions with polyepitopic specificity.

The term "codon optimized sequences" refers to nucleotide sequences that have been optimized for a particular host species by replacing any codons having a usage frequency of less than about 20%. Nucleotide sequences that have been optimized for expression in a given host species by elimination of spurious polyadenylation sequences, elimination of exon/intron splicing signals, elimination of transposon-like repeats and/or optimization of GC content in addition to codon optimization are referred to herein as an "expression enhanced sequences."

A "combinatorial library" is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library, such as a polypeptide (e.g., mutein) library, is formed by combining a set of chemical building blocks called amino acids in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Numerous chemical compounds are synthesized through such combinatorial mixing of chemical building blocks (Gallop et al., J. Med. Chem. 37(9): 1233-1251 (1994)).

Preparation and screening of combinatorial libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Patent No. 5,010,175, Furka, Pept. Prot. Res. 37:487-493 (1991), Houghton et al., Nature, 354:84-88 (1991)), peptoids (PCT Publication No WO 91/19735), encoded peptides (PCT Publication WO 93/20242), random bio- oligomers (PCT Publication WO 92/00091), benzodiazepines (U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs et al., Proc. Nat. Acad. Sci. USA 90:6909-6913 (1993)), vinylogous polypeptides (Hagihara et al., J. Amer. Chem. Soc. 114:6568 (1992)), nonpeptidic peptidomimetics with a Beta-D-Glucose scaffolding (Hirschmann et al., J. Amer. Chem. Soc. 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen et al., J. Amer. Chem. Soc. 116:2661 (1994)), oligocarbarnates (Cho, et al., Science 261:1303 (1993)), and/or peptidyl phosphonates (Campbell et al., J. Org. Chem. 59:658 (1994)). See, generally, Gordon et al., J. Med. Chem. 37:1385 (1994), nucleic acid libraries (see, e.g., Stratagene, Corp.), peptide nucleic acid libraries (see, e.g., U.S. Patent 5,539,083), antibody libraries (see, e.g., Vaughn et al., Nature Biotechnology 14(3): 309-314 (1996), and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al., Science 274:1520-1522 (1996), and U.S. Patent No. 5,593,853), and small organic molecule libraries (see, e.g., benzodiazepines, Baum, C&EN, Jan 18, page 33 (1993); isoprenoids, U.S. Patent No. 5,569,588; thiazolidinones and metathiazanones, U.S. Patent No. 5,549,974; pyrrolidines, U.S. Patent Nos. 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent No. 5,506, 337; benzodiazepines, U.S. Patent No. 5,288,514; and the like).

Devices for the preparation of combinatorial libraries are commercially available (see, e.g., 357 NIPS, 390 NIPS, Advanced Chem Tech, Louisville KY; Symphony, Rainin, Woburn, MA; 433A, Applied Biosystems, Foster City, CA; 9050,

Plus, Millipore, Bedford, NIA). A number of well-known robotic systems have also been developed for solution phase chemistries. These systems include automated workstations such as the automated synthesis apparatus developed by Takeda Chemical Industries, LTD. (Osaka, Japan) and many robotic systems utilizing robotic arms (Zymate H, Zymark Corporation, Hopkinton, Mass.; Orca, Hewlett-Packard, Palo Alto, Calif.), which mimic the manual synthetic operations performed by a chemist. Any of the above devices are suitable for use with the present invention. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art. In addition, numerous combinatorial libraries are themselves commercially available (see, e.g., ComGenex, Princeton, NJ; Asinex, Moscow, RU; Tripos, Inc., St. Louis, MO; ChemStar, Ltd, Moscow, RU; 3D Pharmaceuticals, Exton, PA; Martek Biosciences, Columbia, MD; etc.).

The term "cytotoxic agent" refers to a substance that inhibits or prevents the expression activity of cells, function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof. Examples of cytotoxic agents include, but are not limited to auristatins, auromycins, maytansinoids, yttrium, bismuth, ricin, ricin A-chain, combrestatin, duocarmycins, dolostatins, doxorubicin, daunorubicin, taxol, cisplatin, cc1065, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha-sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, *Saponaia officinalis* inhibitor, and glucocorticoid and other chemotherapeutic agents, as well as radioisotopes such as At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹² or ²¹³, P³² and radioactive isotopes of Lu including Lu¹⁷⁷. Antibodies may also be conjugated to an anti-cancer pro-drug activating enzyme capable of converting the pro-drug to its active form.

The "gene product" is sometimes referred to herein as a protein or mRNA. For example, a "gene product of the invention" is sometimes referred to herein as a "cancer amino acid sequence", "cancer protein", "protein of a cancer listed in Table I", a "cancer mRNA", "mRNA of a cancer listed in Table I", etc. In one embodiment, the cancer protein is encoded by a nucleic acid of Figure 2. The cancer protein can be a fragment, or alternatively, be the full-length protein to the fragment encoded by the nucleic acids of Figure 2. In one embodiment, a cancer amino acid sequence is used to determine sequence identity or similarity. In another embodiment, the sequences are naturally occurring allelic variants of a protein encoded by a nucleic acid of Figure 2. In another embodiment, the sequences are sequence variants as further described herein.

"High throughput screening" assays for the presence, absence, quantification, or other properties of particular nucleic acids or protein products are well known to those of skill in the art. Similarly, binding assays and reporter gene assays are similarly well known. Thus, e.g., U.S. Patent No. 5,559,410 discloses high throughput screening methods for proteins; U.S. Patent No. 5,585,639 discloses high throughput screening methods for nucleic acid binding (i.e., in arrays); while U.S. Patent Nos. 5,576,220 and 5,541,061 disclose high throughput methods of screening for ligand/antibody binding.

In addition, high throughput screening systems are commercially available (see, e.g., Amersham Biosciences, Piscataway, NJ; Zymark Corp., Hopkinton, MA; Air Technical Industries, Mentor, OH; Beckman Instruments, Inc. Fullerton, CA; Precision Systems, Inc., Natick, MA; etc.). These systems typically automate entire procedures, including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. The manufacturers of such systems provide detailed protocols for various high throughput systems. Thus, e.g., Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like.

The term "homolog" refers to a molecule which exhibits homology to another molecule, by for example, having sequences of chemical residues that are the same or similar at corresponding positions.

"Human Leukocyte Antigen" or "HLA" is a human class I or class II Major Histocompatibility Complex (MHC) protein (see, e.g., Stites, *et al.*, IMMUNOLOGY, 8TH ED., Lange Publishing, Los Altos, CA (1994).

The terms "hybridize", "hybridizing", "hybridizes" and the like, used in the context of polynucleotides, are meant to refer to conventional hybridization conditions, preferably such as hybridization in 50% formamide/6XSSC/0.1% SDS/100 µg/ml ssDNA, in which temperatures for hybridization are above 37 degrees C and temperatures for washing in 0.1XSSC/0.1% SDS are above 55 degrees C.

The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany the material as it is found in its native state. Thus, isolated peptides in accordance with the invention preferably do not contain materials normally associated with the peptides in their *in situ* environment. For example, a polynucleotide is said to be "isolated" when it is substantially separated from contaminant polynucleotides that correspond or are complementary to genes other than the 161P2F10B genes or that encode polypeptides other than 161P2F10B gene product or fragments thereof. A skilled artisan can readily employ nucleic acid isolation procedures to obtain an isolated 161P2F10B polynucleotide. A protein is said to be "isolated," for example, when physical, mechanical or chemical methods are employed to remove the 161P2F10B proteins from cellular constituents that are normally associated with the protein. A skilled artisan can readily employ standard purification methods to obtain an isolated 161P2F10B protein. Alternatively, an isolated protein can be prepared by chemical means.

The term "mammal" refers to any organism classified as a mammal, including mice, rats, rabbits, dogs, cats, cows, horses and humans. In one embodiment of the invention, the mammal is a mouse. In another embodiment of the invention, the mammal is a human.

The terms "metastatic prostate cancer" and "metastatic disease" mean prostate cancers that have spread to regional lymph nodes or to distant sites, and are meant to include stage D disease under the AUA system and stage TxNxM+ under the TNM system. As is the case with locally advanced prostate cancer, surgery is generally not indicated for patients with metastatic disease, and hormonal (androgen ablation) therapy is a preferred treatment modality. Patients with metastatic prostate cancer eventually develop an androgen-refractory state within 12 to 18 months of treatment initiation. Approximately half of these androgen-refractory patients die within 6 months after developing that status. The most common site for prostate cancer metastasis is bone. Prostate cancer bone metastases are often osteoblastic rather than osteolytic (i.e., resulting in net bone formation). Bone metastases are found most frequently in the spine, followed by the femur, pelvis, rib cage, skull and humerus. Other common sites for metastasis include lymph nodes, lung, liver and brain. Metastatic prostate cancer is typically diagnosed by open or laparoscopic pelvic lymphadenectomy, whole body radionuclide scans, skeletal radiography, and/or bone lesion biopsy.

The term "modulator" or "test compound" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for the capacity to directly or indirectly alter the cancer phenotype or the expression of a cancer sequence, e.g., a nucleic acid or protein sequences, or effects of cancer sequences (e.g., signaling, gene expression, protein interaction, etc.) In one aspect, a modulator will neutralize the effect of a cancer protein of the invention. By "neutralize" is meant that an activity of a protein is inhibited or blocked, along with the consequent effect on the cell. In another aspect, a modulator will neutralize the effect of a gene, and its corresponding protein, of the invention by normalizing levels of said protein. In preferred embodiments, modulators alter expression profiles, or expression profile nucleic acids or proteins provided herein, or downstream effector pathways. In one embodiment, the modulator suppresses a cancer phenotype, e.g. to a normal tissue fingerprint. In another embodiment, a modulator induced a cancer phenotype. Generally, a plurality of assay mixtures is run in parallel with

different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e., at zero concentration or below the level of detection.

Modulators, drug candidates or test compounds encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 Daltons. Preferred small molecules are less than 2000, or less than 1500 or less than 1000 or less than 500 D. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Modulators also comprise biomolecules such as peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides. One class of modulators are peptides, for example of from about five to about 35 amino acids, with from about five to about 20 amino acids being preferred, and from about 7 to about 15 being particularly preferred. Preferably, the cancer modulatory protein is soluble, includes a non-transmembrane region, and/or, has an N-terminal Cys to aid in solubility. In one embodiment, the C-terminus of the fragment is kept as a free acid and the N-terminus is a free amine to aid in coupling, i.e., to cysteine. In one embodiment, a cancer protein of the invention is conjugated to an immunogenic agent as discussed herein. In one embodiment, the cancer protein is conjugated to BSA. The peptides of the invention, e.g., of preferred lengths, can be linked to each other or to other amino acids to create a longer peptide/protein. The modulatory peptides can be digests of naturally occurring proteins as is outlined above, random peptides, or "biased" random peptides. In a preferred embodiment, peptide/protein-based modulators are antibodies, and fragments thereof, as defined herein.

Modulators of cancer can also be nucleic acids. Nucleic acid modulating agents can be naturally occurring nucleic acids, random nucleic acids, or "biased" random nucleic acids. For example, digests of prokaryotic or eukaryotic genomes can be used in an approach analogous to that outlined above for proteins.

The term "monoclonal antibody" refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the antibodies comprising the population are identical except for possible naturally occurring mutations that are present in minor amounts.

A "motif", as in biological motif of a 161P2F10B-related protein, refers to any pattern of amino acids forming part of the primary sequence of a protein, that is associated with a particular function (e.g. protein-protein interaction, protein-DNA interaction, etc) or modification (e.g. that is phosphorylated, glycosylated or amidated), or localization (e.g. secretory sequence, nuclear localization sequence, etc.) or a sequence that is correlated with being immunogenic, either humorally or cellularly. A motif can be either contiguous or capable of being aligned to certain positions that are generally correlated with a certain function or property. In the context of HLA motifs, "motif" refers to the pattern of residues in a peptide of defined length, usually a peptide of from about 8 to about 13 amino acids for a class I HLA motif and from about 6 to about 25 amino acids for a class II HLA motif, which is recognized by a particular HLA molecule. Peptide motifs for HLA binding are typically different for each protein encoded by each human HLA allele and differ in the pattern of the primary and secondary anchor residues.

A "pharmaceutical excipient" comprises a material such as an adjuvant, a carrier, pH-adjusting and buffering agents, tonicity adjusting agents, wetting agents, preservative, and the like.

"Pharmaceutically acceptable" refers to a non-toxic, inert, and/or composition that is physiologically compatible with humans or other mammals.

The term "polynucleotide" means a polymeric form of nucleotides of at least 10 bases or base pairs in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide, and is meant to include single and double

stranded forms of DNA and/or RNA. In the art, this term is often used interchangeably with "oligonucleotide". A polynucleotide can comprise a nucleotide sequence disclosed herein wherein thymidine (T), as shown for example in Figure 2, can also be uracil (U); this definition pertains to the differences between the chemical structures of DNA and RNA, in particular the observation that one of the four major bases in RNA is uracil (U) instead of thymidine (T).

The term "polypeptide" means a polymer of at least about 4, 5, 6, 7, or 8 amino acids. Throughout the specification, standard three letter or single letter designations for amino acids are used. In the art, this term is often used interchangeably with "peptide" or "protein".

An HLA "primary anchor residue" is an amino acid at a specific position along a peptide sequence which is understood to provide a contact point between the immunogenic peptide and the HLA molecule. One to three, usually two, primary anchor residues within a peptide of defined length generally defines a "motif" for an immunogenic peptide. These residues are understood to fit in close contact with peptide binding groove of an HLA molecule, with their side chains buried in specific pockets of the binding groove. In one embodiment, for example, the primary anchor residues for an HLA class I molecule are located at position 2 (from the amino terminal position) and at the carboxyl terminal position of a 8, 9, 10, 11, or 12 residue peptide epitope in accordance with the invention. Alternatively, in another embodiment, the primary anchor residues of a peptide binds an HLA class II molecule are spaced relative to each other, rather than to the termini of a peptide, where the peptide is generally of at least 9 amino acids in length. The primary anchor positions for each motif and supermotif are set forth in Table IV. For example, analog peptides can be created by altering the presence or absence of particular residues in the primary and/or secondary anchor positions shown in Table IV. Such analogs are used to modulate the binding affinity and/or population coverage of a peptide comprising a particular HLA motif or supermotif.

"Radioisotopes" include, but are not limited to the following (non-limiting exemplary uses are also set forth):

Examples of Medical Isotopes:

Isotope

Description of use

Actinium-225

(Ac-225)

See Thorium-229 (Th-229)

Actinium-227

(Ac-227)

Parent of Radium-223 (Ra-223) which is an alpha emitter used to treat metastases in the skeleton resulting from cancer (i.e., breast and prostate cancers), and cancer radioimmunotherapy

Bismuth-212

(Bi-212)

See Thorium-228 (Th-228)

Bismuth-213

(Bi-213)

See Thorium-229 (Th-229)

Cadmium-109

(Cd-109)

Cancer detection

Cobalt-60

(Co-60)

Radiation source for radiotherapy of cancer, for food irradiators, and for sterilization of medical supplies

Copper-64

(Cu-64)

A positron emitter used for cancer therapy and SPECT imaging

Copper-67

(Cu-67)

Beta/gamma emitter used in cancer radioimmunotherapy and diagnostic studies (i.e., breast and colon cancers, and lymphoma)

Dysprosium-166

(Dy-166)

Cancer radioimmunotherapy

Erbium-169

(Er-169)

Rheumatoid arthritis treatment, particularly for the small joints associated with fingers and toes

Europium-152

(Eu-152)

Radiation source for food irradiation and for sterilization of medical supplies

Europium-154

(Eu-154)

Radiation source for food irradiation and for sterilization of medical supplies

Gadolinium-153

(Gd-153)

Osteoporosis detection and nuclear medical quality assurance devices

Gold-198

(Au-198)

Implant and intracavity therapy of ovarian, prostate, and brain cancers

Holmium-166

(Ho-166)

Multiple myeloma treatment in targeted skeletal therapy, cancer radioimmunotherapy, bone marrow ablation, and rheumatoid arthritis treatment

Iodine-125

(I-125)

Osteoporosis detection, diagnostic imaging, tracer drugs, brain cancer treatment, radiolabeling, tumor imaging, mapping of receptors in the brain, interstitial radiation therapy, brachytherapy for treatment of prostate cancer, determination of glomerular filtration rate (GFR), determination of plasma volume, detection of deep vein thrombosis of the legs

Iodine-131

(I-131)

Thyroid function evaluation, thyroid disease detection, treatment of thyroid cancer as well as other non-malignant thyroid diseases (i.e., Graves disease, goiters, and hyperthyroidism), treatment of leukemia, lymphoma, and other forms of cancer (e.g., breast cancer) using radioimmunotherapy

Iridium-192

(Ir-192)

Brachytherapy, brain and spinal cord tumor treatment, treatment of blocked arteries (i.e., arteriosclerosis and restenosis), and implants for breast and prostate tumors

Lutetium-177

(Lu-177)

Cancer radioimmunotherapy and treatment of blocked arteries (i.e., arteriosclerosis and restenosis)

Molybdenum-99

(Mo-99)

Parent of Technetium-99m (Tc-99m) which is used for imaging the brain, liver, lungs, heart, and other organs. Currently, Tc-99m is the most widely used radioisotope used for diagnostic imaging of various cancers and diseases involving the brain, heart, liver, lungs; also used in detection of deep vein thrombosis of the legs

Osmium-194

(Os-194)

Cancer radioimmunotherapy

Palladium-103

(Pd-103)

Prostate cancer treatment

Platinum-195m

(Pt-195m)

Studies on biodistribution and metabolism of cisplatin, a chemotherapeutic drug

Phosphorus-32

(P-32)

Polycythemia rubra vera (blood cell disease) and leukemia treatment, bone cancer diagnosis/treatment; colon, pancreatic, and liver cancer treatment; radiolabeling nucleic acids for in vitro research, diagnosis of superficial tumors, treatment of blocked arteries (i.e., arteriosclerosis and restenosis), and intracavity therapy

Phosphorus-33

(P-33)

Leukemia treatment, bone disease diagnosis/treatment, radiolabeling, and treatment of blocked arteries (i.e., arteriosclerosis and restenosis)

Radium-223

(Ra-223)

See Actinium-227 (Ac-227)

Rhenium-186

(Re-186)

Bone cancer pain relief, rheumatoid arthritis treatment, and diagnosis and treatment of lymphoma and bone, breast, colon, and liver cancers using radioimmunotherapy

Rhenium-188

(Re-188)

Cancer diagnosis and treatment using radioimmunotherapy, bone cancer pain relief, treatment of rheumatoid arthritis, and treatment of prostate cancer

Rhodium-105

(Rh-105)

Cancer radioimmunotherapy

Samarium-145

(Sm-145)

Ocular cancer treatment

Samarium-153

(Sm-153)

Cancer radioimmunotherapy and bone cancer pain relief

Scandium-47

(Sc-47)

Cancer radioimmunotherapy and bone cancer pain relief

Selenium-75

(Se-75)

Radiotracer used in brain studies, imaging of adrenal cortex by gamma-scintigraphy, lateral locations of steroid secreting tumors, pancreatic scanning, detection of hyperactive parathyroid glands, measure rate of bile acid loss from the endogenous pool

Strontium-85

(Sr-85)

Bone cancer detection and brain scans

Strontium-89

(Sr-89)

Bone cancer pain relief, multiple myeloma treatment, and osteoblastic therapy

Technetium-99m

(Tc-99m)

See Molybdenum-99 (Mo-99)

Thorium-228

(Th-228)

Parent of Bismuth-212 (Bi-212) which is an alpha emitter used in cancer radioimmunotherapy

Thorium-229

(Th-229)

Parent of Actinium-225 (Ac-225) and grandparent of Bismuth-213 (Bi-213) which are alpha emitters used in cancer radioimmunotherapy

Thulium-170

(Tm-170)

Gamma source for blood irradiators, energy source for implanted medical devices

Tin-117m

(Sn-117m)

Cancer immunotherapy and bone cancer pain relief

Tungsten-188

(W-188)

Parent for Rhenium-188 (Re-188) which is used for cancer diagnostics/treatment, bone cancer pain relief, rheumatoid arthritis treatment, and treatment of blocked arteries (i.e., arteriosclerosis and restenosis)

Xenon-127

(Xe-127)

Neuroimaging of brain disorders, high resolution SPECT studies, pulmonary function tests, and cerebral blood flow studies

Ytterbium-175

(Yb-175)

Cancer radioimmunotherapy

Yttrium-90

(Y-90)

Microseeds obtained from irradiating Yttrium-89 (Y-89) for liver cancer treatment

Yttrium-91

(Y-91)

A gamma-emitting label for Yttrium-90 (Y-90) which is used for cancer radioimmunotherapy (i.e., lymphoma, breast, colon, kidney, lung, ovarian, prostate, pancreatic, and inoperable liver cancers)

By "randomized" or grammatical equivalents as herein applied to nucleic acids and proteins is meant that each nucleic acid and peptide consists of essentially random nucleotides and amino acids, respectively. These random peptides (or nucleic acids, discussed herein) can incorporate any nucleotide or amino acid at any position. The synthetic process can be designed to generate randomized proteins or nucleic acids, to allow the formation of all or most of the possible combinations over the length of the sequence, thus forming a library of randomized candidate bioactive proteinaceous agents.

In one embodiment, a library is "fully randomized," with no sequence preferences or constants at any position. In another embodiment, the library is a "biased random" library. That is, some positions within the sequence either are held constant, or are selected from a limited number of possibilities. For example, the nucleotides or amino acid residues are randomized within a defined class, e.g., of hydrophobic amino acids, hydrophilic residues, sterically biased (either small or large) residues, towards the creation of nucleic acid binding domains, the creation of cysteines, for cross-linking, prolines for SH-3 domains, serines, threonines, tyrosines or histidines for phosphorylation sites, etc., or to purines, etc.

A "recombinant" DNA or RNA molecule is a DNA or RNA molecule that has been subjected to molecular manipulation *in vitro*.

Non-limiting examples of small molecules include compounds that bind or interact with 161P2F10B, ligands including hormones, neuropeptides, chemokines, odorants, phospholipids, and functional equivalents thereof that bind and preferably inhibit 161P2F10B protein function. Such non-limiting small molecules preferably have a molecular weight of less than about 10 kDa, more preferably below about 9, about 8, about 7, about 6, about 5 or about 4 kDa. In certain embodiments, small molecules physically associate with, or bind, 161P2F10B protein; are not found in naturally occurring metabolic pathways; and/or are more soluble in aqueous than non-aqueous solutions

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured nucleic acid sequences to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel *et al.*, Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, are identified by, but not limited to, those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42 °C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42 °C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55 °C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55 °C. "Moderately stringent conditions" are described by, but not limited to, those in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

An HLA "supermotif" is a peptide binding specificity shared by HLA molecules encoded by two or more HLA alleles. Overall phenotypic frequencies of HLA-supertypes in different ethnic populations are set forth in Table IV (F). The non-limiting constituents of various supetypes are as follows:

A2: A*0201, A*0202, A*0203, A*0204, A*0205, A*0206, A*6802, A*6901, A*0207

A3: A3, A11, A31, A*3301, A*6801, A*0301, A*1101, A*3101

B7: B7, B*3501-03, B*51, B*5301, B*5401, B*5501, B*5502, B*5601, B*6701, B*7801, B*0702, B*5101, B*5602

B44: B*3701, B*4402, B*4403, B*60 (B*4001), B61 (B*4006)

A1: A*0102, A*2604, A*3601, A*4301, A*8001

A24: A*24, A*30, A*2403, A*2404, A*3002, A*3003

B27: B*1401-02, B*1503, B*1509, B*1510, B*1518, B*3801-02, B*3901, B*3902, B*3903-04, B*4801-02, B*7301, B*2701-08

B58: B*1516, B*1517, B*5701, B*5702, B58

B62: B*4601, B52, B*1501 (B62), B*1502 (B75), B*1513 (B77)

Calculated population coverage afforded by different HLA-supertype combinations are set forth in Table IV (G).

As used herein "to treat" or "therapeutic" and grammatically related terms, refer to any improvement of any consequence of disease, such as prolonged survival, less morbidity, and/or a lessening of side effects which are the byproducts of an alternative therapeutic modality; full eradication of disease is not required.

A "transgenic animal" (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A "transgene" is a DNA that is integrated into the genome of a cell from which a transgenic animal develops.

As used herein, an HLA or cellular immune response "vaccine" is a composition that contains or encodes one or more peptides of the invention. There are numerous embodiments of such vaccines, such as a cocktail of one or more individual peptides; one or more peptides of the invention comprised by a polyepitopic peptide; or nucleic acids that encode such individual peptides or polypeptides, e.g., a minigene that encodes a polyepitopic peptide. The "one or more peptides" can include any whole unit integer from 1-150 or more, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, or 150 or more peptides of the invention. The peptides or polypeptides can optionally be modified, such as by lipidation, addition of targeting or other sequences. HLA class I peptides of the invention can be admixed with, or linked to, HLA class II peptides, to facilitate activation of both cytotoxic T lymphocytes and helper T lymphocytes. HLA vaccines can also comprise peptide-pulsed antigen presenting cells, e.g., dendritic cells.

The term "variant" refers to a molecule that exhibits a variation from a described type or norm, such as a protein that has one or more different amino acid residues in the corresponding position(s) of a specifically described protein (e.g. the 161P2F10B protein shown in Figure 2 or Figure 3. An analog is an example of a variant protein. Splice isoforms and single nucleotides polymorphisms (SNPs) are further examples of variants.

The "161P2F10B-related proteins" of the invention include those specifically identified herein, as well as allelic variants, conservative substitution variants, analogs and homologs that can be isolated/generated and characterized without undue experimentation following the methods outlined herein or readily available in the art. Fusion proteins that combine parts of different 161P2F10B proteins or fragments thereof, as well as fusion proteins of a 161P2F10B protein and a heterologous polypeptide are also included. Such 161P2F10B proteins are collectively referred to as the 161P2F10B-related proteins, the proteins of the invention, or 161P2F10B. The term "161P2F10B-related protein" refers to a polypeptide fragment or a 161P2F10B protein sequence of 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more than 25 amino acids; or, at least 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, or 664 or more amino acids.

II.) 161P2F10B Polynucleotides

One aspect of the invention provides polynucleotides corresponding or complementary to all or part of a 161P2F10B gene, mRNA, and/or coding sequence, preferably in isolated form, including polynucleotides encoding a 161P2F10B-related protein and fragments thereof, DNA, RNA, DNA/RNA hybrid, and related molecules, polynucleotides or oligonucleotides complementary to a 161P2F10B gene or mRNA sequence or a part thereof, and polynucleotides or oligonucleotides that hybridize to a 161P2F10B gene, mRNA, or to a 161P2F10B encoding polynucleotide (collectively, "161P2F10B polynucleotides"). In all instances when referred to in this section, T can also be U in Figure 2.

Embodiments of a 161P2F10B polynucleotide include: a 161P2F10B polynucleotide having the sequence shown in Figure 2, the nucleotide sequence of 161P2F10B as shown in Figure 2 wherein T is U; at least 10 contiguous nucleotides of a polynucleotide having the sequence as shown in Figure 2; or, at least 10 contiguous nucleotides of a polynucleotide having the sequence as shown in Figure 2 where T is U. For example, embodiments of 161P2F10B nucleotides comprise, without limitation:

- (I) a polynucleotide comprising, consisting essentially of, or consisting of a sequence as shown in Figure 2, wherein T can also be U;
- (II) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2A, from nucleotide residue number 44 through nucleotide residue number 2671, including the stop codon, wherein T can also be U;
- (III) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2B, from nucleotide residue number 44 through nucleotide residue number 2671, including the stop codon, wherein T can also be U;
- (IV) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2C, from nucleotide residue number 44 through nucleotide residue number 2671, including the a stop codon, wherein T can also be U;
- (V) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2D, from nucleotide residue number 44 through nucleotide residue number 2671, including the stop codon, wherein T can also be U;
- (VI) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2E, from nucleotide residue number 44 through nucleotide residue number 2671, including the stop codon, wherein T can also be U;
- (VII) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2F, from nucleotide residue number 84 through nucleotide residue number 2711, including the stop codon, wherein T can also be U;
- (VIII) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in Figure 2G, from nucleotide residue number 276 through nucleotide residue number 2801, including the stop codon, wherein T can also be U;

- (IX) a polynucleotide that encodes a 161P2F10B-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% homologous to an entire amino acid sequence shown in Figure 2A-G;
- (X) a polynucleotide that encodes a 161P2F10B-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% identical to an entire amino acid sequence shown in Figure 2A-G;
- (XI) a polynucleotide that encodes at least one peptide set forth in Tables VIII-XXI and XXII-XLIX;
- (XII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3A-D in any whole number increment up to 875 that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of Figure 5;
- (XIII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3A-D in any whole number increment up to 875 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydropathicity profile of Figure 6;
- (XIV) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3A-D in any whole number increment up to 875 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of Figure 7;
- (XV) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3A-DF in any whole number increment up to 875 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of Figure 8;
- (XVI) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3A-D in any whole number increment up to 875 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of Figure 9;
- (XVII) a polynucleotide that encodes a peptide-region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3E in any whole number increment up to 841 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of Figure 5;

(XVIII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3E in any whole number increment up to 841 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydropathicity profile of Figure 6;

(XIX) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3E in any whole number increment up to 841 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of Figure 7;

(XX) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3E in any whole number increment up to 841 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of Figure 8;

(XXI) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of Figure 3E in any whole number increment up to 841 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of Figure 9

(XXII) a polynucleotide that encodes monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)15 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

XXIII) a polynucleotide that encodes monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)29 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXIV) a polynucleotide that encodes monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)37 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXV) a polynucleotide that encodes monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(4)6 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXVI) a polynucleotide that encodes monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)17 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXVII) a polynucleotide that encodes monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)50 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXVIII) a polynucleotide that is fully complementary to a polynucleotide of any one of (I)-(XXVII).

(XXIX) a peptide that is encoded by any of (I) to (XXVII); and

(XXX) a composition comprising a polynucleotide of any of (I)-(XXVII) or peptide of (XXIX) together with a pharmaceutical excipient and/or in a human unit dose form.

(XXXI) a method of using a polynucleotide of any (I)-(XXVII) or peptide of (XXIX) or a composition of (XXX) in a method to modulate a cell expressing 161P2F10b,

(XXXII) a method of using a polynucleotide of any (I)-(XXVII) or peptide of (XXIX) or a composition of (XXX) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 161P2F10b

(XXXIII) a method of using a polynucleotide of any (I)-(XXVII) or peptide of (XXIX) or a composition of (XXX) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 161P2F10b, said cell from a cancer of a tissue listed in Table I;

(XXXIV) a method of using a polynucleotide of any (I)-(XLII) or peptide of (XXIX) or a composition of (XXX) in a method to diagnose, prophylax, prognose, or treat a a cancer;

(XXXV) a method of using a polynucleotide of any (I)-(XLII) or peptide of (XXIX) or a composition of (XXX) in a method to diagnose, prophylax, prognose, or treat a a cancer of a tissue listed in Table I; and,

(XXXVI) a method of using a polynucleotide of any (I)-(XLII) or peptide of (XXIX) or a composition of (XXX) in a method to identify or characterize a modulator of a cell expressing 161P2F10b.

As used herein, a range is understood to disclose specifically all whole unit positions thereof.

Typical embodiments of the invention disclosed herein include 161P2F10B polynucleotides that encode specific portions of 161P2F10B mRNA sequences (and those which are complementary to such sequences) such as those that encode the proteins and/or fragments thereof, for example: .

(a) 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 860, 870, 875 or more contiguous amino acids of 161P2F10B variant 1; the maximal lengths relevant for other variants are: variant 2, 875 amino acids; variant 3, 875 amino acids, variant 4, 875 amino acids, and variant 7, 841 amino acids.

For example, representative embodiments of the invention disclosed herein include: polynucleotides and their encoded peptides themselves encoding about amino acid 1 to about amino acid 10 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 10 to about amino acid 20 of the 161P2F10B protein shown

in Figure 2 or Figure 3, polynucleotides encoding about amino acid 20 to about amino acid 30 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 30 to about amino acid 40 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 40 to about amino acid 50 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 50 to about amino acid 60 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 60 to about amino acid 70 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 70 to about amino acid 80 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 80 to about amino acid 90 of the 161P2F10B protein shown in Figure 2 or Figure 3, polynucleotides encoding about amino acid 90 to about amino acid 100 of the 161P2F10B protein shown in Figure 2 or Figure 3, in increments of about 10 amino acids, ending at the carboxyl terminal amino acid set forth in Figure 2 or Figure 3. Accordingly, polynucleotides encoding portions of the amino acid sequence (of about 10 amino acids), of amino acids, 100 through the carboxyl terminal amino acid of the 161P2F10B protein are embodiments of the invention. Wherein it is understood that each particular amino acid position discloses that position plus or minus five amino acid residues.

Polynucleotides encoding relatively long portions of a 161P2F10B protein are also within the scope of the invention. For example, polynucleotides encoding from about amino acid 1 (or 20 or 30 or 40 etc.) to about amino acid 20, (or 30, or 40 or 50 etc.) of the 161P2F10B protein "or variant" shown in Figure 2 or Figure 3 can be generated by a variety of techniques well known in the art. These polynucleotide fragments can include any portion of the 161P2F10B sequence as shown in Figure 2.

Additional illustrative embodiments of the invention disclosed herein include 161P2F10B polynucleotide fragments encoding one or more of the biological motifs contained within a 161P2F10B protein "or variant" sequence, including one or more of the motif-bearing subsequences of a 161P2F10B protein "or variant" set forth in Tables VIII-XXI and XXII-XLIX. In another embodiment, typical polynucleotide fragments of the invention encode one or more of the regions of 161P2F10B protein or variant that exhibit homology to a known molecule. In another embodiment of the invention, typical polynucleotide fragments can encode one or more of the 161P2F10B protein or variant N-glycosylation sites, cAMP and cGMP-dependent protein kinase phosphorylation sites, casein kinase II phosphorylation sites or N-myristoylation site and amidation sites.

Note that to determine the starting position of any peptide set forth in Tables VIII-XXI and Tables XXII to XLIX (collectively HLA Peptide Tables) relative to its parental protein, e.g., variant 1, variant 2, etc., reference is made to three factors: the particular variant, the length of the peptide in an HLA Peptide Table, and the Search Peptides listed in Table LVII. Generally, a unique Search Peptide is used to obtain HLA peptides for a particular variant. The position of each Search Peptide relative to its respective parent molecule is listed in Table VII. Accordingly, if a Search Peptide begins at position "X", one must add the value "X minus 1" to each position in Tables VIII-XXI and Tables XXII-IL to obtain the actual position of the HLA peptides in their parental molecule. For example if a particular Search Peptide begins at position 150 of its parental molecule, one must add 150 - 1, i.e., 149 to each HLA peptide amino acid position to calculate the position of that amino acid in the parent molecule.

II.A.) Uses of 161P2F10B Polynucleotides

II.A.1.) Monitoring of Genetic Abnormalities

The polynucleotides of the preceding paragraphs have a number of different specific uses. The human 161P2F10B gene maps to the chromosomal location set forth in the Example entitled "Chromosomal Mapping of 161P2F10B." For example, because the 161P2F10B gene maps to this chromosome, polynucleotides that encode different regions of the 161P2F10B proteins are used to characterize cytogenetic abnormalities of this chromosomal locale, such as abnormalities that are identified as being associated with various cancers. In certain genes, a variety of chromosomal abnormalities including rearrangements have been identified as frequent cytogenetic abnormalities in a number of different

cancers (see e.g. Krajinovic *et al.*, *Mutat. Res.* 382(3-4): 81-83 (1998); Johansson *et al.*, *Blood* 86(10): 3905-3914 (1995) and Finger *et al.*, *P.N.A.S.* 85(23): 9158-9162 (1988)). Thus, polynucleotides encoding specific regions of the 161P2F10B proteins provide new tools that can be used to delineate, with greater precision than previously possible, cytogenetic abnormalities in the chromosomal region that encodes 161P2F10B that may contribute to the malignant phenotype. In this context, these polynucleotides satisfy a need in the art for expanding the sensitivity of chromosomal screening in order to identify more subtle and less common chromosomal abnormalities (see e.g. Evans *et al.*, *Am. J. Obstet. Gynecol* 171(4): 1055-1057 (1994)).

Furthermore, as 161P2F10B was shown to be highly expressed in bladder and other cancers, 161P2F10B polynucleotides are used in methods assessing the status of 161P2F10B gene products in normal versus cancerous tissues. Typically, polynucleotides that encode specific regions of the 161P2F10B proteins are used to assess the presence of perturbations (such as deletions, insertions, point mutations, or alterations resulting in a loss of an antigen etc.) in specific regions of the 161P2F10B gene, such as regions containing one or more motifs. Exemplary assays include both RT-PCR assays as well as single-strand conformation polymorphism (SSCP) analysis (see, e.g., Marrogi *et al.*, *J. Cutan. Pathol.* 26(8): 369-378 (1999), both of which utilize polynucleotides encoding specific regions of a protein to examine these regions within the protein.

II.A.2.) Antisense Embodiments

Other specifically contemplated nucleic acid related embodiments of the invention disclosed herein are genomic DNA, cDNAs, ribozymes, and antisense molecules, as well as nucleic acid molecules based on an alternative backbone, or including alternative bases, whether derived from natural sources or synthesized, and include molecules capable of inhibiting the RNA or protein expression of 161P2F10B. For example, antisense molecules can be RNAs or other molecules, including peptide nucleic acids (PNAs) or non-nucleic acid molecules such as phosphorothioate derivatives that specifically bind DNA or RNA in a base pair-dependent manner. A skilled artisan can readily obtain these classes of nucleic acid molecules using the 161P2F10B polynucleotides and polynucleotide sequences disclosed herein.

Antisense technology entails the administration of exogenous oligonucleotides that bind to a target polynucleotide located within the cells. The term "antisense" refers to the fact that such oligonucleotides are complementary to their intracellular targets, e.g., 161P2F10B. See for example, Jack Cohen, *Oligodeoxynucleotides, Antisense Inhibitors of Gene Expression*, CRC Press, 1989; and *Synthesis* 1:1-5 (1988). The 161P2F10B antisense oligonucleotides of the present invention include derivatives such as S-oligonucleotides (phosphorothioate derivatives or S-oligos, see, Jack Cohen, *supra*), which exhibit enhanced cancer cell growth inhibitory action. S-oligos (nucleoside phosphorothioates) are isoelectronic analogs of an oligonucleotide (O-oligo) in which a nonbridging oxygen atom of the phosphate group is replaced by a sulfur atom. The S-oligos of the present invention can be prepared by treatment of the corresponding O-oligos with 3H-1,2-benzodithiol-3-one-1,1-dioxide, which is a sulfur transfer reagent. See, e.g., Iyer, R. P. *et al.*, *J. Org. Chem.* 55:4693-4698 (1990); and Iyer, R. P. *et al.*, *J. Am. Chem. Soc.* 112:1253-1254 (1990). Additional 161P2F10B antisense oligonucleotides of the present invention include morpholino antisense oligonucleotides known in the art (see, e.g., Partridge *et al.*, 1996, *Antisense & Nucleic Acid Drug Development* 6: 169-175).

The 161P2F10B antisense oligonucleotides of the present invention typically can be RNA or DNA that is complementary to and stably hybridizes with the first 100 5' codons or last 100 3' codons of a 161P2F10B genomic sequence or the corresponding mRNA. Absolute complementarity is not required, although high degrees of complementarity are preferred. Use of an oligonucleotide complementary to this region allows for the selective hybridization to 161P2F10B mRNA and not to mRNA specifying other regulatory subunits of protein kinase. In one embodiment, 161P2F10B antisense oligonucleotides of the present invention are 15 to 30-mer fragments of the antisense DNA molecule that have a sequence that hybridizes to 161P2F10B mRNA. Optionally, 161P2F10B antisense oligonucleotide is a 30-mer oligonucleotide that is

complementary to a region in the first 10 5' codons or last 10 3' codons of 161P2F10B. Alternatively, the antisense molecules are modified to employ ribozymes in the inhibition of 161P2F10B expression, see, e.g., L. A. Couture & D. T. Stinchcomb; *Trends Genet* 12: 510-515 (1996).

II.A.3.) Primers and Primer Pairs

Further specific embodiments of these nucleotides of the invention include primers and primer pairs, which allow the specific amplification of polynucleotides of the invention or of any specific parts thereof, and probes that selectively or specifically hybridize to nucleic acid molecules of the invention or to any part thereof. Probes can be labeled with a detectable marker, such as, for example, a radioisotope, fluorescent compound, bioluminescent compound, a chemiluminescent compound, metal chelator or enzyme. Such probes and primers are used to detect the presence of a 161P2F10B polynucleotide in a sample and as a means for detecting a cell expressing a 161P2F10B protein.

Examples of such probes include polypeptides comprising all or part of the human 161P2F10B cDNA sequence shown in Figure 2. Examples of primer pairs capable of specifically amplifying 161P2F10B mRNAs are also described in the Examples. As will be understood by the skilled artisan, a great many different primers and probes can be prepared based on the sequences provided herein and used effectively to amplify and/or detect a 161P2F10B mRNA.

The 161P2F10B polynucleotides of the invention are useful for a variety of purposes, including but not limited to their use as probes and primers for the amplification and/or detection of the 161P2F10B gene(s), mRNA(s), or fragments thereof; as reagents for the diagnosis and/or prognosis of prostate cancer and other cancers; as coding sequences capable of directing the expression of 161P2F10B polypeptides; as tools for modulating or inhibiting the expression of the 161P2F10B gene(s) and/or translation of the 161P2F10B transcript(s); and as therapeutic agents.

The present invention includes the use of any probe as described herein to identify and isolate a 161P2F10B or 161P2F10B related nucleic acid sequence from a naturally occurring source, such as humans or other mammals, as well as the isolated nucleic acid sequence *per se*, which would comprise all or most of the sequences found in the probe used.

II.A.4.) Isolation of 161P2F10B-Encoding Nucleic Acid Molecules

The 161P2F10B cDNA sequences described herein enable the isolation of other polynucleotides encoding 161P2F10B gene product(s), as well as the isolation of polynucleotides encoding 161P2F10B gene product homologs, alternatively spliced isoforms, allelic variants, and mutant forms of a 161P2F10B gene product as well as polynucleotides that encode analogs of 161P2F10B-related proteins. Various molecular cloning methods that can be employed to isolate full length cDNAs encoding a 161P2F10B gene are well known (see, for example, Sambrook, J. *et al.*, *Molecular Cloning: A Laboratory Manual*, 2d edition, Cold Spring Harbor Press, New York, 1989; *Current Protocols in Molecular Biology*. Ausubel *et al.*, Eds., Wiley and Sons, 1995). For example, lambda phage cloning methodologies can be conveniently employed, using commercially available cloning systems (e.g., Lambda ZAP Express, Stratagene). Phage clones containing 161P2F10B gene cDNAs can be identified by probing with a labeled 161P2F10B cDNA or a fragment thereof. For example, in one embodiment, a 161P2F10B cDNA (e.g., Figure 2) or a portion thereof can be synthesized and used as a probe to retrieve overlapping and full-length cDNAs corresponding to a 161P2F10B gene. A 161P2F10B gene itself can be isolated by screening genomic DNA libraries, bacterial artificial chromosome libraries (BACs), yeast artificial chromosome libraries (YACs), and the like, with 161P2F10B DNA probes or primers.

II.A.5.) Recombinant Nucleic Acid Molecules and Host-Vector Systems

The invention also provides recombinant DNA or RNA molecules containing a 161P2F10B polynucleotide, a fragment, analog or homologue thereof, including but not limited to phages, plasmids, phagemids, cosmids, YACs, BACs, as well as various viral and non-viral vectors well known in the art, and cells transformed or transfected with such recombinant DNA or RNA molecules. Methods for generating such molecules are well known (see, for example, Sambrook *et al.*, 1989, *supra*).

The invention further provides a host-vector system comprising a recombinant DNA molecule containing a 161P2F10B polynucleotide, fragment, analog or homologue thereof within a suitable prokaryotic or eukaryotic host cell.

Examples of suitable eukaryotic host cells include a yeast cell, a plant cell, or an animal cell, such as a mammalian cell or an insect cell (e.g., a baculovirus-infectible cell such as an Sf9 or HighFive cell). Examples of suitable mammalian cells include various prostate cancer cell lines such as DU145 and TsuPr1, other transfectable or transducible prostate cancer cell lines, primary cells (PrEC), as well as a number of mammalian cells routinely used for the expression of recombinant proteins (e.g., COS, CHO, 293, 293T cells). More particularly, a polynucleotide comprising the coding sequence of 161P2F10B or a fragment, analog or homolog thereof can be used to generate 161P2F10B proteins or fragments thereof using any number of host-vector systems routinely used and widely known in the art.

A wide range of host-vector systems suitable for the expression of 161P2F10B proteins or fragments thereof are available, see for example, Sambrook *et al.*, 1989, *supra*; Current Protocols in Molecular Biology, 1995, *supra*). Preferred vectors for mammalian expression include but are not limited to pcDNA 3.1 myc-His-tag (Invitrogen) and the retroviral vector pSR α tkneo (Muller *et al.*, 1991, MCB 11:1785). Using these expression vectors, 161P2F10B can be expressed in several prostate cancer and non-prostate cell lines, including for example 293, 293T, rat-1, NIH 3T3 and TsuPr1. The host-vector systems of the invention are useful for the production of a 161P2F10B protein or fragment thereof. Such host-vector systems can be employed to study the functional properties of 161P2F10B and 161P2F10B mutations or analogs.

Recombinant human 161P2F10B protein or an analog or homolog or fragment thereof can be produced by mammalian cells transfected with a construct encoding a 161P2F10B-related nucleotide. For example, 293T cells can be transfected with an expression plasmid encoding 161P2F10B or fragment, analog or homolog thereof, a 161P2F10B-related protein is expressed in the 293T cells, and the recombinant 161P2F10B protein is isolated using standard purification methods (e.g., affinity purification using anti-161P2F10B antibodies). In another embodiment, a 161P2F10B coding sequence is subcloned into the retroviral vector pSR α MSVtkneo and used to infect various mammalian cell lines, such as NIH 3T3, TsuPr1, 293 and rat-1 in order to establish 161P2F10B expressing cell lines. Various other expression systems well known in the art can also be employed. Expression constructs encoding a leader peptide joined in frame to a 161P2F10B coding sequence can be used for the generation of a secreted form of recombinant 161P2F10B protein.

As discussed herein, redundancy in the genetic code permits variation in 161P2F10B gene sequences. In particular, it is known in the art that specific host species often have specific codon preferences, and thus one can adapt the disclosed sequence as preferred for a desired host. For example, preferred analog codon sequences typically have rare codons (i.e., codons having a usage frequency of less than about 20% in known sequences of the desired host) replaced with higher frequency codons. Codon preferences for a specific species are calculated, for example, by utilizing codon usage tables available on the INTERNET such as at URL dna.affrc.go.jp/~nakamura/codon.html.

Additional sequence modifications are known to enhance protein expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon/intron splice site signals, transposon-like repeats, and/or other such well-characterized sequences that are deleterious to gene expression. The GC content of the sequence is adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. Where possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures. Other useful modifications include the addition of a translational initiation consensus sequence at the start of the open reading frame, as described in Kozak, *Mol. Cell Biol.*, 9:5073-5080 (1989). Skilled artisans understand that the general rule that eukaryotic ribosomes initiate translation exclusively at the 5' proximal AUG codon is abrogated only under rare conditions (see, e.g., Kozak PNAS 92(7): 2662-2666, (1995) and Kozak NAR 15(20): 8125-8148 (1987)).

III.) 161P2F10B-related Proteins

Another aspect of the present invention provides 161P2F10B-related proteins. Specific embodiments of 161P2F10B proteins comprise a polypeptide having all or part of the amino acid sequence of human 161P2F10B as shown

in Figure 2 or Figure 3. Alternatively, embodiments of 161P2F10B proteins comprise variant, homolog or analog polypeptides that have alterations in the amino acid sequence of 161P2F10B shown in Figure 2 or Figure 3.

Embodiments of a 161P2F10B polypeptide include: a 161P2F10B polypeptide having a sequence shown in Figure 2, a peptide sequence of a 161P2F10B as shown in Figure 2 wherein T is U; at least 10 contiguous nucleotides of a polypeptide having the sequence as shown in Figure 2; or, at least 10 contiguous peptides of a polypeptide having the sequence as shown in Figure 2 where T is U. For example, embodiments of 161P2F10B peptides comprise, without limitation:

- (I) a protein comprising, consisting essentially of, or consisting of an amino acid sequence as shown in Figure 2A-G or Figure 3A-E;
- (II) a 161P2F10B-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% homologous to an entire amino acid sequence shown in Figure 2A-G;
- (III) a 161P2F10B-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% identical to an entire amino acid sequence shown in Figure 2A-G or 3A-E;
- (IV) a protein that comprises at least one peptide set forth in Tables VIII to XLIX, optionally with a *proviso* that it is not an entire protein of Figure 2;
- (V) a protein that comprises at least one peptide set forth in Tables VIII-XXI, collectively, which peptide is also set forth in Tables XXII to XLIX, collectively, optionally with a *proviso* that it is not an entire protein of Figure 2;
- (VI) a protein that comprises at least two peptides selected from the peptides set forth in Tables VIII-XLIX, optionally with a *proviso* that it is not an entire protein of Figure 2;
- (VII) a protein that comprises at least two peptides selected from the peptides set forth in Tables VIII to XLIX collectively, with a *proviso* that the protein is not a contiguous sequence from an amino acid sequence of Figure 2;
- (VIII) a protein that comprises at least one peptide selected from the peptides set forth in Tables VIII-XXI; and at least one peptide selected from the peptides set forth in Tables XXII to XLIX, with a *proviso* that the protein is not a contiguous sequence from an amino acid sequence of Figure 2;
- (IX) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of Figure 3A, 3B, 3C, 3D, or 3E in any whole number increment up to 875, 875, 875, 875, or 841 respectively that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of Figure 5;
- (X) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of Figure 3A, 3B, 3C, 3D, or 3E in any whole number increment up to 875, 875, 875, 875, or 841 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydropathicity profile of Figure 6;

- (XI) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of Figure 3A, 3B, 3C, 3D, or 3E in any whole number increment up to 875, 875, 875, 875, or 841 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of Figure 7;
- (XII) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of Figure 3A, 3B, 3C, 3D, or 3E in any whole number increment up to 875, 875, 875, 875, or 841 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of Figure 8;
- (XIII) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of Figure 3A, 3B, 3C, 3D, or 3E in any whole number increment up to 875, 875, 875, 875, or 841 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of Figure 9;
- (XIV) a peptide that occurs at least twice in Tables VIII-XXI and XXII to XLIX, collectively;
- (XV) a peptide that occurs at least three times in Tables VIII-XXI and XXII to XLIX, collectively;
- (XVI) a peptide that occurs at least four times in Tables VIII-XXI and XXII to XLIX, collectively;
- (XVII) a peptide that occurs at least five times in Tables VIII-XXI and XXII to XLIX, collectively;
- (XVIII) a peptide that occurs at least once in Tables VIII-XXI, and at least once in tables XXII to XLIX;
- (XIX) a peptide that occurs at least once in Tables VIII-XXI, and at least twice in tables XXII to XLIX;
- (XX) a peptide that occurs at least twice in Tables VIII-XXI, and at least once in tables XXII to XLIX;
- (XXI) a peptide that occurs at least twice in Tables VIII-XXI, and at least twice in tables XXII to XLIX;
- (XXII) a peptide which comprises one two, three, four, or five of the following characteristics, or an oligonucleotide encoding such peptide:
- i) a region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Hydrophilicity profile of Figure 5;
 - ii) a region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or less than 0.5, 0.4, 0.3, 0.2, 0.1, or having a value equal to 0.0, in the Hydropathicity profile of Figure 6;
 - iii) a region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Percent Accessible Residues profile of Figure 7;
 - iv) a region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Average Flexibility profile of Figure 8; or,

v) a region of at least 5 amino acids of a particular peptide of Figure 3, in any whole number increment up to the full length of that protein in Figure 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Beta-turn profile of Figure 9;

(XXIII) a monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)15 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXIV) a monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)29 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXV) a monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)37 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXVI) a monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(4)6 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXVII) a monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)17 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXVIII) a monolonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)50 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____;

(XXIX) a composition comprising a peptide of (I)-(XXII) or an antibody or binding region thereof of (XXIII to XXVIII) together with a pharmaceutical excipient and/or in a human unit dose form.

(XXX) a method of using a peptide of (I)-(XXII), or an antibody or binding region thereof of (XXIII to XXVIII) or a composition of (XXIX) in a method to modulate a cell expressing 161P2F10b,

(XXXI) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof of (XXIII to XXVIII) or a composition of (XXIX) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 161P2F10b

(XXXII) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof of (XXIII to XXVIII) or a composition (XXIX) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 161P2F10b, said cell from a cancer of a tissue listed in Table I;

(XXXIII) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof of (XXIII to XXVIII) or a composition of (XXIX) in a method to diagnose, prophylax, prognose, or treat a cancer;

(XXXIV) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof of (XXIII to XXVIII) or a composition of (XXIX) in a method to diagnose, prophylax, prognose, or treat a cancer of a tissue listed in Table I; and,

(XXXV) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof of (XXIII to XXVIII) or a composition (XXIX) in a method to identify or characterize a modulator of a cell expressing 161P2F10b.

As used herein, a range is understood to specifically disclose all whole unit positions thereof.

Typical embodiments of the invention disclosed herein include 161P2F10B polynucleotides that encode specific portions of 161P2F10B mRNA sequences (and those which are complementary to such sequences) such as those that encode the proteins and/or fragments thereof, for example:

(a) 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 860, 870, 875 or more contiguous amino acids of 161P2F10B variant 1; the maximal lengths relevant for other variants are: variant 2, 875 amino acids; variant 3, 875 amino acids, variant 4, 875, and variant 7, 841 amino acids..

In general, naturally occurring allelic variants of human 161P2F10B share a high degree of structural identity and homology (e.g., 90% or more homology). Typically, allelic variants of a 161P2F10B protein contain conservative amino acid substitutions within the 161P2F10B sequences described herein or contain a substitution of an amino acid from a corresponding position in a homologue of 161P2F10B. One class of 161P2F10B allelic variants are proteins that share a high degree of homology with at least a small region of a particular 161P2F10B amino acid sequence, but further contain a radical departure from the sequence, such as a non-conservative substitution, truncation, insertion or frame shift. In comparisons of protein sequences, the terms, similarity, identity, and homology each have a distinct meaning as appreciated in the field of genetics. Moreover, orthology and paralogy can be important concepts describing the relationship of members of a given protein family in one organism to the members of the same family in other organisms.

Amino acid abbreviations are provided in Table II. Conservative amino acid substitutions can frequently be made in a protein without altering either the conformation or the function of the protein. Proteins of the invention can comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 conservative substitutions. Such changes include substituting any of isoleucine (I), valine (V), and leucine (L) for any other of these hydrophobic amino acids; aspartic acid (D) for glutamic acid (E) and vice versa; glutamine (Q) for asparagine (N) and vice versa; and serine (S) for threonine (T) and vice versa. Other substitutions can also be considered conservative, depending on the environment of the particular amino acid and its role in the three-dimensional structure of the protein. For example, glycine (G) and alanine (A) can frequently be interchangeable, as can alanine (A) and valine (V). Methionine (M), which is relatively hydrophobic, can frequently be interchanged with leucine and isoleucine, and sometimes with valine. Lysine (K) and arginine (R) are frequently interchangeable in locations in which the significant feature of the amino acid residue is its charge and the differing pK's of these two amino acid residues are not significant. Still other changes can be considered "conservative" in particular environments (see, e.g. Table III herein; pages 13-15 "Biochemistry" 2nd ED. Lubert Stryer ed (Stanford University); Henikoff *et al.*, PNAS 1992 Vol 89 10915-10919; Lei *et al.*, J Biol Chem 1995 May 19; 270(20):11882-6).

Embodiments of the invention disclosed herein include a wide variety of art-accepted variants or analogs of 161P2F10B proteins such as polypeptides having amino acid insertions, deletions and substitutions. 161P2F10B variants can be made using methods known in the art such as site-directed mutagenesis, alanine scanning, and PCR mutagenesis.

Site-directed mutagenesis (Carter *et al.*, *Nucl. Acids Res.*, 13:4331 (1986); Zoller *et al.*, *Nucl. Acids Res.*, 10:6487 (1987)), cassette mutagenesis (Wells *et al.*, *Gene*, 34:315 (1985)), restriction selection mutagenesis (Wells *et al.*, *Philos. Trans. R. Soc. London SerA*, 317:415 (1986)) or other known techniques can be performed on the cloned DNA to produce the 161P2F10B variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence that is involved in a specific biological activity such as a protein-protein interaction. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions (Creighton, *The Proteins*, (W.H. Freeman & Co., N.Y.); Chothia, *J. Mol. Biol.*, 150:1 (1976)). If alanine substitution does not yield adequate amounts of variant, an isosteric amino acid can be used.

As defined herein, 161P2F10B variants, analogs or homologs, have the distinguishing attribute of having at least one epitope that is "cross reactive" with a 161P2F10B protein having an amino acid sequence of Figure 3. As used in this sentence, "cross reactive" means that an antibody or T cell that specifically binds to a 161P2F10B variant also specifically binds to a 161P2F10B protein having an amino acid sequence set forth in Figure 3. A polypeptide ceases to be a variant of a protein shown in Figure 3, when it no longer contains any epitope capable of being recognized by an antibody or T cell that specifically binds to the starting 161P2F10B protein. Those skilled in the art understand that antibodies that recognize proteins bind to epitopes of varying size, and a grouping of the order of about four or five amino acids, contiguous or not, is regarded as a typical number of amino acids in a minimal epitope. See, e.g., Nair *et al.*, *J. Immunol* 2000 165(12): 6949-6955; Hebbes *et al.*, *Mol Immunol* (1989) 26(9):865-73; Schwartz *et al.*, *J Immunol* (1985) 135(4):2598-608.

Other classes of 161P2F10B-related protein variants share 70%, 75%, 80%, 85% or 90% or more similarity with an amino acid sequence of Figure 3, or a fragment thereof. Another specific class of 161P2F10B protein variants or analogs comprises one or more of the 161P2F10B biological motifs described herein or presently known in the art. Thus, encompassed by the present invention are analogs of 161P2F10B fragments (nucleic or amino acid) that have altered functional (e.g. immunogenic) properties relative to the starting fragment. It is to be appreciated that motifs now or which become part of the art are to be applied to the nucleic or amino acid sequences of Figure 2 or Figure 3.

As discussed herein, embodiments of the claimed invention include polypeptides containing less than the full amino acid sequence of a 161P2F10B protein shown in Figure 2 or Figure 3. For example, representative embodiments of the invention comprise peptides/proteins having any 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or more contiguous amino acids of a 161P2F10B protein shown in Figure 2 or Figure 3.

Moreover, representative embodiments of the invention disclosed herein include polypeptides consisting of about amino acid 1 to about amino acid 10 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 10 to about amino acid 20 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 20 to about amino acid 30 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 30 to about amino acid 40 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 40 to about amino acid 50 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 50 to about amino acid 60 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 60 to about amino acid 70 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 70 to about amino acid 80 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 80 to about amino acid 90 of a 161P2F10B protein shown in Figure 2 or Figure 3, polypeptides consisting of about amino acid 90 to about amino acid 100 of a 161P2F10B protein shown in

Figure 2 or Figure 3, etc. throughout the entirety of a 161P2F10B amino acid sequence. Moreover, polypeptides consisting of about amino acid 1 (or 20 or 30 or 40 etc.) to about amino acid 20, (or 130, or 140 or 150 etc.) of a 161P2F10B protein shown in Figure 2 or Figure 3 are embodiments of the invention. It is to be appreciated that the starting and stopping positions in this paragraph refer to the specified position as well as that position plus or minus 5 residues.

161P2F10B-related proteins are generated using standard peptide synthesis technology or using chemical cleavage methods well known in the art. Alternatively, recombinant methods can be used to generate nucleic acid molecules that encode a 161P2F10B-related protein. In one embodiment, nucleic acid molecules provide a means to generate defined fragments of a 161P2F10B protein (or variants, homologs or analogs thereof).

III.A.) Motif-bearing Protein Embodiments

Additional illustrative embodiments of the invention disclosed herein include 161P2F10B polypeptides comprising the amino acid residues of one or more of the biological motifs contained within a 161P2F10B polypeptide sequence set forth in Figure 2 or Figure 3. Various motifs are known in the art, and a protein can be evaluated for the presence of such motifs by a number of publicly available Internet sites (see, e.g., URL addresses: pfam.wustl.edu/; searchlauncher.bcm.tmc.edu/seq-search/struc-predict.html; psort.ims.u-tokyo.ac.jp/; cbs.dtu.dk/; ebi.ac.uk/interpro/scan.html; expasy.ch/tools/scnpsit1.html; Epimatrix™ and Epimer™, Brown University, brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html; and BIMAS, bimas.dcrf.nih.gov/).

Motif bearing subsequences of all 161P2F10B variant proteins are set forth and identified in Tables VIII-XXI and XXII-XLIX.

Table V sets forth several frequently occurring motifs based on pfam searches (see URL address pfam.wustl.edu/). The columns of Table V list (1) motif name abbreviation, (2) percent identity found amongst the different member of the motif family, (3) motif name or description and (4) most common function; location information is included if the motif is relevant for location.

Polypeptides comprising one or more of the 161P2F10B motifs discussed above are useful in elucidating the specific characteristics of a malignant phenotype in view of the observation that the 161P2F10B motifs discussed above are associated with growth dysregulation and because 161P2F10B is overexpressed in certain cancers (See, e.g., Table I). Casein kinase II, cAMP and camp-dependent protein kinase, and Protein Kinase C, for example, are enzymes known to be associated with the development of the malignant phenotype (see e.g. Chen *et al.*, Lab Invest., 78(2): 165-174 (1998); Gaiddon *et al.*, Endocrinology 136(10): 4331-4338 (1995); Hall *et al.*, Nucleic Acids Research 24(6): 1119-1126 (1996); Peterziel *et al.*, Oncogene 18(46): 6322-6329 (1999) and O'Brian, Oncol. Rep. 5(2): 305-309 (1998)). Moreover, both glycosylation and myristoylation are protein modifications also associated with cancer and cancer progression (see e.g. Dennis *et al.*, Biochem. Biophys. Acta 1473(1):21-34 (1999); Raju *et al.*, Exp. Cell Res. 235(1): 145-154 (1997)). Amidation is another protein modification also associated with cancer and cancer progression (see e.g. Treston *et al.*, J. Natl. Cancer Inst. Monogr. (13): 169-175 (1992)).

In another embodiment, proteins of the invention comprise one or more of the immunoreactive epitopes identified in accordance with art-accepted methods, such as the peptides set forth in Tables VIII-XXI and XXII-XLIX. CTL epitopes can be determined using specific algorithms to identify peptides within a 161P2F10B protein that are capable of optimally binding to specified HLA alleles (e.g., Table IV; Epimatrix™ and Epimer™, Brown University, URL brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html; and BIMAS, URL bimas.dcrf.nih.gov/.) Moreover, processes for identifying peptides that have sufficient binding affinity for HLA molecules and which are correlated with being immunogenic epitopes, are well known in the art, and are carried out without undue experimentation. In addition, processes for identifying peptides that are immunogenic epitopes, are well known in the art, and are carried out without undue experimentation either *in vitro* or *in vivo*.

Also known in the art are principles for creating analogs of such epitopes in order to modulate immunogenicity. For example, one begins with an epitope that bears a CTL or HTL motif (see, e.g., the HLA Class I and HLA Class II motifs/super motifs of Table IV). The epitope is analoged by substituting out an amino acid at one of the specified positions, and replacing it with another amino acid specified for that position. For example, on the basis of residues defined in Table IV, one can substitute out a deleterious residue in favor of any other residue, such as a preferred residue; substitute a less-preferred residue with a preferred residue; or substitute an originally-occurring preferred residue with another preferred residue. Substitutions can occur at primary anchor positions or at other positions in a peptide; see, e.g., Table IV.

A variety of references reflect the art regarding the identification and generation of epitopes in a protein of interest as well as analogs thereof. See, for example, WO 97/33602 to Chesnut *et al.*; Sette, Immunogenetics 1999 50(3-4): 201-212; Sette *et al.*, J. Immunol. 2001 166(2): 1389-1397; Sidney *et al.*, Hum. Immunol. 1997 58(1): 12-20; Kondo *et al.*, Immunogenetics 1997 45(4): 249-258; Sidney *et al.*, J. Immunol. 1996 157(8): 3480-90; and Falk *et al.*, Nature 351: 290-6 (1991); Hunt *et al.*, Science 255:1261-3 (1992); Parker *et al.*, J. Immunol. 149:3580-7 (1992); Parker *et al.*, J. Immunol. 152:163-75 (1994); Kast *et al.*, 1994 152(8): 3904-12; Borrás-Cuesta *et al.*, Hum. Immunol. 2000 61(3): 266-278; Alexander *et al.*, J. Immunol. 2000 164(3): 1625-1633; Alexander *et al.*, PMID: 7895164, UI: 95202582; O'Sullivan *et al.*, J. Immunol. 1991 147(8): 2663-2669; Alexander *et al.*, Immunity 1994 1(9): 751-761 and Alexander *et al.*, Immunol. Res. 1998 18(2): 79-92.

Related embodiments of the invention include polypeptides comprising combinations of the different motifs set forth in Table VI, and/or, one or more of the predicted CTL epitopes of Tables VIII-XXI and XXII-XLIX, and/or, one or more of the predicted HTL epitopes of Tables XLVI-XLIX, and/or, one or more of the T cell binding motifs known in the art. Preferred embodiments contain no insertions, deletions or substitutions either within the motifs or within the intervening sequences of the polypeptides. In addition, embodiments which include a number of either N-terminal and/or C-terminal amino acid residues on either side of these motifs may be desirable (to, for example, include a greater portion of the polypeptide architecture in which the motif is located). Typically, the number of N-terminal and/or C-terminal amino acid residues on either side of a motif is between about 1 to about 100 amino acid residues, preferably 5 to about 50 amino acid residues.

161P2F10B-related proteins are embodied in many forms, preferably in isolated form. A purified 161P2F10B protein molecule will be substantially free of other proteins or molecules that impair the binding of 161P2F10B to antibody, T cell or other ligand. The nature and degree of isolation and purification will depend on the intended use. Embodiments of a 161P2F10B-related proteins include purified 161P2F10B-related proteins and functional, soluble 161P2F10B-related proteins. In one embodiment, a functional, soluble 161P2F10B protein or fragment thereof retains the ability to be bound by antibody, T cell or other ligand.

The invention also provides 161P2F10B proteins comprising biologically active fragments of a 161P2F10B amino acid sequence shown in Figure 2 or Figure 3. Such proteins exhibit properties of the starting 161P2F10B protein, such as the ability to elicit the generation of antibodies that specifically bind an epitope associated with the starting 161P2F10B protein; to be bound by such antibodies; to elicit the activation of HTL or CTL; and/or, to be recognized by HTL or CTL that also specifically bind to the starting protein.

161P2F10B-related polypeptides that contain particularly interesting structures can be predicted and/or identified using various analytical techniques well known in the art, including, for example, the methods of Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis, or based on immunogenicity. Fragments that contain such structures are particularly useful in generating subunit-specific anti-161P2F10B antibodies or T cells or in identifying cellular factors that bind to 161P2F10B. For example, hydrophilicity profiles can be generated, and immunogenic peptide fragments identified, using the method of Hopp, T.P. and Woods, K.R., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828. Hydrophobicity profiles can be generated, and immunogenic peptide fragments identified, using the method of Kyte, J. and

Doolittle, R.F., 1982, J. Mol. Biol. 157:105-132. Percent (%) Accessible Residues profiles can be generated, and immunogenic peptide fragments identified, using the method of Janin J., 1979, Nature 277:491-492. Average Flexibility profiles can be generated, and immunogenic peptide fragments identified, using the method of Bhaskaran R., Ponnuswamy P.K., 1988, Int. J. Pept. Protein Res. 32:242-255. Beta-turn profiles can be generated, and immunogenic peptide fragments identified, using the method of Deleage, G., Roux B., 1987, Protein Engineering 1:289-294.

CTL epitopes can be determined using specific algorithms to identify peptides within a 161P2F10B protein that are capable of optimally binding to specified HLA alleles (e.g., by using the SYFPEITHI site at World Wide Web URL syfpeithi.bmi-heidelberg.com/; the listings in Table IV(A)-(E); Epimatrix™ and Epimer™, Brown University, URL (brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html); and BIMAS, URL bimas.dcrf.nih.gov/). Illustrating this, peptide epitopes from 161P2F10B that are presented in the context of human MHC Class I molecules, e.g., HLA-A1, A2, A3, A11, A24, B7 and B35 were predicted (see, e.g., Tables VIII-XXI, XXII-XLIX). Specifically, the complete amino acid sequence of the 161P2F10B protein and relevant portions of other variants, i.e., for HLA Class I predictions 9 flanking residues on either side of a point mutation or exon junction, and for HLA Class II predictions 14 flanking residues on either side of a point mutation or exon junction corresponding to that variant, were entered into the HLA Peptide Motif Search algorithm found in the Bioinformatics and Molecular Analysis Section (BIMAS) web site listed above; in addition to the site SYFPEITHI, at URL syfpeithi.bmi-heidelberg.com/.

The HLA peptide motif search algorithm was developed by Dr. Ken Parker based on binding of specific peptide sequences in the groove of HLA Class I molecules, in particular HLA-A2 (see, e.g., Falk *et al.*, Nature 351: 290-6 (1991); Hunt *et al.*, Science 255:1261-3 (1992); Parker *et al.*, J. Immunol. 149:3580-7 (1992); Parker *et al.*, J. Immunol. 152:163-75 (1994)). This algorithm allows location and ranking of 8-mer, 9-mer, and 10-mer peptides from a complete protein sequence for predicted binding to HLA-A2 as well as numerous other HLA Class I molecules. Many HLA class I binding peptides are 8-, 9-, 10 or 11-mers. For example, for Class I HLA-A2, the epitopes preferably contain a leucine (L) or methionine (M) at position 2 and a valine (V) or leucine (L) at the C-terminus (see, e.g., Parker *et al.*, J. Immunol. 149:3580-7 (1992)). Selected results of 161P2F10B predicted binding peptides are shown in Tables VIII-XXI and XXII-XLIX herein. In Tables VIII-XXI and XXII-XLVII, selected candidates, 9-mers and 10-mers, for each family member are shown along with their location, the amino acid sequence of each specific peptide, and an estimated binding score. In Tables XLVI-XLIX, selected candidates, 15-mers, for each family member are shown along with their location, the amino acid sequence of each specific peptide, and an estimated binding score. The binding score corresponds to the estimated half time of dissociation of complexes containing the peptide at 37°C at pH 6.5. Peptides with the highest binding score are predicted to be the most tightly bound to HLA Class I on the cell surface for the greatest period of time and thus represent the best immunogenic targets for T-cell recognition.

Actual binding of peptides to an HLA allele can be evaluated by stabilization of HLA expression on the antigen-processing defective cell line T2 (see, e.g., Xue *et al.*, Prostate 30:73-8 (1997) and Peshwa *et al.*, Prostate 36:129-38 (1998)). Immunogenicity of specific peptides can be evaluated *in vitro* by stimulation of CD8+ cytotoxic T lymphocytes (CTL) in the presence of antigen presenting cells such as dendritic cells.

It is to be appreciated that every epitope predicted by the BIMAS site, Epimer™ and Epimatrix™ sites, or specified by the HLA class I or class II motifs available in the art or which become part of the art such as set forth in Table IV (or determined using World Wide Web site URL syfpeithi.bmi-heidelberg.com/, or BIMAS, bimas.dcrf.nih.gov/) are to be "applied" to a 161P2F10B protein in accordance with the invention. As used in this context "applied" means that a 161P2F10B protein is evaluated, e.g., visually or by computer-based patterns finding methods, as appreciated by those of skill in the relevant art. Every subsequence of a 161P2F10B protein of 8, 9, 10, or 11 amino acid residues that bears an HLA Class I motif, or a subsequence of 9 or more amino acid residues that bear an HLA Class II motif are within the scope of the invention.

III.B.) Expression of 161P2F10B-related Proteins

In an embodiment described in the examples that follow, 161P2F10B can be conveniently expressed in cells (such as 293T cells) transfected with a commercially available expression vector such as a CMV-driven expression vector encoding 161P2F10B with a C-terminal 6XHis and MYC tag (pcDNA3.1/mycHIS, Invitrogen or Tag5, GenHunter Corporation, Nashville TN). The Tag5 vector provides an IgGK secretion signal that can be used to facilitate the production of a secreted 161P2F10B protein in transfected cells. The secreted HIS-tagged 161P2F10B in the culture media can be purified, e.g., using a nickel column using standard techniques.

III.C.) Modifications of 161P2F10B-related Proteins

Modifications of 161P2F10B-related proteins such as covalent modifications are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a 161P2F10B polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of a 161P2F10B protein. Another type of covalent modification of a 161P2F10B polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of a protein of the invention. Another type of covalent modification of 161P2F10B comprises linking a 161P2F10B polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The 161P2F10B-related proteins of the present invention can also be modified to form a chimeric molecule comprising 161P2F10B fused to another, heterologous polypeptide or amino acid sequence. Such a chimeric molecule can be synthesized chemically or recombinantly. A chimeric molecule can have a protein of the invention fused to another tumor-associated antigen or fragment thereof. Alternatively, a protein in accordance with the invention can comprise a fusion of fragments of a 161P2F10B sequence (amino or nucleic acid) such that a molecule is created that is not, through its length, directly homologous to the amino or nucleic acid sequences shown in Figure 2 or Figure 3. Such a chimeric molecule can comprise multiples of the same subsequence of 161P2F10B. A chimeric molecule can comprise a fusion of a 161P2F10B-related protein with a polyhistidine epitope tag, which provides an epitope to which immobilized nickel can selectively bind, with cytokines or with growth factors. The epitope tag is generally placed at the amino- or carboxyl- terminus of a 161P2F10B protein. In an alternative embodiment, the chimeric molecule can comprise a fusion of a 161P2F10B-related protein with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a 161P2F10B polypeptide in place of at least one variable region within an Ig molecule. In a preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see, e.g., U.S. Patent No. 5,428,130 issued June 27, 1995.

III.D.) Uses of 161P2F10B-related Proteins

The proteins of the invention have a number of different specific uses. As 161P2F10B is highly expressed in prostate and other cancers, 161P2F10B-related proteins are used in methods that assess the status of 161P2F10B gene products in normal versus cancerous tissues, thereby elucidating the malignant phenotype. Typically, polypeptides from specific regions of a 161P2F10B protein are used to assess the presence of perturbations (such as deletions, insertions, point mutations etc.) in those regions (such as regions containing one or more motifs). Exemplary assays utilize antibodies or T cells targeting 161P2F10B-related proteins comprising the amino acid residues of one or more of the biological motifs

contained within a 161P2F10B polypeptide sequence in order to evaluate the characteristics of this region in normal versus cancerous tissues or to elicit an immune response to the epitope. Alternatively, 161P2F10B-related proteins that contain the amino acid residues of one or more of the biological motifs in a 161P2F10B protein are used to screen for factors that interact with that region of 161P2F10B.

161P2F10B protein fragments/subsequences are particularly useful in generating and characterizing domain-specific antibodies (e.g., antibodies recognizing an extracellular or intracellular epitope of a 161P2F10B protein), for identifying agents or cellular factors that bind to 161P2F10B or a particular structural domain thereof, and in various therapeutic and diagnostic contexts, including but not limited to diagnostic assays, cancer vaccines and methods of preparing such vaccines.

Proteins encoded by the 161P2F10B genes, or by analogs, homologs or fragments thereof, have a variety of uses, including but not limited to generating antibodies and in methods for identifying ligands and other agents and cellular constituents that bind to a 161P2F10B gene product. Antibodies raised against a 161P2F10B protein or fragment thereof are useful in diagnostic and prognostic assays, and imaging methodologies in the management of human cancers characterized by expression of 161P2F10B protein, such as those listed in Table I. Such antibodies can be expressed intracellularly and used in methods of treating patients with such cancers. 161P2F10B-related nucleic acids or proteins are also used in generating HTL or CTL responses.

Various immunological assays useful for the detection of 161P2F10B proteins are used, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), immunocytochemical methods, and the like. Antibodies can be labeled and used as immunological imaging reagents capable of detecting 161P2F10B-expressing cells (e.g., in radioscintigraphic imaging methods). 161P2F10B proteins are also particularly useful in generating cancer vaccines, as further described herein.

IV.) 161P2F10B Antibodies

Another aspect of the invention provides antibodies that bind to 161P2F10B-related proteins. Preferred antibodies specifically bind to a 161P2F10B-related protein and do not bind (or bind weakly) to peptides or proteins that are not 161P2F10B-related proteins. For example, antibodies that bind 161P2F10B can bind 161P2F10B-related proteins such as the homologs or analogs thereof.

161P2F10B antibodies of the invention are particularly useful in cancer (see, e.g., Table I) diagnostic and prognostic assays, and imaging methodologies. Similarly, such antibodies are useful in the treatment, diagnosis, and/or prognosis of other cancers, to the extent 161P2F10B is also expressed or overexpressed in these other cancers. Moreover, intracellularly expressed antibodies (e.g., single chain antibodies) are therapeutically useful in treating cancers in which the expression of 161P2F10B is involved, such as advanced or metastatic prostate cancers.

The invention also provides various immunological assays useful for the detection and quantification of 161P2F10B and mutant 161P2F10B-related proteins. Such assays can comprise one or more 161P2F10B antibodies capable of recognizing and binding a 161P2F10B-related protein, as appropriate. These assays are performed within various immunological assay formats well known in the art, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), and the like.

Immunological non-antibody assays of the invention also comprise T cell immunogenicity assays (inhibitory or stimulatory) as well as major histocompatibility complex (MHC) binding assays.

In addition, immunological imaging methods capable of detecting prostate cancer and other cancers expressing 161P2F10B are also provided by the invention, including but not limited to radioscintigraphic imaging methods using labeled 161P2F10B antibodies. Such assays are clinically useful in the detection, monitoring, and prognosis of 161P2F10B expressing cancers such as prostate cancer.

161P2F10B antibodies are also used in methods for purifying a 161P2F10B-related protein and for isolating 161P2F10B homologues and related molecules. For example, a method of purifying a 161P2F10B-related protein comprises incubating a 161P2F10B antibody, which has been coupled to a solid matrix, with a lysate or other solution containing a 161P2F10B-related protein under conditions that permit the 161P2F10B antibody to bind to the 161P2F10B-related protein; washing the solid matrix to eliminate impurities; and eluting the 161P2F10B-related protein from the coupled antibody. Other uses of 161P2F10B antibodies in accordance with the invention include generating anti-idiotypic antibodies that mimic a 161P2F10B protein.

Various methods for the preparation of antibodies are well known in the art. For example, antibodies can be prepared by immunizing a suitable mammalian host using a 161P2F10B-related protein, peptide, or fragment, in isolated or immunoconjugated form (Antibodies: A Laboratory Manual, CSH Press, Eds., Harlow, and Lane (1988); Harlow, Antibodies, Cold Spring Harbor Press, NY (1989)). In addition, fusion proteins of 161P2F10B can also be used, such as a 161P2F10B GST-fusion protein. In a particular embodiment, a GST fusion protein comprising all or most of the amino acid sequence of Figure 2 or Figure 3 is produced, then used as an immunogen to generate appropriate antibodies. In another embodiment, a 161P2F10B-related protein is synthesized and used as an immunogen.

In addition, naked DNA immunization techniques known in the art are used (with or without purified 161P2F10B-related protein or 161P2F10B expressing cells) to generate an immune response to the encoded immunogen (for review, see Donnelly *et al.*, 1997, Ann. Rev. Immunol. 15: 617-648).

The amino acid sequence of a 161P2F10B protein as shown in Figure 2 or Figure 3 can be analyzed to select specific regions of the 161P2F10B protein for generating antibodies. For example, hydrophobicity and hydrophilicity analyses of a 161P2F10B amino acid sequence are used to identify hydrophilic regions in the 161P2F10B structure. Regions of a 161P2F10B protein that show immunogenic structure, as well as other regions and domains, can readily be identified using various other methods known in the art, such as Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis. Hydrophilicity profiles can be generated using the method of Hopp, T.P. and Woods, K.R., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828. Hydropathicity profiles can be generated using the method of Kyte, J. and Doolittle, R.F., 1982, J. Mol. Biol. 157:105-132. Percent (%) Accessible Residues profiles can be generated using the method of Janin J., 1979, Nature 277:491-492. Average Flexibility profiles can be generated using the method of Bhaskaran R., Ponnuswamy P.K., 1988, Int. J. Pept. Protein Res. 32:242-255. Beta-turn profiles can be generated using the method of Deleage, G., Roux B., 1987, Protein Engineering 1:289-294. Thus, each region identified by any of these programs or methods is within the scope of the present invention. Methods for the generation of 161P2F10B antibodies are further illustrated by way of the examples provided herein. Methods for preparing a protein or polypeptide for use as an immunogen are well known in the art. Also well known in the art are methods for preparing immunogenic conjugates of a protein with a carrier, such as BSA, KLH or other carrier protein. In some circumstances, direct conjugation using, for example, carbodiimide reagents are used; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, IL, are effective. Administration of a 161P2F10B immunogen is often conducted by injection over a suitable time period and with use of a suitable adjuvant, as is understood in the art. During the immunization schedule, titers of antibodies can be taken to determine adequacy of antibody formation.

161P2F10B monoclonal antibodies can be produced by various means well known in the art. For example, immortalized cell lines that secrete a desired monoclonal antibody are prepared using the standard hybridoma technology of Kohler and Milstein or modifications that immortalize antibody-producing B cells, as is generally known. Immortalized cell lines that secrete the desired antibodies are screened by immunoassay in which the antigen is a 161P2F10B-related protein. When the appropriate immortalized cell culture is identified, the cells can be expanded and antibodies produced either from *in vitro* cultures or from ascites fluid.

The antibodies or fragments of the invention can also be produced, by recombinant means. Regions that bind specifically to the desired regions of a 161P2F10B protein can also be produced in the context of chimeric or complementarity-determining region (CDR) grafted antibodies of multiple species origin. Humanized or human 161P2F10B antibodies can also be produced, and are preferred for use in therapeutic contexts. Methods for humanizing murine and other non-human antibodies, by substituting one or more of the non-human antibody CDRs for corresponding human antibody sequences, are well known (see for example, Jones *et al.*, 1986, *Nature* 321: 522-525; Riechmann *et al.*, 1988, *Nature* 332: 323-327; Verhoeyen *et al.*, 1988, *Science* 239: 1534-1536). See also, Carter *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* 89: 4285 and Sims *et al.*, 1993, *J. Immunol.* 151: 2296.

Methods for producing fully human monoclonal antibodies include phage display and transgenic methods (for review, see Vaughan *et al.*, 1998, *Nature Biotechnology* 16: 535-539). Fully human 161P2F10B monoclonal antibodies can be generated using cloning technologies employing large human Ig gene combinatorial libraries (i.e., phage display) (Griffiths and Hoogenboom, Building an *in vitro* immune system: human antibodies from phage display libraries. In: Protein Engineering of Antibody Molecules for Prophylactic and Therapeutic Applications in Man, Clark, M. (Ed.), Nottingham Academic, pp 45-64 (1993); Burton and Barbas, Human Antibodies from combinatorial libraries. *Id.*, pp 65-82). Fully human 161P2F10B monoclonal antibodies can also be produced using transgenic mice engineered to contain human immunoglobulin gene loci as described in PCT Patent Application WO98/24893, Kucherlapati and Jakobovits *et al.*, published December 3, 1997 (see also, Jakobovits, 1998, *Exp. Opin. Invest. Drugs* 7(4): 607-614; U.S. patents 6,162,963 issued 19 December 2000; 6,150,584 issued 12 November 2000; and, 6,114,598 issued 5 September 2000). This method avoids the *in vitro* manipulation required with phage display technology and efficiently produces high affinity authentic human antibodies.

Reactivity of 161P2F10B antibodies with a 161P2F10B-related protein can be established by a number of well known means, including Western blot, immunoprecipitation, ELISA, and FACS analyses using, as appropriate, 161P2F10B-related proteins, 161P2F10B-expressing cells or extracts thereof. A 161P2F10B antibody or fragment thereof can be labeled with a detectable marker or conjugated to a second molecule. Suitable detectable markers include, but are not limited to, a radioisotope, a fluorescent compound, a bioluminescent compound, chemiluminescent compound, a metal chelator or an enzyme. Further, bi-specific antibodies specific for two or more 161P2F10B epitopes are generated using methods generally known in the art. Homodimeric antibodies can also be generated by cross-linking techniques known in the art (e.g., Wolff *et al.*, *Cancer Res.* 53: 2560-2565).

V.) 161P2F10B Cellular Immune Responses

The mechanism by which T cells recognize antigens has been delineated. Efficacious peptide epitope vaccine compositions of the invention induce a therapeutic or prophylactic immune responses in very broad segments of the world-wide population. For an understanding of the value and efficacy of compositions of the invention that induce cellular immune responses, a brief review of immunology-related technology is provided.

A complex of an HLA molecule and a peptidic antigen acts as the ligand recognized by HLA-restricted T cells (Buus, S. *et al.*, *Cell* 47:1071, 1986; Babbitt, B. P. *et al.*, *Nature* 317:359, 1985; Townsend, A. and Bodmer, H., *Annu. Rev. Immunol.* 7:601, 1989; Germain, R. N., *Annu. Rev. Immunol.* 11:403, 1993). Through the study of single amino acid substituted antigen analogs and the sequencing of endogenously bound, naturally processed peptides, critical residues that correspond to motifs required for specific binding to HLA antigen molecules have been identified and are set forth in Table IV (see also, e.g., Southwood, *et al.*, *J. Immunol.* 160:3363, 1998; Rammensee, *et al.*, *Immunogenetics* 41:178, 1995; Rammensee *et al.*, SYFPEITHI, access via World Wide Web at URL (134.2.96.221/scripts.hlaserver.dll/home.htm); Sette, A. and Sidney, J. *Curr. Opin. Immunol.* 10:478, 1998; Engelhard, V. H., *Curr. Opin. Immunol.* 6:13, 1994; Sette, A. and Grey, H. M., *Curr. Opin. Immunol.* 4:79, 1992; Sinigaglia, F. and Hammer, J. *Curr. Biol.* 6:52, 1994; Ruppert *et al.*, *Cell* 74:929-937,

1993; Kondo *et al.*, *J. Immunol.* 155:4307-4312, 1995; Sidney *et al.*, *J. Immunol.* 157:3480-3490, 1996; Sidney *et al.*, *Human Immunol.* 45:79-93, 1996; Sette, A. and Sidney, J. *Immunogenetics* 1999 Nov; 50(3-4):201-12, Review).

Furthermore, x-ray crystallographic analyses of HLA-peptide complexes have revealed pockets within the peptide binding cleft/groove of HLA molecules which accommodate, in an allele-specific mode, residues borne by peptide ligands; these residues in turn determine the HLA binding capacity of the peptides in which they are present. (See, e.g., Madden, D.R. *Annu. Rev. Immunol.* 13:587, 1995; Smith, *et al.*, *Immunity* 4:203, 1996; Fremont *et al.*, *Immunity* 8:305, 1998; Stern *et al.*, *Structure* 2:245, 1994; Jones, E.Y. *Curr. Opin. Immunol.* 9:75, 1997; Brown, J. H. *et al.*, *Nature* 364:33, 1993; Guo, H. C. *et al.*, *Proc. Natl. Acad. Sci. USA* 90:8053, 1993; Guo, H. C. *et al.*, *Nature* 360:364, 1992; Silver, M. L. *et al.*, *Nature* 360:367, 1992; Matsumura, M. *et al.*, *Science* 257:927, 1992; Madden *et al.*, *Cell* 70:1035, 1992; Fremont, D. H. *et al.*, *Science* 257:919, 1992; Saper, M. A., Bjorkman, P. J. and Wiley, D. C., *J. Mol. Biol.* 219:277, 1991.)

Accordingly, the definition of class I and class II allele-specific HLA binding motifs, or class I or class II supermotifs allows identification of regions within a protein that are correlated with binding to particular HLA antigen(s).

Thus, by a process of HLA motif identification, candidates for epitope-based vaccines have been identified; such candidates can be further evaluated by HLA-peptide binding assays to determine binding affinity and/or the time period of association of the epitope and its corresponding HLA molecule. Additional confirmatory work can be performed to select, amongst these vaccine candidates, epitopes with preferred characteristics in terms of population coverage, and/or immunogenicity.

Various strategies can be utilized to evaluate cellular immunogenicity, including:

- 1) Evaluation of primary T cell cultures from normal individuals (see, e.g., Wentworth, P. A. *et al.*, *Mol. Immunol.* 32:603, 1995; Celis, E. *et al.*, *Proc. Natl. Acad. Sci. USA* 91:2105, 1994; Tsai, V. *et al.*, *J. Immunol.* 158:1796, 1997; Kawashima, I. *et al.*, *Human Immunol.* 59:1, 1998). This procedure involves the stimulation of peripheral blood lymphocytes (PBL) from normal subjects with a test peptide in the presence of antigen presenting cells *in vitro* over a period of several weeks. T cells specific for the peptide become activated during this time and are detected using, e.g., a lymphokine- or ⁵¹Cr-release assay involving peptide sensitized target cells.
- 2) Immunization of HLA transgenic mice (see, e.g., Wentworth, P. A. *et al.*, *J. Immunol.* 26:97, 1996; Wentworth, P. A. *et al.*, *Int. Immunol.* 8:651, 1996; Alexander, J. *et al.*, *J. Immunol.* 159:4753, 1997). For example, in such methods peptides in incomplete Freund's adjuvant are administered subcutaneously to HLA transgenic mice. Several weeks following immunization, splenocytes are removed and cultured *in vitro* in the presence of test peptide for approximately one week. Peptide-specific T cells are detected using, e.g., a ⁵¹Cr-release assay involving peptide sensitized target cells and target cells expressing endogenously generated antigen.
- 3) Demonstration of recall T cell responses from immune individuals who have been either effectively vaccinated and/or from chronically ill patients (see, e.g., Rehmann, B. *et al.*, *J. Exp. Med.* 181:1047, 1995; Doolan, D. L. *et al.*, *Immunity* 7:97, 1997; Bertoni, R. *et al.*, *J. Clin. Invest.* 100:503, 1997; Threlkeld, S. C. *et al.*, *J. Immunol.* 159:1648, 1997; Diepolder, H. M. *et al.*, *J. Virol.* 71:6011, 1997). Accordingly, recall responses are detected by culturing PBL from subjects that have been exposed to the antigen due to disease and thus have generated an immune response "naturally", or from patients who were vaccinated against the antigen. PBL from subjects are cultured *in vitro* for 1-2 weeks in the presence of test peptide plus antigen presenting cells (APC) to allow activation of "memory" T cells, as compared to "naive" T cells. At the end of the culture period, T cell activity is detected using assays including ⁵¹Cr release involving peptide-sensitized targets, T cell proliferation, or lymphokine release.

Nucleic acids that encode a 161P2F10B-related protein can also be used to generate either transgenic animals or "knock out" animals that, in turn, are useful in the development and screening of therapeutically useful reagents. In accordance with established techniques, cDNA encoding 161P2F10B can be used to clone genomic DNA that encodes 161P2F10B. The cloned genomic sequences can then be used to generate transgenic animals containing cells that express DNA that encode 161P2F10B. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 issued 12 April 1988, and 4,870,009 issued 26 September 1989. Typically, particular cells would be targeted for 161P2F10B transgene incorporation with tissue-specific enhancers.

Transgenic animals that include a copy of a transgene encoding 161P2F10B can be used to examine the effect of increased expression of DNA that encodes 161P2F10B. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this aspect of the invention, an animal is treated with a reagent and a reduced incidence of a pathological condition, compared to untreated animals that bear the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of 161P2F10B can be used to construct a 161P2F10B "knock out" animal that has a defective or altered gene encoding 161P2F10B as a result of homologous recombination between the endogenous gene encoding 161P2F10B and altered genomic DNA encoding 161P2F10B introduced into an embryonic cell of the animal. For example, cDNA that encodes 161P2F10B can be used to clone genomic DNA encoding 161P2F10B in accordance with established techniques. A portion of the genomic DNA encoding 161P2F10B can be deleted or replaced with another gene, such as a gene encoding a selectable marker that can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector (see, e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected (see, e.g., Li *et al.*, *Cell*, 69:915 (1992)). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras (see, e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal, and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock out animals can be characterized, for example, for their ability to defend against certain pathological conditions or for their development of pathological conditions due to absence of a 161P2F10B polypeptide.

VII.) Methods for the Detection of 161P2F10B

Another aspect of the present invention relates to methods for detecting 161P2F10B polynucleotides and 161P2F10B-related proteins, as well as methods for identifying a cell that expresses 161P2F10B. The expression profile of 161P2F10B makes it a diagnostic marker for metastasized disease. Accordingly, the status of 161P2F10B gene products provides information useful for predicting a variety of factors including susceptibility to advanced stage disease, rate of progression, and/or tumor aggressiveness. As discussed in detail herein, the status of 161P2F10B gene products in patient samples can be analyzed by a variety of protocols that are well known in the art including immunohistochemical analysis, the variety of Northern blotting techniques including *in situ* hybridization, RT-PCR analysis (for example on laser capture micro-dissected samples), Western blot analysis and tissue array analysis.

More particularly, the invention provides assays for the detection of 161P2F10B polynucleotides in a biological sample, such as serum, bone, prostate, and other tissues, urine, semen, cell preparations, and the like. Detectable 161P2F10B

polynucleotides include, for example, a 161P2F10B gene or fragment thereof, 161P2F10B mRNA, alternative splice variant 161P2F10B mRNAs, and recombinant DNA or RNA molecules that contain a 161P2F10B polynucleotide. A number of methods for amplifying and/or detecting the presence of 161P2F10B polynucleotides are well known in the art and can be employed in the practice of this aspect of the invention.

In one embodiment, a method for detecting a 161P2F10B mRNA in a biological sample comprises producing cDNA from the sample by reverse transcription using at least one primer; amplifying the cDNA so produced using a 161P2F10B polynucleotides as sense and antisense primers to amplify 161P2F10B cDNAs therein; and detecting the presence of the amplified 161P2F10B cDNA. Optionally, the sequence of the amplified 161P2F10B cDNA can be determined.

In another embodiment, a method of detecting a 161P2F10B gene in a biological sample comprises first isolating genomic DNA from the sample; amplifying the isolated genomic DNA using 161P2F10B polynucleotides as sense and antisense primers; and detecting the presence of the amplified 161P2F10B gene. Any number of appropriate sense and antisense probe combinations can be designed from a 161P2F10B nucleotide sequence (see, e.g., Figure 2) and used for this purpose.

The invention also provides assays for detecting the presence of a 161P2F10B protein in a tissue or other biological sample such as serum, semen, bone, prostate, urine, cell preparations, and the like. Methods for detecting a 161P2F10B-related protein are also well known and include, for example, immunoprecipitation, immunohistochemical analysis, Western blot analysis, molecular binding assays, ELISA, ELIFA and the like. For example, a method of detecting the presence of a 161P2F10B-related protein in a biological sample comprises first contacting the sample with a 161P2F10B antibody, a 161P2F10B-reactive fragment thereof, or a recombinant protein containing an antigen-binding region of a 161P2F10B antibody; and then detecting the binding of 161P2F10B-related protein in the sample.

Methods for identifying a cell that expresses 161P2F10B are also within the scope of the invention. In one embodiment, an assay for identifying a cell that expresses a 161P2F10B gene comprises detecting the presence of 161P2F10B mRNA in the cell. Methods for the detection of particular mRNAs in cells are well known and include, for example, hybridization assays using complementary DNA probes (such as *in situ* hybridization using labeled 161P2F10B riboprobes, Northern blot and related techniques) and various nucleic acid amplification assays (such as RT-PCR using complementary primers specific for 161P2F10B, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like). Alternatively, an assay for identifying a cell that expresses a 161P2F10B gene comprises detecting the presence of 161P2F10B-related protein in the cell or secreted by the cell. Various methods for the detection of proteins are well known in the art and are employed for the detection of 161P2F10B-related proteins and cells that express 161P2F10B-related proteins.

161P2F10B expression analysis is also useful as a tool for identifying and evaluating agents that modulate 161P2F10B gene expression. For example, 161P2F10B expression is significantly upregulated in prostate cancer, and is expressed in cancers of the tissues listed in Table I. Identification of a molecule or biological agent that inhibits 161P2F10B expression or over-expression in cancer cells is of therapeutic value. For example, such an agent can be identified by using a screen that quantifies 161P2F10B expression by RT-PCR, nucleic acid hybridization or antibody binding.

VIII.) Methods for Monitoring the Status of 161P2F10B-related Genes and Their Products

Oncogenesis is known to be a multistep process where cellular growth becomes progressively dysregulated and cells progress from a normal physiological state to precancerous and then cancerous states (see, e.g., Alers *et al.*, Lab Invest. 77(5): 437-438 (1997) and Isaacs *et al.*, Cancer Surv. 23: 19-32 (1995)). In this context, examining a biological sample for evidence of dysregulated cell growth (such as aberrant 161P2F10B expression in cancers) allows for early detection of such aberrant physiology, before a pathologic state such as cancer has progressed to a stage that therapeutic options are more limited and or the prognosis is worse. In such examinations, the status of 161P2F10B in a biological

sample of interest can be compared, for example, to the status of 161P2F10B in a corresponding normal sample (e.g. a sample from that individual or alternatively another individual that is not affected by a pathology). An alteration in the status of 161P2F10B in the biological sample (as compared to the normal sample) provides evidence of dysregulated cellular growth. In addition to using a biological sample that is not affected by a pathology as a normal sample, one can also use a predetermined normative value such as a predetermined normal level of mRNA expression (see, e.g., Grever *et al.*, J. Comp. Neurol. 1996 Dec 9; 376(2): 306-14 and U.S. Patent No. 5,837,501) to compare 161P2F10B status in a sample.

The term "status" in this context is used according to its art accepted meaning and refers to the condition or state of a gene and its products. Typically, skilled artisans use a number of parameters to evaluate the condition or state of a gene and its products. These include, but are not limited to the location of expressed gene products (including the location of 161P2F10B expressing cells) as well as the level, and biological activity of expressed gene products (such as 161P2F10B mRNA, polynucleotides and polypeptides). Typically, an alteration in the status of 161P2F10B comprises a change in the location of 161P2F10B and/or 161P2F10B expressing cells and/or an increase in 161P2F10B mRNA and/or protein expression.

161P2F10B status in a sample can be analyzed by a number of means well known in the art, including without limitation, immunohistochemical analysis, *in situ* hybridization, RT-PCR analysis on laser capture micro-dissected samples, Western blot analysis, and tissue array analysis. Typical protocols for evaluating the status of a 161P2F10B gene and gene products are found, for example in Ausubel *et al.* eds., 1995, Current Protocols In Molecular Biology, Units 2 (Northern Blotting), 4 (Southern Blotting), 15 (Immunoblotting) and 18 (PCR Analysis). Thus, the status of 161P2F10B in a biological sample is evaluated by various methods utilized by skilled artisans including, but not limited to genomic Southern analysis (to examine, for example perturbations in a 161P2F10B gene), Northern analysis and/or PCR analysis of 161P2F10B mRNA (to examine, for example alterations in the polynucleotide sequences or expression levels of 161P2F10B mRNAs), and, Western and/or immunohistochemical analysis (to examine, for example alterations in polypeptide sequences, alterations in polypeptide localization within a sample, alterations in expression levels of 161P2F10B proteins and/or associations of 161P2F10B proteins with polypeptide binding partners). Detectable 161P2F10B polynucleotides include, for example, a 161P2F10B gene or fragment thereof, 161P2F10B mRNA, alternative splice variants, 161P2F10B mRNAs, and recombinant DNA or RNA molecules containing a 161P2F10B polynucleotide.

The expression profile of 161P2F10B makes it a diagnostic marker for local and/or metastasized disease, and provides information on the growth or oncogenic potential of a biological sample. In particular, the status of 161P2F10B provides information useful for predicting susceptibility to particular disease stages, progression, and/or tumor aggressiveness. The invention provides methods and assays for determining 161P2F10B status and diagnosing cancers that express 161P2F10B, such as cancers of the tissues listed in Table I. For example, because 161P2F10B mRNA is so highly expressed in prostate and other cancers relative to normal prostate tissue, assays that evaluate the levels of 161P2F10B mRNA transcripts or proteins in a biological sample can be used to diagnose a disease associated with 161P2F10B dysregulation, and can provide prognostic information useful in defining appropriate therapeutic options.

The expression status of 161P2F10B provides information including the presence, stage and location of dysplastic, precancerous and cancerous cells, predicting susceptibility to various stages of disease, and/or for gauging tumor aggressiveness. Moreover, the expression profile makes it useful as an imaging reagent for metastasized disease. Consequently, an aspect of the invention is directed to the various molecular prognostic and diagnostic methods for examining the status of 161P2F10B in biological samples such as those from individuals suffering from, or suspected of suffering from a pathology characterized by dysregulated cellular growth, such as cancer.

As described above, the status of 161P2F10B in a biological sample can be examined by a number of well-known procedures in the art. For example, the status of 161P2F10B in a biological sample taken from a specific location in the body can be examined by evaluating the sample for the presence or absence of 161P2F10B expressing cells (e.g. those that

express 161P2F10B mRNAs or proteins). This examination can provide evidence of dysregulated cellular growth, for example, when 161P2F10B-expressing cells are found in a biological sample that does not normally contain such cells (such as a lymph node), because such alterations in the status of 161P2F10B in a biological sample are often associated with dysregulated cellular growth. Specifically, one indicator of dysregulated cellular growth is the metastases of cancer cells from an organ of origin (such as the prostate) to a different area of the body (such as a lymph node). In this context, evidence of dysregulated cellular growth is important for example because occult lymph node metastases can be detected in a substantial proportion of patients with prostate cancer, and such metastases are associated with known predictors of disease progression (see, e.g., Murphy *et al.*, Prostate 42(4): 315-317 (2000); Su *et al.*, Semin. Surg. Oncol. 18(1): 17-28 (2000) and Freeman *et al.*, J Urol 1995 Aug 154(2 Pt 1):474-8).

In one aspect, the invention provides methods for monitoring 161P2F10B gene products by determining the status of 161P2F10B gene products expressed by cells from an individual suspected of having a disease associated with dysregulated cell growth (such as hyperplasia or cancer) and then comparing the status so determined to the status of 161P2F10B gene products in a corresponding normal sample. The presence of aberrant 161P2F10B gene products in the test sample relative to the normal sample provides an indication of the presence of dysregulated cell growth within the cells of the individual.

In another aspect, the invention provides assays useful in determining the presence of cancer in an individual, comprising detecting a significant increase in 161P2F10B mRNA or protein expression in a test cell or tissue sample relative to expression levels in the corresponding normal cell or tissue. The presence of 161P2F10B mRNA can, for example, be evaluated in tissues including but not limited to those listed in Table I. The presence of significant 161P2F10B expression in any of these tissues is useful to indicate the emergence, presence and/or severity of a cancer, since the corresponding normal tissues do not express 161P2F10B mRNA or express it at lower levels.

In a related embodiment, 161P2F10B status is determined at the protein level rather than at the nucleic acid level. For example, such a method comprises determining the level of 161P2F10B protein expressed by cells in a test tissue sample and comparing the level so determined to the level of 161P2F10B expressed in a corresponding normal sample. In one embodiment, the presence of 161P2F10B protein is evaluated, for example, using immunohistochemical methods. 161P2F10B antibodies or binding partners capable of detecting 161P2F10B protein expression are used in a variety of assay formats well known in the art for this purpose.

In a further embodiment, one can evaluate the status of 161P2F10B nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules. These perturbations can include insertions, deletions, substitutions and the like. Such evaluations are useful because perturbations in the nucleotide and amino acid sequences are observed in a large number of proteins associated with a growth dysregulated phenotype (see, e.g., Marrogi *et al.*, 1999, J. Cutan. Pathol. 26(8):369-378). For example, a mutation in the sequence of 161P2F10B may be indicative of the presence or promotion of a tumor. Such assays therefore have diagnostic and predictive value where a mutation in 161P2F10B indicates a potential loss of function or increase in tumor growth.

A wide variety of assays for observing perturbations in nucleotide and amino acid sequences are well known in the art. For example, the size and structure of nucleic acid or amino acid sequences of 161P2F10B gene products are observed by the Northern, Southern, Western, PCR and DNA sequencing protocols discussed herein. In addition, other methods for observing perturbations in nucleotide and amino acid sequences such as single strand conformation polymorphism analysis are well known in the art (see, e.g., U.S. Patent Nos. 5,382,510 issued 7 September 1999, and 5,952,170 issued 17 January 1995).

Additionally, one can examine the methylation status of a 161P2F10B gene in a biological sample. Aberrant demethylation and/or hypermethylation of CpG islands in gene 5' regulatory regions frequently occurs in immortalized and transformed cells, and can result in altered expression of various genes. For example, promoter hypermethylation of the pi-class

glutathione S-transferase (a protein expressed in normal prostate but not expressed in >90% of prostate carcinomas) appears to permanently silence transcription of this gene and is the most frequently detected genomic alteration in prostate carcinomas (De Marzo *et al.*, Am. J. Pathol. 155(6): 1985-1992 (1999)). In addition, this alteration is present in at least 70% of cases of high-grade prostatic intraepithelial neoplasia (PIN) (Brooks *et al.*, Cancer Epidemiol. Biomarkers Prev., 1998, 7:531-536). In another example, expression of the LAGE-I tumor specific gene (which is not expressed in normal prostate but is expressed in 25-50% of prostate cancers) is induced by deoxy-azacytidine in lymphoblastoid cells, suggesting that tumoral expression is due to demethylation (Lethe *et al.*, Int. J. Cancer 76(6): 903-908 (1998)). A variety of assays for examining methylation status of a gene are well known in the art. For example, one can utilize, in Southern hybridization approaches, methylation-sensitive restriction enzymes that cannot cleave sequences that contain methylated CpG sites to assess the methylation status of CpG islands. In addition, MSP (methylation specific PCR) can rapidly profile the methylation status of all the CpG sites present in a CpG island of a given gene. This procedure involves initial modification of DNA by sodium bisulfite (which will convert all unmethylated cytosines to uracil) followed by amplification using primers specific for methylated versus unmethylated DNA. Protocols involving methylation interference can also be found for example in Current Protocols In Molecular Biology, Unit 12, Frederick M. Ausubel *et al.* eds., 1995.

Gene amplification is an additional method for assessing the status of 161P2F10B. Gene amplification is measured in a sample directly, for example, by conventional Southern blotting or Northern blotting to quantitate the transcription of mRNA (Thomas, 1980, Proc. Natl. Acad. Sci. USA, 77:5201-5205), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies are employed that recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn are labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Biopsied tissue or peripheral blood can be conveniently assayed for the presence of cancer cells using for example, Northern, dot blot or RT-PCR analysis to detect 161P2F10B expression. The presence of RT-PCR amplifiable 161P2F10B mRNA provides an indication of the presence of cancer. RT-PCR assays are well known in the art. RT-PCR detection assays for tumor cells in peripheral blood are currently being evaluated for use in the diagnosis and management of a number of human solid tumors. In the prostate cancer field, these include RT-PCR assays for the detection of cells expressing PSA and PSM (Verkaik *et al.*, 1997, Urol. Res. 25:373-384; Ghossein *et al.*, 1995, J. Clin. Oncol. 13:1195-2000; Heston *et al.*, 1995, Clin. Chem. 41:1687-1688).

A further aspect of the invention is an assessment of the susceptibility that an individual has for developing cancer. In one embodiment, a method for predicting susceptibility to cancer comprises detecting 161P2F10B mRNA or 161P2F10B protein in a tissue sample, its presence indicating susceptibility to cancer, wherein the degree of 161P2F10B mRNA expression correlates to the degree of susceptibility. In a specific embodiment, the presence of 161P2F10B in prostate or other tissue is examined, with the presence of 161P2F10B in the sample providing an indication of prostate cancer susceptibility (or the emergence or existence of a prostate tumor). Similarly, one can evaluate the integrity 161P2F10B nucleotide and amino acid sequences in a biological sample, in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like. The presence of one or more perturbations in 161P2F10B gene products in the sample is an indication of cancer susceptibility (or the emergence or existence of a tumor).

The invention also comprises methods for gauging tumor aggressiveness. In one embodiment, a method for gauging aggressiveness of a tumor comprises determining the level of 161P2F10B mRNA or 161P2F10B protein expressed by tumor cells, comparing the level so determined to the level of 161P2F10B mRNA or 161P2F10B protein expressed in a corresponding normal tissue taken from the same individual or a normal tissue reference sample, wherein the degree of 161P2F10B mRNA or 161P2F10B protein expression in the tumor sample relative to the normal sample indicates the degree of aggressiveness. In a

specific embodiment, aggressiveness of a tumor is evaluated by determining the extent to which 161P2F10B is expressed in the tumor cells, with higher expression levels indicating more aggressive tumors. Another embodiment is the evaluation of the integrity of 161P2F10B nucleotide and amino acid sequences in a biological sample, in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like. The presence of one or more perturbations indicates more aggressive tumors.

Another embodiment of the invention is directed to methods for observing the progression of a malignancy in an individual over time. In one embodiment, methods for observing the progression of a malignancy in an individual over time comprise determining the level of 161P2F10B mRNA or 161P2F10B protein expressed by cells in a sample of the tumor, comparing the level so determined to the level of 161P2F10B mRNA or 161P2F10B protein expressed in an equivalent tissue sample taken from the same individual at a different time, wherein the degree of 161P2F10B mRNA or 161P2F10B protein expression in the tumor sample over time provides information on the progression of the cancer. In a specific embodiment, the progression of a cancer is evaluated by determining 161P2F10B expression in the tumor cells over time, where increased expression over time indicates a progression of the cancer. Also, one can evaluate the integrity 161P2F10B nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like, where the presence of one or more perturbations indicates a progression of the cancer.

The above diagnostic approaches can be combined with any one of a wide variety of prognostic and diagnostic protocols known in the art. For example, another embodiment of the invention is directed to methods for observing a coincidence between the expression of 161P2F10B gene and 161P2F10B gene products (or perturbations in 161P2F10B gene and 161P2F10B gene products) and a factor that is associated with malignancy, as a means for diagnosing and prognosticating the status of a tissue sample. A wide variety of factors associated with malignancy can be utilized, such as the expression of genes associated with malignancy (e.g. PSA, PSCA and PSM expression for prostate cancer etc.) as well as gross cytological observations (see, e.g., Bocking *et al.*, 1984, *Anal. Quant. Cytol.* 6(2):74-88; Epstein, 1995, *Hum. Pathol.* 26(2):223-9; Thorson *et al.*, 1998, *Mod. Pathol.* 11(6):543-51; Baisden *et al.*, 1999, *Am. J. Surg. Pathol.* 23(8):918-24). Methods for observing a coincidence between the expression of 161P2F10B gene and 161P2F10B gene products (or perturbations in 161P2F10B gene and 161P2F10B gene products) and another factor that is associated with malignancy are useful, for example, because the presence of a set of specific factors that coincide with disease provides information crucial for diagnosing and prognosticating the status of a tissue sample.

In one embodiment, methods for observing a coincidence between the expression of 161P2F10B gene and 161P2F10B gene products (or perturbations in 161P2F10B gene and 161P2F10B gene products) and another factor associated with malignancy entails detecting the overexpression of 161P2F10B mRNA or protein in a tissue sample, detecting the overexpression of PSA mRNA or protein in a tissue sample (or PSCA or PSM expression), and observing a coincidence of 161P2F10B mRNA or protein and PSA mRNA or protein overexpression (or PSCA or PSM expression). In a specific embodiment, the expression of 161P2F10B and PSA mRNA in prostate tissue is examined, where the coincidence of 161P2F10B and PSA mRNA overexpression in the sample indicates the existence of prostate cancer, prostate cancer susceptibility or the emergence or status of a prostate tumor.

Methods for detecting and quantifying the expression of 161P2F10B mRNA or protein are described herein, and standard nucleic acid and protein detection and quantification technologies are well known in the art. Standard methods for the detection and quantification of 161P2F10B mRNA include *in situ* hybridization using labeled 161P2F10B riboprobes, Northern blot and related techniques using 161P2F10B polynucleotide probes, RT-PCR analysis using primers specific for 161P2F10B, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like. In a specific embodiment, semi-quantitative RT-PCR is used to detect and quantify 161P2F10B mRNA expression. Any number of primers capable of amplifying 161P2F10B can be used for this purpose, including but not limited to the various primer sets specifically

described herein. In a specific embodiment, polyclonal or monoclonal antibodies specifically reactive with the wild-type 161P2F10B protein can be used in an immunohistochemical assay of biopsied tissue.

IX.) Identification of Molecules That Interact With 161P2F10B

The 161P2F10B protein and nucleic acid sequences disclosed herein allow a skilled artisan to identify proteins, small molecules and other agents that interact with 161P2F10B, as well as pathways activated by 161P2F10B via any one of a variety of art accepted protocols. For example, one can utilize one of the so-called interaction trap systems (also referred to as the "two-hybrid assay"). In such systems, molecules interact and reconstitute a transcription factor which directs expression of a reporter gene, whereupon the expression of the reporter gene is assayed. Other systems identify protein-protein interactions *in vivo* through reconstitution of a eukaryotic transcriptional activator, see, e.g., U.S. Patent Nos. 5,955,280 issued 21 September 1999, 5,925,523 issued 20 July 1999, 5,846,722 issued 8 December 1998 and 6,004,746 issued 21 December 1999. Algorithms are also available in the art for genome-based predictions of protein function (see, e.g., Marcotte, *et al.*, Nature 402: 4 November 1999, 83-86).

Alternatively one can screen peptide libraries to identify molecules that interact with 161P2F10B protein sequences. In such methods, peptides that bind to 161P2F10B are identified by screening libraries that encode a random or controlled collection of amino acids. Peptides encoded by the libraries are expressed as fusion proteins of bacteriophage coat proteins, the bacteriophage particles are then screened against the 161P2F10B protein(s).

Accordingly, peptides having a wide variety of uses, such as therapeutic, prognostic or diagnostic reagents, are thus identified without any prior information on the structure of the expected ligand or receptor molecule. Typical peptide libraries and screening methods that can be used to identify molecules that interact with 161P2F10B protein sequences are disclosed for example in U.S. Patent Nos. 5,723,286 issued 3 March 1998 and 5,733,731 issued 31 March 1998.

Alternatively, cell lines that express 161P2F10B are used to identify protein-protein interactions mediated by 161P2F10B. Such interactions can be examined using immunoprecipitation techniques (see, e.g., Hamilton B.J., *et al.* Biochem. Biophys. Res. Commun. 1999, 261:646-51). 161P2F10B protein can be immunoprecipitated from 161P2F10B-expressing cell lines using anti-161P2F10B antibodies. Alternatively, antibodies against His-tag can be used in a cell line engineered to express fusions of 161P2F10B and a His-tag (vectors mentioned above). The immunoprecipitated complex can be examined for protein association by procedures such as Western blotting, ³⁵S-methionine labeling of proteins, protein microsequencing, silver staining and two-dimensional gel electrophoresis.

Small molecules and ligands that interact with 161P2F10B can be identified through related embodiments of such screening assays. For example, small molecules can be identified that interfere with protein function, including molecules that interfere with 161P2F10B's ability to mediate phosphorylation and de-phosphorylation, interaction with DNA or RNA molecules as an indication of regulation of cell cycles, second messenger signaling or tumorigenesis. Similarly, small molecules that modulate 161P2F10B-related ion channel, protein pump, or cell communication functions are identified and used to treat patients that have a cancer that expresses 161P2F10B (see, e.g., Hille, B., Ionic Channels of Excitable Membranes 2nd Ed., Sinauer Assoc., Sunderland, MA, 1992). Moreover, ligands that regulate 161P2F10B function can be identified based on their ability to bind 161P2F10B and activate a reporter construct. Typical methods are discussed for example in U.S. Patent No. 5,928,868 issued 27 July 1999, and include methods for forming hybrid ligands in which at least one ligand is a small molecule. In an illustrative embodiment, cells engineered to express a fusion protein of 161P2F10B and a DNA-binding protein are used to co-express a fusion protein of a hybrid ligand/small molecule and a cDNA library transcriptional activator protein. The cells further contain a reporter gene, the expression of which is conditioned on the proximity of the first and second fusion proteins to each other, an event that occurs only if the hybrid ligand binds to target sites on both hybrid proteins. Those cells that express the reporter gene are selected and the unknown small molecule or

the unknown ligand is identified. This method provides a means of identifying modulators, which activate or inhibit 161P2F10B.

An embodiment of this invention comprises a method of screening for a molecule that interacts with a 161P2F10B amino acid sequence shown in Figure 2 or Figure 3, comprising the steps of contacting a population of molecules with a 161P2F10B amino acid sequence, allowing the population of molecules and the 161P2F10B amino acid sequence to interact under conditions that facilitate an interaction, determining the presence of a molecule that interacts with the 161P2F10B amino acid sequence, and then separating molecules that do not interact with the 161P2F10B amino acid sequence from molecules that do. In a specific embodiment, the method further comprises purifying, characterizing and identifying a molecule that interacts with the 161P2F10B amino acid sequence. The identified molecule can be used to modulate a function performed by 161P2F10B. In a preferred embodiment, the 161P2F10B amino acid sequence is contacted with a library of peptides.

X.) Therapeutic Methods and Compositions

The identification of 161P2F10B as a protein that is normally expressed in a restricted set of tissues, but which is also expressed in prostate and other cancers, opens a number of therapeutic approaches to the treatment of such cancers. As contemplated herein, 161P2F10B functions as a transcription factor involved in activating tumor-promoting genes or repressing genes that block tumorigenesis.

Accordingly, therapeutic approaches that inhibit the activity of a 161P2F10B protein are useful for patients suffering from a cancer that expresses 161P2F10B. These therapeutic approaches generally fall into two classes. One class comprises various methods for inhibiting the binding or association of a 161P2F10B protein with its binding partner or with other proteins. Another class comprises a variety of methods for inhibiting the transcription of a 161P2F10B gene or translation of 161P2F10B mRNA.

X.A.) Anti-Cancer Vaccines

The invention provides cancer vaccines comprising a 161P2F10B-related protein or 161P2F10B-related nucleic acid. In view of the expression of 161P2F10B, cancer vaccines prevent and/or treat 161P2F10B-expressing cancers with minimal or no effects on non-target tissues. The use of a tumor antigen in a vaccine that generates humoral and/or cell-mediated immune responses as anti-cancer therapy is well known in the art and has been employed in prostate cancer using human PSMA and rodent PAP immunogens (Hodge *et al.*, 1995, *Int. J. Cancer* 63:231-237; Fong *et al.*, 1997, *J. Immunol.* 159:3113-3117).

Such methods can be readily practiced by employing a 161P2F10B-related protein, or a 161P2F10B-encoding nucleic acid molecule and recombinant vectors capable of expressing and presenting the 161P2F10B immunogen (which typically comprises a number of antibody or T cell epitopes). Skilled artisans understand that a wide variety of vaccine systems for delivery of immunoreactive epitopes are known in the art (see, e.g., Heryln *et al.*, *Ann Med* 1999 Feb 31(1):66-78; Maruyama *et al.*, *Cancer Immunol Immunother* 2000 Jun 49(3):123-32). Briefly, such methods of generating an immune response (e.g. humoral and/or cell-mediated) in a mammal, comprise the steps of: exposing the mammal's immune system to an immunoreactive epitope (e.g. an epitope present in a 161P2F10B protein shown in Figure 3 or analog or homolog thereof) so that the mammal generates an immune response that is specific for that epitope (e.g. generates antibodies that specifically recognize that epitope). In a preferred method, a 161P2F10B immunogen contains a biological motif, see e.g., Tables VIII-XXI and XXII-XLIX, or a peptide of a size range from 161P2F10B indicated in Figure 5, Figure 6, Figure 7, Figure 8, and Figure 9.

The entire 161P2F10B protein, immunogenic regions or epitopes thereof can be combined and delivered by various means. Such vaccine compositions can include, for example, lipopeptides (e.g., Vitiello, A. *et al.*, *J. Clin. Invest.*

95:341, 1995), peptide compositions encapsulated in poly(DL-lactide-co-glycolide) ("PLG") microspheres (see, e.g., Eldridge, *et al.*, *Molec. Immunol.* 28:287-294, 1991; Alonso *et al.*, *Vaccine* 12:299-306, 1994; Jones *et al.*, *Vaccine* 13:675-681, 1995), peptide compositions contained in immune stimulating complexes (ISCOMS) (see, e.g., Takahashi *et al.*, *Nature* 344:873-875, 1990; Hu *et al.*, *Clin Exp Immunol.* 113:235-243, 1998), multiple antigen peptide systems (MAPs) (see e.g., Tam, J. P., *Proc. Natl. Acad. Sci. U.S.A.* 85:5409-5413, 1988; Tam, J.P., *J. Immunol. Methods* 196:17-32, 1996), peptides formulated as multivalent peptides; peptides for use in ballistic delivery systems, typically crystallized peptides, viral delivery vectors (Perkus, M. E. *et al.*, In: *Concepts in vaccine development*, Kaufmann, S. H. E., ed., p. 379, 1996; Chakrabarti, S. *et al.*, *Nature* 320:535, 1986; Hu, S. L. *et al.*, *Nature* 320:537, 1986; Kieny, M.-P. *et al.*, *AIDS Bio/Technology* 4:790, 1986; Top, F. H. *et al.*, *J. Infect. Dis.* 124:148, 1971; Chanda, P. K. *et al.*, *Virology* 175:535, 1990), particles of viral or synthetic origin (e.g., Kofler, N. *et al.*, *J. Immunol. Methods* 192:25, 1996; Eldridge, J. H. *et al.*, *Sem. Hematol.* 30:16, 1993; Falo, L. D., Jr. *et al.*, *Nature Med.* 7:649, 1995), adjuvants (Warren, H. S., Vogel, F. R., and Chedid, L. A. *Annu. Rev. Immunol.* 4:369, 1986; Gupta, R. K. *et al.*, *Vaccine* 11:293, 1993), liposomes (Reddy, R. *et al.*, *J. Immunol.* 148:1585, 1992; Rock, K. L., *Immunol. Today* 17:131, 1996), or, naked or particle absorbed cDNA (Ulmer, J. B. *et al.*, *Science* 259:1745, 1993; Robinson, H. L., Hunt, L. A., and Webster, R. G., *Vaccine* 11:957, 1993; Shiver, J. W. *et al.*, In: *Concepts in vaccine development*, Kaufmann, S. H. E., ed., p. 423, 1996; Cease, K. B., and Berzofsky, J. A., *Annu. Rev. Immunol.* 12:923, 1994 and Eldridge, J. H. *et al.*, *Sem. Hematol.* 30:16, 1993). Toxin-targeted delivery technologies, also known as receptor mediated targeting, such as those of Avant Immunotherapeutics, Inc. (Needham, Massachusetts) may also be used.

In patients with 161P2F10B-associated cancer, the vaccine compositions of the invention can also be used in conjunction with other treatments used for cancer, e.g., surgery, chemotherapy, drug therapies, radiation therapies, *etc.* including use in combination with immune adjuvants such as IL-2, IL-12, GM-CSF, and the like.

Cellular Vaccines:

CTL epitopes can be determined using specific algorithms to identify peptides within 161P2F10B protein that bind corresponding HLA alleles (see e.g., Table IV; Epimer™ and Epimatrix™, Brown University (URL brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html); and, BIMAS, (URL bimas.dcrt.nih.gov/; SYFPEITHI at URL syfpeithi.bmi-heidelberg.com/). In a preferred embodiment, a 161P2F10B immunogen contains one or more amino acid sequences identified using techniques well known in the art, such as the sequences shown in Tables VIII-XXI and XXII-XLIX or a peptide of 8, 9, 10 or 11 amino acids specified by an HLA Class I motif/supermotif (e.g., Table IV (A), Table IV (D), or Table IV (E)) and/or a peptide of at least 9 amino acids that comprises an HLA Class II motif/supermotif (e.g., Table IV (B) or Table IV (C)). As is appreciated in the art, the HLA Class I binding groove is essentially closed ended so that peptides of only a particular size range can fit into the groove and be bound, generally HLA Class I epitopes are 8, 9, 10, or 11 amino acids long. In contrast, the HLA Class II binding groove is essentially open ended; therefore a peptide of about 9 or more amino acids can be bound by an HLA Class II molecule. Due to the binding groove differences between HLA Class I and II, HLA Class I motifs are length specific, i.e., position two of a Class I motif is the second amino acid in an amino to carboxyl direction of the peptide. The amino acid positions in a Class II motif are relative only to each other, not the overall peptide, i.e., additional amino acids can be attached to the amino and/or carboxyl termini of a motif-bearing sequence. HLA Class II epitopes are often 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 amino acids long, or longer than 25 amino acids.

Antibody-based Vaccines

A wide variety of methods for generating an immune response in a mammal are known in the art (for example as the first step in the generation of hybridomas). Methods of generating an immune response in a mammal comprise exposing the mammal's immune system to an immunogenic epitope on a protein (e.g. a 161P2F10B protein) so that an immune response is generated. A typical embodiment consists of a method for generating an immune response to 161P2F10B in a host, by contacting the host with a sufficient amount of at least one 161P2F10B B cell or cytotoxic T-cell epitope or analog

thereof; and at least one periodic interval thereafter re-contacting the host with the 161P2F10B B cell or cytotoxic T-cell epitope or analog thereof. A specific embodiment consists of a method of generating an immune response against a 161P2F10B-related protein or a man-made multi-epitopic peptide comprising: administering 161P2F10B immunogen (e.g. a 161P2F10B protein or a peptide fragment thereof, a 161P2F10B fusion protein or analog etc.) in a vaccine preparation to a human or another mammal. Typically, such vaccine preparations further contain a suitable adjuvant (see, e.g., U.S. Patent No. 6,146,635) or a universal helper epitope such as a PADRE™ peptide (Epimmune Inc., San Diego, CA; see, e.g., Alexander *et al.*, J. Immunol. 2000 164(3): 1625-1633; Alexander *et al.*, Immunity 1994 1(9): 751-761 and Alexander *et al.*, Immunol. Res. 1998 18(2): 79-92). An alternative method comprises generating an immune response in an individual against a 161P2F10B immunogen by: administering *in vivo* to muscle or skin of the individual's body a DNA molecule that comprises a DNA sequence that encodes a 161P2F10B immunogen, the DNA sequence operatively linked to regulatory sequences which control the expression of the DNA sequence; wherein the DNA molecule is taken up by cells, the DNA sequence is expressed in the cells and an immune response is generated against the immunogen (see, e.g., U.S. Patent No. 5,962,428). Optionally a genetic vaccine facilitator such as anionic lipids; saponins; lectins; estrogenic compounds; hydroxylated lower alkyls; dimethyl sulfoxide; and urea is also administered. In addition, an anti-idiotypic antibody can be administered that mimics 161P2F10B, in order to generate a response to the target antigen.

Nucleic Acid Vaccines:

Vaccine compositions of the invention include nucleic acid-mediated modalities. DNA or RNA that encode protein(s) of the invention can be administered to a patient. Genetic immunization methods can be employed to generate prophylactic or therapeutic humoral and cellular immune responses directed against cancer cells expressing 161P2F10B. Constructs comprising DNA encoding a 161P2F10B-related protein/immunogen and appropriate regulatory sequences can be injected directly into muscle or skin of an individual, such that the cells of the muscle or skin take-up the construct and express the encoded 161P2F10B protein/immunogen. Alternatively, a vaccine comprises a 161P2F10B-related protein. Expression of the 161P2F10B-related protein immunogen results in the generation of prophylactic or therapeutic humoral and cellular immunity against cells that bear a 161P2F10B protein. Various prophylactic and therapeutic genetic immunization techniques known in the art can be used (for review, see information and references published at Internet address genweb.com). Nucleic acid-based delivery is described, for instance, in Wolff *et al.*, *Science* 247:1465 (1990) as well as U.S. Patent Nos. 5,580,859; 5,589,466; 5,804,566; 5,739,118; 5,736,524; 5,679,647; WO 98/04720. Examples of DNA-based delivery technologies include "naked DNA", facilitated (bupivacaine, polymers, peptide-mediated) delivery, cationic lipid complexes, and particle-mediated ("gene gun") or pressure-mediated delivery (see, e.g., U.S. Patent No. 5,922,687).

For therapeutic or prophylactic immunization purposes, proteins of the invention can be expressed via viral or bacterial vectors. Various viral gene delivery systems that can be used in the practice of the invention include, but are not limited to, vaccinia, fowlpox, canarypox, adenovirus, influenza, poliovirus, adeno-associated virus, lentivirus, and sindbis virus (see, e.g., Restifo, 1996, *Curr. Opin. Immunol.* 8:658-663; Tsang *et al.* *J. Natl. Cancer Inst.* 87:982-990 (1995)). Non-viral delivery systems can also be employed by introducing naked DNA encoding a 161P2F10B-related protein into the patient (e.g., intramuscularly or intradermally) to induce an anti-tumor response.

Vaccinia virus is used, for example, as a vector to express nucleotide sequences that encode the peptides of the invention. Upon introduction into a host, the recombinant vaccinia virus expresses the protein immunogenic peptide, and thereby elicits a host immune response. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Patent No. 4,722,848. Another vector is BCG (Bacille Calmette Guérin). BCG vectors are described in Stover *et al.*, *Nature* 351:456-460 (1991). A wide variety of other vectors useful for therapeutic administration or immunization of the

peptides of the invention, e.g. adeno and adeno-associated virus vectors, retroviral vectors, *Salmonella typhi* vectors, detoxified anthrax toxin vectors, and the like, will be apparent to those skilled in the art from the description herein.

Thus, gene delivery systems are used to deliver a 161P2F10B-related nucleic acid molecule. In one embodiment, the full-length human 161P2F10B cDNA is employed. In another embodiment, 161P2F10B nucleic acid molecules encoding specific cytotoxic T lymphocyte (CTL) and/or antibody epitopes are employed.

Ex Vivo Vaccines

Various ex vivo strategies can also be employed to generate an immune response. One approach involves the use of antigen presenting cells (APCs) such as dendritic cells (DC) to present 161P2F10B antigen to a patient's immune system. Dendritic cells express MHC class I and II molecules, B7 co-stimulator, and IL-12, and are thus highly specialized antigen presenting cells. In prostate cancer, autologous dendritic cells pulsed with peptides of the prostate-specific membrane antigen (PSMA) are being used in a Phase I clinical trial to stimulate prostate cancer patients' immune systems (Tjoa *et al.*, 1996, Prostate 28:65-69; Murphy *et al.*, 1996, Prostate 29:371-380). Thus, dendritic cells can be used to present 161P2F10B peptides to T cells in the context of MHC class I or II molecules. In one embodiment, autologous dendritic cells are pulsed with 161P2F10B peptides capable of binding to MHC class I and/or class II molecules. In another embodiment, dendritic cells are pulsed with the complete 161P2F10B protein. Yet another embodiment involves engineering the overexpression of a 161P2F10B gene in dendritic cells using various implementing vectors known in the art, such as adenovirus (Arthur *et al.*, 1997, Cancer Gene Ther. 4:17-25), retrovirus (Henderson *et al.*, 1996, Cancer Res. 56:3763-3770), lentivirus, adeno-associated virus, DNA transfection (Ribas *et al.*, 1997, Cancer Res. 57:2865-2869), or tumor-derived RNA transfection (Ashley *et al.*, 1997, J. Exp. Med. 186:1177-1182). Cells that express 161P2F10B can also be engineered to express immune modulators, such as GM-CSF, and used as immunizing agents.

X.B.) 161P2F10B as a Target for Antibody-based Therapy

161P2F10B is an attractive target for antibody-based therapeutic strategies. A number of antibody strategies are known in the art for targeting both extracellular and intracellular molecules (see, e.g., complement and ADCC mediated killing as well as the use of intrabodies). Because 161P2F10B is expressed by cancer cells of various lineages relative to corresponding normal cells, systemic administration of 161P2F10B-immunoreactive compositions are prepared that exhibit excellent sensitivity without toxic, non-specific and/or non-target effects caused by binding of the immunoreactive composition to non-target organs and tissues. Antibodies specifically reactive with domains of 161P2F10B are useful to treat 161P2F10B-expressing cancers systemically, either as conjugates with a toxin or therapeutic agent, or as naked antibodies capable of inhibiting cell proliferation or function.

161P2F10B antibodies can be introduced into a patient such that the antibody binds to 161P2F10B and modulates a function, such as an interaction with a binding partner, and consequently mediates destruction of the tumor cells and/or inhibits the growth of the tumor cells. Mechanisms by which such antibodies exert a therapeutic effect can include complement-mediated cytotoxicity, antibody-dependent cellular cytotoxicity, modulation of the physiological function of 161P2F10B, inhibition of ligand binding or signal transduction pathways, modulation of tumor cell differentiation, alteration of tumor angiogenesis factor profiles, and/or apoptosis.

Those skilled in the art understand that antibodies can be used to specifically target and bind immunogenic molecules such as an immunogenic region of a 161P2F10B sequence shown in Figure 2 or Figure 3. In addition, skilled artisans understand that it is routine to conjugate antibodies to cytotoxic agents (see, e.g., Slevens *et al.* Blood 93:11 3678-3684 (June 1, 1999)). When cytotoxic and/or therapeutic agents are delivered directly to cells, such as by conjugating them to antibodies specific for a molecule expressed by that cell (e.g. 161P2F10B), the cytotoxic agent will exert its known biological effect (i.e. cytotoxicity) on those cells.

A wide variety of compositions and methods for using antibody-cytotoxic agent conjugates to kill cells are known in the art. In the context of cancers, typical methods entail administering to an animal having a tumor a biologically effective amount of a conjugate comprising a selected cytotoxic and/or therapeutic agent linked to a targeting agent (e.g. an anti-161P2F10B antibody) that binds to a marker (e.g. 161P2F10B) expressed, accessible to binding or localized on the cell surfaces. A typical embodiment is a method of delivering a cytotoxic and/or therapeutic agent to a cell expressing 161P2F10B, comprising conjugating the cytotoxic agent to an antibody that immunospecifically binds to a 161P2F10B epitope, and, exposing the cell to the antibody-agent conjugate. Another illustrative embodiment is a method of treating an individual suspected of suffering from metastasized cancer, comprising a step of administering parenterally to said individual a pharmaceutical composition comprising a therapeutically effective amount of an antibody conjugated to a cytotoxic and/or therapeutic agent.

Cancer immunotherapy using anti-161P2F10B antibodies can be done in accordance with various approaches that have been successfully employed in the treatment of other types of cancer, including but not limited to colon cancer (Arlen *et al.*, 1998, Crit. Rev. Immunol. 18:133-138), multiple myeloma (Ozaki *et al.*, 1997, Blood 90:3179-3186, Tsunenari *et al.*, 1997, Blood 90:2437-2444), gastric cancer (Kasprzyk *et al.*, 1992, Cancer Res. 52:2771-2776), B-cell lymphoma (Funakoshi *et al.*, 1996, J. Immunother. Emphasis Tumor Immunol. 19:93-101), leukemia (Zhong *et al.*, 1996, Leuk. Res. 20:581-589), colorectal cancer (Moun *et al.*, 1994, Cancer Res. 54:6160-6166; Velders *et al.*, 1995, Cancer Res. 55:4398-4403), and breast cancer (Shepard *et al.*, 1991, J. Clin. Immunol. 11:117-127). Some therapeutic approaches involve conjugation of naked antibody to a toxin or radioisotope, such as the conjugation of Y⁹¹ or I¹³¹ to anti-CD20 antibodies (e.g., ZevalinTM, IDEC Pharmaceuticals Corp. or BexxarTM, Coulter Pharmaceuticals), while others involve co-administration of antibodies and other therapeutic agents, such as HerceptinTM (trastuzumab) with paclitaxel (Genentech, Inc.). The antibodies can be conjugated to a therapeutic agent. To treat prostate cancer, for example, 161P2F10B antibodies can be administered in conjunction with radiation, chemotherapy or hormone ablation. Also, antibodies can be conjugated to a toxin such as calicheamicin (e.g., MylotargTM, Wyeth-Ayerst, Madison, NJ, a recombinant humanized IgG₄ kappa antibody conjugated to antitumor antibiotic calicheamicin) or a maytansinoid (e.g., taxane-based Tumor-Activated Prodrug, TAP, platform, ImmunoGen, Cambridge, MA, also see e.g., US Patent 5,416,064).

Although 161P2F10B antibody therapy is useful for all stages of cancer, antibody therapy can be particularly appropriate in advanced or metastatic cancers. Treatment with the antibody therapy of the invention is indicated for patients who have received one or more rounds of chemotherapy. Alternatively, antibody therapy of the invention is combined with a chemotherapeutic or radiation regimen for patients who have not received chemotherapeutic treatment. Additionally, antibody therapy can enable the use of reduced dosages of concomitant chemotherapy, particularly for patients who do not tolerate the toxicity of the chemotherapeutic agent very well. Fan *et al.* (Cancer Res. 53:4637-4642, 1993), Prewett *et al.* (International J. of Onco. 9:217-224, 1996), and Hancock *et al.* (Cancer Res. 51:4575-4580, 1991) describe the use of various antibodies together with chemotherapeutic agents.

Although 161P2F10B antibody therapy is useful for all stages of cancer, antibody therapy can be particularly appropriate in advanced or metastatic cancers. Treatment with the antibody therapy of the invention is indicated for patients who have received one or more rounds of chemotherapy. Alternatively, antibody therapy of the invention is combined with a chemotherapeutic or radiation regimen for patients who have not received chemotherapeutic treatment. Additionally, antibody therapy can enable the use of reduced dosages of concomitant chemotherapy, particularly for patients who do not tolerate the toxicity of the chemotherapeutic agent very well.

Cancer patients can be evaluated for the presence and level of 161P2F10B expression, preferably using immunohistochemical assessments of tumor tissue, quantitative 161P2F10B imaging, or other techniques that reliably indicate the presence and degree of 161P2F10B expression. Immunohistochemical analysis of tumor biopsies or surgical

specimens is preferred for this purpose. Methods for immunohistochemical analysis of tumor tissues are well known in the art.

Anti-161P2F10B monoclonal antibodies that treat prostate and other cancers include those that initiate a potent immune response against the tumor or those that are directly cytotoxic. In this regard, anti-161P2F10B monoclonal antibodies (mAbs) can elicit tumor cell lysis by either complement-mediated or antibody-dependent cell cytotoxicity (ADCC) mechanisms, both of which require an intact Fc portion of the immunoglobulin molecule for interaction with effector cell Fc receptor sites on complement proteins. In addition, anti-161P2F10B mAbs that exert a direct biological effect on tumor growth are useful to treat cancers that express 161P2F10B. Mechanisms by which directly cytotoxic mAbs act include: inhibition of cell growth, modulation of cellular differentiation, modulation of tumor angiogenesis factor profiles, and the induction of apoptosis. The mechanism(s) by which a particular anti-161P2F10B mAb exerts an anti-tumor effect is evaluated using any number of *in vitro* assays that evaluate cell death such as ADCC, ADMMC, complement-mediated cell lysis, and so forth, as is generally known in the art.

In some patients, the use of murine or other non-human monoclonal antibodies, or human/mouse chimeric mAbs can induce moderate to strong immune responses against the non-human antibody. This can result in clearance of the antibody from circulation and reduced efficacy. In the most severe cases, such an immune response can lead to the extensive formation of immune complexes which, potentially, can cause renal failure. Accordingly, preferred monoclonal antibodies used in the therapeutic methods of the invention are those that are either fully human or humanized and that bind specifically to the target 161P2F10B antigen with high affinity but exhibit low or no antigenicity in the patient.

Therapeutic methods of the invention contemplate the administration of single anti-161P2F10B mAbs as well as combinations, or cocktails, of different mAbs. Such mAb cocktails can have certain advantages inasmuch as they contain mAbs that target different epitopes, exploit different effector mechanisms or combine directly cytotoxic mAbs with mAbs that rely on immune effector functionality. Such mAbs in combination can exhibit synergistic therapeutic effects. In addition, anti-161P2F10B mAbs can be administered concomitantly with other therapeutic modalities, including but not limited to various chemotherapeutic agents, androgen-blockers, immune modulators (e.g., IL-2, GM-CSF), surgery or radiation. The anti-161P2F10B mAbs are administered in their "naked" or unconjugated form, or can have a therapeutic agent(s) conjugated to them.

Anti-161P2F10B antibody formulations are administered via any route capable of delivering the antibodies to a tumor cell. Routes of administration include, but are not limited to, intravenous, intraperitoneal, intramuscular, intratumor, intradermal, and the like. Treatment generally involves repeated administration of the anti-161P2F10B antibody preparation, via an acceptable route of administration such as intravenous injection (IV), typically at a dose in the range of about 0.1, .2, .3, .4, .5, .6, .7, .8, .9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or 25 mg/kg body weight. In general, doses in the range of 10-1000 mg mAb per week are effective and well tolerated.

Based on clinical experience with the Herceptin™ mAb in the treatment of metastatic breast cancer, an initial loading dose of approximately 4 mg/kg patient body weight IV, followed by weekly doses of about 2 mg/kg IV of the anti-161P2F10B mAb preparation represents an acceptable dosing regimen. Preferably, the initial loading dose is administered as a 90-minute or longer infusion. The periodic maintenance dose is administered as a 30 minute or longer infusion, provided the initial dose was well tolerated. As appreciated by those of skill in the art, various factors can influence the ideal dose regimen in a particular case. Such factors include, for example, the binding affinity and half life of the Ab or mAbs used, the degree of 161P2F10B expression in the patient, the extent of circulating shed 161P2F10B antigen, the desired steady-state antibody concentration level, frequency of treatment, and the influence of chemotherapeutic or other agents used in combination with the treatment method of the invention, as well as the health status of a particular patient.

Optionally, patients should be evaluated for the levels of 161P2F10B in a given sample (e.g. the levels of circulating 161P2F10B antigen and/or 161P2F10B expressing cells) in order to assist in the determination of the most effective dosing regimen, etc. Such evaluations are also used for monitoring purposes throughout therapy, and are useful to gauge therapeutic success in combination with the evaluation of other parameters (for example, urine cytology and/or ImmunoCyt levels in bladder cancer therapy, or by analogy, serum PSA levels in prostate cancer therapy).

Anti-idiotypic anti-161P2F10B antibodies can also be used in anti-cancer therapy as a vaccine for inducing an immune response to cells expressing a 161P2F10B-related protein. In particular, the generation of anti-idiotypic antibodies is well known in the art; this methodology can readily be adapted to generate anti-idiotypic anti-161P2F10B antibodies that mimic an epitope on a 161P2F10B-related protein (see, for example, Wagner *et al.*, 1997, Hybridoma 16: 33-40; Foon *et al.*, 1995, J. Clin. Invest. 96:334-342; Herlyn *et al.*, 1996, Cancer Immunol. Immunother. 43:65-76). Such an anti-idiotypic antibody can be used in cancer vaccine strategies.

X.C.) 161P2F10B as a Target for Cellular Immune Responses

Vaccines and methods of preparing vaccines that contain an immunogenically effective amount of one or more HLA-binding peptides as described herein are further embodiments of the invention. Furthermore, vaccines in accordance with the invention encompass compositions of one or more of the claimed peptides. A peptide can be present in a vaccine individually. Alternatively, the peptide can exist as a homopolymer comprising multiple copies of the same peptide, or as a heteropolymer of various peptides. Polymers have the advantage of increased immunological reaction and, where different peptide epitopes are used to make up the polymer, the additional ability to induce antibodies and/or CTLs that react with different antigenic determinants of the pathogenic organism or tumor-related peptide targeted for an immune response. The composition can be a naturally occurring region of an antigen or can be prepared, e.g., recombinantly or by chemical synthesis.

Carriers that can be used with vaccines of the invention are well known in the art, and include, e.g., thyroglobulin, albumins such as human serum albumin, tetanus toxoid, polyamino acids such as poly L-lysine, poly L-glutamic acid, influenza, hepatitis B virus core protein, and the like. The vaccines can contain a physiologically tolerable (*i.e.*, acceptable) diluent such as water, or saline, preferably phosphate buffered saline. The vaccines also typically include an adjuvant. Adjuvants such as incomplete Freund's adjuvant, aluminum phosphate, aluminum hydroxide, or alum are examples of materials well known in the art. Additionally, as disclosed herein, CTL responses can be primed by conjugating peptides of the invention to lipids, such as tripalmitoyl-S-glycerylcysteinylserine (P₃CSS). Moreover, an adjuvant such as a synthetic cytosine-phosphorothiolated-guanine-containing (CpG) oligonucleotides has been found to increase CTL responses 10- to 100-fold. (see, e.g. Davila and Celis, J. Immunol. 165:539-547 (2000))

Upon immunization with a peptide composition in accordance with the invention, via injection, aerosol, oral, transdermal, transmucosal, intrapleural, intrathecal, or other suitable routes, the immune system of the host responds to the vaccine by producing large amounts of CTLs and/or HTLs specific for the desired antigen. Consequently, the host becomes at least partially immune to later development of cells that express or overexpress 161P2F10B antigen, or derives at least some therapeutic benefit when the antigen was tumor-associated.

In some embodiments, it may be desirable to combine the class I peptide components with components that induce or facilitate neutralizing antibody and/or helper T cell responses directed to the target antigen. A preferred embodiment of such a composition comprises class I and class II epitopes in accordance with the invention. An alternative embodiment of such a composition comprises a class I and/or class II epitope in accordance with the invention, along with a cross reactive HTL epitope such as PADRE™ (Epimmune, San Diego, CA) molecule (described e.g., in U.S. Patent Number 5,736,142).

A vaccine of the invention can also include antigen-presenting cells (APC), such as dendritic cells (DC), as a vehicle to present peptides of the invention. Vaccine compositions can be created *in vitro*, following dendritic cell mobilization and harvesting, whereby loading of dendritic cells occurs *in vitro*. For example, dendritic cells are transfected, e.g., with a minigene in accordance with the invention, or are pulsed with peptides. The dendritic cell can then be administered to a patient to elicit immune responses *in vivo*. Vaccine compositions, either DNA- or peptide-based, can also be administered *in vivo* in combination with dendritic cell mobilization whereby loading of dendritic cells occurs *in vivo*.

Preferably, the following principles are utilized when selecting an array of epitopes for inclusion in a polyepitopic composition for use in a vaccine, or for selecting discrete epitopes to be included in a vaccine and/or to be encoded by nucleic acids such as a minigene. It is preferred that each of the following principles be balanced in order to make the selection. The multiple epitopes to be incorporated in a given vaccine composition may be, but need not be, contiguous in sequence in the native antigen from which the epitopes are derived.

1.) Epitopes are selected which, upon administration, mimic immune responses that have been observed to be correlated with tumor clearance. For HLA Class I this includes 3-4 epitopes that come from at least one tumor associated antigen (TAA). For HLA Class II a similar rationale is employed; again 3-4 epitopes are selected from at least one TAA (see, e.g., Rosenberg *et al.*, *Science* 278:1447-1450). Epitopes from one TAA may be used in combination with epitopes from one or more additional TAAs to produce a vaccine that targets tumors with varying expression patterns of frequently-expressed TAAs.

2.) Epitopes are selected that have the requisite binding affinity established to be correlated with immunogenicity: for HLA Class I an IC_{50} of 500 nM or less, often 200 nM or less; and for Class II an IC_{50} of 1000 nM or less.

3.) Sufficient supermotif bearing-peptides, or a sufficient array of allele-specific motif-bearing peptides, are selected to give broad population coverage. For example, it is preferable to have at least 80% population coverage. A Monte Carlo analysis, a statistical evaluation known in the art, can be employed to assess the breadth, or redundancy of, population coverage.

4.) When selecting epitopes from cancer-related antigens it is often useful to select analogs because the patient may have developed tolerance to the native epitope.

5.) Of particular relevance are epitopes referred to as "nested epitopes." Nested epitopes occur where at least two epitopes overlap in a given peptide sequence. A nested peptide sequence can comprise B cell, HLA class I and/or HLA class II epitopes. When providing nested epitopes, a general objective is to provide the greatest number of epitopes per sequence. Thus, an aspect is to avoid providing a peptide that is any longer than the amino terminus of the amino terminal epitope and the carboxyl terminus of the carboxyl terminal epitope in the peptide. When providing a multi-epitopic sequence, such as a sequence comprising nested epitopes, it is generally important to screen the sequence in order to insure that it does not have pathological or other deleterious biological properties.

6.) If a polyepitopic protein is created, or when creating a minigene, an objective is to generate the smallest peptide that encompasses the epitopes of interest. This principle is similar, if not the same as that employed when selecting a peptide comprising nested epitopes. However, with an artificial polyepitopic peptide, the size minimization objective is balanced against the need to integrate any spacer sequences between epitopes in the polyepitopic protein. Spacer amino acid residues can, for example, be introduced to avoid junctional epitopes (an epitope recognized by the immune system, not present in the target antigen, and only created by the man-made juxtaposition of epitopes), or to facilitate cleavage between epitopes and thereby enhance epitope presentation. Junctional epitopes are generally to be avoided because the recipient may generate an immune response to that non-native epitope. Of particular concern is a junctional epitope that is a "dominant epitope." A dominant epitope may lead to such a zealous response that immune responses to other epitopes are diminished or suppressed.

7.) Where the sequences of multiple variants of the same target protein are present, potential peptide epitopes can also be selected on the basis of their conservancy. For example, a criterion for conservancy may define that the entire sequence of an HLA class I binding peptide or the entire 9-mer core of a class II binding peptide be conserved in a designated percentage of the sequences evaluated for a specific protein antigen.

X.C.1. Minigene Vaccines

A number of different approaches are available which allow simultaneous delivery of multiple epitopes. Nucleic acids encoding the peptides of the invention are a particularly useful embodiment of the invention. Epitopes for inclusion in a minigene are preferably selected according to the guidelines set forth in the previous section. A preferred means of administering nucleic acids encoding the peptides of the invention uses minigene constructs encoding a peptide comprising one or multiple epitopes of the invention.

The use of multi-epitope minigenes is described below and in, Ishioka *et al.*, *J. Immunol.* 162:3915-3925, 1999; An, L. and Whitton, J. L., *J. Virol.* 71:2292, 1997; Thomson, S. A. *et al.*, *J. Immunol.* 157:822, 1996; Whitton, J. L. *et al.*, *J. Virol.* 67:348, 1993; Hanke, R. *et al.*, *Vaccine* 16:426, 1998. For example, a multi-epitope DNA plasmid encoding supermotif- and/or motif-bearing epitopes derived 161P2F10B, the PADRE® universal helper T cell epitope or multiple HTL epitopes from 161P2F10B (see e.g., Tables VIII-XXI and XXII to XLIX), and an endoplasmic reticulum-translocating signal sequence can be engineered. A vaccine may also comprise epitopes that are derived from other TAAs.

The immunogenicity of a multi-epitopic minigene can be confirmed in transgenic mice to evaluate the magnitude of CTL induction responses against the epitopes tested. Further, the immunogenicity of DNA-encoded epitopes *in vivo* can be correlated with the *in vitro* responses of specific CTL lines against target cells transfected with the DNA plasmid. Thus, these experiments can show that the minigene serves to both: 1.) generate a CTL response and 2.) that the induced CTLs recognized cells expressing the encoded epitopes.

For example, to create a DNA sequence encoding the selected epitopes (minigene) for expression in human cells, the amino acid sequences of the epitopes may be reverse translated. A human codon usage table can be used to guide the codon choice for each amino acid. These epitope-encoding DNA sequences may be directly adjoined, so that when translated, a continuous polypeptide sequence is created. To optimize expression and/or immunogenicity, additional elements can be incorporated into the minigene design. Examples of amino acid sequences that can be reverse translated and included in the minigene sequence include: HLA class I epitopes, HLA class II epitopes, antibody epitopes, a ubiquitination signal sequence, and/or an endoplasmic reticulum targeting signal. In addition, HLA presentation of CTL and HTL epitopes may be improved by including synthetic (e.g. poly-alanine) or naturally-occurring flanking sequences adjacent to the CTL or HTL epitopes; these larger peptides comprising the epitope(s) are within the scope of the invention.

The minigene sequence may be converted to DNA by assembling oligonucleotides that encode the plus and minus strands of the minigene. Overlapping oligonucleotides (30-100 bases long) may be synthesized, phosphorylated, purified and annealed under appropriate conditions using well known techniques. The ends of the oligonucleotides can be joined, for example, using T4 DNA ligase. This synthetic minigene, encoding the epitope polypeptide, can then be cloned into a desired expression vector.

Standard regulatory sequences well known to those of skill in the art are preferably included in the vector to ensure expression in the target cells. Several vector elements are desirable: a promoter with a down-stream cloning site for minigene insertion; a polyadenylation signal for efficient transcription termination; an *E. coli* origin of replication; and an *E. coli* selectable marker (e.g. ampicillin or kanamycin resistance). Numerous promoters can be used for this purpose, e.g., the human cytomegalovirus (hCMV) promoter. See, e.g., U.S. Patent Nos. 5,580,859 and 5,589,466 for other suitable promoter sequences.

Additional vector modifications may be desired to optimize minigene expression and immunogenicity. In some cases, introns are required for efficient gene expression, and one or more synthetic or naturally-occurring introns could be incorporated into the transcribed region of the minigene. The inclusion of mRNA stabilization sequences and sequences for replication in mammalian cells may also be considered for increasing minigene expression.

Once an expression vector is selected, the minigene is cloned into the polylinker region downstream of the promoter. This plasmid is transformed into an appropriate *E. coli* strain, and DNA is prepared using standard techniques. The orientation and DNA sequence of the minigene, as well as all other elements included in the vector, are confirmed using restriction mapping and DNA sequence analysis. Bacterial cells harboring the correct plasmid can be stored as a master cell bank and a working cell bank.

In addition, immunostimulatory sequences (ISSs or CpGs) appear to play a role in the immunogenicity of DNA vaccines. These sequences may be included in the vector, outside the minigene coding sequence, if desired to enhance immunogenicity.

In some embodiments, a bi-cistronic expression vector which allows production of both the minigene-encoded epitopes and a second protein (included to enhance or decrease immunogenicity) can be used. Examples of proteins or polypeptides that could beneficially enhance the immune response if co-expressed include cytokines (e.g., IL-2, IL-12, GM-CSF), cytokine-inducing molecules (e.g., I α IF), costimulatory molecules, or for HTL responses, pan-DR binding proteins (PADRE™, Epimmune, San Diego, CA). Helper (HTL) epitopes can be joined to intracellular targeting signals and expressed separately from expressed CTL epitopes; this allows direction of the HTL epitopes to a cell compartment different than that of the CTL epitopes. If required, this could facilitate more efficient entry of HTL epitopes into the HLA class II pathway, thereby improving HTL induction. In contrast to HTL or CTL induction, specifically decreasing the immune response by co-expression of immunosuppressive molecules (e.g. TGF- β) may be beneficial in certain diseases.

Therapeutic quantities of plasmid DNA can be produced for example, by fermentation in *E. coli*, followed by purification. Aliquots from the working cell bank are used to inoculate growth medium, and grown to saturation in shaker flasks or a bioreactor according to well-known techniques. Plasmid DNA can be purified using standard bioseparation technologies such as solid phase anion-exchange resins supplied by QIAGEN, Inc. (Valencia, California). If required, supercoiled DNA can be isolated from the open circular and linear forms using gel electrophoresis or other methods.

Purified plasmid DNA can be prepared for injection using a variety of formulations. The simplest of these is reconstitution of lyophilized DNA in sterile phosphate-buffer saline (PBS). This approach, known as "naked DNA," is currently being used for intramuscular (IM) administration in clinical trials. To maximize the immunotherapeutic effects of minigene DNA vaccines, an alternative method for formulating purified plasmid DNA may be desirable. A variety of methods have been described, and new techniques may become available. Cationic lipids, glycolipids, and fusogenic liposomes can also be used in the formulation (see, e.g., as described by WO 93/24640; Mannino & Gould-Fogerite, *BioTechniques* 6(7): 682 (1988); U.S. Pat No. 5,279,833; WO 91/06309; and Felgner, *et al.*, *Proc. Nat'l Acad. Sci. USA* 84:7413 (1987). In addition, peptides and compounds referred to collectively as protective, interactive, non-condensing compounds (PINC) could also be complexed to purified plasmid DNA to influence variables such as stability, intramuscular dispersion, or trafficking to specific organs or cell types.

Target cell sensitization can be used as a functional assay for expression and HLA class I presentation of minigene-encoded CTL epitopes. For example, the plasmid DNA is introduced into a mammalian cell line that is suitable as a target for standard CTL chromium release assays. The transfection method used will be dependent on the final formulation. Electroporation can be used for "naked" DNA, whereas cationic lipids allow direct *in vitro* transfection. A plasmid expressing green fluorescent protein (GFP) can be co-transfected to allow enrichment of transfected cells using fluorescence activated cell sorting (FACS). These cells are then chromium-51 (⁵¹Cr) labeled and used as target cells for

epitope-specific CTL lines; cytolysis, detected by ^{51}Cr release, indicates both production of, and HLA presentation of, minigene-encoded CTL epitopes. Expression of HTL epitopes may be evaluated in an analogous manner using assays to assess HTL activity.

In vivo immunogenicity is a second approach for functional testing of minigene DNA formulations. Transgenic mice expressing appropriate human HLA proteins are immunized with the DNA product. The dose and route of administration are formulation dependent (e.g., IM for DNA in PBS, intraperitoneal (i.p.) for lipid-complexed DNA). Twenty-one days after immunization, splenocytes are harvested and restimulated for one week in the presence of peptides encoding each epitope being tested. Thereafter, for CTL effector cells, assays are conducted for cytolysis of peptide-loaded, ^{51}Cr -labeled target cells using standard techniques. Lysis of target cells that were sensitized by HLA loaded with peptide epitopes, corresponding to minigene-encoded epitopes, demonstrates DNA vaccine function for *in vivo* induction of CTLs. Immunogenicity of HTL epitopes is confirmed in transgenic mice in an analogous manner.

Alternatively, the nucleic acids can be administered using ballistic delivery as described, for instance, in U.S. Patent No. 5,204,253. Using this technique, particles comprised solely of DNA are administered. In a further alternative embodiment, DNA can be adhered to particles, such as gold particles.

Minigenes can also be delivered using other bacterial or viral delivery systems well known in the art, e.g., an expression construct encoding epitopes of the invention can be incorporated into a viral vector such as vaccinia.

X.C.2. Combinations of CTL Peptides with Helper Peptides

Vaccine compositions comprising CTL peptides of the invention can be modified, e.g., analoged, to provide desired attributes, such as improved serum half life, broadened population coverage or enhanced immunogenicity.

For instance, the ability of a peptide to induce CTL activity can be enhanced by linking the peptide to a sequence which contains at least one epitope that is capable of inducing a T helper cell response. Although a CTL peptide can be directly linked to a T helper peptide, often CTL epitope/HTL epitope conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under physiological conditions. The spacers are typically selected from, e.g., Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two residues, more usually three to six residues and sometimes 10 or more residues. The CTL peptide epitope can be linked to the T helper peptide epitope either directly or via a spacer either at the amino or carboxy terminus of the CTL peptide. The amino terminus of either the immunogenic peptide or the T helper peptide may be acylated.

In certain embodiments, the T helper peptide is one that is recognized by T helper cells present in a majority of a genetically diverse population. This can be accomplished by selecting peptides that bind to many, most, or all of the HLA class II molecules. Examples of such amino acid bind many HLA Class II molecules include sequences from antigens such as *tetanus toxoid* at positions 830-843 (QYIKANSKFIGITE; SEQ ID NO: 25), *Plasmodium falciparum* circumsporozoite (CS) protein at positions 378-398 (DIEKKIAKMEKASSVFNVVNS; SEQ ID NO: 26), and *Streptococcus* 18kD protein at positions 116-131 (GAVDSILGGVATYGAA; SEQ ID NO: 27). Other examples include peptides bearing a DR 1-4-7 supermotif, or either of the DR3 motifs.

Alternatively, it is possible to prepare synthetic peptides capable of stimulating T helper lymphocytes, in a loosely HLA-restricted fashion, using amino acid sequences not found in nature (see, e.g., PCT publication WO 95/07707). These synthetic compounds called Pan-DR-binding epitopes (e.g., PADRE™, Epimmune, Inc., San Diego, CA) are designed, most preferably, to bind most HLA-DR (human HLA class II) molecules. For instance, a pan-DR-binding epitope peptide having the formula: aKXVAAWTLKAAa (SEQ ID NO: 28), where "X" is either cyclohexylalanine, phenylalanine, or tyrosine, and a is

either D-alanine or L-alanine, has been found to bind to most HLA-DR alleles, and to stimulate the response of T helper lymphocytes from most individuals, regardless of their HLA type. An alternative of a pan-DR binding epitope comprises all "L" natural amino acids and can be provided in the form of nucleic acids that encode the epitope.

HTL peptide epitopes can also be modified to alter their biological properties. For example, they can be modified to include D-amino acids to increase their resistance to proteases and thus extend their serum half life, or they can be conjugated to other molecules such as lipids, proteins, carbohydrates, and the like to increase their biological activity. For example, a T helper peptide can be conjugated to one or more palmitic acid chains at either the amino or carboxyl termini.

X.C.3. Combinations of CTL Peptides with T Cell Priming Agents

In some embodiments it may be desirable to include in the pharmaceutical compositions of the invention at least one component which primes B lymphocytes or T lymphocytes. Lipids have been identified as agents capable of priming CTL *in vivo*. For example, palmitic acid residues can be attached to the ϵ - and α - amino groups of a lysine residue and then linked, e.g., via one or more linking residues such as Gly, Gly-Gly-, Ser, Ser-Ser, or the like, to an immunogenic peptide. The lipidated peptide can then be administered either directly in a micelle or particle, incorporated into a liposome, or emulsified in an adjuvant, e.g., incomplete Freund's adjuvant. In a preferred embodiment, a particularly effective immunogenic composition comprises palmitic acid attached to ϵ - and α - amino groups of Lys, which is attached via linkage, e.g., Ser-Ser, to the amino terminus of the immunogenic peptide.

As another example of lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P₃CSS) can be used to prime virus specific CTL when covalently attached to an appropriate peptide (see, e.g., Deres, *et al.*, *Nature* 342:561, 1989). Peptides of the invention can be coupled to P₃CSS, for example, and the lipopeptide administered to an individual to prime specifically an immune response to the target antigen. Moreover, because the induction of neutralizing antibodies can also be primed with P₃CSS-conjugated epitopes, two such compositions can be combined to more effectively elicit both humoral and cell-mediated responses.

X.C.4. Vaccine Compositions Comprising DC Pulsed with CTL and/or HTL Peptides

An embodiment of a vaccine composition in accordance with the invention comprises *ex vivo* administration of a cocktail of epitope-bearing peptides to PBMC, or isolated DC therefrom, from the patient's blood. A pharmaceutical to facilitate harvesting of DC can be used, such as Progenipoietin™ (Pharmacia-Monsanto, St. Louis, MO) or GM-CSF/IL-4. After pulsing the DC with peptides and prior to reinfusion into patients, the DC are washed to remove unbound peptides. In this embodiment, a vaccine comprises peptide-pulsed DCs which present the pulsed peptide epitopes complexed with HLA molecules on their surfaces.

The DC can be pulsed *ex vivo* with a cocktail of peptides, some of which stimulate CTL responses to 161P2F10B. Optionally, a helper T cell (HTL) peptide, such as a natural or artificial loosely restricted HLA Class II peptide, can be included to facilitate the CTL response. Thus, a vaccine in accordance with the invention is used to treat a cancer which expresses or overexpresses 161P2F10B.

X.D. Adoptive Immunotherapy

Antigenic 161P2F10B-related peptides are used to elicit a CTL and/or HTL response *ex vivo*, as well. The resulting CTL or HTL cells, can be used to treat tumors in patients that do not respond to other conventional forms of therapy, or will not respond to a therapeutic vaccine peptide or nucleic acid in accordance with the invention. *Ex vivo* CTL or HTL responses to a particular antigen are induced by incubating in tissue culture the patient's, or genetically compatible, CTL or HTL precursor cells together with a source of antigen-presenting cells (APC), such as dendritic cells, and the appropriate immunogenic peptide. After an appropriate incubation time (typically about 7-28 days), in which the precursor cells are activated and expanded into effector cells, the cells are infused back into the patient, where they will destroy (CTL) or

facilitate destruction (HTL) of their specific target cell (e.g., a tumor cell). Transfected dendritic cells may also be used as antigen presenting cells.

X.E. Administration of Vaccines for Therapeutic or Prophylactic Purposes

Pharmaceutical and vaccine compositions of the invention are typically used to treat and/or prevent a cancer that expresses or overexpresses 161P2F10B. In therapeutic applications, peptide and/or nucleic acid compositions are administered to a patient in an amount sufficient to elicit an effective B cell, CTL and/or HTL response to the antigen and to cure or at least partially arrest or slow symptoms and/or complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, e.g., the particular composition administered, the manner of administration, the stage and severity of the disease being treated, the weight and general state of health of the patient, and the judgment of the prescribing physician.

For pharmaceutical compositions, the immunogenic peptides of the invention, or DNA encoding them, are generally administered to an individual already bearing a tumor that expresses 161P2F10B. The peptides or DNA encoding them can be administered individually or as fusions of one or more peptide sequences. Patients can be treated with the immunogenic peptides separately or in conjunction with other treatments, such as surgery, as appropriate.

For therapeutic use, administration should generally begin at the first diagnosis of 161P2F10B-associated cancer. This is followed by boosting doses until at least symptoms are substantially abated and for a period thereafter. The embodiment of the vaccine composition (*i.e.*, including, but not limited to embodiments such as peptide cocktails, polyepitopic polypeptides, minigenes, or TAA-specific CTLs or pulsed dendritic cells) delivered to the patient may vary according to the stage of the disease or the patient's health status. For example, in a patient with a tumor that expresses 161P2F10B, a vaccine comprising 161P2F10B-specific CTL may be more efficacious in killing tumor cells in patient with advanced disease than alternative embodiments.

It is generally important to provide an amount of the peptide epitope delivered by a mode of administration sufficient to stimulate effectively a cytotoxic T cell response; compositions which stimulate helper T cell responses can also be given in accordance with this embodiment of the invention.

The dosage for an initial therapeutic immunization generally occurs in a unit dosage range where the lower value is about 1, 5, 50, 500, or 1,000 μg and the higher value is about 10,000; 20,000; 30,000; or 50,000 μg . Dosage values for a human typically range from about 500 μg to about 50,000 μg per 70 kilogram patient. Boosting dosages of between about 1.0 μg to about 50,000 μg of peptide pursuant to a boosting regimen over weeks to months may be administered depending upon the patient's response and condition as determined by measuring the specific activity of CTL and HTL obtained from the patient's blood. Administration should continue until at least clinical symptoms or laboratory tests indicate that the neoplasia, has been eliminated or reduced and for a period thereafter. The dosages, routes of administration, and dose schedules are adjusted in accordance with methodologies known in the art.

In certain embodiments, the peptides and compositions of the present invention are employed in serious disease states, that is, life-threatening or potentially life threatening situations. In such cases, as a result of the minimal amounts of extraneous substances and the relative nontoxic nature of the peptides in preferred compositions of the invention, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these peptide compositions relative to these stated dosage amounts.

The vaccine compositions of the invention can also be used purely as prophylactic agents. Generally the dosage for an initial prophylactic immunization generally occurs in a unit dosage range where the lower value is about 1, 5, 50, 500, or 1000 μg and the higher value is about 10,000; 20,000; 30,000; or 50,000 μg . Dosage values for a human typically range from about 500 μg to about 50,000 μg per 70 kilogram patient. This is followed by boosting dosages of between about 1.0

μg to about 50,000 μg of peptide administered at defined intervals from about four weeks to six months after the initial administration of vaccine. The immunogenicity of the vaccine can be assessed by measuring the specific activity of CTL and HTL obtained from a sample of the patient's blood.

The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral, nasal, intrathecal, or local (e.g. as a cream or topical ointment) administration. Preferably, the pharmaceutical compositions are administered parentally, e.g., intravenously, subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration which comprise a solution of the immunogenic peptides dissolved or suspended in an acceptable carrier, preferably an aqueous carrier.

A variety of aqueous carriers may be used, e.g., water, buffered water, 0.8% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well-known sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration.

The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH-adjusting and buffering agents, tonicity adjusting agents, wetting agents, preservatives, and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

The concentration of peptides of the invention in the pharmaceutical formulations can vary widely, *i.e.*, from less than about 0.1%, usually at or at least about 2% to as much as 20% to 50% or more by weight, and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected.

A human unit dose form of a composition is typically included in a pharmaceutical composition that comprises a human unit dose of an acceptable carrier, in one embodiment an aqueous carrier, and is administered in a volume/quantity that is known by those of skill in the art to be used for administration of such compositions to humans (see, e.g., Remington's Pharmaceutical Sciences, 17th Edition, A. Gennaro, Editor, Mack Publishing Co., Easton, Pennsylvania, 1985). For example a peptide dose for initial immunization can be from about 1 to about 50,000 μg , generally 100-5,000 μg , for a 70 kg patient. For example, for nucleic acids an initial immunization may be performed using an expression vector in the form of naked nucleic acid administered IM (or SC or ID) in the amounts of 0.5-5 mg at multiple sites. The nucleic acid (0.1 to 1000 μg) can also be administered using a gene gun. Following an incubation period of 3-4 weeks, a booster dose is then administered. The booster can be recombinant fowlpox virus administered at a dose of $5 \cdot 10^7$ to $5 \cdot 10^9$ pfu.

For antibodies, a treatment generally involves repeated administration of the anti-161P2F10B antibody preparation, via an acceptable route of administration such as intravenous injection (IV), typically at a dose in the range of about 0.1 to about 10 mg/kg body weight. In general, doses in the range of 10-500 mg mAb per week are effective and well tolerated. Moreover, an initial loading dose of approximately 4 mg/kg patient body weight IV, followed by weekly doses of about 2 mg/kg IV of the anti- 161P2F10B mAb preparation represents an acceptable dosing regimen. As appreciated by those of skill in the art, various factors can influence the ideal dose in a particular case. Such factors include, for example, half life of a composition, the binding affinity of an Ab, the immunogenicity of a substance, the degree of 161P2F10B expression in the patient, the extent of circulating shed 161P2F10B antigen, the desired steady-state concentration level, frequency of treatment, and the influence of chemotherapeutic or other agents used in combination with the treatment method of the invention, as well as the health status of a particular patient. Non-limiting preferred human unit doses are, for example, 500 μg - 1mg, 1mg - 50mg, 50mg - 100mg, 100mg - 200mg, 200mg - 300mg, 400mg - 500mg, 500mg - 600mg, 600mg - 700mg, 700mg - 800mg, 800mg - 900mg, 900mg - 1g, or 1mg - 700mg. In certain embodiments, the dose is in a range of 2-5 mg/kg body weight, e.g., with follow on weekly doses of 1-3 mg/kg; 0.5mg, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10mg/kg body weight followed, e.g., in two, three or four weeks by weekly doses; 0.5 - 10mg/kg body weight, e.g., followed in two, three or four

weeks by weekly doses; 225, 250, 275, 300, 325, 350, 375, 400mg m² of body area weekly; 1-600mg m² of body area weekly; 225-400mg m² of body area weekly; these does can be followed by weekly doses for 2, 3, 4, 5, 6, 7, 8, 9, 19, 11, 12 or more weeks.

In one embodiment, human unit dose forms of polynucleotides comprise a suitable dosage range or effective amount that provides any therapeutic effect. As appreciated by one of ordinary skill in the art a therapeutic effect depends on a number of factors, including the sequence of the polynucleotide, molecular weight of the polynucleotide and route of administration. Dosages are generally selected by the physician or other health care professional in accordance with a variety of parameters known in the art, such as severity of symptoms, history of the patient and the like. Generally, for a polynucleotide of about 20 bases, a dosage range may be selected from, for example, an independently selected lower limit such as about 0.1, 0.25, 0.5, 1, 2, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400 or 500 mg/kg up to an independently selected upper limit, greater than the lower limit, of about 60, 80, 100, 200, 300, 400, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000 or 10,000 mg/kg. For example, a dose may be about any of the following: 0.1 to 100 mg/kg, 0.1 to 50 mg/kg, 0.1 to 25 mg/kg, 0.1 to 10 mg/kg, 1 to 500 mg/kg, 100 to 400 mg/kg, 200 to 300 mg/kg, 1 to 100 mg/kg, 100 to 200 mg/kg, 300 to 400 mg/kg, 400 to 500 mg/kg, 500 to 1000 mg/kg, 500 to 5000 mg/kg, or 500 to 10,000 mg/kg. Generally, parenteral routes of administration may require higher doses of polynucleotide compared to more direct application to the nucleotide to diseased tissue, as do polynucleotides of increasing length.

In one embodiment, human unit dose forms of T-cells comprise a suitable dosage range or effective amount that provides any therapeutic effect. As appreciated by one of ordinary skill in the art, a therapeutic effect depends on a number of factors. Dosages are generally selected by the physician or other health care professional in accordance with a variety of parameters known in the art, such as severity of symptoms, history of the patient and the like. A dose may be about 10⁴ cells to about 10⁶ cells, about 10⁶ cells to about 10⁸ cells, about 10⁸ to about 10¹¹ cells, or about 10⁸ to about 5 x 10¹⁰ cells. A dose may also about 10⁶ cells/m² to about 10¹⁰ cells/m², or about 10⁶ cells/m² to about 10⁸ cells/m².

Proteins(s) of the invention, and/or nucleic acids encoding the protein(s), can also be administered via liposomes, which may also serve to: 1) target the proteins(s) to a particular tissue, such as lymphoid tissue; 2) to target selectively to diseases cells; or, 3) to increase the half-life of the peptide composition. Liposomes include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. In these preparations, the peptide to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to a receptor prevalent among lymphoid cells, such as monoclonal antibodies which bind to the CD45 antigen, or with other therapeutic or immunogenic compositions. Thus, liposomes either filled or decorated with a desired peptide of the invention can be directed to the site of lymphoid cells, where the liposomes then deliver the peptide compositions. Liposomes for use in accordance with the invention are formed from standard vesicle-forming lipids, which generally include neutral and negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, e.g., liposome size, acid lability and stability of the liposomes in the blood stream. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka, *et al.*, *Ann. Rev. Biophys. Bioeng.* 9:467 (1980), and U.S. Patent Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369.

For targeting cells of the immune system, a ligand to be incorporated into the liposome can include, e.g., antibodies or fragments thereof specific for cell surface determinants of the desired immune system cells. A liposome suspension containing a peptide may be administered intravenously, locally, topically, etc. in a dose which varies according to, *inter alia*, the manner of administration, the peptide being delivered, and the stage of the disease being treated.

For solid compositions, conventional nontoxic solid carriers may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is

formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95% of active ingredient, that is, one or more peptides of the invention, and more preferably at a concentration of 25%-75%.

For aerosol administration, immunogenic peptides are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of peptides are about 0.01%-20% by weight, preferably about 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from about 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute about 0.1%-20% by weight of the composition, preferably about 0.25-5%. The balance of the composition is ordinarily propellant. A carrier can also be included, as desired, as with, e.g., lecithin for intranasal delivery.

XI.) Diagnostic and Prognostic Embodiments of 161P2F10B.

As disclosed herein, 161P2F10B polynucleotides, polypeptides, reactive cytotoxic T cells (CTL), reactive helper T cells (HTL) and anti-polypeptide antibodies are used in well known diagnostic, prognostic and therapeutic assays that examine conditions associated with dysregulated cell growth such as cancer, in particular the cancers listed in Table I (see, e.g., both its specific pattern of tissue expression as well as its overexpression in certain cancers as described for example in the Example entitled "Expression analysis of 161P2F10B in normal tissues, and patient specimens").

161P2F10B can be analogized to a prostate associated antigen PSA, the archetypal marker that has been used by medical practitioners for years to identify and monitor the presence of prostate cancer (see, e.g., Merrill *et al.*, J. Urol. 163(2): 503-5120 (2000); Polascik *et al.*, J. Urol. Aug; 162(2):293-306 (1999) and Fortier *et al.*, J. Nat. Cancer Inst. 91(19): 1635-1640(1999)). A variety of other diagnostic markers are also used in similar contexts including p53 and K-ras (see, e.g., Tulchinsky *et al.*, Int J Mol Med 1999 Jul 4(1):99-102 and Minimoto *et al.*, Cancer Detect Prev 2000;24(1):1-12). Therefore, this disclosure of 161P2F10B polynucleotides and polypeptides (as well as 161P2F10B polynucleotide probes and anti-161P2F10B antibodies used to identify the presence of these molecules) and their properties allows skilled artisans to utilize these molecules in methods that are analogous to those used, for example, in a variety of diagnostic assays directed to examining conditions associated with cancer.

Typical embodiments of diagnostic methods which utilize the 161P2F10B polynucleotides, polypeptides, reactive T cells and antibodies are analogous to those methods from well-established diagnostic assays, which employ, e.g., PSA polynucleotides, polypeptides, reactive T cells and antibodies. For example, just as PSA polynucleotides are used as probes (for example in Northern analysis, see, e.g., Sharief *et al.*, Biochem. Mol. Biol. Int. 33(3):567-74(1994)) and primers (for example in PCR analysis, see, e.g., Okegawa *et al.*, J. Urol. 163(4): 1189-1190 (2000)) to observe the presence and/or the level of PSA mRNAs in methods of monitoring PSA overexpression or the metastasis of prostate cancers, the 161P2F10B polynucleotides described herein can be utilized in the same way to detect 161P2F10B overexpression or the metastasis of prostate and other cancers expressing this gene. Alternatively, just as PSA polypeptides are used to generate antibodies specific for PSA which can then be used to observe the presence and/or the level of PSA proteins in methods to monitor PSA protein overexpression (see, e.g., Stephan *et al.*, Urology 55(4):560-3 (2000)) or the metastasis of prostate cells (see, e.g., Alanen *et al.*, Pathol. Res. Pract. 192(3):233-7 (1996)), the 161P2F10B polypeptides described herein can be utilized to generate antibodies for use in detecting 161P2F10B overexpression or the metastasis of prostate cells and cells of other cancers expressing this gene.

Specifically, because metastases involves the movement of cancer cells from an organ of origin (such as the lung or prostate gland etc.) to a different area of the body (such as a lymph node), assays which examine a biological sample for the presence of cells expressing 161P2F10B polynucleotides and/or polypeptides can be used to provide evidence of

metastasis. For example, when a biological sample from tissue that does not normally contain 161P2F10B-expressing cells (lymph node) is found to contain 161P2F10B-expressing cells such as the 161P2F10B expression seen in LAPC4 and LAPC9, xenografts isolated from lymph node and bone metastasis, respectively, this finding is indicative of metastasis.

Alternatively 161P2F10B polynucleotides and/or polypeptides can be used to provide evidence of cancer, for example, when cells in a biological sample that do not normally express 161P2F10B or express 161P2F10B at a different level are found to express 161P2F10B or have an increased expression of 161P2F10B (see, e.g., the 161P2F10B expression in the cancers listed in Table I and in patient samples etc. shown in the accompanying Figures). In such assays, artisans may further wish to generate supplementary evidence of metastasis by testing the biological sample for the presence of a second tissue restricted marker (in addition to 161P2F10B) such as PSA, PSCA etc. (see, e.g., Alanen *et al.*, *Pathol. Res. Pract.* 192(3): 233-237 (1996)).

Just as PSA polynucleotide fragments and polynucleotide variants are employed by skilled artisans for use in methods of monitoring PSA, 161P2F10B polynucleotide fragments and polynucleotide variants are used in an analogous manner. In particular, typical PSA polynucleotides used in methods of monitoring PSA are probes or primers which consist of fragments of the PSA cDNA sequence. Illustrating this, primers used to PCR amplify a PSA polynucleotide must include less than the whole PSA sequence to function in the polymerase chain reaction. In the context of such PCR reactions, skilled artisans generally create a variety of different polynucleotide fragments that can be used as primers in order to amplify different portions of a polynucleotide of interest or to optimize amplification reactions (see, e.g., Caetano-Anolles, G. *Biotechniques* 25(3): 472-476, 478-480 (1998); Robertson *et al.*, *Methods Mol. Biol.* 98:121-154 (1998)). An additional illustration of the use of such fragments is provided in the Example entitled "Expression analysis of 161P2F10B in normal tissues, and patient specimens," where a 161P2F10B polynucleotide fragment is used as a probe to show the expression of 161P2F10B RNAs in cancer cells. In addition, variant polynucleotide sequences are typically used as primers and probes for the corresponding mRNAs in PCR and Northern analyses (see, e.g., Sawai *et al.*, *Fetal Diagn. Ther.* 1996 Nov-Dec 11(6):407-13 and *Current Protocols In Molecular Biology*, Volume 2, Unit 2, Frederick M. Ausubel *et al.* eds., 1995)). Polynucleotide fragments and variants are useful in this context where they are capable of binding to a target polynucleotide sequence (e.g., a 161P2F10B polynucleotide shown in Figure 2 or variant thereof) under conditions of high stringency.

Furthermore, PSA polypeptides which contain an epitope that can be recognized by an antibody or T cell that specifically binds to that epitope are used in methods of monitoring PSA. 161P2F10B polypeptide fragments and polypeptide analogs or variants can also be used in an analogous manner. This practice of using polypeptide fragments or polypeptide variants to generate antibodies (such as anti-PSA antibodies or T cells) is typical in the art with a wide variety of systems such as fusion proteins being used by practitioners (see, e.g., *Current Protocols In Molecular Biology*, Volume 2, Unit 16, Frederick M. Ausubel *et al.* eds., 1995). In this context, each epitope(s) functions to provide the architecture with which an antibody or T cell is reactive. Typically, skilled artisans create a variety of different polypeptide fragments that can be used in order to generate immune responses specific for different portions of a polypeptide of interest (see, e.g., U.S. Patent No. 5,840,501 and U.S. Patent No. 5,939,533). For example it may be preferable to utilize a polypeptide comprising one of the 161P2F10B biological motifs discussed herein or a motif-bearing subsequence which is readily identified by one of skill in the art based on motifs available in the art. Polypeptide fragments, variants or analogs are typically useful in this context as long as they comprise an epitope capable of generating an antibody or T cell specific for a target polypeptide sequence (e.g. a 161P2F10B polypeptide shown in Figure 3).

As shown herein, the 161P2F10B polynucleotides and polypeptides (as well as the 161P2F10B polynucleotide probes and anti-161P2F10B antibodies or T cells used to identify the presence of these molecules) exhibit specific properties that make them useful in diagnosing cancers such as those listed in Table I. Diagnostic assays that measure the presence of 161P2F10B gene products, in order to evaluate the presence or onset of a disease condition described herein, such as

prostate cancer, are used to identify patients for preventive measures or further monitoring, as has been done so successfully with PSA. Moreover, these materials satisfy a need in the art for molecules having similar or complementary characteristics to PSA in situations where, for example, a definite diagnosis of metastasis of prostatic origin cannot be made on the basis of a test for PSA alone (see, e.g., Alanen *et al.*, Pathol. Res. Pract. 192(3): 233-237 (1996)), and consequently, materials such as 161P2F10B polynucleotides and polypeptides (as well as the 161P2F10B polynucleotide probes and anti-161P2F10B antibodies used to identify the presence of these molecules) need to be employed to confirm a metastases of prostatic origin.

Finally, in addition to their use in diagnostic assays, the 161P2F10B polynucleotides disclosed herein have a number of other utilities such as their use in the identification of oncogenetic associated chromosomal abnormalities in the chromosomal region to which the 161P2F10B gene maps (see the Example entitled "Chromosomal Mapping of 161P2F10B" below). Moreover, in addition to their use in diagnostic assays, the 161P2F10B-related proteins and polynucleotides disclosed herein have other utilities such as their use in the forensic analysis of tissues of unknown origin (see, e.g., Takahama K Forensic Sci Int 1996 Jun 28;80(1-2): 63-9).

Additionally, 161P2F10B-related proteins or polynucleotides of the invention can be used to treat a pathologic condition characterized by the over-expression of 161P2F10B. For example, the amino acid or nucleic acid sequence of Figure 2 or Figure 3, or fragments of either, can be used to generate an immune response to a 161P2F10B antigen. Antibodies or other molecules that react with 161P2F10B can be used to modulate the function of this molecule, and thereby provide a therapeutic benefit.

XII.) Inhibition of 161P2F10B Protein Function

The invention includes various methods and compositions for inhibiting the binding of 161P2F10B to its binding partner or its association with other protein(s) as well as methods for inhibiting 161P2F10B function.

XII.A.) Inhibition of 161P2F10B With Intracellular Antibodies

In one approach, a recombinant vector that encodes single chain antibodies that specifically bind to 161P2F10B are introduced into 161P2F10B expressing cells via gene transfer technologies. Accordingly, the encoded single chain anti-161P2F10B antibody is expressed intracellularly, binds to 161P2F10B protein, and thereby inhibits its function. Methods for engineering such intracellular single chain antibodies are well known. Such intracellular antibodies, also known as "intrabodies", are specifically targeted to a particular compartment within the cell, providing control over where the inhibitory activity of the treatment is focused. This technology has been successfully applied in the art (for review, see Richardson and Marasco, 1995, TIBTECH vol. 13). Intrabodies have been shown to virtually eliminate the expression of otherwise abundant cell surface receptors (see, e.g., Richardson *et al.*, 1995, Proc. Natl. Acad. Sci. USA 92: 3137-3141; Beerli *et al.*, 1994, J. Biol. Chem. 269: 23931-23936; Deshane *et al.*, 1994, Gene Ther. 1: 332-337).

Single chain antibodies comprise the variable domains of the heavy and light chain joined by a flexible linker polypeptide, and are expressed as a single polypeptide. Optionally, single chain antibodies are expressed as a single chain variable region fragment joined to the light chain constant region. Well-known intracellular trafficking signals are engineered into recombinant polynucleotide vectors encoding such single chain antibodies in order to target precisely the intrabody to the desired intracellular compartment. For example, intrabodies targeted to the endoplasmic reticulum (ER) are engineered to incorporate a leader peptide and, optionally, a C-terminal ER retention signal, such as the KDEL amino acid motif. Intrabodies intended to exert activity in the nucleus are engineered to include a nuclear localization signal. Lipid moieties are joined to intrabodies in order to tether the intrabody to the cytosolic side of the plasma membrane. Intrabodies can also be

targeted to exert function in the cytosol. For example, cytosolic intrabodies are used to sequester factors within the cytosol, thereby preventing them from being transported to their natural cellular destination.

In one embodiment, intrabodies are used to capture 161P2F10B in the nucleus, thereby preventing its activity within the nucleus. Nuclear targeting signals are engineered into such 161P2F10B intrabodies in order to achieve the desired targeting. Such 161P2F10B intrabodies are designed to bind specifically to a particular 161P2F10B domain. In another embodiment, cytosolic intrabodies that specifically bind to a 161P2F10B protein are used to prevent 161P2F10B from gaining access to the nucleus, thereby preventing it from exerting any biological activity within the nucleus (e.g., preventing 161P2F10B from forming transcription complexes with other factors).

In order to specifically direct the expression of such intrabodies to particular cells, the transcription of the intrabody is placed under the regulatory control of an appropriate tumor-specific promoter and/or enhancer. In order to target intrabody expression specifically to prostate, for example, the PSA promoter and/or promoter/enhancer can be utilized (See, for example, U.S. Patent No. 5,919,652 issued 6 July 1999).

XII.B.) Inhibition of 161P2F10B with Recombinant Proteins

In another approach, recombinant molecules bind to 161P2F10B and thereby inhibit 161P2F10B function. For example, these recombinant molecules prevent or inhibit 161P2F10B from accessing/binding to its binding partner(s) or associating with other protein(s). Such recombinant molecules can, for example, contain the reactive part(s) of a 161P2F10B specific antibody molecule. In a particular embodiment, the 161P2F10B binding domain of a 161P2F10B binding partner is engineered into a dimeric fusion protein, whereby the fusion protein comprises two 161P2F10B ligand binding domains linked to the Fc portion of a human IgG, such as human IgG1. Such IgG portion can contain, for example, the C_H2 and C_H3 domains and the hinge region, but not the C_H1 domain. Such dimeric fusion proteins are administered in soluble form to patients suffering from a cancer associated with the expression of 161P2F10B, whereby the dimeric fusion protein specifically binds to 161P2F10B and blocks 161P2F10B interaction with a binding partner. Such dimeric fusion proteins are further combined into multimeric proteins using known antibody linking technologies.

XII.C.) Inhibition of 161P2F10B Transcription or Translation

The present invention also comprises various methods and compositions for inhibiting the transcription of the 161P2F10B gene. Similarly, the invention also provides methods and compositions for inhibiting the translation of 161P2F10B mRNA into protein.

In one approach, a method of inhibiting the transcription of the 161P2F10B gene comprises contacting the 161P2F10B gene with a 161P2F10B antisense polynucleotide. In another approach, a method of inhibiting 161P2F10B mRNA translation comprises contacting a 161P2F10B mRNA with an antisense polynucleotide. In another approach, a 161P2F10B specific ribozyme is used to cleave a 161P2F10B message, thereby inhibiting translation. Such antisense and ribozyme based methods can also be directed to the regulatory regions of the 161P2F10B gene, such as 161P2F10B promoter and/or enhancer elements. Similarly, proteins capable of inhibiting a 161P2F10B gene transcription factor are used to inhibit 161P2F10B mRNA transcription. The various polynucleotides and compositions useful in the aforementioned methods have been described above. The use of antisense and ribozyme molecules to inhibit transcription and translation is well known in the art.

Other factors that inhibit the transcription of 161P2F10B by interfering with 161P2F10B transcriptional activation are also useful to treat cancers expressing 161P2F10B. Similarly, factors that interfere with 161P2F10B processing are useful to treat cancers that express 161P2F10B. Cancer treatment methods utilizing such factors are also within the scope of the invention.

XII.D.) General Considerations for Therapeutic Strategies

Gene transfer and gene therapy technologies can be used to deliver therapeutic polynucleotide molecules to tumor cells synthesizing 161P2F10B (i.e., antisense, ribozyme, polynucleotides encoding intrabodies and other 161P2F10B inhibitory molecules). A number of gene therapy approaches are known in the art. Recombinant vectors encoding 161P2F10B antisense polynucleotides, ribozymes, factors capable of interfering with 161P2F10B transcription, and so forth, can be delivered to target tumor cells using such gene therapy approaches.

The above therapeutic approaches can be combined with any one of a wide variety of surgical, chemotherapy or radiation therapy regimens. The therapeutic approaches of the invention can enable the use of reduced dosages of chemotherapy (or other therapies) and/or less frequent administration, an advantage for all patients and particularly for those that do not tolerate the toxicity of the chemotherapeutic agent well.

The anti-tumor activity of a particular composition (e.g., antisense, ribozyme, intrabody), or a combination of such compositions, can be evaluated using various *in vitro* and *in vivo* assay systems. *In vitro* assays that evaluate therapeutic activity include cell growth assays, soft agar assays and other assays indicative of tumor promoting activity, binding assays capable of determining the extent to which a therapeutic composition will inhibit the binding of 161P2F10B to a binding partner, etc.

In vivo, the effect of a 161P2F10B therapeutic composition can be evaluated in a suitable animal model. For example, xenogenic prostate cancer models can be used, wherein human prostate cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice (Klein *et al.*, 1997, Nature Medicine 3: 402-408). For example, PCT Patent Application WO98/16628 and U.S. Patent 6,107,540 describe various xenograft models of human prostate cancer capable of recapitulating the development of primary tumors, micrometastasis, and the formation of osteoblastic metastases characteristic of late stage disease. Efficacy can be predicted using assays that measure inhibition of tumor formation, tumor regression or metastasis, and the like.

In vivo assays that evaluate the promotion of apoptosis are useful in evaluating therapeutic compositions. In one embodiment, xenografts from tumor bearing mice treated with the therapeutic composition can be examined for the presence of apoptotic foci and compared to untreated control xenograft-bearing mice. The extent to which apoptotic foci are found in the tumors of the treated mice provides an indication of the therapeutic efficacy of the composition.

The therapeutic compositions used in the practice of the foregoing methods can be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery method. Suitable carriers include any material that when combined with the therapeutic composition retains the anti-tumor function of the therapeutic composition and is generally non-reactive with the patient's immune system. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like (see, generally, Remington's Pharmaceutical Sciences 16th Edition, A. Osal., Ed., 1980).

Therapeutic formulations can be solubilized and administered via any route capable of delivering the therapeutic composition to the tumor site. Potentially effective routes of administration include, but are not limited to, intravenous, parenteral, intraperitoneal, intramuscular, intratumor, intradermal, intraorgan, orthotopic, and the like. A preferred formulation for intravenous injection comprises the therapeutic composition in a solution of preserved bacteriostatic water, sterile unpreserved water, and/or diluted in polyvinylchloride or polyethylene bags containing 0.9% sterile Sodium Chloride for Injection, USP. Therapeutic protein preparations can be lyophilized and stored as sterile powders, preferably under vacuum, and then reconstituted in bacteriostatic water (containing for example, benzyl alcohol preservative) or in sterile water prior to injection.

Dosages and administration protocols for the treatment of cancers using the foregoing methods will vary with the method and the target cancer, and will generally depend on a number of other factors appreciated in the art.

XIII.) Identification, Characterization and Use of Modulators of 161P2F10b**Methods to Identify and Use Modulators**

In one embodiment, screening is performed to identify modulators that induce or suppress a particular expression profile, suppress or induce specific pathways, preferably generating the associated phenotype thereby. In another embodiment, having identified differentially expressed genes important in a particular state; screens are performed to identify modulators that alter expression of individual genes, either increase or decrease. In another embodiment, screening is performed to identify modulators that alter a biological function of the expression product of a differentially expressed gene. Again, having identified the importance of a gene in a particular state, screens are performed to identify agents that bind and/or modulate the biological activity of the gene product.

In addition, screens are done for genes that are induced in response to a candidate agent. After identifying a modulator (one that suppresses a cancer expression pattern leading to a normal expression pattern, or a modulator of a cancer gene that leads to expression of the gene as in normal tissue) a screen is performed to identify genes that are specifically modulated in response to the agent. Comparing expression profiles between normal tissue and agent-treated cancer tissue reveals genes that are not expressed in normal tissue or cancer tissue, but are expressed in agent treated tissue, and vice versa. These agent-specific sequences are identified and used by methods described herein for cancer genes or proteins. In particular these sequences and the proteins they encode are used in marking or identifying agent-treated cells. In addition, antibodies are raised against the agent-induced proteins and used to target novel therapeutics to the treated cancer tissue sample.

Modulator-related Identification and Screening Assays:**Gene Expression-related Assays**

Proteins, nucleic acids, and antibodies of the invention are used in screening assays. The cancer-associated proteins, antibodies, nucleic acids, modified proteins and cells containing these sequences are used in screening assays, such as evaluating the effect of drug candidates on a "gene expression profile," expression profile of polypeptides or alteration of biological function. In one embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent (e.g., Davis, GF, et al, J Biol Screen 7:69 (2002); Zlokarnik, et al., Science 279:84-8 (1998); Heid, Genome Res 6:986-94, 1996).

The cancer proteins, antibodies, nucleic acids, modified proteins and cells containing the native or modified cancer proteins or genes are used in screening assays. That is, the present invention comprises methods for screening for compositions which modulate the cancer phenotype or a physiological function of a cancer protein of the invention. This is done on a gene itself or by evaluating the effect of drug candidates on a "gene expression profile" or biological function. In one embodiment, expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring after treatment with a candidate agent, see Zlokarnik, supra.

A variety of assays are executed directed to the genes and proteins of the invention. Assays are run on an individual nucleic acid or protein level. That is, having identified a particular gene as up regulated in cancer, test compounds are screened for the ability to modulate gene expression or for binding to the cancer protein of the invention. "Modulation" in this context includes an increase or a decrease in gene expression. The preferred amount of modulation will depend on the original change of the gene expression in normal versus tissue undergoing cancer, with changes of at least 10%, preferably 50%, more preferably 100-300%, and in some embodiments 300-1000% or greater. Thus, if a gene exhibits a 4-fold increase in cancer tissue compared to normal tissue, a decrease of about four-fold is often desired; similarly, a 10-fold

decrease in cancer tissue compared to normal tissue a target value of a 10-fold increase in expression by the test compound is often desired. Modulators that exacerbate the type of gene expression seen in cancer are also useful, e.g., as an upregulated target in further analyses.

The amount of gene expression is monitored using nucleic acid probes and the quantification of gene expression levels, or, alternatively, a gene product itself is monitored, e.g., through the use of antibodies to the cancer protein and standard immunoassays. Proteomics and separation techniques also allow for quantification of expression.

Expression Monitoring to Identify Compounds that Modify Gene Expression

In one embodiment, gene expression monitoring, i.e., an expression profile, is monitored simultaneously for a number of entities. Such profiles will typically involve one or more of the genes of Figure 2. In this embodiment, e.g., cancer nucleic acid probes are attached to biochips to detect and quantify cancer sequences in a particular cell. Alternatively, PCR can be used. Thus, a series, e.g., wells of a microtiter plate, can be used with dispensed primers in desired wells. A PCR reaction can then be performed and analyzed for each well.

Expression monitoring is performed to identify compounds that modify the expression of one or more cancer-associated sequences, e.g., a polynucleotide sequence set out in Figure 2. Generally, a test modulator is added to the cells prior to analysis. Moreover, screens are also provided to identify agents that modulate cancer, modulate cancer proteins of the invention, bind to a cancer protein of the invention, or interfere with the binding of a cancer protein of the invention and an antibody or other binding partner.

In one embodiment, high throughput screening methods involve providing a library containing a large number of potential therapeutic compounds (candidate compounds). Such "combinatorial chemical libraries" are then screened in one or more assays to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds," as compounds for screening, or as therapeutics.

In certain embodiments, combinatorial libraries of potential modulators are screened for an ability to bind to a cancer polypeptide or to modulate activity. Conventionally, new chemical entities with useful properties are generated by identifying a chemical compound (called a "lead compound") with some desirable property or activity, e.g., inhibiting activity, creating variants of the lead compound, and evaluating the property and activity of those variant compounds. Often, high throughput screening (HTS) methods are employed for such an analysis.

As noted above, gene expression monitoring is conveniently used to test candidate modulators (e.g., protein, nucleic acid or small molecule). After the candidate agent has been added and the cells allowed to incubate for a period, the sample containing a target sequence to be analyzed is, e.g., added to a biochip.

If required, the target sequence is prepared using known techniques. For example, a sample is treated to lyse the cells, using known lysis buffers, electroporation, etc., with purification and/or amplification such as PCR performed as appropriate. For example, an in vitro transcription with labels covalently attached to the nucleotides is performed. Generally, the nucleic acids are labeled with biotin-FITC or PE, or with cy3 or cy5.

The target sequence can be labeled with, e.g., a fluorescent, a chemiluminescent, a chemical, or a radioactive signal, to provide a means of detecting the target sequence's specific binding to a probe. The label also can be an enzyme, such as alkaline phosphatase or horseradish peroxidase, which when provided with an appropriate substrate produces a product that is detected. Alternatively, the label is a labeled compound or small molecule, such as an enzyme inhibitor, that binds but is not catalyzed or altered by the enzyme. The label also can be a moiety or compound, such as, an epitope tag or biotin which specifically binds to streptavidin. For the example of biotin, the streptavidin is labeled as described above, thereby, providing a detectable signal for the bound target sequence. Unbound labeled streptavidin is typically removed prior to analysis.

As will be appreciated by those in the art, these assays can be direct hybridization assays or can comprise "sandwich assays", which include the use of multiple probes, as is generally outlined in U.S. Patent Nos. 5, 681,702; 5,597,909; 5,545,730; 5,594,117; 5,591,584; 5,571,670; 5,580,731; 5,571,670; 5,591,584; 5,624,802; 5,635,352; 5,594,118; 5,359,100; 5,124, 246; and 5,681,697. In this embodiment, in general, the target nucleic acid is prepared as outlined above, and then added to the biochip comprising a plurality of nucleic acid probes, under conditions that allow the formation of a hybridization complex.

A variety of hybridization conditions are used in the present invention, including high, moderate and low stringency conditions as outlined above. The assays are generally run under stringency conditions which allow formation of the label probe hybridization complex only in the presence of target. Stringency can be controlled by altering a step parameter that is a thermodynamic variable, including, but not limited to, temperature, formamide concentration, salt concentration, chaotropic salt concentration pH, organic solvent concentration, etc. These parameters may also be used to control non-specific binding, as is generally outlined in U.S. Patent No. 5,681,697. Thus, it can be desirable to perform certain steps at higher stringency conditions to reduce non-specific binding.

The reactions outlined herein can be accomplished in a variety of ways. Components of the reaction can be added simultaneously, or sequentially, in different orders, with preferred embodiments outlined below. In addition, the reaction may include a variety of other reagents. These include salts, buffers, neutral proteins, e.g. albumin, detergents, etc. which can be used to facilitate optimal hybridization and detection, and/or reduce nonspecific or background interactions. Reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may also be used as appropriate, depending on the sample preparation methods and purity of the target. The assay data are analyzed to determine the expression levels of individual genes, and changes in expression levels as between states, forming a gene expression profile.

Biological Activity-related Assays

The invention provides methods identify or screen for a compound that modulates the activity of a cancer-related gene or protein of the invention. The methods comprise adding a test compound, as defined above, to a cell comprising a cancer protein of the invention. The cells contain a recombinant nucleic acid that encodes a cancer protein of the invention. In another embodiment, a library of candidate agents is tested on a plurality of cells.

In one aspect, the assays are evaluated in the presence or absence or previous or subsequent exposure of physiological signals, e.g. hormones, antibodies, peptides, antigens, cytokines, growth factors, action potentials, pharmacological agents including chemotherapeutics, radiation, carcinogenics, or other cells (i.e., cell-cell contacts). In another example, the determinations are made at different stages of the cell cycle process. In this way, compounds that modulate genes or proteins of the invention are identified. Compounds with pharmacological activity are able to enhance or interfere with the activity of the cancer protein of the invention. Once identified, similar structures are evaluated to identify critical structural features of the compound.

In one embodiment, a method of modulating (e.g., inhibiting) cancer cell division is provided; the method comprises administration of a cancer modulator. In another embodiment, a method of modulating (e.g., inhibiting) cancer is provided; the method comprises administration of a cancer modulator. In a further embodiment, methods of treating cells or individuals with cancer are provided; the method comprises administration of a cancer modulator.

In one embodiment, a method for modulating the status of a cell that expresses a gene of the invention is provided. As used herein status comprises such art-accepted parameters such as growth, proliferation, survival, function, apoptosis, senescence, location, enzymatic activity, signal transduction, etc. of a cell. In one embodiment, a cancer inhibitor is an

antibody as discussed above. In another embodiment, the cancer inhibitor is an antisense molecule. A variety of cell growth, proliferation, and metastasis assays are known to those of skill in the art, as described herein.

High Throughput Screening to Identify Modulators

The assays to identify suitable modulators are amenable to high throughput screening. Preferred assays thus detect enhancement or inhibition of cancer gene transcription, inhibition or enhancement of polypeptide expression, and inhibition or enhancement of polypeptide activity.

In one embodiment, modulators evaluated in high throughput screening methods are proteins, often naturally occurring proteins or fragments of naturally occurring proteins. Thus, e.g., cellular extracts containing proteins, or random or directed digests of proteinaceous cellular extracts, are used. In this way, libraries of proteins are made for screening in the methods of the invention. Particularly preferred in this embodiment are libraries of bacterial, fungal, viral, and mammalian proteins, with the latter being preferred, and human proteins being especially preferred. Particularly useful test compound will be directed to the class of proteins to which the target belongs, e.g., substrates for enzymes, or ligands and receptors.

Use of Soft Agar Growth and Colony Formation to Identify and Characterize Modulators

Normal cells require a solid substrate to attach and grow. When cells are transformed, they lose this phenotype and grow detached from the substrate. For example, transformed cells can grow in stirred suspension culture or suspended in semi-solid media, such as semi-solid or soft agar. The transformed cells, when transfected with tumor suppressor genes, can regenerate normal phenotype and once again require a solid substrate to attach to and grow. Soft agar growth or colony formation in assays are used to identify modulators of cancer sequences, which when expressed in host cells, inhibit abnormal cellular proliferation and transformation. A modulator reduces or eliminates the host cells' ability to grow suspended in solid or semisolid media, such as agar.

Techniques for soft agar growth or colony formation in suspension assays are described in Freshney, *Culture of Animal Cells a Manual of Basic Technique* (3rd ed., 1994). See also, the methods section of Garkavtsev et al. (1996), *supra*.

Evaluation of Contact Inhibition and Growth Density Limitation to Identify and Characterize Modulators

Normal cells typically grow in a flat and organized pattern in cell culture until they touch other cells. When the cells touch one another, they are contact inhibited and stop growing. Transformed cells, however, are not contact inhibited and continue to grow to high densities in disorganized foci. Thus, transformed cells grow to a higher saturation density than corresponding normal cells. This is detected morphologically by the formation of a disoriented monolayer of cells or cells in foci. Alternatively, labeling index with (³H)-thymidine at saturation density is used to measure density limitation of growth, similarly an MTT or Alamar blue assay will reveal proliferation capacity of cells and the the ability of modulators to affect same. See Freshney (1994), *supra*. Transformed cells, when transfected with tumor suppressor genes, can regenerate a normal phenotype and become contact inhibited and would grow to a lower density.

In this assay, labeling index with (³H)-thymidine at saturation density is a preferred method of measuring density limitation of growth. Transformed host cells are transfected with a cancer-associated sequence and are grown for 24 hours at saturation density in non-limiting medium conditions. The percentage of cells labeling with (³H)-thymidine is determined by incorporated cpm.

Contact independent growth is used to identify modulators of cancer sequences, which had led to abnormal cellular proliferation and transformation. A modulator reduces or eliminates contact independent growth, and returns the cells to a normal phenotype.

Evaluation of Growth Factor or Serum Dependence to Identify and Characterize Modulators

Transformed cells have lower serum dependence than their normal counterparts (see, e.g., Temin, *J. Natl. Cancer Inst.* 37:167-175 (1966); Eagle et al., *J. Exp. Med* 131:836-879 (1970)); Freshney, *supra*. This is in part due to release of

various growth factors by the transformed cells. The degree of growth factor or serum dependence of transformed host cells can be compared with that of control. For example, growth factor or serum dependence of a cell is monitored in methods to identify and characterize compounds that modulate cancer-associated sequences of the invention.

Use of Tumor-specific Marker Levels to Identify and Characterize Modulators

Tumor cells release an increased amount of certain factors (hereinafter "tumor specific markers") than their normal counterparts. For example, plasminogen activator (PA) is released from human glioma at a higher level than from normal brain cells (see, e.g., Gullino, *Angiogenesis, Tumor Vascularization, and Potential Interference with Tumor Growth*, in *Biological Responses in Cancer*, pp. 178-184 (Mihich (ed.) 1985)). Similarly, Tumor Angiogenesis Factor (TAF) is released at a higher level in tumor cells than their normal counterparts. See, e.g., Folkman, *Angiogenesis and Cancer*, *Sem Cancer Biol.* (1992)), while bFGF is released from endothelial tumors (Ensoli, B et al).

Various techniques which measure the release of these factors are described in Freshney (1994), *supra*. Also, see, Unkless et al., *J. Biol. Chem.* 249:4295-4305 (1974); Strickland & Beers, *J. Biol. Chem.* 251:5694-5702 (1976); Whur et al., *Br. J. Cancer* 42:305-312 (1980); Gullino, *Angiogenesis, Tumor Vascularization, and Potential Interference with Tumor Growth*, in *Biological Responses in Cancer*, pp. 178-184 (Mihich (ed.) 1985); Freshney, *Anticancer Res.* 5:111-130 (1985). For example, tumor specific marker levels are monitored in methods to identify and characterize compounds that modulate cancer-associated sequences of the invention.

Invasiveness into Matrigel to Identify and Characterize Modulators

The degree of invasiveness into Matrigel or an extracellular matrix constituent can be used as an assay to identify and characterize compounds that modulate cancer associated sequences. Tumor cells exhibit a positive correlation between malignancy and invasiveness of cells into Matrigel or some other extracellular matrix constituent. In this assay, tumorigenic cells are typically used as host cells. Expression of a tumor suppressor gene in these host cells would decrease invasiveness of the host cells. Techniques described in *Cancer Res.* 1999; 59:6010; Freshney (1994), *supra*, can be used. Briefly, the level of invasion of host cells is measured by using filters coated with Matrigel or some other extracellular matrix constituent. Penetration into the gel, or through to the distal side of the filter, is rated as invasiveness, and rated histologically by number of cells and distance moved, or by prelabeling the cells with ¹²⁵I and counting the radioactivity on the distal side of the filter or bottom of the dish. See, e.g., Freshney (1984), *supra*.

Evaluation of Tumor Growth *In Vivo* to Identify and Characterize Modulators

Effects of cancer-associated sequences on cell growth are tested in transgenic or immune-suppressed organisms. Transgenic organisms are prepared in a variety of art-accepted ways. For example, knock-out transgenic organisms, e.g., mammals such as mice, are made, in which a cancer gene is disrupted or in which a cancer gene is inserted. Knock-out transgenic mice are made by insertion of a marker gene or other heterologous gene into the endogenous cancer gene site in the mouse genome via homologous recombination. Such mice can also be made by substituting the endogenous cancer gene with a mutated version of the cancer gene, or by mutating the endogenous cancer gene, e.g., by exposure to carcinogens.

To prepare transgenic chimeric animals, e.g., mice, a DNA construct is introduced into the nuclei of embryonic stem cells. Cells containing the newly engineered genetic lesion are injected into a host mouse embryo, which is re-implanted into a recipient female. Some of these embryos develop into chimeric mice that possess germ cells some of which are derived from the mutant cell line. Therefore, by breeding the chimeric mice it is possible to obtain a new line of mice containing the introduced genetic lesion (see, e.g., Capecchi et al., *Science* 244:1288 (1989)). Chimeric mice can be derived according to US Patent 6,365,797, issued 2 April 2002; US Patent 6,107,540 issued 22 August 2000; Hogan et al.,

Manipulating the Mouse Embryo: A laboratory Manual, Cold Spring Harbor Laboratory (1988) and Teratocarcinomas and Embryonic Stem Cells: A Practical Approach, Robertson, ed., IRL Press, Washington, D.C., (1987).

Alternatively, various immune-suppressed or immune-deficient host animals can be used. For example, a genetically athymic "nude" mouse (see, e.g., Giovanella et al., J. Natl. Cancer Inst. 52:921 (1974)), a SCID mouse, a thymectomized mouse, or an irradiated mouse (see, e.g., Bradley et al., Br. J. Cancer 38:263 (1978); Selby et al., Br. J. Cancer 41:52 (1980)) can be used as a host. Transplantable tumor cells (typically about 10^6 cells) injected into isogenic hosts produce invasive tumors in a high proportion of cases, while normal cells of similar origin will not. In hosts which developed invasive tumors, cells expressing cancer-associated sequences are injected subcutaneously or orthotopically. Mice are then separated into groups, including control groups and treated experimental groups) e.g. treated with a modulator). After a suitable length of time, preferably 4-8 weeks, tumor growth is measured (e.g., by volume or by its two largest dimensions, or weight) and compared to the control. Tumors that have statistically significant reduction (using, e.g., Student's T test) are said to have inhibited growth.

In Vitro Assays to Identify and Characterize Modulators

Assays to identify compounds with modulating activity can be performed in vitro. For example, a cancer polypeptide is first contacted with a potential modulator and incubated for a suitable amount of time, e.g., from 0.5 to 48 hours. In one embodiment, the cancer polypeptide levels are determined in vitro by measuring the level of protein or mRNA. The level of protein is measured using immunoassays such as Western blotting, ELISA and the like with an antibody that selectively binds to the cancer polypeptide or a fragment thereof. For measurement of mRNA, amplification, e.g., using PCR, LCR, or hybridization assays, e. g., Northern hybridization, RNase protection, dot blotting, are preferred. The level of protein or mRNA is detected using directly or indirectly labeled detection agents, e.g., fluorescently or radioactively labeled nucleic acids, radioactively or enzymatically labeled antibodies, and the like, as described herein.

Alternatively, a reporter gene system can be devised using a cancer protein promoter operably linked to a reporter gene such as luciferase, green fluorescent protein, CAT, or P-gal. The reporter construct is typically transfected into a cell. After treatment with a potential modulator, the amount of reporter gene transcription, translation, or activity is measured according to standard techniques known to those of skill in the art (Davis GF, supra; Gonzalez, J. & Negulescu, P. Curr. Opin. Biotechnol. 1998: 9:624).

As outlined above, in vitro screens are done on individual genes and gene products. That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of the expression of the gene or the gene product itself is performed.

In one embodiment, screening for modulators of expression of specific gene(s) is performed. Typically, the expression of only one or a few genes is evaluated. In another embodiment, screens are designed to first find compounds that bind to differentially expressed proteins. These compounds are then evaluated for the ability to modulate differentially expressed activity. Moreover, once initial candidate compounds are identified, variants can be further screened to better evaluate structure activity relationships.

Binding Assays to Identify and Characterize Modulators

In binding assays in accordance with the invention, a purified or isolated gene product of the invention is generally used. For example, antibodies are generated to a protein of the invention, and immunoassays are run to determine the amount and/or location of protein. Alternatively, cells comprising the cancer proteins are used in the assays.

Thus, the methods comprise combining a cancer protein of the invention and a candidate compound such as a ligand, and determining the binding of the compound to the cancer protein of the invention. Preferred embodiments utilize

the human cancer protein; animal models of human disease of can also be developed and used. Also, other analogous mammalian proteins also can be used as appreciated by those of skill in the art. Moreover, in some embodiments variant or derivative cancer proteins are used.

Generally, the cancer protein of the invention, or the ligand, is non-diffusibly bound to an insoluble support. The support can, e.g., be one having isolated sample receiving areas (a microtiter plate, an array, etc.). The insoluble supports can be made of any composition to which the compositions can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports can be solid or porous and of any convenient shape.

Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharide, nylon, nitrocellulose, or Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition to the support is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusible. Preferred methods of binding include the use of antibodies which do not sterically block either the ligand binding site or activation sequence when attaching the protein to the support, direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or ligand/binding agent to the support, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

Once a cancer protein of the invention is bound to the support, and a test compound is added to the assay. Alternatively, the candidate binding agent is bound to the support and the cancer protein of the invention is then added. Binding agents include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc.

Of particular interest are assays to identify agents that have a low toxicity for human cells. A wide variety of assays can be used for this purpose, including proliferation assays, cAMP assays, labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

A determination of binding of the test compound (ligand, binding agent, modulator, etc.) to a cancer protein of the invention can be done in a number of ways. The test compound can be labeled, and binding determined directly, e.g., by attaching all or a portion of the cancer protein of the invention to a solid support, adding a labeled candidate compound (e.g., a fluorescent label), washing off excess reagent, and determining whether the label is present on the solid support. Various blocking and washing steps can be utilized as appropriate.

In certain embodiments, only one of the components is labeled, e.g., a protein of the invention or ligands labeled. Alternatively, more than one component is labeled with different labels, e.g., ^{125} I, for the proteins and a fluorophor for the compound. Proximity reagents, e.g., quenching or energy transfer reagents are also useful.

Competitive Binding to Identify and Characterize Modulators

In one embodiment, the binding of the "test compound" is determined by competitive binding assay with a "competitor." The competitor is a binding moiety that binds to the target molecule (e.g., a cancer protein of the invention). Competitors include compounds such as antibodies, peptides, binding partners, ligands, etc. Under certain circumstances, the competitive binding between the test compound and the competitor displaces the test compound. In one embodiment, the test compound is labeled. Either the test compound, the competitor, or both, is added to the protein for a time sufficient to allow binding. Incubations are performed at a temperature that facilitates optimal activity, typically between four and 40°C.

Incubation periods are typically optimized, e.g., to facilitate rapid high throughput screening; typically between zero and one hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

In one embodiment, the competitor is added first, followed by the test compound. Displacement of the competitor is an indication that the test compound is binding to the cancer protein and thus is capable of binding to, and potentially modulating, the activity of the cancer protein. In this embodiment, either component can be labeled. Thus, e.g., if the competitor is labeled, the presence of label in the post-test compound wash solution indicates displacement by the test compound. Alternatively, if the test compound is labeled, the presence of the label on the support indicates displacement.

In an alternative embodiment, the test compound is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor indicates that the test compound binds to the cancer protein with higher affinity than the competitor. Thus, if the test compound is labeled, the presence of the label on the support, coupled with a lack of competitor binding, indicates that the test compound binds to and thus potentially modulates the cancer protein of the invention.

Accordingly, the competitive binding methods comprise differential screening to identify agents that are capable of modulating the activity of the cancer proteins of the invention. In this embodiment, the methods comprise combining a cancer protein and a competitor in a first sample. A second sample comprises a test compound, the cancer protein, and a competitor. The binding of the competitor is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to the cancer protein and potentially modulating its activity. That is, if the binding of the competitor is different in the second sample relative to the first sample, the agent is capable of binding to the cancer protein.

Alternatively, differential screening is used to identify drug candidates that bind to the native cancer protein, but cannot bind to modified cancer proteins. For example the structure of the cancer protein is modeled and used in rational drug design to synthesize agents that interact with that site, agents which generally do not bind to site-modified proteins. Moreover, such drug candidates that affect the activity of a native cancer protein are also identified by screening drugs for the ability to either enhance or reduce the activity of such proteins.

Positive controls and negative controls can be used in the assays. Preferably control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples occurs for a time sufficient to allow for the binding of the agent to the protein. Following incubation, samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples can be counted in a scintillation counter to determine the amount of bound compound.

A variety of other reagents can be included in the screening assays. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc. which are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., can be used. The mixture of components is added in an order that provides for the requisite binding.

Use of Polynucleotides to Down-regulate or Inhibit a Protein of the Invention.

Polynucleotide modulators of cancer can be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand-binding molecule, as described in WO 91/04753. Suitable ligand-binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide

or its conjugated version into the cell. Alternatively, a polynucleotide modulator of cancer can be introduced into a cell containing the target nucleic acid sequence, e.g., by formation of a polynucleotide-lipid complex, as described in WO 90/10448. It is understood that the use of antisense molecules or knock out and knock in models may also be used in screening assays as discussed above, in addition to methods of treatment.

Inhibitory and Antisense Nucleotides

In certain embodiments, the activity of a cancer-associated protein is down-regulated, or entirely inhibited, by the use of antisense polynucleotide or inhibitory small nuclear RNA (snRNA), i.e., a nucleic acid complementary to, and which can preferably hybridize specifically to, a coding mRNA nucleic acid sequence, e.g., a cancer protein of the invention, mRNA, or a subsequence thereof. Binding of the antisense polynucleotide to the mRNA reduces the translation and/or stability of the mRNA.

In the context of this invention, antisense polynucleotides can comprise naturally occurring nucleotides, or synthetic species formed from naturally occurring subunits or their close homologs. Antisense polynucleotides may also have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur containing species which are known for use in the art. Analogs are comprised by this invention so long as they function effectively to hybridize with nucleotides of the invention. See, e.g., Isis Pharmaceuticals, Carlsbad, CA; Sequitor, Inc., Natick, MA.

Such antisense polynucleotides can readily be synthesized using recombinant means, or can be synthesized in vitro. Equipment for such synthesis is sold by several vendors, including Applied Biosystems. The preparation of other oligonucleotides such as phosphorothioates and alkylated derivatives is also well known to those of skill in the art.

Antisense molecules as used herein include antisense or sense oligonucleotides. Sense oligonucleotides can, e.g., be employed to block transcription by binding to the anti-sense strand. The antisense and sense oligonucleotide comprise a single stranded nucleic acid sequence (either RNA or DNA) capable of binding to target mRNA (sense) or DNA (antisense) sequences for cancer molecules. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment generally at least about 12 nucleotides, preferably from about 12 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, e.g., Stein & Cohen (Cancer Res. 48:2659 (1988) and van der Krol et al. (BioTechniques 6:958 (1988)).

Ribozymes

In addition to antisense polynucleotides, ribozymes can be used to target and inhibit transcription of cancer-associated nucleotide sequences. A ribozyme is an RNA molecule that catalytically cleaves other RNA molecules. Different kinds of ribozymes have been described, including group I ribozymes, hammerhead ribozymes, hairpin ribozymes, RNase P, and axhead ribozymes (see, e.g., Castanotto et al., Adv. in Pharmacology 25: 289-317 (1994) for a general review of the properties of different ribozymes).

The general features of hairpin ribozymes are described, e.g., in Hampel et al., Nucl. Acids Res. 18:299-304 (1990); European Patent Publication No. 0360257; U.S. Patent No. 5,254,678. Methods of preparing are well known to those of skill in the art (see, e.g., WO 94/26877; Ojwang et al., Proc. Natl. Acad. Sci. USA 90:6340-6344 (1993); Yamada et al., Human Gene Therapy 1:39-45 (1994); Leavitt et al., Proc. Natl. Acad. Sci. USA 92:699- 703 (1995); Leavitt et al., Human Gene Therapy 5: 1151-120 (1994); and Yamada et al., Virology 205: 121-126 (1994)).

Use of Modulators in Phenotypic Screening

In one embodiment, a test compound is administered to a population of cancer cells, which have an associated cancer expression profile. By "administration" or "contacting" herein is meant that the modulator is added to the cells in such a manner as to allow the modulator to act upon the cell, whether by uptake and intracellular action, or by action at the cell

surface. In some embodiments, a nucleic acid encoding a proteinaceous agent (i.e., a peptide) is put into a viral construct such as an adenoviral or retroviral construct, and added to the cell, such that expression of the peptide agent is accomplished, e.g., PCT US97/01019. Regulatable gene therapy systems can also be used. Once the modulator has been administered to the cells, the cells are washed if desired and are allowed to incubate under preferably physiological conditions for some period. The cells are then harvested and a new gene expression profile is generated. Thus, e.g., cancer tissue is screened for agents that modulate, e.g., induce or suppress, the cancer phenotype. A change in at least one gene, preferably many, of the expression profile indicates that the agent has an effect on cancer activity. Similarly, altering a biological function or a signaling pathway is indicative of modulator activity. By defining such a signature for the cancer phenotype, screens for new drugs that alter the phenotype are devised. With this approach, the drug target need not be known and need not be represented in the original gene/protein expression screening platform, nor does the level of transcript for the target protein need to change. The modulator inhibiting function will serve as a surrogate marker

As outlined above, screens are done to assess genes or gene products. That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of either the expression of the gene or the gene product itself is performed.

Use of Modulators to Affect Peptides of the Invention

Measurements of cancer polypeptide activity, or of the cancer phenotype are performed using a variety of assays. For example, the effects of modulators upon the function of a cancer polypeptide(s) are measured by examining parameters described above. A physiological change that affects activity is used to assess the influence of a test compound on the polypeptides of this invention. When the functional outcomes are determined using intact cells or animals, a variety of effects can be assessed such as, in the case of a cancer associated with solid tumors, tumor growth, tumor metastasis, neovascularization, hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., by Northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as cGNIP.

Methods of Identifying Characterizing Cancer-associated Sequences

Expression of various gene sequences is correlated with cancer. Accordingly, disorders based on mutant or variant cancer genes are determined. In one embodiment, the invention provides methods for identifying cells containing variant cancer genes, e.g., determining the presence of, all or part, the sequence of at least one endogenous cancer gene in a cell. This is accomplished using any number of sequencing techniques. The invention comprises methods of identifying the cancer genotype of an individual, e.g., determining all or part of the sequence of at least one gene of the invention in the individual. This is generally done in at least one tissue of the individual, e.g., a tissue set forth in Table I, and may include the evaluation of a number of tissues or different samples of the same tissue. The method may include comparing the sequence of the sequenced gene to a known cancer gene, i.e., a wild-type gene to determine the presence of family members, homologies, mutations or variants. The sequence of all or part of the gene can then be compared to the sequence of a known cancer gene to determine if any differences exist. This is done using any number of known homology programs, such as BLAST, Bestfit, etc. The presence of a difference in the sequence between the cancer gene of the patient and the known cancer gene correlates with a disease state or a propensity for a disease state, as outlined herein.

In a preferred embodiment, the cancer genes are used as probes to determine the number of copies of the cancer gene in the genome. The cancer genes are used as probes to determine the chromosomal localization of the cancer genes. Information such as chromosomal localization finds use in providing a diagnosis or prognosis in particular when chromosomal abnormalities such as translocations, and the like are identified in the cancer gene locus.

XIV.) Kits/Articles of Manufacture

For use in the diagnostic and therapeutic applications described herein, kits are also within the scope of the invention. Such kits can comprise a carrier, package or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in the method. For example, the container(s) can comprise a probe that is or can be detectably labeled. Such probe can be an antibody or polynucleotide specific for a Figure 2-related protein or a Figure 2 gene or message, respectively. Where the method utilizes nucleic acid hybridization to detect the target nucleic acid, the kit can also have containers containing nucleotide(s) for amplification of the target nucleic acid sequence and/or a container comprising a reporter-means, such as a biotin-binding protein, such as avidin or streptavidin, bound to a reporter molecule, such as an enzymatic, florescent, or radioisotope label. The kit can include all or part of the amino acid sequences in Figure 2 or Figure 3 or analogs thereof, or a nucleic acid molecules that encodes such amino acid sequences.

The kit of the invention will typically comprise the container described above and one or more other containers comprising materials desirable from a commercial and user standpoint, including buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and package inserts with instructions for use.

A label can be present on the container to indicate that the composition is used for a specific therapy or non-therapeutic application, such as a diagnostic or laboratory application, and can also indicate directions for either *in vivo* or *in vitro* use, such as those described herein. Directions and or other information can also be included on an insert(s) or label(s) which is included with or on the kit.

The terms "kit" and "article of manufacture" can be used as synonyms.

In another embodiment of the invention, an article(s) of manufacture containing compositions, such as amino acid sequence(s), small molecule(s), nucleic acid sequence(s), and/or antibody(s), e.g., materials useful for the diagnosis, prognosis, prophylaxis and/or treatment of neoplasias of tissues such as those set forth in Table I is provided. The article of manufacture typically comprises at least one container and at least one label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers can be formed from a variety of materials such as glass or plastic. The container can hold amino acid sequence(s), small molecule(s), nucleic acid sequence(s), and/or antibody(s), in one embodiment the container holds a polynucleotide for use in examining the mRNA expression profile of a cell, together with reagents used for this purpose.

The container can alternatively hold a composition which is effective for treating, diagnosis, prognosing or prophylaxing a condition and can have a sterile access port (for example the container can be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agents in the composition can be an antibody capable of specifically binding 161P2F10B and modulating the function of 161P2F10B.

The label can be on or associated with the container. A label can be on a container when letters, numbers or other characters forming the label are molded or etched into the container itself; a label can be associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. The label can indicate that the composition is used for diagnosing, treating, prophylaxing or prognosing a condition, such as a neoplasia of a tissue set forth in Table I. The article of manufacture can further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and/or dextrose solution. It can further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, stirrers, needles, syringes, and/or package inserts with indications and/or instructions for use.

EXAMPLES:

Various aspects of the invention are further described and illustrated by way of the several examples that follow, none of which are intended to limit the scope of the invention.

Example 1: SSH-Generated Isolation of cDNA Fragment of the STEAP Gene

To isolate genes that are over-expressed in kidney cancer we used the Suppression Subtractive Hybridization (SSH) procedure using cDNA derived from kidney cancer patient tissues.

The 161P2F10B SSH cDNA sequence was derived from a subtraction consisting of a kidney cancer minus normal kidney and a mixture of 9 normal tissues: stomach, skeletal muscle, lung, brain, liver, kidney, pancreas, small intestine and heart. By RT-PCR, the 161P2F10B cDNA was identified as highly expressed in kidney cancer pool, with lower expression detected in prostate cancer xenograft pool, prostate cancer pool, colon cancer pool, lung cancer pool, ovary cancer pool, breast cancer pool, metastasis cancer pool, pancreas cancer pool, 2 different prostate cancer metastasis to lymph node, VP1 and VP2. (Figure 14).

The 161P2F10B SSH cDNA sequence of 182 bp matches the cDNA for phosphodiesterase I/nucleotide pyrophosphatase 3 (PDNP3). The full-length 161P2F10B cDNA and ORF are described in Figure 2 with the protein sequence listed in Figure 3.

Materials and Methods**RNA Isolation:**

Tumor tissues were homogenized in Trizol reagent (Life Technologies, Gibco BRL) using 10 ml/ g tissue or 10 ml/ 10⁸ cells to isolate total RNA. Poly A RNA was purified from total RNA using Qiagen's Oligotex mRNA Mini and Midi kits. Total and mRNA were quantified by spectrophotometric analysis (O.D. 260/280 nm) and analyzed by gel electrophoresis.

Oligonucleotides:

The following HPLC purified oligonucleotides were used.

DPNCDN (cDNA synthesis primer):

5'TTTTGATCAAGCTT₃₀3' (SEQ ID NO: 29)

Adaptor 1:

5'CTAATACGACTCACTATAGGGCTCGAGCGGCCGCCCCGGGCAG3' (SEQ ID NO: 30)

3'GGCCCGTCCTAG5' (SEQ ID NO: 31)

Adaptor 2:

5'GTAATACGACTCACTATAGGGCAGCGTGGTCGCGGCCGAG3' (SEQ ID NO: 32)

3'CGGCTCCTAG5' (SEQ ID NO: 33)

PCR primer 1:

5'CTAATACGACTCACTATAGGGC3' (SEQ ID NO: 34)

Nested primer (NP)1:

5'TCGAGCGGCCGCCCCGGGCAGGA3' (SEQ ID NO: 35)

Nested primer (NP)2:

5'AGCGTGGTCGCGGCCGAGGA3' (SEQ ID NO: 36)

Suppression Subtractive Hybridization:

Suppression Subtractive Hybridization (SSH) was used to identify cDNAs corresponding to genes that may be differentially expressed in prostate cancer. The SSH reaction utilized cDNA from kidney cancer patient specimens. The gene 161P2F10B was derived from kidney cancer patient tissues minus normal kidney and a mixture of 9 normal tissues: stomach, skeletal muscle, lung, brain, liver, kidney, pancreas, small intestine and heart. The SSH DNA sequence (Figure 1) was identified.

The cDNA derived from kidney cancer patient tissues was used as the source of the "driver" cDNA, while the cDNA from normal tissues was used as the source of the "tester" cDNA. Double stranded cDNAs corresponding to tester and driver cDNAs were synthesized from 2 µg of poly(A)⁺ RNA isolated from the relevant tissue, as described above, using CLONTECH's PCR-Select cDNA Subtraction Kit and 1 ng of oligonucleotide DPNCDN as primer. First- and second-strand synthesis were carried out as described in the Kit's user manual protocol (CLONTECH Protocol No. PT1117-1, Catalog No. K1804-1). The resulting cDNA was digested with Dpn II for 3 hrs at 37°C. Digested cDNA was extracted with phenol/chloroform (1:1) and ethanol precipitated.

Tester cDNA was generated by diluting 1 µl of Dpn II digested cDNA from the relevant tissue source (see above) (400 ng) in 5 µl of water. The diluted cDNA (2 µl, 160 ng) was then ligated to 2 µl of Adaptor 1 and Adaptor 2 (10 µM), in separate ligation reactions, in a total volume of 10 µl at 16°C overnight, using 400 u of T4 DNA ligase (CLONTECH). Ligation was terminated with 1 µl of 0.2 M EDTA and heating at 72°C for 5 min.

The first hybridization was performed by adding 1.5 µl (600 ng) of driver cDNA to each of two tubes containing 1.5 µl (20 ng) Adaptor 1- and Adaptor 2- ligated tester cDNA. In a final volume of 4 µl, the samples were overlaid with mineral oil, denatured in an MJ Research thermal cycler at 98°C for 1.5 minutes, and then were allowed to hybridize for 8 hrs at 68°C. The two hybridizations were then mixed together with an additional 1 µl of fresh denatured driver cDNA and were allowed to hybridize overnight at 68°C. The second hybridization was then diluted in 200 µl of 20 mM Hepes, pH 8.3, 50 mM NaCl, 0.2 mM EDTA, heated at 70°C for 7 min. and stored at -20°C.

PCR Amplification, Cloning and Sequencing of Gene Fragments Generated from SSH:

To amplify gene fragments resulting from SSH reactions, two PCR amplifications were performed. In the primary PCR reaction 1 µl of the diluted final hybridization mix was added to 1 µl of PCR primer 1 (10 µM), 0.5 µl dNTP mix (10 µM), 2.5 µl 10 x reaction buffer (CLONTECH) and 0.5 µl 50 x Advantage cDNA polymerase Mix (CLONTECH) in a final volume of 25 µl. PCR 1 was conducted using the following conditions: 75°C for 5 min., 94°C for 25 sec., then 27 cycles of 94°C for 10 sec, 66°C for 30 sec, 72°C for 1.5 min. Five separate primary PCR reactions were performed for each experiment. The products were pooled and diluted 1:10 with water. For the secondary PCR reaction, 1 µl from the pooled and diluted primary PCR reaction was added to the same reaction mix as used for PCR 1, except that primers NP1 and NP2 (10 µM) were used instead of PCR primer 1. PCR 2 was performed using 10-12 cycles of 94°C for 10 sec, 68°C for 30 sec, and 72°C for 1.5 minutes. The PCR products were analyzed using 2% agarose gel electrophoresis.

The PCR products were inserted into pCR2.1 using the T/A vector cloning kit (Invitrogen). Transformed *E. coli* were subjected to blue/white and ampicillin selection. White colonies were picked and arrayed into 96 well plates and were grown in liquid culture overnight. To identify inserts, PCR amplification was performed on 1 ml of bacterial culture using the conditions of PCR1 and NP1 and NP2 as primers. PCR products were analyzed using 2% agarose gel electrophoresis.

Bacterial clones were stored in 20% glycerol in a 96 well format. Plasmid DNA was prepared, sequenced, and subjected to nucleic acid homology searches of the GenBank, dBest, and NCI-CGAP databases.

RT-PCR Expression Analysis:

First strand cDNAs can be generated from 1 µg of mRNA with oligo (dT)12-18 priming using the Gibco-BRL Superscript Preamplification system. The manufacturer's protocol was used which included an incubation for 50 min at 42°C with reverse transcriptase followed by RNase H treatment at 37°C for 20 min. After completing the reaction, the volume can be increased to 200 µl with water prior to normalization. First strand cDNAs from 16 different normal human tissues can be obtained from Clontech.

Normalization of the first strand cDNAs from multiple tissues was performed by using the primers 5'atatcgccgcgctcgtcgtcgacaa3' (SEQ ID NO: 37) and 5'agccacacgcagctcattgtagaagg 3' (SEQ ID NO: 38) to amplify β-actin. First strand cDNA (5 µl) were amplified in a total volume of 50 µl containing 0.4 µM primers, 0.2 µM each dNTPs, 1XPCR buffer (Clontech, 10 mM Tris-HCl, 1.5 mM MgCl₂, 50 mM KCl, pH8.3) and 1X KlenTaq DNA polymerase (Clontech). Five µl of the PCR reaction can be removed at 18, 20, and 22 cycles and used for agarose gel electrophoresis. PCR was performed using an MJ Research thermal cycler under the following conditions: Initial denaturation can be at 94°C for 15 sec, followed by a 18, 20, and 22 cycles of 94°C for 15, 65°C for 2 min, 72°C for 5 sec. A final extension at 72°C was carried out for 2 min. After agarose gel electrophoresis, the band intensities of the 283 bp β-actin bands from multiple tissues were compared by visual inspection. Dilution factors for the first strand cDNAs were calculated to result in equal β-actin band intensities in all tissues after 22 cycles of PCR. Three rounds of normalization can be required to achieve equal band intensities in all tissues after 22 cycles of PCR.

To determine expression levels of the 161P2F10B gene, 5 µl of normalized first strand cDNA were analyzed by PCR using 26, and 30 cycles of amplification. Semi-quantitative expression analysis can be achieved by comparing the PCR products at cycle numbers that give light band intensities.

A typical RT-PCR expression analysis is shown in Figure 14. RT-PCR expression analysis was performed on first strand cDNAs generated using pools of tissues from multiple samples. The cDNAs were shown to be normalized using beta-actin PCR. Strong expression of 161P2F10B was observed in kidney cancer pool. Expression was also detected in VP1, prostate cancer xenograft pool, prostate cancer pool and colon cancer pool. Low expression was observed in VP2, lung cancer pool, ovary cancer pool, breast cancer pool, metastasis pool, pancreas cancer pool, and in the 2 different prostate cancer metastasis to lymph node.

Example 2: Isolation of Full Length 161P2F10B Encoding cDNA

To isolate genes that are involved in kidney cancer, an experiment was conducted using kidney cancer patient specimens. The gene 161P2F10B was derived from a subtraction consisting of kidney cancer specimens, minus normal kidney mixed with a cocktail of 9 normal tissues: stomach, skeletal muscle, lung, brain, liver, kidney, pancreas, small intestine and heart. The SSH DNA sequence (Figure 1) was designated 161P2F10B. cDNA clone 161P2F10B was cloned from kidney cancer specimens (Figure 2 and Figure 3). 161P2F10B showed homology to the gene ENPP3. The amino acid alignment of 161P2F10B with ENPP3 is shown in Figure 4 (also, see, e.g., Buhning, et al., Blood 97:3303-3305 (2001)).

Example 3: Chromosomal Mapping of 161P2F10B

Chromosomal localization can implicate genes in disease pathogenesis. Several chromosome mapping approaches are available including fluorescent *in situ* hybridization (FISH), human/hamster radiation hybrid (RH) panels (Walter et al., 1994; Nature Genetics 7:22; Research Genetics, Huntsville AL), human-rodent somatic cell hybrid panels such as is available from the Coriell Institute (Camden, New Jersey), and genomic viewers utilizing BLAST homologies to sequenced and mapped genomic clones (NCBI, Bethesda, Maryland).

161P2F10B maps to chromosome 6q22, using 161P2F10B sequence and the NCBI BLAST tool located at the World Wide Web at (ncbi.nlm.nih.gov/genome/seq/page.cgi?F=HsBlast.html&&ORG=Hs).

Example 4: Expression Analysis of 161P2F10B

To compare expression of 161P2F10B in normal versus patient cancer tissues, RT-PCR experiment was performed using normal and patient cancer tissues (Figure 14). First strand cDNA was generated from normal stomach, normal brain, normal heart, normal liver, normal skeletal muscle, normal testis, normal prostate, normal bladder, normal kidney, normal colon, normal lung, normal pancreas, and a pool of cancer specimens from prostate cancer patients, bladder cancer patients, kidney cancer patients, colon cancer patients, lung cancer patients, pancreas cancer patients, a pool of prostate cancer xenografts (LAPC-4AD, LAPC-4AI, LAPC-9AD and LAPC-9AI), and a pool of 2 patient prostate metastasis to lymph node. Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 161P2F10B, was performed at 26 and 30 cycles of amplification. Samples were run on an agarose gel, and PCR products were quantitated using the Alphasizer software. Results show strong expression in prostate cancer, bladder cancer, kidney cancer, colon cancer, lung cancer, pancreas cancer, bone cancer, lymphoma cancer, uterus cancer, compared to all normal tissues tested. Strong expression was also detected in the xenograft pool as well as the prostate cancer metastasis to lymph node specimens.

Figure 15 & Table LIX shows expression of 161P2F10B in a panel of kidney cancer clear cell carcinoma (A), kidney cancer papillary carcinoma (B), and in uterus patient cancer specimens (C). First strand cDNA was prepared from the patient specimens. Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 161P2F10B, was performed at 26 and 30 cycles of amplification. Samples were run on an agarose gel, and PCR products were quantitated using the Alphasizer software. Expression was recorded as absent, low, medium or strong. Results show expression of 161P2F10B in 94.7% of clear cell renal carcinoma, 62.5% of papillary renal cell carcinoma, and in 61.5% of uterus cancer.

The restricted expression of 161P2F10B in normal tissues and the upregulation detected in kidney cancer, in kidney cancer metastasis, as well as in prostate, bladder, colon, lung, pancreas, bone, lymphoma, uterus, breast, and ovary cancers, suggest that 161P2F10B is a potential therapeutic target and a diagnostic marker for human cancers.

Example 5: Transcript Variants of 161P2F10B

Transcript variants are variants of mature mRNA from the same gene, which arise by alternative transcription or alternative splicing. Alternative transcripts are transcripts from the same gene but start transcription at different points. Splice variants are mRNA variants spliced differently from the same transcript. In eukaryotes, when a multi-exon gene is transcribed from genomic DNA, the initial RNA is spliced to produce functional mRNA, which has only exons and is used for translation into an amino acid sequence. Accordingly, a given gene can have zero to many alternative transcripts and each transcript can have zero to many splice variants. Each transcript variant has a unique exon makeup, and can have different coding and/or non-coding (5' or 3' end) portions, from the original transcript. Transcript variants can code for similar or different proteins with the same or a similar function or can encode proteins with different functions, and can be expressed in the same tissue at the same time, or in different tissues at the same time, or in the same tissue at different times, or in different tissues at different times. Proteins encoded by transcript variants can have similar or different cellular or extracellular localizations, e.g., secreted versus intracellular.

Transcript variants are identified by a variety of art-accepted methods. For example, alternative transcripts and splice variants are identified by full-length cloning experiment, or by use of full-length transcript and EST sequences. First, all human ESTs were grouped into clusters which show direct or indirect identity with each other. Second, ESTs in the same

cluster were further grouped into sub-clusters and assembled into a consensus sequence. The original gene sequence is compared to the consensus sequence(s) or other full-length sequences. Each consensus sequence is a potential splice variant for that gene. Even when a variant is identified that is not a full-length clone, that portion of the variant is very useful for antigen generation and for further cloning of the full-length splice variant, using techniques known in the art.

Moreover, computer programs are available in the art that identify transcript variants based on genomic sequences. Genomic-based transcript variant identification programs include FgenesH (A. Salamov and V. Solovyev, "Ab initio gene finding in Drosophila genomic DNA," *Genome Research*. 2000 April; 10(4):516-22); Grail (URL compbio.ornl.gov/Grail-bin/EmptyGrailForm) and GenScan (URL genes.mit.edu/GENSCAN.html). For a general discussion of splice variant identification protocols see., e.g., Southan, C., A genomic perspective on human proteases, *FEBS Lett*. 2001 Jun 8; 498(2-3):214-8; de Souza, S.J., *et al.*, Identification of human chromosome 22 transcribed sequences with ORF expressed sequence tags, *Proc. Natl Acad Sci U S A*. 2000 Nov 7; 97(23):12690-3.

To further confirm the parameters of a transcript variant, a variety of techniques are available in the art, such as full-length cloning, proteomic validation, PCR-based validation, and 5' RACE validation, etc. (see e.g., Proteomic Validation: Brennan, S.O., *et al.*, Albumin banks peninsula: a new termination variant characterized by electrospray mass spectrometry, *Biochem Biophys Acta*. 1999 Aug 17; 1433(1-2):321-6; Ferranti P, *et al.*, Differential splicing of pre-messenger RNA produces multiple forms of mature caprine alpha(s1)-casein, *Eur J Biochem*. 1997 Oct 1; 249(1):1-7. For PCR-based Validation: Wellmann S, *et al.*, Specific reverse transcription-PCR quantification of vascular endothelial growth factor (VEGF) splice variants by LightCycler technology, *Clin Chem*. 2001 Apr; 47(4):654-60; Jia, H.P., *et al.*, Discovery of new human beta-defensins using a genomics-based approach, *Gene*. 2001 Jan 24; 263(1-2):211-8. For PCR-based and 5' RACE Validation: Brigle, K.E., *et al.*, Organization of the murine reduced folate carrier gene and identification of variant splice forms, *Biochem Biophys Acta*. 1997 Aug 7; 1353(2): 191-8).

It is known in the art that genomic regions are modulated in cancers. When the genomic region to which a gene maps is modulated in a particular cancer, the alternative transcripts or splice variants of the gene are modulated as well. Disclosed herein is that 161P2F10B has a particular expression profile related to cancer. Alternative transcripts and splice variants of 161P2F10B may also be involved in cancers in the same or different tissues, thus serving as tumor-associated markers/antigens.

Using the full-length gene and EST sequences, two transcript variants were identified, designated as 161P2F10B v.6 and v.7. Compared with 161P2F10B v.1, transcript variant 161P2F10B v.6 has extra 40 bases to the 5' starting site of variant 161P2F10B v.1 transcript and has a different 3' end portion, which is on the same chromosome as other exons in the current version of human genome. Variant 161P2F10B v.7 inserted 130 bases in between positions 121 and 122 of variant 161P2F10B v.1. Theoretically, each different combination of exons in spatial order, e.g. exons 2 and 3, is a potential splice variant. Due to the incorrect assembly of the chromosome region in the current version of human genome, the transcript structure cannot be derived computationally.

Tables LI through LVIII are set forth on a variant by variant bases. Tables LI and LV show the nucleotide sequence of the transcript variant. Tables LII and LVI show the alignment of the transcript variant with nucleic acid sequence of 161P2F10B v.1. Tables LIII and LVII lay out amino acid translation of the transcript variant for the identified reading frame orientation. Tables LIV and LVIII display alignments of the amino acid sequence encoded by the splice variant with that of 161P2F10B v.1.

Example 6: Single Nucleotide Polymorphisms of 161P2F10B

A Single Nucleotide Polymorphism (SNP) is a single base pair variation in a nucleotide sequence at a specific location. At any given point of the genome, there are four possible nucleotide base pairs: A/T, C/G, G/C and T/A. Genotype

refers to the specific base pair sequence of one or more locations in the genome of an individual. Haplotype refers to the base pair sequence of more than one location on the same DNA molecule (or the same chromosome in higher organisms), often in the context of one gene or in the context of several tightly linked genes. SNPs that occur on a cDNA are called cSNPs. These cSNPs may change amino acids of the protein encoded by the gene and thus change the functions of the protein. Some SNPs cause inherited diseases; others contribute to quantitative variations in phenotype and reactions to environmental factors including diet and drugs among individuals. Therefore, SNPs and/or combinations of alleles (called haplotypes) have many applications, including diagnosis of inherited diseases, determination of drug reactions and dosage, identification of genes responsible for diseases, and analysis of the genetic relationship between individuals (P. Nowotny, J. M. Kwon and A. M. Goate, " SNP analysis to dissect human traits," *Curr. Opin. Neurobiol.* 2001 Oct; 11(5):637-641; M. Pirmohamed and B. K. Park, "Genetic susceptibility to adverse drug reactions," *Trends Pharmacol. Sci.* 2001 Jun; 22(6):298-305; J. H. Riley, C. J. Allan, E. Lai and A. Roses, " The use of single nucleotide polymorphisms in the isolation of common disease genes," *Pharmacogenomics.* 2000 Feb; 1(1):39-47; R. Judson, J. C. Stephens and A. Windemuth, "The predictive power of haplotypes in clinical response," *Pharmacogenomics.* 2000 feb; 1(1):15-26).

SNPs are identified by a variety of art-accepted methods (P. Bean, "The promising voyage of SNP target discovery," *Am. Clin. Lab.* 2001 Oct-Nov; 20(9):18-20; K. M. Weiss, "In search of human variation," *Genome Res.* 1998 Jul; 8(7):691-697; M. M. She, "Enabling large-scale pharmacogenetic studies by high-throughput mutation detection and genotyping technologies," *Clin. Chem.* 2001 Feb; 47(2):164-172). For example, SNPs are identified by sequencing DNA fragments that show polymorphism by gel-based methods such as restriction fragment length polymorphism (RFLP) and denaturing gradient gel electrophoresis (DGGE). They can also be discovered by direct sequencing of DNA samples pooled from different individuals or by comparing sequences from different DNA samples. With the rapid accumulation of sequence data in public and private databases, one can discover SNPs by comparing sequences using computer programs (Z. Gu, L. Hillier and P. Y. Kwok, "Single nucleotide polymorphism hunting in cyberspace," *Hum. Mutat.* 1998; 12(4):221-225). SNPs can be verified and genotype or haplotype of an individual can be determined by a variety of methods including direct sequencing and high throughput microarrays (P. Y. Kwok, "Methods for genotyping single nucleotide polymorphisms," *Annu. Rev. Genomics Hum. Genet.* 2001; 2:235-258; M. Kokoris, K. Dix, K. Moynihan, J. Mathis, B. Erwin, P. Grass, B. Hines and A. Duesterhoeft, "High-throughput SNP genotyping with the Masscode system," *Mol. Diagn.* 2000 Dec; 5(4):329-340). Using the methods described above, four SNPs were identified in the original transcript, 161P2F10B v.1, at positions 408 (A/G), 2502 (A/G), 2663 (A/C) and 3233 (A/C). The transcripts or proteins with alternative alleles were designated as variants 161P2F10B v.2, v.3, v.4, and v.5, respectively. Figure 10 shows the schematic alignment of the SNP variants. Figure 11 shows the schematic alignment of protein variants, corresponding to nucleotide variants. Nucleotide variants that code for the same amino acid sequence as variant 1 are not shown in Figure 11. These alleles of the SNPs, though shown separately here, can occur in different combinations (haplotypes) and in any one of the transcript variants (such as 161P2F10B v.7) that contains the sequence context of the SNPs.

Example 7: Production of Recombinant 161P2F10B in Prokaryotic Systems

To express recombinant 161P2F10B in prokaryotic cells, the full or partial length 161P2F10B cDNA sequences can be cloned into any one of a variety of expression vectors known in the art. One or more of the following regions of 161P2F10B are expressed in these constructs, amino acids 1 to 875; or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 161P2F10B, variants, or analogs thereof.

A. *In vitro* transcription and translation constructs:

pCRII: To generate 161P2F10B sense and anti-sense RNA probes for RNA *in situ* investigations, pCRII constructs (Invitrogen, Carlsbad CA) are generated encoding either all or fragments of the 161P2F10B cDNA. The pCRII vector has

Sp6 and T7 promoters flanking the insert to drive the transcription of 161P2F10B RNA for use as probes in RNA *in situ* hybridization experiments. These probes are used to analyze the cell and tissue expression of 161P2F10B at the RNA level. Transcribed 161P2F10B RNA representing the cDNA amino acid coding region of the 161P2F10B gene is used in *in vitro* translation systems such as the TnT™ Coupled Reticulolysate System (Promega, Corp., Madison, WI) to synthesize 161P2F10B protein.

B. Bacterial Constructs:

pGEX Constructs: To generate recombinant 161P2F10B proteins in bacteria that are fused to the Glutathione S-transferase (GST) protein, all or parts of the 161P2F10B cDNA protein coding sequence are fused to the GST gene by cloning into pGEX-6P-1 or any other GST- fusion vector of the pGEX family (Amersham Pharmacia Biotech, Piscataway, NJ). These constructs allow controlled expression of recombinant 161P2F10B protein sequences with GST fused at the amino-terminus and a six histidine epitope (6X His) at the carboxyl-terminus. The GST and 6X His tags permit purification of the recombinant fusion protein from induced bacteria with the appropriate affinity matrix and allow recognition of the fusion protein with anti-GST and anti-His antibodies. The 6X His tag is generated by adding 6 histidine codons to the cloning primer at the 3' end, e.g., of the open reading frame (ORF). A proteolytic cleavage site, such as the PreScission™ recognition site in pGEX-6P-1, may be employed such that it permits cleavage of the GST tag from 161P2F10B-related protein. The ampicillin resistance gene and pBR322 origin permits selection and maintenance of the pGEX plasmids in *E. coli*.

pMAL Constructs: To generate, in bacteria, recombinant 161P2F10B proteins that are fused to maltose-binding protein (MBP), all or parts of the 161P2F10B cDNA protein coding sequence are fused to the MBP gene by cloning into the pMAL-c2X and pMAL-p2X vectors (New England Biolabs, Beverly, MA). These constructs allow controlled expression of recombinant 161P2F10B protein sequences with MBP fused at the amino-terminus and a 6X His epitope tag at the carboxyl-terminus. The MBP and 6X His tags permit purification of the recombinant protein from induced bacteria with the appropriate affinity matrix and allow recognition of the fusion protein with anti-MBP and anti-His antibodies. The 6X His epitope tag is generated by adding 6 histidine codons to the 3' cloning primer. A Factor Xa recognition site permits cleavage of the pMAL tag from 161P2F10B. The pMAL-c2X and pMAL-p2X vectors are optimized to express the recombinant protein in the cytoplasm or periplasm respectively. Periplasm expression enhances folding of proteins with disulfide bonds.

pET Constructs: To express 161P2F10B in bacterial cells, all or parts of the 161P2F10B cDNA protein coding sequence are cloned into the pET family of vectors (Novagen, Madison, WI). These vectors allow tightly controlled expression of recombinant 161P2F10B protein in bacteria with and without fusion to proteins that enhance solubility, such as NusA and thioredoxin (Trx), and epitope tags, such as 6X His and S-Tag™ that aid purification and detection of the recombinant protein. For example, constructs are made utilizing pET NusA fusion system 43.1 such that regions of the 161P2F10B protein are expressed as amino-terminal fusions to NusA.

C. Yeast Constructs:

pESC Constructs: To express 161P2F10B in the yeast species *Saccharomyces cerevisiae* for generation of recombinant protein and functional studies, all or parts of the 161P2F10B cDNA protein coding sequence are cloned into the pESC family of vectors each of which contain 1 of 4 selectable markers, HIS3, TRP1, LEU2, and URA3 (Stratagene, La Jolla, CA). These vectors allow controlled expression from the same plasmid of up to 2 different genes or cloned sequences containing either Flag™ or Myc epitope tags in the same yeast cell. This system is useful to confirm protein-protein interactions of 161P2F10B. In addition, expression in yeast yields similar post-translational modifications, such as glycosylations and phosphorylations, that are found when expressed in eukaryotic cells.

pESP Constructs: To express 161P2F10B in the yeast species *Saccharomyces pombe*, all or parts of the 161P2F10B cDNA protein coding sequence are cloned into the pESP family of vectors. These vectors allow controlled high level of expression of a 161P2F10B protein sequence that is fused at either the amino terminus or at the carboxyl terminus to

GST which aids purification of the recombinant protein. A Flag™ epitope tag allows detection of the recombinant protein with anti-Flag™ antibody.

Example 8: Production of Recombinant 161P2F10B in Higher Eukaryotic Systems

A. Mammalian Constructs:

To express recombinant 161P2F10B in eukaryotic cells, the full or partial length 161P2F10B cDNA sequences can be cloned into any one of a variety of expression vectors known in the art. One or more of the following regions of 161P2F10B are expressed in these constructs, amino acids 1 to 875; or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 161P2F10B, variants, or analogs thereof.

The constructs were transfected into any one of a wide variety of mammalian cells such as 293T cells or kidney cancer cell lines. Transfected 293T cell lysates were probed with the anti-161P2F10B polyclonal serum and monoclonal antibodies, described herein.

pcDNA3.1/MyHis Constructs: To express 161P2F10B in mammalian cells, the 161P2F10B ORF, or portions thereof, of 161P2F10B with a consensus Kozak translation initiation site were cloned into pcDNA3.1/MyHis Version A (Invitrogen, Carlsbad, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter. The recombinant proteins have the myc epitope and 6X His epitope fused to the carboxyl-terminus. The pcDNA3.1/MyHis vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability, along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Neomycin resistance gene can be used, as it allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*.

The pcDNA3.1/mycHis encoding 161P2F10B was transfected in 293T cells. Cells were harvested 24 hours later and analyzed showing cell surface expression of 161P2F10B driven from the pcDNA3.1/mycHis vector (Figure 29).

pTag5: The 161P2F10B ORF, or portions thereof, of 161P2F10B were cloned into pTag-5. This vector is similar to pAPtag but without the alkaline phosphatase fusion. This construct generates 161P2F10B protein with an amino-terminal IgGκ signal sequence and myc and 6X His epitope tags at the carboxyl-terminus that facilitate detection and affinity purification. The resulting recombinant 161P2F10B protein was optimized for secretion into the media of transfected mammalian cells, and was used as immunogen or ligand to identify proteins such as ligands or receptors that interact with the 161P2F10B proteins. Protein expression is driven from the CMV promoter. The Zeocin resistance gene present in the vector allows for selection of mammalian cells expressing the protein, and the ampicillin resistance gene permits selection of the plasmid in *E. coli*. Figures 31 and 32 show expression and enzymatic activity of the soluble pTag5 expressing 161P2F10B.

PsecFc: The 161P2F10B ORF, or portions thereof, of 161P2F10B were cloned into psecFc. The psecFc vector was assembled by cloning the human immunoglobulin G1 (IgG) Fc (hinge, CH2, CH3 regions) into pSecTag2 (Invitrogen, California). This construct generates an IgG1 Fc fusion at the amino-terminus of the 161P2F10B proteins. 161P2F10B fusions utilizing the murine IgG1 Fc region was also generated and expressed. The resulting recombinant 161P2F10B proteins are optimized for secretion into the media of transfected mammalian cells, and can be used as immunogens or to identify proteins such as ligands or receptors that interact with the 161P2F10B protein. Protein expression is driven from the CMV promoter. The hygromycin resistance gene present in the vector allows for selection of mammalian cells that express the recombinant protein, and the ampicillin resistance gene permits selection of the plasmid in *E. coli*.

pSRα Constructs: To generate mammalian cell lines that express 161P2F10B constitutively, 161P2F10B ORF, or portions thereof, of 161P2F10B are cloned into pSRα constructs. Amphotropic and ecotropic retroviruses were generated by transfection of pSRα constructs into the 293T-10A1 packaging line or co-transfection of pSRα and a helper plasmid

(containing deleted packaging sequences) into the 293 cells. The retrovirus is used to infect a variety of mammalian cell lines, resulting in the integration of the cloned gene, 161P2F10B, into the host cell-lines. Protein expression is driven from a long terminal repeat (LTR). The Neomycin resistance gene present in the vector allows for selection of mammalian cells that express the protein, and the ampicillin resistance gene and ColE1 origin permit selection and maintenance of the plasmid in *E. coli*. The retroviral vectors were thereafter used for infection and generation of various cell lines using, for example, NIH 3T3, 293 Rat-1 cells or kidney cancer cell lines such as Caki and 769 cells. Figures 16 and 30 show cell surface expression of 161P2F10B driven from the pSRa construct in Caki and NIH3T3 cells respectively.

Additional pSRa constructs were generated encoding 3 different mutants of 161P2F10B. The first mutant is D80E, converted the D amino acid residue of the RGD domain at position 80 into E. The other mutants are mutants of the active site of 161P2F10B, converting the T205 amino acid residue at position 205 into either A (T205A), or S (T205S). The 3 mutant pSRa constructs were transfected into a variety of mammalian cell lines such as 293T cells and CaKi kidney cancer cells. Expression was confirmed using anti-161P2F10B monoclonal antibody and phosphodiesterase enzyme activity was tested (Figure 30).

pcDNA4/HisMax Constructs: To express 161P2F10B in mammalian cells, the 161P2F10B ORF, or portions thereof, of 161P2F10B are cloned into pcDNA4/HisMax Version A (Invitrogen, Carlsbad, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter and the SP16 translational enhancer. The recombinant protein has Xpress™ and six histidine (6X His) epitopes fused to the amino-terminus. The pcDNA4/HisMax vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Zeocin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*.

pcDNA3.1/CT-GFP-TOPO Construct: To express 161P2F10B in mammalian cells and to allow detection of the recombinant proteins using fluorescence, the 161P2F10B ORF, or portions thereof, of 161P2F10B with a consensus Kozak translation initiation site are cloned into pcDNA3.1/CT-GFP-TOPO (Invitrogen, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter. The recombinant proteins have the Green Fluorescent Protein (GFP) fused to the carboxyl-terminus facilitating non-invasive, in vivo detection and cell biology studies. The pcDNA3.1/CT-GFP-TOPO vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Neomycin resistance gene allows for selection of mammalian cells that express the protein, and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*. Additional constructs with an amino-terminal GFP fusion are made in pcDNA3.1/NT-GFP-TOPO spanning the entire length of the 161P2F10B proteins.

PAPtag: The 161P2F10B ORF, or portions thereof, of 161P2F10B are cloned into pAPtag-5 (GenHunter Corp. Nashville, TN). This construct generates an alkaline phosphatase fusion at the carboxyl-terminus of the 161P2F10B proteins while fusing the IgGκ signal sequence to the amino-terminus. Constructs are also generated in which alkaline phosphatase with an amino-terminal IgGκ signal sequence is fused to the amino-terminus of 161P2F10B proteins. The resulting recombinant 161P2F10B proteins are optimized for secretion into the media of transfected mammalian cells and can be used to identify proteins such as ligands or receptors that interact with the 161P2F10B proteins. Protein expression is driven from the CMV promoter and the recombinant proteins also contain myc and 6X His epitopes fused at the carboxyl-terminus that facilitates detection and purification. The Zeocin resistance gene present in the vector allows for selection of mammalian cells expressing the recombinant protein and the ampicillin resistance gene permits selection of the plasmid in *E. coli*.

Additional Viral Vectors: Additional constructs are made for viral-mediated delivery and expression of 161P2F10B. High virus titer leading to high-level expression of 161P2F10B is achieved in viral delivery systems such as adenoviral vectors and herpes amplicon vectors. The 161P2F10B coding sequences or fragments thereof are amplified by PCR and subcloned into the AdEasy shuttle vector (Stratagene). Recombination and virus packaging are performed according to the manufacturer's instructions to generate adenoviral vectors. Alternatively, 161P2F10B coding sequences or fragments thereof are cloned into the HSV-1 vector (Imgenex) to generate herpes viral vectors. The viral vectors are thereafter used for infection of various cell lines such as PC3, NIH 3T3, 293 or rat-1 cells.

Regulated Expression Systems: To control expression of 161P2F10B in mammalian cells, coding sequences of 161P2F10B, or portions thereof, are cloned into regulated mammalian expression systems such as the T-Rex System (Invitrogen), the GeneSwitch System (Invitrogen) and the tightly-regulated Ecdysone System (Stratagene). These systems allow the study of the temporal and concentration dependent effects of recombinant 161P2F10B. These vectors are thereafter used to control expression of 161P2F10B in various cell lines such as PC3, NIH 3T3, 293 or rat-1 cells.

B. Baculovirus Expression Systems

To generate recombinant 161P2F10B proteins in a Baculovirus expression system, 161P2F10B ORF, or portions thereof, are cloned into the Baculovirus transfer vector pBlueBac 4.5 (Invitrogen), which provides a His-tag at the N-terminus. Specifically, pBlueBac-161P2F10B is co-transfected with helper plasmid pBac-N-Blue (Invitrogen) into SF9 (*Spodoptera frugiperda*) insect cells to generate recombinant Baculovirus (see Invitrogen instruction manual for details). Baculovirus is then collected from cell supernatant and purified by plaque assay.

Recombinant 161P2F10B protein is then generated by infection of HighFive insect cells (Invitrogen) with purified Baculovirus. Recombinant 161P2F10B protein can be detected using anti-161P2F10B or anti-His-tag antibody. 161P2F10B protein can be purified and used in various cell-based assays or as immunogen to generate polyclonal and monoclonal antibodies specific for 161P2F10B.

Example 9: Antigenicity Profiles and Secondary Structure

Figure 5, Figure 6, Figure 7, Figure 8, and Figure 9 depict graphically five amino acid profiles of the 161P2F10B amino acid sequence, each assessment available by accessing the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) on the ExPasy molecular biology server.

These profiles: Figure 5, Hydrophilicity, (Hopp T.P., Woods K.R., 1981. Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828); Figure 6, Hydropathicity, (Kyte J., Doolittle R.F., 1982. J. Mol. Biol. 157:105-132); Figure 7, Percentage Accessible Residues (Janin J., 1979 Nature 277:491-492); Figure 8, Average Flexibility, (Bhaskaran R., and Ponnuswamy P.K., 1988. Int. J. Pept. Protein Res. 32:242-255); Figure 9, Beta-turn (Deleage, G., Roux B. 1987 Protein Engineering 1:289-294); and optionally others available in the art, such as on the ProtScale website, were used to identify antigenic regions of the 161P2F10B protein. Each of the above amino acid profiles of 161P2F10B were generated using the following ProtScale parameters for analysis: 1) A window size of 9; 2) 100% weight of the window edges compared to the window center; and, 3) amino acid profile values normalized to lie between 0 and 1.

Hydrophilicity (Figure 5), Hydropathicity (Figure 6) and Percentage Accessible Residues (Figure 7) profiles were used to determine stretches of hydrophilic amino acids (i.e., values greater than 0.5 on the Hydrophilicity and Percentage Accessible Residues profile, and values less than 0.5 on the Hydropathicity profile). Such regions are likely to be exposed to the aqueous environment, be present on the surface of the protein, and thus available for immune recognition, such as by antibodies.

Average Flexibility (Figure 8) and Beta-turn (Figure 9) profiles determine stretches of amino acids (i.e., values greater than 0.5 on the Beta-turn profile and the Average Flexibility profile) that are not constrained in secondary structures

such as beta sheets and alpha helices. Such regions are also more likely to be exposed on the protein and thus accessible to immune recognition, such as by antibodies.

Antigenic sequences of the 161P2F10B protein indicated, e.g., by the profiles set forth in Figure 5, Figure 6, Figure 7, Figure 8, and/or Figure 9 are used to prepare immunogens, either peptides or nucleic acids that encode them, to generate therapeutic and diagnostic anti-161P2F10B antibodies. The immunogen can be any 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more than 50 contiguous amino acids, or the corresponding nucleic acids that encode them, from the 161P2F10B protein. In particular, peptide immunogens of the invention can comprise, a peptide region of at least 5 amino acids of Figure 2 in any whole number increment up to 875 that includes an amino acid position having a value greater than 0.5 in the Hydrophilicity profile of Figure 5; a peptide region of at least 5 amino acids of Figure 2 in any whole number increment up to 875 that includes an amino acid position having a value less than 0.5 in the Hydropathicity profile of Figure 6; a peptide region of at least 5 amino acids of Figure 2 in any whole number increment up to 875 that includes an amino acid position having a value greater than 0.5 in the Percent Accessible Residues profile of Figure 7; a peptide region of at least 5 amino acids of Figure 2 in any whole number increment up to 875 that includes an amino acid position having a value greater than 0.5 in the Average Flexibility profile on Figure 8; and, a peptide region of at least 5 amino acids of Figure 2 in any whole number increment up to 875 that includes an amino acid position having a value greater than 0.5 in the Beta-turn profile of Figure 9. Peptide immunogens of the invention can also comprise nucleic acids that encode any of the forgoing.

All immunogens of the invention, peptide or nucleic acid, can be embodied in human unit dose form, or comprised by a composition that includes a pharmaceutical excipient compatible with human physiology.

The secondary structure of 161P2F10B, namely the predicted presence and location of alpha helices, extended strands, and random coils, is predicted from the primary amino acid sequence using the HNN - Hierarchical Neural Network method (Guermeur, 1997, http://pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_nn.html), accessed from the ExPasy molecular biology server (<http://www.expasy.ch/tools/>). The analysis indicates that 161P2F10B is composed 31.31% alpha helix, 11.31% extended strand, and 57.37% random coil (Figure 19A).

Analysis for the potential presence of transmembrane domains in 161P2F10B was carried out using a variety of transmembrane prediction algorithms accessed from the ExPasy molecular biology server (<http://www.expasy.ch/tools/>). The programs predict the presence of 1 transmembrane domain in 161P2F10B, consistent with that of a Type II cell surface protein. Shown graphically in Figure 19 are the results of analysis using the TMpred (Figure 19B) and TMHMM (Figure 19C) prediction programs depicting the location of the transmembrane domain.

Example 10: Generation of 161P2F10B Polyclonal Antibodies

Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. In addition to immunizing with the full length 161P2F10B protein, computer algorithms are employed in design of immunogens that, based on amino acid sequence analysis contain characteristics of being antigenic and available for recognition by the immune system of the immunized host (see the Example entitled "Antigenicity Profiles"). Such regions would be predicted to be hydrophilic, flexible, in beta-turn conformations, and be exposed on the surface of the protein (see, e.g., Figure 5, Figure 6, Figure 7, Figure 8, or Figure 9 for amino acid profiles that indicate such regions of 161P2F10B).

For example, 161P2F10B recombinant bacterial fusion proteins or peptides containing hydrophilic, flexible, beta-turn regions of the 161P2F10B, in which numerous regions are found in the predicted extracellular domain coded by amino acids 45-870, are used as antigens to generate polyclonal antibodies in New Zealand White rabbits. For example, such

immunoprecipitation, fluorescent microscopy, and flow cytometric techniques using cells that endogenously express 161P2F10B are carried out to test specificity.

The anti-serum from the Tag5 161P2F10B immunized rabbit is affinity purified by passage over a column composed of the Tag5 antigen covalently coupled to Affigel matrix (BioRad, Hercules, Calif.). The serum is then further purified by protein G affinity chromatography to isolate the IgG fraction. Serum from rabbits immunized with fusion proteins, such as GST and MBP fusion proteins, are purified by depletion of antibodies reactive to the fusion partner sequence by passage over an affinity column containing the fusion partner either alone or in the context of an irrelevant fusion protein. Sera from other His-tagged antigens and peptide immunized rabbits as well as fusion partner depleted sera are affinity purified by passage over a column matrix composed of the original protein immunogen or free peptide.

Example 11: Generation of 161P2F10B Monoclonal Antibodies (mAbs)

The use of agents to identify the presence of 161P2F10B in biopsy specimens or to neutralize the effect of 161P2F10B has a beneficial effect in diagnosis, prognosis, prophylaxis and/or therapy. One particularly useful class of anti 161P2F10B agents is antibodies, in particular monoclonal antibodies (Mabs) to 161P2F10B. Anti 161P2F10B Abs, such as Mabs, are generated that react with the epitopes of the 161P2F10B protein such that they either indicate its presence, disrupt or modulate its biological function (for example those that would disrupt the interaction with ligands or proteins that mediate or are involved in its biological activity) or are able to carry a toxin to the cell which is expressing 161P2F10B.

The term anti 161P2F10B antibody as used herein is to be understood to cover antibodies to any epitope of the 161P2F10B gene product. Diagnostic Mabs, e.g. those used for imaging or immunocytochemistry, comprise those that specifically bind epitopes of 161P2F10B protein and thus demonstrate its presence. Therapeutic Mabs include those that are useful for diagnosis but also comprise those that specifically bind epitopes of 161P2F10B exposed on the cell surface and thus are useful to modulate growth and survival of cells expressing 161P2F10B by disrupting the function of a cell expressing 161P2F10B and/or disrupting the interaction of cells expressing 161P2F10B and the ligand for 161P2F10B.

Preferred antibodies which form one aspect of the invention include but are not limited to antibodies entitled X41(4)6, X41(3)17, X41(3)50, X41(3)15, X41(3)29 and X41(3)37 secreted by a hybridoma deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No's. _____; and derivatives thereof, the production of which is described herein.

Pathological conditions which are characterized by the presence of 161P2F10B expression include, but are not restricted to, neoplasms of tissues such as those listed in Table I. One aspect of the invention provides a method of detecting the presence of 161P2F10B. A further aspect of the invention provides a method of treatment of conditions characterized by the presence of 161P2F10B, comprising administering an effective amount of an anti 161P2F10B antibody. The administration of anti-161P2F10B antibody is particularly advantageous in the treatment of conditions characterized by the presence of 161P2F10B.

The antibodies against 161P2F10B for use according to the invention can be from any species, and can belong to any immunoglobulin class. Thus, for example, the anti 161P2F10B antibody for use according to the invention can be an immunoglobulin G (IgG), Immunoglobulin M (IgM), immunoglobulin A (IgA), Immunoglobulin E (IgE) or immunoglobulin D (IgD).

The anti 161P2F10B antibody can be from an animal, for example mammalian or avian origin, and can be for example of murine, rat or human origin. The antibody can be a whole immunoglobulin, or a fragment thereof, for example a fragment derived by proteolytic cleavage of a whole antibody, such as F(ab')₂, Fab' or Fab fragments or fragments obtained by recombinant DNA techniques, for example Fv fragments.

Particularly useful antibodies for use according to the invention include humanized or fully human anti 161P2F10B antibodies and fragments thereof. These antibodies are produced by any suitable procedure including, but not restricted to, mammalian cell and bacterial cell fermentation systems.

The anti 161P2F10B Mabs are prepared by immunological techniques employing 161P2F10B antigens. Thus, for example, any suitable host can be injected (immunized) with a suitable reagent which makes 161P2F10B available as an immunogen. Examples of reagents which make 161P2F10B available as an immunogen are purified protein (e.g. the whole extra-cellular domain (ecd) or fragments thereof), peptides designed using the full length protein as a template (e.g. peptides encompassing the catalytic domain), DNA vectors encoding all or truncated fragments of the ecd, recombinant cells expressing 161P2F10B (e.g. Rat-1, Mouse 3T3, Mouse 300.19, and mouse NSO), Cell lines with endogenous 161P2F10B expression (e.g. human UT-7) or xenografts (i.e. patient derived clear cell and papillary xenografts).

Immune cells, for example splenocytes or lymphocytes, are recovered from the immunized host and immortalized, using for example the method of Kohler et al, Eur. J. Immunol 6, 511 (1976), or their immunoglobulin genes can be isolated and transferred to an appropriate DNA vector for expression in an appropriate cell type. The resulting cells, generated by either technique, will be selected to obtain a single genetic line producing a single unique type of antibody more commonly known as a monoclonal antibody. Antibody fragments can be produced using techniques such as enzymatic digestion of whole antibodies e.g. with pepsin (Parham, J. Immunol 131:2895 (1983)) or papain (Lamoyi and Nisonoff, J. Immunol Meth. 56:235 (1983)), or by recombinant DNA techniques.

Suitable hosts for the production of Mab's to 161P2F10B include mice, rats, hamsters and rabbits. For example, mice are immunized with a number of different reagents which make 161P2F10B available as a source of antigenic material (immunogen). The route and timing of the immunizations will depend on the source and/or embodiment of the immunogen. Sources of immunogen for 161P2F10B include, but are not restricted to, peptide, protein, fusion protein, DNA, RNA, cells or cell membranes as detailed above.. These can be used separately as immunogens or in combination to produce a specific immune reaction to 161P2F10B. The use and application of these various immunogens is described fully in the accompanying examples.

Example 12: HLA Class I and Class II Binding Assays

HLA class I and class II binding assays using purified HLA molecules are performed in accordance with disclosed protocols (e.g., PCT publications WO 94/20127 and WO 94/03205; Sidney *et al.*, *Current Protocols in Immunology* 18.3.1 (1998); Sidney, *et al.*, *J. Immunol.* 154:247 (1995); Sette, *et al.*, *Mol. Immunol.* 31:813 (1994)). Briefly, purified MHC molecules (5 to 500 nM) are incubated with various unlabeled peptide inhibitors and 1-10 nM ¹²⁵I-radiolabeled probe peptides as described. Following incubation, MHC-peptide complexes are separated from free peptide by gel filtration and the fraction of peptide bound is determined. Typically, in preliminary experiments, each MHC preparation is titrated in the presence of fixed amounts of radiolabeled peptides to determine the concentration of HLA molecules necessary to bind 10-20% of the total radioactivity. All subsequent inhibition and direct binding assays are performed using these HLA concentrations.

Since under these conditions $[label] < [HLA]$ and $IC_{50} \geq [HLA]$, the measured IC_{50} values are reasonable approximations of the true K_D values. Peptide inhibitors are typically tested at concentrations ranging from 120 µg/ml to 1.2 ng/ml, and are tested in two to four completely independent experiments. To allow comparison of the data obtained in different experiments, a relative binding figure is calculated for each peptide by dividing the IC_{50} of a positive control for inhibition by the IC_{50} for each tested peptide (typically unlabeled versions of the radiolabeled probe peptide). For database purposes, and inter-experiment comparisons, relative binding values are compiled. These values can subsequently be converted back into IC_{50} nM values by dividing the IC_{50} nM of the positive controls for inhibition by the relative binding of the

peptide of interest. This method of data compilation is accurate and consistent for comparing peptides that have been tested on different days, or with different lots of purified MHC.

Binding assays as outlined above may be used to analyze HLA supermotif and/or HLA motif-bearing peptides (see Table IV).

Example 13: Identification of HLA Supermotif- and Motif-Bearing CTL Candidate Epitopes

HLA vaccine compositions of the invention can include multiple epitopes. The multiple epitopes can comprise multiple HLA supermotifs or motifs to achieve broad population coverage. This example illustrates the identification and confirmation of supermotif- and motif-bearing epitopes for the inclusion in such a vaccine composition. Calculation of population coverage is performed using the strategy described below.

Computer searches and algorithms for identification of supermotif and/or motif-bearing epitopes

The searches performed to identify the motif-bearing peptide sequences in the Example entitled "Antigenicity Profiles" and Tables VIII-XXI and XXII-XLIX employ the protein sequence data from the gene product of 161P2F10B set forth in Figures 2 and 3, the specific search peptides used to generate the tables are listed in Table VII.

Computer searches for epitopes bearing HLA Class I or Class II supermotifs or motifs are performed as follows. All translated 161P2F10B protein sequences are analyzed using a text string search software program to identify potential peptide sequences containing appropriate HLA binding motifs; such programs are readily produced in accordance with information in the art in view of known motif/supermotif disclosures. Furthermore, such calculations can be made mentally.

Identified A2-, A3-, and DR-supermotif sequences are scored using polynomial algorithms to predict their capacity to bind to specific HLA-Class I or Class II molecules. These polynomial algorithms account for the impact of different amino acids at different positions, and are essentially based on the premise that the overall affinity (or ΔG) of peptide-HLA molecule interactions can be approximated as a linear polynomial function of the type:

$$"\Delta G" = a_{1i} \times a_{2i} \times a_{3i} \dots \times a_{ni}$$

where a_{ji} is a coefficient which represents the effect of the presence of a given amino acid (j) at a given position (i) along the sequence of a peptide of n amino acids. The crucial assumption of this method is that the effects at each position are essentially independent of each other (i.e., independent binding of individual side-chains). When residue j occurs at position i in the peptide, it is assumed to contribute a constant amount j_i to the free energy of binding of the peptide irrespective of the sequence of the rest of the peptide.

The method of derivation of specific algorithm coefficients has been described in Gulukota *et al.*, *J. Mol. Biol.* 267:1258-126, 1997; (see also Sidney *et al.*, *Human Immunol.* 45:79-93, 1996; and Southwood *et al.*, *J. Immunol.* 160:3363-3373, 1998). Briefly, for all i positions, anchor and non-anchor alike, the geometric mean of the average relative binding (ARB) of all peptides carrying j is calculated relative to the remainder of the group, and used as the estimate of j_i . For Class II peptides, if multiple alignments are possible, only the highest scoring alignment is utilized, following an iterative procedure. To calculate an algorithm score of a given peptide in a test set, the ARB values corresponding to the sequence of the peptide are multiplied. If this product exceeds a chosen threshold, the peptide is predicted to bind. Appropriate thresholds are chosen as a function of the degree of stringency of prediction desired.

Selection of HLA-A2 supertype cross-reactive peptides

Protein sequences from 161P2F10B are scanned utilizing motif identification software, to identify 8-, 9- 10- and 11-mer sequences containing the HLA-A2-supermotif main anchor specificity. Typically, these sequences are then scored using the protocol described above and the peptides corresponding to the positive-scoring sequences are synthesized and tested

for their capacity to bind purified HLA-A*0201 molecules *in vitro* (HLA-A*0201 is considered a prototype A2 supertype molecule).

These peptides are then tested for the capacity to bind to additional A2-supertype molecules (A*0202, A*0203, A*0206, and A*6802). Peptides that bind to at least three of the five A2-supertype alleles tested are typically deemed A2-supertype cross-reactive binders. Preferred peptides bind at an affinity equal to or less than 500 nM to three or more HLA-A2 supertype molecules.

Selection of HLA-A3 supermotif-bearing epitopes

The 161P2F10B protein sequence(s) scanned above is also examined for the presence of peptides with the HLA-A3-supermotif primary anchors. Peptides corresponding to the HLA A3 supermotif-bearing sequences are then synthesized and tested for binding to HLA-A*0301 and HLA-A*1101 molecules, the molecules encoded by the two most prevalent A3-supertype alleles. The peptides that bind at least one of the two alleles with binding affinities of ≤ 500 nM, often ≤ 200 nM, are then tested for binding cross-reactivity to the other common A3-supertype alleles (e.g., A*3101, A*3301, and A*6801) to identify those that can bind at least three of the five HLA-A3-supertype molecules tested.

Selection of HLA-B7 supermotif bearing epitopes

The 161P2F10B protein(s) scanned above is also analyzed for the presence of 8-, 9-, 10-, or 11-mer peptides with the HLA-B7-supermotif. Corresponding peptides are synthesized and tested for binding to HLA-B*0702, the molecule encoded by the most common B7-supertype allele (*i.e.*, the prototype B7 supertype allele). Peptides binding B*0702 with IC_{50} of ≤ 500 nM are identified using standard methods. These peptides are then tested for binding to other common B7-supertype molecules (e.g., B*3501, B*5101, B*5301, and B*5401). Peptides capable of binding to three or more of the five B7-supertype alleles tested are thereby identified.

Selection of A1 and A24 motif-bearing epitopes

To further increase population coverage, HLA-A1 and -A24 epitopes can also be incorporated into vaccine compositions. An analysis of the 161P2F10B protein can also be performed to identify HLA-A1- and A24-motif-containing sequences.

High affinity and/or cross-reactive binding epitopes that bear other motif and/or supermotifs are identified using analogous methodology.

Example 14: Confirmation of Immunogenicity

Cross-reactive candidate CTL A2-supermotif-bearing peptides that are identified as described herein are selected to confirm *in vitro* immunogenicity. Confirmation is performed using the following methodology:

Target Cell Lines for Cellular Screening:

The .221A2.1 cell line, produced by transferring the HLA-A2.1 gene into the HLA-A, -B, -C null mutant human B-lymphoblastoid cell line 721.221, is used as the peptide-loaded target to measure activity of HLA-A2.1-restricted CTL. This cell line is grown in RPMI-1640 medium supplemented with antibiotics, sodium pyruvate, nonessential amino acids and 10% (v/v) heat inactivated FCS. Cells that express an antigen of interest, or transfectants comprising the gene encoding the antigen of interest, can be used as target cells to confirm the ability of peptide-specific CTLs to recognize endogenous antigen.

Primary CTL Induction Cultures:

Generation of Dendritic Cells (DC): PBMCs are thawed in RPMI with 30 $\mu\text{g/ml}$ DNase, washed twice and resuspended in complete medium (RPMI-1640 plus 5% AB human serum, non-essential amino acids, sodium pyruvate, L-glutamine and penicillin/streptomycin). The monocytes are purified by plating 10×10^6 PBMC/well in a 6-well plate. After 2 hours at 37°C , the non-adherent cells are removed by gently shaking the plates and aspirating the supernatants. The wells are washed a total of three times with 3 ml RPMI to remove most of the non-adherent and loosely adherent cells. Three ml of complete medium containing 50 ng/ml of GM-CSF and 1,000 U/ml of IL-4 are then added to each well. $\text{TNF}\alpha$ is added to the DCs on day 6 at 75 ng/ml and the cells are used for CTL induction cultures on day 7.

Induction of CTL with DC and Peptide: CD8^+ T-cells are isolated by positive selection with Dynal immunomagnetic beads (Dynabeads® M-450) and the detach-bead® reagent. Typically about $200\text{--}250 \times 10^6$ PBMC are processed to obtain 24×10^6 CD8^+ T-cells (enough for a 48-well plate culture). Briefly, the PBMCs are thawed in RPMI with 30 $\mu\text{g/ml}$ DNase, washed once with PBS containing 1% human AB serum and resuspended in PBS/1% AB serum at a concentration of 20×10^6 cells/ml. The magnetic beads are washed 3 times with PBS/AB serum, added to the cells (140 μl beads/ 20×10^6 cells) and incubated for 1 hour at 4°C with continuous mixing. The beads and cells are washed 4x with PBS/AB serum to remove the nonadherent cells and resuspended at 100×10^6 cells/ml (based on the original cell number) in PBS/AB serum containing 100 $\mu\text{l/ml}$ detach-bead® reagent and 30 $\mu\text{g/ml}$ DNase. The mixture is incubated for 1 hour at room temperature with continuous mixing. The beads are washed again with PBS/AB/DNase to collect the CD8^+ T-cells. The DC are collected and centrifuged at 1300 rpm for 5-7 minutes, washed once with PBS with 1% BSA, counted and pulsed with 40 $\mu\text{g/ml}$ of peptide at a cell concentration of $1\text{--}2 \times 10^6$ /ml in the presence of 3 $\mu\text{g/ml}$ β_2 -microglobulin for 4 hours at 20°C . The DC are then irradiated (4,200 rads), washed 1 time with medium and counted again.

Setting up induction cultures: 0.25 ml cytokine-generated DC (at 1×10^5 cells/ml) are co-cultured with 0.25 ml of CD8^+ T-cells (at 2×10^6 cell/ml) in each well of a 48-well plate in the presence of 10 ng/ml of IL-7. Recombinant human IL-10 is added the next day at a final concentration of 10 ng/ml and rhuman IL-2 is added 48 hours later at 10 IU/ml.

Restimulation of the induction cultures with peptide-pulsed adherent cells: Seven and fourteen days after the primary induction, the cells are restimulated with peptide-pulsed adherent cells. The PBMCs are thawed and washed twice with RPMI and DNase. The cells are resuspended at 5×10^6 cells/ml and irradiated at ~ 4200 rads. The PBMCs are plated at 2×10^6 in 0.5 ml complete medium per well and incubated for 2 hours at 37°C . The plates are washed twice with RPMI by tapping the plate gently to remove the nonadherent cells and the adherent cells pulsed with 10 $\mu\text{g/ml}$ of peptide in the presence of 3 $\mu\text{g/ml}$ β_2 -microglobulin in 0.25 ml RPMI/5% AB per well for 2 hours at 37°C . Peptide solution from each well is aspirated and the wells are washed once with RPMI. Most of the media is aspirated from the induction cultures (CD8^+ cells) and brought to 0.5 ml with fresh media. The cells are then transferred to the wells containing the peptide-pulsed adherent cells. Twenty four hours later recombinant human IL-10 is added at a final concentration of 10 ng/ml and recombinant human IL2 is added the next day and again 2-3 days later at 50 IU/ml (Tsai *et al.*, *Critical Reviews in Immunology* 18(1-2):65-75, 1998). Seven days later, the cultures are assayed for CTL activity in a ^{51}Cr release assay. In some experiments the cultures are assayed for peptide-specific recognition in the *in situ* IFN γ ELISA at the time of the second restimulation followed by assay of endogenous recognition 7 days later. After expansion, activity is measured in both assays for a side-by-side comparison.

Measurement of CTL lytic activity by ^{51}Cr release.

Seven days after the second restimulation, cytotoxicity is determined in a standard (5 hr) ^{51}Cr release assay by assaying individual wells at a single E:T. Peptide-pulsed targets are prepared by incubating the cells with 10 $\mu\text{g/ml}$ peptide overnight at 37°C .

Adherent target cells are removed from culture flasks with trypsin-EDTA. Target cells are labeled with 200 μCi of ^{51}Cr sodium chromate (Dupont, Wilmington, DE) for 1 hour at 37°C . Labeled target cells are resuspended at 10^6 per ml and

diluted 1:10 with K562 cells at a concentration of 3.3×10^6 /ml (an NK-sensitive erythroblastoma cell line used to reduce non-specific lysis). Target cells (100 μ l) and effectors (100 μ l) are plated in 96 well round-bottom plates and incubated for 5 hours at 37°C. At that time, 100 μ l of supernatant are collected from each well and percent lysis is determined according to the formula:

$$[(\text{cpm of the test sample} - \text{cpm of the spontaneous } ^{51}\text{Cr release sample}) / (\text{cpm of the maximal } ^{51}\text{Cr release sample} - \text{cpm of the spontaneous } ^{51}\text{Cr release sample})] \times 100.$$

Maximum and spontaneous release are determined by incubating the labeled targets with 1% Triton X-100 and media alone, respectively. A positive culture is defined as one in which the specific lysis (sample- background) is 10% or higher in the case of individual wells and is 15% or more at the two highest E:T ratios when expanded cultures are assayed.

In situ Measurement of Human IFN γ Production as an Indicator of Peptide-specific and Endogenous Recognition

Immulon 2 plates are coated with mouse anti-human IFN γ monoclonal antibody (4 μ g/ml 0.1M NaHCO₃, pH8.2) overnight at 4°C. The plates are washed with Ca²⁺, Mg²⁺-free PBS/0.05% Tween 20 and blocked with PBS/10% FCS for two hours, after which the CTLs (100 μ l/well) and targets (100 μ l/well) are added to each well, leaving empty wells for the standards and blanks (which received media only). The target cells, either peptide-pulsed or endogenous targets, are used at a concentration of 1×10^6 cells/ml. The plates are incubated for 48 hours at 37°C with 5% CO₂.

Recombinant human IFN-gamma is added to the standard wells starting at 400 pg or 1200pg/100 microliter/well and the plate incubated for two hours at 37°C. The plates are washed and 100 μ l of biotinylated mouse anti-human IFN-gamma monoclonal antibody (2 microgram/ml in PBS/3%FCS/0.05% Tween 20) are added and incubated for 2 hours at room temperature. After washing again, 100 microliter HRP-streptavidin (1:4000) are added and the plates incubated for one hour at room temperature. The plates are then washed 6x with wash buffer, 100 microliter/well developing solution (TMB 1:1) are added, and the plates allowed to develop for 5-15 minutes. The reaction is stopped with 50 microliter/well 1M H₃PO₄ and read at OD450. A culture is considered positive if it measured at least 50 pg of IFN-gamma/well above background and is twice the background level of expression.

CTL Expansion.

Those cultures that demonstrate specific lytic activity against peptide-pulsed targets and/or tumor targets are expanded over a two week period with anti-CD3. Briefly, 5×10^4 CD8⁺ cells are added to a T25 flask containing the following: 1×10^6 irradiated (4,200 rad) PBMC (autologous or allogeneic) per ml, 2×10^5 irradiated (8,000 rad) EBV- transformed cells per ml, and OKT3 (anti-CD3) at 30ng per ml in RPMI-1640 containing 10% (v/v) human AB serum, non-essential amino acids, sodium pyruvate, 25 μ M 2-mercaptoethanol, L-glutamine and penicillin/streptomycin. Recombinant human IL2 is added 24 hours later at a final concentration of 200IU/ml and every three days thereafter with fresh media at 50IU/ml. The cells are split if the cell concentration exceeds 1×10^6 /ml and the cultures are assayed between days 13 and 15 at E:T ratios of 30, 10, 3 and 1:1 in the ⁵¹Cr release assay or at 1×10^6 /ml in the *in situ* IFN γ assay using the same targets as before the expansion.

Cultures are expanded in the absence of anti-CD3⁺ as follows. Those cultures that demonstrate specific lytic activity against peptide and endogenous targets are selected and 5×10^4 CD8⁺ cells are added to a T25 flask containing the following: 1×10^6 autologous PBMC per ml which have been peptide-pulsed with 10 μ g/ml peptide for two hours at 37°C and irradiated (4,200 rad); 2×10^5 irradiated (8,000 rad) EBV-transformed cells per ml RPMI-1640 containing 10%(v/v) human AB serum, non-essential AA, sodium pyruvate, 25mM 2-ME, L-glutamine and gentamicin.

Immunogenicity of A2 supermotif-bearing peptides

A2-supermotif cross-reactive binding peptides are tested in the cellular assay for the ability to induce peptide-specific CTL in normal individuals. In this analysis, a peptide is typically considered to be an epitope if it induces peptide-specific CTLs in at least individuals, and preferably, also recognizes the endogenously expressed peptide.

Immunogenicity can also be confirmed using PBMCs isolated from patients bearing a tumor that expresses 161P2F10B. Briefly, PBMCs are isolated from patients, re-stimulated with peptide-pulsed monocytes and assayed for the ability to recognize peptide-pulsed target cells as well as transfected cells endogenously expressing the antigen.

Evaluation of A*03/A11 immunogenicity

HLA-A3 supermotif-bearing cross-reactive binding peptides are also evaluated for immunogenicity using methodology analogous for that used to evaluate the immunogenicity of the HLA-A2 supermotif peptides.

Evaluation of B7 immunogenicity

Immunogenicity screening of the B7-supertype cross-reactive binding peptides identified as set forth herein are confirmed in a manner analogous to the confirmation of A2-and A3-supermotif-bearing peptides.

Peptides bearing other supermotifs/motifs, e.g., HLA-A1, HLA-A24 etc. are also confirmed using similar methodology

Example 15: Implementation of the Extended Supermotif to Improve the Binding Capacity of Native Epitopes by Creating Analogs

HLA motifs and supermotifs (comprising primary and/or secondary residues) are useful in the identification and preparation of highly cross-reactive native peptides, as demonstrated herein. Moreover, the definition of HLA motifs and supermotifs also allows one to engineer highly cross-reactive epitopes by identifying residues within a native peptide sequence which can be analoged to confer upon the peptide certain characteristics, e.g. greater cross-reactivity within the group of HLA molecules that comprise a supertype, and/or greater binding affinity for some or all of those HLA molecules. Examples of analoging peptides to exhibit modulated binding affinity are set forth in this example.

Analoging at Primary Anchor Residues

Peptide engineering strategies are implemented to further increase the cross-reactivity of the epitopes. For example, the main anchors of A2-supermotif-bearing peptides are altered, for example, to introduce a preferred L, I, V, or M at position 2, and I or V at the C-terminus.

To analyze the cross-reactivity of the analog peptides, each engineered analog is initially tested for binding to the prototype A2 supertype allele A*0201, then, if A*0201 binding capacity is maintained, for A2-supertype cross-reactivity.

Alternatively, a peptide is confirmed as binding one or all supertype members and then analoged to modulate binding affinity to any one (or more) of the supertype members to add population coverage.

The selection of analogs for immunogenicity in a cellular screening analysis is typically further restricted by the capacity of the parent wild type (WT) peptide to bind at least weakly, *i.e.*, bind at an IC₅₀ of 5000nM or less, to three or more A2 supertype alleles. The rationale for this requirement is that the WT peptides must be present endogenously in sufficient quantity to be biologically relevant. Analoged peptides have been shown to have increased immunogenicity and cross-reactivity by T cells specific for the parent epitope (see, e.g., Parkhurst *et al.*, *J. Immunol.* 157:2539, 1996; and Pogue *et al.*, *Proc. Natl. Acad. Sci. USA* 92:8166, 1995).

In the cellular screening of these peptide analogs, it is important to confirm that analog-specific CTLs are also able to recognize the wild-type peptide and, when possible, target cells that endogenously express the epitope.

Analoging of HLA-A3 and B7-supermotif-bearing peptides

Analogues of HLA-A3 supermotif-bearing epitopes are generated using strategies similar to those employed in analoging HLA-A2 supermotif-bearing peptides. For example, peptides binding to 3/5 of the A3-supertype molecules are engineered at primary anchor residues to possess a preferred residue (V, S, M, or A) at position 2.

The analog peptides are then tested for the ability to bind A*03 and A*11 (prototype A3 supertype alleles). Those peptides that demonstrate ≤ 500 nM binding capacity are then confirmed as having A3-supertype cross-reactivity.

Similarly to the A2- and A3- motif bearing peptides, peptides binding 3 or more B7-supertype alleles can be improved, where possible, to achieve increased cross-reactive binding or greater binding affinity or binding half life. B7 supermotif-bearing peptides are, for example, engineered to possess a preferred residue (V, I, L, or F) at the C-terminal primary anchor position, as demonstrated by Sidney *et al.* (*J. Immunol.* 157:3480-3490, 1996).

Analoging at primary anchor residues of other motif and/or supermotif-bearing epitopes is performed in a like manner.

The analog peptides are then be confirmed for immunogenicity, typically in a cellular screening assay. Again, it is generally important to demonstrate that analog-specific CTLs are also able to recognize the wild-type peptide and, when possible, targets that endogenously express the epitope.

Analoging at Secondary Anchor Residues

Moreover, HLA supermotifs are of value in engineering highly cross-reactive peptides and/or peptides that bind HLA molecules with increased affinity by identifying particular residues at secondary anchor positions that are associated with such properties. For example, the binding capacity of a B7 supermotif-bearing peptide with an F residue at position 1 is analyzed. The peptide is then analoged to, for example, substitute L for F at position 1. The analoged peptide is evaluated for increased binding affinity, binding half life and/or increased cross-reactivity. Such a procedure identifies analoged peptides with enhanced properties.

Engineered analogs with sufficiently improved binding capacity or cross-reactivity can also be tested for immunogenicity in HLA-B7-transgenic mice, following for example, IFA immunization or lipopeptide immunization. Analoged peptides are additionally tested for the ability to stimulate a recall response using PBMC from patients with 161P2F10B-expressing tumors.

Other analoging strategies

Another form of peptide analoging, unrelated to anchor positions, involves the substitution of a cysteine with α -amino butyric acid. Due to its chemical nature, cysteine has the propensity to form disulfide bridges and sufficiently alter the peptide structurally so as to reduce binding capacity. Substitution of α -amino butyric acid for cysteine not only alleviates this problem, but has been shown to improve binding and crossbinding capabilities in some instances (see, e.g., the review by Sette *et al.*, In: Persistent Viral Infections, Eds. R. Ahmed and I. Chen, John Wiley & Sons, England, 1999).

Thus, by the use of single amino acid substitutions, the binding properties and/or cross-reactivity of peptide ligands for HLA supertype molecules can be modulated.

Example 16: Identification and confirmation of 161P2F10B-derived sequences with HLA-DR binding motifs

Peptide epitopes bearing an HLA class II supermotif or motif are identified and confirmed as outlined below using methodology similar to that described for HLA Class I peptides.

Selection of HLA-DR-supermotif-bearing epitopes.

To identify 161P2F10B-derived, HLA class II HTL epitopes, a 161P2F10B antigen is analyzed for the presence of sequences bearing an HLA-DR-motif or supermotif. Specifically, 15-mer sequences are selected comprising a DR-supermotif, comprising a 9-mer core, and three-residue N- and C-terminal flanking regions (15 amino acids total).

Protocols for predicting peptide binding to DR molecules have been developed (Southwood *et al.*, *J. Immunol.* 160:3363-3373, 1998). These protocols, specific for individual DR molecules, allow the scoring, and ranking, of 9-mer core regions. Each protocol not only scores peptide sequences for the presence of DR-supermotif primary anchors (i.e., at

position 1 and position 6) within a 9-mer core, but additionally evaluates sequences for the presence of secondary anchors. Using allele-specific selection tables (see, e.g., Southwood *et al.*, *ibid.*), it has been found that these protocols efficiently select peptide sequences with a high probability of binding a particular DR molecule. Additionally, it has been found that performing these protocols in tandem, specifically those for DR1, DR4w4, and DR7, can efficiently select DR cross-reactive peptides.

The 161P2F10B-derived peptides identified above are tested for their binding capacity for various common HLA-DR molecules. All peptides are initially tested for binding to the DR molecules in the primary panel: DR1, DR4w4, and DR7. Peptides binding at least two of these three DR molecules are then tested for binding to DR2w2 β 1, DR2w2 β 2, DR6w19, and DR9 molecules in secondary assays. Finally, peptides binding at least two of the four secondary panel DR molecules, and thus cumulatively at least four of seven different DR molecules, are screened for binding to DR4w15, DR5w11, and DR8w2 molecules in tertiary assays. Peptides binding at least seven of the ten DR molecules comprising the primary, secondary, and tertiary screening assays are considered cross-reactive DR binders. 161P2F10B-derived peptides found to bind common HLA-DR alleles are of particular interest.

Selection of DR3 motif peptides

Because HLA-DR3 is an allele that is prevalent in Caucasian, Black, and Hispanic populations, DR3 binding capacity is a relevant criterion in the selection of HTL epitopes. Thus, peptides shown to be candidates may also be assayed for their DR3 binding capacity. However, in view of the binding specificity of the DR3 motif, peptides binding only to DR3 can also be considered as candidates for inclusion in a vaccine formulation.

To efficiently identify peptides that bind DR3, target 161P2F10B antigens are analyzed for sequences carrying one of the two DR3-specific binding motifs reported by Geluk *et al.* (*J. Immunol.* 152:5742-5748, 1994). The corresponding peptides are then synthesized and confirmed as having the ability to bind DR3 with an affinity of 1 μ M or better, i.e., less than 1 μ M. Peptides are found that meet this binding criterion and qualify as HLA class II high affinity binders.

DR3 binding epitopes identified in this manner are included in vaccine compositions with DR supermotif-bearing peptide epitopes.

Similarly to the case of HLA class I motif-bearing peptides, the class II motif-bearing peptides are analogized to improve affinity or cross-reactivity. For example, aspartic acid at position 4 of the 9-mer core sequence is an optimal residue for DR3 binding, and substitution for that residue often improves DR 3 binding.

Example 17: Immunogenicity of 161P2F10B-derived HTL epitopes

This example determines immunogenic DR supermotif- and DR3 motif-bearing epitopes among those identified using the methodology set forth herein.

Immunogenicity of HTL epitopes are confirmed in a manner analogous to the determination of immunogenicity of CTL epitopes, by assessing the ability to stimulate HTL responses and/or by using appropriate transgenic mouse models. Immunogenicity is determined by screening for: 1.) *in vitro* primary induction using normal PBMC or 2.) recall responses from patients who have 161P2F10B-expressing tumors.

Example 18: Calculation of phenotypic frequencies of HLA-supertypes in various ethnic backgrounds to determine breadth of population coverage

This example illustrates the assessment of the breadth of population coverage of a vaccine composition comprised of multiple epitopes comprising multiple supermotifs and/or motifs.

In order to analyze population coverage, gene frequencies of HLA alleles are determined. Gene frequencies for each HLA allele are calculated from antigen or allele frequencies utilizing the binomial distribution formulae $gf=1-(\text{SQRT}(1-$

af)) (see, e.g., Sidney *et al.*, *Human Immunol.* 45:79-93, 1996). To obtain overall phenotypic frequencies, cumulative gene frequencies are calculated, and the cumulative antigen frequencies derived by the use of the inverse formula $[af=1-(1-Cgf)^2]$.

Where frequency data is not available at the level of DNA typing, correspondence to the serologically defined antigen frequencies is assumed. To obtain total potential supertype population coverage no linkage disequilibrium is assumed, and only alleles confirmed to belong to each of the superotypes are included (minimal estimates). Estimates of total potential coverage achieved by inter-loci combinations are made by adding to the A coverage the proportion of the non-A covered population that could be expected to be covered by the B alleles considered (e.g., $total=A+B*(1-A)$). Confirmed members of the A3-like supertype are A3, A11, A31, A*3301, and A*6801. Although the A3-like supertype may also include A34, A66, and A*7401, these alleles were not included in overall frequency calculations. Likewise, confirmed members of the A2-like supertype family are A*0201, A*0202, A*0203, A*0204, A*0205, A*0206, A*0207, A*6802, and A*6901. Finally, the B7-like supertype-confirmed alleles are: B7, B*3501-03, B51, B*5301, B*5401, B*5501-2, B*5601, B*6701, and B*7801 (potentially also B*1401, B*3504-06, B*4201, and B*5602).

Population coverage achieved by combining the A2-, A3- and B7-superotypes is approximately 86% in five major ethnic groups. Coverage may be extended by including peptides bearing the A1 and A24 motifs. On average, A1 is present in 12% and A24 in 29% of the population across five different major ethnic groups (Caucasian, North American Black, Chinese, Japanese, and Hispanic). Together, these alleles are represented with an average frequency of 39% in these same ethnic populations. The total coverage across the major ethnicities when A1 and A24 are combined with the coverage of the A2-, A3- and B7-supertype alleles is >95%, see, e.g., Table IV (G). An analogous approach can be used to estimate population coverage achieved with combinations of class II motif-bearing epitopes.

Immunogenicity studies in humans (e.g., Bertoni *et al.*, *J. Clin. Invest.* 100:503, 1997; Doolan *et al.*, *Immunity* 7:97, 1997; and Threlkeld *et al.*, *J. Immunol.* 159:1648, 1997) have shown that highly cross-reactive binding peptides are almost always recognized as epitopes. The use of highly cross-reactive binding peptides is an important selection criterion in identifying candidate epitopes for inclusion in a vaccine that is immunogenic in a diverse population.

With a sufficient number of epitopes (as disclosed herein and from the art), an average population coverage is predicted to be greater than 95% in each of five major ethnic populations. The game theory Monte Carlo simulation analysis, which is known in the art (see e.g., Osborne, M.J. and Rubinstein, A. "A course in game theory" MIT Press, 1994), can be used to estimate what percentage of the individuals in a population comprised of the Caucasian, North American Black, Japanese, Chinese, and Hispanic ethnic groups would recognize the vaccine epitopes described herein. A preferred percentage is 90%. A more preferred percentage is 95%.

Example 19: CTL Recognition Of Endogenously Processed Antigens After Priming

This example confirms that CTL induced by native or analoged peptide epitopes identified and selected as described herein recognize endogenously synthesized, *i.e.*, native antigens.

Effector cells isolated from transgenic mice that are immunized with peptide epitopes, for example HLA-A2 supermotif-bearing epitopes, are re-stimulated *in vitro* using peptide-coated stimulator cells. Six days later, effector cells are assayed for cytotoxicity and the cell lines that contain peptide-specific cytotoxic activity are further re-stimulated. An additional six days later, these cell lines are tested for cytotoxic activity on ^{51}Cr labeled Jurkat-A2.1/K ^{b} target cells in the absence or presence of peptide, and also tested on ^{51}Cr labeled target cells bearing the endogenously synthesized antigen, *i.e.* cells that are stably transfected with 161P2F10B expression vectors.

The results demonstrate that CTL lines obtained from animals primed with peptide epitope recognize endogenously synthesized 161P2F10B antigen. The choice of transgenic mouse model to be used for such an analysis depends upon the epitope(s) that are being evaluated. In addition to HLA-A*0201/K ^{b} transgenic mice, several other transgenic mouse models including mice with human A11, which may also be used to evaluate A3 epitopes, and B7 alleles

have been characterized and others (e.g., transgenic mice for HLA-A1 and A24) are being developed. HLA-DR1 and HLA-DR3 mouse models have also been developed, which may be used to evaluate HTL epitopes.

Example 20: Activity Of CTL-HTL Conjugated Epitopes In Transgenic Mice

This example illustrates the induction of CTLs and HTLs in transgenic mice, by use of a 161P2F10B-derived CTL and HTL peptide vaccine compositions. The vaccine composition used herein comprise peptides to be administered to a patient with a 161P2F10B-expressing tumor. The peptide composition can comprise multiple CTL and/or HTL epitopes. The epitopes are identified using methodology as described herein. This example also illustrates that enhanced immunogenicity can be achieved by inclusion of one or more HTL epitopes in a CTL vaccine composition; such a peptide composition can comprise an HTL epitope conjugated to a CTL epitope. The CTL epitope can be one that binds to multiple HLA family members at an affinity of 500 nM or less, or analogs of that epitope. The peptides may be lipidated, if desired.

Immunization procedures: Immunization of transgenic mice is performed as described (Alexander *et al.*, *J. Immunol.* 159:4753-4761, 1997). For example, A2/K^b mice, which are transgenic for the human HLA A2.1 allele and are used to confirm the immunogenicity of HLA-A*0201 motif- or HLA-A2 supermotif-bearing epitopes, and are primed subcutaneously (base of the tail) with a 0.1 ml of peptide in Incomplete Freund's Adjuvant, or if the peptide composition is a lipidated CTL/HTL conjugate, in DMSO/saline, or if the peptide composition is a polypeptide, in PBS or Incomplete Freund's Adjuvant. Seven days after priming, splenocytes obtained from these animals are restimulated with syngenic irradiated LPS-activated lymphoblasts coated with peptide.

Cell lines: Target cells for peptide-specific cytotoxicity assays are Jurkat cells transfected with the HLA-A2.1/K^b chimeric gene (e.g., Vitiello *et al.*, *J. Exp. Med.* 173:1007, 1991)

In vitro CTL activation: One week after priming, spleen cells (30x10⁶ cells/flask) are co-cultured at 37°C with syngeneic, irradiated (3000 rads), peptide coated lymphoblasts (10x10⁶ cells/flask) in 10 ml of culture medium/T25 flask. After six days, effector cells are harvested and assayed for cytotoxic activity.

Assay for cytotoxic activity: Target cells (1.0 to 1.5x10⁶) are incubated at 37°C in the presence of 200 µl of ⁵¹Cr. After 60 minutes, cells are washed three times and resuspended in R10 medium. Peptide is added where required at a concentration of 1 µg/ml. For the assay, 10⁴ ⁵¹Cr-labeled target cells are added to different concentrations of effector cells (final volume of 200 µl) in U-bottom 96-well plates. After a six hour incubation period at 37°C, a 0.1 ml aliquot of supernatant is removed from each well and radioactivity is determined in a Micromedic automatic gamma counter. The percent specific lysis is determined by the formula: percent specific release = 100 x (experimental release - spontaneous release)/(maximum release - spontaneous release). To facilitate comparison between separate CTL assays run under the same conditions, % ⁵¹Cr release data is expressed as lytic units/10⁶ cells. One lytic unit is arbitrarily defined as the number of effector cells required to achieve 30% lysis of 10,000 target cells in a six hour ⁵¹Cr release assay. To obtain specific lytic units/10⁶, the lytic units/10⁶ obtained in the absence of peptide is subtracted from the lytic units/10⁶ obtained in the presence of peptide. For example, if 30% ⁵¹Cr release is obtained at the effector (E): target (T) ratio of 50:1 (i.e., 5x10⁵ effector cells for 10,000 targets) in the absence of peptide and 5:1 (i.e., 5x10⁴ effector cells for 10,000 targets) in the presence of peptide, the specific lytic units would be: [(1/50,000)-(1/500,000)] × 10⁶ = 18 LU.

The results are analyzed to assess the magnitude of the CTL responses of animals injected with the immunogenic CTL/HTL conjugate vaccine preparation and are compared to the magnitude of the CTL response achieved using, for example, CTL epitopes as outlined above in the Example entitled "Confirmation of Immunogenicity." Analyses similar to this may be performed to confirm the immunogenicity of peptide conjugates containing multiple CTL epitopes and/or multiple HTL epitopes. In accordance with these procedures, it is found that a CTL response is induced, and concomitantly that an HTL response is induced upon administration of such compositions.

Example 21: Selection of CTL and HTL epitopes for inclusion in a 161P2F10B-specific vaccine.

This example illustrates a procedure for selecting peptide epitopes for vaccine compositions of the invention. The peptides in the composition can be in the form of a nucleic acid sequence, either single or one or more sequences (*i.e.*, minigene) that encodes peptide(s), or can be single and/or polyepitopic peptides.

The following principles are utilized when selecting a plurality of epitopes for inclusion in a vaccine composition. Each of the following principles is balanced in order to make the selection.

Epitopes are selected which, upon administration, mimic immune responses that are correlated with 161P2F10B clearance. The number of epitopes used depends on observations of patients who spontaneously clear 161P2F10B. For example, if it has been observed that patients who spontaneously clear 161P2F10B-expressing cells generate an immune response to at least three (3) epitopes from 161P2F10B antigen, then at least three epitopes should be included for HLA class I. A similar rationale is used to determine HLA class II epitopes.

Epitopes are often selected that have a binding affinity of an IC_{50} of 500 nM or less for an HLA class I molecule, or for class II, an IC_{50} of 1000 nM or less; or HLA Class I peptides with high binding scores from the BIMAS web site, at URL bimas.dcrt.nih.gov/.

In order to achieve broad coverage of the vaccine through out a diverse population, sufficient supermotif bearing peptides, or a sufficient array of allele-specific motif bearing peptides, are selected to give broad population coverage. In one embodiment, epitopes are selected to provide at least 80% population coverage. A Monte Carlo analysis, a statistical evaluation known in the art, can be employed to assess breadth, or redundancy, of population coverage.

When creating polyepitopic compositions, or a minigene that encodes same, it is typically desirable to generate the smallest peptide possible that encompasses the epitopes of interest. The principles employed are similar, if not the same, as those employed when selecting a peptide comprising nested epitopes. For example, a protein sequence for the vaccine composition is selected because it has maximal number of epitopes contained within the sequence, *i.e.*, it has a high concentration of epitopes. Epitopes may be nested or overlapping (*i.e.*, frame shifted relative to one another). For example, with overlapping epitopes, two 9-mer epitopes and one 10-mer epitope can be present in a 10 amino acid peptide. Each epitope can be exposed and bound by an HLA molecule upon administration of such a peptide. A multi-epitopic, peptide can be generated synthetically, recombinantly, or via-cleavage from the native source. Alternatively, an analog can be made of this native sequence, whereby one or more of the epitopes comprise substitutions that alter the cross-reactivity and/or binding affinity properties of the polyepitopic peptide. Such a vaccine composition is administered for therapeutic or prophylactic purposes. This embodiment provides for the possibility that an as yet undiscovered aspect of immune system processing will apply to the native nested sequence and thereby facilitate the production of therapeutic or prophylactic immune response-inducing vaccine compositions. Additionally such an embodiment provides for the possibility of motif-bearing epitopes for an HLA makeup that is presently unknown. Furthermore, this embodiment (absent the creating of any analogs) directs the immune response to multiple peptide sequences that are actually present in 161P2F10B, thus avoiding the need to evaluate any junctional epitopes. Lastly, the embodiment provides an economy of scale when producing nucleic acid vaccine compositions. Related to this embodiment, computer programs can be derived in accordance with principles in the art, which identify in a target sequence, the greatest number of epitopes per sequence length.

A vaccine composition comprised of selected peptides, when administered, is safe, efficacious, and elicits an immune response similar in magnitude to an immune response that controls or clears cells that bear or overexpress 161P2F10B.

Example 22: Construction of "Minigene" Multi-Epitope DNA Plasmids

This example discusses the construction of a minigene expression plasmid. Minigene plasmids may, of course, contain various configurations of B cell, CTL and/or HTL epitopes or epitope analogs as described herein.

A minigene expression plasmid typically includes multiple CTL and HTL peptide epitopes. In the present example, HLA-A2, -A3, -B7 supermotif-bearing peptide epitopes and HLA-A1 and -A24 motif-bearing peptide epitopes are used in conjunction with DR supermotif-bearing epitopes and/or DR3 epitopes. HLA class I supermotif or motif-bearing peptide epitopes derived 161P2F10B, are selected such that multiple supermotifs/motifs are represented to ensure broad population coverage. Similarly, HLA class II epitopes are selected from 161P2F10B to provide broad population coverage, *i.e.* both HLA DR-1-4-7 supermotif-bearing epitopes and HLA DR-3 motif-bearing epitopes are selected for inclusion in the minigene construct. The selected CTL and HTL epitopes are then incorporated into a minigene for expression in an expression vector.

Such a construct may additionally include sequences that direct the HTL epitopes to the endoplasmic reticulum. For example, the li protein may be fused to one or more HTL epitopes as described in the art, wherein the CLIP sequence of the li protein is removed and replaced with an HLA class II epitope sequence so that HLA class II epitope is directed to the endoplasmic reticulum, where the epitope binds to an HLA class II molecules.

This example illustrates the methods to be used for construction of a minigene-bearing expression plasmid. Other expression vectors that may be used for minigene compositions are available and known to those of skill in the art.

The minigene DNA plasmid of this example contains a consensus Kozak sequence and a consensus murine kappa Ig-light chain signal sequence followed by CTL and/or HTL epitopes selected in accordance with principles disclosed herein. The sequence encodes an open reading frame fused to the Myc and His antibody epitope tag coded for by the pcDNA 3.1 Myc-His vector.

Overlapping oligonucleotides that can, for example, average about 70 nucleotides in length with 15 nucleotide overlaps, are synthesized and HPLC-purified. The oligonucleotides encode the selected peptide epitopes as well as appropriate linker nucleotides, Kozak sequence, and signal sequence. The final multiepitope minigene is assembled by extending the overlapping oligonucleotides in three sets of reactions using PCR. A Perkin/Elmer 9600 PCR machine is used and a total of 30 cycles are performed using the following conditions: 95°C for 15 sec, annealing temperature (5° below the lowest calculated T_m of each primer pair) for 30 sec, and 72°C for 1 min.

For example, a minigene is prepared as follows. For a first PCR reaction, 5 µg of each of two oligonucleotides are annealed and extended: In an example using eight oligonucleotides, *i.e.*, four pairs of primers, oligonucleotides 1+2, 3+4, 5+6, and 7+8 are combined in 100 µl reactions containing *Pfu* polymerase buffer (1x= 10 mM KCL, 10 mM (NH₄)₂SO₄, 20 mM Tris-chloride, pH 8.75, 2 mM MgSO₄, 0.1% Triton X-100, 100 µg/ml BSA), 0.25 mM each dNTP, and 2.5 U of *Pfu* polymerase. The full-length dimer products are gel-purified, and two reactions containing the product of 1+2 and 3+4, and the product of 5+6 and 7+8 are mixed, annealed, and extended for 10 cycles. Half of the two reactions are then mixed, and 5 cycles of annealing and extension carried out before flanking primers are added to amplify the full length product. The full-length product is gel-purified and cloned into pCR-blunt (Invitrogen) and individual clones are screened by sequencing.

Example 23: The Plasmid Construct and the Degree to Which It Induces Immunogenicity.

The degree to which a plasmid construct, for example a plasmid constructed in accordance with the previous Example, is able to induce immunogenicity is confirmed *in vitro* by determining epitope presentation by APC following transduction or transfection of the APC with an epitope-expressing nucleic acid construct. Such a study determines "antigenicity" and allows the use of human APC. The assay determines the ability of the epitope to be presented by the APC in a context that is recognized by a T cell by quantifying the density of epitope-HLA class I complexes on the cell surface. Quantitation can be performed by directly measuring the amount of peptide eluted from the APC (see, *e.g.*, Sijts *et al.*, *J.*

Immunol. 156:683-692, 1996; Demotz *et al.*, *Nature* 342:682-684, 1989); or the number of peptide-HLA class I complexes can be estimated by measuring the amount of lysis or lymphokine release induced by diseased or transfected target cells, and then determining the concentration of peptide necessary to obtain equivalent levels of lysis or lymphokine release (see, e.g., Kageyama *et al.*, *J. Immunol.* 154:567-576, 1995).

Alternatively, immunogenicity is confirmed through *in vivo* injections into mice and subsequent *in vitro* assessment of CTL and HTL activity, which are analyzed using cytotoxicity and proliferation assays, respectively, as detailed e.g., in Alexander *et al.*, *Immunity* 1:751-761, 1994.

For example, to confirm the capacity of a DNA minigene construct containing at least one HLA-A2 supermotif peptide to induce CTLs *in vivo*, HLA-A2.1/K^b transgenic mice, for example, are immunized intramuscularly with 100 µg of naked cDNA. As a means of comparing the level of CTLs induced by cDNA immunization, a control group of animals is also immunized with an actual peptide composition that comprises multiple epitopes synthesized as a single polypeptide as they would be encoded by the minigene.

Splenocytes from immunized animals are stimulated twice with each of the respective compositions (peptide epitopes encoded in the minigene or the polyepitopic peptide), then assayed for peptide-specific cytotoxic activity in a ⁵¹Cr release assay. The results indicate the magnitude of the CTL response directed against the A2-restricted epitope, thus indicating the *in vivo* immunogenicity of the minigene vaccine and polyepitopic vaccine.

It is, therefore, found that the minigene elicits immune responses directed toward the HLA-A2 supermotif peptide epitopes as does the polyepitopic peptide vaccine. A similar analysis is also performed using other HLA-A3 and HLA-B7 transgenic mouse models to assess CTL induction by HLA-A3 and HLA-B7 motif or supermotif epitopes, whereby it is also found that the minigene elicits appropriate immune responses directed toward the provided epitopes.

To confirm the capacity of a class II epitope-encoding minigene to induce HTLs *in vivo*, DR transgenic mice, or for those epitopes that cross react with the appropriate mouse MHC molecule, I-A^b-restricted mice, for example, are immunized intramuscularly with 100 µg of plasmid DNA. As a means of comparing the level of HTLs induced by DNA immunization, a group of control animals is also immunized with an actual peptide composition emulsified in complete Freund's adjuvant. CD4⁺ T cells, *i.e.* HTLs, are purified from splenocytes of immunized animals and stimulated with each of the respective compositions (peptides encoded in the minigene). The HTL response is measured using a ³H-thymidine incorporation proliferation assay, (see, e.g., Alexander *et al.* *Immunity* 1:751-761, 1994). The results indicate the magnitude of the HTL response, thus demonstrating the *in vivo* immunogenicity of the minigene.

DNA minigenes, constructed as described in the previous Example, can also be confirmed as a vaccine in combination with a boosting agent using a prime boost protocol. The boosting agent can consist of recombinant protein (e.g., Barnett *et al.*, *Aids Res. and Human Retroviruses* 14, Supplement 3:S299-S309, 1998) or recombinant vaccinia, for example, expressing a minigene or DNA encoding the complete protein of interest (see, e.g., Hanke *et al.*, *Vaccine* 16:439-445, 1998; Sedegah *et al.*, *Proc. Natl. Acad. Sci USA* 95:7648-53, 1998; Hanke and McMichael, *Immunol. Letters* 66:177-181, 1999; and Robinson *et al.*, *Nature Med.* 5:526-34, 1999).

For example, the efficacy of the DNA minigene used in a prime boost protocol is initially evaluated in transgenic mice. In this example, A2.1/K^b transgenic mice are immunized IM with 100 µg of a DNA minigene encoding the immunogenic peptides including at least one HLA-A2 supermotif-bearing peptide. After an incubation period (ranging from 3-9 weeks), the mice are boosted IP with 10⁷ pfu/mouse of a recombinant vaccinia virus expressing the same sequence encoded by the DNA minigene. Control mice are immunized with 100 µg of DNA or recombinant vaccinia without the minigene sequence, or with DNA encoding the minigene, but without the vaccinia boost. After an additional incubation period of two weeks, splenocytes from the mice are immediately assayed for peptide-specific activity in an ELISPOT assay.

Additionally, splenocytes are stimulated *in vitro* with the A2-restricted peptide epitopes encoded in the minigene and recombinant vaccinia, then assayed for peptide-specific activity in an alpha, beta and/or gamma IFN ELISA.

It is found that the minigene utilized in a prime-boost protocol elicits greater immune responses toward the HLA-A2 supermotif peptides than with DNA alone. Such an analysis can also be performed using HLA-A11 or HLA-B7 transgenic mouse models to assess CTL induction by HLA-A3 or HLA-B7 motif or supermotif epitopes. The use of prime boost protocols in humans is described below in the Example entitled "Induction of CTL Responses Using a Prime Boost Protocol."

Example 24: Peptide Compositions for Prophylactic Uses

Vaccine compositions of the present invention can be used to prevent 161P2F10B expression in persons who are at risk for tumors that bear this antigen. For example, a polyepitopic peptide epitope composition (or a nucleic acid comprising the same) containing multiple CTL and HTL epitopes such as those selected in the above Examples, which are also selected to target greater than 80% of the population, is administered to individuals at risk for a 161P2F10B-associated tumor.

For example, a peptide-based composition is provided as a single polypeptide that encompasses multiple epitopes. The vaccine is typically administered in a physiological solution that comprises an adjuvant, such as Incomplete Freund's Adjuvant. The dose of peptide for the initial immunization is from about 1 to about 50,000 µg, generally 100-5,000 µg, for a 70 kg patient. The initial administration of vaccine is followed by booster dosages at 4 weeks followed by evaluation of the magnitude of the immune response in the patient, by techniques that determine the presence of epitope-specific CTL populations in a PBMC sample. Additional booster doses are administered as required. The composition is found to be both safe and efficacious as a prophylaxis against 161P2F10B-associated disease.

Alternatively, a composition typically comprising transfecting agents is used for the administration of a nucleic acid-based vaccine in accordance with methodologies known in the art and disclosed herein.

Example 25: Polyepitopic Vaccine Compositions Derived from Native 161P2F10B Sequences

A native 161P2F10B polyprotein sequence is analyzed, preferably using computer algorithms defined for each class I and/or class II supermotif or motif, to identify "relatively short" regions of the polyprotein that comprise multiple epitopes. The "relatively short" regions are preferably less in length than an entire native antigen. This relatively short sequence that contains multiple distinct or overlapping, "nested" epitopes can be used to generate a minigene construct. The construct is engineered to express the peptide, which corresponds to the native protein sequence. The "relatively short" peptide is generally less than 250 amino acids in length, often less than 100 amino acids in length, preferably less than 75 amino acids in length, and more preferably less than 50 amino acids in length. The protein sequence of the vaccine composition is selected because it has maximal number of epitopes contained within the sequence, *i.e.*, it has a high concentration of epitopes. As noted herein, epitope motifs may be nested or overlapping (*i.e.*, frame shifted relative to one another). For example, with overlapping epitopes, two 9-mer epitopes and one 10-mer epitope can be present in a 10 amino acid peptide. Such a vaccine composition is administered for therapeutic or prophylactic purposes.

The vaccine composition will include, for example, multiple CTL epitopes from 161P2F10B antigen and at least one HTL epitope. This polyepitopic native sequence is administered either as a peptide or as a nucleic acid sequence which encodes the peptide. Alternatively, an analog can be made of this native sequence, whereby one or more of the epitopes comprise substitutions that alter the cross-reactivity and/or binding affinity properties of the polyepitopic peptide.

The embodiment of this example provides for the possibility that an as yet undiscovered aspect of immune system processing will apply to the native nested sequence and thereby facilitate the production of therapeutic or prophylactic immune response-inducing vaccine compositions. Additionally, such an embodiment provides for the possibility of motif-

bearing epitopes for an HLA makeup(s) that is presently unknown. Furthermore, this embodiment (excluding an analoged embodiment) directs the immune response to multiple peptide sequences that are actually present in native 161P2F10B, thus avoiding the need to evaluate any junctional epitopes. Lastly, the embodiment provides an economy of scale when producing peptide or nucleic acid vaccine compositions.

Related to this embodiment, computer programs are available in the art which can be used to identify in a target sequence, the greatest number of epitopes per sequence length.

Example 26: Polyepitopic Vaccine Compositions from Multiple Antigens

The 161P2F10B peptide epitopes of the present invention are used in conjunction with epitopes from other target tumor-associated antigens, to create a vaccine composition that is useful for the prevention or treatment of cancer that expresses 161P2F10B and such other antigens. For example, a vaccine composition can be provided as a single polypeptide that incorporates multiple epitopes from 161P2F10B as well as tumor-associated antigens that are often expressed with a target cancer associated with 161P2F10B expression, or can be administered as a composition comprising a cocktail of one or more discrete epitopes. Alternatively, the vaccine can be administered as a minigene construct or as dendritic cells which have been loaded with the peptide epitopes *in vitro*.

Example 27: Use of peptides to evaluate an immune response

Peptides of the invention may be used to analyze an immune response for the presence of specific antibodies, CTL or HTL directed to 161P2F10B. Such an analysis can be performed in a manner described by Ogg *et al.*, *Science* 279:2103-2106, 1998. In this Example, peptides in accordance with the invention are used as a reagent for diagnostic or prognostic purposes, not as an immunogen.

In this example highly sensitive human leukocyte antigen tetrameric complexes ("tetramers") are used for a cross-sectional analysis of, for example, 161P2F10B HLA-A*0201-specific CTL frequencies from HLA A*0201-positive individuals at different stages of disease or following immunization comprising a 161P2F10B peptide containing an A*0201 motif. Tetrameric complexes are synthesized as described (Musey *et al.*, *N. Engl. J. Med.* 337:1267, 1997). Briefly, purified HLA heavy chain (A*0201 in this example) and β 2-microglobulin are synthesized by means of a prokaryotic expression system. The heavy chain is modified by deletion of the transmembrane-cytosolic tail and COOH-terminal addition of a sequence containing a BirA enzymatic biotinylation site. The heavy chain, β 2-microglobulin, and peptide are refolded by dilution. The 45-kD refolded product is isolated by fast protein liquid chromatography and then biotinylated by BirA in the presence of biotin (Sigma, St. Louis, Missouri), adenosine 5' triphosphate and magnesium. Streptavidin-phycoerythrin conjugate is added in a 1:4 molar ratio, and the tetrameric product is concentrated to 1 mg/ml. The resulting product is referred to as tetramer-phycoerythrin.

For the analysis of patient blood samples, approximately one million PBMCs are centrifuged at 300g for 5 minutes and resuspended in 50 μ l of cold phosphate-buffered saline. Tri-color analysis is performed with the tetramer-phycoerythrin, along with anti-CD8-Tricolor, and anti-CD38. The PBMCs are incubated with tetramer and antibodies on ice for 30 to 60 min and then washed twice before formaldehyde fixation. Gates are applied to contain >99.98% of control samples. Controls for the tetramers include both A*0201-negative individuals and A*0201-positive non-diseased donors. The percentage of cells stained with the tetramer is then determined by flow cytometry. The results indicate the number of cells in the PBMC sample that contain epitope-restricted CTLs, thereby readily indicating the extent of immune response to the 161P2F10B epitope, and thus the status of exposure to 161P2F10B, or exposure to a vaccine that elicits a protective or therapeutic response.

Example 28: Use of Peptide Epitopes to Evaluate Recall Responses

The peptide epitopes of the invention are used as reagents to evaluate T cell responses, such as acute or recall responses, in patients. Such an analysis may be performed on patients who have recovered from 161P2F10B-associated disease or who have been vaccinated with a 161P2F10B vaccine.

For example, the class I restricted CTL response of persons who have been vaccinated may be analyzed. The vaccine may be any 161P2F10B vaccine. PBMC are collected from vaccinated individuals and HLA typed. Appropriate peptide epitopes of the invention that, optimally, bear supermotifs to provide cross-reactivity with multiple HLA supertype family members, are then used for analysis of samples derived from individuals who bear that HLA type.

PBMC from vaccinated individuals are separated on Ficoll-Histopaque density gradients (Sigma Chemical Co., St. Louis, MO), washed three times in HBSS (GIBCO Laboratories), resuspended in RPMI-1640 (GIBCO Laboratories) supplemented with L-glutamine (2mM), penicillin (50U/ml), streptomycin (50 µg/ml), and Hepes (10mM) containing 10% heat-inactivated human AB serum (complete RPMI) and plated using microculture formats. A synthetic peptide comprising an epitope of the invention is added at 10 µg/ml to each well and HBV core 128-140 epitope is added at 1 µg/ml to each well as a source of T cell help during the first week of stimulation.

In the microculture format, 4×10^5 PBMC are stimulated with peptide in 8 replicate cultures in 96-well round bottom plate in 100 µl/well of complete RPMI. On days 3 and 10, 100 µl of complete RPMI and 20 U/ml final concentration of rIL-2 are added to each well. On day 7 the cultures are transferred into a 96-well flat-bottom plate and restimulated with peptide, rIL-2 and 10^5 irradiated (3,000 rad) autologous feeder cells. The cultures are tested for cytotoxic activity on day 14. A positive CTL response requires two or more of the eight replicate cultures to display greater than 10% specific ^{51}Cr release, based on comparison with non-diseased control subjects as previously described (Rehermann, *et al.*, *Nature Med.* 2:1104,1108, 1996; Rehermann *et al.*, *J. Clin. Invest.* 97:1655-1665, 1996; and Rehermann *et al.* *J. Clin. Invest.* 98:1432-1440, 1996).

Target cell lines are autologous and allogeneic EBV-transformed B-LCL that are either purchased from the American Society for Histocompatibility and Immunogenetics (ASHI, Boston, MA) or established from the pool of patients as described (Guilhot, *et al. J. Virol.* 66:2670-2678, 1992).

Cytotoxicity assays are performed in the following manner. Target cells consist of either allogeneic HLA-matched or autologous EBV-transformed B lymphoblastoid cell line that are incubated overnight with the synthetic peptide epitope of the invention at 10 µM, and labeled with 100 µCi of ^{51}Cr (Amersham Corp., Arlington Heights, IL) for 1 hour after which they are washed four times with HBSS.

Cytolytic activity is determined in a standard 4-h, split well ^{51}Cr release assay using U-bottomed 96 well plates containing 3,000 targets/well. Stimulated PBMC are tested at effector/target (E/T) ratios of 20-50:1 on day 14. Percent cytotoxicity is determined from the formula: $100 \times [(\text{experimental release} - \text{spontaneous release}) / (\text{maximum release} - \text{spontaneous release})]$. Maximum release is determined by lysis of targets by detergent (2% Triton X-100; Sigma Chemical Co., St. Louis, MO). Spontaneous release is <25% of maximum release for all experiments.

The results of such an analysis indicate the extent to which HLA-restricted CTL populations have been stimulated by previous exposure to 161P2F10B or a 161P2F10B vaccine.

Similarly, Class II restricted HTL responses may also be analyzed. Purified PBMC are cultured in a 96-well flat bottom plate at a density of 1.5×10^5 cells/well and are stimulated with 10 µg/ml synthetic peptide of the invention, whole 161P2F10B antigen, or PHA. Cells are routinely plated in replicates of 4-6 wells for each condition. After seven days of culture, the medium is removed and replaced with fresh medium containing 10U/ml IL-2. Two days later, 1 µCi ^3H -thymidine is added to each well and incubation is continued for an additional 18 hours. Cellular DNA is then harvested on glass fiber mats and analyzed for ^3H -thymidine incorporation. Antigen-specific T cell proliferation is calculated as the ratio of ^3H -thymidine incorporation in the presence of antigen divided by the ^3H -thymidine incorporation in the absence of antigen.

Example 29: Induction Of Specific CTL Response In Humans

A human clinical trial for an immunogenic composition comprising CTL and HTL epitopes of the invention is set up as an IND Phase I, dose escalation study and carried out as a randomized, double-blind, placebo-controlled trial. Such a trial is designed, for example, as follows:

A total of about 27 individuals are enrolled and divided into 3 groups:

Group I: 3 subjects are injected with placebo and 6 subjects are injected with 5 µg of peptide composition;

Group II: 3 subjects are injected with placebo and 6 subjects are injected with 50 µg peptide composition;

Group III: 3 subjects are injected with placebo and 6 subjects are injected with 500 µg of peptide composition.

After 4 weeks following the first injection, all subjects receive a booster inoculation at the same dosage.

The endpoints measured in this study relate to the safety and tolerability of the peptide composition as well as its immunogenicity. Cellular immune responses to the peptide composition are an index of the intrinsic activity of this the peptide composition, and can therefore be viewed as a measure of biological efficacy. The following summarize the clinical and laboratory data that relate to safety and efficacy endpoints.

Safety: The incidence of adverse events is monitored in the placebo and drug treatment group and assessed in terms of degree and reversibility.

Evaluation of Vaccine Efficacy: For evaluation of vaccine efficacy, subjects are bled before and after injection. Peripheral blood mononuclear cells are isolated from fresh heparinized blood by Ficoll-Hypaque density gradient centrifugation, aliquoted in freezing media and stored frozen. Samples are assayed for CTL and HTL activity.

The vaccine is found to be both safe and efficacious.

Example 30: Phase II Trials In Patients Expressing 161P2F10B

Phase II trials are performed to study the effect of administering the CTL-HTL peptide compositions to patients having cancer that expresses 161P2F10B. The main objectives of the trial are to determine an effective dose and regimen for inducing CTLs in cancer patients that express 161P2F10B, to establish the safety of inducing a CTL and HTL response in these patients, and to see to what extent activation of CTLs improves the clinical picture of these patients, as manifested, e.g., by the reduction and/or shrinking of lesions. Such a study is designed, for example, as follows:

The studies are performed in multiple centers. The trial design is an open-label, uncontrolled, dose escalation protocol wherein the peptide composition is administered as a single dose followed six weeks later by a single booster shot of the same dose. The dosages are 50, 500 and 5,000 micrograms per injection. Drug-associated adverse effects (severity and reversibility) are recorded.

There are three patient groupings. The first group is injected with 50 micrograms of the peptide composition and the second and third groups with 500 and 5,000 micrograms of peptide composition, respectively. The patients within each group range in age from 21-65 and represent diverse ethnic backgrounds. All of them have a tumor that expresses 161P2F10B.

Clinical manifestations or antigen-specific T-cell responses are monitored to assess the effects of administering the peptide compositions. The vaccine composition is found to be both safe and efficacious in the treatment of 161P2F10B-associated disease.

Example 31: Induction of CTL Responses Using a Prime Boost Protocol

A prime boost protocol similar in its underlying principle to that used to confirm the efficacy of a DNA vaccine in transgenic mice, such as described above in the Example entitled "The Plasmid Construct and the Degree to Which It

Induces Immunogenicity," can also be used for the administration of the vaccine to humans. Such a vaccine regimen can include an initial administration of, for example, naked DNA followed by a boost using recombinant virus encoding the vaccine, or recombinant protein/polypeptide or a peptide mixture administered in an adjuvant.

For example, the initial immunization may be performed using an expression vector, such as that constructed in the Example entitled "Construction of "Minigene" Multi-Epitope DNA Plasmids" in the form of naked nucleic acid administered IM (or SC or ID) in the amounts of 0.5-5 mg at multiple sites. The nucleic acid (0.1 to 1000 μ g) can also be administered using a gene gun. Following an incubation period of 3-4 weeks, a booster dose is then administered. The booster can be recombinant fowlpox virus administered at a dose of $5 \cdot 10^7$ to $5 \cdot 10^9$ pfu. An alternative recombinant virus, such as an MVA, canarypox, adenovirus, or adeno-associated virus, can also be used for the booster, or the polyepitopic protein or a mixture of the peptides can be administered. For evaluation of vaccine efficacy, patient blood samples are obtained before immunization as well as at intervals following administration of the initial vaccine and booster doses of the vaccine. Peripheral blood mononuclear cells are isolated from fresh heparinized blood by Ficoll-Hypaque density gradient centrifugation, aliquoted in freezing media and stored frozen. Samples are assayed for CTL and HTL activity.

Analysis of the results indicates that a magnitude of response sufficient to achieve a therapeutic or protective immunity against 161P2F10B is generated.

Example 32: Administration of Vaccine Compositions Using Dendritic Cells (DC)

Vaccines comprising peptide epitopes of the invention can be administered using APCs, or "professional" APCs such as DC. In this example, peptide-pulsed DC are administered to a patient to stimulate a CTL response *in vivo*. In this method, dendritic cells are isolated, expanded, and pulsed with a vaccine comprising peptide CTL and HTL epitopes of the invention. The dendritic cells are infused back into the patient to elicit CTL and HTL responses *in vivo*. The induced CTL and HTL then destroy or facilitate destruction, respectively, of the target cells that bear the 161P2F10B protein from which the epitopes in the vaccine are derived.

For example, a cocktail of epitope-comprising peptides is administered *ex vivo* to PBMC, or isolated DC therefrom. A pharmaceutical to facilitate harvesting of DC can be used, such as Progenipoiectin™ (Monsanto, St. Louis, MO) or GM-CSF/IL-4. After pulsing the DC with peptides, and prior to reinfusion into patients, the DC are washed to remove unbound peptides.

As appreciated clinically, and readily determined by one of skill based on clinical outcomes, the number of DC reinfused into the patient can vary (see, e.g., *Nature Med.* 4:328, 1998; *Nature Med.* 2:52, 1996 and *Prostate* 32:272, 1997). Although $2 \cdot 50 \times 10^6$ DC per patient are typically administered, larger number of DC, such as 10^7 or 10^8 can also be provided. Such cell populations typically contain between 50-90% DC.

In some embodiments, peptide-loaded PBMC are injected into patients without purification of the DC. For example, PBMC generated after treatment with an agent such as Progenipoiectin™ are injected into patients without purification of the DC. The total number of PBMC that are administered often ranges from 10^8 to 10^{10} . Generally, the cell doses injected into patients is based on the percentage of DC in the blood of each patient, as determined, for example, by immunofluorescence analysis with specific anti-DC antibodies. Thus, for example, if Progenipoiectin™ mobilizes 2% DC in the peripheral blood of a given patient, and that patient is to receive 5×10^6 DC, then the patient will be injected with a total of 2.5×10^8 peptide-loaded PBMC. The percent DC mobilized by an agent such as Progenipoiectin™ is typically estimated to be between 2-10%, but can vary as appreciated by one of skill in the art.

***Ex vivo* activation of CTL/HTL responses**

Alternatively, *ex vivo* CTL or HTL responses to 161P2F10B antigens can be induced by incubating, in tissue culture, the patient's, or genetically compatible, CTL or HTL precursor cells together with a source of APC, such as DC, and

immunogenic peptides. After an appropriate incubation time (typically about 7-28 days), in which the precursor cells are activated and expanded into effector cells, the cells are infused into the patient, where they will destroy (CTL) or facilitate destruction (HTL) of their specific target cells, *i.e.*, tumor cells.

Example 33: An Alternative Method of Identifying and Confirming Motif-Bearing Peptides

Another method of identifying and confirming motif-bearing peptides is to elute them from cells bearing defined MHC molecules. For example, EBV transformed B cell lines used for tissue typing have been extensively characterized to determine which HLA molecules they express. In certain cases these cells express only a single type of HLA molecule. These cells can be transfected with nucleic acids that express the antigen of interest, *e.g.* 161P2F10B. Peptides produced by endogenous antigen processing of peptides produced as a result of transfection will then bind to HLA molecules within the cell and be transported and displayed on the cell's surface. Peptides are then eluted from the HLA molecules by exposure to mild acid conditions and their amino acid sequence determined, *e.g.*, by mass spectral analysis (*e.g.*, Kubo *et al.*, *J. Immunol.* 152:3913, 1994). Because the majority of peptides that bind a particular HLA molecule are motif-bearing, this is an alternative modality for obtaining the motif-bearing peptides correlated with the particular HLA molecule expressed on the cell.

Alternatively, cell lines that do not express endogenous HLA molecules can be transfected with an expression construct encoding a single HLA allele. These cells can then be used as described, *i.e.*, they can then be transfected with nucleic acids that encode 161P2F10B to isolate peptides corresponding to 161P2F10B that have been presented on the cell surface. Peptides obtained from such an analysis will bear motif(s) that correspond to binding to the single HLA allele that is expressed in the cell.

As appreciated by one in the art, one can perform a similar analysis on a cell bearing more than one HLA allele and subsequently determine peptides specific for each HLA allele expressed. Moreover, one of skill would also recognize that means other than transfection, such as loading with a protein antigen, can be used to provide a source of antigen to the cell.

Example 34: Complementary Polynucleotides

Sequences complementary to the 161P2F10B-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring 161P2F10B. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using, *e.g.*, OLIGO 4.06 software (National Biosciences) and the coding sequence of 161P2F10B. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to a 161P2F10B-encoding transcript.

Example 35: Purification of Naturally-occurring or Recombinant 161P2F10B Using 161P2F10B-Specific Antibodies

Naturally occurring or recombinant 161P2F10B is substantially purified by immunoaffinity chromatography using antibodies specific for 161P2F10B. An immunoaffinity column is constructed by covalently coupling anti-161P2F10B antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing 161P2F10B are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of 161P2F10B (*e.g.*, high ionic strength buffers in the presence of

detergent). The column is eluted under conditions that disrupt antibody/161P2F10B binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GCR.P is collected.

Example 36: Identification of Molecules Which Interact with 161P2F10B

161P2F10B, or biologically active fragments thereof, are labeled with 121 I Bolton-Hunter reagent. (See, e.g., Bolton *et al.* (1973) *Biochem. J.* 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled 161P2F10B, washed, and any wells with labeled 161P2F10B complex are assayed. Data obtained using different concentrations of 161P2F10B are used to calculate values for the number, affinity, and association of 161P2F10B with the candidate molecules.

Example 37: *In Vivo* Assay for 161P2F10B Tumor Growth Promotion

The effect of the 161P2F10B protein on tumor cell growth is evaluated *in vivo* by gene overexpression in tumor-bearing mice. For example, SCID mice are injected subcutaneously on each flank with 1×10^6 of either PC3, TSUPR1, or DU145 cells containing tkNeo empty vector or 161P2F10B. At least two strategies may be used: (1) Constitutive 161P2F10B expression under regulation of a promoter such as a constitutive promoter obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), or from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, provided such promoters are compatible with the host cell systems, and (2) regulated expression under control of an inducible vector system, such as ecdysone, tet, etc., provided such promoters are compatible with the host cell systems. Tumor volume is then monitored at the appearance of palpable tumors and followed over time and determines that 161P2F10B-expressing cells grow at a faster rate and/or tumors produced by 161P2F10B-expressing cells demonstrate characteristics of altered aggressiveness (e.g. enhanced metastasis, vascularization, reduced responsiveness to chemotherapeutic drugs).

Additionally, mice can be implanted with 1×10^5 of the same cells orthotopically to determine that 161P2F10B has an effect on local growth in the prostate and/or on the ability of the cells to metastasize, e.g., to lungs, lymph nodes, and bone marrow.

The assay is also useful to determine the 161P2F10B-inhibitory effect of candidate therapeutic compositions, such as for example, small molecule drugs, 161P2F10B intrabodies, 161P2F10B antisense molecules and ribozymes.

Example 38: 161P2F10B Monoclonal Antibody-mediated Inhibition of Prostate Tumors *In Vivo*.

The significant expression of 161P2F10B, in cancer tissues, together with its restricted expression in normal tissues along with its cell surface expression makes 161P2F10B an excellent target for antibody therapy. Similarly, 161P2F10B is a target for T cell-based immunotherapy. Thus, the therapeutic efficacy of anti-161P2F10B mAbs is evaluated, e.g., in human prostate cancer xenograft mouse models using androgen-independent LAPC-4 and LAPC-9 xenografts (Craft, N., *et al.*, *Cancer Res.* 1999. 59(19): p. 5030-6), kidney cancer xenografts (AGS-K3, AGS-K6), kidney cancer metastases to lymph node (AGS-K6 met) xenografts, and kidney cancer cell lines transfected with 161P2F10B, such as 769P-161P2F10B, A498-161P2F10B.

Antibody efficacy on tumor growth and metastasis formation is studied, e.g., in mouse orthotopic prostate cancer xenograft models and mouse kidney xenograft models. The antibodies can be unconjugated, as discussed in this Example, or can be conjugated to a therapeutic modality, as appreciated in the art. Anti-161P2F10B mAbs inhibit formation of both the androgen-dependent LAPC-9 and androgen-independent PC3-161P2F10B tumor xenografts. Anti-161P2F10B mAbs also retard the growth of established orthotopic tumors and prolonged survival of tumor-bearing mice. These results indicate the

utility of anti-161P2F10B mAbs in the treatment of local and advanced stages of prostate cancer. (See, e.g., Saffran, D., et al., PNAS 10:1073-1078 or www.pnas.org/cgi/doi/10.1073/pnas.051624698). Similarly, anti-161P2F10B mAbs can inhibit formation of AGS-K3 and AGS-K6 tumors in SCID mice, and prevent or retard the growth A498-161P2F10B tumor xenografts. These results indicate the use of anti-161P2F10B mAbs in the treatment of prostate and/or kidney cancer.

Administration of the anti-161P2F10B mAbs leads to retardation of established orthotopic tumor growth and inhibition of metastasis to distant sites, resulting in a significant prolongation in the survival of tumor-bearing mice. These studies indicate that 161P2F10B is an attractive target for immunotherapy and demonstrate the therapeutic use of anti-161P2F10B mAbs for the treatment of local and metastatic prostate cancer. This example demonstrates that unconjugated 161P2F10B monoclonal antibodies are effective to inhibit the growth of human prostate tumor xenografts and human kidney xenografts grown in SCID mice.

Tumor inhibition using multiple unconjugated 161P2F10B mAbs

Materials and Methods

161P2F10B Monoclonal Antibodies:

Monoclonal antibodies are raised against 161P2F10B as described in the Example entitled "Generation of 161P2F10B Monoclonal Antibodies (mAbs)" or are obtained commercially, e.g., 97A6 (Coulter Immunotech). The antibodies are characterized by ELISA, Western blot, FACS, and immunoprecipitation for their capacity to bind 161P2F10B. Epitope mapping data for the anti-161P2F10B mAbs, as determined by ELISA and Western analysis, recognize epitopes on the 161P2F10B protein. The 97A6 antibody binds to amino acids 393-405 of the 161P2F10B protein shown in Figure 2. Immunohistochemical analysis of cancer tissues and cells is performed with these antibodies.

The monoclonal antibodies are purified from ascites or hybridoma tissue culture supernatants by Protein-G Sepharose chromatography, dialyzed against PBS, filter sterilized, and stored at -20°C. Protein determinations are performed by a Bradford assay (Bio-Rad, Hercules, CA). A therapeutic monoclonal antibody or a cocktail comprising a mixture of individual monoclonal antibodies is prepared and used for the treatment of mice receiving subcutaneous or orthotopic injections of LAPC-9 prostate tumor xenografts.

Cancer Xenografts and Cell Lines

The LAPC-9 xenograft, which expresses a wild-type androgen receptor and produces prostate-specific antigen (PSA), is passaged in 6- to 8-week-old male ICR-severe combined immunodeficient (SCID) mice (Taconic Farms) by s.c. trocar implant (Craft, N., *et al.*, *supra*). The AGS-K3 and AGS-K6 kidney xenografts are also passaged by subcutaneous implants in 6- to 8- week old SCID mice. Single-cell suspensions of tumor cells are prepared as described in Craft, *et al.* The prostate carcinoma cell line PC3 (American Type Culture Collection) is maintained in RPMI supplemented with L-glutamine and 10% FBS, and the kidney carcinoma line A498 (American Type Culture Collection) is maintained in DMEM supplemented with L-glutamine and 10% FBS.

PC3-161P2F10B and A498-161P2F10B cell populations are generated by retroviral gene transfer as described in Hubert, R.S., et al., STEAP: A Prostate-specific Cell-surface Antigen Highly Expressed in Human Prostate Tumors, *Proc Natl Acad Sci U S A*, 1999, 96(25): p. 14523-8. Anti-161P2F10B staining is detected by using an FITC-conjugated goat anti-mouse antibody (Southern Biotechnology Associates) followed by analysis on a Coulter Epics-XL flow cytometer.

Xenograft Mouse Models.

Subcutaneous (s.c.) tumors are generated by injection of 1×10^6 LAPC-9, AGS-K3, AGS-K6, PC3, PC3-161P2F10B, A498 or A498-161P2F10B cells mixed at a 1:1 dilution with Matrigel (Collaborative Research) in the right flank of male SCID mice. To test antibody efficacy on tumor formation, i.p. antibody injections are started on the same day as tumor-cell injections. As a control, mice are injected with either purified mouse IgG (ICN) or PBS; or a purified monoclonal

antibody that recognizes an irrelevant antigen not expressed in human cells. In preliminary studies, no difference is found between mouse IgG or PBS on tumor growth. Tumor sizes are determined by vernier caliper measurements, and the tumor volume is calculated as length x width x height. Mice with s.c. tumors greater than 1.5 cm in diameter are sacrificed. PSA levels are determined by using a PSA ELISA kit (Anogen, Mississauga, Ontario). Circulating levels of anti-161P2F10B mAbs are determined by a capture ELISA kit (Bethyl Laboratories, Montgomery, TX). (See, e.g., (Saffran, D., et al., PNAS 10:1073-1078 or www.pnas.org/cgi/doi/10.1073/pnas.051624698)

Orthotopic prostate injections are performed under anesthesia by using ketamine/xylazine. For prostate orthotopic studies, an incision is made through the abdominal muscles to expose the bladder and seminal vesicles, which then are delivered through the incision to expose the dorsal prostate. LAPC-9 cells (5×10^5) mixed with Matrigel are injected into each dorsal lobe in a 10- μ l volume. To monitor tumor growth, mice are bled on a weekly basis for determination of PSA levels. For kidney orthotopic models, an incision is made through the abdominal muscles to expose the kidney. AGS-K3 or AGS-K6 cells mixed with Matrigel are injected under the kidney capsule. The mice are segregated into groups for the appropriate treatments, with anti-161P2F10B or control mAbs being injected i.p.

Anti-161P2F10B mAbs Inhibit Growth of 161P2F10B-Expressing Xenograft-Cancer Tumors

The effect of anti-161P2F10B mAbs on tumor formation is tested by using LAPC-9 and/or AGS-K3 orthotopic models. As compared with the s.c. tumor model, the orthotopic model, which requires injection of tumor cells directly in the mouse prostate or kidney, respectively, results in a local tumor growth, development of metastasis in distal sites, deterioration of mouse health, and subsequent death (Saffran, D., et al., PNAS supra; Fu, X., et al., Int J Cancer, 1992. 52(6): p. 987-90; Kubota, T., J Cell Biochem, 1994. 56(1): p. 4-8). The features make the orthotopic model more representative of human disease progression and allowed for tracking of the therapeutic effect of mAbs on clinically relevant end points.

Accordingly, tumor cells are injected into the mouse prostate or kidney, and 2 days later, the mice are segregated into two groups and treated with either: a) 200-500 μ g, of anti-161P2F10B Ab, or b) PBS three times per week for two to five weeks.

A major advantage of the orthotopic prostate-cancer model is the ability to study the development of metastases. Formation of metastasis in mice bearing established orthotopic tumors is studied by IHC analysis on lung sections using an antibody against a prostate-specific cell-surface protein STEAP expressed at high levels in LAPC-9 xenografts (Hubert, R.S., et al., Proc Natl Acad Sci U S A, 1999. 96(25): p. 14523-8) or anti-G250 antibody for kidney cancer models.

Mice bearing established orthotopic LAPC-9 tumors are administered 1000 μ g injections of either anti-161P2F10B mAb or PBS over a 4-week period. Mice in both groups are allowed to establish a high tumor burden (PSA levels greater than 300 ng/ml), to ensure a high frequency of metastasis formation in mouse lungs. Mice then are killed and their prostate/kidney and lungs are analyzed for the presence of tumor cells by IHC analysis.

These studies demonstrate a broad anti-tumor efficacy of anti-161P2F10B antibodies on initiation and progression of prostate and kidney cancer in xenograft mouse models. Anti-161P2F10B antibodies inhibit tumor formation of both androgen-dependent and androgen-independent prostate tumors as well as retarding the growth of already established tumors and prolong the survival of treated mice. Moreover, anti-161P2F10B mAbs demonstrate a dramatic inhibitory effect on the spread of local prostate tumor to distal sites, even in the presence of a large tumor burden. Similar therapeutic effects are seen in the kidney cancer model. Thus, anti-161P2F10B mAbs are efficacious on major clinically relevant end points (tumor growth), prolongation of survival, and health.

Example 39: Therapeutic and Diagnostic use of Anti-161P2F10B Antibodies in Humans.

Anti-161P2F10B monoclonal antibodies are safely and effectively used for diagnostic, prophylactic, prognostic and/or therapeutic purposes in humans. Western blot and immunohistochemical analysis of cancer tissues and cancer xenografts with anti-161P2F10B mAb show strong extensive staining in carcinoma but significantly lower or undetectable levels in normal tissues. Detection of 161P2F10B in carcinoma and in metastatic disease demonstrates the usefulness of the mAb as a diagnostic and/or prognostic indicator. Anti-161P2F10B antibodies are therefore used in diagnostic applications such as immunohistochemistry of kidney biopsy specimens to detect cancer from suspect patients.

As determined by flow cytometry, anti-161P2F10B mAb specifically binds to carcinoma cells. Thus, anti-161P2F10B antibodies are used in diagnostic whole body imaging applications, such as radioimmunoscinigraphy and radioimmunotherapy, (see, e.g., Potamianos S., et. al. *Anticancer Res* 20(2A):925-948 (2000)) for the detection of localized and metastatic cancers that exhibit expression of 161P2F10B. Shedding or release of an extracellular domain of 161P2F10B into the extracellular milieu, such as that seen for alkaline phosphodiesterase B10 (Meerson, N. R., *Hepatology* 27:563-568 (1998)), allows diagnostic detection of 161P2F10B by anti-161P2F10B antibodies in serum and/or urine samples from suspect patients.

Anti-161P2F10B antibodies that specifically bind 161P2F10B are used in therapeutic applications for the treatment of cancers that express 161P2F10B. Anti-161P2F10B antibodies are used as an unconjugated modality and as conjugated form in which the antibodies are attached to one of various therapeutic or imaging modalities well known in the art, such as a prodrugs, enzymes or radioisotopes. In preclinical studies, unconjugated and conjugated anti-161P2F10B antibodies are tested for efficacy of tumor prevention and growth inhibition in the SCID mouse cancer xenograft models, e.g., kidney cancer models AGS-K3 and AGS-K6, (see, e.g., the Example entitled "161P2F10B Monoclonal Antibody-mediated Inhibition of Bladder and Lung Tumors *In Vivo*"). Either conjugated and unconjugated anti-161P2F10B antibodies are used as a therapeutic modality in human clinical trials either alone or in combination with other treatments as described in following Examples.

Example 40: Human Clinical Trials for the Treatment and Diagnosis of Human Carcinomas through use of Human Anti-161P2F10B Antibodies *In vivo*

Antibodies are used in accordance with the present invention which recognize an epitope on 161P2F10B, and are used in the treatment of certain tumors such as those listed in Table I. Based upon a number of factors, including 161P2F10B expression levels, tumors such as those listed in Table I are presently preferred indications. In connection with each of these indications, three clinical approaches are successfully pursued.

I.) **Adjunctive therapy:** In adjunctive therapy, patients are treated with anti-161P2F10B antibodies in combination with a chemotherapeutic or antineoplastic agent and/or radiation therapy. Primary cancer targets, such as those listed in Table I, are treated under standard protocols by the addition anti-161P2F10B antibodies to standard first and second line therapy. Protocol designs address effectiveness as assessed by reduction in tumor mass as well as the ability to reduce usual doses of standard chemotherapy. These dosage reductions allow additional and/or prolonged therapy by reducing dose-related toxicity of the chemotherapeutic agent. Anti-161P2F10B antibodies are utilized in several adjunctive clinical trials in combination with the chemotherapeutic or antineoplastic agents adriamycin (advanced prostate carcinoma), cisplatin (advanced head and neck and lung carcinomas), taxol (breast cancer), and doxorubicin (preclinical).

II.) **Monotherapy:** In connection with the use of the anti-161P2F10B antibodies in monotherapy of tumors, the antibodies are administered to patients without a chemotherapeutic or antineoplastic agent. In one embodiment,

monotherapy is conducted clinically in end stage cancer patients with extensive metastatic disease. Patients show some disease stabilization. Trials demonstrate an effect in refractory patients with cancerous tumors.

III.) Imaging Agent: Through binding a radionuclide (e.g., iodine or yttrium (I^{131} , Y^{90}) to anti-161P2F10B antibodies, the radiolabeled antibodies are utilized as a diagnostic and/or imaging agent. In such a role, the labeled antibodies localize to both solid tumors, as well as, metastatic lesions of cells expressing 161P2F10B. In connection with the use of the anti-161P2F10B antibodies as imaging agents, the antibodies are used as an adjunct to surgical treatment of solid tumors, as both a pre-surgical screen as well as a post-operative follow-up to determine what tumor remains and/or returns. In one embodiment, a (^{111}In)-161P2F10B antibody is used as an imaging agent in a Phase I human clinical trial in patients having a carcinoma that expresses 161P2F10B (by analogy see, e.g., Divgi *et al. J. Natl. Cancer Inst.* 83:97-104 (1991)). Patients are followed with standard anterior and posterior gamma camera. The results indicate that primary lesions and metastatic lesions are identified

Dose and Route of Administration

As appreciated by those of ordinary skill in the art, dosing considerations can be determined through comparison with the analogous products that are in the clinic. Thus, anti-161P2F10B antibodies can be administered with doses in the range of 5 to 400 mg/m², with the lower doses used, e.g., in connection with safety studies. The affinity of anti-161P2F10B antibodies relative to the affinity of a known antibody for its target is one parameter used by those of skill in the art for determining analogous dose regimens. Further, anti-161P2F10B antibodies that are fully human antibodies, as compared to the chimeric antibody, have slower clearance; accordingly, dosing in patients with such fully human anti-161P2F10B antibodies can be lower, perhaps in the range of 50 to 300 mg/m², and still remain efficacious. Dosing in mg/m², as opposed to the conventional measurement of dose in mg/kg, is a measurement based on surface area and is a convenient dosing measurement that is designed to include patients of all sizes from infants to adults.

Three distinct delivery approaches are useful for delivery of anti-161P2F10B antibodies. Conventional intravenous delivery is one standard delivery technique for many tumors. However, in connection with tumors in the peritoneal cavity, such as tumors of the ovaries, biliary duct, other ducts, and the like, intraperitoneal administration may prove favorable for obtaining high dose of antibody at the tumor and to also minimize antibody clearance. In a similar manner, certain solid tumors possess vasculature that is appropriate for regional perfusion. Regional perfusion allows for a high dose of antibody at the site of a tumor and minimizes short term clearance of the antibody.

Clinical Development Plan (CDP)

Overview: The CDP follows and develops treatments of anti-161P2F10B antibodies in connection with adjunctive therapy, monotherapy, and as an imaging agent. Trials initially demonstrate safety and thereafter confirm efficacy in repeat doses. Trials are open label comparing standard chemotherapy with standard therapy plus anti-161P2F10B antibodies. As will be appreciated, one criteria that can be utilized in connection with enrollment of patients is 161P2F10B expression levels in their tumors as determined by biopsy.

As with any protein or antibody infusion-based therapeutic, safety concerns are related primarily to (i) cytokine release syndrome, i.e., hypotension, fever, shaking, chills; (ii) the development of an immunogenic response to the material (i.e., development of human antibodies by the patient to the antibody therapeutic, or HAHA response); and, (iii) toxicity to normal cells that express 161P2F10B. Standard tests and follow-up are utilized to monitor each of these safety concerns. Anti-161P2F10B antibodies are found to be safe upon human administration.

Example 41: Human Clinical Trial Adjunctive Therapy with Human Anti-161P2F10B Antibody and Chemotherapeutic Agent

A phase I human clinical trial is initiated to assess the safety of six intravenous doses of a human anti-161P2F10B antibody in connection with the treatment of a solid tumor, e.g., a cancer of a tissue listed in Table I. In the study, the safety of single doses of anti-161P2F10B antibodies when utilized as an adjunctive therapy to an antineoplastic or chemotherapeutic agent as defined herein, such as, without limitation: cisplatin, topotecan, doxorubicin, adriamycin, taxol, or the like, is assessed. The trial design includes delivery of six single doses of an anti-161P2F10B antibody with dosage of antibody escalating from approximately about 25 mg/m² to about 275 mg/m² over the course of the treatment in accordance with the following schedule:

	Day 0	Day 7	Day 14	Day 21	Day 28	Day 35
mAb Dose	25	75	125	175	225	275
	mg/m ²	mg/m ²	mg/m ²	mg/m ²	mg/m ²	mg/m ²
Chemotherapy (standard dose)	+	+	+	+	+	+

Patients are closely followed for one-week following each administration of antibody and chemotherapy. In particular, patients are assessed for the safety concerns mentioned above: (i) cytokine release syndrome, i.e., hypotension, fever, shaking, chills; (ii) the development of an immunogenic response to the material (i.e., development of human antibodies by the patient to the human antibody therapeutic, or HAHA response); and, (iii) toxicity to normal cells that express 161P2F10B. Standard tests and follow-up are utilized to monitor each of these safety concerns. Patients are also assessed for clinical outcome, and particularly reduction in tumor mass as evidenced by MRI or other imaging.

The anti-161P2F10B antibodies are demonstrated to be safe and efficacious, Phase II trials confirm the efficacy and refine optimum dosing.

Example 42: Human Clinical Trial: Monotherapy with Human Anti-161P2F10B Antibody

Anti-161P2F10B antibodies are safe in connection with the above-discussed adjunctive trial, a Phase II human clinical trial confirms the efficacy and optimum dosing for monotherapy. Such trial is accomplished, and entails the same safety and outcome analyses, to the above-described adjunctive trial with the exception being that patients do not receive chemotherapy concurrently with the receipt of doses of anti-161P2F10B antibodies.

Example 43: Human Clinical Trial: Diagnostic Imaging with Anti-161P2F10B Antibody

Once again, as the adjunctive therapy discussed above is safe within the safety criteria discussed above, a human clinical trial is conducted concerning the use of anti-161P2F10B antibodies as a diagnostic imaging agent. The protocol is designed in a substantially similar manner to those described in the art, such as in Divgi *et al. J. Natl. Cancer Inst.* 83:97-104 (1991). The antibodies are found to be both safe and efficacious when used as a diagnostic modality.

Example 44: Homology Comparison of 161P2F10B to Known Sequences

The 161P2F10B gene is identical to a previously cloned and sequenced gene, namely ectonucleotide pyrophosphatase/phosphodiesterase 3 (gi 4826896) (Jin-Hua P et al, Genomics 1997, 45:412), also known as phosphodiesterase-I beta; gp130RB13-6; E-NPP3 (ENPP3), PDNP3 and CD203c. The 161P2F10B protein shows 100% identity to human ectonucleotide pyrophosphatase/phosphodiesterase 3 (gi 4826896), and 81% homology and 89% identity

to rat alkaline phosphodiesterase (gi 1699034). The 161P2F10B protein consists of 875 amino acids, with calculated molecular weight of 100.09 kDa, and pI of 6.12. 161P2F10B is a cell surface protein as shown by immunostaining in basophils (Buhning HJ et al, Blood 2001, 97:3303) and in epithelial tumor cells as shown in the example entitled "Expression of 161P2F10B protein in kidney cancer xenograft tissues". Some localization to the golgi-endoplasmic fraction has also been observed (Geoffroy V et al, Arch Biochem Biophys. 2001, 387:154). Two isoforms of phosphodiesterase 3 have been identified, with one protein containing an additional 145 aa at its amino-terminus (Choi YH et al, Biochem J. 2001, 353:41). In addition, two variants of 161P2F10B have been identified. The first variant contains a point mutation at amino acid 122 of the 161P2F10B protein, changing a lysine to an arginine at that position. The second variant contains a single nucleotide polymorphisms, identified at position 383, resulting in a change in amino acid from threonine to proline at that position see URL located on the World Wide Web at (ncbi.nlm.nih.gov/SNP/snp_ref.cgi?type=rs&rs=1054222) (figure 4A and 4B). In addition, we have recently identified another variant of 161P2F10B, namely 161P2F10B v.7. This variant differs from v.1 at its N-terminus as it lacks the first 34 aa found in v.1. The loss of the N-terminal 34 aa affects the localization of 161P2F10B v.7. PSort analysis revealed that, while 161P2F10B v.1 is primarily located at the plasma membrane, 161P2F10B v.7 primarily localizes to the cytoplasm (52%) with some localization to the nucleus (17%).

Motif analysis revealed the presence of several known motifs, including 2-3 somatostatin B domains located at the amino terminus of the 161P2F10B protein, a phosphodiesterase domain and an endonuclease domain at the C-terminus. 161P2F10B belongs to a family of closely related phosphodiesterases, consisting of PDNP1, -2, and -3 (Bollen M et al, Crit. Rev. Biochem Mol. Biol. 2000, 35: 393). All three members of this family are type II proteins, with a short N-terminus domain located intracellularly. They contain one transmembrane domain, a catalytic phosphodiesterase domain and a C-terminal nuclease domain.

Phosphodiesterase 3 expression has been detected in human neoplastic submandibular cells, glioma cells, and transformed lymphocytes (Murata T et al, Anticancer Drugs 2001, 12:79; Andoh K et al, Biochim Biophys Acta 1999, 1446:213; Ekholm D et al, Biochem Pharmacol 1999, 58: 935).

Phosphodiesterase 3 plays an important role in several biological processes, including release of nucleotides, cell differentiation, metabolism, cell growth, survival, angiogenesis and cell motility (Bollen M et al, Crit. Rev. Biochem Mol. Biol. 2000, 35: 393; Rawadi G et al, Endocrinol 2001, 142:4673; DeFouw L et al, Microvasc Res 2001, 62:263). In addition, Phosphodiesterase 3 regulates gene expression in epithelial cells, including the expression of key adhesion molecules such as VCAM-1 (Blease K et al, Br J Pharmacol. 1998, 124:229).

This information indicates that 161P2F10B plays a role in the growth of mammalian cells, supports cell survival and motility, and regulate gene transcription by regulating events in the nucleus. Accordingly, when 161P2F10B functions as a regulator of cell transformation, tumor formation, or as a modulator of transcription involved in activating genes associated with inflammation, tumorigenesis or proliferation, 161P2F10B is used for therapeutic, diagnostic, prognostic and/or preventative purposes. In addition, when a molecule, such as a variant, polymorphism or SNP of 161P2F10B is expressed in cancerous tissues, such as those listed in Table I, they are used for therapeutic, diagnostic, prognostic and/or preventative purposes.

Example 45: Regulation of Transcription

The cell surface localization of 161P2F10B and ability to regulate VCAM expression indicate that 161P2F10B is effectively used as a modulator of the transcriptional regulation of eukaryotic genes. Regulation of gene expression is confirmed, e.g., by studying gene expression in cells expressing or lacking 161P2F10B. For this purpose, two types of experiments are performed.

In the first set of experiments, RNA from parental and 161P2F10B-expressing cells are extracted and hybridized to commercially available gene arrays (Clontech) (Smid-Koopman E et al. Br J Cancer. 2000. 83:246). Resting cells as well as cells treated with FBS or androgen are compared. Differentially expressed genes are identified in accordance with procedures known in the art. The differentially expressed genes are then mapped to biological pathways (Chen K et al. Thyroid. 2001. 11:41.).

In the second set of experiments, specific transcriptional pathway activation is evaluated using commercially available (Stratagene) luciferase reporter constructs including: NFkB-luc, SRE-luc, ELK1-luc, ARE-luc, p53-luc, and CRE-luc. These transcriptional reporters contain consensus binding sites for known transcription factors that lie downstream of well-characterized signal transduction pathways, and represent a good tool to ascertain pathway activation and screen for positive and negative modulators of pathway activation.

Thus, 161P2F10B plays a role in gene regulation, and it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 46: Identification and Confirmation of Potential Signal Transduction Pathways

Many mammalian proteins have been reported to interact with signaling molecules and to participate in regulating signaling pathways. (J Neurochem. 2001; 76:217-223). In particular, GPCRs have been reported to activate MAK cascades as well as G proteins, and been associated with the EGFR pathway in epithelial cells (Naor, Z., et al, Trends Endocrinol Metab. 2000, 11:91; Vacca F et al, Cancer Res. 2000, 60:5310; Della Rocca GJ et al, J Biol Chem. 1999, 274:13978). Using immunoprecipitation and Western blotting techniques, proteins are identified that associate with 161P2F10B and mediate signaling events. Several pathways known to play a role in cancer biology can be regulated by 161P2F10B, including phospholipid pathways such as PI3K, AKT, etc, adhesion and migration pathways, including FAK, Rho, Rac-1, etc, as well as mitogenic/survival cascades such as ERK, p38, etc (Cell Growth Differ. 2000,11:279; J Biol Chem. 1999, 274:801; Oncogene. 2000, 19:3003, J. Cell Biol. 1997, 138:913.).

To confirm that 161P2F10B directly or indirectly activates known signal transduction pathways in cells, luciferase (luc) based transcriptional reporter assays are carried out in cells expressing individual genes. These transcriptional reporters contain consensus-binding sites for known transcription factors that lie downstream of well-characterized signal transduction pathways. The reporters and examples of these associated transcription factors, signal transduction pathways, and activation stimuli are listed below.

1. NFkB-luc, NFkB/Rel; Ik-kinase/SAPK; growth/apoptosis/stress
2. SRE-luc, SRF/TCF/ELK1; MAPK/SAPK; growth/differentiation
3. AP-1-luc, FOS/JUN; MAPK/SAPK/PKC; growth/apoptosis/stress
4. ARE-luc, androgen receptor; steroids/MAPK; growth/differentiation/apoptosis
5. p53-luc, p53; SAPK; growth/differentiation/apoptosis
6. CRE-luc, CREB/ATF2; PKA/p38; growth/apoptosis/stress

Gene-mediated effects can be assayed in cells showing mRNA expression. Luciferase reporter plasmids can be introduced by lipid-mediated transfection (TFX-50, Promega). Luciferase activity, an indicator of relative transcriptional activity, is measured by incubation of cell extracts with luciferin substrate and luminescence of the reaction is monitored in a luminometer.

Signaling pathways activated by 161P2F10B are mapped and used for the identification and validation of therapeutic targets. When 161P2F10B is involved in cell signaling, it is used as target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 47: Involvement in Tumor Progression

The 161P2F10B gene can contribute to the growth of cancer cells. The role of 161P2F10B in tumor growth is confirmed in a variety of primary and transfected cell lines including prostate, colon, bladder and kidney cell lines, as well as NIH 3T3 cells engineered to stably express 161P2F10B. Parental cells lacking 161P2F10B and cells expressing 161P2F10B are evaluated for cell growth using a well-documented proliferation assay (Fraser SP, Grimes JA, Djamgoz MB. Prostate. 2000;44:61, Johnson DE, Ochieng J, Evans SL. Anticancer Drugs. 1996, 7:288).

To confirm the role of 161P2F10B in the transformation process, its effect in colony forming assays is investigated. Parental NIH-3T3 cells lacking 161P2F10B are compared to NIH-3T3 cells expressing 161P2F10B, using a soft agar assay under stringent and more permissive conditions (Song Z. et al. Cancer Res. 2000;60:6730).

To confirm the role of 161P2F10B in invasion and metastasis of cancer cells, a well-established assay is used, e.g., a Transwell Insert System assay (Becton Dickinson) (Cancer Res. 1999; 59:6010). Control cells, including prostate, colon, bladder and kidney cell lines lacking 161P2F10B are compared to cells expressing 161P2F10B. Cells are loaded with the fluorescent dye, calcein, and plated in the top well of the Transwell insert coated with a basement membrane analog. Invasion is determined by fluorescence of cells in the lower chamber relative to the fluorescence of the entire cell population. Using this approach we have demonstrated that 161P2F10B induces the invasion of 3T3 cells through the basement membrane analog matrigel (Figure 22). As shown in figure 22, 3T3-neo cells that do not express 161P2F10B exhibit negligible levels of invasion through matrigel. Compared to 3T3-neo cells, 3T3-161P2F10B cells, which express abundant levels of 161P2F10B (Figure 16), invade through matrigel and migrate to the lower chamber of the transwell system in a manner similar to that observed with cells expressing the strong protooncogene 12V-Ras.

161P2F10B can also play a role in cell cycle and apoptosis. Parental cells and cells expressing 161P2F10B are compared for differences in cell cycle regulation using a well-established BrdU assay (Abdel-Malek ZA. J Cell Physiol. 1988, 136:247). In short, cells are grown under both optimal (full serum) and limiting (low serum) conditions are labeled with BrdU and stained with anti-BrdU Ab and propidium iodide. Cells are analyzed for entry into the G1, S, and G2M phases of the cell cycle.

In contrast to normal cells, cancer cells have been shown to withstand stress, growth factor deprivation and pro-apoptotic signals, thereby providing tumors with a growth and survival advantage. The effect of stress on apoptosis is evaluated in control parental cells and cells expressing 161P2F10B, including normal and tumor prostate, colon and lung cells using standard assays methods including annexin V binding and caspase activation (Moore A, et al, Methods Cell Biol. 1998;57:265; Porter AG, Janicke RU. Cell Death Differ. 1999; 6:99). Engineered and parental cells were treated with various chemotherapeutic agents, such as etoposide, doxorubicin, kinase inhibitors such as staurosporine, DNA damaging agents such as UV, hypoxia and protein synthesis inhibitors, such as cycloheximide. Cells were stained with annexin V-FITC and cell death measured by FACS analysis. Figure 20 shows that expression of 161P2F10B prevent the apoptosis of 3T3 cells exposed to staurosporine or UV irradiation. While 64% and 62% of 3T3-neo cells underwent apoptosis in response to staurosporine and UV irradiation, respectively, only 14% and 30% of 161P2F10B-expressing 3T3 cells died under the same conditions. Similar results were obtained in another experiment comparing the effect of staurosporine and UV irradiation on 3T3-neo cells and clonal populations of 3T3-161P2F10B cell lines (Figure 19). As with the population of 3T3-161P2F10B, clones 3T3-161P2F10B-C and 3T3-161P2F10B-10B were resistant to staurosporine-induced cell death. Since caspase activation is a hallmark of apoptosis and serves to distinguish apoptosis from other forms of cell death, we investigated the

effect of to chemotherapeutic agent and, staurosporine on the apoptosis of kidney cancer cells using caspase activation as assay read out (Figure 21). The 769 kidney tumor cells that normally lack 161P2F10B were engineered to express the 161P2F10B protein as describe in example 8, Production of Recombinant 161P2F10B in Higher Eukaryotic Systems, above. The cells were treated with chemotherapeutic agents or staurosporine, lysed and analyzed for caspase activity. Figure 21 shows that expression of 161P2F10B prevents caspase activation in 161P2F10B-expressing kidney cancer cells treated with doxorubicin or staurosporine. These results show that 161P2F10B imparts resistance to the chemotherapeutic drug doxorubicin and to staurosporine-induced cell death in kidney cancer cells.

A characteristic that distinguishes cancer cells from normal cells is their ability to become serum independent and survive in low serum conditions. The effect of serum deprivation on the survival of 161P2F10B expressing cells was studied using caspase activation as a read out. The fibroblast cell line Rat-1 becomes growth arrested when serum deprived, thereby mimicking normal non-transformed cells (James L, Eisenman RN. Proc Natl Acad Sci U S A. 2002, 99:10429). Rat-1 cells expressing c-Myc (Rat-Myc) undergo apoptosis under serum deprivation conditions (James L, Eisenman RN. Proc Natl Acad Sci U S A. 2002, 99:10429). Rat-1 and Rat-Myc cells were engineered to stably express 161P2F10B. The cells were grown in 0.1% or 10% FBS and examined for apoptosis by microscopy and caspase activity (Figures 17 and 18). When 161P2F10B is stably expressed in Rat-Myc cells, it inhibits Myc-induced apoptosis and reduces caspase activity to background levels. The inhibition of cell death by 161P2F10B plays a critical role in regulating tumor progression and tumor load.

When 161P2F10B plays a role in cell growth, transformation, invasion or apoptosis, it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 48: Involvement in Angiogenesis

Angiogenesis or new capillary blood vessel formation is necessary for tumor growth (Hanahan D, Folkman J. Cell. 1996, 86:353; Folkman J. Endocrinology. 1998 139:441). Based on the effect of phosphodiesterase inhibitors on endothelial cells, and the homology of 161P2F10B to other ENPP family members, 161P2F10B plays a role in angiogenesis (DeFouw L et al, Microvasc Res 2001, 62:263). Several assays have been developed to measure angiogenesis *in vitro* and *in vivo*, such as the tissue culture assays endothelial cell tube formation and endothelial cell proliferation. Using these assays as well as *in vitro* neo-vascularization, the role of 161P2F10B in angiogenesis, enhancement or inhibition, is confirmed.

For example, endothelial cells engineered to express 161P2F10B are evaluated using tube formation and proliferation assays. The effect of 161P2F10B is also confirmed in animal models *in vivo*. For example, cells either expressing or lacking 161P2F10B are implanted subcutaneously in immunocompromised mice. Endothelial cell migration and angiogenesis are evaluated 5-15 days later using immunohistochemistry techniques. Similarly, the secreted extracellular portion of 161P2F10B can function as an angiogenic factor and enhance the proliferation and tube formation of endothelial cells. The effect of the extracellular domain of 161P2F10B on angiogenesis is supported by its similarity to other ENPPs, with biologically active secreted extracellular domain. When 161P2F10B affects angiogenesis, and it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 49: Involvement in Protein-Protein Interactions

Several phosphodiesterases have been shown to interact with other proteins, thereby regulating gene transcription, as well as cell growth (Butt E et al, Mol Pharmacol. 1995, 47:340). Using immunoprecipitation techniques as well as two yeast hybrid systems, proteins are identified that associate with 161P2F10B. Immunoprecipitates from cells expressing 161P2F10B and cells lacking 161P2F10B are compared for specific protein-protein associations.

Studies are performed to confirm the extent of association of 161P2F10B with effector molecules, such as nuclear proteins, transcription factors, kinases, phosphates etc. Studies comparing 161P2F10B positive and 161P2F10B negative cells as well as studies comparing unstimulated/resting cells and cells treated with epithelial cell activators, such as cytokines, growth factors, androgen and anti-integrin Ab reveal unique interactions.

In addition, protein-protein interactions are confirmed using two yeast hybrid methodology (Curr Opin Chem Biol. 1999, 3:64). A vector carrying a library of proteins fused to the activation domain of a transcription factor is introduced into yeast expressing a 161P2F10B-DNA-binding domain fusion protein and a reporter construct. Protein-protein interaction is detected by colorimetric reporter activity. Specific association with effector molecules and transcription factors directs one of skill to the mode of action of 161P2F10B, and thus identifies therapeutic, prognostic, preventative and/or diagnostic targets for cancer. This and similar assays are also used to identify and screen for small molecules that interact with 161P2F10B. Thus it is found that 161P2F10B associates with proteins and small molecules. Accordingly, 161P2F10B and these proteins and small molecules are used for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 50: Involvement in Cell Adhesion

Cell adhesion plays a critical role in tissue colonization and metastasis. 161P2F10B can participate in cellular organization, and as a consequence cell adhesion and motility. To confirm that 161P2F10B regulates cell adhesion, control cells lacking 161P2F10B are compared to cells expressing 161P2F10B, using techniques previously described (see, e.g., Haier et al, Br. J. Cancer. 1999, 80:1867; Lehr and Pienta, J. Natl. Cancer Inst. 1998, 90:118). Briefly, in one embodiment, cells labeled with a fluorescent indicator, such as calcein, are incubated on tissue culture wells coated with media alone or with matrix proteins. Adherent cells are detected by fluorimetric analysis and percent adhesion is calculated. In another embodiment, cells lacking or expressing 161P2F10B are analyzed for their ability to mediate cell-cell adhesion using similar experimental techniques as described above. Both of these experimental systems are used to identify proteins, antibodies and/or small molecules that modulate cell adhesion to extracellular matrix and cell-cell interaction. Cell adhesion plays a critical role in tumor growth, progression, and, colonization, and 161P2F10B is involved in these processes. Thus, it serves as a diagnostic, prognostic, preventative and/or therapeutic modality.

Example 51: Phosphodiesterase Activity of 161P2F10B expressing recombinant cell lines.

In order to delineate the function 161P2F10B, several cell lines that lack 161P2F10B were transduced with 161P2F10B-encoding retrovirus as described in example 8, Production of Recombinant 161P2F10B in Higher Eukaryotic Systems, above. Cell lines were characterized for 161P2F10B cell surface expression by FACS analysis (figures 28, 29, 30, and 16). cDNA was stably introduced into the fibroblast lines NIH 3T3 and Rat-1, myeloma NSO cells, and kidney cancer CaKi cells. The cells were immunostained with anti-CD203c mAb and analyzed by flow cytometry. Figures 28, 29, 30, and 16 show that while parental cells fail to express 161P2F10B, engineered lines demonstrate abundant expression of 161P2F10B on their cell surface. Expression of 161P2F10B in engineered cells was compared to that in UT7, a cell line that expresses 161P2F10B endogenously (Figure 28). Our results show that engineered Rat-1 and 3T3 cells express 161P2F10B at levels comparable to UT7 cells.

Since 161P2F10B is identical to the ecto-enzyme ENPP3 phosphodiesterase, and members of the ENPP family possess pyrophosphatase activities, the recombinant cell lines were also characterized for phosphodiesterase activity (figures 28, 29, 30, and 16). Control and 161P2F10B-expressing cells were lysates or intact cells were incubated for at 37 degrees in 20mM Tris/HCL, pH 9.6 containing 5 mM MgCl₂ and 1mM p-nitrophenyl thymidine-5'-L-monophosphate. The reaction was terminated by the addition of 0.1 N NaOH and the reaction product quantified by reading absorbance at 410nm. Figures 28, 29, 30, and 16 show that 161P2F10B expression parallels phosphodiesterase activity. Using CaKi cells

expressing either wild type or mutant 161P2F10B, we show that mutation of T205 inhibits phosphodiesterase activity (figure 30). When 161P2F10B shows phosphodiesterase activity, it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

In addition to phosphodiesterase activity, members of the ENPP family exhibit lysophospholipase D (lysoPLD) activity (Umezū-Goto M et al, J Cell Biol. 2002, 158: 227). ENPP-2 (aka autotoxin) in particular was found to act on lysophosphatidylcholine (LPC) to generate lysophosphatidic acid (LPA) (Umezū-Goto M et al, J Cell Biol. 2002, 158: 227; Tokumura A et al, J Biol. Chem 2002, 277:39436). LPA is involved in various biological functions associated with tumor development, including cell proliferation and invasion (Gschwind A, Prenzel N, Ullrich A. Cancer Res. 2002, 62:6329). Based on the homology of 161P2F10B to other ENPP family members, 161P2F10B has lysoPLD activity. The lysoPLD activity of 161P2F10B expressing cells is compared to cells lacking 161P2F10B using a standard choline release assay. In short, cell lysates are incubated with LPC for 1 hr at 37°C. Liberated choline is detected by fluoremetry following the addition of choline oxidase. When 161P2F10B shows lysoPLD activity, it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 52: RNA interference (RNAi)

Several methods of reducing or abolishing the expression of specific genes have been used for confirming the importance of said genes in tumor growth and progression. These methods include antisense oligonucleotides, morpholino, ribozyme, etc that function in a sequence specific manner to prevent gene transcription or translation. More recently, RNA interference by duplexes of short nucleotide RNAs has been shown to inhibit gene expression in a sequence specific manner in mammalian cells (Elbashir S et al, Nature 2001, 411:494). RNA interference (RNAi) makes use of sequence specific double stranded RNA known as small interfering RNAs (siRNAs) to prevent gene expression. Small interfering RNA (siRNA) are transfected into mammalian cells and thereby mediate sequence specific mRNA degradation. (Elbashir, et al, Nature, 2001; 411: 494). Similarly, siRNA have been used to generate stable vector systems that can be delivered *in vitro* and *in vivo* to mammalian cells, thereby providing therapeutic use for siRNAs (Lee N et al, Nature Biotechnol 2002, 19:500). Several siRNAs can be used to modulate the expression of 161P2F10B in mammalian cells, including for example the following siRNA oligonucleotide sequences:

161P2F10B (1) target: GAAUCUACGUUGACUUUAG (corresponding to nucleotides 4-23 of 161P2F10B ORF)
(SEQ ID NO: 39)

The sense strand of 161P2F10B (1) can be labeled at 3' with fluorescein, 6-FAM (ABS 494nm, EMM 525 nm, green) for easy detection. The siRNA is dissolved in RNA-free sterile buffer (100mM KOAc, 30 mM HEPES KOH, 2mM MOAc, at pH 7.4) to make 20 µM stock (200x). The siRNA is transfected into various normal and tumor cells, including UT7, 3T3-161P2F10B, CaKi-161P2F10B and Rat-161P2F10B cells. Control, non-specific oligonucleotide is used as a control to rule out any non-specific effect of 161P2F10B siRNA oligonucleotides

Protein expression is determined 24-48 hours after transfection by immunostaining followed by flow cytometry. In addition, confirmation of altered gene expression is performed by Western blotting. Cells transfected with control or 161P2F10B-specific siRNAi are compared using functional assays described above, including invasion, proliferation, colony formation and response to apoptotic stimuli. Therefore, the RNA oligonucleotide sequences are used to assess how modulating the expression of a 161P2F10B gene affects function of cancer cells and/or tissues.

Accordingly, the RNA oligonucleotide sequences are used in therapeutic and prophylactic applications.

Example 53: Generation of antibodies to 161P2F10B using peptide encoding the catalytic domain of 161P2F10B as the immunogen.

In one embodiment peptides of 22 amino acids encompassing the 161P2F10B catalytic domain (Threonine (T) at position 205), CGIHSKYMRAMYPTKTFPNHYT (SEQ ID NO: 40) were generated. These were synthesized and the peptides were coupled to KLH through the N-terminal cysteine residue.

Balb/c mice were immunized intraperitoneally (i.p.) with 10 µg of peptide every 2 weeks over a 4 week period. The initial immunization was given i.p. in Complete Freund's Adjuvant (CFA) and the subsequent two immunizations were given i.p. in Incomplete Freund's Adjuvant (IFA).

To determine the specificity of the response following immunization, mice were bled 10 days after the final immunization. Reactivity was determined by Enzyme Linked Immunosorbent Assay (ELISA) using non-KLH conjugated (free) peptide as a target. Mice with the highest titers were given a final boost of 10 µg peptide in PBS and sacrificed for fusion 3 days later. Spleen cells from the immunized mice were fused with mouse Sp2/0 myeloma cells using PEG 1500 according to standard protocols (Kohler et al, Eur. J. Immunol 6: 511 (1976)). Fused cells were plated in 10 96 well microtiter plates and hybridomas were selected using HAT media supplement. Supernatants from fusion wells were screened 10-17 days later by ELISA against 161P2F10B peptide, and clones were then checked for the ability of the monoclonal antibody to recognize cell membrane 161P2F10B by FACS on 161P2F10 expressing Rat-1 cells.

Example 54: Generation of antibodies to 161P2F10B using protein encoding the whole extra cellular domain (aa 1-975) of 161P2F10B as the immunogen.

In one embodiment the whole extra cellular domain of 161P2F10B fused at the C' terminal with 6 Histidines (6-His for purification) was purified for use as an immunogen.

Balb/c mice were immunized intraperitoneally (i.p.) with 10 µg of protein every 2 weeks over a 4 week period. The initial immunization was given i.p. in Complete Freund's Adjuvant (CFA) and the subsequent two immunizations were given i.p. in Incomplete Freund's Adjuvant (IFA).

To determine the specificity of the response following immunization, mice were bled 10 days after the final immunization. Reactivity was determined by Enzyme Linked Immunosorbent Assay (ELISA) using purified protein as a screening agent.

Mice with the highest titers were given a final boost of 10 µg protein in PBS and sacrificed for fusion 3 days later. Spleen cells from the immunized mice were fused with mouse Sp2/0 myeloma cells using PEG 1500 according to standard protocols (Kohler et al, Eur. J. Immunol 6: 511 (1976)). Fused cells were plated in 10 96 well microtiter plates and hybridomas were selected using HAT media supplement. Supernatants from fusion wells were screened 10-17 days later by ELISA against 161P2F10B protein, and clones were then checked for the ability of the monoclonal antibody to recognize cell membrane 161P2F10B by FACS on 161P2F10 expressing Rat-1 cells.

Example 55: Generation of Mabs to 161P2F10B Using DNA Immunization with a vector encoding 161P2F10B fused at the C' terminus with human IgG Fc.

In another embodiment, a vector was constructed that encodes the 975 amino acids of the 161P2F10 extra cellular domain fused at the C-terminus to the human immunoglobulin G1 (IgG) Fc (hinge, CH2, CH3 regions). This construct was used in a DNA based immunization strategy.

Balb/c mice were immunized intra-dermally (ID) at the base of their tail. Three immunizations were given to each mouse of 100 µg of DNA in PBS over a two-week period. To increase the immune response, each mouse was given an i.p. boost of 2 µg of 161P2F10B-Fc protein in tissue culture media 10 days after the final DNA immunization. Bleeds were

collected 10 days after the final immunization and reactivity in the sera to the middle loop of 161P2F10B was tested by ELISA using 161P2F10B-Fc fusion protein as a target (test 1). In parallel the sera were also tested on an unrelated human Fc fusion protein (test 2). Specific reactivity to the 161P2F10B portion of the fusion protein was indicated.

All mice were sacrificed and fusions and hybridoma selection was carried out as described in Example 54. Hybridoma supernatants were screened 10-17 days later by ELISA using 161P2F10B-Fc protein as target. 161P2F10B-Fc positives were subsequently cross-screened on irrelevant Fc proteins to identify 161P2F10 specific clones. Monoclonal antibodies were tested for specificity and reactivity to cell surface 161p2F10B using recombinant Rat 1 cells. Several antibodies were identified this way including X41(4)6, X41(3)15, X41(3)17, X41(3)29, X41(3)37 and X41(3)50. These antibodies or binding region thereof secreted by a hybridoma entitled X41(3)15/29/37, X41(4)6, X41(3)17, and X41(3)50 were deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____. FACS data for these monoclonal antibodies is shown on (Figure 40).

Example 56: Generation of Mabs to 161P2F10B Using DNA Immunization with a vector encoding 161P2F10B fused at the C' terminus with the myc His Tag.

In another embodiment, a vector was constructed that encodes the 975 amino acids of the 161P2F10 extra cellular domain fused at the C-terminus to the myc-His tag. This construct was used in a DNA based immunization strategy.

Balb/c mice were immunized intra-dermally (ID) at the base of their tail. Three immunizations were given to each mouse of 100µg of DNA in PBS over a two-week period. To increase the immune response, each mouse was given an i.p. boost of 2 µg of 161P2F10B-Fc protein in tissue culture media 10 days after the final DNA immunization. Bleeds were collected 10 days after the final immunization and reactivity in the sera to the middle loop of 161P2F10B was tested by ELISA using 161P2F10B-Fc fusion protein as a target (test 1). In parallel the sera were also tested on an unrelated human Fc fusion protein (test 2). Specific reactivity to the 161P2F10B portion of the fusion protein was indicated. All mice were sacrificed and fusions and hybridoma selection was carried out as described in Example 11. Hybridoma supernatants were screened 10-17 days later by ELISA using 161P2F10B-Fc protein as target. 161P2F10B-Fc positives were subsequently cross-screened on irrelevant Fc proteins to identify 161P2F10 specific clones. Monoclonal antibodies were tested for specificity and reactivity to cell surface 161p2F10B using recombinant Rat 1 cells.

Example 57: Generation of Monoclonal Antibodies specific for 161P2F10B using UT7 cells endogenously expressing 161P2F10B.

It has been reported in the literature that antibodies to 161P2F10B can be made by immunization with the human erythro-megakaryoblastic cell line UT-7 cultured with IL3 (Buhring et.al. Blood 94(7): 2343. 1999). Antibodies described in this publication are available commercially and have been used as controls in the invention described here.

In another embodiment, mice were immunized intra-peritoneally with UT-7 cells, 10⁶ cells per immunization. A total of 5 immunizations were given approximately 2 weeks apart with the final injection being given three days before mice were sacrificed for fusions. Mice were bled 10 days after the third injection and the 161P2F10B specific titer of the sera was determined by ELISA using 161P2F10 as a screening agent. Mice with high titers were then used for fusions as described in Example 11. Monoclonal antibodies generated in this way were selected by ELISA and their ability to recognize cells surface 161P2F10B was confirmed by FACS on Rat 1 cells expressing 161P2F10B.

Example 58: Generation of Monoclonal Antibodies specific for 161P2F10B using the recombinant cell line 3T3 expressing 161P2F10B.

In another embodiment, mice were immunized intra-peritoneally with 3T3 cells expressing 161P2F10B, 10^6 cells per immunization. A total of 5 immunizations were given approximately 2 weeks apart with the final injection being given three days before mice were sacrificed for fusions. Mice were bled 10 days after the third injection and the 161P2F10B specific titer of the sera was determined by ELISA using 161P2F10 as a screening agent. Mice with high titers were then used for fusions as described in Example 11. Monoclonal antibodies generated in this way were selected by ELISA and their ability to recognize cells surface 161P2F10B was confirmed by FACS on Rat 1 cells expressing 161P2F10B.

Example 59: Generation of Monoclonal Antibodies specific for 161P2F10B using the recombinant cell line Rat 1 expressing 161P2F10B.

In another embodiment mice were immunized with Rat-1 cells expressing 161P2F10B. Mice were then used for fusions as described in Example 11. Monoclonal antibodies generated in this way were selected by ELISA and their ability to recognize cells surface 161P2F10B was confirmed by FACS on Rat 1 cells expressing 161P2F10B.

Example 60: Detection of 161P2F10B protein in kidney cancer patient specimens

To confirm the expression of 161P2F10B protein, kidney cancer specimens were obtained from kidney cancer patients, and stained using the commercially available antibody 97A6 specific for ENPP3 protein (also called anti-CD203c) (Immunotech, Marseilles, France). Briefly, frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with PE-labeled mouse monoclonal anti-ENPP3 antibody for 3 hours (Figure 24 A-C), or isotype control antibody (Figure 44 G-I). The slides were washed three times in buffer, and either analyzed by fluorescence microscopy (Figure 44 A, B and C), or further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour (Figure 44 D, E, and F FIG 24 A-C). The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy (Figure 44 D, E and F). The results showed strong expression of 161P2F10B in the renal carcinoma patient tissue (Figure 44A and D) and the kidney cancer metastasis to lymph node tissue (Figure 44 C and F), but weakly in normal kidney (Figure 44 B and E). The expression was detected mostly around the cell periphery in renal clear cell carcinoma (FIG 44 A and D, FIG 24 A and B) and was strongly expressed throughout the cells with an apparent predisposition towards the cell periphery in renal papillary carcinoma (FIG 24 C) indicating that 161P2F10B is membrane associated in kidney cancer tissues. The weak expression detected in normal kidney was localized to the kidney tubules. The sections stained with the isotype control antibody were negative showing the specificity of the anti- ENPP3 antibody (Figure 44 G-I). Kidney cancer specimens were obtained from patients with different types of renal tumor including renal clear cell carcinoma; papillary cell carcinoma; renal cell carcinoma, chromophobe type; transitional cell carcinoma and oncocytoma and were stained for 161P2F10B using the commercially available antibody 97A6 specific for ENPP3 protein (also called anti-CD203c) (Immunotech, Marseilles, France). All tissue specimens for renal clear cell carcinoma and papillary cell carcinoma were positive for 161P2F10B (Table LIX).

Figure 45 shows expression of 161P2F10B in human patient cancers by Western blot analysis. Cell lysates from kidney cancer tissues (KiCa), kidney cancer metastasis to lymph node (KiCa Met), as well as normal kidney (NK) were subjected to western analysis using an anti-161P2F10B mouse monoclonal antibody. Briefly, tissues (~25 µg total protein) were solubilized in SDS-PAGE sample buffer and separated on a 10-20% SDS-PAGE gel and transferred to nitrocellulose. Blots were blocked in Tris-buffered saline (TBS) + 3% non-fat milk and then probed with purified anti-161P2F10B antibody in TBS + 0.15% Tween-20 + 1% milk. Blots were then washed and incubated with a 1:4,000 dilution of anti-mouse IgG-HRP conjugated secondary antibody. Following washing, anti-161P2F10B immunoreactive bands were developed and visualized by enhanced chemiluminescence and exposure to autoradiographic film. The specific anti-161P2F10B immunoreactive

bands represent a monomeric form of the 161P2F10B protein, which runs at approximately 130kDa. These results demonstrate that 161P2F10B is useful as a diagnostic and therapeutic target for kidney cancers, metastatic cancers and other human cancers that express this protein.

The strong expression of 161P2F10B in kidney cancer tissues and its restricted expression in normal kidney as well as its membrane localization show that 161P2F10B is a target, e.g., for kidney cancer diagnosis and therapy. The expression detected in kidney cancer metastatic tissue indicates that 161P2F10B is also a target for metastatic disease. As disclosed herein, Western blot and immunohistochemical analysis of kidney cancer tissues and kidney cancer xenografts with mAb 97A6 showed strong extensive staining of ENPP3 in clear cell kidney carcinoma but significantly lower or undetectable levels in normal kidney (Figures 44, 45, 46, and 24). Detection of 161P2F10B (ENPP3) in high grade clear cell carcinoma and in metastatic disease.

Example 61: Detection of 161P2F10B protein in colon cancer patient specimens

Tissue specimens of colon adenocarcinoma were obtained from nine different colon cancer patients. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours. The slides were washed three times in buffer, and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed strong expression of 161P2F10B in two of the nine colon cancer patient tissues, one of which is illustrated (Figure 26). 161P2F10B was most strongly expressed on the tumor cells with a luminal cell surface but was also expressed throughout all the tumor tissue.

Example 62: Detection of 161P2F10B protein by immunohistochemistry in a prostate cancer patient specimens.

Tissue specimens of prostate adenocarcinoma were obtained from eight different prostate cancer patients. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours. The slides were washed three times in buffer, and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed expression of 161P2F10B in six of the eight prostate cancer patient tissues, one of which is illustrated (Figure 25). 161P2F10B was expressed on the tumor cells with an apparent proclivity towards the luminal cell surface.

Example 63: Detection of 161P2F10B protein by immunohistochemistry in normal tissue specimens.

Normal tissue specimens from a number of organs were obtained either from patients undergoing surgery or from autopsy. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated with mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours. The slides were washed three times in buffer, and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed weak expression of 161P2F10B in some of the tubules in all of the kidney specimens and weak staining of some glandular epithelium in half of the prostate tissues. There was no expression of 161P2F10B in any of the other normal tissues studied except for expression in a very few cells within one lung, one bladder and two colon samples which could be

mast cells (TABLE LX). As disclosed herein, Western blot and immunohistochemical analysis of kidney cancer tissues and kidney cancer xenografts with mAb 97A6 showed strong extensive staining of ENPP3 in clear cell kidney carcinoma but significantly lower or undetectable levels in normal kidney (Figures 44, 45, 46, and 24). Detection of 161P2F10B (ENPP3) in high grade clear cell carcinoma and in metastatic disease.

Example 64: Detection by immunohistochemistry of 161P2F10B protein expression in kidney clear cell cancer patient specimens by specific binding of mouse monoclonal antibodies.

Renal clear cell carcinoma tissue and its matched normal adjacent were obtained from a kidney cancer patient. Frozen tissues were cut into 4 micron sections and fixed in acetone for 10 minutes. The sections were then incubated either mouse monoclonal anti-ENPP3 antibody (Coulter-Immunotech, Marseilles, France) for 3 hours (Figure 27 panels A, D), or mouse monoclonal antibody X41(3)50 (Figure 27 panels B, E), or mouse monoclonal antibody X41(3)37 (Figure 27 panels C, F). The slides were washed three times in buffer and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-mouse secondary antibody (DAKO Corporation, Carpinteria, CA) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy (Figure 27 panels A-F). The results showed strong expression of 161P2F10B in the renal clear cell carcinoma patient tissue (Figure 27 panels A-C), but weakly in normal kidney (Figure 27 panels D-F). The expression was predominantly around the cell periphery indicating that 161P2F10B is membrane associated in kidney cancer tissues. The weak expression detected in normal kidney was localized to the kidney proximal tubules. As disclosed herein, Western blot and immunohistochemical analysis of kidney cancer tissues and kidney cancer xenografts with mAb 97A6 showed strong extensive staining of ENPP3 in clear cell kidney carcinoma but significantly lower or undetectable levels in normal kidney (Figures 44, 45, 46, and 24). Detection of 161P2F10B (ENPP3) in high grade clear cell carcinoma and in metastatic disease.

Example 65: Characteristics and utility of anti-161P2F10b MAbs.

Using a variety of immunization strategies as described in Example 11, a panel of MAbs that specifically bind 161P2F10b protein was generated. The characteristics of this panel is summarized in Figure 39. These antibodies specifically bind with high affinity to 161P2F10b on the surface of endogenously-expressing and recombinant cell lines as determined by flow cytometry (Figures 28 and 40). Upon engagement of surface 161P2F10b, these MAbs mediate internalization of the MAb-protein complex (Figures 33, 34, and 35). These MAbs are thus useful as a specific targeting modality for toxin-conjugates, as exemplified by the growth inhibition and induction of apoptosis of Caki-161P2F10b cells by MAb X41.50 with a saporin toxin-conjugated secondary Ab (Figure 36). Treatment of 161P2F10-expressing cancerous cells with the naked MAb also has a therapeutic effect in vivo as exemplified by the inhibition of UGK3 tumor formation in SCID mice injected with MAb X41.50 (Figure 23).

161P2F10b encodes phosphodiesterase enzymatic activity that is easily monitored both in recombinant purified protein (Figure 31) and on cells (Figure 32). The relevance of the enzymatic activity to the function of 161P2F10b may be monitored by utilization of mutants that disrupt this activity (Figure 30). Engagement of 161P2F10b with MAbs may alter, disrupt, block, or downregulate 161P2F10 enzymatic activity, which may serve as a potential therapeutic mechanism for targeting 161P2F10b-expressing cancers and diseased tissues. Engagement of cell surface 161P2F10b cells with a subset of the MAbs listed in Figure 39 does mediate internalization and marked downregulation of cell surface enzymatic activity (Figure 37 and 38) thus demonstrating the utility of the MAbs for disrupting the function of 161P2F10b in cells and tissues.

161P2F10b protein and the MAbs that bind it are useful in the diagnosis of 161P2F10b-expressing cancer and diseased tissues. Immunohistochemical analysis of the panel of MAbs, as summarized in Figure 39, specifically stain (to varying degrees) a variety of kidney cancer samples with little to no staining of adjacent normal tissues. These MAbs are

thus useful as diagnostic reagents for a variety of 161P2F10b-expressing cancers by immunohistochemistry and are potentially useful as imaging reagents in patients. In addition, the MAbs were used (specifically X48.54 and X41.29, but others that do not compete for the same epitope are also used) to demonstrate the shedding and/or secretion of the protein from 161P2F10b-expressing cancer cells and tissues (Figures 42 and 43). This supports the utility of 161P2F10b as a serum and/or urine diagnostic marker and the MAbs as reagents to quantitatively measure serum and/or urine concentrations of 161P2F10b protein.

Throughout this application, various website data content, publications, patent applications and patents are referenced. (Websites are referenced by their Uniform Resource Locator, or URL, addresses on the World Wide Web.) The disclosures of each of these references are hereby incorporated by reference herein in their entireties.

The present invention is not to be limited in scope by the embodiments disclosed herein, which are intended as single illustrations of individual aspects of the invention, and any that are functionally equivalent are within the scope of the invention. Various modifications to the models and methods of the invention, in addition to those described herein, will become apparent to those skilled in the art from the foregoing description and teachings, and are similarly intended to fall within the scope of the invention. Such modifications or other embodiments can be practiced without departing from the true scope and spirit of the invention.

TABLES:

TABLE I: Tissues that Express 161P2F10B:

- a. Exemplary Normal Tissues:
 - Prostate
 - Kidney
- b. Malignant Tissues
 - Kidney
 - Uterus
 - Pancreas
 - Prostate
 - Colon
 - Lung
 - Bone
 - Lymphoma
 - Breast
 - Ovary

TABLE II: Amino Acid Abbreviations

SINGLE LETTER	THREE LETTER	FULL NAME
F	Phe	phenylalanine
L	Leu	leucine
S	Ser	serine
Y	Tyr	tyrosine
C	Cys	cysteine
W	Trp	tryptophan
P	Pro	proline
H	His	histidine
Q	Gln	glutamine
R	Arg	arginine
I	Ile	isoleucine
M	Met	methionine
T	Thr	threonine
N	Asn	asparagine
K	Lys	lysine
V	Val	valine
A	Ala	alanine
D	Asp	aspartic acid
E	Glu	glutamic acid
G	Gly	glycine

TABLE III: Amino Acid Substitution Matrix

Adapted from the GCG Software 9.0 BLOSUM62 amino acid substitution matrix (block substitution matrix). The higher the value, the more likely a substitution is found in related, natural proteins. (See world wide web URL ikp.unibe.ch/manual/blosum62.html)

[illegible]

TABLE IV:
HLA Class I/II Motifs/Supermotifs

TABLE IV (A): HLA Class I Supermotifs/Motifs

SUPERMOTIF	POSITION	POSITION	POSITION
	2 (Primary Anchor)	3 (Primary Anchor)	C Terminus (Primary Anchor)
A1	T ILVMS		F WY
A2	LIV M ATQ		IV M ATL
A3	V S MATLI		R K
A24	Y F WIVLMT		FIYW L M
B7	P		VILFMW Y A
B27	R H K		F Y LWMIVA
B44	E D		FW Y LIMVA
B58	A T S		FW Y LIVMA
B62	Q L IVMP		FW Y MIVLA
MOTIFS			
A1	T S M		Y
A1		D E A S	Y
A2.1	L M VQ I A T		V L I M A T
A3	L M VISAT F CG D		K Y R H F A
A11	V T M L ISAG N C D F		K R Y H
A24	Y F W M		F L I W
A*3101	M V T A L I S		R K
A*3301	M V AL F I S T		R K
A*6801	A V T M S L I		R K
B*0702	P		L M F W Y A I V
B*3501	P		L M F W Y I V A
B51	P		L I V F W Y A M
B*5301	P		I M F W Y A L V
B*5401	P		A T I V L M F W Y

Bolded residues are preferred, italicized residues are less preferred: A peptide is considered motif-bearing if it has primary anchors at each primary anchor position for a motif or supermotif as specified in the above table.

TABLE IV (B): HLA Class II Supermotif

1	6	9
W, F, Y, V, .I, L	A, V, I, L, P, C, S, T	A, V, I, L, C, S, T, M, Y

TABLE IV (C): HLA Class II Motifs

MOTIFS		1° anchor 1	2	3	4	5	1° anchor 6	7	8	9
DR4	preferred deleterious	FMYLIVW	M	T	W	I	VSTCPALIM	MH R		MH WDE
DR1	preferred deleterious	MFLIVWY	C	CH	PAMQ FD	CWD	VMATSPLIC	M GDE	D	AVM
DR7	preferred deleterious	MFLIVWY	M C	W	A G		IVMSACTPL	M GRD		IV G
DR3	<u>MOTIFS</u>	1° anchor 1	2	3	1° anchor 4	5	1° anchor 6			
Motif a preferred		LIVMFY			D					
Motif b preferred		LIVMFAY			DNQEST		KRH			
DR Supermotif		MFLIVWY					VMSTACPLI			

Italicized residues indicate less preferred or "tolerated" residues

TABLE IV (D): HLA Class I Supermotifs

	POSITION:	1	2	3	4	5	6	7	8	C-terminus
<u>SUPER-MOTIFS</u>										
A1			1° Anchor TILVMS							1° Anchor FWY
A2			1° Anchor LIVMATQ							1° Anchor LIVMAT
A3	Preferred		1° Anchor VSMATLI	YFW (4/5)			YFW (3/5)	YFW (4/5)	P (4/5)	1° Anchor RK
	deleterious	DE (3/5); P (5/5)		DE (4/5)						
A24			1° Anchor YFWIVLMT							1° Anchor FIYWLM
B7	Preferred	FWY (5/5)	1° Anchor P	FWY (4/5)					FWY (3/5)	1° Anchor VILFMWYA
	deleterious	DE (3/5); P(5/5); G(4/5); A(3/5); QN(3/5)			DE (3/5)	G (4/5)	QN (4/5)		DE (4/5)	
B27			1° Anchor RHK							1° Anchor FYLWMI/VA
B44			1° Anchor ED							1° Anchor FWYLIMVA
B58			1° Anchor ATS							1° Anchor FWYLIVMA
B62			1° Anchor QLIVMP							1° Anchor FWYMIVLA

Italicized residues indicate less preferred or "tolerated" residues

TABLE IV (E): HLA Class I Motifs

	POSITION 1	2	3	4	5	6	7	8	9	C-terminus
									or C-terminus	
A1 9-mer	preferred GFYW	<u>1°Anchor</u> STM	DEA	YFW		P	DEQN	YFW	<u>1°Anchor</u> Y	
	deleterious DE		RHKLIVMP	A	G	A				
A1 9-mer	preferred GRHK	ASTCLIVM	<u>1°Anchor</u> DEAS	GSTC		ASTC	LIVM	DE	<u>1°Anchor</u> Y	
	deleterious A	RHKDEPYFW		DE	PQN	RHK	PG	GP		
A1 10-mer	preferred YFW	<u>1°Anchor</u> STM	DEAQN	A	YFWQN		PASTC	GDE	P	<u>1°Anchor</u> Y
	deleterious GP		RHKGLIVM	DE	RHK	QNA	RHKYFW	RHK	A	
A1 10-mer	preferred YFW	STCLIVM	<u>1°Anchor</u> DEAS	A	YFW		PG	G	YFW	<u>1°Anchor</u> Y
	deleterious RHK	RHKDEPYFW			P	G		PRHK	QN	
A2.1 9-mer	preferred YFW	<u>1°Anchor</u> LM/VQAT	YFW	STC	YFW		A	P	<u>1°Anchor</u> VLIMAT	
	deleterious DEP		DERKH			RKH	DERKH			
	POSITION: 1	<u>2</u>	3	4	5	6	7	8	9	C-Terminus
A2.1 10-mer	preferred AYFW	<u>1°Anchor</u> LM/VQAT	LVIM	G		G		FYWL VIM		<u>1°Anchor</u> VLIMAT
	deleterious DEP		DE	RKHA	P		RKH	DERK H	RKH	
A3	preferred RHK	<u>1°Anchor</u> LMVISATFCGD	YFW	PRHKYF W	A	YFW		P	<u>1°Anchor</u> KYRHFA	
	deleterious DEP		DE							
A11	preferred A	<u>1°Anchor</u> VTLMISAGNCD F	YFW	YFW	A	YFW	YFW	P	<u>1°Anchor</u> KRYH	
	deleterious DEP						A	G		
A24 9-mer	preferred YFWRHK	<u>1°Anchor</u> YFWM		STC			YFW	YFW	<u>1°Anchor</u> FLIW	
	deleterious DEG		DE	G	QNP	DERH K	G	AQN		
A24 10-mer	Preferred	<u>1°Anchor</u> YFWM		P	YFWP		P			<u>1°Anchor</u> FLIW
	Deleterious		GDE	QN	RHK	DE	A	QN	DEA	
A310 1	Preferred RHK	<u>1°Anchor</u> MVTALIS	YFW	P		YFW	YFW	AP	<u>1°Anchor</u> RK	
	Deleterious DEP		DE		ADE	DE	DE	DE		
A330 1	Preferred	<u>1°Anchor</u> MVALFIST	YFW				AYFW		<u>1°Anchor</u> RK	
	Deleterious GP		DE							
A680 1	Preferred YFWSTC	<u>1°Anchor</u> AVTMSLI			YFWLIV M		YFW	P	<u>1°Anchor</u> RK	
	deleterious GP		DEG		RHK			A		
B070 2	Preferred RHKFWY	<u>1°Anchor</u> P	RHK		RHK	RHK	RHK	PA	<u>1°Anchor</u> LMFWYA/ V	
	deleterious DEQNP		DEP	DE	DE	GDE	QN	DE		

	POSITION 1	2	3	4	5	6	7	8	9	C-terminus
									or C-terminus	
A1 9-mer	preferred GFYW	<u>1°Anchor</u> STM	DEA	YFW		P	DEQN	YFW	<u>1°Anchor</u> Y	
	deleterious DE		RHKLIVMP	A	G	A				
A1 9-mer	preferred GRHK	ASTCLIVM	<u>1°Anchor</u> DEAS	GSTC		ASTC	LIVM	DE	<u>1°Anchor</u> Y	
	deleterious A	RHKDEPYFW		DE	PQN	RHK	PG	GP		
B350 1	Preferred FWYLIVM	<u>1°Anchor</u> P	FWY				FWY		<u>1°Anchor</u> LMFWY/V A	
	deleterious AGP				G	G				
B51	Preferred LIVMFWY	<u>1°Anchor</u> P	FWY	STC	FWY		G	FWY	<u>1°Anchor</u> LIVFWYA M	
	deleterious AGPDER HKSTC				DE	G	DEQN	GDE		
B530 1	preferred LIVMFWY	<u>1°Anchor</u> P	FWY	STC	FWY		LIVMFW Y	FWY	<u>1°Anchor</u> IMFWYAL V	
	deleterious AGPQN					G	RHKQN	DE		
B540 1	preferred FWY	<u>1°Anchor</u> P	FWYLIVM		LIVM		ALIVM	FWYA P	<u>1°Anchor</u> ATIVLMF WY	
	deleterious GPQNDE		GDESTC		RHKDE	DE	QNDGE	DE		

TABLE IV (F):

Summary of HLA-supertypes								
Overall phenotypic frequencies of HLA-supertypes in different ethnic populations								
Specificity			Phenotypic frequency					
Supertype	Position 2	C-Terminus	Caucasian	N.A. Black	Japanese	Chinese	Hispanic	Average
B7	P	AILMVFWY	43.2	55.1	57.1	43.0	49.3	49.5
A3	AILMVST	RK	37.5	42.1	45.8	52.7	43.1	44.2
A2	AILMVT	AILMVT	45.8	39.0	42.4	45.9	43.0	42.2
A24	YF (WIVLMT)	FI (YWLM)	23.9	38.9	58.6	40.1	38.3	40.0
B44	E (D)	FWYLIMVA	43.0	21.2	42.9	39.1	39.0	37.0
A1	TI (LVMS)	FWY	47.1	16.1	21.8	14.7	26.3	25.2
B27	RHK	FYL (WMI)	28.4	26.1	13.3	13.9	35.3	23.4
B62	QL (IVMP)	FWY (MIV)	12.6	4.8	36.5	25.4	11.1	18.1
B58	ATS	FWY (LIV)	10.0	25.1	1.6	9.0	5.9	10.3

TABLE IV (G):

Calculated population coverage afforded by different HLA-supertype combinations

HLA-supertypes		Phenotypic frequency					
		Caucasian	N.A Blacks	Japanese	Chinese	Hispanic	Average
A2, A3 and B7 A2, A3, B7, A24, B44 and A1 A2, A3, B7, A24, B44, A1, B27, B62, and B 58		83.0	86.1	87.5	88.4	86.3	86.2
		99.5	98.1	100.0	99.5	99.4	99.3
		99.9	99.6	100.0	99.8	99.9	99.8

Motifs indicate the residues defining supertype specificities. The motifs incorporate residues determined on the basis of published data to be recognized by multiple alleles within the supertype. Residues within brackets are additional residues also predicted to be tolerated by multiple alleles within the supertype.

Table V: Frequently Occurring Motifs

Name	avrg. % identity	Description	Potential Function
zf-C2H2	34%	Zinc finger, C2H2 type	Nucleic acid-binding protein functions as transcription factor, nuclear location probable
cytochrome_b_N	68%	Cytochrome b(N-terminal)/b6/petB	membrane bound oxidase, generate superoxide
Ig	19%	Immunoglobulin domain	domains are one hundred amino acids long and include a conserved intradomain disulfide bond.
WD40	18%	WD domain, G-beta repeat	tandem repeats of about 40 residues, each containing a Trp-Asp motif. Function in signal transduction and protein interaction
PDZ	23%	PDZ domain	may function in targeting signaling molecules to sub-membranous sites
LRR	28%	Leucine Rich Repeat	short sequence motifs involved in protein-protein interactions
Pkinase	23%	Protein kinase domain	conserved catalytic core common to both serine/threonine and tyrosine protein kinases containing an ATP binding site and a catalytic site

PH	16%	PH domain	pleckstrin homology involved in intracellular signaling or as constituents of the cytoskeleton
EGF	34%	EGF-like domain	30-40 amino-acid long found in the extracellular domain of membrane-bound proteins or in secreted proteins
Rvt	49%	Reverse transcriptase (RNA-dependent DNA polymerase)	
Ank	25%	Ank repeat	Cytoplasmic protein, associates integral membrane proteins to the cytoskeleton
Oxidored_q1	32%	NADH-Ubiquinone/plastoquinone (complex I), various chains	membrane associated. Involved in proton translocation across the membrane
Efhand	24%	EF hand	calcium-binding domain, consists of a 12 residue loop flanked on both sides by a 12 residue alpha-helical domain
Rvp	79%	Retroviral aspartyl protease	Aspartyl or acid proteases, centered on a catalytic aspartyl residue
Collagen	42%	Collagen triple helix repeat (20 copies)	extracellular structural proteins involved in formation of connective tissue. The sequence consists of the G-X-Y and the polypeptide chains forms a triple helix.
Fn3	20%	Fibronectin type III domain	Located in the extracellular ligand-binding region of receptors and is about 200 amino acid residues long with two pairs of cysteines involved in disulfide bonds
7tm_1	19%	7 transmembrane receptor (rhodopsin family)	seven hydrophobic transmembrane regions, with the N-terminus located extracellularly while the C-terminus is cytoplasmic. Signal through G proteins

Table VI: Motifs and Post-translational Modifications of 161P2F10B

N-glycosylation site:

Number of matches: 10

- 1 236-239 NFSL (SEQ ID NO: 41)
- 2 279-282 NGSF (SEQ ID NO: 42)
- 3 290-293 NGSV (SEQ ID NO: 43)
- 4 426-429 NLSC (SEQ ID NO: 44)
- 5 533-536 NGTH (SEQ ID NO: 45)
- 6 582-585 NSTQ (SEQ ID NO: 46)
- 7 594-597 NLTQ (SEQ ID NO: 47)
- 8 687-690 NITH (SEQ ID NO: 48)
- 9 699-702 NRTS (SEQ ID NO: 49)
- 10 789-792 NKSH (SEQ ID NO: 50)

cAMP- and cGMP-dependent protein kinase phosphorylation site

14-17 KKNT (SEQ ID NO: 51)

Protein kinase C phosphorylation site

WO 03/040340

PCT/US02/36002

142

Number of matches: 13

- 1 17-19 TLK
- 2 53-55 SCR
- 3 428-430 SCR
- 4 62-64 SFR
- 5 92-94 STR
- 6 240-242 SSK
- 7 335-337 SAR
- 8 53-55 SCR
- 9 428-430 SCR
- 10 502-504 SFK
- 11 603-605 TVK
- 12 676-678 SQK
- 13 698-700 SNR

Casein kinase II phosphorylation site

Number of matches: 15

- 1 88-91 TCVE (SEQ ID NO: 52)
- 2 106-109 TRLE (SEQ ID NO: 53)
- 3 114-117 SCSD (SEQ ID NO: 54)
- 4 138-141 SWLE (SEQ ID NO: 55)
- 5 240-243 SSKE (SEQ ID NO: 56)
- 6 502-505 SFKE (SEQ ID NO: 57)
- 7 507-510 TEVE (SEQ ID NO: 58)
- 8 551-554 SHAE (SEQ ID NO: 59)
- 9 584-587 TQLE (SEQ ID NO: 60)
- 10 596-599 TQEE (SEQ ID NO: 61)
- 11 660-663 TVPD (SEQ ID NO: 62)
- 12 704-707 SQYD (SEQ ID NO: 63)
- 13 813-816 TNVE (SEQ ID NO: 64)
- 14 817-820 SCPE (SEQ ID NO: 65)
- 15 846-849 TGLD (SEQ ID NO: 66)

Tyrosine kinase phosphorylation site

700-706 RTSDSQY (SEQ ID NO: 67)

N-myristoylation site

Number of matches: 11

- 1 38-43 GLGLGL (SEQ ID NO: 68)
- 2 40-45 GLGLGL (SEQ ID NO: 69)
- 3 38-43 GLGLGL (SEQ ID NO: 70)
- 4 40-45 GLGLGL (SEQ ID NO: 71)

- 5 65-70 GLENCR (SEQ ID NO: 72)
- 6 222-227 GIIDNN (SEQ ID NO: 73)
- 7 263-268 GLKAAT (SEQ ID NO: 74)
- 8 273-278 GSEVAI (SEQ ID NO: 75)
- 9 280-285 GSFPSI (SEQ ID NO: 76)
- 10 331-336 GGPVSA (SEQ ID NO: 77)
- 11 374-379 GMDQTY (SEQ ID NO: 78)

Cell attachment sequence

78-80 RGD

Somatomedin B domain signature

Number of matches: 2

- 1 69-89 CRCDVACKDRGDCCWDFEDTC (SEQ ID NO: 79)
- 2 113-133 CSCSDDCLQKKDCCADYKSVC (SEQ ID NO: 80)

Table VII:
Search Peptides**161P2F10B variant 1 (SEQ ID NO: 81)**

1 MESTLTATE QPVKKNTLKK YKIACIVLLA LLVIMSLGLG LGLGLRKLEK QGSCRKKCFD
61 ASFRGLENCR CDVACKDRGD CCWDFEDTCV ESTRIWMCNK FRCGETRLEA SLCSCSDDCL
121 QKKDCCADYK SVCQGETSWL EENCDDTAQQS QCPEGFDLPP VILFSMDGFR AEYLYTWDTL
181 MPNINKLKTC GIHSKYMRAM YPTKTFPNHY TIVTGLYPES HGIIDNNMYD VNLNKNFSL
241 SKEQNNPAWW HGQPMWLTAM YQGLKAATYF WPGSEVAING SFPSIYMPYN GSVPFEEERIS
301 TLLKWLDLPK AERPRFYTM YFEEDSSGHA GGPVSARVIK ALQVVDHAFG MLMEGLKQRN
361 LHNCVNIILL ADHGMDQTYC NKMEYMTDYF PRINFFYMYE GPAPRIRAHN IPHDFFSFNS
421 EEIVRNLSR KPDQHFKPYL TPDLPKRLHY AKNVRIDKVH LFVDQQLAV RSKSNTNCGG
481 GNHGYNNEFR SMEAIFLAHG PSFKEKTEVE PFENIEVYNL MCDLLRIQPA PNNGTHGSLN
541 HLLKVPFYEP SHAEVSKFS VCGFANPLPT ESLDCFCPHL QNSTQLEQVN QMLNLTQEEI
601 TATVKVNLPF GRPRVLQKNV DHCLLYHREY VSGFGKAMRM PMWSSYTVPQ LGDTSPLPPT
661 VPDCLRADVR VPPSESQKCS FYLADKNITH GFLYPPASNR TSDSQYDALI TSNLVPMYEE
721 FRKMWDYFHS VLLIKHATER NGVNVVSGPI FDYNYDGHFD APDEITKHLA NTDVPIPTHY
781 FVVLTSCKNK SHTPENCPGW LDVLPFIIPH RPTNVESCPE GKPEALWVEE RFTAHARVR
841 DVELLTGLDF YQDKVQPVSE ILQLKTYLPT FETTI

Variant 2

9-mers

SCSDDCLQ⁸¹KKDCCADYK (SEQ ID NO: 82)

10-mers

SCSDDCLQ⁸¹KKDCCADYKS (SEQ ID NO: 83)

15-mers

LEASLCSCSDDCLQ⁸¹KKDCCADYKSVQGE (SEQ ID NO: 84)**Variant 3**

9-mers

PTNVESCPGGKPEALWV (SEQ ID NO: 85)

10-mers

RPTNVESCPGGKPEALWVE (SEQ ID NO: 86)

15-mers

FIIPHRPTNVESCPGGKPEALWVEERFTA (SEQ ID NO: 87)

Variant 4

9-mers

TYLPTFETPI (SEQ ID NO: 88)

10-mers

KTYLPTFETPI (SEQ ID NO: 89)

15-mers

EILQLKTYLPTFETPI (SEQ ID NO: 90)

WO 03/040340

PCT/US02/36002

145

Table VIII-V1-HLA-A1-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
105	SMDGFRAEY	25.000
754	NVESCPEGK	18.000
55	CSDDCLQKK	15.000
446	KTEVEPFEN	11.250
245	WLDLPKAER	10.000
798	VSEILQLKT	6.750
317	QTYCNKMEY	6.250
371	KPDQHFKPY	6.250
10	RCDVACKDR	5.000
314	GMDQTYCNK	5.000
322	KMEYMTDYF	4.500
454	NIEVYNLMC	4.500
29	CVESTRIWM	4.500
650	ITSNLVPMY	2.500
118	DTLMPNINK	2.500
685	VVSGPIFDY	2.500
559	NVDHCLLYH	2.500
711	NTDVPIPTH	2.500
740	WLDVLPFII	2.500
402	FVDQQWLAV	2.500
431	SMEAIFLAH	2.250
610	RVPPSESQK	2.000
379	YLTPDLPKR	2.000
613	PSESQKCSF	1.350
213	GSEVAINGS	1.350
359	NSEEIVRNL	1.350
128	KTCGIHSKY	1.250
326	MTDYFPRIN	1.250
426	NNEFRSMEA	1.125
632	FLYPPASNR	1.000
163	IIDNNMYDV	1.000
512	SLDCFCPHL	1.000
66	CADYKSVCQ	1.000
653	NLVPMYEEF	1.000
606	RADVRVPPS	1.000
54	SCSDDCLQK	1.000
767	WVEERFTAH	0.900
448	EVEPFENIE	0.900
525	QLEQVNQML	0.900
79	WLEENCDTA	0.900
492	HAEEVSKFS	0.900
537	QEEITATVK	0.900
5	GLENCRCDV	0.900
47	RLEASLCSC	0.900
641	TSDSQYDAL	0.750
626	KNITHGFLY	0.625
558	KNVDHCLLY	0.625
783	ELLTGLDFY	0.500
62	KKDCCADYK	0.500
310	LADHGMDQT	0.500

Table VIII-V1-HLA-A1-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
167	NMYDVNLNK	0.500
562	HCLLYHREY	0.500
197	LTAMYQGLK	0.500
699	FDAPDEITK	0.500
787	GLDFYQDKV	0.500
83	NCDTAQQSQ	0.500
593	DTSPLPPTV	0.500
284	VVDHAFGML	0.500
144	KTFPNHYTI	0.500
417	NCGGGNHGY	0.500
350	NIPHDFFSF	0.500
97	DLPPVILFS	0.500
742	DVLPFIIPH	0.500
250	KAERPRFYT	0.450
292	LMEGLKQRN	0.450
677	ATERNGVNV	0.450
261	FEEPDSSEGH	0.450
618	KCSFYLADK	0.400
571	VSGFGKAMR	0.300
536	TQEEITATV	0.270
193	QPMWLTAMY	0.250
687	SGPIFDYNY	0.250
772	FTAHARVR	0.250
655	VPMYEEFRK	0.250
476	HGSLNHLLK	0.250
712	TDVPIPTHY	0.250
550	FGRPRVLQK	0.250
601	VPDCLRADV	0.250
578	MRMPMWSSY	0.250
546	VNLPGFRPR	0.250
461	MCDLLRIQP	0.250
395	RIDKVHLFV	0.250
586	YTVPQLGDT	0.250
96	FDLPPVILF	0.250
509	PTESLDCFC	0.225
758	CPEGKPEAL	0.225
733	TPENCPGWL	0.225
493	AEVSKFSV	0.225
43	CGETRLEAS	0.225
722	VVLTSCKNK	0.200
746	FIIPHRPTN	0.200
436	FLAHGPSFK	0.200
743	VLPFIIPHR	0.200
133	HSKYMARAMY	0.150
201	YQGLKAATY	0.150
231	GSVPFEERI	0.150
686	VSGPIFDYN	0.150
382	PDLPKRLHY	0.125
191	HGQPMWLTA	0.125
701	APDEITKHL	0.125

Table VIII-V2-HLA-A1-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
2	CSDDCLQRK	15.000
9	RKDCCADYK	0.500
1	SCSDDCLQR	0.500
5	DCLQRKDCC	0.010
8	QRKDCCADY	0.005
3	SDDCLQRKD	0.003
6	CLQRKDCCA	0.001
4	DDCLQRKDC	0.001
7	LQRKDCCAD	0.000

Table VIII-V3-HLA-A1-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
3	NVESCPEGK	18
6	SCPGGKPEA	0.02
5	ESCPGGKPE	0.015
9	GGKPEALWV	0.013
2	TNVESCPGG	0.005
7	CPGGKPEAL	0.003
1	PTNVESCPG	0.003
4	VESCPGGKP	0
8	PGGKPEALW	0

Table VIII-V4-HLA-A1-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
2	YLPTFETPI	0.01
1	TYLPTFETP	0.001

Table IX-V1-HLA-A1-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		

Start	Subsequence	Score
798	VSEILQLKTY	67.500
711	NTDVPIPTHY	62.500
326	MTDYFPRINF	62.500
781	DVELLTGLDF	45.000
448	EVEPFENIEV	45.000
220	GSFPSIYMPY	37.500
381	TPDLPKRLHY	31.250
310	LADHGMDQTY	25.000
686	VSGPIFDYNY	15.000
213	GSEVAINGSF	13.500
402	FVDQQWLAVR	10.000
559	NVDHCLLYHR	10.000
613	PSESQKCSFY	6.750
525	QLEQVNQMLN	4.500
567	HREYVSGFGK	4.500
492	HAEEVSKFSV	4.500
536	TQEEITATVK	2.700
684	NVVS GPIFDY	2.500
698	HFDAPDEITK	2.500
284	VVDHAFGMLM	2.500
658	YEEFRKMWDY	2.250
446	KTEVEPFENI	2.250
762	KPEALWVEER	2.250
742	DVLPFIIPHR	2.000
119	TLMPNINKLK	2.000
767	WVEERFTAH	1.800
53	CSCSDDCLQK	1.500
641	TSDSQYDALI	1.500
104	FSMDGFRAEY	1.500
359	NSEEIVRNLS	1.350
349	HNIPHDFFSF	1.250
328	DYFPRINFFY	1.250
601	VPDCLRADVR	1.250
95	GFDLPPVILF	1.250
157	YPESHGIIDN	1.125
654	LVP MYEEFRK	1.000
653	NLVP MYEEFR	1.000
649	LITSNLVPMY	1.000
47	RLEASLCSCS	0.900
29	CVESTRIWMC	0.900
5	GLENCRCDA	0.900
110	RAEYLYTWD	0.900
79	WLEENCDAQ	0.900
431	SMEAIFLAHG	0.900
356	FSFNSEEIVR	0.750
785	LTGLDFYQDK	0.500
623	LADKNITHGF	0.500
83	NCDTAQQSQ	0.500
66	CADYKSVCQG	0.500
541	TATVKVNLP	0.500
10	RCDVACKDRG	0.500
794	KVQPVSEILQ	0.500
787	GLDFYQDKVQ	0.500
163	IIDNNMYDVN	0.500
512	SLDCFCPHLQ	0.500
461	MCDLLRIQPA	0.500
217	AINGSFPSIY	0.500
97	DLPPVILFSM	0.500

Table IX-V1-HLA-A1-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
292	LMEGLKQRNL	0.450
487	FYEP SHAEEV	0.450
92	CPEGFDLPPV	0.450
43	CGETRLEASL	0.450
235	FEERISTLLK	0.450
454	NIEVYNLMCD	0.450
261	FEEDSSGHA	0.450
758	CPEGKPEALW	0.450
338	MYEGPAPRIR	0.450
316	DQTYCNKMEY	0.375
756	ESCEGKPEA	0.300
263	EPDSSGHAGG	0.250
740	WLDVLPFIIP	0.250
586	YTPVQLGDS	0.250
314	GMDQTYCNKM	0.250
479	LNHLLKVPHY	0.250
139	AMYPTKTFPN	0.250
168	MYDVNLKNF	0.250
25	FEDTCVEST	0.250
377	KPYLTPDLPK	0.250
105	SMDGFRAEYL	0.250
591	LGDSPLPPT	0.250
166	NNMYDVNLNK	0.250
144	KTFPNHYTIV	0.250
371	KPDQHFKPYL	0.250
690	IFDYN YDGHF	0.250
426	NNEFRSMEAI	0.225
74	QGETSWLEEN	0.225
360	SEEIVRNLS	0.225
677	ATERNGVNVV	0.225
570	YVSGFGKAMR	0.200
129	TCGIHSKYMR	0.200
721	FVLTSCCKNK	0.200
682	GVNVVSGPIF	0.200
196	WLTAMYQGLK	0.200
434	AIFLAHGPSF	0.200
54	SCSDDCLQKK	0.200
478	SLNHLLKVPF	0.200
437	LAHGPSFKEK	0.200
250	KAERPRFYTM	0.180
231	GSVPFEERIS	0.150
192	GQPMWLTAMY	0.150

Table IX-V2-HLA-A1-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
1	CSCSDDCLQR	0.750
2	SCSDDCLQRK	0.200
3	CSDDCLQRKD	0.075
10	RKDCCADYKS	0.050
4	SDDCLQRKDC	0.025
6	DCLQRKDCCA	0.010
8	LQRKDCCADY	0.002
9	QRKDCCADYK	0.001
5	DDCLQRKDCC	0.001
7	CLQRKDCCAD	0.000

Table IX-V3-HLA-A1-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
6	ESCPGGKPEA	0.3
4	NVESCPGGKP	0.09
3	TNVESCPGGK	0.05
7	SCPGGKPEAL	0.01
2	PTNVESCPGG	0.005
8	CPGGKPEALW	0.005
1	RPTNVESCPG	0.003
10	GGKPEALWVE	0
9	PGGKPEALWV	0
5	VESCPGGKPE	0

Table IX-V4-HLA-A1-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	TYLPTFETPI	0.005
1	KTYLPTFETP	0.003

Table X-V1-HLA-A0201-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position		

plus eight.		
Start	Subsequence	Score
663	KMWDYFHSV	11367.476
563	CLLYHREYV	693.538
325	YMTDYFPRI	270.002
807	YLPTFETTI	182.365
119	TLMPNINKL	181.794
196	WLTAMYQGL	147.401
459	NLMCDLLRI	88.783
113	YLYTWDTLM	73.129
547	NLPFGRPRV	69.552
765	ALWVEERFT	68.037
740	WLDVLPFII	45.649
238	RISTLLKWL	37.157
155	GLYPESHGI	33.385
512	SLDCFCPHL	32.471
579	RMPMWSSYT	29.601
199	AMYQGLKAA	26.408
395	RIDKVHLFV	21.039
402	FVDQQWLAV	19.036
524	TQLEQVNQM	17.575
747	IIPHRPTNV	16.258
163	IIDNNMYDV	14.957
400	HLFVDQQWL	14.781
787	GLDFYQDKV	13.632
90	SQCPEGFDL	12.562
693	YNYDGHFDA	11.352
283	QVVDHAFGM	10.337
300	NLHNCVNII	9.838
555	VLQKNVDHC	9.518
532	MLNLTQEEI	8.691
570	YVSGFGKAM	7.599
802	LQLKTYLPT	7.129
500	SVCGFANPL	7.103
805	KTYLPTFET	6.723
430	RSMEAIFLA	6.563
277	RVIKALQVV	5.739
171	VNLNKNFSL	5.087
59	CLQKKDCCA	4.968
534	NLTQEEITA	4.968
383	DLPKRLHYA	4.713
800	EILQLKTYL	4.483
5	GLENCRCDV	4.451
452	FENIEVYNL	4.395
307	IILLADHGM	4.297
714	VPIPTHYFV	4.245
477	GSLNHLLKV	3.864
111	AEYLYTWDT	3.478
488	YEPShAEV	3.048
79	WLEENCDA	2.938
580	MPMWSSYTV	2.856
30	VESTRIWMC	2.833
217	AINGSFPSI	2.726
649	LITSNLVPM	2.671
51	SLCSCSDDC	2.434
670	SVLLIKHAT	2.413
449	VEPFENIEV	2.299
380	LTPDLPKRL	2.068
21	CCWDFEDTC	2.055

Table X-V1-HLA-A0201-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
144	KTFPNHYTI	1.876
297	KQRNLHNCV	1.876
240	STLLKWLDL	1.866
536	TQEEITATV	1.850
535	LTQEEITAT	1.659
356	FSFNSEEIV	1.552
528	QVNQMLNLT	1.500
583	WSSYTPVQL	1.475
622	YLADKNITH	1.405
525	QLEQVNQML	1.367
794	KVQPVSEIL	1.314
72	VCQGETSWL	1.304
467	IQPAPNNGT	1.284
233	VPFEERIST	1.255
192	GQPMWLTAM	1.159
456	EVYNLMCDL	1.032
131	GIHSKYMRA	1.025
280	KALQVVDHA	1.007
291	MLMEGLKQR	0.884
427	NEFRSMEAI	0.846
784	LLTGLDFYQ	0.808
447	TEVEPFENI	0.774
715	PIPTHYFVV	0.750
250	KAERPRFYT	0.740
98	LPPVILFSM	0.735
47	RLEASLCSC	0.731
330	FPRINFFYM	0.687
474	GTHGSLNHL	0.682
337	YMYEGPAPR	0.650
274	VSARVIKAL	0.545
521	QNSTQLEQV	0.512
540	ITATVKVNL	0.504
493	AEVSKFSV	0.502
270	AGGPVSARV	0.454
665	WDYFHSVLL	0.437
790	FYQDKVQPV	0.419
44	GETRLEASL	0.415
190	WHGQPMWLT	0.411
656	PMYEEFRKM	0.394
436	FLAHGPSFK	0.377
527	EQVNQMLNL	0.374
115	YTWDTLMPN	0.373
708	HLANTDVPI	0.355

Table X-V2-A0201-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position

for each peptide is the start position plus eight.

Start	Subsequence	Score
6	CLQRKDCCA	4.968
5	DCLQRKDCC	0.004
4	DDCLQRKDC	0.001
1	SCSDDCLQR	0.000
2	CSDDCLQRK	0.000
7	LQRKDCCAD	0.000
9	RKDCCADYK	0.000
3	SDDCLQRKD	0.000
8	QRKDCCADY	0.000

Table X-V3-HLA-A2-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
9	GGKPEALWV	0.087
7	CPGGKPEAL	0.068
6	SCPGGKPEA	0.032
2	TNVESCPGG	0.002
4	VESCPGGKP	0
1	PTNVESCPG	0
3	NVESCPGGK	0
8	PGGKPEALW	0
5	ESCPGGKPE	0

Table X-V4-HLA-A2-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 9 each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
2	YLPTFETPI	182.365
1	TYLPTFETP	0

Table XI-V1-HLA-A0201-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
663	KMWDYFHSV	2862.980
337	YMYEGPAPRI	454.740
765	ALWVEERFTA	239.160
102	ILFSMDGFRA	181.243
632	FLYPPASNRT	109.693
379	YLTPDLPKRL	98.267

WO 03/040340

148

PCT/US02/36002

Table XI-V1-HLA-A0201-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
162	GIIDNNMYDV	90.183
309	LLADHGMDQT	58.537
115	YTWDTLMPNI	52.169
579	RMPMWSSYTV	50.232
746	FIIPHRPTNV	43.992
555	VLQKNVDHCL	36.316
407	WLAVRSKSN	34.279
34	RIWMCNKFR	32.884
524	TQLEQVNQML	32.857
600	TVPDCLRADV	24.952
801	ILQLKTYLPT	19.003
199	AMYQGLKAAT	17.222
534	NLTQEEITAT	17.140
105	SMDGFRAEYL	16.632
21	CCWDFEDTCV	15.450
531	QMLNLTQEEI	13.661
520	LQNSTQLEQV	13.511
614	SESQKCSFY	13.251
648	ALITSNLVPM	11.426
51	SLCSCSDDCL	10.468
387	RLHYAKNVRI	10.433
71	SVCQGETSWL	10.281
120	LMPNINKLKT	9.149
300	NLHNCVNIL	8.759
795	VQPVSEILQL	8.469
233	VPFEERISTL	8.271
144	KTFPNHYTIV	7.693
4	RGLENCRCDV	6.887
535	LTQEEITATV	6.733
97	DLPPVILFSM	4.970
282	LQVVDHAFGM	4.966
400	HLFVDQQWLA	4.687
767	WVEERFTHAI	4.187
371	KPDQHFKPYL	4.080
224	SIYMPYNGSV	3.978
786	TGLDFYQDKV	3.375
622	YLADKNITHG	3.233
207	ATYFWPGSEV	3.091
546	VNLPFGRPRV	2.856
714	VPIPTHYFVV	2.753
458	YNLMCDLLRI	2.666
532	MLNLTQEEIT	2.545
713	DVPIPTHYFV	2.510
499	FSVCGFANPL	2.438
799	SEILQLKTYL	2.285
291	MLMEGLKQRN	1.922
283	QVVDHAFGML	1.893
136	YMRAMYPTKT	1.882
554	RVLQKNVDHC	1.813
155	GLYPESHGII	1.779

Table XI-V1-HLA-A0201-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
547	NLPFGRPRVL	1.752
455	IEVYNLMCDL	1.624
526	LEQVNQMLNL	1.624
167	NMYDVNLNKN	1.624
484	KVPFYEPSHA	1.521
314	GMDQTYCNKM	1.435
123	NINKLKTCGI	1.435
317	QTYCNKMEYM	1.369
128	KTCGIHSKYM	1.328
508	LPTESLDCFC	1.243
358	FNSEEIVRNL	1.210
284	VVDHAFGMLM	1.123
111	AEYLYTWDTL	1.107
805	KTYLPTFETT	1.079
466	RIQPAPNNGT	1.025
139	AMYPTKTFPN	0.999
329	YFPRINFFYM	0.962
640	RTSDSQYDAL	0.894
170	DVNLNKNFSL	0.813
89	QSQCPEGFDL	0.809
28	TCVESTRIWM	0.731
716	IPTHYFWLT	0.723
306	NIILLADHGM	0.683
242	LLKWLDLPKA	0.680
662	RKMWDYFHSV	0.679
504	FANPLPTESL	0.669
452	FENIEVYNLM	0.667
232	SVPFEERIST	0.652
425	YNNEFRSMEA	0.612
564	LLYHREYVSG	0.608
511	ESLDCFCPHL	0.603
198	TAMYQGLKAA	0.587
401	LFVDQQWLAV	0.572
399	VHLFVDQQWL	0.513
63	KDCCADYKSV	0.507
665	WDYFHSVLLI	0.491
392	KNVRIDKVHL	0.488
29	CVESTRIWMC	0.480
216	VAINGSFPSI	0.468
440	GPSFKEKTEV	0.454
773	TAHIARVRDV	0.444
339	YEGPAPRIRA	0.444
646	YDALITSNLV	0.444
610	RVPPSESQKC	0.435

Table XI-V2-A0201-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10		

amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	CLQRKDCCAD	0.015
6	DCLQRKDCCA	0.009
4	SDDCLQRKDC	0.003
8	LQRKDCCADY	0.001
2	SCSDDCLQRK	0.001
5	DDCLQRKDCC	0.000
1	CSCSDDCLQR	0.000
10	RKDCCADYKS	0.000
3	CSDDCLQRKD	0.000
9	QRKDCCADYK	0.000

Table XI-V3-HLA-A2-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	SCPGGKPEAL	0.068
9	PGGKPEALWV	0.055
6	ESCPGGKPEA	0.002
5	VESCPGGKPE	0
8	CPGGKPEALW	0
1	RPTNVESCPG	0
3	TNVESCPGGK	0
10	GGKPEALWVE	0
2	PTNVESCPGG	0
4	NVESCPGGKP	0

Table XI-V4-HLA-A2-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	TYLPTFETPI	0.02
1	KTYLPTFETP	0.002

Table XII-V1-HLA-A3-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
167	NMYDVNLNKN	300.000
314	GMDQTYCNK	60.000
632	FLYPPASNR	45.000

Table XII-V1-HLA-A3-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
242	LLKWLDLPK	40.000
337	YMYEGPAPR	30.000
663	KMWDYFHSV	27.000
436	FLAHGPSFK	20.000
136	YMRAMYPTK	20.000
105	SMDGFRAEY	18.000
120	LMPNINKLK	15.000
155	GLYPESHGI	13.500
379	YLTPDLPKR	9.000
803	QLKTYLPTF	9.000
743	VLPFIIPHR	9.000
291	MLMEGLKQR	6.750
245	WLDLPKAER	6.000
322	KMEYMTDYF	6.000
102	ILFSMDGFR	6.000
672	LLIKHATER	6.000
325	YMTDYFPRI	5.400
653	NLVPMYEEF	4.500
32	STRIWMCNK	4.500
685	VVSGPIFDY	4.050
387	RLHYAKNVR	4.000
610	RVPPSESQK	3.000
400	HLFVDQQWL	3.000
281	ALQVVDHAF	3.000
113	YLYTWD TLM	3.000
459	NLMCDLLRI	2.700
618	KCSFYLADK	2.700
783	ELLTGLDFY	2.700
119	TLMPNINKL	2.025
144	KTFPNHYTI	2.025
317	QTYCNKMEY	2.000
754	NVESCPEGK	2.000
363	IVRNLSCRK	2.000
350	NIPHDFFSF	1.800
431	SMEAIFLAH	1.800
203	GLKAATYFW	1.800
512	SLDCFCPHL	1.800
300	NLHNCVNII	1.800
807	YLPTFETTI	1.800
740	WLDVLPFII	1.800
787	GLDFYQDKV	1.800
722	VVLTSCKNK	1.500
128	KTCGIHSKY	1.350
118	DTLMPNINK	1.350
654	LVP MYEEFR	1.200
34	RIWMCNKFR	1.000
272	GPVSARVIK	0.900
655	VPMYEEFRK	0.900
405	QQWLAVRSK	0.900
197	LTAMYQGLK	0.900

Table XII-V1-HLA-A3-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
568	REYVSGFGK	0.900
525	QLEQVNQML	0.900
199	AMYQGLKAA	0.750
797	PVSEILQLK	0.675
532	MLNLTQEEI	0.600
5	GLENCRCDV	0.600
8	NCRCDVACK	0.600
564	LLYHREYVS	0.600
555	VLQKNVDHC	0.600
384	LPKRLHYAK	0.600
708	HLANTDVPI	0.600
650	ITSNLVPMY	0.600
196	WLTAMYQGL	0.600
390	YAKNVRIDK	0.600
542	ATVKVNLPF	0.450
101	VILFSMDGF	0.450
794	KVQPVSEIL	0.405
534	NLTQEEITA	0.400
54	SCSDDCLQK	0.400
622	YLADKNITH	0.400
805	KTYLPTFET	0.338
79	WLEENCDTA	0.300
579	RMPMWSSYT	0.300
47	RLEASLCSC	0.300
563	CLLYHREYV	0.300
577	AMRMPMWSS	0.270
362	EIVRNLSCR	0.270
217	AINGSFPSI	0.270
482	LLKVPFYEP	0.270
500	SVCGFANPL	0.270
269	HAGGPVSAR	0.270
126	KLKTCGIHS	0.240
474	GTHGSLNHL	0.203
51	SLCSCSDDC	0.200
258	TMYFEEPDS	0.200
547	NLPFGRPRV	0.200
59	CLQKKDCCA	0.200
550	FGRPRVLQK	0.180
290	GMLMEGLKQ	0.180
484	KVPFYEPSH	0.180
371	KPDQHFKPY	0.180
55	CSDDCLQKK	0.150
139	AMYPTKTFP	0.150
450	EPFENIEVY	0.135
481	HLLKVPFYE	0.135
241	TLLKWLDLP	0.135
456	EVYNLMCDL	0.135

Table XII-V2-HLA-A3-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
6	CLQRKDCCA	0.200
2	CSDDCLQRK	0.150
1	SCSDDCLQR	0.080
9	RKDCCADYK	0.020
8	QRKDCCADY	0.004
5	DCLQRKDCC	0.001
7	LQRKDCCAD	0.001
4	DDCLQRKDC	0.000
3	SDDCLQRKD	0.000

Table XII-V3-HLA-A3-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
3	NVESC PG GK	0.6
7	CPGGKPEAL	0.009
6	SCPGGKPEA	0.003
9	GGKPEALWV	0.002
1	PTNVESCPG	0
2	TNVESCPGG	0
8	PGGKPEALW	0
4	VESCPGGKP	0
5	ESCPGGKPE	0

Table XII-V4-HLA-A3-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
2	YLPTFETPI	1.8
1	TYLPTFETP	0

Table XIII-V1-HLA-A3-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
126	KLKTCGIHSK	90.000

WO 03/040340

PCT/US02/36002

150

Table XIII-V1-HLA-A3-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
241	TLLKWLDLPK	60.000
119	TLMPNINKLK	33.750
663	KMWDYFHSVL	27.000
653	NLVPMYEEFR	27.000
383	DLPKRLHYAK	18.000
196	WLTAMYQGLK	18.000
290	GMLMEGLKQR	13.500
377	KPYLTPDLPK	9.000
337	YMYEGPAPRI	6.750
654	LVPMYEEFRK	6.000
671	VLLIKHATER	6.000
155	GLYPESHGII	4.050
684	NVSGPIFDY	4.050
577	AMRMPMWSSY	4.000
102	ILFSMDGFRA	3.000
785	LTGLDFYQDK	3.000
765	ALWVEERFTA	3.000
400	HLFVDQQWLA	3.000
478	SLNHLLKVPF	2.000
226	YMPYNGSVPF	2.000
559	NVDHCLLYHR	1.800
314	GMDQTYCNKM	1.800
300	NLHNCVNIL	1.800
402	FVDQQWLAVR	1.800
217	AINGSFPSIY	1.800
721	FVVLTSCKNK	1.500
220	GSFPSIYMPY	1.350
543	TVKVNLPFGR	1.200
649	LITSNLVPMY	1.200
330	FPRINFFYMY	1.080
762	KPEALWVEER	1.080
434	AIFLAHGPSF	1.000
531	QMLNLTQEEI	0.900
105	SMDGFRAEYL	0.900
295	GLKQRNLHNC	0.900
555	VLQKNVDHCL	0.900
362	EIVRNLSRK	0.900
536	TQEEITATVK	0.900
632	FLYPPASNRT	0.750
796	QPVSEILQLK	0.675
97	DLPPVILFSM	0.608
742	DVLPIIPHR	0.608
51	SLCSCSDDCL	0.600
682	GVNVVSGPIF	0.600
579	RMPMWSSYTV	0.600
5	GLENCRCDA	0.600
387	RLHYAKNARI	0.600
247	DLPKAERPRF	0.600
570	YVSGFGKAMR	0.600
393	NVRIDKVHLF	0.600

Table XIII-V1-HLA-A3-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
199	AMYQGLKAAT	0.500
139	AMYPTKTFPN	0.450
648	ALITSNLVPM	0.450
379	YLTPDLPKRL	0.450
437	LAHGSPFKEK	0.450
802	LQLKTYLPTF	0.405
162	GIIDNNMYDV	0.405
481	HLLKVPFYEP	0.405
446	KTEVEPFENI	0.405
545	KVNLPFGRPR	0.360
192	GQPMWLTAMY	0.360
34	RIWMCNKFRC	0.300
326	MTDYFPRINF	0.300
711	NTDVPIPTHY	0.300
54	SCSDDCLQKK	0.300
507	PLPTESLDCF	0.300
136	YMRAMYPTKT	0.300
242	LLKWLDLPKA	0.300
686	VSGPIFDYNY	0.270
784	LLTGLDFYQD	0.270
767	WVEERFTAH	0.270
172	NLNKNFSLSS	0.240
656	PMYEEFRKMW	0.225
115	YTWDTLMPNI	0.225
805	KTYLPTFETT	0.225
144	KTFPNHYTIV	0.225
366	NLSRKPDQH	0.200
801	ILQLKTYLPT	0.200
356	FSFNSEEIVR	0.200
53	CSCSDDCLQK	0.200
120	LMPNINKLKT	0.200
258	TMYFEEDSS	0.200
368	SCRKPDQHFK	0.200
113	YLYTWDTLMP	0.200
166	NNMYDVNLNK	0.180
495	EVSKEFVCGF	0.180
740	WLDVLPFIIP	0.180
323	MEYMTDYFPR	0.180
563	CLLYHREYVS	0.180
101	VILFSMDGFR	0.180
203	GLKAATYFWP	0.180
322	KMEYMTDYFP	0.180
534	NLTQEEITAT	0.150
167	NMYDVNLNKN	0.150
309	LLADHGMDQT	0.150
596	PLPPTVPDCL	0.135
280	KALQVVDHAF	0.135
31	ESTRIWMCNK	0.135
474	GTHGSLNHLL	0.135

Table XIII-V2-HLA-A3-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	SCSDDCLQRK	0.300
8	LQRKDCCADY	0.120
1	CSCSDDCLQR	0.040
7	CLQRKDCCAD	0.020
9	QRKDCCADYK	0.020
6	DCLQRKDCCA	0.001
10	RKDCCADYKS	0.000
4	SDDCLQRKDC	0.000
5	DDCLQRKDCC	0.000
3	CSDDCLQRKD	0.000

Table XIII-V3-HLA-A3-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
3	TNVECPGGK	0.027
7	SCPGGKPEAL	0.009
8	CPGGKPEALW	0.005
4	NVECPGGKPE	0.001
6	ESCPGGKPEA	0
10	GGKPEALWVE	0
1	RPTNVECPG	0
2	PTNVECPGG	0
9	PGGKPEALWV	0
5	VECPGGKPE	0

Table XIII-V4-HLA-A3-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
1	KTYLPTFETP	0.045
2	TYLPTFETPI	0.004

Table XIV-V1-HLA-A11-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		

Start	Subsequence	Score
610	RVPPSESQK	6.000
363	IVRNLSCKR	2.000
754	NVESCPEGK	2.000
167	NMYDVNLNK	1.600
722	VVLTSCKNK	1.500
314	GMDQTYCNK	1.200
655	VPMYEEFRK	1.200
568	REYVSGFGK	1.080
32	STRIWMCNK	1.000
197	LTAMYQGLK	1.000
272	GPVSARVIK	0.900
118	DTLMPNINK	0.900
242	LLKWLDLPK	0.800
618	KCSFYLDK	0.600
390	YAKNVRIDK	0.400
654	LVPMYEEFR	0.400
136	YMRAMYPTK	0.400
384	LPKRLHYAK	0.400
436	FLAHGPSFK	0.400
54	SCSDDCLQK	0.400
667	YFHSVLLIK	0.400
720	YFVVLTSCK	0.300
387	RLHYAKNVR	0.240
34	RIWMCNKFR	0.240
797	PVSEILQLK	0.200
8	NCRCDVACK	0.200
120	LMPNINKLK	0.200
632	FLYPPASNR	0.160
337	YMYEGPAPR	0.160
102	ILFSMDGFR	0.160
324	EYMTDYFPR	0.144
405	QQWLAVRSK	0.120
378	PYLTPDLPK	0.120
144	KTFPNHYTI	0.120
672	LLIKHATER	0.120
554	RVLQKNVDH	0.090
283	QVVDHAFGM	0.090
277	RVIKALQVV	0.090
743	VLPIIIPHR	0.080
291	MLMEGLKQR	0.080
357	SFNSEEIVR	0.080
379	YLTPDLPKR	0.080
245	WLDLPKAER	0.080
62	KKDCCADYK	0.060
794	KVQPVSEIL	0.060
537	QEEITATVK	0.060
682	GVNVVSGPI	0.060
484	KVPFYEPSH	0.060
10	RCDVACKDR	0.060
685	VVSGPIFDY	0.060
640	RTSDSQYDA	0.060
2	SFRGLENCR	0.040
476	HGSLNHLLK	0.040
289	FGMLMEGLK	0.040
559	NVDHCLLYH	0.040
317	QTYCNKMEY	0.040
699	FDAPDEITK	0.040
29	CVESTRIWM	0.040

Table XIV-V1-HLA-A11-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
550	FGRPRVLQK	0.040
402	FVDQQWLAV	0.040
269	HAGGPVSAR	0.040
236	EERISTLLK	0.036
362	EIVRNLSCKR	0.036
786	TGLDFYQDK	0.030
240	STLLKWLDL	0.030
128	KTCGIHSKY	0.030
474	GTHGSLNHL	0.030
542	ATVKVNLPF	0.030
458	YNLMCDLLR	0.024
663	KMWDFHHSV	0.024
131	GIHSKYMRA	0.024
345	RIRAHNIPH	0.024
155	GLYPESHGI	0.024
203	GLKAATYFW	0.024
395	RIDKVHLFV	0.024
393	NVRIDKVHL	0.020
284	VVDHAFGML	0.020
500	SVCGFANPL	0.020
772	FTAHARVR	0.020
369	CRKPDQHFV	0.020
71	SVCQGETSW	0.020
55	CSDDCLQKK	0.020
127	LKTCGIHSK	0.020
767	WVEERFTAH	0.020
174	NKNFSLSSK	0.020
544	VKVNLFPGR	0.018
742	DVLPFIIPH	0.018
805	KTYLPTFET	0.018
192	GQPMWLTAM	0.018
90	SQCPEGFDL	0.018
297	KQRNLHNCV	0.018
459	NLMCDLLRI	0.016
130	CGIHSKYMR	0.012
5	GLENCRCDV	0.012
42	RCGETRLEA	0.012
253	RPRFYTMFY	0.012
740	WLDVLPFII	0.012
787	GLDFYQDKV	0.012
616	SQKCSFYLA	0.012
350	NIPHDFFSF	0.012

Table XIV-V2-HLA-A11-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
1	SCSDDCLQR	0.080
9	RKDCCADYK	0.060
2	CSDDCLQRK	0.020
6	CLQRKDCCA	0.004
7	LQRKDCCAD	0.001
8	QRKDCCADY	0.000
5	DCLQRKDCC	0.000
4	DDCLQRKDC	0.000
3	SDDCLQRKD	0.000

Table XIV-V3-HLA-A11-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
3	NVESCPCGGK	2
7	CPGGKPEAL	0.002
6	SCPGGKPEA	0.002
9	GGKPEALWV	0.001
1	PTNVESCPG	0
2	TNVESCPGG	0
4	VESCPGGKP	0
8	PGGKPEALW	0
5	ESCPGGKPE	0

Table XIV-V4-HLA-A11-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
2	YLPTFETPI	0.004
1	TYLPTFETP	0.001

Table XV-V1-HLA-A11-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
654	LVPMYEEFRK	6.000

Table XV-V1-HLA-A11-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
377	KPYLTPDLPK	2.400
135	KYMRAMYPTK	2.400
721	FVVLTSCKNK	1.500
543	TVKVNLPFGR	1.200
241	TLLKWLDLPK	1.200
126	KLKTCGIHSK	1.200
785	LTGLDFYQDK	1.000
559	NVDHCLLYHR	0.800
719	HYFVVLTSCK	0.800
389	HYAKNVRIDK	0.800
536	TQEEITATVK	0.600
666	DYFHSVLLIK	0.480
196	WLTAMYQGLK	0.400
402	FVDQQWLAVR	0.400
570	YVSGFGKAMR	0.400
119	TLMPNINKLK	0.400
698	HFDAPDEITK	0.400
435	IFLAHGPFK	0.300
796	QPVSEILQLK	0.300
383	DLPKRLHYAK	0.240
54	SCSDDCLQKK	0.200
288	AFGMLMEGLK	0.200
368	SCRKPDQHFK	0.200
742	DVLPFIIPHR	0.180
631	GFLYPASNR	0.180
290	GMLMEGLKQR	0.180
362	EIVRNLSCKR	0.180
336	FYMYEGPAPR	0.160
457	VYNLMCDLLR	0.160
166	NNMYDVNLNK	0.160
671	VLLIKHATER	0.120
235	FEERISTLLK	0.120
545	KVNLPFGRPR	0.120
762	KPEALWVEER	0.120
653	NLVPMEEFR	0.120
101	VILFSMDGFR	0.120
437	LAHGPFKEK	0.100
684	NVVGPIFDY	0.090
129	TCGIHSKYMR	0.080
323	MEYMTDYFPR	0.072
567	HREYVSGFGK	0.060
753	TNVECPGK	0.060
271	GGPVSAVRIK	0.060
682	GVNVSGPIF	0.060
489	EPSHAEVSK	0.060
484	KVPFYEPSHA	0.060
144	KTFPNHYTIV	0.060
398	KVHLFVDQQW	0.060
475	THGSLNHLLK	0.040
117	WDTLMPNINK	0.040

Table XV-V1-HLA-A11-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
173	LNKNFSLSSK	0.040
313	HGMDQTYCNK	0.040
597	LPPTVPDCLR	0.040
284	VVDHAFGMLM	0.040
53	CSCSDDCLQK	0.040
601	VPDCLRADVR	0.040
549	PFGPRVLQK	0.040
162	GIIDNNMYDV	0.036
609	VRVPPSESQK	0.030
283	QVVDHAFGML	0.030
640	RTSDSQYDAL	0.030
446	KTEVEPFENI	0.030
474	GTHGSLNHLL	0.030
282	LQVVDHAFGM	0.027
421	GNHGYNNEFR	0.024
663	KMWDFHSL	0.024
765	ALWVEERFTA	0.024
579	RMPMWSSYTV	0.024
155	GLYPESHGII	0.024
102	ILFSMDGFRA	0.024
152	IVTGLYPESH	0.020
71	SVCQGETSWL	0.020
304	CVNIILLADH	0.020
207	ATYFWPGSEV	0.020
767	WVEERFTAH	0.020
197	LTAMYQGLKA	0.020
600	TVPDCLRADV	0.020
115	YTWDTLMPNI	0.020
617	QKCSFYLADK	0.020
326	MTDYFPRINF	0.020
317	QTYCNKMEYM	0.020
61	QKKDCCADYK	0.020
393	NVRIDKVHLF	0.020
386	KRLHYAKNVR	0.018
244	KWLDLPKAER	0.018
192	GQPMWLTAMY	0.018
272	GPVSARVIKA	0.018
170	DVNLNKNFSL	0.018
404	DQQWLAVRSK	0.018
400	HLFVDQQWLA	0.016
356	FSFNSEEIVR	0.016
128	KTCGIHSKYM	0.015
378	PYLTPDLPKR	0.012
771	RFTAHARVR	0.012
268	GHAGGPVSAR	0.012
250	KAERPRFYTM	0.012
108	GFRAEYLYTW	0.012
573	GFGKAMRMPM	0.012
387	RLHYAKNVRI	0.012

Table XV-V2-HLA-A11-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	SCSDDCLQRK	0.200
9	QRKDCCADYK	0.020
1	CSCSDDCLQR	0.008
8	LQRKDCCADY	0.006
6	DCLQRKDCCA	0.001
7	CLQRKDCCAD	0.000
10	RKDCCADYKS	0.000
4	SDDCLQRKDC	0.000
5	DDCLQRKDCC	0.000
3	CSDDCLQRKD	0.000

Table XV-V3-HLA-A11-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
3	TNVECPGGK	0.06
8	CPGGKPEALW	0.002
7	SCPGGKPEAL	0.002
4	NVECPGGKPK	0.002
1	RPTNVECPG	0.001
10	GGKPEALWVE	0
2	PTNVECPGG	0
6	ESCPGGKPEA	0
9	PGGKPEALWV	0
5	VESCPGGKPE	0

Table XV-V4-HLA-A11-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	TYLPTFETPI	0.006
1	KTYLPTFETP	0.006

Table XVI-V1-HLA-A24-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		

WO 03/040340

PCT/US02/36002

153

Start	Subsequence	Score
112	EYLYTWDTL	300.000
457	VYNLMCDLL	300.000
328	DYFPRINFF	144.000
156	LYPESHGII	90.000
338	MYEGPAPRI	75.000
666	DYFHSVLLI	50.000
424	GYNNEFRSM	45.000
40	KFRCGETRL	40.000
318	TYCNKMEYM	25.000
288	AFGMLMEGL	24.000
794	KVQPVEIL	20.160
95	GFDLPPVIL	20.000
435	IFLAHGPSF	15.000
135	KYMRAMYPT	15.000
633	LYPPASNRT	10.800
790	FYQDKVQPV	10.800
806	TYLPTFETT	10.800
359	NSEEIVRNL	10.080
525	QLEQVNQML	10.080
660	EFRKMWDYF	10.000
428	EFRSMEAIF	10.000
569	EYVSGFGKA	9.900
238	RISTLLKWL	9.600
119	TLMPNINKL	9.504
657	MYEEFRKMW	9.000
621	FYLADKNIT	9.000
225	IYMPYNGSV	9.000
200	MYQGLKAAT	9.000
380	LTPDLPKRL	8.640
597	LPPTVPDCL	8.400
140	MYPTKTFPN	7.500
800	EILQLKTYL	7.200
149	HYTIVTGLY	7.000
719	HYFVVLTS	7.000
701	APDEITKHL	6.720
168	MYDVNLNKN	6.600
736	NCPGWLDVL	6.000
259	MYFEEDSS	6.000
138	RAMYPTKTF	6.000
758	CPEGKPEAL	6.000
733	TPENCPGWL	6.000
527	EQVNQMLNL	6.000
240	STLLKWLDL	6.000
615	ESQKCSFYL	6.000
165	DNNMYDVNL	6.000
72	VCQGETSWL	6.000
322	KMEYMTDYF	6.000
645	QYDALITSN	6.000
796	QPVEILQL	6.000
171	VNLNKNFSL	6.000
505	ANPLPTESL	6.000
556	LQKNVDHCL	5.600
347	RAHNIPHDF	5.600
540	ITATVKVNL	5.600
274	VSARVIKAL	5.600
208	TYFWPGSEV	5.500
355	FFSFNSEEI	5.500
620	SFYLADKNI	5.000

Table XVI-V1-HLA-A24-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
474	GTHGSLNHL	4.800
456	EVYNLMCDL	4.800
716	IPTHYFVVL	4.800
196	WLTAMYQGL	4.800
90	SQCPEGFDL	4.800
500	SVCGFANPL	4.800
400	HLFVDQQWL	4.800
284	VVDHAFGML	4.800
776	IARVRDVEL	4.400
542	ATVKVNLPF	4.200
764	EALWVEERF	4.200
281	ALQVVDHAF	4.200
189	WWHGQPMWL	4.000
512	SLDCFCPHL	4.000
393	NVRIDKVHL	4.000
583	WSSYTVPQL	4.000
302	HNCVNIILL	4.000
664	MWDYFHSLV	4.000
548	LPFGRPRVL	4.000
52	LCSCSDDCL	4.000
253	RPRFYTMFY	4.000
641	TSDSQYDAL	4.000
653	NLVPMYEEF	3.960
234	PFEERISTL	3.600
350	NIPHDFFSF	3.600
101	VILFSMDGF	3.000
299	RNLHNCVNI	3.000
713	DVPIPTHYF	3.000
683	VNVVSGPIF	3.000
202	QGLKAATYF	3.000
421	GNHGYNNEF	2.640
739	GWLDVLPFI	2.520
144	KTFPNHYTI	2.400
368	SCRKPDQHF	2.400
479	LNHLLKVPF	2.400
508	LPTESLDCF	2.400
682	GVNVVSGPI	2.100
88	QQSQCPGEF	2.000
227	MPYNGSVPF	2.000
496	VSKFSVCGF	2.000
248	LPKAERPRF	2.000
803	QLKTYLPTF	2.000

Table XVI-V2-HLA-A24-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
5	DCLQRKDCC	0.150
6	CLQRKDCCA	0.150
2	CSDDCLQRK	0.014
8	QRKDCCADY	0.012
1	SCSDDCLQR	0.012
4	DDCLQRKDC	0.010
7	LQRKDCCAD	0.010
9	RKDCCADYK	0.002
3	SDDCLQRKD	0.001

Table XVI-V3-HLA-A24-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
7	CPGGKPEAL	4
6	SCPGGKPEA	0.165
9	GGKPEALWV	0.12
2	TNVECPGG	0.018
3	NVECPGGK	0.015
5	ESCPGGKPE	0.012
8	PGGKPEALW	0.01
1	PTNVECPG	0.002
4	VESCPGGKP	0.001

Table XVI-V4-HLA-A24-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
2	YLPTFETPI	1.5
1	TYLPTFETP	1.08

Table XVII-V1-HLA-A24-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
645	QYDALITSNL	280.000

Table XVII-V1-HLA-A24-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
168	MYDVNLNKNF	120.000
565	LYHREYVSGF	100.000
806	TYLPTFETTI	90.000
324	EYMTDYFPRI	90.000
569	EYVSGFGKAM	37.500
112	EYLYTWD TLM	37.500
375	HFKP YLTPDL	28.800
428	EFRSMEAIFL	20.000
790	FYQDKVQPV S	12.600
524	TQLEQVNQML	12.096
392	KNVRIDKVHL	12.000
700	DAPDEITKHL	10.080
690	IFDYN YDGHF	10.000
95	GFDLPPVILF	10.000
487	FYEP SHAEV	9.900
663	KMWDYFHSVL	9.600
640	RTSDS QYDAL	9.600
633	LYPPASNR TS	9.000
283	QVVDHAFGML	8.640
328	DYFPRINFFY	8.400
280	KALQVVDHAF	8.400
555	VLQKNVDHCL	8.400
371	KPDQHFKPYL	8.000
118	DTLMPNINKL	7.920
200	MYQGLKAATY	7.500
692	DYN YDGHFDA	7.500
757	SCPEGKPEAL	7.200
504	FANPLPTESL	7.200
511	ESLDCFCPHL	7.200
195	MWLTAMYQGL	7.200
732	HTPENCPGWL	7.200
499	FSVCGFANPL	7.200
43	CGETRLEASL	7.200
358	FNSEEIVRNL	6.720
547	NLPFGRPRVL	6.000
170	DVNLNKNFSL	6.000
795	VQPVSEILQL	6.000
89	QSQCPEGFDL	6.000
470	APNNGTHGSL	6.000
209	YFWPGSEVAI	6.000
588	VPQLGDT SPL	6.000
292	LMEGLKQRNL	6.000
379	YLTPDL PKRL	5.760
300	NLHNCVNIL	5.600
539	EITATVKVNL	5.600
68	DYKSVCQGET	5.500
354	DFFSFNSEEI	5.500
234	PFEERISTLL	5.040
114	LYTWD TLM PN	5.000
318	TYCNKMEYMT	5.000

Table XVII-V1-HLA-A24-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
208	TYFWPGSEVA	5.000
585	SYTVPQLGDT	5.000
735	ENCPGWLDVL	4.800
287	HAFGMLMEGL	4.800
473	NGTHGSLNHL	4.800
474	GTHGSLNHLL	4.800
233	VPFEERISTL	4.800
94	EGFDLPPVIL	4.800
329	YFPRINFFYM	4.500
775	HIARVRDVEL	4.400
349	HNIPHDFFSF	4.320
213	GSEVAINGSF	4.200
51	SLCSCSDDCL	4.000
71	SVCQGETSWL	4.000
239	ISTLLKWLDL	4.000
517	CPHLQNSTQL	4.000
582	MWSSYTVPQL	4.000
188	AWWHGQPMWL	4.000
776	IARVRDVELL	4.000
347	RAHNIPHDF	4.000
105	SMDGFRAEYL	4.000
556	LQKNVDHCLL	4.000
456	EVYNLMCDLL	4.000
664	MWDYFHSVLL	4.000
420	GGNHGYNNEF	3.960
478	SLNHLLKVPF	3.600
299	RNLHNCVNII	3.600
446	KTEVEPFENI	3.600
652	SNLVPMYEEF	3.300
247	DLPKAERPRF	3.000
802	LQLKTYLPTF	3.000
87	AQQSQCEPGF	3.000
226	YMPYNGSVPF	3.000
451	PFENIEVYNL	3.000
682	GVNVVSGPIF	3.000
781	DVELLTGLDF	3.000
623	LADKNITHGF	2.800
541	TATVKVNL PF	2.800
32	STRIWMCNKF	2.640
573	GFGKAMRMPM	2.500
367	LSCRKPDQHF	2.400
739	GWLDVLPFII	2.160
681	NGNVVSGPI	2.100
434	AIFLAHG PSF	2.000
393	NVRIDKVHLF	2.000
737	CPGWLDVLPF	2.000
495	EVSKFSVCGF	2.000
326	MTDYFPRINF	2.000
201	YQGLKAATYF	2.000

Table XVII-V2-HLA-A24-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
6	DCLQRKDCCA	0.150
8	LQRKDCCADY	0.100
10	RKDCCADYKS	0.022
3	CSDDCLQRKD	0.016
7	CLQRKDCCAD	0.015
2	SCSDDCLQRK	0.014
4	SDDCLQRKDC	0.010
1	CSCSDDCLQR	0.010
5	DDCLQRKDCC	0.010
9	QRKDCCADYK	0.001

Table XVII-V3-HLA-A24-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7 each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	SCPGGKPEAL	6
6	ESCPGGKPEA	0.132
8	CPGGKPEALW	0.1
1	RPTNVESCPG	0.02
3	TNVESCPGGK	0.018
4	NVESCPGGKP	0.017
10	GGKPEALWVE	0.012
9	PGGKPEALWV	0.01
2	PTNVESCPGG	0.002
5	VESCPGGKPE	0.001

Table XVII-V4-HLA-A24-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	TYLPTFETPI	90
1	KTYLPTFETP	0.024

Table XVIII-V1-HLA-B7-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		

WO 03/040340

PCT/US02/36002

155

Start	Subsequence	Score
343	APRIRAHNI	240.000
330	FPRINFFYM	200.000
393	NVRIDKVHL	200.000
776	IARVRDVEL	120.000
548	LPFGRPRVL	80.000
716	IPTHYFVVL	80.000
796	QPVSEILQL	80.000
597	LPPTVPDCL	80.000
701	APDEITKHL	72.000
552	RPRVLQKNV	40.000
758	CPEGKPEAL	24.000
733	TPENCPGWL	24.000
500	SVCGFANPL	20.000
456	EVYNLMCDL	20.000
98	LPPVILFSM	20.000
794	KVQPVSEIL	20.000
505	ANPLPTESL	18.000
580	MPMWSSYTV	12.000
119	TLMPNINKL	12.000
284	VVDHAFGML	6.000
283	QVVDHAFGM	5.000
570	YVSGFGKAM	5.000
778	RVRDVELLT	5.000
274	VSARVIKAL	4.000
240	STLLKWLDL	4.000
736	NCPGWLDVL	4.000
90	SQCPEGFDL	4.000
714	VPIPTHYFV	4.000
800	EILQLKTYL	4.000
196	WLTAMYQGL	4.000
540	ITATVKVNL	4.000
238	RISTLLKWL	4.000
253	RPRFYTMF	4.000
72	VCQGETSWL	4.000
474	GTHGSLNHL	4.000
40	KFRCGETRL	4.000
400	HLFVDQQWL	4.000
615	ESQKCSFYL	4.000
171	VNLNKNFSL	4.000
527	EQVNQMLNL	4.000
165	DNNMYDVNL	4.000
583	WSSYTPQQL	4.000
380	LTPDLPKRL	4.000
556	LQKNVDHCL	4.000
52	LCSCSDDCL	4.000
302	HNCVNIILL	4.000
251	AERPRFYTM	3.000
233	VPFEERIST	3.000
409	AVRSKSNTN	3.000
29	CVESTRIWM	2.250
146	FPNHYTIVT	2.000
682	GVNVVSGPI	2.000
485	VPFYEPSHA	2.000
611	VPPSESQKC	2.000
297	KQRNLHNCV	2.000
121	MPNINKLKT	2.000
601	VPDCLRADV	1.800
608	DVRVPPSES	1.500

Table XVIII-V1-HLA-B7-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
574	FGKAMRMPM	1.500
219	NGSFPSIYM	1.500
288	AFGMLMEGL	1.200
459	NLMCDLLRI	1.200
525	QLEQVNQML	1.200
512	SLDCFCPHL	1.200
193	QPMWLTAMY	1.200
217	AINGSFPSI	1.200
641	TSDSQYDAL	1.200
777	ARVRDVELL	1.200
359	NSEEIVRNL	1.200
470	APNNGTHGS	1.200
453	ENIEVYNLM	1.000
307	IILLADHGM	1.000
113	YLYTWD TLM	1.000
572	SGFGKAMRM	1.000
524	TQLEQVNQM	1.000
129	TCGIHSKYM	1.000
192	GQPMWLTAM	1.000
649	LITSNLVPM	1.000
45	ETRLEASLC	1.000
277	RVIKALQVV	1.000
198	TAMYQGLKA	0.900
577	AMRMPMWSS	0.900
248	LPKAERPRF	0.600
655	VPMYEEFRK	0.600
647	DALITSNLV	0.600
270	AGGPVSARV	0.600
528	QVNQMLNLT	0.500
363	IVRNLSCRK	0.500
670	SVLLIKHAT	0.500
13	VACKDRGDC	0.450
589	PQLGDT SPL	0.400
708	HLANTDVPI	0.400
299	RNLHNCVNI	0.400
807	YLPTFETTI	0.400
475	THGSLNHLL	0.400
452	FENIEVYNL	0.400
94	EGFDLPPVI	0.400
227	MPYNGSVPF	0.400
471	PNNGTHGSL	0.400
211	WPGSEVAIN	0.400

Table XVIII-V2-HLA-B7-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
5	DCLQRKDCC	0.100
6	CLQRKDCCA	0.100
7	LQRKDCCAD	0.100
4	DDCLQRKDC	0.015
1	SCSDDCLQR	0.010
2	CSDDCLQRK	0.003
8	QRKDCCADY	0.002
3	SDDCLQRKD	0.000
9	RKDCCADYK	0.000

Table XVIII-V3-HLA-B7-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
7	CPGGKPEAL	80
9	GGKPEALWV	0.2
6	SCPGGKPEA	0.1
3	NVESC PGGK	0.015
5	ESCPGGKPE	0.01
2	TNVE SCPGG	0.01
8	PGGKPEALW	0.003
4	VE SCPGGKP	0.002
1	PTNVE SCPG	0.001

Table XVIII-V4-HLA-B7-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
2	YLPTFETPI	0.4
1	TYLPTFETP	0.001

Table XIX-V1-HLA-B7-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
470	APNNGTHGSL	240.000

Table XIX-V1-HLA-B7-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
776	IARVRDVELL	120.000
588	VPQLGDT SPL	80.000
517	CPHLQNSTQL	80.000
233	VPFEERISTL	80.000
655	VPMYEEFRKM	60.000
371	KPDQHF KPYL	24.000
283	QVVDHAF GML	20.000
186	NPAWWHGQPM	20.000
170	DVNLNKNFSL	20.000
456	EVYNLMCDLL	20.000
71	SVCQGETSWL	20.000
504	FANPLPTESL	18.000
409	AVRSKSN TNC	15.000
287	HAFGMLMEGL	12.000
700	DAPDEITKHL	12.000
94	EGFDLPPVIL	6.000
343	APRIRAHNIP	6.000
275	SARVIKALQV	6.000
300	NLHNCVNIIL	4.000
89	QSQCPEGFDL	4.000
775	HIARVRDVEL	4.000
239	ISTLLKWLDL	4.000
392	KNVRIDKVHL	4.000
440	GPSFKEKTEV	4.000
547	NLPFGRPRVL	4.000
795	VQPVSEILQL	4.000
732	HTPENCPGWL	4.000
473	NGTHGSLNHL	4.000
555	VLQKNVDHCL	4.000
511	ESLDCFCPHL	4.000
51	SLCSCSDDCL	4.000
714	VPIPTHYFVV	4.000
757	SCPEGKPEAL	4.000
330	FPRINFFYMY	4.000
663	KMWDYFHSVL	4.000
640	RTSDSQYDAL	4.000
524	TQLEQVNQML	4.000
428	EFRSMEAIFL	4.000
358	FNSEEIVRNL	4.000
735	ENCPGWLDVL	4.000
118	DTLMPNINKL	4.000
539	EITATVKVNL	4.000
556	LQKNVDHCLL	4.000
474	GTHGSLNHLL	4.000
499	FSVCGFANPL	4.000
379	YLTPDL PKRL	4.000
744	LPFIIPHRPT	3.000
648	ALITSNLVPM	3.000
716	IPTHYFV VLT	2.000
552	RPRVLQKNVD	2.000

Table XIX-V1-HLA-B7-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
595	SPLPPTVPDC	2.000
272	GPVSARVIKA	2.000
273	PVSARVIKAL	2.000
508	LPTESLDCFC	2.000
253	RPRFYTMYFE	2.000
121	MPNINKLKTC	2.000
506	NPLPTESLDC	2.000
701	APDEITKHLA	1.800
600	TVPDCLRADV	1.500
218	INGSFPSIYM	1.500
28	TCVESTRIWM	1.500
284	VVDHAFGMLM	1.500
43	CGETRLEASL	1.200
270	AGGPVSARVI	1.200
292	LMEGLKQRNL	1.200
105	SMDGFRAEYL	1.200
624	ADKNITHGFL	1.200
188	AWWHGQPMWL	1.200
216	VAINGSFPSI	1.200
92	CPEGFDLPPV	1.200
111	AEYLYTWDTL	1.200
571	VSGFGKAMRM	1.000
423	HGYNNEFRSM	1.000
713	DVPIPTHYFV	1.000
191	HGQPMWLTAM	1.000
393	NVRIDKVHLF	1.000
159	ESHGIIDNNM	1.000
523	STQLEQVNQM	1.000
282	LQVVDHAFGM	1.000
97	DLPPVILFSM	1.000
136	YMRAMYPTKT	1.000
306	NIILLADHGM	1.000
317	QTYCNKMEYM	1.000
131	GIHSKYMRAM	1.000
128	KTCGIHSKYM	1.000
207	ATYFWPGSEV	0.900
198	TAMYQGLKAA	0.900
250	KAERPRFYTM	0.900
12	DVACKDRGDC	0.750
232	SVPFEERIST	0.750
608	DVRVPPSESQ	0.750
337	YMYEGPAPRI	0.600
676	HATERNGVNV	0.600
390	YAKNVRIDKV	0.600
773	TAHIARVRDV	0.600
193	QPMWLTAMYQ	0.600
767	WVEERF TAHI	0.600
580	MPMWSSYTV P	0.600
269	HAGGPVSARV	0.600

Table XIX-V2-HLA-B7-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
8	LQRKDCCADY	0.200
6	DCLQRKDCCA	0.100
7	CLQRKDCCAD	0.010
1	CSCSDDCLQR	0.010
5	DDCLQRKDCC	0.010
2	SCSDDCLQRK	0.010
4	SDDCLQRKDC	0.004
3	CSDDCLQRKD	0.003
9	QRKDCCADYK	0.001
10	RKDCCADYKS	0.001

Table XIX-V3-HLA-B7-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	SCPGGKPEAL	4
8	CPGGKPEALW	0.6
1	RPTNVESCPG	0.2
6	ESCPGGKPEA	0.1
4	NVESCPGGKP	0.023
9	PGGKPEALWV	0.02
10	GGKPEALWVE	0.01
3	TNVESCPGGK	0.01
2	PTNVESCPGG	0.001
5	VESCPGGKPE	0.001

Table XIX-V4-HLA-B7-10mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	TYLPTFETPI	0.04
1	KTYLPTFETP	0.01

Table XX-V1-HLA-B3501-9mers-161P2F10B		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		

Start	Subsequence	Score
253	RPRFYTMYP	120.000
330	FPRINFFYM	120.000
248	LPKAERPRF	90.000
450	EPFENIEVY	80.000
98	LPPVILFSM	40.000
508	LPTESLDCF	40.000
193	QPMWLTAMY	40.000
133	HSKYMRAMY	30.000
796	QPVSEILQL	30.000
552	RPRVLQKNV	24.000
371	KPDQHFQPY	24.000
343	APRIRAHNI	24.000
597	LPPTVPDCL	20.000
548	LPFGRPRVL	20.000
716	IPTHYFVVL	20.000
227	MPYNGSVPF	20.000
496	VSKFSVCGF	15.000
701	APDEITKHL	12.000
776	IARVRDVEL	9.000
558	KNVDHCLLY	8.000
733	TPENCPGWL	6.000
758	CPEGKPEAL	6.000
347	RAHNIPHDF	6.000
138	RAMYPTKTF	6.000
638	SNRTSDSQY	6.000
233	VPFEERIST	6.000
574	FGKAMRMPM	6.000
583	WSSYTVPQL	5.000
615	ESQKCSFYL	5.000
274	VSARVIKAL	5.000
393	NVRIDKVHL	4.500
453	ENIEVYNLM	4.000
128	KTCGIHSKY	4.000
580	MPMWSSYTV	4.000
714	VPIPTHYFV	4.000
351	IPHDFFSFN	4.000
626	KNITHGFLY	4.000
283	QVVDHAFGM	4.000
524	TQLEQVNQM	4.000
803	QLKTYLPTF	3.000
611	VPPSESQKC	3.000
368	SCRKPDQHF	3.000
359	NSEEIVRNL	3.000
556	LQKNVDHCL	3.000
764	EALWVEERF	3.000
485	VPFYEPSHA	3.000
161	HGIIDNNMY	3.000
180	SSKEQNNPA	3.000
211	WPGSEVAIN	3.000
238	RISTLLKWL	2.000
222	FPSIYMPYN	2.000
562	HCLLYHREY	2.000
685	VVSGPIFDY	2.000
649	LITSNLVPM	2.000
650	ITSNLVPMY	2.000
146	FPNHYTIVT	2.000
218	INGSFPSIY	2.000
634	YPPASNRTS	2.000

Table XX-V1-HLA-B3501-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
107	DGFRAEYLY	2.000
783	ELLTGLDFY	2.000
570	YVSGFGKAM	2.000
121	MPNINKLKT	2.000
113	YLYTWDTLM	2.000
317	QTYCNKMEY	2.000
231	GSVPFEERI	2.000
417	NCGGGNHGY	2.000
219	NGSFPSIYM	2.000
129	TCGIHSKYM	2.000
201	YQGLKAATY	2.000
307	IILLADHGM	2.000
687	SGPIFDYNY	2.000
192	GQPMWLTAM	2.000
489	EPSHAEVS	2.000
794	KVQPVSEIL	2.000
470	APNNGTHGS	2.000
572	SGFGKAMRM	2.000
430	RSMEAIFLA	2.000
380	LTPDLPKRL	2.000
641	TSDSQYDAL	1.500
203	GLKAATYFW	1.500
350	NIPHDFFSF	1.500
400	HLFVDQQWL	1.500
90	SQCPEGFDL	1.500
72	VCQGETSWL	1.500
124	INKLKTCGI	1.200
61	QKKDCCADY	1.200
778	RVRDVELLT	1.200
601	VPDCLRADV	1.200
297	KQRNLHNCV	1.200
736	NCPGWLDVL	1.000
732	HTPENCPGW	1.000
196	WLTAMYQGL	1.000
28	TCVESTRIW	1.000
683	VNVVSGPIF	1.000
202	QGLKAATYF	1.000
500	SVCGFANPL	1.000
527	EQVNQMLNL	1.000
88	QQSQCEPGE	1.000
713	DVPIPTHYF	1.000
302	HNCVNIILL	1.000

Table XX-V2-HLA-B35-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
8	QRKDCCADY	1.200
5	DCLQRKDCC	0.100
6	CLQRKDCCA	0.100
7	LQRKDCCAD	0.045
1	SCSDDCLQR	0.030
2	CSDDCLQRK	0.030
4	DDCLQRKDC	0.010
9	RKDCCADYK	0.001
3	SDDCLQRKD	0.000

Table XX-V3-HLA-B35-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 7 each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
7	CPGGKPEAL	20
9	GGKPEALWV	0.9
6	SCPGGKPEA	0.1
8	PGGKPEALW	0.05
5	ESCPGGKPE	0.05
2	TNVEPCPGG	0.02
3	NVEPCPGGK	0.003
1	PTNVEPCPG	0.002
4	VEPCPGGKP	0.001

Table XX-V4-HLA-B35-9mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Start	Subsequence	Score
2	YLPTFETPI	0.4
1	TYLPTFETP	0.001

Table XXI-V1-HLA-B35-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
248	LPKAERPRFY	120.000

Table XXI-V1-HLA-B35-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
330	FPRINFFYMY	120.000
655	VPMYEEFRKM	60.000
233	VPFEERISTL	40.000
141	YPTKTFPNHY	40.000
186	NPAWWHGQPM	40.000
737	CPGWLDVLPF	30.000
588	VPQLGDT SPL	30.000
104	FSMDGFRAEY	20.000
470	APNNGTHGSL	20.000
517	CPHLQNSTQL	20.000
180	SSKEQNNPAW	15.000
776	IARVRDVELL	13.500
371	KPDQHFKPYL	12.000
381	TPDLPKRLHY	12.000
686	VSGPIFDYNY	10.000
220	GSFPSIYMPY	10.000
571	VSGFGKAMRM	10.000
511	ESLDCFCPHL	10.000
159	ESHGIIDNNM	10.000
637	ASNRTSDSQY	10.000
89	QSQCPEGFDL	7.500
280	KALQVVDHAF	6.000
700	DAPDEITKHL	6.000
347	RAHNIPHDF	6.000
577	AMRMPMWSSY	6.000
320	CNKMEYMTDY	6.000
440	GPSFKEKTEV	6.000
384	LPKRLHYAKN	6.000
60	LQKKDCCADY	6.000
499	FSVCGFANPL	5.000
239	ISTLLKWLDL	5.000
367	LSCRKPDQHF	5.000
70	KSVCQGETSW	5.000
556	LQKNVDHCLL	4.500
612	PPSESQKCSF	4.000
640	RTSDSQYDAL	4.000
351	IPHDFFSFNS	4.000
508	LPTESLDCFC	4.000
663	KMWDYFHSV L	4.000
714	VPIPTHYFVV	4.000
28	TCVESTRIWM	4.000
450	EPFENIEVYN	4.000
128	KTCGIHSKYM	4.000
250	KAERPRFYTM	3.600
287	HAFGMLMEGL	3.000
32	STRIWMCNKF	3.000
393	NVRIDKVHLF	3.000
14	ACKDRGDCCW	3.000
392	KNVRIDKVHL	3.000

Table XXI-V1-HLA-B35-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
758	CPEGKPEALW	3.000
541	TATVKVNL PF	3.000
423	HGYNNEFRSM	3.000
506	NPLPTESLDC	3.000
798	VSEILQLKTY	3.000
504	FANPLPTESL	3.000
97	DLPPVILFSM	2.000
316	DQTYCNKMEY	2.000
595	SPLPPTVPDC	2.000
272	GPVSARVIKA	2.000
648	ALITSNLVPM	2.000
479	LNHLLKVPFY	2.000
283	QVVDHAFGML	2.000
416	TNCGGGNHGY	2.000
131	GIHSKYMRAM	2.000
649	LITSNLVPMY	2.000
684	NVVS GPIFDY	2.000
306	NIILLADHGM	2.000
121	MPNINKLKTC	2.000
358	FNSEEIVRNL	2.000
218	INGSFPSIYM	2.000
192	GQPMWLTAMY	2.000
317	QTYCNKMEYM	2.000
757	SCPEGKPEAL	2.000
611	VPPSESQKCS	2.000
744	LPFIIPHRPT	2.000
217	AINGSFPSIY	2.000
191	HGQPMWLTAM	2.000
341	GPAPRIRAHN	2.000
732	HTPENCPGWL	2.000
282	LQVVDHAFGM	2.000
716	IPTHYFVLT	2.000
94	EGFDLPPVIL	2.000
523	STQLEQVNQM	2.000
619	CSFYLADKNI	2.000
524	TQLEQVNQML	2.000
748	IPHRPTNVES	2.000
390	YAKNVRIDKV	1.800
310	LADHGMDQTY	1.800
275	SARVIKALQV	1.800
92	CPEGFDLPPV	1.800
71	SVCQGETSWL	1.500
213	GSEVAINGSF	1.500
349	HNIPHDFFSF	1.500
795	VQPVSEILQL	1.500
574	FGKAMRMPMW	1.500
496	VSKFSVCGFA	1.500
247	DLPKAERPRF	1.500
552	RPRVLQKNVD	1.200
676	HATERNGVNV	1.200

Table XXI-V2-HLA-B35-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 5; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
8	LQRKDCCADY	6.000
6	DCLQRKDCCA	0.100
1	CSCSDDCLQR	0.075
3	CSDDCLQRKD	0.030
2	SCSDDCLQRK	0.020
7	CLQRKDCCAD	0.015
5	DDCLQRKDCC	0.010
9	QRKDCCADYK	0.006
10	RKDCCADYKS	0.006
4	SDDCLQRKDC	0.003

Table XXI-V3-B35-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
8	CPGGKPEALW	10
7	SCPGGKPEAL	1
1	RPTNVESCPG	0.6
6	ESCPGGKPEA	0.5
9	PGGKPEALWV	0.03
10	GGKPEALWVE	0.03
3	TNVESCPGGK	0.02
4	NVESCPGGKP	0.003
2	PTNVESCPGG	0.001
5	VESCPGGKPE	0.001

Table XXI-V4-HLA-B35-10mers-161P2F10B

Each peptide is a portion of SEQ ID NO: 9; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Start	Subsequence	Score
2	TYLPTFETPI	0.04
1	KTYLPTFETP	0.02

Tables XXII-XLIX:

Table XXII-V1-HLA-A1-9mers-161P2F10B		
Pos	123456789	score
165	SMDGFRAEY	29
431	KPDQHFKPY	28
442	PDLPKRLHY	26
188	KTCGIHSKY	24
618	KNVDHCLLY	23
710	ITSNLVPMY	23
858	VSEILQLKT	23
8	ATEQPVKKN	21
193	HSKYMRAMY	21
202	PTKTFPNHY	21
377	QTYCNKMEY	21
491	SMEAIFLAH	21
209	HYTIVTGLY	19
391	PRINFFYMY	19
462	FVDQQWLAV	19
477	NCGGGNHGY	19
514	NIEVYNLMC	19
638	MRMPMWSSY	19
772	TDVPIPTHY	19
859	SEILQLKTY	19
13	VKKNLKKY	18
389	YFPRINFFY	18
455	RIDKVHLFV	18
686	KNITHGFLY	18
737	ATERNGVNV	18
745	VVSGPIFDY	18
843	ELLTGLDFY	18
152	CPEGFDLPP	17
167	DGFRAEYLY	17
217	YPESHGIID	17
221	HGIIDNNMY	17
261	YQGLKAATY	17
281	SFPSIYMPY	17
371	ADHGMDQTY	17
420	SEEIVRNLS	17
674	SESQKCSFY	17
698	SNRTSDSQY	17
747	SGPIFDYNY	17
771	NTDVPIPTH	17
839	VRDVELLTG	17
115	CSDDCLQKK	16
134	QGETSWLEE	16
253	QPMWLTAMY	16
312	ERPRFYTM	16
381	NKMEYMTDY	16
386	MTDYFPRIN	16
506	KTEVEPFEN	16
510	EPFENIEVY	16
540	NHLLKVPFY	16
569	PTESLDCFC	16
602	ATVKVNLPF	16
619	NVDHCLLYH	16
622	HCLLYHREY	16
651	LGDTSPLEP	16

Table XXII-V1-HLA-A1-9mers-161P2F10B		
Pos	123456789	score
659	PTVPDCLRA	16
673	PSESQKCSF	16
701	TSDSQYDAL	16
89	CVESTRIWM	15
121	QKKDCCADY	15
210	YTIVTGLYP	15
278	INGSFPSIY	15
309	PKAERPRFY	15
321	FEEDSSGH	15
344	VVDHAFGML	15
419	NSEEIVRN	15
521	MCDLLRIQP	15
547	FYEPSHAEE	15
585	QLEQVNQML	15
719	EEFRKMWDY	15
851	YQDKVQPV	15
47	KLEKQGSCR	14
116	SDDCLQKKD	14
273	GSEVAINGS	14
300	STLLKWLDL	14
508	EVEPFENIE	14
646	YTVPQLGDT	14
754	NYDGHFDAP	14
847	GLDFYQDKV	14

Table XXII-V2-HLA-A1-9mers-161P2F10B		
Pos	123456789	score
8	QRKDCCADY	15
2	CSDDCLQRK	14
3	SDDCLQRKD	14
9	RKDCCADYK	10
1	SCSDDCLQR	8

Table XXII-V3-HLA-A1-9mers-161P2F10B		
Pos	123456789	score
5	ESCPGGKPE	11
3	NVESC PGK	10
9	GGKPEALWV	10
1	PTNVESCPG	6
4	VESCPGGKP	5

Table XXII-V4-HLA-A1-9mers-161P2F10B		
Pos	123456789	score
1	TYLPTFETP	7
2	YLPTFETPI	3

Table XXIII-V1-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
179	TLMPNINKL	30
40	GLGLGLRKL	29

Table XXIII-V1-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
723	KMWDYFHSV	25
27	VLLALLVIM	24
29	LALLVIMSL	24
31	LLVIMSLGL	24
215	GLYPESHGI	24
277	AINGSFPSI	24
519	NLMCDLLRI	24
572	SLDCFCPHL	24
33	VIMSLGLGL	23
360	NLHNCVNII	23
847	GLDFYQDKV	23
867	YLPTFETTI	23
23	IACIVLLAL	22
298	RISTLLKWL	22
534	GTHGSLNHL	22
607	NLPFGRPRV	22
768	HLANTDVPI	22
24	ACIVLLALL	21
26	IVLLALLVI	21
223	IIDNNMYDV	21
259	AMYQGLKAA	21
460	HLFVDQQWL	21
592	MLNLTQEEI	21
623	CLLYHREYV	21
21	YKIACIVLL	20
22	KIACIVLLA	20
25	CIVLLALLV	20
65	GLENCRCDV	20
256	WLTAMYQGL	20
300	STLLKWLDL	20
337	RVIKALQVV	20
385	YMTDYFPRI	20
439	YLTPDLPKR	20
537	GSLNHLLKV	20
560	SVCGFANPL	20
600	ITATVKVNL	20
807	IIPHRPTNV	20
836	IARVRDVEL	20
28	LLALLVIMS	19
37	LGLGLGLGL	19
204	KTFPNHYTI	19
451	AKNVRIDKV	19
455	RIDKVHLFV	19
585	QLEQVNQML	19
653	DTSPLPPTV	19
709	LITSNLVPM	19
800	WLDVLPFII	19
5	LTATEQPV	18
36	SLGLGLGLG	18
285	IYMPYNGSV	18
351	MLMEGLKQR	18
367	IILLADHGM	18
443	DLPKRLHYA	18
682	YLADKNITH	18
731	VLLIKHATE	18
834	AHIARVRDV	18

WO 03/040340

PCT/US02/36002

160

Table XXIII-V1-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
840	RDVELLTGL	18
860	EILQLKTYL	18
35	MSLGLGLGL	17
38	GLGLGLGLR	17
139	WLEENCDTA	17
165	SMDGFRAEY	17
173	YLYTWD TLM	17
334	VSARVIKAL	17
397	YMYEGPAPR	17
440	LTPDLPKRL	17
462	FVDQQWLAV	17
512	FENIEVYNL	17
538	SLNHLLKVP	17
565	ANPLPTESL	17
595	LTQEEITAT	17
596	TQEEITATV	17
737	ATERNGVNV	17
738	TERNGVNVV	17
799	GWLDVLPFI	17
825	ALWVEERFT	17
850	FYQDKVQPV	17
854	KVQPVSEIL	17
863	QLKTYLPTF	17
34	IMSLGLGLG	16
157	DLPPVILFS	16
208	NHYTIVTGL	16
231	VNLNKNFSL	16
301	TLLKWLDLP	16
330	AGGPVSARV	16
453	NVRIDKVHL	16
520	LMCDLLRIQ	16
646	YTVPLGDT	16
706	YDALITSNL	16
707	DALITSNLV	16
732	LLIKHATER	16
774	VPIPTHYFV	16
796	NCPGWLDVL	16
806	FIIPHRPTN	16
837	ARVRDVELL	16
861	ILQLKTYLP	16
18	LKKYKIACI	15
104	GETRLEASL	15
107	RLEASLCSC	15
119	CLQKKDCCA	15
132	VCQGETSWL	15
211	TIVTGLYPE	15
305	WLDLPKAER	15
328	GHAGGPVSA	15
340	KALQVVDHA	15
344	VVDHAFGML	15
369	LLADHGMDQ	15
419	NSEEIVRNL	15
496	FLAHGPSFK	15
542	LLKVPFYEP	15
591	QMLNLTQEE	15
594	NLTQEEITA	15

Table XXIII-V1-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
598	EEITATVKV	15
608	LPFGRPRVL	15
615	VLQKNVDHC	15
692	FLYPPASNR	15
708	ALITSNLVP	15
735	KHATERNGV	15
761	APDEITKHL	15
775	PIPTHYFVV	15
803	VLPFIIPHR	15
831	RFTAHIA RV	15
843	ELLTGLDFY	15
856	QPVSEILQL	15
865	KTYLPTFET	15
30	ALLVIMSLG	14
42	GLGLRKLEK	14
150	SQCPEGFDL	14
155	GFDLPPVIL	14
191	GIHSKYMRA	14
268	TYFWPGSEV	14
270	FWPGSEVAI	14
284	SIYMPYNGS	14
294	PFEERISTL	14
303	LKWLDLPKA	14
336	ARVIKALQV	14
341	ALQVVDHAF	14
348	AFGMLMEGL	14
350	GMLMEGLKQ	14
362	HNCVNIILL	14
368	ILLADHGMD	14
436	FKPYLTPDL	14
448	LHYAKNVRI	14
524	LLRIQPAPN	14
548	YEPSHAEEV	14
581	QNSTQLEQV	14
588	QVNQMLNLT	14
624	LLYHREYVS	14
640	MPMWSSYTV	14
643	WSSYTVPQL	14
656	PLPPTVPDC	14
661	VPDCLRADV	14
664	CLRADVRVP	14
688	ITHGFLYPP	14
713	NLVPMYEEF	14
716	PMYEEFRKM	14
730	SVLLIKHAT	14
742	GVNVVSGPI	14
764	EITKHLANT	14
776	IPTHYFVVL	14
844	LLTGLDFYQ	14
853	DKVQPVSEI	14

Table XXIII-V2-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
6	CLQRKDCCA	15
2	CSDDCLQRK	5

Table XXIII-V2-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
3	SDDCLQRKD	5

Table XXIII-V3-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
7	CPGGKPEAL	14
6	SCPGGKPEA	13
9	GGKPEALWV	13

Table XXIII-V4-HLA-A0201-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	21

Table XXIV-V1-HLA-A0203-9mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XXIV-V2-HLA-A0203-9mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XXIV-V3-HLA-A0203-9mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XXIV-V4-HLA-A0203-9mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XXV-V1-HLA-A3-9mers-161P2F10B		
Pos	123456789	score
670	RVPPSESQK	33
496	FLAHGPSFK	27
692	FLYPPASNR	27
42	GLGLRKLEK	26
423	IVRNLSCKR	26
6	TLATEQPVK	25
614	RVLQKNVDH	25
12	PVKKNTLKK	24
302	LLKWLDLPK	24
610	FGRPRVLQK	24
47	KLEKQGSCR	23
708	ALITSNLVP	23
26	IVLLALLVI	22
447	RLHYAKNVR	22
782	VVLTSCKNK	22
838	RVRDVELLT	22
337	RVIKALQVV	21
405	RIRAHNIPH	21

Table XXV-V1-HLA-A3-9mers-161P2F10B		
Pos	123456789	score
469	AVRSKSNTN	21
597	QEEITATVK	21
628	REYVSGFGK	21
682	YLADKNITH	21
731	VLLIKHATE	21
732	LLIKHATER	21
814	NVESCPEGK	21
857	PVSEILQLK	21
863	QLKTYLPTF	21
39	LGLGLGLRK	20
351	MLMEGLKQR	20
624	LLYHREYVS	20
843	ELLTGLDFY	20
196	YMRAMYPTK	19
227	NMYDVNLNK	19
338	VIKALQVVD	19
341	ALQVVDHAF	19
544	KVPFYEPSH	19
664	CLRADVRVP	19
802	DVLPFIIPH	19
806	FIIPHRPTN	19
827	WVEERFTAH	19
27	VLLALLVIM	18
30	ALLVIMSLG	18
38	GLGLGLGLR	18
107	RLEASLCSC	18
114	SCSDDCLQK	18
261	YQGLKAATY	18
275	EVAINGSFP	18
305	WLDLPKAER	18
368	ILLADHGMD	18
422	EIVRNLSR	18
453	NVRIDKVHL	18
524	LLRIQPAPN	18
550	PSHAEVSK	18
745	VVSGPIFDY	18
759	FDAPDEITK	18
835	HIARVRDVE	18
157	DLPPVILFS	17
162	ILFSMDGFR	17
186	KLKTCGIHS	17
215	GLYPESHGI	17
296	EERISTLLK	17
439	YLTPDLPKR	17
458	KVHLFVDQQ	17
619	NVDHCLLYH	17
647	TVPQLGDT	17
668	DVRVPPSES	17
713	NLVPMYEEF	17
854	KVQPVSEIL	17
7	LATEQPVKK	16
44	GLRKLEKQG	16
68	NCRCDVACK	16
94	RIWMCNKFR	16
122	KKDCCADYK	16
131	SVCQGETSW	16

Table XXV-V1-HLA-A3-9mers-161P2F10B		
Pos	123456789	score
343	QVVDHAFGM	16
355	GLKQRNLHN	16
369	LLADHGMDQ	16
442	PDLPKRLHY	16
455	RIDKVHLFV	16
498	AHGSPFKEK	16
523	DLLRIQPAP	16
526	RIQPAPNNG	16
560	SVCGFANPL	16
630	YVSGFGKAM	16
678	KCSFYLADK	16
698	SNRTSDSQY	16
768	HLANTDVPI	16

Table XXV-V2-HLA-A3-9mers-161P2F10B		
Pos	123456789	score
9	RKDCCADYK	17
6	CLQRKDCCA	14
8	QRKDCCADY	13
1	SCSDDCLQR	12
2	CSDDCCLQRK	11

Table XXV-V3-HLA-A3-9mers-161P2F10B		
Pos	123456789	score
3	NVESCPEGK	22
9	GGKPEALWV	11

Table XXV-V4-HLA-A3-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	11

Table XXVI-V1-HLA-A26-9mers-161P2F10B		
Pos	123456789	score
510	EPFENIEVY	31
516	EVYNLMCDL	29
773	DVPIPTHYF	26
388	DYFPRINFF	25
719	EEFRKMWDY	25
422	EIVRNLSR	24
710	ITSNLVPMY	24
745	VVSGPIFDY	24
802	DVLPFIIPH	24
843	ELLTGLDFY	24
275	EVAINGSFP	23
587	EQVNQMLNL	23
860	EILQLKTYL	23
167	DGFRAEYLY	22
188	KTCGIHSKY	22
297	ERISTLLKW	21
312	ERPRFYTMY	21
508	EVEPFENIE	21

Table XXVI-V1-HLA-A26-9mers-161P2F10B		
Pos	123456789	score
602	ATVKVNLPF	21
10	EQPVKKNTL	20
344	VVDHAFGML	20
534	GTHGSLNHL	20
555	EVSKFSVCG	20
841	DVELLTGLD	20
859	SEILQLKTY	20
32	LVIMSLGLG	19
72	DVACKDRGD	19
105	ETRLEASLC	19
136	ETSWLEENC	19
172	EYLYTWDTL	19
230	DVNLNKNFS	19
300	STLLKWLDL	19
377	QTYCNKMEY	19
488	EFRSMEAIF	19
560	SVCGFANPL	19
599	EITATVKVN	19
653	DTSPLPPTV	19
668	DVRVPPSES	19
764	EITKHLANT	19
840	RDVELLTGL	19
854	KVQPVSEIL	19
856	QPVSEILQL	19
202	PTKTFPNHY	18
337	RVIKALQVV	18
364	CVNIILLAD	18
440	LTPDLPKRL	18
453	NVRIDKVHL	18
454	VRIDKVHLF	18
646	YTVPQLGDT	18
675	ESQKCSFYL	18
720	EFRKMWDYF	18
772	TDVPIPTHY	18
824	EALWVEERF	18
857	PVSEILQLK	18
24	ACIVLLALL	17
77	DRGDCCWDF	17
178	DTLMPNINK	17
274	SEVAINGSF	17
298	RISTLLKWL	17
551	SHAEVSKF	17
554	EEVSKFSVC	17
629	EYVSGFGKA	17
660	TVPDCLRAD	17
685	DKNITHGFL	17
726	DYFHSVLLI	17
853	DKVQPVSEI	17
2	ESTLTATE	16
21	YKIACIVLL	16
33	VIMSLGLGL	16
87	DTCVESTRI	16
145	DTAQQSQCP	16
154	EGFDLPPVI	16
179	TLMPNINKL	16
219	ESHGIIDNN	16

WO 03/040340

PCT/US02/36002

162

Table XXVI-V1-HLA-A26-9mers-161P2F10B		
Pos	123456789	score
225	DNNMYDVNL	16
294	PFEERISTL	16
333	PVSARVIKA	16
419	NSEEIVRNL	16
421	EEIVRNLSC	16
513	ENIEVYNLM	16
600	ITATVKVNL	16
603	TVKVNLPFG	16
618	KNVDHCLLY	16
744	NVVS GPIFD	16
830	ERFTHAIAR	16
837	ARVRDVELL	16
13	VKKNTLKKY	15
26	IVLLALLVI	15
204	KTFPNHYTI	15
221	HGIIDNNMY	15
343	QVVDHAFGM	15
391	PRINFFYMY	15
458	KVHLFVDQQ	15
493	EAIFLAHGP	15
540	NHLLKVPFY	15
568	LPTESLDCF	15
595	LTQEEITAT	15
598	EEITATVKV	15
619	NVDHCLLYH	15
626	YHREYVSGF	15
686	KNITHGFLY	15
763	DEITKHLAN	15
781	FVVLTSCKN	15

Table XXVI-V2-HLA-A26-9mers-161P2F10B		
Pos	123456789	score
8	QRKDCCADY	11
4	DDCLQRKDC	8
5	DCLQRKDCC	8
2	CSDDCLQRK	5

Table XXVI-V3-HLA-A26-9mers-161P2F10B		
Pos	123456789	score
3	NVESC PGK	12
5	ESCPG GKPE	12
7	CPGGKPEAL	10
1	PTNVE SC PG	8
2	TNVE SC PG	7

Table XXVI-V4-HLA-A26-9mers-161P2F10B		
Pos	123456789	score
1	TYLPTFETP	3

Table XXVII-V1-HLA-B0702-9mers-161P2F10B		
Pos	123456789	score

Table XXVII-V1-HLA-B0702-9mers-161P2F10B		
Pos	123456789	score
776	IPTHYFVVL	25
761	APDEITKHL	24
608	LPFGRPRVL	23
818	CPEGKPEAL	23
856	QPVSEILQL	22
313	RPRFYTMFY	20
403	APRIRAHNI	20
657	LPPTVPDCL	20
793	TPENC PGWL	20
287	MPYNGSVPF	19
308	LPKAERPRF	19
390	FPRINFFYM	19
612	RPRVLQKNV	19
661	VPDCLRADV	19
181	MPNINKLKT	18
206	FPNHYTIVT	18
655	SPLPPTVPD	18
774	VPIPTHYFV	18
293	VPFEERIST	17
640	MPMWSSYTV	17
155	GFDLPPVIL	16
158	LPPVILFSM	16
545	VPFYEPSHA	16
565	ANPLPTESL	16
568	LPTESLDCF	16
808	IPHRPTNVE	16
35	MSLGLGLGL	15
37	LGLGLGLGL	15
152	CPEGFDLPP	15
836	IARVRDVEL	15
23	IACIVLLAL	14
33	VIMSLGLGL	14
100	KFRCGETRL	14
323	EPDSSGHAG	14
431	KPDQHFKPY	14
441	TPDLPKRLH	14
453	NVRIDKVHL	14
549	EP SHAEVS	14
600	ITATVKVNL	14
643	WSSYTVPQL	14
649	PQLGDT SPL	14
822	KPEALWVEE	14
21	YKIACIVLL	13
24	ACIVLLALL	13
58	CFDASFRGL	13
132	VCQGETSWL	13
201	YPTKTFPNH	13
253	QPMWLTAMY	13
271	WPGSEVAIN	13
282	FPSIYMPYN	13
298	RISTLLKWL	13
332	GPVSARVIK	13
348	AFGMLMEGL	13
401	GPAPRIRAH	13
411	IPHDFFSFN	13
489	FRSMEAIFL	13

Table XXVII-V1-HLA-B0702-9mers-161P2F10B		
Pos	123456789	score
530	APNNGTHGS	13
560	SVCGFANPL	13
587	EQVNQMLNL	13
658	PPTVPDCLR	13
695	PPASNRTSD	13
796	NCPGWLDVL	13
797	CPGWLDVLP	13
837	ARVRDVELL	13
1	MESTLTLAT	12
11	QPVKKNTLK	12
20	KYKIACIVL	12
31	LLVIMSLGL	12
112	LCSCSDDCL	12
150	SQCPEGFDL	12
179	TLMPNINKL	12
198	RAMYPTKTF	12
208	NHYTIVTGL	12
225	DNNMYDVNL	12
249	WWHGQPMWL	12
277	AINGSFPSI	12
300	STLLKWLDL	12
311	AERPRFYTM	12
334	VSARVIKAL	12
344	VVDHAFGML	12
398	MYEGPAPRI	12
400	EGPAPRIRA	12
408	AHNIPHDFE	12
428	SCRKPDQHF	12
436	FKPYLTPDL	12
455	RIDKVHLFV	12
500	GPSFKEKTE	12
510	EPFENIEVY	12
528	QPAPNNGTH	12
531	PNNGTHGSL	12
534	GTHGSLNHL	12
572	SLDCFCPHL	12
602	ATVKVNLPF	12
630	YVSGFGKAM	12
672	PPSESQKCS	12
675	ESQKCSFYL	12
701	TSDSQYDAL	12
706	YDALITSNL	12
724	MWDYFHSV L	12
725	WDYFHSVLL	12
840	RDVELLTGL	12
854	KVQPVSEIL	12
860	EILQLKTYL	12

Table XXVII-V2-B0702-9mers-161P2F10B		
Pos	123456789	score
6	CLQRKDCCA	6
1	SCSDDCLQR	4
7	LQRKDCCAD	4
9	RKDCCADYK	2
2	CSDDCLQRK	1

WO 03/040340

PCT/US02/36002

163

Table XXVII-V2-B0702-9mers-161P2F10B		
Pos	123456789	score
3	SDDCLQRKD	1
4	DDCLQRKDC	1
8	QRKDCCADY	1

Table XXVII-V3-HLA-B0702-9mers-161P2F10B		
Pos	123456789	score
7	CPGGKPEAL	24

Table XXVII-V4-HLA-B0702-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	8

Table XXVIII-V1-HLA-B08-9mers-161P2F10B		
Pos	123456789	score
836	IARVRDVEL	32
18	LKKYKIACI	28
184	INKLKTCGI	28
818	CPEGKPEAL	27
355	GLKQRNLHN	26
403	APRIRAHNI	26
502	SFKEKTEVE	24
608	LPFGRPRVL	24
10	EQPVKKNTL	23
300	STLLKWLDL	23
231	VNLNKNFSL	22
308	LPKAERPRF	22
786	SCKNKSHTP	22
863	QLKTYLPTF	22
294	PFEERISTL	21
353	MEGLKQRNL	21
20	KYKIACIVL	20
616	LQKNVDHCL	20
98	CNKFRGET	19
761	APDEITKHL	19
40	GLGLGLRKL	18
53	SCRKKCFDA	18
74	ACKDRGDCC	18
166	MDGFRAEYL	18
238	SLSSKEQNN	18
313	RPRFYTMYP	18
448	LHYAKNVRI	18
454	VRIDKVHLF	18
460	HLFVDQQWL	18
500	GPSFKEKTE	18
542	LLKVPFYEP	18
556	VSKFSVCGF	18
572	SLDCFCPHL	18
776	IPTHYFVVL	18
861	ILQLKTYLP	18
11	QPVKKNTLK	17
15	KNTLKKYKI	17
17	TLKKYKIAC	17

Table XXVIII-V1-HLA-B08-9mers-161P2F10B		
Pos	123456789	score
42	GLGLRKLEK	17
179	TLMPNINKL	17
263	GLKAATYFW	17
401	GPAPRIRAH	17
444	LPKRLHYAK	17
453	NVRIDKVHL	17
504	KEKTEVEPF	17
585	QLEQVNQML	17
682	YLADKNITH	17
731	VLLIKHATE	17
856	QPVSEILQL	17
23	IACIVLLAL	16
29	LALLVIMSL	16
31	LLVIMSLGL	16
55	RKKCFDASF	16
100	KFRCGETRL	16
119	CLQKKDCCA	16
126	CADYKSVQCQ	16
186	KLKTCGIHS	16
256	WLTAMYQGL	16
302	LLKWLDLPK	16
338	VIKALQVVD	16
467	WLAVRSKSN	16
469	AVRSKSNTN	16
601	TATVKVNLP	16
610	FGRPRVLQK	16
657	LPPTVPDCL	16
684	ADKNITHGF	16
736	HATERNGVN	16
793	TPENCPGWL	16
860	EILQLKTYL	16
33	VIMSLGLGL	15
51	QGSCRKKCF	15
191	GIHSKYMRA	15
240	SSKEQNNPA	15
298	RISTLLKWL	15
388	DYFPRINFF	15
626	YHREYVSGF	15

Table XXVIII-V2-B08-9mers-161P2F10B		
Pos	123456789	score
6	CLQRKDCCA	16
8	QRKDCCADY	10
5	DCLQRKDCC	8
7	LQRKDCCAD	7

Table XXVIII-V3-HLA-B08-9mers-161P2F10B		
Pos	123456789	score
7	CPGGKPEAL	27

Table XXVIII-V4-HLA-B08-9mers-161P2F10B		
Pos	123456789	score

Table XXVIII-V4-HLA-B08-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	12

Table XXIX-V1-HLA-B1510-9mers-161P2F10B		
Pos	123456789	score
192	IHSKYMRA	24
578	PHLQNSTQL	22
208	NHYTIVTGL	21
535	THGSLNHLL	21
361	LHNCVNIL	20
551	SHAEVSKF	19
626	YHREYVSGF	19
328	GHAGGPVSA	17
408	AHNIPHDF	17
220	SHGIIDNNM	16
600	ITATVKVNL	16
608	LPFGRPRVL	16
776	IPTHYFVVL	16
836	IARVRDVEL	16
155	GFDLPPVIL	15
179	TLMPNINKL	15
419	NSEEIVRNL	15
809	PHRPTNVES	15
10	EQPVKKNTL	14
21	YKIACIVL	14
40	GLGLGLRKL	14
294	PFEERISTL	14
440	LTPDLPKRL	14
448	LHYAKNVRI	14
757	GHFDAPDEI	14
818	CPEGKPEAL	14
23	IACIVLLAL	13
100	KFRCGETRL	13
250	WHGQPMWLT	13
334	VSARVIKAL	13
346	DHAFGMLME	13
362	HNCVNIILL	13
434	QHFKPYLTP	13
453	NVRIDKVHL	13
498	AHGPFKEK	13
643	WSSYTPVQL	13
767	KHLANTDVP	13
778	THYFVVLTS	13
793	TPENCPGWL	13
796	NCPGWLDVL	13
834	AHIARVRDV	13
860	EILQLKTYL	13
20	KYKIACIVL	12
58	CFDASFRGL	12
150	SQCPEGFDL	12
172	EYLYTWDTL	12
225	DNNMYDVNL	12
249	WWHGQPMWL	12
353	MEGLKQRNL	12
432	PDQHFKPYL	12
516	EVYNLMCDL	12

WO 03/040340

PCT/US02/36002

164

Table XXIX-V1-HLA-B1510-9mers-161P2F10B		
Pos	123456789	score
534	GTHGSLNHL	12
540	NHLLKVPFY	12
585	QLEQVNQML	12
587	EQVNQMLNL	12
621	DHCLLYHRE	12
675	ESQKCSFYL	12
689	THGFLYPFA	12
701	TSDSQYDAL	12
706	YDALITSNL	12
724	MWDYFHSV	12
725	WDYFHSVLL	12
728	FHSVLLIKH	12
735	KHATERNGV	12
791	SHTPENCPG	12
854	KVQPVSEIL	12
24	ACIVLLALL	11
29	LALLVIMSL	11
35	MSLGLGLGL	11
37	LGLGLGLGL	11
104	GETRLEASL	11
112	LCSCSDDCL	11
132	VCQGETSWL	11
295	FEERISTLL	11
298	RISTLLKWL	11
348	AFGMLMEGL	11
436	FKPYLTPLD	11
460	HLFVDQQWL	11
482	NHGYNNEFR	11
489	FRSMEAIFL	11
512	FENIEVYNL	11
531	PNNGTHGSL	11
560	SVCGFANPL	11
565	ANPLPTESL	11
572	SLDCFCPHL	11
617	QKNVDHCLL	11
649	PQLGDTSP	11
657	LPPTVPDCL	11
761	APDEITKHL	11
837	ARVRDVELL	11
840	RDVELLTGL	11
856	QPVSEILQL	11

Table XXIX-V2-B1510-9mers:161P2F10B		
Pos	123456789	score
2	CSDDCLQRK	3
1	SCSDDCLQR	2
3	SDDCLQRKD	2
5	DCLQRKDCC	2
7	LQRKDCCAD	2
4	DDCLQRKDC	1
8	QRKDCCADY	1

Table XXIX-V3-HLA-B1510-9mers-161P2F10B		
Pos	123456789	score
7	CPGGKPEAL	13

Table XXIX-V4-HLA-B1510-9mers-161P2F10B		
Pos	123456789	score
1	TYLPTFETP	4
2	YLPTFETPI	1

Table XXX-V1-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
93	TRIWMCNKF	25
454	VRIDKVHLF	24
830	ERFTHAIAR	24
429	CRKPDQHF	23
489	FRSMEAIFL	23
837	ARVRDVELL	23
77	DRGDCCWDF	21
391	PRINFFYMY	21
638	MRMPMWSSY	21
312	ERPRFYTM	20
721	FRKMWDYFH	20
840	RDVELLTGL	20
29	LALLVIMSL	19
40	GLGLGLRKL	19
525	LRIQPAPNN	19
614	RVLQKNVDH	19
39	LGLGLGLRK	18
100	KFRCGETRL	18
388	DYFPRINFF	18
470	VRKSNTNC	18
481	GNHGYNNEF	18
534	GTHGSLNHL	18
578	PHLQNSTQL	18
632	SGFGKAMRM	18
670	RVPPSESQK	18
692	FLYPPASNR	18
732	LLIKHATER	18
860	EILQLKTYL	18
11	QPVKKNTLK	17
38	GLGLGLGLR	17
42	GLGLRKLEK	17
48	LEKQGSCRK	17
55	RKKCFDASF	17
178	DTLMPNINK	17
179	TLMPNINKL	17
227	NMYDVNLNK	17
262	QGLKAATYF	17
374	GMDQTYCNK	17
446	KRLHYAKNV	17
447	RLHYAKNVR	17
602	ATVKVNLPF	17
611	GRPRVLQKN	17
628	REYVSGFGK	17
856	QPVSEILQL	17

Table XXX-V1-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
21	YKIACIVLL	16
24	ACIVLLALL	16
35	MSLGLGLGL	16
37	LGLGLGLGL	16
104	GETRLEASL	16
155	GFDLPPVIL	16
156	FDLPPVILF	16
187	LKTCGIH	16
188	KTCGIH	16
198	RAMYPTKTF	16
208	NHYTIVTGL	16
287	MPYNGSVPF	16
294	PFEERISTL	16
297	ERISTLLKW	16
298	RISTLLKWL	16
336	ARVIKALQV	16
353	MEGLKQRNL	16
387	TDYFPRINF	16
397	YMYEGPAPR	16
510	EPFENIEVY	16
551	SHAEVSKF	16
600	ITATVKVNL	16
608	LPFGRPRVL	16
739	ERNGVNVVS	16
760	DAPDEITKH	16
6	TLATEQPVK	15
12	PVKKNTLKK	15
47	KLEKQGSCR	15
49	EKQGSCRKK	15
63	FRGLENCRC	15
94	RIWMCNKFR	15
106	TRLEASLCS	15
162	ILFSMDGFR	15
190	CGIHSKYMR	15
229	YDVNLNKNF	15
231	VNLNKNFSL	15
300	STLLKWLDL	15
305	WDLPLKAER	15
313	RPRFYTM	15
329	HAGGPVSAR	15
332	GPVSARVIK	15
351	MLMEGLKQR	15
359	RNLHNCVNI	15
401	GPAPRIRAH	15
406	IRAHNIPHD	15
407	RAHNIPHDF	15
419	NSEEIVRNL	15
423	IVRNLS	15
438	PYLTPDLPK	15
439	YLTPDLPKR	15
460	HLFVDQQWL	15
463	VDQQWLAVR	15
495	IFLAHG	15
512	FENIEVYNL	15
613	PRVLQKNVD	15
649	PQLGDTSP	15

WO 03/040340

PCT/US02/36002

165

Table XXX-V1-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
669	VRVPPSEQ	15
706	YDALITSNL	15
782	VVLTSCKNK	15
802	DVLPFIIPH	15
803	VLPFIIPHR	15
824	EALWVEERF	15
854	KVQPVSEIL	15
7	LATEQPVKK	14
10	EQPVKKNTL	14
14	KKNTLKKYK	14
15	KNTLKKYKI	14
20	KYKIACIVL	14
23	IACIVLLAL	14
56	KKCFDASFR	14
62	SFRGLENCR	14
69	CRCDVACKD	14
70	RCDVACKDR	14
99	NKFRCGETR	14
132	VCQGETSWL	14
169	FRAEYLYTW	14
197	MRAMYPTKT	14
204	KTFPNHYTI	14
215	GLYPESHGI	14
225	DNNMYDVNL	14
234	NKNFSLSSK	14
252	GQPMWLTAM	14
274	SEVAINGSF	14
295	FEERISTLL	14
308	LPKAERPRF	14
314	PRFYTMFE	14
334	VSARVIKAL	14
362	HNCVNIILL	14
405	RIRAHNIPH	14
424	VRNLSCRKP	14
428	SCRKPDQHF	14
452	KNVRIDKVH	14
498	AHGPFKEK	14
504	KEKTEVEPF	14
518	YNLMCDLLR	14
533	NGTHGSLNH	14
539	LNHLLKVPF	14
540	NHLLKVPFY	14
560	SVCGFANPL	14
565	ANPLPTESL	14
584	TQLEQVNQM	14
585	QLEQVNQML	14
587	EQVNQMLNL	14
597	QEEITATVK	14
610	FGRPRVLQK	14
631	VSGFGKAMR	14
678	KCSFYLADK	14
716	PMYEEFRKM	14
725	WDYFHSVLL	14
745	VVSGPIFDY	14
771	NTDVPIPTH	14
796	NCPGWLDVL	14

Table XXX-V1-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
799	GWLDVLPFI	14
836	IARVRDVEL	14
839	VRDVELLTG	14
842	VELLTGLDF	14
26	IVLLALLVI	13
27	VLLALLVIM	13
31	LLVIMSLGL	13
45	LRKLEKQGS	13
86	EDTCVESTR	13
101	FRCGETRLE	13
115	CSDDCLQKK	13
122	KKDCCADYK	13
161	VILFSMDGF	13
172	EYLYTWDTL	13
173	YLYTWDTLM	13
201	YPTKTFPNH	13
220	SHGIIDNNM	13
249	WWHGQPMWL	13
261	YQGLKAATY	13
291	GSVPFEERI	13
302	LLKWLDLPK	13
341	ALQVVDHAF	13
354	EGLKQRNLH	13
358	QRNLHNCVN	13
361	LHNCVNIIL	13
365	VNIILLADH	13
367	IILLADHGM	13
371	ADHGMDQTY	13
377	QTYCNKMEY	13
398	MYEGPAPRI	13
404	PRIRAHNIP	13
417	SFNSEEIVR	13
422	EIVRNLSCR	13
440	LTPDLPKRL	13
442	PDLPKRLHY	13
448	LHYAKNVRI	13
465	QQWLAVRSK	13
477	NCGGGNHGY	13
484	GYNNEFRSM	13
496	FLAHGPSFK	13
513	ENIEVYNLM	13
516	EVYNLMCDL	13
531	PNNGTHGSL	13
550	PSHAEVSK	13
568	LPTESLDCF	13
606	VNLPFGRPR	13
620	VDHCLLYHR	13
626	YHREYVSGF	13
643	WSSYTPVQL	13
662	PDCLRADVR	13
675	ESQKCSFYL	13
713	NLVPMYEEF	13
715	VPMYEEFRK	13
719	EEFRKMWDY	13
727	YFHSVLLIK	13
728	FHSVLLIKH	13

Table XXX-V1-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
743	VNVVSGPIF	13
751	FDYNYDGHF	13
757	GHFDAPDEI	13
759	FDAPDEITK	13
761	APDEITKHL	13
772	TDVPIPTHY	13
776	IPTHYFVVL	13
780	YFVLTSCCK	13
798	PGWLDVLPF	13
818	CPEGKPEAL	13
823	PEALWVEER	13
832	FTAHARVR	13
843	ELLTGLDFY	13
853	DKVQPVSEI	13
857	PVSEILQLK	13
859	SEILQLKTY	13
863	QLKTYLPTF	13
33	VIMSLGLGL	12
54	CRKKCFDAS	12
68	NCRCDVACK	12
87	DTCVESTRI	12
92	STRIWMCNK	12
112	LCSCSDDCL	12
114	SCSDDCLQK	12
148	QQSQCPGEF	12
150	SQCPEGFDL	12
154	EGFDLPPVI	12
165	SMDGFRAEY	12
167	DGFRAEYLY	12
180	LMPNINKLK	12
185	NKLKTCGIH	12
196	YMRAMYPTK	12
221	HGIIDNNMY	12
257	LTAMYQGLK	12
277	AINGSFPSI	12
280	GSFPSIYMP	12
290	NGSVPFEER	12
296	EERISTLLK	12
307	DLPKAERPR	12
311	AERPRFYTM	12
331	GGPV SARVI	12
339	IKALQVVDH	12
348	AFGMLMEGL	12
382	KMEYMTDYF	12
408	AHNIPHDF	12
431	KPDQHFKPY	12
432	PDQHFKPYL	12
436	FKPYLTPDL	12
444	LPKRLHYAK	12
450	YAKNVRIDK	12
453	NVRIDKVHL	12
475	NTNCGGGNH	12
535	THGSLNHLL	12
536	HGSLNHLLK	12
544	KVPFYEPSH	12
556	VSKFSVCGF	12

WO 03/040340

PCT/US02/36002

166

Table XXX-V1-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
604	VKVNLPFGR	12
616	LQKNVDHCL	12
618	KNVDHCLLY	12
622	HCLLYHREY	12
627	HREYVSGFG	12
658	PPTVPDCLR	12
665	LRADVRVPP	12
673	PSESQKCSF	12
674	SESQKCSFY	12
680	SFYLADKNI	12
682	YLADKNITH	12
684	ADKNITHGF	12
686	KNITHGFLY	12
698	SNRTSDSQY	12
710	ITSNLVPMY	12
720	EFRKMWDYF	12
724	MWDYFHSVL	12
726	DYFHSVLLI	12
747	SGPIFDYNY	12
810	HRPTNVESC	12
846	TGLDFYQDK	12

Table XXX-V2-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
8	QRKDCCADY	19
9	RKDCCADYK	15
2	CSDDCLQRK	13
1	SCSDDCLQR	12

Table XXX-V3-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
7	CPGGKPEAL	13
3	NVESCPCGK	10
6	SCPCGKPEA	7

Table XXX-V4-HLA-B2705-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	8
1	TYLPTFETP	4

Table XXXI-V1-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
446	KRLHYAKNV	24
837	ARVRDVELL	24
336	ARVIKALQV	23
489	FRSMEAIFL	21
454	VRIDKVHLF	20
93	TRIWMCNKF	19
77	DRGDCCWDF	18
359	RNLHNCVNI	16
840	RDVELLTGL	15

Table XXXI-V1-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
19	KKYKIACIV	14
104	GETRLEASL	14
155	GFDLPPVIL	14
204	KTFPNHYTI	14
208	NHYTIVTGL	14
215	GLYPESHGI	14
291	GSVPFEERI	14
298	RISTLLKWL	14
525	LRIQPAPNN	14
537	GSLNHLLKV	14
799	GWLDVLPI	14
831	RFTAHIAV	14
856	QPVSEILQL	14
35	MSLGLGLGL	13
37	LGLGLGLGL	13
40	GLGLGLRKL	13
100	KFRCGETRL	13
106	TRLEASLCS	13
297	ERISTLLKW	13
300	STLLKWLDL	13
314	PRFYTMFE	13
337	RVIKALQVV	13
407	RAHNIPHDF	13
419	NSEEIVRNL	13
455	RIDKVHLFV	13
460	HLFVDQQWL	13
512	FENIEVYNL	13
534	GTHGSLNHL	13
578	PHLQNSTQL	13
587	EQVNQMLNL	13
600	ITATVKVNL	13
611	GRPRVLQKN	13
612	RPRVLQKNV	13
643	WSSYTPVQL	13
649	PQLGDTSP	13
699	NRTSDSQYD	13
725	WDYFHSVLL	13
757	GHFDAPDEI	13
776	IPTHYFVVL	13
854	KVQPVSEIL	13
15	KNTLK KYKI	12
20	KYKIACIVL	12
21	YKIACIVLL	12
23	IACIVLLAL	12
24	ACIVLLALL	12
26	IVLLALLVI	12
29	LALLVIMSL	12
31	LLVIMSLGL	12
33	VIMSLGLGL	12
55	RKKCFDASF	12
156	FDLPPVILF	12
172	EYLYTWDTL	12
198	RAMYPTKTF	12
225	DNNMYDVNL	12
231	VNLNKNFSL	12
313	RPRFYTMFE	12

Table XXXI-V1-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
331	GGPV SARVI	12
391	PRINFFYMY	12
406	IRAHNIPHD	12
429	CRKPDQHFK	12
448	LHYAKNVRI	12
453	NVRIDKVHL	12
495	IFLAHGPSF	12
516	EVYNLMCDL	12
584	TQLEQVNQM	12
608	LPFGRPRVL	12
632	SGFGKAMRM	12
663	DCLRADVRV	12
669	VRVPPSESQ	12
706	YDALITSNL	12
726	DYFHSVLLI	12
761	APDEITKHL	12
830	ERFTAHIAV	12
836	IARVRDVEL	12
860	EILQLKTYL	12
25	CIVLLALLV	11
27	VLLALLVIM	11
63	FRGLENCRC	11
65	GLENCRCDV	11
69	CRCDVACKD	11
150	SQCPEGFDL	11
154	EGFDLPPVI	11
166	MDGFRAEYL	11
169	FRAEYLYTW	11
179	TLMPNINKL	11
252	GQPMWLTAM	11
256	WLTAMYQGL	11
262	QGLKAATYF	11
287	MPYNGSVPF	11
295	FEERISTLL	11
311	AERPRFYTM	11
330	AGGPVSARV	11
348	AFGMLMEGL	11
353	MEGLKQRNL	11
367	IILLADHGM	11
385	YMTDYFPRI	11
387	TDYFPRINF	11
388	DYFPRINFF	11
398	MYEGPAPRI	11
403	APRIRAHNI	11
404	PRIRAHNIP	11
432	PDQHFKPYL	11
436	FKPYLTPDL	11
440	LTPDLPKRL	11
470	VRSKSNTNC	11
481	GNHGYNNEF	11
484	GYNNEFRSM	11
504	KEKTEVEPF	11
519	NLMCDLLRI	11
535	THGSLNHLL	11
560	SVCGFANPL	11
565	ANPLPTESL	11

WO 03/040340

PCT/US02/36002

167

Table XXXI-V1-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
572	SLDCFCPHL	11
602	ATVKVNLPF	11
613	PRVLQKNVD	11
617	QKNVDHCLL	11
638	MRMPMWSSY	11
665	LRADVRVPP	11
680	SFYLADKNI	11
701	TSDSQYDAL	11
709	LITSNLVPM	11
721	FRKMWDYFH	11
737	ATERNGVNV	11
739	ERNGVNVVS	11
742	GVNVVSGPI	11
774	VIPTHYFV	11
798	PGWLDVLPF	11
810	HRPTNVESC	11
820	EGKPEALWV	11
824	EALWVEERF	11
834	AHIARVRDV	11
839	VRDVELLTG	11
842	VELLTGLDF	11
847	GLDFYQDKV	11

Table XXXI-V2-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
8	QRKDCCADY	10
9	RKDCCADYK	5

Table XXXI-V3-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
9	GGKPEALWV	14
7	CPGGKPEAL	10

Table XXXI-V4-HLA-B2709-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	8
1	TYLPTFETP	3

Table XXXII-V1-HLA-B4402-9mers-161P2F10B		
Pos	123456789	score
859	SEILQLKTY	29
719	EEFRKMWDY	25
242	KEQNNPAWW	24
487	NEFRSMEAI	24
295	FEERISTLL	23
504	KEKTEVEPF	23
842	VELLTGLDF	23
274	SEVAINGSF	22
353	MEGLKQRNL	22
674	SESQKCSFY	22
104	GETRLEASL	21

Table XXXII-V1-HLA-B4402-9mers-161P2F10B		
Pos	123456789	score
512	FENIEVYNL	21
819	PEGKPEALW	21
179	TLMPNINKL	20
388	DYFPRINFF	20
828	VEERFTAH	20
311	AERPRFYTM	19
507	TEVEPFENI	19
510	EPFENIEVY	19
598	EEITATVKV	19
21	YKIACIVLL	18
24	ACIVLLALL	18
297	ERISTLLKW	18
334	VSARVIKAL	18
761	APDEITKHL	18
156	FDLPPVILF	17
421	EEIVRNLS	17
454	VRIDKVHLF	17
608	LPFGRPRVL	17
684	ADKNITHGF	17
763	DEITKHLAN	17
1	MESTLTLAT	16
90	VESTRIWMC	16
204	KTFPNHYTI	16
341	ALQVVDHAF	16
362	HNCVNIILL	16
442	PDLPKRLHY	16
565	ANPLPTESL	16
686	KNITHGFLY	16
837	ARVRDVELL	16
9	TEQPVKKNT	15
10	EQPVKKNTL	15
13	VKKNTLKKY	15
29	LALLVIMSL	15
93	TRIWMCNKF	15
154	EGFDLPPVI	15
171	AEYLYTWD	15
198	RAMYPTKTF	15
218	PESHGIIDN	15
294	PFEERISTL	15
296	EERISTLLK	15
298	RISTLLKWL	15
371	ADHGMDQTY	15
391	PRINFFYMY	15
399	YEGPAPRIR	15
431	KPDQHFKPY	15
440	LTPDLPKRL	15
745	VVSGPIFDY	15
772	TDVPIPTHY	15
796	NCPGWLDVL	15
843	ELLTGLDFY	15
856	QPVSEILQL	15
23	IACIVLLAL	14
40	GLGLGLRKL	14
51	QGSCRKKCF	14
141	EENCDDTAQQ	14
150	SQCPEGFDL	14

Table XXXII-V1-HLA-B4402-9mers-161P2F10B		
Pos	123456789	score
155	GFDLPPVIL	14
165	SMDGFRAEY	14
167	DGFRAEYLY	14
172	EYLYTWDTL	14
188	KTCGIHSKY	14
208	NHYTIVTGL	14
221	HGIIDNNMY	14
241	SKEQNNPAW	14
300	STLLKWLDL	14
312	ERPRFYTMY	14
322	EEPSSGHA	14
403	APRIRAHNI	14
407	RAHNIPHDF	14
419	NSEEIVRNL	14
420	SEEIVRNLS	14
540	NHLLKVPFY	14
554	EEVSKFSVC	14
560	SVCGFANPL	14
587	EQVNQMLNL	14
602	ATVKVNLPF	14
618	KNVDHCLLY	14
638	MRMPMWSSY	14
713	NLVPMYEEF	14
738	TERNGVNVV	14
818	CPEGKPEAL	14
829	EERFTAHIA	14
854	KVQPVSEIL	14

Table XXXII-V2-HLA-B4402-9mers-161P2F10B		
Pos	123456789	score
8	QRKDCCADY	11
1	SCSDDCLQR	5

Table XXXII-V3-HLA-B4402-9mers-161P2F10B		
Pos	123456789	score
7	CPGGKPEAL	14
4	VESCPGGKP	12
8	PGGKPEALW	11
5	ESCPGGKPE	7

Table XXXII-V4-HLA-B4402-9mers-161P2F10B		
Pos	123456789	score
2	YLPTFETPI	10
1	TYLPTFETP	6

Table XXXIII-V1-HLA-B5101-9mers-161P2F10B		
Pos	123456789	score
707	DALITSNLV	26
608	LPFGRPRVL	25
776	IPTHYFVVL	24
29	LALLVIMSL	23

Table XXXIII-V1-HLA-B5101-9mers-
161P2F10B

Pos	123456789	score
331	GGPV SARVI	23
657	LPPTV PDCL	23
23	IACIVLLAL	22
154	EGFDLPPIVI	22
640	MPMWSSYTV	22
403	APRIRAHNI	21
448	LHYAKNVRI	21
760	DAPDEITKH	20
761	APDEITKHL	20
774	VPIPTHYFV	20
26	IVLLALLVI	19
612	RPRVLQKNV	19
661	VPDCLRADV	19
694	YPPASNRTS	19
726	DYFHSVLLI	19
818	CPEGKPEAL	19
836	IARVRDVEL	19
7	LATEQPVKK	18
37	LGLGLGLGL	18
793	TPENC PGWL	18
856	QPVSEILQL	18
330	AGGPVSARV	17
340	KALQVVDHA	17
510	EPFENIEVY	17
867	YLPTFETTI	17
87	DTCVESTRI	16
206	FPNHYTIVT	16
216	LYPESHGII	16
287	MPYNGSVPF	16
385	YMTDYFPRI	16
437	KPYLTPDLP	16
568	LPTESLDCF	16
663	DCLRADV RV	16
683	LADKNITHG	16
736	HATERNGVN	16
799	GWLDVLPFI	16
808	IPHRPTNVE	16
820	EGKPEALWV	16
853	DKVQP VSEI	16
60	DASFRGLEN	15
201	YPTKTFPNH	15
208	NHYTIVTGL	15
270	FWPGSEVAI	15
293	VPFEERIST	15
450	YAKNVRIDK	15
497	LAHGPSFKE	15
552	HAEEVSKFS	15
601	TATVKVNLP	15
653	DTSPLPPTV	15
680	SFYLADKNI	15
738	TERNGVNVV	15
769	LANTDVPIPI	15
804	LPFIIPHRP	15
824	EALWVEERF	15
5	LTLATEQPV	14
11	QPVKKNTLK	14

Table XXXIII-V1-HLA-B5101-9mers-
161P2F10B

Pos	123456789	score
18	LKKYKIACI	14
19	KKYKIACIV	14
43	LGLRKLEKQ	14
124	DCCADYKSV	14
158	LPPVILFSM	14
181	MPNINKLKT	14
215	GLYPESHGI	14
258	TAMYQGLKA	14
308	LPKAERPRF	14
347	HAFGMLMEG	14
360	NLHNCVNII	14
398	MYEGPAPRI	14
411	IPHDFFSFN	14
500	GPSFKEKTE	14
507	TEVEPFENI	14
519	NLMCDLLRI	14
537	GSLNHLLKV	14
545	VPFYEPSHA	14
566	NPLPTESLD	14
671	VPPSESQKC	14
672	PPSESQKCS	14
15	KNTLK KYKI	13
39	LGLGLGLRK	13
184	INKLKTCGI	13
198	RAMYPTKTF	13
204	KTFPNHYTI	13
217	YPESHGIID	13
265	KAATYFWPG	13
277	AINGSFPSI	13
329	HAGGPVSAR	13
337	RVIKALQVV	13
359	RNLHNCVNI	13
440	LTPDLPKRL	13
446	KRLHYAKNV	13
468	LAVRSKSNT	13
487	NEFRSMEAI	13
528	QPAPNNGTH	13
529	PAPNNGTHG	13
549	EP SHAEVS	13
564	FANPLPTES	13
577	CPHLQNSTQ	13
596	TQEEITATV	13
655	SPLPPTVPD	13
757	GHFDAPDEI	13
775	PIPTHYFVV	13
796	NCPGWLDVL	13
807	IIPHRPTNV	13
833	TAHIARVRD	13
850	FYQDKVQPV	13
41	LGLGLRKLE	12
73	VACKDRGDC	12
126	CADYKSV CQ	12
146	TAQQSQCPE	12
152	CPEGFDLPP	12
176	TWDTLMPNI	12
225	DNNMYDVNL	12

Table XXXIII-V1-HLA-B5101-9mers-
161P2F10B

Pos	123456789	score
271	WPGSEVAIN	12
272	PGSEVAING	12
282	FPSIYMPYN	12
285	IYMPYNGSV	12
361	LHNCVNIIIL	12
370	LADHGMDQT	12
390	FPRINFFYM	12
401	GPAPRIRAH	12
402	PAPRIRAHN	12
419	NSEEIVRNL	12
431	KPDQHF KPY	12
441	TPDLPKRLH	12
444	LPKRLHYAK	12
455	RIDKVHLFV	12
509	VEPFENIEV	12
548	YEP SHAEV	12
592	MLNLTQEEI	12
598	EEITATVKV	12
600	ITATVKVNL	12
636	KAMRMPMWS	12
648	VPQLGDTSP	12
651	LGDTSP LPP	12
666	RADVRVPPS	12
715	VPMYEEFRK	12
723	KMWDFH SV	12
742	GVNVVSGPI	12
768	HLANTDVPI	12
778	THYFVVLTS	12
811	RPTNVESCP	12
822	KPEALWEE	12
828	VEERF TAHI	12
834	AHIARVRDV	12

Table XXXIII-V2-HLA-B5101-9mers-
161P2F10B

Pos	123456789	score
5	DCLQRKDCC	6
4	DDCLQRKDC	5
2	CSDDCLQRK	3
3	SDDCLQRKD	3
7	LQRKDCCAD	3
1	SCSDDCLQR	2

Table XXXIII-V3-HLA-B5101-9mers-
161P2F10B

Pos	123456789	score
7	CPGGKPEAL	19
9	GGKPEALWV	16

Table XXXIII-V4-HLA-B5101-9mers-
161P2F10B

Pos	123456789	score
2	YLPTFETPI	15
1	TYLPTFETP	7

Table XXXIV-V1-HLA-A1-10MERS-161P2F10B		
Pos	1234567890	score
441	TPDLPKRLHY	32
771	NTDVPIPTHY	31
858	VSEILQLKTY	30
673	PSESQKCSFY	29
718	YEEFRKMWDY	27
370	LADHGMDQTY	25
746	VSGPIFDYNY	24
280	GSFPSIYMPY	23
617	QKNVDHCLLY	23
164	FSMDGFRAEY	21
311	AERPRFYTMY	21
386	MTDYFPRINF	21
388	DYFPRINFFY	20
390	FPRINFFYMY	20
697	ASNRTSDSQY	20
220	SHGIIDNNMY	19
295	FEERISTLLK	19
420	SEEIVRNLS	19
506	KTEVEPFENI	19
508	EVEPFENIEV	19
737	ATERNGVNVV	19
430	RKPDQHFKPY	18
685	DKNITHGFLY	18
800	WLDVLPFIIP	18
8	ATEQPVKKNT	17
12	PVKKNTLKKY	17
155	GFDLPPVILF	17
166	MDGFRAEYLY	17
187	LKTCGIHSKY	17
192	IHSKYMRAMY	17
201	YPTKTFPNHY	17
208	NHYTIVTGLY	17
277	AINGSFPSIY	17
419	NSEEIVRNLS	17
476	TNCGGNGHGY	17
509	VEPFENIEVY	17
709	LITSNLVPMY	17
841	DVELLTGLDF	17
115	CSDDCLQKKD	16
217	YPESHGIIDN	16
252	GQPMWLTAMY	16
260	MYQGLKAATY	16
539	LNHLLKVPFY	16
569	PTESLDCFCP	16
637	AMRMPMWSSY	16
701	TSDSQYDALI	16
762	PDEITKHLAN	16
842	VELLTGLDFY	16
120	LQKKDCCADY	15
257	LTAMYQGLKA	15
273	GSEVAINGSF	15
308	LPKAERPRFY	15
321	FEEDSSGHA	15
376	DQTYCNKMEY	15
380	CNKMEYMTDY	15
547	FYEPShAEV	15

Table XXXIV-V1-HLA-A1-10MERS-161P2F10B		
Pos	1234567890	score
553	AEEVSKFSVC	15
572	SLDCFCPHLQ	15
621	DHCLLYHREY	15
744	NVSGPIFDY	15

Table XXXIV-V2-HLA-A1-10mers-161P2F10B		
Pos	1234567890	score
3	CSDDCLQRKD	16
8	LQRKDCCADY	15
4	SDDCLQRKDC	13
10	RKDCCADYKS	12
1	CSCSDDCLQR	10

Table XXXIV-V3-HLA-A1-10mers-161P2F10B		
Pos	1234567890	score
4	NVESCPGGKP	13
6	ESCPGGKPEA	8
2	PTNVESCPGG	6
9	PGGKPEALWV	6

Table XXXIV-V4-HLA-A1-10mers-161P2F10B		
Pos	1234567890	score
1	KTYLPTFETP	9
2	TYLPTFETPI	4

Table XXXV-V1-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
28	LLALLVIMSL	30
22	KIACIVLLAL	27
30	ALLVIMSLGL	26
36	SLGLGLGLGL	24
708	ALITSNLVPM	24
835	HIARVRDVEL	24
17	TLKKYKIACI	23
222	GIIDNNMYDV	23
284	SIYMPYNGSV	23
806	FIIPHRPTNV	23
23	IACIVLLALL	22
34	IMSLGLGLGL	22
39	LGLGLGLRKL	22
111	SLCSCSDDCL	22
165	SMDGFRAEYL	22
178	DTLMPNINKL	22
215	GLYPESHGII	22
302	LLKWLDLPKA	22
397	YMYEGPAPRI	22
439	YLTPDLPKRL	22
595	LTQEEITATV	22
615	VLQKNVDHCL	22
25	CIVLLALLVI	21
157	DLPPVILFSM	21

Table XXXV-V1-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
352	LMEGLKQRNL	21
450	YAKNVRIDKV	21
639	RMPMWSSYTV	21
723	KMWDFHSHVL	21
32	LVIMSLGLGL	20
293	VPFEERISTL	20
360	NLHNCVNIL	20
564	FANPLPTESL	20
591	QMLNLTQEEI	20
737	ATERNGVNVV	20
836	IARVRDVELL	20
4	TLTLATEQPV	19
369	LLADHGMDQT	19
447	RLHYAKNVRI	19
607	NLPFGRPRVL	19
656	PLPPTVPDCL	19
682	YLADKNITHG	19
692	FLYPPASNRT	19
825	ALWVEERFTA	19
26	IVLLALLVIM	18
183	NINKLKTCGI	18
204	KTFPNHYTIV	18
259	AMYQGLKAAT	18
267	ATYFWPGSEV	18
276	VAINGSFPSI	18
329	HAGGPVSARV	18
347	HAFGMLMEGL	18
534	GTHGSLNHLL	18
594	NLTQEEITAT	18
833	TAHIARVRDV	18
27	VLLALLVIMS	17
42	GLGLRKLEKQ	17
162	ILFSMDGFRA	17
175	YTWDTLMPNI	17
180	LMPNINKLKT	17
186	KLKTCGIHSK	17
227	NMYDVNLNKN	17
335	SARVIKALQV	17
361	LHNCVNILL	17
418	FNSEEIVRN	17
518	YNLMCDLLRI	17
536	HGSLNHLLKV	17
552	HAEEVSKFSV	17
580	LQNSTQLEQV	17
624	LLYHREYVSG	17
660	TVPDCLRADV	17
736	HATERNGVNV	17
768	HLANTDVPIP	17
775	PIPTHYFVVL	17
861	ILQLKTYLPT	17
6	TLATEQPVKK	16
20	KYKIACIVLL	16
31	LLVIMSLGLG	16
131	SVCQGETSWL	16
152	CPEGFDLPPV	16
196	YMRAMYPTKT	16

WO 03/040340

PCT/US02/36002

170

Table XXXV-V1-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
230	DVNLNKNFSL	16
269	YFWPGSEVAI	16
336	ARVIKALQVV	16
366	NIILLADHGM	16
374	GMDQTYCNKM	16
467	WLAVRSKSNT	16
526	RIQPAPNNGT	16
599	EITATVKVNL	16
606	VNLPFGRPRV	16
700	RTSDSQYDAL	16
792	HTPENCPGWL	16
839	VRDVELLTGL	16
855	VQPVSEILQL	16
9	TEQPVKKNTL	15
24	ACIVLLALLV	15
171	AEYLYTWDTL	15
224	IDNNMYDVNL	15
257	LTAMYQGLKA	15
333	PVSARVIKAL	15
343	QVVDHAFGML	15
355	GLKQRNLHNC	15
359	RNLHNCVNII	15
454	VRIDKVHLFV	15
460	HLFVDQQWLA	15
461	LFVDQQWLAV	15
491	SMEAIFLAHG	15
496	FLAHGPSFKE	15
519	NLMCDLLRIQ	15
530	APNNGTHGSL	15
538	SLNHLLKVPF	15
541	HLLKVPFYEP	15
583	STQLEQVNQM	15
592	MLNLTQEEIT	15
642	MWSSYTVPQL	15
706	YDALITSNLV	15
731	VLLIKHATER	15
765	ITKHLANTDV	15
774	VPIPTHYFVV	15
817	SCPEGKPEAL	15
827	WVEERFTAH	15
846	TGLDFYQDKV	15
33	VIMSLGLGLG	14
57	KCFDASFRGL	14
64	RGLENCRCDV	14
65	GLENCRCDA	14
173	YLYTWDTLMP	14
191	GIHSKYMRAM	14
232	NLNKNFSLSS	14
338	VIKALQVVDH	14
341	ALQVVDHAFG	14
350	GMLMEGLKQR	14
368	ILLADHGMDQ	14
494	AIFLAHGPSF	14
508	EVEPFENIEV	14
533	NGTHGSLNHL	14
547	FYEPSHAEEV	14

Table XXXV-V1-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
584	TQLEQVNQML	14
709	LITSNLVPMY	14
722	RKMWDYFHSV	14
732	LLIKHATERN	14
734	IKHATERNGV	14
773	DVPIPTHYFV	14
830	ERFTAHIAHV	14
844	LLTGLDFYQD	14
849	DFYQDKVQPV	14
859	SEILQLKTYL	14
866	TYLPTFETTI	14

Table XXXV-V2-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
7	CLQRKDCCAD	11
2	SCSDDCLQRK	7
6	DCLQRKDCCA	5

Table XXXV-V3-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
7	SCPGGKPEAL	15
9	PGGKPEALWV	8

Table XXXV-V4-HLA-A0201-10mers-161P2F10B		
Pos	1234567890	score
2	TYLPTFETPI	12
1	KTYLPTFETP	8

Table XXXVI-V1-HLA-A0203-10mers-161P2F10B		
Pos	1234567890	score
258	TAMYQGLKAA	19
259	AMYQGLKAAT	17
15	KNTLKKYKIA	10
21	YKIACIVLLA	10
52	GSCRKKCFDA	10
65	GLENCRCDA	10
101	FRCGETRLEA	10
118	DCLQKKDCCA	10
138	SWLEENCDA	10
162	ILFSMDGFRA	10
190	CGIHSKYMRA	10
239	LSSKEQNNPA	10
250	WHGQPMWLTA	10
257	LTAMYQGLKA	10
268	TYFWPGSEVA	10
302	LLKWLDLPKA	10
321	FEEDSSGHA	10
327	SGHAGGPVSA	10
332	GPVSARVIKA	10
339	IKALQVVDHA	10
362	HNCVNIILLA	10

Table XXXVI-V1-HLA-A0203-10mers-161P2F10B		
Pos	1234567890	score
394	NFFYMYEGPA	10
399	YEGPAPRIRA	10
442	PDLPKRLHYA	10
460	HLFVDQQWLA	10
485	YNNEFRSMEA	10
489	FRSMEAIFLA	10
521	MCDLLRIQPA	10
544	KVPFYEPSHA	10
556	VSKFSVCGFA	10
593	LNLTQEEITA	10
628	REYVSGFGKA	10
658	PPTVPDCLRA	10
675	ESQKCSFYLA	10
688	ITHGFLYPPA	10
699	NRTSDSQYDA	10
728	FHSVLLIKHA	10
752	DYNYDGHFDA	10
761	APDEITKHLA	10
816	ESCPEGKPEA	10
825	ALWVEERFTA	10
828	VEERFTAHIA	10
16	NTLKKYKIAC	9
22	KIACIVLLAL	9
53	SCRKKCFDAS	9
66	LENCRCDVAC	9
102	RCGETRLEAS	9
119	CLQKKDCCAD	9
139	WLEENCDAQ	9
163	LFSMDGFRAE	9
191	GIHSKYMRAM	9
240	SSKEQNNPAW	9
251	HGQPMWLTAM	9
269	YFWPGSEVAI	9
303	LKWLDLPKAE	9
322	EEPDSGSHAG	9
328	GHAGGPVSAR	9
333	PVSARVIKAL	9
340	KALQVVDHAF	9
363	NCVNIILLAD	9
395	FFYMYEGPAP	9
400	EGPAPRIRAH	9
443	DLPKRLHYAK	9
461	LFVDQQWLAV	9
486	NNEFRSMEAI	9
490	RSMEAIFLAH	9
522	CDLLRIQPAP	9
545	VPFYEPSHAE	9
557	SKFSVCGFAN	9
594	NLTQEEITAT	9
629	EYVSGFGKAM	9
659	PTVPDCLRAD	9
676	SQKCSFYLAD	9
689	THGFLYPPAS	9
700	RTSDSQYDAL	9
729	HSVLLIKHAT	9
753	YNYDGHFDAP	9

Table XXXVI-V1-HLA-A0203-10mers-161P2F10B		
Pos	1234567890	score
762	PDEITKHLAN	9
817	SCPEGKPEAL	9
826	LWVEERFTAH	9
829	EERFTAHAR	9

Table XXXVI-V2-HLA-A0203-10mers-161P2F10B		
Pos	1234567890	score
6	DCLQRKDCCA	10
7	CLQRKDCCAD	9
8	LQRKDCCADY	8

Table XXXVI-V3-HLA-A0203-10mers-161P2F10B		
Pos	1234567890	score
6	ESCPGGKPEA	10
7	SCPGGKPEAL	9
8	CPGGKPEALW	8

Table XXXVI-V4-HLA-A0203-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XXXVII-V1-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
38	GLGLGLGLRK	27
6	TLATEQPVKK	26
186	KLKTCGIHSK	26
301	TLLKWLDLPK	25
337	RVIKALQVVD	25
838	RVRDVELLTG	25
47	KLEKQGSCRK	23
179	TLMPNINKLK	23
256	WLTAMYQGLK	23
277	AINGSFPSY	23
212	IVTGLYPESH	21
422	EIVRNLSCKR	21
437	KPYLTDPDPK	21
443	DLPKRLHYAK	21
462	FVDQQWLAVR	21
494	AIFLAHGPF	21
605	KVNLPFGRPR	21
730	SVLLIKHATE	21
731	VLLIKHATER	21
781	FVVLTSCKNK	21
841	DVELLTGLDF	21
195	KYMRAMYPTK	20
368	ILLADHGMDQ	20
495	IFLAHGPF	20
624	LLYHREYVSG	20
630	YVSGFGKAMR	20
664	CLRADVRVPP	20
708	ALITSNLVPM	20

Table XXXVII-V1-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
714	LVPMEEFK	20
26	IVLLALLVIM	19
32	LVIMSLGLGL	19
107	RLEASLCSCS	19
338	VIKALQVVDH	19
523	DLLRIQPAPN	19
538	SLNHLLKVPF	19
588	QVNQMLNLTQ	19
596	TQEEITATVK	19
609	PFGRPRVLQK	19
669	VRVPPSESQK	19
802	DVLPFIIPHR	19
5	LTATEQPVK	18
30	ALLVIMSLGL	18
36	SLGLGLGLGL	18
160	PVILFSMDGF	18
173	YLYTWDTLMP	18
215	GLYPESHGII	18
260	MYQGLKAATY	18
364	CVNIILLADH	18
426	NLSCRKPDQH	18
447	RLHYAKNVRI	18
453	NVRIDKVHLF	18
697	ASNRTSDSQY	18
825	ALWVEERFTA	18
12	PVKKNTLKKY	17
41	LGLGLRKLEK	17
46	RKLEKQGSCR	17
121	QKKDCCADYK	17
233	LNKNFSLSSK	17
311	AERPRFYTM	17
405	RIRAHNIPHD	17
458	KVHLFVDQW	17
469	AVRSKSNTNC	17
524	LLRIQPAPNN	17
544	KVPFYEPSHA	17
637	AMRMPMWSSY	17
650	QLGDTSPPLP	17
670	RVPPSESQKC	17
692	FLYPASNRT	17
713	NLVPMYEEFR	17
742	GVNVSGPIF	17
831	RFTAHARVR	17
22	KIACIVLLAL	16
25	CIVLLALLVI	16
67	ENCRCDVACK	16
113	CSCSDCLQK	16
284	SIYMPYNGSV	16
295	FEERISTLLK	16
341	ALQVVDHAFG	16
423	IVRNLSCKRP	16
455	RIDKVHLFVD	16
549	EPSHAEVSK	16
603	TVKVNLPFGR	16
614	RVLQKNVDHC	16
623	CLLYHREYVS	16

Table XXXVII-V1-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
744	NVVS GPIFDY	16
758	HFDAPDEITK	16
775	PIPTHYFVVL	16
783	VLTSCKNKSH	16
806	FIIPHRPTNV	16
835	HIARVRDVEL	16
854	KVQPVSEILQ	16
861	ILQLKTYLPT	16
11	QPVKKNTLKK	15
27	VLLALLVIMS	15
28	LLALLVIMSL	15
55	RKKCFDASFR	15
65	GLENCRCDDA	15
131	SVCQGETSWL	15
157	DLPPVILFSM	15
161	VILFSMDGFR	15
162	ILFSMDGFRA	15
223	IIDNNMYDVN	15
232	NLNKNFSLSS	15
285	IYMPYNGSVP	15
307	DLPKAERPRF	15
331	GGPV SARVIK	15
343	QVVDHAFGML	15
367	IILLADHGMD	15
369	LLADHGMDQT	15
516	EVYNLMCDLL	15
526	RIQPAPNNGT	15
607	NLPFGRPRVL	15
660	TVPDCLRADV	15
668	DVRVPPSESQ	15
677	QKCSFYLADK	15
709	LITSNLVPMY	15
764	EITKHLANTD	15
844	LLTGLDFYQD	15
863	QLKTYLPTFE	15
17	TLKKYKIACI	14
263	GLKAATYFWP	14
292	SVPFEERIST	14
305	WLDLPKAERP	14
344	VVDHAFGMLM	14
348	AFGMLMEGLK	14
428	SCRKPDQHF	14
439	YLTDPDLPKRL	14
467	WLAVRSKSNT	14
555	EVSKFSVCGF	14
567	PLPTESLDCF	14
579	HLQNSTQLEQ	14
599	EITATVKVNL	14
619	NVDHCLLYHR	14
656	PLPPTVPDCL	14
807	IIPHRPTNVE	14
827	WVEERFTAH	14
847	GLDFYQDKVQ	14
857	PVSEILQLKT	14
44	GLRKLEKQGS	13
72	DVACKDRGDC	13

Table XXXVII-V1-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
91	ESTRIWMCNK	13
139	WLEENCDAQ	13
192	IHSKYMRAMY	13
226	NNMYDVNLNK	13
230	DVNLNKNFSL	13
267	ATYFWPGSEV	13
304	KWLDLPKAER	13
330	AGGPVSARVI	13
333	PVSARVIKAL	13
335	SARVIKALQV	13
446	KRLHYAKNVR	13
490	RSMEAIFLAH	13
508	EVEPFENIEV	13
527	IQPAPNNGTH	13
535	THGSLNHLLK	13
560	SVCGFANPLP	13
585	QLEQVNQMLN	13
627	HREYVSGFGK	13
647	TVPQLGDTSP	13
682	YLADKNITHG	13
726	DYFHSVLLIK	13
732	LLIKHATERN	13
749	PIFDYNYDGH	13
779	HYFVLTSCK	13
834	AHIARVRDVE	13

Table XXXVII-V2-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
9	QRKDCCADYK	17
7	CLQRKDCCAD	14
2	SCSDDCLQRK	12
8	LQRKDCCADY	12
1	CSCSDDCLQR	10

Table XXXVII-V3-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
4	NVESCPCGGKP	14
3	TNVESCPCGGK	13
10	GGKPEALWVE	8
9	PGGKPEALWV	7

Table XXXVII-V4-HLA-A3-10mers-161P2F10B		
Pos	1234567890	score
1	KTYLPTFETP	12
2	TYLPTFETPI	6

Table XXXVIII-V1-HLA-A26-10mers-161P2F10B		
Pos	1234567890	score
555	EVSKEFSVCGF	32
178	DTLMPNINKL	29
744	NVVS GPIFDY	29

Table XXXVIII-V1-HLA-A26-10mers-161P2F10B		
Pos	1234567890	score
516	EVYNLMCDLL	28
841	DVELLTGLDF	28
230	DVNLNKNFSL	27
297	ERISTLLKWL	27
12	PVKKNTLKKY	25
343	QVVDHAFGML	25
32	LVIMSLGLGL	24
154	EGFDLPPVIL	23
333	PVSARVIKAL	23
388	DYFPRINFFY	23
599	EITATVKVNL	23
795	ENCPGWLDVL	23
802	DVLPFIIPHR	23
160	PVILFSMDGF	22
453	NVRIDKVHLF	22
508	EVEPFENIEV	22
719	EEFRKMWDYF	22
709	LITSNLVPMY	21
131	SVCQGETSWL	20
275	EVAINGSFPS	20
293	VPFEERISTL	20
488	EFRSMEAIFL	20
853	DKVQPVSEIL	20
72	DVACKDRGDC	19
105	ETRLEASLCS	19
376	DQTYCNKMEY	19
421	EEIVRNLSR	19
571	ESLDCFCPHL	19
668	DVRVPPSESQ	19
685	DKNITHGFLY	19
742	GVNVVSGPIF	19
792	HTPENCPGWL	19
22	KIACIVLLAL	18
92	STRIWMCNKF	18
136	ETSWLEENCD	18
312	ERPRFYTMFY	18
422	EIVRNLSRCK	18
494	AIFLAHGPSF	18
513	ENIEVYNLMC	18
587	EQVNQMLNLT	18
598	EEITATVKVN	18
621	DHCLLYHREY	18
659	PTVPDCLRAD	18
700	RTSDSQYDAL	18
771	NTDVPIPTHY	18
773	DVPIPTHYFV	18
830	ERFTAHIARV	18
386	MTDYFPRINF	17
457	DKVHLFVDQQ	17
629	EYVSGFGKAM	17
653	DTSPLPPTVP	17
726	DYFHSVLLIK	17
760	DAPDEITKHL	17
838	RVRDVELLTG	17
87	DTCVESTRIW	16
89	CVESTRIWMC	16

Table XXXVIII-V1-HLA-A26-10mers-161P2F10B		
Pos	1234567890	score
145	DTAQSQSQPE	16
210	YTIVTGLYPE	16
296	EERISTLLKW	16
307	DLPKAERPRF	16
510	EPFENIEVYN	16
534	GTHGSLNHLL	16
574	DCFCPHLQNS	16
602	ATVKVNLPGF	16
763	DEITKHLANT	16
764	EITKHLANTD	16
277	AINGSFPSY	15
280	GSFPSIYMPY	15
400	EGPAPRIRAH	15
493	EAIFLAHGPS	15
554	EEVSKFSVCG	15
646	YTVPQLGDT	15
739	ERNGVNVVSG	15
820	EGKPEALWVE	15
858	VSEILQLKTY	15

Table XXXVIII-V2-HLA-A26-10mers-161P2F10B		
Pos	1234567890	score
8	LQRKDCCADY	10
5	DDCLQRKDCC	8
6	DCLQRKDCCA	8
2	SCSDDCLQRK	6

Table XXXVIII-V3-HLA-A26-10mers-161P2F10B		
Pos	1234567890	score
4	NVESCPCGGKP	14
6	ESCPGGKPEA	11
7	SCPCGGKPEAL	10
2	PTNVESCPCGG	8
3	TNVESCPCGGK	8

Table XXXVIII-V4-HLA-A26-10mers-161P2F10B		
Pos	1234567890	score
1	KTYLPTFETP	9

Table XXXIX-V1-HLA-B0702-10mers-161P2F10B		
Pos	1234567890	score
431	KPDQHF KPYP	23
530	APNNGTHGSL	22
648	VPQLGDTSP	22
577	CPHLQNSTQL	21
797	CPGWLDVLPF	21
152	CPEGFDLPPV	20
293	VPFEERISTL	20
776	IPTHYFVLT	20
500	GPSFKEKTEV	19
761	APDEITKHLA	19

Table XXXIX-V1-HLA-B0702-10mers-161P2F10B		
Pos	1234567890	score
246	NPAWWHGQPM	18
658	PPTVPDCLRA	18
672	PPSESQKCSF	18
715	VPMYEEFRKM	18
774	VPIPTHYFVV	18
332	GPVSARVIKA	17
804	LPFIIPHRPT	17
22	KIACIVLLAL	15
34	IMSLGLGLGL	15
549	EPSHAEVSK	15
608	LPFGRPRVLQ	15
11	QPVKKNTLKK	14
30	ALLVIMSLGL	14
313	RPRFYTMYFE	14
333	PVSARVIKAL	14
401	GPAPRIRAHN	14
403	APRIRAHNIP	14
437	KPYLTPDLPK	14
441	TPDLPKRLHY	14
488	EFRSMEAIFL	14
599	EITATVKVNL	14
612	RPRVLQKNVD	14
642	MWSSYTPVQL	14
700	RTSDSQYDAL	14
775	PIPTHYFVVL	14
836	IARVRDVELL	14
19	KKYKIACIVL	13
20	KYKIACIVLL	13
36	SLGLGLGLGL	13
154	EGFDLPPVIL	13
165	SMDGFRAEYL	13
224	IDNNMYDVNL	13
287	MPYNGSVPFE	13
299	ISTLLKWLDL	13
323	EPDSSGHAGG	13
390	FPRINFFYMY	13
418	FNSEEIVRNL	13
435	HFKPYLTPDL	13
452	KNVRIDKVHL	13
510	EPFENIEVYN	13
528	QPAPNNGTHG	13
566	NPLPTESLDC	13
640	MPMWSSYTVP	13
655	SPLPPTVPDC	13
684	ADKNITHGFL	13
694	YPPASNRTSD	13
705	QYDALITSNL	13
724	MWDYFHSVLL	13
795	ENCPGWLDVL	13
808	IPHRPTNVES	13
818	CPEGKPEALW	13
822	KPEALWVEER	13
835	HIARVRDVEL	13
23	IACIVLLALL	12
32	LVIMSLGLGL	12
39	LGLGLGLRKL	12

Table XXXIX-V1-HLA-B0702-10mers-161P2F10B		
Pos	1234567890	score
57	KCFDASFRGL	12
171	AEYLYTWDTL	12
206	FPNHYTIVTG	12
207	PNHYTIVTGL	12
248	AWWHGQPMWL	12
282	FPSIYMPYNG	12
308	LPKAERPRFY	12
347	HAFGMLMEGL	12
352	LMEGLKQRNL	12
534	GTHGSLNHLL	12
564	FANPLPTESL	12
571	ESLDCFCPHL	12
586	LEQVNQMLNL	12
661	VPDCLRADVR	12
674	SESQKCSFYL	12
695	PPASNRTSDS	12
723	KMWDYFHSVL	12
811	RPTNVESCPE	12
839	VRDVELLTGL	12
855	VQPVSEILQL	12
9	TEQPVKKNTL	11
28	LLALLVIMSL	11
76	KDRGDCCWDF	11
99	NKFRCGETRL	11
101	FRCGETRLEA	11
103	CGETRLEASL	11
131	SVCQGETSWL	11
158	LPPVILFSMD	11
201	YPTKTFPNHY	11
253	QPMWLTAMYQ	11
271	WPGSEVAING	11
294	PFEERISTLL	11
297	ERISTLLKWL	11
325	DSSGHAGGPV	11
330	AGGPVSARVI	11
343	QVVDHAFGML	11
411	IPHDFFSFNS	11
439	YLTPDLPKRL	11
444	LPKRLHYAKN	11
511	PFENIEVYNL	11
515	IEVYNLMCDL	11
516	EVYNLMCDLL	11
545	VPFYEPSHAE	11
555	EVSKFSVCGF	11
559	FSVCGFANPL	11
568	LPTESLDCFC	11
607	NLPFGRPRVL	11
615	VLQKNVDHCL	11
656	PLPPTVPDCL	11
688	ITHGFLYPPA	11
708	ALITSNLVPM	11
760	DAPDEITKHL	11
767	KHLANTDVPI	11
817	SCPEGKPEAL	11
859	SEILQLKTYL	11

Table XXXIX-V2-HLA-B0702-10mers-161P2F10B		
Pos	1234567890	score
6	DCLQRKDCCA	6
8	LQRKDCCADY	4
1	CSCSDDCLQR	2
2	SCSDDCLQRK	2
4	SDDCLQRKDC	2
10	RKDCCADYKS	2
3	CSDDCLQRKD	1
7	CLQRKDCCAD	1

Table XXXIX-V3-HLA-B0702-10mers-161P2F10B		
Pos	1234567890	score
8	CPGGKPEALW	14
7	SCPGGKPEAL	13
1	RPTNVESCPG	12
9	PGGKPEALWV	10
6	ESCPGGKPEA	9

Table XXXIX-V4-HLA-B0702-10mers-161P2F10B		
Pos	1234567890	score
2	TYLPTFETPI	9

Table XL-V1-HLA-B08-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XL-V2-HLA-B08-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XL-V3-HLA-B08-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XL-V4-HLA-B08-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLI-V1-HLA-B1510-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLI-V2-HLA-B1510-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

WO 03/040340

PCT/US02/36002

174

Table XLI-V3-HLA-B1510-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLI-V4-HLA-B1510-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLII-V1-HLA-B2705-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLII-V2-HLA-B2705-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLII-V3-HLA-B2705-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLII-V4-HLA-B2705-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLIII-V1-HLA-B2709-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLIII-V2-HLA-B2709-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLIII-V3-HLA-B2709-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLIII-V4-HLA-B2709-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLIV-V1-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
311	AERPRFYTM	26
9	TEQPVKKNTL	25
171	AEYLYTWDTL	25
296	EERISTLLKW	25

Table XLIV-V1-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
509	VEPFENIEVY	25
719	EEFRKMWDYF	25
859	SEILQLKTYL	25
487	NEFRSMEAIF	24
842	VELLTGLDFY	23
674	SESQKCSFYL	22
586	LEQVNQMLNL	21
823	PEALWVEERF	21
153	PEGFDLPPVI	20
515	IEVYNLMCDL	20
598	EEITATVKVN	20
718	YEEFRKMWDY	20
297	ERISTLLKWL	18
333	PVSARVIKAL	18
154	EGFDLPPVIL	17
322	EEPDSGSHAG	17
330	AGGPVSARVI	17
409	HNIPHDFFSF	17
795	ENC PGWLDVL	17
20	KYKIACIVLL	16
57	KCFDASFRGL	16
178	DTLMPNINKL	16
293	VPFEERISTL	16
421	EEIVRNLSR	16
439	YLTPDLPKRL	16
494	AIFLAHGPF	16
855	VQPVSEILQL	16
22	KIACIVLLAL	15
30	ALLVIMSLGL	15
39	LGLGLGLRKL	15
155	GFDLPPVILF	15
277	AINGSFPSIY	15
280	GSFPSIYMPY	15
312	ERPRFYTM	15
340	KALQVVDHAF	15
361	LHNCVNILL	15
388	DYFPRINFFY	15
399	YEGPAPRIRA	15
430	RKPDQHFHPY	15
553	AEEVSKFSVC	15
607	NLPFGRPRVL	15
656	PLPPTVPDCL	15
697	ASNRTSDSQY	15
760	DAPDEITKHL	15
763	DEITKHLANT	15
772	TDVPIPTHYF	15
775	PIPTHYFVVL	15
817	SCPEGKPEAL	15
829	EERFTAHIAR	15
1	MESTLTATE	14
12	PVKKNTLKKY	14
28	LLALLVIMSL	14
32	LVIMSLGLGL	14
74	ACKDRGDCCW	14
99	NKFRCGETRL	14
141	EENCDAQQS	14

Table XLIV-V1-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
147	AQQSQCEPGEF	14
160	PVILFSMDGF	14
165	SMDGFRAEYL	14
197	MRAMYPTKTF	14
208	NHYTIVTGLY	14
218	PESHGIIDNN	14
248	AWWHGQPMWL	14
269	YFWPGSEVAI	14
295	FEERISTLLK	14
353	MEGLKQRNLH	14
387	TDYFPRINFF	14
435	HFKPYLTPDL	14
441	TPDLPKRLHY	14
453	NVRIDKVHLF	14
488	EFRSMEAIFL	14
516	EVYNLMCDLL	14
530	APNNGTHGSL	14
555	EVSKEFSVCGF	14
564	FANPLPTESL	14
571	ESLDCFCPHL	14
684	ADKNITHGFL	14
700	RTSDSQYDAL	14
744	NVVS GPIFDY	14
791	SHTPENCPGW	14
858	VSEILQLKTY	14
19	KKYKIACIVL	13
34	IMSLGLGLGL	13
36	SLGLGLGLGL	13
50	KQGSCRKKCF	13
66	LENCRCDVAC	13
85	FEDTCVETR	13
92	STRIWMCNKF	13
104	GETRLEASLC	13
140	LEENCDAQQ	13
164	FSMDGFRAEY	13
228	MYDVNLNKNF	13
240	SSKEQNNPAW	13
242	KEQNNPAWWH	13
260	MYQGLKAATY	13
276	VAINGSFPSI	13
321	FEEDSSGHA	13
347	HAFGMLMEGL	13
360	NLHNCVNIL	13
381	NKMEYMTDYF	13
384	EYMTDYFPRI	13
386	MTDYFPRINF	13
418	FNSEEIVRNL	13
452	KNVRIDKVHL	13
458	KVHLFVDQQW	13
533	NGTHGSLNHL	13
534	GTHGSLNHLL	13
554	EEVSKFSVCG	13
567	PLPTESLDCF	13
570	TESLDCFCPH	13
597	QEEITATVKV	13
599	EITATVKVNL	13

Table XLIV-V1-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
637	AMRMPMWSSY	13
642	MWSSYTVPQL	13
683	LADKNITHGF	13
716	PMYEEFRKMW	13
723	KMWDYFHSVL	13
724	MWDYFHSVLL	13
738	TERNGVNVVS	13
746	VSGPIFDYNY	13
771	NTDVPIPTHY	13
815	VESCPEGKPE	13
828	VEERFTAHIA	13
862	LQLKTYLPTF	13
866	TYLPTFETTI	13
23	IACIVLLALL	12
25	CIVLLALLVI	12
48	LEKQGSCRKK	12
76	KDRGDCCWDF	12
87	DTCVESTRIW	12
90	VESTRIWMCN	12
131	SVCQGETSWL	12
168	GFRAEYLYTW	12
192	IHSKYMRAMY	12
201	YPTKTFPNHY	12
220	SHGIIDNNMY	12
224	IDNNMYDVNL	12
241	SKEQNNPAWW	12
252	GQPMWLTAMY	12
255	MWLTAMYQGL	12
261	YQGLKAATYF	12
262	QGLKAATYFW	12
286	YMPYNGSVPF	12
299	ISTLLKWLDL	12
307	DLPKAERPRF	12
308	LPKAERPRFY	12
343	QVVDHAFGML	12
352	LMEGLKQRNL	12
370	LADHGMDQTY	12
402	PAPRIRAHNI	12
406	IRAHNIPHDF	12
420	SEEIVRNLSC	12
431	KPDQHFKPYL	12
476	TNCGGGNHGY	12
492	MEAIFLAHGP	12
504	KEKTEVEPFE	12
511	PFENIEVYNL	12
512	FENIEVYNLM	12
538	SLNHLLKVPF	12
539	LNHLLKVPFY	12
548	YEPShAEEVS	12
550	PSHAEVSKF	12
559	FSVCGFANPL	12
577	CPHLQNSTQL	12
616	LQKNVDHCLL	12
617	QKNVDHCLLY	12
621	DHCLLYHREY	12
625	LYHREYVSGF	12

Table XLIV-V1-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
628	REYVSGFGKA	12
648	VPQLGDT SPL	12
679	CSFY LADKNI	12
705	QYDALITSNL	12
712	SNLVPMYEEF	12
792	HTPENCPGWL	12
797	CPGWLDVLPF	12
818	CPEGKPEALW	12
819	PEGKPEALWV	12
835	HIARVRDVEL	12
836	IARVRDVELL	12
839	VRDVELLTGL	12
841	DVELLTGLDF	12

Table XLIV-V2-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
8	LQRKDCCADY	10
4	SDDCLQRKDC	5
2	SCSDDCLQRK	4
3	CSDDCLQRKD	4

Table XLIV-V3-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
7	SCPGGKPEAL	16
5	VESCPGGKPE	13
8	CPGGKPEALW	12

Table XLIV-V4-HLA-B4402-10mers-161P2F10B		
Pos	1234567890	score
2	TYLPTFETPI	12

Table XLV-V1-HLA-B5101-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLV-V2-HLA-B5101-10mers-161P2F10B		
Pos	1234567890	score
No Results Found.		

Table XLV-V3-HLA-B5101-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLV-V4-HLA-B5101-10mers-161P2F10B		
Pos	123456789	score
No Results Found.		

Table XLVI-V1-DRB1-0101-15mers- 161P2F10B		
Pos	123456789012345	score
207	PNHYTIVTGLYPESH	36
181	MPNINKLKTCGIHSK	35
486	NNEFRSMEAIFLAHG	35
839	VRDVELLTGLDFYQD	33
703	DSQYDALITSNLVPM	32
42	GLGLRKLEKQGSCRK	31
740	RNGVNVVSGPIFDYN	31
28	LLALLVIMSLGLGLG	30
160	PVILFSMDGFRAEYL	30
797	CPGWLDVLPFIIPHR	30
858	VSEILQLKTYLPTFE	30
801	LDVLPFIIPHRPTNV	29
393	INFFYMYEGPAPRIR	28
421	EEIVRNLSCKRPDQH	28
627	HREYVSGFGKAMRMP	28
30	ALLVIMSLGLGLGLG	27
32	LVIMSLGLGLGLGLR	27
34	IMSLGLGLGLGLRKL	27
364	CVNIILLADHGMDQT	27
522	CDLLRIQPAPNNGTH	27
23	IACIVLLALLVIMSL	26
258	TAMYQGLKAATYFWP	26
273	GSEVAINGSFPSIYM	26
283	PSIYMPYNGSVPFEE	26
300	STLLKWLDLPAERP	26
545	VPFYEP SHAEEVSKF	26
645	SYTVPQLGDTSPPLP	26
778	THYFVVLTSCKNKSH	26
26	IVLLALLVIMSLGLG	25
230	DVNLNKNFSLSSKEQ	25
266	AATYFWPGSEVAING	25
292	SVPFEERISTLLKWL	25
303	LKWLDLPAERPRFY	25
350	GMLMEGLKQRNLHNC	25
433	DQHFKPYLTPDLPKR	25
459	VHLFVDQQWLAVRSK	25
492	MEAIFLAHGSPFKEK	25
509	VEPFENIEVYNLMCD	25
536	HGSLNHLLKVPFYEP	25
561	VCGFANPLPTESLDC	25
22	KIACIVLLALLVIMS	24
25	CIVLLALLVIMSLGL	24
163	LFSMDGFRAEYLYTW	24
174	LYTWDTLMPNINKLK	24
194	SKYMRAMYPTKTFPN	24
246	NPAWWHGQPMWLTAM	24
313	RPRFYTMFEEDSS	24
316	FYTMFEEDSSGHA	24
333	PVSARVIKALQVVDH	24
342	LQVVDHAFGMLMEGL	24
387	TDYFPRINFFYMYEG	24
392	RINFFYMYEGPAPRI	24
395	FFYMYEGPAPRIRAH	24
412	PHDFFSFNSEEIVRN	24
437	KPYLTPDLPKRLHYA	24
514	NIEVYNLMCDLLRIQ	24

Table XLVI-V1-DRB1-0101-15mers- 161P2F10B		
Pos	123456789012345	score
518	YNLMCDLLRIQPAPN	24
539	LNHLLKVPFYEPSHA	24
542	LLKVPFYEPSHAEEV	24
589	VNQMLNLTQEEITAT	24
605	KVNLPGFRPRVLQKN	24
722	RKMWDYFHSVLLIKH	24
798	PGWLDVLPFIIPHRP	24
45	LRKLEKQGSCRKKCF	23
84	DFEDTCVESTRIWMC	23
129	YKSVCQGETSWLEEN	23
191	GIHSKYMRAMYPTKT	23
210	YTIVTGLYPESHGII	23
272	PGSEVAINGSFPSIY	23
284	SIYMPYNGSVPFEEER	23
328	GHAGGPVSARVIKAL	23
464	DQQWLAVRSKSNTNC	23
562	CGFANPLPTESLDCF	23
637	AMRMPMWSSYTPQL	23
644	SSYTPQLGDTSPPLP	23
651	LGDTSPPLPTVPDCL	23
687	NITHGFLYPASNRT	23
804	LPIIPHRPTNVESC	23
1	MESTLTLATEQPVKK	22
18	LKKYKIACIVLLALL	22
20	KYKIACIVLLALLVI	22
145	DTAQSQSQCEGFDLP	22
152	CPEGFDLPPVILFSM	22
155	GFDLPPVILFSMDGF	22
171	AEYLYTWDTLMPNIN	22
380	CNKMEYMTDYFPRIN	22
400	EGPAPRIRAHNIPHD	22
434	QHFKPYLTPDLPKRL	22
575	CFCPHLQNSTQLEQV	22
648	VPQLGDTSPPLPTVP	22
654	TSPLPPTVPDCLRAD	22
777	PTHYFVVLTSCKNKS	22
825	ALWVEERFTAHIARV	22
833	TAHIARVRDVELLTG	22
60	DASFRGLENCRCDDVA	21
142	ENCDTAQSQSQCEGF	21
186	KLKTCGIHSKYMRAM	21
451	AKNVRIDKVHLFVDQ	21
490	RSMEAIFLAHGSPFK	21
665	LRADVRVPPSESQKC	21
704	SQYDALITSNLVPMY	21
739	ERNGVNVVSGPIFDY	21
770	ANTDVPIPTHYFVVL	21
98	CNKFRCGETRLEASL	20
253	QPMWLTAMYQGLKAA	20
443	DLPKRLHYAKNVRID	20
521	MCDLLRIQPAPNNGT	20
553	AAEVSKFSVCGFANP	20
623	CLLYHREYVSGFGKA	20
631	VSGFGKAMRMPMWSS	20
718	YEEFRKMWDYFHSVL	20
729	HSVLLIKHATERNGV	20
750	IFDYNVDGHFDAPDE	20

Table XLVI-V1-DRB1-0101-15mers- 161P2F10B		
Pos	123456789012345	score
829	EERFTAHARVRDVE	20
15	KNTLK KYKIACIVLL	19
37	LGLGLGLRLKLEKQ	19
153	PEGFDLPPVILFSMD	19
158	LPPVILFSMDGFRAE	19
170	RAEYLYTWDTLMPNI	19
226	NNMYDVNLNKNFSL	19
305	WLDLPKAERPRFYT	19
340	KALQVVDHAFGMLME	19
346	DHAFGMLMEGLKQRN	19
363	NCVNIILLADHGMDQ	19
413	HDFFSFNSEEIVRNL	19
500	GPSFKEKTEVEPFEN	19
556	VSKFSVCGFANPLPT	19
603	TVKVNLPFGRPRVLQ	19
691	GFLYPASNRTSDSQ	19
762	PDEITKHLANTDVPI	19
771	NTDVPIPTHYFVLT	19
817	SCPEGKPEALWVEER	19
2	ESTLTATEQPVKKN	18
12	PVKKNTLK KYKIACI	18
29	LALLVIMSLGLGLGL	18
102	RCGETRLEASLCSCS	18
136	ETSWLEENCDAQQS	18
157	DLPPVILFSMDGFRA	18
183	NINKLKTCGIHSKYM	18
214	TGLYPESHGIIDNNM	18
228	MYDVNLNKNFSLSSK	18
255	MWLTAMYQGLKAATY	18
259	AMYQGLKAATYFWPG	18
267	ATYFWPGSEVAINGS	18
297	ERISTLLKWLDLPKA	18
302	LLKWLDLPKAERPRF	18
318	TMYFEEDSSGHAGG	18
322	EEPSSGHAGGPVSA	18
362	HNCVNIILLADHGMD	18
382	KMEYMTDYFPRINFF	18
515	IEVYNLMCDLLRIQP	18
595	LTQEEITATVKVNL	18
613	PRVLQKNVDHCLLYH	18
640	MPMWSSYTPQLGDT	18
662	PDCLRADVRVPPSES	18
681	FYLADKNITHGFLYP	18
689	THGFLYPASNRTSD	18
748	GPIFDYNYDGHFDAP	18
763	DEITKHLANTDVPI	18
847	GLDFYQDKVQPVSEI	18
848	LDFYQDKVQPVSEIL	18
16	NLTKKYKIACIVLLA	17
21	YKIACIVLLALLVIM	17
24	ACIVLLALLVIMSLG	17
31	LLVIMSLGLGLGLGL	17
36	SLGLGLGLRLKLEK	17
80	DCCWDFEDTCVEST	17
103	CGETRLEASLCSCSD	17
126	CADYKSVCQGETSWL	17
139	WLEENCDAQQSQCP	17

Table XLVI-V1-DRB1-0101-15mers- 161P2F10B		
Pos	123456789012345	score
147	AQQSQCEGFDLPPV	17
189	TCGIHSKYMRAMYPT	17
209	HYTIVTGLYPESHGI	17
212	IVTGLYPESHGIIDN	17
227	NMYDVNLNKNFSLSS	17
236	NFSLSSKEQNNPAWW	17
261	YQGLKAATYFWPGSE	17
265	KAATYFWPGSEVAIN	17
282	FPSIYMPYNGSVPF	17
286	YMPYNGSVPFEEIS	17
324	PDSSGHAGGPVSARV	17
331	GGPVSARVIKALQV	17
336	ARVIKALQVVDHAFG	17
341	ALQVVDHAFGMLMEG	17
347	HAFGMLMEGLKQRNL	17
372	DHGMDQTYCNKMEYM	17
394	NFFYMYEGPAPRIRA	17
396	FYMYEGPAPRIRAHN	17
491	SMEAIFLAHGSPFKE	17
493	EAIFLAHGSPFKEKT	17
506	KTEVEPFENIEVYNL	17
519	NLMCDLLRIQPAPNN	17
527	IQPAPNNGTHGSLNH	17
540	NHLLKVPFYEPSHAE	17
557	SKFSVCGFANPLPTE	17
597	QEEITATVKVNLPG	17
601	TATVKVNLPGRPRV	17
602	ATVKVNLPGRPRVL	17
629	EYVSGFGKAMRMPMW	17
677	QKCSFYLADKNITHG	17
685	DKNITHGFLYPASN	17
725	WDYFHSVLLIKHATE	17
726	DYFHSVLLIKHATER	17
734	IKHATERNGVNVVSG	17
780	YFVLTSCKNKSHTP	17
812	PTNVESCEGKPEAL	17
823	PEALWVEERFTAHIA	17
836	IARVRDVELLTGLDF	17
842	VELLTGLDFYQDKVQ	17

Table XLVI-V2-DRB1-0101-15mers- 161P2F10B		
Pos	123456789012345	score
2	EASLCSCSDDCLQRK	16
7	SCSDDCLQRKDCCAD	9
12	CLQRKDCCADYKSVC	9
13	LQRKDCCADYKSVCQ	9
14	QRKDCCADYKSVCQG	9
9	SDDCLQRKDCCADYK	8
10	DDCLQRKDCCADYKS	8
8	CSDDCLQRKDCCADY	7

Table XLVI-V3-HLA-DRB1-0101-15mers- 161P2F10B		
Pos	123456789012345	score
7	PTNVESCPGGKPEAL	25
12	SCPGGKPEALWVEER	19

Table XLVI-V3-HLA-DRB1-0101-15mers-161P2F10B		
Pos	123456789012345	score
4	PHRPTNVESCPGGKP	14
10	VESCPGGKPEALWVE	14

Table XLVI-V4-HLA-DRB1-0101-15mers-161P2F10B		
Pos	123456789012345	score
2	ILQLKTYLPTFETPI	16
1	EILQLKTYLPTFETP	10

Table XLVII-V1-HLA-DRB1-0301-15mers-161P2F10B		
Pos	123456789012345	score
437	KPYLTPDLPKRLHYA	31
451	AKNVRIDKVHLFVDQ	31
662	PDCLRADVRVPPSES	28
458	KVHLFVDQQWLAVRS	27
597	QEEITATVKVNLPGF	27
228	MYDVNLNKNFSLSSK	26
350	GMLMEGLKQRNLHNC	26
536	HGSLNHLLKVPFYEP	26
305	WLDLPKAERPRFYTM	25
358	QRNLHNCVNIILLAD	24
30	ALLVIMSLGLGLGLG	22
366	NIILLADHGMDQTYC	22
28	LLALLVIMSLGLGLG	21
32	LVIMSLGLGLGLRLR	21
34	IMSLGLGLGLRLKL	21
158	LPPVILFSMDGFRAE	21
161	VILFSMDGFRAEYLY	21
517	VYNLMCDLLRIQPAP	21
707	DALITSNLVPMYEEF	21
742	GVNVVSGPIFDYNYD	21
10	EQPVKKNTLKKYKIA	20
20	KYKIACIVLLALLVI	20
109	EASLCSCSDDCLQKK	20
118	DCLQKKDCCADYKSV	20
236	NFSLSSKEQNNPAWW	20
284	SIYMPYNGSVPFEER	20
408	AHNIPHDFFSFNSEE	20
467	WLAVRSKSNTNCGGG	20
615	VLQKNVDHCLLYHRE	20
635	GKAMRMPMWSSYTP	20
780	YFVLTSCKNKSHTP	20
839	VRDVELLTGLDFYQD	20
847	GLDFYQDKVQPVSEI	20
855	VQPVSEILQLKTYLP	20
2	ESTLTATEQPVKKN	19
26	IVLLALLVIMSLGLG	19
38	GLGLGLGLRKLEKQG	19
42	GLGLRKLEKQGSCRK	19
74	ACKDRGDCCWDFEDT	19
129	YKSVCQGETSWLEEN	19
153	PEGFDLPPVILFSMD	19
257	LTAMYQGLKAATYFW	19
331	GGPV SARVIKALQVV	19
335	SARVIKALQVDHAF	19
341	ALQVVDHAFGMLMEG	19

Table XLVII-V1-HLA-DRB1-0301-15mers-161P2F10B		
Pos	123456789012345	score
349	FGMLMEGLKQRNLHN	19
433	DQHFKPYLTPDLPKR	19
486	NNEFRSMEAIFLAHG	19
492	MEAIFLAHGSPFKEK	19
514	NIEVYNLMCDLLRIQ	19
524	LLRIQPAPNNGTHGS	19
553	AEVSKFSVCGFANP	19
565	ANPLPTESLDCFCPH	19
654	TSPLPPTVPDCLRAD	19
679	CSFYLADKNITHGFL	19
697	ASNRTSDSQYDALIT	19
740	RNGVNVVSGPIFDYN	19
773	DVPIPTHYFVLTSC	19
781	FVLTSCKNKSHTPE	19
812	PTNVESCEPGKPEAL	19
833	TAHIARVRDVELLTG	19
835	HIARVRDVELLTGLD	19
852	QDKVQPVSEILQLKT	19
163	LFSMDGFRAEYLYTW	18
177	WDTLMPNINKLKTCG	18
178	DTLMPNINKLKTCGI	18
224	IDNNMYDVNLNKNFS	18
254	PMWLTAMYQGLKAAT	18
267	ATYFWPGSEVAINGS	18
275	EVAINGSFPSIYMPY	18
292	SVPFEERISTLLKWL	18
296	EERISTLLKWLDLPK	18
445	PKRLHYAKNVRIDKV	18
465	QQWLAVRSKSNTNCG	18
506	KTEVEPFENIEVYNL	18
509	VEPFENIEVYNLMCD	18
582	NSTQLEQVNQMLNLT	18
586	LEQVNQMLNLTQEEI	18
589	VNQMLNLTQEEITAT	18
605	KVNLPFGRPRVLQKN	18
613	PRVLQKNVDHCLLYH	18
621	DHCLLYHREYVSGFG	18
628	REYVSGFGKAMRMPM	18
658	PPTVPDCLRADVRVP	18
711	TSNLVPMYEEFRKMW	18
712	SNLVPMYEEFRKMWD	18
721	FRKMWDYFHSVLLIK	18
731	VLLIKHATERNGVNV	18
7	LATEQPVKKNTLKKY	17
92	STRIWMCNKFCRCGET	17
219	ESHGIIDNNMYDVNL	17
282	FPSIYMPYNGSVPFE	17
290	NGSVPFEERISTLLK	17
439	YLTPDLPKRLHYAKN	17
457	DKVHLFVDQQWLAVR	17
500	GPSFKEKTEVEPFEN	17
518	YNLMCDLLRIQPAPN	17
558	KFSVCGFANPLPTES	17
573	LDCFCPHLQNSTQLE	17
577	CPHLQNSTQLEQVNO	17
583	STQLEQVNQMLNLTQ	17
590	NQMLNLTQEEITATV	17

Table XLVII-V1-HLA-DRB1-0301-15mers-161P2F10B		
Pos	123456789012345	score
714	LVPMYEEFRKMWDYF	17
746	VSGPIFDYNYDGHFD	17
823	PEALWVEERFTAHIA	17
857	PVSEILQLKTYLPTF	17
60	DASFRGLENCRCDDVA	16
70	RCDVACKDRGDCCWD	16
78	RGDCCWDFEDTCVES	16
94	RIWMCNKFRGGETRL	16
101	FRCGETRLEASLCSC	16
122	KKDCCADYKSVCQGE	16
226	NNMYDVNLNKNFSL	16
234	NKNFSLSSKEQNNPA	16
346	DHAFGMLMEGLKQRN	16
384	EYMTDYFPRINFFYM	16
450	YAKNVRIDKVHLFVD	16
532	NNGTHGSLNHLLKVP	16
599	EITATVKVNLFPGRP	16
678	KCSFYLDKNITHGF	16
748	GPIFDYNYDGHFDAP	16
24	ACIVLLALLVIMSLG	15
48	LEKQGSCRKKCFDAS	15
56	KKCFDASFRGLENCR	15
97	MCNKFRGGETRLEAS	15
162	ILFSMDGFRAEYLYT	15
164	FSMDGFRAEYLYTWD	15
190	CGHSKYMRAMYPTK	15
218	PESHGIIDNNMYDVN	15
374	GMDQTYCNKMEYMTD	15
386	MTDYFPRINFFYMYE	15
413	HDFFSFNSEEIVRN	15
474	SNTNCGGGNHGYNNE	15
478	CGGGNHGYNNEFRSM	15
485	YNNEFRSMEAIFLAH	15
670	RVPPSESQKCSFYLA	15
728	FHSVLLIKHATERNG	15
803	VLPFIIPHRPTNVES	15
821	GKPEALWVEERFTAH	15

Table XLVII-V2-DRB1-0301-15mers-161P2F10B		
Pos	123456789012345	score
2	EASLCSCSDDCLQRK	20
11	DCLQRKDCCADYKSV	20
15	RKDCCADYKSVCQGE	16
4	SLCSCSDDCLQRKDC	12
5	LCSCSDDCLQRKDCC	11
10	DDCLQRKDCCADYKS	11

Table XLVII-V3-HLA-DRB1-0301-15mers-161P2F10B		
Pos	123456789012345	score
7	PTNVECPGGKPEAL	13
10	VECPGGKPEALWVE	10
11	ESCPGGKPEALWVEE	9
9	NVECPGGKPEALWV	8
1	FIIPHRPTNVECPG	7
3	IPHRPTNVECPGGK	6

Table XLVII-V4-HLA-DRB1-0301-15mers-161P2F10B		
Pos	123456789012345	score
2	ILQLKTYLPTFETPI	12
1	EILQLKTYLPTFETP	10

Table XLVIII-V1-HLA-DRB1-0401-15mers-161P2F10B		
Pos	123456789012345	score
126	CADYKSVCQGETSWL	28
412	PHDFFSFNSEEIVRN	28
464	DQQWLAVRSKSNINC	28
631	VSGFGKAMRMPMWSS	28
691	GFLYPASNRTSDSQ	28
703	DSQYDALITSNLVPM	28
722	RKMWDYFHSVLLIKH	28
750	IFDYNIDGHFDAPDE	28
777	PTHYFVLTSCKNKS	28
778	THYFVLTSCKNKSH	28
28	LLALLVIMSLGLGLG	26
181	MPNINKLKTGCIHSK	26
213	VTGLYPESHGIIDNN	26
290	NGSVPFEERISTLLK	26
350	GMLMEGLKQRNLHNC	26
445	PKRLHYAKNVRIDKV	26
458	KVHLFVDQQWLAVRS	26
506	KTEVEPFENIEVYNL	26
524	LLRIQPAPNNGTHGS	26
583	STQLEQVNQMLNLTQ	26
589	VNQMLNLTQEEITAT	26
823	PEALWVEERFTAHIA	26
855	VQPVSEILQLKTYLP	26
18	LKKYKIACIVLLALL	22
60	DASFRGLENCRCDDVA	22
80	DCCWDFEDTCVESTR	22
136	ETSWLEENCDDTAQQS	22
172	EYLYTWDTLMPNINK	22
174	LYTWDTLMPNINKLK	22
193	HSKYMRAMYPTKTFP	22
203	TKTFPNHYTIVTGLY	22
253	QPMWLTAMYQGLKAA	22
266	AATYFWPGSEVAING	22
279	NGSFPSIYMPYNGSV	22
346	DHAFGMLMEGLKQRN	22
382	KMEYMTDYFPRINFF	22
387	TDYFPRINFFYMYEG	22
415	FFSFNSEEIVRNLS	22
433	DQHFKPYLTPDLPKR	22
447	RLHYAKNVRIDKVHL	22
482	NHGYNNEFRSMEAIF	22
509	VEPFENIEVYNLMCD	22
573	LDCFCPHLQNSTQLE	22
678	KCSFYLDKNITHGF	22
679	CSFYLDKNITHGFL	22
797	CPGWLDVLPFIIPHR	22
803	VLPFIIPHRPTNVES	22
847	GLDFYQDKVQPVSEI	22
2	ESTLTLATEQPVKKN	20
4	TLTLATEQPVKKNL	20

Table XLVIII-V1-HLA-DRB1-0401-15mers-161P2F10B		
Pos	123456789012345	score
20	KYKIACIVLLALLVI	20
23	IACIVLLALLVIMSL	20
24	ACIVLLALLVIMSLG	20
25	CIVLLALLVIMSLGL	20
26	IVLLALLVIMSLGLG	20
42	GLGLRKLEKQGSCRK	20
45	LRKLEKQGSCRKKCF	20
129	YKSVCQGETSWLEEN	20
155	GFDLPPVILFMDGF	20
163	LFSMDGFRAEYLYTW	20
177	WDTLMPNINKLKTCG	20
178	DTLMPNINKLKTCGI	20
197	MRAMYPTKTFPNHYT	20
225	DNNMYDVNLNKNFSL	20
228	MYDVNLNKNFSLSSK	20
230	DVNLNKNFSLSSKEQ	20
254	PMWLTAMYQGLKAAT	20
273	GSEVAINGSFPSIYM	20
282	FPSIYMPYNGSVPF	20
300	STLLKWLDLPKAERP	20
305	WDLPLKAERPFRYTM	20
316	FYTMFEEDSSGHA	20
335	SARVIKALQVVDHAF	20
339	IKALQVVDHAFGMLM	20
342	LQVVDHAFGMLMEGL	20
358	QRNLHNCVNIILLAD	20
364	CVNIILLADHGMDQT	20
365	VNIILLADHGMDQTY	20
366	NIILLADHGMDQTYC	20
408	AHNIPHDFSFNSEE	20
420	SEEIVRNLSRKPQDQ	20
437	KPYLTPDLPKRLHYA	20
514	NIEVYNLMCDLLRIQ	20
517	VYNLMCDLLRIQPAP	20
518	YNLMCDLLRIQPAPN	20
521	MCDLLRIQPAPNNGT	20
558	KFSVCGFANPLPTES	20
570	TESLDCFCPHLQNST	20
577	CPHLQNSTQLEQVNQ	20
586	LEQVNQMLNLTQEEI	20
592	MLNLTQEEITATVKV	20
637	AMRMPMWSSYTPQL	20
639	RMPMWSSYTPQLGD	20
658	PPTVPDCLRADVRVP	20
662	PDCLRADVRVPPSES	20
666	RADVRVPPSESQKCS	20
690	HGFLYPASNRTSDS	20
714	LVPMYEEFRKMWDYF	20
721	FRKMWDYFHSVLLIK	20
728	FHSVLLIKHATERNG	20
730	SVLLIKHATERNGVN	20
771	NTDVPIPTHYFVLT	20
833	TAHIARVRDVELLTG	20
852	QDKVQPVSEILQLKT	20
858	VSEILQLKTYLPTFE	20
84	DFEDTCVESTRIWMC	18
99	NKFRCGETRLEASLC	18

Table XLVIII-V1-HLA-DRB1-0401-15mers-161P2F10B		
Pos	123456789012345	score
103	CGETRLEASLCSCSD	18
135	GETSWLEENCDTAQQ	18
168	GFRAEYLYTWDTLMP	18
186	KLKTCGIHISKYMRAM	18
200	MYPTKTFPNHYTIVT	18
224	IDNNMYDVNLNKNFS	18
233	LNKNFSLSSKEQNNP	18
237	FSLSSKEQNNPAWWH	18
276	VAINGSFPSIYMPYN	18
293	VPFEERISTLLKWLD	18
319	MYFEEDSSGHAGGP	18
355	GLKQRNLHNCVNIIL	18
370	LADHGMDQTYCNKME	18
397	YMYEGPAPRIAHNI	18
409	HNIPHDFSFNSEEI	18
417	SFNSEEIVRNLSRCK	18
457	DKVHLFVDQQWLAVR	18
463	VDQQWLAVRSKSNNTN	18
483	HGYNNEFRSMEAIFL	18
491	SMEAIFLAHGSPFKE	18
533	NGTHGSLNHLLKVPF	18
549	EPShAEEVSKFSVCG	18
562	CGFANPLPTESLDCF	18
574	DCFCPHLQNSTQLEQ	18
580	LQNSTQLEQVNQMLN	18
593	LNLTQEEITATVKVN	18
604	VKNLFPGRPRVLQK	18
609	PFGRPRVLQKNVDHC	18
614	RVLQKNVDHCLLYHR	18
669	VRVPPSESQKCSFYL	18
682	YLADKNITHGFLYPP	18
697	ASNRTSDSQYDALIT	18
704	SQYDALITSNLVPMY	18
733	LIKHATERNGVNVVS	18
739	ERNGVNVVSGPIFDY	18
758	HFDAPDEITKHLANT	18
763	DEITKHLANTDVPIP	18
802	DVLPFIIPHRPTNVE	18
844	LLTGLDFYQDKVQPV	18
292	SVPFEEERISTLLKW	17
848	LDYQDKVQPVSEIL	17
56	KKCFDASFRGLENCR	16
82	CWDFEDTCVESTRIW	16
98	CNKFRCGETRLEASL	16
161	VILFMDGFRAEYLY	16
166	MDGFRAEYLYTWDTL	16
170	RAEYLYTWDTLMPNI	16
207	PNHYTIVTGLYPESH	16
226	NNMYDVNLNKNFSLS	16
234	NKNFSLSSKEQNNPA	16
246	NPAWWHGQPMWLTAM	16
247	PAWWHGQPMWLTAMY	16
258	TAMYQGLKAATYFWP	16
267	ATYFWPGSEVAINGS	16
268	TYFWPGSEVAINGSF	16
302	LLKWLDLPKAERPRF	16
317	YTMFEEDSSGHAG	16

Table XLVIII-V1-HLA-DRB1-0401-15mers-161P2F10B		
Pos	123456789012345	score
318	TMYFEEDSSGHAGG	16
392	RINFFYMYEGPAPRI	16
394	NFFYMYEGPAPRIRA	16
413	HDFFSFNSEEIVRNL	16
459	VHLFVDQQWLAVRSK	16
486	NNEFRSMEAIFLAHG	16
493	EAIFLAHGPSFKEKT	16
544	KVPFYEPSHAEEVSK	16
545	VPFYEPSHAEEVSKF	16
561	VCGFANPLPTESLDC	16
607	NLPFGRPRVLQKNVD	16
623	CLLYHREYVSGFGKA	16
627	HREYVSGFGKAMRMP	16
640	MPMWSSYTPQLGDT	16
643	WSSYTPQLGDTSP	16
715	VPMYEEFRKMWDYFH	16
724	MWDYFHSVLLIKHAT	16
725	WDYFHSVLLIKHATE	16
756	DGHFDAPDEITKHLA	16
824	EALWVEERFTAHIAR	16
441	TPDLPKRLHYAKNVR	15
825	ALWVEERFTAHIARV	15
15	KNTLKKYKIACIVLL	14
30	ALLVIMSLGLGLGLG	14
31	LLVIMSLGLGLGLGL	14
32	LVIMSLGLGLGLGLR	14
34	IMSLGLGLGLGLRKL	14
36	SLGLGLGLGLRKLEK	14
38	GLGLGLGLRKLEKQG	14
63	FRGLENCRCDVACKD	14
105	ETRLEASLCSCSDDC	14
109	EASLCSCSDDCLQKK	14
137	TSWLEENCDTAQQSQ	14
158	LPPVILFSMDGFRAE	14
160	PVILFSMDGFRAEYL	14
171	AEYLYTWDTLMPNIN	14
184	INKLKTCGIHSKYMR	14
194	SKYMRAMYPTKTFPN	14
209	HYTIVTGLYPESHGI	14
210	YTIVTGLYPESHGII	14
220	SHGIIDNNMYDVNLN	14
221	HGIIDNNMYDVNLNK	14
257	LTAMYQGLKAATYFW	14
261	YQGLKAATYFWPGSE	14
284	SIYMPYNGSVPFEER	14
296	EERISTLLKWLDLPK	14
299	ISTLLKWLDLPKAER	14
336	ARVIKALQVVDHAFG	14
341	ALQVVDHAFGMLMEG	14
348	AFGMLMEGLKQRNLH	14
349	FGMLMEGLKQRNLHN	14
362	HNCVNIILLADHGMD	14
367	IILLADHGMDQTYCN	14
372	DHGMDQTYCNKMEYM	14
383	MEYMTDYFPRINFFY	14
390	FPRINFFYMYEGPAP	14
395	FFYMYEGPAPRIRAH	14

Table XLVIII-V1-HLA-DRB1-0401-15mers-161P2F10B		
Pos	123456789012345	score
421	EEIVRNLSCKRPDQH	14
451	AKNVRIDKVHLFVDQ	14
453	NVRIDKVHLFVDQQW	14
456	IDKVHLFVDQQWLAV	14
489	FRSMEAIFLAHGSPF	14
492	MEAIFLAHGSPFKEK	14
494	AIFLAHGSPFKEKTE	14
522	CDLLRIQPAPNNGTH	14
536	HGSLNHLLKVPFYEP	14
539	LNHLLKVPFYEPSHA	14
540	NHLLKVPFYEPSHAE	14
542	LLKVPFYEPSHAEEV	14
553	AAEVSKFSVCGFANP	14
565	ANPLPTESLDCFCPH	14
590	NQMLNLTQEEITATV	14
597	QEEITATVKVNLPGF	14
605	KVNLPGFRPRVLQKN	14
613	PRVLQKNVDHCLLYH	14
621	DHCLLYHREYVSGFG	14
628	REYVSGFGKAMRMPM	14
645	SYTPQLGDTSP	14
654	TSPLPTVPDCLRAD	14
668	DVRVPPSESQKCSFY	14
706	YDALITSNLVPMYEE	14
712	SNLVPMYEEFRKMWD	14
731	VLLIKHATERNGVNV	14
740	RNGVNVVSGPIFDYN	14
743	VNVVSGPIFDYNYDG	14
773	DVPIPTHYFVLTSC	14
779	HYFVLTSCKNKSHT	14
780	YFVLTSCKNKSHTP	14
781	FVLTSCKNKSHTPE	14
798	PGWLDVLPFIIPHRP	14
800	WLDVLPFIIPHRPTN	14
801	LDVLPFIIPHRPTNV	14
804	LPFIIPHRPTNVECS	14
836	IARVRDVELLTGLDF	14
839	VRDVELLTGLDFYQD	14
841	DVELLTGLDFYQDKV	14
842	VELLTGLDFYQDKVQ	14
845	LTGLDFYQDKVQPV	14

Table XLVIII-V2-DR1-0401-15mers-161P2F10B		
Pos	123456789012345	score
2	EASLCSCSDDCLQRK	14
5	LCSCSDDCLQRKDCC	12
7	SCSDDCLQRKDCCAD	12
14	QRKDCCADYKSVCCG	12
1	LEASLCSCSDDCLQR	6
3	ASLCSCSDDCLQRKD	6
4	SLCSCSDDCLQRKDC	6
6	CSCSDDCLQRKDCCA	6
11	DCLQRKDCCADYKSV	6
12	CLQRKDCCADYKSV	6
13	LQRKDCCADYKSVCC	6
15	RKDCCADYKSVCCGE	6

Table XLIX-V1-HLA-DRB1-1101-15mers- 161P2F10B		
Pos	123456789012345	score
339	IKALQVVDHAFGMLM	27
42	GLGLRKLEKQGSCRK	26
518	YNLMCDLLRIQPAPN	26
207	PNHYTIVTGLYPESH	24
302	LLKWLDLPKAERPRF	24
750	IFDYNVDGHFDAPDE	24
392	RINFFYMYEGPAPRI	23
417	SFNSEEIVRNLSCKR	23
313	RPRFYTMFEEDSS	22
662	PDCLRADVRVPPSES	22
160	PVILFSMDGFRAEYL	21
178	DTLMPNINKLKTGCI	21
296	EERISTLLKWLDLPK	21
759	FDAPDEITKHLANTD	21
780	YFVLTSCNKSHTP	21
823	PEALWVEERFTAHIA	21
227	NMYDVNLKNFSLSS	20
421	EEIVRNLSCKPDQH	20
447	RLHYAKNVRIDKVLH	20
491	SMEAIFLAHGSPFKE	20
536	HGSLNHLLKVPFYEP	20
728	FHSVLLIKHATERNG	20
798	PGWLDVLPFIIPHRP	20
801	LDVLPFIIPHRPTNV	20
29	LALLVIMSLGLGLGL	19
31	LLVIMSLGLGLGLGL	19
56	KKCFDASFRGLENCR	19
300	STLLKWLDLPKAERP	19
678	KCSFYLDKNITHGF	19
848	LDFYQDKVQPVSEIL	19
858	VSEILQLKTYLPTFE	19
25	CIVLLALLVIMSLGL	18
60	DASFRGLENCRCDDA	18
234	NKNFSLSSKEQNNPA	18
266	AATYFWPGSEVAING	18
380	CNKMEYMTDYFPRIN	18
482	NHGYNNEFRSMEAIF	18
489	FRSMEAIFLAHGSPF	18
539	LNHLLKVPFYEPSHA	18
544	KVPFYEPSHAEEVSK	18
597	QEEITATVKVNLPGF	18
631	VSGFGKAMRMPMWSS	18
645	SYTVPQLGDTSPPLP	18
715	VPMYEEFRKMWDYFH	18
777	PTHYFVLTSCNKS	18
778	THYFVLTSCNKS	18
803	VLPFIIPHRPTNVS	18
836	IARVRDVELLTGLDF	18
68	NCRCDVACKDRGDCC	17
93	TRIWMCNKFRGETR	17
174	LYTWDTLMPNINKLK	17
283	PSIYMPYNGSVPFEE	17
317	YTMFEEDSSGHAG	17
346	DHAFGMLMEGLKQRN	17
463	VDQQWLAVRSKNTN	17
515	IEVYNLMCDLLRIQP	17

Table XLIX-V1-HLA-DRB1-1101-15mers- 161P2F10B		
Pos	123456789012345	score
556	VSKFSVCGFANPLPT	17
691	GFLYPASNRTSDSQ	17
725	WDYFHSVLLIKHATE	17
797	CPGWLDVLPFIIPHR	17
92	STRIWMCNKFRGET	16
126	CADYKSVCCGETSWL	16
136	ETSWLENCDDAQQS	16
258	TAMYQGLKAATYFWP	16
279	NGSFPSIYMPYNGSV	16
305	WLDLPKAERPRFYTM	16
349	FGMLMEGLKQRNLHN	16
387	TDYFPRINFFYMYEG	16
393	INFFYMYEGPAPRIR	16
397	YMYEGPAPRIRAHNI	16
438	PYLTPDLPKRLHYAK	16
464	DQQWLAVRSKNTNC	16
486	NNEFRSMEAIFLAHG	16
500	GPSFKEKTEVEPFEN	16
509	VEPFENIEVYNLMCD	16
561	VCGFANPLPTESLDC	16
610	FGRPRVLQKNVDHCL	16
703	DSQYDALITSNLVPM	16
718	YEEFRKMWDYFHSV	16
802	DVLPFIIPHRPTNVE	16
830	ERFTAHARVRDVEL	16
12	PVKKNTLKKYKIACI	15
38	GLGLGLGLRKLEKQG	15
213	VTGLYPESHGIIDNN	15
254	PMWLTAMYQGLKAAT	15
332	GPVSARVIKALQVVD	15
401	GPAPRIRAHNIPHDF	15
427	LSCRKPDQHFQPYLT	15
458	KVHLFVDQQWLAVRS	15
465	QQWLAVRSKNTNCG	15
533	NGTHGSLNHLLKVPF	15
614	RVLQKNVDHCLLYHR	15
619	NVDHCLLYHREYVSG	15
721	FRKMWDYFHSVLLIK	15
846	TGLDFYQDKVQPVSE	15
7	LATEQPVKKNTLKKY	14
20	KYKIACIVLLALLVI	14
26	IVLLALLVIMSLGLG	14
39	LGLGLGLRKLEKQGS	14
70	RCDVACKDRGDCCWD	14
94	RIWMCNKFRGETRL	14
99	NKFRGETRLEASLC	14
114	SCSDDCLQKKDCCAD	14
180	LMPNINKLKTGCIHS	14
196	YMRAMYPTKTFPNHY	14
210	YTIVTGLYPESHGII	14
243	EQNNPAWWHGQPMWL	14
257	LTAMYQGLKAATYFW	14
290	NGSVPFEEERISTLLK	14
303	LKWLDLPKAERPRFY	14
321	FEEDSSGHAGGPVS	14
365	VNIILLADHGMDQTY	14
405	RIRAHNIPHDFFSFN	14

Table XLIX-V1-HLA-DRB1-1101-15mers-161P2F10B		
Pos	123456789012345	score
441	TPDLPKRLHYAKNVR	14
450	YAKNVRIDKVHLFVD	14
550	PSHAEEVSKFSVCGF	14
602	ATVKVNLPFGRPRVL	14
628	REYVSGFGKAMRMPM	14
658	PPTVPDCLRADVRVP	14
682	YLADKNITHGFLYPP	14
685	DKNITHGFLYPPASN	14
714	LVPMYEEFRKMWDYF	14
724	MWDYFHSVLLIKHAT	14
727	YFHSVLLIKHATERN	14
740	RNGVNVVSGPIFDYN	14
771	NTDVPIPTHYFVLT	14
814	NVESCPEGKPEALWV	14
829	EERFTAHARVRDVE	14
842	VELLTGLDFYQDKVQ	14
23	IACIVLLALLVIMSL	13
27	VLLALLVIMSLGLGL	13
28	LLALLVIMSLGLGLG	13
33	VIMSLGLGLGLLRK	13
63	FRGLENCRCDVACKD	13
194	SKYMRAMYPTKTFPN	13
225	DNNMYDVNLNKNFSL	13
350	GMLMEGLKQRNLHNC	13
434	QHFKPYLTPDLPKRL	13
453	NVRIDKVHLFVDQQW	13
514	NIEVYNLMCDLLRIQ	13
540	NHLLKVPFYEPSHAE	13
558	KFSVCGFANPLPTES	13
583	STQLEQVNQMLNLTQ	13
598	EEITATVKVNLPFGR	13
621	DHCLLYHREYVSGFG	13
643	WSSYTPQLGDTSP	13
651	LGDTSPLPPTVPDCL	13
726	DYFHSVLLIKHATER	13
731	VLLIKHATERNGVNV	13
766	TKHLANTDVPIPTHY	13
812	PTNVESCEPGKPEAL	13
852	QDKVQPVSEILQLKT	13
855	VQPVSEILQLKTYLP	13

Table XLVIII-V3-HLA-DR1-0401-15mers-161P2F10B		
Pos	123456789012345	score
1	FIIPHRPTNVESCPG	12
4	PHRPTNVESCPGGKP	12
7	PTNVESCPGGKPEAL	8
5	HRPTNVESCPGGKPE	6
6	RPTNVESCPGGKPEA	6
9	NVESCPGGKPEALWV	6
10	VESCPGGKPEALWVE	6
12	SCPGGKPEALWVEER	6
13	CPGGKPEALWVEERF	6
14	PGGKPEALWVEERFT	6
15	GGKPEALWVEERFTA	6

Table XLVIII-V4-HLA-DR1-0401-15mers-161P2F10B		
Pos	123456789012345	score
2	ILQLKTYLPTFETPI	8

Table XLIX-V2-DRB1-1101-15mers-161P2F10B		
Pos	123456789012345	score
7	SCSDDCLQRKDCCAD	14
15	RKDCCADYKSVCCGE	9
8	CSDDCLQRKDCCADY	8
13	LQRKDCCADYKSVCCQ	7
2	EASLCSCSDDCLQRK	6
9	SDDCLQRKDCCADYK	6
10	DDCLQRKDCCADYKS	6

Table XLIX-V3-HLA-DRB1-1101-15mers-161P2F10B		
Pos	123456789012345	score
4	PHRPTNVESCPGGKP	14
9	NVESCPGGKPEALWV	14
7	PTNVESCPGGKPEAL	13
3	IPHRPTNVESCPGGK	8
5	HRPTNVESCPGGKPE	7
1	FIIPHRPTNVESCPG	6
15	GGKPEALWVEERFTA	6

Table XLIX-V4-HLA-DRB1-1101-15mers-161P2F10B		
Pos	123456789012345	score
2	ILQLKTYLPTFETPI	6

Table L: Properties of 161P2F10B

	Bioinformatic Program	URL (Located on the World Wide Web at)	Outcome
Feature	Bioinformatic Program		Outcome
ORF (includes stop codon)	ORF finder	.ncbi.nlm.nih.gov/	
# of amino acids			875
Transmembrane region	TM Pred	.ch.embnet.org/	
			One TM, aa 23-41
	HMMTop	.enzim.hu/hmmtop/	
			One TM, aa 23-45
	Sosui	.genome.ad.jp/SOSui/	
Signal Peptide	TMHMM	.cbs.dtu.dk/services/TMHMM	
	Signal P	.cbs.dtu.dk/services/SignalP/	
			One TM, aa 23-45
pI	pI/MW tool	.expasy.ch/tools/	None
Molecular weight			6.12
	pI/MW tool	.expasy.ch/tools/	
Localization			100.09 kDa
	PSORT	psort.nibb.ac.jp/	Plasma membrane 74% Golgi 30%
	PSORT II	psort.nibb.ac.jp/	Endoplasmic 30.4% Golgi 21.7%
Motifs	Pfam	.sanger.ac.uk/Pfam/	Somatomedin B, Type I phosphodiesterase / nucleotide pyrophosphatase
	Prints	.biochem.ucl.ac.uk/	Cell Attachment RGD
	Blocks	.blocks.fhcrc.org/	Somatomedin B, DNA/RNA non-specific endonuclease,
	Prosite	.genome.ad.jp/	Somatomedin B

Table LI. Nucleotide sequence of transcript variant 161P2F10B v.6 (SEQ ID NO: 91)

```

atacagtttc tctttgccag actagactaa agaaggagca ctactttatt ctgataaaac      60
aggtctatgc agctaccagg acaatggaat ctacgttgac tttagcaacg gaacaacctg      120
ttaagaagaa cactcttaag aaatataaaa tagcttgcac tgttcttctt gctttgctgg      180
tgatcatgtc acttgatta ggcctggggc ttggactcag gaaactggaa aagcaaggca      240
gctgcaggaa gaagtgcttt gatgcatcat ttagaggact ggagaactgc cgggtgtgatg      300
tggcatgtaa agaccgaggt gattgctgct gggattttga agacacctgt gtggaatcaa      360
ctcgaatatg gatgtgcaat aaatttcggt gtggagagac cagattagag gccagccttt      420
gctcttgttc agatgactgt ttgcagagga aagattgctg tgctgactat aagagtgttt      480
gccaaggaga aacctcatgg ctggaagaaa actgtgacac agcccagcag tctcagtgcc      540
cagaagggtt tgacctgcca ccagttatct tgttttctat ggatggattt agagctgaat      600
atztatcac atgggatact ttaatgcaa atatcaataa actgaaaaca tgtggaattc      660
attcaaaata catgagagct atgtatccta ccaaaacctt cccaaatcat tacaccattg      720
tcacgggctt gtatccggag tcacatggca tcattgacaa taatatgtat gatgtaaatc      780
tcaacaagaa tttttcactt tcttcaaagg aacaaaataa tccagcctgg tggcatgggc      840
aaccaatgtg gctgacagca atgtatcaag gtttaaaagc cgctacctac ttttggcccg      900
gatcagaagt ggctataaat ggctcctttc cttccatata catgccttac aacggaagtg      960
tcccatttga agagaggatt tctacactgt taaaatggct ggacctgccc aaagctgaga     1020
gacctcaggt ttataccatg ttttttgaag aacctgattc ctctggacat gcaggtggac     1080
cagtcagtgc cagagtaatt aaagccttac aggtagtaga tcatgctttt gggatgttga     1140
tggaaggcct gaagcagcgg aatttgcaca actgtgtcaa tatcatcctt ctggctgacc     1200

```

```

atggaatgga ccagacttat tgtaacaaga tgggaatacat gactgattat tttcccagaa 1260
taaacttctt ctacatgtac gaagggcctg cccccgcgt cagagctcat aatatacctc 1320
atgacttttt tagttttaat tctgaggaaa ttgttagaaa cctcagttgc cgaaaacctg 1380
atcagcattt caagccctat ttgactcctg atttgccaaa gcgactgcac tatgccaaga 1440
acgtcagaat cgacaaagtt catctctttg tggatcaaca gtggctggct gttaggagta 1500
aatcaaatac aaattgtgga ggaggcaacc atgggtataa caatgagttt aggagcatgg 1560
aggctatctt tctggcacat ggacccagtt ttaaagagaa gactgaagtt gaaccatttg 1620
aaaatattga agtctataac ctaatgtgtg atcttctacg cattcaacca gcaccaaaca 1680
atggaacca tggtagttta aaccatcttc tgaagggtgcc tttttatgag ccatcccatg 1740
cagaggaggt gtcaaagttt tctgtttgtg gctttgctaa tccattgccc acagagtctc 1800
ttgactgttt ctgccctcac ctacaaaata gtactcagct ggaacaagtg aatcagatgc 1860
taaactctac ccaagaagaa ataacagcaa cagtgaagtt aaatttgcca tttgggagggc 1920
ctaggggtact gcagaagaac gtggaccact gtctccttta ccacaggga tatgtcagtg 1980
gatttggaag agctatgagg atgcccattg ggagttcata cacagtcctc cagttgggag 2040
acacatcgcc tctgcctccc actgtcccag actgtctgct ggctgatgtc agggttcctc 2100
cttctgagag ccaaaaatgt tccttctatt tagcagacaa gaatatcacc cacggcttcc 2160
tctatctctc tgccagcaat agaacatcag atagccaata tgatgcttta attactagca 2220
atgttggtacc tatgtatgaa gaattcagaa aaatgtggga ctacttccac agtgttcttc 2280
ttataaaaca tgccacagaa agaaatggag taaatgtggt tagtggacca atatttgatt 2340
ataattatga tggccatttt gatgtctccag atgaaattac caaacattta gccaacactg 2400
atgttcccat cccaacacac tactttgtgg tgctgaccag ttgtaaaaac aagagccaca 2460
caccggaaaa ctgccctggg tggctggatg tcctaccctt tatcatccct caccgacctc 2520
ccaacgtgga gagctgtcct gaaggtaaac cagaagctct ttgggttgaa gaaagattta 2580
cagctcacat tgcccgggtc cgtgatgtag aacttctcac tgggcttgac ttctatcagg 2640
ataaagtgca gctgtctct gaaattttgc aactaaagac atatttacca acatttgaaa 2700
ccactattta acttaataat gtctacttaa tatataattt actgtataaa gtaatttttg 2760
caaaatataa gtgatttttt tctggagaat tgtaaaataa agttttctat ttttccttaa 2820
gtcccctaaa agccataatt tttattattc ctttttctct tttttcaatt ctatgaatat 2880
gtattatttt aaagttatat ttttcacaca gagatgatgc tatattacac cttccctttt 2940
ttgttggttt cttaaactct aatctcatga cagattatac cttccttatt acttgtttta 3000
tcttactcag aatctttgaa tatatttttc tgcccagaat tatctaaaca aaaggagaa 3060
caaaagaagt atgtctcact tgggaactga atcaactcta aatcagtttt gtcacaaaac 3120
tttttgtatt tgactggcaa tgctgattaa aattaaaaat gcaca 3165

```

Table LII. Nucleotide sequence alignment of 161P2F10B v.1 (SEQ ID NO: 92) and 161P2F10B v.6 (SEQ ID NO: 93)

Score = 5301 bits (2757), Expect = 0.0 Identities = 2774/2780 (99%), Gaps = 1/2780 (0%) Strand = Plus / Plus

```

v.1: 1      ctactttattctgataaaaacaggtctatgcagctaccaggacaatggaatctacgttgac 60
          ||||||||||||||||||||||||||||||||||||||||||||||||||||||||
v.6: 41     ctactttattctgataaaaacaggtctatgcagctaccaggacaatggaatctacgttgac 100

v.1: 61     tttagcaacggaacaacctgttaagaagaacactcttaagaaatataaaaatagcttgcat 120
          ||||||||||||||||||||||||||||||||||||||||||||||||||||||||
v.6: 101    tttagcaacggaacaacctgttaagaagaacactcttaagaaatataaaaatagcttgcat 160

v.1: 121    tgttcttcttgctttgctgggtgatcatgtcacttggttaggcctggggcttgactcag 180
          ||||||||||||||||||||||||||||||||||||||||||||||||||||||||
v.6: 161    tgttcttcttgctttgctgggtgatcatgtcacttggttaggcctggggcttgactcag 220

v.1: 181    gaaactggaaaagcaaggcagctgcaggaagaagtgctttgatgcatcatttagaggact 240
          ||||||||||||||||||||||||||||||||||||||||||||||||||||||||
v.6: 221    gaaactggaaaagcaaggcagctgcaggaagaagtgctttgatgcatcatttagaggact 280

v.1: 241    ggagaactgccggtgtgatgtggcatgtaaagaccgaggtgattgctgctgggattttga 300
          ||||||||||||||||||||||||||||||||||||||||||||||||||||||||
v.6: 281    ggagaactgccggtgtgatgtggcatgtaaagaccgaggtgattgctgctgggattttga 340

v.1: 301    agacacctgtgtggaatcaactcgaatatggatgtgcaataaatttcgttgaggagagac 360
          ||||||||||||||||||||||||||||||||||||||||||||||||||||||||
v.6: 341    agacacctgtgtggaatcaactcgaatatggatgtgcaataaatttcgttgaggagagac 400

```

v.1: 361 cagattagaggccagcctttgctcttggttcagatgactgtttgcagaagaaagattgctg 420
|||||
v.6: 401 cagattagaggccagcctttgctcttggttcagatgactgtttgcagaggaaagattgctg 460
|||||

v.1: 421 tgctgactataagagtgtttgccaaggagaaacctcatggctggaagaaaactgtgacac 480
|||||
v.6: 461 tgctgactataagagtgtttgccaaggagaaacctcatggctggaagaaaactgtgacac 520
|||||

v.1: 481 agcccagcagtcctcagtgcccagaagggtttgacctgccaccagttatcttggttttctat 540
|||||
v.6: 521 agcccagcagtcctcagtgcccagaagggtttgacctgccaccagttatcttggttttctat 580
|||||

v.1: 541 ggatggatttagagctgaatatattatacacatgggatactttaatgccaaatatcaataa 600
|||||
v.6: 581 ggatggatttagagctgaatatattatacacatgggatactttaatgccaaatatcaataa 640
|||||

v.1: 601 actgaaaacatgtggaattcattcaaaatacatgagagctatgtatcctacaaaacctt 660
|||||
v.6: 641 actgaaaacatgtggaattcattcaaaatacatgagagctatgtatcctacaaaacctt 700
|||||

v.1: 661 cccaaatcattacaccattgtcacgggcttgatccagagtcacatggcatcattgacaa 720
|||||
v.6: 701 cccaaatcattacaccattgtcacgggcttgatccggagtcacatggcatcattgacaa 760
|||||

v.1: 721 taatatgtatgatgtaaattctcaacaagaatttttcttcttcaaaggaacaaaataa 780
|||||
v.6: 761 taatatgtatgatgtaaattctcaacaagaatttttcttcttcaaaggaacaaaataa 820
|||||

v.1: 781 tccagcctggtggcatgggcaaccaatgtggctgacagcaatgtatcaaggtttaaaagc 840
|||||
v.6: 821 tccagcctggtggcatgggcaaccaatgtggctgacagcaatgtatcaaggtttaaaagc 880
|||||

v.1: 841 cgctacctacttttggcccggatcagaagtggctataaatggctcctttccttccatata 900
|||||
v.6: 881 cgctacctacttttggcccggatcagaagtggctataaatggctcctttccttccatata 940
|||||

v.1: 901 catgccttacaacggaagtgtcccatttgaagagaggatttctacactgttaaaatggct 960
|||||
v.6: 941 catgccttacaacggaagtgtcccatttgaagagaggatttctacactgttaaaatggct 1000
|||||

v.1: 961 ggacctgccc aaagctgaaagacccaggttttataccatgtattttgaagaacctgattc 1020
|||||
v.6: 1001 ggacctgccc aaagctgagagacccaggttttataccatgtatttttgaagaacctgattc 1060
|||||

v.1: 1021 ctctggacatgcaggtggaccagtcagtgccagagtaattaaagccttacaggtagtaga 1080
|||||
v.6: 1061 ctctggacatgcaggtggaccagtcagtgccagagtaattaaagccttacaggtagtaga 1120
|||||

v.1: 1081 tcatgcttttgggatgttgatggaaggcctgaagcagcggaatttgcacaactgtgtcaa 1140
|||||
v.6: 1121 tcatgcttttgggatgttgatggaaggcctgaagcagcggaatttgcacaactgtgtcaa 1180
|||||

v.1: 1141 tatcatccttctggctgacctggaatggaccagacttattgtaacaagatggaatacat 1200
|||||
v.6: 1181 tatcatccttctggctgacctggaatggaccagacttattgtaacaagatggaatacat 1240
|||||

v.1: 1201 gactgattatccccagaataaacttcttctacatgtacgaagggcctgccccccgcat 1260
|||||
v.6: 1241 gactgattatccccagaataaacttcttctacatgtacgaagggcctgccccccgcat 1300
|||||

v.1: 1261 ccgagctcataataacatcatgacttttttagttttaattctgaggaaattggttagaaa 1320
|||||
v.6: 1301 ccgagctcataataacatcatgacttttttagttttaattctgaggaaattggttagaaa 1360
|||||

v.1: 1321 cctcagttgccgaaaacatgatcagcatttcaagccctatttgactcctgatttgccaaa 1380
|||||
v.6: 1361 cctcagttgccgaaaacatgatcagcatttcaagccctatttgactcctgatttgccaaa 1420
|||||

v.1: 1381 gcgactgcactatgccaagaacgtcagaatcgacaaagttcatctcttctgtggatcaaca 1440
|||||
v.6: 1421 gcgactgcactatgccaagaacgtcagaatcgacaaagttcatctcttctgtggatcaaca 1480
|||||

v.1: 1441 gtggctggctgttaggagtaaatacaatacaaatgtggaggaggcaaccatgggtataa 1500
|||||
v.6: 1481 gtggctggctgttaggagtaaatacaatacaaatgtggaggaggcaaccatgggtataa 1540
|||||

v.1: 1501 caatgagtttaggagcatggaggctatcttctggcacatggaccagttttaagagaa 1560
|||||
v.6: 1541 caatgagtttaggagcatggaggctatcttctggcacatggaccagttttaagagaa 1600
|||||

v.1: 1561 gactgaagttgaaccatttgaaaatattgaagtctataacctaattgtgtgatcttctacg 1620
|||||
v.6: 1601 gactgaagttgaaccatttgaaaatattgaagtctataacctaattgtgtgatcttctacg 1660
|||||

v.1: 1621 cattcaaccagcaccaacaatggaacccatggtagtttaaacatcttctgaagggtgcc 1680
|||||
v.6: 1661 cattcaaccagcaccaacaatggaacccatggtagtttaaacatcttctgaagggtgcc 1720
|||||

v.1: 1681 tttttatgagccatcccatgcagaggaggtgtcaaagtcttctgttggcttctgctaa 1740
|||||
v.6: 1721 tttttatgagccatcccatgcagaggaggtgtcaaagtcttctgttggcttctgctaa 1780
|||||

v.1: 1741 tccattgcccacagagtctcttgactgttctgcccctcacctacaaaatagtactcagct 1800
|||||
v.6: 1781 tccattgcccacagagtctcttgactgttctgcccctcacctacaaaatagtactcagct 1840
|||||

v.1: 1801 ggaacaagtgaatcagatgctaaatctcacccaagaagaataacagcaacagtgaaggt 1860
|||||
v.6: 1841 ggaacaagtgaatcagatgctaaatctcacccaagaagaataacagcaacagtgaaggt 1900
|||||

v.1: 1861 aaatttgccatttgggaggcctagggtagtgcagaagaacgtggaccactgtctccttta 1920
|||||
v.6: 1901 aaatttgccatttgggaggcctagggtagtgcagaagaacgtggaccactgtctccttta 1960
|||||

v.1: 1921 ccacagggaatatgtcagtggtttggaaaagctatgaggatgcccattgtggagttcata 1980
|||||
v.6: 1961 ccacagggaatatgtcagtggtttggaaaagctatgaggatgcccattgtggagttcata 2020
|||||

v.1: 1981 cacagtccccagttgggagacacatcgccctctgcctcccactgtcccagactgtctgcg 2040
|||||
v.6: 2021 cacagtccccagttgggagacacatcgccctctgcctcccactgtcccagactgtctgcg 2080
|||||

v.1: 2041 ggctgatgtcaggggttcctccttctgagagccaaaaatgttccttctatttagcagacaa 2100
 |||||
 v.6: 2081 ggctgatgtcaggggttcctccttctgagagccaaaaatgttccttctatttagcagacaa 2140

v.1: 2101 gaatatcaccacggcttcctctatcctcctgccagcaatagaacatcagatagccaata 2160
 |||||
 v.6: 2141 gaatatcaccacggcttcctctatcctcctgccagcaatagaacatcagatagccaata 2200

v.1: 2161 tgatgctttaattactagcaatttggtacctatgtatgaagaattcagaaaaatgtggga 2220
 |||||
 v.6: 2201 tgatgctttaattactagcaatttggtacctatgtatgaagaattcagaaaaatgtggga 2260

v.1: 2221 ctacttccacagtgttcttcttataaaacatgccacagaaagaaatggagtaaattgtggt 2280
 |||||
 v.6: 2261 ctacttccacagtgttcttcttataaaacatgccacagaaagaaatggagtaaattgtggt 2320

v.1: 2281 tagtggaccaatatttgattataattatgatggccattttgatgctccagatgaaattac 2340
 |||||
 v.6: 2321 tagtggaccaatatttgattataattatgatggccattttgatgctccagatgaaattac 2380

v.1: 2341 caaacatttagccaacactgatgttcccatcccaacacactactttgtggtgctgaccag 2400
 |||||
 v.6: 2381 caaacatttagccaacactgatgttcccatcccaacacactactttgtggtgctgaccag 2440

v.1: 2401 ttgtaaaaacaagagccacacaccggaaaactgccctgggtggctggatgtcctaccctt 2460
 |||||
 v.6: 2441 ttgtaaaaacaagagccacacaccggaaaactgccctgggtggctggatgtcctaccctt 2500

v.1: 2461 tatcatccctcaccgacctaccaacgtggagagctgtcctgaaggtaaaccagaagctct 2520
 |||||
 v.6: 2501 tatcatccctcaccgacctaccaacgtggagagctgtcctgaaggtaaaccagaagctct 2560

v.1: 2521 ttgggttgaagaaagatttacagctcacattgcccgggtccgtgatgtagaacttctcac 2580
 |||||
 v.6: 2561 ttgggttgaagaaagatttacagctcacattgcccgggtccgtgatgtagaacttctcac 2620

v.1: 2581 tgggcttgacttctatcaggataaagtgcagcctgtctctgaaattttgcaactaaagac 2640
 |||||
 v.6: 2621 tgggcttgacttctatcaggataaagtgcagcctgtctctgaaattttgcaactaaagac 2680

v.1: 2641 atatttaccacatttgaaaccactatttaacttaataatgtctacttaatatataattt 2700
 |||||
 v.6: 2681 atatttaccacatttgaaaccactatttaacttaataatgtctacttaatatataattt 2740

v.1: 2701 actgtataaagtaattttggcaaaatataagtga-ttttttctggagaattgtaaaataa 2759
 |||||
 v.6: 2741 actgtataaagtaattttggcaaaatataagtgaatttttctggagaattgtaaaataa 2800

v.1: 2760 agttttctatttttccttaa 2779
 |||||
 v.6: 2801 agttttctatttttccttaa 2820

Table LIII. Peptide sequences of protein coded by 161P2F10B v.6 (SEQ ID NO: 94)

MESTLTATE QPVKNTLKK YKIACIVLLA LLVIMSLGLG LGLGLRKLEK QGSCRKKCFD	60
ASFRGLENCR CDVACKDRGD CCWDFEDTCV ESTRIWMCNK FRCGETRLEA SLCSCSDDCL	120
QKKDCCADYK SVCQGETSWL EENCDTAQQS QCPEGFDLPP VILFSMDGFR AEYLYTWDTL	180

MPNINKLKTC	GIH SKYMRAM	YPTKTFPNHY	TIVTGLYPES	HGIIDNNMYD	VNLNKNFSL	240
SKEQNNPAWW	HGQPMWLTAM	YQGLKAATYF	WPGSEVAING	SFPSIYMPYN	GSVPFEERIS	300
TLLKWLDLPK	AERPRFYTM	FEEPDSGHA	GGPV SARVIK	ALQVVDHAFG	MLMEGLKQRN	360
LHNCVNIILL	ADHGMDQTYC	NKMEYMTDYF	PRINFFYMYE	GPAPRIRAHN	IPHDFFSFNS	420
EEIVRNLSR	KPDQHF KPYL	TPDL PKRLHY	AKNVRIDKVH	LFVDQQWLAV	RSKSNTNCGG	480
GNHGYNNEFR	SMEAI FLAHG	PSFKEKTEVE	PFENIEVYNL	MCDLLRIQPA	PNNNGTHGSLN	540
HLLKVPFYEP	SHAEVSKFS	VCGFANPLPT	ESLDCFCPHL	QNSTQLEQVN	QMLNLTQEEI	600
TATVKVNL PF	GRPRVLQKNV	DHCLLYHREY	VSGFGKAMRM	PMWSSYTPVQ	LGDT SPLPPT	660
VPDCLRADVR	VPPSESQKCS	FYLADKNITH	GFLYP PASNR	TSDSQYDALI	TSNLVPMYEE	720
FRKMWDYFHS	VLLIKHATER	NGVNVVSGPI	FDYNYDGHFD	APDEITKHLA	NTDVPIPTHY	780
FVVLTSCKNK	SHTPENCPGW	LDVLPFIIPH	RPTNVE SCPE	GKPEALWVEE	RFTAH IARVR	840
DVELLTGLDF	YQDKVQPVSE	ILQLKTYLPT	FETTI			875

Table LIV. Amino acid sequence alignment of 161P2F10Bv.1 v.1 (SEQ ID NO: 95) and 161P2F10B v.6 (SEQ ID NO: 96)

Score = 1855 bits (4804), Expect = 0.0 Identities = 875/875 (100%), Positives = 875/875 (100%)

161P2F10Bv.1: 1	MESTLTLATEQPVKKNTLKKYKIACIVLLALLVIMSLGLGLGLRLKLEKQGSCRKKCFD	60
	MESTLTLATEQPVKKNTLKKYKIACIVLLALLVIMSLGLGLGLRLKLEKQGSCRKKCFD	
161P2F10Bv.6: 1	MESTLTLATEQPVKKNTLKKYKIACIVLLALLVIMSLGLGLGLRLKLEKQGSCRKKCFD	60
161P2F10Bv.1: 61	ASFRGLENCRCDVACKDRGDCCWDFEDTCVESTRIWMCNKFRCGETRLEASLCSCSDDCL	120
	ASFRGLENCRCDVACKDRGDCCWDFEDTCVESTRIWMCNKFRCGETRLEASLCSCSDDCL	
161P2F10Bv.6: 61	ASFRGLENCRCDVACKDRGDCCWDFEDTCVESTRIWMCNKFRCGETRLEASLCSCSDDCL	120
161P2F10Bv.1: 121	QKKDCCADYKSVCQGETSWLEENCDAQQSQCEGFDLPPVILFSMDGFRAEYLYTWDTL	180
	QKKDCCADYKSVCQGETSWLEENCDAQQSQCEGFDLPPVILFSMDGFRAEYLYTWDTL	
161P2F10Bv.6: 121	QKKDCCADYKSVCQGETSWLEENCDAQQSQCEGFDLPPVILFSMDGFRAEYLYTWDTL	180
161P2F10Bv.1: 181	MPNINKLKTCGIH SKYMRAMYPTKTFPNHYTIVTGLYPESHGIIIDNNMYDVNLNKNFSL	240
	MPNINKLKTCGIH SKYMRAMYPTKTFPNHYTIVTGLYPESHGIIIDNNMYDVNLNKNFSL	
161P2F10Bv.6: 181	MPNINKLKTCGIH SKYMRAMYPTKTFPNHYTIVTGLYPESHGIIIDNNMYDVNLNKNFSL	240
161P2F10Bv.1: 241	SKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGSEVAINGSFPSIYMPYNGSVPFEEERIS	300
	SKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGSEVAINGSFPSIYMPYNGSVPFEEERIS	
161P2F10Bv.6: 241	SKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGSEVAINGSFPSIYMPYNGSVPFEEERIS	300
161P2F10Bv.1: 301	TLLKWLDLPKAERPRFYTMFEEPDSGHAAGGPVSARVIKALQVVDHAFGMLMEGLKQRN	360
	TLLKWLDLPKAERPRFYTMFEEPDSGHAAGGPVSARVIKALQVVDHAFGMLMEGLKQRN	
161P2F10Bv.6: 301	TLLKWLDLPKAERPRFYTMFEEPDSGHAAGGPVSARVIKALQVVDHAFGMLMEGLKQRN	360
161P2F10Bv.1: 361	LHNCVNIILLADHGMDQTYCNKMEYMTDYFPRINFFYMYEGPAPRIRAHNIPHDFFSFNS	420
	LHNCVNIILLADHGMDQTYCNKMEYMTDYFPRINFFYMYEGPAPRIRAHNIPHDFFSFNS	
161P2F10Bv.6: 361	LHNCVNIILLADHGMDQTYCNKMEYMTDYFPRINFFYMYEGPAPRIRAHNIPHDFFSFNS	420
161P2F10Bv.1: 421	EEIVRNLSRKPQDQHF KPYLTPDL PKRLHYAKNVRIDKVHLFVDQQWLAVRSKSNTNCGG	480
	EEIVRNLSRKPQDQHF KPYLTPDL PKRLHYAKNVRIDKVHLFVDQQWLAVRSKSNTNCGG	
161P2F10Bv.6: 421	EEIVRNLSRKPQDQHF KPYLTPDL PKRLHYAKNVRIDKVHLFVDQQWLAVRSKSNTNCGG	480
161P2F10Bv.1: 481	GNHGYNNEFRSMEAI FLAHGPSFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLN	540
	GNHGYNNEFRSMEAI FLAHGPSFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLN	
161P2F10Bv.6: 481	GNHGYNNEFRSMEAI FLAHGPSFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLN	540
161P2F10Bv.1: 541	HLLKVPFYEP SHAEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEI	600
	HLLKVPFYEP SHAEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEI	
161P2F10Bv.6: 541	HLLKVPFYEP SHAEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEI	600
161P2F10Bv.1: 601	TATVKVNL PFGRPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDT SPLPPT	660
	TATVKVNL PFGRPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDT SPLPPT	
161P2F10Bv.6: 601	TATVKVNL PFGRPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDT SPLPPT	660
161P2F10Bv.1: 661	VPDCLRADVRVPPSESQKCSFYLA DNITHGFLYP PASNRTSDSQYDALITSNLVPMYEE	720
	VPDCLRADVRVPPSESQKCSFYLA DNITHGFLYP PASNRTSDSQYDALITSNLVPMYEE	
161P2F10Bv.6: 661	VPDCLRADVRVPPSESQKCSFYLA DNITHGFLYP PASNRTSDSQYDALITSNLVPMYEE	720
161P2F10Bv.1: 721	FRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHY	780
	FRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHY	
161P2F10Bv.6: 721	FRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHY	780

161P2F10Bv.1: 781 FVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVR 840
 FVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVR
 161P2F10Bv.6: 781 FVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVR 840
 161P2F10Bv.1: 841 DVELLTGLDFYQDKVQPVSEILQLKTYLPTFETI 875
 DVELLTGLDFYQDKVQPVSEILQLKTYLPTFETI
 161P2F10Bv.6: 841 DVELLTGLDFYQDKVQPVSEILQLKTYLPTFETI 875

Table LV. Nucleotide sequence of transcript variant 161P2F10B v.7 (SEQ ID NO: 97)

ctactttatt	ctgataaaac	aggtctatgc	agctaccagg	acaatggaat	ctacgttgac	60
tttagcaacg	gaacaacctg	ttaagaagaa	cactcttaag	aaatataaaa	tagcttgcat	120
tacaggggtct	ctctcctttg	ggatctcacc	tcaccacaac	ctctgtttcc	caggctcaag	180
tgatcctcct	gcctcagcct	cctgagtagc	ttggaccaca	ggcacatgcc	acaaggctca	240
gctaagtttt	tgttcttctt	gctttgctgg	tgatcatgtc	acttggatta	ggcctggggc	300
ttggactcag	gaaactggaa	aagcaaggca	gctgcaggaa	gaagtgcttt	gatgcatcat	360
ttagaggact	ggagaactgc	cgggtgtgatg	tggcatgtaa	agaccgaggt	gattgctgct	420
gggattttga	agacacctgt	gtggaatcaa	ctcgaatatg	gatgtgcaat	aaatttcggt	480
gtggagagac	cagattagag	gccagccttt	gctcttgctc	agatgactgt	ttgcagaaga	540
aagattgctg	tgctgactat	aagagtgttt	gccaaggaga	aacctcatgg	ctggaagaaa	600
actgtgacac	agcccagcag	tctcagtgcc	cagaagggtt	tgacctgcca	ccagttatct	660
tgttttctat	ggatggattt	agagctgaat	atttatacac	atgggatact	ttaatgccaa	720
atatcaataa	actgaaaaca	tgtggaattc	attcaaaaata	catgagagct	atgtatccta	780
ccaaaacctt	cccaaatcat	tacaccattg	tcacgggctt	gtatccagag	tcacatggca	840
tcattgacaa	taatatgtat	gatgtaaatc	tcaacaagaa	tttttcactt	tcttcaaagg	900
aacaaaataa	tccagcctgg	tggcatgggc	aaccaatgtg	gctgacagca	atgtatcaag	960
gtttaaaagc	cgctacctac	ttttggcccg	gatcagaagt	ggctataaat	ggctcctttc	1020
cttccatata	catgccttac	aacggaagtg	tcccatttga	agagaggatt	tctacactgt	1080
taaaatggct	ggacctgccc	aaagctgaaa	gacctcaggt	ttataccatg	tattttgaag	1140
aacctgattc	ctctggacat	gcaggtggac	cagtcagtgc	cagagtaatt	aaagccttac	1200
aggtagtaga	tcattgctttt	gggatgttga	tgggaaggcct	gaagcagcgg	aatttgcaca	1260
actgtgtcaa	tatcatcctt	ctggctgacc	atggaatgga	ccagacttat	tgtaacaaga	1320
tggaatacat	gactgattat	tttcccagaa	taaacttctt	ctacatgtac	gaagggcctg	1380
ccccccgcac	ccgagctcat	aatatacctc	atgacttttt	tagttttaat	tctgaggaaa	1440
ttgttagaaa	cctcagttgc	cgaaaacctg	atcagcattt	caagccctat	ttgactcctg	1500
atlttgccaaa	gcgactgcac	tatgccaaga	acgtcagaat	cgacaaagtt	catctctttg	1560
tggatcaaca	gtggctggct	gttaggagta	aatcaaatac	aaattgtgga	ggaggcaacc	1620
atggttataa	caatgagttt	aggagcatgg	aggctatctt	tctggcacat	ggacctcagt	1680
ttaaagagaa	gactgaagtt	gaaccatttg	aaaatattga	agtctataac	ctaattgtgtg	1740
atcttctacg	cattcaacca	gcaccaaaca	atggaaccca	tggtagttaa	aacctcttct	1800
tgaagggtgcc	tttttatgag	ccatcccatg	cagaggaggt	gtcaaagttt	tctgtttgtg	1860
gctttgctaa	tccattgccc	acagagtctc	ttgactgttt	ctgccctcac	ctacaaaata	1920
gtactcagct	ggaacaagtg	aatcagatgc	taaatctcac	ccaagaagaa	ataacagcaa	1980
cagtgaaggt	aaatttgcca	tttgggaggc	ctagggtact	gcagaagaac	gtggaccact	2040
gtctccttta	ccacagggaa	tatgtcagtg	gatttggaag	agctatgagg	atgcccattgt	2100
ggagttcata	cacagtcccc	cagttgggag	acacatcgcc	tctgcctccc	actgtcccag	2160
actgtctgcg	ggctgatgtc	agggttcctc	cttctgagag	ccaaaaatgt	tccttctatt	2220
tagcagacaa	gaatatcacc	cacggcttcc	tctatcctcc	tgccagcaat	agaacatcag	2280
atagccaata	tgatgcttta	attactagca	atlttggtacc	tatgtatgaa	gaattcagaa	2340
aaatgtggga	ctacttccac	agtgttcttc	ttataaaaaca	tgccacagaa	agaaatggag	2400
taaatgtggg	tagtggacca	atatttgatt	ataattatga	tggccatttt	gatgctccag	2460
atgaaattac	caaacattta	gccaacactg	atgttcccat	cccaacacac	tactttgtgg	2520
tgctgaccag	ttgtaaaaac	aagagccaca	caccggaaaa	ctgccctggg	tggctggatg	2580
tcctaccctt	tatcatccct	caccgacctc	ccaacgtgga	gagctgtcct	gaaggtaaag	2640
cagaagctct	ttgggttgaa	gaaagattta	cagctcacat	tgcccgggtc	cgtgatgtag	2700
aacttctcac	tgggcttgac	ttctatcagg	ataaagtga	gcctgtctct	gaaattttgc	2760
aactaaagac	atattttacca	acatttgaaa	ccactattta	acttaataat	gtctacttaa	2820
tatataattt	actgtataaa	gtaatttttg	caaaaataaa	gtgatttttt	ctggagaatt	2880
gtaaaataaa	gttttctatt	tttctttaa	aaaaaaaccg	gaattccggg	cttgggaggc	2940
tgaggcagga	gactcgcttg	aaccggggag	gcagagggtg	cagtgaagca	agattgcgcc	3000
attgcactcc	agagcctggg	tgacagagca	agactacatc	tcaaaaaata	aataaataaa	3060
ataaaagtaa	caataaaaaat	aaaaagaaca	gcagagagaa	tgagcaagga	gaaatgtcac	3120
aaactattgc	aaaatactgt	tacactgggt	tggctctcca	agaagatact	ggaatctctt	3180
cagccatttg	cttttcagaa	gtagaaacca	gcaaaccacc	tctaagcgga	gaacatacga	3240
ttctttatta	agtagctctg	gggaaggaaa	gaataaaaagt	tgatagctcc	ctgattggga	3300
aaaaatgcac	aattaataaa	gaatgaagat	gaaagaaagc	atgcttatgt	tgtaacacaa	3360
aaaaaattca	caaacgttgg	tgggaaggaaa	acagtataga	aaacattact	ttactaaaaa	3420
gctggaaaaa	ttttcagttg	ggatgcgact	gacaaaaaga	acgggatttc	caggcataaa	3480
gttggcgtga	gctacagagg	gcaccatgtg	gctcagtggg	agacccttca	agattcaaag	3540

```

ttccatttga cagagcaaag gcacttcgca aggagaaggg tttaaattat ggggtccaaaa 3600
gccaagtggg aaagcgagca atttgcagca taactgcttc tcctagacag ggctgagtgg 3660
gcaaaatacg acagtacaca cagtgactat tagccactgc cagaaacagg ctgaacagcc 3720
ctgggagaca agggaaggca ggtgggtggga gttgttcatt gagagaaagg agagttttag 3780
aaccagcaca tccactggag atgctggggc accagacccc tcccagtcaa taaagtctgg 3840
tgcctcattt gatctcagcc tcatcatgac cctggagaga ccttgatacc atctgccagt 3900
ccccgacagc ttaggcactc cttgccatca acctgacccc ccgagtgggt ctccaggctc 3960
cctgccccac ccattcaggc cggaattc 3988

```

Table LVI. Nucleotide sequence alignment of 161P2F10B v.1 (SEQ ID NO:98) and 161P2F10B v.7 (SEQ ID NO: 99)

Score = 233 bits (121), Expect = 2e-57Identities = 121/121 (100%) Strand = Plus / Plus

```

v.1: 1 ctactttattctgataaaacaggtctatgcagctaccaggacaatggaatctacgttgac 60
      |||
v.7: 1 ctactttattctgataaaacaggtctatgcagctaccaggacaatggaatctacgttgac 60

```

```

v.1: 61 tttagcaacggaacaacctgttaagaagaacactcttaagaaatataaaatagcttgcat 120
      |||
v.7: 61 tttagcaacggaacaacctgttaagaagaacactcttaagaaatataaaatagcttgcat 120

```

```

v.1: 121 t 121
      |
v.7: 121 t 121

```

Score = 7189 bits (3739), Expect = 0.0Identities = 3739/3739 (100%) Strand = Plus / Plus

```

v.1: 120 ttgttcttcttgctttgctggtgatcatgtcacttggattaggcctggggcttggactca 179
      |||
v.7: 250 ttgttcttcttgctttgctggtgatcatgtcacttggattaggcctggggcttggactca 309

```

```

v.1: 180 ggaaactggaaaagcaaggcagctgcaggaagaagtgctttgatgcatcatttagaggac 239
      |||
v.7: 310 ggaaactggaaaagcaaggcagctgcaggaagaagtgctttgatgcatcatttagaggac 369

```

```

v.1: 240 tggagaactgccggtgtgatgtggcatgtaaagaccgaggtgattgctgctgggattttg 299
      |||
v.7: 370 tggagaactgccggtgtgatgtggcatgtaaagaccgaggtgattgctgctgggattttg 429

```

```

v.1: 300 aagacacctgtgtggaatcaactcgaatatggatgtgcaataaatttcgttgtggagaga 359
      |||
v.7: 430 aagacacctgtgtggaatcaactcgaatatggatgtgcaataaatttcgttgtggagaga 489

```

```

v.1: 360 ccagattagaggccagcctttgctcttgcttcagatgactgtttgcagaagaaagattgct 419
      |||
v.7: 490 ccagattagaggccagcctttgctcttgcttcagatgactgtttgcagaagaaagattgct 549

```

```

v.1: 420 gtgctgactataagagtgtttgccaaggagaaacctcatggctggaagaaaactgtgaca 479
      |||
v.7: 550 gtgctgactataagagtgtttgccaaggagaaacctcatggctggaagaaaactgtgaca 609

```

```

v.1: 480 cagcccagcagtctcagtgccagaagggtttgacctgccaccagttatcttgttttcta 539
      |||
v.7: 610 cagcccagcagtctcagtgccagaagggtttgacctgccaccagttatcttgttttcta 669

```

```

v.1: 540 tggatggatttagagctgaatatattatacacatgggatactttaatgccaaatatcaata 599
      |||
v.7: 670 tggatggatttagagctgaatatattatacacatgggatactttaatgccaaatatcaata 729

```

v.1: 600 aactgaaaacatgtggaattcattcaaaatacatgagagctatgtatcctacaaaacct 659
|||||
v.7: 730 aactgaaaacatgtggaattcattcaaaatacatgagagctatgtatcctacaaaacct 789
|||||

v.1: 660 tcccaaatacattacaccattgtcacgggcttgtatccagagtcacatggcatcattgaca 719
|||||
v.7: 790 tcccaaatacattacaccattgtcacgggcttgtatccagagtcacatggcatcattgaca 849
|||||

v.1: 720 ataatatgtatgatgtaaatctcaacaagaatttttctactttcttcaaaggaacaaaata 779
|||||
v.7: 850 ataatatgtatgatgtaaatctcaacaagaatttttctactttcttcaaaggaacaaaata 909
|||||

v.1: 780 atccagcctggtggcatgggcaaccaatgtggctgacagcaatgtatcaaggttttaaag 839
|||||
v.7: 910 atccagcctggtggcatgggcaaccaatgtggctgacagcaatgtatcaaggttttaaag 969
|||||

v.1: 840 ccgctacctacttttggcccggatcagaagtggtataaatggctcctttccttccatat 899
|||||
v.7: 970 ccgctacctacttttggcccggatcagaagtggtataaatggctcctttccttccatat 1029
|||||

v.1: 900 acatgccttacaacggaagtgtcccatttgaagagaggatttctacactgttaaaatggc 959
|||||
v.7: 1030 acatgccttacaacggaagtgtcccatttgaagagaggatttctacactgttaaaatggc 1089
|||||

v.1: 960 tggacctgccc aaagctgaaagacccagggtttataccatgtattttgaagaacctgatt 1019
|||||
v.7: 1090 tggacctgccc aaagctgaaagacccagggtttataccatgtattttgaagaacctgatt 1149
|||||

v.1: 1020 cctctggacatgcaggtggaccagtcagtgccagagtaattaaagccttacaggtagtag 1079
|||||
v.7: 1150 cctctggacatgcaggtggaccagtcagtgccagagtaattaaagccttacaggtagtag 1209
|||||

v.1: 1080 atcatgcttttgggatgttgatggaaggcctgaagcagcggaaatttgcaactgtgtca 1139
|||||
v.7: 1210 atcatgcttttgggatgttgatggaaggcctgaagcagcggaaatttgcaactgtgtca 1269
|||||

v.1: 1140 atatcatccttctggctgaccatggaatggaccagacttattgtaacaagatggaataca 1199
|||||
v.7: 1270 atatcatccttctggctgaccatggaatggaccagacttattgtaacaagatggaataca 1329
|||||

v.1: 1200 tgactgattattttcccagaataaacttcttctacatgtacgaagggcctgcccccgca 1259
|||||
v.7: 1330 tgactgattattttcccagaataaacttcttctacatgtacgaagggcctgcccccgca 1389
|||||

v.1: 1260 tccgagctcataataatacctcatgacttttttagttttaattctgaggaaattgtagaa 1319
|||||
v.7: 1390 tccgagctcataataatacctcatgacttttttagttttaattctgaggaaattgtagaa 1449
|||||

v.1: 1320 acctcagttgccgaaaacctgatcagcatttcaagccctatttgactcctgatttgccaa 1379
|||||
v.7: 1450 acctcagttgccgaaaacctgatcagcatttcaagccctatttgactcctgatttgccaa 1509
|||||

v.1: 1380 agcgactgcactatgccaaagaacgtcagaatcgacaaagttcatctctttgtggatcaac 1439
|||||
v.7: 1510 agcgactgcactatgccaaagaacgtcagaatcgacaaagttcatctctttgtggatcaac 1569
|||||

v.1: 1440 agtggctggctgtaggagtaaatacaatacaaatgtggaggaggcaaccatgggtata 1499
|||||
v.7: 1570 agtggctggctgtaggagtaaatacaatacaaatgtggaggaggcaaccatgggtata 1629

v.1: 1500 acaatgagtttaggagcatggaggctatctttctggcacatggaccagttttaagaga 1559
|||||
v.7: 1630 acaatgagtttaggagcatggaggctatctttctggcacatggaccagttttaagaga 1689

v.1: 1560 agactgaagttgaaccatttgaaaatattgaagtctataacctaattgtgtgatcttctac 1619
|||||
v.7: 1690 agactgaagttgaaccatttgaaaatattgaagtctataacctaattgtgtgatcttctac 1749

v.1: 1620 gcattcaaccagcaccaacaatggaacccatggtagtttaaacctcttctgaagggtgc 1679
|||||
v.7: 1750 gcattcaaccagcaccaacaatggaacccatggtagtttaaacctcttctgaagggtgc 1809

v.1: 1680 ctttttatgagccatcccatgcagaggaggtgtcaaagttttctgtttgtggccttgcta 1739
|||||
v.7: 1810 ctttttatgagccatcccatgcagaggaggtgtcaaagttttctgtttgtggccttgcta 1869

v.1: 1740 atccattgcccacagagtctcttgactgtttctgccctcacctacaaaatagtactcagc 1799
|||||
v.7: 1870 atccattgcccacagagtctcttgactgtttctgccctcacctacaaaatagtactcagc 1929

v.1: 1800 tggaacaagtgaatcagatgctaaatctcacccaagaagaataacagcaacagtgaag 1859
|||||
v.7: 1930 tggaacaagtgaatcagatgctaaatctcacccaagaagaataacagcaacagtgaag 1989

v.1: 1860 taaatttgccatttgggaggcctagggtactgcagaagaacgtggaccactgtctcctt 1919
|||||
v.7: 1990 taaatttgccatttgggaggcctagggtactgcagaagaacgtggaccactgtctcctt 2049

v.1: 1920 accacaggaatatgtcagtggatttggaaaagctatgaggatgcccatgtggagttcat 1979
|||||
v.7: 2050 accacaggaatatgtcagtggatttggaaaagctatgaggatgcccatgtggagttcat 2109

v.1: 1980 acacagtcccccagttgggagacacatcgctctgcctcccactgtcccagactgtctgc 2039
|||||
v.7: 2110 acacagtcccccagttgggagacacatcgctctgcctcccactgtcccagactgtctgc 2169

v.1: 2040 gggctgatgtcagggttcctccttctgagagccaaaatgttccttctatttagcagaca 2099
|||||
v.7: 2170 gggctgatgtcagggttcctccttctgagagccaaaatgttccttctatttagcagaca 2229

v.1: 2100 agaatatcaccacaggcttcctctatcctcctgccagcaatagaacatcagatagccaat 2159
|||||
v.7: 2230 agaatatcaccacaggcttcctctatcctcctgccagcaatagaacatcagatagccaat 2289

v.1: 2160 atgatgctttaattactagcaatttggtacctatgtatgaagaattcagaaaaatgtggg 2219
|||||
v.7: 2290 atgatgctttaattactagcaatttggtacctatgtatgaagaattcagaaaaatgtggg 2349

v.1: 2220 actacttccacagtgttcttcttataaaaacatgccacagaaagaaatggagtaaattgtgg 2279
|||||
v.7: 2350 actacttccacagtgttcttcttataaaaacatgccacagaaagaaatggagtaaattgtgg 2409

v.1: 2280 ttagtggaccaatatttgattataattatgatggccattttgatgctccagatgaaatta 2339
|||||
v.7: 2410 ttagtggaccaatatttgattataattatgatggccattttgatgctccagatgaaatta 2469

v.1: 2340 ccaaacatttagccaacactgatgttcccatcccaacacactactttgtggtgctgacca 2399
|||||
v.7: 2470 ccaaacatttagccaacactgatgttcccatcccaacacactactttgtggtgctgacca 2529

v.1: 2400 gttgtaaaaacaagagccacacaccggaaaactgccctgggtggctggatgtcctaccct 2459
|||||
v.7: 2530 gttgtaaaaacaagagccacacaccggaaaactgccctgggtggctggatgtcctaccct 2589

v.1: 2460 ttatcatccctcaccgacctaccaacgtggagagctgtcctgaaggtaaaccagaagctc 2519
|||||
v.7: 2590 ttatcatccctcaccgacctaccaacgtggagagctgtcctgaaggtaaaccagaagctc 2649

v.1: 2520 tttgggttgaagaaagatttacagctcacattgcccggtccgtgatgtagaacttctca 2579
|||||
v.7: 2650 tttgggttgaagaaagatttacagctcacattgcccggtccgtgatgtagaacttctca 2709

v.1: 2580 ctgggcttgacttctatcaggataaagtgcagcctgtctctgaaattttgcaactaaaga 2639
|||||
v.7: 2710 ctgggcttgacttctatcaggataaagtgcagcctgtctctgaaattttgcaactaaaga 2769

v.1: 2640 catatttaccaacatttgaaaccactatttaacttaataatgtctacttaatatataatt 2699
|||||
v.7: 2770 catatttaccaacatttgaaaccactatttaacttaataatgtctacttaatatataatt 2829

v.1: 2700 tactgtataaagtaattttggcaaaatataagtgattttttctggagaattgtaaaataa 2759
|||||
v.7: 2830 tactgtataaagtaattttggcaaaatataagtgattttttctggagaattgtaaaataa 2889

v.1: 2760 agttttctatttttccttaaaaaaaaaaaccggaattccgggcttgggaggctgaggcagg 2819
|||||
v.7: 2890 agttttctatttttccttaaaaaaaaaaaccggaattccgggcttgggaggctgaggcagg 2949

v.1: 2820 agactcgcttgaacccgggaggcagaggttgagtgagccaagattgcgccattgcactc 2879
|||||
v.7: 2950 agactcgcttgaacccgggaggcagaggttgagtgagccaagattgcgccattgcactc 3009

v.1: 2880 cagagcctgggtgacagagcaagactacatctcaaaaaataaataaaaataaaaagta 2939
|||||
v.7: 3010 cagagcctgggtgacagagcaagactacatctcaaaaaataaataaaaataaaaagta 3069

v.1: 2940 acaataaaaaataaaaagaacagcagagagaatgagcaaggagaaatgtcacaaactattg 2999
|||||
v.7: 3070 acaataaaaaataaaaagaacagcagagagaatgagcaaggagaaatgtcacaaactattg 3129

v.1: 3000 caaaatactgttacactgggttggctctccaagaagatactggaatctcttcagccattt 3059
|||||
v.7: 3130 caaaatactgttacactgggttggctctccaagaagatactggaatctcttcagccattt 3189

v.1: 3060 gcttttcagaagtagaaaccagcaaaccacctctaagcggagaacatacgattctttatt 3119
|||||
v.7: 3190 gcttttcagaagtagaaaccagcaaaccacctctaagcggagaacatacgattctttatt 3249

v.1: 3120 aagtagctctggggaaggaaagaataaaaagttgatagctccctgattgggaaaaaatgca 3179
 |||||
 v.7: 3250 aagtagctctggggaaggaaagaataaaaagttgatagctccctgattgggaaaaaatgca 3309

v.1: 3180 caattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaaaaaattc 3239
 |||||
 v.7: 3310 caattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaaaaaattc 3369

v.1: 3240 acaaacgttggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaa 3299
 |||||
 v.7: 3370 acaaacgttggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaa 3429

v.1: 3300 attttcagttgggatgcgactgacaaaaagaacgggatttccaggcataaagttggcgtg 3359
 |||||
 v.7: 3430 attttcagttgggatgcgactgacaaaaagaacgggatttccaggcataaagttggcgtg 3489

v.1: 3360 agctacagagggcaccatgtggctcagtggagacccttcaagattcaaagttccatttg 3419
 |||||
 v.7: 3490 agctacagagggcaccatgtggctcagtggagacccttcaagattcaaagttccatttg 3549

v.1: 3420 acagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaagccaagtgg 3479
 |||||
 v.7: 3550 acagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaagccaagtgg 3609

v.1: 3480 taaagcgagcaatttgcagcataactgcttctcctagacagggctgagtgggcaaaatac 3539
 |||||
 v.7: 3610 taaagcgagcaatttgcagcataactgcttctcctagacagggctgagtgggcaaaatac 3669

v.1: 3540 gacagtacacacagtgactattagccactgccagaaacaggctgaacagccctgggagac 3599
 |||||
 v.7: 3670 gacagtacacacagtgactattagccactgccagaaacaggctgaacagccctgggagac 3729

v.1: 3600 aagggaaggcaggtggtgggagttgttcattggagagaaaggagagttttagaaccagcac 3659
 |||||
 v.7: 3730 aagggaaggcaggtggtgggagttgttcattggagagaaaggagagttttagaaccagcac 3789

v.1: 3660 atccactggagatgctgggcccaccagaccctcccagtcaataaagtctggtgcctcatt 3719
 |||||
 v.7: 3790 atccactggagatgctgggcccaccagaccctcccagtcaataaagtctggtgcctcatt 3849

v.1: 3720 tgatctcagcctcatcatgaccctggagagaccctgataccatctgccagtccccgacag 3779
 |||||
 v.7: 3850 tgatctcagcctcatcatgaccctggagagaccctgataccatctgccagtccccgacag 3909

v.1: 3780 cttaggcactccttgccatcaacctgaccccccgagtgggttctccaggctccctgccccca 3839
 |||||
 v.7: 3910 cttaggcactccttgccatcaacctgaccccccgagtgggttctccaggctccctgccccca 3969

v.1: 3840 cccattcaggccggaattc 3858
 |||||
 v.7: 3970 cccattcaggccggaattc 3988

Table LVII. Peptide sequences of protein coded by 161P2F10B v.7 (SEQ ID NO: 100)

MSLGLGLGLG	LRKLEKQGSC	RKKCFDASFR	GLENCRCDDVA	CKDRGDCCWD	FEDTCVEST	60
IWMCNKFRCG	ETRLASLCS	CSDDCLQRKD	CCADYKSVQC	GETSWLEENC	DTAQSQCP	120
GFDLPPVILF	SMDGFRAEYL	YTWDTLMPNI	NKLKTCGIHS	KYMRAMYPTK	TFPNHYTIVT	180

GLYPESHGII	DNNMYDVNLN	KNFSLSSKEQ	NNPAWWHGQP	MWLTAMYQGL	KAATYFWPGS	240
EVAINGSFPS	IYMPYNGSVP	FEERISTLLK	WLDLPKAERP	RFYTMFFEEP	DSSGHAGGPV	300
SARVIKALQV	VDHAFGMLME	GLKQRNLHNC	VNIILLADHG	MDQTYCNKME	YMTDYFPRIN	360
FFYMYEGPAP	RVRAHNIPHD	FFSFNSEEIV	RNLSCRKPDQ	HFKPYLTPDL	PKRLHYAKNV	420
RIDKVHLFVD	QQWLAVRSKS	NTNCGGGNHG	YNNEFRSMEA	IFLAHGPSFK	EKTEVEPFEN	480
IEVYNLMCDL	LRIQPAPNNG	THGSLNHLLK	VPFYEPSHAE	EVSKEFSVCGF	ANPLPTESLD	540
CFCPHLQNST	QLEQVNQMLN	LTQEEITATV	KVNLPFGRPR	VLQKNVDHCL	LYHREYVSGF	600
GKAMRMPMWS	SYTVPQLGDT	SPLPPTVPDC	LRADVRVPPS	ESQKCSFYLA	DKNITHGFLY	660
PPASNRTSDS	QYDALITSNL	VPMYEEFRKM	WDYFHSVLLI	KHATERNGVN	VVSGPIFDYN	720
YDGHFDAPDE	ITKHLANTDV	PIPTHYFVVL	TSCKNKSHTP	ENCPGWLDVL	PFIIPHRPTN	780
VESCEPGKPE	ALWVEERFTA	HIARVRDVEL	LTGLDFYQDK	VQPVSEILQL	KTYLPTFETT	840
I						841

Table LVIII. Amino acid sequence alignment of 161P2F10Bv.1 v.1 (SEQ ID NO: 101) and 161P2F10B v.7 (SEQ ID NO: 102)

Score = 1789 bits (4634), Expect = 0.0 Identities = 838/841 (99%), Positives = 841/841 (99%)

161P2F10Bv.1: 35	MSLGLGLGLGLRKLEKQGSCRKKCFDASFRGLENCRCDVACKDRGDCCWDFEDTCVEST	94
161P2F10Bv.7: 1	MSLGLGLGLGLRKLEKQGSCRKKCFDASFRGLENCRCDVACKDRGDCCWDFEDTCVEST	60
161P2F10Bv.1: 95	IWMCNKFRGGETRLEASLCSCSDDCLQKDCADYKSVQGETSWLEENCDAQQSQCP	154
161P2F10Bv.7: 61	IWMCNKFRGGETRLEASLCSCSDDCLQKDCADYKSVQGETSWLEENCDAQQSQCP	120
161P2F10Bv.1: 155	GFDLPPVILFSMDGFRAEYLYTWDTLMPNINKLKTGCIH SKYMRAMYPTKTFPNHYTIVT	214
161P2F10Bv.7: 121	GFDLPPVILFSMDGFRAEYLYTWDTLMPNINKLKTGCIH SKYMRAMYPTKTFPNHYTIVT	180
161P2F10Bv.1: 215	GLYPESHGII DNNMYDVNLNKNFSLSSKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGS	274
161P2F10Bv.7: 181	GLYPESHGII DNNMYDVNLNKNFSLSSKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGS	240
161P2F10Bv.1: 275	EVAINGSFPSIYMPYNGSVPFEERISTLLKWL DLPKAERPRFYTMFFEEP DSSGHAGGPV	334
161P2F10Bv.7: 241	EVAINGSFPSIYMPYNGSVPFEERISTLLKWL DLPKAERPRFYTMFFEEP DSSGHAGGPV	300
161P2F10Bv.1: 335	SARVIKALQVVDHAFGMLMEGLKQRNLHNCVNI ILLADHGMDQTYCNKMEYMTDYFPRIN	394
161P2F10Bv.7: 301	SARVIKALQVVDHAFGMLMEGLKQRNLHNCVNI ILLADHGMDQTYCNKMEYMTDYFPRIN	360
161P2F10Bv.1: 395	FFYMYEGPAPRI RAHNIPHDFFSFNSEEIVRNLS CRKPDQHFKPYLTPDL PKRLHYAKNV	454
161P2F10Bv.7: 361	FFYMYEGPAPRVRAHNIPHDFFSFNSEEIVRNLS CRKPDQHFKPYLTPDL PKRLHYAKNV	420
161P2F10Bv.1: 455	RIDKVHLFVDQQWLAVRSKSNTNCGGGNHGYNNEFRSMEAIFLAHGPSFKEKTEVEPFEN	514
161P2F10Bv.7: 421	RIDKVHLFVDQQWLAVRSKSNTNCGGGNHGYNNEFRSMEAIFLAHGPSFKEKTEVEPFEN	480
161P2F10Bv.1: 515	IEVYNLMCDLLRIQPAPNNGTHGSLNHLLKVPFYEPSHAEVSKFSVCGFANPLPTESLD	574
161P2F10Bv.7: 481	IEVYNLMCDLLRIQPAPNNGTHGSLNHLLKVPFYEPSHAEVSKFSVCGFANPLPTESLD	540
161P2F10Bv.1: 575	CFCPHLQNSTQLEQVNQMLNLTQEEITATVKVNLPFGRPRVLQKNVDHCLLYHREYVSGF	634
161P2F10Bv.7: 541	CFCPHLQNSTQLEQVNQMLNLTQEEITATVKVNLPFGRPRVLQKNVDHCLLYHREYVSGF	600
161P2F10Bv.1: 635	GKAMRMPMWSSYTVPQLGDTSPPLPPTVPDC LRADVRVPPSESQKCSFYLA DKNITHGFLY	694
161P2F10Bv.7: 601	GKAMRMPMWSSYTVPQLGDTSPPLPPTVPDC LRADVRVPPSESQKCSFYLA DKNITHGFLY	660
161P2F10Bv.1: 695	PPASNRTSDSQYDALITSNLVPMYEEFRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYN	754
161P2F10Bv.7: 661	PPASNRTSDSQYDALITSNLVPMYEEFRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYN	720
161P2F10Bv.1: 755	YDGHFDAPDEITKHLANTDVPIPTHYFVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTN	814
	YDGHFDAPDEITKHLANTDVPIPTHYFVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTN	

161P2F10Bv.7: 721 YDGHFDAPDEITKHLANTDVPIPTHYFVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTN 780

161P2F10Bv.1: 815 VESCPEGKPEALWVEERFTAHIARVRDVELLTGLDFYQDKVQPVSEILQLKTYLPTFETT 874
 VESCPEGKPEALWVEERFTAHIARVRDVELLTGLDFYQDKVQPVSEILQLKTYLPTFETT

161P2F10Bv.7: 781 VESCPEGKPEALWVEERFTAHIARVRDVELLTGLDFYQDKVQPVSEILQLKTYLPTFETT 840

161P2F10Bv.1: 875 I 875
 I

161P2F10Bv.7: 841 I 841

Table LIX: 161P2F10B Expression in Kidney Cancer

RNA

analysis:

Clear	Papillary	Chromophob	Transitiona	Oncocytoma
cell	e		l	
33/34	16/19	2/3 (67%)	3/7 (42%)	0/3 (0%)
(97%)	(84%)			

Protein

analysis:

Clear	Papillary	Chromophob	Transitiona	Oncocytoma
cell	e		l	
12/12	5/5	1/3 (33%)	0/3 (0%)	0/2 (0%)
(100%)	(100%)			

Table LX: 161P2F10B protein expression in normal tissues

<u>TISSUE</u>	<u>FREQUENCY</u>
Kidney	5/5
Prostate	4/8
Bladder	1/4*
Colon	2/5*
Lung	1/4*
Brain	0/1
Breast	0/2
Heart	0/1
Liver	0/3
Ovary	0/1
Pancreas	0/2
Placenta	0/1
Skin	0/1
Spleen	0/1
Testis	0/4
Thymus	0/1
Uterus	0/1

CLAIMS:

1. A composition comprising:
a substance that a) modulates the status of a protein of Figure 2, or b) a molecule that is modulated by a protein of Figure 2, whereby the status of a cell that expresses a protein of Figure 2 is modulated.
2. A composition of claim 1, further comprising a physiologically acceptable carrier.
3. A pharmaceutical composition that comprises the composition of claim 1 in a human unit dose form.
4. A composition of claim 1 wherein the substance comprises an antibody or fragment thereof that specifically binds to a protein that is related to a protein of Figure 2.
5. An antibody or fragment thereof of claim 4, which is monoclonal.
6. An antibody of claim 4, which is a human antibody, a humanized antibody or a chimeric antibody.
7. A non-human transgenic animal that produces an antibody of claim 4.
8. A hybridoma that produces an antibody of claim 5.
9. A method of delivering a cytotoxic agent or a diagnostic agent to a cell that expresses a protein of Figure 2, said method comprising:
providing the cytotoxic agent or the diagnostic agent conjugated to an antibody or fragment thereof of claim 4; and,
exposing the cell to the antibody-agent or fragment-agent conjugate.
10. A composition of claim 1 wherein the substance comprises a polynucleotide that encodes an antibody or fragment thereof, either of which immunospecifically bind to a protein of Figure 2.
11. A composition of claim 1 wherein the substance comprises a protein related to a protein of Figure 2.
12. A protein of claim 11 that is at least 90% homologous to an entire amino acid sequence shown in Figure 2.
13. A composition of claim 1 wherein the substance comprises:
 - a) a peptide of eight, nine, ten, or eleven contiguous amino acids of a protein of Figure 2;
 - b) a peptide of Tables VIII-XXI;
 - c) a peptide of Tables XXII to XLV; or,
 - d) a peptide of Tables XLVI to XLIX.

14. A composition of claim 1 wherein the substance comprises a CTL polypeptide or an analog thereof, from the amino acid sequence of a protein of Figure 2.

15. A composition of claim 14 further limited by a *proviso* that the epitope is not an entire amino acid sequence of Figure 2.

16. A composition of claim 14 wherein the substance comprises a CTL polypeptide set forth in Tables VIII-XXI.

17. A composition of claim 16 further limited by a *proviso* that the polypeptide is not an entire amino acid sequence of a protein of Figure 2.

18. A composition of claim 1 wherein the substance comprises an antibody polypeptide epitope from an amino acid sequence of Figure 2.

19. A composition of claim 18 further limited by a *proviso* that the epitope is not an entire amino acid sequence of Figure 2.

20. A composition of claim 18 wherein the antibody epitope comprises a peptide region of at least 5 amino acids of Figure 2 in any whole number increment up to the end of said peptide, wherein the epitope comprises an amino acid position selected from:

- a) an amino acid position having a value greater than 0.5 in the Hydrophilicity profile of Figure 5,
- b) an amino acid position having a value less than 0.5 in the Hydrophobicity profile of Figure 6;
- c) an amino acid position having a value greater than 0.5 in the Percent Accessible Residues profile of Figure 7;
- d) an amino acid position having a value greater than 0.5 in the Average Flexibility profile of Figure 8;
- e) an amino acid position having a value greater than 0.5 in the Beta-turn profile of Figure 9;
- f) a combination of at least two of a) through e);
- g) a combination of at least three of a) through e);
- h) a combination of at least four of a) through e); or
- i) a combination of five of a) through e).

21. A composition of claim 20 further limited by a *proviso* that the epitope is not an entire amino acid sequence of Figure 2.

22. A polynucleotide that encodes a protein of claim 11.

23. A polynucleotide of claim 22 that comprises a nucleic acid molecule set forth in Figure 2.

24. A polynucleotide of claim 22 further limited by a *proviso* that the encoded protein is not an entire amino acid sequence of Figure 2.

25. A polynucleotide of claim 22 wherein T is substituted with U.
26. A composition of claim 1 wherein the substance comprises a polynucleotide that comprises a coding sequence of a nucleic acid sequence of Figure 2.
27. A polynucleotide of claim 22 that further comprises an additional nucleotide sequence that encodes an additional protein of claim 11.
28. A composition comprising a polynucleotide that is fully complementary to a polynucleotide of claim 22.
29. A composition comprising a polynucleotide that is fully complementary to a polynucleotide of claim 25.
30. A composition comprising a polynucleotide that is fully complementary to a polynucleotide of claim 27.
31. A composition of claim 1 wherein the substance comprises a) a ribozyme that cleaves a polynucleotide having a 161P2F10B coding sequence, or b) a nucleic acid molecule that encodes the ribozyme; and, a physiologically acceptable carrier.
32. A composition of claim 1 wherein the substance comprises human T cells, wherein said T cells specifically recognize a 161P2F10B peptide subsequence in the context of a particular HLA molecule.
33. A method of inhibiting growth of cancer cells that express a protein of Figure 2, the method comprising:
administering to the cells the composition of claim 1.
34. A method of claim 33 of inhibiting growth of cancer cells that express a protein of Figure 2, the method comprising steps of:
administering to said cells an antibody or fragment thereof, either of which specifically bind to a 161P2F10B-related protein.
35. A method of claim 33 of inhibiting growth of cancer cells that express a protein of Figure 2, the method comprising steps of:
administering to said cells a 161P2F10B-related protein.
36. A method of claim 33 of inhibiting growth of cancer cells that express a protein of Figure 2, the method comprising steps of:
administering to said cells a polynucleotide comprising a coding sequence for a 161P2F10B-related protein or comprising a polynucleotide complementary to a coding sequence for a 161P2F10B-related protein.

37. A method of claim 33 of inhibiting growth of cancer cells that express a protein of Figure 2, the method comprising steps of:

administering to said cells a ribozyme that cleaves a polynucleotide that encodes a protein of Figure 2.

38. A method of claim 33 of inhibiting growth of cancer cells that express a protein of Figure 2 and a particular HLA molecule, the method comprising steps of:

administering human T cells to said cancer cells, wherein said T cells specifically recognize a peptide subsequence of a protein of Figure 2 while the subsequence is in the context of the particular HLA molecule.

39. A method of claim 33, the method comprising steps of:

administering a vector that delivers a nucleotide that encodes a single chain monoclonal antibody, whereby the encoded single chain antibody is expressed intracellularly within cancer cells that express a protein of Figure 2.

40. A method of generating a mammalian immune response directed to a protein of Figure 2, the method comprising:

exposing cells of the mammal's immune system to a portion of

- a) a 161P2F10B-related protein and/or
- b) a nucleotide sequence that encodes said protein,

whereby an immune response is generated to said protein.

41. A method of generating an immune response of claim 40, said method comprising:

providing a 161P2F10B-related protein that comprises at least one T cell or at least one B cell epitope; and,

contacting the epitope with a mammalian immune system T cell or B cell respectively, whereby the T cell or B cell is activated.

42. A method of claim 41 wherein the immune system cell is a B cell, whereby the induced B cell generates antibodies that specifically bind to the 161P2F10B-related protein.

43. A method of claim 41 wherein the immune system cell is a T cell that is a cytotoxic T cell (CTL), whereby the activated CTL kills an autologous cell that expresses the 161P2F10B-related protein.

44. A method of claim 41 wherein the immune system cell is a T cell that is a helper T cell (HTL), whereby the activated HTL secretes cytokines that facilitate the cytotoxic activity of a cytotoxic T cell (CTL) or the antibody-producing activity of a B cell.

45. A method for detecting, in a sample, the presence of a 161P2F10B-related protein or a 161P2F10B-related polynucleotide, comprising steps of:

contacting the sample with a substance of claim 1 that specifically binds to the 161P2F10B-related protein or to the 161P2F10B-related polynucleotide, respectively; and,

determining that there is a complex of the substance with the 161P2F10B-related protein or the substance with the 161P2F10B-related polynucleotide, respectively.

46. A method of claim 45 for detecting the presence of a 161P2F10B-related protein in a sample comprising steps of:

contacting the sample with an antibody or fragment thereof either of which specifically bind to the 161P2F10B-related protein; and,

determining that there is a complex of the antibody or fragment thereof and the 161P2F10B-related protein.

47. A method of claim 45 further comprising a step of:
taking the sample from a patient who has or who is suspected of having cancer.

48. A method of claim 45 for detecting the presence of a protein of Figure 2 mRNA in a sample comprising:

producing cDNA from the sample by reverse transcription using at least one primer;

amplifying the cDNA so produced using 161P2F10B polynucleotides as sense and antisense primers, wherein the 161P2F10B polynucleotides used as the sense and antisense primers serve to amplify a 161P2F10B cDNA; and,

detecting the presence of the amplified 161P2F10B cDNA.

49. A method of claim 45 for monitoring one or more 161P2F10B gene products in a biological sample from a patient who has or who is suspected of having cancer, the method comprising:

determining the status of one or more 161P2F10B gene products expressed by cells in a tissue sample from an individual;

comparing the status so determined to the status of one or more 161P2F10B gene products in a corresponding normal sample; and,

identifying the presence of one or more aberrant gene products of 161P2F10B in the sample relative to the normal sample.

50. The method of claim 49 further comprising a step of determining if there are one or more elevated gene products of a 161P2F10B mRNA or a 161P2F10B protein, whereby the presence of one or more elevated gene products in the test sample relative to the normal tissue sample indicates the presence or status of a cancer.

51. A method of claim 50 wherein the cancer occurs in a tissue set forth in Table I.

52. An antibody of claim 4 which is a monoclonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)15 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____; a monoclonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)29 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____; a monoclonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)37 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No.

_____; a monoclonal antibody or binding region thereof secreted by a hybridoma entitled X41(4)6 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____; a monoclonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)17 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____; or, a monoclonal antibody or binding region thereof secreted by a hybridoma entitled X41(3)50 deposited with the American Type Culture Collection (ATCC; 10801 University Blvd., Manassas, VA 20110-2209 USA) on 07 November 2002 and assigned as Accession No. _____.

1/67

Figure 1: 161P2F10B SSH sequence of 182 nucleotides (SEQ ID NO: 1)

```
1  GATCACACAT TAGGTTATNG ACTTCAATAT TTTCAAATGG TTCAACTTCA GTCTTCTCTT
61  TAAAACTGGG TCCATGTGCC AAGAAAGATA GCCTCCATGC TCCTAAACTC ATTGTTATAA
121 CCATGGTTGC CTCCTCCACA ATTTGTATTT GATTACTCC TAACAGCCAG CCACTGTTGA
181 TC
```

Figure 2:

Figure 2A. The cDNA (SEQ ID NO : 2) and amino acid sequence (SEQ ID NO : 3) of 161P2F10B variant 1. The 3858 nucleotide sequence of 161P2F10B variant 1 is shown. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

1 M E S T L T
1 ctacttttattctgataaaacaggtctatgcagctaccaggacaATGGAATCTACGTTGAC
7 L A T E Q P V K K N T L K K Y K I A C I
61 TTTAGCAACGGAACAACCTGTTAAGAAGAACACTCTTAAGAAATATAAAATAGCTTGCAT
27 V L L A L L V I M S L G L G L G L G L R
121 TGTTCTTCTTGCTTTGCTGGTGATCATGTCACTTGGATTAGGCCTGGGGCTTGGACTCAG
47 K L E K Q G S C R K K C F D A S F R G L
181 GAAACTGGAAAAGCAAGGCAGCTGCAGGAAGAAGTGCTTTGATGCATCATTTAGAGGACT
67 E N C R C D V A C K D R G D C C W D F E
241 GGAGAACTGCCGGTGTGATGTGGCATGTAAAGACCGAGGTGATTGCTGCTGGGATTTTGA
87 D T C V E S T R I W M C N K F R C G E T
301 AGACACCTGTGTGGAATCAACTCGAATATGGATGTGCAATAAATTTTCGTTGTGGAGAGAC
107 R L E A S L C S C S D D C L Q K K D C C
361 CAGATTAGAGGCCAGCCTTTGCTCTTGTTTCAGATGACTGTTTGCAGAAGAAAGATTGCTG
127 A D Y K S V C Q G E T S W L E E N C D T
421 TGCTGACTATAAGAGTGTTTGCCAAGGAGAAACCTCATGGCTGGAAGAAACTGTGACAC
147 A Q Q S Q C P E G F D L P P V I L F S M
481 AGCCCAGCAGTCTCAGTGCCCAGAAGGGTTTGACCTGCCACCAGTTATCTTGTTTTCTAT
167 D G F R A E Y L Y T W D T L M P N I N K
541 GGATGGATTTAGAGCTGAATATTTATACACATGGGATACTTTAATGCCAAATATCAATAA
187 L K T C G I H S K Y M R A M Y P T K T F
601 ACTGAAAACATGTGGAATTCATTCAAATACATGAGAGCTATGTATCCTACCAAAACCTT
207 P N H Y T I V T G L Y P E S H G I I D N
661 CCCAAATCATTACACCATTTGTCACGGGCTTGTATCCAGAGTCACATGGCATCATTGACAA
227 N M Y D V N L N K N F S L S S K E Q N N
721 TAATATGTATGATGTAAATCTCAACAAGAATTTTTCACTTTCTTCAAAGGAACAAAATAA
247 P A W W H G Q P M W L T A M Y Q G L K A
781 TCCAGCCTGGTGGCATGGGCAACCAATGTGGCTGACAGCAATGTATCAAGGTTTAAAAGC
267 A T Y F W P G S E V A I N G S F P S I Y
841 CGCTACCTACTTTTGGCCCGGATCAGAAGTGGCTATAAATGGCTCCTTTCCTTCCATATA
287 M P Y N G S V P F E E R I S T L L K W L
901 CATGCCTTACAACGGAAGTGTCCCATTTGAAGAGAGGATTTCTACACTGTAAATGGCT
307 D L P K A E R P R F Y T M Y F E E P D S

Figure 2A-1

3/67

961 GGACCTGCCCAAAGCTGAAAGACCCAGGTTTTATACCATGTATTTTGAAGAACCTGATTC
327 S G H A G G P V S A R V I K A L Q V V D
1021 CTCTGGACATGCAGGTGGACCAGTCAGTGCCAGAGTAATTAAAGCCTTACAGGTAGTAGA
347 H A F G M L M E G L K Q R N L H N C V N
1081 TCATGCTTTTGGGATGTTGATGGAAGGCCTGAAGCAGCGGAATTTGCACAACCTGTGTCAA
367 I I L L A D H G M D Q T Y C N K M E Y M
1141 TATCATCCTTCTGGCTGACCATGGAATGGACCAGACTTATTGTAACAAGATGGAATACAT
387 T D Y F P R I N F F Y M Y E G P A P R I
1201 GACTGATTATTTTCCCAGAATAAACTTCTTCTACATGTACGAAGGGCCTGCCCCCGCAT
407 R A H N I P H D F F S F N S E E I V R N
1261 CCGAGCTCATAATATACCTCATGACTTTTTTAGTTTTTAATTCTGAGGAAATTGTTAGAAA
427 L S C R K P D Q H F K P Y L T P D L P K
1321 CCTCAGTTGCCGAAAACCTGATCAGCATTTCAAGCCCTATTTGACTCCTGATTTGCCAAA
447 R L H Y A K N V R I D K V H L F V D Q Q
1381 GCGACTGCACTATGCCAAGAACGTCAGAATCGACAAAGTTCATCTCTTTGTGGATCAACA
467 W L A V R S K S N T N C G G G N H G Y N
1441 GTGGCTGGCTGTTAGGAGTAAATCAAATACAAATTGTGGAGGAGGCAACCATGGTTATAA
487 N E F R S M E A I F L A H G P S F K E K
1501 CAATGAGTTTAGGAGCATGGAGGCTATCTTTCTGGCACATGGACCCAGTTTTAAAGAGAA
507 T E V E P F E N I E V Y N L M C D L L R
1561 GACTGAAGTTGAACCATTTGAAAATATTGAAGTCTATAACCTAATGTGTGATCTTCTACG
527 I Q P A P N N G T H G S L N H L L K V P
1621 CATTCAACCAGCACCAAACAATGGAACCCATGGTAGTTTAAACCATCTTCTGAAGGTGCC
547 F Y E P S H A E E V S K F S V C G F A N
1681 TTTTATGAGCCATCCCATGCAGAGGAGGTGTCAAAGTTTTCTGTTTGTGGCTTTGCTAA
567 P L P T E S L D C F C P H L Q N S T Q L
1741 TCCATTGCCCACAGAGTCTCTTGACTGTTTCTGCCCTCACCTACAAAATAGTACTCAGCT
587 E Q V N Q M L N L T Q E E I T A T V K V
1801 GGAACAAGTGAATCAGATGCTAAATCTCACCCAAGAAGAAATAACAGCAACAGTGAAAGT
607 N L P F G R P R V L Q K N V D H C L L Y
1861 AAATTTGCCATTTGGGAGGCCTAGGGTACTGCAGAAGAACGTGGACCACTGTCTCCTTTA
627 H R E Y V S G F G K A M R M P M W S S Y
1921 CCACAGGGAATATGTCAGTGGATTTGGAAAAGCTATGAGGATGCCCATGTGGAGTTCATA
647 T V P Q L G D T S P L P P T V P D C L R
1981 CACAGTCCCCCAGTTGGGAGACACATCGCCTCTGCCTCCCACTGTCCCAGACTGTCTGCG
667 A D V R V P P S E S Q K C S F Y L A D K
2041 GGCTGATGTCAGGGTTCCTCCTTCTGAGAGCCAAAAATGTTCTTCTATTTAGCAGACAA
687 N I T H G F L Y P P A S N R T S D S Q Y

Figure 2A-2
SUBSTITUTE SHEET (RULE 26)

4/67

2101 GAATATCACCCACGGCTTCCTCTATCCTCCTGCCAGCAATAGAACATCAGATAGCCAATA
707 D A L I T S N L V P M Y E E F R K M W D
2161 TGATGCTTTAATTACTAGCAATTTGGTACCTATGTATGAAGAATTCAGAAAAATGTGGGA
727 Y F H S V L L I K H A T E R N G V N V V
2221 CTACTTCCACAGTGTTCCTTCTTATAAAACATGCCACAGAAAGAAATGGAGTAAATGTGGT
747 S G P I F D Y N Y D G H F D A P D E I T
2281 TAGTGGACCAATATTTGATTATAATTATGATGGCCATTTTGATGCTCCAGATGAAATTAC
767 K H L A N T D V P I P T H Y F V V L T S
2341 CAAACATTTAGCCAACACTGATGTTCCCATCCCAACACACTACTTTGTGGTGCTGACCAG
787 C K N K S H T P E N C P G W L D V L P F
2401 TTGTAAAAACAAGAGCCACACACCGGAAACTGCCCTGGGTGGCTGGATGTCCTACCCTT
807 I I P H R P T N V E S C P E G K P E A L
2461 TATCATCCCTCACCGACCTACCAACGTGGAGAGCTGTCCTGAAGGTAAACCAGAAGCTCT
827 W V E E R F T A H I A R V R D V E L L T
2521 TTGGGTTGAAGAAAGATTTACAGCTCACATTGCCCGGGTCCGTGATGTAGAACTTCTCAC
847 G L D F Y Q D K V Q P V S E I L Q L K T
2581 TGGGCTTGACTTCTATCAGGATAAAGTGCAGCCTGTCTCTGAAATTTTGCAACTAAAGAC
867 Y L P T F E T T I *
2641 ATATTTACCAACATTTGAAACCACTATTTAAActtaataatgtctacttaatatataattt
2701 actgtataaagtaatttttgcaaaatataagtgattttttctggagaattgtaaaataaa
2761 gttttctatttttccttaaaaaaaaaaccggaattccgggcttgggaggctgaggcagga
2821 gactcgcttgaacccgggaggcagaggttgcagtgagccaagattgcgccattgcactcc
2881 agagcctgggtgacagagcaagactacatctcaaaaaataaataaataaaaagtaa
2941 caataaaaataaaaagaacagcagagagaatgagcaaggagaaatgtcacaactattgc
3001 aaaataactgttacactgggttggctctccaagaagatactggaatctcttcagccatttg
3061 cttttcagaagtagaaaccagcaaaccacctctaagcggagaacatacgattctttatta
3121 agtagctctggggaaggaaagaataaaaagttgatagctccctgattgggaaaaaatgcac
3181 aattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaaaaattca
3241 caaacgttgggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaaa
3301 ttttcagttgggatgcgactgacaaaaagaacgggatttccaggcataaagttggcgtga
3361 gctacagagggcaccatgtggctcagtggaagacccttcaagattcaaagttccatttga
3421 cagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaagccaagtgg
3481 aaagcgagcaatttgcagcataactgcttctcctagacagggtgagtgggcaaaatagc
3541 acagtacacacagtgactattagccactgccagaaacaggctgaacagccctgggagaca
3601 agggaaggcaggtgggtgggagttgttcatggagagaaaggagagttttagaaccagcaca
3661 tccactggagatgctgggccaccagacccctcccagtcaataaagtctgggtgcctcattt
3721 gatctcagcctcatcatgaccctggagagaccctgataccatctgccagtccccgacagc

Figure 2A-3

SUBSTITUTE SHEET (RULE 26)

5/67

3781 ttaggcactccttgccatcaacctgaccccccgagtgggttctccaggctccctgccccac
 3841 ccattcaggccggaattc

Figure 2A-4

Figure 2B: The cDNA (SEQ ID NO: 4) and amino acid sequence (SEQ ID NO: 5) of 161P2F10B variant 2. The 3858 nucleotide sequence of 161P2F10B variant 2 is shown. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

1	M E S T L T
1	ctactttattctgataaaacaggtctatgcagctaccaggacaATGGAATCTACGTTGAC
7	L A T E Q P V K K N T L K K Y K I A C I
61	TTTAGCAACGGAACAACCTGTTAAGAAGAACAACCTCTTAAGAAATATAAAATAGCTTGCAT
27	V L L A L L V I M S L G L G L G L G L R
121	TGTTCTTCTTGCTTTGCTGGTGATCATGTCACTTGGATTAGGCCTGGGGCTTGGACTCAG
47	K L E K Q G S C R K K C F D A S F R G L
181	GAAACTGGAAAAGCAAGGCAGCTGCAGGAAGAAGTGCTTTGATGCATCATTTAGAGGACT
67	E N C R C D V A C K D R G D C C W D F E
241	GGAGAACTGCCGGTGTGATGTGGCATGTAAAGACCGAGGTGATTGCTGCTGGGATTTTGA
87	D T C V E S T R I W M C N K F R C G E T
301	AGACACCTGTGTGGAATCAACTCGAATATGGATGTGCAATAAATTTTCGTTGTGGAGAGAC
107	R L E A S L C S C S D D C L Q R K D C C
361	CAGATTAGAGGCCAGCCTTTGCTCTTGTTTCAGATGACTGTTTGCAGAGGAAAGATTGCTG
127	A D Y K S V C Q G E T S W L E E N C D T
421	TGCTGACTATAAGAGTGTTTGCCAAGGAGAAACCTCATGGCTGGAAGAAACTGTGACAC
147	A Q Q S Q C P E G F D L P P V I L F S M
481	AGCCCAGCAGTCTCAGTGCCCAGAAGGGTTTGACCTGCCACCAGTTATCTTGTTTTCTAT
167	D G F R A E Y L Y T W D T L M P N I N K
541	GGATGGATTTAGAGCTGAATATTTATACACATGGGATACTTTAATGCCAAATATCAATAA
187	L K T C G I H S K Y M R A M Y P T K T F
601	ACTGAAAACATGTGGAATTCATTCAAATAACATGAGAGCTATGTATCCTACCAAACCTT
207	P N H Y T I V T G L Y P E S H G I I D N
661	CCCAAATCATTACACCATTTGTCACGGGCTTGTATCCAGAGTCACATGGCATCATTGACAA
227	N M Y D V N L N K N F S L S S K E Q N N
721	TAATATGTATGATGTAAATCTCAACAAGAATTTTCACTTTCTTCAAAGGAACAAAATAA
247	P A W W H G Q P M W L T A M Y Q G L K A
781	TCCAGCCTGGTGGCATGGGCAACCAATGTGGCTGACAGCAATGTATCAAGGTTTAAAGC
267	A T Y F W P G S E V A I N G S F P S I Y
841	CGCTACCTACTTTTGGCCCGGATCAGAAGTGGCTATAAATGGCTCCTTTCCTTCCATATA
287	M P Y N G S V P F E E R I S T L L K W L
901	CATGCCTTACAACGGAAGTGTCCCATTTGAAGAGAGGATTTCTACACTGTTAAATGGCT
307	D L P K A E R P R F Y T M Y F E E P D S
961	GGACCTGCCCAAAGCTGAAAGACCCAGGTTTTATACCATGTATTTTGAAGAACCTGATTC

Figure 2B-1

6/67

327 S G H A G G P V S A R V I K A L Q V V D
1021 CTCTGGACATGCAGGTGGACCAGTCAGTGCCAGAGTAATTAAAGCCTTACAGGTAGTAGA
347 H A F G M L M E G L K Q R N L H N C V N
1081 TCATGCTTTTGGGATGTTGATGGAAGGCCTGAAGCAGCGGAATTTGCACAACCTGTGTCAA
367 I I L L A D H G M D Q T Y C N K M E Y M
1141 TATCATCCTTCTGGCTGACCATGGAATGGACCAGACTTATTGTAACAAGATGGAATACAT
387 T D Y F P R I N F F Y M Y E G P A P R I
1201 GACTGATTATTTTCCCAGAATAAACTTCTTCTACATGTACGAAGGGCCTGCCCCCGCAT
407 R A H N I P H D F F S F N S E E I V R N
1261 CCGAGCTCATAATATACCTCATGACTTTTTTTAGTTTAAATTCTGAGGAAATTGTTAGAAA
427 L S C R K P D Q H F K P Y L T P D L P K
1321 CCTCAGTTGCCGAAAACCTGATCAGCATTTCAAGCCCTATTTGACTCCTGATTTGCCAAA
447 R L H Y A K N V R I D K V H L F V D Q Q
1381 GCGACTGCACTATGCCAAGAACGTCAGAATCGACAAAGTTCATCTCTTTGTGGATCAACA
467 W L A V R S K S N T N C G G G N H G Y N
1441 GTGGCTGGCTGTTAGGAGTAAATCAAATACAAATTGTGGAGGAGGCAACCATGGTTATAA
487 N E F R S M E A I F L A H G P S F K E K
1501 CAATGAGTTTAGGAGCATGGAGGCTATCTTTCTGGCACATGGACCCAGTTTAAAGAGAA
507 T E V E P F E N I E V Y N L M C D L L R
1561 GACTGAAGTTGAACCATTTGAAAATATTGAAGTCTATAACCTAATGTGTGATCTTCTACG
527 I Q P A P N N G T H G S L N H L L K V P
1621 CATTCAACCAGCACCAACAATGGAACCCATGGTAGTTTAAACCATCTTCTGAAGGTGCC
547 F Y E P S H A E E V S K F S V C G F A N
1681 TTTTATGAGCCATCCCATGCAGAGGAGGTGTCAAAGTTTTCTGTTTGTGGCTTTGCTAA
567 P L P T E S L D C F C P H L Q N S T Q L
1741 TCCATTGCCCACAGAGTCTCTTGACTGTTTCTGCCCTCACCTACAAAATAGTACTCAGCT
587 E Q V N Q M L N L T Q E E I T A T V K V
1801 GGAACAAGTGAATCAGATGCTAAATCTCACCCAAGAAGAAATAACAGCAACAGTGAAAGT
607 N L P F G R P R V L Q K N V D H C L L Y
1861 AAATTTGCCATTTGGGAGGCCTAGGGTACTGCAGAAGAACGTGGACCACTGTCTCCTTTA
627 H R E Y V S G F G K A M R M P M W S S Y
1921 CCACAGGGAATATGTCAGTGGATTTGGAAAAGCTATGAGGATGCCCATGTGGAGTTCATA
647 T V P Q L G D T S P L P P T V P D C L R
1981 CACAGTCCCCCAGTTGGGAGACACATCGCCTCTGCCTCCCACTGTCCCAGACTGTCTGCG
667 A D V R V P P S E S Q K C S F Y L A D K
2041 GGCTGATGTCAGGGTTCCTCCTTCTGAGAGCCAAAATGTTCCCTTCTATTTAGCAGACAA
697 N I T H G F L Y P P A S N R T S D S Q Y
2101 GAATATCACCCACGGCTTCCTCTATCCTCCTGCCAGCAATAGAACATCAGATAGCCAATA
707 D A L I T S N L V P M Y E E F R K M W D
2161 TGATGCTTTAATTACTAGCAATTTGGTACCTATGTATGAAGAATTCAGAAAAATGTGGGA
727 Y F H S V L L I K H A T E R N G V N V V
2221 CTA CTTCACAGTGTTCTTCTTATAAAACATGCCACAGAAAGAAATGGAGTAAATGTGGT

Figure 2B-2
SUBSTITUTE SHEET (RULE 26)

7/67

```

747 S G P I F D Y N Y D G H F D A P D E I T
2281 TAGTGGACCAATATTTGATTATAATTATGATGGCCATTTTGATGCTCCAGATGAAATTAC
767 K H L A N T D V P I P T H Y F V V L T S
2341 CAAACATTTAGCCAACACTGATGTTCCCATCCCAACACACTACTTTGTGGTGCTGACCAG
787 C K N K S H T P E N C P G W L D V L P F
2401 TTGTAAAAACAAGAGCCACACACCGGAAAACCTGCCCTGGGTGGCTGGATGTCCTACCCTT
807 I I P H R P T N V E S C P E G K P E A L
2461 TATCATCCCTCACCGACCTACCAACGTGGAGAGCTGTCCTGAAGGTAAACCAGAAGCTCT
827 W V E E R F T A H I A R V R D V E L L T
2521 TTGGGTTGAAGAAAGATTTACAGCTCACATTGCCCGGGTCCGTGATGTAGAACTTCTCAC
847 G L D F Y Q D K V Q P V S E I L Q L K T
2581 TGGGCTTGACTTCTATCAGGATAAAGTGCAGCCTGTCTCTGAAATTTTGCAACTAAAGAC
867 Y L P T F E T T I *
2641 ATATTTACCAACATTTGAAACCACTATTTAACTtaataatgtctacttaatatataattt
2701 actgtataaagtaatttttgcaaaatataagtgattttttctggagaattgtaaaataaa
2761 gtttttctatttttcttaaaaaaaaaaacgggaattccgggcttgggaggctgaggcagga
2821 gactcgcttgaacccgggaggcagaggttgcagtgagccaagattgcgccattgcactcc
2881 agagcctgggtgacagagcaagactacatctcaaaaaataaataaaaaataaaaagtaa
2941 caataaaaaataaaaagaacagcagagagaatgagcaaggagaaatgtcacaaactattgc
3001 aaaatactgttacactgggttggctctccaagaagatactggaatctcttcagccatttg
3061 cttttcagaagtagaaaccagcaaaccacctctaagcgggagaacatacgattctttatta
3121 agtagctctggggaaggaaagaataaaaagttgatagctccctgattgggaaaaaatgcac
3181 aattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaaaaattca
3241 caaacgttgggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaaa
3301 ttttcagttgggatgagactgacaaaaagaacgggatttccaggcataaagttggcgtga
3361 gctacagagggcaccatgttggtcagtggaagacccttcaagattcaaagttccatttga
3421 cagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaagccaagtgggt
3481 aaagcgagcaatttgcagcataactgcttctcctagacagggtgagtgggcaaaatacg
3541 acagtacacacagtgactattagccactgccagaaacaggctgaacagccctgggagaca
3601 aggggaaggcaggtgggtgggagttgttcatggagagaaaggagagttttagaaccagcaca
3661 tccactggagatgctgggccaccagacccctcccagtcataaaagtctgggtgcctcattt
3721 gatctcagcctcatcatgaccctggagagaccctgataccatctgccagtccccgacagc
3781 ttaggcactccttgccatcaacctgaccccccgagtgggttctccaggctccctgccccac
3841 ccattcaggccggaattc

```

Figure 2B-3

Figure 2C: The cDNA (SEQ ID NO: 6) and amino acid sequence (SEQ ID NO: 7) of 161P2F10B variant 3. The 3858 nucleotide sequence of 161P2F10B variant 3 is shown. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

```

1 M E S T L T
1 ctactttatttctgataaaacaggtctatgcagctaccaggacaATGGAATCTACGTTGAC
7 L A T E Q P V K K N T L K K Y K I A C I

```

Figure 2C-1

8/67

61 TTTAGCAACGGAACAACCTGTTAAGAAGAACTCTTAAGAAATATAAAATAGCTTGCAT
27 V L L A L L V I M S L G L G L G L G L R
121 TGTTCCTTCTTGCTTTGCTGGTGATCATGTCACCTGGATTAGGCCTGGGGCTTGGACTCAG
47 K L E K Q G S C R K K C F D A S F R G L
181 GAAACTGGAAAAGCAAGGCAGCTGCAGGAAGAAGTGCTTTGATGCATCATTTAGAGGACT
67 E N C R C D V A C K D R G D C C W D F E
241 GGAGAACTGCCGGTGTGATGTGGCATGTAAAGACCGAGGTGATTGCTGCTGGGATTTTGA
87 D T C V E S T R I W M C N K F R C G E T
301 AGACACCTGTGTGGAATCAACTCGAATATGGATGTGCAATAAATTTTCGTTGTGGAGAGAC
107 R L E A S L C S C S D D C L Q K K D C C
361 CAGATTAGAGGCCAGCCTTTGCTCTTGTTTCAGATGACTGTTTGCAGAAGAAAGATTGCTG
127 A D Y K S V C Q G E T S W L E E N C D T
421 TGCTGACTATAAGAGTGTTTGCCAAGGAGAAACCTCATGGCTGGAAGAAAACCTGTGACAC
147 A Q Q S Q C P E G F D L P P V I L F S M
481 AGCCCAGCAGTCTCAGTGCCCAAGGGTTTGACCTGCCACCAGTTATCTTGTTTTCTAT
167 D G F R A E Y L Y T W D T L M P N I N K
541 GGATGGATTTAGAGCTGAATATTTATACACATGGGATACTTTAATGCCAAATATCAATAA
187 L K T C G I H S K Y M R A M Y P T K T F
601 ACTGAAAACATGTGGAATTCATTCAAATACATGAGAGCTATGTATCCTACCAAAACCTT
207 P N H Y T I V T G L Y P E S H G I I D N
661 CCCAAATCATTACACCATTGTCACGGGCTTGTATCCAGAGTCACATGGCATCATTGACAA
227 N M Y D V N L N K N F S L S S K E Q N N
721 TAATATGTATGATGTAAATCTCAACAAGAATTTTTTCACTTTTCTTCAAAGGAACAAAATAA
247 P A W W H G Q P M W L T A M Y Q G L K A
781 TCCAGCCTGGTGGCATGGGCAACCAATGTGGCTGACAGCAATGTATCAAGGTTTAAAGC
267 A T Y F W P G S E V A I N G S F P S I Y
841 CGCTACCTACTTTTGGCCCGGATCAGAAGTGGCTATAAATGGCTCCTTTTCTTCCATATA
287 M P Y N G S V P F E E R I S T L L K W L
901 CATGCCTTACAACGGAAGTGTCCCATTTGAAGAGAGGATTTCTACACTGTAAATGGCT
307 D L P K A E R P R F Y T M Y F E E P D S
961 GGACCTGCCCAAAGCTGAAAGACCCAGGTTTTTATACCATGTATTTTGAAGAACCTGATTC
327 S G H A G G P V S A R V I K A L Q V V D
1021 CTCTGGACATGCAGGTGGACCAGTCAGTGCCAGAGTAATTAAAGCCTTACAGGTAGTAGA
347 H A F G M L M E G L K Q R N L H N C V N
1081 TCATGCTTTTGGGATGTTGATGGAAGGCCTGAAGCAGCGGAATTTGCACAACCTGTGTCAA
367 I I L L A D H G M D Q T Y C N K M E Y M
1141 TATCATCCTTCTGGCTGACCATGGAATGGACCAGACTTATTGTAACAAGATGGAATACAT
387 T D Y F P R I N F F Y M Y E G P A P R I
1201 GACTGATTATTTTCCCAGAATAAACTTCTTCTACATGTACGAAGGGCCTGCCCCCGCAT
407 R A H N I P H D F F S F N S E E I V R N
1261 CCGAGCTCATAATATACCTCATGACTTTTTTGTAGTTTTAATTCTGAGGAAATTGTTAGAAA
427 L S C R K P D Q H F K P Y L T P D L P K
1321 CCTCAGTTGCCGAAAACCTGATCAGCATTTCAAGCCCTATTTGACTCCTGATTGCCCCAAA
447 R L H Y A K N V R I D K V H L F V D Q Q
1381 GCGACTGCACTATGCCAAGAACGTCAGAATCGACAAAGTTCATCTCTTTGTGGATCAACA
467 W L A V R S K S N T N C G G G N H G Y N
1441 GTGGCTGGCTGTTAGGAGTAAATCAAATACAAATTGTGGAGGAGGCAACCATGGTTATAA
487 N E F R S M E A I F L A H G P S F K E K
1501 CAATGAGTTTAGGAGCATGGAGGCTATCTTTCTGGCACATGGACCCAGTTTTTAAAGAGAA
507 T E V E P F E N I E V Y N L M C D L L R
1561 GACTGAAGTTGAACCATTTGAAAATATTGAAGTCTATAACCTAATGTGTGATCTTCTACG
527 I Q P A P N N G T H G S L N H L L K V P
1621 CATTCAACCAGCACCAACAATGGAACCCATGGTAGTTTAAACCATCTTCTGAAGGTGCC
547 F Y E P S H A E E V S K F S V C G F A N
1681 TTTTATGAGCCATCCCATGCAGAGGAGGTGTCAAAGTTTTCTGTTTGTGGCTTTGCTAA
567 P L P T E S L D C F C P H L Q N S T Q L
1741 TCCATTGCCACAGAGTCTCTTGACTGTTTCTGCCCTCACCTACAAAATAGTACTCAGCT

Figure 2C-2

9/67

```

587  E Q V N Q M L N L T Q E E I T A T V K V
1801 GGAACAAGTGAATCAGATGCTAAATCTCACCCAAGAAGAAATAACAGCAACAGTGAAAGT
607  N L P F G R P R V L Q K N V D H C L L Y
1861 AAATTTGCCATTTGGGAGGCCTAGGGTACTGCAGAAGAACGTGGACCACTGTCTCCTTTA
627  H R E Y V S G F G K A M R M P M W S S Y
1921 CCACAGGGAATATGTCAGTGGATTTGGAAAAGCTATGAGGATGCCCATGTGGAGTTCATA
647  T V P Q L G D T S P L P P T V P D C L R
1981 CACAGTCCCCCAGTTGGGAGACACATCGCCTCTGCCTCCCAGTGTCCCAGACTGTCTGCG
667  A D V R V P P S E S Q K C S F Y L A D K
2041 GGCTGATGTCAGGGTTCCTCCTTCTGAGAGCCAAAAATGTTCTTCTATTTAGCAGACAA
687  N I T H G F L Y P P A S N R T S D S Q Y
2101 GAATATCACCCACGGCTTCCTCTATCCTCCTGCCAGCAATAGAACATCAGATAGCCAATA
707  D A L I T S N L V P M Y E E F R K M W D
2161 TGATGCTTTAATTACTAGCAATTTGGTACCTATGTATGAAGAATTCAGAAAAATGTGGGA
727  Y F H S V L L I K H A T E R N G V N V V
2221 CTACTTCCACAGTGTCTTCTTATAAAACATGCCACAGAAAGAAATGGAGTAAATGTGGT
747  S G P I F D Y N Y D G H F D A P D E I T
2281 TAGTGGACCAATATTTGATTATAATTATGATGGCCATTTTGATGCTCCAGATGAAATTAC
767  K H L A N T D V P I P T H Y F V V L T S
2341 CAAACATTTAGCCAACACTGATGTTCCCATCCCAACACACTACTTTGTGGTGCTGACCAG
787  C K N K S H T P E N C P G W L D V L P F
2401 TTGTAAAAACAAGAGCCACACACCGGAAACTGCCCTGGGTGGCTGGATGTCCTACCCTT
807  I I P H R P T N V E S C P G G K P E A L
2461 TATCATCCCTCACCGACCTACCAACGTGGAGAGCTGTCCTGGAGGTAAACCAGAAGCTCT
827  W V E E R F T A H I A R V R D V E L L T
2521 TTGGGTTGAAGAAAGATTTACAGCTCACATTGCCCGGGTCCGTGATGTAGAACTTCTCAC
847  G L D F Y Q D K V Q P V S E I L Q L K T
2581 TGGGCTTGACTTCTATCAGGATAAAGTGCAGCCTGTCTCTGAAATTTTGCAACTAAAGAC
867  Y L P T F E T T I *
2641 ATATTTACCAACATTTGAAACCCTATTTAAActtaataatgtctacttaatatataat
2701 actgtataaagtaattttggcaaaatataagtgatttttttctggagaattgtaaaataaa
2761 gttttctatttttccttaaaaaaaaaaacgggaattccgggcttgggaggctgaggcagga
2821 gactcgcttgaacccgggaggcagaggttgcagtgagccaagattgcgccattgcactcc
2881 agagcctgggtgacagagcaagactacatctcaaaaaataaataaataaataaagtaa
2941 caataaaaaataaaaaagaacagcagagagaatgagcaaggagaaatgtcacaactattgc
3001 aaaatactgttacactgggttggctctccaagaagatactggaatctcttcagccatttg
3061 cttttcagaagtagaaaccagcaaacacctctaagcggagaacatacagattctttatta
3121 agtagctctggggaaggaaagaataaaagttgatagctccctgattgggaaaaaatgcac
3181 aattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaaaaattca
3241 caaacgttggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaaa
3301 ttttcagttgggatgagactgacaaaaagaacgggatttccaggcataaagttggcgtga
3361 gctacagagggcaccatgtggctcagtggaagacccttcaagattcaaagttccatttga
3421 cagagcaaaggcacttcgcaaggagaagggttttaaattatgggtccaaaagccaagtgg
3481 aaagcgagcaatttgcagcataaactgcttctcctagacagggtgagtgaggcaaaatacg
3541 acagtacacacagtgactattagccactgccagaaacaggctgaacagccctgggagaca
3601 aggggaaggcaggtggtgggagttgttcatggagagaaaggagagttttagaaccagcaca
3661 tccactggagatgctgggccaccagaccctcccagtcataaagtcgtggtgcctcattt
3721 gatctcagcctcatcatgaccctggagagaccctgataccatctgccagtccccgacagc
3781 ttaggcactccttgccatcaacctgaccccccgagtggttctccaggctccctgccccac
3841 ccattcaggccggaattc

```

Figure 2C-3

Figure 2D: The cDNA (SEQ ID NO: 8) and amino acid sequence (SEQ ID NO: 9) of 161P2F10B variant 4. The 3858 nucleotide sequence of 161P2F10B variant 4 is shown. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

10/67

1 M E S T L T
1 ctactttattctgataaaacaggtctatgcagctaccaggacaATGGAATCTACGTTGAC
7 L A T E Q P V K K N T L K K Y K I A C I
61 TTTAGCAACGGAACAACCTGTTAAGAAGAACAACCTCTTAAGAAATATAAAATAGCTTGCAT
27 V L L A L L V I M S L G L G L G L G L R
121 TGTTCTTCTTGCTTTGCTGGTGATCATGTCACTTGGATTAGGCCTGGGGCTTGGACTCAG
47 K L E K Q G S C R K K C F D A S F R G L
181 GAAACTGGAAAAGCAAGGCAGCTGCAGGAAGAAGTGCTTTGATGCATCATTTAGAGGACT
67 E N C R C D V A C K D R G D C C W D F E
241 GGAGAACTGCCGGTGTGATGTGGCATGTAAAGACCGAGGTGATTGCTGCTGGGATTTTGA
87 D T C V E S T R I W M C N K F R C G E T
301 AGACACCTGTGTGGAATCAACTCGAATATGGATGTGCAATAAATTTTCGTTGTGGAGAGAC
107 R L E A S L C S C S D D C L Q K K D C C
361 CAGATTAGAGGCCAGCCTTTGCTCTTGTTTCAGATGACTGTTTGCAGAAGAAAGATTGCTG
127 A D Y K S V C Q G E T S W L E E N C D T
421 TGCTGACTATAAGAGTGTTTGCCAAGGAGAAACCTCATGGCTGGAAGAAAACCTGTGACAC
147 A Q Q S Q C P E G F D L P P V I L F S M
481 AGCCCAGCAGTCTCAGTGCCCAGAAGGGTTTGACCTGCCACCAGTTATCTTGTTTTCTAT
167 D G F R A E Y L Y T W D T L M P N I N K
541 GGATGGATTTAGAGCTGAATATTTATACACATGGGATACTTTAATGCCAAATATCAATAA
187 L K T C G I H S K Y M R A M Y P T K T F
601 ACTGAAAACATGTGGAATTCATTCAAATAACATGAGAGCTATGTATCCTACCAAACCTT
207 P N H Y T I V T G L Y P E S H G I I D N
661 CCCAAATCATTACACCATTGTCACGGGCTTGTATCCAGAGTCACATGGCATCATTGACAA
227 N M Y D V N L N K N F S L S S K E Q N N
721 TAATATGTATGATGTAAATCTCAACAAGAATTTTTCACTTTCTTCAAAGGAACAAAATAA
247 P A W W H G Q P M W L T A M Y Q G L K A
781 TCCAGCCTGGTGGCATGGGCAACCAATGTGGCTGACAGCAATGTATCAAGGTTTAAAAGC
267 A T Y F W P G S E V A I N G S F P S I Y
841 CGCTACCTACTTTTGGCCCGGATCAGAAGTGGCTATAAATGGCTCCTTTCCTTCCATATA
287 M P Y N G S V P F E E R I S T L L K W L
901 CATGCCTTACAACGGAAGTGTCCCATTTGAAGAGAGGATTTCTACACTGTTAAATGGCT
307 D L P K A E R P R F Y T M Y F E E P D S
961 GGACCTGCCCAAAGCTGAAAGACCCAGGTTTTATACCATGTATTTTGAAGAACCTGATTC
327 S G H A G G P V S A R V I K A L Q V V D
1021 CTCTGGACATGCAGGTGGACCAGTCAGTGCCAGAGTAATTAAAGCCTTACAGGTAGTAGA
347 H A F G M L M E G L K Q R N L H N C V N
1081 TCATGCTTTTGGGATGTTGATGGAAGGCCTGAAGCAGCGGAATTTGCACAACTGTGTCAA

Figure 2D-1

SUBSTITUTE SHEET (RULE 26)

11/67

367 I I L L A D H G M D Q T Y C N K M E Y M
1141 TATCATCCTTCTGGCTGACCATGGAATGGACCAGACTTATTGTAACAAGATGGAATACAT
387 T D Y F P R I N F F Y M Y E G P A P R I
1201 GACTGATTATTTTCCCAGAATAAACTTCTTCTACATGTACGAAGGGCCTGCCCCCGCAT
407 R A H N I P H D F F S F N S E E I V R N
1261 CCGAGCTCATAATATACCTCATGACTTTTTTAGTTTTAATTCTGAGGAAATTGTTAGAAA
427 L S C R K P D Q H F K P Y L T P D L P K
1321 CCTCAGTTGCCGAAAACCTGATCAGCATTTCAAGCCCTATTTGACTCCTGATTTGCCAAA
447 R L H Y A K N V R I D K V H L F V D Q Q
1381 GCGACTGCACTATGCCAAGAACGTCAGAATCGACAAAGTTCATCTCTTTGTGGATCAACA
467 W L A V R S K S N T N C G G G N H G Y N
1441 GTGGCTGGCTGTTAGGAGTAAATCAAATACAAATTGTGGAGGAGGCAACCATGGTTATAA
487 N E F R S M E A I F L A H G P S F K E K
1501 CAATGAGTTTAGGAGCATGGAGGCTATCTTTCTGGCACATGGACCCAGTTTTAAAGAGAA
507 T E V E P F E N I E V Y N L M C D L L R
1561 GACTGAAGTTGAACCATTTGAAAATATTGAAGTCTATAACCTAATGTGTGATCTTCTACG
527 I Q P A P N N G T H G S L N H L L K V P
1621 CATTCAACCAGCACCAAACAATGGAACCCATGGTAGTTTAAACCATCTTCTGAAGGTGCC
547 F Y E P S H A E E V S K F S V C G F A N
1681 TTTTATGAGCCATCCCATGCAGAGGAGGTGTCAAAGTTTTCTGTTTGTGGCTTTGCTAA
567 P L P T E S L D C F C P H L Q N S T Q L
1741 TCCATTGCCCACAGAGTCTCTTGACTGTTTCTGCCCTCACCTACAAAATAGTACTCAGCT
587 E Q V N Q M L N L T Q E E I T A T V K V
1801 GGAACAAGTGAATCAGATGCTAAATCTCACCCAAGAAGAAATAACAGCAACAGTGAAAGT
607 N L P F G R P R V L Q K N V D H C L L Y
1861 AAATTTGCCATTTGGGAGGCCTAGGGTACTGCAGAAGAACGTGGACCACTGTCTCCTTTA
627 H R E Y V S G F G K A M R M P M W S S Y
1921 CCACAGGGAATATGTCAGTGGATTTGGAAAAGCTATGAGGATGCCCATGTGGAGTTCATA
647 T V P Q L G D T S P L P P T V P D C L R
1981 CACAGTCCCCCAGTTGGGAGACACATCGCCTCTGCCTCCCAGTGTCCCAGACTGTCTGCG
667 A D V R V P P S E S Q K C S F Y L A D K
2041 GGCTGATGTCAGGGTTCCTCCTTCTGAGAGCCAAAATGTTTCCTTCTATTTAGCAGACAA
687 N I T H G F L Y P P A S N R T S D S Q Y
2101 GAATATCACCCACGGCTTCCTCTATCCTCCTGCCAGCAATAGAACATCAGATAGCCAATA
707 D A L I T S N L V P M Y E E F R K M W D
2161 TGATGCTTTAATTACTAGCAATTTGGTACCTATGTATGAAGAATTCAGAAAAATGTGGGA
727 Y F H S V L L I K H A T E R N G V N V V
2221 CTACTTCCACAGTGTTCTTCTTATAAAACATGCCACAGAAAGAAATGGAGTAAATGTGGT

Figure 2D-2
SUBSTITUTE SHEET (RULE 26)

12/67

747 S G P I F D Y N Y D G H F D A P D E I T
 2281 TAGTGGACCAATATTTGATTATAATTATGATGGCCATTTTGATGCTCCAGATGAAATTAC
 767 K H L A N T D V P I P T H Y F V V L T S
 2341 CAAACATTTAGCCAACACTGATGTTCCCATCCCAACACACTACTTTGTGGTGCTGACCAG
 787 C K N K S H T P E N C P G W L D V L P F
 2401 TTGTAAAAACAAGAGCCACACACCGGAAACTGCCCTGGGTGGCTGGATGTCCTACCCTT
 807 I I P H R P T N V E S C P E G K P E A L
 2461 TATCATCCCTCACCGACCTACCAACGTGGAGAGCTGTCCTGAAGGTAAACCAGAAGCTCT
 827 W V E E R F T A H I A R V R D V E L L T
 2521 TTGGGTTGAAGAAAGATTTACAGCTCACATTGCCCGGGTCCGTGATGTAGAACTTCTCAC
 847 G L D F Y Q D K V Q P V S E I L Q L K T
 2581 TGGGCTTGACTTCTATCAGGATAAAGTGCAGCCTGTCTCTGAAATTTTGCAACTAAAGAC
 867 Y L P T F E T P I *
 2641 ATATTTACCAACATTTGAAACCCCTATTTAAActtaataatgtctacttaatatataattt
 2701 actgtataaagtaatttttgcaaaatataagtgatTTTTTctggagaattgtaaaataaa
 2761 gTTTTctatttttcttaaaaaaaaaaacggaattccgggcttgggaggctgaggcagga
 2821 gactcgcttgaacccgggaggcagaggttgcagtgagccaagattgcgccattgcactcc
 2881 agagcctgggtgacagagcaagactacatctcaaaaaataaataaaaaataaaagtaa
 2941 caataaaaaataaaaaagaacagcagagagaatgagcaaggagaaatgtcacaaactattgc
 3001 aaaatactgttacactgggttggctctccaagaagatactggaatctcttcagccatttg
 3061 cttttcagaagtagaaaccagcaaaccacctctaagcggagaacatacgattctttatta
 3121 agtagctctggggaaggaaagaataaaaagttgatagctccctgattgggaaaaaatgcac
 3181 aattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaaaaaaattca
 3241 caaacgttgggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaaa
 3301 ttttcagttgggatgagactgacaaaaagaacgggatttccaggcataaagttggcgtga
 3361 gctacagagggcaccatgtggctcagtggaagacccttcaagattcaaagttccatttga
 3421 cagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaagccaagtgg
 3481 aaagcgagcaatttgcagcataactgcttctcctagacagggtgagtgggcaaaatagc
 3541 acagtacacacagtgactattagccactgccagaaacagggtgaacagccctgggagaca
 3601 agggaaggcaggtggtgggagttgttcatggagagaaaggagagttttagaaccagcaca
 3661 tccactggagatgctgggccaccagacccctcccagtcaataaagtctggtgcctcattt
 3721 gatctcagcctcatcatgaccctggagagaccctgataccatctgccagtccccgacagc
 3781 ttaggcactccttgccatcaacctgaccccccgagtgggttctccaggctccctgccccac
 3841 ccattcaggccggaattc

Figure 2D-3

Figure 2E: The cDNA (SEQ ID NO: 10) and amino acid sequence (SEQ ID NO: 11) of 161P2F10B variant 5. The 3858 nucleotide sequence of 161P2F10B variant 5 is shown. The open reading frame extends from nucleic acid 44-2671 including the stop codon.

13/67

1 M E S T L T
1 ctactttattctgataaaacaggtctatgcagctaccaggacaATGGAATCTACGTTGAC
7 L A T E Q P V K K N T L K K Y K I A C I
61 TTTAGCAACGGAACAACCTGTTAAGAAGAACTCTTAAGAAATATAAAATAGCTTGCAT
27 V L L A L L V I M S L G L G L G L G L R
121 TGTTCTTCTTGCTTTGCTGGTGATCATGTCACCTGGATTAGGCCTGGGGCTTGGACTCAG
47 K L E K Q G S C R K K C F D A S F R G L
181 GAAACTGGAAAAGCAAGGCAGCTGCAGGAAGAAGTGCTTTGATGCATCATTTAGAGGACT
67 E N C R C D V A C K D R G D C C W D F E
241 GGAGAACTGCCGGTGTGATGTGGCATGTAAAGACCGAGGTGATTGCTGCTGGGATTTTGA
87 D T C V E S T R I W M C N K F R C G E T
301 AGACACCTGTGTGGAATCAACTCGAATATGGATGTGCAATAAATTTTCGTTGTGGAGAGAC
107 R L E A S L C S C S D D C L Q K K D C C
361 CAGATTAGAGGCCAGCCTTTGCTCTTGTTTCAGATGACTGTTTGCAGAAGAAAGATTGCTG
127 A D Y K S V C Q G E T S W L E E N C D T
421 TGCTGACTATAAGAGTGTTTGCCAAGGAGAAACCTCATGGCTGGAAGAAAACCTGTGACAC
147 A Q Q S Q C P E G F D L P P V I L F S M
481 AGCCCAGCAGTCTCAGTGCCCAGAAGGGTTTGACCTGCCACCAGTTATCTTGTTTTCTAT
167 D G F R A E Y L Y T W D T L M P N I N K
541 GGATGGATTTAGAGCTGAATATTTATACACATGGGATACTTTAATGCCAAATATCAATAA
187 L K T C G I H S K Y M R A M Y P T K T F
601 ACTGAAAACATGTGGAATTCATTCAAATACATGAGAGCTATGTATCCTACCAAACCTT
207 P N H Y T I V T G L Y P E S H G I I D N
661 CCCAAATCATTACACCATTTGTCACGGGCTTGTATCCAGAGTCACATGGCATCATTGACAA
227 N M Y D V N L N K N F S L S S K E Q N N
721 TAATATGTATGATGTAAATCTCAACAAGAATTTTTCACTTTCTTCAAAGGAACAAAATAA
247 P A W W H G Q P M W L T A M Y Q G L K A
781 TCCAGCCTGGTGGCATGGGCAACCAATGTGGCTGACAGCAATGTATCAAGGTTTAAAGC
267 A T Y F W P G S E V A I N G S F P S I Y
841 CGCTACCTACTTTTGGCCCCGATCAGAAGTGGCTATAAATGGCTCCTTTCCTTCCATATA
287 M P Y N G S V P F E E R I S T L L K W L
901 CATGCCTTACAACGGAAGTGTCCCATTTGAAGAGAGGATTTCTACACTGTTAAAATGGCT
307 D L P K A E R P R F Y T M Y F E E P D S
961 GGACCTGCCCAAAGCTGAAAGACCCAGGTTTTATACCATGTATTTTGAAGAACCTGATTC
327 S G H A G G P V S A R V I K A L Q V V D
1021 CTCTGGACATGCAGGTGGACCAGTCAGTGCCAGAGTAATTAAAGCCTTACAGGTAGTAGA
347 H A F G M L M E G L K Q R N L H N C V N
1081 TCATGCTTTTGGGATGTTGATGGAAGGCCTGAAGCAGCGGAATTTGCACAACTGTGTCAA

Figure 2E-1

SUBSTITUTE SHEET (RULE 26)

14/67

367 I I L L A D H G M D Q T Y C N K M E Y M
1141 TATCATCCTTCTGGCTGACCATGGAATGGACCAGACTTATTGTAACAAGATGGAATACAT
387 T D Y F P R I N F F Y M Y E G P A P R I
1201 GACTGATTATTTTCCCAGAATAAACTTCTTCTACATGTACGAAGGGCCTGCCCCCGCAT
407 R A H N I P H D F F S F N S E E I V R N
1261 CCGAGCTCATAATATACCTCATGACTTTTTTAGTTTTAATTCTGAGGAAATTGTTAGAAA
427 L S C R K P D Q H F K P Y L T P D L P K
1321 CCTCAGTTGCCGAAAACCTGATCAGCATTTCAAGCCCTATTTGACTCCTGATTTGCCAAA
447 R L H Y A K N V R I D K V H L F V D Q Q
1381 GCGACTGCACTATGCCAAGAACGTCAGAATCGACAAAGTTCATCTCTTTGTGGATCAACA
467 W L A V R S K S N T N C G G G N H G Y N
1441 GTGGCTGGCTGTTAGGAGTAAATCAAATACAAATTGTGGAGGAGGCAACCATGGTTATAA
487 N E F R S M E A I F L A H G P S F K E K
1501 CAATGAGTTTAGGAGCATGGAGGCTATCTTTCTGGCACATGGACCCAGTTTTAAAGAGAA
507 T E V E P F E N I E V Y N L M C D L L R
1561 GACTGAAGTTGAACCATTTGAAAATATTGAAGTCTATAACCTAATGTGTGATCTTCTACG
527 I Q P A P N N G T H G S L N H L L K V P
1621 CATTCAACCAGCACCAACAATGGAACCCATGGTAGTTTAAACCATCTTCTGAAGGTGCC
547 F Y E P S H A E E V S K F S V C G F A N
1681 TTTTATGAGCCATCCCATGCAGAGGAGGTGTCAAAGTTTTCTGTTTGTGGCTTTGCTAA
567 P L P T E S L D C F C P H L Q N S T Q L
1741 TCCATTGCCCACAGAGTCTCTTGACTGTTTCTGCCCTCACCTACAAAATAGTACTCAGCT
587 E Q V N Q M L N L T Q E E I T A T V K V
1801 GGAACAAGTGAATCAGATGCTAAATCTCACCCAAGAAGAAATAACAGCAACAGTGAAAGT
607 N L P F G R P R V L Q K N V D H C L L Y
1861 AAATTTGCCATTTGGGAGGCCTAGGGTACTGCAGAAGAACGTGGACCACTGTCTCCTTTA
627 H R E Y V S G F G K A M R M P M W S S Y
1921 CCACAGGGAATATGTCAGTGGATTTGGAAAAGCTATGAGGATGCCCATGTGGAGTTCATA
647 T V P Q L G D T S P L P P T V P D C L R
1981 CACAGTCCCCCAGTTGGGAGACACATCGCCTCTGCCTCCCACTGTCCCAGACTGTCTGCG
667 A D V R V P P S E S Q K C S F Y L A D K
2041 GGCTGATGTCAGGGTTCCTCCTTCTGAGAGCCAAAAATGTTCTTCTATTTAGCAGACAA
687 N I T H G F L Y P P A S N R T S D S Q Y
2101 GAATATCACCCACGGCTTCCTCTATCCTCCTGCCAGCAATAGAACATCAGATAGCCAATA
707 D A L I T S N L V P M Y E E F R K M W D
2161 TGATGCTTTAATTACTAGCAATTTGGTACCTATGTATGAAGAATTCAGAAAAATGTGGGA
727 Y F H S V L L I K H A T E R N G V N V V
2221 CTACTTCCACAGTGTTCTTCTTATAAAACATGCCACAGAAAGAAATGGAGTAAATGTGGT

Figure 2E-2

SUBSTITUTE SHEET (RULE 26)

15/67

```

747  S G P I F D Y N Y D G H F D A P D E I T
2281 TAGTGGACCAATATTTGATTATAATTATGATGGCCATTTTGATGCTCCAGATGAAATTAC
767  K H L A N T D V P I P T H Y F V V L T S
2341 CAAACATTTAGCCAACACTGATGTTCCCATCCCAACACACTACTTTGTGGTGCTGACCAG
787  C K N K S H T P E N C P G W L D V L P F
2401 TTGTAAAAACAAGAGCCACACACCGGAAACTGCCCTGGGTGGCTGGATGTCCTACCCTT
807  I I P H R P T N V E S C P E G K P E A L
2461 TATCATCCCTCACCGACCTACCAACGTGGAGAGCTGTCCTGAAGGTAAACCAGAAGCTCT
827  W V E E R F T A H I A R V R D V E L L T
2521 TTGGGTTGAAGAAAGATTTACAGCTCACATTGCCCCGGTCCGTGATGTAGAACTTCTCAC
847  G L D F Y Q D K V Q P V S E I L Q L K T
2581 TGGGCTTGACTTCTATCAGGATAAAGTGCAGCCTGTCTCTGAAATTTTGCAACTAAAGAC
867  Y L P T F E T T I *
2641 ATATTTACCAACATTTGAAACCACTATTTAAActtaataatgtctacttaatatataattt
2701 actgtataaagtaatttttgcaaaatataagtgattttttctggagaattgtaaaataaa
2761 gttttctatttttccttaaaaaaaaaaacggaattccgggcttgggaggctgaggcagga
2821 gactcgcttgaacccgggaggcagaggttgacgtgagccaagattgcgccattgcactcc
2881 agagcctgggtgacagagcaagactacatctcaaaaaataaataaaaaataaaaagtaa
2941 caataaaaaataaaaaagaacagcagagagaatgagcaaggagaaatgtcacaaactattgc
3001 aaaataactgttacactgggttggtctctccaagaagatactggaatctcttcagccatttg
3061 cttttcagaagtagaaaaccagcaaaccacctctaagcggagaacatacgattctttatta
3121 agtagctctggggaaggaaagaataaaaagttgatagctccctgattgggaaaaaatgcac
3181 aattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaaaacaaattca
3241 caaacgttggtggaaggaaaacagtatagaaaacattactttaactaaaagctggaaaaa
3301 ttttcagttgggatgacgactgacaaaaagaacgggatttccaggcataaagttggcgtga
3361 gctacagagggcaccatgtggctcagtggagacccttcaagattcaaagttccatttga
3421 cagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaagccaagtgggt
3481 aaagcgagcaatttgcagcataactgcttctcctagacagggtgagtgggcaaaaatacg
3541 acagtacacacagtgactattagccactgccagaaacaggctgaacagccctgggagaca
3601 aggggaaggcaggtgggtgggagttgttcacatggagagaaaggagagttttagaaccagcaca
3661 tccactggagatgctgggcccaccagacccctcccagtcaataaagtctggtgcctcattt
3721 gatctcagcctcatcatgaccctggagagaccctgataccatctgccagtcctccgacagc
3781 ttaggcactccttgccatcaacctgaccccccgagtggttctccaggctccctgccccac
3841 ccattcaggccggaattc

```

Figure 2E-3

Figure 2F: The cDNA (SEQ ID NO: 12) and amino acid sequence (SEQ ID NO: 13) of 161P2F10B variant 6. The 3165 nucleotide sequence of 161P2F10B variant 6 is shown. The open

16/67

reading frame extends from nucleic acid 84-2711 including the stop codon.

```

1  atacagtttctcttttgccagactagactaaagaaggagcactactttattctgataaaac
1      M E S T L T L A T E Q P V
61  aggtctatgcagctaccaggacaATGGAATCTACGTTGACTTTAGCAACGGAACAACCTG
14      K K N T L K K Y K I A C I V L L A L L V
121 TTAAGAAGAACACTCTTAAGAAATATAAAATAGCTTGCATTGTTCTTCTTGCTTTGCTGG
34      I M S L G L G L G L G L R K L E K Q G S
181 TGATCATGTCACTTGGATTAGGCCTGGGGCTTGGACTCAGGAACTGGAAAAGCAAGGCA
54      C R K K C F D A S F R G L E N C R C D V
241 GCTGCAGGAAGAAGTGCTTTGATGCATCATTTAGAGGACTGGAGAACTGCCGGTGTGATG
74      A C K D R G D C C W D F E D T C V E S T
301 TGGCATGTAAAGACCGAGGTGATTGCTGCTGGGATTTTGAAGACACCTGTGTGGAATCAA
94      R I W M C N K F R C G E T R L E A S L C
361 CTCGAATATGGATGTGCAATAAATTCGTTGTGGAGAGACCAGATTAGAGGCCAGCCTTT
114     S C S D D C L Q R K D C C A D Y K S V C
421 GCTCTTGTTTCAGATGACTGTTTGCAGAGGAAAGATTGCTGTGCTGACTATAAGAGTGTTT
134     Q G E T S W L E E N C D T A Q Q S Q C P
481 GCCAAGGAGAAACCTCATGGCTGGAAGAAAACCTGTGACACAGCCCAGCAGTCTCAGTGCC
154     E G F D L P P V I L F S M D G F R A E Y
541 CAGAAGGGTTTGACCTGCCACCAGTTATCTTGTTTTCTATGGATGGATTTAGAGCTGAAT
174     L Y T W D T L M P N I N K L K T C G I H
601 ATTTATACACATGGGATACTTTAATGCCAAATATCAATAAACTGAAAACATGTGGAATTC
194     S K Y M R A M Y P T K T F P N H Y T I V
661 ATTCAAATACATGAGAGCTATGTATCCTACCAAACCTTCCCAAATCATTACACCATTG
214     T G L Y P E S H G I I D N N M Y D V N L
721 TCACGGGCTTGTATCCGGAGTCACATGGCATCATTGACAATAATATGTATGATGTAAATC
234     N K N F S L S S K E Q N N P A W W H G Q
781 TCAACAAGAATTTTTCACCTTCTTCAAAGGAACAAAATAATCCAGCCTGGTGGCATGGGC
254     P M W L T A M Y Q G L K A A T Y F W P G
841 AACCAATGTGGCTGACAGCAATGTATCAAGGTTTAAAAGCCGCTACCTACTTTTGGCCCG
274     S E V A I N G S F P S I Y M P Y N G S V
901 GATCAGAAGTGGCTATAAATGGCTCCTTTCCTTCCATATACATGCCTTACAACGGAAGTG
294     P F E E R I S T L L K W L D L P K A E R
961 TCCCATTTGAAGAGAGGATTCTACACTGTTAAATGGCTGGACCTGCCCAAAGCTGAGA
314     P R F Y T M F F E E P D S S G H A G G P
1021 GACCCAGGTTTTATACCATGTTTTTTGAAGAACCTGATTCCTCTGGACATGCAGGTGGAC
334     V S A R V I K A L Q V V D H A F G M L M
1081 CAGTCAGTGCCAGAGTAATTAAAGCCTTACAGGTAGTAGATCATGCTTTTGGGATGTTGA

```

Figure 2F-1

SUBSTITUTE SHEET (RULE 26)

17/67

354 E G L K Q R N L H N C V N I I L L A D H
1141 TGGAAGGCCTGAAGCAGCGGAATTTGCACAACTGTGTCAATATCATCCTTCTGGCTGACC
374 G M D Q T Y C N K M E Y M T D Y F P R I
1201 ATGGAATGGACCAGACTTATTGTAACAAGATGGAATACATGACTGATTATTTTCCCAGAA
394 N F F Y M Y E G P A P R V R A H N I P H
1261 TAAACTTCTTCTACATGTACGAAGGGCCTGCCCCCGCGTCCGAGCTCATAATATACCTC
414 D F F S F N S E E I V R N L S C R K P D
1321 ATGACTTTTTTAGTTTTAATTCTGAGGAAATTGTTAGAAACCTCAGTTGCCGAAAACCTG
434 Q H F K P Y L T P D L P K R L H Y A K N
1381 ATCAGCATTTCAAGCCCTATTTGACTCCTGATTTGCCAAAGCGACTGCACTATGCCAAGA
454 V R I D K V H L F V D Q Q W L A V R S K
1441 ACGTCAGAATCGACAAAGTTCATCTCTTTGTGGATCAACAGTGGCTGGCTGTTAGGAGTA
474 S N T N C G G G N H G Y N N E F R S M E
1501 AATCAAATACAAATTGTGGAGGAGGCAACCATGGTTATAACAATGAGTTTAGGAGCATGG
494 A I F L A H G P S F K E K T E V E P F E
1561 AGGCTATCTTTCTGGCACATGGACCCAGTTTTAAAGAGAAGACTGAAGTTGAACCATTTG
514 N I E V Y N L M C D L L R I Q P A P N N
1621 AAAATATTGAAGTCTATAACCTAATGTGTGATCTTCTACGCATTCAACCAGCACCAAACA
534 G T H G S L N H L L K V P F Y E P S H A
1681 ATGGAACCCATGGTAGTTTAAACCATCTTCTGAAGGTGCCTTTTTATGAGCCATCCCATG
554 E E V S K F S V C G F A N P L P T E S L
1741 CAGAGGAGGTGTCAAAGTTTTCTGTTTGTGGCTTTGCTAATCCATTGCCCACAGAGTCTC
574 D C F C P H L Q N S T Q L E Q V N Q M L
1801 TTGACTGTTTCTGCCCTCACCTACAAAATAGTACTCAGCTGGAACAAGTGAATCAGATGC
594 N L T Q E E I T A T V K V N L P F G R P
1861 TAAATCTCACCCAAGAAGAAATAACAGCAACAGTGAAAGTAAATTTGCCATTTGGGAGGC
614 R V L Q K N V D H C L L Y H R E Y V S G
1921 CTAGGGTACTGCAGAAGAACGTGGACCACTGTCTCCTTTACCACAGGGAATATGTCAGTG
634 F G K A M R M P M W S S Y T V P Q L G D
1981 GATTTGGAAGCTATGAGGATGCCCATGTGGAGTTCATACACAGTCCCCCAGTTGGGAG
654 T S P L P P T V P D C L R A D V R V P P
2041 ACACATCGCCTCTGCCTCCCACTGTCCCAGACTGTCTGCGGGCTGATGTCAGGGTTCCTC
674 S E S Q K C S F Y L A D K N I T H G F L
2101 CTTCTGAGAGCCAAAAATGTTTCCTTCTATTTAGCAGACAAGAATATCACCCACGGCTTCC
694 Y P P A S N R T S D S Q Y D A L I T S N
2161 TCTATCCTCCTGCCAGCAATAGAACATCAGATAGCCAATATGATGCTTTAATTACTAGCA
714 L V P M Y E E F R K M W D Y F H S V L L
2221 ATTTGGTACCTATGTATGAAGAATTCAGAAAAATGTGGGACTACTTCCACAGTGTTCTTC

Figure 2F-2

SUBSTITUTE SHEET (RULE 26)

18/67

```

734   I K H A T E R N G V N V V S G P I F D Y
2281 TTATAAAACATGCCACAGAAAGAAATGGAGTAAATGTGGTTAGTGGACCAATATTTGATT
754   N Y D G H F D A P D E I T K H L A N T D
2341 ATAATTATGATGGCCATTTTGATGCTCCAGATGAAATTACCAAACATTTAGCCAACACTG
774   V P I P T H Y F V V L T S C K N K S H T
2401 ATGTTCCCATCCCAACACACTACTTTGTGGTGCTGACCAGTTGTAAAAACAAGAGCCACA
794   P E N C P G W L D V L P F I I P H R P T
2461 CACCGGAAAACCTGCCCTGGGTGGCTGGATGTCCTACCCTTTATCATCCCTCACCGACCTA
814   N V E S C P E G K P E A L W V E E R F T
2521 CCAACGTGGAGAGCTGTCCTGAAGGTAAACCAGAAGCTCTTTGGGTTGAAGAAAGATTTA
834   A H I A R V R D V E L L T G L D F Y Q D
2581 CAGCTCACATTGCCCGGGTCCGTGATGTAGAACTTCTCACTGGGCTTGACTTCTATCAGG
854   K V Q P V S E I L Q L K T Y L P T F E T
2641 ATAAAGTGCAGCCTGTCTCTGAAATTTTGCAACTAAAGACATATTTACCAACATTTGAAA
874   T I *
2701 CCACTATTTAAActtaataatgtctacttaatatataattttactgtataaagtaatttttg
2761 caaaatataagtgatttttttctggagaattgtaaaataaagttttctatttttccttaa
2821 gtcccctaaaagccataattttttattattccttttttctcttttttcaattctatgaatat
2881 gtattatttttaagttatattttttcacacagagatgatgctatattacaccttcctttt
2941 ttgttggttttcttaaaactctaattctcatgacagattataaccttccttattacttgtttta
3001 tcttactcagaatctttgaatatatttttctgcccagaattatctaaacaaaagggagaa
3061 caaaagaagtatgtctcacttggggaactgaatcaactctaaatcagttttgtcacaaaac
3121 tttttgtatttgactggcaatgctgattaaaattaaaaatgcaca

```

Figure 2F-3

Figure 2G: The cDNA (SEQ ID NO: 14) and amino acid sequence (SEQ ID NO: 15) of 161P2F10B variant 7. The 3988 nucleotide sequence of 161P2F10B variant 7 is shown. The open reading frame extends from nucleic acid 276-2801 including the stop codon.

```

1  ctactttatttctgataaaacaggtctatgcagctaccaggacaatggaatctacgttgac
61  tttagcaacggaacaacctgttaagaagaacactcttaagaaatataaaatagcttgcat
121  tacagggtctctctcctttgggatctcacctcaccacaacctctgtttcccaggctcaag
181  tgatcctcctgcctcagcctcctgagtagcttgaccacaggcacatgccacaaggctca
1      M S L G L G L G L
241  gctaagtttttggttcttcttgctttgctgggtgatcATGTCACCTGGATTAGGCCTGGGGC
10   G L R K L E K Q G S C R K K C F D A S F
301  TTGGACTCAGGAACTGGAAAAGCAAGGCAGCTGCAGGAAGAAGTGCTTTGATGCATCAT
30   R G L E N C R C D V A C K D R G D C C W
361  TTAGAGGACTGGAGAACTGCCGGTGTGATGTGGCATGTAAAGACCGAGGTGATTGCTGCT
50   D F E D T C V E S T R I W M C N K F R C
421  GGGATTTTGAAGACACCTGTGTGGAATCAACTCGAATATGGATGTGCAATAAATTCGTT
70   G E T R L E A S L C S C S D D C L Q K K
481  GTGGAGAGACCAGATTAGAGGCCAGCCTTTGCTCTTGTTTCAGATGACTGTTTGCAGAAGA
90   D C C A D Y K S V C Q G E T S W L E E N
541  AAGATTGCTGTGCTGACTATAAGAGTGTTTGCCAAGGAGAAACCTCATGGCTGGAAGAAA

```

Figure 2G-1**SUBSTITUTE SHEET (RULE 26)**

19/67

110 C D T A Q Q S Q C P E G F D L P P V I L
601 ACTGTGACACAGCCCAGCAGTCTCAGTGCCCAGAAGGGTTTGACCTGCCACCAGTTATCT
130 F S M D G F R A E Y L Y T W D T L M P N
661 TGTTTTCTATGGATGGATTTAGAGCTGAATATTTATACACATGGGATACTTTAATGCCAA
150 I N K L K T C G I H S K Y M R A M Y P T
721 ATATCAATAAACTGAAAACATGTGGAATTCATTCAAAATACATGAGAGCTATGTATCCTA
170 K T F P N H Y T I V T G L Y P E S H G I
781 CCAAAACCTTCCCAAATCATTACACCATTGTCACGGGCTTGATCCAGAGTCACATGGCA
190 I D N N M Y D V N L N K N F S L S S K E
841 TCATTGACAATAATATGTATGATGTAAATCTCAACAAGAATTTTTCACTTTCTTCAAAGG
210 Q N N P A W W H G Q P M W L T A M Y Q G
901 AACAAAATAATCCAGCCTGGTGGCATGGGCAACCAATGTGGCTGACAGCAATGTATCAAG
230 L K A A T Y F W P G S E V A I N G S F P
961 GTTTAAAAGCCGCTACCTACTTTTGGCCCGGATCAGAAGTGGCTATAAATGGCTCCTTTC
250 S I Y M P Y N G S V P F E E R I S T L L
1021 CTTCCATATACATGCCTTACAACGGAAGTGTCCCATTTGAAGAGAGGATTTCTACACTGT
270 K W L D L P K A E R P R F Y T M Y F E E
1081 TAAAATGGCTGGACCTGCCCAAAGCTGAAAGACCCAGGTTTTATACCATGTATTTTGAAG
290 P D S S G H A G G P V S A R V I K A L Q
1141 AACCTGATTCCTCTGGACATGCAGGTGGACCAGTCAGTGCCAGAGTAATTAAAGCCTTAC
310 V V D H A F G M L M E G L K Q R N L H N
1201 AGGTAGTAGATCATGCTTTTGGGATGTTGATGGAAGGCCTGAAGCAGCGGAATTTGCACA
330 C V N I I L L A D H G M D Q T Y C N K M
1261 ACTGTGTCAATATCATCCTTCTGGCTGACCATGGAATGGACCAGACTTATTGTAACAAGA
350 E Y M T D Y F P R I N F F Y M Y E G P A
1321 TGAATACATGACTGATTATTTTCCAGAATAAACTTCTTCTACATGTACGAAGGGCCTG
370 P R I R A H N I P H D F F S F N S E E I
1381 CCCCCGCATCCGAGCTCATAATATACCTCATGACTTTTTTAGTTTTAATTCTGAGGAAA
390 V R N L S C R K P D Q H F K P Y L T P D
1441 TTGTTAGAAACCTCAGTTGCCGAAAACCTGATCAGCATTTCAAGCCCTATTTGACTCCTG
410 L P K R L H Y A K N V R I D K V H L F V
1501 ATTTGCCAAAGCGACTGCACTATGCCAAGAACGTCAGAATCGACAAAGTTCATCTCTTTG
430 D Q Q W L A V R S K S N T N C G G G N H
1561 TGGATCAACAGTGGCTGGCTGTTAGGAGTAAATCAAATACAAATTGTGGAGGAGGCAACC
450 G Y N N E F R S M E A I F L A H G P S F
1621 ATGGTTATAACAATGAGTTTAGGAGCATGGAGGCTATCTTTCTGGCACATGGACCCAGTT
470 K E K T E V E P F E N I E V Y N L M C D
1681 TTAAAGAGAAGACTGAAGTTGAACCATTTGAAAATATTGAAGTCTATAACCTAATGTGTG
490 L L R I Q P A P N N G T H G S L N H L L
1741 ATCTTCTACGCATTCAACCAGCACCAAACAATGGAACCCATGGTAGTTTAAACCATCTTC
510 K V P F Y E P S H A E E V S K F S V C G
1801 TGAAGGTGCCTTTTTATGAGCCATCCCATGCAGAGGAGGTGTCAAAGTTTTCTGTTTGTG
530 F A N P L P T E S L D C F C P H L Q N S
1861 GCTTTGCTAATCCATTGCCACAGAGTCTCTTGACTGTTTCTGCCCTCACCTACAAAATA
550 T Q L E Q V N Q M L N L T Q E E I T A T
1921 GTACTCAGCTGGAACAAGTGAATCAGATGCTAAATCTCACCCAAGAAGAAATAACAGCAA
570 V K V N L P F G R P R V L Q K N V D H C
1981 CAGTGAAAGTAAATTTGCCATTTGGGAGGCCTAGGGTACTGCAGAAGAACGTGGACCACT
590 L L Y H R E Y V S G F G K A M R M P M W
2041 GTCTCCTTTACCACAGGGAATATGTCAGTGGATTTGGAAAAGCTATGAGGATGCCCATGT
610 S S Y T V P Q L G D T S P L P P T V P D
2101 GGAGTTCATACACAGTCCCCAGTTGGGAGACACATCGCCTCTGCCTCCCACTGTCCCAG
630 C L R A D V R V P P S E S Q K C S F Y L
2161 ACTGTCTGCGGGCTGATGTCAGGGTTCCTCCTTCTGAGAGCCAAAATGTTCTTCTATT
650 A D K N I T H G F L Y P P A S N R T S D
2221 TAGCAGACAAGAATATACCCACGGCTTCCTCTATCCTCCTGCCAGCAATAGAACATCAG
670 S Q Y D A L I T S N L V P M Y E E F R K

Figure 2G-2

20/67

2281 ATAGCCAATATGATGCTTTAATTACTAGCAATTTGGTACCTATGTATGAAGAATTCAGAA
690 M W D Y F H S V L L I K H A T E R N G V
2341 AAATGTGGGACTACTTCCACAGTGTTCTTCTTATAAAACATGCCACAGAAAGAAATGGAG
710 N V V S G P I F D Y N Y D G H F D A P D
2401 TAAATGTGGTTAGTGGACCAATATTTGATTATAATTATGATGGCCATTTTGATGCTCCAG
730 E I T K H L A N T D V P I P T H Y F V V
2461 ATGAAATTACCAAACATTTAGCCAACACTGATGTTCCCATCCCAACACACTACTTTGTGG
750 L T S C K N K S H T P E N C P G W L D V
2521 TGCTGACCAGTTGTAAAAACAAGAGCCACACACCGGAAAACCTGCCCTGGGTGGCTGGATG
770 L P F I I P H R P T N V E S C P E G K P
2581 TCCTACCCTTTATCATCCCTCACCGACCTACCAACGTGGAGAGCTGTCCTGAAGGTAAAC
790 E A L W V E E R F T A H I A R V R D V E
2641 CAGAAGCTCTTTGGGTTGAAGAAAGATTTACAGCTCACATTGCCCGGGTCCGTGATGTAG
810 L L T G L D F Y Q D K V Q P V S E I L Q
2701 AACTTCTCACTGGGCTTGACTTCTATCAGGATAAAGTGCAGCCTGTCTCTGAAATTTTGC
830 L K T Y L P T F E T T I *
2761 AACTAAAGACATATTTACCAACATTTGAAACCACTATTTAAActtaataatgtctacttaa
2821 tatataatttactgtataaagtaattttggcaaaatataagtgattttttctggagaatt
2881 gtaaaataaagttttctattttttccttaaaaaaaaaaacggaattccgggcttgggaggc
2941 tgaggcaggagactcgcttgaacccgggaggcagagggttgcagtgagccaagattgcgcc
3001 attgcactccagagcctgggtgacagagcaagactacatctcaaaaaataaataaataaa
3061 ataaaagtaacaataaaaaataaaaaagaacagcagagagaatgagcaaggagaaatgtcac
3121 aaactattgcaaaatactgttacactgggttggctctccaagaagatactggaatctctt
3181 cagccatttgctttttcagaagtagaaaccagcaaaccacctctaagcggagaaacatacga
3241 ttctttattaagtagctctggggaaggaaagaataaaaagttgatagctccctgattggga
3301 aaaaatgcacaattaataaagaatgaagatgaaagaaagcatgcttatgttgtaacacaa
3361 aaaaaattcacaaacgttggtggaaggaaaacagtatagaaaacattactttaactaaaa
3421 gctggaaaaattttcagttgggatgagactgacaaaaagaacgggatttccaggcataaa
3481 gttggcgtgagctacagagggcaccatgtggctcagtggaagacccttcaagattcaaag
3541 ttccatttgacagagcaaaggcacttcgcaaggagaagggtttaaattatgggtccaaaa
3601 gccaaagtggtaaaagcgagcaatttgcagcataactgcttctcctagacagggctgagtgg
3661 gcaaaatacgacagtacacacagtgactattagccactgccagaaacaggctgaacagcc
3721 ctgggagacaagggaaggcagggtggtgggagttgttcatggagagaaaggagagttttag
3781 aaccagcacatccactggagatgctgggccaccagacccctcccagtcaataaagtctgg
3841 tgcctcatttgatctcagcctcatcatgaccctggagagaccctgataccatctgccagt
3901 ccccgacagcttaggcactccttgccatcaacctgaccccccgagtggttctccaggctc
3961 cctgccccacccattcaggccggaattc

Figure 2G-3

21/67

Figure 3A. Amino acid sequence of 161P2F10B variant 1 (SEQ ID NO: 16). The 161P2F10B variants 1 protein has 875 amino acids.

```

1  MESTLTLATE QPVKKNTLKK YKIACIVLLA LLVIMSLGLG LGLGLRKLEK QGSCRKKCFD
61 ASFRGLENCR CDVACKDRGD CCWDFEDTCV ESTRIWMCNK FRCGETRLEA SLCSCSDDCL
121 QKKDCCADYK SVCQGETSWL EENCDDTAQQS QCPEGFDLPP VILFSMDGFR AEYLYTWDTL
181 MPNINKLKTC GIHSKYMRAM YPTKTFPNHY TIVTGLYPES HGIIDNNMYD VNLNKNFSL
241 SKEQNNPAWW HGQPMWLTAM YQGLKAATYF WPGSEVAING SFPSIYMPYN GSVPFEEERIS
301 TLLKWLDLPK AERPRFYTM YFEEDSSGHA GGPVSARVIK ALQVVDHAFG MLMEGLKQRN
361 LHNCVNIILL ADHGMDQTYC NKMEYMTDYF PRINFFYMYE GPAPRIRAHN IPHDEFFSFNS
421 EEIVRNLSR KPDQHFKPYL TPDLPKRLHY AKNVRIDKVH LFVDQQWLAV RSKSNTNCGG
481 GNHGYNNEFR SMEAIFLAHG PSFKEKTEVE PFENIEVYNL MCDLLRIQPA PNNGTHGSLN
541 HLLKVPFYEP SHAEVSKFS VCGFANPLPT ESLDCFCPHL QNSTQLEQVN QMLNLTQEEI
601 TATVKVNLPF GRPRVLQKNV DHCLLYHREY VSGFGKAMRM PMWSSYTVPQ LGDTSPLPPT
661 VPDCLRADVR VPPSESQKCS FYLADKNITH GFLYPPASNR TSDSQYDALI TSNLVPMYEE
721 FRKMWDYFHS VLLIKHATER NGVNVVSGPI FDYNYDGHFD APDEITKHLA NTDVPIPTHY
781 FVVLTSCKNK SHTPENCPGW LDVLPFIIPH RPTNVESCPE GKPEALWVEE RFTAHIAVR
841 DVELLTGLDF YQDKVQPVSE ILQLKTYLPT FETTI

```

Figure 3B. Amino acid sequence of 161P2F10B variant 2 (SEQ ID NO: 17). The 161P2F10B variant 2 protein has 875 amino acids.

```

1  MESTLTLATE QPVKKNTLKK YKIACIVLLA LLVIMSLGLG LGLGLRKLEK QGSCRKKCFD
61 ASFRGLENCR CDVACKDRGD CCWDFEDTCV ESTRIWMCNK FRCGETRLEA SLCSCSDDCL
121 QRKDCCADYK SVCQGETSWL EENCDDTAQQS QCPEGFDLPP VILFSMDGFR AEYLYTWDTL
181 MPNINKLKTC GIHSKYMRAM YPTKTFPNHY TIVTGLYPES HGIIDNNMYD VNLNKNFSL
241 SKEQNNPAWW HGQPMWLTAM YQGLKAATYF WPGSEVAING SFPSIYMPYN GSVPFEEERIS
301 TLLKWLDLPK AERPRFYTM YFEEDSSGHA GGPVSARVIK ALQVVDHAFG MLMEGLKQRN
361 LHNCVNIILL ADHGMDQTYC NKMEYMTDYF PRINFFYMYE GPAPRIRAHN IPHDEFFSFNS
421 EEIVRNLSR KPDQHFKPYL TPDLPKRLHY AKNVRIDKVH LFVDQQWLAV RSKSNTNCGG
481 GNHGYNNEFR SMEAIFLAHG PSFKEKTEVE PFENIEVYNL MCDLLRIQPA PNNGTHGSLN
541 HLLKVPFYEP SHAEVSKFS VCGFANPLPT ESLDCFCPHL QNSTQLEQVN QMLNLTQEEI
601 TATVKVNLPF GRPRVLQKNV DHCLLYHREY VSGFGKAMRM PMWSSYTVPQ LGDTSPLPPT
661 VPDCLRADVR VPPSESQKCS FYLADKNITH GFLYPPASNR TSDSQYDALI TSNLVPMYEE
721 FRKMWDYFHS VLLIKHATER NGVNVVSGPI FDYNYDGHFD APDEITKHLA NTDVPIPTHY
781 FVVLTSCKNK SHTPENCPGW LDVLPFIIPH RPTNVESCPE GKPEALWVEE RFTAHIAVR
841 DVELLTGLDF YQDKVQPVSE ILQLKTYLPT FETTI

```

22/67

Figure 3C. Amino acid sequence of 161P2F10B variant 3 (SEQ ID NO: 18). The 161P2F10B variant 3 protein has 875 amino acids.

```

1  MESTLTATE QPVKKNTLKK YKIACIVLLA LLVIMSLGLG LGLGLRKLEK QGSCRKKCFD
61 ASFRGLENCR CDVACKDRGD CCWDFEDTCV ESTRIWMCNK FRCGETRLEA SLCSCSDDCL
121 QKKDCCADYK SVCQGETSWL EENCDDTAQQS QCPEGFDLPP VILFSMDGFR AEYLYTWDTL
181 MPNINKLKTC GIHSKYMRAM YPTKTFPNHY TIVTGLYPES HGIIDNNMYD VNLNKNFSL
241 SKEQNNPAWW HGQPMWLTAM YQGLKAATYF WPGSEVAING SFPSIYMPYN GSVPFEEERIS
301 TLLKWLDLPK AERPRFYTMY FEEDSSGHA GGPVSARVIK ALQVVDHAFG MLMEGLKQRN
361 LHNCVNIILL ADHGMDQTYC NKMEYMTDYF PRINFFYMYE GPAPRIRAHN IPHDFFSFNS
421 EEIVRNLSR KPDQHFKPYL TPDLPKRLHY AKNVRIDKVH LFVDQQWLAV RSKSNTNCGG
481 GNHGYNNEFR SMEAIFLAHG PSFKEKTEVE PFENIEVYNL MCDLLRIQPA PNNGTHGSLN
541 HLLKVPFYEP SHAEVSKFS VCGFANPLPT ESLDCFCPHL QNSTQLEQVN QMLNLTQEEI
601 TATVKVNLPF GRPRVLQKNV DHCLLYHREY VSGFGKAMRM PMWSSYTVPQ LGDTSPLPPT
661 VPDCLRADVR VPPSESQKCS FYLADKNITH GFLYPPASNR TSDSQYDALI TSNLVPMYEE
721 FRKMWDYFHS VLLIKHATER NGVNVVSGPI FDYNYDGHFD APDEITKHLA NTDVPIPTHY
781 FVVLTSCKNK SHTPENCPGW LDVLPFIIPH RPTNVESCPG GKPEALWVEE RFTAHIAVR
841 DVELLTGLDF YQDKVQPVSE ILQLKTYLPT FETTI

```

Figure 3D. Amino acid sequence of 161P2F10B variant 4 (SEQ ID NO: 19). The 161P2F10B variant 4 protein has 875 amino acids.

```

1  MESTLTATE QPVKKNTLKK YKIACIVLLA LLVIMSLGLG LGLGLRKLEK QGSCRKKCFD
61 ASFRGLENCR CDVACKDRGD CCWDFEDTCV ESTRIWMCNK FRCGETRLEA SLCSCSDDCL
121 QKKDCCADYK SVCQGETSWL EENCDDTAQQS QCPEGFDLPP VILFSMDGFR AEYLYTWDTL
181 MPNINKLKTC GIHSKYMRAM YPTKTFPNHY TIVTGLYPES HGIIDNNMYD VNLNKNFSL
241 SKEQNNPAWW HGQPMWLTAM YQGLKAATYF WPGSEVAING SFPSIYMPYN GSVPFEEERIS
301 TLLKWLDLPK AERPRFYTMY FEEDSSGHA GGPVSARVIK ALQVVDHAFG MLMEGLKQRN
361 LHNCVNIILL ADHGMDQTYC NKMEYMTDYF PRINFFYMYE GPAPRIRAHN IPHDFFSFNS
421 EEIVRNLSR KPDQHFKPYL TPDLPKRLHY AKNVRIDKVH LFVDQQWLAV RSKSNTNCGG
481 GNHGYNNEFR SMEAIFLAHG PSFKEKTEVE PFENIEVYNL MCDLLRIQPA PNNGTHGSLN
541 HLLKVPFYEP SHAEVSKFS VCGFANPLPT ESLDCFCPHL QNSTQLEQVN QMLNLTQEEI
601 TATVKVNLPF GRPRVLQKNV DHCLLYHREY VSGFGKAMRM PMWSSYTVPQ LGDTSPLPPT
661 VPDCLRADVR VPPSESQKCS FYLADKNITH GFLYPPASNR TSDSQYDALI TSNLVPMYEE
721 FRKMWDYFHS VLLIKHATER NGVNVVSGPI FDYNYDGHFD APDEITKHLA NTDVPIPTHY
781 FVVLTSCKNK SHTPENCPGW LDVLPFIIPH RPTNVESCPE GKPEALWVEE RFTAHIAVR
841 DVELLTGLDF YQDKVQPVSE ILQLKTYLPT FETPI

```

23/67

Figure 3E. Amino acid sequence of 161P2F10B variant 7 (SEQ ID NO: 20). The 161P2F10B variant 7 protein has 841 amino acids.

```
1 MSLGLGLGLG LRKLEKQGSC RKKCFDASFR GLENCRCDDVA CKDRGDCCWD FEDTCVESTF
61 IWMCNKFRCG ETRLEASLCS CSDDCLQRKD CCADYKSVCQ GETSWLEENC DTAQQSQCPF
121 GFDLPPVILF SMDGFRAEYL YTWDTLMPNI NKLKTCGIHS KYMRAMYPTK TFPNHYTIVT
181 GLYPESHGII DNNMYDVNLN KNFSLSSKEQ NNPAWWHGQP MWLTAMYQGL KAATYFWPGS
241 EVAINGSFPS IYMPYNGSVP FEERISTLLK WDLPLKAERP RFYTMFFEEP DSSGHAGGPV
301 SARVIKALQV VDHAFGMLME GLKQRNLHNC VNIILLADHG MDQTYCNKME YMTDYFPRIN
361 FFYMYEGPAP RVRAHNIPHD FFSFNSEEIV RNLSCRKPDQ HFKPYLTPDL PKRLHYAKNV
421 RIDKVHLEVD QQWLAVRSKS NTNCGGGNHG YNNEFRSMEA IFLAHGPSFK EKTEVEPFEN
481 IEVYNLMCDL LRIQPAPNNG THGSLNHLK VPFYEPSHAE EVSKFSVCGF ANPLPTESLD
541 CFCPHLQNST QLEQVNQMLN LTQEEITATV KVNLPFGRPR VLQKNVDHCL LYHREYVSGF
601 GKAMRMPMWS SYTVPQLGDT SPLPPTVPDC LRADVVRVPPS ESQKCSFYLA DKNITHGFLY
661 PPASNRTSDS QYDALITSNL VPMYEEFRKM WDYFHSVLLI KHATERNGVN VVSGPIFDYN
721 YDGHFDAPDE ITKHLANTDV PIPTHYFVVL TSCKNKSHTP ENCPGWLDVL PFIIPHRPTN
781 VESCPEGKPE ALWVEERETA HIARVRDVEL LTGLDFYQDK VQPVSEILQL KTYLPTFETT
841 I
```

Figure 4: Comparison of 161P2F10B with known genes: identification of 2 variants for 161P2F10B.

A) Alignment of 161P2F10 (SEQ ID NO: 21) with variant 1 (SEQ ID NO: 22) carrying a K to R mutation.

```

161P2: 1  MESTLTLATEQPVKNTLKKYKIACIVLLALLVIMSLGLGLGLGLRKLEKQGSCRKKCFD 60
          MESTLTLATEQPVKNTLKKYKIACIVLLALLVIMSLGLGLGLGLRKLEKQGSCRKKCFD
Sbjct: 1  MESTLTLATEQPVKNTLKKYKIACIVLLALLVIMSLGLGLGLGLRKLEKQGSCRKKCFD 60

Query: 61  ASFRGLENCRCDVACKDRGDCCWDFEDTCVESTRIWMCNKFRGGETRLEASLCSCSDDCL 120
          ASFRGLENCRCDVACKDRGDCCWDFEDTCVESTRIWMCNKFRGGETRLEASLCSCSDDCL
Sbjct: 61  ASFRGLENCRCDVACKDRGDCCWDFEDTCVESTRIWMCNKFRGGETRLEASLCSCSDDCL 120

Query: 121 QKKDCCADYKSVCQGETSWLEENCDTAQQSQCEGFDLPPVILFSMDGFRAEYLYTWDTL 180
          Q+KDCCADYKSVCQGETSWLEENCDTAQQSQCEGFDLPPVILFSMDGFRAEYLYTWDTL
Sbjct: 121 QRKDCCADYKSVCQGETSWLEENCDTAQQSQCEGFDLPPVILFSMDGFRAEYLYTWDTL 180

Query: 181 MPNINKLKTCGIHSKYMRAMYPTKTFPNHYTIVTGLYPESHGIIDNNMYDVNLNKNFSL 240
          MPNINKLKTCGIHSKYMRAMYPTKTFPNHYTIVTGLYPESHGIIDNNMYDVNLNKNFSL
Sbjct: 181 MPNINKLKTCGIHSKYMRAMYPTKTFPNHYTIVTGLYPESHGIIDNNMYDVNLNKNFSL 240

Query: 241 SKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGSEVAINGSFPSIYMPYNGSVPFEERIS 300
          SKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGSEVAINGSFPSIYMPYNGSVPFEERIS
Sbjct: 241 SKEQNNPAWWHGQPMWLTAMYQGLKAATYFWPGSEVAINGSFPSIYMPYNGSVPFEERIS 300

Query: 301 TLLKWLDLPKAERPRFYTMFYFEEPDSGHAGGPVSARVIKALQVVDHAFGMLMEGLKQRN 360
          TLLKWLDLPKAERPRFYTMFYFEEPDSGHAGGPVSARVIKALQVVDHAFGMLMEGLKQRN
Sbjct: 301 TLLKWLDLPKAERPRFYTMFYFEEPDSGHAGGPVSARVIKALQVVDHAFGMLMEGLKQRN 360

Query: 361 LHNCVNIILLADHGMDQTYCNKMEYMTDYFPRINFFYMYEGPAPRIRAHNIPHDFFSFNS 420
          LHNCVNIILLADHGMDQTYCNKMEYMTDYFPRINFFYMYEGPAPRIRAHNIPHDFFSFNS
Sbjct: 361 LHNCVNIILLADHGMDQTYCNKMEYMTDYFPRINFFYMYEGPAPRIRAHNIPHDFFSFNS 420

Query: 421 EEIVRNLSCKRPDQHFKPYPYLTPLDPKRLHYAKNVRIDKVHLFVDQQWLAVRSKSNNTNCGG 480
          EEIVRNLSCKRPDQHFKPYPYLTPLDPKRLHYAKNVRIDKVHLFVDQQWLAVRSKSNNTNCGG
Sbjct: 421 EEIVRNLSCKRPDQHFKPYPYLTPLDPKRLHYAKNVRIDKVHLFVDQQWLAVRSKSNNTNCGG 480

Query: 481 GNHGYNNEFRSMEAIFLAHGSPFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLN 540
          GNHGYNNEFRSMEAIFLAHGSPFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLN
Sbjct: 481 GNHGYNNEFRSMEAIFLAHGSPFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLN 540

Query: 541 HLLKVPFYEP SHAEEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEI 600
          HLLKVPFYEP SHAEEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEI
Sbjct: 541 HLLKVPFYEP SHAEEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEI 600

Query: 601 TATVKVNLPFGRPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDTSPPLPPT 660
          TATVKVNLPFGRPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDTSPPLPPT
Sbjct: 601 TATVKVNLPFGRPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDTSPPLPPT 660

Query: 661 VPDCLRADVRVPPSESQKCSFYLDKNITHGFLYPPASNRTSDSQYDALITSNLVPMYEE 720
          VPDCLRADVRVPPSESQKCSFYLDKNITHGFLYPPASNRTSDSQYDALITSNLVPMYEE
Sbjct: 661 VPDCLRADVRVPPSESQKCSFYLDKNITHGFLYPPASNRTSDSQYDALITSNLVPMYEE 720

Query: 721 FRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHY 780
          FRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHY
Sbjct: 721 FRKMWDYFHSVLLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHY 780

Query: 781 FVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVR 840
          FVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVR
Sbjct: 781 FVVLTSCKNKSHTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVR 840

Query: 841 DVELLTGLDFYQDKVQPVSEILQLKTYLPTFETTI 875
          DVELLTGLDFYQDKVQPVSEILQLKTYLPTFETTI
Sbjct: 841 DVELLTGLDFYQDKVQPVSEILQLKTYLPTFETTI 875

```

Figure 4A

25/67

B) Alignment of 161P2F10B (SEQ ID NO: 23) and SNP variant (SEQ ID NO: 24) carrying a T to P mutation.

```

161P2: 492 MEAIFLAHGPSFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLNHLLKVPFYEPS 551
          MEAIFLAHGPSFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLNHLLKVPFYEPS
Sbjct: 1   MEAIFLAHGPSFKEKTEVEPFENIEVYNLMCDLLRIQPAPNNGTHGSLNHLLKVPFYEPS 60

Query: 552 HAEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEITATVKVNLPPFG 611
          HAEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEITATVKVNLPPFG
Sbjct: 61  HAEVSKFSVCGFANPLPTESLDCFCPHLQNSTQLEQVNQMLNLTQEEITATVKVNLPPFG 120

Query: 612 RPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDTSPPLPPTVPDCLRADVRV 671
          RPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDTSPPLPPTVPDCLRADVRV
Sbjct: 121 RPRVLQKNVDHCLLYHREYVSGFGKAMRMPMWSSYTPQLGDTSPPLPPTVPDCLRADVRV 180

Query: 672 PPSESQKCSFYLAADKNITHGFLYPASNRTSDSQYDALITSNLVPMYEEFRKMWDYFHSV 731
          PPSESQKCSFYLAADKNITHGFLYPASNRTSDSQYDALITSNLVPMYEEFRKMWDYFHSV
Sbjct: 181 PPSESQKCSFYLAADKNITHGFLYPASNRTSDSQYDALITSNLVPMYEEFRKMWDYFHSV 240

Query: 732 LLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHYFVVLTSCKNKS 791
          LLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHYFVVLTSCKNKS
Sbjct: 241 LLIKHATERNGVNVVSGPIFDYNYDGHFDAPDEITKHLANTDVPIPTHYFVVLTSCKNKS 300

Query: 792 HTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVRDVELLTGLDFY 851
          HTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVRDVELLTGLDFY
Sbjct: 301 HTPENCPGWLDVLPFIIPHRPTNVESCPEGKPEALWVEERFTAHIARVRDVELLTGLDFY 360


Query: 852 QDKVQPVSEILQLKTYLPTFETI 875
          QDKVQPVSEILQLKTYLPTFET I
Sbjct: 361 QDKVQPVSEILQLKTYLPTFETPI 384

```

Figure 4B


26/67

Figure 5: 161P2F10B Hydrophilicity profile
(Hopp T.P., Woods K.R., 1981.
Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828)




27/67

Figure 6: 161P2F10B Hydropathicity Profile
(Kyte J., Doolittle R.F., 1982. J. Mol. Biol. 157:105-132)




28/67

Figure 7: 161P2F10B % Accessible Residues Profile
(Janin J., 1979. Nature 277:491-492)




29/67

Figure 8: 161P2F10B Average Flexibility Profile
(Bhaskaran R., Ponnuswamy P.K., 1988.
Int. J. Pept. Protein Res. 32:242-255)




30/67

Figure 9: 161P2F10B Beta-turn Profile
(Deleage, G., Roux B. 1987. Protein Engineering 1:289-294)



31/67

Figure 10

32/67



Figure 11

Figure 13A


```
c: random coil (31.31%)
e: extended strand (11.31%)
h: alpha helix (57.37%)
```

34/67

Figure 13B




C



35/67

Figure 14 161P2F10B Expression in Normal and Patient Cancer Tissues



XP = Prostate xenograft pool
PMLN(2) = Pool of 2 prostate metastasis to lymph node

Figure 15 161P2F10B Expression in Kidney and Uterus
Patient Cancer Specimens






Figure 16: Phosphodiesterase Activity of 3T3-161P2F10B Stable Cells



37/67

Cell surface phosphodiesterase activity is assayed on 3T3 and 3T3-161P2F10B using the substrate p-nitrophenyl thymidine -5'-L-monophosphate.

Figure 17: Protection from Apoptosis by 161P2F10B






Figure 18 161P2F10B Protects from Apoptotic Signals



•Rat1(E) cells were analyzed using Apo1 caspase assay

Figure 19: 161P2F10B Protects from Staurosporine and UV-Induced Apoptosis



Treatment	Time	% apoptosis			
		3T3-Neo	AGS-16	AGS-16/9C	AGS16/9A
control (DMEM)	24h	10.62	8.28	3.74	8.45
UV (300 mj)	24h	28.57	9.84		
Stauro 2uM	24h	51.47	21.57	24.54	17.91

Figure 20: 161P2F10B Expression Protects Cells from Drug and UV-Induced Apoptosis

Cell Line	Treatment	Time	% Apoptosis
3T3 neo	control	24h	3.62
	Staurosporine, 2.14 uM	24h	64.93
	UV (300 mjoules/cm ²)	24h	62.92
3T3 -AGS16	control	24h	1.72
	Staurosporine, 2.14 uM	24h	14.76
	UV (300 mjoules/cm ²)	24h	30.8

NIH 3T3 cells were treated with the staurosporine or UV, stained with Annexin V-FITC and propidium iodide, and analyzed by FACS.

Figure 21: 161P2F10B Protects from Apoptosis By Chemotherapeutic Agents





Figure 22: Effect of 161P2F10B on *In Vitro* Invasion




- 3T3 cells stably expressing 161P2F10B were assayed using a Transwell Insert System.
- Invasion was determined by measuring the fluorescence of cells in the lower chamber relative to the fluorescence of the entire cell population.
- The assay was performed in triplicate.

Figure 23: 161P2F10B MAb Attenuates the Growth of Human Kidney Cancer
Xenograft in SCID Mice

Group	Mouse #	Tumor volume (mm ³)			
		10/28/02	10/30/02	11/4/02	
PBS control	6738	0.00	0.00	40.00	
	6739	20.25	20.25	31.50	
	6740	0.00	13.50	50.63	
	6741	0.00	0.00	4.00	
	6742	0.00	0.00	0.00	
	Mean	4.05	6.75	25.23	
X41(3)50 MAb	SD	9.1	9.5	22.3	
	6733	0.00	4.00	4.00	
	6734	0.00	0.00	0.00	
	6735	4.00	4.00	40.00	
	6736	6.00	13.50	6.00	
	6737	4.00	4.00	1.00	
	Mean	2.80	5.10	10.20	
	SD	2.7	5.0	16.8	



44/67



- ◆ Tumor injection/first MAb injection – 10/11/02
- ◆ 2.5 x 10⁶ UG-K3 tumor cells were injected s.c. into the mouse flank. MAb (500 µl) injected i.p. twice weekly.


45/67

Figure 24: 161P2F10B protein expression in kidney cancer specimens




46/67

Figure 25: 161P2F10B protein expression in prostate cancer specimen



47/67

Figure 26: 161P2F10B protein expression in colon cancer specimen



48/67

Figure 27: Immunohistochemistry showing expression of 161P2F10B protein to kidney clear cell cancer patient specimens by specific binding of mouse monoclonal antibodies





Figure 28: Expression of 161P2F10b in recombinant cell lines



Cell line	Relative enzyme activity (OD)
Rat1-neo	0.097
Rat1-161P2F10b	0.269
3T3-neo	0.083
3T3-161P2F10b	0.250
300.19-neo	0.014
300.19-161P2F10b	0.112
NSO-neo	0.415
NSO-161P2F10b	0.469
UT7	0.104


Figure 29: Surface expression and phosphodiesterase activity of 161P2F10b

A

Cell surface expression of
161P2F10b in transfected cells

B

Cell surface phosphodiesterase activity
of 161P2F10b in transfected cells



161P2F10b transfected 293T cells were stained with PE-conjugated anti-CD-203c MAb and examined by fluorescent microscopy.

161P2F10b and vector transfected 293T cells were incubated in assay buffer containing the phosphodiesterase-1 colorimetric substrate p-nitrophenyl thymidine-5'-L-monophosphate and optical densities (O.D) were obtained at 405nm.

Figure 30: Relative expression and enzymatic activity of 161P2F10b mutants in recombinant Caki kidney cancer cells




Figure 31: Purification of the recombinant extracellular domain (ECD) of 161P2F10b




Figure 32: 161P2F10b enzymatic assays utilizing P-nitrophenyl-thymidine monophosphate (p-nTMP)






Figure 33:
Analysis of the internalization of anti-161P2F10b MAb X41.6



55/67

Figure 34: Internalization of selected anti-161P2F10b
murine MAbs



56/67

Figure 35: Antibody engagement of 161P2F10b results in its internalization





Figure 36: Effects of X41.50 MAb-saporin toxin conjugate on Caki-161P2F10b cells



58/67

Figure 37: Internalization mediated downregulation of 161P2F10 protein by MAb X41.50





Figure 38: Anti-161P2F10b MAbs downregulate surface 161P2F10b enzymatic activity



Rat1-161P2F10b cells were treated with and without various concentrations of the indicated MAbs for 48 hours and then assayed for surface enzymatic activity.

Figure 39: Characteristics of mouse 161P2F10b MAbs

MAb Clone	Isotype	Relative Affinity (nM)	Surface Staining	Internalization (FACS)	Downregulation of enzyme activity	Specific IHC staining	Epitope family
X41(4)6	IgG1	0.09	s	++	m	s	1
X41(3)15	IgG2b	0.23	m	+	-	s	1
X41(3)17	IgG1	0.12	s	+++	s	m	2
X41(3)29	IgG2a	169.9	w	+	-	s	ND
X41(3)37	IgG1	0.24	m	+++	s	s	2
X41(3)50	IgG1	0.09	s	+++	s	s	2
97A6(CD203c)	IgG1	0.28	s	+++	ND	s	2

60/67

s = strong = +++
m = moderate = ++
w = weak = +
Negative = -
ND=not determined

61/67


Figure 40: Surface staining of selected anti-161P2F10b MAbs

Figure 41: Anti-161P2F10b MAbs X41.6 does not cross-react with ENPP1




Figure 42: Detection of 161P2F10b in the conditioned media of 161P2F10b-expressing cells

SUBSTITUTE SHEET (RULE 26)






Figure 43: Detection of secreted 161P2F10b in the serum of mice bearing UGK3 human kidney cancer xenografts

A

SUBSTITUTE SHEET (RULE 26)




B



Mice were implanted with UGK3 xenograft cells and monitored for tumor size at the indicated times and bled to measure 161P2F10b expression in the serum by an 161P2F10b capture ELISA.

65/67

Figure 44: Expression of 161P2F10B Protein by Immunohistochemistry in Kidney Cancer Patient Specimens



66/67

Figure 45 Expression of 161P2F10B in Human Patient Cancers
by Western Blot





Figure 46 Expression of 161P2F10B in Human Xenograft
Tissues by Western Blot



KiCa Xeno = Xenograft of kidney cancer, clear cell carcinoma
Met Xeno = Xenograft from Kidney cancer metastasis to lymph node
NK = Normal kidney

1 GATCACACAT TAGGTTATNG ACTTCAATAT TTTCAAATGG TTCAACTTCA GTCTTCTCTT
61 TAAAACTGGG TCCATGTGCC AAGAAAGATA GCCTCCCATGC TCCTAAACTC ATTGTTATAA
121 CCATGGTTGC CTCCTCCACA ATTTGTATTT GATTTACTCC TAACAGCCAG CCACTGTTGA
181 TC