

62791/86

FORM 1

REGULATION 9

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

APPLICATION FOR A STANDARD PATENT

We, WARNER-LAMBERT COMPANY, a Corporation organised under the laws of the State of Delaware, United States of America, of 2800 Plymouth Road, Ann Arbor, Michigan, United States of America, hereby apply for the grant of a Standard Patent for an invention entitled:-

"(N-SUBSTITUTED-2-HYDROXY) BENZAMIDES AND N-SUBSTITUTED-2-HYDOXY-a-

OXO-BENZENEACETAMIDES, PHARMACEUTICAL COMPOSITIONS AND METHODS OF

USE THEREOF HAVING ACTIVITY AS MODULATORS OF THE ARACHIDONIC ACID

CASCADE."

which is described in the accompanying Complete Specification.

Details of basic application:-

Number: 782,763

APPLICATION ACCEPTED AND AMENDMENTS

Country: U.S.A.

ALLOWED 27-11-90

Date: 1st October, 1985

Our address for service is:

SHELSTON WATERS

55 Clarence Street

SYDNEY, N.S.W. 2000.

DATED this 15th day of September, 1986

WARNER-LAMBERT COMPANY


by *Robert J. Shelston*

Fellow Institute of Patent Attorneys of Australia
of SHELSTON WATERS

To: Patent Commissioner of Patents
MODERN OFFICE T. 2606

File: DOB 367 91

Fee: Collected Public Moneys



(CONVENTION — Company)

PD-3374-1-JT
FORM 8 — REGULATION 12 (2)

COMMONWEALTH OF AUSTRALIA PATENTS ACT, 1952-1973
DECLARATION IN SUPPORT OF A CONVENTION APPLICATION FOR A PATENT

In support of the Convention Application No. made

(a) Here Insert (In full)
Name of Company.

by ^(a) WARNER-LAMBERT COMPANY

(b) Here Insert Title of
Invention.

(hereinafter referred to as "Applicant") for a patent for an invention entitled:

^(b) (N-SUBSTITUTED-2-HYDROXY) BENZAMIDES AND N-SUBSTITUTED-2-HYDROXY- α -OXO-BENZENEACETAMIDES, PHARMACEUTICAL COMPOSITIONS AND METHODS OF USE THEREOF HAVING ACTIVITY AS MODULATORS OF THE ARACHIDONIC ACID CASCADE

(c) and (d) Here Insert
Full Name and Address
of Company Official
authorised to make
declaration.

^(c) Christine A. Trautwein, Assistant Secretary

^(d) Warner-Lambert Company, 2800 Plymouth Rd.

Ann Arbor, Michigan 48105, United States of America.

do solemnly and sincerely declare as follows:

1. I am authorised by Applicant to make this declaration on its behalf.

2. The basic Application(s) as defined by section 141 of the Act was/were made in ^(e) U.S.A. 782,763 on the 1st day of October 1985

on the day of 19 ..

^(f) by David T. Connor, Daniel L. Flynn, Wlaczslaw A. Cetenko, Elizabeth A. Johnson, Jagadish C. Sircar, Charles Schwender, Roderick J. Sorenson, Paul C. Unangst.


^(g) 3. David T. Connor, 2453 Antietam, Ann Arbor, MI 48105; Daniel L. Flynn, 2700 Lakehurst, Ann Arbor, MI 48105; Wlaczslaw A. Cetenko, 3565 Greerbrier, Ann Arbor, MI 48105; Jagadish C. Sircar, 3615 Charter Place, Ann Arbor, MI 48105; Charles F. Schwender, Box 587C, Philhower Rd. RR#2 Califon, NJ 07830; (SEE ATTACHED SHEET FOR ADDITIONAL ADDRESSES) the actual Inventor(s) of the invention and the facts upon which Applicant is entitled to make the Application are as follows:

Applicant is the Assignee of the said Inventor(s).

4. The basic Application(s) referred to in paragraph 2 of this Declaration was/were the first Application(s) made in a Convention country in respect of the invention, the subject of the Application.

DECLARED at ... WARNER-LAMBERT COMPANY, Ann Arbor, Michigan
this 21st day of ... August 19 86

(h) Personal Signature
of Declarant (c) (no seal,
witness or legalisation).

^(h) 
Christine A. Trautwein
(Signature of Declarant)
Assistant Secretary

To THE COMMISSIONER OF PATENTS.

SHELSTON WATERS
PATENT ATTORNEYS
55 CLARENCE STREET, SYDNEY
AUSTRALIA
Cables: 'Valid' Sydney Telex: 24422

AUSTRALIA, FORM 8

PD-3374-1-JT

ADDITIONAL ADDRESSES OF INVENTORS:

Elizabeth A. Johnson, 711 Meadowsweet Drive, Apt. 207
Corte Madera, CA 94925; Roderick J. Sorenson, 2820 Briarcliff,
Ann Arbor, MI 48105; Paul C. Unangst, 3659 Middleton, Ann Arbor,
MI 48105.


(12) PATENT ABRIDGMENT (11) Document No. AU-B-62791/86
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 606848

- (54) Title
2-OH-BENZAMIDES AND 2-OH-ALPHA OXO-BENZENE ACETAMIDES
- International Patent Classification(s)
(51)^d C07C 103/76 C07C 103/78 C07C 103/85 A61K 031/165
- (21) Application No. : 62791/86 (22) Application Date : 17.09.86
- (30) Priority Data
- (31) Number (32) Date (33) Country
782763 01.10.85 US UNITED STATES OF AMERICA
- (43) Publication Date : 02.04.87
- (44) Publication Date of Accepted Application : 21.02.91
- (71) Applicant(s)
WARNER-LAMBERT COMPANY
- (72) Inventor(s)
DAVID T. CONNOR; DANIEL L. FLYNN; WIACZESLAW A. CETENKO; ELIZABETH A. JOHNSON; JAGADISH C. SIRCAR; CHARLES F. SCHWENDER; RODERICK J. SORENSON; PAUL C. UNANGST
- (74) Attorney or Agent
SHELSTON WATERS, 55 Clarence Street, SYDNEY NSW 2000
- (56) Prior Art Documents
US 3646199
AU 67806/87 C07C 103/76, C07D 295/18, A61K 31/40
AU 67959/87 C07C 103/78, C07C 295/18, A61K 31/165

- (57) The present invention relates to novel compounds, pharmaceutical compositions, and methods of use for the treatment of diseases in which products of lipoxxygenase enzyme activity or the action of leukotrienes contribute to the pathological condition. Selected novel intermediates are also the present invention. The novel 2-hydroxybenzamides and N-substituted-2-hydroxy- α -oxo-benzeneacetamides of the present invention are lipoxxygenase inhibitors providing activity useful for treating asthma, allergies, cardiovascular diseases, migraines, and immunoinflammatory conditions.

CLAIM

6. A compound having the general Formula I



(11) AU-B-62791/86

-2-

(10) 606848

and pharmaceutically acceptable salts thereof, wherein y is one or two; b is zero, one, two, three, or four; R_1 is selected from a group consisting of alkyl of from one to four carbons, inclusive; alkoxy of from one to four carbons, inclusive; thioalkoxy of from one to four carbon atoms, inclusive; carboalkoxy of from two to four carbons, inclusive; alkanoyl of from one to four carbons; phenyl; hydroxy; halogen; hydrogen; nitro; amino; mono- and di-alkylamino having each alkyl the same or different and of from one to four carbons, inclusive; carbalkoxyamido, of from one to four carbons, inclusive; alkylsulfinyl of from one to four carbons, inclusive; alkylsulfonyl of from one to four carbons, inclusive; and when b is one, then R_1 may be $-(CH=CH-CH=CH)-$ taken together with an adjacent ring carbon to form a benzo radical; R_5 is hydrogen; halogen; alkyl of from one to four carbons, inclusive; alkoxy of from one to four carbons, inclusive; carboalkoxy of from two to four carbons, inclusive; hydroxy; halogen; or $-(CH=CH-CH=CH)-$ taken together with adjacent carbons to form a benzo radical; R_6 is alkyl of from six to twenty carbons, $-CH=CH-R_4$, $-(CH_2)_nCOR_4$, or $-(CH_2)_n-R_4$ wherein n is zero to four, inclusive; and R_4 is phenyl, unsubstituted or substituted at the two through six positions by carboalkoxy, having alkoxy of from one to four carbons, inclusive; alkyl of from one to four carbons; alkoxy, or thioalkoxy of from one to four carbons, inclusive; phenalkoxy of from one to four carbons inclusive in the alkoxy group; amino, monoalkyl and dialkyl amino having the alkyl of from one to four carbons, inclusive; alkanoylamino of from two to six

(11) AU-B-62791/86

-2-

(10) 606848

carbons, inclusive; carboxyl; benzo; halogen; hydroxy;

hydroxyalkyl of from

one to four carbons, inclusive; alkanoyl of from one to

four carbons, inclusive; nitro, alkanesulfonamido of from


one to four carbons, inclusive; X is hydrogen or lower

alkyl of from one to four carbons, inclusive, with the

proviso that, when R_1 is halogen and b is one or two

or when b is zero then R_6 cannot be alkyl of from six

to twenty carbons or cannot be



optionally substituted by alkyl of one to four carbons

when R_5 is hydroxy.

COMMONWEALTH OF AUSTRALIA

FORM 10

PATENTS ACT 1952

C O M P L E T E S P E C I F I C A T I O N
606848

FOR OFFICE USE:

Application Number:
Lodged:

62791/86

Class

Int. Class

Complete Specification Lodged:
Accepted:
Published:

Priority:

Related Art:

This document contains the
amendments made under
Section 49 and is correct for
printing.

Name of Applicant:

WARNER-LAMBERT COMPANY

Address of Applicant:

2800 PLYMOUTH ROAD,
ANN ARBOR,
MICHIGAN 48105,
U.S.A.

Actual Inventor:

DAVID T. CONNOR, DANIEL L. FLYNN,
WIACZESLAW A. CETENKO, ELIZABETH A. JOHNSON
JAGADISH C. SIRCAR, CHARLES SCHWENDER,
RODERICK J. SORENSON, AND PAUL C. UNANGST.

Address for Service: SHELSTON WATERS, 55 Clarence Street, Sydney

Complete Specification for the Invention entitled:

"(N-SUBSTITUTED-2-HYDROXY) BENZAMIDES AND N-SUBSTITUTED-2-HYDROXY-a-
OXO-BENZENEACETAMIDES, PHARMACEUTICAL COMPOSITIONS AND METHODS OF USE
THEREOF HAVING ACTIVITY AS MODULATORS OF THE ARACHIDONIC ACID
CASCADE"

The following statement is a full description of this invention,
including the best method of performing it known to me/us:-

BACKGROUND ON THE INVENTION

The present invention relates to novel compounds, pharmaceutical compositions, and methods of use for the treatment of diseases in which products of lipoxxygenase enzyme activity or the action of leukotrienes contribute to the pathological condition. Selected novel intermediates are also the present invention. The novel 2-hydroxybenzamides and N-substituted-2-hydroxy- α -oxo-benzeneacetamides of the present invention are lipoxxygenase inhibitors providing activity useful for treating asthma, allergies, cardiovascular diseases, migraines, and immunoinflammatory conditions.

More particularly, this invention concerns certain novel 2-hydroxybenzamides and novel 2-hydroxy- α -oxo-benzeneacetamides having the Formula I as defined below, pharmaceutical compositions having the novel 2-hydroxybenzamides and novel 2-hydroxy- α -oxobenzeneacetamides therein, and methods of use therefore in the treatment or amelioration of diseases in which products of lipoxxygenase enzyme activity or the reaction of leukotrienes contribute to the pathological condition. Lipoxxygenase enzymes are part of the arachidonic acid cascade.

Arachidonic acid serves as the biological precursor for a family of physiologically active eicosanoids. These include products derived from the action of cyclooxygenase such as the class of prostaglandin-E and -F compounds, thromboxanes, and prostacyclin, and products derived from the action of lipoxxygenase enzymes such as hydroxy- and hydroperoxyeicosatetraenoic acids and the leukotrienes.

Lipoxxygenase pathway products such as the leukotrienes B₄, C₄, D₄, and E₄, 5-hydroxyeicosatetraenoic acid, 5-hydroperoxyeicosatetraenoic acid, and 12-hydroxyeicosatetraenoic acid are involved in the

condition recognized as inflammation, and in allergic and immune responses.

5 These lipxygenase products have been shown to be highly potent stereospecific inducers of polymor-
phonuclear leukocyte migration or chemotaxis,
lysosomal enzyme release, and degranulation.
Additionally, these products induce the contraction
of smooth muscle such as vascular and pulmonary
10 tissue, and induce the generation of additional
inflammogens such as thromboxane A2 and prostacyclin.
Lipxygenase products also interact with vasodilator
prostanoids and other mediators, leading to the
enhancement or amplification of the inflammatory
response.

15 Leukotrienes and the hydroxy- and hydroperoxy-
eicosatetraenoic acids play a major role in the
pathogenesis of many disease conditions. These
compounds have been found in synovial fluid of
rheumatoid joints, in involved skin of psoriatic
20 patients, in inflamed colonic tissue, and at
elevated levels in ischemic myocardial tissue. They
are also mediators of allergic and asthmatic
conditions.


25 Compounds and pharmaceutical compositions in
accordance with the present invention inhibit lipoxy-
genase or the biosynthesis or biochemical action of
leukotrienes and, therefore, are useful in the
treatment or amelioration of a number of diseases
whose pathogenesis involves the production of the
30 leukotrienes and other lipxygenase-derived products.
These lipxygenase inhibitors aid in the prevention
of tissue damage and inflammation which result from
infiltration of leukocytes, release of tissue-
digesting lysosomal enzymes, and changes in the
35 permeability and contractile state of smooth muscle
tissue.

Specific conditions in which such lipxygenase-inhibiting compounds and pharmaceutical compositions in accordance with the present invention are useful include allergy; asthma; arthritis; skin disorders including psoriasis and acne; inflammation; inflammatory bowel diseases; pain; and cardiovascular disorders including myocardial ischemia and infarction, angina, arrhythmias, stroke, and atherosclerosis.

"Derivatives of 3-, 4-, and 5-phenylsalicyl-amides" by H. Jules, et al, J. Am. Pharm. Assoc., Sci. Ed. 45, 277-81 (1956) as reviewed in CA50:16715 describes selected phenylsalicylamides having a phenyl substituent on the phenyl moiety of the phenylsalicylamides and thus differing from the present invention.

Also falling within the scope of the present invention are the pharmaceutically acceptable acid and base addition salts of the compounds of the present invention.

The present invention are compounds of the Formula I



I

and pharmaceutically acceptable salts thereof wherein:

- (1) y is one or two;
- (2) b is zero, one, two, three, or four;
- (3) R₁ is selected from a group consisting of alkyl of from one to four carbons, inclusive, alkoxy of from one to four carbons, inclusive; thioalkoxy of from one to four carbons, inclusive; carboalkoxy of from two to four carbons, inclusive; alkanoyl of from one to four carbons; phenyl; hydroxy, halogen, nitro, amino, mono- and di- alkylamino having each alkyl the same or different from one to four carbons, inclusive, _____



carboalkoxyamido of from one to four carbons,
inclusive, alkylsulfinyl of from one to four carbons,
inclusive, alkylsulfonyl of from one to four carbons,
inclusive, and where b is one then R_1 may also be
5 $-(CH=CH-CH=CH)-$ taken together with adjacent ring
carbons to form a benzo radical;

(4) R_5 is hydrogen; alkyl of from one to four
carbons, inclusive; alkoxy of from one to four
carbons, inclusive; carboalkoxy of from two to four
10 carbons, inclusive; hydroxy, halogen, or
 $-(CH=CH-CH=CH)-$ taken together with adjacent carbons
to form a benzo radical;

(5) R_6 is (a) alkyl of from six to twenty
carbons, (b) $-CH=CH-R_4$, (c) $-(CH_2)_nCOR_4$ or (d)
15 $-(CH_2)_n-R_4$ wherein n is zero to four, inclusive,
and R_4 is phenyl optionally substituted at the two
through six positions by carboalkoxy
having alkoxy of from one to four carbons,
inclusive, alkyl of from one to four carbons, alkoxy
20 or thioalkoxyl of from one to four carbons, inclusive,
phenalkoxy, amino, monoalkyl or dialkyl amino having
the alkyl of from one to four carbons, inclusive,
alkanoylamino of from two to six carbons, inclusive,
carboxyl, halogen, hydroxy, hydroxyalkyl of from one
25 to four carbons, inclusive, alkanoyl of from one to
four carbons, inclusive, nitro, or alkanesulfonamido
of from one to four carbons;

(6) X is hydrogen or lower alkyl of from one
to four carbons, inclusive.

30 One group of preferred compounds of Formula I
include compounds wherein R_1 is hydrogen, y is 1, R_5
is H, and R_6 is alkyl of from 6 to 20 carbons,
inclusive, or $-(CH_2)_n R_4$ wherein n is two and R_4
is phenyl optionally substituted by carboxyl,
35 carboalkoxy of from one to four carbons, inclusive,
chloro, alkoxy of from one to four carbons, inclusive,



hydroxy, or phenyl; or the pharmaceutically acceptable acid or base addition salts thereof.

Another group of preferred compounds of Formula I include compounds wherein R_1 is hydrogen, y
5 is 2, R_5 is hydrogen or the benzo radical; R_6 is alkyl of from 6 to 20 carbons, inclusive, or $-(CH_2)_n-R_4$ wherein n is 2 and R_4 is phenyl optionally substituted by
10 carboxyl, carboalkoxy wherein the alkoxy is from one to four carbons, inclusive, alkoxy of from one to four carbons, inclusive, hydroxy, or pharmaceutically acceptable acid or base addition salts.

Thus, the more preferred compounds of Formula I are:

15 N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy- α -oxo-benzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy- α -oxo-benzeneacetamide.

20 N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-benzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methylbenzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-3-(1,1-dimethylethyl)-2-hydroxy- α -oxobenzeneacetamide.

25 N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxybenzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-N,4-dimethylbenzamide.

The present invention is also a pharmaceutical
30 composition comprising an effective amount of a compound having the Formula I as defined above together with a pharmaceutically acceptable carrier. An effective amount is the amount useful for treating or ameliorating a number of diseases or conditions
35 comprising an inhibition of a lipxygenase effect. The diseases or conditions are readily recognized for



the pathogenesis affected by lipoxxygenase effect, and are recited specifically above.

Thus, in accordance with the present invention, another aspect of the invention, provides a method of administering to mammals, including humans, in need of treatment or amelioration of diseases or conditions an amount effective for treatment of the diseases or conditions as denoted above of a compound or composition having the Formula I as defined above.

The antiasthma and antiallergic activity provides methods of treatment for hypersensitivity reaction having broad symptoms. For example, the symptoms may include dermatitis, lacrimation, nasal discharge, coughing, sneezing, nausea, vomiting, diarrhea, difficulty in breathing, pain, inflammation, and in severe cases, anaphylatic shock and circulatory collapse. The symptoms may be found in man as well as other animals suffering from bronchial asthma, seasonal pollinosis (e.g., hay-fever), allergic rhinitis, urticaria, allergic conjunctivitis, food allergies, and anaphylactoid reactions.

Likewise, the activity of the compounds of Formula I provides a method of treatment for cardiovascular disorders, particularly ischemia and myocardial infarctions. The symptoms of a subject having a cardiovascular disorder may be determined by special diagnostic procedures directed to subjects having a history, general physical appearance and then detailed deviations from normal appearances suggesting a cardiovascular disorder. Such disorders are also found in man as well as other mammals. Symptoms of the disorders are described extensively in The Merck Manual 14th ed, (1982).

Further, method of treatment is provided by the compounds of Formula I herein for migraine and

inflammation. The symptoms requiring treatment for these purposes are also readily recognized, particularly for migraine in man and/or inflammation in man as well as other mammals.

5 Pharmaceutical compositions which also are the present invention are prepared from the compound of Formula I and salts thereof described as the present invention having inert pharmaceutical carriers. The compositions may be either solid or liquid.

10 A physician or veterinarian of ordinary skill readily determines a subject who is exhibiting symptoms described herein. Regardless of the route of administration selected, the compounds of the present invention are formulated into pharmaceu-
15 tically acceptable dosage forms by conventional methods known to the pharmaceutical art.

 The compounds can be administered in such oral unit dosage forms such as tablets, capsules, pills, powders, or granules. They also may be administered
20 rectally or vaginally in such forms as suppositories or bougies; they may also be introduced parenterally (e.g., subcutaneously, intravenously, or intramuscularly), using forms known to the pharmaceutical art. They are also introduced directly
25 to an affected area (e.g., in the form of eye drops or by inhalation). For the treatment of asthma or allergies such as erythema, and inflammatory skin disorders (psoriasis), the compounds of the present invention may also be administered topically in the
30 form of ointments, creams, gels, or the like.

 An effective but nontoxic quantity of the compound is employed in treatment. The ordinarily skilled physician or veterinarian will readily determine and prescribe the effective amount of
35 the compound to prevent or arrest the progress of the condition for which treatment is indicated.

In so proceeding, the physician or veterinarian could employ relatively low dosages at first, subsequently increasing the dose until a maximum response is obtained.

5 Initial dosages of the compounds of the invention having Formula I are ordinarily in the area of 10 mg up to 2 g per day orally, preferably 10 mg to 500 mg per dose orally, given from one to four times daily or as needed. When other forms
10 of administration are employed equivalent doses are administered.

The compounds of the invention are capable of forming both pharmaceutically acceptable acid addition and/or base salts. Base salts are formed with metals
15 or amines, such as ammonium, alkali, and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine,
20 choline, diethanolamine, ethylenediamine, N-methylflucamine, and procaine.

Pharmaceutically acceptable acid addition salts are formed with organic and inorganic acids.

Examples of suitable acids for salt formation
25 are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic, salicylic, malic, gluconic, fumaric, succinic, ascorbic, maleic, methanesulfonic, arginine, and the like. The salts are prepared by contacting the free base form with a sufficient amount of the
30 desired acid to produce either a mono or di, etc salt in the conventional manner. The free base forms may be regenerated by treating the salt form with a base. For example, dilute solutions of aqueous base may be utilized. Dilute aqueous sodium hydroxide, potassium
35 carbonate, ammonia, and sodium bicarbonate solutions are suitable for this purpose. The free base forms

differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but the salts are otherwise equivalent to their respective free base forms for purposes of the invention.

The compounds of the invention can exist in unsolvated as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms and the like are equivalent to the unsolvated forms for purposes of the invention.

Finally, the methods of preparation and selected novel intermediates for preparation for compounds of Formula I as defined above are also the present invention.

Generally, a method of preparation of the compounds of Formula I as defined above can be accomplished as shown in Scheme I wherein R_1 , b , Y , X , and R_5 , and R_6 are as defined above and R is hydrogen, lower alkyl, or phenyl.

When R is hydrogen preparation of the compound of Formula I wherein y is one, is shown in Scheme I (A). The preparation may be accomplished by reacting the salicylic acid of Formula II wherein R is hydrogen with dicyclohexylcarbodiimide or carbonyldiimidazole and the desired compound of Formula III in an inactive solvent, such as tetrahydrofuran, methylene chloride, or ethylene dichloride or mixtures thereof under nitrogen at from about 0°C to about room temperature for from 50 minutes to 24 hours. Optimum conditions vary within reasonable experimentation depending upon the reactants.

Alternatively, when R is lower alkyl or phenyl the preparation of the compound of Formula I wherein y is one or two shown in Scheme I (A) may be accomplished by reacting the ester of Formula II wherein R is lower alkyl or phenyl in the presence of

butyl lithium, diisopropylamine, and the desired aniline of Formula III. An inert organic solvent such as tetrahydrofuran is used in the reaction which is maintained at ice bath temperature with an ice bath
5 for from ten minutes to two hours. See, for example, K. W. Yank, et al, Tetrahedron Letters, 1791 (1970).

According to Scheme I (B) a slight excess of the compound of the Formula II₁ wherein R₁ and b are as defined above, is heated with the compound of
10 Formula III at from 24° to 240°C, preferably of from 140° to 210°C under argon for about two to five hours.

Additionally, the compound of Formula I can be prepared by the method shown in Scheme I (C) where a slight excess of the compound of Formula II₂ is
15 reacted with a compound of Formula III in a nonprotic solvent such as tetrahydrofuran, and the like.

Specific variations within the above general description may include, for example, preparation of compounds of Formula I wherein R₄ includes phenyl optionally substituted by at least one hydroxy group
20 by treatment of corresponding methoxy groups with boron tribromide, hydrobromic acid or trimethylsilyl-iodide using appropriate conditions. Preferred solvents are dichloroethane or dichloromethane. For
25 example, see also M. V. Bhatt and S. U. Kulkarmi, Synthesis (4), 249 (1983) for a review of the cleavage of ethers.

The intermediates of Formula III wherein R₆ is alkyl of from six to twenty carbons, inclusive, are
30 known or can be readily prepared by an ordinarily skilled artisan. However, the novel intermediate of Formula III wherein R₆ is -CH=CH-R₄ and -(CH₂)_n-R₄ or (CH₂)_nCOR₄ are prepared by a synthetic
35 sequence as shown for III₂, III₃, and III₄ in Schemes III or IV, respectively. More specifically, the compound of Formula IV₁, wherein R₇ are the

optional substituents for the phenyl as defined above for R₄, b is an integer of from zero to five, and R₅ is as defined above; is prepared in a manner shown in Scheme II which is analogous to the method disclosed by P. Pfeiffer and S. Sergiewskaya, Ber., 44:1109 (1911). Subsequent reduction of compounds of Formula IV₁ is accomplished by either H₂ and Raney nickel or iron and hydrochloric acid or dithionite to produce the compound of Formula III₂ or Formula III₃, respectively, wherein R₇ and R₅ are as defined above. The reduction is carried out in conditions within the ranges known for the reagents. Reduction of IV₁ by catalytic hydrogenation using a Raney nickel catalyst within the range of conditions known for this reduction produces compounds of Formula III₂ reducing both the nitro-moiety and unsaturation of the hydrocarbon chain in -CH=CH-R₄ of the R₆ definition with the compound of Formula I above. Reduction of IV₁ with iron and HCl or dithionite selectively reduces the nitro moiety.

Intermediate compounds of Formula III₄ wherein R₅ and R₇ are as defined above are obtained by catalytic addition of H₂ to the compound of Formula IV₂ over a palladium/carbon catalyst using conditions within those known or without unreasonable experimentation for hydrogenation using H₂ with these catalysts. Scheme IV shows the hydrogenation of the intermediate precursor having Formula IV₂ to obtain III₄. The compounds of Formula IV₂ having R₅ and R₇ as defined above are prepared in a manner analogous to known Friedel-Crafts acylation methods as disclosed by Tadkod, et al, J. Karntack Univ., 3: 78-80 (1958).

The intermediates of Formula II wherein R₁, b, and R are as defined above and y is one are known or are synthesized by a process analogous to those

known in the art. The intermediates of Formula II wherein R_1 , b, and R are as defined above and y is two, generally, are prepared by reacting a salicylaldehyde type compound of Formula XXII with trimethylsilyl cyanide in the presence of a trace amount of zinc iodide at a temperature of about 0°C to +25°C, preferably 0° to 10°C for about four to twelve hours in an inert atmosphere. The treatment of salicylaldehyde is analogous to the Showalter and Haskell, J. Heterocyclic Chem., 18, 367 (1981), disclosure. The resulting $\alpha,2$ -bistrimethylsiloxybenzeneacetonitrile having an $(R_1)_b$ substituent defined above and as shown by Formula XII is added at the rate of one equivalent over a 20 to 30 minute period to hexamethyldisilazane which is previously treated at about 0°C under an inert atmosphere with from one to slightly more than one equivalent of n-butyllithium and after the treated hexamethyldisilazane is stirred at about 10°C for from 10 to 30 minutes and then cooled to at least about -78°C. The mixture of $\alpha,2$ -bistrimethylsiloxybenzeneacetonitrile with treated and stirred hexamethyldisilazane is stirred for an additional hour. Lower alkyl, preferably methyl or ethyl chloroformate is added to the mixture, stirred, and then warmed. See Scheme V.

Compounds of Formula I wherein X is lower alkyl may be prepared by a process step analogous to known methods.

Under certain circumstances it is necessary to protect either the N or O of intermediates in the above noted process with suitable protecting groups which are known. Introduction and removal of such suitable oxygen and nitrogen protecting groups are well known in the art of organic chemistry; see for example "Protective Groups in Organic Chemistry," J. F. W. McOmie, ed., (New York, 1973), pages 43ff, 95ff;

J. F. W. McOmie, Advances in Organic Chemistry, Vol. 3, 191-281 (1963); R. A. Borssenas, Advances in Organic Chemistry, Vol. 3, 159-190 (1963); and J. F. W. McOmie, Chem. & Ind., 603 (1979).

5 Examples of suitable oxygen protecting groups are benzyl, t-butyldimethylsilyl, ethoxyethyl, and the like. Protection of an N-H containing moiety is necessary for some of the processes described herein for the preparation of compounds of this
10 invention. Suitable nitrogen protecting groups are benzyl, triphenylmethyl, trialkylsilyl, trichloroethylcarbamate, trichloroethoxycarbonyl, vinyloxy-carbamate, and the like.

15 Under certain circumstances it is necessary to protect two different oxygens with dissimilar protecting groups such that one can be selectively removed while leaving the other in place. The benzyl and t-butyldimethylsilyl groups are used in this way; either is removable in the presence of the other,
20 benzyl being removed by catalytic hydrogenolysis, and t-butyldimethylsilyl being removed by reaction with, for example, tetra-n-butylammonium fluoride.

25 In the process described herein for the preparation of compounds of this invention the requirements for protective groups are generally well recognized by one skilled in the art of organic chemistry, and accordingly the use of appropriate protecting groups is necessarily implied by the processes of the charts herein, although not expressly illustrated.

30 The products of the reactions described herein are isolated by conventional means such as extraction, distillation, chromatography, and the like.

35 The salts of compounds of Formula I described above are prepared by reacting the appropriate base or acid with a stoichiometric equivalent of the acid phenol or N base compounds of Formula I,

respectively, to obtain pharmaceutically acceptable salts thereof.

DETAILED DESCRIPTION OF THE INVENTION

By the term, "alkyl of from 6 to 20 carbons, inclusive" is meant any branched or unbranched saturated hydrocarbon grouping having the noted number of carbons, such as hexyl, heptyl, octyl, nonyl, decyl, dodecyl, and the like, and isomers thereof.

The term "alkoxy of from one to four carbons, inclusive" means methoxy, ethoxy, propoxy, or butoxy, and isomers thereof attached to the parent molecular residue through an oxygen atom. Thioalkoxy of from one to four carbons, inclusive, is the same except attached through a sulfur atom.

The term "monoalkyl- or dialkyl-amino having of from one to four carbons, inclusive," means respectively, one or two alkyl groups, as previously defined for of from one to four carbons, inclusive, attached to the parent molecular residue through a nitrogen atom.

The term "alkanoyl of from one to four carbons, inclusive," means a branched or unbranched alkyl, as previously defined for of from one to four carbons, inclusive, attached to the parent molecule residue through the carbonyl group.

The term "hydroxyalkyl of from one to four carbons, inclusive," is an hydroxy attached through an alkyl group, as previously defined for of from one to four carbons, to the parent molecular residue.

The term "alkanoylamino of from two to six carbons, inclusive," means an alkanoyl, as previously defined by including also pentyl or hexyl and isomers thereof among the alkyl attached to the parent molecule residue through the amino group.

The term "carboxyalkoxy having alkoxy of from one to four carbons, inclusive," means an alkyl, as previously defined for alkyl of from one to four carbons, inclusive, attached to the oxygen atom of an ester group, through which the alkyl is attached to the parent molecular residue.

"Halogen" means fluorine, chlorine, bromine, iodine, or trifluoromethyl.

"Carboalkoxyamide of from one to four carbons, inclusive," means an alkyl, as defined above for of from one to four carbons, inclusive, attached to the oxygen atom of an urethane group which is in turn attached to the parent molecule residue through an amino group.

"Alkyl sulfinyl" and "alkyl sulfonyl" are respectively, an alkyl attached to the parent residue molecule through a sulfinyl and sulfonyl group.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention is further elaborated by the representative examples as follows. Such examples are not meant to be limiting thereto.

I. Preparation of compounds of Formula IV

A. For compounds of Formula IV, see Scheme II.

PREPARATION A

1,2-Dimethoxy-4-[2-(4-nitrophenyl)ethenyl]benzene
(See Scheme II, Formula IV), R₇ is 1,2-dimethoxy,
R₅ is hydrogen)

A mixture of 272 g (1.5 mole) of p-nitrophenyl-acetic acid and 249 g (1.5 mole) of 3,4-dimethoxy-benzaldehyde in a 2.0 l nitrogen-filled flask is

heated to 60°C (temperature of reaction mixture) on the steam bath. Piperidine (150 ml; 129 g, 1.52 mole) is added to the warm reaction mixture in small portions over 15 minutes. After ~50 ml of piperidine is added, a mild exotherm developed, and the temperature of the reaction mixture rose to 95°C without external heating. The steam bath is replaced by a heating mantle, and the mixture is heated to reflux over 15 minutes, then maintained at 110-120°C for four hours. The reaction mixture is cooled to 70°C and stirred vigorously while 500 ml of methanol is added. After cooling the mixture in ice, the precipitate that formed is filtered, stirred in 1.0 l of fresh methanol, and refiltered. There is obtained 219 g (51% yield) of olefin product, mp 132-134°C.

PREPARATION B

1,2-Dichloro-4-[2-(4-nitrophenyl)ethenyl]benzene
(See Scheme II, Formula IV, wherein R_7 is
1,2-dichloro, and R_5 is hydrogen)

Prepared by the procedure described in Preparation A, from p-nitrophenylacetic acid (125 g, 0.69 mole) and 3,4-dichlorobenzaldehyde (121 g, 0.69 mole). There was obtained 70 g (35% yield) of the product, mp 197-199°C.

In an manner analogous to that found in above preparation using appropriate starting materials, the following compounds are prepared (see Scheme II).

PREPARATION C

4-[2-[(4-Nitrophenyl)ethenyl][1,1'-biphenyl] mp 238-239°C

PREPARATION D

1-Methoxy-4-[2-(4-nitrophenyl)ethenyl]-2-(phenyl-methoxy)benzene, mp 139-144°C

PREPARATION E

5 1,2-Dimethyl-4-[2-(4-nitrophenyl)ethenyl]benzene, mp 113-115°C

PREPARATION F

1,3-Dimethoxy-5-[2-(4-nitrophenyl)ethenyl]benzene, mp 145-146°C

10

PREPARATION G

2-[2-(4-Nitrophenyl)ethenyl]naphthalene, mp 168-170°C

PREPARATION H

1,2,3-Trimethoxy-5-[2-(4-nitrophenyl)ethenyl]benzene, mp 192-195°C

15

PREPARATION I

1,2-Dimethoxy-3-[2-(4-nitrophenyl)ethenyl]benzene, mp 143-145°C

PREPARATION J

20

2,4-Dimethoxy-1-[2-(4-nitrophenyl)ethenyl]benzene, mp 107-110°C

PREPARATION K

1,2-Dimethoxy-4-[2-(2-nitrophenyl)ethenyl]benzene,
mp 134-137°C

PREPARATION L

- 5 1,2-Dibenzyloxy-4-[2-(4-nitrophenyl)ethenyl]benzene
p-Nitrophenyl acetic acid (29.2 g, 161 mmole)
and 3,4-dibenzyloxybenzaldehyde (51.9 g, 163 mmole)
are mixed with piperidine (16 ml), and heated for three
hours under a dean stark trap. The product is
10 recrystallized from methanol to afford 36.0 g
(51%) of 1,2-dibenzyloxy-4-[nitrophenyl)ethenyl]-
benzene, mp 138-141°C.

B. For compounds of Formula IV₂ see Scheme IV

PREPARATION M

- 15 N-[2-Methoxy-5-[(4-nitrophenyl)acetyl]phenyl]acetamide
(See Scheme IV, Formula IV₂ Wherein R₇ is 2-methoxy
and Acetamide, n is one, and R₅ is hydrogen)
A mixture of anhydrous AlCl₃ (36 g, 270 mmol)
and 50 ml of CH₂Cl₂ is cooled to 0° in an ice bath.
20 2-Acetylanisidine (33 g, 200 mmol) is added to the
stirring mixture. A solution of 39.0 g (200 mmol) of
4-nitrophenylacetyl chloride in 130 ml of CH₂Cl₂
is added slowly to the cooled reaction mixture. The
reaction mixture is stirred at 0°C for 0.75 hour and
25 22 hours at room temperature. The reaction mixture
is poured onto a mixture of 800 ml ice and 40 ml
concentrated hydrochloric acid and allowed to stir
for 1.25 hours before extraction with CH₂Cl₂. The
CH₂Cl₂ extract is evaporated to a dark oily residue
30 which crystallized from MeOH to give 28 g (52%) of a

yellow solid. Further recrystallization from MeOH gave the pure product, mp 200-203°C.

In a manner analogous to that found above in Preparation M using appropriate starting materials the following compounds of Formula IV₂ are prepared.

PREPARATION N

1-(3,4-Dimethoxyphenyl)-3-(4-nitrophenyl)propanone,
mp-126-132°C

PREPARATION O

10 1-(3,4-Dimethoxyphenyl)-4-(4-nitrophenyl)butanone,
mp 109-112°C

II. Preparation of Compounds of Formula III

A. For compounds of Formula III₂ and III₃ see Scheme III.

15 PREPARATION 1

4-[2-(3,4-Dimethoxyphenyl)ethyl]benzeneamine (See Scheme III Formula III₂ Wherein R₇ is 3,4-dimethoxy, and R₅ is Hydrogen

20 A mixture of 19.4 g (0.068 mole) of 1,2-dimethoxy-4-[2-(4-nitrophenyl)ethenyl]benzene as prepared in Preparation A above, and 0.20 g 10% Pd/C catalyst in 200 ml of N,N-dimethylformamide is hydrogenated at 55 psig H₂ pressure for 16 hours. The catalyst is removed by filtration, and the filtrate is
25 evaporated. Recrystallization of the residue from methanol yielded 12.3 g (70% yield) of the amine product, mp 116-117°C.

PREPARATION 2

4-[2-(3,4-Dichlorophenyl)ethyl]benzenamine (See Scheme III, Formula III₂ Wherein R₇ is 3,4-dichloro, b is two, and R₅ is Hydrogen)

5 A mixture of 62.3 g (0.21 mole) of 1,2-dichloro-4-[2-(4-nitrophenyl)ethenyl]benzene as prepared in Preparation B above, and 2.0 g of Raney Nickel catalyst in 935 ml of tetrahydrofuran is hydrogenated at 65 psig H₂ pressure for 20 hours. The catalyst
10 is removed by filtration, and the filtrate is evaporated. Recrystallization of the residue from hexane/dichloromethane yields 49 g (87% yield) of the amine product, mp 73-75°C.

15 In a manner analogous to that found above in Preparations 1 and 2 using appropriate starting materials. The following compounds of Formula III₂ are prepared.

PREPARATION 3

20 4-[2-(1,1'-Biphenyl)-4-ylethyl]benzenamine, mp 109-111°C

PREPARATION 4

4-[2-(2-Naphthyl)ethyl]benzeneamine, mp 123-125°C

PREPARATION 5

25 4-[2-(3-Hydroxy-4-methoxyphenyl)ethyl]benzenamine, mp 152-154°C

PREPARATION 6

4-[2-(3-Methoxyphenyl)ethyl]benzenamine, mp 49-51°C

PREPARATION 7

4-[2-(2,3-Dimethoxyphenyl)ethyl]benzenamine·HCl,
mp 135-136°C

- 5 The starting material, 1,2-dimethoxy-3-[2-(4-nitrophenyl)ethenyl]benzene, is as prepared in Preparation I above.

PREPARATION 8

4-[2-(2,4-Dimethoxyphenyl)ethyl]benzenamine, mp 56-58°C

- 10 The starting material, 2,4-dimethoxy-1-[2-(4-nitrophenyl)ethenyl]benzene, is as prepared in Preparation J above.

PREPARATION 9

- 15 4-[2-(3,4,5-Trimethoxyphenyl)ethyl]benzenamine, mp 91-93°C

 The starting material, 1,2,3-trimethoxy-5-[2-(4-nitrophenyl)ethenyl]benzene, is as prepared in Preparation H above.

PREPARATION 10

- 20 4-[2-(3,5-Dimethoxyphenyl)ethyl]benzenamine·HCl,
mp 155-157°C

 The starting material, 1,3-dimethoxy-4-[2-(4-nitrophenyl)ethenyl]benzene, is prepared in a manner analogous to Preparations A through K.

PREPARATION 11

4-[2-(2-Chlorophenyl)ethyl]benzenamine·HCl, mp 208-211°C

- 5 The starting material, 2-chloro-1-[2-(4-nitrophenyl)ethenyl]benzene, is prepared in a manner analogous to Preparations A through K.

PREPARATION 12

4-[2-(2-Methylphenyl)ethyl]benzenamine·HCl, mp 171-173°C

- 10 The starting material, 2-methyl-1-[2-(4-nitrophenyl)ethenyl]benzene, is prepared in a manner analogous to Preparations A through K.

PREPARATION 13

4-[2-(4-Butoxyphenyl)ethyl]benzenamine, mp 58-59°C

- 15 The starting material, 4-butoxy-1-[2-(4-nitrophenyl)ethenyl]benzene, is prepared in a manner analogous to Preparations A through K.

PREPARATION 14

2-[2-(3,4-Dimethoxyphenyl)ethyl]benzenamine, mp 58-60°C

- 20 The starting material, 1,2-dimethoxy-4-[2-(2-nitrophenyl)ethenyl]benzene, is as prepared in Preparation K above.

PREPARATION 15

N-[2-methoxy-5-[(4-aminophenyl)ethyl]phenyl]acetamide,
mp 135-140°C

- 5 The starting material, N-[2-methoxy-5-[(4-nitro-phenyl)ethenyl]phenyl]acetamide, is prepared in a manner analogous to the methods of Preparations A through K.

PREPARATION 16

10 4-[3-(3,4-dimethoxyphenyl)propyl]benzeamine, mp 54-57°C

The starting material, 1,2-dimethoxy-4-[3-(4-nitrophenyl)prop-2-enyl]benzene, is prepared in a manner analogous to Preparations A through K above.

PREPARATION 17

15 4-[4-(3,4-Dimethoxyphenyl)butyl]benzamine, mp 97-100°C

The starting material, 1,2-dimethyl-4-[4-(4-nitrophenyl)but-3-enyl]benzene is prepared in a manner analogous to Preparations A through K above.

- 20 B. An alternate method of preparation for a compound of Formula III wherein R₆ is (CH₂)_n-R₄ wherein n is one or two is as follows.

PREPARATION 18

4-[(3,4-Dimethoxyphenyl)methyl]aniline

- 25 Mixture of glacial acetic acid (100 ml), 20% Pd/C catalyst (0.5 g) and 3,4-dimethoxy-4'-nitrobenzo-phenone (Tadkod, Kulkarni, and Nargund, J. Karnatak

Univ., 3, 78-80 (1958)) (5.4 g, 18.8 mmol) is hydrogenated at 52 psi for about five hours.

Concentrated H₂SO₄ (1.1 ml) and additional 20% Pd/C (0.5 g) are added and the hydrogenation is continued until five equivalents are consumed (21.2 hours). Potassium acetate (2 g, 20 mmol) is added to the mixture and the catalyst is removed by filtration through celite. The filtrate is acidified with concentrated HCl (1.7 ml), concentrated in vacuo to a residual oil and dissolved in 10% HCl (400 ml). The acidic solution is washed with Et₂O (2 x 400 ml) and CH₂Cl₂ (1 x 100 ml) and then basified with Na₂CO₃. The aqueous fraction was extracted with CH₂Cl₂ and the CH₂Cl₂ extract was dried with Na₂SO₄. Evaporation of the volatile solvent in vacuo gave 4.4 g (96%) of crude oily product which crystallized upon standing. The analytical amine was obtained by column chromatography; yield, 1.58 g (35%), mp 101-104.

C. For a salt of a compound of Formula III.

PREPARATION 19

4-[2-(3,4-Dihydroxyphenyl)ethyl]benzenamine as an acetate salt, mp 216-218°C

A mixture of 20 g (78 mmol) of 4-[2-(3,4-dimethoxyphenyl)ethyl]aniline which is prepared in Preparation 1 above and 300 ml of 48% hydrobromic acid is stirred at reflux under nitrogen for seven hours and at room temperature overnight. The resultant precipitate is collected, washed with ether, and redissolved in 1 N NaOH. The solution is acidified to pH 6 with glacial HOAc and the resultant precipitate is collected as crude product. Recrystallization from

H₂O and then from MeOH yields the 4-[2-(3,4-dihydroxyphenyl)ethyl]benzenamine as an acetate salt; yield, 13.4 g (76%), mp 216-218°C.

- 5 D. For protected substituents on compounds of Formula III.

PREPARATION 20

4-(3,4-Trimethylsilyloxyphenethyl)aniline

- 10 A mixture of 4-(3,4-dihydroxyphenethyl)aniline (34.39 g, 0.15 mole) and hexamethyldisilazane (24.2 g, 0.15 mole) is heated in a wax bath at 120-160°C for 3.75 hours under nitrogen, to give dark colored oily residue, which is chromatographed on silica gel (160 g). Elution with chloroform gives oily product 15 (47.1 g, 84%) of satisfactory purity for the next step.

PREPARATION 21

4-[2-(3,4-dibenzyloxyphenyl)ethyl]benzeneamine

- 20 The 1,2-dibenzyloxy-4-[2-(4-nitrophenyl)ethenyl]-benzene (10 g, 22 mmole) is dissolved in methanol (50 ml), THF (100 ml) and reacted with Ra-N₁ (1.5 g). Reduction under a pressure of 512 psi at 19.5°C affords 5.9 g (60%) of 4-[2-(3,4-dibenzyloxyphenyl)ethyl]-benzenamine, mp 97-101°C.

- 25 PREPARATION 22

N-formyl-4-[2-(3,4-dibenzyloxyphenyl)ethyl]benzenamine

- 30 The 4-[2-(3,4-dibenzyloxyphenyl)ethyl]benzenamine (4.5 g, 11 mmole) is dissolved in toluene (75 ml) containing formic acid (0.51 g, 11 mmole) and refluxed for two hours. The reaction mixture is evaporated to

dryness, and the residue is recrystallized from toluene to afford 4.7 g (97%) of N-formyl-4-[2-(3,4-dibenzyloxyphenyl)ethyl]benzeneamine, mp 119-122°C.

PREPARATION 23

5 N-Methyl-4-[2-(3,4-dibenzyloxyphenyl)ethyl]benzeneamine

LAH (0.25 g, 6.5 mmole) is added to dry THF (20 ml) under an inert atmosphere, then cooled to ~4°C (ice/water bath). N-formyl-4-[2-(3,4-dibenzyloxyphenyl)ethyl]benzeneamine (2.9 g, 6.5 mmole)
10 is dissolved in dry THF (20 ml) and added via cannula dropwise to the LAH/THF suspension. The reaction is stirred for 20 hours at ambient temperature under an inert atmosphere. Water (0.25 ml) followed by 15% NaOH (0.25 ml) and finally
15 water (0.75 ml) is added to the reaction mixture. The mixture is filtered, diluted with 0.5 volume of ether, and the organics are washed with brine and then dried (Na₂SO₄). Concentration affords 1.9 g (70%) of N-methyl-4-[2-(3,4-dibenzyloxyphenyl)ethyl]-
20 benzeneamine, mp 61-65°C, of sufficient purity for further use.

PREPARATION 24

N-formyl-4-[2-(3,4-dimethoxyphenyl)ethyl]benzeneamine

The 4-[2-(3,4-dimethoxyphenyl)ethyl]benzenamine
25 (9.9 g, 38.9 mmole) is dissolved in toluene (125 ml) containing formic acid (5.0 g, 108 mole) and refluxed for one hour. The reaction mixture is cooled to room temperature and evaporated to dryness. Recrystalliza-
tion of the residue from toluene affords 11.0 g (82%)
30 of N-formyl-4-[2-(3,4-dimethoxyphenyl)ethyl]benzen-amine, mp 128-130°C.

PREPARATION 25

N-methyl-4-[2-(3,4-dimethoxyphenyl)ethyl]benzeneamine

LAH (1.2 g, 31.6 mole) is added to dry THF (35 ml) under an inert atmosphere then cooled to about 4°C (ice/water bath). N-formyl-4-[2-(3,4-dimethoxyphenyl)ethyl]benzenamine (9.0 g, 31.5 mole) is dissolved in dry THF (35 ml) and added dropwise via a cannula to the LAH/THF suspension. The reaction is stirred for 20 hours at ambient temperature. Water (1.2 ml) is added followed by 15% NaOH (1.2 ml) and finally water (3.6 ml). The mixture is filtered and the filtrate is diluted with 0.5 volume of ether, washed with brine, dried (Na₂SO₄), and concentrated to afford 6.8 g (79%) of N-methyl-4-[2-(3,4-dimethoxyphenyl)ethyl]-benzenamine, mp 82-84°C.

III. Preparation of Compounds of Formula II

PREPARATION I

2,2,7-Trimethyl-4H-1,3,2-benzodioxasilin-4-one. See Scheme I (B) Compound of Formula II₁, Wherein R₁ is methyl

A solution of 4-methylsalicylic acid (5.0 g, 33 mmoles) in CHCl₃ (30 mls) is stirred under argon. Pyridine (8.0 mls, 99 mmoles) is added followed by dichlorodimethylsilane (4.0 mls, 33 mmoles) and the mixture is heated to reflux. After 2.5 hours additional pyridine (1 ml, 12 mmoles) and dichlorodimethylsilane (0.5 mls, 4 mmoles) is added and reflux continued for another 30 minutes. The mixture is then cooled and the solvent removed under reduced pressure. The residue is treated with pet ether, filtered off, and rinsed five times with pet ether.

The filtrate is evaporated to leave the product, (4.2 g) mp 69-72°C, suitably pure for subsequent reactions.

PREPARATION II

5 Step I

5-Chloro- α ,2-[(trimethylsilyl)oxy]-benzeneacetonitrile
(Formula XII, Scheme 5)

A mixture of 5-chlorosalicylaldehyde (12.15 g, 77.6 mmol), trimethylsilylcyanide (16.94 g, 170.7 mmol),
10 and zinc iodide (2 mg) is stirred at 0°C for four hours under argon atmosphere. The mixture is then allowed to warm to ambient temperature overnight (12 hours). The viscous oil is vacuum-distilled to afford 5-chloro- α ,2-[(trimethylsilyl)oxy]-benzeneacetonitrile, bp 120-
15 122°C/0.27 mmHg, in 56% yield.

Step II

Ethyl 5-chloro-2-hydroxy-2-oxobenzeneacetate

Hexamethyldisilazane (3.11 g, 19.27 mmol) is dissolved in tetrahydrofuran (20 ml) and cooled to 0°C
20 under an argon atmosphere. n-Butyllithium (2.3 M, 8.4 ml, 19.27 mmol) is added, and the solution is stirred at 10°C for 20 minutes followed by cooling to -78°C. At this time, 5-chloro- α ,2-[(trimethylsilyl)oxy]-benzeneacetonitrile (6.00 g, 18.35 mmol) as
25 prepared in Step I above is added over a 30 minute period. After stirring for an additional hour, ethyl chloroformate (1.95 ml, 20.19 mmol) is added dropwise. The solution is stirred for one hour and then allowed to rise to 10°C over a 90 minute period. The reaction
30 is quenched by pouring the contents into saturated ammonium chloride solution followed by extraction into dichloromethane. The organics are washed (saturated ammonium chloride solution followed by brine), dried

(sodium sulfate), and concentrated to afford a residue. The residue is dissolved in tetrahydrofuran (60 ml). Triethylamine hydrofluoride (5.93 g, 48.99 mmol) is added, and the solution is stirred at 0°C for 5 90 minutes. The solution is then concentrated and redissolved in dichloromethane. The organics are washed (1% hydrochloric acid followed by brine), dried (sodium sulfate), and concentrated to give 3.62 g of an oil. Chromatography (Kieselgel 60, 10 dichloromethane) affords ethyl 5-chloro-2-hydroxy- α -oxobenzeneacetate (0.82 g) as a light yellow oil.

PREPARATION III

Following the procedure of Preparation II, 15 Step I, α ,2-[(trimethylsilyl)oxy]benzeneacetonitrile is prepared (bp 103-104°C/1.8 mmHg, 86% yield).

PREPARATION IV

Following the procedure of Preparation II, 20 Step II, ethyl 2-hydroxy- α -oxo-benzeneacetate is prepared (70%).

PREPARATION V

7-(1,1-dimethylethyl)-2,3-benzofurandione (Zwanenburg, Synthesis, 624 (1976)) (See Scheme IC, Compound II₂)

A mixture of 2-tert-butylphenol (15 g, 25 0.1 mmol) and 4-dimethylaminopyridine (0.5 g) is stirred under nitrogen in 300 ml of dichloromethane. Oxalyl chloride (20 ml, 0.22 moles) is added dropwise, then the mixture is heated to reflux. After ten hours the mixture is cooled and the 30 solvent is removed under reduced pressure. The

residue is taken up in 100 ml of 1,2-dichloroethane and added dropwise under nitrogen to a suspension of aluminum chloride (40 g, 0.3 mmol) in 300 ml of 1,2-chloroethane. After 20 hours at room temperature the mixture is slowly diluted with water until all solids dissolve. The organic layer is separated and dried over molecular sieves, and the solvent is evaporated to leave a syrup which is taken up in chloroform and filtered through a short column of silica gel. The filtrate is stripped of solvent under reduced pressure to leave the product (9.8 g) as a syrup suitably pure for subsequent reactions.

IV. Preparation of Compounds of Formula I Wherein y is one

EXAMPLE 1

Benzamide, N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxy

A mixture of 4-methoxysalicylic acid (1.00 g, 5.95 mmol), 4-[2-(3,4-dimethoxyphenyl)ethyl]benzenamine (1.53 g, 5.95 mmol), and dicyclohexylcarbodiimide (1.23 g, 5.95 mmol) in dichloromethane (50 ml) is stirred for 12 hours at ambient temperature. The insoluble dicyclohexylurea is removed by filtration, and the filtrate is concentrated to afford 2.63 g of solid residue. Chromatography (Merck Kieselgel 60, chloroform:ethyl acetate 19:1) of the residue gives N-[4-[2-[(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxybenzamide (0.89 g, 40%); mp 146-148°C after recrystallization from 2-propanol.

Examples 2-7 are prepared by the method of Example 1 and are summarized in Table 1.

TABLE 1

Compounds of Formula I wherein y is one, R₅ is hydrogen, b is one, and R₁ and R₆ are as shown.

5	Example	R ₁ Position				R ₆	Yield	mp (°C)
		3	4	5	6			
	2	H OCH ₃	H H			4-[<u>n</u> -decyl]	47%	116-117
	3	H OCH ₃	H H			4-[2-(3,4-bistri-	21%	
						methylsiloxo-		
10						phenyl)ethyl]		
	4	H Ph*	H H			4-[2-(3,4-dimeth-	20%	190-192
						oxyphenyl)ethyl]		
	5	H CH ₃	H H			4-[<u>n</u> -decyl]	27%	141-142
	6	H Ph	H H			4-[<u>n</u> -decyl]	25%	154-156
15	7	H Cl	H H			4-[<u>n</u> -decyl]	13%	159-160

*Ph is phenyl

EXAMPLE 8

N-[4-[2-(3,4-Dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxybenzamide

20 To a solution of N-[4-[2-(3,4-bis(trimethyl-siloxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxybenzamide (0.65 g, 1.24 mmol) in methanol (30 ml) is added seven drops of concentrated hydrochloric acid. The solution is then heated to 40-50°C for ten

25 minutes. Removal of volatiles gives 0.491 g of a white solid. Recrystallization (2-propanol) affords N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxy benzamide (0.42 g, 90%), mp 179-180°C, 195-196°C (double mp).

EXAMPLE 9

[1,1'-Biphenyl]-3-carboxamide, N-(4-decylphenyl)-2-hydroxy

To a tetrahydrofuran (20 ml) solution of 4-decylaniline (3.07 g, 13.14 mmol), cooled to 0-5°C (inert atmosphere), is added n-butyllithium (2.3 M, 13.14 mmol). The deeply colored solution is stirred for ten minutes, after which a tetrahydrofuran (20 ml) solution of methyl 3-phenylsalicylate (1.00 g, 4.38 mmol) is added. The temperature is allowed to rise to 25°C over a 30 minute period. The reaction is quenched by pouring the contents into 10% hydrochloric acid (100 ml). The organics are extracted into ethyl acetate, washed with 10% hydrochloric acid, dried (sodium sulfate), and concentrated to give 2.77 g of crude solid. Chromatography (Kieselgel 60, dichloromethane) affords N-(4-decylphenyl)-2-hydroxy-[1,1'-biphenyl]-3-carboxamide (1.69 g, 90%), mp 74-75°C.

Similarly, Examples 10-20 are prepared by the method of Example 9 and are found in Table 2.

TABLE 2

A compound of formula I wherein y is one, R₅ is hydrogen, b is one, and R₁ and R₆ are as shown:

5	Example	R ₁ Position				R ₆	Yield	mp (°C)
		3	4	5	6			
10	10	H	H	Ph*	H	4-(<u>n</u> -decyl)	90%	179-180
	11	H	Ph	H	H	4-(<u>n</u> -decyl)	35%	155-157
	12	H	H	Br	H	4-(<u>n</u> -decyl)	74%	172-174
	13	H	H	H	H	4-(<u>n</u> -decyl)	45%	94-95
15	14	H	H	Br	H	4-[2-(3,4-dimethoxy-phenyl)ethyl]	76%	154-156
	15	H	Cl	H	H	4-[2-(3,4-dimethoxy-phenyl)ethyl]	80%	166
	16	H	H	Cl	H	4-(<u>n</u> -decyl)	80%	165
20	17	H	H	CH ₃	H	4-(<u>n</u> -decyl)	73%	127-128
	18	NO ₂	H	H	H	4-(<u>n</u> -decyl)	79%	101-102
	21	H	H	NO ₂	H	4-(<u>n</u> -decyl)	87%	137-138
	22	CH ₃	H	H	H	4-(<u>n</u> -decyl)	71%	90-91
25	23	Cl	H	H	H	4-[2-(3,4-dimethoxyphenyl)ethyl]	45%	125
	24	Cl	Cl	H	H	4-[2-(3,4-dimethoxyphenyl)ethyl]	63%	175-177
	25	H	H	H	OH	4-(<u>n</u> -decyl)	38%	121-123
30	26	H	Me	H	H	4-[2-(3-methoxy-4-hydroxyphenyl)-ethyl]	82%	170
	27	Cl	Cl	H	H	4-(<u>n</u> -decyl)	81	154-155
	28	Cl	H	H	H	4-(<u>n</u> -decyl)	73	124
	29	H	CH ₃	H	H	4-[2-(3,4-dibenzoyloxyphenyl)ethyl]; N-methyl	71	non-crystalline

TABLE 2 (CONT'D)

Example	R ₁	R ₆	% Yield	mp (°C)
30	3,4 	4-[2-(4-chloro phenyl)-ethyl]	61	160-164
31	3,4 	4-2-(3,4-dichloro- phenyl)ethyl]	37	166-168

5

EXAMPLE 32

Benzamide, N-(4-n-decylphenyl)-2-hydroxy-5-amino·hydrochloride

A methanol (75 ml) solution of N-(4-n-decylphenyl)-2-hydroxy-5-nitro-benzamide (890 mg, 2.23 mmol) and Raney-nickel (200 mg) is stirred at ambient temperature until the calculated pressure change is realized. The contents are filtered and acidified with concentrated HCl (0.2 ml). Removal of solvents affords 831 mg of the desired N-(4-n-decylphenyl)-2-hydroxy-5-aminobenzamide-hydrochloride, dc = 240-245°C.

15

EXAMPLE 33

3,5-Dichloro-N-(4-decylphenyl)-2-hydroxybenzamide

Under an argon atmosphere, a tetrahydrofuran (20 ml) solution of diisopropylamine (1.59 ml, 11.31 mmol) is cooled to 0°C. n-Butyllithium (2.3 M, 4.9 ml, 11.31 mmol) is added, after which the solution is allowed to stir an additional ten minutes. n-Decylaniline (2.64 g, 11.31 mmol) is then added and the solution allowed to stir for 15 minutes at ambient temperature. A tetrahydrofuran 925 ml) solution of methyl 3,5-dichlorosalicylate (1.00 g, 4.52 mmol) is

20

25

added and the resulting solution is stirred for 45 minutes at ambient temperature. The contents are then poured into 10% hydrochloric acid (100 ml) and extracted into diethyl ether. The organics are washed with 10% hydrochloric acid, dried (sodium sulfate), and concentrated to give 3.47 g of a residue. Recrystallization from 2-propanol/water gives 3,5-dichloro-N-(4-decylphenyl)-2-hydroxybenzamide (1.32 g, 69%), mp 90-91°C.

Similarly prepared are Examples 34-42, which are found in Table 3.


TABLE 3

A compound of Formula I wherein y is one, R₅ is hydrogen, b is one, x is hydrogen and R₁ and R₆ are as shown:

Example	R ₁ Position			R ₆	Yield	mp (°C)
	3	4	5			
34	Cl	H	H	4-[2-(3,4-dichloro-phenyl)ethyl]	44%	166-167
35	H	NO ₂	H	4-(n-decyl)	40%	180-181
36	H	Cl	H	4-[2-(3,4-dichloro-phenyl)ethyl]	34%	203-204
37	H	Cl	H	4-[2-(3,4-dichloro-phenyl)ethyl]	57%	214-215

TABLE 2 (CONT'D)

Example	R ₁ Position			R ₆	% Yield	mp (°C)
	3	4	5			
5	38	4,5		4-[2-(3,4-dimethoxy-phenyl)ethyl]	38	201-202
	39	3,4		4-[2-(3,4-dimethoxy-phenyl)ethyl]	48	179-180
	40	3,4		4-[2-(1,1'-biphenyl-4-yl)ethyl]*	39	196-198
10	41	5,6		4-[2-(3,4-dimethoxy-phenyl)ethyl]	74	148-150
	42	H	H	4-[2-(3,4-dichloro-phenyl)ethyl]		170



15

EXAMPLE 43

4-Methyl-N-methyl-N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxybenzamide

To an ethyl acetate:methanol (1:1, 40 ml) solution of 4-methyl-N-methyl-N-[4-[2-(3,4-dibenzoyloxyphenyl)ethyl]phenyl]-2-hydroxybenzamide (1.00 g, 1.80 mmol) is added 10% Pd/C (200 mg). The mixture is stirred under hydrogen atmosphere (1 atmosphere) for 12 hours. The catalyst is removed by filtration (Celite®), and the filtrate then concentrated to give 740 mg of crude material. Flash chromatography (SiO₂: 95:5 CHCl₃:MeOH) affords 660 mg (97%) of

4-methyl-N-methyl-[4-(2-(3,4-dihydroxyphenyl)ethyl)-phenyl]-2-hydroxybenzamide as a noncrystalline semisolid.

EXAMPLE 44

5 4-Chloro-N-[4-[2-(3,4-dihydroxyphenyl)ethyl]-2-hydroxybenzamide

A dichloromethane (40 ml) solution of 4-chloro-N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxybenzamide (0.50 g, 1.21 mmol) is cooled to
10 -78°C under an argon atmosphere. Boron tribromide (1.0 M solution in dichloromethane, 5.5 ml, 5.5 mmol) is added, and the mixture is stirred at -78°C for four hours, followed by stirring at ambient temperature for two hours. The solution is then recooled to -20°C
15 and quenched with water (5.5 ml). The mixture is allowed to stir at ambient temperature for 12 hours, after which additional water (10 ml) is added. Filtration affords 4-chloro-N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxybenzamide (0.42 g,
20 91%), mp 229-231°C after recrystallization from methanol/water.

EXAMPLE 45

N-[4-[2-(3,4-Dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methylbenzamide

25 A mixture of 2,2,7-trimethyl-4H-1,3,2-benzodioxasilin-4-one (2.0 g, 10 mmole) and 4-[2-[3,4-bis-[(trimethylsilyl)oxy]phenyl]ethyl]benzenamine (3.0 g, 8 mmol) is heated under argon to 180°C. After three hours the mixture is cooled, triturated with pentane
30 and filtered. The collected solid is rinsed several times with pentane and dried. Recrystallization from isopropanol gave the pure product (1.4 g), mp 201-202°C.

EXAMPLE 46

N-[4-[2-(3,4-Dimethoxyphenyl)ethyl]phenyl]-2-hydroxy-4-methylbenzamide

5 The preparation is as described for Example 45 using 2,2,7-trimethyl-4H-1,3,2-benzodioxasilin-4-one (2.0 g, 10 mmol) and 4-[2-(3,4-dimethoxyphenyl)ethyl]-benzenamine (2.5 g, 10 mmol). Recrystallization from methanol/DMF gives the pure product (1.9 g) mp 162-164°C.

10

EXAMPLE 47

Benzamide, N-[4-[2-(3,4-dihydroxyphenyl)ethyl]-phenyl]-2-hydroxy

15 Prepared by the method described in Example 44 from benzamide, N-[4-[2-(3,4-dimethoxyphenyl)-ethyl]phenyl]-2-hydroxy. Recrystallization from water/2-propanol gives the product, mp 156-158°C.

EXAMPLE 48

N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy benzamide

20 A mixture of 2-acetoxybenzoylchloride (6.5 g, 0.033 mole) and 4-[2-(3,4-dimethoxyphenyl)ethyl]-benzamine (6.5 g; 0.025 mmol) in xylene (300 ml) is stirred at room temperature and then heated to reflux for two hours when a clear solution is formed.
25 The solvent is evaporated off under vacuum and the oil taken up in CH₂Cl₂, washed with sodium bicarbonate solution, with water and dried. The methylene chloride is distilled off. The residual oil is dissolved in hot isodipropyl ether to give the
30 acetate derivative which is removed by filtration. The filtrate is evaporated off to give an oil. The

crude oil is dissolved in methanol (150 ml) and 1(N) NaOH solvent (50 ml) and is heated to reflux for two hours. The reaction mixture is concentrated, diluted with cold water, and then acidified with 4N HCl (30 ml) when the product crystallized out. The crude product is recrystallized from methanol to give analytical sample (1.9 g), mp 149-151.

EXAMPLE 49

10 2-Naphthalenecarboxamide, N-(4-decylphenyl)-3-hydroxy

A suspension of 1.0 g (0.021 mole) of 50% sodium hydride/mineral oil in 25 ml of dimethyl sulfoxide under a nitrogen atmosphere is cooled in a cold water bath while 4.7 g (0.020 mole) of 4-(n-decyl)-aniline is added. The mixture is stirred at room temperature for one hour, then treated in portions over 15 minutes with 2.0 g (0.0099 mole) of 3-hydroxy-2-naphthalene-carboxylic acid methyl ester. An additional 50 ml of dimethyl sulfoxide is added, and the mixture is stirred at room temperature for 45 hours. The reaction mixture is added to 500 g of ice/water and acidified with 4.0 N hydrochloric acid. The gelatinous precipitate is filtered and distributed between water (300 ml) and dichloromethane (100 ml). The layers are separated and the aqueous layer is washed with fresh dichloromethane (2 x 150 ml). The combined organic layers are washed with water (1 x 250 ml), 1.0 N hydrochloric acid (2 x 250 ml), and water again. The organic layer is dried (anhydrous sodium sulfate) and evaporated. Recrystallization of the residue from aqueous 2-propanol yield 1.4 g (35% yield) of the amide product. An additional recrystallization as above yields an analytically pure sample, mp 171-173°C.

V. Preparation of Compounds of Formula I Wherein y
is 2

EXAMPLE 50

N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy-
5 5-chloro- α -oxo-benzeneacetamide

Under an argon atmosphere, a tetrahydrofuran
(20 ml) solution of 4-[2-(3,4-dimethoxyphenyl)ethyl]-
benzenamine (2.50 g, 9.72 mmol) is cooled to 0°C.
n-Butyllithium (2.3 M, 4.2 ml, 9.72 mmol) is added,
10 and the resulting solution is stirred for 15 minutes.
A tetrahydrofuran (10 ml) solution of ethyl 5-chloro-2-
hydroxy- α -oxobenzeneacetate (0.74 g, 3.24 mmol) is
then added, after which the solution is allowed to
warm to ambient temperature over a 30 minute period.
15 The contents are then poured into 5% hydrochloric acid
(100 ml) and extracted into ethyl acetate. The
organics are washed with 10% sodium bicarbonate and
brine, then dried (sodium sulfate), and concentrated
to give 1.51 g of a crude solid. Chromatography
20 (Kieselgel 60, dichloromethane) affords N-[4-[2-
(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy-5-chloro-
 α -oxobenzeneacetamide (0.86 g, 61%), mp 128-130°C.

EXAMPLE 51

N-Methyl-N-[4-(2-(3,4-dimethoxyphenyl)ethyl)phenyl]-2-
25 hydroxy- α -oxo-benzeneacetamide

Following the procedure of Example 50, N-methyl-
N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy-
 α -oxobenzeneacetamide is prepared from N-methyl-N-
4-[2-(3,4-dimethoxyphenyl)ethyl]benzenamine and ethyl-
30 2-hydroxy- α -oxo-benzeneacetate in 22% yield.

EXAMPLE 52

N-[4-(2-(3,4-dimethoxyphenyl)ethyl)phenyl]-2-hydroxy-
α-oxo-Benzeneacetamide

A mixture of 2,3-benzofurandione (Fries and
5 Pfaffendorf, Ber, 45, 156 (1912); Valentine, Titoff,
Muller, and Reichstein, Helv. Chim. Acta, 20, 883
(1937)) (10 g, 0.0675 mol) and 4-[2-(4-aminophenyl)-
ethyl]-1,2-dimethoxybenzene (15.6 g, 0.0606 mol) in
dry tetrahydrofuran is stirred at room temperature
10 under nitrogen for 18 hours in the dark. The
solvent is removed under reduced pressure on a
rotary evaporator below 35° and the resulting solid
is recrystallized from tetrahydrofuran-ethanol to
give 22.2 g (90.6%) of a light-yellow solid, mp 124-
15 125°C.

EXAMPLE 53

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]-phenyl]-2-hydroxy-
α-oxo-benzeneacetamide

A mixture of 2,3-benzofurandione (4.56 g,
20 0.0307 mole) and 1,2-benzenediol, 4-[2-(4-aminophenyl)-
ethyl] (7.05 g, 0.0307 mole) in dry tetrahydrofuran
is stirred at room temperature under nitrogen for
19 hours in the dark. The solvent is removed under
reduced pressure on a rotary evaporator below 50°C
25 and the resulting solid is purified by column
chromatography on silica-gel (260 g). Elution with
ethyl acetate gave 10.7 g of a solid. Recrystallization
from acetonitrile with ice cooling gave 6.9 g (59.6%)
of a light-yellow solid, mp 165-167°C.

EXAMPLE 54

N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-3-(1,1-dimethylethyl)-2-hydroxy- α -oxo-benzeneacetamide

- 5 A solution of 7-(1,1-dimethylethyl)-2,3-benzofurandione (1.0 g, 5 mmol) and 4-[2-(3,4-dimethoxyphenyl)ethyl]benzenamine (1.0 g, 4 mmol) in 10 ml of dichloromethane is stirred at room temperature for 48 hours. The solvent is evaporated and the residue crystallized from ether/hexane.
- 10 Recrystallization from acetonitrile gave the pure product, (1.0 g) mp 143-144°C.

EXAMPLE 55

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-3-(1,1-dimethylethyl)-2-hydroxy- α -oxobenzenacetamide

- 15 A mixture of 7-(1,1-dimethylethyl)-2,3-benzofurandione (3.7 g, 18 mmols) and 4-[2-(4-aminophenyl)ethyl]-1,2-benzenediol (3.4 g, 15 mmols) in 30 ml of dichloromethane is stirred at room temperature for 24 hours. The solution is diluted with ether and
- 20 filtered. The filtrate is evaporated under reduced pressure to leave the product as a syrup which eventually crystallizes. Recrystallization from ether/pet ether gave the pure product (2.0 g) mp 138-149°C.

- 25 The compounds in Tables 4 and 5 are prepared by the method described in Example 53.

TABLE 4

A compound of Formula I wherein y is two, b is one, and R₁ is hydrogen, and R₅ and R₆ are as shown.

5	Example	R5	R6	mp °C	Yield %	Recrystallization Solvent
	56	3-Cl	4-Cl	138-140	67	Tetrahydrofuran/ acetonitrile
	57	H	4-Cl	145-146	69	Tetrahydrofuran/ acetonitrile

10

TABLE 5

	Example	Compound	mp °C	Description
15	58	3-chloro-N-[4-[2-(3,4-dihydroxyphenyl)ethyl]-phenyl]-6-hydroxy-2,4-dimethyl- α -oxo-benzene-acetamide	140-142	Deep yellow solid
20	59	3-chloro-N-[4-[2-(3,4-dimethoxyphenyl)ethyl]-phenyl]-6-hydroxy-2,4-dimethyl- α -oxo-benzene-acetamide	163-164	Fluffy off-white solid
25	60	3-chloro-N-[4-[2-(3,4-dimethoxyphenyl)ethyl]-phenyl]-2-hydroxy- α -oxo-benzenacetamide	125	Yellow solid

The preparation of 2,3-benzofurandione, 5-chloro-4,6-dimethyl is analogous to that described by R. Stolle and E. Knebel Ber. 59, 1216 (1921).

The usefulness of the compounds of the present invention as inhibitors of lipoxxygenase enzyme or antagonists of leukotriene or other related biochemical actions is demonstrated by their effectiveness in various standard pharmacological test procedures. A description of each procedure follows.

10 Human Leukocyte Lipoxxygenase Assay (LDA-H)

Whole blood is collected from normal volunteers and spun in a refrigerated centrifuge for four minutes at 1-6°C at 3800 g. The buffy coat is manually separated and washed twice with chilled 0.83% NH₄Cl and centrifuged at 1000 RPM for ten minutes at 4°C. The white cell is suspended in culture media-EMEM supplemented with 6% Agamma human serum, tricine buffer, and neomycin and recentrifuged at 1000 g to yield a pellet containing the leukocytes used for the preparation of the acetone pentane powder.

The acetone-pentane powder is prepared utilizing a modification of the procedure reported for human platelet lipoxxygenase. See Siegel, et al, Arachidonate Metabolism via Lipoxxygenase and 12-L-hydroperoxy-5-Eicostetraenoic acid Peroxidase Sensitive to Antiinflammatory Drugs, Proc. Natl. Acad. Sci., USA 77:308, 1980 and D.P. Wallach and V.R. Brown, a novel Preparation of Human Platlet Lipoxxygenase, Biochem. Biophys. Acta. 663:361, 1981. Buffy coat prepared above is resuspended in 5-7 volumes of cold 0.1 M Tris buffer, pH 7.4 containing 0.154 M NaCl. The suspension is centrifuged at 13,300 g for ten minutes at 4°C. The resultant pellet was retained, resuspended in five volumes of cold acetone, recentrifuged at 13,300 g and resuspended in five volumes of cold pentane. The pentane suspension is centrifuged for ten minutes at

13,300 g to give a pellet which is dried in the cold under vacuum with periodic pulverization. The dry powder is stable for several weeks when stored at -88°C.

Enzyme stock solution is prepared in the following manner. About 15 mg of the acetone-pentane powder is suspended in 4 ml of cold tris buffer (0.1 M, pH 7.4), allowed to stand for five minutes, and homogenized thoroughly. The homogenate is sonicated three times for 15 seconds each time, diluted to 7 ml with cold tris buffer (0.1 M, pH 7.4), and centrifuged at 4°C for 60 minutes at 13,300 g. The supernatant is retained and diluted to a total of 10 ml with cold tris buffer (0.1 M, pH 7.4) to give the stock enzyme solution. Additional dilutions of 2-50 fold are done as necessary to locate optimal enzyme reaction rate in the assay described below.

Substrate solution is prepared at 100 μ M or 1.0 μ M concentrations of arachidonic acid or linoleic acid in 0.1 M tris buffer, pH 9.0 containing 20% ethanol.

The enzyme reaction is followed spectrophotometrically by the appearance of a conjugated diene product at 234 nm. The reaction is monitored at 24°C using a Gilford Model 2600 spectrophotometer. Each assay had a total volume of 1.0 ml and contained substrate, tris buffer (0.1 M, pH 9.0), 2% ethanol, and sufficient enzyme to give an easily measurable initial rate of reaction. The effects of inhibitors on the reaction are compared with control reactions run under identical conditions. Routinely, each compound of the present invention is incubated with the enzyme for five minutes prior to addition of substrate to initiate the reaction. Inhibition expressed as IC₅₀ as molar concentration of the compound required to reduce reaction rate to 50% control.

To Evaluate the Effect of Each Compounds
as a 5-Lipoxygenase Inhibitor in Comparison to
Standard Reference Agents in Human Leukocytes (5LOA1)

5 The purpose of this assay is to evaluate the
activity of each compound as an inhibitor of human
leukocyte 5-lipoxygenase.

Arachidonic acid and calcium ionophore A23187
are obtained from Sigma (St. Louis, MO). Silica
gel plates, GF are obtained from Analtech (Newark, DE).
10 Arachidonic acid, (1-¹⁴C) and 5-HETE (³H), 5 (S)-
hydroxy-6-trans, 8,11,14-cis eicosatetraenoic acid,
are obtained from New England Nuclear (Boston, MA).
Six percent Dextran-70 in 0.9% NaCl is obtained from
Cutter Labs (Berkeley, CA).

15 Preparation of Leukocytes

Fresh blood from normal adult men who had not
received any drugs for at least the previous five
days is obtained by the Community Research Clinic
(WL/PD) using venipuncture and collected into
20 heparinized vacutainer tubes. To every 100 ml of
pooled blood is added 25 ml of dextran solution
(6% dextran -70 in 0.9% sodium chloride containing 3%
dextrose) and this is mixed gently in a plastic
cylinder. The mixture is left to stand at room
25 temperature for at least 90 minutes. The upper
layer which is rich in leukocytes and platelets is
then carefully decanted into 50 ml plastic tubes and
centrifuged at about 100 x g for eight minutes in
an IEC centrifuge and rotor number 269 (about ,
30 600 rpm). The supernatant fluid is discarded and
the pellet is resuspended in 10 ml of 0.87%
ammonium chloride for exactly two minutes. This
procedure is to lyse completely contaminating red
blood cells. Leukocytes are then separated by

centrifugation for ten minutes. The pellet is washed three times by suspension in 20 ml PBS (sodium chloride, 7.1 g; Na_2HPO_4 , 1.15 g; KH_2PO_4 , 0.2 g, and KCl , 0.2 g/L) and centrifuged as before. The final pellet is suspended in PBS containing 0.87 mM CaCl_2 . Viability of the cells is then checked using trypan blue exclusion method and is found to be over 90%.

5-Lipoxygenase Enzyme Assay

10 Leukocyte cells in suspension (0.98 ml) are incubated with or without test compounds for five minutes at 37°C in a shaking water bath. At this time a 17 μl mixture is prepared per 1 ml of cell suspension: 100 mM arachidonic acid, 1 μl ,
15 0.05 μCi ^{14}C -arachidonic acid in 5 μl ; 1 mM calcium ionophore A23187, 10 μl (1). This mixture is added and the incubation continued for five minutes. The reaction is stopped by adding four volumes of absolute ethanol and the mixture is kept in
20 ice for 30 minutes. The flocculated precipitate is separated by centrifugation at about 37,000 x g for 20 minutes (Beckman Instruments rotor number 40). The alcohol extract is taken to dryness under a stream of nitrogen and the residue is dissolved in 100-200 μl
25 absolute ethanol. At the time any turbidity is removed by centrifugation. An aliquot (25-50 μl) is applied onto 20 x 20 cm silica gel TLC plate and developed using the following solvent system: diethyl ether, petroleum ether (20:40°C), acetic acid (50:50:
30 1 v/v). Zones of 1 cm apart are scraped from the TLC plate and transferred to mini-vials. Methanol (0.5 ml) is added to dissolve the radioactivity adsorbed to the silica gel and scintillation fluid (H.P., Beckman), 5 ml is then added and vials are
35 counted in a liquid scintillation counter. A sample

of ^3H -5-HETE is applied and used for the identification of the formed 5-HETE. Total radioactivity in the test as well as the control samples are normalized and the amount of 5-HETE present is calculated accordingly.

IC₅₀ values are defined as the concentrations of test agents which caused a 50% inhibition of the formation of 5-HETE as compared to control and are determined by inspection of the concentration-response curves.

5-Lipoxygenase Assay Using Isolated
Human Leukocytes (5LOA₂)

The formation of 5-HETE in human leukocytes is considered a measure of 5-lipoxygenase activity. The protocol is described in the following.

Fresh heparinized or EDTA treated human blood is mixed with 6% dextran-3% dextrose in isotonic saline in the ratio 0.25 ml dextran solution per 1.0 ml blood. After mixing the blood is allowed to sit at room temperature for about 90 minutes while the RBC's settle. During this period, the plasma is removed with a plastic pipette to nalgens tubes.

The plasma is centrifuged at 800 rpm (125 kg) on the Beckman Td-b refrigerated centrifuge to remove the platelets (which remain in the supernatant). The pellet, consisting of leukocytes and erythrocytes, is treated with 10 ml 0.87% ammonium chloride at room temperature for four minutes, lysing the red cells. At the end of four minutes the cells are diluted with a 2x volume of phosphate buffered saline, pH 7.4, and centrifuged for ten minutes. The cells are washed three times with the phosphate buffered saline. Any of the pelleted cell matter which is not easily resuspended is discarded during the washings - the material contains platelets (12-lipoxygenase activity).

After washing, the cells are resuspended in phosphate buffered saline containing 1.0 mM calcium and 0.5 mM magnesium. After counting the cells are diluted to $1.5-2.0 \times 10^7$ leukocytes per milliliter.

5 To each polypropylene reaction tube is added 0.48 ml leukocytes in Ca-Mg phosphate buffered saline, pH 7.4; 1-5 μ l test compound dissolved in DMSO and buffer; or DMSO for control tubes.

The tubes preincubate at 37°C for five minutes.

10 The reaction is started by adding 20 μ l of the following, 0.5 μ l 20 mM arachidonic acid - final concentration = 20 μ M; 1 μ l 5 mM calcium ionophore A23187 - final concentration = 10 μ M; and 18.5 μ l buffer.

15 The reaction proceeds for five minutes, then is stopped by adding 0.5 ml 0.5 mM ice cold Tris buffer, pH 8.0. The tubes are chilled on ice for ten minutes and then extracted three times with a total of 3.5 ml ethyl acetate (3.0 ml removed).

20 The tubes can be stored at this point. For extended storage, the tubes should be filled with nitrogen.

The ethyl acetate is evaporated with a Sorvall Speed-Vac. The residue is dissolved in ethanol. The
25 tubes can also be stored at this point at -20°C under nitrogen.

A portion of the ethanol solution is injected into the HPLC system for 5-HETE quantitation.

30 The HPLC system consists of Hewlett-Packard 1040A UV spectrophotometry system with an HP85 computer. Injections are made automatically with a Waters WISP 710B. The pump is a Spectra Physics SP8700. Peaks are measured with a Hewlett Packard 3390A integrator. An RP C-18 column is used. The
35 solvent system is isocratic; the solvent is 70% methanol and 30% 0.01M sodium acetate, pH 5.7, pumped

at 1.0 ml/min. The flow is monitored at 235 nm for 5-HETE quantitation. Using a 15 cm Alltech Nucleosil C-18 5 μ M column provides for a sample turnaround time of about 16 minutes.

- 5 IC_{50} is calculated as the amount of test agent that causes 50% inhibition of the formation of 5-HETE relative to the control.

When tested by the above described procedures and shown by the notations of the acronym for each test, various compounds of the Formula I as defined above indicated activity at the highest dose tested as shown in Table 6.

TABLE 6

15		Concentration (M)	% Inhibition
	Example 49		
	5LOA	5.00 E* -6	11.6
		2.00 E -5	17.9
	LDAH	2.50 E -5	0.0
	Example 52		
20	5LOA	5.00 E -6	50.0 IC_{50}
	LDAH	1.84 E -7	50.0 IC_{50}
	Example 53		
	5LOA	3.38 E -6	50.0 IC_{50}
	LDAH	1.70 E -5	50.0 IC_{50}
25	Example 48		
	5LOA	4.20 E -6	50.0 IC_{50}
		3.50 E -6	50.0 IC_{50}
	LDAH	2.50 E -5	0.0
	Example 47		
30	LDAH	2.50 E -5	0.0

TABLE 6 (Cont'd)

		Concentration (M)			% Inhibition	
5	Example 46					
	5LOA	5.00	E	-7	50.0	IC50
	5LOA2	1.69	E	-6	50.00	
	LDAH	1.40	E	-5	50.00	IC50
10	Example 54					
	5LOA	1.00	E	-5	6.5	
		2.00	E	-5	6.0	
		4.00	E	-5	14.5	
	LDAH	2.80	E	-7	50.0	IC50
15	Example 55					
	5LOA	7.10	E	-6	50.0	IC50
		8.17	E	-6	50.0	IC50
	LDAH	3.90	E	-6	50.0	IC50
20	Example 2					
	5LOA	1.00	E	-5	11.3	
		4.00	E	-5	13.3	
	Example 8					
	5LOA	7.10	E	-6	50.0	IC50
25	Example 7					
	5LOA	4.00	E	-5	18.9	
		1.00	E	-5	16.4	
	LDAH	2.80	E	-6	50.0	IC50
	Example 12					
	LDAH	2.50	E	-5	0.0	

TABLE 6 (Cont'd)

		Concentration (M)			% Inhibition
5	Example 13				
	5LOA	1.00	E -4		+1.3
		1.00	E -4		+12.2
	LDAH	2.50	E -5		0.0
10	Example 36				
	LDAH	2.50	E -5		0.0
	Example 37				
	LDAH	5.45	E -7		50.0 IC50
15	Example 34				
	5LOA	3.92	E -5		50.0 IC50
	Example 38				
	LDAH	2.50	E -5		0.0
20	Example 9				
	5LOA	5.00	E -6		6.5
		1.00	E -5		20.9
25		2.00	E -5		21.3
		1.00	E -4		+9.6
		1.00	E -4		+3.3
	Example 45				
	5LOA	5.00	E -6		32.9
		2.00	E -5		46.0
	Example 56				
	5LOA	5.00	E -6		5.0
		2.00	E -5		13.3
		5.00	E -6		+18.4
		1.50	E -5		+20.3
		2.50	E -5		+2.5

TABLE 6 (Cont'd)

		Concentration (M)			% Inhibition
5	Example 57				
	5LOA	5.00	E -6		4.6
		1.00	E -5		7.2
		2.00	E -5		5.0
10	Example 35				
	5LOA	5.00	E -6		1.7
		2.00	E -5		6.0
15	Example 21				
	5LOA	2.08	E -5		50.0 IC50
	Example 22				
	LDAH	6.80	E -7		50.0 IC50
20	Example 60				
	LDAH	1.20	E -4		50.0 IC50
	Example 23				
	LDAH	4.10	E -6		50.0 IC50
	Example 25				
	LDAH	8.00	E -5		0.0
	Example 26				
	5LOA	5.00	E -6		0.6
		2.00	E -5		7.6
	LDAH	2.40	E -5		50.0 IC50
	Example 32				
	5LOA	1.00	E -4		16.4
		1.00	E -4		11.5

TABLE 6 (Cont'd)

	Concentration (M)	% Inhibition
Example 43		
5LOA	5.30 E -6	50.0 IC50
Example 51		
5LOA	1.11 E -5	50.0 IC50
Example 42		
5LOA2	5.00 E -6	21.1
	2.00 E -5	30.9

*The notation E -number means "X 10^{-no.}".

Accordingly, the present invention also includes a pharmaceutical composition for treating one of the above diseases or conditions comprising an antidisease or anticondition effective amount of a compound of the Formula I as defined above together with a pharmaceutically acceptable carrier.

The present invention further includes a method for treating one of the above named diseases or conditions in mammals, including man, suffering therefrom comprising administering to such mammals either orally or parenterally a corresponding pharmaceutical composition containing a compound of Formula I as defined above in appropriate unit dosage form.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories. A solid carrier can be one or more substances which may also act as

diluents, flavoring agents, solublizers, lubricants, suspending agents, binders or tablet disintegrating agents; it can also be encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active compound. In the tablet the active compound is mixed with carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from 5 or 10 to about 70 percent of the active ingredient. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component (with or without other carriers) is surrounded by carrier, which is thus in association with it. Similarly, cachets are included. Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides or cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions. As an example may be mentioned water or water propylene glycol solutions for parenteral injection. Liquid preparations can also be formulated in solution in aqueous polyethylene glycol solution. Aqueous solutions suitable for oral

use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by
5 dispersing the finely divided active component in water with viscous material, i.e., natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

10 Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions, and emulsions. These particular solid
15 form preparations are most conveniently provided in unit dose form and as such are used to provide a single liquid dosage unit. Alternately, sufficient solid may be provided so that after conversion to liquid form, multiple individual liquid doses may be
20 obtained by measuring predetermined volumes of the liquid form preparation as with a syringe, teaspoon, or other volumetric container. When multiple liquid doses are so prepared, it is preferred to maintain the unused portion of said liquid doses at low temperature
25 (i.e., under refrigeration) in order to retard possible decomposition. The solid form preparations intended to be converted to liquid form may contain, in addition to the active material, flavorants, colorants, stabilizers, buffers, artificial and
30 natural sweeteners, dispersants, thickeners, solubilizing agents, and the like. The liquid utilized for preparing the liquid form preparation may be water, isotonic water, ethanol, glycerine, propylene glycol, and the like as well as mixtures
35 thereof. Naturally, the liquid utilized will be chosen with regard to the route of administration, for

example, liquid preparations containing large amounts of ethanol are not suitable for parenteral use.


Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself or it can be the appropriate number of any of these in packaged form.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from 10 mg to 2 g preferably to 10 to 500 mg according to the particular application and the potency of the active ingredient. The compositions can, if desired, also contain other compatible therapeutic agents.

In therapeutic use as described above, the dosages may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed.


Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

FORMULA




I


SCHEME I



or




wherein $y = 1$




wherein $y = 2$

-61-


SCHEME II




SCHEME III



SCHEME IV



SCHEME V




wherein $y = 2$

The word "Celite" as used herein is an Australian Registered Trade Mark.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A pharmaceutical composition comprising a therapeutically effective amount of a compound having the Formula 1



I

and pharmaceutically acceptable salts thereof, wherein y is one or two; b is zero, one, two, three, or four; R_1 is selected from a group consisting of alkyl of from one to four carbons, inclusive; alkoxy of from one to four carbons, inclusive; thioalkoxy of from one to four carbon atoms, inclusive; carboalkoxy of from two to four carbons, inclusive; alkanoyl of from one to four carbons; phenyl; hydroxy; halogen; hydrogen; nitro; amino; mono- and di-alkylamino having each alkyl the same or different and of from one to four carbons, inclusive; carbalkoxyamido, of from one to four carbons, inclusive; alkylsulfinyl of from one to four carbons, inclusive; alkylsulfonyl of from one to four carbons, inclusive; and when b is one, the R_1 may be $-(CH=CH-CH=CH)-$ taken together with an adjacent ring carbon to form a benzo radical; R_5 is hydrogen; halogen; alkyl of from one to four carbons, inclusive; alkoxy of from one to four carbons, inclusive; carboalkoxy



of from two to four carbons, inclusive; hydroxy; or
-(CH=CH-CH=CH)- taken together with adjacent carbons to
form a benzo radical; R_6 is alkyl of from six to
twenty carbons, $-CH=CH-R_4$, $-(CH_2)_nCOR_4$, or
 $-(CH_2)_n-R_4$ wherein n is zero to four, inclusive;
and R_4 is phenyl, unsubstituted or substituted at the
two through six positions by carboalkoxy, having alkoxy of
from one to four carbons, inclusive; alkyl of from one to
four carbons; alkoxy, or thioalkoxy of from one to four
carbons, inclusive; phenalkoxy of from one to four
carbons, inclusive, in the alkoxy group; amino, monoalkyl
and dialkyl amino having the alkyl of from one to four
carbons, inclusive; alkanoylamino of from two to six
carbons, inclusive; carboxyl; benzo; halogen; hydroxy;
hydroxyalkyl of from one to four carbons, inclusive;
alkanoyl of from one to four carbons, inclusive; nitro,
alkanesulfonamido of from one to four carbons, inclusive;
X is hydrogen or lower alkyl of from one to four carbons,
inclusive, together with a pharmaceutically acceptable
carrier.

2. A pharmaceutical composition according to Claim 1,
wherein in Formula I, y is one or two; b is zero, one, or
two; R_1 is hydrogen, halogen, nitro, hydroxy, amino,
phenyl, straight or branched chained alkyl of from one to
four carbons, inclusive; alkoxy of from one to four
carbons, inclusive, or -(CH=CH-CH=CH)- taken together with



adjacent ring carbons to form a benzo radical; R_5 is hydrogen or halogen; R_6 is halogen; alkyl of from six to twenty carbons; or $-(CH_2)_n-R_4$ wherein n is two; and R_4 is phenyl, unsubstituted or substituted at the two through six positions by halogen; hydroxy; phenyl; alkoxy of from one to four carbons, inclusive; carboxyl; benzyloxy; X is hydrogen or lower alkyl of from one to four carbons, inclusive.

10 3. A pharmaceutical composition according to Claim 1, wherein in Formula I, R_1 is hydrogen, y is 1, R_5 is H, and R_6 is alkyl of from 6 to 20 carbons, inclusive, or $-(CH_2)_n-R_4$ wherein n is two and R_4 is phenyl optionally substituted by carboxyl, carboalkoxy of from one to four carbons, inclusive, chloro, alkoxy of from one to four carbons, inclusive, hydroxy, or phenyl.

4. A pharmaceutical composition according to Claim 1, wherein in Formula I, R_1 is hydrogen, y is 2, R_5 is hydrogen or the benzo radical; R_6 is alkyl of from 6 to 20 carbons, inclusive, or $-(CH_2)_n-R_4$ wherein n is 2 and R_4 is phenyl optionally substituted by lower alkoxy carbonyl; carboxyl, carboalkoxy wherein the alkoxy is from one to four carbons, inclusive, alkoxy of from one to four carbons, inclusive, hydroxy.

5. A pharmaceutical composition according to Claim 1,



wherein the compound is a member of the group selected of

N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy- α -oxo-benzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy- α -oxo-benzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-benzamide.


N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methylbenzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-3-(1,1-dimethylethyl)-2-hydroxy- α -oxobenzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxybenzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-N,4-dimethylbenzamide.

6. A compound having the general Formula I




and pharmaceutically acceptable salts thereof, wherein y is one or two; b is zero, one, two, three, or four; R₁ is selected from a group consisting of alkyl of from one to four carbons, inclusive; alkoxy of from one to four carbons, inclusive; thioalkoxy of from one to four carbon atoms, inclusive; carboalkoxy of from two to four carbons, inclusive; alkanoyl of from one to four carbons; phenyl;



hydroxy; halogen; hydrogen; nitro; amino; mono- and di-
10 alkylamino having each alkyl the same or different and of
from one to four carbons, inclusive; carbalkoxyamido, of
from one to four carbons, inclusive; alkylsulfinyl of from
one to four carbons, inclusive; alkylsulfonyl of from one
to four carbons, inclusive; and when b is one, then R_1
may be $-(CH=CH-CH=CH)-$ taken together with an adjacent
ring carbon to form a benzo radical; R_5 is hydrogen;
halogen; alkyl of from one to four carbons, inclusive;
alkoxy of from one to four carbons, inclusive; carboalkoxy
of from two to four carbons, inclusive; hydroxy; halogen;
20 or $-(CH=CH-CH=CH)-$ taken together with adjacent carbons to
form a benzo radical; R_6 is alkyl of from six to twenty
carbons, $-CH=CH-R_4$, $-(CH_2)_nCOR_4$, or
 $-(CH_2)_n-R_4$ wherein n is zero to four, inclusive;
and R_4 is phenyl, unsubstituted or substituted at the
two through six positions by carboalkoxy, having alkoxy of
from one to four carbons, inclusive; alkyl of from one to
four carbons; alkoxy, or thioalkoxy of from one to four
carbons, inclusive; phenalkoxy of from one to four carbons
inclusive in the alkoxy group; amino, monoalkyl and
dialkyl amino having the alkyl of from one to four
carbons, inclusive; alkanoylamino of from two to six
carbons, inclusive; carboxyl; benzo; halogen; hydroxy;
hydroxyalkyl of from



one to four carbons, inclusive; alkanoyl of from one to four carbons, inclusive; nitro, alkanesulfonamido of from one to four carbons, inclusive; X is hydrogen or lower alkyl of from one to four carbons, inclusive, with the proviso that, when R_1 is halogen and b is one or two or when b is zero then R_6 cannot be alkyl of from six to twenty carbons or cannot be



optionally substituted by alkyl of one to four carbons when R_5 is hydroxy.

7. A compound of claim 6 wherein in Formula I, R_1 is hydrogen, y is 1, R_5 is H, and R_6 is alkyl of from 6 to 20 carbons, inclusive, or $-(CH_2)_nR_4$ wherein n is two and R_4 is phenyl optionally substituted by carboxyl, carboalkoxy of from one to four carbons, inclusive, chloro, alkoxy of from one to four carbons, inclusive, hydroxy, or phenyl; or the pharmaceutically acceptable acid or base addition salts.

8. A compound of claim 6 wherein in Formula I, R_1 is hydrogen, y is 2, R_5 is hydrogen or the benzo radical; R_6 is alkyl of from 6 to 20 carbons, inclusive, or $-(CH_2)_nR_4$ wherein n is 2 and R_4 is phenyl optionally substituted by carboxyl, carboalkoxy wherein the alkoxy is from one to four carbons, inclusive, alkoxy of from one to



four carbons, inclusive, hydroxy, or pharmaceutically acceptable acid or base addition salts.

9. A compound of Claim 6 being a member of the group consisting of

N-[4-[2-(3,4-dimethoxyphenyl)ethyl]phenyl]-2-hydroxy- α -oxo-benzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy- α -oxo-benzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-benzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methylbenzamide.


N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-3-(1,1-dimethylethyl)-2-hydroxy- α -oxobenzeneacetamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-4-methoxybenzamide.

N-[4-[2-(3,4-dihydroxyphenyl)ethyl]phenyl]-2-hydroxy-N,4-dimethylbenzamide.

10


10. A process for the preparation of compounds according to Claims 6 to 9 which comprises reacting a compound of general Formula II



II



wherein R_1 , b, and y are as defined above and R is lower alkyl or phenyl; with a compound of general formula III



III

wherein X, R₅, and R₆ are as defined above, to obtain the compound having the Formula I as defined above.

11. A method of treating allergies or immuno-inflammatory conditions in a mammal suffering therefrom which comprises administering to such mammals a compound as defined by formula I, in claim 1.

12. A compound as defined in Claim 6 substantially as herein described with reference to the Examples.

DATED this 6th day of NOVEMBER, 1990

WARNER-LAMBERT COMPANY

Attorney: WILLIAM S. LLOYD

Fellow Institute of Patent Attorneys of Australia
of SHELSTON WATERS

