

FASCICULE DE BREVET D'INVENTION

A61K 31/47 (2018.01);
A61K 47/02 (2018.01);
A61K 47/12 (2018.01);
A61K 47/26 (2018.01);
A61K 9/10 (2018.01);
A61K 9/14 (2018.01);
A61P 31/06 (2018.01);
A61P 31/08 (2018.01)

21 Numéro de dépôt : 1202200532
PCT/EP2021/068956

22 Date de dépôt : 08/07/2021

30 Priorité(s) :
EP n° 20185105.2 du 09/07/2020

24 Délivré le : 04/07/2023

45 Publié le : 09.10.2023

73 Titulaire(s) :
JANSSEN PHARMACEUTICA NV,
Turnhoutseweg 30,
2340 BEERSE (BE)

72 Inventeur(s) :
HOLM, René (BE);
VERVOORT, Iwan, Caroline, F (BE);
DONG, Wenyu (BE);
COLOMBO, Miriam (BE)


74 Mandataire : S.C.P AKKUM, AKKUM & Associates,
n° 1777, Rue 6.261, Auditorium Jean Paul II, Quartier
Mbankolo, B.P. 4966, YAOUNDE (CM).

54 Titre : Long-Acting Formulations.

57 Abrégé :

This invention concerns pharmaceutical compositions for administration via intramuscular or subcutaneous injection, comprising micro- or nanoparticles of the anti-TB compound bedaquiline, suspended in an aqueous pharmaceutically acceptable carrier, and comprising PEG4000 as a surface modifier, and the use of such pharmaceutical compositions in the treatment and prophylaxis of a pathogenic mycobacterial infection.

Fig. 1



Long-acting formulations

Field of the Invention

This invention concerns pharmaceutical compositions for administration via intramuscular or subcutaneous injection, comprising micro- or nanoparticles of the ATP synthase inhibitor compound, bedaquiline (marketed as Sirturo®, where bedaquiline is in the form of its fumarate salt), suspended in an aqueous pharmaceutically acceptable carrier, and the use of such pharmaceutical compositions in the treatment of bacterial infections, e.g. tuberculosis and the like.

Background of the Invention

Bedaquiline is a known anti-tuberculosis drug used in various combinations. It may be formulated in the form of a pharmaceutically acceptable salt, such as in the form of bedaquiline fumarate, marketed as Sirturo®. It is thought to act as an ATP synthase inhibitor, possessing a selectivity index of more than 20000 for mycobacterial ATP synthase versus eukaryotic mitochondrial ATP synthase.

Bedaquiline has already been reported as being useful in the treatment of mycobacterial infections, as well as being useful in killing dormant, latent, persistent mycobacteria, in particular *Mycobacterium tuberculosis*, and can consequently be used to treat latent TB. Such use of bedaquiline has been described in several publications including international patent documents WO 2004/011436 and WO 2006/067048. It is also known that bedaquiline is bactericidal against mycobacterium leprae, for example as described in "Bacterial Activities of R207910 and other Antimicrobial Agents against *Mycobacterium leprae* in Mice", Antimicrobial agents and Chemotherapy, April 2006, p 1558, and "The Diarylquinolone R207910 is Bactericidal against *Mycobacterium leprae* in mice and at Low Dose Administered Intermittently", Antimicrobial agents and Chemotherapy, Sept 2009, p3989.

The goal of long-acting formulations can be to reduce drug burden. This is particularly useful for treatment regimens that may last several months.

The number and/or volume of dosage forms that need to be administered are commonly referred to as "pill burden". A high pill burden is undesirable for many reasons, such as the frequency of intake, often combined with the inconvenience of having to swallow large dosage forms, as well as the need to store and transport a large number or volume of pills. A high pill burden increases the risk of patients not taking their entire dose, thereby failing to comply with the prescribed dosage regimen. As well as reducing the

effectiveness of the treatment, this may also lead to the emergence of resistance (e.g. in the case of bedaquiline, bacterial resistance).

5 It would be attractive to provide therapy involving the administration of dosage forms at long time intervals such as one week or longer, or even one month or longer.

Various formulations are known in the art, including long-acting ones. For instance, micro- and nano-suspension technology is known for achieving long-acting formulations in the field of anti-HIV drugs, for instance as described in international patent applications WO 2007/147882 and WO 2012/140220. Further, nanoparticles
10 known in the prior art have been described, for example, in EP-A-0 499 299. Such particles have an average particle size in the submicron range and consist of particles of a crystalline drug substance having a surface modifier adsorbed on their surface. Nanoparticles have also been used to formulate poorly water-soluble active ingredients.

15 Long-acting formulations of the anti-tuberculosis drug bedaquiline are also described in international patent application WO 2019/012100.

The importance of long-acting formulations relates to the intermittent administration of
20 these micro- or nanoparticle formulations at time intervals of one week or longer that result in plasma levels that may be sufficient to suppress the growth of the mycobacterial infection. This allows for a reduced number of administrations thereby being beneficial in terms of pill burden and drug compliance of the patient. Micro- or nanoparticle formulations of bedaquiline therefore may be useful in the long-term
25 treatment of mycobacterial infections (e.g. tuberculosis, including latent tuberculosis, and leprosy).

The intermittent administration of micro- or nanoparticle formulations of bedaquiline at time intervals of one week or longer furthermore results in plasma levels that may be
30 sufficient to provide prevention against transmission of mycobacterial infection. Also in this instance, a reduced number of administrations is required, which again is advantageous in terms of pill burden and drug compliance of the individual at risk of being infected.

35 A challenge relating to the manufacture and suitability of such long-acting formulations relates to fact that they have to be sterilized (which is important for injectables, for instance if they are intended to be administered intravenously, intramuscularly or subcutaneously). There are a number of different ways to sterilize such long-acting

formulations, including by heat sterilization, autoclaving and gamma-radiation (γ -radiation). An example of some methods are described in e.g. US patents/applications US 5,298,262, US 5,346,702 and US 2010/255102. For heat sterilization and autoclaving, it is important to be able to select excipients (e.g. surface modifiers or surfactants) that are autoclavable, e.g. do not degrade. Further challenges arise after such sterilization, which are linked to desired stability of the long-acting formulation, undesired aggregation of particles of the active pharmaceutical ingredient (API) within that formulation and the desired re-suspendability of the formulation (after sterilization, e.g. autoclaving). US patents US 5,298,262 and US 5,346,702 disclose the use of cloud point modifiers to prevent particle aggregation during sterilization. The cloud point is the temperature above which the surfactant (or surface modifier) phase-separates and precipitates out of the solution. Heat sterilization or autoclaving of suspensions must be performed below the cloud point of the surfactant / surface modifier as otherwise they would phase-separate and precipitate when heated above their cloud temperature due to a solubility change. This would leave the particle (of the active pharmaceutical ingredient) surface free and the particles would thereby aggregate. The idea of a cloud point modifier (or booster) is to allow the temperature of the sterilization or autoclaving process to be higher and thereby preventing or limiting particle aggregation. The cloud point modifiers mentioned in US 5,298,262 and US 5,346,702 include ionic and non-ionic cloud point modifiers, such as sodium dodecyl sulfate, dodecyltrimethylammonium bromide, polyethylene glycol and propylene glycol. The polyethylene glycols mentioned as cloud point modifiers include PEG300, PEG400, PEG1000 and PEG2000, with PEG400 indicated as being preferred, and in the examples specifically PEG400 and PEG1000 were shown to raise cloud point (of Tetronic 908). Other cloud point modifiers or boosters are also described in a number of other documents.

Now further alternative and/or improved long acting formulations are described, and the invention relates to such formulations.

30 Summary of the Invention

The present invention is concerned with a pharmaceutical composition for administration by intramuscular or subcutaneous injection, comprising a therapeutically effective amount of bedaquiline, or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nanoparticles comprising:

- 35 (a) bedaquiline, or a pharmaceutically acceptable salt thereof, in micro- or nanoparticle form, and a surface modifier; and
(b) a pharmaceutically acceptable aqueous carrier,
which is characterised in that the surface modifier comprises PEG4000 or the like,

wherein such a composition may be referred to herein as “composition(s) of the invention”.

PEG4000, or, polyethylene glycol 4000, is a known high-molecular weight polymer where the 4000 refers to the approximate average molecular weight in daltons. PEG4000 is commercially available from sources such as Sigma-Aldrich and hence why it is used as such. However, embraced within the scope of the invention (e.g. when the term “PEG4000” or “PEG4000, or the like” is used) are other high-molecular weight polyethylene glycols, for instance those above 1000 and up to 8000 (e.g. PEG1000 to PEG8000, for instance PEG2000 to PEG6000), even though in a particular embodiment the PEG group when referred to herein in the context of the invention is PEG3000 to PEG5000 (e.g. PEG3500 to PEG4500). As indicated herein, the number next to the PEG represents average molecule weight in daltons, as it is understood that most PEGs include molecules with a distribution of molecular weights, i.e. they are polydisperse.

The composition of the invention is a suspension, by which we mean that the bedaquiline active ingredient is suspended in the pharmaceutically acceptable aqueous carrier.

The composition of the invention (i.e. the suspension) contains a surface modifier, which may be adsorbed onto the surface of the active ingredient bedaquiline. As indicated, the surface modifier comprises PEG4000, or the like (and may also contain other surface modifiers, such as those described hereinafter).

In an embodiment, the present invention may therefore concern a pharmaceutical composition for administration by intramuscular or subcutaneous injection, comprising a therapeutically effective amount of bedaquiline, or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nanoparticles comprising:

- (a) bedaquiline, or a pharmaceutically acceptable salt thereof, in micro- or nanoparticle form, having a surface modifier adsorbed to the surface thereof; and
 - (b) a pharmaceutically acceptable aqueous carrier; wherein the bedaquiline active ingredient is suspended,
- and wherein the surface modifier comprises PEG4000, or the like.

The invention further concerns a method of treating a subject infected with pathogenic mycobacteria such as *Mycobacterium tuberculosis*, *M. bovis*, *M. leprae*, *M. avium* and *M. marinum*. In an embodiment, the mycobacteria is *Mycobacterium tuberculosis*

(including the latent or dormant form) or *Mycobacterium leprae*. The compositions of the invention may be particularly suitable for the treatment of *Mycobacterium leprae* and the latent or dormant form of *Mycobacterium tuberculosis*. This is because for treating these specific infections, a lower concentration of bedaquiline in the plasma may be effective against such infection, for instance as described in Antimicrobial Agents and Chemotherapy, Sept 2009, p. 3989-3991 by Robert Gelber, Koen Andries *et al* (the contents of which are hereby incorporated by reference, and wherein, essentially, it is reported that low and intermittent dosing with bedaquiline holds promise for leprosy patients; whereas minimal dose killing 99% of bacilli for *M. tuberculosis* is 30 mg/kg/wk, for *M. lepra* it is < 5.0 mg/kg/wk, and hence dosing once a month may be as efficient as 5 days a week; other publications of the effect of bedaquiline on *Mycobacterium leprae* in mice include Antimicrobial Agents and Chemotherapy, April 2006, p. 1558-1560 by Baohong Ji, Koen Andries *et al* – the contents of which are also hereby incorporated by reference). Hence, the compositions of the invention may be particularly suitable in a method of treating a subject infected with *Mycobacterium leprae* or the latent/dormant form of *Mycobacterium tuberculosis*. Such methods of treating a subject infected with pathogenic mycobacteria comprise the administration, by intramuscular or subcutaneous injection, of a therapeutically effective amount of a pharmaceutical composition as specified above or hereinafter. Or, alternatively, the invention concerns the use of a pharmaceutical composition as specified above or hereinafter, for the manufacture of a medicament for treating pathogenic mycobacteria infection (or for using such medicament in a particular treatment regime as described herein). In one embodiment, the composition is for the long-term treatment of pathogenic mycobacteria infection. In an embodiment, the pathogenic mycobacterial infection may such as described above or hereinafter, such as an infection that requires long-term treatment (in a further embodiment, an infection that further may be treated at relatively low plasma concentration levels of bedaquiline or its active metabolite, for instance latent/dormant *Mycobacterium tuberculosis* or, in a particular embodiment, *Mycobacterium leprae*).

The invention further concerns a method of treating a subject infected with pathogenic mycobacteria such as *Mycobacterium tuberculosis*, and by this we also include multi-drug resistant tuberculosis. The term “drug resistant” (DR) is a term well understood by the person skilled in microbiology. A drug resistant *Mycobacterium* is a *Mycobacterium* which is no longer susceptible to at least one previously effective drug; which has developed the ability to withstand antibiotic attack by at least one previously effective drug. A drug resistant strain may relay that ability to withstand to its progeny. Said resistance may be due to random genetic mutations in the bacterial cell that alters

its sensitivity to a single drug or to different drugs. Multi-drug resistant (MDR) tuberculosis is a specific form of drug resistant tuberculosis due to a bacterium resistant to at least isoniazid and rifampicin (with or without resistance to other drugs), which are at present the two most powerful anti-TB drugs. Thus, whenever used hereinbefore or hereinafter “drug resistant” includes multi drug resistant. The compositions of the invention are also useful for the treatment of MDR-TB.

In another aspect, there is provided a method for the long term treatment of a subject infected with pathogenic mycobacteria such as *Mycobacterium tuberculosis*, *M. bovis*, *M. leprae*, *M. avium* and *M. marinum*, said method comprising the administration of an effective amount of a pharmaceutical composition as specified above or hereinafter, for administration by intramuscular or subcutaneous injection; wherein the composition is administered or is to be administered intermittently at a time interval that is in the range of one week to one year, or one week to two years. Or, alternatively, the invention concerns the use of a pharmaceutical composition as specified above or hereinafter, for the manufacture of a medicament for the long term treatment of a subject infected with pathogenic mycobacteria such as *Mycobacterium tuberculosis*, *M. bovis*, *M. leprae*, *M. avium* and *M. marinum*, for administration by intramuscular or subcutaneous injection, wherein the composition is administered or is to be administered intermittently at a time interval that is in the range of one week to one year, or one week to two years. Hence, it will be understood that the term “long term treatment” refers to treatment where one dose or one administration (e.g. by intramuscular or subcutaneous injection) will have a persistent therapeutic effect over a time period, as described herein, for instance a persistent therapeutic effect over several hours, weeks or months (e.g. in an embodiment, over a period of at least or up to one month, three months or six months); see examples. Put another way, long term treatment may refer to, where there is more than one dose/administration, the long period of time (as described herein) between the doses/administrations, i.e. the intervals are a long period of time as described herein.

In another aspect, there is provided a method for the long term treatment of a subject infected with pathogenic mycobacteria (e.g. of any of the types as described here), as described herein (e.g. above) wherein one dose or administration (e.g. of the amount described herein, e.g. hereinafter) is provided/required (and has a persistent effect, e.g. over a time period described herein). In another aspect, there is provided such a long term treatment regime, where two such doses or administrations are provided/required, which doses/administrations are given at intervals, wherein the interval time period is that as described herein, e.g. a period of at least or up to one month, three months or six

months – for instance for a period of time in which persistent therapeutic effect lasts). In a further embodiment, there is provided such a long term treatment regime, in which three such doses or administrations are provided/required at such intervals as herein described. In yet a further embodiment, there is provided a long term treatment regime
5 as herein described but which is preceded with a lead-in treatment phase (that is not a long term treatment regime, e.g. a once-daily administration course, lasting for one week, two weeks, three weeks or one month).

The invention further concerns a method for the prevention of a pathogenic
10 mycobacterial infection in a subject at risk of being infected by a pathogenic mycobacterial infection, said method comprising administering an amount, effective in preventing a pathogenic mycobacterial infection, of a pharmaceutical composition as specified above or as further specified hereinafter, to said subject. Or alternatively, the invention concerns the use of a pharmaceutical composition as specified above or as
15 further specified hereinafter for the manufacture of a medicament for the prevention of a pathogenic mycobacterial infection in a subject at risk of being infected by a pathogenic mycobacterial infection.

In another aspect the invention relates to a method for the long term prevention of a
20 pathogenic mycobacterial infection in a subject at risk of being infected by a pathogenic mycobacterial infection, said method comprising administering to said subject an effective amount of a pharmaceutical composition as specified above or as further specified hereinafter, wherein the composition is administered or is to be administered intermittently at a time interval that is in the range of one week to one
25 year, or one week to two years.

The present invention furthermore relates to the use of a pharmaceutical composition as specified above or as further specified hereinafter, for the manufacture of a medicament for the long term prevention for the long term prevention of a pathogenic mycobacterial
30 infection in a subject at risk of being infected by a pathogenic mycobacterial infection, wherein the composition is administered or is to be administered intermittently at a time interval that is in the range of one week to one year or one week to two years.

In one embodiment the invention concerns a use or a method as specified herein,
35 wherein the pharmaceutical composition is administered or is to be administered at a time interval that is in the range of one week to one month, or in the range of one month to three months, or in the range of three months to six months, or in the range of six months to twelve months, or in the range of 12 months to 24 months.

In another embodiment the invention concerns a use or a method as specified herein, wherein the pharmaceutical composition is administered or is to be administered once every two weeks, or once every month, or once every three months.

5

Further pharmaceutical compositions, methods of treatment or prevention, as well as uses for the manufacture of medicaments based on these compositions will be described hereinafter and are meant to be part of the present invention.

10 The invention is also described with reference to the following figures:

Figure 1: PSD measurements of Reference Example A, at time zero and at 1 month, where "Concept 7" refers to Reference Example A

Figure 2: PSD measurements for Reference Examples B and C under various conditions (including after autoclaving), and where Concept 3 refers to Reference

15 Example B and Concept 4 refers to Reference Example C

Figure 3: PSD of the micro-suspension of Example 1, before and after autoclaving

Figure 4: PSD of the micro-suspension of Example 1, under various conditions including after autoclaving and after further time (and at varying temperatures)

Figure 5: PSD of the micro-suspension of Example 1, under various other conditions, including up to 3 months at 60°C

20

Figure 6: "Plasma kinetics of TMC207 in male rats when administered IM or SC with 200 mg/ml micro-formulation (see Example 1, Formulation 1B i.e. the micro-suspension) at a dose of 40 mg/kg" and "Plasma kinetics of TMC207 in male rats when administered IM or SC with 200 mg/ml nano-formulation (see Example 1, Formulation 1A, i.e. the nano-suspension) at a dose of 40 mg/kg"

25

Figure 7: Plasma concentration versus time profiles of subcutaneous administered bedaquiline LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats; data represent means with SD

Figure 8: Plasma concentration versus time profiles of bedaquiline (BDQ) metabolite after subcutaneous administration of BDQ LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats; data represent means with SD

30

Figure 9: Plasma concentration versus time profiles of intramuscular administered bedaquiline LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats; data represent means with SD

Figure 10: Plasma concentration versus time profiles of bedaquiline (BDQ) metabolite after intramuscular administration of BDQ LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats; data represent means with SD

Detailed Description of the Invention

10 The compound used in the invention is the compound TMC207, also referred to as bedaquiline.

Bedaquiline can be used in its non-salt form or as a suitable pharmaceutically acceptable salt form, such as an acid addition salt form or base addition salt form. In an
15 embodiment, bedaquiline is in its non-salt form in compositions of the invention.

The pharmaceutically acceptable acid addition salts are defined to comprise the therapeutically active non-toxic acid addition salt forms which bedaquiline is able to form. Said acid addition salts can be obtained by treating the free form of bedaquiline
20 with appropriate acids, for example inorganic acids, for example hydrohalic acid, in particular hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid and phosphoric acid ; organic acids, for example acetic acid, hydroxyacetic acid, propanoic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, fumaric acid, malic acid, tartaric acid, citric acid, methanesulfonic acid, ethanesulfonic acid,
25 benzenesulfonic acid, p-toluenesulfonic acid, cyclamic acid, salicylic acid, p-aminosalicylic acid and pantoic acid. In particular, the fumarate salt is considered, given that this is the form employed in the already-marketed product Sirturo®.

Possible therapeutically active non-toxic base addition salt forms may be prepared by
30 treatment with appropriate organic and inorganic bases. Appropriate base salts forms comprise, for example, the ammonium salts, the alkaline and earth alkaline metal salts, in particular lithium, sodium, potassium, magnesium and calcium salts, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hybamine salts, and salts with amino acids, for example arginine and lysine.

35 Conversely, said acid or base addition salt forms can be converted into the free forms by treatment with an appropriate base or acid.

The term addition salt as used in the framework of this application also comprises the solvates which bedaquiline as well as the salts thereof, are able to form. Such solvates are, for example, hydrates and alcoholates.

5

Whenever reference to bedaquiline (or TMC207) is employed herein, we refer to the single stereoisomeric form that is employed in the marketed product Sirturo®, and which is disclosed in WO2004/011436 as an antimycobacterial agent.

- 10 It has been found that the physico-chemical properties of bedaquiline allow for the manufacture of micro- or nanoparticle suspensions that have unique pharmacokinetic properties in that they can be used for the long term treatment of a pathogenic mycobacterial infection as well as in the long term prevention of a pathogenic mycobacterial infection and to this purpose only a limited number of drug
- 15 administrations is required. This is beneficial in terms of pill-burden as well as patient compliance with the prescribed dose regimen.

As used herein the term “treatment of a pathogenic mycobacterial infection” relates to the treatment of a subject being infected with a pathogenic mycobacterial infection.

- 20 Such mycobacterial infection may be mycobacterium tuberculosis or multi-drug resistance mycobacterium tuberculosis.

The term “prevention of a pathogenic mycobacterial infection” relates to the prevention or avoidance of a subject becoming infected with a pathogenic mycobacterial infection.

- 25 The source of infection can be various, for instance a material containing a pathogenic mycobacterial infection.

The terms “therapeutically effective amount”, “an amount, effective in preventing a pathogenic mycobacterial infection”, and similar terms, refer to amounts, or

30 concentrations, of the compositions of the invention (or amounts/concentrations of active ingredient bedaquiline within such compositions) that result in efficacious plasma levels. With “efficacious plasma levels” it is meant those plasma levels of bedaquiline that provide effective treatment or effective prevention of a pathogenic mycobacterial infection. This is because amount/dose/administration given may be

35 linked to the desired exposure levels or desired plasma levels for the effective treatment/prevention, for instance as described herein (see e.g. the examples).

The term “subject” in particular relates to a human being.

- The term "micro- or nanoparticles" refers to particles in the micrometer or nanometer range. The size of the particles should be below a maximum size above which administration by subcutaneous or intramuscular injection becomes impaired or is even no longer possible. Said maximum size depends for example on the limitations imposed by the needle diameter or by adverse reactions of the body to large particles, or both. In one embodiment, the pharmaceutical compositions of the invention comprise bedaquiline in microparticle form. In another embodiment, the pharmaceutical compositions of the invention comprise bedaquiline in nanoparticle form.
- The average effective particle size of the micro- or nanoparticles of the present invention may be below about 50 μm , or below about 20 μm , or below about 10 μm , or below about 1000 nm, or below about 500 nm, or below about 400 nm, or below about 300 nm, or below about 200 nm. The lower limit of the average effective particle size may be low, e.g. as low as about 100 nm or as low as about 50 nm. In one embodiment, the average effective particle size is in the range of about 50 nm to about 50 μm , or about 50 nm to about 20 μm , or about 50 nm to about 10 μm , or about 50 nm to about 1000 nm, about 50 nm to about 500 nm, or about 50 nm to about 400 nm, or about 50 nm to about 300 nm, or about 50 nm to about 250 nm, or about 100 nm to about 250 nm, or about 150 nm to about 220 nm, or 100 to 200 nm, or about 150 nm to about 200 nm, e.g. about 130 nm, or about 150 nm. For instance, both after preparation and after a period of time of up to 3 months (e.g. when stored at temperatures of about 5°C, 25°C and 40°C) generally:
- the micro-suspensions may have, in an embodiment, a D90 of between about 3 and 10 μm (e.g. about 3.5, 4 or 5 μm) and a D50 of between about 2 and 4 μm (e.g. about 3 μm)
 - the nano-suspensions may have, in an embodiment, a D90 of between about 0.5 and 1.5 μm (e.g. about, or less than 1 μm or about, or less than about 1000 nm) and a D50 of between about 0.1 and 0.5 μm (e.g. about, or less than, about 0.3 μm , or less than about 300 nm).
- In an embodiment, the micro-particles are employed, wherein the average effective particle size, as measured by D10, D50 and/or D90 (in an embodiment as measured by D50) is below about 50 μm , or below about 20 μm , and above about 0.1 μm (100 nm). In an embodiment the range for such micro-particles employed in the compositions of the invention is between about 20 μm and about 0.1 μm (in a further embodiment between about 15 μm , and above about 0.2 μm (200 nm) and in a further embodiment between about 10 μm , and above 0.5 μm (500 nm), for instance between about 10 μm , and above 1 μm or above about 1000 nm, or above about 500 nm, or above about

400 nm, or above about 300 nm, or above about 200 nm. The foregoing values refer to measurements after preparation. They may also, however, in an embodiment, refer to measurements after a period of time up to 3 months (e.g. after 5 days, one week, two weeks, one month, two months or three months) and stored at various temperatures (e.g. at temperatures of about 5°C, 25°C and 40°C).

As used herein, the term average effective particle size has its conventional meaning as known to the person skilled in the art and can be measured by art-known particle size measuring techniques such as, for example, sedimentation field flow fractionation, photon correlation spectroscopy, laser diffraction or disk centrifugation. The average effective particle sizes mentioned herein may be related to volume distributions of the particles. In that instance, by "an effective average particle size of less than about 50 µm" it is meant that at least 50% of the volume of the particles has a particle size of less than the effective average of 50 µm, and the same applies to the other effective particle sizes mentioned. In a similar manner, the average effective particle sizes may be related to weight distributions of the particles but usually this will result in the same or about the same value for the average effective particle size.

The pharmaceutical compositions of the present invention provide release of the active ingredient bedaquiline over a prolonged period of time and therefore they can also be referred to as sustained or delayed release compositions. After administration, the compositions of the invention stay in the body and steadily release bedaquiline, keeping such levels of this active ingredient in the patient's system for a prolonged period of time, thereby providing, during said period, the appropriate treatment or prevention of a pathogenic mycobacterial infection. Because of the fact that the pharmaceutical compositions of the invention stay in the body and steadily release bedaquiline (and its active metabolite, referred to as M2 herein; see hereinafter, the methyl-substituted metabolite), they can be referred to as pharmaceutical compositions suitable as long-acting (or depot) formulations.

As used herein with the term "prolonged period of time", there is meant a term (or time period) that may be in the range of one week up to one year or up to two years, or a term in the range of one to two weeks, or two to three weeks, or three to four weeks, or a term in the range of one to two months, or two to three months, or three to four months, or three to six months, or six months to 12 months, or 12 months to 24 months, or a term that is in the range of several days, e.g. 7, 10 or 12 days, or several weeks, e.g. 2, 3 or 4 weeks, or one month, or several months, e.g. 2, 3, 4, 5 or six months or even longer, e.g. 7, 8, 9 or 12 months.

The pharmaceutical compositions of this invention may be applied in the long-term treatment or the long-term prevention of a pathogenic mycobacterial infection, or with other words they may be used in the treatment of a pathogenic mycobacterial infection, or in the prevention of a pathogenic mycobacterial infection, during a prolonged period of time. The compositions of the invention are effective in the treatment or prevention of a pathogenic mycobacterial infection for a prolonged period of time, for example for at least about one week or longer, or for about 1 month or longer. By the expression "effective for at least about one week or longer", one means that the plasma level of the active ingredient, bedaquiline (and/or its active metabolite M2), should be above a threshold value. In case of therapeutic application said threshold value is the lowest plasma level at which bedaquiline (and/or its active metabolite M2) provides effective treatment of a pathogenic mycobacterial infection. In case of application in the prevention of a pathogenic mycobacterial infection said threshold value is the lowest plasma level at which bedaquiline (and/or its active metabolite M2) is effective in preventing transmission of a pathogenic mycobacterial infection.

With "long term" for example as used in relation to "long term prevention of a pathogenic mycobacterial infection" or "long term treatment of a pathogenic mycobacterial infection", or similar terminology, there are meant terms that may be in the range of one week up to one year or up to two years, or longer, such as five or 10 years. In particular in the case of treatment of a pathogenic mycobacterial infection, such terms will be long, in the order of one to several months, one year or longer. Such terms may also be relatively short, in particular in the case of prevention. Shorter terms are those of several days, e.g. 7, 10 or 12 days, or several weeks, e.g. 2, 3 or 4 weeks, or one month, or several months, e.g. 2, 3, 4, 5 or six months or even longer, e.g. 7, 8, 9 or 12 months. In one embodiment the methods and uses in accordance with the present invention are for the prevention of a pathogenic mycobacterial infection during one month, or several months, e.g. 2, 3, 4, 5 or six months or even longer, e.g. 7, 8, 9 or 12 months.

The pharmaceutical compositions of the present invention can be administered at various time intervals. When used in the prevention of a pathogenic mycobacterial infection, the pharmaceutical compositions of this invention can be administered only once or a limited number of times such as twice, three, four, five or six times, or more. This may be recommendable where prevention is required during a limited period of time, such as the period during which there is a risk of infection.

The pharmaceutical compositions of the present invention can be administered at the time intervals mentioned above, such as at a time interval that is in the range of one week to one month, or in the range of one month to three months, or in the range of three months to six months, or in the range of six months to twelve months. In one
5 embodiment, the pharmaceutical composition can be administered once every two weeks, or once every month, or once every three months. In another embodiment the time interval is in the range of one to two weeks, or two to three weeks, or three to four weeks, or the time interval is in the range of one to two months, or two to three months, or three to four months, or three to six months, or six months to 12 months, or
10 12 months to 24 months. The time interval may be at least one week, but may also be several weeks, e.g. 2, 3, 4, 5 or 6 weeks, or at time intervals of one month, or of several months, e.g. 2, 3, 4, 5 or 6 months or even longer, e.g. 7, 8, 9 or 12 months. In one embodiment, the pharmaceutical compositions of the present invention are administered at a time interval of one, two or three months. These longer periods
15 between each administration of the pharmaceutical compositions of the invention provide further improvements in terms of pill burden and compliance. To further improve compliance, patients can be instructed to take their medication at a certain day of the week, where the composition is administered on a weekly schedule, or at a certain day of the month in case of a monthly schedule.

20 The length of the time intervals between each administration of a composition of the present invention may vary. For example said time intervals may be selected in function of the plasma levels. The intervals may be shorter where the plasma levels of bedaquiline (and/or its active metabolite M2) are deemed too low, e.g. when these
25 approach the minimum plasma level specified hereinafter. The intervals may be longer where the plasma levels of bedaquiline (and/or its active metabolite M2) are deemed too high. In one embodiment, the compositions of the invention are administered at equal time intervals. The compositions may be administered without any interjacent additional administrations, or with other words, the compositions may be administered
30 at particular points in time separated from one another by a time period of varying or equal length, e.g. a time period of at least one week, or any other time period specified herein, during which no further bedaquiline is administered. Having time intervals of the same length has the advantage that the administration schedule is simple, e.g. administration takes place at the same day in the week, or the same day in the month.
35 Such administration schedule therefore involves limited "pill burden" thereby contributing beneficially to the patient's compliance to the prescribed dosing regimen.

The concentration (or “C”) of bedaquiline (and/or its active metabolite M2) in the plasma of a subject treated therewith is generally expressed as mass per unit volume, typically nanograms per milliliter (ng/ml). For convenience, this concentration may be referred to herein as “plasma drug concentration” or “plasma concentration”.

5

The dose (or amount) of bedaquiline administered, depends on the amount of bedaquiline in the pharmaceutical compositions of the invention, or on the amount of a given composition that is administered. Where higher plasma levels are desired, either or both of a composition of higher bedaquiline concentration, or more of a given composition, may be administered. This applies vice versa if lower plasma levels are desired. Also a combination of varying time intervals and varying dosing may be selected to attain certain desired plasma levels.

10

The dose (or amount) of bedaquiline administered also depends on the frequency of the administrations (i.e. the time interval between each administration). Usually, the dose will be higher where administrations are less frequent. All these parameters can be used to direct the plasma levels to desired values

15

The dosing regimen also depends on whether prevention or treatment of the pathogenic mycobacterial infection is envisaged. In case of therapy, the dose of bedaquiline administered or the frequency of dosing, or both, are selected so that the plasma concentration of bedaquiline is kept above a minimum plasma level. The term “minimum plasma level” (or C_{min}) in this context refers to the plasma level of bedaquiline (and/or its active metabolite M2) that provides effective treatment of the pathogenic mycobacterial infection. In particular, the plasma level of bedaquiline (and/or its active metabolite M2) is kept at a level above a minimum plasma level of about 10 ng/ml, or above about 15 ng/ml, or above about 20 ng/ml, or above about 40 ng/ml. The plasma level of bedaquiline (and/or its active metabolite M2) may be kept above a minimum plasma level that is higher, for example above about 50 ng/ml, or above about 90 ng/ml, or above about 270 ng/ml, or above about 540 ng/ml. In one embodiment, the plasma level of bedaquiline (and/or its active metabolite M2) is kept above a level of about 13.5 ng/ml, or is kept above a level of about 20 ng/ml. Or the plasma level of bedaquiline (and/or its active metabolite M2) may be kept within certain ranges, in particular ranges starting from a minimum plasma level selected from those mentioned above and ending at a higher plasma levels selected from those mentioned above and selected from 500 ng/ml and 1000 ng/ml (e.g. from 10 to 15, 10 to 20, 10 to 40, etc., or from 15 to 20, or 15 to 40, or 15 to 90, etc., or 20 to 40, 20 to 90, or 20 to 270, etc., or 40 to 90, 40 to 270, or 40 -540, etc., each time from about the

20

25

30

35

indicated value in ng/ml to about the indicated value in ng/ml). In one embodiment said range is from about 10 to about 20, from about 20 to about 90, from 90 to 270, from 270 to 540, from 540 to 1000, each time from about the indicated value in ng/ml to about the indicated value in ng/ml.

5

The plasma levels of bedaquiline (and/or its active metabolite M2) should be kept above the above-mentioned minimum plasma levels because at lower levels the bacteria may no longer be sufficiently suppressed so that it can multiply with the additional risk of the emergence of mutations.

10

In the instance of prevention, the term “minimum plasma level” (or C_{min}) refers to the lowest plasma level of bedaquiline (and/or its active metabolite M2) that provides effective treatment/prevention of infection.

15

In particular, in the instance of prevention, the plasma level of bedaquiline (and/or its active metabolite M2) can be kept at a level above a minimum plasma level mentioned above in relation to therapy. However in prevention the plasma level of bedaquiline (and/or its active metabolite M2) can be kept at a lower level, for example at a level above about 4 ng/ml, or about 5 ng/ml, or about 8 ng/ml. The plasma levels of bedaquiline (and/or its active metabolite M2) should preferably be kept above these minimum plasma levels because at lower levels the drug may no longer be effective thereby increasing the risk of transmission of infection. Plasma levels of bedaquiline (and/or its active metabolite M2) may be kept at somewhat higher levels to have a safety margin. Such higher levels start from about 50 ng/ml or more. The plasma level of bedaquiline (and/or its active metabolite M2) can be kept at a level that is in the ranges mentioned above in relation to therapy, but where the lower limits include the plasma levels of about 4 ng/ml, or about 5 ng/ml, or about 8 ng/ml.

20

25

30

35

An advantage of bedaquiline (and/or its active metabolite M2) is that it may be used up to relatively high plasma levels without any significant side effects. The plasma concentrations of bedaquiline (and/or its active metabolite M2) may reach relatively high levels, but as with any drug should not exceed a maximum plasma level (or C_{max}), which is the plasma level where bedaquiline (and/or its active metabolite M2) causes significant side effects. Additionally, compound-release from the tissue should also be taken into account, which is not counted for within plasma levels. As used herein, the term “significant side effects” means that the side effects are present in a relevant patient population to an extent that the side effects affect the patients’ normal functioning. In an embodiment, the amount and the frequency of administrations of

bedaquiline (and/or its active metabolite M2) to be administered are selected such that the plasma concentrations are kept during a long term at a level comprised between a maximum plasma level (or $C_{\max}$ as specified above) and a minimum plasma level (or $C_{\min}$ as specified above).

5

In certain instances it may be desirable to keep the plasma levels of bedaquiline (and/or its active metabolite M2) at relatively low levels, e.g. as close as possible to the minimum plasma levels specified herein. This will allow reducing the frequency of the administrations and/or the quantity of bedaquiline (and/or its active metabolite M2) administered with each administration. It will also allow avoiding undesirable side effects, which will contribute to the acceptance of the dosage forms in most of the targeted population groups who are healthy people at risk of being infected and therefore are less inclined to tolerate side effects. The plasma levels of bedaquiline (and/or its active metabolite M2) may be kept at relatively low levels in the instance of prevention. One embodiment concerns uses or methods for prevention of infection, as specified above or hereinafter, wherein the minimum plasma level of bedaquiline (and/or its active metabolite M2) is as specified herein and the maximum plasma level is about equal to the lowest plasma level that causes the active ingredient to act therapeutically, also as specified herein.

20

In other embodiments, the plasma level of bedaquiline (and/or its active metabolite M2) is kept at a level below a lower maximum plasma level of about 10 ng/ml, more in particular about 15 ng/ml, further in particular about 20 ng/ml, still more in particular about 40 ng/ml. In a particular embodiment, the plasma level of bedaquiline (and/or its active metabolite M2) is kept below a level of about 13.5 ng/ml. In one embodiment, the plasma level of bedaquiline (and/or its active metabolite M2) is kept in an interval of the lower maximum blood level specified above, and the minimum plasma levels mentioned in relation to prevention. For example the plasma levels of bedaquiline (and/or its active metabolite M2) are kept below about 10 ng/ml and above a minimum level of about 4 ng/ml.

30

In other instances it may be desirable to keep the plasma levels of bedaquiline (and/or its active metabolite M2) at relatively higher levels, for example where there is a high risk of infection and more frequent and/or higher doses are not an issue. In these instances the minimum plasma level may be equal to the lowest plasma level of bedaquiline (and/or its active metabolite M2) that provides effective treatment of a pathogenic mycobacterial infection, such as the specific levels mentioned herein.

35

In the instance of prevention, the dose to be administered should be calculated on a basis of about 0.2 mg/day to about 50 mg/day, or 0.5 mg/day to about 50 mg/day, or of about 1 mg/day to about 10 mg/day, or about 2 mg/day to about 5 mg/day, e.g. about 3 mg/day. This corresponds to a weekly dose of about 1.5 mg to about 350 mg, in particular of about 3.5 mg to about 350 mg, in particular of about 7 mg to about 70 mg, or about 14 mg to about 35 mg, e.g. about 35 mg, or to a monthly dose of from 6 mg to about 3000 mg, in particular about 15 mg to about 1,500 mg, more in particular of about 30 mg to about 300 mg, or about 60 mg to about 150 mg, e.g. about 150 mg. Doses for other dosing regimens can readily be calculated by multiplying the daily dose with the number of days between each administration.

In the instance of therapy, the dose to be administered should be somewhat higher and should be calculated on a basis of about 1 mg/day to about 150 mg/day, or of about 2 mg/day to about 100 mg/day, or of about 5 mg/day to about 50 mg/day, or about 10 mg/day to about 25 mg/day, e.g. about 15 mg/day. The corresponding weekly or monthly doses can be calculated as set forth above. For applications in prevention, the doses may be lower although the same dosing as for therapeutic applications may be used. In an embodiment, the dose/administration is given at monthly intervals or three-monthly or six-monthly intervals, with the total treatment duration being three, six or 12 months. In the instances where the dose/administration is monthly, three monthly or six-monthly, in an embodiment, the dose given (e.g. in human subjects) is calculated on the basis of a 400 mg daily dose given for 2 weeks. Hence, the total amount of bedaquiline given per dose may be about 5600 mg (e.g. in the range of 3000 and 8000 mg), but it may be up to one fifth of such an amount (e.g. in the range of 500 and 2000 mg, e.g. between about 1000 and 1500 mg).

In another embodiment, in the case of prevention or in particular therapy, the doses may also be expressed in mg/kg. For instance, in the examples, certain doses may be administered based on weight (of e.g. the mammal, and as shown in the examples here, in mouse) and hence doses between 1 mg/kg and 1000 mg/kg may be employed (e.g. 40 mg/kg, 80 mg/kg, 160 mg/kg, 320 mg/kg or 480 mg/kg may be employed) and such doses may remain effective for a period of 4 weeks, 8 weeks or 12 weeks (for example as shown in the examples). For instance, one dose may be taken every 4 weeks (effectively seen as a 12 week treatment regimen, i.e. three doses in total) or one single dose may be taken, which effectively provides sufficient treatment (e.g. as defined by reduction in CFUs, see examples) as may be evidenced by monitoring over a 12 week period. Hence, in an aspect, in order to treat the bacterial infection one dose may be taken (e.g. between 1 mg/kg and 1000 mg/kg, for instance between 2 mg/kg and

500 mg/kg) or one such dose may be taken every 4 weeks (e.g. two or three such doses may be taken). Such dose depends on the bacterial infection to be treated. For instance, in the treatment of latent tuberculosis or leprosy, lower doses may be required (compared to e.g. multi-drug resistant tuberculosis) given that a lower amount of

5 bedaquiline is required to control the bacteria. An example of this may be described hereinafter, wherein it is indicated that in mice one dose of 160 mg/kg may sufficiently reduce CFUs in the mouse model of latent tuberculosis infection – it was also seen that two or three doses of 160 mg/kg (the second and the third doses administered at 4 and 8 weeks, respectively) were also effective in that model.

10

It has been found that, once administered, the plasma levels of bedaquiline (and/or its active metabolite M2) are more or less stable, i.e. they fluctuate within limited margins. The plasma levels have been found to approach more or less a steady state mode or to approximate more or less a zero order release rate during a prolonged period of time.

15

By “steady state” is meant the condition in which the amount of drug present in the plasma of a subject stays at more or less the same level over a prolonged period of time. The plasma levels of bedaquiline (and/or its active metabolite M2) generally do not show any drops below the minimum plasma level at which the drug is effective. The term “stays at more or less the same level” does not exclude that there can be small

20 fluctuations of the plasma concentrations within an acceptable range, e.g. fluctuations within a range of about $\pm 30\%$, or about $\pm 20\%$, or about $\pm 10\%$, or about $\pm 10\%$.

20

In some instances there may be an initial plasma concentration peak after administration, after which the plasma levels achieve a “steady-state”, as mentioned

25 hereinafter.

25

The compositions of the invention show good local tolerance and ease of administration. Good local tolerance relates to minimal irritation and inflammation at the site of injection; ease of administration refers to the size of needle and length of

30 time required to administer a dose of a particular drug formulation. In addition, the compositions of the invention show good stability and have an acceptable shelf life.

30

The micro- or nanoparticles of the present invention have a surface modifier adsorbed on the surface thereof. The function of the surface modifier is to act as a wetting agent

35 as well as a stabilizer of the colloidal suspension.

35

In one embodiment, the micro- or nanoparticles in the compositions of the invention mainly comprise crystalline bedaquiline or a salt thereof; and a surface modifier, the

combined amount of which may at least comprise about 50%, or at least about 80%, or at least about 90%, or at least about 95%, or at least about 99% of the micro- or nano particles. As indicated herein, in an embodiment, bedaquiline is in its non-salt form (or in its "free form") and in a further embodiment it is in a crystalline non-salt (or free) form. In this respect, as mentioned herein, bedaquiline may be prepared as such using the procedures described in international patent application WO 2004/011436 (or in WO 2006/125769, which describes an optical resolution with a chiral reagent).

Following such procedure, the bedaquiline is obtained by precipitation from toluene/ethanol and it is indicated that the product crystallises. Such form of bedaquiline may be used in the preparation of the compositions of the invention and, further, such form may be a single crystalline polymorph with the following characterising features:

- (i) a melting endotherm at 181.5°C (endotherm onset) and DSC curve showing melting of the product at about 182.5°C (immediately followed by decomposition; measured by differential scanning calorimetry (DSC) by transfer of about 3 mg of compound into a standard aluminum TA-Instrument sample pan, sample pan closed with the appropriate cover and DSC curve recorded on a TA-Instruments Q2000 MTDSC equipped with a RCS cooling unit using the following parameters – initial temperature 25°C; heating range 10°C/min; final temperature 300°C, nitrogen flow 50 ml/min);
- (ii) infrared (IR) spectrum peaks at *inter alia* about 1600 cm⁻¹, about 1450 cm⁻¹, about 1400 cm⁻¹, about 1340 cm⁻¹, and about 1250 cm⁻¹ (where a sample is analysed using a suitable microATR accessory deploying 32 scans, 1 cm⁻¹ resolution, Thermo Nexus 670 FTIR spectrometer, a DTGS with KBr windows detector, Ge on KBr beamsplitter and a micro ATR accessory (Harrick Split Pea with Si crystal); and/or
- (iii) X-ray powder diffraction (XRPD) with characteristic peaks at about 11.25° 2-Theta, about 18° 2-Theta, about 18.5° 2-Theta, about 19° 2-Theta, about 20.25° 2-Theta, about 21.25° 2-Theta, about 22.25° 2-Theta, about 24.5° 2-Theta and about 27° 2-Theta, showing diffraction peaks without the presence of a halo indicating crystallinity of the product (where the analysis was carried out on a PANalytical (Philips) X'PertPRO MPD diffractometer, and the instrument is equipped with a Cu LFF X-ray tube and the compound was spread on a zero background sample holder; the Instrument Parameters were: generator voltage – 45 kV; generator amperage – 40 mA; geometry – Bragg-Brentano; stage – spinner stage; scan mode – continuous; scan range 3 to 50° 2θ; step size 0.02°/step; counting time 30 sec/step; spinner revolution time – 1 sec; radiation type CuKα).

Hence, in an embodiment, the bedaquiline employed in a process to prepare compositions of the invention (i.e. before conversion to micro/nano-particles) is a crystalline form (e.g. of the specific form characterised above). In a further embodiment of the invention, the bedaquiline employed in the compositions of the invention (i.e. after conversion to micro/nano-particles, for instance by milling) is also in a crystalline form (e.g. of the specific form characterised above).

In a further aspect, the present invention is concerned with a pharmaceutical composition for administration by intramuscular or subcutaneous injection, comprising a therapeutically effective amount of bedaquiline, or a pharmaceutically acceptable salt thereof, in the form of a suspension of particles consisting essentially of:

- (1) bedaquiline, or a pharmaceutically acceptable salt thereof in micro- or nanoparticle form, having a surface modifier adsorbed to the surface thereof; and
- (2) a pharmaceutically acceptable aqueous carrier; wherein the active ingredient is suspended,

which is characterised in that the surface modifier comprises PEG4000 or the like.

It is indicated that the formulations of the invention contain PEG4000 (or the like), and for the avoidance of doubt, this may be in combination with another suitable surface modifier.

Suitable surface modifiers (that may be used in combination with PEG4000, or the like) can be selected from known organic and inorganic pharmaceutical excipients, including various polymers, low molecular weight oligomers, natural products and surfactants.

Particular surface modifiers include nonionic and anionic surfactants. Representative examples of surface modifiers include gelatin, casein, lecithin, salts of negatively charged phospholipids or the acid form thereof (such as phosphatidyl glycerol, phosphatidyl inositol, phosphatidyl serine, phosphatic acid, and their salts such as alkali metal salts, e.g. their sodium salts, for example egg phosphatidyl glycerol sodium, such as the product available under the tradename Lipoid™ EPG), gum acacia, stearic acid, benzalkonium chloride, polyoxyethylene alkyl ethers, e.g., macrogol ethers such as cetomacrogol 1000, polyoxyethylene castor oil derivatives; polyoxyethylene stearates, colloidal silicon dioxide, sodium dodecylsulfate, carboxymethylcellulose sodium, bile salts such as sodium taurocholate, sodium desoxytaurocholate, sodium desoxycholate; methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, magnesium aluminate silicate, polyvinyl alcohol (PVA), poloxamers, such as Pluronic™ F68, F108 and F127 which are block copolymers of ethylene oxide and propylene oxide; tyloxapol; Vitamin E-TGPS (α -tocopheryl polyethylene glycol

succinate, in particular α -tocopheryl polyethylene glycol 1000 succinate); poloxamines, such as TetronicTM 908 (T908) which is a tetrafunctional block copolymer derived from sequential addition of ethylene oxide and propylene oxide to ethylenediamine; dextran; lecithin; dioctyl ester of sodium sulfosuccinic acid such as the products sold under the tradename Aerosol OTTM (AOT); sodium lauryl sulfate (DuponolTM P); alkyl aryl polyether sulfonate available under the tradename TritonTM X-200; polyoxyethylene sorbitan fatty acid esters (TweensTM 20, 40, 60 and 80); sorbitan esters of fatty acids (SpanTM 20, 40, 60 and 80 or ArlacelTM 20, 40, 60 and 80); sucrose stearate and sucrose distearate mixtures such as the product available under the tradename CrodestaTM F110 or CrodestaTM SL-40; hexyldecyl trimethyl ammonium chloride (CTAC); polyvinylpyrrolidone (PVP). If desired, two or more surface modifiers can be used in combination (with PEG4000 or the like).

Particular surface modifiers that may be employed in combination with PEG4000 (or the like) are selected from poloxamers, α -tocopheryl polyethylene glycol succinates, polyoxyethylene sorbitan fatty acid esters, and salts of negatively charged phospholipids or the acid form thereof. More in particular the surface modifiers are selected from PluronicTM F108, Vitamin E TGPS, TweenTM 80, and LipoidTM EPG (and, in a particular embodiment, it is Vitamin E TPGS). One or more of these surface modifiers may be used. PluronicTM F108 corresponds to poloxamer 338 and is the polyoxyethylene, polyoxypropylene block copolymer that conforms generally to the formula $\text{HO}-[\text{CH}_2\text{CH}_2\text{O}]_x-[\text{CH}(\text{CH}_3)\text{CH}_2\text{O}]_y-[\text{CH}_2\text{CH}_2\text{O}]_z-\text{H}$ in which the average values of x, y and z are respectively 128, 54 and 128. Other commercial names of poloxamer 338 are Hodag NonionicTM 1108-F and SynperonicTM PE/F108. In one embodiment, the surface modifier comprises a combination of a polyoxyethylene sorbitan fatty acid ester and a phosphatidyl glycerol salt (in particular egg phosphatidyl glycerol sodium).

The optimal relative amount of bedaquiline in relation to the surface modifier depends on the surface modifier selected, the specific surface area of the bedaquiline suspension which is determined by the average effective particle size and the bedaquiline concentration, the critical micelle concentration of the surface modifier if it forms micelles, etc. The relative amount (w/w) of bedaquiline to the surface modifier preferably is in the range of 1 : 2 to about 20 : 1, in particular in the range of 1 : 1 to about 10 : 1, e.g. in the range of 2 : 1 to about 10 : 1, for instance about 4 : 1.

As indicated, the surface modifier contains PEG4000, but may also contain a further surface modifier (for instance, a surface modifier mentioned hereinbefore). In various

embodiments of the invention, the compositions of the invention comprise a surface modifier that contains PEG4000 and one or more other surface modifiers in the following w/w ratios:

- at least 1 : 10 of PEG4000 : one or more other surface modifiers
- 5 - between 1 : 10 and 100 : 1 (e.g. between about 1 : 10 and 20 : 1) of PEG4000 : one or more other surface modifiers
- about 10 : 1 PEG4000 : one or more other surface modifiers

Hence, when the surface modifier of the compositions of the invention comprises a ratio of at least 1 : 10 w/w of PEG4000 (or the like) : one or more other surface modifiers, then it may contain 5 mg/mL PEG4000 and 50 mg/ml of one or more other surface modifier (e.g. Vitamin E TPGS, also referred to herein as simply "TPGS"). In an embodiment, given that the relative amount of bedaquinine to the surface modifier may be between 1 : 1 and 10 : 1 (e.g. about 4 : 1), then the bedaquinine may be present in about 200 mg/ml in such instances (which may form a particular injectable formulation or dose). While the compositions of the invention are distinguished as they contain PEG4000 (or the like) and it may be a relatively small amount, in an embodiment, the surface modifier comprises at least 25% by weight, for example at least 50% by weight PEG4000 or the like (and the remainder being one or more other suitable surface modifiers as described herein, for example Vitamin E TPGS). For instance, in an embodiment, the surface modifier of the compositions of the invention comprise a ratio of at least 1 : 1 w/w of PEG4000 or the like : one or more other suitable surface modifiers. In a further embodiment, the surface modifier comprises at least 75% by weight PEG4000 or the like (and the remainder being one or more other suitable surface modifiers as described herein, for example Vitamin E TPGS). Hence, in an embodiment, the surface modifier of the compositions of the invention comprise a ratio of at least 3 : 1 w/w of PEG4000 or the like : one or more other suitable surface modifiers. In yet further embodiments, the surface modifier of the compositions of the invention comprise at least 85% by weight PEG4000 or the like or between about 85% and about 95% PEG4000 or the like (and in each case, the remainder is one or more other suitable surface modifier as described herein, e.g. Vitamin E TPGS). Hence, in an embodiment the surface modifier of the compositions of the invention comprise a ratio of at least 8 : 1 w/w of PEG4000 or the like : one or more other suitable surface modifiers (for instance a ratio of between 8 : 1 and 12 : 1 w/w of PEG4000 or the like : one or more other suitable surface modifiers).

The ratios of PEG4000 (and the like) and one or more other surface modifiers may also depend on the other surface modifiers being used; for instance when the one or more

other surface modifiers comprises Vitamin E TPGS and/or Tween (a polyoxyethylene polyether sulfonate), the ratios hereinabove may be applicable, and for instance the surface modifier comprises at least 60% by weight PEG4000 and, in an embodiment at least 75%; in the case where the one or more other surface modifiers comprises a poloxamer then the ratio may be between 1:10 to 10:1 (of PEG : one or more other surface modifier), for instance between 1:5 and 5:1 and, in an embodiment between 1:2 and 2:1, and, in an embodiment, the surface modifier in this instance comprises at least 30% PEG4000, for instance, at least 40% (and, in a specific embodiment about 50%). In certain instances, at least 10% PEG4000 is required, but the upper limit may be 60% (e.g. when the one or more other surface modifier is a poloxamer).

As indicated, the compositions of the invention comprise a surface modifier that contain PEG4000 or the like. In an embodiment, the surface modifier may consist essentially of PEG4000 or the like. However, in an alternative embodiment, the surface modifier also contains another suitable surface modifier as described herein.

Where one or more other surface modifiers are employed in compositions of the invention, then those other surface modifiers may, in a particular embodiment, be selected from Vitamin E TPGS or a poloxamer. For instance, the other surface modifier may be Vitamin E TPGS. Hence, as indicated herein, the w/w ratio of bedaquiline to surface modifier may be in the range 2:1 to 10:1 (e.g. about 4:1) and hence, when 200 mg/ml of bedaquiline is employed (e.g. for a single injectable dose), then that may contain between 100 mg/ml and 20 mg/ml surface modifier. In this instance, and again as indicated, the amount of surface modifier may contain PEG4000 (or the like) and one or more other suitable surface modifiers in a ratio of, for example, at least 3 : 1 (or at least 75% by weight PEG4000). Hence, when there is 100 mg/ml surface modifier present, then that may consist of at least 75 mg/ml PEG4000 or the like, with any remainder consisting of one or more other suitable surface modifiers (e.g. Vitamin E TPGS) and when there is 20 mg/ml surface modifier then this may consist of at least 15 mg/ml PEG4000 or the like, and any remainder consisting of one or more other suitable surface modifier. As it is indicated hereinbefore that the ratio of bedaquiline to surface modifier may be about 4 : 1, then when there is 200 mg/ml bedaquiline (e.g. as one injectable dose), then the amount of surface modifier may be between about 35 mg/ml and 60 mg/ml (for instance about 55 mg/ml, in which case the surface modifier may contain about 50 mg/ml PEG4000 or the like, and about 5 mg/ml of one or more other surface modifier, e.g. Vitamin E TPGS).

The compositions of the invention may need to be sterile so that they can be administered to patients. Achieving sterile compositions may be done in a number of ways, including manufacturing such compositions in a sterile process or environment. However, such a method has a number of drawbacks, challenges and is associated with higher costs. A preferred alternative is to undergo sterilization without having to conform to an entire sterile process, and heat sterilization, autoclaving and gamma radiations are sterilization steps that can achieve that. Advantageously, compositions of the invention can be autoclaved, i.e. are autoclavable, and that can be done without substantial degradation or decomposition of the compositions.

Further challenges arise after sterilization, which are linked to desired stability of the long-acting formulation, undesired aggregation of particles of the active pharmaceutical ingredient (API) within that formulation and the desired re-suspendability of the formulation (after sterilization, e.g. autoclaving).

In this case, the compositions of the invention may be sterilized, for instance by heat sterilization, autoclaving or gamma radiation (in an embodiment, the sterilization is performed by autoclaving), even though the cloud point may be below the temperature at which autoclaving takes place. Advantageously, the compositions of the invention may be easily resuspended after sterilization (even if the cloud point is exceeded during the sterilization process, in particular the autoclaving process).

Hence, in a further aspect of the invention, there is provided:

(a) a process for sterilizing the compositions of the invention (for instance, autoclaving the compositions);

(b) followed by re-suspending such compositions of the invention, which process may be referred to as a "process of the invention". The examples show that the PEG4000 may be key in re-suspending. It will be understood that after sterilization (e.g. heat sterilization or autoclaving), there may be some particle aggregation (especially if the sterilization process is performed at a temperature above the cloud point), for instance due to phase separation. Given that the compositions of the invention should essentially be a suspension, then the re-suspending step may be necessary (such re-suspending step may also be performed at a later point in time, e.g. when the suspension is being prepared for its end use). The compositions of the invention start as a suspension, with the bedaquiline particles suspended in the pharmaceutically acceptable carrier and the surface modifier (i.e. PEG4000 containing surface modifier as hereinbefore defined) may be adsorbed onto the surface of the bedaquiline – after autoclaving there may be disassociation between the surface

modifier (also referred to herein as wetting agent) and the bedaquiline and/or bedaquiline particle aggregation. Hence, re-suspending back to the original suspension is essential and may be effected by swirling or shaking the composition of the invention (after sterilization, e.g. autoclaving). The re-suspending (of bedaquiline in the carrier) may occur by allowing the surface modifier (i.e. PEG4000 and one or more other suitable surface modifiers) to adsorb onto the surface of bedaquiline.

As indicated, the re-suspendability after sterilization (e.g. autoclaving) may be linked to the presence of PEG4000. Additionally or alternatively, the use of PEG4000 as a surface modifier may be advantageous as it may replace a surface modifier that may be as efficient (e.g. with similar properties allowing for suspension and/or re-suspendability after sterilization) but where that surface modifier being replaced may not be tolerated (e.g. in humans) above a certain dose or quantity (e.g. as an injectable). For instance, other surface modifiers such as Vitamin E TPGS may not be tolerated above a certain dose as an injectable in humans and hence would either need to be replaced entirely or the dose/amount reduced.

A micro- or nano-suspension (not containing PEG4000) may be sterilized by autoclaving and may be adequately re-suspendable (for example, re-suspendable under conditions defined herein, especially by swirling for less than 40 seconds) in which case PEG4000 may not be needed. However, in an embodiment where the re-suspendability is not adequate (for instance, takes longer than 40 seconds), then the use of PEG4000, or the like, in such a micro- or nano-suspension may assist in improving the re-suspendability (i.e. by making it easier, including by reducing the time taken to less than 40 seconds), for instance after auto-claving. The US Pharmacopoeia indicates that suspensions should be re-dispersible in case they settle upon storage, etc, and a goal is to have a suspension in general where the time taken to re-suspend is as short as possible; in this respect, and in an aspect of the invention thus, PEG4000 (or the like) can assist.

Hence, in view of the above, in further embodiments of the invention, there is provided:

- PEG4000, or the like, for use as a surface modifier in a pharmaceutical composition for administration by intramuscular or subcutaneous injection, wherein said composition comprises an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, characterised in that the PEG4000 assists in re-suspending said composition for instance after sterilization (e.g. autoclaving)

- PEG4000, or the like, for use in re-suspending a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, for instance wherein said composition has undergone sterilization (e.g. autoclaving)
- PEG4000, or the like, for use as a resuspendability aid in a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, for instance wherein said composition has undergone sterilization (e.g. autoclaving)
- PEG4000, or the like, for use to increase (or improve) the resuspendability of a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, for instance wherein said composition has undergone sterilization (e.g. autoclaving)

In all cases above, the PEG4000, or the like, may be for such uses in pharmaceutical compositions described herein. Resuspendability may in certain circumstances be compared to the pharmaceutical composition without the PEG4000.

In an alternative further embodiments, there is provided:

- the use of PEG4000, or the like, as a surface modifier in a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, wherein the PEG4000 assists in re-suspending said composition, for instance after sterilization (e.g. autoclaving)
- the use of PEG4000, or the like, in re-suspending a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, for instance wherein said composition has undergone sterilization (e.g. autoclaving)
- the use of PEG4000, or the like, as a resuspendability aid in a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, for instance wherein said composition has undergone sterilization (e.g. autoclaving)
- the use of PEG4000, or the like, to increase (or improve) the resuspendability of a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a

suspension of micro- or nano-particles, for instance wherein said composition has undergone sterilization (e.g. autoclaving)

In all cases above, the use of PEG4000, or the like, may be in pharmaceutical compositions described herein. Again, resuspendability may in certain circumstances
5 be compared to the pharmaceutical composition without the PEG4000.

The particles of this invention can be prepared by means of micronization/particle size reduction/nanonization by mechanical means and by controlled precipitation from a supersaturated solution, or by using supercritical fluids such as in the GAS technique
10 (“gas anti-solvent”), or any combination of such techniques. In one embodiment a method is used comprising the steps of dispersing bedaquiline in a liquid dispersion medium and applying mechanical means in the presence of grinding media to reduce the particle size of bedaquiline to an average effective particle size of less than about 50 μm , in particular less than about 1,000 nm. The particles can be reduced in size in
15 the presence of a surface modifier.

A general procedure for preparing the particles of this invention comprises

- (a) obtaining bedaquiline in micronized form;
- (b) adding the micronized bedaquiline to a liquid medium to form a
20 premix/predispersion; and
- (c) subjecting the premix to mechanical means in the presence of a grinding medium to reduce the average effective particle size.

In a particular embodiment, there is provided a process for preparing a pharmaceutical
25 composition comprising

- (a) obtaining an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in micronized form;
- (b) adding the micronized active ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, to a liquid medium to form a
30 premix/predispersion, characterised in that the liquid medium contains a surface modifier comprising PEG4000, or the like, as per any one of claims 1, 2, 3 or 4;
- (c) subjecting the premix to mechanical means in the presence of a grinding medium to reduce the average effective particle size;
- 35 (d) sterilization (e.g. autoclaving); and
- (e) re-suspending (e.g. if needed).

In such instances, the re-suspending may be performed by swirling for less than 40 seconds. In a particular embodiment, there is provided the use of PEG4000 in such (a) process(es).

- 5 Bedaquiline in micronized form is prepared using techniques known in the art. It is preferred that the average effective particle size of the bedaquiline active agent in the predispersion be less than about 100 μm as determined by sieve analysis. Where the average effective particle size of the micronized bedaquiline is greater than about 100 μm , it is preferred that the particles of the bedaquiline compound be reduced in size
10 to less than 100 μm (for example to a size or size range as described herein).

- The micronized bedaquiline can then be added to a liquid medium in which it is essentially insoluble to form a predispersion. The concentration of bedaquiline in the liquid medium (weight by weight percentage) can vary widely and depends on the
15 selected surface modifier and other factors. Suitable concentrations of bedaquiline in compositions vary between about 0.1% to about 60%, or between about 1% to about 60%, or between about 10% to about 50%, or between about 10% to about 30%, e.g. about 10%, 20% or 30% (each % in this paragraph relating to w/v).

- 20 The premix can be used directly by subjecting it to mechanical means to reduce the effective average effective particle size in the dispersion to less than 2,000 nm. It is preferred that the premix be used directly when a ball mill is used for attrition. Alternatively, bedaquiline and, optionally, the surface modifier, can be dispersed in the liquid medium using suitable agitation such as, for example, a roller mill, until a
25 homogeneous dispersion is achieved.

- The mechanical means applied to reduce the effective average effective particle size of bedaquiline conveniently can take the form of a dispersion mill. Suitable dispersion mills include a ball mill, an attritor/attrition mill, a vibratory mill, a planetary mill,
30 media mills, such as a sand mill and a bead mill. A media mill is preferred due to the relatively shorter milling time required to provide the desired reduction in particle size. The beads preferably are ZrO_2 beads. For instance, for the nanoparticles, the ideal bead size is about 0.5 mm and, for the microparticles, the ideal bead size is about 2 mm.

- 35 The grinding media for the particle size reduction step can be selected from rigid media preferably spherical or particulate in form having an average size less than 3 mm and, more preferably, less than 1 mm (as low as 200 μm beads). Such media desirably can provide the particles of the invention with shorter processing times and impart less

wear to the milling equipment. Examples of grinding media are ZrO_2 such as 95% ZrO_2 stabilized with magnesia or stabilized with yttrium, zirconium silicate, glass grinding media, polymeric beads, stainless steel, titania, alumina and the like. Preferred grinding media have a density greater than 2.5 g/cm^3 and include 95% ZrO_2 stabilized with magnesia and polymeric beads.

The attrition time can vary widely and depends primarily upon the particular mechanical means and processing conditions selected. For rolling mills, processing times of up to two days or longer may be required.

The particles should be reduced in size at a temperature that does not significantly degrade the bedaquiline compound. Processing temperatures of less than 30 to 40°C are ordinarily preferred. If desired, the processing equipment may be cooled with conventional cooling equipment. The method is conveniently carried out under conditions of ambient temperature and at processing pressures, which are safe and effective for the milling process.

The pharmaceutical compositions according to the present invention contain an aqueous carrier that preferably is pharmaceutically acceptable. Said aqueous carrier comprises sterile water optionally in admixture with other pharmaceutically acceptable ingredients. The latter comprise any ingredients for use in injectable formulations. Such ingredients are optional. These ingredients may be selected from one or more of a suspending agent, a buffer, a pH adjusting agent, a preservative, an isotonicizing agent, and the like ingredients. In one embodiment, said ingredients are selected from one or more of a suspending agent, a buffer, a pH adjusting agent, and optionally, a preservative and an isotonicizing agent. Particular ingredients may function as two or more of these agents simultaneously, e.g. behave like a preservative and a buffer, or behave like a buffer and an isotonicizing agent.

Suitable optional buffering agents and pH adjusting agents should be used in amount sufficient to render the dispersion neutral to very slightly basic (up to pH 8.5), preferably in the pH range of 7 to 7.5. Particular buffers are the salts of weak acids. Buffering and pH adjusting agents that can be added may be selected from tartaric acid, maleic acid, glycine, sodium lactate/lactic acid, ascorbic acid, sodium citrates/citric acid, sodium acetate/acetic acid, sodium bicarbonate/carbonic acid, sodium succinate/succinic acid, sodium benzoate/benzoic acid, sodium phosphates, tris(hydroxymethyl)aminomethane, sodium bicarbonate/sodium carbonate, ammonium hydroxide, benzene sulfonic acid, benzoate sodium/acid, diethanolamine, glucono delta

lactone, hydrochloric acid, hydrogen bromide, lysine, methanesulfonic acid, monoethanolamine, sodium hydroxide, tromethamine, gluconic, glyceric, glutamic, ethylene diamine tetraacetic (EDTA), triethanolamine, including mixtures thereof. In an embodiment, the compositions of the invention do not contain a buffering agent. In an embodiment, especially when pH lowers, the compositions of the invention do contain a buffer, for example a citrate-phosphate buffer.

Suitable optional preservatives comprise antimicrobials and anti-oxidants which can be selected from the group consisting of benzoic acid, benzyl alcohol, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), chlorbutol, a gallate, a hydroxybenzoate, EDTA, phenol, chlorocresol, metacresol, benzethonium chloride, myristyl- γ -piccolinium chloride, phenylmercuric acetate and thimerosal. Radical scavengers include BHA, BHT, Vitamin E and ascorbyl palmitate, and mixtures thereof. Oxygen scavengers include sodium ascorbate, sodium sulfite, L-cysteine, acetylcysteine, methionine, thioglycerol, acetone sodium bisulfite, isoascorbic acid, hydroxypropyl cyclodextrin. Chelating agents include sodium citrate, sodium EDTA and malic acid. In an embodiment of the invention, the compositions of the invention do not contain a perseverative.

An isotonizing agent or isotonifier may be present to ensure isotonicity of the pharmaceutical compositions of the present invention, and includes sugars such as glucose, dextrose, sucrose, fructose, trehalose, lactose; polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol. Alternatively, sodium chloride, sodium sulfate, or other appropriate inorganic salts may be used to render the solutions isotonic. These isotonifiers can be used alone or in combination. The suspensions conveniently comprise from 0 to 10% (w/v), in particular 0 to 6% of isotonizing agent. Of interest are nonionic isotonifiers, e.g. glucose, as electrolytes may affect colloidal stability. In an embodiment of the invention, the compositions of the invention contain an isotonizing agent or isotonifier, which, in a further embodiment is a nonionic isotonifier, such as a suitable sugar such as mannitol. The amount of the isotonizing agent is as hereinbefore described, but may also be added in a certain ratio compared to bedaquiline, for instance the w/w ratio of bedaquiline and isotonizing agent (e.g. mannitol) may be between 1 : 1 and 10 : 1, for instance between about 2 : 1 and 8 : 1, especially between about 3 : 1 and 6 : 1 (e.g. about 4 : 1).

A desirable feature for a pharmaceutical composition of the invention relates to the ease of administration. The viscosity of the pharmaceutical compositions of the invention

should be sufficiently low to allow administration by injection. In particular they should be designed so that they can be taken up easily in a syringe (e.g. from a vial), injected through a fine needle (e.g. a 20 G 1½, 21 G 1½, 22 G 2 or 22 G 1¼ needle) in not too long a time span. In one embodiment the viscosity of the compositions of the invention is below about 75 mPa·s, or below 60 mPa·s. Aqueous suspensions of such viscosity or lower usually meet the above-mentioned criteria.

Ideally, the aqueous suspensions according to the present invention will comprise as much bedaquiline (or pharmaceutically acceptable salt thereof) as can be tolerated so as to keep the injected volume to a minimum, in particular from 3 to 70% (w/v), or from 3 to 60% (w/v), or from 3 to 40% (w/v), or from 10 to 40% (w/v), of bedaquiline (or pharmaceutically acceptable salt thereof). In one embodiment the aqueous suspensions of the invention contain about 50% - 70% (w/v) of bedaquiline (or pharmaceutically acceptable salt thereof), or about 40% - 60% (w/v) of bedaquiline (or pharmaceutically acceptable salt thereof), or about 30% - 50% (w/v) of bedaquiline (or pharmaceutically acceptable salt thereof).

In one embodiment, the aqueous suspensions may comprise by weight, based on the total volume of the composition:

- (a) from 10% to 70% (w/v), or from 20% to 60% (w/v), or from 20% to 50% (w/v), or from 20% to 40% (w/v) of bedaquiline (or pharmaceutically acceptable salt thereof);
- (b) from 0.5% to 20 % (w/v), or from 2% to 15% or 20% (w/v), or from 5% to 15% (w/v) of a wetting agent (also referred to herein as a surface modifier);
- (c) from 0% to 10% (w/v), or from 0% to 5% (w/v), or from 0% to 2% (w/v), or from 0% to 1% (w/v) of one or more buffering agents;
- (d) from 0% to 20 % (w/v), or from 2% to 15% or 20% (w/v), or from 5% to 15% (w/v) of a isotonizing agent
- (e) from 0% to 2% (w/v) preservatives; and
- (f) water for injection q.s. ad 100%.

In one embodiment, the aqueous suspensions may comprise by weight, based on the total volume of the composition:

- (a) from 3% to 50% (w/v), or from 10% to 40% (w/v), or from 10% to 30% (w/v), of bedaquiline (or pharmaceutically acceptable salt thereof);
- (b) from 0.5% to 10 % (w/v), or from 0.5% to 2% (w/v) of a wetting agent;
- (c) from 0% to 10% (w/v), or from 0% to 5% (w/v), or from 0% to 2% (w/v), or from 0% to 1% (w/v) of one or more buffering agents;

- (d) from 0% to 10 % (w/v), or from 0% to 6% (w/v) of a isotonizing agent
- (e) from 0% to 2% (w/v) preservatives; and
- (f) water for injection q.s. ad 100%.

5 To the suspensions may optionally be added an amount of acid or base to bring the pH to a value of about pH 7. Suitable acids or bases are any of those that are physiologically acceptable, e.g. HCl, HBr, sulfuric acid, alkali metal hydroxides such as NaOH. In an embodiment, such acid or base need not be added to the compositions of the invention.

10

The administration of bedaquiline (or pharmaceutically acceptable salt thereof) as in the present invention may suffice to treat a pathogenic mycobacterial infection although in a number of cases it may be recommendable to co-administer other anti-TB drugs.

15 In certain instances, the treatment of a pathogenic mycobacterial infection may be limited to only the administration of a composition of bedaquiline (and/or its metabolite thereof) in accordance with this invention, i.e. as monotherapy without co-administration of further anti-TB drugs. This option may be recommended, for example, for certain mycobacterial infections where a low concentration of the active
20 ingredient may treat the bacteria (e.g. for latent/dormant TB or for *Mycobacterium leprae*).

In a further aspect the present invention relates to the use of a pharmaceutical composition comprising an effective amount of bedaquiline or a pharmaceutically
25 acceptable salt thereof, in accordance with the present invention, for the manufacture of a medicament for maintenance therapy of a subject being infected with a pathogenic mycobacterial infection, wherein the composition is administered or is to be administered intermittently at a time interval that is in the range of one week to one year, or one week to two years.

30

Thus in a further aspect, the present invention provides a method for the long term treatment of a patient being infected with a pathogenic mycobacterial infection (e.g. drug-resistant or latent/dormant mycobacteria), said method comprising

- (i) the treatment of said patient with a combination of anti-TB drugs; followed by
- 35 (ii) the intermittent administration of a pharmaceutical composition comprising an effective amount of bedaquiline or a pharmaceutically acceptable salt thereof, in accordance with the present invention, wherein the composition is administered at a time interval of at least one week.

Where the treatment is directed towards *Mycobacterium leprae*, then again the treatment regime might be given as monotherapy or in combination with existing drugs useful for the treatment of *Mycobacterium leprae* (e.g. rifapentin). The composition of the invention might be administered by injection once, or up to three times, e.g. as monthly intervals. Advantages are associated with compliance, no resistance by avoiding dapsone, no stigma by avoiding clofazimine.

The present invention also concerns a pharmaceutical composition as described hereinbefore for use as a medicament in the treatment or prophylaxis of a pathogenic mycobacterial infection.

In addition, the present invention concerns the use of a pharmaceutical composition as described herein for the preparation of a medicament for the prophylaxis or treatment of a pathogenic mycobacterial infection.

The present invention further concerns a method of treating a subject infected with a pathogenic mycobacterial infection, said method comprising the administration of a therapeutically effective amount of a pharmaceutical composition as described herein.

As used herein, the word "substantially" does not exclude "completely" e.g. a composition which is "substantially free" from Y may be completely free from Y. Where necessary, the word "substantially" may be omitted from the definition of the invention. The term "about" in connection with a numerical value is meant to have its usual meaning in the context of the numerical value. Where necessary the word "about" may be replaced by the numerical value $\pm 10\%$, or $\pm 5\%$, or $\pm 2\%$, or $\pm 1\%$. All documents cited herein are incorporated by reference in their entirety.

The following examples are intended to illustrate the present invention and should not be construed as limiting the invention thereto.

EXAMPLES

Process Example: preparation of micro- and nano-suspensions

The active ingredient bedaquiline may be used as such or may be converted into a pharmaceutically acceptable salt thereof, such as a fumarate salt (for example the form used in the marketed product Sirturo®). Where referred to herein, bedaquiline is used in its non-salt form unless otherwise specified.

The prototype of the bedaquiline formulation is as follows:

Preparation of 200 and 100 mg/mL nano- and micro-suspensions.

5 Materials used:

Zirconium beads 0.5 mm (to aid process)

Sterile water for injection (Viaflo)

Bedaquiline (not milled/ground)

Surface modifier, including PEG4000 (or the like) and one or more other suitable

10 surface modifiers (e.g. Tocopheryl PEG 1000 succinate) – excipient(s)

Zirconium beads 2 mm (to aid process)

Mannitol (parenteral) – an excipient

Buffer (if needed) e.g. citrate-phosphate buffer

15 Glass bottles and ZrO₂ beads (either 0.5 mm or 2 mm, depending on the desired nano- or micro-suspensions), used as the milling media, were sterilized in an autoclave. The drug substance (quantity depending on the formulation to be prepared; see e.g. formulation/suspension below) was put into the glass bottle as well as a solution of surface modifier (e.g. PEG 4000 and tocopheryl PEG 1000 succinate) in water

20 (quantity depending on the concentration required/desired; see e.g. formulation/suspension below) for injection. ZrO₂-beads with an average particle size of 500 µm or 2 mm (depending on whether a micro- or nano-suspension is required/desired) were added. The bottle was placed on a roller mill. The suspension was micronized/nanonized at 100 rpm for a period of time up to 72 hours. For

25 instance, micronizing may be performed at 100 rpm for a period of 3 hours (or up to 3 hours) and nanonizing may be performed at 100 rpm for a period of up to 46 hours (e.g. about 40 hours). At the end of the milling process the concentrated micro- or nano-suspension was removed with a syringe and filled into vials. The resulting formulations (based on the nano-suspension and micro-suspension) are described in the following

30 tables. Determination of the concentration was done by HPLC/UV. If needed, a dilution was made to a final concentration of 200 mg/ml of active ingredient bedaquiline. The resulting suspension was shielded from light. Other concentrations were also made and tested, including 300 mg/ml and 100 mg/ml nano- and micro-

35 formulations.

Such formulations were (and will be) dosed intramuscular and subcutaneous in animals for PK study to investigate a possible long-acting effect (e.g. in treatment of leprosy).

Physical stability of the suspensions will be followed up by measuring particle size after different storage conditions.

Certain embodiments of the formulation(s) have the following features:

- 5 - Micro-suspension by using 2 mm Zr beads
- Milling at 200 mg/mL (otherwise the concentration may be too high, e.g. with 300 mg/ml)
- Longer milling, resulting in nano-suspension
- 10 - A suitable surface modifier, for instance selected based on physical stability, e.g. a surface modifier or wetting agent as described herein.

Reference examples of bedaquiline micro-suspensions

- 200 mg/ml micro-suspension referred to herein as Reference Example A (without
15 buffer) and Reference Examples B and C (with buffer)

Reference Example A

	mg/ml
Bedaquiline	200
TPGS	50
Mannitol	50
Sterile water for injection	q.s.

- 20 Particle Size Distribution (PSD)

Storage time	Storage temperature (°C)	D10 (µm)	D50 (µm)	D90 (µm)
0		0.820	1.99	4.96
1 month	25°C	0.704	1.59	3.64

- The PSD measurements after 1 month indicate that formulation remains relatively stable, and the Volume Density % is also depicted in Figure 1 (where “Concept 7”
25 refers to Reference Example A).

Stability Test using HPLC:

An HPLC test method was used to determine how stable the long acting injectable formulation of Reference Example A is. The purpose was to measure the amount of bedaquiline relative to two known degradants after certain periods of time at room temperature.

- 5 HPLC Procedure: Column – ProntoSIL 120-3-C18 SH, 100 mm length x 3.0 mm i.d., 3 µm particle size, or equivalent; column temperature 35°C; auto-sampler temperature 5°C; Flow rate 0.5 mL/min; Detection UV; Wavelength 230 nm; Data Collection Time 50 minutes; Analysis Run Time 60 minutes; Injection volume 10 µl; Mobile Phase A is 0.03 M Hydrochloric Acid in Water; Mobile Phase B is Methanol/Acetonitrile/2-Propanol – 45/45/10 (v/v/v).

Time (at room temperature)	Degradant A (% w/w)	Degradant B (% w/w)	Bedaquiline (% w/w)
Time zero	0.49	0.05	102.5
3 months	0.50	0.06	102.3
6 months	0.50	0.08	103.2

- The HPLC purity test shows that the formulation of Reference Example A is relatively stable for a long period (given that the relative amounts of degradants and bedaquiline remained stable).

Reference Example B and C

	Reference Example B mg/ml (where applicable)	Reference Example C mg/ml (where applicable)
Bedaquiline	200	200
TPGS	50	50
Buffer – citrate-phosphate	0.01N (pH 6)	0.05N (pH 6)
Mannitol	50	50
Sterile water for injection	q.s.	q.s.

- 20 Particle Size Distribution (PSD)

Reference Example	Conditions	D10 (µm)	D50 (µm)	D90 (µm)
B	After 3hrs milling	0.856	2.28	5.38

Reference Example	Conditions	D10 (μm)	D50 (μm)	D90 (μm)
C	After 3hrs milling	0.969	2.54	6.29
B	After autoclaving	1.13	2.29	4.72
C	After autoclaving	1.15	2.37	5.10
B	After 1 month at 25°C	1.13	2.29	4.68
C	After 1 month at 25°C	1.16	2.36	4.96
B	After autoclaving and 1 month at 5°C	1.12	2.27	4.65
C	After autoclaving and 1 month at 5°C	1.15	2.35	4.98

The PSD for these formulations under various conditions (including after autoclaving) indicate that the formulations remain relatively stable. This is shown in Figure 2, where Concept 3 refers to Reference Example B and Concept 4 refers to Reference Example C.

Example 1 – a micro-suspension of the invention

The suspensions of the Reference Examples all contain Vitamin E TPGS, which may not be tolerated parenterally, e.g. intramuscularly, particularly in the quantities specified (e.g. 50 mg/ml). The suspensions of the invention advantageously reduce the quantity of Vitamin E TPGS (as surface modifier), although it need not be completely replaced (e.g. as 5 mg/ml may be tolerated parenterally). PEG4000 (or polyethylene glycol 4000) is used, which can be supplied from Clariant GmbH. PEG4000 is a hydrophilic agent that can be used to increase the viscosity of the suspending vehicle and can act as a suspending agent.

Example 1 formulation:

	mg/ml
--	-------

Bedaquiline	200
PEG4000	50
TPGS	5
Buffer – citrate-phosphate	0.01N (pH 6)
Mannitol	50
Sterile water for injection	q.s.

In this case a buffer was added to avoid a drop in pH.

Particle Size Distribution (PSD)

Conditions	D10 (µm)	D50 (µm)	D90 (µm)
Before autoclaving	1.27	3.29	10.4
After autoclaving	1.22	2.74	6.72

5

The PSD of the micro-suspension of Example 1 shows that the formulation remains relatively stable after autoclaving. This is shown in Figure 3.

10

The approximate cloud point of the formulation of Example 1 was calculated to be about 105 to 110°C.

The autoclaving of the micro-suspension of Example 1 was conducted in a Systec autoclave (VX/VE series), where the present parameters are:

Sterilization temperature: 121°C (above the calculated cloud point)

15

Sterilization time: 15 minutes

Unloading temperature: 80°C

20

A typical autoclaving cycle – the steam generator builds up the required steam pressure and the steam flows into the sterilization chamber, after the sterilization temperature has been reached, it then remains constant for the duration of the sterilization period, and after the period has elapsed the cycles with the optional built-in cooling apparatus are cooled down until the unloading temperature has been reached.

25

Given that the autoclaving temperature is higher than the measured cloud point, it could be expected that particle aggregation would be seen, for instance due to phase-separation.

Further Data on Micro-suspension of Example 1

Particle Size Distribution (PSD) after suspension is subjected to certain conditions

Conditions (all after autoclaving)	D10 (μm)	D50 (μm)	D90 (μm)	Resuspendability
18 days at 5°C	1.132	2.399	5.510	Not tested
1 month at 5°C	1.116	2.384	5.488	30 seconds
1 month at 25°C	1.116	2.406	5.700	30 seconds
1 month at 40°C	1.114	2.384	5.474	30 seconds
1 month at 60°C	1.142	2.430	5.624	30 seconds

- 5 The above data on PSD also show that the micro-suspension of Example 1 remains relatively stable after autoclaving and after further time (and at varying temperatures), which is also outlined in Figure 4.

- Re-suspendability: after autoclaving, particles can be seen at the bottom of the vessel,
 10 which must therefore be shaken. Advantageously, it was seen that, where tested, the formulation of Example 1 could easily be resuspended after shaking.

Further Data on Micro-suspension of Example 1

- 15 Particle Size Distribution (PSD) after suspension is subjected to certain further conditions

Conditions	D10 (μm)	D50 (μm)	D90 (μm)
After 4 hours milling, before autoclaving	1.10	3.40	10.6
After autoclaving, 18 days at 5°C	1.13	2.4	5.51
1 month at 25°C	1.12	2.41	5.70
3 months at 25°C	1.12	2.40	5.55
3 months at 60°C	1.20	2.52	5.82

- It can be seen that even after autoclaving, there is a stable PSD, even up to 3 months at
 20 60°C, which is outlined in Figure 5.

The HPLC test method above was used to determine how stable the long acting injectable formulation of Example 1 is. Again, the purpose was to measure the amount of bedaquiline relative to known degradants/impurities after certain periods of time at room temperature and it gave the following results:

5

Conditions (all after autoclaving)	RRT0.15 (%w/w)	Degradant A (%w/w)	Bedaquiline (%w/w)	RRT1.24 (%w/w)	Degradant B (%w/w)	RRT1.64 (%w/w)
Time zero, 5°C	<0.05%	0.13%	73.0%	<0.05%	0.14%	<0.05%
14 days at 5°C	<0.05%	0.13%	72.8%	<0.05%	0.13%	<0.05%
14 days at 40°C	<0.05%	0.13%	72.8%	<0.05%	0.13%	<0.05%
14 days at 60°C	<0.05%	0.13%	72.6%	<0.05%	0.16%	<0.05%
1 month at 5°C	<0.05%	0.12%	72.0%	<0.05%	0.14%	<0.05%
1 month at 40°C	<0.05%	0.12%	72.5%	<0.05%	0.14%	<0.05%
1 month at 60°C	<0.05%	0.12%	72.6%	<0.05%	0.17%	<0.05%
3 months at 5°C	<0.05%	0.12%	72.06%	<0.05%	0.13%	<0.05%
3 months at 40°C	<0.05%	0.12%	71.76%	<0.05%	0.13%	<0.05%
3 months at 60°C	<0.05%	0.11%	71.43%	<0.05%	0.24%	<0.05%

Conclusions

A key conclusion is that, the suspensions of the Reference Example and of Example 1 are stable, as determined by PSD and purity determination in the HPLC test method, even after autoclaving, after storage for a certain amount of time and at high temperatures.

10

A further key conclusion was that the suspensions of Example 1 were easily re-suspendable after autoclaving, even after storage for a certain amount of time and at high temperatures.

5 Further re-suspendability data

Composition	PEG4000 mg/ml	Auto- claved	Storage temp	Re- suspendability
200 mg/ml bedaquiline; 10 mg/ml TPGS; 50 mg/ml mannitol; 0.01N citrate-phosphate buffer pH 6.0	0	Yes	25°C	Difficult
200 mg/ml bedaquiline; 10 mg/ml TPGS; 45 mg/ml mannitol, 0.01N citrate-phosphate buffer pH 6.0	50	Yes	25°C	Easy
200 mg/ml bedaquiline; 5 mg/ml TPGS, 50 mg/ml mannitol, 0.01N citrate-phosphate buffer pH 6.0	50	Yes	25°C	Easy

The re-suspendability of the above compositions was objectively tested, after autoclaving in the above examples, by swirling the relevant composition. It was found that the composition without PEG4000 was difficult to re-suspend (here, it took more than 40 seconds to re-suspend, and required swirling and shaking), whereas the compositions with PEG4000 were relatively easy to re-suspend (here, they required less than 40 seconds of gentle swirling).

15 **Example 1A (microsuspension)**

	Example 1A mg/ml (where applicable)
Bedaquiline	200
TPGS	10
PEG4000	50
Mannitol	QS
Buffer – Citrate phosphate	0.05 M (pH 6)

Particle Size Distribution (PSD) and resuspendability (Example 1A)

	Storage temperature	Time point	PSD			Resuspendability
			Dv(10)	Dv(50)	Dv(90)	
Before autoclavation		T0	0.953	2.351	5.069	<u>Good (< 10 sec shaking)</u>
	40°C	3M	1.120	2.546	5.782	<u>Good (< 10 sec shaking)</u>
After autoclavation		T0	1.093	2.375	5.196	<u>Good (< 10 sec shaking)</u>
	25°C	6M	1.121	2.384	5.112	<u>Good (< 10 sec shaking)</u>
	40°C	1M	1.159	2.450	5.640	<u>Good (< 10 sec shaking)</u>
	40°C	3M	1.104	2.378	4.832	<u>Good (< 10 sec shaking)</u>
	40°C	6M	1.139	2.403	4.962	<u>Good (< 10 sec shaking)</u>
	50°C	1M	1.180	2.491	5.800	<u>Good (< 10 sec shaking)</u>

Example 1B (microsuspension)

5

	Example 1B mg/ml (where applicable)
Bedaquiline	200
TPGS	20
PEG4000	50
Mannitol	QS
Buffer -- Citrate phosphate	0.05 M (pH 6)

Particle Size Distribution (PSD) and resuspendability (Example 1B)

	Storage temperature	Time point	PSD			Resuspendability
			Dv(10)	Dv(50)	Dv(90)	
Before autoclavation						<u>Good (< 10 sec shaking)</u>
	40°C	3M	0.849	2.238	4.915	<u>Good (< 10 sec shaking)</u>
After autoclavation		T0	0.895	2.117	4.716	<u>Good (< 10 sec shaking)</u>
	25°C	6M	0.882	2.074	4.376	<u>Good (< 10 sec shaking)</u>
	40°C	1M	0.920	2.120	4.713	<u>Good (< 10 sec shaking)</u>
	40°C	3M	0.867	2.098	4.784	<u>Good (< 10 sec shaking)</u>
	40°C	6M	0.889	2.072	4.254	<u>Good (< 10 sec shaking)</u>
	50°C	1M	0.940	2.144	4.856	<u>Good (< 10 sec shaking)</u>

Example 1C (microsuspension)

5

	Example 1C mg/ml (where applicable)
Bedaquiline	200
TPGS	10
PEG4000	25
Mannitol	QS
Buffer – Citrate phosphate	0.05 M (pH 6)

Particle Size Distribution (PSD) and resuspendability (Example 1C)

			PSD			Resuspendability
	Storage temperature	Time point	<u>Dv(10)</u>	<u>Dv(50)</u>	<u>Dv(90)</u>	
Before autoclavation		T0	0.889	2.216	4.490	<u>Good (< 10 sec shaking)</u>
After autoclavation		T0	1.027	2.240	4.804	<u>Good (< 10 sec shaking)</u>
	25°C	6M	1.031	2.210	4.529	<u>Good (< 10 sec shaking)</u>
	40°C	1M	1.075	2.276	4.993	<u>Good (< 10 sec shaking)</u>
	40°C	3M	1.025	2.253	4.721	<u>Good (< 10 sec shaking)</u>
	40°C	6M	1.058	2.247	4.540	<u>Good (< 10 sec shaking)</u>
	50°C	1M	1.084	2.276	4.796	<u>Good (< 10 sec shaking)</u>

Example 1D (microsuspension)

5

	Example 1D mg/ml (where applicable)
Bedaquiline	200
Poloxamer 338	50
PEG4000	50
Mannitol	QS
Buffer – Citrate phosphate	0.05 M (pH 6)

Particle Size Distribution (PSD) and resuspendability (Example 1D)

			PSD			Resuspendability
	Storage temperature	Time point	<u>Dv(10)</u>	<u>Dv(50)</u>	<u>Dv(90)</u>	
Before autoclavation		T0	1.155	2.733	6.537	<u>Good (< 10 sec shaking)</u>

			PSD			Resuspendability
	Storage temperature	Time point	<u>Dv(10)</u>	<u>Dv(50)</u>	<u>Dv(90)</u>	
After autoclavation		T0	1.438	3.067	7.269	<u>Good (< 10 sec shaking)</u>
	25 °C	6M	1.456	3.027	6.319	<u>Good (< 10 sec shaking)</u>
	40 °C	6M	1.493	3.137	7.005	<u>Good (< 10 sec shaking)</u>

Example 1E (microsuspension)

	Example 1E mg/ml (where applicable)
Bedaquiline	200
TPGS	5
Tween 20	5
PEG4000	50
Mannitol	QS
Buffer – Citrate phosphate	0.05 M (pH 6)

5 **Particle Size Distribution (PSD) and resuspendability (Example 1E)**

			PSD			Resuspendability
	Storage temperature	Time point	<u>Dv(10)</u>	<u>Dv(50)</u>	<u>Dv(90)</u>	
Before autoclavation		T0	0.867	2.002	4.406	<u>Good (< 10 sec shaking)</u>
After autoclavation		T0	0.978	2.035	4.319	<u>Good (< 10 sec shaking)</u>
	25 °C	6M	0.985	2.006	3.961	<u>Good (< 10 sec shaking)</u>
	40 °C	6M	1.007	2.022	3.928	<u>Good (< 10 sec shaking)</u>

Reference Example 1F (microsuspension)

	Example 1F mg/ml (where applicable)
Bedaquiline	200
TPGS	20
PEG4000	50
Sodium chloride	QS
Buffer – Citrate phosphate	0.05 M (pH 6)

Particle Size Distribution (PSD) and resuspendability (Example 1F)

			PSD			Resuspendability
			<u>Dv(10)</u>	<u>Dv(50)</u>	<u>Dv(90)</u>	
Before autoclavation		T0	0.766	1.976	4.275	<u>Good (< 10 sec shaking)</u>
	40°C	3M	0.898	2.119	4.363	<u>Good (< 10 sec shaking)</u>
After autoclavation		T0	0.912	2.011	4.230	<u>Good (< 10 sec shaking)</u>
	25°C	6M	0.923	2.013	4.207	<u>Good (< 10 sec shaking)</u>
	40°C	1M	0.961	2.039	4.200	<u>Good (< 10 sec shaking)</u>
	40°C	3M	0.911	2.028	4.221	<u>Good (< 10 sec shaking)</u>
	40°C	6M	0.938	2.033	4.201	<u>Good (< 10 sec shaking)</u>
	50°C	1M	0.967	2.038	4.134	<u>Good (< 10 sec shaking)</u>

5

Example 2: pharmacokinetic studies**Pharmacokinetic studies in mouse, rat and beagle dog**

A number of studies in mouse, rat and beagle dog are described in international patent application WO 2019/012100, which generally demonstrate that a sustained plasma concentration of bedaquiline and/or its active metabolite M2 were seen over certain

10

periods of time (including 1 month, 3 months and 6 months) using e.g. the formulation of Reference Example A.

Pharmacokinetic profile in rats

- 5 Formulations of concentrations 200 mg/mL were used in this study, and the micro-suspension of Reference Example A was used, i.e. using, in addition to the 200 mg/ml concentration of micro-particles (of the active bedaquiline), TPGS (4:1 bedaquiline : TPGS) and 50 mg/ml Mannitol in WFI (water for injection), without buffer. Bedaquiline is also referred to as TMC207.

10

These studies demonstrate that Reference Example A resulted in stable plasma levels over a prolonged period of time in male rats, when administered subcutaneously (SC) and intramuscularly (IM).

Male Rats

The first experiment was performed on male rats, where each relevant 200 mg/ml nano-suspension and micro-suspension referred to above were administered subcutaneously (SC) and intramuscularly (IM) at a concentration of 40 mg/kg (0.2 mL/kg). An interim analysis was performed at 3 months and the results were followed-up at 6 months.

- 20 Twelve rats were used in the study. Three rats were dosed intramuscularly (IM) with the 200 mg/ml micro-suspension (see Reference Example A). Three rats were dosed subcutaneously (SC) with the 200 mg/ml micro-suspension (see Reference Example A).

Phase 1 of the Results – up to 2200 hours

Figure 6 “Plasma kinetics of TMC207 in male rats when administered IM or SC with 200 mg/ml micro-formulation (see Reference Example A) at a dose of 40 mg/kg”

The following parameters were calculated for TMC207 (see Figure):

30

	Microsuspension SC	Microsuspension IM
n	3	3
C_{max} (ng/ml)	68.1 ± 17.6	215 ± 66.7
T_{max}^a (h)	24 (24.00 - 24.00)	18 (7.00 - 24.00)
T_{last}^a = around 3 mths (h)	2184 (2184 - 2184)	2184 (2184 - 2184)

	Microsuspension SC	Microsuspension IM
AUC _{0-2184h} (3 mths) (ng.h/ml)	34700 ± 1770	91500 ± 13200

where applicable mean values are given (with min → max in parentheses)

Phase 2 of the Results – up to 4400 hours

In all cases the plasma concentration of BDQ or M2 is calculated as the mean of the three rats in the relevant study.

Study in rats: for formulation of Reference Example A, i.e. the micro-suspension of 200 mg/ml concentration, and dosed SC at 40 mg/kg (StDev = standard deviation) and IM at 40 mg/kg

Time (h)	Plasma concentration of bedaquiline (BDQ) or its metabolite (M2)							
	SC at 40 mg/kg				IM at 40 mg/kg			
	BDQ	StDev	M2	StDev	BDQ	StDev	M2	StDev
1	22.1	5.36	0.589	NC	139	33.2	5.11	2.24
4	36.9	7.42	6.62	1.69	172	47.2	18.8	5.98
7	40.7	6.27	8.59	1.46	185	24.2	28.7	7.57
24	68.1	17.6	24.0	1.14	212	70.6	77.3	23.5
168	16.5	4.84	9.03	1.81	98.0	19.1	91.5	33.1
336	18.1	3.30	9.28	2.10	69.7	10.2	54.9	16.8
504	22.8	3.96	9.68	2.85	52.2	4.05	37.9	16.5
672	14.7	0.964	7.32	1.34	42.6	6.85	29.5	14.8
840	15.1	1.74	7.40	2.46	33.6	6.39	22.3	10.6
1008	14.7	3.47	6.82	2.06	28.5	6.24	20.2	11.2
1176	13.2	2.99	5.96	1.76	24.1	7.04	16.6	9.37
1344	12.4	2.34	6.10	1.79	20.7	3.07	14.1	8.88
1512	12.0	0.917	5.81	1.81	19.7	5.98	13.4	7.16
1680	12.3	1.95	5.42	2.01	18.4	3.30	11.5	5.77
1848	10.6	0.83	5.18	1.51	14.3	1.35	11.4	6.39
2016	9.83	2.06	4.30	2.03	14.9	1.75	9.86	3.75
2184	10.2	2.42	4.55	1.36	12.6	0.755	8.87	3.37
2520	9.45	2.16	5.54	1.82	11.6	2.06	9.27	3.53
2856	8.26	0.737	4.78	1.61	10.5	2.65	7.49	3.02
3192	6.82	1.38	4.04	1.02	8.67	2.71	6.28	2.52

Time (h)	Plasma concentration of bedaquiline (BDQ) or its metabolite (M2)							
	SC at 40 mg/kg				IM at 40 mg/kg			
	BDQ	StDev	M2	StDev	BDQ	StDev	M2	StDev
3528	6.83	2.27	4.02	1.05	6.92	2.09	5.68	2.79
3864	6.69	0.794	3.95	0.866	5.90	2.21	5.00	2.53
4200	6.41	1.72	3.49	0.987	4.41	2.04	3.74	2.03
≈ CV%	8 – 29		NC – 47		6 – 46		26 – 63	
T max (h)	24		24		18		120	83
Cmax (ng/mL)	68.1	17.6	24.0	1.14	215	66.7	94.2	33.3
T last (h)	4200		4200		4200		4200	
AUClast (ng*h/mL)	50200	4240	24800	5520	109000	12300	75200	28700
AUC ₀₋₂₈₅₆ (ng*h/mL)	41000	2880	19300	4150	99200	13200	67600	26100
AUCinf (ng*h/mL)	NC		NC		121000	11400	85500	28800

Plasma Concentration vs Time – profiles for Reference Example A and Example 1

In the following figures, it is shown that the plasma concentration versus time profiles of the Reference Example A (labelled F4) and Example 1 (labelled F1) were studied in rats after SC injection of 40 mg/kg. The concentration of bedaquiline and its active metabolite M2 were measured.

Sustained plasma concentrations of the parent compound above the LLOQ (lower limit of quantification) were observed in all animals in all groups for the duration of the study. Within the first 28 days after SC administration, 2 plasma concentration peaks (C_{max}) of the parent compound were observed for both formulations F1 and F4. After 28 days, general converging of drug plasma concentrations to similar profiles and concentrations over time occurred for both formulations.

Figure 7 shows plasma concentration versus time profiles of subcutaneous administered bedaquiline LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats.

Regarding plasma concentration-time profiles of the M2 metabolite, again sustained plasma concentrations of M2 above the LLOQ were observed in all animals for both F1 and F4.

Figure 8 shows plasma concentration versus time profiles of bedaquiline (BDQ) metabolite after subcutaneous administration of BDQ LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats.

After intramuscular administration, again sustained plasma concentrations of the parent
5 above the LLOQ was observed in all animals for both F1 and F4 formulations for the duration of the study.

Figure 9 shows plasma concentration versus time profiles of intramuscular administered bedaquiline LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats.

10 Similarly, sustained plasma concentrations were achieved for the metabolite after intramuscular administration.

Figure 10 shows plasma concentration versus time profiles of bedaquiline (BDQ) metabolite after intramuscular administration of BDQ LAI microsuspensions containing different surfactants (PEG 4000 combined with TPGS, and TPGS) in rats.

15 **Conclusion:** both formulations of Reference Example A (F4) and Example 1 (F1) were effective, in their own way, in achieving sustained release of both drug and an active metabolite (M2) and were both therefore considered to be suitable for such purpose.

Example 3

20 Evaluation of an injectable, long-acting bedaquiline formulation in the paucibacillary mouse model of latent tuberculosis infection

The objective of this study was to use the paucibacillary mouse model of latent tuberculosis infection (LTBI) to compare the bactericidal activity of a long-acting bedaquiline (B_{LA}) formulation administered every 4 weeks for a total of 1, 2, or 3 doses
25 to the activity of daily (5 days per week) oral dosing of B at the standard 25 mg/kg dose or lower doses matching to total drug doses administered as B_{LA}. The original study scheme is presented in **Table 1**. The B_{LA} used for this study is that described above in Reference Example A, i.e. the microsuspension at a concentration of 200 mg/ml). The primary outcome was the decline in *Mycobacterium tuberculosis* lung CFU counts
30 during treatment.

Table 1. Original study scheme to evaluate the bactericidal activity of B_{LA} in a mouse model of paucibacillary LTBI.

LTBI treatment regimen*	Number of mice sacrificed for lung CFU counts at the following time points:						Total mice	Total dose B over 12 weeks (mg/kg)
	BCG immunization	<i>M.tb.</i> challenge	Treatment initiation	During treatment				
	Week -12	Week -6	Day 0	Week 4	Week 8	Week 12		
Untreated	5	5	5	5	5	5	30	na
R ₁₀ (5/7)				5	5	5	15	na
P ₁₅ H ₅₀ (1/7)				5	5	5	15	na
B ₂₅ (5/7)				5	5	5	15	1500
B ₈ (5/7)				5	5	5	15	480
B _{5.33} (5/7)				5	5	5	15	320
B _{2.67} (5/7)				5	5	5	15	160
B _{LA-160} (1/28) x 3						5	5	480
B _{LA-160} (1/28) x 2					5	5	10	320
B _{LA-160} (1/28) x 1				5	5	5	15	160
Total mice	5	5	5	40	45	50	150	

- 5 *R, rifampin; P, rifapentine; H, isoniazid; B, bedaquiline; B_{LA}, long-acting bedaquiline formulation. All drug doses in mg/kg indicated by subscript. Fractions in parentheses indicate dosing frequency, in days. B_{LA} is administered by intramuscular injection; all other drugs are administered by gavage. na, not applicable.

10 **Justification of the regimens**

- Untreated mice were used to determine the level and stability of the paucibacillary infection.
- R₁₀ (5/7) is an alternative regimen for treatment of LTBI in the US and Canada, administered for 4 months. It was used here as a control to qualify the model.
- 15 ○ P₁₅H₅₀ (1/7) is an alternative regimen for treatment of LTBI in the US, administered once weekly for 3 months (12 doses). It proved at least as efficacious as 9 months of isoniazid. It is the most intermittent of currently recommended regimens and serves as a second control.
- B₂₅ (5/7) is daily B at the human equivalent dose previously studied in the
- 20 paucibacillary model. It provides a total dose of 500 mg/kg every 28 days.

- B₈ (5/7) is daily B at a dose that is reduced to provide the same total dose (480 mg/kg) as the B_{LA} formulation dose (*i.e.*, 160 mg/kg) administered every 28 days × 3 doses.
- B_{5.33} (5/7) is daily B at a dose that is reduced to provide the same total dose (320 mg/kg) as the B_{LA} formulation dose (*i.e.*, 160 mg/kg) administered every 28 days × 2 doses.
- B_{2.67} (5/7) is daily B at a dose that is reduced to provide the same total dose (160 mg/kg) as the B_{LA} formulation dose (*i.e.*, 160 mg/kg) administered once.
- B_{LA-160} (1/28) × 3 is the B_{LA} formulation administered as 160 mg/kg every 28 days for a total of 3 doses. Thus, the total B dose will match that of the B₈ (5/7) group at each 28-day interval.
- B_{LA-160} (1/28) × 2 is the B_{LA} formulation administered as 160 mg/kg every 28 days for a total of 2 doses, beginning on Day 0. Thus, the total B dose administered by Week 12 (320 mg/kg) will be the same as that of the B_{5.33} (5/7) group.
- B_{LA-160} (1/28) × 1 is the B_{LA} formulation administered as 160 mg/kg just once on Day 0. Thus, the total B dose administered by Week 12 (160 mg/kg) will be the same as that in the B_{2.67} (5/7) group.

FINAL RESULTS

- All CFU count data are finalized and presented below in **Table 2**. Due to delays in finalizing institutional agreements and obtaining the B_{LA} supply, treatment was not initiated until approximately 13 weeks after the *M. tuberculosis* challenge infection, and the time line in **Table 2** has been adjusted accordingly. For comparison between different treatment groups, statistical significance was assessed using one-way ANOVA adjusted with Bonferroni's multiple comparisons test.

Table 2. Final *M. tuberculosis* lung CFU count data.

LTBI treatment regimen*	Mean (SD) log ₁₀ <i>M. tuberculosis</i> CFU/lung at the following time points:						Total dose B over 12 weeks (mg/kg)
	BCG immunization	<i>M.tb.</i> challenge	Treatment initiation	During treatment			
	Week -19	Week -13	Day 0	Week 4	Week 8	Week 12	
Untreated	na	2.11 (0.09)	4.75 (0.27)	4.71 (0.48)	4.60 (0.27)	4.94 (0.29)	na
R ₁₀ (5/7)				3.39 (0.46)	2.74 (0.62)	1.27 (0.85)	na
P ₁₅ H ₅₀ (1/7)				2.67 (0.25)	0.79 (0.80)	0.28 (0.41)	na
B ₂₅ (5/7)				3.01 (0.45)	0.82 (0.49)	0.07 (0.09)	1500
B ₈ (5/7)				3.30 (0.12)	2.42 (0.26)	0.69 (0.43)	480
B _{5.33} (5/7)				3.83 (0.25)	3.15 (0.47)	1.98 (0.17)	320
B _{2.67} (5/7)				3.96 (0.35)	3.52 (0.38)	3.16 (0.24)	160
B _{LA-160} (1/28) x 3						1.23 (0.16)	480
B _{LA-160} (1/28) x 2					2.31 (0.40)	1.63 (0.40)	320
B _{LA-160} (1/28) x 1				3.55 (0.32)	3.31 (0.38)	1.83 (0.34)	160

*R, rifampin; P, rifapentine; H, isoniazid; B, bedaquiline, B_{LA}, long-acting bedaquiline formulation. All drug doses in mg/kg indicated by subscript. Fractions in parentheses indicate dosing frequency, in days. SD, standard deviation. na, not applicable.

BCG immunization. One-hundred fifty female BALB/c mice were infected by aerosol with *M. bovis* rBCG30. A culture suspension with an OD₆₀₀ of 1.03 was diluted 10-fold and then used for aerosol infection. The concentration of the bacterial suspension was 6.88 log₁₀ CFU/mL, which resulted in a mean implantation of 3.05 (SD 0.10) log₁₀ CFU/lung. Six weeks later, at the time of the *M. tuberculosis* challenge infection, the mean BCG burden in the mouse lungs was 4.95 (SD 0.11) log₁₀ CFU. By Day 0, the BCG burden had decreased and stabilized at 3.27 (SD 0.45) log₁₀ CFU/lung, with similar lung burdens observed in the untreated mice at Weeks 4, 8, and 12. Thus, a low-level, stable BCG infection was established in the lungs of these mice as expected.

***M. tuberculosis* challenge.** Six weeks after BCG immunization, mice were infected by aerosol with *M. tuberculosis* H37Rv. A culture suspension with an OD₆₀₀ of 0.850 was diluted ~100-fold and then used for aerosol infection. The concentration of the bacterial suspension was 4.73 log₁₀ CFU/mL, which resulted in a mean implantation of 2.11 (SD 0.09) log₁₀ CFU/lung. This implantation was approximately 1 log₁₀ CFU higher than was intended. By Day 0, the *M. tuberculosis* burden had stabilized at around

4.8 log₁₀ CFU/lung, with similar lung burdens observed in the untreated mice at Weeks 4, 8, and 12. Thus, despite the higher implantation, a stable *M. tuberculosis* infection was established in the lungs of these mice, with the stabilized lung CFU burden correspondingly nearly 1 log₁₀ CFU higher than observed in previous experiments (1-3).

Assessment of bactericidal activity (Table 2). Compared to the *M. tuberculosis* CFU counts in the lungs of untreated mice, the R₁₀ (5/7) control regimen reduced the mean CFU count by approximately 1, 2 and 3 log₁₀ CFU/lung after 4, 8, and 12 weeks of treatment, respectively. The P₁₅H₅₀ (1/7) control regimen resulted in reductions of about 2, 3, and 4.5 log₁₀ CFU after 4, 8, and 12 weeks of treatment, respectively. The relative magnitudes of the decline in lung CFU counts for both control regimens are as expected based on previous studies (1,2). B₂₅ (5/7) resulted in a reduction of about 1.7, 4.0, and 4.9 log₁₀ CFU/lung after 4, 8, and 12 weeks of treatment, results which were also expected based on previous studies (1-2). Thus, the higher implantation and Day 0 CFU counts did not affect the relative activity of the drugs against this stabilized bacterial population in the mouse lungs.

For all B test regimens, there was increasing activity with increasing dose observed at Weeks 4, 8, and 12. For mice that received one or two doses of B_{LA-160} (1/28), the decrease in lung CFU counts relative to untreated mice was equivalent to the decrease in mice that received the same total dose administered as a daily oral regimen, B₈ (5/7), for 4 or 8 weeks, respectively ($p > 0.05$ at both time points). One dose of B_{LA-160}, delivering 160 mg/kg at Day 0, resulted in a decline of about 1.3 log₁₀ CFU/lung, and four weeks of B₈ (5/7) resulted in a decline of about 1.5 log₁₀ CFU/lung. After two doses of B_{LA-160} (1/28) or 8 weeks of B₈ (5/7), the CFU counts in the lungs decreased by an additional 1 log₁₀ in mice that received either of these regimens. After 12 weeks of treatment, the CFU counts in the lungs were lower in mice that had received one dose of B_{LA-160} than in the mice that received the same total dose of bedaquiline (160 mg/kg) via daily dosing with B_{2.67} (5/7) ($p = 0.0002$), with the former regimen resulting in a decline of about 3 log₁₀ CFU/lung and the latter resulting in a decline of 1.7 log₁₀ CFU/lung, compared to the lung counts in the untreated control mice. In mice that received a total bedaquiline dose of 320 mg/kg, either through two doses of B_{LA-160} or through daily dosing of B_{5.33} (5/7), the decline in lung CFU counts was the same at about 3 log₁₀ CFU/lung ($p > 0.05$). For mice that received a total bedaquiline dose of 480 mg/kg via three doses of B_{LA-160} (1/28), the lung CFU counts were higher than in mice that received the equivalent total dose through daily dosing with B₈ (5/7), although the difference was not statistically significant.

After 12 weeks of treatment, nearly all test regimens had equivalent bactericidal activity as the R₁₀ (5/7) control regimen, with only the B_{2.67} (5/7) regimen being significantly less bactericidal than this control ($p < 0.0001$). The test regimen B₈ (5/7) demonstrated equivalent bactericidal activity to both the P₁₅H₅₀ (1/7) and B₂₅ (5/7) control regimens, while all other test regimens were significantly less bactericidal than either of these control regimens at Week 12. However, CFU data recorded at the Week 12 time point may not reflect the overall efficacy of long-acting bedaquiline regimens. In mice that received a single dose of B_{LA-160} on Day 0, bacterial killing was still observed 12 weeks after administration. Thus, it is conceivable that the bacterial burden in the lungs of mice that received 2 and 3 doses of B_{LA-160} would still further decrease for at least 12 weeks after administration of the last dose (if not longer). Also of interest is that the single dose of B_{LA-160} seemed to exert greater bactericidal activity from weeks 1 to 4 and from weeks 9 to 12, compared to weeks 5 to 8 post-administration, suggesting the possibility of biphasic B release kinetics from the long-acting vehicle.

CONCLUSIONS

- Despite a higher bacterial implantation than anticipated, a stable *M. tuberculosis* infection was established in BALB/c mice that was suitable for evaluation of LTBI treatment regimens.
- After 12 weeks of treatment, once-monthly dosing with B_{LA-160} demonstrated superior or equivalent bactericidal activity compared to daily dosing for total bedaquiline doses of 160 or 320 and 480 mg/kg, respectively.
- The bactericidal activity observed from a single dose of B_{LA-160} was evident for at least 12 weeks after administration, and likely CFU counts would continue to decrease in the lungs of mice that received 2 and 3 doses. Taken together with the higher-than-expected baseline bacterial burden in this experiment, these findings suggest that cure after 2 or 3 injections may be possible. Thus, it will be critical to evaluate the sterilizing activity of these B_{LA} regimens over longer time periods to truly understand their potential for use in LTBI treatment.

REFERENCES

- 1) Zhang, T., Li, S., Williams, K., Andries, K., Nuermberger, E. 2011. Short-course chemotherapy with TMC207 and rifapentine in a murine model of latent tuberculosis infection. *Am. J Respir. Crit. Care Med.* 184:732-737.
- 5 2) Lanoix, J.P., Betoudji, F., Nuermberger, E. 2014. Novel regimens identified in mice for treatment of latent tuberculosis infection in contacts of multidrug-resistant tuberculosis cases. *Antimicrob. Agents Chemother.* 58:2316-2321.
- 3) Zhang, T., M. Zhang, I. M. Rosenthal, J. H. Grosset, and E. L. Nuermberger. 2009. Short-course therapy with daily rifapentine in a murine model of latent tuberculosis
10 infection. *Am.J Respir.Crit Care Med.* 180:1151-1157.

Claims

1. A pharmaceutical composition for administration by intramuscular or
subcutaneous injection, comprising a therapeutically effective amount of
5 bedaquiline, or a pharmaceutically acceptable salt thereof, in the form of a
suspension of micro- or nanoparticles comprising:
 (a) bedaquiline, or a pharmaceutically acceptable salt thereof, in micro- or
 nanoparticle form, and a surface modifier; and
 (b) a pharmaceutically acceptable aqueous carrier
10 which is characterised in that the surface modifier comprises PEG4000 or the
 like.
2. A composition according to claim 1, wherein the surface modifier comprises at
least 75% by weight PEG4000 or the like and the remainder is one or more other
15 suitable surface modifiers
3. A composition according to claim 2, wherein the one or more other suitable
surface modifiers are selected from the group of poloxamers, α -tocopheryl
polyethylene glycol succinates, polyoxyethylene sorbitan fatty acid esters, and
20 salts of negatively charged phospholipids.
4. A composition according to claim 3, wherein the other surface modifiers represent
one surface modifier that is an α -tocopheryl polyethylene glycol succinate (TPGS).
- 25 5. A composition according to any of claims 1 to 4, wherein bedaquiline is in its non-
salt or free form or in the form of a fumarate salt.
6. A composition according to any of claims 1 to 5, wherein the average effective
particle size of the bedaquiline, or a pharmaceutically acceptable salt thereof,
30 micro- or nanoparticles is below about 50 μm , in particular below about 200 nm.
7. A composition according to claims 1 or 2, comprising by weight based on the total
volume of the composition:
 (a) from 10% to 70% (w/v), or from 20% to 60% (w/v), or from 20% to 50%
35 (w/v), or from 20% to 40% (w/v) of bedaquiline (or pharmaceutically
acceptable salt thereof; but where the w/v is calculated on the basis of its
non-salt form);

- (b) from 0.5% to 20 % (w/v), or from 2% to 15% or 20% (w/v), or from 5% to 15% (w/v) of a wetting agent (or surface modifier, i.e. comprising PEG4000 or the like);
- (c) from 0% to 10% (w/v), or from 0% to 5% (w/v), or from 0% to 2% (w/v), or from 0% to 1% (w/v) of one or more buffering agents;
- (d) from 0% to 20 % (w/v), or from 2% to 15% or 20% (w/v), or from 5% to 15% (w/v) of a isotonizing agent
- (e) from 0% to 2% (w/v) preservatives; and
- (f) water for injection q.s. ad 100%.
8. The use of a pharmaceutical composition as defined in any of claims 1 to 7, for the manufacture of a medicament for the treatment of a pathogenic mycobacterial infection.
9. The use of claim 8 wherein the medicament is for the long-term treatment of *Mycobacterium tuberculosis* (such as the drug-resistant or latent/dormant form) or *Mycobacterium leprae*.
10. The use according to claim 8 wherein the medicament is for administration by intramuscular or subcutaneous injection; wherein the composition is administered intermittently at a time interval of one week to two years.
11. The use according to claim 8 wherein the pharmaceutical composition is administered at an interval of at least one month to one year.
12. The use according to claim 8, wherein the pharmaceutical composition is administered at a time interval that is in the range of one week to one month, or in the range of one month to three months, or in the range of three months to six months, or in the range of six months to twelve months, or in the range of 12 months to 24 months.
13. The use according to claim 8, wherein the pharmaceutical composition is administered once every two weeks, or once every month, or once every three months.

14. A process for preparing a pharmaceutical composition as defined in any of claims 1 to 7, comprising
- (a) obtaining bedaquiline, or a pharmaceutically acceptable salt thereof, in micronized form;
 - 5 (b) adding the micronized bedaquiline, or a pharmaceutically acceptable salt thereof, to a liquid medium to form a premix/predispersion; and
 - (c) subjecting the premix to mechanical means in the presence of a grinding medium to reduce the average effective particle size.
- 10 15. A process as claimed in claim 14, which is followed by sterilization, for instance autoclaving.
16. A process as claimed in claim 15, which is followed by re-suspending.
- 15 17. A process as claimed in claim 16, wherein the re-suspending consists of swirling the composition after sterilization for less than 40 seconds.
18. PEG4000, or the like, for use as a surface modifier in a pharmaceutical composition for administration by intramuscular or subcutaneous injection, wherein
- 20 said composition comprises an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, characterised in that the PEG4000 assists in re-suspending said composition after sterilization (e.g. autoclaving).
- 25 19. PEG4000, or the like, for use in re-suspending a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, wherein said composition has undergone sterilization (e.g. autoclaving).
- 30 20. PEG4000, or the like, for use as claimed in claim 18 or claim 19, wherein the pharmaceutical composition is as claimed in any one of claims 1 to 7.
21. The use of PEG4000, or the like, as a surface modifier in a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a
- 35 pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, wherein the PEG4000 assists in re-suspending said composition after sterilization (e.g. autoclaving).

22. The use of PEG4000, or the like, in re-suspending a pharmaceutical composition comprising an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in the form of a suspension of micro- or nano-particles, wherein said composition has undergone sterilization (e.g. autoclaving).

5

23. A use as claimed in claim 21 or claim 22, wherein the pharmaceutical composition is as claimed in any one of claims 1 to 7.

24. A process for preparing a pharmaceutical composition comprising

10

(a) obtaining an active pharmaceutical ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, in micronized form;

(b) adding the micronized active ingredient (e.g. bedaquiline), or a pharmaceutically acceptable salt thereof, to a liquid medium to form a premix/predispersion, characterised in that the liquid medium contains a surface modifier comprising PEG4000, or the like, as per any one of claims 1, 2, 3 or 4;

15

(c) subjecting the premix to mechanical means in the presence of a grinding medium to reduce the average effective particle size;

(d) sterilization (e.g. autoclaving); and

20

(e) re-suspending (if needed).

25. A process according to claim 24, wherein the re-suspending is performed by swirling for less than 40 seconds.

25

26. The use of PEG4000 in a process as claimed in claim 24 or claim 25.

FIGURE 1




FIGURE 2

Reference Examples B and C




FIGURE 3




FIGURE 4

Example 1





FIGURE 5

Example 1



6/10

FIGURE 6





FIGURE 7



8/10

FIGURE 8




FIGURE 9





FIGURE 10



Abstract**Long-acting formulations**

5 This invention concerns pharmaceutical compositions for administration via intramuscular or subcutaneous injection, comprising micro- or nanoparticles of the anti-TB compound bedaquiline, suspended in an aqueous pharmaceutically acceptable carrier, and comprising PEG4000 as a surface modifier, and the use of such pharmaceutical compositions in the treatment and prophylaxis of a pathogenic mycobacterial infection.

