



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2676010 C 2016/05/17

(11)(21) **2 676 010**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2008/01/18
(87) Date publication PCT/PCT Publication Date: 2008/07/24
(45) Date de délivrance/Issue Date: 2016/05/17
(85) Entrée phase nationale/National Entry: 2009/07/17
(86) N° demande PCT/PCT Application No.: US 2008/051466
(87) N° publication PCT/PCT Publication No.: 2008/089426
(30) Priorité/Priority: 2007/01/19 (IS8593/2007)

(51) Cl.Int./Int.Cl. *A61K 47/34* (2006.01),
A61K 31/55 (2006.01), *A61K 31/5517* (2006.01),
A61K 47/10 (2006.01), *A61K 9/00* (2006.01)

(72) Inventeurs/Inventors:
GIZURARSON, SVEINBJORN, IS;
ARP, JAMES, US;
GHADERI, RAOUF, US;
LUSTY, MICHAEL E., US;
...

(73) Propriétaires/Owners:
HANANJA EHF, IS;
UNIVERSITY OF ICELAND, IS;
IKANO THERAPEUTICS INC., US

(74) Agent: DEETH WILLIAMS WALL LLP

(54) Titre : PROCEDES ET COMPOSITIONS POUR L'ADMINISTRATION D'UN AGENT THERAPEUTIQUE
(54) Title: METHODS AND COMPOSITIONS FOR THE DELIVERY OF A THERAPEUTIC AGENT

(57) Abrégé/Abstract:

The present invention provides a liquid pharmaceutical composition comprising a therapeutic agent and an alkoxy-polyethylene glycol, for example, methoxy-polyethylene glycol, for administration of the therapeutic agent to the mammal. The compositions can be applied to a membrane, for example, a nasal membrane during intranasal administration. The invention also provides methods of administering such compositions to a mammal.

(72) Inventeurs(suite)/Inventors(continued): PLUCINSKI, GREGORY G., US; YAZDI, MEHDI, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
24 July 2008 (24.07.2008)

PCT

(10) International Publication Number
WO 2008/089426 A3

(51) International Patent Classification:

A61K 47/34 (2006.01) *A61K 9/00* (2006.01)
A61K 31/55 (2006.01) *A61K 47/10* (2006.01)
A61K 31/5517 (2006.01)

Mehdi [US/US]; 34 Mallet Hill Road, Columbia, SC 29223 (US).

(21) International Application Number:

PCT/US2008/051466

(74) Agents: GREENHALGH, Duncan, A. et al.; Goodwin Procter LLP, Exchange Place, Boston, MA 02109 (US).

(22) International Filing Date: 18 January 2008 (18.01.2008)

(25) Filing Language: English

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:

8593/2007 19 January 2007 (19.01.2007) IS

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (for all designated States except US): HANANJA EHF [IS/IS]; Aflagrandi 7, IS-107 Reykjavik (IS). UNIVERSITY OF ICELAND [IS/IS]; Sudurgata, IS-101 Reykjavik (IS). INTRANASAL THERAPEUTICS [US/US]; 210 Summit Avenue, Suite C2, Montvale, NJ 07645 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report: 12 February 2009

(54) Title: METHODS AND COMPOSITIONS FOR THE DELIVERY OF A THERAPEUTIC AGENT

(57) Abstract: The present invention provides a liquid pharmaceutical composition comprising a therapeutic agent and an alkoxy-polyethylene glycol, for example, methoxy-polyethylene glycol, for administration of the therapeutic agent to the mammal. The compositions can be applied to a membrane, for example, a nasal membrane during intranasal administration. The invention also provides methods of administering such compositions to a mammal.

**METHODS AND COMPOSITIONS FOR THE
DELIVERY OF A THERAPEUTIC AGENT**

[0001]

5

FIELD OF THE INVENTION

[0002] This invention relates generally to compositions for the delivery of a therapeutic agent and to related methods, and more particularly relates to 10 compositions containing one or more alkoxy-polyethylene glycols for the delivery of a therapeutic agent and to related methods.

BACKGROUND

[0003] The administration of a therapeutic agent by injection (e.g., intravenous, intramuscular or subcutaneous injection) typically is regarded as the most convenient way 15 of administration when the purpose is to achieve a rapid and strong systemic effect, for example, within 3-10 minutes, when the agent is not absorbed by the gastrointestinal tract, or when the agent is inactivated in the gastrointestinal tract or by first-pass hepatic metabolism. However, administration by injection presents a range of disadvantages. For example, sterile syringes must be used and injections cannot be administered by 20 untrained personnel. Furthermore, this mode of administration may cause pain and/or irritation, especially in the case of repeated injections at the same site.

[0004] Mucosal administration, such as, intranasal, buccal, sublingual, rectal and pulmonal administration, is receiving particular interest as it avoids many of the 25 disadvantages of injecting a therapeutic agent while, at the same time, still providing a strong and rapid systemic effect. In order to be an attractive alternative to injection, mucosal administration, for example, intranasal administration, should neither cause significant pain, discomfort or irritation nor cause any irreversible damage to the mucosal surface. However, in the case of acute health threatening indications, a relatively high local irritation to the mucosa may be acceptable.

- 2 -

[0005] In mucosal administration, such as during nasal, buccal or rectal administration, the therapeutic agent should be applied to the mucosa in a vehicle that permits it to penetrate, or be absorbed through, the mucosa. In order to penetrate the mucus, the vehicle should be biocompatible with mucus and hence have a certain degree of hydrophilicity. However, the 5 vehicle should preferably also possess lipophilic properties to dissolve a clinically relevant amount of the therapeutic agent of interest.

[0006] The extensive network of blood capillaries under the mucosal surface, especially in the nasal mucosa, is well suited to provide a rapid and effective systemic absorption of drugs, vaccines and biologicals. Moreover, the nasal epithelial membrane in effect contains a single 10 layer of epithelial cells (*pseudostratified epithelium*) and, therefore, is more suited for drug administration than other mucosal surfaces having squamous epithelial layers, such as, the mouth and vagina.

[0007] It has been hypothesized that the usefulness of nasal administration can be limited if the therapeutic agent has limited solubility in water (Proctor, D. F. (1985) *Nasal Physiology in 15 Intranasal Drug Administrations*, in Chien, Y. W. (Ed.) *TRANSNASAL SYSTEMIC MEDICATIONS, FUNDAMENTALS, DEVELOPMENTAL CONCEPTS AND BIOMEDICAL ASSESSMENTS*, ELSEVIER Science Publishers, Amsterdam, pp. 101-105). As a result, this hypothesis, if correct, may limit the delivery of certain therapeutic agents that are sparingly soluble in water.

[0008] To facilitate delivery to the nasal cavity, an effective amount of the therapeutic 20 agent should be dissolved in a small volume, for example, less than about 1000 µL, preferably less than 300 µL, and more preferably less than 150 µL. Larger volumes drain out anteriorly through the nostrils or posteriorly toward the pharynx where excess liquid is swallowed. As a result, if large volumes are administered, a portion of the therapeutic agent can be lost from the absorption site, and it can be difficult if not impossible to reproducibly administer the correct 25 dose of the therapeutic agent.

[0009] A variety of delivery systems have been developed for the nasal administration of therapeutic agents. Lau and Slattery studied the absorption characteristics of diazepam and lorazepam following their intranasal administration for the treatment of epilepticus (Lau, S. W. 30 J. & Slattery, J. T. (1989), *Absorption of Diazepam and Lorazepam Following Intranasal Administration*, *INT. J. PHARM.*, 54, 171-174). In order to solubilize the therapeutic agent, a non-ionic surfactant - polyoxyethylated castor oil - was selected as the least irritating solvent of

- 3 -

several solvents studied, including polyethylene glycol 400 (PEG 400). Diazepam absorption was 84 % and 72 %, respectively, in two adults measured over a period of 60 hours. However, the peak concentration was not observed until 1.4 hours after the nasal administration and was only about 27 % with reference to intravenous administration, suggesting that most of the
5 absorption had taken place after the test substance passed down to pharynx and swallowed. Similar results were obtained for lorazepam but with an even longer time to peak (2.3 hours). The authors concluded that the intranasal route of administration had limited potential for the acute treatment of epileptic seizures.

[0010] Wilton *et al.* attempted to administer midazolam to 45 children to achieve pre-anesthetic sedation (Wilton *et al.* (1988) *Preanaesthetic Sedation of Preschool Children Using Intranasal Midazolam*, ANESTHESIOLOGY, 69, 972-975). However, the volumes used were impractical and exceeded the maximal volume required for efficient administration. This resulted in coughing and sneezing with expulsion of at least part of the dose.

[0011] Morimoto *et al.* studied a gel preparation for nasal application in rats of nifedipine containing the gelling agent carbopol (polyacrylic acid) in PEG 400, for achieving prolonged action and high bioavailability of the therapeutic agent (Morimoto *et al.* (1987) *Nasal Absorption of Nifedipine from Gel Preparations in Rats*, CHEMICAL AND PHARMACEUTICAL BULLETINS, 35, No. 7, 3041-3044). A mixture of equal amounts of carbopol and PEG 400 was preferred. It was shown that nasal application provided higher bioavailability of nifedipine
15 than after peroral administration, but the peak plasma concentration was not observed until 30 minutes after administration.

[0012] Danish Patent Application No. 2586/87 discloses a pharmaceutical composition comprising an anti-inflammatory steroid, water, 2 to 10 % (v/v) propylene glycol, 10 to 25 % (v/v) PEG 400, and 1 to 4 % (v/v) Tween 20.

25 [0013] U.S. Patent No. 4,153,689 discloses a stable aqueous solution of insulin intended for intranasal administration. The solutions had a pH not more than 4.7, and contained from 0.1 to 20 % by weight of a stabilizing agent including (a) one or more non-ionic surface active agents whose hydrophile-lipophile balance value was in the range of 9 to 22, and/or (b) polyethylene glycol whose molecular weight was in the range of from 200 to 7500. Exemplary non-ionic
30 surface active agents included polyoxyethylene fatty acid ester, a polyoxyethylene higher

alcohol ether, a polyoxyethylene alkylphenyl ether, or a polyoxyethylene alkylphenyl ether, or a polyoxyethylene hydrogenated castor oil.

[0014] International Patent Publication No. *WO 91/16929* discloses the nasal 5 administration of therapeutic agents, for example, benzodiazepines, in compositions containing n-glycofurol, a derivative of polyethyleneglycol, for mucosal administration. The application discloses the nasal administration of therapeutic agents, for example, benzodiazepines, in formulations containing at least 30% n-ethyleneglycols ranging from 1-8 ethylene glycol, for example, polyethylene glycol 200 (PEG 200).

10 [0015] U.S. Patent No. 5,693,608 discloses a method of administering a therapeutic agent via the nasal mucosa of a mammal, where the agent is dissolved or suspended in an n-ethyleneglycol containing vehicle where the n-ethyleneglycol is represented by the formula, $H(OCH_2CH_2)_pOH$, wherein p is a number from 1 to 8.

[0016] Notwithstanding, there is still a need for compositions deliverable through 15 mucosal membranes that produce therapeutic plasma concentrations of the therapeutic agent as fast as or nearly as fast as by intravenous administration but without causing irritation and/or unacceptable damage to the mucosal membrane.

SUMMARY OF THE INVENTION


[0017] The invention is based, in part, upon the discovery that the inclusion of one or 20 more alkoxy-polyethylene glycols in a formulation provides certain advantages when the resulting composition is to be applied, for example, to a mucosal surface. For example, it has been discovered that when alkoxy-polyethylene glycol is used in such formulations, the therapeutic agent can be still be solubilized (which is especially useful for poorly soluble therapeutic agents) but the resulting formulations 25 are less viscous and cause less irritation to mucosal membranes because the amount of other potentially viscous and irritable excipients, for example, polyethylene glycol or propylene glucol, can be reduced or eliminated altogether. As a result, the lower viscosity formulations, when converted into droplets, for example, by a nasal sprayer during intranasal delivery, can produce a spray pattern optimized for delivering the therapeutic 30 agent to the mucosal membrane. In addition, formulations containing alkoxy-polyethylene glycols create less irritation (burning sensation) when applied to a mucosal surface, for example, a nasal membrane following nasal administration. In addition, when

- 5 -

administered intranasally, the compositions of the invention minimize undesirable after taste (for example, a petroleum-like after taste) that can be associated with certain other excipients.

[0018] In one aspect, the invention provides a liquid pharmaceutical composition comprising a therapeutic agent and an alkoxy-polyethylene glycol represented by Formula I:

5




wherein,

R is methyl, ethyl, n-propyl, isopropyl, or cyclopropyl; and

n, which is the average number of oxyethylene repeating units, is a number in the range of from about 1 to about 25.

10 **[0019]** In another aspect, the invention provides a liquid formulation for solubilizing a poorly soluble therapeutic agent, for example, a poorly soluble organic therapeutic agent. The composition comprises a poorly soluble therapeutic agent and an alkoxy-polyethylene glycol represented by Formula I:



15

wherein,

R is (C₁-C₆)alkyl; and

n, which is the average number of oxyethylene repeating units, is a number in the range of from about 1 to about 25.

20 **[0020]** In another aspect, the invention provides methods of delivering a therapeutic agent of interest to a mammal, for example, a human, using an alkoxy-polyethylene glycol containing composition described herein. The composition is particularly useful when the composition is applied to a mucosal membrane, for example, a nasal membrane during intranasal drug delivery.

25 **[0021]** These and other aspects and advantages of the invention will become apparent upon consideration of the following detailed description and claims.

DETAILED DESCRIPTION OF THE INVENTION

[0022] The invention is based, in part, upon the discovery that the inclusion of one or more alkoxy-polyethylene glycols into formulations provides certain advantages over other excipients when the formulations are applied, for example, to a mucosal surface. For example, 5 it has been discovered that when an alkoxy-polyethylene glycol is used in such a formulation, the therapeutic agent (for example, a poorly soluble therapeutic agent) can be solubilized more easily and in larger amounts than when other excipients, for example, polyethylene glycol (more particularly PEG 400), are used. However, the resulting formulations are less viscous and cause less irritation to mucosal membranes as the amount of other viscous and irritable 10 excipients can be reduced or eliminated altogether. As a result, the lower viscosity formulations, when converted into droplets, for example, by a nasal sprayer during intranasal delivery, produce a spray pattern optimized for delivering the therapeutic agent to the mucosal membrane. In addition, formulations containing one or more alkoxy-polyethylene glycols 15 cause less irritation (for example, a burning sensation) when applied to a mucosal surface, for example, a nasal membrane during intranasal administration. In addition, when administered intranasally, the compositions of the invention have less undesirable after taste (for example, a petroleum-like after taste) than when other excipients, for example, propylene glycol, are used.

[0023] Under certain circumstances, the alkoxy-group also increases the bioadhesion of the 20 composition to the site of administration on the mucosal surface thereby prolonging the duration of the composition at the site of administration. This can increase the amount of therapeutic agent that is ultimately absorbed.

I - Formulations

In one aspect, the invention provides a liquid pharmaceutical composition comprising a therapeutic agent and an alkoxy-polyethylene glycol represented by Formula I:



wherein,


R is methyl, ethyl, n-propyl, isopropyl, or cyclopropyl; and

n is the average number of oxyethylene repeating units and is a number in the range of from about 1 to about 25.

- 7 -

[0024] In another aspect, the invention provides a liquid formulation for solubilizing a poorly soluble therapeutic agent. The liquid pharmaceutical composition comprises a poorly soluble therapeutic agent, for example, a poorly soluble organic therapeutic agent, and an alkoxy-polyethylene glycol represented by Formula I:

5




wherein,

R is (C₁-C₆)alkyl; and

n is the average number of oxyethylene repeating units and is a number in the range of from about 1 to about 25. The formulations typically are in liquid form at 20°C, 25°C, 30°C, 10 35°C, or 40°C. Certain formulations preferably are liquid formulations at 37°C.

[0025] The term "poorly soluble therapeutic agent" refers to a compound having biological activity and a solubility in water of less than about 1 mg/mL at pH 7 and 20°C. In certain embodiments, the poorly soluble therapeutic agent is an organic compound that has a molecular weight of less than 1500 g/mol, and preferably less than 500 g/mol. In certain embodiments, 15 the poorly soluble therapeutic agent is a compound, for example, an organic compound, having an aqueous solubility of less than about 0.5 mg/mL, less than about 0.3 mg/mL, or less than about 0.1 mg/mL, at pH 7 and 20°C.

[0026] In addition, the term "alkyl" is art-recognized, and includes saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. 20 The term "(C₁-C₆)alkyl" refers to an alkyl group having between 1 and 6 carbon atoms. Representative alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, pentyl, hexyl, cyclopropyl, cyclobutyl, cyclopropylmethylen, cyclopentyl, cyclobutylmethylen, cyclobutylethylene, cyclohexyl, cyclopropylpropylene, 25 cyclobutylethylene, and cyclopentylmethylen. The term cyclopropylmethylen, for example, is art-recognized and refers to a radical having the following formula:



[0027] In certain embodiments, the alkoxy-polyethylene glycol can comprise from about 0.1% (v/v) to about 80% (v/v), or from about 0.5 % (v/v) to about 70 % (v/v), of the composition. In certain other embodiments, the alkoxy-polyethylene glycol can comprise from about 5% (v/v) to about 80% (v/v), or from about 30 % (v/v) to about 75 % (v/v) or from about 5 40% (v/v) to about 70% (v/v), of the composition. For certain hydrophilic drugs, the alkoxy-polyethylene glycol can comprise from about 0.1 % (v/v) to about 80% (v/v), or from about 0.5 % (v/v) to about 70 % (v/v), or from about 1 % (v/v) to about 60% of the composition. For certain lipophilic drugs, the alkoxy-polyethylene glycol can comprise from about 1 % (v/v) to about 80% (v/v), or from about 2 % (v/v) to about 65 % (v/v), or from about 5 % (v/v) to about 10 50% of the composition. Furthermore, the therapeutic agent can comprise from about 0.001 % (w/v) to about 20 % (w/v) of the composition, or from about 0.1 % (w/v) to about 10 % (w/v) of the composition.

[0028] The pharmaceutical composition can have a pH in the range of from about 4.5 to about 8.5, or from about 4.5 to about 7.5, or from about 4.5 to about 6.5, or from about 5.5 to 15 about 8.5, or from about 6.5 to about 8.5, or from about 5.5 to about 7.5.

[0029] As discussed, one of the advantages of using an alkoxy-polyethylene glycol is that it can be used in place of or can be used to reduce the amount of other excipients, for example, certain polyethylene glycols and propylrnr glycol, so as to reduce the viscosity of the resulting formulation. By reducing the viscosity of the resulting formation it is possible to create sprays 20 that have more uniform spray characteristics (for example, more uniform droplet sizes and/or plume geometries) for the intranasal administration of therapeutic agent. The resulting pharmaceutical composition at a temperature of 20°C has a viscosity in the range of about 1.5 cP to about 60 cP, or from about 2 cP to about 50 cP, or from about 3 cP to about 40 cP, or from about 4 cP to about 30 cP, or from about 5 cP to about 25 cP.

25 [0030] Exemplary alkoxy-polyethylene glycols, therapeutic agents, and other excipients useful in creating compositions of the invention are described in the following sections.

- 9 -

A. Alkoxy-polyethylene Glycol

[0031] Useful alkoxy-polyethylene glycol excipients useful in the practice of the invention are represented by Formula (I):



5 wherein, n, as the average number of oxyethylene repeating units, is a number in the range of from about 1 to about 25. Accordingly, n can be a number about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25. In certain embodiments, n is an number in the range of from about 2 to about 15, or from about 2 to about 14, or from about 2 to about 13, or from about 2 to about 12, or from about 2 to about 11, or from about 2 to about 10, or from about 3 to about 15, or from about 3 to about 14, or from about 3 to about 13, or from about 3 to about 12, or from about 3 to about 11, or from about 3 to about 10.

10 In certain embodiments, R is (C₁-C₆)alkyl. For example, as discussed above, R can be methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, pentyl, hexyl, cyclopropyl, cyclobutyl, cyclopropylmethylene, cyclopentyl, cyclobutylmethylene, 15 cyclobutylethylene, cyclohexyl, cyclopropylpropylene, cyclobutylethylene, or cyclopentylmethylene. In certain embodiments, R is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, and cyclopropyl.

[0032] In a preferred embodiment, the alkoxy-polyethylene glycol is methoxy-polyethylene glycol where R is methyl and n is an number from about 1 to about 25, or from about 2 to about 20 12, or from about 3 to about 10.

[0033] Useful methoxy-polyethylene glycols include, for example, methoxy-diethyleneglyol (m2EG), methoxy-triethylene glycol (m3EG), methoxy-tetraethylene glycol (m4EG), methoxy-pentaethylene glycol (m5EG), methoxy-hexaethylene glycol (m6EG), methoxy-heptaethylene glycol (m7EG), methoxy-octaethylene glycol (m8EG), methoxy-nonaethylene glycol (m9EG), methoxy-decaethylene glycol (m10EG), methoxy-undecaethylene glycol (m11EG), methoxy-dodecaethylene glycol (m12EG), methoxy-tridecaethylene glycol (m13EG) and methoxy-tetradecaethylene glycol (m14EG). The ethylene glycols may be used in the form of the single compounds or as a mixture of two or more methoxy-n-ethylene glycols.

- 10 -

[0034] In certain embodiments, the alkoxy-polyethylene glycol is methoxy-polyethylene glycol 350 (mPEG 350) or is methoxy-polyethylene glycol 550 (mPEG 550) or is methoxy-polyethylene glycol 750 (mPEG 750). The term “mPEG 350” is understood to mean methoxy polyethylene glycol having an average molecular weight of about 350, and in certain 5 embodiments “n,” as denoted in Formula I, is 7.2. The term “mPEG550” is understood to mean methoxy polyethylene glycol having an average molecular weight of about 550, and in certain embodiments “n,” as denoted in Formula I, is 11.8. The term “mPEG750” is understood to mean methoxy polyethylene glycol having an average molecular weight of about 750, and in certain embodiments “n,” as denoted in Formula I, is 16.3.

[0035] Certain, preferred alkoxy-polyethylene glycols include CarbowaxTM mPEG 350, CarbowaxTM mPEG 550 or CarbowaxTM mPEG 750, which are available commercially from Dow Chemical Company. Both mPEG350 and mPEG550 are colorless liquids that are miscible with water, alcohols, such as methanol, ethanol, n-propanol, glycerol and various oils in all proportions, and have a boiling point about 155°C. It is understood that alkoxy- 15 polyethylene glycols are known by other names, where, for example, methoxy-polyethylene glycol is also known as mono-methyl polyethylene glycol and poly(ethylene glycol) methyl ether.

[0036] By using one or more of the alkoxy-polyethylene glycols described herein, the resulting pharmaceutical compositions can be optimized, for example, with respect to 20 bioadhesion, viscosity and sprayability. For example, mPEG 350, at an equivalent concentration as PEG 200, can still solubilize a therapeutic agent but the resulting composition has a lower viscosity. As a result, this substitution has a surprisingly positive effect on the sprayability compared with lower molecular weight PEG 200, which is important where the formulation is to be sprayed.

25 **B. Therapeutic Agent**

[0037] The pharmaceutical composition of the invention may comprise one or more therapeutic agents (also referred to as biologically active substances) selected from the group consisting of hydrophobic therapeutic agents, hydrophilic therapeutic agents, and combinations thereof.

[0038] The alkoxy-polyethylene glycol excipients are surprisingly capable of solubilizing and delivering a wide variety of hydrophilic and hydrophobic therapeutic agents. The hydrophobic drugs have little or no water solubility. It is understood that the excipients described herein can be used to solubilize therapeutic agents that have a solubility in water of less than about 1.0 mg/mL, less than about 0.5 mg/mL, less than about 0.3 mg/mL, or less than about 0.1 mg/mL, or less than about 0.01 mg/mL, at pH 7 and 20°C. Such therapeutic agents can be any agents having therapeutic or other value when administered to a mammal, for example, a human, and can include organic molecules (for example, small molecule drugs having a molecular weight of less than 1,500 g/mol., or less than 500 g/mol.), proteins, peptides, immungens (e.g. vaccines, cytokines, etc.), nutrients, and cosmetics (cosmeceuticals).

[0039] In certain embodiments, the therapeutic agent is an analgesic agent, an anti-inflammatory agent, an anti-arrhythmic agent, an anti-asthma agent, an anti-bacterial agent, an anti-viral agent, an anti-coagulant, an anti-depressant, an anti-diabetic, an anti-epileptic, an anti-fungal agent, an anti-hypertensive agent, an anti-malarial, an anti-migraine agent, an anti-muscarinic agent, an anti-neoplastic agent, an immunosuppressant, an anti-protozoal agent, an anti-thyroid agent, an anxiolytic agent, a sedative, a hypnotic agent, a neuroleptic agent, a beta-Blocker, a cardiac inotropic agent, a corticosteroid, a diuretic agent, an anti-Parkinsonian agent, a gastrointestinal agent, an anti-histamine, a histamine-receptor antagonist, a lipid regulating agent, a muscle relaxant, nitrate and other anti-anginal agent, a nutritional agent, an opioid analgesic, sex hormone, stimulant, cytokine, peptidomimetic, peptide, protein, toxoid, sera, antibody, vaccine, nucleoside, nucleotide, nucleic acid and peptidyl-nucleic acid.

[0040] Specific non-limiting examples of hydrophobic therapeutic agents that can be used in the pharmaceutical compositions of the present invention include the following representative compounds, as well as their pharmaceutically acceptable salts, isomers, esters, ethers and other derivatives including, for example: (1) analgesics and anti-inflammatory agents, such as, aloxiprin, auranofin, azapropazone, benorylate, capsaicin, celecoxib, diclofenac, diflunisal, etodolac, fenbufen, fenoprofen calcium, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, leflunomide, meclofenamic acid, mefenamic acid, nabumetone, naproxen, oxaprozin, oxyphenbutazone, phenylbutazone, piroxicam, refcoxib, sulindac, tetrahydrocannabinol, tramadol and tromethamine; (2) anti-arrhythmic agents, such as, amiodarone HCl, disopyramide, flecainide acetate and quinidine sulfate; (3) anti-asthma

agents, such as, zileuton, zafirlukast, montelukast, and albuterol; (4) anti-bacterial agents, such as, baclofen, benzathine penicillin, cinoxacin, clarithromycin, clofazimine, cloxacillin, demeclocycline, dirithromycin, doxycycline, erythromycin, ethionamide, furazolidone, grepafloxacin, imipenem, levofloxacin, lorefloxacin, moxifloxacin HCl, nalidixic acid, 5 nitrofurantoin, norfloxacin, ofloxacin, rifampicin, rifabutine, rifapentine, sparfloxacin, spiramycin, sulphabenzamide, sulphadoxine, sulphamerazine, sulphacetamide, sulphadiazine, sulphafurazole, sulphamethoxazole, sulphapyridine, tetracycline, trimethoprim and trovafloxacin; (5) anti-viral agents, such as, abacavir, amprenavir, delavirdine, efavirenz, indinavir, lamivudine, nelfinavir, nevirapine, ritonavir, saquinavir, and stavudine; (6) anti- 10 coagulants, such as, cilostazol, clopidogrel, dicumarol, dipyridamole, nicoumalone, oprelvekin, phenindione, ticlopidine, and tirofiban; (7) anti-depressants, such as amoxapine, bupropion, citalopram, clomipramine, maprotiline HCl, mianserin HCl, nortriptyline HCl, paroxetine HCl, sertraline HCl, trazodone HCl, trimipramine maleate, and venlafaxine HCl; (8) anti-diabetics, such as, acetohexamide, chlorpropamide, glibenclamide, gliclazide, glipizide, glimepiride, 15 miglitol, pioglitazone, repaglinide, rosiglitazone, tolazamide, tolbutamide and troglitazone; (9) anti-epileptics, such as, beclamide, carbamazepine, clonazepam, ethotoin, felbamate, fosphenytoin sodium, lamotrigine, methoin, methsuximide, methylphenobarbitone, oxcarbazepine, paramethadione, phenacetin, phenobarbitone, phenytoin, phensuximide, primidone, sulthiame, tiagabine HCl, topiramate, valproic acid, and vigabatrin; (10) anti-fungal 20 agents, such as, amphotericin, butenafine HCl, butoconazole nitrate, clotrimazole, econazole nitrate, fluconazole, flucytosine, griseofulvin, itraconazole, ketoconazole, miconazole, natamycin, nystatin, sulconazole nitrate, oxiconazole, terbinafine HCl, terconazole, tioconazole and undecenoic acid; (11) anti-hypertensive agents, such as, amlodipine, benidipine, 25 benazepril, candesartan, captopril, darodipine, diltiazem HCl, diazoxide, doxazosin HCl, elanapril, eposartan, losartan mesylate, felodipine, fenoldopam, fosenopril, guanabenz acetate, irbesartan, isradipine, lisinopril, minoxidil, nicardipine HCl, nifedipine, nimodipine, nisoldipine, phenoxybenzamine HCl, prazosin HCl, quinapril, reserpine, terazosin HCl, telmisartan, and valsartan; (12) anti-malarials, such as, amodiaquine, chloroquine, chlorproguanil HCl, halofantrine HCl, mefloquine HCl, proguanil HCl, pyrimethamine and 30 quinine sulfate; (13) anti-migraine agents, such as, dihydroergotamine mesylate, ergotamine tartrate, frovatriptan, methysergide maleate, naratriptan HCl, pizotyline malate, rizatriptan benzoate, sumatriptan succinate, and zolmitriptan; (14) anti-muscarinic agents, such as,

atropine, benzhexol HCl, biperiden, ethopropazine HCl, hyoscyamine, oxyphencyclimine HCl and tropicamide; (15) anti-neoplastic agents and immunosuppressants, such as, aminoglutethimide, amsacrine, azathioprine, bicalutamide, bisantrene, busulfan, camptothecin, chlorambucil, cyclosporin, dacarbazine, ellipticine, estramustine, etoposide, irinotecan, 5 lomustine, melphalan, mercaptopurine, mitomycin, mitotane, mitoxantrone, mofetil mycophenolate, nilutamide, paclitaxel, procarbazine HCl, sirolimus, tacrolimus, tamoxifen citrate, teniposide, testolactone, topotecan HCl, and toremifene citrate; (16) anti-protozoal agents, such as, atovaquone, benznidazole, clioquinol, decoquinate, diiodohydroxyquinoline, diloxanide furoate, dinitolmide, furazolidone, metronidazole, nimorazole, nitrofurazone, 10 ornidazole and tinidazole; (17) anti-thyroid agents, such as, carbimazole, paracalcitol, and propylthiouracil; anti-tussives, such as, benzonatate; (18) anxiolytics, sedatives, hypnotics and neuroleptics, such as, alprazolam, amylobarbitone, barbitone, bentazepam, bromazepam, bromperidol, brotizolam, butobarbitone, carbromal, chlordiazepoxide, chlormethiazole, chlorpromazine, chlorprothixene, clonazepam, clobazam, clotiazepam, clozapine, diazepam, 15 droperidol, ethinamate, flunanisone, flunitrazepam, triflupromazine, fluphenthixol decanoate, fluphenazine decanoate, flurazepam, gabapentin, haloperidol, lorazepam, lormetazepam, medazepam, meprobamate, mesoridazine, methaqualone, methylphenidate, midazolam, molindone, nitrazepam, olanzapine, oxazepam, pentobarbitone, perphenazine pimozide, prochlorperazine, pseudoephedrine, quetiapine, rispiridone, sertindole, sulpiride, temazepam, 20 thioridazine, triazolam, zolpidem, and zopiclone; (19) beta-Blockers, such as, acebutolol, alprenolol, labetalol, metoprolol, nadolol, oxprenolol, pindolol and propranolol; (20) cardiac inotropic agents, such as, amrinone, digitoxin, digoxin, enoximone, lanatoside C and medigoxin; (21) a corticosteroid, such as, beclomethasone, betamethasone, budesonide, cortisone acetate, desoxymethasone, dexamethasone, fludrocortisone acetate, flunisolide, 25 fluocortolone, fluticasone propionate, hydrocortisone, methylprednisolone, prednisolone, prednisone and triamcinolone; (22) diuretics, such as, acetazolamide, amiloride, bendroflumethiazide, bumetanide, chlorothiazide, chlorthalidone, ethacrynic acid, frusemide, metolazone, spironolactone and triamterene; (23) anti-Parkinsonian agents, such as, bromocriptine mesylate, lysuride maleate, pramipexole, ropinirole HCl, and tolcapone; (24) 30 gastrointestinal agents, such as bisacodyl, cimetidine, cisapride, diphenoxylate HCl, domperidone, famotidine, lansoprazole, loperamide, mesalazine, nizatidine, omeprazole, ondansetron HCl, rabeprazole sodium, ranitidine HCl and sulphasalazine; (25) anti-histamines

and histamine-receptor antagonists, such as, acrivastine, astemizole, chlorpheniramine, cinnarizine, cetrizine, clemastine fumarate, cyclizine, cyproheptadine HCl, dexchlorpheniramine, dimenhydrinate, fexofenadine, flunarizine HCl, loratadine, meclizine HCl, oxatomide, and terfenadine; (26) lipid regulating agents, such as, atorvastatin, bezafibrate, 5 cerivastatin, ciprofibrate, clofibrate, fenofibrate, fluvastatin, gemfibrozil, pravastatin, probucol, and simvastatin; (27) muscle relaxants, such as, dantrolene sodium and tizanidine HCl; (28) nitrates and other anti-anginal agents, such as, amyl nitrate, glyceryl trinitrate, isosorbide dinitrate, isosorbide mononitrate and pentaerythritol tetranitrate; (29) nutritional agents, such as, calcitriol, carotenes, dihydrotachysterol, essential fatty acids, non-essential fatty acids, 10 phytonadiol, vitamin A, vitamin B₂, vitamin D, vitamin E and vitamin K; (30) opioid analgesics, such as, codeine, dextropropoxyphene, diamorphine, dihydrocodeine, fentanyl, meptazinol, methadone, morphine, hydromorphone, nalbuphine and pentazocine; (31) sex hormones, such as, clomiphene citrate, cortisone acetate, danazol, dehydroepiandrosterone, ethynodiol diacetate, finasteride, fludrocortisone, fluoxymesterone, medroxyprogesterone acetate, 15 megestrol acetate, mestranol, methyltestosterone, norethisterone, norgestrel, oestradiol, conjugated estrogens, progesterone, rimexolone, stanozolol, stilbestrol, testosterone and tibolone; and (32) stimulants, such as, amphetamine, dexamphetamine, dextroamphetamine, dextroamphetamine, fenfluramine and mazindol; and others, such as, becaplermin, donepezil HCl, L-thyroxine, methoxsalen, verteporfin, physostigmine, pyridostigmine, raloxifene HCl, sibutramine HCl, 20 sildenafil citrate, tacrine, tamsulosin HCl, and tolterodine.

[0041] Certain exemplary hydrophobic therapeutic agents include sildenafil citrate, amlodipine, tramadol, celecoxib, rofecoxib, oxaprozin, nabumetone, ibuprofen, terbenafine, itraconazole, zileuton, zafirlukast, cisapride, fenofibrate, tizanidine, nizatidine, fexofenadine, loratadine, famotidine, paricalcitol, atovaquone, nabumetone, alprazolam, bromazepam, 25 chlorpromazine, clonazepam, diazepam, flunitrazepam, flurazepam, haloperidol, lorazepam, lormetazepam, midazolam, nitrazepam, oxazepam, pseudoephedrine, temazepam, triazolam, zolpidem, zopiclone, tetrahydrocannabinol, testosterone, megestrol acetate, repaglinide, progesterone, rimexolone, cyclosporin, tacrolimus, sirolimus, teniposide, paclitaxel, 30 pseudoephedrine, troglitazone, rosiglitazone, finasteride, vitamin A, vitamin D, vitamin E, and pharmaceutically acceptable salts, isomers and derivatives thereof. It should be appreciated that the listing of hydrophobic therapeutic agents and their therapeutic classes is merely

illustrative. It is understood, that mixtures of hydrophobic therapeutic agents may also be used where desired.

[0042] An advantage of using alkoxy-polyethylene glycol as an excipient in the pharmaceutical composition is that, for example, highly lipophilic substances, such as, 5 lorazepam, midazolam, clonazepam, alprazolam and other compounds belonging to the benzodiazepines, as well as water soluble substances, for example, peptides and proteins, such as, the pancreatic hormones can be solubilized in a clinically relevant volume (for example, 25 - 300 μ L) for delivery to a human subject. By way of comparison, clinically relevant doses of midazolam, lorazepam, alprazolam, diazepam and clonazepam would have to be dissolved in at 10 least 5 mL of water.

[0043] Although the alkoxy-polyethylene glycol excipients are particularly useful for the delivery of hydrophobic agents, alkoxy-polyethylene glycol can also be used to deliver a variety of hydrophilic therapeutic agents. Alkoxy-polyethylene glycals, under certain 15 circumstances, may prolong the duration of the therapeutic agent at the absorption site thereby increasing the amount of agent ultimately delivered. Exemplary hydrophilic therapeutic agents include hydrophilic drugs (i.e., conventional non-peptidic drugs), hydrophilic macromolecules, such as, cytokines, peptides, proteins, peptidomimetics, toxoids, sera, antibodies, vaccines, nucleosides, nucleotides, nucleic acids, and genetic material. The hydrophilic therapeutic agent can be administered alone or in combination with other agents, for example, a hydrophobic 20 therapeutic agent discussed hereinabove or a second, different hydrophilic therapeutic agent.

[0044] Without limitation, exemplary hydrophilic therapeutic agents that can be delivered using the compositions and methods of the present invention, include the following compounds as well as their pharmaceutically acceptable salts, isomers, esters, ethers and other derivatives, for example: acarbose; acyclovir; acetyl cysteine; acetylcholine chloride; alatrofloxacin; 25 alendronate; alglucerase; amantadine hydrochloride; ambenonium; amifostine; aminocaproic acid; antihemophilic factor (human); antihemophilic factor (porcine); antihemophilic factor (recombinant); aprotinin; asparaginase; atenolol; atracurium besylate; azithromycin; aztreonam; BCG vaccine; bacitracin; becadermin; belladonna; bepridil hydrochloride; bleomycin sulfate; calcitonin human; calcitonin salmon; carboplatin; capecitabine; capreomycin sulfate; 30 cefamandole nafate; cefazolin sodium; cefepime hydrochloride; cefixime; cefonicid sodium; cefoperazone; cefotetan disodium; cefotaxime; cefoxitin sodium; ceftizoxime; ceftriaxone;

cefuroxime axetil; cephalexin; cephapirin sodium; cholera vaccine; chorionic gonadotropin; cidofovir; cisplatin; cladribine; clidinium bromide; clindamycin and clindamycin derivatives; ciprofloxacin; clondronate; colistimethate sodium; colistin sulfate; corticotropin; cosyntropin; cromalyn sodium; cytarabine; daltaperin sodium; danaproid; deforoxamine; denileukin diftitox; 5 desmopressin; diatrizoate meglumine and diatrizoate sodium; dicyclomine; didanosine; dopamine hydrochloride; dornase alpha; doxacurium chloride; doxorubicin; editronate disodium; elanaprilat; enkephalin; enoxacin; enoxaprin sodium; ephedrine; epinephrine; epoetin alpha; esmol hydrochloride; Factor IX; famiciclovir; fludarabine; fluoxetine; foscarnet sodium; ganciclovir; granulocyte colony stimulating factor; granulocyte-macrophage 10 stimulating factor; growth hormones (human or bovine); gentamycin; glucagon; glycopyrrolate; gonadotropin releasing hormone and synthetic analogs thereof; GnRH; gonadorelin; grepafloxacin; hemophilus B conjugate vaccine; Hepatitis A virus vaccine inactivated; Hepatitis B virus vaccine inactivated; heparin sodium; indinavir sulfate; influenza virus vaccine; interleukin-2; interleukin-3; insulin-human; insulin-porcine; insulin NPH; insulin 15 aspart; insulin glargine; insulin detemir; interferon- α ; interferon- β ; ipratropium bromide; isofosfamide; japanese encephalitis virus vaccine; leucovorin calcium; leuprolide acetate; levofloxacin; lincomycin and lincomycin derivatives; lobucavir; lomefloxacin; loracarbef; mannitol; measles virus vaccine; meningococcal vaccine; menotropins; mephenzolate bromide; mesalmine; methanamine; methotrexate; methscopolamine; metformin hydrochloride; 20 metroprolol; mezocillin sodium; mivacurium chloride; mumps viral vaccine; nedocromil sodium; neostigmine bromide; neostigmine methyl sulfate; neutontin; norfloxacin; octreotide acetate; olpadronate; oxytocin; pamidronate disodium; pancuronium bromide; paroxetine; pefloxacin; pentamidine isethionate; pentostatin; pentoxifylline; periciclovir; pentagastrin; phentolamine mesylate; phenylalanine; physostigmine salicylate; plague vaccine; piperacillin 25 sodium; platelet derived growth factor-human; pneumococcal vaccine polyvalent; poliovirus vaccine (live or inactivated); polymixin B sulfate; pralidoxine chloride; pramlintide; pregabalin; propofenone; propenthaline bromide; pyridostigmine bromide; rabies vaccine; residronate; ribavarin; rimantadine hydrochloride; rotavirus vaccine; salmetrol xinafoate; sicalide; small pox vaccine; solatol; somatostatin; sparfloxacin; spectinomycin; stavudine; 30 streptokinase; streptozocin; suxamethonium chloride; tacrine hydrochloride; terbutaline sulfate; thiopeta; ticarcillin; tiludronate; timolol; tissue type plasminogen activator; TNFR:Fc; TNK-tPA; trandolapril; trimetrexate gluconate; trospectinomycin; trovafloxacin; tubocurarine

chloride; tumor necrosis factor; typhoid vaccine live; urea; urokinase; vancomycin; valaciclovir; varicella virus vaccine live; vasopressin and vasopessin derivatives; vecuronium bromide; vinblastine; vincristine; vinorelbine; warfarin - sodium; yellow fever vaccine; zalcitabine; zanamivir; zolandronate; and zidovudine.

5 [0045] Other therapeutic agents that can be administered in this formulation may comprise adrenal hormones, corticosteroids and derivatives, such as, ACTH and analogs thereof, teracosactrin, alsactide, cortisone, hydrocortisone alcohol, hydrocortisone acetate, hydrocortisone hemisuccinate, prednisolone terbutate, 9-alpha-fluoroprednisolone, triamcinolone acetonide, dexamethasone phosphate, flurisolide, toxicorol pivalate; anorectics, 10 such as, benzphetamine HCl chlorphentermine HCl; antibiotics, such as, tetracycline HCl, tyrothricin, cephalosporine, aminoglycosides, streptomycin, gentamycin, leucomycin, penicillin and derivatives; anti-allergic agents; antibodies, such as, monoclonal or polyclonal antibodies; anti-cholinergic agents; anti-depressants, such as, amitriptyline HCl, imipramine HCl; anti-emetics, such as, neuroleptica, for example, metopimazin, anti-emetics having a regulatory 15 effect on the motility of the intestine, such as, domperidone; anti-histaminic agents and histaminic agents, such as, diphenhydramine HCl, chlorpheniramine maleate, histamine, prophenpyridamine maleate, chlorprophenpyridamine maleate, disodium cromoglycate, meclizine; anti-hypertensive agents, such as, clonidine HCl; anti-inflammatory agents (enzymatic), such as, chymotrypsin, bromelain seratiopeptidase; anti-inflammatory agents 20 (non-steroidal), such as, acetaminophen, aspirin, aminopyrine, phenylbutazone, colchicine, probenecid; anti- inflammatory agents (steroidal), such as, fluticasone, prednisolone, triaxncinolone acetonide; anti-neoplastic agents, such as, actinomycin C.; anti-septics, such as, chlorhexidine HCl, hexylresorcinol, dequalinium chloride, ethacridine; anti-tussive expectorant (asthmatic agents), such as, sodium cromoglycate, isoproterenol HCl; anti-viral and anti-cancer 25 agents such as interferons (such as alpha-2 interferon for treatment of common colds), phenyl-p-guanidino benzoate, enviroxime, etc.; beta-adrenergic blocking agents, such as, propranolol HCl; blood factors, such as, factor VII, factor VIII; bone metabolism controlling agents, such as, vitamin D₃; bronchoisters, such as, clenbuterol HCl, bitolterol mesylate; cardiotonics such as digitalis; cardiovascular regulatory hormones, drugs and derivatives, such as, bradykin 30 antagonists, atrial natriuretic peptide and derivatives, such as, hydralazine, angiotensin II antagonist, nitroglycerin, propranolol, clofilium rosylate; chemotherapeutic agents, such as,

sulphathiazole, nitrofurazone; CNS-stimulants, such as, lidocaine, cocaine; corticosteroids, such as, lacicortone, hydrocorticeone, fluocinolone acetonide, triamcinolone acetonide; enzymes, such as, lysozyme chloride, dextranase; gastrointestinal hormones and derivatives, such as, secretin, substance P; hypothalamus hormones and derivatives, such as, LHRH and

5 analogues (such as naferelin, buserelin, zolidex), TRH (thyrotropin releasing hormone); hypotensives; local anaesthetics, such as, benzocaine; migraine treatment substances, such as, dihydroergot amine, ergometrine, ergotamine, pizotizin; pancreatic hormones and derivatives, such as, insulin (hexameric/dimeric/ monomeric forms); parasympathomimetics, such as, nicotine, methacholine; parasympatholytics, such as, scopolamine, atropine, ipratropium;

10 Parkinson's disease substances, such as, apomorphin; pituitary gland hormones and derivatives, such as, growth hormone (e.g. human), vasopressin and analogues (DDAVP, Lypressin); prostaglandins, such as, PGA and derivatives thereof, PGE₁ and derivatives thereof, PGE₂ and derivatives thereof, PGF₁ and derivatives thereof, dinoprost trometamol; protease inhibitors, such as, citrate, or α_1 -antitrypsin; sex-hormones, such as, ethinyloestradiol, levonorgestrel,

15 FSH, LH, LTH, estradiol-17-beta, norethindrone; sympathomimetics, such as, phenylephrine, xylometazoline, tramazoline, dopamine, dobutamine; sleep-aids, such as granistron and ramelteon, tranquilizers, such as, brotizolam, camazepam, chlorazepic acid, cloxazolam, delorazepam, estazolam, ethyl loflazepate, fludiazepam, flutazolam, halazepam, haloxazolam, ketazolam, loprazolam, lormetazepam, nimetazepam, nitrazepam, nordiazepam, oxazepam,

20 pinazepam, prazepam, temazepam, tetrazepam, tofisopam; vaccines, such as, AIDS-vaccines, parainfluenza virus, polio, rhinovirus type 13, respiratory syncytial virus; vasoconstrictors, such as, phenylephrine HCl, tetrahydrozoline HCl, naphazoline nitrate, oxymetazoline HCl, tramazoline HCl; vasodilators, such as, papaverine HCl, Substance P, vasoactive intestinal peptide (VIP).

25 [0046] Certain exemplary formulations comprising alkoxy-polyethylene glycol also comprise one or more therapeutic agents selected from the group consisting of peptide drugs, such as, oxytocin, vasopression (desmopresin), insulin, calcitonin, elcatonin, cyanocobalmin B₁₂, and glucagon-like protein-1 (GLP-1), and small organic molecules, such as, dinoprsoone, misoprostol, apomorphine, fentanyl, metoclopramide, butorphanol, and midazolam

C. Other Excipients

[0047] It is understood that the compositions of the invention, in addition to the alkoxy-polyethylene glycol and the therapeutic agent, can comprise a number of other excipients known to those skilled in the art, including absorption promoters, buffering agents, water absorbing polymers, alcohols, lipids, osmotic pressure controlling agents, pH-controlling agents, preservatives, propellants, surfactants, enzyme inhibitors, excipients for adjusting hydrophilic-lipophilic balance (HLB) and stabilizers.

[0048] Exemplary surfactants, include, for example nonoxynol, octoxynol, tweens, spans, sodium lauryl sulfate, and sorbitan monopalmitate. Exemplary absorption promoters include, for example, bile salts and derivatives thereof, fusidic acid and derivatives thereof, oleic acid, lecithin, lysolechitins, dodecanoyle phosphatidylcholine (DDPC), sucrose monododecanoate, n-dodecyl- β -D-maltopyranoside, pectin, chitosan, α -, β - and γ -cyclodextrins and derivatives thereof, pegylated caprylic-/capric glycerides and derivatives thereof, such as, Softigen and Labrasol. Exemplary water absorbing polymers include, for example, polyethylene glycols having an average molecular weight ranging from 200 to 7500, propylene glycol, or mixtures thereof, or single ethylene glycols such as tetraethylene glycol and pentaethylene glycol. Exemplary alcohols include, for example, ethanol, isopropyl alcohol. Exemplary lipids include, for example, vegetable oil, soyabean oil, peanut oil, coconut oil, maize oil, olive oil, sunflower oil, monoglycerides, diglycerides, mono/diglycerides, mono/di/triglycerides. Exemplary osmotic pressure controlling agents include, for example, glycerol, dextrose, maltose, sucrose, mannitol, xylitol, various salts (for example, sodium chloride). Exemplary pH-controlling agents include, for example, buffers, acids (for example, nitric acid, phosphoric acid, or acetic acid). Exemplary preservatives include, for example, methyl paraoxybenzoate, phenyl ethyl alcohol or benzoic acid. Exemplary propellants, include, for example, butane or air displacement such as nitrogen. Excipients adjusting the HLB of the formulation include, for example, Tween 20, 25, 40, 45, 65, 85, Span 20-80, Brij 30-98, acacia. Exemplary enzyme inhibitors include, for example aprotinin and other peptidase inhibitors, diisopropylfluorophosphate (DFP), carbopol. Exemplary stabilizers include, for example, cyclodextrins.

- 20 -

[0049] Although it is understood that the alkoxy-polyethylene glycols described herein, for example, methoxy-polyethylene glycol, can solubilize poorly soluble therapeutic agents, under certain circumstances, it may be helpful to include additional compounds that enhance the solubility of the therapeutic agent. Examples of such solubilizers include, for example, 5 alcohols and polyols, such as ethanol, isopropanol, butanol, benzyl alcohol, ethylene glycol, propylene glycol, butanediols and isomers thereof, glycerol, pentaerythritol, sorbitol, mannitol, transcutol, dimethyl isosorbide, polyethylene glycol, polypropylene glycol, pegylated-mono/di-caprylic/capric glycerides, polyvinylalcohol, hydroxypropyl methylcellulose and other cellulose derivatives, cyclodextrins (for example, α -, β -, or γ -cyclodextrins) and cyclodextrin 10 derivatives; ethers of polyethylene glycols having an average molecular weight of about 200 to about 6000 or tetrahydrofurfuryl alcohol PEG ether (glycofurool, available commercially from BASF under the trade name Tetraglycol); surfactants, such as, sodium lauryl sulfate, oleic acid, linoleic acid, monoolein, lecithin, lysolecithin, deoxycholate, taurodeoxycholate, glycochenodeoxycholate, polyoxyethylene X-lauryl ether, where X is from 9 to 20, sodium 15 tauro-24, 25-dihydrofusidate, polyoxyethylene ether, polyoxyethylene sorbitan esters, p-t-octylphenoxy polyoxyethylene, N-lauryl- β -D-maltopyranoside, 1-dodecylazacycloheptane-2-azone; amides, such as, 2-pyrrolidone, 2-piperidone, caprolactam, N-alkylpyrrolidone, N-hydroxyalkylpyrrolidone, N-alkylpiperidone, N-alkylcaprolactam, dimethylacetamide, and polyvinylpyrrolidone; esters, such as, ethyl propionate, tributylcitrate, acetyl triethylcitrate, 20 acetyl tributyl citrate, triethylcitrate, ethyl oleate, ethyl caprylate, ethyl butyrate, triacetin, propylene glycol monoacetate, propylene glycol diacetate, caprolactone and isomers thereof, valerolactone and isomers thereof, β -butyrolactone and isomers thereof; and other solubilizers 25 known in the art, such as dimethyl acetamide, dimethyl isosorbide (Arlasolve DMI (ICI)), N-methyl pyrrolidones (Pharmasolve (ISP)), mono-octanoin, and diethylene glycol monoethyl ether (available from Gattefosse under the trade name Transcutol).

[0050] Preferred additional solubilizers include triacetin, triethylcitrate, ethyl oleate, ethyl caprylate, dimethylacetamide, N-methylpyrrolidone, N-hydroxyethylpyrrolidone, polyvinylpyrrolidone, hydroxypropyl methylcellulose, hydroxypropyl cyclodextrins, ethanol, polyethylene glycol 200-1000, PEG 300, PEG 400, Transcutol, and dimethyl isosorbide, 30 sorbitol, glycerol, triacetin, glycofurool and propylene glycol. Typically, the solubilizer, if present, is present in an amount of from about 0.1 % (w/v) to about 50 % (w/v), from about 1

% (w/v) to about 40 % (w/v) or from about 2 % (w/v) to about 25 % (w/v). In addition, the liquid pharmaceutical composition can comprise water, for example, from about 2 % (v/v) to about 99 % (v/v), from about 10 % (v/v) to about 95 % (v/v), or from about 20 % (v/v) to about 90 % (v/v), of the liquid composition.

5 [0051] As discussed, the composition can comprise a preservative. In addition or in the alternative, the composition can be sterilized. Sterilization can be achieved by filter sterilization, autoclaving, exposure to ionizing radiation, for example, gamma radiation, UV irradiation, and chemical sterilization. In one embodiment, the sterile composition has a sterility assurance level of at least about 10^3 . The resulting liquid compositions preferably are
10 stable at room temperature, such that less than 5%, 4%, 3%, 2% or 1 % by weight of the therapeutic agent degrades after storage for 30 days, or more preferably 6 months, at 20°C.

15 [0052] In addition, the formulations may also include a sweetener or flavoring agent. Exemplary sweeteners or flavoring agents include, for example, acacia syrup, acesulfame potassium, anethole, anise oil, aromatic elixir, aspartame, benzaldehyde, benzaldehyde elixir, cyclodextrins, caraway, caraway oil, cardamom oil, cardamom seed, cardamom spirit, cardamom tincture, cherry juice, cherry syrup, cinnamon, cinnamon oil, cinnamon water, citric acid, citric acid syrup, clove oil, cocoa, cocoa syrup, coriander oil, dextrose, eriodictyon, eriodictyon fluid extract, eriodictyon syrup, aromatic, ethylacetate, ethyl vanillin, fennel oil, ginger, ginger fluid extract, ginger oleoresin, glucose, sugar, maltodextrin, glycerin, glycyrrhiza, glycyrrhiza elixir, glycyrrhiza extract, glycyrrhiza extract pure, glycyrrhiza fluid extract, glycyrrhiza syrup, honey, iso-alcoholic elixir, lavender oil, lemon oil, lemon tincture, maltodextrin, maltose, mannitol, methyl salicylate, menthol, nutmeg oil, orange bitter, elixir, orange bitter, oil, orange flower oil, orange flower water, orange oil, orange peel, bitter, orange peel sweet, tincture, orange spirit, orange syrup, peppermint, peppermint oil, peppermint spirit, 20 peppermint water, phenylethyl alcohol, raspberry juice, raspberry syrup, rosemary oil, rose oil, rose water, saccharin, saccharin calcium, saccharin sodium, sarsaparilla syrup, sarsaparilla compound, sorbitol solution, spearmint, spearmint oil, sucrose, sucralose, syrup, thyme oil, tolu balsam, tolu balsam syrup, wintergreen oil, vanilla, vanilla tincture, vanillin, wild cherry syrup, xylitol, or combinations thereof.

- 22 -

[0053] In addition, the formulations optionally can contain a taste masking agents.

Exemplary masking agents include, for example, cyclodextrins, cyclodextrin emulsions, cyclodextrin particles, cyclodextrin complexes, or combinations thereof.

[0054] The foregoing list of excipients and additives is by no means complete, and it is
5 understood that a person of ordinary skill in the art can choose other excipients and additives
from the GRAS (generally regarding as safe) list of chemicals used in pharmaceutical
preparations and those that are currently allowed in topical and parenteral formulations.

[0055] Exemplary liquid compositions of the invention contain, for example, the active
ingredient (for example, midazolam), 40 % (v/v) to 70 % (v/v) methoxy-polyethylene glycol
10 (for example, mPEG 350), 0 % (v/v) to 20 % (v/v) polyethylene glycol (for example, PEG
400), 0 % (v/v) to 10 % (v/v) propylene glycol, and 0 % (v/v) to 5 % (v/v) ethanol. Other
exemplary liquid compositions of the invention contain, for example, the active ingredient
dissolved in 50 % (v/v) to 70 % (v/v) mPEG 350, 1 % (v/v) to 4 % (v/v) propylene glycol, and
1 % (v/v) to 4 % (v/v) ethanol.

[0056] It is understood that the choice and amounts of each of the therapeutic agents,
alkoxy-polyethylene glycol and other excipients combined to produce the compositions of the
invention will depend upon the ultimate use of the composition, and the intended therapy and
mode of administration. When the liquid compositions ultimately are administered to a patient,
the amount of a given excipient, unless the circumstances dictate otherwise, preferably is
20 limited to a bioacceptable amount, which is readily determined by one of skill in the art.
Furthermore, it is understood that the liquid compositions of the invention can be formulated
using techniques known to those skilled in the art. A thorough discussion of formulations and
the selection of pharmaceutically acceptable carriers, stabilizers, etc. can be found, for
example, in *Remington's Pharmaceutical Sciences* (18th Ed.), Mack Publishing Company,
25 Eaton, PA.

[0057] Although the therapeutic agent and the alkoxy-polyethylene glycol are combined in
the liquid formulations of the invention, they are not covalently linked to one another. In
certain embodiments, the liquid formulations of the invention are free or are substantially free
of chitosan. In certain embodiments, the compositions of the invention may be prepared in a
30 powder form.

II Modes of Administration and Pharmacokinetics

[0058] The compositions of the invention are particularly useful in delivering one or more therapeutic agents to a mucosal membrane or the skin of a mammal, for example, a human. The mucosal membrane to which the pharmaceutical preparation of the invention is administered may be any mucosal membrane of the mammal to which the therapeutic agent is to be applied, for example, the nose (for example, via a nasal membrane), vagina, eye (for example, via an ocular membrane), ear (for example, via a tympanic membrane), mouth (for example, via the buccal membrane), lungs (for example, via the pulmonal membrane), or rectum (for example, via the rectal membrane). The compositions are particularly useful in delivering a therapeutic agent to the mucosa of the nose, mouth (buccal, gingual, sublingual or to the hard palate), or the vagina.

[0059] It is understood that the compositions of the invention are particularly useful in the intranasal delivery of a therapeutic agent. When the composition is applied to the nasal mucosa, the volume of the pharmaceutical composition applied typically is in the range of 1-1000 μ L, preferably not more than 700 μ L, more preferably 50-150 μ L per nostril, and most preferably about 100 μ L/nostril.

[0060] It is understood that when administered intranasally, the compositions are delivered via a spray device that produces a plume of spray droplets that contact the nasal mucosa. It is contemplated that the compositions can be delivered using commercially available spray devices available from, for example, Pfeiffer of America, Princeton, NJ; Valois of America, Inc., Greenwich, CT; or Becton Dickinson, Franklin Lakes, NJ. Furthermore, these devices are easily operable by the patient or care giver, and leave little or no residual formulation in the device after use.

[0061] Such devices can be filled with single or multi-dose amounts of the desired formulation. The container holding the pharmaceutical composition and its sealing means are sterilizable. At least the parts of the device that are in contact with the pharmaceutical composition should be constructed and assembled in a configuration that can be sterilized. Exemplary delivery devices with one or more unit-dose(s) are described, for example, in U.S. Patent Nos. 4,946,069; 5,307,953; 6,948,492; and 6,446,839. Individual devices can be packaged, sterilized and shipped; alternatively, entire shipping and storage packages can be

sterilized at once, and the devices removed individually for dispensing, without affecting the sterility of the remaining units.

[0062] The mucosal epithelium in the nasal cavity is covered with many hair-like cilia that provide an important defense mechanism against inhaled dust, allergens and microorganisms.

5 The normal half-time for non-absorbed substances administered to the nasal cavity is about 15 minutes due to the mucociliary clearance removing foreign particles and excess mucus toward the pharynx. For this reason it is preferred that the absorption occurs rapidly and preferably within 0.5 to 20 minutes. However, in the current invention due to bioadhesive properties of the invention, the preferred absorption may occur within 0.5 to 300 minutes (e.g., for vaccines 10 and biologicals), preferably between 0.5 to 60 minutes (e.g., for large molecules) and more preferably between 0.5 and 20 minutes, for example, within 2, 3, 4, 5, 10, 15 or 20 minutes post administration. The composition can be formulated so that upon administration to a subject, for example, via intranasal administration to the subject, the therapeutic agent has a peak concentration (T_{max}) in the blood of the subject within 30, 25, 20, 15, 10 8, 5, 3 or 2 minutes 15 after administration of the therapeutic agent.

[0063] In addition to administration to humans, the compositions of the invention can be used to deliver the therapeutic agent to an animal, for example: pets, for example, dogs, cats, rabbits, and guinea pigs; and farm animals, for example, horses, sheep, pigs, cattle, and chickens.

20 **[0064]** Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where processes are described as having, including, or comprising specific process steps, the processes also consist essentially of, or consist of, the recited processing steps. Except where indicated otherwise, the order of steps or 25 order for performing certain actions are immaterial so long as the invention remains operable. Moreover, unless otherwise noted, two or more steps or actions may be conducted simultaneously.

EXAMPLES

[0065] The invention is explained in more detail with reference to the following Examples, which are to be considered as illustrative and not to be construed so as to limit the scope of the invention as set forth in the appended claims.

5 EXAMPLE 1 – Exemplary Lorazepam Containing Formulation

[0066] In this Example, 10 mg lorazepam is dissolved in 2 mL of mPEG 350 using ultrasound to obtain a solution containing 5 mg/mL of lorazepam. 50 μ L of the resulting composition is administered into each nasal cavity of male New Zealand White rabbits held in a supine position during, and then one minute after application. An Eppendorf pipette is used 10 for each application. After administration, blood samples then are harvested from a marginal ear vein at 0, 2, 5, 10, 15, 30 and 60 minutes, and the lorazepam concentration determined by high performance liquid chromatography (HPLC). The pharmacokinetics of lorazepam delivery via intranasal administration can then be compared with the pharmacokinetics of lorazepam delivery by intravenous administration. It is contemplated that the pharmacokinetics 15 of the intranasally administered lorazepam will be comparable to those of the intravenously administered lorazepam.

EXAMPLE 2 – Exemplary Midazolam Containing Formulation

[0067] In this Example, 10 mg of midazolam is dissolved in 2 mL of mPEG 350 using ultrasound to obtain a solution containing 5 mg/mL midazolam. 50 μ L of the resulting 20 composition is then administered into each nasal cavity of male New Zealand White rabbits, held in a supine position during, and then one minute after application. After administration, blood samples then are harvested from a marginal ear vein at 0, 2, 5, 10, 15, 30 and 60 minutes, and the midazolam concentration determined by HPLC. The pharmacokinetics of midazolam delivery via intranasal administration can then be compared with the pharmacokinetics of 25 midazolam delivery by intravenous administration. It is contemplated that the pharmacokinetics of the intranasally administered midazolam will be comparable to those of the intravenously administered midazolam.

- 26 -

EXAMPLE 3 – Pharmacokinetics of Exemplary Midazolam Formulations

[0068] This Example describes a variety of formulations containing methoxy-polyethylene glycol that demonstrate comparable pharmacokinetic properties to a control formulation containing polyethylene glycol and propylene. In addition to having comparable 5 pharmacokinetic properties, the methoxy-polyethylene glycol formulations had a lower viscosity than the control formulation.

[0069] Table 1 lists the composition of a first test formulation, Table 2 lists the composition of a second test formulation, and Table 3 lists the composition of a control formulation.

10

TABLE 1 – Test Formulation 1 Containing 50 mg/mL Midazolam base

Component	Percent of Final (% v/v)
mPEG 350	47
PEG 400	18
Propylene Glycol	5
Ethanol	7
Water	21

TABLE 2 – Test Formulation 2 Containing 50 mg/mL Midazolam base

Component	Percent of Final (% v/v)
mPEG 350	47
PEG 400	18
Propylene Glycol	10
Polysorbate 80	0.1
Water	22

- 27 -

TABLE 3 – Control Containing 50 mg/mL Midazolam base

Component	Percent of Final (% v/v)
PEG 400	18
Propylene Glycol	78
Butylatedhydroxytoluene	0.01

[0070] Each of the test and control formulations contained 50 mg/mL midazolam. Once made, 5 mg of midazolam in 0.1 mL was delivered intranasally to the right nostril of each dog 5 (3 dogs per set) for each formulation in a cross over manner. Blood was harvested from the dogs predose and at 0.03, 0.08, 0.16, 0.25, 0.5, 0.75, 1, 2, and 4 hours after administration, and the concentration of midazolam in the blood measured by HPLC. The pharmacokinetic properties are summarized in Table 4.

TABLE 4 – Pharmacokinetic Properties

Formulation	T _{max} (hr)	C _{max} (ng/mL)	AUC _{last} (hr*ng/mL)	AUC _{INF} (hr*ng/mL)
Test Formulation 1	0.11 ± 0.05	2220 ± 289	506 ± 94	508 ± 95
Test Formulation 2	0.05 ± 0.03	1880 ± 682	399 ± 75	403 ± 76
Control Formulation	0.07 ± 0.03	2350 ± 796	506 ± 72	509 ± 71

10

In Table 4, C_{max} refer to the maximum plasma concentration, T_{max} refers to the time to reach C_{max}, AUC_{last} refers to area under the concentration curve from time zero to the last measurable plasma concentration, and AUC_{INF} refers to the area under the concentration curve from time zero to infinity.

15 [0071] The results demonstrate that the test formulations, when administered intranasally to the dogs, produced comparable pharmacokinetics to the control formulation. The test formulations, however, had a lower viscosity than the control formulations. Lower viscosity correlates with better spray pattern characteristics.

EXAMPLE 4 – Additional Exemplary Methoxy–Polyethylene Glycol Containing Formulations

[0072] This example describes the benefit of incorporating methoxy-polyethylene glycol into certain intranasal formulations. Table 5 describes a formulation containing methoxy-

- 28 -

polyethylene glycol and PEG 400, Table 6 describes a formulation containing methoxy-polyethylene glycol without PEG 400, Table 7 describes a formulation containing methoxy-polyethylene glycol without PEG 400 (with reduced ethanol), and Table 8 describes a control formulation containing PEG 400 but no methoxy-polyethylene glycol.

5

TABLE 5

Component	Viscosity (cP)	% (w/w)
Propylene glycol	40	5
PEG 400	80	18
mPEG 350	29	47
Ethanol	-	6
Water	1	18
Midazolam	-	7

TABLE 6

Component	Viscosity (cP)	% (w/w)
Propylene glycol	40	2
mPEG 350	29	64
Ethanol	-	5
Water	1	23
Midazolam	-	7

10

TABLE 7

Component	75 mg Midazolam/dose (% w/w)	50 mg Midazolam/dose (% w/w)	25 mg Midazolam/dose (% w/w)
MPEG 350	64.8	64.8	64.8
Propylene glycol	2.0	2.0	2.0
Ethanol	2.5	2.5	2.5
Water	23.7	26.0	28.3
Phenethyl Alcohol	0.025	0.025	0.025
Midazolam	6.9	4.6	2.3

TABLE 8 – Control Formulation (Without Methoxy-polyethylene Glycol)

Component	Viscosity (cP)	% (w/w)
PEG 400	89	19
Propylene Glycol	40	76
Butylatedhydroxytoluene	-	0.01
Midazolam	-	5

[0073] The formulation of Table 5 was prepared by mixing the propylene glycol, PEG 400, mPEG 350, and ethanol. The midazolam was then added to the mixture, and then after the 5 midazolam had dissolved following mixing the water was added to the formulation. The formulations in Tables 6 and 7 were prepared as follows. The midazolam was weighed in a container and the ethanol was added to wet the active ingredient. After mixing, about 65% of the mPEG 350 was added, and the resulting mixture mixed for 2 minutes. Thereafter, the additional organic phase was added. The remaining 35% of the mPEG 350 was mixed with 10 water, and the diluted mPEG 350 then was gradually added with mixing to the midazolam solution until a clear solution was produced. The viscosity of the resulting formulations and spray patterns were tested.

[0074] The viscosity of the resulting formulations was tested at 25°C using a viscometer (Brookfield DV-II PRO). The viscosity of the control formulation without methoxy-polyethylene glycol (Table 8) was found to be about 42 cP, whereas the viscosity of the test 15 formulation of Table 5 was found to be about 30 cP, and the viscosity of the test formulation of Table 6 was found to be about 23 cP. The reduced viscosity of the test formulations relative to the control formulation permitted the creation of more reproducible spray patterns.

[0075] The spray patterns produced by the two test formulations of Tables 5 and 6 were 20 tested using a Proveris spray view instrument. The resulting parameters of the resulting spray patterns were found to be more reproducible as the viscosity of the formulation decreased.

[0076] In addition, 100 µL of placebo formulations (no midazolam) corresponding to the formulations set forth in Tables 7 and 8 were administered intranasally to three healthy test subjects using a commercially available spray device from Pfeiffer. The placebo formulation 25 corresponding to Table 7 containing methoxy-polyethylene glycol had a less noticeable taste relative to the placebo formulation corresponding to Table 8.

[0077] These collective results demonstrate that mPEG 350 is a suitable excipient for nasal administration, for example, with midazolam.

EXAMPLE 5 – Sprayability of Methoxy-Polyethylene Glycol Containing Formulations

5 [0078] This Example demonstrates that methoxy-polyethylene glycol-based formulations produce superior spray plumes relative to polyethylene glycol. Solutions containing 100 % methoxy-polyethylene glycol 350 (MPEG 350) from Sigma-Aldrich Chemie GmbH (St. Louis, MO, USA) and 100 % polyethylene glycol 300 (PEG 300) from Croda Chemicals Europe Ltd. (Goole, UK) were placed into Pfeiffer 20 mL bottles (Pfeiffer 34473) and attached
10 to certain pumps from Valois or Pfeiffer (see Table 8). Each spray device was placed 25 cm under a sheet of absorbent paper and the sprayability was measured as the diameter across of the paper wetted by the resulting spray. The results are summarized in Table 9.

TABLE 9

Pump Type	Formulation	Spray diameter (cm)
Pfeiffer 71514	PEG 300	2.0-3.2 cm
Pfeiffer 71514	mPEG 350	22-26 cm
Valois VP6/100	PEG 300	5.0-7.5 cm
Valois VP6/100	mPEG 350	15-21 cm


15 [0079] The results show that there is a clear difference in the sprayability of mPEG 350 compared with PEG 300. The required spray angle is not achieved using PEG 300, whereas a clinically relevant spray-angle can be achieved using mPEG 350.

[0080]

[0081] The scope of the claims that follow is not limited by the embodiments set forth in the description. The claims should be given the broadest purposive construction consistent with the description as a whole.

WHAT IS CLAIMED IS:

1. A liquid pharmaceutical composition formulated for administration through a nasal mucosal membrane comprising a therapeutic agent and an alkoxypolyethylene glycol represented by Formula I:



wherein:

R is methyl; and

n is a number in the range of from about 1 to about 25.

2. The liquid pharmaceutical composition of claim 1, wherein the therapeutic agent is an organic compound having a molecular weight of less than 1500 g/mol.
3. The liquid pharmaceutical composition according to any one of claims 1 and 2, wherein the alkoxy-polyethylene glycol comprises from about 0.5% (v/v) to about 70% (v/v) of the composition.
4. The liquid pharmaceutical composition according to any one of claims 1, 2 and 3, wherein the alkoxy-polyethylene glycol comprises from about 1% (v/v) to about 60% (v/v) of the composition.
5. The liquid pharmaceutical composition according to any one of claims 1, 2 and 3, wherein the alkoxy-polyethylene glycol comprises from about 5% (v/v) to about 50% (v/v) of the composition.
6. The liquid pharmaceutical composition according to any one of claims 1-5, wherein the therapeutic agent comprises from about 0.001% (w/v) to about 20 % (w/v) of the composition.
7. The liquid pharmaceutical composition according to any one of claims 1-5, wherein the therapeutic agent comprises about 0.01% (w/v) to about 10 % (w/v) of the composition.
8. The liquid pharmaceutical composition according to any one of claims 1-7, further comprising water.
9. The liquid pharmaceutical composition of claim 9, wherein the water comprises about 10% (v/v) to about 95 % (v/v) of the composition.

10. The liquid pharmaceutical composition of any one of claims 1-9, further comprising a buffer.
11. The liquid pharmaceutical composition of any one of claims 1-10, further comprising a preservative.
12. The liquid pharmaceutical composition of any one of claims 1-11, wherein the composition at a temperature of 20°C has a viscosity in the range of from about 1.5 cP to about 60 cP.
13. The liquid pharmaceutical composition of any one of claims 1-11, wherein the composition at a temperature of 20°C has a viscosity in the range of from about 5 cP to about 25 cP.
14. The liquid pharmaceutical composition of any one of claims 1-13, wherein the composition has a pH in the range of from about 4.5 to about 8.5.
15. The liquid pharmaceutical composition of any one of claims 1-13, wherein the composition has a pH in the range of from about 5.5 to about 7.5.
16. The liquid pharmaceutical composition of any one of claims 1-15, wherein the composition is sterile.
17. The liquid pharmaceutical composition of any one of claims 1-16, wherein the composition has a sterility assurance level of at least about 10^3 .
18. The liquid pharmaceutical composition of any one of claims 1-17, wherein the composition is formulated so that upon intranasal administration to a subject, the therapeutic agent has a peak concentration (T_{max}) in the blood of the subject within 30 minutes after administration of the therapeutic agent.
19. The liquid pharmaceutical composition of any one of 1-17, wherein the composition is formulated so that upon intranasal administration to a subject, the therapeutic agent has a peak concentration (T_{max}) in the blood of the subject within 10 minutes after administration of the therapeutic agent.
20. The liquid pharmaceutical composition according to any one of claims 1-19, wherein n is 3-15.

21. The liquid pharmaceutical composition of any one of claims 1-20, wherein the alkoxy-polyethylene glycol is mPEG 350, mPEG 550, or a combination thereof.
22. The liquid pharmaceutical composition of any one of claims 1-20 wherein the alkoxy-polyethylene glycol is methoxy-triethylene glycol (m3EG), methoxy-tetraethylene glycol (m4EG), methoxy-pentaethylene glycol (m5EG), methoxy-hexaethylene glycol (m6EG), methoxy-heptaethylene glycol (m7EG), methoxy-octaethylene glycol (m8EG), methoxy-nonaethylene glycol (m9EG), methoxy-decaethylene glycol (m10EG), methoxy-undecaethylene glycol (m11EG), methoxy-dodecaethylene glycol (m12EG), methoxy-tridecaethylene glycol (m13EG), or methoxy-tetradecaethylene glycol (m14EG).
23. The liquid pharmaceutical composition of any one of claims 1-22, wherein the therapeutic agent has a molecular weight of less than 500 g/mol.
24. The liquid pharmaceutical composition according any one of claims 1-22, wherein the therapeutic agent has an aqueous solubility of less than about 0.3 mg/mL at pH 7 and 20°C.
25. The liquid pharmaceutical composition of any one of claims 1-24, wherein the therapeutic agent has an aqueous solubility of less than about 0.1 mg/mL at pH 7 and 20°C.
26. The liquid pharmaceutical composition of any one of claims 1-25, wherein the therapeutic agent is a benzodiazepine or a pharmaceutically acceptable salt thereof.
27. The liquid pharmaceutical composition according to claim 26, wherein the therapeutic agent is midazolam or a pharmaceutically acceptable salt thereof.
28. A use of the liquid pharmaceutical composition of any one of claims 1-27 in the preparation of a medicine for administration of a therapeutic agent to a mucosal surface of a mammal.
29. The use of claim 28, wherein the therapeutic agent has a peak concentration (T_{max}) in the blood of the mammal within 30 minutes after administration of the therapeutic agent.
30. The use of claim 28, wherein the therapeutic agent has a peak concentration (T_{max}) in the blood of the mammal within 10 minutes after administration of the therapeutic agent.
31. The use of any one of claims 28, 29 and 30, wherein the liquid pharmaceutical composition has a volume of from about 1 μ L to about 1000 μ L.

-35-

32. The use of claim 31, wherein the liquid pharmaceutical composition has a volume from about 50 μ L to about 300 μ L.

33. The use of any one of claims 28-32, wherein the mammal is a human.