

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 4058474 T3**

-
- (51) Int.Cl.: **C 07 K 14/705 (2006.01)** **A 61 K 35/17 (2015.01)** **A 61 P 35/00 (2006.01)**
C 07 K 16/28 (2006.01) **C 12 N 5/10 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-07-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-05-08**
- (86) Europæisk ansøgning nr.: **21769291.2**
- (86) Europæisk indleveringsdag: **2021-08-19**
- (87) Den europæiske ansøgnings publiceringsdag: **2022-09-21**
- (86) International ansøgning nr.: **US2021046733**
- (87) Internationalt publikationsnr.: **WO2022040444**
- (30) Prioritet: **2020-08-20 US 202063068249 P** **2020-10-26 US 202063105639 P**
2021-08-06 US 202163230632 P
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **A2 Biotherapeutics, Inc., 30301 Agoura Road Suite 210, Agoura Hills CA 91301, USA**
- (72) Opfinder: **OH, Julyun, , Agoura Hills, California 91301, USA**
XU, Han, , Agoura Hills, California 91301, USA
KAMB, Carl Alexander, , Agoura Hills, California 91301, USA
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **SAMMENSÆTNINGER OG FREMGANGSMÅDER TIL BEHANDLING AF EGFR-POSITIVE KRÆFTFORMER**
- (56) Fremdragne publikationer:
WO-A1-2015/136001
WO-A1-2019/068007
WO-A2-2020/065406
PAUCEK RICHARD D ET AL: "The Cellular Immunotherapy Revolution: Arming the Immune System for Precision Therapy", TRENDS IN IMMUNOLOGY, vol. 40, no. 4, 1 April 2019 (2019-04-01), pages 292-309, XP085646047, ISSN: 1471-4906, DOI: 10.1016/J.IT.2019.02.002
XUNLEI KANG ET AL: "Inhibitory leukocyte immunoglobulin-like receptors: Immune checkpoint proteins and tumor sustaining factors", CELL CYCLE, vol. 15, no. 1, 4 December 2015 (2015-12-04), pages 25-40, XP55439503, US ISSN: 1538-4101, DOI: 10.1080/15384101.2015.1121324
YU SHENGAN ET AL: "Chimeric antigen receptor T cells: a novel therapy for solid tumors", JOURNAL OF HEMATOLOGY & ONCOLOGY, vol. 10, no. 1, 1 December 2017 (2017-12-01), XP055872101, DOI: 10.1186/s13045-017-0444-9 Retrieved from the Internet: URL:https://jhoonline.biomedcentral.com/track/pdf/10.1186/s13045-017-0444-9.pdf>
Xunlei Kang ET AL: "Inhibitory leukocyte immunoglobulin-like receptors: Immune checkpoint proteins and

Fortsættes ...

tumor sustaining factors", CELL CYCLE, vol. 15, no. 1, 4 December 2015 (2015-12-04), pages 25-40,
XP55439503, US ISSN: 1538-4101, DOI: 10.1080/15384101.2015.1121324

Description**RELATED APPLICATIONS**

5 **[0001]** This application claims priority to U.S. Provisional Application No. 63/068,249, filed on August 20, 2020; U.S. Provisional Application No. 63/105,639, filed on October 26, 2020; and U.S. Provisional Application No. 63/230,632, filed on August 6, 2021.

TECHNICAL FIELD

10 **[0002]** This application relates generally to engineered receptors and immune cell compositions comprising said receptors for use in cell therapy applications.

REFERENCE TO SEQUENCE LISTING

15 **[0003]** The sequence listing paragraph application contains a Sequence Listing, which has been submitted in ASCII format via EFS-WEB. Said ASCII copy, created on August 17, 2021 is named A2BI_021_02WO_SeqList_ST25.txt and is 675 KB in size.

BACKGROUND

20 **[0004]** Cell therapy is a powerful tool for the treatment of various diseases, particularly cancers. In conventional adoptive cell therapies, immune cells are engineered to express specific receptors, for example chimeric antigen receptors (CARs) or T cell receptors (TCRs), which direct the activity of the immune cells to cellular targets via interaction of the
 25 receptor with a ligand expressed by the target cell. For example, WO 2020/065406 A2 provides a method for identifying a target pair comprising i) an inhibitory chimeric antigen receptor (iCAR) or a protective chimeric antigen receptor (pCAR), wherein the iCAR or pCAR target is directed to a target extracellular polymorphic epitope, and ii) an activating chimeric antigen receptor (aCAR), wherein the aCAR is directed to a target non-polymorphic cell surface epitope of a protein, as
 30 well as methods of making and use of such pairs in the treatment of cancer. However, identification of suitable target molecules remains challenging, as many targets are expressed in normal tissues. This expression can lead to toxicity when the transplanted cells target normal tissues expressing target molecules. There is thus a need in the art for compositions and methods useful in the treatment of disease, particularly cancers, by adoptive cell therapy.

SUMMARY

35 **[0005]** The disclosure provides compositions and methods for increasing the specificity of immune cells used in adoptive cell therapy. The disclosure provides immune cells comprising a two-receptor system that increases the specificity of the immune cells for target cells expressing a target antigen. The immune cells comprise a first, activator receptor that activates the immune cells in response to binding of the first receptor by the target antigen. The immune cells further
 40 comprise a second, inhibitory receptor specific to a non-target antigen. This second receptor inhibits activation of the immune cells when the second receptor is bound by the non-target antigen, even when the first receptor is bound by the target antigen.

[0006] The present invention provides an immune cell, comprising: a.) a first receptor, comprising an extracellular ligand binding domain specific to Epidermal Growth Factor Receptor (EGFR); and b.) a second receptor, comprising an
 45 extracellular ligand binding domain specific to a non-target antigen lost in EGFR+ cancer due to loss of heterozygosity, wherein the first receptor is an activator receptor responsive to EGFR; and wherein the second receptor is an inhibitory receptor responsive to the non-target antigen, wherein the extracellular ligand binding domain of the second receptor specifically binds HLA-A*02, and wherein the extracellular ligand binding domain of the first receptor comprises complementarity determining regions (CDRs) CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3 of SEQ ID NOs:
 50 154, 155, 156, 151, 152, and 153, and wherein the extracellular ligand binding domain of the second receptor comprises complementarity determining regions CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 of the humanized scFv binding domain sequence of C-002162 PA2.1.14 as depicted in Table 4, wherein the CDR-L1 sequence is RSSQSIVH-SNGNTYLE (SEQ ID NO: 101), the CDR-L2 sequence is KVSNRFSQVGPARG, the CDR-L3 sequence is FQGSHPVPRP
 55 (SEQ ID NO: 103), the CDR-H1 sequence is ASGYTFTSYHIH (SEQ ID NO: 104), the CDR-H2 sequence is WIYPGN-VNTEYNEKFKGK (SEQ ID NO: 105), and the CDR-H3 sequence is EEITYAMDY (SEQ ID NO: 106), wherein the immune cell is a CD8+ CD4- T cell or a CD8-CD4+ T cell.

[0007] It is described herein that the extracellular ligand binding domain of the second receptor specifically binds an allelic variant of an MHC protein. It is described herein that the extracellular ligand binding domain of the second receptor

specifically binds an allelic variant of an HLA-A, HLA-B, or HLA-C protein. According to the present invention, the extracellular ligand binding domain of the second receptor specifically binds HLA-A*02. It is described herein that the extracellular ligand binding domain of the second receptor specifically binds HLA-A*01, HLA-A*03, HLA-A*11, HLA-C*07, or HLA-B*07.

[0008] It is described herein that the extracellular ligand binding domain of the second receptor comprises complementarity determining regions (CDRs) CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 as disclosed Table 5; or CDR sequences having at most 1, 2, or 3 substitutions, deletions, or insertion relative to the CDRs of Table 5.

[0009] It is described herein that the extracellular ligand binding domain of the second receptor comprises complementarity determining regions (CDRs) CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 of SEQ ID NOs: 101-106 or of SEQ ID NOs: 106-112; or CDR sequences having at most 1, 2, or 3 substitutions, deletions, or insertion relative to the CDRs of Table 5. According to the present invention, the extracellular ligand binding domain of the second receptor comprises complementarity determining regions CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 of the humanized scFv binding domain sequence of C-002162 PA2.1.14 as depicted in Table 4, wherein the CDR-L1 sequence is RSSQSIVHSNGNTYLE (SEQ ID NO: 101), the CDR-L2 sequence is KVSNRFGSGVPAR, the CDR-L3 sequence is FQGSHVPRT (SEQ ID NO: 103), the CDR-H1 sequence is ASGYTFTSYHIH (SEQ ID NO: 104), the CDR-H2 sequence is WIYPGNVNTHEYNEKFKGK (SEQ ID NO: 105), and the CDR-H3 sequence is EEITYAMDY (SEQ ID NO: 106).

[0010] It is described herein that the extracellular ligand binding domain of the second receptor comprises a polypeptide sequence selected from the polypeptide sequence disclosed in Table 4; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto. It is described herein that the extracellular ligand binding domain of the second receptor comprises any one of SEQ ID NOs: 89-100; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto. In an embodiment of the present invention, the extracellular ligand binding domain of the second receptor comprises SEQ ID NO: 93; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0011] In some embodiments of the present invention, the first receptor is a chimeric antigen receptor (CAR).

[0012] It is further described herein that the extracellular ligand binding domain of the first receptor comprises a variable heavy (VH) portion comprising a set of heavy chain complementarity determining regions (HC-CDRs) selected from the group of sequences set forth in Table 3; and/or a variable light (VL) portion comprising a set of light chain complementarity determining regions (LC-CDRs) from the group of sequences set forth in Table 3; or CDR sequences having at most 1, 2, 3, 4 substitutions, insertions, or deletions in each CDR.

[0013] It is further described herein that the extracellular ligand binding domain of the first receptor comprises a variable heavy (VH) portion having a sequence selected from the VH sequence set forth in Table 2; and/or a variable light (VL) portion comprising a sequence set forth in Table 2; or sequences having at least 70%, at least 85%, at least 90%, or at least 95% identity thereto.

[0014] In one embodiment of the present invention, the extracellular ligand binding domain of the first receptor comprises a variable heavy (VH) portion having a sequence selected from the VH sequence set forth in SEQ ID NO: 149; and/or a variable light (VL) portion comprising a sequence set forth in SEQ ID NO: 150; or sequences having at least 70%, at least 85%, at least 90%, or at least 95% identity thereto.

[0015] It is further described herein that the extracellular ligand binding domain of the first receptor comprising a sequence selected from the group of sequences set forth in Table 1; or a sequence having at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% identity thereto.

[0016] It is further described herein that the extracellular ligand binding domain of the first receptor comprises an scFv sequence selected from the group consisting of SEQ ID NO: 9-18; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0017] In one embodiment of the present invention, the extracellular ligand binding domain of the first receptor comprises the scFv sequence of SEQ ID NO: 18; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0018] In some embodiments of the present invention, the second receptor comprises a LILRB1 intracellular domain or a functional variant thereof. In some embodiments of the present invention, the LILRB1 intracellular domain comprises a sequence at least 90%, at least 95%, at least 97%, at least 99%, or is identical to SEQ ID NO: 129.

[0019] In some embodiments of the present invention, the second receptor comprises a LILRB1 transmembrane domain or a functional variant thereof. In some embodiments of the present invention, the LILRB1 transmembrane domain or a functional variant thereof comprises a sequence at least 90%, at least 95%, at least 97%, at least 99% or is identical to SEQ ID NO: 133.

[0020] In some embodiments of the present invention, the second receptor comprises a LILRB1 hinge domain or functional variant thereof. In some embodiments of the present invention, the LILRB1 hinge domain comprises a sequence at least 90%, at least 95%, at least 97%, at least 99% or is identical to SEQ ID NO: 132, SEQ ID NO: 125, SEQ ID NO: 126.

[0021] In some embodiments of the present invention, the second receptor comprises a LILRB1 intracellular domain, a LILRB1 transmembrane domain, a LILRB1 hinge domain, a functional variant of any of these, or combinations thereof.

In some embodiments of the present invention, the LILRB1 intracellular domain and LILRB1 transmembrane domain comprises SEQ ID NO: 128 or a sequence at least 95% identical to SEQ ID NO: 128.

[0022] In some embodiments of the present invention, the immune cell of the present invention is for use in the treatment of EGFR+ cancer. In some embodiments of the present invention, the EGFR+ cancer is an EGFR+ tumor having loss of heterozygosity at a locus encoding a non-target antigen and/or the EGFR+ cancer comprises EGFR+ cancer cells having loss of heterozygosity at an MHC class I locus. In some embodiments of the present invention, the EGFR+ cancer cell is a lung cancer cell, a small cell lung cancer cell, a non-small cell lung cancer cell, a pancreatic ductal carcinoma cell, a colorectal cancer cell, a head and neck cancer cell, a esophagus and gastric adenocarcinoma cell, an ovarian cancer cell, a glioblastoma multiforme cell, a cervical squamous cell carcinoma cell, a kidney cancer cell, a papillary kidney cancer cell, a kidney renal clear cell carcinoma cell, a bladder cancer cell, a breast cancer cell, a bile duct cancer cell, a liver cancer cell, a prostate cancer cell, a sarcoma cell, a thyroid cancer cell, a thymus cancer cell, a stomach cancer cell, or a uterine cancer cell.

[0023] In some embodiments of the present invention, the EGFR+ cancer cell is an EGFR+/HLA-A*02- cancer cell that does not express HLA-A*02; or a cancer cell derived from an HLA-A*02+ individual, which does not express HLA-A*02. In some embodiments of the present invention, the EGFR+/HLA-A*02- cancer cell is derived from an EGFR+/HLA-A*02+ cell by loss of heterozygosity at HLA-A leading to loss of HLA-A*02.

[0024] In some embodiments, the first receptor and the second receptor together specifically activate the immune cell in the presence of the EGFR/HLA-A*02- cancer cell having loss of heterozygosity.

[0025] In some embodiments, the first receptor and the second receptor together do not specifically activate the immune cell in the presence of an EGFR+ cell that has not lost HLA-A*02 by loss of heterozygosity.

[0026] It is also described herein that the immune cell is a T cell, macrophage, NK cell, iNKT cell, or a gamma delta T cell. According to the present invention, the T cell is a CD8+ CD4- T cell or a CD8- CD4+ T cell.

[0027] In some embodiments of the present invention, the immune cell further comprises reduced or eliminated expression and/or function of an MHC Class I gene.

[0028] In some embodiments of the present invention, the MHC Class I gene is beta-2-microglobulin (B2M).

[0029] In some embodiments of the present invention, the immune cell further comprises a polynucleotide comprising an interfering RNA, the interfering RNA comprising a sequence complementary to a sequence of a B2M mRNA (SEQ ID NO: 172). In some embodiments of the present invention, the interfering RNA is capable of inducing RNAi-mediated degradation of the B2M mRNA. In some embodiments of the present invention, the interfering RNA is a short hairpin RNA (shRNA). In some embodiments of the present invention, the shRNA comprises: a.) A first sequence, having from 5'-end to 3'-end a sequence complementary to a sequence of the B2M mRNA; and b.) A second sequence, having from 5'-end to 3'-end a sequence complementary to the first sequence, wherein the first sequence and the second sequence form the shRNA.

[0030] In some embodiments of the present invention, the immune cell comprises one or more modifications to a sequence encoding B2M (SEQ ID NO: 170), wherein the one or more modifications reduce the expression and/or eliminate the function of B2M. In some embodiments of the present invention, the one or more modifications comprise one or more inactivating mutations of the endogenous gene encoding B2M. In some embodiments, the one or more inactivating mutations comprise a deletion, an insertion, a substitution, or a frameshift mutation. In some embodiments of the present invention, the one or more inactivating mutations are introduced with a nucleic acid guided endonuclease in a complex with at least one guide nucleic acid (gNA) that specifically targets a sequence of the endogenous gene encoding B2M (SEQ ID NO: 170).

[0031] In an embodiment, the MHC Class I gene is HLA-A*02.

[0032] It is described herein that the immune cell further comprises a polynucleotide comprising an interfering RNA, comprising a sequence complementary to a sequence of an HLA-A*02 mRNA (SEQ ID NO: 171). It is further described herein that the interfering RNA is capable of inducing RNA interference (RNAi)-mediated degradation of the HLA-A*02 mRNA. It is further described herein that the interfering RNA is a short hairpin RNA (shRNA) comprising: a.) a first sequence, having from 5'-end to 3'-end a sequence complementary to a sequence of the HLA-A*02 mRNA; and b.) a second sequence, having from 5'-end to 3'-end a sequence complementary to the first sequence, wherein the first sequence and the second sequence form the shRNA.

[0033] It is described herein that the immune cell comprises one or more modifications to a sequence of an endogenous gene encoding HLA-A*02 (SEQ ID NO: 169), wherein the one or more modifications reduce the expression and/or eliminate the function of HLA-A*02. It is further described herein that the one or more modifications comprise one or more inactivating mutations of the endogenous gene encoding HLA-A*02. It is further described herein that the one or more inactivating mutations comprise a deletion, an insertion, a substitution, or a frameshift mutation. It is further described herein that the one or more inactivating mutations are introduced with a nucleic acid guided endonuclease in a complex with at least one guide nucleic acid (gNA) that specifically targets a sequence of the endogenous gene encoding HLA-A*02.

[0034] In some embodiments of the present invention, the first receptor comprises SEQ ID NO: 177, or a sequence

sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 174 or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0035] It is also disclosed herein that the first receptor comprises SEQ ID NO: 177, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0036] It is also disclosed herein that the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 174, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0037] It is also disclosed herein that the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0038] It is also disclosed herein that the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0039] It is also disclosed herein that the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0040] It is also disclosed herein that the immune cell further comprises a T2A self-cleaving peptide, wherein the T2A self-cleaving peptide comprises SEQ ID NO: 178, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0041] In some embodiments of the present invention, the immune cell further comprises an interfering RNA, wherein the interfering RNA comprises SEQ ID NO: 179, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0042] In some embodiments of the present invention, the immune cell is autologous. In some embodiments of the present invention, the immune cell is allogeneic.

[0043] In one aspect, the present invention provides a pharmaceutical composition, comprising a therapeutically effective amount of the immune cells according to the present invention. In some embodiments, the immune cell expresses both the first receptor and the second receptor. In some embodiments, at least about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% of the immune cells express both the first receptor and the second receptor. In some embodiments, at least 90% of the immune cells express both the first receptor and the second receptor. In some embodiments, the pharmaceutical composition further comprises a pharmaceutically acceptable carrier, diluent or excipient. In some embodiments of the present invention, the pharmaceutical composition according to the present invention is for use in the treatment of EGFR+ cancer.

[0044] Further described herein is a polynucleotide or polynucleotide system, comprising one or more polynucleotides comprising polynucleotide sequences encoding: a.) a first receptor, comprising an extracellular ligand binding domain specific to Endothelial Growth Factor Receptor (EGFR); and b.) a second receptor, comprising an extracellular ligand binding domain specific to a non-target antigen that has been lost in the EGFR+ cancer cell due to loss of heterozygosity, wherein the first receptor is an activator receptor responsive to EGFR on the EGFR+ cancer cell; and wherein the second receptor is an inhibitory receptor responsive to the non-target antigen.

[0045] In one aspect, the present invention provides a polynucleotide comprising polynucleotide sequences encoding the first receptor and the second receptor for generating the immune cells according to the present invention.

[0046] In some embodiments of the polynucleotide, the polynucleotide comprises a sequence encoding a shRNA specific to B2M. In some embodiments, the sequences encoding the first receptor, the second receptor, and the shRNA specific to B2M are encoded by the same polynucleotide.

[0047] In one aspect, the present invention provides a vector, comprising polynucleotide according to the present invention.

[0048] In one aspect, the present invention further provides the immune cell according to the present invention for use in the treatment of EGFR+ cancer. In an embodiment, the EGFR+ cancer is an EGFR+ tumor having loss of heterozygosity at a locus encoding a non-target antigen. In one embodiment, the EGFR+ cancer comprises EGFR+ cancer cells having loss of heterozygosity at an MHC class I locus.

[0049] In some embodiments, the EGFR+ cancer cell is a lung cancer cell, a small cell lung cancer cell, a non-small cell lung cancer cell, a pancreatic ductal carcinoma cell, a colorectal cancer cell, a head and neck cancer cell, a esophagus

and gastric adenocarcinoma cell, an ovarian cancer cell, a glioblastoma multiforme cell, a cervical squamous cell carcinoma cell, a kidney cancer cell, a papillary kidney cancer cell, a kidney renal clear cell carcinoma cell, a bladder cancer cell, a breast cancer cell, a bile duct cancer cell, a liver cancer cell, a prostate cancer cell, a sarcoma cell, a thyroid cancer cell, a thymus cancer cell, a stomach cancer cell, or a uterine cancer cell. In some embodiments, the EGFR+ cancer cell is a lung cancer cell.

[0050] In some embodiments of the present invention, the EGFR+ cancer cell is an EGFR+/HLA-A*02- cancer cell that does not express HLA-A*02; or a cancer cell derived from an HLA-A*02+ individual, which does not express HLA-A*02. In some embodiments of the present invention, the EGFR+/HLA-A*02- cancer cell is derived from an EGFR+/HLA-A*02+ cell by loss of heterozygosity at HLA-A leading to loss of HLA-A*02.

[0051] In one aspect, the present invention provides an immune cell according to the present invention for use in the treatment of EGFR+ cancer. In one embodiment, the EGFR+ cancer is an EGFR+ tumor having loss of heterozygosity at a locus encoding a non-target antigen. In one embodiment, the EGFR+ cancer comprises EGFR+ cancer cells having loss of heterozygosity at an MHC class I locus.

[0052] It is described herein that the subject is a heterozygous HLA-A*02 patient with a malignancy that expresses EGFR and has lost HLA-A*02 expression.

[0053] It is described herein that the subject is a heterozygous HLA-A*02 patient with recurrent unresectable or metastatic solid tumors that express EGFR and have lost HLA-A*02 expression.

[0054] Described herein is a method of treating a cancer in a subject comprising: a.) determining the genotype or expression level of a non-target antigen in non-malignant cells and cancer cells of the subject; b.) determining the expression level of EGFR in cancer cells of the subject; and c.) administering to the subject an effective amount of the immune cell described herein or the pharmaceutical composition described herein if the non-malignant cells express the non-target antigen and the cancer cells do not express the non-target antigen, and the cancer cells are EGFR positive.

[0055] It is described herein that administration of the immune cell described herein or the pharmaceutical composition described herein reduces the size of a tumor in the subject.

[0056] It is described herein that the tumor is reduced by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, or about 100%. It is described herein that the tumor is eliminated.

[0057] It is described herein that administration of the immune cell or the pharmaceutical composition arrests the growth of a tumor in the subject. It is described herein that administration of the immune cell described herein or the pharmaceutical composition described herein reduces the number of tumors in the subject.

[0058] It is described herein that administration of the immune cell or the pharmaceutical composition results in selective killing of a cancer cell, but not a normal cell in the subject. It is described herein that at least about 60% of the cells killed are cancer cells, at least about 65% of the cells killed are cancer cells, at least about 70% of the cells killed are cancer cells, at least about 75% of the cells killed are cancer cells, at least about 80% of the cells killed are cancer cells, at least about 85% of the cells killed are cancer cells, at least about 90% of the cells killed are cancer cells, at least about 95% of the cells killed are cancer cells, or about 100% of the cells killed are cancer cells.

[0059] It is described herein that administration of the immune cell or pharmaceutical composition results in the killing of about 40%, about 50%, about 60%, about 70%, about 80%, about 90% or all of the cancer cells of the subject. In some embodiments, the cancer cell comprises a lung cancer cell, a small cell lung cancer cell, a non-small cell lung cancer cell, a pancreatic ductal carcinoma cell, a colorectal cancer cell, a head and neck cancer cell, a esophagus and gastric adenocarcinoma cell, an ovarian cancer cell, a glioblastoma multiforme cell, a cervical squamous cell carcinoma cell, a kidney cancer cell, a papillary kidney cancer cell, a kidney renal clear cell carcinoma cell, a bladder cancer cell, a breast cancer cell, a bile duct cancer cell, a liver cancer cell, a prostate cancer cell, a sarcoma cell, a thyroid cancer cell, a thymus cancer cell, a stomach cancer cell, or a uterine cancer cell. It is described herein that administration of the immune cell or the pharmaceutical composition results in fewer side effects for the subject than administration of an otherwise equivalent immune cell comprising the first activator receptor, but no second inhibitory receptor.

[0060] Further described herein is a method of making a plurality of immune cells, comprising: a.) providing a plurality of immune cells, and b.) transforming the plurality of immune cells with the polynucleotide system described herein, or the vector described herein.

[0061] In one aspect, the disclosure provides a kit comprising the immune cell described herein or the pharmaceutical composition described herein. As described herein, the kit further comprises instructions for use.

The present invention is defined in the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0062] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

FIG. 1A is a plot showing that Jurkat cells expressing an EGFR CAR activator and an HLA-A*02 LIR-1 blocker are activated by EGFR+/HLA-A*02-HeLa target cells, but not EGFR+/HLA-A*02+-HeLa target cells.

FIG. 1B is a pair of fluorescence activated cell sorting (FACS) plots showing expression of activator (CT479) and blocker (C1765) receptors on Jurkat cells.

FIG. 1C is a pair of plots showing expression of activator antigen (EGFR) and blocker antigen (HLA-A*02) on HeLa cells. HLA-A*02 expression was detected using BB7.2 antibody.

FIG. 2A shows the expression of HLA-A*02 on HeLa cells transduced with HLA-A*02, and HCT116 cells. HeLa and HCT116 cells were labeled with the anti-HLA-A2 antibody BB7.2 and FACS sorted. Green: unlabeled HeLa; orange: unlabeled HCT116; blue: wild type HCT116 labeled with BB7.2; red: HeLa cells transduced with HLA-A*02 and labeled with BB7.2.

FIG. 2B shows expression of EGFR on HeLa cells and HCT116 cells. HeLa and HCT116 cells were labeled with anti-EGFR antibody. Green: unlabeled HeLa; orange: unlabeled HCT116; blue: wild type HCT116 labeled with anti-EGFR; red: HeLa cells transduced with HLA-A*02 and labeled with anti-EGFR.

FIG. 3A shows EGFR CAR activation of Jurkat cells expressing an EGFR CAR, when co-cultured with HCT116 target cells.

FIG. 3B shows that EGFR CAR activation of Jurkat cells can be blocked by an HLA-A*02 LIR-1 inhibitory receptor. Co-expression of the EGFR CAR and HLA-A*02 LIR-1 inhibitory receptor by Jurkat cells leads to a shift in the $CARE_{MAX}$ of approximately 1.8x when Jurkat cells are presented with HCT116 target cells expressing EGFR and HLA-A*02.

FIG. 3C shows EGFR CAR activation of Jurkat cells expressing an EGFR CAR, when co-cultured with HeLa target cells.

FIG. 3D shows that EGFR CAR activation of Jurkat cells can be blocked by an HLA-A*02 LIR-1 inhibitory receptor. Co-expression of the EGFR CAR and HLA-A*02 LIR-1 inhibitory receptor by Jurkat cells leads to a shift in the $CARE_{MAX}$ of approximately 5.4x when Jurkat cells are presented with HeLa target cells expressing EGFR and HLA-A*02.

FIG. 4A shows titration of activator antigen in a bead-based assay to determine the optimal amount of activator antigen.

FIG. 4B shows titration of blocker antigen in a bead based assay to determine the optimal ratio of activator to blocker antigen.

FIG. 5 is a series of plots and a table that show that a different degree of blocking is observed when an HLA-A*02 scFv LIR1 inhibitor is used with different EGFR scFv CAR activators in primary T cells.

FIG. 6A is a series of plots that show that a different degree of blocking is observed when two different HLA-A*02 scFv LIR1 inhibitors are used with different EGFR scFv CAR activators in primary T cells.

FIG. 6B is a series of plots that show the sensitivity of activating and inhibitory receptors when different pairs of activating and inhibitory receptors are co-expressed. Amount of activator or non-target antigen are varied on the target cells by mRNA titration.

FIG. 6C is a series of charts showing the maximum specific killing (%) of target cells by cells expressing an EGFR targeting activator receptor and either a humanized inhibitory receptor or a murine inhibitory receptor. The selectivity ratio of target tumor cells vs. normal cells is also shown for each pair of activator and inhibitory receptors.

FIG. 7 is a series of plots showing cytotoxicity of T cells expressing either EGFR scFv CAR, or EGFR scFv CAR and HLA-A*02 scFv LIR1 inhibitory receptors that were co-cultured with HeLa cells at a 1:1 ratio. A Tumor Cells: HeLa cells expressing EGFR and not HLA-A*02; AB Normal cells: HeLa cells expression EGFR and HLA-A*02; A- B- normal cells: HeLa cells that did not express either EGFR or HLA-A*02; B normal cells: HeLa cells expressing HLA-A*02, but not EGFR. Y axis shows green area, which is proportional to healthy HeLa cells in the assay. UTD: untransduced.

FIG. 8 is a series of plots showing the enrichment of primary T cells expressing blocker and activator receptors (top row), and the cytotoxic activity of these T cells expressing EGFR scFv CAR, or EGFR scFv CAR and HLA-A*02 scFv LIR1 inhibitory receptor, were mixed 1: 1 with HeLa cells expressing either EGFR (Target A), or EGFR and HLA-A*02 (Target AB).

FIG. 9 is a series of plots showing cytotoxicity of T cells expressing either EGFR scFv CAR, or EGFR scFv CAR and HLA-A*02 scFv LIR1 inhibitory receptor, which were mixed 1: 1 with HeLa cells expressing either EGFR, HLA-A*02, or both. A Tumor Cells: HeLa cells expressing EGFR and not HLA-A*02; AB Normal cells: HeLa cells expression EGFR and HLA-A*02; B normal cells: HeLa cells expressing HLA-A*02, but not EGFR. Y axis shows green area, which is proportional to healthy HeLa cells in the assay. UTD: untransduced.

FIG. 10 is a series of plots showing cytotoxicity of T cells expressing either EGFR scFv CAR, or EGFR scFv CAR and HLA-A*02 scFv LIR1 inhibitory receptor, which were mixed 1: 1 with HCT.116 cells expressing either EGFR (Target A) or EGFR and HLA-A*02 (Target AB).

FIG. 11A is a series of fluorescence activated cell sorting (FACS) plots showing expression of EGFR scFv CAR activator receptor by T cells following incubation of T cells expressing different EGFR scFv CAR and an HLA-A*02 scFv LIR1 inhibitor with HeLa cells expressing EGFR activator alone (Target A), inhibitor target alone (Target B) or

activator and inhibitor targets (Target AB).

FIG. 11B is a plot showing quantification activator receptor expression before exposure to target cells, and after 120 hours co-culture with target cells expressing activator ligand alone (Target A), or target cells expressing both activator and blocker ligands (Target AB).

FIG. 12A is a plot showing cell surface expression of the activator receptor on T cells expressing an EGFR scFv CAR (CT-482) activator and HLA-A*02 scFv LIR1 inhibitor (C1765) following co-culture with populations of HeLa cells expressing EGFR (Target A), HLA-A*02 (Target B), a combination of EGFR and HLA-A*02 on the same cell (Target AB), a mixed population of HeLa cells expressing Target A and Target AB on different cells, or a mixed population of HeLa cells expressing Target B and Target AB on different cells.

FIG. 12B is a plot showing cell surface expression of the inhibitory receptor on T cells expressing an EGFR scFv CAR (CT-482) activator and HLA-A*02 scFv LIR1 inhibitor (C1765) following co-culture with populations of HeLa cells expressing EGFR (Target A), HLA-A*02 (Target B), a combination of EGFR and HLA-A*02 on the same cell (Target AB), a mixed population of HeLa cells expressing Target A and Target AB on different cells, or a mixed population of HeLa cells expressing Target B and Target AB on different cells.

FIG. 13 is a diagram of an experiment to determine if cytotoxicity or blocking activity by T cells was reversible.

FIG. 14A is a series of plots showing that cytotoxicity and blocking activity is reversible and correspond to activator surface expression. The T cells were co-cultured with EGFR+/HLA-A2+ target cells in round 1, then switched to alternating target cells in the following rounds. At top: percent killing of target HeLa cells by T cells is shown. At bottom: activator and inhibitory receptor expression as assayed by FACS.

FIG. 14B is a series of plots showing that cytotoxicity and blocking activity is reversible and correspond to activator surface expression. The T cells were co-cultured with EGFR+/HLA-A2-target cells in round 1, then switched to alternating target cells in the following rounds. At top: percent killing of target HeLa cells by T cells is shown. At bottom: activator and inhibitory receptor expression as assayed by FACS.

FIG. 15A is a plot showing that expression of EGFR activator CAR on the cell surface is reversibly downregulated by the presence of target cells expressing both activator and inhibitor ligands.

FIG. 15B is a series of plots showing that EGFR activator CAR expression on the cell surface is quickly restored when activator only target cells are present, or target cells expressing activator and blocker targets are not present.

FIG. 16 shows that activation of Jurkat cells expressing an EGFR scFv CAR using a bead based assay can be blocked by a pMHC HLA-A*02 scFv LIR-1 based inhibitory receptor when the activator and inhibitor antigens are present on beads in *cis*, but not when the activator and inhibitor antigens are present on the beads in *trans*.

FIG. 17A shows that activation of Jurkat cells by an EGFR scFv CAR can be blocked by a pMHC HLA-A*02 scFv LIR-1 based inhibitory receptor using SiHa target cells expressing HLA-A*02 (SiHa A02), but not by SiHa cells that do not express HLA-A*02 (SiHa WT).

FIG. 17B shows that activation of Jurkat cells by an EGFR scFv CAR can be blocked by a pMHC HLA-A*02 scFv LIR-1 based inhibitory receptor using HeLa target cells expressing HLA-A*02 (HeLa A02), but not by HeLa cells that do not express HLA-A*02 (HeLa WT).

FIG. 18 is a diagram showing the bioinformatics search process used to identify potential non-target antigen candidate genes.

FIG. 19 is a diagram showing the experimental design for determining efficacy and selectivity of immune cells expressing an EGFR targeting activator receptor and an inhibitory receptor *in vivo*.

FIG. 20A is a plot showing Jurkat NFAT Luciferase (JNL) cell inhibition in an mRNA titration assay using HeLa cells transfected with mRNA encoding HLA-A*11. HeLa cells were transfected with serially diluted HLA-A*11 mRNA, and JNL cells were transiently transfected with an EGFR activating CAR +/- HLA-A*11 inhibitory receptor with scFv HLA-A*11 #4. The functional response was assessed after a 6 hour co-culture.

FIG. 20B shows the molecule/cell sensitivity (IC₅₀) of the inhibitory receptor with HLA-A*11 #4.

FIG. 21 is a plot showing that an HLA-B*07-scFv-LIR1 inhibitory receptor inhibits immune cell activation mediated by a co-expressed EGFR targeting activator receptor.

FIG. 22 is a series of plots showing the expression of EGFR activator receptors (EGFR-CAR only) and EGFR activator/HLA-A inhibitory receptor pairs in primary T cells comprising vectors disclosed herein. Percent of cells expressing one or both receptors are shown before enrichment (pre-enrichment) and after enrichment (post-enrichment), where the enrichment selects for cells expressing the receptors. Receptor pairs are expressed from either two separate vectors (dual vector) or one vector (single vector).

FIG. 23A is a series of plots showing the percentage of specific killing of HeLa target cells or HCT116 target cells by immune cells expressing an EGFR activator receptor and an HLA-A*02 scFv LIR1 inhibitor receptor. The target cells express the EGFR antigen (A), or both the EGFR antigen and the HLA-A*02 non-target antigen (B). Immune cells expressing the activator/inhibitor receptor pair were incubated with target cells at a 1:1 ratio.

FIG. 23B is a plot and series of images showing specific and selective killing of target cells by immune cells expressing an EGFR activator receptor and an HLA-A*02 scFv LIR1 inhibitor receptor in a mixed culture of cancer cells and

normal cells. UTD = immune cells no activator or inhibitor receptors; CAR = immune cells expressing only activator receptor; Tmod = immune cells expressing both activator and inhibitor receptors. In the images on the right: green = normal cells; red = tumor cells.

FIG. 24 is a chart showing selective cell activation in Jurkat cells (effectors) expressing EGFR activator receptor (CAR) or EGFR activator receptor and HLA-A*02 inhibitor receptor (Tmod). The Jurkat cells were incubated with a panel of target cell lines that express HLA-A*02 with or without EGFR expression. Tmod Jurkat effector cells showed no off-target activation when co-cultured with target cells that do not express EGFR or cells that express HLA-A*02.

DETAILED DESCRIPTION

[0063] Described herein are compositions and methods for treating cancers using immune cells comprising a two receptor system responsive to differences in gene expression of a ligand between cancer and normal cells. These differences in expression can be due to loss of heterozygosity in the cancer cells. Alternatively, the differences in expression can be, because the gene expression is not expressed in cancer cells, or is expressed in cancer cells at a lower level than normal cells. The two-receptor system is expressed in immune cells, for example immune cells used in adoptive cell therapy, and targets activity of these immune cells to cancer cells exhibiting loss of heterozygosity or expression differences. In this two receptor system, the first receptor (an activator receptor, sometimes referred to herein as an A module) activates, or promote activation of the immune cells, while the second receptor (an inhibitory receptor, sometimes referred to herein as a blocker, or inhibitor receptor, or a B module) acts to inhibit activation of the immune cells by the first receptor. Each receptor contains a ligand-binding domain (LBD) that binds a specific ligand. Signals from the two receptors upon ligand binding are integrated by the immune cell. Differential expression of ligands for the first and second receptors in cancer and normal cells, for example through loss of heterozygosity of the locus encoding the inhibitory ligand in cancer cells, or differences in transcription levels, mediates activation of immune cells by target cancer cells that express the first activator ligand, but not the second inhibitory ligand.

[0064] In particular embodiments of the compositions and methods described herein, immune cells comprising the two receptor system described herein are used to treat epidermal growth factor receptor (EGFR) positive cancers. This includes lung cancers, glioblastoma, breast cancers, head and neck cancer and colorectal cancers. In the case of EGFR-positive cancers, the target antigen of the activator receptor is EGFR, or a peptide antigen thereof, in a complex with a major histocompatibility complex class I (MHC-I). EGFR is expressed in a wide variety of normal tissues, such as epithelial, mesenchymal and neuronal tissues, and plays a major role in normal cellular processes such as cellular proliferation and differentiation, as well as development. Because of its expression in certain tumors, EGFR is an attractive tumor-specific antigen that could mediate selective killing of EGFR+ tumors if these cancer cells could be specifically targeted with an appropriate therapeutic. However, normal EGFR expression in non-cancer (non-target) cells has prevented the effective use of EGFR for targeted therapies such as adoptive cell therapies. Skin and gastrointestinal toxicities have prevented effective targeting of EGFR in EGFR positive cancers. By pairing an EGFR activator receptor with an inhibitory receptor, the methods provided herein increase the specificity of EGFR targeted adoptive cell therapies and decrease harmful effects associated with these therapies, such as dose-limited toxicity.

[0065] In some embodiments, the ligand for the activator is an EGFR peptide complexed with MHC class I. In the methods described herein, this EGFR targeted activator receptor is paired with an inhibitory receptor, which increases the safety window of the activator by blocking its cytolytic effect on normal EGFR-positive tissues. However, the activator receptor still directs the targeted killing of tumor cells by immune cells comprising the two-receptor system, as the tumor cells do not express the ligand for the inhibitor, or blocker, receptor. The target for the second, inhibitory receptor is expressed by EGFR positive tissues such as epithelial tissues, but is not expressed in cancer cells, and the inhibitory receptor recognizes this "non-target antigen" as an inhibitory stimulus. An exemplary target for the second inhibitory receptor is expressed by lung epithelial tissue, and is lost from EGFR positive cancer cells due to heterozygosity (LOH), leaving a single allelic form in cancer cells that can be distinguished from other alleles via an allele-specific ligand binding domain on the inhibitory receptor. Exemplary targets of the inhibitory receptor include Major Histocompatibility Complex (MHC) proteins such as human leukocyte antigen A (HLA-A), HLA-B, HLA-C, and other HLAs. HLAs are encoding by variant genes, such as HLA-A*01, HLA-A*02, HLA-A*03, HLA-C*07, and others, which can be lost from EGFR positive cancer cells through loss of heterozygosity. Alternatively, further exemplary targets of the inhibitory receptor include collectin subfamily member 12 COLEC12, APC downregulated 1 (APCDD1) and C-X-C motif chemokine ligand 16 (CXCL16). Each of these has a common nonsynonymous variant form, with the amino-acid alteration in its extracellular domain accessible to antibodies, which can be used as a B module target for a cellular integrator designed to safely treat patients with EGFR positive cancers with engineered T cells activated by an activator receptor such as a EGFR or EGFR pMHC responsive activator receptor. The compositions and methods of the disclosure can reduce or eliminate DLT caused by expression of EGFR by normal tissue. The disclosure provides methods of targeting EGFR in cancer cells to treat EGFR positive cancers using adoptive cell therapies by adding a second inhibitory receptor that blocks activation of the adoptive immune cells in the presence of a second ligand (a ligand other than EGFR, termed a "non-

target antigen"). Using the compositions and methods described herein, tumor cells that express EGFR are attacked by the adoptive immune cells expressing the two receptors, because these tumor cells express only the activator ligand, EGFR. In contrast, normal cells that express EGFR plus the non-target antigen (alternatively termed a "blocker antigen") are protected from the adoptive immune cells. The inhibitory receptor response to the non-target antigen on normal cells and thereby prevents activation of immune cells by the EGFR-targeted activator receptor. This dual-targeting approach creates the therapeutic window that will allow an EGFR-directed cell therapy to be dosed safely and effectively in EGFR-positive cancer patients.

[0066] The disclosure provides methods and compositions that allow the use of potent EGFR CAR and TCRs that induce on-target toxicity, and renders these EGFR targeted receptors useful as a therapeutic by mitigating their toxicity.

[0067] In variations, the compositions and methods described herein may be used to kill target cells and/or treat subjects, in which expression of the non-target antigen is partially or completely decreased by causes other than loss of heterozygosity, including partial gene deletion, epigenetic silence, point mutations, truncations.

Definitions

[0068] Prior to setting forth this disclosure in more detail, it may be helpful to an understanding thereof to provide definitions of certain terms to be used herein.

[0069] As used herein, the term "about" or "approximately" refers to a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2% or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length. As described herein, the term "about" or "approximately" refers a range of quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length $\pm 15\%$, $\pm 10\%$, $\pm 9\%$, $\pm 8\%$, $\pm 7\%$, $\pm 6\%$, $\pm 5\%$, $\pm 4\%$, $\pm 3\%$, $\pm 2\%$, or $\pm 1\%$ about a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

[0070] As used herein, the term "isolated" means material that is substantially or essentially free from components that normally accompany it in its native state. As used herein, the term "obtained" or "derived" is used synonymously with isolated.

[0071] The terms "subject," "patient" and "individual" are used interchangeably herein to refer to a vertebrate, preferably a mammal, more preferably a human. Tissues, cells, and their progeny of a biological entity obtained in vivo or cultured in vitro are also encompassed. A "subject," "patient" or "individual" as used herein, includes any animal that exhibits pain that can be treated with the vectors, compositions, and methods contemplated herein. Suitable subjects (e.g., patients) include laboratory animals (such as mouse, rat, rabbit, or guinea pig), farm animals, and domestic animals or pets (such as a cat or dog). Non-human primates and, preferably, human patients, are included.

[0072] As used herein "treatment" or "treating," includes any beneficial or desirable effect, and may include even minimal improvement in symptoms. "Treatment" does not necessarily indicate complete eradication or cure of the disease or condition, or associated symptoms thereof.

[0073] As used herein, "prevent," and similar words such as "prevented," "preventing" etc., indicate an approach for preventing, inhibiting, or reducing the likelihood of a symptom of disease. It also refers to delaying the onset or recurrence of a disease or condition or delaying the occurrence or recurrence of the symptoms of a disease. As used herein, "prevention" and similar words also includes reducing the intensity, effect, symptoms and/or burden of disease prior to onset or recurrence.

[0074] As used herein, the term "amount" refers to "an amount effective" or "an effective amount" of a virus to achieve a beneficial or desired prophylactic or therapeutic result, including clinical results.

[0075] A "therapeutically effective amount" of a virus or cell may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the virus or cell to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the virus or cell are outweighed by the therapeutically beneficial effects. The term "therapeutically effective amount" includes an amount that is effective to "treat" a subject (e.g., a patient).

[0076] An "increased" or "enhanced" amount of a physiological response, e.g., electrophysiological activity or cellular activity, is typically a "statistically significant" amount, and may include an increase that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1, e.g., 1.5, 1.6, 1.7, 1.8, etc.) the level of activity in an untreated cell.

[0077] A "decreased" or "reduced" amount of a physiological response, e.g., electrophysiological activity or cellular activity, is typically a "statistically significant" amount, and may include an decrease that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1, e.g., 1.5, 1.6, 1.7, 1.8, etc.) the level of activity in an untreated cell.

[0078] By "maintain" or "preserve" or "maintenance" or "no change" or "no substantial change" or "no substantial decrease" refers generally to a physiological response that is comparable to a response caused by either vehicle, or a

control molecule/composition. A comparable response is one that is not significantly different or measurable different from the reference response.

[0079] In general, "sequence identity" or "sequence homology" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Typically, techniques for determining sequence identity include determining the nucleotide sequence of a polynucleotide and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. Two or more sequences (polynucleotide or amino acid) can be compared by determining their "percent identity". The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences divided by the length of the shorter sequences and multiplied by 100. Percent identity may also be determined, for example, by comparing sequence information using the advanced BLAST computer program, including version 2.2.9, available from the National Institutes of Health. The BLAST program is based on the alignment method of Karlin and Altschul, *Proc. Natl. Acad. Sci. USA* 87:2264-2268 (1990) and as discussed in Altschul, et al., *J. Mol. Biol.* 215:403-410 (1990); Karlin and Altschul, *Proc. Natl. Acad. Sci. USA* 90:5873-5877 (1993); and Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997). Briefly, the BLAST program defines identity as the number of identical aligned symbols (generally nucleotides or amino acids), divided by the total number of symbols in the shorter of the two sequences. The program may be used to determine percent identity over the entire length of the proteins being compared. Default parameters are provided to optimize searches with short query sequences in, for example, with the blastp program. The program also allows use of an SEG filter to mask-off segments of the query sequences as determined by the SEG program of Wootton and Federhen, *Computers and Chemistry* 17:149-163 (1993). Ranges of desired degrees of sequence identity are approximately 80% to 100% and integer values therebetween. Typically, the percent identities between a disclosed sequence and a claimed sequence are at least 80%, at least 85%, at least 90%, at least 95%, or at least 98%.

[0080] As used herein, a "polynucleotide system" refers to one or more polynucleotides. The one or more polynucleotides may be designed to work in concert for a particular application, or to produce a desired transformed cell.

[0081] The term "exogenous" is used herein to refer to any molecule, including nucleic acids, protein or peptides, small molecular compounds, and the like that originate from outside the organism. In contrast, the term "endogenous" refers to any molecule that originates from inside the organism (i.e., naturally produced by the organism).

[0082] The term "MOI" is used herein to refer to multiplicity of infection, which is the ratio of agents (e.g. viral particles) to infection targets (e.g. cells).

[0083] In the present description, any concentration range, percentage range, ratio range, or integer range is to be understood to include the value of any integer within the recited range and, when appropriate, fractions thereof (such as one tenth and one hundredth of an integer), unless otherwise indicated.

[0084] As used herein, a "target cell" refers to cell that is targeted by an adoptive cell therapy. For example, a target cell can be a cancer cell, which can be killed by the transplanted T cells of the adoptive cell therapy. Target cells of the disclosure express a target antigen, as described herein, and do not express a non-target antigen.

[0085] As used herein, a "non-target cell" refers to a cell that is not targeted by an adoptive cell therapy. For example, in an adoptive cell targeting cancer cells, normal, healthy, non-cancerous cells are non-target cells. Some, or all, non-target cells in a subject may express both the target antigen and the non-target antigen. Non-target cells in a subject may express the non-target antigen irrespective of whether or not these cells also express the target antigen.

[0086] As used herein, a "target antigen," whether referred to using the term antigen or the name of a specific antigen, refers to an antigen expressed by a target cell, such as a cancer cell. Expression of target antigen is not limited to target cells. Target antigens may be expressed by both cancer cells and normal, non-cancer cells in a subject.

[0087] As used herein, a "non-target antigen" (or "blocker antigen") whether referred to using the term antigen or the name of a specific antigen, refers to an antigen that is expressed by normal, non-cancer cells and is not expressed in cancer cells. This difference in expression allows the inhibitory receptor to inhibit immune cell activation in the presence of non-target cells, but not in the presence of target cells.

[0088] Polymorphism refers to the presence of two or more variants of a nucleotide sequence in a population. A polymorphism may comprise one or more base changes, an insertion, a repeat, or a deletion. A polymorphism includes e.g. a simple sequence repeat (SSR) and a single nucleotide polymorphism (SNP), which is a variation, occurring when a single nucleotide of adenine (A), thymine (T), cytosine (C) or guanine (G) is altered.

[0089] As used herein, "affinity" refers to strength of binding of a ligand to a single ligand binding site on a receptor, for example an antigen for the antigen binding domain of any of the receptors described herein. Ligand binding domains can have a weaker interaction (low affinity) with their ligand, or a stronger interaction (high affinity).

[0090] K_d, or dissociation constant, is a type of equilibrium constant that measures the propensity of a larger object to separate reversibly into smaller components, such as, for example, when a macromolecular complex comprising receptor and its cognate ligand separates into the ligand and the receptor. When the K_d is high, it means that a high concentration of ligand is needed to occupy the receptor, and the affinity of the receptor for the ligand is low. Conversely, a low K_d means that the ligand has a high affinity for the receptor.

[0091] As used herein, a receptor that is "responsive" or "responsive to" refers to a receptor comprising an intracellular domain, that when bound by a ligand (*i.e.* antigen) generates a signal corresponding to the known function of the intracellular domain. An activator receptor bound to a target antigen can generate a signal that causes activation of an immune cell expressing the activator receptor. An inhibitory receptor bound to a non-target antigen can generate an inhibitory signal that prevents or reduces activation of an immune cell expressing the activator receptor. Responsiveness of receptors, and their ability to activate or inhibit immune cells expressing the receptors, can be assayed by any means known in the art and described herein, including reporter assays and cytotoxicity assays.

[0092] As used herein, "activation" of an immune cell or an immune cell that is "activated" is an immune cell that can carry out one or more functions characteristic of an immune response. These functions include proliferation, release of cytokines, and cytotoxicity, *i.e.* killing of a target cell. Activated immune cells express markers that will be apparent to persons of skill in the art. For example, activated T cells can express one or more of CD69, CD71, CD25 and HLA-DR. An immune cell expressing an activator receptor (*e.g.* an EGFR CAR) can be activated by the activator receptor when it becomes responsive to the binding of a target antigen (*e.g.* EGFR). A "target antigen" can also be referred to as an "activator antigen" and may be isolated or expressed by a target cell. Activation of an immune cell expressing an inhibitory receptor can be reduced or prevented when the inhibitory receptor becomes responsive to the binding of a non-target antigen (*e.g.* HLA-A*02), even when the activator receptor is bound to the target activator ligand. A "non-target antigen" can also be referred to as an "inhibitory ligand" or a "blocker", and may be isolated or expressed by a target cell.

[0093] Receptor expression on an immune cell can be verified by assays that report the presence of the activator receptors and inhibitory receptors described herein. For example, a population of immune cells can be stained with a labeled molecule (*e.g.* a fluorophore labeled receptor-specific antibody or a fluorophore-labeled receptor-specific ligand), and quantified using fluorescence activated cell sorting (FACS) flow cytometry. This method allows a percentage of immune cells in a population of immune cells to be characterized as expressing an activator receptor, an inhibitory receptor, or both receptors. The ratio of activator receptor and inhibitory receptors expressed by the immune cells described herein can be determined by, for example, digital droplet PCR. These approaches can be used to characterize the population of cells for the production and manufacturing of the immune cells, pharmaceutical compositions, and kits described herein. For the immune cells, pharmaceutical compositions, and kits described herein, it is understood that a suitable percentage of immune cells expressing both an activator receptor and an inhibitory receptor is determined specifically for the methods described herein. For example, a suitable percentage of immune cells expressing both an activator receptor and an inhibitory receptor can be at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95%. For example, a suitable percentage of immune cells expressing both an activator receptor and an inhibitory receptor can be at most 50%, at most 55%, at most 60%, at most 65%, at most 70%, at most 75%, at most 80%, at most 85%, at most 90%, or at most 95%. For example, a suitable ratio of activator receptor and inhibitory receptor in an immune cell can be about 5:1, about 4:1, about 3:1, about 2:1, about 1:1, about 1:2, about 1:3, about 1:4, or about 1:5. It is understood that purification, enrichment, and/or depletion steps can be used on populations of immune cells to meet suitable values for the immune cells, pharmaceutical compositions, and kits described herein.

[0094] A responsive receptor expressed by the immune cells described herein can be verified by assays that measure the generation of a signal expected to be generated by the intracellular domain of the receptor. Reporter cell lines, such as Jurkat-Luciferase NFAT cells (Jurkat cells), can be used to characterize a responsive receptor. Jurkat cells are derived from T cells and comprise a stably integrated nuclear factor of activated T-cells (NFAT)-inducible luciferase reporter system. NFAT is a family of transcription factors required for immune cell activation, whose activation can be used as a signaling marker for T cell activation. Jurkat cells can be transduced or transfected with the activator receptors and/or inhibitory receptors described herein. The activator receptor is responsive to the binding of a ligand if the Jurkat cell expresses a luciferase reporter gene, and the level of responsiveness can be determined by the level of reporter gene expression. The presence of luciferase can be determined using any known luciferase detection reagent, such as luciferin. An inhibitory receptor is responsive to the binding of a ligand if, when co-expressed with an activator receptor in Jurkat cells, it prevents a normally responsive immune cell from expressing luciferase in response to the activator receptor. For example, the responsiveness of an inhibitory receptor can be determined and quantified in a Jurkat cell expressing both an activator and an inhibitor by observing the following: 1) the Jurkat cell expresses luciferase in the presence of activator receptor ligand and absence of inhibitory receptor ligand; and 2) luciferase expression in the Jurkat cell is reduced or eliminated in the presence of both an activator receptor ligand and an inhibitory receptor ligand. This approach can be used to determine the sensitivity, potency, and selectivity of activator receptors and specific pairs of activator receptors and inhibitory receptors. The sensitivity, potency, and selectivity can be quantified by EC50 or IC50 values using dose-response experiments, where an activator receptor ligand and/or inhibitory receptor ligand is titrated into a culture of Jurkat cells expressing an activator receptor or a specific pair of activator and inhibitory receptors. Alternatively, the EC50 and IC50 values can be determined in a co-culture of immune cells (*e.g.* Jurkat cells or primary immune cells) expressing an activator receptor or a specific pair of activator and inhibitory receptors and target cells expressing an increasing amount of an activator ligand or inhibitor ligand. An increasing amount of activator ligand or inhibitor ligand

can be accomplished in the target cell by, for example, titration of activator ligand or inhibitor ligand encoding mRNA into target cells, or use of target cells that naturally express different levels of the target ligands. Exemplary suitable EC50 and IC50 values for the activator and inhibitory receptors as determined used target cells expressing varying amounts of the target and non-target ligands include an EC50 of 260 transcripts per million (TPM) or less for the activator receptor, for example an EC50 of between 10 and 260 TPM, and an IC50 of 10 TPM or less for the inhibitory receptor, for example an IC50 of 1-5 TPM.

[0095] Activation of the immune cells described herein that express an activator receptor or specific pairs of activator and inhibitory receptors can be further determined by assays that measure the viability of a target cell following co-incubation with said immune cells. The immune cells, sometimes referred to as effector cells, are co-incubated with target cells that express an activator receptor ligand, an inhibitory receptor ligand, or both an activator and inhibitory receptor ligand. Following co-incubation, viability of the target cell is measured using any method to measure viability in a cell culture. For example, viability can be determined using a mitochondrial function assay that uses a tetrazolium salt substrate to measure active mitochondrial enzymes. Viability can also be determined using imaging based methods. Target cells can express a fluorescent protein, such as green fluorescent protein or red fluorescent protein. Reduction in total cell fluorescence indicates a reduction in viability of the target cell. A reduction in viability of the target cell following incubation with immune cells expressing an activator receptor or a specific pair of activator and inhibitory receptors is interpreted as target cell-mediated activation of the immune cell. A measure of the selectivity of the immune cells can also be determined using this approach. The immune cell expressing a pair of activator and inhibitory receptors is selective if the following is observed: 1) viability is reduced in target cells expressing the activator receptor ligand, but not the inhibitory receptor ligand; 2) viability is not reduced in target cells expressing both an activator receptor ligand and an inhibitory receptor ligand. From these measurements, a "specific killing" value can be derived that quantifies the percentage of immune cell activation based on the reduction in viability of target cell as a percentage of a negative control (immune cells that do not express an activator receptor). Further, from these measurements a "selectivity ratio" value can be derived that represents the ratio of the specific killing observed in target cells expressing an activator receptor ligand in the absence of inhibitory receptor ligand to the specific killing observed in target cells expressing both an activator receptor ligand and an inhibitory receptor ligand. This approach can be used to characterize the population of cells for the production and manufacturing of the immune cells, pharmaceutical compositions, and kits described herein. A suitable specific killing value for the immune cells, pharmaceutical compositions, and kits can be, for example, the following criteria: 1) at least 50% specific killing following a 48 hour co-incubation of immune cells and target cells expressing activator receptor ligand in the absence of inhibitory receptor ligand; 2) less than or equal to 20% specific killing of target cell expressing both an activator receptor ligand and an inhibitory receptor ligand. As a further example, the immune cells are capable of killing at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 97% or at least 99% of target cells expressing the activator ligand and not the inhibitor ligand over a period of 6 hours, 12 hours, 18 hours, 24 hours, 30 hours, 36 hours, 42 hours, 48 hours, 54 hours, or 60 hours, while killing less than 40%, less than 30%, less than 20%, less than 10%, less than 5%, less than 3% or less than 1% of target cells expressing the activator and inhibitor ligands over the same time period.

[0096] A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at least about 50% to at least about 95%. A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95%. A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at most about 50%, at most about 55%, at most about 60%, at most about 65%, at most about 70%, at most about 75%, at most about 80%, at most about 85%, at most about 90%, or at most about 95%. A suitable specific killing value of target cells expressing both an activator receptor ligand and an inhibitory receptor ligand for the immune cells, pharmaceutical compositions, and kits can be less than about 50%, less than about 45%, less than about 40%, less than about 35%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, or less than about 5%. The suitable specific killing value for the immune cells, pharmaceutical compositions, and kits can be determined following about 6 hours, about 12 hours, about 18 hours, about 24 hours, about 30 hours, about 36 hours, about 42 hours, about 48 hours, about 54 hours, about 60 hours, about 66 hours, or about 72 hours of co-incubation of immune cells with target cells.

[0097] A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at least about 50% to at least about 95%. A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95%. A suitable specific killing value of the target cell

expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at most about 50%, at most about 55%, at most about 60%, at most about 65%, at most about 70%, at most about 75%, at most about 80%, at most about 85%, at most about 90%, or at most about 95%. A suitable specific killing value of target cells expressing both an activator receptor ligand and an inhibitory receptor ligand for the immune cells, pharmaceutical compositions, and kits can be less than about 50%, less than about 45%, less than about 40%, less than about 35%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, or less than about 5%. The suitable specific killing value for the immune cells, pharmaceutical compositions, and kits can be determined following about 6 hours, about 12 hours, about 18 hours, about 24 hours, about 30 hours, about 36 hours, about 42 hours, about 48 hours, about 54 hours, about 60 hours, about 66 hours, or about 72 hours of co-incubation of immune cells with target cells.

[0098] A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at least about 50% to at least about 95%. A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95%. A suitable specific killing value of the target cell expressing an activator ligand in the absence of an inhibitory ligand value for the immune cells, pharmaceutical compositions, and kits can be, for example, at most about 50%, at most about 55%, at most about 60%, at most about 65%, at most about 70%, at most about 75%, at most about 80%, at most about 85%, at most about 90%, or at most about 95%. A suitable specific killing value of target cells expressing both an activator receptor ligand and an inhibitory receptor ligand for the immune cells, pharmaceutical compositions, and kits can be less than about 50%, less than about 45%, less than about 40%, less than about 35%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, or less than about 5%. The suitable specific killing value for the immune cells, pharmaceutical compositions, and kits can be determined following about 6 hours, about 12 hours, about 18 hours, about 24 hours, about 30 hours, about 36 hours, about 42 hours, about 48 hours, about 54 hours, about 60 hours, about 66 hours, or about 72 hours of co-incubation of immune cells with target cells.

[0099] As used herein, the term "functional variant" refers to a protein that has one or more amino-acid substitutions, insertions, or deletions as compared to a parental protein, and which retains one or more desired activities of the parental protein. A functional variant may be a fragment of the protein (*i.e.* a variant having N- and/or C-terminal deletions) that retain the one or more desired activities of the parental protein.

[0100] As used herein, a "non-target allelic variant" refers to an allele of a gene whose product is expressed by non-target cells, but is not expressed by target cells. For example, a non-target allelic variant is an allele of a gene that is expressed by normal, non-cancer cells of a subject, but not expressed by cancer cells of the subject. The expression of the non-target allelic variant can be lost in the cancer cells by any mechanism, including loss of heterozygosity, mutation, or epigenetic modification of the gene encoding the non-target allelic variant.

[0101] As used herein, "specific to" or "specifically binds to" when used with respect to a ligand binding domain, such as an antigen binding domain, refers to a ligand binding domain that has a high specificity for a named target. Antibody specificity can be viewed as a measure of the goodness of fit between the ligand binding domain and the corresponding ligand, or the ability of the ligand binding domain to discriminate between similar or even dissimilar ligands. In comparison with specificity, affinity is a measure of the strength of the binding between the ligand binding domain and ligand, such that a low-affinity ligand binding domain binds weakly and high-affinity ligand binding domain binds firmly. A ligand binding domain that is specific to a target allele is one that can discriminate between different alleles of a gene. For example, a ligand binding domain that is specific to HLA-A*02 will not bind, or bind only weakly to, other HLA-A alleles such as HLA-A*01 or HLA-A*03. In the context of the activator and inhibitory receptors described herein, the ligand binding domain mediates specific activation or inhibition of the immune cell response. The person of skill in the art will appreciate that a ligand binding domain can be said to be specific to a particular target, and yet still have low levels of binding to one or more additional targets that do not affect its function in the receptor systems described herein.

[0102] In case of conflict, the present application, including any definitions herein, will control. However, mention of any reference, article, publication, patent, patent publication, and patent application cited herein is not, and should not be taken as an acknowledgment, or any form of suggestion, that they constitute valid prior art or form part of the common general knowledge in any country in the world.

Activator Receptors

[0103] The disclosure provides a first receptor, comprising a first extracellular ligand binding domain specific to a target antigen comprising a cancer cell-specific antigen, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I). The first receptor is an activator receptor, and mediates activation of an immune cell expressing the first receptor upon binding of the target antigen by the extracellular ligand binding domain of the first receptor. The

first receptor is responsive to a target antigen (*i.e.* activator ligand). For example, when a target antigen binds to or contacts the first receptor, the first receptor is responsive and activates an immune cell expressing the first receptor upon binding of the target antigen by the extracellular ligand binding domain of the first receptor. In some embodiments, the first receptor is a chimeric antigen receptor (CAR). In some embodiments, the first receptor is a T cell receptor (TCR).

[0104] In some embodiments, the first receptor is humanized. As used herein, "humanized" refers to the replacement of a sequence or a subsequence in a transgene that has been isolated or derived from a non-human species with a homologous, or functionally equivalent, human sequence. For example, a humanized antibody can be created by grafting mouse CDRs into human framework sequences, followed by back substitution of certain human framework residues for the corresponding mouse residues from the source antibody.

Activator Targets

[0105] According to the present invention, the target antigen for the first receptor is epidermal growth factor receptor (EGFR), or a peptide antigen of EGFR in a complex with a major histocompatibility complex class I (MHC-I).

[0106] The major histocompatibility complex class I (MHC-I) is a protein complex that displays antigens to cells of the immune system, triggering an immune response. The Human Leukocyte Antigens (HLAs) corresponding to MHC-I are HLA-A, HLA-B and HLA-C.

[0107] Cancer cell-specific pMHC antigens comprising any of HLA-A, HLA-B, HLA-C, HLA-E, HLA-F or HLA-G are described herein (not according to the invention). The cancer cell-specific antigen may comprise HLA-A (not according to the invention). HLA-A receptors are heterodimers comprising a heavy α chain and smaller β chain. The α chain is encoded by a variant of HLA-A, while the β chain (β 2-microglobulin) is an invariant (not according to the invention). There are several thousand variant HLA-A genes (not according to the invention). The MHC-I may comprise a human leukocyte antigen A*02 allele (HLA-A*02) (not according to the invention).

[0108] As described herein, but not according to the invention, the cancer cell-specific antigen comprises HLA-B. Hundreds of versions (alleles) of the HLA-B gene are known, each of which is given a particular number (such as HLA-B27).

[0109] As described herein, but not according to the invention, the cancer cell-specific antigen comprises HLA-C. HLA-C belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (β 2-microglobulin). Over one hundred HLA-C alleles are known in the art.

[0110] As described herein, but not according to the invention, the cancer cell-specific antigen is a lung cancer antigen, a glioblastoma antigen, a breast cancer antigen, a head and neck cancer antigen or a colorectal cancer antigen. As described herein, but not according to the invention, the cancer cell-specific antigen is a colorectal cancer antigen. As described herein, but not according to the invention, the cancer cell-specific antigen is EGFR or a peptide antigen thereof.

[0111] As described herein, but not according to the invention, the cancer cell-specific antigen is EGFR, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I). EGFR is a transmembrane protein that is receptor form members of the epidermal growth factor (EGF) family of extracellular protein ligands. EGFR is widely expressed in normal tissues, such as mesenchymal, epithelial and neuronal tissues, and plays an important role in cell proliferation, differentiation, and development. EGFR is also highly expressed in a variety of solid tumors, and EGFR expression correlates with tumor progression, resistance to chemotherapy, and poor prognosis.

[0112] All isoforms of EGFR and clinically relevant mutations of EGFR (*e.g.* EGFRvIII, S468/492R) are envisaged as cancer cell-specific antigens of the disclosure. EGFR isoform a is described in NCBI record number NP_005219.2. In some embodiments, EGFR comprises an amino acid sequence of:

1 MRPSGTAGAA LLALLAALCP ASRALEEEKV CQGTSNKLTQ LGTFEDHFLS LQRMFNNCEV
 61 VLGNLEITYV QRNYDLSFLK TIQEVAGYVL IALNTVERIP LENLQIIRGN MYYENSYALA
 121 VLSNYDANKT GLKELPMRNL QEILHGAVRF SNNPALCNVE SIQWRDIVSS DFLSNMSMDF
 181 QNHLGSCQKC DPSCPNGSCW GAGEENCQKL TKIICAQQCS GRCRGKSPSD CCHNQCAAGC
 241 TGPRESDECLV CRKFRDEATC KDTCPPLMLY NPTTYQMDVN PEGKYSFGAT CVKKCPRNYV
 301 VTDHGSCVRA CGADSYEMEE DGVRKCKKCE GPCRKVCNGI GIGEFKDSLS INATNIKHFK
 361 NCTSISGDLH ILPVAFRGDS FTHTPPLDPQ ELDILKTVE ITGFLLIQAW PENRTDLHAF
 421 ENLEIIRGRT KQHGOFS LAV VSLNITSLGL RSLKEISDGD VIISGNKNLC YANTINWKKL
 481 FGTSQGQTKI ISNRGENSCK ATGQVCHALC SPEGCWGPEP RDCVSCRNVS RGRECVDKCN
 541 LLEGEPPREFV ENSECIQCHP ECLPQAMNIT CTGRGPDNCI QCAHYIDGPH CVKTCPAGVM
 601 GENNTLVWKY ADAGHVCHLC HPNCTYGCTG PGLEGCPNTG PKIPSIATGM VGALLLLLVV
 661 ALGIGLFMR R RHIVRKRTL RLLQERELVE PLTPSGEAPN QALLRILKET EFKKIKVLGS
 721 GAFGTVYKGL WIPEGEKVKI PVAIKELREA TSPKANKEIL DEAYVMASVD NPHVCRLLGI
 781 CLTSTVQLIT QLMFFGCLLD YVREHKDNIG SQYLLNWCVQ IAKGMNYLED RRLVHRDLAA
 841 RNVLVKTPQH VKITDFGLAK LLGAEKEYH AEGGKVPKW MALESILHRI YTHQSDVWSY
 901 GVTWELMTF GSKPYDGIPA SEISSILEKG ERLPQPPICT IDVYMIMVKC WMIDADSRPK
 961 FRELIIEFSK MARDPQRYLV IQGDERMHL P SPTDSNFYRA LMDEEDMDV VDADEYLIPQ
 1021 QGFFSSPSTS RTPLSSLSA TSNNSTVACI DRNGLQSCPI KEDSFLQRY S SDPTGALTED
 1081 SIDDFTFLPVP EYINQSVPKR PAGSVQNPVY HNQPLNPAPS RDPHYQDPHS TAVGNPEYLN
 1141 TVQPTCVNST FDSFAHWAQK GSHQISLDNP DYQQDFFPKE AKPNGIFKGS TAENAEYLRV
 1201 APQSSEFIGA (SEQ ID NO: 1).

EGFR isoform b is described in NCBI record number NP_958439.1. In some embodiments, EGFR comprises an amino acid sequence of:

1 MRPSGTAGAA LLALLAALCP ASRALEEEKV CQGTSNKLTQ LGTFEDHFLS LQRMFNNCEV
 61 VLGNLEITYV QRNYDLSFLK TIQEVAGYVL IALNTVERIP LENLQIIRGN MYYENSYALA
 121 VLSNYDANKT GLKELPMRNL QEILHGAVRF SNNPALCNVE SIQWRDIVSS DFLSNMSMDF
 181 QNHLGSCQKC DPSCPNGSCW GAGEENCQKL TKIICAQQCS GRCRGKSPSD CCHNQCAAGC
 241 TGPRESDECLV CRKFRDEATC KDTCPPLMLY NPTTYQMDVN PEGKYSFGAT CVKKCPRNYV
 301 VTDHGSCVRA CGADSYEMEE DGVRKCKKCE GPCRKVCNGI GIGEFKDSLS INATNIKHFK
 361 NCTSISGDLH ILPVAFRGDS FTHTPPLDPQ ELDILKTVE ITGFLLIQAW PENRTDLHAF
 421 ENLEIIRGRT KQHGOFS LAV VSLNITSLGL RSLKEISDGD VIISGNKNLC YANTINWKKL
 481 FGTSQGQTKI ISNRGENSCK ATGQVCHALC SPEGCWGPEP RDCVSCRNVS RGRECVDKCN
 541 LLEGEPPREFV ENSECIQCHP ECLPQAMNIT CTGRGPDNCI QCAHYIDGPH CVKTCPAGVM
 601 GENNTLVWKY ADAGHVCHLC HPNCTYGS (SEQ ID NO: 2).

EGFR isoform c is described in NCBI record number NP_958440.1. In some embodiments, EGFR comprises an amino acid sequence of:

1 MRPSGTAGAA LLALLAALCP ASRALEEEKV CQGTSNKLTQ LGTFEDHFLS LQRMFNNCEV
 61 VLGNLEITYV QRNYDLSFLK TIQEVAGYVL IALNTVERIP LENLQIIRGN MYYENSYALA
 121 VLSNYDANKT GLKELPMRNL QEILHGAVRF SNNPALCNVE SIQWRDIVSS DFLSNMSMDF
 181 QNHLGSCQKC DPSCPNGSCW GAGEENCQKL TKIICAQQCS GRCRGKSPSD CCHNQCAAGC
 241 TGPRESDECLV CRKFRDEATC KDTCPPLMLY NPTTYQMDVN PEGKYSFGAT CVKKCPRNYV
 301 VTDHGSCVRA CGADSYEMEE DGVRKCKKCE GPCRKVCNGI GIGEFKDSLS INATNIKHFK
 361 NCTSISGDLH ILPVAFRGDS FTHTPPLDPQ ELDILKTVE ITGLS (SEQ ID NO: 3).

EGFR isoform d is described in NCBI record number NP_958441.1. In some embodiments, EGFR comprises an amino acid sequence of:

50

55

1 MRPSGTAGAA LLALLAALCP ASRALEEKKV CQGTSNKLTQ LGTFEDHFLS LQRMFNANCEV
 61 VLGNIETIYV QRNYDLSFLK TIQEVAGYVL IALNTVERIP LENLQIIRGN MYEENSYALA
 121 VLSNYDANKT GLKELPMRNL QEILHGAVRF SNNPALCNVE SIQWRDIVSS DFLSNMSMDF
 181 QNHLGSCQKC DPSCPNGSCW GAGEENCQKL TKIICAQQCS GRCRGKSPSD CCHNQCAAGC
 241 TGPRESCLV CRKFRDEATC KDTCPPLMLY NPPTYQMDVN PEGKYSFGAT CVKKCPRNYV
 301 VTDHGSCVRA CGADSYEMEE DGVRKCKKCE GPCRKVCNGI GIGEFKDSLS INATNIKHFK
 361 NCTISGDLH ILPVAFRGDS FTHTPPLDPQ ELDILKTVE ITGFLLIQAW PENRTDLHAF
 421 ENLEIIRGRT KQHGOFS LAV VSLNITSLGL RSLKEISDGD VIISGNKNLC YANTINWKKL
 481 FGTSGQKTKI ISNRGENSCK ATGQVCHALC SPEGCWGPEP RDCVSCRNVS RGRECVDKCN
 541 LLEGEPRFV ENSECIQCHP ECLPQAMNIT CTGRGPDNCI QCAHYIDGPH CVKTCFAGVM
 601 GENNTLVWKY ADAGHVCHLC HPNCTYGGPN ESLKAMLFCL FKLSSCNQSN DGSVSHQSGS
 661 PAAQESCLGW IPSLLPSEFQ LGWGGCSHLH AWPSASVIIT ASSCH (SEQ ID NO: 4).

EGFR isoform e is described in NCBI record number NP_001333826.1. In some embodiments, EGFR comprises an amino acid sequence of:

1 MRPSGTAGAA LLALLAALCP ASRALEEKKV CQGTSNKLTQ LGTFEDHFLS LQRMFNANCEV
 61 VLGNIETIYV QRNYDLSFLK TIQEVAGYVL IALNTVERIP LENLQIIRGN MYEENSYALA
 121 VLSNYDANKT GLKELPMRNL QGQKCDPSCP NGSCWGAGEE NCQKLTKIIC AQQCSGRCRG
 181 KSPSDCCHNQ CAAGCTGPRE SDCLVCRKFR DEATCKDTCPLMLYNPPTY QMDVNPEGKY
 241 SFGATCVKKC PRNYVVDHG SCVRACGADS YEMEEDGVRK CKKCEGPCRK VCNIGIGIEF
 301 KDSLSINATN IKHFKNCTSI SGDLHILPVA FRGDSFTHTP PLDPQELDIL KTVKEITGFL
 361 LIQAWPENRT DLHAFENLEI IRGRTKQHGO FSLAVVSLNI TSLGLRSLKE ISDGDVIISG
 421 NKNLCYANTI NWKKLFGTSG QKTKIISNRG ENSCKATGQV CHALCSPEGC WGPEPRDCVS
 481 CRNVSRGREG VDKCNLLEGE PREFVENSEC IQCHPECLPQ AMNITCTGRG PDNCIQCAHY
 541 IDGPHCVKTC PAGVMGENNT LVWKYADAGH VCHLCHPNCT YGCTGPGLEG CPTNGPKIPS
 601 IATGMVGALL LLLVVALGIG LFMRRRHIVR KRTLRLRLQE RELVEPLTPS GEAPNQALLR
 661 ILKETEFKKI KVLGSGAFGT VYKGLWIPEG EKVKIPVAIK ELREATSPKA NKEILDEAYV
 721 MASVDNPHVC RLLGICLTST VQLITQLMPF GCLLDYVREH KDNIGSQYLL NWCVQIAKGM
 781 NYLEDRLRLVH RDLAARNVLV KTPQHVKITD FGLAKLLGAE EKEYHAEGGK VPIKWMALLES
 841 ILHRIYTHQS DVWSYGVTWV ELMTFGSKPY DGIPASEISS ILEKGERLPQ PPICTIDVYM
 901 IMVKCWMIDA DSRPKFRELI IEFKSMARDP QRYLVIQGDE RMHLPSPTDS NFYRALMDEE
 961 DMDDVVDADE YLIPQQGFFS SPSTSRTPLL SLSATSNNNS TVACIDRNLG QSCPIKEDSF
 1021 LQRYSSDPTG ALTEDSIDDT FLPVPGEWLV WKQSCSSTSS THSAAASLQC PSQVLPPASP
 1081 EGETVADLQT Q (SEQ ID NO: 5).

EGFR isoform f is described in NCBI record number NP_001333827.1. In some embodiments, EGFR comprises an amino acid sequence of:

40

45

50

55

1 MRPSGTAGAA LLALLAALCP ASRALEEKKV CQGTSNKLTQ LGTFEDHFLS LQRMFNNCEV
 61 VLG NLEITYV Q RNYDLSFLK TIQEVAGYVL IALNTVERIP LENLQIIRGN MYYENSYALA
 121 VLSNYDANKT GLKELPMRNL QEILHGAVRF SNNPALCNVE SIQWRDIVSS DFLSNMSMDF
 181 QNHLGSCQKC DPSPNGSCW GAGEENCQKL TKIICAQQCS GRCRGKSPSD CCHNQCAAGC
 5 241 TGPRES DCLV CRKFRDEATC KDTCPPLMLY NPPTYQMDVN PEGKYSFGAT CVKKCPRNYV
 301 VTDHGSCVRA CGADSYEMEE DGVRKCKKCE GPCRKCNGI GIGEFKDSLS INATNIKHF
 361 NCTSI SGLH ILPVAFRGDS FTHTPPLDPQ ELDILKTVKE ITGFLLIQAW PENRTDLHAF
 421 ENLEIIRGRT KQHGQFSLAV VSLNITSLGL RSLKEISDGD VIISGNKNLC YANTINWKKL
 481 FGTSGQKTKI ISNRGENSCK ATGQVCHALC SPEGCWGPEP RDCVSCRNV S RGRECVDKCN
 10 541 LLEGEPRFV ENSECIQCHP ECLPQAMNIT CTGRGPDNCI QCAHYIDGPH CVKTC PAGVM
 601 GENNTLVWKY ADAGHVCHLC HPNCTYGCTG PGLEGCP TNG PKIPSIATGM VGALLLLLVV
 661 ALGIGLFMR RHIVRKRTL RLLQERELVE PLTPSGEAPN QALLRILKET EFKKIKVLGS
 721 GAFGT VYKGL WIPEGEKVKI PVAIKELREA TSPKANKEIL DEAYVMASVD NPHVCRLGI
 781 CLTSTVQLIT QLMFP GCLLD YVREHKDNIG SQYLLNWC VQ IAKGMNYLED RRLVHRDLAA
 841 RNVLVKTPQH VKITDFGLAK LLGAE EKEYH AEGGKVP I KW MALESILHRI YTHQSDVWSY
 15 901 GVTVWELMTF GSKPYDGIPA SEISSILEKG ERLPQPPICT IDVYMIMVKC WMIDADSRPK
 961 FRELIIEFSK MARDPQRYLV IQGDERMHL P SPTDSNFYRA LMDEEDMDDV VDADEYLIPQ
 1021 QGFFSSPSTS RTPLLSSLSA TSNNSTVACI DRNGLQSCPI KEDSFLQ RYS SDPTGALTED
 1081 SIDD TFLPVP GEWL VWKQSC SSTSS THSAA ASLQCP SQVL PPASPEGETV ADLQTQ (SEQ ID
 NO: 6).

EGFR isoform h is described in NCBI record number NP_001333829.1. In some embodiments, EGFR comprises an amino acid sequence of:

1 MFNNCEVVLG NLEITYVQRN YDLSFLKTIQ EVAGYVLIAL NTVERIPL EN LQIIRGNMY Y
 61 ENSYALAVLS NYDANKTGLK ELPMRNLQEI LHGAVRFSNN PALCNVESIQ WRDIVSSDFL
 121 SNMSMDFQNH LGSCQKCDPS CPNGSCWGAG EENCQKLTKI ICAQQCSGRC RGKSPSDCCH
 181 NQCAAGCTGP RESDCLVCRK FRDEATCKDT CPPLMLYNPT TYQMDVNPEG KYSFGATCVK
 241 KCPRNYVVD HGS CVRACGA DSYEMEEDGV RKCKKCEGPC RKVCNGIGIG EFKDSLSINA
 30 301 TNIKHFKNCT SISGDLHILP VAFRGDSFTH TPPLDPQELD ILKTVKEITG FLLIQAWPEN
 361 RTDLHAFENL EIIRGRTKQH GQFSLAVVSL NITSLGLRSL KEISDGDVII SGNKNLCYAN
 421 TINWKLFGT SGQKTKIISN RGENSCKATG QVCHALCSPE GCWGPEPRDC VSCRNVSRGR
 481 ECVDKCNLLE GEPREFVENS ECIQCHPECL PQAMNITCTG RGPDN CIQCA HYIDGPHCVK
 541 TCPAGVMGEN NTLVWKYADA GHVCHLCHPN CTYGCTGPGL EGCPTNGPKI PSIATGMVGA
 601 LLLLLVVALG IGLFMRRRHI VRKRTLRLRL QERELVEPLT PSGEAPNQAL LRILKETEFK
 35 661 KIKVLGSGAF GTVYKGLWIP EGEKV KIPVA IKELREATSP KANKEILDEA YVMASVDNPH
 721 VCRLLGICLT STVQLITQLM PFGCLLDYVR EHKNIGSQY LLNWC VQIAK GMNYLED RRL
 781 VHRDLAARNV LVKTPQHVKI TDFGLAKLLG AEEKEYHAEG GKVPIKWMAL ESILHRIYTH
 841 QSDVWSYGV T VWELMTFGSK PYDGIPASEI SSILEKGERL PQQPICTIDV YMIMVKC WMI
 901 DADSRPKFRE LIIEFSKMAR DPQRYLVIQ G DERMHLPSPT DSNFYRALMD EEDMDDV VDA
 40 961 DEYLIPQQGF FSSPSTSRTF LLSSLSATSN NSTVACIDRN GLQSCPIKED SFLQRYSSDP
 1021 TGALTEDSID DTFLPVPEYI NQSVPKRPAG SVQNPVYHNQ PLNPAPSRDP HYQDPHSTAV
 1081 GNPEYLN TVQ PTCVNSTFDS PAHWAQKGSH QISLDNPDYQ QDFFPK EAKP NGIFKGSTAE
 1141 NAEYLRVAPQ SSEFIGA (SEQ ID NO: 7).

EGFR isoform i is described in NCBI record number NP_001333870.1. In some embodiments, EGFR comprises an amino acid sequence of:

1 MRPSGTAGAA LLALLAALCP ASRALEEKKG NYVVTDHGSC VRACGADSYE MEEDGVRKCK
 61 KCEGPCRKVC NGIGIGEFKD SLSINATNIK HFKNCTSIG DLHILPVAFR GDSFTHTPPL
 121 DPQELDILKT VKEITGFLLI QAWPENRTDL HAFENLEIIR GRTKQHGQFS LAVVSLNITS
 181 LGLRSLKEIS DGDVIISGNK NLCYANTINW KKLFGTSGQK TKIISNRGEN SCKATGQVCH
 241 ALCSPEGCWG PEPRDCVSCR NVSRGRECDV KCNLLGEPR EFVENSECIQ CHPECLPQAM
 301 NITCTGRGPD NCIQCAHYID GPHCVKTCPA GVMGENNTLV WKYADAGHVC HLCHPNCTYG
 361 CTGPGLEGCP TNGPKIPSIA TGMVGALLLL LVVALGIGLF MRRRHIVRKR TLRRLLQERE
 421 LVEPLTPSGE APNQALLRIL KETEFKKIKV LGSGAFGTvy KGLWIPEGEK VKIPVAIKEL
 481 REATSPKANK EILDEAYVMA SVDNPHVCRL LGICLTSTVQ LITQLMPFGC LLDYVREHKD
 541 NIGSQYLLNW CVQIAKGMNY LEDRRLVHRD LAARNVLVKT PQHVKITDFG LAKLLGAEEK
 601 EYHAEGGKVP IKWMALESIL HRIYTHQSDV WSYGVTWEL MTFGSKPYDG IPASEISSIL
 661 EKGERLPQPP ICTIDVYMIM VKCWMIDADS RPKFRELIIE FSKMARDPQR YLVIQGDERM
 721 HLPSPDTSNF YRALMDEEDM DDVVDAD EYL IPQGGFFSSP STSRTPLLSS LSATSNNSTV
 781 ACIDRNLQSS CPIKEDSFLQ RYSSDPTGAL TEDSIDDTFL PVPEYINQSV PKRPAGSVQN
 841 PVYHNQPLNP APSRDPHYQD PHSTAVGNPE YLNTVQPTCV NSTFDSPAHW AQKGSHQISL
 901 DNPDYQQDFF PKEAKPNGIF KGSTAENAEY LRVAPQSSEF IGA (SEQ ID NO: 8).

In some embodiments, the cancer cell-specific antigen is a peptide antigen derived from EGFR. For example, the peptide antigen comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to a sequence or subsequence of any one of SEQ ID NOs: 1-8. In some embodiments, the peptide antigen comprises a sequence identical to a subsequence of any one of SEQ ID NOs: 1-8 or SEQ ID NOs: 918-919.

Extracellular Ligand Binding Domain

[0113] The disclosure provides a first receptor, comprising a first extracellular ligand binding domain specific to a target antigen. In some embodiments, the target antigen comprises a cancer cell-specific antigen.

[0114] In some embodiments, the cancer cell-specific antigen is EGFR or a EGFR-derived peptide antigen complexed with MHC-I, and the ligand binding domain of the first receptor recognizes and binds to the EGFR antigen. In some embodiments, the cancer cell-specific antigen is a clinically relevant mutant of EGFR. For example, a mutated variant of EGFR can be a substitution mutation (*e.g.* L858R in exon 21), a deletion mutation (*e.g.* EGFRvIII or deletions in exon 19 or exon 23), or amplifications mutations to the EGFR gene.

[0115] Not according to the invention: The ligand binding domain may be an antigen binding domain. Exemplary antigen binding domains include, *inter alia*, scFv, SdAb, V β -only domains, and TCR antigen binding domains derived from the TCR α and β chain variable domains.

[0116] For example, the first extracellular ligand binding domain may be part of a contiguous polypeptide chain including, for example, a V β -only domain, a single domain antibody fragment (sdAb) or heavy chain antibodies HCAb, a single chain antibody (scFv) derived from a murine, humanized or human antibodies (Harlow et al., 1999, In: Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, N.Y.; Harlow et al., 1989, In: Antibodies: A Laboratory Manual, Cold Spring Harbor, N.Y.; Houston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; Bird et al., 1988, Science 242:423-426). In some aspects, the first extracellular ligand binding domain comprises an antigen binding domain that comprises an antibody fragment. In further aspects, the first extracellular ligand binding domain comprises an antibody fragment that comprises a scFv or an sdAb.

[0117] The term "antibody," as used herein, refers to a protein, or polypeptide sequences derived from an immunoglobulin molecule, which specifically binds to an antigen. Antibodies can be intact immunoglobulins of polyclonal or monoclonal origin, or fragments thereof and can be derived from natural or from recombinant sources.

[0118] The terms "antibody fragment" or "antibody binding domain" refer to at least one portion of an antibody, or recombinant variants thereof, that contains the antigen binding domain, *i.e.*, an antigenic determining variable region of an intact antibody, that is sufficient to confer recognition and specific binding of the antibody fragment to a target, such as an antigen and its defined epitope. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments, single-chain (sc)Fv ("scFv") antibody fragments, linear antibodies, single domain antibodies (abbreviated "sdAb") (either VL or VH), camelid VHH domains, and multi-specific antibodies formed from antibody fragments.

[0119] The term "scFv" refers to a fusion protein comprising at least one antibody fragment comprising a variable region of a light chain and at least one antibody fragment comprising a variable region of a heavy chain, wherein the light and heavy chain variable regions are contiguously linked via a short flexible polypeptide linker, and capable of being expressed as a single polypeptide chain, and wherein the scFv retains the specificity of the intact antibody from which it is derived.

[0120] "Heavy chain variable region" or "VH" (or, in the case of single domain antibodies, *e.g.*, nanobodies, "VHH")

with regard to an antibody refers to the fragment of the heavy chain that contains three CDRs interposed between flanking stretches known as framework regions, these framework regions are generally more highly conserved than the CDRs and form a scaffold to support the CDRs.

[0121] Unless specified, as used herein a scFv may have the VL and VH variable regions in either order, *e.g.*, with respect to the N-terminal and C-terminal ends of the polypeptide, the scFv may comprise VL-linker-VH or may comprise VH-linker-VL.

[0122] In some embodiments, the antigen binding domain of the activator and/or inhibitory receptor comprises an scFv. In some embodiments, the scFv may comprise a VL and VH region joined by a linker. In some embodiments, the linker may comprise a glycine serine linker, for example GGGGSGGGGSGGGGSGG (SEQ ID NO: 136). In some embodiments, the scFv may further comprise a signal sequence at the N terminus of the scFv. Exemplary signal sequences include MDMRVPAQLLGLLLWLRGARC (SEQ ID NO: 137), which is encoded by

ATGGACATGAGGGTCCCCGCTCAGCTCCTGGGGCTCCTGCTACTCTGGCTCCGAGGT
GCCAGATGT (SEQ ID NO: 138)

or

ATGGATATGAGAGTGCCTGCCAGCTGCTCGGACTGCTCCTTCTGTGGTTGAGAGGA
GCTCGGTGC (SEQ ID NO: 917).

[0123] The term "antibody light chain" refers to the smaller of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations. Kappa ("K") and lambda ("λ") light chains refer to the two major antibody light chain isotypes.

[0124] The term "recombinant antibody" refers to an antibody that is generated using recombinant DNA technology, such as, for example, an antibody expressed by a bacteriophage or yeast expression system. The term should also be construed to mean an antibody, which has been generated by the synthesis of a DNA molecule encoding the antibody and which DNA molecule expresses an antibody protein, or an amino acid sequence specifying the antibody, wherein the DNA or amino acid sequence has been obtained using recombinant DNA or amino acid sequence technology, which is available and well known in the art.

[0125] The term "Vβ domain", "Vβ-only domain", "β chain variable domain" or "single variable domain TCR (svd-TCR)" refers to an antigen binding domain that consists essentially of a single T Cell Receptor (TCR) beta variable domain that specifically binds to an antigen in the absence of a second TCR variable domain. The Vβ-only domain engages antigen using complementaritydetermining regions (CDRs). Each Vβ-only domain contains three complement determining regions (CDR1, CDR2, and CDR3). Additional elements may be combined provided that the Vβ domain is configured to bind the epitope in the absence of a second TCR variable domain.

[0126] In some embodiments, the extracellular ligand binding domain of the first receptor comprises an antibody fragment, a single chain Fv antibody fragment (scFv), or a β chain variable domain (Vβ).

[0127] In some embodiments, the extracellular ligand binding domain of the first receptor comprises a TCR α chain variable domain and a TCR β chain variable domain.

[0128] In some embodiments, the extracellular ligand binding domain of the first receptor comprises an scFv antigen binding domain. Exemplary EGFR scFv are shown in Table 1 below.

Table 1. EGFR scFv domains

Amino Acid Sequence	DNA Sequence
CT-478 EGFR (VH-VL scFv Format): QVQLVESGGGVWQPGRSLRLSC AASGFTFTYGMHWVRQAPG KGLEWVAWIWDDGSYKYYGDS VKGRFTISRDNKNTLYLQMNS LRAEDTAVYYCARDGITMVRGV MKDYFDYWGGTLTVVSSGG GGSGGGSGGGGGGAIQLTQ SPSSLSASVGDRTVITCRASQDIS SALVWYQQKPGKAPKLLIYDAS SLESGVPSRFSGSEGTDFLTIS SLQPEDFATYCCQFNYSPLTF GGGTKVEIK (SEQ ID NO: 9)	CT-478 EGFR: CAGGTGCAGCTGGTGGAGTCTGGGGGAGGCGTGGTCCAGCCTGGGAGGT CCCTGAGACTCTCTGTGCAGCGTCTGGATTACCTTCAGTACCTATGGCA TGCACTGGTCCGCCAGGCTCCAGGCAAGGGCTGGAGTGGTGGCAGT TATATGGGATGATGGAAGTTATAAATACTATGGAGACTCCGTGAAGGGCC GATTCACCATCTCCAGAGACAATCCAAAGAACACGCTGTATCTGCAAAATGA ACAGCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGATGG TATTACTATGTTCCGGGAGTTATGAAGGACTACTTTGACTACTGGGGCC AGGAAACCTGGTCACCGTCTCTCAGGCGAGGTGGAAGCGGAGGGGG AGGATCTGCGGGGAGGAGCGGAGCGCCATCCAGTTGACCCAGTCT CCATCTCCCTGCTGCACTCTGTAGGAGACAGAGTCAACCATCACTTGCCGG GCAAGTCAGGACATTAGCAGTGCTTGTAGTCTGGTATCAGCAGAAACAGG GAAAGCTCTAAGCTCTGATCTATGATGCCCTCCAGTTGGAAAGTGGGG TCCCATCAAGGTTTCAGCGGAGTGAATCTGGACAGATTCACCTCACCA TCAGCAGCTGCAGCTGAAGATTTTGCAACTTATTACTGTCAACAGTTTA ATAGTTACCCGCTCACTTTCGGCGGAGGGACCAAGGTGGAGATCAAAA (SEQ ID NO: 139)
CT-479 EGFR (VL-VH scFv Format): AIQLTQSPSSLSASVGDRTVITC RASQDISSALVWYQQKPGKAP KLLIYDASSLESVPSRFSGSESG TDFLTISLQPEDFATYCCQF NSYPLTFGGGKVEIKGGGSG GGSGGGGGGGVQLVESGG GVVQPGRSRLSCAASGFTFTY GMHWVRQAPGKGLEWVAVI WDDGSYKYYGDSVKGRFTISR	CT-479 EGFR: GCCATCCAGTTGACCCAGTCTCCATCCTCCTGTCTGCATCTGTAGGAGAC AGAGTCACCATCACTTCCGGGCAAGTCAGGACATTAGCAGTGCTTTAGT CTGGTATCAGCAGAAACAGGAAAGCTCCTAAGCTCCTGATCTATGATG CCTCCAGTTTGGAAAGTGGGTCCCATCAAGTTTCAGCGGAGTGAATCT GGGACAGATTTCACTCTACCATCAGCAGCCTGCAGCCTGAAGATTTTGA ACTTATTACTGTCAACAGTTAATAGTTACCCGCTCACTTCGGCGGAGGG ACCAAGGTGGAGATCAAAAGCGGAGGTGGAAGCGGAGGGGAGGATCT GGCGCGGAGGAAGCGGAGGCCAGGTGCAGTGGTGGAGTCTGGGGGA GGCGTGGTCCAGCTGGGAGGTCCCTGAGACTCTCTGTGCAGCGTCTGG ATTACCTTCAGTACCTATGGCATGCACTGGGTCCGCCAGGCTCCAGGCAA

(continued)

Amino Acid Sequence	DNA Sequence
NSKNTLYLQMNSLRAEDTAVVY CARDGITMVRGVMMKDYFDYW GQGTLLVTYSS (SEQ ID NO: 10)	GGGGCTGGAGTGGGTGGCAGTTATATGGGATGATGGAAGTTATAAATAC TATGGAGACTCCGTGAAGGGCCGATTACCATCTCCAGAGACAAATCCAA GAACACGCTGATCTGCAAAATGAACAGCCTGAGAGCCGAGGACACGGCT GTGATTACTGTGCGAGAGATGGTATTACTATGTTTCGGGGAGTTATGAA GGACTACTTTGACTACTGGGGCCAGGGAACCTGGTCAACCGTCTCCTCA (SEQ ID NO: 140)
CT-480 EGFR (VH-VL scFv format): QIQLVQSGPELKKPGETVKISCK ASGYTFEYPIHWVKQAPGKGF KWMGMIYTDIGKPTYAEFEKGF RFAFSLETSASTAYLQINNLIKNE DTATYFCVRDRYDSLFDYWQ GTTLTVSSGGGGSGGGSGGG GSGGDVVMQTQPLSLPVSIGD QASISCRSSQSLVHSNGNTYLH WYLQKPGQSPKLLIKVSNRFS GVPDRFSGSGSGTDFTLKISRVE AEDLGVYFCFSQSTHPVPTFGG GTKLEIK (SEQ ID NO: 11)	CT-480 EGFR: CAGATCCAGTTGGTGCAGTCTGGACCTGAGCTGAAGAACCTGGAGAGA CAGTCAAGATCTCCTGCAAGGCCTCTGGGTATACCTTCACAGAAATATCCAA TACACTGGGTGAAGCAGGCTCCAGGAAAGGTTTCAAGTGGATGGGCAT GATATACCCGACATTGAAAGCCAAACATATGCTGAAGAGTTCAAGGGAC GGTTTGCCTTCTTTGGAGACCTCTGCCAGCACTGCCTATTTGCAGATCAA CAACCTCAAAGAATGAGGACACGCTACATATTTCTGTGTAAGAGATCGAT ATGATTCCTCTTTGACTACTGGGGCCAAAGGCACCACTCTCACAGTCTCCTC AGGCGGAGGTGGAAGCGGAGGGGGAGGATCTGGCGGCGGAGGAAGCG GAGCGGATGTTGTGATGACCCAAACTCCACTCTCCCTGCCTGTCACTTTG GAGATCAAGCCTCCATCTTGCAGATCTAGTCAGAGCCTTGTACACAGTA ATGGAACACCTATTTACATTTGGTACCTGCAGAACCCAGGCCAGTCTCCAA AGCTCCTGATCTACAAAGTTTCCAACCGATTTTCTGGGGTCCCAGACAGGT TCAGTGGCAGTGGATCAGGGACAGATTTACACTCAAGATCAGCAGAGTG GAGGCTGAGGATCTGGGAGTTTATTTCTGCTCTCAAAAGTACACATGTTCCG TGGACGTTCCGTGGAGGCACCAAGCTGGAAATCAAA (SEQ ID NO: 141)
CT-481 EGFR (VL-VH scFv Format):	CT-481 EGFR:

(continued)

Amino Acid Sequence	DNA Sequence
DVVMTQTPLSLPVSLGDDQASIS CRSSQSLVHSNGNTYLHWYLQ KPGQSPKLLIYKVSNRFSVPDR FSGSGGTDFTLKISRVEAEDLG VFCSQSTHVPWTFGGGKLEI KGGGSGGGSGGGGGGGQI QLVQSGPELKPGETVKISKAS GYTFEYPIHWVKQAPGKGFK WMGMIYTDIGKPTYAEFEKGR FAFSLETSASTAYLQINILKNE TATYFCVRDRYDSLFDYWGGQ TTLTVSS (SEQ ID NO: 12)	GATGTTGTGATGACCCAACTCCACTCTCCCTGCCTGTGAGTCTTGGAGAT CAAGCCTCCATCTCTTGAGATCTAGTCAGAGCCTTGACACAGTAATGGA AACACCTATTACATTGGTACCTGCAGAAAGCCAGGCTCTCCAAAGCTC CTGATCTACAAAGTTTCCAACCGATTTTCTGGGGTCCAGACAGGTTCACT GGCAGTGGATCAGGGACAGATTTCACACTCAAGATCAGCAGAGTGGAGG CTGAGGATCTGGGAGTTTATTCTGCTCTCAAAATACACATGTTCCGTGGA CGTTCGGTGGAGGCACCAAGCTGAAATCAAAGCGGAGGTGGAAGCG GAGGGGAGGATCTGGCGGGGAGGAAAGCGGAGGCCAGATCCAGTTGG TGCACTCTGGACCTGAGCTGAAGAACCTGGAGAGACAGTCAAGATCTCC TGCAAGGCCTCTGGGTATACCTTCACAGAATATCCAAATACACTGGGTGAA GCAGGCTCCAGGAAAGGTTTCAAGTGGATGGCATGATATACACCGAC ATTGAAAAGCCAAACATATGCTGAAGAGTTCAAGGACGGTTGCTTCTC TTTGGAGACCTCTGCCAGACTGCCTATTTGCAGATCAACAACCTCAAGAA TGAGGACACGGCTACATATTTCTGTGTAAGAGATCGATATGATTCCTCTT TGACTACTGGGGCCCAAGGCACCACTCTCACAGTCTCCTCA (SEQ ID NO: 142)
CT-482 EGFR (VH-VL scFv Format): EMQLVESGGGFVKPGGSLKLS AASGFAFSHYDMSWVRQTPK QRLEWVAYIASGGDITYYADTV KGRFTISRDNQNTLYLQMSSL KSEDAMFYCSRSSYGNGDAL	CT-482 EGFR: GAAATGCAGCTGGTGGAGTCTGGGGAGGCTTCGTGAAGCCTGGAGGGT CCCTGAAACTCTCATGTGCAGCCTCTGGATTGCTTTCACTACTATGACAT GTCTTGGGTTCCGACAGACTCCGAAGCAGAGGCTGGAGTGGTCCGATACA TTGCTAGTGGTGGTGATATCACTACTATGCAGACACTGTGAAGGGCCGA TTCACCATCTCAGAGACAATGCCAGAACCCCTGTACCTGCAAATGAGC AGTCTGAAGTCTGAGGACACAGCCATGTTTTACTGTTACGATCCTCCTAT

(continued)

Amino Acid Sequence	DNA Sequence
DFWGQGTSTVTVSSGGGSGG GGSGGGGGDVVMTQPLSL PVS LGDQASISCRSSQSLVHSN GNTYLHWY LQKPGQSPKLLYK VSNRFGVPDRFSGSGSGTDFT LKISRVEAEDLGVYFCSQSTHVL TFGSGTKLEIK (SEQ ID NO: 13)	GGTAACAACGGAGATGCCCTGGACTTCTGGGGTCAAGGTACCTCAGTCAC CGTCTCTCAGGCGGAGGTGGAAGCGGAGGGGAGGATCTGGCGGGG AGGAAGCGGAGGCGATGTTGTATGACCCAACTCCACTCTCCCTGCCTG TCAGTCTTGAGATCAAGCCTCCATCTCTTGAGATCTAGTCAGAGCCTTG TTCACAGTAATGGAACACCTATTACATTGGTACCTGCAGAAGCCAGGCC AGTCCAAAGCTCTGATCTACAAAGTTTCCAACCGATTTCTGGGGTCC CAGACAGTTTCAGTGGCAGTGGATCAGGGACAGATTTCACACTCAAGATC AGCAGATGGAGGCTGAGGATCTGGGAGTTTATTCTGCTCTCAAAGTAC ACATGTTCTCACGTTGCGCTCGGGACAAAAGTTGGAAATAAAA (SEQ ID NO: 143)
CT-483 EGFR (VL-VH scFv Format): DVVMTQTPLSLPVSLG DQASIS CRSSQSLVHSNGNTYLHWY LQ KPGQSPKLLYK VSNRFGVPDR FSGSGGTDFTLKISRVEAEDLG VYFCSQSTHVLTFGSGTKLEIKG GGSGGGGGSGGGSGGEMQL VESGGGFVKPGGSLKLSCAASG FAFHYDMSWVRQTPKQRLE WVAYIASGGDITYADTVKGRF TISRDNQNTLYLQMSSLKSED TAMFYCSRSSYGNGDLDLFW GQGTSTVTVSS (SEQ ID NO: 14)	CT-483 EGFR: GATGTTGTGATGACCCAACTCCACTCTCCCTGCCTGTGAGTCTGGAGAT CAAGCCTCATCTCTTGAGATCTAGTCAGAGCCTTGTTCACAGTAATGGA AACACCTATTTACATTGGTACCTGCAGAAAGCCAGGCTCCAAAGCTC CTGATCTACAAAGTTTCCAACCGATTTCTGGGGTCCCAGACAGGTTTCAGT GGCAGTGGATCAGGGACAGATTTCACACTCAAGATCAGCAGAGTGGAGG CTGAGGATCTGGGAGTTTATTCTGCTCTCAAAGTACACATGTTCTCAGTT CGGCTCGGGACAAAGTTGGAATAAAAGCGGAGGTGGAAGCGGAGG GGGAGGATCTGCGGCGGAGGAAAGCGGAGGCGAAATGCAGCTGGTGGA GTCTGGGGGAGGCTTCGTGAAGCCTGGAGGGTCCCTGAAACTCTCATGTG CAGCCTCTGGATTCTGCTTTCAGTCACTATGACATGCTTGGGTTCCGCGA CTCCGAAGCAGAGGCTGGAGTGGGTCGCATACATTGCTAGTGGTGGTAT ATCAGCTACTATGCAGACACTGTGAAGGGCCGATTACCATCTCCAGAGAC AATGCCAAGAACACCTGTACTGCAAAATGAGCAGTCTGAAGTCTGAGGA CACAGCCATGTTTACTGTTACGATCTCCTATGTTGAACAAACGGAGATGC CCTGGACTTCTGGGGTCAAGGTACCTCAGTCACCGTCTCTCTCA (SEQ ID NO: 144)
CT-486 EGFR (VH-VL scFv Format):	CT-486 EGFR:

(continued)

Amino Acid Sequence	DNA Sequence
QVQLKQSGPGLVQPSSLSITC TVSGFSLTNYGVHWVRQSPGK GLEWLGVIWSSGNTDYNTPFT SRLSINKDNSKQVFFKMINSIQ SNDTAIYYCARALTYDYEFAY WGQGLTVTSAGGSGSGGG SGGGSGGDILLTQSPVILSVSP GERVSFSCRASQSIGTNIHWYQ QRTNGSPRLLIKYASESISGIPSR FSGSGSGTDFTLSINSVESEDIAD YYCQQNNNWPTTFGAGTKLEL K (SEQ ID NO: 15)	CAGGTGCAGCTGAAGCAGTCCGGCCCCGGCCTGGTGCAGCCCTCCCAGTC CCTGTCCATCACCTGCACCGTGTCCGGCTTCTCCTGACCAACTACGGCGT GCACTGGTGCAGTCCCGGCAAGGGCTGAGTGGTGGCGGTG ATCTGTCCGGCGGCAACACCGACTACAACACCCCTTACCTCCCGGCTG TCCATCAACAAGGACAACCTCAAGTCCAGGTGTTCTTCAAGATGAACCTC CTGCAGTCCAACGACACCGCCATCTACTGCGCCGGCCCTGACCTAC TAGACTACGAGTTCGCTACTGGGCCAGGGCACCTGGTGACCGTGTC CGCCGGGAGGTGAAGGGAGGGGAGGATCTGGCGGCGGAGGAA GCGGAGGCGACATCCTGCTGACCCAGTCCCGCTGATCCTGTCGCTGCC CCGGGAGCGGGTGTCTTCTCCTGCGGGCTCCAGTCCATCGGCACC AACATCCACTGTACAGCAGCGGACCAACGGTCCCCCGGCTGCTGAT CAAGTACGCTCCGAGTCACTCCGGCATCCCCTCCGGTTCCTCCGGCTC CGGCTCGGCACCGACTTACCTGTCCATCAACTCCGTGGAGTCCGAGG ACATGCCGACTACTGCGCAGCAGAAACAACACTGGCCACCACTTCG GCGCCGGCACCAAGCTGGAGTGAAG (SEQ ID NO: 145)
CT-487 EGFR (VH-VL scFv Format): QVQLVQSGAEVKKPGASVKVS CKASGYTFTSHWMHWVRQAP GQGLEWIGEFNPSNGRTNYNE KFKSKATMTVDSTNTAYMELS SLRSEDTAVYYCASRDYDGR YFDYWGQGLTVTVSSGGGSGG GGSGGGSGGDIQMTQSPSS LSASVGDRTITCSASSVTMY WYQQKPGKAPKLLIYDTSNLAS GVPSRFSGSGSDTYFTISSIQ PEDIATYYCQQWSSSHITFGQG TKVEIK (SEQ ID NO: 16)	CT-487 EGFR: CAGGTGCAGCTGGTGCAGTCCGGCGCCGAGGTGAAGAAGCCCGGCCT CCGTGAAGGTGTCTGCAAGGCTCCGGCTACACCTTCACTCCCACTGGA TGCACTGGGTGCGGCAGGCCCCCGGCGAGGCTGGAGTGGATCGGCGA GTTCAACCCCTCCAACGGCGGACCACTACAACGAGAAGTTCAAGTCCA AGGCCACCATGACCGTGGACACTCCACCAACACCGCTACATGGAGCTG TCTCCTCGGTCCGAGGACACCGCGTGTAAGTACTGCGCTCCCGGAC TACGACTACGACGGCGGTACTTCTGACTACTGGGCCAGGGCACCTGGT GACCGTGTCTCCGGCGGAGGTGAAGCGGAGGGGAGGATCTGGCGG CGGAGGAAGCGGAGGCGACATCCAGATGACCCAGTCCCCCTCCTCCCTGT CCGCTCCGTGGCGACCGGTGACCATCACTGCTCCGCCTCCTCCTCCG TGACCTACATGTAAGTACCAGCAGAAAGCCGCAAGGCCCAAGCTG CTGATCTACGACACCTCCAACCTGGCTCCGGCGTCCCTCCCGTTCTCC GGCTCCGGTCCGGCACCGACTACACCTTACCATCTCCTCCCTGACGCC GAGGACATCGCCACCTACTACTGCCAGCAGTGGTCTCCACACATCTTACC TTCGGCCAGGGCACCAAGGTGGAGATCAAG (SEQ ID NO: 146)
CT-488 EGFR (VH-VL scFv Format):	CT-488 EGFR:

(continued)

Amino Acid Sequence	DNA Sequence
QVQLQESGPGLVKPSETLSLTCT VSGSVSSGDYYWWTWIRQSPG KGLEWIGHIYYSGNTNYPNPSLKS RLTISIDTSKTQFSLKLSVTAAD TAIYYCVRDRV/TGAFDIW/GQGT MVTVSSGGGGSGGGSGGGG SGGDIQMTQSPSSLSASV/GDRV TITCQASQDISNLYLNWYQQKPG KAPKLLIYDASNLETGVP/PSRFSG SGSGTDFITISLQPEDIA/TYFC QHFDHLPLAFGGGKVEIK (SEQ ID NO: 17)	CAGGTGCAGTGCAGGAGTCCGGCCCCGGCCTGGTGAAGCCCTCCGAGA CCCTGTCCCTGACCTGCACCGTGTCCGGGGCTCCGTGCTCCTCCGGCGACT ACTACTGGACCTGGATCCGGCAGTCCCCGGCAAGGGCCTGGAAGTGGATC GGCCACATCTACTACTCGGGCAACACCACTACAACCCCTCCCTGAAGTCC CGGTGACCATCTCCATCGACACCTCCAAGACCCAGTTCTCCCTGAAGCTG TCCTCCGTACCGCCCGACACCGCCATCTACTACTGCGTGC GGACCG GGTAGCGGCGCTTCGACATCTGGGGCAGGGCACCACCATGGTGACCGTGT CCTCCGGCGAGGTGGAAGCGGAGGGGAGGATCTGGCGGGGAGGAA GCGGAGCGACATCCAGATGACCCAGTCCCGCTCCTCCCTGTC CGCTCCG TGGCGACCGGGTGACCATCACTGCCAGGCTCCAGGACATCTCCAAC TACCTGAACCTGGTACCAGAGAAGCCCGCAAGGCCCAAGCTGCTGAT CTACGACGCTCCAACCTGGAGACCGGGTGCCCTCCCGGTTCTCCGGCTC CGGTCCGGCACCGACTTCACCTTCACCATCTCCTCCCTGCAGCCCCGAGGA CATCGCCACCTACTTCTGCCAGCACTTCGACCACCTGCCCTGCGCTTCGGC GGCGGCACCAAGGTGGAGATCAAG (SEQ ID NO: 147)
CT-489 EGFR scFv: QVQLQESGPGLVKPSQTLSLTC TVSGSISGDYYWSWIRQPPG KGLEWIGIYYSGSTDYNPSLKS RVTMSVDTSKNQFSLKVNSTVA ADTAVYCARVSIFGVGTFDYW GQGTLVTVSSGGGGSGGGSG GGSGGEIVMTQSPATLSLSPG ERATLSCRASQSVSYLAWYQQ KPGQAPRLIYDASNRATGIPAR FSGSGGTDFTLTISLEPEDFAV YYCHQYGSTPLTFGGGTKAEIK (SEQ ID NO: 18)	CAGGTCCAGTCCAAGAGTCAGGACCAGGTCTGGTCAAAACCTCACAGAC GCTCTCTTACTTGACAGATACAGCGGATCCATTAGTTCGGTGATTAT TACTGGTCTTGGATCCGCCAGCCTCCTGTAAGGGGCTGGAGTGGATCGG ATACATATACTACAGTGGTTCACCTGATTATAATCCGTCACTCAAGTCCCGA GTAACATATGCCGTGGATACTAGCAAAAATCAGTTCAGCCTCAAAGTCAAT TCCGTAACGCCGCCGACACCGGGTCTATTATTGTGCGGGGTGAGCAT CTTCGGCTCGGTACGTTTCGATTACTGGGGACAAGAACTCTCGTAACAG TAAGCTCTGGCGAGGTGGAAGCGGAGGGGAGGATCTGGCGGGGAG GAAAGCGAGGGCGAGATTGTAATGACCCAAAGCCCTGCCACTCTTCTCTG TCTCCGGGGGAACGAGCGACTCTGTCTATGTAGGGCGTCCCAGAGTGCTC TAGTACCTCGCATGGTATCAGCAGAAACCCGGACAAGCCCCCTCGGCTCTT GATTACGATGCCTCAAACAGAGCAACGGGGAATCCGGCACGGTTCAGCG GGTCTGGTAGCGGGACAGACTTCACACTGACGATCTCTTCACTCGAACCA GAAGATTTTGAGTCTACTATTGCCATCAGTATGGTTCTACGCCACTTACAT TTGGCGGGGCCACCAAGGGCGGAGATAAAA (SEQ ID NO: 148)

[0129] In some embodiments, the activator ligand is EGFR or a peptide antigen thereof, and the activator ligand binding domain comprises an EGFR binding domain. In some embodiments, the EGFR ligand binding domain comprises an scFv domain. For example, the EGFR ligand binding domain may comprise a sequence of any one of SEQ ID NOs: 9-18. For example, the EGFR ligand binding domain may comprise a sequence at least 90%, at least 95% or at least 99% identical to any one of SEQ ID NOs: 9-18. In some embodiments, the EGFR ligand binding domain is encoded by a sequence selected from the group of sequences in Table 1. For example, the EGFR ligand binding domain is encoded by a sequence having at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity or at least 99% identity to a sequence from Table 1. It is understood that the scFv domains of the disclosure that bind EGFR can crossreact with one or more variants of EGFR. For example, an scFv may bind at least one of a normal EGFR, a variant of EGFR, or a mutated EGFR (e.g. EGFRvIII).

[0130] For example, the extracellular ligand binding domain of the first receptor comprises an scFv antigen binding domain having at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, at least 97% identity or at least 99% identity to any one of SEQ ID NOs: 9-18. As described herein, the extracellular ligand binding domain of the first receptor comprises an ScFv antigen binding domain comprising a sequence of any one of SEQ ID NOs: 9-18. As described herein, the extracellular ligand binding domain of the first receptor consists essentially of a sequence selected from the group consisting of SEQ ID NOs: 9-18. In one embodiment of the present invention, the extracellular ligand binding domain of the first receptor comprises the scFv sequence of SEQ ID NO: 18; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

Table 2. EGFR Variable Heavy (VH) and Variable Light (VL) domains

EGFR VH	EGFR VL
CT478, CT479: QVQLVESGGGVVQPGRLSLRSCAASGFTSTYGMHWVRQ APGKLEWVAVIWDGSKYKYGDSVKGRFTISRDNSKNTLY LQMNSLRAEDTAVYICARDGITMVRGVMDYFDYWGGG TLVTVSS (SEQ ID NO: 19) (not according to the invention)	CT478, CT479: AIQLTQSPSSLSASVGDRTTITCRASQDISSALVWY QQKPGKAPKLLIYDASSLESIGVPSRFSGESGTDFTL TISLQPEDFATYCCQFNQSYPLTFGGGKVEIK (SEQ ID NO: 25) (not according to the invention)
CT480, CT481: QIQLVQSGPELKKPGETVKISKASGYTFTEYPHWVKQAPG KGFKWMGMIYTDIGKPTYAEFEKGRFAFSLETSASTAYLQIN NLKNEDATYFCVDRDRYDSLFDYWGGGTTLVSS (SEQ ID NO: 20) (not according to the invention)	CT480, CT481: DVVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGN TYLHWYLQKPGQSPKLLIYKVSNRFSGVPDRFSGSG SGTDFTLKISRVEAEDLGVYFCSQSTHV/PWTFGGG TKLEIK (SEQ ID NO: 26) (not according to the invention)
CT482, CT483: EMQLVESGGGFVKPGGSLKLSCAASGFAFSHYDMSWVRQT PKQRLEWVAYIASGGDITYADTVKGRFTISRDNAQNTLYLQ MSSLKSEDTAMFYCSRSSYGNGDALDFWGGGTSVTVSS (SEQ ID NO: 21) (not according to the invention)	CT482, CT483: DVVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGN TYLHWYLQKPGQSPKLLIYKVSNRFSGVPDRFSGSG SGTDFTLKISRVEAEDLGVYFCSQSTHVLTFGSGTKL EIK (SEQ ID NO: 27) (not according to the invention)
CT486: QVQLKQSGPGLVQPQSLSITCTVSGFSLTNYGVHWVRQSP GKGLEWLGVIWSSGGNTDYNTPFTSRLSINKDNSKSQVFFKM NSLQSNDAIYICARALTYDYEFAYWGGGTLVTVA (SEQ ID NO: 22) (not according to the invention)	CT486: DILLTQSPVILSVSPGERVSFSCRASQSIGTNIHWYQ QRTNGSPRLLIYASESISGIPSRFSGSGGTDFTLSI NSVESEDIADYCCQNNNNWPTTFGAGTKLELK (SEQ ID NO: 28) (not according to the invention)
CT487:	CT487:

(continued)

EGFR VH	EGFR VL
QVQLVQSGAEVKKPGASVKVCKASGYTFTSHWMHWVRQ APGQGLEWIGEFNPSNGRNTYNEKFKSKATMTVDTSNTA YMESSLRSEDYAVYYCASRDYDYGRTFDYWGQGTLVTVS S (SEQ ID NO: 23) (not according to the invention)	DIQMTQSPSSLSASVGDRVTITCSASSSVTYMYWY QQKPGKAPKLLIYDTSNLASGVPSRFGSGSGTDYT FTISLQPEDIATYYCQQWSSHIFTFGGQTKVEIK (SEQ ID NO: 29) (not according to the invention)
CT488: QVQLQESGPGGLVKPSETLSLTCTVSGGSVSGDYWTWIRQ SPGKGLEWIGHIYYSGNTNYPNPSLKSRLTISIDTSKTQFSLKLS SVTAADTAIYYCVRDRVTGAFDIWGQGTMVTVSS (SEQ ID NO: 24) (not according to the invention)	CT488: DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNW YQKPGKAPKLLIYDASNLETGVPSRFGSGSGTDF TFTISLQPEDIATYFCQHFDHPLAFGGGQTKVEIK (SEQ ID NO: 30) (not according to the invention)
CT489: QVQLQESGPGGLVKPSQTLSTCTVSGGSISSGDYVWSWIRQ PPGKGLEWIGIYYSGSTDYNPSLKSRTMSVDTSKNQFSLK VNSVTAADTAIYYCARVSIFGVGTFDYWGQGTLVTVSS (SEQ ID NO: 149) (according to the invention)	CT489: EIVMTQSPATLSLSPGERATLSCRASQSVSSYLAWY QQKPGQAPRLLIYDASNRTATGIPARFGSGSGTDF TLTISLEPEDFAVYYCHQYGSTPLTFGGGQTKAEIK (SEQ ID NO: 150) (according to the invention)

[0131] In some embodiments, the activator ligand is EGFR or a peptide antigen thereof, and the activator ligand binding domain comprises an EGFR ligand binding domain. As described herein, the EGFR binding domain comprises a VH and/or a VL domain selected from the group disclosed in Table 2 or a sequence having at least 90% identity, at least 95% identity, at least 97% identity or at least 99% identity thereto. As described herein, the EGFR ligand binding domain comprises a VH domain selected from the group consisting of SEQ ID NOs: 19-24. As described herein, the EGFR ligand binding domain comprises a VH selected from the group consisting of SEQ ID NOs: 19-24 or a sequence having at least 90%, at least 95% or at least 99% identity thereto. As described herein, the EGFR ligand binding domain comprises a VL domain selected from the group consisting of SEQ ID NOs: 25-30. As described herein, the EGFR ligand binding domain comprises a VL selected from the group consisting of SEQ ID NOs: 25-30 or a sequence having at least 90%, at least 95% or at least 99% identity thereto.

Table 3. EGFR antigen binding domain CDRs.

CDR H1	CDR H2	CDR H3	CDR L1	CDR L2	CDR L3
TYGMH (SEQ ID NO: 31) (not according to the invention)	VIWDDGSYKYYG DSVKG (SEQ ID NO: 37) (not according to the invention)	DGITMVRGVMKDY FDY (SEQ ID NO: 43) (not according to the invention)	RASQDISSALV (SEQ ID NO: 49) (not according to the invention)	DASSLES (SEQ ID NO: 55) (not according to the invention)	QQFNSYPLT (SEQ ID NO: 60) (not according to the invention)
EYPIH (SEQ ID NO: 32) (not according to the invention)	MIYTDIGKPTYAE EFKG (SEQ ID NO: 38) (not according to the invention)	DRYDSLFDY (SEQ ID NO: 44) (not according to the invention)	RSSQSLVHSNGNT YLH (SEQ ID NO: 50) (not according to the invention)	KVSNRFS (SEQ ID NO: 56) (not according to the invention)	SQSTHV/PW T (SEQ ID NO: 61) (not according to the invention)
HYDMS (SEQ ID NO: 33) (not according to the invention)	YIASGGDITYYAD TVKG (SEQ ID NO: 39) (not according to the invention)	SSYG N NG DALDF (SEQ ID NO: 45) (not according to the invention)	RSSQSLVHSNGNT YLH (SEQ ID NO: 51) (not according to the invention)	KVSNRFS (SEQ ID NO: 56) (not according to the invention)	SQSTHVL (SEQ ID NO: 62) (not according to the invention)
NYGVH (SEQ ID NO: 34) (not according to the invention)	VIWSGGNTDYN TPFTS (SEQ ID NO: 40) (not according to the invention)	ALTYDYEFAY (SEQ ID NO: 46) (not according to the invention)	RASQSIGTNIH (SEQ ID NO: 52) (not according to the invention)	YASESIS (SEQ ID NO: 57) (not according to the invention)	QQNNNWP TT (SEQ ID NO: 63) (not according to the invention)
SHWMH (SEQ ID NO: 35) (not according to the invention)	EFNPSNGRTNYN EKFKS (SEQ ID NO: 41) (not according to the invention)	RDYDYDGRYFDY (SEQ ID NO: 47) (not according to the invention)	SASSSVTYMY (SEQ ID NO: 53) (not according to the invention)	DTSNLAS (SEQ ID NO: 58) (not according to the invention)	QQWSSHIFT (SEQ ID NO: 64) (not according to the invention)

(continued)

CDR H1	CDR H2	CDR H3	CDR L1	CDR L2	CDR L3
SGDYWWT (SEQ ID NO: 36) (not according to the invention)	HIYSGNTNYNP SLKS (SEQ ID NO: 42) (not according to the invention)	DRVTGAFDI (SEQ ID NO: 48) (not according to the invention)	QASQDISNYLN (SEQ ID NO: 54) (not according to the invention)	DASNLET (SEQ ID NO: 59) (not according to the invention)	QHFDHLPLA (SEQ ID NO: 65) (not according to the invention)
SGDYWWS (SEQ ID NO: 151) (according to the invention)	YIYSGSTDYNPS LKS (SEQ ID NO: 152) (according to the invention)	VSIFGVGTFDY (SEQ ID NO: 153) (according to the invention)	RASQSVSSYLA (SEQ ID NO: 154) (according to the invention)	DASNRAT (SEQ ID NO: 155) (according to the invention)	HQYGSTPLT (SEQ ID NO: 156) (according to the invention)

[0132] In some embodiments, the activator ligand is EGFR or a peptide antigen thereof, and the activator ligand binding domain is an EGFR ligand binding domain. As described herein, the EGFR binding domain comprises complementarity determining region (CDRs) selected from the group of CDRs disclosed in Table 3. As also described herein, the EGFR ligand binding domain comprises CDRs having at least 95% sequence identity to CDRs disclosed in Table 3. As also described herein, the EGFR ligand binding domain comprises CDRs selected from SEQ ID NOs: 31-65. As also described herein, the EGFR ligand binding domain comprises a heavy chain CDR 1 (CDR H1) selected from the group consisting of SEQ ID NOs: 31-36. As also described herein, the EGFR ligand binding domain comprises a heavy chain CDR 2 (CDR H2) selected from the group consisting of SEQ ID NOs: 37-42. As also described herein, the EGFR ligand binding domain comprises a heavy chain CDR 3 (CDR H3) selected from the group consisting of SEQ ID NOs: 43-48. As also described herein, the EGFR ligand binding domain comprises a light chain CDR 1 (CDR L1) selected from the group consisting of SEQ ID NOs: 49-54. As also described herein, the EGFR ligand binding domain comprises a light chain CDR 2 (CDR L2) selected from the group consisting of SEQ ID NOs: 55-59. As also described herein, the EGFR ligand binding domain comprises a light chain CDR 3 (CDR L3) selected from the group consisting of SEQ ID NOs: 60-65. As also described herein, the EGFR ligand binding domain comprises a CDR H1 selected from SEQ ID NOs: 31-36, a CDR H2 selected from SEQ ID NOs: 37-42, a CDR H3 selected from SEQ ID NOs: 43-48, a CDR L1 selected from SEQ ID NOs: 49-54, a CDR L2 selected from SEQ ID NOs: 55-59, and a CDR L3 selected from SEQ ID NOs: 60-65.

[0133] It is further described herein that one or more (e.g., 1, 2, 3, 4, 5, or 6) amino acid residues in a CDR of the antigen binding domains provided herein are substituted with another amino acid. The substitution may be "conservative" in the sense of being a substitution within the same family of amino acids. The naturally occurring amino acids may be divided into the following four families and conservative substitutions will take place within those families: (1) amino acids with basic side chains: lysine, arginine, histidine; (2) amino acids with acidic side chains: aspartic acid, glutamic acid; (3) amino acids with uncharged polar side chains: asparagine, glutamine, serine, threonine, tyrosine; and (4) amino acids with nonpolar side chains: glycine, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan, cysteine. By varying the amino acid sequence of the CDRs of an antibody by addition, deletion or substitution of amino acids, various effects such as increased binding affinity for the target antigen may be obtained.

Chimeric Antigen Receptors (CARs)

[0134] The disclosure provides a first, activator receptor and immune cells comprising same. In some embodiments of the present invention, the first receptor is a chimeric antigen receptor.

[0135] It is described herein that the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. It is also described herein that the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. In some embodiments of the present invention, the first receptor comprises SEQ ID NO: 177, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0136] The term "chimeric antigen receptors (CARs)" as used herein, may refer to artificial receptors derived from T-cell receptors and encompasses engineered receptors that graft an artificial specificity onto a particular immune effector cell. CARs may be employed to impart the specificity of a monoclonal antibody onto a T cell, thereby allowing a large number of specific T cells to be generated, for example, for use in adoptive cell therapy. In specific embodiments, CARs direct specificity of the cell to a tumor associated antigen, for example. Exemplary CARs comprise an intracellular activation domain, a transmembrane domain, and an extracellular domain comprising a tumor associated antigen binding region. In some embodiments, CARs further comprise a hinge domain. In particular aspects, CARs comprise fusions of single-chain variable fragments (scFv) derived from monoclonal antibodies, fused to a CD3 transmembrane domain and endodomain. The specificity of other CAR designs may be derived from ligands of receptors (e.g., peptides). In certain cases, CARs comprise domains for additional co-stimulatory signaling, such as CD3, 4-1BB, FcR, CD27, CD28, CD137, DAP10, and/or OX40. In some cases, molecules can be co-expressed with the CAR, including co-stimulatory molecules, reporter genes for imaging, gene products that conditionally ablate the T cells upon addition of a pro-drug, homing receptors, cytokines, and cytokine receptors.

[0137] In some embodiments, the extracellular ligand binding domain of the first receptor is fused to the extracellular domain of a CAR.

[0138] In some embodiments, the CARs of the present disclosure may comprise an extracellular hinge region. Incorporation of a hinge region can affect cytokine production from CAR-T cells and improve expansion of CAR-T cells *in vivo*. Exemplary hinges can be isolated or derived from IgD and CD8 domains, for example IgG1. For example, the hinge may be isolated or derived from CD8 α or CD28.

[0139] For example, the hinge may be isolated or derived from CD8 α or CD28. The CD8 α hinge may comprise an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of

TTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACD (SEQ ID NO: 66). The CD8 α hinge may comprise SEQ ID NO: 66. The CD8 α hinge may consist essentially of SEQ ID NO: 66. The CD8 α hinge may be encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of

ACCACGACGCCAGCGCCGCGACCAACACCGGCGCCACCATCGCGTCGCAGCCCCCTGTCCCTGCGCCCAGAGGC
GTGCCGCGCCAGCGGCGGGGGGCGCAGTGCACACGAGGGGGCTGGACTTCGCCTGTGAT (SEQ ID NO: 67).

The CD8 α hinge may be encoded by SEQ ID NO: 67.

[0140] The CD28 hinge comprises an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of CTIEVMYPPPYLDNEKSNGTIIHVKGKHLCP-SPLFPGPSKP (SEQ ID NO: 68). The CD28 hinge may comprise or consist essentially of SEQ ID NO: 68. The CD28 hinge may be encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of

TGTACCATTGAAGTTATGTATCCTCCTCCTTACCTAGACAATGAGAAGAGCAATGGAAC-
CATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCC (SEQ ID NO: 69). The CD28 hinge may be encoded by SEQ ID NO: 69.

[0141] The CARs of the present disclosure can be designed to comprise a transmembrane domain that is fused to the extracellular domain of the CAR. The transmembrane domain that naturally is associated with one of the domains in the CAR may be used. For example, a CAR comprising a CD28 co-stimulatory domain might also use a CD28 transmembrane domain. The transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins to minimize interactions with other members of the receptor complex.

[0142] The transmembrane domain may be derived either from a natural or from a synthetic source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein. Transmembrane regions may be isolated or derived from (i.e. comprise at least the transmembrane region(s) of) the alpha, beta or zeta chain of the T-cell receptor, CD28, CD3 epsilon, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154, or from an immunoglobulin such as IgG4. Alternatively, the transmembrane domain may be synthetic, in which case it will comprise predominantly hydrophobic residues such as leucine and valine. For example, a triplet of phenylalanine, tryptophan and valine will be found at each end of a synthetic transmembrane domain. Optionally, a short oligo- or polypeptide linker, preferably between 2 and 10 amino acids in length may form the linkage between the transmembrane domain and the cytoplasmic signaling domain of the CAR. A glycine-serine doublet provides a particularly suitable linker.

[0143] The CARs of the disclosure may comprise a CD28 transmembrane domain. For example, the CD28 transmembrane domain comprises an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of FWVLVWGGVLACYLLVTVAFIIFWV (SEQ ID NO: 70). As described herein, the CD28 transmembrane domain comprises or consists essentially of SEQ ID NO: 70. As also described herein, the CD28 transmembrane domain is encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of
TTCTGGGTGCTGGTCTGTGGGCGGCGTGCTGGCCTGCTACAGCCTGCTGGTGACAGT-
GGCCTTCATCATCTTTTGGTG (SEQ ID NO: 71). In some embodiments, the CD28 transmembrane domain is encoded by SEQ ID NO: 71.

[0144] The CARs of the disclosure may comprise an IL-2Rbeta transmembrane domain. For example, the IL-2Rbeta transmembrane domain may comprise an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of IPWLGHLLVGLSGAFGIILVYLLI (SEQ ID NO: 72). For example, the IL-2Rbeta transmembrane domain may comprise or consist essentially of SEQ ID NO: 72. For example, the IL-2Rbeta transmembrane domain may be encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of
ATTCCGTGGC TCGGCCACCT CTCGTGGGC CTCAGCGGGG CTTTGGCTT CATCATCTTA GTGTACTTGC
TGATC (SEQ ID NO: 73). For example, the IL-2Rbeta transmembrane domain may be encoded by SEQ ID NO: 73.

[0145] The cytoplasmic domain or otherwise the intracellular signaling domain of the CARs of the instant disclosure is responsible for activation of at least one of the normal effector functions of the immune cell in which the CAR has been placed. The term "effector function" refers to a specialized function of a cell. Thus the term "intracellular signaling domain" refers to the portion of a protein, which transduces the effector function signal and directs the cell to perform a specialized function. While usually the entire intracellular signaling domain can be employed, in many cases it is not necessary to use the entire domain. To the extent that a truncated portion of the intracellular signaling domain is used, such truncated portion may be used in place of the intact chain as long as it transduces the effector function signal. In

some cases, multiple intracellular domains can be combined to achieve the desired functions of the CAR-T cells of the instant disclosure. The term intracellular signaling domain is thus meant to include any truncated portion of one or more intracellular signaling domains sufficient to transduce the effector function signal.

[0146] Examples of intracellular signaling domains for use in the CARs of the instant disclosure include the cytoplasmic sequences of the T cell receptor (TCR) and co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any derivative or variant of these sequences and any synthetic sequence that has the same functional capability.

[0147] Accordingly, the intracellular domain of CARs of the instant disclosure comprises at least one cytoplasmic activation domain. As described herein, the intracellular activation domain ensures that there is T-cell receptor (TCR) signaling necessary to activate the effector functions of the CAR T-cell. For example, the at least one cytoplasmic activation is a CD247 molecule (CD3 ζ) activation domain, a stimulatory killer immunoglobulin-like receptor (KIR) KIR2DS2 activation domain, or a DNAX-activating protein of 12 kDa (DAP12) activation domain.

[0148] For example, the CD3 ζ activation domain may comprise an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of

RVKFSRSADAPAYKQGQNQLYNELNLGRREEYDVLDRGRDPEMGGKPRRKNPQEG
LYNELQKDKMAEAYSEIGMKGERRRGKGDGLYQGLSTATKDTYDALHMQALPPR (SEQ ID NO: 74). For example, the CD3 ζ activation domain may comprise or consist essentially of SEQ ID NO: 74. For example, the CD3 ζ activation domain may be encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of

AGAGTGAAGTTCAGCAGGAGCGCAGACGCCCGCGTACAAGCAGGGCCAGAACCAGCTCTATAACGAGCTCA
ATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGCGTAGAGGCCGGGACCCTGAGATGGGGGGAAAGCC
GAGAAGGAAGAACCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAG-
GCCTACAGTGAGATTGGGATGAAA GGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGACT-
CAGTACAGCCACCAAGGACACCTACGACGC CCTTCACATGCAGGCCCTGCCCCCTCGC (SEQ ID NO: 75). As de-
scribed herein, the CD3 ζ activation domain may be encoded by SEQ ID NO: 75.

[0149] It is known that signals generated through the TCR alone are often insufficient for full activation of the T cell and that a secondary or co-stimulatory signal is also required. Thus, T cell activation can be said to be mediated by two distinct classes of cytoplasmic signaling sequence: those that initiate antigen-dependent primary activation through the TCR (primary cytoplasmic signaling sequences) and those that act in an antigen-independent manner to provide a secondary or co-stimulatory signal (secondary cytoplasmic signaling sequences).

[0150] Primary cytoplasmic signaling sequences regulate primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary cytoplasmic signaling sequences that act in a stimulatory manner may contain signaling motifs, which are known as immunoreceptor tyrosine-based activation motifs or ITAMs. As described herein, the ITAM may contain a tyrosine separated from a leucine or an isoleucine by any two other amino acids (YxxL/I) (SEQ ID NO: 157). As also described herein, the cytoplasmic domain may contain 1, 2, 3, 4 or 5 ITAMs. An exemplary ITAM containing cytoplasmic domain is the CD3 ζ activation domain. Further examples of ITAM containing primary cytoplasmic signaling sequences that can be used in the CARs of the instant disclosure may include those derived from TCR ζ , FcR γ , FcR β , CD3 γ , CD3 δ , CD3 ϵ , CD3 ζ , CD5, CD22, CD79a, CD79b, and CD66d.

[0151] For example, the CD3 ζ activation domain comprising a single ITAM may comprise an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of

RVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLHMQALPPR (SEQ ID NO: 76). For example, the CD3 ζ activation domain may comprise SEQ ID NO: 76. For example, the CD3 ζ activation domain comprising a single ITAM may consist essentially of an amino acid sequence of RVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLHMQALPPR (SEQ ID NO: 76). For example, the CD3 ζ activation domain comprising a single ITAM may be encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or may be identical to a sequence of

AGAGTGAAGT TCAGCAGGAG CGCAGACGCC CCCGCGTACC AGCAGGGCCA GAACCAGCTC TATAACGAGC
TCAATCTAGG ACGAAGAGAG GAGTACGATG TTTTGCACAT GCAGGCCCTG CCCCCTCGC (SEQ ID NO: 77). As
described herein, the CD3 ζ activation domain may be encoded by SEQ ID NO: 77.

[0152] The cytoplasmic domain of the CAR can be designed to comprise the CD3 ζ signaling domain by itself or combined with any other desired cytoplasmic domain(s) useful in the context of the CAR of the instant disclosure. For example, the cytoplasmic domain of the CAR can comprise a CD3 ζ chain portion and a co-stimulatory domain. The co-stimulatory domain refers to a portion of the CAR comprising the intracellular domain of a costimulatory molecule. A costimulatory molecule is a cell surface molecule other than an antigen receptor or its ligands that is required for an efficient response of lymphocytes to an antigen. Examples of such molecules include the co-stimulatory domain is selected from the group consisting of IL-2R β , Fc Receptor gamma (FcR γ), Fc Receptor beta (FcR β), CD3g molecule gamma (CD3 γ), CD3 δ , CD3 ϵ , CD5 molecule (CD5), CD22 molecule (CD22), CD79a molecule (CD79a), CD79b molecule

(CD79b), carcinoembryonic antigen related cell adhesion molecule 3 (CD66d), CD27 molecule (CD27), CD28 molecule (CD28), TNF receptor superfamily member 9 (4-1BB), TNF receptor superfamily member 4 (OX40), TNF receptor superfamily member 8 (CD30), CD40 molecule (CD40), programmed cell death 1 (PD-1), inducible T cell costimulatory (ICOS), lymphocyte function-associated antigen-1 (LFA-1), CD2 molecule (CD2), CD7 molecule (CD7), TNF superfamily member 14 (LIGHT), killer cell lectin like receptor C2 (NKG2C) and CD276 molecule (B7-H3) c-stimulatory domains, or functional fragments thereof. The intracellular domains of CARs of the instant disclosure comprise at least one co-stimulatory domain. For example, the co-stimulatory domain is isolated or derived from CD28.

[0153] The intracellular domains of CARs of the instant disclosure comprise at least one costimulatory domain. For example, the co-stimulatory domain is isolated or derived from CD28. For example, the CD28 co-stimulatory domain comprises an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of

RSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAYRS (SEQ ID NO: 78). For example, the CD28 co-stimulatory domain comprises or consists essentially of SEQ ID NO: 78. For example, the CD28 co-stimulatory domain is encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of

AGGAGCAAGCGGAGCAGACTGCTGCACACGCGACTACATGAACATGACCCCCCGGAG
GCCTGGCCCCACCCGGAAGCACTACCAGCCCTACGCCCTCCCAGGGATTTCGCCGC CTACCGGAGC (SEQ ID NO: 79). As described herein, the CD28 co-stimulatory domain is encoded by SEQ ID NO: 79. The cytoplasmic domains within the cytoplasmic signaling portion of the CARs of the instant disclosure may be linked to each other in a random or specified order. Optionally, a short oligo- or polypeptide linker, for example between 2 and 10 amino acids in length may form the linkage. A glycine-serine doublet provides an example of a suitable linker.

[0154] For example, the co-stimulatory domain is isolated or derived from 4-1BB. For example, the 4-1BB co-stimulatory domain comprises an amino acid sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of KRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL (SEQ ID NO: 159). As described herein, the 4-1BB co-stimulatory domain comprises or consists essentially of KRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL (SEQ ID NO: 159). For example, the 4-1BB co-stimulatory domain is encoded by a nucleotide sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity or is identical to a sequence of

aaacggggcagaagaactcctgtatatattcaacaaccatttatgaggccagtacaaactactcaagaggaagatggctgtagctgcc
gattccagaagaagaaggaggatgtgaactg (SEQ ID NO: 160).

[0155] The intracellular domain of the CAR may comprise a CD28 co-stimulatory domain, a 4-1BB costimulatory domain, and a CD3 ζ activation domain. For example, the intracellular domain of the CAR may comprise a sequence of RSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAYRSKRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCELRVKFSRSADAPAYKQGQNQLYNELNLGRREEYDV LDKRRGRDPEMGGKPRRKNPQEGLYNELQDKMAEAYSEIGMKGERRRGKGHDGLY QGLSTATKDTYDALHMQALPPR (SEQ ID NO: 158), or a sequence having at least 80% identity, at least 90% identity, at least 95% identity, at least 99% identity thereto.

[0156] The cytoplasmic domains within the cytoplasmic signaling portion of the CARs of the instant disclosure may be linked to each other in a random or specified order. Optionally, a short oligo- or polypeptide linker, for example between 2 and 10 amino acids in length may form the linkage. A glycine-serine doublet provides an example of a suitable linker. An exemplary linker comprises a sequence of GGGGSGGGGSGGGSGG (SEQ ID NO: 136).

[0157] The cytoplasmic domains within the cytoplasmic signaling portion of the CARs of the instant disclosure may be linked to each other in a random or specified order. Optionally, a short oligo- or polypeptide linker, for example between 2 and 10 amino acids in length may form the linkage. A glycine-serine doublet provides an example of a suitable linker.

T Cell Receptors (TCRs)

[0158] The disclosure provides a first, activator receptor and immune cells comprising same. The first receptor may be a T cell receptor (TCR).

[0159] Exemplary TCRs comprising intracellular domains for use in the instant disclosure are described in PCT/US2020/045250 filed on September 6, 2020.

[0160] As used herein, a "TCR", sometimes also called a "TCR complex" or "TCR/CD3 complex" refers to a protein complex comprising a TCR alpha chain, a TCR beta chain, and one or more of the invariant CD3 chains (zeta, gamma, delta and epsilon), sometimes referred to as subunits. The TCR alpha and beta chains can be disulfide-linked to function

as a heterodimer to bind to peptide-MHC complexes. Once the TCR alpha/beta heterodimer engages peptide-MHC, conformational changes in the TCR complex in the associated invariant CD3 subunits are induced, which leads to their phosphorylation and association with downstream proteins, thereby transducing a primary stimulatory signal. In an exemplary TCR complex, the TCR alpha and TCR beta polypeptides form a heterodimer, CD3 epsilon and CD3 delta form a heterodimer, CD3 epsilon and CD3 gamma form a heterodimer, and two CD3 zeta form a homodimer.

[0161] Any suitable ligand binding domain may be fused to an extracellular domain, hinge domain or transmembrane of the TCRs described herein. For example, the ligand binding domain can be an antigen binding domain of an antibody or TCR, or comprise an antibody fragment, a V β only domain, a linear antibody, a single-chain variable fragment (scFv), or a single domain antibody (sdAb).

[0162] The ligand binding domain may be fused to one or more extracellular domains or transmembrane domains of one or more TCR subunits. The TCR subunit can be TCR alpha, TCR beta, CD3 delta, CD3 epsilon, CD3 gamma or CD3 zeta. For example, the ligand binding domain can be fused to TCR alpha, or TCR beta, or portions of the ligand binding can be fused to two subunits, for example portions of the ligand binding domain can be fused to both TCR alpha and TCR beta.

[0163] TCR subunits include TCR alpha, TCR beta, CD3 zeta, CD3 delta, CD3 gamma and CD3 epsilon. Any one or more of TCR alpha, TCR beta chain, CD3 gamma, CD3 delta, CD3 epsilon, or CD3 zeta, or fragments or derivative thereof, can be fused to one or more domains capable of providing a stimulatory signal of the disclosure, thereby enhancing TCR function and activity.

[0164] TCR transmembrane domains isolated or derived from any source are envisaged as within the scope of the disclosure. The transmembrane domain may be derived either from a natural or from a recombinant source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein.

[0165] The transmembrane domain may be capable of signaling to the intracellular domain(s) whenever the TCR complex has bound to a target. A transmembrane domain of particular use in this disclosure may include at least the transmembrane region(s) of e.g., the alpha, beta or zeta chain of the TCR, CD3 delta, CD3 epsilon or CD3 gamma, CD28, CD3 epsilon, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154.

[0166] The transmembrane domain can be attached to the extracellular region of a polypeptide of the TCR, e.g., the antigen binding domain of the TCR alpha or beta chain, via a hinge, e.g., a hinge from a human protein. For example, the hinge can be a human immunoglobulin (Ig) hinge, e.g., an IgG4 hinge, or a CD8a hinge. The hinge may be isolated or derived from CD8 α or CD28.

[0167] The extracellular ligand binding domain may be attached to one or more transmembrane domains of the TCR. The transmembrane domain may comprise a TCR alpha transmembrane domain, a TCR beta transmembrane domain, or both. The transmembrane may comprise a CD3 zeta transmembrane domain.

[0168] A transmembrane domain can include one or more additional amino acids adjacent to the transmembrane region, e.g., one or more amino acid associated with the extracellular region of the protein from which the transmembrane was derived (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or up to 15 amino acids of the extracellular region) and/or one or more additional amino acids associated with the intracellular region of the protein from which the transmembrane protein is derived (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or up to 15 amino acids of the intracellular region).

[0169] The transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins, e.g., to minimize interactions with other members of the receptor complex.

[0170] When present, the transmembrane domain may be a natural TCR transmembrane domain, a natural transmembrane domain from a heterologous membrane protein, or an artificial transmembrane domain. The transmembrane domain may be a membrane anchor domain. Without limitation, a natural or artificial transmembrane domain may comprise a hydrophobic α -helix of about 20 amino acids, often with positive charges flanking the transmembrane segment. The transmembrane domain may have one transmembrane segment or more than one transmembrane segment. Prediction of transmembrane domains/segments may be made using publicly available prediction tools (e.g. TMHMM, Krogh et al. Journal of Molecular Biology 2001; 305(3):567-580; or TMPred, Hofmann & Stoffel Biol. Chem. Hoppe-Seyler 1993; 347: 166). Non-limiting examples of membrane anchor systems include platelet derived growth factor receptor (PDGFR) transmembrane domain, glycosylphosphatidylinositol (GPI) anchor (added post-translationally to a signal sequence) and the like.

[0171] The transmembrane domain may comprise a TCR alpha transmembrane domain. The TCR alpha transmembrane domain may comprise an amino acid sequence having at least 85% identity, at least 90% identity, at least 95% identity, at least 96% identity, at least 97% identity, at least 98% identity, at least 99% identity or may be identical to a sequence of: VIGFRILLKLVAGFNLLMTLRLW (SEQ ID NO: 80). The TCR alpha transmembrane domain may comprise, or consist essentially of, SEQ ID NO: 80. The TCR alpha transmembrane domain may be encoded by a sequence of

GTGATTGGGTTC CGAATCCTCCTCCTGAAAGTGGCCGGGTTTAATCTGCTCATGACG
CTGCGGCTGTGG (SEQ ID NO: 81).

5

[0172] The transmembrane domain may comprise a TCR beta transmembrane domain. The TCR beta transmembrane domain may comprise an amino acid sequence having at least 85% identity, at least 90% identity, at least 95% identity, at least 96% identity, at least 97% identity, at least 98% identity, at least 99% identity or may be identical to a sequence of: TILYEILLGKATLYAVLVLSALVL (SEQ ID NO : 82). The TCR beta transmembrane domain may comprise, or consist essentially of, SEQ ID NO: 82. The TCR beta transmembrane domain may be encoded by a sequence of ACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCTTGATGCCGTGCTGGTCAGTGCCCTCGTGCTG (SEQ ID NO: 83).

10

[0173] TCRs of the disclosure can comprise one or more intracellular domains. The intracellular domain may comprise one or more domains capable of providing a stimulatory signal to a transmembrane domain. The intracellular domain may comprise a first intracellular domain capable of providing a stimulatory signal and a second intracellular domain capable of providing a stimulatory signal. The intracellular domain may comprise a first, second and third intracellular domain capable of providing a stimulatory signal. The intracellular domains capable of providing a stimulatory signal may be selected from the group consisting of a CD28 molecule (CD28) domain, a LCK proto-oncogene, Src family tyrosine kinase (Lck) domain, a TNF receptor superfamily member 9 (4-1BB) domain, a TNF receptor superfamily member 18 (GITR) domain, a CD4 molecule (CD4) domain, a CD8a molecule (CD8a) domain, a FYN proto-oncogene, Src family tyrosine kinase (Fyn) domain, a zeta chain of T cell receptor associated protein kinase 70 (ZAP70) domain, a linker for activation of T cells (LAT) domain, lymphocyte cytosolic protein 2 (SLP76) domain, (TCR) alpha, TCR beta, CD3 delta, CD3 gamma and CD3 epsilon intracellular domains.

15

20

[0174] An intracellular domain may comprise at least one intracellular signaling domain. An intracellular signaling domain generates a signal that promotes a function a cell, for example an immune effector function of a TCR containing cell, e.g., a TCR-expressing T-cell. The intracellular domain of the first receptor of the disclosure may include at least one intracellular signaling domain. For example, the intracellular domains of CD3 gamma, delta or epsilon may comprise signaling domains.

25

[0175] The extracellular domain, transmembrane domain and intracellular domain may be isolated or derived from the same protein, for example T-cell receptor (TCR) alpha, TCR beta, CD3 delta, CD3 gamma, CD3 epsilon or CD3 zeta.

30

[0176] Examples of intracellular domains for use in activator receptors of the disclosure may include the cytoplasmic sequences of the TCR alpha, TCR beta, CD3 zeta, and 4-1BB, and the intracellular signaling co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any derivative or variant of these sequences and any recombinant sequence that has the same functional capability.

35

[0177] The intracellular signaling domain may comprise a primary intracellular signaling domain. Exemplary primary intracellular signaling domains may include those derived from the proteins responsible for primary stimulation, or antigen dependent stimulation.

[0178] The intracellular domain may comprise a CD3 delta intracellular domain, a CD3 epsilon intracellular domain, a CD3 gamma intracellular domain, a CD3 zeta intracellular domain, a TCR alpha intracellular domain or a TCR beta intracellular domain.

40

[0179] The intracellular domain may comprise a TCR alpha intracellular domain. A TCR alpha intracellular domain may comprise Ser-Ser. A TCR alpha intracellular domain may be encoded by a sequence of TCCAGC.

[0180] The intracellular domain may comprise a TCR beta intracellular domain. The TCR beta intracellular domain may comprise an amino acid sequence having at least 80% identity, at least 90% identity, or may be identical to a sequence of: MAMVKRKDSR (SEQ ID NO: 84). The TCR beta intracellular domain may comprise, or consist essentially of SEQ ID NO: 84. The TCR beta intracellular domain may be encoded by a sequence of ATGGCCATGGTCAAGAGAAAGGATTCCAGA (SEQ ID NO: 85).

45

[0181] The intracellular signaling domain may comprise at least one stimulatory intracellular domain. The intracellular signaling domain may comprise a primary intracellular signaling domain, such as a CD3 delta, CD3 gamma and CD3 epsilon intracellular domain, and one additional stimulatory intracellular domain, for example a co-stimulatory domain. The intracellular signaling domain may comprise a primary intracellular signaling domain, such as a CD3 delta, CD3 gamma and CD3 epsilon intracellular domain, and two additional stimulatory intracellular domains.

50

[0182] Exemplary co-stimulatory intracellular signaling domains include those derived from proteins responsible for co-stimulatory signals, or antigen independent stimulation. Co-stimulatory molecules include an MHC class I molecule, BTLA, a Toll ligand receptor, as well as DAP10, DAP12, CD30, LIGHT, OX40, CD2, CD27, CDS, ICAM-1, LFA-1 (CD11a/CD18) 4-1BB (CD137, TNF receptor superfamily member 9), and CD28 molecule (CD28). A co-stimulatory protein can be represented in the following protein families: TNF receptor proteins, Immunoglobulin-like proteins, cytokine receptors, integrins, signaling lymphocytic activation molecules (SLAM proteins), and activating NK cell receptors. Ex-

55

amples of such molecules include CD27, CD28, 4-1BB (CD137), OX40, GITR, CD30, CD40, ICOS, BAFFR, HVEM, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3, a ligand that specifically binds with CD83, CD4, and the like. The co-stimulatory domain can comprise the entire intracellular portion, or the entire native intracellular signaling domain, of the molecule from which it is derived, or a functional variant thereof.

[0183] The stimulatory domain may comprise a co-stimulatory domain. The co-stimulatory domain may comprise a CD28 or 4-1BB co-stimulatory domain. CD28 and 4-1BB are well characterized co-stimulatory molecules required for full T cell activation and known to enhance T cell effector function. For example, CD28 and 4-1BB have been utilized in chimeric antigen receptors (CARs) to boost cytokine release, cytolytic function, and persistence over the first generation CAR containing only the CD3 zeta signaling domain. Likewise, inclusion of co-stimulatory domains, for example CD28 and 4-1BB domains, in TCRs can increase T cell effector function and specifically allow co-stimulation in the absence of co-stimulatory ligand, which is typically down-regulated on the surface of tumor cells. As also described herein, the stimulatory domain comprises a CD28 intracellular domain or a 4-1BB intracellular domain.

Inhibitory Receptors

[0184] The disclosure provides a second receptor, comprising an extracellular ligand binding domain specific to a non-target antigen that has been lost in a cancer cell, such as an allelic variant of a gene. The non-target allelic variant can be lost in the cancer cell through any mechanism, such as, without limitation, epigenetic changes that effect non-target allelic variant expression, mutations to the gene encoding the non-target allelic variant, disruption of cellular signaling that regulates expression of the non-target allelic variant, chromosome loss, partial or complete deletion of the genomic locus, gene silencing through modification of nucleic acids or heterochromatin, or loss of expression through other mechanisms. In variations of the compositions and methods disclosed herein, the cells or subject treated may exhibit a loss of expression of the non-target allelic variant because of non-genetic changes. Accordingly the disclosure provides compositions and methods for killing cells and/or treating subject lacking expression of the non-target antigen from any cause, including but not limited to, loss of heterozygosity.

[0185] The non-target antigen can be a protein, or an antigen peptide thereof in a complex with a major histocompatibility complex class I (MHC-I), where the non-target antigen comprises a polymorphism. Because the non-target antigen is polymorphic, loss of a single copy of the gene encoding the non-target antigen, which may occur through loss of heterozygosity in a cancer cell, yields a cancer cell that retains the other polymorphic variant of gene, but has lost the non-target antigen. For example, a subject having HLA-A*02 and HLA-A*01 alleles at the HLA locus may have a cancer in which only the HLA-A*02 allele is lost. In such a subject, the HLA-A*01 protein remains present, but is not recognized by the inhibitory receptor of immune cells encountering the cancer cell, because the inhibitor receptor is designed to be specific to the HLA-A*02 (or other non-target antigen). In normal non-malignant cells, the HLA-A*02 (or other non-target antigen) is present and inhibits activation of the engineered immune cell. In cancer cells having loss of heterozygosity, the HLA-A*02 allelic variant (or other non-target antigen) is lost. Immune cells engineered to express the inhibitory receptor do not receive an inhibitory signal from the inhibitory receptor, as the inhibitory receptor only responds to the HLA-A*02 (or other non-target antigen), which is absent on cancer cells. By this mechanism, the immune cell is selectively activated, and selectively kills, cancer cells expressing EGFR, but having lost HLA-A*02 (or another non-target antigen) due to loss-of-heterozygosity. HLA-A is used here as an example. Similar polymorphic variation occurs in the population at other MHC genes and in other non-MHC genes as well. Accordingly, the non-target antigen may comprise a polymorphic variant of COLEC12, APCDD1 or CXCL16, or HLA-A*02. The non-target antigen may be an HLA class I allele or a minor histocompatibility antigen (MiHA). The HLA Class I allele may comprise HLA-A, HLA-B, HLA-C, or HLA-E. In some embodiments, the HLA class I allele is HLA-A*02. In some embodiments, the HLA-A*02 non-target antigen is expressed by healthy cells of a subject. In some embodiments, the non-target antigen is a non-target allelic variant. In some embodiments, the non-target antigen is not expressed in a cancer cell of the subject. In some embodiments, the non-target antigen is not expressed in a fraction of the cells in a tumor in the subject. In some embodiments, a cancer cell in a subject has lost expression of the non-target antigen. Loss of expression or lack of expression of the non-target antigen in a cell can be by any mechanism, such as, epigenetic changes that effect non-target gene expression, mutations to the gene encoding the non-target antigen, or disruption of cellular signaling that regulates expression of the non-target gene.

[0186] In some embodiments, the second receptor is an inhibitory chimeric antigen receptor (*i.e.* inhibitory receptor). In some embodiments, the second receptor is an inhibitory receptor. In some embodiments, the second receptor is humanized.

[0187] As described herein, the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As described herein, the second receptor comprises SEQ ID NO: 174 or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0188] The disclosure provides a second receptor, which is an inhibitory receptor, comprising an extracellular ligand binding that can discriminate between single amino-acid variant alleles of a non-target antigen. This ability to discriminate between allelic variants of a non-target antigen allows the second receptor to inhibit activation of immune cells comprising the second receptor in the presence of non-target cells that express that the allele recognized by the ligand binding domain. However, activation of immune cells is not inhibited in the presence of target cells that have lost the allele, for example cancer cells that have lost one allele of a gene through loss of heterozygosity.

[0189] The disclosure provides a second receptor, which is an inhibitory receptor, comprising an extracellular ligand binding that can discriminate between different levels of expression of a non-target antigen. This allows the second receptor to inhibit activation of immune cells comprising the second receptor in the presence of non-target cells that express the ligand for the second receptor, but to allow activation of immune cells in the presence of cancer cells that express low levels, or have no expression, of the ligand for the second receptor.

Inhibitor Ligands

[0190] In some embodiments, the non-target antigen is not expressed by the target cells, and is expressed by non-target cells. In some embodiments, the non-target antigen is expressed by healthy cells, i.e. cells that are not cancer cells. In some embodiments, the target cells are a plurality of cancer cells that have lost expression of the non-target antigen through loss of heterozygosity (LOH). In some embodiments, the non-target cells are a plurality of healthy cells (i.e., non-cancer, normal, or healthy cells), that express both the target and the non-target antigen.

[0191] Any cell surface molecule expressed by the non-target cells that is not expressed by target cells may be a suitable non-target antigen for the second receptor extracellular ligand binding domain. For example, a cell adhesion molecule, a cell-cell signaling molecule, an extracellular domain, a molecule involved in chemotaxis, a glycoprotein, a G protein-coupled receptor, a transmembrane, a receptor for a neurotransmitter or a voltage gated ion channel can be used as a non-target antigen.

[0192] In some embodiments, the target antigen is a peptide antigen of a cancer cell-specific antigen in a complex with a major histocompatibility complex class I (MHC-I).

[0193] In some embodiments, the non-target antigen is lost in the cancer cells due to loss of heterozygosity. Exemplary non-target antigens lost in cancer cells due to loss of heterozygosity include COLEC12, APCDD1, CXCL16 and HLA-A*02. As described herein, the non-target antigen is selected from the group consisting of a polymorphic variant of COLEC12, APCDD1 and CXCL166, or a peptide antigen thereof a complex with a major histocompatibility complex class I (MHC-I), or HLA-A*02. As described herein, the non-target antigen is an antigen peptide comprising a polymorphic residue of COLEC12, APCDD1 and CXCL16 in a complex with a major histocompatibility complex class I (MHC-I).

[0194] Non-target MHC-1 (MHC) antigens comprising any of HLA-A, HLA-B or HLA-C are described herein (not according to the invention). The non-target antigen may comprise HLA-A (not according to the invention). The non-target antigen may comprise a human leukocyte antigen A*02 allelic product (HLA-A*02) (not according to the invention). As described herein, but not according to the invention, the non-target antigen comprises human leukocyte antigen A*69. As also described herein, but not according to the invention, the non-target antigen comprises HLA-B. As also described herein, but not according to the invention, the non-target antigen comprises HLA-C.

[0195] In some embodiments, the non-target antigen comprises HLA-A*02.

[0196] As described herein, but not according to the invention, the non-target antigen comprises C-X-C motif chemokine ligand 16 (CXCL16) or an antigen peptide thereof in a complex with MHC-I. Human CXCL16 precursor is described in NCBI record number NP_001094282.1. As described herein, CXCL16 comprises an amino acid sequence of:

```

1 MSGSQSEVAP SPQSPRSPEM GRDLRPGSRV LLLLLLLLLLV YLTQPGNGNE GSVTGSCYCG
61 KRISSDSPPS VQFMNRLRKH LRAYHRCLEY TRFQLLSWSV CGGNKDPWVQ ELMSCLDLKE
121 CGHAYSGIVA HQKHLLPTSP PISQASEGAS SDIHTPAQML LSTLQSTQRP TLPVGSLSDD
181 KELTRPNETT IHTAGHSLAA GPEAGENQKQ PEKNAGPTAR TSATVPVLCL LAIIFILTAA
241 LSYVLCKRRR GQSPQSSPDL PVHYIPVAPD SNT (SEQ ID NO: 86).

```

[0197] As described herein, but not according to the invention, the non-target antigen comprises a polymorphism of CXCL16. For example, the non-target antigen comprises a peptide derived from CXCL16 comprising a polymorphic residue of CXCL16. Polymorphic residues of CXCL16 include positions 142 and 200 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising amino acid 142 or 200 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising an A at amino acid 200 of SEQ ID NO: 86. As also described herein, but

not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising a V at amino acid 200 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising an I at amino acid 142 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising a T at amino acid 142 of SEQ ID NO: 86.

[0198] As also described herein, but not according to the invention, the non-target antigen comprises a polymorphism of CXCL16. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising an A at amino acid 200 of SEQ ID NO: 86, and the second receptor comprises a ligand binding domain with a higher affinity for a CXCL16 ligand with an A at position 200 of SEQ ID NO: 86 than for a CXCL16 ligand with a V at position 200 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising a V at amino acid 200 of SEQ ID NO: 86, and the second receptor comprises a ligand binding domain with a higher affinity for a CXCL16 ligand with a V at position 200 of SEQ ID NO: 86 than for a CXCL16 ligand with an A at position 200 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising an I at amino acid 142 of SEQ ID NO: 86, and the second receptor comprises a ligand binding domain with a higher affinity for a CXCL16 ligand with an I at position 142 of SEQ ID NO: 86 than for a CXCL16 ligand with a T at position 142 of SEQ ID NO: 86. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of CXCL16 comprising a T at amino acid 142 of SEQ ID NO: 86, and the second receptor comprises a ligand binding domain with a higher affinity for a CXCL16 ligand with a T at position 142 of SEQ ID NO: 86 than for a CXCL16 ligand with an I at position 142 of SEQ ID NO: 86.

[0199] As described herein, but not according to the invention, the non-target antigen comprises collectin subfamily member 12 (COLEC12) or an antigen peptide thereof in a complex with MHC-I. Human COLEC12 is described in NCBI record number NP_569057.2. As also described herein, but not according to the invention, COLEC12 comprises an amino acid sequence of:

```

1  MKDDFAEEEE VQSFQYKRFQ IQEGTQCTKC KNNWALKFSI ILLYILCALL TITVAILGYK
61  VVEKMDNVTG GMETSRQTYD DKLTAVESDL KKLGDQTGKK AISTNSELSL FRSDILDRLQ
121 QLRTEIKTS KNKDTLEKLQ ASGDALVDRQ SQLKETLENN SFLITTVNKT LQAYNGYVTN
181 LQQDTSVLQG NLQNQMYSHN VVIMNLLNLLN LTQVQQRNLI TNLQRSVDDT SQAIQRIKND
241 FQNLQQVFLQ AKKDTDWLKE KVQSLQTLAA NNSALAKANN DTLEDMNSQL NSFTGQMENI
301 TTISQANEQN LKDLQDLHKD AENRTAIKEN QLEERFQLFE TDIVNIISNI SYTAHHLRTL
361 TSNLNEVRTT CDTLTLTKHTD DLTSLNNTLA NIRLDSVSLR MQQDLMRSRL DTEVANLSVI
421 MEEMKLVDSK HGQLIKNFTI LQGPPGPRGP RGDRGSQGPP GPTGNKGQKG EKGEPPGPPG
481 AGERGPIGPA GPPGERGGKG SKGSQGPKGS RGSFGKPGPQ GSSGDPGPPG PPGKEGLPGP
541 QGPPGFQGLQ GTVGEPPGVP PRGLPGLPGV PGMPGPKGPP GPPGPSGAVV PLALQNEPTP
601 APEDNGCPPH WKNFTDKCYY FSVEKEIFED AKLFCEDKSS HLVFINTREE QQWIKKQMVG
661 RESHWIGLTD SERENEWKWL DGTSPDYKNW KAGQPDNWDH GHGPGEDCAG LIYAGQWNDF
721 QCEDVNNFIC EKDRETVLSS AL (SEQ ID NO: 87).

```

[0200] As described herein, but not according to the invention, the non-target antigen comprises a polymorphism of COLEC12. For example, the non-target antigen comprises a peptide derived from COLEC12 comprising a polymorphic residue of COLEC12. Polymorphic residues of COLEC12 include position 522 of SEQ ID NO: 87. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of COLEC12 comprising amino acid 522 of SEQ ID NO: 87. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of COLEC12 comprising an S at amino acid 522 of SEQ ID NO: 87. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of COLEC12 comprising a P at amino acid 522 of SEQ ID NO: 87.

[0201] As described herein, but not according to the invention, the non-target antigen comprises APC down-regulated 1 (APCDD1) or an antigen peptide thereof in a complex with MHC-I. An exemplary human APCDD1 is described in UniProtKB record number Q8J025. As also described herein, but not according to the invention, APCDD1 comprises an amino acid sequence of:

1 MSWPRRLLLR YLFPALLLHG LGECSALLHP DSRSHPRSLE KSAWRAFKE QCHHMLKHLH
61 NGARITVQMP PTIEGHWVST GCEVRSGPEF ITRSYRFYHN NTFKAYQFY GSNRCTNPTY

5 121 TLIIRGKIRL RQASWIIRGG TEADYQLHN**V** QVICHTEAVA EKLGGQVNRT CPGFLADGGP
181 WVQDVAYDLW REENGCECTK AVNFAMHELQ LIRVEKQYLH HNL DHLVEEL FLGDIHTDAT
241 QRMFYRPSSY QPPLQNAKNH DHACIACRII YRSDEHHFPI LPPKADLTIG LHGEWVSQRC
10 301 EVRPEVLFLT RHFIFHDNNN TWEGHYHYHS DPVCKHPTFS IYARGRYSRG VLSSRVMGGT
361 EFVFKVNHMK VTPMDAATAS LLNVFNGNEC GAEGSWQVGI QQDVTHTNGC VALGIKLPHT
421 EYEIFKMEQD ARGRYLLFNG QRPSDGSSPD RPEKRATSYQ MPLVQCASSS PRAEDLAEDS
481 GSSLYGRAPG RHTWSLLLLA LACLVPLLHW NIRR (SEQ ID NO: 88).

15 **[0202]** As described herein, but not according to the invention, the non-target antigen comprises a polymorphism of APCDD1. Exemplary polymorphisms of APCDD1 include rs3748415, which can be a V, I or L at position 150 of SEQ ID NO: 88. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising amino acid 150 of SEQ ID NO: 88. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising an V at amino acid 150 of SEQ ID NO: 88. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising an I at amino acid 150 of SEQ ID NO: 88. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising an L at amino acid 150 of SEQ ID NO: 88.

20 **[0203]** A further exemplary human APCDD1 is described in UniProtKB record number V9GY82. As described herein, but not according to the invention, APCDD1 comprises an amino acid sequence of:

1 XDVAIDLWRE ENGCECTKAV NFAMHELQLI RVEKQYLHHN LDHLVEELFL GDIHTDATQR
61 MFYRPSSYQP PLQNAKCAAE SSGSFQILPQ DSSEKEQNGL SHWCLSRPGH QKDWALCAHA
30 121 GPATAGCPSC LWPPAETGRK AGRTSSKTVH ACPGEAGTSS FELF**F**FPNCW SIETKLKISL
181 NAKLSFKPRA SAPLETGHRV KIETLSQLVF LSFQILCCEV QSPLANK (SEQ ID NO: 134).

35 **[0204]** Exemplary polymorphisms of APCDD1 include rs1786683, which can be a Y or S at position 165 of SEQ ID NO: 133. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising amino acid 165 of SEQ ID NO: 134. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising a Y at amino acid 165 of SEQ ID NO: 134. As also described herein, but not according to the invention, the non-target antigen comprises a peptide of APCDD1 comprising an S at amino acid 165 of SEQ ID NO: 134.

40 **[0205]** A further exemplary human APCDD1 is described in UniProt record number J3QSE3. As described herein, APCDD1 comprises an amino acid sequence of:

1 PEDVLPALQL PAPSACQVE MGFHHVG**Q**DG LQLPTSSDPP ALASQSAGIT GVSHRPPGRH
45 61 LSNDLRTTMM PASPVGSSIG QTSTTLPSCP QRQT (SEQ ID NO: 135).

50 **[0206]** Exemplary polymorphisms of APCDD1 include rs9952598, which can be a Q or R at position 28 of SEQ ID NO: 135. As described herein, the non-target antigen comprises a peptide of APCDD1 comprising amino acid 28 of SEQ ID NO: 135. As also described herein, the non-target antigen comprises a peptide of APCDD1 comprising a Q at amino acid 28 of SEQ ID NO: 135. As also described herein, the non-target antigen comprises a peptide of APCDD1 comprising an R at amino acid 28 of SEQ ID NO: 135.

[0207] For example, APCDD1 comprises a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, at least 94%, at least 97%, at least 98%, or at least 99% identity to any one of SEQ ID NOs: 88 or 134-135. Polymorphic residues of APCDD1 are marked as bold and underlined in SEQ ID NOs: 88 or 134-135.

55 **[0208]** In some embodiments, the non-target antigen comprises HLA-A*02. Various single variable domains known in the art or disclosed herein that bind to and recognize HLA-A*02 are suitable for use in embodiments.

[0209] As described herein, but not according to the invention, the non-target antigen comprises HLA-B*07. Various single variable domains known in the art or disclosed herein that bind to and recognize HLA-B*07 are suitable.

[0210] As also described herein, but not according to the invention, the non-target antigen comprises HLA-A*11. Various single variable domains known in the art or disclosed herein that bind to and recognize HLA-A* 11 are suitable.

[0211] As also described herein, but not according to the invention, the non-target antigen comprises HLA-A*01. Various single variable domains known in the art or disclosed herein that bind to and recognize HLA-A*01 are suitable.

[0212] As also described herein, but not according to the invention, the non-target antigen comprises HLA-A*03. Various single variable domains known in the art or disclosed herein that bind to and recognize HLA-A*03 are suitable.

[0213] As also described herein, but not according to the invention, the non-target antigen comprises HLA-C*07. Various single variable domains known in the art or disclosed herein that bind to and recognize HLA-C*07 are suitable.

[0214] Such scFvs include, for example, the following mouse and humanized scFv antibodies that bind non-target antigens (e.g. HLA-A*02, HLA-A*01, HLA-A*03, HLA-A*11, HLA-B*07, or HLA-C*07) in a peptide-independent way shown in Table 4 below (where indicated, complementarity determining regions underlined):

Table 4. scFv binding domains

HLA-A*02 antigen binding domains	
C-001765 PA2.1 scFv (mouse) (not according to the invention)	DVLMTQTPLSLPVSLGDQASISCRSSQSIVHSNGNTYLEWYL QKPGQSPKLLIYKVSNRFSQVPDRFSGSGSGTDFTLKISRVE AEDLGVYYCFQGSHVPRTSGGGTKLEIKGGGGSGGGGSGG GGSGGQVQLQQSGPELVKPGASVRISCKASGYTFTSYHHW VKQRPQGQLEWIGWIYPGNVNTEYNEKFKGKATLTADKSS STAYMHLSSLTSEDSAVYFCAREEITYAMDYWGQGTSTV SS (SEQ ID NO: 89)
C-002159 PA2.1.8 scFv (humanized) (not according to the invention)	DVLMTQTPLSLPVSLGDQASISCRSSQSIVHSNGNTYLEWYL QKPGQSPKLLIYKVSNRFSQVPDRFSGSGSGTDFTLKISRVE AEDLGVYYCFQGSHVPRTSGGGTKLEIKGGGGSGGGGSGG GGSGGQVQLQQSGPELVKPGASVRISCKASGYTFTSYHHW VKQRPQGQLEWIGWIYPGNVNTEYNEKFKGKATLTADKSS STAYMHLSSLTSEDSAVYFCAREEITYAMDYWGQGTSTV SS (SEQ ID NO: 89) QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHHWVRQA PGQGLEWMGWIYPGNVNTEYNEKFKGKATITADKSTSTAY MELSSLRSEDYAVYYCAREEITYAMDYWGQGTSTVTVSSGG GGSGGGGSGGGGSGGEIVLTQSPGTLSPGERATLSCRSSQ SIVHSNGNTYLEWYQQKPGQAPRLLIYKVSNRFSQIPDRFSG SGSGTDFTLTISRLEPEDFAVYYCFQGSHVPRTFGGGTKVEI K (SEQ ID NO: 90)
C-002160 PA2.1.9 scFv (humanized) (not according to the invention)	C-002160 PA2.1.9 scFv (humanized): QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHHWVRQA PGQGLEWMGWIYPGNVNTEYNEKFKGKATITADKSTSTAY MELSSLRSEDYAVYYCAREEITYAMDYWGQGTSTVTVSSGG GGSGGGGSGGGGSGGDIVMTQTPLSLPVTPGEPASISCRSSQ SIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSQVPDRFSG SGSGTDFTLKISRVEAEDVGVYYCFQGSHVPRTFGGGTKVEI K (SEQ ID NO: 91)

(continued)

HLA-A*02 antigen binding domains	
5 10 15 20 25 30	C-002161 PA2.1.10 scFv (humanized) (not according to the invention) QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHHWVRQA PGQGLEWMGWIYPGNVNTEYNEKFKGKATITADKSTSTAY MELSSLRSEDTAVYYCAREEITYAMDYWGQGTTVTVSSGG GGSGGGGSGGGGSGGDIVMTQTPLSLPVTGPGEASISCRSSQ SIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNNRFSGVDPDRFSG SGSGTDFTLKISRVEADVGVYYCFQGSHPVPTFGGGTKVEI K (SEQ ID NO: 91) EVQLVESGGGLVKPGGSLRLSCAASGYTFTSYHHWVRQAP GKGLEWVGWIYPGNVNTEYNEKFKGRFTISRDDSKNTLYL QMNSLKTEDTAVYYCAREEITYAMDYWGQGTTVTVSSGG GGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRTITCRSS QSIVHSNGNTYLEWYQKPKGAPKLLIYKVSNNRFSGVPSRF SGSGGTDFTLTISSLQPEDFATYYCFQGSHPVPTFGGGTKV EIK (SEQ ID NO: 92) QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHHWVRQA PGQGLEWIGWIYPGNVNTEYNEKFKGKATITADESTNTAY MELSSLRSEDTAVYYCAREEITYAMDYWGQGTLLTVSSGG GGSGGGGSGGGGSGGDIQMTQSPSTLSASVGDRTITCRSS QSIVHSNGNTYLEWYQKPKGAPKLLIYKVSNNRFSGVPARF SGSGGTDFTLTISSLQPDDEFATYYCFQGSHPVPTFGQGTKV EVK (SEQ ID NO: 93)
35 40	C-002162 PA2.1.14 scFv (humanized) (according to the invention) QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHHWVRQA PGQGLEWIGWIYPGNVNTEYNEKFKGKATITADESTNTAY MELSSLRSEDTAVYYCAREEITYAMDYWGQGTLLTVSSGG GGSGGGGSGGGGSGGDIQMTQSPSTLSASVGDRTITCRSS QSIVHSNGNTYLEWYQKPKGAPKLLIYKVSNNRFSGVPARF SGSGGTDFTLTISSLQPDDEFATYYCFQGSHPVPTFGQGTKV EVK (SEQ ID NO: 93)
45	C-002163 PA2.1.18 scFv (humanized) (not according to the invention) QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHMHWVRQ APGQGLEWIGYIYPGNVNTEYNEKFKGKATLTADKSTNTA YMELSSLRSEDTAVYFCAREEITYAMDYWGQGTLLTVSSG GGSGGGGSGGGGSGGDVQMTQSPSTLSASVGDRTITCSS SQSIVHSNGNTYMEWYQKPKGAPKLLIYKVSNNRFSGVDPDR FSGSGGTDFTLTISSLQPDDEFATYYCHQGSHPVPTFGQGTK VEVK (SEQ ID NO: 94)
HLA-A*02 antigen binding domains derived from BB7.2 mAb	
50 55	C-002164 BB7.2 scFv (mouse): (not according to the invention) QVQLQQSGPELVKPGASVKMSCKASGYTFTSYHIQWVKQR PGQGLEWIGWIYPGDGSTQYNEKFKGKTTLTADKSSSTAY MLLSSLTSEDSAIYFCAREGTYYAMDYWGQGTSTTVSSGG GGSGGGGSGGGGSGGDVLMQTPLSLPVSLGDQVVISCRSS QSIVHSNGNTYLEWYLQKPGQSPKLLIYKVSNNRFSGVDPDRFS GSGGTDFTLKISRVEADLGVYYCFQGSHPVPTFGGGTKL EIK (SEQ ID NO: 95)

(continued)

HLA-A*02 antigen binding domains derived from BB7.2 mAb	
5 10	C-002165 BB7.2.1 scFv (humanized) (not according to the invention) QLQLQESGPGLVKPSETLSLTCTVSGYTFTSYHIQWIRQPPG KGLEWIGWIYPGDGSTQYNEKFKGRATISVDTSKNQFSLNL DSVSAADTAIYYCAREGTYYAMDYWGKGSTVTVSSGGGG SGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRSSQSI VHSNGNTYLEWYQQKPGKAPKLLIYKVSNRFSGVPSRFSGS GSGTDFTFITSSLPEDIATYYCFQGSHVPRTFGPGTKVDIK (SEQ ID NO: 96)
15 20	C-002166 BB7.2.2 scFv (humanized) (not according to the invention) EVQLVQSGAELKKPGSSVKVSCKASGYTFTSYHIQWVKQA PGQGLEWIGWIYPGDGSTQYNEKFKGKATLTVDKSTNTAY MELSSLRSEDТАVYYCAREGTYYAMDYWGQGTЛTVSSG GGSGGGGGSGGGSGGGDIQMTQSPSTLSASVGDRVITICRS SQSIVHSNGNTYLEWYQQKPGKAPKLLIYKVSNRFSGVPSR FSGSGSGTDFTLTISLQPDDEFATYYCFQGSHVPRTFGQGTK VEVK (SEQ ID NO: 97)
25 30	C-002167 BB7.2.3 scFv (humanized): (not according to the invention) QVQLVQSGAEVKKPGSSVKVSCKASGYTFTSYHIQWVRQA PGQGLEWMGWIPGDGSTQYNEKFKGRVTITADKSTSTAY MELSSLRSEDТАVYYCAREGTYYAMDYWGQGTТTVTVSSG GGSGGGGGSGGGSGGGEIVLTQSPGTLSPGERATLSCRSS QSIVHSNGNTYLEWYQQKPGQAPRLLIYKVSNRFSGIPDRFS GSGSGTDFTLTISRLEPEDFAVYYCFQGSHVPRTFGGGTKVE IK (SEQ ID NO: 98)
35 40	C-002168 BB7.2.5 scFv (humanized) (not according to the invention) QVTLKQSGAEVKKPGSSVKVSCТАSGYTFTSYHVSWVRQA PGQGLEWLGRIPGDGSTQYNEKFKGKVITITADKSMDTSF MELTSLTSEDТАVYYCAREGTYYAMDЛWGQGTЛTVTVSSG GGSGGGGGSGGGSGGGEIVLTQSPGTLSPGERATLSCRSS QSIVHSNGNTYLAWYQQKPGQAPRLISKVSNRFSGVPDRLF SGSGSGTDFTLTISRLEPEDFAVYYCQQGSHPRTFGGGTKV EIK (SEQ ID NO: 99)
45	C-002169 BB7.2.6 scFv (humanized) (not according to the invention) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYHMHWVRQ APGQRLEWMGWIPGDGSTQYNEKFKGKVITITRDTSASTA YMELSSLRSEDТАVYYCAREGTYYAMDYWGQGTЛTVTVSS GGGGSGGGSGGGSGGGDIVMTQTPLSLPVTPGEPASISCRS SQSIVHSNGNTYLDWYLQKPGQSPQLLIYKVSNRFSGVPDRL FSGSGSGTDFTLKISRVEADVGVYYCМQGSHPRTFGGGT KVEIK (SEQ ID NO: 100)
HLA-B*07 antigen binding domains	
50 55	BB7.1.10 scFv (not according to the invention) QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYSWHWIRQPP KGLEWIGYIHFSGSTHYHPSLKSРVTISVDTSKNQFSLKLSS VTAADТАVYYCARGGVVSHYAMDCWGQGTТTVTVSSGGG GSGGGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRASE NIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGSG TDFTLTISLQPEDFATYYCQHFVWTPYTFGGGTКVEIK (SEQ ID NO: 199)

(continued)

HLA-B*07 antigen binding domains	
5 BB7.1.9 scFv (not according to the invention)	EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ APGKGLEWVSYIHFSGSTHYHPSLKSRTISRDNAKNSLYLQ MNSLRAEDTAVYYCARGGVVSHYAMDCWGQGTTVTVSSG GGGSGGGGSGGGGSGGDIQMTQSPSSVSASVGDRVITTCRA 10 SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSG SGTDFTLTISSLQPEDFATYYCQHFVWVTPYTFGGGTKVEIK (SEQ ID NO: 200)

15

20

25

30

35

40

45

50

55

(continued)

HLA-B*07 antigen binding domains

BB7.1.8 scFv (not
according to the invention)

QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYSWHWIRQPP
GKGLEWIGYIHFSGSTHYHPSLKSRTISVDTSKNQFSLKLSS
VTAADTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGGG
GGGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITITCRASE
NIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGSG
TDFTLTISLQPEDFATYYCQHFVWTPYTFGGGGTKVEIK
(SEQ ID NO: 199)

EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ
APGKGLEWVSYIHFSGSTHYHPSLKSRTISRDNANKSLYLQ
MNSLRAEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSG
GGGSGGGGSGGGGSGGDIQMTQSPSSVSASVGDRVITITCRA
SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSG
SGTDFTLTISLQPEDFATYYCQHFVWTPYTFGGGGTKVEIK
(SEQ ID NO: 200)

EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ
APGKGLEWVGYYIHFSGSTHYHPSLKSRTISRDDSKNTLYLQ
MNSLKTEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSG
GGGSGGGGSGGGGSGGEIVLTQSPATLSLSPGERATLSCRAS
ENIYSNLAWYQQKPGQAPRLIYAATYLPDGIPARFSGSGSG
TDFTLTISLQPEDFAVYYCQHFVWTPYTFGGGGTKVEIK
(SEQ ID NO: 201)

QVQLQQSGPGLVKPSQTLSTCAISGYSITSGYSWHWIRQSP
SRGLEWLGYIHFSGSTHYHPSLKSRTINPDTSKNQFSLQLNS
VTPEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGGGG
SGGGGSGGGGSGGEIVLTQSPATLSLSPGERATLSCRASENI
YSNLAWYQQKPGQAPRLIYAATYLPDGIPARFSGSGSGTD
FTLTISRLEPEDFAVYYCQHFVWTPYTFGGGGTKVEIK (SEQ
ID NO: 202)

EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ
APGKGLEWVGYYIHFSGSTHYHPSLKSRTISRDDSKNTLYLQ
MNSLKTEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSG
GGGSGGGGSGGGGSGGDIQMTQSPSSVSASVGDRVITITCRA
SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSG
SGTDFTLTISLQPEDFATYYCQHFVWTPYTFGGGGTKVEIK
(SEQ ID NO: 203)

EVQLVESGGGLVQPGGSLRLSCAASGYSITSGYSWHWVRQ
APGKGLEWVSYIHFSGSTHYHPSLKSRTISRDNANKNTLYLQ
MNSLRAEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSG
GGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITITCRA
SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSG

(continued)

HLA-B*07 antigen binding domains	
5 10	<u>BB7.1.7 scFv (not according to the invention)</u> QVQLQQSGPGLVKPSQTLSTCAISGYSITSGYSWHWIRQSP SRGLEWLGYIHFSGSTHYHPSLKSRTINPDTSKNQFSLQLNS VTPEDTAVYYCARGGVVSHYAMDCWGQGTTVTVSSGGGG SGGGGSGGGSGGGEIVLTQSPATLSLSPGERATLSCRA SENIYSNLAWYQQKPGQAPRLLIYAATYLPDGI PARFSGSGSGTDFTLTISRLEPEDFAVYYCQHF WVTPYTFGGGGTKVEIK (SEQ ID NO: 202)
15 20	<u>BB7.1.6 scFv (not according to the invention)</u> EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ APGKGLEWVGYIHFSGSTHYHPSLKSRTISRDDSKNTLYLQ MNSLKTEDTAVYYCARGGVVSHYAMDCWGQGTTVTVSSG GGGSGGGSGGGSGGDIQMTQSPSSVSASVGDRVTITCRA SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFS GSGSGTDFTLTISLQPEDFATYYCQHFVWVTPYTFGGG GTKVEIK (SEQ ID NO: 203)
25 30	<u>BB7.1.5 scFv (not according to the invention)</u> EVQLVESGGGLVQPGGSLRLSCAASGYSITSGYSWHWVRQ APGKGLEWVSYIHFSGSTHYHPSLKSRTISRDN SKNTLYLQMNSLRAEDTAVYYCARGGVVSHYAMDCWG QGTTVTVSSGGGGSGGGSGGGSGGDIQMTQSPSSLSA SVGDRVTITCRA SENIYSNLAWYQQKPGKAPKLLIYA ATYLPDGVPSRFSGSGSGTDFTLTISLQPEDFATYYC QHFVWVTPYTFGGGGTKVEIK (SEQ ID NO: 204)
35 40	<u>BB7.1.4 scFv (not according to the invention)</u> EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ APGKGLEWVGYIHFSGSTHYHPSLKSRTISRDDSKNTLYLQ MNSLKTEDTAVYYCARGGVVSHYAMDCWGQGTTVTVSSG GGGSGGGSGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFS GSGSGTDFTLTISLQPEDFATYYCQHFVWVTPYTFGGG GTKVEIK (SEQ ID NO: 205)

40

45

50

55

(continued)

HLA-B*07 antigen binding domains	
5 BB7.1.3 scFv (not according to the invention)	SGTDFTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 204) EVQLVESGGGLVKPGGSLRLSCAASGYSITSGYSWHWVRQ APGKGLEWVGYIHFSGSTHYHPSLKSRTISRDDSKNTLYLQ 10 MNSLKTEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSG GGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITTCRA SENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSG SGTDFTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 205) 15 QVQLQQWGAGLLKPSETLSLTCAVYGYSITSGYSWHWIRQ PPGKGLEWIGYIHFSGSTHYHPSLKSRTISVDTSKNQFSLKL SSVTAADTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGG GGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITTCRAS 20 ENIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGS GTDFTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 206) 25 QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYSWHWIRQH PGKGLEWIGYIHFSGSTHYHPSLKSRTISVDTSKNQFSLKLS SVTAADTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGGG GSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITTCRASE NIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGSG 30 TDFTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 207) 35 QVQLQQSGPGLVKPSQTLSTCAISGYSITSGYSWHWIRQSP SRGLEWLGYIHFSGSTHYHPSLKSRTINPDTSKNQFSLQLNS VTPEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGGGG SGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITTCRASENI YSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 208)
40 BB7.1.2 scFv (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYSWHWIRQH PGKGLEWIGYIHFSGSTHYHPSLKSRTISVDTSKNQFSLKLS SVTAADTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGGG 45 GSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITTCRASE NIYSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGSG TDFTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 207)
50 BB7.1.1 scFv (not according to the invention)	QVQLQQSGPGLVKPSQTLSTCAISGYSITSGYSWHWIRQSP SRGLEWLGYIHFSGSTHYHPSLKSRTINPDTSKNQFSLQLNS VTPEDTAVYYCARGGVVSHYAMDCWGQGTTTVTVSSGGGG 55 SGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITTCRASENI YSNLAWYQQKPGKAPKLLIYAATYLPDGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQHFVWTPYTFGGGKVEIK (SEQ ID NO: 208)

(continued)

HLA-A*11 antigen binding domains	
5 10	<u>9 (not according to the invention)</u> QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRTISVDTSKNQFSLKL SSVTAADTAVYYCARHYYYYSMDEVWGKGTTVTVSSGGGG SGGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRASQSI SSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 209)
15 20	<u>8 (not according to the invention)</u> QITLKESGPTLVKPTQTLTLCTFSGFSLSTSGVGVGWIRQPP GKALEWLALIYWND DKRYSPSLKSRLTITKDTSKNQVVL MTNMDPVDATATYYCAHRHMLSCFDYWGQGLTVTVSSGG GGGSGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRAS QSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSG TDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 210)
25 30	<u>7 (not according to the invention)</u> QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYAMHWVRQ APGQRLEWMGWINAGNGNTKYSQKFQGRVTITRDTSASTA YMELSSLRSEDATVYYCAREGNGANPDADFIDWGQGMVTV SSGGGSGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITIT CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSG SSGSGTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 211)
35 40	<u>6 (not according to the invention)</u> EVQLVESGGGLVQP GGSRLSCAASGFTFSSYDMHWVRQA TGKGLEWVSAIGTAGDTYYPGSVKGRFTISRENAKNSLYLQ MNSLRAGDTAVYYCARDLPGSYWFYDLWGRGTLTVTVSSG GGGSGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS GTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 212)
45 50	<u>5 (not according to the invention)</u> QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRTISVDTSKNQFSLKL SSVTAADTAVYYCARHYYYYYLDVWGKGTTVTVSSGGGG SGGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRASQSI SSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 213)
55	<u>4 (not according to the invention)</u> EVQLVESGGGLVQP GGSRLSCAASGFTFSSYWMHWVRQA PGKGLVWVSRINSDGSSTSYADSVKGRFTISRDN AKNTLYL QMNSLRAEDTAVYYCCLGVLLYNWFDWPWGQGLTVTVSSG GGGSGGGSGGGSGGGDIQMTQSPSSLSASVGDRVITICRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS GTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 214)

(continued)

HLA-A*11 antigen binding domains	
5 10	<u>3 (not according to the invention)</u> QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRLTISVDTSKNQFSLKL SSVTAADTAVYYCARHYYYYYMDVWGKGTTVTVSSGGGGG GGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDF TLTISSLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 215)
15 20	<u>2 (not according to the invention)</u> QITLKESGPTLVKPTQTLTLCTFSGFSLSTSGVGVGWIRQPP GKALEWLALIYWDDKRYSPSLKSRLTITKDTSKNQVVL MTNMDPVDATYYCAHKTTSFYFDYWGGQGLTVTVSSGGG GSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SSSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGT DFTLTISSLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 216)
25 30	<u>1 (not according to the invention)</u> QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRLTISVDTSKNQFSLKL SSVTAADTAVYYCARHYYYYYMDVWGKGTTVTVSSGGG GSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SSSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGT DFTLTISSLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 217)
HLA-C*07 antigen binding domains	
35	<u>C7-45 (not according to the invention)</u> EVQLLESGLLVQPGGSLRLSCAASGFTSSYAMSWVRQAPGKLEWVSAISGS GGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAVSFDWFDPWG QGTLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SSSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFA TYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 218)
40	<u>C7-44 (not according to the invention)</u> QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRLTISVDTSKNQFSLKLSSVTAADTAVYYCARERSISPYYYYYMD VWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 219)
45 50	<u>C7-43 (not according to the invention)</u> QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYYWGWIRQPPGKLEWIGSIYY GSTYYNPSLKSRLTISVDTSKNQFSLKLSSVTAADTAVYYCARDSEIVWYFDPWG QGTLTVTVSSGGGGSGGGGSGGGGSGGQSVLTQPPSASGTPGQRTVITCSGSSSN IGSNTVNWYQQLPGTAPKLLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDE ADYYCAAWDDSLNGWVFGGKTLTVL (SEQ ID NO: 220)
55	<u>C7-42 (not according to the invention)</u> QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRLTISVDTSKNQFSLKLSSVTAADTAVYYCAREILPRLSYYYYYMD VWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 221)

(continued)

HLA-C*07 antigen binding domains	
5 C7-41 (not according to the invention)	EVQLLES GGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGS GGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAVSFDWFDPWG QGT LVT VSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SIS SYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFA TYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 218)
10	QVQLQESG PGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCARERSISPYYYYYMD VWGKGTTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 219)
15	QLQLQESG PGLVKPSETLSLTCTVSGGSISSSSYYWG WIRQPPGKGLEWIGSIYY GSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCARD SVIWWYFDPWG QGT LVT VSSGGGGSGGGGSGGGGSGGQSVLTQPPSASGTPGQRTVISCSSSSN IGSNTVNWYQQLPGTAPKLLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDE ADYYCAAWDDSLNGWVFGGGTKLTVL (SEQ ID NO: 220)
20	QVQLQESG PGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCAREILPRLSYYYYYMD VWGKGTTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 221)
25	QVQLVQSGSELKPGASVKVSC KASGYTFTSYAMN WVRQAPGQGLEWMGWI NTNTGNPTYAQGFTGRFVFSFDTSVSTAYLQICSLKAEDTAVYYCARGGRAHSSW YFDLWGRGTLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 222)
30	QVQLQESG PGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCARDRIKLPRLGYYYY MDVWVGKTTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 223)
35	QVQLQESG PGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCARDTVIHYYYYMDV WGKGTTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 224)
40	QVQLQESG PGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCARDVIVEVFLSYYYY MDVWVGKTTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGG TKVEIK (SEQ ID NO: 225)
45	QVQLQESG PGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSL KSRVTISVDT SKNQFSLKLSSVTAADTAVYYCARDIFIHYYYYMDVW GKGTTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRAS QSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDF
50	

55

(continued)

HLA-C*07 antigen binding domains	
5 C7-40 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDRIKILPRLGYYYY MDVWVGKGTTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 223)
10 C7-39 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDTVIHYYYYMDV WGKGTTVTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 224)
15 C7-38 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDVIVEVFLSYYYY MDVWVGKGTTVTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 225)
20 C7-37 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDIFIHYYYYMDVW GKGTTVTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTITCRAS QSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDF ATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 226)
25 C7-36 (not according to the invention)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSYISS SSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDGTFYSYSPYYF DYWGQGTTLTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE EDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 227)
30 C7-35 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAREWIKILPRLGYYYY MDVWVGKGTTVTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 228)
35 C7-34 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDRLYYYYYMDV WGKGTTVTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 229)
40 C7-33 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDKILAPNYYYYMD VWVGKGTTVTVTVSSGGGGSGGGGSGGGGSGGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE DFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 230)

(continued)

HLA-C*07 antigen binding domains	
5 C7-32 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCAREKSWKYFYYYYYY MDVWVGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISL QPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 231)
10 C7-31 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARENTSTIPYYYYYM DVWVGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQP EDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 232)
15 C7-30 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCAREDVDKNTSTIYYY YYYMDVWVGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTIS SLQPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 233)
20 C7-29 (not according to the invention)	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWIRQAPGKGLEWVSYISS GSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDGGDIVSSSAIY WYFDLWGRGTLTVTVSSGGGGSGGGGSGGGGSGGAIQLTQSPSSLSASVGDRVT ITCRASQGISSALAWYQQKPGKAPKLLIYDASSLESGVPSRFSGSGSGTDFLTISL QPEDFATYYCQQFNSYPLTFGGGTKVEIK (SEQ ID NO: 234)
25 C7-28 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDLILPPYYYYYMD VWVGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPE DFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 235)
30 C7-27 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARETWIKILPRYYYYY YYMDVWVGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTIS SLQPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 236)
35 C7-26 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDLSRYYYYYYMDV WGKGTTVTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPED FATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 237)
40 C7-25 (not according to the invention)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSYISS SSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAREHIVLCFDYWG QGTTLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ GISSWLAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPEDFA TYYCQQYNSYPLTFGGGTKVEIK (SEQ ID NO: 238)

(continued)

HLA-C*07 antigen binding domains	
5 C7-24 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDKILPRPYYYYYYM DVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPE EDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 239)
10 C7-23 (not according to the invention)	QVQLVQSGAEVKKPGASVKVCKASGYTFTSYGISWVRQAPGQGLEWMGWIS AYNGNTNYAQKLQGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARGSNEYFQ HWGQGTTLTVTVSSGGGGSGGGGSGGGGSGGQSALTQPPSASGSPGQSVTISCT GTSSDVGGYNYVSWYQQHPGKAPKLLMIYEVSKRPSGVPDRFSGSKSGNTASLTV SGLQAEDEADYYCSSYAGSNWVFGGGTKLTVL (SEQ ID NO: 240)
15 C7-22 (not according to the invention)	QVQLVQSGSELKKPGASVKVCKASGYTFTSYAMNWRQAPGQGLEWMGWI NTNTGNPTYAQGFTGRFVSFDTSVSTAYLQICSLKAEDTAVYYCARGTSYWFYD LWGRGTLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPE DFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 241)
20 C7-21 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAREEIVEFYIYYMD VWGKGTTVTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPE DFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 242)
25 C7-20 (not according to the invention)	EVQLLESGLLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGS GGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKVDYYFDYWG QGTLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPEDFA TYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 243)
30 C7-19 (not according to the invention)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYWMHWVRQAPGKGLVWVSRINS DGSSTSYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAWSTNILLSYTKA FDIWGQGTMTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISL QPEDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 244)
35 C7-18 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDKTYYYYYYMDV WGKGTTVTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTISLQPED FATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 245)
40 C7-17 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAREKYFHDYFHDYY YYYMDVWGKGTTVTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFLTIS SLQPEDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 246)

(continued)

HLA-C*07 antigen binding domains	
5 C7-16 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDTSVYYYYYMDV WGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 247)
10 C7-15 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAREKILPYYYYYYMD VWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE DFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 248)
15 C7-14 (not according to the invention)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSYISS SSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAIQWIIYINPRGFIF LHDAFDIWGQGTMTVTVSSGGGGSGGGGSGGGGSGGQSVLTQPPSASGTPGQ RVTISCSGSSSNIGSNTVNWYQQLPGTAPKLLIYSNNQRPSGVPDRFSGSKSGTSA SLAISGLQSEADYYCAAWDDSLNGWVFGGGTKLTVL (SEQ ID NO: 249)
20 C7-13 (not according to the invention)	QVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIRQSPSRGLEWLGRTY YRSKWYNDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDTAVYYCAKEDVDFHHD AFDIWGGQGTMTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVT ITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 250)
25 C7-12 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAREGVDKNTSTIYY YYYMDVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTIS SLQPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 251)
30 C7-11 (not according to the invention)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSYISS SSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRRGYFDLWGR GTLVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQGI SSWLAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFAT YYCQQYNSYPLTFGGGTKVEIK (SEQ ID NO: 252)
35 C7-10 (not according to the invention)	EVQLVQSGAEVKKPGATVKISKVSGYFTDYMHVWVQAPGKGLEWMGLVD PEDGETIYAEKFQGRVTITADTSTDYAMELSSLRSEDVAVYYCATGIHVDIRSME DWFDWPWGQGLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDR VTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTIS SLQPEDFATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 253)
40 C7-9 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDIGTSYYYYYMDV WGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQSYSTPLTFGGGTKVEIK (SEQ ID NO: 254)

(continued)

HLA-C*07 antigen binding domains	
5 C7-8 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAREVVEFLYYYYYM DVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE EDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 255)
10 C7-7 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDLYYYYYYMDV WGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 256)
15 C7-6 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARESWKYFYPRGSIFI HYYYYYMDVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVG DRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTL TISLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 257)
20 C7-5 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGYIYY SGSTYYNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDRIVEVFYYYYMD VWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE DFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 258)

30

35

40

45

50

55

(continued)

HLA-C*07 antigen binding domains	
5 C7-4 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCAREVVEFLYYYYYM DVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQ 10 EDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 255) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDLYYYYYYMDV WGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPED 15 FATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 256) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCAREWKYFYPRGSIFI HYYYYYMDVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVG DRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTL 20 TISLQPEDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 257) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDRIVEVFYYYYMD VWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCR 25 ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPE DFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 258) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCAREKYFHDWLYYYYY MDVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI 30 TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 259) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDLVDKNTSYYYY 35 MDVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 260) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQAPGQGLEWMGWIS AYNGNTNYAQKLQGRVTMTTDTSTAYMELRSLRSDDTAVYYCARVQNEYFQ 40 HWGQGLTVTVSSGGGGSGGGGSGGGGSGGQSALTQPPSASGSPGQSVTISCT GTSSDVGGYNYVSWYQQHPGKAPKLMIEVSKRPSGVPDRFSGSKSGNTASLTV SGLQAEDEADYYCSSYAGSNNWVFGGGTKLTVL (SEQ ID NO: 261) QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWIRQAPGKGLEWVSYISS 45 GSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCATANWFDPWGQ GTLVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSVSASVGDRVTITCRASQGI SSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFAT YYCQQAQNSFPLTFGGGTKVEIK (SEQ ID NO: 262)
50 C7-3 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQPPGKGLEWIGIYY SGSTYYNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDLVDKNTSYYYY MDVWGKGTTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTI TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISL 55 QPEDFATYYCQSYSTPLTFGGGTKVEIK (SEQ ID NO: 260)

(continued)

HLA-C*07 antigen binding domains	
5 C7-2 (not according to the invention)	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQAPGQGLEWMGWIS AYNGNTNYAQKLQGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARVQNEYFQ HWGQGLTVTVSSGGGGSGGGGSGGGGSGGQSALTQPPSASGSPGQSVTISCT GTSSDVGGYNVSWYQQHPGKAPKLMIEVSKRPSGVPDRFSGSKSGNTASLTV SGLQAEDEADYYCSSYAGSNNWVFGGGTKLTVL (SEQ ID NO: 261)
10 C7-1 (not according to the invention)	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWIRQAPGKGLEWVSYISS GSTIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCATANWFDPWGQ GTLTVTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSVSASVGDRVTITCRASQGI SSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFAT YYCQQANSFPLTFGGGKVEIK (SEQ ID NO: 262)
HLA-A*03 scFv Sequences	
20 15 (not according to the invention)	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMELRSLRSDDTAVYYCARERVSQRGAFDIWGQGTMTVTS SGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 263)
30 16 (not according to the invention)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQA PGKGLEWVSYISSSSSTIYYADSVKGRFTISRDNAKNSLYLQ MNSLRAEDTAVYYCARGNPDKDPFDYWGQGLTVTVSSGG GGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRAS QSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSG TDFTLTISLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 264)
35 17 (not according to the invention)	QVQLQESGPGLVKPSETLSLTCTVSGGSVSSGSYYWSWIRQ PPGKGLEWIGYIYYSGSTNYPNPSLKSRVTISVDTSKNQFSLK LSSVTAADTAVYYCARDFYCTNWFYFDLWGRGTLTVTVSSG GGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRA SQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS GTDFTLTISLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 265)
40 18 (not according to the invention)	QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIRQPPG KGLEWIGYIYYSGSTNYPNPSLKSRVTISVDTSKNQFSLKLSS VTAADTAVYYCARESSSGSYWYFDLWGRGTLTVTVSSGGG GSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVTITCRASQ SISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGT DFTLTISLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 266)

(continued)

HLA-A*03 scFv Sequences	
5 10	19 (not according to the invention) EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMP GKGLEWMGIIYPGDS DTRYSPSFQGGVTTISADKSISTAYLQW SSLKASDTAMY YCARD SGYKYNLYYYYYYMDVWGKGT VTVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRV TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSR FSGSGSGTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGK VEIK (SEQ ID NO: 267)
15 20	20 (not according to the invention) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMERLSLRSDDTAVYYCARGGDL SHYYYYYMDVWGKGT VTVSSGGGGSGGGGSGGGGSGGQTVVTQEPLTVSPGGTV TLTCASSTGAVTSGYYPNWFQQKPGQAPRALIYSTSNKHSW TPARFSGSLLGGKAALTL SGVQPEDEAEYYCLLYGGAQW VFGGGKLTVL (SEQ ID NO: 268)
25 30	21 (not according to the invention) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMERLSLRSDDTAVYYCARENRRYNSCYFDYWGQGTLV TVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRV ITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRF SGSGSGTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGK EIK (SEQ ID NO: 269)
35 40	22 (not according to the invention) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMERLSLRSDDTAVYYCARGGDL SHYYYYYLDVWGKGT TVSSGGGGSGGGGSGGGGSGGQTVVTQEPLTVSPGGTVTL TCASSTGAVTSGYYPNWFQQKPGQAPRALIYSTSNKHSWTP ARFSGSLLGGKAALTL SGVQPEDEAEYYCLLYGGAQWVF GGGKLTVL (SEQ ID NO: 270)
45 50	23 (not according to the invention) EVQLVESGGGLVQP GGSRLSCAASGFTVSSNYMSWVRQA PGKGLEWVSVIYSGGSTYYADSVKGRFTISRDN SKNTLYLQ MNSLRAEDTAVYYCARATLLSLSYDAFDIWGQGTMTVSS GGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRV TITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSG SGTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 271)
55	24 (not according to the invention) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMERLSLRSDDTAVYYCARGGDL SHYYYYYMDVWGKGT TVTVSSGGGGSGGGGSGGGGSGGQTVVTQEPLTVSPGGTVTLT CASSTGAVTSGYYPNWFQQKPGQAPRALIYSTSNKHSWTPA RFSGSLLGGKAALTL SGVQPEDEAEYYCLLYGGAQWVFG GGKLTVL (SEQ ID NO: 272)

(continued)

HLA-A*03 scFv Sequences	
5 25 (not according to the invention)	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMP GKGLEWMGIIYPGDS DTRYSPSFQGGQVTISADKSISTAYLQW SSLKASDTAMY YCARE RDRWFD PWGQGLTVTVSSGGGGGS GGGGSGGGGSGGDIQMTQSPSSLSASVGDRTITCRASQSS SYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDF 10 TLTISSLQPEDFATYYCQQSYSTPLTFGGGKVEIK (SEQ ID NO: 273)
15 26 (not according to the invention)	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMERLSLRSDDTAVYYCARETPPSLGAFDIWGGTMVTVS SGGGGSGGGGSGGGGSGGQSALTQPPSASGSPGQSVTISCT GTSSDVGGYNYVSWYQQHPGKAPKLMIEVSKRPSGVPDR 20 FSGSKSGNTASLTVSGLQAEDADYYCSSYAGSNNWVFGG GTKLTVL (SEQ ID NO: 274)
25 27 (not according to the invention)	QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYYWGWRQP PGKGLEWIGSIYYSGSTYYNPSLKSRTISVDTSKNQFSLKLS SVTAADTAVYYCAREAYCLSDSYWYFDLWGRGTLTVTVSS GGGGSGGGGSGGGGSGGQSVLTQPPSASGTPGQRTISCSG SSSNIGSNTVNWYQQLPGTAPKLLIYSNNQRPSGVPDRFSGS KSGTSASLAISGLQSEDEADYYCAAWDDSLNGWVFGGGTK 30 LTVL (SEQ ID NO: 275)
35 28 (not according to the invention)	QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRTISVDTSKNQFSLKL SSVTAADTAVYYCARESWKYFYPRGYMDVWGKGTITVTVS SGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRTITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGKVEIK 35 (SEQ ID NO: 276)
HLA-A*01 scFv Sequences	
40 A1-9 (not according to the invention)	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQA PGQGLEWMGWISAYNGNTNYAQKLQGRVTMTTDTSTSTA YMERLSLRSDDTAVYYCARGGWTAWYYYMDVWGKGTITV 45 TVSSGGGGSGGGGSGGGGSGGQTVVTQEPSLTVSPGGTVTL TCASSTGAVTSGYYPNWFQQKPGQAPRALIYSTSNKHSWTP ARFSGSLLGGKAALTLSGVQPEDEAEYYCLLYGGAQWVF GGGKTLTVL (SEQ ID NO: 277)
50 A1-8 (not according to the invention)	EVQLVESGGGLVQP GGSRLSCAASGFTFSSYSMNWVRQA PGKGLEWVSYISSSSSTIYYADSVKGRFTISRDNKNSLYLQ MNSLRAEDTAVYYCARAKYYYMDVWGKGTITVTVSSGGG 55 GSGGGGSGGGGSGGQSVLTQPPSASGTPGQRTISCSGSSN IGSNTVNWYQQLPGTAPKLLIYSNNQRPSGVPDRFSGSKSGT SASLAISGLQSEDEADYYCAAWDDSLNGWVFGGGKTLTVL (SEQ ID NO: 278)

(continued)

HLA-A*01 scFv Sequences	
5 10	A1-7 (not according to the invention) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRTISVDTSKNQFSLKL SSVTAADTAVYYCARDQVDKNTYYYYMDVWGKGTITVTV SSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITIT CRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSG SGSGTDFTLTISLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 279)
15 20	A1-6 (not according to the invention) QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWIRQAP GKGLEWVSYISSSGSTIYYADSVKGRFTISRDNALNSLYLQ MNSLRADTAVYYCARACQLAEYFQHWGQGTLVTVSSGG GGSGGGGSGGGGSGGDIQMTQSPSSVSASVGDRVITITCRAS QGISSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS GTDFTLTISLQPEDFATYYCQQANSFPLTFGGGGTKVEIK (SEQ ID NO: 280)
25 30	A1-5 (not according to the invention) QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRTISVDTSKNQFSLKL SSVTAADTAVYYCARDRVDKNTSYYMDVWGKGTITVTVS SGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITITC RASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGS GSGTDFTLTISLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 281)
35 40	A1-4 (not according to the invention) QVQLQESGPGLVKPSDTLSLTCAVSGYSISSNWWGWIRQP PGKGLEWIGYIYYSGSTYYNPSLKSRTTMSVDTSKNQFSLK LSSVTAVDTAVYYCARRVQLKLVHWFDWPWGQGLTVTVSS GGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITITCR ASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSG SGTDFTLTISLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 282)
45 50	A1-3 (not according to the invention) QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYDINWVRQA TGQGLEWMGMNPNSGNTGYAQKFQGRVTMTNRTSISTA YMELSSLRSEDVAVYYCATYYDYVTVFYFQHWGQGLTVT VSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVIT TCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFS GSGSGTDFTLTISLQPEDFATYYCQQSYSTPLTFGGGGTKVEI K (SEQ ID NO: 283)
55	A1-2 (not according to the invention) QLQLQESGSLVKPSQTLSTCAVSGGSISSGGYSWSWIRQP PGKGLEWIGYIYHSGSTYYNPSLKSRTISVDRSKNQFSLKL SSVTAADTAVYYCARESYPSFYAFDIWGQGTMTVTVSSGGG GSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVITITCRASQ SISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGSGT DFTLTISLQPEDFATYYCQQSYSTPLTFGGGGTKVEIK (SEQ ID NO: 284)

(continued)

HLA-A*01 scFv Sequences	
5 10 A1-1 (not according to the invention)	QITLKESGPTLVKPTQTLTLCTFSGFSLSTSGVGVGWIRQPP GKALEWLALIYWNDKRYSPSLKSRLTITKDTSKNQVVL MTNMDPVDATATYYCAHSNMWSYSLNDYYFDYWGQGLV TVSSGGGGSGGGGSGGGGSGGDIQMTQSPSSLSASVGDRVT ITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRF SGSGSGTDFTLTISSLQPEDFATYYCQQSYSTPLTFGGGTKV EIK (SEQ ID NO: 285)

[0215] Exemplary heavy chain and light chain CDRs (CDR-H1, CDR-H2 and CDR-H3, or CDR-L1, CDR-L2 and CDR-L3, respectively) for non-target ligand binding domains are shown in table 5 below.

Table 5. CDRs corresponding to antigen binding domains

CDR-L1	CDR-L2	CDR-L3	CDR-H1	CDR-H2	CDR-H3
HLA-A*02 CDRs					
RSSQSIVHSN GNTYLE (SEQ ID NO: 101) (according to the invention)	KVSNRFGV P DR (SEQ ID NO: 102) (not according to the invention) KVSNRFGV P AR (according to the invention)	FQGSHPRT (SEQ ID NO: 103) (according to the invention)	ASGYFTSYHI H (SEQ ID NO: 104) (according to the invention)	WIYPGNVNT EYNEKFKGK (SEQ ID NO: 105) (according to the invention)	EEITYAMDY (SEQ ID NO: 106) (according to the invention)
RSSQSIVHSN GNTYLD (SEQ ID NO: 107) (not according to the invention)	KVSNRFGV P DR (SEQ ID NO: 108) (not according to the invention)	MQGSHVPT (SEQ ID NO: 109) (not according to the invention)	SGYFTSYHM H (SEQ ID NO: 110) (not according to the invention)	WIYPGDGSTQYNEKFKG (SEQ ID NO: 111) (not according to the invention)	EGTYAMDY (SEQ ID NO: 112) (not according to the invention)
HLA-A*03 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	ERVSRGAFD I (SEQ ID NO: 345)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYSMN (SEQ ID NO: 306)	YISSSSSTIYA DSVKG (SEQ ID NO: 327)	GNPDKDPFD Y (SEQ ID NO: 346)

(continued)

HLA-A*03 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGSYWS (SEQ ID NO: 307)	YIYSGSTNYN PSLKS (SEQ ID NO: 328)	DFYCTNWYF DL (SEQ ID NO: 347)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYWWS (SEQ ID NO: 308)	YIYSGSTNYN PSLKS (SEQ ID NO: 328)	ESSSGSYWYF DL (SEQ ID NO: 348)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYWIG (SEQ ID NO: 309)	IYPGDSDTRY SPSFQG (SEQ ID NO: 329)	DSGYKYNLYY YYYMDV (SEQ ID NO: 349)
ASSTGAVTSG YYPN (SEQ ID NO: 287)	STSNKHS (SEQ ID NO: 294)	LLYYGGAQW V (SEQ ID NO: 299)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	GGDLSHYYY MDV (SEQ ID NO: 350)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	ENRRYNSCYY FDY (SEQ ID NO: 351)

(continued)

HLA-A*03 CDRs (not according to the invention)					
ASSTGAVTSG YYPN (SEQ ID NO: 287)	STSNKHS (SEQ ID NO: 294)	LLYYGGAQW V (SEQ ID NO: 299)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	GGDLSHYYY LDV (SEQ ID NO: 352)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SNYMS (SEQ ID NO: 310)	VIYSGGSTYYA DSVKG (SEQ ID NO: 330)	ATLLSLSYDAF DI (SEQ ID NO: 353)
ASSTGAVTSG YYPN (SEQ ID NO: 287)	STSNKHS (SEQ ID NO: 294)	LLYYGGAQW V (SEQ ID NO: 299)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	GGDLSHYYY MDV (SEQ ID NO: 354)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYWIG (SEQ ID NO: 309)	IYPGDS DTRY SPSFQG (SEQ ID NO: 329)	ERDRWFDP (SEQ ID NO: 355)

(continued)

HLA-A*03 CDRs (not according to the invention)					
TGTSSDVGGY NVVS (SEQ ID NO: 288)	EVSKRPS (SEQ ID NO: 295)	SSYAGSNNW V (SEQ ID NO: 300)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	ETPPSLGAFDI (SEQ ID NO: 356)
SGSSSNGSNT VN (SEQ ID NO: 289)	SNNQRPS (SEQ ID NO: 296)	AAWDDSLNG WV (SEQ ID NO: 301)	SSSYWVG (SEQ ID NO: 311)	SIYSGSTWYN PSLKS (SEQ ID NO: 331)	EAYCLSDSYW YFDL (SEQ ID NO: 357)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTWYN PSLKS (SEQ ID NO: 332)	ESWKYFYPRG YMDV (SEQ ID NO: 358)
HLA-B*07 CDRs (not according to the invention)					
RASENIYSNLA (SEQ ID NO: 290)	AATYLPD (SEQ ID NO: 297)	QHFWWTPYT (SEQ ID NO: 302)	SGYSWH (SEQ ID NO: 313)	YIHFGSTHYH PSLKS (SEQ ID NO: 333)	GGVWSHYAM DC (SEQ ID NO: 359)
HLA-A*11 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTWYN PSLKS (SEQ ID NO: 332)	HYYYM DV (SEQ ID NO: 360)

(continued)

HLA-A*11 CDRs (not according to the invention)						
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	TSGVGVG (SEQ ID NO: 314)	LIYWDDKRY SPSLKS (SEQ ID NO: 334)	KTTSFYFDY (SEQ ID NO: 361)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	HYYYMDV (SEQ ID NO: 362)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYWMH (SEQ ID NO: 315)	RINSDGSSTSY ADSVKG (SEQ ID NO: 335)	GVLNWFDP (SEQ ID NO: 363)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	HYYYLDV (SEQ ID NO: 364)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYDMH (SEQ ID NO: 316)	AIGTAGDTYY PGSVKG (SEQ ID NO: 336)	DLPGSWYFD L (SEQ ID NO: 365)	

(continued)

HLA-A*11 CDRs (not according to the invention)						
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYAMH (SEQ ID NO: 317)	WINAGNGNT KYSQKFQG (SEQ ID NO: 337)	EGNGANPDA FDI (SEQ ID NO: 366)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	TSGVGVG (SEQ ID NO: 314)	LIYWDDKRY SPSLKS (SEQ ID NO: 334)	RHMLSCFDY (SEQ ID NO: 367)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTYNN PSLKS (SEQ ID NO: 332)	HYYYSM DV (SEQ ID NO: 368)	
HLA-C*07 CDRs (not according to the invention)						
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYAMS (SEQ ID NO: 318)	AISGGGGSTYY ADSVKG (SEQ ID NO: 338)	SFDWFDP (SEQ ID NO: 369)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTYNN PSLKS (SEQ ID NO: 332)	ERSISPYYYY MDV (SEQ ID NO: 370)	

(continued)

HLA-C*07 CDRs (not according to the invention)					
SGSSNIGSNT VN (SEQ ID NO: 289)	SNNQRPS (SEQ ID NO: 296)	AAWDDSLNG WV (SEQ ID NO: 301)	SSSYWYG (SEQ ID NO: 311)	SIYSGSTWYN PSLKS (SEQ ID NO: 331)	DSVIWYWF P (SEQ ID NO: 371)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTWYN PSLKS (SEQ ID NO: 332)	EEILPRLSYWY MDV (SEQ ID NO: 372)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYAMN (SEQ ID NO: 319)	WINTNTGNP TYAQGFTG (SEQ ID NO: 339)	GGRAHSSWY FDL (SEQ ID NO: 373)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTWYN PSLKS (SEQ ID NO: 332)	DRIKILPRLGY YYMDV (SEQ ID NO: 374)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTWYN PSLKS (SEQ ID NO: 332)	DTVIHYWY DV (SEQ ID NO: 375)

(continued)

HLA-C*07 CDRs (not according to the invention)						
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	DVIVEFLSY YMDV (SEQ ID NO: 376)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	DIFIHYYYM DV (SEQ ID NO: 377)	DGTFYSYSPYY FDY (SEQ ID NO: 378)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYSMN (SEQ ID NO: 306)	YISSSSSTIYYA DSVKG (SEQ ID NO: 327)	DGTFYSYSPYY FDY (SEQ ID NO: 378)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	EWIKILPRLGY YYMDV (SEQ ID NO: 379)	

(continued)

HLA-C*07 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DRSLYYYYM DV (SEQ ID NO: 380)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DKILAPNYYY MDV (SEQ ID NO: 381)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	EKSWKYFYY YYMDV (SEQ ID NO: 382)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	ENTSTIPYYY YMDV (SEQ ID NO: 383)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	EDVDKNTSTI YYYYYMDV (SEQ ID NO: 384)

(continued)

HLA-C*07 CDRs (not according to the invention)						
RASQGISSAL A (SEQ ID NO: 291)	DASSLES (SEQ ID NO: 55)	QQFNSYPLT (SEQ ID NO: 60)	DYYMS (SEQ ID NO: 320)	YISSSGSTIYYA DSVKG (SEQ ID NO: 340)	DGGDIVSSAI YWYFDL (SEQ ID NO: 385)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTIYYN PSLKS (SEQ ID NO: 332)	DLILPPYVYY MDV (SEQ ID NO: 386)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTIYYN PSLKS (SEQ ID NO: 332)	ETWIKILPRYY YVYYMDV (SEQ ID NO: 387)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWWS (SEQ ID NO: 312)	YIYSGSTIYYN PSLKS (SEQ ID NO: 332)	DLSRYVYYM DV (SEQ ID NO: 388)	
RASQGISSWL A (SEQ ID NO: 292)	AASSLQS (SEQ ID NO: 293)	QQYNSYPLT (SEQ ID NO: 303)	SYSMN (SEQ ID NO: 306)	YISSSSSTIYYA DSVKG (SEQ ID NO: 327)	EHIVLCFDY (SEQ ID NO: 389)	

(continued)

HLA-C*07 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	DKILRPYYYY YMDV (SEQ ID NO: 390)
TGTSSDVGGY NWS (SEQ ID NO: 288)	EVSKRPS (SEQ ID NO: 295)	SSYAGSNNW V (SEQ ID NO: 300)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQLQGG (SEQ ID NO: 326)	GSNEYFQH (SEQ ID NO: 391)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYAMN (SEQ ID NO: 319)	WININIGNP TYAQQFTG (SEQ ID NO: 339)	GTSYWFYDL (SEQ ID NO: 392)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	EEIVEVFYYY MDV (SEQ ID NO: 393)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYAMS (SEQ ID NO: 318)	AISGSGGSTYY ADSVKG (SEQ ID NO: 338)	VDDYFDDY (SEQ ID NO: 394)

(continued)

HLA-C*07 CDRs (not according to the invention)						
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYWMH (SEQ ID NO: 315)	RINSDGSSTSY ADSVKG (SEQ ID NO: 335)	STNILLSYTKA FDI (SEQ ID NO: 395)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DKTYYYYYYMM DV (SEQ ID NO: 396)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	EKYFHDKYFH DYYYYYMDV (SEQ ID NO: 397)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DTSVYYYYYMM DV (SEQ ID NO: 398)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	EKILPYYYYY MDV (SEQ ID NO: 399)	

(continued)

HLA-C*07 CDRs (not according to the invention)					
SGSSSNIGSNT VN (SEQ ID NO: 289)	SNNQRPS (SEQ ID NO: 296)	AAWDDSLG WV (SEQ ID NO: 301)	SYSMN (SEQ ID NO: 306)	YISSSSSTIYYA DSVKG (SEQ ID NO: 327)	QWIYIYNPR GFIFLHDAFDI (SEQ ID NO: 400)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SNSAAWN (SEQ ID NO: 321)	RTYRSKWYN DYAVSVKS (SEQ ID NO: 341)	EDVDFHHDA FDI (SEQ ID NO: 401)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	EGVDKNTSTI YYYYYMDV (SEQ ID NO: 402)
RASQGISSWL A (SEQ ID NO: 292)	AASSLQS (SEQ ID NO: 293)	QQYNSYPLT (SEQ ID NO: 303)	SYSMN (SEQ ID NO: 306)	YISSSSSTIYYA DSVKG (SEQ ID NO: 327)	DRRGYFDL (SEQ ID NO: 403)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	DYYMH (SEQ ID NO: 322)	LVDPEGETIY AEKFQG (SEQ ID NO: 342)	GIHVDIRSME DWFDP (SEQ ID NO: 404)

(continued)

HLA-C*07 CDRs (not according to the invention)						
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DIGTSYYYYM DV (SEQ ID NO: 405)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	EVVEVFLYYY YMDV (SEQ ID NO: 406)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DLYYYYYYM DV (SEQ ID NO: 407)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	ESWKYFYPRG SIFIHYYYMD V (SEQ ID NO: 408)	
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYWN PSLKS (SEQ ID NO: 332)	DRIVEVYYYY MDV (SEQ ID NO: 409)	

(continued)

HLA-C*07 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	EKYFHDWLYY YYMDV (SEQ ID NO: 410)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYWS (SEQ ID NO: 312)	YIYSGSTYYN PSLKS (SEQ ID NO: 332)	DLVDKNTSYY YYMDV (SEQ ID NO: 411)
TGTSSDVGGY NYVS (SEQ ID NO: 288)	EVSKRPS (SEQ ID NO: 295)	SSYAGSNNW V (SEQ ID NO: 300)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	VQNEYFQH (SEQ ID NO: 412)
RASQGISSWL A (SEQ ID NO: 292)	AASSLQS (SEQ ID NO: 293)	QQANSFPLT (SEQ ID NO: 304)	DYYMS (SEQ ID NO: 320)	YISSSGSTIYYA DSVKG (SEQ ID NO: 340)	ANWFDP (SEQ ID NO: 413)

(continued)

HLA-A*01 CDRs (not according to the invention)					
ASSTGAVTSG YYPN (SEQ ID NO: 287)	STSNKHS (SEQ ID NO: 294)	LLYYGGAQW V (SEQ ID NO: 299)	SYGIS (SEQ ID NO: 305)	WISAYNGNT NYAQKLQG (SEQ ID NO: 326)	GGWTAWYYY MDV (SEQ ID NO: 414)
SGSSNIGSNT VN (SEQ ID NO: 289)	SNNQRPS (SEQ ID NO: 296)	AAWDDSLNG WV (SEQ ID NO: 301)	SYSMN (SEQ ID NO: 306)	YISSSSSTIYYA DSVKG (SEQ ID NO: 327)	AKYYYMDV (SEQ ID NO: 415)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYYS (SEQ ID NO: 312)	YIYSGSTIYYN PSLKS (SEQ ID NO: 332)	DQVDKNTYYY YMDV (SEQ ID NO: 416)
RASQGISSWL A (SEQ ID NO: 292)	AASSLQS (SEQ ID NO: 293)	QQANSFPLT (SEQ ID NO: 304)	DYYMS (SEQ ID NO: 320)	YISSSGSTIYYA DSVKG (SEQ ID NO: 340)	ACQLAEYFQH (SEQ ID NO: 417)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYYS (SEQ ID NO: 312)	YIYSGSTIYYN PSLKS (SEQ ID NO: 332)	DRVDKNTSY YMDV (SEQ ID NO: 418)

(continued)

HLA-A*01 CDRs (not according to the invention)					
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SSNWWG (SEQ ID NO: 323)	YIYSGSTYNN PSLKS (SEQ ID NO: 332)	RVQLKLVHW FDP (SEQ ID NO: 419)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SYDIN (SEQ ID NO: 324)	WMNPNSGN TGQAQKFQG (SEQ ID NO: 343)	YYDYVTVFYF QH (SEQ ID NO: 420)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	SGGYSWS (SEQ ID NO: 325)	YIYHSGSTYNN PSLKS (SEQ ID NO: 344)	ESYPSFYAFDI (SEQ ID NO: 421)
RASQSISSYLN (SEQ ID NO: 286)	AASSLQS (SEQ ID NO: 293)	QQSYSTPLT (SEQ ID NO: 298)	TSGVGVG (SEQ ID NO: 314)	LIYWDDDKRY SPSLKS (SEQ ID NO: 334)	SNMWSYSLN DYYFDY (SEQ ID NO: 422)

[0216] In some embodiments, the non-target antigen comprising HLA-A*02, and the ligand binding domain of the second receptor comprises an HLA-A*02 ligand binding domain. In some embodiments, the ligand binding domain binds HLA-A*02 independent of the peptide in a pMHC complex comprising HLA-A*02. In some embodiments, the HLA-A*02 ligand binding domain comprises an scFv domain. In some embodiments, the HLA-A*02 ligand binding domain comprises a sequence of any one of SEQ ID NOS: 89-100. In some embodiments, the HLA-A*02 ligand binding domain comprises a sequence at least 90%, at least 95% or at least 99% identical to a sequence of any one of SEQ ID NOS: 89-100.

[0217] As described herein, the HLA-A*02 scFv comprises the complementarity determined regions (CDRs) of any one of SEQ ID NOS: 101-112. As described herein, the scFv comprises a sequence at least 95% identical to any one of SEQ ID NOS: 101-112. As described herein, the scFv comprises a sequence identical to any one of SEQ ID NOS: 101-112. As described herein, the heavy chain of the antigen binding domain comprises the heavy chain CDRs of any one of SEQ ID NOS: 101-112, and wherein the light chain of the antigen binding domain comprises the light chain CDRs of any one of SEQ ID NOS: 101-112. As described herein, the HLA-A*02 antigen binding domain comprises a heavy chain and a light chain, and the heavy chain comprises CDRs selected from SEQ ID NOS: 104-106 and 110-112 and the light chain comprises CDRs selected from SEQ ID NOS: 101-103 and 107-109. It is described herein for any of the ligand binding domains that each CDR sequence may have 1, 2, 3 or more substitutions, insertions, or deletions. CDR sequences may tolerate substitutions, deletions, or insertions. Using sequence alignment tools, routine experimentation, and known assays, those of skill in the art may generate and test variant sequences having 1, 2, 3, or more substitutions, insertions, or deletions in CDR sequences without undue experimentation.

[0218] As described herein, the HLA-A*02 antigen binding domain comprises a heavy chain and a light chain, and the heavy chain comprises a sequence at least 95% identical to the heavy chain portion of any one of SEQ ID NOS: 89-100, and the light chain comprises a sequence at least 95% identical to the light chain portion of any one of SEQ ID NOS: 89-100.

[0219] As described herein, the heavy chain comprises a sequence identical to the heavy chain portion of any one of SEQ ID NOS: 89-100, and wherein the light chain comprises a sequence identical to the light chain portion of any one of SEQ ID NOS: 89-100.

[0220] Not according to the invention: As also described herein, the non-target antigen comprises HLA-B*07, and the ligand binding domain of the second receptor comprises an HLA-B*07 ligand binding domain. As also described herein, the ligand binding domain binds HLA-B*07 independent of the peptide in a pMHC complex comprising HLA-B*07. As also described herein, the HLA-B*07 ligand binding domain comprises an scFv domain. As also described herein, the HLA-B*07 ligand binding domain comprises a sequence of any one of the indicated B7 binding domains in Table 4. As also described herein, the HLA-B*07 ligand binding domain comprises a sequence at least 90%, at least 95% or at least 99% identical to a sequence of any one of the indicated HLA-B*07 binding domains in Table 4. As also described herein, the HLA-B*07 scFv comprises any one of the indicated HLA-B*07 complementarity determined regions (CDRs) in Table 5. As also described herein, the scFv comprises a sequence at least 95% identical to any one of the indicated HLA-B*07 scFv sequences in Table 4. As also described herein, the scFv comprises a sequence identical to any one of the indicated HLA-B*07 scFv sequences in Table 4.

[0221] Not according to the invention: As also described herein, the non-target antigen comprises HLA-A*11, and the ligand binding domain of the second receptor comprises an HLA-A*11 ligand binding domain. As also described herein, the ligand binding domain binds HLA-A*11 independent of the peptide in a pMHC complex comprising HLA-A*11. As also described herein, the HLA-A*11 ligand binding domain comprises an scFv domain. As also described herein, the HLA-A*11 ligand binding domain comprises a sequence of any one of the indicated HLA-A*11 binding domains in Table 4. As also described herein, the HLA-A*11 ligand binding domain comprises a sequence at least 90%, at least 95% or at least 99% identical to a sequence of any one of the indicated HLA-A*11 binding domains in Table 4. As also described herein, the HLA-A*11 scFv comprises any one of the indicated HLA-A*11 complementarity determined regions (CDRs) in Table 5. As also described herein, the scFv comprises a sequence at least 95% identical to any one of the indicated HLA-A*11 scFv sequences in Table 4. As also described herein, the scFv comprises a sequence identical to any one of the indicated HLA-A*11 scFv sequences in Table 4.

[0222] Not according to the invention: As described herein, the non-target antigen comprises HLA-A*01, and the ligand binding domain of the second receptor comprises an HLA-A*01 ligand binding domain. As also described herein, the ligand binding domain binds HLA-A*01 independent of the peptide in a pMHC complex comprising HLA-A*01. As also described herein, the HLA-A*01 ligand binding domain comprises an scFv domain. As also described herein, the HLA-A*01 ligand binding domain comprises a sequence of any one of the indicated HLA-A*01 binding domains in Table 4. As also described herein, the HLA-A*01 ligand binding domain comprises a sequence at least 90%, at least 95% or at least 99% identical to a sequence of any one of the indicated HLA-A*01 binding domains in Table 4. As also described herein, the HLA-A*01 scFv comprises any one of the indicated HLA-A*01 complementarity determined regions (CDRs) in Table 5. As also described herein, the scFv comprises a sequence at least 95% identical to any one of the indicated HLA-A*01 scFv sequences in Table 4. As also described herein, the scFv comprises a sequence identical to any one of the indicated HLA-A*01 scFv sequences in Table 4.

[0223] Not according to the invention: As also described herein, the non-target antigen comprises HLA-A*03, and the

ligand binding domain of the second receptor comprises an HLA-A*03 ligand binding domain. As also described herein, the ligand binding domain binds HLA-A*03 independent of the peptide in a pMHC complex comprising HLA-A*03. As also described herein, the HLA-A*03 ligand binding domain comprises an scFv domain. As also described herein, the HLA-A*03 ligand binding domain comprises a sequence of any one of the indicated HLA-A*03 binding domains in Table 4. As also described herein, the HLA-A*03 ligand binding domain comprises a sequence at least 90%, at least 95% or at least 99% identical to a sequence of any one of the indicated HLA-A*03 binding domains in Table 4. As also described herein, the HLA-A*03 scFv comprises any one of the indicated HLA-A*03 complementarity determined regions (CDRs) in Table 5. As also described herein, the scFv comprises a sequence at least 95% identical to any one of the indicated HLA-A*03 scFv sequences in Table 4. As also described herein, the scFv comprises a sequence identical to any one of the indicated HLA-A*03 scFv sequences in Table 4.

[0224] Not according to the invention: As also described herein, the non-target antigen comprises HLA-C*07, and the ligand binding domain of the second receptor comprises an HLA-C*07 ligand binding domain. As also described herein, the ligand binding domain binds HLA-C*07 independent of the peptide in a pMHC complex comprising HLA-C*07. As also described herein, the HLA-C*07 ligand binding domain comprises an scFv domain. As also described herein, the HLA-C*07 ligand binding domain comprises a sequence of any one of the indicated HLA-C*07 binding domains in Table 4. As also described herein, the HLA-C*07 ligand binding domain comprises a sequence at least 90%, at least 95% or at least 99% identical to a sequence of any one of the indicated HLA-C*07 binding domains in Table 4. As also described herein, the HLA-C*07 scFv comprises any one of the indicated HLA-C*07 complementarity determined regions (CDRs) in Table 5. As also described herein, the scFv comprises a sequence at least 95% identical to any one of the indicated HLA-C*07 scFv sequences in Table 4. As also described herein, the scFv comprises a sequence identical to any one of the indicated HLA-C*07 scFv sequences in Table 4.

Inhibitory Receptors

[0225] The disclosure provides a second receptor that is an inhibitory chimeric antigen receptor (*i.e.* inhibitory receptor). The inhibitory receptor may comprise an extracellular ligand binding domain that binds to and recognizes the non-target antigen or a peptide derivative thereof in an MHC-1 complex.

[0226] Exemplary inhibitory receptors are described in PCT/US2020/045228 filed on September 6, 2020, PCT/US2020/064607, filed on December 11, 2020, PCT/US2021/029907, filed on April 29, 2021 and PCT/US2020/059856 filed on November 10, 2020.

[0227] The non-target antigen can be a non-target allelic variant. A "non-target allelic variant" as used herein refers to an allele (*e.g.* an allelic variant of an MHC Class I gene), whose expression is reduced or eliminated in a target cell (*e.g.* a cancer cell) due to loss of heterozygosity, or another mechanism of expression loss, but is expressed or detectably expressed in a normal or healthy cell.

[0228] The term "inhibitory chimeric antigen receptor" or "inhibitory receptor" as used herein refers to an antigen-binding domain that is fused to an intracellular signaling domain capable of transducing an inhibitory signal that inhibits or suppresses the immune activity of an immune cell. Inhibitory receptors have immune cell inhibitory potential, and are distinct and distinguishable from CARs, which are receptors with immune cell activating potential. For example, CARs are activating receptors as they include intracellular stimulatory and/or co-stimulatory domains. Inhibitory receptors are inhibiting receptors that contain intracellular inhibitory domains.

[0229] As used herein "inhibitory signal" refers to signal transduction or changes in protein expression in an immune cell resulting in suppression of an immune response (*e.g.*, decrease in cytokine production or reduction of immune cell activation). Inhibition or suppression of an immune cell can be selective and/or reversible, or not selective and/or reversible. Inhibitory receptors are responsive to non-target antigens (*e.g.* HLA-A*02). For example, when a non-target antigen (*e.g.* HLA-A*02) binds to or contacts the inhibitory receptor, the inhibitory receptor is responsive and activates an inhibitory signal in the immune cell expressing the inhibitory receptor upon binding of the non-target antigen by the extracellular ligand binding domain of the inhibitory receptor.

[0230] Inhibitory receptors of the disclosure may comprise an extracellular ligand binding domain.

[0231] The ligand binding domain may be an antigen binding domain (not according to the invention). Exemplary antigen binding domains include, *inter alia*, scFv, SdAb, V β -only domains, and TCR antigen binding domains derived from the TCR α and β chain variable domains.

[0232] As described herein, the extracellular ligand binding domain of the second receptor binds to and recognizes a polymorphic variant of COLEC12, APCDD1, CXCL16 or an antigen peptide thereof in a complex with a major histocompatibility complex class I (MHC-I), or HLA-A*02. In some embodiments, the extracellular ligand binding domain of the second receptor is an scFv.

[0233] In some embodiments, the extracellular ligand binding domain of the second receptor is fused to the extracellular domain of an inhibitory receptor.

[0234] In some embodiments, the inhibitory receptors of the present disclosure comprise an extracellular hinge region.

Exemplary hinges can be isolated or derived from IgD and CD8 domains, for example IgG1. The hinge may be isolated or derived from CD8 α or CD28.

[0235] The inhibitory receptors of the present disclosure can be designed to comprise a transmembrane domain that is fused to the extracellular domain of the inhibitory receptors. In some instances, the transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins to minimize interactions with other members of the receptor complex.

[0236] The transmembrane domain may be derived either from a natural or from a synthetic source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein. Transmembrane regions may be isolated or derived from (i.e. comprise at least the transmembrane region(s) of) the alpha, beta or zeta chain of the T-cell receptor, CD28, CD3 epsilon, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154, or from an immunoglobulin such as IgG4. Alternatively, the transmembrane domain may be synthetic, in which case it will comprise predominantly hydrophobic residues such as leucine and valine. In some embodiments, a triplet of phenylalanine, tryptophan and valine will be found at each end of a synthetic transmembrane domain. Optionally, a short oligo- or polypeptide linker, preferably between 2 and 10 amino acids in length may form the linkage between the transmembrane domain and the intracellular domain of the inhibitory receptors. A glycine-serine doublet provides a particularly suitable linker.

[0237] The disclosure provides inhibitory receptors comprising an intracellular domain. The intracellular domain of the inhibitory receptors of the instant disclosure is responsible for inhibiting activation of the immune cells comprising the inhibitory receptors, which would otherwise be activated in response to activation signals by the first receptor. The inhibitory intracellular domain may comprise an immunoreceptor tyrosine-based inhibitory motif (ITIM). The inhibitory intracellular domain comprising an ITIM can be isolated or derived from an immune checkpoint inhibitor such as CTLA-4 and PD-1. CTLA-4 and PD-1 are immune inhibitory receptors expressed on the surface of T cells, and play a pivotal role in attenuating or terminating T cell responses.

[0238] An inhibitory intracellular domain may be isolated from human tumor necrosis factor related apoptosis inducing ligand (TRAIL) receptor and CD200 receptor 1. The TRAIL receptor comprises TR10A, TR10B or TR10D.

[0239] An inhibitory intracellular domain may be isolated from phosphoprotein membrane anchor with glycosphingolipid microdomains 1 (PAG1). In some embodiments of the present invention, an inhibitory intracellular domain is isolated from leukocyte immunoglobulin like receptor B1 (LILRB 1).

[0240] The inhibitory domain may be isolated or derived from a human protein, for example a human TRAIL receptor, CTLA-4, PD-1, PAG1 or LILRB1 protein.

[0241] In some embodiments, the inhibitory domain comprises an intracellular domain, a transmembrane or a combination thereof. In some embodiments, the inhibitory domain comprises an intracellular domain, a transmembrane domain, a hinge region or a combination thereof.

[0242] The inhibitory domain may be isolated or derived from killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 2 (KIR3DL2), killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 3 (KIR3DL3), leukocyte immunoglobulin like receptor B1 (LIR1, also called LIR-1 and LILRB1), programmed cell death 1 (PD-1), Fc gamma receptor IIB (Fc γ R1IB), killer cell lectin like receptor K1 (NKG2D), CTLA-4, a domain containing a synthetic consensus ITIM, a ZAP70 SH2 domain (e.g., one or both of the N and C terminal SH2 domains), or ZAP70 KI_K369A (kinase inactive ZAP70).

[0243] In some embodiments, the inhibitory domain is isolated or derived from a human protein.

[0244] In some embodiments, the second, inhibitory receptor comprises an inhibitory domain. In some embodiments, the second, inhibitory receptor comprises an inhibitory intracellular domain and/or an inhibitory transmembrane domain. In some embodiments, the inhibitory intracellular domain is fused to an intracellular domain of an inhibitory receptor. In some embodiments, the inhibitory intracellular domain is fused to the transmembrane domain of an inhibitory receptor.

[0245] In some embodiments, the second, inhibitory receptor comprises a cytoplasmic domain, a transmembrane domain, and an extracellular domain or a portion thereof isolated or derived isolated or derived from the same protein, for example an ITIM containing protein. In some embodiments, the second, inhibitory receptor comprises a hinge region isolated or derived from the same protein as the intracellular domain and/or transmembrane domain, for example an ITIM containing protein.

[0246] In some embodiments, the second receptor is a TCR comprising an inhibitory domain (an inhibitory TCR). In some embodiments, the inhibitory TCR comprises an inhibitory intracellular domain and/or an inhibitory transmembrane domain. In some embodiments, the inhibitory intracellular domain is fused to the intracellular domain of TCR alpha, TCR beta, CD3 delta, CD3 gamma or CD3 epsilon or a portion thereof a TCR. In some embodiments, the inhibitory intracellular domain is fused to the transmembrane domain of TCR alpha, TCR beta, CD3 delta, CD3 gamma or CD3 epsilon.

[0247] In some embodiments, the second receptor is a TCR comprising an inhibitory domain (an inhibitory TCR). In some embodiments, the inhibitory domain is isolated or derived from LILRB1.

LILRB1 Inhibitory receptors

[0248] The disclosure provides a second, inhibitory receptor comprising a LILRB 1 inhibitory domain, and optionally, a LILRB1 transmembrane and/or hinge domain, or functional variants thereof. The inclusion of the LILRB1 transmembrane domain and/or the LILRB1 hinge domain in the inhibitory receptor may increase the inhibitory signal generated by the inhibitory receptor compared to a reference inhibitory receptor having another transmembrane domain or another hinge domains. The second, inhibitory receptor comprising the LILRB 1 inhibitory domain may be a CAR or TCR, as described herein. Any suitable ligand binding domain, as described herein, may be fused to the LILRB1-based second, inhibitory receptors.

[0249] Leukocyte immunoglobulin-like receptor subfamily B member 1 (LILRB1), also known as Leukocyte immunoglobulin-like receptor B1, as well as ILT2, LIR1, MIR7, PIRB, CD85J, ILT-2 LIR-1, MIR-7 and PIR-B, is a member of the leukocyte immunoglobulin-like receptor (LIR) family. The LILRB1 protein belongs to the subfamily B class of LIR receptors. These receptors contain two to four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The LILRB 1 receptor is expressed on immune cells, where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. LILRB 1 is thought to regulate inflammatory responses, as well as cytotoxicity, and to play a role in limiting auto-reactivity. Multiple transcript variants encoding different isoforms of LILRB1 exist.

[0250] The inhibitory receptor may comprise one or more domains isolated or derived from LILRB1. For the receptors having one or more domains isolated or derived from LILRB1, the one or more domains of LILRB 1 may comprise an amino acid sequence that is at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or may be identical to a sequence or subsequence of SEQ ID NO: 113. The one or more domains of LILRB 1 may comprise an amino acid sequence that is identical to a sequence or subsequence of SEQ ID NO: 113. The one or more domains of LILRB 1 may consist of an amino acid sequence that is at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or may be identical to a sequence or subsequence of SEQ ID NO: 113. The one or more domains of LILRB 1 may consist of an amino acid sequence that is identical to a sequence or subsequence of SEQ ID NO: 113.

[0251] For the receptors having one or more domains isolated or derived from LILRB1, the one or more domains of LILRB 1 may be encoded by a polynucleotide sequence that is at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or may be identical to a sequence or subsequence of

```

1  aaatgagttt  taaaaaggct  tgtccaggaa  gcacatatgg  gagctgggtca  ctctgcattt
61  tgggccctcc  tggaggtggt  tagaccttcc  gagagagaaa  ctgagacaca  tgagagggaa
121  gaaatgactc  agtgggtgag  ccctgtggag  tcccacccac  aaccagcaca  ctgtgaccca
181  ctgcacaaac  ctctagccca  cagctcactt  cctcctttta  gaagagaaga  gaaaagagga
241  gaggagagga  ggaacagaaa  agaaaagaaa  agaaaagtgt  ggaaacaaat  aatctaagaa
301  tgaggagaaa  gcaagaagag  tgacccctt  gtgggcactc  cattggtttt  atggcgctc
361  tactttctgg  agtttgtgta  aaacaaaaat  attatgggtc  ttgtgcacat  ttacatcaag
421  ctgagcctgg  gcggcacagc  cagatgctg  atgctctct  gctgatctga  gtctgcctgc
481  agcatggacc  tgggtcttcc  ctgaagcatc  tccagggtg  gagggacgac  tgccatgcac
541  cgagggtc  tccatccaca  gagcagggca  gtgggaggag  acgcatgac  ccccatcctc
601  acggtcctga  tctgtctcgg  gctgagtctg  ggcccccgga  cccacgtgca  ggcagggcac
661  ctccccaagc  ccaccctctg  ggctgaacca  ggctctgtga  tcaccagggt  gagtcctgtg
721  accctcaggt  gtcagggggg  ccaggagacc  caggagtacc  gtctatatag  agaaaagaaa
781  acagcaccct  ggattacacg  gatccacacg  gagcttgtga  agaagggtca  gttccccatc
841  ccatccatca  cctgggaaca  cacaggcggt  tatcgctgtt  actatggtag  cgacactgca
901  ggccgctcag  agagcagtga  cccctgggag  ctggtggtga  caggagccta  catcaaacc
961  accctctcag  cccagcccag  cccgtggtg  aactcaggag  ggaatgtaac  cctccagtgt
1021  gactcacagg  tggcatttga  tggcttcatt  ctgtgtaagg  aaggagaaga  tgaacacca

```

```

1081 caatgcctga actcccagcc ccatgcccgt gggtcgtccc gcgccatctt ctccgtgggc
1141 cccgtgagcc cgagtcgcag gtgggtgtac aggtgctatg ctatgactc gaactctccc
1201 tatgagtggt ctctaccag tgatctcctg gagctcctgg tcctaggtgt ttctaagaag
1261 ccatcactct cagtgcagcc aggtcctatc gtggcccctg aggagaccct gactctgcag
5 1321 tgtggctctg atgctggcta caacagattt gttctgtata aggacgggga acgtgacttc
1381 cttcagctcg ctggcgca caacagattt gttctgtata aggacgggga acgtgacttc
1441 cctgtgagcc gctcctacgg gggccagtag agatgctacg gtgcacacaa cctctcctcc
1501 gagtggtcgg ccccagcga cccctggac atcctgatcg caggacagtt ctatgacaga
1561 gtctccctct cgggtgcagcc gggcccacg gtggcctcag gagagaacgt gaccctgctg
10 1621 tgtcagtcac agggatggat gcaaaacttt cttctgacca aggagggggc agctgatgac
1681 ccatggcgct taagatcaac gtaccaatct caaaaatacc aggtggaatt ccccatgggt
1741 cctgtgacct cagcccatgc ggggacctac aggtgctacg gctcacagag ctccaaaccc
1801 tacctgctga ctacccccag tgacccctcg gagctcgtgg tctcaggacc gtctgggggc
1861 cccagctccc cgacaacagg cccacacctc acatctggcc ctgaggacca gcccctcacc
15 1921 cccaccgggt cggatcccca gagtggctcg ggaaggcacc tgggggttgt gatcggcatc
1981 ttgggtggccg tcactcctact gctcctcctc ctctcctcc tcttctcat cctccgacat
2041 cgacgtcagg gcaaacactg gacatcgacc cagagaaagg ctgatttcca acatcctgca
2101 ggggctgtgg ggccagagcc cacagacaga ggctgcagt ggaggtccag cccagctgcc
2161 gatgccagga aagaaaacct ctatgctgac gtgaagcaca cacagcctga ggatgggggtg
2221 gagatggaca ctgggagccc acacgatgaa gacccccagg cagtgcagta tgccgaggtg
20 2281 aaacactcca gacctaggag agaaatggcc tctcctcctt cccactgtgc tggggaattc
2341 ctggacacaa aggacagaca ggcggaagag gacaggcaga tggactgca ggctgctgca
2401 tctgaagccc cccaggatgt gacctacgcc cagctgcaca gcttgaccct cagacgggag
2461 gcaactgagc ctctccatc ccaggaaggg ccctctccag ctgtgcccag catctacgcc
25 2521 actctggcca tccactagcc caggggggga cgcagacccc aactccatg gactctggaa
2581 tgcattggag ctgccccccc agtggacacc attggacccc accagcctg gatctacccc
2641 aggagactct gggaaactttt aggggtcact caattctgca gtataaataa ctaatgtctc
2701 tacaattttt aaataaagca acagacttct caataatcaa tgaagtagct gagaaaacta
2761 agtcagaaag tgcattaaac tgaatcacia tgtaaatatt acacatcaag cgatgaaact
2821 ggaaaactac aagccacgaa tgaatgaatt aggaagaaa aaaagtagga aatgaatgat
30 2881 cttggctttt ctataagaaa tttagggcag ggcacggtgg ctacgcctg taattccagc
2941 actttgggag gccgaggcgg gcagatcacg agttcaggag atcgagacca tcttgcccaa
3001 catggtgaaa cctgtctct ctaaaaaata caaaaattag ctggatgtgg tggcagtgcc
3061 tgtaatccca gctatttggg aggtgaggc aggagaatcg cttgaaccag ggagtcagag
3121 gtttcagtga gccaagatcg caccactgct ctccagcctg gcgacagagg gagactccat
3181 ctcaaattaa aaaaaaaaaa aaaaaagaaa gaaaaaaaaa aaaaaaaaaa (SEQ ID NO: 114).
35

```

[0252] For the receptors having one or more domains of LILRB 1, the one or more domains of LILRB 1 may be encoded by a polynucleotide sequence that is identical to a sequence or subsequence of SEQ ID NO: 114.

[0253] An inhibitory receptor may comprise a polypeptide, wherein the polypeptide comprises one or more of: an LILRB1 hinge domain or functional fragment or variant thereof; an LILRB 1 transmembrane domain or a functional variant thereof; and an LILRB 1 intracellular domain or an intracellular domain comprising at least one, or at least two immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAEV (SEQ ID NO: 116), VTAYL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0254] As used herein an "immunoreceptor tyrosine-based inhibitory motif" or "ITIM" may refer to a conserved sequence of amino acids with a consensus sequence of S/I/V/LxYxxI/V/L (SEQ ID NO: 423), that is found in the cytoplasmic tails of many inhibitory receptors of the immune system. After ITIM-possessing inhibitory receptors interact with their ligand, the ITIM motif may be phosphorylated, allowing the inhibitory receptor to recruit other enzymes, such as the phosphotyrosine phosphatases SHP-1 and SHP-2, or the inositol-phosphatase called SHIP.

[0255] In some embodiments, the polypeptide may comprise an intracellular domain comprising at least one immunoreceptor tyrosine-based inhibitory motif (ITIM), at least two ITIMs, at least 3 ITIMs, at least 4 ITIMs, at least 5 ITIMs or at least 6 ITIMs. In some embodiments, the intracellular domain may have 1, 2, 3, 4, 5, or 6 ITIMs.

[0256] In some embodiments, the polypeptide may comprise an intracellular domain comprising at least one ITIM selected from the group of ITIMs consisting of NLYAAV (SEQ ID NO: 115), VTAEV (SEQ ID NO: 116), VTAYL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0257] In further particular embodiments, the polypeptide may comprise an intracellular domain comprising at least two immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM is independently selected from NLYAAV (SEQ ID NO: 115), VTAEV (SEQ ID NO: 116), VTAYL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0258] In some embodiments, the intracellular domain may comprise both ITIMs NLYAAV (SEQ ID NO: 115) and VTAEV (SEQ ID NO: 116). In some embodiments, the intracellular domain may comprise a sequence at least 95%

identical to SEQ ID NO: 119. In some embodiments, the intracellular domain may comprise or consist essentially of a sequence identical to SEQ ID NO: 119.

[0259] In some embodiments, the intracellular domain may comprise both ITIMs VTAAEV (SEQ ID NO: 116) and VTAAQL (SEQ ID NO: 117). In some embodiments, the intracellular domain may comprise a sequence at least 95% identical to SEQ ID NO: 120. In some embodiments, the intracellular domain may comprise or consist essentially of a sequence identical to SEQ ID NO: 120.

[0260] In some embodiments, the intracellular domain may comprise both ITIMs VTAAQL (SEQ ID NO: 117) and SIYATL (SEQ ID NO: 118). In some embodiments, the intracellular domain may comprise a sequence at least 95% identical to SEQ ID NO: 121. In some embodiments, the intracellular domain may comprise or consist essentially of a sequence identical to SEQ ID NO: 121.

[0261] In some embodiments, the intracellular domain may comprise the ITIMs NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), and VTAAQL (SEQ ID NO: 117). In some embodiments, the intracellular domain may comprise a sequence at least 95% identical to SEQ ID NO: 122. In some embodiments, the intracellular domain may comprise or consist essentially of a sequence identical to SEQ ID NO: 122.

[0262] In some embodiments, the intracellular domain may comprise the ITIMs VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118). In some embodiments, the intracellular domain may comprise a sequence at least 95% identical to SEQ ID NO: 123. In some embodiments, the intracellular domain comprises or consists essentially of a sequence identical to SEQ ID NO: 123.

[0263] In some embodiments, the intracellular domain may comprise the ITIMs NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118). In some embodiments, the intracellular domain may comprise a sequence at least 95% identical to SEQ ID NO: 124. In some embodiments, the intracellular domain comprises or consists essentially of a sequence identical to SEQ ID NO: 124.

[0264] In some embodiments of the present invention, the intracellular domain comprises a sequence at least 95% identical to the LILRB1 intracellular domain (SEQ ID NO: 129). In some embodiments of the present invention, the intracellular domain comprises or consists essentially of a sequence identical to the LILRB 1 intracellular domain (SEQ ID NO: 129).

[0265] LILRB 1 intracellular domains or functional variants thereof of the disclosure can have at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, or at least 8 ITIMs. In some embodiments, the LILRB 1 intracellular domain or functional variant thereof has 2, 3, 4, 5, or 6 ITIMs.

[0266] In particular embodiments, the intracellular domain may comprise two, three, four, five, or six immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0267] In particular embodiments, the intracellular domain may comprise at least three immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0268] In particular embodiments, the intracellular domain may comprise three immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0269] In particular embodiments, the intracellular domain may comprise four immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0270] In particular embodiments, the intracellular domain may comprise five immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0271] In particular embodiments, the intracellular domain may comprise six immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0272] In particular embodiments, the intracellular domain may comprise at least seven immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0273] The LILRB1 protein has four immunoglobulin (Ig) like domains termed D1, D2, D3 and D4. In some embodiments, the LILRB1 hinge domain may comprise an LILRB1 D3D4 domain or a functional variant thereof. In some embodiments, the LILRB 1 D3D4 domain comprises a sequence at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or identical to SEQ ID NO: 125. In some embodiments, the LILRB1 D3D4 domain comprises or consists essentially of SEQ ID NO: 125.

[0274] In some embodiments of the present invention, the polypeptide comprises the LILRB 1 hinge domain or functional fragment or variant thereof. In some embodiments of the present invention, the LILRB 1 hinge domain or functional fragment or variant thereof comprises a sequence at least 95%, at least 96%, at least 97%, at least 98%, at least 99% identical

or identical to SEQ ID NO: 132, SEQ ID NO: 125, or SEQ ID NO: 126. In embodiments of the present invention, the LILRB 1 hinge domain or functional fragment or variant thereof comprises a sequence at least 95% identical to SEQ ID NO: 132, SEQ ID NO: 125, or SEQ ID NO: 126.

[0275] In some embodiments of the present invention, the LILRB1 hinge domain comprises a sequence identical to SEQ ID NO: 132, SEQ ID NO: 125, or SEQ ID NO: 126.

[0276] In some embodiments of the present invention, the LILRB 1 hinge domain consists essentially of a sequence identical to SEQ ID NO: 132, SEQ ID NO: 125, or SEQ ID NO: 126.

[0277] In some embodiments of the present invention, the transmembrane domain is a LILRB 1 transmembrane domain or a functional variant thereof. In some embodiments of the present invention, the LILRB 1 transmembrane domain or a functional variant thereof comprises a sequence at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical or at least 99% to SEQ ID NO: 133. In some embodiments of the present invention, the LILRB 1 transmembrane domain or a functional variant thereof comprises a sequence at least 95% identical to SEQ ID NO: 133. In some embodiments of the present invention, the LILRB 1 transmembrane domain comprises a sequence identical to SEQ ID NO: 133. In embodiments of the present invention, the LILRB1 transmembrane domain consists essentially of a sequence identical to SEQ ID NO: 133.

[0278] In some embodiments, the transmembrane domain can be attached to the extracellular region of the second, inhibitory receptor, *e.g.*, the antigen binding domain or ligand binding domain, via a hinge, *e.g.*, a hinge from a human protein. For example, in some embodiments, the hinge can be a human immunoglobulin (Ig) hinge, *e.g.*, an IgG4 hinge, a CD8a hinge or an LILRB 1 hinge.

[0279] In some embodiments, the second, inhibitory receptor comprises an inhibitory domain. In some embodiments, the second, inhibitory receptor comprises an inhibitory intracellular domain and/or an inhibitory transmembrane domain. In some embodiments, the inhibitory domain is isolated or derived from LILRB1.

Inhibitory Receptors Comprising Combinations of LILRB1 Domains

[0280] In some embodiments, the LILRB1-based inhibitory receptors of the disclosure comprise more than one LILRB1 domain or functional equivalent thereof. For example, in some embodiments, the inhibitory receptor comprises an LILRB1 transmembrane domain and intracellular domain, or an LILRB 1 hinge domain, transmembrane domain and intracellular domain.

[0281] In particular embodiments of the present invention, the inhibitory receptor comprises an LILRB1 hinge domain or functional fragment or variant thereof, and the LILRB1 transmembrane domain or a functional variant thereof. The polypeptide may comprise a sequence at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical, at least 99% identical or may be identical to SEQ ID NO: 127. The polypeptide may comprise a sequence at least 95% identical to SEQ ID NO: 127. The polypeptide may comprise a sequence identical to SEQ ID NO: 127.

[0282] The inhibitory receptor may comprise: the LILRB 1 transmembrane domain or a functional variant thereof, and an LILRB 1 intracellular domain and/or an intracellular domain comprising at least one immunoreceptor tyrosine-based inhibitory motif (ITIM), wherein the ITIM may be selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118). The polypeptide may comprise the LILRB 1 transmembrane domain or a functional variant thereof, and an LILRB 1 intracellular domain and/or an intracellular domain comprising at least two ITIM, wherein each ITIM may be independently selected from NLYAAV (SEQ ID NO: 115), VTAAEV (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0283] In some embodiments of the present invention, the inhibitory receptor comprises a LILRB 1 transmembrane domain and intracellular domain. In some embodiments of the present invention, the polypeptide comprises a sequence at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical, at least 99% identical or identical to SEQ ID NO: 128. In some embodiments of the present invention, the polypeptide comprises a sequence at least 95% identical to SEQ ID NO: 128. In some embodiments of the present invention, the polypeptide comprises a sequence identical to SEQ ID NO: 128.

[0284] In preferred embodiments of the present invention, the inhibitory receptor comprises: an LILRB 1 hinge domain or functional fragment or variant thereof; an LILRB 1 transmembrane domain or a functional variant thereof; and an LILRB1 intracellular domain. The inhibitory receptor may comprise an intracellular domain comprising at least two immunoreceptor tyrosine-based inhibitory motifs (ITIMs), wherein each ITIM is independently selected from LYAAV (SEQ ID NO: 115), VTAAE (SEQ ID NO: 116), VTAAQL (SEQ ID NO: 117), and SIYATL (SEQ ID NO: 118).

[0285] The inhibitory receptor may comprise a sequence at least 95% identical to SEQ ID NO: 130 or SEQ ID NO: 131, or at least 99% identical to SEQ ID NO: 130 or SEQ ID NO: 131, or identical to SEQ ID NO: 130 or SEQ ID NO: 131.

[0286] The polypeptide may comprise a sequence at least 99% identical to SEQ ID NO: 127, or at least 99% identical to SEQ ID NO: 127, or identical to SEQ ID NO: 127.

[0287] In some embodiments of the present invention, the polypeptide comprises a sequence at least 99% identical to SEQ ID NO: 128, or at least 99% identical to SEQ ID NO: 128, or identical to SEQ ID NO: 128.

Table 6. Polypeptide sequences for illustrative LILRB1-based inhibitory receptors

Name	Sequence
LILRB 1	MTPILTVLICLGLSLGPRTHVQAGHLPKPTLWAEPGSVITQ GSPVTLRCQGGQETQEYRLYREKKTALWITRIPQELVKKG QFPIPSITWEHAGRYRCYYGSDTAGRSESSDPLELVVTGA YIKPTLSAQSPVVNSGGNVILQCDSQVAFDGFSLCKEGED EHPQCLNSQPHARGSSRAIFSVGPVSPSRRWWYRCYAYDS NSPYEWSLPSDLLELLVLGVSKKPSLSVQPGPIVAPEETLT LQCGSDAGYNRFVLYKDGERDFLQLAGAQPQAGLSQANF TLGPVSRSYGGQYRCYGAHNLSSEWSAPSDPLDILIAGQF YDRVSLSVQPGPTVASGENVTLLCQSQGWMTFLLTKEG AADDPWRLRSTYQSQKYQAEFPMGPV TSAHAGTYRC <u>YG</u> <u>SQSSKPYLLTHPSDPLEL</u> VVSGPSGGPSSPTTGPTSTSGPE DQPLTPTGSDPQSGLGRLGVVIGILVAVILLLLLLLLLLFLI LRHRRQGKHWTSTQRKADFQHPAGAVGPEPTDRGLQWR SSPAADAQEEN <u>LYAAVKHT</u> QPEDGVEMDTRSPHDEDPA <u>VTYAEVKHSRPRREMASPPSPLSGEFLDTKDRQAEEDRQ</u> MDTEAAASEAPQD <u>VTYAQLHSL</u> TLRREATEPPPSQEGPSP AVP <u>SIYATL</u> AIHPSQEGPSPAVPSIYATLAIH SEQ ID NO: 113 <u>YGSQSSKPYLLTHPSDPLEL</u> VVSGPSGGPSSPTTGPTSTSG
LILRB 1 hinge-	<u>YGSQSSKPYLLTHPSDPLEL</u> VVSGPSGGPSSPTTGPTSTSG
transmembrane-intracellular domain	PEDQPLTPTGSDPQSGLGRLGVVIGILVAVILLLLLLLLLL <u>FLILR</u> HRRQGKHWTSTQRKADFQHPAGAVGPEPTDRGLQ WRSSPAADAQEENLYAAVKHTQPEDGVEMDTRSPHDEDP QAVTYAEVKHSRPRREMASPPSPLSGEFLDTKDRQAEEDR QMDTEAAASEAPQDV <u>TYAQLHSL</u> TLRREATEPPPSQEGPS PAVPSIYATLAIH SEQ ID NO: 130

(continued)

	Name	Sequence
5	LILRB 1 hinge-transmembrane-intracellular domain (w/o YGS Q S SKP YLLTHPSD PLEL)	VVSGPSGGPSSPTTGPTSTSGPEDQPLTPTGSDPQSGLGRH LGVVIGILVAVILLLLLLLLLLLFLILRHRRQGKHWTSTQRK ADFQHPAGAVGPEPTDRGLQWRSSPAADAQEENLYAAV KHTQPEDGVEMDTRSPHDEDPQAVTYAEVKHSRPRREMA SPPSPLSGEFLDTKDRQAEEDRQMDTEAAASEAPQDVTYA QLHSLTLRREATEPPPSQEGPSPAVPSIYATLAIH SEQ ID NO: 131
10	LILRB 1 hinge domain	<u>YGSQSSKP YLLTHPSD PLEL</u> VVSGPSGGPSSPTTGPTSTSG PEDQPLTPTGSDPQSGLGRHLG SEQ ID NO: 132
20	LILRB 1 transmembrane domain	VVIGILVAVILLLLLLLLLLLFLIL SEQ ID NO: 133
25	LILRB 1 intracellular domain	RHRRQGKHWTSTQRKADFQHPAGAVGPEPTDRGLQWRS SPAADAQEENLYAAVKHTQPEDGVEMDTRSPHDEDPQAV TYAEVKHSRPRREMASPPSPLSGEFLDTKDRQAEEDRQM DTEAAASEAPQDVTYAQLHSLTLRREATEPPPSQEGPSPA VPSIYATLAIH SEQ ID NO: 129
30	ITIM1	NLYAAV SEQ ID NO: 115
35	ITIM2	VTYAEV SEQ ID NO: 116
40	ITIM3	VTYAQL SEQ ID NO: 117
45	ITIM4	SIYATL SEQ ID NO: 118
50	ITIM 1-2	<u>NLYAAV</u> KHTQPEDGVEMDTRSPHDEDPQAV <u>VTYAEV</u> SEQ ID NO: 119
55	ITIM2-3	<u>VTYAEV</u> KHSRPRREMASPPSPLSGEFLDTKDRQAEEDRQ MDTEAAASEAPQD <u>VTYAQL</u> SEQ ID NO: 120
	ITIM3-4	<u>VTYAQL</u> HSLTLRREATEPPPSQEGPSPAV <u>PSIYATL</u> SEQ ID NO: 121

(continued)

	Name	Sequence
5	ITIM1-3	<u>NLYAAVKHTQ</u> PEDGVEMDTRSPHDEDPQA <u>VTYAEVKHS</u> RPRREMASPPSPLSGEFLDTKDRQAEEDRQMDTEAAASEA PQD <u>VTYAQL</u>
10		SEQ ID NO: 122
15	ITIM2-4	<u>VTYAEVKHS</u> RPRREMASPPSPLSGEFLDTKDRQAEEDRQ MDTEAAASEAPQD <u>VTYAQL</u> HSLTLRREATEPPPSQEGPSP AVP <u>SIYATL</u>
20		SEQ ID NO: 123
25	ITIM 1-4	<u>NLYAAVKHTQ</u> PEDGVEMDTRSPHDEDPQA <u>VTYAEVKHS</u> RPRREMASPPSPLSGEFLDTKDRQAEEDRQMDTEAAASEA PQD <u>VTYAQL</u> HSLTLRREATEPPPSQEGPSPAVP <u>SIYATL</u>
30		SEQ ID NO: 124
35	D3D4 domain	YGSQSSKPYLLTHPSDPLEL SEQ ID NO: 125
40	Short hinge	VVSGPSGGPSSPTTGPTSTSGPEDQPLTPTGSDPQSGLGRH LG
45		SEQ ID NO: 126
50	Hinge (iTIM hinge)	YGSQSSKPYLLTHPSDPLELVVSGPSGGPSSPTTGPTSTSGP EDQPLTPTGSDPQSGLGRHLGV (SEQ ID NO: 424)
55	Short hinge 2	VVSGPSGGPSSPTTGPTSTSGPEDQPLTPTGSDPQSGLGRH LGV (SEQ ID NO: 425)
	Long hinge 1	AGSGGSGGSGGSPVPSTPPTPSPSTPPTPSPSGGSGNSSGSG GSPVPSTPPTPSPSTPPTPSPSASV (SEQ ID NO: 426)
	Long hinge 2	AGSGGSGGSGGSPVPSTPPTNSSSTPPTPSPSPVPSTPPTNSS STPPTPSPSPVPSTPPTNSSSTPPTPSPSASV (SEQ ID NO: 427)
	2x short hinge	VVSGPSGGPSSPTTGPTSTSGPEDQPLTPTGSDPQSGLGRH VVSGPSGGPSSPTTGPTSTSGPEDQPLTPTGSDPQSGLGRH LGV (SEQ ID NO: 428)
	Hinge (truncated)	TTGPTSTSGPEDQPLTPTGSDPQSGLGRHLGV (SEQ ID NO: 429)

(continued)

	Name	Sequence
5	Hinge-transmembrane	<u>YGSQSSKPYLLTHPSDPLEL</u> VVSGPSGGPSSPTTGPTSTSG PEDQPLTPTGSDPQSGGLGRHLGV VVIGILVAVILL LLLLLLLL FLIL
10		SEQ ID NO: 127
15	Transmembrane-intracellular domain.	VVIGILVAVILL LLLLLLLL FLIL RHRRQGKHWTSTQRKA DFQHPAGAVGPEPTDRGLQWRSSPAADAQEENLYAAVK HTQPEDGVEMDTRSPHDEDPQAVTYAEVKHSRPRREMAS PPSPLSGEFLDTKDRQAEEDRQMDTEAAASEAPQDVTYA QLHSLTLRREATEPPPSQEGPSPAVPSIYATLAIH
20		SEQ ID NO: 128

Polynucleotides and Vectors

25 **[0288]** The present invention provides polynucleotides comprising polynucleotide sequences encoding the first and second receptors as used in the present invention. In some embodiments, the polynucleotides may further comprise an shRNA described herein. The present invention provides vectors comprising the polynucleotides according to the present invention. The disclosure provides immune cells comprising the polynucleotides and vectors described herein.

[0289] As described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 180, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 181, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 182, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 183, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 184, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0290] As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 185, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 187, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 189, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 191, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the polynucleotide or polynucleotide system comprises SEQ ID NO: 193, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0291] As described herein, the sequence of the first and/or second receptor is operably linked to a promoter. As also described herein, the sequence encoding the first receptor is operably linked to a first promoter, and the sequence encoding the second receptor is operably linked to a second promoter.

50 [0292] The present invention provides vectors comprising the polynucleotides according to the present invention.

[0293] In some embodiments, the first receptor is encoded by a first vector and the second receptor is encoded by second vector. In some embodiments, both receptors are encoded by a single vector.

[0294] In some embodiments, the first and second receptors are encoded by a single vector. Methods of encoding multiple polypeptides using a single vector will be known to persons of ordinary skill in the art, and include, *inter alia*, encoding multiple polypeptides under control of different promoters, or, if a single promoter is used to control transcription of multiple polypeptides, use of sequences encoding internal ribosome entry sites (IRES) and/or self-cleaving peptides. Exemplary self-cleaving peptides include T2A, P2A, E2A and F2A self-cleaving peptides. As described herein, the T2A self-cleaving peptide comprises a sequence of EGRGSLTCDGVEENPGP (SEQ ID NO: 430). As also described herein,

the P2A self-cleaving peptide comprises a sequence of ATNFSLLKQAGDVEENPGP (SEQ ID NO: 431). As also described herein, the E2A self-cleaving peptide comprises a sequence of QCTNYALLKLAGDVESNPGP (SEQ ID NO: 432). As also described herein, the F2A self-cleaving peptide comprises a sequence of VKQTLNFDLLKLAGDVESNPGP (SEQ ID NO: 433). As also described herein, the T2A self-cleaving peptide comprises a sequence of EGRGSLTTCGD-VEENPGP (SEQ ID NO: 430). Any of the foregoing can also include an N terminal GSG linker. For example, a T2A self-cleaving peptide can also comprise a sequence of GSGEGRGSLTTCGDVEENPGP (SEQ ID NO: 178), which can be encoded by a sequence of GGATCCGGAGAGGGCAGAGGCAGCCTGCTGACATGTGGCGACGTGGAAGAGAACCCTGGCCCC (SEQ ID NO: 434).

[0295] As described herein, the vector comprises SEQ ID NO: 180, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 181, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 182, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 183, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 184, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0296] As also described herein, the vector comprises SEQ ID NO: 185, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 187, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 189, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 191, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector comprises SEQ ID NO: 193, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0297] As described herein, the vector encodes a polypeptide sequence comprising SEQ ID NO: 186, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector encodes a polypeptide sequence comprising SEQ ID NO: 188, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector encodes a polypeptide sequence comprising SEQ ID NO: 190, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector encodes a polypeptide sequence comprising SEQ ID NO: 192, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the vector encodes a polypeptide sequence comprising SEQ ID NO: 194, or a sequence that shares at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0298] In some embodiments, the vector is an expression vector, i.e. for the expression of the first and/or second receptor in a suitable cell.

[0299] Vectors derived from retroviruses such as the lentivirus are suitable tools to achieve long-term gene transfer since they allow long-term, stable integration of a transgene and its propagation in daughter cells. Lentiviral vectors have the added advantage over vectors derived from onco-retroviruses such as murine leukemia viruses in that they can transduce non-proliferating cells, such as hepatocytes. They also have the added advantage of low immunogenicity.

[0300] The expression of natural or synthetic nucleic acids encoding receptors is typically achieved by operably linking a nucleic acid encoding the receptor or portions thereof to a promoter, and incorporating the construct into an expression vector. The vectors can be suitable for replication and integration eukaryotes. Typical cloning vectors contain transcription and translation terminators, initiation sequences, and promoters useful for regulation of the expression of the desired nucleic acid sequence.

[0301] The polynucleotides encoding the receptors can be cloned into a number of types of vectors. For example, the polynucleotides can be cloned into a vector including a plasmid, a phagemid, a phage derivative, an animal virus, and a cosmid. Vectors of particular interest include expression vectors, replication vectors, probe generation vectors, and sequencing vectors.

[0302] Further, the expression vector may be provided to cells, such as immune cells, in the form of a viral vector. Viral vector technology is well known in the art and is described, for example, in Sambrook et al. (2001, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York), and in other virology and molecular biology manuals. Viruses, which are useful as vectors include retroviruses, adenoviruses, adeno-associated viruses, herpes viruses, and lentiviruses. In general, a suitable vector contains an origin of replication functional in at least one organism, a promoter sequence, convenient restriction endonuclease sites, and one or more selectable markers, (e.g., WO 01/96584; WO 01/29058; and U.S. Pat. No. 6,326,193).

[0303] A number of viral based systems have been developed for gene transfer into mammalian cells. For example, retroviruses provide a convenient platform for gene delivery systems. A selected gene can be inserted into a vector and

packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to cells of the subject either in vivo or ex vivo. A number of retroviral systems are known in the art. In some embodiments, adenovirus vectors are used. A number of adenovirus vectors are known in the art. In one embodiment, lentivirus vectors are used.

[0304] Additional promoter elements, e.g., enhancers, regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 basepairs (bp) upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the thymidine kinase (tk) promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription.

[0305] One example of a suitable promoter is the immediate early cytomegalovirus (CMV) promoter sequence. This promoter sequence is a strong constitutive promoter sequence capable of driving high levels of expression of any polynucleotide sequence operatively linked thereto. Another example of a suitable promoter is Elongation Growth Factor-1 α (EF-1 α). However, other constitutive promoter sequences may also be used, including the simian virus 40 (SV40) early promoter, mouse mammary tumor virus (MMTV), human immunodeficiency virus (HIV) long terminal repeat (LTR) promoter, MoMuLV promoter, an avian leukemia virus promoter, an Epstein-Barr virus immediate early promoter, a Rous sarcoma virus promoter, U6 promoter, as well as human gene promoters such as the actin promoter, the myosin promoter, the hemoglobin promoter, and the creatine kinase promoter. Further, the disclosure includes the use of constitutive promoters. Inducible promoters are also contemplated as part of the disclosure. The use of an inducible promoter provides a molecular switch capable of turning on expression of the polynucleotide sequence, which it is operatively linked when such expression is desired, or turning off the expression, when expression is not desired. Examples of inducible promoters include a metallothionein promoter, a glucocorticoid promoter, a progesterone promoter, and a tetracycline promoter.

[0306] In order to assess the expression of a receptor, the expression vector to be introduced into a cell can also contain either a selectable marker gene or a reporter gene or both to facilitate identification and selection of expressing cells from the population of cells sought to be transfected or infected through viral vectors. In other aspects, the selectable marker may be carried on a separate piece of DNA and used in a co-transfection procedure. Both selectable markers and reporter genes may be flanked with appropriate regulatory sequences to enable expression in the host cells. Useful selectable markers include, for example, antibiotic-resistance genes, such as neo.

[0307] Reporter genes are used for identifying potentially transfected or transduced cells and for evaluating the functionality of regulatory sequences. In general, a reporter gene is a gene that is not present in or expressed by the recipient organism or tissue and that encodes a polypeptide whose expression is manifested by some easily detectable property, e.g., enzymatic activity. Expression of the reporter gene is assayed at a suitable time after the DNA has been introduced into the recipient cells. Suitable reporter genes may include genes encoding luciferase, beta-galactosidase, chloramphenicol acetyl transferase, secreted alkaline phosphatase, or the green fluorescent protein gene (e.g., Ui-Tei et al., 2000 FEBS Letters 479: 79-82). Suitable expression systems are well known and may be prepared using known techniques or obtained commercially. In general, the construct with the minimal 5' flanking region showing the highest level of expression of reporter gene is identified as the promoter. Such promoter regions may be linked to a reporter gene and used to evaluate agents for the ability to modulate promoter-driven transcription.

[0308] Methods of introducing and expressing genes into a cell are known in the art. In the context of an expression vector, the vector can be readily introduced into a host cell, e.g., mammalian, bacterial, yeast, or insect cell by any method in the art. For example, the expression vector can be transferred into a host cell by physical, chemical, or biological means.

[0309] Physical methods for introducing a polynucleotide into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, and electroporation. Methods for producing cells comprising vectors and/or exogenous nucleic acids are well-known in the art. See, for example, Sambrook et al. (2001, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York). One method for the introduction of a polynucleotide into a host cell is calcium phosphate transfection.

[0310] Biological methods for introducing a polynucleotide of interest into a host cell include the use of DNA and RNA vectors. Viral vectors, and especially retroviral vectors, have become the most widely used method for inserting genes into mammalian, e.g., human cells. Other viral vectors can be derived from lentivirus, poxviruses, herpes simplex virus I, adenoviruses and adeno-associated viruses. See, for example, U.S. Pat. Nos. 5,350,674 and 5,585,362.

[0311] Chemical means for introducing a polynucleotide into a host cell include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. An exemplary colloidal system for use as a delivery vehicle in vitro and in vivo is a liposome (e.g., an artificial membrane vesicle).

[0312] Regardless of the method used to introduce exogenous nucleic acids into a host cell or otherwise expose a

cell to the inhibitor of the present disclosure, in order to confirm the presence of the recombinant DNA sequence in the host cell, a variety of assays may be performed. Such assays include, for example, "molecular biological" assays well known to those of skill in the art, such as Southern and Northern blotting, RT-PCR and PCR; "biochemical" assays, such as detecting the presence or absence of a particular peptide, e.g., by immunological means (ELISAs and Western blots) or by assays described herein to identify agents falling within the scope of the disclosure.

Immune Cells

[0313] The present invention provides immune cells according to the appended claims.

[0314] As disclosed herein, the immune cells comprise: (a) first receptor, comprising a first extracellular ligand binding domain specific to a target antigen selected from: (i) a cancer cell-specific antigen, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); or (ii) EGFR, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); and (b) a second receptor, comprising a second extracellular ligand binding specific to a non-target antigen whose expression that has been lost from a cancer cell due to loss of heterozygosity or another mechanism. As described herein, the non-target antigen is selected from CXCL16, COLEC12 and APCDD1, HLA-A*01, HLA-A*02, HLA-A*03, HLA-A*1 1, HLA-B*07, HLA-C*07, or an antigen peptide thereof in a complex with a major histocompatibility complex class I (MHC-I). In some embodiments, the non-target antigen is HLA-A*02, or an allelic variant thereof. In some embodiments, the first receptor is a CAR or TCR. In some embodiments, the second receptor is an inhibitory receptor.

[0315] In some embodiments, the immune cell comprises an shRNA. In some embodiments, the shRNA reduces or eliminates expression of B2M. In some embodiments, the shRNA reduces or eliminates expression of HLA-A*02.

[0316] In some embodiments, the immune cell comprises a modification to an MHC Class I gene or a *B2M* gene. In some embodiments, the MHC Class I gene is HLA-A* 02, or an allelic variant thereof. In some embodiments, the modification reduces or eliminates expression of the MHC Class I gene or the *B2M* gene.

[0317] In some embodiments, the immune cell comprises a polynucleotide according to the present invention. In some embodiments, the polynucleotide comprises a sequence encoding a first receptor as used in the present invention. In some embodiments, the polynucleotide comprises a sequence encoding a second receptor as used in the present invention. In some embodiments, the polynucleotide encodes a shRNA as described herein. In some embodiments, the polynucleotide comprises a sequence encoding a first receptor as used in the present invention and a second receptor as used in the present invention. In some embodiments, the polynucleotide comprises a sequence encoding a first receptor as used in the present invention, a second receptor as used in the present invention, and a shRNA as described herein.

[0318] As used herein, the term "immune cell" refers to a cell involved in the innate or adaptive (acquired) immune systems. Exemplary innate immune cells include phagocytic cells such as neutrophils, monocytes and macrophages, Natural Killer (NK) cells, polymorphonuclear leukocytes such as neutrophils eosinophils and basophils and mononuclear cells such as monocytes, macrophages and mast cells. Immune cells with roles in acquired immunity include lymphocytes such as T-cells and B-cells.

[0319] In some embodiments of the present invention, the first receptor comprises SEQ ID NO: 177, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 174 or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As described herein, the first receptor comprises SEQ ID NO: 177, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 174, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto. As also described herein, the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and the second receptor comprises SEQ ID NO: 174, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0320] As also described herein, the immune cell further comprises a T2A self-cleaving peptide, wherein the T2A self-

cleaving peptide comprises SEQ ID NO: 178, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0321] In some embodiments of the present invention, the immune cell further comprises an interfering RNA, wherein the interfering RNA comprises SEQ ID NO: 179, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0322] As used herein, a "T-cell" refers to a type of lymphocyte that originates from a bone marrow precursor that develops in the thymus gland. There are several distinct types of T-cells, which develop upon migration to the thymus, which include, helper CD4+ T-cells, cytotoxic CD8+ T cells, memory T cells, regulatory CD4+ T-cells, NK T cells, $\gamma\delta$ T cells, Mucosal-associated invariant T (MAIT) cells, and stem memory T-cells. Different types of T-cells can be distinguished by the ordinarily skilled artisan based on their expression of markers. Methods of distinguishing between T-cell types will be readily apparent to the ordinarily skilled artisan.

[0323] As described herein, the immune cell is selected from the group consisting of T cells, B cells and Natural Killer (NK) cells. As also described herein, the immune cell is a T cell. As also described herein, the immune cell is a B cell. As also described herein, the immune cell is a Natural Killer (NK) cell. As also described herein, the immune cell is CD8-. As also described herein, the immune cell is CD8+. As also described herein, the immune cell is CD4+. As also described herein, the immune cell is CD4-. According to the present invention, the immune cell is a CD8- CD4+ T cell or a CD8+ CD4- T cell.

[0324] As described herein, the immune cell is a gamma delta ($\gamma\delta$) T cell. As also described herein, the immune cell is an invariant T cell. As also described herein, the immune cell is an invariant natural killer T cell (iNKT cell).

[0325] As described herein, the first receptor and the second receptor together specifically activate the immune cell in the presence of the target cell.

[0326] The immune cell may be non-natural. The immune cell may be isolated.

[0327] Methods transforming populations of immune cells, such as T cells, with the vectors of the instant disclosure will be readily apparent to the person of ordinary skill in the art. For example, CD3+ T cells can be isolated from PBMCs using a CD3+ T cell negative isolation kit (Miltenyi), according to manufacturer's instructions. T cells can be cultured at a density of 1×10^6 cells/mL in X-Vivo 15 media supplemented with 5% human A/B serum and 1% Pen/strep in the presence of CD3/28 Dynabeads (1:1 cell to bead ratio) and 300 Units/mL of IL-2 (Miltenyi). After 2 days, T cells can be transduced with viral vectors, such as lentiviral vectors using methods known in the art. The viral vector may be transduced at a multiplicity of infection (MOI) of 5. Cells can then be cultured in IL-2 or other cytokines such as combinations of IL-7/15/21 for an additional 5 days prior to enrichment. Methods of isolating and culturing other populations of immune cells, such as B cells, or other populations of T cells, will be readily apparent to the person of ordinary skill in the art. Although this method outlines a potential approach it should be noted that these methodologies are rapidly evolving. For example excellent viral transduction of peripheral blood mononuclear cells can be achieved after 5 days of growth to generate a >99% CD3+ highly transduced cell population.

[0328] Methods of activating and culturing populations of T cells comprising the TCRs, CARs, inhibitory receptors, receptors or vectors encoding same, will be readily apparent to the person of ordinary skill in the art.

[0329] Whether prior to or after genetic modification of T cells to express a TCR, the T cells can be activated and expanded generally using methods as described, for example, in U.S. Pat. Nos. 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; 6,867,041, 10040846; and U.S. Pat. Appl. Pub. No. 2006/0121005.

[0330] T cells of the instant disclosure may be expanded and activated in vitro. Generally, the T cells of the instant disclosure are expanded in vitro by contact with a surface having attached thereto an agent that stimulates a CD3/TCR complex associated signal and a ligand that stimulates a co-stimulatory molecule on the surface of the T cells. In particular, T cell populations may be stimulated as described herein, such as by contact with an anti-CD3 antibody. For co-stimulation of an accessory molecule on the surface of the T cells, a ligand that binds the accessory molecule is used. For example, a population of T cells can be contacted with an anti-CD3 antibody and an anti-CD28 antibody, under conditions appropriate for stimulating proliferation of the T cells. To stimulate proliferation of either CD4+ T cells or CD8+ T cells, an anti-CD3 antibody and an anti-CD28 antibody can be used. Examples of an anti-CD28 antibody include 9.3, B-T3, XR-CD28 (Diacione, Besançon, France) can be used as can other methods commonly known in the art (Berg et al., Transplant Proc. 30(8):3975-3977, 1998; Haanen et al., J. Exp. Med. 190(9):13191328, 1999; Garland et al., J. Immunol Meth. 227(1-2):53-63, 1999).

[0331] The primary stimulatory signal and the co-stimulatory signal for the T cell may be provided by different protocols. For example, the agents providing each signal may be in solution or coupled to a surface. When coupled to a surface, the agents may be coupled to the same surface (i.e., in "cis" formation) or to separate surfaces (i.e., in "trans" formation). Alternatively, one agent may be coupled to a surface and the other agent in solution. The agent providing the co-stimulatory signal may be bound to a cell surface and the agent providing the primary activation signal may be in solution or coupled to a surface. Both agents may be in solution. The agents may be in soluble form, and then cross-linked to a surface, such as a cell expressing Fc receptors or an antibody or other binding agent, which will bind to the agents. In

this regard, see for example, U.S. Patent Application Publication Nos. 20040101519 and 20060034810 for artificial antigen presenting cells (aAPCs) that are contemplated for use in activating and expanding T cells in the present disclosure.

[0332] The two agents are immobilized on beads, either on the same bead, i.e., "cis," or to separate beads, i.e., "trans". By way of example, the agent providing the primary activation signal is an anti-CD3 antibody or an antigen-binding fragment thereof and the agent providing the co-stimulatory signal is an anti-CD28 antibody or antigen-binding fragment thereof; and both agents are co-immobilized to the same bead in equivalent molecular amounts. A 1:1 ratio of each antibody bound to the beads for CD4+ T cell expansion and T cell growth may be used. The ratio of CD3:CD28 antibody bound to the beads may range from 100:1 to 1:100 and all integer values there between. In one aspect of the present disclosure, more anti-CD28 antibody is bound to the particles than anti-CD3 antibody, i.e., the ratio of CD3:CD28 is less than one. The ratio of anti CD28 antibody to anti CD3 antibody bound to the beads may be greater than 2:1.

[0333] Ratios of particles to cells from 1: 500 to 500:1 and any integer values in between may be used to stimulate T cells or other target cells. As those of ordinary skill in the art can readily appreciate, the ratio of particles to cells may depend on particle size relative to the target cell. For example, small sized beads could only bind a few cells, while larger beads could bind many. The ratio of cells to particles may range from 1:100 to 100:1 and any integer values in-between and in further embodiments the ratio may comprise 1:9 to 9:1 and any integer values in between, can also be used to stimulate T cells. A ratio of 1:1 cells to beads may be used. One of skill in the art will appreciate that a variety of other ratios may be suitable for use in the present disclosure. In particular, ratios will vary depending on particle size and on cell size and type.

[0334] In the present disclosure, the cells, such as T cells, may be combined with agent-coated beads, the beads and the cells are subsequently separated, and then the cells are cultured. Prior to culture, the agent-coated beads and cells may not be separated but may be cultured together. The beads and cells may be first concentrated by application of a force, such as a magnetic force, resulting in increased ligation of cell surface markers, thereby inducing cell stimulation.

[0335] By way of example, cell surface proteins may be ligated by allowing paramagnetic beads to which anti-CD3 and anti-CD28 are attached to contact the T cells. The cells (for example, CD4+ T cells) and beads (for example, DYNABEADS CD3/CD28 T paramagnetic beads at a ratio of 1:1) may be combined in a buffer. Again, those of ordinary skill in the art can readily appreciate any cell concentration may be used. It may be desirable to significantly decrease the volume, in which particles and cells are mixed together (i.e., increase the concentration of cells), to ensure maximum contact of cells and particles. For example, a concentration of about 2 billion cells/ml may be used. Greater than 100 million cells/ml may be used. A concentration of cells of 10, 15, 20, 25, 30, 35, 40, 45, or 50 million cells/ml may be used. A concentration of cells from 75, 80, 85, 90, 95, or 100 million cells/ml may be used. Concentrations of 125 or 150 million cells/ml can be used. Cells that are cultured at a density of 1×10^6 cells/mL may be used.

[0336] The mixture may be cultured for several hours (about 3 hours) to about 14 days or any hourly integer value in between. The beads and T cells may be cultured together for 2-3 days. Conditions appropriate for T cell culture include an appropriate media (e.g., Minimal Essential Media or RPMI Media 1640 or, X-vivo 15, (Lonza)) that may contain factors necessary for proliferation and viability, including serum (e.g., fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, GM-CSF, IL-10, IL-12, IL-15, TGF β , and TNF- α or any other additives for the growth of cells known to the skilled artisan. Other additives for the growth of cells include surfactant, plasmanate, and reducing agents such as N-acetyl-cysteine and 2-mercaptoethanol. Media can include RPMI 1640, AIM-V, DMEM, MEM, α -MEM, F-12, X-Vivo 15, and X-Vivo 20, Optimizer, with added amino acids, sodium pyruvate, and vitamins, either serum-free or supplemented with an appropriate amount of serum (or plasma) or a defined set of hormones, and/or an amount of cytokine(s) sufficient for the growth and expansion of T cells. The media may comprise X-VIVO-15 media supplemented with 5% human A/B serum, 1% penicillin/streptomycin (pen/strep) and 300 Units/ml of IL-2 (Miltenyi).

[0337] The T cells may be maintained under conditions necessary to support growth, for example, an appropriate temperature (e.g., 37° C.) and atmosphere (e.g., air plus 5% CO₂).

[0338] The T cells comprising TCRs, CARs and inhibitory receptors of the disclosure may be autologous. Prior to expansion and genetic modification, a source of T cells may be obtained from a subject. Immune cells such as T cells can be obtained from a number of sources, including peripheral blood mononuclear cells, bone marrow, lymph node tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In the present disclosure, any number of T cell lines available in the art, may be used. In the present disclosure, T cells can be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled artisan, such as Ficoll™ separation.

[0339] Cells from the circulating blood of an individual may be obtained by apheresis. The apheresis product typically may contain lymphocytes, including T cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. The cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing steps. The cells may be washed with phosphate buffered saline (PBS). The wash solution may lack calcium and may lack magnesium or may lack many if not all divalent cations. As those of ordinary skill in the art would readily appreciate, a washing step may be accomplished by methods

known to those in the art, such as by using a semi-automated "flow-through" centrifuge (for example, the Cobe 2991 cell processor, the Baxter CytoMate, or the Haemonetics Cell Saver 5) according to the manufacturer's instructions. After washing, the cells may be resuspended in a variety of biocompatible buffers, such as, for example, Ca²⁺-free, Mg²⁺-free PBS, PlasmaLyte A, or other saline solution with or without buffer. Alternatively, the undesirable components of the apheresis sample may be removed and the cells may be directly resuspended in culture media.

[0340] Immune cells such as T cells may be isolated from peripheral blood lymphocytes by lysing the red blood cells and depleting the monocytes, for example, by centrifugation through a PERCOLL[™] gradient or by counterflow centrifugal elutriation. Specific subpopulations of immune cells, such as T cells, B cells, or CD4⁺ T cells can be further isolated by positive or negative selection techniques. For example, T cells may be isolated by incubation with anti-CD4-conjugated beads, for a time period sufficient for positive selection of the desired T cells.

[0341] Enrichment of an immune cell population, such as a T cell population, by negative selection can be accomplished with a combination of antibodies directed to surface markers unique to the negatively selected cells. One method is cell sorting and/or selection via negative magnetic immune-adherence or flow cytometry that uses a cocktail of monoclonal antibodies directed to cell surface markers present on the cells negatively selected. For example, to enrich for CD4⁺ cells by negative selection, a monoclonal antibody cocktail typically includes antibodies to CD14, CD20, CD11b, CD16, HLA-DR, and CD8.

[0342] For isolation of a desired population of immune cells by positive or negative selection, the concentration of cells and surface (e.g., particles such as beads) can be varied. It may be desirable to significantly decrease the volume, in which beads and cells are mixed together (i.e., increase the concentration of cells), to ensure maximum contact of cells and beads.

[0343] The cells may be incubated on a rotator for varying lengths of time at varying speeds at either 2-10°C or at room temperature.

[0344] T cells for stimulation, or PBMCs from which immune cells such as T cells are isolated, can also be frozen after a washing step. Wishing not to be bound by theory, the freeze and subsequent thaw step provides a more uniform product by removing granulocytes and to some extent monocytes in the cell population. After the washing step that removes plasma and platelets, the cells may be suspended in a freezing solution. While many freezing solutions and parameters are known in the art and will be useful in this context, one method involves using PBS containing 20% DMSO and 8% human serum albumin, or culture media containing 10% Dextran 40 and 5% Dextrose, 20% Human Serum Albumin and 7.5% DMSO, or 31.25% Plasmalyte-A, 31.25% Dextrose 5%, 0.45% NaCl, 10% Dextran 40 and 5% Dextrose, 20% Human Serum Albumin, and 7.5% DMSO or other suitable cell freezing media containing for example, Hespan and PlasmaLyte A, the cells then are frozen to -80°C at a rate of 1° per minute and stored in the vapor phase of a liquid nitrogen storage tank. Other methods of controlled freezing may be used as well as uncontrolled freezing immediately at -20°C or in liquid nitrogen.

[0345] The disclosure provides an immune cell expressing the activator and/or inhibitory receptors described herein, wherein the immune cell has reduced expression and/or function of the major histocompatibility (MHC) class I complex.

[0346] In some embodiments of the present invention, the immune cell is autologous. For example, the immune cells is isolated or derived from the same subject who will receive the cell as part of a therapeutic regimen. It can be advantageous to modify autologous immune cells to have reduced expression and/or function of MHC class I with the inhibitory receptor being specific to an MHC class I antigen. Without wishing to be bound by theory, modification of autologous immune cells having reduced expression and/or function of MHC class I reduces binding of the inhibitory receptor by MHC class I expressed by the immune cells, either in cis or in trans.

[0347] In some embodiments of the present invention, the immune cell is all allogeneic. Allogeneic immune cells can be derived from a donor other than the subject to which the immune cells will be administered. Allogeneic immune cells have been commonly referred to in cell therapy as "off-the-shelf" or "universal", because of the possibility for allogeneic cells to be prepared and stored for use in subjects of a variety of genotypes.

[0348] Any suitable methods of reducing expression and/or function of the MHC class I complex are envisaged as within the scope of the instant disclosure, and include, inter alia, expression of interfering RNAs that knock down one or more RNAs encoding MHC class I components, or modifications of genes encoding MHC class I components.

[0349] The major histocompatibility complex (MHC) is a locus on the vertebrate genome that encodes a set of polypeptides required for the adaptive immune system. Among these are MHC class I polypeptides that include HLA-A, HLA-B, and HLA-C and alleles thereof. MHC class I alleles are highly polymorphic and expressed in all nucleated cells. MHC class I polypeptides encoded by HLA-A, HLA-B, and HLA-C and alleles thereof form heterodimers with β 2 microglobulin (B2M) and present in complex with antigens on the surface of cells. As referred to herein, an MHC class I gene or polypeptide may refer to any polypeptide found in the the MHC or the corresponding gene encoding said polypeptide. The immune cells of the disclosure may be inactivated by an inhibitor ligand comprising an MHC class I polypeptide, e.g. HLA-A, HLA-B, and HLA-C and alleles thereof. HLA-A alleles can be, for example, HLA-A*02, HLA-A*02:01, HLA-A*02:01:01, HLA-A*02:01:01:01, and/or any gene that encodes protein identical or similar to HLA-A*02 protein. Thus, to prevent autocrine signaling/binding as described herein, it is desirable to eliminate or reduce expression of polypeptides

encoded by HLA-A, HLA-B, and HLA-C and alleles thereof in the immune cells.

Immune Cells with Reduced MHC Class I Polypeptide Expression

[0350] The immune cells described herein may be modified to inactivate, or reduce or eliminate expression or function of an endogenous gene encoding an allele of an endogenous MHC class I polypeptide. As described herein, the gene encoding the MHC class I polypeptide is HLA-A, HLA-B, and/or HLA-C. HLA-A, HLA-B and HLA-C are encoded by the HLA-A, HLA-B and HLA-C loci. Each of HLA-A, HLA-B and HLA-C includes many variant alleles. As described herein, the gene encoding the MHC class I polypeptide is HLA-A. In some embodiments of the present invention, the gene encoding the MHC class I polypeptide is HLA-A*02. In some embodiments of the present invention, the gene encoding the MHC class I polypeptide is HLA-A*02:01. In some embodiments of the present invention, the gene encoding the MHC class I polypeptide is HLA-A*02:01:01. In some embodiments of the present invention, the gene encoding the MHC class I polypeptide is HLA-A*02:01:01:01.

[0351] In some embodiments of the present invention, the genetically engineered immune cells are modified to reduce or eliminate expression of the B2M gene product. The beta-2 microglobulin (B2M) gene encodes a protein that associates with the major histocompatibility complex (MHC) class I, i.e. MHC-I complex. The MHC-I complex is required for presentation of antigens on the cell surface. The MHC-I complex is disrupted and non-functional when the B2M is deleted (Wang D et al. Stem Cells Transl Med. 4:1234-1245 (2015)). Furthermore, the B2M gene can be disrupted with high efficiency using gene editing techniques known in the art (Ren et al. Clin. Cancer Res. 23:2255-2266 (2017)). Reducing or eliminating B2M can reduce, or eliminate functional MHC I on the surface of the immune cell.

[0352] The disclosure provides gene editing systems for editing an endogenous target gene in an immune cell. The disclosure provides interfering RNAs specific to sequences of target genes. Gene editing systems such as CRISPR/Cas systems, TALENs and zinc fingers can be used to generate double strand breaks, which, through gene repair mechanisms such as homology directed repair or non-homologous end joining (NHEJ), can be used to introduce mutations. NHEJ after resection of the ends of the break, or improper end joining, can be used to introduce deletions. In some embodiments, the target gene comprises a gene encoding a subunit of the MHC-I complex.

[0353] Target gene sequences include, promoters, enhancers, introns, exons, intron/exon junctions, transcription products (pre-mRNA, mRNA, and splice variants), and/or 3' and 5' untranslated regions (UTRs). Any gene element or combination of gene elements may be targeted for the purpose of genetic editing in the immune cells described herein. Modifications to the target genes can be accomplished using any method known in the art to edit the target gene that results in altered or disrupted expression or function the target gene or gene product.

[0354] In some embodiments, modifying the gene encoding the MHC class I polypeptide comprises deleting all or a portion of the gene. In some embodiments, modifying the gene encoding the MHC class I polypeptide comprises introducing a mutation in the gene. In some embodiments, the mutation comprises a deletion, insertion, substitution, or frameshift mutation. In some embodiments, modifying the gene comprises using a nucleic acid guided endonuclease.

[0355] Gene sequences for the target genes described herein are known in the art. The sequences can be found at public databases, such as NCBI GenBank or the NCBI nucleotide database. Sequences may be found using gene identifiers, for example, the HLA-A gene has NCBI Gene ID: 3105, the HLA-B gene has NCBI Gene ID: 3106, the HLA-C gene has NCBI Gene ID: 3107, and the B2M gene has NCBI Gene ID: 567 and NCBI Reference Sequence: NC_000015.10. Gene sequences may also be found by searching public databases using keywords. For example, HLA-A alleles may be found in the NCBI nucleotide database by searching keywords, "HLA-A*02", "HLA-A*02:01", "HLA-A*02:01:01", or "HLA-A*02:01:01:01". These sequences can be used for targeting in various gene editing techniques known in the art. Table 7 provides nonlimiting illustrative sequences of HLA-A allele and B2M gene sequences targeted for modification as described herein.

Table 7. Exemplary Target Gene Sequences

B2M mRNA	
	AUUCUGAAGCUGACAGCAUUCGGGCCGAGAUGUCUCGCUCCGUGGCCUAGCUGUGCUCGCGCU
	ACUCUCUCUUUCUGGCCUGGAGGCUAUCCAGCGUACUCCAAAGAUUCAGGUUUACUCACGUCauc
	CAGCAGAGAAUGGAAAGUCAAAUUCUGAAUUGCUAUGUGUCUGGGUUCAUCCAUCGACAUU
	GAAGUUGACUUAUGAAGAAUGGAGAGAGAAUUGAAAAAGUGGAGCAUUCAGACUUGUCUUUCAG
	CAAGGACUGGUCUUUCUAUCUCUUGUACUACACUGAAUUCACCCACUGAAAAAGUAGUAUG
	CCUGCCGUGUGAACCAUGUGACUUUGUCACAGCCCAAGAUAGUUAAGUGGGAUCGAGACAUGUAA
	GCAGCAUCAUGGAGGUUUGAAGAUGCCGCAUUUGGAUUGGAUGAAUUCCAAUUCUGCUUGCUUG
	CUUUUUAAUUAUUGAUUGCUUAUACACUUAACACUUUAUGCACAAGUAGGGUUUAUAAUUAUGU
	UACAUGGACAUGAUCUUCUUUAUAAUUCUACUUUGAGUGCUGUCUCCAUUGUUUGAUGUAUCUGA
	GCAGGUUGCUCCACAGGUAGCUCUAGGAGGGCUGGCAACUUAGAGGUGGGAGCAGAGAAUUCUC
	UUAUCCAACAUCAACAUCUUGGUCAGAUUUGAACUCUUCUCAAUCUCUUGCACUCAAGCUUGUUA

(continued)

5	GAUAGUUAAGCGUGCAUAAGUUAACUUCCAAUUUACAUAUCUCUGCUUAGAAUUUUGGGGAAAAUU UAGAAAUAAUUGACAGGAUUAUUGGAAAUUUUUUUAUAAUGAAUGAAACAUUUUGUCAUAUAAG AUUCAUAUUUACUUCUUAUACAUAUUGAUAAAGUAAGGCAUGGUUGUGGUUAAUCUGGUUUUUUUU UGUUCACAAGUUAUAAUACAUAUAAACUUGA (SEQ ID NO: 172)
<i>B2M</i> Gene (GenBank: 567) 10 15 20 25 30 35 40 45 50 55	AAAGTGGAGGCGTCGCGCTGGCGGGCATTCCCTGAAGCTGACAGCATTTCGGGCCGAGAT GTCTCGCTCCGTGGCCTTAGCTGTGCTCGCGCTACTCTCTCTTTCTGGCCTGGAGGCTA TCCAGCGTGAGTCTCTCTACCCTCCCGCTCTGGTCCTTCCTCTCCCGCTCTGCACCCT CTGTGGCCCTCGCTGTGCTCTCTCGCTCCGTGACTTCCCTTCTCCAAGTTCTCCTTGGT GGCCCGCCGTGGGGCTAGTCCAGGGCTGGATCTCGGGGAAGCGGCGGGGTGGCCTG GGAGTGGGGAAGGGGGTGGCGACCCGGGACGCGCGCTACTTGCCCTTTTCGGCGGGG AGCAGGGGAGACCTTTGGCCTACGGCGACGGGAGGGTCGGGACAAAGTTTAGGGCG TCGATAAGCGTCAGAGCGCCGAGGTTGGGGGAGGGTTTCTCTCCGCTCTTTCGCGG GGCCTCTGGCTCCCCAGCGCAGCTGGAGTGGGGGACGGGTAGGCTCGTCCCAAAGG CGCGGCGCTGAGGTTTGTGAACGCGTGAGAGGGCGCTTGGGGTCTGGGGGAGGCGTC GCCC GGTAAGCCTGTCTGCTGCGGCTCTGCTTCCCTTAGACTGGAGAGCTGTGGACT TCGTCTAGGCGCCCGCTAAGTTCGCATGTCCTAGCACCTCTGGGTCTATGTGGGGCCA CACCGTGGGGAGGAAACAGCACGCGACGCTTTGTAGAATGCTTGGCTGTGATACAAAG CGTTTTCGAATAATTAACCTTATTTGTTCCCATCACATGTCACCTTTTAAAAAATTATAA GAACTACCCGTTATTGACATCTTTCTGTGTGCCAAGGACTTTATGTGCTTTGCGTCATT TAATTTTGA AAAACAGTTATCTTCCGCCATAGATAACTACTATGGTTATCTTCTGCCTCT CACAGATGAAGAACTAAGGCACCGAGATTTTAAGAACTTAATTACACAGGGGAT AAATGGCAGCAATCGAGATTGAAGTCAAGCCTAACCAGGGCTTTTGC GGAGCGCAT GCCTTTTGGCTGTAATTCGTGCATTTTTTTTTTAAGAAAAACGCCTGCCTTCTGCGTGAG ATTCTCCAGAGCAAACCTGGGCGGCATGGGCCCTGTGGTCTTTTCGTACAGAGGGCTTC CTCTTTGGCTCTTTGCTGTTGTTTCCAAGATGTACTGTGCCTCTTACTTTTCGGTTTT GAAAACATGAGGGGGTGGGCGTGGTAGCTTACGCCTGTAATCCAGCCTAGGGA GGCCGAGGCGGGAGGATGGCTTAGAGTCCGTAGTTGAGACCAGCCTGGCCAACTATG GTGAAGCCTGGTCTCTACAAAAAATAAATAACAAAAATTAGCCGGGTGTGGTGCTCG TGCTGTGGTCCCAGCTGCTCCGGTGGCTGAGGCGGGAGGATCTCTTGAGCTTAGGCT TTTGAGCTATCATGGCGCCAGTGCCTCCAGCGTGGGCAACAGAGCGAGACCCTGTC TCTCAAAAAAGAAAAAAGAAAAAAGAAAAAGAAAAAGAAAAAGAAAGT GAAGTTTGTGTCAGTCAGGGGAGCTGTAAACCATTAAATAAAGATAATCCAAGATGGT TACCAAGACTGTTGAGGACGCCAGAGATCTTGAGCACTTTCTAAGTACCTGGCAATA CACTAAGCGCGCTCACCTTTTCTCTGGCAAAACATGATCGAAAGCAGAATGTTTTGA TCATGAGAAAATTGCATTTAATTTGAATACAATTTATTTACAACATAAAGGATAATGT ATATATCACCACCATTACTGGTATTTGCTGGTTATGTTAGATGTCATTTAAAAAATA ACAATCTGATATTTAAAAAAAATCTTATTTTGA AAAATTTCCAAAGTAATACATGCCA TGCATAGACCATTTCTGGAAGATAACCACAAGAAACATGTAATGATGATTGCCTCTGA AGGTCTATTTTCTCCTCTGACCTGTGTGTGGGTTTTGTTTTGTTTTACTGTGGGCAT AAATTAATTTTTCAGTTAAGTTTTGGAAGCTTAAATAACTCTCCAAAAGTCATAAAGC CAGTAAGTGGTTGAGCCCAAATTCAAACCCAGCCTGTCTGATACTTGTCTCTTCTTA GAAAAGATTACAGTGATGCTCTCAAAAATCTTGCCGCCTTCCCTCAAAACAGAGAGT TCCAGGCAGGATGAATCTGTGCTCTGATCCCTGAGGCATTTAATATGTTCTTATTATT AGAAGCTCAGATGCAAAGAGCTCTCTAGCTTTTAATGTTATGAAAAAATCAGGTC TTCATTAGATTCCCAATCCACCTCTTGTAGGGGCTAGTAGCCTTTCCTTAATGATAG GGTGTCTTAGAGAGATATATCTGGTCAAGGTGGCCTGGTACTCCTCTTCTCCAC AGCCTCCCAGACAAGGAGGAGTAGCTGCCCTTTAGTGATCATGTACCTGAATATAA GTGTATTTAAAAGAATTTTATACACATATTTAGTGTCAATCTGTATATTTAGTAGC ACTAACACTTCTCTTCAATTTTCAATGAAAAATATAGAGTTTATAATTTTCTTCCAC TTCCCATGGATGGTCTAGTCATGCCTCTCATTTTGGAAAGTACTGTTCTGAAACAT TAGGCAATATATTCCCAACCTGGCTAGTTTACAGCAATCACCTGTGGATGCTAATTAA AACGCAAATCCCAGTGCATGCACTTCCATTTGATCATAATGGAAAAGTATGTTTC TGTCCCATTTGCCATAGTCCTCACCTATCCCTGTTGTATTTTATCGGGTCCAACCTCAAC CATTTAAGGTATTTGCCAGCTCTTGTATGCATTTAGGTTTGTCTTTGTTTTTAGCT CATGAAATTAGGTACAAAGTCAGAGAGGGGTCTGGCATATAAAACCTCAGCAGAAA TAAAGAGGTTTTGTTGTTTGGTAAGAACATACCTTGGGTTGGTTGGGCACGGTGGCTC

(continued)

5	GTGCCCTGTAATCCCCAACACTTTGGGAGGCCAAGGCAGGCTGATCACTTGAAGTTGGG AGTTCAAGACCAGCCTGGCCAACATGGTGAAATCCCGTCTCTACTGAAAATACAAAA ATTAACCAGGCATGGTGGTGTGTGCCTGTAGTCCCAGGAATCACTTGAACCCAGGAG GCGGAGGTTGCAGTGAGCTGAGATCTCACCCTGCACACTGCACTCCAGCCTGGGCA ATGGAATGAGATTCCATCCCCAAAAAATAAAAAAATAAAAAAATAAAGAACATACCT TGGGTTGATCCACTTAGGAACCTCAGATAATAACATCTGCCACGTATAGAGCAATTG CTATGTCCCAGGCACTCTACTAGACACTTCATACAGTTTAGAAAATCAGATGGGTGTA GATCAAGGCAGGAGCAGGAACCAAAAAGAAAGGCATAAACATAAGAAAAAAAATG GAAGGGGTGGAACAGAGTACAATAACATGAGTAATTTGATGGGGGCTATTATGAA CTGAGAAATGAACCTTTGAAAAGTATCTTGGGGCCAAATCATGTAGACTCTTGAGTG TGTGTTAAGGAATGCTATGAGTGCTGAGAGGGCATCAGAAGTCCTTGAGAGCCTCCA GAGAAAGGCTCTTAAAAATGCAGCGCAATCTCCAGTGACAGAAGATACTGCTAGAA ATCTGCTAGAAAAAAAACAAAAAAGGCATGTATAGAGGAATTATGAGGGAAAGATA CCAAGTCACGGTTTATTCTTCAAAATGGAGGTGGCTTGTGGGAAGGTGGAAGCTCA TTTGGCCAGAGTGGAATGGAATTGGGAGAAATCGATGACCAAATGTAAACACTTG TGCCTGATATAGCTTGACACCAAGTTAGCCCCAAGTGAAATACCCTGGCAATATTAA TGTGTCTTTTCCCGATATTCCTCAGGTAATCCAAAGATTCAGGTTTACTCACGTCATCC AGCAGAGAATGGAAAGTCAAATTTCTGAATTGCTATGTGTCTGGGTTTCATCCATCC GACATTGAAGTTGACTTACTGAAGAATGGAGAGAGAATTGAAAAAGTGGAGCATT AGACTTGTCTTTCAGCAAGGACTGGTCTTTCTATCTCTGTACTACACTGAATTCACCC CCACTGAAAAAGATGAGTATGCCTGCCGTGTGAACCATGTGACTTTGTACAGCCCA AGATAGTTAAGTGGGGTAAGTCTTACATTCTTTTGTAAAGCTGCTGAAAGTTGTGTATG AGTAGTCATATCATAAAGCTGCTTTGATATAAAAAAGGTCTATGGCCATACTACCCTG AATGAGTCCCATCCCATCTGATATAACAATCTGCATATTGGGATTGTGAGGGAATGT TCTTAAAGATCAGATTAGTGGCACCTGCTGAGATACTGATGCACAGCATGGTTTCTGA ACCAGTAGTTTCCCTGCAGTTGAGCAGGGAGCAGCAGCAGCACTTGCACAAATACAT ATACACTCTTAACACTTCTTACCTACTGGCTTCTCTAGCTTTTGTGGCAGCTTCAGGT ATATTTAGCACTGAACGAACATCTCAAGAAGGTATAGGCCTTTGTTTGTAAAGTCCTGC TGTCTAGCATCCTATAATCCTGGACTTCTCCAGTACTTTCTGGCTGGATTGGTATCTG AGGCTAGTAGGAAGGGCTTGTTCCTGCTGGGTAGCTCTAAACAATGTATTTCATGGGT AGGAACAGCAGCCTATTCTGCCAGCCTTATTTCTAACCATTITTAGACATTTGTTAGTA CATGGTATTTTAAAAGTAAAACCTAATGTCTTCTTTTTTTTTCTCCACTGTCTTTTTCAT AGATCGAGACATGTAAGCAGCATCATGGAGGTAAGTTTTTGACCTTGAGAAAATGTT TTTGTCTTCACTGTCTGAGGACTATTTATAGACAGCTCTAACATGATAACCCTCACTA TGTGGAGAACATTGACAGAGTAACATTTTAGCAGGGAAAGAAGAATCCTACAGGGTC ATGTTCCCTTCTCCTGTGGAGTGGCATGAAGAAGGTGTATGGCCCCAGGTATGGCCAT ATTACTGACCTCTACAGAGAGGGCAAAGGAACCTGCCAGTATGGTATTGCAGGATAA AGGCAGGTGGTTACCCACATTACCTGCAAGGCTTTGATCTTTCTCTGCCATTTCCAC ATTGGACATCTCTGCTGAGGAGAGAAAAATGAACCACTTTTTCTTTGTATAATGTGTG TTTTATTCTTCAGACAGAAGAGAGGAGTTATACAGCTCTGCAGACATCCCATTCTGT ATGGGGACTGTGTTTGCCTCTTAGAGGTTCCAGGCCACTAGAGGAGATAAAGGGAA ACAGATTGTTATAAATTGATATAATGATACTATAATAGATGTAACATAAGGAGCTC CAGAAGCAAGAGAGAGGGAGGAACCTGGACTTCTCTGCATCTTTAGTTGGAGTCCAA AGGCTTTTCAATGAAATTCTACTGCCAGGGTACATTGATGCTGAAACCCCATTCAAA TCTCCTGTTATATTCTAGAACAGGGAATTGATTTGGGAGAGCATCAGGAAGGTGGAT GATCTGCCCAGTCACACTGTTAGTAAATTGTAGAGCCAGGACCTGAACCTAATATA GTCATGTGTTACTTAATGACGGGGACATGTTCTGAGAAATGCTTACACAAACCTAGG TGTTGTAGCCTACTACACGCATAGGCTACATGGTATAGCCTATTGCTCCTAGACTACA AACCTGTACAGCCTGTTACTGTACTGAATACTGTGGGCAGTTGTAACACAATGGTAA GTATTTGTGTATCTAAACATAGAAGTTGCAGTAAAAATATGCTATTTTAATCTTATGA GACCACTGTCATATATACAGTCCATCATTGACCAAAACATCATATCAGCATTTTTTCT TCTAAGATTTTGGGAGCACCAAAGGGATACACTAACAGGATATACTCTTTATAATGG GTTTGGAGAACTGTCTGCAGCTACTTCTTTAAAAAAGGTGATCTACACAGTAGAAATT AGACAAGTTTGGTAATGAGATCTGCAATCCAAATAAAATAAATTCATTGCTAACCTTT TTCTTTTCTTTTTCAGGTTTGAAGATGCCGCATTTGGATTGGATGAATTCCAAATCTGC TTGCTTGCTTTTAAATATTGATATGCTTATACACTTACACTTTATGCACAAAATGTAGG
---	--

55

(continued)

5		GTTATAATAATGTAAACATGGACATGATCTTCTTTATAATTCTACTTTGAGTGCTGTCT CCATGTTTGATGTATCTGAGCAGGTTGCTCCACAGGTAGCTCTAGGAGGGCTGGCAA CTTAGAGGTGGGGAGCAGAGAATTCTCTTATCCAACATCAACATCTTGGTCAGATTTG AACTCTTCAATCTCTTGCACCTCAAAGCTTGTTAAGATAGTTAAGCGTGCATAAGTTAA CTTCCAATTTACATACTCTGCTTAGAATTTGGGGGAAAATTTAGAAATATAATTGACA GGATTATTGGAAATTTGTTATAATGAATGAAACATTTTGTGCATATAAGATTTCATATTT ACTTCTTATACATTTGATAAAGTAAGGCATGGTTGTGGTTAATCTGGTTTATTTTTGTT CCACAAGTTAAATAAATCATAAAACTTGA (SEO ID NO: 170)
10		
15	HLA-A*02:01: 01: 01 sequence encoding mRNA	CAGAAGCAGAGGGGTCAGGGCGAAGTCCCAGGGCCCCAGGCGTGGCTCTCAGGGTCTCAGGCCCC GAAGGCGGTGTATGGATTGGGGAGTCCCAGCCTTGGGGATTCCCCAAGTCCGCGAGTTTCTTTTCT CCCTCTCCCCAACCTATGTAGGGTCTTCTTCTCGGATACTCACGACGCGGAGCCAGTTCTCACTC CCATTGGGTGTGGGTTTCCAGAGAAGCCAATCAGTGTCTGTCGCGGTGCGGGTTCTAAAGTCCGC ACGCACCCACCGGGACTCAGATTCTCCCCAGACGCCGAGGATGGCCGTCATGGCAGCCCCGAACCC TCGTCTTGTACTCTCGGGGGCTCTGGCCCTGACCCAGACCTGGGCGGGCTCTCACTCCATGAGG TATTTCTTACATCCGTGTCCCGGCCCCGCGCGGGGAGCCCCGCTTCATCGCAGTGGGCTACGT GGACGACACGCAGTTCGTGCGGTTTCGACAGCGACGCCGCGAGCCAGAGGATGGAGCCGCGGGCGC CGTGGATAGAGCAGGAGGGTCCGGAGTATTGGGACGGGGAGACACGGAAGTGAAGGCCCACTCA CAGACTCACCGAGTGGACCTGGGGACCTGCGCGGCTACTACAACAGAGCGAGGCCGGTTCTCA CACCGTCCAGAGGATGTATGGCTGCGACGTGGGGTCCGACTGGCGCTTCTCCGCGGGTACCACC AGTACGCCTACGACGGCAAGGATTACATCGCCCTGAAAGAGGACCTGCGCTCTTGGACCGCGGCG GACATGGCAGCTCAGACCACCAAGCACAAAGTGGGAGCGGCCCATGTGGCGGAGCAGTTGAGAGC CTACCTGGAGGGCACGTGCGTGGAGTGGCTCCGCAGATACCTGGAGAACGGGAGGAGACGCTGC AGCGCACGGACGCCCCAAAACGCATATGACTCACCACGCTGTCTCTGACCATGAAGCCACCCTG AGGTGCTGGGCCCTGAGCTTCTACCCTGCGGAGATCACACTGACCTGGCAGCGGGATGGGGAGGA CCAGACCCAGGACACGGAGCTCGTGGAGACCAGGCCTGCAGGGGATGGAACCTTCCAGAAGTGGG CGGCTGTGGTGGTGCCCTTCTGGACAGGAGCAGAGATACACCTGCCATGTGCAGCATGAGGGTTTG CCCCAAGCCCCCTCACCTGAGATGGGAGCCGTCTTCCCAGCCCCACCATCCCCATCGTGGGCATCAT TGCTGGCCTGGTTCTCTTTGGAGCTGTGATCACTGGAGCTGTGGTCTGCTGTGTATGTGGAGGA GGAAGAGCTCAGATAGAAAAGGAGGGAGCTACTCTCAGGCTGCAAGCAGTGACAGTGCCCCAGGGC TCTGATGTGTCTCTCACAGCTTGTAAAGTGTGAGACAGCTGCCTTGTGTGGGACTGAGAGGCAAG AGTTGTTCTCTGCCCTTCCCTTTGTGACTTGAAGAACCCTGACTTTGTTTCTGCAAAAGGCACCTGC ATGTGTCTGTGTTCGTGTAGGCATAATGTGAGGAGGTGGGGAGACCACCCACCCCCATGTCCAC CATGACCCCTCTTCCCACGCTGACCTGTGCTCCCTCCCCAATCATCTTCTGTTCCAGAGAGGTG GGCTGAGGTGTCTCCATCTCTGCTCAACTTCATGGTGCAGTGAAGTCTTCTCTTCTCC TATTAATAA (SEQ ID NO: 171)
20		
25		
30		
35		
40	HLA-A*02 (GenBank: LK021978.1)	ATGGCCGTCATGGCGCCCCGAACCCTCGTCTCTGCTACTCTCGGGGGCTCTGGCCCTGA CCCAGACCTGGGCGGGTGAGTGCGGGGTGCGGAGGGAAACGGCCTCTGTGGGGAGA AGCAACGGGCCCCGCTGGCGGGGGCGCAGGACCCGGGAAGCCGCGCCGGGAGGAGG GTGCGGGCGGGTCTCAGCCACTCCTCGTCCCCAGGCTCTCACTCCATGAGGTATTTCTT CACATCCGTGTCCCGGCCCCGCGCGGGGAGCCCCGCTTCATCGCAGTGGGCTACGT GGACGACACGCAGTTTCGTGCGGTTTCGACAGCGACGCCGCGAGCCAGAGGATGGAGC CGCGGGCGCCGTGGATAGAGCAGGAGGGTCCGGAGTATTGGGACGGGGAGACACGG AAAGTGAAGGCCCACTCACAGACTACCGAGTGGACCTGGGGACCCTGCGCGGCTAC TACAACCAGAGCGAGGCCGGTGAGTGACCCCGGCCCGGGGCGCAGGTACGACCTCT CATCCCCACGGACGGGCCAGGTGCCCCACAGTCTCCGGGTCCGAGATCCGCCCCGA AGCCGCGGGACCCCCGAGACCCTTGCCCCGGGAGAGGCCAGGCGCCTTTACCCGGTT TCATTTTCAGTTTAGGCCAAAAATCCCCCAGGTTGGTGGGGCGGGGCGGGGCTCG GGGGACCGGGCTGACCGCGGGGTCCGGGCCAGGTTCTCACACCGTCCAGAGGATGTA TGGCTGCGACGTGGGGTCCGACTGGCGCTTCTCCGCGGGTACCACCAGTACGCCTA CGACGGCAAGGATTACATCGCCCTGAAAGAGGACCTGCGCTCTTTGACCGCGGCGGA CATGGCAGCTCAGACCACCAAGCACAAAGTGGGAGGCGGCCCATGTGGCGGAGCAGT TGAGAGCCTACCTGGAGGGCACGTGCGTGGAGTGGCTCCGCGAGATACCTGGAGAACG GGAAGGAGACGCTGCAGCGCACGGGTACCAGGGGCCACGGGGCGCCTCCCTGATCG CCTGTAGATCTCCCGGGCTGGCCTCCACAAAGAGGGGAGACAATTGGGACCAACAC TAGAATATCGCCCTCCCTCTGCTCTGAGGGAGAGGAATCCTCCTGGGTTTCCAGATC CTGTACCAGAGAGTGACTCTGAGGTTCCGCCCTGCTCTCTGACACAATTAAGGGATA
45		
50		
55		

(continued)

5	AAATCTCTGAAGGAATGACGGGAAGACGATCCCTCGAATACTGATGAGTGGTTCCCT
	TTGACACACACAGGCAGCAGCCTTGGGCCCGTGACTTTTCCTCTCAGGCCTTGTTCTC
	TGCTTCACTCAATGTGTGTGGGGTCTGAGTCCAGCACTTCTGAGTCCCTTCAGCCT
	CCACTCAGGTCAGGACCAGAAGTCGCTGTTCCCTCTTCAGGGACTAGAATTTCCACG
	GAATAGGAGATTATCCCAGGTGCCTGTGTCCAGGCTGGTGTCTGGGTTCTGTGCTCCC
	TTCCCCATCCCAGGTGTCCTGTCCATTCTCAAGATAGCCACATGTGTGCTGGAGGAGT
	GTCCCATGACAGATGCAAAATGCCTGAATGATCTGACTCTTCCTGACAGACGCCCCC
10	AAAACGCATATGACTCACCACGCTGTCTCTGACCATGAAGCCACCCTGAGGTGCTGG
	GCCCTGAGCTTCTACCCTGCGGAGATCACACTGACCTGGCAGCGGGATGGGGAGGAC
	CAGACCCAGGACACGGAGCTCGTGGAGACCAGGCCTGCAGGGGATGGAACCTTCCA
	GAAGTGGGCGGCTGTGGTGGTGCCTTCTGGACAGGAGCAGAGATACACCTGCCATGT
	GCAGCATGAGGGTTTGCCCAAGCCCCCTCACCTGAGATGGGGTAAGGAGGGAGACG
	GGGGTGTCTGTCTTTTAGGGAAAGCAGGAGCCTCTCTGACCTTTAGCAGGGTCAGG
15	GCCCCCTACCTTCCCCTCTTTTCCCAGAGCCGTCTTCCCAGCCCACCATCCCCATCGTG
	GGCATCATTGCTGGCCTGGTTCTCTTTGGAGCTGTGATCACTGGAGCTGTGGTGCCTG
	CTGTGATGTGGAGGAGGAAGAGCTCAGGTGGGGAAGGGGTGAAGGGTGGGTCTGAG
	ATTTCTTGTCTCACTGAGGGTTCCAAGACCCAGGTAGAAAGTGTGCCCTGCCTCGTTAC
	TGGGAAGCACCAACCAATTATGGGCCTACCCAGCCTGGGCCCTGTGTGCCAGCAC
20	TTACTCTTTTGTAAAGCACCTGTTAAATGAAGGACAGATTTATCACCTTGATTACAG
	CGGTGATGGGACCTGATCCCAGCAGTCACAAGTCACAGGGGAAGGTCCCTGAGGACC
	TTCAGGAGGGCGGTTGGTCCAGGACCCACACCTGCTTCTTCATGTTTCCTGATCCCG
	CCCTGGGTCTGCAGTCACACATTTCTGGAACTTCTCTGAGGTCCAAGACTTGGAGGT
	TCCTCTAGGACCTTAAGGCCCTGACTCCTTTCTGGTATCTCACAGGACATTTTCTTCCC
	ACAGATAGAAAAGGAGGGAGCTACTCTCAGGCTGCAAGTAAGTATGAAGGAGGCTG
25	ATGCCCTGAGGTCTTGGGATATGTGTTTGGGAGCCCATGGGGGAGCTCACCCACCC
	CACAATTCCTCCTCTAGCCACATCTTCTGTGGGATCTGACCAGGTCTGTTTTGTCT
	ACCCACAGGCAGTGACAGTGCCAGGGCTCTGATGTGTCTCTCACAGCTTGTAAGGT
	GAGAGCCTGGAGGGCCTGATGTGTGTTGGGTGTTGGGCGGAACAGTGGACACAGCTG
	TGCTATGGGGTTTCTTTCCATTGGATGTATTGAGCATGCGATGGGCTGTTTAAAGTGT
30	GACCCCTCACTGTGACAGATACGAATTTGTTTCATGAATATTTTTTCTATAGTGTGAG
	ACAGCTGCCTTGTGTGGGACTGAGAGGCAAGAGTTGTTCTGCCCTTCCCTTTGTGAC
	TTGAAGAACCCTGACTTTGTTTCTGCAAAGGCACCTGCATGTGTCTGTGTTTCGTGTAG
	GCATAATGTGAGGAGGTGGGGAGACCACCCACCCCATGTCCACCATGACCCTCTT
	CCCACGCTGACCTGTGCTCCCTCCCCAATCATCTTTCCTGTTCCAGAGAGGTGGGGCT
35	GAGGTGTCTCCATCTCTGTCTCAACTTCATGGTGCCTGAGCTGTAACCTCTTCCTTCC
	CTATTAATAA (SEQ ID NO: 169)

[0356] The person of ordinary skill in the art will appreciate that T can be substituted for U to convert an RNA sequence to a DNA sequence and vice versa, and both are envisaged as target gene sequences of the disclosure.

[0357] A target gene is edited in the immune cells described herein using a nucleic acid guided endonuclease. Exemplary nucleic acid guided endonucleases include Class II endonucleases, such as CRISPR/Cas9.

[0358] "CRISPR" or "CRISPR gene editing" as used herein refers to a set of clustered regularly interspaced short palindromic repeats, or a system comprising such a set of repeats. "Cas", as used herein, refers to a CRISPR-associated protein. A "CRISPR/Cas" system refers to a system derived from CRISPR and Cas, which can be used to silence, knock out, or mutate a target gene. This system is a type of prokaryotic immune system that confers resistance to foreign genetic elements such as plasmids and phages and provides a form of acquired immunity. The CRISPR/Cas system has been modified for use in gene editing. This is accomplished by introducing into the eukaryotic cell a one or more specifically designed guide nucleic acids (gNAs), typically guide RNAs (gRNAs), and an appropriate Cas endonuclease, which forms a ribonucleoprotein complex with the gNA. The gNA guides the gNA-endonuclease protein complex to a target genomic location, and the endonuclease introduces a strand break at the target genomic location. This double strand break can be repaired by cellular mechanisms such as non-homologous end joining (leading to deletions) or homologous repair (which can generate insertions), thereby introducing genetic modifications into the host cell genome.

[0359] CRISPR/Cas systems are classified by class and by type. Class 2 systems currently represent a single interference protein that is categorized into three distinct types (types II, V and VI). Any class 2 CRISPR/Cas system suitable for gene editing, for example a type II, a type V or a type VI system, is envisaged as within the scope of the instant disclosure. Exemplary Class 2 type II CRISPR systems include Cas9, Csn2 and Cas4. Exemplary Class 2, type V CRISPR systems include, Cas12, Cas12a (Cpf1), Cas12b (C2c1), Cas12c (C2c3), Cas12d (CasY), Cas12e (CasX), Cas12f, Cas12g, Cas12h, Cas12i and Cas12k (C2c5). Exemplary Class 2 Type VI systems include Cas13, Cas13a

(C2c2) Cas13b, Cas13c and Cas13d.

[0360] The CRISPR sequence, sometimes called a CRISPR locus, comprises alternating repeats and spacers. In a naturally-occurring CRISPR, the spacers usually comprise sequences foreign to the bacterium such as a plasmid or phage sequence. As described herein, spacer sequences may also be referred to as "targeting sequences". In CRISPR/Cas systems for a genetic engineering, the spacers are derived from the target gene sequence (the gNA).

[0361] An exemplary Class 2 type II CRISPR system relies on the protein Cas9, which is a nuclease with two active cutting sites, one for each strand of the double helix. Combining Cas9 and modified CRISPR locus RNA can be used in a system for gene editing. Pennisi (2013) Science 341: 833-836. The Cas protein may be used to modify the immune cells is Cas9.

[0362] The CRISPR/Cas system can thus be used to edit a target gene, such as a gene targeted for editing in the immune cells described herein, by adding or deleting a base pair, or introducing a premature stop, which thus decreases expression of the target. The CRISPR/Cas system can alternatively be used like RNA interference, turning off a target gene in a reversible fashion. In a mammalian cell, for example, the RNA can guide the Cas protein to a target gene promoter, sterically blocking RNA polymerases.

[0363] A Cas protein may be derived from any bacterial or archaeal Cas protein. Any suitable CRISPR/Cas system is envisaged as within the scope of the instant disclosure. In other aspects, Cas protein comprises one or more of Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9, Cas10, Cas12a (Cpf1), Cas13, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, CasX, CasY, homologs thereof, or modified versions thereof. The Cas protein may be a Cas9 protein, a Cpf1 protein, a C2c1 protein, a C2c2 protein, a C2c3 protein, Cas3, Cas3-HD, Cas 5, Cas7, Cas8, Cas10, or combinations or complexes of these. The Cas protein may be a Cas9 protein.

[0364] Artificial CRISPR/Cas systems can be generated which inhibit a target gene, using technology known in the art, *e.g.*, that described in U.S. Publication No. 20140068797, and Cong (2013) Science 339: 819-823. Other artificial CRISPR/Cas systems that are known in the art may also be generated, which inhibit a target gene, *e.g.*, that described in Tsai (2014) Nature Biotechnol., 32:6 569-576, U.S. Pat. Nos. 8,871,445; 8,865,406; 8,795,965; 8,771,945; and 8,697,359. Methods of designing suitable gNAs for a particular Cas protein will be known by persons of ordinary skill in the art.

[0365] The present disclosure provides gene-targeting guide nucleic acids (gNAs) that can direct the activities of an associated polypeptide (*e.g.*, nucleic acid guided endonuclease) to a specific target gene sequence within a target nucleic acid genome. The genome-targeting nucleic acid can be an RNA. A genome-targeting RNA is referred to as a "guide RNA" or "gRNA" herein. A guide RNA can comprise at least a targeting sequence that hybridizes to a target nucleic acid sequence of interest, and a CRISPR repeat sequence. In some Type II systems, the gRNA also comprises a second RNA called the tracrRNA sequence, also referred to herein as a "scaffold" sequence. In the Type II guide RNA (gRNA), the CRISPR repeat sequence and scaffold sequence hybridize to each other to form a duplex. In the Type V guide RNA (gRNA), the crRNA forms a duplex. In both systems, the duplex can bind a site-directed polypeptide, such that the guide RNA and site-directed polypeptide form a complex. The gene-targeting nucleic acid can provide target specificity to the complex by virtue of its association with the site-directed polypeptide. The gene-targeting nucleic acid thus can direct the activity of the site-directed polypeptide.

[0366] The disclosure provides a guide RNA comprising a targeting sequence and a guide RNA scaffold sequence, wherein the targeting sequence is complementary to the sequence of a target gene.

[0367] Exemplary guide RNAs include targeting sequences of about 15-20 bases. As is understood by the person of ordinary skill in the art, each gRNA can be designed to include a targeting sequence complementary to its genomic target sequence. For example, each of the targeting sequences, *e.g.*, the RNA version of the DNA sequences presented in Table 8, minus the three 3' nucleotides which represent that PAM site, can be put into a single RNA chimera or a crRNA.

[0368] The gene targeting nucleic acid can be a double-molecule guide RNA. The gene targeting nucleic acid can be a single-molecule guide RNA. The gene targeting nucleic acid can be any known configuration of guide RNA known in the art, such as, for example, including paired gRNA, or multiple gRNAs used in a single step. Although it is clear from genomic sequences where the coding sequences and splice junctions are, other features required for gene expression may be idiosyncratic and unclear.

[0369] A double-molecule guide RNA can comprise two strands of RNA. The first strand comprises a sequence in the 5' to 3' direction, an optional spacer extension sequence, a targeting sequence and a minimum CRISPR repeat sequence. The second strand can comprise a minimum tracrRNA sequence (complementary to the minimum CRISPR repeat sequence), a 3' tracrRNA sequence and an optional tracrRNA extension sequence.

[0370] A single-molecule guide RNA (sgRNA) in a Type II system can comprise, in the 5' to 3' direction, an optional spacer extension sequence, a targeting sequence, a minimum CRISPR repeat sequence, a single-molecule guide linker, a minimum tracrRNA sequence, a 3' tracrRNA sequence and an optional tracrRNA extension sequence. The optional tracrRNA extension can comprise elements that contribute additional functionality (*e.g.*, stability) to the guide RNA. The

single-molecule guide linker can link the minimum CRISPR repeat and the minimum tracrRNA sequence to form a hairpin structure. The optional tracrRNA extension can comprise one or more hairpins.

[0371] Guide RNA or single-molecule guide RNA (sgRNA) can comprise a targeting sequence and a scaffold sequence. The scaffold sequence may be a Cas9 gRNA sequence. The scaffold sequence may be encoded by a DNA sequence that comprises a sequence that shares at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to GTTTTAGAGCTAGAAATAGCAAGTTAAATAAGGCTAGTCCGTTATCAACTTGAAAA AGTGGCAC-CGAGTCGGTGCTTTTTTTT (SEQ ID NO: 435). The scaffold sequence may be encoded by a DNA sequence that comprises GTTTTAGAGCTAGAAATAGCAAGTTAAA ATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCAC-CGAGTCGGTGCTTTTTTTT (SEQ ID NO: 435).

[0372] For example, those embodiments where the CRISPR/Cas system is a Cas9 system, the sgRNA can comprise a 20 nucleotide targeting sequence at the 5' end of the sgRNA sequence. The sgRNA can comprise a less than a 20 nucleotide targeting sequence at the 5' end of the sgRNA sequence. The sgRNA can comprise a more than 20 nucleotide targeting sequence at the 5' end of the sgRNA sequence. The sgRNA can comprise a variable length targeting sequence with 17-30 nucleotides at the 5' end of the sgRNA sequence.

[0373] Suitable scaffold sequences, and arrangement of scaffold targeting sequences, will depend on choice of endonuclease, and will be known to persons of skill in the art.

[0374] A single-molecule guide RNA (sgRNA) in a Type II system, e.g. Cas9, can comprise, in the 5' to 3' direction, a minimum CRISPR repeat sequence and a targeting sequence.

[0375] By way of illustration, guide RNAs used in the CRISPR/Cas9 or CRISPR/Cpf1 system, or other smaller RNAs can be readily synthesized by chemical means, as illustrated below and described in the art. While chemical synthetic procedures are continually expanding, purifications of such RNAs by procedures such as high performance liquid chromatography (HPLC, which avoids the use of gels such as PAGE) tends to become more challenging as polynucleotide lengths increase significantly beyond a hundred or so nucleotides. One approach used for generating RNAs of greater length is to produce two or more molecules that are ligated together. Much longer RNAs, such as those encoding a Cas9 or Cpf1 endonuclease, are more readily generated enzymatically. Various types of RNA modifications can be introduced during or after chemical synthesis and/or enzymatic generation of RNAs, e.g., modifications that enhance stability, reduce the likelihood or degree of innate immune response, and/or enhance other attributes, as described in the art.

[0376] The targeting sequence of a gRNA hybridizes to a sequence in a target nucleic acid of interest. The targeting sequence of a genome-targeting nucleic acid can interact with a target nucleic acid in a sequence-specific manner via hybridization (i.e., base pairing). The nucleotide sequence of the targeting sequence can vary depending on the sequence of the target nucleic acid of interest.

[0377] In a Cas9 system described herein, the targeting sequence can be designed to hybridize to a target nucleic acid that is located 5' of the reverse complement of a PAM of the Cas9 enzyme used in the system. The targeting sequence may perfectly match the target sequence or may have mismatches. Each CRISPR/Cas system protein may have a particular PAM sequence, in a particular orientation and position, that it recognizes in a target DNA. For example, *S. pyogenes* Cas9 recognizes in a target nucleic acid a PAM that comprises the sequence 5'-NRG-3', where R comprises either A or G, where N is any nucleotide and N is immediately 3' of the target nucleic acid sequence targeted by the targeting sequence. Selection of appropriate PAM sequences will be apparent to the person of ordinary skill in the art.

[0378] The target sequence is complementary to, and hybridizes with, the targeting sequence of the gRNA. The target nucleic acid sequence can comprise 20 nucleotides. The target nucleic acid can comprise less than 20 nucleotides. The target nucleic acid can comprise more than 20 nucleotides. The target nucleic acid can comprise at least: 5, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30 or more nucleotides. For example those embodiments where the CRISPR/Cas system is a Cas9 system, the target nucleic acid sequence can comprise 20 nucleotides immediately 5' of the first nucleotide of the reverse complement of the PAM sequence. This target nucleic acid sequence is often referred to as the PAM strand or a target strand, and the complementary nucleic acid sequence is often referred to the non-PAM strand or non-target strand. One of skill in the art would recognize that the targeting sequence hybridizes to the non-PAM strand of the target nucleic acid, see e.g., US20190185849A1.

[0379] In some examples, the percent complementarity between the targeting sequence and the target nucleic acid is at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 97%, at least about 98%, at least about 99%, or 100%. In some examples, the percent complementarity between the targeting sequence and the target nucleic acid is at most about 30%, at most about 40%, at most about 50%, at most about 60%, at most about 65%, at most about 70%, at most about 75%, at most about 80%, at most about 85%, at most about 90%, at most about 95%, at most about 97%, at most about 98%, at most about 99%, or 100%. In some examples, the percent complementarity between the targeting sequence and the target nucleic acid is 100% over the six contiguous 5'-most nucleotides of the target sequence of the complementary strand of the target nucleic acid. The percent complementarity between the targeting sequence and the target nucleic acid can be at least 60% over about 20 contiguous

nucleotides. The length of the targeting sequence and the target nucleic acid can differ by 1 to 6 nucleotides, which may be thought of as a bulge or bulges.

[0380] The targeting sequence can be designed or chosen using computer programs known to persons of ordinary skill in the art. The computer program can use variables, such as predicted melting temperature, secondary structure formation, predicted annealing temperature, sequence identity, genomic context, chromatin accessibility, % GC, frequency of genomic occurrence (e.g., of sequences that are identical or are similar but vary in one or more spots as a result of mismatch, insertion or deletion), methylation status, presence of SNPs, and the like. Available computer programs can take as input NCBI gene IDs, official gene symbols, Ensembl Gene IDs, genomic coordinates, or DNA sequences, and create an output file containing sgRNAs targeting the appropriate genomic regions designated as input. The computer program may also provide a summary of statistics and scores indicating on- and off-target binding of the sgRNA for the target gene (Doench et al. Nat Biotechnol. 34:184-191 (2016)). The disclosure provides guide RNAs comprising a targeting sequence. The guide RNA may further comprise a guide RNA scaffold sequence. As described herein, the targeting sequence is complementary to the sequence of a target gene selected from the group consisting of HLA-A, HLA-B, HLA-C, B2M or an allele thereof. As also described herein, the target gene is an HLA-A gene. As also described herein, the target gene is an HLA-B gene. As also described herein, the target gene is an HLA-C gene. As also described herein, the target gene is HLA-A, HLA-B, HLA-C, or a combination thereof. The targeting sequence may comprise a sequence that shares about 90%, about 95%, about 96%, about 97%, about 98%, about 99% identity to or may be identical to a sequence disclosed in Table 7.

[0381] As described herein, the gNAs specifically target the sequence of an endogenous HLA-A locus. As also described herein, the gNAs that specifically target the sequence of an HLA-A locus comprise a sequence that shares about 90%, about 95%, about 96%, about 97%, about 98%, or about 99% identity to a sequence selected from the sequences disclosed in Table 8. As also described herein, the gNAs that specifically target the sequence of an HLA-A locus comprise a sequence selected from the sequences disclosed in Table 8.

[0382] In some embodiments, the gNAs specifically target a sequence of HLA-A*02 alleles. For example, the gRNAs specifically target, and hybridize to, a sequence shared by all HLA-A*02 alleles, but that is not shared by HLA-A*02 and HLA-A*03 alleles. In some embodiments, the gNAs specifically target a sequence of HLA-A*02:01 alleles. In some embodiments, the gNAs specifically target a sequence of HLA-A*02:01:01 alleles. In some embodiments, the gNAs specifically target a sequence of HLA-A*02:01:01:01 alleles. In some embodiments, the gNAs specifically target a sequence of HLA-A*02:01:01:01 alleles.

[0383] In some embodiments, the gNAs specifically target a coding DNA sequence of HLA-A*02.

[0384] In some embodiments, the gNAs specifically target a coding DNA sequence that is shared by more than 1000 HLA-A*02 alleles. In some embodiments, the gNAs that specifically target a coding DNA sequence in greater than 1000 HLA-A*02 alleles comprise a sequence that shares about 90%, about 95%, about 96%, about 97%, about 98%, about 99% identity or is identical to a sequence selected from SEQ ID NOs: 400-465.

Table 8. Illustrative sequences targeting HLA-A and HLA-A alleles

SEQ ID NO	Guide Nucleic Acid Targeting Sequences
436	TGGACGACACGCAGTTCGTG
437	CAGATACCTGGAGAACGGGA
438	TCCCGTTCTCCAGGTATCTG
439	CCGCCGCGGTCCAAGAGCGC
440	CCTGCGCTCTTGACCGCGG
441	GGACCTGCGCTCTTGACCGC
442	AAGGAGACGCTGCAGCGCACGGG
443	GAAGGAGACGCTGCAGCGCACGG
444	GCGGGCGCCGTGGATAGAGCAGG
445	TGCTCTATCCACGGCGCCCGCGG
446	CGATGAAGCGGGGCTCCCCGCGG
447	CGTGTCGGCGCCCGGCCGCGGGG
448	CGGCTCCATCCTCTGGCTCGCGG
449	GATGTAATCCTTGCCGTCGTAGG

(continued)

	SEQ ID NO	Guide Nucleic Acid Targeting Sequences
5	450	ACAGCGACGCCGCGAGCCAGAGG
	451	GGATGGAGCCGCGGGCGCCGTGG
	452	GGCGCCGTGGATAGAGCAGGAGG
	453	GCGCCGTGGATAGAGCAGGAGGG
10	454	CGGCTACTACAACCAGAGCGAGG
	455	CTGGTTGTAGTAGCCGCGCAGGG
	456	TACTACAACCAGAGCGAGGCCGG
15	457	CTACCTGGAGGGCACGTGCGTGG
	458	CACGCACGTGCCCTCCAGGTAGG
	459	GCAGGGTCCCCAGGTCCACTCGG
	460	GTGGACCTGGGGACCCTGCGCGG
20	461	TGGAGGGCACGTGCGTGGAGTGG
	462	GTATGGCTGCGACGTGGGGTCGG
	463	CTGAGCTGCCATGTCCGCCGCGG
25	464	GGATTACATCGCCCTGAAAGAGG
	465	CAAGTGGGAGGCGGCCCATGTGG
	466	GTGGGAGGCGGCCCATGTGGCGG
	467	CAGTTGAGAGCCTACCTGGAGGG
30	468	GCAGTTGAGAGCCTACCTGGAGG
	469	TACCACCAGTACGCCTACGACGG
	470	TGCCGTCGTAGGCGTACTGGTGG
35	471	CCAGTACGCCTACGACGGCAAGG
	472	GGATGTGAAGAAATACCTCATGG
	473	ATTTCTTCACATCCGTGTCCCGG
	474	AGGCGTACTGGTGGTACCCGCGG
40	475	CGTACTGGTGGTACCCGCGGAGG
	476	GAGGATGTATGGCTGCGACGTGG
	477	GGATGTATGGCTGCGACGTGGGG
45	478	CTCAGACCACCAAGCACAAGTGG
	479	TCAGACCACCAAGCACAAGTGGG
	480	CACCAAGCACAAGTGGGAGGCGG
50	481	GACCACCAAGCACAAGTGGGAGG
	482	GAGCCCCGCTTCATCGCAGTGGG
	483	GTAGCCCACTGCGATGAAGCGGG
	484	TAGCCCACTGCGATGAAGCGGGG
55	485	CGTAGCCCACTGCGATGAAGCGG
	486	CTTCATCGCAGTGGGCTACGTGG
	487	GGAGCCCCGCTTCATCGCAGTGG

(continued)

5

10

15

20

25

30

35

40

45

50

55

SEQ ID NO	Guide Nucleic Acid Targeting Sequences
488	CGGGGAGACACGGAAAGTGAAGG
489	AGTATTGGGACGGGGAGACACGG
490	AGGGTCCGGAGTATTGGGACGGG
491	GAGGGTCCGGAGTATTGGGACGG
492	GGACCCTCCTGCTCTATCCACGG
493	GTGGATAGAGCAGGAGGGTCCGG
494	AGACTCACCGAGTGGACCTGGGG
495	CACTCGGTGAGTCTGTGAGTGGG
496	CAGACTCACCGAGTGGACCTGGG
497	CCACTCACAGACTCACCGAGTGG
498	CCACTCGGTGAGTCTGTGAGTGG
499	TCGGACTGGCGCTTCTCCGCGG
500	GCAGCCATACATCCTCTGGACGG
501	TCTCAACTGCTCCGCCACATGGG
502	ACCCTCATGCTGCACATGGCAGG
503	ACCTGCCATGTGCAGCATGAGGG
504	CACCTGCCATGTGCAGCATGAGG
505	GGAGGACCAGACCCAGGACACGG
506	GGATGGGGAGGACCAGACCCAGG
507	GACCTGGCAGCGGGATGGGGAGG
508	AGATCACACTGACCTGGCAGCGG
509	GATCACACTGACCTGGCAGCGGG
510	AGGTCAGTGTGATCTCCGCAGGG
511	AAGCCCCTCACCTGAGATGGGG
512	CTGCGGAGATCACACTGACCTGG
513	CAGCAATGATGCCCACGATGGGG
514	CCAGCAATGATGCCCACGATGGG
515	GCCAGCAATGATGCCCACGATGG
516	GGATGGAACCTTCCAGAAGTGGG
517	GGGATGGAACCTTCCAGAAGTGG
518	ATGCCCACGATGGGGATGGTGGG
519	CAGCCCACCATCCCCATCGTGGG
520	CCAGCCCACCATCCCCATCGTGG
521	GATGCCCACGATGGGGATGGTGG
522	CAGGGCCCAGCACCTCAGGGTGG
523	AATGATGCCCACGATGGGGATGG
524	GGCCCTGACCCAGACCTGGGCGG
525	GACCCAGGACACGGAGCTCGTGG

(continued)

SEQ ID NO	Guide Nucleic Acid Targeting Sequences
526	ACACGGAGCTCGTGGAGACCAGG
527	CGTGGAGACCAGGCCTGCAGGGG
528	TCGTGGAGACCAGGCCTGCAGGG
529	AGCTGTGATCACTGGAGCTGTGG
530	AAAAGGAGGGAGCTACTCTCAGG
531	ATGTGGAGGAGGAAGAGCTCAGG
532	GTGTCTCTCACAGCTTGTAAGG
533	GAGAGACACATCAGAGCCCTGGG
534	CTCCGCAGGGTAGAAGCTCAGGG
535	GGCCCTGAGCTTCTACCCTGCGG
536	GCTCAGGGCCCAGCACCTCAGGG
537	TATCTCTGCTCCTGTCCAGAAGG
538	AGTAGCAGGACGAGGGTTCGGGG
539	CCCCGAGAGTAGCAGGACGAGGG
540	CCCTCGTCCTGCTACTCTCGGGG
541	CCTCGTCCTGCTACTCTCGGGG
542	CTGTGGTCGCTGCTGTGATGTGG
543	TCGCTGCTGTGATGTGGAGGAGG
544	TGGTCGCTGCTGTGATGTGGAGG
545	CACAGCCGCCCACTTCTGGAAGG
546	CCAGAAAGTGGGCGGCTGTGGTGG
547	TGGAACCTTCCAGAAGTGGGCGG
548	TCACAGCTCCAAAGAGAACCAGG
549	CTGACCATGAAGCCACCCTGAGG
550	GCAAACCCTCATGCTGCACATGG
551	TGAAGCCACCCTGAGGTGCTGGG
552	GGTGAGTCATATGCGTTTTGGGG
553	GTGAGTCATATGCGTTTTGGGGG
554	CTTCATGGTCAGAGACAGCGTGG
555	TCTGGCCCTGACCCAGACCTGGG

[0385] The sequences disclosed in Table 8 include the corresponding genomic sequences, inclusive of the PAM sequence. The skilled artisan will understand that the targeting sequence of the gRNA does not include three 3' terminal nucleotides of the sequences in Table 8, which represent the corresponding PAM site for the gRNA.

[0386] The disclosure provides gNAs comprising a targeting sequence specific to the *B2M* gene. In some embodiments, the gNAs specifically target the coding sequence (CDS) sequence of the *B2M* gene. In some embodiments, the gNA comprises a sequence that targets the *B2M* gene promoter sequence.

[0387] In some embodiments the gNA comprise a targeting sequence and a gNA scaffold sequence. In some embodiments, the targeting sequence comprises a sequence set forth in Table 9, or a sequence shares about 90%, about 95%, about 96%, about 97%, about 98%, about 99% identity thereto.

[0388] In some embodiments, the targeting sequence is complementary to a sequence of the *B2M* gene. In some

embodiments, the *B2M* gene comprises a sequence that shares about 90%, about 95%, about 96%, about 97%, about 98%, about 99% identity to the B2M sequence set forth in Table 7.

Table 9. Illustrative sequences targeting *B2M*

SEQ ID NO	Sequence
556	CGCGAGCACAGCTAAGGCCA
557	GAGTAGCGCGAGCACAGCTA
558	AGGGTAGGAGAGACTCACGC
559	CTGAATCTTTGGAGTACCTG
560	TCACGTCATCCAGCAGAGAA
561	TCCTGAATTGCTATGTGTCT
562	AAGTCAACTTCAATGTCGGA
563	GTCTTTTCCCGATATTCCTC
564	TGGAGTACCTGAGGAATATC
565	CAGCCCAAGATAGTTAAGTG
566	ACAAAGTCACATGGTTCACA
567	ACTCTCTCTTCTGGCCTGG
568	TGGGCTGTGACAAAGTCACA
569	GGCCGAGATGTCTCGCTCCG
570	CAGTAAGTCAACTTCAATGT
571	ACTCACGCTGGATAGCCTCC
572	CATACTCATCTTTTTCAGTG
573	CACAGCCCAAGATAGTTAAG
574	TTCAGACTTGCTTTTCAGCA
575	AGTCACATGGTTCACACGGC
576	ATACTCATCTTTTTCAGTGG
577	GGCATACTCATCTTTTTCAG
578	ACAGCCCAAGATAGTTAAGT
579	GCTACTCTCTTTTCTGGCC
580	TGGAGAGAGAATTGAAAAAG
581	ACTTGTCTTTCAGCAAGGAC
582	GAAGTTGACTTACTGAAGAA
583	GGCCACGGAGCGAGACATCT
584	GCATACTCATCTTTTTCAGT
585	CGTGAGTAAACCTGAATCTT
586	TTACCCCACTTAACTATCTT
587	TTGGAGTACCTGAGGAATAT
588	ACCCAGACACATAGCAATTC
589	TTTGACTTTCATTCTCTGC
590	TTCCTGAATTGCTATGTGTC
591	CTCAGGTACTCCAAAGATTC

(continued)

5

10

15

20

25

30

35

40

45

50

55

SEQ ID NO	Sequence
592	CTTACCCCACTTAACTATCT
593	CTCGCGCTACTCTCTCTTTC
594	TCGATCTATGAAAAAGACAG
595	GAGACATGTAAGCAGCATCA
596	ACATGTAAGCAGCATCATGG
597	GAAGTCCTAGAATGAGCGCC
598	GAGCGCCCGGTGTCCCAAGC
599	AGCGCCCGGTGTCCCAAGCT
600	GCGCCCGGTGTCCCAAGCTG
601	CTGGGGCGCGCACCCCAGAT
602	GGGCGCGCACCCCAGATCGG
603	GGCGCGCACCCCAGATCGGA
604	CATCACGAGACTCTAAGAAA
605	TAAGAAAAGGAAACTGAAAA
606	AAGAAAAGGAAACTGAAAAC
607	GAAAGTCCCTCTCTTAACC
608	CTAACCTGGCACTGCGTCGC
609	CTGGCACTGCGTCGCTGGCT
610	TGCGTCGCTGGCTTGGAGAC
611	GCTGGCTTGGAGACAGGTGA
612	GAGACAGGTGACGGTCCCTG
613	AGACAGGTGACGGTCCCTGC
614	CCTGCGGGCCTTGTCTGAT
615	CGGGCCTTGTCTGATTGGC
616	GGGCCTTGTCTGATTGGCT
617	GGGCACGCGTTTAATATAAG
618	CACGCGTTTAATATAAGTGG
619	TATAAGTGGAGGCGTCGCGC
620	AAGTGGAGGCGTCGCGCTGG
621	AGTGGAGGCGTCGCGCTGGC
622	TTCCTGAAGCTGACAGCATT
623	TCCTGAAGCTGACAGCATTC
624	GCCCGAATGCTGTCAGCTTC
625	AAACGCGTGCCCAGCCAATC
626	GTGCCCAGCCAATCAGGACA
627	CCAATCAGGACAAGGCCCGC
628	CAATCAGGACAAGGCCCGCA
629	CAAGCCAGCGACGCAGTGCC

(continued)

5

10

15

20

25

30

35

40

45

50

55

SEQ ID NO	Sequence
630	CGCAGTGCCAGGTTAGAGAG
631	GCAGTGCCAGGTTAGAGAGA
632	GAGTCTCGTGATGTTTAAGA
633	TAAGAAGGCATGCACTAGAC
634	AAGAAGGCATGCACTAGACT
635	TGAGTTTGCTGTCTGTACAT
636	TACATCGGCGCCCTCCGATC
637	ACATCGGCGCCCTCCGATCT
638	CATCGGCGCCCTCCGATCTG
639	CTGGGGTGCGCGCCCCAGCT
640	TGGGGTGCGCGCCCCAGCTT
641	CGCGCCCCAGCTTGGGACAC
642	GCGCCCCAGCTTGGGACACC
643	CAAGTCACTTAGCATCTCTG
644	ACAGAAGTTCTCCTTCTGCT
645	ATTCAAAGATCTTAATCTTC
646	TTCAAAGATCTTAATCTTCT
647	TTTTCTCGAATGAAAAATGC
648	TGCAGGTCCGAGCAGTTAAC
649	GGTCCGAGCAGTTAACTGGC
650	GTCCGAGCAGTTAACTGGCT
651	TCCGAGCAGTTAACTGGCTG
652	AGCAAGTCACTTAGCATCTC
653	GCAAGTCACTTAGCATCTCT
654	TGGGGCCAGTCTGCAAAGCG
655	GGGGCCAGTCTGCAAAGCGA
656	GGGCCAGTCTGCAAAGCGAG
657	GGCCAGTCTGCAAAGCGAGG
658	GGACACCGGGCGCTCATTCT
659	GGCGCTCATTCTAGGACTTC
660	CTCATTCTAGGACTTCAGGC
661	ATTCTAGGACTTCAGGCTGG
662	TTCAGGCTGGAGGCACATTA
663	TGCCCCCTCGCTTTGCAGAC
664	GATGCTAAGTGAATTGCTAA
665	GCCCCAGCCAGTTAACTGCT
666	GCATTTTTCATTGAGAAAA
667	TTTGAATGCTACCTAGCAGA

(continued)

SEQ ID NO	Sequence
668	TTCTGTTTATAACTACAGCT
669	TCTGTTTATAACTACAGCTT

[0389] The immune cells described herein may be edited using TALEN gene editing.

[0390] "TALEN" or "TALEN gene editing" refers to a transcription activator-like effector nuclease, which is an artificial nuclease used to edit a target gene.

[0391] TALENs are produced artificially by fusing a TAL effector DNA binding domain to a DNA cleavage domain. Transcription activator-like effectors (TALEs) derived from *Xanthomonas* bacteria can be engineered to bind any desired DNA sequence, including a portion of target genes such as TCR subunits, MHC class I complex components, or CD52. By combining an engineered TALE with a DNA cleavage domain, a restriction enzyme can be produced, which is specific to any desired DNA sequence, including a target gene sequence. These can then be introduced into a cell, wherein they can be used for genome editing.

[0392] To produce a TALEN, a TALE protein is fused to a nuclease (N), which is a wild-type or mutated FokI endonuclease. Several mutations to FokI have been made for its use in TALENs; these, for example, improve cleavage specificity or activity.

[0393] The FokI domain functions as a dimer, requiring two constructs with unique DNA binding domains for sites in the target genome with proper orientation and spacing. Both the number of amino acid residues between the TALE DNA binding domain and the FokI cleavage domain and the number of bases between the two individual TALEN binding sites appear to be important parameters for achieving high levels of activity.

[0394] TALENs specific to sequences in a target gene can be constructed using any method known in the art, including various schemes using modular components.

[0395] A target gene may be edited in the immune cells described herein using ZFN gene editing.

[0396] "ZFN" or "Zinc Finger Nuclease" or "ZFN gene editing" refer to a zinc finger nuclease, an artificial nuclease, which can be used to edit a target gene.

[0397] Like a TALEN, a ZFN comprises a FokI nuclease domain (or derivative thereof) fused to a DNA-binding domain. In the case of a ZFN, the DNA-binding domain comprises one or more zinc fingers.

[0398] A zinc finger is a small protein structural motif stabilized by one or more zinc ions. A zinc finger can comprise, for example, Cys2His2, and can recognize an approximately 3-bp sequence. Various zinc fingers of known specificity can be combined to produce multi-finger polypeptides, which recognize about 6, 9, 12, 15 or 18-bp sequences. Various selection and modular assembly techniques are available to generate zinc fingers (and combinations thereof) recognizing specific sequences, including phage display, yeast one-hybrid systems, bacterial one-hybrid and two-hybrid systems, and mammalian cells.

[0399] Like a TALEN, a ZFN must dimerize to cleave DNA. Thus, a pair of ZFNs are required to target non-palindromic DNA sites. The two individual ZFNs must bind opposite strands of the DNA with their nucleases properly spaced apart.

[0400] Also like a TALEN, a ZFN can create a double-stranded break in the DNA, which can create a frame-shift mutation if improperly repaired, leading to a decrease in the expression and amount of a target gene or gene product in a cell. ZFNs can also be used with homologous recombination to mutate in a target gene.

[0401] ZFNs specific to sequences in a target gene can be constructed using any method known in the art.

[0402] The expression and function of one or more MCH-I components may be reduced using RNA interference. "RNAi" or "RNA interference" refers to the process of sequence-specific post-transcriptional gene silencing, mediated by double-stranded RNA (dsRNA). Duplex RNAs such as siRNA (small interfering RNA), miRNA (micro RNA), shRNA (short hairpin RNA), ddRNA (DNA-directed RNA), piRNA (Piwi-interacting RNA), or rasiRNA (repeat associated siRNA) and modified forms thereof are all capable of mediating RNA interference. These dsRNA molecules may be commercially available or may be designed and prepared based on known sequence information. The anti-sense strand of these molecules can include RNA, DNA, PNA, or a combination thereof. DNA/RNA chimeric polynucleotides may include, a double-strand polynucleotide composed of DNA and RNA that inhibits the expression of a target gene. dsRNA molecules can also include one or more modified nucleotides, as described herein, which can be incorporated on either or both strands.

[0403] In RNAi gene silencing or knockdown, dsRNA comprising a first (anti-sense) strand that is complementary to a portion of a target gene and a second (sense) strand that is fully or partially complementary to the first anti-sense strand is introduced into an organism. After introduction into the organism, the target gene-specific dsRNA is processed into relatively small fragments (siRNAs) and can subsequently become distributed throughout the organism, decrease messenger RNA of target gene, leading to a phenotype that may come to closely resemble the phenotype arising from

a complete or partial deletion of the target gene.

[0404] Certain dsRNAs in cells can undergo the action of Dicer enzyme, a ribonuclease III enzyme. Dicer can process the dsRNA into shorter pieces of dsRNA, i.e. siRNAs. RNAi also involves an endonuclease complex known as the RNA induced silencing complex (RISC). Following cleavage by Dicer, siRNAs enter the RISC complex and direct cleavage of a single stranded RNA target having a sequence complementary to the anti-sense strand of the siRNA duplex. The other strand of the siRNA is the passenger strand. Cleavage of the target RNA takes place in the middle of the region complementary to the anti-sense strand of the siRNA duplex. siRNAs can thus down regulate or knock down gene expression by mediating RNA interference in a sequence-specific manner.

[0405] As used herein with respect to RNA interference, "target gene" or "target sequence" refers to a gene or gene sequence whose corresponding RNA is targeted for degradation through the RNAi pathway using dsRNAs or siRNAs as described herein. Exemplary target gene sequences are shown in Table 7. To target a gene, for example using an siRNA, the siRNA comprises an anti-sense region complementary to, or substantially complementary to, at least a portion of the target gene or sequence, and sense strand complementary to the anti-sense strand. Once introduced into a cell, the siRNA directs the RISC complex to cleave an RNA comprising a target sequence, thereby degrading the RNA. The disclosure provides interfering RNAs. The double stranded RNA molecule of the disclosure may be in the form of any type of RNA interference molecule known in the art. In some embodiments, the double stranded RNA molecule is a small interfering RNA (siRNA). In other embodiments, the double stranded RNA molecule is a short hairpin RNA (shRNA) molecule. In other embodiments, the double stranded RNA molecule is a Dicer substrate that is processed in a cell to produce an siRNA. In other embodiments the double stranded RNA molecule is part of a microRNA precursor molecule.

[0406] In some embodiments, the shRNA is a length to be suitable as a Dicer substrate, which can be processed to produce a RISC active siRNA molecule. See, e.g., Rossi et al., US2005/0244858.

[0407] A Dicer substrate double stranded RNA (e.g. a shRNA) can be of a length sufficient that it is processed by Dicer to produce an active siRNA, and may further include one or more of the following properties: (i) the Dicer substrate shRNA can be asymmetric, for example, having a 3' overhang on the anti-sense strand, (ii) the Dicer substrate shRNA can have a modified 3' end on the sense strand to direct orientation of Dicer binding and processing of the dsRNA to an active siRNA, for example the incorporation of one or more DNA nucleotides, and (iii) the first and second strands of the Dicer substrate dsRNA can be from 21-30 bp in length.

[0408] In some embodiments, the interfering RNAs comprise a sequence complementary to a sequence of a B2M mRNA. In some embodiments, the interfering RNA is capable of inducing RNAi-mediated degradation of the B2M mRNA. In some embodiments, the B2M mRNA sequence comprises a coding sequence. In some embodiments, the B2M mRNA sequence comprises an untranslated region.

[0409] In some embodiments, the interfering RNAs comprise a sequence complementary to a sequence of an HLA-A*02 mRNA. In some embodiments, the interfering RNA is capable of inducing RNAi-mediated degradation of the HLA-A*02 mRNA. In some embodiments, the HLA-A*02 mRNA sequence comprises a coding sequence. In some embodiments, the HLA-A*02 mRNA sequence comprises an untranslated region.

[0410] In some embodiments, the interfering RNA is a short hairpin RNA (shRNA). In some embodiments, the shRNA comprises a first sequence, having from 5' to 3' end a sequence complementary to the B2M mRNA; and a second sequence, having from 5' to 3'-end a sequence complementary to the first sequence, wherein the first sequence and second sequence form the shRNA.

[0411] The first sequence may have 18, 19, 20, 21, or 22 nucleotides. The first sequence may be complementary to a sequence selected from the sequences set forth in Tables 10 and 11. The first sequence may have a GC content greater than or equal to 25% and less than 60%. The first sequence may be complementary to a sequence selected from the sequences set forth in Tables 10 and 11. The first sequence may not comprise four nucleotides of the same base or a run of seven C or G nucleotide bases. The first sequence may be 21 nucleotides.

[0412] Illustrative target B2M sequences complementary to the first sequence are shown in Table 10.

[0413] In some cases, the first sequence may have 100% identity, i.e. complete identity, homology, complementarity to the target nucleic acid sequence. In other cases, there may be one or more mismatches between the first sequence and the target nucleic acid sequence. For example, there may be 1, 2, 3, 4, 5, 6, or 7 mismatches between the sense region and the target nucleic acid sequence.

[0414] The sequences set forth in Table 10 are presented as DNA sequences. In all sequences set forth in Table 10, thymine (T) may be replaced by uracil (U) to arrive at the sequence of the target mRNA sequence.

Table 10. Illustrative target B2M sequences complementary to first sequence.

SEQ ID NO	Sequence
670	AGAGAATGGAAAGTCAAATTT
671	ATGGACATGATCTTCTTTATA

(continued)

	SEQ ID NO	Sequence
5	672	TGGACATGATCTTCTTTATAA
	673	GGACATGATCTTCTTTATAAT
	674	TGACAGGATTATTGGAAATTT
	675	TTGTGGTTAATCTGGTTTATT
10	676	TGTGGTTAATCTGGTTTATTT
	677	GCAGAGAATGGAAAGTCAAAT
	678	CAGAGAATGGAAAGTCAAATT
15	679	GAGAATGGAAAGTCAAATTTT
	680	GTCACAGCCCAAGATAGTTAA
	681	TGCTTATACACTTACACTTTA
	682	GCTTATACACTTACACTTTAT
20	683	CTTATACACTTACACTTTATG
	684	ACATGGACATGATCTTCTTTA
	685	CATGGACATGATCTTCTTTAT
25	686	ATCAACATCTTGGTCAGATTT
	687	CTTGCACTCAAAGCTTGTTAA
	688	AGTTAAGCGTGCATAAGTTAA
	689	GCATAAGTTAACTTCCAATTT
30	690	TACATACTCTGCTTAGAATTT
	691	ACATACTCTGCTTAGAATTTG
	692	TTGACAGGATTATTGGAAATT
35	693	GACAGGATTATTGGAAATTTG
	694	TAAGGCATGGTTGTGGTTAAT
	695	GTTGTGGTTAATCTGGTTTAT
	696	GTTCCACAAGTTAAATAAATC
40	697	TCCAGCGTACTCCAAAGATTC
	698	TACTCCAAAGATTCAGGTTTA
	699	ACTCCAAAGATTCAGGTTTAC
45	700	CACGTCATCCAGCAGAGAATG
	701	GGTTTCATCCATCCGACATTG
	702	CCGACATTGAAGTTGACTTAC
	703	TGAAGAATGGAGAGAGAATTG
50	704	GAGCATTGAGACTTGTCTTTC
	705	TTCAGCAAGGACTGGTCTTTC
	706	GCAAGGACTGGTCTTCTATC
55	707	CGTGTGAACCATGTGACTTTG
	708	CTTTGTCACAGCCCAAGATAG
	709	TCACAGCCCAAGATAGTTAAG

(continued)

	SEQ ID NO	Sequence
5	710	AGTGGGATCGAGACATGTAAG
	711	AGGTTTGAAGATGCCGCATTT
	712	GGTTTGAAGATGCCGCATTTG
	713	TTGATATGCTTATACACTTAC
10	714	TGAGTGCTGTCTCCATGTTTG
	715	TGTCTCCATGTTTGATGTATC
	716	TCAACATCTTGGTCAGATTTG
15	717	TCAGATTTGAACCTCTTCAATC
	718	TTCAATCTCTTGCACTCAAAG
	719	TTGCACTCAAAGCTTGTTAAG
	720	ACTCAAAGCTTGTTAAGATAG
20	721	AGATAGTTAAGCGTGCATAAG
	722	TGCATAAGTTAACTTCCAATT
	723	GTAACTTCCAATTTACATAC
25	724	ATTGACAGGATTATTGGAAAT
	725	GTAAGGCATGGTTGTGGTTAA
	726	GGTTGTGGTTAATCTGGTTTA
	727	TTCTGAAGCTGACAGCATTG
30	728	GCTATCCAGCGTACTCCAAAG
	729	CATCCAGCAGAGAATGGAAAG
	730	CAAATTCCTGAATTGCTATG
35	731	ATTGCTATGTGTCTGGGTTTC
	732	GAAGATGCCGCATTTGGATTG
	733	CAATTTACATACTCTGCTTAG
	734	TATCCAGCGTACTCCAAAGAT
40	735	ATCCAGCGTACTCCAAAGATT
	736	CTCCAAAGATTGAGTTTACT
	737	TGCTATGTGTCTGGGTTTCAT
45	738	TTTCATCCATCCGACATTGAA
	739	GAAGTTGACTTACTGAAGAAT
	740	GAAGAATGGAGAGAGAATTGA
	741	AGAATGGAGAGAGAATTGAAA
50	742	CAGCAAGGACTGGTCTTTCTA
	743	AGCAAGGACTGGTCTTTCTAT
	744	ACTTTGTACAGCCCAAGATA
55	745	TTGTCACAGCCCAAGATAGTT
	746	TGTCACAGCCCAAGATAGTTA
	747	CACAGCCCAAGATAGTTAAGT

(continued)

5

10

15

20

25

30

35

40

45

50

55

SEQ ID NO	Sequence
748	GCAGCATCATGGAGGTTTGAA
749	CCGCATTTGGATTGGATGAAT
750	TTGAGTGCTGTCTCCATGTTT
751	AGTGCTGTCTCCATGTTTGAT
752	CTGTCTCCATGTTTGATGTAT
753	TCTAGGAGGGCTGGCAACTTA
754	CAACATCTTGGTCAGATTTGA
755	GTCAGATTTGAACTCTTCAAT
756	TCTTGCACTCAAAGCTTGTTA
757	TGCACTCAAAGCTTGTTAAGA
758	GCACTCAAAGCTTGTTAAGAT
759	CACTCAAAGCTTGTTAAGATA
760	TCAAAGCTTGTTAAGATAGTT
761	CAAAGCTTGTTAAGATAGTTA
762	GATAGTTAAGCGTGCATAAGT
763	ATAGTTAAGCGTGCATAAGTT
764	TAGTTAAGCGTGCATAAGTTA
765	TTAAGCGTGCATAAGTTAACT
766	TAAGCGTGCATAAGTTAACTT
767	ATTTACATACTCTGCTTAGAA
768	TTTACATACTCTGCTTAGAAT
769	ACAGGATTATTGGAAATTTGT
770	CAGGATTATTGGAAATTTGTT
771	AGGCATGGTTGTGGTTAATCT
772	CAGCAGAGAATGGAAAGTCAA
773	TCCGACATTGAAGTTGACTTA
774	CTGGTCTTTCTATCTCTTGTA
775	CCGTGTGAACCATGTGACTTT
776	CCCAAGATAGTTAAGTGGGAT
777	GGTTGCTCCACAGGTAGCTCT
778	GCTCCACAGGTAGCTCTAGGA
779	GGGAGCAGAGAATTCTCTTAT
780	GGAGCAGAGAATTCTCTTATC
781	GAGCAGAGAATTCTCTTATCC
782	GAGAATTCTCTTATCCAACAT
783	GAATTCTCTTATCCAACATCA
784	AAGTGGAGCATTGAGACTTGT
785	AAGGACTGGTCTTTCTATCTC

(continued)

SEQ ID NO	Sequence
786	AAGCTTGTTAAGATAGTTAAG
787	AAGCGTGCATAAGTTAACTTC
788	AAGATGCCGCATTTGGATTGG
789	AAGAATGGAGAGAGAATTGAA
790	AACATCAACATCTTGGTCAGA
791	AAGGCATGGTTGTGGTTAATC
792	AAGCAGCATCATGGAGGTTTG
793	AAGATGAGTATGCCTGCCGTG
794	AAGTTGACTTACTGAAGAATG
795	AAGATAGTTAAGCGTGCATAA
796	AACTTCCAATTTACATACTCT
797	AACATCTTGGTCAGATTTGAA
798	AACTCTCAATCTCTTGCACT
799	AATTCCTGAATTGCTATGTG
800	AATGGAAAGTCAAATTCCTG
801	AACCATGTGACTTTGTCACAG
802	AATTGACAGGATTATTGGAAA
803	AATTCTTATCCAACATCAA
804	AAAGTGGAGCATTGAGCTTG
805	AAAGTCAAATTCCTGAATTG
806	GTTGCTCCACAGGTAGCTCTA
807	AATTTACATACTCTGCTTAGA

[0415] In some embodiments, the interfering RNAs comprise a sequence complementary to a sequence of an HLA-A*02 mRNA. In some embodiments, the interfering RNA is capable of inducing RNAi-mediated degradation of the HLA-A*02 mRNA. In some embodiments, the HLA-A*02 mRNA sequence comprises a coding sequence. In some embodiments, the HLA-A*02 mRNA sequence comprises an untranslated region.

[0416] In one embodiment of the immune cell of the present invention, a sequence encoding a B2M shRNA comprises a sequence of GCACTCAAAGCTTGTTAAGATCGAAATCTTAACAAGCTTTGAGTGC (SEQ ID NO: 179), or a sequence having at least 90%, at least 95%, at least 97% or at least 99% identity thereto. A further exemplary sequence encoding a B2M shRNA comprises a sequence of GTTAACTTCCAATTTACATACCGAAGTATGTAAATTGGAAGTTAAC (SEQ ID NO: 808), or a sequence having at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0417] In one embodiment of the immune cell of the present invention, the interfering RNA is a short hairpin RNA (shRNA). In one embodiment of the immune cell of the present invention, the shRNA comprises a first sequence, having from 5'- to 3'-end a sequence complementary to the HLA-A*02 mRNA; and a second sequence, having from 5'- to 3'-end a sequence complementary to the first sequence, wherein the first sequence and second sequence form the shRNA.

[0418] Illustrative target HLA sequences complementary to the first sequence are shown in Table 11.

Table 11. Illustrative target HLA sequences complementary to first sequence

SEQ ID NO	Sequence
809	CTTCTTCCTTCCCTATTAAAA
810	TCTCACTCCATGAGGTATTTT
811	CTCTCACTCCATGAGGTATTT

(continued)

	SEQ ID NO	Sequence
5	812	GAGGAGGAAGAGCTCAGATAG
	813	GCTCTCACTCCATGAGGTATT
	814	AGGATTACATCGCCCTGAAAG
	815	ACACCGTCCAGAGGATGTATG
10	816	AGGGTCCTTCTTCCTGGATAC
	817	CCTACGACGGCAAGGATTACA
	818	TCACTCCATGAGGTATTTCTT
15	819	CTACGACGGCAAGGATTACAT
	820	CTCACTCCATGAGGTATTTCT
	821	GGAGGAAGAGCTCAGATAGAA
	822	CACACCGTCCAGAGGATGTAT
20	823	CACGCTGTCTCTGACCATGAA
	824	CTGGACAGGAGCAGAGATACA
	825	TGGAGGAGGAAGAGCTCAGAT
25	826	GGCTCTCACTCCATGAGGTAT
	827	CATCTCTGTCTCAACTTCATG
	828	TACGACGGCAAGGATTACATC
	829	GGATTACATCGCCCTGAAAGA
30	830	GATTACATCGCCCTGAAAGAG
	831	CTCAGACCACCAAGCACAAAGT
	832	TCACACCGTCCAGAGGATGTA
35	833	ACTCCATGAGGTATTTCTTCA
	834	CACTCCATGAGGTATTTCTTC
	835	CCATGAGGTATTTCTTCACAT
	836	ACTTCTTCCTTCCCTATTAAA
40	837	GTGTCTCTCACAGCTTGTAAG
	838	CTGTGTTTCGTGTAGGCATAAT
	839	TGTGTTTCGTGTAGGCATAATG
45	840	TAACTTCTTCCTTCCCTATTAA
	841	TCTGGACAGGAGCAGAGATAC
	842	TTGCTGGCCTGGTTCTCTTTG
	843	TGTCTCTCACAGCTTGTAAG
50	844	ACTTGAAGAACCCTGACTTTTG
	845	GAAGAACCCTGACTTTGTTTC
	846	TCTGTGTTTCGTGTAGGCATAA
55	847	CATGGTGCACTGAGCTGTAAC
	848	GTAACCTTCTTCCTTCCCTATT
	849	CATGTGCAGCATGAGGGTTTG

(continued)

	SEQ ID NO	Sequence
5	850	TTGTTCCCTG CCCTTCCCTTTG
	851	ACCCAGTTCTCACTCCCATTG
	852	GGGTTTCCAGAGAAGCCAATC
	853	TTCTCCCTCTCCCAACCTATG
10	854	GTCTCTCACAGCTTGTAAGT
	855	TGTGTCTCTCACAGCTTGTA
	856	GAGGAAGAGCTCAGATAGAAA
15	857	TGAAGAACCCTGACTTTGTTT
	858	TTGAAGAACCCTGACTTTGTT
	859	GTGTTTCGTGTAGGCATAATGT
	860	TGGTGCACTGAGCTGTAACCTT
20	861	CTCCCTCTCCCAACCTATGTA
	862	AGGAGGAAGAGCTCAGATAGA
	863	ACCTATGTAGGGTCCTTCTTC
25	864	GGGTCCTTCTTCCTGGATACT
	865	GGTCCTTCTTCCTGGATACTC
	866	GTCCTTCTTCCTGGATACTCA
	867	AAGCCAATCAGTGTCTGTCGCG
30	868	AAGAGGACCTGCGCTCTTGGA
	869	AAGTGTGAGACAGCTGCCTTG
	870	AAG GCACCTG CATGTGTCTGT
35	871	AATCATCTTTCCTGTTCCAGA
	872	AAAG G CACCTGCATGTGTCTG
	873	AAAGAGGACCTGCGCTCTTG
	874	AAACGCATATGACTCACCACG
40	875	GGAAGAGCTCAGATAGAAA
	876	GGGAGACACGGAAAGTGAA
	877	CACCTGCCATGTGCAGCATGA
45	878	GGAGATCACACTGACCTGGCA
	879	GGATTACATCGCCCTGAAAG
	880	GCAGGAGGGTCCGGAGTATT
	881	GGACGGGGAGACACGGAAAG
50	882	GAAAGTGAAGGCCCACTCA
	883	GATACCTGGAGAACGGGAAG
	884	GCTGTGGTGGTGCCTTCTGG
55	885	GCTACTACAACCAGAGCGAG
	886	GTG G CTCCG CAGATACCTG
	887	GCCAATCAGTGTCTGTCGCG

(continued)

SEQ ID NO	Sequence
888	GAGGACCTGCGCTCTTGA
889	GTGTGAGACAG CTG CCTTG
890	GGCACCTGCATGTGTCTGT
891	TCATCTTTCCTGTTCCAGA
892	AGGCACCTGCATGTGTCTG
893	AGAGGACCTGCGCTCTTGG
894	ACGCATATGACTCACCACG

[0419] In some embodiments, the first sequence and second sequence are separated by a linker, sometimes referred to as a loop. In some embodiments, both the first sequence and the second sequence are encoded by one single-stranded RNA or DNA vector. In some embodiments, the loop is between the first and second sequences. In these embodiments, the first sequence and the second sequence hybridize to form a duplex region. The first sequence and second sequence are joined by a linker sequence, forming a "hairpin" or "stem-loop" structure. The shRNA can have complementary first sequences and second sequences at opposing ends of a single stranded molecule, so that the molecule can form a duplex region with the complementary sequence portions, and the strands are linked at one end of the duplex region by a linker (*i.e.* loop sequence). The linker, or loop sequence, can be either a nucleotide or non-nucleotide linker. The linker can interact with the first sequence, and optionally, second sequence through covalent bonds or noncovalent interactions.

[0420] Any suitable nucleotide loop sequence is envisaged as within the scope of the disclosure. An shRNA of this disclosure may include a nucleotide, non-nucleotide, or mixed nucleotide/non-nucleotide linker that joins the first sequence of the shRNA to the second sequence of the shRNA. A nucleotide loop sequence can be ≥ 2 nucleotides in length, for example about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 nucleotides in length. Illustrative loop sequences are disclosed in Table 12.

[0421] In some embodiments, the shRNA further comprises a 5'-flank sequence and a 3'-flank sequence. In some embodiments, wherein the 5'-flank sequence is joined to the 5'-end of the first sequence, and wherein the 3'-flank sequence is joined to the 3'-end of the second sequence.

[0422] Without wishing to be bound by theory, it is thought that flanking shRNA stem loop sequence with 5'- and 3'-sequences similar to those found in microRNAs can target the shRNA for processing by the endogenous microRNA processing machinery, increasing the effectiveness of shRNA processing. Alternatively, or in addition, flanking sequences may increase shRNA compatibility with polymerase II or polymerase III promoters, leading to more effective regulation of shRNA expression.

[0423] The 5'-flank sequence may be selected from the sequences set forth in Table 12. Illustrative flank sequence are shown in Table 12.

Table 12. Illustrative flank sequences

SEQ ID NO	5' Flank Sequence
895	GG
896	ACACCAUGUUGCCAGUCUCUAGG
897	UGAUAGCAAUGUCAGCAGUGCCU
898	UAUUGCUGUUGACAGUGAGCGAC
SEQ ID NO	3' Flank Sequence
899	UGGCGUCUGGCCCAACCACAC
900	GUAAGGUUGACCAUACUCUAC

[0424] The first and second sequence may be present on a single stranded polynucleotide, wherein the first sequence and second sequence may be separated by 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 nucleotides, wherein the 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 nucleotides form a loop region in the shRNA. The loop region may comprise a sequence

selected from the sequences set forth in Table 13

Table 13. Illustrative loop region sequences

SEQ ID NO	Loop Region Sequence
901	CGAA
902	UUCAAGA
903	AUAUUCA
904	UGUGCUGUC
905	CUCGAG
906	CUUCCUGUCAGA
907	CUUCCCUUGUCAGA
908	GUGUUAUUCUUG
909	GUGUCUAAUUG
910	GUGUUAGUCUUG
911	UCAAGAG
912	GGACAUCCAGGG
913	GUGAAGCCACAGAUG
914	GAUUCUAAAA

[0425] shRNAs of the disclosure may be generated exogenously by chemical synthesis, by in vitro transcription, or by cleavage of longer double-stranded RNA with Dicer or another appropriate nuclease with similar activity. Chemically synthesized siRNAs, produced from protected ribonucleoside phosphoramidites using a conventional DNA/RNA synthesizer, may be obtained from commercial suppliers such as Millipore Sigma (Houston, Tex.), Ambion Inc. (Austin, Tex.), Invitrogen (Carlsbad, Calif.), or Dharmacon (Lafayette, Colo.). siRNAs can be purified by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof, for example. Alternatively, siRNAs may be used with little if any purification to avoid losses due to sample processing.

[0426] The shRNAs of the disclosure can be produced using an expression vector into which a nucleic acid encoding the double stranded RNA has been cloned, for example under control of a suitable promoter.

Pharmaceutical Compositions

[0427] The disclosure provides pharmaceutical compositions comprising immune cells comprising the first and second receptors of the disclosure and a pharmaceutically acceptable diluent, carrier or excipient.

[0428] Such compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; and preservatives.

[0429] The immune cell may express both the first receptor and the second receptor. At least about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% of the immune cells may express both the first receptor and the second receptor. At least 90% of the immune cells may express both the first receptor and the second receptor.

Treating Cancer

[0430] Provided herein are methods of killing a plurality of cancer cells, or treating cancer, in a subject, comprising administering to the subject a therapeutically effective amount of a composition comprising immune cells comprising the first and second receptors of the disclosure. The immune cells express both receptors in the same cell.

[0431] Cancer is a disease in which abnormal cells divide without control and spread to nearby tissue. The cancer may comprise a liquid tumor or a solid tumor. Cancers can arise in virtually any organ in the body, including epithelial tissues. For example, EGFR positive cancers that can be treated using the methods described herein (not according to the present invention) include lung cancer, small cell lung cancer, non-small cell lung cancer, pancreatic ductal carcinoma,

colorectal cancer, head and neck cancer, esophagus and gastric adenocarcinoma, ovarian cancer, glioblastoma multiforme, cervical squamous cell carcinoma, kidney cancer, papillary kidney cancer, kidney renal clear cell carcinoma, bladder cancer, breast cancer, bile duct cancer, liver cancer, prostate cancer, sarcoma, thyroid cancer, thymus cancer, stomach cancer, or uterine cancer, and all-other EGFR target expressing tumors.

[0432] In some embodiments, the plurality of cancer cells express the target antigen. In some embodiments, the plurality of cancer cells of the subject express EGFR. EGFR positive cancers include, lung cancer, small cell lung cancer, non-small cell lung cancer, pancreatic ductal carcinoma, colorectal cancer, head and neck cancer, esophagus and gastric adenocarcinoma, ovarian cancer, glioblastoma multiforme, cervical squamous cell carcinoma, kidney cancer, papillary kidney cancer, kidney renal clear cell carcinoma, bladder cancer, breast cancer, bile duct cancer, liver cancer, prostate cancer, sarcoma, thyroid cancer, thymus cancer, stomach cancer, or uterine cancer.

[0433] As described herein, a plurality of cancer cells do not express a polymorphic allele of COLEC12, APCDD1 or CXCL16. As described herein, a plurality of cancer cells do not express a non-target antigen. As described herein, a plurality of cancer cells do not express HLA-A*02. As described herein, the plurality of cancer cells do not express an allelic variant of HLA-A*02. As described herein, the plurality of cancer cells do not express HLA-A*01. As also described herein, the plurality of cancer cells do not express HLA-A*03. As also described herein, the plurality of cancer cells do not express HLA-A*11. As also described herein, the plurality of cancer cells do not express HLA-B*07. As also described herein, the cancer cells have lost an allele of COLEC12, APCDD1 or CXCL16, or HLA-A*02 through loss of heterozygosity at that locus.

[0434] Administration of the immune cells or pharmaceutical compositions described herein can reduce the size of a tumor in the subject. The size of the tumor may be reduced by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, or about 100%, relative to the size of the tumor before administration of the immune cells or pharmaceutical compositions. The tumor may be eliminated.

[0435] Administration of the immune cells or pharmaceutical compositions described herein can arrest the growth of a tumor in the subject. For example, the immune cells or pharmaceutical compositions can kill tumor cells, so that the tumor stops growing, or is reduced in size. In some cases, immune cells or pharmaceutical compositions can prevent formation of additional tumors, or reduce the total number of tumors in the subject.

[0436] Administration of the immune cells or pharmaceutical compositions described herein can result in selective killing of a cancer cell, but not a normal cell in the subject. About 60% of the cells killed may be cancer cells, about 65% of the cells killed may be cancer cells, about 70% of the cells killed may be cancer cells, about 75% of the cells killed may be cancer cells, about 80% of the cells killed may be cancer cells, about 85% of the cells killed may be cancer cells, about 90% of the cells killed may be cancer cells, about 95% of the cells killed may be cancer cells, or about 100% of the cells killed may be cancer cells.

[0437] Administration of the immune cells or pharmaceutical compositions described herein can results in the killing of about 40%, about 50%, about 60%, about 70%, about 80%, about 90% or all of the cancer cells of the subject.

[0438] Administration of the immune cells or pharmaceutical compositions described herein can result in fewer side effects for the subject than administration of an otherwise equivalent immune cell comprising the first activator receptor, but no second inhibitory receptor. For example, administering the immune cells or pharmaceutical compositions described herein can reduce dose limited toxicity relative to the EGFR CAR, or EGFR TCR administered without the second inhibitory receptor.

[0439] The disclosure provides methods of treating a cancer in a subject comprising: (a) determining the genotype of normal cells and a plurality of cancer cells of the subject at a polymorphic locus selected from the group consisting of a polymorphic locus of COLEC12, a polymorphic locus of APCDD1 and a polymorphic locus of CXCL16, or the HLA-A*02 locus; (b) determining the expression of EGFR in a plurality of cancer cells; and (c) administering a plurality of immune cells to the subject if the normal cells are heterozygous for the polymorphic locus and the plurality of cancer cells are hemizygous for the polymorphic locus or have lost HLA-A* 02, and the plurality of cancer cells are EGFR positive, wherein the plurality of immune cells comprise: (i) a first receptor, optionally a chimeric antigen receptor (CAR) or T cell receptor (TCR), comprising an extracellular ligand binding domain specific to EGFR, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); and (ii) a second receptor, optionally an inhibitory chimeric antigen receptor (*i.e.* inhibitory receptors), comprising an extracellular ligand binding specific to a non-target antigen selected from COLEC12, APCDD1, and CXCL16, or an antigen peptide thereof in a complex with an a major histocompatibility complex class I (MHC-I), or HLA-A*02, wherein the non-target antigen comprises a polymorphism.

[0440] Methods of genotyping cancer cells and normal cells from a subject for the presence or absence of SNPs will be readily apparent to persons of ordinary skill in the art. SNP genotyping methods include, *inter alia*, PCR based methods such as dual-probe TaqMan assays, array based hybridization methods and sequencing.

[0441] Methods of measuring the expression of the target antigen in cancer or normal cells from a subject will be readily apparent to persons of ordinary skill in the art. These include, *inter alia*, methods of measuring RNA expression such as RNA sequencing and reverse transcription polymerase chain reaction (RT-PCR), as well as methods of meas-

uring protein expression such as immunohistochemistry based methods. Methods of measuring loss of heterozygosity in a plurality of cancer cells, include, *inter alia*, high throughput sequencing of genomic DNA extracted from cancer cells using methods known in the art.

[0442] Methods of measuring the expression of the target antigen in cancer or normal cells from a subject will be readily apparent to persons of ordinary skill in the art. These include, *inter alia*, methods of measuring RNA expression such as RNA sequencing and reverse transcription polymerase chain reaction (RT-PCR), as well as methods of measuring protein expression such as immunohistochemistry based methods.

[0443] As described herein, the immune cells are T cells.

[0444] In some embodiments, the immune cells are allogeneic or autologous.

[0445] In some embodiments, the second receptor increases the specificity of the immune cells for the EGFR positive cancer cells compared to immune cells that express the first receptor but do not express the second receptor. In some embodiments, the immune cells have reduced side effects compared to immune cells that express the first receptor but do not express the second receptor.

[0446] Treating cancer can result in a reduction in size of a tumor. A reduction in size of a tumor may also be referred to as "tumor regression". After treatment, tumor size may be reduced by 5% or greater relative to its size prior to treatment; the tumor size may be reduced by 10% or greater; reduced by 20% or greater; reduced by 30% or greater; reduced by 40% or greater; reduced by 50% or greater; or reduced by greater than 75% or greater. Size of a tumor may be measured by any reproducible means of measurement. The size of a tumor may be measured as a diameter of the tumor.

[0447] Treating cancer can result in a reduction in tumor volume. After treatment, tumor volume may be reduced by 5% or greater relative to its size prior to treatment; the tumor volume may be reduced by 10% or greater; reduced by 20% or greater; reduced by 30% or greater; reduced by 40% or greater; reduced by 50% or greater; or reduced by greater than 75% or greater. Tumor volume may be measured by any reproducible means of measurement.

[0448] Treating cancer results in a decrease in number of tumors. After treatment, tumor number may be reduced by 5% or greater relative to number prior to treatment; tumor number may be reduced by 10% or greater; reduced by 20% or greater; reduced by 30% or greater; reduced by 40% or greater; reduced by 50% or greater; or reduced by greater than 75%. Number of tumors may be measured by any reproducible means of measurement. The number of tumors may be measured by counting tumors visible to the naked eye or at a specified magnification. The specified magnification may be 2x, 3x, 4x, 5x, 10x, or 50x.

[0449] Treating cancer can result in a decrease in number of metastatic lesions in other tissues or organs distant from the primary tumor site. After treatment, the number of metastatic lesions may be reduced by 5% or greater relative to number prior to treatment; the number of metastatic lesions may be reduced by 10% or greater; reduced by 20% or greater; reduced by 30% or greater; reduced by 40% or greater; reduced by 50% or greater; or by greater than 75%. The number of metastatic lesions may be measured by any reproducible means of measurement. The number of metastatic lesions may be measured by counting metastatic lesions visible to the naked eye or at a specified magnification. The specified magnification may be 2x, 3x, 4x, 5x, 10x, or 50x.

[0450] Treating cancer can result in an increase in average survival time of a population of treated subjects in comparison to a population receiving carrier alone. The average survival time may be increased by more than 30 days; by more than 60 days; by more than 90 days; or by more than 120 days. An increase in average survival time of a population may be measured by any reproducible means. An increase in average survival time of a population may be measured, for example, by calculating for a population the average length of survival following initiation of treatment with an active compound. An increase in average survival time of a population may also be measured, for example, by calculating for a population the average length of survival following completion of a first round of treatment with an active compound.

[0451] Treating cancer can result in an increase in average survival time of a population of treated subjects in comparison to a population of untreated subjects. The average survival time may be increased by more than 30 days; by more than 60 days; by more than 90 days; or by more than 120 days. An increase in average survival time of a population may be measured by any reproducible means. An increase in average survival time of a population may be measured, for example, by calculating for a population the average length of survival following initiation of treatment with an active compound. An increase in average survival time of a population may also be measured, for example, by calculating for a population the average length of survival following completion of a first round of treatment with an active compound.

[0452] Treating cancer can result in increase in average survival time of a population of treated subjects in comparison to a population receiving monotherapy with a drug that is not a compound of the present disclosure, or a pharmaceutically acceptable salt, prodrug, metabolite, analog or derivative thereof. The average survival time may be increased by more than 30 days; by more than 60 days; by more than 90 days; or by more than 120 days. An increase in average survival time of a population may be measured by any reproducible means. An increase in average survival time of a population may be measured, for example, by calculating for a population the average length of survival following initiation of treatment with an active compound. An increase in average survival time of a population may also be measured, for example, by calculating for a population the average length of survival following completion of a first round of treatment with an active compound.

[0453] Treating cancer can result in a decrease in the mortality rate of a population of treated subjects in comparison to a population receiving carrier alone. Treating cancer can result in a decrease in the mortality rate of a population of treated subjects in comparison to an untreated population. Treating cancer can result in a decrease in the mortality rate of a population of treated subjects in comparison to a population receiving monotherapy with a drug that is not a compound of the present disclosure, or a pharmaceutically acceptable salt, prodrug, metabolite, analog or derivative thereof. As described herein, the mortality rate may be decreased by more than 2%; by more than 5%; by more than 10%; or by more than 25%. A decrease in the mortality rate of a population of treated subjects may be measured by any reproducible means. A decrease in the mortality rate of a population may be measured, for example, by calculating for a population the average number of disease-related deaths per unit time following initiation of treatment with an active compound. A decrease in the mortality rate of a population may also be measured, for example, by calculating for a population the average number of disease-related deaths per unit time following completion of a first round of treatment with an active compound.

[0454] Treating cancer can result in a decrease in tumor growth rate. After treatment, tumor growth rate may be reduced by at least 5% relative to the number prior to treatment; tumor growth rate may be reduced by at least 10%; reduced by at least 20%; reduced by at least 30%; reduced by at least 40%; reduced by at least 50%; reduced by at least 50%; or reduced by at least 75%. Tumor growth rate may be measured by any reproducible means of measurement. Tumor growth rate can be measured according to a change in tumor diameter per unit time.

[0455] Treating cancer can result in a decrease in tumor regrowth. After treatment, tumor regrowth may be less than 5%; tumor regrowth may be less than 10%; less than 20%; less than 30%; less than 40%; less than 50%; less than 50%; or less than 75%. Tumor regrowth may be measured by any reproducible means of measurement. Tumor regrowth may be measured, for example, by measuring an increase in the diameter of a tumor after a prior tumor shrinkage that followed treatment. A decrease in tumor regrowth is indicated by failure of tumors to reoccur after treatment has stopped.

[0456] Treating or preventing a cancer can result in a reduction in the rate of cellular proliferation. After treatment, the rate of cellular proliferation may be reduced by at least 5%; by at least 10%; by at least 20%; by at least 30%; by at least 40%; by at least 50%; by at least 50%; or by at least 75%. The rate of cellular proliferation may be measured by any reproducible means of measurement. The rate of cellular proliferation is measured, for example, by measuring the number of dividing cells in a tissue sample per unit time.

[0457] Treating or preventing cancer can result in a reduction in the proportion of proliferating cells. After treatment, the proportion of proliferating cells may be reduced by at least 5%; by at least 10%; by at least 20%; by at least 30%; by at least 40%; by at least 50%; by at least 50%; or by at least 75%. The proportion of proliferating cells may be measured by any reproducible means of measurement. The proportion of proliferating cells may be measured, for example, by quantifying the number of dividing cells relative to the number of nondividing cells in a tissue sample. The proportion of proliferating cells can be equivalent to the mitotic index.

[0458] Treating or preventing cancer can result in a decrease in size of an area or zone of cellular proliferation. After treatment, size of an area or zone of cellular proliferation may be reduced by at least 5% relative to its size prior to treatment; reduced by at least 10%; reduced by at least 20%; reduced by at least 30%; reduced by at least 40%; reduced by at least 50%; reduced by at least 50%; or reduced by at least 75%. Size of an area or zone of cellular proliferation may be measured by any reproducible means of measurement. The size of an area or zone of cellular proliferation may be measured as a diameter or width of an area or zone of cellular proliferation.

[0459] Treating or preventing cancer can result in a decrease in the number or proportion of cells having an abnormal appearance or morphology. After treatment, the number of cells having an abnormal morphology may be reduced by at least 5% relative to its size prior to treatment; reduced by at least 10%; reduced by at least 20%; reduced by at least 30%; reduced by at least 40%; reduced by at least 50%; reduced by at least 50%; or reduced by at least 75%. An abnormal cellular appearance or morphology may be measured by any reproducible means of measurement. An abnormal cellular morphology can be measured by microscopy, e.g., using an inverted tissue culture microscope. An abnormal cellular morphology can take the form of nuclear pleiomorphism.

Dosage and Administration

[0460] The immune cells and of the present disclosure may be administered in a number of ways depending upon whether local or systemic treatment is desired.

[0461] In general, administration may be parenteral.

[0462] Methods for administration of cells for adoptive cell therapy are known and may be used in connection with the provided methods and compositions. For example, adoptive T cell therapy methods are described, e.g., in US Patent Application Publication No. 2003/0170238 to Gruenberg et al and U.S. Pat. No. 4,690,915 to Rosenberg.

[0463] The compositions of the disclosure are suitable for parenteral administration. As used herein, "parenteral administration" of a pharmaceutical composition includes any route of administration characterized by physical breaching of a tissue of a subject and administration of the pharmaceutical composition through the breach in the tissue, thus

generally resulting in the direct administration into the blood stream, into muscle, or into an internal organ. Parenteral administration thus includes administration of a pharmaceutical composition by injection of the composition, by application of the composition through a surgical incision, by application of the composition through a tissue-penetrating non-surgical wound. In particular, parenteral administration is contemplated to include subcutaneous, intraperitoneal, intramuscular, intrasternal, intravenous, intraarterial, intrathecal, intraventricular, intraurethral, intracranial, intratumoral, intrasynovial injection or infusions; and kidney dialytic infusion techniques. As described herein, parenteral administration of the compositions of the present disclosure comprises intravenous or intraarterial administration.

[0464] The disclosure provides pharmaceutical compositions comprising a plurality of immune cells of the disclosure, and a pharmaceutically acceptable carrier, diluent or excipient.

[0465] Formulations of a pharmaceutical composition suitable for parenteral administration typically generally comprise of immune cells combined with a pharmaceutically acceptable carrier, such as sterile water or sterile isotonic saline. Such formulations may be prepared, packaged, or sold in a form suitable for bolus administration or for continuous administration. Injectable formulations may be prepared, packaged, or sold in unit dosage form, such as in ampoules or in multi-dose containers containing a preservative. Formulations for parenteral administration include suspensions, solutions, emulsions in oily or aqueous vehicles, and pastes. Such formulations may further comprise one or more additional ingredients including suspending, stabilizing, or dispersing agents. Parenteral formulations also include aqueous solutions, which may contain excipients such as salts, carbohydrates and buffering agents. Exemplary parenteral administration forms include solutions or suspensions in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired. Formulations for parenteral administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

[0466] As described herein, the formulated composition comprising the immune cells is suitable for administration via injection. As also described herein, the formulated composition comprising the immune cells is suitable for administration via infusion.

[0467] The pharmaceutical compositions of the present disclosure, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the immune cells with the pharmaceutical carrier(s) or excipient(s), such as liquid carriers.

[0468] Aqueous suspensions may further contain substances that increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0469] The compositions of the present disclosure may additionally contain other adjunct components conventionally found in pharmaceutical compositions. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present disclosure, such as dyes, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the immune cells of the compositions of the present disclosure.

[0470] The formulation or composition may also contain more than one active ingredient useful for the particular indication, disease, or condition being treated with the immune cells, where the respective activities do not adversely affect one another. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended. Thus, in some embodiments, the pharmaceutical composition further includes other pharmaceutically active agents or drugs, such as chemotherapeutic agents.

[0471] The pharmaceutical composition in some aspects can employ time-released, delayed release, and sustained release delivery systems such that the delivery of the composition occurs prior to, and with sufficient time to cause, sensitization of the site to be treated. Many types of release delivery systems are available and known. Such systems can avoid repeated administrations of the composition, thereby increasing convenience to the subject and the physician.

[0472] Administration can be effected in one dose, continuously or intermittently throughout the course of treatment. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician.

[0473] The pharmaceutical composition in some embodiments contains the immune cells in amounts effective to treat or prevent a cancer, such as a therapeutically effective or prophylactically effective amount. Therapeutic or prophylactic efficacy in some embodiments is monitored by periodic assessment of treated subjects. For repeated administrations over days, weeks or months, depending on the condition, the treatment can be repeated until a desired suppression of cancer signs or symptoms occurs. However, other dosage regimens may be useful and can be determined. The desired dosage can be delivered by a single bolus administration or infusion of the composition or by multiple bolus administrations or infusions of the composition.

[0474] As described herein, the cells or population of cells can be administered in one or more doses. As described herein, an effective amount of cells can be administered as a single dose. As described herein, an effective amount of

cells can be administrated as more than one doses over a period time. Timing of administration is within the judgment of a managing physician and depends on the clinical condition of the patient.

[0475] The cells or population of cells may be obtained from any source, such as a blood bank or a donor, or the patient themselves.

[0476] An effective amount means an amount which provides a therapeutic or prophylactic benefit. The dosage administered will be dependent upon the age, health and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment and the nature of the effect desired. As described herein, an effective amount of cells or composition comprising those cells are administrated parenterally. As described herein, administration can be an intravenous administration. As described herein, administration can be directly done by injection within a tumor.

[0477] For purposes of the disclosure, an assay, which comprises, for example, comparing the extent to which target cells are lysed or one or more cytokines are secreted by immune cells expressing the receptors, upon administration of a given dose of such immune cells to a mammal, among a set of mammals of which is each given a different dose of the immune cells, can be used to determine a starting dose to be administered to a mammal.

[0478] As described herein, the cells are administered as part of a combination treatment, such as simultaneously with or sequentially with, in any order, another therapeutic intervention, such as an antibody or engineered cell or receptor or agent, such as a cytotoxic or therapeutic agent. The immune cells of the disclosure are co-administered with one or more additional therapeutic agents or in connection with another therapeutic intervention, either simultaneously or sequentially in any order. In some contexts, the immune cells are co-administered with another therapy sufficiently close in time such that the immune cell populations enhance the effect of one or more additional therapeutic agents, or vice versa. As described herein, the immune cells are administered prior to the one or more additional therapeutic agents. As described herein, the immune cells are administered after the one or more additional therapeutic agents.

[0479] As described herein, a lympho-depleting chemotherapy is administered to the subject prior to, concurrently with, or after administration (e.g., infusion) of adoptive immune cells. In an example, the lympho-depleting chemotherapy is administered to the subject prior to administration of the immune cells. For example, the lympho-depleting chemotherapy ends 1-4 days (e.g., 1, 2, 3, or 4 days) prior to adoptive cell infusion. As described herein, multiple doses of adoptive cells are administered. As described herein, a lympho-depleting chemotherapy is administered to the subject prior to, concurrently with, or after administration (e.g., infusion) of the immune cells described herein. Examples of lympho-depletion include non-myeloablative lympho-depleting chemotherapy, myeloablative lympho-depleting chemotherapy, total body irradiation, etc. Examples of lympho-depleting agents include antithymocyte globulin, anti-CD3 antibodies, anti-CD4 antibodies, anti-CD8 antibodies, anti-CD52 antibodies, anti-CD2 antibodies, TCR $\alpha\beta$ blockers, anti-CD20 antibodies, anti-CD19 antibodies, Bortezomib, rituximab, anti-CD 154 antibodies, rapamycin, CD3 immunotoxin, fludarabine, cyclophosphamide, busulfan, melphalan, Mabthera, Tacrolimus, alefacept, alemtuzumab, OKT3, OKT4, OKT8, OKT11, fingolimod, anti-CD40 antibodies, anti-BR3 antibodies, Campath-1H, anti-CD25 antibodies, calcineurin inhibitors, mycophenolate, and steroids, which may be used alone or in combination. As described herein, a lympho-depletion regimen can include, administration of alemtuzumab, cyclophosphamide, benduamustin, rituximab, pentostatin, and/or fludarabine. As described herein, lympho-depletion regimen can be administered in one or more cycles until the desired outcome of reduced circulating immune cells. As described herein, the lympho-depletion comprises administering an agent that specifically targets, and reduces or eliminates CD52+ cells in the subject, and the immune cells are modified to reduce or eliminate CD52 expression.

[0480] As described herein, an immune stimulating therapy is administered to the subject prior to, concurrently with, or after administration (e.g. infusion) of adoptive immune cells. As also described herein, the immune stimulating therapy comprises homeostatic cytokines. As also described herein, the immune stimulating therapy comprises immune-stimulatory molecules. As also described herein, the immune stimulating therapy comprises IL-2, IL-7, IL-12, IL-15, IL-21, IL-9, or a functional fragment thereof. As also described herein, the immune stimulating therapy comprises IL-2, IL-7, IL-12, IL-15, IL-21, IL-9, or combinations thereof. As also described herein, the immune stimulating therapy comprises IL-2, or a functional fragment thereof.

[0481] Methods for adoptive cell therapy using autologous cells includes isolating immune cells from patient blood, performing a series of modifications on the isolated cells including transducing the cells with one or more vectors encoding the dual receptor system described herein, and administering the cells to a patient. Providing immune cells from a subject suffering from or at risk for cancer or a hematological malignancy requires isolation of immune cell from the patient's blood, and can be accomplished through methods known in the art, for example, by leukapheresis. During leukapheresis, blood from a subject is extracted and the peripheral blood mononuclear cells (PBMCs) are separated, and the remainder of the blood is returned to the subject's circulation. The PBMCs are stored either frozen or cryopreserved as a sample of immune cells and provided for further processing steps, such as, e.g. the modifications described herein.

[0482] As described herein, the method of treating a subject described herein comprises modifications to immune cells from the subject comprising a series of modifications comprising enrichment and/or depletion, activation, genetic modification, expansion, formulation, and cryopreservation.

[0483] The disclosure provides enrichment and/or depletion steps that can be, for example, washing and fractionating

methods known in the art for preparation of subject PBMCs for downstream procedures, *e.g.* the modifications described herein. For example, methods can include devices to remove gross red blood cells and platelet contaminants, systems for size-based cell fractionation for the depletion of monocytes and the isolation of lymphocytes, and/or systems that allow the enrichment or depletion of specific subsets of T cells, such as, *e.g.* CD4+, CD8+, CD25+, or CD62L+ T cells. Following the enrichment steps, a target sub-population of immune cells will be isolated from the subject PMBCs for further processing. Those skilled in the art will appreciate that enrichment steps, as provided herein, may also encompass any newly discovered method, device, reagent or combination thereof.

[0484] The disclosure provides activation steps that can be any method known in the art to induce activation of immune cells, *e.g.* T cells, required for their *ex vivo* expansion. Immune cell activation can be achieved, for example, by culturing the subject immune cells in the presence of dendritic cells, culturing the subject immune cells in the presence of artificial antigen-presenting cells (AAPCs), or culturing the immune cells in the presence of irradiated K562-derived AAPCs. Other methods for activating subject immune cells can be, for example, culturing the immune cells in the presence of isolated activating factors and compositions, *e.g.* beads, surfaces, or particles functionalized with activating factors. Activating factors can include, for example, antibodies, *e.g.* anti-CD3 and/or anti-CD28 antibodies. Activating factors can also be, for example, cytokines, *e.g.* interleukin (IL)-2 or IL-21. Activating factors can also be costimulatory molecules, such as, for example, CD40, CD40L, CD70, CD80, CD83, CD86, CD137L, ICOSL, GITRL, and CD134L. Those skilled in the art will appreciate that activating factors, as provided herein, may also encompass any newly discovered activating factor, reagent, composition, or combination thereof that can activate immune cells.

[0485] The disclosure provides genetic modification steps for modifying the subject immune cells. For example, the genetic modification comprises transducing the immune cell with a vector comprising a shRNA described herein complementary to B2M or HLA-A. The genetic modification may comprise modifying the genome of the immune cells to induce mutations in B2M or HLA-A using CRISPR/Cas mediated genome engineering. For example, the method comprises transducing the immune cell with one or more vectors encoding the activator and inhibitory receptors, thereby producing immune cells expressing the activator and inhibitory receptors.

[0486] The disclosure provides expansion steps for the genetically modified subject immune cells. Genetically modified subject immune cells can be expanded in any immune cell expansion system known in the art to generate therapeutic doses of immune cells for administration. For example, bioreactor bags for use in a system comprising controller pumps, and probes that allow for automatic feeding and waste removal can be used for immune cell expansion. Cell culture flasks with gas-permeable membranes at the base may be used for immune cell expansion. Any such system known in the art that enables expansion of immune cells for clinical use is encompassed by the expansion step provided herein. Immune cells are expanded in culture systems in media formulated specifically for expansion. Expansion can also be facilitated by culturing the immune cell of the disclosure in the presence of activation factors as described herein. Those skilled in the art will appreciate that expansion steps, as provided herein, may also encompass any newly discovered culture systems, media, or activating factors that can be used to expand immune cells.

[0487] The disclosure provides formulation and cryopreservation steps for the expanded genetically modified subject immune cells. Formulation steps provided include, for example, washing away excess components used in the preparation and expansion of immune cells of the methods of treatment described herein (not according to the invention). Any pharmaceutically acceptable formulation medium or wash buffer compatible with immune cell known in the art may be used to wash, dilute/concentrate immune cells, and prepare doses for administration. Formulation medium can be acceptable for administration of the immune cells, such as, for example crystalloid solutions for intravenous infusion.

[0488] Cryopreservation can optionally be used to store immune cells long-term. Cryopreservation can be achieved using known methods in the art, including for example, storing cells in a cryopreservation medium containing cryopreservation components. Cryopreservation components can include, for example, dimethyl sulfoxide or glycerol. Immune cells stored in cryopreservation medium can be cryopreserved by reducing the storage temperature to -80°C to -196 °C.

[0489] As described herein, but not according to the invention, the method of treatment comprises determining the HLA germline type of the subject. As described herein, the HLA germline type is determined in bone marrow.

[0490] As described herein, but not according to the invention, the method of treatment comprises determining the level of expression of EGFR. As described herein, but not according to the invention, the level of expression of EGFR is determined in tumor tissue samples from the subject. As described herein, but not according to the invention, the expression level of EGFR is determined using next generation sequencing. As described herein, but not according to the invention, the expression level of EGFR is determined using RNA sequencing. As described herein, but not according to the invention, the level of EGFR is determined using immunohistochemistry.

[0491] As described herein, but not according to the invention, the method of treatment comprises administering a therapeutically effective dose of immune cells comprising an HLA-A*02 inhibitory receptor to a subject in need thereof, wherein the subject is determined to be HLA germline HLA-A*02 heterozygous and have cancer cells with loss of HLA-A*02. As described herein, but not according to the invention, the method of treatment comprises administering a therapeutically effective dose of immune cells comprising an HLA-A*01 inhibitory receptor to a subject in need thereof, wherein the subject is determined to be HLA germline HLA-A*01 heterozygous and have cancer cells with loss of HLA-

A*01. As described herein, but not according to the invention, the method of treatment comprises administering a therapeutically effective dose of immune cells comprising an HLA-A*03 to a subject in need thereof, wherein the subject is determined to be HLA germline HLA-A*03 heterozygous and have cancer cells with loss of HLA-A*03. As described herein, but not according to the invention, the method of treatment comprises administering a therapeutically effective dose of immune cells comprising an HLA-A*07 inhibitory receptor to a subject in need thereof, wherein the subject is determined to be HLA germline HLA-A*07 heterozygous and have cancer cells with loss of HLA-A*07. As described herein, but not according to the invention, the method of treatment comprises administering a therapeutically effective dose of immune cells comprising an HLA-C*07 inhibitory receptor to a subject in need thereof, wherein the subject is determined to be HLA germline HLA-C*07 heterozygous and have cancer cells with and loss of HLA-C*07. As described herein, but not according to the invention, the method of treatment comprises administering a therapeutically effective dose of immune cells comprising an HLA-B*07 inhibitory receptor in a subject in need thereof, wherein the subject is determined to be HLA germline HLA-B*07 heterozygous and have cancer cells with loss of HLA-B*07.

[0492] Described herein, but not according to the present invention is a method of treatment of heterozygous HLA-A*02 patients with malignancies that express EGFR and have lost HLA-A*02 expression; and/or of treatment of heterozygous HLA-A*02 adult patients with recurrent unresectable or metastatic solid tumors that express EGFR and have lost HLA-A*02 expression.

[0493] As described herein, a therapeutically effective dose of the immune cells described herein are administered. As described herein, the immune cells of the disclosure are administered by intravenous injection. As described herein, the immune cells of the disclosure are administered by intraperitoneal injection. As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells, about 1×10^6 cells, about 2×10^6 cells, about 3×10^6 cells, 4×10^6 cells, about 5×10^6 cells, about 6×10^6 cells, about 7×10^6 cells, about 8×10^6 cells, about 9×10^6 cells, about 1×10^7 , about 2×10^7 , about 3×10^7 , about 4×10^7 , about 5×10^7 , about 6×10^7 , about 7×10^7 , about 8×10^7 , about 9×10^7 , about 1×10^8 cells, about 2×10^8 cells, about 3×10^8 cells, about 4×10^8 cells, about 5×10^8 cells, about 6×10^8 cells, about 7×10^8 cells, about 8×10^8 cells, about 9×10^8 cells, about 1×10^9 cells, about 2×10^9 cells, about 3×10^9 cells, about 3×10^9 cells, about 4×10^9 cells, about 5×10^9 cells, about 5×10^9 cells, about 6×10^9 cells, about 7×10^9 cells, about 8×10^9 cells, about 9×10^9 cells, about 1×10^{10} cells, about 2×10^{10} cells, about 3×10^{10} cells, about 4×10^{10} cells, about 5×10^{10} cells, about 6×10^{10} cells, about 7×10^{10} cells, about 8×10^{10} cells, or about 9×10^{10} cells.

[0494] As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells to about 9×10^{10} cells, about 1×10^6 cells to about 5×10^{10} cells, about 2×10^6 cells to about 5×10^9 cells, about 3×10^6 cells to about 5×10^9 cells, about 4×10^6 cells to about 3×10^9 cells, about 5×10^6 cells to about 2×10^9 cells, about 6×10^6 cells to about 1×10^9 cells, 0.5×10^6 cells to about 6×10^9 cells, about 1×10^6 cells to about 5×10^9 cells, about 2×10^6 cells to about 5×10^9 cells, about 3×10^6 cells to about 4×10^9 cells, about 4×10^6 cells to about 3×10^9 cells, about 5×10^6 cells to about 2×10^9 cells, about 6×10^6 cells to about 1×10^9 cells, 0.5×10^6 cells to about 6×10^8 cells, about 1×10^6 cells to about 5×10^8 cells, about 2×10^6 cells to about 5×10^8 cells, about 3×10^6 cells to about 4×10^8 cells, about 4×10^6 cells to about 3×10^8 cells, about 5×10^6 cells to about 2×10^8 cells, about 6×10^6 cells to about 1×10^8 cells, about 7×10^6 cells to about 9×10^8 cells, about 8×10^6 cells to about 8×10^8 cells, about 9×10^6 cells to about 7×10^8 cells, about 1×10^7 cells to about 6×10^8 cells, about 2×10^7 cells to about 5×10^8 cells, about 7×10^6 cells to about 9×10^7 cells, about 8×10^6 cells to about 8×10^7 cells, about 9×10^6 cells to about 7×10^7 cells, about 1×10^7 cells to about 6×10^7 cells, or about 2×10^7 cells to about 5×10^7 cells.

[0495] As described herein, a therapeutically effective dose comprises about 0.5×10^5 cells to about 9×10^{10} cells. As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells to about 1×10^{10} cells. As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells to about 5×10^9 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells to about 1×10^9 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells to about 6×10^8 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^6 cells to about 9×10^{10} cells. As described herein, a therapeutically effective dose comprises about 0.5×10^7 cells to about 1×10^{10} cells. As described herein, a therapeutically effective dose comprises about 0.5×10^7 cells to about 5×10^9 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^7 cells to about 1×10^9 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^7 cells to about 6×10^8 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^8 cells to about 9×10^{10} cells. As described herein, a therapeutically effective dose comprises about 0.5×10^8 cells to about 1×10^{10} cells. As described herein, a therapeutically effective dose comprises about 0.5×10^8 cells to about 5×10^9 cells. As described herein, a therapeutically effective dose comprises about 0.5×10^8 cells to about 1×10^9 cells. The term "about" as referred to in a therapeutically dose, can be, for example, $\pm 0.5 \times 10^6$ cells, $\pm 0.5 \times 10^7$ cells, or $\pm 0.5 \times 10^8$ cells.

Kits and Articles of Manufacture

[0496] The disclosure provides kits and articles of manufacture comprising the polynucleotides and vectors encoding the receptors described herein, and immune cells comprising the receptors described herein. As described herein, the

kit comprises articles such as vials, syringes and instructions for use.

[0497] As also described herein, the kit comprises a polynucleotide or vector comprising a sequence encoding one or more receptors of the disclosure.

[0498] As also described herein, the kit comprises a plurality of immune cells comprising the first and second receptors as described herein. The plurality of immune cells may comprise a plurality of T cells.

[0499] As also described herein, the kit further comprises instructions for use.

ENUMERATED ITEMS (NOT ACCORDING TO THE INVENTION)

[0500] The present invention is defined in the appended claims.

[0501] Item 1. An immune cell, comprising:

a.) a first receptor, comprising an extracellular ligand binding domain specific to Epidermal Growth Factor Receptor (EGFR); and

b.) a second receptor, comprising an extracellular ligand binding domain specific to a non-target antigen lost in EGFR+ cancer due to loss of heterozygosity,

wherein the first receptor is an activator receptor responsive to EGFR; and wherein the second receptor is an inhibitory receptor responsive to the non-target antigen.

[0502] Item 2. The immune cell of Item 1, wherein the extracellular ligand binding domain of the second receptor specifically binds an allelic variant of an MHC protein. Item 3. The immune cell of Item 1, wherein the extracellular ligand binding domain of the second receptor specifically binds an allelic variant of an HLA-A, HLA-B, or HLA-C protein.

[0503] Item 4. The immune cell of Item 1, wherein the extracellular ligand binding domain of the second receptor specifically binds HLA-A*02.

[0504] Item 5. The immune cell of Item 1, wherein the extracellular ligand binding domain of the second receptor specifically binds HLA-A*01, HLA-A*03, HLA-A*11, HLA-C*07, or HLA-B*07.

[0505] Item 6. The immune cell of any one of Items 3-5, wherein the extracellular ligand binding domain of the second receptor comprises complementarity determining regions (CDRs) CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 as disclosed in Table 5; or CDR sequences having at most 1, 2, or 3 substitutions, deletions, or insertions relative to the CDRs of Table 5.

[0506] Item 7. The immune cell of any one of Items 3-5, wherein the extracellular ligand binding domain of the second receptor comprises complementarity determining regions (CDRs) CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 of SEQ ID NOs: 101-106 or of SEQ ID NOs: 106-112; or CDR sequences having at most 1, 2, or 3 substitutions, deletions, or insertions relative to the CDRs of Table 5.

[0507] Item 8. The immune cell of any one of Items 3-5, wherein the extracellular ligand binding domain of the second receptor comprises a polypeptide sequence selected from the polypeptide sequence disclosed in Table 4; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0508] Item 9. The immune cell of Item 4, wherein the extracellular ligand binding domain of the second receptor comprises any one of SEQ ID NOs: 89-100; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0509] Item 10. The immune cell of any one of Items 1-9, wherein the first receptor is a chimeric antigen receptor (CAR).

[0510] Item 11. The immune cell of any one of Items 1-10, wherein the extracellular ligand binding domain of the first receptor comprises a variable heavy (VH) portion comprising a set of heavy chain complementarity determining regions (HC-CDRs) selected from the group of sequences set forth in Table 3; and/or a variable light (VL) portion comprising a set of light chain complementarity determining regions (LC-CDRs) from the group of sequences set forth in Table 3; or CDR sequences having at most 1, 2, 3, 4 substitutions, insertions, or deletions in each CDR.

[0511] Item 12. The immune cell of any one of Items 1-11, wherein the extracellular ligand binding domain of the first receptor comprises a variable heavy (VH) portion having a sequence selected from the VH sequence set forth in Table 2; and/or a variable light (VL) portion comprising a sequence set forth in Table 2; or sequences having at least 70%, at least 85%, at least 90%, or at least 95% identity thereto.

[0512] Item 13. The immune cell of any one of Items 1-12, wherein the extracellular ligand binding domain of the first receptor comprising a sequence selected from the group of sequences set forth in Table 1; or a sequence having at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% identity thereto. Item 14. The immune cell of any one of Items 1-13, wherein the extracellular ligand binding domain of the first receptor comprises an scFv sequence selected from the group consisting of SEQ ID NO: 9-18; or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto. Item 15. The immune cell of any one of Items 1-14, wherein the second receptor comprises a LILRB1 intracellular domain or a functional variant thereof.

[0513] Item 16. The immune cell of Item 15, wherein the LILRB1 intracellular domain comprises a sequence at least

90%, at least 95%, at least 97%, at least 99%, or is identical to SEQ ID NO: 129.

[0514] Item 17. The immune cell of any one of Items 1-16, wherein the second receptor comprises a LILRB1 transmembrane domain or a functional variant thereof.

[0515] Item 18. The immune cell of Item 17, wherein the LILRB1 transmembrane domain or a functional variant thereof comprises a sequence at least 90%, at least 95%, at least 97%, at least 99% or is identical to SEQ ID NO: 133.

[0516] Item 19. The immune cell of any one of Items 1-18, wherein the second receptor comprises a LILRB1 hinge domain or functional variant thereof.

[0517] Item 20. The immune cell of Item 19, wherein the LILRB1 hinge domain comprises a sequence at least 90%, at least 95%, at least 97%, at least 99% or is identical to SEQ ID NO: 132, SEQ ID NO: 125, SEQ ID NO: 126.

[0518] Item 21. The immune cell of any one of Items 1-20, wherein the second receptor comprises a LILRB1 intracellular domain, a LILRB1 transmembrane domain, a LILRB1 hinge domain, a functional variant of any of these, or combinations thereof.

[0519] Item 22. The immune cell of Item 21, wherein the LILRB1 intracellular domain and LILRB1 transmembrane domain comprises SEQ ID NO: 128 or a sequence at least 95% identical to SEQ ID NO: 128.

[0520] Item 23. The immune cell of any one of Items 1-22, wherein the EGFR+ cancer cell is a lung cancer cell, a small cell lung cancer cell, a non-small cell lung cancer cell, a pancreatic ductal carcinoma cell, a colorectal cancer cell, a head and neck cancer cell, a esophagus and gastric adenocarcinoma cell, an ovarian cancer cell, a glioblastoma multiforme cell, a cervical squamous cell carcinoma cell, a kidney cancer cell, a papillary kidney cancer cell, a kidney renal clear cell carcinoma cell, a bladder cancer cell, a breast cancer cell, a bile duct cancer cell, a liver cancer cell, a prostate cancer cell, a sarcoma cell, a thyroid cancer cell, a thymus cancer cell, a stomach cancer cell, or a uterine cancer cell.

[0521] Item 24. The immune cell of Item 23, wherein the EGFR+ cancer cell is a lung cancer cell.

[0522] Item 25. The immune cell of any one of Items 1-24, wherein the EGFR+ cancer cell is an EGFR+/HLA-A*02- cancer cell that does not express HLA-A*02; or a cancer cell derived from an HLA-A*02+ individual, which does not express HLA-A*02.

[0523] Item 26. The immune cell of Item 25, wherein the EGFR+/HLA-A*02- cancer cell is derived from an EGFR+/HLA-A*02+ cell by loss of heterozygosity at HLA-A leading to loss of HLA-A*02.

[0524] Item 27. The immune cell of any one of Items 1-26, wherein the first receptor and the second receptor together specifically activate the immune cell in the presence of the EGFR/HLA-A*02- cancer cell having loss of heterozygosity.

[0525] Item 28. The immune cell of any one of Items 1-27, wherein the first receptor and the second receptor together do not specifically activate the immune cell in the presence of an EGFR+ cell that has not lost HLA-A* 02 by loss of heterozygosity.

[0526] Item 29. The immune cell of any one of Items 1-28, wherein the immune cell is a T cell, macrophage, NK cell, iNKT cell, or a gamma delta T cell. Item 30. The immune cell of Item 29, wherein the T cell is a CD8+ CD4- T cell.

[0527] Item 31. The immune cell of any one of Items 1-30, wherein expression and/or function of an MHC Class I gene has been reduced or eliminated.

[0528] Item 32. The immune cell of Item 30, wherein the MHC Class I gene is beta-2-microglobulin (B2M).

[0529] Item 33. The immune cell of Item 32, further comprising a polynucleotide comprising an interfering RNA, the interfering RNA comprising a sequence complementary to a sequence of a B2M mRNA (SEQ ID NO: 172).

[0530] Item 34. The immune cell of Item 33, wherein the interfering RNA is capable of inducing RNAi-mediated degradation of the B2M mRNA.

[0531] Item 35. The immune cell of Item 33, wherein the interfering RNA is a short hairpin RNA (shRNA).

[0532] Item 36. The immune cell of Item 35, wherein the shRNA comprises:

- a.) a first sequence, having from 5'-end to 3'-end a sequence complementary to a sequence of the B2M mRNA; and
- b.) a second sequence, having from 5'-end to 3'-end a sequence complementary to the first sequence,

wherein the first sequence and the second sequence form the shRNA.

[0533] Item 37. The immune cell of Item 32, comprising one or more modifications to a sequence encoding B2M (SEQ ID NO: 170), wherein the one or more modifications reduce the expression and/or eliminate the function of B2M.

[0534] Item 38. The immune cell of Item 37, wherein the one or more modifications comprise one or more inactivating mutations of the endogenous gene encoding B2M.

[0535] Item 39. The immune cell of Item 38, wherein the one or more inactivating mutations comprise a deletion, an insertion, a substitution, or a frameshift mutation.

[0536] Item 40. The immune cell of any one of Items 38-39, wherein the one or more inactivating mutations are introduced with a nucleic acid guided endonuclease in a complex with at least one guide nucleic acid (gNA) that specifically targets a sequence of the endogenous gene encoding B2M (SEQ ID NO: 170).

[0537] Item 41. The immune cell of Item 31, wherein the MHC Class I gene is HLA-A*02. Item 42. The immune cell

of Item 41, further comprising a polynucleotide comprising an interfering RNA, comprising a sequence complementary to a sequence of an HLA-A*02 mRNA (SEQ ID NO: 171).

[0538] Item 43. The immune cell of Item 42, wherein the interfering RNA is capable of inducing RNA interference (RNAi)-mediated degradation of the HLA-A*02 mRNA.

[0539] Item 44. The immune cell of Item 43, wherein the interfering RNA is a short hairpin RNA (shRNA) comprising:

- a.) a first sequence, having from 5'-end to 3'-end a sequence complementary to a sequence of the HLA-A*02 mRNA; and
- b.) a second sequence, having from 5'-end to 3'-end a sequence complementary to the first sequence,

wherein the first sequence and the second sequence form the shRNA.

[0540] Item 45. The immune cell of Item 41, comprising one or more modifications to a sequence of an endogenous gene encoding HLA-A*02 (SEQ ID NO: 169), wherein the one or more modifications reduce the expression and/or eliminate the function of HLA-A*02. Item 46. The immune cell of Item 45, wherein the one or more modifications comprise one or more inactivating mutations of the endogenous gene encoding HLA-A*02.

[0541] Item 47. The immune cell of Item 45 or Item 46, wherein the one or more inactivating mutations are introduced with a nucleic acid guided endonuclease in a complex with at least one guide nucleic acid (gNA) that specifically targets a sequence of the endogenous gene encoding HLA-A*02.

[0542] Item 48. The immune cell of Item 1, wherein:

- a.) the first receptor comprises SEQ ID NO: 177, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and
- b.) the second receptor comprises SEQ ID NO: 174 or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0543] Item 49. The immune cell of Item 1, wherein:

- a.) the first receptor comprises SEQ ID NO: 177, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and
- b.) the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0544] Item 50. The immune cell of Item 1, wherein:

- a.) the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and
- b.) the second receptor comprises SEQ ID NO: 174, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0545] Item 51. The immune cell of Item 1, wherein:

- a.) the first receptor comprises SEQ ID NO: 175, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and
- b.) the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0546] Item 52. The immune cell of Item 1, wherein:

- a.) the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and
- b.) the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0547] Item 53. The immune cell of Item 1, wherein:

- a.) the first receptor comprises SEQ ID NO: 176, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto; and
- b.) the second receptor comprises SEQ ID NO: 173, or a sequence sharing at least 75%, at least 80%, at least 85%,

at least 90%, at least 95%, or at least 98% identity thereto.

[0548] Item 54. The immune cell of any one of Items 48 to 53, further comprising a T2A self-cleaving peptide, wherein the T2A self-cleaving peptide comprises SEQ ID NO: 178, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0549] Item 55. The immune cell of any one of Items 48 to 54, further comprising an interfering RNA, wherein the interfering RNA comprises SEQ ID NO: 179, or a sequence sharing at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% identity thereto.

[0550] Item 56. The immune cell of any one of Items 1-55, wherein the immune cell is autologous.

[0551] Item 57. The immune cell of any one of Items 1-55, wherein the immune cell is allogeneic.

[0552] Item 58. A pharmaceutical composition, comprising a therapeutically effective amount of the immune cells of any one of Items 1-57.

[0553] Item 59. The pharmaceutical composition of Item 58, wherein the immune cell expresses both the first receptor and the second receptor.

[0554] Item 60. The pharmaceutical composition of Item 59, wherein at least about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% of the immune cells express both the first receptor and the second receptor.

[0555] Item 61. The pharmaceutical composition of Item 59, wherein at least 90% of the immune cells express both the first receptor and the second receptor.

[0556] Item 62. The pharmaceutical composition of any one of Items 58-61, further comprising a pharmaceutically acceptable carrier, diluent or excipient.

[0557] Item 63. The pharmaceutical composition of any one of Items 58-62, for use as a medicament in the treatment of EGFR+ cancer.

[0558] Item 64. A polynucleotide or polynucleotide system, comprising one or more polynucleotides comprising polynucleotide sequences encoding:

a.) a first receptor, comprising an extracellular ligand binding domain specific to Endothelial Growth Factor Receptor (EGFR); and

b.) a second receptor, comprising an extracellular ligand binding domain specific to a non-target antigen that has been lost in the EGFR+ cancer cell due to loss of heterozygosity,

wherein the first receptor is an activator receptor responsive to EGFR on the EGFR+ cancer cell; and wherein the second receptor is an inhibitory receptor responsive to the non-target antigen.

[0559] Item 65. A polynucleotide or polynucleotide system, comprising one or more polynucleotides comprising polynucleotide sequences encoding the first receptor and the second receptor for use in generating the immune cells of any one of Items 1-57.

[0560] Item 66. A vector, comprising the one or more polynucleotides of Item 64 or 65.

[0561] Item 67. A method of killing EGFR+ cancer cell having loss of heterozygosity at an MHC class I locus, comprising administering to a subject an effective amount of the immune cell of any one of Items 1-57 or the pharmaceutical composition of any one of Items 58-62.

[0562] Item 68. The method of Item 67, wherein the EGFR+ cancer cell is a lung cancer cell, a small cell lung cancer cell, a non-small cell lung cancer cell, a pancreatic ductal carcinoma cell, a colorectal cancer cell, a head and neck cancer cell, a esophagus and gastric adenocarcinoma cell, an ovarian cancer cell, a glioblastoma multiforme cell, a cervical squamous cell carcinoma cell, a kidney cancer cell, a papillary kidney cancer cell, a kidney renal clear cell carcinoma cell, a bladder cancer cell, a breast cancer cell, a bile duct cancer cell, a liver cancer cell, a prostate cancer cell, a sarcoma cell, a thyroid cancer cell, a thymus cancer cell, a stomach cancer cell, or a uterine cancer cell.

[0563] Item 69. The method of Item 67, wherein the EGFR+ cancer cell is a lung cancer cell.

[0564] Item 70. The method of Item 67, wherein the EGFR+ cancer cell is an EGFR+/HLA-A*02- cancer cell that does not express HLA-A*02; or a cancer cell derived from an HLA-A*02+ individual, which does not express HLA-A*02.

[0565] Item 71. The method of Item 70, wherein the EGFR+/HLA-A*02- cancer cell is derived from an EGFR+/HLA-A*02+ cell by loss of heterozygosity at HLA-A leading to loss of HLA-A*02.

[0566] Item 72. A method of treating EGFR+ cancer in a subject having an EGFR+ tumor having loss of heterozygosity at a locus encoding a non-target antigen, comprising administering to the subject an effective amount of the immune cell of any one of Items 1-57 or the pharmaceutical composition of any one of Items 58-62.

[0567] Items 73. The method of Item 72, wherein the subject is a heterozygous HLA-A*02 patient with a malignancy that expresses EGFR and has lost HLA-A*02 expression.

[0568] Item 74. The method of Item 72, wherein the subject is a heterozygous HLA-A*02 patient with recurrent unresectable or metastatic solid tumors that express EGFR and have lost HLA-A*02 expression.

[0569] Item 75. A method of treating a cancer in a subject comprising:

- a.) determining the genotype or expression level of a non-target antigen in non-malignant cells and cancer cells of the subject;
- b.) determining the expression level of EGFR in cancer cells of the subject; and
- c.) administering to the subject an effective amount of the immune cell of any one of Items 1-57 or the pharmaceutical composition of any one of Items 58-62 if the non-malignant cells express the non-target antigen and the cancer cells do not express the non-target antigen, and the cancer cells are EGFR positive.

[0570] Item 76. The method of any one of Items 67-75, wherein administration of the immune cell of any one of Items 1-57 or the pharmaceutical composition of any one of Items 58-62 reduces the size of a tumor in the subject.

[0571] Item 77. The method of Items 76, wherein the tumor is reduced by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, or about 100%.

[0572] Item 78. The method of Item 76, wherein the tumor is eliminated.

[0573] Item 79. The method of any one of Items 67-75, wherein administration of the immune cell or the pharmaceutical composition arrests the growth of a tumor in the subject.

[0574] Item 80. The method of any one of Items 67-75, wherein administration of the immune cell of any one of Items 1-57 or the pharmaceutical composition of any one of Items 58-62 reduces the number of tumors in the subject.

[0575] Item 81. The method of any one of Items 67-80, wherein administration of the immune cell or the pharmaceutical composition results in selective killing of a cancer cell, but not a normal cell in the subject.

[0576] Item 82. The method of Item 81, wherein at least about 60% of the cells killed are cancer cells, at least about 65% of the cells killed are cancer cells, at least about 70% of the cells killed are cancer cells, at least about 75% of the cells killed are cancer cells, at least about 80% of the cells killed are cancer cells, at least about 85% of the cells killed are cancer cells, at least about 90% of the cells killed are cancer cells, at least about 95% of the cells killed are cancer cells, or about 100% of the cells killed are cancer cells.

[0577] Item 83. The method of Item 81, wherein administration of the immune cell or pharmaceutical composition results in the killing of about 40%, about 50%, about 60%, about 70%, about 80%, about 90% or all of the cancer cells of the subject.

[0578] Item 84. The method of any one of Items 67-83, wherein administration of the immune cell or the pharmaceutical composition results in fewer side effects for the subject than administration of an otherwise equivalent immune cell comprising the first activator receptor, but no second inhibitory receptor.

[0579] Item 85. A method of making a plurality of immune cells, comprising:

- a.) providing a plurality of immune cells, and
- b.) transforming the plurality of immune cells with the polynucleotide system of Item 64 or Item 65, or the vector of Item 66.

[0580] Item 86. A kit comprising the immune cell of any one of Items 1-57 or the pharmaceutical composition of any one of Items 58-62.

[0581] Item 87. The kit of Item 86, further comprising instructions for use.

[0582] Item 88. An immune cell responsive to loss of heterozygosity in a cancer cell, comprising:

- a. a first receptor, optionally a chimeric antigen receptor (CAR) or T cell receptor (TCR), comprising an extracellular ligand binding domain specific to a target antigen selected from:

- i. a cancer cell-specific antigen, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); or

- ii. epidermal growth factor receptor (EGFR), or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); and

- b. a second receptor, optionally an inhibitory receptor, comprising an extracellular ligand binding domain specific to a non-target antigen that has been lost in the cancer cell due to loss of heterozygosity.

[0583] Item 89. The method of Item 88, wherein the non-target antigen comprises a polymorphism.

[0584] Item 90. The immune cell of Item 89, wherein the non-target antigen comprises a COLEC12 antigen that shares at least 95% identity to SEQ ID NO: 87 and the polymorphism comprises an S or P at position 522 of SEQ ID NO: 87.

[0585] Item 91. The immune cell of Item 89, wherein the non-target antigen comprises an APCDD1 antigen that shares

at least 95% identity to SEQ ID NO: 88, SEQ ID NO: 134 or SEQ ID NO: 135, and the polymorphism is selected from the group consisting of:

- a. A, V, I or L at position 150 of SEQ ID NO: 88;
- b. a Y or S at position 165 of SEQ ID NO: 134; and
- c. A, Q or R at position 28 of SEQ ID NO: 135.

[0586] Item 92. The immune cell of Item 89, wherein the non-target antigen comprises a CXCL16 antigen that shares at least 95% identity to SEQ ID NO: 86 and the polymorphism is selected from the group consisting of:

- a. an I or T at position 142 of SEQ ID NO: 86; and
- b. an A or V at position 200 of SEQ ID NO: 86.

[0587] Item 93. The immune cell of Item 88 or 89, wherein the non-target antigen comprises HLA-A*02.

[0588] Item 94. The immune cell of any one of Items 88-93, wherein the target antigen is expressed by the cancer cell.

[0589] Item 95. The immune cell of any one of Items 88-94, wherein the non-target antigen is not expressed by the cancer cell.

[0590] Item 96. The immune cell of any one of Items 88-95, wherein the non-target antigen is expressed by a non-target cell.

[0591] Item 97. The immune cell of any one of Items 88-96, wherein the non-target cell expresses both target antigen and the non-target antigen.

[0592] Item 98. The immune cell of any one of Items 88-97, wherein the target antigen is a cancer cell-specific antigen.

[0593] Item 99. The immune cell of any one of Items 88-98, wherein the target antigen is a peptide antigen of a cancer cell-specific antigen in a complex with a major histocompatibility complex class I (MHC-I).

[0594] Item 100. The immune cell of any one of Items 88-99, wherein the cancer cell is a lung cancer cell, a glioblastoma cell, a breast cancer cell, a head and neck cancer cell or a colorectal cancer cell.

[0595] Item 101. The immune cell of any one of Items 88-100, wherein the cancer cell expresses EGFR.

[0596] Item 102. The immune cell of any one of Items 88-101, wherein the first receptor and the second receptor together specifically activate the immune cell in the presence of the cancer cell.

[0597] Item 103. The immune cell of Item 102, wherein the immune cell is a T cell, macrophage, NK cell, iNKT cell, or a gamma delta T cell.

[0598] Item 104. The immune cell of Item 103, wherein the T cell is a CD8⁺ CD4⁻ T cell.

[0599] Item 105. The immune cell of any one of Items 88-104, wherein the EGFR comprises a sequence that shares at least 95% identity to any one of SEQ ID NOs: 1-8.

[0600] Item 106. The immune cell of any one of Items 88-105, wherein the first receptor is a T cell receptor (TCR).

[0601] Item 107. The immune cell of any one of Items 88-106, wherein the first receptor is a chimeric antigen receptor (CAR).

[0602] Item 108. The immune cell of Item 106 or 107, wherein the extracellular ligand binding domain of the first receptor comprises an antibody fragment, a single chain Fv antibody fragment (scFv), a β chain variable domain (V β), or a TCR α chain variable domain and a TCR β chain variable domain.

[0603] Item 109. The immune cell of Item 106 or 107, wherein the extracellular ligand binding domain of the first receptor comprises an scFv.

[0604] Item 110. The immune cell of Item 109, wherein the scFv comprises a sequence selected from the group consisting of SEQ ID NOs: 9-18, or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0605] Item 111. The immune cell of Item 109, wherein the scFv comprises or consists essentially of a sequence selected from the group consisting of SEQ ID NOs: 9-18.

[0606] Item 112. The immune cell of Item 106 or 107, wherein the extracellular ligand binding domain of the first receptor comprises VH and VL domain.

[0607] Item 113. The immune cell of Item 112, wherein the VH domain comprises a sequence selected from the group consisting of SEQ ID NOs: 19-24, or a VH sequence disclosed in Table 2, or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0608] Item 114. The immune cell of Item 112, wherein the VH domain comprises or consists essentially of a sequence selected from the group consisting of SEQ ID NOs: 19-24, or a VH sequence disclosed in Table 2.

[0609] Item 115. The immune cell of any one of Items 112-114, wherein the VL domain comprises a sequence selected from the group consisting of SEQ ID NOs: 25-30, or a VL sequence disclosed in Table 2, or a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity thereto.

[0610] Item 116. The cell of any one of Items 112-114, wherein the VL domain comprises or consists essentially of a

sequence selected from the group consisting of SEQ ID NOs: 25-30, or a VL sequence disclosed in Table 2.

[0611] Item 117. The immune cell of any one of Items 105-107, wherein the extracellular ligand binding domain of the first receptor comprises complementarity-determining regions (CDRs) selected from the group consisting of SEQ ID NOs: 31-65, or a CDR sequence disclosed in Table 3.

[0612] Item 118. The immune cell of any one of Items 88-117, wherein the extracellular ligand binding domain of the second receptor comprises an antibody fragment, a single chain Fv antibody fragment (scFv), a β chain variable domain ($V\beta$), or a TCR α chain variable domain and a TCR β chain variable domain.

[0613] Item 119. The immune cell of any one of Items 88 or 93-118, wherein the extracellular ligand binding domain of the second receptor comprises an scFv.

[0614] Item 120. The immune cell of Item 119, wherein the scFv comprises a sequence having at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to any one of SEQ ID NOs: 89-100.

[0615] Item 121. The immune cell of Item 119, wherein the scFv comprises or consists essentially of a sequence of any one of SEQ ID NOs: 89-100.

[0616] Item 122. The immune cell of any one of Items 88 or 93-118, wherein the extracellular ligand binding domain of the second receptor comprises CDRs selected from the group consisting of SEQ ID NOs: 101-112.

[0617] Item 123. The immune cell of any one of Items 88-122, wherein the second receptor comprises a LILRB1 intracellular domain or a functional variant thereof.

[0618] Item 124. The immune cell of Item 123, wherein the LILRB1 intracellular domain comprises a sequence at least 95% identical to SEQ ID NO: 124.

[0619] Item 125. The immune cell of any one of Items 88-124, wherein the second receptor comprises a LILRB1 transmembrane domain or a functional variant thereof.

[0620] Item 126. The immune cell of Item 125, wherein the LILRB1 transmembrane domain or a functional variant thereof comprises a sequence at least 95% identical to SEQ ID NO: 133.

[0621] Item 127. The immune cell of any one of Items 88-126, wherein the second receptor comprises a LILRB1 hinge domain or functional fragment or variant thereof.

[0622] Item 128. The immune cell of Item 127, wherein the LILRB1 hinge domain comprises a sequence at least 95% identical to SEQ ID NO: 132, SEQ ID NO: 125 or SEQ ID NO: 126.

[0623] Item 129. The immune cell of any one of Items 88-128, wherein the second receptor comprises a LILRB1 intracellular domain and a LILRB1 transmembrane domain, or a functional variant thereof.

[0624] Item 130. The immune cell of Item 129, wherein the LILRB1 intracellular domain and LILRB1 transmembrane domain comprises SEQ ID NO: 128 or a sequence at least 95% identical to SEQ ID NO: 128.

[0625] Item 131. The immune cell of any one of Item 88-130, wherein the immune cell is a T cell, macrophage, NK cell, iNKT cell, or a gamma delta T cell.

[0626] Item 132. The immune cell of Item 131, wherein the T cell is a CD8⁺ CD4⁻ T cell.

[0627] Item 133. A pharmaceutical composition, comprising a therapeutically effective amount of the immune cells of any one of Items 88-132.

[0628] Item 134. The pharmaceutical composition of Item 133, further comprising a pharmaceutically acceptable carrier, diluent or excipient.

[0629] Item 135. The pharmaceutical composition of Items 133 or 134, for use as a medicament in the treatment of cancer.

[0630] Item 136. A polynucleotide system, comprising polynucleotides comprising polynucleotide sequences encoding:

a. a first receptor, optionally a chimeric antigen receptor (CAR) or T cell receptor (TCR), comprising an extracellular ligand binding domain specific to a target antigen selected from:

- i. a cancer cell-specific antigen, or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); or
- ii. epidermal growth factor receptor (EGFR), or a peptide antigen thereof in a complex with a major histocompatibility complex class I (MHC-I); and

b. a second receptor, optionally an inhibitory receptor, comprising an extracellular ligand binding domain specific to a non-target antigen that has been lost in the cancer cell due to loss of heterozygosity.

[0631] Item 137. The polynucleotide system of Item 136, wherein the non-target antigen comprises a polymorphism.

[0632] Item 138. The polynucleotide system of Item 137, wherein the non-target antigen is an COLEC12 antigen that shares at least 95% identity to SEQ ID NO: 87 and the polymorphism comprises an S or P at position 522 of SEQ ID NO: 87.

[0633] Item 139. The polynucleotide system of Item 137, wherein the non-target antigen is a APCDD1 antigen that shares at least 95% identity to SEQ ID NO: 88, SEQ ID NO: 134 or SEQ ID NO: 135, and the polymorphism is selected

from the group consisting of:

- a. A, V, I or L at position 150 of SEQ ID NO: 88;
- b. a Y or S at position 165 of SEQ ID NO: 134; and
- c. A, Q or R at position 28 of SEQ ID NO: 135.

[0634] Item 140. The polynucleotide system of Item 137, wherein the non-target antigen is a CXCL16 antigen that shares at least 95% identity to SEQ ID NO: 86 and the polymorphism is selected from the group consisting of:

- a. an I or T at position 142 of SEQ ID NO: 86; and
- b. an A or V at position 200 of SEQ ID NO: 86.

[0635] Item 141. The polynucleotide system of Items 136 or 137, wherein the non-target antigen is HLA-A*02.

[0636] Item 142. A vector, comprising the one or more polynucleotides of any one of Items 136-141.

[0637] Item 143. A method of killing a plurality of cancer cell and/or treating cancer in a subject, comprising administering to the subject an effective amount of the immune cell of any one of Items 88-132 or the pharmaceutical composition of any one of Items 46-48.

[0638] Item 144. The use of the immune cell in the method of Item 143, wherein a plurality of cancer cells express the target antigen.

[0639] Item 145. The method of Items 143 or 144, wherein a plurality of cancer cells do not express the non-target antigen.

[0640] Item 146. A method of making a plurality of immune cells, comprising:

- a. providing a plurality of immune cells, and
- b. transforming the plurality of immune cells with the polynucleotide system of any one of Items 136-141, or the vector of Item 142.

[0641] Item 147. A kit comprising the immune cell of any one of Items 88-132 or the pharmaceutical composition of any one of Items 133-135.

[0642] Item 148. The kit of Item 147, further comprising instructions for use.

EXAMPLES

[0643] Throughout the examples, the term "blocker antigen" is used to describe a non-target antigen.

Example 1: Identification of Blocker Targets Lost in Cancer Cells Due to Loss of Heterozygosity

[0644] A bioinformatics pipeline was used to identify candidate blocker targets. The set of human genes was searched for genes with common nonsynonymous variants in extracellular domains that have high loss of heterozygosity (greater than 0.5) in colorectal cancers. Genes with nonsynonymous variants were searched in dbSNP, a database of single nucleotide polymorphisms, that also includes small-scale insertions and deletions along with publication, population frequency, molecular consequence, and genomic mapping information. Common variations were defined as having a minor allele frequency (MAF) of greater than or equal to 0.01 in at least one major population and with at least two unrelated individuals having the minor allele in NCBI. MAF of greater than or equal to 0.1 as criterion for common variations. The focus was on chromosomes 17 and 18, as these chromosomes have high LOH in colorectal cancers. Genes were filtered for membrane proteins, colon expression, and common nonsynonymous variants in the extracellular domain, as described above. A summary of the search process is shown in FIG. 18.

[0645] Additional databases used in this analysis include the following: Uniprot (The Universal Protein Resource), which was used as resource for protein sequence and annotation data hosted by EMBL-EBI, SIB and PIR. GTEx (The Genotype-Tissue Expression) was used as a public resource for tissue-specific gene expression and regulation. It contains samples from 54 non-diseased tissue sites across nearly 1000 individuals. TCGA (The Cancer Genome Atlas) was used as a resource for over 20,000 primary cancer and matched normal samples spanning 33 cancer types. The TCGA-COADREAD dataset is a Colon Adenocarcinoma and Rectum Adenocarcinoma dataset. CCLE (Cancer cell line Encyclopedia) contains information on 57 Colorectal Cancer (CRC) cell lines.

[0646] RNASeqDB is a database of processed data from the GTEx and TCGA using the same pipeline, which allows comparative studies from Memorial Sloan Kettering Cancer Center. 372 TCGA-COADREAD samples and 339 normal colon samples from GTEx were analyzed.

[0647] COLEC12, CXCL16 and APCDD1 were identified as potential blocker targets. Table 14 summarizes the ex-

pression data for these genes in colorectal cancers. Expression data from UCSC Xena browser (for TCGA) and CCLE samples.

Table 14. Expression

Gene name	TCGA-Colorectal Adenocarcinoma (Median-FPKM (383 samples))	Median expression _CCLE_ colorectal_RPKM (57 cell lines)	Colon - Sigmoid	Colon - Transverse
CXCL16	5.7509	25.88884	11.51	15.44
COLEC12	-0.6416	0.01964	27.25	11.24
APCDD1	4.1498	0.58982	12.26	11.22

[0648] Table 15 summarizes the variants and minor allele frequencies.

Table 15. Position, Characteristics and Variation

Gene names	All variants	TCGA-Colorectal Adenocarcinoma_ Frequency of heterozygous deletion (score=-1)	Protein position of Amino Acid Change	Amino acid Change	MAF
CXCL16	rs2277680, rs1050998	0.569805195	1. 200, 2. 142	1. A/V, 2. I/T	1. 0.4615, 2. 0.4633
COLEC12	rs2305025	0.584415584	522	S/P	0.6252
APCDD1	rs1786683	0.600649351	165	Y/S	0.2496

Table 16. LOH Frequencies in Various Cancers

	LOH Freq		
	COLEC12	CXCL16	APCDD1
All cancers	0.23	0.36	0.23
CRC	0.59	0.58	0.6
Lung	0.3	0.58	0.29
Pancreatic	0.3	0.48	0.28
Ovarian	0.39	0.74	0.36
DBCL	0.15	0.23	0.13
Blood	0.06	0.11	0.05
variant	S/P	I/T	Y/S
MAF	0.63	0.46	0.25

Example 2: Identification of Blocker Ligand Binding Domains

[0649] Publicly available antibodies to candidate blocker antigens are sequenced, if CDR sequences are unknown. If no antibodies to candidate blocker targets are available, these antibodies are generated by immunization of mice, rats, or rabbits with purified protein (e.g., COLEC12, CXCL16 and other targets described in the Examples). Sera from immunized animals is used to screen for mAbs for binding to blocker targets. Antibodies to blocker targets are also generated using the huTARG system. Antibodies with the desired specificity are then isolated and sequenced to determine CDR sequences.

[0650] CDR sequence from antibodies to blocker targets are used to generate scFv using standard molecular biology techniques. Candidate scFv are fused to inhibitory receptor hinge or transmembrane domains to generate inhibitory receptors using standard molecular biology techniques. Candidate scFv are also fused to activator receptor hinge or

transmembrane domains (e.g., CAR) to generate full length activator receptors to use as a positive control for scFv binding to target antigens. The ability of candidate scFv to work in the context of an inhibitory receptor is assayed in Jurkat cells using the NFAT-luciferase reporter assay.

5 **Example 3: Methods for Jurkat and Primary T Cell Activation Experiments**

Cell culture

10 **[0651]** Jurkat cells encoding an NFAT Luciferase reporter were obtained from BPS Bioscience. In culture, Jurkat cells were maintained in RPMI media supplemented with 10% FBS, 1% Pen/Strep and 0.4mg/mL G418/Geneticin. All other cell lines used in this study were obtained from ATCC, and maintained as suggested by ATCC.

Jurkat cell transfection

15 **[0652]** Jurkat cells were transiently transfected via 100uL format Neon electroporation system (Lonza) according to manufacturer's protocol using the following settings: 3 pulses, 1500V, 10 msec. Cotransfection was performed with 1-3ug of activator CAR or TCR construct and 1-3ug of blocker constructs or empty vector per 1e6 cells and recovered in RPMI media supplemented with 20% heat-inactivated FBS and 0.1% Pen/Strep.

20 Jurkat-NFAT-luciferase activation studies

[0653] Jurkat cells were resuspended in 15uL of RPMI supplemented with 10% heat-inactivated FBS and 0.1% Pen/Strep, added to the peptide-loaded beads and co-cultured for 6 hours. ONE-Step Luciferase Assay System (BPS Bioscience) was used to evaluate Jurkat luminescence. Assays were performed in technical duplicates.

25

Primary T cell transduction, expansion, and enrichment

30 **[0654]** Frozen PBMCs were thawed in 37°C water bath and cultured at 1e6 cells/mL in LymphoONE (Takara) with 1% human serum and activated using 1:100 of T cell TransAct (Miltenyi) supplemented with 300 IU/ml IL-2. After 24 hours, lentivirus was added to PBMCs at a MOI of 5. PBMCs were cultured for 2-3 additional days to allow cells to expand under TransAct stimulation. Post expansion, activator and blocker transduced primary T cells were enriched using anti-PE microbeads (Miltenyi) according to manufacturer's instructions. Briefly, primary T cells were incubated with EGFR-Fc (R&D Systems) at 1:25 dilution for 30 minutes at room temperature in MACS buffer (0.5% BSA + 2mM EDTA in PBS). Cells were washed 3 times in MACS buffer and incubated in secondary antibody (1:1200) for 30 minutes at room temperature in MACS buffer. Cells were then incubated in anti-PE microbeads and passed through the LS column (Miltenyi).

35

Primary T cell in vitro cytotoxicity studies

40 **[0655]** For cytotoxicity studies with pMHC targets, enriched primary T cells were incubated with HeLa or HCT116 target cells that express GFP or RFP. For "tumor" cells, WT HeLa or HLA-A*02 knocked-out HCT116 were used and for "normal" cells, HLA-A*02 transduced HeLa or WT HCT116 were used. The cocultures were imaged using an IncuCyte live cell imager and target cell fluorescence area was used to quantify live target cells.

45 **Example 4: HLA-A*02 Inhibitory Receptor Blocks EGFR-Mediated Activation of Jurkat Cells**

[0656] The ability of an inhibitory receptor with an HLA-A-A* 02 antigen binding domain and a LIR-1 ICD (C1765 (not according to the invention) or C2162 (according to the invention)) to block activation of Jurkat cells expressing an activator CAR with an EGFR antigen binding domain (e.g. CT479 (not according to the invention), CT486 (not according to the invention), or CT489 (according to the invention)) was assayed using the NFAT-luciferase reporter system as previously described. Normal HeLa tumor cells, which were EGFR+ and HLA-A*02-, were used as target cells. EGFR+/ HLA-A*02- HeLa cells were also transduced with a polynucleotide encoding HLA-A*02+ to generate EGFR+ /HLA-A*02+ HeLa cells to use as target cells expressing both activator and blocker antigens.

50

[0657] As shown in FIG. 1A, expression of the HLA-A*02 LIR-1 blocker (C1765, not according to the invention) in Jurkat cells expressing the EGFR CAR (CT479, not according to the invention) shifts the CAR E_{MAX} by >5 fold compared to the CAR E_{max} of Jurkat cells that do not express the blocker.

55

[0658] Furthermore, lower blocking was observed with lower HLA-A2 expression levels on target cells. Normal HCT116 cells are EGFR+ and HLA-A*02+. The levels of EGFR and HLA-A*02 were assayed in HCT116 cells and HeLa cells

transduced with polynucleotides encoding the HLA-A*02 polynucleotides using an anti-EGFR antibody and an anti-HLA-A*02 antibody (BB7.2) followed by FACs sorting. As shown in FIGS. 2A & 2B, HCT116 cells have lower levels of blocker HLA-A*02 antigen than transduced HeLa cells. When Jurkat cells expressing the EGFR CAR and HLA-A*02 LIR-1 blocker was presented with HCT116 target cells expressing EGFR and HLA-A*02 antigens, presence of the HLA-A*02 LIR-1 blocker shifted the E_{MAX} of the EGFR CAR 1.8 fold (FIG. 3B). In contrast, transduced HeLa cells, which expressed a higher level of HLA-A*02 antigen, were able to mediate an EGFR CAR E_{MAX} shift of >5 fold (FIG. 1B, FIG. 17B, and FIG. 3D). As a control, there was minimal activation by EGFR knockout HCT116 cells (FIG. 3A).

[0659] The ratio of blocker to activator necessary to achieve 50% blocking using the EGFR CAR and HLA-A*02 LIR-1 blocker was assayed using a bead based system, which is shown in FIG. 4A and FIG. 4B.

[0660] To determine the EC50 of the activator antigen, activator beads were coated with activator antigen at different concentrations to determine EC50. An irrelevant protein was added to each concentration so that the total protein concentration was the same, and a constant amount of beads was added to Jurkat effector cells expressing the EGFR CAR (FIG. 4A).

[0661] To determine blocker antigen IC50, beads were coated with activator antigen at the EC50 concentration (determined in FIG. 4A), and coated with blocker antigen at different concentrations. An irrelevant protein was added at each concentration so that the total protein concentration remained the same, and a constant amount of beads was added to Jurkat effector cells expressing either the EGFR CAR or the EGFR CAR and the HLA-A*02 LIR-1 blocker (FIG. 4B).

Example 5: Characterization of Specific Receptor Pairs

[0662] T cells transfected with either an EGFR scFv CAR activator (CT-479 (not according to the invention), CT-482 (not according to the invention), CT-486 (not according to the invention), CT-487 (not according to the invention), CT-488 (not according to the invention), or CT-489 (according to the invention), as indicated in FIGS. 5 and 7-9), or with EGFR scFv CAR activator and an HLA-A*02 PA2.1 scFv LIR1 inhibitor (C1765 (not according to the invention) or C2162 (according to the invention)) were co-cultured with HeLa target cells. Normal HeLa cell lines express EGFR, but not HLA-A*02, were transduced to express the HLA-A*02 inhibitory receptor target. Cells were co-cultured at a 1:1 ratio of effector to target (E:T). In the lower right of FIG. 5, effector cell receptor expression is indicated first, while HeLa cell expression is in parentheses. As can be seen from FIG. 5, a different degree of blocking is observed when the same HLA-A*02 PA2.1 scFv LIR1 inhibitor was used with different EGFR activator receptors.

[0663] The ability of the HLA-A*02 PA2.1 scFv LIR1 inhibitor (C1765, not according to the invention) to block EGFR scFv CAR activation of T cells co-cultured with HCT. 116 cells expressing either EGFR (Target A) or EGFR and HLA-A*02 (Target AB) was also assayed (FIG. 10). As with HeLa cells, different degree of blocking is observed when the same HLA-A*02 PA2.1 scFv LIR1 inhibitor was used with different EGFR activator receptors.

[0664] The ability of the HLA-A*02 PA2.1 scFv LIR1 inhibitor (C1765, not according to the invention) to block EGFR scFv CAR activation of Jurkat cells when Jurkat cells were co-cultured with HeLa target cells was also assayed. The results are shown in FIG. 2. Similar to the results in primary T cells, a different degree of blocking is observed when the same HLA-A*02 PA2.1 scFv LIR1 inhibitor was used with different EGFR activator receptors.

[0665] Immune cell activation was measured in immune cells expressing an activator or a specific activator/blocker pair in the presence of a target cell expressing only an activating ligand or a target cell expressing both an activator ligand and a blocking ligand (FIG. 6A). Selectivity ratios (SR) were calculated for tested pairs of CAR activators and blockers. SR is determined by calculating the ratio of the activation of an immune cell expressing an activator/blocker pair in the presence of a target cell expressing only an activating ligand to the activation of an immune cell expressing the activator/blocker pair in the presence of a target cell expressing both an activator ligand and a blocking ligand. Selected EGFR targeting CAR activators were expressed as a pair with either a humanized blocker (C2162, according to the invention) or a mouse blocker (C1765, not according to the invention). Among the EGFR targeting CAR activators paired with the humanized blocker, the selectivity ratio for CT489 (according to the invention) reached 24.2; the selectivity ratio for CT486 (not according to the invention) reached 17; the selectivity ratio for CT487 (not according to the invention) reached 5.0; and the selectivity ratio for CT487 (not according to the invention) reached 5.0. Among the EGFR targeting CAR activators paired with the mouse blocker, the selectivity ratio for CT489 (according to the invention) was 21.6; the selectivity ratio for CT486 (not according to the invention) was 15.7; the selectivity ratio for CT487 (not according to the invention) was 18.4; and the selectivity ratio for CT487 (not according to the invention) was 21.2 (FIG. 6A). The maximum specific killing was also determined for each of the EGFR targeting activator and inhibitory receptors pairs. For all pairings, the specific killing exceeded about 60%, indicating that choice of blocker did not significantly impact T cell activation by the EGFR targeting activator receptors. However, the pairing of a specific activator and specific blocker pair had an impact on the selectivity ratio, indicating that off-target killing by facilitated a specific activator/inhibitory receptor pair is not predictable.

[0666] Activator and blocker sensitivity was determined in immune cells expressing the CAR activator alone or the

CAR activator and either a humanized blocker or a mouse blocker (FIG. 6B). To determine activator sensitivity immune cells were incubated in the presence of EGFR negative target cells. Prior to incubation, target cells were titrated with EGFR mRNA, resulting in increasing levels EGFR expression (FIG. 6B, left panel). To determine blocker sensitivity immune cells were incubated in the presence of HLA-A*02 negative target cells that express EGFR (about 53,000 EGFR molecules per cell). Prior to incubation, target cells were titrated with HLA-A*02 mRNA, resulting in increasing levels HLA-A*02 expression (FIG. 6B, right panel). EC₅₀ values were calculated for each CAR activator and the blocker in combination with different CAR activators (Table 5.1). The results further confirm that the specific activator receptor has an impact on inhibitory receptor sensitivity.

Table 5.1

Activator	Activator EC ₅₀ (ng)	Mouse Blocker EC ₅₀ (ng)	Human Blocker EC ₅₀ (ng)
CT486	5,300	4,700	4,000
CT487	3,500	3,500	2,900
CT488	4,700	4,700	3,500
CT489	3,100	4,200	3,200

Example 6: Inhibitory Receptors Reversibly Decrease Surface Level of Activator Receptors in T Cells

[0667] Primary T cells from two HLA-A*02 negative donors were transduced with an EGFR scFv CAR activator (CT-479 (not according to the invention), CT-482 (not according to the invention), CT-486 (not according to the invention), CT-487 (not according to the invention) or CT-488 (not according to the invention)) and an HLA-A*02 PA2.1 scFv LIR1 inhibitor (C1765, not according to the invention). Transduced cells were enriched by FACS sorting on the blocker and activator receptors, or by double column purification on the blocker and activator receptors. Transduced T cells were co-cultured with HeLa target cells. Normal HeLa cell lines express EGFR, but not HLA-A*02, but were transduced to express the HLA-A*02 inhibitory receptor target. Cells were co-cultured at a 1:1 ratio of effector to target (E:T). Surface expression of the EGFR CAR activator was assayed after 120 hours using labeled peptides that bound the activator and inhibitory receptors, and fluorescence activated cell sorting. The change in activator surface level following co-culture with HeLa cells expressing both activator and blocker ligands corresponded to the T cells' ability to kill target cells (compare FIG. 5 and FIG. 11A).

[0668] T cells expressing the CT-482 EGFR scFv CAR activator (not according to the invention) and HLA-A*02 PA2.1 scFv LIR1 inhibitor (C1765, not according to the invention) combination, were co-cultured with HeLa cells expressing EGFR (Target A), HLA-A* 02 (Target B), a combination of EGFR and HLA-A* 02 on the same cell (Target AB), a mixed population of HeLa cells expressing Target A and Target AB on different cells, or a mixed population of HeLa cells expressing Target B and Target AB on different cells (FIGS. 12A-12B). T cells were cultured with HeLa target cells at a ratio of 1:1 effector cell to target cell. When T cells were co-cultured with a Target A plus Target AB population of HeLa cells, levels of activator decreased, then recovered (FIG. 12A). Furthermore, the activator and blocker antigens must be present together on the same cell to trigger activator surface expression loss on effector T cells. In contrast to the activator, blocker expression was largely unchanged (FIG. 12B).

[0669] FIG. 13 shows a schematic for an experiment to determine if loss of cytotoxicity, blocking activity, and expression of activator receptor by T cells was reversible. T cells expressing EGFR scFv CAR activator receptor (CT-487, not according to the invention) and HLA-A*02 PA2.1 scFv LIR1 (C1765, not according to the invention) inhibitory receptor were co-cultured with HeLa target cells expressing both the activator and inhibitory receptor targets (AB). Following 3 days co-culture, HeLa cells were removed using an anti-EGFR column, and the T cells were either stained for the activator and inhibitory receptors, or co-cultured with HeLa cells expressing EGFR activator target only for an additional 3 days. After the additional 3 days co-culture, HeLa cells were again removed using an anti-EGFR column, and the T cells were either stained for the activator and inhibitory receptors, or were co-cultured for an additional 3 days with HeLa cells expressing EGFR activator target only, or both the activator and blocker targets (AB), before staining. T cells were assayed for the presence of activator and inhibitory receptors (stained) using labeled EGFR and A2 probes, and the levels of receptor expression were quantified using fluorescence activated cell sorting. Results of the experiment are shown in FIGS. 14A-14B. As shown in FIGS. 14A-14B, co-culture of T cells with HeLa cells expressing both activator and inhibitor targets reduces EGFR activator staining (FIGS. 14A-14B, left panel). When T cells are co-cultured with HeLa cells expressing activator (Target A only) at round 2, expression of EGFR activator increases. Thus, activator surface loss is reversible and tracks with T cell cytotoxicity.

[0670] FIGS. 15A-15B show the results of an additional experiment showing that loss of activator receptor expression is rapidly reversible. Activator CAR cell surface expression is diminished in T cells expressing both activator and inhibitory

receptors when these T cells were co-cultured with model "normal" cells expressing both activator and inhibitor ligand. However, when T cells were switched to co-culture with model "cancer" cells that only express the activator ligand, the T cells were able to quickly (within hours) restore cell surface expression of the activator CAR. The absence of model "normal" cells expressing both activator and inhibitor ligand could also restore cell surface expression of the activator CAR, regardless of presence of tumor cells, within similar time frame.

Example 7: HLA-A*02 LIR-1 based inhibitory receptors can block activation by an EGFR CAR activator

[0671] Activation of Jurkat effector cells expressing an EGFR CAR activator and a pMHC HLA-A*02 scFv LIR-1 based inhibitory receptor (comprising a LIR-1 hinge, transmembrane and ICD) was assayed using the NFAT Luciferase assay described in Example 3.

[0672] Jurkat cells were transfected with activator and inhibitory receptor DNA, and activation was assayed in a cell-free bead-based assay (FIG. 16). Beads were loaded with either activator antigen, blocker antigen, or activator and inhibitor antigens, and the ratio of beads to Jurkat cells was varied. In the cell-free bead based assay, the HLA-A*02 scFv LIR-1 based inhibitory receptor was able to block activation of the Jurkat cells when cells were contacted with beads carrying the HLA-A*02 blocker and EGFR activator in *cis*, but not when the HLA-A*02 blocker and EGFR activator were in *trans* (on different beads). Presence of the HLA-A*02 blocker on the beads was able to shift E_{MAX} of EGFR CAR by greater than or equal to 9X (FIG. 16).

[0673] Activation of Jurkat cells expressing the EGFR CAR activator and an HLA-A*02 scFv LIR-1 based inhibitory receptor was also assayed using HeLa and SiHa cells as target cells. Normal HeLa and SiHa cell lines express EGFR, but not HLA-A*02 (SiHa WT and HeLa WT), but were transduced to express the HLA-A*02 inhibitory receptor target (SiHa A02 and HeLa A02). As can be seen in FIG. 17A and FIG. 17B, the HLA-A*02 scFv LIR-1 based inhibitory receptor was able to shift the EGFR E_{MAX} by greater than 4X using SiHa target cells (FIG. 17A) and greater than 5X using HeLa target cells (FIG. 17B).

Example 8: In vivo efficacy of EGFR CAR activator and inhibitory receptor expressing T cells

[0674] Efficacy and tumor vs. normal cell selectivity for immune cells expressing activator receptors (EGFR CAR) and inhibitory receptor (Blocker) pairs are established in an in vivo environment (FIG. 19). A murine xenograft model in NSG (NOD.Cg-Prkdc^{scid}Il2rg^{tm1Wjl}/SzJ) using human EGFR+/HLA-A*02+ HeLa cells representing "normal" cells and isogenic EGFR+/HLA-A*02(-) knockout HeLa cells as "tumor" cells. The in vivo therapeutic window is established by comparing the effect of the activator/inhibitory receptor pairs on normal vs. tumor cells. These data provide in vivo support for an on-target, off-tumor therapeutic window. Optimal doses for activator/inhibitory receptor pairs are established and optimized for sensitivity and selectivity.

[0675] The in vivo model involves subcutaneous xenografts of "normal" cells in the right flank and "tumor" cells in the left flank. Immune cells expressing the activator receptor, an activator/inhibitory receptor pair, or untransduced control cells are administered to the animals at a range of time points.

[0676] Animals are monitored for general health via clinical observations and effects on body weight. Tumor xenotransplants are assessed for size by caliper and imaging. At each time point, human T cells are enumerated by flow cytometry and assessed for surface markers including CD3, CD4, CD8, activator receptor expression, and inhibitory receptor expression. Serum cytokine levels are determined, including IFN γ and interleukin-2 (IL-2). In post-life analysis, protocol-specified tissues, including the injection site and the tumor, are assessed by histology for tumor infiltrating T-cells and activator receptor antigen (e.g. EGFR) and blocker antigen expression (e.g. HLA-A*02) by an ACVP board-certified pathologist.

Example 9: B7 and HLA-A*11 are inhibitory receptor ligands (not according to the invention)

[0677] An HLA-A*011 inhibitory receptor was created by fusing scFv #4 (Table) to a hinge, TM and ICD derived from LIR-1. Gene segments were combined using Golden Gate cloning and inserted downstream of a human EF1 α promoter contained in a lentivirus expression plasmid.

[0678] Jurkat NFAT-Luciferase effector cells were transformed as described above with a CAR activator, or a CAR activator and the scFv #4 HLA-A*11 inhibitory (blocker) receptor. Sequences of the HLA-A*011 inhibitory receptor are shown in Table 9.1, below and co-cultured with HeLa cells expressing both the activator ligand and HLA-A*011:01. NFAT-Luciferase assays were carried out using HeLa instead of T2 target cells.

[0679] Results are shown in FIG. 20A and FIG. 20B. As shown in FIG. 20A, scFv #4, when fused to LIR-1's hinge, TM and ICD domains, is capable of inhibiting activation of Jurkat effector cells co-cultured with target cells expressing both activator ligand and HLA-A* 011.

Table 9.1. HLA-A*011 inhibitory receptor sequences.

Name	Protein Sequence	DNA Sequence
5 HLA-A*11 Blocker 4	MDMRVPAQLL GLLLLWLRGA RCEVQLVESGG <u>GLVQPGGSLRLS</u> <u>CAASGFTFSSYW</u> 10 <u>MHWVRQAPGK</u> <u>GLVWVSRINSDG</u> <u>SSTSYADSVKGR</u> <u>FTISRDNANKNTL</u> 15 <u>YLOMNSLRAED</u> <u>TAVYYCCLGVL</u> <u>LYNWFDPPWGQG</u> <u>TLVTVSSGGGG</u> SGGGGSGGGGS GGDIQMTQSPSS 20 LSASVGDRVIT CRASQSISSYL WYQQKPGKAPK LLIYAASSLQSG 25 VPSRFSGSGSGT DFTLTISLQPED FATYYCQSYST PLTFGGGTKVEI KYGSQSSKPYLL 30 THPSDPLELVVS GPSGGPSSPTTGP TSTSGPEDQPLTP TGSDPQSGLGRH 35 LGVVIGILVAVIL LLLLLLLLFLILR HRRQGKHWTST QRKADFQHPAG AVGPEPTDRGLQ 40 WRSSPAADAQE ENLYAAVKHTQ PEDGVEMDTRSP HDEDPAVITYA EVKHSRPRREM 45 ASPPSPLSGEFLD TKDRQAEEDRQ MDTEAAASEAP QDVTYAQLHSL TLRREATEPPPS 50 QEGPSPAVPSIY ATLAIH (SEQ ID NO: 915)	ATGGACATGAGGGTCCCCGCTCAGCTCCTGGGGCTC CTGCTACTCTGGCTCCGAGGTGCCAGATGTGAAGTG CAGCTGGTGGAAAGCGGCGGAGGCCTGGTGCAGCC TGGCGGCAGCCTGAGACTGTCTTGCGCCGCCAGCGG CTTCACCTTCAGCAGCTACTGGATGCACTGGGTCCG CCAGGCCCCTGGCAAGGGACTGGTCTGGGTGTCTCG AATCAACAGCGACGGCAGCAGCACCAGCTACGCCG ACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGAC AACGCCAAGAACACCCTGTACCTGCAGATGAACAG CCTGCGGGCCGAGGACACCGCCGTGTATTACTGTTG TTTGGGTGTTTTATTATACAACCTGGTTCGACCCCTGG GGCCAGGGAACCCTGGTCACCGTGTCTCAGGCGG AGGTGGAAGCGGAGGGGGAGGATCTGGCGGCGGAG GAAGCGGAGGCGACATCCAGATGACCCAGTCTCCA TCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC ATCACTTGCCGGGCAAGTCAGAGCATTAGCAGCTAT TTAAATTGGTATCAGCAGAAACCAGGGAAAGCCCC TAAGCTCCTGATCTATGCTGCATCCAGTTTGCAAAG TGGGGTCCCATCAAGGTTCAAGTGGCAGTGGATCTGG GACAGATTTCACTCTCACCATCAGCAGTCTGCAACC TGAAGATTTTGCAACTTACTACTGTCAACAGAGTTA CAGTACCCCTCTCACTTTTCGGCGGCGGAACAAAGGT GGAGATCAAGTACGGCTCACAGAGCTCCAAACCCT ACCTGCTGACTCACCCAGTGACCCCTGGAGCTCG TGGTCTCAGGACCGTCTGGGGGGCCCCAGCTCCCCGA CAACAGGCCCCACCTCCACATCTGGCCCTGAGGACC AGCCCCCTACCCCCACCGGGTCGGATCCCCAGAGTG GTCTGGGAAGGCACCTGGGGGTTGTGATCGGCATCT TGGTGGCCGTATCCTACTGCTCCTCCTCCTCCTCCT CCTCTTCCTCATCCTCCGACATCGACGTCAGGGCAA ACACTGGACATCGACCCAGAGAAAGGCTGATTTCC AACATCCTGCAGGGGCTGTGGGGCCAGAGCCCACA GACAGAGGCCTGCAGTGGAGGTCCAGCCCAGCTGC CGATGCCCAGGAAGAAAACCTCTATGCTGCCGTGA AGCACACACAGCCTGAGGATGGGGTGGAGATGGAC ACTCGGAGCCACACGATGAAGACCCCCAGGCAGT GACGTATGCCGAGGTGAAACACTCCAGACCTAGGA GAGAAATGGCCTCTCCTCCTTCCCCACTGTCTGGGG AATTCCTGGACACAAAGGACAGACAGGCGGAAGAG GACAGGCAGATGGACACTGAGGCTGCTGCATCTGA AGCCCCCAGGATGTGACCTACGCCCAGCTGCACAG CTTGACCCTCAGACGGGAGGCAACTGAGCCTCCTCC ATCCCAGGAAGGGCCCTCTCCAGCTGTGCCCAGCAT CTACGCCACTCTGGCCATCCAC (SEQ ID NO: 916)

55 **[0680]** Jurkat NFAT luciferase (JNL) cell activation was measured in an mRNA titration assay using HeLa cells transfected with mRNA. HeLa cells were transfected with serially-diluted mRNA encoding HLA-B*07:02. JNL cells were transiently transfected with the EGFR activating CAR and HLA-B*07 blocker (circle) or negative control vector (triangle). The functional response was assessed after a 6 hour co-culture. The results shown in FIG. 21 demonstrate that HLA-

B*07 blocker blocks activation of the EGFR targeting CARs in the presence of titrated HLA-B*07:02, but not in the absence of blocker.

Example 10: Characterization of EGFR activator and HLA-A*02 inhibitory receptor expression in primary T cells

[0681] To stain for EGFR activator, primary T cells were incubated with EGFR-Fc (R&D Systems) at 1:25 dilution for 30 minutes at room temperature in FACS buffer (1% BSA in PBS). Cells were washed 2 times in FACS buffer and incubated in anti-Fc (1:100) for 30 minutes at room temperature in FACS buffer. To stain for HLA-A*02 inhibitory receptor, T cells were incubated with HLA-A*02 tetramer probe for 30 minutes at room temperature in FACS buffer. Receptor expressions were analyzed by flow cytometry after staining.

[0682] Primary T cells were transduced with polynucleotides encoding an EGFR targeting activator receptor and either a human or mouse HLA-A*02 targeting inhibitory receptor. Several specific pairs of EGFR targeting activators and HLA-A*02 targeting inhibitory receptors were tested. Transduction was performed with either a single vector encoding both the activator receptor and inhibitory receptor or two vectors, one encoding the activator receptor and one encoding the inhibitory receptor. Cells were analyzed with and without enrichment to determine the percentage of cells expressing both the activator and inhibitor receptor (FIG. 22).

[0683] In primary T cells transduced with two vectors, one encoding the activator receptor and one encoding the inhibitory receptor, the percentage of cells in the transduced T cell population expressing both receptors was about 22.5% (FIG. 22A, left panel) prior to enrichment. Following enrichment, the percentage of cells in the transduced T cell population expressing both receptors increased to about 91.6%.

[0684] In primary T cells transduced with the single vector encoding both the activator receptor and inhibitory receptor, the percentage of cells in the transduced T cell population expressing both receptors ranged between about 6.29% and 13.2% prior to enrichment. Following enrichment, the percentage of cells expressing both receptors ranged from 22.6% to 82.8%. The percentage of cells expressing both receptors was dependent on the specific activator and inhibitory receptor pair.

Example 11: Characterization of immune cell activation selectivity in primary T cells expressing an activator and inhibitory receptor

[0685] For selective cytotoxicity studies, enriched primary T cells were incubated with HeLa or HCT116 target cells that express GFP or RFP. For "tumor" cells, WT HeLa or HLA-A*02 knocked-out HCT116 were used and for "normal" cells, HLA-A*02 transduced HeLa or WT HCT116 were used. The cocultures were imaged using an IncuCyte live cell imager and target cell fluorescence area was used to quantify live target cells (FIG. 23A). For measuring selective cytotoxicity in mixed target culture, RFP expressing "tumor" WT HeLa cells and GFP expressing "normal" HeLa cells transduced with HLA-A*02 were mixed at a 1:1 ratio. Enriched primary T cells were added to the mixed target culture and imaged using an IncuCyte live cell imager and target cell fluorescence area was used to quantify live target cells (FIG. 23B).

[0686] FIG. 23A is a series of plots showing the percentage of specific killing of HeLa target cells or HCT116 target cells by immune cells expressing an EGFR activator receptor and an HLA-A*02 scFv LIR1 inhibitor receptor. The target cells express the EGFR antigen (A), the HLA-A*02 non-target antigen (B), or both (AB). Immune cells expressing the activator/inhibitor receptor pair were incubated with target cells at a 1:1 ratio.

[0687] FIG. 23B is a plot and series of images showing specific and selective killing of target cells by immune cells expressing an EGFR activator receptor and an HLA-A*02 scFv LIR1 inhibitor receptor in a mixed culture of cancer cells and normal cells. UTD = immune cells no activator or inhibitor receptors; CAR = immune cells expressing only activator receptor; Tmod = immune cells expressing both activator and inhibitor receptors.

Example 12: Characterization of immune cell activation selectivity in Jurkat cells expressing an activator and inhibitory receptor

[0688] FIG. 24 is a chart showing immune cell activation in Jurkat cells (effectors) expressing EGFR activator receptor (CAR) or EGFR activator receptor and HLA-A*02 inhibitor receptor (Tmod). The Jurkat cells were incubated with a panel of target cell lines that express HLA-A*02 either without EGFR expression or express HLA-A*02 with EGFR expression. EGFR negative cell lines were generated by knocking out EGFR using CRISPR/Cas9. Tmod Jurkat effector cells showed no off-target activation when cocultured with EGFR negative cells.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 63068249 [0001]
- US 63105639 [0001]
- US 63230632 [0001]
- WO 2020065406 A2 [0004]
- US 2020045250 W [0159]
- US 2020045228 W [0226]
- US 2020064607 W [0226]
- US 2021029907 W [0226]
- US 2020059856 W [0226]
- WO 0196584 A [0302]
- WO 0129058 A [0302]
- US 6326193 B [0302]
- US 5350674 A [0310]
- US 5585362 A [0310]
- US 6352694 B [0329]
- US 6534055 B [0329]
- US 6905680 B [0329]
- US 6692964 B [0329]
- US 5858358 A [0329]
- US 6887466 B [0329]
- US 6905681 B [0329]
- US 7144575 B [0329]
- US 7067318 B [0329]
- US 7172869 B [0329]
- US 7232566 B [0329]
- US 7175843 B [0329]
- US 5883223 A [0329]
- US 6905874 B [0329]
- US 6797514 B [0329]
- US 6867041 B [0329]
- US 10040846 B [0329]
- US 20060121005 [0329]
- US 20040101519 [0331]
- US 20060034810 A [0331]
- US 20140068797 A [0364]
- US 8871445 B [0364]
- US 8865406 B [0364]
- US 8795965 B [0364]
- US 8771945 B [0364]
- US 8697359 B [0364]
- US 20190185849 A1 [0378]
- US 20050244858 A, Rossi [0406]
- US 20030170238, Gruenberg [0462]
- US 4690915 A, Rosenberg [0462]

Non-patent literature cited in the description

- KARLIN ; ALTSCHUL. *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 2264-2268 [0079]
- ALTSCHUL et al. *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0079]
- KARLIN ; ALTSCHUL. *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 5873-5877 [0079]
- ALTSCHUL et al. *Nucleic Acids Res.*, 1997, vol. 25, 3389-3402 [0079]
- WOOTTON ; FEDERHEN. *Computers and Chemistry*, 1993, vol. 17, 149-163 [0079]
- HARLOW et al. *Using Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1999 [0116]
- HARLOW et al. *Antibodies: A Laboratory Manual*. Cold Spring Harbor, 1989 [0116]
- HOUSTON et al. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 5879-5883 [0116]
- BIRD et al. *Science*, 1988, vol. 242, 423-426 [0116]
- KROGH et al. *Journal of Molecular Biology*, 2001, vol. 305 (3), 567-580 [0170]
- HOFMANN ; STOFFEL. *Biol. Chem.*, 1993, vol. 347, 166 [0170]
- SAMBROOK et al. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory, 2001 [0302]
- UI-TEI et al. *FEBS Letters*, 2000, vol. 479, 79-82 [0307]
- BERG et al. *Transplant Proc.*, 1998, vol. 30 (8), 3975-3977 [0330]
- HAANEN et al. *J. Exp. Med.*, 1999, vol. 190 (9), 1319-1328 [0330]
- GARLAND et al. *J. Immunol Meth.*, 1999, vol. 227 (1-2), 53-63 [0330]
- WANG D et al. *Stem Cells Transl Med.*, 2015, vol. 4, 1234-1245 [0351]
- REN et al. *Clin. Cancer Res.*, 2017, vol. 23, 2255-2266 [0351]
- PENNISI. *Science*, 2013, vol. 341, 833-836 [0361]
- CONG. *Science*, 2013, vol. 339, 819-823 [0364]
- TSAI. *Nature Biotechnol.*, 2014, vol. 32 (6), 569-576 [0364]
- DOENCH et al. *Nat Biotechnol.*, 2016, vol. 34, 184-191 [0380]

Patentkrav**1. Immuncelle, omfattende:**

- 5 a. en første receptor, omfattende et ekstracellulært ligandbindingsdomæne specifikt for epidermal vækstfaktorreceptor (EGFR); og
- b. en anden receptor, omfattende et ekstracellulært ligandbindingsdomæne specifikt for et ikke-målantigen tabt i EGFR+-kræft på grund af tab af heterozygositet,
- 10 hvor den første receptor er en aktivatorreceptor responsiv over for EGFR; og hvor den anden receptor er en inhiberende receptor responsiv over for ikke-målantigenet, hvor det ekstracellulære ligandbindingsdomæne af den anden receptor specifikt binder HLA-A*02, og
- 15 hvor det ekstracellulære ligandbindingsdomæne af den første receptor omfatter komplementaritetsbestemmende regioner (CDR'er) CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2 og CDR-H3 af SEQ ID NOs: 154, 155, 156, 151, 152 og 153, og
- 20 hvor det ekstracellulære ligandbindingsdomæne af den anden receptor omfatter komplementaritetsbestemmende regioner CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, CDR-H3 af den humaniserede scFv-bindingsdomænesekvens af C-002162 PA2.1.14 som afbildet i tabel 4, hvor CDR-L1-sekvensen er RSSQSIVHSNGNTYLE (SEQ ID NO: 101), CDR-L2-sekvensen er KVSNRFSGVPAR, CDR-L3-sekvensen er FQGSHVPRT (SEQ ID NO: 103), CDR-H1-sekvensen er ASGYTFTSYHIH (SEQ ID NO: 104), CDR-H2-sekvensen er WIYPGNVNTEYNEKFKGK (SEQ ID NO: 105), og CDR-H3-sekvensen er EEITYAMDY (SEQ ID NO: 106),
- 25 hvor immuncellen er en CD8+ CD4- T-celle eller en CD8- CD4+ T-celle.

2. Immuncellen ifølge krav 1,

- hvor det ekstracellulære ligandbindingsdomæne af den anden receptor omfatter
- 30 SEQ ID NO: 93; eller en sekvens med mindst 85%, mindst 90%, mindst 95%, mindst 97% eller mindst 99% identitet dermed.

3. Immuncellen ifølge et hvilket som helst af kravene 1 eller 2, hvor den første receptor er en kimærisk antigenreceptor (CAR);

hvor det ekstracellulære ligandbindingsdomæne af den første receptor omfatter en variabel tung (VH) del med en sekvens valgt fra VH-sekvensen angivet i SEQ ID NO: 149; og/eller en variabel let (VL) del omfattende en sekvens angivet i SEQ ID NO: 150; eller sekvenser med mindst 70%,
 5 mindst 85%, mindst 90% eller mindst 95% identitet dermed; og/eller hvor det ekstracellulære ligandbindingsdomæne af den første receptor omfatter scFv-sekvensen af SEQ ID NO: 18; eller en sekvens med mindst 85%, mindst 90%, mindst 95%, mindst 97% eller mindst 99% identitet dermed.

10

4. Immuncellen ifølge et hvilket som helst af kravene 1-3, hvor den anden receptor omfatter et LILRB1-intracellulært domæne eller en funktionel variant deraf, eventuelt hvor det LILRB1-intracellulære domæne omfatter en sekvens mindst 90%, mindst 95%, mindst 97%, mindst 99% eller er identisk med SEQ ID
 15 NO: 129;

hvor den anden receptor omfatter et LILRB1-transmembrandomæne eller en funktionel variant deraf, eventuelt hvor LILRB1-transmembrandomænet eller en funktionel variant deraf omfatter en sekvens mindst 90%, mindst 95%, mindst 97%, mindst 99% eller er identisk med SEQ ID NO: 133;

20

og/eller hvor den anden receptor omfatter et LILRB1-hængselsdomæne eller funktionel variant deraf, eventuelt hvor LILRB1-hængselsdomænet omfatter en sekvens mindst 90%, mindst 95%, mindst 97%, mindst 99% eller er identisk med SEQ ID NO: 132, SEQ ID NO: 125 eller SEQ ID NO:
 25 126.

5. Immuncellen ifølge et hvilket som helst af kravene 1-4, hvor den anden receptor omfatter et LILRB1-intracellulært domæne, et LILRB1-transmembrandomæne, et LILRB1-hængselsdomæne, en funktionel variant af en hvilken som
 30 helst af disse, eller kombinationer deraf, eventuelt hvor det LILRB1-intracellulære domæne og LILRB1-transmembrandomænet omfatter SEQ ID NO: 128 eller en sekvens mindst 95% identisk med SEQ ID NO: 128.

6. Immuncellen ifølge et hvilket som helst af kravene 1-5, hvor ekspression
 35 og/eller funktion af et gen fra MHC klasse I er blevet reduceret eller elimineret,

eventuelt hvor genet fra MHC klasse I er beta-2-mikroglobulin (B2M); og/eller hvor immuncellen omfatter et polynukleotid omfattende et interfererende RNA, idet det interfererende RNA omfatter en sekvens komplementær til en sekvens af et B2M mRNA (SEQ ID NO: 172), eventuelt hvor det interfererende RNA er i stand
5 til at inducere RNAi-medieret nedbrydning af B2M mRNA'et.

7. Immuncellen ifølge krav 6, hvor det interfererende RNA er et kort hårnåle-RNA (shRNA), og eventuelt hvor shRNA'et omfatter:

- a. en første sekvens, med fra 5'-ende til 3'-ende en sekvens
10 komplementær til en sekvens af B2M mRNA'et; og
- b. en anden sekvens, med fra 5'-ende til 3'-ende en sekvens
komplementær til den første sekvens,
hvor den første sekvens og den anden sekvens danner shRNA'et.

15 **8.** Immuncellen ifølge krav 6, omfattende en eller flere modifikationer af en sekvens der koder for B2M (SEQ ID NO: 170), eventuelt hvor den ene eller flere modifikationer omfatter en eller flere inaktiverende mutationer af det endogene gen, der koder for B2M, eventuelt hvor den ene eller flere inaktiverende mutationer indføres med en nukleinsyre-guided endonuklease i et kompleks med
20 mindst en guide-nukleinsyre (gNA), der er specifikt rettet mod en sekvens af det endogene gen, der koder for B2M (SEQ ID NO: 170).

9. Immuncellen ifølge et hvilket som helst af kravene 1-8, hvor:

- a. den første receptor omfatter SEQ ID NO: 177, eller en sekvens der deler
25 mindst 75%, mindst 80%, mindst 85%, mindst 90%, mindst 95% eller mindst 98% identitet dermed; og
- b. den anden receptor omfatter SEQ ID NO: 174 eller en sekvens der deler mindst 75%, mindst 80%, mindst 85%, mindst 90%, mindst 95% eller mindst 98% identitet dermed.

30

10. Immuncellen ifølge et hvilket som helst af kravene 1-9, yderligere omfattende et interfererende RNA, hvor det interfererende RNA omfatter SEQ ID NO: 179, eller en sekvens der deler mindst 75%, mindst 80%, mindst 85%, mindst 90%, mindst 95% eller mindst 98% identitet dermed, eventuelt hvor immuncellen er

autolog, eller hvor immuncellen er allogen.

11. Farmaceutisk sammensætning, omfattende en terapeutisk effektiv mængde af immuncellerne ifølge et hvilket som helst af kravene 1-10.

5

12. Polynukleotid omfattende polynukleotidsekvenser, der koder for den første receptor og den anden receptor til generering af immuncellerne ifølge et hvilket som helst af kravene 1-10 eller en vektor, omfattende polynukleotidet.

- 10 **13.** Immuncellen ifølge et hvilket som helst af kravene 1-10 til anvendelse i behandlingen af EGFR+-kræft, eventuelt hvor EGFR+-kræftformen er en EGFR+-tumor med tab af heterozygositet ved et locus, der koder for et ikke-målantigen, og/eller hvor EGFR+-kræftformen omfatter EGFR+-kræftceller med tab af heterozygositet ved et locus i MHC klasse I,
- 15 eventuelt hvor EGFR+-kræftcellen er en lungekræftcelle, en småcellet lungekræftcelle, en ikke-småcellet lungekræftcelle, en pankreatisk duktalt karcinomcelle, en kolorektal kræftcelle, en hoved- og halskræftcelle, en spiserørs- og mave-adenokarcinomcelle, en æggestokkræftcelle, en glioblastom multiformet celle, en cervikal pladeetiphelcellekarcinomcelle, en nyrekræftcelle, en papillær
- 20 nyrekræftcelle, en klar-cellet nyrecellekarcinomcelle, en blærekræftcelle, en brystkræftcelle, en galdegangskræftcelle, en leverkræftcelle, en prostatakkræftcelle, en sarkomcelle, en skjoldbruskkirtelkræftcelle, en thymuskirtelkræftcelle, en mavekræftcelle eller en livmoderkræftcelle.

- 25 **14.** Immuncellen til anvendelse ifølge krav 13, hvor EGFR+-kræftcellen er en lungekræftcelle;

hvor EGFR+-kræftcellen er en EGFR+/HLA-A*02-kræftcelle, der ikke udtrykker HLA-A*02; eller en kræftcelle afledt fra et HLA-A*02+-individ, som ikke udtrykker HLA-A*02; eventuelt

- 30 hvor EGFR+/HLA-A*02-kræftcellen er afledt fra en EGFR+/HLA-A*02+-celle ved tab af heterozygositet ved HLA-A, som fører til tab af HLA-A*02.

DRAWINGS

Drawing




FIG. 1B




FIG. 1C




FIG. 2A





FIG. 2B



Sample Name
HeLa_unstained_001.fcs
HCT116_unstained_005.fcs
HCT116_WT_007.fcs
HeLa_+A02_003.fcs

FIG. 3A




FIG. 3B




FIG. 3C





FIG. 3D



Hela

- WT
- +HLA-A2
- EGFR KO

FIG. 4A




FIG. 4B




FIG. 5




FIG. 6B




FIG. 6C




FIG. 7




FIG. 8

Enrichment by double-column purification on blocker & activator





FIG. 9




FIG. 10




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B




FIG. 13




FIG. 14A




FIG. 15A




FIG. 15B




FIG. 16




FIG. 17B




FIG. 17A





FIG. 18

FIG. 19




FIG. 20A




FIG. 20B

Construct	On-target mRNA (ng) IC50	On-target # of molecules IC50
Empty vector	-	-
Activator CAR	-	-
Activator CAR + HLA-A*11 blocker-4	31.5 ng A*11 mRNA	~23,000 (molecules per cell)

FIG. 21




FIG. 22




FIG. 23A




FIG. 23B





FIG. 24



SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

