



US 20210079048A1

(19) **United States**(12) **Patent Application Publication**
NORRIS(10) **Pub. No.: US 2021/0079048 A1**(43) **Pub. Date: Mar. 18, 2021**(54) **VMP-LIKE SEQUENCES OF PATHOGENIC
BORRELIA SPECIES AND STRAINS**(60) Provisional application No. 60/435,077, filed on Dec.
20, 2002.(71) Applicant: **BOARD OF REGENTS OF THE
UNIVERSITY OF TEXAS SYSTEM,**
Austin, TX (US)**Publication Classification**(72) Inventor: **Steven J. NORRIS,** Houston, TX (US)(51) **Int. Cl.****C07K 14/20** (2006.01)**G01N 33/569** (2006.01)**C07K 17/08** (2006.01)**C07K 17/12** (2006.01)**C08G 69/48** (2006.01)**G01N 33/68** (2006.01)(21) Appl. No.: **16/939,100**(22) Filed: **Jul. 27, 2020**(52) **U.S. Cl.**CPC **C07K 14/20** (2013.01); **G01N 33/56911**
(2013.01); **C07K 17/08** (2013.01); **A61K 39/02**
(2013.01); **C08G 69/48** (2013.01); **G01N**
33/6854 (2013.01); **C07K 17/12** (2013.01)**Related U.S. Application Data**(60) Continuation of application No. 16/412,913, filed on
May 15, 2019, now Pat. No. 10,759,833, which is a
continuation of application No. 15/609,188, filed on
May 31, 2017, now Pat. No. 10,323,070, which is a
continuation of application No. 14/804,882, filed on
Jul. 21, 2015, now Pat. No. 9,670,254, which is a
continuation of application No. 14/257,613, filed on
Apr. 21, 2014, now Pat. No. 9,115,193, which is a
continuation of application No. 13/645,950, filed on
Oct. 5, 2012, now Pat. No. 8,722,871, which is a
division of application No. 13/324,357, filed on Dec.
13, 2011, now Pat. No. 8,283,458, which is a division
of application No. 12/962,154, filed on Dec. 7, 2010,
now Pat. No. 8,076,470, which is a division of
application No. 10/539,956, filed on Apr. 6, 2006,
now Pat. No. 7,847,084, filed as application No.
PCT/US03/41182 on Dec. 22, 2003.

(57)

ABSTRACT

The present invention relates to DNA sequences encoding Vmp-like polypeptides of pathogenic *Borrelia*, the use of the DNA sequences in recombinant vectors to express polypeptides, the encoded amino acid sequences, application of the DNA and amino acid sequences to the production of polypeptides as antigens for immunoprophylaxis, immunotherapy, and immunodiagnosis. Also disclosed are the use of the nucleic acid sequences as probes or primers for the detection of organisms causing Lyme disease, relapsing fever, or related disorders, and kits designed to facilitate methods of using the described polypeptides, DNA segments and antibodies.

Specification includes a Sequence Listing.




FIG. 1A




FIG. 1B

	VR-I										VR-II										VR-III									
cassette1	ESAVDGVSKWLEEMIKAAKEAATK	---	GGG	---	NN	---	SD	K	---	---	NAL	TG	KE	VA	DEA	---	GA	NE	AG	KLF	AG	---	AGG	---	DADDAN	---	---	---	---	---
cassette2	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette3	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette4	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette5	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette6	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette7	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette8	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette9	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette10	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette11	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette12	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette13	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
cassette14	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
B31 VisE1	G.IKE.EL.DKIV	V.---	A	---	E	ASS	TAA	E.V.DADA	AK	...	A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...

	VR-IV										VR-V										VR-VI									
cassette1	IAKAAGAVTAVSGEQILKAIVD--CAGGAA-QDGKAA-EAKNPIAAAIGADAAG--ADFG-DMKK-SDTAALVLRGVAKSCKFAVAAA--KRESVKSAAV										TNDD.--E.K.G.--N										G									
cassette2	--A.A.N.....D										AG--N										NDS.A.....									
cassette3	--A..D-A...D-A										V										A.ANDNS.A.....									
cassette4	--A.D-D-A.....										N.DNA.A.K.E.--										D									
cassette5	--D.N.A.....										TNE.--E.--										D									
cassette6	--A.DK.N.....D										AD.AA.A.NEN.--										D									
cassette7	--A.A.N.A.....D										TADD.--E.K.--										D									
cassette8	--A.A.D-A.AA.G-A										GA--EEFK.E.--										G									
cassette9	--A.A.D-A.A.D-A										TNE.--A.K-G.RN.N										A.AD-AG.A.R.*RVL									
cassette10	EAA.D.N.A...D-										T.D.--A.Q										G									
cassette11	--A.....V.AG.G-A.T										AGD.--K.E.RN.										D									
cassette12	E-A.A.D-A.V.E										T.D.A-E.EN.-N.N										ANANDAG.....									
cassette13	E--A.DK.N.V.V..GA.T										T.DDNAA.FDK-.E.SN.										D									
cassette14	-----										GAG.--A.EN.RN.N										D									
B31 VlseI	AS.....S.....S.....PE-										DKDG.--E.Q.E.-D.Q										A.M.D									

FIG. 2A

	VR - I	VR - II	VR - III
cassette10	KGTV ----KNAVDMTKAAVAASA ----ASAATGAAIGDVVNGVDG-AAKGGDAASVNGIAKGIGIVDAAEKADAKEGKLN	VAGACAGAEAGNEAAGKLFVKKNAGDHGGEAGDAG	
cassette1	E.AITEIS.WLD.A.A... ---E. ---E. ---G. -.....E	D...D. -GAGGG.A -----.GA..I~	
cassette2	E.AITEIS.WLD.A.A... ---E. ---E. ---G. -.....E	D...D. -KD.....E...N...	
cassette3	E.....A.A...E... ---E. ---E. ---G. -.....E	D...D. -TT.VNV.....N.-NE..D.S...	
cassette4	A.A...E... ---E. ---E. ---G. -.....E		
cassette5	A.A...E... ---E. ---E. ---G. -.....E	D...D. -T.KD.....D..D...	
cassette6	A.EE... ---E. ---E. ---G. -.....E	DAT.E.-TT.VN.....RA.D.G.DADDAGK	
cassette7	II.E... ---E. ---E. ---G. -.....E	A...E.-TT.AD.....-NV.....	
cassette8	GSATNKT.V.I.NLVKTVQA.LKKV...KNSE---D...	D...D. -T.KD.....N.-NE..D.I...	
cassette9	II.T... ---E. ---E. ---G. -.....E	D...D. -T.KD.....-NE...N...	
cassette11	E.AI---KEVSELD.LVKAVKT. ---EG.SS.T...E.ADA. ---VA.K...K	KAVA...KG.N.KG.....G.AG.AA.-.DSEA.S	

	VR - IV	VR - V	VR - VI
cassette10	RAAAVAVSCEIILKAIVDAAKD GGDRQG-KKAEDAENIPIDAAIGSTGADNNAFAF --AATMKKDDQJAAAWVIRGMAKDGQFALK --DAAHNDH		
cassette1	K.....E.D....-A..T...E...GAD.GA.-.....NK.....D.AAH~		
cassette2	K.....VTDVK..T...E...--TN..D.A.--D...K.~		
cassette3	T...-V.A...--AD..-NKEG.....TNNA...~		
cassette4	K.....T.VTDVK..T...--SA..AN...--DK.....NND....		
cassette5	VA.....G.N.-ND..A.--		
cassette6	K.....T.KNNVK...E...SADA.A.FNK--EG.....TND...~		
cassette7	K.....D....-.....T...GA..GA..A..--NN.....T--NN.T...		
cassette8	AAEQD.-.P.E.K...A....D--K.GG.EFGQ--DE		
cassette9	IA.....K.VI--GEKEKA		
cassette11			
B31 VISe1			

FIG. 2B




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13

VMP-LIKE SEQUENCES OF PATHOGENIC BORRELIA SPECIES AND STRAINS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation of U.S. patent application Ser. No. 16/412,913, filed May 15, 2019, which is a continuation of U.S. patent application Ser. No. 15/609,188, filed May 31, 2019, now U.S. Pat. No. 10,323,070, which is a continuation of U.S. patent application Ser. No. 14/804,882, filed Jul. 21, 2015, now U.S. Pat. No. 9,670,254, which is a continuation of U.S. patent application Ser. No. 14/257,613, filed Apr. 21, 2014, now U.S. Pat. No. 9,115,193, which is a continuation of U.S. patent application Ser. No. 13/645,950, filed Oct. 5, 2012, now U.S. Pat. No. 8,722,871, issued May 13, 2014, which is a divisional of U.S. patent application Ser. No. 13/324,357, filed Dec. 13, 2011, now U.S. Pat. No. 8,283,458, issued Oct. 9, 2012, which is a divisional of U.S. patent application Ser. No. 12/962,154, filed Dec. 7, 2010, now U.S. Pat. No. 8,076,470, issued Dec. 13, 2011, which is a divisional of U.S. patent application Ser. No. 10/539,956, filed on Apr. 6, 2006, now U.S. Pat. No. 7,847,084, issued on Dec. 7, 2010, which is a U.S. national phase application under 35 U.S.C. § 371 of International Application No. PCT/US03/041182, filed Dec. 22, 2003, which claims priority to U.S. Provisional Patent Application No. 60/435,077, filed Dec. 20, 2002. The entire text of each of the above-referenced disclosures is specifically incorporated by reference.

BACKGROUND OF THE INVENTION

[0002] This invention was made with government support under AI37277 awarded by the National Institutes of Health. The government has certain rights in the invention.

A. FIELD OF THE INVENTION

[0003] The invention relates to the field of molecular biology; in particular, to immunogenic compositions and recombinant VMP-like genes useful for treatment and diagnosis of Lyme disease. Also included are methods for the determination of virulence factors in Lyme disease.

B. DESCRIPTION OF RELATED ART

[0004] Lyme disease is a bacterial infection caused by pathogenic spirochetes of the genus *Borrelia*. The infection can occur in humans, dogs, deer, mice and other animals, and is transmitted by arthropod vectors, most notably ticks of the genus *Ixodes*. *Borrelia burgdorferi*, the most common cause of Lyme disease in North America, was first cultured in 1982. *B. garinii* and *B. afzelii* are the most common infectious agents of Lyme disease in Europe, and another species, *B. japonicum*, has been described in Japan. These organisms are closely related and cause similar manifestations with multiple stages: an expanding rash at the site of the tick bite (erythema migrans); fever, lymphadenopathy, fatigue, and malaise; effects of disseminated infection, including carditis, meningoradiculitis, and polyarthritis; and chronic manifestations including arthritis and neurologic disorders.

[0005] Lyme disease is often difficult to diagnose because of shared manifestations with other disorders, and it can also be refractory to treatment during late stages of the disease. It is most common in areas such as suburban regions of

upstate New York and Connecticut, where large populations of deer and white-footed mice serve as the principal mammalian hosts and reservoirs of infection. Approximately 20,000 cases of Lyme disease in humans are reported per year in the United States, and it is also a significant veterinary problem due to a high infection rate of dogs and other domestic animals in endemic regions.

[0006] The pathogenic *Borrelia* that cause Lyme disease are able to persist for years in patients or animals despite the presence of an active immune response. Antigenic variation is a mechanism by which members of the genus *Borrelia* may be able to evade the host immune response (Zhang, 1997). Antigenic variation has been defined as changes in the structure or expression of antigenic proteins that occurs during infection at a frequency greater than the usual mutation rate (Borst and Geaves, 1987; Robertson and Meyer, 1992; Seifert and So, 1988).

[0007] Relapsing fever is another disease caused by pathogenic *Borrelia*. It has both epidemic and endemic forms. The epidemic form is caused by *B. recurrentis* and is transmitted between humans by lice. It was a major source of morbidity and mortality during World War I, but has been rare since then due largely to public health measures. Endemic relapsing fever is an epizootic infection caused by several *Borrelia* species, including *B. hermsii*. It occurs sporadically among hunters, spelunkers, and others who come in contact with infected soft-bodied ticks of the genus *Ornithodoros*. Relapsing fever is characterized by two or more episodes or "relapses" of high bacteremia (up to 10^8 /ml). The first wave of infection is caused by *Borreliae* expressing a certain Variable Major Protein (VMP) on their surface (e.g. Vmp21). The gene encoding this VMP is located at a promoter site in the expression plasmid, whereas over 24 nonexpressed copies of different VMP genes are present on the so-called silent plasmid. When the host develops antibodies against the expressed VMP, the organisms of that serotype are destroyed and the patient improves. However, a small proportion of organisms have undergone antigenic switching to a different serotype. Nonreciprocal recombination occurs between the expression plasmid and the silent plasmid, resulting in the insertion of a different VMP gene in the expression site (e.g., Vmp7). The organisms expressing Vmp7 are not affected by the anti-Vmp21 antibodies, and therefore multiply in the host and cause a second episode of the disease. Up to five of these 3-5 day episodes can occur, separated by 1-2 week intervals.

[0008] Such well-demarcated episodes of infection do not occur during Lyme disease, and fewer organisms are present in the blood at any stage. However, there are reasons to suspect that similar mechanisms of antigenic variation may occur in *B. afzelii* and other Lyme disease *Borreliae* such as *B. garinii* and *B. burgdorferi*. The infection, if untreated, commonly persists for months to years despite the occurrence of host antibody and cellular responses; this observation indicates effective evasion of the immune response. Lyme disease may be disabling (particularly in its chronic form), and thus there is a need for effective therapeutic and prophylactic treatment.

[0009] Genetic loci analogous to the VMP antigenic variation system have been detected in North American and European Lyme disease *Borrelia* by Southern hybridization and PCR analysis (Wang et al., 2001). In addition, sequences from fragments of vls (VMP-like sequence) silent cassettes have been reported for the *Borrelia burgdorferi* strains 297

and N40, and the *Borrelia garinii* strains Ip90 and A87S (Liang and Philipp, 1999; Wang et al., 2001), (S. Feng and S. W. Barthold, unpublished data). VMP-like sequences of *B. burgdorferi* have been described and patented in U.S. Pat. No. 6,437,116.

[0010] Open reading frames in a *B. burgdorferi* plasmid that encode hypothetical proteins resembling the VMP proteins of relapsing fever organisms have been identified (Zhang et al., 1997). The inventors have found that the presence of the plasmid containing these VMP-like sequences in *B. burgdorferi* clones correlates strongly with infectivity (Zhang et al., 1997; Purser and Norris, 2000). Thus it is likely that the proteins encoded by the VMP-like sequences are important in immunoprotection and pathogenesis. Proteins encoded by the VMP-like sequences of *B. burgdorferi* may provide protection when used either alone or in combination with other antigens. They may also be useful for immunodiagnosis.

[0011] Greater than 90% of Lyme disease patients beyond the erythema migrans stage from North America and Europe express antibodies against VlsE (Liang et al., 1999; Liang et al., 2000). In addition, mice infected experimentally with *Borrelia afzelii* and *Borrelia garinii* strains express anti-VlsE antibodies (Liang et al., 2000). Finally, a protein product of ~35 kDa expressed by *Borrelia garinii* Ip90 reacts with antibodies against IR6, a peptide corresponding to invariant region 6 of the VlsE cassette region (Liang et al., 1999a). Portions of several vls silent cassettes from *Borrelia garinii* strain A87S have been published (Wang et al., 2001). Further, several amino acid sequences of *Borrelia garinii* Ip90 have been previously characterized by Liang et al. (1999a).

[0012] There is a commercial demand for vaccines and diagnostic kits for Lyme disease, both for human and veterinary use. Several companies have active research and development programs in these areas.

SUMMARY OF THE INVENTION

[0013] Partial and complete DNA sequences have been determined for several recombinant clones containing DNA encoding VMP-like sequences. The identification and characterization of these sequences now allows: (1) identification of the expressed gene(s) or DNA segments containing open reading frames in several *Borreliae*; (2) expression of these gene(s) by a recombinant vector in a host organism such as *E. coli*; (3) immunization of laboratory animals with the resulting polypeptide, and determination of protective activity against *Borreliae* infection; (4) use of antibodies against the expressed protein to identify the reactive polypeptide(s) in *Borreliae* cells; (5) use of the expressed protein (s) to detect antibody responses in infected humans and animals; (6) determination of the presence, sequence differences, and expression of the VMP-like DNA sequences in other Lyme disease *Borreliae*.

[0014] The invention is contemplated to be useful in the immunoprophylaxis, diagnosis, or treatment of Lyme disease, relapsing fever, or related diseases in humans or animals. It is expected that recombinant or native proteins expressed by the VMP-like genes (or portions thereof) will be useful for (a) immunoprophylaxis against Lyme disease, relapsing fever, or related disorders in humans and animals; (b) immunotherapy of existing Lyme disease, relapsing fever, or related illnesses, by way of immunization or injection of antibodies directed against VMP-like proteins;

and (c) immunodiagnosis of Lyme disease, relapsing fever, or related diseases, including their use in kits in which the VMP-like proteins are the sole antigen or one of multiple antigens. The DNA may be employed in: (a) production of recombinant DNA plasmids or other vectors capable of expressing recombinant polypeptides; and (b) design and implementation of nucleic acid probes or oligonucleotides for detection and/or amplification of VMP-like sequences. The latter is expected to have application in the diagnosis of infection with *Borrelia* organisms.

[0015] Another aspect of the invention is the method for identification of possible virulence factors. This approach entails subtractive hybridization of target DNA from high infectivity organisms with driver DNA from low-infectivity strains or clones. This procedure greatly enriches for sequences which differ between the high- and low-infectivity strains and thus may encode proteins important in virulence. Of particular utility is the use of closely related isogenic clones that differ in their infectivity; in this case, the DNA differences should be restricted more stringently to those related to infectivity.

[0016] The invention is considered to include DNA segments corresponding to 10, 20, 30, and 40 base pairs of the VMP-like sequences; DNA segments inclusive of the entire open reading frames of the VMP-like sequences; shorter DNA segments containing portions of these open reading frames; amino acid sequences corresponding to both conserved and variable regions of the VMP-like sequences; recombinant vectors encoding an antigenic protein corresponding to the above amino acid sequences; recombinant cells where extrachromosomal DNA expresses a polypeptide encoded by the DNA encoding *Borrelia* VMP-like sequences; a recombinant *Borreliae* or *E. coli* cell containing the DNA encoding VMP-like sequences; methods of preparing transformed bacterial host cells using the DNA encoding the VMP-like polypeptides; methods using the plasmid or another vector to transform the bacterial host cell to express *Borreliae* polypeptides encoded by the DNA sequences; and methods for immunization of humans or animals with the native *Borreliae* polypeptides, polypeptides expressed by recombinant cells that include DNA encoding the VMP-like polypeptides, or synthetic peptides that include VMP-like sequences.

[0017] Also included in the invention are primer sets capable of priming amplification of the VMP-like DNA sequences; kits for the detection of *Borreliae* nucleic acids in a sample, the kits containing a nucleic acid probe specific for the VMP-like sequences, together with a means for detecting a specific hybridization with the probe; kits for detection of antibodies against the VMP-like sequences of *Borreliae* and kits containing a native, recombinant, or synthetic VMP-like polypeptide, together with means for detecting a specific binding of antibodies to the antigen.

[0018] A preferred embodiment of the present invention is an isolated nucleic acid comprising a nucleotide sequence that encodes an antigenic peptide of *Borrelia garinii* or *B. afzelii*. More preferably, the present invention provides an isolated nucleic acid that encodes at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 75, 100, 125, 150, 175, 200 or more contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID

NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97. Further, the invention contemplates any range derivable between any of the above-described integers.

[0019] In another embodiment, the present invention provides an isolated nucleic acid comprising 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300, 400, 500 or more contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96. Further, the invention contemplates any range derivable between any of the above-described integers.

[0020] In yet another embodiment, the isolated nucleic acid comprises a complement to or a degenerate variant of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300, 400, 500 or more contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96. Further, the invention contemplates any range derivable between any of the above-described integers.

[0021] In some embodiments the isolated nucleic acid is a DNA molecule. In other embodiments the isolated nucleic acid is an RNA molecule.

[0022] In certain embodiments the invention provides an isolated nucleic acid obtained by amplification from a template nucleic acid using a primer selected from the group consisting of SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, and SEQ ID NO:107.

[0023] The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, enhancers, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like.

[0024] A preferred embodiment of the present invention is an isolated polypeptide comprising 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 75, 100, 125, 150, 175, 200 or more contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97. Further, the invention contemplates any range derivable between any of the above-described integers.

[0025] In one aspect, the present invention provides for an isolated polypeptide or an isolated nucleic acid encoding a polypeptide having between about 70% and about 75%; or more preferably between about 75% and about 80%; or more preferably between about 80% and about 90%; or even more preferably between about 90% and about 99% of amino acids that are identical to the amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 or fragments thereof. The percent identity or homology is determined with regard to the length of the relevant amino acid sequence. Therefore, if a polypeptide of the present invention is comprised within a larger polypeptide, the percent homology is determined with regard only to the portion of the polypeptide that corresponds to the polypeptide of the present invention and not the percent homology of the entirety of the larger polypeptide.

[0026] In addition, the present invention encompasses fragments of polypeptides or nucleic acids encoding fragments of polypeptides that have between about 70% and about 75%; or more preferably between about 75% and about 80%; or more preferably between about 80% and about 90%; or even more preferably between about 90% and about 99% of amino acids that are identical to the amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 even if the particular fragment itself does not have between about 70% and about 75%; or more preferably between about 75% and about 80%; or more preferably between about 80% and about 90%; or even more preferably between about 90% and about 99% amino acid homology with the polypeptides of the present invention.

[0027] In another embodiment the invention provides an isolated polypeptide that binds immunologically with antibodies raised against an antigenic polypeptide of *Borrelia garinii* or *B. afzelii*. In a preferred embodiment the antibodies are raised against an antigenic polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 75, 100, 125, 150, 175, 200 or more contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97. Further, the invention contemplates any range derivable between any of the above-described integers.

[0028] The polypeptides of the present invention may be fused with other proteins or peptides. Such fusion polypeptides may be useful for purification or immunodetection purposes, for example. In a preferred embodiment the polypeptides of the invention are expressed as fusions with β -galactosidase, avidin, ubiquitin, *Schistosoma japonicum* glutathione S-transferase, multiple histidines, epitope-tags and the like.

[0029] Another aspect of the invention comprises vectors that comprise a nucleic acid encoding all or part of a polypeptide of the present invention. The vectors may, for example, be cloning or expression vectors.

[0030] In certain embodiments, it is contemplated that particular advantages will be gained by positioning the nucleic acid sequences of the present invention under the control of a promoter. The promoter may be the promoter

that is normally associated with the nucleic acid sequence in its natural environment or it may be a recombinant or heterologous promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with a vls gene in its natural environment. Such promoters may include those normally associated with other *Borrelia* polypeptide genes, or promoters isolated from any other bacterial, viral, eukaryotic, or mammalian cell. Naturally, it will be important to employ a promoter that effectively directs the expression of the nucleic acid in the particular cell being used.

[0031] The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level or regulated expression of the introduced nucleic acid. In preferred embodiments the promoters are lac, T7, Ara, CMV, RSV LTR, the SV40 promoter alone, or the SV40 promoter in combination with the SV40 enhancer.

[0032] Another embodiment is a method of preparing a protein composition comprising growing a recombinant host cell comprising a vector that encodes a polypeptide of the present invention under conditions permitting nucleic acid expression and protein production followed by recovering the protein so produced. The host cell, conditions permitting nucleic acid expression, protein production and recovery, will be known to those of skill in the art, in light of the present disclosure of the vls gene. The recombinant host cell may be a prokaryotic cell or a eukaryotic cell.

[0033] VMP-like related proteins and functional variants are also considered part of the invention. Thus it is expected that truncated and mutated versions of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 will afford more convenient and effective forms of polypeptides for treatment regimens. Thus, any functional version of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97, such as truncated species or homologs, and mutated versions of VMP-like protein are considered as part of the invention.

[0034] Another aspect of the invention comprises the recombination of the 14 silent vls cassettes of *B. afzelii* in numerous combinations, providing for example a cocktail of peptide compositions for use as immunogens to develop vaccines for use in Lyme disease and related conditions. Likewise, the 11 silent vls cassettes of *B. garinii* and the 15

silent vls cassettes of *B. burgdorferi* may be recombined in numerous combinations. It is further contemplated by the present invention that these cassettes may be recombined among strains, as well as species of *Borrelia*, providing a cocktail of peptide compositions for use as immunogens to develop vaccines for use in Lyme disease and related conditions.

[0035] Pharmaceutical compositions prepared in accordance with the present invention find use in preventing or ameliorating conditions associated with *Borrelia* infections, particularly Lyme disease.

[0036] Such methods generally involve administering a pharmaceutical composition comprising an effective amount of a VMP-like antigen of *Borrelia*, such as SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 or various epitopes thereof.

[0037] In certain embodiments of the invention a vaccine may comprise a polynucleotide encoding an antigenic polypeptide. In more specific embodiments the polynucleotide may have a sequence of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96 or various fragments thereof. The vaccines of the present invention may comprise multiple polypeptides and/or polynucleotides.

[0038] It will also be understood that, if desired, the nucleic acid segment or gene encoding a VMP-like protein could be administered in combination with further agents, such as, proteins or polypeptides or various pharmaceutically active agents. There is virtually no limit to other components which may also be included, given that the additional agents do not cause a significant adverse effect upon contact with the target cells or host tissues.

[0039] Therapeutic kits comprising a polypeptide or nucleic acid of the present invention comprise another aspect of the invention. Such kits will generally contain, in suitable container means, a pharmaceutically acceptable formulation of a polypeptide or nucleic acid of the present invention. The kit may have a single container means that contains a polypeptide or nucleic acid of the present invention or it may have distinct container means for the polypeptide or nucleic acid of the present invention and other reagents that may be included within such kits.

[0040] The components of the kit may be provided as liquid solution(s), or as dried powder(s). When the components are provided in a liquid solution, the liquid solution is an aqueous solution, with a sterile aqueous solution being particularly preferred. When reagents or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container means.

[0041] In another embodiment, the invention provides diagnostic kits. The diagnostic kits may comprise reagents for detecting VMP-like polypeptides or anti-VMP-like antibodies in a sample, such as required for immunoassay. The immunodetection reagent will typically comprise a label associated with the antibody or antigen, or associated with a secondary binding ligand. Exemplary ligands might include a secondary antibody directed against the first antibody or antigen or a biotin or avidin (or streptavidin) ligand having an associated label. Of course, a number of exemplary labels are known in the art and all such labels may be employed in connection with the present invention. The kits may contain antibody-label conjugates either in fully conjugated form, in the form of intermediates, or as separate moieties to be conjugated by the user of the kit.

[0042] The container means will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which the antigen or antibody may be placed, and preferably suitably aliquoted. Where a second binding ligand is provided, the kit will also generally contain a second vial or other container into which this ligand or antibody may be placed. The kits of the present invention will also typically include a means for containing the antibody, antigen, and reagent containers in close confinement for commercial sale. Such containers may include injection or blow-molded plastic containers into which the desired vials are retained.

[0043] In another aspect, the present invention contemplates an antibody that is immunoreactive with a polypeptide of the invention. An antibody can be a polyclonal or a monoclonal antibody. In a preferred embodiment, an antibody is a monoclonal antibody.

[0044] Antibodies, both polyclonal and monoclonal, specific for VMP-like polypeptides and particularly those represented by SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 or variants and epitopes thereof, may be prepared using conventional immunization techniques, as will be generally known to those of skill in the art.

[0045] In related embodiments, the invention provides methods of using the antibodies of the invention. In preferred embodiments, the antibodies may be used in immunochemical procedures, such as ELISA and Western blot methods. In other embodiments, the antibodies may be used in purifying native or recombinant VMP-like polypeptides, inhibition studies, and immunolocalization studies.

[0046] Table 1 below provides the SEQ ID NO, the GenBank accession number, if any, and a brief description of the sequences described herein.

TABLE 1

SEQ ID NO.	GENBANK NO.	DESCRIPTION
SEQ ID NO: 1	U76405	<i>B. burgdorferi</i> vlsE gene allele vlsE1
SEQ ID NO: 2	AAC45733	Translation of <i>B. burgdorferi</i> vlsE1 gene
SEQ ID NO: 3	L04788	<i>B. hermsii</i> vmp17 gene
SEQ ID NO: 4	AAA22963	Translation of <i>B. hermsii</i> vmp17 gene
SEQ ID NO: 5	AY100629	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2622 vlsE
SEQ ID NO: 6	AAM77200	Translation of AY100629
SEQ ID NO: 7	AY100630	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2624a vlsE
SEQ ID NO: 8	AAM77201	Translation of AY100630
SEQ ID NO: 9	AY100631	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2624b vlsE
SEQ ID NO: 10	AAM77202	Translation of AY100631
SEQ ID NO: 11	AY100632	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2625 vlsE
SEQ ID NO: 12	AAM77203	Translation of AY100632
SEQ ID NO: 13	AY100634	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 17 vlsE
SEQ ID NO: 14	AAM77204	Translation of AY100634
SEQ ID NO: 15	AY100635	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 20 vlsE
SEQ ID NO: 16	AAM77205	Translation of AY100635
SEQ ID NO: 17	AY100636	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 21 vlsE
SEQ ID NO: 18	AAM77206	Translation of AY100636
SEQ ID NO: 19	AY100637	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 23 vlsE
SEQ ID NO: 20	AAM77207	Translation of AY100637
SEQ ID NO: 21	N/A	Primer 4540 (Wang et al., 2001)
SEQ ID NO: 22	N/A	Primer 4548 (Wang et al., 2001)
SEQ ID NO: 23	N/A	Primer 4545 (Wang et al., 2001)
SEQ ID NO: 24	N/A	Primer 4587 (Wang et al., 2001)
SEQ ID NO: 25	N/A	Primer 4588 (Wang et al., 2001)
SEQ ID NO: 26	N/A	Primer 4470 (Wang et al., 2001)
SEQ ID NO: 27	N/A	Primer 4471 (Wang et al., 2001)
SEQ ID NO: 28	AY100633	<i>B. garinii</i> vls silent cassette locus
SEQ ID NO: 29	AY100633	<i>B. garinii</i> upstream ORF
SEQ ID NO: 30	AAN87823	Translation of <i>B. garinii</i> upstream ORF
SEQ ID NO: 31	AY100633	<i>B. garinii</i> 5' vlsE homolog
SEQ ID NO: 32	AAN87824	Translation of <i>B. garinii</i> 5' vlsE homolog
SEQ ID NO: 33	AY100633	<i>B. garinii</i> vls1
SEQ ID NO: 34	AAN87825	Translation of <i>B. garinii</i> vls1
SEQ ID NO: 35	AY100633	<i>B. garinii</i> vls2
SEQ ID NO: 36	AAN87826	Translation of <i>B. garinii</i> vls2
SEQ ID NO: 37	AY100633	<i>B. garinii</i> vls3
SEQ ID NO: 38	AAN87827	Translation of <i>B. garinii</i> vls3
SEQ ID NO: 39	AY100633	<i>B. garinii</i> vls4
SEQ ID NO: 40	AAN87828	Translation of <i>B. garinii</i> vls4
SEQ ID NO: 41	AY100633	<i>B. garinii</i> vls5
SEQ ID NO: 42	AAN87829	Translation of <i>B. garinii</i> vls5
SEQ ID NO: 43	AY100633	<i>B. garinii</i> vls6
SEQ ID NO: 44	AAN87830	Translation of <i>B. garinii</i> vls6
SEQ ID NO: 45	AY100633	<i>B. garinii</i> vls7
SEQ ID NO: 46	AAN87831	Translation of <i>B. garinii</i> vls7
SEQ ID NO: 47	AY100633	<i>B. garinii</i> vls8
SEQ ID NO: 48	AAN87832	Translation of <i>B. garinii</i> vls8
SEQ ID NO: 49	AY100633	<i>B. garinii</i> vls9
SEQ ID NO: 50	AAN87833	Translation of <i>B. garinii</i> vls9
SEQ ID NO: 51	AY100633	<i>B. garinii</i> vls10
SEQ ID NO: 52	AAN87834	Translation of <i>B. garinii</i> vls10
SEQ ID NO: 53	AY100633	<i>B. garinii</i> vls11
SEQ ID NO: 54	AAN87835	Translation of <i>B. garinii</i> vls11
SEQ ID NO: 55	AY100633	<i>B. garinii</i> truncated gene

TABLE 1-continued

SEQ ID NO.	GENBANK NO.	DESCRIPTION
SEQ ID NO: 56	AAN87823	Translation of <i>B. garinii</i> truncated gene
SEQ ID NO: 57	AY100628	vis silent cassette locus of <i>B. afzelii</i>
SEQ ID NO: 58	AY100628	<i>B. afzelii</i> vls1
SEQ ID NO: 59	AAN87809	Translation of <i>B. afzelii</i> vls1
SEQ ID NO: 60	AY100628	<i>B. afzelii</i> vls2
SEQ ID NO: 61	AAN87810	Translation of <i>B. afzelii</i> vls2
SEQ ID NO: 62	AY100628	<i>B. afzelii</i> vls3
SEQ ID NO: 63	AAN87811	Translation of <i>B. afzelii</i> vls3
SEQ ID NO: 64	AY100628	<i>B. afzelii</i> vls4
SEQ ID NO: 65	AAN87812	Translation of <i>B. afzelii</i> vls4
SEQ ID NO: 66	AY100628	<i>B. afzelii</i> vls5
SEQ ID NO: 67	AAN87813	Translation of <i>B. afzelii</i> vls5
SEQ ID NO: 68	AY100628	<i>B. afzelii</i> vls6
SEQ ID NO: 69	AAN87814	Translation of <i>B. afzelii</i> vls6
SEQ ID NO: 70	AY100628	<i>B. afzelii</i> vls7
SEQ ID NO: 71	AAN87815	Translation of <i>B. afzelii</i> vls7
SEQ ID NO: 72	AY100628	<i>B. afzelii</i> vls8
SEQ ID NO: 73	AAN87816	Translation of <i>B. afzelii</i> vls8
SEQ ID NO: 74	AY100628	<i>B. afzelii</i> vls9a
SEQ ID NO: 75	AAN87817	Translation of <i>B. afzelii</i> vls9a
SEQ ID NO: 76	AY100628	<i>B. afzelii</i> vls10
SEQ ID NO: 77	AAN87818	Translation of <i>B. afzelii</i> vls10
SEQ ID NO: 78	AY100628	<i>B. afzelii</i> vls11
SEQ ID NO: 79	AAN87819	Translation of <i>B. afzelii</i> vls11
SEQ ID NO: 80	AY100628	<i>B. afzelii</i> vls12
SEQ ID NO: 81	AAN87820	Translation of <i>B. afzelii</i> vls12
SEQ ID NO: 82	AY100628	<i>B. afzelii</i> vls13
SEQ ID NO: 83	AAN87821	Translation of <i>B. afzelii</i> vls13
SEQ ID NO: 84	AY100628	<i>B. afzelii</i> vls14
SEQ ID NO: 85	AAN87822	Translation of <i>B. afzelii</i> vls14
SEQ ID NO: 86	AY100628	<i>B. afzelii</i> conserved protein
SEQ ID NO: 87	AAN87823	Translation of <i>B. afzelii</i> conserved protein
SEQ ID NO: 88	N/A	Nucleotides 1-2775 of AY100633 (<i>B. garinii</i>)
SEQ ID NO: 89	N/A	Nucleotides 3823-5897 of AY100633 (<i>B. garinii</i>)
SEQ ID NO: 90	N/A	Fragment of <i>B. garinii</i> vls5
SEQ ID NO: 91	N/A	Amino acids 1-184 of AAN87829 (<i>B. garinii</i>)
SEQ ID NO: 92	N/A	Fragment of <i>B. garinii</i> vls8
SEQ ID NO: 93	N/A	Amino acids 56-195 of AAN87832 (<i>B. garinii</i>)
SEQ ID NO: 94	N/A	Expressed ORF in pBG-10-1
SEQ ID NO: 95	N/A	Protein sequence expressed by pBG-10-1
SEQ ID NO: 96	N/A	Expressed ORF in pBA-13-1
SEQ ID NO: 97	N/A	Protein sequence expressed by pBA-13-1
SEQ ID NO: 98	N/A	Primer
SEQ ID NO: 99	N/A	Primer
SEQ ID NO: 100	N/A	Primer
SEQ ID NO: 101	N/A	Primer
SEQ ID NO: 102	N/A	Primer
SEQ ID NO: 103	N/A	Primer
SEQ ID NO: 104	N/A	Primer
SEQ ID NO: 105	N/A	Primer
SEQ ID NO: 106	N/A	17-bp direct repeat of <i>B. burgdorferi</i>
SEQ ID NO: 107	N/A	EcoRI linker

BRIEF DESCRIPTION OF THE DRAWINGS

[0047] FIGS. 1A-1B. Arrangement of vls silent cassette regions of *B. garinii* Ip90 and *B. afzelii* ACAI. The orientation of the silent cassettes is indicated by a dashed arrow. Direct repeats are indicated by heavily weighted lines between silent cassettes. The location and orientation of conserved hypothetical protein genes are indicated at the 3' end of each locus. Restriction sites used for cloning and

sequencing are also shown. (FIG. 1A) *B. garinii* Ip90. The cross-hatched bar indicates the location of P7-1 clone (Liang and Philipp, 1999) in the vls locus of Ip90. The locations of the telomeric repeat sequences (TRS) and the vlsE-like sequence are shown. (FIG. 1B) *B. afzelii* ACAI. The location and orientation of the vls cassettes and other features of this region are indicated as described above.

[0048] FIGS. 2A-2B. Alignment of predicted amino acid sequences of vls silent cassettes of *B. afzelii* ACAI (FIG. 2A) and *B. garinii* Ip90 (FIG. 2B) with the cassette region of *B. burgdorferi* B31 vlsE. Alignment for *B. afzelii* ACAI is based on cassette 1 and for *B. garinii* Ip90 based on cassette 10. The underlined residues at the end of cassette 9 in panel A are a continuation of the cassette following a frameshift. Identical amino acid sequences are shown as periods. The variable regions are indicated by shaded boxes and the lines under the shaded boxes represent the corresponding variable regions of *B. burgdorferi* B31. Gaps and predicted stop codons are indicated as dashes and asterisks, respectively. Cassette 1 (SEQ ID NO:59), cassette 2 (SEQ ID NO:61), cassette 3 (SEQ ID NO:63), cassette 4 (SEQ ID NO:65), cassette 5 (SEQ ID NO:67), cassette 6 (SEQ ID NO:69), cassette 7 (SEQ ID NO:71), cassette 8 (SEQ ID NO:73), cassette 9 (SEQ ID NO:75), cassette 10 (SEQ ID NO:77), cassette 11 (SEQ ID NO:79), cassette 12 (SEQ ID NO:81), cassette 13 (SEQ ID NO:83), cassette 14 (SEQ ID NO:85), cassette B31 vlsE (SEQ ID NO:108).

[0049] FIG. 3. RT-PCR of vlsE sequences, using RNA from *B. afzelii* ACAI (lanes 1 and 2) and *B. garinii* Ip90 (lanes 3 and 4) as template. Lanes 2 and 4, with reverse transcriptase; lanes 1 and 3, controls without reverse transcriptase. DNA marker sizes (bp) are indicated on the left.

[0050] FIGS. 4A-4B. Alignment of the predicted amino acid sequences based on RT-PCR products from vlsE variants of *B. afzelii* ACAI (FIG. 4A) and *B. garinii* Ip90 (FIG. 4B). Alignments for *B. afzelii* ACAI and *B. garinii* Ip90 are based on the sequences of clones 2622 and 17, respectively. The variable regions labeled VR-I through VR-VI (FIG. 4A) and VR-II through VR-V (FIG. 4B) are indicated by boxes. Only portions of VR-I and VR-VI are shown for ACAI. Identical amino acid sequences and gaps are shown as periods and dashes, respectively. Solid and dotted bars indicate the predicted minimum and maximum possible recombination events, respectively, resulting in the given vlsE variant. Solid lines indicate 100% sequence identity between the given position in the variant and silent cassette (s) indicated. Dashed lines mark the limits of maximum recombination. Asterisks above certain residues indicate sites of possible point mutations, as explained in the text. In regions where more than one silent cassette matches the variant amino acid sequence, the matches were further analyzed at the nucleotide level. ACAI VlsE Clone 2622 (SEQ ID NO:109), ACAI VlsE Clone 2624a (SEQ ID NO:110), ACAI VlsE Clone 2624b (SEQ ID NO:111), ACAI VlsE Clone 2625 (SEQ ID NO:112), Ip90 VlsE Clone 17 (SEQ ID NO:113), Ip90 VlsE Clone 20 (SEQ ID NO:114), Ip90 VlsE Clone 21 (SEQ ID NO:115), Ip90 VlsE Clone 23 (SEQ ID NO:116).

[0051] FIG. 5. Hybridization of plasmid DNA of *B. afzelii* ACAI and *B. garinii* Ip90 with pJRZ53 probe. Lane 1, ACAI plasmid DNA; lane 2, ACAI plasmid DNA digested with EcoRI; lane 3, Ip90 plasmid DNA; and lane 4, Ip90 plasmid DNA digested with EcoRI. The size of EcoRI fragments containing vls sequences are indicated by arrows at left.

[0052] FIG. 6. Reactivity of human Lyme disease serum pool and a normal human serum pool with recombinant *Borrelia afzelii* Vls protein VLS-BA13.

[0053] FIG. 7. Effect of VLS-BA13 protein concentration on enzyme immunoassay reactivity of serum pools from Lyme disease human subjects and normal human subjects.

[0054] FIG. 8. Reactivity of mouse anti-*Borrelia burgdorferi* serum and normal mouse serum with recombinant *Borrelia afzelii* Vls protein VLS-BA13. The reactivity of normal mouse serum was below background levels.

[0055] FIG. 9. Effect of VLS-BA13 protein concentration on enzyme immunoassay reactivity of mouse anti-*B. burgdorferi* antiserum and normal mouse serum. The reactivity of normal mouse serum was below background levels.

[0056] FIG. 10. Reactivity of human Lyme disease serum pool and a normal human serum pool with recombinant *Borrelia garinii* Vls protein VLS-BG10.

[0057] FIG. 11. Effect of VLS-BG10 protein concentration on enzyme immunoassay reactivity of serum pools from Lyme disease human subjects and normal human subjects.

[0058] FIG. 12. Reactivity of mouse anti-*Borrelia burgdorferi* serum and normal mouse serum with recombinant *Borrelia garinii* Vls protein VLS-BG10. The reactivity of normal mouse serum was below background levels.

[0059] FIG. 13. Effect of VLS-BG10 protein concentration on enzyme immunoassay reactivity of mouse anti-*B. burgdorferi* antiserum and normal mouse serum. The reactivity of normal mouse serum was below background levels.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0060] The present work discloses the identification and characterization of an elaborate genetic system in the Lyme disease spirochete *Borrelia* that promotes extensive antigenic variation of a surface-exposed lipoprotein, VlsE.

[0061] Hybridization with the *B. burgdorferi* B31 vls silent cassette sequence in recombinant plasmid pJRZ53 was used in identifying the plasmids and DNA fragments containing vls sequences in *B. garinii* Ip90 and *B. afzelii* ACAI. The pJRZ53 probe hybridized exclusively to plasmids with an approximate size of 28 kb in both ACAI and Ip90. DNA fragments from these *B. garinii* Ip90 and *B. afzelii* ACAI plasmids were inserted into a recombinant lambda bacteriophage vector (lambda-DashI) and sequenced. The results showed *B. garinii* Ip90 to consist of 11 vls silent cassettes and *B. afzelii* ACAI of 14 vls silent cassettes.

[0062] With the exception of the junctions at vls3/4 and vls6/7, the 11 vls silent cassettes of Ip90 are flanked by 18 bp direct repeat sequences in the 6 kb region. However, several of these cassettes (vls1, vls4, vls6, and vls11) are truncated (189 to 288 bp in length) relative to the other, full-length cassettes ranging in size from 573 to 594 bp. Unlike Ip90 and B31, the ACAI vls locus is located on an internal EcoRI fragment of a 28-kb linear plasmid, and its location relative to the plasmid telomeres is not known. The ACAI vls locus contained 13 complete and 1 partial silent cassettes with each cassette being flanked by an 18 bp direct repeat sequence.

[0063] These silent cassettes share 90% to 97% nucleotide sequence identity with one another within the Ip90 vls locus and 84% to 91% within the ACAI vls locus. Amino acid similarity to the B31 silent cassettes ranges from 51% to 62% for the Ip90 vls silent cassettes and from 50% to 65% for the ACAI vls silent cassettes. The nucleotide sequence

and predicted amino acid sequence of vlsE in *B. burgdorferi* is provided in SEQ ID NO:1 and SEQ ID NO:2, respectively. The vlsE expression sites of Ip90 and ACAI have not been isolated, but transcripts of vlsE have been detected by reverse transcriptase PCR for both Ip90 and ACAI. In addition, the occurrence of sequence variation within the vlsE cassette region of these transcripts was verified. Mice infected experimentally with *B. garinii* and *B. afzelii* strains have been shown to express anti-VlsE antibodies (Liang et al., 2000a). Additionally, a protein product of ~35 kDa expressed by *B. garinii* Ip90 reacts with antibodies against IR6, a peptide corresponding to invariant region 6 of the VlsE cassette region (Liang et al., 1999a). The characteristics of the vls loci present in *B. garinii* Ip90 and *B. afzelii* ACAI are therefore similar to those found in *B. burgdorferi* B31.

[0064] Genetic variation involved in multi-gene families has been described in several other pathogenic microorganisms (Borst and Geaves, 1987; Borst et al., 1995; Donelson, 1995). In the context of combinatorial recombination, the genetic variation at the vlsE site is similar to that of the pilin-encoding genes of *Neisseria gonorrhoeae* (Seifert and So, 1988). The gonococcal pilus is primarily composed of repeating subunits of an 18-kilodalton pilin protein and is required for adherence of the bacterium to a variety of human cells (Swanson and Koomey, 1989). While the complete pilin genes are expressed only at two expression sites (pilE1 and pilE2), multiple silent copies (pilS) containing portions of the pilin genes are found over a wide range on the gonococcal chromosome (Haas and Meyer, 1986). Through multiple combinatorial recombination events, a single gonococcal clone expressing one pilin stereotype can give rise to a large number of progeny that express antigenically distinctive pilin variants (Meyer et al., 1982; Hagblom et al., 1985; Segal et al., 1986). The recombination between the expression and silent loci occurs predominantly through a non-reciprocal gene conversion mechanism (Haas and Meyer, 1986; Koomey et al., 1987).

[0065] The coding sequences of the *Neisseria* pilin variants are divided into constant, semi-variable, and hypervariable regions (Haas and Meyer, 1986), which are analogous to the conserved and variable regions of the vls cassettes. The constant regions and a conserved DNA sequence (Sma/Cla repeat) located at the 3' end of all pilin loci are thought to pair the regions involved in recombination events (Wainwright et al., 1994). In this context, the 18-bp direct repeats and the conserved regions of the vls cassettes in *B. garinii* and *B. afzelii* may play a similar role in recombination events. The silent loci of gonococcal pilin genes contain different regions of the complete pilin genes (Haas and Meyer, 1986), whereas the silent vls cassettes of *Borrelia* represent only the central cassette region of the vlsE gene.

[0066] Non-reciprocal recombinations also occur between the expressed and the silent genes encoding variant surface glycoproteins (VsGs) in African trypanosomes (Donelson, 1995). Based on similarities between the vls locus and the multi-gene families of the other pathogenic microorganisms and experimental data (Zhang and Norris, 1998b), it is likely that a unidirectional gene conversion mechanism is also active in the vls locus. The exact mechanism of vls recombination remains to be determined.

[0067] Variation of *Borrelia* surface proteins such as VlsE may also affect the organism's virulence and its ability to adapt to different micro-environments during infection of

the mammalian host. Recent studies of a *Borrelia turicatae* mouse infection model that resembles Lyme disease showed that one serotype expressing VmpB exhibited more severe arthritic manifestations, whereas another expressing VmpA had more severe central nervous system involvement. The numbers of *Borreliae* present in the joints and blood of serotype B-infected mice were much higher than those of mice infected with serotype A, consistent with a relationship between Vmp serotype and disease severity. Antigenic variation of *Neisseria* pilin (Lambden et al., 1980; Rudel et al., 1992; Nassif et al., 1993; Jonsson et al., 1994) and Opa proteins (Kupsch et al., 1993) is known to affect adherence of the organisms to human leukocytes and epithelial cells.

A. Antigenic Variation in *B. hermsii*

[0068] A complex antigenic variation mechanism has been characterized in *Borrelia hermsii*, a relative of *B. afzelii* and *B. garinii* that causes relapsing fever (Balmelii and Piffatetti, 1996; Barbour, 1993; Donelson, 1995). Surface-exposed lipoproteins called variable major proteins (Vmps) are encoded by homologous genes located in 28- to 32-kb linear plasmids with covalently closed telomeres (Barbour and Garon, 1987; Kitten and Barbour, 1990). The vmp genes have been subdivided into two groups: small and large (Restrepo et al., 1992). Large vmp genes such as vmp7 and vmp17 and small vmp genes such as vmp1 and vmp3 are approximately 1 kb and 0.6 kb in size, respectively. Each organism contains both small and large vmp genes in an unexpressed (silent) form in the so-called storage plasmids (Plasterk et al., 1985). Only one vmp gene located near one of the telomeres of a different plasmid (called the expression plasmid) is expressed in each organism (Kitten and Barbour, 1990; Barbour et al., 1991a). The nucleotide sequence and predicted amino acid sequence of an expressed vmp gene of *B. hermsii* are provided in SEQ ID NO:3 and SEQ ID NO:4, respectively. Antigenic variation occurs when the expressed vmp is replaced completely or partially by one of the silent vmp genes at the telomeric expression site through interplasmic recombination (Meier et al., 1985; Plasterk et al., 1985; Barbour et al., 1991b), intraplasmic recombination (Restrepo et al., 1994), and post-switch rearrangement (Restrepo and Barbour, 1994). The antigenic switch occurs spontaneously at a frequency of 10^{-3} to 10^{-4} per generation (Stoenner et al., 1982).

B. Identification of Vls

[0069] The present invention discloses a repetitive DNA sequence ~500 bp in length, which is present in multiple, nonidentical copies in a 28-kb linear plasmid of infectious *Borrelia burgdorferi*, *Borrelia garinii*, and *Borrelia afzelii*, the causative agents of Lyme disease. These DNA sequences encode polypeptides that have sequence similarity to the Variable Major Proteins (VMPs) of relapsing fever *Borreliae* (such as *B. hermsii*). VMPs are highly antigenic surface proteins, which the relapsing fever *Borreliae* are able to change through a genetic recombination mechanism, thereby evading the immune response. Antibodies against a particular VMP protein are protective, resulting in rapid clearance of bacteria of the corresponding serotype. In *Borrelia burgdorferi*, *Borrelia garinii*, and *Borrelia afzelii*, VMP-like sequences (vls) are present on a 28-kb linear plasmid, and this plasmid appears to encode virulence factor(s) required for infectivity.

C. ELISAs

[0070] ELISAs may be used in conjunction with the invention. In an ELISA assay, proteins or peptides incorporating *Borrelia* Vls antigenic sequences are immobilized onto a selected surface, preferably a surface exhibiting a protein affinity such as the wells of a polystyrene microtiter plate. The antigenic proteins or peptides may be isolated or comprised within larger polypeptides. For example, an antigenic Vls peptide may be comprised within a larger polypeptide that also includes a moiety that is useful for anchoring the polypeptide to the selected surface. The anchoring moiety may be an amino acid sequence. Virtually any amino acid sequence may be added to the antigenic Vls sequence so long as it does not confound the results of the ELISA assay. Those of skill in the art would know how to select amino acid sequences that are antigenically neutral with regard to antibodies in the biological sample (including, but not limited to, whole blood, plasma, serum, cerebrospinal fluid, other body fluids, or tissue extracts) that is being tested.

[0071] After washing to remove incompletely adsorbed material, it is desirable to bind or coat the assay plate wells with a nonspecific protein that is known to be antigenically neutral with regard to the biological sample such as bovine serum albumin (BSA), casein or solutions of powdered milk. This allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of antibodies in the biological sample onto the surface.

[0072] After binding of antigenic material to the well, coating with a non-reactive material to reduce background, and washing to remove unbound material, the immobilizing surface is contacted with the antisera or clinical or biological sample to be tested in a manner conducive to immune complex (antigen/antibody) formation. Such conditions preferably include diluting the sample with diluents such as BSA, solution or phosphate buffered saline (PBS)/Tween®. These added agents also tend to assist in the reduction of nonspecific background. The layered biological sample preparation is then allowed to incubate in the well for from about 1 to about 4 hr, at temperatures preferably on the order of about 25° to about 37° C. Following incubation with the diluted or undiluted biological sample, the antisera-contacted surface is washed so as to remove non-immunocomplexed material. A preferred washing procedure includes washing with a solution such as PBS/Tween®.

[0073] Following formation of specific immunocomplexes between the test sample and the bound antigen, and subsequent washing, the occurrence and even amount of immunocomplex formation may be determined by subjecting same to a second antibody having specificity for the first. To provide a detecting means, the second antibody will preferably have an associated enzyme that will generate a color development upon incubating with an appropriate chromogenic substrate. Thus, for example, one will desire to contact and incubate the antisera-bound surface with a urease, alkaline phosphatase or peroxidase-conjugated anti-human IgG for a period of time and under conditions which favor the development of immunocomplex formation (e.g., incubation for 2 hr at room temperature in a PBS-containing solution such as PBS/Tween®).

[0074] After incubation with the second enzyme-tagged antibody, and subsequent to washing to remove unbound material, the amount of label is quantified by incubation with

a chromogenic substrate such as urea and bromocresol purple or 2,2'-azino-di-(3-ethyl-benzthiazoline)-6-sulfonic acid (ABTS) and H₂O₂, in the case of peroxidase as the enzyme label. Quantitation is then achieved by measuring the degree of color generation, e.g., using a visible spectrum spectrophotometer.

[0075] Alternatively, the ELISA assay may be performed where antibodies that bind immunologically to *Borrelia* Vls antigenic sequences are immobilized onto a selected surface. After binding of the antibody to the surface, coating with a non-reactive material to reduce background, and washing to remove unbound material, the immobilizing surface is contacted with the biological sample to be tested in a manner conducive to immune complex (antigen/antibody) formation. Following formation of specific immunocomplexes between the test sample and the bound antibody, and subsequent washing, immunocomplex formation may be determined using a second, labeled antibody. This approach enables the detection of an antigen in a biological sample.

D. Epitopic Core Sequences

[0076] The present invention is also directed to protein or peptide compositions, free from total cells and other peptides, which comprise a purified protein or peptide which incorporates an epitope that is immunologically cross-reactive with one or more anti-*Borrelia* VMP-like antibodies.

[0077] As used herein, the term "incorporating an epitope (s) that is immunologically cross-reactive with one or more anti-VMP-like antibodies" is intended to refer to a peptide or protein antigen which includes a primary, secondary or tertiary structure similar to an epitope located within a *Borrelia* VMP-like polypeptide. The level of similarity will generally be to such a degree that polyclonal antibodies directed against the *Borrelia* VMP-like polypeptide will also bind to, react with, or otherwise recognize, the cross-reactive peptide or protein antigen. Various immunoassay methods may be employed in conjunction with such antibodies, such as, for example, Western blotting, ELISA, RIA, and the like, all of which are known to those of skill in the art.

[0078] The identification of *Borrelia* VMP-like epitopes, and/or their functional equivalents, suitable for use in vaccines is a relatively straightforward matter. For example, one may employ the methods of Hopp, as taught in U.S. Pat. No. 4,554,101, incorporated herein by reference, which teaches the identification and preparation of epitopes from amino acid sequences on the basis of hydrophilicity. The methods described in several other papers, and software programs based thereon, can also be used to identify epitopic core sequences (see, for example, U.S. Pat. No. 4,554,101). The amino acid sequence of these "epitopic core sequences" may then be readily incorporated into peptides, either through the application of peptide synthesis or recombinant technology.

[0079] Preferred peptides for use in accordance with the present invention will generally be on the order of about 5 to about 50 amino acids in length, and more preferably about 8 to about 40 amino acids in length. Such peptides may be isolated or comprised within a larger polypeptide. It is proposed that shorter antigenic *Borrelia* VMP-like-derived peptide sequences will provide advantages in certain circumstances, for example, in the preparation of vaccines or in immunologic detection assays. Exemplary advantages include the ease of preparation and purification, the relatively low cost and improved reproducibility of production, and advantageous biodistribution.

[0080] It is proposed that particular advantages of the present invention may be realized through the preparation of synthetic peptides which include modified and/or extended epitopic/immunogenic core sequences which result in a “universal” epitopic peptide directed to *Borrelia* VMP-like and *Borrelia* VMP-like-related sequences. It is proposed that these regions represent those which are most likely to promote T-cell or B-cell stimulation in an animal, and, hence, elicit specific antibody production in such an animal.

[0081] An epitopic core sequence, as used herein, is a relatively short stretch of amino acids that is “complementary” to, and therefore will bind, antigen binding sites on vls protein-specific antibodies. Additionally or alternatively, an epitopic core sequence is one that will elicit antibodies that are cross-reactive with antibodies directed against the peptide compositions of the present invention. It will be understood that in the context of the present disclosure, the term “complementary” refers to amino acids or peptides that exhibit an attractive force towards each other. Thus, certain epitope core sequences of the present invention may be operationally defined in terms of their ability to compete with or perhaps displace the binding of the desired protein antigen with the corresponding protein-directed antisera.

[0082] In general, the size of the polypeptide antigen is not believed to be particularly crucial, so long as it is at least large enough to carry the identified core sequence or sequences. The smallest useful core sequence expected by the present disclosure would generally be on the order of about 5 amino acids in length, with sequences on the order of 8 or 25 being more preferred. Thus, this size will generally correspond to the smallest peptide antigens prepared in accordance with the invention. However, the size of the antigen may be larger where desired, so long as it contains a basic epitopic core sequence.

[0083] The identification of epitopic core sequences is known to those of skill in the art, for example, as described in U.S. Pat. No. 4,554,101, incorporated herein by reference, which teaches the identification and preparation of epitopes from amino acid sequences on the basis of hydrophilicity. Moreover, numerous computer programs are available for use in predicting antigenic portions of proteins. Computerized peptide sequence analysis programs (e.g., DNASTar® software, DNASTar, Inc., Madison, Wis.) may also be useful in designing synthetic *Borrelia* VMP-like peptides and peptide analogs in accordance with the present disclosure. In addition, epitope mapping may be performed, in which overlapping peptides corresponding to all regions of the protein are synthesized and tested for reactivity with antibodies directed against vls sequences. Reactivity of serum from animals or humans infected with Lyme disease *Borrelia*, and nonreactivity with serum from animals or patients that do not have Lyme disease would help to define those peptides that react sensitively and specifically with antibodies against Lyme disease *Borrelia*.

[0084] An epitopic core sequence may be comprised within a larger polypeptide. For example, an epitopic core sequence of the present invention may be comprised in a larger polypeptide, which also comprises a moiety that is useful for anchoring the polypeptide to the selected surface. The anchoring moiety may be an amino acid sequence. These polypeptides would be particularly useful in the various immunoassay methods of the present invention. In a particular example, a peptide or polypeptide of the present invention may have a cysteine added at one end of the amino

acid sequence to permit the addition of biotin. The biotinylated peptides or polypeptides could then be captured on streptavidin-coated surfaces. Those of skill in the art would know how to identify which polypeptides react sensitively and specifically with antibodies against Lyme disease *Borrelia*. For example, reactivity of serum from animals or humans infected with Lyme disease *Borrelia*, and nonreactivity with serum from animals or patients that do not have Lyme disease would help to define those polypeptides that react sensitively and specifically with antibodies against Lyme disease *Borrelia*.

[0085] Syntheses of epitopic sequences, or peptides which include an antigenic epitope within their sequence, are readily achieved using conventional synthetic techniques such as the solid phase method (e.g., through the use of commercially available peptide synthesizer such as an Applied Biosystems Model 430A Peptide Synthesizer). Peptide antigens synthesized in this manner may then be aliquoted in predetermined amounts and stored in conventional manners, such as in aqueous solutions or, even more preferably, in a powder or lyophilized state pending use.

[0086] In general, due to the relative stability of peptides, they may be readily stored in aqueous solutions for fairly long periods of time if desired, e.g., up to six months or more, in virtually any aqueous solution without appreciable degradation or loss of antigenic activity. However, where extended aqueous storage is contemplated it will generally be desirable to include agents including buffers such as Tris or phosphate buffers to maintain a pH of about 7.0 to about 7.5. Moreover, it may be desirable to include agents which will inhibit microbial growth, such as sodium azide or Merthiolate. For extended storage in an aqueous state it will be desirable to store the solutions at 4° C., or more preferably, frozen. Of course, where the peptides are stored in a lyophilized or powdered state, they may be stored virtually indefinitely, e.g., in metered aliquots that may be rehydrated with a predetermined amount of water (preferably distilled) or buffer prior to use.

E. Antibodies

[0087] Means for preparing and characterizing antibodies are well known in the art. An antibody can be a polyclonal or a monoclonal antibody.

[0088] The methods for generating monoclonal antibodies (mAbs) generally begin along the same lines as those for preparing polyclonal antibodies. Briefly, a polyclonal antibody is prepared by immunizing an animal with an immunogenic composition in accordance with the present invention and collecting antisera from that immunized animal. A wide range of animal species can be used for the production of antisera. Typically the animal used for production of anti-antisera is a rabbit, a mouse, a rat, a hamster, a guinea pig or a goat. Because of the relatively large blood volume of rabbits, a rabbit is a preferred choice for production of polyclonal antibodies.

[0089] As is well known in the art, a given composition may vary in its immunogenicity. It is often necessary therefore to boost the host immune system, as may be achieved by coupling a peptide or polypeptide immunogen to a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin or rabbit serum albumin can also be used as carriers. Means for conjugating a polypeptide to a carrier protein are well

known in the art and include glutaraldehyde, m-maleimido-benzoyl-N-hydroxysuccinimide ester, carbodiimide and bis-biazotized benzidine.

[0090] As is also well known in the art, the immunogenicity of a particular immunogen composition can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Exemplary and preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvant and aluminum hydroxide adjuvant.

[0091] The amount of immunogen composition used in the production of polyclonal antibodies varies upon the nature of the immunogen as well as the animal used for immunization. A variety of routes can be used to administer the immunogen (subcutaneous, intramuscular, intradermal, intravenous and intraperitoneal). The production of polyclonal antibodies may be monitored by sampling blood of the immunized animal at various points following immunization. A second, booster, injection may also be given. The process of boosting and titering is repeated until a suitable titer is achieved. When a desired level of immunogenicity is obtained, the immunized animal can be bled and the serum isolated and stored, and/or the animal can be used to generate mAbs.

[0092] mAbs may be readily prepared through use of well-known techniques, such as those exemplified in U.S. Pat. No. 4,196,265, incorporated herein by reference. Typically, this technique involves immunizing a suitable animal with a selected immunogen composition, e.g., a purified or partially purified LCRF protein, polypeptide or peptide. The immunizing composition is administered in a manner effective to stimulate antibody producing cells. Rodents such as mice and rats are preferred animals, however, the use of rabbit, sheep, or frog cells is also possible. The use of rats may provide certain advantages, but mice are preferred, with the BALB/c mouse being most preferred as this is most routinely used and generally gives a higher percentage of stable fusions.

[0093] Following immunization, somatic cells with the potential for producing antibodies, specifically B-lymphocytes (B-cells), are selected for use in the mAb generating protocol. These cells may be obtained from biopsied spleens, tonsils or lymph nodes, or from a peripheral blood sample. Spleen cells and peripheral blood cells are preferred, the former because they are a rich source of antibody-producing cells that are in the dividing plasmablast stage, and the latter because peripheral blood is easily accessible. Often, a panel of animals will have been immunized and the spleen of animal with the highest antibody titer will be removed and the spleen lymphocytes obtained by homogenizing the spleen with a syringe. Typically, a spleen from an immunized mouse contains approximately 5×10^7 to 2×10^8 lymphocytes.

[0094] The antibody-producing B lymphocytes from the immunized animal are then fused with cells of an immortal myeloma cell, generally one of the same species as the animal that was immunized. Myeloma cell lines suited for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the desired fused cells (hybridomas).

[0095] Any one of a number of myeloma cells may be used, as are known to those of skill in the art. For example, where the immunized animal is a mouse, one may use P3-X63/Ag8, X63-Ag8.653, NS1/1-Ag41, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XX0 Bul; for rats, one may use R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210; and U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6 are all useful in connection with human cell fusions.

[0096] One preferred murine myeloma cell is the NS-1 myeloma cell line (also termed P3-NS-1-Ag4-1), which is readily available from the NIGMS Human Genetic Mutant Cell Repository by requesting cell line repository number GM3573. Another mouse myeloma cell line that may be used is the 8-azaguanine-resistant mouse murine myeloma SP2/0 non-producer cell line.

[0097] Methods for generating hybrids of antibody-producing spleen or lymph node cells and myeloma cells usually comprise mixing somatic cells with myeloma cells in a 2:1 ratio, though the ratio may vary from about 20:1 to about 1:1, respectively, in the presence of an agent or agents (chemical or electrical) that promote the fusion of cell membranes. Fusion methods using Sendai virus have been described, and those using polyethylene glycol (PEG), such as 37% (v/v) PEG. The use of electrically induced fusion methods is also appropriate.

[0098] Fusion procedures usually produce viable hybrids at low frequencies, about 1×10^{-6} to 1×10^{-8} . However, this does not pose a problem, as the viable, fused hybrids are differentiated from the parental, unfused cells (particularly the unfused myeloma cells that would normally continue to divide indefinitely) by culturing in a selective medium. The selective medium is generally one that contains an agent that blocks the de novo synthesis of nucleotides in the tissue culture media. Exemplary and preferred agents are aminopterin, methotrexate, and azaserine. Aminopterin and methotrexate block de novo synthesis of both purines and pyrimidines, whereas azaserine blocks only purine synthesis. Where aminopterin or methotrexate is used, the media is supplemented with hypoxanthine and thymidine as a source of nucleotides (HAT medium). Where azaserine is used, the media is supplemented with hypoxanthine.

[0099] The preferred selection medium is HAT. Only cells capable of operating nucleotide salvage pathways are able to survive in HAT medium. The myeloma cells are defective in key enzymes of the salvage pathway, e.g., hypoxanthine phosphoribosyl transferase (HPRT), and they cannot survive. The B-cells can operate this pathway, but they have a limited life span in culture and generally die within about two weeks. Therefore, the only cells that can survive in the selective media are those hybrids formed from myeloma and B-cells.

[0100] This culturing provides a population of hybridomas from which specific hybridomas are selected. Typically, selection of hybridomas is performed by culturing the cells by single-clone dilution in microtiter plates, followed by testing the individual clonal supernatants (after about two to three weeks) for the desired reactivity. The assay should be sensitive, simple and rapid, such as radioimmunoassays, enzyme immunoassays, cytotoxicity assays, plaque assays, dot immunobinding assays, and the like.

[0101] The selected hybridomas would then be serially diluted and cloned into individual antibody-producing cell lines, which clones can then be propagated indefinitely to

provide mAbs. The cell lines may be exploited for mAb production in two basic ways. A sample of the hybridoma can be injected (often into the peritoneal cavity) into a histocompatible animal of the type that was used to provide the somatic and myeloma cells for the original fusion. The injected animal develops tumors secreting the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can then be tapped to provide mAbs in high concentration. The individual cell lines could also be cultured *in vitro*, where the mAbs are naturally secreted into the culture medium from which they can be readily obtained in high concentrations. mAbs produced by either means may be further purified, if desired, using filtration, centrifugation and various chromatographic methods such as HPLC or affinity chromatography.

F. Immunoprecipitation

[0102] The antibodies of the present invention are particularly useful for the isolation of antigens by immunoprecipitation. Immunoprecipitation involves the separation of the target antigen-antibody complexes from a complex mixture, and is used to discriminate or isolate minute amounts of protein. For the isolation of membrane proteins cells must be solubilized into detergent micelles. Nonionic detergents are preferred, since other agents, such as bile salts, precipitate at acid pH or in the presence of bivalent cations.

[0103] In an alternative embodiment the antibodies of the present invention are useful for the close juxtaposition of two antigens. This is particularly useful for increasing the localized concentration of antigens, e.g., enzyme-substrate pairs.

G. Western Blots

[0104] The compositions of the present invention will find great use in immunoblot or western blot analysis. The anti-*Borrelia* VMP-like antibodies may be used as high-affinity primary reagents for the identification of proteins immobilized onto a solid support matrix, such as nitrocellulose, nylon or combinations thereof. In conjunction with immunoprecipitation, followed by gel electrophoresis, these may be used as a single step reagent for use in detecting antigens against which secondary reagents used in the detection of the antigen cause an adverse background. This is especially useful when the antigens studied are immunoglobulins (precluding the use of immunoglobulins binding bacterial cell wall components), the antigens studied cross-react with the detecting agent, or they migrate at the same relative molecular weight as a cross-reacting signal.

[0105] Immunologically-based detection methods for use in conjunction with Western blotting include enzymatically-, radiolabel-, or fluorescently-tagged secondary antibodies against the toxin moiety are considered to be of particular use in this regard.

H. Vaccines

[0106] An important aspect of the invention is the recognition that *Borrelia* VMP-like sequences recombine at the *vlsE* site, with the result that antigenic variation is virtually limitless. Multiclonal populations therefore can exist in an infected patient so that immunological defenses are severely tested if not totally overwhelmed. Thus there is now the opportunity to develop more effective combinations of

immunogens for protection against *Borrelia* infections or as preventive inoculations such as in the form of cocktails of multiple antigenic variants based on a series of combinatorial VMP-like antigens.

[0107] VMP-like protein preparations may be administered in several ways, either locally or systemically in pharmaceutically acceptable formulations. Amounts appropriate for administration are determined on an individual basis depending on such factors as age and sex of the subject, as well as physical condition and weight. Such determinations are well within the skill of the practitioner in the medical field.

[0108] Other methods of administration may include injection of *Borrelia* VMP-like DNAs into vaccine recipients (human or animal) driven by an appropriate promoter such as CMV, (so called DNA vaccines). Such preparations could be injected subcutaneously or intramuscularly, administered orally, or introduced into the skin on metal particles propelled by high-pressure gas. DNA vaccination techniques are currently well past the initial development stage and have shown promise as vaccination strategies.

[0109] The present invention contemplates vaccines for use in both active and passive immunization embodiments. Immunogenic compositions, proposed to be suitable for use as a vaccine, may be prepared most directly from immunogenic *Borrelia* VMP-like peptides prepared in a manner disclosed herein. Preferably the antigenic material is extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle.

[0110] The preparation of vaccines which contain *Borrelia* VMP-like peptide or polypeptide sequences as active ingredients is generally well understood in the art, as exemplified by U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all incorporated herein by reference. Typically, such vaccines are prepared as injectables. Either as liquid solutions or suspensions: solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccines.

[0111] Vaccines may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Vaccines may also be administered orally. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10 to about 95% of active ingredient, preferably about 25 to about 70%.

[0112] The *Borrelia* VMP-like-derived peptides or polypeptides of the present invention may be formulated into the vaccine as neutral or salt forms. It is anticipated that many VMP-like-derived peptides or polypeptides with different sequences could be incorporated into a single vaccine, in effect producing a combinatorial vaccine. Pharmaceutically-

acceptable salts, include the acid addition salts (formed with the free amino groups of the peptide) and those which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

[0113] The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including, e.g., the capacity of the individual's immune system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by subsequent inoculations or other administrations.

[0114] The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application on a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection or the like. The dosage of the vaccine will depend on the route of administration and will vary according to the size of the host.

[0115] Various methods of achieving adjuvant effect for the vaccine includes use of agents such as aluminum hydroxide or phosphate (alum), commonly used as about 0.05 to about 0.1% solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol®) used as an about 0.25% solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between about 70° to about 101° C. for a 30-second to 2-minute period, respectively. Aggregation by reactivating with pepsin treated (Fab) antibodies to albumin, mixture with bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of Gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA®) used as a block substitute may also be employed.

[0116] In many instances, it will be desirable to have multiple administrations of the vaccine, usually not exceeding six vaccinations, more usually not exceeding four vaccinations and preferably one or more, usually at least about three vaccinations. The vaccinations will normally be at from two to twelve week intervals, more usually from three to five week intervals. Periodic boosters at intervals of 1-5 years, usually three years, will be desirable to maintain protective levels of the antibodies. The course of the immunization may be followed by assays for antibodies for the supernatant antigens. The assays may be performed by labeling with conventional labels, such as radionucleotides, enzymes, fluorescents, and the like. These techniques are well known and may be found in a wide variety of patents, such as U.S. Pat. Nos. 3,791,932; 4,174,384 and 3,949,064, as illustrative of these types of assays.

I. Nucleic Acids

[0117] The present invention provides the nucleotide sequences of the vls gene in *B. garinii* and *B. afzelii*. It is contemplated that the isolated nucleic acids of the present invention may be put under the control of a promoter. The promoter may be the promoter that is naturally associated with the vls gene or it may be a recombinant or heterologous promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with a DNA segment encoding a *Borrelia* VMP-like peptide in its natural environment. Such promoters may include promoters normally associated with other genes, and/or promoters isolated from any viral, prokaryotic (e.g., bacterial), eukaryotic (e.g., fungal, yeast, plant, or animal) cell.

[0118] Naturally, it will be important to employ a promoter that effectively directs the expression of the DNA segment in the cell type, organism, or even animal, chosen for expression. The use of promoter and cell type combinations for protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook et al., 2001. The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins or peptides. Appropriate promoter/expression systems contemplated for use in high-level expression include, but are not limited to, the *Pichia* expression vector system (Pharmacia LKB Biotechnology), a baculovirus system for expression in insect cells, or any suitable yeast or bacterial expression system.

[0119] In connection with expression embodiments to prepare recombinant proteins and peptides, it is contemplated that longer DNA segments will most often be used, with DNA segments encoding the entire peptide sequence being most preferred. However, it will be appreciated that the use of shorter DNA segments to direct the expression of *Borrelia* VMP-like peptides or epitopic core regions, such as may be used to generate anti-*Borrelia* VMP-like antibodies, also falls within the scope of the invention. DNA segments that encode *Borrelia* VMP-like peptide antigens from about 10 to about 100 amino acids in length, or more preferably, from about 20 to about 80 amino acids in length, or even more preferably, from about 30 to about 70 amino acids in length are contemplated to be particularly useful.

[0120] In addition to their use in directing the expression of *Borrelia* VMP-like peptides of the present invention, the nucleic acid sequences contemplated herein also have a variety of other uses. For example, they also have utility as probes or primers in nucleic acid hybridization embodiments. As such, it is contemplated that nucleic acid segments that comprise a sequence region that consists of at least about a 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, an about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 nucleotide long contiguous DNA segment of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66,

SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96 will find particular utility. Longer contiguous identical or complementary sequences, e.g., those of about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300, 400, 500, (including all intermediate lengths) and those up to and including full-length sequences will also be of use in certain embodiments.

[0121] The ability of such nucleic acid probes to specifically hybridize to *Borrelia* VMP-like-encoding sequences will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are envisioned, including the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructions.

[0122] Nucleic acid molecules having sequence regions consisting of contiguous nucleotide stretches of about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300, 400, 500 or more, identical or complementary to the DNA sequence of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96, are particularly contemplated as hybridization probes for use in, e.g., Southern and Northern blotting. Smaller fragments will generally find use in hybridization embodiments, wherein the length of the contiguous complementary region may be varied, such as between about 10-14 and up to about 100 nucleotides, but larger contiguous complementary stretches may be used, according to the length complementary sequences one wishes to detect.

[0123] The use of a hybridization probe of about 14 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 14 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design

nucleic acid molecules having gene-complementary stretches of about 15 to about 20 contiguous nucleotides, or even longer where desired.

[0124] Of course, fragments may also be obtained by other techniques such as, e.g., by mechanical shearing or by restriction enzyme digestion. Small nucleic acid segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of nucleic acid reproduction technology, such as PCR™, by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

[0125] Accordingly, the nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of DNA fragments. Depending on the application envisioned, one will desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the hybrids, e.g., conditions of high stringency where one will select relatively low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50° C. to about 70° C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating *Borrelia* VMP-like-encoding DNA segments. Detection of DNA segments via hybridization is well-known to those of skill in the art, and the teachings of U.S. Pat. Nos. 4,965,188 and 5,176,995 (each incorporated herein by reference) are exemplary of the methods of hybridization analyses.

[0126] Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template or where one seeks to isolate *Borrelia* VMP-like-encoding sequences from related species, functional equivalents, or the like, less stringent hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ conditions such as about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20° C. to about 55° C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

[0127] In certain embodiments, it will be advantageous to employ nucleic acid sequences of the present invention in combination with an appropriate means, such as a label, for determining hybridization. A wide variety of appropriate indicator means are known in the art, including fluorescent, radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of giving a detectable signal. In preferred embodiments, one will likely desire to employ a fluorescent label or an enzyme tag, such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other

environmental undesirable reagents. In the case of enzyme tags, colorimetric indicator substrates are known that can be employed to provide a means visible to the human eye or spectrophotometrically, to identify specific hybridization with complementary nucleic acid-containing samples.

[0128] In general, it is envisioned that the hybridization probes described herein will be useful both as reagents in solution hybridization as well as in embodiments employing

and so would be expected to be structurally and functionally equivalent to VMP-like proteins isolated from a spirochete. However, substitutions of preferred human codons may improve expression in the human host, thereby improving the efficiency of potential DNA vaccines. This method may also be useful in achieving improved expression of the recombinant VMP-like protein in *E. coli* or any of a variety of prokaryotic and eukaryotic cells.

TABLE 2

Codon Frequency in <i>Homo sapiens</i>											
Codon	v^b	Total # ^a	Codon	v^b	Total # ^a	Codon	v^b	Total # ^a	Codon	v^b	Total # ^a
UUU	16.6	72711	UCU	14.0	62953	UAU	12.3	55039	UGU	9.5	42692
UUC	21.4	95962	UCC	17.7	79482	UAC	17.0	76480	UGC	12.8	57368
UUA	6.3	28202	UCA	10.7	48225	UAA	0.7	2955	UGA	1.2	5481
UUG	11.5	51496	UCG	4.4	19640	UAG	0.5	2181	UGG	13.5	59982
CUU	11.7	52401	CCU	16.7	74975	CAU	9.6	43193	CGU	4.6	20792
CUC	19.5	87696	CCC	20.0	89974	CAC	14.6	65533	CGC	11.0	49507
CUA	6.3	28474	CCA	16.2	72711	CAA	11.4	51146	CGA	5.9	26551
CUG	40.6	182139	CCG	6.9	30863	CAG	33.7	151070	CGG	11.3	50682
AUU	15.7	70652	ACU	12.8	57288	AAU	16.6	74401	AGU	11.1	49894
AUC	23.7	106390	ACC	21.1	94793	AAC	21.1	94725	AGC	19.1	85754
AUA	6.7	30139	ACA	14.7	66136	AAA	23.2	104221	AGA	10.8	48369
AUG	22.6	101326	ACG	6.7	30059	AAG	33.9	152179	AGG	10.9	48882
GUU	10.6	47805	GCU	18.7	83800	GAU	22.0	98712	GCU	11.2	50125
GUC	15.6	70189	GCC	29.2	130966	GAC	27.0	121005	GGC	24.0	107571
GUA	6.6	29659	GCA	15.3	68653	GAA	27.8	124852	GGA	16.9	75708
GUG	30.0	134750	GCG	7.5	33759	GAG	40.8	182943	GGG	16.7	74859

Coding GC 52.96%

1st letter GC 55.98%

2nd letter GC 42.29%

3rd letter GC 60.60%

^aTotal 4489120

^b v = Frequency per 1000

a solid phase. In embodiments involving a solid phase, the test DNA (or RNA) is adsorbed or otherwise affixed to a selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to specific hybridization with selected probes under desired conditions. The selected conditions will depend on the particular circumstances based on the particular criteria required (depending, for example, on the G+C content, type of target nucleic acid, source of nucleic acid, size of hybridization probe, etc.). Following washing of the hybridized surface so as to remove non-specifically bound probe molecules, specific hybridization is detected, or even quantitated, by means of the label.

[0129] Isolated nucleic acids encoding vls or vls-related genes are contemplated to be particularly useful in connection with this invention. Any recombinant vls combining any of the vlsE expression site loci and/or silent vls cassette would likewise be very useful with the methods of the invention.

[0130] Isolation of the DNA encoding VMP-like polypeptides allows one to use methods well known to those of skill in the art, and as herein described, to make changes in the codons for specific amino acids such that the codons are “preferred usage” codons for a given species. Thus for example, preferred codons will vary significantly for bacterial species as compared with mammalian species; however, there are preferences even among related species. Shown below is a preferred codon usage table for humans. Isolation of spirochete DNA encoding VMP-like proteins will allow substitutions for preferred human codons, although expressed polypeptide product from human DNA is expected to be homologous to bacterial VMP-like proteins

[0131] The definition of a “VMP-like sequence” or “VMP-related gene” as used herein, is a gene that hybridizes, under relatively stringent hybridization conditions (see, e.g., Sambrook et al., 2001), to DNA sequences presently known to include related gene sequences.

[0132] To prepare a VMP-like gene segment or cDNA one may follow the teachings disclosed herein and also the teachings of any patents or scientific documents specifically referenced herein. One may obtain a rVMP- or other related-encoding DNA segments using molecular biological techniques, such as polymerase chain reaction (PCRTM) or screening of a cDNA or genomic library, using primers or probes with sequences based on the above nucleotide sequence. Such single- or double-stranded DNA segments may be readily prepared by, for example, directly synthesizing the fragments by chemical means, by application of nucleic acid reproduction technology, such as the PCRTM technology of U.S. Pat. Nos. 4,683,195 and 4,683,202 (herein incorporated by reference). The practice of these techniques is a routine matter for those of skill in the art, as taught in various scientific texts (see e.g., Sambrook et al., 2001), incorporated herein by reference. Certain documents further particularly describe suitable mammalian expression vectors, e.g., U.S. Pat. No. 5,168,050, incorporated herein by reference. The VMP-like genes and DNA segments that are particularly preferred for use in certain aspects of the present methods are those encoding VMP-like and VMP-related polypeptides.

[0133] It is also contemplated that one may clone other additional genes or cDNAs that encode a VMP-like or

VMP-related peptide, protein or polypeptide. The techniques for cloning DNA molecules, i.e., obtaining a specific coding sequence from a DNA library that is distinct from other portions of DNA, are well known in the art. This can be achieved by, for example, screening an appropriate DNA library which relates to the cloning of a vls gene such as from the variable region of that gene. The screening procedure may be based on the hybridization of oligonucleotide probes, designed from a consideration of portions of the amino acid sequence of known DNA sequences encoding related *Borrelia* proteins. The operation of such screening protocols is well known to those of skill in the art and are described in detail in the scientific literature, for example, see Sambrook et al., 2001.

[0134] Techniques for introducing changes in nucleotide sequences that are designed to alter the functional properties of the encoded proteins or polypeptides are well known in the art, e.g., U.S. Pat. No. 4,518,584, incorporated herein by reference, which techniques are also described in further detail herein. Such modifications include the deletion, insertion or substitution of bases, which may or may not result in changes in the amino acid sequence. Changes may be made to increase the activity of a protein, to increase its biological stability or half-life, to change its glycosylation pattern, and the like. All such modifications to the nucleotide sequences are encompassed by this invention.

I. Biological Functional Equivalents

[0135] Modification and changes may be made in the structure of the peptides of the present invention and DNA segments which encode them and still obtain a functional molecule that encodes a protein or peptide with desirable characteristics. The following is a discussion based upon changing the amino acids of a protein to create an equivalent, or even an improved, second-generation molecule. The amino acid changes may be achieved by changing the codons of the DNA sequence, according to the following codon table:

TABLE 3

Amino Acids		Codons					
Alanine	Ala	A	GCA	GCC	GCG	GCU	
Cysteine	Cys	C	UGC	UGU			
Aspartic acid	Asp	D	GAC	GAU			
Glutamic acid	Glu	E	GAA	GAG			
Phenylalanine	Phe	F	UUC	UUU			
Glycine	Gly	G	GGA	GGC	GGG	GGU	
Histidine	His	H	CAC	CAU			
Isoleucine	Ile	I	AUA	AUC	AUU		
Lysine	Lys	K	AAA	AAG			
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG CUU
Methionine	Met	M	AUG				
Asparagine	Asn	N	AAC	AAU			
Proline	Pro	P	CCA	CCC	CCG	CCU	
Glutamine	Gln	Q	CAA	CAG			
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG CGU
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU	
Valine	Val	V	GUA	GUC	GUG	GUU	
Tryptophan	Trp	W	UGG				
Tyrosine	Tyr	Y	UAC	UAU			

[0136] For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of anti-

bodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated by the inventors that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

[0137] In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art. It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

[0138] Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics, these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

[0139] It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, i.e., still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

[0140] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Pat. No. 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

[0141] As detailed in U.S. Pat. No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (−0.5 $\pm$ 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−1.5); leucine (−1.8); isoleucine (−1.8); tyrosine (−2.3); phenylalanine (−2.5); tryptophan (−3.4).

[0142] It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

[0143] As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydro-

phobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions which take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

J. Site-Specific Mutagenesis

[0144] Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent proteins or peptides, through specific mutagenesis of the underlying DNA. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 1 to 10 residues on both sides of the junction of the sequence being altered.

[0145] In general, the technique of site-specific mutagenesis is well known in the art, as exemplified by various publications. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

[0146] In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

[0147] The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis is provided as a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

H. Expression of VMP-Like Proteins

[0148] A particular aspect of this invention provides novel ways in which to utilize VMP-like DNA segments and

recombinant vectors comprising vls DNA segments. As is well known to those of skill in the art, many such vectors are readily available, one particular detailed example of a suitable vector for expression in mammalian cells is that described in U.S. Pat. No. 5,168,050, incorporated herein by reference. However, there is no requirement that a highly purified vector be used, so long as the coding segment employed encodes a VMP-like protein and does not include any coding or regulatory sequences that would have an adverse effect on cells. Therefore, it will also be understood that useful nucleic acid sequences may include additional residues, such as additional non-coding sequences flanking either of the 5' or 3' portions of the coding including, for example, promoter regions, or may include various internal sequences, i.e., introns, which are known to occur within genes.

[0149] After identifying an appropriate VMP-encoding gene or DNA molecule, it may be inserted into any one of the many vectors currently known in the art, so that it will direct the expression and production of the VMP-like protein when incorporated into a host cell. In a recombinant expression vector, the coding portion of the DNA segment is positioned under the control of a promoter. The promoter may be in the form of the promoter which is naturally associated with a VMP-encoding gene, as may be obtained by isolating the 5' non-coding sequences located upstream of the coding segment, for example, using recombinant cloning and/or PCR™ technology, in connection with the compositions disclosed herein.

[0150] The use of recombinant promoters to achieve protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook et al., (2001).

[0151] For the expression of VMP-like proteins, once a suitable (full-length if desired) clone or clones have been obtained, whether they be cDNA based or genomic, one may proceed to prepare an expression system for the recombinant preparation of VMP-like proteins. The engineering of DNA segment(s) for expression in a prokaryotic or eukaryotic system may be performed by techniques generally known to those of skill in recombinant expression. It is believed that virtually any expression system may be employed in the expression of VMP-like proteins.

[0152] VMP-like proteins may be successfully expressed in eukaryotic expression systems, however, it is also envisioned that bacterial expression systems may be preferred for the preparation of VMP-like proteins for all purposes. The DNA or cDNA encoding VMP-like proteins may be separately expressed in bacterial systems, with the encoded proteins being expressed as fusions with beta-galactosidase, ubiquitin, *Schistosoma japonicum* glutathione S-transferase, green fluorescent protein, polyhistidine and the like. It is believed that bacterial expression will ultimately have advantages over eukaryotic expression in terms of ease of use and quantity of materials obtained thereby.

[0153] It is proposed that transformation of host cells with DNA segments encoding VMP-like proteins will provide a convenient means for obtaining VMP-like peptides. Both cDNA and genomic sequences are suitable for eukaryotic expression, as the host cell will, of course, process the genomic transcripts to yield functional mRNA for translation into protein.

[0154] It is similarly believed that almost any eukaryotic expression system may be utilized for the expression of

VMP-like proteins, e.g., baculovirus-based, glutamine synthase-based or dihydrofolate reductase-based systems could be employed. However, in preferred embodiments, it is contemplated that plasmid vectors incorporating an origin of replication and an efficient eukaryotic promoter, as exemplified by the eukaryotic vectors of the pCMV series, such as pCMV5, will be of most use.

[0155] For expression in this manner, one would position the coding sequences adjacent to and under the control of the promoter. It is understood in the art that to bring a coding sequence under the control of such a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame of the protein between about 1 and about 50 nucleotides "downstream" of (i.e., 3' of) the chosen promoter.

[0156] Where eukaryotic expression is contemplated, one will also typically desire to incorporate into the transcriptional unit which includes VMP-like protein, an appropriate polyadenylation site (e.g., 5'-AATAAA-3') if one was not contained within the original cloned segment. Typically, the poly A addition site is placed about 30 to 2000 nucleotides "downstream" of the termination site of the protein at a position prior to transcription termination.

[0157] Translational enhancers may also be incorporated as part of the vector DNA. Thus the DNA constructs of the present invention should also preferably contain one or more 5' non-translated leader sequences which may serve to enhance expression of the gene products from the resulting mRNA transcripts. Such sequences may be derived from the promoter selected to express the gene or can be specifically modified to increase translation of the RNA. Such regions may also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic gene sequence.

[0158] Such "enhancer" sequences may be desirable to increase or alter the transcription of translational efficiency of the resultant mRNA. The present invention is not limited to constructs where the enhancer is derived from the native 5'-nontranslated promoter sequence, but may also include non-translated leader sequences derived from other non-related promoters such as other enhancer transcriptional activators or genes.

[0159] It is contemplated that virtually any of the commonly employed host cells can be used in connection with the expression of VMPs in accordance herewith. Examples include cell lines typically employed for eukaryotic expression such as 239, AtT-20, HepG2, VERO, HeLa, CHO, WI 38, BHK, COS-7, RIN and MDCK cell lines.

[0160] It is contemplated that VMP-like protein may be "overexpressed", i.e., expressed in increased levels relative to its natural expression in *Borrelia* cells, or even relative to the expression of other proteins in a recombinant host cell containing VMP-encoding DNA segments. Such overexpression may be assessed by a variety of methods, including radio-labeling and/or protein purification. However, simple and direct methods are preferred, for example, those involving SDS/PAGE and protein staining or Western blotting, followed by quantitative analyses, such as densitometric scanning of the resultant gel or blot. A specific increase in the level of the recombinant protein or peptide in comparison to the level in natural VMP-producing animal cells is indicative of overexpression, as is a relative abundance of the specific protein in relation to the other proteins produced by the host cell and, e.g., visible on a gel.

[0161] As used herein, the term "engineered" or "recombinant" cell is intended to refer to a cell into which a recombinant gene, such as a gene encoding a VMP-like peptide has been introduced. Therefore, engineered cells are distinguishable from naturally occurring cells which do not contain a recombinantly introduced gene. Engineered cells are thus cells having a gene or genes introduced through the hand of man. Recombinantly introduced genes will either be in the form of a cDNA gene (i.e., they will not contain introns), a copy of a genomic gene, or will include genes positioned adjacent to a promoter not naturally associated with the particular introduced gene.

[0162] It will be understood that recombinant VMP-like proteins may differ from naturally produced VMP-like proteins in certain ways. In particular, the degree of post-translational modifications, such as, for example, lipidation, glycosylation and phosphorylation may be different between the recombinant VMP-like and the VMP-like polypeptide purified from a natural source, such as *Borrelia*.

[0163] After identifying an appropriate DNA molecule by any or a combination of means as described above, the DNA may then be inserted into any one of the many vectors currently known in the art and transferred to a prokaryotic or eukaryotic host cell where it will direct the expression and production of the so-called "recombinant" version of the protein. The recombinant host cell may be selected from a group consisting of *S. mutans*, *E. coli*, *S. cerevisiae*, *Bacillus* sp., *Lactococci* sp., *Enterococci* sp., or *Salmonella* sp. In certain preferred embodiments, the recombinant host cell will have a recA phenotype.

[0164] Where the introduction of a recombinant version of one or more of the foregoing genes is required, it will be important to introduce the gene such that it is under the control of a promoter that effectively directs the expression of the gene in the cell type chosen for engineering. In general, one will desire to employ a promoter that allows constitutive (constant) expression of the gene of interest. The use of these constitutive promoters will ensure a high, constant level of expression of the introduced genes. The level of expression from the introduced genes of interest can vary in different clones, probably as a function of the site of insertion of the recombinant gene in the chromosomal DNA. Thus, the level of expression of a particular recombinant gene can be chosen by evaluating different clones derived from each transfection study; once that line is chosen, the constitutive promoter ensures that the desired level of expression is permanently maintained. It may also be possible to use promoters that are subject to regulation, such as those regulated by the presence of lactose analog or by the expression of bacteriophage T7 DNA polymerase.

[0165] Technology for introduction of DNA into cells is well-known to those of skill in the art. Five general methods for delivering a gene into cells have been described: (1) chemical methods; (2) physical methods such as microinjection, electroporation and the gene gun; (3) viral vectors; (4) receptor-mediated mechanisms; and (5) direct injection of purified DNA into human or animals.

G. Liposomes and Nanocapsules

[0166] The formation and use of liposomes is generally known to those of skill in the art (see for example, Couvreur et al., *Pharm Res.*: 8(9), 1079-86, 1991, which describes the use of liposomes and nanocapsules in the targeted antibiotic therapy of intracellular bacterial infections and diseases).

Recently, liposomes were developed with improved serum stability and circulation half-times of substances, including DNA (Gabizon and Papahadjopoulos, *Proc Natl Acad Sci USA*, 85(18): 6949-53, 1988; Allen and Chonn, *FEBS Lett.*, 223(1): 42-6, 1987). The following is a brief description of this and other DNA delivery modes.

[0167] Nanocapsules can generally entrap compounds in a stable and reproducible way (Henry-Michelland et al., *Int J Pharm.*, 35: 121-127, 1987). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μm) should be designed using polymers able to be degraded in vivo. Biodegradable polyalkyl-cyanoacrylate nanoparticles that meet these requirements are contemplated for use in the present invention, and such particles may be easily made, as described (Couvreur, *J. Pharm Belg.*, 39(4): 249-54, 1984; Couvreur et al., *Bull Mem Acad R Med Belg.*, 143(7-9): 378-88, 1988).

[0168] Liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs)). MLVs generally have diameters ranging from 25 μm to 4 μm . Sonication of MLVs results in the formation of small unilamellar vesicles (SUVs) with diameters in the range of 200 to 500 $\text{\AA}$, containing an aqueous solution in the core.

[0169] The following information may be utilized in generating liposomal formulations. Phospholipids can form a variety of structures other than liposomes when dispersed in water, depending on the molar ratio of lipid to water. At low ratios the liposome is the preferred structure. The physical characteristics of liposomes depend on pH, ionic strength and the presence of divalent cations. Liposomes can show low permeability to ionic and polar substances, but at elevated temperatures undergo a phase transition which markedly alters their permeability. The phase transition involves a change from a closely packed, ordered structure, known as the gel state, to a loosely packed, less-ordered structure, known as the fluid state. This occurs at a characteristic phase-transition temperature and results in an increase in permeability to ions, sugars and drugs.

[0170] Liposomes interact with cells via four different mechanisms: Endocytosis by phagocytic cells of the reticuloendothelial system such as macrophages and neutrophils; adsorption to the cell surface, either by nonspecific weak hydrophobic or electrostatic forces, or by specific interactions with cell-surface components; fusion with the plasma cell membrane by insertion of the lipid bilayer of the liposome into the plasma membrane, with simultaneous release of liposomal contents into the cytoplasm; and by transfer of liposomal lipids to cellular or subcellular membranes, or vice versa, without any association of the liposome contents. It often is difficult to determine which mechanism is operative and more than one may operate at the same time.

L. Pharmaceutical Compositions

[0171] The pharmaceutical compositions disclosed herein may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compounds may be incorporated with excipients and used in the form of ingestible tablets,

buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of the unit. The amount of active compounds in such therapeutically useful compositions is such that a suitable dosage will be obtained.

[0172] The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compounds sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

[0173] The active compounds may also be administered parenterally or intraperitoneally. Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0174] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0175] Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the

appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0176] As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

[0177] The phrase “pharmaceutically acceptable” refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human. The preparation of an aqueous composition that contains a protein as an active ingredient is well understood in the art. Typically, such compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified.

[0178] The composition can be formulated in a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

[0179] Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms such as injectable solutions, drug release capsules and the like.

[0180] For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, “Remington’s Pharmaceutical Sciences” 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine

the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

EXAMPLES

[0181] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Experimental Procedures

[0182] Bacterial Strains

[0183] *B. garinii* Ip90 was initially isolated from ticks collected in eastern Russia (Kriuchechnikov et al., 1988). *B. afzelii* ACAI was cultured from a patient in Sweden with acrodermatitis chronica atrophicans (Asbrink et al., 1984). Both strains were graciously provided by Dr. Alan Barbour, University of California at Irvine School of Medicine, and had been passed through C3H/HeN mice to assure infectivity. Strains were passaged in vitro fewer than 5 times following mouse infection.

[0184] DNA Cloning and Sequencing

[0185] Plasmid DNA was purified from the *Borrelia* strains as described previously (Purser and Norris, 2000). λ DASH II libraries of plasmid DNA fragments were prepared as described by Zhang et al. (Zhang et al., 1997), with minor modifications. Thirty micrograms of plasmid DNA was treated with 30 units of mung bean nuclease at 30° C. for 30 min to hydrolyze hairpin loops in telomeres, and an EcoRI linker (5'-CCGGAATTCCGG-3'; SEQ. ID. NO:107) was then ligated to the treated plasmid DNA using T₄ DNA ligase at 15° C. overnight. This preparation was then digested to completion with EcoRI, and the resulting DNA fragments were fractionated by agarose gel electrophoresis. EcoRI-treated DNA fragments ranging in size from 8 kb to 25 kb were used to create libraries in EcoRI pre-treated λ DASH II vector arms as described in the manufacturer’s instructions (Stratagene, La Jolla, Calif., USA). Recombinant phages were screened by plaque hybridization using *B. burgdorferi* B31 vls silent cassette clone pJRZ53 (Zhang et al., 1997) as probe; hybridization with pJRZ53 was confirmed by secondary phage plaque screening as well as Southern blot hybridization. Selected phage clones were expanded, phage were purified, and DNA was prepared by standard techniques. The λ phage clones Ip90.1A1 and ACAI.2A1, each containing a 15 kb *Borrelia* DNA insert, were selected for analysis.

[0186] To sequence the DNA insert of Ip90.1A1, the phage DNA was digested with EcoRI and HindIII and a 6 kb EcoRI/HindIII fragment containing vls-like sequence was then cloned into pBluescript II SK(-) (Stratagene). The plasmid DNA of the pBluescript clone was digested with EcoRI and HindIII, and the 6 kb DNA fragment was isolated

by agarose gel electrophoresis followed by electroelution, partially digested with DNase I and cloned into EcoRV treated pBluescript II SK (–) to create random DNase I library as described previously (Zhang et al., 1997). Clones with insert DNA ranging in size from 500 to 1,000 bp from the DNase I library were selected for sequencing using primers specific for the vector T7 and T3 sequences. To facilitate sequencing of the ACAI.2A1 clone, the phage DNA was treated with XbaI and EcoRI, and one 8 kb EcoRI/XbaI fragment containing vls-like sequence was isolated from an agarose gel. This 8 kb EcoRI/XbaI fragment was digested separately with RsaI and PstI and then cloned into pBluescript II SK (–) to generate RsaI and PstI libraries. Clones from both libraries were selected for sequencing at the Department of Microbiology and Molecular Genetics Sequencing Facility. Primer walking and PCR (see below) were utilized as needed to fill gaps, establish clone order, and confirm and extend the sequences. DNA sequences were assembled using DNASTAR software (DNASTAR, Inc., Madison, Wis.).

[0187] Southern Hybridization

[0188] Fifty nanograms of DNA was digested with the indicated restriction enzymes, subjected to agarose electrophoresis in 1× TAE buffer at 100V for 2 hr, and transferred to Amersham Hybond N⁺ membranes using standard alkaline transfer techniques. Hybridization with pJRZ53 as probe was performed by enhanced chemiluminescence techniques following the manufacturer's protocol (Amersham Gene Images, Amersham, Piscataway, N.J., USA).

[0189] PCR and RT-PCR

[0190] PCR was utilized to amplify vls sequences beyond the end of the 8 kb EcoRI/XbaI fragment from ACAI, and thereby extend the sequence beyond the cloned region. The specific primer 4540 (5'-CCA GCA AAC AAC TTC CCC GCC-3'—SEQ ID NO:21), based on a variable region, and the nonspecific primer 4548 (5'-ATC CTT AAA CTC CGC CCC ATC ATC-3'—SEQ ID NO:22), based on an invariant 5' region of the vls silent cassettes of ACAI, were used as primers. Primer 4545 (5'-GAG TGC TGT GGA GAG TGC TGT TGA TGA G-3'—SEQ ID NO:23), based on the direct repeat sequence, was also used in some PCR studies. *B. afzelii* ACAI plasmid DNA was used as the template in these reactions.

[0191] RT-PCR was used to detect transcription of vlsE in *B. garinii* Ip90 and *B. afzelii* ACAI. Forward primer 4587 (5'-GGG GAT AAA GGG GAT TGT TGAT GCT GC-3'—SEQ ID NO:24) and reverse primer 4588 (5'-GCA AAC TGC CCA TCC TTA GCC ATT CC-3'—SEQ ID NO:25) were designed based on the invariable regions of vls silent cassettes of Ip90; the forward primer 4470 (5'-AAG GGG ATT GCG AAG GGG ATA AAG G-3'—SEQ ID NO:26) and reverse primer 4471 (5'-TTA GCA GCAAACCTT CCA TCC TTA GCC-3'—SEQ ID NO:27) were used for ACAI. Total RNA was isolated from late log-phase cultures of Ip90 and ACAI using an RNA purification kit (Amersham). RT-PCR was carried out using the Promega Access RT-PCR kit according to manufacturer's instructions. Briefly, reverse transcription was carried out for 50 min at 48° C. followed by an initial denaturation at 94° C. for 3 min, and 30 cycles consisting of denaturation at 94° C. for 30 sec, annealing at 68° C. for 1.5 min, and extension at 68° C. for 1.5 min.

[0192] Cloning and Sequencing vlsE RT-PCR Products

[0193] As mentioned above, both *B. afzelii* ACAI and *B. garinii* Ip90 used in these studies were first cloned by colony

formation and then passaged through mice. To determine whether vlsE sequence variation was present following mouse infection, *B. afzelii* ACAI was grown from a frozen stock and cloned by colony formation on BSKY plates (Dever et al., 1992). RT-PCR of individual clones was performed as described in a previous section, and cDNA was ligated into pCR 2.1 TOPO TA cloning vector (Invitrogen, Carlsbad, Calif., USA). Each vlsE variant was sequenced with the M13 forward and reverse primers. *B. garinii* Ip90 RNA was isolated from an uncloned population following mouse infection, and thus contained a mixture of variants. RT-PCR and cDNA cloning were performed using the method described for ACAI. Sequences were aligned with the multiple alignment program (Smith et al., 1996). The alignment output was formatted using Boxshade 3.21 (Hofmann and Baron, 1996).

[0194] Accession Numbers

[0195] The sequence of the vls silent cassette region of *B. afzelii* ACAI is provided at the United States National Center for Biomedical Information with GenBank accession number AY100628 (SEQ ID: NO:57). The *B. garinii* Ip90 silent cassette region is listed as AY100633 (SEQ ID NO:28). The RT-PCR product sequences obtained are listed as AY100629-AY100632 (SEQ ID: NOS:5-12) and AY100634-AY100637 (SEQ ID NOS:13-20) for ACAI and Ip90, respectively.

Example 2

[0196] Identification of Vls Loci in *B. garinii* Ip90 and *B. afzelii* ACAI

[0197] Hybridization with the *B. burgdorferi* B31 vls silent cassette sequence in recombinant plasmid pJRZ53 was used as a means of identifying the plasmids and DNA fragments containing vls sequences in *B. garinii* Ip90 and *B. afzelii* ACAI. The pJRZ53 probe hybridized exclusively to plasmids with an approximate size of 28 kb in both ACAI and Ip90. Following treatment of plasmid preparations with restriction enzymes, the major hybridizing DNA segments were identified as a 15 kb EcoRI fragment of ACAI DNA and a 20 kb EcoRI fragment of Ip90 plasmid DNA. Libraries of plasmid DNA EcoRI fragments were prepared in Lambda Dash II using a technique that permits the cloning of telomere-containing as well as internal fragments through treatment of the hairpin loop telomeres with mung bean nuclease followed by ligation with EcoRI linkers (Zhang et al., 1997). The phage libraries were screened by hybridization with pJRZ53, and clones Ip90.1A1 and ACAI.2A1, each containing 15 kb of insert DNA, were used for further analysis.

Example 3

Organization of Vls Silent Cassette Loci

[0198] The overall organization of the vls silent cassette loci of Ip90 and ACAI is shown in FIG. 1. As was the case in *B. burgdorferi* B31, the silent cassette loci in each strain was composed of a contiguous array of multiple cassettes. The loci in Ip90 and ACAI consisted largely of contiguous, uninterrupted open reading frames, with one frameshift present at the 3' end of cassette 9 in ACAI. The B31 vls silent cassette locus contained one stop codon and two frame shifts (Zhang et al., 1997).

Example 4

Structure of the Ip90 Vls Silent Cassette Locus

[0199] In Ip90, the vls array consisted of 11 regions with homology to the vls cassettes of B31 (FIG. 1A). With the exception of the junctions at vls3/4 and vls6/7, the 11 vls silent cassettes are flanked by 18 bp direct repeat sequences in the 6 kb region. However, several of these cassettes (vls1, vls4, vls6, and vls11) were truncated (189 to 288 bp in length) relative to the other, full-length cassettes ranging in size from 573 to 594 bp. By comparison with the vls expression cassette of B31, cassette 1 is truncated at the 3' region, containing only 92 amino acid codons; cassette 4 lacks 125 codons in its 5' region; cassette 6 contains only 89 codons and is missing most of the 3' region; and cassette 11 has 86 codons, but is missing the 3' region. A portion of the silent cassette locus from the last 3 bp of cassette 5 to the first 165 bp of cassette 8 is identical to the P7-1 clone previously characterized by Liang et al. (Liang and Philipp, 1999) (FIG. 1A). The 3' end of the Ip90 silent cassette locus possessed a truncated pseudogene of a conserved hypothetical protein belonging to gene family 144 of *B. burgdorferi* B31 (TIGR, 2002).

[0200] The 5' end of the locus also contained a region homologous to the 5', unique (non-cassette) portion of B31 expression site, vlsE (FIG. 1A). However, this gene segment is lacking a promoter region and the first 59 codons of vlsE, and also contains segments that are non-homologous to B31 vlsE. Therefore, this 'vlsE-like' sequence appears to be a pseudogene, although it is in frame with the cassette 1 of the vls silent cassette array and could conceivably encode a vlsE-like product. It is of interest to note that vlsE of *B. burgdorferi* B31 is located close to the telomere of Ip28-1, but is oriented in the opposite direction (i.e. is transcribed toward the telomere) relative to the vlsE-like sequence of Ip90. In addition, the reading frame of the vls silent cassette array in Ip90 runs away from, rather than toward (as is the case with the silent cassettes in B31), the nearest telomere (FIG. 1) (Zhang et al., 1997). Therefore, the B31 and Ip90 versions of the silent cassette loci have likely undergone large-scale rearrangements during evolution from a common ancestral organism, and it is unlikely that the Ip90 vlsE-like pseudogene evolved directly from a functional telomeric copy of vlsE. Based on other evidence, we believe that a functional vlsE gene is located elsewhere on the 28 kb plasmid of Ip90 (see below).

[0201] Portions of several vls silent cassettes from *Borrelia garinii* strain A87S were published previously (Wang et al., 2001). Each putative silent cassette in the longest available A87S sequence (GenBank Accession No. AF274070) was compared to its corresponding cassette among the Ip90 silent cassettes. The A87S sequence shared only 63 to 68% nucleotide identity to Ip90 sequences, and amino acid similarity ranged from 51 to 57%. An amino acid alignment between the A87S and Ip90 silent cassettes reveals that the heterogeneity exists largely within invariable region 1 (IR1), found upstream of VR-I (data not shown). There are also considerable differences in IR4 and IR6, but to a lesser extent when compared to IR1. The sequence differences between the vls silent cassettes sequences of Ip90 and A87S indicates that a considerable degree of heterogeneity exists among vls sequences within this species, as also appears to be the case with *Borrelia burgdorferi* strains.

[0202] An unusual feature of the Ip90 telomere region upstream of the vls cassettes is the presence of a set of 6 complete and 1 partial copies of a 41 bp direct repeat sequence. The telomere itself was identified by its location in the lambda clone insert next to the EcoRI linker used to clone mung bean nuclease-treated telomere regions. Because mung bean nuclease potentially could remove terminal nucleotides as well as disrupting the hairpin loop 5'-3' bond, it is not known whether this sequence represents the absolute end of the telomere sequence. The telomeric repeat sequences (TRS) begin 52 bp from the end of the telomere sequence, and are present as six 41-bp repeats (TRS-A through TRS-F) followed by a 32-bp truncated version of the 41-bp sequence (TRS-G) in a contiguous array. These direct repeats differ at only one position in TRS-B, and are otherwise identical. The telomeric direct repeat has no significant homology with vls sequences or any other *Borrelia* sequence reported previously. Although the direct repeats obviously arose through duplication events, their origin and significance are unknown at this time.

Example 5

Structure of the ACAI Vls Silent Cassette Locus

[0203] The overall arrangement of the *B. afzelii* ACAI vls silent cassette locus is depicted in FIG. 1B. Unlike Ip90 and B31, the ACAI vls locus was located on an internal EcoRI fragment of a 28-kb linear plasmid, and its location relative to the plasmid telomeres is not known. The ACAI vls locus contains 13 complete and 1 partial silent cassettes and each cassette is also flanked by an 18 bp direct repeat sequence. Twelve of the cassettes appear to represent 'full-length' sequences (ranging from 591 to 630 bp in length), whereas cassette 11 contains an internal deletion and cassette 14 has an internal deletion and a short, 3' truncation relative to the other cassette sequences (FIG. 1B). The 3' end of the silent cassette locus is demarcated by a complete copy of a conserved hypothetical protein gene belonging to gene family 57 of *B. burgdorferi* B31 (TIGR, 2002). We were unable to obtain additional sequence 5' of cassette 1, and it is possible that additional vls sequences are localized upstream of the region we have characterized thus far.

Example 6

Direct Repeats in the Silent Cassette Loci

[0204] In *B. burgdorferi* B31, both the central cassette of vlsE and the homologous vls silent cassettes are flanked by a 17 bp direct repeat sequence (5'-TGAGGGGGCTAT-TAAGG-3' (SEQ ID NO:106)). This sequence is generally well-conserved in the vlsE expression site and the silent cassettes; it is absent from the 5'-truncated cassette 1, and only 10 of 17 nucleotides are present at the junction between vls9 and vls10 (Zhang et al., 1997). Based on the location and high degree of conservation of the 17 bp direct repeat, it was hypothesized previously that these sequences may play an important role in the vls gene conversion process. However, the 17 bp sequence is not highly conserved in the *B. garinii* Ip90 and *B. afzelii* ACAI vls silent cassette sequences (data not shown). A comparison of 17 bp consensus sequences from Ip90 and ACAI to the B31 17 bp sequence shows that the Ip90 and ACAI sequences are more similar to each other than to the B31 sequence. Nevertheless,

the higher degree of variability in the Ip90 and ACAI 17 bp sequences compared to the B31 sequence suggests that the 17 bp sequence is not as important in the gene conversion process as previously thought (Zhang et al., 1997).

Example 7

Similarity of Vls Silent Cassette Loci

[0205] Alignment of the vls cassette sequences from Ip90, ACAI, and B31 indicates a high degree of sequence conservation both within and between each strain (FIG. 2). The Ip90 cassettes share 90 to 97% nucleotide sequence identity with one another, whereas the ACAI silent cassettes have from 84 to 91% nucleotide sequence identity (data not shown). The Ip90 vls silent cassettes are also highly homologous with *B. burgdorferi* vls sequences; for example, sequence identities with the B31 allele vlsE1 (Zhang et al., 1997) range from 64% to 73% on the nucleotide level and from 53% to 62% in predicted amino acid sequence (FIG. 2A). The identities between the ACAI vls silent cassettes and B31 vlsE1 likewise range from 65% to 73% on the nucleotide level and from 50% to 65% in predicted amino acid sequence (FIG. 2B). Each complete silent cassette of Ip90 and ACAI contains six variable regions interspersed by six invariable regions similar to those found in the vls sequences of B31 (FIG. 2).

[0206] SEQ ID NO:28 is the *B. garinii* Ip90 vls locus silent cassette nucleic acid sequence. SEQ ID NO:30 is a translation of an upstream open reading frame of SEQ ID NO:28, which is contiguous with the open reading frame of the silent cassettes of the *B. garinii* Ip90 vls locus. SEQ ID NO:32 is a translation of a vlsE-like sequence of SEQ ID NO:28. SEQ ID NOS:33-54 are nucleotide and amino acid sequences of silent cassette Nos. 1-11 of the *B. garinii* Ip90 vls locus as set forth in FIG. 2B. SEQ ID NO:55 and 56 are the nucleotide and amino acid sequences of a truncated pseudogene in the *B. garinii* Ip90 vls locus with 85% similarity to amino acids 70-140 of the *B. burgdorferi* B31 ORF-10 predicted product, GenBank Accession No. AA 34908.

[0207] SEQ ID NO:57 is the *B. afzelii* ACAI vls silent cassette locus nucleic acid sequence. SEQ ID NOS:58-85 are the nucleotide and amino acid sequences of silent cassette Nos. 1-14 of the *B. afzelii* ACAI silent cassette locus as set forth in FIG. 2A. SEQ ID NOS:86 and 87 are the nucleotide and amino acid sequences of a portion of the *B. afzelii* ACAI vls silent cassette locus which encodes a member of protein family PF02414, a conserved hypothetical protein family thought to be involved in *Borrelia* plasmid partitions of replication.

Example 8

[0208] Transcription of VlsE of *B. garinii* Ip90 and *B. afzelii* ACAI

[0209] We have thus far been unsuccessful in cloning a complete vlsE expression site from either Ip90 or ACAI using a variety of approaches (data not shown). To determine whether vls expression sites are present in Ip90 and ACAI, RT-PCR was carried out using total RNA from in vitro cultured *B. garinii* Ip90 and *B. afzelii* ACAI. Primers corresponding to invariant regions in the vls silent cassette regions of each organism were utilized. We observed a positive RT-PCR result in ethidium bromide-stained agarose

gels for both *B. garinii* Ip90 and *B. afzelii* ACAI, but no products were observed if reverse transcriptase was omitted in the RT reaction (FIG. 3). The RT-PCR products containing vls-like sequence were confirmed by sequencing, confirming that both organisms have vls expression sites. In *B. burgdorferi* B31, vlsE is located only 160 bp from the vls silent cassette array (Hudson et al., 2001; Zhang et al., 1997). Based on our studies, the vls expression sites of ACAI and Ip90 do not appear to be located in close proximity to the vls silent cassettes.

Example 9

[0210] Sequence Analysis of VlsE Variants of *B. afzelii* ACAI and *B. garinii* Ip90

[0211] Both ACAI and Ip90 were passaged through mice prior to analysis. In previous studies with *B. burgdorferi* B31, extensive sequence variation due to apparent gene conversion events occurred within the vlsE cassette region during mouse infection (Zhang and Norris, 1998a, b). To determine whether similar sequence variation occurred in ACAI and Ip90, individual RT-PCR products from each mouse-passaged strain were cloned and sequenced.

[0212] An alignment of the predicted VlsE protein sequences of ACAI and Ip90 (FIG. 4) demonstrated that sequence variation occurred within each strain. Moreover, the changes observed were consistent with gene conversion involving segments of the silent cassettes, as had been seen previously with B31. As with B31, the sequence differences were predictably localized primarily within the variable regions.

[0213] Using the sequences from the silent cassettes of each organism (FIG. 2), we determined the silent cassette sequences that were most likely involved in the gene conversion events within ACAI and Ip90 vlsE genes (FIG. 4). The theoretical minimum and maximum recombination events are indicated by solid and dotted lines, respectively. In FIG. 4A, silent cassette amino acid sequences matching regions of each variant are noted for all ACAI vlsE variants except clone 2622. The variation seen in clones 2624a and 2624b can be attributed to two silent cassettes each. In clone 2624a, vls8 matched the sequence found in a portion of variable region I (VR-I) and the entire sequence within VR-II, while vls7 matched the sequence found in VR-III, VR-IV, and VR-V. In clone 2624b, vls10 matched the sequence found in a portion of VR-I and the entire sequence within VR-II and VR-III, while vls12 matched the sequence found in VR-IV and VR-V. While both vls5 and vls6 match large portions of sequence in clone 2625, it seems more likely that vls5 was exclusively involved in the gene conversion events leading to the variation seen in clone 2625 since it contains sequence identity to VR-II, VR-III, VR-IV, and VR-V. It was difficult to ascertain which silent cassettes most likely contributed to the variation seen in clone 2622. Most silent cassettes matches spanned only a few residues in clone 2622. The nature of the sequence in clone 2622 suggests that it may be an artifactual PCR product.

[0214] Minimal recombination regions, indicated by solid lines in FIG. 4, were defined as the range of a vlsE RT-PCR product sequence that matched only a single silent cassette sequence. These commonly extend over several variable regions, as was also the case with *B. burgdorferi* B31 in previous studies (Zhang et al., 1997). In some cases, there are two or more silent cassettes that contain the same sequence within the same range. Therefore, it is only pos-

sible to predict the most likely silent cassette sequences involved (Indest et al., 2001). Maximum recombination regions commonly extend from a variable region and continue into the flanking invariant region of the corresponding matching silent cassette (FIG. 4). The extension of the maximum recombination region ends at the first position of sequence non-identity between the vlsE sequence of the clone and the given silent cassette. The degree of variation appears to be less than observed previously with *B. burgdorferi* B31, but an analysis of vlsE at different times during mammalian infection (Zhang and Norris, 1998b) is required to provide an accurate measure of the kinetics.

[0215] There are two instances of what we believe to be point mutations in the Ip90 clones (FIG. 4B). The first instance lies two residues upstream of VR-II in clone 21, where there is an arginine residue not encoded in the silent cassettes. We believe a point mutation was responsible for changing the AAG codon in the silent cassettes to AGG in clone 21. The second example of a possible point mutation is the lone threonine after VR-V in clone 20. All of the silent cassette sequences possess a GCT codon at that position, while ACT is present in clone 20.

[0216] In conclusion, our results verify previous indications that both *B. garinii* and *B. afzelii* contain plasmid-encoded vls silent cassette loci similar to those of *B. burgdorferi*. In addition, RT-PCR results indicate that a vls product is expressed by both species, and that sequence variation occurs and hence may contribute to antigenic variation. Taken together, these and previous findings confirm that the vls sequence variation system is a common feature of Lyme disease *Borrelia*, and hence is likely to be important in the pathogenesis of these organisms.

Example 10

[0217] Reactivity of Sera from Human Lyme Disease Patients and Infected Mice with *Borrelia afzelii* Protein.

[0218] A recombinant DNA vector comprising a nucleotide sequence encoding the predicted amino acid sequence of the *B. afzelii* ACA-I vls cassette 13 (SEQ ID NOs:96 and 97) has been constructed. Briefly, DNA containing the coding sequence of the cassette region was amplified using a two-step polymerase chain reaction (PCR) method. During the first amplification, specific primers flanking the *B. afzelii* ACA-I vls cassette 13 (5'-CGGAATTCACCTCGCCTTACTATTATC-3' (SEQ ID NO:98) and 5'-CGG-GATCCGAGAGTGCTGTTGATGAGGTT-3' (SEQ ID NO:99)) were used with *B. afzelii* ACA-I DNA as template to amplify a fragment containing the desired cassette. Then a second PCR was performed using primers specific for the cassette region itself (5'-CGGGATCCAAGAGTGCTGTGGATGAGGCTAGCAAG-3' (SEQ ID NO:100) and 5'-TTCTGCAGCACACTCGCCTTACTATTATCATTAGC-3' (SEQ ID NO:101)) and the purified product of the first reaction as the DNA template. The two primers contained BamHI and PstI sites, respectively (underlined); the PCR product was treated with these two enzymes and ligated into the expression vector pQE30 cut with the same two enzymes. The sequence of the insert was analyzed and found to be the correct sequence. The resulting recombinant plasmid, pBA-13-1 was used to transform *E. coli* cells, and expression was induced by incubation of a transformed *E. coli* clone to 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) for 3 hours. The *E. coli* cells were lysed by sonica-

tion and centrifuged to remove cellular debris. The recombinant, His6-tagged protein (VLS-BA13) was purified by liquid chromatography over a nickel affinity column, elution of bound protein with imidazole, and further purification using a heparin-Sepharose column. The purity of the protein was determined to be >90% by sodium-dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), and the concentration determined by a Bradford protein assay.

[0219] The purified recombinant protein VLS-BA13 was tested for reactivity with antibodies from humans using a pool of sera from patients fulfilling CDC criteria for Lyme disease, acquired in the North Central United States. A pool of negative control sera was obtained from human blood donors in Houston, Tex. Enzyme-linked immunosorbent assays (ELISAs) were performed as described (Lawrenz et al., *J. Clin. Microbiol.*, 37(12): 3997-4004, 1999), except that protein and serum concentrations were varied to determine the optimal concentrations. As shown in FIG. 6, VLS-BA13 protein (50 nanograms per well) consistently yielded higher absorbance readings with the Lyme disease serum pool than with the normal serum pool, up to a serum dilution of 1:6400. Differences in absorbance between the two serum preparations (1:200 dilution) were observed with VLS-BA13 protein concentrations as low as 3.13 nanograms per well (FIG. 7). Very similar results were obtained with sera from mice infected experimentally with *Borrelia burgdorferi* and sera from uninfected mice (FIGs. C and D). Taken together, these results provide evidence that amino acid sequences corresponding to *B. afzelii* Vls protein sequences react in a specific and sensitive manner with serum antibodies from Lyme disease patients or from *B. burgdorferi* infected mice.

Example 11

[0220] Reactivity of Sera from Human Lyme Disease Patients and Infected Mice with *Borrelia garinii* Protein.

[0221] A recombinant DNA vector comprising a nucleotide sequence encoding the predicted amino acid sequence of the *B. garinii* Ip90 vls cassette 10 (SEQ ID NOs:94 and 95) has been constructed. Briefly, DNA containing the coding sequence of the cassette region was amplified using a two-step polymerase chain reaction (PCR) method. During the first amplification, specific primers flanking the *B. garinii* Ip90 vls cassette 10 (5'-CGGGATCCGCTGTTGGGAGTYGCAAC-3' (SEQ ID NO:102) and 5'-AACTGCA-GATTATCATGAGCAGCATCCTTC-3' (SEQ ID NO:103)) were used with *B. garinii* Ip90 DNA as template to amplify a fragment containing the desired cassette. Then a second PCR was performed using primers specific for the cassette region itself (5'-CGGGATCCAAGGGGACTGTTAAGAATGCTGTTG-3' (SEQ ID NO:104) and 5'-TTCTGCAGATGATTATCATGAGCAGCATCCTTCA-3' (SEQ ID NO:105)) and the purified product of the first reaction as the DNA template. The two primers contained BamHI and PstI sites, respectively (underlined); the PCR product was treated with these two enzymes and ligated into the expression vector pQE30 cut with the same two enzymes. The sequence of the insert was analyzed and found to be the correct sequence. The resulting recombinant plasmid, pBG-10-1 was used to transform *E. coli* cells, and expression was induced by incubation of a transformed *E. coli* clone to 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) for 3 hours. The *E. coli* cells were lysed by sonica-

tion and centrifuged to remove cellular debris. The recombinant, His6-tagged protein (VLS-BG10) was purified by liquid chromatography over a nickel affinity column, elution of bound protein with imidazole, and further purification using a heparin-Sepharose column. The purity of the protein was determined to be >90% by sodium-dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), and the concentration determined by a Bradford protein assay.

[0222] The purified recombinant protein VLS-BG10 was tested for reactivity with antibodies from humans using a pool of sera from patients fulfilling CDC criteria for Lyme disease, acquired in the North Central United States. A pool of negative control sera was obtained from human blood donors in Houston, Tex. Enzyme-linked immunosorbent assays (ELISAs) were performed as described (Lawrenz et al., *J. Clin. Microbiol.*, 37(12): 3997-4004, 1999), except that protein and serum concentrations were varied to determine the optimal concentrations. In the examples shown, the antigen (VLS-BG10) was used to coat the wells, and the measured parameter was the amount of antibody bound as determined by addition of either goat anti-human IgG (alkaline phosphatase conjugate) or goat anti-mouse IgG (alkaline phosphatase conjugate), followed by washing and addition of a suitable substrate. As shown in FIG. 10, VLS-BG10 protein (10 nanograms per well) consistently yielded higher absorbance readings with the Lyme disease serum pool than with the normal serum pool, up to a serum dilution of 1:6400. Differences in absorbance between the two serum preparations (1:200 dilution) were observed with VLS-BG10 protein concentrations as low as 0.031 micrograms per well (FIG. 11). Very similar results were obtained with sera from mice infected experimentally with *Borrelia burgdorferi* and sera from uninfected mice (FIGS. 12 and 13). Taken together, these results provide evidence that amino acid sequences corresponding to *B. garinii* Vls protein sequences react in a specific and sensitive manner with serum antibodies from Lyme disease patients or from *B. burgdorferi* infected mice.

[0223] All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

- [0224] The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.
- [0225] U.S. Pat. No. 3,791,932
- [0226] U.S. Pat. No. 3,949,064
- [0227] U.S. Pat. No. 4,174,384
- [0228] U.S. Pat. No. 4,196,265
- [0229] U.S. Pat. No. 4,518,584
- [0230] U.S. Pat. No. 4,554,101
- [0231] U.S. Pat. No. 4,578,770
- [0232] U.S. Pat. No. 4,596,792
- [0233] U.S. Pat. No. 4,599,230
- [0234] U.S. Pat. No. 4,599,231
- [0235] U.S. Pat. No. 4,601,903
- [0236] U.S. Pat. No. 4,608,251
- [0237] U.S. Pat. No. 4,683,195
- [0238] U.S. Pat. No. 4,683,202
- [0239] U.S. Pat. No. 5,155,022
- [0240] U.S. Pat. No. 5,168,050
- [0241] U.S. Pat. No. 5,178,859
- [0242] U.S. Pat. No. 5,187,065
- [0243] U.S. Pat. No. 5,217,872
- [0244] U.S. Pat. No. 5,246,844
- [0245] U.S. Pat. No. 5,279,938
- [0246] U.S. Pat. No. 5,283,175
- [0247] U.S. Pat. No. 5,324,630
- [0248] U.S. Pat. No. 5,385,826
- [0249] U.S. Pat. No. 5,403,718
- [0250] U.S. Pat. No. 5,434,077
- [0251] U.S. Pat. No. 5,436,000
- [0252] U.S. Pat. No. 6,437,116
- [0253] Altschul, Gish, Miller, Myers, Lipman, "Basic local alignment search tool," *J. Mol. Biol.*, 215:403-410, 1990.
- [0254] Asbrink, Hederstedt, Hovmark, "The spirochetal etiology of erythema chronicum migrans *Afzelius*," *Acta Derm. Venerol.*, 64:291-295, 1984.
- [0255] Balmelii and Piffatetti, "Analysis of the genetic polymorphism of *Borrelia burgdorferi* sensu lato by multilocus enzyme electrophoresis," *Int. J. Syst. Bacteriol.*, 46:167-172, 1996.
- [0256] Barbour, "Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent," *J. Clin. Microbiol.*, 42:475-478, 1988.
- [0257] Barbour, "Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent," *J. Clin. Microbiol.*, 26:475-478, 1988.
- [0258] Barbour, "Linear DNA of *Borrelia* species and antigenic variation," *Trends Microbiol.*, 1:236-239, 1993.
- [0259] Barbour and Garon, "Linear plasmids of the bacterium *Borrelia burgdorferi* have covalently closed ends," *Science*, 237:409-411, 1987.
- [0260] Barbour, Burman, Carter, Kitten, Bergstrom, "Variable antigen genes of the relapsing fever agent *Borrelia hermsii* are activated by promoter addition," *Mol. Microbiol.*, 5:489-493, 1991a.
- [0261] Barbour, Carter, Burman, Freitag, Garon, Bergstrom, "Tandem insertion sequence-like elements define the expression site for variable antigen genes of *Borrelia hermsii*," *Infect. Immun.*, 59:390-397, 1991b.
- [0262] Barbour et al., "Structural analysis of the variable major proteins of *Borrelia hermsii*," *J. Exp. Med.*, 158: 2127-2140, 1983.
- [0263] Barbour et al., "Variable major proteins of *Borrelia hermsii*," *J. Exp. Med.*, 156:1312-1324, 1982.
- [0264] Barstad et al., "Variable major proteins of *Borrelia hermsii*. Epitope mapping and partial sequence analysis of CNBr peptides," *J. Exp. Med.*, 161:1302-1314, 1985.
- [0265] Barthold, "Antigenic stability of *Borrelia burgdorferi* during chronic infections of immunocompetent mice," *Infect. Immun.*, 61:4955-4961, 1993.

- [0266] Barthold, Moody, Beck, "Susceptibility of laboratory rats to isolates of *Borrelia burgdorferi* from different geographic areas," *Am. J. Trop. Med. Hyg.*, 42:596-600, 1990.
- [0267] Borst and Geaves, "Programmed gene rearrangements altering gene expression," *Science*, 235:658-667, 1987.
- [0268] Borst, Bitter, McCulloch, Leeuwen, Rudenko, "Antigenic variation in malaria," *Cell*, 82:104, 1995.
- [0269] Burgdorfer, Barbour, Hayes, Benach, Grunwaldt, Davis, "Lyme disease, a tick-borne spirochetosis?," *Science*, 216: 1317-1319, 1982.
- [0270] Carroll and Gheradini, "Membrane protein variations associated with in vitro passage of *Borrelia burgdorferi*," *Infect. Immun.*, 64:392-398, 1996.
- [0271] Carter et al., "A family of surface-exposed proteins of 20 kilodaltons in the genus *Borrelia*," *Infect. Immun.*, 62:2792-2799, 1994.
- [0272] Casjens, Delange III, Ley, Rosa, Huang, "Linear chromosomes of Lyme disease agent spirochetes: genetic diversity and conservation of gene order," *J. Bacteriol.*, 177:2769-2780, 1995.
- [0273] Demolis, Mallet, Bussereau, Jacquet, "Improved strategy for large-scale DNA sequencing using DNase I cleavage for generating random subclones," *Biotechniques*, 18:453-457, 1995.
- [0274] Dever, Jorgensen, Barbour, "In vitro antimicrobial susceptibility testing of *Borrelia burgdorferi*: a microdilution MIC method and time-kill studies," *J. Clin. Microbiol.*, 30:2692-2697, 1992.
- [0275] Donelson, "Mechanisms of antigenic variation in *Borrelia hermsii* and African trypanosomes," *J. Biol. Chem.*, 270:7783-7786, 1995.
- [0276] Fuchs, Jauris, Lottspeich, Preacmursic, Wilskie, Soutschek, "Molecular analysis and expression of a *Borrelia burgdorferi* gene encoding a 22 kDa protein (pC) in *E. coli*," *Mol. Microbiol.*, 6:503-509, 1992.
- [0277] Haas and Meyer, "The repertoire of silent pilus genes in *Neisseria gonorrhoeae*: evidence for gene conversion," *Cell*, 44:107-115, 1986.
- [0278] Hagblom, Segal, Billyard, So, "Intragenic recombination leads to pilus antigenic variation in *Neisseria gonorrhoeae*," *Nature*, 315:156-158, 1985.
- [0279] Hinnebusch, Bergstrom, Barbour, "Cloning and sequence analysis of linear plasmid telomeres of the bacterium *Borrelia burgdorferi*," *Mol. Microbiol.*, 4:811-820, 1990.
- [0280] Hofman and Baron, "Boxshade 3.21 [WWW Document] accessed on the World Wide Web at isrec.isbsib.ch:8080/software/BOX_form.html, 1996.
- [0281] Hudson, Frye, Quinn, Gherardini, "Increased expression of *Borrelia burgdorferi* vlsE in response to human endothelial cell membranes," *Mol. Microbiol.*, 41:229-239, 2001.
- [0282] Hughes and Johnson, "Methylated DNA in *Borrelia* species," *J. Bacteriol.*, 172:6602-6604, 1990.
- [0283] Indest, Howell, Jacobs, School-Meker, Norris, Philipp, "Analysis of *Borrelia burgdorferi* vlsE gene expression and recombination in the tick vector," *Infect. Immun.*, 69:7083-7090, 2001.
- [0284] Johnson et al., "Infection of Syrian hamsters with Lyme disease spirochetes," *J. Clin. Microbiol.*, 20:1099-1101, 1984.
- [0285] Jonsson, Ilver, Falk, Pepose, Normark, "Sequence changes in the pilus subunit lead to tropism variation of *Neisseria gonorrhoeae* to human tissue," *Mol. Microbiol.*, 13:403-416, 1994.
- [0286] Kitten and Barbour, "Juxtaposition of expressed variable antigen genes with a conserved telomere in the bacterium *Borrelia hermsii*," *Proc. Natl. Acad. Sci. USA*, 87:6077-6081, 1990.
- [0287] Kitten and Barbour, "The relapsing fever agent *Borrelia hermsii* has multiple copies of its chromosome and linear plasmids," *Genetics*, 132:311-324, 1992.
- [0288] Koomey, Gotschlich, Robbins, Bergstrom, Swanson, "Effects of recA mutations on pilus antigenic variation and phase transitions in *Neisseria gonorrhoeae*," *Genetics*, 117:391-398, 1987.
- [0289] Kriuchechnikov, Korenberg, Shcherbakov, Kovalenskii, Levin, "Identification of *Borrelia* isolated in the USSR from *Ixodes persulcatus* Schulze ticks," *Zh Mikrobiol. Epidemiol. Immunobiol.*, 12:41-44, 1988.
- [0290] Kupsch, Knepper, Kuroki, Heuer, Meyer, "Variable opacity (Opa) outer membrane proteins account for the cell tropisms displayed by *Neisseria gonorrhoeae* for human leukocytes and epithelial cells," *EMBO. J.*, 12:641-650, 1993.
- [0291] Lambden, Robertson, Watt, "Biological properties of two distinct pilus types produced by isogenic variants of *Neisseria gonorrhoeae* P9," *J. Bacteriol.*, 141:393-396, 1980.
- [0292] Liang and Philipp, "Analysis of antibody response to invariable regions of VlsE, the variable surface antigen of *Borrelia burgdorferi*," *Infect. Immun.*, 67:6702-6706, 1999.
- [0293] Liang, Alvarez, Gu, Nowling, Ramamoorthy, Philipp, "An immunodominant conserved region within the variable domain of VlsE, the variable surface antigen of *Borrelia burgdorferi*," *J. Immunol.*, 163:5566-5573, 1999a.
- [0294] Liang, Aberer, Cinco, Gem, Hu, Lobet, Ruscio, Voet, Jr., Weynants, Philipp, "Antigenic conservation of an immunodominant invariable region of the VlsE lipoprotein among European pathogenic genospecies of *Borrelia burgdorferi* SL," *J. Infect. Dis.*, 182:1455-1462, 2000a.
- [0295] Livey, Gibbs, Schuster, Dorner, "Evidence for lateral transfer and recombination in OspC variation in Lyme disease *Borrelia*," *Mol. Microbiol.*, 18:257-269, 1995.
- [0296] Marconi, Konkell, Garon, "Variability of osp genes and gene products among species of Lyme disease spirochetes," *Infect. Immun.*, 61:2611-2617, 1993.
- [0297] Marconi, Samuels, Landry, Garon, "Analysis of the distribution and molecular heterogeneity of the ospD gene among the Lyme disease spirochetes: evidence for lateral gene exchange," *J. Bacteriol.*, 176:4572-4582, 1994.
- [0298] Margolis et al., "Homology between *Borrelia burgdorferi* OspC and members of the family of *Borrelia hermsii* variable major proteins," *Gene*, 143:105-110, 1994.
- [0299] Meier, Simon, Barbour, "Antigenic variation is associated with DNA rearrangements in a relapsing fever *Borrelia*," *Cell*, 41:403-409, 1985.
- [0300] Meyer, Mlawer, So, "Pilus expression in *Neisseria gonorrhoeae* involves chromosomal rearrangement," *Cell*, 30:45-52, 1982.

- [0301] Moody et al., "Lyme borreliosis in laboratory animals: effect of host species and in vitro passage of *Borrelia burgdorferi*," *Am. J. Trop. Med. Hyg.*, 43:87-92, 1990.
- [0302] Nassif, Lowry, Stenberg, O'Gaora, Ganji, So, "Antigenic variation of pilin regulates adhesion of *Neisseria meningitidis* to human epithelial cells," *Mol. Microbiol.*, 8:719-725, 1993.
- [0303] Norris, Carter, Howell, Barbour, "Low-passage-associated proteins of *Borrelia burgdorferi* B31: characterization and molecular cloning of OspD, a surface-exposed, plasmid-encoded lipoprotein," *Infect. Immun.*, 60:4662-4672, 1992.
- [0304] Norris et al., "High- and low-infectivity phenotypes of clonal populations of in vitro-cultured *Borrelia burgdorferi*," *Infect. Immun.*, 63:2206-2212, 1995.
- [0305] Norris et al., "Low-passage-associated proteins of *Borrelia burgdorferi* B31: characterization and molecular cloning of OspD, a surface exposed, plasmid-encoded lipoprotein," *Infect. Immun.*, 60:4662-4672, 1992.
- [0306] Persing, Mathiesen, Podzorski, Barthold, "Genetic stability of *Borrelia burgdorferi* recovered from chronically infected immunocompetent mice," *Infect. Immun.*, 62:3521-3527, 1994.
- [0307] Plasterk et al., "Transposition of structural genes to an expression sequence on a linear plasmid causes antigenic variation in the bacterium *Borrelia hermsii*," *Nature*, 318:257-263, 1985.
- [0308] Purser and Norris, "Correlation between plasmid content and infectivity in *Borrelia burgdorferi*," *Proc. Natl. Acad. Sci. USA*, 97:13865-13870, 2000.
- [0309] Restrepo and Barbour, "Antigen diversity in the bacterium *B. hermsii* through 'somatic' mutations in rearranged vmp genes," *Cell*, 78:867-876, 1994.
- [0310] Restrepo, Carter, Barbour, "Activation of a vmp pseudogene in *Borrelia hermsii*: an alternate mechanism of antigenic variation during relapsing fever," *Mol. Microbiol.*, 13:287-299, 1994.
- [0311] Restrepo, Kitten, Carter, Infante, Barbour, "Subtelomeric expression regions of *Borrelia hermsii* linear plasmids are highly polymorphic," *Mol. Microbiol.*, 6:3299-3311, 1992.
- [0312] Robertson and Meyer, "Genetic variation in pathogenic bacteria," *Trends Genet.*, 8:422-427, 1992.
- [0313] Rosa, Samuels, Hogan, Stevenson, Casjens, Tilly, "Directed insertion of a selectable marker into a circular plasmid of *Borrelia burgdorferi*," *J. Bacteriol.*, 178:5946-5953, 1996.
- [0314] Rosa, Schwan, Hogen, "Recombination between genes encoding major surface proteins A and B of *Borrelia burgdorferi*," *Mol. Microbiol.*, 6:3031-3040, 1992.
- [0315] Rudel, Van Putten, Gibbs, Haas, Meyer, "Interaction of two variable proteins (PilE and PilC) required for pilus-mediated adherence of *Neisseria gonorrhoeae* to human epithelial cells," *Mol. Microbiol.*, 6:3439-3450, 1992.
- [0316] Sadziene, Rosa, Thompson, Hogan, Barbour, "Antibody-resistant mutations of *Borrelia burgdorferi*: in vitro selection and characterization," *J. Exp. Med.*, 176:799-809, 1992.
- [0317] Sambrook, Fritsch, Maniatis, "Molecular cloning: a laboratory manual," Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press, 1989.
- [0318] Sambrook, Fritsch, Maniatis, "Molecular cloning: a laboratory manual," Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press, 2001.
- [0319] Samuels, Mach, Garon, "Genetic transformation of the Lyme disease agent *Borrelia burgdorferi* with coumarin-resistant *gyrB*," *J. Bacteriol.*, 176:6045-6049, 1994.
- [0320] Schutzer, "Lyme disease: Molecular and immunologic approaches. In: *Current communications in cell and molecular biology*," J. Inglis and J. A. Witkowski, eds. (Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press).
- [0321] Schwann et al., "Changes in infectivity and plasmid profile of the Lyme disease spirochete, *Borrelia burgdorferi*, as a result of in vitro cultivation," *Infect. Immun.*, 56:1831-1836, 1988a.
- [0322] Schwann, Burgdorfer, Schruppg, Karstens, "The urinary bladder, a consistent source of *Borrelia burgdorferi* in experimentally infected white-footed mice (*Peromyscus leucopus*)," *J. Clin. Microbiol.*, 26:893-895, 1988b.
- [0323] Schwann, Karstens, Schrupf, Simpson, "Changes in antigenic reactivity of *Borrelia burgdorferi*, the Lyme disease spirochete, during persistent infection of mice," *Can. J. Microbiol.*, 37:450-454, 1991.
- [0324] Seal, Jackson, Daniels, "Isolation of a *Pseudomonas solanacearum*-specific DNA probe by subtraction hybridization and construction of species-specific oligonucleotide primers for sensitive detection by the polymerase chain reaction," *Appl. Environ. Microbiol.*, 58:3751-3758, 1992.
- [0325] Segal, Hagblom, Seifert, So, "Antigenic variation of gonococcal pilus involves assembly of separated silent gene segments," *Proc. Natl. Acad. Sci. USA*, 83:2177-2181, 1986.
- [0326] Seifert and So, "Genetic mechanisms of bacterial antigenic variation," *Microbiol. Rev.*, 52:327-336, 1988.
- [0327] Smith, Wiese, Wojzynski, Davison, Worley, "BCM Search Launcher—an integrated interface to molecular biology data base search and analysis services available on the World Wide Web," *Genome Res.*, 6:454-462, 1996.
- [0328] Steere, "Lyme disease," *N. Engl. J. Med.*, 321:586-596, 1989.
- [0329] Stevenson, Bockenstedt, Barthold, "Expression and gene sequence of outer surface protein C of *Borrelia burgdorferi* reisolated from chronically infected mice," *Infect. Immun.*, 62:3568-3571, 1994.
- [0330] Stoenner, Dodd, Larsen, "Antigenic variation of *Borrelia hermsii*," *J. Exp. Med.*, 156:1297-1311, 1982.
- [0331] Swanson and Koomey, "Mechanisms for variation of pili and outer membrane protein II in *Neisseria gonorrhoeae*," D. E. Berg and M. M. Howe, Eds. (Washington, D.C.: American Society for Microbiology).
- [0332] Thiessen et al., "Evolution of the *Borrelia burgdorferi* outer surface protein OspC," *J. Bacteriol.*, 177:3036-3044, 1995.
- [0333] TIGR, The Institute for Genomic Research (accessed on the World Wide Web at tigr.org), 2002.
- [0334] Wainwright, Pritchard, Seifert, "A conserved DNA sequence is required for efficient gonococcal pilin antigenic variation," *Mol. Microbiol.*, 13:75-87, 1994.

- [0335] Walker, Howell, You, Hoffmaster, Heath, Weinstock, Norris, "Physical map of the genome of *Treponema pallidum* subsp. *pallidum* (Nichols)," *J. Bacteriol.*, 177:1797-1804, 1995.
- [0336] Wang, van Dam, Dankert, "Analysis of a VMP-like sequence (vls) locus in *Borrelia garinii* and Vls homologues among four *Borrelia burgdorferi* sensu lato species," *FEMS Microbiol. Lett.*, 3199:9-45, 2001.
- [0337] Wilske, Barbour, Bergstrom, Burman, Restrepo, Rosa, Schwan, Soutschek, Wallich, "Antigenic variation and strain heterogeneity in *Borrelia* spp.," *Res. Microbiol.*, 143:583-596, 1992.
- [0338] Wu and Tokunaga, "Biogenesis of lipoproteins in bacteria," *Curr. Top. Microbiol. Immunol.*, 125:127-157, 1986.
- [0339] Xu, Kodner, Coleman, Johnson, "Correlation of plasmids with infectivity of *Borrelia burgdorferi* sensu stricto type strain B31," *Infect. Immun.*, 64:3870-3876, 1996.
- [0340] Xu and Johnson, "Analysis and comparison of plasmid profile of *Borrelia burgdorferi* sensu lato strains," *J. Clin. Microbiol.*, 33:2679-2685, 1995.
- [0341] Zhang, Hardman, Barbour, Norris, "Antigenic variation in Lyme disease *Borreliae* by promiscuous recombination of VMP-like sequence cassettes," *Cell*, 89:275-285, 1997.
- [0342] Zhang and Norris, "Genetic variation of the *Borrelia burgdorferi* gene vlsE involves cassette-specific, segmental gene conversion," *Infect. Immun.*, 66:3698-3704, 1998a.
- [0343] Zhang and Norris, "Kinetics and in vivo induction of genetic variation of vlsE in *Borrelia burgdorferi*," *Infect. Immun.*, 66:3689-3697, 1998b.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 107

<210> SEQ ID NO 1

<211> LENGTH: 1227

<212> TYPE: DNA

<213> ORGANISM: *Borrelia burgdorferi*

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (75) .. (1142)

<400> SEQUENCE: 1

```

acctacactt gttaaaactc tctttttgag ttaagatgat aacttatact ttctattata      60
aggagacgat gaat atg aaa aaa att tca agt gca agt tta tta aca act      110
              Met Lys Lys Ile Ser Ser Ala Ser Leu Leu Thr Thr
              1              5              10

ttc ttt gtt ttt att aat tgt aaa agc caa gtt gct gat aag gac gac      158
Phe Phe Val Phe Ile Asn Cys Lys Ser Gln Val Ala Asp Lys Asp Asp
              15              20              25

cca aca aac aaa ttt tac caa tct gtc ata caa tta ggt aac gga ttt      206
Pro Thr Asn Lys Phe Tyr Gln Ser Val Ile Gln Leu Gly Asn Gly Phe
              30              35              40

ctt gat gta ttc aca tct ttt ggt ggg tta gta gca gag gct ttt gga      254
Leu Asp Val Phe Thr Ser Phe Gly Gly Leu Val Ala Glu Ala Phe Gly
              45              50              55              60

ttt aaa tca gat cca aaa aaa tct gat gta aaa acc tat ttt act act      302
Phe Lys Ser Asp Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Thr Thr
              65              70              75

gta gct gcc aaa ttg gaa aaa aca aaa acc gat ctt aat agt ttg cct      350
Val Ala Ala Lys Leu Glu Lys Thr Lys Thr Asp Leu Asn Ser Leu Pro
              80              85              90

aag gaa aaa agc gat ata agt agt acg acg ggg aaa cca gat agt aca      398
Lys Glu Lys Ser Asp Ile Ser Ser Thr Thr Gly Lys Pro Asp Ser Thr
              95              100              105

ggg tct gtt gga act gcc gtt gag ggg gct att aag gaa gtt agc gag      446
Gly Ser Val Gly Thr Ala Val Glu Gly Ala Ile Lys Glu Val Ser Glu
              110              115              120

ttg ttg gat aag ctg gta aaa gct gta aag aca gct gag ggg gct tca      494
Leu Leu Asp Lys Leu Val Lys Ala Val Lys Thr Ala Glu Gly Ala Ser
              125              130              135              140

agt ggt act gct gca att gga gaa gtt gtg gct gat gct gat gct gca      542
Ser Gly Thr Ala Ala Ile Gly Glu Val Val Ala Asp Ala Asp Ala Ala
              145              150              155

```

-continued

aag gtt gct gat aag gcg agt gtg aag ggg att gct aag ggg ata aag Lys Val Ala Asp Lys Ala Ser Val Lys Gly Ile Ala Lys Gly Ile Lys 160 165 170	590
gag att gtt gaa gct gct ggg ggg agt gaa aag ctg aaa gct gtt gct Glu Ile Val Glu Ala Ala Gly Gly Ser Glu Lys Leu Lys Ala Val Ala 175 180 185	638
gct gct aaa ggg gag aat aat aaa ggg gca ggg aag ttg ttt ggg aag Ala Ala Lys Gly Glu Asn Asn Lys Gly Ala Gly Lys Leu Phe Gly Lys 190 195 200	686
gct ggt gct gct gct cat ggg gac agt gag gct gct agc aag gcg gct Ala Gly Ala Ala Ala His Gly Asp Ser Glu Ala Ala Ser Lys Ala Ala 205 210 215 220	734
ggt gct gtt agt gct gtt agt ggg gag cag ata tta agt gcg att gtt Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Ser Ala Ile Val 225 230 235	782
acg gct gct gat gcg gct gag cag gat gga aag aag cct gag gag gct Thr Ala Ala Asp Ala Ala Glu Gln Asp Gly Lys Lys Pro Glu Glu Ala 240 245 250	830
aaa aat ccg att gct gct gct att ggg gat aaa gat ggg ggt gcg gag Lys Asn Pro Ile Ala Ala Ala Ile Gly Asp Lys Asp Gly Gly Ala Glu 255 260 265	878
ttt ggt cag gat gag atg aag aag gat gat cag att gct gct gct att Phe Gly Gln Asp Glu Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Ile 270 275 280	926
gct ttg agg ggg atg gct aag gat gga aag ttt gct gtg aag gat ggt Ala Leu Arg Gly Met Ala Lys Asp Gly Lys Phe Ala Val Lys Asp Gly 285 290 295 300	974
gag aaa gag aag gct gag ggg gct att aag gga gct gct gag tct gca Glu Lys Glu Lys Ala Glu Gly Ala Ile Lys Gly Ala Ala Glu Ser Ala 305 310 315	1022
gtt cgc aaa gtt tta ggg gct att act ggg cta ata gga gac gcc gtg Val Arg Lys Val Leu Gly Ala Ile Thr Gly Leu Ile Gly Asp Ala Val 320 325 330	1070
agt tcc ggg cta agg aaa gtc ggt gat tca gtg aag gct gct agt aaa Ser Ser Gly Leu Arg Lys Val Gly Asp Ser Val Lys Ala Ala Ser Lys 335 340 345	1118
gaa aca cct cct gcc ttg aat aag tgatttaatt aagtgtatgg acacgactat Glu Thr Pro Pro Ala Leu Asn Lys 350 355	1172
gccctcatga ttgaggaaat agtcgagaga tatatatact aaaagataat aaata	1227
<210> SEQ ID NO 2	
<211> LENGTH: 356	
<212> TYPE: PRT	
<213> ORGANISM: Borrelia burgdorferi	
<400> SEQUENCE: 2	
Met Lys Lys Ile Ser Ser Ala Ser Leu Leu Thr Thr Phe Phe Val Phe 1 5 10 15	
Ile Asn Cys Lys Ser Gln Val Ala Asp Lys Asp Asp Pro Thr Asn Lys 20 25 30	
Phe Tyr Gln Ser Val Ile Gln Leu Gly Asn Gly Phe Leu Asp Val Phe 35 40 45	
Thr Ser Phe Gly Gly Leu Val Ala Glu Ala Phe Gly Phe Lys Ser Asp 50 55 60	
Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Thr Thr Val Ala Ala Lys 65 70 75 80	

[illegible]

```
<210> SEQ ID NO 3
<211> LENGTH: 1141
<212> TYPE: DNA
<213> ORGANISM: Borrelia hermsii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1062)
```

<400> SEQUENCE: 3

atg	aga	aaa	aga	ata	agt	gca	ata	ata	atg	act	tta	ttt	atg	gta	tta	48
Met	Arg	Lys	Arg	Ile	Ser	Ala	Ile	Ile	Met	Thr	Leu	Phe	Met	Val	Leu	
1				5					10					15		
gta	agc	tgt	aat	agc	ggg	ggg	gtt	gcg	gaa	gat	cct	aaa	act	gtg	tat	96
Val	Ser	Cys	Asn	Ser	Gly	Gly	Val	Ala	Glu	Asp	Pro	Lys	Thr	Val	Tyr	
			20					25					30			
tta	aca	tct	ata	gct	aat	tta	ggg	aaa	gga	ttt	tta	gat	gtt	ttt	gtg	144
Leu	Thr		Ser	Ile	Ala	Asn	Leu	Gly	Lys	Gly	Phe	Leu	Asp	Val	Phe	Val
		35					40					45				

-continued

act ttt gga gat atg gtt act gga gct ttt ggt att aag gca gat act	192
Thr Phe Gly Asp Met Val Thr Gly Ala Phe Gly Ile Lys Ala Asp Thr	
50 55 60	
aag aaa agt gat ata ggg aag tat ttt act gat att gag agc act atg	240
Lys Lys Ser Asp Ile Gly Lys Tyr Phe Thr Asp Ile Glu Ser Thr Met	
65 70 75 80	
aca tca gtt aaa aag aag ttg caa gat gaa gtt gct aag aat ggt aac	288
Thr Ser Val Lys Lys Lys Leu Gln Asp Glu Val Ala Lys Asn Gly Asn	
85 90 95	
tat cca aag gta aag aca gct gtt gac gaa ttt gtt gca atc tta gga	336
Tyr Pro Lys Val Lys Thr Ala Val Asp Glu Phe Val Ala Ile Leu Gly	
100 105 110	
aag atc gag aaa gga gca aaa gaa gca tct aaa ggg gct act ggt gat	384
Lys Ile Glu Lys Gly Ala Lys Glu Ala Ser Lys Gly Ala Thr Gly Asp	
115 120 125	
gtt att att ggg aat act gtt aag aat ggt gat gct gta cct gga gaa	432
Val Ile Ile Gly Asn Thr Val Lys Asn Gly Asp Ala Val Pro Gly Glu	
130 135 140	
gca aca agt gtc aat tct ctt gtt aaa gga att aaa gaa ata gtt ggg	480
Ala Thr Ser Val Asn Ser Leu Val Lys Gly Ile Lys Glu Ile Val Gly	
145 150 155 160	
gta gtc ttg aag gaa ggt aag gca gat gct gat gct act aaa gat gat	528
Val Val Leu Lys Gly Lys Ala Asp Ala Asp Ala Thr Lys Asp Asp	
165 170 175	
agt aag aaa gat att ggt aaa tta ttt acc gca acc act gat gcg aat	576
Ser Lys Lys Asp Ile Gly Lys Leu Phe Thr Ala Thr Thr Asp Ala Asn	
180 185 190	
aga gct gat aat gcg gca gct caa gca gct gca gcg tca ata gga gca	624
Arg Ala Asp Asn Ala Ala Ala Gln Ala Ala Ala Ser Ile Gly Ala	
195 200 205	
gtg aca ggt gct gat atc ttg caa gct ata gta caa tct aag gaa aat	672
Val Thr Gly Ala Asp Ile Leu Gln Ala Ile Val Gln Ser Lys Glu Asn	
210 215 220	
cct gtt gca aat agt act gat gga att gaa aaa gca aca gat gca gct	720
Pro Val Ala Asn Ser Thr Asp Gly Ile Glu Lys Ala Thr Asp Ala Ala	
225 230 235 240	
gag att gca gtt gct cca gct aaa gat aat aaa aaa gag att aaa gat	768
Glu Ile Ala Val Ala Pro Ala Lys Asp Asn Lys Lys Glu Ile Lys Asp	
245 250 255	
gga gca aaa aaa gac gca gtt att gct gca ggc att gca ctg cga gca	816
Gly Ala Lys Lys Asp Ala Val Ile Ala Ala Gly Ile Ala Leu Arg Ala	
260 265 270	
atg gct aag aat ggt aca ttt tct att aaa aac aat gaa gat gcg gct	864
Met Ala Lys Asn Gly Thr Phe Ser Ile Lys Asn Asn Glu Asp Ala Ala	
275 280 285	
gta acg acg ata aat agt gca gca gca agc gca gtg aac aag att tta	912
Val Thr Thr Ile Asn Ser Ala Ala Ala Ser Ala Val Asn Lys Ile Leu	
290 295 300	
agc act cta ata ata gca ata agg aat aca gtt gat agt ggt tta aaa	960
Ser Thr Leu Ile Ile Ala Ile Arg Asn Thr Val Asp Ser Gly Leu Lys	
305 310 315 320	
aca ata aat gag gct ctt gct aca gtt aaa caa gaa gat aaa tct gta	1008
Thr Ile Asn Glu Ala Leu Ala Thr Val Lys Gln Glu Asp Lys Ser Val	
325 330 335	
gaa gca act aat act gca gaa gca aca act agt ggt cag caa gcg aaa	1056
Glu Ala Thr Asn Thr Ala Glu Ala Thr Thr Ser Gly Gln Gln Ala Lys	
340 345 350	

-continued

```

aac tag ttaagggttaa atataaagga taaagttatt gtaagggaag agcttttctt    1112
Asn
gtttttaatg caggaatgta gtttctctg                                     1141

```

<210> SEQ ID NO 4

<211> LENGTH: 353

<212> TYPE: PRT

<213> ORGANISM: *Borrelia hermsii*

<400> SEQUENCE: 4

```

Met Arg Lys Arg Ile Ser Ala Ile Ile Met Thr Leu Phe Met Val Leu
1          5          10          15
Val Ser Cys Asn Ser Gly Gly Val Ala Glu Asp Pro Lys Thr Val Tyr
20          25          30
Leu Thr Ser Ile Ala Asn Leu Gly Lys Gly Phe Leu Asp Val Phe Val
35          40          45
Thr Phe Gly Asp Met Val Thr Gly Ala Phe Gly Ile Lys Ala Asp Thr
50          55          60
Lys Lys Ser Asp Ile Gly Lys Tyr Phe Thr Asp Ile Glu Ser Thr Met
65          70          75          80
Thr Ser Val Lys Lys Lys Leu Gln Asp Glu Val Ala Lys Asn Gly Asn
85          90          95
Tyr Pro Lys Val Lys Thr Ala Val Asp Glu Phe Val Ala Ile Leu Gly
100         105         110
Lys Ile Glu Lys Gly Ala Lys Glu Ala Ser Lys Gly Ala Thr Gly Asp
115         120         125
Val Ile Ile Gly Asn Thr Val Lys Asn Gly Asp Ala Val Pro Gly Glu
130         135         140
Ala Thr Ser Val Asn Ser Leu Val Lys Gly Ile Lys Glu Ile Val Gly
145         150         155         160
Val Val Leu Lys Glu Gly Lys Ala Asp Ala Asp Ala Thr Lys Asp Asp
165         170         175
Ser Lys Lys Asp Ile Gly Lys Leu Phe Thr Ala Thr Thr Asp Ala Asn
180         185         190
Arg Ala Asp Asn Ala Ala Ala Gln Ala Ala Ala Ser Ile Gly Ala
195         200         205
Val Thr Gly Ala Asp Ile Leu Gln Ala Ile Val Gln Ser Lys Glu Asn
210         215         220
Pro Val Ala Asn Ser Thr Asp Gly Ile Glu Lys Ala Thr Asp Ala Ala
225         230         235         240
Glu Ile Ala Val Ala Pro Ala Lys Asp Asn Lys Lys Glu Ile Lys Asp
245         250         255
Gly Ala Lys Lys Asp Ala Val Ile Ala Ala Gly Ile Ala Leu Arg Ala
260         265         270
Met Ala Lys Asn Gly Thr Phe Ser Ile Lys Asn Asn Glu Asp Ala Ala
275         280         285
Val Thr Thr Ile Asn Ser Ala Ala Ala Ser Ala Val Asn Lys Ile Leu
290         295         300
Ser Thr Leu Ile Ile Ala Ile Arg Asn Thr Val Asp Ser Gly Leu Lys
305         310         315         320
Thr Ile Asn Glu Ala Leu Ala Thr Val Lys Gln Glu Asp Lys Ser Val
325         330         335

```

-continued

Glu Ala Thr Asn Thr Ala Glu Ala Thr Thr Ser Gly Gln Gln Ala Lys
 340 345 350

Asn

<210> SEQ ID NO 5
 <211> LENGTH: 416
 <212> TYPE: DNA
 <213> ORGANISM: *Borrelia afzelii*
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(414)

<400> SEQUENCE: 5

aag ggg att gcg aag ggg ata aag ggg att gtt gcg gct gct ggg aag 48
 Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala Gly Lys
 1 5 10 15

gct ttt ggc aag gat ggt gat gcg ctg aca ggt gtt gca aaa gct gct 96
 Ala Phe Gly Lys Asp Gly Asp Ala Leu Thr Gly Val Ala Lys Ala Ala
 20 25 30

gag aat gat gct aac aag gat gcg ggg aag ttg ttt gct ggt aag aat 144
 Glu Asn Asp Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
 35 40 45

ggt aat gct ggt gct gct gac att gcg aag gcg gct gct gct gtt act 192
 Gly Asn Ala Gly Ala Ala Asp Ile Ala Lys Ala Ala Ala Val Thr
 50 55 60

gcg gtt agt ggg gag cag ata cta aaa gct att gtt gag gcg gct ggt 240
 Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala Ala Gly
 65 70 75 80

gat gcg gat cag gcg ggt gta aag gct gat gcg gct aag aat ccg att 288
 Asp Ala Asp Gln Ala Gly Val Lys Ala Asp Ala Ala Lys Asn Pro Ile
 85 90 95

gca gct gcg att ggg act gct gat gat ggt gct gcg ttt ggt aag gat 336
 Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Ala Phe Gly Lys Asp
 100 105 110

gag atg aag aag aga aat gat aag att gtt gca gct att gtt ttg agg 384
 Glu Met Lys Lys Arg Asn Asp Lys Ile Val Ala Ala Ile Val Leu Arg
 115 120 125

ggg gtg cct aag gat gga aag ttt gct gct aa 416
 Gly Val Pro Lys Asp Gly Lys Phe Ala Ala
 130 135

<210> SEQ ID NO 6
 <211> LENGTH: 138
 <212> TYPE: PRT
 <213> ORGANISM: *Borrelia afzelii*

<400> SEQUENCE: 6

Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala Gly Lys
 1 5 10 15

Ala Phe Gly Lys Asp Gly Asp Ala Leu Thr Gly Val Ala Lys Ala Ala
 20 25 30

Glu Asn Asp Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
 35 40 45

Gly Asn Ala Gly Ala Ala Asp Ile Ala Lys Ala Ala Ala Val Thr
 50 55 60

Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala Ala Gly
 65 70 75 80

Asp Ala Asp Gln Ala Gly Val Lys Ala Asp Ala Ala Lys Asn Pro Ile

-continued

85	90	95	
Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Ala Phe Gly Lys Asp			
100	105	110	
Glu Met Lys Lys Arg Asn Asp Lys Ile Val Ala Ala Ile Val Leu Arg			
115	120	125	
Gly Val Pro Lys Asp Gly Lys Phe Ala Ala			
130	135		
 <210> SEQ ID NO 7			
<211> LENGTH: 413			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia afzelii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(411)			
 <400> SEQUENCE: 7			
aag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag			48
Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys			
1	5	10	15
gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt gca aaa gtt gct			96
Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val Ala			
20	25	30	
gat gat gat aac aag gat gcg ggg aag ttg ttt gct ggt aag aat ggt			144
Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn Gly			
35	40	45	
ggt gct ggt gct gct gat gcg att ggg aag gcg gct gct gct gtt act			192
Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Val Thr			
50	55	60	
gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gat gct gct ggt			240
Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Gly			
65	70	75	80
gct gca gct aat cag gcg ggt aaa aag gct gcg gat gct aag aat ccg			288
Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn Pro			
85	90	95	
att gcg gct gcg att ggg act gct gat gat ggg gcg gag ttt aag gat			336
Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys Asp			
100	105	110	
gat atg aag aag agt gat aat att gct gcg gct att gtt ttg agg ggg			384
Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg Gly			
115	120	125	
gtg cct aag gat gga aag ttt gct gct aa			413
Val Pro Lys Asp Gly Lys Phe Ala Ala			
130	135		
 <210> SEQ ID NO 8			
<211> LENGTH: 137			
<212> TYPE: PRT			
<213> ORGANISM: Borrelia afzelii			
 <400> SEQUENCE: 8			
Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys			
1	5	10	15
Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val Ala			
20	25	30	
Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn Gly			
35	40	45	
Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Val Thr			

-continued

50	55	60	
Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Gly			
65	70	75	80
Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn Pro			
	85	90	95
Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys Asp			
	100	105	110
Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg Gly			
	115	120	125
Val Pro Lys Asp Gly Lys Phe Ala Ala			
	130	135	
<210> SEQ ID NO 9			
<211> LENGTH: 428			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia afzelii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(426)			
<400> SEQUENCE: 9			
aag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag			48
Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys			
1	5	10	15
gct ttt ggt aag gag ggt gat gcg ctg aag gat gtt gca aaa gtt gct			96
Ala Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala			
	20	25	30
gat gag aat ggg gat aac aag gat gcg ggg aag ttg ttt gct ggt gag			144
Asp Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu			
	35	40	45
aat ggt aat gct ggt ggt gct gct gat gct gac att gcg aag gcg gct			192
Asn Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala			
	50	55	60
gct gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt			240
Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val			
65	70	75	80
gag gcg gct ggt gct gcg gat cag gcg ggt gta aag gct gag gag gct			288
Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala Glu Glu Ala			
	85	90	95
aag aat ccg att gca gct gcg att ggg act gat gat gct ggt gcg gcg			336
Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly Ala Ala			
	100	105	110
gag ttt ggt gag aat gat atg aag aag aat gat aat att gct gcg gct			384
Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile Ala Ala Ala			
	115	120	125
att gtt ttg agg ggg gtg cct aag gat gga aag ttt gct gct aa			428
Ile Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala			
	130	135	140
<210> SEQ ID NO 10			
<211> LENGTH: 142			
<212> TYPE: PRT			
<213> ORGANISM: Borrelia afzelii			
<400> SEQUENCE: 10			
Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys			
1	5	10	15
Ala Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala			

-continued

20	25	30	
Asp Glu Asn Gly Asp Asn Lys	Asp Ala Gly Lys Leu Phe	Ala Gly Glu	
35	40	45	
Asn Gly Asn Ala Gly Gly Ala	Ala Asp Ala Asp Ile	Ala Lys Ala Ala	
50	55	60	
Ala Ala Val Thr Ala Val Ser	Gly Glu Gln Ile Leu Lys	Ala Ile Val	
65	70	75	80
Glu Ala Ala Gly Ala Ala Asp	Gln Ala Gly Val Lys	Ala Glu Glu Ala	
85	90	95	
Lys Asn Pro Ile Ala Ala Ala	Ile Gly Thr Asp Asp	Ala Gly Ala Ala	
100	105	110	
Glu Phe Gly Glu Asn Asp Met	Lys Lys Asn Asp Asn	Ile Ala Ala Ala	
115	120	125	
Ile Val Leu Arg Gly Val Pro	Lys Asp Gly Lys Phe	Ala Ala	
130	135	140	
 <210> SEQ ID NO 11			
<211> LENGTH: 426			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia afzelii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (3)..(425)			
 <400> SEQUENCE: 11			
ag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag			47
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys			
1 5 10 15			
gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt aaa aca gtt gct			95
Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr Val Ala			
20 25 30			
gct gag aat gag gct aac aag gat gcg ggg aag ttg ttt gct ggt aag			143
Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys			
35 40 45			
aat ggt aat gct gat gct gct gat gct gct gac att gcg aag gcg gct			191
Asn Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys Ala Ala			
50 55 60			
ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa gct att gtt			239
Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val			
65 70 75			
gat ggt gct ggt gat gca gct aat cag gcg ggt aaa aag gct gct gag			287
Asp Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Glu			
80 85 90 95			
gct aag aat ccg att gcg gct gcg att ggg act aat gaa gct ggg gcg			335
Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala			
100 105 110			
gag ttt ggt gat gat atg aag aag aga aat gat aag att gct gcg gct			383
Glu Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala			
115 120 125			
att gtt ttg agg ggg gtg cct aag gat gga aag ttt gct gct a			426
Ile Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala			
130 135 140			
 <210> SEQ ID NO 12			
<211> LENGTH: 141			
<212> TYPE: PRT			
<213> ORGANISM: Borrelia afzelii			

-continued

<400> SEQUENCE: 12

```

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala
1      5      10      15
Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr Val Ala Ala
20     25     30
Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
35     40     45
Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys Ala Ala Gly
50     55     60
Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp
65     70     75     80
Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Glu Ala
85     90     95
Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala Glu
100    105    110
Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala Ile
115    120    125
Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala
130    135    140

```

<210> SEQ ID NO 13

<211> LENGTH: 396

<212> TYPE: DNA

<213> ORGANISM: Borrelia garinii

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (2)..(394)

<400> SEQUENCE: 13

```

g ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa      49
  Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
  1      5      10      15
ggg aag ttg aat gct gct ggt gct gag ggt acg act aac gcg gat gct      97
Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala Asp Ala
20     25     30
ggg aag ttg ttt gtg aag aat gct ggt aat gtg ggt ggt gaa gca ggt      145
Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu Ala Gly
35     40     45
gat gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag      193
Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln
50     55     60
ata tta aaa gcg att gtt gat gct gct aag gat ggt ggt gag aag cag      241
Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu Lys Gln
65     70     75     80
ggt aag aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg      289
Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly
85     90     95
ggt gcg ggt gat aat gat gct gct gcg gcg ttt gct act atg aag aag      337
Gly Ala Gly Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys
100    105    110
gat gat cag att gct gct gct atg gtt ctg agg gga atg gct aag gat      385
Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp
115    120    125
ggg cag ttt gcg      396
Gly Gln Phe
130

```

-continued

<210> SEQ ID NO 14
 <211> LENGTH: 131
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 14

Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
 1 5 10 15
 Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala Asp Ala
 20 25 30
 Gly Lys Leu Phe Val Lys Asn Ala Glu Asn Val Gly Gly Glu Ala Gly
 35 40 45
 Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln
 50 55 60
 Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu Lys Gln
 65 70 75 80
 Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly
 85 90 95
 Gly Ala Gly Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys
 100 105 110
 Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp
 115 120 125
 Gly Gln Phe
 130

<210> SEQ ID NO 15
 <211> LENGTH: 390
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (2)..(388)

<400> SEQUENCE: 15

g ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa 49
 Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
 1 5 10 15
 ggg aag ttg gat gtg gct ggt gat gct ggt gaa act aac aag gat gct 97
 Gly Lys Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala
 20 25 30
 ggg aag ttg ttt gtg aag aag aat aat gag ggt ggt gaa gca aat gat 145
 Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp
 35 40 45
 gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata 193
 Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile
 50 55 60
 tta aaa gcg att gtt gat gct gct gag ggt ggt gag aag cag ggt aag 241
 Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys
 65 70 75 80
 aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg ggt gcg 289
 Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala
 85 90 95
 ggt gat aat gat gct gct gcg gcg ttt gct act atg aag aag gat gat 337
 Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp
 100 105 110
 cag att gct act gct atg gtt ctg agg gga atg gct aag gat ggg cag 385
 Gln Ile Ala Thr Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln

-continued

115	120	125	
ttt gc			390
Phe			
<210> SEQ ID NO 16			
<211> LENGTH: 129			
<212> TYPE: PRT			
<213> ORGANISM: Borrelia garinii			
<400> SEQUENCE: 16			
Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu			
1 5 10 15			
Gly Lys Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala			
20 25 30			
Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp			
35 40 45			
Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile			
50 55 60			
Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys			
65 70 75 80			
Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala			
85 90 95			
Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp			
100 105 110			
Gln Ile Ala Thr Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln			
115 120 125			
Phe			
<210> SEQ ID NO 17			
<211> LENGTH: 390			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia garinii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (2)..(388)			
<400> SEQUENCE: 17			
g ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa			49
Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu			
1 5 10 15			
ggg agg ttg gat gtg gct ggt gat gct ggt gaa act aac aag gat gct			97
Gly Arg Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala			
20 25 30			
ggg aag ttg ttt gtg aag aag aat aat gag ggt ggt gaa gca aat gat			145
Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp			
35 40 45			
gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata			193
Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile			
50 55 60			
tta aaa gcg att gtt gat gct gct gag ggt ggt gag aag cag ggt aag			241
Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys			
65 70 75 80			
aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg ggt gcg			289
Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala			
85 90 95			
ggt gat aat gat gct gct gcg gcg ttt gct act atg aag aag gat gat			337
Gly Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp			
100 105 110			

-continued

```

cag att gct gct gct atg gtt ctg agg gga atg gct aag gat ggg cag      385
Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln
      115              120              125

```

```

ttt gc      390
Phe

```

```

<210> SEQ ID NO 18
<211> LENGTH: 129
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

```

```

<400> SEQUENCE: 18

```

```

Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
1           5           10           15
Gly Arg Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala
      20           25           30
Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp
      35           40           45
Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile
      50           55           60
Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys
      65           70           75           80
Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala
      85           90           95
Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp
      100          105          110
Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln
      115          120          125
Phe

```

```

<210> SEQ ID NO 19
<211> LENGTH: 339
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (2)..(337)

```

```

<400> SEQUENCE: 19

```

```

g ggg ata aag ggg att gtt gat gct gct ggt gaa act aac aag gat gct      49
  Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Glu Thr Asn Lys Asp Ala
    1           5           10           15
ggg aag ttg ttt gtg aag aag aat aat gag ggt ggt gaa gca aat gat      97
Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp
      20           25           30
gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata      145
Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile
      35           40           45
tta aaa gcg att gtt gat gct gct gag ggt ggt gag aag cag ggt aag      193
Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys
      50           55           60
aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg ggt aca      241
Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Thr
      65           70           75           80
aat gat aat gat gct gcg gcg ttt gct act atg aag aag gat gat cag      289
Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp Gln
      85           90           95

```

-continued

att gct gct gct atg gtt ctg agg gga atg gct aag gat ggg cag ttt	337
Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe	
100 105 110	

gc	339
----	-----

<210> SEQ ID NO 20
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: *Borrelia garinii*

<400> SEQUENCE: 20

Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Glu Thr Asn Lys Asp Ala	
1 5 10 15	
Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp	
20 25 30	
Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile	
35 40 45	
Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys	
50 55 60	
Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Thr	
65 70 75 80	
Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp Gln	
85 90 95	
Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe	
100 105 110	

<210> SEQ ID NO 21
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 21

ccagcaaaca acttccccgc c	21
-------------------------	----

<210> SEQ ID NO 22
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 22

atccttaaac tccgccccat catc	24
----------------------------	----

<210> SEQ ID NO 23
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 23

gagtgctgtg gagagtgtg ttgatgag	28
-------------------------------	----

<210> SEQ ID NO 24
<211> LENGTH: 27

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 24

ggggataaag gggattgttg atgctgc 27

<210> SEQ ID NO 25
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 25

gcaaactgcc catccttagc cattcc 26

<210> SEQ ID NO 26
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 26

aaggggattg cgaaggggat aaagg 25

<210> SEQ ID NO 27
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 27

ttagcagcaa actttccatc cttagcc 27

<210> SEQ ID NO 28
<211> LENGTH: 5897
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 28

cggaaatcaa gccaccta aaacttccc aaaagtttct caaaaaatat tatattcagc 60
agtaaattct ataagtcatt aattatttaa tactattcaa cagtaaatc tataagtcac 120
taattattta atactattca gcagtaaat ctataagtca ttaattattt aatactattc 180
agcagtaaat tctataagtc attaattatt taatactatt cagcagtaaa ttctataagt 240
cattaattat ttaatactat tcagcagtaa attctataag tcattaatta tttaatacta 300
ttcagcagta aattctataa gtcattaatt caattaggta acggattctt agatgtattc 360
acctcttttg gtggattagt tgcagatgca ttgggggtta aagctgatcc aaaaaaatct 420
gatgtaaaaa cttattttga atctctagct aaaaaattag aagaacaaa agatgggtta 480
actaagttgt ccaaaggtaa tgacgggtgat actggaaagg ctggtgatgc tgggtgggct 540
ggtggtggcg ctagtgtgc aggtggcgct ggtgggattg agggcgctat aacagagatt 600

-continued

agcaaatggt tagatgatat ggcaaaagct gctgcggaag ctgcaagtgc tgctactggt	660
aatgcagcaa ttggggatgt tgttaatggt aatggtggag cagcaaaagg tggatgatgcg	720
gagagtgtta atgggattgc taaggggata aaggggattg ttgatgctgc tgagaaggct	780
gatgcgaagg aaggaagtt ggatgtggct ggtgatgctg gtggggctgg tggtgccgct	840
ggtgctgcag gtggcgctgg tgggattgag ggcgctataa cagagattag caaatggtta	900
gatgatatgg caaaagctgc tgcggttgct gcaagtgcctg caagtgcctg tactggtaat	960
gcagcaattg gggatgttgt taatggtaat gatggagcag caaaagggtg tgatgcggcg	1020
agtgttaatg ggattgctaa ggggataaag gggattgttg atgctgctga gaaggctgat	1080
gcgaagggaag ggaagttgga tgtggctggg gatgctgggt agggtaacaa ggatgctggg	1140
aagctgtttg tgaagaagaa tgctggtgat gaggggtggg aagcaaatga tgctgggaag	1200
gctgctgctg cgggtgctgc tgttagtggg gagcagatat taaaagcgat tgttgatgct	1260
gctgaggggtg atgataagca gggtaagaag gctgcggatg ctacaaatcc gattgagggc	1320
gctattgggg gtgcggatgc ggggtgcta atgctaggcg ttaataagat gaagaaggat	1380
gatcagattg ctgctgctat ggttctgagg ggaatggcta aggatgggca gtttgccttg	1440
aaggatgatg ctgctgctca tgaagggact gtttaagaat ctgttgatat ggcaaaggcc	1500
gctgcggaag ctgcaagtgc tgcaagtgc gctactggtg gtacaacgat tggagatgtt	1560
gttaagagtg gtgaggcaaa agatggtgat gcggcgagtg ttaatgggat tgctaagggg	1620
ataaagggga ttgttgatgc tgcctgagaag gctgatgcga aggaaggga gttggatgtg	1680
gctggtgctg ctggtacgac taactgta atgtgggaag tgtttgtgaa gaataatggt	1740
aatgaggggtg gtgatgcaag tgatgctggg aaagctgctg ctgcggttgc tgctgttagt	1800
ggggagcaga tattaaaagc gattgttgat gctgctaaag atggtgataa gcaggggggtt	1860
actgatgtaa aggatgctac aaatccgatt gaggcggcta ttgggggtac aaatgataat	1920
gatgctgcgg cgtttgctac tatgaagaag gatgatcaga ttgctgctgc tatggttctg	1980
aggggaatgg ctaaggatgg gcagtttgct ttgaaggatg atgctgctaa ggatggtgat	2040
aaaacggggg ttgctgcgga tgcgtgaaaat ccgattgacg cggctattgg gggctcggat	2100
gctgatgctg cggcgcttaa taaggagggg atgaagaagg atgatcagat tgctgctgct	2160
atggttctga ggggaatggc taaggatggg cagtttgctt tgacgaataa tgctgctgct	2220
catgaaggga ctgttaagaa tgctgttgat atggcaaaag ctgctgcggt tgctgcaagt	2280
gctgctactg gcaatgcagc aattggggat gttgttaaga gtaatggtgg agcagcagca	2340
aaagtggtg atgcggcgag tgttaatggg attgctaagg ggataaaggg gattgttgat	2400
gctgctgaga aggctgatgc gaaggaaggg aagttggatg tggctggtgc tgctggtgaa	2460
actaacaagg atgctgggaa gttgtttgtg aagaagaatg gtgatgatgg tggatgatgca	2520
ggtgatgctg ggaaggctgc tgcgcgggtt gctgctgtta gtggggagca gatattaaaa	2580
gcgattgttg atgctgctaa agatggtgat aagacggggg ttactgatgt aaaggatgct	2640
acaaatccga ttgacgcggc tattgggggg agtgcggtatg ctaatgctga ggcgtttgat	2700
aagatgaaga aggatgatca gattgctgct gctatggttc tgaggggaat ggctaaggat	2760
gggcagtttg ctttgaagaa taatgatcat gataatcata aggggactgt taagaatgct	2820
gttgatatgg caaaggccgc tgaggaagct gcaagtgcctg caagtgcctg tactggtaat	2880

-continued

gcagcgattg	gggatgttgt	taagaatagt	ggggcagcag	caaaaggttg	tgaggcggcg	2940
agtgttaatg	ggattgctaa	ggggataaag	gggatgttg	atgctgctgg	aaaggctgat	3000
gcgaaggaag	ggaagtgtga	tgctactggg	gctgagggta	cgactaacgt	gaatgctggg	3060
aagttgtttg	tgaagagggc	ggctgatgat	ggtggtgatg	cagatgatgc	tggaagggct	3120
gctgctgcgg	ttgctgcaag	tgctgctact	ggtaatgcag	cgattggaga	tgtgtttaat	3180
ggtgatgtgg	caaaagcaaa	agggtggtgat	gcggcgagtg	ttaatgggat	tgctaagggg	3240
ataaagggga	ttgttgatgc	tgctgagaag	gctgatgcga	aggaagggaa	gttgaatgct	3300
gctggtgctg	agggctacgac	taacgcggat	gctgggaagt	tgtttgtgaa	gaatgctggt	3360
aatgtgggtg	gtgaagcagg	tgatgctggg	aaggctgctg	ctgcggttgc	tgctgttagt	3420
ggggagcaga	tattaaaagc	gattgttgat	gctgctaagg	atggtggtga	gaagcagggg	3480
aagaaggctg	cggatgctac	aaatccgatt	gacgcggcta	ttgggggtac	aaatgataat	3540
gatgctgctg	cggcggttgc	tactatgaag	aaggatgatc	agattgctgc	tgctatggtt	3600
ctgaggggaa	tggtcaagga	tgggcaattt	gctttgaagg	atgctgctgc	tgctcatgaa	3660
gggactgtta	agaatgctgt	tgatataata	aaggctgctg	cggaagctgc	aagtgctgca	3720
agtgtgctta	ctggttagtgc	agcaattggg	gatgttggtta	atggtaatgg	agcaacagca	3780
aaaggtggtg	atgcgaagag	tgtaaatggg	attgctaagg	ggataaaggg	gattgttgat	3840
gctgctgaga	aggctgatgc	gaaggaaggg	aagttggatg	tggtggtga	tgctggtgaa	3900
actaacaagg	atgctgggaa	gttggtttgtg	aagaacaatg	gtaatgaggg	tggtgatgca	3960
gatgatgctg	ggaaggctgc	tgctgcggtt	gctgctgtta	gtggggagca	gatattaaaa	4020
gcgattgttg	atgctgctaa	gggtggtgat	aagacgggta	agaataatgt	gaaggatgct	4080
gaaaatccga	ttgaggcggc	tattgggagt	agtgcggatg	ctgatgctgc	ggcggttaat	4140
aaggagggga	tgaagaagga	tgatcagatt	gctgctgcta	tggttctgag	gggaatggct	4200
aaggatgggc	agtttgcttt	gacgaatgat	gctgctgctc	atgaagggac	tgtaagaat	4260
gctgttggga	gtgcaacaaa	taagaccggt	gttgctttgg	ctaacttggt	tcgaaagacc	4320
gtgcaagctg	gggtgaagaa	gggtggggat	gttgtaaga	atagtgaggc	aaaagatggt	4380
gatgcggcga	gtgttaatgg	gattgctaag	gggataaagg	ggattgttga	tgctgctgag	4440
aaggctgatg	cgaaggaagg	gaagtgggat	gtggctggtg	ctgctggtga	aactaacaag	4500
gatgctggga	agttgtttgt	gaagaagaat	aatgaggggtg	gtgaagcaaa	tgatgctggg	4560
aaggctgctg	ctgcggttgc	tgctgttagt	ggggagcaga	tattaaaagc	gattgttgat	4620
gctgctaagg	atggtgatga	taagcagggt	aagaaggctg	aggatgctac	aaatccgatt	4680
gacgcggcta	ttgggggtgc	aggctcgggt	gctaatactg	ctgcggcggt	taataatatg	4740
aagaaggatg	atcagattgc	tgctgctatg	gttctgaggg	gaatggctaa	ggatgggcag	4800
tttgctttga	cgaataatgc	tcatactaata	cataagggga	ctgttaagaa	tgctgttgat	4860
atgacaaaag	ctgctgcggg	tgctgcaagt	gctgcaagtg	ctgctactgg	taatgcagca	4920
attggggatg	ttgttaatgg	taatgatgga	gcagcaaaag	gtggtgatgc	ggcgagtgtt	4980
aatgggattg	ctaaggggat	aaaggggatt	gttgatgctg	ctgagaaggc	tgatgcgaag	5040
gaaggggaagt	tgaatgtggc	tggtgctgct	ggtgctgagg	gtaacgaggc	tgctgggaag	5100
ctgtttgtga	agaagaatgc	tggtgatcat	ggtggtgaag	caggtgatgc	tggaagggct	5160

-continued

```

gctgctgctg ttgctgctgt tagtggggag cagatattaa aagcgattgt tgatgctgct 5220
aaggatgggtg gtgataagca gggtaagaag gctgaggatg ctgaaaatcc gattgacgcg 5280
gctattggga gtacgggtgc ggatgataat gctgctgagg cgtttgctac tatgaagaag 5340
gatgatcaga ttgctgctgc tatggttctg aggggaatgg ctaaggatgg gcagtttgct 5400
ttgaaggatg ctgctcatga taatcataag gggactgtta agaagtctgt tgatataata 5460
aaggctactg cggttgctgc aagtgtctgt actggtagta caacgattgg ggatgttggt 5520
aagaatgggtg aggcaaaagg tggtagggcg aagagtgtta atgggattgc taaggggata 5580
aaggggattg ttgatgctgc tggaaaggct gatgcgaagg aagggaagtt gaatgtggct 5640
ggtgctgctg gtgagggtaa cgaggctgct gggaaagctgt ttgtgtaaat tactatagga 5700
ttagaactag tgtacgatat gagtccttg gttattttgc agctgctaata gaatttgaaa 5760
taagtgaagt taaaattgct gatgttaatg gaacacattt tattgctaca aaagagaaag 5820
aatattata tgattcactt gatttaaggg ctctgggaaa aatatttgaa ataacttcaa 5880
agcgaatggt taagctt 5897

```

```

<210> SEQ ID NO 29
<211> LENGTH: 300
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(300)

```

```

<400> SEQUENCE: 29

```

```

gct att aat tat tta ata cta ttc agc agt aaa ttc tat aag tca tta 48
Val Ile Asn Tyr Leu Ile Leu Phe Ser Ser Lys Phe Tyr Lys Ser Leu
1 5 10 15

att caa tta ggt aac gga ttc tta gat gta ttc acc tct ttt ggt gga 96
Ile Gln Leu Gly Asn Gly Phe Leu Asp Val Phe Thr Ser Phe Gly Gly
20 25 30

tta gtt gca gat gca ttg ggg ttt aaa gct gat cca aaa aaa tct gat 144
Leu Val Ala Asp Ala Leu Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp
35 40 45

gta aaa act tat ttt gaa tct cta gct aaa aaa tta gaa gaa aca aaa 192
Val Lys Thr Tyr Phe Glu Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys
50 55 60

gat ggt tta act aag ttg tcc aaa ggt aat gac ggt gat act gga aag 240
Asp Gly Leu Thr Lys Leu Ser Lys Gly Asn Asp Gly Asp Thr Gly Lys
65 70 75 80

gct ggt gat gct ggt ggg gct ggt ggt ggc gct agt gct gca ggt ggc 288
Ala Gly Asp Ala Gly Gly Ala Gly Gly Gly Ala Ser Ala Ala Gly Gly
85 90 95

gct ggt ggg att 300
Ala Gly Gly Ile
100

```

```

<210> SEQ ID NO 30
<211> LENGTH: 100
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

```

```

<400> SEQUENCE: 30

```

```

Val Ile Asn Tyr Leu Ile Leu Phe Ser Ser Lys Phe Tyr Lys Ser Leu
1 5 10 15

```

-continued

```

Ile Gln Leu Gly Asn Gly Phe Leu Asp Val Phe Thr Ser Phe Gly Gly
      20              25              30

Leu Val Ala Asp Ala Leu Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp
      35              40              45

Val Lys Thr Tyr Phe Glu Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys
      50              55              60

Asp Gly Leu Thr Lys Leu Ser Lys Gly Asn Asp Gly Asp Thr Gly Lys
      65              70              75              80

Ala Gly Asp Ala Gly Gly Ala Gly Gly Gly Ala Ser Ala Ala Gly Gly
      85              90              95

Ala Gly Gly Ile
      100

```

```

<210> SEQ ID NO 31
<211> LENGTH: 102
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(102)

```

```

<400> SEQUENCE: 31

```

```

ggg ttt aaa gct gat cca aaa aaa tct gat gta aaa act tat ttt gaa      48
Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Glu
1              5              10              15

tct cta gct aaa aaa tta gaa gaa aca aaa gat ggt tta act aag ttg      96
Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys Asp Gly Leu Thr Lys Leu
      20              25              30

tcc aaa      102
Ser Lys

```

```

<210> SEQ ID NO 32
<211> LENGTH: 34
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

```

```

<400> SEQUENCE: 32

```

```

Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Glu
1              5              10              15

Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys Asp Gly Leu Thr Lys Leu
      20              25              30

Ser Lys

```

```

<210> SEQ ID NO 33
<211> LENGTH: 288
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(288)

```

```

<400> SEQUENCE: 33

```

```

gag ggc gct ata aca gag att agc aaa tgg tta gat gat atg gca aaa      48
Glu Gly Ala Ile Thr Glu Ile Ser Lys Trp Leu Asp Asp Met Ala Lys
1              5              10              15

gct gct gcg gaa gct gca agt gct gct act ggt aat gca gca att ggg      96
Ala Ala Ala Glu Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly
      20              25              30

gat gtt gtt aat ggt aat ggt gga gca gca aaa ggt ggt gat gcg gag      144

```

-continued

Asp	Val	Val	Asn	Gly	Asn	Gly	Gly	Ala	Ala	Lys	Gly	Gly	Asp	Ala	Glu	
	35						40					45				
agt	gtt	aat	ggg	att	gct	aag	ggg	ata	aag	ggg	att	gtt	gat	gct	gct	192
Ser	Val	Asn	Gly	Ile	Ala	Lys	Gly	Ile	Lys	Gly	Ile	Val	Asp	Ala	Ala	
	50					55				60						
gag	aag	gct	gat	gcg	aag	gaa	ggg	aag	ttg	gat	gtg	gct	ggt	gat	gct	240
Glu	Lys	Ala	Asp	Ala	Lys	Glu	Gly	Lys	Leu	Asp	Val	Ala	Gly	Asp	Ala	
65				70					75					80		
ggt	ggg	gct	ggt	ggc	gct	ggt	gct	gca	ggt	ggc	gct	ggt	ggg	att		288
Gly	Gly	Ala	Gly	Gly	Gly	Ala	Gly	Ala	Ala	Gly	Gly	Ala	Gly	Gly	Ile	
			85					90					95			
<210> SEQ ID NO 34																
<211> LENGTH: 96																
<212> TYPE: PRT																
<213> ORGANISM: Borrelia garinii																
<400> SEQUENCE: 34																
Glu	Gly	Ala	Ile	Thr	Glu	Ile	Ser	Lys	Trp	Leu	Asp	Asp	Met	Ala	Lys	
1			5						10					15		
Ala	Ala	Ala	Glu	Ala	Ala	Ser	Ala	Ala	Thr	Gly	Asn	Ala	Ala	Ile	Gly	
			20					25					30			
Asp	Val	Val	Asn	Gly	Asn	Gly	Gly	Ala	Ala	Lys	Gly	Gly	Asp	Ala	Glu	
	35					40					45					
Ser	Val	Asn	Gly	Ile	Ala	Lys	Gly	Ile	Lys	Gly	Ile	Val	Asp	Ala	Ala	
50						55				60						
Glu	Lys	Ala	Asp	Ala	Lys	Glu	Gly	Lys	Leu	Asp	Val	Ala	Gly	Asp	Ala	
65					70				75					80		
Gly	Gly	Ala	Gly	Gly	Gly	Ala	Gly	Ala	Ala	Gly	Gly	Ala	Gly	Gly	Ile	
			85					90						95		
<210> SEQ ID NO 35																
<211> LENGTH: 594																
<212> TYPE: DNA																
<213> ORGANISM: Borrelia garinii																
<220> FEATURE:																
<221> NAME/KEY: CDS																
<222> LOCATION: (1)..(594)																
<400> SEQUENCE: 35																
gag	ggc	gct	ata	aca	gag	att	agc	aaa	tgg	tta	gat	gat	atg	gca	aaa	48
Glu	Gly	Ala	Ile	Thr	Glu	Ile	Ser	Lys	Trp	Leu	Asp	Asp	Met	Ala	Lys	
1			5						10					15		
gct	gct	gcg	gtt	gct	gca	agt	gct	gca	agt	gct	gct	act	ggt	aat	gca	96
Ala	Ala	Ala	Val	Ala	Ala	Ser	Ala	Ala	Ser	Ala	Ala	Thr	Gly	Asn	Ala	
			20					25					30			
gca	att	ggg	gat	gtt	gtt	aat	ggt	aat	gat	gga	gca	gca	aaa	ggt	ggt	144
Ala	Ile	Gly	Asp	Val	Val	Asn	Gly	Asn	Asp	Gly	Ala	Ala	Lys	Gly	Gly	
	35						40				45					
gat	gcg	gcg	agt	gtt	aat	ggg	att	gct	aag	ggg	ata	aag	ggg	att	gtt	192
Asp	Ala	Ala	Ser	Val	Asn	Gly	Ile	Ala	Lys	Gly	Ile	Lys	Gly	Ile	Val	
	50					55				60						
gat	gct	gct	gag	aag	gct	gat	gcg	aag	gaa	ggg	aag	ttg	gat	gtg	gct	240
Asp	Ala	Ala	Glu	Lys	Ala	Asp	Ala	Lys	Glu	Gly	Lys	Leu	Asp	Val	Ala	
65					70				75					80		
ggt	gat	gct	ggt	gag	ggt	aac	aag	gat	gct	ggg	aag	ctg	ttt	gtg	aag	288
Gly	Asp	Ala	Gly	Gly	Gly	Asn	Lys	Asp	Ala	Gly	Lys	Leu	Phe	Val	Lys	
			85					90					95			

-continued

aag aat gct ggt gat gag ggt ggt gaa gca aat gat gct ggg aag gct Lys Asn Ala Gly Asp Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala 100 105 110	336
gct gct gcg gtt gct gct gtt agt ggg gag cag ata tta aaa gcg att Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile 115 120 125	384
gtt gat gct gct gag ggt gat gat aag cag ggt aag aag gct gcg gat Val Asp Ala Ala Glu Gly Asp Asp Lys Gln Gly Lys Lys Ala Ala Asp 130 135 140	432
gct aca aat ccg att gag gcg gct att ggg ggt gcg gat gcg ggt gct Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala Asp Ala Gly Ala 145 150 155 160	480
aat gct gag gcg ttt aat aag atg aag aag gat gat cag att gct gct Asn Ala Glu Ala Phe Asn Lys Met Lys Lys Asp Asp Gln Ile Ala Ala 165 170 175	528
gct atg gtt ctg agg gga atg gct aag gat ggg cag ttt gct ttg aag Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys 180 185 190	576
gat gat gct gct gct cat Asp Asp Ala Ala Ala His 195	594

<210> SEQ ID NO 36

<211> LENGTH: 198

<212> TYPE: PRT

<213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 36

Glu Gly Ala Ile Thr Glu Ile Ser Lys Trp Leu Asp Asp Met Ala Lys 1 5 10 15
Ala Ala Ala Val Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala 20 25 30
Ala Ile Gly Asp Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly 35 40 45
Asp Ala Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val 50 55 60
Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala 65 70 75 80
Gly Asp Ala Gly Glu Gly Asn Lys Asp Ala Gly Lys Leu Phe Val Lys 85 90 95
Lys Asn Ala Gly Asp Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala 100 105 110
Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile 115 120 125
Val Asp Ala Ala Glu Gly Asp Asp Lys Gln Gly Lys Lys Ala Ala Asp 130 135 140
Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala Asp Ala Gly Ala 145 150 155 160
Asn Ala Glu Ala Phe Asn Lys Met Lys Lys Asp Asp Gln Ile Ala Ala 165 170 175
Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys 180 185 190
Asp Asp Ala Ala Ala His 195

-continued

```

<210> SEQ ID NO 37
<211> LENGTH: 573
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(573)

<400> SEQUENCE: 37

gaa ggg act gtt aag aat gct gtt gat atg gca aag gcc gct gcg gaa      48
Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Ala Glu
1          5          10          15

gct gca agt gct gca agt gct gct act ggt agt aca acg att gga gat      96
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp
          20          25          30

gtt gtt aag agt ggt gag gca aaa gat ggt gat gcg gcg agt gtt aat     144
Val Val Lys Ser Gly Glu Ala Lys Asp Gly Asp Ala Ala Ser Val Asn
          35          40          45

ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct     192
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala
          50          55          60

gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct gct ggt acg act     240
Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Thr Thr
          65          70          75          80

aac gtg aat gtt ggg aag ttg ttt gtg aag aat aat ggt aat gag ggt     288
Asn Val Asn Val Gly Lys Leu Phe Val Lys Asn Asn Gly Asn Glu Gly
          85          90          95

ggt gat gca agt gat gct ggg aaa gct gct gct gcg gtt gct gct gtt     336
Gly Asp Ala Ser Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val
          100          105          110

agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aaa gat ggt     384
Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly
          115          120          125

gat aag cag ggg gtt act gat gta aag gat gct aca aat ccg att gag     432
Asp Lys Gln Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Glu
          130          135          140

gcg gct att ggg ggt aca aat gat aat gat gct gcg gcg ttt gct act     480
Ala Ala Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr
          145          150          155          160

atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg     528
Met Lys Lys Asp Asp Gln Ile Ala Ala Met Val Leu Arg Gly Met
          165          170          175

gct aag gat ggg cag ttt gct ttg aag gat gat gct gct aag gat         573
Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp Asp Ala Ala Lys Asp
          180          185          190

<210> SEQ ID NO 38
<211> LENGTH: 191
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 38

Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Ala Glu
1          5          10          15

Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp
          20          25          30

Val Val Lys Ser Gly Glu Ala Lys Asp Gly Asp Ala Ala Ser Val Asn
          35          40          45

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala

```

-continued

50	55	60	
Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Thr Thr			
65	70	75	80
Asn Val Asn Val Gly Lys Leu Phe Val Lys Asn Asn Gly Asn Glu Gly			
	85	90	95
Gly Asp Ala Ser Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val			
	100	105	110
Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly			
	115	120	125
Asp Lys Gln Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Glu			
	130	135	140
Ala Ala Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr			
	145	150	155
Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met			
	165	170	175
Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp Asp Ala Ala Lys Asp			
	180	185	190

<210> SEQ ID NO 39
 <211> LENGTH: 189
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(189)

<400> SEQUENCE: 39

ggt gat aaa acg ggg gtt gct gcg gat gct gaa aat ccg att gac gcg	48
Gly Asp Lys Thr Gly Val Ala Ala Asp Ala Glu Asn Pro Ile Asp Ala	
1	5 10 15
gct att ggg ggt gcg gat gct gat gct gcg gcg ttt aat aag gag ggg	96
Ala Ile Gly Gly Ala Asp Ala Asp Ala Ala Phe Asn Lys Glu Gly	
	20 25 30
atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg	144
Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met	
	35 40 45
gct aag gat ggg cag ttt gct ttg acg aat aat gct gct gct cat	189
Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn Ala Ala Ala His	
	50 55 60

<210> SEQ ID NO 40
 <211> LENGTH: 63
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 40

Gly Asp Lys Thr Gly Val Ala Ala Asp Ala Glu Asn Pro Ile Asp Ala	
1	5 10 15
Ala Ile Gly Gly Ala Asp Ala Asp Ala Ala Ala Phe Asn Lys Glu Gly	
	20 25 30
Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met	
	35 40 45
Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn Ala Ala Ala His	
	50 55 60

<210> SEQ ID NO 41
 <211> LENGTH: 576

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(576)

<400> SEQUENCE: 41

gaa ggg act gtt aag aat gct gtt gat atg gca aaa gct gct gcg gtt      48
Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Ala Val
1          5          10          15

gct gca agt gct gct act ggc aat gca gca att ggg gat gtt gtt aag      96
Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys
20          25          30

agt aat ggt gga gca gca gca aaa ggt ggt gat gcg gcg agt gtt aat     144
Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn
35          40          45

ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct     192
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala
50          55          60

gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct gct ggt gaa act     240
Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr
65          70          75          80

aac aag gat gct ggg aag ttg ttt gtg aag aag aat ggt gat gat ggt     288
Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly
85          90          95

ggt gat gca ggt gat gct ggg aag gct gct gct gcg gtt gct gct gtt     336
Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val
100         105         110

agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aaa gat ggt     384
Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly
115        120        125

gat aag acg ggg gtt act gat gta aag gat gct aca aat ccg att gac     432
Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp
130        135        140

gcg gct att ggg ggg agt gcg gat gct aat gct gag gcg ttt gat aag     480
Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys
145        150        155        160

atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg     528
Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met
165        170        175

gct aag gat ggg cag ttt gct ttg aag aat aat gat cat gat aat cat     576
Ala Lys Asp Gly Gln Phe Ala Leu Lys Asn Asn Asp His Asp Asn His
180        185        190

<210> SEQ ID NO 42
<211> LENGTH: 192
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 42

Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Ala Val
1          5          10          15

Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys
20          25          30

Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn
35          40          45

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala
50          55          60

```

-continued

Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr
 65 70 75 80
 Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly
 85 90 95
 Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val
 100 105 110
 Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly
 115 120 125
 Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp
 130 135 140
 Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys
 145 150 155 160
 Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met
 165 170 175
 Ala Lys Asp Gly Gln Phe Ala Leu Lys Asn Asn Asp His Asp Asn His
 180 185 190

<210> SEQ ID NO 43
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(336)

<400> SEQUENCE: 43

aag ggg act gtt aag aat gct gtt gat atg gca aag gcc gct gag gaa	48
Lys Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Glu Glu	
1 5 10 15	
gct gca agt gct gca agt gct gct act ggt aat gca gcg att ggg gat	96
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp	
20 25 30	
gtt gtt aag aat agt ggg gca gca gca aaa ggt ggt gag gcg gcg agt	144
Val Val Lys Asn Ser Gly Ala Ala Lys Gly Gly Glu Ala Ala Ser	
35 40 45	
gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct gga	192
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly	
50 55 60	
aag gct gat gcg aag gaa ggg aag ttg gat gct act ggt gct gag ggt	240
Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Ala Thr Gly Ala Glu Gly	
65 70 75 80	
acg act aac gtg aat gct ggg aag ttg ttt gtg aag agg gcg gct gat	288
Thr Thr Asn Val Asn Ala Gly Lys Leu Phe Val Lys Arg Ala Ala Asp	
85 90 95	
gat ggt ggt gat gca gat gat gct ggg aag gct gct gct gcg gtt gct	336
Asp Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Val Ala	
100 105 110	

<210> SEQ ID NO 44
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 44

Lys Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Glu Glu
 1 5 10 15
 Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp
 20 25 30

-continued

Val Val Lys Asn Ser Gly Ala Ala Ala Lys Gly Gly Glu Ala Ala Ser
 35 40 45

Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
 50 55 60

Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Ala Thr Gly Ala Glu Gly
 65 70 75 80

Thr Thr Asn Val Asn Ala Gly Lys Leu Phe Val Lys Arg Ala Ala Asp
 85 90 95

Asp Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Ala Val Ala
 100 105 110

<210> SEQ ID NO 45
 <211> LENGTH: 522
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(522)

<400> SEQUENCE: 45

gca agt gct gct act ggt aat gca gcg att gga gat gtt gtt aat ggt Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Asn Gly 1 5 10 15	48
gat gtg gca aaa gca aaa ggt ggt gat gcg gcg agt gtt aat ggg att Asp Val Ala Lys Ala Lys Gly Gly Asp Ala Ala Ser Val Asn Gly Ile 20 25 30	96
gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala 35 40 45	144
aag gaa ggg aag ttg aat gct gct ggt gct gag ggt acg act aac gcg Lys Glu Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala 50 55 60	192
gat gct ggg aag ttg ttt gtg aag aat gct ggt aat gtg ggt ggt gaa Asp Ala Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu 65 70 75 80	240
gca ggt gat gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg Ala Gly Asp Ala Gly Lys Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly 85 90 95	288
gag cag ata tta aaa gcg att gtt gat gct gct aag gat ggt ggt gag Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu 100 105 110	336
aag cag ggt aag aag gct gcg gat gct aca aat ccg att gac gcg gct Lys Gln Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Asp Ala Ala 115 120 125	384
att ggg ggt aca aat gat aat gat gct gct gcg gcg ttt gct act atg Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met 130 135 140	432
aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg gct Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala 145 150 155 160	480
aag gat ggg caa ttt gct ttg aag gat gct gct gct gct cat Lys Asp Gly Gln Phe Ala Leu Lys Asp Ala Ala Ala Ala His 165 170	522

<210> SEQ ID NO 46
 <211> LENGTH: 174
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

-continued

<400> SEQUENCE: 46

```

Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Asn Gly
1          5          10          15
Asp Val Ala Lys Ala Lys Gly Gly Asp Ala Ala Ser Val Asn Gly Ile
          20          25          30
Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala
          35          40          45
Lys Glu Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala
          50          55          60
Asp Ala Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu
          65          70          75          80
Ala Gly Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly
          85          90          95
Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu
          100          105          110
Lys Gln Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Asp Ala Ala
          115          120          125
Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met
          130          135          140
Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala
          145          150          155          160
Lys Asp Gly Gln Phe Ala Leu Lys Asp Ala Ala Ala Ala His
          165          170

```

<210> SEQ ID NO 47

<211> LENGTH: 585

<212> TYPE: DNA

<213> ORGANISM: Borrelia garinii

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(585)

<400> SEQUENCE: 47

```

gaa ggg act gtt aag aat gct gtt gat ata ata aag gct gct gcg gaa      48
Glu Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Ala Ala Glu
1          5          10          15
gct gca agt gct gca agt gct gct act ggt agt gca gca att ggg gat      96
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Ala Ala Ile Gly Asp
          20          25          30
gtt gtt aat ggt aat gga gca aca gca aaa ggt ggt gat gcg aag agt      144
Val Val Asn Gly Asn Gly Ala Thr Ala Lys Gly Gly Asp Ala Lys Ser
          35          40          45
gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag      192
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu
          50          55          60
aag gct gat gcg aag gaa ggg aag ttg gat gtg gct ggt gat gct ggt      240
Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Asp Ala Gly
          65          70          75          80
gaa act aac aag gat gct ggg aag ttg ttt gtg aag aac aat ggt aat      288
Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Asn Asn Gly Asn
          85          90          95
gag ggt ggt gat gca gat gat gct ggg aag gct gct gct gcg gtt gct      336
Glu Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Val Ala
          100          105          110
gct gtt agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aag      384

```

-continued

Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys	
115 120 125	
ggt ggt gat aag acg ggt aag aat aat gtg aag gat gct gaa aat ccg	432
Gly Gly Asp Lys Thr Gly Lys Asn Asn Val Lys Asp Ala Glu Asn Pro	
130 135 140	
att gag gcg gct att ggg agt agt gcg gat gct gat gct gcg gcg ttt	480
Ile Glu Ala Ala Ile Gly Ser Ser Ala Asp Ala Asp Ala Ala Phe	
145 150 155 160	
aat aag gag ggg atg aag aag gat gat cag att gct gct gct atg gtt	528
Asn Lys Glu Gly Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val	
165 170 175	
ctg agg gga atg gct aag gat ggg cag ttt gct ttg acg aat gat gct	576
Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asp Ala	
180 185 190	
gct gct cat	585
Ala Ala His	
195	

<210> SEQ ID NO 48
 <211> LENGTH: 195
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 48

Glu Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Ala Ala Glu	
1 5 10 15	
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Ala Ala Ile Gly Asp	
20 25 30	
Val Val Asn Gly Asn Gly Ala Thr Ala Lys Gly Gly Asp Ala Lys Ser	
35 40 45	
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu	
50 55 60	
Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Asp Ala Gly	
65 70 75 80	
Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Asn Asn Gly Asn	
85 90 95	
Glu Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Val Ala	
100 105 110	
Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys	
115 120 125	
Gly Gly Asp Lys Thr Gly Lys Asn Asn Val Lys Asp Ala Glu Asn Pro	
130 135 140	
Ile Glu Ala Ala Ile Gly Ser Ser Ala Asp Ala Asp Ala Ala Phe	
145 150 155 160	
Asn Lys Glu Gly Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val	
165 170 175	
Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asp Ala	
180 185 190	
Ala Ala His	
195	

<210> SEQ ID NO 49
 <211> LENGTH: 591
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:

-continued

<221> NAME/KEY: CDS

<222> LOCATION: (1) .. (591)

<400> SEQUENCE: 49

gaa ggg act gtt aag aat gct gtt ggg agt gca aca aat aag acc gtt 48
 Glu Gly Thr Val Lys Asn Ala Val Gly Ser Ala Thr Asn Lys Thr Val
 1 5 10 15

gtt gct ttg gct aac ttg gtt cga aag acc gtg caa gct ggg ttg aag 96
 Val Ala Leu Ala Asn Leu Val Arg Lys Thr Val Gln Ala Gly Leu Lys
 20 25 30

aag gtt ggg gat gtt gtt aag aat agt gag gca aaa gat ggt gat gcg 144
 Lys Val Gly Asp Val Val Lys Asn Ser Glu Ala Lys Asp Gly Asp Ala
 35 40 45

gcg agt gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct 192
 Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala
 50 55 60

gct gag aag gct gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct 240
 Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala
 65 70 75 80

gct ggt gaa act aac aag gat gct ggg aag ttg ttt gtg aag aag aat 288
 Ala Gly Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn
 85 90 95

aat gag ggt ggt gaa gca aat gat gct ggg aag gct gct gct gcg gtt 336
 Asn Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala Ala Ala Val
 100 105 110

gct gct gtt agt ggg gag cag ata tta aaa gcg att gtt gat gct gct 384
 Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala
 115 120 125

aag gat ggt gat gat aag cag ggt aag aag gct gag gat gct aca aat 432
 Lys Asp Gly Asp Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Thr Asn
 130 135 140

cgc att gac gcg gct att ggg ggt gca ggt gcg ggt gct aat gct gct 480
 Pro Ile Asp Ala Ala Ile Gly Gly Ala Gly Ala Gly Ala Asn Ala Ala
 145 150 155 160

gcg gcg ttt aat aat atg aag aag gat gat cag att gct gct gct atg 528
 Ala Ala Phe Asn Asn Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met
 165 170 175

gtt ctg agg gga atg gct aag gat ggg cag ttt gct ttg acg aat aat 576
 Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn
 180 185 190

gct cat act aat cat 591
 Ala His Thr Asn His
 195

<210> SEQ ID NO 50

<211> LENGTH: 197

<212> TYPE: PRT

<213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 50

Glu Gly Thr Val Lys Asn Ala Val Gly Ser Ala Thr Asn Lys Thr Val
 1 5 10 15

Val Ala Leu Ala Asn Leu Val Arg Lys Thr Val Gln Ala Gly Leu Lys
 20 25 30

Lys Val Gly Asp Val Val Lys Asn Ser Glu Ala Lys Asp Gly Asp Ala
 35 40 45

Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala
 50 55 60

-continued

Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala
65 70 75 80

Ala Gly Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn
85 90 95

Asn Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala Ala Ala Val
100 105 110

Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala
115 120 125

Lys Asp Gly Asp Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Thr Asn
130 135 140

Pro Ile Asp Ala Ala Ile Gly Gly Ala Gly Ala Gly Ala Asn Ala Ala
145 150 155 160

Ala Ala Phe Asn Asn Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met
165 170 175

Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn
180 185 190

Ala His Thr Asn His
195

<210> SEQ ID NO 51
 <211> LENGTH: 594
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(594)

<400> SEQUENCE: 51

aag ggg act gtt aag aat gct gtt gat atg aca aaa gct gct gcg gtt Lys Gly Thr Val Lys Asn Ala Val Asp Met Thr Lys Ala Ala Ala Val 1 5 10 15	48
gct gca agt gct gca agt gct gct act ggt aat gca gca att ggg gat Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp 20 25 30	96
gtt gtt aat ggt aat gat gga gca gca aaa ggt ggt gat gcg gcg agt Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly Asp Ala Ala Ser 35 40 45	144
gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu 50 55 60	192
aag gct gat gcg aag gaa ggg aag ttg aat gtg gct ggt gct gct ggt Lys Ala Asp Ala Lys Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly 65 70 75 80	240
gct gag ggt aac gag gct gct ggg aag ctg ttt gtg aag aag aat gct Ala Glu Gly Asn Glu Ala Ala Gly Lys Leu Phe Val Lys Lys Asn Ala 85 90 95	288
ggt gat cat ggt ggt gaa gca ggt gat gct ggg agg gct gct gct gcg Gly Asp His Gly Gly Glu Ala Gly Asp Ala Gly Arg Ala Ala Ala Ala 100 105 110	336
gtt gct gct gtt agt ggg gag cag ata tta aaa gcg att gtt gat gct Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala 115 120 125	384
gct aag gat ggt ggt gat aag cag ggt aag aag gct gag gat gct gaa Ala Lys Asp Gly Gly Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Glu 130 135 140	432
aat ccg att gac gcg gct att ggg agt acg ggt gcg gat gat aat gct	480

-continued

Asn Pro Ile Asp Ala Ala Ile Gly Ser Thr Gly Ala Asp Asp Asn Ala	
145 150 155 160	
gct gag gcg ttt gct act atg aag aag gat gat cag att gct gct gct	528
Ala Glu Ala Phe Ala Thr Met Lys Lys Asp Asp Gln Ile Ala Ala Ala	
165 170 175	
atg gtt ctg agg gga atg gct aag gat ggg cag ttt gct ttg aag gat	576
Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp	
180 185 190	
gct gct cat gat aat cat	594
Ala Ala His Asp Asn His	
195	

<210> SEQ ID NO 52
 <211> LENGTH: 198
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 52

Lys Gly Thr Val Lys Asn Ala Val Asp Met Thr Lys Ala Ala Val	
1 5 10 15	
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp	
20 25 30	
Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly Asp Ala Ala Ser	
35 40 45	
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu	
50 55 60	
Lys Ala Asp Ala Lys Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly	
65 70 75 80	
Ala Glu Gly Asn Glu Ala Ala Gly Lys Leu Phe Val Lys Lys Asn Ala	
85 90 95	
Gly Asp His Gly Gly Glu Ala Gly Asp Ala Gly Arg Ala Ala Ala Ala	
100 105 110	
Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala	
115 120 125	
Ala Lys Asp Gly Gly Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Glu	
130 135 140	
Asn Pro Ile Asp Ala Ala Ile Gly Ser Thr Gly Ala Asp Asp Asn Ala	
145 150 155 160	
Ala Glu Ala Phe Ala Thr Met Lys Lys Asp Asp Gln Ile Ala Ala Ala	
165 170 175	
Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp	
180 185 190	
Ala Ala His Asp Asn His	
195	

<210> SEQ ID NO 53
 <211> LENGTH: 261
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(261)

<400> SEQUENCE: 53

aag ggg act gtt aag aat gct gtt gat ata ata aag gct act gcg gtt	48
Lys Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Thr Ala Val	
1 5 10 15	

-continued

```

gct gca agt gct gct act ggt agt aca acg att ggg gat gtt gtt aag      96
Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp Val Val Lys
                20                25                30

aat ggt gag gca aaa ggt ggt gag gcg aag agt gtt aat ggg att gct      144
Asn Gly Glu Ala Lys Gly Gly Glu Ala Lys Ser Val Asn Gly Ile Ala
                35                40                45

aag ggg ata aag ggg att gtt gat gct gct gga aag gct gat gcg aag      192
Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala Asp Ala Lys
                50                55                60

gaa ggg aag ttg aat gtg gct ggt gct gct ggt gag ggt aac gag gct      240
Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly Glu Gly Asn Glu Ala
        65                70                75                80

gct ggg aag ctg ttt gtg taa
Ala Gly Lys Leu Phe Val
                85

```

```

<210> SEQ ID NO 54
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

```

```

<400> SEQUENCE: 54

```

```

Lys Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Thr Ala Val
1          5          10          15

Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp Val Val Lys
20          25          30

Asn Gly Glu Ala Lys Gly Gly Glu Ala Lys Ser Val Asn Gly Ile Ala
35          40          45

Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala Asp Ala Lys
50          55          60

Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly Glu Gly Asn Glu Ala
65          70          75          80

Ala Gly Lys Leu Phe Val
85

```

```

<210> SEQ ID NO 55
<211> LENGTH: 213
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(213)

```

```

<400> SEQUENCE: 55

```

```

gta aat tac tat agg att aga act agt gta cga tat gag tcc ttt ggt      48
Val Asn Tyr Tyr Arg Ile Arg Thr Ser Val Arg Tyr Glu Ser Phe Gly
1          5          10          15

tat ttt gca gct gct aat gaa ttt gaa ata agt gaa gtt aaa att gcg      96
Tyr Phe Ala Ala Ala Asn Glu Phe Glu Ile Ser Glu Val Lys Ile Ala
20          25          30

gat gtt aat gga aca cat ttt att gct aca aaa gag aaa gaa ata tta      144
Asp Val Asn Gly Thr His Phe Ile Ala Thr Lys Glu Lys Glu Ile Leu
35          40          45

tat gat tca ctt gat tta agg gct cgt gga aaa ata ttt gaa ata act      192
Tyr Asp Ser Leu Asp Leu Arg Ala Arg Gly Lys Ile Phe Glu Ile Thr
50          55          60

tca aag cga atg ttt aag ctt
Ser Lys Arg Met Phe Lys Leu
213

```

-continued

65

70

<210> SEQ ID NO 56
 <211> LENGTH: 71
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 56

Val	Asn	Tyr	Tyr	Arg	Ile	Arg	Thr	Ser	Val	Arg	Tyr	Glu	Ser	Phe	Gly
1				5					10					15	
Tyr	Phe	Ala	Ala	Ala	Asn	Glu	Phe	Glu	Ile	Ser	Glu	Val	Lys	Ile	Ala
		20						25					30		
Asp	Val	Asn	Gly	Thr	His	Phe	Ile	Ala	Thr	Lys	Glu	Lys	Glu	Ile	Leu
		35					40					45			
Tyr	Asp	Ser	Leu	Asp	Leu	Arg	Ala	Arg	Gly	Lys	Ile	Phe	Glu	Ile	Thr
	50					55					60				
Ser	Lys	Arg	Met	Phe	Lys	Leu									
65					70										

<210> SEQ ID NO 57
 <211> LENGTH: 8762
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 57

gagagtgtctg	ttgatgggggt	tagcaagtgg	ttagaagaga	tgataaaagc	tgctaaggag	60
gctgtctacaa	aggggtgttac	tggtgtgtgt	agcgaaaaga	ttggggatgt	tggtgtctgct	120
aataatcagg	gtgctgtagc	tgataaggac	agtgttaagg	ggattgcgaa	ggggataaag	180
gggattgttg	atgctgtctgg	gaaggctttt	ggtaaggatg	gtaatgcgct	gacaggtgta	240
aaagaagtgtg	ctgatgaggc	tggggctaac	gaggatgcgg	ggaagttgtt	tgctggtaat	300
gctgtgtaatg	ctgctgtctgc	tgacattgcg	aaggcggtcg	gtgctgttac	tgcggttagt	360
ggggagcaga	tactgaaagc	tattgttgat	ggtgctgggtg	gtgcggctca	agatggtaaa	420
aaggctgcgg	aggctaagaa	tccgattgca	gctgcgattg	gggctgatgc	tgctggtgcg	480
gattttgtgtg	atgatatgaa	gaagatgat	aagattgctg	cggtatttgt	tttgaggggg	540
gtggctaaga	gtggaaagt	tgctgttgct	aatgctgcta	agaaggagag	tgtgaagagt	600
gctgtggaga	gtgctgttga	tgaggttagc	aagtggttag	aagagatgat	aaaagctgct	660
gggtggggctg	ctaaggggtg	tactggtggt	aataacgaaa	agattgggga	ttctgataat	720
aataaggggtg	ctgtagctga	taaggacagt	gttaagggga	ttgcgaaggg	gataaagggg	780
attgttgatg	ctgctgggaa	ggcttttgg	aaggatggtg	atgcgctgaa	ggatgttgca	840
aaagttgctg	atgatgcggc	tggggctaac	gcgaatgcag	ggaagttgtt	tgctggtaat	900
gctgtctgggtg	gtgccgctga	tgctgatgat	gctaacattg	cgaaggcggc	tggtgtctgtt	960
agtgcggtta	gtggggagca	gatactgaaa	gctattgttg	atgctgctgg	tgctgtctgct	1020
aatcaggatg	gtaagaaggc	tcgggatgct	aagaatccga	ttgcagctgc	gattgggact	1080
aatgatgatg	gggagggtt	taaggatgga	atgaagaaga	gtgataatat	tgctgcagct	1140
attgttttga	gggggggtgg	taagggtgga	aagtttctg	ttgctaattgc	tgctaattgat	1200
agtaaggcga	gtgtgaagag	tgctgtggag	agtgtgttg	atgaggttag	caagtggtta	1260
gaagagatga	taacagctgc	tggtgaggct	gctacaaagg	gtggtgatgc	tggtggtggt	1320

-continued

gctgataaga	ttggggatgt	tgggtgctgct	aataatgggtg	ctgtagctga	tgcgagcagt	1380
gttaaggaga	ttgcgaaggg	gataaagggg	attgttgatg	ctgctgggaa	ggcttttggc	1440
aaggatggta	atgcgctgaa	ggatgttgca	gaagttgctg	atgataagaa	ggaggcgggg	1500
aagttgtttg	ctggtaatgc	tgggtggtgct	gttgctgatg	ctgctgcgat	tggaagggcg	1560
gctggtgctg	ttactgcggg	tagtggggag	cagatactga	aagctattgt	tgatgctgct	1620
ggtggtgctg	atcaggcggg	taagaaggct	gatgcggcta	agaatccgat	tgcagctgctg	1680
attggggctg	atgctgctgg	tgctggtgctg	gattttggta	atgatatgaa	gaagagaaat	1740
gataagattg	ttgcggctat	tgttttgagg	ggggtggcta	aggatggaaa	gtttgctgct	1800
gctgctaattg	atgataatag	taaggcgagt	gtgaagagtg	ctgtgggagag	tgctgttgat	1860
gaggttagca	agtgggttaga	agagatgata	acagctgctg	atggggctgc	taaagggtggt	1920
actggtggta	atagcgaaaa	gattggggat	gctggtgata	ataataatgg	tgctgtagct	1980
gatgagaaca	gtgttaagga	gattgcaaa	gggataaagg	ggattgttgc	ggctgctggg	2040
aaggcttttg	gcaaggatgg	caaggatggt	gatgcgctga	aggatgttga	aacagttgct	2100
gctgagaatg	aggctaacaa	ggatgcgggg	aagttgtttg	ctggtgctaa	tgtaaatgct	2160
ggtgctgctg	ttggtgacat	tgcgaaggcg	gctgctgctg	ttactgcggg	tagtggggag	2220
cagatactaa	aagctattgt	tgatgctgct	ggtgatgcgg	atcaggcggg	taagaaggct	2280
gctgaggcta	agaatccgat	tgcagctgctg	attggggcta	atgctgctga	taatgcggcg	2340
gcgtttggta	aggatgagat	gaagaagagt	gataagattg	ctgcagctat	tgttttgagg	2400
ggggtggcta	aggatggaaa	gtttgctggt	gctaagtcta	atgatgataa	gaaggcgagt	2460
gtgaagagtg	ctgtgggagag	tgctgtggat	gagggttagca	agtgggttaga	agagatgata	2520
acagctgcta	aggaggctgc	tacaaagggt	ggtactggtg	gtaataacga	aaagattgga	2580
gattctgatg	ctaataatgg	tgcgaaggct	gatgcgagca	gtgttaatgg	gattgcgaat	2640
gggataaagg	ggattgttga	tgctgctggg	aaggcttttg	gcaaggaggg	tagtgcgctg	2700
aaggatgtta	aaacagttgc	tgctgagaat	gaggctaaca	aggatgcggg	gaagttgttt	2760
gctggttaaga	atggtaatgc	tgatgctgct	gatgctgctg	acattgcgaa	ggcggtggt	2820
gctgttagtg	cggtagtggtg	ggagcagata	ctgaaagcta	ttgttgatgg	tgctggtgat	2880
gcagctaatac	aggcgggttaa	aaaggctgct	gaggctaaga	atccgattgc	ggctgcgatt	2940
gggactaatg	aagctggggc	ggagtttggt	gatgatatga	agaagagaaa	tgataagatt	3000
gctgcggcta	ttgttttgag	gggggtggct	aaggatggaa	agtttgctgt	tgctaattgct	3060
gctgctgata	atagtaaggc	gagtggtgaag	agtgctgttg	atgaggttag	caagtggtta	3120
gaagagatga	taaaggctgc	tggtgaggct	gctacaaagg	gtggtgatgc	tggtggtggt	3180
gctgataaga	ttggggatgc	tggtgataag	ggtgctgtag	ctgatgcgag	cagtgttaag	3240
gagattgcga	atgggataaaa	gggattgtt	gatgctgctg	ggaaggcttt	tggaaggag	3300
ggtagtgctg	tgaaggatgt	taaaacagtt	gctgctgaga	atgaggctaa	caaggatgcg	3360
gggaagttgt	ttgctggtaa	tgctggtaat	ggtgctgctg	atgacattgc	gaaggcggct	3420
gctgctgtta	ctgcgggttag	tggggagcag	atactgaaag	ctattgttga	tgctgctggt	3480
gataaggcta	atcaggatgg	taaaaaggct	gcggatgcta	agaatccgat	tgcggctgctg	3540
attggggctg	ctgatgctgg	tgctgcggcg	gcgtttaatg	agaatgatat	gaagaagagt	3600

-continued

gataagattg	ctgcagctat	tgttttgagg	ggggtggcta	aggatggaaa	gtttgctgct	3660
gctgatgctg	atgctaataa	tagtaaggcg	agcgtgaaga	gtgctgttgg	tgaggttagc	3720
aagtgggttag	aagagatgat	aaaagctgct	ggtgaggctg	caaaagttgg	tggtactggt	3780
ggtagcgaag	agattgggga	tgctgataat	aataagggtg	ctgtagctga	tgcgagcagt	3840
gttaatggga	ttgcgaatgg	gataaagggg	attgttgatg	ctgctgggaa	ggcttttggt	3900
aaggatgggtg	cgctggcagg	tgttgagcgt	gctgctgaga	atgatgataa	gaaggatgctg	3960
gggaagtgtg	ttgctggtaa	gaatggtggt	gctggtgctg	ctgatgcgat	tggaaggcgg	4020
gctgctgctg	ttactgcggt	tagtggggag	cagatactga	aagctattgt	tgatgctgct	4080
ggtgctgcag	ctaactcaggc	gggtaaaaag	gctgcggatg	ctaagaatcc	gattgcggct	4140
gcgattggga	ctgctgatga	tggggcggag	tttaaggatg	atatgaagaa	gagtgataat	4200
attgctgcgg	ctattgtttt	gaggggggtg	gctaaggatg	gaaagtgtgc	tgttgctaata	4260
gctgatgata	ataaggcgag	tgtgaagagt	gctgtggaga	gtgctgttga	tgaggttagc	4320
aagtgggttag	aagagatgat	aacagctgct	ggtgaggctg	caaaagttgg	tgctggtggt	4380
ggtgctgata	agattgggga	tgctgctaata	aatcagggtg	cgaaggctga	tgagagcagt	4440
gttaatggaa	ttgcaaaagg	gataaagggg	attgttgatg	ctgctgggaa	ggcttttggt	4500
aaggagggtg	gtgcgctgaa	ggatgttgca	aaagtgtgctg	atgatgataa	caaggatgctg	4560
gggaagtgtg	ttgctggtaa	tgctggtggt	ggtgctggtg	ctgatattgc	gaaggcggct	4620
gctgctgtta	ctgcggttag	tggggagcag	atactgaaag	ctattgttga	tgctgctggt	4680
gctgcggatc	aggcgggtgc	agctgctggt	gcggctaaga	atccgattgc	ggctgcgatt	4740
ggggctgatg	ctggtgctgc	ggaggagttt	aaggatgaga	tgaagaagag	tgataagatt	4800
gctgcggcta	ttgttttgag	gggggtggct	aagggtggaa	agtttgctgt	tgctgctaata	4860
gatgctgcaa	atgtgaagag	tgctgtggag	agtgtgttgg	gtgaggtttag	cgcatgggtta	4920
gaagagatga	taacagctgc	tagtgaggct	gctacaaagg	gtggtactgg	tggtactggt	4980
ggtgatagtg	aaaagattgg	ggattctgat	gctaataatg	gtgctgtagc	tgatgcgagc	5040
agtgttaagg	agattgcgaa	ggggataaag	gggattgttg	atgctgctgg	gaaggctttt	5100
ggtaaggatg	gtaatgcgct	gaaggatgtt	gcagaagttg	ctgatgatga	ggctaacgcg	5160
gatgcgggga	agttgtttgc	tggtaatgct	ggtaatgctg	ctgctgctga	cgttgcaag	5220
gcggctggtg	ctgttactgc	ggttagtggg	gagcagatac	tgaaagctat	tgttgatgct	5280
gctggtgctg	cggatcaggc	gggtgcaaag	gctgatgcgg	ctaagaatcc	gattgcagct	5340
gcgattggga	ctaataaagc	tggggcggcg	tttaaggatg	gaatgaagaa	gagaaatgat	5400
aatattgctg	cggctattgt	tttgaggggg	gtggctaaga	gtggaaagtt	tgctgttgct	5460
gctgctgatg	ctggtaaagg	gagagtgtga	agagtgtgt	ggagagtgt	gttgatgagg	5520
ttagcaagtg	gttagaagag	atgataacag	ctgctagtga	ggctgcaaaa	gttggtgctg	5580
gtggtgatga	taagattggg	gattctgcta	ataatggtgc	tgtagctgat	gcgggcagtg	5640
ttaagggaat	tgcaaggggg	ataaagggga	ttgttgatgc	tgctgggaag	gcttttggtta	5700
aggagggtga	tgcgctgaag	gatgttgcaa	aagttgctga	tgagaatggg	gataacaagg	5760
atgcggggaa	gttgtttgct	ggtgagaatg	gtaatgctgg	tggtgctgct	gatgctgaca	5820
ttgcgaaggc	ggctgctgct	gttactgcgg	ttagtgggga	gcagatactg	aaagctattg	5880

-continued

ttgaggctgc	tgggtgctggt	gatgcagcta	atcaggcggg	taagaaggct	gatgaggcta	5940
agaatccgat	tgcggctgcg	attgggactg	atgatgctgg	ggcggcgttt	ggtcaggatg	6000
atatgaagaa	gagaaatgat	aatattgctg	cggctattgt	tttgaggggg	gtggctaagg	6060
gtggaaagt	tgtgttgct	aatgctgcta	atgatagtaa	ggcgagtgtg	aagagtgtctg	6120
tggagagtgc	tgttgatgag	gttagcaagt	ggtagaaga	gataataaca	gctactggga	6180
aggcttttgg	taaggatggt	aatgcgctgg	caggtgttgc	aaaagttgct	gatgatgagg	6240
ctaacgcgga	tgcggggaag	ttgtttgctg	gtgagaatgg	taatgctggt	gctgctgcga	6300
ttgggaaggc	ggctgctgct	gttactgcgg	ttagtgggga	gcagatactg	aaagctattg	6360
ttgatgctgc	tgggtggtgcg	gctcagggtg	gtgctgggtc	tgggtgcggc	acgaatccga	6420
ttgcagctgc	gattggggct	gctgggtgat	gtgcggattt	tggttaaggat	gagatgaaga	6480
agagaaatga	taagattgct	gcggctattg	ttttgagggg	ggtggctaag	gatggaaagt	6540
ttgctgctgc	tgctaattgat	agtaaggcga	gtgtgaagag	tgctgtggag	agtgtgtt	6600
atgaggttag	caagtgttga	gaagagatga	taacagctgc	tgatgctgct	gctgctaaag	6660
ttggcgatgc	tgggtggtggt	gctgataaga	ttggggatgt	tgggtgctgct	aataagggtg	6720
cgaaggctga	tgcgagcagt	gttaaggaga	ttgcgaaggg	gataaagggg	attgttgatg	6780
ctgctgggaa	ggcttttgg	ggtgatgcgc	tgaaggatgt	taaagctgct	ggtgatgata	6840
acaaggaggc	agggaaagttg	tttctggtg	ctaattggtaa	tgctggtgct	aatgctgctg	6900
ctgctgatga	cattgcgaag	gcggctggtg	ctgttagtgc	ggttagtggg	gagcagatac	6960
tgaagctat	tgttgaggcg	gctggtgctg	cggatcaggc	gggtgtaaag	gctgaggagg	7020
ctaagaatcc	gattgcagct	gcgattggga	ctgatgatgc	tgggtgcggcg	gagtttgggtg	7080
agaatgatat	gaagaagaat	gataatattg	ctgcggctat	tgttttgagg	ggggtggcta	7140
agagtggaaa	gtttgctgct	aatgctaattg	atgctggtaa	gaaggagagt	gtgaagagt	7200
ctgtggatga	ggctagcaag	tggttagaag	agatgataac	agctgctggt	gaggctgcta	7260
caaagggtgg	tactggtgaa	gctagcgaaa	agattgggga	tggtggtgat	aataatcatg	7320
gtgctgtagc	tgatgcggac	agtgttaagg	ggattgcgaa	ggggataaag	gggattgtt	7380
atgctgctgg	gaaggctttt	ggttaaggatg	gtgcgctgaa	ggatgttgca	gctgctgctg	7440
gtgatgaggc	taacaaggat	gcgggggaag	tgtttctggtg	tcaggatggt	ggtggtgctg	7500
atggtgacat	tgcgaaggcg	gctgctgctg	ttactgcggt	tagtggggag	cagatactga	7560
aagctattgt	tgaggctgct	ggtgataagg	ctaatacagg	gggtgtaaag	gctgctggtg	7620
cggctacgaa	tccgattgca	gctgcgattg	ggactgatga	tgataatgcg	gcggcgttt	7680
ataaggatga	gatgaagaag	agtaatgata	agattgctgc	ggctattgtt	ttgagggggg	7740
tggctaagga	tggaaagttt	gctgctaattg	ctaataataa	tagtaaggcg	agtgtgaaga	7800
gtgctgtgga	tgaggttagc	aagtgttag	aagagatgat	aacagctgct	agtgtgctg	7860
ctacaaagg	tggtactggt	gaagctagcg	aaaagattgg	ggattctgat	gctaataagg	7920
gtgctggtgc	tggggcgcg	tttggtgaga	atgatatgaa	gaagagaaat	gataatattg	7980
ctgcagctat	tgttttgagg	gggtgggcta	aggatggaaa	gtttgctgtt	aaggaggatt	8040
attgaactca	gctttatagg	ggaacagcaa	ttcgctagaa	aatgattaaa	aagcttaact	8100
tcgactggtt	cttgccctaa	ttttattcct	ttgttattat	ttatcaatta	aattcacttc	8160

-continued

```

ggtttgcttt taaattaatt ctggtatact atgtatacta gatacacaaa ttaaggagaa 8220
gtgaaatgga aaaaatagaa aaatttaaaa acaaatgtca acataaacta caacataaac 8280
taatcgatt agtatcaaca ctttgctata taaacaataa aaataaaaaa tattcacaaa 8340
gcaacatcct ttattatctt aatgaaaatt taaaagaaa tgggcaaacc cctattaaaa 8400
taaaaacatt acaaaactat ctttataaac tggaaaaaga atttgaagta acaactaatt 8460
attataaaca cttggggggtt aattgtggaa ccgaaattta ctataaactt aaatatcaaa 8520
aacaaaaatg ctatcataaa ataaaccaat attttaaaaa gaaaaaagaa attaaattta 8580
acttaagagt aagtgcattt ttttaaaaaa aacactcaaa aaaaggaggt gtagaattaa 8640
aggaatgtaa taataataat aataataaag agaaagaac atcccaaaaa attgaaattt 8700
tacaacaaaa agtctatgcc aaaaaatgta aatttttgac aaactactat actaaaattt 8760
ta 8762

```

```

<210> SEQ ID NO 58
<211> LENGTH: 606
<212> TYPE: DNA
<213> ORGANISM: Borrelia afzelii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(606)

```

```

<400> SEQUENCE: 58

```

```

gag agt gct gtt gat ggg gtt agc aag tgg tta gaa gag atg ata aaa 48
Glu Ser Ala Val Asp Gly Val Ser Lys Trp Leu Glu Glu Met Ile Lys
1 5 10 15

gct gct aag gag gct gct aca aag ggt ggt act ggt ggt ggt agc gaa 96
Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Gly Ser Glu
20 25 30

aag att ggg gat gtt ggt gct gct aat aat cag ggt gct gta gct gat 144
Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gln Gly Ala Val Ala Asp
35 40 45

aag gac agt gtt aag ggg att gcg aag ggg ata aag ggg att gtt gat 192
Lys Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp
50 55 60

gct gct ggg aag gct ttt ggt aag gat ggt aat gcg ctg aca ggt gta 240
Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Thr Gly Val
65 70 75 80

aaa gaa gtt gct gat gag gct ggg gct aac gag gat gcg ggg aag ttg 288
Lys Glu Val Ala Asp Glu Ala Gly Ala Asn Glu Asp Ala Gly Lys Leu
85 90 95

ttt gct ggt aat gct ggt aat gct gct gct gct gac att gcg aag gcg 336
Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Asp Ile Ala Lys Ala
100 105 110

gct ggt gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att 384
Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile
115 120 125

gtt gat ggt gct ggt ggt gcg gct caa gat ggt aaa aag gct gcg gag 432
Val Asp Gly Ala Gly Gly Ala Ala Gln Asp Gly Lys Lys Ala Ala Glu
130 135 140

gct aag aat ccg att gca gct gcg att ggg gct gat gct gct ggt gcg 480
Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala
145 150 155 160

gat ttt ggt gat gat atg aag aag agt gat aag att gct gcg gct att 528
Asp Phe Gly Asp Asp Met Lys Lys Ser Asp Lys Ile Ala Ala Ala Ile

```

-continued

165	170	175	
gtt ttg agg ggg gtg gct aag agt gga aag ttt gct gtt gct aat gct			576
Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Asn Ala			
180	185	190	
gct aag aag gag agt gtg aag agt gct gtg			606
Ala Lys Lys Glu Ser Val Lys Ser Ala Val			
195	200		

<210> SEQ ID NO 59
 <211> LENGTH: 202
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 59

Glu Ser Ala Val Asp Gly Val Ser Lys Trp Leu Glu Glu Met Ile Lys	
1 5 10 15	
Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Gly Ser Glu	
20 25 30	
Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gln Gly Ala Val Ala Asp	
35 40 45	
Lys Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp	
50 55 60	
Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Thr Gly Val	
65 70 75 80	
Lys Glu Val Ala Asp Glu Ala Gly Ala Asn Glu Asp Ala Gly Lys Leu	
85 90 95	
Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Asp Ile Ala Lys Ala	
100 105 110	
Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile	
115 120 125	
Val Asp Gly Ala Gly Gly Ala Ala Gln Asp Gly Lys Lys Ala Ala Glu	
130 135 140	
Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala	
145 150 155 160	
Asp Phe Gly Asp Asp Met Lys Lys Ser Asp Lys Ile Ala Ala Ala Ile	
165 170 175	
Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Asn Ala	
180 185 190	
Ala Lys Lys Glu Ser Val Lys Ser Ala Val	
195 200	

<210> SEQ ID NO 60
 <211> LENGTH: 621
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(621)

<400> SEQUENCE: 60

gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aaa	48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys	
1 5 10 15	
gct gct ggt ggg gct gct aag ggt ggt act ggt ggt aat aac gaa aag	96
Ala Ala Gly Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Asn Glu Lys	
20 25 30	

-continued

att ggg gat tct gat aat aat aag ggt gct gta gct gat aag gac agt	144
Ile Gly Asp Ser Asp Asn Asn Lys Gly Ala Val Ala Asp Lys Asp Ser	
35 40 45	
ggt aag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg	192
Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly	
50 55 60	
aag gct ttt ggt aag gat ggt aat gcg ctg aag gat gtt gca aaa gtt	240
Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val Ala Lys Val	
65 70 75 80	
gct gat gat gcg gct ggg gct aac gcg aat gca ggg aag ttg ttt gct	288
Ala Asp Asp Ala Ala Gly Ala Asn Ala Asn Ala Gly Lys Leu Phe Ala	
85 90 95	
ggt aat gct gct ggt ggt gcc gct gat gct gat gat gct aac att gcg	336
Gly Asn Ala Ala Gly Gly Ala Ala Asp Ala Asp Asp Ala Asn Ile Ala	
100 105 110	
aag gcg gct ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa	384
Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys	
115 120 125	
gct att gtt gat gct gct ggt gct gct gct aat cag gat ggt aag aag	432
Ala Ile Val Asp Ala Ala Gly Ala Ala Ala Asn Gln Asp Gly Lys Lys	
130 135 140	
gct gcg gat gct aag aat ccg att gca gct gcg att ggg act aat gat	480
Ala Ala Asp Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Asp	
145 150 155 160	
gat ggg gcg gag ttt aag gat gga atg aag aag agt gat aat att gct	528
Asp Gly Ala Glu Phe Lys Asp Gly Met Lys Lys Ser Asp Asn Ile Ala	
165 170 175	
gca gct att gtt ttg agg ggg gtg gct aag ggt gga aag ttt gct gtt	576
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val	
180 185 190	
gct aat gct gct aat gat agt aag gcg agt gtg aag agt gct gtg	621
Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val	
195 200 205	
<210> SEQ ID NO 61	
<211> LENGTH: 207	
<212> TYPE: PRT	
<213> ORGANISM: Borrelia afzelii	
<400> SEQUENCE: 61	
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys	
1 5 10 15	
Ala Ala Gly Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Asn Glu Lys	
20 25 30	
Ile Gly Asp Ser Asp Asn Asn Lys Gly Ala Val Ala Asp Lys Asp Ser	
35 40 45	
Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly	
50 55 60	
Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val Ala Lys Val	
65 70 75 80	
Ala Asp Asp Ala Ala Gly Ala Asn Ala Asn Ala Gly Lys Leu Phe Ala	
85 90 95	
Gly Asn Ala Ala Gly Gly Ala Ala Asp Ala Asp Asp Ala Asn Ile Ala	
100 105 110	
Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys	
115 120 125	

-continued

Ala Ile Val Asp Ala Ala Gly Ala Ala Ala Asn Gln Asp Gly Lys Lys	
130 135 140	
Ala Ala Asp Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Asp	
145 150 155 160	
Asp Gly Ala Glu Phe Lys Asp Gly Met Lys Lys Ser Asp Asn Ile Ala	
165 170 175	
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val	
180 185 190	
Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val	
195 200 205	
 <210> SEQ ID NO 62	
<211> LENGTH: 618	
<212> TYPE: DNA	
<213> ORGANISM: Borrelia afzelii	
<220> FEATURE:	
<221> NAME/KEY: CDS	
<222> LOCATION: (1)..(618)	
 <400> SEQUENCE: 62	
gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca	48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr	
1 5 10 15	
gct gct ggt gag gct gct aca aag ggt ggt gat gct ggt ggt ggt gct	96
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala	
20 25 30	
gat aag att ggg gat gtt ggt gct gct aat aat ggt gct gta gct gat	144
Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gly Ala Val Ala Asp	
35 40 45	
gcg agc agt gtt aag gag att gcg aag ggg ata aag ggg att gtt gat	192
Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Asp	
50 55 60	
gct gct ggg aag gct ttt ggc aag gat ggt aat gcg ctg aag gat gtt	240
Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val	
65 70 75 80	
gca gaa gtt gct gat gat aag aag gag gcg ggg aag ttg ttt gct ggt	288
Ala Glu Val Ala Asp Asp Lys Lys Glu Ala Gly Lys Leu Phe Ala Gly	
85 90 95	
aat gct ggt ggt gct gtt gct gat gct gct gcg att ggg aag gcg gct	336
Asn Ala Gly Gly Ala Val Ala Asp Ala Ala Ile Gly Lys Ala Ala	
100 105 110	
ggt gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt	384
Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val	
115 120 125	
gat gct gct ggt ggt gcg gat cag gcg ggt aag aag gct gat gcg gct	432
Asp Ala Ala Gly Gly Ala Asp Gln Ala Gly Lys Lys Ala Asp Ala Ala	
130 135 140	
aag aat ccg att gca gct gcg att ggg gct gat gct gct ggt gct ggt	480
Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala Gly	
145 150 155 160	
gcg gat ttt ggt aat gat atg aag aag aga aat gat aag att gtt gcg	528
Ala Asp Phe Gly Asn Asp Met Lys Lys Arg Asn Asp Lys Ile Val Ala	
165 170 175	
gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct gct	576
Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala	
180 185 190	
gct aat gat gat aat agt aag gcg agt gtg aag agt gct gtg	618
Ala Asn Asp Asp Asn Ser Lys Ala Ser Val Lys Ser Ala Val	

-continued

195	200	205	
<210> SEQ ID NO 63			
<211> LENGTH: 206			
<212> TYPE: PRT			
<213> ORGANISM: Borrelia afzelii			
<400> SEQUENCE: 63			
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr			
1 5 10 15			
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala			
20 25 30			
Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gly Ala Val Ala Asp			
35 40 45			
Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Asp			
50 55 60			
Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val			
65 70 75 80			
Ala Glu Val Ala Asp Asp Lys Lys Glu Ala Gly Lys Leu Phe Ala Gly			
85 90 95			
Asn Ala Gly Gly Ala Val Ala Asp Ala Ala Ala Ile Gly Lys Ala Ala			
100 105 110			
Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val			
115 120 125			
Asp Ala Ala Gly Gly Ala Asp Gln Ala Gly Lys Lys Ala Asp Ala Ala			
130 135 140			
Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala Gly			
145 150 155 160			
Ala Asp Phe Gly Asn Asp Met Lys Lys Arg Asn Asp Lys Ile Val Ala			
165 170 175			
Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala			
180 185 190			
Ala Asn Asp Asp Asn Ser Lys Ala Ser Val Lys Ser Ala Val			
195 200 205			
<210> SEQ ID NO 64			
<211> LENGTH: 630			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia afzelii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(630)			
<400> SEQUENCE: 64			
gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca			48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr			
1 5 10 15			
gct gct gat ggg gct gct aaa ggt ggt act ggt ggt aat agc gaa aag			96
Ala Ala Asp Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Ser Glu Lys			
20 25 30			
att ggg gat gct ggt gat aat aat aat ggt gct gta gct gat gag aac			144
Ile Gly Asp Ala Gly Asp Asn Asn Asn Gly Ala Val Ala Asp Glu Asn			
35 40 45			
agt gtt aag gag att gca aag ggg ata aag ggg att gtt gcg gct gct			192
Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala			
50 55 60			

-continued

ggg aag gct ttt ggc aag gat ggc aag gat ggt gat gcg ctg aag gat	240
Gly Lys Ala Phe Gly Lys Asp Gly Lys Asp Gly Asp Ala Leu Lys Asp	
65 70 75 80	
ggt gaa aca gtt gct gct gag aat gag gct aac aag gat gcg ggg aag	288
Val Glu Thr Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys	
85 90 95	
ttg ttt gct ggt gct aat ggt aat gct ggt gct gct gtt ggt gac att	336
Leu Phe Ala Gly Ala Asn Gly Asn Ala Gly Ala Ala Val Gly Asp Ile	
100 105 110	
gcg aag gcg gct gct gct gtt act gcg gtt agt ggg gag cag ata cta	384
Ala Lys Ala Ala Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu	
115 120 125	
aaa gct att gtt gat gct gct ggt gat gcg gat cag gcg ggt aag aag	432
Lys Ala Ile Val Asp Ala Ala Gly Asp Ala Asp Gln Ala Gly Lys Lys	
130 135 140	
gct gct gag gct aag aat ccg att gca gct gcg att ggg gct aat gct	480
Ala Ala Glu Ala Lys Asn Pro Ile Ala Ala Ile Gly Ala Asn Ala	
145 150 155 160	
gct gat aat gcg gcg gcg ttt ggt aag gat gag atg aag aag agt gat	528
Ala Asp Asn Ala Ala Ala Phe Gly Lys Asp Glu Met Lys Lys Ser Asp	
165 170 175	
aag att gct gca gct att gtt ttg agg ggg gtg gct aag gat gga aag	576
Lys Ile Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys	
180 185 190	
ttt gct gtt gct aat gct aat gat gat aag aag gcg agt gtg aag agt	624
Phe Ala Val Ala Asn Ala Asn Asp Lys Lys Ala Ser Val Lys Ser	
195 200 205	
gct gtg	630
Ala Val	
210	

<210> SEQ ID NO 65

<211> LENGTH: 210

<212> TYPE: PRT

<213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 65

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr
1 5 10 15

Ala Ala Asp Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Ser Glu Lys
20 25 30

Ile Gly Asp Ala Gly Asp Asn Asn Asn Gly Ala Val Ala Asp Glu Asn
35 40 45

Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala
50 55 60

Gly Lys Ala Phe Gly Lys Asp Gly Lys Asp Gly Asp Ala Leu Lys Asp
65 70 75 80

Val Glu Thr Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys
85 90 95

Leu Phe Ala Gly Ala Asn Gly Asn Ala Gly Ala Ala Val Gly Asp Ile
100 105 110

Ala Lys Ala Ala Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu
115 120 125

Lys Ala Ile Val Asp Ala Ala Gly Asp Ala Asp Gln Ala Gly Lys Lys
130 135 140

Ala Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asn Ala

-continued

145	150	155	160
Ala Asp Asn Ala Ala Ala Phe Gly Lys Asp Glu Met Lys Lys Ser Asp	165	170	175
Lys Ile Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys	180	185	190
Phe Ala Val Ala Asn Ala Asn Asp Asp Lys Lys Ala Ser Val Lys Ser	195	200	205
Ala Val			
210			
<210> SEQ ID NO 66			
<211> LENGTH: 612			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia afzelii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(612)			
<400> SEQUENCE: 66			
gag agt gct gtg gat gag gtt agc aag tgg tta gaa gag atg ata aca			48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr			
1 5 10 15			
gct gct aag gag gct gct aca aag ggt ggt act ggt ggt aat aac gaa			96
Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Asn Asn Glu			
20 25 30			
aag att gga gat tct gat gct aat aat ggt gcg aag gct gat gcg agc			144
Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Lys Ala Asp Ala Ser			
35 40 45			
agt gtt aat ggg att gcg aat ggg ata aag ggg att gtt gat gct gct			192
Ser Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala			
50 55 60			
ggg aag gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt aaa aca			240
Gly Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr			
65 70 75 80			
gtt gct gct gag aat gag gct aac aag gat gcg ggg aag ttg ttt gct			288
Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala			
85 90 95			
ggt aag aat ggt aat gct gat gct gct gat gct gct gac att gcg aag			336
Gly Lys Asn Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys			
100 105 110			
gcg gct ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa gct			384
Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala			
115 120 125			
att gtt gat ggt gct ggt gat gca gct aat cag gcg ggt aaa aag gct			432
Ile Val Asp Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala			
130 135 140			
gct gag gct aag aat ccg att gcg gct gcg att ggg act aat gaa gct			480
Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala			
145 150 155 160			
ggg gcg gag ttt ggt gat gat atg aag aag aga aat gat aag att gct			528
Gly Ala Glu Phe Gly Asp Asp Met Lys Arg Asn Asp Lys Ile Ala			
165 170 175			
gcg gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gtt			576
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val			
180 185 190			
gct aat gct gct gct gat aat agt aag gcg agt gtg			612
Ala Asn Ala Ala Ala Asp Asn Ser Lys Ala Ser Val			
195 200			

-continued

<210> SEQ ID NO 67
 <211> LENGTH: 204
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 67

```

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr
1          5          10          15
Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Asn Asn Glu
20          25          30
Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Lys Ala Asp Ala Ser
35          40          45
Ser Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala
50          55          60
Gly Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr
65          70          75          80
Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala
85          90          95
Gly Lys Asn Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys
100         105         110
Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala
115         120         125
Ile Val Asp Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala
130         135         140
Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala
145         150         155         160
Gly Ala Glu Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala
165         170         175
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val
180         185         190
Ala Asn Ala Ala Ala Asp Asn Ser Lys Ala Ser Val
195         200

```

<210> SEQ ID NO 68
 <211> LENGTH: 609
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(609)

<400> SEQUENCE: 68

```

aag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aag      48
Lys Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys
1          5          10          15
gct gct ggt gag gct gct aca aag ggt ggt gat gct ggt ggt ggt gct      96
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala
20          25          30
gat aag att ggg gat gct ggt gat aag ggt gct gta gct gat gcg agc      144
Asp Lys Ile Gly Asp Ala Gly Asp Lys Gly Ala Val Ala Asp Ala Ser
35          40          45
agt gtt aag gag att gcg aat ggg ata aag ggg att gtt gat gct gct      192
Ser Val Lys Glu Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala
50          55          60
ggg aag gct ttt gcc aag gag ggt agt gcg ctg aag gat gtt aaa aca      240

```

-continued

Gly	Lys	Ala	Phe	Gly	Lys	Glu	Gly	Ser	Ala	Leu	Lys	Asp	Val	Lys	Thr	
65					70					75					80	
ggt	gct	gct	gag	aat	gag	gct	aac	aag	gat	gcg	ggg	aag	ttg	ttt	gct	288
Val	Ala	Ala	Glu	Asn	Glu	Ala	Asn	Lys	Asp	Ala	Gly	Lys	Leu	Phe	Ala	
				85					90					95		
ggt	aat	gct	ggt	aat	ggt	gct	gct	gat	gac	att	gcg	aag	gct	gct	gct	336
Gly	Asn	Ala	Gly	Asn	Gly	Ala	Ala	Asp	Asp	Ile	Ala	Lys	Ala	Ala	Ala	
			100					105					110			
gct	ggt	act	gct	ggt	agt	ggg	gag	cag	ata	ctg	aaa	gct	att	ggt	gat	384
Ala	Val	Thr	Ala	Val	Ser	Gly	Glu	Gln	Ile	Leu	Lys	Ala	Ile	Val	Asp	
			115				120					125				
gct	gct	ggt	gat	aag	gct	aat	cag	gat	ggt	aaa	aag	gct	gct	gat	gct	432
Ala	Ala	Gly	Asp	Lys	Ala	Asn	Gln	Asp	Gly	Lys	Lys	Ala	Ala	Asp	Ala	
		130				135					140					
aag	aat	ccg	att	gct	gct	gct	att	ggg	gct	gct	gat	gct	ggt	gct	gct	480
Lys	Asn	Pro	Ile	Ala	Ala	Ala	Ile	Gly	Ala	Ala	Asp	Ala	Gly	Ala	Ala	
145				150						155				160		
gct	gct	ttt	aat	gag	aat	gat	atg	aag	aag	agt	gat	aag	att	gct	gca	528
Ala	Ala	Phe	Asn	Glu	Asn	Asp	Met	Lys	Lys	Ser	Asp	Lys	Ile	Ala	Ala	
			165					170						175		
gct	att	ggt	ttg	agg	ggg	gtg	gct	aag	gat	gga	aag	ttt	gct	gct	gct	576
Ala	Ile	Val	Leu	Arg	Gly	Val	Ala	Lys	Asp	Gly	Lys	Phe	Ala	Ala	Ala	
			180					185					190			
gat	gct	gat	gct	aat	aat	agt	aag	gct	agc	gtg						609
Asp	Ala	Asp	Ala	Asn	Asn	Ser	Lys	Ala	Ser	Val						
			195				200									

<210> SEQ ID NO 69

<211> LENGTH: 203

<212> TYPE: PRT

<213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 69

Lys	Ser	Ala	Val	Asp	Glu	Val	Ser	Lys	Trp	Leu	Glu	Glu	Met	Ile	Lys	
1				5					10					15		
Ala	Ala	Gly	Glu	Ala	Ala	Thr	Lys	Gly	Gly	Asp	Ala	Gly	Gly	Gly	Ala	
			20					25					30			
Asp	Lys	Ile	Gly	Asp	Ala	Gly	Asp	Lys	Gly	Ala	Val	Ala	Asp	Ala	Ser	
		35				40					45					
Ser	Val	Lys	Glu	Ile	Ala	Asn	Gly	Ile	Lys	Gly	Ile	Val	Asp	Ala	Ala	
	50				55						60					
Gly	Lys	Ala	Phe	Gly	Lys	Glu	Gly	Ser	Ala	Leu	Lys	Asp	Val	Lys	Thr	
65					70					75					80	
Val	Ala	Ala	Glu	Asn	Glu	Ala	Asn	Lys	Asp	Ala	Gly	Lys	Leu	Phe	Ala	
			85						90					95		
Gly	Asn	Ala	Gly	Asn	Gly	Ala	Ala	Asp	Asp	Ile	Ala	Lys	Ala	Ala	Ala	
		100						105					110			
Ala	Val	Thr	Ala	Val	Ser	Gly	Glu	Gln	Ile	Leu	Lys	Ala	Ile	Val	Asp	
		115					120					125				
Ala	Ala	Gly	Asp	Lys	Ala	Asn	Gln	Asp	Gly	Lys	Lys	Ala	Ala	Asp	Ala	
	130					135					140					
Lys	Asn	Pro	Ile	Ala	Ala	Ala	Ile	Gly	Ala	Ala	Asp	Ala	Gly	Ala	Ala	
145				150						155				160		
Ala	Ala	Phe	Asn	Glu	Asn	Asp	Met	Lys	Lys	Ser	Asp	Lys	Ile	Ala	Ala	
			165					170						175		

-continued

Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala
180 185 190

Asp Ala Asp Ala Asn Asn Ser Lys Ala Ser Val
195 200

<210> SEQ ID NO 70
 <211> LENGTH: 600
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(600)

<400> SEQUENCE: 70

aag agt gct gtt ggt gag gtt agc aag tgg tta gaa gag atg ata aaa 48
 Lys Ser Ala Val Gly Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys
 1 5 10 15

gct gct ggt gag gct gca aaa gtt ggt ggt act ggt ggt agc gaa aag 96
 Ala Ala Gly Glu Ala Ala Lys Val Gly Gly Thr Gly Gly Ser Glu Lys
 20 25 30

att ggg gat gct gat aat aat aag ggt gct gta gct gat gcg agc agt 144
 Ile Gly Asp Ala Asp Asn Asn Lys Gly Ala Val Ala Asp Ala Ser Ser
 35 40 45

gtt aat ggg att gcg aat ggg ata aag ggg att gtt gat gct gct ggg 192
 Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
 50 55 60

aag gct ttt ggt aag gat ggt gcg ctg gca ggt gtt gca gct gct gct 240
 Lys Ala Phe Gly Lys Asp Gly Ala Leu Ala Gly Val Ala Ala Ala Ala
 65 70 75 80

gag aat gat gat aag aag gat gcg ggg aag ttg ttt gct ggt aag aat 288
 Glu Asn Asp Asp Lys Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
 85 90 95

ggt ggt gct ggt gct gct gat gcg att ggg aag gcg gct gct gct gtt 336
 Gly Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Val
 100 105 110

act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gat gct gct 384
 Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala
 115 120 125

ggt gct gca gct aat cag gcg ggt aaa aag gct gcg gat gct aag aat 432
 Gly Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn
 130 135 140

ccg att gcg gct gcg att ggg act gct gat gat ggg gcg gag ttt aag 480
 Pro Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys
 145 150 155 160

gat gat atg aag aag agt gat aat att gct gcg gct att gtt ttg agg 528
 Asp Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg
 165 170 175

ggg gtg gct aag gat gga aag ttt gct gtt gct aat gct gat gat aat 576
 Gly Val Ala Lys Asp Gly Lys Phe Ala Val Ala Asn Ala Asp Asp Asn
 180 185 190

aag gcg agt gtg aag agt gct gtg 600
 Lys Ala Ser Val Lys Ser Ala Val
 195 200

<210> SEQ ID NO 71
 <211> LENGTH: 200
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 71

-continued

Lys Ser Ala Val Gly Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys
 1 5 10 15
 Ala Ala Gly Glu Ala Ala Lys Val Gly Gly Thr Gly Gly Ser Glu Lys
 20 25 30
 Ile Gly Asp Ala Asp Asn Asn Lys Gly Ala Val Ala Asp Ala Ser Ser
 35 40 45
 Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
 50 55 60
 Lys Ala Phe Gly Lys Asp Gly Ala Leu Ala Gly Val Ala Ala Ala Ala
 65 70 75 80
 Glu Asn Asp Asp Lys Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
 85 90 95
 Gly Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Ala Val
 100 105 110
 Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala
 115 120 125
 Gly Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn
 130 135 140
 Pro Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys
 145 150 155 160
 Asp Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg
 165 170 175
 Gly Val Ala Lys Asp Gly Lys Phe Ala Val Ala Asn Ala Asp Asp Asn
 180 185 190
 Lys Ala Ser Val Lys Ser Ala Val
 195 200

<210> SEQ ID NO 72
 <211> LENGTH: 592
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(591)

<400> SEQUENCE: 72

gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca	48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr	
1 5 10 15	
gct gct ggt gag gct gca aaa gtt ggt gct ggt ggt gct gat aag	96
Ala Ala Gly Glu Ala Ala Lys Val Gly Ala Gly Gly Ala Asp Lys	
20 25 30	
att ggg gat gct gct aat aat cag ggt gcg aag gct gat gag agc agt	144
Ile Gly Asp Ala Ala Asn Asn Gln Gly Ala Lys Ala Asp Glu Ser Ser	
35 40 45	
gtt aat gga att gca aag ggg ata aag ggg att gtt gat gct gct ggg	192
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly	
50 55 60	
aag gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt gca aaa gtt	240
Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val	
65 70 75 80	
gct gat gat gat aac aag gat gcg ggg aag ttg ttt gct ggt aat gct	288
Ala Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Asn Ala	
85 90 95	
ggg ggt ggt gct ggt gct gat att gcg aag gcg gct gct gct gtt act	336

-continued

Gly	Gly	Gly	Ala	Gly	Ala	Asp	Ile	Ala	Lys	Ala	Ala	Ala	Ala	Val	Thr	
			100					105						110		
gcg	gtt	agt	ggg	gag	cag	ata	ctg	aaa	gct	att	gtt	gat	gct	gct	ggt	384
Ala	Val	Ser	Gly	Glu	Gln	Ile	Leu	Lys	Ala	Ile	Val	Asp	Ala	Ala	Gly	
			115				120					125				
gct	gcg	gat	cag	gcg	ggt	gca	gct	gct	ggt	gcg	gct	aag	aat	ccg	att	432
Ala	Ala	Asp	Gln	Ala	Gly	Ala	Ala	Ala	Gly	Ala	Ala	Lys	Asn	Pro	Ile	
			130			135						140				
gcg	gct	gcg	att	ggg	gct	gat	gct	ggt	gct	gcg	gag	gag	ttt	aag	gat	480
Ala	Ala	Ala	Ile	Gly	Ala	Asp	Ala	Gly	Ala	Ala	Glu	Glu	Phe	Lys	Asp	
			145			150				155					160	
gag	atg	aag	aag	agt	gat	aag	att	gct	gcg	gct	att	gtt	ttg	agg	ggg	528
Glu	Met	Lys	Lys	Ser	Asp	Lys	Ile	Ala	Ala	Ala	Ile	Val	Leu	Arg	Gly	
				165				170						175		
gtg	gct	aag	ggt	gga	aag	ttt	gct	gtt	gct	gct	aat	gat	gct	gca	aat	576
Val	Ala	Lys	Gly	Gly	Lys	Phe	Ala	Val	Ala	Ala	Asn	Asp	Ala	Ala	Asn	
			180				185							190		
gtg	aag	agt	gct	gtg	g											592
Val	Lys	Ser	Ala	Val												
			195													

<210> SEQ ID NO 73
 <211> LENGTH: 197
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 73

Glu	Ser	Ala	Val	Asp	Glu	Val	Ser	Lys	Trp	Leu	Glu	Glu	Met	Ile	Thr	
1				5				10						15		
Ala	Ala	Gly	Glu	Ala	Ala	Lys	Val	Gly	Ala	Gly	Gly	Gly	Ala	Asp	Lys	
			20					25					30			
Ile	Gly	Asp	Ala	Ala	Asn	Asn	Gln	Gly	Ala	Lys	Ala	Asp	Glu	Ser	Ser	
		35				40						45				
Val	Asn	Gly	Ile	Ala	Lys	Gly	Ile	Lys	Gly	Ile	Val	Asp	Ala	Ala	Gly	
	50					55					60					
Lys	Ala	Phe	Gly	Lys	Glu	Gly	Ser	Ala	Leu	Lys	Asp	Val	Ala	Lys	Val	
	65				70				75					80		
Ala	Asp	Asp	Asp	Asn	Lys	Asp	Ala	Gly	Lys	Leu	Phe	Ala	Gly	Asn	Ala	
			85					90						95		
Gly	Gly	Gly	Ala	Gly	Ala	Asp	Ile	Ala	Lys	Ala	Ala	Ala	Ala	Val	Thr	
			100				105							110		
Ala	Val	Ser	Gly	Glu	Gln	Ile	Leu	Lys	Ala	Ile	Val	Asp	Ala	Ala	Gly	
			115				120					125				
Ala	Ala	Asp	Gln	Ala	Gly	Ala	Ala	Ala	Gly	Ala	Ala	Lys	Asn	Pro	Ile	
			130			135						140				
Ala	Ala	Ala	Ile	Gly	Ala	Asp	Ala	Gly	Ala	Ala	Glu	Glu	Phe	Lys	Asp	
			145			150				155					160	
Glu	Met	Lys	Lys	Ser	Asp	Lys	Ile	Ala	Ala	Ala	Ile	Val	Leu	Arg	Gly	
				165				170						175		
Val	Ala	Lys	Gly	Gly	Lys	Phe	Ala	Val	Ala	Ala	Asn	Asp	Ala	Ala	Asn	
			180				185						190			
Val	Lys	Ser	Ala	Val												
			195													

<210> SEQ ID NO 74

-continued

```

<211> LENGTH: 597
<212> TYPE: DNA
<213> ORGANISM: Borrelia afzelii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(597)

<400> SEQUENCE: 74

gag agt gct gtt ggt gag gtt agc gca tgg tta gaa gag atg ata aca      48
Glu Ser Ala Val Gly Glu Val Ser Ala Trp Leu Glu Glu Met Ile Thr
1          5          10          15

gct gct agt gag gct gct aca aag ggt ggt act ggt ggt act ggt ggt      96
Ala Ala Ser Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Thr Gly Gly
20          25          30

gat agt gaa aag att ggg gat tct gat gct aat aat ggt gct gta gct      144
Asp Ser Glu Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Val Ala
35          40          45

gat gcg agc agt gtt aag gag att gcg aag ggg ata aag ggg att gtt      192
Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val
50          55          60

gat gct gct ggg aag gct ttt ggt aag gat ggt aat gcg ctg aag gat      240
Asp Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp
65          70          75          80

gtt gca gaa gtt gct gat gat gag gct aac gcg gat gcg ggg aag ttg      288
Val Ala Glu Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu
85          90          95

ttt gct ggt aat gct ggt aat gct gct gct gct gac gtt gcg aag gcg      336
Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Ala Asp Val Ala Lys Ala
100         105         110

gct ggt gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att      384
Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile
115         120         125

gtt gat gct gct ggt gct gcg gat cag gcg ggt gca aag gct gat gcg      432
Val Asp Ala Ala Gly Ala Ala Asp Gln Ala Gly Ala Lys Ala Asp Ala
130         135         140

gct aag aat ccg att gca gct gcg att ggg act aat gaa gct ggg gcg      480
Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala
145         150         155         160

gcg ttt aag gat gga atg aag aag aga aat gat aat att gct gcg gct      528
Ala Phe Lys Asp Gly Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala
165         170         175

att gtt ttg agg ggg gtg gct aag agt gga aag ttt gct gtt gct gct      576
Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Ala
180         185         190

gct gat gct ggt aag gcg aga      597
Ala Asp Ala Gly Lys Ala Arg
195

```

```

<210> SEQ ID NO 75
<211> LENGTH: 199
<212> TYPE: PRT
<213> ORGANISM: Borrelia afzelii

```

```

<400> SEQUENCE: 75

Glu Ser Ala Val Gly Glu Val Ser Ala Trp Leu Glu Glu Met Ile Thr
1          5          10          15

Ala Ala Ser Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Thr Gly Gly
20          25          30

Asp Ser Glu Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Val Ala

```

-continued

35	40	45	
Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val			
50	55	60	
Asp Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp			
65	70	75	80
Val Ala Glu Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu			
	85	90	95
Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Ala Asp Val Ala Lys Ala			
	100	105	110
Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile			
	115	120	125
Val Asp Ala Ala Gly Ala Ala Asp Gln Ala Gly Ala Lys Ala Asp Ala			
	130	135	140
Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala			
	145	150	155
Ala Phe Lys Asp Gly Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala			
	165	170	175
Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Ala			
	180	185	190
Ala Asp Ala Gly Lys Ala Arg			
	195		
<210> SEQ ID NO 76			
<211> LENGTH: 621			
<212> TYPE: DNA			
<213> ORGANISM: Borrelia afzelii			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(621)			
<400> SEQUENCE: 76			
gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca			48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr			
1	5	10	15
gct gct agt gag gct gca aaa gtt ggt gct ggt gat gat aag att			96
Ala Ala Ser Glu Ala Ala Lys Val Gly Ala Gly Gly Asp Asp Lys Ile			
	20	25	30
ggg gat tct gct aat aat ggt gct gta gct gat gcg ggc agt gtt aag			144
Gly Asp Ser Ala Asn Asn Gly Ala Val Ala Asp Ala Gly Ser Val Lys			
	35	40	45
gga att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag gct			192
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala			
	50	55	60
ttt ggt aag gag ggt gat gcg ctg aag gat gtt gca aaa gtt gct gat			240
Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala Asp			
	65	70	75
gag aat ggg gat aac aag gat gcg ggg aag ttg ttt gct ggt gag aat			288
Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu Asn			
	85	90	95
ggt aat gct ggt ggt gct gct gat gct gac att gcg aag gcg gct gct			336
Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala Ala			
	100	105	110
gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gag			384
Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu			
	115	120	125
gct gct ggt gct ggt gat gca gct aat cag gcg ggt aag aag gct gat			432

-continued

Ala Ala Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Asp	
130 135 140	
gag gct aag aat ccg att gcg gct gcg att ggg act gat gat gct ggg	480
Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly	
145 150 155 160	
gcg gcg ttt ggt cag gat gat atg aag aag aga aat gat aat att gct	528
Ala Ala Phe Gly Gln Asp Asp Met Lys Lys Arg Asn Asp Asn Ile Ala	
165 170 175	
gcg gct att gtt ttg agg ggg gtg gct aag ggt gga aag ttt gct gtt	576
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val	
180 185 190	
gct aat gct gct aat gat agt aag gcg agt gtg aag agt gct gtg	621
Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val	
195 200 205	

<210> SEQ ID NO 77
 <211> LENGTH: 207
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 77

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr	
1 5 10 15	
Ala Ala Ser Glu Ala Ala Lys Val Gly Ala Gly Gly Asp Asp Lys Ile	
20 25 30	
Gly Asp Ser Ala Asn Asn Gly Ala Val Ala Asp Ala Gly Ser Val Lys	
35 40 45	
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala	
50 55 60	
Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala Asp	
65 70 75 80	
Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu Asn	
85 90 95	
Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala Ala	
100 105 110	
Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu	
115 120 125	
Ala Ala Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Asp	
130 135 140	
Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly	
145 150 155 160	
Ala Ala Phe Gly Gln Asp Asp Met Lys Lys Arg Asn Asp Asn Ile Ala	
165 170 175	
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val	
180 185 190	
Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val	
195 200 205	

<210> SEQ ID NO 78
 <211> LENGTH: 459
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(459)

<400> SEQUENCE: 78

-continued

gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag ata ata aca	48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Ile Ile Thr	
1 5 10 15	
gct act ggg aag gct ttt ggt aag gat ggt aat gcg ctg gca ggt gtt	96
Ala Thr Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Ala Gly Val	
20 25 30	
gca aaa gtt gct gat gat gag gct aac gcg gat gcg ggg aag ttg ttt	144
Ala Lys Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu Phe	
35 40 45	
gct ggt gag aat ggt aat gct ggt gct gct gcg att ggg aag gcg gct	192
Ala Gly Glu Asn Gly Asn Ala Gly Ala Ala Ala Ile Gly Lys Ala Ala	
50 55 60	
gct gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt	240
Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val	
65 70 75 80	
gat gct gct ggt ggt gcg gct cag gtg ggt gct ggt gct ggt gcg gct	288
Asp Ala Ala Gly Gly Ala Ala Gln Val Gly Ala Gly Ala Gly Ala Ala	
85 90 95	
acg aat ccg att gca gct gcg att ggg gct gct ggt gat ggt gcg gat	336
Thr Asn Pro Ile Ala Ala Ala Ile Gly Ala Ala Gly Asp Gly Ala Asp	
100 105 110	
ttt ggt aag gat gag atg aag aag aga aat gat aag att gct gcg gct	384
Phe Gly Lys Asp Glu Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala	
115 120 125	
att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct gct gct	432
Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala Ala	
130 135 140	
aat gat agt aag gcg agt gtg aag agt	459
Asn Asp Ser Lys Ala Ser Val Lys Ser	
145 150	

<210> SEQ ID NO 79

<211> LENGTH: 153

<212> TYPE: PRT

<213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 79

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Ile Ile Thr
1 5 10 15
Ala Thr Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Ala Gly Val
20 25 30
Ala Lys Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu Phe
35 40 45
Ala Gly Glu Asn Gly Asn Ala Gly Ala Ala Ala Ile Gly Lys Ala Ala
50 55 60
Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val
65 70 75 80
Asp Ala Ala Gly Gly Ala Ala Gln Val Gly Ala Gly Ala Gly Ala Ala
85 90 95
Thr Asn Pro Ile Ala Ala Ala Ile Gly Ala Ala Gly Asp Gly Ala Asp
100 105 110
Phe Gly Lys Asp Glu Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala
115 120 125
Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala Ala
130 135 140

-continued

Asn Asp Ser Lys Ala Ser Val Lys Ser
145 150

<210> SEQ ID NO 80

<211> LENGTH: 612

<212> TYPE: DNA

<213> ORGANISM: Borrelia afzelii

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(612)

<400> SEQUENCE: 80

gct gtg gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg 48
Ala Val Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met
1 5 10 15

ata aca gct gct gat gct gct gct gct aaa gtt ggc gat gct ggt ggt 96
Ile Thr Ala Ala Asp Ala Ala Ala Ala Lys Val Gly Asp Ala Gly Gly
20 25 30

ggt gct gat aag att ggg gat gtt ggt gct gct aat aag ggt gcg aag 144
Gly Ala Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Lys Gly Ala Lys
35 40 45

gct gat gcg agc agt gtt aag gag att gcg aag ggg ata aag ggg att 192
Ala Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile
50 55 60

gtt gat gct gct ggg aag gct ttt ggt ggt gat gcg ctg aag gat gtt 240
Val Asp Ala Ala Gly Lys Ala Phe Gly Gly Asp Ala Leu Lys Asp Val
65 70 75 80

aaa gct gct ggt gat gat aac aag gag gca ggg aag ttg ttt gct ggt 288
Lys Ala Ala Gly Asp Asp Asn Lys Glu Ala Gly Lys Leu Phe Ala Gly
85 90 95

gct aat ggt aat gct ggt gct aat gct gct gct gct gat gac att gcg 336
Ala Asn Gly Asn Ala Gly Ala Asn Ala Ala Ala Asp Asp Ile Ala
100 105 110

aag gcg gct ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa 384
Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys
115 120 125

gct att gtt gag gcg gct ggt gct gcg gat cag gcg ggt gta aag gct 432
Ala Ile Val Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala
130 135 140

gag gag gct aag aat ccg att gca gct gcg att ggg act gat gat gct 480
Glu Glu Ala Lys Asn Pro Ile Ala Ala Ile Gly Thr Asp Asp Ala
145 150 155 160

ggt gcg gcg gag ttt ggt gag aat gat atg aag aag aat gat aat att 528
Gly Ala Ala Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile
165 170 175

gct gcg gct att gtt ttg agg ggg gtg gct aag agt gga aag ttt gct 576
Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala
180 185 190

gct aat gct aat gat gct ggt aag aag gag agt gtg 612
Ala Asn Ala Asn Asp Ala Gly Lys Lys Glu Ser Val
195 200

<210> SEQ ID NO 81

<211> LENGTH: 204

<212> TYPE: PRT

<213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 81

Ala Val Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met
1 5 10 15

-continued

```

Ile Thr Ala Ala Asp Ala Ala Ala Ala Lys Val Gly Asp Ala Gly Gly
      20                      25                      30
Gly Ala Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Lys Gly Ala Lys
      35                      40                      45
Ala Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile
      50                      55                      60
Val Asp Ala Ala Gly Lys Ala Phe Gly Gly Asp Ala Leu Lys Asp Val
      65                      70                      75                      80
Lys Ala Ala Gly Asp Asp Asn Lys Glu Ala Gly Lys Leu Phe Ala Gly
      85                      90                      95
Ala Asn Gly Asn Ala Gly Ala Asn Ala Ala Ala Ala Asp Asp Ile Ala
      100                     105                     110
Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys
      115                     120                     125
Ala Ile Val Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala
      130                     135                     140
Glu Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala
      145                     150                     155                     160
Gly Ala Ala Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile
      165                     170                     175
Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala
      180                     185                     190
Ala Asn Ala Asn Asp Ala Gly Lys Lys Glu Ser Val
      195                     200

```

```

<210> SEQ ID NO 82
<211> LENGTH: 603
<212> TYPE: DNA
<213> ORGANISM: Borrelia afzelii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(603)

```

```

<400> SEQUENCE: 82

```

```

aag agt gct gtg gat gag gct agc aag tgg tta gaa gag atg ata aca      48
Lys Ser Ala Val Asp Glu Ala Ser Lys Trp Leu Glu Glu Met Ile Thr
1                      5                      10                      15

gct gct ggt gag gct gct aca aag ggt ggt act ggt gaa gct agc gaa      96
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu
20                     25                     30

aag att ggg gat gtt ggt gat aat aat cat ggt gct gta gct gat gcg      144
Lys Ile Gly Asp Val Gly Asp Asn Asn His Gly Ala Val Ala Asp Ala
35                     40                     45

gac agt gtt aag ggg att gcg aag ggg ata aag ggg att gtt gat gct      192
Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala
50                     55                     60

gct ggg aag gct ttt ggt aag gat ggt gcg ctg aag gat gtt gca gct      240
Ala Gly Lys Ala Phe Gly Lys Asp Gly Ala Leu Lys Asp Val Ala Ala
65                     70                     75                     80

gct gct ggt gat gag gct aac aag gat gcg ggg aag ttg ttt gct ggt      288
Ala Ala Gly Asp Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly
85                     90                     95

cag gat ggt ggt ggt gct gat ggt gac att gcg aag gcg gct gct gct      336
Gln Asp Gly Gly Glu Ala Asp Gly Asp Ile Ala Lys Ala Ala Ala Ala
100                    105                    110

```

-continued

ggt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gag gct	384
Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala	
115 120 125	
gct ggt gat aag gct aat cag gtg ggt gta aag gct gct ggt gcg gct	432
Ala Gly Asp Lys Ala Asn Gln Val Gly Val Lys Ala Ala Gly Ala Ala	
130 135 140	
acg aat ccg att gca gct gcg att ggg act gat gat gat aat gcg gcg	480
Thr Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Asp Asn Ala Ala	
145 150 155 160	
gcg ttt gat aag gat gag atg aag aag agt aat gat aag att gct gcg	528
Ala Phe Asp Lys Asp Glu Met Lys Lys Ser Asn Asp Lys Ile Ala Ala	
165 170 175	
gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct aat	576
Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Asn	
180 185 190	
gct aat gat aat agt aag gcg agt gtg	603
Ala Asn Asp Asn Ser Lys Ala Ser Val	
195 200	

<210> SEQ ID NO 83
 <211> LENGTH: 201
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 83

Lys Ser Ala Val Asp Glu Ala Ser Lys Trp Leu Glu Glu Met Ile Thr	
1 5 10 15	
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu	
20 25 30	
Lys Ile Gly Asp Val Gly Asp Asn Asn His Gly Ala Val Ala Asp Ala	
35 40 45	
Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala	
50 55 60	
Ala Gly Lys Ala Phe Gly Lys Asp Gly Ala Leu Lys Asp Val Ala Ala	
65 70 75 80	
Ala Ala Gly Asp Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly	
85 90 95	
Gln Asp Gly Gly Gly Ala Asp Gly Asp Ile Ala Lys Ala Ala Ala Ala	
100 105 110	
Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala	
115 120 125	
Ala Gly Asp Lys Ala Asn Gln Val Gly Val Lys Ala Ala Gly Ala Ala	
130 135 140	
Thr Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Asp Asn Ala Ala	
145 150 155 160	
Ala Phe Asp Lys Asp Glu Met Lys Lys Ser Asn Asp Lys Ile Ala Ala	
165 170 175	
Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Asn	
180 185 190	
Ala Asn Asp Asn Ser Lys Ala Ser Val	
195 200	

<210> SEQ ID NO 84
 <211> LENGTH: 249
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii

-continued

```

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(249)

<400> SEQUENCE: 84

aag agt gct gtg gat gag gtt agc aag tgg tta gaa gag atg ata aca      48
Lys Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr
1          5          10          15

gct gct agt gat gct gct aca aag ggt ggt act ggt gaa gct agc gaa      96
Ala Ala Ser Asp Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu
          20          25          30

aag att ggg gat tct gat gct aat aag ggt gct ggt gct ggg gcg gcg     144
Lys Ile Gly Asp Ser Asp Ala Asn Lys Gly Ala Gly Ala Gly Ala Ala
          35          40          45

ttt ggt gag aat gat atg aag aag aga aat gat aat att gct gca gct     192
Phe Gly Glu Asn Asp Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala
          50          55          60

att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gtt aag gag     240
Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val Lys Glu
65          70          75          80

gat tat tga
Asp Tyr
249

<210> SEQ ID NO 85
<211> LENGTH: 82
<212> TYPE: PRT
<213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 85

Lys Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr
1          5          10          15

Ala Ala Ser Asp Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu
          20          25          30

Lys Ile Gly Asp Ser Asp Ala Asn Lys Gly Ala Gly Ala Gly Ala Ala
          35          40          45

Phe Gly Glu Asn Asp Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala
          50          55          60

Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val Lys Glu
65          70          75          80

Asp Tyr

<210> SEQ ID NO 86
<211> LENGTH: 537
<212> TYPE: DNA
<213> ORGANISM: Borrelia afzelii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(537)

<400> SEQUENCE: 86

atg gaa aaa ata gaa aaa ttt aaa aac aaa tgt caa cat aaa cta caa      48
Met Glu Lys Ile Glu Lys Phe Lys Asn Lys Cys Gln His Lys Leu Gln
1          5          10          15

cat aaa cta atc gta tta gta tca aca ctt tgc tat ata aac aat aaa      96
His Lys Leu Ile Val Leu Val Ser Thr Leu Cys Tyr Ile Asn Asn Lys
          20          25          30

aat aaa aaa tat tca caa agc aac atc ctt tat tat ttt aat gaa aat     144
Asn Lys Lys Tyr Ser Gln Ser Asn Ile Leu Tyr Tyr Phe Asn Glu Asn
          35          40          45

```

-continued

tta aaa aga aat ggg caa acc cct att aaa ata aaa aca tta caa aac Leu Lys Arg Asn Gly Gln Thr Pro Ile Lys Ile Lys Thr Leu Gln Asn 50 55 60	192
tat ctt tat aaa ctg gaa aaa gaa ttt gaa gta aca act aat tat tat Tyr Leu Tyr Lys Leu Glu Lys Glu Phe Glu Val Thr Thr Asn Tyr Tyr 65 70 75 80	240
aaa cac ttg ggg gtt aat tgt gga acc gaa att tac tat aaa ctt aaa Lys His Leu Gly Val Asn Cys Gly Thr Glu Ile Tyr Tyr Lys Leu Lys 85 90 95	288
tat caa aaa caa aaa tgc tat cat aaa ata aac caa tat ttt aaa aag Tyr Gln Lys Gln Lys Cys Tyr His Lys Ile Asn Gln Tyr Phe Lys Lys 100 105 110	336
aaa aaa gaa att aaa ttt aac tta aga gta agt gca ttt ttt aat aaa Lys Lys Glu Ile Lys Phe Asn Leu Arg Val Ser Ala Phe Phe Asn Lys 115 120 125	384
aaa cac tca aaa aaa ggg agt gta gaa tta aag gaa tgt aat aat aat Lys His Ser Lys Lys Gly Ser Val Glu Leu Lys Glu Cys Asn Asn Asn 130 135 140	432
aat aat aat aaa gag aaa gaa aca tcc caa aaa att gaa att tta caa Asn Asn Asn Lys Glu Lys Glu Thr Ser Gln Lys Ile Glu Ile Leu Gln 145 150 155 160	480
aca aaa gtc tat gcc aaa aaa tgt aaa ttt ttg aca aac tac tat act Thr Lys Val Tyr Ala Lys Lys Cys Lys Phe Leu Thr Asn Tyr Tyr Thr 165 170 175	528
aaa att tta Lys Ile Leu	537

<210> SEQ ID NO 87
 <211> LENGTH: 179
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 87

Met Glu Lys Ile Glu Lys Phe Lys Asn Lys Cys Gln His Lys Leu Gln 1 5 10 15
His Lys Leu Ile Val Leu Val Ser Thr Leu Cys Tyr Ile Asn Asn Lys 20 25 30
Asn Lys Lys Tyr Ser Gln Ser Asn Ile Leu Tyr Tyr Phe Asn Glu Asn 35 40 45
Leu Lys Arg Asn Gly Gln Thr Pro Ile Lys Ile Lys Thr Leu Gln Asn 50 55 60
Tyr Leu Tyr Lys Leu Glu Lys Glu Phe Glu Val Thr Thr Asn Tyr Tyr 65 70 75 80
Lys His Leu Gly Val Asn Cys Gly Thr Glu Ile Tyr Tyr Lys Leu Lys 85 90 95
Tyr Gln Lys Gln Lys Cys Tyr His Lys Ile Asn Gln Tyr Phe Lys Lys 100 105 110
Lys Lys Glu Ile Lys Phe Asn Leu Arg Val Ser Ala Phe Phe Asn Lys 115 120 125
Lys His Ser Lys Lys Gly Ser Val Glu Leu Lys Glu Cys Asn Asn Asn 130 135 140
Asn Asn Asn Lys Glu Lys Glu Thr Ser Gln Lys Ile Glu Ile Leu Gln 145 150 155 160
Thr Lys Val Tyr Ala Lys Lys Cys Lys Phe Leu Thr Asn Tyr Tyr Thr 165 170 175

Lys Ile Leu

-continued

<210> SEQ ID NO 88

<211> LENGTH: 2775

<212> TYPE: DNA

<213> ORGANISM: *Borrelia garinii*

<400> SEQUENCE: 88

```
cggaaatcaa gccaccta aa acaacttccc aaaagtttct caaaaaatat tatattcagc      60
agtaaatctc ataagtcatt aattatttaa tactattcaa cagtaaatcc tataagtcac      120
taattattta atactattca gcagtaaat ctataagtc ttaattattt aatactattc      180
agcagtaaat tctataagtc attaattatt taatactatt cagcagtaaa ttctataagt      240
cattaattat ttaatactat tcagcagtaa attctataag tcattaatta ttttaacta      300
ttcagcagta aattctataa gtcattaatt caattaggta acggattctt agatgtattc      360
acctcttttg gtggattagt tgcagatgca ttgggggtta aagctgatcc aaaaaaatct      420
gatgtaaaaa cttattttga atctctagct aaaaaattag aagaacaaa agatgggtta      480
actaagttgt ccaaaggtaa tgacggtgat actggaaagg ctggtgatgc tgggtgggct      540
ggtggtggcg ctagtgtgc aggtggcgct ggtgggattg agggcgctat aacagagatt      600
agcaaatggt tagatgatat ggcaaaagct gctgcggaag ctgcaagtgc tgctactggt      660
aatgcagcaa ttggggatgt tgttaatggt aatggtggag cagcaaaagg tgggtgatgcg      720
gagagtgtta atgggattgc taaggggata aaggggattg ttgatgtgc tgagaaggct      780
gatgcgaagg aagggaagtt ggtatgtggc ggtgatgctg gtggggctgg tgggtggcgct      840
ggtgtgctgc gtggcgctgg tgggattgag ggcgctataa cagagattag caaatgggta      900
gatgatatgg caaaagctgc tgcggttgct gcaagtgtgc caagtgtgc tactggtaat      960
gcagcaattg gggatgttgt taatggtaat gatggagcag caaaaggtgg tgatgcggcg      1020
agtgttaatg ggattgtcaa ggggataaag gggattgttg atgctgctga gaaggctgat      1080
gcgaaggaa ggaagttgga tgtggctggc gatgctggcg agggtaacaa ggatgctggg      1140
aagctgtttg tgaagaagaa tctgtgtgat gaggggtggc aagcaaatga tgctgggaag      1200
gctgtgctgc cgggtgtgct tgttagtggg gagcagatat taaaagcgat tgttgatgct      1260
gctgagggcg atgataagca gggtaagaag gctgcggatg ctacaaatcc gattgaggcg      1320
gctattgggg gtgcggatgc ggggtgctaat gctgaggcgt ttaataagat gaagaaggat      1380
gatcagattg ctgctgctat ggttctgagg ggaatggcta aggatgggca gtttgccttg      1440
aaggatgatg ctgctgctca tgaagggact gtaagaatg ctgttgatat ggcaaaggcc      1500
gctgcggaag ctgcaagtgc tgcaagtgc gctactgcta gtacaacgat tggagatgtt      1560
gttaagagtg gtgaggcaaa agatggtgat gcggcgagtg ttaatgggat tgctaagggg      1620
ataaagggga ttgttgatgc tgctgagaag gctgatgcga aggaagggaa gttggatgtg      1680
gctggtgctg ctggtacgac taacgtgaat gttgggaagt tgtttgtgaa gaataatggt      1740
aatgagggcg gtgatgcaag tgatgctggg aaagctgctg ctgcggttgc tgctgttagt      1800
ggggagcaga tattaagc gattgttgat gctgctaaag atggtgataa gcaggggggtt      1860
actgatgtaa aggatgctac aaatccgatt gagggcgcta ttgggggtac aaatgataat      1920
gatgctgcgg cgtttgctac tatgaagaag gatgatcaga ttgctgctgc tatggttctg      1980
aggggaatgg ctaaggatgg gcagtttgct ttgaaggatg atgctgctaa ggatggtgat      2040
```

-continued

```

aaaacggggg ttgctgcgga tgctgaaaat cggattgacg cggctattgg gggcgcgcat 2100
gctgatgctg cggcggttaa taaggagggg atgaagaagg atgatcagat tgctgctgct 2160
atggttctga ggggaatggc taaggatggg cagtttgctt tgacgaataa tgctgctgct 2220
catgaaggga ctgttaagaa tgctgttgat atggcaaaag ctgctgcggt tgctgcaagt 2280
gctgctactg gcaatgcagc aattggggat gttgttaaga gtaatggtag agcagcagca 2340
aaagtggtg atgcggcgag tgttaatggg attgctaagg ggataaaggg gattgttgat 2400
gctgctgaga aggctgatgc gaaggaaggg aagttggatg tggctggtgc tgctggtgaa 2460
actaacaagg atgctgggaa gttgtttgtg aagaagaatg gtgatgatgg tggatgatgca 2520
ggtgatgctg ggaaggctgc tgctgcggtt gctgctgcta gtggggagca gatattaaaa 2580
gcgattgttg atgctgctaa agatggatg atgacggggg ttactgatgt aaaggatgct 2640
acaaatccga ttgacgcggc tattgggggg agtgcggatg ctaatgctga ggcgtttgat 2700
aagatgaaga aggatgatca gattgctgct gctatgggtc tgaggggaat ggctaaggat 2760
gggcagtttg ctttg 2775

```

```

<210> SEQ ID NO 89
<211> LENGTH: 2075
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<400> SEQUENCE: 89

```

```

ataaagggga ttgttgatgc tgctgagaag gctgatgcga aggaagggaa gttggatgtg 60
gctggtgatg ctggtgaaac taacaaggat gctgggaagt tgtttgtgaa gaacaatggt 120
aatgaggggt gtgatgcaga tgatgctggg aaggctgctg ctgcggttgc tgctgttagt 180
ggggagcaga tattaaaagc gattgttgat gctgctaagg gtggtgataa gacgggtaag 240
aataatgtga aggatgctga aaatccgatt gaggcggcta ttgggagtag tgcggatgct 300
gatgctgcgg cgtttaataa ggaggggatg aagaaggatg atcagattgc tgctgctatg 360
gttctgaggg gaattggctaa ggtatggcag ttgtcttga cgaatgatgc tgctgctcat 420
gaagggactg ttaagaatgc tgttgggagt gcaacaaata agaccgttgt tgccttggtc 480
aacttggttc gaaagaccgt gcaagctggg ttgaagaagg ttggggatgt tgtaagaat 540
agtgaaggca aagatggtga tgcggcgagt gttaatggga ttgctaaggg gataaagggg 600
attgttgatg ctgctgagaa ggtgatgctg aaggaaggga agttggatgt ggctggtgct 660
gctggtgaaa ctaacaagga tgctgggaag ttgtttgtga agaagaataa tgaggggtgt 720
gaagcaaatg atgctgggaa ggtgctgctg gcggttgctg ctgttagtgg ggagcagata 780
ttaaaagcga ttgttgatgc tgctaaggat ggtgatgata agcagggtaa gaaggctgag 840
gatgctacaa atccgattga cgcggctatt gggggtgcag gtgcgggtgc taatgctgct 900
gcggcggtta ataatatgaa gaaggatgat cagattgctg ctgctatggt tctgagggga 960
atggctaagg atgggcagtt tgctttgacg aataatgctc atactaatca taaggggact 1020
gttaagaatg ctgttgatat gacaaaagct gctgcggttg ctgcaagtgc tgcaagtgct 1080
gctactggtg atgcagcaat tggggatggt gttaatggta atgatggagc agcaaaagg 1140
ggtgatgcgg cgagtgttaa tgggattgct aaggggataa aggggattgt tgatgctgct 1200
gagaaggctg atgcgaagga agggaagttg aatgtggctg gtgctgctgg tgctgaggg 1260

```

-continued

```

aacgaggctg ctgggaagct gtttgtgaag aagaatgctg gtgatcatgg tggagaagca 1320
ggtgatgctg ggagggctgc tgctgcggtt gctgctgtta gtggggagca gatattaaaa 1380
gcgattgttg atgctgctaa ggatggtggt gataagcagg gtaagaaggc tgaggatgct 1440
gaaaatccga ttgacgcggc tattgggagt acgggtgcgg atgataatgc tgctgaggcg 1500
tttgctacta tgaagaagga tgatcagatt gctgctgcta tggttctgag gggaatggct 1560
aaggatgggc agtttgcttt gaaggatgct gctcatgata atcataaggg gactgttaag 1620
aatgctgttg atataataaa ggctactgcg gttgctgcaa gtgctgctac tggtagtaca 1680
acgattgggg atgttggttaa gaatggtgag gcaaaagggtg gtgaggcgaa gagtgttaat 1740
gggattgcta aggggataaa ggggattggt gatgctgctg gaaaggctga tgcgaaggaa 1800
gggaagttag atgtggctgg tgctgctggt gagggtaacg aggctgctgg gaagctgttt 1860
gtgtaaatta ctataggatt agaactagtg tacgatatga gtcctttggt tattttgcag 1920
ctgctaataa atttgaataa agtgaagtta aaattgcgga tggttaatga acacatttta 1980
ttgctacaaa agagaaagaa atattatatg attcacttga ttttaagggt cgtggaaaaa 2040
tatttgaaat aacttcaaag cgaatgttta agctt 2075

```

```

<210> SEQ ID NO 90
<211> LENGTH: 552
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(552)

```

```

<400> SEQUENCE: 90

```

```

gaa ggg act gtt aag aat gct gtt gat atg gca aaa gct gct gcg gtt 48
Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Val
1 5 10 15

gct gca agt gct gct act ggc aat gca gca att ggg gat gtt gtt aag 96
Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys
20 25 30

agt aat ggt gga gca gca gca aaa ggt ggt gat gcg gcg agt gtt aat 144
Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn
35 40 45

ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct 192
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala
50 55 60

gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct gct ggt gaa act 240
Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr
65 70 75 80

aac aag gat gct ggg aag ttg ttt gtg aag aag aat ggt gat gat ggt 288
Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly
85 90 95

ggt gat gca ggt gat gct ggg aag gct gct gct gcg gtt gct gct gtt 336
Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val
100 105 110

agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aaa gat ggt 384
Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly
115 120 125

gat aag acg ggg gtt act gat gta aag gat gct aca aat ccg att gac 432
Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp
130 135 140

```

-continued

```

gcg gct att ggg ggg agt gcg gat gct aat gct gag gcg ttt gat aag      480
Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys
145                      150                      155                      160

```

```

atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg      528
Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met
                      165                      170                      175

```

```

gct aag gat ggg cag ttt gct ttg      552
Ala Lys Asp Gly Gln Phe Ala Leu
180

```

```

<210> SEQ ID NO 91
<211> LENGTH: 184
<212> TYPE: PRT
<213> ORGANISM: Borrelia garinii

```

```

<400> SEQUENCE: 91

```

```

Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Val
1      5      10      15

```

```

Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys
20      25      30

```

```

Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn
35      40      45

```

```

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala
50      55      60

```

```

Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr
65      70      75      80

```

```

Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly
85      90      95

```

```

Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val
100     105     110

```

```

Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly
115     120     125

```

```

Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp
130     135     140

```

```

Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys
145     150     155     160

```

```

Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met
165     170     175

```

```

Ala Lys Asp Gly Gln Phe Ala Leu
180

```

```

<210> SEQ ID NO 92
<211> LENGTH: 420
<212> TYPE: DNA
<213> ORGANISM: Borrelia garinii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(420)

```

```

<400> SEQUENCE: 92

```

```

ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa ggg      48
Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu Gly
1      5      10      15

```

```

aag ttg gat gtg gct ggt gat gct ggt gaa act aac aag gat gct ggg      96
Lys Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala Gly
20      25      30

```

```

aag ttg ttt gtg aag aac aat ggt aat gag ggt ggt gat gca gat gat      144

```

-continued

Lys	Leu	Phe	Val	Lys	Asn	Asn	Gly	Asn	Glu	Gly	Gly	Asp	Ala	Asp	Asp		
	35						40					45					
gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata 192																	
Ala	Gly	Lys	Ala	Ala	Ala	Ala	Val	Ala	Ala	Val	Ser	Gly	Glu	Gln	Ile		
	50						55				60						
tta aaa gcg att gtt gat gct gct aag ggt ggt gat aag acg ggt aag 240																	
Leu	Lys	Ala	Ile	Val	Asp	Ala	Ala	Lys	Gly	Gly	Asp	Lys	Thr	Gly	Lys		
65					70					75				80			
aat aat gtg aag gat gct gaa aat ccg att gag gcg gct att ggg agt 288																	
Asn	Asn	Val	Lys	Asp	Ala	Glu	Asn	Pro	Ile	Glu	Ala	Ala	Ile	Gly	Ser		
			85					90					95				
agt gcg gat gct gat gct gcg gcg ttt aat aag gag ggg atg aag aag 336																	
Ser	Ala	Asp	Ala	Asp	Ala	Ala	Ala	Phe	Asn	Lys	Glu	Gly	Met	Lys	Lys		
			100					105					110				
gat gat cag att gct gct gct atg gtt ctg agg gga atg gct aag gat 384																	
Asp	Asp	Gln	Ile	Ala	Ala	Ala	Met	Val	Leu	Arg	Gly	Met	Ala	Lys	Asp		
		115					120					125					
ggg cag ttt gct ttg acg aat gat gct gct gct cat 420																	
Gly	Gln	Phe	Ala	Leu	Thr	Asn	Asp	Ala	Ala	Ala	His						
	130					135					140						

<210> SEQ ID NO 93
 <211> LENGTH: 140
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 93

Ile	Lys	Gly	Ile	Val	Asp	Ala	Ala	Glu	Lys	Ala	Asp	Ala	Lys	Glu	Gly		
1			5					10					15				
Lys	Leu	Asp	Val	Ala	Gly	Asp	Ala	Gly	Glu	Thr	Asn	Lys	Asp	Ala	Gly		
	20							25				30					
Lys	Leu	Phe	Val	Lys	Asn	Asn	Gly	Asn	Glu	Gly	Gly	Asp	Ala	Asp	Asp		
	35						40					45					
Ala	Gly	Lys	Ala	Ala	Ala	Ala	Val	Ala	Ala	Val	Ser	Gly	Glu	Gln	Ile		
	50						55				60						
Leu	Lys	Ala	Ile	Val	Asp	Ala	Ala	Lys	Gly	Gly	Asp	Lys	Thr	Gly	Lys		
65					70					75				80			
Asn	Asn	Val	Lys	Asp	Ala	Glu	Asn	Pro	Ile	Glu	Ala	Ala	Ile	Gly	Ser		
			85					90					95				
Ser	Ala	Asp	Ala	Asp	Ala	Ala	Ala	Phe	Asn	Lys	Glu	Gly	Met	Lys	Lys		
	100							105					110				
Asp	Asp	Gln	Ile	Ala	Ala	Ala	Met	Val	Leu	Arg	Gly	Met	Ala	Lys	Asp		
	115						120					125					
Gly	Gln	Phe	Ala	Leu	Thr	Asn	Asp	Ala	Ala	Ala	His						
	130					135					140						

<210> SEQ ID NO 94
 <211> LENGTH: 942
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia garinii

<400> SEQUENCE: 94

atgagaggat	cgcacaccca	tcaccatcac	ggatccaagg	ggactgttaa	gaatgctgtt		60
gatatgacaa	aagctgctgc	gggtgctgca	agtgctgcaa	gtgctgctac	tggtaatgca		120
gcaattgggg	atgttggttaa	tggtaatgat	ggagcagcaa	aagtggtga	tcggcgagat		180

-continued

```

gttaatggga ttgctaaggg gataaagggg attgttgatg ctgctgagaa ggctgatgcg 240
aaggaagggga agttgaatgt ggctggtgct gctggtgctg agggtaacga ggctgctggg 300
aagctgtttg tgaagaagaa tgctggtgat catggtggtg aagcaggtga tgctgggagg 360
gctgctgctg cggttgctgc tgttagtggg gagcagatat taaaagcgat tgttgatgct 420
gctaaggatg gtggtgataa gcagggtaag aaggctgagg atgctgaaaa tccgattgac 480
gcggtctattg ggagtacggg tgcggtgat aatgctgctg aggcgtttgc tactatgaag 540
aaggatgatc agattgctgc tgctatggtt ctgaggggaa tggctaagga tgggcagttt 600
gctttgaagg atgctgctca tgataatcat ctgcagccaa gcttaattag ctgagcttgg 660
actcctgttg atagatccag taatgacctc agaactccat ctggatttgt tcagaacgct 720
cggttgccgc cgggcgtttt ttattggtga gaatccaagc tagcttggcg agattttcag 780
gagctaagga agctaaaatg gagaaaaaat cactggatat accaccgttg atatatccca 840
atggcatcgt aaagaacatt ttgaggcatt tcagtcagtt gctcaatgta cctataacca 900
gaccgttcag ctggatatta cggccttttt aaagaccgta ag 942

```

<210> SEQ ID NO 95

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: *Borrelia garinii*

<400> SEQUENCE: 95

```

Met Arg Gly Ser His His His His His Gly Ser Lys Gly Thr Val
1           5           10          15
Lys Asn Ala Val Asp Met Thr Lys Ala Ala Ala Val Ala Ala Ser Ala
20          25          30
Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Asn Gly
35          40          45
Asn Asp Gly Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn Gly Ile
50          55          60
Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala
65          70          75          80
Lys Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly Ala Glu Gly Asn
85          90          95
Glu Ala Ala Gly Lys Leu Phe Val Lys Lys Asn Ala Gly Asp His Gly
100         105         110
Gly Glu Ala Gly Asp Ala Gly Arg Ala Ala Ala Val Ala Ala Val
115         120         125
Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly
130         135         140
Gly Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Glu Asn Pro Ile Asp
145         150         155         160
Ala Ala Ile Gly Ser Thr Gly Ala Asp Asp Asn Ala Ala Glu Ala Phe
165         170         175
Ala Thr Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg
180         185         190
Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp Ala Ala His Asp
195         200         205
Asn His Leu Gln Pro Ser Leu Ile Ser
210         215

```

-continued

<210> SEQ ID NO 96
 <211> LENGTH: 663
 <212> TYPE: DNA
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 96

```

atgagaggat cgcataacca tcaccatcac ggatccaaga gtgctgtgga tgaggctagc      60
aagtggtagg aagagatgat aacagctgct ggtgaggctg ctacaaaggg tggtagtggt      120
gaagctagcg aaaagattgg ggatgttggt gataataatc atggtgctgt agctgatgcg      180
gacagtgtta aggggattgc gaaggggata aaggggattg ttgatgctgc tgggaaggct      240
tttgtaagg atggtgctgc gaaggatgtt gcagctgctg ctggtgatga ggctaacaag      300
gatgcgggga agttgtttgc tggtcaggat ggtggtggtg ctgatggtga cattgcgaag      360
gcggctgctg ctgttactgc ggttagtggg gagcagatac tgaaagctat tgttgaggct      420
gctggtgata aggctaatac ggtgggtgta aaggctgctg gtgcggctac gaatccgatt      480
gcagctgcga ttgggactga tgatgataat gcggcggcgt ttgataagga tgagatgaag      540
aagagtaatg ataagattgc tcgggctatt gttttgaggg ggggtggctaa ggatggaaag      600
tttctgctga atgctaatag taatagtaag gcgagtgtgc tgcagccaag cttaattagc      660
tga                                                                                   663

```

<210> SEQ ID NO 97
 <211> LENGTH: 220
 <212> TYPE: PRT
 <213> ORGANISM: Borrelia afzelii

<400> SEQUENCE: 97

```

Met Arg Gly Ser His His His His His Gly Ser Lys Ser Ala Val
1           5           10          15
Asp Glu Ala Ser Lys Trp Leu Glu Glu Met Ile Thr Ala Ala Gly Glu
20          25          30
Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu Lys Ile Gly Asp
35          40          45
Val Gly Asp Asn Asn His Gly Ala Val Ala Asp Ala Asp Ser Val Lys
50          55          60
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala
65          70          75          80
Phe Gly Lys Asp Gly Ala Leu Lys Asp Val Ala Ala Ala Ala Gly Asp
85          90          95
Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Gln Asp Gly Gly
100         105         110
Gly Ala Asp Gly Asp Ile Ala Lys Ala Ala Ala Val Thr Ala Val
115         120         125
Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala Ala Gly Asp Lys
130         135         140
Ala Asn Gln Val Gly Val Lys Ala Ala Gly Ala Ala Thr Asn Pro Ile
145         150         155         160
Ala Ala Ala Ile Gly Thr Asp Asp Asp Asn Ala Ala Ala Phe Asp Lys
165         170         175
Asp Glu Met Lys Lys Ser Asn Asp Lys Ile Ala Ala Ala Ile Val Leu
180         185         190

```

-continued

Arg	Gly	Val	Ala	Lys	Asp	Gly	Lys	Phe	Ala	Ala	Asn	Ala	Asn	Asp	Asn
		195					200					205			

Ser	Lys	Ala	Ser	Val	Leu	Gln	Pro	Ser	Leu	Ile	Ser
	210				215					220	

<210> SEQ ID NO 98
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 98

cggaattcac tcgccttact attatc 26

<210> SEQ ID NO 99
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 99

cgggatccga gagtgcgtgtt gatgaggtt 29

<210> SEQ ID NO 100
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 100

cgggatccaa gagtgcgtgtg gatgaggcta gcaag 35

<210> SEQ ID NO 101
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 101

ttctgcagca cactcgccctt actattatca ttagc 35

<210> SEQ ID NO 102
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 102

cgggatccgc tggtgggagt ygcaac 26

<210> SEQ ID NO 103
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

<400> SEQUENCE: 103

aactgcagat tatcatgagc agcatccttc                               30

<210> SEQ ID NO 104
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

<400> SEQUENCE: 104

cgggatccaa ggggactgtt aagaatgctg ttg                             33

<210> SEQ ID NO 105
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

<400> SEQUENCE: 105

ttctgcagat gattatcatg agcagcatcc ttca                             34

<210> SEQ ID NO 106
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Borrelia burgdorferi

<400> SEQUENCE: 106

tgagggggct attaagg                                                17

<210> SEQ ID NO 107
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

<400> SEQUENCE: 107

ccggaattcc gg                                                       12

```

1. An recombinant nucleic acid comprising (i) a nucleotide sequence operably linked to a heterologous promoter, wherein said nucleotide sequence encodes at least 12 contiguous amino acids of SEQ ID NO:32.

2. The recombinant nucleic acid of claim 1, wherein the nucleotide sequence encodes at least 15 contiguous amino acids of SEQ ID NO:32.

3. The recombinant nucleic acid of claim 1, wherein the nucleotide sequence encodes at least 20 contiguous amino acids of SEQ ID NO:32.

4. The recombinant nucleic acid of claim 1, wherein the nucleotide sequence encodes a peptide comprising SEQ ID NO:32.

5. The recombinant nucleic acid of claim 1, wherein the nucleic acid is an RNA segment.

6-20. (canceled)

21. A method of using an isolated nucleic acid that encodes at least 12 contiguous amino acids of SEQ ID NO:32 or 50 contiguous amino acids of SEQ ID NO:39, comprising the steps of: (a) preparing a recombinant vector in which the isolated nucleic acid is positioned under the control of a promoter; (b) introducing said recombinant vector into a host cell; (c) culturing said host cell under conditions effective to allow expression of the polypeptide; and (d) collecting said expressed polypeptide.

22. A method of assaying for *Borrelia* infection comprising: (a) contacting a sample obtained from a subject with an isolated polypeptide, said isolated polypeptide being immobilized on a surface and comprising: (i) at least 15 contiguous amino acids of SEQ ID NO:32; or (ii) at least 50

contiguous amino acids of SEQ ID NO:39; and (b) determining whether immunologic binding occurs between the isolated polypeptide and an antibody in the sample, wherein immunologic binding is indicative of *Borrelia* infection.

23. The method of claim 22, wherein the isolated polypeptide sequence comprises at least 15 contiguous amino acids of SEQ ID NO:32.

24. The method of claim 22, wherein the isolated polypeptide sequence comprises at least 50 contiguous amino acids of SEQ ID NO:39.

25. The method of claim 22, wherein the isolated polypeptide comprises a label.

26. The method of claim 22, wherein the isolated polypeptide is a fusion protein.

27. The method of claim 22, further defined as comprising performing an ELISA assay.

28. A recombinant nucleic acid comprising (i) a nucleotide sequence operably linked to a heterologous promoter, wherein said nucleotide sequence encodes at least 50 contiguous amino acids of SEQ ID NO:39.

29. The recombinant nucleic acid of claim 1, wherein the nucleotide sequence encodes at least 75 contiguous amino acids of SEQ ID NO:39.

30. The recombinant nucleic acid of claim 1, wherein the nucleotide sequence encodes at least 100 contiguous amino acids of SEQ ID NO:39.

31. The recombinant nucleic acid of claim 1, wherein the nucleotide sequence encodes a peptide comprising SEQ ID NO:39.

32. The recombinant nucleic acid of claim 28, wherein the nucleic acid is an RNA segment.

* * * * *