

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(10) International Publication Number

WO 2018/187423 A1

(43) International Publication Date
11 October 2018 (11.10.2018)(51) International Patent Classification:
G01N 33/68 (2006.01) *C12Q 1/533* (2006.01)
G01N 33/566 (2006.01) *A61K 47/42* (2017.01)
(21) International Application Number:
PCT/US2018/026014(22) International Filing Date:
04 April 2018 (04.04.2018)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/482,018 05 April 2017 (05.04.2017) US

(71) Applicant: WARP DRIVE BIO, INC. [US/US]; 400 Technology Square, Cambridge, MA 02139 (US).

(72) Inventors: MULVIHILL, Mark Joseph; 4 Webster Circle, Sudbury, MA 01776 (US). JIN, Meizhong; 34 Priscilla Circle, Wellesley, MA 02481 (US). PERL, Nicholas; 7 Gibbens St, Somerville, MA 02143 (US).

(74) Agent: BELLIVEAU, Michael, J.; Clark & Elbing LLP, 101 Federal Street, 15th Floor, Boston, MA 02110 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:
— of inventorship (Rule 4.17(iv))

(54) Title: METHODS AND REAGENTS FOR ANALYZING PROTEIN-PROTEIN INTERFACES

FIG. 9A



(57) Abstract: The present disclosure provides methods and reagents useful for analyzing protein-protein interfaces such as interfaces between a presenter protein (e.g., a member of the FKBP family, a member of the cyclophilin family, or PIN1) and a target protein. In some embodiments, the target and/or presenter proteins are intracellular proteins. In some embodiments, the target and/or presenter proteins are mammalian proteins.

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

METHODS AND REAGENTS FOR ANALYZING PROTEIN-PROTEIN INTERFACES**Background**

The vast majority of small molecule drugs act by binding a functionally important pocket on a target protein, thereby modulating the activity of that protein. For example, the cholesterol-lowering drugs statins bind the enzyme active site of HMG-CoA reductase, thus preventing the enzyme from engaging with its substrates. The fact that many such drug/target interacting pairs are known may have misled some into believing that a small molecule modulator could be discovered for most, if not all, proteins provided a reasonable amount of time, effort, and resources. This is far from the case. Current estimates hold that only about 10% of all human proteins are targetable by small molecules. The other 90% are currently considered refractory or intractable toward small molecule drug discovery. Such targets are commonly referred to as “undruggable.” These undruggable targets include a vast and largely untapped reservoir of medically important human proteins. Thus, there exists a great deal of interest in discovering new molecular modalities capable of modulating the function of such undruggable targets.

15

Summary

Small molecules are limited in their targeting ability because their interactions with the target are driven by adhesive forces, the strength of which is roughly proportional to contact surface area. Because of their small size, the only way for a small molecule to build up enough intermolecular contact surface area to effectively interact with a target protein is to be literally engulfed by that protein. Indeed, a large body of both experimental and computational data supports the view that only those proteins having a hydrophobic “pocket” on their surface are capable of binding small molecules. In those cases, binding is enabled by engulfment.

Nature has evolved a strategy that allows a small molecule to interact with target proteins at sites other than hydrophobic pockets. This strategy is exemplified by naturally occurring immunosuppressive drugs cyclosporine A, rapamycin, and FK506. The biological activity of these drugs involves the formation of a high-affinity complex of the small molecule with a small presenting protein. The composite surface of the small molecule and the presenting protein engages the target. Thus, for example, the binary complex formed between cyclosporin A and cyclophilin A targets calcineurin with high affinity and specificity, but neither cyclosporin A or cyclophilin A alone binds calcineurin with measurable affinity.

The present inventors have developed compounds and conjugates useful for identifying presenter protein and target protein pairs, and probing the interfaces between them for use in the development of small molecules capable of modulating these interactions.


Accordingly, the present disclosure provides methods and reagents useful for analyzing protein-protein interfaces such as interfaces between a presenter protein (e.g., a member of the FKBP family, a member of the cyclophilin family, or PIN1) and a target protein. Such analysis is useful in aiding the design of small molecules that are capable of binding simultaneously to both a presenter protein and a target protein, such that the resulting small molecule-presenter protein complexes can bind to and modulate the activity of the target protein. In some embodiments, the target and/or presenter proteins are intracellular proteins. In some embodiments, the target and/or presenter proteins are mammalian proteins.

In some aspects, the disclosure provides compounds that may be used as cross-linking substrates. These compounds may include a protein binding moiety capable of covalent or non-covalent binding to a protein (e.g., a target protein or a presenter protein) and at least one cross-linking group capable of a chemoselective reaction with an amino acid of a different protein than that which binds to the protein binding moiety. In some embodiments, the compounds include only one cross-linking group.

Accordingly, in an aspect, the disclosure provides a compound including a protein binding moiety (e.g., a presenter protein binding moiety or a target protein binding moiety) and a cross-linking group (e.g., a moiety capable of a chemoselective reaction with an amino acid of a different protein than that which binds to the protein binding moiety). The protein binding moiety is capable of binding (covalently or non-covalently) to a protein (e.g., a presenter protein or target protein, depending upon whether it is a presenter protein binding moiety or a target protein binding moiety), while the cross-linking group is capable of forming a covalent bond with a protein (e.g., a presenter protein, a target protein, or another compound that is capable of binding such other protein). In some embodiments, when the compound includes a presenter protein binding moiety, the compound does not include a target protein binding moiety. In some embodiments, when the compound includes a target protein binding moiety, the compound does not include a presenter protein binding moiety.

In some embodiments, the cross-linking group is a sulfhydryl-reactive cross-linking group (e.g., the cross-linking group includes a mixed disulfide, a maleimide, vinyl sulfone, vinyl ketone, or an alkyl halide), an amino-reactive cross-linking group, a carboxyl-reactive cross-linking group, a carbonyl-reactive cross-linking group, or a triazole-forming cross-linking group.

In some embodiments, the cross-linking group includes a mixed disulfide, e.g., the cross-linking group includes the structure of Formula Ia:




Formula Ia

wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound; and

R^A is 0, 1, or 2;


R^A is optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_6 - C_{10} aryl, or optionally substituted C_2 - C_9 heteroaryl.

In some embodiments, R^A is optionally substituted C_2 - C_9 heteroaryl (e.g., pyridyl). In some embodiments, the cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.


In some embodiments, R^A is optionally substituted C_1 - C_6 heteroalkyl (e.g., N, N-dimethylethyl). In some embodiments, the cross-linking group includes the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound. In some embodiments, the cross-linking group includes the structure:




5 In some embodiments, R^A is optionally substituted C₁-C₆ alkyl (e.g., methyl). In some embodiments, the cross-linking group includes the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

10 In some embodiments, the cross-linking group includes a carbon-based cross-linking group (e.g., a cross-linking group that forms a carbon-sulfide bond upon reaction with a thiol).

In some embodiments, the cross-linking group includes a maleimide, e.g., the cross-linking group includes the structure of formula Ib, Ic, Id, or Ie:



Formula Ib

Formula Ic

Formula Id

Formula Ie

15 wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^A is -C(O)- or -SO₂-;

X^B is -C(O)- or CR^ER^F;

20 R^B and R^C are, independently, hydrogen, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl;

25 RD is hydrogen, hydroxyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and


30 RE and RF are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heterocyclyl

35 and optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl;

heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.


In some embodiments, the cross-linking group includes the structure of Formula Ib. In some embodiments, X^A is -C(O)-. In some embodiments, X^B is -C(O)-. In some embodiments, R^B and R^C are 5 hydrogen or optionally substituted C₁-C₆ alkyl (e.g., methyl).

In some embodiments, the cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

10 In some embodiments, the cross-linking group includes the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

In some embodiments, the cross-linking group includes a structure of formula If, Ig, Ih, or II:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^C is -C(O)- or -SO₂-;


20 X^D is absent, NR^JR^K, or OR^L;

R^G, R^H, and R^I are, independently, hydrogen, nitrile, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally 25 substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and

30 R^J, R^K, and R^L are, independently, absent, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.


In some embodiments, the cross-linking group includes the structure of Formula I^f. In some embodiments, X^D is absent. In some embodiments, R^G, R^H, and R^I are hydrogen. In some embodiments, X^C is $-\text{C}(\text{O})-$. In some embodiments, X^C is $-\text{SO}_2-$.

5 In some embodiments, the cross-linking group includes a vinyl sulfone, e.g., the cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

10 In some embodiments, the cross-linking group includes a vinyl sulfone, e.g., the cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

15 In some embodiments, the cross-linking group includes a vinyl ketone, e.g., the cross-linking group includes the structures:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

20 In some embodiments, the cross-linking group includes a vinyl ketone, e.g., the cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

25 In some embodiments, the cross-linking group includes a vinyl ketone, e.g., the cross-linking group includes the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

30 In some embodiments, the cross-linking group includes an ynone such as a structure of formula I^j or I^k:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound:


5 X^E is absent, NR^NR^O , or OR^P ;

R^M is hydrogen, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl; and

10 substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and


R^N , R^O , and R^P are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

In some embodiments, the cross-linking group includes a vinyl ketone, e.g., the cross-linking group includes the structure:



20 H 
wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

In some embodiments, the cross-linking group includes a structure of formula I_m or I_n:



25 Formula 1m

Formula In

wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound:

X^F is absent, NR^SRT , or OR^U :

X^G is absent or -C(O)-;


30 Y is a leaving group:

R^Q and R^R are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉


heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and

R^S, R^T, and R^U are, independently, absent, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

In some embodiments, Y is a halogen (e.g., fluoro, chloro, bromo, or iodo), a mesylate, a tosylate, or a triflate. In some embodiments, Y is a nitrile. In some embodiments, X^F and X^G are absent. In some embodiments, R^Q and R^R are hydrogen. In some embodiments, the cross-linking group includes an alkyl halide such as an alkyl chloride, e.g., the cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound. In some embodiments, the cross-linking group includes an alkyl halide such as an alkyl chloride or alkyl fluoride, e.g., the cross-linking group includes the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

In some embodiments, the cross-linking group includes an epoxide, e.g., the cross-linking group includes a structure of formula I_o:




Formula I_o

wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

R^V, R^W, and R^X are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

In some embodiments, the cross-linking group includes a structure of formula I_p:



Formula I(p)

wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

5 the dotted lines represent optional double bonds included as necessary for the structure to be aromatic;

b is 0, 1, or 2;

Y is a leaving group;

R^Y and R^Z are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol,

10 optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl

15 C₁-C₆ alkyl;

each of X^H, X^I, X^J, X^K, and X^L are, independently, absent, NR^{AA}, or CR^{AB}, wherein at least five of X^H, X^I, X^J, X^K, and X^L are NR^{AA}, or CR^{AB};

R^{AA} is absent or hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally


20 substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and

25 R^{AB} is hydrogen, nitrile, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

30 In some embodiments, at least one R^{AB} is an electron withdrawing group. In some embodiments, one to three R^{AB} are electron withdrawing groups.


In some embodiments, Y is a nitrile. In some embodiments, Y is a halogen (e.g., fluoro, chloro, bromo, or iodo), a mesylate, a tosylate, or a triflate.

In some embodiments, the cross-linking group includes the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.


In some embodiments, the cross-linking group includes the structure:



5


wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

In some embodiments, the cross-linking group includes the structure:



10 wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

In some embodiments, the cross-linking group is an internal cross-linking group, e.g., the cross-linking group includes a structure of formula Iq, Ir, or Is:



15

Formula Iq

Formula Ir

Formula Is

wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^M is $-\text{C}(\text{O})-$ or $-\text{SO}_2-$;

X^N is absent, NR^{AE} , or O ;

20 R^{AC} and R^{AD} are, independently, hydrogen, nitrile, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted $\text{C}_1\text{-}\text{C}_6$ alkyl, optionally substituted $\text{C}_2\text{-}\text{C}_6$ alkenyl, optionally substituted $\text{C}_2\text{-}\text{C}_6$ alkynyl, optionally substituted $\text{C}_3\text{-}\text{C}_{10}$ carbocyclyl, optionally substituted $\text{C}_6\text{-}\text{C}_{10}$ aryl, optionally substituted $\text{C}_6\text{-}\text{C}_{10}$ aryl $\text{C}_1\text{-}\text{C}_6$ alkyl, optionally substituted $\text{C}_2\text{-}\text{C}_9$ heteroaryl, optionally substituted $\text{C}_2\text{-}\text{C}_9$ heteroaryl $\text{C}_1\text{-}\text{C}_6$ alkyl, optionally substituted $\text{C}_2\text{-}\text{C}_9$ heterocyclyl, or optionally substituted $\text{C}_2\text{-}\text{C}_9$ heterocyclyl $\text{C}_1\text{-}\text{C}_6$ alkyl; and


5 R^{AE} is hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.


In some embodiments, X^N is NR^{AE} , wherein R^{AE} is hydrogen. In some embodiments, R^{AC} and R^{AD} are hydrogen. In some embodiments, X^M is $-C(O)-$. In some embodiments, X^M is $-SO_2-$.


10 In some embodiments of any of the foregoing compounds, the protein binding moiety portion is capable of non-covalent interaction with a protein. In some embodiments of any of the foregoing compounds, the protein binding moiety portion is capable of covalent interaction with a protein.

In some aspects, the disclosure provides a compound including a presenter protein binding moiety and a cross-linking group. In some embodiments, the protein binding moiety and the cross-linking group are attached through a linker.

15 In some aspects, the disclosure provides a compound having the structure:







In some aspects, the disclosure provides conjugates, methods for their synthesis, and uses thereof, including a presenter protein binding moiety capable of covalent or non-covalent binding to a 5 presenter protein conjugated to a target protein through a linker.

Accordingly, in another aspect, the disclosure provides a conjugate including a presenter protein binding moiety conjugated to a target protein. In some embodiments, the presenter protein binding moiety portion of the conjugate is capable of non-covalent interaction with a presenter protein. In some 10 embodiments, the presenter protein binding moiety portion of the conjugate is capable of covalent interaction with a presenter protein.

In some aspects, the disclosure provides a method of producing a conjugate including a presenter protein binding moiety conjugated to a target protein. This method includes reacting (a) a compound including a presenter protein binding moiety and a cross-linking group with (b) a target protein under conditions that permit production of the conjugate.

15 In some aspects, the disclosure provides a method of producing a conjugate including a presenter protein binding moiety conjugated to a target protein. This method includes providing (a) a compound including a presenter protein binding moiety and a cross-linking group; (b) a target protein; and (c) a presenter protein; and reacting the compound with the target protein under conditions that permit production of the conjugate.

20 In some aspects, the disclosure provides complexes, methods for their production, and uses thereof, including a presenter protein and a conjugate including a presenter protein binding moiety and a target protein.

Accordingly, in another aspect, the disclosure provides a complex including (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein.

25 In some aspects, the disclosure provides a method of producing a complex including (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter

protein. This method includes combining a conjugate including a presenter protein binding moiety conjugated to a target protein and a presenter protein under conditions that permit production of the complex.

5 In some aspects, the disclosure provides a method of producing a complex including (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein. This method includes providing (a) a compound including a presenter protein binding moiety and a cross-linking group; (b) a target protein; and (c) a presenter protein; and reacting the compound with the target protein under conditions that permit production of the complex.

10 In some embodiments of the foregoing methods, the presenter protein binds to the compound in the absence of the target protein. In some embodiments of the foregoing methods, the presenter protein does not substantially bind to the compound in the absence of the target protein. In some embodiments of the foregoing methods, the compound and the target protein do not substantially react in the absence of the presenter protein. In some embodiments of the foregoing methods, the compound and the target protein react in the absence of the presenter protein. In some embodiments of the foregoing methods, 15 the conditions do not include a reducing reagent. In some embodiments of the foregoing methods, the conditions include an excess of presenter protein.

15 In some embodiments, detectable binding between the compound and the presenter protein is observed in the absence of the target protein. In some embodiments, however detectable binding between the compound and the presenter protein is not observed (e.g., the presenter protein does not 20 substantially bind to the compound) in the absence of the target protein. In some embodiments, significant reaction between the cross-linking group and the target protein (e.g., significant conjugate formation) is not observed in the absence of the presenter protein. In some embodiments, however, significant reaction between the cross-linking group and the target protein may be observed even in the absence of the presenter protein. In some embodiments, rate and/or extent of such reaction (e.g., rate 25 and/or amount of conjugate formation) may differ in a given assay when presenter protein is present as compared with when it is absent (e.g., the rate and/or amount of conjugate formation is 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, or 100-fold greater in the presence of the presenter protein).

In some embodiments, conjugate production as described herein is performed under conditions that do not include (e.g., are substantially free of) a reducing reagent.

30 In some embodiments, the present invention provides a complex comprising (i) a presenter protein; (ii) a compound as described herein (e.g., compound whose structure includes a presenter protein binding moiety and a cross-linking group); and (iii) a target protein. In some embodiments, such complex is exposed to and/or maintained under conditions that permit reaction of the cross-linking moiety with the target protein, so that a cross-link therebetween is formed. In some embodiments, the cross-link 35 is with a heteroatom in an amino acid (e.g., in an amino acid side chain) of the target protein. In some embodiments, the cross-link is with an –S- atom in a cysteine in the target protein. In some embodiments, the target protein is a variant of a natural target protein; in some such embodiments, the variant has an amino acid sequence that shows a high degree (e.g., 80%, 81%, 82%; 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, 99% or higher) with the natural 40 target protein but differs by substitution or addition of at least one amino acid susceptible to participation

in a cross-link with the cross-linking group (e.g., whose amino acid side chain includes a heteroatom that can participate in such a cross-link).

5 In some aspects, the disclosure provides conjugates, methods for their synthesis, and uses thereof, including a target protein binding moiety capable of covalent or non-covalent binding to a target protein conjugated to a presenter protein through a linker.

10 Accordingly, in another aspect, the disclosure provides a conjugate including a target protein binding moiety conjugated to a presenter protein. In some embodiments, the target protein binding moiety portion of the conjugate is capable of non-covalent interaction with a target protein. In some embodiments, the target protein binding moiety portion of the conjugate is capable of non-covalent interaction with target protein. In some embodiments, the target protein binding moiety and the presenter protein are conjugated through a linker.

15 In some aspects, the disclosure provides a method of producing a conjugate including a target protein binding moiety conjugated to a presenter protein. This method includes reacting (a) a compound including a target protein binding moiety and a cross-linking group with (b) a presenter protein under conditions that permit production of the conjugate.

20 In some aspects, the disclosure provides a method of producing a conjugate including a target protein binding moiety conjugated to a presenter protein. This method includes providing (a) a compound including a target protein binding moiety and a cross-linking group; (b) a presenter protein; and (c) a target protein; and reacting the compound with the presenter protein under conditions that permit production of the conjugate.

25 In some embodiments, detectable binding between the compound and the target protein is observed absence the presenter protein. In some embodiments, however detectable binding between the compound and the target protein is not observed (e.g., the presenter protein does not substantially bind to the compound) in the absence of the presenter protein. In some embodiments, significant reaction between the cross-linking group and the presenter protein (e.g., significant conjugate formation) is not observed in the absence of the target protein. In some embodiments, however, significant reaction between the cross-linking group and the presenter protein may be observed even in the absence of the target protein. In some embodiments, the rate and/or extent of such reaction (e.g., the rate and/or amount of conjugate formation) may differ in a given assay when presenter protein is present as compared with when it is absent (e.g., the rate and/or amount of conjugate formation is 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 100-fold greater in the presence of the presenter protein).

30 In some embodiments, the target protein binds to the compound in the absence of the presenter protein. In some embodiments, the target protein does not substantially bind to the compound in the absence of the presenter protein. In some embodiments, the presenter protein does not substantially bind to the compound in the absence of the target protein. In some embodiments, reaction between the cross-linking group and the target protein (e.g., conjugate formation) is not observed in the absence of the presenter protein. In some embodiments, however, reaction between the cross-linking group and the target protein is observed even in the absence of the presenter protein. In some embodiments, conjugate production as described herein is performed under conditions that do not include (e.g., are substantially free of) a reducing agent.

In some embodiments, the present invention provides a complex comprising (i) a presenter protein; (ii) a compound as described herein (e.g., compound whose structure includes a presenter protein binding moiety and a cross-linking group); and (iii) a target protein. In some embodiments, such complex is exposed to and/or maintained under conditions that permit reaction of the cross-linking moiety with the target protein, so that a cross-link therebetween is formed. In some embodiments, the cross-link is with a heteroatom in an amino acid (e.g., in an amino acid side chain) of the target protein. In some embodiments, the cross-link is with an –S- atom in a cysteine in the target protein. In some embodiments, the target protein is a variant of a natural target protein; in some such embodiments, the variant has an amino acid sequence that shows a high degree (e.g., 80%, 81%, 82%; 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, 99% or higher) with the natural target protein but differs by substitution or addition of at least one amino acid susceptible to participation in a cross-link with the cross-linking group (e.g., whose amino acid side chain includes a heteroatom that can participate in such a cross-link).

In some aspects, the disclosure provides complexes, methods for their production, and uses thereof, including a target protein and a conjugate including a target protein binding moiety conjugated to a presenter protein through a linker.

In some aspects, the disclosure provides a complex including (i) a conjugate including a target protein binding moiety conjugated to a presenter protein; (ii) a target protein; and (iii) a presenter protein. In some embodiments, such complex is exposed to and/or maintained under conditions that permit reaction of the cross-linking moiety with the presenter protein, so that a cross-link therebetween is formed. In some embodiments, the cross-link is with a heteroatom in an amino acid (e.g., in an amino acid side chain) of the presenter protein. In some embodiments, the cross-link is with an –S- atom in a cysteine in the presenter protein. In some embodiments, the presenter protein is a variant of a natural presenter protein; in some such embodiments, the variant has an amino acid sequence that shows a high degree (e.g., 80%, 81%, 82%; 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, 99% or higher) with the natural presenter protein but differs by substitution or addition of at least one amino acid susceptible to participation in a cross-link with the cross-linking group (e.g., whose amino acid side chain includes a heteroatom that can participate in such a cross-link).

In some aspects, the disclosure provides a method of producing a complex including (i) a conjugate including a target protein binding moiety conjugated to a presenter protein and (ii) a target protein. This method includes combining a conjugate including a target protein binding moiety conjugated to a presenter protein and a target protein under conditions that permit production of the complex.

In some aspects, the invention features a method of producing a complex including (i) a conjugate as described herein (e.g., a conjugate including a target protein binding moiety and a presenter protein) and (ii) a target protein. In some such embodiments, a provided method includes combining the conjugate and target protein under conditions that permit production of the complex. Alternatively or additionally, in some embodiments, such a method includes, for example, (i) combining (a) a compound (e.g., a compound whose structure includes a target protein binding moiety and a cross-linking group); (b) a target protein; and (c) a presenter protein with one another; and (ii) exposing the combination to and/or maintaining the combination under conditions that permit production of the complex. In some such

embodiments, the conditions permit reaction of the cross-linking group with the presenter protein so that a conjugate is produced.

In some aspects, the disclosure provides a method of producing a complex including (i) a conjugate including a target protein binding moiety conjugated to a presenter protein and (ii) a target protein. This method includes providing (a) a compound including a target protein binding moiety and a cross-linking group; (b) a presenter protein; and (c) a target protein; and reacting the compound with the presenter protein under conditions that permit production of the complex.

In some such embodiments, the conditions are such that the compound, presenter protein, and/or target protein are characterized in that detectable binding between the compound and the target protein is observed in the absence of the presenter protein. In some embodiments, however, detectable binding between the compound and the target protein is not observed (e.g., the target protein does not substantially bind to the compound) under the conditions in the absence of the presenter protein. In some embodiments, significant reaction between the cross-linking group and the presenter protein is not observed in the absence of the target protein under the conditions. In some embodiments, however, significant reaction between the cross-linking group and the presenter protein may be observed even in the absence of the target protein under the conditions. In some embodiments, the conditions do not include a reducing reagent. In some embodiments, the conditions include an excess of presenter protein.

In some embodiments, the target protein binds to the compound in the absence of the presenter protein. In some embodiments, the target protein does not substantially bind to the compound in the absence of the presenter protein. In some embodiments, the compound and the presenter protein do not substantially react in the absence of the target protein. In some embodiments, the compound and the presenter protein react in the absence of the target protein. In some embodiments, the conditions do not include a reducing reagent. In some embodiments, the conditions include an excess of target protein.


In some aspects, the disclosure provides compounds including a presenter protein binding moiety capable on non-covalent interaction with a presenter protein and a target protein binding moiety capable of covalent or non-covalent interaction with a target protein. In some embodiments, the presenter protein binding moiety and the target protein binding moiety are attached via a linker.

Accordingly, in some aspects, the disclosure provides a compound having the structure of Formula VII:

30 A-L-B

Formula VII

wherein A includes the structure of Formula VIIa or VIIb:



35 wherein b and c are independently 0, 1, or 2;
d is 0, 1, 2, 3, 4, 5, 6, or 7;

X¹ and X² are each, independently, absent, CH₂, O, S, SO₂, or NR¹³;

each R¹ and R² are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl (e.g., optionally substituted C₂-C₉ heteroaryl), optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl (e.g., optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl), or R¹ and R² combine with the carbon atom to which they are bound to form C=O or R¹ and R² combine to form an optionally substituted C₃-C₁₀ carbocyclyl or 10 optionally substituted C₂-C₉ heterocyclyl;

each R³ is, independently, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl (e.g., optionally substituted C₂-C₉ heteroaryl), or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl (e.g., optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl) or two R⁸ combine to form an optionally substituted C₃-C₁₀ carbocyclyl, 15 optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heterocyclyl, e.g., optionally substituted C₂-C₉ heteroaryl;

20 R⁴ is optionally substituted C₁-C₆ alkyl;

L is an optional linker; and

B is a target protein binding moiety.

In some embodiments of a compound of Formula VII, the target protein binding moiety, B, is capable of non-covalent interaction with a target protein. In some embodiments of a compound of 25 Formula VII, the target protein binding moiety, B, is capable of covalent interaction with a target protein. In some embodiments of a compound of Formula VII, the linker, L, is present. In some embodiments of a compound of Formula VII, the linker, L, is absent.

In some aspects, the disclosure provides ternary complexes, methods for their production, and uses thereof, including a presenter protein, a target protein, and a compound including a presenter 30 protein binding moiety and a target protein binding moiety.

Accordingly, in another aspect, the disclosure provides a complex including (i) a compound of Formula VII; (ii) a target protein; and (iii) a presenter protein.

In some embodiments, the compounds, conjugates, and complexes of the present invention may be useful for the identification of conjugates including a presenter protein binding moiety and a target 35 protein that are capable of forming complexes with presenter proteins.

In some aspects, the invention features a method of identifying and/or characterizing a conjugate as described herein (e.g., in which a compound whose structure includes a presenter protein binding moiety and a cross-linking group, is conjugated to a target protein) that is capable of forming a complex with a presenter protein. In some embodiments, such a method includes steps of: (a) providing (i) such 40 a conjugate (e.g., in which a compound whose structure includes a presenter protein binding moiety and a cross-linking group, conjugated to a target protein) and (ii) a presenter protein; (b) combining the

conjugate and the presenter protein under conditions suitable to permit complex formation if the conjugate is capable of forming a complex with the presenter protein; and (c) determining whether a complex comprising the conjugate and the presenter protein is formed, wherein formation of the complex indicates that the conjugate is one that is capable of forming a complex with a presenter protein.

5 Accordingly, in some aspects, the disclosure provides a method of identifying and/or characterizing a conjugate that is capable of forming a complex with a presenter protein. This method includes the steps of: (a) providing (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein; (b) combining the conjugate and the presenter protein under conditions suitable to permit complex formation if the conjugate is capable of forming a complex 10 with the presenter protein; and (c) determining whether a complex comprising the conjugate and the presenter protein is formed, wherein formation of the complex indicates that the conjugate is one that is capable of forming a complex with a presenter protein.

15 In some embodiments, the compounds, conjugates, and complexes of the present invention may be useful for the identification of target proteins capable of forming covalent bonds to compounds in the presence of a presenter protein.

20 Accordingly, in another aspect, the disclosure provides a method of identifying and/or characterizing a target protein capable of reacting with a compound in the presence of a presenter protein, wherein the compound includes a presenter protein binding moiety and a cross-linking moiety. This method includes the steps of: (a) providing (i) a compound including a presenter protein binding 25 moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter protein under conditions suitable for to permit complex formation if the conjugate is capable of forming a complex with the presenter protein; and (c) determining whether the target protein and the compound react during formation of the complex to form a conjugate, wherein if the target protein and the compound form a conjugate, the target protein is identified as capable of reacting with the compound in the presence of a presenter protein.

30 In some embodiments, the compounds, conjugates, and complexes of the invention may be useful for the identification of target proteins capable of forming complexes with presenter proteins.

35 Accordingly, in another aspect, the disclosure provides a method of identifying and/or characterizing a target protein which binds to a presenter protein. This method includes the steps of: (a) providing (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein; (b) combining the conjugate and the presenter protein under conditions suitable to permit complex formation if the conjugate is capable of forming a complex with the presenter protein; and (c) determining whether the target protein binds to the presenter protein in the complex, wherein if the target protein binds to the presenter protein, the target protein is identified as binding to the presenter protein.

40 In some aspects, the disclosure provides a method of identifying and/or characterizing a target protein which binds to a presenter protein. This method includes the steps of: (a) providing (i) a compound including a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter protein under conditions suitable for to permit complex formation if the conjugate is capable of forming a complex with the presenter protein; and (c) determining whether the target protein binds to the presenter protein in

the complex, wherein if the target protein binds to the presenter protein, the target protein is identified as a target protein that binds to a presenter protein.

In some aspects, the disclosure provides a method of identifying and/or characterizing a target protein capable of forming a complex with a presenter protein. This method includes the steps of: (a) providing (i) a compound of Formula VII; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter protein under conditions suitable to permit complex formation if the conjugate is capable of forming a complex with the presenter protein; and (c) determining if the compound, the target protein, and the presenter protein form a complex, wherein if the compound, the target protein, and the presenter protein form a complex, the target protein is identified as a target protein capable of forming a complex with a presenter protein.

In some aspects, the disclosure provides a method of identifying and/or characterizing a target protein which binds to a presenter protein. This method includes the steps of: (a) providing (i) a compound of Formula VII; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter protein under conditions suitable for to permit complex formation if the compound is capable of forming a complex with the presenter protein; and (c) determining whether the target protein binds to the presenter protein in the complex, wherein if the target protein binds to the presenter protein, the target protein is identified as a target protein that binds to a presenter protein.

In some aspects, the disclosure provides a method of identifying a target protein capable of forming a complex with a presenter protein by (a) providing (i) one or more target proteins, (ii) any of the foregoing compounds; and (iii) a presenter protein that includes a tag (e.g., an affinity tag); (b) combining the one or more target proteins, the compound, and the presenter protein under conditions suitable to permit complex formation if one or more of the target proteins is capable of forming a complex with the presenter protein; and (c) determining whether one or more target proteins form a complex with the compound and the presenter protein; wherein target proteins that form a complex with the presenter protein are identified as a target protein capable of forming a complex with a presenter protein.

In some embodiments, the determining step comprises utilizing the tag of said presenter protein to selectively isolate target proteins which have formed a complex with the presenter protein (e.g., by use in a pull down experiment). In some embodiments, the complex includes a target protein, a presenter protein, and a compound of the invention. In some embodiments, the complex includes a conjugate including a target protein and a presenter protein binding moiety (e.g., a conjugate formed by reaction between a cross-linking group of a compound of the invention and a reactive amino acid of a target protein) and a presenter protein. In some embodiments, the method further comprises (d) identifying the target protein (e.g., determining the structure of the target protein) in a complex formed between one or more target proteins, the compound, and the presenter protein. In some embodiments, the identifying of the structure of the target protein comprises performing mass spectrometry on the complex. In some embodiments, determination of whether the target protein and presenter protein form a complex and/or target protein binds to the presenter protein in the complex may be carried out using pull down experiments wherein either the target protein or the presenter protein is labeled (e.g., wherein a complex may be selectively pulled down in the presence of target proteins and/or presenter proteins which are not in a complex).

In some aspects, the disclosure provides a method of identifying a target protein capable of forming a complex with a presenter protein, by (a) providing (i) two or more target proteins; (ii) any of the foregoing compounds; and (iii) a presenter protein including an affinity tag; (b) combining the two or more target proteins, the compound, and the presenter protein under conditions suitable to permit complex formation if said target protein is capable of forming a complex with the presenter protein; (c) selectively isolating one or more complexes of a target protein, the compound, and the presenter protein formed in step (b); and (d) identifying the target protein (e.g., determining the structure of the target protein) in the one or more complexes isolated in step (c) by mass spectrometry; thereby identifying a target protein capable of forming a complex with a presenter protein.

10 In some embodiments, the determining step comprises utilizing the tag of said presenter protein to selectively isolate target proteins which have formed a complex with the presenter protein (e.g., by use in a pull down experiment). In some embodiments, the complex includes a target protein, a presenter protein, and a compound of the invention. In some embodiments, the complex includes a conjugate including a target protein and a presenter protein binding moiety (e.g., a conjugate formed by reaction 15 between a cross-linking group of a compound of the invention and a reactive amino acid of a target protein) and a presenter protein. In some embodiments, determination of whether the target protein and presenter protein form a complex and/or target protein binds to the presenter protein in the complex may be carried out using pull down experiments wherein either the target protein or the presenter protein is labeled (e.g., wherein a complex may be selectively pulled down in the presence of target proteins and/or 20 presenter proteins which are not in a complex).

In some embodiments, the compounds, conjugates, and complexes of the present invention may be useful to identify locations on target proteins to attach presenter protein binding moieties which result in conjugates capable of forming complexes with presenter proteins.

Accordingly, in another aspect, the disclosure provides a method of identifying and/or 25 characterizing a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein. This method includes the steps of: (a) providing (i) a conjugate including a presenter protein binding moiety conjugated to a target protein at a location and (ii) a presenter protein; (b) combining the conjugate and the presenter protein; (c) determining if the conjugate and the presenter protein form a complex; and (d) optionally repeating 30 steps (a) to (c) with the presenter protein binding moiety conjugated at different locations on the target protein until a conjugate and the presenter protein form a complex, wherein a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein is identified if the conjugate and the presenter protein form a complex. In some embodiments, the presenter protein is a variant of a naturally occurring target protein.

35 In some aspects, the disclosure provides a method of identifying and/or characterizing a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein. This method includes the steps of: (a) providing (i) a compound including a presenter protein binding moiety and a cross-linking group; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound with the target protein in the presence of the 40 presenter protein under conditions that permit the formation of a conjugate including a presenter protein binding moiety conjugated to a target protein at a location; (c) determining if the conjugate and the

presenter protein form a complex; and (d) optionally repeating steps (a) to (c) wherein the presenter protein binding moiety is conjugated at different locations on the target protein until a conjugate and the presenter protein form a complex; wherein a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein is identified if the conjugate and the presenter protein form a complex, thereby identifying a location on a target protein to form a conjugate capable of forming a complex with a presenter protein. In some embodiments, the target protein is a variant of a naturally occurring target protein.

5 In some embodiments, the compounds, conjugates, and complexes of the present invention may be useful for identifying compounds capable of forming covalent bonds to target proteins in the presence 10 of presenter proteins. In some embodiments, the compounds identified selectively form covalent bonds with target proteins in the presence of presenter proteins.

15 Accordingly, in another aspect, the disclosure provides a method of identifying and/or characterizing a compound capable of covalently binding to a target protein in the presence of a presenter protein. This method includes the steps of: (a) providing a sample including (i) a compound including a presenter protein binding moiety and a cross-linking group; (ii) a target protein; and (iii) a presenter protein; and (b) determining if the compound and the target protein form a covalent bond via the cross-linking group in said compound in the sample, wherein a compound is identified as covalently 20 binding to a target protein in the presence of a presenter protein if the compound and the target protein react in the sample.

25 In some aspects, the disclosure provides a method of identifying and/or characterizing a compound capable of selective and covalent binding to a target protein in the presence of a presenter protein. This method includes the steps of: (a) providing a first sample including (i) a compound including a presenter protein binding moiety and a cross-linking group; (ii) a target protein; and (iii) a presenter protein and a second sample including (i) the same compound including a presenter protein binding moiety and a cross-linking group as in the first sample and (ii) the same target protein as in the first sample; and (b) determining the extent to which the compound and the target protein react in the first sample compared to the second sample, wherein a compound is identified as selectively covalently 30 binding to a target protein in the presence of a presenter protein if the compound and the target protein reacts in the first sample more than in the second sample.

35 In some embodiments, a compound is identified as selectively covalently binding to a target protein in the presence of a presenter protein if the compound and the target protein reacts in the first sample at least 5-fold more than in the second sample. In some embodiments, a compound is identified as selectively covalently binding to a target protein in the presence of a presenter protein if the compound and the target protein reacts in the first sample, but does not substantially react in the second sample.

40 In some embodiments, the compounds, conjugates, and complexes of the present invention may be useful in identifying conjugates including a target protein and a presenter protein binding moiety capable of forming complexes with presenter proteins.

45 Accordingly, in another aspect, the disclosure provides a method of identifying and/or characterizing a conjugate capable of forming a complex with a presenter protein. This method includes the steps of: (a) providing (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein; and (b) combining the conjugate and the presenter protein

under conditions suitable for forming a complex; (c) determining if the conjugate and the presenter protein form a complex, wherein a conjugate is identified as capable of forming a complex with a presenter protein if the conjugate and the presenter protein form a complex, thereby identifying a conjugate capable of forming a complex with a presenter protein.

5 In some embodiments, binding between a conjugate and a protein may be determined by a method that includes a ternary time-resolved fluorescence energy transfer assay, a ternary amplified luminescent proximity homogeneous assay, a isothermal titration calorimetry, surface plasmon resonance, or nuclear magnetic resonance.

10 In some embodiments, the compounds, conjugates, and complexes of the present invention may be useful for the determining the structure of protein-protein interfaces between presenter proteins and target proteins.

15 Accordingly, in another aspect, the disclosure provides a method of determining the structure of and/or assessing one or more structural features of an interface in a complex including a presenter protein and a target protein. This method includes the steps of: (a) providing (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein; (b) contacting the conjugate with a presenter protein to form a complex (e.g., in a vial); and (c) determining the crystal structure of the complex, wherein the structure of the interface includes at least the portion of the crystal structure between the presenter protein and the target protein, thereby determining the structure of an interface in a complex including a presenter protein and a target protein.

20 In some aspects, the disclosure provides a method of determining the structure of and/or assessing one or more structural features of an interface in a complex including a presenter protein and a target protein. This method includes the steps of: (a) providing (i) a compound including a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter protein under conditions suitable for 25 forming a conjugate between the compound and target protein and for the formation of a complex between said conjugate and said presenter protein (e.g., in a vial); and (c) determining the crystal structure of the complex, wherein the structure of the interface includes at least the portion of the crystal structure between the presenter protein and the target protein, thereby determining the structure of an interface in a complex including a presenter protein and a target protein.

30 In some aspects, the disclosure provides a method of determining the structure of and/or assessing one or more structural features of an interface in a complex including a presenter protein and a target protein. This method includes the steps of: (a) providing (i) a compound of Formula VII; (ii) a target protein; and (iii) a presenter protein; (b) forming a complex including the compound, the target protein, and the presenter protein (e.g., in a vial); and (c) determining the crystal structure of the complex, wherein 35 the structure of the interface includes at least the portion of the crystal structure between the presenter protein and the target protein, thereby determining the structure of an interface in a complex including a presenter protein and a target protein.

40 In some aspects, the disclosure provides a method of determining the structure of and/or assessing one or more structural features of a protein-protein interface in a complex including a presenter protein and a target protein. This method includes the steps of: (a) providing a crystal of any of the foregoing complexes; and (b) determining the structure of the crystal, wherein the structure of the

interface includes at least the portion of the crystal structure between the presenter protein and the target protein, thereby determining the structure of a protein-protein interface in a complex including a presenter protein and a target protein.

In some aspects, the disclosure provides a method of identifying and/or characterizing

5 compounds capable of modulating the biological activity of a target protein. This method includes the steps of: (a) providing the structure of a protein-protein interface in a complex including a presenter protein and a target protein (e.g., a structure determined by any of the foregoing methods); and (b) determining the structure of compounds capable of binding at the interface, thereby identifying compounds capable of modulating the biological activity of a target protein. In some embodiments, the 10 structure of compounds capable of binding at the interface is determined using computational methods. In some embodiments, the structure of compounds capable of binding at the interface is determined by screening of compounds including a presenter protein binding moiety described herein for complex formation in the presence of a target protein and a presenter protein.

In some aspects, the disclosure provides a method of obtaining X-ray crystal coordinates for a

15 complex. This method includes the steps of: (a) providing (i) a conjugate including a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein; (b) combining the conjugate and the presenter protein under conditions suitable for to permit complex formation if the conjugate is capable of forming a complex with the presenter protein; and (c) determining the crystal structure of the complex, thereby obtaining X-ray crystal coordinates for the complex.

20 In some aspects, the disclosure provides a method of obtaining X-ray crystal coordinates for a complex. This method includes the steps of: (a) providing (i) a compound including a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter protein under conditions suitable for to permit complex formation if the compound is capable of forming a complex with the presenter protein; and (c) 25 determining the crystal structure of the complex, thereby obtaining X-ray crystal coordinates for the complex.

In some aspects, the disclosure provides a method of obtaining X-ray crystal coordinates for a complex. This method includes the steps of: (a) providing (i) a compound of the invention; (ii) a target protein; and (iii) a presenter protein; (b) combining the compound, the target protein, and the presenter 30 protein under conditions suitable for to permit complex formation if the compound is capable of forming a complex with the presenter protein; and (c) determining the crystal structure of the complex, thereby obtaining X-ray crystal coordinates for the complex.

In some aspects, the disclosure provides a method of determining the residues on a target protein that participate in binding with a presenter protein. This method includes the steps of: (a)

35 providing X-ray crystal coordinates of a complex obtained by a method of the invention; (b) identifying the residues of the target protein which include an atom within 4 Å of an atom on the presenter protein; thereby determining the residues on a target protein that participate in binding with a presenter protein.

In some aspects, the disclosure provides a method of determining biochemical and/or biophysical properties of any of the presenter protein/target protein complexes described herein. This method

40 includes the steps of: (a) providing X-ray crystal coordinates of a complex described herein obtained by a method described herein; (b) calculating a biochemical and/or biophysical property of the complex;

thereby determining biochemical and/or biophysical properties of a presenter protein/target protein complex.

In some embodiments, the biochemical and/or biophysical properties include the free energy of binding of a complex, the K_d of a complex, the K_i of a complex, the K_{inact} of a complex, and/or the K_i/K_{inact} of a complex. In some embodiments, the biochemical and/or biophysical properties are determined by isothermal titration calorimetry, surface plasmon resonance, and/or mass spectrometry.

5 In some embodiments, the interface in a complex including a presenter protein and a target protein is or comprises a binding pocket.

10 In some aspects, the disclosure provides compositions including any of the foregoing compounds, a target protein, and a presenter protein in solution.

In some aspects, the disclosure provides a pharmaceutical composition including any of the compounds, conjugates, or complexes of the invention and a pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition is in unit dosage form.

15 In some aspects, the disclosure provides a method of modulating a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target proteins or a prokaryotic target protein such as a bacterial target protein). In some embodiments, such a method includes steps of contacting the target protein with a modulating (e.g., positive or negative modulation) amount of any of the compounds (e.g., in the presence of a presenter protein), conjugates including a target protein binding moiety, or compositions of the invention.

20 In some aspects, the disclosure provides a method of modulating (e.g., positively or negatively modulating) a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target proteins or a prokaryotic target protein such as a bacterial target protein). In some embodiments, such a method includes steps of contacting a cell expressing the target protein and a presenter protein with an effective amount of a compound or composition of the invention under 25 conditions wherein the compound can form a complex with the presenter protein and the resulting complex can bind to the target protein, thereby modulating (e.g., positively or negatively modulating) the target protein.


30 In some aspects, the disclosure provides a method of modulating (e.g., positively or negatively modulating) a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target proteins or a prokaryotic target protein such as a bacterial target protein). In some embodiments, such a method includes steps of contacting the target protein with conjugate of the invention including a target protein binding moiety, thereby modulating the target protein.

35 In some aspects, the disclosure provides a method of inhibiting prolyl isomerase activity. In some embodiments, such a method includes contacting a cell expressing the prolyl isomerase with a compound or composition of the invention under conditions that permit the formation of a complex between the compound and the prolyl isomerase, thereby inhibiting the prolyl isomerase activity.

40 In some aspects, the disclosure provides a method of forming a presenter protein/compound complex in a cell. In some embodiments, such a method includes steps of contacting a cell expressing the presenter protein with a compound or composition of the invention under conditions that permit the formation of a complex between the compound and the presenter protein.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety is capable of binding a protein encoded by any one of the genes of Table 1. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety is a prolyl isomerase binding moiety. In 5 some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety is a FKBP binding moiety (e.g., the presenter protein binding moiety is capable of binding FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52), a cyclophilin binding moiety (e.g., the presenter protein binding moiety is capable of binding PP1A, CYPB, CYPc, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, 10 RANBP2, or PPWD1), or a PIN1 binding moiety. In some embodiments of any of the foregoing methods, the presenter protein is known to bind to the presenter protein binding moiety.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety is a FKBP binding moiety (e.g., a selective FKBP binding moiety or a non-selective FKBP binding moiety). In some embodiments of any of the foregoing 15 compounds, conjugates, complexes, compositions, or methods, the FKBP binding moiety includes the structure of Formula IIa or IIb:



Formula IIa Formula IIb

wherein Z¹ and Z² are each, independently, optionally substituted C₁-C₆ alkyl, optionally 20 substituted C₁-C₆ heteroalkyl, or Z¹ and Z² combine to form, with the atoms to which they are attached, an optionally substituted 10 to 40 member macrocycle; and wherein at least one of Z¹ or Z² includes a point of attachment to the cross-linking group;

b and c are independently 0, 1, or 2;

d is 0, 1, 2, 3, 4, 5, 6, or 7;

25 X¹ and X² are each, independently, absent, CH₂, O, S, SO₂, or NR⁴;

each R¹ and R² are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ 30 aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl (e.g., optionally substituted C₂-C₉ heteroaryl), optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl (e.g., optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl), or R¹ and R² combine with the carbon atom to which they are bound to form C=O or R¹ and R² combine to form an optionally substituted C₃-C₁₀ carbocyclyl or optionally substituted C₂-C₉ heterocyclyl;


35 each R³ is, independently, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally

substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, , optionally substituted C₂-C₉ heterocyclyl (e.g., optionally substituted C₂-C₉ heteroaryl), or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl (e.g., optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl) or two R⁸ combine to form an optionally substituted C₃-C₁₀ carbocyclyl,


5 optionally substituted C₆-C₁₀ aryl, e.g., optionally substituted C₂-C₉ heteroaryl; and

each R⁴ is, independently, hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted aryl, C₃-C₇ carbocyclyl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, and optionally substituted C₃-C₇ carbocyclyl C₁-C₆ alkyl.

10 In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety includes the structure:



15 In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety is a cyclophilin binding moiety (e.g., a selective cyclophilin binding moiety or a non-selective cyclophilin binding moiety). In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the cyclophilin binding moiety includes the structure of Formula III or IV:



Formula III

Formula IV

wherein Z^3 , Z^4 , Z^5 , and Z^6 are each, independently, hydroxyl, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, or Z^3 and Z^4 or Z^5 and Z^6 combine to form, with the atoms to

5 which they are attached, an optionally substituted 10 to 40 member macrocycle;


at least one of Z^3 , Z^4 , Z^5 , Z^6 , or R^5 includes a point of attachment to the cross-linking group;
 e is 0, 1, 2, 3, or 4;

R^5 and R^7 are, independently, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally 10 substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl;

R^6 is optionally substituted C_1 - C_6 alkyl; and

15 R^8 is hydrogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted aryl, C_3 - C_7 carbocyclyl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, and optionally substituted C_3 - C_7 carbocyclyl C_1 - C_6 alkyl.


In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the cyclophilin binding moiety includes the structure of Formula IVa:



Formula IVa

wherein each R^7 is, independently, hydroxyl, cyano, optionally substituted amino, halogen, thiol, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted 25 C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl (e.g., optionally substituted C_2 - C_9 heteroaryl), or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl (e.g., optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl).

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein binding moiety includes the structure:



5 In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs. In some embodiments of any of the foregoing compounds, conjugates, 10 complexes, compositions, or methods, the target protein includes an undruggable surface. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the target protein does not have a traditional binding pocket.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the amino acid sequence of the target protein has been modified to substitute at least one 15 native amino acid with a reactive amino acid (e.g., a natural amino acid such as a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine, or a non-natural amino acid). In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the amino acid sequence of the target protein has been modified to substitute at least one native reactive amino acid (e.g., a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine) with a non-reactive amino acid 20 (e.g., a natural amino acid such as a serine, valine, alanine, isoleucine, threonine, tyrosine, aspartic acid, glutamic acid, or leucine, or a non-natural amino acid). In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the at least one native reactive amino acid is a solvent exposed amino acid. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the amino acid sequence of the target protein is 25 modified to substitute all reactive amino acids with a non-reactive amino acid. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the substitution is a


conservative substitution. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the target protein includes only one solvent exposed reactive amino acid.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein is a protein encoded by any one of the genes of Table 1. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the presenter protein is a prolyl isomerase. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the prolyl isomerase is a member of the FKBP family (e.g., FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52), a member of the cyclophilin family (e.g., PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, or PPWD1), or PIN1.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the amino acid sequence of the presenter protein has been modified to substitute at least one native amino acid with a reactive amino acid (e.g., a natural amino acid such as a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine, or a non-natural amino acid). In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the amino acid sequence of the presenter protein has been modified to substitute at least one native reactive amino acid (e.g., a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine) with a non-reactive amino acid (e.g., a natural amino acid such as a serine, valine, alanine, isoleucine, threonine, tyrosine, aspartic acid, glutamic acid, or leucine, or a non-natural amino acid). In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the at least one native reactive amino acid is a solvent exposed amino acid. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the amino acid sequence of the presenter protein is modified to substitute all reactive amino acids with a non-reactive amino acid. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the substitution is a conservative substitution.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the linker is 1 to 20 atoms in length. In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the linker is 1.5 to 30 angstroms in length.

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the linker has the structure of Formula V:




Formula V

wherein A¹ is a bond between the linker and protein binding moiety; A² is a bond between the cross-linking group and the linker; B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁-C₂ alkyl, optionally substituted C₁-C₃ heteroalkyl, O, S, and NR^N; R^N is hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted C₂₋₄ alkenyl, optionally substituted C₂₋₄ alkynyl, optionally substituted C₂₋₆ heterocycl, optionally substituted C₆₋₁₂ aryl, or optionally substituted C₁₋₇ heteroalkyl; C¹ and C² are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f, g, h, i, j, and k are each, independently, 0 or 1; and D is optionally substituted C₁₋₁₀ alkyl, optionally substituted

C₂₋₁₀ alkenyl, optionally substituted C₂₋₁₀ alkynyl, optionally substituted C₂₋₆ heterocyclyl, optionally substituted C₆₋₁₂ aryl, optionally substituted C_{2-C₁₀} polyethylene glycol, or optionally substituted C₁₋₁₀ heteroalkyl, or a chemical bond linking A¹-(B¹)_f-(C¹)_g-(B²)_h to -(B³)_i-(C²)_j-(B⁴)_k-A².

In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions,

5 or methods, the linker includes the structure of Formula VI:



Formula VI

wherein A¹ is a bond between the linker and protein binding moiety;

A^2 is a bond between the cross-linking group and the linker;

10 $|z|$ is 0, 1, 2, or 3;

m is 0 or 1;

n is 0, 1, or 2; and


X^3 , X^4 , and X^5 are each, independently, absent, O, S, $-C\equiv C-$, CR^9R^{10} or NR^{11} ; and

each R⁹, R¹⁰, and R¹¹ are, independently, hydrogen, optionally substituted C₁-C₆ alkyl, optionally

15 substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted aryl, C₃-C₇

carbocyclyl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, and optionally substituted C₃-C₇ carbocyclyl C₁-C₆ alkyl. In some embodiments, each R⁹, R¹⁰, and R¹¹ are, independently, hydrogen, unsubstituted C₁-C₆ alkyl, unsubstituted C₂-C₆ alkenyl, unsubstituted C₂-C₆ alkynyl, unsubstituted aryl, C₃-C₇ carbocyclyl, unsubstituted C₆-C₁₀ aryl C₁-C₆ alkyl, and unsubstituted C₃-C₇ carbocyclyl C₁-C₆ alkyl.

20 In some embodiments of any of the foregoing compounds, conjugates, complexes, compositions, or methods, the linker includes the structure:



25


Chemical Terms

Those skilled in the art will appreciate that certain compounds described herein can exist in one or more different isomeric (e.g., stereoisomers, geometric isomers, tautomers) and/or isotopic (e.g., in which one or more atoms has been substituted with a different isotope of the atom, such as hydrogen substituted for deuterium) forms. Unless otherwise indicated or clear from context, a depicted structure can be understood to represent any such isomeric or isotopic form, individually or in combination.

Compounds described herein can be asymmetric (e.g., having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present disclosure that contain asymmetrically substituted carbon atoms can be

isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically active starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the 5 present disclosure. Cis and trans geometric isomers of the compounds of the present disclosure are described and may be isolated as a mixture of isomers or as separated isomeric forms.

In some embodiments, one or more compounds depicted herein may exist in different tautomeric forms. As will be clear from context, unless explicitly excluded, references to such compounds 10 encompass all such tautomeric forms. In some embodiments, tautomeric forms result from the swapping of a single bond with an adjacent double bond and the concomitant migration of a proton. In certain embodiments, a tautomeric form may be a prototropic tautomer, which is an isomeric protonation states having the same empirical formula and total charge as a reference form. Examples of moieties with 15 prototropic tautomeric forms are ketone – enol pairs, amide – imidic acid pairs, lactam – lactim pairs, amide – imidic acid pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, such as, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H- isoindole, and 1H- and 2H-pyrazole. In some embodiments, tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution. In certain embodiments, 20 tautomeric forms result from acetal interconversion, e.g., the interconversion illustrated in the scheme below:



Those skilled in the art will appreciate that, in some embodiments, isotopes of compounds 25 described herein may be prepared and/or utilized in accordance with the present invention. "Isotopes" refers to atoms having the same atomic number but different mass numbers resulting from a different number of neutrons in the nuclei. For example, isotopes of hydrogen include tritium and deuterium. In some embodiments, an isotopic substitution (e.g., substitution of hydrogen with deuterium) may alter the physicochemical properties of the molecules, such as metabolism and/or the rate of racemization of a chiral center.

As is known in the art, many chemical entities (in particular many organic molecules and/or many 30 small molecules) can adopt a variety of different solid forms such as, for example, amorphous forms and/or crystalline forms (e.g., polymorphs, hydrates, solvates, etc). In some embodiments, such entities may be utilized in any form, including in any solid form. In some embodiments, such entities are utilized in a particular form, for example in a particular solid form.

In some embodiments, compounds described and/or depicted herein may be provided and/or utilized in salt form. 35 In certain embodiments, compounds described and/or depicted herein may be provided and/or utilized in hydrate or solvate form.

At various places in the present specification, constituents of compounds of the present disclosure are disclosed in groups or in ranges. It is specifically intended that the present disclosure

include each and every individual subcombination of the members of such groups and ranges. For example, the term "C₁₋₆ alkyl" is specifically intended to individually disclose methyl, ethyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, and C₆ alkyl. Furthermore, where a compound includes a plurality of positions at which substituents are disclosed in groups or in ranges, unless otherwise indicated, the present disclosure is

5 intended to cover individual compounds and groups of compounds (e.g., genera and subgenera) containing each and every individual subcombination of members at each position.

Herein a phrase of the form "optionally substituted X" (e.g., optionally substituted alkyl) is intended to be equivalent to "X, wherein X is optionally substituted" (e.g., "alkyl, wherein said alkyl is optionally substituted"). It is not intended to mean that the feature "X" (e.g. alkyl) *per se* is optional.

10 The term "alkyl," as used herein, refers to saturated hydrocarbon groups containing from 1 to 20 (e.g., from 1 to 10 or from 1 to 6) carbons. In some embodiments, an alkyl group is unbranched (i.e., is linear); in some embodiments, an alkyl group is branched. Alkyl groups are exemplified by methyl, ethyl, n- and iso-propyl, n-, sec-, iso- and tert-butyl, neopentyl, and the like, and may be optionally substituted with one, two, three, or, in the case of alkyl groups of two carbons or more, four substituents

15 independently selected from the group consisting of: (1) C₁₋₆ alkoxy; (2) C₁₋₆ alkylsulfinyl; (3) amino, as defined herein (e.g., unsubstituted amino (i.e., -NH₂) or a substituted amino (i.e., -N(R^{N1})₂, where R^{N1} is as defined for amino); (4) C₆₋₁₀ aryl-C₁₋₆ alkoxy; (5) azido; (6) halo; (7) (C₂₋₉ heterocyclyl)oxy; (8) hydroxyl, optionally substituted with an *O*-protecting group; (9) nitro; (10) oxo (e.g., carboxyaldehyde or acyl); (11) C₁₋₇ spirocyclyl; (12) thioalkoxy; (13) thiol; (14) -CO₂R^A, optionally substituted with an *O*-protecting group

20 and where R^A is selected from the group consisting of (a) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c) C₆₋₁₀ aryl, (d) hydrogen, (e) C₁₋₆ alk-C₆₋₁₀ aryl, (f) amino-C₁₋₂₀ alkyl, (g) polyethylene glycol of -(CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h) amino-polyethylene glycol of -

25 NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl; (15) -C(O)NR^B'R^C', where each of R^B' and R^C' is, independently, selected from the group consisting of (a) hydrogen, (b) C₁₋₆ alkyl, (c) C₆₋₁₀ aryl, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (16) -SO₂R^D', where R^D' is selected from

30 the group consisting of (a) C₁₋₆ alkyl, (b) C₆₋₁₀ aryl, (c) C₁₋₆ alk-C₆₋₁₀ aryl, and (d) hydroxyl; (17) -SO₂NRE'R^F', where each of R^E' and R^F' is, independently, selected from the group consisting of (a) hydrogen, (b) C₁₋₆ alkyl, (c) C₆₋₁₀ aryl and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (18) -C(O)R^G', where R^G' is selected from the group consisting of (a) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c) C₆₋₁₀ aryl, (d) hydrogen, (e) C₁₋₆ alk-C₆₋₁₀ aryl, (f) amino-C₁₋₂₀ alkyl, (g) polyethylene glycol of -

35 (CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h) amino-polyethylene glycol of -NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl; (19) -NR^HC(O)R^I', where R^H is selected from the group consisting of (a1) hydrogen and (b1) C₁₋₆ alkyl,

and R^l is selected from the group consisting of (a2) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b2) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c2) C₆₋₁₀ aryl, (d2) hydrogen, (e2) C₁₋₆ alk-C₆₋₁₀ aryl, (f2) amino-C₁₋₂₀ alkyl, (g2) polyethylene glycol of -(CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h2) amino-polyethylene glycol of -NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl; (20) -NR^JC(O)OR^K, wherein R^J is selected from the group consisting of (a1) hydrogen and (b1) C₁₋₆ alkyl, and R^K is selected from the group consisting of (a2) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b2) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c2) C₆₋₁₀ aryl, (d2) hydrogen, (e2) C₁₋₆ alk-C₆₋₁₀ aryl, (f2) amino-C₁₋₂₀ alkyl, (g2) polyethylene glycol of -(CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h2) amino-polyethylene glycol of -NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl; (21) amidine; and (22) silyl groups such as trimethylsilyl, t-butyldimethylsilyl, and tri-isopropylsilyl. In some embodiments, each of these groups can be further substituted as described herein. For example, the alkylene group of a C₁-alkaryl can be further substituted with an oxo group to afford the respective aryloyl substituent.

The term "alkylene" and the prefix "alk-," as used herein, represent a saturated divalent hydrocarbon group derived from a straight or branched chain saturated hydrocarbon by the removal of two hydrogen atoms, and is exemplified by methylene, ethylene, isopropylene, and the like. The term "C_{x-y} alkylene" and the prefix "C_{x-y} alk-" represent alkylene groups having between x and y carbons. Exemplary values for x are 1, 2, 3, 4, 5, and 6, and exemplary values for y are 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, or 20 (e.g., C₁₋₆, C₁₋₁₀, C₂₋₂₀, C₂₋₆, C₂₋₁₀, or C₂₋₂₀ alkylene). In some embodiments, the alkylene can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for an alkyl group.

The term "alkenyl," as used herein, represents monovalent straight or branched chain groups of, unless otherwise specified, from 2 to 20 carbons (e.g., from 2 to 6 or from 2 to 10 carbons) containing one or more carbon-carbon double bonds and is exemplified by ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, and the like. Alkenyls include both cis and trans isomers. Alkenyl groups may be optionally substituted with 1, 2, 3, or 4 substituent groups that are selected, independently, from amino, aryl, cycloalkyl, or heterocyclyl (e.g., heteroaryl), as defined herein, or any of the exemplary alkyl substituent groups described herein.

The term "alkynyl," as used herein, represents monovalent straight or branched chain groups from 2 to 20 carbon atoms (e.g., from 2 to 4, from 2 to 6, or from 2 to 10 carbons) containing a carbon-carbon triple bond and is exemplified by ethynyl, 1-propynyl, and the like. Alkynyl groups may be optionally substituted with 1, 2, 3, or 4 substituent groups that are selected, independently, from aryl, cycloalkyl, or heterocyclyl (e.g., heteroaryl), as defined herein, or any of the exemplary alkyl substituent groups described herein.

The term "amino," as used herein, represents $-N(R^{N1})_2$, wherein each R^{N1} is, independently, H, OH, NO_2 , $N(R^{N2})_2$, SO_2OR^{N2} , SO_2R^{N2} , SOR^{N2} , an *N*-protecting group, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, cycloalkyl, alkycycloalkyl, carboxyalkyl (e.g., optionally substituted with an *O*-protecting group, such as optionally substituted arylalkoxycarbonyl groups or any described herein), sulfoalkyl, acyl (e.g., 5 acetyl, trifluoroacetyl, or others described herein), alkoxycarbonylalkyl (e.g., optionally substituted with an *O*-protecting group, such as optionally substituted arylalkoxycarbonyl groups or any described herein), heterocyclyl (e.g., heteroaryl), or alk heterocyclyl (e.g., alk heteroaryl), wherein each of these recited R^{N1} groups can be optionally substituted, as defined herein for each group; or two R^{N1} combine to form a heterocyclyl or an *N*-protecting group, and wherein each R^{N2} is, independently, H, alkyl, or aryl. The 10 amino groups of the invention can be an unsubstituted amino (i.e., $-NH_2$) or a substituted amino (i.e., $-N(R^{N1})_2$). In a preferred embodiment, amino is $-NH_2$ or $-NHR^{N1}$, wherein R^{N1} is, independently, OH, NO_2 , NH_2 , NR^{N2}_2 , SO_2OR^{N2} , SO_2R^{N2} , SOR^{N2} , alkyl, carboxyalkyl, sulfoalkyl, acyl (e.g., acetyl, trifluoroacetyl, or others described herein), alkoxycarbonylalkyl (e.g., t-butoxycarbonylalkyl) or aryl, and each R^{N2} can be H, C_{1-20} alkyl (e.g., C_{1-6} alkyl), or C_{6-10} aryl.

15 The term "amino acid," as described herein, refers to a molecule having a side chain, an amino group, and an acid group (e.g., a carboxy group of $-CO_2H$ or a sulfo group of $-SO_3H$), wherein the amino acid is attached to the parent molecular group by the side chain, amino group, or acid group (e.g., the side chain). As used herein, the term "amino acid" in its broadest sense, refers to any compound and/or substance that can be incorporated into a polypeptide chain, e.g., through formation of one or more 20 peptide bonds. In some embodiments, an amino acid has the general structure $H_2N-C(H)(R)-COOH$. In some embodiments, an amino acid is a naturally-occurring amino acid. In some embodiments, an amino acid is a synthetic amino acid; in some embodiments, an amino acid is a D-amino acid; in some 25 embodiments, an amino acid is an L-amino acid. "Standard amino acid" refers to any of the twenty standard L-amino acids commonly found in naturally occurring peptides. "Nonstandard amino acid" refers to any amino acid, other than the standard amino acids, regardless of whether it is prepared synthetically or obtained from a natural source. In some embodiments, an amino acid, including a carboxy- and/or amino-terminal amino acid in a polypeptide, can contain a structural modification as compared with the general structure above. For example, in some embodiments, an amino acid may be modified by 30 methylation, amidation, acetylation, and/or substitution as compared with the general structure. In some embodiments, such modification may, for example, alter the circulating half life of a polypeptide containing the modified amino acid as compared with one containing an otherwise identical unmodified amino acid. In some embodiments, such modification does not significantly alter a relevant activity of a polypeptide containing the modified amino acid, as compared with one containing an otherwise identical 35 unmodified amino acid. As will be clear from context, in some embodiments, the term "amino acid" is used to refer to a free amino acid; in some embodiments it is used to refer to an amino acid residue of a polypeptide. In some embodiments, the amino acid is attached to the parent molecular group by a carbonyl group, where the side chain or amino group is attached to the carbonyl group. In some 40 embodiments, the amino acid is an α -amino acid. In certain embodiments, the amino acid is a β -amino acid. In some embodiments, the amino acid is a γ -amino acid. Exemplary side chains include an optionally substituted alkyl, aryl, heterocyclyl, alkaryl, alk heterocyclyl, aminoalkyl, carbamoylalkyl, and carboxyalkyl. Exemplary amino acids include alanine, arginine, asparagine, aspartic acid, cysteine,

glutamic acid, glutamine, glycine, histidine, hydroxynorvaline, isoleucine, leucine, lysine, methionine, norvaline, ornithine, phenylalanine, proline, pyrrolysine, selenocysteine, serine, taurine, threonine, tryptophan, tyrosine, and valine. Amino acid groups may be optionally substituted with one, two, three, or, in the case of amino acid groups of two carbons or more, four substituents independently selected

5 from the group consisting of: (1) C₁₋₆ alkoxy; (2) C₁₋₆ alkylsulfinyl; (3) amino, as defined herein (e.g., unsubstituted amino (i.e., -NH₂) or a substituted amino (i.e., -N(R^{N1})₂, where R^{N1} is as defined for amino); (4) C₆₋₁₀ aryl-C₁₋₆ alkoxy; (5) azido; (6) halo; (7) (C₂₋₉ heterocyclyl)oxy; (8) hydroxyl; (9) nitro; (10) oxo (e.g., carboxyaldehyde or acyl); (11) C₁₋₇ spirocyclyl; (12) thioalkoxy; (13) thiol; (14) -CO₂R^{A'}, where R^{A'} is selected from the group consisting of (a) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c) C₆₋₁₀ aryl, (d) hydrogen, (e) C₁₋₆ alk-C₆₋₁₀ aryl, (f) amino-C₁₋₂₀ alkyl, (g) polyethylene glycol of -

10 (CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h) amino-polyethylene glycol of -NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl;

15 (15) -C(O)NR^{B'}R^{C'}, where each of R^{B'} and R^{C'} is, independently, selected from the group consisting of (a) hydrogen, (b) C₁₋₆ alkyl, (c) C₆₋₁₀ aryl, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (16) -SO₂R^{D'}, where R^{D'} is selected from the group consisting of (a) C₁₋₆ alkyl, (b) C₆₋₁₀ aryl, (c) C₁₋₆ alk-C₆₋₁₀ aryl, and (d) hydroxyl; (17) -

20 SO₂NR^{E'}R^{F'}, where each of R^{E'} and R^{F'} is, independently, selected from the group consisting of (a) hydrogen, (b) C₁₋₆ alkyl, (c) C₆₋₁₀ aryl and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (18) -C(O)R^{G'}, where R^{G'} is selected from the group consisting of (a) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c) C₆₋₁₀ aryl, (d) hydrogen, (e) C₁₋₆ alk-C₆₋₁₀ aryl, (f) amino-C₁₋₂₀ alkyl, (g) polyethylene glycol of -

25 (CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h) amino-polyethylene glycol of -NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl;

30 (19) -NR^{H'}C(O)R^{I'}, wherein R^{H'} is selected from the group consisting of (a1) hydrogen and (b1) C₁₋₆ alkyl, and R^{I'} is selected from the group consisting of (a2) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b2) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c2) C₆₋₁₀ aryl, (d2) hydrogen, (e2) C₁₋₆ alk-C₆₋₁₀ aryl, (f2) amino-C₁₋₂₀ alkyl, (g2) polyethylene glycol of -(CH₂)_{s2}(OCH₂CH₂)_{s1}(CH₂)_{s3}OR', wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C₁₋₂₀ alkyl, and (h2) amino-polyethylene glycol of -

35 NR^{N1}(CH₂)_{s2}(CH₂CH₂O)_{s1}(CH₂)_{s3}NR^{N1}, wherein s1 is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of s2 and s3, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C₁₋₆ alkyl;

40 (20) -NR^{J'}C(O)OR^{K'}, wherein R^{J'} is selected from the group consisting of (a1) hydrogen and (b1) C₁₋₆ alkyl, and R^{K'} is selected from the group consisting of (a2) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl), (b2) C₂₋₂₀ alkenyl (e.g., C₂₋₆ alkenyl), (c2) C₆₋₁₀ aryl, (d2) hydrogen, (e2) C₁₋₆ alk-C₆₋₁₀ aryl, (f2) amino-C₁₋₂₀ alkyl, (g2) polyethylene

glycol of $-(CH_2)_{s2}(OCH_2CH_2)_{s1}(CH_2)_{s3}OR'$, wherein $s1$ is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of $s2$ and $s3$, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and R' is H or C_{1-20} alkyl, and (h2) amino-polyethylene glycol of $-NR^{N1}(CH_2)_{s2}(CH_2CH_2O)_{s1}(CH_2)_{s3}NR^{N1}$, wherein $s1$ is an integer from 1 to 10 (e.g., from 1 to 6 or from 1 to 4), each of $s2$ and $s3$, independently, is an integer from 0 to 10 (e.g., from 0 to 4, from 0 to 6, from 1 to 4, from 1 to 6, or from 1 to 10), and each R^{N1} is, independently, hydrogen or optionally substituted C_{1-6} alkyl; and (21) amidine. In some embodiments, each of these groups can be further substituted as described herein.

The term "N-alkylated amino acids" as used herein, refers to amino acids containing an optionally substituted C_1 to C_6 alkyl on the nitrogen of the amino acid that forms the peptidic bond. N-alkylated amino acids include, but are not limited to, N-methyl amino acids, such as N-methyl-alanine, N-methyl-threonine, N-methyl-phenylalanine, N-methyl-aspartic acid, N-methyl-valine, N-methyl-leucine, N-methyl-glycine, N-methyl-isoleucine, N(α)-methyl-lysine, N(α)-methyl-asparagine, and N(α)-methyl-glutamine.

The term "aryl," as used herein, represents a mono-, bicyclic, or multicyclic carbocyclic ring system having one or two aromatic rings and is exemplified by phenyl, naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, anthracenyl, phenanthrenyl, fluorenyl, indanyl, indenyl, and the like, and may be optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of: (1) C_{1-7} acyl (e.g., carboxyaldehyde); (2) C_{1-20} alkyl (e.g., C_{1-6} alkyl, C_{1-6} alkoxy- C_{1-6} alkyl, C_{1-6} alkylsulfinyl- C_{1-6} alkyl, amino- C_{1-6} alkyl, azido- C_{1-6} alkyl, (carboxyaldehyde)- C_{1-6} alkyl, halo- C_{1-6} alkyl (e.g., perfluoroalkyl), hydroxy- C_{1-6} alkyl, nitro- C_{1-6} alkyl, or C_{1-6} thioalkoxy- C_{1-6} alkyl); (3) C_{1-20} alkoxy (e.g., C_{1-6} alkoxy, such as perfluoroalkoxy); (4) C_{1-6} alkylsulfinyl; (5) C_{6-10} aryl; (6) amino; (7) C_{1-6} alk-C $_6$ - C_{1-10} aryl; (8) azido; (9) C_{3-8} cycloalkyl; (10) C_{1-6} alk-C $_3$ - C_{8} cycloalkyl; (11) halo; (12) C_{1-12} heterocyclyl (e.g., C_{1-12} heteroaryl); (13) (C_{1-12} heterocyclyl)oxy; (14) hydroxyl; (15) nitro; (16) C_{1-20} thioalkoxy (e.g., C_{1-6} thioalkoxy); (17) $-(CH_2)_qCO_2R^A$, where q is an integer from zero to four, and R^A is selected from the group consisting of (a) C_{1-6} alkyl, (b) C_{6-10} aryl, (c) hydrogen, and (d) C_{1-6} alk-C $_6$ - C_{1-10} aryl; (18) $-(CH_2)_qCONR^B R^C$, where q is an integer from zero to four and where R^B and R^C are independently selected from the group consisting of (a) hydrogen, (b) C_{1-6} alkyl, (c) C_{6-10} aryl, and (d) C_{1-6} alk-C $_6$ - C_{1-10} aryl; (19) $-(CH_2)_qSO_2R^D$, where q is an integer from zero to four and where R^D is selected from the group consisting of (a) alkyl, (b) C_{6-10} aryl, and (c) alk-C $_6$ - C_{1-10} aryl; (20) $-(CH_2)_qSO_2NR^E R^F$, where q is an integer from zero to four and where each of R^E and R^F is, independently, selected from the group consisting of (a) hydrogen, (b) C_{1-6} alkyl, (c) C_{6-10} aryl, and (d) C_{1-6} alk-C $_6$ - C_{1-10} aryl; (21) thiol; (22) C_{6-10} aryloxy; (23) C_{3-8} cycloalkoxy; (24) C_{6-10} aryl- C_{1-6} alkoxy; (25) C_{1-6} alk-C $_3$ - C_{12} heterocyclyl (e.g., C_{1-6} alk-C $_3$ - C_{12} heteroaryl); (26) C_{2-20} alkenyl; and (27) C_{2-20} alkynyl. In some embodiments, each of these groups can be further substituted as described herein. For example, the alkylene group of a C_1 -alkaryl or a C_1 -alkheterocyclyl can be further substituted with an oxo group to afford the respective aryloyl and (heterocyclyl)oyl substituent group.

The "arylalkyl" group, which as used herein, represents an aryl group, as defined herein, attached to the parent molecular group through an alkylene group, as defined herein. Exemplary unsubstituted arylalkyl groups are from 7 to 30 carbons (e.g., from 7 to 16 or from 7 to 20 carbons, such as C_{1-6} alk-C $_6$ - C_{1-10} aryl, C_{1-10} alk-C $_6$ - C_{1-10} aryl, or C_{1-20} alk-C $_6$ - C_{1-10} aryl). In some embodiments, the alkylene and the aryl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective groups.

Other groups preceded by the prefix “alk-” are defined in the same manner, where “alk” refers to a C₁₋₆ alkylene, unless otherwise noted, and the attached chemical structure is as defined herein.

The term “azido” represents an –N₃ group, which can also be represented as –N=N=N.

The terms “carbocyclic” and “carbocyclyl,” as used herein, refer to an optionally substituted C₃₋₁₂

5 monocyclic, bicyclic, or tricyclic non-aromatic ring structure in which the rings are formed by carbon atoms. Carbocyclic structures include cycloalkyl, cycloalkenyl, and cycloalkynyl groups.

10 The “carbocyclylalkyl” group, which as used herein, represents a carbocyclic group, as defined herein, attached to the parent molecular group through an alkylene group, as defined herein. Exemplary unsubstituted carbocyclylalkyl groups are from 7 to 30 carbons (e.g., from 7 to 16 or from 7 to 20 carbons, such as C₁₋₆ alk-C₆₋₁₀ carbocyclyl, C₁₋₁₀ alk-C₆₋₁₀ carbocyclyl, or C₁₋₂₀ alk-C₆₋₁₀ carbocyclyl). In some embodiments, the alkylene and the carbocyclyl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective groups. Other groups preceded by the prefix “alk-” are defined in the same manner, where “alk” refers to a C₁₋₆ alkylene, unless otherwise noted, and the attached chemical structure is as defined herein.

15 The term “carbonyl,” as used herein, represents a C(O) group, which can also be represented as C=O.

The term “carboxy,” as used herein, means –CO₂H.

The term “cyano,” as used herein, represents an –CN group.

20 The term “cycloalkyl,” as used herein, represents a monovalent saturated or unsaturated non-aromatic cyclic hydrocarbon group from three to eight carbons, unless otherwise specified, and is exemplified by cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclic heptyl, and the like. When the cycloalkyl group includes one carbon-carbon double bond, the cycloalkyl group can be referred to as a “cycloalkenyl” group. Exemplary cycloalkenyl groups include cyclopentenyl, cyclohexenyl, and the like. The cycloalkyl groups of this invention can be optionally substituted with: (1) C₁₋₇ acyl (e.g., 25 carboxyaldehyde); (2) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₁₋₆ alkylsulfinyl-C₁₋₆ alkyl, amino-C₁₋₆ alkyl, azido-C₁₋₆ alkyl, (carboxyaldehyde)-C₁₋₆ alkyl, halo-C₁₋₆ alkyl (e.g., perfluoroalkyl), hydroxy-C₁₋₆ alkyl, nitro-C₁₋₆ alkyl, or C₁₋₆ thioalkoxy-C₁₋₆ alkyl); (3) C₁₋₂₀ alkoxy (e.g., C₁₋₆ alkoxy, such as 30 perfluoroalkoxy); (4) C₁₋₆ alkylsulfinyl; (5) C₆₋₁₀ aryl; (6) amino; (7) C₁₋₆ alk-C₆₋₁₀ aryl; (8) azido; (9) C₃₋₈ cycloalkyl; (10) C₁₋₆ alk-C₃₋₈ cycloalkyl; (11) halo; (12) C₁₋₁₂ heterocyclyl (e.g., C₁₋₁₂ heteroaryl); (13) (C₁₋₁₂ 35 heterocyclyl)oxy; (14) hydroxyl; (15) nitro; (16) C₁₋₂₀ thioalkoxy (e.g., C₁₋₆ thioalkoxy); (17) –(CH₂)_qCO₂R^A, where q is an integer from zero to four, and R^A is selected from the group consisting of (a) C₁₋₆ alkyl, (b) C₆₋₁₀ aryl, (c) hydrogen, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (18) –(CH₂)_qCONR^BR^C, where q is an integer from zero to four and where R^B and R^C are independently selected from the group consisting of (a) hydrogen, (b) C₆₋₁₀ alkyl, (c) C₆₋₁₀ aryl, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (19) –(CH₂)_qSO₂R^D, where q is an integer from zero to four and where R^D is selected from the group consisting of (a) C₆₋₁₀ alkyl, (b) C₆₋₁₀ aryl, and (c) C₁₋₆ alk-C₆₋₁₀ aryl; (20) –(CH₂)_qSO₂NR^ER^F, where q is an integer from zero to four and where each of R^E and R^F is, independently, selected from the group consisting of (a) hydrogen, (b) C₆₋₁₀ alkyl, (c) C₆₋₁₀ aryl, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (21) thiol; (22) C₆₋₁₀ aryloxy; (23) C₃₋₈ cycloalkoxy; (24) C₆₋₁₀ aryl-C₁₋₆ alkoxy; (25) C₁₋₆ alk-C₁₋₁₂ heterocyclyl (e.g., C₁₋₆ alk-C₁₋₁₂ heteroaryl); (26) oxo; (27) C₂₋₂₀ alkenyl; and (28) C₂₋₂₀ alkynyl. In 40 some embodiments, each of these groups can be further substituted as described herein. For example,

the alkylene group of a C₁-alkaryl or a C₁-alkheterocyclyl can be further substituted with an oxo group to afford the respective aryloyl and (heterocyclyl)oyl substituent group.

The “cycloalkylalkyl” group, which as used herein, represents a cycloalkyl group, as defined herein, attached to the parent molecular group through an alkylene group, as defined herein (e.g., an alkylene group of from 1 to 4, from 1 to 6, from 1 to 10, or from 1 to 20 carbons). In some embodiments, the alkylene and the cycloalkyl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective group.

The term “diastereomer,” as used herein, means stereoisomers that are not mirror images of one another and are non-superimposable on one another.

10 The term “enantiomer,” as used herein, means each individual optically active form of a compound of the invention, having an optical purity or enantiomeric excess (as determined by methods standard in the art) of at least 80% (i.e., at least 90% of one enantiomer and at most 10% of the other enantiomer), preferably at least 90% and more preferably at least 98%.

15 The term “halo,” as used herein, represents a halogen selected from bromine, chlorine, iodine, or fluorine.

20 The term “heteroalkyl,” as used herein, refers to an alkyl group, as defined herein, in which one or two of the constituent carbon atoms have each been replaced by nitrogen, oxygen, or sulfur. In some embodiments, the heteroalkyl group can be further substituted with 1, 2, 3, or 4 substituent groups as described herein for alkyl groups. The terms “heteroalkenyl” and heteroalkynyl,” as used herein refer to alkenyl and alkynyl groups, as defined herein, respectively, in which one or two of the constituent carbon atoms have each been replaced by nitrogen, oxygen, or sulfur. In some embodiments, the heteroalkenyl and heteroalkynyl groups can be further substituted with 1, 2, 3, or 4 substituent groups as described herein for alkyl groups.

25 The term “heteroaryl,” as used herein, represents that subset of heterocycls, as defined herein, which are aromatic: i.e., they contain $4n+2$ pi electrons within the mono- or multicyclic ring system. Exemplary unsubstituted heteroaryl groups are of 1 to 12 (e.g., 1 to 11, 1 to 10, 1 to 9, 2 to 12, 2 to 11, 2 to 10, or 2 to 9) carbons. In some embodiment, the heteroaryl is substituted with 1, 2, 3, or 4 substituents groups as defined for a heterocyclyl group.


30 The term “heteroarylalkyl” refers to a heteroaryl group, as defined herein, attached to the parent molecular group through an alkylene group, as defined herein. Exemplary unsubstituted heteroarylalkyl groups are from 2 to 32 carbons (e.g., from 2 to 22, from 2 to 18, from 2 to 17, from 2 to 16, from 3 to 15, from 2 to 14, from 2 to 13, or from 2 to 12 carbons, such as C₁₋₆ alk-C₁₋₁₂ heteroaryl, C₁₋₁₀ alk-C₁₋₁₂ heteroaryl, or C₁₋₂₀ alk-C₁₋₁₂ heteroaryl). In some embodiments, the alkylene and the heteroaryl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective group.

35 Heteroarylalkyl groups are a subset of heterocyclalkyl groups.

40 The term “heterocycl,” as used herein, represents a 5-, 6- or 7-membered ring, unless otherwise specified, containing one, two, three, or four heteroatoms independently selected from the group consisting of nitrogen, oxygen, and sulfur. The 5-membered ring has zero to two double bonds, and the 6- and 7-membered rings have zero to three double bonds. Exemplary unsubstituted heterocycl groups are of 1 to 12 (e.g., 1 to 11, 1 to 10, 1 to 9, 2 to 12, 2 to 11, 2 to 10, or 2 to 9) carbons. The term “heterocycl” also represents a heterocyclic compound having a bridged

multicyclic structure in which one or more carbons and/or heteroatoms bridges two non-adjacent members of a monocyclic ring, e.g., a quinuclidinyl group. The term “heterocyclyl” includes bicyclic, tricyclic, and tetracyclic groups in which any of the above heterocyclic rings is fused to one, two, or three carbocyclic rings, e.g., an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, 5 a cyclopentene ring, or another monocyclic heterocyclic ring, such as indolyl, quinolyl, isoquinolyl, tetrahydroquinolyl, benzofuryl, benzothienyl and the like. Examples of fused heterocyclyls include tropanes and 1,2,3,5,8,8a-hexahydroindolizine. Heterocyclics include pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, piperidinyl, homopiperidinyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, 10 isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, indazolyl, quinolyl, isoquinolyl, quinoxalinyl, dihydroquinoxalinyl, quinazolinyl, cinnolinyl, phthalazinyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, benzothiadiazolyl, furyl, thienyl, thiazolidinyl, isothiazolyl, triazolyl, tetrazolyl, oxadiazolyl (e.g., 1,2,3-oxadiazolyl), purinyl, thiadiazolyl (e.g., 1,2,3-thiadiazolyl), tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, dihydrothienyl, 15 dihydroindolyl, dihydroquinolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, dihydroisoquinolyl, pyranyl, dihydropyranol, dithiazolyl, benzofuranyl, isobenzofuranyl, benzothienyl, and the like, including dihydro and tetrahydro forms thereof, where one or more double bonds are reduced and replaced with hydrogens. Still other exemplary heterocyclyls include: 2,3,4,5-tetrahydro-2-oxo-oxazolyl; 2,3-dihydro-2-oxo-1H-imidazolyl; 2,3,4,5-tetrahydro-5-oxo-1H-pyrazolyl (e.g., 2,3,4,5-tetrahydro-2-phenyl-20 5-oxo-1H-pyrazolyl); 2,3,4,5-tetrahydro-2,4-dioxo-1H-imidazolyl (e.g., 2,3,4,5-tetrahydro-2,4-dioxo-5-methyl-5-phenyl-1H-imidazolyl); 2,3-dihydro-2-thioxo-1,3,4-oxadiazolyl (e.g., 2,3-dihydro-2-thioxo-5-phenyl-1,3,4-oxadiazolyl); 4,5-dihydro-5-oxo-1H-triazolyl (e.g., 4,5-dihydro-3-methyl-4-amino 5-oxo-1H-triazolyl); 1,2,3,4-tetrahydro-2,4-dioxopyridinyl (e.g., 1,2,3,4-tetrahydro-2,4-dioxo-3,3-diethylpyridinyl); 2,6-dioxo-piperidinyl (e.g., 2,6-dioxo-3-ethyl-3-phenylpiperidinyl); 1,6-dihydro-6-25 oxopyridinyl; 1,6-dihydro-4-oxopyrimidinyl (e.g., 2-(methylthio)-1,6-dihydro-4-oxo-5-methylpyrimidin-1-yl); 1,2,3,4-tetrahydro-2,4-dioxopyrimidinyl (e.g., 1,2,3,4-tetrahydro-2,4-dioxo-3-ethylpyrimidinyl); 1,6-dihydro-6-oxo-pyridazinyl (e.g., 1,6-dihydro-6-oxo-3-ethylpyridazinyl); 1,6-dihydro-6-oxo-1,2,4-triazinyl (e.g., 1,6-dihydro-5-isopropyl-6-oxo-1,2,4-triazinyl); 2,3-dihydro-2-oxo-1H-indolyl (e.g., 3,3-dimethyl-2,3-dihydro-2-oxo-1H-indolyl and 2,3-dihydro-2-oxo-3,3'-spiropropane-30 1H-indol-1-yl); 1,3-dihydro-1-oxo-2H-iso-indolyl; 1,3-dihydro-1,3-dioxo-2H-iso-indolyl; 1H-benzopyrazolyl (e.g., 1-(ethoxycarbonyl)-1H-benzopyrazolyl); 2,3-dihydro-2-oxo-1H-benzimidazolyl (e.g., 3-ethyl-2,3-dihydro-2-oxo-1H-benzimidazolyl); 2,3-dihydro-2-oxo-benzoxazolyl (e.g., 5-chloro-2,3-dihydro-2-oxo-benzoxazolyl); 2,3-dihydro-2-oxo-benzoxazolyl; 2-oxo-2H-benzopyranol; 1,4-benzodioxanyl; 1,3-benzodioxanyl; 2,3-dihydro-3-oxo,4H-1,3-benzothiazinyl; 3,4-dihydro-4-oxo-3H-35 quinazolinyl (e.g., 2-methyl-3,4-dihydro-4-oxo-3H-quinazolinyl); 1,2,3,4-tetrahydro-2,4-dioxo-3H-quinazolinyl (e.g., 1-ethyl-1,2,3,4-tetrahydro-2,4-dioxo-3H-quinazolinyl); 1,2,3,6-tetrahydro-2,6-dioxo-7H-purinyl (e.g., 1,2,3,6-tetrahydro-1,3-dimethyl-2,6-dioxo-7H-purinyl); 1,2,3,6-tetrahydro-2,6-dioxo-1H-purinyl (e.g., 1,2,3,6-tetrahydro-3,7-dimethyl-2,6-dioxo-1H-purinyl); 2-oxobenz[c,d]indolyl; 1,1-dioxo-2H-naphth[1,8-c,d]isothiazolyl; and 1,8-naphthylenedicarboxamido. Additional heterocyclics 40 include 3,3a,4,5,6,6a-hexahydro-pyrrolo[3,4-b]pyrrol-(2H)-yl, and 2,5-diazabicyclo[2.2.1]heptan-2-yl,

homopiperazinyl (or diazepanyl), tetrahydropyranyl, dithiazolyl, benzofuranyl, benzothienyl, oxepanyl, thiepanyl, azocanyl, oxecanyl, and thiocanyl. Heterocyclic groups also include groups of the formula



E' is selected from the group consisting of -N- and -CH-; F' is selected from the group consisting

5 of -N=CH-, -NH-CH₂-, -NH-C(O)-, -NH-, -CH=N-, -CH₂-NH-, -C(O)-NH-, -CH=CH-, -CH₂-, -CH₂CH₂-, -CH₂O-, -OCH₂-, -O-, and -S-; and G' is selected from the group consisting of -CH- and -N-. Any of the heterocycl groups mentioned herein may be optionally substituted with one, two, three, four or five substituents independently selected from the group consisting of: (1) C₁₋₇ acyl (e.g., carboxyaldehyde); (2) C₁₋₂₀ alkyl (e.g., C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₁₋₆ alkylsulfinyl-C₁₋₆ alkyl, amino-C₁₋₆ alkyl, azido-C₁₋₆ alkyl, (carboxyaldehyde)-C₁₋₆ alkyl, halo-C₁₋₆ alkyl (e.g., perfluoroalkyl), hydroxy-C₁₋₆ alkyl, nitro-C₁₋₆ alkyl, or C₁₋₆ thioalkoxy-C₁₋₆ alkyl); (3) C₁₋₂₀ alkoxy (e.g., C₁₋₆ alkoxy, such as perfluoroalkoxy); (4) C₁₋₆ alkylsulfinyl; (5) C₆₋₁₀ aryl; (6) amino; (7) C₁₋₆ alk-C₆₋₁₀ aryl; (8) azido; (9) C₃₋₈ cycloalkyl; (10) C₁₋₆ alk-C₃₋₈ cycloalkyl; (11) halo; (12) C₁₋₁₂ heterocycl (e.g., C₂₋₁₂ heteroaryl); (13) (C₁₋₁₂ heterocycl)oxy; (14) hydroxyl; (15) nitro; (16) C₁₋₂₀ thioalkoxy (e.g., C₁₋₆ thioalkoxy); (17) -(CH₂)_qCO₂R^A, where q is an integer

10 from zero to four, and R^A is selected from the group consisting of (a) C₁₋₆ alkyl, (b) C₆₋₁₀ aryl, (c) hydrogen, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (18) -(CH₂)_qCONR^BR^C, where q is an integer from zero to four and where R^B and R^C are independently selected from the group consisting of (a) hydrogen, (b) C₁₋₆ alkyl, (c) C₆₋₁₀ aryl, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (19) -(CH₂)_qSO₂R^D, where q is an integer from zero to four and where R^D is selected from the group consisting of (a) C₁₋₆ alkyl, (b) C₆₋₁₀ aryl, and (c) C₁₋₆ alk-C₆₋₁₀ aryl; (20) -(CH₂)_qSO₂NR^ER^F, where q is an integer from zero to four and where each of R^E and R^F is, independently, selected from the group consisting of (a) hydrogen, (b) C₁₋₆ alkyl, (c) C₆₋₁₀ aryl, and (d) C₁₋₆ alk-C₆₋₁₀ aryl; (21) thiol; (22) C₆₋₁₀ aryloxy; (23) C₃₋₈ cycloalkoxy; (24) arylalkoxy; (25) C₁₋₆ alk-C₁₋₁₂ heterocycl (e.g., C₁₋₆ alk-C₁₋₁₂ heteroaryl); (26) oxo; (27) (C₁₋₁₂ heterocycl)imino; (28) C₂₋₂₀ alkenyl; and (29) C₂₋₂₀ alkynyl. In some embodiments, each of these groups can be further substituted as described herein. For example, the alkylene group of a C₁-alkaryl or a C₁-alkheterocycl can be further substituted with an oxo group to afford the respective aryloyl and (heterocycl)oyl substituent group.

The “heterocyclalkyl” group, which as used herein, represents a heterocycl group, as defined herein, attached to the parent molecular group through an alkylene group, as defined herein. Exemplary unsubstituted heterocyclalkyl groups are from 2 to 32 carbons (e.g., from 2 to 22, from 2 to 18, from 2 to 17, from 2 to 16, from 3 to 15, from 2 to 14, from 2 to 13, or from 2 to 12 carbons, such as C₁₋₆ alk-C₁₋₁₂ heterocycl, C₁₋₁₀ alk-C₁₋₁₂ heterocycl, or C₁₋₂₀ alk-C₁₋₁₂ heterocycl). In some embodiments, the alkylene and the heterocycl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective group.

The term “hydrocarbon,” as used herein, represents a group consisting only of carbon and hydrogen atoms.

The term “hydroxyl,” as used herein, represents an -OH group. In some embodiments, the hydroxyl group can be substituted with 1, 2, 3, or 4 substituent groups (e.g., O-protecting groups) as defined herein for an alkyl.

The term "isomer," as used herein, means any tautomer, stereoisomer, enantiomer, or diastereomer of any compound of the invention. It is recognized that the compounds of the invention can have one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (i.e., geometric E/Z isomers) or diastereomers (e.g., enantiomers (i.e., (+) or (-)) or 5 cis/trans isomers). According to the invention, the chemical structures depicted herein, and therefore the compounds of the invention, encompass all of the corresponding stereoisomers, that is, both the stereomerically pure form (e.g., geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures, e.g., racemates. Enantiomeric and stereoisomeric mixtures of 10 compounds of the invention can typically be resolved into their component enantiomers or stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and stereoisomers can also be obtained from stereomerically or 15 enantiomerically pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.

15 The term "N-protected amino," as used herein, refers to an amino group, as defined herein, to which is attached one or two N-protecting groups, as defined herein.

The term "N-protecting group," as used herein, represents those groups intended to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, "Protective Groups in Organic Synthesis," 3rd Edition (John Wiley & Sons, New York, 1999), which is incorporated herein by reference. N-protecting groups include acyl, 20 aryl, or carbamyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxyacetyl, α -chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and chiral auxiliaries such as protected or unprotected D, L or D, L-amino acids such as alanine, leucine, phenylalanine, and the like; sulfonyl-containing groups 25 such as benzenesulfonyl, p-toluenesulfonyl, and the like; carbamate forming groups such as benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1- 30 methylethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxy carbonyl, t-butylloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxy carbonyl, 4-nitrophenoxy carbonyl, fluorenyl-9-methoxycarbonyl, cyclopentyloxycarbonyl, adamantlyloxycarbonyl, cyclohexyloxycarbonyl, 35 phenylthiocarbonyl, and the like, alkaryl groups such as benzyl, triphenylmethyl, benzyloxymethyl, and the like and silyl groups, such as trimethylsilyl, and the like. Preferred N-protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, alanyl, phenylsulfonyl, benzyl, t-butyloxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).

The term "nitro," as used herein, represents an $-NO_2$ group.

40 The term "O-protecting group," as used herein, represents those groups intended to protect an oxygen containing (e.g., phenol, hydroxyl, or carbonyl) group against undesirable reactions during synthetic procedures. Commonly used O-protecting groups are disclosed in Greene, "Protective Groups

in Organic Synthesis," 3rd Edition (John Wiley & Sons, New York, 1999), which is incorporated herein by reference. Exemplary *O*-protecting groups include acyl, aryloyl, or carbamyl groups, such as formyl, acetyl, propionyl, pivaloyl, *t*-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxyacetyl, α -chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, *t*-butyldimethylsilyl, tri-*iso*-propylsilyloxymethyl, 4,4'-dimethoxytrityl, isobutyryl, phenoxyacetyl, 4-isopropylpehenoxyacetyl, dimethylformamidino, and 4-nitrobenzoyl; alkylcarbonyl groups, such as acyl, acetyl, propionyl, pivaloyl, and the like; optionally substituted arylcarbonyl groups, such as benzoyl; silyl groups, such as trimethylsilyl (TMS), *tert*-butyldimethylsilyl (TBDMS), tri-*iso*-propylsilyloxymethyl (TOM), triisopropylsilyl (TIPS), and the like; ether-forming groups with the hydroxyl, such methyl, methoxymethyl, tetrahydropyranyl, benzyl, p-methoxybenzyl, trityl, and the like; alkoxy carbonyls, such as methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, n-isopropoxycarbonyl, n-butyloxycarbonyl, isobutyloxycarbonyl, sec-butyloxycarbonyl, *t*-butyloxycarbonyl, 2-ethylhexyloxycarbonyl, cyclohexyloxycarbonyl, methyloxycarbonyl, and the like; alkoxyalkoxycarbonyl groups, such as methoxymethoxycarbonyl, ethoxymethoxycarbonyl, 2-methoxyethoxycarbonyl, 2-ethoxyethoxycarbonyl, 2-butoxyethoxycarbonyl, 2-methoxyethoxymethoxycarbonyl, allyloxycarbonyl, propargyloxycarbonyl, 2-butenoxycarbonyl, 3-methyl-2-butenoxycarbonyl, and the like; haloalkoxycarbonyls, such as 2-chloroethoxycarbonyl, 2-chloroethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, and the like; optionally substituted arylalkoxycarbonyl groups, such as benzyloxycarbonyl, p-methylbenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2,4-dinitrobenzyloxycarbonyl, 3,5-dimethylbenzyloxycarbonyl, p-chlorophenoxy carbonyl, 2-chloro-4-nitrophenoxy carbonyl, and the like); substituted alkyl, aryl, and alkaryl ethers (e.g., trityl; methylthiomethyl; methoxymethyl; benzyloxymethyl; siloxymethyl; 2,2,2-trichloroethoxymethyl; tetrahydropyranyl; tetrahydrofuran; ethoxyethyl; 1-[2-(trimethylsilyl)ethoxy]ethyl; 2-trimethylsilylethyl; *t*-butyl ether; p-chlorophenyl, p-methoxyphenyl, p-nitrophenyl, benzyl, p-methoxybenzyl, and nitrobenzyl); silyl ethers (e.g., trimethylsilyl; triethylsilyl; triisopropylsilyl; dimethylisopropylsilyl; *t*-butyldimethylsilyl; *t*-butyldiphenylsilyl; tribenzylsilyl; triphenylsilyl; and diphenylmethylsilyl); carbonates (e.g., methyl, methoxymethyl, 9-fluorenylmethyl; ethyl; 2,2,2-trichloroethyl; 2-(trimethylsilyl)ethyl; vinyl, allyl, nitrophenyl; benzyl; methoxybenzyl; 3,4-dimethoxybenzyl; and nitrobenzyl); carbonyl-protecting groups (e.g., acetal and ketal groups, such as dimethyl acetal, 1,3-dioxolane, and the like; acylal groups; and dithiane groups, such as 1,3-dithianes, 1,3-dithiolane, and the like); carboxylic acid-protecting groups (e.g., ester groups, such as methyl ester, benzyl ester, *t*-butyl ester, orthoesters, and the like; and oxazoline groups).

The term "oxo" as used herein, represents =O.

The prefix "perfluoro," as used herein, represents anyl group, as defined herein, where each hydrogen radical bound to the alkyl group has been replaced by a fluoride radical. For example, perfluoroalkyl groups are exemplified by trifluoromethyl, pentafluoroethyl, and the like.

The term "protected hydroxyl," as used herein, refers to an oxygen atom bound to an *O*-protecting group.

The term "spirocyclyl," as used herein, represents a C₂₋₇ alkylene diradical, both ends of which are bonded to the same carbon atom of the parent group to form a spirocyclic group, and also a C₁₋₆ heteroalkylene diradical, both ends of which are bonded to the same atom. The heteroalkylene radical forming the spirocyclyl group can contain one, two, three, or four heteroatoms independently selected from the group consisting of nitrogen, oxygen, and sulfur. In some embodiments, the spirocyclyl group includes one to seven carbons, excluding the carbon atom to which the diradical is attached. The spirocyclyl groups of the invention may be optionally substituted with 1, 2, 3, or 4 substituents provided herein as optional substituents for cycloalkyl and/or heterocyclyl groups.

The term "stereoisomer," as used herein, refers to all possible different isomeric as well as conformational forms which a compound may possess (e.g., a compound of any formula described herein), in particular all possible stereochemically and conformationally isomeric forms, all diastereomers, enantiomers and/or conformers of the basic molecular structure. Some compounds of the present invention may exist in different tautomeric forms, all of the latter being included within the scope of the present invention.

The term "sulfonyl," as used herein, represents an -S(O)₂- group.

The term "thiol," as used herein, represents an -SH group.

Definitions

In this application, unless otherwise clear from context, (i) the term "a" may be understood to mean "at least one"; (ii) the term "or" may be understood to mean "and/or"; (iii) the terms "comprising" and "including" may be understood to encompass itemized components or steps whether presented by themselves or together with one or more additional components or steps; and (iv) the terms "about" and "approximately" may be understood to permit standard variation as would be understood by those of ordinary skill in the art; and (v) where ranges are provided, endpoints are included.

As is known in the art, "affinity" is a measure of the tightness with which a particular ligand binds to its partner. Affinities can be measured in different ways. In some embodiments, affinity is measured by a quantitative assay. In some such embodiments, binding partner concentration may be fixed to be in excess of ligand concentration so as to mimic physiological conditions. Alternatively or additionally, in some embodiments, binding partner concentration and/or ligand concentration may be varied. In some such embodiments, affinity may be compared to a reference under comparable conditions (e.g., concentrations).

As used herein, the terms "approximately" and "about" are each intended to encompass normal statistical variation as would be understood by those of ordinary skill in the art as appropriate to the relevant context. In certain embodiments, the terms "approximately" or "about" each refer to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of a stated value, unless otherwise stated or otherwise evident from the context (e.g., where such number would exceed 100% of a possible value).

It will be understood that the term "binding" as used herein, typically refers to association (e.g., non-covalent or covalent) between or among two or more entities. "Direct" binding involves physical contact between entities or moieties; indirect binding involves physical interaction by way of physical

contact with one or more intermediate entities. Binding between two or more entities can typically be assessed in any of a variety of contexts – including where interacting entities or moieties are studied in isolation or in the context of more complex systems (e.g., while covalently or otherwise associated with a carrier entity and/or in a biological system or cell).

5 The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_D). Affinity can be measured by common methods known in the art, including those described herein. Specific illustrative and exemplary embodiments for measuring binding affinity are described below. The term “ K_D ,” as used herein, is intended to refer to the dissociation equilibrium constant of a particular compound-protein or complex-protein interaction. Typically, the compounds of the invention
10 bind to presenter proteins with a dissociation equilibrium constant (K_D) of less than about 10^{-6} M, such as less than approximately 10^{-7} M, 10^{-8} M, 10^{-9} M, or 10^{-10} M or even lower, e.g., when determined by surface plasmon resonance (SPR) technology using the presenter protein as the analyte and the compound as the ligand. The presenter protein/compound complexes of the invention bind to target
15 proteins (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target protein or a prokaryotic target protein such as a bacterial target protein) with a dissociation equilibrium constant (K_D) of less than about 10^{-6} M, such as less than approximately 10^{-7} M, 10^{-8} M, 10^{-9} M, or 10^{-10} M or even lower, e.g., when determined by surface plasmon resonance (SPR) technology using the target protein as the analyte and the complex as the ligand.

As used herein, the term “cross-linking group” refers to a group comprising a reactive functional group capable of chemically attaching to specific functional groups (e.g., primary amines, sulfhydryls) on proteins or other molecules. A “moiety capable of a chemoselective reaction with an amino acid,” as used herein refers to a moiety comprising a reactive functional group capable of chemically attaching to a functional group of a natural or non-natural amino acid (e.g., primary and secondary amines, sulfhydryls, alcohols, carboxyl groups, carbonyls, or triazole forming functional groups such as azides or alkynes).
20 Examples of cross-linking groups include sulfhydryl-reactive cross-linking groups (e.g., groups comprising maleimides, haloacetyls, pyridyldisulfides, thiosulfonates, or vinylsulfones), amine-reactive cross-linking groups (e.g., groups comprising esters such as NHS esters, imidoesters, and pentafluorophenyl esters, or hydroxymethylphosphine), carboxyl-reactive cross-linking groups (e.g., groups comprising primary or secondary amines, alcohols, or thiols), carbonyl-reactive cross-linking groups (e.g., groups comprising
25 hydrazides or alkoxyamines), and triazole-forming cross-linking groups (e.g., groups comprising azides or alkynes).
30

As used herein, the term “complex” refers to a group of two or more compounds and/or proteins which are bound together through a binding interaction (e.g., a non-covalent interaction, such as a hydrophobic effect interaction, an electrostatic interaction, a van der Waals interaction, or π -effect
35 interaction). Examples of complexes are “presenter protein/conjugate complex” and “target protein/conjugate complex” which include a conjugate of the invention bound to a presenter protein or a target protein.

As used herein, the term “conjugate” refers to a compound formed by the joining (e.g., via a covalent bond forming reaction) of two or more chemical compounds (e.g., a compound including a cross-linking group and a protein such as a target protein or a presenter protein).
40

As used herein, the term “electron withdrawing group” refers to a functional group which removes electron density from a π system. Examples of electron withdrawing groups include, but are not limited to, halides (e.g., fluoride, chloride, bromide, iodide), aldehydes, ketones, carboxylic acids, acyl chlorides, esters, amides, trihalides (e.g., trifluoromethyl, trichloromethyl), nitriles, sulfonates, and nitro groups.

5 As used herein, the term “leaving group” refers to a molecular fragment that departs with a pair of electrons in a heterolytic bond cleavage. Examples of leaving groups include, but are not limited to halides (e.g., fluoride, chloride, bromide, iodide), carboxylates, tosylates, mesylates, perfluoroalkylsulfonates (e.g., triflate), nitrates, and phosphates.

10 As used herein, an atom that “participates in binding” is within 4 \AA of the entity to which they bind or connects to an atom that is within 4 \AA of the entity to which they bind.

15 The term “presenter protein” refers to a protein that binds to a small molecule to form a complex that binds to and modulates the activity of a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target protein or a prokaryotic target protein such as a bacterial target protein). In some embodiments, the presenter protein is a relatively abundant protein (e.g., the presenter protein is sufficiently abundant that participation in a tripartite complex does not substantially impact the biological role of the presenter protein in a cell and/or viability or other attributes of the cell). In certain embodiments, the presenter protein is a protein that has chaperone activity within a cell. In some embodiments, the presenter protein is a protein that has multiple natural interaction partners within a cell. In certain embodiments, the presenter protein is one which is known to bind a small molecule to form a 20 binary complex that is known to or suspected of binding to and modulating the biological activity of a target protein.

25 The term “presenter protein binding moiety” refers to a group of atoms and the moieties attached thereto (e.g., atoms within 20 atoms such as, atoms within 15 atoms, atoms within 10, atoms within 5 atoms) that participate in binding to a presenter protein such that the compound specifically binds to said presenter protein, for example, with a K_D of less than 10 μM (e.g., less than 5 μM , less than 1 μM , less than 500 nM, less than 200 nM, less than 100 nM, less than 75 nM, less than 50 nM, less than 25 nM, less than 10 nM) or inhibits the peptidyl-prolyl isomerase activity of the presenter protein, for example, with an IC_{50} of less than 1 μM (e.g., less than 0.5 μM , less than 0.1 μM , less than 0.05 μM , less than 0.01 μM). It will be understood that the presenter protein binding moiety does not necessarily encompass the 30 entirety of atoms in the compound that interact with the presenter protein. It will also be understood that one or more atoms of the presenter protein binding moiety may be within the target protein binding moiety (e.g., eukaryotic target protein binding moiety such as mammalian target protein binding moiety or fungal target protein binding moiety or prokaryotic target protein binding moiety such as a bacterial target protein binding moiety).

35 As used herein, “FKBP binding moiety” refers to a presenter protein binding moiety that is selective for presenter proteins in the FKBP family of proteins (e.g., FKBP12, FKBP12.6, FKBPP13, FKBP25, FKBP51, or FKBP52). A “selective FKBP binding moiety,” as used herein, refers to a binding moiety that is specific for one or more (e.g., two, three, four, five) members of the FKBP family over all other members of the FKBP family. A “non-selective FKBP binding moiety,” as used herein, refers to a 40 binding moiety that has comparable affinity (within 2-fold, within 3-fold, within 4-fold, within 5-fold, within 10-fold) for all members of the FKBP family.

The term "protein binding moiety" refers to a group of atoms and the moieties attached thereto (e.g., atoms within 20 atoms such as, atoms within 15 atoms, atoms within 10, atoms within 5 atoms) that participate in binding to a protein (e.g., a presenter protein or a target protein) such that the compound specifically binds to said protein, for example, with a K_D of less than 10 μM (e.g., less than 5 μM , less than 1 μM , less than 500 nM, less than 200 nM, less than 100 nM, less than 75 nM, less than 50 nM, less than 25 nM, less than 10 nM) or inhibits the peptidyl-prolyl isomerase activity of the presenter protein, for example, with an IC_{50} of less than 1 μM (e.g., less than 0.5 μM , less than 0.1 μM , less than 0.05 μM , less than 0.01 μM). It will be understood that the protein binding moiety does not necessarily encompass the entirety of atoms in the compound that interact with the protein.

10 As used herein, the term "react" refers to a process in which atoms of the same or different elements rearrange themselves to form a new substance. For example, the formation of a covalent bond between two atoms such as the reaction between a reactive amino acid on a protein and a cross-linking group to form a covalent bond. A reaction may be measured by any method known in the art, for example, formation of a reaction product can be determined by LC-MS or NMR.

15 As used herein, the term "reactive amino acid" refers to a natural or non-natural amino acid comprising a functional group (e.g., a nucleophilic functional group) capable of chemically attaching to specific functional groups (e.g., a cross-linking group). Examples of reactive amino acids include cysteine, lysine, serine, and amino acids having azides on the side chain. "Non-reactive amino acids" refers to natural or non-natural amino acids that do not contain a functional group capable of chemically 20 attaching to specific functional groups. Examples of non-reactive amino acids include valine, alanine, isoleucine, threonine, and leucine.

The term "reference" is often used herein to describe a standard or control compound, individual, population, sample, sequence or value against which a compound, individual, population, sample, sequence or value of interest is compared. In some embodiments, a reference compound, individual, 25 population, sample, sequence or value is tested and/or determined substantially simultaneously with the testing or determination of the compound, individual, population, sample, sequence or value of interest. In some embodiments, a reference compound, individual, population, sample, sequence or value is a historical reference, optionally embodied in a tangible medium. Typically, as would be understood by those skilled in the art, a reference compound, individual, population, sample, sequence or value is 30 determined or characterized under conditions comparable to those utilized to determine or characterize the compound, individual, population, sample, sequence or value of interest.

As used herein, the term "solvent exposed amino acid" refers to an amino acid that is accessible to the solvent surrounding the protein. In some embodiments, a solvent exposed amino acid is an amino acid that when substituted does not substantially change the three-dimensional structure of the protein.

35 As used herein, the terms "specific binding" or "specific for" or "specific to" refer to an interaction between a binding agent and a target entity. As will be understood by those of ordinary skill, an interaction is considered to be "specific" if it is favored in the presence of alternative interactions, for example, binding with a K_D of less than 10 μM (e.g., less than 5 μM , less than 1 μM , less than 500 nM, less than 200 nM, less than 100 nM, less than 75 nM, less than 50 nM, less than 25 nM, less than 10 nM). In many embodiments, specific interaction is dependent upon the presence of a particular structural 40 feature of the target entity (e.g., an epitope, a cleft, a binding site). It is to be understood that specificity

need not be absolute. In some embodiments, specificity may be evaluated relative to that of the binding agent for one or more other potential target entities (e.g., competitors). In some embodiments, specificity is evaluated relative to that of a reference specific binding agent. In some embodiments specificity is evaluated relative to that of a reference non-specific binding agent.

5 The term "specific" when used with reference to a compound having an activity, is understood by those skilled in the art to mean that the compound discriminates between potential target entities or states. For example, in some embodiments, a compound is said to bind "specifically" to its target if it binds preferentially with that target in the presence of one or more competing alternative targets. In many 10 embodiments, specific interaction is dependent upon the presence of a particular structural feature of the target entity (e.g., an epitope, a cleft, a binding site). It is to be understood that specificity need not be absolute. In some embodiments, specificity may be evaluated relative to that of the binding agent for one or more other potential target entities (e.g., competitors). In some embodiments, specificity is evaluated relative to that of a reference specific binding agent. In some embodiments specificity is evaluated relative to that of a reference non-specific binding agent. In some embodiments, the agent or entity does 15 not detectably bind to the competing alternative target under conditions of binding to its target entity. In some embodiments, binding agent binds with higher on-rate, lower off-rate, increased affinity, decreased dissociation, and/or increased stability to its target entity as compared with the competing alternative target(s).

20 The term "substantially" refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term "substantially" is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical phenomena.

25 The term "does not substantially bind" to a particular protein as used herein can be exhibited, for example, by a molecule or portion of a molecule having a K_D for the target of 10^{-4} M or greater, alternatively 10^{-5} M or greater, alternatively 10^{-6} M or greater, alternatively 10^{-7} M or greater, alternatively 10^{-8} M or greater, alternatively 10^{-9} M or greater, alternatively 10^{-10} M or greater, alternatively 10^{-11} M or greater, alternatively 10^{-12} M or greater, or a K_D in the range of 10^{-4} M to 10^{-12} M or 10^{-6} M to 10^{-10} M or 10^{-7} M to 10^{-9} M.

30 The term "target protein" refers to any protein that participates in a biological pathway associated with a disease, disorder or condition. In some embodiments, the target protein is not mTOR or calcineurin. In some embodiments, the target protein is capable of forming a tripartite complex with a presenter protein and a small molecule. In some embodiments, a target protein is a naturally-occurring protein; in some such embodiments, a target protein is naturally found in certain mammalian cells (e.g., a 35 mammalian target protein), fungal cells (e.g., a fungal target protein), bacterial cells (e.g., a bacterial target protein) or plant cells (e.g., a plant target protein). In some embodiments, a target protein is characterized by natural interaction with one or more natural presenter protein/natural small molecule complexes. In some embodiments, a target protein is characterized by natural interactions with a plurality of different natural presenter protein/natural small molecule complexes; in some such embodiments some 40 or all of the complexes utilize the same presenter protein (and different small molecules). In some embodiments, a target protein does not substantially bind to a complex of cyclosporin, rapamycin, or

FK506 and a presenter protein (e.g., FKBP). Target proteins can be naturally occurring, e.g., wild type. Alternatively, the target protein can vary from the wild type protein but still retain biological function, e.g., as an allelic variant, a splice mutant or a biologically active fragment. Exemplary mammalian target proteins are GTPases, GTPase activating protein, Guanine nucleotide-exchange factor, heat shock proteins, ion channels, coiled-coil proteins, kinases, phosphatases, ubiquitin ligases, transcription factors, chromatin modifier/remodelers, proteins with classical protein-protein interaction domains and motifs, or any other proteins that participate in a biological pathway associated with a disease, disorder or condition.

5 In some embodiments, the target protein is a modified target protein. A modified target protein can include an amino acid insertion, deletion, or substitution, either conservative or non-conservative (e.g., D-amino acids, desamino acids) in the protein sequence (e.g., where such changes do not substantially alter the biological activity of the polypeptide). In particular, the addition of one or more cysteine residues to the amino or carboxy terminus of any of the polypeptides of the invention can facilitate conjugation of these proteins by, e.g., disulfide bonding. In some embodiments, one or more reactive amino acid residues (e.g., cysteines) are removed to decrease the number of possible 10 conjugation sites on the protein. Amino acid substitutions can be conservative (i.e., wherein a residue is replaced by another of the same general type or group) or non-conservative (i.e., wherein a residue is replaced by an amino acid of another type). In addition, a naturally occurring amino acid can be substituted for a non-naturally occurring amino acid (i.e., non-naturally occurring conservative amino acid 15 substitution or a non-naturally occurring non-conservative amino acid substitution).

20 The term “target protein binding moiety” refers to a group of ring atoms and the moieties attached thereto (e.g., atoms within 20 atoms such as, atoms within 15 atoms, atoms within 10 atoms, within 5 atoms) that participate in binding to a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target protein or a prokaryotic target protein such as a bacterial target protein) when the compound is in a complex with a presenter protein. It will be understood that the 25 target protein binding moiety does not necessarily encompass the entirety of atoms in the compound that interact with the target protein. It will also be understood that one or more atoms of the presenter protein binding moiety may also be present in the target protein binding moiety.

30 The term “traditional binding pocket” refers to cavities or pockets on a protein structure with physiochemical and/or geometric properties comparable to proteins whose activity has been modulated by one or more small molecules. In some embodiments, a traditional binding pocket is a well-defined 35 pocket with a volume greater than 1000 Å³. Those of ordinary skill in the art are familiar with the concept of a traditional binding pocket and, moreover are aware of its relationship to “druggability”. In certain embodiments, a protein is considered to not have a traditional binding pocket if it is undruggable, as defined herein.

35 The term “undruggable target” refers to proteins that are not members of a protein family which is known to be targeted by drugs and/or does not possess a binding site that is suitable for high-affinity 40 binding to a small molecule. Methods for determining whether a target protein is undruggable are known in the art. For example, whether a target protein is undruggable may be determined using an structure-based algorithm, such as those used by the program DOGSITESCORER® (Universitat Hamburg, Hamburg, Germany) that assesses druggability based on parameters computed for binding pockets on a protein including volume, surface area, lipophilic surface area, depth, and/or hydrophobic ratio.

Brief Description of the Drawings

FIG. 1 is an image illustrating SDS-PAGE analysis of KRAS_{GTP/S39C} lite/C2-FK506 conjugates.

Lane 1: KRAS_{GTP/S39C} lite; Lane 2: KRAS_{GTP/S39C} lite/C2-FK506 reaction mixture; Lane 3: KRAS_{GTP/S39C} lite/C2-FK506 reaction mixture + 100 mM DTT.

FIG. 2 is an image illustrating SDS-PAGE analysis of KRAS_{GTP/G12C} lite/SFAX9DS conjugates.

FIGS. 3A and 3B are images illustrating SEC and SDS-PAGE Analysis of KRAS_{GTP/S39C} lite/C2Holt/FKBP12 Complex Formation. FIG. 3A) SEC purification profile. Dashed blue lines indicate the peak corresponding to elution of the KRAS_{GTP/S39C} lite/C2Holt/FKBP12 ternary complex; FIG. 3B) SDS-PAGE analysis of SEC elution peaks. Dashed blue lines correspond with the fractions collected for the KRAS_{GTP/S39C} lite/C2Holt/FKBP12 elution peak.

FIG. 4 is an image illustrating SEC profile and SDS-PAGE analysis of the elution peaks confirm the formation of KRAS_{GDP/S39C} lite/SFAC4DS/CypAc_{52S} complex.

FIGS. 5A and 5B are images illustrating the SEC profile and SDS-PAGE analysis of free PTP1B_{S187C} lite and FKBP12 proteins and the PTP1B_{S187C} lite/C3SLF/FKBP12 complex.

FIG. 6 is an image illustrating crosslinking efficiency of C3 and C4SLF by SDS-PAGE.

FIGS. 7A and 7B is an image illustrating the crystal structure of FKBP12-Compound 1-KRAS_{GTP/S39C} complex. FIG. 7A) Ribbon representation showing FKBP12, KRAS_{GTP/S39C} and the ligand. Fo-Fc electron density at 3 σ shown is shown for the ligand in the close-up view. FIG. 7B) Surface representation of the complex with atoms within 4 \AA proximity to either ligand or partner protein colored in red.

FIG. 8 is an image illustrating the crystal structure of CypAc_{52S}-SFAC4DS-KRAS_{GDP/S39C}.

FIGS. 9A and 9B are images illustrating the crystal structure of FKBP12-C3SLF-PTP1B_{S187C}. FIG. 9A illustrates that the crystal contains two complex molecules of FKBP12-C3SLF-PTP1B_{S187C} in the asymmetric unit. FIG. 9B illustrates that the buried surface area of PTP1B_{S187C} is 427 \AA^2 and the buried surface area of C3SLF is 615 \AA^2 .

FIG. 10 is an image illustrating the crystal structure of MCL1_{S245C}/C3SLF/FKBP52.

FIG. 11 is an image illustrating the binding curve of W21487 dependent complex formation of CYPA-W21487- KRAS_{G12C-GTP} ternary complex.

FIG. 12 is an image illustrating the binding curve of W21487 dependent complex formation of CYPA-W21487- KRAS_{G12C-GTP} ternary complex.

FIG. 13 is an image illustrating ITC measurements for the binding of FKBP12-Compound 1 and FKBP12-Compound 2 binary complexes to CEP250.

FIG. 14 is an image illustrating SPR sensorgrams for the binding of FKBP12/Compound 1 to CEP250_{11.4} and CEP250_{29.2}.

FIG. 15 is an image illustrating sensogram and steady state fitting curves for the binding of CYPA/ Compound 3 to KRAS_{G12C-GTP}.

FIG. 16 is an image illustrating fluorescence polarization curves for CypA:C3DS:KRAS complex formation.

FIGS. 17A-17C are images illustrating the 2D 1H-15N TROSY-HSQC spectrum of KRAS_{G12C-GTP} (FIG. 17A), the addition of a stoichiometric amount of CYPA (FIG. 17B), and KRAS and CYPA alone (FIG. 17C).

5

Detailed Description of Certain Embodiments

Small molecules are limited in their targeting ability because their interactions with the target are driven by adhesive forces, the strength of which is roughly proportional to contact surface area. Because of their small size, the only way for a small molecule to build up enough intermolecular contact surface area to effectively interact with a target protein is to be literally engulfed by that protein. Indeed, a large 10 body of both experimental and computational data supports the view that only those proteins having a hydrophobic “pocket” on their surface are capable of binding small molecules. In those cases, binding is enabled by engulfment.

Nature has evolved a strategy that allows a small molecule to interact with target proteins at sites other than hydrophobic pockets. This strategy is exemplified by naturally occurring immunosuppressive 15 drugs cyclosporine A, rapamycin, and FK506. The biological activity of these drugs involves the formation of a high-affinity complex of the small molecule with a small presenting protein. The composite surface of the small molecule and the presenting protein engages the target. Thus, for example, the binary complex formed between cyclosporin A and cyclophilin A targets calcineurin with high affinity and specificity, but neither cyclosporin A or cyclophilin A alone binds calcineurin with measurable affinity.

20 Many important therapeutic targets exert their function by complexation with other proteins. The protein/protein interaction surfaces in many of these systems contain an inner core of hydrophobic side chains surrounded by a wide ring of polar residues. The hydrophobic residues contribute nearly all of the energetically favorable contacts, and hence this cluster has been designated as a “hotspot” for engagement in protein-protein interactions. Importantly, in the aforementioned complexes of naturally 25 occurring small molecules with small presenting proteins, the small molecule provides a cluster of hydrophobic functionality akin to a hotspot, and the protein provides the ring of mostly polar residues. In other words, presented small molecule systems mimic the surface architecture employed widely in natural protein/protein interaction systems.

Nature has demonstrated the ability to reprogram the target specificity of presented small 30 molecules—portable hotspots—through evolutionary diversification. In the best characterized example, the complex formed between FK506 binding protein (FKBP) and FK506 targets calcineurin. However, FKBP can also form a complex with the related molecule rapamycin, and that complex interacts with a completely different target, TorC1. To date, no methodology has been developed to reprogram the binding and modulating ability of presenter protein/ligand interfaces so that they can interact with and 35 modulate other target proteins that have previously been considered undruggable.

In addition, it is well established that some drug candidates fail because they modulate the activity of both the intended target and other non-intended proteins as well. The problem is particularly daunting when the drug binding site of the target protein is similar to binding sites in non-target proteins. The insulin like growth factor receptor (IGF-1R), whose ATP binding pocket is structurally similar to the 40 binding pocket of the non-target insulin receptor (IR), is one such example. Small molecule development candidates that were designed to target IGF-1R typically have the unacceptable side effect of also

modulating the insulin receptor. However, structural dissimilarities do exist between these two proteins in the regions surrounding the ATP binding pocket. Despite such knowledge, no methodology exists to date to take advantage of those differences and develop drugs that are specific to IGF-1R over IR.

The present disclosure provides methods and reagents useful for analyzing protein-protein interfaces such as the interface between a presenter protein (e.g., a member of the FKBP family, a member of the cyclophilin family, or PIN1) and a target protein. In some embodiments, the target and/or presenter proteins are intracellular proteins. In some embodiments, the target and/or presenter proteins are mammalian proteins. In some embodiments, these methods and reagents may be useful for identifying target proteins amenable to inhibition or activation by forming a complex with a presenter protein and a small molecule. In some embodiments, these methods and reagents may be useful in identifying compounds capable of inhibiting or activating target proteins by forming a complex with a presenter protein and the target protein.

Compounds and Conjugates

The disclosure provides compounds including a protein binding moiety (e.g., a presenter protein binding moiety or target protein binding moiety) and a cross-linking group. The invention also features conjugates including a protein binding moiety conjugated to a protein, e.g., a presenter protein binding moiety conjugated to a target protein or a target protein binding moiety conjugated to a presenter protein.


The invention also features compounds of Formula VII:

20

A-L-B


Formula VII


wherein A comprises the structure of Formula VIII:




25

In some embodiments, the compound of the invention is:







Cross-linking groups

5 In some embodiments, compounds of the invention include a cross-linking group. A cross-linking group refers to a group comprising a reactive functional group capable of chemically attaching to specific functional groups (e.g., primary amines, sulfhydryls) on proteins or other molecules. Examples of cross-linking groups include sulfhydryl-reactive cross-linking groups (e.g., groups comprising maleimides, haloacetyls, pyridyldisulfides, thiosulfonates, or vinylsulfones), amine-reactive cross-linking groups (e.g., groups comprising esters such as NHS esters, imidoesters, and pentafluorophenyl esters, or hydroxymethylphosphine), carboxyl-reactive cross-linking groups (e.g., groups comprising primary or secondary amines, alcohols, or thiols), carbonyl-reactive cross-linking groups (e.g., groups comprising hydrazides or alkoxyamines), and triazole-forming cross-linking groups (e.g., groups comprising azides or alkynes).


10 15 Exemplary cross-linking groups include 2'-pyridyldisulfide, 4'-pyridyldisulfide iodoacetyl, maleimides, thioesters, alkyl disulfides, alkylamine disulfides, nitrobenzoic acid disulfide, anhydrides, NHS esters, aldehydes, alkyl chlorides, alkynes, Michael acceptor groups (e.g., α,β -unsubstituted ketones or sulfones), epoxides, heteroaryl nitriles, and azides.

In some embodiments, compounds of the invention include a presenter protein binding moiety. In some embodiments, a presenter protein binding moiety includes a group of atoms (e.g., 5 to 20 atoms, 5 to 10 atoms, 10 to 20 atoms) and may include any moieties attached thereto (e.g., atoms within 20 atoms, atoms within 15 atoms, atoms within 10 atoms, atoms within 5 atoms) that participate in binding to a 5 presenter protein such that a provided compound specifically binds to said presenter protein, for example, with a K_D of less than 10 μM (e.g., less than 5 μM , less than 1 μM , less than 500 nM, less than 200 nM, less than 100 nM, less than 75 nM, less than 50 nM, less than 25 nM, less than 10 nM) or inhibits the 10 peptidyl-prolyl isomerase activity of the presenter protein, for example, with an IC_{50} of less than 1 μM (e.g., less than 0.5 μM , less than 0.1 μM , less than 0.05 μM , less than 0.01 μM). In some embodiments, 15 the presenter protein binding moiety does not encompass the entirety of atoms in a provided compound that interact with the presenter protein. In certain embodiments, one or more atoms of the presenter protein binding moiety do not interact with the presenter protein.


In some embodiments, a presenter protein binding moiety includes a N-acyl proline moiety, a N- 20 acyl-pipeolic acid moiety, a N-acyl 3-morpholino-carboxylic acid moiety, and/or a N-acyl piperzic acid moiety (e.g., with acylation on either nitrogen atom). In certain embodiments, a presenter protein binding 15 moiety includes a N-acyl-pipeolic acid moiety. In some embodiments, a presenter protein binding moiety includes a N-acyl proline moiety. In certain embodiments, a presenter protein binding moiety includes a N-acyl 3-morpholino-carboxylic acid moiety. In some emobodiments, a presenter protein binding moiety includes a N-acyl piperzic acid moiety.


In some embodiments, at least one atom of a presenter protein binding moiety participates in 25 binding with one or more (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or fifteen) of Tyr 27, Phe 37, Asp 38, Arg 41, Phe 47, Gln 54, Glu 55, Val 56, Ile 57, Trp 60, Ala 82, Try 83, His 88, Ile 92, and/or Phe 100 of FKBP12. In some embodiments, at least one at of a presenter protein binding moiety participates in binding with at least one (e.g., two, three, or four) of Arg 41, Gln 54, Glu 55, and/or Ala 82 of FKBP12.


In some embodiments, a presenter protein binding moiety has a structure according to Formula II-IV:




In some embodiments, a presenter protein binding moiety includes or consists of the structure:









or a stereoisomer thereof.

A presenter protein can bind to an atom in a presenter protein binding moiety. Alternatively or 5 additionally, a presenter protein can bind to two or more atoms in a presenter protein binding moiety. In another alternative, a presenter protein bind can to a substituent attached to one or more atoms in a presenter protein binding moiety. Furthermore, in some embodiments, a presenter protein can bind to an atom in a presenter protein binding moiety and to a substituent attached to one or more atoms in a presenter protein binding moiety. In some embodiments, a presenter protein binds to a group that mimics 10 a natural ligand of a presenter protein and wherein the group that mimics a natural ligand of a presenter protein is attached to a presenter protein binding moiety. In some embodiments, a presenter protein binds to a presenter protein and affinity of a presenter protein for a presenter protein in the binary complex is increased relative to the affinity of a presenter protein for a presenter protein in the absence of the complex. Binding in such examples is typically through, but not limited to non-covalent interactions of 15 a presenter protein to a presenter protein binding moiety.

Target Protein Binding Moieties

In some embodiments, compounds of the invention include a target protein binding moiety (e.g., a eukaryotic target protein binding moiety such as a mammalian target protein binding moiety or a fungal 20 target protein binding moiety or a prokaryotic target protein binding moiety such as a bacterial target protein binding moiety). In some embodiments, the target protein binding moiety includes a group of atoms (e.g., 5 to 20 atoms, 5 to 10 atoms, 10 to 20 atoms) and may include any moieties attached thereto (e.g., atoms within 20 atoms, atoms within 15 atoms, atoms within 10 atoms, atoms within 5 atoms) that specifically bind to a target protein. In some embodiments, a target protein binding moiety comprises a 25 plurality of the atoms in the compound that interact with the target protein. In certain embodiments, one or more atoms of a target protein binding moiety do not interact with the target protein.

A target protein can bind to an atom in a target protein binding moiety. Alternatively or additionally, a target protein can bind to two or more atoms in a target protein binding moiety. In another

alternative, a target protein bind can to a substituent attached to one or more atoms in a target protein binding moiety. In another alternative, a target protein can bind to an atom in a target protein binding moiety and to a substituent attached to one or more atoms in a target protein binding moiety. In another alternative, a target protein binds to a group that mimics a natural ligand of a target protein and wherein the group that mimics a natural ligand of a target protein is attached to a target protein binding moiety. In yet another alternative, a target protein binds to a presenter protein and the affinity of a target protein for a presenter protein in the binary complex is increased relative to the affinity of a target protein for a presenter protein in the absence of the complex. Binding in these examples is typically through, but not limited to non-covalent interactions of a target protein to a target protein binding moiety.

10 In some embodiments, the target protein binding moiety includes a cross-linking group (e.g., an internal cross-linking group).

Linkers

15 The compounds of the invention include a linker (e.g., moiety linker joining a protein binding moiety (e.g., a presenter protein binding moiety or a target protein binding moiety) to a cross-linking group or a linker joining a protein binding moiety to a protein (e.g., a presenter protein or target protein). The linker component of the invention is, at its simplest, a bond, but may also provide a linear, cyclic, or branched molecular skeleton having pendant groups covalently linking two moieties.

20 In some embodiments, at least one atom of a linker participates in binding to the presenter protein and/or the target protein. In certain embodiments, at least one atom of a linker does not participate in binding to the presenter protein and/or the target protein.

25 Thus, a linker, when included in a compound and/or conjugate as described herein, achieves linking of two (or more) moieties by covalent means, involving bond formation with one or more functional groups located on either moiety. Examples of chemically reactive functional groups which may be employed for this purpose include, without limitation, amino, hydroxyl, sulfhydryl, carboxyl, carbonyl, carbohydrate groups, vicinal diols, thioethers, 2-aminoalcohols, 2-aminothiols, guanidinyl, imidazolyl, and phenolic groups.

30 In some embodiments, such covalent linking of two (or more) moieties may be effected using a linker that contains reactive moieties capable of reaction with such functional groups present in either moiety. For example, an amine group of a moiety may react with a carboxyl group of the linker, or an activated derivative thereof, resulting in the formation of an amide linking the two.

35 Examples of moieties capable of reaction with sulfhydryl groups include α -haloacetyl compounds of the type XCH_2CO- (where $X=Br$, Cl , or I), which show particular reactivity for sulfhydryl groups, but which can also be used to modify imidazolyl, thioether, phenol, and amino groups as described by Gurd, *Methods Enzymol.* 11:532 (1967). N-Maleimide derivatives are also considered selective towards sulfhydryl groups, but may additionally be useful in coupling to amino groups under certain conditions. Reagents such as 2-iminothiolane (Traut et al., *Biochemistry* 12:3266 (1973)), which introduce a thiol group through conversion of an amino group, may be considered as sulfhydryl reagents if linking occurs through the formation of disulfide bridges.

40 Examples of reactive moieties capable of reaction with amino groups include, for example, alkylating and acylating agents. Representative alkylating agents include:

(i) α -haloacetyl compounds, which show specificity towards amino groups in the absence of reactive thiol groups and are of the type XCH_2CO^- (where $X=Br$, Cl , or I), for example, as described by Wong *Biochemistry* 24:5337 (1979);

5 (ii) N-maleimide derivatives, which may react with amino groups either through a Michael type reaction or through acylation by addition to the ring carbonyl group, for example, as described by Smyth et al., *J. Am. Chem. Soc.* 82:4600 (1960) and *Biochem. J.* 91:589 (1964);

(iii) aryl halides such as reactive nitrohaloaromatic compounds;

(iv) alkyl halides, as described, for example, by McKenzie et al., *J. Protein Chem.* 7:581 (1988);

10 (v) aldehydes and ketones capable of Schiff's base formation with amino groups, the adducts formed usually being stabilized through reduction to give a stable amine;

(vi) epoxide derivatives such as epichlorohydrin and bisoxiranes, which may react with amino, sulfhydryl, or phenolic hydroxyl groups;

(vii) chlorine-containing derivatives of s-triazines, which are very reactive towards nucleophiles such as amino, sulfhydryl, and hydroxyl groups;

15 (viii) aziridines based on s-triazine compounds detailed above, e.g., as described by Ross, *J. Adv. Cancer Res.* 2:1 (1954), which react with nucleophiles such as amino groups by ring opening;

(ix) squaric acid diethyl esters as described by Tietze, *Chem. Ber.* 124:1215 (1991); and

20 (x) α -haloalkyl ethers, which are more reactive alkylating agents than normal alkyl halides because of the activation caused by the ether oxygen atom, as described by Benneche et al., *Eur. J. Med. Chem.* 28:463 (1993).

Representative amino-reactive acylating agents include:

(i) isocyanates and isothiocyanates, particularly aromatic derivatives, which form stable urea and thiourea derivatives respectively;

(ii) sulfonyl chlorides, which have been described by Herzig et al., *Biopolymers* 2:349 (1964);

25 (iii) acid halides;

(iv) active esters such as nitrophenylesters or N-hydroxysuccinimidyl esters;

(v) acid anhydrides such as mixed, symmetrical, or N-carboxyanhydrides;

(vi) other useful reagents for amide bond formation, for example, as described by M. Bodansky, *Principles of Peptide Synthesis*, Springer-Verlag, 1984;

30 (vii) acylazides, e.g., wherein the azide group is generated from a preformed hydrazide derivative using sodium nitrite, as described by Wetz et al., *Anal. Biochem.* 58:347 (1974);

(viii) imidoesters, which form stable amidines on reaction with amino groups, for example, as described by Hunter and Ludwig, *J. Am. Chem. Soc.* 84:3491 (1962); and

(ix) haloheteroaryl groups such as halopyridine or halopyrimidine.

35 Aldehydes and ketones may be reacted with amines to form Schiff's bases, which may advantageously be stabilized through reductive amination. Alkoxyamino moieties readily react with ketones and aldehydes to produce stable alkoxamines, for example, as described by Webb et al., in *Bioconjugate Chem.* 1:96 (1990).

40 Examples of reactive moieties capable of reaction with carboxyl groups include diazo compounds such as diazoacetate esters and diazoacetamides, which react with high specificity to generate ester groups, for example, as described by Herriot, *Adv. Protein Chem.* 3:169 (1947). Carboxyl modifying

reagents such as carbodiimides, which react through O-acylurea formation followed by amide bond formation, may also be employed.

It will be appreciated that functional groups in either moiety may, if desired, be converted to other functional groups prior to reaction, for example, to confer additional reactivity or selectivity. Examples of 5 methods useful for this purpose include conversion of amines to carboxyls using reagents such as dicarboxylic anhydrides; conversion of amines to thiols using reagents such as N-acetylhomocysteine thiolactone, S-acetylmercaptosuccinic anhydride, 2-iminothiolane, or thiol-containing succinimidyl derivatives; conversion of thiols to carboxyls using reagents such as α -haloacetates; conversion of thiols to amines using reagents such as ethylenimine or 2-bromoethylamine; conversion of carboxyls to amines 10 using reagents such as carbodiimides followed by diamines; and conversion of alcohols to thiols using reagents such as tosyl chloride followed by transesterification with thioacetate and hydrolysis to the thiol 15 using sodium acetate.

So-called zero-length linkers, involving direct covalent joining of a reactive chemical group of one moiety with a reactive chemical group of the other without introducing additional linking material may, if 15 desired, be used in accordance with the invention.

More commonly, however, the linker includes two or more reactive moieties, as described above, 20 connected by a spacer element. The presence of such a spacer permits bifunctional linkers to react with specific functional groups within either moiety, resulting in a covalent linkage between the two. The reactive moieties in a linker may be the same (homobifunctional linker) or different (heterobifunctional linker, or, where several dissimilar reactive moieties are present, heteromultifunctional linker), providing a diversity of potential reagents that may bring about covalent attachment between the two moieties.


Spacer elements in the linker typically consist of linear or branched chains and may include a C₁-₁₀ alkyl, C₂-₁₀ alkenyl, C₂-₁₀ alkynyl, C₂-₆ heterocycl, C₆-₁₂ aryl, C₇-₁₄ alkaryl, C₃-₁₀ alkylheterocycl, C₂-₁₀₀ polyethylene glycol, or C₁-₁₀ heteroalkyl.

25 In some instances, the linker is described by Formula V.

Examples of homobifunctional linkers useful in the preparation of conjugates of the invention include, without limitation, diamines and diols selected from ethylenediamine, propylenediamine and hexamethylenediamine, ethylene glycol, diethylene glycol, propylene glycol, 1,4-butanediol, 1,6-hexanediol, cyclohexanediol, and polycaprolactone diol.

30 In some embodiments, the linker is a bond or a linear chain of up to 10 atoms, independently selected from carbon, nitrogen, oxygen, sulfur or phosphorous atoms, wherein each atom in the chain is optionally substituted with one or more substituents independently selected from alkyl, alkenyl, alkynyl, aryl, heteroaryl, chloro, iodo, bromo, fluoro, hydroxyl, alkoxy, aryloxy, carboxy, amino, alkylamino, dialkylamino, acylamino, carboxamido, cyano, oxo, thio, alkylthio, arylthio, acylthio, alkylsulfonate, 35 arylsulfonate, phosphoryl, and sulfonyl, and wherein any two atoms in the chain may be taken together with the substituents bound thereto to form a ring, wherein the ring may be further substituted and/or fused to one or more optionally substituted carbocyclic, heterocyclic, aryl, or heteroaryl rings.

In some embodiments, a linker has the structure of Formula XIX:



Formula XIX

where A^1 is a bond between the linker and presenter protein binding moiety; A^2 is a bond between the mammalian target interacting moiety and the linker; B^1 , B^2 , B^3 , and B^4 each, independently, is selected from optionally substituted C_{1-2} alkyl, optionally substituted C_{1-3} heteroalkyl, O, S, and NR^N ; R^N is hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted C_{2-4} alkenyl, optionally substituted C_{2-4} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, or optionally substituted C_{1-7} heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; a , b , c , d , e , and f are each, independently, 0 or 1; and D is optionally substituted C_{1-10} alkyl, optionally substituted C_{2-10} alkenyl, optionally substituted C_{2-10} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, optionally substituted $C_{2-C_{10}}$ polyethylene glycol, or optionally substituted C_{1-10} heteroalkyl, or a chemical bond linking $A^1-(B^1)_a-(C^1)_b-(B^2)_c-$ to $-(B^3)_d-(C^2)_e-(B^4)_f-A^2$.

15

Proteins

Presenter Proteins

Presenter proteins can bind a small molecule to form a complex, which can bind to and modulate the activity of a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target protein or a prokaryotic target protein such as a bacterial target protein). In some embodiments, the presenter protein is a mammalian presenter protein (e.g., a human presenter protein). In some embodiments, the presenter protein is a fungal presenter protein. In certain embodiments, the presenter protein is a bacterial presenter protein. In some embodiments, the presenter protein is a plant presenter protein. In some embodiments, the presenter protein is a relatively abundant protein (e.g., the presenter protein is sufficiently abundant that participation in a tripartite complex does not materially negatively impact the biological role of the presenter protein in a cell and/or viability or other attributes of the cell). In some embodiments, the presenter protein is more abundant than the target protein. In certain embodiments, the presenter protein is a protein that has chaperone activity within a cell. In some embodiments, the presenter protein has multiple natural interaction partners within a cell. In certain embodiments, the presenter protein is one which is known to bind a small molecule to form a binary complex that is known to or suspected of binding to and modulating the biological activity of a target protein. Immunophilins are a class of presenter proteins which are known to have these functions and include FKBP and cyclophilins. In some embodiments, a reference presenter protein exhibits peptidyl prolyl isomerase activity; in some embodiments, a presenter protein shows comparable activity to the reference presenter protein. In certain embodiments, the presenter protein is a member of the FKBP family (e.g., FKBP12, FKBP12.6, FKBP13, FKBP19, FKBP22, FKBP23, FKBP25, FKBP36, FKBP38, FKBP51, FKBP52, FKBP60, FKBP65, and FKBP133), a member of the cyclophilin family (e.g., PP1A, CYPB, CYPc, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, PPWD1, PPIAL4A, PPIAL4B, PPIAL4C, PPIAL4D, or PPIAL4G), or PIN1. The "FKBP family" is a family of proteins that have prolyl isomerase activity and function as protein folding chaperones for proteins containing proline residues. Genes that encode proteins in this family include

AIP, AIPL1, FKBP1A, FKBP1B, FKBP2, FKBP3, FKBP4, FKBP5, FKBP6, FKBP7, FKBP8, FKBP9, FKBP9L, FKBP10, FKBP11, FKBP14, FKBP15, and LOC541473.

The “cyclophilin family” is a family of proteins that bind to cyclosporine. Genes that encode proteins in this family include PPIA, PPIB, PPIC, PPID, PPIE, PPIF, PPIG, PPIH, SDCCAG-10, PPIL1, 5 PPIL2, PPIL3, PPIL4, P270, PPWD1, and COAS-2. Exemplary cyclophilins include PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, PPWD1, PPIAL4A, PPIAL4B, PPIAL4C, PPIAL4D, and PPIAL4G.

In some embodiments, a presenter protein is a chaperone protein such as GRP78/BiP, GRP94, GRP170, calnexin, calreticulin, HSP47, ERp29, Protein disulfide isomerase (PDI), and ERp57.

10 In some embodiments, a presenter protein is an allelic variant or splice variant of a FKBP or cyclophilin disclosed herein.

15 In some embodiments, a presenter protein is a polypeptide whose amino acid sequence i) shows significant identity with that of a reference presenter protein; ii) includes a portion that shows significant identity with a corresponding portion of a reference presenter protein; and/or iii) includes at least one characteristic sequence found in presenter protein. In many embodiments, identity is considered “significant” for the purposes of defining an presenter protein if it is above 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher. In some embodiments, the portion showing significant identity has a length of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 20 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 300, 350, 450, 500, 550, 600 amino acids or more.

25 Representative presenter proteins are encoded by the genes or homologs thereof listed in Table 1; in some embodiments, a reference presenter protein is encoded by a gene set forth in Table 1. Also, those of ordinary skill in the art, referring to Table 1, can readily identify sequences that are characteristic of presenter proteins generally, and/or of particular subsets of presenter proteins.

Table 1. Genes that Encode Selected Presenter Proteins

Gene Name	Uniprot Accession Number
AIP	O00170
AIPL1	Q9NZN9
FKBP1A	P62942
FKBP1B	P68106
FKBP2	P26885
FKBP3	Q00688
FKBP4	Q02790
FKBP5	Q13451
FKBP6	O75344
FKBP7	Q9Y680
FKBP8	Q14318

FKBP9	O95302
FKBP9L	Q75LS8
FKBP10	Q96AY3
FKBP11	Q9NYL4
FKBP14	Q9NWM8
FKBP15	Q5T1M5
LOC541473	-
PPIA	Q567Q0
PPIB	P23284
PPIC	P45877
PPID	Q08752
PPIE	Q9UNP9
PPIG	Q13427
PPIH	O43447
PPIL1	Q9Y3C6
PPIL2	Q13356
PPIL3	Q9H2H8
PPIL4	Q8WUA2
PPIL5	Q32Q17
PPIL6	Q8IXY8
PPWD1	Q96BP3

Target Proteins

A target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target protein or a prokaryotic target protein such as a bacterial target protein) is a protein which mediates a disease condition or a symptom of a disease condition. As such, a desirable therapeutic effect can be achieved by modulating (inhibiting or increasing) its activity. Target proteins useful in the complexes and methods of the invention include those which do not naturally associate with a presenter protein, e.g., those which have an affinity for a presenter protein in the absence of a binary complex with a compound of the invention of greater than 1 μ M, preferably greater than 5 μ M, and more preferably greater than 10 μ M. Alternatively, target proteins which do not naturally associate with a presenter protein are those which have an affinity for a compound of the invention in the absence of a binary complex greater than 1 μ M, preferably greater than 5 μ M, and more preferably greater than 10 μ M. In another alternative, target proteins which do not naturally associate with a presenter protein are those which have an affinity for a binary complex of cyclosporine, rapamycin, or FK506 and a presenter protein (e.g., FKBP) of greater than 1 μ M, preferably greater than 5 μ M, and more preferably greater than 10 μ M. In yet another alternative, target proteins that do not naturally associate with a presenter protein are those which are other than calcineurin or mTOR. The selection of suitable target proteins for the complexes and methods of the invention may depend on the presenter protein. For example, target proteins that have low affinity for a cyclophilin may have high affinity for an FKBP and would not be used together with the latter.

Target proteins can be naturally occurring, e.g., wild type. Alternatively, a target protein can vary from the wild type protein but still retain biological function, e.g., as an allelic variant, a splice mutant or a biologically active fragment.

In some embodiments, a target protein is a transmembrane protein. In some embodiments, a target protein has a coiled coil structure. In certain embodiments, a target protein is one protein of a dimeric complex.

In some embodiments, a target protein of the invention includes one or more surface sites (e.g., a flat surface site) characterized in that, in the absence of forming a presenter protein/compound complex, small molecules typically demonstrate low or undetectable binding to the site(s). In some embodiments, a target protein includes one or more surface sites (e.g., a flat surface site) to which, in the absence of forming a presenter protein/compound complex, a particular small molecule (e.g., the compound) shows low or undetectable binding (e.g., binding at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100 fold or more lower than that observed with a presenter protein/compound complex involving the same compound). In some embodiments, a target protein has a surface characterized by one or more sites (and, in some embodiments, an entire surface) that lack(s) any traditional binding pocket, for example, a cavity or pocket on the protein structure with physiochemical and/or geometric properties comparable to proteins whose activity has been modulated by one or more small molecules. In certain embodiments, a target protein has a traditional binding pocket and a site for a protein-protein interaction. In some embodiments, a target protein is an undruggable target, for example, a target protein is not a member of a protein family which is known to be targeted by drugs and/or does not possess a binding site that is expected (e.g., according to art-accepted understanding, as discussed herein) to be suitable for binding to a small molecule. In some embodiments, the protein includes at least one reactive cysteine.

In some embodiments, the target protein is a GTPase such as DIRAS1, DIRAS2, DIRAS3, ERAS, GEM, HRAS, KRAS, MRAS, NKIRAS1, NKIRAS2, NRAS, RALA, RALB, RAP1A, RAP1B, RAP2A, RAP2B, RAP2C, RASD1, RASD2, RASL10A, RASL10B, RASL11A, RASL11B, RASL12, REM1, REM2, RERG, RERGL, RRAD, RRAS, RRAS2, RHOA, RHOB, RHOBTB1, RHOBTB2, RHOBTB3, RHOC, RHOD, RHOF, RHOG, RHOH, RHOJ, RHOQ, RHOU, RHOV, RND1, RND2, RND3, RAC1, RAC2, RAC3, CDC42, RAB1A, RAB1B, RAB2, RAB3A, RAB3B, RAB3C, RAB3D, RAB4A, RAB4B, RAB5A, RAB5B, RAB5C, RAB6A, RAB6B, RAB6C, RAB7A, RAB7B, RAB7L1, RAB8A, RAB8B, RAB9, RAB9B, RABL2A, RABL2B, RABL4, RAB10, RAB11A, RAB11B, RAB12, RAB13, RAB14, RAB15, RAB17, RAB18, RAB19, RAB20, RAB21, RAB22A, RAB23, RAB24, RAB25, RAB26, RAB27A, RAB27B, RAB28, RAB2B, RAB30, RAB31, RAB32, RAB33A, RAB33B, RAB34, RAB35, RAB36, RAB37, RAB38, RAB39, RAB39B, RAB40A, RAB40AL, RAB40B, RAB40C, RAB41, RAB42, RAB43, RAP1A, RAP1B, RAP2A, RAP2B, RAP2C, ARF1, ARF3, ARF4, ARF5, ARF6, ARL1, ARL2, ARL3, ARL4, ARL5, ARL5C, ARL6, ARL7, ARL8, ARL9, ARL10A, ARL10B, ARL10C, ARL11, ARL13A, ARL13B, ARL14, ARL15, ARL16, ARL17, TRIM23, ARL4D, ARFRP1, ARL13B, RAN, RHEB, RHEBL1, RRAD, GEM, REM, REM2, RIT1, RIT2, RHOT1, or RHOT2. In some embodiments, the target protein is a GTPase activating protein such as NF1, IQGAP1, PLEXIN-B1, RASAL1, RASAL2, ARHGAP5, ARHGAP8, ARHGAP12, ARHGAP22, ARHGAP25, BCR, DLC1, DLC2, DLC3, GRAF, RALBP1, RAP1GAP, SIPA1, TSC2, AGAP2, ASAP1, or ASAP3. In some embodiments, the target protein is a Guanine nucleotide-exchange factor such as CNRASGEF, RASGEF1A, RASGRF2, RASGRP1, RASGRP4, SOS1, RALGDS, RGL1, RGL2, RGR, ARHGEF10, ASEF/ARHGEF4, ASEF2, DBS, ECT2, GEF-H1, LARG, NET1, OBSCURIN, P-REX1, P-REX2, PDZ-RHOGEF, TEM4, TIAM1, TRIO, VAV1, VAV2, VAV3, DOCK1, DOCK2, DOCK3, DOCK4, DOCK8, DOCK10, C3G, BIG2/ARFGEF2, EFA6, FBX8, or GEP100. In certain embodiments, the target

protein is a protein with a protein-protein interaction domain such as ARM; BAR; BEACH; BH; BIR; BRCT; BROMO; BTB; C1; C2; CARD; CC; CALM; CH; CHROMO; CUE; DEATH; DED; DEP; DH; EF-hand; EH; ENTH; EVH1; F-box; FERM; FF; FH2; FHA; FYVE; GAT; GEL; GLUE; GRAM; GRIP; GYF; HEAT; HECT; IQ; LRR; MBT; MH1; MH2; MIU; NZF; PAS; PB1; PDZ; PH; POLO-Box; PTB; PUF; 5 PWWP; PX; RGS; RING; SAM; SC; SH2; SH3; SOCS; SPRY; START; SWIRM; TIR; TPR; TRAF; SNARE; TUBBY; TUDOR; UBA; UEV; UIM; VHL; VHS; WD40; WW; SH2; SH3; TRAF; Bromodomain; or TPR. In some embodiments, the target protein is a heat shock protein such as Hsp20, Hsp27, Hsp70, Hsp84, alpha B crystalline, TRAP-1, hsf1, or Hsp90. In certain embodiments, the target protein is an ion channel such as Cav2.2, Cav3.2, IKACH, Kv1.5, TRPA1, Nav1.7, Nav1.8, Nav1.9, P2X3, or P2X4. In 10 some embodiments, the target protein is a coiled-coil protein such as geminin, SPAG4, VAV1, MAD1, ROCK1, RNF31, NEDP1, HCCM, EEA1, Vimentin, ATF4, Nemo, SNAP25, Syntaxin 1a, FYCO1, or CEP250. In certain embodiments, the target protein is a kinase such as Cyclin D1, ABL, ALK, AXL, BTK, EGFR, FMS, FAK, FGFR1, 2, 3, 4, FLT3, HER2/ErbB2, HER3/ErbB3, HER4/ErbB4, IGF1R, INSR, JAK1, JAK2, JAK3, KIT, MET, PDGFRA, PDGFRB, RET RON, ROR1, ROR2, ROS, SRC, SYK, TIE1, TIE2, 15 TRKA, TRKB, KDR, AKT1, AKT2, AKT3, PDK1, PKC, RHO, ROCK1, RSK1, RKS2, RKS3, ATM, ATR, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8, CDK9, CDK10, ERK1, ERK2, ERK3, ERK4, GSK3A, GSK3B, JNK1, JNK2, JNK3, AurA, AurB, PLK1, PLK2, PLK3, PLK4, IKK, KIN1, cRaf, PKN3, c-Src, Fak, PyK2, or AMPK. In some embodiments, the target protein is a phosphatase such as WIP1, SHP2, SHP1, PRL-3, PTP1B, or STEP. In certain embodiments the target protein is a ubiquitin or 20 ubiquitin-like protein (such as NEDD8, ATG8 proteins, SUMO proteins, ISG15), activating enzyme (E1's such as UBA1, UBA2, UBA3, UBA5, UBA6, UBA7, ATG7, NAE1, SAE1), conjugation enzyme (E2's such as UBE proteins, ATG3, BIRC6), ligation enzyme (E3's such as BMI-1, MDM2, NEDD4-1, Beta-TRCP, SKP2, E6AP, CBL-B, or APC/C), and ubiquitin or ubiquitin-like protein protease. In some embodiments, the target protein is a chromatin modifier/remodeler such as a chromatin modifier/remodeler encoded by 25 the gene BRG1, BRM, ATRX, PRDM3, ASH1L, CBP, KAT6A, KAT6B, MLL, NSD1, SETD2, EP300, KAT2A, or CREBBP. In some embodiments, the target protein is a transcription factor such as a transcription factor encoded by the gene EHF, ELF1, ELF3, ELF4, ELF5, ELK1, ELK3, ELK4, ERF, ERG, ETS1, ETV1, ETV2, ETV3, ETV4, ETV5, ETV6, FEV, FLI1, GAVPA, SPDEF, SPI1, SPIC, SPIB, E2F1, E2F2, E2F3, E2F4, E2F7, E2F8, ARNTL, BHLHA15, BHLHB2, BHLHB3, BHLHE22, BHLHE23, 30 BHLHE41, CLOCK, FIGLA, HAS5, HES7, HEY1, HEY2, ID4, MAX, MESP1, MLX, MLXIPL, MNT, MSC, MYF6, NEUROD2, NEUROG2, NHLH1, OLIG1, OLIG2, OLIG3, SREBF2, TCF3, TCF4, TFAP4, TFE3, TFEB, TFEC, USF1, ARF4, ATF7, BATF3, CEBPB, CEBPD, CEBPG, CREB3, CREB3L1, DBP, HLF, JDP2, MAFF, MAFK, NRL, NFE2, NFIL3, TEF, XBP1, PROX1, TEAD1, TEAD3, TEAD4, ONECUT3, ALX3, ALX4, ARX, BARHL2, BARX, BSX, CART1, CDX1, CDX2, DLX1, DLX2, DLX3, DLX4, 35 DLX5, DLX6, DMBX1, DPRX, DRGX, DUXA, EMX1, EMX2, EN1, EN2, ESX1, EVX1, EVX2, GBX1, GBX2, GSC, GSC2, GSX1, GSX2, HESX1, HMX1, HMX2, HMX3, HNF1A, HNF1B, HOMEZ, HOXA1, HOXA10, HOXA13, HOXA2, HOXB13, HOXB2, HOXB3, HOXB5, HOXC10, HOXC11, HOXC12, HOXC13, HOXD11, HOXD12, HOXD13, HOXD8, IRX2, IRX5, ISL2, ISX, LBX2, LHX2, LHX6, LHX9, LMX1A, LMX1B, MEIS1, MEIS2, MEIS3, MEOX1, MEOX2, MIXL1, MNX1, MSX1, MSX2, NKX2-3, 40 NKX2-8, NKX3-1, NKX3-2, NKX6-1, NKX6-2, NOTO, ONECUT1, ONECUT2, OTX1, OTX2, PDX1, PHOX2A, PHOX2B, PITX1, PITX3, PKNOX1, PROP1, PRRX1, PRRX2, RAX, RAXL1, RHOXF1, SHOX,

SHOX2, TGIF1, TGIF2, TGIF2LX, UNCX, VAX1, VAX2, VENTX, VSX1, VSX2, CUX1, CUX2, POU1F1, POU2F1, POU2F2, POU2F3, POU3F1, POU3F2, POU3F3, POU3F4, POU4F1, POU4F2, POU4F3, POU5F1P1, POU6F2, RFX2, RFX3, RFX4, RFX5, TFAP2A, TFAP2B, TFAP2C, GRHL1, TFCP2, NFIA, NFIB, NFIX, GCM1, GCM2, HSF1, HSF2, HSF4, HSFY2, EBF1, IRF3, IRF4, IRF5, IRF7, IRF8, IRF9, 5 MEF2A, MEF2B, MEF2D, SRF, NRF1, CPEB1, GMEB2, MYBL1, MYBL2, SMAD3, CENPB, PAX1, PAX2, PAX9, PAX3, PAX4, PAX5, PAX6, PAX7, BCL6B, EGR1, EGR2, EGR3, EGR4, GLIS1, GLIS2, GLI2, GLIS3, HIC2, HINFP1, KLF13, KLF14, KLF16, MTF1, PRDM1, PRDM4, SCRT1, SCRT2, SNAI2, SP1, SP3, SP4, SP8, YY1, YY2, ZBED1, ZBTB7A, ZBTB7B, ZBTB7C, ZIC1, ZIC3, ZIC4, ZNF143, ZNF232, ZNF238, ZNF282, ZNF306, ZNF410, ZNF435, ZBTB49, ZNF524, ZNF713, ZNF740, ZNF75A, 10 ZNF784, ZSCAN4, CTCF, LEF1, SOX10, SOX14, SOX15, SOX18, SOX2, SOX21, SOX4, SOX7, SOX8, SOX9, SRY, TCF7L1, FOXO3, FOXB1, FOXC1, FOXC2, FOXD2, FOXD3, FOXG1, FOXI1, FOXJ2, FOXJ3, FOXK1, FOXL1, FOXO1, FOXO4, FOXO6, FOXP3, EOMES, MGA, NFAT5, NFATC1, NFkB1, NFkB2, TP63, RUNX2, RUNX3, T, TBR1, TBX1, TBX15, TBX19, TBX2, TBX20, TBX21, TBX4, TBX5, AR, ESR1, ESRRA, ESRRB, ESRRG, HNF4A, NR2C2, NR2E1, NR2F1, NR2F6, NR3C1, NR3C2, 15 NR4A2, RARA, RARB, RARG, RORA, RXRA, RXRB, RXRG, THRA, THRB, VDR, GATA3, GATA4, or GATA5; or C-myc, Max, Stat3, Stat4, Stat6, androgen receptor, C-Jun, C-Fox, N-Myc, L-Myc, MITF, Hif-1alpha, Hif-2alpha, Bcl6, E2F1, NF-kappaB, Stat5, or ER(coact). In certain embodiments, the target protein is TrkA, P2Y14, mPEGS, ASK1, ALK, Bcl-2, BCL-XL, mSIN1, RORyt, IL17RA, eIF4E, TLR7 R, PCSK9, IgE R, CD40, CD40L, Shn-3, TNFR1, TNFR2, IL31RA, OSMR, IL12beta1,2, Tau, FASN, KCTD 6, KCTD 9, Raptor, Rictor, RALGAPA, RALGAPB, Annexin family members, BCOR, NCOR, beta catenin, AAC 11, PLD1, PLD2, Frizzled7, RaLP, ,MLL-1, Myb, Ezh2, RhoGD12, EGFR, CTLA4R, GCGC (coact), Adiponectin R2, GPR 81, IMPDH2, IL-4R, IL-13R, IL-1R, IL2-R, IL-6R, IL-22R, TNF-R, TLR4, MyD88, Keap1, or Nrlp3.

25 *Protein Variants*

A protein or polypeptide variant, as described herein, generally has an amino acid sequence that shows significant (e.g., 80% or more, i.e., 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more) identity with that of a reference polypeptide (e.g., a presenter protein or target protein as described herein such as for example a mammalian 30 presenter protein or target protein) but includes a limited number of particular amino acid changes (e.g., insertions, deletions, or substitutions, either conservative or non-conservative and/or including one or more amino acid variants or analogs (e.g., D-amino acids, desamino acids) relative to the reference polypeptide. In certain embodiments, a variant shares a relevant biological activity (e.g., binding to a particular compound or moiety thereof) with the reference polypeptide; in some such embodiments, the 35 variant displays such activity at a level that is not less than about 50% of that of the reference polypeptide and/or is not less than about 0.5 fold below that of the reference polypeptide.

In some embodiments, a variant polypeptide has an amino acid sequence that differs from that of a reference polypeptide at least (or only) in that the variant has a larger number of cysteine residues and/or has one or more cysteine residues at a position corresponding to a non-cysteine residue in the 40 reference polypeptide. For example, in some embodiments, addition of one or more cysteine residues to

the amino or carboxy terminus of any of a polypeptide (e.g., of a presenter protein and/or of a target protein) as described herein can facilitate conjugation of such polypeptide by, e.g., disulfide bonding. In some embodiments, amino acid substitutions can be conservative (i.e., wherein a residue is replaced by another of the same general type or group) or non-conservative (i.e., wherein a residue is replaced by 5 an amino acid of another type). In some embodiments, a naturally occurring amino acid can be substituted for a non-naturally occurring amino acid (i.e., non-naturally occurring conservative amino acid substitution or a non-naturally occurring non-conservative amino acid substitution), or vice versa.

Polypeptides made synthetically can include substitutions of amino acids not naturally encoded by DNA (e.g., non-naturally occurring or unnatural amino acid). Examples of non-naturally occurring 10 amino acids include D-amino acids, an amino acid having an azide-containing side chain, an amino acid having an acetylaminomethyl group attached to a sulfur atom of a cysteine, a pegylated amino acid, the omega amino acids of the formula $\text{NH}_2(\text{CH}_2)_n\text{COOH}$ wherein n is 2-6, neutral nonpolar amino acids, such as sarcosine, t-butyl alanine, t-butyl glycine, N-methyl isoleucine, and norleucine. Phenylglycine may 15 substitute for Trp, Tyr, or Phe; citrulline and methionine sulfoxide are neutral nonpolar, cysteic acid is acidic, and ornithine is basic. Proline may be substituted with hydroxyproline and retain the conformation conferring properties.

Analogs may be generated by substitutional mutagenesis and retain the structure (e.g., a local 20 structure or global structure) of the original protein. Examples of substitutions identified as "conservative substitutions" are shown in Table 2. If such substitutions result in a change not desired, then other type of substitutions, denominated "exemplary substitutions" in Table 2, or as further described herein in reference to amino acid classes, are introduced and the products screened.

Substantial modifications in function or immunological identity are accomplished by selecting 25 substitutions that differ significantly in their effect on maintaining (a) the structure of the protein backbone in the area of the substitution, for example, as a sheet or helical conformation. (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side chain properties:

- (1) hydrophobic: norleucine, methionine (Met), Alanine (Ala), Valine (Val), Leucine (Leu), Isoleucine (Ile), Histidine (His), Tryptophan (Trp), Tyrosine (Tyr), Phenylalanine (Phe),
- (2) neutral hydrophilic: Cysteine (Cys), Serine (Ser), Threonine (Thr)
- (3) acidic/negatively charged: Aspartic acid (Asp), Glutamic acid (Glu)
- (4) basic: Asparagine (Asn), Glutamine (Gln), Histidine (His), Lysine (Lys), Arginine (Arg)
- (5) residues that influence chain orientation: Glycine (Gly), Proline (Pro);
- (6) aromatic: Tryptophan (Trp), Tyrosine (Tyr), Phenylalanine (Phe), Histidine (His),
- (7) polar: Ser, Thr, Asn, Gln
- (8) basic positively charged: Arg, Lys, His, and;
- (9) charged: Asp, Glu, Arg, Lys, His

Other amino acid substitutions are listed in Table 2.

Table 2. Amino acid substitutions

Original residue	Exemplary substitution	Conservative substitution
Ala (A)	Val, Leu, Ile	Val
Arg (R)	Lys, Gln, Asn	Lys
Asn (N)	Gln, His, Lys, Arg	Gln
Asp (D)	Glu	Glu
Cys (C)	Ser	Ser
Gln (Q)	Asn	Asn
Glu (E)	Asp	Asp
Gly (G)	Pro	Pro
His (H)	Asn, Gln, Lys, Arg	Arg
Ile (I)	Leu, Val, Met, Ala, Phe, norleucine	Leu
Leu (L)	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys (K)	Arg, Gln, Asn	Arg
Met (M)	Leu, Phe, Ile	Leu
Phe (F)	Leu, Val, Ile, Ala	Leu
Pro (P)	Gly	Gly
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser	Phe
Val (V)	Ile, Leu, Met, Phe, Ala, norleucine	Leu

Protein Variants with Altered Reactive Amino Acid Profiles

In some embodiments, a protein or polypeptide variant may include the addition of one or more reactive amino acid residues (e.g., cysteines) to a protein (e.g., at the amino or carboxy terminus of any of the proteins described herein) can facilitate conjugation of these proteins by, e.g., disulfide bonding. In some embodiments, one or more reactive amino acids (e.g., cysteines) may be removed to decrease the number of possible conjugation sites on a protein. Amino acid substitutions can be conservative (i.e., wherein a residue is replaced by another of the same general type or group) or non-conservative (i.e., wherein a residue is replaced by an amino acid of another type). In addition, a naturally occurring amino acid can be substituted for a non-naturally occurring amino acid (i.e., non-naturally occurring conservative amino acid substitution or a non-naturally occurring non-conservative amino acid substitution).

As is known in the art, e.g., as described in Chin, J. W., Expanding and Reprogramming the Genetic Code of Cells and Animals, Annual Review of Biochemistry, Vol. 83: 379-408, unnatural amino acids may be incorporated into proteins made *in vitro*. For example, in one system, UAG amber (stop) codons have been used to incorporate pyrrolysine via an archeal tRNA synthetase and tRNA and these can also be used to incorporate azides and alkynes via feeding. Other side chains on unnatural amino acids that have been demonstrated in the art include cyclopropene, trans-cyclooctene, bicyclo[6.1.0]nonyne-lysine, coumarins, p-azidophenylalanine, N6-[(2-propynloxy)carbonyl]-L-Lysine, bicyclo[6.1.0]non-4-yn-9-ylmethanol (BCN), N-5-norbornene-2-yloxycarbonyl-L-lysine, N-tert-

butyloxycarbonyl-L-lysine, N-2-azidoethyloxycarbonyl-L-lysine, N-L-thiaprolyl-L-lysine, N-D-cysteinyl-L-lysine, N-L-cysteinyl-L-lysine, N6-[(2-propynyloxy)carbonyl]-L-lysine, N6-[(2-azidoethoxy)carbonyl]-L-lysine, benzophenone, 4-(6-methyl-s-tetrazin-3-yl)aminophenylalanine, and cyclooctynes.

5 **Complexes**

In naturally occurring protein-protein interactions, binding events are typically driven largely by hydrophobic residues on flat surface sites of the interacting proteins, in contrast to many small molecule-protein interactions which are driven by interactions between the small molecule in a cavity or pocket on the protein. Commonly, hydrophobic residues on a protein's flat surface site form a hydrophobic hot spot 10 wherein most of the binding interactions between or among interacting proteins are van der Waals interactions. In some situations, a small molecule may provide a "portable hotspot" (or portion thereof) in that it participates in or generates such as a hydrophobic interaction site on a protein (e.g., a presenter proteins) where such does not exist absent the small molecule; aspects of the present disclosure are particularly applicable to such situations. For example, in some embodiments, a compound (and/or a 15 tagged form thereof) as described herein forms a complex with a protein (e.g., a presenter protein/compound complex) and participates in pseudo protein-protein interactions (e.g., forming a tripartite complex with a target protein).

Many mammalian proteins are able to bind to any of a plurality of different partners; in some cases, such alternative binding interactions contribute to biological activity of the proteins. Many of these 20 proteins adapt the inherent variability of the hot spot protein regions to present the same residues in different structural contexts. More specifically, the protein-protein interactions can be mediated by a class of natural products produced by a select group of fungal and bacterial species. These molecules exhibit both a common structural organization and resultant functionality that provides the ability to modulate 25 protein-protein interaction. These molecules contain a presenter protein binding moiety that is highly conserved and a target protein interacting moiety that exhibits a high degree of variability among the different natural products. The presenter protein binding moiety confers specificity for the presenter protein and allows the molecule to bind to the presenter protein to form a complex; the mammalian target protein binding moiety confers specificity for the target protein and allows the binary complex to bind to the target protein, typically modulating (e.g., positively or negatively modulating) its activity. In the 30 present invention, a binary complex (e.g., between a compound and presenter protein or a compound and a target protein) is mimicked by conjugating a presenter protein binding moiety to a target protein or a target protein binding moiety to presenter protein. The resulting conjugates of the invention may then bind to a presenter protein or target protein forming a complex that mimics the tripartite complex. These 35 complexes may be used, e.g., to determine the structure of the interface between the presenter protein and the target protein. Furthermore, by simplifying the formation of the complex, e.g., by conjugated a presenter protein binding moiety to a target protein, the compounds of the invention may be used, e.g., to identify target proteins capable of binding to presenter proteins.

Uses

40 *Identification of Target Proteins*

In some embodiments, the compounds, conjugates, complexes, compositions, and/or methods of the present invention may be useful to identify target proteins capable of forming complexes with presenter proteins (e.g., in the presence of a small molecule). The target proteins may be identified by formation of conjugates including a presenter protein binding moiety conjugated to a target moiety and 5 determining if the conjugate forms a complex with a presenter protein.

Most target proteins known in the art to form ternary complexes with presenter proteins and small molecules were identified fortuitously during determination of the mechanism of action of the small molecule. The present methods allow for the rational identification of target proteins capable of forming complexes with presenter proteins in the presence of small molecules by covalently conjugating a 10 presenter protein binding moiety to the target molecule, allowing formation of a complex prior to identification of a compound capable of binding both the presenter protein and target protein simultaneously.

Screening of small molecules for their ability to facilitate complex formation between the presenter protein and identified target protein could then be carried out to identify potential therapeutics 15 capable of modulating the biological activity of the target protein.

In some embodiments, the compounds of the invention may be used to identify target proteins capable of forming complexes with presenter proteins. For example, target proteins may be identified by combining one or more target proteins with a labeled presenter protein (e.g., labeled with biotin) in the presence of a compound of the invention under conditions suitable to allow for formation of a presenter 20 protein/target protein complex. The target proteins which do not form complexes with presenter proteins may then be removed (e.g., washed out) and the target proteins which form complexes may then be pulled down using the label on the presenter protein and analyzed. In some embodiments, the pulled down target proteins may be analyzed by mass spectrometry to determine their identity.

25 *Compound Design*

In some embodiments, the compounds, conjugates, complexes, compositions, and/or methods of the present invention may be useful for the design of compounds capable of modulating the biological activity of target proteins for use in the treatment of disease.

For example, formation of complexes of presenter proteins and conjugates of the invention can 30 facilitate determination of the structure of the protein-protein interface between a presenter protein and a target protein by crystallization and crystal structure determination of the complex. Once the crystal structure of a complex of the invention is determined, methods known in the art for rational drug design may be used to develop small molecules capable of facilitating complex formation between the presenter 35 protein and the target protein such as computational chemistry methods to build structures *de novo* and/or fragment based drug design using methods such as fragment soaking the crystals of complexes of the invention and determining the resulting structure.

The compounds designed as described above may then be screened to determine their ability to modulate the biological activity of the target protein and modified using medicinal chemistry techniques, as necessary, to produce therapeutically useful compounds.

In some embodiments, the compounds, conjugates, complexes, compositions, and/or methods of the present invention may be useful for identifying compounds capable of modulating the biological activity of target proteins through covalent interaction.

For example, the compounds of the inventions may be screened for their ability to covalently bind

5 to target proteins in the presence and absence of presenter proteins to identify compounds capable of selectively binding to target proteins only in the presence of a presenter protein. These compounds may then be tested for their ability to modulate biological activity of the target protein and modified using medicinal chemistry techniques, as necessary, to produce therapeutically useful compounds.

10 *Determination of Biochemical and/or Biophysical Properties*

In some embodiments, the compounds, conjugates, complexes, compositions, and/or methods of the invention may be useful for determining biochemical and/or biophysical properties of a protein or complex.

For example, the free energy of binding between a conjugate including a presenter protein

15 binding moiety and a target protein and a presenter protein may be determined, e.g., by isothermal titration calorimetry. The K_d of a conjugate including a presenter protein binding moiety and a target protein for a presenter protein may be determined, e.g., by surface plasmon resonance. The K_i , K_{inact} , and/or K_i/K_{inact} for a compound and a presenter protein for a target protein may be determined, e.g., by mass spectrometry.

20

Treatment of diseases or disorders

Compounds, conjugates, and complexes described herein may be useful in the methods of treating diseases or disorders related to the target proteins described herein, and, while not bound by theory, are believed to exert their desirable effects through their ability to modulate (e.g., positively or 25 negatively modulate) the activity of a target protein (e.g., a eukaryotic target protein such as a mammalian target protein or a fungal target protein or a prokaryotic target protein such as a bacterial target protein), through interaction with presenter proteins and the target protein.

Kits

30 In some embodiments, the present invention relates to a kit for conveniently and effectively carrying out the methods in accordance with the present invention. In general, the pharmaceutical pack or kit comprises one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Such kits are especially suited for the delivery of solid oral forms such as tablets or capsules. Such a kit preferably includes a number of unit dosages, and may also include a 35 card having the dosages oriented in the order of their intended use. If desired, for instance if the subject suffers from Alzheimer's disease, a memory aid can be provided, for example in the form of numbers, letters, or other markings or with a calendar insert, designating the days in the treatment schedule in which the dosages can be administered. Alternatively, placebo dosages, or calcium dietary supplements, either in a form similar to or distinct from the dosages of the pharmaceutical compositions, can be 40 included to provide a kit in which a dosage is taken every day. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture,

use or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Pharmaceutical Compositions

5 For use as treatment of human and animal subjects, the compounds and conjugates of the invention can be formulated as pharmaceutical or veterinary compositions. Depending on the subject to be treated, the mode of administration, and the type of treatment desired—e.g., prevention, prophylaxis, or therapy—the compounds are formulated in ways consonant with these parameters. A summary of such techniques is found in *Remington: The Science and Practice of Pharmacy, 21st Edition*, Lippincott 10 Williams & Wilkins, (2005); and *Encyclopedia of Pharmaceutical Technology*, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York, each of which is incorporated herein by reference.

Compounds described herein may be present in amounts totaling 1-95% by weight of the total weight of the composition. The composition may be provided in a dosage form that is suitable for intraarticular, oral, parenteral (e.g., intravenous, intramuscular), rectal, cutaneous, subcutaneous, topical, 15 transdermal, sublingual, nasal, vaginal, intravesicular, intraurethral, intrathecal, epidural, aural, or ocular administration, or by injection, inhalation, or direct contact with the nasal, genitourinary, reproductive or oral mucosa. Thus, the pharmaceutical composition may be in the form of, e.g., tablets, capsules, pills, powders, granulates, suspensions, emulsions, solutions, gels including hydrogels, pastes, ointments, creams, plasters, drenches, osmotic delivery devices, suppositories, enemas, injectables, implants, 20 sprays, preparations suitable for iontophoretic delivery, or aerosols. The compositions may be formulated according to conventional pharmaceutical practice.

In general, for use in treatment, compounds described herein may be used alone, or in combination with one or more other active agents. An example of other pharmaceuticals to combine with the compounds described herein would include pharmaceuticals for the treatment of the same indication. 25 Another example of a potential pharmaceutical to combine with compounds described herein would include pharmaceuticals for the treatment of different yet associated or related symptoms or indications. Depending on the mode of administration, compounds are formulated into suitable compositions to permit facile delivery. Each compound of a combination therapy may be formulated in a variety of ways that are known in the art. For example, the first and second agents of the combination therapy may be formulated 30 together or separately. Desirably, the first and second agents are formulated together for the simultaneous or near simultaneous administration of the agents.

Compounds of the invention may be prepared and used as pharmaceutical compositions comprising an effective amount of a compound described herein and a pharmaceutically acceptable carrier or excipient, as is well known in the art. In some embodiments, a composition includes at least 35 two different pharmaceutically acceptable excipients or carriers.

Formulations may be prepared in a manner suitable for systemic administration or topical or local administration. Systemic formulations include those designed for injection (e.g., intramuscular, intravenous or subcutaneous injection) or may be prepared for transdermal, transmucosal, or oral administration. A formulation generally include diluents as well as, in some cases, adjuvants, buffers, 40 preservatives and the like. Compounds can be administered also in liposomal compositions or as microemulsions.

For injection, formulations can be prepared in conventional forms as liquid solutions or suspensions or as solid forms suitable for solution or suspension in liquid prior to injection or as emulsions. Suitable excipients include, for example, water, saline, dextrose, glycerol and the like. Such compositions may also contain amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, such as, for example, sodium acetate, sorbitan monolaurate, and so forth.

5 Various sustained release systems for drugs have also been devised. See, for example, U.S. patent No. 5,624,677, which is herein incorporated by reference.

10 Systemic administration may also include relatively noninvasive methods such as the use of suppositories, transdermal patches, transmucosal delivery and intranasal administration. Oral administration is also suitable for compounds of the invention. Suitable forms include syrups, capsules, and tablets, as is understood in the art.

15 Each compound of a combination therapy, as described herein, may be formulated in a variety of ways that are known in the art. For example, the first and second agents of the combination therapy may be formulated together or separately.

20 The individually or separately formulated agents can be packaged together as a kit. Non-limiting examples include, but are not limited to, kits that contain, e.g., two pills, a pill and a powder, a suppository and a liquid in a vial, two topical creams, etc. The kit can include optional components that aid in the administration of the unit dose to subjects, such as vials for reconstituting powder forms, syringes for injection, customized IV delivery systems, inhalers, etc. Additionally, the unit dose kit can contain instructions for preparation and administration of the compositions. The kit may be manufactured as a single use unit dose for one subject, multiple uses for a particular subject (at a constant dose or in which the individual compounds may vary in potency as therapy progresses); or the kit may contain multiple doses suitable for administration to multiple subjects ("bulk packaging"). The kit components may be 25 assembled in cartons, blister packs, bottles, tubes, and the like.

30 Formulations for oral use include tablets containing the active ingredient(s) in a mixture with non-toxic pharmaceutically acceptable excipients. These excipients may be, for example, inert diluents or fillers (e.g., sucrose, sorbitol, sugar, mannitol, microcrystalline cellulose, starches including potato starch, calcium carbonate, sodium chloride, lactose, calcium phosphate, calcium sulfate, or sodium phosphate); granulating and disintegrating agents (e.g., cellulose derivatives including microcrystalline cellulose, 35 starches including potato starch, croscarmellose sodium, alginates, or alginic acid); binding agents (e.g., sucrose, glucose, sorbitol, acacia, alginic acid, sodium alginate, gelatin, starch, pregelatinized starch, microcrystalline cellulose, magnesium aluminum silicate, carboxymethylcellulose sodium, methylcellulose, hydroxypropyl methylcellulose, ethylcellulose, polyvinylpyrrolidone, or polyethylene glycol); and lubricating agents, glidants, and antiadhesives (e.g., magnesium stearate, zinc stearate, stearic acid, silicas, hydrogenated vegetable oils, or talc). Other pharmaceutically acceptable excipients can be colorants, flavoring agents, plasticizers, humectants, buffering agents, and the like.

40 Two or more compounds may be mixed together in a tablet, capsule, or other vehicle, or may be partitioned. In one example, the first compound is contained on the inside of the tablet, and the second compound is on the outside, such that a substantial portion of the second compound is released prior to the release of the first compound.

Formulations for oral use may also be provided as chewable tablets, or as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluents (e.g., potato starch, lactose, microcrystalline cellulose, calcium carbonate, calcium phosphate or kaolin), or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin, 5 or olive oil. Powders, granulates, and pellets may be prepared using the ingredients mentioned above under tablets and capsules in a conventional manner using, e.g., a mixer, a fluid bed apparatus or a spray drying equipment.

Dissolution or diffusion controlled release can be achieved by appropriate coating of a tablet, capsule, pellet, or granulate formulation of compounds, or by incorporating the compound into an 10 appropriate matrix. A controlled release coating may include one or more of the coating substances mentioned above and/or, e.g., shellac, beeswax, glycowax, castor wax, carnauba wax, stearyl alcohol, glyceryl monostearate, glyceryl distearate, glycerol palmitostearate, ethylcellulose, acrylic resins, di-polylactic acid, cellulose acetate butyrate, polyvinyl chloride, polyvinyl acetate, vinyl pyrrolidone, polyethylene, polymethacrylate, methylmethacrylate, 2-hydroxymethacrylate, methacrylate hydrogels, 1,3 butylene glycol, ethylene glycol methacrylate, and/or polyethylene glycols. In a controlled release matrix 15 formulation, the matrix material may also include, e.g., hydrated methylcellulose, carnauba wax and stearyl alcohol, carbopol 934, silicone, glyceryl tristearate, methyl acrylate-methyl methacrylate, polyvinyl chloride, polyethylene, and/or halogenated fluorocarbon.

The liquid forms in which the compounds and compositions of the present invention can be 20 incorporated for administration orally include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.


Generally, when administered to a human, the oral dosage of any of the compounds of the 25 combination of the invention depends on the nature of the compound, and can readily be determined by one skilled in the art. Typically, such dosage is normally about 0.001 mg to 2000 mg per day, desirably about 1 mg to 1000 mg per day, and more desirably about 5 mg to 500 mg per day. Dosages up to 200 mg per day may be necessary.

Administration of each drug in a combination therapy, as described herein, can, independently, 30 be one to four times daily for one day to one year, and may even be for the life of the subject. Chronic, long-term administration may be indicated.

EXAMPLES

Example 1: Synthesis of Certain Cross-linking Reagents

Synthesis of acrylamide-containing cyclosporine analogs



All reagents and solvents were purchased from Sinopharm Chemical Reagent Co. Ltd. Fmoc-amino acids, HATU, HOAT, H-Ala-2-Cl-(Trt) resin (0.36 mmol/g), H-Leu-2-Cl-(Trt) resin (0.30 mmol/g), H-Phe-2-Cl-(Trt) resin (0.35 mmol/g) and H-Thr(tBu)-2-Cl-(Trt) resin (0.36 mmol/g) were purchased from GL Biochem (Shanghai) Ltd.

The coupling of linear peptides was carried out using standard Fmoc SPSS procedure on automated synthesizer.

15 **General method A:** The linear peptides were synthesized by using a TETRAS™ synthesizer with a scale of 0.025 mmol of resin. A general protocol is as follows: 2 x NMP, 30 s; 1 x 20% (vol/vol) piperidine in NMP, 15 min; 5 x NMP, 30 s; amino acids (3 eq) solution in NMP was added into the vessel containing resin followed by the addition of a solution of HATU and DIEA in DMF respectively, coupling for 45 min; 3 x NMP, 30 s. A double coupling strategy was applied for all the amino acids.

20

General method B: attaching of Boc-7mer

The coupling was accomplished on TETRAS™ synthesizer by using the same general protocol as for amino acid coupling except that the amount of Boc-7mer is 1.5 equivalents. Only one coupling was needed.

5 **General method C:** Removal of ivDde protecting group.

The removal of ivDde protecting group on Dap side chain was accomplished on TETRAS™ synthesizer. General protocol: A solution of 20% (vol/vol) hydrazine monohydrate in NMP was added in the vessel containing the resin. The vessel was shook for 30 min. The resin was drained and rinsed with 5 x 5 mL (30 s) of NMP.

10

General method D: attaching of acrylic acid on side-chain amino group of Dap.

The coupling was accomplished on TETRAS™ synthesizer by using the same general protocol as for amino acid coupling. A double coupling strategy was applied.

15 **General method E:** Deprotection of side chain protecting groups and final cleavage from resin.

Global deprotection and cleavage from resin was achieved by TFA cocktails for 1-2 hr at room temperature. The cleavage cocktails (TFA/TIPS/H₂O, 95/2.5/2.5) or (TFA/DCM/TIPS, 40:60:1) can be used for the final cleavage. Most solvent was removed under reduced pressure and the residue was concentrated under vacuum to remove the trace solvent. The resulting residue was used in the next cyclization directly without further purification.

20


General method F: Cyclization of linear peptides

The crude linear peptide was dissolved in dry DCM to get the final concentration of 0.1 M. Then HATU (3eq), HOAt (3eq) and DIEA (6eq) was added. The reaction mixture was stirred overnight and then

25 monitored by ESI-LCMS. The solvent was concentrated under reduced pressure and the residue was dissolved in NMP and purified by preparative HPLC. Cyclo-peptides were identified by ESI-LCMS.

30 **Reversed-Phase HPLC.** Accucore C18 column (2.6 µm, 2.1 mm x 50 mm) with a 1 mL/min flow rate was used for analytical RP-HPLC. Xselect Peptide CSH column (5um, 19 mm x 150mm) was used for preparative RP-HPLC. Mobile phase A: water (0.1% formic acid), Mobile phase B: ACN; Flow rate: 20 mL/min; Gradient: 25% B to 95% B in 16 min.


Synthesis of cyclo[7mer-NMALA-Dap-d-nmala-LEU]:



The linear peptide-chain assembly was performed using the general methods described above with 850 mg of H-Leu-2-Cl-(Trt) resin (0.3 mmol/g, 0.025 mmol scale). D-N-methyl Ala, Dap and L-N-


5 methyl Ala residue were attached using general method A. Then Boc-7mer was assembled with general method B. The removal of side-chain ivDde protecting group was accomplished by using general method C. Then acrylic acid was attached on the free amino group on the side chain of Dap with general method D. After global deprotection and cleavage from the resin in one step (general method E), the crude linear peptide was submitted to the cyclization (general method F). After final preparative HPLC, 17 mg of final 10 compound was obtained as a white solid (56.6% calculated by resin loading). ESI-MS: [M+1]⁺=1202, [M+Na]⁺=1224, [M/2 + 1]⁺=602.

Synthesis of cyclo[7mer-NMALA-Dap-d-nmala-PHE]:




15 2.8 mg of final compound was obtained as a white solid (9.1% calculated by resin loading) with the similar method as the synthesis of cyclo[7mer-NMALA-Dap-d-nmala-LEU]. ESI-MS: [M+1]⁺=1236, [M+Na]⁺=1258, [M/2 + 1]⁺=619.

Synthesis of cyclo[7mer-NMALA-Dap-d-nmala-THR]:



3.4 mg of final compound was obtained as a white solid (11.4% calculated by resin loading) with the similar method as the synthesis of cyclo[7mer-NMALA-Dap-d-nmala-LEU]. ESI-MS: $[M+1]^+=1190$,
5 $[M+Na]^+=1212$, $[M/2 + 1]^+=596$.

Synthesis of acrylamide-containing FKBP12 ligands



10 Boc-3mer

All reagents and solvents were purchased from Sinopharm Chemical Reagent Co. Ltd. Fmoc-amino acids, HATU, HOAT, 2-Cl-(Trt)-Cl resin (0.9 mmol/g based on active site) and Ala loaded 2-Cl-(Trt)-Cl resin (0.36 mmol/g) were purchased from GL Biochem (Shanghai) Ltd.

To attach the first amino acid on 2-Cl-(Trt)-Cl resin.

15 In a vessel for SPSS, 5 g of resin was swelled in 50 mL of dry NMP for 40 min. Drain the resin and rinse it with DCM (5 x 30 mL) then with NMM 2%/DCM (3 x 30 mL). A solution of Fmoc-Dap(ivDde)-OH (3.0 mmol, 1.6g) with NMM (4 mmol, 400 mg) in 50 mL of DCM was added. The vessel was shook overnight. Then 2 mL of a solution of 25% NMM/MeOH was added and the vessel was shook for one more hour. The resin was drained and rinsed with DCM, NMP, MeOH and EtOH (3 x 50 mL each). The
20 resin was dried under vacuum at room temperature.

The loading was measured by Fmoc cleavage photometric measurement (0.32 mmol/g).

The coupling of linear peptides was carried out using standard Fmoc SPSS procedure on an automated synthesizer.

5 **General method A:** The linear peptides were synthesized by using a TETRAS™ synthesizer with a scale of 0.025 mmol of resin. A general protocol as follows: 2 x NMP, 30 s; 1 x 20% (vol/vol) piperidine in NMP, 15 min; 5 x NMP, 30 s; amino acids (3 eq) solution in NMP was added into the vessel containing resin followed by the addition of a solution of HATU and DIEA in DMF respectively, coupling for 45 min; 3 x NMP, 30 s. A double coupling strategy was applied for all the amino acids.

10

General method B: attaching of Boc-3mer

The coupling was accomplished on TETRAS™ synthesizer by using the same general protocol as for amino acids coupling except that the amount of Boc-3mer is 1.5 equivalents. Only once coupling was needed.

15

General method C: Removal of ivDde protecting group.

The removal of ivDde protecting group on Dap side chain was accomplished on TETRAS™ synthesizer. General protocol: A solution of 20% (vol/vol) hydrazine monohydrate in NMP was added in the vessel containing the resin. The vessel was shook for 30 min. The resin was drained and rinsed with 5 x 5 mL (30 s) of NMP.

20

General method D: attaching of acrylic acid on side-chain amino group of Dap.

The coupling was accomplished on TETRAS™ synthesizer by using the same general protocol as for amino acids coupling. A double coupling strategy was applied.

25

General method E: Deprotection of side chain protecting groups and final cleavage from resin.

Global deprotection and cleavage from resin was achieved by TFA cocktails for 1-2 hr at room temperature. The cleavage cocktails (TFA/TIPS/H₂O, 95/2.5/2.5) or (TFA/DCM/TIPS, 40:60:1) can be used for the final cleavage. Most solvent was removed under reduced pressure and the residue was concentrated under vacuum to remove the trace solvent. The resulting residue was used in the next cyclization directly without further purification.


30

General method F: cyclization of linear peptides

The crude linear peptide was dissolved in dry DCM to get the final concentration of 0.1 M. Then HATU (3eq), HOAt (3eq) and DIEA (6eq) was added. The reaction mixture was stirred overnight and then monitored by ESI-LCMS. The solvent was concentrated under reduced pressure and the residue was dissolved in NMP and purified by preparative HPLC. Cyclo-peptides were identified by ESI-LCMS.


Reversed-Phase HPLC. Accucore C18 column (2.6 μ m, 2.1 mm x 50 mm) with a 1 mL/min flow rate was used for analytical RP-HPLC. Xselect Peptide CSH column (5um, 19 mm x 150mm) was used for preparative RP-HPLC. Mobile phase A: water (0.1% formic acid), Mobile phase B: ACN; Flow rate: 20 mL/min; Gradient: 25% B to 95% B in 16 min.

Synthesis of cyclo[diamine-Dap-ALA-ALA]:



The linear peptide-chain assembly was performed using the general methods described above with 700 mg of H-Ala-2-Cl-(Trt) resin (0.025 mmol scale). Ala and Dap residue was attached by using general method A. Then Boc-3mer was assembled with general method B. The removal of side-chain ivDde protecting group was accomplished by using general method C. Then acrylic acid was attached on the free amino group on the side chain of Dap with general method D. After global deprotection and cleavage from the resin in one step (general method E), the crude linear peptide was submitted to the cyclization (general method F). After final preparative HPLC, 3.1 mg of final compound was obtained as a white solid (14.5% calculated by resin loading). ESI-MS: $[M+1]^+=857$, $[M+Na]^+=879$.


Synthesis of cyclo[diamine-ALA-ALA-Dap]



2.2 mg of final compound was obtained as a white solid (10.3% calculated by resin loading) with the similar method as the synthesis of cyclo[diamine-Dap-ALA-ALA]. ESI-MS: $[M+1]^+=857$, $[M+Na]^+=879$.


5

Synthesis of cyclo[diamine-Dap-ALA]:




2.7 mg of final compound was obtained as a white solid (12.6% calculated by resin loading) with the similar method as the synthesis of cyclo[diamine-Dap-ALA-ALA]. ESI-MS: $[M+1]^+=786$.

Synthesis of sanglefehrin analogs:



Method A may be used to prepare compounds of Formula IV as shown below in Scheme 1.

15 Scheme 1




where R^7 , R^8 , Z^5 , and Z^6 are as previously defined, PG^1 is a suitable amine protecting group including but not limited to Boc, Cbz, Alloc, and Fmoc, and X is either OH, Cl, F, or some other group suitable for displacement or activation followed by displacement.

5 In a typical procedure, protected amine IV is reacted with an appropriate reagent familiar to those skilled in the art to remove the protecting group PG^1 . The isolated crude product may be reacted with acylating agent $Z^5C(O)X$ in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBROP, HATU, HBTU, COMU) known to those skilled in the art. The acid and amine coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, 10 dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or slightly elevated temperature.

15 Alternatively, the deprotected amine may be directly reacted with the acylating reagent $Z^5C(O)X$ when X is a halogen in a suitable solvent (including but not limited to DMF, dichloromethane, DME, acetonitrile, and tetrahydrofuran) in the presence of base (including but not limited to pyridine, DIPEA, triethylamine, and NMM) in the temperature range of -78 °C to about 120 °C, preferably between -20 °C and 50 °C.

Compounds of Formula IVb of Scheme 1 may be prepared as shown below in Scheme 2.

Scheme 2




where R⁷, R⁸, and Z⁶ are as previously defined and PG¹ and PG² are each independently suitable amine protecting groups including but not limited to Boc, Cbz, Alloc, and Fmoc.

In a typical procedure, protected amine IVc is reacted with the appropriate reagent familiar to those skilled in the art to remove the protecting group PG² to produce the corresponding amine, which is then reacted with the appropriate amino acid in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBOP, HATU, HBTU, COMU) known to those skilled in the art. The acid and amine coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or slightly elevated temperature to deliver compounds of Formula IVb.

Compounds of Formula IVc of Scheme 2 may be prepared as shown below in Scheme 3.

Scheme 3




where R⁷ and Z⁶ are as previously defined and PG² is a suitable amine protecting group including but not limited to Boc, Cbz, Alloc, and Fmoc.

In a typical procedure, protected amine IVd is reacted with the piperazic acid derivative IVe in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBOP, HATU, HBTU, COMU) known to those skilled in the art. The acid and amine coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or slightly elevated temperature.

Alternatively, compounds of Formula IVb may be synthesized from compounds of formula II-B as described in Scheme 4.

Scheme 4




where R⁷, R⁸, and Z⁶ are as previously defined, PG¹ is a suitable amine protecting group including but not limited to Boc, Cbz, Alloc, and Fmoc, and PG³ is an alkyl or aryl group included but not limited to methyl, ethyl, benzyl, allyl, phenyl, and the like that can be removed by methods known to those skilled in the art.

In a typical procedure, compound IVe is reacted at room temperature under hydrolysis conditions using base (e.g., lithium hydroxide, sodium hydroxide, sodium carbonate) in a solvent system comprised of organic (e.g., methanol, THF, dioxane) with or without water. It will be appreciated by those skilled in the art that, depending on the identity of PG³, removal of PG³ can also occur under hydrogenolysis

conditions using an appropriate catalyst, or under de-allylation conditions using a palladium catalyst, for example $\text{Pd}(\text{PPh}_3)_4$ and a basic scavenger (e.g., piperidine, morpholine, piperidine) Following deprotection to give the resulting carboxylic acid, Z^6 may be introduced in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBROP, HATU, HBTU, COMU) known to those skilled in the art. The carboxylic acid and the coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or slightly elevated temperature to give product IVb.

Alternatively, compounds of Formula IVb may be synthesized as shown below in Scheme 5.


10 Scheme 5



where R^7 , R^8 , and Z^6 are as previously defined and PG^1 is a suitable amine protecting group including but not limited to Boc, Cbz, Alloc, and Fmoc.

In a typical procedure, protected amine VI is reacted with reagent V in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBROP, HATU, HBTU, COMU) known to those skilled in the art. The acid and amine coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, NMM) at room temperature or slightly elevated temperature.

Synthesis of Intermediate I-8



To a room temperature solution of **Intermediate I-1** (2.00 g, 7.11 mmol) in DMF (13 mL) was added cesium carbonate (4.75 g, 14.58 mmol) and benzyl bromide (2.49 g, 14.58 mmol, 1.73 mL) at 25 5 °C. The reaction mixture was stirred for two hours and then diluted with ethyl acetate (100 mL) and washed with brine (50 mL x 3). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a crude residue. The crude product was purified by column chromatography with silica gel (eluent: petroleum ether/ethyl acetate → 20:1 to 10:1) to afford **Intermediate I-2** (3.20 g, 96% yield) as colorless oil. ¹H NMR (400MHz, CDCl₃) δ 7.45 - 7.30 (m, 10 H), 7.20 - 7.10 (m, 1 H), 6.95 - 6.85 (m, 1 H), 6.74 (s, 1 H), 6.70 - 6.60 (m, 1 H), 5.20 - 5.10 (m, 2 H), 4.99 (s, 2 H), 4.70 - 4.60 (m, 1 H), 3.15 - 3.05 (m, 2 H), 1.43 (s, 9 H). ESI-MS m/z = 484.1 [M+Na]⁺; Calculated MW: 461.55.

To a solution of **Intermediate I-2** (3.20 g, 6.93 mmol) in tetrahydrofuran (15 mL) was added lithium hydroxide (1 M in water, 10 mL) at 0°C. The reaction mixture was stirred at room 15 temperature for 1 hour. The reaction mixture was adjusted to pH~6 with HCl (1 M in water) at 0°C and extracted with ethyl acetate (100 mL x 3). The combined organic layer was washed with brine (20 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure to give a crude residue. The crude product was dissolved in aqueous sodium bicarbonate (7 mL) and extracted with MTBE (100 mL x 3). The aqueous layer was adjusted to pH~6 with hydrochloric acid (1 M in H₂O) and extracted with ethyl 20 acetate (50 mL x 3). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to afford **Intermediate I-3** (2.27 g, 88% yield) as a colorless oil. ¹H NMR (400MHz, CDCl₃) δ 7.45 - 7.30 (m, 5 H), 7.25 - 7.18 (m, 1 H), 6.90 - 6.85 (m, 1 H), 6.83 - 6.75 (m, 2 H), 5.05 (s, 2 H), 4.94 (d, J=8.00 Hz, 1H), 4.60 - 4.50 (m, 1 H), 3.20 - 3.12 (m, 1 H), 3.10 - 3.00 (m, 1 H), 1.43 (s, 9 H). ESI-MS m/z = 394.3 [M+Na]⁺; Calculated MW: 371.43.

25 To a solution of **Intermediate I-3** (888 mg, 2.39 mmol) in dichloromethane (15 mL) was added N-methyl morpholine (967 mg, 9.56 mmol), HOBr (65 mg, 478 umol), (S)-methyl hexahydropyridazine-3-

carboxylate as the TFA salt (890 mg, 2.39 mmol), and EDCI (917 mg, 4.78 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 1 hour. The reaction mixture was diluted with dichloromethane (50 mL) and adjusted to pH~6 with 5% aqueous citric acid. The aqueous layer was extracted with dichloromethane (20 mL x 2). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a crude product. The crude product was purified by column chromatography with silica gel (eluent: petroleum ether/ethyl acetate → 10:1, 5:1 to 3:1) to afford **Intermediate I-4** (910 mg, 77% yield) as colorless oil. ¹HNMR (400MHz, CDCl₃) δ 7.50 - 7.35 (m, 5 H), 7.20 - 7.13 (m, 1 H), 6.90 - 6.80 (m, 3 H), 5.60 - 5.50 (m, 1 H), 5.30 - 5.20 (m, 1 H), 5.05 - 5.00 (m, 2 H), 4.40 - 4.30 (m, 1 H), 3.65 (s, 3 H), 3.55 (d, *J*=11.20 Hz, 1H), 3.00 - 2.93 (m, 1 H), 2.90 - 2.80 (m, 1 H), 2.75 - 2.65 (m, 1 H), 2.35 - 2.25 (m, 1 H), 1.80 - 1.72 (m, 2 H), 1.42 (s, 9 H). ESI-MS m/z = 520.1 [M+Na]⁺. Calculated MW: 497.58.

To a solution **Intermediate I-4** (450 mg, 904 umol) in ethyl acetate (4 mL) was added hydrochloric acid in ethyl acetate (4 M, 8.00 mL) at 0 °C. The reaction mixture was stirred at 25 °C for 1 hour. The reaction mixture was concentrated under reduced pressure to afford the HCl salt of **Intermediate I-5** (390 mg, 100% yield) as light yellow solid and used for next step without purification. ESI-MS m/z = 398.0 [M+H]⁺, 420.0 [M+Na]⁺, Calculated MW: 397.47.


To a solution of **Intermediate I-5** (195 mg, 899 umol) in dichloromethane (5.00 mL) was added *N*-methylmorpholine (273 mg, 2.70 mmol), HOEt (24.29 mg, 179.75 umol), the HCl salt of (S)-methyl 1-((S)-2-amino-3-(3-(benzyloxy)phenyl)propanoyl)hexahydropyridazine-3-carboxylate (390 mg, 899 umol) and EDCI (345 mg, 1.80 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 1 hour. The reaction mixture was diluted with dichloromethane (20 mL) and adjusted to pH~6 with 5% aqueous citric acid. The aqueous layer was extracted with dichloromethane (20 mL x 2). The combined organic layers were washed with brine (20 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a crude product. The crude product was purified by column chromatography with silica gel (eluent: petroleum ether/ethyl acetate → 5:1, 2:1, 1:1) to afford **Intermediate I-6** (750 mg, 70% yield) as a white solid. ¹H NMR (400MHz, CDCl₃) δ 7.50 - 7.35 (m, 5 H), 7.23 - 7.15 (m, 1 H), 6.90 - 6.80 (m, 3 H), 6.13 - 6.08 (m, 1 H), 5.83 - 5.73 (m, 1 H), 5.10 - 5.00 (m, 3 H), 4.30 - 4.20 (m, 1 H), 4.00 - 3.90 (m, 1 H), 3.65 (s, 3 H), 3.50 (d, *J*=11.20 Hz, 1H), 3.05 - 2.97 (m, 1 H), 2.93 - 2.83 (m, 1 H), 2.80 - 2.70 (m, 1 H), 2.35 - 2.25 (m, 1 H), 2.15 - 2.10 (m, 1 H), 1.75 - 1.65 (m, 4 H), 1.45 (s, 9 H), 0.94 (d, *J*=6.80 Hz, 3H), 0.88 (d, *J*=6.80 Hz, 3H). ESI-MS m/z = 597.1 [M+H]⁺. Calculated MW: 596.71.

To a solution of **Intermediate I-6** (3.00g, 6.03 mmol) in methanol (300 mL) was added palladium on carbon (2 grams, 10% loading) under a nitrogen atmosphere. The suspension was degassed and purged with hydrogen gas and the mixture was stirred under hydrogen (1 atm) at 20 °C for 3 hours. The mixture was filtered and concentrated under reduced pressure to give **Intermediate I-7** (2.3 g, 5.64 mmol, 94% yield) as a white solid. ESI-MS m/z = 430.1 [M+Na]⁺. Calculated MW: 407.46

To a mixture of **Intermediate I-7** (2.3 g, 5.64 mmol) in dioxane (10 mL) was added hydrochloric acid in dioxane (4 M, 30 mL) in one portion at 20 °C. The mixture was stirred at 20 °C for 3 hours. The mixture was adjusted to pH ~7-8 with saturated aqueous NaHCO₃ and extracted with ethyl acetate (100 mL x 3). The combined organic phase was washed with brine (100 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give **Intermediate I-8** (1.3 g, 3.63 mmol, 64% yield, 86% purity) as a light yellow solid. ¹H NMR (400MHz, CDCl₃) δ 7.15-7.02 (m, 1 H), 6.75-6.5 (m, 3 H), 4.90-

4.77 (m, 1 H), 4.50-4.37 (m, 1 H), 4.00-3.90 (m, 1 H), 3.80-3.70 (m, 1 H), 3.68-3.65 (m, 3 H), 2.95-2.80 (m, 1 H), 2.77-2.62 (m, 2 H), 2.50-2.35 (m, 1 H), 1.97-1.85 (m, 1 H), 1.82-1.70 (m, 1 H), 1.60 – 1.45 (m, 1 H), 1.42-1.28 (m, 1 H). ESI-MS m/z = 308.2 [M+Na]⁺. Calculated MW: 307.34.


5 Synthesis of Intermediate I-11



To a solution of 2-(dimethylamino)ethanethiol **I-9** (500 mg, 3.53 mmol) in methanol (10 mL) was added a solution of 1,2-di(pyridin-2-yl)disulfide (1.17 g, 5.3 mmol) in methanol (10 mL) at 0 °C. The mixture was stirred at 25 °C for 15 hours. The reaction mixture was concentrated *in vacuo*. The residue 10 was purified by silica gel column (eluent: petroleum ether/ethyl acetate → 3:1, 1:1, then dichloromethane/ethyl acetate → 2:1, then dichloromethane/methanol → 10:1) which was combined with a previous batch to afford **Intermediate I-10** (1.05 g, 58% yield) as light yellow solid. ¹H NMR (400MHz, CD₃OD) δ 8.56 (d, *J*=4.0 Hz, 1 H), 7.85-7.75 (m, 1 H), 7.69 (d, *J*=8.0 Hz, 1 H), 7.36-7.26 (m, 1 H), 3.48-3.40 (m, 2 H), 3.28-3.20 (m, 2 H), 2.93 (s, 6 H). ESI-MS m/z = 214.9 [M+H]⁺. Calculated MW: 214.35.

To a solution of 4-mercaptobutanoic acid (50 mg, 416 umol) in methanol (1 mL) was added a solution of **Intermediate I-10** (125 mg, 499 umol) in methanol (1.5 mL) at 25 °C. The mixture was stirred at 25 °C for 15 hours. The reaction mixture was concentrated at low pressure. The residue was purified by column chromatography with silica gel (eluent: petroleum ether/ethyl acetate → 2:1, 1:1, then dichloromethane/methanol → 10:1) to afford **Intermediate I-11** (80 mg, 86% yield) as a white gum. ¹H NMR (400MHz, CD₃OD) δ 3.55-3.45 (m, 2 H), 3.08-3.00 (m, 2 H), 2.93 (s, 6H), 2.83 (t, *J*=7.2 Hz, 2 H), 2.43 (t, *J*=7.2 Hz, 2 H), 2.05-1.94 (m, 2 H).

Synthesis of Intermediate I-15




To a solution of tert-butyl 2-mercaptoacetate **I-12** (800 mg, 5.4 mmol) in *N,N*-dimethylformamide (15 mL) was added potassium carbonate (1.49 g, 10.8 mmol) and 1-bromo-2-chloroethane (2.32 g, 16.2 mmol). The mixture was stirred at 25 °C for 2 hours. The mixture was diluted with ethyl acetate (100 mL) and washed with water (50 mL x 3). The organic layer was dried over anhydrous sodium sulfate and concentrated to give **Intermediate I-13** (1.00 g, 79% yield) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 3.64 - 3.71 (m, 2 H), 3.14 - 3.17 (m, 2 H), 2.95 - 3.01 (m, 2 H), 1.47 (s, 9 H).

To a solution of **Intermediate I-13** (1.00 g, 4.75 mmol) in dichloromethane (10 mL) was added a solution of *m*-CPBA (4.61 g, 21.4 mmol, 80% purity) in dichloromethane (10 mL) at 0 °C. The mixture was warmed to room temperature stirred for 2 hours. The mixture was diluted with dichloromethane (80 mL) and washed with saturated aqueous sodium sulfite (50 mL) and saturated aqueous sodium bicarbonate (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated to give a crude residue that was purified by silica gel chromatography (eluent: petroleum ether/ethyl acetate → 5/1) to give **Intermediate I-14** (700 mg, 60% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 3.99 (s, 2 H), 3.90 - 3.95 (m, 2 H), 3.69 - 3.75 (m, 2 H), 1.50 - 1.53 (m, 9 H).

To a solution of **Intermediate I-14** (650 mg, 2.68 mmol) in dichloromethane (12 mL) was added trifluoroacetic acid (6.00 mL). The mixture was stirred at 45 °C for 2 hours. Then the mixture was concentrated to give **Intermediate I-15** (360.00 mg, 72% yield) as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 4.34 (s, 2 H), 3.89 - 4.01 (m, 2 H), 3.71 - 3.81 (m, 2 H).


Synthesis of Compound-1



To a solution of the HCl salt of **Intermediate I-8** (31 mg, 119 umol) and **Intermediate I-11** (44 mg, 99 umol) in dichloromethane (1.5 mL) was added HOBr (1.34 mg, 9.9 umol), *N*-methylmorpholine (38.2 μ L) and EDCI (27 mg, 139 umol). The mixture was stirred at 25 °C for one hour. The reaction mixture was diluted with dichloromethane (10 mL) and washed with water (5 mL). The aqueous layer was extracted with dichloromethane (5 mL x 2). The combined organic layers were dried over anhydrous


sodium sulfate, filtered, and concentrated. The residue was purified by reverse phase HPLC (HCl additive) and repurified using reverse phase HPLC (HCOOH additive) and lyophilized to afford **Compound-1** (5.8 mg, 9% yield) as a light yellow oil. ^1H NMR (400MHz, MeOD) δ 8.53 (br.s., 1 H), 7.12-7.04 (m, 1 H), 6.72-6.62 (m, 3H), 5.75 -5.65 (m, 1 H), 4.20-4.10 (m, 2H), 3.70 (s, 3H), 3.08- 3.00 (m, 2H), 2.95-2.76 (m, 5 H), 2.76-2.70 (m, 2 H), 2.60 (s, 6 H), 2.42-2.32 (m, 3 H), 2.10-1.98 (m, 3 H), 1.85-1.70 (m, 2 H), 1.55-1.40 (m, 2 H), 1.02-0.85 (m, 6 H). ESI-MS m/z = 612.1 [M+H] $^+$, 634.5 [M+Na] $^+$. Calculated MW: 611.82.

Synthesis of Compound-2 (SFAC4DS)



A solution of **Intermediate I-8** (27 mg, 60 umol) in dichloromethane was treated with *N,N*-diisopropylethylamine (24 mg, 180 umol), **Intermediate I-16** (14 mg, 60 umol), HOBr (1.6 mg, 2 umol) and then finally EDCI (18mg, 90 umol). After stirring for 15 hours, the solution was poured into water and ethyl acetate. The layers were separated and the aqueous layer was extracted with ethyl acetate. The organics were dried over magnesium sulfate, filtered, and the solvent removed was in *vacuo*. Purification by reverse phase HPLC (acetonitrile in water with 0.1% formic acid) and lyophilization provided **Compound-2 (SFAC4DS)** (16mg, 42% yield) as a white solid. ^1H NMR (400MHz, CDCl_3) 8.47 (d, 1H), 8.16 (br s, 1H), 7.69-7.66 (m, 1H), 7.63-7.60 (m, 1H), 7.13 (app t, 1H), 7.09-7.05 (m, 1H), 5.85-5.79 (m, 1H), 4.72 (m, 1H), 4.31 (br s, 1H), 3.70 (s, 3H), 3.42 (d, 1H), 3.00 (dd, 1H), 2.87-2.78 (m, 3H), 2.52-2.43 (m, 2H), 2.28 (br s, 1H), 2.14 -2.04 (m, 3H), 1.83-1.73 (m, 2H), 1.62-1.43 (m, 5H), 0.95 (d, 3H), 0.90 (d, 3H). ESI-MS m/z = 617.9 $[\text{M}+\text{H}]^+$. Calculated MW: 617.78.


Synthesis of Compound-3



25 To a solution of the HCl salt of **Intermediate I-15** (31.8 mg, 0.17 mmol), HOBt (2.3 mg, 0.017 mmol) and NMM (54 μ L, 0.54 mmol) in DCM (0.75 mL) was added the HCl salt of **I-8** (86mg, 0.17 mmol,) and EDCI (76 mg, 0.34 mmol) at room temperature. The reaction mixture was stirred at room temperature for 2 hours. The mixture was diluted with dichloromethane (2 mL), washed with citric acid (pH~3, 2 mL),


saturated aqueous sodium bicarbonate (2 mL), and saturated aqueous sodium chloride (2 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure at 15 °C to give a product as yellow oil. The residue was purified by prep-TLC with silica gel (eluent: DCM/MeOH → 10:1) followed by preparative HPLC (column: Phenomenex Gemini C18 250 x 50 10u; mobile phase: [water (0.225%FA)-5 ACN]; B%: 26%-56%, 11.2 min) to afford **Compound-3** (8.5 mg, 8% yield) as white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.02 (s, 1 H), 8.80 (s, 1 H), 8.42 (s, 1 H), 7.01 - 7.08 (m, 1 H), 6.82 (dd, *J*=16.3, 9.9 Hz, 1 H), 6.74 (d, *J*=7.5 Hz, 1 H), 6.64 - 6.70 (m, 2 H), 6.36 (d, *J*=16.5 Hz, 1 H), 6.12 – 6.16 (m, 2 H), 4.81 – 4.85 (m, 1 H), 4.40 – 4.46 (m, 2 H), 4.21 – 4.25 (m, 1 H), 4.09 - 4.13 (m, 1 H), 3.57 (s, 3 H), 2.81 - 3.08 (m, 2 H), 2.63 – 2.65 (m, 1 H), 2.15 – 2.19 (m, 1 H), 1.21 - 1.60 (m, 4 H), 0.88 – 0.92 (m, 6 H). ESI-MS 10 m/z = 539.1 [M+H]⁺. Calculated MW: 538.61.

Synthesis of Compound-4




Compound-4 was prepared as described in the preparation of **Compound-3** by using the HCl salt of **Intermediate I-8** (34 mg, 77 umol) and **2-(3-methyl-2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)acetic acid I-17** (13 mg, 77 umol) as starting materials. The resulting crude product was combined with previously synthesized crude material and purified to give **Compound-4** (29.0 mg, 51.28 umol, 45% yield) as a white solid following lyophilization. ¹H NMR (400MHz, CDCl₃) δ 8.55-8.40 (s, 1 H), 8.20-8.07 (m, 1 H), 7.10-6.97 (m, 2 H), 6.80-6.73 (m, 1 H), 6.73-6.63 (m, 1H), 6.63-6.50 (m, 1 H), 6.47-6.37 (m, 1 H), 5.90-5.75 (m, 1 H), 4.80-4.67 (m, 1 H), 4.35-4.15 (m, 3 H), 3.70-3.57 (m, 3 H), 3.45-3.30 (d, 1 H), 3.10-3.00 (m, 1 H), 3.00-2.70 (m, 2 H), 2.20-2.00 (m, 5 H), 1.85-1.60 (m, 2 H), 1.60-1.45 (m, 1 H), 1.45-1.30 (m, 1 H), 1.05-0.80 (m, 6 H). ESI-MS m/z = 558.3 [M+H]⁺; Calculated MW: 557.60.

Synthesis of Compound-5




Compound-5 was prepared as described in the preparation of **Compound-3** by using the HCl salt of **Intermediate I-8** (34 mg, 77 umol) and **2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)acetic acid I-18** (13 mg, 77 umol) as starting materials to give **Compound-5**. ESI-MS m/z = 544.0 [M+H]⁺; Calculated MW: 543.58.


Synthesis of methyl (S)-1-((S)-3-(3-hydroxyphenyl)-2-((S)-3-methyl-2-(3-(2-(pyridin-2-yl)disulfanyl)ethoxy)propanamido)butanamido)propanoyl)hexahydropyridazine-3-carboxylate (SFAX6):



Carboxylic acid **2** (70mg, 0.270mmol) and HBTU (204mg, 0.540mmol, 2.00eq) were mixed in 5 3mL of acetonitrile, and the resulting suspension was stirred at room temperature for 15min. Following this period, amine **1** (Intermediate I-8) (110mg, 0.270mmol, 1.00eq) was added followed by triethylamine (113µL, 0.810mmol, 3.00eq) and the mixture was stirred at room temperature for 18h. The mixture was then treated with 20mL of saturated sodium bicarbonate and extracted with 2x30mL portions of ethyl acetate. The pooled organic extracts were washed with 2x20mL portions of brine, dried over saturated 10 sodium sulfate, filtered and concentrated under vacuum. The residue was purified using silica gel chromatography, eluting with dichloromethane:MeOH, 100 : 1 to 50 : 1, affording 70mg (40%) of the product as a colorless oil. R_f =0.31 (dichloromethane:MeOH, 20 : 1). MS (ESI) calc=648.2 (M+H), obs = 648.2.

Synthesis of SFAX9DS:

SFAX9DS was synthesized according to a similar procedure described above for the synthesis of SFAX6.


5 **Synthesis of Compound-6**

Compound-6 was prepared starting with **Intermediate I-19** to give **Compound-6**. ESI-MS m/z = 631.0 $[M+H]^+$; Calculated MW: 630.82.

10 **Synthesis of Compounds of Formula IVf**

Synthesis of Compounds of Formula IVf may be synthesized as shown below in Scheme 6.


Scheme 6



where R^7 , R^8 , and Z^6 are as previously defined, PG^1 is a suitable amine protecting group including but not limited to Boc, Cbz, Alloc, and Fmoc, and PG^4 is an alkyl or aryl group included but not limited to methyl, ethyl, benzyl, allyl, and phenyl that can be removed by methods known to those skilled in the art.


In a typical procedure, compound IVd is reacted with reagent IVg in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBOP, HATU, HBTU, COMU) familiar to those skilled in the art. The acid and amine coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature

or slightly elevated temperature. Following amide formation, the protecting group PG⁴ is then removed using conditions described for removing of PG³ of compound of formula IVe (Scheme 4).



Method B may be used to prepare compounds of Formula IIa as shown below in Scheme 7.

5 Scheme 7




wherein R^1 , R^2 , R^3 , X^1 , X^2 , Z^1 and Z^2 are as previously defined.

In a typical procedure, carboxylic acid IIc is reacted with intermediate XI in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBOP, HATU, HBTU, COMU) known to those skilled in the art. The acid IIc and coupling partner is combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, DCE, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or slightly elevated temperature.

Compounds of Formula IIc may be synthesized as shown below in Scheme 8.

15

Scheme 8



where R¹, R², R³, X¹, and Z² are as previously defined, PG⁵ is an alkyl or aryl group included but not limited to methyl, ethyl, benzyl, allyl, and phenyl that can be removed by methods known to those skilled in the art, and LG¹ is a group such as OH that can be activated and displaced by the amine of compound IIe. Alternatively, LG¹ may be an appropriate halide (such as F, Cl, and Br) that can be displaced by a nucleophile.


In a typical procedure, IIe is reacted with II^f (LG¹= halide) in the presence of a suitable base (including but not limited to pyridine, triethylamine, DIPEA, and NMM) in an appropriate solvent (including but not limited to THF, DCM, and DMF) at temperatures ranging from -78 °C to 120 °C, but optimally from -20 °C to 50 °C. Alternatively, if LG¹ is OH, XII is reacted with reagent II^f in the presence of standard

coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBROP, HATU, HBTU, COMU) known to those skilled in the art. The acid and amine coupling partners are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or 5 slightly elevated temperature. It will be appreciated by those skilled in the art that compound II^f (LG¹= halide), may be readily made from the corresponding carboxylic acid by treatment with a suitable halogenating reagent. Following amide formation between II^e and II^f, the protecting group PG⁵ is then removed using conditions described for removing of PG³ of compound of formula IV^e (scheme 4) to give compound II^c.

10

Method B-2 may be used to prepare compounds of Formula II^a as shown below in Scheme 9.


Scheme 9



where R¹, R², R³, X¹, X², Z¹ and Z² are as previously defined. The reaction may be conducted under 15 conditions described for coupling reaction between compound II^g and compound II^f (scheme 8)

Compounds of Formula II^g may be prepared as shown below in Scheme 11.


Scheme 11



where R³, X¹, X², Z¹ are as previously defined and PG⁶ is a suitable amine protecting group 20 including but not limited to Boc, Cbz, Alloc, and Fmoc.

In a typical procedure, carboxylic acid II^g is reacted with an appropriate coupling partner containing a -X²-Z¹ moiety in the presence of standard coupling agents (e.g., EDC/HOBt, DCC, PyBOP, PyBROP, HATU, HBTU, COMU) known to those skilled in the art. The acid and the coupling partner are combined with the coupling agent in an organic solvent (including but not limited to DMF, dichloromethane, acetonitrile, and tetrahydrofuran) in the presence of a base (including but not limited to DIPEA, triethylamine, and NMM) at room temperature or slightly elevated temperature. PG⁶ is then removed by reacting the resulting intermediate with the appropriate reagent familiar to those skilled in the art.


Synthesis of Compound-7 (C3SLF)



To a solution of **Intermediate I-24** (18 mg, 0.085 mmol) in tetrahydrofuran (1.2 mL) was added *N*-methylmorpholine (10 μ L, 0.085 mmol). The solution was cooled to -20 °C and then

5 isobutylchloroformate (11 μ L, 0.085 mmol) was added dropwise. The solution was immediately warmed to 0 °C. After stirring for 45 minutes at 0 °C, the TFA salt of **Intermediate I-23** (37 mg, 0.057 mmol) was mixed with *N*-methylmorpholine (7 μ L, 0.057 mmol) in anhydrous tetrahydrofuran (0.5 mL) and added dropwise to the 0 °C solution of mixed anhydride over the course of 5 minutes. The resulting solution was stirred for 1.5 hours at 0 °C, at which point methanol (1 mL) was added, and the solution was immediately 10 warmed to room temperature and stirred for 5 minutes. The solution was diluted with dichloromethane (75 mL) and saturated aqueous sodium bicarbonate (50 mL). The layers were separated and the aqueous layer was extracted with dichloromethane (2 x 30 mL). The organics were combined, washed with brine (20 mL), dried over magnesium sulfate, filtered, and the solvent was removed in vacuo. 15 Purification by silica gel chromatography (0-80% ethyl acetate in hexanes) provided **Compound 7** (C3SLF) as a white foam (20 mg, 50% yield). ^1H NMR (400MHz, CDCl_3) δ 9.49 (s, 1H), 8.55 (d, 1H), 7.77 (app t, 1H), 7.71-7.65 (m, 2H), 7.60-7.58 (m, 1H), 7.20-7.17 (m, 1H), 7.01 (d, 1H), 6.78-6.76 (m, 1H) 20 6.70-6.67 (m, 2H), 5.77 (dd, 1H), 5.33 (d, 1H), 3.86 (s, 3H), 3.85 (s, 3H), 3.35 (d, 1H), 3.25 (t, 2H), 3.15 (dt, 1H), 2.92 (t, 2H), 2.64-2.50 (m, 2H), 2.38 (d, 1H), 2.29-2.15 (m, 1H), 2.10-2.00 (m, 1H), 1.78-1.63 (m, 3H), 1.53-1.37 (m, 3H), 1.22 (s, 6H), 0.89 (t, 3H, rotamer 1), 0.80 (t, 3H, rotamer 2). ESI-MS m/z = 722.0 [M+H] $^+$, 744.0 [M+Na] $^+$; Calculated MW: 721.93.

Synthesis of Compound-8




To a solution of **Compound 7** (22 mg, 0.031 mmol) in dichloromethane (0.5 mL) was added 25 triethylamine (62 μ L, 0.55 mmol) followed by 2-(dimethylamino)ethanethiol (0.0mL, 0.0500mmol) (4.8 mg, 0.046mmol). After stirring at room temperature for 30 minutes, the solution was poured into dichloromethane (30 mL) and saturated aqueous sodium bicarbonate (30 mL). The layers were separated and the aqueous layer was extracted with dichloromethane (2 x 20 mL). The organics were dried over magnesium sulfate, filtered, and the solvent was removed in vacuo. Purification by silica gel 30 chromatography (0-10% methanol in dichloromethane) provided impure material that was repurified by reverse phase HPLC (acetonitrile in water with 0.1% formic acid) to give the formic acid salt of **Compound-8** (7.1 mg, 21% yield) as a white solid. ^1H NMR (400MHz, CDCl_3) δ 12.53 (br s, 1H), 9.50 (s, 1H), 8.00 (s, 1H), 7.75 (d, 1H), 7.68 (s, 1H), 7.29 (t, 1H), 7.02 (d, 1H), 6.79-6.77 (m, 1H), 6.70-6.67 (m,

2H), 5.76 (dd, 1H), 5.29 (d, 1H), 3.86 (s, 3H), 3.85 (s, 3H), 3.36-3.27 (m, 2H), 3.23-3.11 (m, 4H), 2.90 (t, 2H), 2.79-2.71 (m, 8H), 2.62-2.50 (m, 2H), 2.55 (d, 1H), 2.28-2.17 (m, 1H), 2.12-2.03 (m, 1H), 1.72-1.61 (m, 3H), 1.49-1.36 (m, 3H), 1.22 (s, 3H), 1.21 (s, 3H), 0.88 (t, 3H, rotamer 1), 0.79 (t, 3H, rotamer 2). ESI-MS m/z = 716.1 [M+H]⁺; Calculated MW: 715.97.


5

Synthesis of Compound-9



To a room temperature solution of **Intermediate I-23** (15 mg, 0.029 mmol), 3-(2,5-dioxopyrrol-1-yl)propanoic acid (5.8mg, 0.034 mmol), and DMAP (0.4 mg, 0.003 mmol), in dichloromethane was added 10 *N,N*-dicyclohexylcarbodiimide (9.4 mg, 0.046 mmol). After stirring at room temperature for 15 hours, the resulting solids were filtered through a syringe filter and washed with dichloromethane. The solvent was removed in *vacuo* and the resulting residue was brought up in ethyl acetate. After cooling to -20 °C, the suspension was filtered through a syringe filter and washed with cold ethyl acetate. The filtrate was 15 diluted with ethyl acetate (30 mL) and water (30 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 20 mL). The organics were dried over magnesium sulfate, filtered, and the solvent was removed *in vacuo*. Purification by silica gel chromatography (0-90% ethyl acetate in hexanes) provided **Compound-9** (11 mg, 60% yield) as an oil. ¹H NMR (400MHz, CDCl₃) δ 8.25 (br s, 1H), 7.79 (d, 1H), 7.38 (s, 1H), 7.29 (t, 1H), 6.98 (d, 1H), 6.82 (s, 2H), 6.78-6.76 (m, 1H), 6.70-6.66 (m, 2H), 5.83 (dd, 1H), 5.37 (d, 1H), 4.43 (d, 1H), 4.37 (d, 1H), 3.86 (s, 3H), 3.85 (s, 3H), 3.32 (d, 1H), 2.96 (t, 20 1H), 2.55 (t, 2H), 2.35 (d, 1H), 2.29-2.15 (m, 1H), 2.11-2.01 (m, 1H), 1.81-1.60 (m, 4H), 1.49-1.42 (m, 3H), 1.27 (s, 3H), 1.25 (s, 3H), 0.93 (t, 3H, rotamer 1), 0.82 (t, 3H, rotamer 2). ESI-MS m/z = 662.0 [M+H]⁺; Calculated MW: 661.75.

Synthesis of (*R*)-3-(3,4-dimethoxyphenyl)-1-(3-(4-(pyridin-2-ylsulfanyl)butanamido)phenyl)propyl (*S*)-1-(3,3-dimethyl-2-oxopentanoyl)piperidine-2-carboxylate (C4-SLF):



To a solution of Aniline **1** (90mg, 172 μ mol, 1eq), disulfide **2** (79mg, 343 μ mol, 2eq) and diisopropylethylamine (149 μ L, 111mg, 858 μ mol, 5eq) in DMF (3mL) was added HATU (130mg, 343 μ mol, 2eq) and the reaction mixture was stirred at room temperature for 24h. The reaction mixture was diluted with water and extracted with ethyl acetate (3x). The organic extracts were washed with water, saturated sodium chloride, dried over magnesium sulfate and evaporated. The residue was purified on Silica gel

gradient elution (20% ethyl acetate : 80% heptane → 100% ethyl acetate) to provide the title compound C4-SLF (98mg, 77%). MS (ESI) calc=736.3 (M+H), obs=736.3.

Example 2: Synthesis of Certain Conjugates

5 *General Protocol:* This protocol describes a method for the formation of target protein-compound conjugates.


Reagents: Compound in 100% DMSO (in-house) and mammalian target protein (in-house)

10 *Equipment:* Mini-PROTEAN TGX Gel (Bio-Rad)

15 *Experimental Protocol:* A 1:2 molar ratio of target protein and compound are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Cross-linking efficiency is assessed by SDS-PAGE gel. Conjugates migrate slower than non-cross-linked target protein. For thiol reactive compounds, the Cys specific attachment of the compound to the target protein can be further confirmed by SDS-PAGE after the addition of 100 mM DTT to the reaction mixture, which reduces the conjugate back into its components.

20 **A. Formation of KRAS_{GTP/S39C} lite/C2-FK506 conjugates**

Reagents: C2-FK506 in 100% DMSO (in-house), KRAS_{GTP/S39C} lite (in house; residues 1-169 containing G12V/S39C/C51S/C80L/C118S).



C2-FK506

25

Equipment: Mini-PROTEAN TGX Gel (Bio-Rad)

30 *Experimental Protocol:* A 1:2 molar ratio of KRAS_{GTP/S39C} lite and C2-FK506 are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Cross-linking efficiency is assessed by SDS-PAGE gel. Attachment of C2-FK506 to Cysteine 39 on KRAS_{GTP/S39C} lite is also assessed by incubation of the reaction mixture with 100 mM DTT.

35 *Results:* C2-FK506 cross-links efficiently with KRAS_{GTP/S39C} lite and is specific for Cysteine 39 (FIG. 1).

B. Formation of KRAS_{GTP/G12C} lite/SFAX9DS conjugates

Reagents: SFAX9DS in 100% DMSO (in-house), KRAS_{GTP/G12C} lite (in house; residues 1-169 containing G12C/C51S/C80L/C118S).

5 *Equipment:* Mini-PROTEAN TGX Gel (Bio-Rad)

Experimental Protocol: A 1:2 molar ratio of KRAS_{GTP/G12C} lite and SFAX9DS are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Cross-linking efficiency is assessed by 10 SDS-PAGE gel. Wild-type CypA also cross-linked with the compound. Cysteine 52 as a reactive Cysteine on CypA and was mutated to Serine to abrogate presenter cross-linking.

15 Results: SFAX9DS cross-links efficiently with KRAS_{GTP/G12C} lite protein and CypAc_{52S} does not cross-link to SFAX9DS (FIG. 2).

Example 3: Formation of Certain Complexes

General Protocol: This protocol describes two methods for the formation and isolation of complexes comprised of presenter protein, compound, and mammalian target protein.

20 *Reagents:* Compound in 100% DMSO (in-house), presenter protein (in-house) and mammalian target protein (in-house)

Equipment: Mini-PROTEAN TGX Gel (Bio-Rad), Superdex 75 (GE Healthcare, CV 120 mL)

25 *Experimental Protocol A: Pre-conjugated compound and Protein*

A 1:2 molar ratio of conjugate and presenter protein are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Pure complex is isolated by Size Exclusion Chromatography (SEC) purification. The reaction mixture is directly injected on a Superdex 75 column (CV 120mL) pre-equilibrated with buffer containing 12.5 mM HEPES pH 7.4, 75 mM NaCl. Complex elutes at a higher molecular weight than unreacted target protein and presenter protein. To confirm presence of complex in the elution peak, samples are assessed by SDS-PAGE.


35 *Experimental Protocol B: Cross-linking reagent, presenter protein and target protein*

A 1:2:2 molar ratio of compound, presenter protein, and target protein are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Pure complex is isolated by Size Exclusion Chromatography (SEC) purification. The reaction mixture is directly injected on a Superdex 75 column (CV 120mL) pre-equilibrated with buffer containing 12.5 mM HEPES pH 7.4, 75 mM NaCl. Complex elutes at a higher molecular weight than unreacted target protein and presenter protein. To confirm presence of complex in the elution peak, samples are assessed by SDS-PAGE.

A. Formation of KRAS_{GTP/S39C} lite/C2Holt/FKBP12 ternary complex

Reagents: C2Holt in 100% DMSO (in-house), KRAS_{GTP/S39C} lite (in house; residues 1-169 containing G12V/S39C/C51S/C80L/C118S), and FKBP12 (in-house).

5



C2Holt

Equipment: Mini-PROTEAN TGX Gel (Bio-Rad), Superdex 75 (GE Healthcare, CV 120 mL)

10 *Experimental Protocol:* A 1:2:2 molar ratio of C2-Holt, FKBP12, and KRAS_{GTP/S39C} lite are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Pure complex is isolated by Size Exclusion Chromatography (SEC) purification. The reaction mixture is directly injected on a Superdex 75 column (CV 120mL) pre-equilibrated with buffer containing 12.5 mM HEPES pH 7.4, 75 mM 15 NaCl. Complex elutes at around 69 mL post injection and unreacted KRAS_{GTP/S39C} lite and FKBP12 elutes at around 75 mL and 87 mL post injection respectively. To confirm the presence of KRAS_{GTP/S39C} lite and FKBP12 in the elution peak, samples are also assessed by SDS-PAGE.

20 *Results:* The SEC profile and SDS-PAGE analysis of the elution peaks confirm the formation of KRAS_{GTP/S39C} lite/C2Holt/FKBP12 complex (FIG. 3A and 3B).

B. Formation of KRAS_{GDP/S39C} lite/SFAC4DS/CypAc_{52S} ternary complex

Reagents: SFAC4DS in 100% DMSO (in-house), KRAS_{GDP/S39C} lite (in house; residues 1-169 containing G12V/S39C/C51S/C80L/C118S), and CypAc_{52S} (in-house).

25

Equipment: Mini-PROTEAN TGX Gel (Bio-Rad), Superdex 75 (GE Healthcare, CV 120 mL)

30 *Experimental Protocol:* A 1:2:2 molar ratio of SFAC4DS, CypAc_{52S}, and KRAS_{GDP/S39C} lite are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 2% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Pure complex is isolated by Size Exclusion Chromatography (SEC) purification. The reaction mixture is directly injected on a Superdex 75 column (CV 120mL) pre-equilibrated with buffer containing 12.5 mM HEPES pH 7.4, 75 mM NaCl. Complex elutes at around 69 mL post injection and unreacted KRAS_{GDP/S39C} lite and CypAc_{52S} elutes at around 75 mL and 80 mL post injection respectively. To confirm the presence of KRAS_{GDP/S39C} 35 lite and CypAc_{52S} in the elution peak, samples are also assessed by SDS-PAGE.

Results: The SEC profile and SDS-PAGE analysis of the elution peaks confirm the formation of KRAS_{GDP/S39C} lite/SFAC4DS/CypAc_{52S} complex (FIG. 4).

C. Formation of PTP1B_{S187C} lite/C3SLF/FKBP12 ternary complex

Reagents: C3SLF in 100% DMSO (in-house), PTP1B_{E186C} lite (in house; residues 1-293 containing C32S/C92V/C121S/S187C), and FKBP12 (in-house).

5 *Equipment:* Mini-PROTEAN TGX Gel (Bio-Rad), Superdex 75 (GE Healthcare, CV 120 mL)

Experimental Protocol: A 1:3:3 molar ratio of C3SLF, FKBP12, and PTP1B_{S187C} lite are mixed together in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer containing 4% DMSO. The reaction is incubated at 37°C for 30 min, followed by overnight incubation at room temperature. Pure complex is isolated by
10 Size Exclusion Chromatography (SEC) purification. The reaction mixture is directly injected on a Superdex 75 column (CV 120mL) pre-equilibrated with buffer containing 12.5 mM HEPES pH 7.4, 75 mM NaCl. Complex elutes at around 62 mL post injection and unreacted FKBP12 elutes at around 75 mL (Dimer) and 90 mL (Monomer) post injection respectively. To confirm the presence of PTP1B_{S187C} lite and
15 FKBP12 in the elution peak, samples are also assessed by SDS-PAGE. Free PTP1B_{S187C} lite and elution time.

Results: The SEC profile and SDS-PAGE analysis of free PTP1B_{S187C} lite and FKBP12 proteins (FIG. 5A) confirming the free PTP1B_{S187C} lite elutes at around 64-65 ml. The SEC profile and SDS-PAGE analysis of the elution peaks confirm the formation of PTP1B_{S187C} lite/C3SLF/FKBP12 complex, which
20 elutes at around 61 ml (FIG. 5B).

Example 4: Conjugate Formation When Presenter Protein is Present, but not When it is Absent

This protocol describes methods to analyzing cross-linking efficiency using mass spectrometry and gel shift assay in effort to assess the presenter dependency of conjugate formation.

25 *Reagents:* Compound in 100% DMSO (in-house), FKBP12 (in-house), KRAS_{GTP/G12C} (in-house, residues 1-169).

Experimental protocol: In order to follow the kinetics of disulfide crosslinking reactions, Agilent 6230 TOF-LC/MS and Agilent 1260 HPLC instruments employing a AdvanceBio RP-mAb C4 column (2.1 x 100 mm, 3.5 µm) and equipped with an auto-sampler were used. HPLC grade acetonitrile and water
30 (each containing 1.0 mM ammonium formate and 1% formic acid by volume) were used as mobile phase with the following ramp: 0.6 ml/min flow rate, water:acetonitrile = 95:5 from 0.0 to 13.0 min ramping to water:acetonitrile = 5:95 from 13.0 to 17.0 min. Total time = 17.0 min.

All crosslinking reactions were performed in 1.5 mL amber colored glass vials equipped with 0.5 mL glass inserts. A water-soluble peptide (SEQ ID NO: 1: YQNLLVGRNRGEEILD) was employed as an
35 internal standard. Although the actual sequence of the internal standard is inconsequential, choice of amino acid residues were critical to avoid interference in the crosslinking assays. Hence, proline (interfering with FKBP12) and cysteine (interfering with disulfide bond formation) residues were excluded. All reactions and standard solutions were prepared in HEPES (pH 7.4, 1.0 mM MgCl₂) buffer.

Prior to every reaction, a standard curve was obtained for the individual components using a
40 series of standard solutions (an example of standard curve analysis for FKBP12 is shown in Table 3 below). Using the data from standard curve, µmol of protein samples were plotted against the ratio of

areas (sample:std) and a linear fit ($y = mx+c$) was employed to obtain the slope and intercept. The value of slope and intercept involved in these standard curves were accounted for during evaluation of the substrate and product concentration before and during the course of the reaction. For every substrate/product, an initial injection was followed up by a blank injection to verify presence of any residual protein/reagents. Based on this analysis the sequence of auto-sampler can be adjusted to include appropriate number of blank injections for removal of residual components, if any. The MS spectra were analyzed using Agilent MassHunter v B.07.0 software.

Table 3. Standard curve for FKBP12. Slope = 6.53, Intercept = -0.64, $R^2 = 0.999$

Solution	Concentration (mM)	Area under internal standard ($R_t = 7.1$ min)	Area under FKBP-12 ($R_t = 8.9$ min)	Ratio of Sample:Std
1	100	2615940.1	40314760.1	15.41
2	10	5244417.4	8718886.5	1.66
3	1	5868006.9	1745344.5	0.30
4	0.5	5407354.8	910748.2	0.17
5	0.1	5626447.0	237074.5	0.04

10 In a representative experiment to assess the presenter dependency of ligand cross-linking on the target protein, KRAS_{GTP/G12C} and either C3 or C4SLF ligand were incubated in the presence or absence of FKBP12 at concentrations of 2 μ M KRAS, 10 μ M FKBP12, 10 μ M C3 or C4SLF for 4 hours at room temperature in 12.5 mM HEPES pH 7.4, 75 mM NaCl, 1 mM MgCl₂, 3% DMSO. Analysis of the amount 15 of KRAS undergoing disulfide cross-linking with the ligand using the method above. As shown in Table 4, 5- to 10-fold increased cross-linking efficiency was observed in the presence of the presenter:

Table 4. Crosslinking efficiency of C3- and C4SLF analyzed MS

	C3SLF		C4SLF	
FKBP12	-	+	-	+
%xlink to KRAS	9.6	47.5	7.0	69.8

20 In parallel to mass spectrometry analysis, cross-linking reactions with C3 or C4SLF were subjected to gel shift assay using 12% SDS-PAGE in the presence or absence of FKBP12 at the same experimental condition described above except that cross-linking reactions were set up at higher concentrations (60 μ M KRAS, 180 μ M FKBP12, and 180 μ M C3 or C4SLF) and they were quenched by MMTS to terminate the reaction. Similar to the MS data, the ligand cross-linking efficiency was boosted significantly in the presence of FKBP12, which is more pronounced for C4SLF (FIG. 6).

Example 5: Determination of Presenter Protein/Target Protein Interface Structure by X-ray Analysis

This protocol describes the crystallization and structure determination method for the crystal structure of a ternary complex of FKBP12-C2Holt-KRAS_{GTP/S39C}.

5

A. Crystal structure determination of FKBP12-C2Holt-KRAS_{GTP/S39C} ternary complex

Reagents: Ligand (C2Holt) in 100% DMSO (in-house), FKBP12 (in-house), KRAS_{GTP/S39C} lite (in-house, residues 1-169 containing G12V/S39C/C51S/C80L/C118S).

10 **Equipment:** Superdex 75 (GE Healthcare)

Experimental Protocol: C Holt and FKBP12 were added to KRAS_{GTP/S39C} lite at 3:1 and 1.5:1 molar excess in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer, 1 mM MgCl₂, 2% DMSO, and incubated overnight at 20°C or 36-72 hours at 4°C. Pure complex was isolated by size exclusion chromatography 15 on a Superdex 75 column in 12.5 mM HEPES pH 7.4, 75 mM NaCl, and 1 mM MgCl₂. Purified complex (at 15-20 mg/ml) was subjected to crystallization screening at 20°C using sitting drop vapor diffusion method. Crystals were grown in a well solution containing 0.1 M MES pH 6.5, 20-22% PEG 20,000. For data collection crystals were transferred to a solution containing mother liquor supplemented with 15% glycerol, and then frozen in liquid nitrogen. Diffraction datasets were collected at the Advanced Photon 20 Source (APS) and processed with the HKL program. Molecular replacement solutions were obtained using the program PHASER in the CCP4 suite, using the published structure of FKBP12 (PDB-ID 1FKD) and KRAS (PDB-ID 3GFT) as search models. Subsequent model building and refinement were performed according to standard protocols with the software packages CCP4 and COOT.

25 **Results:** Overall structure of FKBP12-C2Holt -KRAS_{GTP/S39C}: The crystal contains one heterodimer of FKBP12 and KRAS_{GTP/S39C} in the asymmetric unit (FIG. 7). The model comprised residues Met1 to Glu108 of FKBP12 and Met1 to Lys169 of KRAS_{GTP/S39C}. The resulting electron density shows unambiguous binding mode, including the orientation and conformation of the ligand. The continuous 30 electron density was observed for the disulfide generated from the cysteine of the protein and the sulfur from the ligand.

The KRAS_{GTP/S39C} residues involved in binding C2Holt (4 Å distance cut-off) are Glu37, Cys39, Leu56, and Met67. The KRAS_{GTP/S39C} residues involved in binding to FKBP12 are Glu3, Lys5, Ile36, Cys39, Tyr40, Arg41, Asp54, Glu63, Tyr64, Met67, and Arg73. The FKBP12 residues involved in binding KRAS_{GTP/S39C} are Arg43, Lys53, Gln54, Glu55, Thr86, Pro89, Gly90, and Ile92. The FKBP12 residues 35 involved in binding C2Holt are Tyr27, Phe37, Asp38, Phe47, Glu55, Val56, Ile57, Trp60, Tyr83, His88, Ile91, Ile92, and Phe100.

The total buried surface area of the complex is 1,947 Å². The buried surface area of KRAS_{GTP/S39C} is 600 Å² of which 501 Å² is contributed by FKBP12 (83%), and 99 Å² contributed by C2Holt (17%). The buried surface area of FKBP12 is 762 Å² of which 500 Å² contributed by KRAS_{GTP/S39C} (66%) 40 and 262 Å² contributed by C2Holt (34%). The buried surface area of C2 Holt is 584 Å² of which 132 Å² contributed by KRAS_{GTP/S39C} (23%) and 452 Å² contributed by FKBP12 (77%). The protein-protein

interface between KRAS_{GTP/S39C} and FKBP12 is formed by both hydrophobic and polar interactions, including three intermolecular H-bonds. The binding interface between C2 Holt and FKBP12 is largely contributed by hydrophobic interactions, but also contributed by three H-bonds between three carbonyl groups of the ligand and Tyr27, Ile57, and Tyr83 of FKBP12. C2 Holt forms minimal contact with 5 KRAS_{GTP/S39C} by design (99 Å²) but forms one H-bond with Glu37 of KRAS_{GTP/S39C}. Data collection and refinement statistics of the final structure are listed in Table 5 below.

B. Crystal structure determination of KRAS_{GDP/S39C}/SFAC4DS/CypA_{C52S} ternary complex

10 *Reagents:* Ligand (SFAC4DS) in 100% DMSO (in-house), CypA_{C52S} (in-house), KRAS_{GDP/S39C} lite (in-house, residues 1-169 containing G12V/S39C/C51S/C80L/C118S).

Equipment: Superdex 75 (GE Healthcare)

15 *Experimental Protocol:* SFAC4DS and CypA_{C52S} were added to KRAS_{GDP/S39C} lite at 2:1 and 2:1 molar excess in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer, 1 mM MgCl₂, 2% DMSO, and incubated overnight at 20°C. Pure complex was isolated by size exclusion chromatography on a Superdex 75 column in 12.5 mM HEPES pH 7.4, 75 mM NaCl, and 1 mM MgCl₂. Purified complex (at 15 mg/ml) was subjected to crystallization screening at 20°C using sitting drop vapor diffusion method. Crystals were 20 grown in a well solution containing 0.1 M Bis-Tris pH 6.5, 25 % PEG 3350. For data collection, crystals were transferred to a solution containing mother liquor supplemented with extra PEG 3350 to make it 40 % of PEG, and then frozen in liquid nitrogen. Diffraction datasets were collected at the Advanced Light 25 Source (ALS) and processed with the HKL program. Molecular replacement solutions were obtained using the program PHASER in the CCP4 suite, using the published structure of CypA (PDB-ID 1CWA) and KRAS (PDB-ID 3GFT) as search models. Subsequent model building and refinement were performed according to standard protocols with the software packages CCP4 and COOT.

30 *Results:* Overall structure of CypA_{C52S}-SFAC4DS -KRAS_{GDP/S39C}: The crystal contains one heterodimer of CypA_{C52S} and KRAS_{GDP/S39C} in the asymmetric unit (FIG. 8). The model comprised residues Met1 to Glu165 of CypA and Met1 to Lys169 of KRAS_{GDP/S39C}. The resulting electron density shows unambiguous binding mode, including the orientation and conformation of the ligand. The continuous electron density was observed for the disulfide generated from the cysteine of the protein and the sulfur from the ligand.

35 The KRAS_{GDP/S39C} residues involved in binding SFAC4DS (4 Å distance cut-off) are Glu3, Lys5, Cys39, Arg41, Leu52, Asp54, Ile55 and Leu56. The KRAS_{GDP/S39C} residues involved in binding to CypA_{C52S} are Glu37, Asp38, Cys39, Arg41, Gln43, Leu56, Ala66, Met67, Gln70, and Thr74. The CypA_{C52S} residues involved in binding KRAS_{GDP/S39C} are Arg55, Ile57, Arg69, Asn71, Thr73, Ala81, Ala103, Arg148, and Asn149. The CypA_{C52S} residues involved in binding SFAC4DS are Arg55, Phe60, Met61, Gln63, Gly72, Ala101, Asn102, Gln111, Phe113, and His126.

40 The total buried surface area of this complex cannot be calculated due to partial structural disorder at the protein-protein interface. Excluding the disordered region for calculation, the buried surface area at the protein-protein interface is greater than 1,350 Å², of which over 30% is contributed by

SFAC4DS (443 Å²). The protein-protein interface between KRAS_{GDP/S39C} and CypA_{C52S} is formed by both hydrophobic and polar interactions, including two intermolecular H-bonds. The binding interface between SFAC4DS and CypA is contributed both by hydrophobic and polar interactions. There are six H-bonds between carbonyl and N-H groups of the ligand and residues Arg55, Gln63, Asn102, and His126 of 5 CypA_{C52S}. SFAC4DS forms minimal direct contact with KRAS_{GDP/S39C} but forms one H-bond with Arg41 of KRAS_{GDP/S39C}. Data collection and refinement statistics of the final structure are listed in Table 5 below.

C. Crystal structure determination of PTP1B_{S187C}/C3SLF/FKBP12 ternary complex

Reagents: Ligand (C3SLF) in 100% DMSO (in-house), FKBP12 (in-house), PTP1B_{S187C} lite (in-10 house, residues 1-169 containing C32S/C92V/C121S/S187C).

Equipment: Superdex 75 (GE Healthcare), Gryphon (Art Robbins Instruments)

Experimental Protocol: C3SLF and FKBP12 were added to PTP1B_{S187C} lite at 3:1 molar excess in 15 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer, 4% DMSO, and incubated 36-72 hours at 4°C. Pure complex was isolated by size exclusion chromatography on a Superdex 75 column in 12.5 mM HEPES pH 7.4 and 75 mM NaCl. Purified complex (at 15 mg/ml) was subjected to crystallization screening at 20°C using sitting drop vapor diffusion method. Crystals were grown in a well solution containing 0.2 M Magnesium Acetate, 20 % w/v PEG 3350. For data collection, crystals were transferred to a solution 20 containing mother liquor supplemented with 25% PEG400, and then frozen in liquid nitrogen. Diffraction datasets were collected at the Advanced Photon Source (APS) and processed with the XDS program. Molecular replacement solutions were obtained using the program PHASER in the CCP4 suite, using the published structure of FKBP12 (PDB-ID 2PPN) and PTP1B (PDB-ID 2NT7) as search models. Subsequent model building and refinement were performed according to standard protocols with the 25 software packages CCP4 and COOT.

Results: Overall structure of FKBP12-C3SLF-PTP1B_{S187C}: The crystal contains two complex molecules of FKBP12-C3SLF-PTP1B_{S187C} in the asymmetric unit (FIG. 9A). The model comprised residues Gly2 to Glu108 of FKBP12 and Glu6 to Phe280 of PTP1B_{S187C}. The resulting electron density 30 shows unambiguous binding mode, including the orientation and conformation of the ligand. The continuous electron density was observed for the disulfide generated from the cysteine of the protein and the sulfur from the ligand.

The total buried surface area of the complex is 1,042 Å². The buried surface area of PTP1B_{S187C} is 427 Å². The buried surface area of C3SLF is 615 Å² (FIG. 9B). The protein-protein interface between 35 PTP1B_{S187C} and FKBP12 is formed by both hydrophobic and polar interactions.

D. Crystal structure determination of MCL1_{S245C}/C3SLF/FKBP52 ternary complex

Reagents: Ligand (C3SLF) in 100% DMSO (in-house), FKBP52 (in-house, residues 1-140), MCL1_{S245C} lite (in-house, residues 172-327 containing S245C/C286S).

40

Equipment: Superdex 75 (GE Healthcare), Gryphon (Art Robbins Instruments)

Experimental Protocol: C3SLF and FKBP52 were mixed with MCL1_{S245C} lite at 3:1 molar excess in 12.5 mM HEPES pH 7.4, 75 mM NaCl buffer, 2% DMSO, and incubated 24-48 hours at 4°C. Pure complex was isolated by size exclusion chromatography on a Superdex 75 column in 12.5 mM HEPES pH 7.4 and 75 mM NaCl. Purified complex (at 15 mg/ml) was subjected to crystallization screening at 20°C using sitting drop vapor diffusion method. Crystals were grown in a well solution containing 2.1 M Malic acid. For data collection, crystals were transferred to a solution containing mother liquor supplemented with 20% glycerol, and then flash-frozen in liquid nitrogen. 3.0 Å resolution diffraction dataset was measured at the Advanced Photon Source (APS) and processed with the XDS program.

10 Molecular replacement solutions were obtained using the program PHASER in the CCP4 suite, using the published structure of FKBP52 (PDB-ID 1N1A) and PTP1B (PDB-ID 3MK8) as search models. Subsequent model building and refinement were performed according to standard protocols with the software packages CCP4 and COOT.

15 *Results:* The crystal contains one complex molecule of MCL1_{S245C}/C3SLF/FKBP52 in the asymmetric unit (FIG. 10). The resulting electron density revealed unambiguous binding between two proteins, including the orientation and conformation of the ligand. The continuous electron density was observed for the disulfide generated from the cysteine of the protein and the sulfur from the ligand. The total buried surface area of the complex is 1,410 Å², of which approximately 60% is contributed by

20 FKBP52 (804 Å²), and approximately 40% by C3SLF (606 Å²). Due to the limited resolution, the detailed analysis in the protein-protein and protein-ligand interaction was not feasible.

Table 5. Data collection and refinement statistics

.	Structure A	Structure B	Structure C	Structure D
Resolution [Å]	47.8-1.4	70.1-1.6	49.0-2.4	65.9-3.0
5 Number of reflections (working/test)	50,557/2,708	34,377/1,790	30,175/1,543	5,343/245
R_{cryst} [%]	18.9	17.9	23.0	33.2
$R_{\text{free}}^{[1]}$ [%]	21.7	21.2	28.1	37.8
Total number of atoms:				
10 Protein	2,202	2,528	6,192	2,035
Water	268	171	38	0
Ligands	69	63	43	43
Ions	1	1	0	0
Deviation from ideal geometry:				
15 Bond lengths [Å]	0.007	0.011	0.004	0.010
Bond angles [°]	1.33	1.49	0.87	1.30
Ramachandran plot:				
20 Most favoured regions [%]	93.8	96.5	93.3	95.2
Allowed regions [%]	6.2	3.1	4.9	4.4
Disallowed region [%]	0.0	0.3	1.8	0.4

¹Test-set contains 5% of measured reflections

²Root mean square deviations from geometric target values

³Calculated with RAMPAGE

25 **Example 6: Determination of Complex Formation by TR-FRET**

TR-FRET technology (LANCE, Perkin Elmer) is a standard method to detect the binary association of two fusion-tagged proteins, e.g., protein 1/tag A and protein 2/tag B, where A and B can be any of glutathione-S-transferase (GST), hexahistidine (His₆), FLAG, biotin-avi, Myc, and Hemagglutinin (HA). In this example, the technology is used to measure the compound-facilitated association of a 30 presenter protein with a target protein. A mixture of a presenter protein/tag A and a target protein/tag B are added to a 384-well assay plate containing compounds of the invention and incubated for 15 minutes. A mixture of anti-fusion tag A or B Europium-chelate donor and anti-fusion tag A or B allophycocyanin acceptor or Ulight acceptor reagents are added and the reactions are incubated for 240 minutes. The 35 TR-FRET signal is read on an Envision microplate reader (Perkin Elmer) using excitation = 320 nm, emission = 665/615 nm. Compounds that facilitate ternary complex formation are identified as those eliciting an increase in the TR-FRET ratio relative to DMSO control wells.

Determination of CYPA-Compound 3-KRAS_{G12C-GTP} Complex Formation by TR-FRET

Avi-tagged Cyclophilin A and His-tagged KRAS_{G12C-GTP} were mixed with increasing concentration of ligand (Compound 3) and incubated at room temperature for 15 minutes to allow formation of ternary complex. A pre-mixture of Anti-His Eu-W1024 and Streptavidin APC were then added and incubated for 5 60 minutes. TR-FRET signal is read on an EnVision microplate reader (Perkin Elmer, Ex 320 nm, Em 665/615 nm). A counter screen without presenter and target protein is also run to rule out the contribution of compounds alone.

Reagents and Instrument

- 10 ▪ His6-KRAS_{G12C-GTP} (in house; residues 1-169); 1.2 mM in PBS buffer, pH 7.4
- Avi-CYPA (in house; residues 1-165); 556 μ M in PBS buffer, pH 7.4
- Anti-His Eu-W1024 (Perkin Elmer)
- Streptavidin APC (Perkin Elmer)
- Ligand (W21487), 10 mM in 100% DMSO
- 15 ▪ EnVision (Perkin Elmer)
- Combi Mutidrop liquid dispenser with 8-channel small volume cassette
- 384-w ProxiPlate (black)

Experimental Protocol

1. Use Mosquito to dispense 100 nL/well of compounds (varying concentration in DMSO) into 384-w 20 black ProxiPlate to make assay-ready-plate (ARP).
2. Make 2 x assay buffer containing 40 mM Hepes pH 8.0, 200 mM NaCl, 2 mM MgCl₂, 0.1% BSA and 0.004% Tween-20.
3. Make 2x PRE-MIX A: 100 nM of His6-KRas G12C-GTP (1-169) and 1000 nM of Avi-CypA (1-165) in 1X assay buffer.
- 25 4. Use MutiDrop Combi to dispense 2x PRE-MIX A into ARP, 5 μ l/well. Incubate 15 min at RT.
5. Make 2X PRE-MIX B: 10 nM of anti-His Eu-W1024 and 40 nM of SA APC.
6. Use MutiDrop Combi to dispense 2x PRE-MIX B into ARP, 5 μ l/well. Shake briefly on Combi and incubate 60 min at RT.
7. Read on EnVision (Ex: 320 nm; Em1: 615 nm; Em2: 665 nm).
- 30 8. Data is processed using Dotmatics. Curves are fit using a 4 parameter non-linear fit to determine the EC50 value for formation of the ternary complex.

Results: The binding curve (FIG. 11) demonstrates Compound 3 dependent complex formation of CYPA-Compound 3- KRAS_{G12C-GTP} ternary complex, with a calculated EC50 value of 2.1 μ M

35

Example 7: Determination of Complex Formation by Amplified Luminescent Proximity Homogeneous Assay

AlphaScreen technology (Perkin Elmer) is a standard method to detect the binary association of two fusion-tagged proteins, e.g., protein 1/tag A and protein 2/tag B, where A and B can be any of 40 glutathione-S-transferase (GST), hexahistidine (His₆), FLAG, biotin-avi, Myc, and Hemagglutinin (HA). In

this example, the technology is used to measure the compound-facilitated association of a presenter protein with a target protein. A mixture of presenter protein/tag A and target protein/tag B are added to a 384-well assay plate containing compounds of the invention and incubated for 15 minutes. A mixture of anti-fusion tag A or B AlphaScreen donor beads and anti-fusion tag A or B AlphaScreen acceptor beads 5 are added and the reactions are incubated for 240 minutes. The AlphaScreen signal is read on an Envision microplate reader (Perkin Elmer) using excitation = 680 nm, emission = 585 nm. Compounds that facilitate ternary complex formation are identified as those eliciting an increase in the AlphaScreen signal relative to DMSO control wells.

10 **Determination of CYPA-Compound 3- KRAS_{G12C-GTP} Complex Formation by alpha LISA**

Avi-tagged Cyclophilin A and His-tagged KRAS_{G12C-GTP} were mixed with increasing concentration of ligand (Compound 3) and incubated at room temperature for 60 minutes to allow formation of ternary complex. A pre-mixture of Nickel chelate donor beads and Streptavidin acceptor beads were then added and incubated for 60 minutes. AlphaLISA signal is read on an EnVision microplate reader (Perkin Elmer, 15 Ex 680 nm, Em 615 nm). A counter screen without presenter and target protein is also run to rule out the contribution of compounds alone.

Reagents and instrument:

- His6-KRAS_{G12C-GTP} (in house; residues 1-169); 1.2 mM in PBS buffer, pH 7.4
- 20 ▪ Avi-CYPA (in house; residues 1-165); 556 uM in PBS buffer, pH 7.4
- Nickel chelate donor beads (Perkin Elmer)
- Streptavidin acceptor beads (Perkin Elmer)
- Ligand (W21487), 10 mM in 100% DMSO
- EnVision (Perkin Elmer)
- 25 ▪ Combi Mutidrop liquid dispenser with 8-channel small volume cassette
- alphaPlate-384 plate (white)

Experimental Protocol:

1. Use Mosquito to dispense 100 nL/well of compounds (varying concentration in DMSO) into 384-well black ProxiPlate to make assay-ready-plate (ARP).
- 30 2. Make 2X assay buffer containing 40 mM Hepes pH 8.0, 200 mM NaCl, 2 mM MgCl₂ and 0.004% Tween-20.
3. Make 2x PRE-MIX A: 300 nM of His6-KRas G12C-GTP (1-169) and 300 nM of Avi-CypA (1-165) in 1X assay buffer.
- 35 4. Use MutiDrop Combi to dispense 2x PRE-MIX A into ARP, 5 uL/well. Incubate 60 min at RT.
5. Make 2X PRE-MIX B: 30 uM/ml of streptavidin acceptor beads and 30 uM/ml of Nickel chelate donor beads.
6. Use MutiDrop Combi to dispense 2x PRE-MIX B into ARP, 5 uL/well. Shake briefly on Combi and incubate 60 min at RT.
- 40 7. Read on EnVision (Ex: 680 nm; Em1: 615nm).

8. Data is processed using Dotmatics. Curves are fit using a 4 parameter non-linear fit to determine the EC50 value for formation of the ternary complex.

Results: The binding curve (FIG. 12) demonstrates Compound 3 dependent complex formation of

5 CYPA-Compound 3- KRAS_{G12C-GTP} ternary complex, with a calculated EC50 value of 0.99 μ M.

Example 8: Determination of Complex Formation by Isothermal Titration Calorimetry

Isothermal Titration Calorimetry (ITC) is an established biophysical technique used to directly measure the heat change associated with the binary interaction of two proteins or protein to a ligand. Measurement of the heat change allows accurate determination of association constants (K_a), reaction 10 stoichiometry (N), and the change in binding enthalpy (ΔH). Gibbs energy changes (ΔG) and entropy changes (ΔS) can also be determined using the relationship: $\Delta G = -RT\ln K_a = \Delta H - T\Delta S$ (where R is the gas constant and T is the absolute temperature). In this example, the method is used to measure binding (e.g., non-covalent or covalent binding) of a compound or conjugate of the invention to a presenter protein.

15

Determination of Kinetics and Thermodynamics of Binding between FKBP12-Compound 1 and CEP250 by ITC

Reagents: Compound 1 and Compound 2 in 100% DMSO (in-house), Protein Buffer (10 mM HEPES, pH 7.5, 75 mM NaCl, 0.5 mM TCEP), assay buffer (protein buffer + 1% DMSO), FKBP12 (in-20 house), CEP250_{29.4} (in-house, residues 1982-2231) and CEP250_{11.4} (in-house, residues 2134-2231).

Equipment: MicroCalTM ITC₂₀₀ (GE Healthcare). Instrument parameters are shown in Table 6.

Table 6. Isothermal Titration Calorimetry instrument parameters

Experimental device	MicroCal TM ITC ₂₀₀ (GE Healthcare)
Sample cell volume (μ l)	270
Injector volume (μ l)	40
Experimental parameters	
Total number of Injections	19
Cell Temperature ($^{\circ}$ C)	25
Reference Power (μ Cal/s)	5
Initial Delay (s)	200
Stirring Speed (rpm)	750
Injection parameters	
Volume (μ l)	2
Duration (s)	4
Spacing (s)	170-200
Filter Period (s)	5
Feedback Mode/Gain	High

Experimental Protocol: FKBP12 stock solution is diluted to 10 μ M in assay buffer (1% DMSO final). Compound is added to FKBP12 to 20 μ M (1% DMSO final), and binary complex is filled into the reaction cell of the ITC device after 5-10 min pre-incubation time. CEP250 protein stocks are diluted to 50 μ M in assay buffer and supplemented with 20 μ M compound (1% DMSO final) before being filled into the injection syringe. A control experiment in the absence of compound is also run to determine the heat associated with operational artifacts and the dilution of titrant as it is injected from the syringe into the reaction cell. More detailed experimental parameters are shown in Table 7.

Table 7. Final protein and ligand concentrations

Experiment	Cell content	Syringe content	Ligand	DMSO conc. (%)
3	FKBP12, 10 μ M	CEP250 _{29.2} , 50 μ M	None	1.0
4	FKBP12, 10 μ M	CEP250 _{11.4} , 50 μ M	None	1.0
5	FKBP12, 10 μ M	CEP250 _{29.2} , 118 μ M	Compound 1b, 20 μ M	1.0
6	FKBP12, 10 μ M	CEP250 _{29.2} , 118 μ M	Compound 2, 20 μ M	1.0
7	FKBP12, 10 μ M	CEP250 _{11.4} , 68 μ M	Compound 1b, 20 μ M	1.0
8	FKBP12, 10 μ M	CEP250 _{11.4} , 68 μ M	Compound 2, 20 μ M	1.0

10

Data Fitting: Data were fitted with the Origin ITC200 software according to the following procedure:

- 1) Read raw data
- 2) In "mRawITC": adjust integration peaks and baseline, integrate all peaks
- 3) In "Delta H" - data control: remove bad data (injection #1 and other artifacts), subtract straight line (background subtraction)
- 4) In "Delta H"- model fitting: select *one set of sites* model, perform fitting with Levenberg-Marquardt algorithm until Chi Square is not reduced further, finish with "done" (parameters N, K_a and ΔH are calculated based on fitting)

20

Results: ITC measurements for the binding of FKBP12-Compound 1 and FKBP12-Compound 2 binary complexes to CEP250 are summarized in Table 8 and FIG. 13. Overall, the data for FKBP12-Compound 1 and FKBP12-Compound 2 binary complexes binding to CEP250_{11.4} and CEP250_{29.4} show similar interaction parameters. K_d values were similar for all combinations. All interactions show an almost identical thermodynamic profile in which binding is characterized by a purely enthalpic binding mode ($-T^* \Delta S$ term is positive and does not contribute to the Gibbs free energy). Binding stoichiometries for all interactions were N=0.5-0.6 and support a 1:2 binding ratio for 1 CEP250 homodimer binding to 2 FKBP12 molecules, as evidenced in the crystal structure of CEP250_{11.4}/Compound 1/FKBP12.

25

Table 8. Determination of FKBP12-Compound 1-CEP250 ternary complex formation by ITC

Experiment	T (K)	N	K _d (μ M)*	ΔH (kJ*mol ⁻¹)**	$-T^* \Delta S$ (kJ*mol ⁻¹)***	ΔH (kJ*mol ⁻¹)****
3	298	N.D.	N.D.	N.D.	N.D.	N.D.
4	298	N.D.	N.D.	N.D.	N.D.	N.D.

5	298	0.50	0.19	-52.21	13.80	-38.41
6	298	0.57	0.36	-58.48	21.73	-36.74
7	298	0.56	0.07	-49.37	8.62	-40.75
8	298	0.54	0.08	-47.78	7.41	-40.36

* K_d (calculated from $K_a = 1/K_d$)

** ΔH

*** $T^* \Delta S$ (calculated from $-T\Delta S = \Delta G - \Delta H$)

**** $\Delta G = -RT \ln K_a = RT \ln K_d$

5

Example 9: Determination of Kinetics of Binding between Conjugates and Proteins by Surface Plasmon Resonance

Surface Plasmon Resonance (SPR) is a biophysical technique used to measure the kinetics associated with the binary interaction of either two proteins or a protein to a ligand. Typically, one component of the binary interacting pair is immobilized on a flow cell of an activated sensor chip via a fusion tag. Increasing concentrations of the second component (the analyte) are then injected over the active surface for a fixed time. An increase in SPR signal (expressed in resonance units, RU) during the association phase and decrease in SPR signal during the dissociation phase is indicative of an interaction and can be fit to a binding model to determine associated K_D , K_a , K_d values. In this example, the method is used to measure kinetics for the binding of a conjugate of the invention to a presenter protein, in which either (i) the conjugate is immobilized on the chip via fusion tag and a presenter protein is injected over the surface, or (ii) a presenter protein is immobilized on the chip via a fusion tag and a conjugate is injected over the surface.

20 Determination of kinetics of binding between FKBP12-Compound 1 and CEP250 by SPR

This protocol utilizes Surface Plasmon Resonance (SPR) as a method to determine kinetics (K_D , K_a , K_d) for the binding of CEP250 (analyte) to immobilized FKBP12-Compound 1 binary complex (ligand).

25 *Reagents:* Compound 1 in 100% DMSO (in-house), 10 X HBS-P+ buffer (GE Healthcare BR-1006-71), Assay buffer (1 X HBS-P+ buffer, 1% DMSO, 1 μ M Compound 1), 12 x HIS tagged FKBP12 (in-house), CEP250_{29.2} (residues 1982-2231) and CEP250_{11.4} (residues 2134-2231) (in-house).

Equipment: BIACORETM X100 (GE Healthcare)

30 *Supplies:* NTA Sensor chip (GE Healthcare BR-1000-34)

Experimental Protocol: Experiments are performed at 25°C. Stock solution of 12XHIS tagged FKBP12 is diluted to 100 nM in assay buffer containing 1 μ M Compound 1 (1% DMSO final). Approximately 200-400 RU of FKBP12 is immobilized on one of two flow cells of an activated NTA chip. The second flow cell is not activated as a reference for non-specific interaction of the analyte to the sensor chip. Various concentrations of CEP250 (1 nM-1 μ M range), serially diluted into the same assay buffer containing 1 μ M Compound 1 (1% DMSO final), are injected onto the FKBP12 surface and

reference surface at a flow rate of 10 μ l/min. The surface is regenerated between analyte injections with 350 mM EDTA.

5 *Data Fitting:* The BiaEvaluation software program is used for data fitting. All data is reference subtracted against both the reference flow cell and a buffer injection. For kinetic analyses, data is locally fit to a 1:1 interaction model.

10 *Results:* SPR sensorgrams and are shown in FIG. 14. Dissociation constants (K_D) of 5.4 nM and 0.29 nM were determined for the binding of FKBP12/Compound 1 to CEP250_{11.4} and CEP250_{29.2}, respectively.

Example 10: Determination of Kinetics of Binding between Conjugates and Proteins by Biolayer Interferometry

15 Biolayer Interferometry (BLI) is a biophysical technique used to measure the kinetics associated with the binary interaction of either two proteins or a protein to a ligand. Typically, one component of the binary interacting pair is immobilized on a biosensor tip via a fusion tag. Increasing concentrations of the second component (the analyte) are then injected over the biosensor tip for a fixed time. An increase in BLI signal (expressed in optical thickness, nm) during the association phase and decrease in BLI signal during the dissociation phase is indicative of an interaction and can be fit to a binding model to determine 20 associated K_D , K_a , K_d values. In this example, the method is used to measure kinetics for the binding of a conjugate of the invention to a presenter protein, in which either (i) the conjugate is immobilized on the tip via fusion tag and a presenter protein is injected over the surface, or (ii) a presenter protein is immobilized on the tip via a fusion tag and a conjugate is injected over the surface.

25 Determination of kinetics of binding between CYPA-Compound 3 and KRAS_{G12C-GTP} by BLI

This protocol utilizes Biolayer Interferometry (BLI) as a method to determine the dissociation constant (K_D) for the binding of KRAS_{G12C-GTP} (analyte) to immobilized CYPA-Compound 3 binary complex (ligand).

30 *Reagents:* Compound 3 in 100% DMSO (in-house), ForteBio Kinetic Buffer (ForteBio Inc., Menlo Park, CA), Assay Buffer (Kinetic Buffer, 1% DMSO, 2 μ M Compound 3), Avi-tagged CYPA (in-house), KRAS_{G12C-GTP} (residues 1-169) (in-house).

35 *Equipment:* Octet Red 96 instrument (ForteBio Inc., Menlo Park, CA)

Supplies: Streptavidin (SA) biosensors (ForteBio)

40 *Experimental Protocol:* Streptavidin (SA) biosensors were coated in a solution containing 10 μ M Avi-CYPA protein at 25 °C to a loading signal of 0.6 nm. The loading of the protein showed stability over time and an absence of baseline drift. The formation of the ternary complex was evaluated in dose-response experiments with KRAS_{G12C-GTP} protein concentrations starting from 200 μ M in a 1:2 dilution

series. For negative control, sensors coated with Avi-CYPA protein were dipped into wells containing only the screening buffer (supplemented with 2 μ M Compound 3). Corrected binding response sensograms were recorded and analyzed.

5 *Data Fitting:* Analysis on the *FortéBio Octet RED* instrument was performed using the *FortéBio* software. The analysis accounts for non-specific binding, background, and signal drift and minimizes well based and sensor variability. Dose-dependent formation of the ternary complex was observed and the corresponding equilibrium dissociation constants (K_D) were determined.

10 *Results:* Sensogram and steady state fitting curves are shown in FIG. 15. A dissociation constant (K_D) of 44 μ M was determined for the binding of CYPA/Compound 3 to KRAS_{G12C-GTP}.

Example 11: Proteomic Identification of FKBP12 bound Target Proteins for Cross-linking Reagents

15 *Reagents:* Compound in 100% DMSO (in-house), N-terminal biotin-FKBP12 (in-house), HEK293T cell lysate (in-house).

20 *Experimental protocol:* HEK293T cell lysate was prepared using a lysis buffer consisting of 40 mM HEPES, pH 7.3, 120 mM NaCl, 2 mM MgCl₂, 2 mM CaCl₂, 0.5% octyl-β-glucoside, and EDTA free protease inhibitor cocktail (Roche) using sonication (4, 10 second pulses at 20% power) on ice. The lysate was first cleared via centrifugation and the resulting supernatant is passed through a 0.2 mm syringe filter on ice. N-terminal biotin labeled FKBP12 was added to 500ml of the lysate to a final concentration of 4mM, mixed via pipetting, and then compound was added to a final concentration of 10 mM with the reaction being mixed via pipetting. 60mL of 50% slurry agarose-Streptavidin resin (pre-equilibrated in the lysis buffer) was added and the reaction is allowed to proceed at 4 °C with gentle rocking for 1 hour. After incubation, the resin was gently pelleted, washed 4 times with 1 mL of lysis buffer on ice via addition, centrifugation, and aspiration and then washed another 4 times with 1mL of lysis buffer without detergent in the same physical manner. Retained proteins were eluted from the resin using 8M Urea, pH 8.0 in HEPES buffer, diluted to 7M Urea with 100 mM HEPES, pH 8.0 and 25 Endoproteinase Lys-C added for protein digestion at 37°C for 2 hours. Next, the sample was diluted to 0.8M Urea using 100 mM HEPES, pH8.0, Trypsin was added, and the sample digested for an additional 16 hours at 37 °C. After digestion was complete, the sample was prepared for LC-MS/MS analysis using a C18 SPE filter onto which the sample was loaded, washed, eluted, desiccated in a speed-vac, and finally suspended in 10ml of 5% acetonitrile, 5% formic acid buffer for LC-MS/MS analysis. LC-MS/MS 30 analysis was performed on a Thermo-Fisher LTQ-Velos-Pro OrbiTrap mass spectrometer using a top 20 data dependent acquisition method and 8-35% acetonitrile gradient for the HPLC. Peptide sequences 35 were assigned using the Sequest algorithm and identified proteins are compared to control samples (DMSO only) in order to identify candidate target proteins.

40 *Results:* Using the above protocol, >100 target proteins have been identified as being capable of binding to a presenter protein in the presence of a cross-linking compound. The identified target proteins


include kinases, phosphatases, ubiquitin ligases, DNA binding proteins, heat shock proteins, DNA helicases, GTPase activating proteins, nucleotide binding proteins, and miscellaneous protein binding proteins.

5 **Example 12: Determination of Binding between Conjugates and Proteins by Fluorescence Polarization**

The technique of fluorescence polarization (FP) is based on the observation that when a fluorescently labeled molecule is excited by polarized light, it emits light with a degree of polarization that is inversely proportional to the rate of molecular rotation. Small molecules rotate quickly during the 10 excited state, and upon emission, have low polarization values. Large complexes, formed by binding of a labeled molecule to a second molecule, rotate little during the excited state, and therefore have high polarization values. This property of fluorescence can be used to measure the interaction of a labeled ligand with a larger protein and provides a basis for direct and competition binding assays. In this 15 example, the method is used to measure the binding of compound or conjugate of the invention to the presenter protein and establish ternary complex formation with a target protein.

Determination of CypA:C3DS:KRAS complex formation by FP

Reagents: C3DS in 100% DMSO (in-house), Protein Buffer (12.5 mM HEPES pH = 7.4, 1 mM 20 MgCl₂), assay buffer (25 mM HEPES, pH 7.3, 0.002% Tween 20, 0.1% BSA, 10 mM NaCl, 1 mM MgCl₂), CYPA (in-house), Mant-GMP-PNP loaded KRAS (1-169 residues).



Equipment: SpectraMax

25 *Experimental Protocol:* KRAS stock solution is loaded to final concentration of 0.8 μM in assay buffer (1% DMSO final). Compound (C3DS) is added to a final concentration of 10 μM and the reaction mixture is dispensed to a 384 well Costar black plate. CYPA is serially diluted into the wells of the plate and allowed to incubate for 15 mins at room temperature. A control experiment in the absence of compound is also run to determine the association of the CYPA to KRAS in the absence of compound. 30 The reaction mixtures are excited at 355nm and the emission signal is recorded at 455 nm. The signals are measured at perpendicular and parallel planes and the polarization is recorded using the following equation.

35
$$FP \text{ (polarization units} \times 10^{-3}) = \frac{\text{Signal (Parallel)} - \text{Signal(Perpendicular)}}{[\text{Signal (Parallel)} + \text{Signal(Perpendicular)}]}$$

Results: A representative curve and is shown in FIG. 16 and a table listing the EC50 (concentration required to enhance the FP signal of the KRAS by 50%) is listed below. The curves were fit to a four-parameter equation and the EC50s obtained indicate the effect of the ligand C3DS towards enhancing the binding between CYPA and KRAS.

5

	CypA:Kras	CypA:C3DS:Kras
EC50	30 μ M	2.3 μ M

Example 13: Determination of Binding between Conjugates and Proteins by Nuclear Magnetic Resonance

Nuclear Magnetic Resonance (NMR) spectroscopy is a technique used to solve three-dimensional structures and study the dynamics of proteins and protein-ligand complexes. In addition, it can be used to identify the ligand binding site in protein-ligand interaction. Out of several available NMR approaches, protein structure based ligand screening (highly sensitive 2D 1 H- 15 N TROSY-HSQC spectrum) and identification of critical residues involved in ligand (drug) binding is the most sensitive method for such studies. Addition of sequentially increasing ligand concentration into protein's NMR sample and collection of 2D 1 H- 15 N TROSY-HSQC provides atomic level highly resolved residue perturbation information, called chemical shift perturbation (CSP), that directly provides more accurate information on identification of ligand binding site not possible by any other biophysical techniques available. With this approach weak, intermediate, and strong affinity of ligand binding to a protein or binary protein complex can be studied, and this information can be directly linked to existing structural, dynamic, and kinetic information. In this example, the method is used to demonstrate binding (e.g. non-covalent or covalent) of a compound (drug) or conjugate of the invention to a presenter protein.

Determination of KRAS(G12C)-Cyclophilin-Compound 3 binding in ternary complex by solution NMR spectroscopy

25 *Reagents:* Compound 3 in 100% DMSO (in-house), Protein Buffer (50 mM TRIS-d₁₁, 50 mM NaCl, pH 7.0, 1 mM TCEP-d₁₆, 1 mM MgCl₂), Additives in NMR sample of KRAS (100 μ M DSS in 93% H₂O and 7% D₂O), assay buffer (protein buffer + increasing equivalents of drug in DMSO (\leq 5%)), GMP- 15 N-KRAS(G12C) -16 (N-His, residues 1-169, in-house), unlabeled (UL) Cyclophilin (CYPA; residues 1-165) (in-house).

30

Equipment: Bruker Avance 800 MHz Spectrometer equipped with 5mm CPTCI 1 H- 13 C/ 15 N/D Z-GRD Z44909/0026 cryoprobe (Bruker). High precision 5 mm NMR tubes are used in these experiments.

NMR Data processing and analysis: Linux computers running Topspin v3.1 and NMRPipe/NMRDraw for processing, and CCPNMR “analysis” program for data analysis.

35

Experimental protocol: GMP- 15 N-KRAS(G12C) -16 stock solution of 0.72 mM in protein buffer was used to prepare 0.18 mM NMR sample in 600 μ l (including NMR additives). DSS was used as internal standard (1 H peak at 0.0 ppm) for chemical shift referencing. 2D 1 H- 15 N TROSY-HSQC spectrum of 15 N KRAS was collected (data size 2048x128). One equivalent (0.18 mM) of CYPA in protein buffer

was added (from stock solution of 0.4 mM) into NMR sample and agitated for 10 minutes. Final NMR sample volume was maintained to 600 μ l. 2D ^1H - ^{15}N TROSY-HSQC spectrum of binary complex (^{15}N -KRAS + UL-CYPA) was collected (data size 2048x128) keeping other acquisition parameters the same (only KRAS ^1H - ^{15}N correlation crosspeaks are visible in the spectrum). A stock solution of 20 mM

5 Compound 3 in 100% DMSO was used for NMR titrations. Compound 3 was sequentially added into NMR sample to obtain its 0.5, 1.0, 2.5, and 5.0 equivalents (to that of ^{15}N -KRAS concentration) in the NMR sample of binary complex (^{15}N -KRAS + UL-CYPA). At each stages sample volume of 600 μ l was maintained while keeping the acquisition parameters the same. At each stages of Compound 3 addition, 2D ^1H - ^{15}N TROSY-HSQC spectrum was acquired to investigate chemical shift perturbation (CSP) of

10 KRAS residues. All spectra were superimposed to each other. Effective CSP at each Compound 3 titration point is determined using the difference of chemical shifts of each residue of KRAS in ternary complex (KRAS+CYPA+ Compound 3) versus the binary complex (KRAS+CYPA). Subsequently, weighted average chemical shift ($\Delta\delta_{\text{weighted}}$) of each KRAS residue are determined using the below formula:

15
$$\Delta\delta_{\text{weighted}} = [(\Delta^1\text{H})^2 + (\Delta^{15}\text{N}/5)^2]^{1/2}$$

Residues eliciting $\Delta\delta_{\text{weighted}}$ greater than one standard deviation from overall average are considered significantly perturbed and used in the binding site mapping. In separate titration experiments, we have collected 2D ^1H - ^{15}N TROSY-HSQC spectra on binary complex (KRAS+CYPA) by sequentially adding 20 DMSO equivalents (to meet equivalent solvent concentration as in above experiment) to subtract contribution from DMSO addition. In second control experiment, we have collected series of 2D ^1H - ^{15}N TROSY-HSQC spectra of ^{15}N -KRAS titrated with Compound 3 at different equivalents (in absence of CYPA).

The effective CSP is tabulated and analyzed. Drug binding residues of KRAS (in presence of 25 CYPA) is mapped onto the protein surface. Dissociation constant, K_D , is determined.

Experimental and processing parameters:

Spectrum data size: 2048 (1H dimension) x128 (15N-dimension)

Number of scans: 4

Temperature: 298 K

5 Quadrature Detection Mode: DQD (1H) and Echo-AntiEcho (15N)

Data sizes were extended by applying forward-backward linear prediction within the indirect dimension.

Data sets were extrapolated by zero filling once in each dimension prior to Fourier transformation.

Results: 2D 1H-15N TROSY-HSQC spectrum of KRAS_{G12C-GTP} is shown in FIG. 17A. Adding a

10 stoichiometric amount of CYPA has no effect on KRAS amide backbone crosspeaks (FIG. 17B),

indicating that KRAS and CYPA do not interact directly. Titration of W21487 into a 1:1 sample of CYPA:KRAS elicits distinct chemical shifts (FIG. 17C), indicative of a direct interaction with KRAS.

Example 14: Determination of Binding between Conjugates and Proteins by Microscale**Thermophoresis**

Microscale Thermophoresis (MST) is a technique for characterization of biomolecular interactions by correlating any changes in the molecular properties of the molecule such as size, conformation to its mobility in a directed temperature gradient. The generation of the gradient is induced by an infrared laser. The movement of the biomolecule is frequently characterized by labeling the molecule using covalently attached fluorophores or even intrinsic fluorescence. In this example, the method is used to measure the binding of compound or conjugate of the invention to the presenter protein and establish ternary complex formation with a target protein, in which either (i) the conjugate is labelled with a fluorophore and a presenter protein is titrated in, or (ii) a presenter protein is labelled with a fluorophore and a conjugate is titrated in.

25

Example 15: Determination of Binding between Conjugates and Proteins by Second Harmonic Generation Technology

Second Harmonic Generation (SHG) is an optical phenomenon that can be used to measure conformational changes in aqueous solution in real time. SHG signal intensity is sensitive to average angular orientation of dye labeled to a protein tethered to a surface and magnitude of signal change directly correlates to amount of angular change. Different conformations can be classified by magnitude of signal change upon binding, signal relative to baseline (more vertical orientations relative to surface produce positive signal changes and vice versa) and kinetics. In this example, the method is used to measure the binding of compound or conjugate of the invention to the presenter protein and establish ternary complex formation with a target protein, in which either (i) a dye labelled conjugate is immobilized on the surface via a fusion tag and a presenter protein is injected over the surface, or (ii) a dye labelled presenter protein is immobilized on the surface via a fusion tag and a conjugate is injected over the surface.

Example 16: Determination of Binding between Conjugates and Proteins by Differential Scanning Fluorimetry

Differential Scanning Fluorimetry (DSF) is a solution based biophysical technique used to measure the melting temperature (T_m) of a protein. In a typical experiment, protein of interest is subject to increasing heat (typically from 4°C-95°C) in the presence of a fluorescent dye (e.g. SYPRO orange). Fluorescent intensities are plotted as a function of temperature and the T_m is calculated from the negative derivative minimum of the fluorescence signal. For a target protein, the thermal shift (ΔT_m) in the presence of a small molecule can be measured to assess whether the small molecule binds to and stabilizes the protein. In this example, the method is used to measure the thermal shift (e.g., non-covalent or covalent binding) of a compound or conjugate of the invention to a presenter protein, in which either (i) the conjugate is labelled with a fluorescent dye and a presenter protein is titrated in, or (ii) a presenter protein is labelled with a fluorescent dye and a conjugate is titrated in.

Example 17: Determination of Binding between Conjugates and Proteins by NanoDSF

NanoDSF is an advanced DSF method for measuring the T_m of a protein using intrinsic tryptophan or tyrosine fluorescence. In a typical experiment, protein of interest is subject to increasing heat (typically from 4°C-95°C) and fluorescent intensities of intrinsic tryptophan or tyrosine residues are monitored as a function of temperature. T_m can be calculated from the changes in tryptophan fluorescence intensity, or from the ratio of tryptophan emission at 330 and 350 nm, which describes the shift of tryptophan emission upon unfolding. For a target protein, ΔT_m in the presence of a small molecule can be measured to assess whether the small molecule binds to and stabilizes the protein. In this example, the method is used to measure the thermal shift (e.g., non-covalent or covalent binding) of a compound or conjugate of the invention to a presenter protein, in which either (i) the fluorescence of the conjugate is measured and a presenter protein is titrated in, or (ii) the fluorescence of a presenter protein is monitored and a conjugate is titrated in.

Example 18: Determination of Complex Formation by Differential Light Scattering

Dynamic light scattering (DLS) is an established biophysics method used to measures time-dependent fluctuations in the scattering intensity arising from particles undergoing random Brownian motion. Diffusion coefficient and particle size information can be obtained from the analysis of these fluctuations. More specifically, the method provides the ability to measure size characteristics including radius and molecular weight of proteins in aqueous solution. In this example, the method is used to measure change in radius or molecular weight in either (i) the presenter protein upon binding of conjugate of the invention or (ii) the conjugate of the invention upon binding to a presenter protein.

Example 19: Determination of Binding between Conjugates and Proteins by Sonic Wave Acoustic Technology

Surface Acoustic Wave (SAW) technology is a biophysical method used for the real-time detection of binding-induced conformational changes through monitoring the shift in the phase of surface acoustic waves that travel along the biosensor. It can be used to measure the kinetics associated with

the binary interaction of either two proteins or a protein to a ligand. Typically, one component of the binary interacting pair is immobilized on the biosensor via a fusion tag. Increasing concentrations of the second component (the analyte) are then injected over the biosensor for a fixed time. An increase in signal (measured either through a change in wave phase or amplitude) during the association phase and 5 decrease in signal during the dissociation phase is indicative of an interaction and can be fit to a binding model to determine associated K_D , K_a , K_d values. In this example, the method is used to measure kinetics for the binding of a conjugate of the invention to a presenter protein, in which either (i) the conjugate is immobilized on the biosensor chip via fusion tag and a presenter protein is injected over the surface, or (ii) a presenter protein.

10 **Example 20: Determination of Complex Formation by Small-Angle X-Ray Scattering**

Small-Angle X-Ray Scattering (SAXS) is a solution based method used to determine the structure of a protein in terms of average particle size and shape. It is capable of delivering structural information in the resolution range between 1 and 25 nm, and of repeat distances in partially ordered systems of up to 150 nm in size. Ultra small-angle scattering (USAS) can resolve even larger dimensions. In a typical 15 scattering experiment, a solution of protein or protein complex are exposed to X-rays (with wavelength λ typically around 0.15 nm). The scattered intensity $I(s)$ is recorded as a function of momentum transfer s ($s=4\pi\sin\theta/\lambda$, where 2θ is the angle between the incident and scattered radiation). From the intensity of the solution the scattering from only the solvent is subtracted. An X-ray scattering curve (intensity versus scattering angle) is then used to create a low-resolution model of a protein or protein complex. In this 20 example, the method is used to identify existence of a ternary complex (e.g., non-covalent or covalent binding) of a compound or conjugate of the invention to a presenter protein.

Other Embodiments

It is to be understood that while the present disclosure has been described in conjunction with the 25 detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the present disclosure, which is defined by the scope of the appended claims. Other aspects, advantages, and alterations are within the scope of the following claims.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments in accordance with the invention 30 described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the appended claims.

In the claims, articles such as "a," "an," and "the" may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the 35 group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

It is also noted that the term "comprising" is intended to be open and permits but does not require the inclusion of additional elements or steps. When the term "comprising" is used herein, the term "consisting of" is thus also encompassed and disclosed.


Where ranges are given, endpoints are included. Furthermore, it is to be understood that unless 5 otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or subrange within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

In addition, it is to be understood that any particular embodiment of the present invention that falls 10 within the prior art may be explicitly excluded from any one or more of the claims. Since such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the compositions of the invention (e.g., any polynucleotide or protein encoded thereby; any method of production; any method of use) can be excluded from any one or more claims, for any reason, whether or not related to the existence of prior 15 art.

What is claimed is:

CLAIMS

1. A compound comprising a protein binding moiety and a cross-linking group.
2. The compound of claim 1, wherein said cross-linking group is a moiety capable of a chemoselective reaction with an amino acid.
3. The compound of claim 1 or 2, wherein said cross-linking group is a sulphydryl-reactive cross-linking group, an amino-reactive cross-linking group, a carboxyl-reactive cross-linking group, a carbonyl-reactive cross-linking group, or a triazole-forming cross-linking group.
4. The compound of claim 3, wherein said cross-linking group is a sulphydryl-reactive cross-linking group.
5. The compound of any one of claims 1 to 4, wherein said cross-linking group comprises a mixed disulfide.
6. The compound of claim 5, wherein said cross-linking group comprises the structure of Formula I:




Formula I

wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound; and

a is 0, 1, or 2;

R^A is optionally substituted $\text{C}_1\text{-}\text{C}_6$ alkyl, optionally substituted $\text{C}_1\text{-}\text{C}_6$ heteroalkyl, optionally substituted $\text{C}_6\text{-}\text{C}_{10}$ aryl, or optionally substituted $\text{C}_2\text{-}\text{C}_9$ heteroaryl.

7. The compound of claim 6, wherein R^A is optionally substituted $\text{C}_2\text{-}\text{C}_9$ heteroaryl.
8. The compound of claim 7, wherein said optionally substituted $\text{C}_2\text{-}\text{C}_9$ heteroaryl is pyridyl.
9. The compound of claim 6, wherein said cross-linking group comprises the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

10. The compound of claim 6, wherein R^A is optionally substituted $\text{C}_1\text{-}\text{C}_6$ heteroalkyl.


11. The compound of claim 10, wherein said optionally substituted C₁-C₆ heteroalkyl is N,N-dimethyl-ethylene.

12. The compound of claim 6, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

13. The compound of claim 12, wherein said cross-linking group comprises the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

14. The compound of claim 6, wherein R^A is optionally substituted C₁-C₆ alkyl.

15. The compound of claim 10, wherein said optionally substituted C₁-C₆ alkyl is methyl.

16. The compound of claim 14 or 15, wherein a is 2.


17. The compound of claim 6, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

18. The compound of any one of claims 1 to 4, wherein said cross-linking group comprises a maleimide.

19. The compound of claim 18, wherein the cross-linking group comprises the structure of formula Ib, Ic, Id, or Ie:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^A is -C(O)- or -SO₂-;

X^B is -C(O)- or CR^ERF;

R^B and R^C are, independently, hydrogen, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl;

R^D is hydrogen, hydroxyl, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl; and

R^E and R^F are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl.


20. The compound of claim 19, wherein the cross-linking group comprises the structure of Formula Ib.

21. The compound of claim 20, wherein X^A is $-C(O)-$.

22. The compound of claim 20 or 21, wherein X^B is $-C(O)-$.


23. The compound of any one of claims 20 to 22, wherein R^B and R^C are hydrogen or optionally substituted C_1 - C_6 alkyl.

24. The compound of claim 23, wherein the cross-linking group comprises the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

25. The compound of claim 24, wherein the cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

26. The compound of any one of claim 1 to 4, wherein the cross-linking group comprises the structure of formula If, Ig, Ih, or li:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^C is $-C(O)-$ or $-SO_2-$;

X^D is absent, NR^JRK^L , or OR^L ;

R^G , R^H , and R^I are, independently, hydrogen, nitrile, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl; and

R^J , R^K , and R^L are, independently, absent, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl.

27. The compound of claim 26, wherein the cross-linking group comprises the structure of Formula If.


28. The compound of claim 26 or 27, wherein X^D is absent.

29. The compound of any one of claims 26 to 28, wherein R^G , R^H , and R^I are hydrogen.

30. The compound of any one of claims 26 to 29, wherein X^C is $-SO_2-$.


31. The compound of any one of claims 26 to 30, wherein said cross-linking group comprises a vinyl sulfone

32. The compound of claim 26, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

33. The compound of claim 32, wherein said cross-linking group comprises the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

34. The compound of any one of claims 26 to 29, wherein X^C is $-C(O)-$.


35. The compound of any one of claims 26 to 29, wherein said cross-linking group includes a vinyl ketone.

36. The compound of claim 35, wherein said cross-linking group includes the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

37. The compound of claim 36, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.


38. The compound of claim 36, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

39. The compound of any one of claims 26 to 29, wherein said cross-linking group comprises an ynnone.

40. The compound of claim 39, wherein the cross-linking group comprises a structure of formula I_j or I_k:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;


X^E is absent, $NR^N R^O$, or OR^P ;

R^M is hydrogen, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and

R^N , R^O , and R^P are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.


41. The compound of claim 40, wherein the cross-linking group comprises a structure of formula I.

42. The compound of claim 40 or 41, wherein the cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

43. The compound of any one of claims 1 to 4, wherein the cross-linking group comprises a structure of formula I_m or I_n:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^F is absent, $NR^S R^T$, or OR^U ;

X^G is absent or $-C(O)-$;

Y is a leaving group;

R^Q and R^R are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and

R^S , R^T , and R^U are, independently, absent, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

44. The compound of claim 43, wherein Y is a halogen.

45. The compound of claim 44, wherein said halogen is a chloride or fluoride.

46. The compound of claim 43, wherein Y is a nitrile.


47. The compound of any one of claims 43 to 46, wherein X^F and X^G are absent.

48. The compound of any one of claims 43 to 47, wherein R^Q and R^R are hydrogen.

49. The compound of claim 43, wherein said cross-linking group comprises an alkyl halide.


50. The compound of claim 49, wherein said alkyl halide is an alkyl chloride or alkyl fluoride.

51. The compound of claim 50, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.


52. The compound of claim 51, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

53. The compound of any one of claims 1 to 4, wherein said cross-linking group comprises an epoxide.


54. The compound of claim 53, wherein said cross-linking group comprises a structure of formula I_o:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

R^V, R^W, and R^X are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

55. The compound of any one of claims 1 to 4, wherein said cross-linking group comprises a structure of formula I_p:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

the dotted lines represent optional double bonds included as necessary for the structure to be aromatic;

b is 0, 1, or 2;

Y is a leaving group;

R^Y and R^Z are, independently, hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl;

each of X^H, X^I, X^J, X^K, and X^L are, independently, absent, NR^{AA}, or CR^{AB}, wherein at least five of X^H, X^I, X^J, X^K, and X^L are NR^{AA}, or CR^{AB};

R^{AA} is absent or hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and


R^{AB} is hydrogen, nitrile, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

56. The compound of claim 55, wherein at least one R^{AB} is an electron withdrawing group.

57. The compound of claim 56, wherein one to three R^{AB} are electron withdrawing groups.


58. The compound of any one of claims 55 to 57, wherein Y is a nitrile, a halogen, a mesylate, a tosylate, or a triflate.

59. The compound of any one of claims 55 to 58, wherein said cross-linking group comprises the structure:




wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

60. The compound of claim 59, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.


61. The compound of claim 60, wherein said cross-linking group comprises the structure:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound.

62. The compound of any one of claims 1 to 4, wherein said cross-linking group is an internal cross-linking group.

63. The compound of claim 62, wherein said cross-linking group comprises a structure of formula Iq, Ir, or Is:



wherein the wavy line illustrates the point of attachment of the cross-linking group to the remainder of the compound;

X^M is $-\text{C}(\text{O})-$ or $-\text{SO}_2-$;

X^N is absent, NR^{AE} , or O ;

R^{AC} and R^{AD} are, independently, hydrogen, nitrile, halogen, optionally substituted hydroxyl, optionally substituted amino, optionally substituted $\text{C}_1\text{-C}_6$ alkyl, optionally substituted $\text{C}_2\text{-C}_6$ alkenyl, optionally substituted $\text{C}_2\text{-C}_6$ alkynyl, optionally substituted $\text{C}_3\text{-C}_{10}$ carbocyclyl, optionally substituted $\text{C}_6\text{-C}_{10}$ aryl, optionally substituted $\text{C}_6\text{-C}_{10}$ aryl $\text{C}_1\text{-C}_6$ alkyl, optionally substituted $\text{C}_2\text{-C}_9$ heteroaryl, optionally substituted $\text{C}_2\text{-C}_9$ heteroaryl $\text{C}_1\text{-C}_6$ alkyl, optionally substituted $\text{C}_2\text{-C}_9$ heterocyclyl, or optionally substituted $\text{C}_2\text{-C}_9$ heterocyclyl $\text{C}_1\text{-C}_6$ alkyl; and

R^{AE} is hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted $\text{C}_1\text{-C}_6$ alkyl, optionally substituted $\text{C}_2\text{-C}_6$ alkenyl, optionally substituted $\text{C}_2\text{-C}_6$ alkynyl, optionally substituted $\text{C}_1\text{-C}_6$ heteroalkyl, optionally substituted $\text{C}_2\text{-C}_6$ heteroalkenyl, optionally substituted $\text{C}_2\text{-C}_6$ heteroalkynyl,

optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl.

64. The compound of claim 63, wherein X^N is NR^{AE}, wherein R^{AE} is hydrogen.

65. The compound of claim 63 or 64, wherein R^{AC} and R^{AD} are hydrogen.

66. The compound of any one of claims 63 to 65, wherein X^M is -C(O)-.

67. The compound of any one of claims 63 to 65, wherein X^M is -SO₂-.

68. The compound of any one of claims 1 to 67, wherein the interaction between said protein binding moiety and a protein is non-covalent.

69. The compound of any one of claims 1 to 67, wherein the interaction between said protein binding moiety and a protein is covalent.

70. A compound comprising a presenter protein binding moiety and a cross-linking group.

71. The compound of claim 70, wherein said presenter protein binding moiety is capable of binding a protein encoded by any one of the genes of Table 1.

72. The compound of claim 70, wherein said presenter protein binding moiety is a prolyl isomerase binding moiety.

73. The compound of any one of claims 70 to 72, wherein said presenter protein binding moiety is a FKBP binding moiety, a cyclophilin binding moiety, or a PIN1 binding moiety.


74. The compound of claim 73, wherein said presenter protein binding moiety is a FKBP binding moiety.

75. The compound of claim 74, wherein said presenter protein binding moiety is capable of binding FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52.

76. The compound of claim 74 or 75, wherein said FKBP binding moiety is a selective FKBP binding moiety.

77. The compound of claim 74 or 75, wherein said FKBP binding moiety is a non-selective FKBP binding moiety.

78. The compound of any one of claims 74 to 77, wherein said FKBP binding moiety comprises the structure of Formula IIa or IIb:



wherein Z¹ and Z² are each, independently, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, or Z¹ and Z² combine to form, with the atoms to which they are attached, an optionally substituted 10 to 40 member macrocycle; and wherein at least one of Z¹ or Z² comprises a point of attachment to the cross-linking group;

b and c are independently 0, 1, or 2;

d is 0, 1, 2, 3, 4, 5, 6, or 7;


X^1 and X^2 are each, independently, absent, CH_2 , O , S , SO , SO_2 , or NR^4 ;

each R¹ and R² are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl, or R¹ and R² combine with the carbon atom to which they are bound to form C=O or R¹ and R² combine to form an optionally substituted C₃-C₁₀ carbocyclyl or optionally substituted C₂-C₉ heterocyclyl;

each R³ is, independently, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl or two R⁸ combine to form an optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, or optionally substituted C₂-C₉ heteroaryl; and

each R⁴ is, independently, hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted aryl, C₃-C₇ carbocyclyl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, and optionally substituted C₃-C₇ carbocyclyl C₁-C₆ alkyl.

79. The compound of claim 78, wherein said presenter protein binding moiety comprises the structure:




80. The compound of claim 73, wherein said presenter protein binding moiety is a cyclophilin binding moiety.

81. The compound of claim 80, wherein said presenter protein binding moiety is capable of binding PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, or PPWD1.

82. The compound of claim 80 or 81, wherein said cyclophilin binding moiety is a selective cyclophilin binding moiety.

83. The compound of claim 80 or 81, wherein said cyclophilin binding moiety is a non-selective cyclophilin binding moiety.

84. The compound of any one of claims 80 to 83, wherein said cyclophilin binding moiety comprises the structure of Formula III or IV:



Formula III

Formula IV

wherein Z^3 , Z^4 , Z^5 , and Z^6 are each, independently, hydroxyl, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, or Z^3 and Z^4 or Z^5 and Z^6 combine to form, with the atoms to which they are attached, an optionally substituted 10 to 40 member macrocycle;

at least one of Z^3 , Z^4 , Z^5 , Z^6 , or R^5 comprises a point of attachment to the cross-linking group; e is 0, 1, 2, 3, or 4;


R^5 is optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl;

R^6 is optionally substituted C_1 - C_6 alkyl;

each R^7 is, independently, hydroxyl, cyano, optionally substituted amino, halogen, thiol, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl, or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl; and

R^8 is hydrogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted aryl, C_3 - C_7 carbocyclyl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, and optionally substituted C_3 - C_7 carbocyclyl C_1 - C_6 alkyl.


85. The compound of claim 84, wherein the cyclophilin binding moiety includes the structure of Formula IVa:



Formula IVa

wherein each R^7 is, independently, hydroxyl, cyano, optionally substituted amino, halogen, thiol, optionally substituted C_1 - C_6 alkyl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 alkynyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_2 - C_6 heteroalkenyl, optionally substituted C_2 - C_6 heteroalkynyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_6 - C_{10} aryl C_1 - C_6 alkyl, optionally substituted C_2 - C_9 heterocyclyl (e.g., optionally substituted C_2 - C_9 heteroaryl), or optionally substituted C_2 - C_9 heterocyclyl C_1 - C_6 alkyl (e.g., optionally substituted C_2 - C_9 heteroaryl C_1 - C_6 alkyl).

86. The compound of claim 84 or 85, wherein said presenter protein binding moiety comprises the structure:



87. A compound comprising a target protein binding moiety and a cross-linking group.

88. The compound of claim 87, wherein said target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs.


89. The compound of claim 87 or 88, wherein said target protein comprises an undruggable surface.

90. The compound of any one of claims 87 to 89, wherein said target protein does not have a traditional binding pocket.

91. The compound of any one of claims 1 to 90, wherein said protein binding moiety and said cross-linking group are joined through a linker.

92. The compound of claim 91, wherein said linker is 1 to 20 atoms in length.


93. The compound of claim 91 or 92, wherein said linker has the structure of Formula V:



Formula V

wherein A^1 is a bond between the linker and protein binding moiety; A^2 is a bond between the cross-linking group and the linker; B^1 , B^2 , B^3 , and B^4 each, independently, is selected from optionally substituted C_{1-2} alkyl, optionally substituted C_{1-3} heteroalkyl, O, S, and NR^N ; R^N is hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted C_{2-4} alkenyl, optionally substituted C_{2-4} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, or optionally substituted C_{1-7} heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f , g , h , i , j , and k are each, independently, 0 or 1; and D is optionally substituted C_{1-10} alkyl, optionally substituted C_{2-10} alkenyl, optionally substituted C_{2-10} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, optionally substituted C_{2-10} polyethylene glycol, or optionally substituted C_{1-10} heteroalkyl, or a chemical bond linking $A^1-(B^1)_f-(C^1)_g-(B^2)_h-$ to $-(B^3)_i-(C^2)_j-(B^4)_k-A^2$

94. The compound of any one of claims 91 to 93, wherein said linker comprises the structure of Formula VI:



Formula VI

wherein A^1 is a bond between the linker and protein binding moiety;

A^2 is a bond between the cross-linking group and the linker;

f is 0, 1, 2, or 3;


g is 0 or 1;

h is 0, 1, or 2; and


X^3 , X^4 , and X^5 are each, independently, absent, O, S, $-C\equiv C-$, CR^9R^{10} or NR^{11} ; and


each R^9 , R^{10} , and R^{11} are, independently, hydrogen, optionally substituted C_{1-6} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted aryl, C_{3-7} carbocycl, optionally substituted C_{6-10} aryl C_{1-6} alkyl, and optionally substituted C_{3-7} carbocycl C_{1-6} alkyl.

95. The compound of claim 94, wherein said linker comprises the structure:



96. A compound having the structure:





97. A conjugate comprising a presenter protein binding moiety conjugated to a target protein.

98. The conjugate of claim 97, wherein the presenter protein binding moiety portion of said conjugate is capable of non-covalent interaction with a presenter protein.

99. The conjugate of claim 97, wherein the presenter protein binding moiety portion of said conjugate is capable of covalent interaction with a presenter protein.

100. The conjugate of any one of claims 97 to 99, wherein said presenter protein binding moiety is capable of binding a protein encoded by any one of the genes of Table 1.

101. The conjugate of any one of claims 97 to 99, wherein said presenter protein binding moiety is a prolyl isomerase binding moiety.

102. The conjugate of any one of claims 97 to 101, wherein said presenter protein binding moiety is a FKBP binding moiety, a cyclophilin binding moiety, or a PIN1 binding moiety.


103. The conjugate of claim 102, wherein said presenter protein binding moiety is a FKBP binding moiety.

104. The conjugate of claim 103, wherein said presenter protein binding moiety is capable of binding FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52.

105. The conjugate of claim 103 or 104, wherein said FKBP binding moiety is a selective FKBP binding moiety.

106. The conjugate of claim 103 or 104, wherein said FKBP binding moiety is a non-selective FKBP binding moiety.

107. The conjugate of any one of claims 103 to 106, wherein said FKBP binding moiety comprises the structure of Formula IIa or IIb:



wherein Z¹ and Z² are each, independently, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, or Z¹ and Z² combine to form, with the atoms to which they are attached, an optionally substituted 10 to 30 member macrocycle; and wherein at least one of Z¹ or Z² comprises a point of attachment to the target protein;

b and c are independently 0, 1, or 2;

d is 0, 1, 2, 3, 4, 5, 6, or 7;


X¹ and X² are each, independently, absent, CH₂, O, S, SO, SO₂, or NR¹³;

each R¹ and R² are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀

aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl, or R¹ and R² combine with the carbon atom to which they are bound to form C=O or R¹ and R² combine to form an optionally substituted C₃-C₁₀ carbocyclyl or optionally substituted C₂-C₉ heterocyclyl; and

each R³ is, independently, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl or two R⁸ combine to form an optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, or optionally substituted C₂-C₉ heteroaryl.

108. The conjugate of claim 107, wherein said presenter protein binding moiety comprises the structure:




109. The compound of claim 102, wherein said presenter protein binding moiety is a cyclophilin binding moiety.

110. The compound of claim 109, wherein said presenter protein binding moiety is capable of binding PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, or PPWD1.

111. The compound of claim 109 or 110, wherein said cyclophilin binding moiety is a selective cyclophilin binding moiety.

112. The compound of claim 109 or 110, wherein said cyclophilin binding moiety is a non-selective cyclophilin binding moiety.

113. The compound of any one of claims 109 to 112, wherein said cyclophilin binding moiety comprises the structure of Formula III or IV:



Formula III

Formula IV

wherein Z³, Z⁴, Z⁵, and Z⁶ are each, independently, hydroxyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, or Z³ and Z⁴ or Z⁵ and Z⁶ combine to form, with the atoms to which they are attached, an optionally substituted 10 to 40 member macrocycle;

at least one of Z³, Z⁴, Z⁵, Z⁶, or R⁵ comprises a point of attachment to the cross-linking group;

d is 0, 1, 2, 3, or 4;


R⁵ is optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl;

R⁶ is optionally substituted C₁-C₆ alkyl;

each R⁷ is, independently, hydroxyl, cyano, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl; and

R⁸ is hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted aryl, C₃-C₇ carbocyclyl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, and optionally substituted C₃-C₇ carbocyclyl C₁-C₆ alkyl.

114. The compound of claim 113, wherein said presenter protein binding moiety comprises the structure:



115. The conjugate of any one of claims 97 to 114, wherein said target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs.

116. The conjugate of any one of claims 97 to 115, wherein said target protein comprises an undruggable surface.

117. The conjugate of any one of claims 97 to 116, wherein said target protein does not have a traditional binding pocket.

118. The conjugate of any one of claims 97 to 117, wherein the amino acid sequence of said target protein has been modified to substitute at least one native amino acid with a reactive amino acid.

119. The conjugate of claim 118, wherein said reactive amino acid is a natural amino acid.

120. The conjugate of claim 119, wherein said reactive amino acid is a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine.

121. The conjugate of claim 118, wherein said reactive amino acid is a non-natural amino acid.

122. The conjugate of any one of claims 97 to 121, wherein the amino acid sequence of said target protein has been modified to substitute at least one native reactive amino acid with a non-reactive amino acid.

123. The conjugate of claim 122, wherein said at least one native reactive amino acid is a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine.

124. The conjugate of claim 122 or 123, wherein said at least one native reactive amino acid is a solvent exposed amino acid.

125. The conjugate of any one of claims 122 to 124, wherein the amino acid sequence of said target protein is modified to substitute all reactive amino acids with a non-reactive amino acid.

126. The conjugate of any one of claims 122 to 125, wherein said non-reactive amino acid is a natural amino acid.

127. The conjugate of any one of claims 118 to 126, wherein said substitution is a conservative substitution.


128. The conjugate of any one of claims 121 to 127, wherein said reactive amino acid is substituted with a serine, valine, alanine, isoleucine, threonine, tyrosine, aspartic acid, glutamic acid, or leucine.

129. The conjugate of any one of claims 121 to 124, wherein said non-reactive amino acid is a non-natural amino acid.

130. The conjugate of any one of claims 97 to 129, wherein said presenter protein binding moiety and said target protein are conjugated through a linker.

131. The conjugate of claim 130, wherein said linker is 1 to 20 atoms in length.

132. The conjugate of claim 130 or 131, wherein said linker has the structure of Formula III:




Formula V

wherein A^1 is a bond between the linker and protein binding moiety; A^2 is a bond between the cross-linking group and the linker; B^1 , B^2 , B^3 , and B^4 each, independently, is selected from optionally substituted C_{1-2} alkyl, optionally substituted C_{1-3} heteroalkyl, O, S, and NR^N ; R^N is hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted C_{2-4} alkenyl, optionally substituted C_{2-4} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, or optionally substituted C_{1-7} heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f , g , h , i , j , and k are each, independently, 0 or 1; and D is optionally substituted C_{1-10} alkyl, optionally substituted

C₂₋₁₀ alkenyl, optionally substituted C₂₋₁₀ alkynyl, optionally substituted C₂₋₆ heterocycl, optionally substituted C₆₋₁₂ aryl, optionally substituted C₂-C₁₀ polyethylene glycol, or optionally substituted C₁₋₁₀ heteroalkyl, or a chemical bond linking A¹-(B¹)_f-(C¹)_g-(B²)_h- to -(B³)_i-(C²)_j-(B⁴)_k-A².


133. The conjugate of any one of claims 130 to 132, wherein said linker comprises the structure of Formula IV:



Formula VI

wherein A¹ is a bond between the linker and protein binding moiety;
 A² is a bond between the cross-linking group and the linker;
 l is 0, 1, 2, or 3;
 m is 0 or 1;
 n is 0, 1, or 2; and
 X³, X⁴, and X⁵ are each, independently, absent, O, S, -C≡C-, CR⁹R¹⁰ or NR¹¹; and
 each R⁹, R¹⁰, and R¹¹ are, independently, hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted aryl, C₃-C₇ carbocycl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, and optionally substituted C₃-C₇ carbocycl C₁-C₆ alkyl.

134. The conjugate of claim 133, wherein said linker comprises the structure:



135. A method of producing a conjugate comprising a presenter protein binding moiety conjugated to a target protein, said method comprising reacting (a) a compound comprising a presenter protein binding moiety and a cross-linking group with (b) a target protein under conditions that permit production of said conjugate.

136. A method of producing a conjugate comprising a presenter protein binding moiety conjugated to a target protein, said method comprising
 providing (a) a compound comprising a presenter protein binding moiety and a cross-linking group; (b) a target protein; and (c) a presenter protein; and
 reacting said compound with said target protein under conditions that permit production of said conjugate.

137. The method of claim 136, wherein said presenter protein binds to said compound in the absence of said target protein.

138. The method of claim 136, wherein said presenter protein does not substantially bind to said compound in the absence of said target protein.

139. The method of any one of claims 136 to 138, wherein said compound and said target protein do not substantially react in the absence of said presenter protein.

140. The method of any one of claims 136 to 138, wherein said compound and said target protein react in the absence of said presenter protein.

141. The method of any one of claims 135 to 140, wherein said conditions do not comprise a reducing reagent.

142. A complex comprising (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein.

143. The complex of claim 142, wherein said presenter protein is a protein encoded by any one of the genes of Table 1.

144. The complex of claim 142, wherein said presenter protein is a prolyl isomerase.

145. The complex of any one of claims 142 to 144, wherein said prolyl isomerase is a member of the FKBP family, a member of the cyclophilin family, or PIN1.

146. The complex of claim 145, wherein said member of the FKBP family is FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52.

147. The complex of claim 145, wherein said member of the cyclophilin family is PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, or PPWD1.

148. A method of producing a complex comprising (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein, said method comprising combining a conjugate comprising a presenter protein binding moiety conjugated to a target protein and a presenter protein under conditions that permit production of said complex.

149. A method of producing a complex comprising (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein, said method comprising

providing (a) a compound comprising a presenter protein binding moiety and a cross-linking group; (b) a target protein; and (c) a presenter protein; and

reacting said compound with said target protein under conditions that permit production of said complex.

150. The method of claim 149, wherein said presenter protein binds to said compound in the absence of said target protein.

151. The method of claim 149, wherein said presenter protein does not substantially bind to said compound in the absence of said target protein.

152. The method of any one of claims 149 to 151, wherein said compound and said target protein do not substantially react in the absence of said presenter protein.

153. The method of any one of claims 149 to 151, wherein said compound and said target protein react in the absence of said presenter protein.

154. The method of any one of claims 148 to 153, wherein said conditions do not comprise a reducing reagent.

155. The method of any one of claims 148 to 154, wherein said conditions comprise an excess of presenter protein.

156. A conjugate comprising a target protein binding moiety conjugated to a presenter protein.

157. The conjugate of claim 156, wherein the target protein binding moiety portion of said conjugate is capable of non-covalent interaction with a target protein.

158. The conjugate of claim 156, wherein the target protein binding moiety portion of said conjugate is capable of covalent interaction with a target protein.

159. The conjugate of any one of claims 156 to 158, wherein said target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs.

160. The conjugate of any one of claims 156 to 159, wherein said target protein comprises an undruggable surface.

161. The conjugate of any one of claims 156 to 160, wherein said target protein does not have a traditional binding pocket.

162. The conjugate of any one of claims 156 to 161, wherein said presenter protein is a protein encoded by any one of the genes of Table 1.

163. The conjugate of any one of claims 156 to 161, wherein said presenter protein is a prolyl isomerase.

164. The conjugate of any one of claims 156 to 163, wherein said presenter protein is a member of the FKBP family, a member of the cyclophilin family, or PIN1.

165. The conjugate of claim 164, wherein said member of the FKBP family is FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52.

166. The conjugate of claim 164, wherein said member of the cyclophilin family is PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, or PPWD1.

167. The conjugate of any one of claims 156 to 166, wherein the amino acid sequence of said presenter protein has been modified to substitute at least one amino acid with a reactive amino acid.

168. The conjugate of claim 167, wherein said reactive amino acid is a natural amino acid.

169. The conjugate of claim 168, wherein said reactive amino acid is a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine.

170. The conjugate of claim 167, wherein said reactive amino acid is a non-natural amino acid.

171. The conjugate of any one of claims 156 to 170, wherein the amino acid sequence of said presenter protein has been modified to substitute at least one native reactive amino acid with a non-reactive amino acid.

172. The conjugate of claim 171, wherein said at least one native reactive amino acid is a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine.

173. The conjugate of claim 171 or 172, wherein said at least one native reactive amino acid is a solvent exposed amino acid.

174. The conjugate of any one of claims 171 to 173, wherein the amino acid sequence of said presenter protein is modified to substitute all reactive amino acids with a non-reactive amino acid.

175. The conjugate of any one of claims 171 to 174, wherein said non-reactive amino acid is a natural amino acid.

176. The conjugate of any one of claims 167 to 175, wherein said substitution is a conservative substitution.


177. The conjugate of any one of claims 171 to 176, wherein said reactive amino acid is substituted with a serine, valine, alanine, isoleucine, threonine, tyrosine, aspartic acid, glutamic acid, or leucine.

178. The conjugate of any one of claims 171 to 174, wherein said non-reactive amino acid is a non-natural amino acid.

179. The conjugate of any one of claims 156 to 178, wherein said target protein binding moiety and said presenter protein are joined through a linker.

180. The conjugate of claim 179, wherein said linker is 1 to 20 atoms in length.


181. The conjugate of claim 179 or 180, wherein said linker has the structure of Formula V:



Formula V

wherein A^1 is a bond between the linker and protein binding moiety; A^2 is a bond between the cross-linking group and the linker; B^1 , B^2 , B^3 , and B^4 each, independently, is selected from optionally substituted C_{1-2} alkyl, optionally substituted C_{1-3} heteroalkyl, O, S, and NR^N ; R^N is hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted C_{2-4} alkenyl, optionally substituted C_{2-4} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, or optionally substituted C_{1-7} heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f , g , h , i , j , and k are each, independently, 0 or 1; and D is optionally substituted C_{1-10} alkyl, optionally substituted C_{2-10} alkenyl, optionally substituted C_{2-10} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, optionally substituted $C_{2-C_{10}}$ polyethylene glycol, or optionally substituted C_{1-10} heteroalkyl, or a chemical bond linking $A^1-(B^1)-f-(C^1)_g-(B^2)_h-$ to $-(B^3)_i-(C^2)_j-(B^4)_k-A^2$.

182. The conjugate of any one of claims 179 to 181, wherein said linker comprises the structure of Formula IV:



Formula VI

wherein A^1 is a bond between the linker and protein binding moiety;

A^2 is a bond between the cross-linking group and the linker;

l is 0, 1, 2, or 3;


m is 0 or 1;

n is 0, 1, or 2; and

X³, X⁴, and X⁵ are each, independently, absent, O, S, -C≡C-, CR⁹R¹⁰ or NR¹¹; and

each R⁹, R¹⁰, and R¹¹ are, independently, hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted aryl, C₃-C₇ carbocyclyl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, and optionally substituted C₃-C₇ carbocyclyl C₁-C₆ alkyl.

183. The conjugate of claim 182, wherein said linker comprises the structure:



184. A method of producing a conjugate comprising a target protein binding moiety conjugated to a presenter protein, said method comprising reacting (a) a compound comprising a target protein binding moiety and a cross-linking group with (b) a presenter protein under conditions that permit production of said conjugate.

185. A method of producing a conjugate comprising a target protein binding moiety conjugated to a presenter protein, said method comprising

providing (a) a compound comprising a target protein binding moiety and a cross-linking group; (b) a presenter protein; and (c) a target protein; and

reacting said compound with said presenter protein under conditions that permit production of said conjugate.

186. The method of claim 185, wherein said target protein binds to said compound in the absence of said presenter protein.

187. The method of claim 185, wherein said target protein does not substantially bind to said compound in the absence of said presenter protein.

188. The method of any one of claims 185 to 187, wherein said compound and said presenter protein do not substantially react in the absence of said target protein.

189. The method of any one of claims 185 to 187, wherein said compound and said presenter protein react in the absence of said target protein.

190. The method of any one of claims 184 to 189, wherein said conditions do not comprise a reducing reagent.

191. A complex comprising (i) a conjugate comprising a target protein binding moiety conjugated to a presenter protein and (ii) a target protein.

192. The complex of claim 191, wherein said target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs.

193. The complex of claim 191 or 192, wherein said target protein comprises an undruggable surface.

194. The complex of any one of claims 191 to 193, wherein said target protein does not have a traditional binding pocket.

195. A method of producing a complex comprising (i) a conjugate comprising a target protein binding moiety conjugated to a presenter protein and (ii) a target protein, said method comprising combining a conjugate comprising a target protein binding moiety conjugated to a presenter protein and a target protein under conditions that permit production of said complex.

196. A method of producing a complex comprising (i) a conjugate comprising a target protein binding moiety conjugated to a presenter protein and (ii) a target protein, said method comprising providing (a) a compound comprising a target protein binding moiety and a cross-linking group; (b) a presenter protein; and (c) a target protein; and reacting said compound with said presenter protein under conditions that permit production of said complex.

197. The method of claim 196, wherein said target protein binds to said compound in the absence of said presenter protein.

198. The method of claim 196, wherein said target protein does not substantially bind to said compound in the absence of said presenter protein.

199. The method of any one of claims 196 to 198, wherein said compound and said presenter protein do not substantially react in the absence of said target protein.

200. The method of any one of claims 196 to 198, wherein said compound and said presenter protein react in the absence of said target protein.

201. The method of any one of claims 195 to 200, wherein said conditions do not comprise a reducing reagent.


202. The method of any one of claims 196 to 201, wherein said conditions comprise an excess of target protein.

203. A compound having the structure of Formula VII:

A-L-B

Formula VII

wherein A comprises the structure of Formula VIIa or VIIb:



Formula VIIa Formula VIIb

wherein b and c are independently 0, 1, or 2;

d is 0, 1, 2, 3, 4, 5, 6, or 7;

X¹ and X² are each, independently, absent, CH₂, O, S, SO, SO₂, or NR¹³;

each R¹ and R² are independently hydrogen, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl, or R¹ and R² combine with the carbon atom to which they are bound to form C=O or R¹ and R² combine to form an optionally substituted C₃-C₁₀ carbocyclyl or optionally substituted C₂-C₉ heterocyclyl; and

each R³ is, independently, hydroxyl, optionally substituted amino, halogen, thiol, optionally substituted C₁-C₆ alkyl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ alkynyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₆ heteroalkenyl, optionally substituted C₂-C₆ heteroalkynyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₆-C₁₀ aryl C₁-C₆ alkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkyl or two R³ combine to form an optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl; and

R⁴ is optionally substituted C₁-C₆ alkyl;

L is an optional linker; and

B is a target protein binding moiety.

204. The compound of claim 203, wherein the interaction between said target protein binding moiety and a target protein is non-covalent.


205. The compound of claim 203, wherein the interaction between said target protein binding moiety and a target protein is covalent.

206. The compound of any one of claims 203 to 205, wherein said target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs.

207. The compound of any one of claims 203 to 206, wherein said target protein comprises an undruggable surface.

208. The compound of any one of claims 203 to 207, wherein said target protein does not have a traditional binding pocket.


209. The compound of any one of claims 203 to 208, wherein A comprises the structure:



210. The compound of any one of claims 203 to 209, wherein said target protein binding moiety and said presenter protein are conjugated through a linker.

211. The compound of claim 210, wherein said linker is 1 to 20 atoms in length.


212. The compound of claim 210 or 211, wherein said linker has the structure of Formula V:



wherein A^1 is a bond between the linker and protein binding moiety; A^2 is a bond between the cross-linking group and the linker; B^1 , B^2 , B^3 , and B^4 each, independently, is selected from optionally substituted C_{1-2} alkyl, optionally substituted C_{1-3} heteroalkyl, O, S, and NR^N ; R^N is hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted C_{2-4} alkenyl, optionally substituted C_{2-4} alkynyl, optionally substituted C_{2-6} heterocycl, optionally substituted C_{6-12} aryl, or optionally substituted C_{1-7} heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f , g , h , i , j , and k are each, independently, 0 or 1; and D is optionally substituted C_{1-10} alkyl, optionally substituted C_{2-10} alkenyl, optionally substituted C_{2-10} alkynyl, optionally substituted C_{2-6} heterocycl, optionally

substituted C₆₋₁₂ aryl, optionally substituted C_{2-C₁₀} polyethylene glycol, or optionally substituted C₁₋₁₀ heteroalkyl, or a chemical bond linking A¹-(B¹)_l-(C¹)_g-(B²)_h to -(B³)_i-(C²)_j-(B⁴)_k-A².

213. The compound of any one of claims 210 to 212, wherein said linker comprises the structure of Formula IV:



Formula VI

wherein A¹ is a bond between the linker and protein binding moiety;

A² is a bond between the cross-linking group and the linker;

l is 0, 1, 2, or 3;


m is 0 or 1;

n is 0, 1, or 2; and

X³, X⁴, and X⁵ are each, independently, absent, O, S, -C≡C-, CR⁹R¹⁰ or NR¹¹; and

each R⁹, R¹⁰, and R¹¹ are, independently, hydrogen, optionally substituted C_{1-C₆} alkyl, optionally substituted C_{2-C₆} alkenyl, optionally substituted C_{2-C₆} alkynyl, optionally substituted aryl, C_{3-C₇} carbocyclyl, optionally substituted C_{6-C₁₀} aryl C_{1-C₆} alkyl, and optionally substituted C_{3-C₇} carbocyclyl C_{1-C₆} alkyl.

214. The compound of claim 213, wherein said linker comprises the structure:



215. A complex comprising (i) a compound of claim 203; (ii) a target protein; and (iii) a presenter protein.

216. The complex of claim 215, wherein said target protein is a GTPase, GTPase activating protein, Guanine nucleotide-exchange factor, a heat shock protein, an ion channel, a coiled-coil protein, a kinase, a phosphatase, a ubiquitin ligase, a transcription factor, a chromatin modifier/remodeler, or a protein with classical protein-protein interaction domains and motifs.

217. The complex of claim 215 or 216, wherein said target protein comprises an undruggable surface.

218. The complex of any one of claims 215 to 217, wherein said target protein does not have a traditional binding pocket.

219. The complex of any one of claims 215 to 218, wherein said presenter protein is a protein encoded by any one of the genes of Table 1.

220. The complex of any one of claims 215 to 218, wherein said presenter protein is a prolyl isomerase.

221. The compound of any one of claims 215 to 220, wherein said presenter protein is a member of the FKBP family, a member of the cyclophilin family, or PIN1.

222. The compound of claim 221, wherein said member of the FKBP family is FKBP12, FKBP12.6, FKBP13, FKBP25, FKBP51, or FKBP52.

223. The complex of claim 221, wherein said member of the cyclophilin family is PP1A, CYPB, CYPC, CYP40, CYPE, CYPD, NKTR, SRCyp, CYPH, CWC27, CYPL1, CYP60, CYPJ, PPIL4, PPIL6, RANBP2, or PPWD1.

224. A method of identifying a conjugate comprising a presenter protein binding moiety conjugated to a target protein capable of forming a complex with a presenter protein, said method comprising:

(a) providing (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein;

(b) combining said conjugate and said presenter protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the presenter protein; and

(c) determining if said conjugate and said presenter protein form a complex,

wherein if said conjugate and said presenter protein form a complex, said conjugate is identified as a conjugate capable of forming a complex with a presenter protein,

thereby identifying a conjugate comprising a presenter protein binding moiety conjugated to a target protein capable of forming a complex with a presenter protein.

225. A method of identifying a target protein which binds to a presenter protein, said method comprising:

(a) providing (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein;

(b) combining said conjugate and said presenter protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the presenter protein; and

(c) determining whether said target protein binds to said presenter protein in said complex,

wherein if said target protein binds to said presenter protein, said target protein is identified as binding to the presenter protein,

thereby identifying a target protein which binds to a presenter protein.

226. A method of identifying a target protein capable of reacting with a compound in the presence of a presenter protein, wherein the compound comprises a presenter protein binding moiety and a cross-linking moiety, said method comprising:

(a) providing (i) a compound comprising a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein;

(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and

(c) determining if said target protein and said compound react during formation of said complex to form a conjugate,

wherein if said target protein and said compound react during formation of said complex to form a conjugate, said target protein is identified as a target protein capable of reacting with the compound in the presence of a presenter protein,

thereby identifying a target protein capable of reacting with a compound comprising a presenter protein binding moiety and a cross-linking moiety in the presence of a presenter protein.

227. A method of identifying a target protein which binds to a presenter protein, said method comprising:

(a) providing (i) a compound comprising a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein;

(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and

(c) determining whether said target protein binds to said presenter protein in said complex,

wherein if said target protein binds to said presenter protein, said target protein is identified as a target protein that binds to a presenter protein,

thereby identifying a target protein which binds to a presenter protein.

228. A method of identifying a target protein capable of forming a complex with a presenter protein, said method comprising:

(a) providing (i) a compound of claim 181; (ii) a target protein; and (iii) a presenter protein;

(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and

(c) determining if said compound, said target protein, and said presenter protein form a complex,

wherein if said compound, said target protein, and said presenter protein form a complex, said target protein is identified as a target protein capable of forming a complex with a presenter protein,

thereby identifying a target protein capable of forming a complex with a presenter protein.

229. A method of identifying a target protein which binds to a presenter protein, said method comprising:

- (a) providing (i) a compound of claim 181; (ii) a target protein; and (iii) a presenter protein;
- (b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and
- (c) determining whether said target protein binds to said presenter protein in said complex, wherein if said target protein binds to said presenter protein, said target protein is identified as a target protein that binds to a presenter protein

thereby identifying a target protein which binds to a presenter protein.

230. A method of identifying a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein, said method comprising:

- (a) providing (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein at a location and (ii) a presenter protein;
- (b) combining said conjugate and said presenter protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the presenter protein;
- (c) determining if said conjugate and said presenter protein form a complex; and
- (d) repeating steps (a) to (c) with said presenter protein binding moiety conjugated at different locations on said target protein until a conjugate and said presenter protein form a complex, wherein a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein is identified if said conjugate and said presenter protein form a complex,

thereby identifying a location on a target protein to form a conjugate capable of forming a complex with a presenter protein.

231. A method of identifying a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein, said method comprising:

- (a) providing (i) a compound comprising a presenter protein binding moiety and a cross-linking group; (ii) a target protein; and (iii) a presenter protein;
- (b) combining said compound with said target protein in the presence of said presenter protein under conditions that permit the formation of a conjugate comprising a presenter protein binding moiety conjugated to a target protein at a location;
- (c) determining if said conjugate and said presenter protein form a complex; and
- (d) repeating steps (a) to (c) wherein said presenter protein binding moiety is conjugated at different locations on said target protein until a conjugate and said presenter protein form a complex, wherein a location on a target protein to form a conjugate capable of forming a complex with a presenter protein is identified if said conjugate and said presenter protein form a complex,

thereby identifying a location on a target protein to form a conjugate with a presenter protein binding moiety, which conjugate is capable of forming a complex with a presenter protein.

232. The method of claims 230 or 231, wherein the amino acid sequence of said target protein has been modified to substitute at least one amino acid with a reactive amino acid.

233. The method of claim 232, wherein said reactive amino acid is a natural amino acid.

234. The method of claim 233, wherein said reactive amino acid is a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine.

235. The method of claim 232, wherein said reactive amino acid is a non-natural amino acid.

236. The method of any one of claims 230 to 235, wherein the amino acid sequence of said target protein has been modified to substitute at least one native reactive amino acid with a non-reactive amino acid.

237. The method of claim 236, wherein said at least one native reactive amino acid is a cysteine, lysine, tyrosine, aspartic acid, glutamic acid, or serine.

238. The method of claim 236 or 237, wherein said at least one native reactive amino acid is a solvent exposed amino acid.

239. The method of any one of claims 236 to 238, wherein the amino acid sequence of said target protein is modified to substitute all reactive amino acids with a non-reactive amino acid.

240. The method of any one of claims 236 to 239, wherein said non-reactive amino acid is a natural amino acid.

241. The method of any one of claims 230 to 240, wherein said substitution is a conservative substitution.

242. The method of any one of claims 232 to 241, wherein said reactive amino acid is substituted with a serine, valine, alanine, isoleucine, threonine, tyrosine, aspartic acid, glutamic acid, or leucine.

243. The method of any one of claims 232 to 242, wherein said non-reactive amino acid is a non-natural amino acid.

244. A method of identifying a compound capable of covalently binding to a target protein in the presence of a presenter protein, said method comprising:

(a) providing a sample comprising (i) a compound comprising a presenter protein binding moiety and a cross-linking group; (ii) a target protein; and (iii) a presenter protein; and

(b) determining if said compound and said target protein form a covalent bond via the cross-linking group of said compound in said sample,

wherein a compound is identified as covalently binding to a target protein in the presence of a presenter protein if said compound and said target protein react in said sample.

245. A method of identifying a compound capable of selective and covalent binding to a target protein in the presence of a presenter protein, said method comprising:

(a) providing a first sample comprising (i) a compound comprising a presenter protein binding moiety and a cross-linking group; (ii) a target protein; and (iii) a presenter protein and a second sample comprising (i) the same compound comprising a presenter protein binding moiety and a cross-linking group as in the first sample and (ii) the same target protein as in the first sample; and

(b) determining the extent to which said compound and said target protein react in said first sample as compared to said second sample,

wherein a compound is identified as selectively covalently binding to a target protein in the presence of a presenter protein if said compound and said target protein reacts in said first sample more than in said second sample.

246. The method of claim 245, wherein a compound is identified as selectively covalently binding to a target protein in the presence of a presenter protein if said compound and said target protein reacts in said first sample at least 5-fold more than in said second sample.

247. The method of claim 245 or 246, wherein a compound is identified as selectively covalently binding to a target protein in the presence of a presenter protein if said compound and said target protein reacts in said first sample, but does not substantially react in said second sample.

248. A method of identifying a conjugate capable of forming a complex with a presenter protein, said method comprising:

(a) providing (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein; and

(b) combining said conjugate and said presenter protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the presenter protein;

(c) determining if said conjugate and said presenter protein form a complex,

wherein a conjugate is identified as capable of forming a complex with a presenter protein if said conjugate and said presenter protein form a complex,

thereby identifying a conjugate capable of forming a complex with a presenter protein.

249. A method of determining the structure of an interface in a complex comprising a presenter protein and a target protein, said method comprising:

(a) providing (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein;

(b) combining said conjugate and said presenter protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the presenter protein; and

(c) determining the crystal structure of said complex,

wherein the structure of the interface comprises at least the portion of the crystal structure between said presenter protein and said target protein,

thereby determining the structure of an interface in a complex comprising a presenter protein and a target protein.

250. A method of determining the structure of an interface in a complex comprising a presenter protein and a target protein, said method comprising:

(a) providing (i) a compound comprising a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein;

(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and

(c) determining the crystal structure of said complex,

wherein the structure of the interface comprises at least the portion of the crystal structure between said presenter protein and said target protein,

thereby determining the structure of an interface in a complex comprising a presenter protein and a target protein.

251. A method of determining the structure of an interface in a complex comprising a presenter protein and a target protein, said method comprising:

(a) providing (i) a compound of claim 203; (ii) a target protein; and (iii) a presenter protein;

(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and

(c) determining the crystal structure of said complex,

wherein the structure of the interface comprises at least the portion of the crystal structure between said presenter protein and said target protein;

thereby determining the structure of an interface in a complex comprising a presenter protein and a target protein.

252. The method of any one of claims 249 to 251, wherein said interface in a complex comprising a presenter protein and a target protein is a binding pocket.

253. A method of obtaining X-ray crystal coordinates for a complex, said method comprising:

(a) providing (i) a conjugate comprising a presenter protein binding moiety conjugated to a target protein and (ii) a presenter protein;

(b) combining said conjugate and said presenter protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the presenter protein; and
(c) determining the crystal structure of said complex,
thereby obtaining X-ray crystal coordinates for said complex.

254. A method of obtaining X-ray crystal coordinates for a complex, said method comprising:

(a) providing (i) a compound comprising a presenter protein binding moiety and a cross-linking moiety; (ii) a target protein; and (iii) a presenter protein;
(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and
(c) determining the crystal structure of said complex,
thereby obtaining X-ray crystal coordinates for said complex.

255. A method of obtaining X-ray crystal coordinates for a complex, said method comprising:

(a) providing (i) a compound of claim 203; (ii) a target protein; and (iii) a presenter protein;
(b) combining said compound, said target protein, and said presenter protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the presenter protein; and
(c) determining the crystal structure of said complex,
thereby obtaining X-ray crystal coordinates for said complex.

256. A method of determining the residues on a target protein that participate in binding with a presenter protein, said method comprising:

(a) providing X-ray crystal coordinates of a complex obtained by a method of any one of claims 253 to 255;
(b) identifying the residues of said target protein which comprise an atom within 4 Å of an atom on said presenter protein;
thereby determining the residues on a target protein that participate in binding with a presenter protein.

257. A method of determining biochemical and/or biophysical properties of a complex of any one of claims 142 to 147, 191 to 194, or 215 to 223, said method comprising:

(a) providing X-ray crystal coordinates of a complex of any one of claims 142 to 147, 191 to 194, or 215 to 223 obtained by a method of any one of claims 253 to 255;
(b) calculating a biochemical and/or biophysical property of said complex;
thereby determining biochemical and/or biophysical properties of a presenter protein/target protein complex.

258. The method of claim 257, wherein said biochemical and/or biophysical properties comprise the free energy of binding of said complex, the K_d of said complex, the K_i of said complex, the K_{inact} of said complex, and/or the K_i/K_{inact} of said complex.

259. The method of claim 258, wherein said biochemical and/or biophysical properties comprise the free energy of binding of said complex.

260. The method of claim 259, wherein said free energy of binding of said complex is determined by isothermal titration calorimetry.

261. The method of claim 258, wherein said biochemical and/or biophysical properties comprise the K_d of said complex.

262. The method of claim 261, wherein said K_d of said complex is determined by surface plasmon resonance.

263. The method of claim 258, wherein said biochemical and/or biophysical properties comprise the K_i of said complex, the K_{inact} of said complex, and/or the K_i/K_{inact} of said complex.

264. The method of claim 263, wherein said K_i of said complex, said K_{inact} of said complex, and/or said K_i/K_{inact} of said complex is determined by mass spectrometry.

265. A composition comprising a compound of any one of claims 1 to 98 or 203 to 214 and a suitable carrier.

266. A pharmaceutical composition comprising a compound of any one of claims 1 to 98 or 203 to 214 and a pharmaceutically acceptable carrier.

267. A method of modulating a target protein, said method comprising contacting said target protein with a modulating amount of a compound of any one of claims 1 to 98 or 203 to 214 or a composition of claims 265 or 266.

268. A method of modulating a target protein, said method comprising forming a complex of any one of claims 142 to 147, 191 to 194, or 215 to 223 in a cell by contacting said cell with an effective amount of any one of claims 1 to 98 or 203 to 214 or a composition of claims 265 or 266.

269. A method of modulating a target protein, said method comprising contacting said target protein with a conjugate of any one of claims 155 to 183.

270. A method of inhibiting prolyl isomerase activity, said method comprising contacting a cell expressing said prolyl isomerase with a compound of any one of claims 1 to 98 or 203 to 214 or a

composition of claims 265 or 266 under conditions that permit the formation of a complex between said compound and said prolyl isomerase, thereby inhibiting prolyl isomerase activity.

271. A method of forming a complex of any one of claims 142 to 147, 191 to 194, or 215 to 223 in a cell, said method comprising contacting a cell expressing a presenter protein with a compound of any one of claims 1 to 98 or 203 to 214 or a composition of claims 265 or 266 under conditions that permit the formation of a complex between said compound and said presenter protein.

272. A method of identifying a target protein capable of forming a complex with a presenter protein, said method comprising:

(a) providing (i) one or more target proteins, (ii) a compound of any one of claims 1-98; and (iii) a presenter protein comprising a tag;

(b) combining said one or more target proteins, said compound, and said presenter protein under conditions suitable to permit complex formation if said target protein is capable of forming a complex with said presenter protein; and

(c) determining whether one or more target proteins form a complex with said compound and said presenter protein;

wherein target proteins that form a complex with said compound and said presenter protein are identified as target proteins capable of forming a complex with a presenter protein.

273. The method of claim 272, wherein said presenter protein comprises an affinity tag.

274. The method of claim 272 or 273, wherein said determining step comprises utilizing said tag of said presenter protein to selectively isolate target proteins which have formed a complex with said presenter protein.

275. The method of any one of claims 272 to 274, wherein said method further comprises (d) identifying said target protein in a complex formed between one or more target proteins and said presenter protein.

276. The method of claim 275, wherein said identifying the structure of said target protein comprises performing mass spectrometry on said complex.

277. A method of identifying a target protein capable of forming a complex with a presenter protein, said method comprising:

(a) providing (i) two or more target proteins; (ii) a compound of any one of claims 1-98; and (iii) a presenter protein comprising an affinity tag;

(b) combining said two or more target proteins, said compound, and said presenter protein under conditions suitable to permit complex formation if said target protein is capable of forming a complex with said presenter protein;

- (c) selectively isolating one or more complexes of a target protein, said compound, and said presenter protein formed in step (b); and
- (d) determining the structure of said target protein in said one or more complexes isolated in step (c) by mass spectrometry;
 - thereby identifying a target protein capable of forming a complex with a presenter protein.

FIG. 1




FIG. 2

Compound (SFAX9DS)





FIG. 3A




FIG. 3B

FIG. 4





FIG. 5A




FIG. 5B




FIG. 6






FIG. 7A**FIG. 7B**







FIG. 8

FIG. 9A**FIG. 9B**




FIG. 11




FIG. 12






FIG. 13








FIG. 14 (continued)

FIG. 15







FIG. 16

Kras (Mant-GMP-PNP) binding to Avi-CypA



Fitting View






FIG. 17A

FIG. 17B





FIG. 17C



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/26014

A. CLASSIFICATION OF SUBJECT MATTER

IPC - G01N 33/68, 33/566; C12Q 1/533; A61K 47/42 (2018.01)
 CPC - G01N 33/6845, 33/566; C12Q 1/533; A61K 47/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2016/112279 A1 (WARP DRIVE BIO, INC.) 14 July, 2016; page 1, lines 25-39; page 18, lines 10-17; page 34, lines 14-15; page 61, lines 3-19; page 64, lines 22-36; page 72, lines 5-18	1-2, 3/1-2, 4/3/1-2, 70, 97-99, 135-138, 139/136-138, 140/136-138, 142, 148-151, 152/149-151, 153/149-151, 224-227, 248-250, 252/249-250, 253-254
---		-----
Y		71, 100/97-99, 143, 230-231, 232/230-231, 233/232/230-231, 234/233/232/230-231, 235/232/230-231, 256/253-254
Y	· (MAJUMDER, A et al.) Interaction of Aryl Hydrocarbon Receptor-Interacting Protein-like 1 with the Farnesyl Moiety. The Journal of Biological Chemistry. 04 June 2013; Vol. 288, No. 29; pages 21320-21328; figure 4; page 21324, column 2, paragraph 1; page 21326, column 2, paragraph 2; DOI: 10.1074/jbc.M113.476242	71, 100/97-99, 143
Y	· (SUN, F et al.) Design and Structure-Based Study of New Potential FKBP12 Inhibitors. Biophysical Journal. November, 2003; Vol. 85, No. 5; pages 3194-3201; page 3195, column 2, paragraph 3; DOI: 10.1016/S0006-3495(03)74737-7	256/253-254



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance
 “E” earlier application or patent but published on or after the international filing date
 “L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 “O” document referring to an oral disclosure, use, exhibition or other means
 “P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search

25 July 2018 (25.07.2018)

Date of mailing of the international search report

07 AUG 2018

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
 P.O. Box 1450, Alexandria, Virginia 22313-1450
 Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300
 PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/26014

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2007/0218502 A1 (HAHN, KM et al.) 20 September, 2007; abstract; paragraphs [0008], [0010], [0035]	230-231, 232/230-231, 233/232/230-231, 234/233/232/230-231, 235/232/230-231
P, X	WO 2017/059207 A1 (WARP DRIVE BIO, INC.) 06 April, 2017; whole document	1-2, 3/1-2, 4/3/1-2, 70-71, 97-99, 100/97-99, 135-138, 139/136-138, 140/136-138, 142-143, 148-151, 152/149-151, 153/149-151, 224-227, 230-231, 232/230-231, 233/232/230-231, 234/233/232/230-231, 235/232/230-231, 248-250, 252/249-250, 253-254, 256/253-254

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/26014

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: -**-. Please See Supplemental Page-**-. because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

-**- Please See Supplemental Page-**-

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Group 1+ Claims 1-4, 70, 71 (in-part), 97-99, 100 (in-part), 135-140, 142, 143 (in-part), 148-153, 224-227, 230-235, 248-250, 252 (in-part), 253, 254, and 256 (in-part)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/US18/26014

-***-Continued from Box No. II Observations where certain claims were found unsearchable-***-

Claims Nos: 5-69, 76-79, 82-86, 90-95, 102-134, 141, 154-155, 160-183, 190, 194, 201-202, 207-214, 218-223, 228-229, 236-243, 257-277 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

-***- Continued from Box No. III Observations where unity of invention is lacking -***-

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Groups I+, Claims 1-4, 70-75, 80, 81, 87-89, 96-101, 135-140, 142-153, 156-159, 184-189, 191-193, 195-200, 203-206, 215-217, 224-227, 230-235, 244-256, a presenter protein binding moiety (protein binding moiety), and AIP (presenter protein) are directed toward a compound comprising a protein binding moiety and a cross-linking group; conjugates comprising the compound; complexes comprising the compound, methods of producing the conjugates and complexes; methods associated with the proteins, complexes, conjugates, and the interfaces thereof, or crystal structures thereof.

The compounds, conjugates, complexes and methods will be searched to the extent they encompass a compound encompassing a presenter protein binding moiety (first exemplary protein binding moiety), that binds to a presenter protein encompassing AIP (first exemplary presenter protein). Applicant is invited to elect additional compound(s), with, where applicable, fully specified structure(s) for each compound (i.e. no optional or variable atoms, groups, or substituents), and/or protein binding moiety(ies) and/or specific protein(s) bound by the protein binding moiety(ies) to be searched. Additional compound(s) and/or protein binding moiety(ies) and/or protein(s) bound by the protein binding moiety(ies) will be searched upon the payment of additional fees. It is believed that claims 1-4, 70, 71 (in-part), 97-99, 100 (in-part), 135-140, 142, 143 (in-part), 148-153, 224-227, 230-235, 248-250, 252 (in-part), 253, 254, and 256 (in-part) encompass this first named invention and thus these claims will be searched without fee to the extent that they encompass a presenter protein binding moiety (protein binding moiety), and AIP (presenter protein). Applicants must specify the claims that encompass any additionally elected compound(s), and/or protein binding moiety(ies), and/or specific protein(s) bound by protein binding moiety(ies). Applicants must further indicate, if applicable, the claims which encompass the first named invention, if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched/examined. An exemplary election would be AIP1 (presenter protein). (It should be noted that applicants must identify the protein(s) bound by any elected compound(s) with specified structure(s), wherein the protein bound is associated with the elected compound, and search of said bound protein would not require payment of a further fee (as said protein does not constitute a separately elected species). It should further be noted that, should applicants elect, in general, for the search of an additional protein binding moiety, such as a target protein binding moiety, applicant would be allowed to elect one target protein bound by said moiety as the first embodiment of said election).

No technical features are shared between the compounds and/or crosslinkers and/or proteins of Groups I+ and, accordingly, these groups lack unity a priori.

Additionally, even if Groups I+ were considered to share the technical features including: a compound comprising a protein binding moiety and a cross-linking group; a conjugate comprising protein binding moiety conjugated to a protein; a method of producing a conjugate comprising a protein binding moiety conjugated to a protein, said method comprising reacting (a) a compound comprising a first protein binding moiety and a cross-linking group with (b) a second protein under conditions that permit production of said conjugate; a complex comprising (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein; a method of producing a complex comprising (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein, said method comprising combining a conjugate comprising a first protein binding moiety conjugated to a second protein capable of forming a complex with a first protein, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; and (c) determining if said conjugate and said first protein form a complex, wherein if said conjugate and said first protein form a complex, said conjugate is identified as a conjugate capable of forming a complex with a first protein, thereby identifying a conjugate comprising a first protein binding moiety conjugated to a second protein capable of forming a complex with a first protein; a method of identifying a second protein which binds to a first protein, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; and (c) determining whether said second protein binds to said first protein in said complex, wherein if said second protein binds to said first protein, said second protein is identified as binding to the first protein, thereby identifying a second protein which binds to a first protein; a method of identifying a second protein capable of reacting with a compound in the presence of a first protein, wherein the compound comprises a first protein binding moiety and a cross-linking moiety, said method comprising: (a) providing (i) a compound comprising a first protein binding moiety and a cross-linking moiety; (ii) a second protein; and (iii) a first protein; (b) combining said compound, said second protein, and said first protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the first protein; and (c) determining if said second protein and said compound react during formation of said complex to form a conjugate, said second protein is identified as a second protein capable of reacting with the compound in the presence of a first protein, thereby identifying a second protein capable of reacting with a compound comprising a first protein binding moiety and a cross-linking moiety in the presence of a first protein; a method of identifying a location on a second protein to form a conjugate with a first protein binding moiety, which conjugate is capable of forming a complex with a first protein, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein at a location and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; (c) determining if said conjugate and said first protein form a complex;

-***-Continued Within the Next Supplemental Box-***-

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/26014

-***-Continued from Previous Supplemental Page:

and (d) repeating steps (a) to (c) with said first protein binding moiety conjugated at different locations on said second protein until a conjugate and said first protein form a complex, wherein a location on a second protein to form a conjugate with a first protein binding moiety, which conjugate is capable of forming a complex with a first protein is identified if said conjugate and said first protein form a complex, thereby identifying a location on a second protein to form a conjugate capable of forming a complex with a first protein; a method of identifying a compound capable of selective and covalent binding to a second protein in the presence of a first protein, said method comprising: (a) providing a first sample comprising (i) a compound comprising a first protein binding moiety and a cross-linking group; (ii) a second protein; and (iii) a first protein; and a second sample comprising (i) the same compound comprising a first protein binding moiety and a cross-linking group as in the first sample and (ii) the same second protein as in the first sample; and (b) determining the extent to which said compound and said second protein react in said first sample as compared to said second sample, wherein a compound is identified as selectively covalently binding to a second protein in the presence of a first protein if said compound and said second protein reacts in said first sample more than in said second sample; a method of determining the structure of an interface in a complex comprising a first protein and a second protein, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; and (c) determining the crystal structure of said complex, wherein the structure of the interface comprises at least the portion of the crystal structure between said first protein and said second protein, thereby determining the structure of an interface in a complex comprising a first protein and a second protein; a method of obtaining X-ray crystal coordinates for a complex, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; and (c) determining the crystal structure of said complex, thereby obtaining X-ray crystal coordinates for said complex; and a method of determining the residues on a second protein that participate in binding with a first protein, said method comprising: (a) providing X-ray crystal coordinates of a complex; (b) identifying the residues of said second protein which comprise an atom within 4A of an atom on said first protein; thereby determining the residues on a second protein that participate in binding with a first protein; these shared technical features are previously disclosed by WO 2016/112279 A1 to Warp Drive Bio, Inc., (hereinafter 'Warp') as evidenced by 'Distinction between the weak hydrogen bond and the van der Waals interaction' by Steiner et al. (hereinafter 'Steiner'), and in view of US 2007/0218502 A1 to Hahn et al. (hereinafter 'Hahn').

Warp discloses a compound comprising a protein binding moiety and a cross-linking group (a compound comprising a protein binding moiety and a cross-linking group; page 1, lines 25-39); a conjugate comprising protein binding moiety conjugated to a protein (a compound comprising a linker that participates in binding to a protein covalently (a conjugate comprising protein binding moiety conjugated to a protein) page 60, line 34 - page 61, line 10); a method of producing a conjugate (use of linkers in (a method of) preparation of conjugates; page 63, lines 9-10) comprising a protein binding moiety conjugated to a protein (comprising a protein binding moiety conjugated to a protein; page 60, line 34 - page 61, line 10), said method comprising reacting (a) a compound comprising a first protein binding moiety and a cross-linking group with (b) a second protein under conditions that permit production of said conjugate (the method comprising bond formation with one or more functional groups located on a linker and a second protein (said method comprising reacting (a) a compound comprising a first protein binding moiety and a cross-linking group with (b) a second protein under conditions that permit production of said conjugate); page 60, line 34 - page 61, line 7); a complex (a complex; page 64, lines 11-13) comprising (i) a conjugate comprising a first protein binding moiety conjugated to a second protein (comprising (i) a conjugate comprising a first protein binding moiety conjugated to a second protein; page 60, line 34 - page 61, line 10, page 63, lines 9-10; page 64, lines 11-13) and (ii) a first protein (and a first protein; page 64, lines 11-13); a method of producing a complex (a method of modulating a protein by producing a complex; page 18, lines 18-22, page 60, line 34 - page 61, line 7) comprising (i) a conjugate comprising a first protein binding moiety conjugated to a second protein (comprising (i) a conjugate comprising a first protein binding moiety conjugated to a second protein; page 18, lines 18-22, page 60, line 34 - page 61, line 7) and (ii) a first protein (and a first protein; page 18, lines 18-22), said method comprising combining a conjugate comprising a first protein binding moiety conjugated to a second protein and a first protein (said method comprising combining a conjugate comprising a first protein binding moiety conjugated to a second protein and a first protein; page 18, lines 18-22, page 60, line 34 - page 61, line 7) under conditions that permit production of said complex (under conditions effective to modulate the protein (that permit production of said complex); page 18, lines 18-22); a method of identifying a conjugate comprising a first protein binding moiety conjugated to a second protein (a method of determining binding of conjugate compounds (identifying a conjugate) comprising a first protein binding moiety conjugated to a second protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 86, lines 21-22) capable of forming a complex with a first protein (capable of forming a complex with cyclosporine A or FKBP12 (a first protein); page 60, line 34 - page 61, line 7; page 63; page 86, lines 21-22; Table 5; page 88, lines 3-4, Table 6), said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein (said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 86, lines 21-22) and (ii) a first protein (cyclosporine A or FKBP12 (a first protein); page 86, lines 21-22; Table 5; page 88, lines 3-4, Table 6); (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein (combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 86, lines 34-41; page 88, line 14 - page 89, line 2); and (c) determining if said conjugate and said first protein form a complex (determining if said conjugate and said first protein form a complex; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 86, lines 21-22; Table 5; page 88, lines 3-4, Table 6), wherein if said conjugate and said first protein form a complex, said conjugate is identified as a conjugate capable of forming a complex with a first protein, thereby identifying a conjugate comprising a first protein binding moiety conjugated to a second protein capable of forming a complex with a first protein (wherein if said conjugate and said first protein form a complex, said conjugate is identified as a conjugate capable of forming a complex with a first protein, thereby identifying a conjugate comprising a first protein binding moiety conjugated to a second protein capable of forming a complex with a first protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 86, lines 21-22; Table 5; page 88, lines 3-4, Table 6);

-***-Continued Within the Next Supplemental Box-***-

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/26014

-***-Continued from Previous Supplemental Page:

wherein if said conjugate and said first protein form a complex, said conjugate is identified as a conjugate capable of forming a complex with a first protein, thereby identifying a conjugate comprising a first protein binding moiety conjugated to a second protein capable of forming a complex with a first protein (wherein if said conjugate and said first protein form a complex, said conjugate is identified as a conjugate capable of forming a complex with a first protein, thereby identifying a conjugate comprising a first protein binding moiety conjugated to a second protein capable of forming a complex with a first protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 86, lines 21-22; Table 5; page 88, lines 3-4, Table 6); a method of determining the structure of an interface (determining the residues which interact in a ternary complex comprising a compound and two proteins (a method of determining the structure of an interface in a complex comprising a first protein and a second protein); page 97, lines 29-30; page 98, lines 7-15), said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein (said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 97, line 34 - page 98, line 15) and (ii) a first protein (and a first protein; page 97, line 34 - page 98, line 15); (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein (combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 97, line 34 - page 98, line 15); and (c) determining the crystal structure of said complex (determining the crystal structure of said complex; page 97, line 29 - page 98, line 15), wherein the structure of the interface comprises at least the portion of the crystal structure between said first protein and said second protein (wherein the structure of the interface comprises residues that interact between the first and second proteins, based on the crystal structure (at least the portion of the crystal structure between said first protein and said second protein); page 97, line 29 - page 98, line 15), thereby determining the structure of an interface in a complex comprising a first protein and a second protein (thereby determining the residues that interact between the two proteins in the crystal structure (structure of an interface in a complex comprising a first protein and a second protein); page 97, line 29 - page 98, line 15); a method of obtaining X-ray crystal coordinates for a complex (a method of obtaining X-ray crystal coordinates for a complex; page 97, line 29 - page 98, line 34), said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein (said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 97, line 34 - page 98, line 15) and (ii) a first protein (and a first protein; page 97, line 34 - page 98, line 15); (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; (Combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; page 60, line 34 - page 61, line 7; page 63, lines 9-10; page 97, line 34 - page 98, line 15); and (c) determining the crystal structure of said complex (determining the crystal structure of said complex; page 97, line 29 - page 98, line 34), thereby obtaining X-ray crystal coordinates for said complex (thereby obtaining X-ray crystal coordinates for said complex; page 97, line 34 - page 98, line 15); and a method of determining the residues on a second protein that participate in binding with a first protein (a method of determining the residues on a second protein that participate in binding with a first protein; page 97, line 34 - page 98, line 15), said method comprising: (a) providing X-ray crystal coordinates of a complex (said method comprising: (a) providing X-ray crystal coordinates of a complex; page 97, line 34 - page 98, line 34); (b) identifying the residues of said second protein which comprise an atom within 4A of an atom on said first protein (identifying the residues of said second protein which comprise an atom with van der Waals contacts with atoms of the first protein (within 4A of an atom on said first protein); page 97, line 34 - page 98, line 34; wherein van der Waals contacts are at less than 4A, as evidenced by Steiner [Table 1]); thereby determining the residues on a second protein that participate in binding with a first protein (thereby determining the residues on a second protein that participate in binding with a first protein; page 97, line 34 - page 98, line 34); Warp further discloses wherein a ternary complex is formed by protein-protein interactions (wherein a ternary complex is formed by protein-protein interactions; page 19, lines 6-19; page 72, lines 19-28); and wherein the ternary complex includes polypeptides which do not bind in the absence of the compound, or wherein the first and second proteins bind with at least 5-fold greater affinity in the presence of the compound than the absence of the compound (wherein the ternary complex includes polypeptides which do not bind in the absence of the compound, or wherein the first and second proteins bind with at least 5-fold greater affinity in the presence of the compound than the absence of the compound; page 1, lines 25-34).

Warp does not disclose: a method of identifying a second protein which binds to a first protein, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; and (c) determining whether said second protein binds to said first protein in said complex, wherein if said second protein binds to said first protein, said second protein is identified as binding to the first protein, thereby identifying a second protein which binds to a first protein; a method of identifying a second protein capable of reacting with a compound in the presence of a first protein, wherein the compound comprises a first protein binding moiety and a cross-linking moiety, said method comprising: (a) providing (i) a compound comprising a first protein binding moiety and a cross-linking moiety; (ii) a second protein; and (iii) a first protein; (b) combining said compound, said second protein, and said first protein under conditions suitable for to permit complex formation if said compound is capable of forming a complex with the first protein; and (c) determining if said second protein and said compound react during formation of said complex to form a conjugate, wherein if said second protein and said compound react during formation of said complex to form a conjugate, said second protein is identified as a second protein capable of reacting with the compound in the presence of a first protein, thereby identifying a second protein capable of reacting with a compound comprising a first protein binding moiety and a cross-linking moiety in the presence of a first protein; a method of identifying a location on a second protein to form a conjugate with a first protein binding moiety, which conjugate is capable of forming a complex with a first protein, said method comprising: (a) providing (i) a conjugate comprising a first protein binding moiety conjugated to a second protein at a location and (ii) a first protein; (b) combining said conjugate and said first protein under conditions suitable for to permit complex formation if said conjugate is capable of forming a complex with the first protein; (c) determining if said conjugate and said first protein form a complex; and (d) repeating steps (a) to (c) with said first protein binding moiety conjugated at different locations on said second protein until a conjugate and said first protein form a complex, wherein a location on a second protein to form a conjugate with a first protein binding moiety, which conjugate is capable of forming a complex with a first protein is identified if said conjugate and said first protein form a complex, thereby identifying a location on a second protein to form a conjugate capable of forming a complex with a first protein; and a method of identifying a compound capable of selective and covalent binding to a second protein in the presence of a first protein, said method comprising: (a) providing a first sample comprising (i) a compound comprising a first protein binding moiety and a cross-linking group; (ii) a second protein; and (iii) a first protein; and a second sample comprising (i) the same compound comprising a first protein binding moiety and a cross-linking group

-***-Continued Within the Next Supplemental Box-***-

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/26014

-***-Continued from Box -***-

(ii) a second protein; and (iii) a first protein; and a second sample comprising (i) the same compound comprising a first protein binding moiety and a cross-linking group as in the first sample and (ii) the same second protein as in the first sample; and (b) determining the extent to which said compound and said second protein react in said first sample as compared to said second sample, wherein a compound is identified as selectively covalently binding to a second protein in the presence of a first protein if said compound and said second protein reacts in said first sample more than in said second sample.

Hahn discloses methods for identifying optimum conjugate attachment sites to peptides (methods for identifying optimum conjugate attachment sites to peptides; abstract); including introduction of a conjugation site into the peptide or protein at different locations (including introduction of a conjugation site into the peptide or protein at different locations; paragraphs [0008], [0010]) and assessing the degree to which the different conjugates are active (assessing the degree to which the different conjugates are active; paragraph [0035]).

It would have been obvious to a person of ordinary skill in the art at the time of the invention was made to have modified the disclosure of Warp to have provided a method of identifying a second protein which binds to a first protein by providing a conjugate, as disclosed by Warp, comprising a protein, and a second protein, and determining the crystal structure of the complex to identify the interface between the proteins to a second protein which binds to a first protein. Since Warp further discloses the crosslinking of a compound with a first protein, it further would have been obvious to a person of ordinary skill in the art at the time of the invention was made to have provided the first protein and the compound to which it crosslinks with a second protein, wherein the second protein binds to both the compound and the first protein, as disclosed by Warp, in order to better stabilize the first protein in complex with the compound to more effectively crosslink the compound to the first protein at a desired location. It also would have been obvious to a person of ordinary skill in the art at the time of the invention was made to have modified the disclosure of Warp to have provided a method of identifying a compound capable of selective and covalent binding (i.e. forming a conjugate) to a second protein in the presence of a first protein by comparing the conjugation of the compound to the first protein in the presence of absence of the second protein, wherein, due to the increased affinity in the tripartite complex, the compound reacts and forms a conjugate to a greater degree in the presence of the second protein than in its absence, as a means of specifically identifying compounds that stabilize the tripartite complex with the second protein. Finally, It would have been obvious to a person of ordinary skill in the art at the time of the invention was made to have modified the disclosure of Warp to have used a locatable conjugation site in a protein, as disclosed by Hahn, and moving the site to a plurality of locations on the protein surface, as disclosed by Hahn, in order to determine the best site for the formation of a conjugate capable of producing a tripartite or ternary complex, as disclosed by Warp, with high efficiency.

Since none of the special technical features of the Groups I+ inventions is found in more than one of the inventions, and since all of the shared technical features are previously disclosed by a combination of the Warp and Hahn references, unity of invention is lacking.