

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2005297772 B2**

- (54) Title
Anti-glypican 3 antibody having modified sugar chain
- (51) International Patent Classification(s)
C07K 16/18 (2006.01) **C12N 15/09** (2006.01)
A61K 39/395 (2006.01) **C12P 21/08** (2006.01)
A61P 35/00 (2006.01)
- (21) Application No: **2005297772** (22) Date of Filing: **2005.10.26**
- (87) WIPO No: **WO06/046751**
- (30) Priority Data
- (31) Number (32) Date (33) Country
2004-311356 **2004.10.26** **JP**
- (43) Publication Date: **2006.05.04**
(44) Accepted Journal Date: **2011.06.23**
- (71) Applicant(s)
Chugai Seiyaku Kabushiki Kaisha
- (72) Inventor(s)
Ishiguro, Takahiro;Iijima, Shigeyuki;Nakano, Kiyotaka;Sugimoto, Masamichi;Tanaka, Megumi;Sugo, Izumi
- (74) Agent / Attorney
FB Rice, Level 23 44 Market Street, Sydney, NSW, 2000
- (56) Related Art
WO 2004/022739 A1 (Chugai Pharmaceutical Co., Ltd) 18 March 2004
EP 1331266 A1 (Kyowa Hakko Kogyo KK) 30 July 2003
WO 2006/006693 A1 (Chugai Seiyaku Kabushiki Kaisha) 19 January 2006
EP 1411118 A1 (Chugai Seiyaku KK) 21 April 2004
WO 2003/085119 (Kyowa Hakko Kogyo KK) 16 October 2003

(12) 特許協力条約に基づいて公開された国際出願

(19) 世界知的所有権機関
国際事務局



(43) 国際公開日
2006 年 5 月 4 日 (04.05.2006)

PCT

(10) 国際公開番号
WO 2006/046751 A1

(51) 国際特許分類:

C07K 16/18 (2006.01) C12N 15/09 (2006.01)
A61K 39/395 (2006.01) C12P 21/08 (2006.01)
A61P 35/00 (2006.01)

KAISHA) [JP/JP]; 〒1158543 東京都北区浮間五丁目
5 番 1 号 Tokyo (JP).

(72) 発明者: および

(75) 発明者/出願人 (米国についてのみ): 中野 清孝
(NAKANO, Kiyotaka) [JP/JP]; 〒3004101 茨城県新治
郡新治村永井 1 5 3 - 2 中外製薬株式会社内 Ibaraki
(JP). 周郷 泉 (SUGO, Izumi) [JP/JP]; 〒4128513 静岡県
御殿場市駒門 1 丁目 1 3 5 番地 中外製薬株式会
社内 Shizuoka (JP). 杉本 正道 (SUGIMOTO, Masamichi)
[JP/JP]; 〒2478530 神奈川県鎌倉市梶原 2 0 0 中外製
薬株式会社内 Kanagawa (JP). 石黒 敬弘 (ISHIGURO,
Takahiro) [JP/JP]; 〒2478530 神奈川県鎌倉市梶原
2 0 0 中外製薬株式会社内 Kanagawa (JP). 田中 めぐ
み (TANAKA, Megumi) [JP/JP]; 〒4128513 静岡県御
殿場市駒門 1 丁目 1 3 5 番地 中外製薬株式会
社内

(21) 国際出願番号: PCT/JP2005/020057

(22) 国際出願日: 2005 年 10 月 26 日 (26.10.2005)

(25) 国際出願の言語: 日本語

(26) 国際公開の言語: 日本語

(30) 優先権データ:

特願 2004-311356


2004 年 10 月 26 日 (26.10.2004) JP

(71) 出願人 (米国を除く全ての指定国について): 中
外製薬株式会社 (CHUGAI SEIYAKU KABUSHIKI

[続葉有]

(54) Title: ANTI-GLYPICAN 3 ANTIBODY HAVING MODIFIED SUGAR CHAIN

(54) 発明の名称: 糖鎖改変抗グリピカン 3 抗体



AA...CYTOTOXIC ACTIVITY (%)

BB...ANTIBODY CONCENTRATION (μg/mL)

(57) Abstract: An anti-glypican 3 antibody having a modified sugar chain, more specifically, an anti-glypican 3 antibody lacking fucose. This anti-glypican 3 antibody can be produced by a method for producing an antibody with the sugar chain modification as described above wherein a nucleic acid encoding an anti-glypican 3 antibody is transferred into host cells with lowered ability to add fucose (for example, YB2/0 cells) or fucose transporter-deficient cells and then the host cells are cultured. Because of having a high cytotoxic activity, this anti-glypican 3 antibody having a modified sugar chain is useful as a cell growth inhibitor such as an anticancer agent.

(57) 要約: 糖鎖が改変された抗グリピカン 3 抗体、より具体的にはフコースが欠損した抗グリピカン 3 抗体が開示される。本発明の抗グリピカン 3 抗体は、上記の糖鎖が修飾された抗体の製造方法において、YB2/0 細胞などのフコース付加能が低下した宿主細胞やフコーストランスポーター欠損細胞に抗グリピカン 3 抗体をコードする核酸を導入し、該宿主細胞を培養することにより製造することができる。本発明の糖鎖改変抗グリピカン 3 抗体は、高い細胞障害

[続葉有]

WO 2006/046751 A1



Shizuoka (JP). 飯島 成幸 (IIJIMA, Shigeyuki) [JP/JP];
〒4128513 静岡県御殿場市駒門 1 丁目 1 3 5 番地 中
外製薬株式会社内 Shizuoka (JP).

(74) 代理人: 大野 聖二, 外(OHNO, Seiji et al.); 〒1006036
東京都千代田区霞が関 3 丁目 2 番 5 号 霞が関ビル
3 6 階 大野総合法律事務所 Tokyo (JP).

(81) 指定国 (表示のない限り、全ての種類の国内保護が
可能): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR,
BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX,
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT,
TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) 指定国 (表示のない限り、全ての種類の広域保護が可
能): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD,
SL, SZ, TZ, UG, ZM, ZW), ユーラシア (AM, AZ, BY,
KG, KZ, MD, RU, TJ, TM), ヨーロッパ (AT, BE, BG,
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR),
OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
MR, NE, SN, TD, TG).

添付公開書類:
— 国際調査報告書

2 文字コード及び他の略語については、定期発行される
各 PCT ガゼットの巻頭に掲載されている「コードと略語
のガイダンスノート」を参照。

DESCRIPTION

Anti-Glypican 3 Antibody Having Modified Sugar Chain

TECHNICAL FIELD

The present invention relates to an antibody to glypican 3 antigen (i.e., an anti-glypican 3 antibody) in which cytotoxic activity, especially antibody-dependent cellular cytotoxicity (ADCC) is enhanced, as well as a process for producing the antibody.

BACKGROUND ART

Glypican 3 (GPC3) is a member of the family of heparan sulfate proteoglycans present on the surface of cells. It has been suggested that GPC3 may be involved in cell division upon development and in cancer cell growth, but its function is still not well understood.

It has been discovered that a type of antibody that binds to GPC3 inhibits cell growth due to ADCC (antibody-dependent cellular cytotoxicity) activity and CDC (complement-dependent cytotoxicity) activity (WO 2003/00883). Furthermore, because GPC3 is cleaved in the body and secreted into the blood as a soluble form of GPC3, it has been suggested that cancer can be diagnosed using an antibody that can detect the soluble form of GPC3 (WO 2004/022739, WO 2003/100429, WO 2004/018667).

When developing an anticancer agent that utilizes

antibody cytotoxic activity, preferably the antibody used will have a high level of ADCC activity. Therefore, an anti-GPC3 antibody with a high level of cytotoxic activity has been demanded.

Modification of antibody sugar chains are known to enhance its ADCC activity.

- 5 For example, WO 99/54342 discloses that ADCC activity is enhanced by modifying antibody glycosylation. In addition, WO 00/61739 discloses that ADCC activity is regulated by controlling the presence or absence of fucose in antibody sugar chains. WO 02/31140 discloses producing an antibody having sugar chains that do not contain a-1,6 core fucose by producing that antibody in YB2/0 cells. WO 02/79255 discloses
10 an antibody with sugar chains having bisecting GlcNAc. However, an anti-GPC3 antibody with enhanced ADCC activity due to sugar chain modification has not been disclosed so far.

- Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a
15 context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

DISCLOSURE OF THE INVENTION

- 20 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

- The present invention provides, at least in its preferred form(s), an anti-GPC3
25 antibody composition with enhanced ADCC activity caused by the alteration of the sugar chain component thereof, as well as a process for producing such an antibody.

- After various investigations, the inventors have discovered that an antibody targeting GPC3 with sugar chains lacking a-1,6 core fucose have a high level of cytotoxic activity. Thus, the present invention provides, at least in its preferred
30 form(s), an anti-GPC3 antibody composition wherein the sugar chain component of the antibody has been altered, and more specifically, an antibody composition with a greater fraction of fucose deficient anti-GPC3 antibodies. The sugar chain-modified anti-GPC3 antibody composition of the present invention has a high level of cytotoxic activity, and therefore is useful as a cell growth inhibitor such as an anticancer agent.

The present invention also provides, at least in its preferred form(s), a process for producing an anti-GPC3 antibody composition wherein the sugar chain of the antibody is modified, comprising the steps of: introducing a nucleic acid encoding the anti-GPC3 antibody into a host cell with reduced fucose addition capability such as
5 YB2/0 cells, and culturing the host cell to obtain the antibody. Preferably, the cell with reduced capability of adding fucose to sugar chains is a cell lacking a fucose transporter.

Thus, according to an aspect of the present invention there is provided an anti-glypican 3 antibody composition comprising an antibody having a heavy chain variable
10 region comprising CDR1 having the amino acid sequence set forth in SEQ ID NO: 25, CDR2 having the amino acid sequence set forth in SEQ ID NO: 26, and CDR3 having the amino acid sequence set forth in SEQ ID NO: 27, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting N-
15 acetylglucosamid (GlcNAc).

In a further aspect of the present invention, there is provided an anti-glypican 3 antibody composition comprising an antibody having a light chain variable region comprising CDR1 having the amino acid sequence set forth in SEQ ID NO: 45, CDR2 having the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 having the
20 amino acid sequence set forth in SEQ ID NO: 60, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting N-acetylglucosamid (GlcNAc).

In a further aspect of the present invention, there is provided an anti-glypican 3
25 antibody composition comprising an antibody having a light chain variable region comprising CDR1 having the amino acid sequence set forth in SEQ ID NO: 45, CDR2 having the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 having the amino acid sequence set forth in SEQ ID NO: 60, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient
30 antibodies or an increased ratio of antibodies having an attached bisecting N-acetylglucosamid (GlcNAc).

In a further aspect of the present invention, there is provided an anti-glypican 3 antibody composition wherein the antibody as defined herein further has a light chain variable region comprising CDR1 having the amino acid sequence set forth in SEQ ID
35 NO: 45, CDR2 having the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 having the amino acid sequence set forth in SEQ ID NO: 60.

In a further aspect of the present invention, there is provided an anti-glypican 3 antibody composition comprising an antibody having a heavy chain variable region of any one of (1)-(7):

- (1) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 81;
- (2) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 82;
- (3) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 83;
- 10 (4) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84;
- (5) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85;
- (6) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86; or
- 15 (7) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting GlcNAc.

- 20 In a further aspect of the present invention, there is provided an anti-glypican 3 antibody composition comprising an antibody having a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting
- 25 GlcNAc.

In a further aspect of the present invention, there is provided a process for producing an anti-glypican 3 antibody using a cell, wherein a gene encoding the anti-glypican 3 antibody as defined herein has been introduced into said cell having reduced capability of adding fucose to sugar chains.

- 30 In a further aspect of the present invention, there is provided a process for producing an anti-glypican 3 antibody, comprising the steps of:
- (a) introducing a gene encoding the anti-glypican 3 antibody as defined herein into a cell having reduced capability of adding fucose to sugar chains; and
 - (b) culturing the cell.

- 35 In a further aspect of the present invention, there is provided an anticancer drug having as its active ingredient the antibody composition as defined herein.

In a further aspect of the present invention, there is provided use of a composition as defined herein as a medicament.

In a further aspect of the present invention, there is provided use of a composition as defined herein as an anticancer agent.

- 5 In a further aspect of the present invention, there is provided a method for treating cancer comprising administering a composition as defined herein.

In a further aspect of the present invention, there is provided use of a composition as defined herein for the manufacture of a medicament for the treatment of cancer

10 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 shows the basic structure of N-glycoside linked sugar chains;

Figure 2 shows ADCC activity of a chimeric antibody when HepG2 cells are targeted using human peripheral blood monocytes (PBMC);

Figure 3 shows ADCC activity of a chimeric antibody

when HuH-7 cells are targeted using human PBMC;

Figure 4 shows ADCC activity of antibodies when HuH-7 cells are targeted using human PBMC;

Figure 5 shows normal phase HPLC chromatograms of sugar chains modified by agalactosyl 2-AB prepared from antibodies (a, b, c) produced by FT-KO cells and by CHO cells;

Figure 6 shows the predicted structures for the G(0) and G(0)-Fuc peaks shown in Figure 5; and

Figure 7 shows a differential scanning calorimetry (DSC) measurement plot for the antibodies produced by FT-KO cells (a) and by CHO cells (b).

DETAILED DESCRIPTION OF THE INVENTION

The present invention features an anti-GPC3 antibody composition wherein the sugar chain component of the antibody has been modified. It is known that the structure of the sugar chain linked to an antibody has a significant effect on the expression of antibody cytotoxic activity. Sugar chains that is linked to an antibody include N-glycoside-linked sugar chains, which are attached to a nitrogen atom on the side chain of an asparagine residue on the antibody molecule, and O-glycoside-linked sugar chains, which are attached to a hydroxyl group on the side chain of a serine or threonine residue on the antibody molecule. The present invention

is focused on the presence or absence of fucose in an N-glycoside-linked sugar chain.

Figure 1 shows the basic structure of N-glycoside-linked sugar chains attached to an antibody. As shown in the IgG basic sugar chains (1) and (3) of Figure 1, the N-glycoside linked sugar chains have a basic structure (core) wherein one mannose (Man) and two N-acetylglucosamine (GlcNAc) moieties are linked by β -1,4 linkages [-Man β 1-4GlcNAc β 1-4GlcNAc-]. The "GlcNAc" on the right side of the structure is called the reducing end and the "Man" on the left side of the structure is called the non-reducing end. When a fucose is linked to the reducing end, it usually takes the form of an α -linkage between the 6-position of the N-acetylglucosamine at the reducing end and the 1-position of the fucose. On the other hand, in the sugar chain shown in IgG basic sugar chain (2) of Figure 1, in addition to the aforementioned two sugar chains, one N-acetylglucosamine (GlcNAc) moiety is linked to the non-reducing end of the basic structure (core) via a β 1,4-linkage. This type of N-acetylglucosamine (GlcNAc) is called a "bisecting N-acetylglucosamine." A sugar chain having a bisecting N-acetylglucosamine can be an O-glycoside-linked sugar chain or N-glycoside-linked sugar chain, and it is formed by transfer of N-acetylglucosamine to the sugar chain by N-acetylglucosamine transferase III (GnTIII). The gene

encoding this enzyme has already been cloned, and both the amino acid sequence and the nucleotide sequence of the DNA encoding the enzyme have already been reported (NCBI database (ACCESSION D13789)).

In the present invention, the antibody composition with a modified or altered sugar chain component (sugar chain-modified antibody composition) refers to an antibody composition having a sugar chain component that differs from the antibody composition produced by a host cell serving as a reference standard.

In the present invention, one may determine whether the sugar chain component has been modified or not by using as a reference standard the antibody composition produced by a host cell serving as a reference standard. If an antibody composition has a sugar chain component different from the antibody composition from the reference standard, that antibody composition is considered as an antibody composition with a modified sugar chain component.

The host cell serving as a reference standard in the present invention is CHO DG44 cell. CHO DG44 cell can be obtained, for example, from the Invitrogen Corporation.

Examples of an antibody composition with a modified sugar chain component include, for example, an antibody composition with an increased ratio of fucose (e.g., α -1,6 core fucose)-deficient antibodies in the antibody

composition and an antibody composition with an increased ratio of antibodies having an attached bisecting N-acetylglucosamine (GlcNAc) in the antibody composition.

In a preferred embodiment of the present invention, the antibody composition has a higher ratio of fucose-deficient antibodies than the antibody composition used as a reference standard.

Because some antibodies have a plurality of N-glycoside sugar chains, the fucose-deficient antibody of the present invention encompasses not only antibodies wherein no fucose is attached, but also antibodies wherein the number of fucose moieties attached to the antibody is reduced (an antibody having at least one or more sugar chains wherein fucose is not present).

When manufacturing a sugar chain-modified antibody with host cells, it is often difficult to obtain a composition containing uniform antibodies wherein all antibodies have identical sugar chains. Therefore, if an antibody composition with a modified sugar chain component of the invention is an antibody composition with an increased ratio of fucose-deficient antibodies, for example, then the antibody composition with the modified sugar chain component of the present invention may contain both antibodies deficient in fucose and antibodies not deficient in fucose, but the overall ratio of antibodies deficient in fucose will be higher than in

the antibody composition produced by the host cells serving as a reference standard. The present invention is not particularly limited to a specific ratio of fucose-deficient antibodies in the antibody composition with a high ratio of fucose-deficient antibodies of the present invention, but preferably the ratio is not less than 20%, more preferably not less than 50%, and most preferably not less than 90%.

The present invention is not particularly limited to a specific ratio of bisecting N-acetylglucosamine-added antibodies in the antibody composition having a high ratio of bisecting N-acetylglucosamine-added antibodies of the present invention, but preferably the ratio is not less than 20%, more preferably not less than 50%, and most preferably not less than 90%.

The anti-GPC3 antibody composition with a modified sugar chain component of the present invention can be obtained by methods known to those skilled in the art.

For example, a fucose-deficient antibody can be produced by expressing the anti-GPC3 antibody in host cells either lacking capability or having lower capability to add a-1,6 core fucose.

The present invention is not particularly limited to the host cells lacking capability or having lower capability to add fucose, but host cells with no or reduced fucose transferase activity, host cells with

lower fucose concentration in Golgi bodies, and the like may be used in the present invention. More specifically, examples of host cells include rat myeloma YB2/3HL.P2.G11.16Ag.20 cells (abbreviated as YB2/0 cells) (preserved as ATCC CRL 1662), FTVIII knockout CHO cells (WO 02/31140), Lec13 cells (WO 03/035835) and fucose transporter deficient cells (WO 2005/017155).

As used herein, the term "fucose transporter deficient cell" refers to a cell in which the quantity of fucose transporter in the cell is less than in normal cells, or fucose transporter function is attenuated due to an abnormality in the fucose transporter structure. Examples of fucose transporter deficient cells may include, for example, those cells wherein the fucose transporter gene is knocked out (hereinafter called FT-KO cells), those wherein part of the fucose transporter gene is either lacking or mutated, those deficient in the fucose transporter gene expression system, and the like. The nucleotide sequence of the gene encoding the Chinese hamster fucose transporter and the amino acid sequence thereof are shown in SEQ ID NOS: 126 and 127, respectively.

Moreover, it is possible to obtain the fucose transporter deficient cell of the present invention using RNA interference (RNAi) by utilizing the nucleotide sequence represented by SEQ ID NO: 126. RNAi refers to

the following phenomenon: when double stranded RNA (dsRNA) is introduced into a cell, intracellular mRNA matching that RNA sequence is specifically degraded and cannot be expressed as a protein. Normally dsRNA is used with RNAi, but the present invention is not limited thereto and, for example, double stranded RNA formed by self-complementary single stranded RNA molecules can also be used. With respect to the regions forming the double stranded molecule, the molecule may be double stranded in all regions, or may be single stranded in some regions (for example, one or both ends). The present invention is not limited to a specific length of the oligo-RNA used in RNAi. The length of the oligo-RNA in the present invention can be, for example, 5 to 1000 bases (or 5 to 1000 bp in a double stranded molecule), preferably 10 to 100 bases (or 10 to 100 bp in a double stranded molecule), and most preferably 15 to 25 bases (15 to 25 bp in a double stranded molecule); however, a length of 19 to 23 bases (19 to 23 bp in a double stranded molecule) is especially preferred.

The aforementioned RNAi process utilizes the phenomenon wherein dsRNA consisting of both sense RNA and antisense RNA homologous to a specific gene will destroy the homologous part of the transcript (mRNA) of that gene. dsRNA corresponding to the entire sequence of the fucose transporter gene may be used, or shorter dsRNA (for

example, 21 to 23 bp) corresponding to part of the sequence (small interfering RNA; siRNA) may be used. The dsRNA can be directly transferred into the cell, or a vector producing dsRNA can be prepared and transferred into a host cell, and the dsRNA can then be produced within the cell. For example, all or part of the DNA encoding the fucose transporter gene can be inserted into a vector so that it forms an inverted repeat sequence, and that vector can then be transferred into a host cell. The RNAi procedure can be carried out in accordance with the descriptions in the following references: Fire A. et al., *Nature* (1998), 391, 806-811; Montgomery M. K. et al., *Proc. Natl. Acad. Sci. USA* (1998), 95, 15502-15507; Timmons L. et al., *Nature* (1998), 395, 854; Sánchez A. et al., *Proc. Natl. Acad. Sci. USA* (1999), 96, 5049-5054; Misquitta L. et al., *Proc. Natl. Acad. Sci. USA* (1999), 96, 1451-1456; Kennerdell J. R. et al., *Cell* (1998), 95, 1017-1026; Waterhouse P. M. et al., *Proc. Natl. Acad. Sci. USA* (1998), 95 13959-13964; and Wianny F. et al., *Nature Cell Biol.* (2000), 2, 70-75.

The fucose transporter deficient cells obtained by the RNAi procedure may be screened as indicated by the fucose transporter activity. Screening can also be carried out based on the transcription and expression of the fucose transporter gene indicated by Western blotting or Northern blotting.

An antibody with a bisecting N-acetylglucosamine (GlcNAc) added to the sugar chain can be produced by expressing the anti-GPC3 antibody in a host cell having the capability to form a bisecting N-acetylglucosamine (GlcNAc) structure on the sugar chain.

A method for producing an antibody with a bisecting N-acetylglucosamine-added sugar chain is already known (WO 02/79255). The host cell having the capability to form a bisecting N-acetylglucosamine (GlcNAc) structure on a sugar chain is not particularly limited in the present invention, but may include, for example, a host cell having an expression vector containing DNA encoding GnTIII. Therefore, an anti-GPC3 antibody having a bisecting N-acetylglucosamine-added sugar chain can be produced using a host cell containing both an expression vector with DNA encoding GnTIII and an expression vector encoding the anti-GPC3 antibody. The DNA encoding GnTIII and the gene encoding the anti-GPC3 antibody can both be present on the same vector or can be present on different vectors.

Another method for increasing the ratio of fucose-deficient antibodies or bisecting N-acetylglucosamine-added antibodies in the antibody composition is to increase the ratio of those antibodies in the composition by purifying the fucose-deficient antibodies or bisecting N-acetylglucosamine-added antibodies.

Sugar chain analysis can be carried out by any methods known to those skilled in the art. For example, a sugar chain can be released from an antibody by reacting the antibody with N-glycosidase F (Roche) and the like. Then the sugar chains can be desalted by solid phase extraction using a cellulose cartridge (Shimizu Y. et al., Carbohydrate Research 332(2001), 381-388), concentrated and dried, and fluorescent labeled with 2-aminopyridine (Kondo A. et al., Agricultural and Biological Chemistry 54:8(1990), 2169-2170). The reagent is removed from the pyridylamino-sugar chains (PA-sugar chains) by solid phase extraction with a cellulose cartridge, then the sugar chains are concentrated by centrifugation to obtain purified PA-sugar chains. The sugar chains may be assayed by reverse phase HPLC analysis using an octadecyl silane (ODS) column. The PA-sugar chains thus prepared may be analyzed by two dimensional mapping utilizing a combination of reverse phase HPLC analysis with an ODS column and normal phase HPLC analysis with an amine column.

The sugar chain-modified anti-GPC3 antibody of the present invention is not limited to any specific antibodies, provided it binds to GPC3. Preferably, a binding to GPC3 can be specific. Preferred anti-GPC3 antibodies of the present invention include those antibodies that have the complementarity determining

region (CDR) sequence shown in Table 1 below.

Table 1

Antibody	CDR	Amino acid sequence	SEQ ID NO:
M13B3(H)	CDR1	NYAMS	5
	CDR2	AINNNGDDTYYLDTVKD	6
	CDR3	QGGAY	7
M3B8(H)	CDR1	TYGMGVG	8
	CDR2	NIWWYDAKYNSDLKS	9
	CDR3	MGLAWFAY	10
M11F1(H)	CDR1	IYGMGVG	11
	CDR2	NIWWNDDKYNSALKS	12
	CDR3	IGYFYFDY	13
M5B9(H)	CDR1	GYWMH	14
	CDR2	AIYPGNSDTNYNQKFKG	15
	CDR3	SGDLTGGLAY	16
M6B1(H)	CDR1	SYAMS	17
	CDR2	AINSNGGTTYYPDTMKD	18
	CDR3	HNGGYENYGWFAY	19
M10D2(H)	CDR1	SYWMH	20
	CDR2	EIDPSDSYTYYNQKFRG	21
	CDR3	SNLGDGPHYRFPAPFY	22
L9G11(H)	CDR1	SYWMH	20
	CDR2	TIDPSDSETHYNLQFKD	23
	CDR3	GAFYSSYSYWAUFAY	24
GC33(H)	CDR1	DYEMH	25

	CDR2	ALDPKTGDTAYSQKFKG	26
	CDR3	FYSYTY	27
GC179(H)	CDR1	INAMN	28
	CDR2	RIRSESNNYATYYGDSVKD	29
	CDR3	EVTTSFAY	30
GC194(H)	CDR1	ASAMN	31
	CDR2	RIRSKSNNYAIYYADSVKD	32
	CDR3	DPGYGPNPWFAY	33
GC199(H)	CDR1	DYSMH	34
	CDR2	WINTETGEPTYADDFKG	35
	CDR3	LY	36
GC202(H)	CDR1	TYGMGVG	8
	CDR2	NIWWHDDKYNSALKS	37
	CDR3	IAPRYNKYEGFFAF	38
M13B3(L)	CDR1	KSSQSLLDSDGKTYLN	39
	CDR2	LVSCLDS	40
	CDR3	WQGTHFPLT	41
M3B8(L)	CDR1	KASQDINNYLS	42
	CDR2	RANRLVD	43
	CDR3	LQCDEFPPWT	44
M11F1(L)	CDR1	RSSQSLVHSNGNTYLH	45
	CDR2	KVSNRFS	46
	CDR3	SQSTHVPWT	47
M5B9(L)	CDR1	RSSKSLLSNGITYLY	48
	CDR2	QMSNLAS	49

	CDR3	AQNLELPYT	50
M6B1(L)	CDR1	KASQDINKNII	51
	CDR2	YTSTLQP	52
	CDR3	LQYDNLPR	53
M10D2(L)	CDR1	RASHSISNFLH	54
	CDR2	YASQSIG	55
	CDR3	QQSNIWSLT	56
L9G11(L)	CDR1	RASESVEYYGTSLMQ	57
	CDR2	GASNVES	58
	CDR3	QQSRKVPYT	59
GC33(L)	CDR1	RSSQSLVHSNGNTYLH	45
	CDR2	KVSNRFS	46
	CDR3	SQNTHPPT	60
GC179(L)	CDR1	KSSKSLHSHNGNTYLN	61
	CDR2	WMSNLAS	62
	CDR3	MQHIEYPFT	63
GC194(L)1	CDR1	RSSKSLHSDITYLY	64
	CDR2	QMSNLAS	49
	CDR3	AQNLELPPT	65
GC194(L)2	CDR1	SASSSVSYMY	66
	CDR2	DTSNLAS	67
	CDR3	QQWSSYPLT	68
GC199(L)	CDR1	KSSQSLHSDGKTFLN	69
	CDR2	LVSRLDS	70
	CDR3	CQGTHFPRT	71

GC202(L)	CDR1	RSSQSIVHSNGNTYLE	72
	CDR2	KVSNRFS	46
	CDR3	FQGSHVPWT	73

The antibodies with the CDR sequence listed in the above table have a high level of cytotoxic activity. The antibodies with the CDR sequence listed in the above table recognize epitopes of amino acids 524-563 on GPC3. Because antibodies that recognize epitopes of amino acids 524-563 have a high level of cytotoxic activity, they are preferred as the anti-GPC3 antibody of the present invention.

In one preferred embodiment of the present invention, the antibody composition having a modified sugar chain component of the present invention is characterized by exhibiting enhanced ADCC activity. In the present invention, whether the ADCC activity is enhanced or not may be determined by comparing the ADCC activity of the antibody composition of the present invention with that of the reference standard antibody composition. If the antibody composition of the present invention shows higher ADCC activity than the reference standard, the ADCC activity is said to be enhanced.

ADCC activity can be measured by a method known to those skilled in the art, for example, by mixing the anti-GPC3 antibody with effector cells and target cells,

and then determining the level of ADCC. More specifically, mouse spleen cells, human monocytes isolated from peripheral blood (PBMC) and bone marrow and the like can be used as the effector cells and human cells expressing GPC3 such as human hepatocellular carcinoma cell-line HuH-7 can be used as the target cells. First the target cells are labeled with ^{51}Cr , anti-GPC3 antibody is added, the cells are incubated, and then effector cells in a suitable ratio to the target cells are added, and they are incubated together. After incubation, the supernatant is collected, and the ADCC activity is measured by counting the radioactivity in the supernatant.

Anti-GPC3 Antibody

The anti-GPC3 antibody can be prepared by a method known to those skilled in the art. For example, the antibody can be prepared by using GPC3 as a sensitizing antigen for immunization in accordance with a conventional immunization method, fusing the immune cells with known parent cells by a conventional cell fusion procedure, and screening for monoclonal antibody producing cells by a conventional screening method. More specifically, monoclonal antibodies can be prepared in the following manner. First, the GPC3 to be used as a sensitizing antigen for antibody production is obtained

by expressing GPC3 (MXR7) based on the gene/amino acid sequence disclosed in Lage, H. et al., Gene 188 (1997), 151-156. In other words, the gene sequence encoding GPC3 is inserted into a known expression vector. After suitable host cells are transformed with the vector, the target human glycoprotein 3 protein is purified from the host cells or culture medium supernatant by a known method. Next the purified GPC3 protein is used as a sensitizing antigen. Alternatively, a partial peptide of GPC3 can be used as the sensitizing antigen. In such a process the partial peptide can be obtained by chemical synthesis according to the amino acid sequence of human GPC3. The epitopes on the GPC3 molecule recognized by the anti-GPC3 antibody of the present invention are not limited, but the anti-GPC3 antibody of the present invention may recognize any epitope present on the GPC3 molecule. This is because the anti-GPC antibody exhibits the cell growth inhibitory activity through its ADCC activity, CDC activity, or inhibition of growth factor activity, and because cell growth can also be inhibited by the action of a cytotoxic substance such as a radioactive isotope, chemotherapy drug, bacterial toxin attached to the anti-GPC3 antibody. Therefore, the antigen for preparing the anti-GPC3 antibody of the present invention can be any fragment of GPC3 provided it contains an epitope present on the GPC3 molecule.

In an especially preferred embodiment, a peptide containing amino acids 524-563 can be used as the sensitizing antigen to generate an antibody that recognizes an epitope of amino acids 524-563 of GPC3.

The mammal used for immunization with the sensitizing antigen is not particularly limited in the present invention, but preferably it should be selected in consideration of the compatibility with the parent cells to be used in cell fusion, and may include a rodent, for example, a mouse, rat, or hamster, or a rabbit, monkey, and the like. The animal may be immunized with the sensitizing antigen using a known method. In general, for example, the mammal can be injected intraperitoneally or subcutaneously with sensitizing antigen. More specifically, the sensitizing antigen can be diluted and suspended in a suitable amount of phosphate buffered saline (PBS) or physiological saline, mixed with a suitable amount of conventional adjuvant such as Freund's complete adjuvant if desired, emulsified, and administered to the mammal multiple times every 4 to 21 days. In addition, a suitable vehicle can be used upon immunization with the sensitizing antigen.

After the mammal is immunized in the above manner, and the desired antibody level is detected in the serum, the immune cells are collected from the mammal and subjected to cell fusion. Spleen cells are especially

preferred immune cells for cell fusion. Mammalian myeloma cells are used as the partner parent cells to be fused with the immune cells. Known cell lines suitable for use as the myeloma cells include, for example, P3 (P3x63Ag8.653) (J. Immunol. (1979) 123, 1548-1550), P3x63Ag8U.1 (Current Topics in Microbiology and Immunology (1978) 81, 1-7), NS-1 (Kohler. G. and Milstein, C. Eur. J. Immunol. (1976) 6, 511-519), MPC-11 (Margulies. D.H. et al., Cell (1976) 8, 405-415), SP2/0 (Shulman, M. et al., Nature (1978) 276, 269-270), FO (de St. Groth, S. F. et al., J. Immunol. Methods (1980) 35, 1-21), S194 (Trowbridge, I. S. J. Exp. Med. (1978) 148, 313-323), R210 (Galfre, G. et al., Nature (1979) 277, 131-133). The cell fusion of the immune cells and myeloma cells basically can be carried out in accordance with a known method, for example, the method described by Kohler and Milstein (Kohler. G. and Milstein, C., Methods Enzymol. (1981) 73, 3-46) . More specifically, the cell fusion can be carried out, for example, in a conventional liquid culture medium containing a cell fusion promoter. Examples of the cell fusion promoting chemicals include Polyethylene glycol (PEG) and Sendai virus (HVJ) and the like. If desired, a supplemental agent such as dimethyl sulfoxide and the like may be added to increase fusion efficiency. The ratio of immune cells to myeloma cells

may be established arbitrarily. For example, setting the ratio of immune cells with respect to myeloma cells at 1-fold to 10-fold is preferred. A conventional liquid culture medium used for culturing these types of cells, such as RPMI-1640 liquid medium, MEM liquid medium, or another liquid medium suitable for the growth of the myeloma cell line may be used as the liquid medium in the cell fusion procedure. A serum supplement such as fetal calf serum (FCS) can also be used together. In the cell fusion procedure, specified amounts of the immune cells and myeloma cells are thoroughly mixed in the liquid culture medium, and then PEG solution (for example, average molecular weight of about 1000 to 6000) that has been heated to 37°C is normally added at a concentration of 30 to 60% (w/v) and mixed to allow for forming fused cells (hybridomas). Next, a suitable liquid culture medium is added and centrifuged to remove the supernatant. By repeating this procedure, any cell fusion chemicals unfavorable for the growth of hybridomas are removed. Hybridomas obtained in this manner are selected by culturing them in a conventional liquid selection medium such as HAT medium (a culture medium containing hypoxanthine, aminopterin, and thymidine). Culturing in the HAT medium is continued for a sufficient period of time (normally a few days to a few weeks) until cells other than the target hybridomas (non-fused cells) die

off. Then, a conventional limiting dilution procedure is carried out, followed by screening and monocloning hybridomas that produce the target antibody. In addition to immunizing a non-human animal with the antigen to obtain the hybridomas as above, desired human antibodies having GPC3 binding activity can be obtained by sensitizing human lymphocytes with GPC3 in vitro, and then fusing the sensitized lymphocytes with immortalized human myeloma cells (see Japanese Patent Publication No. H1-59878). In addition, it is possible to administer GPC3 as an antigen to a transgenic animal having the complete repertoire of human antibody genes to generate cells producing anti-GPC3 antibodies, and collect human antibodies to GPC3 from immortalized cells (see International Patent Application No. WO 94/25585, WO 93/12227, WO 92/03918, and WO 94/02602). Hybridomas producing monoclonal antibodies prepared in the above manner can be subcultured in a conventional liquid culture medium, and may be preserved for a long period of time in liquid nitrogen.

Recombinant Antibodies

The monoclonal antibody used in the present invention is a recombinant monoclonal antibody, which can be produced by cloning the antibody gene from a hybridoma, inserting the gene into a suitable vector, and

integrating the vector into a host cell (for example, see Vandamme, A. M. et al., *Eur. J. Biochem.* (1990) 192; 767-775, 1990). More specifically, the mRNA encoding the variable (V) region of the anti-GPC3 antibody is isolated from a hybridoma producing the anti-GPC3 antibody. mRNA can be isolated using a known method such as the guanidine ultracentrifugation method (Chirgwin, J. M. et al., *Biochemistry* (1979) 18, 5294-5299), and the AGPC method (Chomczynski, P., et al., *Anal. Biochem.* (1987) 162, 156-159) to prepare total RNA, and then preparing the target mRNA using an mRNA Purification Kit (Pharmacia) and the like. The mRNA can also be prepared directly by using a QuickPrep mRNA Purification Kit (Pharmacia). Then cDNA of the antibody V region is synthesized from the mRNA thus obtained using reverse transcriptase. cDNA can be synthesized using the AMV Reverse Transcriptase First-strand cDNA Synthesis Kit (Seikagaku Corporation), and the like. Also the 5'-RACE method using the 5'-AmpliFINDER RACE Kit (Clontech) and PCR can be used for cDNA synthesis and amplification (Frohman, M. A. et al., *Proc. Natl. Acad. Sci. USA* (1988) 85, 8998-9002, Belyavsky, A. et al., *Nucleic Acids Res.* (1989) 17, 2919-2932). The target DNA fragment is purified from the PCR product and ligated to the vector DNA. The desired recombinant vector is prepared by inserting those vectors. It is introduced into *E. coli*

and the desired colony is selected to prepare a desired recombinant vector. The nucleotide sequence of the target DNA is confirmed by a known method such as the dideoxynucleotide chain termination method. After DNA encoding the V region of the target anti-GPC3 antibody is obtained, it is inserted into an expression vector containing DNA encoding the desired antibody constant region (C region). For the production of the anti-GPC3 antibody used in the present invention, the antibody gene is inserted into the expression vector so that it will be expressed under the control of an expression control region such as an enhancer, promoter, and the like. Next, the antibody is expressed by transforming a host cell with that expression vector. The antibody gene can be expressed in the host cells by inserting DNA encoding the antibody heavy chain (H chain) and light chain (L chain) into separate expression vectors and simultaneously transforming the host cells, or by inserting DNA encoding both the H chain and L chain into a single expression vector and transforming the host cells (see WO 94/11523). In addition, the recombinant antibody can be produced not only by using the aforementioned host cells, but also by using a transgenic animal. For example, the antibody gene can be inserted into the middle of a gene encoding a protein produced specifically in milk (such as goat β -casein) to prepare a fused gene. Then the DNA fragment

containing the fused gene containing the antibody gene is injected into a goat embryo, and the embryo is implanted in a female goat. The desired antibody can be obtained from the milk produced by the transgenic goat born from the goat implanted with the embryo and the offspring thereof. Furthermore, suitable hormones can be used in the transgenic goat to increase the amount of milk containing the desired antibody that is produced by the transgenic goat (Ebert, K. M. et al., Bio/Technology (1994) 12, 699-702).

Altered Antibodies

In addition to the antibodies as described above, an artificially altered gene recombinant antibody such as a chimeric antibody, humanized antibody, and the like can be used in the present invention for the purpose of reducing the xenoantigenicity to humans. Such modified antibodies can be produced according to known methods. A chimeric antibody can be obtained by ligating DNA encoding the antibody V region obtained as described above with DNA encoding the human antibody C region, and then inserting the DNAs into an expression vector. The vector in which the DNAs are inserted is integrated into host cells to produce the antibody. A chimeric antibody useful in the present invention can be obtained using such a conventional method. A humanized antibody, also

called a reshaped human antibody, comprises the CDR of an antibody from a non-human mammal such as a mouse grafted onto a human antibody CDR. The general genetic engineering methods for obtaining humanized antibodies are known in the art (see EP 125023 and WO 96/02576). More specifically, a DNA sequence designed to link the mouse antibody CDR and the human antibody framework region (FR) is synthesized by PCR using as primers a plurality of oligonucleotides prepared such that they have overlapped CDR and FR terminal regions (the method described in WO 98/13388). The framework region of the human antibody to be linked via the CDR is selected such that the CDR will form a suitable antibody binding site. If necessary, amino acids of the framework region in the variable region of the antibody may be substituted so that the reshaped human antibody CDR will form a suitable antibody binding site (Sato, K. et al., Cancer Res. (1993) 53, 851-856). A human antibody C region is used for the C region of the chimeric antibody and the humanized antibody. For example, Cy1, Cy2, Cy3, and Cy4 can be used in the H chain, and C κ and C λ can be used in the L chain. In addition, the human antibody C region can be modified to improve the stability or productivity of the antibody. The chimeric antibody comprises the variable region of an antibody from a non-human mammal and the constant region from a human antibody. On the

other hand, the humanized antibody comprises the CDR of an antibody from a non-human mammal and the framework region and C region from a human antibody. Because the humanized antibody has lower antigenicity in the human body, it is more useful as the active ingredient in the therapeutic agent of the present invention.

Modified Antibodies

The antibody used in the present invention is not limited to a whole molecule of antibody, but it may be an antibody fragment or a modified form of an antibody, provided it binds to GPC3 and inhibits the activity of GPC3. The present invention also encompasses bivalent antibodies as well as monovalent antibodies. Examples of an antibody fragment include Fab, F(ab')₂, Fv, Fab/c having one Fab and a complete Fc, or a single chain Fv (scFv) wherein the Fv of an H chain or L chain is linked by a suitable ligand. More specifically, to produce an antibody fragment, the antibody can be treated with an enzyme such as papain or pepsin, or a gene encoding such an antibody fragment can be constructed, inserted into an expression vector, and expressed in a suitable host cell (see, for example, Co, M.S. et al., J. Immunol. (1994) 152, 2968-2976, Better, M. & Horwitz, A. H. Methods in Enzymology (1989) 178, 476-496, Academic Press, Inc., Plueckthun, A. & Skerra, A. Methods in Enzymology (1989)

178, 476-496, Academic Press, Inc., Lamoyi, E., Methods in Enzymology (1989) 121, 652-663, Rousseaux, J. et al., Methods in Enzymology (1989) 121, 663-669, Bird, R. E. et al., TIBTECH (1991) 9, 132-137). A scFv can be obtained by joining the H chain V region and L chain V region of an antibody. In a scFv, the H chain V region and L chain V region are joined by a linker, preferably a peptide linker (Huston, J. S. et al., Proc. Natl. Acad. Sci. U.S.A. (1988) 85, 5879-5883). The H chain V region and L chain V region in the scFv may be derived from any antibodies described herein. Any single chain peptide comprising 12 to 19 amino acid residues may be used as the peptide linker joining the V regions. DNA encoding scFv can be obtained by amplifying a fragment by PCR using as a template a DNA portion encoding all or a desired amino acid sequence of the sequences of DNA encoding the H chain or the H chain V region of the above-mentioned antibody and DNA encoding the L chain or the L chain V region of the above-mentioned antibody with a primer pair that defines the both ends thereof. Then the fragment is amplified with a combination of DNA encoding a peptide linker portion and a primer pair which defines both ends to be ligated to the H chain and the L chain. Once DNA encoding scFv is prepared, an expression vector containing the DNA and a host cell transformed with the expression vector can be obtained according to a

standard method. The scFv can be obtained from such a host according to a standard method. These antibody fragments can be produced in a host by obtaining the gene thereof in the same manner as described above and by allowing it to be expressed. In the present invention, the term "antibody" also encompasses a fragment of the antibody. An anti-GPC3 antibody attached to various molecules such as PEG and the like can be used as a modified antibody. In the present invention, the term "antibody" also encompasses such a modified antibody. The modified antibody can be obtained by chemical modification of the antibody obtained as above. Methods for modifying antibodies have already been established in the art.

Furthermore, the antibody used in the present invention can be a bispecific antibody. A bispecific antibody may be an antibody having an antigen binding site that recognizes a different epitope on the GPC3 molecule, or it may be an antibody wherein one antigen binding site recognizes GPC3 and the other antigen binding site recognizes a cytotoxic substance such as a chemotherapy drug and cell-derived toxin. In such a case, the cytotoxic substance will act directly on cells expressing GPC3, and specifically lesion tumor cells, and suppress the growth of tumor cells. A bispecific antibody may be prepared by linking two types of antibody

HL pairs. Also it may be obtained by preparing a bispecific antibody-producing fused cell through the fusion of hybridomas that produce different monoclonal antibodies. A bispecific antibody can also be prepared by genetic engineering methods.

Expression and Production of Recombinant Antibodies or Modified Antibodies

An antibody gene constructed as noted above can be expressed and obtained by known methods. In the case of mammalian cells, a common useful promoter, gene to be expressed, and a poly-A signal sequence downstream on the 3' end can be functionally linked together and expressed. For example, the human cytomegalovirus immediate early promoter/enhancer can be used as the promoter/enhancer. In addition, other promoters/enhancers that can be used to express the antibody of the present invention include viral promoters/enhancers of retrovirus, polyomavirus, adenovirus and simian virus 40 (SV40), or promoters/enhancers from mammalian cells such as human elongation factor 1a (HEF-1a). Antibodies can be readily expressed by the method of Mulligan et al. (Nature (1979) 277, 108) when SV40 promoter/enhancer is used, and by the method of Mizushima et al. (Nucleic Acids Res. (1990) 18, 5322) when HEF1 α promoter/enhancer is used.

The antibody of the present invention may be produced using an eukaryotic expression system having the capability of adding a sugar chain to the expressed antibody. Eukaryotic cells include, for example, established mammalian cell lines and insect cell lines, animal cells, fungal cells, and yeast cells.

Preferably, the antibody of the present invention is expressed in mammalian cells, for example, CHO, COS, myeloma, BHK, Vero, or HeLa cells. The target antibody is produced by culturing the transformed host cells either in vitro or in vivo. The host cells may be cultured using known methods. For example, DMEM, MEM, RPMI-1640, or IMDM may be used as the culture medium, and a serum complement such as fetal calf serum (FCS) may be supplemented.

Isolation and Purification of Antibody

The antibody expressed and produced in the above manner can be separated from the host cells or host animals and purified to homogeneity. The antibody of the present invention can be separated and purified using an affinity column, for example, a protein A column such as Hyper D, POROS and Sepharose F.F. (Pharmacia). In addition, any conventional methods for protein separation and purification may be used in the invention. For example, the antibody can be isolated and purified by

appropriately selecting and combining affinity columns such as Protein A column with chromatography columns, filtration, ultra filtration, salting-out and dialysis procedures (Antibodies A Laboratory Manual, Ed Harlow, David Lane, Cold Spring Harbor Laboratory, 1988). An antibody having a desired sugar chain can be separated with a lectin column by a method known in the art, and the method described in WO 02/30954.

Determination of Antibody Activity

The antigen binding activity (Antibodies: A Laboratory Manual. Ed Harlow, David Lane, Cold Spring Harbor Laboratory, 1988) and ligand receptor binding inhibition (Harada, A. et al., International Immunology (1993) 5, 681-690) of the antigen used in the present invention may be measured by known methods. ELISA (enzyme-linked immunosorbent assay), EIA (enzyme immunoassay), RIA (radioimmunoassay), or the fluorescent antibody technique can be used for measuring antigen binding activity of the anti-GPC3 antibody of the present invention. For example, EIA is carried out as follows. A sample containing the anti-GPC3 antibody, such as culture supernatant of anti-GPC3 antibody producing cells or purified antibody, is added to a plate coated with GPC3. A secondary antibody labeled with an enzyme such as alkaline phosphatase is added, the plate is incubated

and washed, and then the enzyme substrate, such as p-nitrophenyl phosphate, is added and the optical absorption is measured to evaluate the antigen binding activity.

Pharmaceutical Composition

The present invention provides a pharmaceutical composition comprising the anti-GPC3 antibody with a modified sugar chain component of the present invention.

A pharmaceutical composition comprising the antibody composition of the present invention is useful for the prevention and/or treatment of diseases associated with cell growth such as cancer, and is particularly useful for the prevention and/or treatment of liver cancer. The pharmaceutical composition comprising the antibody of the present invention can be formulated by methods known to those skilled in the art. The pharmaceutical composition can be administered parenterally in the form of an injectable formulation comprising a sterile solution or suspension in water or another pharmaceutically acceptable liquid. For example, the pharmaceutical composition can be formulated by suitably combining the antibody with pharmaceutically acceptable vehicles or media, such as sterile water and physiological saline, vegetable oil, emulsifier, suspension agent, surfactant, stabilizer, flavoring excipient, diluent, vehicle,

preservative, binder, followed by mixing in a unit dose form required for generally accepted pharmaceutical practices. The amount of active ingredient included in the pharmaceutical preparations is such that a suitable dose within the designated range is provided.

The sterile composition for injection can be formulated in accordance with conventional pharmaceutical practices using distilled water for injection as a vehicle.

For example, physiological saline or an isotonic solution containing glucose and other supplements such as D-sorbitol, D-mannose, D-mannitol, and sodium chloride may be used as an aqueous solution for injection, optionally in combination with a suitable solubilizing agent, for example, alcohol such as ethanol and polyalcohol such as propylene glycol or polyethylene glycol, and a nonionic surfactant such as polysorbate 80™, HCO-50 and the like.

Examples of oily liquid include sesame oil and soybean oil, and it may be combined with benzyl benzoate or benzyl alcohol as a solubilizing agent. Other items that may be included are a buffer such as a phosphate buffer, or sodium acetate buffer, a soothing agent such as procaine hydrochloride, a stabilizer such as benzyl alcohol or phenol, and an antioxidant. The formulated injection will normally be packaged in a suitable ampule.

Route of administration is preferably parenteral, for example, administration by injection, transnasal administration, transpulmonary administration, transcutaneous administration. Administration may be systemic or local by intravenous injection, intramuscular injection, intraperitoneal injection, subcutaneous injection.

A suitable means of administration can be selected based on the age and condition of the patient. A single dose of the pharmaceutical composition containing the antibody or a polynucleotide encoding the antibody can be selected from a range of 0.001 to 1000 mg/kg of body weight. On the other hand, it is also possible to select a dose in the range of 0.001 to 100000 mg/body, but the present invention is by no means limited to such numerical ranges. The dose and method of administration will vary depending on the weight, age, condition, and the like of the patient, and can be suitably selected as needed by those skilled in the art.

The content of all patents and reference documents expressly cited in the specification of this application are hereby incorporated by reference in its entirety. In addition, the content of the specification and drawings of Japanese Patent Application 2004-311356, which is the basis for the priority claim of this application, are hereby incorporated by reference in its entirety.

EXAMPLES

The present invention is explained in detail through the following examples, but is by no means limited to these examples.

Example 1

Preparation of mouse anti-GPC3 antibody

A soluble GPC3 protein lacking the hydrophobic region of the C terminus (amino acids 564 to 580) was prepared as the immunizing protein for the preparation of the anti-GPC3 antibody for immunization. The MRL/MpJUmCrj-lpr/lpr mouse (hereinafter referred to as MRL/lpr mouse, purchased from Charles River Japan), which is an autoimmune disease mouse, was used as the immunization animal. Immunization was started when the mice were 7 or 8 weeks old, and a preparation for initial immunization was adjusted to a dose of 100 µg/head soluble GPC3. An emulsion was prepared using Freund's complete adjuvant (FCA, Becton Dickinson), and injected subcutaneously. After a series of five immunizations, the final immunization dose was diluted in PBS to 50 µg/head, and injected intravenously via the caudal vein. On day 4 after the final immunization, the spleen cells were resected, mixed with mouse myeloma cells P3-X63Ag8U1 (hereinafter referred to as P3U1, purchased from ATCC) in

a 2:1 ratio, and cell fusion was carried out by gradually adding PEG-1500 (Roche Diagnostics). Hybridomas were screened by ELISA using immunoplates with immobilized soluble GPC3 core protein. Positive clones were monocloned by the limiting dilution procedure. As a result, 11 clones of antibodies with strong GPC3 binding activity were obtained (M3C11, M13B3, M1E7, M3B8, M11F1, L9G11, M19B11, M6B1, M18D4, M5B9, and M10D2).

Among the anti-GPC3 antibodies obtained, M11F1 and M3B8 exhibited particularly strong CDC activity. Thus the GST fusion protein containing the M11F1 and M3B8 epitopes (GC-3), which is the fusion protein containing a peptide from 524 Ala to 563 Lys of GPC3 and GST was used as immunogens for immunization of 3 Balb/c (Charles River Japan) mice and 3 MRL/lpr mice. For the first immunization, a preparation of GC-3 at a concentration of 100 µg/head was emulsified with FCA and was injected subcutaneously. After two weeks, a preparation of 50 µg/head was emulsified with Freund's incomplete adjuvant (FIA) and was injected subcutaneously. After five immunizations, the final immunization (50 µg/head) was injected intravenously to all mice via the caudal vein, and subjected to the cell fusion. Hybridomas were screened by ELISA using immunoplates with immobilized soluble GPC3 core protein lacking the hydrophobic region of the C terminus (amino acids 564-580). Positive clones

were monocloned by the limiting dilution procedure. As a result, 5 clones of antibodies with strong GPC3 binding activity (GC199, GC202, GC33, GC179, and GC194) were obtained.

The H chain and L chain variable regions were cloned and each sequence was determined by standard method. Furthermore, the CDR regions were determined by comparison with a known antibody amino acid sequence database and checking for homology. The sequences of the CDR regions are shown in Tables 1 and 2.

Example 2

Preparation of anti-GPC3 antibody mouse-human chimeric antibody

The variable region sequences of the H chain and L chain of anti-GPC3 antibody GC33 were ligated to the constant region sequences of human IgG1 and κ chain. PCR was carried out using a synthetic oligonucleotide complementary to the 5' terminal nucleotide sequence of the antibody H chain variable region having a Kozak sequence and a synthetic oligonucleotide complementary to the 3' terminal nucleotide sequence having an NheI site. The PCR product thus obtained was cloned into pB-CH vector wherein the human IgG1 constant region has been inserted into pBluescript KS+ vector (Toyobo Co., Ltd.). The mouse H chain variable region and the human H chain

(γ 1 chain) constant region were ligated via the NheI site. The H chain gene fragment thus prepared was cloned into the expression vector pCXND3. Furthermore, PCR was carried out using a synthetic oligonucleotide complementary to the 5' terminal nucleotide sequence of the L variable region of the antibody having a Kozak sequence and a synthetic oligonucleotide complementary to the 3' terminal nucleotide sequence having a BsiWI site. The PCR product thus obtained was cloned into pB-CL vector wherein the human κ chain constant region has been inserted into pBluescript KS+ vector (Toyobo Co., Ltd.). The human L chain variable region and the constant region were ligated via the BsiWI site. The L chain gene fragment thus prepared was cloned into the expression vector pUCAG. The pUCAG vector is a vector prepared by digesting pCXN (Niwa et al., Gene, 1991, 108, 193-200) with the restriction enzyme BamHI to prepare a 2.6 kbp fragment, which was ligated to the restriction enzyme BamHI site of pUC19 vector (Toyobo Co., Ltd.)

To prepare the anti-GPC3 mouse-human chimeric antibody expression vector, a gene fragment was obtained by digesting pUCAG vector, in which the L chain gene fragment was inserted, with the restriction enzyme HindIII (Takara Shuzo Co., Ltd.). This gene fragment was ligated to the restriction enzyme HindIII cleavage site of pCXND3 containing the H chain gene, and then cloned.

The plasmid thus obtained expressed the neomycin resistance gene, DHFR gene, and anti-GPC3 mouse-human chimeric antibody gene in animal cells. (The amino acid sequence of the H chain variable region is shown in SEQ ID NO: 3, and the amino acid sequence of the L chain variable region is shown in SEQ ID NO: 4.)

Example 3

Preparation of low-fucose type anti-GPC3 chimeric antibody

First YB2/0 (ATCC, CRL-1662) cells were cultured as the host cells in RPMI-1640 medium containing 10% FBS. Then 25 µg of the anti-GPC3 chimeric antibody expression vector prepared in Example 2 was introduced into the YB2/0 cells (ATCC CRT-1662) by electroporation at a concentration of 7.5×10^6 cells/0.75 mL PBS(-) at 1.4 kV and 25 µF. After a recovery period of 10 min at room temperature, the cells treated by electroporation were suspended in 40 mL of RPMI-1640 medium containing 10% FBS. A 10-fold dilution was prepared using the same medium, and the cells were aliquoted into a 96-well culture plate at 100 µL/well. After culturing for 24 h in a CO₂ incubator (5% CO₂), Geneticin (Invitrogen Corp.) was added at a concentration of 0.5 mg/mL, and the cells were cultured for 2 weeks. Cell lines with a high level of chimeric antibody expression were screened using sandwich

ELISA with anti-human IgG antibody, and cell lines stably expressing the antibody were established. Each anti-GPC3 mouse-human chimeric antibody was purified using Hi Trap ProteinG HP (Amersham).

Example 4

Measurement of ADCC activity using PBMC from human peripheral blood

Preparation of human PBMC solution

Heparin-added peripheral blood collected from a healthy adult was diluted 2-fold with PBS(-), and layered on Ficoll-Paque™ PLUS (Amersham). After centrifugation (500 ×g, 30 min, 20°C), the intermediate layer, which is the mononuclear cell fraction, was isolated. After the layer was washed 3 times, the cells were suspended in 10% FBS/RPMI to prepare a human PBMC solution.

Preparation of target cells

HepG2 cells (ATCC) and HuH-7 cells (Health Science Research Resources Bank) cultured in 10% FBS/RPMI-1640 medium were detached from the dish using Cell Dissociation Buffer (Invitrogen), aliquoted into a 96-well U-bottomed plate (Falcon) at a concentration of 1×10^4 cells/well, and cultured for one day. After culturing, 5.55 MBq of ^{51}Cr was added, and the cells were cultured for 1 h at 37°C in a 5% CO₂ gas incubator. The cells were washed once with culture medium, and 50 µL of 10%

FBS/RPMI-1640 medium was added to prepare the target cells.

Chromium release assay (ADCC activity)

A volume of 50 μ L of antibody solution prepared at various concentrations was added to the target cells, and allowed for reacting on ice for 15 min. Then 100 μ L of human PBMC solution (5×10^5 cells/well) was added, and the cells were incubated for 4 h at 37°C in a 5% CO₂ gas incubator. After culturing, the plate was centrifuged and the radioactivity in 100 μ L of culture supernatant was measured with a gamma-counter. The specific chromium release rate was determined by the following formula.

$$\text{Specific chromium release rate (\%)} = (A - C) \times 100 / (B - C)$$

In this formula, A represents the mean value of radioactivity (cpm) in each well; B represents the mean value of radioactivity (cpm) in a well wherein 100 μ L of 2% NP-40 aqueous solution (Nonidet P-40, Code No. 252-23, Nacalai Tesque) and 50 μ L of 10% FBS/RPMI medium were added to the target cells; and C represents the mean value of radioactivity (cpm) in a well wherein 150 μ L of 10% FBS/RPMI medium was added to the target cells. The assay was conducted in triplicate, and the means and standard deviations of ADCC activity (%) were calculated.

Figures 2 and 3 show the ADCC activity of the anti-GPC3 chimeric antibody measured using PBMC. In the figures, the vertical axis represents cytotoxic activity

(%) and the horizontal axis represents the concentration ($\mu\text{g/mL}$) of antibody added. Figure 2 shows the results when HepG2 cells are used as the target cells and Figure 3 shows the results for HuH-7 cells. The open circles show the activity of chimeric GC33 antibody produced by CHO cells, and the filled circles show the activity of chimeric GC33 antibody produced by YB2/0 cells. The low fucose type GC33 chimeric antibody produced by the YB2/0 cells shows stronger ADCC activity than the GC33 chimeric antibody produced by CHO cells, clearly indicating that ADCC activity of the anti-GPC3 antibody is enhanced by sugar chain modification.

Example 5

Establishment of antibody producing cells

Hygromycin B was added to SFMII(+) medium at a final concentration of 1 mg/mL, and a fucose transporter deficient cell line (clone 3F2) was subcultured in the medium. A suspension of 3F2 cells in Dulbecco phosphate buffer (8×10^6 cells/0.8 mL) was prepared. To the cell suspension, 25 μg of antibody expression vector was added (Reference Examples 1 and 2), and the cell suspension was transferred to a Gene Pulser Cuvette. After the cuvette was let stand on ice for 10 min, the vector was introduced into the cells by electroporation using a GENE-PULSER II at 1.5 kV and 25 μFD . The cells were

suspended in 40 mL of SFMII(+) medium and transferred to a 96-well flat bottom plate (Iwaki) at 100 μ L/well. After the plate was incubated in a CO₂ incubator for 24 h at 37°C, Geneticin (Invitrogen, Cat. No. 10131-027) was added at a final concentration of 0.5 mg/mL. The amount of antibody produced by the drug-resistant cells was measured to establish humanized anti-GPC3 antibody producing cell lines.

Example 6

Antibody purification

The supernatant from the antibody expressing cell line was collected and loaded on Hitrap™ rProtein A column (Pharmacia Cat. No. 17-5080-01) using a P-1 pump (Pharmacia). After the column was washed with a binding buffer (20 mM sodium phosphate (pH 7.0)), and the protein was eluted with an elution buffer (0.1 M Glycin-HCl (pH 2.7)). The eluate was immediately neutralized with neutralizing buffer (1 M Tris-HCl (pH 9.0)). The antibody elution fractions were selected by DC protein assay (BIO-RAD Cat. No. 500-0111) and pooled, and were concentrated to about 2 mL with a Centriprep-YM10 (Millipore Cat. No. 4304). Next, the antibodies were separated by gel filtration using a Superdex 200 26/60 column (Pharmacia) equilibrated with 20 mM acetate buffer with 150 mM NaCl (pH 6.0). The monomer fraction peaks

were collected, concentrated with Centriprep-YM10, and filtered through MILLEX-GW 0.22 μ m Filter Unit (Millipore Cat. No. SLGV 013SL), and then preserved at 4°C. The absorption at 280 nm was measured and the concentration of purified antibody was calculated from the molar absorption coefficient.

Example 7

In vitro ADCC activity of humanized anti-GPC3 antibody produced by FT-KO cells

Figure 4 shows the in vitro ADCC activity of anti-GPC3 antibody produced by FT-KO cells when human PBMC is used. The method is as described in Example 4. In the figure, the vertical axis represents cytotoxic activity (%) and the horizontal axis represents the concentration (μ g/mL) of antibody added. HuH-7 cells were used as the target cells. The open circles show the activity of anti-GPC3 antibody produced by wild type CHO cells, and the filled circles show the activity of anti-GPC3 antibody produced by FT-KO cells. The low fucose type anti-GPC3 antibody produced by the FT-KO cells shows stronger ADCC activity than the anti-GPC3 antibody produced by wild type CHO cells, clearly indicating that ADCC activity of the anti-GPC3 antibody produced by FT-KO cells is enhanced.

Example 8

Analysis of sugar chains of humanized anti-GPC3 antibody produced by FT-KO cells

1. Preparation of 2-Aminobenzamide-labeled sugar chains (2-AB labeled sugar chains)

The antibodies produced by the FT-KO cells of the present invention and antibodies produced by CHO cells as a control sample were treated with N-Glycosidase F (Roche Diagnostics) to release the sugar chains from the protein (Weitzhandler M. et al., Journal of Pharmaceutical Sciences 83:12 (1994), 1670-1675). After removing the protein with ethanol (Schenk B. et al., The Journal of Clinical Investigation 108:11 (2001), 1687-1695), the sugar chains were concentrated and dried, and fluorescent labeled with 2-aminopyridine (Bigge J. C. et al., Analytical Biochemistry 230:2(1995), 229-238). The reagent was removed from the 2-AB labeled sugar chains by solid phase extraction using a cellulose cartridge, and after concentration by centrifugation, purified 2-AB labeled sugar chains were obtained. Next, the purified 2-AB labeled sugar chains were treated with β -galactosidase (Seikagaku Corp.) to obtain agalactosyl 2-AB labeled sugar chains.

2. Analysis of agalactosyl 2-AB labeled sugar chains by normal phase HPLC

The antibodies produced by the FT-KO cells of the

present invention and the antibodies produced by the CHO cells as a control sample were prepared as agalactosyl 2-AB labeled sugar chains according to the above method, and analyzed by normal phase HPLC using an amide column (Tosoh Corp. TSKgel Amide-80), and the chromatograms were compared. In the antibodies produced by the CHO cells, the main component is G(0), and G(0)-Fuc accounts for about 4% of the peak area. On the other hand, in the antibodies produced by the FT-KO cells, G(0)-Fuc is the main component, and is present at not less than 90% of the peak area in each of the cell lines (Figure 5 and Table 2). Figure 6 shows the putative structures for peaks G(0) and G(0)-Fuc.

Table 2

Relative ratio of sugar chains estimated from normal phase HPLC analysis of agalactosyl 2-AB sugar chains

Sugar chain	CHO	FT-KO-a	FT-KO-b	FT-KO-c
G(0)-Fuc	4.0%	92.4%	92.5%	93.2%
G(0)	96.0%	7.6%	7.5%	6.8%

Example 9

Thermal stability analysis of humanized anti-GPC3

antibody produced by FT-KO cells

1. Preparation of sample solution for DSC measurement

The external dialysis solution was 20 mol/L sodium

acetate buffer (pH 6.0) containing 200 mmol/L sodium chloride. A dialysis membrane filled with 700 µg equivalents of antibody solution was dialyzed by immersing in the external dialysis solution overnight to prepare a sample solution.

2. Measurement of thermal degradation temperature by DSC

After both the sample solution and reference solution (external dialysis solution) were thoroughly degassed, they were each placed in the calorimeter and thermally equilibrated at 20°C. Next DSC measurement was carried out from 20°C to 100°C at a scan rate of approximately 1 K/min. The result is represented by the tip of the degradation peak as a function of temperature (Figure 7). The thermal degradation temperature of the antibodies produced by CHO cells and the antibodies produced by FT-KO cells were found to be equivalent.

Reference Example 1

Humanization of GC33

Antibody sequence data were obtained from the publicly disclosed Kabat Database (<ftp://ftp.ebi.ac.uk/pub/databases/kabat/>) and the ImMunoGeneTics Database (IMGT), and the H chain variable region and L chain variable region were separately subjected to a homology search. It was found that the H chain variable region has a high level of homology with

DN13 (Smithson et al., Mol. Immunol. 1999; 36: 113-124). It was also found that the L chain variable region has a high level of homology with the Homo sapiens IGK mRNA for immunoglobulin kappa light chain VLJ region, partial cds, clone:K64 of Accession Number AB064105. The signal sequence of Accession Number S40357, which has a high level of homology with AB064105, was used as the L chain signal sequence. Then the CDR was grafted to the FR of these antibodies to prepare a humanized antibody.

More specifically, synthetic oligo-DNAs of approximately 50 base were designed in such a manner that approximately 20 bases of them were hybridized each other, and these synthetic oligo-DNAs were assembled by PCR to prepare a gene encoding each variable region. They were digested at the HindIII sequence inserted at the terminus on the 5' end of the synthetic oligo-DNA and the BamHI sequence inserted at the terminus on the 3' end of the synthetic oligo-DNA, and the synthetic oligo-DNA was cloned into an expression vector HEFg λ 1 where the human IgG1 constant region was cloned, or to the expression vector HEFg κ where the human κ -chain constant region was cloned (Sato et al., Mol. Immunol., 1994; 371-381). The H chain and L chain of the humanized GC33 constructed as above were each designated ver.a. The humanized GC33 (ver.a/ver.a) wherein both the H chain and L chain were ver.a had lower binding activity than an antibody with

the mouse GC33 variable regions (mouse/mouse). Chimeric antibodies were prepared by combining mouse GC33 sequences and ver.a sequences for the H chains and L chains (mouse/ver.a, ver.a/mouse), and the binding activity was evaluated. Lower binding activity was found with ver.a/mouse antibody, indicating that the decrease in binding activity is due to amino acid replacement was attributed to the H chain. Then modified H chains designated ver.c, ver.f, ver.h, ver.i, ver.j, and ver.k were prepared. All humanized GC33 antibodies exhibited the same level of binding activity as the chimeric antibody having mouse GC33 variable regions. The nucleotide sequences of the humanized GC33 H chain variable regions ver.a, ver.c, ver.f, ver.h, ver.i, ver.j, and ver.k are shown in SEQ ID NOS:74, 75, 76, 77, 78, 79, and 80, and the amino acid sequences thereof are shown in SEQ ID NOS: 81, 82, 83, 84, 85, 86, and 87, respectively. The nucleotide sequence of the humanized GC33 L chain variable region ver.a is shown in SEQ ID NO: 88, and the amino acid sequence thereof is shown in SEQ ID NO: 89, respectively. In the humanized GC33 H chain variable regions ver.i, ver.j, and ver.k, the sixth glutamic acid was replaced by a glutamine. These antibodies exhibited markedly increased thermal stability.

Reference Example 2

Alteration of humanized GC33 L chain

With respect to protein deamidation, the reaction rate constant of deamidation was known to be dependent on the primary sequence. It is also known that Asn-Gly is particularly susceptible to deamidation (Rocinson et al., Proc. Natl. Acad. Sci. USA 2001; 98: 944-949). Because Asn 33 within CDR1 of the humanized GC33 L chain ver.a variable region of SEQ ID NO: 88 has the primary sequence Asn-Gly, this residue is predicted to be susceptible to deamidation.

To evaluate the effect of deamidation of Asn 33 on binding activity of the antibody, a modified antibody was prepared wherein Asn 33 was replaced with Asp. Quick Change Site-Directed Mutagenesis Kit (Stratagene) was used for introducing a point mutation. More specifically, 50 μ L of reaction solution containing 125 ng of sense primer (CTT GTA CAC AGT GAC GGA AAC ACC TAT: SEQ ID NO: 124), 125 ng of antisense primer (ATA GGT GTT TCC GTC ACT GTG TAC AAG: SEQ ID NO: 125), 5 μ L of 10x reaction buffer, 1 μ L of dNTP mix, 10 ng of HEFgk to which humanized GC33 L chain ver.a had been cloned, and 1 μ L of Pfu Turbo DNA Polymerase was run through 12 cycles consisting of 30 sec at 95°C, 1 min at 55°C, and 9 min at 68°C. The reaction product was digested with the restriction enzyme DpnI for 2 h at 37°C, and introduced into XL1-Blue competent cells to obtain transformants. The variable region was cut out

from the clones containing the correct mutation and cloned again into the expression vector HEFg_K. The expression vector HEFg_{Y1} containing humanized GC33 H chain ver.k was introduced into COS7 cells by using Eugene 6 (Roche). Culture medium supernatant was collected from the cells transiently expressing the modified antibody. The antibody concentration was quantitated by sandwich ELISA using anti-human IgG antibody, and binding activity of the modified antibody was evaluated by ELISA using a plate coated with a soluble GPC3 core protein. Binding activity was lost in the modified antibody (N33D) in which Asn 33 was replaced by Asp, suggesting that the binding activity is significantly affected by deamidation at Asn 33.

Deamidation of Asn 33 was reported to be suppressed by replacing Gly 34 with another amino acid residue (WO 03057881 A1). In accordance with that method, a series of modified antibodies were prepared by replacing G34 with 17 other amino acid residues except for Cys and Met using the Quick Chane Site-Directed Mutagenesis Kit to prepare G34A, G34D, G34E, G34F, G34H, G34N, G34P, G34Q, G34I, G34K, G34L, G34V, G34W, G34Y, G34R, G34S, and G34T. The binding activity of the antibodies was evaluated using culture supernatant of COS7 cells transiently expressing the antibodies. It was revealed that binding activity is maintained even if G34 is replaced with

another amino acid residues other than Pro (G34P), and Val (G34V).

The amino acid sequences of the L chain CDR1 of the modified antibodies are represented by SEQ ID NO: 90 (G34A), SEQ ID NO: 91 (G34D), SEQ ID NO: 92 (G34E), SEQ ID NO: 93 (G34F), SEQ ID NO: 94 (G34H), SEQ ID NO: 95 (G34N), SEQ ID NO: 96 (G34T), SEQ ID NO: 97 (G34Q), SEQ ID NO: 98 (G34I), SEQ ID NO: 99 (G34K), SEQ ID NO: 100 (G34L), SEQ ID NO: 101 (G34S), SEQ ID NO: 102 (G34W), SEQ ID NO: 103 (G34Y), SEQ ID NO: 104 (G34R), SEQ ID NO: 105 (G34V), and SEQ ID NO: 106 (G34P), respectively. The amino acid sequences of the L chain variable regions of the modified antibodies are represented by SEQ ID NO: 107 (G34A), SEQ ID NO: 108 (G34D), SEQ ID NO: 109 (G34E), SEQ ID NO: 110 (G34F), SEQ ID NO: 111 (G34H), SEQ ID NO: 112 (G34N), SEQ ID NO: 113 (G34T), SEQ ID NO: 114 (G34Q), SEQ ID NO: 115 (G34I), SEQ ID NO: 116 (G34K), SEQ ID NO: 117 (G34L), SEQ ID NO: 118 (G34S), SEQ ID NO: 119 (G34W), SEQ ID NO: 120 (G34Y), SEQ ID NO: 121 (G34R), SEQ ID NO: 122 (G34V), and SEQ ID NO: 123 (G34P), respectively.

Reference Example 3

Destruction of fucose transporter gene in CHO cells

1. Construction of targeting vector

(1) Preparation of KO1 vector

The hygromycin resistance gene (Hygr) was

constructed by PCR with Hyg5-BH and Hyg3-NT primers from pcDNA3.1/Hygro (Invitrogen), which has a sequence identical to the 5' portion of the fucose transporter gene start codon by attaching a BamHI site and TGCGC sequence to the 5' portion of the start codon and a NotI site added to the 3' portion containing the region up to the SV40 polyA addition signal, and the Hygr fragment was cut off.

Forward primer

Hyg5-BH 5'-GGA TCC TGC GCA TGA AAA AGC CTG AAC TCA CC-
3' (SEQ ID NO: 128)

Reverse primer

Hyg3-NT 5'-GCG GCC GCC TAT TCC TTT GCC CTC GGA CG-3'
(SEQ ID NO: 129)

The fucose transporter targeting vector ver.1 (hereinafter designated the KO1 vector) was constructed by inserting the 5' portion of the fucose transporter (from the SmaI at base No. 2780 to the BamHI at base No. 4323 of the nucleotide sequence shown in SEQ ID NO: 126), the 3' portion (from base No. 4284 to the SacI at base No. 10934), and an Hygr fragment into pMC1DT-A vector (Yagi T, Proc. Natl. Acad. Sci. USA, Vol. 87, p. 9918-9922, 1990). The characteristic of the KO1 vector is that Hygr will be expressed from the fucose transporter promoter when homologous recombination takes place because no promoter is attached to the Hygr fragment. However, Hygr is not

always expressed to the extent that resistance to hygromycin B is acquired if only one copy of the vector is inserted into a cell by homologous recombination. The KO1 vector was cleaved by NotI and introduced into the cell. It is expected that the fucose transporter will lose 41 base pairs of exon 1 including the start codon by introduction of the KO1 vector, which will result in the loss of its function.

(2) Preparation of pBSK-pgk-1-Hygr

The pBSK-pgk-1 vector was prepared by cutting off the mouse pgk-1 gene promoter from pKJ2 vector (Popo H, Biochemical Genetics, Vol. 28, p. 299-308, 1990) with EcoRI-PstI, and cloning it into the EcoRI-PstI site of pBluescript (Stratagene). By PCR with the Hyg5-AV and Hyg3-BH primers from pcDNA3.1/Hygro, an EcoT22I site and Kozak sequence were attached to the 5' portion of Hygr, and a BamHI site was added to the 3' portion containing the region up to the SV40 poly A addition signal, and then the Hygr fragment was cut off.

Forward primer

Hyg5-AV 5'-ATG CAT GCC ACC ATG AAA AAG CCT GAA CTC ACC
-3' (SEQ ID NO: 130)

Reverse primer

Hyg3-BH 5'-GGA TCC CAG GCT TTA CAC TTT ATG CTT C -3'
(SEQ ID NO: 131)

The pBSK-pgk-1-Hygr vector was prepared by inserting

the Hygr (EcoT22I-BamHI) fragment into the PstI-BamHI site of pBSK-pgk-1.

(3) Preparation of the KO2 vector

The fucose transporter targeting vector ver.2 (hereinafter designated the KO2 vector) was constructed by inserting the 5' portion of the fucose transporter (from the SmaI at base No. 2780 to the BamHI at base No. 4323 of the nucleotide sequence shown in SEQ ID NO: 126), the 3' portion (from base No. 4284 to the SacI at base No. 10934), and pgk-1Hygr fragment into pMC1DT-A vector. Unlike the KO1 vector, KO2 vector will confer resistance to hygromycin B even if only one copy of the vector is inserted by homologous recombination because the pgk-1 gene promoter is attached to Hygr. The KO2 vector was cleaved by NotI and inserted into the cells. It is expected that the fucose transporter will lose 46 base pairs of exon 1 including the start codon by the introduction of the KO2 vector, which will result in the loss of its function.

(4) Preparation of pBSK-pgk-1-Puror

The pBSK-pgk-1-Puror vector was prepared by cleaving pPUR vector (BD Biosciences) with PstI and BamHI, and inserting the digested fragment (Puror) into the PstI-BamHI site of pBSK-pgk-1.

(5) Preparation of the KO3 vector

The fucose transporter targeting vector ver.3

(hereinafter designated the KO3 vector) was constructed by inserting the 5' portion of the fucose transporter (from the SmaI at base No. 2780 to the BamHI at base No. 4323 of the nucleotide sequence shown in SEQ ID NO: 126), the 3' portion (from base No. 4284 to the SacI at base No. 10934), and pgk-1-Puror fragment into pMC1DT-A vector. In addition, a sequence for binding with the primer for screening shown below was attached to the 3' end of pgk-1-Puror. The KO3 vector was cleaved by NotI and inserted into the cells. It is expected that the fucose transporter will lose 46 base pairs of exon 1 including the start codon by the introduction of the KO3 vector, which will result in the loss of its function.

Reverse primer

RSGR-A 5'-GCT GTC TGG AGT ACT GTG CAT CTG C -3' (SEQ ID NO: 132)

The above three species of targeting vectors were used to knock out the fucose transporter gene.

2. Introduction of vectors into CHO cells

HT Supplement (100x) (Invitrogen Cat. No. 11067-030) and penicillin-streptomycin (Invitrogen Cat. No. 15140-122) were added to CHO-S-FMII HT (Invitrogen Cat. No. 12052-098), each at a volume of 1/100 with respect to the volume of CHO-S-SFMII HT. CHO DXB11 cells were subcultured in the culture medium (hereinafter designated SFMII(+)), and this SFMII(+) medium also used for

culturing the cells after gene transfer. The CHO cells were suspended in Dulbecco phosphate buffer (hereinafter designated PBS, Invitrogen Cat. No. 14190-144) at a concentration of 8×10^6 cells/0.8 mL. Then 30 μ g of the targeting vector was added to the cell suspension, and the cell suspension was transferred to a Gene Pulser Cuvette (4 mm) (Bio-Rad, Cat. No. 1652088). After the cuvette was let stand on ice for 10 min, the vector was introduced into the cells by electroporation with a GENE-PULSER II (Bio-Rad, Code No. 340BR) at 1.5 kV and 25 μ FD. After introduction of the vector, the cells were suspended in 200 mL of SFMII(+) medium and transferred to twenty 96-well flat bottomed plates (Iwaki, Cat. No. 1860-096) at 100 μ L/well. The plates were incubated in a CO₂ incubator for 24 h at 37°C, and then the reagent was added.

3. Knockout step 1

Either the KO1 or KO2 vector was introduced into the CHO cells, and after 24 h the cells were selected using hygromycin B (Invitrogen, Cat. No. 10687-010). Hygromycin B was dissolved in the SFMII(+) up to a concentration of 0.3 mg/mL and was added at 100 μ L/well.

4. Screening for homologous recombinants by PCR

(1) Preparation of PCR sample

Screening for homologous recombinants was carried out by PCR. The CHO cells used in screening were

cultured in 96-well plates. After the supernatant was removed, 50 μ L/well of buffer for cytolysis was added, and the cells were first heated at 55°C for 2 h and then 95°C for 15 min to inactivate protease K to prepare PCR template. The buffer for cytolysis consisted of 5 μ L of 10 X LA buffer II (Takara Bio Inc., LA Taq added), 2.5 μ L of 10% NP-40 (Roche, Cat. No. 1 332 473), 4 μ L of proteinase K (20 mg/mL, Takara Bio, Inc. Cat. No. 9033), and 38.5 μ L of distilled water (Nacalai Tesque Cat. No. 36421-35) per well.

(2) PCR conditions

PCR reaction mixture contained 1 μ L of the above PCR sample, 5 μ L of 10 X LA buffer II, 5 μ L of $MgCl_2$ (25 mM), 5 μ L of dNTP (2.5 mM), 2 μ L of primer (10 μ M each), 0.5 μ L of La Taq (5 IU/ μ L, Cat. No. RR002B) and 29.5 μ L of distilled water (50 μ L in total). For screening of cells containing the KO1 vector, TP-F4 and THygro-R1 were used as PCR primers, and for screening of cells containing the KO2 vector, TP-F4 and THygro-F1 were used as PCR primers.

PCR conditions for the cells containing the KO1 vector consisted of preheating at 95°C for 1 min, conducting 40 cycles of the amplification cycle of 30 sec at 95°C, 30 sec at 60°C, and 2 min at 60°C, and then reheating at 72°C for 7 min. PCR conditions for the cells containing the KO2 vector consisted of preheating at 95°C for 1 min, 40 cycles of the amplification cycle

of 30 sec at 95°C and 3 min at 70°C, and then reheating at 70°C for 7 min.

The primers used are listed below. In cell samples wherein homologous recombination occurred with the KO1 vector or the KO2 vector, DNA of approximately 1.6 kb or 2.0 kb will be amplified, respectively. The primer TP-F4 was designed for the 5' genome region of the fucose transporter outside of the vector, and THygro-F1 and THygro-R1 were designed for the Hygr gene inside of the vector.

Forward primer (KO1, KO2)

TP-F4 5'-GGA ATG CAG CTT CCT CAA GGG ACT CGC-3' (SEQ ID NO: 133)

Reverse primer (KO1)

THygro-R1 5'-TGC ATC AGG TCG GAG ACG CTG TCG AAC-3'
(SEQ ID NO: 134)

Reverse primer (KO2)

THygro-F1 5'-GCA CTC GTC CGA GGG CAA AGG AAT AGC-3'
(SEQ ID NO: 135)

5. PCR screening results

A total of 918 cells containing the KO1 vector were analyzed, and 1 cell was appeared to be a homologous recombinant (homologous recombination rate: approximately 0.1%). A total of 537 cells containing the KO2 vector were analyzed and 17 cells were appeared to be homologous recombinants (homologous recombination rate:

approximately 3.2%).

6. Southern blot analysis

Homologous recombination was further confirmed by Southern blot. A total of 10 µg of genomic DNA was prepared from the cultured cells by the standard method to be analyzed in Southern blotting. PCR was conducted using the two primers listed below to prepare a 387 bp probe corresponding to the region of base Nos. 2113 to 2500 of the nucleotide sequence shown in SEQ ID NO: 126, which was used in Southern blotting. The genomic DNA was cleaved by BglII.

Forward primer

Bgl-F: 5'-TGT GCT GGG AAT TGA ACC CAG GAC -3' (SEQ ID NO: 136)

Reverse primer

Bgl-R: 5'-CTA CTT GTC TGT GCT TTC TTC C -3' (SEQ ID NO: 137)

As a result of cleavage with BglII, the blot will show a band of approximately 30 kb from the fucose transporter chromosome, approximately 4.6 kb from the chromosome wherein homologous recombination with the K01 vector occurred, and approximately 5.0 kb from the chromosome wherein homologous recombination with the K02 vector occurred. The experiment comprised 1 cell from homologous recombination with the K01 vector and 7 cells from homologous recombination with the K02 vector. The

only cell obtained from the homologous recombination with the KO1 vector was first designated 5C1, but later analysis revealed that this cell consisted of multiple cell populations. Therefore the cells were cloned by limiting dilution before used in the experiment. One of the cells obtained with the KO2 vector was designated 6E2.

7. Knockout step 2

The three vectors were used to establish cell lines completely defective of fucose transporter gene from the cells wherein homologous recombination with the KO1 and KO2 vectors took place. The combinations of vectors and cells was as follows: Method 1 combined the KO2 vector and 5C1 cells (KO1), Method 2 combined the KO2 vector and 6E2 cells (KO2), and Method 3 combined the KO3 vector and 6E2 cells (KO2). Each vector was introduced into the appropriate cells, and after 24 h selection was started using hygromycin B and puromycin (Nacalai Tesque, Cat. No. 29455-12). The final concentration of hygromycin B was set to 1 mg/mL in Method 1, and 7 mg/mL in Method 2. In Method 3 hygromycin B was added at a final concentration of 0.15 mg/mL and puromycin at a final concentration of 8 µg/mL.

8. Screening for homologous recombinants by PCR

For screening of cells from Method 1, PCR was carried out to detect cells having homologous recombination with both KO1 and KO2 vectors. For

screening of cells from Method 2 the following PCR primers were designed: TPS-F1 was configured in the region from base Nos. 3924 to 3950 of SEQ ID NO: 126, and SHygro-R1 was configured in the region from base Nos. 4248 to 4274. These primers will amplify 350 bp of the fucose transporter gene region containing a deletion due to the KO2 vector. Therefore, in the PCR screening in Method 2, those cells providing no amplification product of 350bp are considered to be completely lacking the fucose transporter gene. The PCR conditions consisted of preheating for 1 min at 95°C, 35 cycles of the amplification cycle of 30 sec at 95°C and 1 min at 70°C, and reheating for 7 min at 70°C.

Forward primer

TPS-F1: 5'-CTC GAC TCG TCC CTA TTA GGC AAC AGC -3' (SEQ ID NO: 138)

Reverse primer

SHygro-R1: 5'-TCA GAG GCA GTG GAG CCT CCA GTC AGC -3' (SEQ ID NO: 139)

For Method 3, TP-F4 was used as the forward primer and RSGR-A was used as the reverse primer. The PCR conditions consisted of preheating for 1 min at 95°C, 35 cycles of the amplification cycle of 30 sec at 95°C, 30 sec at 60°C, and 2 min at 72°C, and reheating for 7 min at 72°C. In the sample of cells having homologous recombination with the KO3 vector, DNA of approximately

1.6 kb will be amplified. This PCR procedure will detect those cells having homologous recombination with the KO3 vector and also those still having homologous recombination with the KO2 vector.

9. PCR screening results

In Method 1, a total of 616 cells were analyzed, and 18 cells were appeared to be homologous recombinants (homologous recombination rate: 2.9%). In Method 2, a total of 524 cells were analyzed, and 2 cells were appeared to be homologous recombinants (homologous recombination rate: 0.4%). In addition, in Method 3, a total of 382 cells were analyzed, and 7 cells were appeared to be homologous recombinants (homologous recombination rate: 1.8%).

10. Southern blot analysis

Southern blotting was carried out according to the method described above. Among the cells analyzed, 1 cell completely lacking the fucose transporter gene was found. In knockout step 1, the analysis results of PCR and Southern blotting were consistent, but not in knockout step 2. Possible causes are as follows: 1. In Method 1 cells having homologous recombination independently with either KO1 or KO2 were mixed together; 2. The fucose transporter gene is present not as one pair (2 genes) but as multiple pairs (or not less than 3 genes); and 3. During the culture of the cell lines established in the

knockout step 1, the copy number of the fucose transporter gene remaining in the subcultured cells increased.

11. Analysis of fucose expression

Fucose expression was analyzed by PCR in 26 cells found to be homologous recombinants. A total of 1×10^6 cells were stained on ice for 1 h with 100 μ L of PBS containing 5 μ g/mL of Lens culinaris Agglutinin, FITC conjugate (Vector Laboratories, Cat. No. FL-1041) 2.5% FBS, and 0.02% sodium azide (hereinafter designated as FACS solution). Then the cells were rinsed 3 times with FACS solution and analyzed with FACSCalibur (Becton Dickinson). The results clearly showed that fucose expression is decreased only in the cells found to be completely lacking the fucose transporter gene in the Southern blot analysis.

The above results have revealed the following:

From the fact that only one cell out of 616 cells screened has complete deletion of the fucose transporter gene, the frequency of homologous recombination was very low at approximately 0.16%. As noted above, there are several possible reasons why the results of PCR and Southern blot analysis in knockout step 2 was not consistent. However, the cell lines obtained in Method 3 may not comprise a mixture of cells having homologous recombination independently with the KO2 and KO3 vectors,

because selection was made using two types of drugs. In addition, it is unlikely that all of the other cell lines having homologous recombination by PCR comprise multiple cell populations. As noted above, if there are 3 or more fucose transporter genes present, targeting of the gene in cells would be particularly difficult. Homologous recombinants could be obtained only by using a vector such as the KO1 vector where Hyger is hardly expressed and by screening a large number of cells.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An anti-glypican 3 antibody composition comprising an antibody having a heavy chain variable region comprising CDR1 having the amino acid sequence set forth in SEQ ID NO: 25, CDR2 having the amino acid sequence set forth in SEQ ID NO: 26, and CDR3 having the amino acid sequence set forth in SEQ ID NO: 27, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting N-acetylglucosamid (GlcNAc).
2. An anti-glypican 3 antibody composition comprising an antibody having a light chain variable region comprising CDR1 having the amino acid sequence set forth in SEQ ID NO: 45, CDR2 having the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 having the amino acid sequence set forth in SEQ ID NO: 60, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting N-acetylglucosamid (GlcNAc).
3. An anti-glypican 3 antibody composition wherein the antibody of claim 1 further has a light chain variable region comprising CDR1 having the amino acid sequence set forth in SEQ ID NO: 45, CDR2 having the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 having the amino acid sequence set forth in SEQ ID NO: 60.
4. An anti-glypican 3 antibody composition comprising an antibody having a heavy chain variable region of any one of (1)-(7):
- (1) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 81;
 - (2) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 82;
 - (3) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 83;
 - (4) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84;
 - (5) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85;

(6) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86; or

(7) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting GlcNAc.

5. An anti-glypican 3 antibody composition comprising an antibody having a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92, wherein the sugar chain component thereof has been modified so as to have an increased ratio of fucose-deficient antibodies or an increased ratio of antibodies having an attached bisecting GlcNAc.

6. The anti-glypican 3 antibody composition according to claim 4, further comprising an antibody having a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92.

7. An anti-glypican 3 antibody composition having an activity equivalent to the activity of the antibody composition of any one of claims 1-6, wherein one or more amino acid residues are substituted, deleted or added and/or inserted from the amino acid sequences set forth in any one of Claims 1-6.

8. The anti-glypican 3 antibody composition according to claim 7, comprising an antibody having a light chain variable region comprising CDRs 1, 2 and 3 of any one of (1)-(15):

(1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 90, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;

(2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 91, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;

(3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 92, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;

(4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 93, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3

- comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 94, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- 5 (6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 95, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 96, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3
- 10 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 97, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 98, CDR2
- 15 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 99, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- 20 (11) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 100, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (12) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 101, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3
- 25 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (13) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 102, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (14) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 103,
- 30 CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60;
- (15) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 104, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 46, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60.

35

9. The anti-glypican 3 antibody composition according to claim 7, comprising a
1226505_1.doc

light chain variable region selected from (1)-(15):

- (1) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 107;
 - (2) a light chain variable region comprising the amino acid sequence set forth in
5 SEQ ID NO: 108;
 - (3) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 109;
 - (4) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 110;
 - 10 (5) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 111;
 - (6) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 112;
 - (7) a light chain variable region comprising the amino acid sequence set forth in
15 SEQ ID NO: 113;
 - (8) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 114;
 - (9) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 115;
 - 20 (10) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 116;
 - (11) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 117;
 - (12) a light chain variable region comprising the amino acid sequence set forth in
25 SEQ ID NO: 118;
 - (13) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 119;
 - (14) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 120; and
 - 30 (15) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 121.
10. The anti-glypican 3 antibody composition according to any of claims 1 to 9,
wherein the sugar chain component thereof has been modified such that the
35 composition comprises more than 20% antibodies lacking fucose.

11. The anti-glypican 3 antibody composition according to any of claims 1 to 9, wherein the sugar chain component thereof has been modified such that the composition comprises more than 50% antibodies lacking fucose.

5 12. The anti-glypican 3 antibody composition according to any of claims 1 to 9, wherein the sugar chain component thereof has been modified such that the composition comprises more than 90% antibodies lacking fucose.

10 13. A process for producing an anti-glypican 3 antibody using a cell, wherein a gene encoding the anti-glypican 3 antibody as defined in any of claims 1 to 9 has been introduced into said cell having reduced capability of adding fucose to sugar chains.

15 14. The process for producing an anti-glypican 3 antibody according to claim 13, wherein the cell having reduced capability of adding fucose to sugar chains is a cell lacking a fucose transporter.

20 15. A process for producing an anti-glypican 3 antibody, comprising the steps of:
(a) introducing a gene encoding the anti-glypican 3 antibody as defined in any one of claims 1 to 9 into a cell having reduced capability of adding fucose to sugar chains; and
(b) culturing the cell.

25 16. An anticancer drug having as its active ingredient the antibody composition according to any of claims 1 to 12.

17. Use of a composition as claimed in any one of claims 1 to 12 as a medicament.

30 18. Use of a composition as claimed in any one of claims 1 to 12 as an anticancer agent.

19. A method for treating cancer comprising administering a composition as claimed in any one of claims 1 to 12.

35 20. Use of a composition as claimed in any one of claims 1 to 12 in the treatment of cancer.

21. Use of a composition as claimed in any one of claims 1 to 12 for the manufacture of a medicament for the treatment of cancer.
- 5 22. A modified anti-glypican 3 antibody composition, process, method or use substantially as hereinbefore described with reference to the accompanying examples.




FIG 1




FIG 2




FIG 3




FIG 4




FIG 5





FIG 6




FIG 7

SEQUENCE LISTING

<110> Chugai Seiyaku Kabushiki Kaisha
 <120> Anti-Glypican 3 Antibodies with Modified Sugars
 <130> PCG-9012W0
 <150> JP 2004-311356
 <151> 2004-10-26
 <160> 139
 <170> PatentIn version 3.1
 <210> 1
 <211> 1743
 <212> DNA
 <213> homo sapiens
 <400> 1

atggccggga ccgtgcgcac cgcgtgcttg gtgggtggcga tgctgctcag cttggacttc	60
ccgggacagg cgcagccccc gccgccgccg ccggacgcca cctgtcacca agtccgctcc	120
ttcttccaga gactgcagcc cggactcaag tgggtgccag aaactcccggt gccaggatca	180
gatttgcaag tatgtctccc taagggccca acatgtctgt caagaaagat ggaagaaaaa	240
taccaactaa cagcagcatt gaacatggaa cagctgcttc agtctgcaag tatggagctc	300
aagttcttaa ttattcagaa tgctgcgggt ttccaagagg cctttgaaat tgttgttcgc	360
catgccaaaga actacaccaa tgccatgttc aagaacaact acccaagcct gactccacaa	420
gcittttgagt ttgtgggtga atttttcaca gatgtgtctc tctacatctt gggttctgac	480
atcaatgtag atgacatggt caatgaattg ttgacagcc tgtttccagt catctatacc	540
cagctaataa acccaggcct gcctgattca gccttggaca tcaatgagtg cctccgagga	600
gcaagacgtg acctgaaagt atttggaat ttccccaagc ttattatgac ccaggtttcc	660
aagtcaactg aagtcactag gatcttcctt caggctctga atcttggaaat tgaagtgatc	720
aacacaactg atcacctgaa gttcagtaag gactgtggcc gaatgctcac cagaatgttg	780
tactgtcttt actgccaggg actgatgatg gttaaaccct gtggcggtta ctgcaatgtg	840
gtcatgcaag gctgtatggc aggtgtgggt gagattgaca agtactggag agaatacatt	900
ctgtcccttg aagaacttgt gaatggcatg tacagaatct atgacatgga gaacgtactg	960
cttggctctt ttccaacaat ccatgattct atccagtatg tccagaagaa tgcaggaaag	1020
ctgaccacca ctattggcaa gttatgtgcc cattctcaac aacgccaata tagatctgct	1080
tattatcctg aagatctctt tattgacaag aaagtattaa aagttgctca tgtagaacat	1140
gaagaaacct tatccagccg aagaaggga ctaattcaga agttgaagtc ttcatcagc	1200
ttctatagtg ctttgcctgg ctacatctgc agccatagcc ctgtggcgga aaacgacacc	1260
ctttgctgga atggacaaga actcgtggag agatacagcc aaaaggcagc aaggaatgga	1320
atgaaaaacc agttcaatct ccatgagctg aaaatgaagg gccctgagcc agtggtcagt	1380

caaattattg acaaactgaa gcacattaac cagctcctga gaaccatgtc tatgccc aaa 1440
 ggtagagttc tggataaaaa cctggatgag gaagggtttg aaagtggaga ctgcggtgat 1500
 gatgaagatg agtgattgg aggctctggt gatggaatga taaaagtga gaatcagctc 1560
 cgcttccttg cagaactggc ctatgatctg gatgtggatg atgcgcctgg aaacagtcag 1620
 caggcaactc cgaaggacaa cgagataagc accittcaca acctcgggaa cgttcattcc 1680
 ccgctgaagc ttctcaccag catggccatc tcggtgggtg gcttcttctt cctggtgac 1740
 tga 1743

<210> 2

<211> 580

<212> PRT

<213> homo sapiens

<400> 2

Met	Ala	Gly	Thr	Val	Arg	Thr	Ala	Cys	Leu	Val	Val	Ala	Met	Leu	Leu
1				5					10					15	
Ser	Leu	Asp	Phe	Pro	Gly	Gln	Ala	Gln	Pro	Pro	Pro	Pro	Pro	Pro	Asp
			20					25					30		
Ala	Thr	Cys	His	Gln	Val	Arg	Ser	Phe	Phe	Gln	Arg	Leu	Gln	Pro	Gly
		35					40					45			
Leu	Lys	Trp	Val	Pro	Glu	Thr	Pro	Val	Pro	Gly	Ser	Asp	Leu	Gln	Val
	50					55					60				
Cys	Leu	Pro	Lys	Gly	Pro	Thr	Cys	Cys	Ser	Arg	Lys	Met	Glu	Glu	Lys
65				70						75				80	
Tyr	Gln	Leu	Thr	Ala	Arg	Leu	Asn	Met	Glu	Gln	Leu	Leu	Gln	Ser	Ala
			85						90					95	
Ser	Met	Glu	Leu	Lys	Phe	Leu	Ile	Ile	Gln	Asn	Ala	Ala	Val	Phe	Gln
			100						105					110	
Glu	Ala	Phe	Glu	Ile	Val	Val	Arg	His	Ala	Lys	Asn	Tyr	Thr	Asn	Ala
		115						120					125		
Met	Phe	Lys	Asn	Asn	Tyr	Pro	Ser	Leu	Thr	Pro	Gln	Ala	Phe	Glu	Phe
	130						135					140			
Val	Gly	Glu	Phe	Phe	Thr	Asp	Val	Ser	Leu	Tyr	Ile	Leu	Gly	Ser	Asp
145				150						155				160	
Ile	Asn	Val	Asp	Asp	Met	Val	Asn	Glu	Leu	Phe	Asp	Ser	Leu	Phe	Pro
			165						170					175	
Val	Ile	Tyr	Thr	Gln	Leu	Met	Asn	Pro	Gly	Leu	Pro	Asp	Ser	Ala	Leu
			180						185					190	

Asp	Ile	Asn	Glu	Cys	Leu	Arg	Gly	Ala	Arg	Arg	Asp	Leu	Lys	Val	Phe
195				200				205							
Gly	Asn	Phe	Pro	Lys	Leu	Ile	Met	Thr	Gln	Val	Ser	Lys	Ser	Leu	Gln
210				215				220							
Val	Thr	Arg	Ile	Phe	Leu	Gln	Ala	Leu	Asn	Leu	Gly	Ile	Glu	Val	Ile
225				230				235				240			
Asn	Thr	Thr	Asp	His	Leu	Lys	Phe	Ser	Lys	Asp	Cys	Gly	Arg	Met	Leu
245				250				255							
Thr	Arg	Met	Trp	Tyr	Cys	Ser	Tyr	Cys	Gln	Gly	Leu	Met	Met	Val	Lys
260				265				270							
Pro	Cys	Gly	Gly	Tyr	Cys	Asn	Val	Val	Met	Gln	Gly	Cys	Met	Ala	Gly
275				280				285							
Val	Val	Glu	Ile	Asp	Lys	Tyr	Trp	Arg	Glu	Tyr	Ile	Leu	Ser	Leu	Glu
290				295				300							
Glu	Leu	Val	Asn	Gly	Met	Tyr	Arg	Ile	Tyr	Asp	Met	Glu	Asn	Val	Leu
305				310				315				320			
Leu	Gly	Leu	Phe	Ser	Thr	Ile	His	Asp	Ser	Ile	Gln	Tyr	Val	Gln	Lys
325				330				335							
Asn	Ala	Gly	Lys	Leu	Thr	Thr	Thr	Ile	Gly	Lys	Leu	Cys	Ala	His	Ser
340				345				350							
Gln	Gln	Arg	Gln	Tyr	Arg	Ser	Ala	Tyr	Tyr	Pro	Glu	Asp	Leu	Phe	Ile
355				360				365							
Asp	Lys	Lys	Val	Leu	Lys	Val	Ala	His	Val	Glu	His	Glu	Glu	Thr	Leu
370				375				380							
Ser	Ser	Arg	Arg	Arg	Glu	Leu	Ile	Gln	Lys	Leu	Lys	Ser	Phe	Ile	Ser
385				390				395				400			
Phe	Tyr	Ser	Ala	Leu	Pro	Gly	Tyr	Ile	Cys	Ser	His	Ser	Pro	Val	Ala
405				410				415							
Glu	Asn	Asp	Thr	Leu	Cys	Trp	Asn	Gly	Gln	Glu	Leu	Val	Glu	Arg	Tyr
420				425				430							
Ser	Gln	Lys	Ala	Ala	Arg	Asn	Gly	Met	Lys	Asn	Gln	Phe	Asn	Leu	His
435				440				445							
Glu	Leu	Lys	Met	Lys	Gly	Pro	Glu	Pro	Val	Val	Ser	Gln	Ile	Ile	Asp
450				455				460							
Lys	Leu	Lys	His	Ile	Asn	Gln	Leu	Leu	Arg	Thr	Met	Ser	Met	Pro	Lys
465				470				475				480			

Gly Arg Val Leu Asp Lys Asn Leu Asp Glu Glu Gly Phe Glu Ser Gly
 485 490 495
 Asp Cys Gly Asp Asp Glu Asp Glu Cys Ile Gly Gly Ser Gly Asp Gly
 500 505 510
 Met Ile Lys Val Lys Asn Gln Leu Arg Phe Leu Ala Glu Leu Ala Tyr
 515 520 525
 Asp Leu Asp Val Asp Asp Ala Pro Gly Asn Ser Gln Gln Ala Thr Pro
 530 535 540
 Lys Asp Asn Glu Ile Ser Thr Phe His Asn Leu Gly Asn Val His Ser
 545 550 555 560
 Pro Leu Lys Leu Leu Thr Ser Met Ala Ile Ser Val Val Cys Phe Phe
 565 570 575
 Phe Leu Val His
 580

<210> 3

<211> 115

<212> PRT

<213> Mus musculus

<400> 3

Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly Ala
 1 5 10 15
 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Glu Met His Trp Val Lys Gln Thr Pro Val His Gly Leu Lys Trp Ile
 35 40 45
 Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

<210> 4

<211> 112

<212> PRT

<213> *Mus musculus*

<400> 4

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly

1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser

20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser

35 40 45

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro

50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile

65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Asn

85 90 95

Thr His Val Pro Pro Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys

100 105 110

<210> 5

<211> 5

<212> PRT

<213> *Mus musculus*

<400> 5

Asn Tyr Ala Met Ser

1 5

<210> 6

<211> 17

<212> PRT

<213> *Mus musculus*

<400> 6

Ala Ile Asn Asn Asn Gly Asp Asp Thr Tyr Tyr Leu Asp Thr Val Lys

1 5 10 15

Asp

<210> 7

<211> 5

<212> PRT

<213> Mus musculus

<400> 7

Gln Gly Gly Ala Tyr

1 5

<210> 8

<211> 7

<212> PRT

<213> Mus musculus

<400> 8

Thr Tyr Gly Met Gly Val Gly

1 5

<210> 9

<211> 16

<212> PRT

<213> Mus musculus

<400> 9

Asn Ile Trp Trp Tyr Asp Ala Lys Tyr Tyr Asn Ser Asp Leu Lys Ser

1 5 10 15

<210> 10

<211> 8

<212> PRT

<213> Mus musculus

<400> 10

Met Gly Leu Ala Trp Phe Ala Tyr

1 5

<210> 11

<211> 7

<212> PRT

<213> Mus musculus

<400> 11

Ile Tyr Gly Met Gly Val Gly

1 5

<210> 12

<211> 16

<212> PRT

<213> Mus musculus

<400> 12

Asn Ile Trp Trp Asn Asp Asp Lys Tyr Tyr Asn Ser Ala Leu Lys Ser

1 5 10 15

<210> 13

<211> 8

<212> PRT

<213> Mus musculus

<400> 13

Ile Gly Tyr Phe Tyr Phe Asp Tyr

1 5

<210> 14

<211> 5

<212> PRT

<213> Mus musculus

<400> 14

Gly Tyr Trp Met His

1 5

<210> 15

<211> 17

<212> PRT

<213> Mus musculus

<400> 15

Ala Ile Tyr Pro Gly Asn Ser Asp Thr Asn Tyr Asn Gln Lys Phe Lys

1 5 10 15

Gly

<210> 16

<211> 10

<212> PRT

<213> Mus musculus

<400> 16

Ser Gly Asp Leu Thr Gly Gly Leu Ala Tyr

1 5 10

<210> 17

<211> 5

<212> PRT

<213> Mus musculus

<400> 17

Ser Tyr Ala Met Ser

1 5

<210> 18

<211> 17

<212> PRT

<213> Mus musculus

<400> 18

Ala Ile Asn Ser Asn Gly Gly Thr Thr Tyr Tyr Pro Asp Thr Met Lys

1 5 10 15

Asp

<210> 19

<211> 13

<212> PRT

<213> Mus musculus

<400> 19

His Asn Gly Gly Tyr Glu Asn Tyr Gly Trp Phe Ala Tyr

1 5 10

<210> 20

<211> 5

<212> PRT

<213> Mus musculus

<400> 20

Ser Tyr Trp Met His

1 5

<210> 21

<211> 17

<212> PRT

<213> Mus musculus

<400> 21

Glu Ile Asp Pro Ser Asp Ser Tyr Thr Tyr Tyr Asn Gln Lys Phe Arg

1 5 10 15

Gly

<210> 22
 <211> 15
 <212> PRT
 <213> Mus musculus
 <400> 22

Ser Asn Leu Gly Asp Gly His Tyr Arg Phe Pro Ala Phe Pro Tyr
 1 5 10 15

<210> 23
 <211> 17
 <212> PRT
 <213> Mus musculus
 <400> 23

Thr Ile Asp Pro Ser Asp Ser Glu Thr His Tyr Asn Leu Gln Phe Lys
 1 5 10 15

Asp

<210> 24
 <211> 15
 <212> PRT
 <213> Mus musculus
 <400> 24

Gly Ala Phe Tyr Ser Ser Tyr Ser Tyr Trp Ala Trp Phe Ala Tyr
 1 5 10 15

<210> 25
 <211> 5
 <212> PRT
 <213> Mus musculus
 <400> 25

Asp Tyr Glu Met His
 1 5

<210> 26
 <211> 17
 <212> PRT
 <213> Mus musculus
 <400> 26

Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe Lys

1 5 10 15

Gly

<210> 27

<211> 6

<212> PRT

<213> Mus musculus

<400> 27

Phe Tyr Ser Tyr Thr Tyr

1 5

<210> 28

<211> 5

<212> PRT

<213> Mus musculus

<400> 28

Ile Asn Ala Met Asn

1 5

<210> 29

<211> 19

<212> PRT

<213> Mus musculus

<400> 29

Arg Ile Arg Ser Glu Ser Asn Asn Tyr Ala Thr Tyr Tyr Gly Asp Ser

1 5 10 15

Val Lys Asp

<210> 30

<211> 8

<212> PRT

<213> Mus musculus

<400> 30

Glu Val Thr Thr Ser Phe Ala Tyr

1 5

<210> 31

<211> 5

<212> PRT

<213> Mus musculus

<400> 31

Ala Ser Ala Met Asn

1 5

<210> 32

<211> 19

<212> PRT

<213> Mus musculus

<400> 32

Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Ile Tyr Tyr Ala Asp Ser

1 5 10 15

Val Lys Asp

<210> 33

<211> 12

<212> PRT

<213> Mus musculus

<400> 33

Asp Pro Gly Tyr Tyr Gly Asn Pro Trp Phe Ala Tyr

1 5 10

<210> 34

<211> 5

<212> PRT

<213> Mus musculus

<400> 34

Asp Tyr Ser Met His

1 5

<210> 35

<211> 17

<212> PRT

<213> Mus musculus

<400> 35

Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe Lys

1 5 10 15

Gly

<210> 36

<211> 2

<212> PRT

<213> Mus musculus

<400> 36

Leu Tyr

1

<210> 37

<211> 16

<212> PRT

<213> Mus musculus

<400> 37

Asn Ile Trp Trp His Asp Asp Lys Tyr Tyr Asn Ser Ala Leu Lys Ser

1

5

10

15

<210> 38

<211> 14

<212> PRT

<213> Mus musculus

<400> 38

Ile Ala Pro Arg Tyr Asn Lys Tyr Glu Gly Phe Phe Ala Phe

1

5

10

<210> 39

<211> 16

<212> PRT

<213> Mus musculus

<400> 39

Lys Ser Ser Gln Ser Leu Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn

1

5

10

15

<210> 40

<211> 7

<212> PRT

<213> Mus musculus

<400> 40

Leu Val Ser Lys Leu Asp Ser

1

5

<210> 41

<211> 9

<212> PRT

<213> Mus musculus

<400> 41

Trp Gln Gly Thr His Phe Pro Leu Thr

1 5

<210> 42

<211> 11

<212> PRT

<213> Mus musculus

<400> 42

Lys Ala Ser Gln Asp Ile Asn Asn Tyr Leu Ser

1 5 10

<210> 43

<211> 7

<212> PRT

<213> Mus musculus

<400> 43

Arg Ala Asn Arg Leu Val Asp

1 5

<210> 44

<211> 10

<212> PRT

<213> Mus musculus

<400> 44

Leu Gln Cys Asp Glu Phe Pro Pro Trp Thr

1 5 10

<210> 45

<211> 16

<212> PRT

<213> Mus musculus

<400> 45

Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His

1 5 10 15

<210> 46

<211> 7
 <212> PRT
 <213> Mus musculus
 <400> 46
 Lys Val Ser Asn Arg Phe Ser
 1 5
 <210> 47
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 47
 Ser Gln Ser Thr His Val Pro Trp Thr
 1 5
 <210> 48
 <211> 16
 <212> PRT
 <213> Mus musculus
 <400> 48
 Arg Ser Ser Lys Ser Leu Leu His Ser Asn Gly Ile Thr Tyr Leu Tyr
 1 5 10 15
 <210> 49
 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 49
 Gln Met Ser Asn Leu Ala Ser
 1 5
 <210> 50
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 50
 Ala Gln Asn Leu Glu Leu Pro Tyr Thr
 1 5
 <210> 51
 <211> 11

<212> PRT

<213> Mus musculus

<400> 51

Lys Ala Ser Gln Asp Ile Asn Lys Asn Ile Ile

1 5 10

<210> 52

<211> 7

<212> PRT

<213> Mus musculus

<400> 52

Tyr Thr Ser Thr Leu Gln Pro

1 5

<210> 53

<211> 9

<212> PRT

<213> Mus musculus

<400> 53

Leu Gln Tyr Asp Asn Leu Pro Arg Thr

1 5

<210> 54

<211> 11

<212> PRT

<213> Mus musculus

<400> 54

Arg Ala Ser His Ser Ile Ser Asn Phe Leu His

1 5 10

<210> 55

<211> 7

<212> PRT

<213> Mus musculus

<400> 55

Tyr Ala Ser Gln Ser Ile Ser

1 5

<210> 56

<211> 9

<212> PRT

<213> Mus musculus

<400> 56

Gln Gln Ser Asn Ile Trp Ser Leu Thr

1 5

<210> 57

<211> 15

<212> PRT

<213> Mus musculus

<400> 57

Arg Ala Ser Glu Ser Val Glu Tyr Tyr Gly Thr Ser Leu Met Gln

1 5 10 15

<210> 58

<211> 7

<212> PRT

<213> Mus musculus

<400> 58

Gly Ala Ser Asn Val Glu Ser

1 5

<210> 59

<211> 9

<212> PRT

<213> Mus musculus

<400> 59

Gln Gln Ser Arg Lys Val Pro Tyr Thr

1 5

<210> 60

<211> 9

<212> PRT

<213> Mus musculus

<400> 60

Ser Gln Asn Thr His Val Pro Pro Thr

1 5

<210> 61

<211> 16

<212> PRT

<213> Mus musculus

<400> 61

Lys Ser Ser Lys Ser Leu Leu His Ser Asn Gly Asn Thr Tyr Leu Asn

1 5 10 15

<210> 62

<211> 7

<212> PRT

<213> Mus musculus

<400> 62

Trp Met Ser Asn Leu Ala Ser

1 5

<210> 63

<211> 9

<212> PRT

<213> Mus musculus

<400> 63

Met Gln His Ile Glu Tyr Pro Phe Thr

1 5

<210> 64

<211> 16

<212> PRT

<213> Mus musculus

<400> 64

Arg Ser Ser Lys Ser Leu Leu His Ser Tyr Asp Ile Thr Tyr Leu Tyr

1 5 10 15

<210> 65

<211> 9

<212> PRT

<213> Mus musculus

<400> 65

Ala Gln Asn Leu Glu Leu Pro Pro Thr

1 5

<210> 66

<211> 10

<212> PRT

<213> Mus musculus

<400> 66

Ser Ala Ser Ser Ser Val Ser Tyr Met Tyr

1 5 10

<210> 67

<211> 7

<212> PRT

<213> Mus musculus

<400> 67

Asp Thr Ser Asn Leu Ala Ser

1 5

<210> 68

<211> 9

<212> PRT

<213> Mus musculus

<400> 68

Gln Gln Trp Ser Ser Tyr Pro Leu Thr

1 5

<210> 69

<211> 16

<212> PRT

<213> Mus musculus

<400> 69

Lys Ser Ser Gln Ser Leu Leu His Ser Asp Gly Lys Thr Phe Leu Asn

1 5 10 15

<210> 70

<211> 7

<212> PRT

<213> Mus musculus

<400> 70

Leu Val Ser Arg Leu Asp Ser

1 5

<210> 71

<211> 9

<212> PRT

<213> Mus musculus

<400> 71

Cys Gln Gly Thr His Phe Pro Arg Thr

1 5
 <210> 72
 <211> 16
 <212> PRT
 <213> Mus musculus
 <400> 72
 Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu
 1 5 10 15
 <210> 73
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 73
 Phe Gln Gly Ser His Val Pro Trp Thr
 1 5
 <210> 74
 <211> 345
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 74
 cagggtgcagc tgggtggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 tcctgcaagg ctctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagct ctgatccta aaactgggta tactgcctac 180
 agtcagaagt tcaagggcag agtcacgatt accgcgagc aatccacgag cacagcctac 240
 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagattctac 300
 tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345
 <210> 75
 <211> 345
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 75
 cagggtgcagc tgggtggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60

tcctgcaagg ctcttgata caccttcacc gactatgaaa tgcactgggt gcgacaggcc	120
cctggacaag ggcttgagt gatgggagct ctgatccta aaactgggtga tactgcctac	180
agtcagaagt tcaagggcag agtcacgctg accgcggacg aatccacgag cacagcctac	240
atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac	300
tcctatactt actggggcca gggaaccctg gtcaccgtct cctca	345

<210> 76

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 76

caggtgcagc tggaggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc	60
tcctgcaagg ctcttgata caccttcacc gactatgaaa tgcactgggt gcgacaggcc	120
cctggacaag ggcttgagt gatgggagct ctgatccta aaactgggtga tactgcctac	180
agtcagaagt tcaagggcag agtcacgctg accgcggaca aatccacgag cacagcctac	240
atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac	300
tcctatactt actggggcca gggaaccctg gtcaccgtct cctca	345

<210> 77

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 77

caggtgcagc tggaggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc	60
tcctgcaagg ctcttgata caccttcacc gactatgaaa tgcactgggt gcgacaggcc	120
cctggacaag ggcttgagt gatgggagct ctgatccta aaactgggtga tactgcctac	180
agtcagaagt tcaagggcag agtcacgctg accgcggaca aatccacgag cacagcctac	240
atggagctga gcagcctgac atctgaggac acggccgtgt attactgtac aagattctac	300
tcctatactt actggggcca gggaaccctg gtcaccgtct cctca	345

<210> 78

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 78

caggtgcagc tggcgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc	60
tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc	120
cctggacaag ggcttgagtg gatgggagct cttgatccta aaactgggtga tactgcctac	180
agtcagaagt tcaagggcag agtcacgctg accgcggacg aatccacgag cacagcctac	240
atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac	300
tcctatactt actggggcca gggaaccctg gtcaccgtct cctca	345

<210> 79

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 79

caggtgcagc tggcgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc	60
tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc	120
cctggacaag ggcttgagtg gatgggagct cttgatccta aaactgggtga tactgcctac	180
agtcagaagt tcaagggcag agtcacgctg accgcggaca aatccacgag cacagcctac	240
atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac	300
tcctatactt actggggcca gggaaccctg gtcaccgtct cctca	345

<210> 80

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 80

caggtgcagc tggcgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc	60
tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc	120
cctggacaag ggcttgagtg gatgggagct cttgatccta aaactgggtga tactgcctac	180
agtcagaagt tcaagggcag agtcacgctg accgcggaca aatccacgag cacagcctac	240
atggagctga gcagcctgac atctgaggac acggccgtgt attactgtac aagattctac	300
tcctatactt actggggcca gggaaccctg gtcaccgtct cctca	345

<210> 81

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 81

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30
Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
50 55 60
Lys Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110
Val Ser Ser
115

<210> 82

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 82

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30
Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Leu Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
Val Ser Ser

115

<210> 83

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 83

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110

Val Ser Ser

115

<210> 84

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 84

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30
Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
50 55 60
Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110
Val Ser Ser
115

<210> 85

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 85

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30
Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
50 55 60
Lys Gly Arg Val Thr Leu Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys

	85	90	95
Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr			
100	105	110	

Val Ser Ser

115

<210> 86

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 86

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val Ser Ser

115

<210> 87

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 87

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala

1	5	10	15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr			
20	25	30	
Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met			
35	40	45	
Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe			
50	55	60	
Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr			
65	70	75	80
Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys			
85	90	95	
Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr			
100	105	110	
Val Ser Ser			

115

<210> 88

<211> 336

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody L chain

<400> 88

gatgttgtga tgactcagtc tccactctcc ctgcccgta cccctggaga gccggcctcc	60
atctcctgca gatctagta gagccttgta cacagtaatg gaaacaccta tttacattgg	120
tacctgcaga agccagggca gtctccacag ctctgatct ataaagtctt caaccgattt	180
tctggggctc ctgacagggt cagtggcagt ggatcaggca cagattttac actgaaaatc	240
agcagagtgg aggctgagga tgttgggggt tattactgct ctcaaaatac acatgttcct	300
cctacgtttg gccaggggac caagctggag atcaaa	336

<210> 89

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody L chain

<400> 89

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1	5	10	15												
Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
	20		25		30										
Asn	Gly	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
	35		40		45										
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
	50		55		60										
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65			70		75									80	
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
		85			90									95	
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys
	100		105		110										

<210> 90

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 90

Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser	Asn	Ala	Asn	Thr	Tyr	Leu	His
1		5		10		15									

<210> 91

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 91

Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser	Asn	Asp	Asn	Thr	Tyr	Leu	His
1		5		10		15									

<210> 92

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 92

Arg Ser Ser Gln Ser Leu Val His Ser Asn Glu Asn Thr Tyr Leu His

1 5 10 15

<210> 93

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 93

Arg Ser Ser Gln Ser Leu Val His Ser Asn Phe Asn Thr Tyr Leu His

1 5 10 15

<210> 94

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 94

Arg Ser Ser Gln Ser Leu Val His Ser Asn His Asn Thr Tyr Leu His

1 5 10 15

<210> 95

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 95

Arg Ser Ser Gln Ser Leu Val His Ser Asn Asn Asn Thr Tyr Leu His

1 5 10 15

<210> 96

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 96

Arg Ser Ser Gln Ser Leu Val His Ser Asn Thr Asn Thr Tyr Leu His

1 5 10 15

<210> 97

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 97

Arg Ser Ser Gln Ser Leu Val His Ser Asn Gln Asn Thr Tyr Leu His

1 5 10 15

<210> 98

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 98

Arg Ser Ser Gln Ser Leu Val His Ser Asn Ile Asn Thr Tyr Leu His

1 5 10 15

<210> 99

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 99

Arg Ser Ser Gln Ser Leu Val His Ser Asn Lys Asn Thr Tyr Leu His

1 5 10 15

<210> 100

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 100

Arg Ser Ser Gln Ser Leu Val His Ser Asn Leu Asn Thr Tyr Leu His

1 5 10 15

<210> 101

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 101

Arg Ser Ser Gln Ser Leu Val His Ser Asn Ser Asn Thr Tyr Leu His

1 5 10 15

<210> 102

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 102

Arg Ser Ser Gln Ser Leu Val His Ser Asn Trp Asn Thr Tyr Leu His

1 5 10 15

<210> 103

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 103

Arg Ser Ser Gln Ser Leu Val His Ser Asn Tyr Asn Thr Tyr Leu His

1 5 10 15

<210> 104

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 104

Arg Ser Ser Gln Ser Leu Val His Ser Asn Arg Asn Thr Tyr Leu His

1 5 10 15

<210> 105

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 105

Arg Ser Ser Gln Ser Leu Val His Ser Asn Val Asn Thr Tyr Leu His

1 5 10 15

<210> 106

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 106

Arg Ser Ser Gln Ser Leu Val His Ser Asn Pro Asn Thr Tyr Leu His

1 5 10 15

<210> 107

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 107

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser

20 25 30

Asn Ala Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser

35 40 45

Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro

50	55	60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile		
65	70	75
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn		
85	90	95
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
100	105	110

<210> 108

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 108

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly		
1	5	10
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser		
20	25	30
Asn Asp Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser		
35	40	45
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro		
50	55	60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile		
65	70	75
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn		
85	90	95
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
100	105	110

<210> 109

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 109

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1	5	10	15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser			
20	25	30	
Asn Glu Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser			
35	40	45	
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro			
50	55	60	
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile			
65	70	75	80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn			
85	90	95	
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys			
100	105	110	

<210> 110

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 110

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly			
1	5	10	15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser			
20	25	30	
Asn Phe Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser			
35	40	45	
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro			
50	55	60	
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile			
65	70	75	80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn			
85	90	95	
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys			
100	105	110	

<210> 111

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 111

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
Asn His Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 112

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 112

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
Asn Asn Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
85 90 95

Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 113

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 113

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30

Asn Thr Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
85 90 95

Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 114

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 114

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30

Asn	Gln	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
		35					40					45			
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
		50					55					60			
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65					70					75					80
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
				85						90					95
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys
			100						105						110

<210> 115

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 115

Asp	Val	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Thr	Pro	Gly
1				5						10				15	
Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
				20						25				30	
Asn	Ile	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
				35						40				45	
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
				50						55				60	
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65					70					75					80
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
					85					90					95
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys
					100					105					110

<210> 116

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 116

Asp	Val	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Thr	Pro	Gly
1				5					10					15	
Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
			20					25					30		
Asn	Lys	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
		35					40					45			
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
	50					55						60			
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65				70					75					80	
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
			85					90						95	
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys
			100					105						110	

<210> 117

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 117

Asp	Val	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Thr	Pro	Gly
1				5					10					15	
Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
			20					25					30		
Asn	Leu	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
		35					40					45			
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
	50					55						60			
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65				70					75					80	
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
			85					90						95	
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys

	100	105	110
<210>	118		
<211>	112		
<212>	PRT		
<213>	Artificial Sequence		
<220>			
<223>	mutant antibody L chain		
<400>	118		
Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly			
1 5 10 15			
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser			
20 25 30			
Asn Ser Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser			
35 40 45			
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro			
50 55 60			
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile			
65 70 75 80			
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn			
85 90 95			
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys			
100 105 110			

<210>	119
<211>	112
<212>	PRT
<213>	Artificial Sequence
<220>	
<223>	mutant antibody L chain
<400>	119
Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly	
1 5 10 15	
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser	
20 25 30	
Asn Trp Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser	
35 40 45	
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro	

50	55	60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile		
65	70	75
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn		
85	90	95
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
100	105	110

<210> 120

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 120

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly		
1	5	10
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser		
20	25	30
Asn Tyr Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser		
35	40	45
Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro		
50	55	60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile		
65	70	75
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn		
85	90	95
Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
100	105	110

<210> 121

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 121

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1	5	10	15												
Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
	20		25		30										
Asn	Arg	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
	35		40		45										
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
	50		55		60										
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65			70		75									80	
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
		85			90									95	
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys
	100		105		110										

<210> 122

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 122

Asp	Val	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Thr	Pro	Gly
1	5		10		15										
Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
	20		25		30										
Asn	Val	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
	35		40		45										
Pro	Gln	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
	50		55		60										
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65			70		75									80	
Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Ser	Gln	Asn
		85			90									95	
Thr	His	Val	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys
	100		105		110										

<210> 123

<211> 112

<212> PRT
 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 123
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Pro Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 124
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer

<400> 124
 cttgtacaca gtgacggaaa cacctat 27

<210> 125
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer

<400> 3
 ataggtgttt ccgtcactgt gtacaag 27

<210> 126

<211> 10939

<212> DNA

<213> *Cricetulus griseus*

<400> 126

gagctcaatt aaccctcact aaaggagtc gaticgatcc ttacagaaa acttgcaaac 60
cctcttggag tagaaaagta gtagtatctg acacaaglat cagcaaaatg caaacttctc 120
cccatcccca gaaaaccatt ataaaaaccc ccatatccta tgcccaactg tagtgatata 180
ttatattatga ttattataaaa ctgtcttaag gattcagaaa gcaaagtcag ccttaagcta 240
tagagaccag gcagtcagtg gtggtacaca cctttaatcc caggactcag gattaagaag 300
tagacggacc tctgttagtt caagctcacc attacctaca caagagtga gagtaaccga 360
tctcatgcct ttgatcccag cagctgggat catgtgcatt caatcccagc attcgggagt 420
tatataagac aggagcaagg tctcagagct ggcattcatt ctccagccac attgaggata 480
ggaaaacatt gaagtgtcag gatgctgagg agaggcagca gtttgaggtt tggtagaacc 540
aggatcacct ttgtgtctga ggtagagtaa gaactgtggc tggctgcttt gcttttctga 600
tcttcagctt gaagcttgaa ctccaatatt tgtctctggg tctattatta tcatgttaca 660
cctaacttta aagctgattt acgcaagaca gttgtaggtg gacctttctt tctgtcccac 720
cagttcccaa ataactgaca cggagactca atattaatta taaatgattg gttaatagct 780
cagtcctgtt actggctaac tcttacattt taaattaact catttccatc cctttacttg 840
ctgccatgtg gttcatggct tgttcaagtc ctgcttcttc tgtctctggc tggtagtgcc 900
tctggttctg ccctttatcc cagaattctc ctagtctggc tctcctgccc agctataggc 960
cagtcagctg ttatttaacc aatgagaata atacatattt atagtgtaca aagattgctc 1020
ctcaacaccc aattttttat gtgcaacctg agaacttga ctcattgccc tcatgcttgc 1080
agaggcggca cccttaccga ctaagccacc ttcttagccc tgttgctttt gttttttgag 1140
acaggttcca ctatgtagcc caggctggcc tcaactgac cattctcctg cctaaacctc 1200
ccgaacactg gaattatagt caaggcctac ctgccctggc attttcacac ttttatttcc 1260
tggctgagtc cattgacttt acactcatca aggttgaacc agttggagt ttattacagt 1320
gccaatcgca ctgaatccca cataatcaaa caacttcaag gaagcaaaaa accagttttt 1380
cctgaagatc aatgtcagct tgcctgattc agaataagacc cccgaaaaaa ggcaaatgct 1440
tgataaccaa ttctttctta ttgttcaatc cctgtctgct gtgtgtgaagc tcttgagaaa 1500
ggacagtaag gggacattca tgatcagaga aagagcccca actccccccc cagccccacc 1560
cccacctgt ccacagctg ttggttttgt ttccccctgg ctgacaccca gaaatcacia 1620
cataatcacc taggtcactg taacaagttc ctttctggaa aatgctacaa atgatattgg 1680
taacatgagt aatgaataat gcctggagtc caactccctt gtgacccagc aatgttttcc 1740
gtgggtgctc ccttccccag ctgcaggcct gacatgtacc ttaaaaagcc tccccaggag 1800
gacagaattt tgtgggtact atagtgttct cacaataact tcccctaata cccttactta 1860
gttaccataa ataacatgca gcccttggtg aggcacacag ggctccaatg tacagcttct 1920

cagacactgc aggaaccttc ctctcctaata gcagcactgg tctcttcagg ctggacagca 1980
ggaacccata ccactccaat cctagtgagg agtagagctg tctacgaaaa ccagcagatc 2040
tatagctaaa tgggtttcaa ttttatgctt tgacaaattg tactgacccc acccccaccc 2100
cttccccctt gctgtgctgg gaattgaacc caggaccttg tgcatgccag gcaagtactc 2160
taacactgag ctatagcccc aatctttcat ccaagtctct atgtgtgccc acactcgctt 2220
tttattttga gacaaaaggt tcttattttg agataaggtc tcactatgtt gccttgactt 2280
tttttttttt ttttttttga acttttgacc ttcttacctc agctgagact acaagtcctt 2340
taccatcagg cccggctgat ggtaaaaata cagtatttga aatagtttaa acacatcatc 2400
ttaatggta accacacaat ttccgaaatg ttgctggctc agtctggggc aaacctgtcc 2460
gccccacat tgggtgctagg aagaaagcac agacaagtag ccttcccagc tcaggagtaa 2520
aagacctgga gggggtgccc cacttcggtc aagttcacgg gatggggagg ggtaccttcc 2580
tccagtagtg gtgtatttg gcagttcctc caccgacgcc ctctggaagc acctgcttgg 2640
acccgcaaag ccaggaatgc agcttccctc agggactcgc cagcgagggt aacaggacag 2700
aggcgtccca agagggtcgg ggcggaagg ggaagacagg gtcggcctta gatagggcaa 2760
aggccttctt ggctgtgttc ccggggtaac cgccttgagc ccgacgtggc 2820
gagcgatggg gacagcgagc aggaagtcgt actggggagg gccgcgtagc agatgcagcc 2880
gagggcggcg ctgccaggta caccgagggt caccgagggt gtgagcgcca ggtccctgaa 2940
ccagccaggc ctccagagcc gaggccggcg gaccgacggt acgttctgga atgggaagg 3000
atccgggaca ccgaattgct gcattgagg gctcagaggt tctgatgtgg gaggccagaa 3060
agggttttat ctaccggagg tgatgtgact tccggcctct ggaagtgtg ttggagtctc 3120
tgggaccttg ggtcctctcg actagggttg gaagggtga aatagggtga gggagaaagg 3180
agaggactgc agcaatgtct tcccgaacga cctgggttcg ggagggtcgg aaggacaagg 3240
ggctgttgtg ggggtcttc agacgagg ggggtgtatt ctattttctg ggaagatggt 3300
gtcgalgcac ttgaccaagt ctatcgatc tgaagaggct aggggaacag acagttagag 3360
aggatggtgg agggagtggc agaaccttc cagaaactgg gagaggctct agcacctgca 3420
accttctccc tggcctccgg ggagtccag aagagggcag gaccatggac acaggtgcat 3480
tcgtgccggc gcgtccggc ctggcgaagg tgcgcctct tggaggccgc gggagggcca 3540
gacgcgcgcc cggagagctg gccctttaag gctaccgga ggcgtgtcag gaaatgcgcc 3600
ctgagccgc cctcccga acgcgcccg agacctggca agctgagacg gaactcggaa 3660
ctagcactcg gctcgggcc tcggtgaggc ctgccccg ccatgcctct gtcattgccc 3720
ctcggccgc ctccctgaac ctccgtgacc gccctgcagt cctccctccc ccccttcgac 3780
tcggcggcg cttccggcg ctccgcagc ccgcccctca cgtagccac acctccctct 3840
cggcgctccg cttcccagc ggtccccgac ctgttcttc ctctccacc ctgcccctct 3900
gtccctctcc ctctcttct cccctgact cgtccctatt aggcaacagc cctgtggtc 3960
cagccggcca tggctgtcaa ggctcacacc cttagctagg ccccttctcc ctccctggg 4020
tcttgtctca tgacccctg ccccgccgg gagcgagcg galgtggagc agtgccctct 4080

gcaagcagaa cttcacccaa gccatgtgac aattgaaggc tglaccccca gaccctaaca 4140
tcttggagcc ctgtagacca gggagtgctt ctggccgtgg ggtgacctag ctcttctacc 4200
accatgaaca gggcccctct gaagcgggcc aggatcctgc gcatggcgct gactggaggc 4260
tccactgcct ctgaggaggc agatgaagac agcaggaaca agccgtttct gctgcgggcg 4320
ctgcagatcg cgttggtcgt ctctctctac tgggtcacct ccatctccat ggtattcctc 4380
aacaagtacc tgctggacag cccctccctg cagctggata cccctatctt cgtcactttc 4440
taccaatgcc tggtagacct tctgctgtgc aagggcctca gcactctggc cacctgtgc 4500
cctggcaccg ttgacttccc caccctgaac ctggacctta aggtggcccg cagcgtgctg 4560
ccactgtcgg tagctttcat tggcatgata agtttcaata acctctgcct caagtacgta 4620
ggggtggcct tctacaacgt ggggcgctcg ctaccaccg tgttcaatgt gcttctgtcc 4680
tacctgtgc tcaaacagac cacttccttc tatgccctgc tcacatgtgg catcatcatt 4740
ggtgagtggg gcccgggggc tgtgggagca ggatgggcat cgaactgaag ccctaaaggt 4800
caacactgta ggtaccttta ctactgtcc caggctccct gcatacagc ttacaggaag 4860
agccctgtag aaaacaaata acttccctat ggtcattcaa caagttaggg acccagccag 4920
ggtgaaaata atgttagcag caactacagc aaagatggct ctgccactt gcatgattaa 4980
aatgtgccag gtactcagat ctaagcattg gatccacatt aactcaacta atccctatta 5040
caaggtaaaa tatatccgaa ttttacagag ggaaaaccaa ggcacagaga ggctaagtag 5100
cttgaccagg atcacacagc taataatcac tgacatagct gggatttaaa cataagcagt 5160
tacctccata gatcacacta tgaccacat gccactgtc ctctcaaga gttccaggat 5220
cctgtctgtc cagttctctt taaagaggac aacacatctg acattgctac cttaggtaa 5280
catttgaaat agtgggtaga catatgtttt aagttttatt ctacttttt atgtgtgtgt 5340
gtttgggggg ccaccacagt gtatgggtgg agataagggg acaacttaag aattggtcct 5400
ttctcccacc acatgggtgc tgaggctga actcaggta tcaggattgg cacaatccc 5460
tttaccact gagccatttc actggtcaa tatatgtgtg cttttaagag gctttaacta 5520
ttttcccaga tgtgaatgtc ctgctgatca ttatccctt ttaccggaa gccctctggg 5580
aggtgccatc cctgtggtcg tctgcataca aatggggaaa ctgcaactca gagaaacaag 5640
gctacttgcc agggcccccac aagtaagata ggctgggatg ccatcccaga ctggccacac 5700
tcctggcct gtgcttcaag ccagtttact ttgttcctgc ccattggaag ttagcatgtt 5760
gcagtcaaac acaataacta caggccaaaa gtgcttttaa attaaagtca gatgaacttt 5820
taaacatcca gagctcctca actgcaggag ttacaacctg attctgcaac catctttgca 5880
gtgcccggta gtcatatgta gctagaggct ctgggctagg acagcatgtg ttaggaaaca 5940
tctggccctg agatcattga attgagtgac tgctgggtga caaagaccaa ggcatccgtt 6000
ccctgagagt cctgggcaag cagcaatgtg accttcattt gtacctactc aggttcttta 6060
tctgtcctgt ttgacctact tagtctctc tgggtgtctc gaggcccagg ctgggtactc 6120
tggaatgcag gatcaggcca atgcgcacat ctgccctaga aatgtccccc tggttgagca 6180
gctcctgaat ccatcggtaa aggtcttgga ccaggaggga gtcagataaa aagctgacag 6240

cactggggga ctccatgggg aactcccacc tgcccccaca catccatcct aagagaactg 6300
 gtattccttg tttcctcttt gtcctacaag gcaccctggg atcccacttc agtctcccag 6360
 ccttgccagg gttagagggc atgagcctcc ttgtggggaa tttagatgca agaaggtaca 6420
 gtcactagag aacctgagct cagatcccca aagtaaccag tacctgatag tgaggcagct 6480
 gagaaccgca gcagcctgcc tgagtggctg aactctgcgg cctccggaac tggccccaac 6540
 tgttgggtct cctcttcctt cctcctgtga gggagggccc atctctgata agtgctgtgg 6600
 ggactctaga gtagggagga ggaggagcaa tctaagcagg ccttactgag aagtccttgc 6660
 tggcatgtgg ctgcctgagg agtacagact gggaacaccc atttgaatga gtaaggtttt 6720
 tcctgaaggc catggggagc cacggaggaa aatcatttta gttacaagac aaagagtaga 6780
 ttggttaaca tgggagcaag gacatggccc caattttcat agatgaagga aattggaact 6840
 cagagagggt aagtaacttc tcccaaatag ctacagcttc aaatcacaga acagtcagag 6900
 tctagatctc tctgatgcct gtgatggicc tgccattcca tgttctgat ccctgtggca 6960
 tcagtaagcc tctacctgtt gggaatgcag gatctaaatg aagagaggaa gtgctggccc 7020
 catgctgtgg tctggaaagc tatgcaggct ctttgagcag agagtgaccc acaagtgaat 7080
 agagtcctat gagactcaaa gcaacatcca cccctaagca gctctaacca aatgctcaca 7140
 ctgagggagc caaagccaag ttagagtcct gtgcttggcc aaggctactt tgcctggccc 7200
 tcctcctata gcaccgtgt tatcttatag ccttcattac agtgattaca attataatta 7260
 gagaggtaac agggccacac tgtccttaca cattccctg ctagattgta gctgggagag 7320
 ggggagatgt aggtggctgg gggagtggga gggaagatgc agattttcat tctgggctct 7380
 acicccctcag ccattttttg gtgtgggagt tagactttgg atatgttgat gatgaggtaa 7440
 gggccacaga acagctgaa ctgtggtatc agaatcctgt ccctctccct ctctcctcat 7500
 ccctcttcac ctgttcactc ctctgtctgc tacagggtgt tcttggtggt gtatagacca 7560
 agaggggagct gagggcaccc tgtccctcat aggcaccatc ttcggggtgc tggccagcct 7620
 ctgcgtctcc ctcaatgcca tctatacca gaaggtgctc ccagcagtgg acaacagcat 7680
 ctggcgcta accttctata acaatgtcaa tgcctgtgtg ctcttcttgc ccctgatgtt 7740
 tctgtgtggt gagctccgtg cctccttga ctttgctcat ctgtacagt cccacttctg 7800
 gctcatgatg acgttgggtg gcctcttcgg ctttgccatt ggctaigtga caggactgca 7860
 gatcaaattc accagtcccc tgaccacaaa tgtatcaggc acagccaagg cctgtgcgca 7920
 gacagtgtg gccgtgctct actatgaaga gactaagagc ttcctgtggt ggacaagcaa 7980
 cctgatggtg ctgggtggct cctcagccta tacctgggtc aggggctggg agatgcagaa 8040
 gacccaagag gaccccagct ccaaagaggg tgagaagagt gctatigggg tgtgagcttc 8100
 ttcagggacc tgggactgaa cccaagtggg gcctacacag cactgaaggc ttcccatgga 8160
 gctagccagt gtggccctga gcaatactgt ttacatctc cttggaatat gatctaagag 8220
 gagccagggt ctttctgtgt aatgtcagaa agctgccaaa tctcctgtct gccccatctt 8280
 gttttgggaa aacctacca ggaatggcac ccctacctgc ctctcctag agcctgtcta 8340
 cctccatatc atctctgggg ttgggaccag ctgcagcctt aaggggctgg attgatgaag 8400

tgatgtcttc tacacaaggg agatgggttg tgatcccact aattgaaggg atttgggtga 8460
 cccacacact ctgggatcca gggcaggtag agtagtagct taggtgctat taacatcagg 8520
 aacacctcag cctgccittg aaggaagtg ggagcttggc caagggagga aatggccatt 8580
 ctgccctctt cagtgtggat gagtatggca gacctgttca tggcagctgc accctgggt 8640
 ggctgataag aaaacattca cctctgcatt tcatatttgc agctctagaa cgggggagag 8700
 ccacacatct tttaggggtt aagtaggggtg atgagctcct ccgcagtcct taaccccagc 8760
 ttacctgcc tggcttcctt tggcccagct acctagctgt actccctttc tgtactcttc 8820
 tcttctccgt catggcctcc cccaacacct ccatctgcag gcaggaagtg gagtccactt 8880
 gtaacctctg ttcccatgac agagcccttt gaatacctga acccctcatg acagtaagag 8940
 acatttatgt tctctggggc tggggctgaa ggagccctact ggttctcact tagcctatct 9000
 ggctcctgtc aaaaaaaaaa aaaaagaaaa aaaaaagca taaaccaagt tactaagaac 9060
 agaagtgggt ttataacgtt ctggggcagc aaagcccaga tgaagggacc catcgacctt 9120
 ctctgtccat atcctcatgc tgcagaagta caggcaagct cctttaagcc tcatatagga 9180
 acactagcct cactcatgag ggttttactc catgacctgt caacctcaaa gccttcaaca 9240
 tgaggactcc aacgtaaatt tggggacaga agcactcaga ccatacccca gcaccacacc 9300
 ctctaacct cagggtagct gtcattctcc tagtctctc tcttgggcct ttagaacccc 9360
 catttccttg gggtaatgtc tgatgttttt gtccctgtca taaaaagatg gagagactgt 9420
 gtccagcctt tgattcctac ttctacaat ccaggttct aatgaagttt gtggggcctg 9480
 atgccctgag ttgtatgtga ttttaataa aaaaagcaag atacagcatg tgtgtggact 9540
 gagttagggc cacagggatc taaaagccaa gtgtgagggg acccagctac agcaggcagc 9600
 atcctgagcc tggaatctct tcaggacaag aattctccat atacctacct actctgggga 9660
 gtaggtggcc agagtccaag cttcccttag taccaactac cactggctgt gctcttactg 9720
 aaggcagaca tggcactgag tgctgtccat ctgtcactca tctccacagc cattccta 9780
 gtgtgggttg ggagccatca ccaaacccca ttttcagata aggacacagg ctccagagagg 9840
 ctgtgttgga gaaaagtagc agcagaattc agagagctgg gtctcctgca gcaccttgga 9900
 ctgccagcag ccacagtgtt tgtcacacag cacatactca aaagaatgcc agccccctca 9960
 gcctagagtg cctggccttt ctttcagatg aggaagaggg tcaaagctgt tagcttgccc 10020
 accatatgac cacatacatg accaacagct tgagggaggg aggattactg tggctcccag 10080
 cctgagaggt gggacacca aatgtattag gtccttgaat cagggtgac ctgtgtattc 10140
 agtcactcct accagaatgc tggggaatgg ggatgcaaaa ggcaaaggag gctttctaag 10200
 gtgtgggtga agataggcat ttctgtcttc atgtacacct gtgagcagag taggaaggcc 10260
 ctgtggagaa tataatccac aaaccagtag ccttctctgg cagtgggtga atactgccac 10320
 cctatacccc tatgcaaggc cagtagaacc acccaacca caacatctag agaaattaca 10380
 ggtcatctta agcctctaaa ttgtggagaa actcgacatg cgcacgattc ctaacctgct 10440
 agcctagggt gcgggttgga taatttaagg aaactgggtt ttcttataga atcggaggct 10500
 ccatgaagtc accctgacaa gaggtcagca atagccagca gcagtggcta ctcttaagcc 10560

tccagacaga gcaccctgtg aatgtacctt attctcacat ctgggtgtct atagggtga 10620
ctgggtcaga tgcacccag gccattgcaa tgggccctta gcccctggg gtgttgggat 10680
agcagccaag cagctcccat gctgagatac tgcctgcagt agactgatgg ataagaaaac 10740
aaggcccaaa atgttttctt tccagacttg atctttcttt gttcaaaaat gctgttttcc 10800
cttaaacttg cccaaacca ttgttttgca gttgaggaaa ataaggcata gaaagattaa 10860
aggaagtffc tgaggttaca gagcaaagta ctggcttcac ctgaaataga cagggtgtgcc 10920
ctgatcctga tttgagctc 10939

<210> 127

<211> 352

<212> PRT

<213> *Cricetulus griseus*

<400> 127

Met	Ala	Leu	Thr	Gly	Gly	Ser	Thr	Ala	Ser	Glu	Glu	Ala	Asp	Glu	Asp
1				5					10					15	
Ser	Arg	Asn	Lys	Pro	Phe	Leu	Leu	Arg	Ala	Leu	Gln	Ile	Ala	Leu	Val
			20					25						30	
Val	Ser	Leu	Tyr	Trp	Val	Thr	Ser	Ile	Ser	Met	Val	Phe	Leu	Asn	Lys
			35					40						45	
Tyr	Leu	Leu	Asp	Ser	Pro	Ser	Leu	Gln	Leu	Asp	Thr	Pro	Ile	Phe	Val
			50					55						60	
Thr	Phe	Tyr	Gln	Cys	Leu	Val	Thr	Ser	Leu	Leu	Cys	Lys	Gly	Leu	Ser
			65					70						75	
Thr	Leu	Ala	Thr	Cys	Cys	Pro	Gly	Thr	Val	Asp	Phe	Pro	Thr	Leu	Asn
								85						90	
Leu	Asp	Leu	Lys	Val	Ala	Arg	Ser	Val	Leu	Pro	Leu	Ser	Val	Val	Phe
								100						105	
Ile	Gly	Met	Ile	Ser	Phe	Asn	Asn	Leu	Cys	Leu	Lys	Tyr	Val	Gly	Val
								115						120	
Ala	Phe	Tyr	Asn	Val	Gly	Arg	Ser	Leu	Thr	Thr	Val	Phe	Asn	Val	Leu
								130						135	
Leu	Ser	Tyr	Leu	Leu	Leu	Lys	Gln	Thr	Thr	Ser	Phe	Tyr	Ala	Leu	Leu
								145						150	
Thr	Cys	Gly	Ile	Ile	Ile	Gly	Gly	Phe	Trp	Leu	Gly	Ile	Asp	Gln	Glu
								165						170	
Gly	Ala	Glu	Gly	Thr	Leu	Ser	Leu	Ile	Gly	Thr	Ile	Phe	Gly	Val	Leu
								180						185	
														190	

Ala Ser Leu Cys Val Ser Leu Asn Ala Ile Tyr Thr Lys Lys Val Leu
 195 200 205
 Pro Ala Val Asp Asn Ser Ile Trp Arg Leu Thr Phe Tyr Asn Asn Val
 210 215 220
 Asn Ala Cys Val Leu Phe Leu Pro Leu Met Val Leu Leu Gly Glu Leu
 225 230 235 240
 Arg Ala Leu Leu Asp Phe Ala His Leu Tyr Ser Ala His Phe Trp Leu
 245 250 255
 Met Met Thr Leu Gly Gly Leu Phe Gly Phe Ala Ile Gly Tyr Val Thr
 260 265 270
 Gly Leu Gln Ile Lys Phe Thr Ser Pro Leu Thr His Asn Val Ser Gly
 275 280 285
 Thr Ala Lys Ala Cys Ala Gln Thr Val Leu Ala Val Leu Tyr Tyr Glu
 290 295 300
 Glu Thr Lys Ser Phe Leu Trp Trp Thr Ser Asn Leu Met Val Leu Gly
 305 310 315 320
 Gly Ser Ser Ala Tyr Thr Trp Val Arg Gly Trp Glu Met Gln Lys Thr
 325 330 335
 Gln Glu Asp Pro Ser Ser Lys Glu Gly Glu Lys Ser Ala Ile Gly Val
 340 345 350

<210> 128

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 128

ggatcctgcg catgaaaaag cctgaactca cc

32

<210> 129

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 129

gcggccgcct attcctttgc cctcggacg	29
<210> 130	
<211> 33	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: Primer	
<400> 130	
atgcatgccca ccatgaaaaa gcctgaactc acc	33
<210> 131	
<211> 28	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: Primer	
<400> 131	
ggatcccagg ctttacactt tatgcttc	28
<210> 132	
<211> 25	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: Primer	
<400> 132	
gctgtctgga gtactgtgca tctgc	25
<210> 133	
<211> 27	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: Primer	
<400> 133	
ggaatgcagc ttctcaagg gactcgc	27
<210> 134	
<211> 27	
<212> DNA	

<213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 134
 tgcacacaggt cggagacgct gtcgaac 27
 <210> 135
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 135
 gcactcgtcc gagggcaaag gaatagc 27
 <210> 136
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 136
 tgtgctggga attgaacca ggac 24
 <210> 137
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 137
 ctacttgtct gtgctttctt cc 22
 <210> 138
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 138

ctcgactcgt ccctattagg caacagc

27

<210> 139

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 139

tcagaggcag tggagcctcc agtcagc

27

1/44

8