



(51) International Patent Classification:

C07K 7/08 (2006.01) *A61K 38/10* (2006.01)
C07K 19/00 (2006.01) *A61P 37/00* (2006.01)

(21) International Application Number:

PCT/US2013/070813

(22) International Filing Date:

19 November 2013 (19.11.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/728,368 20 November 2012 (20.11.2012) US
 61/774,066 7 March 2013 (07.03.2013) US

(71) Applicant: THE U.S.A, AS REPRESENTED BY THE
 SECRETARY, DEPARTMENT OF HEALTH AND
 HUMAN SERVICES [US/US]; National Institutes of
 Health, Office of Technology Transfer, 6011 Executive
 Blvd., Suite 325, MSC 7660, Bethesda, MD 20892-7660
 (US).

(72) Inventors: ASHWELL, Jonathan, D.; 2703 Daniel Road,
 Chevy Chase, MD 20815 (US). ALAM, Muhammad, S.;
 10444 Rockville Pike, Apt # 201, Rockville, MD 20852
 (US).

(74) Agent: CONNOLLY, Jodi, L.; Klarquist Sparkman, LLP,
 One World Trade Center, Suite 1600, 121 Sw Salmon
 Street, Portland, OR 97204 (US).

(81) Designated States (unless otherwise indicated, for every
 kind of national protection available): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
 BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
 DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
 HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
 KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
 OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
 SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
 ZW.

(84) Designated States (unless otherwise indicated, for every
 kind of regional protection available): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
 UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
 TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
 MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
 TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
 KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: INHIBITORS OF THE T CELL-SPECIFIC ALTERNATIVE P38 ACTIVATION PATHWAY AND METHODS OF USE

(57) Abstract: In T lymphocytes, p38 mitogen activated protein kinase (MAPK) can be activated through an alternative pathway that involves phosphorylation at tyrosine 323. Disclosed herein is the identification of a minimal region of the growth arrest and DNA damage-inducible alpha (Gadd45a) protein that is required for binding to and inhibition of tyrosine 323-phosphorylated p38 in T cells. The disclosed Gadd45a polypeptides inhibit proliferation of T cells in response to T cell receptor stimulation, inhibit differentiation of T cells into Th1 or Th17 cells, inhibit the production of proinflammatory cytokines, and reduce tumor formation and growth of inflammatory cancers, such as pancreatic cancer.



WO 2014/081728 A1

INHIBITORS OF THE T CELL-SPECIFIC ALTERNATIVE P38 ACTIVATION PATHWAY AND METHODS OF USE

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of U.S. Provisional Application No. 61/774,066, filed March 7, 2013, and U.S. Provisional Application No. 61/728,368, filed November 20, 2012, both of which are herein incorporated by reference in their entirety.

FIELD

10 This disclosure concerns inhibitors of the T-cell specific alternative p38 activation pathway and their use for the treatment of p38-associated diseases, such as autoimmune and inflammatory disorders, including inflammatory cancers. In particular, this disclosure concerns Gadd45 α polypeptides capable of inhibiting p38 activation in T cells.

BACKGROUND

15 The p38 mitogen-activated protein kinase (MAPK) is a signaling intermediate downstream of proinflammatory cytokine receptors released following environmental stress. This kinase is known to play an important role in inflammatory and autoimmune diseases, including rheumatoid arthritis and multiple sclerosis. The p38 MAPK family includes four separately encoded members
20 – p38 α , p38 β , p38 γ and p38 δ ; p38 α , p38 β and p38 δ are expressed in T cells, while p38 γ is primarily expressed in skeletal muscle (Jirmanova *et al.*, *Blood* 118(12):3280-3289, 2011). The p38 kinase is ubiquitously expressed and is activated in all cells by a series of sequential phosphorylation steps that result in dual phosphorylation at threonine 180 and tyrosine 182. However, in T cells, direct stimulation of the T cells through their antigen receptor results in
25 activation of an alternative p38 pathway, leading to phosphorylation of p38 α and p38 β on tyrosine 323 (Salvador *et al.*, *Nat Immunol* 6(4):396-402, 2005).

 Genetically modified mice that express p38 α and p38 β containing a tyrosine to phenylalanine substitution at residue 323 (Y323F) have been developed and characterized to better understand the role of the T cell-specific alternative p38 activation pathway *in vivo*. Due to the
30 phenylalanine substitution, the modified forms of p38 α and p38 β cannot be phosphorylated at residue 323, which prevents activation of the alternative p38 pathway. The genetically modified mice exhibit a significant reduction in T cell proliferation, proinflammatory cytokine production, as well as reduced susceptibility to animal models of autoimmune diseases such as multiple sclerosis

and rheumatoid arthritis. Studies of these mice established that the alternative p38 activation pathway in T cells plays an important role in T cell activation and the development of autoimmune and inflammatory diseases (Jirmanova *et al.*, *Blood* 118(12):3280-3289, 2011).

Gadd45 (growth arrest and DNA damage-inducible genes) is a family of proteins that play a role in cell growth regulation, differentiation and apoptosis. In mammalian cells, three different Gadd45 family members have been identified – Gadd45 α , Gadd45 β and Gadd45 γ . Using Gadd45 α -deficient mice, prior studies have shown that Gadd45 α inhibits the alternative p38 activation pathway, and the absence of Gadd45 α results in T cell hyperproliferation and autoimmunity (Salvador *et al.*, *Nat Immunol* 6(4):396-402, 2005).

SUMMARY

Provided herein are growth arrest and DNA-damage-inducible alpha (Gadd45 α) polypeptides capable of inhibiting the T-cell specific alternative p38 activation pathway. In some embodiments, the Gadd45 α polypeptide is no more than 25 or no more than 20 amino acids in length and comprises residues 71-85 of human or mouse Gadd45 α . In some cases, the Gadd45 α polypeptides are synthetic or recombinant polypeptides (*i.e.* non-naturally occurring polypeptides). Also provided are fusion proteins comprising a Gadd45 α polypeptide and a heterologous peptide, such as a cell-penetrating peptide. Further provided are compositions comprising the Gadd45 α polypeptides and fusion proteins disclosed herein.

A method is also provided for inhibiting proliferation of T cells in response to T cell receptor stimulation by contacting the T cells with a Gadd45 α polypeptide, fusion protein or composition disclosed herein. Further provided is a method of inhibiting differentiation of T cells into Th1 or Th17 cells by contacting the T cells with a Gadd45 α polypeptide, fusion protein or composition disclosed herein. In some embodiments, the method is an *ex vivo* method wherein the T cells have been isolated from a subject. In other embodiments, the method is an *in vivo* method that includes administering the polypeptide, fusion protein or composition to a subject.

Methods of treating a subject having a T cell- and/or p38-mediated autoimmune or inflammatory disorder are also provided by the present disclosure. In some embodiments, the method includes selecting a subject with a T cell- and/or p38-mediated autoimmune or inflammatory disorder and administering to the subject a therapeutically effective amount of a Gadd45 α polypeptide, fusion protein or composition disclosed herein.

Further provided herein is a method of inhibiting tumor vascularization in a subject. In some embodiments, the method includes selecting a subject with a tumor, for example a pancreatic

tumor, and administering to the subject a therapeutically effective amount of a Gadd45 α polypeptide, fusion protein or composition disclosed herein.

Also provided herein is a method of determining the prognosis of a subject diagnosed with pancreatic ductal adenocarcinoma (PDAC) by calculating the percentage of tumor-infiltrating T cells positive for p38 phosphorylated at tyrosine 323 (p38 pY323) in a tumor sample obtained from the subject; and determining that the subject has a poor prognosis if at least 10% of the T cells are positive for p38 pY323. In some embodiments, the poor prognosis is a decrease in time of survival.

The foregoing and other objects, features, and advantages of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph showing cellular uptake of (11R)-71-85 and (11R)-Ctrl. Purified CD4⁺ T cells were treated with either FITC conjugated (11R)-71-85 or (11R)-Ctrl, or left untreated for 2 hours at 37°C. After incubation, cells were washed and analyzed by flow cytometry for FITC positivity.

FIG. 2 shows inhibition of the p38 alternative pathway activation by (11R)-71-85. Purified p38 proteins were activated by incubating with active Zap70 in kinase buffer for 1 hour. (11R)-71-85 or (11R)-Ctrl was then added and incubated for 30 minutes at 30°C. One μ g ATF2 and 10 μ Ci [³²P]ATP were then added and incubated for 45 minutes at 30°C. Phosphorylated products were resolved by SDS-PAGE, transferred to nitrocellulose membranes, and visualized by phosphorimager.

FIG. 3 is a graph showing T cell proliferation in response to (11R)-71-85. T cells were purified from spleen and treated with different concentrations of (11R)-71-85 or control peptide (11R)-Ctrl. Cells were then washed and stimulated with anti-CD3 and anti-CD28 for 2 days and finally pulsed with 1 μ Ci [³H]-thymidine 14 hours before harvesting. [³H]-thymidine uptake was determined with a Wallac 1450 MicroBeta Liquid Scintillation Counter.

FIGS. 4A-4B are a series flow cytometry plots showing inflammatory cytokine secretion by CD4⁺ (FIG. 4A) and CD8⁺ (FIG. 4B) T cells. T cells were purified from spleen and stimulated with anti-CD3 plus anti-CD28 for 3 days. Cells were then left unstimulated overnight and treated with the indicated concentrations of (11R)-71-85, control peptide (11R)-Ctrl, or DMSO (no peptide). After treatment, cells were restimulated with either anti-CD3 plus anti-CD28, or PMA

plus ionomycin supplemented with monensin. Cytokine expression was measured by flow-cytometry.

FIGS. 5A-5D are a series of flow cytometry plots showing (11R)-71-85 inhibits Th1 differentiation. Naïve CD4⁺ T cells were treated with either 2 µM of (11R)-71-85 or (11R)-Ctrl for two hours and after washing, cells were stimulated with anti-CD3 plus anti-CD28 in Th0 (neutral, FIG. 5A and FIG. 5C) or Th1 polarizing conditions (FIG. 5B and FIG. 5D) for 3 days. Cells were treated again with the same concentration of either medium (first column), (11R)-Ctrl (second column), or (11R)-71-85 (third column) for 2 hours and restimulated with PMA plus ionomycin in the presence of Golgi inhibitor for 4 hours. Expression of T bet and interferon (IFN)-γ (indicators of Th1 cells typical of a pro-inflammatory response) was analyzed by flow cytometry. Whereas cells treated (11R)-Ctrl responded similar to cells treated with medium, the induction of T-bet and IFNγ was blunted by treatment with (11R)-71-85.

FIGS. 6A and 6B show inhibition of Th17 differentiation by full-length Gadd45α. (FIG. 6A) Purified CD4⁺ T cells from WT mice were treated with 4 µM of either TAT-Ctrl, TAT-G45FL or TAT-G45Δ10 and stimulated with plate coated anti-CD3 plus anti-CD28 for 48 hours and blotted for the indicated transcription factors. (FIG. 6B) Naïve CD4⁺ T (CD4⁺CD62L^{hi}) cells were purified from WT mice and treated with either TAT-Ctrl or TAT-G45FL as in (A) and stimulated in neutral (Th0) or Th17 skewing conditions (10 ng/ml IL-6 + 5 ng/ml TGF-β) for 48 hours. The mRNA expression levels of Rorc and Il17a were evaluated. *P* value of ≤ 0.05 was considered as a statistically significant difference and designated with an asterisk (*).

FIG. 7 is a graph showing expression of interferon regulatory factor 4 (Irf4) mRNA following *in vivo* treatment with (11R)-71-85. (11R)-71-85 (40 µg) or vehicle control was injected intraperitoneally into wild type mice. Four hours after injection, splenocytes were collected and stimulated with soluble anti-CD3 plus anti-CD28 for 24 hours. Irf4 mRNA expression was assessed by quantitative real time PCR.

FIG. 8 is a series of graphs showing reduction in expression of Nfatc1 and Irf4 in CD4⁺ T cells following injection of (11R)-71-85. WT C57BL/6 mice were injected i.v. with 50 µg of DMSO, (11R)-Scrambled, or (11R)-71-85 and either total T cells or CD4⁺T cells were purified from spleen at different times. Purified cells were stimulated *in vitro* with anti-CD3/CD28 for 24 hours and the mRNA levels of the indicated transcription factors were determined by quantitative real-time PCR.

FIGS. 9A-9C show injection of (11R)-71-85 stops the growth of established PANC02 tumors. WT C57BL/6 mice were injected with 7.5x10⁵ PANC02 cells. After 16 days, mice

received intra-tumor injections of (11R)-71-85 at the indicated amount or DMSO every other day. (FIG. 9A) Tumor volume from the start of injection (day 16) to day 42 in control mice and mice injected with 40 μ g or 80 μ g of (11R)-71-85. Sacrifice (sac.) of mice before day 42 is indicated. (FIG. 9B) weight and (FIG. 9C) appearance of the control (left) and treated (right) tumors on day 38.

FIGS. 10A-10B are a series of flow cytometry plots demonstrating that intra-tumor injection of (11R)-71-85 inhibits tumor necrosis factor (TNF)- α secretion from tumor-infiltrating T cells. WT C57BL/6 mice were injected with 7.5×10^5 PANC02 cells. After 16 days, mice were injected with 80 μ g of either (11R)-71-85 or (11R)-Scrambled (Scr) every other day for 1 week. Twenty-four hours after the last intra-tumor injection, tumor infiltrating cells were harvested and restimulated with PMA and ionomycin, or left unstimulated, in the presence of Golgi inhibitor for 4 hours. Expression of cytokines (IL-17A and TNF α – FIG. 10A; IL-10 and IFN γ – FIG. 10B) was determined by flow cytometry.

FIG. 11 is a pair of images showing that intra-tumor injection of (11R)-71-85 reduces tumor vascularization. WT C57BL/6 mice were injected with 7.5×10^5 PANC02 cells and 16 days later were injected with 80 μ g of either (11R)-71-85 or (11R)-Scrambled. Mice received intra-tumor injections every other day for 2.5 weeks. Twenty-four hours after the last injection, tumors were collected, fixed in formalin, and stained for the endothelial marker CD31, an indicator of tumor vascularization.

FIG. 12 is a panel of flow cytometry plots showing that intraperitoneal injection of (11R)-71-85 inhibits TNF- α secretion from tumor-infiltrating T cells from KPC mice. KPC mice are genetically engineered to develop pancreatic cancer because of oncogene expression and p53 haploinsufficiency. KPC mice received intraperitoneal injections of 80 μ g of either (11R)-71-85 or (11R)-Scrambled every other day for 1 month. Twenty-four hours after the last injection, tumor infiltrating cells were harvested and restimulated with PMA and ionomycin, or left unstimulated, in the presence of Golgi inhibitor for 4 hours. Expression of intracellular TNF- α was determined by flow cytometry.

FIGS. 13A-13D are graphs showing that (11R)-71-85 inhibits PanIN and cancer formation in KPC mice. (FIG. 13A) KPC mice were sub-lethally irradiated at 6 weeks of age and packed with 10×10^6 hematopoietic bone marrow cells and the survival was monitored (n=5 per group; * $p < 0.05$, Log-rank test). (FIG. 13B) Quantification of normal ducts and PanIN of KPC mice treated as in (A) (n=6 per group). (FIG. 13C) CD4 $^+$ T cells infiltrated in the pancreas were analyzed for TNF α and IFN γ expression by flow cytometry after treatment with (11R) 71-85 or (11R) Scr. Bar

graphs show the average percentage of indicated cytokine secreting cells. (n=8 per group; * $p < 0.05$, *NS*=not significant, unpaired two-tailed Student's *t*-test). (FIG. 13D) KPC mice were treated as in (A) and the expression of different tumor promoting factors was analyzed in isolated pancreatic epithelial cells by quantitative real-time PCR (n=6 per group).

FIGS. 14A-14C are graphs showing higher p38 alternative expression in T cells is associated with shorter survival and tumor neoangiogenesis in human PDAC. (FIG. 14A) Correlation of the patient's survival with lower (0-10%, n=153) versus higher ($\geq 10\%$, n=40) levels of tumor infiltrating T cells in human pancreas activated by the alternative pathway (Log-rank test). (FIG. 14B) Human PDAC sections were stained for CD3 and the absolute number of CD3⁺ cells was analyzed per high power field in all samples of lower (0-10%, n=153) and higher ($\geq 10\%$, n=40) levels of p38 alternatively activated tumor infiltrating T cells (*NS*=not significant, unpaired two-tailed Student's *t*-test). (FIG. 14C) Immunohistochemistry with anti-CD31 antibody for the detection of vascular density of patient's samples with lower (0-10%, n=10) versus higher ($\geq 10\%$, n=10) activation of p38 alternative pathway. (* $p < 0.05$, unpaired two-tailed Student's *t*-test).

SEQUENCE LISTING

The and amino acid sequences listed in the accompanying sequence listing are shown using standard three letter code for amino acids, as defined in 37 C.F.R. 1.822 The Sequence Listing is submitted as an ASCII text file, created on October 29, 2013, 2.61 KB, which is incorporated by reference herein. In the accompanying sequence listing:

SEQ ID NO: 1 is an amino acid consensus sequence of a Gadd45 α polypeptide corresponding to residues 71-85 of Gadd45 α .

SEQ ID NO: 2 is the amino acid sequence of residues 71-85 of human Gadd45 α .

SEQ ID NO: 3 is the amino acid sequence of residues 71-85 of mouse Gadd45 α .

SEQ ID NO: 4 is the amino acid sequence of isoform 1 of the human Gadd45 α protein (GenBank Accession No. NP_001915).

DETAILED DESCRIPTION

I. Abbreviations

ATF2	activating transcription factor 2
CIA	collagen-induced arthritis
CPP	cell-penetrating peptide
EAE	experimental autoimmune encephalomyelitis
Gadd45 α	growth arrest and DNA-damage-inducible alpha
IFN	interferon

	IL	interleukin
	IRF	interferon regulatory factor
	i.v.	intravenous
	LT	lymphotoxin
5	MAPK	mitogen-activated protein kinase
	MS	multiple sclerosis
	NFATc1	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1
	PanIN	pancreatic intraepithelial neoplasia precursor lesions
10	PDAC	pancreatic ductal adenocarcinoma
	RA	rheumatoid arthritis
	TCR	T cell receptor
	TGF	transforming growth factor
	Th	T helper
15	TIL	tumor infiltrating lymphocytes
	TNF	tumor necrosis factor

II. Terms and Methods

Unless otherwise noted, technical terms are used according to conventional usage.

20 Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew *et al.* (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

25 In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

Administer: As used herein, administering a composition (*e.g.* a polypeptide) to a subject means to give, apply or bring the composition into contact with the subject. Administration can be accomplished by any of a number of routes, such as, for example, intravenous, intrathecal, topical,
30 oral, subcutaneous, intramuscular, intraperitoneal, intramuscular or by direct injection into a tissue, such as into a tumor.

Autoimmune disease: A type of disease that arises from inappropriate immune responses targeted against substances or tissues normally present in the body. Autoimmune diseases, include, but are not limited to amyotrophic lateral sclerosis, autoimmune hepatitis, celiac disease, Crohn's
35 disease (inflammatory bowel disease), diabetes type 1, Graves' disease, Guillain-Barre syndrome, Hashimoto's disease, multiple sclerosis, myasthenia gravis, psoriasis, rheumatoid arthritis, Sjögren's syndrome, systemic lupus erythematosus, and ulcerative colitis.

Clinical outcome: Refers to the health status of a patient following treatment for a disease or disorder (such as pancreatic cancer), or in the absence of treatment. Clinical outcomes include,

but are not limited to, a decrease in incidence of cancer, an increase in the length of time until death, a decrease in the length of time until death, an increase in the chance of survival, an increase in the risk of death, survival, disease-free survival, chronic disease, metastasis, advanced or aggressive disease, disease recurrence, death, and favorable or poor response to therapy.

5 **Cell-penetrating peptide (CPP):** Peptides that facilitate the cellular uptake of another protein or molecular cargo linked by a covalent bond or non-covalent interaction. CPPs generally deliver cargo into a cell by endocytosis. In many instances, CPPs have an amino acid composition that is rich in charged amino acids, such as lysine or arginine, or have sequences that contain an alternating pattern of polar/charged amino acids and non-polar/hydrophobic amino acids.

10 **Collagen-induced arthritis (CIA) model:** The most commonly used autoimmune model of rheumatoid arthritis. This model can be used to study disease pathogenesis and to validate therapeutic agents. Autoimmune arthritis is induced in this model by immunization with an emulsion of complete Freund's adjuvant and type II collagen (CII). Protocols for handling and preparation of CII, as well as selection of mouse strains, proper immunization technique and
15 evaluation of the arthritis incidence and severity are known in the art (see, for example, Brand *et al.*, *Nat Protoc* 2(5):1269-1275, 2007). Typically, the first signs of arthritis appear in this model 21-28 days after immunization. The chief pathological features of CIA include a proliferative synovitis with infiltration of polymorphonuclear and mononuclear cells, pannus formation, cartilage degradation, erosion of bone, and fibrosis. As in rheumatoid arthritis in humans, pro-
20 inflammatory cytokines, such as TNF- α and IL-1 β , are abundantly expressed in the arthritic joints of mice with CIA, and blockade of these molecules results in a reduction of disease severity.

Contacting: Placement in direct physical association; includes both in solid and liquid form.

Decrease in survival: As used herein, "decrease in survival" refers to a decrease in the
25 length of time before death of a patient, or an increase in the risk of death for the patient. A decrease in survival also can refer to a decrease in the average time to death in a group, such as a group of patients diagnosed with pancreatic cancer.

Differentiation: Refers to the process by which a cell develops into a specific type of cell. For example, a T cell can develop into a specific T cell subtype, such as Th1, Th2, Th17, T
30 regulatory cell or cytotoxic T cell. The differentiation of T cells into specific T helper subtypes is influenced by the cytokine environment (*e.g.*, IFN- γ and IL-12 promote Th1 differentiation; IL-4 promotes Th2 differentiation; and TGF- β , IL-6, IL-21 and IL-23 promote Th17 differentiation).

Experimental autoimmune encephalomyelitis (EAE): An animal model of MS (for example, see Gold et al., *Brain* 129:1953-1971, 2006). EAE animals exhibit characteristic plaques of tissue injury disseminated throughout the central nervous system. Plaques show infiltration of nervous tissue by lymphocytes, plasma cells, and macrophages, which cause destruction of the myelin sheaths that surround nerve cell axons in the brain and spinal cord. In some cases, EAE is induced by immunization of susceptible animals, such as mice, rats, guinea pigs, or non-human primates, with either myelin or various components of myelin. For example, EAE can be induced by immunization with components of the myelin sheath, such as myelin basic protein, proteolipid protein, or myelin oligodendrocyte glycoprotein (MOG). EAE is a useful and widely accepted model for studying mechanisms of autoimmune CNS tissue injury and for testing potential therapies for MS. EAE also includes “passive EAE” which is induced in the same manner in donor animals, but involves the transfer of activated T-cells harvested from the donor animal’s lymph nodes to naïve recipient animals.

Fusion protein: A protein containing amino acid sequence from at least two different (heterologous) proteins or peptides. In some examples herein, the fusion protein comprises a portion of Gadd45 α and a cell-penetrating peptide. Fusion proteins can be generated, for example, by expression of a nucleic acid sequence engineered from nucleic acid sequences encoding at least a portion of two different (heterologous) proteins. To create a fusion protein, the nucleic acid sequences must be in the same reading frame and contain no internal stop codons. Fusion proteins, particularly short fusion proteins, can also be generated by chemical synthesis.

Growth arrest and DNA-damage-inducible alpha (Gadd45 α): A member of a group of genes whose transcript levels are increased following stressful growth conditions and treatment with DNA-damaging agents. The Gadd45 α protein responds to environmental stresses by mediating activation of the p38/JNK pathway via MTK1/MEKK4 kinase. Gadd45 α has previously been shown to inhibit the alternative p38 activation pathway in T cells and the absence of Gadd45 α results in T cell hyperproliferation and autoimmunity (Salvador *et al.*, *Nat Immunol* 6(4):396-402, 2005). Nucleotide and amino acid sequences for Gadd45 α are publically available, such as in the NCBI database (for example under Gene ID 1647 for human Gadd45 α). An exemplary sequence for human Gadd45 α is set forth herein as SEQ ID NO: 4.

Heterologous: A heterologous protein or polypeptide refers to a protein or polypeptide derived from a different source or species.

Inflammatory disease or disorder: Any disease, disorder or condition that is characterized by the presence of inflammation (*i.e.* redness, heat, swelling and/or pain resulting

from infection, injury, tissue damage, irritation etc.). Examples of inflammatory diseases include, but are not limit to autoimmune diseases (*e.g.* inflammatory bowel disease, rheumatoid arthritis and celiac disease), asthma, vasculitis, chronic prostatitis, pelvic inflammatory disease, cancer, atherosclerosis, ischemic heart disease, reperfusion injury, transplant rejection, sarcoidosis, allergic
 5 hypersensitivity and glomerulonephritis.

Mitogen-activated protein kinase (MAPK) pathway: A cellular signaling pathway that regulates cellular processes such as proliferation, survival, and migration. Mammals express at least four distinctly regulated groups of MAPKs, extracellular signal-related kinases (ERK)-1/2, Jun amino-terminal kinases (JNK1/2/3), p38 proteins (p38alpha/beta/gamma/delta) and ERK5,
 10 which are activated by specific MAPKKs: MEK1/2 for ERK1/2, MKK3/6 for the p38, MKK4/7 (JNKK1/2) for the JNKs, and MEK5 for ERK5. Each MAPKK can be activated by more than one MAPKKK, increasing the complexity and diversity of MAPK signaling. Each MAPK pathway contains a three-tiered kinase cascade comprising a MAP kinase kinase kinase (MAPKKK, MAP3K, MEKK or MKKK), a MAP kinase kinase (MAPKK, MAP2K, MEK or MKK) and the
 15 MAPK. This three-tier module mediates ultrasensitive switch-like responses to stimuli.

MAPK family: A family of serine/threonine protein kinases involved in directing cellular responses to a diverse array of stimuli, including mitogens, osmotic stress, heat shock and proinflammatory cytokines. MAPKs regulate, for example, proliferation, gene expression, differentiation, mitosis, cell survival and apoptosis. Members of the MAPK/ERK family include,
 20 but are not limited to MAPK1 (ERK2), MAPK3 (ERK1), MAPK4 (ERK4), MAPK6 (ERK3), MAPK7 (ERK5), MAPK8 (JNK1), MAPK9 (JNK2), MAPK10 (JNK3), MAPK11 (p38- β), MAPK12 (p38- γ), MAPK13 (p38- δ) and MAPK14 (p38- α).

Metastasis: Refers to the spread of cancer cells from the original tumor to other sites in the body.

Multiple sclerosis: An autoimmune disease classically described as a central nervous system white matter disorder disseminated in time and space that presents as relapsing-remitting illness in 80-85% of patients. Diagnosis can be made by brain and spinal cord magnetic resonance imaging (MRI), analysis of somatosensory evoked potentials, and analysis of cerebrospinal fluid to detect increased amounts of immunoglobulin or oligoclonal bands. MRI is a particularly sensitive
 25 diagnostic tool. MRI abnormalities indicating the presence or progression of MS include hyperintense white matter signals on T2-weighted and fluid attenuated inversion recovery images, gadolinium enhancement of active lesions, hypointensive “black holes” (representing gliosis and axonal pathology), and brain atrophy on T1-weighted studies. Serial MRI studies can be used to
 30

indicate disease progression. **Relapsing-remitting multiple sclerosis (RRMS)** is a clinical course of MS that is characterized by clearly defined, acute attacks with full or partial recovery and no disease progression between attacks. **Secondary-progressive multiple sclerosis (SPMS)** is a clinical course of MS that initially is relapsing-remitting, and then becomes progressive at a variable rate, possibly with an occasional relapse and minor remission. **Primary progressive multiple sclerosis (PPMS)** presents initially in the progressive form.

Neoplasia, malignancy, cancer or tumor: A neoplasm is an abnormal growth of tissue or cells that results from excessive cell division. Neoplastic growth can produce a tumor. The amount of a tumor in an individual is the “tumor burden” which can be measured as the number, volume, or weight of the tumor. A tumor that does not metastasize is referred to as “benign.” A tumor that invades the surrounding tissue and/or can metastasize is referred to as “malignant.”

p38 mitogen-activated protein kinases: A class of MAPKs that are responsive to cellular stresses, such as inflammatory cytokines, ultraviolet radiation, lipopolysaccharide, heat shock, osmotic shock and growth factors. p38 proteins are involved in a number of cellular processes, including cell differentiation, apoptosis and autophagy. Four p38 MAPKs have been identified – p38- α (MAPK14), p38- β (MAPK11), p38- γ (MAPK12) and p38- δ (MAPK13); of these, p38 α , p38 β and p38 δ are expressed in T cells. As used herein, “p38” refers to any of the p38 proteins (p38 α , p38 β , p38 γ and p38 δ). In particular embodiments, “**p38**” refers to any p38 protein expressed in T cells, including p38 α , p38 β and p38 δ .

Pancreatic cancer: A disease in which malignant (cancer) cells are found in the tissues of the pancreas. Also called exocrine cancer.

Pancreatic ductal adenocarcinoma (PDAC): The most common type of pancreatic malignancy. PDAC is the fourth leading cause of cancer death in the United States with a mean survival of less than six months. This type of cancer is capable of rapid dissemination to the lymphatic system and distant organs. PDAC is distinct from other cancers due to the biological barrier the tumor builds around itself. Patients whose disease is caught at an early stage have a better chance of long-term survival, but the pancreas emits few known clues to signal that the carcinogenic process has begun. Therefore, there are currently no early detection tests.

Pharmaceutically acceptable carriers: The pharmaceutically acceptable carriers useful in the methods disclosed herein are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, PA, 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of polypeptides.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (e.g., powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, salts, amino acids, and pH buffering agents and the like, for example sodium or potassium chloride or phosphate, Tween, sodium acetate or sorbitan monolaurate.

Poly-arginine or poly-lysine: A polymer of multiple arginine or lysine residues, respectively. In the context of the present disclosure, “poly-arginine” or “poly-lysine” includes peptides made up of about 4 to about 20 arginine or lysine residues, such as about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 arginine or lysine residues, respectively. In particular embodiments herein, the poly-arginine peptide includes 6 to 12 arginine residues, or the poly-lysine peptide includes 6 to 12 lysine residues. In one non-limiting example, the poly-arginine peptide consists of 11 arginine (11R) residues.

Polypeptide: A polymer in which the monomers are amino acid residues which are joined together through amide bonds. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used. The terms “polypeptide” or “protein” as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term “polypeptide” is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically produced. The term “residue” or “amino acid residue” includes reference to an amino acid that is incorporated into a protein, polypeptide, or peptide.

A conservative substitution in a polypeptide is a substitution of one amino acid residue in a protein sequence for a different amino acid residue having similar biochemical properties. Typically, conservative substitutions have little to no impact on the activity of a resulting polypeptide. For example, a protein or peptide including one or more conservative substitutions (for example no more than 1, 2, 3, 4 or 5 substitutions) retains the structure and function of the wild-type protein or peptide. A polypeptide can be produced to contain one or more conservative substitutions by manipulating the nucleotide sequence that encodes that polypeptide using, for example, standard procedures such as site-directed mutagenesis or PCR. In one example, such

variants can be readily selected by testing antibody cross-reactivity or its ability to induce an immune response. Examples of conservative substitutions are shown below.

	Original Residue	Conservative Substitutions
5	Ala	Ser
	Arg	Lys
	Asn	Gln, His
	Asp	Glu
	Cys	Ser
10	Gln	Asn
	Glu	Asp
	His	Asn; Gln
	Ile	Leu, Val
	Leu	Ile; Val
15	Lys	Arg; Gln; Glu
	Met	Leu; Ile
	Phe	Met; Leu; Tyr
	Ser	Thr
	Thr	Ser
20	Trp	Tyr
	Tyr	Trp; Phe
	Val	Ile; Leu

Conservative substitutions generally maintain (a) the structure of the polypeptide backbone
 25 in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

The substitutions which in general are expected to produce the greatest changes in protein properties will be non-conservative, for instance changes in which (a) a hydrophilic residue, for example, seryl or threonyl, is substituted for (or by) a hydrophobic residue, for example, leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other
 30 residue; (c) a residue having an electropositive side chain, for example, lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, for example, glutamyl or aspartyl; or (d) a residue having a bulky side chain, for example, phenylalanine, is substituted for (or by) one not having a side chain, for example, glycine.

35 **Preventing, treating or ameliorating a disease:** “Preventing” a disease refers to inhibiting the full development of a disease. “Treating” refers to a therapeutic intervention that ameliorates a sign or symptom of a disease or pathological condition after it has begun to develop. “Ameliorating” refers to the reduction in the number or severity of signs or symptoms of a disease, such as a T-cell and/or p38-mediated disease or disorder.

Prognosis: The likelihood of the clinical outcome for a subject afflicted with a specific disease or disorder, such as pancreatic cancer. With regard to cancer, the prognosis is a representation of the likelihood (probability) that the subject will survive (such as for one, two, three, four or five years) and/or respond to anti-cancer treatment, and/or the likelihood (probability) that the tumor will metastasize. In some embodiments, a “poor prognosis” indicates a greater than 50% chance that the subject will not survive to a specified time point (such as one, two, three, four or five years), and/or a greater than 50% chance that the tumor will metastasize. In several examples, a poor prognosis indicates that there is a greater than 60%, 70%, 80%, or 90% chance that the subject will not survive and/or a greater than 60%, 70%, 80% or 90% chance that the tumor will metastasize. Conversely, a “good prognosis” indicates a greater than 50% chance that the subject will survive to a specified time point (such as one, two, three, four or five years), and/or a greater than 50% chance that the tumor will not metastasize. In several examples, a good prognosis indicates that there is a greater than 60%, 70%, 80%, or 90% chance that the subject will survive and/or a greater than 60%, 70%, 80% or 90% chance that the tumor will not metastasize.

Purified: The term purified does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified peptide preparation is one in which the peptide or protein is more enriched than the peptide or protein is in its natural environment within a cell. In one embodiment, a preparation is purified such that the protein or peptide represents at least 50% of the total peptide or protein content of the preparation. Substantial purification denotes purification from other proteins or cellular components. A substantially purified protein is at least 60%, 70%, 80%, 90%, 95% or 98% pure. Thus, in one specific, non-limiting example, a substantially purified protein is 90% free of other proteins or cellular components.

Recombinant: A recombinant nucleic acid or protein is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques. The term recombinant includes nucleic acids and proteins that have been altered solely by addition, substitution, or deletion of a portion of a natural nucleic acid molecule or protein.

Rheumatoid arthritis: A systemic autoimmune disorder characterized by chronic inflammation of the joints. This disease can also include inflammation of tissues in other areas of the body, such as the lungs, heart and eyes.

Sample (or biological sample): A biological specimen containing genomic DNA, RNA, protein, or combinations thereof, obtained from a subject. Examples include, but are not limited to, peripheral blood, urine, saliva, tissue biopsy (such as pancreatic tissue), surgical specimen, and autopsy material. In one example, a sample includes a biopsy of a pancreatic tumor or a sample of
5 normal tissue, such as pancreatic tissue (from a subject not afflicted with a known disease or disorder, such as a cancer-free subject).

Sequence identity/similarity: The identity/similarity between two or more nucleic acid sequences, or two or more amino acid sequences, is expressed in terms of the identity or similarity between the sequences. Sequence identity can be measured in terms of percentage identity; the
10 higher the percentage, the more identical the sequences are. Sequence similarity can be measured in terms of percentage similarity (which takes into account conservative amino acid substitutions); the higher the percentage, the more similar the sequences are. Homologs or orthologs of nucleic acid or amino acid sequences possess a relatively high degree of sequence identity/similarity when aligned using standard methods.

15 Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith & Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman & Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444, 1988; Higgins & Sharp, *Gene*, 73:237-44, 1988; Higgins & Sharp, *CABIOS* 5:151-3, 1989; Corpet *et al.*, *Nuc. Acids Res.* 16:10881-90, 1988; Huang *et al. Computer Appls. in the*
20 *Biosciences* 8, 155-65, 1992; and Pearson *et al.*, *Meth. Mol. Bio.* 24:307-31, 1994. Altschul *et al.*, *J. Mol. Biol.* 215:403-10, 1990, presents a detailed consideration of sequence alignment methods and homology calculations.

The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul *et al.*, *J. Mol. Biol.* 215:403-10, 1990) is available from several sources, including the National Center for Biological
25 Information (NCBI, National Library of Medicine, Building 38A, Room 8N805, Bethesda, MD 20894) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. Additional information can be found at the NCBI web site.

BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. If the two compared sequences share homology, then the designated output
30 file will present those regions of homology as aligned sequences. If the two compared sequences do not share homology, then the designated output file will not present aligned sequences.

Subject: A human or non-human animal. In one embodiment, the subject has a T cell- or p38-mediated disorder.

Synthetic: Produced by artificial means in a laboratory, for example a synthetic polypeptide can be chemically synthesized in a laboratory.

T cell- or p38-mediated disorder: Any disease, disorder or condition that is associated with aberrant or pathological activation of T cells and/or the alternative p38 activation pathway in T cells. T cell- or p38-mediated disorders include, for example, autoimmune disorders (such as, but not limited to multiple sclerosis, rheumatoid arthritis and inflammatory bowel disease) and inflammatory disorders, including tumors that are associated with local inflammatory responses (*e.g.* pancreatic cancer).

T cell receptor (TCR): A molecule expressed on the surface of T lymphocytes that is responsible for recognizing antigens bound to major histocompatibility complex (MHC) molecules. The TCR is a heterodimer consisting of either an alpha (α) and beta (β) chains, or gamma (γ) and delta (δ) chains. In mammals, the TCR is expressed as part of a complex with four CD3 chains (γ , δ , and 2 ϵ chains) and the ζ chain.

T helper (Th) cells: A type of T cell that activates or directs other immune cells but has no intrinsic cytotoxic or phagocytic activity. Th cells play essential roles in B cell antibody class switching, activation and growth of cytotoxic T cells and bactericidal activity of phagocytic cells.

Th1 and Th2 cells are subsets of Th cells that primarily interact with macrophages and B cells, respectively. Th1 cells produce interferon (IFN)- γ , interleukin (IL)-2, lymphotoxin (LT)- α and transforming growth factor (TGF)- β , and function to maximize the killing efficiency of

macrophages and the proliferation of cytotoxic CD8⁺ T cells. Th2 cells produce IL-4, IL-5, IL-6, IL-13 and IL-25, and function by stimulating B cell proliferation, inducing B cell class switching and increasing neutralizing antibody production. **Th17 cells** are another subset of T helper cells that produce the effector cytokines IL-17, IL-21 and IL-22. Th17 cells are considered

developmentally distinct from Th1 and Th2 cells. Excessive numbers of Th17 cells are thought to play a role in several autoimmune diseases. TGF- β , IL-6, IL-21, and IL-23 have been associated with the differentiation of T cells to Th17 cells.

Therapeutically effective amount: A dose sufficient to prevent advancement of a disease, or to cause regression of the disease, or which is capable of reducing symptoms caused by the disease, such as a T cell- or p38-mediated disorder (*e.g.*, an autoimmune disease, such as multiple sclerosis or rheumatoid arthritis, or inflammatory disorder, such as cancer).

Tumor vascularization: The formation of new blood vessels and/or growth of existing blood vessels in a tumor.

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. “Comprising A or B” means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

III. Gadd45 α Polypeptides, Fusion Proteins and Compositions

Disclosed herein is the identification of a minimal region of the growth arrest and DNA damage-inducible alpha (Gadd45 α) protein that is required for binding to p38 and inhibiting the alternative p38 activation pathway in T cells. In particular, the Gadd45 α polypeptides disclosed herein are capable of inhibiting the enzymatic activity of tyrosine 323-phosphorylated p38, inhibiting T cell proliferation in response to TCR activation (but not in response to stimuli that activate the classical p38 activation pathway), inhibiting skewing of T cells to Th1 and Th17 cells and reducing proinflammatory cytokine production. Moreover, administration of the disclosed Gadd45 α polypeptides inhibits activation of the T-cell specific alternative p38 activation pathway *in vivo* and significantly reduces tumor formation and growth of inflammatory pancreatic tumors.

Provided herein are Gadd45 α polypeptides capable of inhibiting the T-cell specific alternative p38 activation pathway. In some embodiments, the Gadd45 α polypeptide is no more than 25 amino acids in length, or no more than 20 amino acids in length, and comprises residues 71-85 of Gadd45 α . In some embodiments, the Gadd45 α is human or mouse Gadd45 α . In some examples, the amino acid sequence of the Gadd45 α polypeptide comprises LQIHFTLIXAFCCEN (SEQ ID NO: 1), wherein X is Q or R. In particular examples, the Gadd45 α polypeptide comprises or consists of the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3). In some instances, the Gadd45 α polypeptide contains 1 to 5, such as 1, 2, 3, 4 or 5 conservative amino acids substitutions in the polypeptide sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3). Conservative

amino acid substitutions are well known in the art and exemplary substitutions are described herein (see Terms and Methods). In some embodiments, the Gadd45 α polypeptide is a synthetic or recombinant (*i.e.* non-naturally occurring) polypeptide.

Also provided herein are fusion proteins comprising a Gadd45 α polypeptide and a heterologous protein or peptide. In some embodiments, the heterologous peptide is a peptide that promotes cellular uptake of the fusion protein, such as a cell-penetrating peptide (CPP).

CPPs are a family of polypeptides that facilitate transduction of proteins, nucleic acids or other compounds across membranes in a receptor-independent manner (Wadia and Dowdy, *Curr. Protein Pept. Sci.* 4(2):97-104, 2003). Typically, CPPs are short polycationic sequences that can facilitate cellular uptake of compounds to which they are linked into endosomes of cells.

The capacity of certain peptides to deliver proteins or nucleic acids into cells was originally described for the HIV-encoded Tat protein, which was shown to cross membranes and initiate transcription. It was then discovered that the portion of the Tat protein that was required for the transduction of the protein was only an 11 amino acid polypeptide, referred to as the Tat peptide.

When fused with other proteins, the Tat peptide has been demonstrated to deliver these proteins, varying in size from 15 to 120 kDa, into cells in tissue culture (Frankel and Pabo, *Cell* 55(6):1189-93, 1988; Green and Loewenstein, *J. Gen. Microbiol.* 134(3):849-55, 1988; Vives *et al.*, *J. Biol. Chem.* 272(25):16010-7, 1997; Yoon *et al.*, *J. Microbiol.* 42(4):328-35, 2004; Cai *et al.*, *Eur. J. Pharm. Sci.* 27(4):311-9, 2006).

Other known CPPs include PENETRATINTM, a 16 amino acid peptide derived from the third helix of the *Drosophila* Antennapedia homeobox gene (U.S. Patent No. 5,888,762; Derossi *et al.*, *J. Biol. Chem.* 269:10444-10450, 1994; Schwarze *et al.*, *Trends Pharmacol. Sci.* 21:45-48, 2000); transportan, a 27 amino acid chimeric peptide comprised of 12 amino acids from the N-terminus of the neuropeptide galanin and the 14-amino acid protein mastoparan, connected via a lysine (U.S. Patent No. 6,821,948; Pooga, *FASEB J.* 12:67-77, 1998; Hawiger, *Curr. Opin. Chem. Biol.* 3:89-94, 1999); peptides from the VP22 protein of herpes simplex virus (HSV) type 1 (Elliott *et al.*, *Cell* 88:223-233, 1997); the UL-56 protein of HSV-2 (U.S. Pre-Grant Publication No. 2006/0099677); and the Vpr protein of HIV-1 (U.S. Pre-Grant Publication No. 2005/0287648). In addition, a number of artificial peptides also are known to function as CPPs, such as poly-arginine, poly-lysine and others (see, for example, U.S. Application Publication Nos. 2006/0106197; 2006/0024331; 2005/0287648; and 2003/0125242; Zhibao *et al.*, *Mol. Ther.* 2:339-347, 2000; and Laus *et al.* *Nature Biotechnol.* 18:1269-1272, 2000).

In some examples, the CPP is rich in charged amino acids, such as lysine or arginine. In other examples, the CPP contains an alternating pattern of polar/charged amino acids and non-polar/hydrophobic amino acids. In particular non-limiting examples, the CPP comprises poly-arginine, such as 6, 7, 8, 9, 10, 11 or 12 arginine residues. In one example, poly-arginine comprises or consists of 11 arginine (11R) residues. In other non-limiting examples, the CPP comprises poly-lysine, such as 6, 7, 8, 9, 10, 11 or 12 lysine residues.

In other embodiments, the heterologous protein or peptide is a protein tag, such as an affinity tag (for example, chitin binding protein, maltose binding protein, glutathione-S-transferase or poly-His), an epitope tag (for example, V5, c-myc, HA or FLAG) or a fluorescent tag (*e.g.*, GFP or another well-known fluorescent protein).

In one embodiment, the fusion protein comprises 11R fused to the N-terminus of a Gadd45 α polypeptide. In some examples, the fusion protein comprises or consists of 11R fused to the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).

Further provided herein are compositions comprising a Gadd45 α polypeptide or Gadd45 α fusion protein disclosed herein. In some embodiments, the composition further comprises a pharmaceutically acceptable carrier.

IV. Use of Gadd45 α Polypeptides

It is disclosed herein that Gadd45 α polypeptides comprising a minimal region that interacts with p38 (Gadd45 α residues 71-85) are capable of inhibiting activation of the T-cell specific alternative p38 activation pathway *in vitro*, *ex vivo* and *in vivo*. In particular, the Gadd45 α polypeptides disclosed herein inhibit T cell proliferation in response to TCR activation, inhibit skewing of T cells to Th1 and Th17 cells and reduce proinflammatory cytokine production. It is also demonstrated herein that administration of the disclosed Gadd45 α polypeptides significantly reduces tumor formation and growth of inflammatory pancreatic tumors. Furthermore, data disclosed herein demonstrates that detection of tumor-infiltrating T cells positive for tyrosine 323-phosphorylated p38 can be used to determine the prognosis of a patient diagnosed with pancreatic ductal adenocarcinoma (PDAC).

Provided herein is a method of inhibiting proliferation of T cells, such as inhibiting proliferation of T cells in response to TCR stimulation. In some embodiments, the method includes contacting the T cells with a Gadd45 α polypeptide, fusion protein or composition disclosed herein, thereby inhibiting proliferation of the T cells. In some examples, the method of inhibiting

proliferation of T cells is an *ex vivo* method in which T cells have been isolated from a subject and the isolated T cells are contacted with the Gadd45 α polypeptide, fusion protein or composition. In other examples, the method of inhibiting proliferation of T cells is an *in vivo* method that includes administering the Gadd45 α polypeptide, fusion protein or composition to the subject. In particular
5 examples, the subject exhibits aberrant proliferation of T cells and/or aberrant activation of the alternative p38 pathway. For example, the subject can be a subject suffering from an autoimmune or inflammatory disease or disorder, such as multiple sclerosis, rheumatoid arthritis or cancer, such as pancreatic cancer (*e.g.* PDAC).

In some examples, T cell proliferation is inhibited at least 10%, at least 20%, at least 30%,
10 at least 40%, at least 50%, at least 60%, at least 70% or at least 80% compared to a control (*e.g.* untreated T cells). Methods for detecting and quantifying proliferation of T cells is well known in the art, including, for example, measuring [³H]-thymidine incorporation or measuring T cell populations using fluorescence activated cell sorting based on T cell-specific markers.

Further provided is a method of inhibiting differentiation of T cells into Th1 or Th17 cells.

15 In some embodiments, the method includes contacting the T cells with a Gadd45 α polypeptide, fusion protein or composition disclosed herein, thereby inhibiting differentiation of the T cells. In some examples, the method of inhibiting differentiation of T cells is an *ex vivo* method in which T cells have been isolated from a subject and the isolated T cells are contacted with the Gadd45 α polypeptide, fusion protein or composition. In other examples, the method of inhibiting
20 differentiation of T cells is an *in vivo* method that includes administering the Gadd45 α polypeptide, fusion protein or composition to a subject. In particular examples, the subject is a subject having a T cell- and/or p38-mediated autoimmune or inflammatory disorder, such as multiple sclerosis, rheumatoid arthritis or cancer (*e.g.* pancreatic cancer).

In some examples, differentiation of T cells into Th1 or Th17 cells is inhibited at least 10%,
25 at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70% or at least 80% relative to a control (*e.g.* untreated T cells). Methods for evaluating differentiation of T cells into particular T helper (Th) subtypes are well known in the art, including the methods described in the Examples. For instance, Th subtypes can be determined by evaluating cytokine expression profiles or subtype-specific transcription factor expression. Th1 cells are characterized by expression of the
30 cytokines including IFN- γ , IL-2 and LT- α , and the transcription factor T-bet; Th2 cells typically produce IL-4, IL-5, IL6 and IL-13; and Th17 cells primarily express IL-17, IL-21 and IL-22 (Zhu and Paul, *Blood* 112(5):1557-1568, 2008).

Also provided herein is a method of treating a subject having a T cell- and/or p38-mediated autoimmune or inflammatory disorder. In some embodiments, the method includes selecting a subject with a T cell- and/or p38-mediated autoimmune or inflammatory disorder and administering to the subject a therapeutically effective amount of a Gadd45 α polypeptide, fusion protein or composition disclosed herein, thereby treating the subject. The T cell- and/or p38-mediated autoimmune or inflammatory disorder is any disease, disorder or condition that is associated with aberrant or pathological activation of T cells and/or the alternative p38 activation pathway in T cells. In some embodiments, the T cell- and/or p38-mediated disorder is an autoimmune disorder, such as, but not limited to amyotrophic lateral sclerosis, autoimmune hepatitis, celiac disease, Crohn's disease (inflammatory bowel disease), diabetes type 1, Graves' disease, Guillain-Barre syndrome, Hashimoto's disease, multiple sclerosis, myasthenia gravis, psoriasis, rheumatoid arthritis, Sjögren's syndrome, systemic lupus erythematosus, and ulcerative colitis. In other embodiments, the T cell- and/or p38-mediated disorder is an inflammatory disorder, such as but not limited to asthma, vasculitis, chronic prostatitis, pelvic inflammatory disease, cancer, atherosclerosis, ischemic heart disease, reperfusion injury, transplant rejection, sarcoidosis, allergic hypersensitivity and glomerulonephritis. In some examples, the T cell- and/or p38-mediated autoimmune or inflammatory disorder comprises multiple sclerosis, rheumatoid arthritis, inflammatory bowel disease or pancreatic cancer (such as, but not limited to, PDAC).

Further provided herein is a method of inhibiting tumor vascularization in a subject. In some embodiments, the method includes selecting a subject with a tumor and administering to the subject a therapeutically effective amount of a Gadd45 α polypeptide, fusion protein or composition disclosed herein. In some examples, the tumor is a pancreatic tumor, such as a PDAC tumor. In some examples, administration comprises intra-tumor injection. In non-limiting examples, tumor vascularization is inhibited at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70% or at least 80% relative to a control. Methods of detecting and evaluating tumor vascularization are known in the art. For example, tumor vascularization can be evaluated by staining tumor cells for an endothelial marker, such as CD31 or CD34.

Methods of administering therapeutic proteins are well known in the art, and include both systemic and local administration. In some embodiments, the Gadd45 α polypeptide, fusion protein or composition is administered intravenously or intraperitoneally. In some examples in which the subject has a tumor, the Gadd45 α polypeptide, fusion protein or composition is administered directly into the tumor. Single or multiple administrations (such as 2, 3, 4, 5 or more

administrations) can be utilized depending on, for example, the type and severity of disease to be treated, and the general health of the subject.

The amount of a Gadd45 α polypeptide, fusion protein or composition administered to a subject is generally an amount sufficient to at least partially inhibit activation of p38 and/or treat or ameliorate one or more signs or symptoms of disease. A therapeutically effective amount of a Gadd45 α polypeptide, fusion protein or composition can be determined by a medical practitioner. In some embodiments, the therapeutically effective amount of a Gadd45 α polypeptide, fusion protein or composition disclosed herein is an amount that provides either subjective relief of one or more symptoms or an objectively identifiable improvement as noted by the clinician or other qualified observer. In some examples, the Gadd45 α polypeptide or fusion protein is administered at a dose of about 20 to about 100 μ g, such as about 20, about 25, about 30, about 35, about 40, about 45, about 50, about 55, about 60, about 65, about 70, about 80, about 90 or about 100 μ g.

Also provided herein is a method of determining the prognosis of a subject diagnosed with PDAC by calculating the percentage of tumor-infiltrating T cells positive for tyrosine 323-phosphorylated p38 (p38 pY323) in a tumor sample obtained from the subject; and determining that the subject has a poor prognosis if at least 10% of the T cells are positive for p38 pY323.

A poor prognosis refers to any negative clinical outcome. For example, in some embodiments, a poor prognosis is a decrease in the time of survival. In some embodiments, a poor prognosis is an increase in the likelihood of metastasis of the cancer. In other embodiments, a poor prognosis refers to failure to respond to therapy, such as radiation therapy or chemotherapy. In particular embodiments, the poor prognosis is a decrease in time of survival. In some examples, a poor prognosis indicates that the subject is likely to survive less than one year.

Methods for detecting tyrosine 323-phosphorylated p38 in a tumor sample are well known in the art, and include, for example, immunohistochemical detection using an antibody specific for tyrosine 323-phosphorylated p38 (see Example 5 below).

V. Additional Embodiments

1. An isolated growth arrest and DNA-damage-inducible alpha (Gadd45 α) polypeptide no more than 25 amino acids in length, wherein the amino acid sequence of the polypeptide comprises LQIHFTLIXAFCCEN (SEQ ID NO: 1), wherein X is Q or R.

2. The isolated Gadd45 α polypeptide of 1 which is no more than 20 amino acids in length.

3. The isolated Gadd45 α polypeptide of 1 or 2, comprising the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).

4. The isolated Gadd45 α polypeptide of 1 or 2, consisting of the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).

5. A fusion protein comprising the Gadd45 α polypeptide of any one of 1-4 and a heterologous peptide.

6. The fusion protein of 5, wherein the heterologous peptide is a cell-penetrating peptide (CPP).

7. The fusion protein of 6, wherein the CPP comprises poly-arginine.

8. The fusion protein of 7, wherein poly-arginine comprises or consists of 11 arginine (11R) residues.

9. The fusion protein of 8, consisting of 11R fused to the N-terminus of the Gadd45 α polypeptide.

10. The fusion protein of 8, consisting of 11R fused to the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).

11. A composition comprising the polypeptide or fusion protein of any one of 1-10 and a pharmaceutically acceptable carrier.

12. An *in vitro* or *in vivo* method of inhibiting proliferation of T cells in response to T cell receptor stimulation, or of inhibiting differentiation of T cells into Th1 or Th17 cells, comprising contacting the T cells with the polypeptide, fusion protein or composition of any one of 1-11, thereby inhibiting proliferation of the T cells.

13. The method of 12 which is an *ex vivo* method, wherein the T cells have been isolated from a subject and the isolated T cells are contacted with the polypeptide or fusion protein.

14. A method of inhibiting proliferation of T cells in response to T cell receptor stimulation, or of inhibiting differentiation of T cells into Th1 or Th17 cells, in a subject comprising administering the polypeptide, fusion protein or composition of any one of 1-11 to the subject.

15. A method of treating a subject with a T cell- and/or p38-mediated autoimmune or inflammatory disorder, comprising selecting a subject with a T cell- and/or p38-mediated autoimmune or inflammatory disorder and administering to the subject a therapeutically effective amount of the polypeptide, fusion protein or composition of any one of 1-11, thereby treating the subject.

16. The method of 16, wherein the T cell- and/or p38-mediated autoimmune or inflammatory disorder comprises multiple sclerosis, rheumatoid arthritis or pancreatic cancer.

17. The method of 15 or 16, wherein administration comprises intravenous, intra-tumor or intraperitoneal administration.

5 18. The method of 17, wherein the T cell- and/or p38-mediated autoimmune or inflammatory disorder comprises pancreatic cancer and administration comprises intra-tumor injection.

10 19. A method of inhibiting tumor vascularization in a subject, comprising selecting a subject with a tumor and administering to the subject a therapeutically effective amount of the polypeptide, fusion protein or composition of any one of 1-11, thereby inhibiting tumor vascularization in the subject.

20. The method of 19, wherein the tumor is a pancreatic tumor.

21. The method of 19 or 20, wherein administration comprises intra-tumor administration.

15 22. The method of 19, wherein the polypeptide or fusion protein is administered at a dose of about 20 to about 100 µg.

23. The method of 14, wherein the polypeptide or fusion protein is administered at a dose of about 20 to about 100 µg.

20 24. The method of 15, wherein the polypeptide or fusion protein is administered at a dose of about 20 to about 100 µg.

25 25. A method of determining the prognosis of a subject diagnosed with pancreatic ductal adenocarcinoma (PDAC), comprising:

calculating the percentage of tumor-infiltrating T cells positive for tyrosine 323-phosphorylated p38 (p38 pY323) in a tumor sample obtained from the subject; and

25 determining that the subject has a poor prognosis if at least 10% of the T cells are positive for p38 pY323.

26. The method of 25, wherein the poor prognosis is a decrease in time of survival.

30 The following examples are provided to illustrate certain particular features and/or embodiments. These examples should not be construed to limit the disclosure to the particular features or embodiments described.

EXAMPLES

Example 1: *In vitro* inhibition of the alternative p38 activation pathway by a Gadd45 α polypeptide

This example describes a Gadd45 α 71-85 polypeptide fused to a cell-penetrating peptide (11R) and its ability to block the T cell-specific alternative p38 activation pathway. The minimal region of Gadd45 α capable of binding to p38 was identified by fusing various truncated portions of Gadd45 α to GST and using the product to pull down recombinant p38 α *in vitro*. The 71-85 fragment pulled down p38 α as well as the full length protein. Furthermore, deletion of 71-80 from the full length Gadd45 α rendered it incapable of pulling down p38 α .

(11R)-71-85

To generate a cell-permeable form of the Gadd45 α 71-85 peptide, 11 arginine residues were added to the N-terminus to produce a fusion protein referred to herein as (11R)-71-85. Cell-permeability of (11R)-71-85 was tested using purified CD4⁺ T cells. (11R)-71-85 and a control peptide fused to 11 arginine residues (referred to as (11R)-Ctrl) were FITC conjugated. Purified CD4⁺ T cells were treated with either FITC conjugated (11R)-71-85 or (11R)-Ctrl, or left untreated for 2 hours at 37°C. After incubation, cells were washed and analyzed by flow cytometry for FITC positivity. As shown in FIG. 1, both 11R-conjugated peptides were taken up by the T cells.

In vitro inhibition of the p38 alternative pathway

To test whether (11R)-71-85 can inhibit the alternative p38 activation pathway *in vitro*, an assay was performed to detect phosphorylation of activating transcription factor 2 (ATF2), a target of activated p38, in the presence and absence of (11R)-71-85. Purified p38 proteins were activated by incubation with active Zap70 in kinase buffer for 1 hour. (11R)-71-85 (0.2, 0.5, 1.0, 2.0 or 4.0 μ g) or (11R)-Ctrl (4.0 μ g) was then added and incubated for 30 minutes at 30°C. One μ g ATF2 and 10 μ Ci [³²P]ATP were added and incubated for 45 minutes at 30°C. Phosphorylated products were resolved by SDS-PAGE, transferred to nitrocellulose membranes, and visualized by phosphorimager. As shown in FIG. 2, incubation with (11R)-71-85, but not (11R)-Ctrl, led to a decrease in phosphorylation of ATF2. In addition, the decrease in ATF2 phosphorylation was dependent on the dose of (11R)-71-85. These results demonstrate that (11R)-71-85 is capable of inhibiting the alternative p38 activation pathway.

Inhibition of T cell proliferation and inflammatory cytokine production

The ability of (11R)-71-85 to inhibit T cell proliferation and inflammatory cytokine production was also evaluated. T cells were purified from spleen and treated with different concentrations (0.2, 0.5, 1.0, 2.0 or 4.0 μ g) of (11R)-71-85 or the control peptide (11R)-Ctrl. Cells were then washed and stimulated with anti-CD3 and anti-CD28 for 2 days and finally pulsed with 1 μ Ci [3 H]-thymidine 14 hours before harvesting. [3 H]-thymidine uptake was determined with a Wallac 1450 MicroBeta Liquid Scintillation Counter. As shown in FIG. 3, treatment with (11R)-71-85, but not (11R)-Ctrl, led to a dose-dependent decrease in T cell proliferation in response to T cell receptor (TCR) stimulation.

To evaluate cytokine production, T cells were purified from spleen and stimulated with anti-CD3 plus anti-CD28 for 3 days. Cells were then left unstimulated overnight and treated with 1 or 2 μ M (11R)-71-85 or control peptide (11R)-Ctrl, or with DMSO only. After treatment, cells were restimulated with either anti-CD3 plus anti-CD28 (to stimulate the TCR and the alternative p38 pathway), or PMA plus ionomycin supplemented with monensin (to stimulate the classical p38 pathway). Cytokine expression was measured by flow-cytometry. As shown in FIGS. 4A-4B, treatment with (11R)-71-85 inhibits the production of TNF- α in response to TCR stimulation, but has no significant effect on cytokine production in response to stimulation with PMA plus ionomycin. The effect of (11R)-71-85 is dose-dependent as treatment with 2 μ M (11R)-71-85 inhibits production of TNF- α and IFN- γ to a greater extent than treatment with 1 μ M (11R)-71-85.

Inhibition of Th1 and Th17 differentiation

Additional studies were carried out to evaluate the effect of (11R)-71-85 on T helper (Th) cell differentiation. Expression of IFN- γ and T-bet was used as an indicator of Th1 cells typical of a pro-inflammatory response. Naïve CD4 $^+$ T cells were treated with either medium or 2 μ M of (11R)-71-85 or (11R)-Ctrl for two hours and after washing, cells were stimulated with anti-CD3 plus anti-CD28 in Th0 (neutral) or Th1 polarizing conditions for 3 days. Cells were treated again with the same concentration of either (11R)-71-85 or (11R)-Ctrl for 2 hours and restimulated with PMA plus ionomycin in the presence of Golgi inhibitor for 4 hours. Expression of T bet and IFN γ was analyzed by flow cytometry (FIGS. 5A-5D). Cells treated with (11R)-71-85 exhibited reduced expression of IFN- γ and T-bet, relative to cells treated with medium or (11R)-Ctrl. These results indicate that treatment with (11R)-71-85 inhibits Th1 differentiation.

The effect of full-length Gadd45 α on Th17 differentiation was also evaluated. Purified CD4⁺ T cells from WT mice were treated with 4 μ M of either TAT-Ctrl, TAT-G45FL or TAT-G45 Δ 10 and stimulated with plate coated anti-CD3 plus anti-CD28 for 48 hours and blotted for the transcription factors NFATc1, IRF4 or IRF8 (FIG. 6A). Treatment of CD4⁺ T cells with full-length Gadd45 α inhibited expression of NFATc1 and IRF4, but not IRF8. Similarly, naïve CD4⁺ T (CD4⁺CD62L^{hi}) cells were purified from WT mice and treated with either TAT-Ctrl or TAT-G45FL and stimulated in neutral (Th0) or Th17 skewing conditions (10 ng/ml IL-6 + 5 ng/ml TGF- β) for 48 hours. The mRNA expression levels of Rorc and Il17a were evaluated (FIG. 6B). Expression of full-length Gadd45 α significantly inhibited expression of both Rorc and Il17a under Th17 skewing conditions. These results indicate that full-length Gadd45 α inhibits Th17 differentiation.

Example 2: 11R-71-85 inhibits expression of T cell activation markers *in vivo*

The studies described in Example 1 demonstrate that a Gadd45 α polypeptide comprising residues 71-85 is capable of inhibiting the alternative p38 activation pathway *in vitro* and *ex vivo*. This example describes the finding that a Gadd45 α polypeptide comprising residues 71-85 can also inhibit the alternative p38 activation pathway *in vivo*.

The ability of (11R)-71-85 to inhibit the p38 alternative pathway *in vivo* was evaluated by assessing expression of interferon regulatory factor 4 (Irf4) mRNA, a transcription factor that is up-regulated following TCR stimulation. (11R)-71-85 (40 μ g) or vehicle control was injected intraperitoneally into wild type mice. Four hours after injection, splenocytes were collected and stimulated with soluble anti-CD3 plus anti-CD28 for 24 hours. Irf4 mRNA expression was assessed by quantitative real time PCR. As shown in FIG. 7, treatment with (11R)-71-85 significantly inhibited expression of Irf4 compared to the control.

Expression of NFATc1 and IRF4, two transcription factors activated upon TCR stimulation, was also evaluated after intravenous (i.v.) injection of (11R)-71-85. WT C57BL/6 mice were injected i.v. with 50 μ g of DMSO, (11R)-Scrambled, or (11R)-71-85, and either total T cells or CD4⁺T cells were purified from spleen at different times. Purified cells were stimulated *in vitro* with anti-CD3/CD28 for 24 hours and the mRNA levels of Nfatc1 and Irf4 were determined by quantitative real-time PCR. As shown in FIG. 8, treatment with (11R)-71-85 inhibits expression of both Nfat1c and Irf4 following TCR stimulation in total T cells and in CD4⁺T cells.

Example 3: Administration of 11R-71-85 inhibits growth and vascularization of pancreatic tumors, and inhibits TNF- α secretion from tumor-infiltrating T cells

The p38 MAPK has been implicated in the growth of some tumors that cause local inflammatory responses, such as pancreatic cancer. Therefore, (11R)-71-85 was tested in a pancreatic tumor model. WT C57BL/6 mice were injected with 7.5×10^5 PANC02 cells. After 16 days, mice received intra-tumor injections of 40 μ g or 80 μ g (11R)-71-85 or DMSO every other day. Tumor volume from the start of injection (day 16) to day 42 in control mice and mice injected with (11R)-71-85 or DMSO was determined (FIG. 9A). Weight (FIG. 9B) and appearance (FIG. 9C) of the control and treated tumors were assessed on day 38. Both tumor volume and tumor weight were significantly decreased in mice treated with (11R)-71-85 relative to control animals. These results demonstrate that injection of (11R)-71-85 stops the growth of established PANC02 tumors.

The effect of intra-tumor injection of (11R)-71-85 on cytokine secretion from tumor-infiltrating T cells was also evaluated in mice with pancreatic tumors. WT C57BL/6 mice were injected with 7.5×10^5 PANC02 cells and 16 days later mice received intra-tumor injections of 80 μ g of (11R)-71-85 or (11R)-Scrambled every other day for 1 week. Twenty-four hours following the last injection, tumor infiltrating cells were harvested and restimulated with PMA and ionomycin, or left unstimulated, in the presence of Golgi inhibitor for 4 hours. Expression of TNF- α , IFN γ , IL-10 and IL-17 was determined by flow cytometry. As shown in FIG. 10A, TNF- α secretion was reduced after (11R)-71-85 injection in unstimulated cells, which represent their status *in vivo*. There was no difference in TNF- α secretion after restimulation, which bypasses TCR signaling and activates the classical MAPK cascade. There was no difference between unstimulated and restimulated cells for the other cytokines tested (IFN γ , IL-10, and IL-17; FIGS. 10A-10B).

Tumor vascularization was also assessed in WT mice treated with (11R)-71-85. WT C57BL/6 mice were injected with 7.5×10^5 PANC02 cells. After 16 days, mice received intra-tumor injections of 80 μ g of (11R)-71-85 or (11R)-Scrambled every other day for 2.5 weeks. Twenty-four hours after the last injection, tumors were collected, fixed in formalin, and stained for the endothelial marker CD31, an indicator for tumor vascularization. As shown in FIG. 11, mice treated with (11R)-71-85 exhibited a significant reduction in CD31 staining compared with control treated mice, indicating that administration of (11R)-71-85 inhibits tumor vascularization.

A second murine model of pancreatic cancer was used to evaluate the effect of (11R)-71-85 on cytokine secretion by tumor cells. KPC mice are genetically engineered to develop pancreatic

cancer because of oncogene expression and p53 haploinsufficiency. KPC mice were administered 80 µg of either (11R)-71-85 or (11R)-Scrambled by intraperitoneal injection every other day for 1 month. Twenty-four hours after the last injection, tumor infiltrating cells were harvested and restimulated with PMA and ionomycin, or left unstimulated, in the presence of Golgi inhibitor for 4 hours. Expression of intracellular TNF-α was determined by flow cytometry (FIG. 12). The results demonstrate that administration of (11R)-71-85 inhibits TNF-α secretion by tumor-infiltrating T cells.

Example 4: (11R)-71-85 inhibits progression of PanIN and cancer formation in KPC mice

KPC mice provide a genetic model of murine pancreatic tumors that closely mimics the human disease. KPC (KrasG12D/+;LSLTrp53R172H/+;Pdx-1-Cre) mice express K-ras in pancreatic epithelium, and when made heterozygous for p53, develop spontaneous tumors starting at approximately 8-10 weeks of age (Hingorani *et al.*, *Cancer Cell* 7(5):469-483, 2005). This involves the formation of pancreatic intraepithelial neoplasia precursor lesions (PanIN) and progression to fully-developed carcinoma with surrounding stromal and inflammatory reactions (Hingorani *et al.*, *Cancer Cell* 7(5):469-483, 2005). Histologic examination of pancreatic tissue from 8-13 week old KPC mice revealed PanIN in various stages of development, defined by histological changes of the pancreatic ductal epithelium as well as the formation of a surrounding fibroinflammatory reaction, as well as areas of fully developed invasive carcinoma. As seen in human disease, infiltrating T cells were found in these lesions. Notably, in T-cell-rich regions one could easily detect mononuclear cells that stained with antibodies specific for p38 pY323, the defining phosphorylation site for the alternative pathway, and TNFα, which were absent in T cell-poor regions.

To ask if T cells in which TCR-signaled p38 activation is involved in tumorigenesis, 6 week old KPC mice were sub-lethally irradiated and reconstituted with WT or DKI bone marrow. The former developed invasive cancer and died with a mean time of survival of 37±9 days (FIG. 13A). In contrast, mice that received DKI bone marrow survived much longer, with a mean of 80±16 days (FIG. 13A). Therefore, genetic ablation of the p38 alternative pathway reduced the progression of oncogene-mediated pancreatic cancer.

Given these findings, further experiments were conducted to determine if inhibition of the alternative pathway in KPC mice would alter disease progression. KPC mice were treated with intravenous (11R)-71-85 or (11R)-Scrambled starting at 9 weeks of age, and the pancreas was histologically evaluated 3.5 weeks later. Thirty-three percent (2 of 6) of mice treated with (11R)-

Scrambled developed invasive carcinoma in the pancreas. In contrast, no mice treated with (11R)-71-85 showed signs of invasiveness. Histologic evaluation of H&E stained sections showed a reduced intrapancreatic fibro-inflammatory reaction in (11R)-71-85 treated compared to (11R)-Scrambled treated tumors. (11R)-71-85 not only prevented the formation of invasive carcinoma, but it also slowed the progression towards high grade PanIN formation (PanIN3, carcinoma *in situ*) (FIG. 13B). Similar to the PANC02 model, tumor infiltrating lymphocytes (TIL) from KPC mice treated with (11R)-71-85 had reduced numbers of CD4⁺ cells expressing TNF α compared to control, whereas the expression of IFN γ was slightly reduced (FIG. 13C). In line with the reduced TNF α secretion, *Irf4* mRNA expression was decreased in CD4⁺ TILs after (11R)-71-85 treatment. Consistent with this, expression of *Ccl2*, *Ccl5*, *Vegf*, and *Tgfb* was also reduced in pancreatic epithelial cells of KPC mice (FIG. 13D). The fact that early stage PanIN formation was present, but later pre-invasive stages were reduced, in (11R)-71-85-treated mice supports a role for T cell-derived TNF α in the progression of PanIN towards invasive carcinoma.

Example 5: High levels of pY323 p38 in human PDAC TIL indicates a poor prognosis

The importance of p38 alternative activation in murine pancreatic cancer, and in particular the oncogene-driven KPC mice, prompted studies to address whether this pathway might be involved in human pancreatic ductal adenocarcinoma (PDAC) as well. Pancreatic tissue from 193 patients with histologically-verified PDAC was analyzed for infiltrating p38 pY323-positive cells. In all samples there was marked infiltration of T cells that stained with antibodies to p38 pY323, but the number of such cells varied from patient to patient. In the majority of cases (n=153, ~80%), less than 10% of the CD3⁺ tumor-infiltrating T cells were also positive for pY323, the remainder having $\geq 10\%$ of their T cells expressing p38 pY323. The p38 pY323-positive T cells also expressed TNF α , as detected in sequential tumor sections. Double immunofluorescence supported these findings, demonstrating that p38 pY323-positive T cells produced TNF α . Patients having infiltrates with more than 10% p38 pY323-positive T cells had a statistically-significant shorter survival (median survival of 296 days post-diagnosis) compared to patients with less than 10% p38 pY323-positive cells (median survival of 613 days post-diagnosis) (FIG. 14A). There was no such correlation with other clinical or pathological findings (age, gender, tumor size, TNM stage, histological grading, UICC stage, number of tumor positive lymph nodes). There was also no difference in the amount of T cell infiltration between the two groups, indicating that activation of the p38 alternative pathway, not simply the presence of infiltrating T cells, correlates with outcome (FIG. 14B). The density of CD31-positive tumor vessels was examined in 10 randomly selected

patients with high and 10 randomly selected patients with low numbers of p38 pY323-positive T cells. The former had a statistically significant increase in vascularization (FIG. 14C), indicating that p38 alternative activation in TIL leads to tumor neovascularization, a known predictor of poor survival in PDAC.

5

Example 6: Evaluation of Gadd45 α polypeptides in an animal model of multiple sclerosis (MS)

This example describes an animal model of multiple sclerosis to evaluate the effect of treatment with a Gadd45 α polypeptide. The experimental autoimmune encephalomyelitis (EAE) mouse model is a recognized model of CNS-relevant demyelinating autoimmune disease. Methods of inducing EAE, and animal models that naturally develop EAE, are well known in the art (see, for example, Matsushita *et al.*, *J. Clin. Invest.* 118(10):3420-3430, 2008; Grewal *et al.*, *Immunity* 14:291-302, 2001; Tran *et al.*, *Eur. J. Immunol.* 30:1410-1415, 2000; or Butzkeuven *et al.*, *Nat. Med.* 8:613-619, 2002; and U.S. Patent Application Publication No. 2008/0166318). The C57BL/6 EAE model is generally representative of chronic and progressive MS.

15

EAE is induced in C57BL/6 female mice by immunization with approximately 200 μ g of myelin oligodendrocyte protein (MOG) 35-55 peptide in complete Freund's adjuvant containing 4 mg/mL of *Mycobacterium tuberculosis* per mouse by subcutaneous injection. Pertussis toxin is administered intraperitoneally at day 0 and day 2 after immunization. At varying stages after immunization, mice are administered a Gadd45 α polypeptide or vehicle alone.

20

Mice are examined for signs of EAE daily. EAE is scored using a 5 point scale (0, no paralysis; 1, limp tail; 2, hind limb paresis and difficulty righting itself; 3, hind limb paralysis; 4, hind limb and forelimb paralysis; and 5, moribund). A decrease in the average score following treatment with the Gadd45 α polypeptide, relative to vehicle alone, indicates that the Gadd45 α polypeptide is effective for promoting myelination and treating MS. Imaging techniques can also be used to assess the degree of myelination at varying stages.

25

Example 7: Evaluation of Gadd45 α polypeptides in MS Patients

This example describes a method to evaluate the use of Gadd45 α polypeptides in humans. The method involves the use of ocular coherence tomography (OCT) and magnetic resonance imaging (MRI) to study myelination of the retinal nerve fiber layer (RNFL).

30

Optic neuritis, a common manifestation of MS, often occurs as the initial manifestation of central nervous system demyelination, or develops during the course of disease. Since the RNFL is

composed only of unmyelinated axons, measuring RNFL thickness represents a viable method of monitoring axonal loss in MS patients. OCT is a noninvasive, noncontact, accurate, and reproducible technique that quantitates the thickness of the peripapillary RNFL, fovea, and macula. RNFL is evaluated in MS patients prior to and following treatment with a Gadd45 α polypeptide.

- 5 An increase in axonal thickness following treatment with the Gadd45 α polypeptide indicates that the Gadd45 α polypeptide is effective for promoting myelination and treatment of MS.

Example 8: Assessment of the efficacy of Gadd45 α polypeptides for the treatment of MS

10 This example describes methods for the assessment of the efficacy of Gadd45 α polypeptides for the treatment of MS.

Subjects having MS are treated with a composition comprising a Gadd45 α polypeptide. One of skill in the art is capable of determining an appropriate dose and dosing schedule of Gadd45 α polypeptide compositions. Subjects are assessed for measures of MS described below prior to initiation of therapy, periodically during the period of therapy and at the end of the course
15 of treatment.

The efficacy of the Gadd45 α polypeptide compositions in subjects with multiple sclerosis can be assessed by the following measures:

1. MRI measures, such as:
 - 20 T₂ lesion load
 - Volume of T₁ hypointensities
 - N-acetyl aspartate (NAA) levels
 - Whole brain atrophy
2. Clinical measures, specifically:
 - 25 Change in EDSS (Expanded Disability Status Scale)
 - Change in SRS (Scripps Neurological Rating Scale)
 - Relapse rate
 - 9-hole peg test
3. Immunologic measures, specifically:
 - 30 Markers of Th1 and Th2 T cell lineages
 - FACS analysis of various T cell markers
 - Cytokine production by T cells *in vitro*
 - Proliferation of T cells

T₂ Lesion Load

The analyses on T₂ lesion load include the following:

- comparison of the mean volume of T₂ lesions during the pre-treatment period to the mean volume of T₂ lesions during the treatment period
- comparison of the mean volume of T₂ lesions during the pre-treatment period to the mean volume of T₂ lesions during the last 4 weeks of the treatment period

Volume of T₁ Hypointensities

The analyses on volume of T₁ hypointensities include the following:

- comparison of the mean volume of T₁ hypointensities during the pre-treatment period to the mean volume of T₁ hypointensities during the treatment period
- comparison of the mean volume of T₁ hypointensities during the pre-treatment period to the mean volume of T₁ hypointensities during the last 4 weeks of the treatment period

EDSS

Baseline EDSS score is determined for subjects prior to start of treatment. EDSS is measured periodically during the course of treatment. The change from baseline EDSS to EDSS during the treatment period is determined. Also, change from baseline scores for SRS and the 9-hole peg test during the treatment period are determined.

Relapses

The frequency of relapses over the 2 years prior to receiving treatment with the Gadd45 α polypeptide are compared to the frequency of relapses while being treated with the Gadd45 α polypeptide.

The effectiveness of therapy comprising administration of Gadd45 α polypeptide compositions in subjects with MS is demonstrated by a reduction in the average number of MS exacerbations per subject in a given period (such as 6, 12, 18 or 24 months) compared with a period prior to treatment. A reduction in the average rate of increase in the subject's disability score over some period (*e.g.*, 6, 12, 18 or 24 months) as measured, for example, by the EDSS score, or even an improvement in the disability score, also demonstrates the effectiveness of the Gadd45 α polypeptide therapy.

Example 9: Evaluation of Gadd45 α polypeptides in an animal model of rheumatoid arthritis

This example describes an animal model of rheumatoid arthritis to evaluate the effect of treatment with a Gadd45 α polypeptide. The collagen-induced arthritis (CIA) model is the most commonly used animal model to study rheumatoid arthritis. Methods of inducing autoimmune arthritis in this model are well known in the art (see, for example, Jirmanova *et al.*, *Blood* 118(12):3280-3289, 2011; Brand *et al.*, *Nat Protoc* 2(5):1269-1275, 2007).

In this example, arthritis is induced by injecting 10 week-old H-2^r mice with 100 μ g bovine type II collagen emulsified in complete Freund's adjuvant (CFA) at the base of the tail, and boosting 21 days later with 100 μ g bovine type II collagen emulsified in incomplete Freund's adjuvant (IFA). After induction of arthritis, mice are administered a Gadd45 α polypeptide or vehicle alone. Each paw is scored as follows: 0, no evidence of erythema; 1, erythema and mild swelling confined to tarsals; 2, erythema and mild swelling extending from the ankle to midfoot; 3, swelling extending from the ankle to metatarsal joints; 4, severe swelling/ankylosis of the whole foot. A decrease in the average score following treatment with the Gadd45 α polypeptide, relative to vehicle alone, indicates that the Gadd45 α polypeptide is effective for the treatment of rheumatoid arthritis.

In view of the many possible embodiments to which the principles of the disclosed invention may be applied, it should be recognized that the illustrated embodiments are only preferred examples of the invention and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

CLAIMS

1. An isolated growth arrest and DNA-damage-inducible alpha (Gadd45 α) polypeptide no more than 25 amino acids in length, wherein the amino acid sequence of the polypeptide
5 comprises LQIHFTLIXAFCCEN (SEQ ID NO: 1), wherein X is Q or R.
2. The isolated Gadd45 α polypeptide of claim 1 which is no more than 20 amino acids in length.
- 10 3. The isolated Gadd45 α polypeptide of claim 1, comprising the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).
4. The isolated Gadd45 α polypeptide of claim 1, consisting of the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).
15
5. A fusion protein comprising the Gadd45 α polypeptide of claim 1 and a heterologous peptide.
6. The fusion protein of claim 5, wherein the heterologous peptide is a cell-penetrating
20 peptide (CPP).
7. The fusion protein of claim 6, wherein the CPP comprises poly-arginine.
8. The fusion protein of claim 7, wherein poly-arginine comprises or consists of 11
25 arginine (11R) residues.
9. The fusion protein of claim 8, consisting of 11R fused to the N-terminus of the Gadd45 α polypeptide.
- 30 10. The fusion protein of claim 8, consisting of 11R fused to the amino acid sequence LQIHFTLIQAFCCEN (SEQ ID NO: 2) or LQIHFTLIRAFCCEN (SEQ ID NO: 3).

11. A composition comprising the polypeptide or fusion protein of any one of claims 1-10 and a pharmaceutically acceptable carrier.

12. An *in vitro* or *ex vivo* method of inhibiting proliferation of T cells in response to T cell receptor stimulation, or of inhibiting differentiation of T cells into Th1 or Th17 cells, comprising contacting the T cells with the polypeptide or fusion protein of any one of claims 1-10, thereby inhibiting proliferation of the T cells, or inhibiting differentiation of the T cells.

13. The method of claim 12 which is an *ex vivo* method, wherein the T cells have been isolated from a subject and the isolated T cells are contacted with the polypeptide or fusion protein.

14. Use of the polypeptide or fusion protein of any one of claims 1-10 in the preparation of a medicament for inhibiting proliferation of T cells in response to T cell receptor stimulation in a subject, or for inhibiting differentiation of T cells into Th1 or Th17 cells in a subject.

15. Use of the polypeptide or fusion protein of any one of claims 1-10 in the preparation of a medicament for treating a subject with a T cell- and/or p38-mediated autoimmune or inflammatory disorder.

16. The use of claim 15, wherein the T cell- and/or p38-mediated autoimmune or inflammatory disorder comprises multiple sclerosis, rheumatoid arthritis or pancreatic cancer.

17. The use of claim 15, wherein the polypeptide or fusion protein is administered by intravenous, intra-tumor or intraperitoneal administration.

18. The use of claim 15, wherein the T cell- and/or p38-mediated autoimmune or inflammatory disorder comprises pancreatic cancer and the polypeptide or fusion protein is administered by intra-tumor injection.

19. Use of the polypeptide or fusion protein of any one of claims 1-10 in the preparation of a medicament for inhibiting tumor vascularization in a subject.

20. The use of claim 19, wherein the tumor is a pancreatic tumor.


21. The use of claim 19, wherein the polypeptide or fusion protein is administered by intra-tumor administration.

5 22. The use of claim 19, wherein the polypeptide or fusion protein is administered at a dose of about 20 to about 100 µg.

23. The use of claim 14, wherein the polypeptide or fusion protein is administered at a dose of about 20 to about 100 µg.

10

24. The use of claim 15, wherein the polypeptide or fusion protein is administered at a dose of about 20 to about 100 µg.

FIG. 1

Sample	%
11R-71-85	88.4
11R-Ctrl	83
No peptide	94.7

FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 5C





FIG. 5D



aCD3+aCD28	-	+	+	+
TAT-Ctrl	-	+	-	-
TAT-G45FL	-	-	+	-
TAT-G45D10	-	-	-	+

FIG. 6A




FIG. 6B




FIG. 7





FIG. 8

T cells



CD4⁺T cells




FIG. 9C




FIG. 9B




FIG. 9A





FIG. 10A








FIG. 11

CD31+ staining

(11R) Scr



(11R) 71-85








FIG. 13A**FIG. 13B**

FIG. 13C




FIG. 13D




FIG. 14B




FIG. 14A





FIG. 14C



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2013/070813**A. CLASSIFICATION OF SUBJECT MATTER****C07K 7/08(2006.01)i, C07K 19/00(2006.01)i, A61K 38/10(2006.01)i, A61P 37/00(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K 7/08; A61K 39/395; C07K 16/40; C07K 19/00; A61K 38/10; A61P 37/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: growth arrest and DNA-damage-inducible alpha (Gadd45a), T cell specific alternative p38 activation pathway, autoimmune

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NCBI, GenBANK accession no.AEI16541.1 (20 June 2011) See the whole document.	1-4, 11
Y		12-15, 17
A		5-10, 16, 18-20
Y	SALVADOR, JESUS M. et al., `The autoimmune suppressor Gadd45a inhibits the T cell alternative p38 activation pathway`, Nature Immunology, 27 February 2005, Vol.6, No.4, pp.396-402 See abstract; p.396, left column, lines 19-20 and p.401, left column, lines 28-29.	12-15, 17
A		1-11, 16, 18-20
X	WO 2005-077983 A2 (GOVWEMENT OF THE UNITED STATE OF AMERICA AS REPRESENTED BY THE SECRETARY OF THE DEPARTMENT OF HEALTH AND HUMAN SERVICES) 25 August 2005 See p.20, line 21 - p.21, line 4; claims 5-6.	1-4, 11
Y		12-15, 17
A		5-10, 16, 18-20

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 February 2014 (24.02.2014)

Date of mailing of the international search report

24 February 2014 (24.02.2014)

Name and mailing address of the ISA/KR


 Korean Intellectual Property Office
 189 Cheongsa-ro, Seo-gu, Daejeon Metropolitan City,
 302-701, Republic of Korea

Facsimile No. +82-42-472-7140

Authorized officer

HEO, Joo Hyung

Telephone No. +82-42-481-8150



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2013/070813

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JIRMANOVA, LUDMILA et al., `Lack of the T cell-specific alternative p38 activation pathway reduces autoimmunity and inflammation`, Blood, 22 September 2011, Vol.118, No.12, pp.3280-3289 See abstract.	1-20
A	SALVADOR, JESUS M. et al., `Alternative p38 activation pathway mediated by T cell receptor-proximal tyrosin kinases`, Nature Immunology, 27 February 2005, Vol.6, No.4, pp.390-395 See abstract.	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2013/070813Patent document
cited in search reportPublication
datePatent family
member(s)Publication
date

WO 2005-077983 A2

25/08/2005

WO 2005-077983 A3

16/02/2006