Innovation, Sciences et Développement économique Canada
Office de la Propriété Intellectuelle du Canada

Innovation, Science and Economic Development Canada
Canadian Intellectual Property Office

CA 3023257 A1 2017/11/09

(21) 3 023 257

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2017/05/05
(87) Date publication PCT/PCT Publication Date: 2017/11/09
(85) Entrée phase nationale/National Entry: 2018/11/02
(86) N° demande PCT/PCT Application No.: US 2017/031301
(87) N° publication PCT/PCT Publication No.: 2017/192993
(30) Priorités/Priorities: 2016/05/05 (US62/332,013); 2016/10/06 (US62/404,960); 2016/12/29 (US62/440,078); 2017/03/16 (US62/472,204)

(51) Cl.Int./Int.Cl. *A61K 9/72* (2006.01), *A61K 31/00* (2006.01), *A61K 31/5575* (2006.01), *A61P 9/12* (2006.01)


(71) Demandeur/Applicant:
LIQUIDIA TECHNOLOGIES, INC., US

(72) Inventeurs/Inventors:
ROSCIGNO, ROBERT FRANK, US;
FARRER, BRIAN T., US;
SPRAGUE, JACOB J., US;
MAYNOR, BENJAMIN, US

(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : POUDRE SECHE DE TREPROSTINIL POUR LE TRAITEMENT DE L'HYPERTENSION PULMONAIRE
(54) Title: DRY POWDER TREPROSTINIL FOR THE TREATMENT OF PULMONARY HYPERTENSION

FIGURE 1



(57) Abrégé/Abstract:
A dry powder inhalation treatment for pulmonary arterial hypertension includes a dose of dry particles comprising greater than 25 micrograms of treprostinil enclosed in a capsule. The dry particles can include treprostinil, a wetting agent, a hydrophobicity modifying agent, a pH modifying agent and a buffer. A method of treating a patient having pulmonary arterial hypertension includes providing a patient a dry powder inhaler, providing the patient at least one capsule for use in the dry powder inhaler, the capsule including at least 25 micrograms of treprostinil.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
09 November 2017 (09.11.2017)

WIPO | PCT

(10) International Publication Number
WO 2017/192993 A1

(51) International Patent Classification:
A61P 9/12 (2006.01) *A61K 31/192* (2006.01)
A61K 9/72 (2006.01) *A61K 31/557* (2006.01)
A61K 31/00 (2006.01) *A61P 9/00* (2006.01)

(21) International Application Number:
PCT/US2017/031301

(22) International Filing Date:
05 May 2017 (05.05.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

62/332,013	05 May 2016 (05.05.2016)	US
62/404,960	06 October 2016 (06.10.2016)	US
62/440,078	29 December 2016 (29.12.2016)	US
62/472,204	16 March 2017 (16.03.2017)	US

(71) Applicant: **LIQUIDIA TECHNOLOGIES, INC.** [US/US]; 419 Davis Drive, Suite 100, Morrisville, NC 27560 (US).

(72) Inventors: **ROSCIGNO, Robert, Frank**; 7415 Aquarina Beach Drive, Unit 403, Melbourne Beach, FL 32951 (US). **FARRER, Brian, T.**; 1204 Fairfax Woods Drive, Apex, NC 27502 (US). **SPRAGUE, Jacob, J.**; 2069 Weston Green Loop, Cary, NC 27513 (US). **MAYNOR, Benjamin**; 3700 Northampton Road, Durham, NC 27707 (US).


(74) Agent: **LEUNG, Richard, L** et al.; Morgan, Lewis & Bockius, LLP, 1701 Market Street, Philadelphia, PA 19103 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: DRY POWDER TREPROSTINIL FOR THE TREATMENT OF PULMONARY HYPERTENSION

FIGURE 1



1 micrometer 'pollen' particle

(57) Abstract: A dry powder inhalation treatment for pulmonary arterial hypertension includes a dose of dry particles comprising greater than 25 micrograms of treprostinil enclosed in a capsule. The dry particles can include treprostinil, a wetting agent, a hydrophobicity modifying agent, a pH modifying agent and a buffer. A method of treating a patient having pulmonary arterial hypertension includes providing a patient a dry powder inhaler, providing the patient at least one capsule for use in the dry powder inhaler, the capsule including at least 25 micrograms of treprostinil.

WO 2017/192993 A1

[Continued on next page]

WO 2017/192993 A1

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

Dry Powder Treprostinil for the Treatment of Pulmonary Hypertension

Cross-Reference to Related Applications

[0001] This application claims priority to and the benefit of U.S. Provisional Patent Application No. 62/332,013, filed May 5, 2016, U.S. Provisional Patent Application No. 62/404,960, filed October 6, 2016, U.S. Provisional Patent Application No. 62/440,078, filed December 29, 2016, and U.S. Provisional Patent Application No. 62/472,204, filed March 16, 2017, all of which are incorporated herein by reference in their entireties.

TECHNICAL FIELD

[0002] The present invention provides an improvement to the treatment of pulmonary hypertension, a condition that deteriorates the lives of many thousands of patients toward an untimely death. The present invention provides, for the first time, a stable, user friendly, uniform dry powder inhaled treprostinil formulation, methods of making, and use thereof in humans.

BACKGROUND

[0003] Pulmonary arterial hypertension (PAH) is a complex, multifactorial, progressive, and life-threatening disease characterized by proliferative and obstructive changes in the pulmonary vasculature and involving numerous biochemical pathways and cell types. The disease is characterized by elevated pulmonary arterial pressure caused by narrowing of the blood vessels in the lungs and, ultimately, right ventricular failure. The disease carries a poor prognosis associated with significant morbidity and mortality, having a historical survival rate less than five years. PAH is a sub-group of pulmonary hypertension (PH), which is elevation of blood pressure in lungs. Endothelial dysfunction is thought to occur early on, leading to cell proliferation and structural changes in the pulmonary vasculature that lead to increased pulmonary arterial pressure (PAP) and resultant right ventricular enlargement and dysfunction. In addition, endothelial dysfunction results in chronically impaired production of vasoactive mediators, such as nitric oxide (NO) and prostacyclin, along with prolonged overexpression of vasoconstrictors, such as endothelin-1.

[0004] PAH affects approximately 15 out of every one million individuals. There are approximately 1,000 new cases of PAH diagnosed in the United States each year. The mean age

at diagnosis is between 50 and 65 years of age, although the disorder may present much earlier in childhood or even infancy. While gender-based prevalence estimates for PAH are variable, estimates for the overall prevalence of pulmonary hypertension (PH) in females is approximately twice that of males.

[0005] PAH is part of a larger classification for pulmonary hypertension which is divided into five groups based on World Health Organization (WHO) criteria (designated as WHO Groups 1 through 5). PAH is used to describe exclusively WHO Group 1. Pulmonary hypertension is used to describe the remaining four groups (WHO Groups 2-5) and also when referring to all 5 groups collectively.

- WHO Group 1 - PAH: Pulmonary arterial hypertension.
- WHO Group 2 - PH: Pulmonary hypertension secondary to left heart disease.
- WHO Group 3 - PH: Pulmonary hypertension secondary to lung diseases or hypoxemia.
- WHO Group 4 - PH: Chronic thromboembolic pulmonary hypertension.
- WHO Group 5 - PH: Pulmonary hypertension with unknown mechanisms.

[0006] PAH initially presents as exertional dyspnea, lethargy, and fatigue and is often confused for other disease states. As PAH progresses and right ventricular failure develops, exertional chest pain (i.e., angina), exertional syncope, and peripheral edema may develop. Following confirmation of diagnosis based on hemodynamic parameters, treatment is recommended to lower pulmonary pressures and treat the symptoms of PAH. Although no cure exists for PAH, treatment of PAH is directed at improving hemodynamic measures, New York Heart Association (NYHA) functional class, the 6 minute walk distance (6MWD), quality of life, and, in some studies, survival.

[0007] The severity of PAH may be classified according to the NYHA heart failure guidelines as follows:

- NYHA Class I: Patients with no limitation of activities; they suffer no symptoms from ordinary activities.
- NYHA Class II: Patients with slight, mild limitation of activity; they are comfortable with rest or with mild exertion.

- NYHA Class III: Patients with marked limitation of activity; they are only comfortable at rest.
- NYHA Class IV: Patients who should be at complete rest, confined to bed or chair; any physical activity brings on discomfort and symptoms occur at rest.

[0008] While the exact underlying cause of PAH is unclear, mutations in the bone morphogenic protein receptor type II (BMPR2) gene account for approximately 75% of familial PAH and up to 25% of apparently sporadic PAH cases. These mutations may promote cell division or prevent cell death, resulting in an overgrowth of cells in smaller pulmonary arteries. This overgrowth increases resistance to blood flow, triggering hypertension. Additional genetic abnormalities may also contribute to PAH.

[0009] Currently Available Treatments

[0010] There are five classes of drugs that have been approved to treat PAH, including endothelin receptor antagonists (ERAs), phosphodiesterase type 5 (PDE5) inhibitors, soluble guanylate cyclase stimulators, prostacyclin receptor agonists, and prostacyclin analogs. Approved PAH therapies and their route of administration include:

- ERA: bosentan (oral) and ambrisentan (oral)
- PDE5: sildenafil (oral, intravenous (IV) and tadalafil (oral)
- Soluble Guanylate Cyclase (sGC) Stimulator: riociguat (oral)
- Prostacyclin Receptor Agonist: selexipag (oral)
- Prostacyclin Analog: epoprostenol (IV) iloprost (inhaled), and treprostinil (oral), (subcutaneous and IV), and (inhaled)

[0011] Treprostinil is a chemically stable tricyclic benzidine prostanoid with vasodilator properties that is capable of reducing pulmonary vasoconstriction with minimal effects on systemic blood pressure. Treprostinil has been approved for the treatment of PAH under the trade names REMODULIN® (United Therapeutics Corporation; subcutaneous or IV infusion) and TYVASO® (United Therapeutics Corporation; inhaled via ultrasonic, pulsed nebulization delivery device). While both have proven effective for PAH, one advantage of TYVASO's

inhaled route of administration is that it brings the drug very near the desired site of action (pulmonary arteries in the lungs).

[0012] Despite the current treatment options for PAH patients, each option includes drawbacks, most notably for the inhaled route of administration Tyvaso requires use of a large, cumbersome nebulization device that requires power, water and user manipulation for cleaning and operating. Moreover, the nebulization device by its nature is not convenient to the patient as compared to carrying a small, concealable dry powder inhalation device such as those used for treating asthma and many other chronic and acute issues. Furthermore, nebulized treprostinil has shown clinical limitations on treprostinil dosing, which may limit the applicability of the inhaled route of administration to a smaller subsector of PAH patients than necessarily treatable via the inhaled route from a dry powder inhaled treprostinil product of the present invention.

SUMMARY OF THE INVENTION

[0013] The present inventors have developed and reduced to practice an inhalation dry powder formulation of treprostinil that is produced using Liquidia's PRINT® Technology (Particle Replication in Nonwetting Templates), LIQUIDIA TECHNOLOGIES, INC. This PRINT particle formulation for dry powder delivery of treprostinil (otherwise referred to as LIQ861) is under clinical evaluation. The present applicants intend to use the same indication (i.e., treatment of pulmonary arterial hypertension [WHO Group 1] in patients with NYHA Class III symptoms, to improve exercise ability) dose and dose regimen (4X/day) as defined in the approved nebulized treatment label (TYVASO® UNITED THERAPEUTICS). In particular, the present invention provides for dosing levels that exceed the maximum tolerated dose delivered through a nebulizer. In some cases the present invention may also treat other indications under the pulmonary hypertension disease states.

[0014] In some embodiments, a dry powder inhalation treatment for pulmonary arterial hypertension according to the present invention includes a dose of dry particles comprising greater than 25 micrograms of treprostinil enclosed in a capsule. In some embodiments, the dose of dry particles comprises from about 25 micrograms to about 400 micrograms of treprostinil. In some embodiments, the dose of dry particles comprises from about 50 micrograms to about 350 micrograms of treprostinil. In some embodiments, the dose of dry

particles comprises from about 75 micrograms to about 300 micrograms of treprostinil. In some embodiments, the dose of dry particles comprises from about 100 micrograms to about 300 micrograms of treprostinil. In some embodiments, the dose of dry particles includes greater than or equal to 100 micrograms of treprostinil. In some embodiments, the dose of dry particles comprises greater than or equal to 150 micrograms of treprostinil. In some embodiments, the dose of dry particles comprises greater than or equal to 200 micrograms of treprostinil. In some embodiments, the dose of dry particles comprises greater than or equal to 250 micrograms of treprostinil. In some embodiments, the dose of dry particles comprises greater than or equal to 300 micrograms of treprostinil. In some embodiments, the dose of dry particles includes greater than or equal to 5 mg of the dry particles. In some embodiments, the dose of dry particles includes greater than or equal to 10 mg of the dry particles. In yet other embodiments, the dose of dry particles includes greater than or equal to 15 mg of the dry particles. In further embodiments, a dry powder treatment for pulmonary arterial hypertension, includes a single capsule enclosing 5 mg or more dry particles comprising 25 micrograms of treprostinil per each 5 mg of the dry particles.

[0015] In some embodiments, a method of treating a patient having pulmonary arterial hypertension includes providing a patient a dry powder inhaler, providing the patient at least one capsule for use in the dry powder inhaler, wherein the capsule comprises at least 25 micrograms of treprostinil, and instructing the patient to utilize the dry powder inhaler to inhale the treprostinil. In some such embodiments, the capsule includes at least 50 micrograms of treprostinil. In some embodiments, the capsule includes at least 100 micrograms of treprostinil. In some embodiments, the capsule comprises at least 150 micrograms of treprostinil. In some embodiments, the capsule comprises greater than or equal to 200 micrograms of treprostinil. In some embodiments, the capsule comprises greater than or equal to 250 micrograms of treprostinil. In some embodiments, the capsule comprises greater than or equal to 300 micrograms of treprostinil. In some embodiments, the capsule comprises from about 25 micrograms to about 400 micrograms of treprostinil. In some embodiments, the capsule comprises from about 50 micrograms to about 350 micrograms of treprostinil. In some embodiments, the capsule comprises from about 75 micrograms to about 300 micrograms of treprostinil. In some embodiments, the capsule comprises from about 100 micrograms to about 300 micrograms of treprostinil. In further embodiments, the patient may be prescribed to use

two capsules per dose cycle per day, generally with PAH requiring 4 times per day dosing. In some embodiments, the patient may be prescribed to use three capsules per day. In some embodiments, the patient may be prescribed to use four capsules per day. In some embodiments, a method of treating a patient having pulmonary arterial hypertension includes dosing the patient having pulmonary arterial hypertension with a dry powder dose of treprostinil, wherein the dose of treprostinil is greater than 85 micrograms (e.g., about 100 micrograms to about 350 micrograms). In some embodiments, the patient may be dosed one, two, three, four, or more times per day. A further method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, greater than 12.5 micrograms of treprostinil to a patient per breath. In another embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, greater than 25 micrograms of treprostinil to a patient per breath. In another embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, from about 12.5 to about 50 micrograms of treprostinil to a patient per breath. In yet another embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, about 25 to about 50 micrograms of treprostinil to a patient per breath. In a further embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, greater than 50 micrograms of treprostinil to a patient per breath. In a further embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, greater than or equal to 100 micrograms of treprostinil to a patient per breath. In a further embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, greater than or equal to 150 micrograms of treprostinil to a patient per breath. In a further embodiment, a method of treating a patient having pulmonary arterial hypertension includes delivering, in dry powder, greater than or equal to 200 micrograms of treprostinil to a patient per breath.

[0016] A dry powder inhalation composition for treating pulmonary arterial hypertension according to a further embodiment includes a plurality of dry powder particles comprising treprostinil, a non-reducing sugar, a wetting agent, a hydrophobicity modifying agent, a pH modifying agent and a buffer. In some such embodiments, the bulking agent comprises trehalose dihydrate. In some embodiments, the wetting agent comprises polysorbate 80. In some embodiments, the hydrophobicity modifying agent comprises L-leucine. In some

embodiments, the pH modifying agent comprises sodium citrate dihydrate. In some embodiments, the buffer comprises sodium chloride. In certain embodiments, the composition comprises less than about 4 percent by weight water. In some embodiments, the composition comprises less than about 2 percent by weight water. In some embodiments, the composition comprises less than about 1 percent by weight water.

[0017] In yet further embodiments, the dry powder particles include particles having a three dimensional shape including a width and length not less than 1 micrometer and not more than 2 micrometers and a depth not less than 0.3 micrometers and not more than 0.8 micrometers. In some embodiments, the dry powder particles comprise a dried solution comprising trehalose dihydrate, L-leucine, treprostinil sodium, polysorbate 80, sodium citrate dihydrate, sodium chloride and water. In some embodiments, the dry powder particles comprise by percent solids about 0.581 percent treprostinil sodium, about 92.32 percent trehelose, about 2.19 percent polysorbate 80, about 4.39 percent L-leucine, about 0.26 percent sodium citrate, and about 0.25 percent sodium chloride.

[0018] A method of making a particle for dry powder delivery to the lung of a patient in need thereof, in some embodiments, includes molding a composition comprising about 12.30 weight percent trehelose dihydrate, about 0.53 weight percent L-leucine, about 0.07 weight percent treprostinil sodium, about 0.26 weight percent polysorbate 80, about 0.04 weight percent sodium citrate dihydrate, about 0.03 weight percent sodium chloride and about 86.78 weight percent water into a particle. In some embodiments, the method of making the particle further includes drying the composition such that the particle comprises less than 4 percent by weight water.

BRIEF DESCRIPTION OF THE FIGURES

[0019] The foregoing summary, as well as the following detailed description of the invention, will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, there are shown in the drawings embodiments which are presently preferred. It should be understood, however, that the invention can be embodied in different forms and thus should not be construed as being limited to the illustrated embodiments set forth herein.

[0020] Figure 1 shows a three-dimension rendering of a pollen particle according to an embodiment of the present invention.

[0021] Figure 2 shows an example NGI distribution for active particles (PAH-1R-0943-010). For each of the three data sets represented for each collection cup, the beginning of the run is the left hand bar (A1), the middle of the run is the center bar (B1), and the end of the run is the right hand bar (C1). Data was obtained using the Monodose Model 8 device (95 L/min, 2 sec).

[0022] Figures 3A and 3B are tables including data for Cohort 1 of a clinical trial. The table shown in Figure 3A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 1. Preliminary non-compartmental PK parameters for treprostinil are summarized in the table shown in Figure 3B.

[0023] Figures 4A and 4B are tables including data for Cohort 2 of a clinical trial. The table shown in Figure 4A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 2. Preliminary non-compartmental PK parameters for treprostinil for Cohort 2 are summarized in the table shown in Figure 4B.

[0024] Figures 5A and 5B are tables including data for Cohort 3 of a clinical trial. The table shown in Figure 5A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 3. Preliminary non-compartmental PK parameters for treprostinil for Cohort 3 are summarized in the table shown in Figure 5B.

[0025] Figures 6A and 6B are tables including data for Cohort 4 of a clinical trial. The table shown in Figure 6A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 4. Preliminary non-compartmental PK parameters for treprostinil for Cohort 4 are summarized in Figure 6B.

[0026] Figures 7A and 7B are tables including data for Cohort 5 for a clinical trial. The table shown in Figure 7A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 5. Preliminary non-compartmental PK parameters for treprostinil for Cohort 5 are summarized in Figure 7B.

[0027] Figures 8A, 8B, and 8C are tables including data for Cohort 6 for a clinical trial. The table shown in Figure 8A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 6-R. Preliminary non-compartmental PK parameters for treprostinil for Cohort 6-R are summarized in Figure 8B. Preliminary non-compartmental PK parameters for treprostinil for Cohort 6-Original are summarized in Figure 8C.

[0028] Figures 8D, 8E, 8F, and 8G contain data for the clinical trial. Mean concentration-time data for each of the six cohorts is displayed on a linear scale in Figure 8D. Plots of the relationship between dose and Cmax and AUCinf are displayed in Figure 8E and Figure 8F, respectively. A plot of the relationship between dose and the oral clearance, CL/F, is shown in Figure 8G.

[0029] Figure 9 is an SEM image showing pollen-shaped particles according to an embodiment of the present invention.

[0030] Figure 10 is a flow diagram showing a process of manufacturing particles according to an embodiment of the present invention.

[0031] Figure 11 shows an example dry powder inhalation device which may be used to deliver particles to a patient in accordance with embodiments of the present invention.

DETAILED DESCRIPTION OF EMBODIMENTS

[0032] Drug Substance

[0033] The drug substance (DS) according to embodiments of the present invention is treprostinil, which is a synthetic analog of prostacyclin (PGI_2). The IUPAC name for treprostinil is (2-[[(1R,2R,3aS,9aS)-2-hydroxy-1-[(3S)-3-hydroxyoctyl]-2,3,3a,4,9,9a-hexahydro-1H-cyclopenta[g]naphthalen-5-yl]oxy]acetic acid).

[0034] Inhalation Powder Drug Product

[0035] The inhalation powder drug product according to certain aspects of the present invention provides a dry powder dosage form of treprostinil and excipients formed into a particle

(drug product intermediate, or DP-intermediate) that is, in some embodiments, filled into a capsule, for example, a hydroxypropyl methylcellulose (HPMC) capsule (size 3) (LIQ861). In some embodiments, the DP-intermediate is a treprostinil/excipient matrix from which particles of precise size and shape are formed according to the methods herein. In one example, the particles of the DP-intermediate comprise a shape corresponding generally to a rounded triangular shape having a volume, where the inner portion of the rounded triangular shape, in size, fits a 1 micrometer equilateral triangle (otherwise referred to as being pollen-shaped). A three-dimensional rendering of such a particle shape is depicted in Figure 1. In another embodiment, the pollen-shape may be trefoil-shaped with an inscribed circle diameter of 1 micrometer, and a prescribed thickness of a value or range between 0.5 and 1 micrometer, or more preferred 0.7 micrometer. In addition, certain embodiments of the present drug product includes particles having 0.5% treprostinil used in a first clinical study to investigate dose levels of 25 mcg, 50 mcg, 75 mcg, 100 mcg, 125 mcg and 150 mcg treprostinil in LIQ861. In further embodiments, a drug product according to the present invention may provide dose levels of 175 mcg, 200 mcg, 225 mcg, 250 mcg, 275 mcg, 300 mcg, 325 mcg, or 350 mcg treprostinil. In further embodiments, a drug product according to the present invention may provide dose levels of 50 mcg treprostinil plus or minus 10 mcg, 9 mcg, 8 mcg, 7 mcg, 6 mcg, 5 mcg, 4 mcg, 3 mcg, 2 mcg or 1 mcg treprostinil loaded into capsules for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide dose levels of 75 mcg treprostinil plus or minus 10 mcg, 9 mcg, 8 mcg, 7 mcg, 6 mcg, 5 mcg, 4 mcg, 3 mcg, 2 mcg or 1 mcg treprostinil loaded into capsules for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide dose levels of 100 mcg treprostinil plus or minus 10 mcg, 9 mcg, 8 mcg, 7 mcg, 6 mcg, 5 mcg, 4 mcg, 3 mcg, 2 mcg or 1 mcg treprostinil loaded into capsules for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide dose levels of 150 mcg treprostinil plus or minus 10 %, 9 %, 8 %, 7 %, 6 %, 5 %, 4 %, 3 %, 2 % or 1 % treprostinil loaded into capsules for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide dose levels of 200 mcg treprostinil plus or minus 10 %, 9 %, 8 %, 7 %, 6 %, 5 %, 4 %, 3 %, 2 % or 1 % mcg treprostinil loaded into capsules for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide dose levels of 300

mcg treprostinil plus or minus 10 %, 9 %, 8 %, 7 %, 6 %, 5 %, 4 %, 3 %, 2 % or 1 % treprostinil loaded into capsules for delivery to a patient in a dry powder.

[0036] According to the present invention, due to the formulation of the present dry powder particles, the particles remain stable for long periods of time at relatively low humidity conditions. In some embodiments, the present invention provides dry powder particles packaged under sealed conditions that remain stable for more than 3 months at 40 degrees Celsius at 75 percent relative humidity. Therefore, the particles can be utilized to provide a patient with a dry powder inhaled drug form of treprostinil, not previously available until the present invention. This invention, in some embodiments, provides a user with a reduction in interaction with drug product by removing the requirements on the patient to reconstitute their drug product for use in a nebulizer device. The patient is also enabled to receive equal dosing with more than 50 percent reduction in breath treatments on a device, and in some embodiments more than 65 percent reduction in breath treatments.

[0037] The present invention, in some embodiments, also provides a dry formulation of treprostinil, which upon delivery to a patient via the inhaled route, becomes soluble and pharmaceutically available in less than 10 seconds. In some embodiments, the dry formulation composition becomes soluble and pharmaceutically available in less than 5 seconds. In some embodiments, the dry formulation composition becomes soluble and pharmaceutically available in less than 2 seconds. In some embodiments, the dry formulation composition becomes soluble and pharmaceutically available in about 1 second. In some embodiments, the dry formulation composition becomes soluble and pharmaceutically available in less than 1 second. In some embodiments, the dry formulation composition becomes soluble and pharmaceutically available in less than about 0.5 seconds. Furthermore, the excipients in the dry particle formulation of the present invention maintain pH and salt gradient during processing such that the active agent remains in a state to become soluble in the lung conditions of a user.

[0038] A detailed description of the LIQ861 formulation, particle composition, particle geometry, packaging, device, delivery, stability, dose, and a description of the use follows.

[0039] In some embodiments, a formulation according to the present invention includes a drug substance (e.g., Treprostinil, Treprostinil Sodium) together with one or more excipients. In some

embodiments, the one or more excipients may include a bulking agent, a wetting agent, a hydrophobicity modifier, a pH modifier, a buffer component, or combinations thereof. Examples of such formulations according to certain specific embodiments are provided in the tables below.

LIQ861 Drug Product-Intermediate Description for Active (LIQ861) and Placebo Formulations (dihydrate form calculations)

Component	Function	Quantity (mg/g) (Active)	Percent Solids
Treprostinil Sodium	Drug Substance	5.3 (5.0 as treprostinil)	0.53
Trehalose Dihydrate	Bulking Agent	930	92.97
Polysorbate 80	Wetting Agent	20	2.00
L-Leucine	Hydrophobicity Modifier	40	4.00
Sodium Citrate Dihydrate	pH Modifier	2.7	0.27
Sodium Chloride	Buffer Component	2.3	0.23

LIQ861 Drug Product-Intermediate Description for Active (LIQ861) and Placebo Formulations (anhydrous form calculations)

Component	Function	Quantity (mg/g) (Active)	Percent Solids	Normalized mg/g
Treprostinil Sodium	Drug Substance	5.3 (5.0 as treprostinil)	0.581	5.81
Trehalose	Bulking Agent	841	92.32	923.23
Polysorbate 80	Wetting Agent	20	2.19	21.94
L-Leucine	Hydrophobicity Modifier	40	4.39	43.89
Sodium Citrate	pH Modifier	2.4	0.26	2.60
Sodium Chloride	Buffer Component	2.3	0.25	2.52

[0040] Inhalation Device

[0041] According to an embodiment of administering the present invention drug particle, LIQ861 is administered using an RS00 Model 8 dry powder inhalation device (Plastiape S.p.A.). The present invention provides for multi-day administration of LIQ861 according to some embodiments.

[0042] Indication

[0043] The present invention, according to an embodiment, is useful for the treatment of pulmonary arterial hypertension (WHO Group 1) in patients with NYHA Class III symptoms, to improve exercise ability.

[0044] Chemistry, Manufacturing, and Controls (CMC)

[0045] Drug Substance (DS)

[0046] The drug substance according to embodiments of the present invention is treprostinil and the salt form used for LIQ861 is treprostinil sodium. Detailed information about treprostinil sodium, including physical and chemical properties, characterization, manufacturing and controls, container closure system, and stability attributes may be found in the Drug Master File (DMF) lodged with the FDA for treprostinil. General information on the DS is provided herein.


[0047] Nomenclature

[0048] The international non-proprietary name (INN) for LIQ861 is treprostinil sodium. The chemical name is 2-((1R,2R,3aS,9aS)-2-hydroxy-1-((S)-3-hydroxyoctyl)-2,3,3a,4,9,9a-hexahydro-1H-cyclopenta[b]naphthalen-5-yloxy)acetic acid, sodium salt. The chemical abstracts service registration number is [289480-64-4].

[0049] Structure

[0050] The structure of treprostinil sodium is depicted herein below. The molecular formula is $C_{23}H_{33}NaO_5$ and it has a molecular weight of 412.49 daltons.

[0051] Chemical Structure of Treprostinil Sodium



[0052] General Properties

[0053] Treprostinil sodium appears as a white or pale yellowish powder. It is very soluble in water and ethanol, very slightly soluble in acetone, and practically insoluble in acetonitrile, n-hexane, and ethyl acetate. The specific optical rotation calculated with reference to the anhydrous and solvent free basis is $[\alpha]_D^{20} = +38.0° \sim +44.0°$. It is hygroscopic. The pKa of treprostinil is 4.5, using aqueous titration with 20% ethanol as a co-solvent. The distribution

coefficient of treprostinil in various buffer solutions at various pH levels indicates distribution into octanol layers at all pH levels.

[0054] Inhalation Particle Drug Product – LIQ861

[0055] Description and Composition of the Drug Product Particle

[0056] The inhalation drug particle product, in some embodiments, includes or consists of a dry powder dosage form of treprostinil and excipients (drug product-intermediate; DP-intermediate; or drug particle) that may be filled into, for example, a HPMC capsule (size 3). The DP-intermediate, in some embodiments, is a treprostinil/excipient matrix from which particles of precise size (e.g., 1 μm) and shape (e.g., "pollen-shaped") are created using Liquidia's PRINT Technology. The "pollen-shaped" particles may also be described as trefoil-shaped, with an inscribed circle diameter of 1 μm, and a thickness of 0.7 μm. A three-dimensional rendering of such a particle shape is depicted in Figure 1. LIQ861 comprised drug product capsule strengths of 25 mcg, 50 mcg, and 75 mcg treprostinil used in the first clinical study to investigate planned dose levels of 25 mcg, 50 mcg, 75 mcg, 100 mcg, 125 mcg and 150 mcg treprostinil. The 100 mcg, 125 mcg and 150 mcg doses may be made up of a combination of lower dose capsules. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 175 mcg, 200 mcg, 225 mcg, 250 mcg, 275 mcg, 300 mcg, 325 mcg, or 350 mcg treprostinil. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 50 mcg treprostinil plus or minus 10 mcg, 9 mcg, 8 mcg, 7 mcg, 6 mcg, 5 mcg, 4 mcg, 3 mcg, 2 mcg or 1 mcg treprostinil for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 75 mcg treprostinil plus or minus 10 mcg, 9 mcg, 8 mcg, 7 mcg, 6 mcg, 5 mcg, 4 mcg, 3 mcg, 2 mcg or 1 mcg treprostinil for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 100 mcg treprostinil plus or minus 10 mcg, 9 mcg, 8 mcg, 7 mcg, 6 mcg, 5 mcg, 4 mcg, 3 mcg, 2 mcg or 1 mcg treprostinil for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 150 mcg treprostinil plus or minus 10 %, 9 %, 8 %, 7 %, 6 %, 5 %, 4 %, 3 %, 2 % or 1 % treprostinil for delivery to a

patient in a dry powder. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 200 mcg treprostinil plus or minus 10 %, 9 %, 8 %, 7 %, 6 %, 5 %, 4 %, 3 %, 2 % or 1 % mcg treprostinil for delivery to a patient in a dry powder. In further embodiments, a drug product according to the present invention may provide capsules with dose levels of 300 mcg treprostinil plus or minus 10 %, 9 %, 8 %, 7 %, 6 %, 5 %, 4 %, 3 %, 2 % or 1 % treprostinil for delivery to a patient in a dry powder. A summary of the LIQ861 formulation, including powder composition, particle geometry, and a description of the dosing unit according to certain exemplary embodiments follows.

LIQ861 Drug Product-Intermediate Description for Active (LIQ861) and Placebo Formulations (dihydrate)

Component	Function	Quantity (mg/g) (Active)	Percent Solids
Treprostinil Sodium	Drug Substance	5.3 (5.0 as treprostinil)	0.53
Trehalose Dihydrate	Bulking Agent	930	92.97
Polysorbate 80	Wetting Agent/Process Aide	20	20
L-Leucine	Hydrophobicity Modifier	40	40
Sodium Citrate Dihydrate	pH Modifier	2.7	0.27
Sodium Chloride	Buffer Component	2.3	0.23

Inhalation Drug Product Dosing Unit Description

Capsule	Size 3 Opaque White HPMC Capsule			
Fill Description	White to Off-White Powder			
Fill Particle Shape	"pollen-shaped"			
Active Strength (μg)	0 (placebo)*	25	50	75
Formulation Powder per Capsule (mg)	15	5	10	15

*Excipients only (no treprostinil). Abbreviations: HPMC, hydroxypropyl methylcellulose

[0057] According to some embodiments of the present invention, drug particles are provided that include a composition having a target dose of 15 – 90 μg of delivered treprostinil to the patient (current TYVASO® label is 18-54 μg). In some embodiments of the present invention the dose of treprostinil provided to the patient can be, for example, 100 micrograms, 125 micrograms or 150 micrograms. In some embodiments of the present invention the dose of treprostinil provided to the patient, for example, can contain about 100 micrograms, about 125 micrograms or about 150 micrograms. In some embodiments, each dose contains greater than or equal to 200 micrograms of treprostinil. In some embodiments, each dose contains greater than or equal to 225 micrograms of treprostinil. In some embodiments, each dose contains greater than or equal to 250 micrograms of treprostinil. In some embodiments, each dose contains greater than or equal to 275 micrograms of treprostinil. In some embodiments, each dose contains greater than or equal to 300 micrograms of treprostinil. In some embodiments, each dose contains from about 10 micrograms to about 15 micrograms, 15 micrograms to about 20 micrograms, 20 micrograms to about 25 micrograms, 25 micrograms to about 30 micrograms, about 30 micrograms to about 35 micrograms, about 35 micrograms to about 40 micrograms, about 40 micrograms to about 45 micrograms, about 45 micrograms to about 50 micrograms, about 50 micrograms to about 55 micrograms, about 55 micrograms to about 60 micrograms, about 60 micrograms to about 65 micrograms, about 65 micrograms to about 70 micrograms, about 70 micrograms to about 75 micrograms, about 75 micrograms to about 80 micrograms, about 80 micrograms to about 85 micrograms, about 85 micrograms to about 90 micrograms, about 90 micrograms to about 95 micrograms, about 95 micrograms to about 100 micrograms, or

about 100 micrograms to about 105 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 110 micrograms, 110 micrograms to about 120 micrograms, 120 micrograms to about 130 micrograms, 130 micrograms to about 140 micrograms, about 140 micrograms to about 150 micrograms, about 150 micrograms to about 160 micrograms, about 160 micrograms to about 170 micrograms, about 170 micrograms to about 180 micrograms, about 180 micrograms to about 190 micrograms, about 190 micrograms to about 200 micrograms, about 200 micrograms to about 210 micrograms, about 210 micrograms to about 220 micrograms, about 220 micrograms to about 230 micrograms, about 230 micrograms to about 240 micrograms, about 240 micrograms to about 250 micrograms, about 250 micrograms to about 260 micrograms, about 260 micrograms to about 270 micrograms, about 270 micrograms to about 280 micrograms, about 280 micrograms to about 290 micrograms, about 290 micrograms to about 300 micrograms, about 300 micrograms to about 310 micrograms, about 310 micrograms to about 320 micrograms, about 320 micrograms to about 330 micrograms, about 330 micrograms to about 340 micrograms, or about 340 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 25 micrograms to about 400 micrograms of treprostinil. In some embodiments, each dose contains from about 25 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 25 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 50 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 200 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 225 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 250 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 275 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 50 micrograms to about 75

micrograms of treprostinil. In some embodiments, each dose contains from about 50 micrograms to about 100 micrograms of treprostinil. In some embodiments, each dose contains from about 50 micrograms to about 150 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 100 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 125 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 150 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 175 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 200 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 225 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 75 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 125 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 150 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 175 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 200 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 225 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 100 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 150 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 175 micrograms of treprostinil. In some embodiments, each dose contains from about 125

micrograms to about 200 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 225 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 125 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 175 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 200 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 225 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 150 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 200 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 225 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 175 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 200 micrograms to about 225 micrograms of treprostinil. In some embodiments, each dose contains from about 200 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 200 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 200 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains

from about 200 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 200 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 225 micrograms to about 250 micrograms of treprostinil. In some embodiments, each dose contains from about 225 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 225 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 225 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 225 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 250 micrograms to about 275 micrograms of treprostinil. In some embodiments, each dose contains from about 250 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 250 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 250 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 275 micrograms to about 300 micrograms of treprostinil. In some embodiments, each dose contains from about 275 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 275 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 300 micrograms to about 325 micrograms of treprostinil. In some embodiments, each dose contains from about 300 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 325 micrograms to about 350 micrograms of treprostinil. In some embodiments, each dose contains from about 350 micrograms to about 375 micrograms of treprostinil. In some embodiments, each dose contains from about 375 micrograms to about 400 micrograms of treprostinil. In some embodiments, a patient may be provided with one, two, three, four, or more doses per day. In some embodiments, a patient may be provided up to one, two, three, or four doses per day. Each dose may be contained in a single capsule according to some embodiments, for example, a HPMC capsule (size 3). In other embodiments, a dose may be made up of a combination of lower dose capsules. In some embodiments, a patient may be provided with four doses per day to match the current treatment cycle (nebulized treprostinil) however the drug dose per treatment cycle under the present invention dry powder provides significantly higher dose levels to be safely administered, such as for example, up to 100 mcg of treprostinil per dosing, up to 125 mcg of treprostinil and up to 150 mcg of treprostinil per dosing

as each were surprisingly demonstrated in the first clinical trial of LIQ861. In alternative embodiments, a patient may be provided with four doses per day to match the current treatment cycle (nebulized treprostinil) however the drug dose per treatment cycle under the present invention dry powder provides significantly higher dose levels to be achieved, such as for example, up to 200 mcg of treprostinil per dosing and up to 300 mcg of treprostinil per dosing as surprisingly demonstrated in pre-clinical toxicology studies using LIQ861.

[0058] Treprostinil itself is poorly soluble in unbuffered water and low pH buffers. However, the solubility improves with increasing pH as the carboxylic acid is deprotonated. The sodium salt was selected for use in this product since it enhances dissolution in aqueous media and facilitates processing.

[0059] Excipients

[0060] According to some embodiments of the present invention, the DP-intermediate (anhydrous) is comprised of particles that include, for example, the following excipients: trehalose, polysorbate 80, L-leucine, sodium citrate, and sodium chloride. In some embodiments, the ratio of treprostinil sodium and excipients is 0.581:92.32:2.19:4.39:0.26:0.25 (wt:wt solids) treprostinil sodium:trehalose:polysorbate 80:leucine:sodium citrate:sodium chloride. A summary of the function, quantity, and compendial status of these excipients is provided herein.

[0061] The excipients were selected based upon the following functional requirements for the formulation:

- Trehalose Dihydrate: Trehalose comprises the bulk of the particle and was selected because it is a non-reducing sugar with a high glass transition temperature. Trehalose is an example of a non-reducing sugar (as opposed to lactose, which is a reducing sugar) that can be used in the present invention. Trehalose is more chemically compatible with compounds containing primary amines, such as leucine.
- Ultra-Pure Polysorbate 80 (Ultra-Pure Tween 80): Polysorbate 80 is added as a processing aide / wetting agent to facilitate particle manufacturing. In some embodiments, Polysorbate 80 is a particle processing aide and enables film generation

during particle manufacture by decreasing dewetting, leading to uniform particle morphology.

- L-leucine: Leucine is added as a hydrophobicity and surface modifier to reduce the hygroscopicity of the particle and improve aerosol efficiency. L-leucine is an example of a formulation additive to reduce hygroscopicity to improve stability of the final drug product powder.
- Sodium chloride and sodium citrate: Sodium citrate and sodium chloride are used to buffer the stock solution used in the PRINT Technology manufacturing process and to help control acidity in the particle. Sodium chloride and sodium citrate are examples of buffers that help maintain pH and control ionization/acidity of the formulation. In some embodiments of the present invention, pH is maintained between about pH 6.0 and 7.2.

[0062] In addition to the active pharmaceutical ingredient the present drug particle comprises a bulking agent, wetting agent, hydrophobicity modifier, pH modifier and buffer. In some embodiments, the present drug particle comprises, along with the active ingredients, a bulking agent, hydrophobicity controlling agent, and a pH controlling agent.

[0063] According to another embodiment of the present invention, LIQ861 contains five excipients as follows: treprostinil sodium:trehalose dihydrate:leucine:polysorbate 80:sodium citrate dihydrate:sodium chloride at ratios of 0.53:92.97:4:2:0.27:0.23. At an example treprostinil dose level of 100 μg/day of the present invention drug particles, a patient would receive the following daily excipient doses:

- 18.6 mg of trehalose dihydrate. Assuming a patient weighs 60 kg and has a lung mass of 1000 g, this is equivalent to 310 μg/kg and 18.6 μg/g of lung.
- 0.4 mg of polysorbate 80. Assuming a patient weighs 60 kg and has a lung mass of 1000 g, this is equivalent to 6.7 μg/kg and 0.4 μg/g of lung.
- 0.8 mg of leucine. Assuming a patient weighs 60 kg and has a lung mass of 1000 g, this is equivalent to 13.3 μg/kg and 0.8 μg/g of lung.

- 0.05 mg of sodium citrate and 0.05 mg of sodium chloride. Assuming a patient weighs 60 kg and has a lung mass of 1000 g, this is equivalent to 0.83 µg/kg for each compound and 0.05 µg/g of lung for each compound.

[0064] Formulation Development

[0065] According to embodiments of the present invention, LIQ861 has been developed as a novel formulation of treprostinil for the treatment of PAH. Treprostinil is currently approved for use in the treatment of PAH by subcutaneous, IV, oral, and inhalation routes of administration. TYVASO is currently the only marketed inhaled formulation of treprostinil and is formulated as a liquid solution for administration using a nebulizer. The nebulized treprostinil is dosed, at maintenance dose, of 6 mcg drug per breath over 9 breaths for a dose of 54 mcg per dosing session. The nebulized treprostinil also has a maximum tolerated dose of 84 mcg over a dosing session with 14 breaths.

[0066] LIQ861 is suitable for inhaled administration using a dry powder inhalation device. The physicochemical properties and performance characteristics, manufacturing process and packaging, and stability characteristics of the DP have been studied, and a suitable formulation has been identified for progression into human studies.

[0067] Physiochemical and Biological Properties

[0068] The "pollen-shaped" LIQ861 particles according to certain embodiments have an aerodynamic size to enable efficient delivery to the pulmonary arterioles ($1 \leq MMAD \leq 5\mu m$) with a high FPF to limit oropharyngeal deposition. A scanning electron microscopy (SEM) image of the "pollen-shaped" feature is provided in Figure 9. The formulation of example particles shown in Figure 9 is: treprostinil:trehalose:leucine:polysorbate 80:sodium citrate: sodium chloride (Batch LKI-1R-983-27). Example aerosol data for the active particles are also provided in the table below.

[0069] During the development of the LIQ861 formulation, the applicants tested other possible particle shapes and sizes (e.g., 1.5 µm donut, 3.0 µm donut). Based upon these studies, the applicants observed that the "pollen-shaped" feature resulted in a greater FPF, reduced

MMAD, acceptable ED, and dose uniformity characteristics when compared to other features both with and without treprostinil.

Representative Aerosol Data (NGI) for Active Particles

Sample	MMAD (μm)	GSD	ED (% nominal)	FPF (% ED)
Treprostinil Sodium: Trehalose:Leucine:Polysorbate 80:Sodium Citrate:Sodium Chloride (“pollen-shaped”)	1.88	1.99	64	83

Abbreviations: NGI, Next Generation Impactor™, MSP Corp.; MMAD, mass median aerodynamic diameter; GSD, geometric standard deviation; ED, emitted dose; FPF, fine particle fraction; wt, weight. Batch LKI-1R-0983-21.

[0070] Manufacture

[0071] The manufacturing of LIQ861 particles according to some embodiments of the present invention is described below. A process flow diagram for the particles (also referred to as DP-intermediate) according to some embodiments is shown in Figure 10.

[0072] In particular embodiments, the particles of the present disclosure are fabricated using PRINT® Technology (Liquidia Technologies, Inc., Morrisville, NC) particle fabrication. In particular, the particles are made by molding the materials intended to make up the particles in mold cavities.

[0073] In some embodiments, the molds can be polymer-based molds and the mold cavities can be formed into any desired shape and dimension. Uniquely, as the particles are formed in the cavities of the mold, the particles are highly uniform with respect to shape, size, and composition. Due to the consistency among the physical and compositional makeup of the particles of the present compositions, the compositions of the present disclosure provide highly uniform release rates and dosing ranges. Methods and materials that may be used for fabricating the particles according to embodiments of the present disclosure are further described and disclosed in issued patents and co-pending patent applications, each of which are incorporated herein by reference in its entirety: U.S. Pat. Nos. 8,518,316; 8,444,907; 8,420,124; 8,268,446; 8,263,129; 8,158,728; 8,128,393; 7,976,759; U.S. Pat. Application Publications Nos. 2013-0249138, 2013-0241107, 2013-0228950, 2013-0202729, 2013-0011618, 2013-0256354, 2012-

0189728, 2010-0003291, 2009-0165320, 2008-0131692; and pending U.S. Application Nos. 13/852,683 filed March 28, 2013 and 13/950,447 filed July 25, 2013.

[0074] Particle Fabrication

[0075] An aqueous stock solution is prepared at the desired total solids concentration. All other excipients are combined with treprostinil and then filtered prior to particle fabrication.

[0076] The stock solution is applied in a thin layer to a continuous polyethylene terephthalate (PET) substrate backing layer. Forced air heat is used to drive off the water resulting in a dry film of treprostinil and excipients. The dried film is then brought into contact with a mold film, having cavities of the desired shape and size which the drug product particles will mimic, at an elevated temperature. The drug/excipient blend flows into the cavities of the mold, conforming to the shape defined by the cavity. The result is a uniform array of particles adhered to a PET backing layer. The particles are then allowed to cool to room temperature as the roll is wound up for later collection.

[0077] In one example of the present drug particles, the following stock solution is used:

Stock solution components used for manufacture of treprostinil particles, according to an embodiment:

Stock component	Target Solution Concentration (Active)	Target Solution Concentration (Placebo)	Target
Trehalose	12%	12.7%	Adjusted based on mass balance of other formulation components
Leucine	0.52%	0.54%	0.52-0.54% (4% solids)
Treprostinil Sodium	0.069%	0 %	0.069% (0.53% solids)
Polysorbate 80	0.26%	0.27%	0.26-0.28% (2% solids)

Sodium Citrate	0.035%	0.037%	Maintain pH stock solution for stability of treprostinil
NaCl	0.030%	0.031%	Maintain tonicity of stock solution
Diluent (water)	87.0%	86.4%	86-91% evaluated; to coat appropriate formulation mass for processing and solubility of excipient component(s)

[0078] Dry Collection and Drying

[0079] Next, the particles are dry collected, the process of removing the molded particles from the PET backing layer and thereby creating a bulk powder. The mold is first separated from the PET backing layer, exposing the particle array attached to the PET backing layer. The particle array is then passed across a blade, in some embodiments a plastic blade, to dislodge the particles from the backing layer. The particles can then be collected into a bulk powder for further processing.

[0080] Humidity is controlled to less than 15% RH during collection, in some embodiments due to the hygroscopicity of the powder. Temperature is maintained at ambient, typically between 15 and 25^{o}C.

[0081] Drying and Bulk Packaging

[0082] The drug particles are dried at less than or equal to 150 mTorr of nitrogen or dry air for at least 2 days in a benchtop lyophilizer at room temperature, according to some embodiments.

[0083] In some embodiments, the particles of the present invention are dried to less than about 10 percent water content. In some embodiments, the particles of the present invention are dried to less than about 5 percent water content. In further embodiments, the particles of the present invention are dried to less than about 4 percent water content. In still further embodiments, the

particles of the present invention are dried to less than about 2 percent water content. In a preferred embodiment, the product is dried to less than about 1 percent water content by Karl Fisher titration.

[0084] Batch-to-Batch Uniformity of Drug Particles

[0085] In some embodiments, the particle uniformity from batch-to-batch provides the present invention with an unexpected and exceptional advantage over the prior art. In certain embodiments, the uniformity within any given batch is unexpected and exceptionally advantageous over the prior art. The present invention includes highly conserved batch uniformity as shown in the following data. See the table below and also Figure 2.

Uniformity**:** Sample aerosol data (NGI) for active particles (PAH-1R-0974-010)

Sample	MMAD	GSD	ED (%rec)	FPF (%ED)	F345 (fill)
PAH-1R-0974-010-A1 First	1.74	1.88	81%	88%	42%
PAH-1R-0974-010-B1 Middle	1.80	1.87	78%	87%	44%
PAH-1R-0974-010-C1 Last	1.72	1.87	90%	89%	40%

In the example shown, fine particle fraction remained within plus/minus 1 percent within a single batch run.

[0086] Capsule Filling and Packaging

[0087] In some embodiments, HPMC capsules are filled with the DP-intermediate in a humidity controlled ISO 8 environment using an XCELODOSE® (Capsugel) instrument. The filled HPMC capsules are packaged in a low humidity environment. Ten capsules are placed in a DESICAP® Vial and closed with a DESICAP® Cap. The closed vial is then placed into a foil bag with a desiccant canister prior to heat sealing the foil bag to form the packaged drug product.

[0088] Stability studies

[0089] According to the formulation of the present drug particle, it was desired to minimize uncontrolled exposure to ambient humidity. The drug particles according to embodiments of the present invention are shown to be stable for at least 9 months when stored under controlled humidity conditions at 25°C/60%RH. In some embodiments, the drug particles are shown to be stable for at least 6 months when stored under controlled humidity conditions at 40°C/75%RH. In some embodiments, the drug particles are shown to be stable for at least 9 months when stored under desiccated conditions at 25°C/60%RH. In some embodiments, the drug particles are shown to be stable for at least 6 months when stored under desiccated conditions at 40°C/75%RH. Studies were conducted to determine the stability of the drug particles at 25 ºC/60% RH and 40 ºC/75% RH.

[0090] Prototype Stability Study

[0091] The purpose of the Prototype Stability Study was to evaluate the stability of drug particles in capsules. Both the 25 and 75 µg strengths were evaluated when stored at 25 ºC/60% RH and 40 ºC/75% RH. For the study, drug particles were placed into size 3 HPMC opaque capsules (Capsugel Vcaps). Ten filled capsules were placed into HDPE vials (Desicap) which were sealed with a stopper. The stoppered vial was placed into a foil overwrap with desiccant sachets.

[0092] Data for the 25 µg dose drug particles stored at 25°C/60%RH is shown in the table below.

Test	Specifications		Time Points				
			Initial	1 Month	3 Months	6 Months	9 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.489	0.520	0.493	0.494	0.486
Aerodynamic Particle Size Distribution	Report Results	MMAD (µm)	1.92	2.11	2.2	2.1	2.1
		GSD (µm)	1.72	1.67	1.6	1.6	1.7
		FPF (%)	87	84	82.9	83.6	83.6

Delivered Dose Uniformity	Report Results	Average (µg)	19.9	21.6	19.65	19.47	18.86

[0093] Data for the 25 µg dose drug particles stored at 40°C/75%RH is shown in the table below.

Test	**Specifications**		**Time Points**			
			Initial	**1 Month**	**3 Months**	**6 Months**
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.489	0.512	0.502	0.492
Aerodynamic Particle Size Distribution	Report Results	MMAD (µm) GSD (µm) FPF (%)	1.92 1.72 87	2.09 1.65 85	2.1 1.6 84.8	2.1 1.6 84.7
Delivered Dose Uniformity	Report Results	Average (µg)	19.9	21.2	18.86	19.28

[0094] Data for the 75 µg dose drug particles stored at 25°C/60%RH is shown in the table below.

Test	**Specifications**		**Time Points**				
			Initial	**1 Month**	**3 Months**	**6 Months**	**9 Months**
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.489	0.500	0.496	0.494	0.487
Aerodynamic Particle Size Distribution	Report Results	MMAD (µm) GSD (µm)	2.13 1.60 85	2.17 1.61 84	2.2 1.6 84.2	2.2 1.6 83.7	2.2 1.6 82.3

		FPF (%)					
Delivered Dose Uniformity	Report Results	Average (µg)	63.6	63.0	60.76	59.62	60.01

[0095] Data for the 75 µg dose drug particles stored at 40°C/75%RH is shown in the table below.

Test	**Specifications**		**Time Points**			
			Initial	**1 Month**	**3 Months**	**6 Months**
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.489	0.509	0.506	0.491
Aerodynamic Particle Size Distribution	Report Results	MMAD (µm)	2.13	2.14	2.2	2.1
		GSD (µm)	1.60	1.61	1.6	1.6
		FPF (%)	85	85	84.8	85.3
Delivered Dose Uniformity	Report Results	Average (µg)	63.6	61.0	59.86	59.42

[0096] Clinical Trial Material Stability Study

[0097] The purpose of the Clinical Trial Material Stability Study was to evaluate the stability of drug particles in capsules. Three strengths were evaluated: 25, 50, and 75 µg active agent doses within capsules. As in the previous study, two storage conditions were evaluated: 25 ºC/60% RH and 40 ºC/75% RH. For the study, drug particles were placed into size 3 HPMC opaque capsules (Capsugel Vcaps). Ten filled capsules were placed into HDPE vials (DESICAP) which were sealed with a stopper. The stoppered vial was placed into a foil overwrap with desiccant sachets.

[0098] Data for the 25 µg dose drug particles stored at 25°C/60%RH is shown in the table below.

Test	Specifications		Time Points			
			Initial	1 Month	3 Months	6 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.520	0.505	0.504	0.504
Aerodynamic Particle Size Distribution	Report Results	MMAD (μm) GSD (μm) FPF (%)	2.3 1.6 84.1	2.3 1.6 82.7	2.2 1.6 83.6	2.1 1.6 85.0
Delivered Dose Uniformity	Report Results (μg)	Average (μg)	19.784	20.48	19.24	19.47

[0099] Data for the 25 μg dose drug particles stored at 40°C/75%RH is shown in the table below.

Test	Specifications		Time Points			
			Initial	1 Month	3 Months	6 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.520	0.518	0.501	0.506
Aerodynamic Particle Size Distribution	Report Results	MMAD (μm) GSD (μm) FPF (%)	2.3 1.6 84.1	2.2 1.6 84.1	2.2 1.6 85.8	2.2 1.6 84.7
Delivered Dose Uniformity	Report Results	Average (μg)	19.784	20.73	18.89	18.80

[00100] Data for the 50 μg dose drug particles stored at 25°C/60%RH is shown in the table below.

Test	Specifications		Time Points			
			Initial	1 Month	3 Months	6 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.515	0.509	0.509	0.506
Aerodynamic Particle Size Distribution	Report Results	MMAD (μm) GSD (μm) FPF (%)	2.2 1.6 86.2	2.2 1.6 85.6	2.2 1.6 85.7	2.2 1.6 85.3
Delivered Dose Uniformity	Report Results (μg)	Average (μg)	40.417	40.75	39.14	40.05

[00101] Data for the 50 μg dose drug particles stored at 40°C/75%RH is shown in the table below.

Test	Specifications		Time Points			
			Initial	1 Month	3 Months	6 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.515	0.520	0.505	0.501
Aerodynamic Particle Size Distribution	Report Results	MMAD (μm) GSD (μm) FPF (%)	2.2 1.6 86.2	2.2 1.6 86.5	2.2 1.6 86.2	2.2 1.6 84.1
Delivered Dose Uniformity	Report Results	Average (μg)	40.417	39.55	38.96	37.50

[00102] Data for the 75 μg dose drug particles stored at 25°C/60%RH is shown in the table below.

Test	Specifications		Time Points			
			Initial	1 Month	3 Months	6 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.517	0.512	0.509	0.508
Aerodynamic Particle Size Distribution	Report Results	MMAD (μm) GSD (μm) FPF (%)	2.3 1.6 84.8	2.3 1.6 84.1	2.3 1.6 84.5	2.2 1.6 85.1
Delivered Dose Uniformity	Report Results (μg)	Average (μg)	61.851	63.91	59.88	60.25

[00103] Data for the 75 μg dose drug particles stored at 40°C/75%RH is shown in the table below.

Test	Specifications		Time Points			
			Initial	1 Month	3 Months	6 Months
Assay	0.450 – 0.550% w/w Treprostinil as free acid (%)		0.517	0.513	0.503	0.495
Aerodynamic Particle Size Distribution	Report Results	MMAD (μm) GSD (μm) FPF (%)	2.3 1.6 84.8	2.3 1.6 85.1	2.2 1.6 85.8	2.2 1.6 84.9
Delivered Dose Uniformity	Report Results	Average (μg)	61.851	61.94	58.61	58.17

[00104] Dry Powder Inhalation Device

[00105] The RS00 Model 8 is a commercially available monodose dry powder inhalation device that is manufactured by Plastiape S.p.A (Italy) in accordance with ISO and FDA standards. The overall design of RS00 Model 8 device is shown in Figure 11.

[00106] A cap, which is retained on the mouthpiece, is designed to prevent ingress of dirt and other foreign material into the inhaler when not in use. The plastic side portions cover the air inlet holes, but the cap does not provide a hermetic seal to the device. The cap does not form part of the actuation process.

[00107] When assembled, the mouthpiece is mounted on the inhaler body, but is removable for cleaning purposes. To assemble the mouthpiece, the off-set peg at the base of the mouthpiece is placed into the corresponding hole in the inhaler body and the mouthpiece is rotated until it snaps closed. The snap closure ensures that the mouthpiece and inhaler body are properly aligned and that no spurious airflow occurs. The mouthpiece contains a mesh that aids particle size reduction and prevent capsule ingestion during inhalation.

[00108] The inhaler body component contains two side buttons, each housing four pins for piercing a capsule. The pins are inserted in the corresponding housing of the pushbuttons and the heads of the pins are retained in their position by a back-plate that is ultrasonically welded to the pushbutton. The buttons and pins are each maintained in their outward position by four small steel springs in each button. A three-component snap-lock system on the inhaler body ensures correct alignment of the mouthpiece when closed.

[00109] A capsule piercing area is located internally, adjacent to the pins. When a capsule is inserted in this area, depressing the buttons causes the button pins to pierce the capsule ends, thereby preparing the capsule for emptying. Above the capsule piercing area, there are 2 tangential air inlets and a circular chamber. These allow the capsule to spin when the patient inhales through the device. Capsule spinning creates a centrifugal effect on the powder that promotes efficient emptying.

[00110] The performance of the premetered dry powder inhaler is a combination of the characteristics of LIQ861 (including the powder and capsule) and the inhalation device itself.

[00111] NONCLINICAL STUDIES

[00112] Treprostinil is a tricyclic benzidine analogue of endogenous PGI_2. The major pharmacologic actions of treprostinil are direct vasodilation of pulmonary and systemic arterial vascular beds and inhibition of platelet aggregation. It was developed for chronic administration as a continuous subcutaneous infusion as a treatment for patients with PAH. PGI_2, an endothelial cell derived substance, is a potent vasodilator and inhibitor of platelet aggregation. The hemodynamic properties of treprostinil are similar to those of PGI_2 but, unlike PGI_2, treprostinil is chemically stable.

[00113] A series of *in vivo* studies were conducted to evaluate pharmokinetics (PK) and toxicology of the present invention dry powder treprostinil formulation.

[00114] Pilot, non-GLP, single-dose, inhalation PK study of treprostinil in dogs (Study 19073)

[00115] This study compared the single dose PK of LIQ861 (administered via DPI) to Tyvaso (administered via nebulizer) at a target lung deposition of 3 μg/kg in 3 beagle dogs. Results showed generally similar treprostinil PK profiles following dosing with LIQ861 compared with Tyvaso. In this pilot single administration PK study, treprostinil (dry powder formulation; estimated lung deposition of 3.0 to 3.4 μg/kg) and treprostinil (nebulized liquid; target lung deposition of 3 μg/kg) were compared in 3 beagle dogs. The results showed generally similar treprostinil PK profiles following dosing with treprostinil (dry powder formulation) compared with treprostinil (nebulized liquid). The study design and results are discussed in more detail herein.

[00116] The applicants conducted a study comparing plasma concentrations and pharmacodynamics (PD) following administration of treprostinil sodium (nebulized liquid versus a dry powder formulation similar to the LIQ861 formulation of the present invention) as a single inhalation exposure (via controlled ventilation) to anesthetized beagle dogs. Treprostinil sodium was prepared as a nebulized liquid from the same DS used to prepare the dry powder formulation. The dry powder formulation was manufactured using PRINT Technology and utilized the same drug substance, treprostinil sodium, but was different in excipient concentrations compared to LIQ861. Importantly, the excipient concentrations of the present invention provide highly consistent and reproducible batch to batch manufacturing of the

LIQ861 product. The formulations used in this study will be referred to as treprostinil (nebulized liquid) and treprostinil (dry powder formulation), respectively, in description of this study. The study design, results, and conclusions are described below.

[00117] In study 19073, 3 dogs received a single inhalation administration of nebulized treprostinil (nebulized liquid; estimated lung deposition of 3.4 μg/kg). After a 2-day washout, the dogs received a single inhalation administration of treprostinil (dry powder formulation; estimated lung deposition of 3.0 to 3.4 μg/kg). Blood was collected for plasma analysis of treprostinil concentrations prior to each administration and at 2, 5, 10, 20, 30, 60, 120, and 180 minutes after the completion of each administration. In addition, 2 different dogs (one assigned to each treprostinil formulation) were used to monitor the following PD endpoints (hemodynamic changes): systemic arterial blood pressures [mean arterial pressure (MAP, mmHg), systolic arterial pressure (mmHg), diastolic arterial pressure (mmHg)], pulmonary artery pressure (PAP, mmHg), right atrial pressure (RAP, mmHg), pulmonary capillary wedge pressure (PCWP, mmHg) or left atrial pressure (mmHg), cardiac output (CO, L/min the average of 3), total peripheral resistance (TPR), pulmonary vascular resistance (PVR), and heart rate (HR). The PD effects were assessed prior to initiation of dose administration and at target times of 5, 10, 20, 30, 60, 120, and 180 minutes after the completion of the administration. In contrast to the first three dogs, the dogs assigned to monitor the PD effects were anesthetized for the duration of data collection. The dog assigned treprostinil (nebulized liquid) received an estimated lung deposition of 4.0 μg/kg and the dog assigned to treprostinil (dry powder formulation) received an estimated lung deposition of 2.5 μg/kg. Blood was collected at the same time points as first 3 dogs.

[00118] Treprostinil (nebulized liquid and dry powder formulation) had no effect on HR, PAP, RAP, PCWP, or CO, but had a slight effect on decreasing and then increasing arterial blood pressure. Treprostinil (nebulized liquid) appeared to decrease stroke volume, increase TPR, and decrease PVR. Treprostinil (dry powder formulation) appeared to increase stroke volume, decrease TPR, and decrease PVR. The Study Director concluded that the pilot data were inconclusive for comparing the potential PD effects of treprostinil (nebulized liquid) to the treprostinil (dry powder formulation) formulation; however, there appeared to be no important differences in PD effects associated with administration of treprostinil in either formulation.

[00119] Pilot, non-GLP, single-dose, inhalation PK study of LIQ861 in male rats (Study 75670)

[00120] This study evaluated the PK of treprostinil in male rats following single inhalation of a range of LIQ861 doses up to a feasible dose. Systemic exposure data from this study was used to determine appropriate doses and blood sampling times for a definitive, comparative PK bridging study of LIQ861 and nebulized treprostinil. Results from this study were used to select dose levels and an optimal blood sampling paradigm for a definitive PK bridging study.

[00121] SUMMARY: STUDY 75670

[00122] The objective of the study was to determine the pharmacokinetic profile of treprostinil in male Sprague Dawley rats when administered as the test item, PRINT Treprostinil dry powder (PRINT-Tre), as a single 4 hour inhalation at targeted dose levels of 0.15, 0.75, and 1.5 mg/kg. Results from this study will be used to determine appropriate dose levels and sampling time points for a definitive PK bridging study.

[00123] The test item was administered once by inhalation to 3 male rats per group as described in the table below:

Group No.	Group Designation	Achieved Mean Total Inhaled Dose Level of Treprostinil (mg/kg/day)	Achieved Aerosol Concentration of Treprostinil (µg/L)	Achieved Aerosol Concentration of Trehalose (µg/L)
1	Low Dose	0.158	1.06	150.46
2	Mid Dose	0.707	4.72	664.85
3	High Dose	1.409	9.39	1298.81

[00124] Assessments of mortality, clinical signs and body weights were performed. Blood samples were collected and analyzed for treprostinil content.

[00125] No mortality occurred. No clinical signs were observed and body weights were unaffected.

[00126] The overall achieved gravimetric and analytical aerosol concentrations for all groups were within 16% of the targeted concentrations. Corresponding average treprostinil dose levels for all groups were within 7% of the targeted dose levels and a clear dose differentiation between groups for each sex was achieved. The gravimetric particle size MMADs from all groups were between 1.2 and 1.6 μm (GSD 2.06 to 2.56). For both treprostinil and trehalose, the chemical determination of particle size distribution ranged from 1.3 to 1.8 μm with the corresponding GSDs between 1.65 and 2.15. The particle size distribution was considered respirable gravimetrically and chemically.

[00127] Mean PK parameters for PRINT-Tre treatment groups obtained by non-compartmental analysis of the mean treprostinil plasma concentration data sets are summarized as follows:

Group		$T_{1/2}$ (hr)	T_{max} (hr)	C_{max} (ng/mL)	$AUC_{0\text{-}Tlast}$ (hr*ng/mL)	AUC_{INF} (hr*ng/mL)
1	Mean	1.01	3.75	6.800	17.320	18.335
	SD	0.521	0.00	0.951	2.281	1.806
	N	3	3	3	3	3
2	Mean	1.68	3.75	31.933	81.289	93.369
	SD	0.967	0.00	9.500	19.478	19.372
	N	3	3	3	3	3
3	Mean	1.48	3.75	46.130	121.285	137.512
	SD	0.619	0.00	20.580	53.331	53.418
	N	3	3	3	3	3

[00128] In conclusion, single inhalation administration for 4 hours of PRINT-Tre at a high average treprostinil dose of 1.409 mg/kg/day by Sprague-Dawley rats was well tolerated as there were no significant test item related findings. The exposure to treprostinil generally increased in a dose proportional manner between the low dose and the mid dose. The exposure between the mid and high dose increased in a slightly less than dose proportional manner. However, animals in the high dose group were exposed to aerosol concentrations far below target for the last 16 to 26 minutes of inhalation, which may account for the less than dose proportional increase in

exposure. Based on these results, similar dose levels are recommended for the following definitive PK study. Blood sampling time points during the test item inhalation period may be adjusted so as to better characterize exposure during test item administration.

[00129] INTRODUCTION

[00130] The objective of the study was to determine the pharmacokinetic profile of treprostinil in male Sprague Dawley rats when administered as the test item, PRINT Treprostinil dry powder (PRINT-Tre), as a single 4 hour inhalation at targeted dose levels of 0.15, 0.75, and 1.5 mg/kg. Results from this study will be used to determine appropriate dose levels and sampling time points for a definitive PK bridging study.

[00131] The study was not performed in compliance with GLP regulations but followed appropriate Standard Operating Procedures (SOPs).

[00132] EXPERIMENTAL DESIGN

[00133] The test item was administered to groups of rats by inhalation administration for one day as described in the table below:

Group No.	Group Designation	Targeted Total Inhaled Dose Level of Treprostinil (mg/kg/day)[a]	Targeted Aerosol Concentration of Treprostinil (µg/L)	Targeted Aerosol Concentration of Trehalose (µg/L)	No. of Animals
					Males
1	Low Dose	0.15	1	130.7	3
2	Mid Dose	0.75	5	653.5	3
3	High Dose	1.5	10	1306.9	3

[a] = Targeted aerosol concentrations were calculated based on an estimated body weight of 0.250 kg

[00134] Following dosing, a series of 6 blood samples for pharmacokinetic evaluation were taken.

- Characterization of Test Item

Test item*:	Identity:	PRINT Treprostinil
	Content:	92.75% of Trehalose, 4% of Leucine, 2% of Tween80, 0.26% of NA Citrate Dihydrate, 0.25% of NaCl : 0.74% of Treprostinil sodium (0.67% treprostinil)
	Storage Conditions:	Cool (2 to 8°C), protect from moisture (e.g., dessicant)
	Handling Precautions:	Standard laboratory precautions. Handle under dry conditions (Relative Humidity ≤ 23%)
	Supplier:	Liquidia Technologies Inc.

[00135] TREATMENT

[00136] Acclimatization to Exposure System

[00137] Before the animals were exposed to the aerosol of the test item, rats were accustomed to the restraint procedure over a period of 3 days. The animals were gradually accustomed to restraint in the dosing tubes used during the exposures up to the duration that was used for aerosol administrations.

[00138] Animal Exposure

Exposure system used:	Flow-past rodent inhalation exposure system
Exposure method:	Inhalation by nose-only exposure
Test Item type:	Dry-Powder formulation
Generation method:	Piston feed/rotating brush generator
Duration of exposure:	240 minutes

[00139] The target aerosol concentrations and dose levels were as follows:

Group No.	Group Designation	Targeted Dose Level of Treprostinil (mg/kg/day)[a]	Targeted Aerosol Concentration of Treprostinil (µg/L)	Targeted Aerosol Concentration of Trehalose (µg/L)
1	Low Dose	0.15	1	130.7
2	Mid Dose	0.75	5	653.5
3	High Dose	1.5	10	1306.9

a = Target aerosol concentrations were calculated based on an estimated body weight of 0.250 kg.

[00140] ESTIMATION OF ACHIEVED DOSE LEVELS

The target dose levels were estimated using the following formula:

$$D_L = \frac{E_c \times RMV \times T}{BW}$$

D_L = Achieved dose levels (mg / kg / day)

E_c = Actual concentration delivered to the animals (mg / L air)

RMV = Respiratory Minute Volume (L / min) according to the method of Bide, Armour and Yee J. App. Toxicol., Vol. 20, 2000: *RMV* (L / min) = 0.499 x *BW* (kg)$^{0.809}$

T = Time, duration of daily exposure (min.)

BW = Mean body weight (kg) during exposure period.

This estimation of total inhaled dose assumed 100% deposition within the respiratory tract.

[00141] Inhalation Exposure System

[00142] The powder aerosol was produced using a piston feed / rotating brush generator. The aerosol produced was diluted as necessary to achieve the target aerosol concentration and discharged through a 40-mm diameter tube into a flow-past inhalation exposure system. The airflow rate through the exposure system was monitored and recorded manually during each aerosol generation period. Airflow to the exposure system was controlled by the absolute volume of air supplying the generation apparatus using variable area flowmeters. Control of the

aerosol exhaust flow from the animal exposure system was achieved using an exhaust valve, and the overall balance of airflows in the exposure system was monitored using pressure gauges. The system provided a minimum of 1.0 L/min to each animal exposure port and was balanced to ensure a slight positive pressure at the site of the proposed animal exposure. This ensured that there was no dilution of the generated aerosol. An equal delivery of aerosol to each proposed exposure position was achieved by employing a distribution network that was identical for each individual exposure position attached to the system.

[00143] Inhalation System Monitoring

[00144] Determinations of aerosol concentration, particle size distribution, oxygen concentration, relative humidity and temperature were measured on samples collected from a representative port of the exposure chamber. The sample flow rates were precisely controlled using variable area flow meters that were calibrated before use using a primary airflow calibrator. The absolute volume of each aerosol concentration sample was measured with a wet type gas meter.

[00145] Oxygen Concentration

[00146] The oxygen concentration of the generated atmosphere was measured once during each aerosol exposure. Oxygen concentrations of the exposure atmospheres were maintained between 19-23%.

[00147] Relative Humidity/Temperature

[00148] The temperature and relative humidity of the generated atmosphere were measured once during each aerosol exposure. Temperatures of the exposure atmospheres were maintained between 19-24°C.

[00149] Determination of Aerosol Concentration

[00150] At least one aerosol concentration filter sample was collected on glass fiber filter and weighed on each day in order to measure the gravimetric concentration of the test item in the generated aerosol. The filter samples were transferred to the analytical chemistry laboratory for

chemical determination of Treprostinil and Trehalose concentrations using an analytical method (Study No. 41609 and Study No. 41635).

[00151] Determination of the Particle Size Distribution and Mass Median Aerodynamic Diameter (MMAD)

[00152] The distribution of particle size in the generated aerosols was measured once during each exposure by collecting samples into a 7-Stage Mercer Cascade Impactor. The MMAD and the Geometric Standard Deviation (GSD) were calculated based on the results obtained from the impactor using a log-probit transformation.

[00153] IN-LIFE OBSERVATIONS

[00154] Mortality

[00155] Mortality checks were performed at least once a day during all phases of the study.

[00156] Clinical Observations

[00157] Cage-side clinical signs (ill health, behavioral changes etc.) were recorded at least once daily during all phases of the study, except on detailed clinical examination days, where the cage-side clinical signs were replaced by a DCE.

[00158] A detailed clinical examination of each rat was performed on arrival as part of the health status, as well as on Day 1, prior to dosing.

[00159] Animal whose health status was judged to warrant additional evaluation was examined by a Clinical Veterinarian.

[00160] Body Weights

[00161] Body weights were recorded for all animals once at arrival as per health status, once prior to group assignment and on Day 1 (prior to dosing).

[00162] Pharmacokinetics

[00163] A series of 6 blood samples (approximately 0.3 mL each) was collected from each rat on Day 1 at -15, 5, 15, 30, 75 and 105 minutes after treatment. Thus a total blood volume of 1.8 mL was taken from each rat during the course of the study. For this purpose, each rat (unanesthetized) was bled by jugular venipuncture and the samples were collected into tubes containing the anticoagulant, K_2EDTA. Tubes were placed on wet ice pending processing.

[00164] Following collection, the samples were centrifuged (2500 rpm for 10 minutes at approximately 4°C) and the resulting plasma was recovered and stored frozen (≤-60°C) in labeled tubes.

[00165] Deviations to the pharmacokinetic time points were noted in the raw data and were made available with the samples. The location of blood withdrawal was noted in the raw data.

[00166] Non-compartmental analysis of treprostinil concentrations in plasma were performed by using the Phoenix WinNonlin 6.3 software.

[00167] The following configuration was used for the analysis:

Sampling Method:	Sparse
AUC Calculation Method:	Linear Trapezoidal with Linear Interpolation
Lambda Z (λ_z) Method:	Best fit for λz, Log regression
Weighting (λ_z calculation):	Uniform

[00168] Pharmacokinetic parameters (including abbreviation and description for each parameter) are described in the following table:

Parameters	Abbreviation	Unit*
Area under the plasma drug concentration-time curve from the time of dosing to the last quantifiable concentration	$AUC_{0\text{-}Tlast}$	µg*hr/mL
Area under the plasma drug concentration-time curve from the time of dosing extrapolated to infinity	AUC_{INF}	µg*hr/mL
Terminal elimination half-life	$T_{1/2}$	hr
The maximum plasma concentration	C_{max}	µg/mL
Time to maximum plasma concentration	T_{max}	hr

*Different units may be presented in the study report

[00169] DATA EVALUATION AND STATISTICS

[00170] Numeric and non-numeric data obtained during the study were reported only as individual values.

[00171] RESULTS

[00172] Aerosol Concentrations

[00173] Achieved gravimetric test atmosphere concentrations were as follows:

Group No.	**Targeted Aerosol Concentration (mg\L)**	**Achieved Mean Aerosol Concentration (mg\L)**	**Coefficient of Variation (%)**	**% of Target**
1	0.156	0.165	17.2	105.9
2	0.781	0.728	14.0	93.2
3	1.563	1.439	43.5	92.0

[00174] Achieved analytical test atmosphere concentrations for treprostinil were as follows:

Group No.	**Targeted Aerosol Concentration (µg\L)**	**Achieved Mean Aerosol Concentration (µg\L)**	**Coefficient of Variation (%)**	**% of Target**
1	1	1.06	17.6	105.9

2	5	4.72	13.9	94.4
3	10	9.39	43.6	93.9

[00175] Achieved analytical test atmosphere concentrations for trehalose were as follows:

Group No.	Targeted Aerosol Concentration (µg\L)	Achieved Mean Aerosol Concentration (µg\L)	Coefficient of Variation (%)	% of Target
1	130.7	150.46	18.8	115.1
2	653.5	664.85	15.0	101.7
3	1306.9	1298.81*	44.1	99.4

*Last 2 aerosol concentrations samples for trehalose were estimated with a 92.79% difference from gravimetric data as analytical results were BLQ

[00176] The overall achieved gravimetric and analytical aerosol concentrations for all groups were within 16% of the targeted concentrations. The generated atmospheres were considered stable over the treatment period as % CV were all below 20%, except for Group 3. The increased % CV for Group 3 was caused by the stoppage of the Rotating Brush Generator (RBG) due to lack of test item remaining in the canister with 26 minutes left in the generation (16 minutes of dosing left for animal 3001A, 21 minutes left for animal 3002A and 26 minutes left for animal 3003A). Though a new test item canister was installed on the RBG apparatus, the aerosol concentrations were much lower than targeted for the last 26 minutes. However, the overall aerosol concentrations were still considered acceptable for the study as there was a significant difference in aerosol concentration between groups.

[00177] Dose Levels

[00178] Overall achieved doses for treprostinil are presented below:

Group No.	Targeted Dose Levels (mg/kg/day)	Duration of Exposure (min)	Animal	Body Weight (kg)	Estimated Achieved Doses (mg/kg/day)	% from Targeted Dose Level
1	0.15	240	1001A	0.326	0.157	104.7
			1002A	0.309	0.159	106.0
			1003A	0.314	0.158	105.3
			Average		*0.158*	*105.3*
2	0.75	240	2001A	0.319	0.703	93.7
			2002A	0.308	0.708	94.4
			2003A	0.304	0.709	94.5
			Average		*0.707*	*94.3*
3	1.5	240	3001A	0.322	1.396	93.1
			3002A	0.321	1.397	93.1
			3003A	0.281	1.433	95.5
			Average		*1.409*	*93.9*

[00179] Average achieved dose levels for all groups were within 7% of the targeted dose levels therefore the dose levels were considered acceptable for the study as a clear dose differentiation between groups for each sex was achieved.

[00180] Particle Size Distribution

[00181] The average gravimetric particle size distribution measurement data were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
1	95.8	92.4	82.4	45.4	32.4	21.5	12.3	0.0	1.2	2.28	93
2	86.8	83.3	76.0	39.4	28.0	15.4	8.9	0.0	1.5	2.56	85

3	94.0	90.9	77.3	30.3	13.4	8.9	5.3	0.0	1.6	2.06	90

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00182] The average chemical determination of particle size distribution for treprostinil were as follows:

Group No.	**Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)**								**Mean**		**% below 4 µm**
	4.60	**3.00**	**2.10**	**1.60**	**1.10**	**0.70**	**0.33**	**0.00**	**MMAD (µm)**	**GSD**	
1	96.4	93.9	83.3	42.8	28.1	16.5	6.9	0.0	1.3	2.06	94
2	91.7	88.5	81.1	37.9	25.8	11.8	4.9	0.0	1.5	2.15	90
3	95.2	91.5	76.6	25.7	8.8	5.1	1.8	0.0	1.7	1.86	91

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00183] The average chemical determination of particle size distribution for trehalose were as follows:

Group No.	**Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)**								**Mean**		**% below 4 µm**
	4.60	**3.00**	**2.10**	**1.60**	**1.10**	**0.70**	**0.33**	**0.00**	**MMAD (µm)**	**GSD**	
1	95.8	91.6	81.5	38.4	24.1	13.0	4.2	0.0	1.4	2.01	93
2	94.4	88.9	83.3	31.5	26.0	11.1	5.6	0.0	1.4	2.11	91
3	95.2	90.4	77.0	25.5	9.6	4.8	0.0	0.0	1.8	1.65	94

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00184] The particle size distribution was considered respirable for this study as the MMADs were below 4 μm and the GSD were within 1.5 and 3.

[00185] Exposure Chamber Conditions

[00186] Exposure chamber conditions from the reported aerosol concentration exposures are summarized below.

Group No.	Humidity (%RH)	Temperature (°C)	Oxygen Concentration (%)
1	31.4	21.2	20.9
2	34.2	21.7	20.9
3	26.2	20.7	20.9

[00187] Exposure atmosphere oxygen concentrations, temperature and relative humidity ranges were considered acceptable on all occasions.

[00188] Mortality

[00189] There were no mortalities during the study.

[00190] Clinical Signs

[00191] There were no adverse clinical signs observed during the study.

[00192] Slight decreased activity, piloerection and partially closed eyes were seen in animal 3001A right before the 15 minute time point. However these were not observed afterwards and were not observed in any other animal therefore were not deemed test item related.

[00193] Body Weight

[00194] Body weights were performed for dose level calculation purposes.

[00195] Pharmacokinetics

[00196] Following the administration of PRINT-Tre at all achieved dose levels, mean C_{max} ranged from 6.800 to 46.133 ng/mL. The mean maximum plasma concentration (T_{max}) was reached at 3.75 hour (15 minutes before end of dosing) for all groups. The mean $AUC_{0\text{-}Tlast}$ (AUC_{INF}) ranged from 17.320 (18.335) to 121.258 (137.512) hr*ng/mL. Following T_{max}, the treprostinil plasma concentrations declined gradually with an estimated mean $T_{1/2}$ ranging from 1.01 to 1.68 hours.

[00197] Over the dose range, exposure to treprostinil (based on $C_{max,}$ $AUC_{0\text{-}Tlast}$ and AUC_{INF}) generally increased in dose proportional manner between the low dose (0.158 mg/kg) and the mid dose (0.707 mg/kg). When dose level increased 4.5-fold from low to mid dose, C_{max} and $AUC_{0\text{-}Tlast}$ increased 4.7-fold. Treprostinil exposure between the mid dose (0.707 mg/kg) and high dose (1.409 mg/kg) increased in a slightly less than dose proportional manner (2-fold increase in dose with a 1.4- (C_{max}) to 1.5-fold ($AUC_{0\text{-}Tlast}$) increase in exposure). However, because animals in the high dose group were exposed to aerosol concentrations far below target for the last 16 to 26 minutes of the exposure period, exposure levels may have been effected and could account for the less than dose proportional increase in exposure.

[00198] CONCLUSION

[00199] Single inhalation administration for 4 hours of PRINT-Tre at a high average treprostinil dose of 1.409 mg/kg/day to Sprague-Dawley rats was well tolerated as there were no significant test item related findings. The exposure to treprostinil generally increased in a dose proportional manner between the low dose and the mid dose. The exposure between the mid and high dose increased in a slightly less than dose proportional manner. However, animals in the high dose group were exposed to aerosol concentrations far below target for the last 16 to 26 minutes of inhalation, which may account for the less than dose proportional increase in exposure. Based on these results, similar dose levels are recommended for the following definitive PK study. Blood sampling time points during the test item inhalation period may be adjusted so as to better characterize exposure during test item administration.

[00200] Non-GLP, single-dose, inhalation, comparative PK study of LIQ861 and nebulized treprostinil in rats (Study 75658)

[00201] This study evaluated and compared the PK profile of LIQ861 to treprostinil (nebulized) to establish a bridge between the two formulations.

[00202] The non-GLP, single administration by inhalation, PK study of treprostinil in rats (Study 75658) has been completed by Liquidia (referred to as the definitive PK bridging study). This study compared the systemic exposure of LIQ861 versus nebulized liquid treprostinil sodium. The observed systemic exposures revealed no meaningful differences between formulations, providing a bridge between the LIQ861 formulation and the marketed Tyvaso formulation and thereby permitting use of Tyvaso nonclinical toxicology studies to support the LIQ861 formulation per the 505(b)(2) pathway.

[00203] In Study 75658, systemic exposure of LIQ861 versus nebulized treprostinil sodium was compared in rats. LIQ861 was delivered over a 4-hour exposure period at total delivered dose levels of 0.273, 0.762, and 1.50 mg/kg body weight. Nebulized treprostinil sodium was delivered at a single dose level (0.785 mg/kg total delivered dose) for the same exposure period (4 hours) as LIQ861. Blood was collected for plasma analysis of treprostinil concentrations at 30 and 60 minutes following the start of administration, immediately post-administration (240 min), and at 5, 15, 30, 75, and 105 minutes following the end of administration.

[00204] Pharmacokinetic parameters from Study 75658. Individual plasma concentrations of treprostinil ranged from 0.345 to 67.4 ng/mL. Maximum plasma concentration was reached 0.5 to 4 hours after the start of the 4-hour exposure period. Maximum concentration (Cmax) and area under the curve (AUC) values were similar between males and females within treatment groups. Dose-related increases in Cmax and AUC values were observed for the three LIQ861 dose groups. Relative bioavailability of LIQ861 compared to nebulized treprostinil based on dose normalized AUC-time curve extrapolated to time infinity (AUCinf) ranged from 1.2 to 2.2.

Summary of Mean Noncompartmental PK Parameters by Treatment and Sex for Study 75658

Type of inhalation	Group	Achieved Mean Dose Level (mg/kg)	Sex	R^2	$t_{1/2}$ (hr)	T_{max} (hr)	C_{max} (ng/mL)	$AUC_{0\text{-}Tlast}$ (hr*ng/mL)	AUC_{INF} (hr*ng/mL)
Treprostinil Sodium (Nebulized)	1	0.785	Female	0.99	0.59	4.00	16.2	62.4	63.5
			Male	0.97	0.77	0.50	16.5	59.3	60.6
Dry Powder (PRINT treprostinil)	2	0.273	Female	0.92	1.77	4.00	5.38	22.2	24.2
			Male	1.00	0.73	1.00	7.18	26.9	27.4
	3	0.762	Female	0.97	0.66	4.00	32.8	107	110
			Male	0.90	0.95	4.00	54.4	144	149
	4	1.498	Female	0.84	0.67	0.50	44.5	174	182
			Male	1.00	0.90	4.00	44.1	143	148

Abbreviations: Cmax, maximal concentration; Tmax, time of maximal concentration; AUClast, area under the concentration-time curve to the last measured timepoint; t½, half-life; AUCinf, area under the concentration-time curve extrapolated to time infinity. PRINT treprostinil = LIQ861 DP-intermediate.

Calculated Relative Bioavailability (Combined Genders) Based on AUCinf (Dose Corrected) from Study 75658

Group	Treatment	Dose Level (mg/kg)	Mean AUCinf (hr*ng/mL)	Frel (PRINT/Treprostinil)
1	Treprostinil	0.785	62.1	NA
2	PRINT-treprostinil Low	0.273	25.8	1.20
3	PRINT-treprostinil Mid	0.762	129	2.15

4	PRINT-treprostinil High	1.498	165	1.39

Abbreviations: AUCinf, area under the concentration-time curve extrapolated to time infinity; Frel, relative bioavailability; NA, not applicable. PRINT treprostinil = LIQ861 DP-intermediate.

[00205] Summary: Study 75658

[00206] The objectives of the study were to determine the pharmacokinetic (PK) profile of Treprostinil in Sprague-Dawley rats when administered as PRINT-Treprostinil (PRINT-Tre) by 4-hour inhalation at 0.15, 0.75, and 1.5 mg/kg, to determine the PK profile of Treprostinil in Sprague-Dawley rats when administered as nebulized Treprostinil sodium in solution (Tre solution) by 4-hour inhalation at 0.75 mg/kg and to compare the PK profiles of Treprostinil when administered as PRINT-Tre and Tre solution.

[00207] The test item was administered once to 6 male and 6 female rats per group by nose-only inhalation for 4 hours as described in the table below:

Group No.	**Group Designation**	**Achieved Mean Total Inhaled Dose Level of Treprostinil (mg/kg/day)**	**Achieved Aerosol Concentration of Treprostinil (µg/L)**	**Achieved Aerosol Concentration of Trehalose (µg/L)**
1	Tre Solution	0.785	5.07	0
2	PRINT-Tre (Low Dose)	0.273	1.76	254.84
3	PRINT-Tre (Mid Dose)	0.762	4.95	719.53
4	PRINT-Tre (High Dose)	1.498	9.73	1394.33

[00208] Assessments of mortality, clinical signs and body weights were performed. Pharmacokinetic samples were collected and the analysis of these samples was performed.

[00209] No mortality occurred and no clinical signs were observed.

[00210] The overall achieved aerosol concentrations for all groups were within 10% of the targeted concentrations gravimetrically and for both treprostinil and trehalose, except for Group

2 which were significantly above the targeted concentrations (76 to 95%). Corresponding average achieved dose levels for all groups were within 5% of the targeted dose levels, except for Group 2 which was 82% above the targeted dose level. However, the dose levels were considered acceptable for the study as a clear dose differentiation between groups for each sex was achieved.

[00211] The particle size MMADs from Groups 2 to 4 were between 1.7 and 2.0 μm gravimetrically (GSD 1.90 to 2.67); for both treprostinil and trehalose, chemical particle size distribution ranged from 1.6 to 1.8 μm with the corresponding GSDs between 1.89 and 2.24. The particle size MMAD for Group 1 was 0.5 μm with a corresponding GSD of 2.60. The particle size distribution was considered respirable.

[00212] With administration of PRINT-Tre at an achieved dose level of 0.273 mg/kg, 0.762 mg/kg or 1.498 mg/kg, plasma exposure to treprostinil was generally similar in both sexes; however, exposure was slightly lower in females than males at the mid-dose level and slightly higher in females than males at the high-dose level.

[00213] Based on $AUC_{0\text{-}Tlast}$, AUC_{INF} and C_{max}, values for both sexes, plasma exposure increased more than proportionally between the low- and mid-dose levels. But between the mid- and high-dose levels, plasma exposure increased less than proportionally for females and there was no increase in the exposure for males. The maximum mean treprostinil plasma concentration (T_{max}) was at the end of inhalation for both sexes, except for low-dose males and high-dose females, where mean T_{max} was at 1 and 0.5 hours after inhalation began, respectively.

[00214] At the low-dose level, mean treprostinil plasma concentration was similar after 0.5, 1, or 4 hours of inhalation exposure to PRINT-Tre, suggesting that steady state was achieved within the first 30 minutes of exposure. The same was true for females at the high-dose level; however, for males at the high-dose level and for both sexes at the mid-dose level, mean treprostinil plasma concentration was greater at the end of inhalation than after one hour of inhalation. When inhalation ended, treprostinil plasma concentrations declined gradually. Given the degree of individual variation, the estimated mean $T_{1/2}$ values were similar at all dose levels and ranged from 0.7 to 1.8 hours in males and 0.7 to 1.0 hours in females.

[00215] For Tre solution, with an administration at 0.785 mg/kg, plasma exposure to treprostinil was generally similar in both sexes. The maximum mean treprostinil plasma concentration (T_{max}) was at the end of inhalation. Mean treprostinil plasma concentration was similar after 0.5, 1, or 4 hours of inhalation exposure to Tre solution, suggesting that steady state was achieved within the first 30 minutes of exposure. When inhalation ended, treprostinil plasma concentrations declined gradually, with estimated mean $T_{1/2}$ values of 0.6 hours in males and 0.8 hours in females.

[00216] Administration of PRINT-Tre and Tre solution at nearly equivalent dose levels (0.76 and 0.79 mg/kg, respectively) resulted in plasma exposures to treprostinil that were greater with PRINT-Tre than with Tre solution. Specifically, mean $AUC_{0\text{-}Tlast}$ was approximately twice as high (126 versus 61 h*ng/mL, respectively) and mean C_{max} was three times as high (44 versus 16 ng/mL, respectively). As would be expected, once treprostinil entered the systemic circulation, it was cleared from plasma at a similar rate, with mean $T_{1/2}$ values of 0.7 to 1.0 hours for PRINT-Tre and 0.6 to 0.8 hours for Tre solution.

[00217] In conclusion, single inhalation administration for 4 hours of PRINT Treprostinil at a high average dose of 1.498 mg/kg/day to Sprague-Dawley rats was well tolerated as there were no test item related findings. At an equivalent dose level, plasma exposures to treprostinil was greater with PRINT-Tre than with Tre solution; specifically, mean $AUC_{0\text{-}Tlast}$ was approximately twice as great and mean C_{max} was three times as great. As would be expected, once treprostinil entered the systemic circulation, it was cleared from plasma at a similar rate regardless of how it was administered.

[00218] The objectives of the study were to:

1. Determine the pharmacokinetic (PK) profile of Treprostinil in Sprague-Dawley rats when administered as PRINT-Treprostinil (PRINT-Tre) by 4-hour inhalation at 0.15, 0.75, and 1.5 mg/kg.

2. Determine the PK profile of Treprostinil in Sprague-Dawley rats when administered as nebulized Treprostinil sodium in solution (Tre solution) by 4-hour inhalation at 0.75 mg/kg.

3. Compare the PK profiles of Treprostinil when administered as PRINT-Tre and Tre solution.

[00219] Experimental Design

[00220] The test items were administered to groups of rats by a 4-hour inhalation administration as described in the table below:

Group No.	Group Designation	Targeted Total Inhaled Dose Level of Treprostinil (mg/kg/day)	Targeted Aerosol Concentration of Treprostinil (µg/L) [a]	Targeted Aerosol Concentration of Trehalose (µg/L) [a]	No. of Animals	
					Males	Females
1	Tre Solution	0.75	5	0	6	6
2	PRINT-Tre	0.15	1	130.7	6	6
3	PRINT-Tre	0.75	5	653.5	6	6
4	PRINT-Tre	1.5	10	1306.9	6	6

[a] = Target aerosol concentrations were calculated based on an estimated body weight of 0.250 kg

[00221] During and after the inhalation period, a series of 8 blood samples for pharmacokinetic evaluation were taken.

[00222] Justification for Selection of Route of Administration, Species and Dose Levels

[00223] The route of administration was chosen because it is the intended human therapeutic route.

[00224] The rat was selected because it is a rodent species recommended by various regulatory authorities. Background data are available. Also, rats were used as the test system for previous toxicity studies with Treprostinil sodium solution that supported development and approval of that product. Using rats in the current study allowed comparison with the previous studies.

[00225] The high-dose level for PRINT-Tre was the feasible dose attainable based on technical aerosol trials with the test item (Study No. 41610).

[00226] The low- and mid-dose levels for PRINT-Tre were selected on the basis of a previous pilot PK study in rats (Study No. 75670).

[00227] The dose level for Tre solution was selected to match the mid-dose level of PRINT-Tre to allow direct comparison.

- Characterization of Test Items

	Content:	92.79% of Trehalose, 4% of Leucine, 2% of Tween80, 0.26% of NA Citrate, 0.24% of NaCl : 0.71% of Treprostinil
	Storage Conditions:	Cool (2 to 8°C), protected from moisture (e.g., dessicant)
	Handling Precautions:	Standard laboratory precautions. Handle under dry conditions (Relative Humidity ≤ 23%)
	Supplier:	Liquidia Technologies Inc.
Test item 2*:	Identity:	Treprostinil Sodium
	Description:	White or pale yellowish powder
	Batch No.:	TN115E010
	Expiry Date:	May 28, 2017
	Purity:	101.49%
	Storage Conditions:	Cool (2 to 8°C)
	Handling Precautions:	Standard laboratory precautions
	Supplier:	Yonsung Fine Chemicals Co., LTD

[00228] Preparation of Test Item

[00229] PRINT-Tre was used as provided by the Sponsor. A glove box under nitrogen was used for handling, aliquoting or packing of the canisters. Relative humidity (RH) inside the glove box was monitored and recorded using a hygrometer and was kept below 23% RH.

[00230] For Group 1, the treprostinil sodium was dissolved in purified water to achieve the desired formulation concentration. A representative sample (0.5mL in duplicate) was collected to verify the formulation concentration of Treprostinil in the formulation.

[00231] Treatment

[00232] Acclimatization to Exposure System

[00233] Before the animals were exposed to the aerosol of the test item, rats were accustomed to the restraint procedure over a period of 3 days. The animals were gradually accustomed to restraint in the dosing tubes used during the exposures up to the duration that was used for aerosol administrations.

[00234] Animal Exposure

Exposure system used:	Flow-past rodent inhalation exposure system
Exposure method:	Inhalation by nose-only exposure
Test Item type:	Solution (Group 1), Dry Powder (Groups 2 to 4)
Generation method:	Nebulization (Group 1) and Piston feed/rotating brush generator (Group 2 to 4)
Duration of exposure:	240 minutes

[00235] The target aerosol concentrations and dose levels were as follows:

Group No.	Group Designation	Targeted Dose Level of Treprostinil (mg/kg/day)	Targeted Aerosol Concentration of Treprostinil (µg/L) [a]	Targeted Aerosol Concentration of Trehalose (µg/L)
1	Tre solution	0.75	5	0
2	PRINT-Tre (Low Dose)	0.15	1	130.7

3	PRINT-Tre (Mid Dose)	0.75	5	653.5
4	PRINT-Tre (High Dose)	1.5	10	1306.9

a = Target aerosol concentrations were calculated based on an estimated body weight of 0.250 kg.

[00236] Estimation of Achieved Dose Levels

The target dose levels were estimated using the following formula:

$$D_L = \frac{E_c \times RMV \times T}{BW}$$

D_L = Achieved dose levels (mg / kg / day)

E_c = Actual concentration delivered to the animals (mg / L air)

RMV = Respiratory Minute Volume (L / min) according to the method of Bide, Armour and Yee 2000 J. App. Toxicol., Vol. 20: *RMV* (L / min) = 0.499 x *BW* (kg)$^{0.809}$

T = Time, duration of daily exposure (min.)

BW = Mean body weight (kg) during exposure period.

This estimation of total inhaled dose assumed 100% deposition within the respiratory tract.

[00237] Inhalation Exposure System

[00238] The powder aerosol for Groups 2 to 4 was produced using a piston feed / rotating brush generator while the liquid aerosol for Group 1 was produced by metering the flow of the formulation to a clinical nebulizer (Sidestream). The aerosol produced was diluted as necessary to achieve the target aerosol concentration and discharged through a 40-mm diameter tube into a flow-past inhalation exposure system. The airflow rate through the exposure system was monitored and recorded manually during each aerosol generation period. Airflow to the exposure system was controlled by the absolute volume of air supplying the generation apparatus using variable area flowmeters. Control of the aerosol exhaust flow from the animal exposure system was achieved using an exhaust valve, and the overall balance of airflows in the exposure system was monitored using pressure gauges. The system provided a minimum of 1.0 L/min to

each animal exposure port and was balanced to ensure a slight positive pressure at the site of the animal exposure. This ensured that there was no dilution of the generated aerosol. An equal delivery of aerosol to each exposure position was achieved by employing a distribution network that was identical for each individual exposure position attached to the system.

[00239] Inhalation System Monitoring

[00240] Determinations of aerosol concentration, particle size distribution, oxygen concentration, relative humidity and temperature were measured on samples collected from a representative port of the exposure chamber. The sample flow rates were precisely controlled using variable area flow meters that were calibrated before use using a primary airflow calibrator. The absolute volume of each aerosol concentration sample was measured with a wet type gas meter.

[00241] Oxygen Concentration

[00242] The oxygen concentration of the generated atmosphere was measured once during each aerosol exposure. Oxygen concentrations of the exposure atmospheres were maintained between 19-23%.

[00243] Relative Humidity/Temperature

[00244] The temperature and relative humidity of the generated atmosphere were measured once during each aerosol exposure. Temperatures of the exposure atmospheres were maintained between 19-24°C.

[00245] Determination of Aerosol Concentration

[00246] At least one aerosol concentration filter sample was collected for all groups on each aerosol generation. The filter samples from Groups 2 to 4 were weighed in order to measure the gravimetric concentration of the test item in the generated aerosol. The filter samples were transferred to the analytical chemistry laboratory for chemical determination of Treprostinil and Trehalose concentrations. The filter samples for Group 1 were not weighed gravimetrically and were only transferred to the analytical laboratory for determination of Treprostinil

concentrations. The analysis in the analytical laboratory was performed using an analytical method (Study No. 41609).

[00247] Determination of the Particle Size Distribution and Mass Median Aerodynamic Diameter (MMAD)

[00248] The distribution of particle size in the generated aerosols was measured once for Groups 1 to 4 by collecting samples into a 7-Stage Mercer Cascade Impactor. All sample substrates obtained from Groups 2 to 4 were weighed gravimetrically and then transferred to the analytical chemistry laboratory for chemical determination of particle size of aerosolized Treprostinil and Trehalose. All sample substrates obtained from Group 1 were only transferred to the analytical laboratory for chemical determination of particle size of aerosolized Treprostinil. The analysis in the analytical laboratory was performed using an analytical method (Study No. 41609).

[00249] The MMAD and the Geometric Standard Deviation (GSD) were calculated based on the results obtained from the impactor using a log-probit transformation.

[00250] Reporting of Analytical Results

[00251] The analytical report containing the results from the filter and particle size distribution sample analyses were prepared. Any samples not employed in the primary analysis or any remaining sample from the primary analysis were retained until it was determined by the analyst and Study Director that it was not be required for confirmatory analysis. These samples were discarded and their disposition recorded in the raw data.

[00252] IN-LIFE OBSERVATIONS

[00253] Mortality

[00254] Mortality checks were performed at least once a day during all phases of the study.

[00255] Clinical Observations

[00256] Cage-side clinical signs (ill health, behavioral changes etc.) were recorded at least once daily during all phases of the study, except on detailed clinical examination days, where the cage-side clinical signs were replaced by a DCE.

[00257] A detailed clinical examination of each rat was performed on arrival as part of the health status, as well as on Day 1, prior to dosing.

[00258] Body Weights

[00259] Body weights were recorded for all animals once at arrival as per health status, once prior to group assignment and on Day 1 (prior to dosing).

[00260] Pharmacokinetics

[00261] A series of 8 blood samples (approximately 0.3 mL each) was collected 30 minutes and 1 hour after exposure began, immediately after exposure ended (IPE), and again at 5, 15, 30, 75 and 105 minutes post-dosing as per the table below. Thus a total blood volume of 1.2 mL was taken from each rat during the course of the study. For this purpose, each rat (unanesthetized) was bled by jugular venipuncture and the samples were collected into tubes containing the anticoagulant, K_2EDTA. Tubes were placed on wet ice pending processing.

Group Number	Number of animals/sex	Toxicokinetic time point							
		30 min post start	1 hour post start	IPE	5 min post end	15 min post end	30 min post end	75 min post end	105 min post end
1	3 +	√		√		√		√	
	3 #		√		√		√		√
2	3 +	√		√		√		√	
	3 #		√		√		√		√
3	3 +	√		√		√		√	
	3 #		√		√		√		√
4	3 +	√		√		√		√	
	3 #		√		√		√		√

\+ animals with the lowest identification numbers

\# animals with the highest identification numbers

[00262] Following collection, the samples were centrifuged (2500 rpm for 10 minutes at approximately 4°C) and the resulting plasma was recovered and stored frozen (≤-60°C) in labeled tubes.

[00263] Deviations to the pharmacokinetic time points were noted in the raw data and were made available with the samples. The location of blood withdrawal was noted in the raw data.

[00264] The plasma analysis was performed and the bioanalytical data was prepared for inclusion in the final report.

[00265] The pharmacokinetic parameters were calculated and the non-compartmental analysis of PRINT-Tre and Tre solution treprostinil concentrations in plasma was performed by using the Phoenix WinNonlin 6.3 software.

[00266] The following configuration was used for the analysis:

Sampling Method:	Sparse
AUC Calculation Method:	Linear Trapezoidal with Linear Interpolation
Lambda Z (λ_z) Method:	Best fit for λz, Log regression
Weighting (λ_z calculation):	Uniform

[00267] Pharmacokinetic parameters (including abbreviation and description for each parameter) were described in the following table:

Parameters	Abbreviation	Unit*
Area under the plasma drug concentration-time curve from the time of dosing to the last quantifiable concentration	$AUC_{0\text{-}Tlast}$	µg*hr/mL
Area under the plasma drug concentration-time curve from the time of dosing extrapolated to infinity	AUC_{INF}	µg*hr/mL
Terminal elimination half-life	$T_{1/2}$	hr
The maximum plasma concentration	C_{max}	µg/mL
Time to maximum plasma concentration	T_{max}	hr

[00268] Data Evaluation and Statistics

[00269] Numeric and non-numeric data obtained during the study were reported only as individual values.

[00270] RESULTS

[00271] Formulation Analysis

[00272] Formulation concentration for Group 1 was as follows:

Group No.	**Average Targeted Concentration (mg/mL)**	**Average Measured Concentration (mg/mL)**	**% of Targeted Concentration**
1	0.50	0.492	98.4

[00273] The formulation concentration for Group 1 was within 2% of the targeted concentration therefore the formulation concentration was considered acceptable for the study.

[00274] Aerosol Concentrations

[00275] Achieved gravimetric test atmosphere concentrations were as follows:

Group No.	Targeted Aerosol Concentration (mg\L)	Achieved Mean Aerosol Concentration (mg\L)	Coefficient of Variation (%)	% of Target
2	0.156	0.283	55.5	181.4
3	0.781	0.814	16.3	104.2
4	1.563	1.548	21.9	99.0

[00276] Achieved chemical test atmosphere concentrations for treprostinil were as follows:

Group No.	Targeted Aerosol Concentration (µg\L)	Achieved Mean Aerosol Concentration (µg\L)	Coefficient of Variation (%)	% of Target
1	5	5.07	3.6	101.4
2	1	1.76	55.2	176.3
3	5	4.95	19.8	99.1
4	10	9.73	22.3	97.3

[00277] Achieved chemical test atmosphere concentrations for trehalose were as follows:

Group No.	Targeted Aerosol Concentration (µg\L)	Achieved Mean Aerosol Concentration (µg\L)	Coefficient of Variation (%)	% of Target
2	130.7	254.84	54.4	195.0
3	653.5	719.53	20.5	110.1
4	1306.9	1394.33	23.2	106.7

[00278] The overall achieved aerosol concentrations for all groups were within 10% of the targeted concentrations gravimetrically and for both treprostinil and trehalose, except for Group 2 which were significantly above the targeted concentrations (76% and 95% for treprostinil and trehalose, respectively). The generated atmospheres were considered stable over the treatment period except for Group 2 (CV % ~54%). However, the overall aerosol concentrations were still considered acceptable for the study as there was a significant difference in aerosol concentration between groups.

[00279] Achieved Dose Levels

[00280] Overall achieved doses for treprostinil are presented below:

Group No.	Targeted Dose Level (mg/kg/day)	Duration of Exposure (min)	Sex	Body Weight (kg)	Estimated Achieved Doses (mg/kg/day)	% from Targeted Dose Level
1	0.75	240	Male	0.308	0.760	101.4
			Female	0.212	0.817	108.9
			Combined	*0.260*	*0.785*	*104.7*
2	0.15	240	Male	0.301	0.265	176.7
			Female	0.211	0.284	189.1
			Combined	*0.256*	*0.273*	*182.3*
3	0.75	240	Male	0.321	0.737	98.2
			Female	0.215	0.795	106.0
			Combined	*0.268*	*0.762*	*101.6*
4	1.5	240	Male	0.317	1.451	96.7
			Female	0.219	1.557	103.8
			Combined	*0.268*	*1.498*	*99.9*

[00281] Average achieved dose levels for all groups were within 5% of the targeted dose levels, except for Group 2 which was 82% above the targeted dose level. However, the dose levels were considered acceptable for the study as a clear dose differentiation between groups for each sex was achieved.

[00282] Particle Size Distribution

[00283] The average gravimetric particle size distribution measurement data were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
2	87.9	80.0	61.0	31.2	21.8	16.2	9.1	0.0	1.7	2.67	80
3	90.7	85.0	66.9	24.9	14.8	9.0	3.0	0.0	1.8	2.12	85
4	91.7	84.0	61.6	23.5	8.1	4.3	0.8	0.0	2.0	1.90	86

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00284] The average chemical determination of particle size distribution for treprostinil were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
1	95.9	95.7	95.4	95.1	85.1	50.1	22.0	0.0	0.5	2.60	98
2	94.0	86.3	64.4	29.3	19.8	14.1	6.1	0.0	1.6	2.24	87
3	95.3	90.3	71.5	25.9	15.2	9.3	2.7	0.0	1.6	1.97	90
4	94.1	88.4	64.3	23.9	8.2	4.4	1.5	0.0	1.8	1.89	88

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00285] The average chemical determination of particle size distribution for trehalose were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
2	94.3	86.7	63.8	26.4	16.8	14.1	6.1	0.0	1.6	2.22	87
3	96.0	92.0	72.3	22.2	12.0	8.0	4.0	0.0	1.6	1.97	90
4	95.7	91.4	68.0	27.9	12.8	8.6	4.3	0.0	1.6	2.00	90

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00286] The particle size distribution was considered respirable for this study as the MMADs were below 4 µm and the GSD were within 1.5 and 3.

[00287] Exposure Chamber Conditions

[00288] Exposure chamber conditions from the reported aerosol concentration exposures are summarized below.

Group No.	Humidity (%RH)	Temperature (°C)	Oxygen Concentration (%)
1	58.5	21.0	20.9
2	35.1	21.6	20.9
3	39.0	21.5	20.9
4	39.4	21.2	20.9

[00289] Exposure atmosphere oxygen concentrations, temperature and relative humidity ranges were considered acceptable on all occasions.

[00290] Mortality

[00291] There were no mortalities during the study.

[00292] Clinical Signs

[00293] There were no clinical signs observed during the study.

[00294] **Body Weight**

[00295] Body weights were performed for dose level calculation purposes.

[00296] **Pharmacokinetics**

[00297] With administration of PRINT-Tre at an achieved dose level of 0.273 mg/kg, 0.762 mg/kg or 1.498 mg/kg, plasma exposure to treprostinil was generally similar in both sexes; however, exposure was slightly lower in females than males at the mid-dose level and slightly higher in females than males at the high-dose level.

[00298] Based on $AUC_{0\text{-}Tlast}$, AUC_{INF} and $C_{max,}$ values for both sexes, plasma exposure increased more than proportionally between the low- and mid-dose levels. But between the mid- and high-dose levels, plasma exposure increased less than proportionally for females and there was no increase in the exposure for males. The maximum mean treprostinil plasma concentration (T_{max}) was at the end of inhalation for both sexes, except for low-dose males and high-dose females, where mean T_{max} was at 1 and 0.5 hours after inhalation began, respectively.

[00299] At the low-dose level, mean treprostinil plasma concentration was similar after 0.5, 1, or 4 hours of inhalation exposure to PRINT-Tre, suggesting that steady state was achieved within the first 30 minutes of exposure. The same was true for females at the high-dose level; however, for males at the high-dose level and for both sexes at the mid-dose level, mean treprostinil plasma concentration was greater at the end of inhalation than after one hour of inhalation. These data are summarized below.

	Males			**Females**		
Dose (mg/kg) =	**0.273**	**0.762**	**1.498**	**0.273**	**0.762**	**1.498**
0.5 hours of inhalation	6.1	22	26	4.2	18	44
1 hour of inhalation	7.2	21	27	5.1	19	35
4 hours of inhalation	5.4	54*	44^	5.4	33**	44

*Individual values were 33, 63, and 67 ng/mL

^Individual values were 34, 48, and 50 ng/mL

**Individual values were 22, 27, and 49 ng/mL

[00300] When inhalation ended, treprostinil plasma concentrations declined gradually. Given the degree of individual variation, the estimated mean $T_{1/2}$ values were similar at all dose levels and ranged from 0.7 to 1.8 hours in males and 0.7 to 1.0 hours in females.

[00301] For Tre solution, with an administration at 0.785 mg/kg, plasma exposure to treprostinil was generally similar in both sexes.

[00302] The maximum mean treprostinil plasma concentration (T_{max}) was at the end of inhalation. Mean treprostinil plasma concentration was similar after 0.5, 1, or 4 hours of inhalation exposure to Tre solution, suggesting that steady state was achieved within the first 30 minutes of exposure. These data are summarized below.

	Males	**Females**
0.5 hours of inhalation	17	12
1 hour of inhalation	11	14
4 hours of inhalation	16	16

[00303] When inhalation ended, treprostinil plasma concentrations declined gradually, with estimated mean $T_{1/2}$ values of 0.6 hours in males and 0.8 hours in females.

[00304] Administration of PRINT-Tre and Tre solution at nearly equivalent dose levels (0.76 and 0.79 mg/kg, respectively) resulted in plasma exposures to treprostinil that were greater with PRINT-Tre than with Tre solution. Specifically, mean $AUC_{0\text{-}Tlast}$ was approximately twice as high (126 versus 61 h*ng/mL, respectively) and mean C_{max} was three times as high (44 versus 16 ng/mL, respectively). As would be expected, once treprostinil entered the systemic circulation, it was cleared from plasma at a similar rate, with mean $T_{1/2}$ values of 0.7 to 1.0 hours for PRINT-Tre and 0.6 to 0.8 hours for Tre solution.

[00305] CONCLUSION

[00306] Single inhalation administration for 4 hours of PRINT Treprostinil at a high average dose of 1.498 mg/kg/day to Sprague-Dawley rats was well tolerated as there were no test item related findings. At an equivalent dose level, plasma exposures to treprostinil was greater with PRINT-Tre than with Tre solution; specifically, mean $AUC_{0\text{-}Tlast}$ was approximately twice as great and mean C_{max} was three times as great. As would be expected, once treprostinil entered the systemic circulation, it was cleared from plasma at a similar rate regardless of how it was administered.

[00307] Non-GLP, 7-day, repeat-dose, range-finding (DRF), inhalation study with LIQ861 in rats (Study 75654)

[00308] Results from the completed comparative PK study will be used to select dose levels to be tested in this DRF study, which will evaluate local toxicity in the respiratory tract as well as systemic treprostinil toxicity. Results will be used to select appropriate dose levels for a 2-week GLP repeat-dose toxicology study in rats.

[00309] Summary: Study 75654

[00310] The objectives of the study were to evaluate the toxicity of the test item, PRINT Treprostinil, and the excipients that make up the control item, PRINT Placebo, when administered to Sprague-Dawley rats by nose-only inhalation for 4 hours a day for 7 days. Results were used to help select dose levels for a subsequent 14-day GLP inhalation toxicology study.

[00311] Groups of 6 rats (3/sex) were exposed by 4-hour inhalation daily for 7 days to air, PRINT Placebo, or PRINT Treprostinil at treprostinil dose levels of approximately 170, 680, or 1370 µg/kg, as described in the table below:

Group No.	Group Designation	Mean Dose Levels and Concentrations[a]				
		Treprostinil		Trehalose		Leucine
		Dose Level (µg/kg/day)	Aerosol Conc. (µg/L)	Dose Level (mg/kg/day)	Aerosol Conc. (µg/L)	Dose Level (mg/kg/day)[b]
1	Air Control	0	0	0	0	0
2	Placebo Control[b]	0	0	281.2	1832.13	12.0
3	PRINT-Tre (Low Dose)	170	1.10	33.1	216.30	1.3
4	PRINT-Tre (Mid Dose)	680	4.44	133.5	869.99	5.1
5	PRINT-Tre (High Dose)	1370	8.94	266.6	1735.84	10.3

a = Based on the mean body weight of each group during the dosing period.

b = Calculated with a content of 4% of Leucine in PRINT Treprostinil and PRINT Placebo and using Trehalose percentage of 93.5% in PRINT Placebo for Group 2 and Treprostinil percentage of 0.53% in PRINT Treprostinil for Groups 3 to 5.

[00312] The particle size MMADs from Groups 2 to 5 were between 1.3 and 2.0 µm gravimetrically (GSD 1.96 to 2.46); for both treprostinil and trehalose, chemical particle size distribution ranged from 1.3 to 2.1 µm with the corresponding GSDs between 1.87 and 1.95. No mortality occurred. No clinical signs were observed while coagulation, clinical chemistry and urinalysis parameters were unaffected and no test item-related findings were seen macroscopically.

[00313] Rats tolerated daily administration of PRINT Placebo or PRINT-Tre at up to 1.37 mg/kg/day by 4-hour inhalation for 7 days.

[00314] Introduction

[00315] The objectives of the study were to:

1. Evaluate the toxicity of the test item, PRINT Treprostinil, when administered to Sprague-Dawley rats by nose-only inhalation for 4 hours a day for 7 days.

2. Evaluate the toxicity of the excipients that make up the control item, PRINT Placebo, when administered to Sprague-Dawley rats by nose-only inhalation for 4 hours a day for 7 days.

3. Determine the dose levels of PRINT Treprostinil for the following 14-day GLP inhalation toxicology study from the results of this dose range-finding study.

[00316] Experimental Design

[00317] Synopsis

[00318] The test and control items were administered to groups of 6 rats (3/sex) by 4-hour inhalation daily for 7 days, as described in the table below. The first day of dosing was designated as Day 1.

Group No.	Test Material	Targeted Aerosol Concentration (µg/L)		Targeted Dose Level (mg/kg/day)		Leucine Dose Level (mg/kg/day)[c]
		Treprostinil	Trehalose	Treprostinil[a]	Trehalose[b]	
1	Air Control	0	0	0	0	0
2	PRINT Placebo[b]	0	1684.6	0	262.9	11.2
3	PRINT-Tre (Low Dose)	1	175.2	0.15	26.3	1.1
4	PRINT-Tre (Mid Dose)	5	876.2	0.75	131.4	5.7
5	PRINT-Tre (High Dose)	10	1752.5	1.5	262.9	11.3

a = Target aerosol concentrations were calculated based on an estimated body weight of 0.250 kg

b = The target dose level for the placebo control was the same dose level as the high dose group (Group 5)

c = Calculated with a content of 4% of Leucine in PRINT Treprostinil and PRINT Placebo (using Trehalose percentage of 93.5% in PRINT Placebo for Group 2 and Treprostinil percentage of 0.53% in PRINT Treprostinil for Groups 3 to 5)

[00319] The high-dose level for PRINT-Tre was the feasible dose attainable based on technical aerosol trials with the test item. (Study No. 41610).

[00320] The low- and mid-dose levels for PRINT-Tre were selected on the basis of a previous PK study in rats (Study No. 75658).

[00321] Test and Control Item Information

[00322] Test Item Action

[00323] Treprostinil, the active ingredient in PRINT-Tre, is a prostacyclin compound approved for treatment of pulmonary arterial hypertension.

[00324] Characterization of Test Item

Content:	92.97% of Trehalose, 4% of Leucine, 2% of Tween80, 0.27% of Sodium Citrate Dihydrate, 0.23% of Sodium Chloride : 0.53% of Treprostinil sodium
Storage Conditions:	Cool (2 to 8°C), protected from moisture (e.g., desiccant)
Handling Precautions:	Standard laboratory precautions. Handled under dry conditions (relative humidity ≤ 23%)
Supplier:	Liquidia Technologies Inc.

[00325] Characterization of Placebo Control Item

Content:	-LKI-1R-983-3: 93.53% of Trehalose, 4% of Leucine, 2% of Tween80, 0.24% of Sodium Citrate Dihydrate, 0.23% of Sodium Chloride -LKI-1R-983-27: 93.5% of Trehalose, 4% of Leucine, 2% of Tween80, 0.27% of Sodium Citrate Dihydrate, 0.23% of Sodium Chloride
Storage Conditions:	Cool (2 to 8°C) , protected from moisture (e.g., desiccant)
Handling Precautions:	Standard laboratory precautions. Handled under dry conditions (relative humidity ≤ 23%)
Supplier:	Liquidia Technologies Inc.

[00326] Characterization of Air Control

Description:	Medical Grade Air (NQ 5710-500/2000)
Supplied By:	Kaeser SM-11 Air Compressor

[00327] Preparation of Test and Control Items

[00328] PRINT-Tre and PRINT Placebo were used as provided by the Sponsor. A glove box under nitrogen was used for handling, aliquoting or packing of the canisters. Relative humidity (RH) inside the glove box was monitored and recorded using a hygrometer and was kept below 23% RH.

[00329] Treatment

[00330] Acclimatization to Exposure System

[00331] Before the rats were presented to exposure atmosphere, rats were accustomed to the restraint procedure over a period of 3 days. The animals were gradually accustomed to restraint in the dosing tubes used during the exposures up to the duration that was used for aerosol administrations.

[00332] Animal Exposure

Exposure system used:	Flow-past rodent inhalation exposure system
Exposure method:	Inhalation by nose-only exposure
Test and Control Item type:	Air (Group 1), Dry Powder (Groups 2 to 5)
Generation method:	Piston feed/rotating brush generator (Groups 2 to 5)
Duration of exposure:	240 minutes

[00333] The target aerosol concentrations and dose levels were as follows:

Group No.	Group Designation	Targeted Total Inhaled Dose Level of Treprostinil (mg/kg/day)[a]	Targeted Aerosol Concentration of Treprostinil (µg/L)	Targeted Total Inhaled Dose Level of Trehalose (mg/kg/day)[a]	Targeted Aerosol Concentration of Trehalose (µg/L)	Estimated Total Inhaled Dose Level of Leucine (mg/kg/day)[c]
1	Air Control	0	0	0	0	0
2	Placebo Control[b]	0	0	262.9	1684.6	11.2
3	PRINT-Tre (Low Dose)	0.15	1	26.3	175.2	1.1
4	PRINT-Tre (Mid Dose)	0.75	5	131.4	876.2	5.7
5	PRINT-Tre (High Dose)	1.5	10	262.9	1752.5	11.3

a = Target aerosol concentrations were calculated based on an estimated body weight of 0.250 kg

b = The target dose level for the placebo control was the dose level as the high dose group (Group 5)

c = Calculated with a content of 4% of Leucine in PRINT Treprostinil and PRINT Placebo (using Trehalose percentage of 93.5% in PRINT Placebo for Group 2 and Treprostinil percentage of 0.53% in PRINT Treprostinil for Groups 3 to 5)

[00334] Estimation of Achieved Dose Levels

The target dose levels were estimated using the following formula:

$$D_L = \frac{E_c \times RMV \times T}{BW}$$

D_L = Achieved dose levels (mg / kg / day)

E_c = Actual concentration delivered to the animals (mg / L air)

RMV = Respiratory Minute Volume (L / min) according to the method of Bide, Armour and Yee. J. App. Toxicol., Vol. 20, 2000: *RMV* (L / min) = 0.499 x *BW* (kg)$^{0.809}$

T = Time, duration of daily exposure (min.)

BW = Mean body weight (kg) during exposure period.

This estimation of total inhaled dose assumed 100% deposition within the respiratory tract.

[00335] Inhalation Exposure System

[00336] The powder aerosol for Groups 2 to 5 was produced using a piston feed / rotating brush generator. The aerosol produced was diluted as necessary to achieve the target aerosol concentration and discharged through a 40-mm diameter tube into a flow-past inhalation exposure system. The airflow rate through the exposure system was monitored and recorded manually during each aerosol generation period. Airflow to the exposure system was controlled by the absolute volume of air supplying the generation apparatus using variable area flowmeters. Control of the aerosol exhaust flow from the animal exposure system was achieved using an exhaust valve, and the overall balance of airflows in the exposure system was monitored using pressure gauges. The system provided a minimum of 1.0 L/min to each animal exposure port and was balanced to ensure a slight positive pressure at the site of the animal exposure. This ensured that there was no dilution of the generated aerosol. An equal delivery of aerosol to each exposure position was achieved by employing a distribution network that was identical for each individual exposure position attached to the system.

[00337] Inhalation System Monitoring

[00338] Determinations of aerosol concentration, particle size distribution, oxygen concentration, relative humidity and temperature were measured on samples collected from a representative port of the exposure chamber, with a collection sample flow-rate of 1 L/min. The sample flow rates were precisely controlled using variable area flow meters that were calibrated before use using a primary airflow calibrator. The absolute volume of each aerosol concentration sample was measured with a wet type gas meter.

[00339] Oxygen Concentration

[00340] The oxygen concentration of the generated atmosphere was measured once during each aerosol exposure. Oxygen concentrations of the exposure atmospheres were maintained between 19-23%.

[00341] Relative Humidity/Temperature

[00342] The temperature and relative humidity of the generated atmosphere were measured once during each aerosol exposure. Temperatures of the exposure atmospheres were maintained between 19-24°C.

[00343] Determination of Aerosol Concentration

[00344] At least one aerosol concentration filter sample was collected for all groups on each aerosol generation. The filter samples from Groups 3 to 5 were weighed in order to measure the gravimetric concentration of the test item in the generated aerosol. The filter samples were transferred to the analytical chemistry laboratory for chemical determination of Treprostinil and Trehalose concentrations. The filter samples from Group 2 were weighed in order to measure the gravimetric concentration of the control item in the generated aerosol. The filter samples were transferred to the analytical chemistry laboratory for chemical determination of Trehalose concentration and to confirm the absence of Treprostinil. The filter samples for Group 1 were not weighed gravimetrically and were only transferred to the analytical laboratory to confirm the absence of Treprostinil and Trehalose. The analysis in the analytical laboratory was performed using an analytical method (Study No. 41609 and 41635).

[00345] Determination of Aerosol Homogeneity

[00346] At least once during the study, atmosphere homogeneity in the exposure system was tested by collecting multiple aerosol samples from the top, middle and bottom tiers of the exposure system of Groups 2 to 5.

[00347] Determination of the Particle Size Distribution and Mass Median Aerodynamic Diameter (MMAD)

[00348] The distribution of particle size in the generated aerosols was measured at least once for Groups 2 to 5 by collecting samples into a 7-Stage Mercer Cascade Impactor. All sample substrates obtained from Groups 3 to 5 were weighed gravimetrically and then transferred to the analytical chemistry laboratory for chemical determination of particle size of aerosolized Treprostinil and Trehalose. All sample substrates obtained from Group 2 were weighed gravimetrically and then transferred to the analytical laboratory for determination of particle size of aerosolized Trehalose. The analysis in the analytical laboratory was performed using an analytical method (Study No. 41609 and 41635).

[00349] The MMAD and the Geometric Standard Deviation (GSD) were calculated based on the results obtained from the impactor using a log-probit transformation.

[00350] Reporting of Analytical Results

[00351] The analytical report containing the results from the filter and particle size distribution sample analyses were prepared. Any samples not employed in the primary analysis or any remaining sample from the primary analysis were retained until it was determined by the analyst and Study Director that it was not required for confirmatory analysis. These samples were then discarded and their disposition was recorded in the raw data.

[00352] Standard Operating Procedures

[00353] All procedures, were performed in accordance with the Standard Operating Procedures and these were kept on file. Deviations to the Standard Operating Procedures were documented in the raw data.

[00354] RESULTS

[00355] Inhalation System Monitoring

[00356] Oxygen Concentration, Temperature, and Relative Humidity

[00357] Exposure chamber conditions from the reported aerosol concentration exposures are summarized below.

Group Number	Humidity (%RH)	Temperature (°C)	Oxygen Concentration (%)
	Average	Average	Average
1	28.6	20.4	20.9
2	39.6	21.2	20.9
3	35.6	21.6	20.9
4	26.2	21.1	20.9
5	31.1	20.8	20.9

[00358] Exposure atmosphere oxygen concentration, temperature and relative humidity were considered acceptable throughout the study.

[00359] Aerosol Concentrations

[00360] Achieved gravimetric test atmosphere concentrations were as follows:

Group No.	Targeted Aerosol Concentration (mg\L)	Achieved Mean Aerosol Concentration (mg\L)	Coefficient of Variation (%)	% of Target
2	2.000*	1.966	34.7	98.3
3	0.200*	0.231	21.4	115.5
4	1.000*	0.928	20.5	92.8
5	2.000*	1.848	26.0	92.4

* Target aerosol concentrations were 0.140mg/L for Group 3, 0.700mg/L for Group 4 and 1.400mg/L for Groups 2 and 5 for the first 2 days of exposure.

[00361] Achieved test atmosphere concentrations for treprostinil were as follows:

Group No.	Targeted Aerosol Concentration (µg\L)	Achieved Mean Aerosol Concentration (µg\L)	Coefficient of Variation (%)	% of Target
3	1.0	1.10	22.1	110.0
4	5.0	4.44	21.0	88.8
5	10.0	8.94	28.1	89.4

[00362] Achieved test atmosphere concentrations for trehalose were as follows:

Group No.	Targeted Aerosol Concentration (µg\L)	Achieved Mean Aerosol Concentration (µg\L)	Coefficient of Variation (%)	% of Target
2	1684.6	1832.13	36.4	108.8
3	175.2	216.30	22.1	123.5
4	876.2	869.99	21.2	99.3
5	1752.5	1735.84	29.4	99.0

[00363] The overall achieved aerosol concentrations for all groups were within 20% of the targeted concentrations gravimetrically and for both treprostinil and trehalose, except for Group

3 for trehalose which was 23.5% greater than the targeted concentration. The generated atmospheres were considered stable over the treatment period even if all % CV were all above 20% as this was due the wrong targeted gravimetric concentrations being applied for the first 2 days of dosing. The overall aerosol concentrations were still considered acceptable for the study as there was a significant difference in aerosol concentration between groups.

[00364] Aerosol Homogeneity

[00365] Achieved gravimetric test atmosphere homogeneity concentrations were as follows:

Group No.	Aerosol Concentration of Top Tier (mg/L)	Aerosol Concentration of Middle Tier (mg/L)	Aerosol Concentration of Bottom Tier (mg/L)	CV (%)
2	1.061	1.029	1.078	2.4
3	0.134	0.136	0.126	4.0
4	0.810	0.877	0.845	4.0
5	1.225	1.263	1.280	2.2

[00366] Achieved test atmosphere homogeneity concentrations for treprostinil were as follows:

Group No.	Aerosol Concentration of Top Tier (µg/L)	Aerosol Concentration of Middle Tier (µg/L)	Aerosol Concentration of Bottom Tier (µg/L)	CV (%)
3	0.63	0.64	0.59	4.3
4	3.90	4.16	4.04	3.2
5	5.73	6.03	6.12	3.4

[00367] Achieved test atmosphere homogeneity concentrations for trehalose were as follows:

Group No.	Aerosol Concentration of Top Tier (µg/L)	Aerosol Concentration of Middle Tier (µg/L)	Aerosol Concentration of Bottom Tier (µg/L)	CV (%)
2	916.21	888.61	919.90	1.9
3	113.82	111.38	102.88	5.3
4	774.67	887.67	780.97	7.8
5	1091.32	1153.45	1160.14	3.3

[00368] Chamber homogeneity of the aerosol concentrations were considered acceptable since the coefficient of variance of aerosol concentration between samples was not greater than 20%.

[00369] Particle Size Distribution

[00370] The average gravimetric particle size distribution measurement data were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
2	89.1	81.8	51.3	19.3	11.6	9.4	7.9	0.0	2.0	2.46	78
3	97.9	95.0	79.6	35.5	21.1	15.1	5.7	0.0	1.3	1.96	94
4	95.5	89.4	60.3	24.1	13.4	9.0	4.8	0.0	1.7	2.07	88
5	96.9	92.3	63.1	28.2	15.7	8.5	4.8	0.0	1.6	1.99	91

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00371] The chemical determinations of particle size distribution for treprostinil were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
3	98.8	95.7	80.3	35.2	20.2	14.2	5.0	0.0	1.3	1.88	96
4	96.3	90.1	59.9	22.2	11.1	6.9	2.9	0.0	1.7	1.95	89
5	97.2	92.8	63.3	26.7	13.6	6.4	2.8	0.0	1.6	1.89	92

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00372] The chemical determinations of particle size distribution for trehalose were as follows:

Group No.	Cumulative % Less Than Stated Effective Cut-Off Diameter (µm)								Mean		% below 4 µm
	4.60	3.00	2.10	1.60	1.10	0.70	0.33	0.00	MMAD (µm)	GSD	
2	89.3	85.2	52.1	17.7	8.1	4.0	0.0	0.0	2.1	1.74	87
3	96.6	93.3	77.4	31.0	15.4	12.0	3.4	0.0	1.5	1.94	93
4	97.3	90.9	59.3	19.9	8.0	5.3	2.7	0.0	1.8	1.88	90
5	97.2	94.4	62.6	25.0	12.1	5.6	2.8	0.0	1.6	1.87	92

MMAD=Mass median aerodynamic diameter

GSD=Geometric standard deviation.

[00373] The particle size distribution was considered respirable for this study as all MMADs were below 4 µm and the GSDs were within 1.5 and 3.

[00374] Estimation of Achieved Dose Levels

[00375] Overall achieved doses for treprostinil are presented below:

Group No.	Targeted Dose Level (mg/kg/day)	Duration of Exposure (min)	Sex	Body Weight (kg)	Estimated Achieved Doses (mg/kg/day)	% of Targeted Dose Level
3	0.15	240	Male	0.324	0.16	106.7
			Female	0.227	0.17	113.3
			Combined	*0.276*	*0.17*	*113.3*
4	0.75	240	Male	0.316	0.66	88.0
			Female	0.229	0.70	93.3
			Combined	*0.273*	*0.68*	*90.7*
5	1.5	240	Male	0.315	1.33	88.7
			Female	0.228	1.42	94.7
			Combined	*0.272*	*1.37*	*91.3*

[00376] Overall achieved doses for trehalose are presented below:

Group No.	Targeted Dose Level (mg/kg/day)	Duration of Exposure (min)	Sex	Body Weight (kg)	Estimated Achieved Doses (mg/kg/day)	% of Targeted Dose Level
2	262.9	240	Male	0.316	273.4	104.0
			Female	0.229	290.8	110.6
			Combined	*0.273*	*281.2*	*107.0*
3	26.3	240	Male	0.324	32.1	122.1
			Female	0.227	34.4	130.8
			Combined	*0.276*	*33.1*	*125.9*
4	131.4	240	Male	0.316	129.8	98.8
			Female	0.229	138.1	105.1
			Combined	*0.273*	*133.5*	*101.6*
5	262.9	240	Male	0.315	259.2	98.6
			Female	0.228	275.7	104.9
			Combined	*0.272*	*266.6*	*101.4*

[00377] Overall achieved doses for leucine are presented below:

Group No.	Targeted Dose Level (mg/kg/day)	Duration of Exposure (min)	Sex	Body Weight (kg)	Estimated Achieved Doses[a] (mg/kg/day)	% of Targeted Dose Level
2	11.2	240	Male	0.316	11.7	104.5
			Female	0.229	12.4	110.7
			Combined	*0.273*	*12.0*	*107.1*
3	1.1	240	Male	0.324	1.2	109.1
			Female	0.227	1.3	118.2
			Combined	*0.276*	*1.3*	*118.2*
4	5.7	240	Male	0.316	5.0	87.7
			Female	0.229	5.3	93.0
			Combined	*0.273*	*5.1*	*89.5*
5	11.3	240	Male	0.315	10.0	88.5
			Female	0.228	10.7	94.7
			Combined	*0.272*	*10.3*	*91.2*

a = Calculated with a content of 4% of Leucine in PRINT-Tre and PRINT Placebo (using Trehalose percentage of 93.5% in PRINT Placebo for Group 2 and Treprostinil percentage of 0.53% in PRINT-Tre for Groups 3 to 5)

[00378] Average achieved dose levels for all groups were within 20% of the targeted dose levels, except for Group 3 for trehalose which was 26% above the targeted dose level; however, the dose levels were considered acceptable for the study as a clear dose differentiation between groups for each sex was achieved.

[00379] Mortality

[00380] There were no mortalities during the study.

[00381] Clinical Signs

[00382] There were no clinical signs observed during the study.

[00383] Body Weight

[00384] The only differences in body weight or weight gain potentially related to administration of the test or control item were slightly less growth (weight gain) in males given PRINT-Tre at 0.68 mg/kg/day and in both sexes given to PRINT-Tre at 1.37 mg/kg/day, relative to the air control group. The pattern of differences implicates the active ingredient treprostinil, not one of the excipients in PRINT-Tre.

[00385] These data are summarized in the table below, with differences potentially related to treprostinil in bold.

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Males					
Starting weight (Day 1) (g)	301	309	315	312	312
After 4 doses (Day 5) (g)	324	326	337	324	322
Change					
Absolute (g)	+23	+17	+22	+12	+10
Relative to Air Control	---	−6	−1	**−11**	**−13**
After 7 doses (fasted Day 8) (g)	300	302	310	301	298
Change					
Absolute (g)*	−1	−7	−5	−10	−14
Relative to Air Control	---	−6	−4	**−9**	**−13**
Females					
Starting weight (Day 1) (g)	220	223	220	227	226
After 4 doses (Day 5) (g)	231	235	232	237	231
Change					
Absolute (g)	+11	+12	+12	+10	+5
Relative to Air Control	---	+1	+1	−1	**−6**
After 7 doses (fasted Day 8) (g)	205	209	204	213	211
Change					
Absolute (g)*	−15	−14	−16	−14	−15
Relative to Air Control	---	+1	−1	−1	0

*All animals were fasted overnight prior to necropsy.

[00386] Remaining differences were considered incidental and of no biological significance.

[00387] Hematology

[00388] The only differences in mean hematology parameters potentially related to administration of the test or control item were greater mean reticulocyte counts in all groups given PRINT-Tre, relative to the air control group. The magnitude of difference was dose-related but statistically significant only in males. The pattern of differences implicates the active ingredient treprostinil, not one of the excipients in PRINT-Tre.

[00389] These data are summarized in the table below, with differences potentially related to treprostinil in bold.

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Reticulocyte count					
Males					
Mean (x 10^{12}/L)	0.226	0.228	0.309	0.333	0.337
Relative to Air Control	---	+1%	**+37%**	**+47%**	**+49%**
Females					
Mean (x 10^{12}/L)	0.179	0.176	0.214	0.290	0.283
Relative to Air Control	---	+1%	**+20%**	**+62%**	**+58%**

[00390] An increase in reticulocyte count is an appropriate response to an increased demand for RBCs. In this study, greater reticulocyte counts were not associated with differences in circulating erythron mass (i.e., no differences in RBC count, haemoglobin concentration, or haematocrit). This suggests that the increased release of reticulocytes was accompanied by, and probably a response to, an increased rate of RBC loss, and that the erythropoietic response was adequate to maintain normal circulating RBC numbers.

[00391] Remaining differences among mean hematology parameters were considered incidental and of no biological significance.

[00392] Coagulation

[00393] There were no differences in mean coagulation parameters that were considered to be related to administration of the test or control item. All differences were considered incidental and of no biological significance.

[00394] Clinical Chemistry

[00395] There were no differences in mean clinical chemistry parameters that were considered to be related to administration of the test or control item. All differences were considered incidental and of no biological significance.

[00396] Urinalysis

[00397] There were no differences in the urinalysis parameters that were considered to be related to administration of the test or control item. All differences were considered incidental and of no biological significance.

[00398] Organ Weights

[00399] Differences in mean organ weight potentially related to administration of the test or control item were noted for lungs, adrenal glands, thymus, and testes.

[00400] Remaining differences in mean organ weight were considered incidental and of no biological significance.

[00401] Lungs

[00402] Mean lungs/trachea weights (absolute and relative to body weight) were greater in all groups given the test or control item, compared to the air control group. The differences were greater with PRINT-Tre than with PRINT Placebo, and the differences were dose-related for PRINT-Tre. This pattern suggests that administration of the excipients (likely trehalose) resulted in a slight (15% to 17%) increase in lung weight, which was exacerbated by co-administration of treprostinil as the lung weights of PRINT-Tre groups were increased compared to the lung weights of the PRINT Placebo group.

[00403] There was a histopathologic finding in the lungs that might have accounted for the greater lung weight; specifically, increased alveolar macrophages with basophilic vacuolated cytoplasm in the lungs of all rats given PRINT Placebo or PRINT-Tre at ≥0.68 mg/kg/day. However, neither the distribution of this histopathologic finding across groups nor the grade of the finding correlated well with the differences in mean lung weight, suggesting that some other factor was responsible. Because lungs were weighed before fixation, it is possible that some material responsible for the greater weight was removed during tissue fixation and processing.

[00404] Lung weight data are summarized in the table below, with differences potentially related to PRINT Placebo and PRINT-Tre in bold.

Mean Lung Weight Data

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Males					
Absolute weight (g)	1.39	1.60	1.70	1.78*	1.90*
Relative to Air Control	---	**+15%**	**+22%**	**+28%**	**+37%**
Relative weight (% body weight)	0.46	0.53	0.55	0.59*	0.64*
Relative to Air Control	---	**+15%**	**+20%**	**+28%**	**+39%**
Females					
Absolute weight (g)	1.09	1.27	1.31*	1.48*	1.44*
Relative to Air Control	---	**+17%**	**+20%**	**+36%**	**+32%**
Relative weight (% body weight)	0.53	0.61	0.64	0.70*	0.68*
Relative to Air Control	---	**+15%**	**+21%**	**+32%**	**+28%**

*Statistically significant compared to air control; Dunnett's 2-sided, $p<0.05$

[00405] Thymus

[00406] Mean thymus weights (absolute and relative to body weight) were slightly lower in both sexes given PRINT-Tre at 0.68 mg/kg/day and in both sexes given to PRINT-Tre at 1.37 mg/kg/day, relative to the air control group (though not statistically significantly different). The pattern of differences implicates the active ingredient treprostinil, not one of the excipients in PRINT-Tre as differences were also seen between PRINT-Tre groups and PRINT Placebo

group. Lower thymus weight was not associated with lower lymphocyte count or with any histopathologic findings.

[00407] Lower thymus weight is one common manifestation of nonspecific physiological or psychological stress (Everds et al., 2013). Because this finding was associated with reduced weight gain (growth) and sometimes also with greater adrenal glands weight, it was most likely secondary to stress and not a direct effect of treprostinil.

[00408] Thymus weight data are summarized in the table below, with differences potentially related to treprostinil in bold.

Mean Thymus Weight Data

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Males					
Absolute weight (mg)	504	593	537	388	429
Relative to Air Control	---	+18%	+7%	**−23%**	**−15%**
Relative weight (% body weight)	0.168	0.198	0.173	0.128	0.143
Relative to Air Control	---	+18%	+3%	**−24%**	**−15%**
Females					
Absolute weight (mg)	469	437	465	428	352
Relative to Air Control	---	−7%	−1%	−9%	**−25%**
Relative weight (% body weight)	0.229	0.210	0.228	0.202	0.166
Relative to Air Control	---	−8%	±0%	−12%	**−28%**

[00409] Adrenal Glands

[00410] Mean adrenal glands weight (absolute and relative to body weight) was greater in males given PRINT-Tre at 0.17 mg/kg/day and in both sexes given to PRINT-Tre at 1.37 mg/kg/day, relative to the air control group (though not statistically significantly different). While the differences may have been due to chance and a consequence of the small group sizes (3/sex), the pattern of differences raises the possibility that they are related to administration of

treprostinil, at least at the high-dose level as differences were also seen between the high dose PRINT-Tre group and the PRINT Placebo group. Greater adrenal glands weight was not associated with any histopathologic findings.

[00411] Greater adrenal glands weight is one common manifestation of nonspecific physiological or psychological stress (Everds et al., 2013). Because this finding was associated with reduced weight gain (growth) and lower thymus weight at the high-dose level, it was most likely secondary to stress and not a direct effect of treprostinil.

[00412] Adrenal glands weight data are summarized in the table below, with differences potentially related to treprostinil in bold.

Mean Adrenal Glands Weight Data

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Males					
Absolute weight (mg)	62	71	76	68	78
Relative to Air Control	---	+15%	**+23%**	+10%	**+26%**
Relative weight (% body weight)	0.021	0.023	0.025	0.023	0.027
Relative to Air Control	---	+10%	**+19%**	+10%	**+29%**
Females					
Absolute weight (mg)	74	73	70	76	89
Relative to Air Control	---	−1%	−5%	+3%	**+20%**
Relative weight (% body weight)	0.036	0.035	0.035	0.036	0.042
Relative to Air Control	---	−3%	−3%	±0%	**+17%**

[00413] Testes

[00414] There was a trend toward slightly lower mean testes weight (absolute and relative to body weight) in groups given PRINT-Tre at ≥0.68 mg/kg/day, relative to the air control group. While the differences may have been due to chance and a consequence of the small group sizes (3/sex), the pattern of differences raises the possibility that they are related to administration of

treprostinil as differences were also seen between the mid and high dose PRINT-Tre groups and the PRINT Placebo group. Slightly lower testes weight was not associated with any histopathologic findings.

[00415] Testes weight data are summarized in the table below, with differences potentially related to treprostinil in bold.

Mean Testes Weight Data

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Absolute weight (g)	3.51	3.40	3.39	3.25	3.15
Relative to Air Control	---	−3%	−3%	**−7%**	**−10%**
Relative weight (% body weight)	1.17	1.13	1.10	1.08	1.06
Relative to Air Control	---	−3%	−6%	**−8%**	**−9%**

[00416] Macroscopic Findings

[00417] There was no evidence of test item-related macroscopic findings at necropsy.

[00418] All findings were considered to be incidental as they were not dose-related, of low incidence, or occurred in the air control, placebo control and treated animals.

[00419] Microscopic Findings

[00420] Treatment-related findings were observed in the lungs, anterior nasal cavity, and nasopharynx. All other microscopic findings were considered to be incidental or procedure-related.

[00421] Lungs

[00422] In the lungs, minimal to mild increased alveolar macrophages with basophilic vacuolated cytoplasm were observed in all rats given PRINT Placebo or PRINT-Tre at ≥0.68 mg/kg/day. The pattern of this finding across groups indicates that it is a response to the

excipients (likely trehalose). There were no associated inflammatory changes in the lungs. Increased alveolar macrophages are a common finding in inhalation toxicity studies with powders. It reflects normal pulmonary clearance of inhaled particles and is not considered to be adverse.

[00423] These data are summarized in the table below, with differences potentially related to test or control item in bold.

Incidence and Grade of Increased Alveolar Macrophages

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Males					
Incidence	0/3	**3/3**	0/3	**3/3**	**3/3**
Mean grade	---	**1.7**	---	**1.0**	**1.7**
Females					
Incidence	0/3	**3/3**	0/3	**3/3**	**3/3**
Mean grade	---	**2.0**	---	**1.0**	**1.7**

[00424] Nasal Cavity and Nasopharynx

[00425] Goblet-cell hypertrophy/hyperplasia was seen in the cranial portion of the nasal cavity and in the nasopharynx of at least one rat in all groups given PRINT Placebo or PRINT-Tre, but the incidence was greater in groups given PRINT-Tre at ≥0.68 mg/kg/day, and the mean grade was greater in the group given PRINT-Tre at 1.37 mg/kg/day. This pattern suggests that administration of the excipients (likely trehalose) resulted in occasional goblet-cell changes, which were exacerbated by co-administration of treprostinil at higher dose levels.

[00426] Goblet cell hypertrophy/hyperplasia in the anterior nasal cavity and nasopharynx is one of the most frequently observed lesions in rodents exposed to irritant compounds. This finding generally is considered a nonspecific protective or adaptive response and not adverse.

[00427] These data are summarized in the table below, with differences potentially related to test or control item in bold.

Incidence and Grade of Goblet-cell Hypertrophy/Hyperplasia

Test Material =	**Air**	**PRINT Placebo**	**PRINT-Tre**		
Treprostinial Dose Level (mg/kg/day) =	**0**	**0**	**0.17**	**0.68**	**1.37**
Trehalose Dose Level (mg/kg/day) =	**0**	**281**	**33**	**134**	**267**
Nasal Cavity					
Males					
Incidence	0/3	**1/3**	**1/3**	**3/3**	**3/3**
Mean grade	---	1.0	1.0	1.0	**1.3**
Females					
Incidence	0/3	0/3	0/3	**1/3**	**2/3**
Mean grade	---	---	---	1.0	**1.5**
Nasopharynx					
Males					
Incidence (all graded minimal)	0/3	0/3	0/3	**3/3**	**3/3**
Females					
Incidence(all graded minimal)	0/3	**1/3**	0/3	**3/3**	**3/3**

[00428] Discussion and Conclusions

[00429] Rats tolerated daily administration of PRINT Placebo or PRINT-Tre at up to 1.37 mg/kg/day by 4-hour inhalation for 7 days.

[00430] The only findings potentially related to administration of excipients (likely trehalose) were:

- Increased alveolar macrophages with basophilic vacuolated cytoplasm in all rats given PRINT Placebo or PRINT-Tre at ≥0.68 mg/kg/day; *i.e.*, in rats given trehalose at ≥134 mg/kg/day. The mean grade of this finding increased with trehalose dose level. This finding was not associated with inflammatory changes in the lungs and was considered to reflect normal pulmonary clearance of inhaled particles. It was not considered to be adverse.
- Greater mean lung weight in groups given PRINT Placebo or PRINT-Tre. The weight differences were unrelated to trehalose dose level. Instead, they were greater with PRINT-Tre than with PRINT Placebo and were dose-related for PRINT-Tre. This pattern suggests that administration of the excipients (likely trehalose) resulted in a slight (15% to 17%) increase in lung weight, which was exacerbated by co-administration of treprostinil. Of note, the pattern of differences in lung weight across groups is distinct from the pattern of increased alveolar macrophages across groups, indicating that the weight differences were not a consequence of increased macrophages. There were no histopathologic findings in the lungs that might have accounted for the greater lung weight. Because lungs were weighed before fixation, it is possible that some material responsible for the greater weight was removed during tissue fixation and processing.
- Minimal goblet-cell hypertrophy/hyperplasia in the cranial portion of the nasal cavity of at least one rat in all groups given PRINT Placebo or PRINT-Tre. The incidence of this finding was unrelated to trehalose dose level. Instead, the incidence was greater with PRINT-Tre at ≥0.68 mg/kg/day, and the mean grade was greater with PRINT-Tre at 1.37 mg/kg/day. This pattern suggests that administration of the excipients (likely trehalose) resulted in occasional goblet-cell changes, which were exacerbated by co-administration of treprostinil at higher dose levels. Goblet-cell hypertrophy/hyperplasia was considered a nonspecific protective or adaptive response and not adverse.

[00431] Besides exacerbating lung weight differences and goblet-cell hypertrophy/hyperplasia in the nasal cavity and nasopharynx, the following other findings were potentially related to administration of treprostinil as PRINT-Tre:

- Slightly less growth (weight gain) in males at 0.68 mg/kg/day and in both sexes at 1.37 mg/kg/day.

- Greater mean reticulocyte counts at all dose levels, with the magnitude of difference increasing with dose level. This was not considered adverse in and of itself; however, it likely reflected an appropriate adaptive response to an increased rate of RBC loss or turnover.
- Greater mean adrenal glands weight in males at 0.17 mg/kg/day, lower mean thymus weight in both sexes at 0.68 mg/kg/day, and greater mean adrenal glands weight and lower mean thymus weight in both sexes at 1.37 mg/kg/day. There were no associated differences in lymphocyte count or histopathologic findings in either organ. These organ weight differences most likely reflected stress and were not a direct effect of treprostinil.

[00432] Based on these results, it is recommended that an upcoming 14-day GLP inhalation toxicology study in rats target similar dose levels as used in the current study.

[00433] CLINICAL STUDY: LIQ861

[00434] Randomized, Placebo-controlled, Single-ascending Dose Study Evaluating Pharmacokinetics (PK) and Safety in Healthy Male and Female Volunteers

[00435] A clinical study was conducted to (1) determine the single-dose safety and tolerability and (2) evaluate the single-dose pharmacokinetics of particles of the invention upon administration to healthy male and female subjects.

[00436] Six cohorts were evaluated: dose levels of 25, 50, 75, 100, 125 and 150 μg of treprostinil respectively. In each cohort, eight subjects were randomly assigned in a 3:1 blinded ratio and received a single dose of either particles of the invention (N = 6) or placebo particles (N = 2).

[00437] Blood was collected for PK evaluation at T = 0, 5, 10, 15, 20, 25, 30, 45, 60, 90, and 120 minutes and 3, 4, 6, and 8 hours post-dose.

[00438] Cohort 1

[00439] Eight subjects were enrolled and dosed in Cohort 1. Six subjects received active treatment and 2 received placebo. Active treatment was administered by dry powder inhalation (DPI) as a single capsule of 25 μg treprostinil strength, and placebo treatments were administered

by DPI as a single capsule of the placebo formulation. All inhalations were administered using the RS00 inhaler.

[00440] Blood samples were obtained prior to dosing and at nominal times of 0.083, 0.167, 0.25, 0.33, 0.417, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 6, and 8 hours after inhalation. Analysis of plasma concentration versus time data for calculation of standard pharmacokinetic (PK) parameters following inhalation was conducted using Phoenix WinNonlin version 6.3 using scheduled blood sampling times. Plasma concentrations were provided only for subjects on active dose and random subject numbers were assigned by the bioanalytical laboratory to retain the study blind.

[00441] The table shown in Figure 3A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 1. Preliminary non-compartmental PK parameters for treprostinil are summarized in the table shown in Figure 3B. The highest concentrations for three of the six subjects occurred at 0.33 hours post-inhalation; one subject each had a Tmax of 0.167, 0.25, and 0.417 hours post-dose. Concentrations subsequently decayed with a single-phase disposition profile, as shown in the log-linear plots. At two hours after inhalation, two of six active subjects had measurable concentrations of treprostinil and only one subject had measurable concentrations at 2.5 and 3 hours after inhalation. No subjects had quantifiable concentrations after the 3 hour timepoint.

[00442] The Cmax averaged 0.364 ng/mL and the most frequent Tmax was 0.33 hours after inhalation. AUCinf values averaged 0.301 h*ng/mL with a CV% of 30.2%. The apparent volume of distribution (Vz/F) averaged 68.1 L. Oral clearance (CL/F) averaged 91.0 L/h and ranged from 59.1 to 150. Variability in the CL/F value had a CV% of 35.8%.

[00443] Cohort 2

[00444] Nine subjects were enrolled and dosed in Cohort 2. At least six subjects received active treatment and at least 2 received placebo; 1 subject withdrew before the 2 hour PK sample and was replaced. Subjects with truncated sampling schedules have been excluded in this interim analysis. Active treatment was administered by dry powder inhalation (DPI) as a single capsule of 50 μg treprostinil strength, and placebo treatments were administered by DPI as a

single capsule of the placebo formulation. All inhalations were administered using the RS00 inhaler.

[00445] Blood samples were obtained prior to dosing and at nominal times of 0.083, 0.167, 0.25, 0.33, 0.417, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 6, and 8 hours after inhalation. Analysis of plasma concentration versus time data for calculation of standard pharmacokinetic (PK) parameters following inhalation was conducted using Phoenix WinNonlin version 6.3 using scheduled blood sampling times. Plasma concentrations were provided only for subjects on active dose and random subject numbers were assigned by the bioanalytical laboratory to retain the study blind.

[00446] The table shown in Figure 4A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 2.

[00447] The highest concentrations for four of the six subjects occurred at 0.17 hours post-inhalation; of the remaining subjects, one had Tmax at 0.083 hours post-dose and one at 0.417 hours post-dose. At 2.5 hours after inhalation, 2 of 6 active subjects had measurable concentrations of treprostinil and only one subject had measurable concentrations at 3 hours after inhalation. No subjects had quantifiable concentrations after the 3 hour timepoint.

[00448] Preliminary non-compartmental PK parameters for treprostinil for Cohort 2 are summarized in the table shown in Figure 4B. The Cmax averaged 0.572 ng/mL and the most frequent Tmax was 0.167 hours after inhalation. AUCinf values averaged 0.422 h*ng/mL with a CV% of 62.8%. The apparent volume of distribution (Vz/F) averaged 110 L. Oral clearance (CL/F) averaged 208 L/h and ranged from 67 to 624. Variability in the CL/F value had a CV% of 101.5%.

[00449] By comparison, the Cmax for Cohort 1 averaged 0.364 ng/mL and the AUCinf values averaged 0.301 h*ng/mL. Thus, a doubling of the treprostinil dose resulted in an approximate 50% increase in exposure. The Vz/F and the CL/F values were considerably higher for Cohort 2 and with greater variability.

[00450] Cohort 3

[00451] Eight subjects were enrolled and dosed in Cohort 3. Six subjects received active treatment and two received placebo. Active treatment was administered by dry powder inhalation (DPI) as a single capsule of 75 μg treprostinil strength and placebo treatments were administered by DPI as a single capsule of the placebo formulation. All inhalations were administered using the RS00 inhaler.

[00452] Blood samples were obtained prior to dosing and at nominal times of 0.083, 0.167, 0.25, 0.33, 0.417, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 6, and 8 hours after inhalation. Analysis of plasma concentration versus time data for calculation of standard pharmacokinetic (PK) parameters following inhalation was conducted using Phoenix WinNonlin version 6.3 using scheduled blood sampling times. Plasma concentrations were provided only for subjects on active dose and random subject numbers were assigned by the bioanalytical laboratory to retain the study blind.

[00453] The table shown in Figure 5A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the 6 active subjects in Cohort 3.

[00454] The highest concentrations for three of the six subjects occurred at 0.25 hours post-inhalation; of the remaining subjects, 1 had Tmax at 0.083 hours post-dose, 1 at 0.17 hours post-dose, and 1 at 0.417 hours post-dose. At 3 hours after inhalation, two of six active subjects had measurable concentrations of treprostinil. No subjects had quantifiable concentrations after the 3 hour timepoint.

[00455] Preliminary non-compartmental PK parameters for treprostinil for Cohort 3 are summarized in the table shown in Figure 5B. The Cmax averaged 0.728 ng/mL and the most frequent Tmax was 0.25 hours after inhalation. AUCinf values averaged 0.757 h*ng/mL with a CV% of 39.4%. The apparent volume of distribution (Vz/F) averaged 97 L. Oral clearance (CL/F) averaged 112 L/h and ranged from 58 to 161. Variability in the CL/F value had a CV% of 39.4%.

[00456] By comparison, the Cmax for Cohort 1 and Cohort 2 averaged 0.364 ng/mL and 572 ng/mL, respectively, while the AUCinf values averaged 0.301 h*ng/mL and 0.422 h*ng/mL. Thus, a tripling of the dose from Cohort 1 resulted in an approximate 100 – 150% increase in

exposure. The CL/F values were for Cohort 3 were more consistent with Cohort 1, and with similar variability, than what was observed in Cohort 2. The results indicate that both Cmax and AUCinf may be increasing proportionately to the increase in the dose and that the CL is independent of dose over the range of 25 to 75 μg treprostinil.

[00457] Cohort 4

[00458] Eight subjects were enrolled and dosed in Cohort 4. Six subjects received active treatment and two received placebo. Active treatment of 100 μg treprostinil was administered by dry powder inhalation (DPI) as 2 capsules of 50 μg treprostinil strength and placebo treatments were administered by DPI as 2 capsules of the placebo formulation. All inhalations were administered using the RS00 inhaler.

[00459] Blood samples were obtained prior to dosing and at nominal times of 0.083, 0.167, 0.25, 0.33, 0.417, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 6, and 8 hours after inhalation. Analysis of plasma concentration versus time data for calculation of standard pharmacokinetic (PK) parameters following inhalation was conducted using Phoenix WinNonlin version 6.3 using scheduled blood sampling times. Plasma concentrations were provided only for subjects on active dose and random subject numbers were assigned by the bioanalytical laboratory to retain the study blind.

[00460] The table shown in Figure 6A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 4. The highest concentrations for 2 of the 6 subjects occurred at 0.25 hours post-inhalation; of the remaining subjects, 2 had Tmax at 0.5 hours post-dose, 1 at 0.17 hours post-dose, and 1 at 0.33 hours post-dose. At 4 hours after inhalation, 3 of 6 active subjects had measurable concentrations of treprostinil. No subjects had quantifiable concentrations at the 6 or 8 hour timepoints.

[00461] Preliminary non-compartmental PK parameters for treprostinil for Cohort 4 are summarized in Figure 6B. The Cmax averaged 1.08 ng/mL and the most frequent Tmax values were observed at 0.25 hours and 0.5 hours after inhalation. AUCinf values averaged 1.22 h*ng/mL with a CV% of 18.4%. The apparent volume of distribution (Vz/F) averaged 96 L.

Oral clearance (CL/F) averaged 84.8 L/h and ranged from 68.3 to 122. Variability (CV%) in the CL/F value was 22.8%.

[00462] By comparison, the Cmax for Cohorts 1, 2, and 3 averaged 0.364 ng/mL, 0.572 ng/mL, and 0.728 ng/mL respectively, while the AUCinf values averaged 0.301 h*ng/mL, 0.422 h*ng/mL, and 0.757 h*ng/mL. Thus, a quadrupling of the dose from Cohort 1 resulted in an approximate 200 – 300% increase in exposure, while a doubling of the dose from Cohort 2 resulted in an approximate 2-fold increase in exposure. The results indicate that both Cmax and AUCinf may be increasing proportionately to the increase in the dose and that the CL/F is independent of dose over the range of 25 to 100 μg treprostinil.

[00463] Cohort 5

[00464] Eight subjects were enrolled and dosed in Cohort 5. Six subjects received active treatment and two received placebo. Active treatment of 125 μg treprostinil was administered by dry powder inhalation (DPI) as 1 capsule of 75 μg and 1 capsule of 50 μg treprostinil strength and placebo treatments were administered by DPI as 2 capsules of the placebo formulation. All inhalations were administered using the RS00 inhaler.

[00465] Blood samples were obtained prior to dosing and at nominal times of 0.083, 0.167, 0.25, 0.33, 0.417, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 6, and 8 hours after inhalation. Analysis of plasma concentration versus time data for calculation of standard pharmacokinetic (PK) parameters following inhalation was conducted using Phoenix WinNonlin version 6.3 using scheduled blood sampling times. Plasma concentrations were provided only for subjects on active dose and random subject numbers were assigned by the bioanalytical laboratory to retain the study blind.

[00466] The table shown in Figure 7A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the 6 active subjects in Cohort 5. The highest concentrations for 3 of the 6 subjects occurred at 0.17 hours post-inhalation; of the remaining subjects, 2 had Tmax at 0.33 hours post-dose, and 1 at 0.42 hours post-dose. At 3.5 and 4 hours after inhalation, only 1 of 6 active subjects had measurable concentrations of treprostinil. No subjects had quantifiable concentrations at the 6 or 8 hour timepoints.

[00467] Preliminary non-compartmental PK parameters for treprostinil for Cohort 5 are summarized in Figure 7B. The Cmax averaged 1.19 ng/mL and the most frequent Tmax values were observed at 0.17 hours after inhalation. AUCinf values averaged 1.15 h*ng/mL with a CV% of 44.9%. The apparent volume of distribution (Vz/F) averaged 101 L. Oral clearance (CL/F) averaged 141 L/h and ranged from 65.7 to 336. Variability (CV%) in the CL/F value was 69.9%.

[00468] By comparison, the Cmax for Cohorts 1, 2, 3, and 4 averaged 0.364 ng/mL, 0.572 ng/mL, 0.728 ng/mL, and 1.08 ng/mL respectively, while the AUCinf values averaged 0.301 h*ng/mL, 0.422 h*ng/mL, 0.757 h*ng/mL, and 1.22 h*ng/mL. Thus, a quintupling of the dose from Cohort 1 resulted in an approximate 220 – 280% increase in exposure. The results indicate that both Cmax and AUCinf may be increasing proportionately to the increase in the dose and that the CL/F is independent of dose over the range of 25 to 125 μg treprostinil.

[00469] Cohort 6

[00470] Cohort 6 was conducted as an original and a repeat. In each Cohort 6 (original and repeat), eight subjects were enrolled and dosed. Six subjects received active treatment and two received placebo. Active treatment of 150 μg treprostinil was administered by dry powder inhalation (DPI) as 2 capsules of 75 μg treprostinil strength and placebo treatments were administered by DPI as 2 capsules of the placebo formulation. All inhalations were administered using the RS00 inhaler. Cohort 6 original included some mechanical device failures and subject non-compliance with instructions, giving rise to Cohort 6 repeat.

[00471] Blood samples were obtained prior to dosing and at nominal times of 0.083, 0.167, 0.25, 0.33, 0.417, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 6, and 8 hours after inhalation. Analysis of plasma concentration versus time data for calculation of standard pharmacokinetic (PK) parameters following inhalation was conducted using Phoenix WinNonlin version 6.3 using scheduled blood sampling times. Plasma concentrations were provided only for subjects on active dose and random subject numbers were assigned by the bioanalytical laboratory to retain the study blind.

[00472] The table shown in Figure 8A contains a summary of the treprostinil concentration-time data for individual subjects with descriptive statistics for the six active subjects in Cohort 6-R. The highest concentrations for 2 of the 6 subjects occurred at 0.25 hours post-inhalation and at 0.33 hours post-inhalation. In the remaining 2 subjects, Tmax occurred at the 0.167 and 0.417 hours post-dose timepoints. At 4 hours after inhalation, 4 of 6 active subjects had measurable concentrations of treprostinil. No subjects had quantifiable concentrations at the 6 or 8 hour timepoints.

[00473] Preliminary non-compartmental PK parameters for treprostinil for Cohort 6-R are summarized in Figure 8B. Preliminary non-compartmental PK parameters for treprostinil for Cohort 6-Original are summarized in Figure 8C. Mean concentration-time data for each of the six cohorts is displayed on a linear scale in Figure 8D. The Cmax averaged 1.45 ng/mL and the most frequent Tmax values were observed at 0.25 and 0.33 hours after inhalation. AUCinf values averaged 1.62 h*ng/mL with a CV% of 68.3%. The apparent volume of distribution (Vz/F) averaged 107 L. Oral clearance (CL/F) averaged 126 L/h and ranged from 51.8 to 245. Variability (CV%) in the CL/F value was 68.3%.

[00474] By comparison, the Cmax for Cohorts 1, 2, 3, 4, 5, and the combined Cohort 6 averaged 0.364 ng/mL, 0.572 ng/mL, 0.728 ng/mL, 1.08 ng/mL, 1.19 ng/mL, and 1.21 ng/mL, respectively (Figure 8E), while the AUCinf values averaged 0.301 h*ng/mL, 0.422 h*ng/mL, 0.757 h*ng/mL, 1.22 h*ng/mL, 1.15 h*ng/mL, and 1.37 h*ng/mL (Figure 8F). Thus, a 6-fold increase of the dose from Cohort 1 to the combined Cohort 6 observations resulted in an approximate 260 – 400% increase in exposure, while tripling from Cohort 2 and doubling from Cohort 3 resulted in approximate increases of exposure by 130 – 255% and 81 – 98%, respectively. Moreover, plots of the relationship between dose and Cmax and AUCinf are displayed in Figure 8E and Figure 8F, respectively. The results indicate that both Cmax and AUCinf may be increasing proportionately to the increase in the dose. It was observed at the CRU during the original 150 μg dosing, however, that there were several apparent device failures that may have resulted in incomplete and/or inefficient exposures. It should be noted that no device failures were noted during the repeat dosing and, while mean values may be higher than in the initial cohort, the variability in the repeated cohort is greater. A plot of the relationship between dose and CL/F (Figure 8G) shows that the CL/F is independent of dose over the range

of 25 to 150 μg treprostinil, which suggests that PK of treprostinil has a proportional relationship to dose over the range of 25 to 150 μg treprostinil.

[00475] CLINICAL CONCLUSIONS

[00476] LIQ861 was dosed at levels of 25, 50, 75, 100, 125 and 150 μg treprostinil from either a single capsule (25, 50 and 75 mcg doses) or a combination of two lower capsular strengths (for 100, 125 and 150 mcg doses), each capsule either undergoing either a single breath or two breaths. According to embodiments of the present invention, novel capsule to particle powder to active ingredient ratios, and breath per capsule and powder per breath ratios for human dosing are included in the following table.

Patient presentation of particle powder and active agent per capsule per breath for particle formulation having 0.5 percent active agent load						
Capsules	1	1	1	2	2	2
Particle Powder in mg	5	10	15	Combination of two 50 mcg capsules or one 25 mcg and one 75 mcg	Combination of, ex., 1 at 50 mcg and 1 at 75 mcg	Combination of, ex., two capsules at 75 mcg
Active Agent Load in mcg	25	50	75	Varies, see above	Varies, see above	Varies, see above
Breaths to Deliver	1 to 2	1 to 2	1 to 2	1 to 2 per capsule	1 to 2 per capsule	1 to 2 per capsule

Particle Powder per Breath in mg	2.5-5	5-10	7.5-15	Varies, up to 15	Varies, up to 15	Varies, up to 15
Active Agent per Breath in mcg	12.5-25	25-50	37.5-75	Varies, up to 75	Varies, up to 75	Varies, up to 75

[00477] According to such embodiments, as shown in the above table, each breath can receive from 2.5 – 15 mg of particle power and from 12.5 – 75 mcg of active agent.

[00478] For the given treprostinil delivered in the given mass of particle powder loaded into a capsule and delivered through a dry powder inhaler results in the human clinical outcomes are included in the following table for LIQ861.

LIQ861 Clinical Outcomes	C_{max} (ng/mL)	T_{max}[a] (h)	$t_{½}$ (h)	AUC_{last} (h*ng/mL)	AUC_{inf} (h*ng/mL)	CL/F (L/h)	Vz/F (L)
25 mcg Treprostinil	0.36 (0.12)	0.33 (0.17, 0.42)	0.52 (0.16)	0.27 (0.09)	0.3 (0.09)	91 (32.6)	68.1 (27.4)
50 mcg Treprostinil	0.57 (0.37)	0.17 (0.08, 0.42)	0.45 (0.12)	0.4 (0.26)	0.42 (0.27)	208 (211)	110 (66.6)
75 mcg Treprostinil	0.73 (0.3)	0.25 (0.08, 0.42)	0.62 (0.18)	0.72 (0.31)	0.76 (0.31)	112 (38.5)	97 (29.1)
100 mcg Treprostinil	1.08 (0.31)	0.29 (0.17, 0.5)	0.78 (0.13)	1.18 (0.22)	1.22 (0.23)	84.8 (19.3)	95.5 (28.2)
125 mcg Treprostinil	1.19 (0.53)	0.25 (0.17, 0.42)	0.53 (0.07)	1.12 (0.51)	1.15 (0.52)	141 (98.8)	101 (58.7)
150 mcg Treprostinil	1.21 (0.3)	0.29 (0.08, 0.42)	0.66 (0.15)	1.33 (0.44)	1.37 (0.42)	119 (35.8)	115 (51.4)
150 mcg Treprostinil	1.45 (0.63)	0.29 (0.17, 0.42)	0.64 (0.11)	1.58 (0.85)	1.62 (0.87)	126 (80.3)	107 (54)

Abbreviations: SD = standard deviation; C_{max} = maximum observed plasma concentration; T_{max} = time to C_{max}; $t_{½}$ = half-life; AUC = area under the curve; CL/F = apparent clearance; Vz/F = apparent volume of distribution

All values except for T_{max} are reported as arithmetic means with SD in parentheses.

[a] T_{max} reports median values with minimum and maximum values in parentheses

[00479] For comparison, TYVASO (United Therapeutics, Inc.) provides the current standard of treatment for inhaled treprostinil. Such treprostinil is delivered through a nebulizer for the treatment of PAH and is limited to deliver 6mcg of treprostinil per breath, utilizing 9 breaths to reach a 54 mcg dose. The current standard of inhaled treatment has shown to be dose limited to a maximum tolerated dose of 84 mcg of treprostinil, which required 14 breaths to reach such dose. See, Channick, R. et al., Inhaled Treprostinil: a therapeutic review, Drug Design, Development and Therapy 2012:6 19-28; and Nelsen AC, et al., Pharmacokinetics Of Inhaled Treprostinil Sodium In Healthy Volunteers. Am J Respir Crit Care Med. 2010; 181:A3348; both of which are incorporated herein by reference in their entirety.

[00480] In alternative embodiments, particles of the present invention may include 1% treprostinil load, as compared to 0.5% treprostinil load of the LIQ861 particles. According to an

embodiment of the present invention, a plurality of 1% treprostinil particles were fabricated from a solution comprising, weight percent solids in water, of: 1.06% treprostinil sodium, 92.44% trehalose dihydrate, 2% polysorbate 80, 4% L leucine, 0.27% sodium citrate dihydrate, and 0.23% sodium chloride.

[00481] According to a 1 percent treprostinil particle formulation of the present invention, particle powder mass and active agent presented to a patient comprise the following novel capsule to particle powder to active ingredient ratios, and breath per capsule and powder per breath ratios for human dosing.

Patient presentation of particle powder and active agent per capsule per breath for particle formulation having 1 percent active agent load						
Capsules	1	1	1	1	1	1
Particle Powder in mg	2.5	5	7.5	10	12.5	15
Active Agent Load in mcg	25	50	75	100	125	150
Breaths to Deliver	1 to 2	1 to 2	1 to 2	1 to 2	1 to 2	1 to 2
Particle Powder per Breath in mg	1.25-2.5	2.5-5	3.75-7.5	5-10	6.25-12.5	7.5-15
Active Agent per Breath in mcg	12.5-25	25-50	37.5-75	50-100	62.5-125	75-150

[00482] According to such embodiments, as shown in the above table, each breath can receive from 1.25 – 15 mg of particle power and from 12.5 – 150 mcg of active agent.

[00483] For the powder mass found acceptable in LIQ861 initial clinical trial associated with delivery of the 150 mcg dose, at a 1% active drug particle a dose of 300 mcg of active drug can be administered in a safe and acceptable powder mass and excipient quantity.

[00484] Kits

[00485] According to embodiments of the present invention the dry powder inhaler device can be combined into a kit with capsules for use therein. The capsules can be packaged in blister packs with or without desiccant to ensure controlled environment for the LIQ861 particle powder while the traveling with a user. The blister packs can include capsules for a single dosing or multiple capsules for a day, week or month of doses. Typically a patient will treat 4 times per day for the PAH indication. The kit can include capsules comprising dosage strengths of 25, 50, 75, 100, 125, 150, 200, 250, 300 mcg or beyond for the treatment of PAH. The particles of the powder in the capsules of the kits can be particles comprising 0.5% treprostinil or 1% treprostinil.

[00486] Abbreviations and Nomenclature Cross-references

6MWD	6 Minute Walk Distance
AE	Adverse Event
AUC	Area Under the Curve
AUCinf	Area Under the Concentration-Time Curve Extrapolated to Time Infinity
AUClast	Area Under the Concentration-Time Curve to the Last Measured Timepoint
AUCext	Percentage of Area Under the Curve Extrapolated Beyond Last Measureable Concentration
AVT	Acute Pulmonary Vasodilator Testing
BA	Bioavailability
BDI	Borg Dyspnea Index
BLQ	Below the Limit of Quantitation
BMPR2	Bone Morphogenic Protein Receptor Type II Gene
BP	British Pharmacopoeia

BTO	Benzidine Triol
CAS	Chemical Abstracts Service
CFR	Code of Federal Regulations
CFU	Colony Forming Unit
cGMP	Current Good Manufacturing Practice
CI	Cardiac Index
CL	Clearance
Cmax	Maximum Concentration
CMC	Chemistry Manufacturing and Controls
CO	Cardiac Output
COPD	Chronic Obstructive Pulmonary Disease
Css	Concentration at Steady State
CTEPH	Chronic Thromboembolic Pulmonary Hypertension
CYP	Cytochrome P450
DMF	Drug Master File
DP	Drug Product
DPI	Dry Powder Inhaler
DPPA	Diphenylphosphinic Acid
DRF	Dose Range Finding
DS	Drug Substance
DUSA	Dosage Unit Sampling Apparatus
eCMH	Extended Cochran-Mantel-Haenszel Test
ECG	Electrocardiogram
ED	Emitted Dose
Emax	Maximum Effect
EP	European Pharmacopoeia
ERA	Endothelin Receptor Agonists
ET3	Polyethylene Terephthalate Cyclic Trimer
EU	Endotoxin Unit
ESRD	End-Stage Renal Disease
F	Bioavailability
FCR	Fluorocur®
FDA	Food and Drug Administration

FPF	Fine Particle Fraction
Frel	Relative Bioavailability
FT-IR	Fourier Transform Infrared Spectroscopy
GC	Gas Chromatography
GLP	Good Laboratory Practice
GMP	Good Manufacturing Practice
GSD	Geometric Standard Deviation
HIAC	High Accuracy Particle Counter
HIV	Human Immunodeficiency Virus
H-L	Hodges-Lehmann
HPLC	High Performance Liquid Chromatography
HPMC	Hydroxypropyl Methylcellulose
HR	Heart Rate
ICH	International Conference on Harmonisation
IM	Intramuscular
IND	Investigational New Drug
INR	International Normalized Ratio
IR	Infrared
ISO	International Organization for Standardization
IV	Intravenous
JP	Japanese Pharmacopoeia
KF	Karl Fischer Titration
LC	Liquid Chromatography
LC-MS	Liquid Chromatography with Mass Spectrometry
LV	Left Ventricular
LVdP/dt	Left Ventricular Contractility
mcg	Micrograms, μg or ug
MDI	Metered Dose Inhaler
MeOH	Methanol
MLWHF	Minnesota Living With Heart Failure Questionnaire
MMAD	Mass Median Aerodynamic Diameter
MTD	Maximum Tolerated Dose
PAPm	Mean Pulmonary Arterial Pressure

NDA	New Drug Application
NF	National Formulary
NGI	Next Generation Impactor™
NMR	Nuclear Magnetic Resonance
NMT	Not More Than
NO	Nitric Oxide
NOAEL	No Observed Adverse Effect Level
NRF	Normal Renal Function
NT	Not Tested
NT-proBNP	N-Terminal of the Prohormone Brain Natriuretic Peptide
NYHA	New York Heart Association
OHSAS	Occupational Health and Safety Advisory Services
OPP	Oriented Polypropylene
PAH	Pulmonary Arterial Hypertension
PAP	Pulmonary Arterial Pressure
PCW	Pulmonary Capillary Wedge pressure
PD	Pharmacodynamics
PDE5	Phosphodiesterase Type 5 Inhibitors
PE	Polyethylene
PET	Polyethylene Terephthalate
PGI_2	Prostaglandin I2 (Prostacyclin)
PH	Pulmonary Hypertension
PK	Pharmacokinetics
PPM	Parts Per Million
PRINT	Particle Replication In Nonwetting Templates
PTFE	Polytetrafluoroethylene
PVR	Pulmonary Vascular Resistance
QID	Quarter in Die (Four Times Daily)
(Q)SAR	Quantitative Structure-Activity Relationship
QTc	Corrected QT Interval
RH	Relative Humidity
RLD	Reference Listed Drug
SAC	Single Actuation Content

SAE	Serious Adverse Event
SAP	Systemic Arterial Pressure
SC	Subcutaneous
SEM	Standard Error of the Mean
SOP	Standard Operating Procedure
SVR	Systemic Vascular Resistance
t½	Half-life
TBD	To Be Determined
TK	Toxicokinetics
Tmax	Time of Maximal Concentration
TMB acid	Trimethylbenzoic Acid
TMB-Ald	Trimethylbenzaldehyde
TMP	Trimethylbenzoyl Diphenylphosphine Oxide
TRIUMPH	Treprostinil Sodium Inhalation Used in the Management of Pulmonary Arterial Hypertension (clinical trial)
TTC	Threshold for Toxicological Concern
IUPAC	International Union of Pure and Applied Chemistry
µg or ug	Micrograms or mcg
US	United States
USP	United States Pharmacopeia
WHO	World Health Organization
WRS	Wilcoxon Rank Sum Test
WT	Weight
XRD	X-ray Diffractometer

Product Nomenclature and Reference Table

Term	Designation	Exemplary Embodiment
Drug Substance	DS or treprostinil	Treprostinil, supplied as treprostinil sodium for manufacture of the LIQ861 drug product-intermediate
Drug Product-Intermediate	DP-intermediate or LIQ861 DP-intermediate	Dry powder particles of precise size and shape containing an integrated matrix of treprostinil and excipients that is produced using Liquidia's PRINT® Technology manufacturing process (bulk dry powder prior to capsule filling)
Placebo Drug Product-Intermediate	Placebo DP-intermediate	Identical formulation as the DP-Intermediate, but treprostinil is replaced with an equal mass of trehalose
Inhalation Powder Drug Product	DP or LIQ861 Drug Product or LIQ861	LIQ861 DP-intermediate filled into Size 3 HPMC capsules for oral inhalation, but prior to integration with Inhalation Device
Placebo Drug Product	Placebo	Placebo DP-intermediate filled into Size 3 hydroxypropyl methylcellulose (HPMC) capsules, but prior to integration with the Inhalation Device
Drug Product Strength or Dose	Treprostinil in LIQ861	Amount of treprostinil in drug product
Packaged Drug Product	None	Drug Product in the Primary Packaging
Inhalation Device	Device	Device that is used to deliver the Drug Product

Premetered Dry Powder Inhaler	DPI	Drug Product integrated with the Inhalation Device; i.e., the final product for patient use

We claim:

1. A dry powder inhalation treatment for pulmonary arterial hypertension, comprising:
a dose of dry particles comprising greater than 25 micrograms of treprostinil enclosed in a capsule.
2. The dry powder inhalation treatment of claim 1, wherein the dose of dry particles comprises greater than or equal to 100 micrograms of treprostinil.
3. The dry powder inhalation treatment of claim 1, wherein the dose of dry particles comprises greater than or equal to 150 micrograms of treprostinil.
4. The dry powder inhalation treatment of claim 1, wherein the dose of dry particles comprises greater than or equal to 5 mg of the dry particles.
5. The dry powder inhalation treatment of claim 2, wherein the dose of dry particles comprises greater than or equal to 10 mg of the dry particles.
6. The dry powder inhalation treatment of claim 3, wherein the dose of dry particles comprises greater than or equal to 15 mg of the dry particles.
7. A dry powder treatment for pulmonary arterial hypertension, comprising:
a single capsule enclosing 5 mg or more dry particles comprising 25 micrograms of treprostinil per each 5 mg of the dry particles.
8. A method of treating a patient having pulmonary arterial hypertension, comprising:
providing a patient a dry powder inhaler;
providing the patient at least one capsule for use in the dry powder inhaler, wherein the capsule comprises at least 25 micrograms of treprostinil; and
instructing the patient to utilize the dry powder inhaler to inhale the treprostinil.
9. The method of claim 8, wherein the capsule comprises at least 50 micrograms of treprostinil.
10. The method of claim 8, wherein the capsule comprises at least 100 micrograms of treprostinil.
11. The method of claim 8, wherein the capsule comprises at least 150 micrograms of treprostinil.
12. A method of treating a patient having pulmonary arterial hypertension, comprising:
dosing the patient having pulmonary arterial hypertension with a dry powder dose of treprostinil, wherein the dose of treprostinil is greater than 85 micrograms.
13. A dry powder inhalation composition for treating pulmonary arterial hypertension, comprising:

a plurality of dry powder particles comprising treprostinil, a non-reducing sugar, a wetting agent, a hydrophobicity modifying agent, a pH modifying agent and a buffer.

14. The dry powder inhalation composition of claim 13, wherein the bulking agent comprises trehalose dihydrate.

15. The dry powder inhalation composition of claim 13, wherein the wetting agent comprises polysorbate 80.

16. The dry powder inhalation composition of claim 13, wherein the hydrophobicity modifying agent comprises L-leucine.

17. The dry powder inhalation composition of claim 13, wherein the pH modifying agent comprises sodium citrate dihydrate.

18. The dry powder inhalation composition of claim 13, wherein the buffer comprises sodium chloride.

19. The dry powder inhalation composition of claim 13, wherein the composition comprises less than about 4 percent by weight water.

20. The dry powder inhalation composition of claim 13, wherein the composition comprises less than about 2 percent by weight water.

21. The dry powder inhalation composition of claim 13, wherein the composition comprises less than about 1 percent by weight water.

22. The dry powder inhalation composition of claim 13, wherein the dry powder particles comprise particles having a three dimensional shape including a width and length not less than 1 micrometer and not more than 2 micrometers and a depth not less than 0.3 micrometers and not more than 0.8 micrometers.

23. The dry powder inhalation composition of claim 13, wherein the dry powder particles comprise a dried solution comprising trehalose dihydrate, L-leucine, treprostinil sodium, polysorbate 80, sodium citrate dihydrate, sodium chloride and water.

24. The dry powder inhalation composition of claim 23, wherein the dry powder particles comprise by percent solids about 0.581 percent treprostinil sodium, about 92.32 percent trehelose, about 2.19 percent polysorbate 80, about 4.39 percent L-leucine, about 0.26 percent sodium citrate, and about 0.25 percent sodium chloride.

25. A method of making a particle for dry powder delivery to the lung of a patient in need thereof, comprising:

molding a composition comprising about 12.30 weight percent trehelose dihydrate, about 0.53 weight percent L-leucine, about 0.07 weight percent treprostinil sodium, about 0.26 weight percent polysorbate 80, about 0.04 weight percent sodium citrate dihydrate, about 0.03 weight percent sodium chloride and about 86.78 weight percent water into a particle; and

drying the composition such that the particle comprises less than 4 percent by weight water.

26. A method of treating a patient having pulmonary arterial hypertension, comprising: delivering greater than 12.5 micrograms of treprostinil to a patient per breath.

27. A method of treating a patient having pulmonary arterial hypertension, comprising: delivering greater than 25 micrograms of treprostinil to a patient per breath.





FIGURE 1

1 micrometer ‘pollen’ particle

FIGURE 2



Example NGI distribution for active particles (PAH-1R-0943-010) made. The beginning of run (A1-red, left bar), middle of run (B1-blue, center bar), end of run (C1-green, right bar). Data obtained using Monodose Model 8 device (95L/min, 2sec).

FIGURE 3A

		Time (h)															
		0.00	**0.08**	**0.17**	**0.25**	**0.33**	**0.42**	**0.50**	**0.75**	**1.00**	**1.50**	**2.00**	**2.50**	**3.00**	**3.50**	**4.00**	**6.00**	**8.00**
Cohort	**Subject Alias**	**Treprostinil (ng/mL)**																
Cohort 1	1-A	0.00	0.164	NR	0.166	0.197	0.150	0.146	0.0992	0.0553	0.0314	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	1-B	0.00	0.243	0.251	0.266	0.326	0.268	0.278	0.136	0.124	0.0688	0.0573	0.0292	0.0328	0.00	0.00	0.00	0.00
	1-C	0.00	0.200	0.272	0.412	0.414	0.251	0.176	0.139	0.108	0.0460	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	1-D	0.00	0.474	0.493	0.458	0.443	0.511	0.383	0.258	0.138	0.0629	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	1-E	0.00	0.200	0.28	0.224	0.255	0.172	0.148	0.114	0.105	0.0519	0.0308	0.00	0.00	0.00	0.00	0.00	0.00
	1-F	0.00	0.402	NR	0.456	0.427	0.313	0.265	0.188	0.106	0.0483	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	N	**6**	**6**	**4**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.00**	**0.281**	**0.324**	**0.330**	**0.344**	**0.278**	**0.233**	**0.156**	**0.106**	**0.0516**	**0.0147**	**0.00487**	**0.00547**	**0.00**	**0.00**	**0.00**	**0.00**
	SD	**0.00**	**0.127**	**0.113**	**0.127**	**0.101**	**0.130**	**0.0934**	**0.0585**	**0.028**	**0.0132**	**0.0242**	**0.0119**	**0.0134**	**0.00**	**0.00**	**0.00**	**0.00**
	Min	**0.00**	**0.164**	**0.251**	**0.166**	**0.197**	**0.150**	**0.146**	**0.0992**	**0.0553**	**0.0314**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Median	**0.00**	**0.222**	**0.276**	**0.339**	**0.370**	**0.260**	**0.221**	**0.138**	**0.107**	**0.0501**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Max	**0.00**	**0.474**	**0.493**	**0.458**	**0.443**	**0.511**	**0.383**	**0.258**	**0.138**	**0.0688**	**0.0573**	**0.0292**	**0.0328**	**0.00**	**0.00**	**0.00**	**0.00**
	CV%	**NC**	**45.1**	**35.0**	**38.6**	**29.5**	**46.7**	**40.1**	**37.6**	**26.4**	**25.6**	**165.1**	**244.9**	**244.9**	**NC**	**NC**	**NC**	**NC**

NC= Not calculated; NR = Not Reported

FIGURE 3B

Cohort	Subject Alias	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 1	1-A	0.197	0.33	0.454	0.146	0.167	150	98.2
	1-B	0.326	0.33	0.795	0.321	0.359	69.7	80.0
	1-C	0.414	0.33	0.460	0.242	0.272	91.8	60.9
	1-D	0.511	0.42	0.377	0.389	0.423	59.1	32.1
	1-E	0.280	0.17	0.648	0.218	0.247	101	94.7
	1-F	0.456	0.25	0.399	0.308	0.336	74.4	42.8
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.364**	**0.306**	**0.522**	**0.271**	**0.301**	**91.0**	**68.1**
	SD	**0.117**	**0.086**	**0.164**	**0.086**	**0.091**	**32.6**	**27.4**
	Min	**0.20**	**0.17**	**0.38**	**0.15**	**0.17**	**59.1**	**32.1**
	Median	**0.37**	**0.33**	**0.46**	**0.28**	**0.30**	**83.1**	**70.4**
	Max	**0.51**	**0.42**	**0.80**	**0.39**	**0.42**	**150**	**98.2**
	CV%	**32.3**	**28.2**	**31.5**	**31.8**	**30.2**	**35.8**	**40.2**
	Geometric Mean	**0.347**	**0.294**	**0.503**	**0.258**	**0.288**	**86.8**	**63.0**
	CV% Geometric Mean	**36.52**	**33.02**	**29.96**	**35.77**	**33.91**	**33.9**	**47.67**

FIGURE 4A

		Time (h)																
		0.00	**0.08**	**0.17**	**0.25**	**0.33**	**0.42**	**0.50**	**0.75**	**1.00**	**1.50**	**2.00**	**2.50**	**3.00**	**3.50**	**4.00**	**6.00**	**8.00**
Cohort	**Subject Alias**	**Treprostinil (ng/mL)**																
Cohort 2	2-A	0.00	NR	0.99	0.93	0.77	0.70	0.58	0.37	0.30	0.12	0.08	0.03	0.00	0.00	0.00	0.00	0.00
	2-B	0.00	0.94	1.03	0.95	0.83	0.64	0.41	0.30	0.18	0.09	0.05	0.03	0.03	0.00	0.00	0.00	0.00
	2-C	0.00	0.48	0.41	0.32	0.31	0.26	0.22	0.13	0.08	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	2-D	0.00	0.45	0.53	0.50	0.46	0.45	0.44	0.30	0.22	0.08	0.04	0.00	0.00	0.00	0.00	0.00	0.00
	2-E	0.00	0.28	0.29	0.26	0.23	0.21	0.20	0.13	0.08	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	2-F	0.00	0.00	0.09	0.11	0.11	0.12	0.11	0.06	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	N	**6**	**5**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.00**	**0.429**	**0.556**	**0.511**	**0.452**	**0.395**	**0.326**	**0.214**	**0.149**	**0.0624**	**0.0277**	**0.0108**	**0.00507**	**0.00**	**0.00**	**0.00**	**0.00**
	SD	**0.00**	**0.342**	**0.382**	**0.355**	**0.295**	**0.240**	**0.180**	**0.123**	**0.101**	**0.0433**	**0.0333**	**0.0168**	**0.0124**	**0.00**	**0.00**	**0.00**	**0.00**
	Min	**0.00**	**0.00**	**0.0853**	**0.111**	**0.108**	**0.116**	**0.110**	**0.0600**	**0.0270**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Median	**0.00**	**0.453**	**0.470**	**0.410**	**0.385**	**0.351**	**0.313**	**0.216**	**0.134**	**0.0603**	**0.0186**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Max	**0.00**	**0.937**	**1.03**	**0.948**	**0.833**	**0.703**	**0.583**	**0.367**	**0.298**	**0.124**	**0.0794**	**0.0332**	**0.0304**	**0.00**	**0.00**	**0.00**	**0.00**
	CV%	**NC**	**79.7**	**68.8**	**69.4**	**65.1**	**60.8**	**55.2**	**57.6**	**68.1**	**69.3**	**120.3**	**155.0**	**244.9**	**NC**	**NC**	**NC**	**NC**

NC= Not calculated; NR = Not Reported

WO 2017/192993 PCT/US2017/031301

CA 03023257 2018-11-02

FIGURE 4B

Cohort	Subject	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 2	2-A	0.991	0.17	0.482	0.723	0.746	67.0	46.6
	2-B	1.03	0.17	0.624	0.663	0.690	72.5	65.2
	2-C	0.477	0.08	0.474	0.254	0.282	177	121
	2-D	0.530	0.17	0.410	0.466	0.488	102	60.5
	2-E	0.288	0.17	0.478	0.212	0.242	206	142
	2-F	0.116	0.42	0.247	0.0706	0.0802	624	222
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.572**	**0.195**	**0.452**	**0.398**	**0.422**	**208**	**110**
	SD	**0.370**	**0.114**	**0.123**	**0.262**	**0.265**	**211**	**66.6**
	Min	**0.116**	**0.0830**	**0.247**	**0.0706**	**0.0802**	**67.0**	**46.6**
	Median	**0.504**	**0.167**	**0.476**	**0.360**	**0.385**	**140**	**93.3**
	Max	**1.03**	**0.417**	**0.624**	**0.723**	**0.746**	**624**	**222**
	CV%	**64.7**	**58.6**	**27.2**	**65.8**	**62.8**	**101.5**	**60.7**
	Geometric Mean	**0.453**	**0.173**	**0.436**	**0.308**	**0.334**	**150**	**94.3**
	CV% Geometric Mean	**98.2**	**54.9**	**31.8**	**107.4**	**100.4**	**100.4**	**65.7**

FIGURE 5A

		NTime (h)															
		0.00	**0.08**	**0.17**	**0.25**	**0.33**	**0.42**	**0.50**	**0.75**	**1.00**	**1.50**	**2.00**	**2.50**	**3.00**	**3.50**	**4.00**	**6.00**	**8.00**
Cohort	**Subject Alias**	**Treprostinil (ng/mL)**																
Cohort 3	3-A	0.00	0.63	0.62	0.55	0.47	0.43	0.42	0.32	0.21	0.11	0.06	0.03	0.00	0.00	0.00	0.00	0.00
	3-B	0.00	0.44	0.52	0.51	0.51	0.53	0.47	0.39	0.26	0.13	0.12	0.03	0.00	0.00	0.00	0.00	0.00
	3-C	0.00	0.75	1.07	1.11	1.01	NR	0.80	0.47	0.35	0.14	0.07	0.05	0.03	0.00	0.00	0.00	0.00
	3-D	0.00	NR	0.40	0.43	0.40	0.37	0.34	0.29	0.20	0.10	0.05	0.00	0.00	0.00	0.00	0.00	0.00
	3-E	0.00	0.45	0.52	0.54	0.57	0.54	0.48	0.36	0.28	0.15	0.11	0.03	0.00	0.00	0.00	0.00	0.00
	3-F	0.00	0.94	1.05	1.10	1.10	1.07	1.03	0.80	0.53	0.30	0.15	0.07	0.04	0.00	0.00	0.00	0.00
	N	**6**	**5**	**6**	**6**	**6**	**5**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.00**	**0.641**	**0.697**	**0.706**	**0.677**	**0.587**	**0.591**	**0.437**	**0.304**	**0.155**	**0.0951**	**0.0349**	**0.0124**	**0.00**	**0.00**	**0.00**	**0.00**
	SD	**0.00**	**0.209**	**0.290**	**0.312**	**0.299**	**0.279**	**0.267**	**0.187**	**0.123**	**0.0755**	**0.0389**	**0.0225**	**0.0192**	**0.00**	**0.00**	**0.00**	**0.00**
	Min	**0.00**	**0.437**	**0.399**	**0.428**	**0.400**	**0.371**	**0.344**	**0.285**	**0.199**	**0.0976**	**0.0547**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Median	**0.00**	**0.630**	**0.571**	**0.545**	**0.540**	**0.532**	**0.474**	**0.379**	**0.269**	**0.136**	**0.0916**	**0.0342**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Max	**0.00**	**0.936**	**1.07**	**1.11**	**1.10**	**1.07**	**1.03**	**0.796**	**0.529**	**0.304**	**0.149**	**0.0682**	**0.0395**	**0.00**	**0.00**	**0.00**	**0.00**
	CV%	**NC**	**32.6**	**41.7**	**44.2**	**44.2**	**47.5**	**45.2**	**42.8**	**40.5**	**48.8**	**40.9**	**64.6**	**155.4**	**NC**	**NC**	**NC**	**NC**

NC= Not calculated; NR = Not Reported

FIGURE 5B

Cohort	Subject Alias	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 3	3-A	0.630	0.08	0.609	0.538	0.568	132	116
	3-B	0.532	0.42	0.568	0.611	0.638	118	96.3
	3-C	1.11	0.25	0.980	0.906	0.955	78.5	111
	3-D	0.428	0.25	0.522	0.411	0.452	166	125
	3-E	0.567	0.33	0.531	0.619	0.639	117	89.9
	3-F	1.10	0.25	0.520	1.26	1.29	58.1	43.6
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.728**	**0.264**	**0.622**	**0.724**	**0.757**	**112**	**97.0**
	SD	**0.299**	**0.111**	**0.179**	**0.309**	**0.310**	**38.5**	**29.1**
	Min	**0.428**	**0.0833**	**0.520**	**0.411**	**0.452**	**58.1**	**43.6**
	Median	**0.599**	**0.250**	**0.550**	**0.615**	**0.639**	**117**	**104**
	Max	**1.11**	**0.417**	**0.980**	**1.26**	**1.29**	**166**	**125**
	CV%	**41.1**	**42.0**	**28.7**	**42.7**	**41.0**	**34.5**	**30.0**
	Geometric Mean	**0.681**	**0.238**	**0.605**	**0.676**	**0.711**	**106**	**92.1**
	CV% Geometric Mean	**41.2**	**60.0**	**24.8**	**41.5**	**39.4**	**39.4**	**40.0**

FIGURE 6A

		NTime (h)															
		0.00	**0.08**	**0.17**	**0.25**	**0.33**	**0.42**	**0.50**	**0.75**	**1.00**	**1.50**	**2.00**	**2.50**	**3.00**	**3.50**	**4.00**	**6.00**	**8.00**
Cohort	**Subject**	**Treprostinil (ng/mL)**																
Cohort 4	4-A	0.00	0.294	0.534	0.648	0.684	0.722	0.763	0.575	0.358	0.150	0.0789	0.0523	0.0332	0.00	0.00	0.00	0.00
	4-B	0.00	0.629	0.872	0.959	1.04	1.02	0.924	0.622	0.495	0.273	0.144	0.0710	0.0449	0.0300	0.00	0.00	0.00
	4-C	0.00	0.552	0.725	0.704	0.718	0.688	0.729	0.625	0.557	0.346	0.205	0.143	0.0908	0.0445	0.0286	0.00	0.00
	4-D	0.00	1.30	1.54	1.44	1.35	1.29	1.08	0.790	0.559	0.290	0.134	0.0709	0.0728	0.0333	0.00	0.00	0.00
	4-E	0.00	0.877	1.14	1.16	1.14	0.969	0.970	0.657	0.513	0.270	0.166	0.0968	0.0524	0.0367	0.0256	0.00	0.00
	4-F	0.00	1.13	1.21	1.23	1.09	0.915	0.826	0.662	0.521	0.253	0.125	0.140	0.0698	0.0324	0.0837	0.00	0.00
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.00**	**0.797**	**1.00**	**1.02**	**1.00**	**0.934**	**0.882**	**0.655**	**0.501**	**0.264**	**0.142**	**0.0957**	**0.0607**	**0.0295**	**0.0230**	**0.00**	**0.00**
	SD	**0.00**	**0.377**	**0.365**	**0.310**	**0.257**	**0.220**	**0.134**	**0.0730**	**0.0742**	**0.0642**	**0.0422**	**0.0382**	**0.0210**	**0.0153**	**0.0326**	**0.00**	**0.00**
	Min	**0.00**	**0.294**	**0.534**	**0.648**	**0.684**	**0.688**	**0.729**	**0.575**	**0.358**	**0.150**	**0.0789**	**0.0523**	**0.0332**	**0.00**	**0.00**	**0.00**	**0.00**
	Median	**0.00**	**0.753**	**1.01**	**1.06**	**1.07**	**0.942**	**0.875**	**0.641**	**0.517**	**0.272**	**0.139**	**0.0839**	**0.0611**	**0.0329**	**0.0128**	**0.00**	**0.00**
	Max	**0.00**	**1.30**	**1.54**	**1.44**	**1.35**	**1.29**	**1.08**	**0.790**	**0.559**	**0.346**	**0.205**	**0.143**	**0.0908**	**0.0445**	**0.0837**	**0.00**	**0.00**
	CV%	**NC**	**47.3**	**36.3**	**30.3**	**25.6**	**23.5**	**15.2**	**11.1**	**14.8**	**24.4**	**29.7**	**40.0**	**34.7**	**51.9**	**141.8**	**NC**	**NC**

NC= Not calculated

FIGURE 6B

Cohort	Subject	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 4	4-A	0.763	0.50	0.801	0.781	0.819	122	141
	4-B	1.04	0.33	0.805	1.13	1.17	85.8	99.6
	4-C	0.729	0.50	0.702	1.18	1.21	82.8	83.8
	4-D	1.54	0.17	0.583	1.44	1.46	68.3	57.4
	4-E	1.16	0.25	0.968	1.26	1.30	77.1	108
	4-F	1.23	0.25	0.794	1.28	1.37	72.8	83.4
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**1.08**	**0.333**	**0.775**	**1.18**	**1.22**	**84.8**	**95.5**
	SD	**0.305**	**0.139**	**0.127**	**0.221**	**0.225**	**19.3**	**28.2**
	Min	**0.729**	**0.167**	**0.583**	**0.781**	**0.819**	**68.3**	**57.4**
	Median	**1.10**	**0.292**	**0.798**	**1.22**	**1.25**	**79.9**	**91.7**
	Max	**1.54**	**0.500**	**0.968**	**1.44**	**1.46**	**122**	**141**
	CV%	**28.3**	**41.8**	**16.4**	**18.7**	**18.4**	**22.8**	**29.5**
	Geometric Mean	**1.04**	**0.309**	**0.766**	**1.16**	**1.20**	**83.2**	**92.0**
	CV% Geometric Mean	**29.5**	**45.5**	**17.0**	**21.2**	**20.7**	**20.7**	**30.8**

FIGURE 7A

		NTime (h)																
		0.00	**0.08**	**0.17**	**0.25**	**0.33**	**0.42**	**0.50**	**0.75**	**1.00**	**1.50**	**2.00**	**2.50**	**3.00**	**3.50**	**4.00**	**6.00**	**8.00**
Cohort	**Subject**	**Treprostinil (ng/mL)**																
Cohort 5	5-A	0.00	1.13	1.96	1.80	1.64	1.54	1.38	0.971	0.752	0.394	0.238	0.136	0.0833	0.0583	0.0345	0.00	0.00
	5-B	0.00	0.562	1.26	1.26	1.45	1.40	1.24	0.876	0.605	0.312	0.195	0.0732	0.0351	0.00	0.00	0.00	0.00
	5-C	0.00	0.478	0.763	0.979	1.02	0.976	0.909	NR	0.409	0.294	0.128	0.0658	0.0433	0.00	0.00	0.00	0.00
	5-D	0.00	1.04	1.42	1.26	1.08	1.02	0.824	0.635	0.467	0.252	0.146	0.0615	0.0304	0.00	0.00	0.00	0.00
	5-E	0.00	0.543	0.728	0.730	0.743	0.754	0.699	0.502	0.397	0.230	0.113	0.0500	0.0262	0.00	0.00	0.00	0.00
	5-F	0.00	0.502	0.510	0.427	0.399	0.333	0.265	0.201	0.120	0.0565	0.0306	0.00	0.00	0.00	0.00	0.00	0.00
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**5**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.00**	**0.709**	**1.11**	**1.08**	**1.06**	**1.00**	**0.886**	**0.637**	**0.458**	**0.256**	**0.142**	**0.0644**	**0.0364**	**0.00972**	**0.00575**	**0.00**	**0.00**
	SD	**0.00**	**0.294**	**0.542**	**0.478**	**0.453**	**0.438**	**0.399**	**0.307**	**0.214**	**0.113**	**0.0714**	**0.0437**	**0.0273**	**0.0238**	**0.0141**	**0.00**	**0.00**
	Min	**0.00**	**0.478**	**0.510**	**0.427**	**0.399**	**0.333**	**0.265**	**0.201**	**0.120**	**0.0565**	**0.0306**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Median	**0.00**	**0.553**	**1.01**	**1.12**	**1.05**	**0.998**	**0.867**	**0.635**	**0.438**	**0.273**	**0.137**	**0.0637**	**0.0328**	**0.00**	**0.00**	**0.00**	**0.00**
	Max	**0.00**	**1.13**	**1.96**	**1.80**	**1.64**	**1.54**	**1.38**	**0.971**	**0.752**	**0.394**	**0.238**	**0.136**	**0.0833**	**0.0583**	**0.0345**	**0.00**	**0.00**
	CV%	**NC**	**41.5**	**48.9**	**44.4**	**43.0**	**43.6**	**45.0**	**48.2**	**46.6**	**44.2**	**50.4**	**67.9**	**74.9**	**244.9**	**244.9**	**NC**	**NC**

NC= Not calculated; NR = Not Reported

WO 2017/192993 PCT/US2017/031301
CA 030232257 2018-11-02
11/22

FIGURE 7B

Cohort	Subject	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 5	5-A	1.96	0.17	0.647	1.87	1.90	65.7	61.3
	5-B	1.45	0.33	0.495	1.42	1.45	86.2	61.6
	5-C	1.02	0.33	0.561	1.05	1.08	115	93.3
	5-D	1.42	0.17	0.526	1.18	1.20	104	78.9
	5-E	0.754	0.42	0.473	0.874	0.891	140	95.6
	5-F	0.510	0.17	0.448	0.352	0.372	336	217
N		**6**	**6**	**6**	**6**	**6**	**6**	**6**
Mean		**1.19**	**0.264**	**0.525**	**1.12**	**1.15**	**141**	**101**
SD		**0.528**	**0.111**	**0.0716**	**0.512**	**0.516**	**98.8**	**58.7**
Min		**0.510**	**0.167**	**0.448**	**0.352**	**0.372**	**65.7**	**61.3**
Median		**1.22**	**0.250**	**0.511**	**1.11**	**1.14**	**110**	**86.1**
Max		**1.96**	**0.417**	**0.647**	**1.87**	**1.90**	**336**	**217**
CV%		**44.5**	**42.0**	**13.6**	**45.6**	**44.9**	**69.9**	**57.9**
Geometric Mean		**1.08**	**0.245**	**0.521**	**1.00**	**1.03**	**121**	**91.3**
CV% Geometric Mean		**52.4**	**44.9**	**13.2**	**62.6**	**60.9**	**60.9**	**49.3**

FIGURE 8A

		NTime (h)															
		0.00	**0.08**	**0.17**	**0.25**	**0.33**	**0.42**	**0.50**	**0.75**	**1.00**	**1.50**	**2.00**	**2.50**	**3.00**	**3.50**	**4.00**	**6.00**	**8.00**
Cohort	**Subject**	**Treprostinil (ng/mL)**																
Cohort 6-R	6-A	0.00	0.244	0.474	0.596	0.622	0.704	0.61	0.459	0.345	0.177	0.085	0.0387	0.00	0.00	0.00	0.00	0.00
	6-B	0.00	0.877	0.907	0.75	0.657	0.538	0.459	0.287	0.166	0.0915	0.0488	0.00	0.00	0.00	0.00	0.00	0.00
	6-C	0.00	0.547	1.47	1.68	1.67	1.59	1.41	1.00	0.762	0.406	0.232	0.144	0.0885	0.0603	0.0376	0.00	0.00
	6-D	0.00	1.41	1.7	1.89	1.86	1.75	1.69	1.24	0.9	0.463	0.22	0.116	0.0673	0.0758	0.0404	0.00	0.00
	6-E	0.00	1.21	1.83	2.12	2.34	2.06	1.9	1.65	1.25	0.699	0.431	0.244	0.14	0.106	0.093	0.00	0.00
	6-F	0.00	0.615	0.938	1.09	1.19	1.12	1.11	0.912	0.682	0.412	0.207	0.122	0.0677	0.0502	0.0325	0.00	0.00
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**0.00**	**0.817**	**1.22**	**1.35**	**1.39**	**1.29**	**1.2**	**0.925**	**0.684**	**0.375**	**0.204**	**0.111**	**0.0606**	**0.0487**	**0.0339**	**0.00**	**0.00**
	SD	**0.00**	**0.436**	**0.529**	**0.631**	**0.688**	**0.605**	**0.58**	**0.501**	**0.389**	**0.217**	**0.135**	**0.0854**	**0.0539**	**0.0422**	**0.0342**	**0.00**	**0.00**
	Min	**0.00**	**0.244**	**0.474**	**0.596**	**0.622**	**0.538**	**0.459**	**0.287**	**0.166**	**0.0915**	**0.0488**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**	**0.00**
	Median	**0.00**	**0.746**	**1.20**	**1.39**	**1.43**	**1.36**	**1.26**	**0.956**	**0.722**	**0.409**	**0.214**	**0.119**	**0.0675**	**0.0553**	**0.0351**	**0.00**	**0.00**
	Max	**0.00**	**1.41**	**1.83**	**2.12**	**2.34**	**2.06**	**1.90**	**1.65**	**1.25**	**0.699**	**0.431**	**0.244**	**0.14**	**0.106**	**0.093**	**0.00**	**0.00**
	CV%	**NC**	**53.4**	**43.4**	**46.6**	**49.5**	**46.8**	**48.4**	**54.2**	**56.8**	**57.8**	**66.1**	**77.1**	**89**	**86.6**	**100.8**	**NC**	**NC**

NC= Not calculated; NR = Not Reported

FIGURE 8B

Cohort	Subject	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 6-R	6-A	0.704	0.42	0.456	0.696	0.721	208	137
	6-B	0.907	0.17	0.566	0.573	0.613	245	200
	6-C	1.68	0.25	0.768	1.80	1.85	81.3	90.1
	6-E	1.89	0.25	0.628	2.10	2.13	70.3	63.6
	6-F	2.34	0.33	0.734	2.80	2.90	51.8	54.9
	6-G	1.19	0.33	0.665	1.50	1.53	97.9	93.8
N		**6**	**6**	**6**	**6**	**6**	**6**	**6**
Mean		**1.45**	**0.292**	**0.636**	**1.58**	**1.62**	**126**	**107**
SD		**0.626**	**0.087**	**0.114**	**0.849**	**0.869**	**80.3**	**54.0**
Min		**0.704**	**0.167**	**0.456**	**0.573**	**0.613**	**51.8**	**54.9**
Median		**1.44**	**0.292**	**0.646**	**1.65**	**1.69**	**89.6**	**92.0**
Max		**2.34**	**0.417**	**0.768**	**2.8**	**2.90**	**245**	**200**
CV%		**43.1**	**30.0**	**18.0**	**53.8**	**53.5**	**63.9**	**50.7**
Geometric Mean		**1.33**	**0.28**	**0.627**	**1.36**	**1.41**	**107**	**96.5**
CV% Geometric Mean		**48.6**	**32.9**	**19.2**	**70**	**68.3**	**68.3**	**50.9**

FIGURE 8C

Cohort	Subject	Cmax (ng/mL)	Tmax (h)	t½ (h)	AUClast (h*ng/mL)	AUCinf (h*ng/mL)	CL/F (L/h)	Vz/F (L)
Cohort 6 - Original	6-A (ORIGINAL)	1.55	0.33	0.607	1.96	1.99	75.6	66.2
	6-B (ORIGINAL)	1.16	0.08	0.946	0.868	0.987	152	208
	6-C (ORIGINAL)	1.17	0.42	0.534	1.28	1.30	115	88.9
	6-D (ORIGINAL)	0.968	0.42	0.649	1.30	1.33	113	105
	6-E (ORIGINAL)	1.55	0.17	0.658	1.68	1.71	87.9	83.5
	6-F (ORIGINAL)	0.835	0.25	0.565	0.871	0.891	168	137
	N	**6**	**6**	**6**	**6**	**6**	**6**	**6**
	Mean	**1.21**	**0.278**	**0.660**	**1.33**	**1.37**	**119**	**115**
	SD	**0.295**	**0.136**	**0.148**	**0.435**	**0.418**	**35.8**	**51.4**
	Min	**0.835**	**0.0833**	**0.534**	**0.868**	**0.891**	**75.6**	**66.2**
	Median	**1.17**	**0.292**	**0.628**	**1.29**	**1.32**	**114**	**97.2**
	Max	**1.55**	**0.417**	**0.946**	**1.96**	**1.99**	**168**	**208**
	CV%	**24.4**	**49.0**	**22.5**	**32.8**	**30.6**	**30.2**	**44.8**
	Geometric Mean	**1.18**	**0.242**	**0.648**	**1.27**	**1.31**	**114**	**107**
	CV% Geometric Mean	**25.2**	**69.5**	**20.4**	**34.2**	**31.4**	**31.4**	**42.4**

FIGURE 8D



Cohort 1
Cohort 2
Cohort 3
Cohort 4
Cohort 5
Cohort 6
kTmax (h)
0
0.5
1
1.5
2
2.5
3
3.5
Mean (ng/mL)
0
0.2
0.4
0.6
0.8
1
1.2
1.4
1.6

FIGURE 8E



Rsq = 0.4785, Intercept = 0.185, Slope = 0.007835
Cmax (ng/mL)
0
2.5
5
7.5
10
12.5
15
0
25
50
75
100
125
150
Dose (ug)

FIGURE 8F



Rsq = 0.5172, Intercept = 0.03113, Slope = 0.009813
AUCinf (h*ng/mL)
Dose (ug)
0
25
50
75
100
125
150
0
1
2
3
4
5

FIGURE 8G



Rsq = 0.001393, Intercept = 133.5, Slope = -0.07923
800
600
400
200
0
CL/F (L/h)
0
25
50
75
100
125
150
Dose (ug)

FIGURE 9



TESCAN
VEGA3 TE
WD: 10.03 mm
View field: 261 um
sem_17_001
SEM MAG
Detector

FIGURE 10





Solvents
Excipients
API
Stock Solution Prep
PARTICLE FABRICATION
Coating
Mold Filling
Drying
Dry Collection
Bulk Packaging
Mold/ Support Films

FIGURE 11



RS00 Model 8 Dry Powder Inhalation Device

FIGURE 1



1 micrometer ‘pollen’ particle