

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Promoter molecules isolated from Brassica napus for use in plants

(51)⁶ International Patent Classification(s)
C12N 15/82 (2006.01) 20060101AFI2006040
C12N 15/82 6BHEP
PCT/US2005/034150

(21) Application No: 2005289727 (22) Application Date: 2005 .09 .23

(87) WIPO No: W006/036787

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/613,081	2004 .09 .24	US

(43) Publication Date : 2006 .04 .06

(71) Applicant(s)
Monsanto Technology LLC

(72) Inventor(s)
Hewitt, Crystal L.; Ruezinsky, Diane M.; Zheng, Wei

(74) Agent/Attorney
EF WELLINGTON & CO, 312 St Kilda Road Southbank, MELBOURNE, VIC, 3006

(56) Related Art
SANDER et al. Plant Mol. Biol., 2001 vol. 46 pages 469-479
Database EMBL entry asseccion no. BH933900

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 April 2006 (06.04.2006)

PCT

(10) International Publication Number
WO 2006/036787 A3

- (51) International Patent Classification:
C12N 15/82 (2006.01)
- (21) International Application Number:
PCT/US2005/034150
- (22) International Filing Date:
23 September 2005 (23.09.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/613,081 24 September 2004 (24.09.2004) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/613,081 (CIP)
Filed on 24 September 2004 (24.09.2004)
- (71) Applicant (for all designated States except US): MON-SANTO TECHNOLOGY LLC [US/US]; 800 N. Lindbergh Blvd., Saint Louis, MO 63167 (US).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): RUEZINSKY, Diane, M. [US/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). HEWITT, Crystal, L. [US/US]; 800 N. Lindbergh Blvd., St. Louis, MO 63167 (US). ZHENG, Wei [CN/US]; 800 N. Lindbergh Blvd., St. Louis, MO 63167 (US).
- (74) Agent: HANSON, Robert, E.; Fulbright & Jaworski, L.L.P., Suite 2400, 600 Congress Avenue, Austin, TX 78701 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SG, SH, SI, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US (patent), UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
31 August 2006
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2006/036787 A3

(54) Title: PROMOTER MOLECULES ISOLATED FROM BRASSICA NAPUS FOR USE IN PLANTS

(57) Abstract: The present invention relates to polynucleotide molecules for regulating gene expression in plants. In particular, the present invention relates to promoters isolated from Brassica napus that are useful for regulating gene expression of heterologous polynucleotide molecules in plants. The present invention also relates to expression constructs and transgenic plants containing the heterologous polynucleotide molecules.

DESCRIPTION**PROMOTER MOLECULES FOR USE IN PLANTS****BACKGROUND OF THE INVENTION**

This application claims the priority of provisional application Serial No. 5 60/613,081, filed September 24, 2004, the disclosure of which is incorporated herein by reference in its entirety.

1. Field of the Invention

The present invention relates to the field of plant molecular biology and plant genetic engineering and more specifically relates to polynucleotide molecules useful 10 for the expression of transgenes in plants.

2. Description of Related Art

One of the goals of plant genetic engineering is to produce plants with agronomically desirable characteristics or traits. Advances in genetic engineering have provided the requisite tools to transform plants to contain and express foreign 15 genes. The technological advances in plant transformation and regeneration have enabled researchers to take an exogenous polynucleotide molecule, such as a gene from a heterologous or native source, and incorporate that polynucleotide molecule into a plant genome. The gene can then be expressed in a plant cell to exhibit the added characteristic or trait. In one approach, expression of a gene in a plant cell or a 20 plant tissue that does not normally express such a gene may confer a desirable phenotypic effect. In another approach, transcription of a gene or part of a gene in an antisense orientation may produce a desirable effect by preventing or inhibiting expression of an endogenous gene.

For production of transgenic plants with various desired characteristics, it 25 would be advantageous to have a variety of promoters to provide gene expression such that a gene is transcribed efficiently in the amount necessary to produce the desired effect. The commercial development of genetically improved germplasm has also advanced to the stage of introducing multiple traits into crop plants, often referred to as a gene stacking approach. In this approach, multiple genes conferring different 30 characteristics of interest can be introduced into a plant. It is often desired when

introducing multiple genes into a plant that each gene is modulated or controlled for optimal expression, leading to a requirement for diverse regulatory elements. In light of these and other considerations, it is apparent that optimal control of gene expression and regulatory element diversity are important in plant biotechnology.

5 Numerous promoters, which are active in plant cells, have been described in the literature. These include the nopaline synthase (nos) promoter and octopine synthase (ocs) promoters carried on tumor-inducing plasmids of *Agrobacterium tumefaciens* and the caulimovirus promoters such as the Cauliflower Mosaic Virus (CaMV) 19S or 35S promoter (U.S. Patent 5,352,605), CaMV 35S promoter with a
10 duplicated enhancer (U.S. Patents 5,164,316; 5,196,525; 5,322,938; 5,359,142; and 5,424,200), and the Figwort Mosaic Virus (FMV) 35S promoter (U.S. Patent 5,378,619). These promoters and numerous others have been used in the creation of constructs for transgene expression in plants. Other useful promoters are described, for example, in U.S. Patents 5,391,725; 5,428,147; 5,447,858; 5,608,144; 5,614,399;
15 5,633,441; 6,232,526; and 5,633,435, all of which are incorporated herein by reference.

While previous work has provided a number of promoters useful to direct transcription in transgenic plants, there is still a need for novel promoters with beneficial expression characteristics. In particular, there is a need for promoters that
20 are capable of directing expression of exogenous genes in dicotyledonous seeds. Many previously identified promoters fail to provide the patterns or levels of expression required to fully realize the benefits of expression of selected genes in transgenic plants. There is, therefore, a need in the art of plant genetic engineering for novel promoters for use in dicots.

25 **Summary of the Invention**

The present invention relates to promoters that are described as silique wall preferred promoters. The phrase "silique wall preferred" refers to promoters that drive expression of operably linked genes to higher levels in the silique wall compared with any other tissue tested. Silique wall preferred promoters are useful for production of
30 transgenic plants with desired seed traits. These include, but are not limited to, altering oil content, protein quality, cell proliferation, or micronutrient quality.

In one embodiment the present invention provides a promoter comprising a polynucleotide sequence substantially homologous to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 7, 8, and 9, or any fragments and variants thereof that are capable of regulating transcription of operably linked polynucleotide molecules, *e.g.*, having promoter activity. In particular embodiments, a fragment of a sequence provided herein is defined as comprising at least about 30, 40, 50, 75, 100, 125, 150, 200, 250, 300, 350, 400, 450, 500, 600, 750, 900, 1000, or more contiguous nucleotides of any of the promoter sequences described herein, including, for example, SEQ ID NO: SEQ ID NOs: 7, 8, and 9.

In another embodiment, the present invention provides a plant expression construct comprising a promoter comprising a polynucleotide sequence substantially homologous to a polynucleotide sequence described herein, for example, selected from the group consisting of SEQ ID NOs: 7, 8, and 9, or any fragments or variants thereof, wherein said promoter is operably linked to a transcribable polynucleotide molecule.

In yet another embodiment, the present invention provides a transgenic plant stably transformed with a plant expression construct comprising a promoter provided by the invention. In one embodiment, the construct comprises a polynucleotide sequence substantially homologous to a polynucleotide sequence selected from the group consisting of SEQ ID NOs: 7, 8, and 9, or any fragments, variants, or regions thereof, wherein said promoter is operably linked to a transcribable polynucleotide molecule.

In another embodiment, the present invention provides a method of making a vegetable oil, comprising the steps of incorporating into the genome of a plant a promoter of the present invention operably linked to a transcribable polynucleotide molecule conferring altered oil and/or protein content, growing the plant to produce seeds, and extracting the oil and/or protein from the seed. In yet another embodiment, the invention provides a method for making food or feed comprising obtaining a plant of the invention and preparing the food or feed from the plant or a part thereof.

The foregoing and other aspects of the present invention will become more apparent from the following detailed description.

Brief Description of the Sequences

- SEQ ID NO: 1 sets forth the polynucleotide sequence for the P-BN.RPC-0:1:1 promoter.
- SEQ ID NO: 2 sets forth the polynucleotide sequence for the primer GSP1.
- 5 SEQ ID NO: 3 sets forth the polynucleotide sequence for the primer AP1.
- SEQ ID NO: 4 sets forth the polynucleotide sequence for the primer GSP2.
- SEQ ID NO: 5 sets forth the polynucleotide sequence for the primer AP2.
- SEQ ID NO: 6 sets forth the polynucleotide sequence for the primer D11 nco.
- SEQ ID NO: 7 sets forth the polynucleotide sequence for the P-BN.SW1-0:1:2 promoter.
- 10 SEQ ID NO: 8 sets forth the polynucleotide sequence for the P-BN.SW2-0:1:2 promoter.
- SEQ ID NO: 9 sets forth the polynucleotide sequence for the P-BN.SW3-0:1:2 promoter.
- 15 SEQ ID NO: 10 sets forth the polynucleotide sequence for the primer #17637.
- SEQ ID NO: 11 sets forth the polynucleotide sequence for the primer #17636.
- SEQ ID NO: 12 sets forth the polynucleotide sequence for the primer #17669.
- SEQ ID NO: 13 sets forth the polynucleotide sequence for the primer #17668.
- SEQ ID NO: 14 sets forth the polynucleotide sequence for the primer #18210.
- 20 SEQ ID NO: 15 sets forth the polynucleotide sequence for the primer #18209.
- SEQ ID NO: 16 sets forth the polynucleotide sequence for the primer #18115.
- SEQ ID NO: 17 sets forth the polynucleotide sequence for the primer #18167.
- SEQ ID NO: 18 sets forth the polynucleotide sequence for the primer #18206.
- SEQ ID NO: 19 sets forth the polynucleotide sequence for the primer #18340.
- 25 SEQ ID NO: 20 sets forth the polynucleotide sequence for the primer #19944.
- SEQ ID NO: 21 sets forth the polynucleotide sequence for the primer #19945.
- SEQ ID NO: 22 sets forth the polynucleotide sequence of the 891 bp fragment containing P-BN.SW3-0:1:2.

30

Detailed Description of the Invention

The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the

present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

As used herein, the phrase "polynucleotide molecule" refers to the single- or double-stranded DNA or RNA of genomic or synthetic origin, *i.e.*, a polymer of deoxyribonucleotide or ribonucleotide bases, respectively, read from the 5' (upstream) end to the 3' (downstream) end.

As used herein, the phrase "polynucleotide sequence" refers to the sequence of a polynucleotide molecule. The nomenclature for DNA bases as set forth at 37 CFR § 1.822 is used.

10 **Promoters**

As used herein, the term "promoter" refers to a polynucleotide molecule that in its native state is located upstream or 5' to a translational start codon of an open reading frame (or protein-coding region) and that is involved in recognition and binding of RNA polymerase II and other proteins (trans-acting transcription factors) to initiate transcription. A "plant promoter" is a native or non-native promoter that is functional in plant cells. Constitutive plant promoters are functional in most or all tissues of a plant throughout plant development. Any plant promoter can be used as a 5' regulatory element for modulating expression of a particular gene or genes operably associated thereto. When operably linked to a transcribable polynucleotide molecule, a promoter typically causes the transcribable polynucleotide molecule to be transcribed in a manner that is similar to that of which the promoter is normally associated. Plant promoters can include promoters produced through the manipulation of known promoters to produce artificial, chimeric, or hybrid promoters. Such promoters can also combine cis-elements from one or more promoters, for example, by adding a heterologous regulatory element to an active promoter with its own partial or complete regulatory elements. Thus, the design, construction, and use of chimeric or hybrid promoters comprising at least one cis-element of SEQ ID NOs: 7, 8, and 9 for modulating the expression of operably linked polynucleotide sequences is encompassed by the present invention.

As used herein, the term "cis-element" refers to a cis-acting transcriptional regulatory element that confers an aspect of the overall control of gene expression. A cis-element may function to bind transcription factors, trans-acting protein factors that

regulate transcription. Some cis-elements bind more than one transcription factor, and transcription factors may interact with different affinities with more than one cis-element. The promoters of the present invention desirably contain cis-elements that can confer or modulate gene expression. Cis-elements can be identified by a number
5 of techniques, including deletion analysis, *i.e.*, deleting one or more nucleotides from the 5' end or internal to a promoter; DNA binding protein analysis using DNase I footprinting, methylation interference, electrophoresis mobility-shift assays, *in vivo* genomic footprinting by ligation-mediated PCR, and other conventional assays; or by DNA sequence similarity analysis with known cis-element motifs by conventional
10 DNA sequence comparison methods. The fine structure of a cis-element can be further studied by mutagenesis (or substitution) of one or more nucleotides or by other conventional methods. Cis-elements can be obtained by chemical synthesis or by isolation from promoters that include such elements, and they can be synthesized with additional flanking nucleotides that contain useful restriction enzyme sites to facilitate
15 subsequence manipulation.

In one embodiment, the promoters of the present invention comprise multiple cis-elements each of which confers a different aspect to the overall control of gene expression. In a preferred embodiment, cis-elements from the polynucleotide molecules of SEQ ID NOs: 7, 8, and 9, are identified using computer programs
20 designed specifically to identify cis-element, domains, or motifs within sequences. Cis-elements may either positively or negatively regulate gene expression, depending on the conditions. The present invention therefore encompasses cis-elements of the disclosed promoters.

As used herein, the phrase "substantially homologous" refers to polynucleotide
25 molecules that generally demonstrate a substantial percent sequence identity with the promoters provided herein. Of particular interest are polynucleotide molecules wherein the polynucleotide molecules function in plants to direct transcription and have at least about 70% sequence identity, at least about 80% sequence identity, at least about 90% sequence identity, or even greater sequence identity, specifically
30 including about 73%, 75%, 78%, 83%, 85%, 88%, 92%, 94, 95 %, 96%, 97%, 98%, 99% or greater sequence identity with the polynucleotide sequences of the promoters described herein, for example, provided in SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9. Polynucleotide molecules that are capable of regulating transcription of

operably linked transcribable polynucleotide molecules and are substantially homologous to the polynucleotide sequences of the promoters provided herein are encompassed within the scope of this invention.

As used herein, the phrase "percent sequence identity" refers to the percentage of identical nucleotides in a linear polynucleotide sequence of a reference polynucleotide molecule (or its complementary strand) as compared to a test polynucleotide molecule (or its complementary strand) when the two sequences are optimally aligned (with appropriate nucleotide insertions, deletions, or gaps totaling less than 20% of the reference sequence over the window of comparison). Optimal alignment of sequences for aligning a comparison window are well known to those skilled in the art and may be conducted by tools such as the local homology algorithm of Smith and Waterman, the homology alignment algorithm of Needleman and Wunsch, the search for similarity method of Pearson and Lipman, and preferably by computerized implementations of these algorithms such as GAP, BESTFIT, FASTA, and TFASTA available as part of the GCG® Wisconsin Package® (Accelrys Inc., San Diego, CA). An "identity fraction" for aligned segments of a test sequence and a reference sequence is the number of identical components which are shared by the two aligned sequences divided by the total number of components in the reference sequence segment, *i.e.*, the entire reference sequence or a smaller defined part of the reference sequence. Percent sequence identity is represented as the identity fraction times 100. The comparison of one or more polynucleotide sequences may be to a full-length polynucleotide sequence or a portion thereof, or to a longer polynucleotide sequence.

As used herein, the term "homology" refers to the level of similarity or percent identity between polynucleotide sequences in terms of percent nucleotide positional identity, *i.e.*, sequence similarity or identity. As used herein, the term homology also refers to the concept of similar functional properties among different polynucleotide molecules, *e.g.*, promoters that have similar function may have homologous cis-elements. Polynucleotide molecules are homologous when under certain conditions they specifically hybridize to form a duplex molecule. Under these conditions, referred to as stringency conditions, one polynucleotide molecule can be used as a probe or primer to identify other polynucleotide molecules that share homology. The phrase "stringent conditions" is functionally defined with regard to the hybridization

of a nucleic-acid probe to a target nucleic acid (*i.e.*, to a particular nucleic-acid sequence of interest) by the specific hybridization procedure discussed in Sambrook *et al.* (2000). Accordingly, the nucleotide sequences of the present invention may be used for their ability to selectively form duplex molecules with complementary stretches of polynucleotide molecule fragments.

Depending on the application envisioned one may desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively high stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50°C to about 70°C. A high stringent condition, for example, is to wash the hybridization filter at least twice with high-stringency wash buffer (0.2X SSC, 0.1% SDS, 65°C). Appropriate moderate stringency conditions that promote DNA hybridization, for example, 6.0 x sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 x SSC at 50°C, are known to those skilled in the art. Additionally, the salt concentration in the wash step can be selected from a low stringency of about 2.0 x SSC at 50°C to a high stringency of about 0.2 x SSC at 50°C. Additionally, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed. Such selective conditions tolerate little mismatch between the probe and the template or target strand. Detection of polynucleotide molecules via hybridization is well known to those of skill in the art, and the teachings of U.S. Patents 4,965,188 and 5,176,995 are exemplary of the methods of hybridization analyses.

Homology can also be determined by computer programs that align polynucleotide sequences and estimate the ability of polynucleotide molecules to form duplex molecules under certain stringency conditions. Polynucleotide molecules from different sources that share a high degree of homology are referred to as "homologues".

Methods well known to one skilled in the art may be used to identify promoters of interest having activity similar to the promoters described herein. For

example, cDNA libraries may be constructed using cells or tissues of interest and screened to identify genes having an expression pattern similar to that of the promoters described herein. The cDNA sequence for the identified gene may then be used to isolate the gene's promoter for further characterization. *See*, for example U.S. Patents 6,096,950; 5,589,583; and 5,898,096, incorporated herein by reference. Alternately, transcriptional profiling or electronic northern techniques may be used to identify genes having an expression pattern similar to that of the promoters described herein. Once these genes have been identified, their promoters may be isolated for further characterization. *See*, for example U.S. Patents 6,506,565 and 6,448,387, incorporated herein by reference. The electronic northern technique refers to a computer-based sequence analysis which allows sequences from multiple cDNA libraries to be compared electronically based on parameters the researcher identifies including abundance in EST populations in multiple cDNA libraries, or exclusively to EST sets from one or combinations of libraries. The transcriptional profiling technique is a high-throughput method used for the systematic monitoring of gene expression profiles for thousands of genes. This DNA chip-based technology arrays thousands of cDNA sequences on a support surface. These arrays are simultaneously hybridized to a population of labeled cDNA probes prepared from RNA samples of different cell or tissue types, allowing direct comparative analysis of expression. This approach may be used for the isolation of regulatory sequences such as promoters associated with those genes.

In another embodiment, the promoter disclosed herein can be modified. Those skilled in the art can create promoters that have variations in the polynucleotide sequence. The polynucleotide sequences of the promoters of the present invention as shown in SEQ ID NOs: 7, 8, and 9 may be modified or altered to enhance their control characteristics. One preferred method of alteration of a polynucleotide sequence is to use PCR to modify selected nucleotides or regions of sequences. These methods are well known to those of skill in the art. Sequences can be modified, for example by insertion, deletion, or replacement of template sequences in a PCR-based DNA modification approach. In the context of the present invention, a "variant" is a promoter containing changes in which one or more nucleotides of an original promoter is deleted, added, and/or substituted, preferably while substantially maintaining promoter function. For example, one or more base pairs may be deleted

from the 5' or 3' end of a promoter to produce a "truncated" promoter. One or more base pairs can also be inserted, deleted, or substituted internally to a promoter. In the case of a promoter fragment, variant promoters can include changes affecting the transcription of a minimal promoter to which it is operably linked. A minimal or basal promoter is a polynucleotide molecule that is capable of recruiting and binding the basal transcription machinery. One example of basal transcription machinery in eukaryotic cells is the RNA polymerase II complex and its accessory proteins. Variant promoters can be produced, for example, by standard DNA mutagenesis techniques or by chemically synthesizing the variant promoter or a portion thereof.

Novel chimeric promoters can be designed or engineered by a number of methods. Many promoters contain cis-elements that activate, enhance, or define the strength and/or specificity of the promoter. For example promoters may contain "TATA" boxes defining the site of transcription initiation and other cis-elements located upstream of the transcription initiation site that modulate transcription levels. For example, a chimeric promoter may be produced by fusing a first promoter fragment containing the activator cis-element from one promoter to a second promoter fragment containing the activator cis-element from another promoter; the resultant chimeric promoter may cause an increase in expression of an operably linked transcribable polynucleotide molecule. Promoters can be constructed such that promoter fragments or elements are operably linked, for example, by placing such a fragment upstream of a minimal promoter. The cis-elements and fragments of the present invention can be used for the construction of such chimeric promoters. Methods for construction of chimeric and variant promoters of the present invention include, but are not limited to, combining control elements of different promoters or duplicating portions or regions of a promoter (*see*, for example, U.S. Patents 4,990,607; 5,110,732; and 5,097,025, all of which are herein incorporated by reference). Those of skill in the art are familiar with the standard resource materials that describe specific conditions and procedures for the construction, manipulation, and isolation of macromolecules (*e.g.*, polynucleotide molecules, plasmids, *etc.*), as well as the generation of recombinant organisms and the screening and isolation of polynucleotide molecules.

In another embodiment, a promoter comprising the polynucleotide sequence shown in SEQ ID NOs: 7, 8, and 9 includes any length of said polynucleotide

sequence that is capable of regulating an operably linked transcribable polynucleotide molecule. For example, the promoters as disclosed in SEQ ID NOs: 7, 8, and 9 may be truncated or portions deleted and still be capable of regulating transcription of an operably linked polynucleotide molecule. In a related embodiment, a cis-element of the disclosed promoters may confer a particular specificity such as conferring enhanced expression of operably linked polynucleotide molecules in certain tissues and therefore is also capable of regulating transcription of operably linked polynucleotide molecules. Consequently, any fragments, portions, or regions of the promoters comprising the polynucleotide sequence shown in SEQ ID NOs: 7, 8, and 9 can be used as regulatory polynucleotide molecules, including but not limited to cis-elements or motifs of the disclosed polynucleotide molecules. Substitutions, deletions, insertions, or any combination thereof can be combined to produce a final construct.

Polynucleotide Constructs

As used herein, the phrase "polynucleotide construct" refers to any recombinant polynucleotide molecule such as a plasmid, cosmid, virus, autonomously replicating polynucleotide molecule, phage, or linear or circular single-stranded or double-stranded DNA or RNA polynucleotide molecule, derived from any source, capable of genomic integration or autonomous replication, comprising a polynucleotide molecule where one or more polynucleotide molecule has been linked in a functionally operative manner. The terms "polynucleotide construct" and "construct" are used interchangeably herein.

As used herein, the phrase "operably linked" refers to a first polynucleotide molecule, such as a promoter, connected with a second transcribable polynucleotide molecule, such as a gene of interest, where the polynucleotide molecules are so arranged that the first polynucleotide molecule affects the function of the second polynucleotide molecule. Preferably, the two polynucleotide molecules are part of a single contiguous polynucleotide molecule and more preferably are adjacent. For example, a promoter is operably linked to a gene of interest if the promoter regulates or mediates transcription of the gene of interest in a cell.

As used herein, the phrase "transcribable polynucleotide molecule" refers to any polynucleotide molecule capable of being transcribed into an RNA molecule.

Methods are known for introducing constructs into a cell in such a manner that the transcribable polynucleotide molecule is transcribed into a functional mRNA molecule that is translated and therefore expressed as a protein product. Constructs may also be constructed to be capable of expressing antisense RNA molecules, in order to inhibit translation of a specific RNA molecule of interest. For the practice of the present invention, conventional compositions and methods for preparing and using constructs and host cells are well known to one skilled in the art, *see* for example, Sambrook *et al.*

Constructs of the present invention would typically contain a promoter operably linked to a transcribable polynucleotide molecule operably linked to a 3' transcription termination polynucleotide molecule. In addition, constructs may include but are not limited to additional regulatory polynucleotide molecules from the 3'-untranslated region (3' UTR) of plant genes (*e.g.*, a 3' UTR to increase mRNA stability of the mRNA, such as the PI-II termination region of potato or the octopine or nopaline synthase 3' termination regions). Constructs may include but are not limited to the 5' untranslated regions (5' UTR) of an mRNA polynucleotide molecule which can play an important role in translation initiation and can also be a genetic component in a plant expression construct. For example, non-translated 5' leader polynucleotide molecules derived from heat shock protein genes have been demonstrated to enhance gene expression in plants (*see*, for example, U.S. Patents 5,659,122 and 5,362,865; and U.S. Published Application No. 2002/0192812, herein incorporated by reference). These additional upstream and downstream regulatory polynucleotide molecules may be derived from a source that is native or heterologous with respect to the other elements present on the promoter construct.

Thus, constructs of the present invention comprise promoters such as those provided in SEQ ID NOs: 7, 8, and 9, or modified as described above, operatively linked to a transcribable polynucleotide molecule so as to direct transcription of said transcribable polynucleotide molecule at a desired level or in a desired tissue or developmental pattern upon introduction of said construct into a plant cell. In some cases, the transcribable polynucleotide molecule comprises a protein-coding region of a gene, and the promoter provides for transcription of a functional mRNA molecule that is translated and expressed as a protein product. Constructs may also be constructed for transcription of antisense RNA molecules or other similar inhibitory

RNA in order to inhibit expression of a specific RNA molecule of interest in a target host cell.

Exemplary transcribable polynucleotide molecules for incorporation into constructs of the present invention include, for example, DNA molecules or genes
5 from a species other than the target gene species, or even genes that originate with or are present in the same species, but are incorporated into recipient cells by genetic engineering methods rather than classical reproduction or breeding techniques. Exogenous gene or genetic element is intended to refer to any gene or DNA molecule that is introduced into a recipient cell. The type of DNA included in the exogenous
10 DNA can include DNA that is already present in the plant cell, DNA from another plant, DNA from a different organism, or a DNA generated externally, such as a DNA molecule containing an antisense message of a gene, or a DNA molecule encoding an artificial or modified version of a gene.

The promoters of the present invention can be incorporated into a construct
15 using marker genes as described and tested in transient analyses that provide an indication of gene expression in stable plant systems. As used herein the phrase "marker gene" refers to any transcribable polynucleotide molecule whose expression can be screened for or scored in some way. Methods of testing for marker gene expression in transient assays are known to those of skill in the art. Transient
20 expression of marker genes has been reported using a variety of plants, tissues, and DNA delivery systems. For example, types of transient analyses can include but are not limited to direct gene delivery via electroporation or particle bombardment of tissues in any transient plant assay using any plant species of interest. Such transient systems would include but are not limited to electroporation of protoplasts from a
25 variety of tissue sources or particle bombardment of specific tissues of interest. The present invention encompasses the use of any transient expression system to evaluate promoters or promoter fragments operably linked to any transcribable polynucleotide molecules, including but not limited to selected reporter genes, marker genes, or genes of agronomic interest. Examples of plant tissues envisioned to test in transients via an
30 appropriate delivery system would include but are not limited to leaf base tissues, callus, cotyledons, roots, endosperm, embryos, floral tissue, pollen, and epidermal tissue.

Any scorable or screenable marker gene can be used in a transient assay. Preferred marker genes for transient analyses of the promoters or promoter fragments of the present invention include a GUS gene (U.S. Patent 5,599,670, herein incorporated by reference) or a GFP gene (U.S. Patent 5,491,084, herein incorporated
5 by reference). The constructs containing the promoters or promoter fragments operably linked to a marker gene are delivered to the tissues and the tissues are analyzed by the appropriate mechanism, depending on the marker. The quantitative or qualitative analyses are used as a tool to evaluate the potential expression profile of the promoters or promoter fragments when operatively linked to genes of agronomic
10 interest in stable plants.

Thus, in one preferred embodiment, a polynucleotide molecule of the present invention as shown in SEQ ID NOs: 7, 8, or 9, or fragments, variants, or derivatives thereof, capable of regulating transcription, is operably linked to a transcribable polynucleotide molecule that provides for a selectable, screenable, or scorable marker.
15 Markers for use in the practice of the present invention include, but are not limited to transcribable polynucleotide molecules encoding β -glucuronidase (GUS), green fluorescent protein (GFP), luciferase (LUC), proteins that confer antibiotic resistance, or proteins that confer herbicide tolerance. Useful antibiotic resistance markers, including those encoding proteins conferring resistance to kanamycin (*nptII*),
20 hygromycin B (*aph IV*), streptomycin or spectinomycin (*aad*, *spec/strep*), and gentamycin (*aac3* and *aacC4*) are known in the art. Herbicides for which transgenic plant tolerance has been demonstrated and the method of the present invention can be applied, include but are not limited to: glyphosate, glufosinate, sulfonyleureas, imidazolinones, bromoxynil, delapone, cyclohezanedione, protoporphyrinogen
25 oxidase inhibitors, and isoxasflutole herbicides. Polynucleotide molecules encoding proteins involved in herbicide tolerance are known in the art, and include, but are not limited to a polynucleotide molecule encoding 5-enolpyruvylshikimate-3-phosphate synthase (EPSP synthase) described in U.S. Patents 5,627,061; 5,633,435; and 6,040,497; and *aroA* described in U.S. Patent 5,094,945 for glyphosate tolerance; a
30 polynucleotide molecule encoding bromoxynil nitrilase (*Bxn*) described in U.S. Patent 4,810,648 for Bromoxynil tolerance; a polynucleotide molecule encoding phytoene desaturase (*crtI*) described in Misawa *et al.* (1993) and Misawa *et al.* (1994) for norflurazon tolerance; a polynucleotide molecule encoding acetohydroxyacid synthase

(AHAS, *aka* ALS) described in Sathasiivan *et al.* (1990) for tolerance to sulfonylurea herbicides; and the *bar* gene described in DeBlock *et al.* (1987) for glufosinate and bialaphos tolerance.

In one preferred embodiment, a polynucleotide molecule of the present invention as shown in SEQ ID NOs: 7, 8, or 9, or fragments, variants, or derivatives thereof, capable of regulating transcription, is operably linked to a transcribable polynucleotide molecule that is a gene of agronomic interest. As used herein, the phrase "gene of agronomic interest" refers to a transcribable polynucleotide molecule that includes but is not limited to a gene that provides a desirable characteristic associated with plant morphology, physiology, growth and development, yield, nutritional enhancement, disease or pest resistance, or environmental or chemical tolerance. The expression of a gene of agronomic interest is desirable in order to confer an agronomically important trait. A gene of agronomic interest that provides a beneficial agronomic trait to crop plants may be, for example, including, but not limited to genetic elements comprising herbicide resistance (U.S. Patents 5,633,435 and 5,463,175), increased yield (U.S. Patent 5,716,837), insect control (U.S. Patents 6,063,597; 6,063,756; 6,093,695; 5,942,664; and 6,110,464), fungal disease resistance (U.S. Patents 5,516,671; 5,773,696; 6,121,436; 6,316,407; and 6,506,962), virus resistance (U.S. Patents 5,304,730 and 6,013,864), nematode resistance (U.S. Patent 6,228,992), bacterial disease resistance (U.S. Patent 5,516,671), starch production (U.S. Patents 5,750,876 and 6,476,295), modified oils production (U.S. Patent 6,444,876), high oil production (U.S. Patents 5,608,149 and 6,476,295), modified fatty acid content (U.S. Patent 6,537,750), high protein production (U.S. Patent 6,380,466), fruit ripening (U.S. Patent 5,512,466), enhanced animal and human nutrition (U.S. Patents 5,985,605 and 6,171,640), biopolymers (U.S. Patent 5,958,745 and U.S. Published Application No. 2003/0028917), environmental stress resistance (U.S. Patent 6,072,103), pharmaceutical peptides (U.S. Patent 6,080,560), improved processing traits (U.S. Patent 6,476,295), improved digestibility (U.S. Patent 6,531,648) low raffinose (U.S. Patent 6,166,292), industrial enzyme production (U.S. Patent 5,543,576), improved flavor (U.S. Patent 6,011,199), nitrogen fixation (U.S. Patent 5,229,114), hybrid seed production (U.S. Patent 5,689,041), and biofuel production (U.S. Patent 5,998,700), the genetic elements and transgenes described in the patents listed above are herein incorporated by reference.

Alternatively, a transcribable polynucleotide molecule can affect the above mentioned phenotypes by encoding a non-translatable RNA molecule that causes the targeted inhibition of expression of an endogenous gene, for example *via* antisense, RNAi, or cosuppression-mediated mechanisms. The RNA could also be a catalytic RNA molecule (*i.e.*, a ribozyme) engineered to cleave a desired endogenous mRNA product. Thus, any polynucleotide molecule that encodes a protein or mRNA that expresses a phenotype or morphology change of interest is useful for the practice of the present invention.

The constructs of the present invention may in one embodiment be double Ti plasmid border DNA constructs that have the right border (RB or AGRtu.RB) and left border (LB or AGRtu.LB) regions of the Ti plasmid isolated from *Agrobacterium tumefaciens* comprising a T-DNA, that along with transfer molecules provided by the *Agrobacterium* cells, permits the integration of the T-DNA into the genome of a plant cell. The constructs also may contain the plasmid backbone DNA segments that provide replication function and antibiotic selection in bacterial cells, for example, an *E. coli* origin of replication such as *ori322*, a broad host range origin of replication such as *oriV* or *oriRi*, and a coding region for a selectable marker such as Spec/Strp that encodes for Tn7 aminoglycoside adenylyltransferase (*aadA*) conferring resistance to spectinomycin or streptomycin, or a gentamicin (Gm, Gent) selectable marker gene. For plant transformation, the host bacterial strain is often *Agrobacterium tumefaciens* ABI, C58, or LBA4404, however, other strains known to those skilled in the art of plant transformation can function in the present invention.

Transformed Plants and Plant Cells

As used herein, the term “transformed” refers to a cell, tissue, organ, or organism into which has been introduced a foreign polynucleotide molecule, such as a construct. Preferably, the introduced polynucleotide molecule is integrated into the genomic DNA of the recipient cell, tissue, organ, or organism such that the introduced polynucleotide molecule is inherited by subsequent progeny. A “transgenic” or “transformed” cell or organism also includes progeny of the cell or organism and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a foreign polynucleotide molecule. A plant transformation construct containing a

promoter of the present invention may be introduced into plants by any plant transformation method. Methods and materials for transforming plants by introducing a plant expression construct into a plant genome in the practice of this invention can include any of the well-known and demonstrated methods including electroporation as
5 illustrated in U.S. Patent 5,384,253; microprojectile bombardment as illustrated in U.S. Patents 5,015,580; 5,550,318; 5,538,880; 6,160,208; 6,399,861; and 6,403,865; *Agrobacterium*-mediated transformation as illustrated in U.S. Patents 5,635,055; 5,824,877; 5,591,616; 5,981,840; and 6,384,301; and protoplast transformation as illustrated in U.S. Patent 5,508,184, all of which are incorporated herein by reference.

10 Methods for specifically transforming dicots are well known to those skilled in the art. Transformation and plant regeneration using these methods have been described for a number of crops including, but not limited to, soybean (*Glycine max*), *Brassica* sp., *Arabidopsis thaliana*, cotton (*Gossypium hirsutum*), peanut (*Arachis hypogae*), sunflower (*Helianthus annuus*), potato (*Solanum tuberosum*), tomato
15 (*Lycopersicon esculentum* L.), and alfalfa (*Medicago sativa*). It is apparent to those of skill in the art that a number of transformation methodologies can be used and modified for production of stable transgenic plants from any number of target crops of interest.

In another embodiment, the present invention provides a method of making a
20 vegetable oil, comprising the steps of incorporating into the genome of a plant a promoter of the present invention operably linked to a transcribable polynucleotide molecule conferring altered oil and/or protein content, growing the plant to produce seeds, and extracting the oil and/or protein from the seed.

The transformed plants may be analyzed for the presence of the genes of
25 interest and the expression level and/or profile conferred by the promoters of the present invention. Those of skill in the art are aware of the numerous methods available for the analysis of transformed plants. For example, methods for plant analysis include, but are not limited to Southern blots or northern blots, PCR-based approaches, biochemical analyses, phenotypic screening methods, field evaluations,
30 and immunodiagnostic assays.

The seeds of this invention can be harvested from fertile transgenic plants and be used to grow progeny generations of transformed plants of this invention. The terms "seeds" and "kernels" are understood to be equivalent in meaning. In the

context of the present invention, the seed refers to the mature ovule consisting of a seed coat, embryo, aleurone, and an endosperm.

The phrase "cell proliferation" refers to cells undergoing mitotic cell divisions, such as in rapidly growing tissues.

5 The phrase "micronutrient content" means the amount of micronutrients, *i.e.*, vitamins or carotenoids, within a seed expressed on a per weight basis.

The phrase "oil content" means the concentration of the oil fraction, expressed as a weight percentage or parts per million basis, which may be determined, for example, by low-resolution ¹H nuclear magnetic resonance (NMR) (Tiwari *et al.*, 10 1974) or Rubel, 1994) or near infrared transmittance (NIT) spectroscopy (Orman *et al.*, 1992; Patrick *et al.*, 1997).

The phrase "protein quality" means the level of one or more essential amino acids, whether free or incorporated in protein, namely histidine, isoleucine, leucine, lysine, methionine, cysteine, phenylalanine, tyrosine, threonine, tryptophan, and 15 valine.

The following examples are included to demonstrate preferred embodiments of the present invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventors to function well in the practice of the present invention. However, those 20 of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the present invention, therefore all matter set forth or shown in the accompanying drawings is to be interpreted as illustrative and not in a limiting sense.

25

Example 1

This example describes the identification and isolation of expressed sequence tags (ESTs) from *Brassica napus*, using an electronic library subtraction method. Proprietary *B. napus* EST libraries were used as the target tissue and proprietary *Arabidopsis* and *B. napus* cDNA libraries as the background tissue. ESTs identified 30 as potentially being expressed only in the silique tissue were further characterized by hybridization.

EST Library Construction

The target tissue EST Library was constructed from a variety of tissues isolated from greenhouse grown *Brassica napus* (cultivar Quantum). The isolated tissues included: silique walls at 10, 20, 30, 40, and 50 days after pollination (DAP), aleurone and seed coat at 25-28 DAP. Tissues were flash frozen in liquid nitrogen and stored at -70°C prior to RNA extraction. RNA was extracted by grinding approximately 500 mg plant tissue in liquid nitrogen with a mortar and pestle. The ground tissue was mixed with 5 ml REC.8 + (Tris-HCl, pH9, 0.8 M NaCl, 10 mM EDTA, 0.5% β-mercaptoethanol, and 0.5% cetyltrimethyl-ammonium bromide) plus 0.1g of polyvinylpyrrolidone. The mixture was homogenized with a mortar and pestle and was centrifuged at 5000g for 5 minutes. The aqueous phase was extracted once with 1.5 ml phenol:chloroform (1:1, v/v) and once with 1.5 ml chloroform. RNA was precipitated from the aqueous layer with an equal volume of 95% ethanol and pelleted at 10,000 x g for 20 minutes. Precipitated RNA was resuspended in water and one volume of 8M LiCl were added. The mixture was incubated at -20°C for 1 hour, followed by centrifugation at 10,000 x g for 20 minutes. Precipitated RNA was washed with 70% ethanol, air dried, and dissolved in water treated with diethylpyrocarbonate (DEPC water). Approximately 25 µg of total RNA was used to prepare double-stranded cDNA using the SuperScript Plasmid System (Invitrogen Corporation Carlsbad, CA) according to the manufacturer's protocol.

EST sequences were generated for each of the six libraries described above. Candidate sequences were identified by using the EST sequences from each library as a query against Monsanto proprietary *Arabidopsis* and *Brassica napus* cDNA libraries and the public GenBank® cDNA collection. GenBank® is the NIH genetic sequence database, an annotated collection of all publicly available DNA sequences (*Nucleic Acids Research*, 32(1):23-6 (2004)). The most highly expressed genes from each library were identified by determining the frequency of appearance in all cDNA collections. An additional BLAST search was completed to identify clones specific for each library.

A second bioinformatics screen was then done using a proprietary Library Subtraction Program. The six libraries described above were used as target libraries. All available Monsanto proprietary, non-silique wall *Brassica napus* cDNA libraries were used as background libraries. The sequences that were present in any of the target libraries but not present in any of the background libraries, were identified as

meeting the criteria for further investigation. The 200 most abundant messages meeting these criteria were characterized further. Fifteen sequences were eliminated for overlap with the initial screen. An additional seven sequences were eliminated for significant identity with retrotransposons. The remaining sequences were then
5 BLAST searched against *Arabidopsis* cDNA libraries derived from leaf, root, stem, flower bud, open flower, and developing seed. Eighty-nine of these sequences were eliminated from consideration for having significant identity to non-silique wall sequences. The remaining 89 EST clones were thus identified from this screening and were further investigated by hybridization screening.

10 Hybridization Screening

The EST sequences identified above were digested with *S*all and *Not*I to release the insert. Fragments were separated on 0.8% agarose gels in 1X TBE. To facilitate transfer, the gels were exposed to short wave uv light for 2–3 minutes. The DNA was denatured in 0.5N NaOH; 1.5 M NaCl and transferred to Nytran Plus
15 (Schleicher & Schuell BioScience, Dassel, Germany) using the TurboBlotter system (Schleicher & Schuell). Blots were hybridized to radiolabelled cDNA derived from silique walls at 10, 20, 30, 40, and 50 DAP, aleurone/seed coat, developing embryo (15-18 DAP), floral buds, open flower, leaf, and root using the SMART cDNA PCR amplification Kit (BD Biosciences, Franklin Lakes, NJ) according to manufacturer's
20 instructions with the addition of radiolabelled dCTP to the final PCR reaction. A number of the clones were eliminated due to hybridization to cDNAs that were derived from developing embryo (15-18 DAP), floral buds, open flower, leaf, and/or root. The remaining 67 clones were used as probes against SMART™ cDNAs (BD Biosciences) derived RNA isolated from silique walls at 10, 15, 20, 25, 30, 35, 40, 45,
25 and 50 DAP, aleurone/seed coat, developing embryo (15-18 DAP), floral buds, open flower, leaf, and root. Six candidates hybridized predominantly to silique wall and/or aleurone. One clone hybridized specifically to root derived cDNA.

Clone LIB4156-008-R1-K1-A4 was characterized as being expressed in the root cells. Clone LIB4153-003-R1-K9-C1 was characterized as being expressed in
30 silique wall, aleurone/seed coat, and open flower. Clone LIB4156-012-R1-K1-D2 was characterized as being expressed in silique wall, aleurone/seed coat, and root. Clone LIB4153-011-R1-K1-E3 was characterized as being expressed in silique wall,

aleurone/seed coat, and open flower. Clone LIB4169-011-Q1-K1-G2 was characterized as being expressed in aleurone/seed coat. Clone LIB4169-008-Q1-K1-D8 was characterized as being expressed in silique wall, aleurone/seed coat, and developing embryo. Clones LIB4153-003-R1-K9-C1, LIB4156-012-R1-K1-D2, 5 LIB4153-011-R1-K1-E3, and LIB4156-008-R1-K1-A4 were submitted for sequence confirmation. All but clone LIB4156-008-R1-K1-A4 were verified as being the expected sequence. The correct clone designation for LIB4156-008-R1-K1-A4 was determined to be LIB4156-011-R1-K1-C3.

10

Example 2

This example describes the isolation of promoter sequences from *Brassica napus* using the EST sequences identified above.

Genome Walker Library Preparation

Brassica napus leaves were harvested and kept frozen in liquid nitrogen until 15 extraction. The tissue was ground to a fine powder using a mortar and pestle while keeping the tissue frozen with liquid nitrogen. Approximately 0.4 grams of ground tissue was transferred to a centrifuge tube. Six milliliters of 65°C SDS extraction buffer (1M Tris; 0.25M EDTA; pH 8.0 with HCL; 20% SDS; 5M NaCl β -mercaptoethanol (7.1:10)) was added just before use. The mixture was vortexed 20 vigorously for 30-60 seconds. Samples were then incubated at 65°C for 45 minutes, with inversion every 15 minutes. Two milliliters of ice temperature 5M potassium acetate solution was added, the samples were inverted gently to mix, and then incubated on ice for 20 minutes. The samples were chloroform extracted by adding 3 25 mls of chloroform to each tube and shaking gently for 10 minutes. The samples were then centrifuged at 9,000 RPM for 20 minutes. The supernatant was filtered through a layer of miracloth and collected in a clean centrifuge tube. The DNA was precipitated by adding 2 ml of 100% isopropanol gently mixed by inverting the tubes 3-4 times, and then centrifuging at 9,000 RPM for 20 minutes. The isopropanol was poured off, and the pellet was resuspended in 200 μ l RNaseA solution (1 μ l of a 100mg/ml stock 30 from Qiagen Inc., Valencia, CA, diluted in 1 ml 50mM Tris-HCL; 10mM EDTA). A volume of 300 μ l of ammonium acetate/100% isopropanol (1:7) was added. Samples were vortexed and spun at 9,000 RPM for 15 minutes. Resulting supernatant was

poured off, and the pellet was washed with 500µl of 80% ethanol. The ethanol was gently decanted and the pellet was allowed to air dry on the bench top. The pellet was resuspended in approximately 200µl TE Buffer (10 mM Tris-HCL pH8, 1 mM EDTA). The resulting DNA was used to prepare Genome Walker™ libraries following the manufacturer's recommended protocol (Clontech, Palo Alto, CA).

Isolation of P-BN.RPC-0:1:1

The P-BN.RPC-0:1:1 root preferred promoter sequence [SEQ ID NO: 1] was isolated from the Genome Walker™ libraries described above. Primers were designed using the manufacturer's parameters for length, melting temperature (T_m), and sequence composition, from the available EST sequence of clone LIB4156-011-R1-K1-C3. A first round PCR reaction was run using the Genome Walker libraries as the template and the following primers:

GSP1

5'-GCTACGGCGATGAGAAGAGAAAATAAAATG-3' [SEQ ID NO: 2], and

AP1

5'-GTAATACGACTCACTATAGGGC-3' [SEQ ID NO: 3] (supplied by BD

Biosciences).

The resulting amplification product was isolated using standard techniques well know in the art. A second PCR reaction was then performed using the first round amplification product as the template and the following primers:

GSP2

5'-CCCACAGATTTTCTATTTCGGTTCTCATAC-3' [SEQ ID NO: 4], and

AP2

5'-ACTATAGGGCACGCGTGGT-3' [SEQ ID NO: 5] (supplied by BD

Biosciences).

The product of the PCR reaction was purified according to standard methodology well known in the art and cloned into pCR2.1 Topo (Invitrogen Corporation, Carlsbad, California) according to manufacturer's instructions. The resulting plasmid was named 2d11-2. The sequence of this clone was determined using standard sequencing methodologies as set forth by PE Applied Biosystems, (Foster City, CA). To facilitate further cloning, an NcoI site was added to the 3' end of the promoter fragment, following the PCR protocol recommended by the enzyme

manufacturer (PE Applied Biosystems). Briefly, approximately 10 nanograms of plasmid 2d11-2 was used as the template and amplified using 30 nanomoles each of the following primers:

- D11 nco (#18337)
 5'-GAGAAACCATGGTGAATAAATGG-3' [SEQ ID NO: 6], and
 AP1 [SEQ ID NO: 3, described above].

Additionally, 10 micromoles each of dATP, dCTP, dGTP, and TTP, and 2.5 units of AmpliTaq Gold in 1X Opti-Prime™ Buffer 3 (Stratagene), were used in the reaction. After an initial incubation at 95°C for 10 minutes, 25 cycles of PCR were performed with a cycle define as 92°C for 30 seconds, 56°C for 30 seconds, and 72°C for 2 minutes, followed by 1 cycle of 72°C for 7 minutes. The product of the PCR reaction was purified according to standard methodology well known in the art and cloned into pCR2.1 Topo (Invitrogen) according to manufacturer's instructions. The resulting plasmid was named pMON69833. The sequence of this clone was determined using standard sequencing methodologies as set forth by PE Applied Biosystems.

Isolation of P-BN.SW1-0:1:2, P-BN.SW2-0:1:2, and P-BN.SW3-0:1:2

The silique wall preferred promoter sequences, P-BN.SW1-0:1:2, P-BN.SW2-0:1:2, and P-BN.SW3-0:1:2 were isolated from the Genome Walker™ libraries using a procedure similar to that described above for P-BN.RPC-0:1:1 with the manufacturer's instruction. For the first round PCR reactions, primers were designed for each target sequence, using the manufacturer's parameters for length, T_m and sequence composition, from the respective EST sequences.

The table below shows the EST clones used as template DNA and the primers used for the first round PCR reactions in the isolation of the listed promoter sequences.

Promoter	EST clone ID	Primer 1	Primer 2
P-BN.SW1-0:1:2	LIB4153-003-R1-K9-C1	#17637	#17636
P-BN.SW2-0:1:2	LIB4156-012-R1-K1-D2	#17669	#17668
P-BN.SW3-0:1:2	LIB4153-011-R1-K1-E3	#18210	#18209

Primer Sequences

- #17637 5'-TTCATATCTGCGTAAGTACGTCCATGTTC-3' [SEQ ID NO: 10]
 #17636 5'-CGAGAAACAGAATGATGAAACTAGAGAGAC-3' [SEQ ID NO: 11]
 #17669 5'-CACCATGCTTTCACGTC'TTTGAGTTG-3' [SEQ ID NO: 12]
 #17668 5'-TCGATGCCACGAAAATGCATAAGAAC-3' [SEQ ID NO: 13]
 5 #18210 5'-CCTCATCCTCATCCTCATCCTCAACC-3' [SEQ ID NO: 14]
 #18209 5'-CCGTACACGGTAAAATAGTTCTTGACGGA-3' [SEQ ID NO: 15]

A second amplification was then performed, similar to that described above for P-BN.RPC-0:1:1, using the first round amplification products from the respective reactions as the template, and the following primers:

- 10 P-BN.SW1-0:1:1 GSP2 [SEQ ID NO: 4] and
 AP2 [SEQ ID NO: 5] (supplied by the manufacturer),
 P-BN.SW2-0:1:1 GSP1 [SEQ ID NO: 2] and
 AP2 [SEQ ID NO: 5],
 15 P-BN.SW3-0:1:1 GSP2 [SEQ ID NO: 4] and
 AP2 [SEQ ID NO: 5].

The products of the respective PCR reactions were purified according to standard methodology well known in the art and cloned into pCR2.1 Topo (Invitrogen) according to manufacturer's instructions. The resulting plasmids were named 3d2-2, 3d3-46, and 4G5-1 for promoters P-BN.SW1-0:1:2, P-BN.SW2-0:1:2, 20 and P-BN.SW3-0:1:2, respectively. The sequences for these clones were determined using standard sequencing methodologies as set forth by PE Applied Biosystems, (Foster City, CA).

Similar to the procedure described above for P-BN.RPC-0:1:1, an NcoI site was added to the 3' end of each promoter fragment to facilitate cloning. As described above, the reaction conditions for the PCR reactions followed the protocol recommended by the enzyme manufacturer (PE Applied Biosystems, Foster City, CA). The primers for the specific reactions were as follows:

- Plasmid DNA 3d2-2 (P-BN.SW1-0:1:2)**
 30 #18115 5'-CTAAAGCCATGGTCTTAGAAAAGTTG-3' [SEQ ID NO: 16], and
 #18167 5'-AATACGACTCACTATAGGGCACGC-3' [SEQ ID NO: 17];
Plasmid DNA 3d3-46 (P-BN.SW2-0:1:2)
 #18206 5'-GAAAAGCCATGGTAAGGCCAATATT-3' [SEQ ID NO: 18], and

API [SEQ ID NO: 3]

Plasmid DNA 4G5-1 (P-BN.SW3-0:1:2)

#18340 5'-CTTCTGCCATGGAAAAGAAAAAGTTG-3' [SEQ ID NO: 19], and

API [SEQ ID NO: 3].

5 The products of the PCR reactions were purified according to standard methodology well known in the art and cloned into pCR2.1 Topo (Invitrogen) according to manufacturer's instructions. The resulting plasmids were named pMON69828, pMON69829, and pMONM69830, containing the nucleic acid sequences for P-BN.SW1-0:1:2, P-BN.SW2-0:1:2, and P-BN.SW3-0:1:2, 10 respectively. The sequences of these clones were determined using standard sequencing methodologies as set forth by PE Applied Biosystems.

Example 3

This example describes the construction of plant transformation vectors containing the promoter sequences described above. The promoter sequences were 15 linked to the marker gene, *E. coli uidA*, resulting in constructs designed to demonstrate expression in *Arabidopsis* and Canola.

PMON69832 (P-BN.RPC-0:1:1::GUS)

A 1661 bp fragment containing P-BN.RPC-0:1:1 [SEQ ID NO: 1] was removed from pMON69833 by digestion with SmaI and Nco I. The fragment was 20 ligated into the plasmid pMON65425, which had also been digested with SmaI and NcoI. The resulting plasmid, containing P-BN.RPC-0:1:1 driving the *E. coli uidA* gene and with the napin 3' UTR was named pMON69832. The nucleic acid sequence was determined using known methodology and confirmed the integrity of the cloning junctions.

25 **PMON69827 (P-BN.SW1-0:1:2::GUS)**

A 1297 bp fragment containing P-BN.SW1-0:1:2 [SEQ ID NO: 7] was removed from pMON69828 by digestion with SmaI and Nco I. The fragment was ligated into the plasmid pMON65425, which had also been digested with SmaI and NcoI. The resulting plasmid, containing P-BN.SW1-0:1:2 driving the *E. coli uidA* 30 gene and with the napin 3' UTR, was named pMON69827. The nucleic acid sequence was determined using known methodology and confirmed the integrity of the cloning junctions.

PMON69825 (P-BN.SW2-0:1:2::GUS)

A 854 bp fragment containing P-BN.SW2-0:1:2 [SEQ ID NO: 8] was removed from pMON69829 by digestion with SmaI and Nco I. The fragment was ligated into pMON65425, which had also been digested with SmaI and NcoI. The resulting plasmid, containing P-BN.SW2-0:1:2 driving the *E. coli uidA* gene and with the napin 3' UTR was named pMON69825. The nucleic acid sequence was determined using known methodology and confirmed the integrity of the cloning junctions.

PMON69831 (P-BN.SW3-0:1:2::GUS)

An 888 bp fragment containing P-BN.SW3-0:1:2 (SEQ ID NO: 9) was removed from pMON69830 by digestion with SmaI and Nco I. The fragment was ligated into pMON65425, which had also been digested with SmaI and NcoI. The resulting plasmid, containing P-BN.SW3-0:1:2 driving the *E. coli uidA* gene and with the napin 3' UTR was named pMON69831. The nucleic acid sequence was determined using known methodology and confirmed the integrity of the cloning junctions.

Example 4

This example sets forth the procedure for transforming *Arabidopsis* plants with the vectors described above. *Arabidopsis* plants were grown by sowing seeds onto 4-inch pots containing reverse osmosis water (ROW) saturated MetroMix 200 (The Scotts Company, Columbus, OH). The plants were vernalized by placing the pots in a covered flat, in a growth chamber at 4-7°C, 8 hours light/day for 4-7 days. The flats were transferred to a growth chamber at 22°C, 55% relative humidity, and 16 hours light/day at an average intensity of 160-200 $\mu\text{Einstein/s/m}^2$. The cover was lifted and slid back 1-inch after germination, and removed when the true leaves had formed. The plants were bottom watered, as needed, with ROW until 2-3 weeks after germination. Plants were then bottom watered, as needed, with Plantex 15-15-18 solution (Plantex Corporation Ottawa, Canada) at 50 ppm N2. Pots were thinned such that 1 plant remained per pot at 2-3 weeks after germination. Once the plants began to bolt, the primary inflorescence was trimmed to encourage the growth of axillary bolts.

Transgenic *Arabidopsis thaliana* plants were obtained as described by Bent *et al.* (1994), or Bechtold *et al.* (1993). Cultures of *Agrobacterium tumefaciens* strain ABI containing either one of the transformation vectors pMON69832, pMON69827,

pMON69825, or pMON69831 were grown overnight in LB (10% bacto-tryptone, 5% yeast extract, and 10% NaCl with kanamycin (75 mg/L), chloramphenicol (25 mg/L), and spectinomycin (100 mg/L)). The bacterial culture was centrifuged and resuspended in 5% sucrose + 0.05% Silwet-77 solution. The aerial portions of whole
5 *Arabidopsis* plants (at about 5-7 weeks of age) were immersed in the resulting solution for 2-3 seconds. The excess solution was removed by blotting the plants on paper towels. The dipped plants were placed on their side in a covered flat and transferred to a growth chamber at 19°C. After 16 to 24 hours the dome was removed and the plants were set upright. When plants had reached maturity, water was
10 withheld for 2-7 days prior to seed harvest. Harvested seed was passed through a stainless steel mesh screen (40 holes/inch) to remove debris. The harvested seed was then stored in paper coin envelopes at room temperature until analysis.

The seeds from the transgenic *Arabidopsis* plants were surfaced sterilized using a vapor phase sterilization protocol. Briefly, an open container of seeds was
15 placed in a dessicator with a beaker containing 100 ml of household bleach. Immediately prior to sealing the dessicator, 3 ml concentrated HCl was added to the bleach solution to generate chlorine gas. The dessicator is sealed and a vacuum was applied to allow sterilization by chlorine fumes. The seeds were incubated for several hours. Sterilized seed were sprinkled onto *Arabidopsis* germination media [MS Salts
20 (1X); sucrose (1%); myo-inositol (100 mg/L); thiamine-HCl (1 mg/L); pyridoxine-HCl (500 mg/L); nicotinic acid (500 mg/L); MES pH 5.7 (0.05%) and Phytagar (0.7%)] supplemented with 50 mg/L glyphosate. Up to 16 glyphosate resistant seedlings were transplanted to 2¼-inch pots containing MetroMix 200, thinned to one seedling per pot, and were grown under the conditions described above until the initial
25 siliques that had formed began to desiccate. Tissue (rosette leaf, cauline leaf, stem, flowers, floral buds, and developing siliques) was removed from each T1 plant for subsequent histochemical staining.

Canola plants were transformed using the protocol described by Moloney and Radke in U.S. Patent 5,720,871. Briefly, seeds of *Brassica napus* cv Ebony were
30 planted in 2-inch pots containing Metro Mix 350 (The Scotts Company, Columbus, OH). The plants are grown in a growth chamber at 24°C, and a 16/8 hour photoperiod, with light intensity of 400 µEm-2 sec-1 (HID lamps). After 2-1/2 weeks, the plants were transplanted into 6-inch pots and grown in a growth chamber at

15/10°C day/night temperature, 16/8 hour photoperiod, light intensity of 800 μEm^{-2} sec⁻¹ (HID lamps). Four terminal internodes from plants just prior to bolting or in the process of bolting but before flowering are removed and surface sterilized in 70% v/v ethanol for 1 minute, 2% w/v sodium hypochlorite for 20 minutes and rinsing 3 times
5 with sterile deionized water. Six to seven stem segments are cut into 5 mm discs, maintaining orientation of basal end. Cultures of *Agrobacterium tumefaciens* strain ABI containing one of the transformation vectors pMON69832, pMON69827, pMON69825, or pMON69831 are grown overnight on a rotator shaker at 24°C in 2 ml of Luria Broth, LB, (10% bacto-tryptone, 5% yeast extract, and 10% NaCl)
10 containing 50 mg/l kanamycin, 24 mg/l chloramphenicol and 100 mg/l spectinomycin. A 1:10 dilution is made in MS media (Murashige and Skoog, 1962) giving approximately 9×10^8 cells per ml. The stem discs (explants) are inoculated with 1.0 ml of *Agrobacterium* and the excess is aspirated from the explants. The explants are placed basal side down in petri plates containing media comprising 1/10 MS salts, B5
15 vitamins (1% inositol; 0.1% thiamine HCl; 0.01% nicotinic acid; 0.01% pyridoxine-HCl), 3% sucrose, 0.8% agar, pH 5.7, 1.0 mg/l 6-benzyladenine (BA). The plates are layered with 1.5 ml of media containing MS salts, B5 vitamins, 3% sucrose, pH 5.7, 4.0 mg/l p-chlorophenoxyacetic acid, 0.005 mg/l kinetin and covered with sterile filter paper. Following a 2 to 3 day co-culture, the explants are transferred to deep dish
20 petri plates containing MS salts, B5 vitamins, 3% sucrose, 0.8% agar, pH 5.7, 1 mg/l BA, 500 mg/l carbenicillin, 50mg/l cefotaxime, and 25 mg/l glyphosate for selection. Seven explants are placed on each plate. After 3 weeks they are transferred to fresh media, 5 explants per plate. The explants are cultured in a growth room at 25°C, continuous light.
25 The transformed plants are grown in a growth chamber at 22°C in a 16-8 hours light-dark cycle with light intensity of 220 $\mu\text{Em}^{-2}\text{s}^{-1}$ for several weeks before transferring to the greenhouse. The plants are then grown in greenhouse conditions until maturity. The resulting mature R1 seeds are collected for analysis. Plants are maintained in a greenhouse under standard conditions. Developing seed is harvested
30 at various stages after pollination and stored at negative 70°C. Mature seed is collected and stored under controlled conditions consisting of about 17°C and 30% humidity.

Example 5

This example describes the analysis of the expression of β -glucuronidase in *Arabidopsis thaliana* plants transformed with pMON69832, pMON69827, pMON69825, or pMON69831 using histochemical staining. The results of this
5 analysis provide a characterization of the expression for the different promoters.

Freshly harvested tissues from transformed *Arabidopsis* plants were incubated for approximately 24 hours at 37°C in a solution containing 50 mM NaPO₄ (pH 7.2); 100 μ M potassium ferricyanide; 100 μ M potassium ferrocyanide, 0.03% Triton X-100; 20% methanol and 2.5 mg/ml 5-bromo-4-chloro-3-indoyl glucuronic acid (X-gluc). In some cases the potassium ferricyanide, potassium ferrocyanide, and methanol were omitted from the staining solution. The stained tissue was cleared of chlorophyll by an overnight incubation in 70% ethanol/30% H₂O at 37°C. Stained tissues were photographed immediately or transferred to a solution of 70% ethanol/30% glycerol (v/v) and stored at 4°C until photographed. The samples were
10 then scored as positive (+) or negative (-) for staining a blue color.

For detection of expression in canola, up to 5 siliques were harvested from individual R0 plants at several time points after pollination. Siliques were scored with an 18 gauge needle to allow the staining solution to contact the developing seed. The siliques were incubated for approximately 24 hours at 37°C in a solution containing 50
20 mM NaPO₄ (pH 7.2); 100 μ M potassium ferricyanide; 100 μ M potassium ferrocyanide, 0.03% Triton X-100; 20% methanol and 2.5 mg/ml 5-bromo-4-chloro-3-indoyl glucuronic acid (X-gluc). The stained tissue was cleared of chlorophyll by an overnight incubation in 70% ethanol/30% H₂O at 37°C. Stained tissues were photographed immediately or transferred to a solution of 70% ethanol/30% glycerol
25 (v/v) and stored at 4°C until photographed. Samples were scored positive (+) or negative (-) for blue color.

For P-BN.RPC-0:1:1, a root preferred promoter contained in pMON69832, none of 10 plants screened had detectable levels of activity in the silique wall (data not shown). For pMON69827, 3 out of 10 plants screened had detectable levels of
30 activity in the silique wall from at least one time point. For pMON69825, 6 out of 10 plants screened had detectable levels of activity in the silique wall from at least one time point. For pMON69831, 5 out of 10 plants screened had detectable levels of

activity in the silique wall from at least one time point. No activity was detected in the seed coat or aleurone of plants transformed with pMON69831. Time course data for individual transformants is presented in Tables 1-3.

Table 1. P-BN.SW1-0:1:2 Expression in Developing Canola Siliques

Days After Pollination

Construct	Event	3	6	9	12	15	20	25	30	35	40
pMON69827	BN_G1427	-	-	-	-	-	-	-	-	-	-
pMON69827	BN_G1428	-	-	-	-	-	-	-	-	-	-
pMON69827	BN_G1429	-	-	-	-	-	-	-	-	-	-
pMON69827	BN_G1430	-	-	-	-	-	-	-	-	-	-
pMON69827	BN_G1480	-	-	-	-	-	-	-	-	-	-
pMON69827	BN_G1481	-	ND	-	ND	+	-	-	-	-	-
pMON69827	BN_G1482	-	-	-	+	-	+	-	-	-	+
pMON69827	BN_G1507	+	+	+	+	+	+	+	+	+	+
pMON69827	BN_G1539	-	-	-	-	-	-	-	-	-	-
pMON69827	BN_G1540	-	-	-	-	-	-	-	-	-	-
Control	SP30052	-	-	-	-	-	-	-	-	-	-

Table 2. P-BN.SW2-0:1:2 Expression in Developing Canola Siliques

Days After Pollination

5

Construct	Event	3	6	9	12	15	20	25	30	35	40
pMON69825	BN_G1474	-	-	-	-	-	-	-	-	-	-
pMON69825	BN_G1475	-	-	-	-	-	-	-	-	-	-
pMON69825	BN_G1476	+	+	+	+	NA	+	+	+	+	+
pMON69825	BN_G1477	-	+	+	+	NA	+	+	-	-	-
pMON69825	BN_G1478	-	-	-	-	-	-	-	-	-	-
pMON69825	BN_G1479	-	ND	+	+	-	-	-	-	-	-
pMON69825	BN_G1536	-	-	-	-	-	-	-	-	-	+
pMON69825	BN_G1537	-	-	-	-	+	+	+	-	-	-
pMON69825	BN_G1538	+	-	-	-	+	+	-	-	-	-
pMON69825	BN_G1580	-	-	-	-	-	-	-	-	-	-
Control	SP30052	-	-	-	-	-	-	-	-	-	-

Table 3. P-BN.SW3-0:1:2 Expression in Developing canola Siliques

Construct	Event	Days After Pollination									
		3	6	9	12	15	20	25	30	35	40
pMON69831	BN_G1839	+	+	+	+	+	-	+	+	+	+
pMON69831	BN_G1901	-	-	-	-	-	-	-	-	-	-
pMON69831	BN_G1902	-	-	-	-	-	-	-	-	-	-
pMON69831	BN_G1903	-	-	-	-	-	-	-	-	-	-
pMON69831	BN_G1904	-	-	-	-	-	-	-	-	-	-
pMON69831	BN_G1905	-	-	-	+	+	+	-	-	-	-
pMON69831	BN_G1906	-	-	+	+	+	+	+	+	+	+
pMON69831	BN_G1907	-	-	-	-	+	+	+	-	-	-
pMON69831	BN_G1908	-	-	-	-	+	-	+	-	-	-
pMON69831	BN_G1951	-	-	-	-	-	-	-	-	-	-
Control	SP30052	-	-	-	-	-	-	-	-	-	-

5 The data in Tables 1-3 indicate that the three promoters, P-BN.SW1-0:1:2 (SEQ ID NO: 7), P-BN.SW2-0:1:2 (SEQ ID NO: 8), and P-BN.SW3-0:1:2 (SEQ ID NO: 9) are capable of driving silique wall expression in *Arabidopsis* and Canola.

Example 6

10 This example describes the construction of plant transformation vectors containing the P-BN.SW3-0:1:2 promoter operably linked to the *Arabidopsis thaliana* isocitrate lyase gene. To facilitate cloning, the P-BN.SW3-0:1:2 promoter was amplified using pMON69830 as a template with the following primers:

19944

5'-CCCGGGCTGGTCTCGCGAAGATTCTCAG-3' [SEQ ID NO: 20], and

15 19945

5'-

CTCGAGCGGCCGCTTCTATGTCGACTGGAAAAGAAAAGTTGTGCCAACAA
AAG-3' [SEQ ID NO: 21].

20 The reactions were run under conditions recommended by the Expand High Fidelity enzyme manufacturer (Roche Applied Biosciences, Indianapolis, IN). The product of the PCR reaction was purified according to standard methodology well

known in the art and cloned into pCR2.1 Topo (Invitrogen) according to the manufacturer's instructions. The resulting plasmid was named pMON79578.

A 910-base pair fragment containing P-Bn.SW3-0:1:2 [SEQ ID NO: 9] was removed from the vector pMON79578 by digestion with SmaI and NotI. The
5 fragment was cloned into pMON82374 which had been digested with PmeI and NotI in place of the pBr.Snap2 promoter. The vector pMON82374 contains a nopaline T-DNA right border sequence and octopine T-DNA left border sequence, with 2 expression cassettes contained between the 2 T-DNA borders. The first cassette contains the pBr.Snap2 promoter and T-Br.Snap2-1 terminator. The second cassette
10 contains with a 35S promoter from the Figwort Mosaic Virus (FMV) between the 2 T-DNA borders, driving the expression of a chimeric EPSP synthase gene containing a chloroplast targeting sequence from the *Arabidopsis* EPSP synthase gene (GenBank identifier number gi:16272) linked to a synthetic EPSP synthase coding region (U.S. Patent 5,633,435) and the 3' untranslated region from the pea *rbcS E9* gene.
15 Additionally, pMON82374 contains recognition sites for cre recombinase. The recombinase sites are 5' of the FMV promoter and 3' of the E9 3' UTR. The resulting plasmid was named pMON79581.

A 2111 fragment containing L-At.ICL, CR-At.ICL, and T-At.ICL was removed from EST clone pMON82360 by digestion with SalI and NotI restriction
20 enzymes and cloned between the pBr.Snap2 promoter and T-Br.Snap2-1 3' UTR in the SalI and Not I digested pMON79581. The resulting plasmid was named pMON79583. The nucleic acid sequence was determined using known methodology and confirmed the integrity of the cloning junctions. pMON79583 was used for the transformation of *Brassica napus*.

An 891-base pair fragment [SEQ ID NO: 22] containing the P-Bn.SW3-0:1:2 promoter sequence [SEQ ID NO: 9], was removed from the vector pMON79578 by
25 digestion with SmaI and SalI. The fragment was cloned into DMRUEZ03.0112 which had been digested with PacI and SalI. Prior to SalI digestion and gel purification, the PacI overhang was blunt ended using Pfu polymerase according to the manufacturer's instructions (Stratagene). The resulting plasmid contains the nopaline T-DNA right
30 border sequence and octopine T-DNA left border sequence, between which are the promoter, 5' UTR and first intron from the *Arabidopsis act7* gene driving the expression of a CP4 EPSP synthase gene. This CP4 EPSP synthase gene comprises a

CTP, linked to a synthetic EPSP synthase coding region and the 3' UTR from the pea rbsc E9 gene; the P-Bn.SW3-0:1:2 promoter operably linked to the *Arabidopsis thaliana* isocitrate lyase gene (as described in PCT Application WO 03/79766) and the napin 3' UTR. The nucleic acid sequence was determined using known methodology
5 and confirmed the integrity of the cloning junctions. The resulting plasmid was named pMON82378 was used for the transformation of soybean.

Example 7

This example sets forth the transformation of soybean plants with the
10 constructs described in Example 5.

Soybean plants were transformed using an *Agrobacterium*-mediated transformation method, as described by Martinell (U.S. Patent 6,384,301). For this method, overnight cultures of *Agrobacterium tumefaciens* containing the plasmid that includes a gene of interest, such as pMON75201, were grown to log phase and then
15 diluted to a final optical density of 0.3 to 0.6 using standard methods known to one skilled in the art. These cultures were used to inoculate the soybean embryo explants prepared as described below. Briefly, the method is a direct germline transformation into individual soybean cells in the meristem of an excised soybean embryo. The soybean embryo is removed after surface sterilization and germination of the seed.
20 The explants are then plated on OR media, a standard MS medium as modified by Barwale *et al.* (1986), plus 3 mg/L BAP, 200 mg/L carbenicillin, 62.5 mg/L cefotaxime, and 60 mg/L benomyl, and stored at 15°C overnight in the dark. The following day the explants are wounded with a scalpel blade and inoculated with the *Agrobacterium* culture prepared as described above. The inoculated explants are then
25 cultured for 3 days at room temperature. Following the post-transformation culture, the meristematic region is then cultured on standard plant tissue culture media in the presence of the herbicide glyphosate (Monsanto Company, St. Louis, MO), which acts as both a selection agent and a shoot inducing hormone. Media compositions and culture lengths are detailed in the aforementioned Martinell patent. After 5 to 6
30 weeks, the surviving explants that have a positive phenotype are transferred to soil and grown under greenhouse conditions until maturity.

The mature seeds of the transformed plants are harvested and analyzed for oil content using standard methodology, for example by near-infrared reflectance (NIR)

2005289727 16 Jul 2007

spectroscopy as described in Williams and Norris (1987). The seeds from plants expressing the *Arabidopsis thaliana* isocitrate lyase gene, driven by the P-Bn.SW3-0:1:2 promoter, have a higher oil content as compared to seeds from non-transformed plants.

With reference to the use of the word(s) "comprise" or "comprises" or "comprising" in the foregoing description and/or in the following claims, unless the context requires otherwise, those words are used on the basis and clear understanding that they are to be interpreted inclusively, rather than exclusively, and that each of those words is to be so interpreted in construing the foregoing description and/or the following claims.

REFERENCES

The references listed below are incorporated herein by reference to the extent that they supplement, explain, provide a background for, or teach methodology, techniques, and/or compositions employed herein.

- U.S. Patent Appl. 2002/0192812
U.S. Patent Appl. 2003/0028917
U.S. Patent 4,810,648
10 U.S. Patent 4,965,188
U.S. Patent 4,990,607
U.S. Patent 5,015,580
U.S. Patent 5,094,945
U.S. Patent 5,097,025
15 U.S. Patent 5,110,732
U.S. Patent 5,164,316
U.S. Patent 5,176,995
U.S. Patent 5,196,525
U.S. Patent 5,229,114
20 U.S. Patent 5,304,730
U.S. Patent 5,322,938
U.S. Patent 5,352,605
U.S. Patent 5,359,142
U.S. Patent 5,362,865
25 U.S. Patent 5,378,619
U.S. Patent 5,384,253
U.S. Patent 5,391,725
U.S. Patent 5,424,200
U.S. Patent 5,428,147
30 U.S. Patent 5,447,858
U.S. Patent 5,463,175
U.S. Patent 5,491,084

- U.S. Patent 5,512,466
- U.S. Patent 5,516,671
- U.S. Patent 5,516,671
- U.S. Patent 5,538,880
- 5 U.S. Patent 5,543,576
- U.S. Patent 5,550,318
- U.S. Patent 5,589,583
- U.S. Patent 5,591,616
- U.S. Patent 5,599,670
- 10 U.S. Patent 5,608,144
- U.S. Patent 5,608,149
- U.S. Patent 5,614,399
- U.S. Patent 5,627,061
- U.S. Patent 5,633,435
- 15 U.S. Patent 5,633,441
- U.S. Patent 5,635,055
- U.S. Patent 5,659,122
- U.S. Patent 5,689,041
- U.S. Patent 5,716,837
- 20 U.S. Patent 5,720,871
- U.S. Patent 5,750,876
- U.S. Patent 5,773,696
- U.S. Patent 5,824,877
- U.S. Patent 5,898,096
- 25 U.S. Patent 5,942,664
- U.S. Patent 5,958,745
- U.S. Patent 5,981,840
- U.S. Patent 5,985,605
- U.S. Patent 5,998,700
- 30 U.S. Patent 6,011,199
- U.S. Patent 6,013,864
- U.S. Patent 6,040,497
- U.S. Patent 6,063,597

- U.S. Patent 6,063,756
U.S. Patent 6,072,103
U.S. Patent 6,080,560
U.S. Patent 6,093,695
5 U.S. Patent 6,096,950
U.S. Patent 6,110,464
U.S. Patent 6,121,436
U.S. Patent 6,160,208
U.S. Patent 6,166,292
10 U.S. Patent 6,171,640
U.S. Patent 6,228,992
U.S. Patent 6,232,526
U.S. Patent 6,316,407
U.S. Patent 6,380,466
15 U.S. Patent 6,384,301
U.S. Patent 6,384,301
U.S. Patent 6,399,861
U.S. Patent 6,403,865
U.S. Patent 6,444,876
20 U.S. Patent 6,448,387
U.S. Patent 6,476,295
U.S. Patent 6,476,295
U.S. Patent 6,476,295
U.S. Patent 6,506,565
25 U.S. Patent 6,506,962
U.S. Patent 6,531,648
U.S. Patent 6,537,750

Barwale *et al.*, *Plants*, 167:473-481, 1986.

- 30 Bechtold *et al.*, *C.R. Acad. Sci. Life Sciences*, 316:1194-1199, 1993.

Bent *et al.*, *Science*, 265:1856-1860, 1994.

DeBlock *et al.*, *EMBO J.*, 6:2513-2519, 1987.

Misawa *et al.*, *Plant J.*, 4:833-840, 1993.

- Misawa *et al.*, *Plant J.*, 6:481-489, 1994.
- Murashige and Skoog, *Physiol. Plant*, 15:473-497, 1962.
- Orman *et al.*, *JAOCS*, 69(10):1036-1038, 1992.
- Patrick *et al.*, *JAOCS*, 74(3):273-276, 1997.
- 5 PCT Appln. WO 03/79766
- Rubel, *JAOCS*, 71:1057-1062, 1994.
- Sambrook *et al.*, In: *Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2000.
- Sathasiivan *et al.*, *Nucl. Acids Res.*, 18:2188-2193, 1990.
- 10 Tiwari *et al.*, *JAOCS*, 51:104-109, 1974.
- Williams and Norris, In: *Near-infrared Technology in the Agricultural and Food Industries*, Amer. Assoc. Cereal Chemists, Inc., St. Paul, MN, 1987.

The claims defining the invention are as follows:

1. An isolated promoter comprising a polynucleotide sequence selected from the group consisting of
 - a) a polynucleotide sequence comprising the nucleic acid sequence of SEQ ID NO:9;
 - b) a polynucleotide sequence having at least about 83% sequence identity to SEQ ID NO:9, and that is capable of regulating transcription of an operably linked transcribable polynucleotide molecule; and
 - c) a fragment of the polynucleotide sequence of a) capable of regulating transcription of an operably linked transcribable polynucleotide molecule.

2. The isolated promoter of claim 1 comprising a polynucleotide sequence selected from the group consisting of
 - a) a polynucleotide sequence comprising the nucleic acid sequence of SEQ ID NO:9; and
 - b) a fragment of the polynucleotide sequence of a) capable of regulating transcription of an operably linked transcribable polynucleotide molecule.

3. A construct comprising the promoter of claim 1 or 2 operably linked to a heterologous transcribable polynucleotide molecule.

4. The construct of claim 3, wherein the transcribable polynucleotide molecule is operably linked to a 3' transcription termination polynucleotide molecule.

5. The construct of claim 3 or 4, wherein said transcribable polynucleotide molecule is a gene of agronomic interest.

6. The construct of claim 3 or 4, wherein said transcribable polynucleotide molecule is a marker gene.

7. A transgenic plant stably transformed with the construct of any one of claims 3 to 6.

8. The transgenic plant of claim 7, wherein the transgenic plant is a dicotyledonous plant.

9. The transgenic plant of claim 8, wherein the dicotyledonous plant is selected from the group consisting of tobacco, tomato, potato, soybean, cotton, canola, sunflower, and alfalfa.

10. The transgenic plant of any one of claims 7 to 9, wherein the transcribable polynucleotide molecule confers altered cell proliferation in the seed of said transgenic plant.

11. The transgenic plant of any one of claims 7 to 9, wherein the transcribable polynucleotide molecule confers altered oil content in the seed of said transgenic plant.

12. The transgenic plant of any one of claims 7 to 9, wherein the transcribable polynucleotide molecule confers altered protein quality in the seed of said transgenic plant.

13. The transgenic plant of any one of claims 7 to 9, wherein said transcribable polynucleotide molecule confers altered micronutrient content to said transgenic plant.

14. A seed of the transgenic plant of any one of claims 7 to 9, wherein the seed comprises said construct.

15. Oil of the transgenic plant of any one of claims 7 to 9 or 11, wherein the oil comprises a detectable nucleic acid comprising the promoter of claim 1 or 2.

16. Meal of the transgenic plant of any one of claims 7 to 13, wherein the meal comprises a detectable nucleic acid comprising the promoter of claim 1 or 2.

17. A transformed plant cell having stably incorporated into its genome the construct of any one of claims 3 to 6.

18. A method of making a vegetable oil, comprising the steps of:
- a) obtaining the seed of claim 14; and
 - b) extracting oil from the seed.
19. A method of making a vegetable protein, comprising the steps of:
- a) obtaining the seed of claim 14; and
 - b) extracting the protein from the seed.
20. A method of directing expression of a transcribable polynucleotide sequence in a plant cell comprising operably linking the promoter of claim 1 or 2 to the polynucleotide sequence and transforming a plant cell with the promoter operably linked to the polynucleotide sequence.
21. The method of claim 20, comprising regenerating a plant from the plant cell.
22. A method of preparing food or feed comprising
- a) obtaining the transgenic plant of any one of claims 7 to 13; and
 - b) preparing food or feed from said plant or a part thereof.

SEQUENCE LISTING

<110> RUEZINSKY, DIANE M.
 HEWITT, CRYSTAL L.
 ZHENG, WEI

<120> PROMOTER MOLECULES FOR USE IN PLANTS

<130> MONS:057WO

<140> UNKNOWN
 <141> 2005-09-23

<150> 60/613,081
 <151> 2004-09-24

<160> 22

<170> PatentIn version 3.2

<210> 1
 <211> 1683
 <212> DNA
 <213> Brassica napus

<400> 1
 ctgttatatg ttattactta tgaggaactg gtattacata tagagaaaaa tatagatcga 60
 gaacacaaa ttatcataaa tatggaactc tgtggttatt aaaaagtgtt gttatacgat 120
 tttatgttct tttaaaagtg tttaatgggg ttgattgtat gatcacaata tatgttatac 180
 atgatcgtgc gttcacccac totgatatga ttgttcattg ggagttttgt atagaattaa 240
 gtgaaattat tagaatttac aataagatca ctaagcagat attttcataa tttttttta 300
 taaaccctat cggctgatcc gatcggtccc gggaggaacg cgtttagtga aatattttca 360
 taatattttt catgacatca actgtgagtg tcacttagaa tctcgaatat aatocaaatc 420
 cgacataaga tatgattctc ttattcttgc attcaaccga ataactctgaa gtgctccaat 480
 cgcaaaactg tttctctgcc catgacctaa aaatggctct ttataataatg cgcataatgat 540
 taatccgcca ccaaagtgtc ttgtgaccac ttatttactt atttatatgc tgataactct 600
 gtgttatgtg taaagtact aattagtta atatttataa gaaagagtaa gattctcatg 660
 ataaacattt aaattagtat ctccagtatg attgtattca ataagtatta gctgaattaa 720
 gctagttagt agtatatcat aattttgaca aatttaacgt aacaattttc ataactattt 780
 actgctgtta gaaacctgtg ttataabbat aactttcctt tttgggtcaa aattacaact 840
 ttotatacat aaaagagata acttaagga aaggcagttg acttgagaat ataaagtcaa 900
 ttggagagca atttgcagga gaaaagtcac aaaagtcgta cattcggcac ttgtaattaa 960
 caagagaatc atcggaaaat ggcatctttg aaaataatat tggtaattta cgtaatacat 1020
 tattggttaa taatttttta aaaaatataa aatttatatt ggtttttaag ccataatattg 1080

tcttttggtta ttaagccact acggttaaga tatgccaaagt tactttatta agatgcatat 1140
 atactcactt ctattacata tatatggtaa tatatatggt totaaaacat agtattatat 1200
 atgcaaacaa agtagaatat tagtcaattt taaaactagt tgcaatttaa aaaaaaaaaa 1260
 ttagacaact ttttacaat gaaaccaatg ggacagggtg aggaatagtg tcttcaatat 1320
 gtgtatagat tagaaaatct tttgaggata tagtcatttg agttagcaa acataaccat 1380
 gacatatgoa tatgatgacc cacaagtgca tactggcagc atatacccg ttagacacct 1440
 acatataact ttgaactaal aalatattaa atgtatatga aaaaccttcg aatgatgaat 1500
 gattaaattt taaaccaaac tatgatctgg cattggctaa tagcttcaca cttcgttggc 1560
 acgtctattt aaagcttcac aaagacgca acttaaatca caaccogaaa acgaaaaata 1620
 tcacaagaaa agaaaaaaaa accatttatt cac 1653

<210> 2
 <211> 27
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 2
 caccatgctt tcacgtcttt tgagttg 27

<210> 3
 <211> 22
 <212> DNA
 <213> artificial

<220>
 <223> primer sequence

<400> 3
 gtaatacgcac tcactatagg gc 22

<210> 4
 <211> 30
 <212> DNA
 <213> artificial

<220>
 <223> primer sequence

<400> 4
 cccacagatt ttctatttcg gttctcatac 30

<210> 5
 <211> 19
 <212> DNA

```

<213> Artificial
<220>
<223> primer sequence

<400> 5
actatagggc acgogtggt 19

<210> 6
<211> 23
<212> DNA
<213> Artificial

<220>
<223> primer sequence

<400> 6
gagaaacccat ggtgaataaa tgg 23

<210> 7
<211> 1288
<212> DNA
<213> Brassica napus

<400> 7
gggctgggtga ttatgattcg atgattgat ccagttagtt aattttgtcg aatcattttc 60
ttotttcttc gtttaaacat ttaacttgca cgaatgggtc tcttgtgatt gaacggaatc 120
tttgaatcaa attaattaat tgataagaag acaaaataag attccatggt ttgataagat 180
taatgatatt gatgcattaa cggcatatct gaaacaaatt cattaatcag caaaaactgag 240
aattttgaac tacgaaatc cagtttctag ttgaaaaat aatgatcgtg gagaacatat 300
tataaaacct cagaaacgaa ctatgtagta gaaattggtt aactgtatct agcaaacaaa 360
aaagcgttaa ctgttaccat aaccattat tttaatttct aaattatgca aattgcaagc 420
ttccaatctt agccttttaa ataaaaaga cgctatggac ttcgaatcag attgtgaaca 480
acttgtggaa ctaattaaca gagaagaaac tgaccagcga tggcagtggg gctagatgag 540
attaaagcct atgcaacaag atttatggag ttttcgatat tgtttatcct tagagctcta 600
aacgtccgtg cagatggaga tgctctcgg aggatggctc gtctgttccc acttgatcta 660
gcctagatgg atagacaaca gaaaaaaaa aaaaatgtcc gtgcacatgg caggcgtctag 720
atcatgagtc ctcaggttcc cctctgtaaa cgggtgttca cccgatggct agtacctaata 780
actggcata actcatacgg atgcagacca gtaagctgga ttgatttttt tcgatgtcaa 840
aaaaaaaaaga cgctatgatc ctaataaaaa cagggtaaac atacataaat gatataagtt 900
tagtagctca aaattgctag tttccaaca aatgttttct gatcctatct tcaggttttt 960
gtaacaaata taatttttaa atgtttccct taatatttgt ttttcaattt tatatcaata 1020

```

aatatatttg tccaaatacc aatcagatc caagttcgg atcgttaagc cttgggagat 1080
 taaagtetaa taggttcgaa ccatgatgag ttctagttat ctacatagat gtataagatg 1140
 atcattactt gtaaaactaa aatactttgt cataatcact tcatagaaca ttaccagtag 1200
 tatcacactt ttttactat aaaacccac tgcgaaacc ttttgagta atcaaaactag 1260
 tatctaaatc cttcaacttt tctaagac 1288

<210> 8
 <211> 848
 <212> DNA
 <213> Brassica napus

<400> 8
 gggctggctc gaaggacaag gatgatcgga tatctgcgtt ggaggctcag atggcagctc 60
 aacagcggg ctatgagacc cagaagaggc tgaacgagca gatgatggac atgatgaaga 120
 ggatgtaccg aacgagacgt tcccgaacat tcaagaccg tagttttttt ttttccaaaa 180
 actctgaaatg ttttatttaa atttgaatat tatctactat gttttatttt atgtttttatt 240
 carttttaat ttttaattta aaaatttta tttttttcca aaaaaaata atttttaaaa 300
 aaaataaatt ttttaaaaa ataattttt tcaaaattcc gagggaatgg aggtcctcgg 360
 ataattccga ggaactttta tctcggtaa attccgacga rcattaagtt cctcggaaat 420
 gtctgagggg aaggattcct cggaaatttc cgagaaactc aattccctcg gaaaattccg 480
 agraatattt cgtcgaaact tccgaggatg gaccatcgga attcctcgg tattttccga 540
 ggaaccttcc gacgaacatt gtgcctcgg agtttctcgg gaattttggt tctcggaaat 600
 tccgtcggaa aattccgacg gaattccgag gaatttgaat tcccgaggag ttatttccga 660
 ggactttttt cgtcggatg tctcggaaat aacgttattc cgacgacata cgcagcattt 720
 tttccctcag tatcccgatg ttttcttga gtgtttatgc attatataaa cccatcgatt 780
 caatcagctc catcatccaa ttaccttaa acttaaacct ctggtttctc aaaaaatatt 840
 ggctttac 848

<210> 9
 <211> 874
 <212> DNA
 <213> Brassica napus

<400> 9
 cctcgaaga ttctcagc acagaccat ctcttttagg gttatttgag atgcccgtat 60
 aaaagaggaa aaggtgggtg acttgaagag gacggcaaca acatcagatt glggagagca 120
 ggtgaaattg attgtcttgg ttaaaagatc gtcgtagcat acctgagcat acatcaatca 180
 agttataaga actagatata tagttcatag ctctattatt cgggtgtaga gtgattagag 240

ctttaacttct cttgattact ctttgactt gagatagtg attgaaagcg tagttcttcc 300
 ccagacgtac cgcaaagga actgggtaac caagccttg tgtcttgtct cctatacaac 360
 aaacaaacaa attgatcaaa gtttaattgat caagcattag ctaaacaaat cttagtaagg 420
 aactaagggt taagacagta atacaatcta agaccttaca atatattgga gaatttatcc 480
 tggtttttta ataactatgc aaatttaagt cgttactaat tacgtaccaa caccaaaatta 540
 tcccattata ctgtagcaaa aatgatgtgt aagtgtatat aaatacttgt gtaatcccat 600
 ccttaatgat taatlaatta ttacaatata gaatccttga cttactcaa acacgttccac 660
 gatggcgcca totgatgcaa cacaaagata taatacttac acgtatata acacacacat 720
 aagtacatat atatacgtgt aaaacatgag attctatttg cgagtctttg tctatataag 780
 ttccaaacttt tctcacagtt gtgtcttctt ttctacctc gccgctcttc aaccaaata 840
 atccctaact tttgttgcca caacttttcc tttc 874

<210> 10
 <211> 29
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 10
 ttcataatctg cgtaagtacg tccatgttc 29

<210> 11
 <211> 30
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 11
 cgagaaacag aatgatgaaa ctagagagac 30

<210> 12
 <211> 27
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 12
 caccatgctt tcacgtcttt tgagttg 27

<210> 13

<211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> primer sequence

 <400> 13
 tcgatgccac gaaaatgcat aagaac 26

 <210> 14
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> primer sequence

 <400> 14
 cctcatcctc atcctcatcc tcaacc 26

 <210> 15
 <211> 29
 <212> DNA
 <213> Artificial

 <220>
 <223> primer sequence

 <400> 15
 ccgtacacgg taaaatagtt ottgacgga 29

 <210> 16
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> primer sequence

 <400> 16
 ctaaagccat ggtottagaa aagttg 26

 <210> 17
 <211> 24
 <212> DNA
 <213> Artificial

 <220>
 <223> primer sequence

 <400> 17
 aatacgactc actatagggc acgc 24

 <210> 18
 <211> 25

<212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 18
 gaaaagccat ggtaaggcca atatt 25

<210> 19
 <211> 25
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 19
 cttctgccat ggaaagaaaa agttg 25

<210> 20
 <211> 28
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 20
 cccgggctgg tctgcgaag attctcag 28

<210> 21
 <211> 53
 <212> DNA
 <213> Artificial

<220>
 <223> primer sequence

<400> 21
 ctcgagcggc cgcttctatg tgcactggaa agaaaaagtt gtgccacaa aag 53

<210> 22
 <211> 891
 <212> DNA
 <213> Brassica napus

<400> 22
 gggctggtcc tgcgaagatt ctcagcacca agaccatct ctttaggggt tatttgagat 60
 gcccgataaa aagaggaaaa ggtgggtgac ttgaagagga cggcaacaac atcagattgt 120
 ggagagcagg tgaagttga ttgtcttggg taaaagatcg tegttagcata cctgagcata 180
 catcaatcaa gttataagaa ctagatatat agttcatagc tctattatct ggtgtagaag 240
 tgattagagc ttacttctc ttgtattaact ctttgtactt gagatagtgg attgaaagcg 300

```
tagttcttcc ccagacgtac ccgaaagggg actgggtaac caaagctttg tgtcttgtct 350
cctatacaac aaacaaacaa attgtatcaa ggtaattga tcaagcatta getaaacaaa 420
tctagctaag gaactaaggt ttaagacagt aatacaatct aagacctac aatataatgg 480
agaatttatt ctggtttttt aataatcatg caaatttaaa gtcttacta attacgtacc 540
aacaccaaat tatccocatta tactgtagca aaaatgatgt gtaagtgtat ataaatactt 600
gtgtaatccc atccttaatg attaattaat tattacaata tcgaatccct tgaattcact 660
caaacacggt cacgatggcg tcactgatg caacacaaga tattaatact tacacgtata 720
tatacacaca cataagtaca tatatatacg tgtaaacat gagattctat ttgcgaagtc 780
tttgtctata taagttcaa cttttctcac agttgtgtct tcttttcta cctcgccgct 840
cttcaaccaa aataatccct aacttttgtt ggcacaactt tttctttcca g 891
```