

A&A Ref: 145250

PUBLICATION PARTICULARS AND ABSTRACT
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

2/1085

7 February 2002

7.5.2003

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

C07D A61K A61P

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

Merck Patent GmbH

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

BATHE, Andreas
BÖTTCHER, Henning

HELFERT, Bernd

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
	33 DE	31 199 32 314.3	32 10 July 1999

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------


Benzofurane derivatives

57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	42
------------------	----

The sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.




(57) Abstract: Benzofurane derivatives of formula (I), wherein R and R' have the meanings given in claim no. (1), and salts thereof are suitable for use as intermediate products in the synthesis of medicaments.

(57) Zusammenfassung: Benzofuranerivate der Formel (I) sowie deren Salze, worin R und R' die in Anspruch (1) angegebenen Bedeutungen haben, eignen sich als Zwischenprodukte bei der Synthese von Arzneimitteln.

Benzofuran derivatives

The invention relates to benzofuran derivatives of the formula I

5



in which

- R is 1-piperazinyl, 4-R¹-piperazinyl or L,
- 10 R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-ylcarbonyl, N,N-di(tert-butyloxy-carbonyl)aminocarbonyl, -CH=C(R⁵R⁶), benzofuran-2-yl-C≡C-, -C(Hal)₃, -CO-C(Hal)₃, 1,4-dihydrobenzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,
- 15 L is Cl, Br, I or a free or reactive functionally modified OH group,
- R¹, R⁴ in each case independently of one another are H, benzyl or another amino protective group,
- 20 R², R³ in each independently of one another are H or alkyl having 1-6 C atoms,
- R⁵, R⁶ in each case independently of one another are alkyl having 1-6 C atoms,
- Hal is F, Cl, Br or I,
- 25 and their salts.

Similar compounds are disclosed in DE 43 33 254 and DE 195 14 567.

30 The invention was based on the object of finding novel compounds which can be used, in particular, as intermediates in the synthesis of medicaments, but can also be used directly for the production of medicaments.

35

It has been found that the compounds of the formula I

and their salts are important intermediates for the production of medicaments and at the same time have pharmacological properties. Thus, they show, for example, effects on the central nervous system.

5

The invention relates to the benzofuran derivatives of the formula I and their salts.

10 Above and below, the radicals R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R , R' , L , Q and Q' have the meanings indicated in the formulae I to V, if not expressly stated otherwise.

In the above formulae, A has 1 to 4, preferably 1, 2 or 3, C atoms. A is preferably methyl or ethyl, furthermore propyl or isopropyl, and additionally also
15 butyl, isobutyl, sec-butyl or tert-butyl.

The radical Ph is phenyl.

In the compounds of the formula [sic] I, II, V, VI and VII, L , Q and Q' are preferably Cl , Br , I or a reactive
20 modified OH group such as, for example, an activated ester, an imidazolidine or alkylsulfonyloxy having 1-6 C atoms (preferably methylsulfonyloxy) or arylsulfonyloxy having 6-10 C atoms (preferably phenyl- or p-tolylsulfonyloxy).

25

The expression "amino protective group" is generally known and relates to groups which are suitable for protecting (for blocking) an amino group from chemical reactions, but which are easily removable after the
30 desired chemical reaction has been carried out at other positions in the molecule. Typical of such groups are, in particular, unsubstituted acyl, aryl, aralkoxymethyl or aralkyl groups. As the amino protective groups are removed after the desired reaction (or reaction
35 sequence), their nature and size is otherwise uncritical; preferred groups, however, are those having 1-20, in particular 1-8 C atoms. The expression "acyl group" is to be interpreted in the widest sense in connection with the present process and the present

compounds. It includes acyl groups derived from aliphatic, araliphatic, aromatic or heterocyclic carboxylic acids or sulfonic acids and also, in particular, alkoxy carbonyl, aryloxy carbonyl and especially aralkoxy carbonyl groups. Examples of acyl groups of this type are alkanoyl such as acetyl, propionyl, butyryl; aralkanoyl such as phenylacetyl; aroyl such as benzoyl or toluyl; aryloxyalkanoyl such as phenoxyacetyl; alkoxy carbonyl such as methoxy carbonyl, ethoxy carbonyl, 2,2,2-trichloroethoxy carbonyl, BOC (tert-butoxy carbonyl), 2-iodoethoxy carbonyl; aralkoxy carbonyl such as CBZ (carbobenzoyl carbonyl, also called "Z"), 4-methoxybenzyloxy carbonyl, FMOC (9-fluorenylmethoxy carbonyl); arylsulfonyl such as Mtr (4-methoxy-2,3,6-trimethylphenylsulfonyl). Preferred amino protective groups are BOC and Mtr, and additionally CBZ or FMOC.

The compounds of the formula I can have one or more chiral centres and therefore occur in various stereoisomeric forms. The formula I includes all these forms.


The invention further relates to a process for the preparation of benzofuran derivatives of the formula I according to Claim 1 and of their salts, characterized in that

a) for the preparation of compounds of the formula I in which

R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl and

R' is 2-R²-5-R³-pyrrol-1-yl carbonyl, 4-R⁴-piperazin-1-yl carbonyl, 1,4-dihydrobenzo[d][1,2]-oxazin-3-yl carbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-yl carbonyl,

a compound of the formula II



II

in which

- 5 R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl
and
Q is Cl, Br, I or a free or reactive functionally
modified OH group,
and R¹ has the meaning indicated in Claim 1

10 is reacted with a compound of the formula III



in which

- 15 R' is 2-R²-5-R³-pyrrol-1-yl, 4-R⁴-piperazin-1-yl,
1,4-dihydrobenzo[d][1,2]oxazin-3-yl or 3,4-
dihydrobenzo-1H-phthalazin-2-yl,
and R², R³ and R⁴ have the meanings indicated in
Claim 1,


20

or

b) for the preparation of compounds of the formula I
in which

- 25 R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl
and
R' is -CH=C(R⁵R⁶), benzofuran-2-yl-C≡C-, -C(Hal)₃
or -CO-C(Hal)₃,
and R¹, R⁵ and R⁶ have the meanings indicated in
30 Claim 1,

i) a compound of the formula IV




IV

in which

R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl,
is reacted with a compound of the formula V

5



V

in which R' is -CH=C(R⁵R⁶), benzofuran-2-yl-C≡C-,
-C(Hal)₃ or -CO-C(Hal)₃,


and Q' is Cl, Br, I or a free or reactive functionally
10 modified OH group,

and R⁵ and R⁶ have the meanings indicated in Claim 1,

or

15

ii) a compound of the formula Va



Va

in which R and R' have the meanings indicated under i)
20 is cyclized,

or

c) a compound of the formula I,

25

in which R is a 1-piperazinyl radical, is converted by
introduction of an amino protective group into another
compound of the formula I in which R is the 4-R¹-
piperazinyl radical,

in which R¹ is an amino protective group,

30

or

d) a compound of the formula I,

in which R is a 4-R¹-piperazinyl group, in which R¹ is
35 benzyl or another amino protective group,

is converted by removal of the benzyl or amino protective group into a compound of the formula I in which R¹ is 1-piperazinyl,

5 or

e) in a compound of the formula I a radical R is converted into another radical R

10 by, for example,

i) replacing a Br atom by OH,

ii) esterifying an OH group or

iii) replacing a Br atom by a 4-R¹-piperazinyl group, in which R¹ is benzyl or an amino protective group,

15

and/or a base of the formula I is converted into one of its salts by treatment with an acid.

The compounds of the formula I and also the starting
20 substances for their preparation are otherwise prepared by methods known per se, such as are described in the literature (e.g. in the standard works such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart),
25 namely under reaction conditions which are known and suitable for the reactions mentioned. In this case, use can also be made of variants which are known per se but not mentioned here in greater detail.

30 If desired, the starting substances can also be formed in situ such that they are not isolated from the reaction mixture, but immediately reacted further to give the compounds of the formula I.

35 In the compounds of the formula II, the radical Q is preferably Cl or Br; however, it can also be I, OH or a reactive modified OH group such as alkylsulfonyloxy having 1-6 C atoms (preferably methylsulfonyloxy) or arylsulfonyloxy having 6-10 C atoms (preferably phenyl-

or p-tolylsulfonyloxy, 1- or 2-naphthalenesulfonyloxy). In the compounds of the formula II, the radical R is preferably Br or 4-benzylpiperazinyl.

5 The compounds of the formula II are known in some cases; the unknown compounds can easily be prepared analogously to the known compounds.

10 The reaction of the compounds of the formula II with compounds of the formula III proceeds according to methods such as are known from the literature for the alkylation of amines. The components can be fused with one another without a solvent being present, if appropriate in a closed tube or in an autoclave.

15 However, it is also possible to react the compounds in the presence of an inert solvent.

Suitable inert solvents are, for example, hydrocarbons such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons such as trichloroethylene, 1,2-dichloroethane, carbon tetrachloride, 20 chloroform or dichloromethane; alcohols such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers such as ethylene glycol monomethyl or 25 monoethyl ether (methyl glycol or ethyl glycol), ethylene glycol dimethyl ether (diglyme); ketones such as acetone or butanone; amides such as acetamide, dimethylacetamide or dimethylformamide (DMF); nitriles such as acetonitrile; sulfoxides such as dimethyl 30 sulfoxide (DMSO); carbon disulfide; nitro compounds such as nitromethane or nitrobenzene; esters such as ethyl acetate, and optionally also mixtures of the solvents mentioned with one another or mixtures with water.

35

The addition of an acid-binding agent, for example of an alkali metal or alkaline earth metal hydroxide, carbonate or bicarbonate or of another salt of a weak acid of the alkali metals or alkaline earth metals,

preferably of potassium, sodium or calcium, or the addition of an organic base such as triethylamine, dimethylamine, pyridine or quinoline or of an excess of the amine component can be favourable. Depending on the conditions used, the reaction time can be between a few minutes and 14 days, and the reaction temperature between 0 and 150°, normally between 20 and 130°C.

In the compounds of the formula V, the radical Q' is preferably Cl or Br; however, it can also be I, OH or a reactive modified OH group such as alkylsulfonyloxy having 1-6 C atoms (preferably methylsulfonyloxy) or arylsulfonyloxy having 6-10 C atoms (preferably phenyl- or p-tolylsulfonyloxy, or 1- or 2-naphthalene-sulphonyloxy).

In the compounds of the formula IV, the radical R is preferably Br or 4-benzylpiperazinyl.

The reaction of the compounds of the formula IV with compounds of the formula V proceeds according to methods such as are known from the literature for the alkylation of phenols.

The compounds of the formula VI are known in some cases; the unknown compounds can easily be prepared analogously to the known compounds. The cyclization is carried out according to generally known methods.

The removal of an amino protective group from a compound of the formula I - depending on the protective group used - is carried out, for example, using strong acids, expediently using TFA (trifluoroacetic acid) or perchloric acid, but also using other strong inorganic acids such as hydrochloric acid or sulfuric acid, strong organic carboxylic acids such as trichloroacetic acid or sulfonic acids such as benzene- or p-toluenesulfonic acid. The presence of an additional inert solvent is possible, but not always necessary.

Suitable inert solvents are preferably organic solvents, for example carboxylic acids such as acetic acid, ethers such as tetrahydrofuran or dioxane, amides such as dimethylformamide, halogenated hydrocarbons such as dichloromethane, in addition also alcohols such as methanol, ethanol or isopropanol, and water. In addition, mixtures of the abovementioned solvents are possible. TFA is preferably used in an excess without addition of a further solvent, perchloric acid in the form of a mixture of acetic acid and 70% perchloric acid in the ratio 9:1. The reaction temperatures are expediently between approximately 0 and approximately 50°; the reaction is preferably carried out between 15 and 30°.

The group BOC is preferably removed using TFA in dichloromethane or using approximately 3 to 5 N hydrochloric acid in dioxane at 15-30°.

Hydrogenolytically removable protective groups (e.g. CBZ or benzyl) can be removed, for example, by treating with hydrogen in the presence of a catalyst (e.g. of a noble metal catalyst such as palladium, expediently on a support such as carbon). Suitable solvents here are those indicated above, in particular, for example, alcohols such as methanol or ethanol or amides such as DMF. The hydrogenolysis is generally carried out at temperatures between approximately 0 and 100° and pressures between approximately 1 and 200 bar, preferably at 20-30° and 1-10 bar.

Compounds of the formula I in which R' is N,N-di(tert-butylloxycarbonyl)aminocarbonyl are preferably obtained by reaction of the unprotected aminocarbonyl compound, in which

R is 4-R¹-piperazinyl or L,

L [lacuna] the meaning indicated in Claim 1 and

R¹ is benzyl or another amino protective group,

with (BOC)₂O in an inert solvent, such as, for example,

THF or dioxane with addition of a base, such as, for example, diethylamine and preferably of a catalytic amount of dimethylaminopyridine.

5 A base of the formula I can be converted into the associated acid addition salt using an acid, for example by reaction of equivalent amounts of the base and of the acid in an inert solvent such as ethanol and subsequent evaporation. Suitable acids for this
10 reaction are in particular those which give physiologically acceptable salts. Thus, inorganic acids can be used, e.g. sulfuric acid, nitric acid, hydrohalic acids such as hydrochloric acid or hydrobromic acid, phosphoric acids such as ortho-
15 phosphoric acid, sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic mono- or polybasic carboxylic, sulfonic or sulfuric acids, e.g. formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid,
20 malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, ethanedisulfonic acid, 2-hydroxy-
25 ethanesulfonic acid, benzenesulfonic acid, p-toluene-sulfonic acid, naphthalenemono- and -disulfonic acids, and laurylsulfuric acid. Salts with physiologically unacceptable acids, e.g. picrates, can be used for the isolation and/or purification of the compounds of the
30 formula I.

On the other hand, compounds of the formula I can be converted using bases (e.g. sodium or potassium hydroxide or carbonate) into the corresponding metal,
35 in particular alkali metal or alkaline earth metal salts, or into the corresponding ammonium salts.

The invention furthermore relates to the use of the compounds of the formula I as intermediates for the

synthesis of medicaments. Corresponding medicaments are described, for example, in DE 4333254.

5 The invention relates in particular to the use of the compounds of the formula I as intermediates for the synthesis of medicaments which exhibit actions on the central nervous system. 1-[4-(5-Cyanoindol-3-yl)butyl]-4-(2-carbamoylbenzofuran-5-yl)piperazine and its salts are very particularly preferably to be mentioned here.

10

The invention accordingly relates in particular to the use of the compounds of the formula I according to Claim 1


in which

15 R is Cl, Br, I or 4-R¹-piperazinyl,
R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-yl carbonyl, 1,4-dihydrobenzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,
20 R¹ is benzyl or another amino protective group,
R⁴ is H, benzyl or another amino protective group,
R², R³ in each case independently of one another are H or alkyl having 1-6 C atoms,

25 in the synthesis of 1-[4-(5-cyanoindol-3-yl)butyl]-4-(2-carbamoylbenzofuran-5-yl)piperazine and its salts, characterized in that

3-R-6-hydroxybenzaldehyde,
30 in which R is Cl, Br or I,

is reacted with a compound of the formula VI



VI


35

in which X is Cl, Br, I or a free or functionally modified OH group,

Q is OH or OR" and

R" is alkyl having 1-6 C atoms,

to give a compound of the formula VII



VII

5

in which

R is Cl, Br or I,

and Q has the meanings indicated,

10 in that, in the compound thus obtained, Q is converted into Cl, Br, I or a functionally modified OH group,

in that the compound thus obtained is reacted with a compound of the formula III

15

R'-H

III

in which

R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-
piperazin-1-ylcarbonyl, 1,4-dihydro-
benzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-
20 dihydrobenzo-1H-phthalazin-2-ylcarbonyl,
and R², R³ and R⁴ have the meanings indicated,

to give a compound of the formula I

in which

25

R is Cl, Br or I,


R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-
piperazin-1-ylcarbonyl, 1,4-dihydro-
benzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-
dihydrobenzo-1H-phthalazin-2-ylcarbonyl,

30

R⁴ is H, benzyl or another amino protective group,
R², R³ in each case independently of one another are H
or alkyl having 1-6 C atoms,

35 in that, in the compound of the formula I thus obtained, the radical R is converted into another radical R,

by reacting under transition metal catalysis with a compound of the formula VIII



in which

5 R¹ is benzyl or another amino protective group,

to give a compound of the formula I

in which

R is 4-R¹-piperazinyl,

10 R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-ylcarbonyl, 1,4-dihydrobenzo[d]-[1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,


R¹ is benzyl or another amino protective group,

15 R⁴ is H, benzyl or another amino protective group,

R², R³ in each case independently of one another are H or alkyl having 1-6 C atoms,

in that the compound thus obtained of the formula I

20 i) is first converted by basic hydrolysis into a compound of the formula IX and/or its acid addition salt




25

in which

R is 4-R¹-piperazinyl and

R¹ is benzyl or another amino protective group,

30 and then converted using ammonia into a compound of the formula X



in which


R is 4-R¹-piperazinyl and

R¹ is benzyl or another amino protective group,

5

or

ii) converted directly using ammonia into a compound of the formula X



10

in which

R is 4-R¹-piperazinyl and

R¹ is benzyl or another amino protective group,

15

in that the compound of the formula X thus obtained is converted into 5-(1-piperazinyl)benzofuran-2-carboxamide or an acid addition salt by removal of the amino protective group R¹, and

20 in that 5-(1-piperazinyl)benzofuran-2-carboxamide is reacted with 3-(4-chlorobutyl)-5-cyanoindole to give 1-[4-(5-cyanoindol-3-yl)butyl]-4-(2-carbamoylbenzofuran-5-yl)piperazine and

optionally converted into its acid addition salt.

25 3-(4-Chlorobutyl)-5-cyanoindole is disclosed in DE 4101686; 1-[4-(5-cyanoindol-3-yl)butyl]-4-(2-carbamoylbenzofuran-5-yl)piperazine is disclosed in DE 4333254.


30 Above and below, all temperatures are indicated in °C. In the following examples "customary working up" means: water is added, if necessary, the solution is adjusted, if necessary, to a pH between 2 and 10 depending on the constitution of the final product, and extracted with ethyl acetate or dichloromethane, the
35 organic phase is separated off, dried over sodium sulfate, evaporated and purified by chromatography on

silica gel and/or by crystallization. R_f values on silica gel.

Example 1

5

(2,5-Dimethylpyrrol-1-yl)-5-bromobenzofuran-2-yl-methanone




10 0.3 g of sodium hydride (60% suspension in paraffin
oil) is introduced into 10 ml of THF and 0.5 mL [sic]
of 2,5-dimethylpyrrole in 10 ml of THF is added
dropwise (5 minutes). After stirring at 50°C for 1
hour, a reddish suspension is present. 1.3 g of
15 5-bromobenzofuran-2-carbonyl chloride in 10 ml of THF
is [sic] added dropwise at 25°C (5 minutes) and the
mixture is subsequently stirred for 2 hours. The
addition of 100 ml of completely deionized water and
100 ml of ethyl acetate follows.

20 The separated organic phase was additionally washed 2 ×
with 100 ml of water and concentrated in vacuo. After
chromatography on silica gel (eluent heptane/ethyl
acetate 4:1), the residual oil gives 700 mg of yellow
crystals (yield 44%), m.p. 115-116°.

25

Example 2

5-Bromobenzofuran-2-carboxylic acid diethylamide




30

1.3 g of 5-bromobenzofuran-2-carbonyl chloride, 1 mL
[sic] of ethyldiisopropylamine and 30 mL [sic] of

toluene are mixed and 0.62 ml of diethylamine is added to the brown-coloured solution with stirring. A precipitate is formed with a slightly exothermic reaction. After 5 minutes, the mixture was treated with 30 ml of completely deionized water and the phases were separated. The organic phase was washed with 1.) 20 ml of 1N HCl 2.) 20 ml of 1N NaOH 3.) 20 ml of water and then with 20 ml of saturated NaCl solution and then freed of solvent components in vacuo. After chromatography on silica gel the residual, yellowish oil forms in the eluent MtB ether/heptane 2:1 [sic]. Final weight: 1.3 g of colourless crystals [sic] (yield: 88%), m.p. 79-81°.

15 Example 3

Di-tert-butyl N-(5-bromobenzofuran-2-carbonyl)imido-dicarbonate




20 12 g of 5-bromobenzofuran-2-carboxamide, 23.5 ml of BOC₂O, 600 mg of DMAP and 8 ml of triethylamine are introduced into 100 ml of THF at 20°C. A clear, orange solution is formed in 3 h in an endothermic reaction.

25 It is warmed to 25°C and treated with 100 ml of water and 100 ml of ethyl acetate. The organic phase is separated off, and washed twice with 100 ml of water and 100 ml of saturated sodium chloride solution. The organic phase is concentrated and forms a mixture of

30 oil and crystals (22 g/yield 40%). After crystallization of the crude product from 160 ml of ethanol, 9.0 g of yellow crystals are obtained, m.p. 138-139°.

Example 4

5-Bromobenzofuran-2-carboxylic acid dibenzylamide



5


A solution of 50 ml of toluene and 7.9 g of dibenzylamine are added dropwise with stirring at 50-60°C in the course of 10 min. to 5.2 g of 5-bromobenzofuran-2-carbonyl chloride in 100 ml of toluene. A colourless solid is obtained. After dropwise addition is complete, the mixture is additionally stirred at 100-110°C for a further 3 hours. After cooling to 10°C, the solid product (dibenzylammonium chloride) is filtered off with suction. The filtrate is then treated with a mixture of 150 ml of water and 10 g of sodium carbonate and thoroughly shaken. The organic phase is separated off, washed again with 100 ml of water, dried using sodium sulfate and filtered. The filtrate is concentrated on a Rotavapor in vacuo to a residue (residue: 9.5 g). After recrystallization from 100 ml of methanol, 6.5 g of product (yield 77%) remain after isolation
m.p. 114-115°.

25

Example 5

5-(4-Benzylpiperazin-1-yl)bromobenzofuran-2-carboxylic acid dibenzylamide

30




0.06 mg of Pd2DBA3 [sic] and 0.007 g of 2-dicyclohexyl-
phosphino-2'-dimethylaminobiphenyl in 40 ml of toluene
5 was stirred under nitrogen at 25°C for 20 min. 1.58 g
of 5-bromo-2,3-dihydrobenzofuran-2-carboxylic acid
dibenzylamide, 0.98 g of 1-benzylpiperazine and 1.43 g
of sodium tert-butylate are then added and the mixture
is stirred at 120°C for 2 hours. The cooled reaction
10 mixture is stirred into a mixture of 150 ml of water
and 5 ml of 37% hydrochloric acid with stirring [sic].
The reaction mixture is neutralized with 1.5 g of
sodium carbonate and the phase is [sic] extracted 3
times with 100 ml of ethyl acetate. The combined
15 organic phases are dried with 5 g of sodium sulfate and
the filtrate is concentrated in vacuo to give a
resinous residue (1.8 g of crude product). The crude
product is dissolved in 100 ml of ethyl acetate,
clarified with activated carbon and filtered. By
20 addition of 25 ml of 2-molar ethanolic hydrochloric
acid, the piperazine product is precipitated as the
hydrochloride, filtered, and the crystals are washed
with 10 ml of ethyl acetate and dried in vacuo at 40°C.
Final weight: 1.5 g/yield 53%, m.p. 196-198°.


25

Example 6

cis/trans-1,2-Di(5-bromobenzofuran)-2-ylethene



A) Preparation of the diether




5

The compound is obtained by reaction of 5-bromo-salicylaldehyde with $\text{Cl-CH}_2\text{-CH=CH-CH}_2\text{-Cl}$.

10 B) The dibenzofuran derivative is obtained by cyclization.


Example 7

15 2,2'-Ethyndiyl-bis-5-bromobenzofuran



A) Preparation of the diether

20




Reaction of 5-bromosalicylaldehyde with $\text{Cl-CH}_2\text{-CC-CH}_2\text{-Cl}$.

25 B) Cyclization of the diether.

Patent claims

1. Benzofuran derivatives of the formula I



in which

- R is 1-piperazinyl, 4-R¹-piperazinyl or L,
 R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-ylcarbonyl, N,N-di(tert-butylloxycarbonyl)aminocarbonyl, -CH=C(R⁵R⁶), benzofuran-2-yl-C≡C-, -C(Hal)₃, -CO-C(Hal)₃, 1,4-dihydrobenzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,
 L is Cl, Br, I or a free or reactive functionally modified OH group,
 R¹, R⁴ in each case independently of one another are H, benzyl or another amino protective group,
 R², R³ in each independently of one another are H or alkyl having 1-6 C atoms,
 R⁵, R⁶ in each case independently of one another are alkyl having 1-6 C atoms,
 Hal is F, Cl, Br or I,
 and their salts.

2. Compounds according to Claim 1

- a) (5-Bromobenzofuran-2-yl)-(2,5-dimethylpyrrol-yl)methanone
 b) (4-Benzylpiperazin-1-yl)-[5-(4-benzylpiperazin-1-yl)benzofuran-2-yl]methanone;
 c) [5-(4-Benzylpiperazin-1-yl)benzofuran-2-yl]-(1,4-dihydrobenzo[d][1,2]oxazin-3-yl)methanone;
 d) [5-(4-Benzylpiperazin-1-yl)benzofuran-2-yl]-(3,4-dihydro-1H-phthalazin-2-yl)methanone;

e) 5-Bromo-2-(2-methylpropenyl)benzofuran;
and their salts.


3. Process for the preparation of benzofuran derivatives of the formula I according to Claim 1 and of their salts, characterized in that

a) for the preparation of compounds of the formula I in which

R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl and

R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-ylcarbonyl, 1,4-dihydrobenzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,

a compound of the formula II



in which

R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl and

Q is Cl, Br, I or a free or reactive functionally modified OH group,

and R¹ has the meaning indicated in Claim 1

is reacted with a compound of the formula III



in which

R' is 2-R²-5-R³-pyrrol-1-yl, 4-R⁴-piperazin-1-yl, 1,4-dihydrobenzo[d][1,2]oxazin-3-yl or 3,4-dihydrobenzo-1H-phthalazin-2-yl,

and R², R³ and R⁴ have the meanings indicated in Claim 1,

or

b) for the preparation of compounds of the formula I


in which

R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl and

R' is -CH=C(R⁵R⁶), benzofuran-2-yl-C≡C-, -C(Hal)₃ or -CO-C(Hal)₃,

and R¹, R⁵ and R⁶ have the meanings indicated in Claim 1,

i) a compound of the formula IV



IV

in which

R is Cl, Br, I, 1-piperazinyl or 4-R¹-piperazinyl,

is reacted with a compound of the formula V




in which R' is -CH=C(R⁵R⁶), benzofuran-2-yl-C≡C-, -C(Hal)₃ or -CO-C(Hal)₃,

and Q' is Cl, Br, I or a free or reactive functionally modified OH group,

and R⁵ and R⁶ have the meanings indicated in Claim 1,

or

ii) a compound of the formula Va



Va

in which R and R' have the meanings indicated under i),

is cyclized,

or

c) a compound of the formula I,
in which R is a 1-piperazinyl radical, is converted by introduction of an amino protective group into another compound of the formula I in which R is the 4-R¹-piperazinyl radical, in which R¹ is an amino protective group,

or

d) a compound of the formula I,
in which R is a 4-R¹-piperazinyl group, in which R¹ is benzyl or another amino protective group, is converted by removal of the benzyl or amino protective group into a compound of the formula I in which R¹ is 1-piperazinyl,

or

e) in a compound of the formula I a radical R is converted into another radical R

by, for example,

- i) replacing a Br atom by OH,
- ii) esterifying an OH group or
- iii) replacing a Br atom by a 4-R¹-piperazinyl group, in which R¹ is benzyl or an amino protective group,

and/or a base of the formula I is converted into one of its salts by treatment with an acid.

4. Use of the compounds of the formula I according to

Claim 1 as intermediates for the synthesis of medicaments.

5. Use of the compounds of the formula I according to Claim 1 as intermediates for the synthesis of medicaments which show effects on the central nervous system.

6. Use of the compounds of the formula I according to Claim 1
in which

R is Cl, Br, I or 4-R¹-piperazinyl,

R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-yl carbonyl, 1,4-dihydrobenzo-[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,

R¹ is benzyl or another amino protective group,

R⁴ is H, benzyl or another amino protective group,

R², R³ in each case independently of one another are H or alkyl having 1-6 C atoms,


in the synthesis of

1-[4-(5-cyanoindol-3-yl)butyl]-4-(2-carbamoylbenzofuran-5-yl)piperazine and its salts, characterized in that

3-R-6-hydroxybenzaldehyde,

in which R is Cl, Br or I,

is reacted with a compound of the formula VI




in which X is Cl, Br, I or a free or functionally modified OH group,

Q is OH or OR" and

R" is alkyl having 1-6 C atoms,

to give a compound of the formula VII




in which

R is Cl, Br or I,

and Q has the meanings indicated,

in that, in the compound thus obtained, Q is converted into Cl, Br, I or a functionally modified OH group,

in that the compound thus obtained is reacted with a compound of the formula III



in which

R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-ylcarbonyl, 1,4-dihydrobenzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl, and R², R³ and R⁴ have the meanings indicated,

to give a compound of the formula I

in which

R is Cl, Br or I,


R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-piperazin-1-ylcarbonyl, 1,4-dihydrobenzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-dihydrobenzo-1H-phthalazin-2-ylcarbonyl,

R⁴ is H, benzyl or another amino protective group,

R², R³ in each case independently of one another are H or alkyl having 1-6 C atoms,

in that, in the compound of the formula I thus obtained, the radical R is converted into another

radical R,
by reacting under transition metal catalysis with
a compound of the formula VIII



in which

R¹ is benzyl or an amino protective group,

to give a compound of the formula I

in which

R is 4-R¹-piperazinyl,

R' is 2-R²-5-R³-pyrrol-1-ylcarbonyl, 4-R⁴-
piperazin-1-ylcarbonyl, 1,4-dihydro-
benzo[d][1,2]oxazin-3-ylcarbonyl or 3,4-
dihydrobenzo-1H-phthalazin-2-ylcarbonyl,


R¹ is benzyl or an amino protective group,

R⁴ is H, benzyl or another amino protective
group,

R², R³ in each case independently of one another
are H or alkyl having 1-6 C atoms,

in that the compound thus obtained of the
formula I

i) is first converted by basic hydrolysis into a
compound of the formula IX and/or its acid
addition salt




IX

in which

R is 4-R¹-piperazinyl and

R¹ is benzyl or another amino protective
group,

and then converted using ammonia into a compound
of the formula X




in which

R is 4-R¹-piperazinyl and

R¹ is benzyl or another amino protective group,

or

ii) converted directly using ammonia into a compound of the formula X



in which

R is 4-R¹-piperazinyl and

R¹ is benzyl or another amino protective group,

in that the compound of the formula X thus obtained is converted into 5-(1-piperazinyl)-benzofuran-2-carboxamide or an acid addition salt by removal of the protective group R¹ and

in that 5-(1-piperazinyl)benzofuran-2-carboxamide is reacted with 3-(4-chlorobutyl)-5-cyanoindole to give 1-[4-(5-cyanoindol-3-yl)butyl]-4-(2-carbamoylbenzofuran-5-yl)piperazine and optionally then converted into its acid addition salt.

7. A compound according to Claim 1, substantially as herein described and illustrated.
8. A process according to Claim 3, substantially as herein described and illustrated.
9. Use according to Claim 4, or Claim 5 or Claim 6, substantially as herein described and illustrated.
10. A new compound, a new process for the preparation of a compound, a new use of a compound of the formula I according to Claim 1 or of a salt thereof, substantially as herein described and illustrated.