




(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2022/05/06
(87) **Date publication PCT/PCT Publication Date:** 2022/11/10
(85) **Entrée phase nationale/National Entry:** 2023/10/24
(86) **N° demande PCT/PCT Application No.:** CA 2022/050712
(87) **N° publication PCT/PCT Publication No.:** 2022/232943
(30) **Priorité/Priority:** 2021/05/07 (US63/185,951)

(51) **Cl.Int./Int.Cl.** **C07D 213/79** (2006.01),
A61K 51/04 (2006.01), **A61K 51/08** (2006.01),
A61K 51/10 (2006.01), **C07K 14/00** (2006.01),
C07K 16/00 (2006.01)
(71) **Demandeur/Applicant:**
THE UNIVERSITY OF BRITISH COLUMBIA, CA
(72) **Inventeurs/Inventors:**
ORVIG, CHRIS, CA;
WHARTON, LUKE, CA
(74) **Agent:** OYEN WIGGS GREEN & MUTALA LLP

(54) **Titre :** CHELATEURS POUR RADIOMETAUX ET LEURS PROCEDES DE FABRICATION ET D'UTILISATION
(54) **Title:** CHELATORS FOR RADIOMETALS AND METHODS OF MAKING AND USING SAME



(57) **Abrégé/Abstract:**

A chelator having the general structure (I) wherein each R1 is independently OH, NH or SH, and X is O, S, or NR3, wherein R3 is H or CH2C(=O)R1. Methods of making and using the chelator, metal chelates, and biological constructs for delivering targeted radiation therapy using the chelator are provided. (I).



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

10 November 2022 (10.11.2022)



(10) International Publication Number

WO 2022/232943 A1

(51) International Patent Classification:

C07D 213/79 (2006.01) A61K 51/10 (2006.01)

A61K 51/04 (2006.01) C07K 14/00 (2006.01)

A61K 51/08 (2006.01) C07K 16/00 (2006.01)

(21) International Application Number:

PCT/CA2022/050712

(22) International Filing Date:

06 May 2022 (06.05.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/185,951 07 May 2021 (07.05.2021) US

(71) Applicant: THE UNIVERSITY OF BRITISH COLUM-

BIA [CA/CA]; University-Industry Liaison Office, 103 - 6190 Agronomy Road, Vancouver, British Columbia V6T 1Z3 (CA).

(72) Inventors: WHARTON, Luke; c/o University-Industry

Liaison Office, 103 - 6190 Agronomy Road, Vancouver, British Columbia V6T 1Z3 (CA). ORVIG, Chris; c/o University-Industry Liaison Office, 103 - 6190 Agronomy Road, Vancouver, British Columbia V6T 1Z3 (CA).

(74) Agent: MARLES, Jennifer A. et al.; 480 - 601 West Cor-

dova Street, Vancouver, British Columbia V6B 1G1 (CA).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH,

KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))
— in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(54) Title: CHELATORS FOR RADIOMETALS AND METHODS OF MAKING AND USING SAME



(57) Abstract: A chelator having the general structure (I) wherein each R₁ is independently OH, NH or SH, and X is O, S, or NR₃, wherein R₃ is H or CH₂C(=O)R₁. Methods of making and using the chelator, metal chelates, and biological constructs for delivering targeted radiation therapy using the chelator are provided. (I).

WO 2022/232943 A1

CHELATORS FOR RADIOMETALS AND METHODS OF MAKING AND USING SAME

Cross-Reference to Related Applications

5 [0001] This application claims priority to, and the benefit of, United States provisional patent application No. 63/185951 filed 7 May 2021, the entirety of which is incorporated by reference herein for all purposes.

Technical Field

10 [0002] Some embodiments relate to improved chelators. Some embodiments relate to improved biological targeting constructs incorporating chelators. Some embodiments relate to chelators coupled to a targeting moiety and capable of binding a radioactive isotope to provide targeted in vivo delivery of the radioactive isotope to a desired location within a mammalian subject.

15 Background


[0003] Radionuclides have potential utility in cancer diagnosis and therapy, particularly if they can be delivered selectively to a target location within the body of a subject. Targeted delivery of radionuclides can be achieved by using constructs that are engineered to both securely retain the radionuclide for in vivo delivery and deliver the radionuclide selectively to
20 a desired location within the body, with a reasonably low level of delivery to non-target regions of the body.

[0004] Targeting constructs have been developed that utilize a targeting moiety that targets a desired region of the body (e.g. a tumor-associated antigen) covalently coupled to a chelator to secure radionuclides for such purposes. The targeting moiety can be coupled to
25 the chelator via a linker. Such targeting constructs may be referred to as radioimmunoconjugates. The radioimmunoconjugate is used to chelate a desired radionuclide for in vivo delivery, for example to provide diagnostic imaging, targeted radionuclide therapy using the construct, or both (i.e. as a theranostic construct). In some cases, the chelator can also be used for in vivo delivery of a suitable companion
30 radionuclide for diagnostics.

[0005] Historically, approaches towards the design of chelating ligands with high metal ion selectivity has been based on several fundamental principles, such as the selection of appropriate donor groups based upon Pearson's Hard and Soft Acids and Bases theory, the influence of cavity size, consideration of the macrocyclic/chelate effect, the optimal chelate ring size, etc. However, the influence of inverted donor group distribution on a common backbone has yet to be investigated. Given the inherent challenges in synthesis of ligands with symmetric donor substitution, and the requirements to often 'break' symmetry in the pursuit of bifunctional derivatives suitable for pharmaceutical application, an assessment of the exact impacts of asymmetric/inverse substitution on metal ion chelation is warranted.

The development of new ligand systems with 'inverted' symmetry offers several advantages with respect to synthesis; protecting group strategies can be circumvented through rational synthetic design and the incorporation of bifunctional handles within the core framework is synthetically more accessible.

[0006] H₄noneunpa having the structure (A) has been previously published by the inventors,⁸ and was also published as Oxyaapa by Hu et al.¹³

H₄noneunpa

(A)

[0007] In the context of nuclear medicine, significant improvements in the treatment of oncogenic disease have been achieved through the advent of several peptide-based radiopharmaceutical agents targeting somatostatin receptor subtype 2 (SSTR2) which is commonly overexpressed in neuroendocrine tumours (NETs).¹⁴ [¹¹¹In][In(DTPA-D-Phe-octreotide)] (OctreoScan™) and [⁶⁸Ga][Ga(DOTATATE)] are two established SSTR2 agonists used clinically for diagnosis of NETs.¹⁴ Most notably, [¹⁷⁷Lu][Lu(DOTATATE)] (Lutathera®) reached clinical FDA-approval as a peptide receptor radionuclide therapy (PRRT) for the treatment of gastroenteropancreatic NETs in 2018.¹⁵ However, as a result of the inherent nuclear decay characteristics of β^- -emitting radionuclides [low linear energy

transfer (~ 0.2 keV/ μm), long-range (0.5 – 10 mm)] treatment scope in this context is limited to primary tumour sites and large metastases.^{4,16}

[0008] Targeted alpha therapy (TAT) has been highlighted as a powerful tool for treatment of both primary and metastatic tumours, with [²²⁵Ac]Ac³⁺ being one of the most prominent candidates for translation into clinical practice, owing to its compatible half-life ($t_{1/2}$ = 10 days), and the high therapeutic potency of its decay progeny (E_{α} = 5–9 MeV, 50–100 μm range).¹⁷ For realisation of this objective, high denticity chelating ligands are required for stable chelation of the [²²⁵Ac]Ac³⁺ ion, which are further capable of sequestering imaging isotopes for diagnostics and assessment of patient suitability prior to radionuclide therapy.

One potential imaging companion is the radio-lanthanide [¹⁵⁵Tb]Tb³⁺, which shares similar bonding characteristics to [²²⁵Ac]Ac³⁺ (e.g. CN = 9; ionic radius = 1.095 Å (Tb³⁺), 1.220 Å (Ac³⁺))^{18,19} and would be suitable for SPECT/CT diagnostics owing to its low-energy gamma emissions (E_{γ} = 87 keV (32%), 105 keV (25%)) and comparable half-life ($t_{1/2}$ = 5.32 days).²⁰ Additionally, [¹⁵⁵Tb]Tb³⁺ is part of the so-called ‘Terbium theranostic quartet’ which includes the radioisotopes ¹⁴⁹Tb, ¹⁵²Tb, ¹⁵⁵Tb, and ¹⁶¹Tb, and encompasses both PET and SPECT imaging modalities, and all three therapeutic decay types (α , β^{-} , Meitner-Auger electrons).^{21–23} Thus, interchange between different Tb³⁺ radioisotopes would provide a true theranostic radiopharmaceutical with identical pharmacokinetic and biodistribution properties.



[0009] There is a general desire for improved chelators that are useful in the targeted in vivo delivery of radiometals using suitable targeting constructs.

[0010] The foregoing examples of the related art and limitations related thereto are intended to be illustrative and not exclusive. Other limitations of the related art will become apparent to those of skill in the art upon a reading of the specification and a study of the drawings.


Summary

[0011] The following embodiments and aspects thereof are described and illustrated in conjunction with systems, tools and methods which are meant to be exemplary and illustrative, not limiting in scope. In various embodiments, one or more of the above-described problems have been reduced or eliminated, while other embodiments are directed to other improvements.

[0012] A novel chelator having the general structure **(I)** is provided wherein each R_1 is independently OH, NH or SH, and X is O, S, or NR_3 , wherein R_3 is H or $CH_2C(=O)R_1$. In some embodiments, the chelator is a bifunctional molecule having the structure **(II)** wherein each R_1 is independently OH, NH or SH or a functional group, each R_2 is independently H or a functional group, and X is O, S, or NR_3 , wherein one or more of the R_1 or R_2 groups is a functional group that allows coupling of the chelator to a biological targeting moiety, and wherein R_3 is H or $CH_2C(=O)R_1$. In some embodiments, the functional group is an ester, an amide, an imide, a thioamide, a thioester, or a guanidinium group. In some embodiments, a radioisotope targeting construct has the structure **(II)**, wherein the chelator is coupled to a biological targeting moiety through one R_1 group or one R_2 group.

**(I)****(II)**


[0013] In some embodiments, the chelator has the structure **(III)** or is a bifunctional molecule having the structure **(III)** wherein each R_1 is independently OH, NH or SH or a functional group; each R_2 is independently H or a functional group; each R_4 is independently H or a functional group, or both R_4 together form a cyclohexyl moiety; each R_5 is independently H or a functional group, or both R_5 together form a cyclohexyl moiety; and X is O, S, or NR_3 , wherein one or more of the R_1 , R_2 , R_4 or R_5 groups is a functional group that allows coupling of the chelator to a biological targeting moiety, and wherein R_3 is H or $CH_2C(=O)R_1$. In some embodiments, the functional group is an ester, an amide, an imide, a thioamide, a thioester, or a guanidinium group. In some embodiments, a radioisotope targeting construct has the structure **(III)**, wherein the chelator is coupled to a biological targeting moiety through one R_1 group, one R_2 group, one R_4 group or one R_5 group.



(III)

[0014] The chelator can be coupled to a biological targeting moiety to facilitate the targeted in vivo delivery of a radioisotope. The radioisotope can be ²²⁷Th, ²²⁵Ac, ¹⁵⁵Tb, ¹⁷⁷Lu, ¹¹¹In, ¹³²La, ²³⁵La, ⁹⁰Y, ⁶⁸Ga, ⁴⁴Sc, ²⁰³Pb, ²¹²Pb, or the like.

[0015] In some embodiments, the chelator has the following structure (6):



(6)

- [0016] Methods of administering an in vivo radioisotope targeting construct comprising the chelator are provided. The in vivo radioisotope targeting construct can be administered to a mammalian subject. The targeting moiety of the in vivo radioisotope targeting construct can be used to enhance accumulation of the radioisotope at a selected location within the body (e.g. the location of cancerous cells), relative to other locations in the body. An imaging procedure can be carried out to evaluate the localization of the in vivo radioisotope targeting construct within the body. The in vivo radioisotope targeting construct can be used to cause cell death at the selected location within the body by exposing the cells to radiation from the radioisotope. The cells can be cancer cells. The mammalian subject can be a human.

[0017] In addition to the exemplary aspects and embodiments described above, further aspects and embodiments will become apparent by reference to the drawings and by study of the following detailed descriptions.

5 Brief Description of the Drawings

[0018] Exemplary embodiments are illustrated in referenced figures of the drawings. It is intended that the embodiments and figures disclosed herein are to be considered illustrative rather than restrictive.

[0019] FIG. 1 shows the structure of an exemplary in vivo targeting chelate construct.

10 [0020] FIG. 2 shows the chemical structure of H₄noneunpaX (**6**) and the ¹H assignment (left side) and ¹³C assignment (right side).

[0021] FIG. 3 shows the ¹H NMR spectrum of H₄noneunpaX·4HCl (**6**) (400 MHz, D₂O, 298 K).

15 [0022] FIG. 4 shows ¹³C {¹H} NMR spectrum of H₄noneunpaX·4HCl (**6**) (75 MHz, D₂O, 298 K).

[0023] FIG. 5 shows ¹H – ¹H COSY NMR spectrum of H₄noneunpaX (**6**) (300 MHz, D₂O, 298 K).

[0024] FIG. 6 shows ¹H – ¹³C HSQC NMR spectrum of H₄noneunpaX 4HCl (**6**) (300 MHz, 75 MHz, D₂O, 298 K).

20 [0025] FIG. 7 shows ¹H – ¹³C HMBC NMR spectrum of H₄noneunpaX 4HCl (**6**) (300 MHz, 75 MHz, D₂O, 298 K).

[0026] FIG. 8 shows stacked ¹H NMR spectra of H₄noneunpaX and corresponding complexes of La³⁺, Lu³⁺ and In³⁺ (400 MHz, D₂O, 298 K, pD 7.0). [In(noneunpaX)]⁺ was characterised at pD=4.5.

25 [0027] FIG. 9 shows ¹H NMR spectrum of [La(noneunpaX)]⁺ (400 MHz, D₂O, 298 K, pH 7), and includes the chemical structure of [La(noneunpaX)]⁺ with ¹H assignments indicated.

[0028] FIG. 10 shows ¹H – ¹H COSY NMR spectrum of [La(noneunpaX)]⁺ (400 MHz, D₂O, 298 K, pH 7.0).

[0029] FIG. 11 shows $^1\text{H} - ^1\text{H}$ COSY NMR spectrum of $[\text{La}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pH 7.0) (alkyl expansion).

[0030] FIG. 12 shows $^1\text{H} - ^1\text{H}$ COSY NMR spectrum of $[\text{La}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pH 7.0) (aromatic expansion).

5 **[0031]** FIG. 13 shows ^1H NMR spectrum of $[\text{Lu}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pH 7.0) and includes the chemical structure of $[\text{Lu}(\text{noneunpaX})]^-$ with ^1H assignments indicated.

[0032] FIG. 14 shows $^1\text{H} - ^1\text{H}$ COSY NMR spectrum of $[\text{Lu}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pH 7.0).

10 **[0033]** FIG. 15 shows $^1\text{H} - ^1\text{H}$ COSY NMR spectrum of $[\text{Lu}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pH 7.0) (alkyl expansion).

[0034] FIG. 16 shows $^1\text{H} - ^1\text{H}$ COSY NMR spectrum of $[\text{Lu}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pH 7.0) (aromatic expansion).

[0035] FIG. 17 shows ^1H NMR spectrum of $[\text{In}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pD 4.5) and includes the chemical structure of $[\text{In}(\text{noneunpaX})]^-$ with ^1H assignments indicated.

15 **[0036]** FIG. 18 shows $^1\text{H} - ^1\text{H}$ COSY NMR spectrum of $[\text{In}(\text{noneunpaX})]^-$ (400 MHz, D_2O , 298 K, pD 4.5).

[0037] FIG. 19 shows calculated absorptivity curves of the eight species of $\text{H}_4\text{noneunpaX}$; determined by combined potentiometric-spectrophotometric titrations ($[\text{L}] = 9.57 \times 10^{-4} \text{ M}$, $l = 0.2 \text{ cm}$), and acidic in-batch spectrophotometric titrations ($[\text{L}] = 1.0 \times 10^{-4} \text{ M}$, $l = 1.0 \text{ cm}$), at
20 $T = 298 \text{ K}$, $I = 0.16 \text{ M NaCl}$.

[0038] FIG. 20 shows a speciation diagram of $\text{H}_4\text{noneunpaX}$ as a function of pH, dotted line indicates pH 7.4.

[0039] FIG. 21 shows a speciation diagram of $\text{H}_4\text{noneunpaX}$ with La^{3+} ($[\text{La}^{3+}] = [\text{H}_4\text{noneunpaX}] = 1 \times 10^{-4} \text{ M}$), dashed lines indicated physiological conditions (pH 7.4), at
25 298 K , $I = 0.16 \text{ M NaCl}$.

[0040] FIG. 22 shows a speciation diagram of $\text{H}_4\text{noneunpaX}$ with Sm^{3+} ($[\text{Sm}^{3+}] = [\text{H}_4\text{noneunpaX}] = 1 \times 10^{-4} \text{ M}$), dashed lines indicated physiological conditions (pH 7.4), at
 298 K , $I = 0.16 \text{ M NaCl}$.

[0041] FIG. 23 shows a speciation diagram of H₄noneunpaX with Gd³⁺ ([Gd³⁺] = [H₄noneunpaX] = 1 × 10⁻⁴ M), dashed lines indicated physiological conditions (pH 7.4), at 298 K, I = 0.16 M NaCl.

[0042] FIG. 24 shows a speciation diagram of H₄noneunpaX with Dy³⁺ ([Dy³⁺] = [H₄noneunpaX] = 1 × 10⁻⁴ M), dashed lines indicated physiological conditions (pH 7.4), at 298 K, I = 0.16 M NaCl.

[0043] FIG. 25 shows a speciation diagram of H₄noneunpaX with Lu³⁺ ([Lu³⁺] = [H₄noneunpaX] = 1 × 10⁻⁴ M), dashed lines indicated physiological conditions (pH 7.4), at 298 K, I = 0.16 M NaCl.

[0044] FIG. 26 shows a speciation diagram of H₄noneunpaX with In³⁺ ([In³⁺] = [H₄noneunpaX] = 1 × 10⁻⁴ M), dashed lines indicated physiological conditions (pH 7.4), at 298 K, I = 0.16 M NaCl.

[0045] FIG. 27 shows a speciation diagram of H₄noneunpaX with Sc³⁺ ([Sc³⁺] = [H₄noneunpaX] = 1 × 10⁻⁴ M), dashed lines indicated physiological conditions (pH 7.4), at 298 K, I = 0.16 M NaCl.

[0046] FIG. 28 shows concentration-dependent radiolabelling studies of H₄noneunpa, H₄noneunpaX, and DOTA with: [⁴⁴Sc]Sc³⁺ (1.2 MBq) in NaOAc (0.1 M, pH 4.5) (panel (A)), [¹¹¹In]In³⁺ (1.0 MBq) in NH₄OAc (0.5 M, pH 5.8) (panel (B)), [¹⁵⁵Tb]Tb³⁺ (40 kBq) in NH₄OAc (0.5 M, pH 6.0) (panel (C)), [¹⁷⁷Lu]Lu³⁺ (150 kBq) in NH₄OAc (0.5 M, pH 6.0) (panel (D)), [²¹³Bi]Bi³⁺ (680 kBq) in MES (1.0 M, pH 5.5), (F) [²²⁵Ac]Ac³⁺ (40 kBq) in NH₄OAc (1.0 M, pH 7.3) (panel (E)).

[0047] FIG. 29 shows human serum stability studies of H₄noneunpa and H₄noneunpaX with: [¹¹¹In]In³⁺ (54 GBq/μmol) (panel (A)), [¹⁵⁵Tb]Tb³⁺ (1.0 GBq/μmol) (panel (B)), [¹⁷⁷Lu]Lu³⁺ (2.0 GBq/μmol) (panel (C)), [²²⁵Ac]Ac³⁺ (134 MBq/μmol) (panel (D)).

[0048] FIG. 30 shows DFT-optimised structures of [La(noneunpaX)]⁻ in panel (A) (left-hand orientation), and panel (B) front orientation, and [Lu(noneunpaX)]⁻ in panel (c) (left-hand orientation), and panel (D) (front orientation). Selected hydrogens have been omitted for clarity.

[0049] FIG. 31 shows concentration – dependent radiolabelling of bifunctional H₄noneunpaX with: **(A)** [⁴⁴Sc]Sc³⁺ (400 kBq) in NaOAc (0.1 M, pH 4.5), **(B)** [¹¹¹In]In³⁺ (1.06

MBq) in NH_4OAc (0.5 M, pH 5.5), (C) $[^{177}\text{Lu}]\text{Lu}^{3+}$ (350 - 500 kBq) in NH_4OAc (0.5 M, pH 6.0), (D) $[^{133/135}\text{La}]\text{La}^{3+}$ (400 kBq) in NH_4OAc (0.2 M, pH 7.0), (E) $[^{155}\text{Tb}]\text{Tb}^{3+}$ (140 kBq) in NH_4OAc (0.5 M, pH 7.0) and (F) $[^{225}\text{Ac}]\text{Ac}^{3+}$ (40 kBq) in NH_4OAc (0.5 M, pH 7.2).

5 **[0050]** FIG. 32 shows a molar activity study of $\text{H}_4\text{noneunpaX-Bn-NH}_2$ with $[^{177}\text{Lu}]\text{LuCl}_3$ (20 MBq) in NH_4OAc (0.5 M, pH 6.0) at RT monitored over 10 minutes. Highest molar activity (250 GBq/ μmol), ligand-to-metal ratio ($[\text{L}]:[\text{M}]$; [174:1]) (left panel). Human serum stability challenge of $[^{177}\text{Lu}][\text{Lu}(\text{noneunpaX-Bn-NH}_2)]$ (5.2 GBq/ μmol) conducted at 37 °C and monitored over 7 days via radio-iTLC ($n = 3$) (right panel).

10 **[0051]** FIG. 33 shows radiolabelling studies of bifunctional $\text{H}_4\text{noneunpaX}$ and corresponding peptide-conjugates: (A) Concentration-dependent radiolabelling with $[^{225}\text{Ac}]\text{Ac}^{3+}$ (40 kBq) in NH_4OAc buffer (0.5 M, pH 7) (RT, 10 min.). (B) Human serum stability challenge of $[^{225}\text{Ac}]\text{Ac}^{3+}$ -labelled compounds. (C) Concentration-dependent radiolabelling with $[^{155}\text{Tb}]\text{Tb}^{3+}$ (120 kBq) in NH_4OAc buffer (0.5 M, pH 6) (RT, 10 min.). (D) Human serum stability challenge of $[^{155}\text{Tb}]\text{Tb}^{3+}$ -labelled bioconjugates.

15 **[0052]** FIG. 34 shows radio-HPLC traces of $[^{155}\text{Tb}]\text{Tb}^{3+}$ control and $[^{155}\text{Tb}][\text{Tb}^{3+}$ labelled $\text{H}_4\text{noneunpaX-Bn-NH}_2$ and $\text{H}_4\text{noneunpaX-Bn-NCS}$. Method: A: H_2O (0.1% TFA), B: MeCN (0.1% TFA); 100% A to 10% B over 15 min., 1 mL/min.

20 **[0053]** FIG. 35 shows radio-HPLC traces for $[^{155}\text{Tb}][\text{Tb}(\text{noneunpaX-Ahx-Tyr}^3\text{-TATE})]$, Method: A: H_2O (0.1% TFA), B: MeCN (0.1% TFA); 100% A to 60% B over 15 min., 1 mL/min.

[0054] FIG. 36 shows radio-HPLC traces for $[^{155}\text{Tb}][\text{Tb}(\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE})]$, Method: A: H_2O (0.1% TFA), B: MeCN (0.1% TFA); 100% A to 60% B over 15 min., 1 mL/min.

25 **[0055]** FIG. 37 shows coronal views of maximum intensity projections (MIPs) from quantitative dynamic SPECT/CT scans at 0 – 1 h post administration of $[^{155}\text{Tb}][\text{Tb}(\text{noneunpaX-Ahx-Tyr}^3\text{-TATE})]$ (top) and $[^{155}\text{Tb}][\text{Tb}(\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE})]$ (bottom) in NRG mice bearing AR42J exocrine tumour xenografts (left shoulder).

[0056] FIG. 38 shows sagittal views of maximum intensity projections (MIPs) from SPECT/CT scans at 1, 2 and 4 h post administration of $[^{155}\text{Tb}][\text{Tb}(\text{noneunpaX-Ahx-Tyr}^3\text{-TATE})]$

TATE)] (top) and [^{155}Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)] (bottom) in NRG mice bearing AR42J pancreatic exocrine tumour xenografts (left shoulder).

[0057] FIG. 39 shows representative time-activity plots of (A) [^{155}Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)] and (B) [^{155}Tb][Tb(noneunpaX-PEG-Tyr³-TATE)] in NRG mice bearing AR42J exocrine/pancreatic tumour xenografts. Standardised uptake values (SUVs) were extracted for ROIs in relevant organs from calibrated SPECT/CT images.


[0058] FIG. 40 shows decay-corrected biodistribution studies of [^{155}Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)] at 2 h and 4 h post-injection and [^{155}Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)] at 4 h post-injection in NRG mice bearing AR42J exocrine/pancreatic tumour xenografts.

Description

[0059] Throughout the following description specific details are set forth in order to provide a more thorough understanding to persons skilled in the art. However, well known elements may not have been shown or described in detail to avoid unnecessarily obscuring the disclosure. Accordingly, the description and drawings are to be regarded in an illustrative, rather than a restrictive, sense.

[0060] The inventors have now developed a novel chelating ligand, H₄noneunpaX, having an inverted arrangement of functional groups, and have compared the characteristics of this novel ligand with H₄noneunpa. The ligand displays preferential complexation with trivalent Ln³⁺ ions, forming single isomeric species in solution, which exhibits high thermodynamic stability and kinetic inertness. Radiolabelling studies of H₄noneunpaX with [^{111}In]In³⁺, [^{155}Tb]Tb³⁺, [^{177}Lu]Lu³⁺ and [^{225}Ac]Ac³⁺ showed excellent compatibility, achieving high molar activities under mild conditions (RT, 10 mins). As a proof of principle, a bifunctional derivative of H₄noneunpaX was prepared through a facile synthetic approach and was conjugated to two Tyr³-Octreotate peptide analogues for targeting of neuroendocrine tumours (NETs). Further radiolabelling studies of these bioconjugates with [^{111}In]In³⁺, [^{177}Lu]Lu³⁺, [$^{133/135}\text{La}$]La³⁺, [^{155}Tb]Tb³⁺ and [^{225}Ac]Ac³⁺, showed promising results for the development of an [^{225}Ac]Ac³⁺/[^{155}Tb]Tb³⁺-theranostic pair. SPECT/CT and biodistribution studies of the [^{155}Tb]Tb³⁺-radiolabelled tracers in tumour bearing mice, showed excellent performance, with good tumour uptake, low residual background organ activity, and no evidence of degradation *in vivo*.

[0061] The objectives of this work were therefore to investigate the influence of donor group substitution about a common backbone and study the impact of asymmetric/inverted displacement on metal ion affinity. H₄noneunpaX having the structure (6) was prepared and comparisons to the symmetric counterpart H₄noneunpa/H₄oxyaapa were assessed. To further assess the utility of this approach, a bifunctional analogue of H₄noneunapX was prepared, to firstly demonstrate the synthetic accessibility of bifunctional asymmetric/inverted chelators, and to investigate their performed *in vivo*.



(6)

[0062] As used herein, the term prophylaxis includes preventing, minimizing the severity of, or preventing a worsening of a condition. As used herein, the terms treat or treatment include reversing or lessening the severity of a condition.

[0063] As used herein, the term antibody includes all forms of antibodies including polyclonal antibodies, monoclonal antibodies, chimeric antibodies, humanized antibodies, single chain antibodies, multimeric antibodies, and the like. The term antigen binding fragment of an antibody refers to any portion of an antibody that is capable of binding to an antigen and includes by way of example only and without limitation Fab fragments, F(ab')₂ fragments, Fv fragments, scFv fragments, minibodies, diabodies, and the like. Reference to a specific antibody includes reference to any antibodies that are determined to be biosimilar to that specific antibody by any regulatory authority.


[0064] As used herein, the term peptidomimetic means a small protein-like molecule designed to mimic a peptide, and includes without limitation modified peptides, peptidic foldamers, structural mimetics and mechanistic mimetics.

[0065] A chelator composition for radiometals is disclosed. A method of using and making the composition is also disclosed. The composition can be used as a therapeutic and/or diagnostic agent.

[0066] The inventors have now determined that chelators having the general structure **(6)** can coordinate radioisotopes under mild conditions and produce a complex that is stable under in vivo conditions, making such chelators particularly suitable for example for application in radiotherapeutic, diagnostic and/or theranostic constructs. The chelator can be coupled directly or via a linker to a biological targeting moiety to create a construct suitable for use in such applications.

[0067] In some embodiments, H₄noneunpaX can be directly coupled to a biological targeting moiety, optionally with a linker interposing the H₄noneunpaX and the biological targeting moiety, by coupling the biological targeting moiety or linker directly to one of the carboxyl groups of structure **(6)**. Furthermore, in some embodiments, one or more of the oxygen atoms of the carboxyl group is substituted by a different heterotatom, e.g. N or S. Thus, in various embodiments, the functional group provided on the bifunctional H₄noneunpaX chelator to couple the chelator to the biological targeting moiety can be a carboxyl, an ester, an amide, an imide, a thioamide, a thioester, a guanidinium, or the like.


[0068] In some embodiments, the chelator has the following structure **(I)** wherein each R₁ is independently OH, NH or SH, and wherein X is O, S, or NR₃, wherein R₃ is H or CH₂C(=O)R₁.



(I)

In some embodiments, the chelator is a bifunctional molecule having the structure **(II)** wherein each R₁ is independently OH, NH or SH or a functional group, each R₂ is independently H or a functional group, and X is O, S, or NR₃, wherein one or more of the R₁

or R_2 groups is a functional group that allows coupling of the chelator to a biological targeting moiety, and wherein R_3 is H or $\text{CH}_2\text{C}(=\text{O})\text{R}_1$.




(II)

5 **[0069]** In some such embodiments, the functional group is a carboxyl, an ester, an amide, an imide, a thioamide, a thioester, a guanidinium, or the like.

[0070] In some embodiments, a radioisotope targeting construct has the structure (II), wherein one of the R_1 groups or one of the R_2 groups is a biological targeting moiety.


[0071] In some embodiments, the chelator has the structure (III) or is a bifunctional molecule having the structure (III) wherein each R_1 is independently OH, NH or SH or a functional group; each R_2 is independently H or a functional group; each R_4 is independently H or a functional group, or both R_4 together form a cyclohexyl moiety; each R_5 is independently H or a functional group, or both R_5 together form a cyclohexyl moiety; and X is O, S, or NR_3 , wherein one or more of the R_1 , R_2 , R_4 or R_5 groups is a functional group that allows coupling of the chelator to a biological targeting moiety, and wherein R_3 is H or $\text{CH}_2\text{C}(=\text{O})\text{R}_1$. In some embodiments, when both R_4 together form a cyclohexyl moiety, each R_5 is independently H or a functional group. In some embodiments, when both R_5 together form a cyclohexyl moiety, each R_4 is independently H or a functional group. In some embodiments, the functional group is an ester, an amide, an imide, a thioamide, a thioester, or a guanidinium group or the like. In some embodiments, a radioisotope targeting construct has the structure (III), wherein the chelator is coupled to a biological targeting moiety through one R_1 group, one R_2 group, one R_4 group or one R_5 group. In some embodiments, a radioisotope targeting construct has the structure (III), wherein one of the

R₁ groups, one of the R₂ groups, one of the R₄ groups or one of the R₅ groups is a biological targeting moiety.



5

[0072] For example, in some embodiments, the chelator has one of the structures shown below:



10 **[0073]** It has been found by other groups, for example Milenic et al.⁵⁸, that DTPA derivatives comprising a cyclohexyl moiety in the backbone of the chelator show comparable specific activities when incorporated into a chelate-antibody conjugate. Accordingly, it can be soundly predicted that compounds incorporating a cyclohexyl moiety in the backbone of the chelator will likely exhibit similar binding properties to NoneunpaX

15 having the structure (6).

[0074] In some embodiments, the chelator is NoneunpaX having the structure (6).

[0075] In some embodiments as shown in FIG. 1, an in vivo targeting chelate construct **120** has a targeting moiety **122** coupled to a chelator **126**. In some embodiments, including the illustrated embodiment, a linker **124** interposes targeting moiety **122** and chelator **126**. Together, targeting moiety **122**, linker **124** (if present) and chelator **126** comprise in vivo targeting construct **130**. Further, chelator **126** is used to chelate a radionuclide **128** that is suitable for in vivo imaging and/or radiotherapy. Radionuclide **128** together with in vivo targeting construct **130** provides an in vivo targeting chelate construct **120** suitable for targeted in vivo delivery of the radionuclide **128** payload as assisted by targeting moiety **122**.

[0076] Any moiety suitable for directing the targeted delivery of in vivo targeting chelate construct **120** in vivo can be used as targeting moiety **122**. In some embodiments, the targeting moiety **122** of the targeting construct **120** is a hapten, antigen, aptamer, affibody molecule, enzyme, protein, peptide, antibody, antigen-binding fragment of an antibody, peptidomimetic, receptor ligand, steroid, hormone, growth factor, cytokine, molecule that recognizes cell surface receptors (including molecules involved in growth, metabolism or function of cells), lipid, lipophilic group, carbohydrate, or any other molecule or targeting component capable of selectively directing a construct to a specific location within the body. The targeting moiety can be produced in any suitable manner, e.g. as a biologic, semisynthetically, or synthetically.

[0077] Examples of targeting moieties that have been developed to deliver radioisotope targeting constructs to desired locations within the body of a mammalian subject in vivo include antibodies targeting specific markers associated with specific types of cancers, peptidomimetics targeting proteins that are highly expressed in cancer cells, and the like. Exemplary non-limiting examples of suitable targeting moieties are listed in Table 1 (Lau, J.; Rousseau, E.; Kwon, D.; Lin, K.; Benard, F.; Chen, X., Insight into the Development of PET Radiopharmaceuticals for Oncology. Cancers 2020, 12, 1312, the entirety of which is incorporated by reference herein). Some targeting moieties selectively interact with biological targets, including antigens, proteins, carbohydrates or other molecules present on the surface of cells that are overexpressed in cancer cells relative to normal cells, e.g. tumor-associated antigens. Exemplary non-limiting examples of suitable targets are listed in Table 1. Suitable targets and/or targeting moieties for radiopharmaceuticals, whether now known or discovered or developed in the future, would be known to a person skilled in

the art. In some embodiments, targeting moiety **122** is an antibody or an antigen-binding fragment of an antibody. In some embodiments, targeting moiety **122** is a peptidomimetic. In some embodiments, the targeting moiety **122** is one of the targeting moieties listed in Table 1, with any chelator present in the referenced molecule replaced by H₄noneunpaX as the chelator. In some embodiments, the targeting moiety **122** interacts selectively with one of the targets listed in Table 1.

Table 1. Exemplary targeting moieties and biological targets for targeted radiation therapy.

Targeting Moiety	Biological Target	Indications
A33 antibody	A33 transmembrane glycoprotein	colorectal cancer
dihydrotestosterone (DHT)	androgen receptor (AR)	prostate cancer
HuMab-5B1	CA19.9	pancreatic cancer, bladder cancer
Girentuximab	carbonic anhydrase 9 (CA-IX)	clear cell renal cell carcinoma
AMG211 bispecific T-cell engager	carcinoembryonic antigen	gastrointestinal adenocarcinoma
IAB22M2C minibody	CD8	melanoma, lung cancer, hepatocarcinoma
Rituximab	CD20	B-cell lymphoma
obinutuzumab	CD20	B-cell lymphoma
U36 antibody	CD44v6	head and neck cancer
Plerixafor	C-X-C chemokine receptor type 4 (CXCR4)	hematological and solid malignancies
Pentixafor	CXCR4	hematological and solid malignancies
NFB	CXCR4	hematological and solid malignancies
Ipilimumab	Cytotoxic T-lymphocyte-associated protein 4 (CTLA-4)	Melanoma
erlotinib	epidermal growth factor receptor (EGFR)	nonsmall cell lung carcinoma; colorectal cancer
PD153035	EGFR	nonsmall cell lung carcinoma; colorectal cancer
Afatinib	EGFR	nonsmall cell lung carcinoma; colorectal cancer
Cetuximab	EGFR	nonsmall cell lung carcinoma; colorectal cancer

panitumumab	EGFR	nonsmall cell lung carcinoma; colorectal cancer
ABY-025 affibody	Epidermal growth factor receptor 2 (ERBB2)	breast cancer
HER2-nanobody	ERBB2	breast cancer
Trastuzumab	ERBB2	breast cancer
Pertuzumab	ERBB2	breast cancer
GSK2849330	epidermal growth factor receptor 3 (ERBB3)	solid malignancies
lumretuzumab	ERBB3	solid malignancies
4FMFES	estrogen receptor (ER)	breast cancer and gynecologic cancers
FAP1-04	fibroblast activation protein α	solid malignancies
FAP1-21	fibroblast activation protein α	solid malignancies
FAP1-46	fibroblast activation protein α	solid malignancies
Galactose	galactose metabolism	hepatocarcinoma
CB-TE2A-AR06 peptide (with H ₄ noneunpaX substituted for DOTA)	gastrin-releasing peptide receptor (GRPR)	prostate cancer, breast cancer, glioma
BAY 864367 peptide (with H ₄ noneunpaX -bound ligand label instead of 18F labeling)	GRPR	prostate cancer, breast cancer, glioma
RM2 peptide (with H ₄ noneunpaX substituted for DOTA)	GRPR	prostate cancer, breast cancer, glioma
SB3 peptide (with H ₄ noneunpaX substituted for DOTA)	GRPR	prostate cancer, breast cancer, glioma
RM26 peptide	GRPR	prostate cancer, breast cancer, glioma
BBN-RGD peptide	GRPR	prostate cancer, breast cancer, glioma
Aca-BBN peptide	GRPR	prostate cancer, breast cancer, glioma
NeoBOMB1 peptide (with H ₄ noneunpaX substituted for DOTA)	GRPR	prostate cancer, breast cancer, glioma
exendin-4 peptide	glucagon-like peptide 1 receptor (GLP-1R)	insulinoma
Glucose	glucose metabolism	neoplasm
Codrituzumab	glypican 3	hepatocarcinoma
EF5	Hypoxia	solid malignancies
MISO	Hypoxia	solid malignancies

AZA	Hypoxia	solid malignancies
HX4	Hypoxia	solid malignancies
ASTM	Hypoxia	solid malignancies
LLP2A peptidomimetic	integrin $\alpha 4 \beta 1$	multiple myeloma
galacto-RGD peptide	integrin $\alpha v \beta 3$	solid malignancies
FPP(RGD) ₂ peptide	integrin $\alpha v \beta 3$	solid malignancies
RGD-K5 peptide	integrin $\alpha v \beta 3$	solid malignancies
Fluciclatide	integrin $\alpha v \beta 3$	solid malignancies
alfatide-I	integrin $\alpha v \beta 3$	solid malignancies
alfatide-II	integrin $\alpha v \beta 3$	solid malignancies
PRGD ₂ peptide	integrin $\alpha v \beta 3$	solid malignancies
$\alpha v \beta 6$ -BP peptide	integrin $\alpha v \beta 6$	head and neck cancer, lung cancer, colorectal cancer, breast cancer, pancreatic cancer
CycMSHhex targeting peptides	melanocortin-1 receptor (MC1R)	melanoma
MMOT0530A antibody	Mesothelin	pancreatic ductal adenocarcinoma and ovarian cancer
SP peptide	Neurokinin1 receptor (NK1R)	glioma
neurotensin	Neurotensin 1 receptor (NTS1R)	prostate cancer
PARPi	poly(ADP-ribose) polymerase 1 (PARP1)	head and neck cancer
PSMA peptidomimetics, e.g. PSMA-1007, PSMA-7 (with H ₄ noneunpaX -bound ligand replacing 18F labelling); PSMA-11 (with H ₄ noneunpaX replacing the HBED chelator); PSMA-617, PSMA-I&T (with H ₄ noneunpaX replacing the DOTA chelator)	Prostate-specific membrane antigen (PSMA)	prostate cancer
DCFPyL	PSMA	prostate cancer
DCFBC	PSMA	prostate cancer
HuJ591 antibody	PSMA	prostate cancer
durvalumab	programmed cell death protein (PD-1)	nonsmall cell lung carcinoma
nivolumab	PD-1	nonsmall cell lung carcinoma
pembrolizumab	PD-1	nonsmall cell lung carcinoma

BMS-986192 adnectin	programmed death-ligand 1 (PD-L1)	nonsmall cell lung carcinoma, bladder cancer, breast cancer
atezolizumab	PD-L1	nonsmall cell lung carcinoma, bladder cancer, breast cancer
MSTP2109A antibody	six-transmembrane epithelial antigen of prostate-1 (STEAP1)	prostate cancer
TATE peptide (octreotate)	somatostatin receptor 2 (SSTR2)	neuroendocrine tumors
TOC peptide	SSTR2	neuroendocrine tumors
NOC peptide	SSTR2	neuroendocrine tumors
JR11	SSTR2	neuroendocrine tumors
thymidine	thymidine kinase (DNA replication)	solid malignancies
fresolimumab	transforming growth factor-beta (TGF- β)	glioma
bevacizumab	vascular endothelial growth factor receptor (VEGFR)	solid malignancies

[0078] Any suitable linker can be used as linker **124** to couple chelator **126** to targeting moiety **122**. For example and by way of illustration only, suitable linkers can include:

- 5 • a hydrocarbon linker containing between 1 and 10 carbon atoms (C₁-C₁₀), including 2, 3, 4, 5, 6, 7, 8 or 9 carbon atoms that is optionally saturated or unsaturated, optionally substituted with one or more heteroatoms or having one or more substituents; the hydrocarbon linker can be linear, cyclic and/or branched, e.g. 8-aminooctanoic acid, 6-aminohexanoic acid;
- 10 • an aromatic linker containing an aromatic moiety such as a benzyl group, e.g. aminophenylacetic acid;
- an amino acid linker having between 1 and 10 amino acid residues, including 2, 3, 4, 5, 6, 7, 8, or 9 amino acid residues, any one or more of which may be naturally occurring amino acid residues, D-amino acid residues or other non-naturally occurring residues, examples of which include GlyGly, GluGluGlu, GlySerGlySer;
- 15 • a cyclized linker, or cyclized ring structure, optionally a cyclized amino acid linker, e.g. aminocyclohexanecarboxylic acid;
- a PEG-linker of any suitable length;
- cationic linkers, whether formed from amino acid residues or other residues, e.g. . Pip, 4-(2-aminoethyl)-1-carboxymethyl-piperazine (Acp);

- anionic linkers, whether formed from amino acid residues or other residues, e.g. AspAsp, GluGlu;
- a carbohydrate containing linker;
- click chemistry linkers (triazoles);
- 5 • any other suitable linker;
- or combinations or modifications of the foregoing.

Examples of linkers that have been developed in the art for other radiopharmaceutical targeting constructs are known to those skilled in the art. Hydrophilic or charged linkers such as PEG-linkers or cationic/anionic linkers may be used to increase the overall water solubility of the targeting construct. Amino acid side chain substitutions and/or inclusion of carbohydrate moieties may be made to improve or alter the solubility and/or pharmacokinetics of the targeting construct. A person skilled in the art could develop and optimize a suitable linker for a particular application if desired. Examples of linkers that have been developed in the art for other radiopharmaceutical targeting constructs are described, by way of example only and without limitation, by Benešová et al., Barnaski et al. and Kuo et al. (See Benešová, M.; Schäfer, M.; Bauder-Wüst, U.; Afshar-Oromieh, A.; Kratochwil, C.; Mier, W.; Haberkorn, U.; Kopka, K.; Eder, M., Preclinical Evaluation of a Tailor-Made DOTA-Conjugated PSMA Inhibitor with Optimized Linker Moiety for Imaging and Endoradiotherapy of Prostate Cancer. *J. Nucl. Med.* **2015**, *56*, 914-920; Baranski, A.-C.; Schäfer, M.; Bauder-Wüst, U.; Wacker, A.; Schmidt, J.; Liolios, C.; Mier, W.; Haberkorn, U.; Eisenhut, M.; Kopka, K.; Eder, M., Improving the Imaging Contrast of ⁶⁸Ga-PSMA-11 by Targeted Linker Design: Charged Spacer Moieties Enhance the Pharmacokinetic Properties. *Bioconjugate Chem.* **2017**, *28*, 2485-2492; Kuo, H.-T.; Pan, J.; Zhang, Z.; Lau, J.; Merckens, H.; Zhang, C.; Colpo, N.; Lin, K.-S.; Benard, F., Effects of linker modification on tumor-to-kidney contrast of ⁶⁸Ga-labeled PSMA-targeted imaging probes. *Mol Pharm.* **2018**, *15*, 3502-3511; Benešová, M.; Bauder-Wüst, U.; Schäfer, M.; Klika, K. D.; Mier, W.; Haberkorn, U.; Kopka, K.; Eder, M., Linker Modification Strategies To Control the Prostate-Specific Membrane Antigen (PSMA)-Targeting and Pharmacokinetic Properties of DOTA-Conjugated PSMA Inhibitors. *J. Med. Chem.* **2016**, *59*, 1761-1775, each of which is incorporated by reference herein in its entirety). A person skilled in the art could develop and optimize a suitable linker for a particular application.

[0079] In some embodiments, a construct such as construct **120** is prepared by carrying out suitable reactions to couple targeting moiety **122** and chelator **126**, for example via suitable chemical reaction, to yield an in vivo targeting construct **130**, optionally with linker **124** interposing targeting moiety **122** and chelator **126**. The radionuclide **128** is then added and bound to chelator **126**, e.g. at a later time and in a hospital or clinic setting, to form the desired in vivo targeting metal chelate construct **120**. In other embodiments, radionuclide **128** could be first chelated with chelator **126**, and then chelator **126** is conjugated with targeting moiety **122** in any suitable manner to yield in vivo targeting chelate construct **120**.

[0080] In some embodiments, the radionuclide **128** is bound to chelator **126** (including as part of construct **130**) under mild temperature conditions, e.g. less than about 65°C, 60°C, 55°C, 50°C, 45°C, 40°C, 35°C or 30°C. In some embodiments, the mild temperature conditions are between about 10°C and 65°C, including any value or subrange therebetween, e.g. 15°C, 16°C, 17°C, 18°C, 19°C, 20°C, 21°C, 22°C, 23°C, 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 35°C, 40°C, 45°C, 50°C, 55°C or 60°C. In some embodiments, the radionuclide **128** is conjugated to chelator **126** or construct **130** at room temperature, i.e. in the range of about 15°C to about 25°C, including any temperature value therebetween, e.g. 16°C, 17°C, 18°C, 19°C, 20°C, 21°C, 22°C, 23°C, or 24°C.

[0081] In some embodiments, the radionuclide **128** or construct **130** is combined with chelator **126** to form a metal chelate under mild pH conditions, e.g. between about 5.0 and about 7.4, including any value or subrange therebetween, e.g. 5.2, 5.4, 5.6, 5.8, 6.0, 6.2, 6.4, 6.6, 6.8, 7.0 or 7.2. In some embodiments the radionuclide **128** is conjugated to chelator **126** at approximately neutral pH, i.e. a pH of approximately 7.0, e.g. between about 6.8 and 7.2 including any value therebetween, e.g. 6.9, 7.0 or 7.1. In some embodiments, the radionuclide **128** is conjugated to chelator **126** at approximately physiological pH, i.e. at approximately pH 7.4, e.g. between about 7.2 and 7.6 including any value therebetween, e.g. 7.3, 7.4 or 7.5. In some embodiments, radionuclide **128** is combined with chelator **126** or construct **130** in aqueous solution. In some embodiments, the aqueous solution is free or substantially free of alcohol such as ethanol.

[0082] In some embodiments, the radionuclide **128** is combined with chelator **126** or construct **130** for an incubation period to allow a chelated metal complex to form. In some embodiments, the incubation period is between about 5 minutes and about 6 hours, including any period therebetween, e.g. 10, 15, 20, 25, 30, 45, 60 or 90 minutes, or 2, 3, 4

or 5 hours. In some embodiments, the incubation period is between about 5 minutes and about 30 minutes.

[0083] In some embodiments, the concentration of chelator **126** or construct **130** that is present when conjugated to radionuclide **128** is between about 10^{-4} to 10^{-7} M, including any value therebetween, e.g. 10^{-5} or 10^{-6} M. The concentration of chelator **126** or construct **130** that is used can be adjusted depending on the complexation kinetics between the particular chelator **126** and radionuclide **128** used in any particular embodiment. Similarly the temperature at which the radionuclide **128** is combined with chelator **126** or construct **130** can be varied depending on the complexation kinetics.

[0084] In some embodiments, in vivo radioisotope targeting chelate construct **120** is present in mammalian serum, optionally in human serum. In some embodiments, in vivo radioisotope targeting chelate construct **120** is stable in mammalian serum, optionally in human serum. In some embodiments, in vivo radioisotope targeting chelate construct **120** is present in mammalian serum within the body of a mammal, optionally in human serum within the body of the human. In some embodiments, in vivo radioisotope targeting chelate construct **120** is present in mammalian blood, optionally in human blood. In some embodiments, in vivo radioisotope targeting chelate construct **120** is present in mammalian blood within the body of a mammal, optionally in human blood in the body of the human. In some embodiments, in vivo radioisotope targeting chelate construct **120** is present within the body of a mammal, optionally the body of a human. In some embodiments, in vivo radioisotope targeting chelate construct **120** is present in a mammalian cell, optionally a human cell.

[0085] In some embodiments, radionuclide **128** is delivered to a selected location within the body of a mammalian subject by administering to the subject an in vivo radioisotope targeting chelate construct **120** incorporating the radionuclide **128** and a targeting moiety **122** that specifically directs the in vivo radioisotope targeting chelate construct **120**, including the bound radionuclide **128**, to the selected location within the body of the subject. In some embodiments, the method includes allowing the targeting moiety **122** to enhance the accumulation of the in vivo radioisotope targeting chelate construct **120** at the selected location within the body relative to other locations in the body to selectively deliver a dose of radiation to the selected location. In some embodiments, the in vivo radioisotope targeting chelate construct **120** is used to cause cell death at the selected location by delivering a

targeted dose of radiation. In some embodiments, the cells that are killed at the selected location are cancer cells. In some embodiments, the radiation is alpha radiation; beta or gamma radiation could be used in other embodiments.

[0086] In some embodiments, in vivo radioisotope targeting chelate construct **120** is

5 internalized by a cell within the mammalian subject, for example by endocytosis or otherwise. Thus in some embodiments, in vivo radioisotope targeting chelate construct **120** is present within a mammalian cell. In some embodiments, the in vivo radioisotope targeting chelate construct **120** is present within a human cell.

[0087] In some embodiments, the in vivo radioisotope targeting chelate construct **120** is

10 prepared prior to administration of construct **120** to a subject by combining an in vivo radioisotope targeting construct **130** having a targeting moiety **122**, a chelator **126** and optionally a linker **124** with a radionuclide **128** to form the in vivo radioisotope targeting chelate construct **120**. In some embodiments, the combining is carried out at a mild temperature, e.g. at a temperature in the range of about 10°C to about 65°C, including any
15 value therebetween e.g. 15°C, 16°C, 17°C, 18°C, 19°C, 20°C, 21°C, 22°C, 23°C, 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 35°C, 40°C, 45°C, 50°C, 55°C or 60°C. In some embodiments, the combining is carried out at a mild pH, e.g. an approximately neutral pH or an approximately physiological pH. In some embodiments, the mild pH is a pH of between about 5.0 and about 7.4, including any value therebetween e.g. 5.2, 5.4, 5.6, 5.8, 6.0, 6.2,
20 6.4, 6.6, 6.8, 7.0 or 7.4. In some embodiments, the mild pH is approximately 6.0. In some embodiments, the combining is carried out a physiological pH, e.g. in the range of about 7.0 to 7.4 including any value therebetween, e.g. 7.1, 7.2 or 7.3. In some embodiments, radionuclide **128** is combined with in vivo radioisotope targeting construct **130** in aqueous solution. In some embodiments, the aqueous solution is free or substantially free of
25 alcohols such as ethanol. In some embodiments, the combining is carried out for a period of between about 5 and about 30 minutes, including any value therebetween e.g. 10, 15, 20 or 25 minutes.

[0088] In some embodiments, in vivo targeting chelate construct **120** is used in diagnostic applications. For example, in vivo targeting chelate construct **120** may be administered to a
30 subject in any suitable manner, and any suitable imaging technology or procedure may be used to evaluate the localization of the targeting chelate construct **120** within the body via targeting moiety **122** by visualizing the location of bound radionuclide **128**, e.g. positron

emission tomography (PET) imaging or single-photon emission computerized tomography (SPECT) imaging. Such imaging procedures can be carried out for example to diagnose a subject as having a particular disorder or type of cancer, or to localize regions of the subject's body affected by the particular disorder or type of cancer. In some embodiments, localization of targeting chelate construct **120** to a target organ, region or plurality of loci within the body as evaluated by such imaging technology may be indicative that the subject has a particular form of cancer, and/or can be used to evaluate the extent of the cancer and or locations within the body wherein cancerous cells are or may be located, and/or can be used to evaluate the extent of metastasis of the cancer.

[0089] In some embodiments, constructs such as targeting chelate construct **120** are used in therapeutic applications, for example to carry out targeted radionuclide therapy. For example, targeting chelate construct **120** may be administered to a subject in any suitable manner, and the targeting effect imparted by targeting moiety **122** can be used to deliver the chelated radionuclide **128** to a desired location within the subject's body. In some embodiments, radiation from radionuclide **128** is used to kill cells at the desired location. In some embodiments, the cells that are killed at the desired location are cancer cells. In some embodiments, targeting construct **120** is used to perform targeted radionuclide therapy. In some embodiments, targeting construct **120** is used to perform targeted alpha therapy.

[0090] In some embodiments, a pharmaceutical composition is provided, the pharmaceutical composition comprising a construct such as targeting construct **120** and a pharmaceutically acceptable carrier. The pharmaceutical composition may include any suitable excipient, vehicle, buffer, diluent, binder, thickener, lubricant, preservative or the like, and may be provided in any desired state, e.g. as a liquid, suspension, emulsion, paste, or the like. In some embodiments, the pharmaceutical composition can be administered in any suitable manner, e.g. orally, intravenously, intramuscularly, subcutaneously, intraperitoneally, intratumorally, by inhalation, or the like.

[0091] In some embodiments, a method of prophylaxis and/or treatment of a subject having or believed to have cancer is provided. In some embodiments, the method comprises administering an in vivo targeting chelate construct **120** or a pharmaceutical composition comprising such a targeting chelate construct **120** to the subject. In some embodiments,

the method comprises administering a therapeutically and/or prophylactically effective amount of the targeting chelate construct **120** to the subject.

[0092] In some embodiments, the subject is a mammal. In some embodiments, the subject is a human. In alternative embodiments, the subject is livestock or a pet, e.g. a horse, cow, sheep, goat, cat, dog, rabbit, or the like. In some embodiments, the subject is a monkey.

[0093] While exemplary embodiments are described herein with reference to the targeting and killing of cancer cells, such constructs can be used for the selective killing and/or ablation of other undesired cell types, for example bacteria, fungi, cells implicated in autoimmune disorders, virus-infected cells, parasites, and so on.

[0094] In some embodiments, the metals that can be used as metal **128** include actinides, lanthanides, rare earth metals, or main group metals. In some embodiments, the lanthanide is La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb or Lu. In some embodiments, the lanthanide is Gd, Lu, Pr, Nd, Ho, Er or Yb. In some embodiments, the lanthanide is a radiolanthanide. In some embodiments, the actinide is Ac, Th, Pa, U, Np, Pu, Am, Cm, Bk, Cf, Es, Fm, Md, No or Lr. In some embodiments, the actinide is Ac, Th or U. In some embodiments, the actinide is a radioactinide. In some embodiments, the rare earth metal is Sc, Y, La, Ce, Pr, Nd, Pm, Sm, Gd, Tb, Dy, Ho, Er, Tm, Yb or Lu. In some embodiments, the metal is a trivalent lanthanide ion.

[0095] In some embodiments, the metal is a radioisotope. In some embodiments, the radioisotope is any desired radioisotope, e.g. ^{225}Ac , ^{227}Th , ^{226}Th , ^{211}At , ^{44}Sc , ^{90}Y , ^{89}Zr , ^{177}Lu , ^{111}In , $^{86/89/90}\text{Y}$, ^{211}At , ^{211}Fr , $^{212/213}\text{Bi}$, ^{153}Sm , $^{161/166}\text{Ho}$, $^{165/166}\text{Dy}$, $^{161/155}\text{Tb}$, ^{140}La , $^{142/143/145}\text{Pr}$, ^{159}Gd , $^{169/175}\text{Yb}$, $^{167/170}\text{Tm}$, ^{169}Er , ^{149}Pm , ^{150}Eu , ^{68}Ga , ^{137}Cs , ^{141}Ce , or the like.

[0096] In some embodiments, the metal is actinium (Ac), lutetium (Lu), bismuth (Bi), gallium (Ga), indium (In), terbium (Tb), thorium (Th), or Caesium (Cs). In some embodiments, the metal is actinium (III) (Ac^{3+}), lutetium (III) (Lu^{3+}), bismuth (III) (Bi^{3+}), gallium (III) (Ga^{3+}), indium (III) (In^{3+}), terbium (III) (Tb^{3+}), thorium (III) (Th^{3+}), or Cesium (I) (Cs^{1+}). In some embodiments, the metal is ^{225}Ac , ^{177}Lu , ^{213}Bi , ^{232}Th , ^{230}Th , ^{228}Th , ^{68}Ga , ^{161}Tb , ^{155}Tb , ^{152}Tb , ^{149}Tb , ^{111}In , or ^{137}Cs . In some embodiments, the metal is ^{227}Th , ^{225}Ac , ^{155}Tb , ^{177}Lu , ^{111}In , ^{132}La , ^{235}La , ^{90}Y , ^{68}Ga , ^{44}Sc , ^{203}Pb , or ^{212}Pb .

[0097] In some embodiments, $\text{H}_4\text{noneunpaX}$ is bound to a metal ion to form a coordination complex. In some embodiments, the coordination complex is referred to as a metal chelate.

In some embodiments, the metal chelate or H₄noneunpaX as the chelating ligand is associated with one or more cations as counter ions, for example Na⁺, K⁺, Ca²⁺ or the like. In some embodiments, the metal chelate or the chelating ligand is fully protonated. In some embodiments, the metal chelate or the chelating ligand is in its free acid form. In some
5 embodiments, the metal chelate or the chelating ligand is in a partially protonated state.


[0098] In some embodiments, the coordination complex is present in mammalian serum, optionally human serum. In some embodiments, the coordination complex is stable in mammalian serum, optionally human serum. In some embodiments, the coordination complex is present in mammalian serum within the body of the mammal, optionally present
10 in human serum within the body of the human. In some embodiments, the coordination complex is present in blood, optionally human blood. In some embodiments, the coordination complex is stable in mammalian blood, optionally human blood. In some embodiments, the coordination complex is present in mammalian blood within the body of the mammal, optionally present in human blood within the body of the human. In some
15 embodiments, the coordination complex is present within the body of a mammal, optionally present within the body of a human. In some embodiments, the coordination complex is present within a cell of a mammalian subject, optionally present within a cell of a human subject.

20 **Examples**

[0099] Further embodiments are described with reference to the following examples, which are intended to be illustrative and not limiting in nature.

Example 1.0 - Synthesis and Characterisation of *H*₄noneunpaX

[0100] The nonadentate chelating ligand, *H*₄noneunpa, has been previously reported^{8,13} and was synthesised following established procedures.⁸ Preparation of *H*₄noneunpaX was carried out in accordance with the linear synthetic route outlined in Scheme 1.



Scheme 1. Synthesis of *H*₄noneunpaX·4HCl·5H₂O from commercially available 2-(2-aminoethoxy)ethan-1-ol, over 5 steps with a cumulative yield of 33%.

[0101] The synthesis of *H*₄noneunpaX proceeds through *N* – alkylation of commercially available 2-(2-aminoethoxy)ethan-1-ol (**1**) with *tert* – butyl bromoacetate (*t*ba) under mild conditions to give di-*tert*-butyl ester (**2**) in high yields. Sequential mesylation of compound (**2**) followed by displacement with sodium azide, proceeded cleanly to give the corresponding azide (**3**) in high yields. Diisopropylethylamine (diPEA) was selected an appropriate base for the mesylation of compound (**2**) in order to precipitate the HCl equivalent generated in the reaction, and thus minimise formation of the corresponding chloride side product. Staudinger reduction of azide (**3**) was employed to generate the corresponding primary amine (**4**) which could be used directly without additional purification. Subsequent *N*-alkylation of amine (**4**) with two equivalents of methyl 6-(bromomethyl)picolinate afforded the protected ligand (**5**) in modest yields. DIPEA was found to be a superior base over K₂CO₃ in this reaction. Attempts to isolate compound (**5**) while using K₂CO₃ as a base were hampered by difficulties in purification, presumably due to the product chelating free K⁺ ions from solution. This difficulty

was mitigated by using an organic base and generated the product in high yield and high purity. Final deprotection of compound **(5)** was achieved through ester hydrolysis in 4 M HCl, followed by reverse phase high performance liquid chromatography (HPLC). Co-evaporation of the purified ligand with 3 M HCl was then used to afford H₄noneunpaX **(6)** as a HCl salt.

- 5 **[0102]** The purified ligand and all synthetic intermediates were fully characterised by NMR spectroscopy (¹H, ¹³C{¹H}, COSY, HSQC) and mass spectrometry (LR/HR-MS). Elemental analysis (EA) and HPLC of the final ligand was performed in order to confirm the purity of the isolated product. Results for H₄noneunpaX are shown in FIGs. 3-7.

- [0103] Di-*tert*-butyl 2,2'-((2-(2-hydroxyethoxy)ethyl)azanediyl)diacetate (2):** *tert*-Butyl bromoacetate (4.30 mL, 5.57 g, 28.6 mmol) was added slowly to a solution of 2-(2-aminoethoxy)ethan-1-ol **(1)** (1.53 g, 14.3 mmol) and diisopropylethylamine (5.00 mL, 3.69 g, 28.6 mmol) dissolved in dry MeCN (150 mL). The reaction mixture was heated to 50 °C and stirred overnight. Upon completion, the volatiles were removed *in vacuo* and the resulting residue re-dissolved in EtOAc (150 mL). After standing for 10 min at RT, the white precipitate formed was removed *via* vacuum filtration, washed with cool EtOAc (50 mL) and discarded. The filtrate was washed with de-ionised water (3 x 150 mL) and the combined aqueous phase back extracted with EtOAc (2 x 150 mL). The combined organics were evaporated *in vacuo* to yield the title compound as a pale-yellow oil (4.69 g, 99%). ¹H NMR (300 MHz, CDCl₃, 298 K) 3.66 (2H, t, *J* = 4.6 Hz, 1 – CH₂), 3.60 (2H, t, *J* = 5.2 Hz, 3 – CH₂), 3.52 (2H, t, *J* = 4.6 Hz, 2 – CH₂), 3.47 (4H, s, 5 – CH₂), 2.96 (1H, br s, 7 – OH), 2.92 (2H, t, *J* = 5.2 Hz, 4 – CH₂), 1.42 (18H, s, 6 – C(CH₃)₃). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 171.0 (6 – C), 81.2 (7 – C), 72.4 (2 – C), 70.2 (3 – C), 61.9 (1 – C), 56.8 (5 – C), 53.4 (4 – C), 28.3 (8 – C). ESI-MS (MeOH) 334.26 [M+H]⁺.
- 10
15
20

- [0104] Di-*tert*-butyl 2,2'-((2-(2-azidoethoxy)ethyl)azanediyl)diacetate (3):** Methanesulphonyl chloride (1.34 mL, 1.98 g, 17.3 mmol) was added slowly to a solution of di-*tert*-butyl 2,2'-((2-(2-hydroxyethoxy)ethyl)azanediyl)diacetate **(2)** (5.26 g, 15.7 mmol) and diisopropylethylamine (3.01 mL, 2.24 g, 17.3 mmol) in EtOAc (15 mL) at 0 °C. After 10 minutes, the suspension was warmed to RT and stirred for a further 3 h. Upon completion, the suspension was cooled to 0 °C and the white solid separated by vacuum filtration. The filtrate was diluted with EtOAc (50 mL) and washed with de-ionised H₂O (3 x 50 mL). The organic phase was dried over Na₂SO₄ and evaporated *in vacuo* to yield the corresponding mesylate as a pale-yellow oil which was used without further purification.
- 25
30

[0105] NaN₃ (3.08 mg, 47.2 mmol) was added to a solution of crude mesylate dissolved in dry DMF (20 mL) and the suspension heated at 80 °C O/N. Upon completion, the solution was cooled to RT and diluted with de-ionised H₂O (50 mL). The aqueous phase was extracted with DCM (3 x 50 mL) and the combined organics washed with 5% LiCl aq. sol.

(50 mL). The volatiles were removed *in vacuo* and the resulting residue purified *via* silica gel chromatography (Combiflash automated purification system; A: DCM, B: MeOH; 100% to 5% B). The title compound was attained as a pale-yellow oil (3.25 g, 58%). ¹H NMR (300 MHz, CDCl₃, 298 K) 3.62 – 3.57 (4H, m, 2 – and 3 – CH₂), 3.46 (4H, s, 5 – CH₂), 3.33 (2H, t, *J* = 5.0 Hz, 1 – CH₂), 2.92 (2H, t, *J* = 5.6 Hz, 4 – CH₂), 1.42 (18H, s, 6 – (C(CH₃)₃)). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 170.9 (6 – C), 81.0 (7 – C), 70.9 (3 – C), 69.8 (2 – C), 56.9 (5 – C), 53.6 (4 – C), 50.8 (1 – C), 28.3 (8 – C). ESI-MS (MeOH) 359.21 [M+H]⁺.

[0106] Di-*tert*-butyl 2,2'-((2-(2-aminoethoxy)ethyl)azanediyl) diacetate (**4**): Di-*tert*-butyl 2,2'-((2-(2-aminoethoxy)ethyl)azanediyl) diacetate (**3**) (2.81 g, 7.85 mmol) was dissolved in dry THF (30 mL) and cooled to 0 °C. PPh₃ (2.47 mg, 9.42 mmol) was added slowly to the reaction mixture under Ar, and the resulting solution allowed to warm to RT and stirred for 5 h. The resulting solution was then added dropwise into de-ionised H₂O (350 mL) and the suspension stirred overnight. The THF was removed *in vacuo* and the off-white suspension left to precipitate at RT for 1 h. The white precipitate was removed *via* vacuum filtration, and the aqueous phase concentrated to ~ 100 mL *in vacuo*. The aqueous phase was extracted with CH₂Cl₂ (4 x 75 mL) and the combined organic phase evaporated *in vacuo* to give the title compound as a pale-yellow oil (2.38 g, 91%). ¹H NMR (300 MHz, CDCl₃, 298 K) 6.72 (3H, br s, 7 – NH₂), 3.71 (2H, t, *J* = 4.5 Hz, 2 – CH₂), 3.58 (2H, t, *J* = 5.1 Hz, 3 – CH₂), 3.10 (2H, t, *J* = 4.5 Hz, 1 – CH₂), 2.91 (2H, t, *J* = 5.1 Hz, 4 – CH₂), 1.42 (18H, s, 6 – (C(CH₃)₃)). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 171.0 (6 – C), 81.6 (7 – C), 68.8 (3 – C), 67.9 (2 – C), 56.2 (5 – C), 53.3 (4 – C), 40.5 (1 – C), 28.3 (8 – C). ESI-MS (MeOH) 333.19 [M+H]⁺.

[0107] (tBu)₂(Me)₂noneunpaX (**5**). Methyl (6-bromomethyl) picolinate (515 mg, 2.824 mmol) was added to a solution of di-*tert*-butyl 2,2'-((2-(2-aminoethoxy)ethyl)azanediyl) diacetate (**4**) (372 mg, 1.12 mmol) in dry MeCN (15 mL) under argon. Diisopropylethylamine (585 µL, 434 mg, 3.36 mmol) was added and the resulting pale-yellow solution stirred for 1 h at ambient temperature. The reaction mixture was subsequently heated to 50 °C and stirred overnight. Upon completion, the volatiles were removed *in vacuo* and the resulting residue was re-dissolved in CH₂Cl₂ (25 mL). The organic phase was washed with de-

ionised H₂O (3 x 25 mL) and brine (25 mL), dried over Na₂SO₄ and evaporated *in vacuo*.

The crude residue was purified *via* flash column chromatography on neutral alumina

(Combiflash automated purification system; A: CH₂Cl₂, B: MeOH; 100% A to 5% B). The

product was attained as a pale-yellow oil (459 mg, 65%). ¹H NMR (300 MHz, CDCl₃, 298 K)

5 7.90 (2H, dd, ³J = 7.3 Hz, ⁴J = 1.1 Hz, 10 – CH), 7.80 (2H, dd, ³J = 8.1 Hz, ⁴J = 1.4 Hz, 8 – CH), 7.75 (2H, t, ³J = 7.5 Hz, 9 – CH), 3.94 (4H, s, 7 – CH₂), 3.90 (6H, s, 11 – CH₃), 3.49 (2H, t, ³J = 5.6 Hz, 2 – CH₂), 3.44 (2H, t, ³J = 5.8 Hz, 3 – CH₂), 3.40 (4H, s, 5 – CH₂), 2.84 (2H, t, ³J = 5.8 Hz, 4 – CH₂), 2.74 (2H, t, ³J = 5.6 Hz, 1 – CH₂), 1.35 (18H, s, 6 – C(CH₃)₃). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 170.8 (8 – C), 165.9 (15 – C), 160.8 (14 – C), 147.3 (10 – C), 137.5 (12 – C), 126.1 (11 – C), 123.6 (13 – C), 80.9 (7 – C), 70.5 (3 – C), 69.4 (2 – C), 60.9 (9 – C), 56.7 (5 – C), 54.1 (1 – C), 53.6 (4 – C), 52.9 (16 – C), 28.2 (8 – C). ESI-MS (MeOH) 653.3 [M+Na]⁺.

[0108] H₄noneunpaX·4HCl·5H₂O (6). (tBu)₂(Me)₂noneunpaX (**5**) (168 mg, 0.257 mmol) was dissolved in 4 M HCl (5 mL) and heated at 60 °C overnight. The volatiles were removed *in*

15 *vacuo* and the resulting off-white solid purified *via* RP-HPLC (A: H₂O (0.1% TFA), B: MeCN; 100% A to 20% B, 30 min., t_R ~ 21.5 min). Pure H₄noneunpaX was attained as a white HCl salt (146 mg, 87%) by co-evaporation with 3 M HCl. ¹H NMR (400 MHz, D₂O, 298 K) 7.95 (2H, d, ³J = 7.7 Hz, 1 – CH), 7.89 (2H, t, ³J = 7.7 Hz, 2 – CH), 7.57 (2H, d, ³J = 7.7 Hz, 3 – CH), 4.76 (4H, s, 3 – CH₂), 4.21 (4H, s, 9 – CH₂), 3.94 (2H, t, ³J = 4.4 Hz, 6 – CH₂), 3.84 (2H, t, ³J = 4.4 Hz, 7 – CH₂), 3.75 (2H, t, ³J = 4.4 Hz, 5 – CH₂), 3.63 (2H, t, ³J = 4.4 Hz, 8 – CH₂). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 167.9 (13 – C), 166.7 (1 – C), 149.7 (6 – C), 146.3 (2 – C), 140.1 (3 – C), 128.5 (5 – C), 125.4 (4 – C), 65.3 (10 – C), 64.6 (9 – C), 58.6 (7 – C), 56.2 (11 – C), 56.0 (8 – C), 55.5 (12 – C). ESI-MS (H₂O) 491.1 [M+H]⁺. HR-ESI-MS (H₂O) calcd. for [C₂₂H₂₆N₄O₉+H]⁺: 491.1700; found [M+H]⁺: 491.1779. Elemental analysis: 25 calcd. % for H₄noneunpaX·4HCl·5H₂O (C₂₂H₂₆N₄O₉·4HCl·5H₂O = 726.178 g mol⁻¹): C 37.59, H 5.36, N 7.97; found: C 37.65, H 5.43, N 7.97.

Example 2.0 - Metal Complexation and NMR Characterisation

[0109] The coordination characteristics of H₄noneunpaX were investigated through

30 complexation of a series of non-radioactive trivalent metal ions (La³⁺, Lu³⁺ and In³⁺) and assessment of the changes in spectral features as a function of metal ionic radius using

NMR spectroscopy (^1H , COSY). Metal complexes of $\text{H}_4\text{noneunpaX}$ were prepared by mixing equimolar quantities of the ligand with the appropriate metal salts in D_2O and adjustment of the pD to neutral using NaOD (0.1 M). The corresponding solutions were then filtered, and analysed directly via ^1H NMR spectroscopy, FIG. 8. Further confirmation of metal complexation was achieved using high resolution electrospray ionisation mass spectrometry (HR-ESI-MS), wherein each metal complex was confirmed by the presence of a monocationic peak corresponding to the $[\text{M}+2\text{H}]^+$ species, Table 2.

Table 2. High-resolution electrospray ionisation mass spectrometry characterisation of metal complexes of $\text{H}_4\text{noneunpaX}$ with La^{3+} , Lu^{3+} and In^{3+} .

Metal Complex	Theoretical Mass $[\text{M}+2\text{H}]^+$	Experimental Mass $[\text{M}+2\text{H}]^+$
$[\text{La}(\text{noneunpaX})]^-$	627.0536	627.0600
$[\text{Lu}(\text{noneunpaX})]^-$	663.0873	663.0946
$[\text{In}(\text{noneunpaX})]^-$	603.0506	603.0572

[0110] Full ^1H NMR characterisation data (^1H , COSY) was obtained.

[0111] $\text{Na}[\text{La}(\text{noneunpaX})]$. Results shown in FIGs. 9-12. $\text{La}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ (2.9 mg, 6.75 μmol , 1.0 equiv.) was added directly to a solution of $\text{H}_4\text{noneunpaX} \cdot 4\text{HCl} \cdot 5\text{H}_2\text{O}$ (4.9 mg, 6.75 μmol , 1.0 equiv.) in D_2O (300 μL). The solution was mixed thoroughly at 1000 rpm using a vortex mixer, and the pD adjusted to ~ 7.0 using dilute NaOD solution to give the corresponding $[\text{La}(\text{noneunpaX})]^-$ complex. The white precipitate was filtered, and the filtrate characterized directly without additional purification. ^1H NMR (400 MHz, D_2O , 298 K, pD 7.0) 7.87 (2H, t, $^3J = 7.7$ Hz, i – and m – CH), 7.77 (2H, d, $^3J = 7.5$ Hz, j – and n – CH), 7.42 (2H, d, $^3J = 7.7$ Hz, h – and l – CH), 4.19 (2H, d, $^2J = 16.4$ Hz, g – and k – CH), 3.93 (2H, d, $^2J = 16.4$ Hz, g' – and k' – CH), 3.53 (2H, t, $^3J = 5.0$ Hz, d – CH_2), 3.48 (2H, d, $^3J = 4.9$ Hz, e – CH_2), 3.37 (2H, d, $^2J = 16.3$ Hz, a – and b – CH), 3.31 (2H, d, $^2J = 16.3$ Hz, a' – and b' – CH), 3.04 (2H, br t, f – CH_2), 2.71 (2H, t, $^3J = 5.0$ Hz, c – CH_2). LR-ESI-MS (H_2O) 627.0 $[\text{M}+2\text{H}]^+$, 624.8 $[\text{M}]^-$. HR-ESI-MS (H_2O) calcd. for $[\text{C}_{22}\text{H}_{22}\text{LaN}_4\text{O}_9+2\text{H}]^+$: 627.0536; found $[\text{M}+2\text{H}]^+$: 627.0600.

[0112] $\text{Na}[\text{Lu}(\text{noneunpaX})]$. Results shown in FIGs. 13-16. $\text{Lu}(\text{NO}_3)_3 \cdot \text{H}_2\text{O}$ (2.4 mg, 6.75 μmol , 1.0 equiv.) was added directly to a solution of $\text{H}_4\text{noneunpaX} \cdot 4\text{HCl} \cdot 5\text{H}_2\text{O}$ (4.9 mg, 6.75

μmol, 1.0 equiv.) in D₂O (300 μL). The solution was mixed thoroughly at 1000 rpm using a vortex mixer, and the pD adjusted to ~ 6.5 using dilute NaOD solution to give the corresponding [Lu(noneunpaX)]⁻ complex. The solution was filtered and analyzed without additional purification. ¹H NMR (400 MHz, D₂O, 298 K, pD 6.5) 8.03 (1H, t, ³J = 7.8 Hz, i – CH), 7.96 (1H, t, ³J = 7.7 Hz, m – CH), 7.90 (1H, d, ³J = 7.7 Hz, n – CH), 7.83 (1H, d, ³J = 7.8 Hz, j – CH), 7.61 (1H, d, ³J = 7.8 Hz, h – CH), 7.43 (1H, d, ³J = 7.7 Hz, l – CH), 4.95 (1H, d, ²J = 15.8 Hz, g – CH), 4.17 (1H, d, ²J = 15.8 Hz, g' – CH), 4.01 (1H, d, ²J = 15.6 Hz, k – CH), 3.97 (1H, d, ²J = 17.4 Hz, a – CH), 3.90 (1H, d, ²J = 17.4 Hz, a' – CH), 3.77 (1H, ddd, c – CH), 3.66 (1H, dd, d – CH), 3.45 (1H, d, ²J = 18.4 Hz, b – CH), 3.43 – 3.40 (1H, m, e – CH), 3.36 – 3.24 (2H, m, e' – and d' – CH), 3.27 (1H, d, ²J = 15.6 Hz, k' – CH), 3.19 (1H, d, ²J = 18.4 Hz, b' – CH), 3.10 (1H, ddd, f – CH), 2.80 (1H, d, ²J = 12.9 Hz, f' – CH), 2.65 (1H, dd, c' – CH). LR-ESI-MS (H₂O) 663.1 [M+2H]⁺, 661.1 [M]⁻. HR-ESI-MS (H₂O) calcd. for [C₂₂H₂₂LuN₄O₉+2H]⁺: 663.0873; found [M+2H]⁺: 663.0946.

[0113] Na[In(noneunpaX)]. Results shown in FIGs. 17 and 18. In(NO₃)₃·H₂O (2.1 mg, 6.75 μmol, 1.0 equiv.) was added directly to a solution of H₄noneunpaX·4HCl·5H₂O (4.9 mg, 6.75 μmol, 1.0 equiv.) in D₂O (300 μL). The solution was mixed thoroughly at 1000 rpm using a vortex mixer, and the pD adjusted to ~ 4.5 using dilute NaOD solution to give the corresponding [In(noneunpaX)]⁻ complex. The solution was filtered and analyzed without additional purification. ¹H NMR (400 MHz, D₂O, 298 K, pD 5.5) 8.32 (2H, t, ³J = 5.2 Hz, i – and m – CH), 8.28 (2H, d, ³J = 5.0 Hz, j – and n – CH), 7.88 (2H, d, ³J = 5.0 Hz, h – and l – CH), 4.82 (2H, d, ²J = 10.7 Hz, g and k – CH₂), 4.28 (2H, d, ²J = 10.7 Hz, g' – and k' – CH₂), 3.74 (4H, s, a/a' – and b/b' – CH₂), 3.56 (4H, br s, d/d' – and e/e' – CH₂), 3.38 (2H, br s, f/f' – CH₂), 2.80 (2H, br s, c/c' – CH₂). LR-ESI-MS (H₂O) 603.0 [M+2H]⁺, 625.0 [M+H+Na]⁺. HR-ESI-MS (H₂O) calcd. for [C₂₂H₂₂InN₄O₉+2H]⁺: 603.0506; found [M+2H]⁺: 603.0572.

[0114] The ¹H NMR spectrum of [La(noneunpaX)]⁻ shows the formation of a single, symmetric isomer in solution, with characteristic diastereotopic splitting for the methylenic protons associated with the four pendent donor arms of the ligand, implying coordination of all four donor groups to the metal centre. Sharply resolved resonances are observed for the protons within the ethylene bridged backbone which indicates the formation of a metal complex with the central ethereal oxygen coordinated. In contrast, the NMR spectra for the Lu³⁺ complex reveal a single, asymmetric isomer, wherein chemically distinct resonances for each picolinic acid donor arm are seen in the aromatic region. Notably, one picolinate

donor arm (H_i , H_m , H_n) maintains the same chemical shift as seen in the La^{3+} complex, while the second picolinate arm (H_h , H_i , H_j) exhibits a significant downfield shift, indicating closer coordination to the Lu^{3+} metal centre and a change in ligand conformation at this position. This aspect is further observed for the adjacent methylenic protons (H_g and $H_{g'}$) which exhibit distinctive diastereotopic splitting, with characteristically large coupling constants ($^2J_{AB} = 15.8$ Hz), and appear in two distinct chemical environments ($\Delta\delta_{g/g'} = 0.78$ ppm). The remaining methylenic protons associated with the pendent donor arms all also become inequivalent on metal ion complexation, appearing as diastereotopic doublets, which can be clearly distinguished in the $^1H - ^1H$ COSY NMR spectrum, FIGs. 14-16. In contrast to $[La(noneunpaX)]^-$, the Lu^{3+} complex shows further splitting for all eight protons in the backbone, with large differences in chemical shift for mutually coupled protons (e.g. H_c and $H_{c'}$) which suggests a rigid complexation environment and close coordination to the metal centre.

[0115] The metal complexes of $H_4noneunpaX$ share similar spectral characteristics to those reported for $H_4noneunpa$; whereby fully saturated metal coordination spheres were achieved with La^{3+} and Lu^{3+} ions, as indicated by diastereotopic splitting of the pendent donor arm resonances and backbone ethylene bridges. Interestingly, the opposite trend in metal complex symmetry was attained for $H_4noneunpa$, wherein the $[La(noneunpa)]^-$ complex appears as an asymmetric isomer, while the $[Lu(noneunpa)]^-$ complex is fully symmetric.⁸

[0116] Analysis of the $[In(noneunpaX)]^-$ complex using NMR spectroscopy was hampered by low solubility at neutral and basic pH, therefore the ^1H NMR spectrum in FIG. 8 was recorded under acidic conditions (pH ~ 4.0). The spectrum obtained shows the formation a different symmetric isomer in which both picolinate donors are coordinated to the metal centre, while the two acetate donor arms remain unbound, as indicated by the presence of a 4H singlet resonance at 3.50 ppm corresponding to $H_{a/a'}$ and $H_{b/b'}$. This assignment is also supported by the resonances observed for the backbone methylene protons, which appear as broad signals in the NMR spectrum. At higher pH values (pH $\sim 5.0 - 8.5$) a second species was observed, which may correspond to the fully coordinated complex, however a complete spectral analysis was not possible.

Example 3.0 - Solution Thermodynamic Stability Studies and Complexation Equilibria with Sc^{3+} , In^{3+} , Lu^{3+} , Dy^{3+} , Gd^{3+} , Sm^{3+} , La^{3+}

[0117] Prior to investigation of the thermodynamic stability of different metal complexes, the protonation constants of the chelating ligand must be determined independently, due to the competition between a given metal ion with a proton for the same coordinating group in the metal complex reaction. Combined potentiometric and UV-vis spectrophotometric titrations were used to evaluate the protonation constants of H₄noneunpaX between pH ~2–11.5, while acidic in-batch UV-vis spectrophotometric titrations were used to determine the protonation constants for the most acidic protons, which were below the threshold of the pH electrode (pH<2). All eight protonation constants for H₄noneunpaX were determined through refinement of the experimental data using HyperSpec²⁴ and Hyperquad²⁵, Table 3.

Table 3. Protonation constants ($\log K_{HqL}$)^a of H₄noneunpaX at 25 °C and $I = 0.16$ M NaCl.

Reaction	$\log \beta$	$\log K$
$L + H^+ \leftrightarrow HL$	8.84 (1)	8.84
$HL + H^+ \leftrightarrow H_2L$	15.92 (1)	7.08
$H_2L + H^+ \leftrightarrow H_3L$	19.71 (1)	3.79
$H_3L + H^+ \leftrightarrow H_4L$	22.54 (2)	2.83
$H_4L + H^+ \leftrightarrow H_5L$	24.92 (3)	2.38
$H_5L + H^+ \leftrightarrow H_6L$	27.06 (4)	2.14
$H_6L + H^+ \leftrightarrow H_7L$	28.07 (4)	1.01
$H_7L + H^+ \leftrightarrow H_8L$	27.35 (6)	-0.72 ^b
$\Sigma \log K_{HqL}$	27.35 (6)	27.35 (6)

^a K_{HqL} defined as $[H_qL]/([H][H_{q-1}L])$. ^b not evaluated at constant $I = 0.16$ M NaCl. Charges omitted for simplicity. Values in brackets are the standard deviations.

[0118] The protonation constants obtained for H₄noneunpaX follow the typical trend observed for similar polyaminocarboxylate-based ligands;^{6,12,13} wherein the first two dissociation events (species H₈L⁴⁺ and H₇L³⁺) can be attributed to the protonated pyridyl nitrogen donors. Deprotonation of each successive pyridine donor, with pK_a -0.72 and 1.01 respectively, is concurrent with spectral changes in the absorbance spectrum of the ligand, where a large decrease in the absorptivity of the ligand is observed for the first dissociation event, FIGs. 19 and 20. Deprotonation of the two acetate carboxylic groups ($\log K_6 = 2.14$

and $\log K_5 = 2.38$), is followed by the two picolinate donors ($\log K_4 = 2.83$ and $\log K_3 = 3.79$), while the final two deprotonations are assigned to dipyridyl ($\log K_2 = 7.08$) and iminodiacetate ($\log K_1 = 8.84$) terminal amines respectively. The protonation constant measured for species HL is comparable to other ligands containing the iminodiacetate moiety (e.g. *N*-(2-hydroxyethyl)iminodiacetic acid (HEIDA), $\log K_1 = 8.68$).²⁶ In contrast, the second terminal amine in H₄noneunpaX is less basic ($\log K_2 = 7.08$) than the equivalent donor group in H₄noneunpa/Oxyaapa ($\log K_2 = 7.63$)¹³ which can be attributed to the stronger electron-withdrawing effect induced by placing two picolinate donors on the same nitrogen atom, in addition to the steric constraints at this position, which may disfavour protonation. The larger difference in basicity between the terminal amines of H₄noneunpaX, in comparison to H₄noneunpa/Oxyaapa, may be advantageous for metal ion coordination, whereby the less basic dipyridyl nitrogen favours complexation at a lower pH, while the more basic iminodiacetate nitrogen will act as a stronger donor group to metal ions.

[0119] To evaluate the affinity of H₄noneunpaX towards metal ions of medicinal importance – e.g. Sc³⁺, In³⁺, Lu³⁺, Dy³⁺, Gd³⁺, Sm³⁺, La³⁺ – the complex formation equilibria were studied using both combined potentiometric-spectrophotometric titrations, and in all cases, acidic in-batch UV-vis spectrophotometric titrations below pH 2. H₄noneunpaX showed a high affinity to all metal ions investigated, with metal complexation observed to begin below pH ≈ 1 . Table 4 presents the stability constants with each metal ion.

Table 4. Stability constants ($\log K_{ML}$, $\log K_{MHL}$ and $\log K_{M(OH)L}$)^a of H₄noneunpaX with M³⁺ = Sc³⁺, In³⁺, Lu³⁺, Dy³⁺, Gd³⁺, Sm³⁺, La³⁺ metal ions at 25 °C and *I* = 0.16 M NaCl.

	Sc ³⁺	In ³⁺	Lu ³⁺	Dy ³⁺	Gd ³⁺	Sm ³⁺	La ³⁺
$\log K_{ML}$	23.57(2)	27.64(2)	21.93(3)	21.99(5)	20.87(4)	21.04(2)	18.45(4)
$\log K_{MHL}$	3.07(1)	6.17(3)	2.28(3)	2.29(2)	2.80(1)	2.77(1)	3.22(2)
$\log K_{M(OH)L}$	10.89(2)	8.58(4)	-	-	-	-	-
$\log K_{M(OH)2L}$	-	9.06(4)	-	-	-	-	-
pM ^b	22.9	26.5	21.3	21.3	20.2	20.4	17.8

^a K_{ML} defined as $[ML]/([M][L])$; K_{MHL} defined as $[MHL]/([ML][H])$; $K_{M(OH)L}$ defined as $[M(OH)L]/([ML][OH])$ (charges omitted for clarity). ^b pM defined as $-\log[M_{free}]$ when $[L] = 10 \mu M$; $[M] = 1 \mu M$ at pH 7.4.

[0120] H₄noneunpaX shows a high thermodynamic preference for complexation with trivalent Ln³⁺ ions, as indicated by the high stability constants (log *K*_{ML}) and p*M* values, Table 4. Notably, these values are directly comparable to those previously determined for H₄noneunpa/ Oxyaapa,¹³ with H₄noneunpaX exhibiting marginally higher stability constants across the Ln³⁺ series, implying a small thermodynamic benefit to this inverted structural arrangement (which may be a result of a degree of preorganisation of the binding cavity of the ligand). Furthermore, the p*M* values obtained for H₄noneunpaX with Ln³⁺ ions exceed those of the current gold-standard chelators DOTA and DTPA by several log units (e.g. p*Lu* = 17.1 and 19.1 respectively).¹¹ While the stability constants and p*M* values are useful parameters for comparing the metal scavenging abilities of different chelating ligands, they do not necessarily correlate well with *in vivo* stability, another aspect for consideration is the speciation behaviour of a given metal complex as a function of pH.

[0121] Speciation plots of the metal complexation of H₄noneunpaX with trivalent lanthanide ions show an interesting relationship, with the formation of the protonated MHL species at acidic pH, which is followed by a single transformation to the dominant [ML]⁻ species across a broad pH range, and the absence of any hydroxo species even under very basic conditions (pH = 11.5), FIGs. 21-27. This is in contrary to many similar chelating ligands with Ln³⁺ ions previously reported – H₄octapa, H₄pypa, H₄octox – which all show the formation of one or more hydroxo species (e.g. [M(OH)L], [M(OH)₂L]) under basic conditions (pH > 9.0) and is typically attributed to deprotonation of a coordinated water molecule in the complex.^{7,9,12} Under physiological conditions (pH 7.4), a single species ([ML]⁻) is observed for all Ln³⁺ complexes of H₄noneunpaX, which is favourable for *in vivo* applications as different isomers can exhibit varying pharmacokinetic characteristics and biodistribution profiles.

[0122] Combined potentiometric-spectrophotometric titrations of H₄noneunpaX with smaller trivalent metal ions show a different relationship to the Ln³⁺ series. Higher stability constants were determined for Sc³⁺ and In³⁺, compared to Ln³⁺, as is typically observed owing to the higher charge density of these metal centres, Table 4. In the case of Sc³⁺, complex formation occurs below pH ≈ 0.5 to give the neutral MHL species, which is deprotonated with increasing pH to give the anionic [ML]⁻ species. This transformation occurs in a similar range to the Ln³⁺ complexes of H₄noneunpaX (p*K*_a = 2.28 – 3.22) and may be attributed to the deprotonation of one of the carboxylate donors. In contrast, the In³⁺ complex shows

different thermodynamic behaviour in solution. The MHL species deprotonates with a higher pK_a to form the anionic $[ML]^-$ species ($pK_a = 6.17(3)$) which is assigned to the tertiary amine of the iminodiacetate moiety, which remains protonated early on in the complexation. This is consistent with the NMR characterisation of the $[In(noneunpaX)]^-$ complex which showed a

5 single species at pH 4.0 in which the two acetate donor groups remain unbound to the metal centre. As in the case of the Sc^{3+} complex, the formation of hydroxo species is observed at higher pH, which without being bound by theory is accounted for by considering the mismatch between the size of these small metal ions and the ligand binding cavity.

[0123] All potentiometric titrations were carried out with a Metrohm Titrand 809 and a

10 Metrohm Dosino 800 with a Ross combined electrode. Direct titrations were recorded using a Varian Cary 60 UV-vis spectrophotometer (200-350 nm spectral range) equipped with an optic dip probe (0.2 cm path length), while acidic in-batch experiments were measured with a standardised glass cuvette (1 cm path length). A temperature-controlled (298 K) 20 mL glass cell with an inlet-outlet adapter for nitrogen gas purging (purified *via* a 10% NaOH

15 solution to exclude CO_2 prior to and during each titration) was used as a titration cell. The electrode was calibrated daily in hydrogen ion concentration by direct titration of HCl with freshly prepared NaOH solution, and the results were analysed with the Gran procedure⁵⁵ to determine the standard potential (E°) and the ionic product of water (pK_w), at 298 K, with 0.16 M NaCl as a supporting electrolyte. Solutions under study were titrated with carbonate-

20 free NaOH solution (~ 0.16 M) which was standardized against freshly recrystallized potassium hydrogen phthalate.

[0124] The protonation equilibria of $H_4noneunpaX$ were evaluated through combined potentiometric-spectrophotometric titrations of solutions containing the ligand ($[L] = 9.57 \times 10^{-4}$ M) in 0.16 M NaCl ($T = 298$ K, $l = 0.2$ cm). Electromotive force (EMF) values and UV-

25 vis spectra were recorded after each addition of NaOH, and the apparatuses synchronized to have consistent time-intervals (30 s) between each addition/equilibration and data acquisition. Further acidic in-batch UV-vis spectrophotometry studies were performed to obtain the protonation constants for the most acidic ionisable protons of $H_4noneunpaX$, which were below the threshold of the pH electrode ($pH < 2$). Separate solutions of the

30 ligand were prepared at constant molarity ($[L] = 1 \times 10^{-4}$ M) with differing amounts of HCl (0.1 M and 3.0 M), while maintaining a constant ionic strength (0.16 M NaCl) wherever possible. In each case, the solution stoichiometry was used to calculate the equilibrium

concentration of H^+ ions, following the Hammett acidity function (H_0) for determination of acidity in high concentration acid solutions. All eight protonation constants for $H_4noneunpaX$ were determined through refinement of the experimental data using HypSpec2014²⁴ and Hyperquad2013²⁵ software.

- 5 **[0125]** The complex formation equilibria of $H_4noneunpaX$ with Sc^{3+} , In^{3+} , Lu^{3+} , Dy^{3+} , Gd^{3+} , Sm^{3+} , and La^{3+} were evaluated using both combined potentiometric-spectrophotometric titrations, and acidic in-batch UV-vis spectrophotometric titrations below pH 2. For combined potentiometric-spectrophotometric titrations (spectral range: 200 – 350 nm), solutions containing a 1:1 molar ratio of metal-to-ligand ($[M^{3+}] \approx [H_4noneunpaX] \approx 8.40 \times 10^{-4}$ M) were
- 10 titrated against NaOH solution at $T = 298$ K, $I = 0.2$ cm. Acidic in-batch UV-vis spectrophotometric measurements (spectral range: 200 – 350 nm) were carried out using sets of solutions containing a 1:1 molar ratio of metal-to-ligand ($[M^{3+}] \approx [H_4noneunpaX] \approx 1 \times 10^{-4}$ M) with differing amounts of HCl (0.1 M and 3.0 M), to give samples ranging from pH 0 – 2. Measurements were recorded at $T = 298$ K, $I = 1.0$ cm, the Hammett acidity function
- 15 (H_0) was used determination of the pH in highly acidic solutions rather than electrode potentials. Metal solutions were prepared from atomic absorption spectroscopy (AAS) standards, which were previously evaluated using Gran's method of titrating equimolar solutions of a given metal ion and Na_2H_2EDTA against NaOH to determine the acid concentration.
- 20 **[0126]** All the potentiometric measurements were processed using the Hyperquad2013²⁵ software, while spectroscopic measurements were analysed using HypSpec2014.²⁴ Proton dissociation constants corresponding to hydrolysis of the Sc^{3+} , In^{3+} , Lu^{3+} , Dy^{3+} , Gd^{3+} , Sm^{3+} , and La^{3+} aqueous ions used in the calculations were taken from Baes and Mesmer. Speciation diagrams were generated with the calculated protonation constants and stability
- 25 constants using Hyss software.⁵⁶

- [0127]** Each species generated in the investigated systems can be defined in accordance with the general equilibrium equation $pM + qH + rL = M_pH_qL_r$ (charges omitted). Wherein, a complex containing a metal ion, M, proton, H and ligand, L, has the general formula $M_pH_qL_r$. The stoichiometric index p may also be 0 in the case of protonation equilibria, and negative
- 30 values of q refers to proton removal from coordinated water, equivalent to hydroxide ion addition during formation of the complex. The overall equilibrium constant for the formation of the complexes $M_pH_qL_r$ from its components is designated as $\log \beta$. Stepwise equilibrium

constants $\log K$ correspond to the difference in log units between the overall constants of sequentially protonated (or hydroxide) species. pM is defined as $(-\log[M^{n+}]_{\text{free}})$ and is calculated from the stability constants obtained for each studied system at $[M^{n+}] = 1 \mu\text{M}$, $[L^{x-}] = 10 \mu\text{M}$, pH 7.4 and 25 °C.⁵⁷

5

Example 4.0 - Radiolabelling Studies

[0128] Radiolabelling studies with $[^{44}\text{Sc}]\text{Sc}^{3+}$, $[^{111}\text{In}]\text{In}^{3+}$, $[^{177}\text{Lu}]\text{Lu}^{3+}$, $[^{155}\text{Tb}]\text{Tb}^{3+}$, $[^{213}\text{Bi}]\text{Bi}^{3+}$ and $[^{225}\text{Ac}]\text{Ac}^{3+}$ have been performed to investigate the variation in metal ion affinity with changes in ionic radii across a broad size range and coordination number. ^{225}Ac ($t_{1/2} = 9.92$ days) is of high interest for applications in targeted alpha therapy (TAT) owing its long half-life and the potency of particulate radiation emitted within its decay scheme. Suitable companion radionuclides for $[^{225}\text{Ac}]\text{Ac}^{3+}$ with comparable coordination characteristics and imageable decay properties are required to perform accurate staging of disease progression and assessment of patient suitability for treatments.

[0129] The relatively long-lived radionuclides $[^{111}\text{In}]\text{In}^{3+}$ ($t_{1/2} = 2.83$ days)²⁷ and $[^{155}\text{Tb}]\text{Tb}^{3+}$ ($t_{1/2} = 5.51$ days)²⁰ are suitable for SPECT imaging owing to the emission of high abundance, low energy γ – rays (171 and 245 keV $[^{111}\text{In}]$ ²⁷; 44, 87 and 105 keV $[^{155}\text{Tb}]$ ²⁰) in their decay schemes, and are well-matched to the long half-life of $[^{225}\text{Ac}]\text{Ac}^{3+}$. $[^{44}\text{Sc}]\text{Sc}^{3+}$ is a promising candidate for PET imaging, owing to its long physical half-life ($t_{1/2} = 3.97$ h) and high positron branching ratio ($E_{\beta^+} = 632$ keV, 94.3%), in addition, the companion radioisotope $[^{47}\text{Sc}]\text{Sc}^3$ ($t_{1/2} = 3.35$ days) may find utility for β^- - therapy ($E_{\beta^-} = 162$ keV, 100%).²⁸

[0130] Concentration-dependent radiolabelling studies have shown $\text{H}_4\text{noneunpaX}$ to be a highly versatile chelator, exhibiting high affinity for all metal ions tested, with the exception of $[^{213}\text{Bi}]\text{Bi}^{3+}$, FIG. 28. All reactions with $\text{H}_4\text{noneunpa}$ and $\text{H}_4\text{noneunpaX}$ were carried out at RT and monitored over 10 minutes, with the exception $[^{213}\text{Bi}]\text{Bi}^{3+}$ which were monitored after 5 minutes, within 10 minutes post-generator elution ($n = 4 - 8$). Reactions with DOTA were carried out at elevated temperatures and monitored over 30 – 60 minutes.

[0131] Significantly, quantitative radiochemical conversion (RCC) was achieved within 10 minutes at room temperature, a notable advantage over the widely applied 'gold-standard' chelator DOTA. Optimal radiochemical yields were obtained under exceptionally mild

conditions (pH 7.0, RT, 10 min.) which are compatible with thermally sensitive biological targeting vectors (monoclonal antibodies). H₄noneunpaX showed comparable coordination characteristics to H₄noneunpa; each chelate was successfully radiolabelled at high molar activities with [¹¹¹In]In³⁺ (54 GBq/μmol), [¹⁵⁵Tb]Tb³⁺ (1.0 GBq/μmol), [¹⁷⁷Lu]Lu³⁺ (2.0 GBq/μmol) and [²²⁵Ac]Ac³⁺ (134 MBq/μmol).

[0132] Concentration-dependent radiolabelling studies of H₄noneunpa and H₄noneunpaX with [⁴⁴Sc]Sc³⁺ showed poor radiometal ion compatibility; whereby low RCYs were achieved at high ligand concentration. These results were anticipated based on the small ionic radius (0.870 Å, CN=8) and chemical hardness of the Sc³⁺ ion, which prefers coordination numbers of 6 – 8.¹⁸ However, these results further suggest no significant difference in metal ion affinity between symmetric and asymmetric ligands based on this framework.

[0133] The distinct contrast in radiolabelling efficiency exhibited for [²¹³Bi]Bi³⁺ by each chelate was somewhat surprising given the similarities in donor atoms and the observed trends for other trivalent metal ions. However, without being bound by theory, this deviation can be rationalised by consideration of the coordination characteristics of the Bi³⁺ ion. Bi³⁺ possesses a similar ionic radius to Ac³⁺ (1.170 Å (CN=8) versus 1.220 Å (CN=9), for Bi³⁺ and Ac³⁺ respectively),^{18,19} but is of intermediate chemical hardness, exhibiting a stronger preference for intermediate hardness donor groups (e.g. nitrogen, pyridine) over hard ionic donors (oxygen, phenolates). Bi³⁺ is also known to exhibit a stereochemically active 6s² lone pair in some of its coordination complexes, which can have significant impacts on the favoured conformational geometries of the chelating ligand and their effective denticities.^{29,30}

Example 5.0 - Human Serum Stability Studies

[0134] A series of serum stability challenge assays were undertaken in order to determine the kinetic inertness of resulting complexes of H₄noneunpaX and H₄noneunpa with [¹¹¹In]In³⁺, [¹⁵⁵Tb]Tb³⁺, [¹⁷⁷Lu]Lu³⁺ and [²²⁵Ac]Ac³⁺ in the presence of competing endogenous metal binding proteins. Results are shown in FIG. 29. In the case of H₄noneunpaX, incubation of the radiolabelled complexes at 37 °C in human serum showed no release of bound radiometal over 5-7 days, indicating high kinetic inertness and potential for *in vivo* application. The radiolabelled complexes of H₄noneunpa exhibited similar kinetic inertness

to H₄noneunpaX, maintaining > 97% radiochemical purity over 5 – 7 days, with the exception of [²²⁵Ac][Ac(noneunpa)]⁻ which showed an initial drop in radiochemical integrity of ~10 % over the course of the study. All experiments were performed at 37 °C and monitored via radio-iTLC (n = 3).

5

Example 6.0 – Computational Studies

[0135] To gain insights into the solution structures of the metal complexes of H₄noneunpaX, density functional theory (DFT) calculations were carried out using Gaussian 16 (revision B.01) in the polarizability continuum model (PCM).^{31,32} Geometrical optimisations were performed with the hybrid Perdew-Burke-Ernzerhof (PBE(0) exchange-correlation functional, using the relatively small core quasi-relativistic effective core potentials (ECP28/60MWB) and associated valence basis-sets for metal ions (La³⁺, Lu³⁺, Bi³⁺), while light atoms (C, H, N, O) were modelled up to Def2TZVP levels of theory.^{33–35} Optimised structures for all complexes, with cartesian coordinates and calculated thermodynamic energy values were determined. Initial geometries were generated using Avogadro (version 1.2.0) to provide input coordinates for calculations.⁵¹ Solvation effects were modelled using the integrated equation formalism polarizability continuum model (IEF-PCM) for all metal complexes.³² Vibrational frequency analysis was carried out on the final optimised geometries to confirm the obtained structures were true energy minima of the potential energy surface. Natural bond orbital (NBO) analysis was performed using NBO (version 3.1) within Gaussian 16.³⁶ Contour plots of the electron-density encompassing the 6s² lone pair in [Bi(noneunpaX)]⁻ were generated using Multiwfn software.⁵²

[0136] The distribution of donor arm substituents within the framework of H₄noneunpaX gives rise to four possible conformational isomers on metal ion coordination (with a coordination number of 9) which arise from the relative orientation of the 5-membered chelate rings formed between the ethylene bridges of the ligand and the metal centre (λ vs. δ). The solution structures of H₄noneunpa/Oxyaapa with La³⁺, Lu³⁺ and In³⁺ have been thoroughly investigated in previous studies.^{8,13}


[0137] Geometric optimisations of the La³⁺ and Lu³⁺ complexes of H₄noneunpaX generated structures with fully saturated metal coordination spheres, whereby all nine donor atoms from the ligand were bound to the metal centres. In the case of [La(noneunpaX)]⁻, geometric

optimisation favoured a single conformational isomer with the symmetric $\delta\delta$ configuration being the lowest in energy, while the $[\text{Lu}(\text{noneunpaX})]^-$ complex favoured a different asymmetric isomer with a twisted $\lambda\delta$ conformation, FIG. 30. Intriguingly, this is in contrast to the complexation reported for $\text{H}_4\text{noneunpa}$, which was found to form an asymmetric arrangement in the La^{3+} complex and a fully symmetric conformation in the Lu^{3+} complex.

[0138] The structure of $[\text{La}(\text{noneunpaX})]^-$ shows a high degree of symmetry, wherein the two picolinic acid donors adopt the same relative arrangement, with both pyridine rings eclipsed (parallel), while the two acetate donors also adopt the same relative arrangement in an antiparallel orientation relative the central plane of symmetry. This characterisation is consistent with the experimental NMR results of the La^{3+} complex which indicated the formation of a single, symmetric isomer in solution. Additionally, from the DFT calculated structure, the methylene protons adjacent to the acetate groups (H_a/H_b and $\text{H}_{a'}/\text{H}_{b'}$) are shown in near identical chemical environments, thus giving rise to two pairs of diastereotopic doublets in the ^1H NMR spectrum with similar chemical shifts ($\delta_{ab} =$ 3.37 ppm, $\delta_{a'b'} = 3.31$ ppm). By contrast, the solution phase structure of $[\text{Lu}(\text{noneunpaX})]^-$ shows a single asymmetric isomer, wherein a change in backbone geometry to the $\lambda\delta$ conformation causes a shift in the relative arrangement of the picolinic acid donors to an orthogonal configuration. This conformational shift is also further supported by the ^1H NMR characterisation of the Lu^{3+} complex, which exhibits sharply resolved, diastereotopic doublets for the four methylenic protons neighbouring the picolinate donor arms. The large differences in chemical shift for mutually coupled protons $\text{H}_g/\text{H}_{g'}$ ($\Delta\delta = 0.78$ ppm) and $\text{H}_k/\text{H}_{k'}$ ($\Delta\delta = 0.74$ ppm) can be clearly accounted for by the distinct local environments adopted in the rigid $[\text{Lu}(\text{noneunpaX})]^-$ complex, FIG. 30, panel (C).

Example 7.0 – Synthesis and Characterization of Bifunctional $\text{H}_4\text{noneunpaX-Bn-NCS}$

[0139] The synthetic approach developed towards bifunctional $\text{H}_4\text{noneunpaX-Bn-NCS}$ having the structure **(13)**



Reaction scheme for the synthesis of **H₄noneunpaX-Bn-NH₂** and **H₄noneunpaX-Bn-NCS**:

Starting material (1): 2-(2-aminoethoxy)ethan-1-ol

Step 1: Reaction with TBBA, DIPEA in MeCN at 50 °C, O/N, > 99% yield, yielding intermediate **(2)**.

Step 2: Reaction with 1. MsCl, DIPEA, EtOAc at 0 °C - RT, 3 h; 2. NaN₃, DMF at 90 °C, O/N; 3. i. PPh₃, THF at 0 °C - RT, 3 h; 3.ii. H₂O at RT, O/N, 91% yield, yielding intermediate **(4)** (87% overall yield).

Step 3: Reaction with NosCl, NEt₃ in CH₂Cl₂ at 0 °C - RT, 3 h, 96% yield, yielding intermediate **(7)**.

Step 4: Reaction with K₂CO₃ in MeCN at 50 °C, O/N, 88% yield, yielding intermediate **(8)**.

Step 5: Reaction with PhSH, K₂CO₃ in MeCN at 50 °C, 2 h, 98% yield, yielding intermediate **(9)**.

Step 6: Reaction with **(S1)** and DIPEA in MeCN at 50 °C, O/N, 90% yield, yielding intermediate **(10)**.

Step 7: Reaction with **(S2)** and Cu(OAc)₂·H₂O, NaAsc in tBuOH/H₂O, MW, 1 h, 85% yield, yielding intermediate **(11)**.


Step 8: Reaction with 3 M HCl at 60 °C, O/N, 90% yield, yielding intermediate **(12)**.

Step 9: Reaction with SCl₂ in CHCl₃/H₂O, 85% yield, yielding the final product **H₄noneunpaX-Bn-NCS** (**13**).


The final product **H₄noneunpaX-Bn-NH₂** is also shown.

Scheme 2. Synthesis of H₄noneunpaX-Bn-NCS.

[0140] The synthesis of bifunctional H₄noneunpaX-Bn-NCS (**13**) proceeds through a pathway analogous to H₄noneunpaX, utilising the same synthetic intermediates (compounds **(1)** – **(4)**) to provide primary amine (**4**) which was protected using 2-nitrobenzenesulphonyl chloride to give the corresponding Nosyl-protected sulfonamide (**7**) in high yield. *N*-alkylation of sulfonamide (**7**) with methyl (6-bromomethyl)picolinate under basic conditions was achieved through mild heating of the reaction mixture overnight, to generate the singly alkylated product, cleanly in high yields. Removal of the Nosyl-protecting group was performed upon treatment of compound (**8**) with thiophenol to give the corresponding secondary amine (**9**).
An excess of thiophenol (3 equiv.) was found to be necessary to achieve complete deprotection of compound (**8**) and thus avoid tedious purification of the desired product. Subsequent *N* – alkylation of secondary amine (**9**) using a structurally modified derivative of the protected picolinic acid electrophile (**S1**) was achieved through mild heating overnight to give the protected bifunctional ligand (**10**) with an alkyne appendage, suitable for click-based bioconjugation. The alkyne-derivatized picolinate electrophile (**S1**) was prepared separately starting from chelidamic acid, as shown in Scheme 3. Alkyne (**10**) was subsequently reacted with the bifunctional azide linker (**S2**) using Husigen's 1,3-dipolar cycloaddition (Click chemistry) using CuSO₄·5H₂O as an in-situ source of Cu(I) catalyst and sodium ascorbate as a reductant. Syntheses for the bifunctional azide linker (**S2**) are also outlined in Scheme 4.
Compound (**11**) was firstly treated with an excess of sodium sulfide to precipitate any traces of Cu(I/II) chelated by the protected ligand prior to deprotection of the methyl, tert-butyl and Boc protected groups using 3 M HCl. Purification of H₄noneunpaX-Bn-NH₂ (**12**) was achieved via RP-HPLC and the product isolated as the corresponding HCl salt by co-evaporation with 3 M HCl. Conversion of the aniline (**12**) to the isothiocyanate (**13**) was carried out using standardised literature approaches involving thiophosgene,³⁷ and the product isolated via RP-HPLC.



Scheme 3. Synthetic scheme of alkyne-functionalised picolinate electrophile (S1).



Scheme 4. Synthetic scheme for the preparation of clickable bifunctional handle (S2).

[0141] Di-tert-butyl 2,2'-((2-(2-((2-nitrophenyl)sulfonamido)ethoxy)ethyl)azanediyl) diacetate (7). Di-tert-butyl 2,2'-((2-(2-aminoethoxy)ethyl)azanediyl)diacetate (4) (2.35 g, 7.16 mmol) was dissolved in dry CH₂Cl₂ (40 mL) and cooled to 0 °C. Triethylamine (2.00 mL, 1.45 g, 14.3 mmol, 2 equiv.) was added to the reaction mixture, followed by slow addition of 2-nitrobenzenesulphonyl chloride (1.59 g, 7.16 mmol). The resulting pale-yellow solution was stirred at 0 °C for 1 h, then allowed to warm to RT and stirred for a further 5 h. Upon completion, the reaction mixture was diluted with CH₂Cl₂ (40 mL) and extracted with de-ionised H₂O (2 x 50 mL) and brine (50 mL). The organic phase was evaporated *in vacuo* and the resulting residue purified via silica gel chromatography (Combiflash automated purification system; A: hexanes, B: EtOAc, 100% A to 40% B). The title compound was attained as a pale-yellow oil (3.51 g, 96%). ¹H NMR (400 MHz, CDCl₃, 298 K) 8.14 – 8.12 (1H, m, 11 – CH), 7.85 – 7.82 (1H, m, 9 – CH), 7.75 – 7.70 (2H, m, 8 – and 11 – CH), 6.13 (1H, t, ³J = 5.6 Hz, 7 – NH), 3.54 – 3.49 (4H, m, 4 – and 5 – CH₂), 3.45 (4H, s, 2 – CH₂), 3.28 (2H, q, ³J = 5.5 Hz, 6 – CH₂), 2.87 (2H, t, ³J = 5.6 Hz, 3 – CH₂), 1.45 (18H, s, 1 – C(CH₃)₃). ¹³C{¹H} NMR (100 MHz, CDCl₃, 298 K) 170.7 (3 – C), 148.1 (14 – C), 134.0 (11 – C), 133.4 (9 – C), 132.6 (12 – C), 130.9 (13 – C), 125.2 (10 – C), 81.1 (2 – C), 70.1 (6 – C), 68.9 (7 –

C), 56.7 (4 – C), 53.2 (5 – C), 43.7 (8 – C), 28.2 (1 – C). ESI-MS (MeOH) 518.7 [M+H]⁺. *R_f* = 0.50 (Hexanes/EtOAc; 2:1).

[0142] Di-tert-butyl 2,2'-((2-(2-((N-((6-(methoxycarbonyl)pyridin-2-yl)methyl)-2-nitrophenyl)sulfonamido)ethoxy)ethyl)azanediyl)diacetate (8): Methyl (6-bromomethyl) picolinate (884 mg, 3.84 mmol) was added to a solution of di-tert-butyl 2,2'-((2-(2-((2-nitrophenyl)sulfonamido)ethoxy)ethyl)azanediyl)diacetate (**7**) (1.98 g, 3.84 mmol) in dry MeCN (60 mL). The solution was stirred for 15 minutes at RT after which K₂CO₃ (1.06 g, 7.68 mmol, 2.0 equiv.) was added and the resulting suspension heated at 50 °C overnight. Upon completion, the inorganic salts were separated *via* centrifuge, washed with CH₂Cl₂ (3 x 10 mL) and the combined organic phase was evaporated in vacuo. The resulting residue was re-dissolved in CH₂Cl₂ (75 mL), washed with de-ionised H₂O (3 x 75 mL) and brine (75 mL) and dried over Na₂SO₄. The organic phase was evaporated *in vacuo* and the resulting residue purified *via* silica gel chromatography (Combiflash automated purification system; A: CH₂Cl₂, B: MeOH; 100% A to 5% B). The title product was attained as a pale-yellow oil (2.26 g, 88%).
¹H NMR (400 MHz, CDCl₃, 298 K) 8.13 – 8.1 (1H, m, 11 – CH), 7.98 (1H, d, ³*J* = 7.1 Hz, 14 – CH), 7.81 (1H, t, ³*J* = 7.1 Hz, 13 – CH), 7.68 (1H, d, ³*J* = 7.1 Hz, 12 – CH), 7.66 – 7.62 (3H, m, 8 – , 9 – and 10 – CH), 4.83 (2H, s, 7 – CH₂), 3.95 (3H, s, 15 – CH₃), 3.54 (2H, t, ³*J* = 5.3 Hz, 6 – CH₂), 3.47 (2H, t, ³*J* = 5.3 Hz, 5 – CH₂), 3.37 (4H, s, 2 – CH₂), 3.35 (2H, t, ³*J* = 6.0 Hz, 4 – CH₂), 2.71 (2H, ³*J* = 6.0 Hz, 3 – CH₂), 1.41 (18H, s, 1 – C(CH₃)₃). ¹³C{¹H} NMR (100 MHz, CDCl₃, 298K) 170.6 (3 – C), 165.4 (21 – C), 157.4 (16 – C), 148.1 (20 – C), 147.4 (14 – C), 137.9 (18 – C), 133.5 (11 – C), 133.4 (9 – C), 131.8 (12 – C), 131.0 (13 – C), 125.4 (17 – C), 124.3 (10 – C), 124.0 (19 – C), 80.9 (2 – C), 70.0 (6 – C), 68.6 (7 – C), 56.5 (4 – C), 53.9 (15 – C), 53.3 (5 – C), 52.8 (22 – C), 48.1 (8 – C), 28.1 (1 – C). ESI – MS (MeOH) 667.3 [M+H]⁺. *R_f* = 0.30 (CH₂Cl₂/MeOH; 99:1).

[0143] Di-tert-butyl 2,2'-((2-(2-(((6-(methoxycarbonyl)pyridin-2-yl)methyl)amino)ethoxy)ethyl)azanediyl)diacetate (9). Thiophenol (1.05 mL, 10.2 mmol, 3.0 equiv.) was added to a suspension of di-tert-butyl 2,2'-((2-(2-((N-((6-(methoxycarbonyl)pyridin-2-yl)methyl)-2-nitrophenyl)sulfonamido)ethoxy)ethyl)azanediyl) diacetate (**8**) (2.26 g, 3.39 mmol) and K₂CO₃ (937 mg, 6.78 mmol, 2.0 equiv.) in dry MeCN (60 mL). The reaction mixture was heated to 50 °C and stirred for 2 h. Upon completion, the inorganic salts were separated *via* centrifuge, washed with CH₂Cl₂ (3 x 10 mL) and the combined organic phase evaporated *in vacuo*. The resulting residue was re-dissolved in CH₂Cl₂ (50 mL) and washed with de-ionised H₂O (2 x 50 mL) and brine (50 mL). The organic phase was evaporated *in vacuo* and

the resulting oil purified *via* silica gel chromatography (Combiflash automated purification system; A: CH₂Cl₂, B: MeOH; 100% A to 10% B) to give the title product as a pale-yellow oil (1.61 g, 98%). ¹H NMR (400 MHz, CDCl₃, 298 K) 8.00 (1H, d, ³J = 7.4 Hz, 11 – CH), 7.81 (1H, t, ³J = 7.4 Hz, 10 – CH), 7.66 (1H, d, ³J = 7.4 Hz, 9 – CH), 4.05 (2H, s, 8 – CH₂), 3.99 (3H, s, 12 – CH₃), 3.61 – 3.56 (4H, m, 4 – and 5 – CH₂), 3.49 (4H, s, 2 – CH₂), 2.94 (2H, t, ³J = 5.8 Hz, 6 – CH₂), 2.83 (2H, t, ³J = 5.2 Hz, 3 – CH₂), 2.26 (1H, br s, 7 – NH), 1.44 (18H, s, 1 – C(CH₃)₃). ¹³C {¹H} NMR (100 MHz, CDCl₃, 298 K) 170.8 (3 – C), 165.8 (15 – C), 160.5 (10 – C), 147.4 (14 – C), 137.5 (12 – C), 125.6 (11 – C), 123.5 (13 – C), 80.9 (2 – C), 70.1 (6 – C), 70.0 (7 – C), 56.6 (4 – C), 54.9 (9 – C), 53.4 (5 – C), 52.8 (16 – C), 48.8 (8 – C), 28.1 (1 – C). ESI – MS (MeOH) 482.2 [M+H]⁺. *R*_f = 0.20 (CH₂Cl₂/MeOH; 9:1).

[0144] (tBu)₂(Me)₂noneunpaX(OCH₂CCH) (10). Methyl 6-(bromomethyl)-4-(prop-2-yn-1-yloxy)picolinate (122 mg, 0.415 mmol) was added to a suspension of di-tert-butyl 2,2'-((2-(2-(((6-(methoxycarbonyl)pyridin-2-yl)methyl)amino)ethoxy)ethyl)azanediyl)diacetate (200 mg, 0.415 mmol) and K₂CO₃ (115 mg, 0.830 mmol) in dry MeCN (10 mL). The reaction mixture was heated to 50 °C and stirred overnight. Upon completion, the reaction mixture was cooled to room temperature, and the inorganic salts removed via centrifuge. The separated salts were washed with CH₂Cl₂ (3 x 10 mL) and the combined organic phase evaporated *in vacuo*. The resulting residue was re-dissolved in CH₂Cl₂ (25 mL) and washed with de-ionised H₂O (2 x 25 mL) and brine (25 mL). The volatiles were removed *in vacuo* and the crude product purified *via* silica gel chromatography (Combiflash automated purification system; A: CH₂Cl₂, B: MeOH; 100% A to 5% B). The title product was attained as a pale-yellow oil (249 mg, 88%). ¹H NMR (400 MHz, CDCl₃, 298 K) 7.98 (1H, dd, ³J = 7.5 Hz, ⁴J = 0.9 Hz, 10 – CH), 7.88 (1H, d, ³J = 7.5 Hz, 8 – CH), 7.78 (1H, t, ³J = 7.5 Hz, 9 – CH), 7.59 (1H, d, ⁴J = 2.5 Hz, 14 – CH), 7.52 (1H, br d, ⁴J = 2.5 Hz, 13 – CH), 4.81 (2H, d, ⁴J = 2.4 Hz, 16 – CH₂) 4.01 (2H, br s, 7 – CH₂), 3.98 (3H, s, 11 – CH₃), 3.97 (5H, br m, 12 – CH₂ and 15 – CH₃), 3.57 (2H, t, ³J = 5.5 Hz, 4 – CH₂), 3.52 (2H, t, ³J = 5.9 Hz, 5 – CH₂), 3.45 (4H, s, 2 – CH₂), 2.91 (2H, t, ³J = 5.9 Hz, 6 – CH₂), 2.81 (2H, t, ³J = 5.5 Hz, 3 – CH₂), 2.60 (1H, t, ⁴J = 2.4 Hz, 17 – CH), 1.42 (18H, s, 1 – C(CH₃)₃). ¹³C {¹H} NMR (100 MHz, CDCl₃, 298 K) 170.7 (3 – C), 165.8 (15 – C), 165.7 (23 – C), 165.1 (20 – C), 162.6 (10 – C), 160.6 (18 – C), 148.9 (22 – C) 147.3 (14 – C), 137.4 (12 – C), 126.1 (11 – C), 123.6 (13 – C), 111.8 (19 – C), 111.0 (21 – C), 80.8 (2 – C), 77.1 (26 – C), 76.9 (27 – C), 70.4 (6 – C), 69.3 (7 – C), 60.8 (9 – C), 60.6 (17 – C), 56.6 (4 – C), 55.9 (25 – C), 54.0 (8 – C), 53.5 (5 – C), 53.0 (16 – C), 52.9 (24 – C), 28.1 (1 – C). ESI – MS (MeOH) 685.3 [M+H]⁺.

[0145] (^tBu)₂(Me)₂noneunpaX(Bn-NHBoc) (11). A solution of Cu(OAc)₂·H₂O (66 mg, 0.329 mmol, 1.0 equiv.) in de-ionised H₂O (1 mL) was added to a mixture containing (^tBu)₂(Me)₂noneunpaX(OCH₂CCH) (225 mg, 0.329 mmol) and *tert*-butyl (4-(2-azidoethyl)phenyl)-carbamate (86 mg, 0.329 mmol) in ^tBuOH (1 mL). Sodium ascorbate (65 mg, 0.329 mmol, 1.0 equiv.) was added to the reaction mixture, which darkened progressively over 10 min to give a red-brown solution. The solution was warmed to 40 °C and stirred for 48 h. Upon completion, the dark brown mixture was treated with a solution Na₂S·9H₂O (790 mg, 3.29 mmol, 10 equiv.) dissolved in de-ionised H₂O (4 mL). Black CuS precipitated immediately on addition, and the suspension was stirred for a further 1 h at RT. The precipitate was separated by centrifuge, and the supernatant collected and evaporated under reduced pressure to remove excess ^tBuOH. The resulting aqueous phase was diluted with de-ionised H₂O (10 mL) and extracted with CH₂Cl₂ (3 x 15 mL). The combined organic phase was washed with de-ionised H₂O (10 mL) and brine (10 mL). The volatiles were removed *in vacuo* and the crude material purified *via* silica gel chromatography (CombiFlash automated purification system; A: CH₂Cl₂, B: MeOH; 100% A to 10% B) to attain the title compound as an white aerated solid (211 mg, 76%). ¹H NMR (400 MHz, CDCl₃, 298 K) 8.00 (1H, d, ³J = 7.5 Hz, 10 – CH), 7.90 (1H, d, ³J = 7.5 Hz, 18 – CH), 7.86 (1H, t, ³J = 7.5 Hz, 9 – CH), 7.60 (1H, d, ⁴J = 2.2 Hz, 14 – CH), 7.53 (1H, d, ⁴J = 2.2 Hz, 13 – CH), 7.45 (1H, s, 17 – CH), 7.27 (2H, d, ³J = 8.2 Hz, 21 – CH), 6.98 (2H, d, ³J = 8.2 Hz, 20 – CH), 6.66 (1H, br s, 22 – NH), 5.28 (2H, s, 25 – CH₂), 4.59 (2H, t, ³J = 7.1 Hz, 18 – CH₂), 4.02 (2H, s, 12 – CH₂), 3.99 (3H, s, 11 – CH₃), 3.98 (3H, s, 15 – CH₃), 3.96 (2H, s, 7 – CH₂), 3.58 (2H, t, ³J = 5.6 Hz, 4 – CH₂), 3.53 (2H, t, ³J = 5.8 Hz, 5 – CH₂), 3.46 (4H, s, 4 – CH₂), 3.17 (2H, t, ³J = 7.1 Hz, 19 – CH₂), 2.91 (2H, t, ³J = 5.8 Hz, 6 – CH₂), 2.81 (2H, t, ³J = 5.6 Hz, 3 – CH₂), 1.52 (9H, s, 23 – C(CH₃)₃), 1.43 (18H, s, 1 – C(CH₃)₃). ESI – MS (MeOH) 947.5 [M+H]⁺.

[0146] H₄noneunpaX–Bn–NH₂ (12). (^tBu)₂(Me)₂noneunpaX(Bn-NHBoc) (200 mg, 0.211 mmol) was dissolved in 4 M HCl (3 mL) and heated at 60 °C overnight. Upon completion, the volatiles were evaporated *in vacuo* and the resulting residue purified *via* RP – HPLC (A: H₂O (0.1% TFA), B: MeCN; 95% A to 30% B over 40 min, R_t = 20.3 - 22.3 min). Purified H₄noneunpaX – triazole – Bn – NH₂ was co-evaporated with 3 M HCl (3 x 2 mL) to give the corresponding HCl as a white solid (120 mg, 81%). ¹H NMR (400 MHz, D₂O, 298 K) 8.03 (1H, s, 14 – CH), 7.94 – 7.83 (2H, m, 8 – and 9 – CH), 7.62 (1H, dd, ³J = 6.3 Hz, ⁴J = 2.0 Hz, 7 – CH), 7.58 (1H, d, ⁴J = 2.7 Hz, 12 – CH), 7.24 (1H, d, ⁴J = 2.7 Hz, 11 – CH), 7.22 (2H, d, ³J = 7.4 Hz, 17 – CH), 7.15 (2H, d, ³J = 7.4 Hz, 18 – CH), 5.29 (2H, s, 13 – CH₂), 4.74 – 4.65 (6H,

m, 6 – , 10 – and 15 – CH_2), 4.22 (4H, s, 1 – CH_2), 3.97 (2H, t, 4 – CH_2), 3.89 (2H, t, 3 – CH_2), 3.76 (2H, t, 5 – CH_2), 3.66 (2H, t, 2 – CH_2), 3.23 (2H, t, 16 – CH_2). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, D_2O , 298 K) 168.1 (17 – C), 168.0 (1 – C), 166.4 (13 – C), 164.7 (20 – C), 150.4 (8 – C), 150.1 (15 – C), 147.3 (12 – C), 146.0 (19 – C), 140.9 (22 – C), 140.7 (10 – C), 138.5 (29 – C), 130.4 (27 – C), 128.5 (26 – C), 128.4 (9 – C), 126.2 (23 – C), 125.5 (11 – C), 123.3 (28 – C), 115.8 (16 – C), 112.8 (18 – C), 65.4 (5 – C), 64.4 (4 – C), 62.1 (21 – C), 58.3 (7 – C), 57.7 (14 – C), 56.4 (3 – C), 56.2 (6 – C), 55.6 (2 – C), 51.9 (24 – C), 35.2 (25 – C). LR-ESI-MS (H_2O) 707.6 $[\text{M}+\text{H}]^+$

[0147] H₄noneunpaX–Bn–NCS (13). Thiophosgene (120 μL , 1.56 mmol) in CHCl_3 (660 μL) was added to a solution of H₄noneunpaX–Bn– $\text{NH}_2\cdot 4\text{HCl}$ (88 mg, 0.103 mmol) in de-ionised H_2O (1 mL). The biphasic mixture was stirred vigorously in the dark overnight at RT, after which the two immiscible layers were allowed to separate. The aqueous phase was collected, washed with CHCl_3 (2 x 1 mL), and then lyophilised to attain the crude isothiocyanate product. The off-white residue was purified via semi-preparation RP-HPLC (A: H_2O (0.1%TFA), B: MeCN (0.1%TFA)): 0 – 6 min. (95% A to 24% B); 6 – 30 min. (24% B to 39% B); 30 - 35 min. (39% B to 100% B); R_t = 25.4 min. Appropriate fractions were pooled and lyophilised to give the purified product as an off-white aerated solid (67 mg, 87%). ESI-MS ($\text{H}_2\text{O}/\text{MeCN}$) 749.3 $[\text{M}+\text{H}]^+$, 747.3 $[\text{M}-\text{H}]^-$.

[0148] Dimethyl-4-hydroxypyridine-2,6-dicarboxylate (S3) Chelidamic acid monohydrate (3.00 g, 14.9 mmol) was suspended in dry MeOH (50 mL) and H_2SO_4 (0.5 mL) added. The reaction mixture was heated at 50 $^\circ\text{C}$ overnight, then cooled to RT and the volatiles removed *in vacuo*. The crude residue was dissolved in sat. NaHCO_3 sol. (150 mL) and extracted with EtOAc (3 x 150 mL). The combined organic phase was evaporated *in vacuo* to attain the title compound as a white solid (3.02 g, 95%). ^1H NMR (300 MHz, CDCl_3 , 298 K) 7.49 (2H, s, 2 – CH), 3.97 (6H, s, 1 – CH_3). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3 , 298 K) 187.5 (5 – C), 163.5 (2 – C), 143.1 (3 – C), 118.6 (4 – C), 53.7 (1 – C).

[0149] Dimethyl 4-(prop-2-yn-1-yloxy)pyridine-2,6-dicarboxylate (S4) Propargyl bromide (80% wt.)(1.75 mL, 15.7 mmol, 1.1 equiv.) was added to a suspension of dimethyl-4-hydroxypyridine-2,6-dicarboxylate (3.02 g, 14.3 mmol) and K_2CO_3 (3.95 g, 28.6 mmol, 2.0 equiv.) in dry DMF (50 mL). The reaction mixture was heated at 50 $^\circ\text{C}$ overnight, then cooled to RT and the salts removed *via* vacuum filtration. The filtrate was evaporated *in vacuo* and the resulting residue purified *via* silica gel chromatography (Combiflash automated purification system; A: CH_2Cl_2 , B: MeOH; 100% A to 5% B). The title compound was attained

as a pale-yellow solid (3.20 g, 90%). ¹H NMR (400 MHz, CDCl₃, 298 K) 7.89 (2H, s, 2 – CH), 4.87 (2H, d, ⁴J = 2.4 Hz, 3 – CH₂), 4.01 (6H, s, 1 – CH₃), 2.62 (1H, t, ⁴J = 2.4 Hz, 4 – CH). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 165.7 (2 – C), 165.1 (5 – C), 150.0 (3 – C), 115.0 (4 – C), 77.7 (7 – C), 76.3 (8 – C), 56.5 (6 – C), 53.4 (1 – C).

- 5 **[0150] Methyl (6-hydroxymethyl)-4-(prop-2-yn-1-yloxy)picolinate (S5)** NaBH₄ (190 mg, 4.92 mmol) was added in small portions to a solution of compound 2 (1.22 g, 4.90 mmol) in dry MeOH/CH₂Cl₂ (40 mL, 1:1) at 0 °C with vigorous stirring. The reaction mixture was stirred for 30 minutes at 0 °C, then allowed to warm to RT and stirred for a further 3.5 h. Upon completion, the reaction solution was cooled to 0 °C and quenched with de-ionised H₂O (25 mL). The volatiles were removed *in vacuo* and the resulting aqueous phase extracted with CH₂Cl₂ (3 x 30 mL). The combined organic phase was dried over Na₂SO₄, filtered and evaporated *in vacuo* to give an off-white solid. The crude material was purified via silica gel chromatography (CombiFlash automated purification system; A: CH₂Cl₂, B: MeOH; 100% A to 5% B) to give the title compound as a white crystalline solid (721 mg, 67%). ¹H NMR (400 MHz, CDCl₃, 298K) 7.57 (1H, d, ⁴J = 2.4 Hz, 2 – CH), 7.16 (1H, d, ⁴J = 2.4 Hz, 5 – CH), 4.79 (2H, s, 6 – CH₂), 4.78 (2H, d, ⁴J = 3 – CH₂), 4.22 (1H, br s, 7 – OH), 3.93 (3H, s, 1 – CH₃), 2.61 (1H, t, ⁴J = 2.4 Hz, 4 – CH). ¹³C {¹H} NMR (100 MHz, CDCl₃, 298 K) 165.4 (2 – C), 165.2 (5 – C) (7 – C), 162.9 (– C), 148.6 (3 – C), 111.2 (4 – C), 109.9 (6 – C), 77.1 (10 – C), 76.8 (11 – C), 64.7 (8 – C), 56.0 (9 – C), 53.0 (1 – C).
- 15 **[0151] Methyl 6-(bromomethyl)-4-(prop-2-yn-1-yloxy)picolinate (S1)** Phosphorus tribromide (85 mL, 245 mg, 0.905 mmol, 1.1 equiv.) was added dropwise to a cooled solution of methyl 6-(hydroxymethyl)-4-(prop-2-yn-1-yloxy)picolinate (182 mg, 0.823 mmol) in dry CHCl₃ (5 mL). The reaction mixture was warmed passively to RT and stirred for 3 h. Upon completion, the yellow reaction mixture was quenched with sat. NH₄Cl (20 mL) and extracted with CH₂Cl₂ (4 x 20 mL). The combined organic phase was evaporated *in vacuo* to yield the title compound as a white solid (230 mg, 99%). ¹H NMR (300 MHz, CDCl₃, 298 K) 7.63 (1H, d, ⁴J = 2.4 Hz, 2 – CH), 7.22 (1H, d, ⁴J = 2.4 Hz, 5 – CH), 4.79 (2H, d, ⁴J = 2.4 Hz, 3 – CH), 4.56 (2H, s, 6 – CH₂), 3.96 (3H, s, 1 – CH₃), 2.60 (1H, t, ⁴J = 2.4 Hz, 4 – CH). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 165.4 (2 – C), 165.3 (5 – C), 159.1 (7 – C), 149.4 (3 – C), 113.7 (6 – C), 111.7 (4 – C), 77.5 (10 – C), 76.5 (11 – C), 56.2 (9 – C), 53.3 (1 – C), 33.3 (8 – C). ESI-MS (MeOH) 284.0.
- 20 **[0152] tert-butyl (4-(2-hydroxyethyl)phenyl)carbamate (S6)** Di-*tert*-butyl dicarbonate (3.50 g, 16.0 mmol, 1.1 equiv.) was added to a solution of 2-(4-aminophenyl)ethanol (2.01 g,
- 25
- 30

14.6 mmol) and triethylamine (2.0 mL, 1.46 g, 14.6 mmol) in dry THF (50 mL). The reaction mixture was stirred overnight at RT, then heated to 50 °C for a further 5 h. Upon completion, the volatiles were removed *in vacuo* to give the title product as an off-white solid (3.35 g, 97%) which was used without further purification. ¹H NMR (300 MHz, CDCl₃, 298 K) 7.29 (2H, d, *J* = 8.4 Hz, 3 – CH), 7.13 (2H, d, *J* = 8.4 Hz, 4 – CH), 6.49 (1H, br s, 2 – NH), 3.80 (2H, t, *J* = 6.6 Hz, 6 – CH₂), 2.80 (2H, t, *J* = 6.6 Hz, 5 – CH₂), 1.50 (9H, s, 1 – C(CH₃)₃). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 153.0 (3 – C), 136.9 (4 – C), 133.2 (7 – C), 129.7 (6 – C), 119.1 (5 – C), 80.6 (2 – C), 63.8 (9 – C), 38.6 (8 – C), 28.5 (1 – C). ESI-MS (MeOH) 260.2 [M+Na]⁺.

[0153] *tert*-butyl (4-(2-azidoethyl)phenyl)carbamate (S2). Methanesulphonyl chloride

(825 mL, 1.22 g, 10.6 mmol, 1.1 equiv.) was added dropwise to a solution of *tert*-butyl (4-(2-hydroxyethyl)phenyl)carbamate (2.29 g, 9.68 mmol) and DIPEA (1.85 mL, 1.37 g, 10.6 mmol, 1.1 equiv.) in dry EtOAc (20 mL) at 0 °C. The solution was stirred for 10 min over ice, then warmed to RT and stirred for a further 1 h. Upon completion, the reaction mixture was cooled to 0 °C, filtered, and the precipitate washed with cold EtOAc (70 mL). The filtrate was concentrated *in vacuo* to yield corresponding mesylate as a pale-yellow oil (**S7**), which was used without further purification. ¹H NMR (300 MHz, CDCl₃, 298 K) 7.31 (2H, d, *J* = 8.5 Hz, 3 – CH), 7.12 (2H, d, *J* = 8.5 Hz, 4 – CH), 6.67 (1H, br s, 2 – NH), 4.34 (2H, t, *J* = 7.1 Hz, 6 – CH₂), 2.96 (2H, t, *J* = 7.1 Hz, 5 – CH₂), 2.83 (3H, s, 7 – CH₃), 1.49 (9H, s, 1 – C(CH₃)₃). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 152.9 (3 – C), 137.5 (4 – C), 130.8 (7 – C), 129.6 (6 – C), 118.9 (5 – C), 80.5 (2 – C), 70.5 (9 – C), 37.4 (10 – C), 35.0 (8 – C), 28.4 (1 – C). R_f = 0.75 (Hexanes/EtOAc, 1:1). 4-((*tert*-butoxycarbonyl)amino)phenethyl methanesulfonate (**S7**) (3.03 g, 9.66 mmol) was dissolved in dry DMF (15 mL) and NaN₃ (703 mg, 10.6 mmol, 1.1 equiv.) added. The reaction mixture was heated to 90 °C and stirred overnight. Upon completion, the volatiles were removed *in vacuo* and the resulting residue partitioned between de-ionised H₂O (50 mL) and EtOAc (50 mL). The aqueous phase was further extracted with EtOAc (2 x 50 mL), and the combined organic phase evaporated *in vacuo* to yield the title compound as a pale-yellow oil (2.18 g, 86%). ¹H NMR (300 MHz, CDCl₃, 298 K) 7.30 (2H, d, ³*J* = 8.4 Hz, 3 – CH), 7.08 (2H, d, ³*J* = 8.4 Hz, 2 – CH), 6.91 (1H, br s, 2 – NH), 3.42 (2H, t, ³*J* = 7.2 Hz, 5 – CH₂), 2.79 (2H, t, ³*J* = 7.2 Hz, 6 – CH₂), 1.48 (9H, s, 1 – C(CH₃)₃). ¹³C {¹H} NMR (75 MHz, CDCl₃, 298 K) 153.0 (3 – C), 137.4 (4 – C), 132.5 (7 – C), 129.3 (6 – C), 118.9 (5 – C), 80.4 (2 – C), 52.6 (9 – C), 34.8 (8 – C), 28.4 (1 – C). ESI-MS (MeOH) 285.16 [M+Na]⁺. R_f = 0.95 (Hexanes/EtOAc, 1:1).

Example 8.0 - Radiolabelling Studies of Bifunctional H₄noneunpaX-Bn-NH₂

[0154] Concentration-dependent radiolabelling studies of H₄noneunpaX-Bn-NH₂ were undertaken to assess the impacts of bifunctionalisation through one picolinic acid donor group on the radiometal ion chelation, FIG. 31. All reactions were conducted at ambient temperature and radiochemical yields (RCYs) determined *via* iTLC using SA-paper plates and EDTA (50 mM, pH 5.0) as eluent (n = 4), except in the case of [²²⁵Ac]Ac³⁺, which used EDTA (50 mM, pH 7.0) as TLC eluent.

[0155] Radiolabelling of H₄noneunpaX-Bn-NH₂ with [⁴⁴Sc]Sc³⁺ (400 kBq) demonstrated comparable results to the original chelating ligand, achieving quantitative RCC within 10 min. at RT, thus implying no significant changes to the coordination environment were incurred through functionalisation using this approach. Without being bound by theory, the moderate improvement in RCC at low ligand concentration (10⁻⁵ M) may be attributed to the bifunctional appendage imposing a degree of preorganisation in the binding cavity, thereby favouring metal complexation under these conditions.

[0156] Radiolabelling studies of H₄noneunpaX-Bn-NH₂ with [¹⁷⁷Lu]Lu³⁺ also showed comparable results to the unmodified chelator; achieving quantitative RCYs over a wide concentration range (10⁻³ to 10⁻⁶ M) at RT within 10 minutes. Further screening of the radiolabelling properties of H₄noneunpaX-Bn-NH₂ with [¹¹¹In]In³⁺, [¹⁷⁷Lu]Lu³⁺, [^{133/135}La]La³⁺, [¹⁵⁵Tb]Tb³⁺ and [²²⁵Ac]Ac³⁺ showed comparable results to the unmodified chelator, with quantitative RCCs being achieved at ligand concentrations of 10⁻⁶ M with each respective radiometal ion. Notably, the amount of radioactivity used for assessment of RCCs with each radionuclide corresponds to similar molar equivalents of radiometal ions thereby allowing more direct comparison of the concentration-dependence with different radionuclides.

[0157] In addition, an evaluation of the maximum molar activity of [¹⁷⁷Lu][Lu(noneunpaX-Bn-NH₂)]⁻ was conducted to determine minimal amount of chelate required to achieve quantitative RCYs using 20 MBq of [¹⁷⁷Lu]Lu³⁺, FIG. 32 left panel. Quantitative radiolabelling of [¹⁷⁷Lu]Lu³⁺ (20 MBq) was achieved using 80 pmol of H₄noneunpa-Bn-NH₂, corresponding to a molar activity of 250 GBq/μmol and a ligand-to-metal ratio of 174:1. Human serum stability studies of [¹⁷⁷Lu][Lu(noneunpaX-Bn-NH₂)]⁻ showed no transchelation of bound radioactivity over 7 days, thereby confirming that modification of the pendent donor arm does not impact the overall stability of the metal complex, FIG. 32 right panel.

Example 9.0 – Bioconjugate Studies

[0158] As a proof of concept to assess the suitability of H₄noneunpaX for *in vivo* applications, two structural analogues of H₄noneunpaX-Bn-NCS were prepared

- 5 incorporating the SSTR2-targeting peptide Tyr³-Octreotate (Tyr³-TATE) for evaluation in mice bearing AR42J exocrine/pancreatic tumour xenografts. Tyr³-TATE was selected as an appropriate model targeting vector due to its ubiquitous use for targeting neuroendocrine tumours (NETs), and to allow direct comparison to existing clinically applied radiotracers, namely [¹⁷⁷Lu][Lu(DOTATATE)].¹⁵

10 *Synthesis of H₄noneunpaX-Ahx/PEG₂-Tyr³-TATE*


[0159] Synthesis of the linear resin-bound Tyr³-Octreotate peptides was carried out using standardised Fmoc-based semi-automated solid-phase peptide synthesis (SPPS) on pre-loaded Wang resin, using an analogous approach to that reported by *Noor et al.*³⁸ The resin-bound linear peptide was prepared according to the sequence: [DPhe-Cys(Acm)-

- 15 Tyr(*t*Bu)-DTrp(*t*Bu)-Lys(Boc)-Thr(*t*Bu)-Cys(Acm)-Thr(*t*Bu)-OH], Scheme 5. After generation of the resin-bound octapeptide sequence, N^α-Fmoc-Ahx-CO₂H or N^α-Fmoc-PEG-CO₂H were incorporated as covalent linkers to give the Fmoc-protected linear-peptides **(14)** and **(15)** respectively. Cyclisation of the linear peptides was achieved by treatment with iodine in DMF, which sequentially removes the acetamido methyl (Acm) protecting groups and
- 20 mediates the formation of the disulfide bridge between Cys² and Cys⁷. Global deprotection and cleavage from the resin was carried out using a standardised TFA cleavage cocktail to give the cyclised Fmoc-protected peptides **(16)** and **(17)** (Fmoc-Ahx-Tyr³-TATE **(16)** and Fmoc-PEG₂-Tyr³-TATE **(17)**), which were isolated by RP-HPLC. ESI-MS (H₂O/MeCN): Fmoc-Ahx-Tyr³-TATE 1384.4 [M+H]⁺, 1382.7 [M-H]⁻; Fmoc-PEG₂-Tyr³-TATE 1416.8 [M+H]⁺,
- 25 1414.7 [M-H]⁻.

- [0160]** To allow selective functionalisation of the N-terminus of each peptide, the side chain primary amine group on Lys⁵ was firstly protected using di-*tert*-butyl dicarbonate prior to N^α-Fmoc cleavage with 20% piperidine in DMF to give compounds **(18)** and **(19)**. This approach was selected to maximise the purity of the Tyr³-Octreotate peptides prior to
- 30 incorporating the bifunctional chelator. ESI-MS (H₂O/MeCN): H₂N-Ahx-Boc(Lys⁵)-Tyr³-

TATE (**18**) 1262.9 [M+H]⁺, 1260.8 [M-H]⁻; H₂N-PEG₂-Boc(Lys⁵)-Tyr³-TATE (**19**) 1294.8 [M+H]⁺, 1292.7 [M-H]⁻.

- [0161]** Conjugation of H₄noneunpaX-Bn-NCS to the free N-terminus of (**18**) and (**19**) was achieved under mild, basic conditions in solution, and final cleavage of N(Boc)-Lys⁵ protecting group gave the corresponding chelate-peptide bioconjugates (**20**) and (**21**).
5 H₄noneunpaX-Ahx-Tyr³-TATE (**20**) and H₄noneunpaX-PEG₂-Tyr³-TATE (**21**) were purified via RP-HPLC and mass spectrometry analysis (ESI/MALDI) performed to confirm isolation of the intended products.



Scheme 5. Synthesis of H₄noneunpaX-Ahx-Tyr³-TATE and H₄noneunpaX-PEG₂-Tyr³-TATE.

Synthesis of H₄noneunpaX-Ahx-Tyr³-TATE

- 5 **[0162]** In more detail, H₄noneunpaX-Bn-NCS (3.7 mg, 4.95 μmol) was added to a solution of H₂N-Ahx-Boc(Lys⁶)-Tyr³-TATE (6.3 mg, 4.95 μmol) in dry DMF (1 mL). The solution was

stirred for 10 min. at RT then DIPEA (10 μ L, 57 μ mol) added. The resulting mixture was stirred O/N at RT, then treated with 20%TFA in CH_2Cl_2 (1 mL) and stirred for a further 2 h. After completion, the volatiles were evaporated under a stream of N_2 gas, and the resulting residue diluted with H_2O (0.1% TFA). The crude bioconjugate was purified via semi-preparative RP-HPLC (A: H_2O (0.1% TFA), B: MeCN (0.1% TFA) (3 mL/min); method: 0 – 6 min. (95% A to 29% B); 6 – 25 min. (ISO 29% B); 25 – 30 min. (29% B to 100% B); R_t = 22.6 min. Appropriate fractions were pooled and lyophilised to give $\text{H}_4\text{noneunpaX-Ahx-Tyr}^3\text{-TATE}$ (**20**) as a white solid. ESI-MS ($\text{H}_2\text{O/MeCN}$ (1:1)) 1911.9 $[\text{M}+\text{H}]^+$.

Synthesis of $\text{H}_4\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE}$

[0163] $\text{H}_4\text{noneunpaX-Bn-NCS}$ (2.3 mg, 3.07 μ mol) was added to a solution of $\text{H}_2\text{N-PEG}_2\text{-Boc(Lys}^6\text{)-Tyr}^3\text{-TATE}$ (4.0 mg, 3.07 μ mol) in dry DMF (500 μ L). The solution was stirred for 10 min. at RT then DIPEA (10 μ L, 57 μ mol) added. The resulting mixture was stirred O/N at RT, then treated with 20%TFA in CH_2Cl_2 (1 mL) and stirred for a further 2 h. After completion, the volatiles were evaporated under a stream of N_2 gas, and the resulting residue diluted with H_2O (0.1% TFA). The crude bioconjugate was purified via semi-preparative RP-HPLC (A: H_2O (0.1% TFA), B: MeCN (0.1% TFA) (3 mL/min); method: 0 – 6 min. (95% A to 29% B); 6 – 25 min. (ISO 29% B); 25 – 30 min. (29% B to 100% B); R_t = 19.1 min. Appropriate fractions were pooled and lyophilised to give $\text{H}_4\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE}$ (**21**) as a white solid. ESI-MS ($\text{H}_2\text{O/MeCN}$ (1:1)) 1943.3 $[\text{M}+\text{H}]^+$, 972.7 $[\text{M}+2\text{H}]^{2+}$.

Radiolabelling studies of chelate-bioconjugates

[0164] The concentration-dependent radiolabelling of $\text{H}_4\text{noneunpaX-Ahx-Tyr}^3\text{-TATE}$ and $\text{H}_4\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE}$ was evaluated with $[\text{}^{155}\text{Tb}]\text{Tb}^{3+}$ and $[\text{}^{255}\text{Ac}]\text{Ac}^{3+}$ as depicted in FIG. 33. Each of the chelate-bioconjugates attained quantitative RCCs at a concentration of 10^{-5} M within 10 min at ambient temperature. The decrease in RCC at lower concentrations in comparison to the free bifunctional chelator is typical for peptide-based bioconjugates, which is attributed to the steric influence of the targeting vector on the metal binding cavity, in addition to the lower solubility of the constructs in aqueous solution and thus availability for metal ion coordination. Assessment of the serum stability for each of the radiolabelled bioconjugates showed excellent stability over the course of the study, with no significant changes in radiochemical purity from the initial time-points.

[0165] For concentration-dependent radiolabelling studies, the following general protocol was applied for radiolabelling with different radiometal ions. Stocks solution of each respective chelating ligand were prepared in ultra-pure de-ionised H₂O at a concentration of 1x10⁻² M. Serial dilution series for each chelator were prepared prior to radiolabelling studies, over a concentration range of 10⁻³ to 10⁻⁶ M. Aliquots (10 µL) of each stock solution were added to NaOAc (0.1 M), NH₄OAc (0.5 or 1.0 M) or MES buffer (1.0 M) (90 µL) to give reaction solutions suitable for study. Aliquots (1- 10 µL) of each respective radionuclide were added, under the following conditions: [⁴⁴Sc]Sc³⁺ (1.2 MBq) in NaOAc (0.1 M, pH 4.5), [¹¹¹In]In³⁺ (1.0 MBq) in NH₄OAc (0.5 M, pH 5.8), [^{132/135}La]La³⁺ (400 kBq) in NH₄OAc (0.2 M, pH 7.0), [¹⁵⁵Tb]Tb³⁺ (40 kBq) in NH₄OAc (0.5 M, pH 6.0), [¹⁷⁷Lu]Lu³⁺ (150 kBq) in NH₄OAc (0.5 M, pH 6.0), [²¹³Bi]Bi³⁺ (680 kBq) in MES (1.0 M, pH 5.5), [²²⁵Ac]Ac³⁺ (40 kBq) in NH₄OAc (1.0 M, pH 7.3). Reactions were carried out at RT and monitored over 10 minutes, with the exception [²¹³Bi]Bi³⁺ which were monitored after 5 minutes, within 10 minutes post-generator elution (n=4-8). Reactions with DOTA were carried out at elevated temperatures (85-90 °C) and monitored over 30-60 minutes. Radiochemical yields (RCYs) determined *via* iTLC using SA-paper plates and EDTA (50 mM, pH 5.0 or 7.0) as eluent (n = 4). In the case of [²¹³Bi]Bi³⁺ RCYs were further confirmed through gamma spectroscopy measurements by analysis of the baseline and solvent front TLC peaks using a high-purity germanium (HPGe) detector and monitoring the 440 keV gamma emission line of ²¹³Bi. In all studies, separate control reactions were performed in parallel by addition of radioactivity to solutions containing buffer (90 µL) and de-ionised H₂O (10 µL).

[0166] Human serum stability studies were carried out by addition of each radiolabelled complex or bioconjugate (100 µL) to vials containing human serum albumin (100 µL) and incubating the resulting solutions at 37 °C over 5-7 days. The radiochemical purity (RCP) was determined by iTLC using SA-paper plates and EDTA (50 mM, pH 5.5 or 7.0) as eluent, whereby transchelated radioactivity migrates with the solvent front (*R_f* = 1.0) while intact metal complexes remain at the baseline (*R_f* = 0). All studies were performed in triplicate, and the average %RCP used for assessment of each compound.

[0167] Dose escalation studies of [¹⁷⁷Lu][Lu(noneunpaX-Bn-NH₂)] were carried out by sequential additions of H₄noneunpaX-Bn-NH₂ (2-8 µL, 10⁻⁴ M, 20-80 pmol) to a solution containing [¹⁷⁷Lu]LuCl₃ (20 MBq) in NH₄OAc buffer (40 µL, 0.5 M, pH 6.0). The reaction solution was allowed to stand for 10 min. at RT between each addition, after which the RCY

was determined by spotting 1 μ L of solution onto SA-paper TLC plates and development with EDTA (50 mM, pH 5.5).

LogD_{7.4} measurements

- [0168]** Prior to investigation of the radiotracer performance *in vivo*, the lipophilicity of the radiolabelled chelate-bioconjugates was determined through measurement of the distribution coefficients between *n*-octanol and PBS (0.01 M, pH 7.4), Table 5. All four radio-labelled tracers are moderately hydrophilic with logD_{7.4} values ranging between -1.933 to -2.585, which are comparable to those reported for [¹⁶¹Tb][Tb(DOTA-LM3)] (logD_{7.4} = -2.5 $\pm$ 0.1), an SSTR2 antagonist analogue of DOTATATE.³⁹ As anticipated, the higher lipophilicity of the aliphatic hexyl covalent linker is reflected in the logD_{7.4} measurements, with both Ahx-Tyr³-TATE tracers being more lipophilic than the PEG₂-Tyr³-TATE derivatives. Notably, interchange between [¹⁵⁵Tb]Tb³⁺ and [²²⁵Ac]Ac³⁺ did not have a significant impact on the logD_{7.4} values for either bioconjugate, which may indicate a comparable biodistribution profile *in vivo* using this theranostic pair.
- [0169]** Aliquots of [¹⁵⁵Tb]Tb³⁺ or [²²⁵Ac]Ac³⁺ radiolabelled H₄noneunpaX-Ahx-Tyr³-TATE or H₄noneunpaX-PEG₂-Tyr³-TATE (10 μ L) were added to a biphasic mixture of *n*-octanol (700 μ L) and PBS (700 μ L, pH 7.4). The mixtures were vortexed for 2 min at RT, and then separated via centrifuge (10 min, 3000 RPM). Aliquots of *n*-octanol (100 μ L) and PBS (100 μ L) were collected, and the activity in each portion determined via gamma spectroscopy. LogD_{7.4} measurements were carried out 5-6 replicates per radiotracer. The LogD_{7.4} is defined as: log₁₀ [(*n*-octanol phase)/(buffer phase)].

Table 5. Distribution coefficients (LogD_{7.4}) for H₄noneunpaX-Ahx-Tyr³-TATE and H₄noneunpaX-PEG-Tyr³-TATE labelled with [²²⁵Ac]Ac³⁺ and [¹⁵⁵Tb]Tb³⁺, measured between PBS and *n*-octanol (n = 5-6).

Compound	LogD _{7.4}
[²²⁵ Ac][Ac(noneunpaX-Ahx-Tyr ³ -TATE)]	-1.933 $\pm$ 0.079
[²²⁵ Ac][Ac(noneunpaX-PEG ₂ -Tyr ³ -TATE)]	-2.507 $\pm$ 0.095
[¹⁵⁵ Tb][Tb(noneunpaX-Ahx-Tyr ³ -TATE)]	-2.006 $\pm$ 0.061

$$[^{155}\text{Tb}][\text{Tb}(\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE})] \quad -2.585 \pm 0.042$$

In vivo SPECT/CT and biodistribution studies

[0170] *In vivo* SPECT/CT studies of [¹⁵⁵Tb]Tb³⁺-labelled H₄noneunpaX-Ahx-Tyr³-TATE and H₄noneunpaX-PEG₂-Tyr³-TATE were performed in NRG mice bearing AR42J tumour xenografts to assess the suitability of [¹⁵⁵Tb]Tb³⁺ as an imaging companion to [²²⁵Ac]Ac³⁺. Each [¹⁵⁵Tb]Tb³⁺-labelled radiotracer was successfully prepared at high molar activity (~23.6 MBq/nmol) with a high radiochemical purity (>98%) as confirmed by iTLC and radio-HPLC (FIGs. 35-37). For *in vivo* SPECT/CT studies, mice were administered with 13.5 MBq (0.572 nmol) of each radiotracer, while biodistribution studies were performed using 0.80 MBq (0.033 nmol) per subject.

[0171] Dynamic SPECT/CT scans of mice bearing AR42J tumour xenografts were acquired over a 1 h period to assess the pharmacokinetics profile of both [¹⁵⁵Tb]Tb-labelled radiotracers after intravenous administration, FIG. 37. Further static SPECT/CT images were recorded at 2 h and 4 h post-administration, and standardised uptake values (SUVs) for regions of interest (ROIs) were extracted from the quantitative image scans to generate time-activity curves for each radiotracer, FIGs. 38 and 39.

[0172] In both cases, the [¹⁵⁵Tb]Tb³⁺-labelled radiotracers exhibit rapid clearance from blood circulation over the first 1 h after administration, with uptake in the tumour beginning within the first 5 minutes p.i. Both tracers followed the typical pharmacokinetic profile observed for hydrophilic octreotate-based bioconjugates, whereby fast clearance and accumulation in the kidneys and bladder is seen.^{40,41} Comparatively, both tracers exhibited very similar overall distribution over the course of the study, with [¹⁵⁵Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)] showing a marginally faster accumulation in the AR42J tumour xenografts, which is accompanied by faster clearance over time compared to [¹⁵⁵Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)]. This observation is clearly in line with the expected trend, accounting for the difference lipophilicity between these bioconjugates. Notably, good contrast in the tumour region is seen after 45 min p.i. with maximum uptake peaking at 2 h p.i.

[0173] Clearance of both radiotracers occurs predominantly through the renal pathway, as observed by high uptake in the kidneys and bladder, and low uptake in the liver. The SPECT/CT images show additional clearance *via* the biliary tract, whereby an increase in

activity is seen in the gallbladder beginning at 2 h p.i., with subsequent uptake in the small intestines (ileum, duodenum) at later time-points. This additional elimination pathway may account for the sustained tumour uptake over 1-2 h, in contrast to [^{177}Lu][Lu(DOTATATE)],⁴⁰ due to reabsorption of the radiotracers in the gastrointestinal tract (enterohepatic circulation) which is commonly seen for drugs exhibiting clearance in the bile.⁴² From assessment of the time-activity curves, the differences in hydrophilicity of each tracer are clearly reflected in the pharmacokinetic profiles; with [^{155}Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)] showing higher renal excretion, while [^{155}Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)] shows greater gallbladder uptake.

[0174] SSTR2 is expressed at low levels in normal healthy tissues, including the pancreas, lungs, stomach, adrenal glands, and kidneys, which are clearly visualised in the SPECT/CT images and biodistribution data determined for each tracer, FIG. 40. However, significantly lower accumulation was observed in the pancreas ($2.39 \pm 0.18\%$ ID/g and $2.43 \pm 0.68\%$ ID/g), lungs ($6.93 \pm 1.60\%$ ID/g and $7.91 \pm 0.47\%$ ID/g) and adrenal glands ($4.84 \pm 1.11\%$ ID/g and $6.28 \pm 1.12\%$ ID/g) at 4h post-injection for [^{155}Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)] and [^{155}Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)] respectively, compared to [^{177}Lu][Lu(DOTATATE)] [pancreas (10.9% ID/g), lungs (17.4% ID/g) and adrenal glands (8.95% ID/g)].⁴⁰ While higher uptake in the kidneys ($19.3 \pm 2.78\%$ ID/g and $20.5 \pm 3.57\%$ ID/g) was seen compared to [^{177}Lu][Lu(DOTATATE)] (6.31% ID/g),⁴⁰ the biodistribution study at 2 h p.i. for [^{155}Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)] and time-activity curves for both tracers shows good clearance rather than retention in the kidneys.

[0175] The relatively low uptake in the AR42J tumours ($11.0 \pm 2.62\%$ ID/g and $8.51 \pm 0.90\%$ ID/g) at 4 h post-administration is primarily a result of the large tumour size used in these studies. As can be visualised from the SPECT/CT images, the tumours appear non-homogeneous, with regions of necrotic tissue and no radiotracer uptake; hence, the %ID/g in the tumours is likely lower than achievable. However, the primary objective of these studies was to evaluate the *in vivo* suitability of H₄noneunpaX as a new bifunctional chelating ligand. Overall, the imaging and biodistribution studies show good initial results for this chelator, with no evidence of degradation *in vivo* or release of bound activity, which would be seen as uptake in the bone over time. In addition, both Tyr³-TATE analogues show improved pharmacokinetics with lower non-target tissue accumulation compared to

[¹⁷⁷Lu][Lu(DOTATATE)] and would be of interest for further development and evaluation *in vivo*.

- [0176]** For the foregoing studies, [¹⁵⁵Tb][Tb(nonenunpaX-Ahx-Tyr³-TATE)] and [¹⁵⁵Tb][Tb(nonenunpaX-PEG₂-Tyr³-TATE)] were prepared with high molar activities (23.6 MBq/nmol and 22.5 MBq/nmol, respectively), suitable for *in vivo* SPECT/CT imaging and biodistribution studies. An aliquot of [¹⁵⁵Tb]Tb³⁺ (28 MBq, 40 µL) was added to a solution of H₄noneunpaX-Ahx-Tyr³-TATE (6 µL, 2x10⁻³ M, 1.2 nmol) in NH₄OAc buffer (10 µL, 0.5 M, pH 5.5). The pH was adjusted to neutral by addition of NaOH (1 M, 1 µL) and the resulting solution incubated at 37 °C for 15 minutes to ensure quantitative incorporation of radioactivity. The same radiolabelling protocol was applied to [¹⁵⁵Tb][Tb(nonenunpaX-PEG₂-Tyr³-TATE)]. Quality control measurements were performed via iTLC measurements and radio-HPLC; method: A: H₂O (0.1% TFA), B: MeCN (0.1% TFA), 100% A to 60% B; 15 min, 1 mL/min., [¹⁵⁵Tb][Tb(nonenunpaX-Ahx-Tyr³-TATE)] (*t_R* = 9.99 min., 99%), [¹⁵⁵Tb][Tb(nonenunpaX-PEG₂-Tyr³-TATE)] (*t_R* = 9.78 min., 96%) (FIGS. 35 and 36). A small aliquot of each radiolabelled tracer was taken for quantification of radioactivity using gamma spectroscopy. No further purification was performed prior to administration of either radiotracer. Each of the radiolabelled bioconjugates was divided into two different stocks and diluted with PBS, to provide doses suitable for SPECT/CT imaging (10.2 – 13.5 MBq per subject) or biodistribution studies (~800 kBq per subject).
- [0177]** Tumour implantation was performed at the BCCRC under the protocol approved by the Animal Care Committee (ACC) of the University of British Columbia (A20-0113). Female NRG mice were anesthetized by inhalation with 2% isoflurane in 2.0 L/min of oxygen and inoculated with a AR42J exocrine pancreatic tumour cell line, subcutaneously on the left shoulder. *In vivo* imaging and biodistribution studies were performed after tumour growth reached ~ 8 – 10 mm in diameter (2-3 weeks post inoculation).

- [0178]** Animal studies were performed in accordance with the Canadian Council on Animal Care (CCAC) using the protocol approved by the Animal Care Committee (ACC) of the University of British Columbia (A20-0132). Female NRG mice bearing AR42J exocrine pancreatic tumour xenografts were anesthetized with 5% isoflurane in an induction chamber and restrained in a tail vein restrainer (Braintree Scientific) while under a continuous stream of 1-1.5% isoflurane. Mice were administered with either [¹⁵⁵Tb][Tb(noneunpaX-Ahx-Tyr³-TATE) (10.2 MBq, 23.6 MBq/nmol) or [¹⁵⁵Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE) (13.5 MBq,

22.5 MBq/nmol) in PBS (100 μ L) via lateral tail vein. Animal subjects were maintained at constant body temperature using a blanket on a heated bed, maintained under continuous stream of 1.5-2% isoflurane, and the respiration rate monitored throughout the duration of each scan. Immediately after injection, whole-body dynamic SPECT/CT imaging scans were acquired over the first 60 min using a multimodal VECTor/CT system (MILabs, Netherlands) equipped with an extra ultra high sensitivity (XUHS) 2-mm pinhole collimator. The mouse whole-body region was centered with a 14 mm axial field of view, and dynamic imaging consisting of six frames of 10 min were acquired over the first 60 min., after which static SPECT/CT scans were recorded at 2 h and 4 h post-injection using single frames of 20 min. acquisitions. Energy windows centred on the 44, 85 and 106 keV photopeaks of ^{155}Tb were applied, with a spectral width of 25%. For quantitative analysis, the SPECT images were reconstructed using pixel-based ordered-subset expectation maximization (POSEM) reconstruction algorithm using a voxel size of 0.4 mm³, 16 subsets with 6 iterations (96 MLEM equivalent). SPECT images were decay corrected and attenuation factors applied based on CT acquisitions at each time-point.⁵³ A calibration factor relating (counts/voxel) to radioactivity concentration was previously determined by measurement of a known source of ^{155}Tb . Spherical volumes of interest (VOIs) (3 mm diameter) were drawn using AMIDE (v. 1.0.4) software to determine the pharmacokinetic profile of the tracer in target organs of interest. Average standardised uptake values (SUVs) were subsequently extracted from the SPECT images. The standardised uptake value was defined according to the equation: SUV (g/mL) = radioactivity concentration (MBq/mL) / [administered dose (MBq) / body weight (g)]. Gaussian filtering (FWHM = 3 mm) and image rendering was carried out post-reconstruction for data visualisation purposes only.

[0179] Biodistribution studies with [^{155}Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)] and [^{155}Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)] were undertaken in NRG mice bearing AR42J exocrine pancreatic tumour xenografts. Prior to administration of each radiotracer (~800 kBq, 0.033 nmol) in PBS (~100 μ L), mice were anaesthetized by inhalation of 2% isoflurane and restrained using a Tail vein restrainer (Braintree Scientific). Intravenous administration of each radiotracer occurred via the lateral tail vein. After injection, mice were allowed to roam freely in their cages and sacrificed at 2 h or 4 h post-injection by CO₂ asphyxiation under 2% isoflurane anesthesia. Cardiac puncture was performed immediately after sacrifice to recover blood, and organs of interest were harvested, rinsed with PBS and

blotted dry. Each organ was weighed, and the radioactivity measured using a calibrated gamma counter (Packard Cobra II Auto-gamma counter, Perkin Elmer, Waltham, MA, USA) with a 1 min acquisition time per sample. All radioactivity measurements were decay corrected to the time of injection, and the injected dose per gram of tissue (%ID/g) calculated based on measured organ weights, with the exception of blood, bone and muscle which were scaled in accordance with literature values.⁵⁴

General Methods and Materials

[0180] All solvents and reagents were purchased from commercial suppliers (Sigma-Aldrich, AK Scientific, Alfa Aesar) and were used directly without further purification.

Analytical thin-layer chromatography (TLC) sheets were purchased from Merck (TLC Silica gel 60 F254, aluminium sheet). Deionized H₂O (18.2 MΩ/cm at 25 °C) was obtained from a PURELAB Ultra water purification system, ELGA LabWater. Flash column chromatography was performed using Siliacflash F60 silica gel (60 Å, 40–63 µm particle size, 230–400 mesh) from Silicycle Inc. Automated column chromatography was performed using a Teledyne Isco (Lincoln, NE) CombiFlash Rf automated purification system equipped with RediSep Rf Gold HP prepacked reusable silica and neutral alumina column cartridges. Low-resolution mass spectrometry (LR-MS) was performed using a Waters 2965 ZQ spectrometer with an electrospray/chemical ionization (ESI/CI) source. High-resolution mass spectrometry (HR-MS) was performed using a Waters Micromass LCT TOF instrument. Elemental analyses (CHN) were carried out using a ThermoFlash 2000 elemental analyzer. ¹H and ¹³C{¹H} NMR spectra were recorded using Bruker AV300 and AV400 spectrometers; all spectra are reported on the delta scale referenced to residual solvent peaks. Analytical and semipreparative high-performance liquid chromatography (HPLC) was carried out using a Waters 600 system equipped with a Waters 2487 dual wavelength absorbance detector monitoring at 254 and 210 nm and a Phenomenex Synergi 250 mm×21.2 mm 4µm hydro-RP80 Å column (10 mL/min). All HPLC methods employed a H₂O/MeCN biphasic solvent system buffered with 0.1% TFA. HPLC solvent system 1 (A: H₂O (0.1%TFA), B: MeCN (0.1%TFA), HPLC solvent system 2: (A: H₂O (0.01%TFA), B: MeCN). [⁴⁴Sc]Sc³⁺ (t_{1/2} = 3.97 hours) was produced at TRIUMF via proton irradiation of ^{nat}Ca targets with 12.8 MeV protons, as previously reported.²⁸ [¹¹¹In]In³⁺ (t_{1/2} = 2.83 days) (purchased from BWX Technologies) was produced by proton irradiation (Advanced Cyclotron Systems, Model TR30) via the ¹¹¹Cd(p,n)¹¹¹In reaction and provided as a 0.05 M HCl solution. [¹⁵⁵Tb]Tb³⁺ (t_{1/2}

= 5.58 days) was produced at TRIUMF via irradiation of tantalum targets with 500 MeV protons, followed by directed isotope separation online (ISOL) and implantation into NH₄Cl layered aluminium disks.⁴³ [^{132/135}La]La³⁺ (t_{1/2} = 19.5 hours) was produced via irradiation of natBa targets with 12.8 MeV protons, using a similar approach to the methods reported by Aluicio-Sarduy *et al.*⁴⁴ [¹⁷⁷Lu]Lu³⁺ (t_{1/2} = 6.67 days) was obtained from ITM Medical Isotopes GmbH Germany as a 0.05 M HCl solution. [²¹³Bi][Bi³⁺ was obtained from an in-house ²²⁵Ac/²¹³Bi generator constructed using AG-MP-50 cation exchange resin, based on established methodologies.⁴⁵ [²²⁵Ac]Ac³⁺ (t_{1/2} = 10.0 days) was produced at TRIUMF via the spallation of ²³²Th targets with 500 MeV protons, and purified as previously reported.⁴⁶

Radiolabelling of compounds was assessed via instant thin-layer chromatography (iTLC) using silicic acid (SA)-impregnated paper TLC plates sourced from Agilent technologies. TLC imaging was performed using an AR-2000 imaging scanner equipped with P-10 gas, and subsequent analysis of radiochemical conversion (RCC) was carried out using WinScan V3_14 software. Radio-HPLC was carried out using an Agilent 1200 instrument equipped with a Phenomenex Synergi 4 µm 250 mm×4.6 mm hydro-RP 80 Å column. Radioactivity was quantified using a calibrated high-purity germanium (HPGe) detector (Mirion Technologies (Canberra)Inc.) with Genie 2000 software. All work with radionuclides at TRIUMF was undertaken in shielded fume hoods to minimize dose to experimenters (and special precautions were used to prevent contamination). Peptides were prepared using an AAPPTec Focus Xi semi-automated solid phase peptide synthesiser. SPECT/CT studies were performed using a multimodal VECTor/CT system (MILabs, Netherlands) in combination with an extra ultra high sensitivity (XUHS) 2-mm pinhole collimator. Image analysis was performed using AMIDE (v. 1.0.4) software.⁴⁷

Conclusions

[0181] A new nonadentate chelating ligand, H₄noneunpaX, was synthesised to investigate the influence of donor group arrangement on metal binding characteristics within the 'NON backbone'. Characterisation of the metal complexation of H₄noneunpaX through NMR spectroscopy, mass spectrometry, radiolabelling, solution thermodynamic stability studies and DFT calculations, showed excellent compatibility with a wide range of large trivalent metal cations, with a particular preference for hard lanthanide ions. The radiolabelling properties of H₄noneunpaX were assessed with [⁴⁴Sc]Sc³⁺, [¹¹¹In]In³⁺, [^{132/235}La]La³⁺, [¹⁵⁵Tb]Tb³⁺, [¹⁷⁷Lu]Lu³⁺, [²¹³Bi]Bi³⁺ and [²²⁵Ac]Ac³⁺, which demonstrated quantitative RCC

within 10 minutes at ambient temperature, and achieved notably high molar activities with $[^{111}\text{In}]\text{In}^{3+}$ (54 GBq/ μmol), $[^{155}\text{Tb}]\text{Tb}^{3+}$ (1.0 GBq/ μmol), $[^{177}\text{Lu}]\text{Lu}^{3+}$ (2.0 GBq/ μmol) and $[^{225}\text{Ac}]\text{Ac}^{3+}$ (134 MBq/ μmol). The reactivity trends were found to be highly comparable to the structural related chelating ligand, $\text{H}_4\text{noneunpa}$, achieving similar RCYs and molar activities. The kinetic inertness of each metal complex was determined using serum challenge studies, which showed excellent *in vitro* stability (>95% RCP over 5-7 days), with the exception of $[^{225}\text{Ac}][\text{Ac}(\text{noneunpa})]$ which showed a reduction in integrity over time (~10% degradation), while $[^{225}\text{Ac}][\text{Ac}(\text{noneunpaX})]$ maintained high radiochemical integrity (>99% RCP over 7 days) thereby showing a small influence between chelate structural arrangements on complex stability. Solution thermodynamic stability studies showed the formation of highly stable metal complexes with $\text{H}_4\text{noneunpaX}$, whereby a single dominant species was observed over a broad pH range (pH = 2.0-11.5) on complexation with the lanthanide series. High stability constants were determined for each of the metal complexes, indicating highly effective metal scavenging ability.

[0182] A bifunctional analogue of $\text{H}_4\text{noneunpaX}$ was prepared using an efficient, versatile/adaptable synthetic approach, and the complexation properties reassessed, showing comparable characteristics to the unmodified chelator. Dose escalation studies of bifunctional $\text{H}_4\text{noneunpaX-Bn-NH}_2$ with $[^{177}\text{Lu}]\text{Lu}^{3+}$ demonstrated that highly competitive radiolabelling with notably high molar activities (250 GBq/ μmol). The bifunctional derivative of $\text{H}_4\text{noneunpaX}$ was conjugated to two different Tyr³-octreotate analogues for targeting of SSTR2 in NER tumours.

[0183] Preliminary *in vivo* SPECT/CT imaging, pharmacokinetics and biodistribution studies were undertaken with $[^{155}\text{Tb}]\text{Tb}^{3+}$ -radiolabelled $\text{H}_4\text{noneunpaX-Ahx-Tyr}^3\text{-TATE}$ and $\text{H}_4\text{noneunpaX-PEG}_2\text{-Tyr}^3\text{-TATE}$ in NRG mice bearing AR42J exocrine pancreatic tumour xenografts. Both radiolabelled bioconjugates showed good *in vivo* performance, with no evident degradation over the course of the study and good tumour uptake ($13.3 \pm 0.18\% \text{ID/g}$ at 2 h). The biodistribution profile of each tracer showed the typical expected clearance profile of SSTR2 targeting tracers (i.e. $[^{177}\text{Lu}][\text{Lu}(\text{DOTATATE})]$), with primarily elimination through the renal pathway, with uptake in the kidneys and bladder, followed by elimination in the urine. Notably, lower accumulation was seen in non-target organs expressing SSTR2 (adrenal glands, lungs, pancreas) compared to $[^{177}\text{Lu}][\text{Lu}(\text{DOTATATE})]$. Ultimately, these studies demonstrate $\text{H}_4\text{noneunpaX}$ as a valuable, new chelating ligand with beneficial

characteristics for e.g. $[^{225}\text{Ac}]\text{Ac}^{3+}$ applications in targeted alpha therapy, in combination with other suitable radionuclides for diagnostic imaging. The mild conditions required for radiolabelling of $\text{H}_4\text{noneunpaX}$ are particularly well suited for combination with thermally sensitive antibody-based targeting vectors and thus it can be soundly predicted that there is a high likelihood that $\text{H}_4\text{noneunpaX}$ will have good utility as a chelator for the targeted in vivo delivery of radiometals.

[0184] The objectives of this work were to investigate the influence of donor group substitution about a common backbone and study the impact of inverted group placement on metal ion affinity. $\text{H}_4\text{noneunpaX}$ was developed to this affect and studied in direct comparison to $\text{H}_4\text{noneunpa}$. An inverted arrangement of donor pendent arms was found to give comparable metal ion binding characteristics to the symmetrically derivatised chelating ligand, with the exception of $[^{213}\text{Bi}]\text{Bi}^{3+}$. High radiochemical yields (RCYs) and molar activities for $\text{H}_4\text{noneunpaX}$ were achieved with $[^{111}\text{In}]\text{In}^{3+}$, $[^{155}\text{Tb}]\text{Tb}^{3+}$, $[^{177}\text{Lu}]\text{Lu}^{3+}$, and $[^{225}\text{Ac}]\text{Ac}^{3+}$ under mild conditions (RT, 10 min), in addition to demonstrating high kinetic inertness when challenge in human serum. Cold complexation studies and DFT simulations indicate highly versatile metal ion chelation with $\text{H}_4\text{noneunpaX}$, with a particular preference for chelation of trivalent lanthanide ions.

[0185] To further assess the utility of this approach, a bifunctional analogue of $\text{H}_4\text{noneunpaX}$ was prepared and radiolabelling studies with $[^{44}\text{Sc}]\text{Sc}^{3+}$ and $[^{177}\text{Lu}]\text{Lu}^{3+}$ performed. Structural modification on the pendent donor groups retained the metal chelation characteristics of the original chelating ligand. Given the synthetic accessibility of bifunctional chelating ligands featuring this inverted donor group arrangement (in contrast to symmetrically derivatised bifunctional chelators), further investigation of $\text{H}_4\text{noneunpaX}$ *in vivo* is of high interest. The versatile coordination characteristics and mild labelling conditions of $\text{H}_4\text{noneunpaX}$, may be particularly well suited to applications in targeted alpha therapy (TAT) involving $[^{225}\text{Ac}]\text{Ac}^{3+}$ in combination with appropriate imaging radionuclides for dosimetry evaluations ($[^{135}\text{La}]\text{La}^{3+}$, $[^{155}\text{Tb}]\text{Tb}^{3+}$, $[^{111}\text{In}]\text{In}^{3+}$, $[^{44}\text{Sc}]\text{Sc}^{3+}$ etc).

[0186] While a number of exemplary aspects and embodiments have been discussed above, those of skill in the art will recognize certain modifications, permutations, additions and sub-combinations thereof. It is therefore intended that the following appended claims and claims hereafter introduced are interpreted to include all such modifications,

permutations, additions and sub-combinations as are consistent with the broadest interpretation of the specification as a whole.

References

- 5 [0187] The following references are of interest with respect to the subject matter described herein. Each of the following references is incorporated by reference in its entirety herein.
- (1) Hancock, R. D.; Martell, A. E. Ligand Design for Selective Complexation of Metal Ions in Aqueous Solution. *Chem. Rev.* **1989**, *89*, 1875–1914. <https://doi.org/10.1021/cr00098a011>.
 - 10 (2) Hancock, R. D.; Wade, P. W.; Ngwenya, M. P.; de Sousa, A. S.; Damu, K. V. Ligand Design for Complexation in Aqueous Solution. 2. Chelate Ring Size as a Basis for Control of Size-Based Selectivity for Metal Ions. *Inorg. Chem.* **1990**, *29*, 1968–1974. <https://doi.org/10.1021/ic00335a039>.
 - (3) Price, E. W.; Orvig, C. Matching Chelators to Radiometals for Radiopharmaceuticals. *Chem. Soc. Rev.* **2014**, *43*, 260–290. <https://doi.org/10.1039/c3cs60304k>.
 - 15 (4) Kostelnik, T. I.; Orvig, C. Radioactive Main Group and Rare Earth Metals for Imaging and Therapy. *Chem. Rev.* **2019**, *119*, 902–956. <https://doi.org/10.1021/acs.chemrev.8b00294>.
 - (5) Price, E. W.; Cawthray, J. F.; Bailey, G. A.; Ferreira, C. L.; Boros, E.; Adam, M. J.; Orvig, C. H 4octapa: An Acyclic Chelator for 111In Radiopharmaceuticals. *J. Am. Chem. Soc.* **2012**, *134*, 8670–8683. <https://doi.org/10.1021/ja3024725>.
 - 20 (6) Spreckelmeyer, S.; Ramogida, C. F.; Rousseau, J.; Arane, K.; Bratanovic, I.; Colpo, N.; Jermilova, U.; Dias, G. M.; Dude, I.; Jaraquemada-Peláez, M. D. G.; Bénard, F.; Schaffer, P.; Orvig, C. P-NO₂-Bn-H₄neunpa and H₄neunpa-Trastuzumab: Bifunctional Chelator for Radiometal pharmaceuticals and 111In Immuno-Single Photon Emission Computed Tomography Imaging. *Bioconjug. Chem.* **2017**, *28*, 2145–2159. <https://doi.org/10.1021/acs.bioconjchem.7b00311>.
 - 25 (7) Li, L.; Jaraquemada-Peláez, M. D. G.; Kuo, H. T.; Merkens, H.; Choudhary, N.; Gitschtaler, K.; Jermilova, U.; Colpo, N.; Uribe-Munoz, C.; Radchenko, V.; Schaffer, P.; Lin, K. S.; Bénard, F.; Orvig, C. Functionally Versatile and Highly Stable Chelator for 111In and 177Lu: Proof-of-Principle Prostate-Specific Membrane Antigen Targeting. *Bioconjug. Chem.* **2019**, *30*, 1539–1553. <https://doi.org/10.1021/acs.bioconjchem.9b00225>.
 - 30 (8) Wharton, L.; Kurakina, E.; Radchenko, V.; Schaffer, P.; Orvig, C. Chemical Promiscuity of Non-Macrocyclic Multidentate Chelating Ligands for Radiometal Ions: H₄neunpa-NH₂ vs H₄noneunpa. *Inorg. Chem.* **2021**, *60*, 4076–4092. <https://doi.org/10.1021/acs.inorgchem.1c00152>.
 - 35 (9) Wang, X.; Jaraquemada-Peláez, M. D. G.; Rodríguez-Rodríguez, C.; Cao, Y.; Buchwalder, C.; Choudhary, N.; Jermilova, U.; Ramogida, C. F.; Saatchi, K.; Häfeli, U. O.; Patrick, B. O.; Orvig, C. H₄octox: Versatile Bimodal Octadentate Acyclic Chelating Ligand for Medicinal Inorganic Chemistry. *J. Am. Chem. Soc.* **2018**, *140* (45), 15487–15500. <https://doi.org/10.1021/jacs.8b09964>.
 - 40 (10) Southcott, L.; Wang, X.; Wharton, L.; Yang, H.; Radchenko, V.; Kubeil, M.; Stephan, H.; de Guadalupe Jaraquemada-Peláez, M.; Orvig, C. High Denticity Oxinate-Linear-

Backbone Chelating Ligand for Diagnostic Radiometal Ions [^{111}In] In^{3+} and [^{89}Zr] Zr^{4+} . *Dalt. Trans.* **2021**, 50 (11), 3874–3886.

<https://doi.org/10.1039/d0dt04230g>.

- 5 (11) Price, E. W.; Zeglis, B. M.; Cawthray, J. F.; Ramogida, C. F.; Ramos, N.; Lewis, J. S.; Adam, M. J.; Orvig, C. H₄octapa-Trastuzumab: Versatile Acyclic Chelate System for ^{111}In and ^{177}Lu Imaging and Therapy. *J. Am. Chem. Soc.* **2013**, 135 (34), 12707–12721. <https://doi.org/10.1021/ja4049493>.
- (12) Jaraquemada-Peláez, M. D. G.; Wang, X.; Clough, T. J.; Cao, Y.; Choudhary, N.; Emler, K.; Patrick, B. O.; Orvig, C. H₄octapa: Synthesis, Solution Equilibria and
10 Complexes with Useful Radiopharmaceutical Metal Ions. *Dalt. Trans.* **2017**, 42, 14647–14658. <https://doi.org/10.1039/c7dt02343j>.
- (13) Hu, A.; Keresztes, I.; MacMillan, S. N.; Yang, Y.; Ding, E.; Zipfel, W. R.; Distasio, R. A.; Babich, J. W.; Wilson, J. J. Oxyaapa: A Picolinate-Based Ligand with Five Oxygen Donors That Strongly Chelates Lanthanides. *Inorg. Chem.* **2020**, 59, 5116–5132.
15 <https://doi.org/10.1021/acs.inorgchem.0c00372>.
- (14) Kwekkeboom, D. J.; Kam, B. L.; Van Essen, M.; Teunissen, J. J. M.; Van Eijck, C. H. J.; Valkema, R.; De Jong, M.; De Herder, W. W.; Krenning, E. P. Somatostatin Receptor-Based Imaging and Therapy of Gastroenteropancreatic Neuroendocrine Tumors. *Endocrine-Related Cancer*. 2010. <https://doi.org/10.1677/ERC-09-0078>.
- 20 (15) Hennrich, U.; Kopka, K. Lutathera®: The First FDA-and EMA-Approved Radiopharmaceutical for Peptide Receptor Radionuclide Therapy. *Pharmaceuticals*. 2019. <https://doi.org/10.3390/ph12030114>.
- (16) Müller, C.; Van Der Meulen, N. P.; Benešová, M.; Schibli, R. Therapeutic Radiometals beyond ^{177}Lu and ^{90}Y : Production and Application of Promising α -Particle, β^- -Particle, and Auger Electron Emitters. *Journal of Nuclear Medicine*. 2017, pp 91S–96S. <https://doi.org/10.2967/jnumed.116.186825>.
- 25 (17) Jain, A. K.; Raut, R.; Tuli, J. K. Nuclear Data Sheets for $A = 225$. *Nucl. Data Sheets* **2009**, 110 (6), 1409–1472. <https://doi.org/10.1016/j.nds.2009.04.003>.
- (18) Shannon, R. D. Revised Effective Ionic Radii and Systematic Studies of Interatomic Distances in Halides and Chalcogenides. *Acta Crystallogr. Sect. A* **1976**, A32, 751–767. <https://doi.org/10.1107/S0567739476001551>.
- 30 (19) Deblonde, G. J. P.; Zavarin, M.; Kersting, A. B. The Coordination Properties and Ionic Radius of Actinium: A 120-Year-Old Enigma. *Coordination Chemistry Reviews*. 2021, p 214130. <https://doi.org/10.1016/j.ccr.2021.214130>.
- 35 (20) Nica, N. Nuclear Data Sheets for $A=155$. *Nucl. Data Sheets* **2019**, 160, 1–404. <https://doi.org/10.1016/j.nds.2019.100523>.
- (21) Webster, B.; Ivanov, P.; Russell, B.; Collins, S.; Stora, T.; Ramos, J. P.; Köster, U.; Robinson, A. P.; Read, D. Chemical Purification of Terbium-155 from Pseudo-Isobaric Impurities in a Mass Separated Source Produced at CERN. *Sci. Rep.* **2019**,
40 9 (1). <https://doi.org/10.1038/s41598-019-47463-3>.
- (22) Müller, C.; Domnanich, K. A.; Umbricht, C. A.; Van Der Meulen, N. P. Scandium and Terbium Radionuclides for Radiotheranostics: Current State of Development towards Clinical Application. *British Journal of Radiology*. 2018. <https://doi.org/10.1259/bjr.20180074>.
- 45 (23) Naskar, N.; Lahiri, S. Theranostic Terbium Radioisotopes: Challenges in Production for Clinical Application. *Frontiers in Medicine*. 2021. <https://doi.org/10.3389/fmed.2021.675014>.
- (24) Gans, P.; Sabatini, A.; Vacca, A. Determination of Equilibrium Constants from Spectrophotometric Data Obtained from Solutions of Known PH: The Program PHab.

Ann. Chim. **1999**, *89* (1–2), 45–49.


- (25) Gans, P.; Sabatini, A.; Vacca, A. Investigation of Equilibria in Solution. Determination of Equilibrium Constants with the HYPERQUAD Suite of Programs. *Talanta* **1996**, *43* (10), 1739–1753. [https://doi.org/10.1016/0039-9140\(96\)01958-3](https://doi.org/10.1016/0039-9140(96)01958-3).
- 5 (26) Martell, A. E.; Smith, R. M. *Critical Stability Constants*; 1982. <https://doi.org/10.1007/978-1-4615-6761-5>.
- (27) Blachot, J. Nuclear Data Sheets for A = 111. *Nucl. Data Sheets* **2009**, *110* (6), 1239–1407. <https://doi.org/10.1016/j.nds.2009.04.002>.
- 10 (28) Kurakina, E. S.; Wharton, L.; Hoehr, C.; Orvig, C.; Magomedbekov, E. P.; Filosofov, D.; Radchenko, V. Improved Separation Scheme for ⁴⁴Sc Produced by Irradiation of NatCa Targets with 12.8 MeV Protons. *Nucl. Med. Biol.* **2022**, *104–105*, 22–27. <https://doi.org/10.1016/j.nucmedbio.2021.11.002>.
- 15 (29) Fiszbein, D. J.; Brown, V.; Thiele, N. A.; Woods, J. J.; Wharton, L.; Macmillan, S. N.; Radchenko, V.; Ramogida, C. F.; Wilson, J. J. Tuning the Kinetic Inertness of Bi³⁺Complexes: The Impact of Donor Atoms on Diaza-18-Crown-6 Ligands as Chelators for ²¹³Bi Targeted Alpha Therapy. *Inorg. Chem.* **2021**, *60* (12), 9199–9211. <https://doi.org/10.1021/acs.inorgchem.1c01269>.
- 20 (30) Pujales-Paradela, R.; Rodríguez-Rodríguez, A.; Gayoso-Padula, A.; Brandariz, I.; Valencia, L.; Esteban-Gómez, D.; Platas-Iglesias, C. On the Consequences of the Stereochemical Activity of the Bi(III) 6s² Lone Pair in Cyclen-Based Complexes. the [Bi(DO3A)] Case. *Dalt. Trans.* **2018**, *47* (39), 13830–13842. <https://doi.org/10.1039/c8dt02602e>.
- 25 (31) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; Li, X.; Caricato, M.; Marenich, A. V.; Bloino, J.; Janesko, B. G.; Gomperts, R.; Mennucci, B.; Hratchian, H. P.; Ortiz, J. V.; Izmaylov, A. F.; Sonnenberg, J. L.; Williams-Young, D.; Ding, F.; Lipparini, F.; Egidi, F.; Goings, J.; Peng, B.; Petrone, A.; Henderson, T.; Ranasinghe, D.; Zakrzewski, V. G.; Gao, J.; Rega, N.; Zheng, G.; Liang, W.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Throssell, K.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M. J.; Heyd, J. J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Millam, J. M.; Klene, M.; Adamo, C.; Cammi, R.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Farkas, O.; Foresman, J. B.; Fox, D. J. Gaussian 16, Rev. B.01. *Gaussian, Inc., Wallingford, CT* **2016**.
- 30 (32) Tomasi, J.; Mennucci, B.; Cammi, R. Quantum Mechanical Continuum Solvation Models. *Chem. Rev.* **2005**, *105*, 2999–3094. <https://doi.org/10.1021/cr9904009>.
- 35 (33) Cao, X.; Dolg, M. Valence Basis Sets for Relativistic Energy-Consistent Small-Core Lanthanide Pseudopotentials. *J. Chem. Phys.* **2001**, *115*, 7348–7355. <https://doi.org/10.1063/1.1406535>.
- 40 (34) Cao, X.; Dolg, M. Segmented Contraction Scheme for Small-Core Lanthanide Pseudopotential Basis Sets. *J. Mol. Struct. THEOCHEM* **2002**, *581*, 139–147. [https://doi.org/10.1016/S0166-1280\(01\)00751-5](https://doi.org/10.1016/S0166-1280(01)00751-5).
- 45 (35) Stoll, H.; Metz, B.; Dolg, M. Relativistic Energy-Consistent Pseudopotentials - Recent Developments. *J. Comput. Chem.* **2002**, *23* (8), 767–778. <https://doi.org/10.1002/jcc.10037>.
- (36) Glendening, E. D.; Reed, A. E.; Carpenter, J. E.; Weinhold, F. NBO 3.0. *QCPE Bull.* **1990**, *10*.

- (37) Chong, H. S.; Song, H. A.; Ma, X.; Milenic, D. E.; Brady, E. D.; Lim, S.; Lee, H.; Baidoo, K.; Cheng, D.; Brechbiel, M. W. Novel Bimodal Bifunctional Ligands for Radioimmunotherapy and Targeted MRI. *Bioconjug. Chem.* **2008**, *19* (7), 1439–1447. <https://doi.org/10.1021/bc800050x>.
- 5 (38) Noor, A.; Van Zuylekom, J. K.; Rudd, S. E.; Roselt, P. D.; Haskali, M. B.; Yan, E.; Wheatcroft, M.; Hicks, R. J.; Cullinane, C.; Donnelly, P. S. Imaging Somatostatin Positive Tumors with Tyr3-Octreotate/Octreotide Conjugated to Desferrioxamine B Squaramide Radiolabeled with Either Zirconium-89 or Gallium-68. *Bioconjug. Chem.* **2021**, *32* (7), 1192–1203. <https://doi.org/10.1021/acs.bioconjchem.1c00109>.
- 10 (39) Borgna, F.; Barritt, P.; Grundler, P. V.; Talip, Z.; Cohrs, S.; Zeevaart, J. R.; Köster, U.; Schibli, R.; van der Meulen, N. P.; Müller, C. Simultaneous Visualization of ¹⁶¹Tb- and ¹⁷⁷Lu-Labeled Somatostatin Analogues Using Dual-Isotope SPECT Imaging. *Pharmaceutics* **2021**, *13* (4). <https://doi.org/10.3390/pharmaceutics13040536>.
- 15 (40) Rousseau, E.; Lau, J.; Zhang, Z.; Uribe, C. F.; Kuo, H. T.; Zhang, C.; Zeisler, J.; Colpo, N.; Lin, K. S.; Bénard, F. Effects of Adding an Albumin Binder Chain on [¹⁷⁷Lu]Lu-DOTATATE. *Nucl. Med. Biol.* **2018**, *66*, 10–17. <https://doi.org/10.1016/j.nucmedbio.2018.08.001>.
- (41) Müller, C.; Fischer, E.; Behe, M.; Köster, U.; Dorrer, H.; Reber, J.; Haller, S.; Cohrs, S.; Blanc, A.; Grünberg, J.; Bunka, M.; Zhernosekov, K.; van der Meulen, N.; Johnston, K.; Türlér, A.; Schibli, R. Future Prospects for SPECT Imaging Using the Radiolanthanide Terbium-155 - Production and Preclinical Evaluation in Tumor-Bearing Mice. *Nucl. Med. Biol.* **2014**, *41* (S). <https://doi.org/10.1016/j.nucmedbio.2013.11.002>.
- 20 (42) Roberts, M. S.; Magnusson, B. M.; Burczynski, F. J.; Weiss, M. Enterohepatic Circulation: Physiological, Pharmacokinetic and Clinical Implications. *Clinical Pharmacokinetics*. 2002, pp 751–790. <https://doi.org/10.2165/00003088-200241100-00005>.
- 25 (43) Minor, G.; Kapalka, J.; Fisher, C.; Paley, W.; Chen, K.; Kinakin, M.; Earle, I.; Moss, B.; Bricault, P.; Gottberg, A. Remote Handling Systems for the ISAC and ARIEL High-Power Fission and Spallation ISOL Target Facilities at TRIUMF. *Nucl. Eng. Technol.* **2020**, 1–12. <https://doi.org/10.1016/j.net.2020.09.024>.
- 30 (44) Aluicio-Sarduy, E.; Hernandez, R.; Olson, A. P.; Barnhart, T. E.; Cai, W.; Ellison, P. A.; Engle, J. W. Production and in Vivo PET/CT Imaging of the Theranostic Pair ¹³²/¹³⁵La. *Sci. Rep.* **2019**, *9* (1). <https://doi.org/10.1038/s41598-019-47137-0>.
- 35 (45) Ma, D.; McDevitt, M. R.; Finn, R. D.; Scheinberg, D. A. Breakthrough of ²²⁵Ac and Its Radionuclide Daughters from an ²²⁵Ac/²¹³Bi Generator: Development of New Methods, Quantitative Characterization, and Implications for Clinical Use. *Appl. Radiat. Isot.* **2001**, *55* (5), 667–678. [https://doi.org/10.1016/S0969-8043\(01\)00062-8](https://doi.org/10.1016/S0969-8043(01)00062-8).
- (46) Robertson, A. K. H.; McNeil, B. L.; Yang, H.; Gendron, D.; Perron, R.; Radchenko, V.; Zeisler, S.; Causey, P.; Schaffer, P. ²³²Th-Spallation-Produced ²²⁵Ac with Reduced ²²⁷Ac Content. *Inorg. Chem.* **2020**, *59* (17), 12156–12165. <https://doi.org/10.1021/acs.inorgchem.0c01081>.
- 40 (47) Loening, A. M.; Gambhir, S. S. AMIDE: A Free Software Tool for Multimodality Medical Image Analysis. *Mol. Imaging* **2003**, *2* (3), 131–137. <https://doi.org/10.1162/153535003322556877>.
- 45 (48) Gran, G. Determination of the Equivalence Point in Potentiometric Titrations. Part II. *Analyst* **1952**, *77* (920), 661–670. <https://doi.org/10.1039/AN9527700661>.
- (49) Huang, P. W. Understanding the Stability Trend Along Light Lanthanide Complexes with an Ethylenediamine-Type Ligand: A Quantum Chemical Study. *ChemistrySelect*

- 2019, 4, 12368–12374. <https://doi.org/10.1002/slct.201902887>.
- (50) Kovács, A.; Varga, Z. Metal–Ligand Interactions in Complexes of Cyclen-Based Ligands with Bi and Ac. *Struct. Chem.* **2021**, 32 (5), 1719–1731. <https://doi.org/10.1007/s11224-021-01816-9>.
- 5 (51) Hanwell, M. D.; Curtis, D. E.; Lonie, D. C.; Vandermeersch, T.; Zurek, E.; Hutchison, G. R. Avogadro: An Advanced Semantic Chemical Editor, Visualization, and Analysis Platform. *J. Cheminform.* **2012**, 4 (8). <https://doi.org/10.1186/1758-2946-4-17>.
- (52) Lu, T.; Chen, F. Multiwfn: A Multifunctional Wavefunction Analyzer. *J. Comput. Chem.* **2012**, 33 (5), 580–592. <https://doi.org/10.1002/jcc.22885>.
- 10 (53) Branderhorst, W.; Vastenhouw, B.; Beekman, F. J. Pixel-Based Subsets for Rapid Multi-Pinhole SPECT Reconstruction. *Phys. Med. Biol.* **2010**, 55 (7), 2023–2034. <https://doi.org/10.1088/0031-9155/55/7/015>.
- (54) Foster, Henry L.; Small, J. David.; Fox, J. G. *The Mouse in Biomedical Research*; Academic Press: New York, 1983.
- 15 (55) Gran, G. Determination of the Equivalence Point in Potentiometric Titrations. Part II. *Analyst* **1952**, 77 (920), 661–670. <https://doi.org/10.1039/AN9527700661>.
- (56) Alderighi, L.; Gans, P.; Ienco, A.; Peters, D.; Sabatini, A.; Vacca, A. Hyperquad Simulation and Speciation (HySS): A Utility Program for the Investigation of Equilibria Involving Soluble and Partially Soluble Species. *Coord. Chem. Rev.* **1999**, 184 (1), 311–318. [https://doi.org/10.1016/S0010-8545\(98\)00260-4](https://doi.org/10.1016/S0010-8545(98)00260-4).
- 20 (57) Harris, W. R.; Carrano, C. J.; Raymond, K. N. Spectrophotometric Determination of the Proton-Dependent Stability Constant of Ferric Enterobactin. *J. Am. Chem. Soc.* **1979**, 101 (8), 2213–2214. <https://doi.org/10.1021/ja00502a053>.
- 25 (58) Milenic, D.E.; Roselli, M.; Mirzadeh, S.; Pippin, C.G.; Gansow, O.A.; Colcher, D.; Brechbiel, M.W.; Schlom, J. In Vivo Evaluation of Bismuth-Labeled Monoclonal Antibody Comparing DTPA-Derived Bifunctional Chelates. *Cancer Biother. & Radiopharm.* **2001**, 16(2), 133–146. <http://doi.org/10.1089/108497801300189227>

CLAIMS:


1. A chelator having the structure (I), wherein each R_1 is independently OH, NH or SH, and X is O, S, or NR_3 , wherein R_3 is H or $CH_2C(=O)R_1$:



5


(I)

2. A chelator as defined in claim 1 having the following structure:




3. A chelator having the structure (III) wherein each R_1 is independently OH, NH or SH or a functional group; each R_2 is independently H or a functional group; each R_4 is independently H or a functional group, or both R_4 together form a cyclohexyl moiety; each R_5 is independently H or a functional group, or both R_5 together form a cyclohexyl moiety; and X is O, S, or NR_3 , wherein one or more of the R_1 , R_2 , R_4 or R_5 groups is a functional group that allows coupling of the chelator to a biological targeting moiety, and wherein R_3 is H or $CH_2C(=O)R_1$:

10



4. A chelator as defined in claim 3 having the structure (II) wherein each R_1 is independently OH, NH or SH or a functional group, each R_2 is independently H or a functional group, and X is O, S, or NR_3 , wherein one or more of the R_1 or R_2 groups is a functional group that allows coupling of the chelator to a biological targeting moiety, and wherein R_3 is H or $CH_2C(=O)R_1$:



5. A chelator as defined in either one of claims 3 or 4, wherein the functional group is a carboxyl, an ester, an amide, an imide, a thioamide, a thioester, or a guanidinium group.
6. A chelate comprising a radiometal and a chelator as defined in any one of claims 1 to 5.

7. An in vivo radioisotope targeting construct comprising a biological targeting moiety and a chelator as defined in claim 3, wherein one R₁ group, one R₂, one R₄ group or one R₅ group is coupled to the biological targeting moiety.
8. An in vivo radioisotope targeting construct as defined in claim 7, further comprising a linker interposing the chelator and the biological targeting moiety, wherein the linker comprises a C₁-C₁₀ hydrocarbon linker that is optionally substituted with one or more heteroatoms or has one or more substituents, an aromatic linker, a cationic linker, an anionic linker, an amino acid linker having between one and ten amino acids, a cyclized amino acid linker, a PEG linker, a cyclized ring linker, an aromatic linker, or a click chemistry linker.
9. An in vivo radioisotope targeting construct as defined in any one of claims 7 or 8, further comprising a radiometal chelated by the chelator.
10. A chelate or an in vivo radioisotope targeting construct as defined in any one of claims 6 to 9, wherein the radiometal comprises ²²⁵Ac, ²²⁷Th, ²²⁶Th, ²¹¹At, ⁴⁴Sc, ⁹⁰Y, ⁸⁹Zr, ¹⁷⁷Lu, ¹¹¹In, ^{86/89/90}Y, ²¹¹At, ²¹¹Fr, ^{212/213}Bi, ¹⁵³Sm, ^{161/166}Ho, ^{165/166}Dy, ^{161/155}Tb, ¹⁴⁰La, ^{142/143/145}Pr, ¹⁵⁹Gd, ^{169/175}Yb, ^{167/170}Tm, ¹⁶⁹Er, ¹⁴⁹Pm, ¹⁵⁰Eu, ⁶⁸Ga, ¹³⁷Cs, or ¹⁴¹Ce.
11. A chelate or an in vivo radioisotope targeting construct as defined in any one of claims 6 to 9, wherein the radiometal comprises ²²⁷Th, ²²⁵Ac, ¹⁵⁵Tb, ¹⁷⁷Lu, ¹¹¹In, ¹³²La, ²³⁵La, ⁹⁰Y, ⁶⁸Ga, ⁴⁴Sc, ²⁰³Pb, or ²¹²Pb.
12. A chelate or an in vivo radioisotope targeting construct as defined in any one of claims 6 to 9, wherein the radiometal comprises ²²⁵Ac, ¹⁵⁵Tb, ¹⁷⁷Lu, ¹¹¹In, ¹³²La, ²³⁵La, or ⁴⁴Sc.
13. A chelate or an in vivo radioisotope targeting construct as defined in any one of claims 6 to 9, wherein the radiometal comprises ²²⁵Ac.
14. An in vivo radioisotope targeting construct as defined in any one of claims 7 to 13, wherein the targeting moiety comprises a hapten, an antigen, an aptamer, an affibody, an enzyme, a protein, a peptide, an antibody, an antigen-binding fragment of an antibody, a peptidomimetic, a receptor ligand, a steroid, a hormone, a growth

factor, a cytokine, a molecule that recognizes cell surface receptors, a lipid, a lipophilic group, or a carbohydrate.

15. An in vivo radioisotope targeting construct as defined in any one of claims 7 to 13, wherein the antigen-binding fragment of an antibody comprises an Fab fragment, an F(ab')₂ fragment, a Fv fragment, an scFv fragment, a minibody, or a diabody.
16. An in vivo radioisotope targeting construct as defined in any one of claims 7 to 13, wherein the biological targeting moiety comprises A33 antibody, dihydrotestosterone (DHT), HuMab-5B1, girentuximab, AMG211 bispecific T-cell engager, IAB22M2C minibody, rituximab, obinutuzumab, U36 antibody, plerixafor, , pentixafor, NFB, ipilimumab, erlotinib, PD153035, afatinib, cetuximab, panitumumab, ABY-025 affibody, HER2-nanobody, trastuzumab, pertuzumab, GSK2849330, lumretuzumab, 4FMFES, FAPI-04, FAPI-21, FAPI-46, galactose, CB-TE2A-AR06 peptide (with H₄noneunpaX substituted for DOTA), BAY 864367 peptide (with H₄noneunpaX-bound ligand label instead of 18F labeling), RM2 peptide (with H₄noneunpaX substituted for DOTA), SB3 peptide (with H₄noneunpaX substituted for DOTA), RM26 peptide, BBN-RGD peptide, Aca-BBN peptide, NeoBOMB1 peptide (with H₄noneunpaX substituted for DOTA), exendin-4 peptide, glucose, codrituzumab, EF5, MISO, AZA, HX4, ASTM, LLP2A, peptidomimetic, galacto-RGD peptide, FPP(RGD)₂ peptide, RGD-K5 peptide, fluciclatide, alfatide-I, alfatide-II, PRGD2 peptide, αβ6-BP peptide, CycMSHhex targeting peptides, MMOT0530A antibody, SP peptide, neurotensin, PARPi, a PSMA peptidomimetic, DCFPyL, DCFBC, HuJ591 antibody, durvalumab, nivolumab, pembrolizumab, BMS-986192 adnectin, atezolizumab, MSTP2109A antibody, TATE peptide (octreotate), TOC peptide, NOC peptide, JR11, thymidine, fresolimumab, or bevacizumab.
17. An in vivo radioisotope targeting construct as defined in any one of claims 7 to 16, wherein a biological target targeted by the in vivo radioisotope targeting construct comprises: a tumor associated antigen, A33 transmembrane glycoprotein, androgen receptor (AR), CA19.9, carbonic anhydrase 9 (CA-IX), carcinoembryonic antigen, CD8, CD20, CD44v6, C-X-C chemokine receptor type 4 (CXCR4), cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), epidermal growth factor receptor (EGFR), epidermal growth factor receptor 2 (ERBB2), epidermal growth factor

receptor 3 (ERBB3), estrogen receptor (ER), fibroblast activation protein α , gastrin-releasing peptide receptor (GRPR), glucagon-like peptide 1 receptor (GLP-1R), glypican 3, integrin $\alpha 4 \beta 1$, integrin $\alpha v \beta 3$, integrin $\alpha v \beta 6$, melanocortin-1 receptor (MC1R), mesothelin, neurokinin1 receptor (NK1R), neurotensin 1 receptor (NTS1R),
 5 poly(ADP-ribose) polymerase 1 (PARP1), prostate-specific membrane antigen (PSMA), programmed cell death protein (PD-1), programmed death-ligand 1 (PD-L1), six-transmembrane epithelial antigen of prostate-1 (STEAP1), somatostatin receptor 2 (SSTR2), thymidine kinase, transforming growth factor-beta (TGF- β), or vascular endothelial growth factor receptor (VEGFR).

- 10 18. An in vivo radioisotope targeting construct as defined in any one of claims 7 to 13, wherein the biological targeting moiety comprises octreotate (TATE).
19. A pharmaceutical composition comprising an in vivo radioisotope targeting construct as defined in any one of claims 7 to 18 and a pharmaceutically acceptable carrier, excipient or vehicle.
- 15 20. A method of delivering a radioisotope to a selected location within the body of a mammalian subject, the method comprising:
 administering an in vivo radioisotope targeting construct as defined in any one
 claims 7 to 18 bearing the radioisotope to the mammalian subject.
- 20 21. A method as defined in claim 20, further comprising allowing the targeting moiety of the in vivo radioisotope targeting construct to enhance the accumulation of the radioisotope at the selected location within the body relative to other locations in the body to selectively deliver radiation to the selected location.
- 25 22. A method as defined in either one of claims 20 or 21, further comprising a step of forming a chelate comprising the radioisotope and the in vivo radioisotope targeting construct prior to the administering step, wherein the step of forming the chelate comprises combining the in vivo radioisotope targeting construct with the radioisotope at a temperature of between about 10°C and about 65°C for an incubation period.

23. A method as defined in claim 22, wherein the temperature is between about 15°C and about 25°C during the incubation period.
24. A method as defined in either one of claims 22 or 23, wherein the incubation period is between about 5 minutes and about 30 minutes.
- 5 25. A method as defined in any one of claims 22 to 24, wherein the combining step is carried out at a pH in the range of about 5.0 to about 7.4.
26. A method as defined in any one of claims 22 to 25, wherein the combining step is carried out in aqueous solution that is substantially free of alcohol.
- 10 27. A method as defined in any one of claims 20 to 26, further comprising carrying out an imaging procedure to evaluate the localization of the in vivo radioisotope targeting construct within the body, wherein the imaging procedure optionally comprises positron emission tomography (PET) imaging or single-photon emission computerized tomography (SPECT) imaging.
- 15 28. A method as defined in any one of claims 20 to 26, wherein the in vivo radioisotope targeting construct is used to cause cell death at the selected location within the body by exposing the cells to radiation from the radioisotope.
29. A method as defined in claim 28, wherein the in vivo radioisotope targeting construct is used to cause death of cancer cells at the selected location within the body.
- 20 30. A method as defined in either one of claims 28 or 29, wherein the radiation comprises alpha radiation or beta radiation.
31. A method as defined in any one of claims 20 to 30, wherein the mammalian subject is a human.

25

1/21




FIG. 1




FIG. 2

2/21




FIG. 3




FIG. 4

3/21




FIG. 5




FIG. 6

4/21




FIG. 7




FIG. 8




FIG. 9






FIG. 10



7/21




FIG. 13






FIG. 14



9/21




FIG. 17






FIG. 18



11/21




FIG. 21




FIG. 22




FIG. 23

12/21




FIG. 24




FIG. 25

13/21




FIG. 26




FIG. 27

14/21




FIG. 28

15/21




FIG. 29




FIG. 30

16/21




FIG. 31




FIG. 32

17/21




FIG. 33




FIG. 34

18/21




FIG. 35





FIG. 36

19/21

[¹⁵⁵Tb][Tb(noneunpaX-Ahx-Tyr³-TATE)]



[¹⁵⁵Tb][Tb(noneunpaX-PEG₂-Tyr³-TATE)]




FIG. 37

20/21




FIG. 38

21/21




FIG. 39





FIG. 40



(I)