

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978**PUBLICATION PARTICULARS AND ABSTRACT**

(Section 32(3)(a) – Regulation 22(1)(g) and 31)

OFFICIAL APPLICATION NO.

LODGING DATE

ACCEPTANCE DATE

21 01 2002 4066

22 22 MAY 2002

43 22.5.2003

INTERNATIONAL CLASSIFICATION

NOT FOR PUBLICATION

51 C07C; C07D

CLASSIFIED BY: WIPO

FULL NAMES OF APPLICANT

71 PHARMACIA AB

FULL NAMES OF INVENTORS

72 1. ANDERSSON, PEHR G
2. HEDBERG, CHRISTIAN

EARLIEST PRIORITY CLAIMED

COUNTRY

NUMBER

DATE

33 SE

31 9904850-6

32 30 DEC 1999

TITLE OF INVENTION

54 PROCESS OF PREPARING TOLTERODINE AND ANALOGUES THEREOF AS WELL AS
INTERMEDIATES PREPARED IN THE PROCESS

57 ABSTRACT (NOT MORE THAN 150 WORDS)

NUMBER OF SHEETS 35

If no classification is finished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

Abstract

The invention relates to a process for the enantioselective preparation of tolterodine and analogues and salts thereof comprising the steps of: a) enantioselectively reducing the carbonyl function in a compound of formula (II), wherein R₁, R₂, and R₃ independently of each other are hydrogen, methyl, methoxy, hydroxy, hydroxymethyl, carbamoyl, sulphamoyl or halogen, to form an enantiomerically enriched compound of formula (IIIa) or (IIIb); b) subjecting the compound of formula (IIIa) or (IIIb) to a sigmatropic rearrangement to form a corresponding enantiomerically enriched compound of formula (IVa) or (IVb); c) subjecting the compound of formula (IVa) or (IVb) to a Baeyer-Villiger oxidation to form a corresponding enantiomerically enriched compound of formula (Va) or (Vb); d) converting the compound of formula (Va) or (Vb) to form the corresponding enantiomerically enriched compound of tolterodine or analogue thereof; and e) optionally converting a compound obtained in base form to a salt thereof, or converting a salt form to the free base. Formulas (IIIa), (IIIb), (IVa), (IVb), (Va) and (Vb) are specified in the application. The invention also relates to novel starting materials and intermediates used in the process.

PROCESS OF PREPARING TOLTERODINE AND ANALOGUES THEREOF AS WELL AS INTERMEDIATES PREPARED IN THE PROCESS

Field of the invention

The present invention relates to a novel process of preparing tolterodine and analogues thereof, as well as to novel intermediates prepared in the process.

Background of the invention

Tolterodine, i.e. (R)-N,N-diisopropyl-3-(2-hydroxy-5-methylphenyl)-3-phenylpropanamine, is useful for treating urinary incontinence. The major, active 10 metabolite of tolterodine, i.e. (R)-N,N-diisopropyl-3-(2-hydroxy-5-hydroxymethylphenyl)-3-phenylpropanamine, contributes significantly to the therapeutic effect of tolterodine. Tolterodine and analogues thereof, including the corresponding (S)-enantiomer, as well as processes for the preparation thereof are disclosed in US-A-5,382,600. The active metabolite and analogues are disclosed in 15 US-A-5,559,269. The (S)-enantiomer and its use in the treatment of urinary and gastrointestinal disorders is further described in WO 98/03067.

One of the processes described in US-A-5,382,600 comprises the steps of preparing the lactone 3,4-dihydro-6-methyl-4-phenyl-2H-benzopyran-2-one, reductively ring-opening the lactone to prepare the corresponding alcohol, reacting the alcohol with 20 isopropylamine, and resolving the racemate formed to isolate tolterodine.



US-A-5,922,914 discloses a modified process for preparing tolterodine by reducing the above-mentioned lactone to the corresponding alcohol, 3,4-dihydro-6-methyl-4-phenyl-2H-benzopyran-2-ol, reductively aminating the alcohol, and resolving the racemate formed to isolate tolterodine.

25 While the above prior art methods thus produce a racemate which has to be resolved to obtain the desired tolterodine enantiomer, Andersson, Pher G. et al., *J. Org. Chem.* 1998, 63, 8067-8070 discloses an enantioselective synthesis of tolterodine which obviates the need of the enantiomer separation step. This method comprises a copper bromide catalyzed asymmetric addition of 2-methoxy-5-methylphenylmagnesium 30 bromide to a 3-phenyl-prop-2-enoyl-oxazolidinone to produce the (5S)-phenyl-(3R)-(2-benzyloxy-5-methylphenyl)-3-phenylpropanoyl-2-oxazolidinone, hydrolyzation of the oxazolidinone to the corresponding propanoic acid, reaction with diisopropylamine to form the amide, and reduction of the amide to tolterodine.

Summary of the invention


The present invention provides an alternate enantioselective synthesis of tolterodine which is more convenient to perform than the prior art method outlined above and which gives a final product of high enantiomeric purity. A key step of the present method is the preparation of the above-mentioned lactone, 3,4-dihydro-6-methyl-4-phenyl-2H-benzopyran-2-one (also referred to as 6-methyl-4-phenyl-chroman-2-one), in an enantiomerically enriched form by enantioselective reactions.

Thus, in a first aspect the present invention provides a process for the enantioselective preparation of a compound of the general formula (Ia) or (Ib):




wherein R₁, R₂ and R₃ independently of each other are hydrogen, methyl, methoxy, hydroxy, hydroxymethyl, carbamoyl, sulphamoyl or halogen, and R₄ and R₅ independently of each other are C₁₋₆-alkyl, or a salt thereof, which process comprises the steps of:

a) enantioselectively reducing the carbonyl function in a compound of formula (II):



wherein R₁, R₂ and R₃ are as defined above, to form an enantiomerically enriched compound of formula (IIIa) or (IIIb):


5



wherein R₁, R₂ and R₃ are as defined above, or a salt thereof;


b) subjecting the compound of formula (IIIa) or (IIIb) to a sigmatropic rearrangement to form a corresponding enantiomerically enriched compound of formula
10 (IVa) or (IVb):

10 (IVa) or (IVb):



wherein R₁, R₂ and R₃ are as defined above, or a salt thereof;

c) subjecting the compound of formula (IVa) or (IVb) to a Baeyer-Villiger oxidation to form a corresponding enantiomerically enriched compound of the general formula (Va) or (Vb):




5

wherein R₁, R₂ and R₃ are as defined above or a salt thereof;

d) converting the compound of formula (Va) or (Vb) to form the corresponding enantiomerically enriched compound of formula (Ia) or (Ib), or a salt thereof; and
 10 e) optionally converting a compound of formula (Ia) or (Ib) in base form to a salt thereof, or converting a salt form to the free base.


In one embodiment of the first aspect of the invention, step d) comprises:


d1) reacting the compound of formula (Va) or (Vb) with an amine of the general formula (VI):



15

wherein R₄ and R₅ are as defined above, to form a corresponding enantiomerically enriched compound of the general formula (VIIa) or (VIIb):






wherein R_1 , R_2 , R_3 , R_4 and R_5 are as defined above; and

5 d2) reducing the carbonyl function in the compound of formula (VIIa) or (VIIb) to form the corresponding enantiomerically enriched compound of formula (Ia) or (Ib).

Optionally, steps d1) and d2) are performed simultaneously in a single step.

In an alternative embodiment, step d) comprises:

d1') reducing the compound of formula (Va) or (Vb) to form a corresponding enantiomerically enriched hydroxy compound of the general formula (VIIa) or (VIIb):





10

wherein R_1 , R_2 and R_3 are as defined in claim 1; and

d2') reductively aminating the hydroxy compound of formula (VIIIa) or (VIIIb) with the amine of formula (VI) to form the corresponding enantiomerically enriched compound of formula (Ia) or (Ib).


15

In second aspect, the present invention provides a process for the enantioselective preparation of a compound of the general formula (Va) or (Vb):




wherein R₁, R₂ and R₃ are as defined above, or a salt thereof, which process comprises the steps of:

5 a) enantioselectively reducing the carbonyl function in a compound of formula (II):





wherein R₁, R₂ and R₃ are as defined above, or a salt thereof, to form an enantiomerically enriched compound of formula (IIIa) or (IIIb):



wherein R₁, R₂ and R₃ are as defined above, or a salt thereof;

b) subjecting the compound of formula (IIIa) or (IIIb) to a sigmatropic rearrangement to form a corresponding enantiomerically enriched compound of formula (IVa) or (IVb):




5

wherein R₁, R₂ and R₃ are as defined above, or a salt thereof; and


c) subjecting the compound of formula (IVa) or (IVb) to a Baeyer-Villiger oxidation to form the corresponding enantiomerically enriched compound of the general formula (Va) or (Vb), or salt thereof.

10 The compound of formula (II) may be prepared by subjecting a compound of the general formula (IX):




wherein R₁, R₂, and R₃ are as defined in claim 1, and Hal is halogen (preferably bromine), or a salt thereof, to a reductive ring closure reaction.

The compound of formula (IX) may be prepared by reacting a compound of the general formula (X):



wherein R_1 and Hal are as defined above, with a compound of the general formula (XI):



5

wherein R_2 and R_3 are as defined above.

Preferably, compounds of formula Ia or Ib are prepared in which R_1 is methyl or hydroxymethyl in 5-position, R_2 and R_3 are hydrogen, and R_4 and R_5 are both isopropyl.

10 In a third aspect, the present invention provides novel compounds of the above of the formulae (II), (IIIa), (IIIb), (IVa), (IVb), (Va), (Vb), and (IX) as defined above and wherein R_1 is methyl or hydroxymethyl in 5-position and R_2 and R_3 are hydrogen and compounds of the formulae (IX) wherein R_1 is hydroxymethyl in 5-position, R_2 and R_3 are hydrogen and halogen is Br, I or F.

15

Detailed description of the invention

A basic concept behind the present invention is the enantioselective reduction of the compound of formula (II) to a compound of formula (IIIa) or (IIIb) in 20 enantiomerically enriched form, which is then rearranged to form the lactone (Va) or (Vb). The respective lactone enantiomers may then be reacted further to tolterodine by methods known *per se* in the art, e.g. as described in the above-mentioned US-A-5,382,600 and US-A-5,922,914.

AMENDED SHEET

2003 -05- 22

The enantioselective reduction of the compound (II) to a compound of formula (IIIa) or (IIIb) may be performed in an organic solvent with a variety of reducing agents and reaction conditions as are known *per se* in the art for enantioselective reduction of carbonyl groups. Such methods are described in, for example, Houben-Weyl, 5 Stereoselective Synthesis, Ed: Günter Helmchen et al., Vol. 7, Chapter 2.3, Thime, Stuttgart-New York 1996. Preferably, the reaction is carried out at from about 0°C to about room temperature. An exemplary method includes the use of a chiral catalyst, such as (R)- or (S)-MeCBS (3,3-diphenyl-1-methyltetrahydro-1H,3H-pyrrolo-[1,2-c][1.3.2]oxazaborole) which is commercially available, a borane complex and a 10 base. The stereochemistry can be directed by using either the R or S enantiomer of the MeCBS oxazaborolidine catalyst in the asymmetric borane reduction of the compound (II). The reduction of a similar substrate is described in, for example, WO 97/17341. The enantioselectivity of asymmetric borane reductions is not very sensitive to 15 stereoelectronic effects.

15 The sigmatropic 1,3-rearrangement (hydride shift) of the compound (IIIa) or (IIIb) to a compound of formula (IVa) or (IVb) may be carried out by treatment with a base, such as triethylamine, and a palladium catalyst, such as Pd(dppe)Cl₂ ([1,2-bis(diphenylphosphino)ethane]palladium (II) chloride) in an organic solvent (see e.g. the above WO 97/17341). Alternatively, the rearrangement reaction may be carried out 20 by treatment with DABCO (1,4-diazabicyclo[2.2.2]octane) and a base, such as triethylamine, in an organic solvent (see Example 1 below). The indanone (IVa) or (IVb) obtained is generally a highly crystalline solid which makes it possible to raise the enantiomeric purity, if desired, by recrystallization from a suitable solvent (for example, an enantiomeric excess (as defined below) of 99% or more may be obtained).

25 The Baeyer-Villiger oxidation of compounds (IVa) and (IVb) may be performed by a variety of oxidizing agents as is well known in the art, e.g. hydrogen peroxide or a peroxy acid, such as 3-chloro-peroxybenzoic acid, preferably in the presence of an acid catalyst, such as p-tolylsulphonic acid (TsOH). The reaction is preferably carried out in an organic solvent and at e.g. from about 0°C to about room temperature.

30 Enantiomeric purity, or enantiomeric enrichment, is usually expressed as "enantiomeric excess", below abbreviated as "ee", and defined as (R-S)/(R+S), where R and S are the amounts of the R- and S-enantiomers, respectively. For the purposes of the

present invention, the enantiomeric purity in the enantioselective process steps is usually at least about 50%, preferably at least about 85%.

Since tolterodine is an amine, it may form salts with both organic and inorganic acids. The pharmaceutically acceptable salts may, depending on the pharmaceutical formulation, be preferred over the corresponding free amines since they produce compounds which are more water soluble and more crystalline. Exemplary pharmaceutically acceptable salts include salts with acids such as methane sulphonic, hydrochloric, hydrobromic, sulphuric, phosphoric, nitric, benzoic, citric, tartaric, fumaric, and maleic acids.

10 The invention will now be illustrated further by the following non-limiting Example.

In the Example:

TLC refers to thin-layer chromatography.

15 MeCBS refers to 3,3-diphenyl-1-methyltetrahydro-1H,3H-pyrrolo-[1,2-c][1.3.2]oxazaborole.

DABCO refers to 1,4-diazabicyclo[2.2.2]octane.

ChiralCel OD-H (trademark) refers to a chiral stationary phase for liquid chromatography consisting of cellulose tris(3,5-dimethylphenyl carbamate) on a silica gel substrate (Daicel Chemical Industries, Ltd).

20 mCPBA refers to 3-chloroperoxybenzoic acid.

"ee" refers to enantiomeric excess as defined above.

EXAMPLE 1:

25 **1-(2-Bromo-4-methyl-phenyl)-3-phenyl-propenone**

To a solution of 2-bromo-4-methylacetophenone (7.20 g, 34.0 mmol) and benzaldehyde (3.65 g, 34.0 mmol) in dry methanol (50 ml) was added freshly prepared sodium methoxide (35.7 mmol) in dry methanol (30 ml) at 0°C. The resulting mixture was stirred at 0°C for 5 h and raised to room temperature over night. 10 ml of HCl (10%) were added slowly and the mixture was evaporated to near dryness under reduced pressure. The residue was suspended in saturated NaHCO₃ (50 ml) and extracted with 3x50 ml diethyl ether, washed with brine and dried over MgSO₄. Purification by flash chromatography eluting with diethyl ether:pentane 5:95, gave 10.1

g (95%) of the title compound. R_f 0.66 (diethyl ether: pentane 20:80). ^1H NMR δ : 2.25 (s, 3H), 6.96 (d, J = 10.2Hz, 1H), 7.15 (d, J = 10.2Hz, 1H), 7.05 (dd, J = 7.6Hz, 2.6Hz, 1H), 7.24 (m, 3H), 7.34 (m, 2H), 7.40 (m, 3H). ^{13}C NMR δ : 21.4, 112.5, 117.3, 122.5, 122.8, 123.7, 124.9, 128.4, 132.2, 133.6, 133.9, 143.6, 145.3, 186.6.

5

5-Methyl-3-phenyl-inden-1-one

To a suspension of anhydrous K_2CO_3 (9.76 g, 70.6 mmol) in dry DMF (100 ml) was added 1-(2-bromo-4-methyl-phenyl)-3-phenyl-propenone (8.40 g, 28.3 mmol), and the mixture was deaerated with dry argon for 15 min. Triphenylphosphine (0.73 g, 2.83 mmol) was added followed by PdCl_2 (0.20 g, 1.13 mmol). The mixture was heated at 80°C until NMR sample indicated disappearance of starting material (5 h). The mixture was reduced to half volume under reduced pressure and poured on ice:water (200 ml). Extractive work-up with CH_2Cl_2 followed by flash chromatography eluting with diethyl ether:pentane 5:95 gave 4.2 g (72%) of the title compound. R_f 0.62 (diethyl ether: pentane 20:80). IR (neat cm^{-1}): 1704, 1606, 1355, 1101, 815, 743. ^1H NMR δ : 2.40 (s, 3H), 5.99 (s, 1H), 7.11 (d, J = 7.2Hz, 1H), 7.18 (s, 1H), 7.43 (d, J = 7.6Hz, 1H), 7.53 (m, 3H), 7.66 (m, 2H). ^{13}C NMR δ : 22.1, 122.7, 122.9, 123.5, 127.4, 128.6, 128.9, 129.2, 129.9, 130.3, 133.2, 143.7, 144.4, 162.4. MS (EI 70 eV) m/z (rel. intensity): 220 (100) [M^+], 205 (75), 191 (51), 177 (10), 165 (15).

20

5-Methyl-3-phenyl-(S)-1H-inden-1-ol

(R)-MeCBS catalyst (0.22 ml, 1 M, 0.22 mmol) was mixed in 5 ml of dry THF, and stirred for 1 h at room temperature. After cooling to 0°C, 2.5 ml of 2 M $\text{BH}_3\text{:Me}_2\text{S}$ (4.99 mmol) in THF were added. 5-Methyl-3-phenyl-inden-1-one (1.00 g, 4.54 mmol) was added as a solution in toluene (2 ml) over 2 h via a syringe pump. The reaction was followed by TLC. After completeness, methanol (0.6 ml, 17 mmol) was added at 0°C and the mixture was evaporated to dryness. Flash chromatography eluting with ethyl acetate:pentane 10:90 gave 0.96 g (95%) of the title compound. R_f 0.35 (ethyl acetate:pentane 20:80) (ChiralCel OD-H) 0.5 ml/min of hexane/isopropanol: 95/5 (S)-isomer 24.53 min, (R)-isomer 27.22 min, 93% ee. IR (neat cm^{-1}): 3300, 1605, 1446, 949, 813. ^1H NMR δ : 1.40 (s, 1H), 2.40 (s, 3H), 5.27 (d, J = 8Hz, 1H), 6.43 (d J = 2Hz, 1H), 7.18 (d, J = 8Hz, 1H), 7.27 (s, 1H), 7.47 (m, 4H), 7.59 (m, 2H). ^{13}C NMR δ : 21.6, 76.2, 121.6, 123.6, 126.9, 127.6, 128.2, 128.6, 134.1, 134.9, 138.2, 142.1, 143.7, 145.6.

MS (EI 70eV) *m/z* (rel. intensity): 220 (100) [M⁺], 207 (71), 178 (66), 144 (42), 116 (23).

5-Methyl-3-(S)-phenyl-indan-1-one

5-Methyl-3-phenyl-(S)-1H-inden-1-ol (750 mg, 3.41 mmol) and DABCO (190 mg, 1.71 mmol) were dissolved in dry THF:triethylamine 20:1 (15 ml) and refluxed for 3 h. The reaction mixture was evaporated to dryness. Flash chromatography eluting with ethyl acetate:pentane 5:95 gave 690 mg (92%) of the title compound. *R_f* 0.62 (ethyl acetate:pentane 20:80) (ChiralCel OD-H) 0.5 ml/min of hexane/isopropanol: 95/5 (*S*)-isomer 19.12 min, (*R*)-isomer 22.33 min, 89% ee. IR (neat cm⁻¹): 3027, 2361, 1710, 1605, 1280, 1238, 1040. ¹H NMR δ: 2.39 (s, 3H), 2.69 (dd, *J* = 3.0, 19.2Hz, 1H), 3.23 (dd, *J* = 8.0, 19.2Hz, 1H), 4.53 (q, *J* = 4Hz, 1H), 7.07 (s, 1H), 7.14 (d, *J* = 8.4Hz, 1H), 7.15 (s, 1H), 7.26 (m, 2H), 7.33 (m, 2H), 7.72 (d, *J* = 7.6Hz, 1H). ¹³C NMR δ: 22.1, 44.3, 46.9, 123.2, 126.9, 127.0, 127.6, 128.9, 134.5, 143.8, 146.3, 158.4, 205.5. 15 MS (EI 70 eV) *m/z* (rel. intensity): 220 (100) [M⁺], 207 (55), 194 (19), 178 (60), 144 (10).

6-Methyl-4-(S)-phenyl-chroman-2-one

5-Methyl-3-(S)-phenyl-indan-1-one (400 mg, 1.8 mmol) and *m*CPBA (98%, 485 mg, 2.8 mmol) were suspended in dry CH₂Cl₂ (6 ml) at 0°C followed by TsOH:H₂O (20 mg). The reaction was kept at 4°C for 48 h. The mixture was diluted with 10 ml of CH₂Cl₂ and washed with 2x10 ml of saturated Na₂SO₃, saturated NaHCO₃ and brine. Flash chromatography eluting with ethyl acetate:pentane 10:90 gave 390 mg (90%) of the title compound. *R_f* 0.83 (ethyl acetate:pentane 20:80) (ChiralCel OD-H) 0.5 mL/min of hexane/isopropanol 95/5 (*S*)-isomer 15.18 min, (*R*)-isomer 17.42 min, 89% ee. IR (neat cm⁻¹): 2900, 2360, 1769, 1495, 1208, 1145. ¹H NMR δ: 2.28 (s, 3H), 3.05 (m, 1H), 4.32 (t, *J* = 6.8Hz, 1H), 6.98 (s, 1H), 7.04 (d, *J* = 8.4Hz, 1H), 7.11 (dd, *J* = 2.0, 8.4Hz, 1H), 7.18 (d, *J* = 8.4Hz, 1H), 7.19 (s, 1H), 7.33 (m, 3H) ¹³C NMR δ: 20.7, 37.1, 40.7, 116.8, 125.3, 127.5, 127.6, 128.6, 129.1, 129.3, 134.3, 140.5, 149.6, 167.8. MS (EI 70 eV) *m/z* (rel. intensity): 238 (55) [M⁺], 220 (57), 195 (100), 181(10), 165 (12), 152 (9).



**(R)-N,N-diisopropyl-3-(2-hydroxy-5-methylphenyl)-3-phenylpropanamine
(tolterodine)**

Tolterodine may be prepared from 6-methyl-4-(S)-phenyl-chroman-2-one as obtained above by method steps corresponding to Examples 3 and 4 of the above-mentioned US 5,922,914 (the full disclosure of which is incorporated by reference herein), i.e. by (i) reducing the lactone 6-methyl-4-(S)-phenyl-chroman-2-one with diisobutylaluminiumhydride in toluene solution at -20 to -25°C to the corresponding hydroxy compound, 6-methyl-4-(S)-phenyl-chroman-2-ol; (ii) reductively aminating the 6-methyl-4-(S)-phenyl-chroman-2-ol in methanol by reaction with diisopropylamine and hydrogenation with palladium on carbon at 45-50 psi and 48°C, and subsequent filtration (solka floc) to obtain the title compound (tolterodine) in substantially enantiomerically pure form.

CLAIMS

1. A process for the enantioselective preparation of a compound of the general formula (Ia) or (Ib):


5




wherein R₁, R₂ and R₃ independently of each other are hydrogen, methyl, methoxy, hydroxy, hydroxymethyl, carbamoyl, sulphamoyl or halogen, and R₄ and R₅ independently of each other are C₁-6-alkyl, or a salt thereof, which process comprises the steps of:

10 a) enantioselectively reducing the carbonyl function in a compound of formula (II):

15




wherein R₁, R₂ and R₃ are as defined above, to form an enantiomerically enriched compound of formula (IIIa) or (IIIb):



5


wherein R_1 , R_2 and R_3 are as defined above;

b) subjecting the compound of formula (IIIa) or (IIIb) to a sigmatropic rearrangement to form a corresponding enantiomerically enriched compound of formula (IVa) or (IVb):



wherein R₁, R₂ and R₃ are as defined above;

5 c) subjecting the compound of formula (IVa) or (IVb) to a Baeyer-Villiger oxidation to form a corresponding enantiomerically enriched compound of the general formula (Va) or (Vb):



wherein R₁, R₂ and R₃ are as defined above;


10 d) converting the compound of formula (Va) or (Vb) to form the corresponding enantiomerically enriched compound of formula (Ia) or (Ib); and

e) optionally converting the compound of formula (Ia) or (Ib) to a salt thereof.

15



2. The process according to claim 1, wherein step d) comprises:

d1) reacting the compound of formula (Va) or (Vb) with an amine of the general formula (VI):



20

wherein R₄ and R₅ are as defined in claim 1, to form a corresponding enantiomerically enriched compound of the general formula (VIIa) or (VIIb):



5 wherein R₁, R₂, R₃, R₄ and R₅ are as defined in claim 1; and

d2) reducing the carbonyl function in the compound of formula (VIIa) or (VIIb) to form the corresponding enantiomerically enriched compound of formula (Ia) or (Ib).

10

3. The process according to claim 2, wherein steps d1) and d2) are performed simultaneously in a single step.


15

4. The process according to claim 1, wherein step d) comprises:

d1') reducing the compound of formula (Va) or (Vb) to form a corresponding enantiomerically enriched hydroxy compound of the general formula (VIIa) or (VIIIb):

AMENDED SHEET


2003 -05- 22



wherein R₁, R₂ and R₃ are as defined in claim 1; and

5 d2') reductively aminating the hydroxy compound of formula (VIIIa) or (VIIIb) with the amine of formula (VI) to form the corresponding enantiomerically enriched compound of formula (Ia) or (Ib).

10 5. A process for the enantioselective preparation of a compound of the general formula (Va) or (Vb):




wherein R₁, R₂ and R₃ are as defined in claim 1, or a salt thereof, which process comprises the steps of:


15 a) enantioselectively reducing the carbonyl function in a compound of formula (II):

AMENDED SHEET

2003 -05- 22




wherein R₁, R₂ and R₃ are as defined above, or a salt thereof, to form an enantiomerically enriched compound of formula (IIIa) or (IIIb):




5

wherein R₁, R₂ and R₃ are as defined above, or a salt thereof;

10 b) subjecting the compound of formula (IIIa) or (IIIb) to a sigmatropic rearrangement to form a corresponding enantiomerically enriched compound of formula (IVa) or (IVb):



(IVa)




(IVb)

wherein R₁, R₂ and R₃ are as defined above, or a salt thereof; and


c) 5 subjecting the compound of formula (IVa) or (IVb) to a Baeyer-Villiger oxidation to form the corresponding enantiomerically enriched compound of the general formula (Va) or (Vb), or salt thereof.

6. 10 The process according to any one of claims 1 to 5, which further comprises preparing the compound of formula (II) by subjecting a compound of the general formula (IX):



15 wherein R₁, R₂, and R₃ are as defined in claim 1, and Hal is halogen, or a salt thereof, to a reductive ring closure reaction.

7. The process according to claim 6, further comprising preparing the compound of formula (IX) by reacting a compound of the general formula (X):




(X)

AMENDED SHEET

2003 -05- 22

wherein R₁ and Hal are as defined in claim 6, with a compound of the general formula (XI):



5 wherein R₂ and R₃ are as defined in claim 1.

8. The process according to any one of claims 1 to 7, wherein R₁ is methyl or hydroxymethyl in 5-position, R₂ and R₃ are hydrogen, and R₄ and R₅ are both iso-propyl.

10

9. The process according to any one of claims 1 to 4 and 6 to 8, wherein tolterodine is prepared.

15

10. Compounds of the formulae (II), (IIIa), (IIIb), (IVa), (IVb), (Va) and (Vb) as defined in claims 1 to 8 and wherein R₁ is methyl or hydroxymethyl in 5-position and R₂ and R₃ are hydrogen and compounds of the formulae (IX) wherein R₁ is hydroxymethyl in 5-position, R₂ and R₃ are hydrogen and halogen is Br, I or F.

20