



US 20050054046A1

(19) **United States**(12) **Patent Application Publication****Presta et al.**(10) **Pub. No.: US 2005/0054046 A1**(43) **Pub. Date: Mar. 10, 2005**(54) **NON-HUMAN PRIMATE FC RECEPTORS  
AND METHODS OF USE****Publication Classification**(75) Inventors: **Leonard G. Presta**, San Francisco, CA  
(US); **Angela K. Namenuk**, Oakland,  
CA (US)(51) **Int. Cl.<sup>7</sup>** ..... **C07H 21/04**; C07K 14/705;  
C12N 5/06(52) **U.S. Cl.** ..... **435/69.1**; 435/320.1; 435/364;  
530/350; 536/23.5

Correspondence Address:

**MERCHANT & GOULD PC****P.O. BOX 2903****MINNEAPOLIS, MN 55402-0903 (US)**(57) **ABSTRACT**(73) Assignee: **Genentech, Inc.**, South San Francisco,  
CA(21) Appl. No.: **10/896,840**(22) Filed: **Jul. 13, 2004****Related U.S. Application Data**(62) Division of application No. 10/027,736, filed on Dec.  
19, 2001.

The invention provides isolated non-human primate Fc receptor polypeptides, the nucleic acid molecules encoding the Fc receptor polypeptides, and the processes for production of recombinant forms of the Fc receptor polypeptides, including fusions, variants, and derivatives thereof. The invention also provides methods for evaluating the safety, efficacy and biological properties of Fc region containing molecules using the non-human primate Fc receptor polypeptides.


**Monomeric IgG Subclass Binding to Human FcγRI  
(Detected with HRP-anti-kappa chain)**

FIG.1A




FIG. 1B




FIG.2




FIG. 3A




FIG. 3B




FIG. 4A




FIG. 4B






FIG. 4C




FIG. 5

Hexameric Complex Binding to Cyno FcγRIIA  
Alanine Variants




FIG. 6




FIG. 7

Hexameric Complex Binding to Cyno FcγRIIIA  
Alanine Variants





FIG. 8



**FIG. 9**





FIG. 10



## NON-HUMAN PRIMATE FC RECEPTORS AND METHODS OF USE

### FIELD OF THE INVENTION

[0001] The invention generally relates to purified and isolated non-human primate Fc receptor polypeptides, the nucleic acid molecules encoding the FcR polypeptides, and the processes for production of non-human primate Fc receptor polypeptides as well as to methods for evaluating the safety, efficacy and biological properties of therapeutic agents.

### BACKGROUND OF THE INVENTION

[0002] Fc receptors (FcRs) are membrane receptors expressed on a number of immune effector cells. Upon interaction with target immunoglobulins, FcRs mediate a number of cellular responses, including, activation of cell mediated killing, induction of mediator release from the cell, uptake and destruction of antibody coated particles, and transport of immunoglobulins. Deo et al., 1997, *Immunology Today* 18:127-135. Further, it has been shown that antigen-presenting cells, e.g., macrophages and dendritic cells, undergo FcR mediated internalization of antigen-antibody complexes, allowing for antigen presentation and the consequent amplification of the immune response. As such, FcRs play a central role in development of antibody specificity and effector cell function. Deo et al., 1997, *Immunology Today* 18:127-135.

[0003] FcRs are defined by their specificity for immunoglobulin isotypes; Fc receptors for IgG antibodies are referred to as FcγR, for IgE as FcεR, for IgA as FcαR and so on. FcRn is a special class of Fc receptor found on neonatal cells and is responsible for, among other things, transporting maternal IgG from milk across the infants intestinal epithelial cells. Three subclasses of human gamma receptors have been identified: FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16). Because each human FcγR subclass is encoded by two or three genes, and alternative RNA splicing leads to multiple transcripts, a broad diversity in Fcγ isoforms exists. The three genes encoding the human FcγRI subclass (FcγRIA, FcγRIB and FcγRIC) are clustered in region 1q21.1 of the long arm of chromosome 1; the genes encoding FcγRII isoforms (FcγRIIA, FcγRIIB and FcγRIIC) and the two genes encoding FcγRIII (FcγRIIIA and FcγRIIIB) are all clustered in region 1q22. FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991); Capel et al., *Immunomethods* 4:25-34 (1994); and de Haas et al., *J Lab. Clin. Med.* 126:330-41 (1995).

[0004] Human FcγRI is a heteroligomeric complex composed of an α-chain and γ-chain. The α-chain is a 70-72 kDa glycoprotein having 3 extracellular C-2 Ig like domains, a 21 amino acid membrane domain and a charged cytoplasmic tail of 61 amino acids. van de Winkel et al., 1993, *Immunology Today* 14:215-221. The γ-chain is a homodimer that is involved in cell surface assembly and cell signaling into the interior of the cell. Each chain of γ homodimer includes a motif involved in cellular activation designated the ITAM motif. Human FcγRI binds monomeric IgG with high affinity ( $10^{-7}$ - $10^{-9}$ M) through the action of the third extracellular C-2 domain.

[0005] FcγRII is a 40 kDa glycoprotein having two C2 set Ig-like extracellular domains, a 27-29 amino acid transmem-

brane domain, and a cytoplasmic domain having variable length, from 44 to 76 amino acids. There are six known isoforms of the human FcγRII, differing for the most part in their heterogeneous cytoplasmic domains. Human FcγRIIA includes an ITAM motif in the cytoplasmic region of the molecule, and upon crosslinking of the receptor this motif is associated with cellular activation. In contrast, human FcγRIIB includes an inhibitory motif in its cytoplasmic region designated ITIM. When the FcγRIIB is crosslinked, cellular activation is inhibited. In general, FcγRII binds monomeric IgG poorly ( $>10^7$  M $^{-1}$ ), but has high affinity for complexed IgG.

[0006] Human FcγRIII has two major isoforms, FcγRIIIA and FcγRIIIB, both isoforms are between 50 to 80 kDa, having two C2 Ig-like extracellular domains. The FcγRIIIA α-chain is anchored to the membrane by a 25 amino acid transmembrane domain, while FcγRIIIB is linked to the membrane via a glycosyl phosphatidyl-inositol (GPI) anchor. Human FcγRIIIA is a heteroligomeric complex with the α-chain complexed with a heterodimeric γ-δ (gamma-delta) chain or γ-γ chain. The γ-chain includes a cytoplasmic tail with an ITAM motif. The α-chain is homologous to the α-chain and is also involved in cell signaling and cell surface assembly. The γ-δ (gamma-delta) chain also includes an ITAM motif in its cytoplasmic region. In both cases, the FcγRIII binds monomeric IgG with low affinity, and binds complexed IgG with high affinity.

[0007] Human FcRn is a heterodimer composed of a β-2 microglobulin chain and a α chain. The β-2 microglobulin chain is approximately 15 kDa and is similar to the β-2 microglobulin chain present in MHC class I heterodimers. The presence of a P-2 microglobulin chain in FcRn makes it the only known Fc receptor to fall within the MHC class I family of proteins. Ghetie et al., 1997 *Immunology Today* 18(12):592-598. The α chain is a 37-40 kDa integral membrane glycoprotein having a single glycosylation site. Evidence suggests that FcRn is involved in transferring maternal IgG across the neonatal gut and in regulating serum IgG levels. FcRn is also found in adults on many tissues.

[0008] As discussed above, human FcγRs, with the exception of FcγRIIB, contain a cytoplasmic ~26 amino acid immunoreceptor tyrosine-based activation motif (ITAM). It is believed that this motif is involved in cell signaling and effector cell function. Crosslinking of FcγRs may lead to the phosphorylation of tyrosine residues within the ITAM motif by src-family tyrosine kinases (PTKs), followed by association and activation of the phosphorylated ITAM motif with syk-family PTKs. Deo et al., 1997, *Immunology Today* 18:127-135. Once activated, a poorly understood signaling cascade is translated into biological responses.

[0009] Human FcγRIIB members contain a distinct 13 amino acid immuno-receptor tyrosine-based inhibitory motif (ITIM) in their cytoplasmic domain. Human FcγRIIB is expressed on B lymphocytes and binds to IgG complexes. However, rather than activating cells, crosslinking of the IIB receptor results in a signal inhibiting B cell activation and antibody secretion. (Camigore et al., 1992, *Cytoplasmic Domain Heterogeneity and Function of IgG Receptors in B Lymphocytes*, *Science* 256:1808.)

[0010] Because of the central role of FcγR as a trigger molecule in numerous immune responses, it has become a target for developing potential therapeutics. For example,



several ongoing clinical trials are based on activating a cancer patient's effector cells by treating the patient with tumor-specific monoclonal antibodies (Mabs). These studies have shown that the tumor-specific antibodies mediate their effects in part through Fc $\gamma$ R binding, and subsequent effector cell activity. Adams et al., 1984, *Proc. Natl. Acad. Sci.* 81:3506-3510; Takahashi et al., 1995, *Gastroenterology* 108:172-182; Riethmeuller et al., 1994, *Lancet* 343:1177-1183, Clynes, R. A., Towers, T. L., Presta, L. G., and Ravetch, J. V., 2000, *Nature Med.* 6:443-446. Further, a novel series of bispecific molecule antibodies (BSMs), molecules engineered to have one arm specific for a tumor cell and the other arm specific for a target Fc $\gamma$ R, are in clinical trials to specifically target a tumor for Fc $\gamma$ R mediated, effector cell destruction of the tumor cells. Valone et al., 1995, *J. Clin. Oncol.* 13:2281-2292; Repp et al., 1995, *Hematother* 4:415-421. In addition, Fc $\gamma$ Rs can be used as therapeutic targets in a number of infectious diseases, and for that matter, a number of autoimmune disorders. With regard to infectious diseases, BSMs are being developed to target any number of microorganisms to a patient's Fc $\gamma$ R expressing effector cells (Deo et al., 1997, *Immunology Today* 18:127-135), while soluble Fc $\gamma$ Rs have been used to inhibit the Arthus reaction, and Fc $\gamma$ R blocking agents have been used to reduce the severity of several autoimmune disorders. Ierino et al., 1993, *J. Exp. Med.* 178:1617-1628; Debre et al., 1993, *Lancet* 342:945-949.

[0011] As antibodies have become increasingly used as therapeutic agents, there is a need to develop animal models for evaluating the toxicity, efficacy and pharmacokinetics of such therapeutic agents. In addition to rodent models for evaluating efficacy of antibody therapeutics, primate models have been used for evaluation of therapeutic antibody pharmacokinetics, toxicity, and efficacy (Anderson, D. R., Grillo-Lopez, A., Varns, C., Chambers, K. S., and Hanna, N. (1997) *Biochem. Soc. Trans.* 25, 705-708). However, there is only sparse information available regarding the interaction of human antibodies with primate Fc $\gamma$  receptors and the effects of this interaction on interpretation of pharmacokinetic, toxicity, and efficacy studies in primates.

[0012] Although many advances have been made in elucidating Fc $\gamma$ R activity and identifying and engineering Fc $\gamma$ R ligands, there still remains a need in the art to identify other Fc $\gamma$ Rs and to identify and engineer other Fc $\gamma$ R ligands, both activating and inhibiting. These new receptors and receptor ligands possess potential therapeutic value in a number of disease states, including, the destruction of tumor cells and infectious material, as well as in blocking portions of the immune response involved in several autoimmune disorders. As antibodies and other Fc $\gamma$ R ligands are used as therapeutic agents, there is also a need to develop models to test the efficacy, toxicity, and pharmacokinetics of these therapeutic agents, especially in vivo.

#### SUMMARY OF INVENTION

[0013] The invention is based upon, among other things, the isolation and sequencing of polynucleotides encoding Fc receptor polypeptides from non-human primates, such as cynomolgus monkeys and chimps. The cynomolgus monkey or chimp FcR polynucleotides and polypeptides of the invention are useful, inter alia, for evaluation of binding of antibodies of any subclass (especially antibodies with prospective therapeutic utility) to cynomolgus or chimpanzee FcR polypeptides prior to in vivo evaluation in a primate.

[0014] The invention provides polynucleotide molecules encoding non-human primate Fc receptor polypeptides. The polynucleotides of the invention encode non-human primate Fc receptor polypeptides with an amino acid sequence of SEQ ID NO: 9, SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 25, SEQ ID NO: 29, SEQ ID NO: 64 or fragments thereof. Fc receptor polynucleotide molecules of the invention include those molecules having a nucleic acid sequence as shown in SEQ ID NOs: 1, 3, 5, 7, 13, 22, and 27, as well as polynucleotides having substantial nucleic acid identity with the nucleic acid sequences of SEQ ID NOs 1, 3, 5, 7, 13, 22, and 27.  $\beta$ -2 microglobulin polynucleotide molecules of the invention also include molecules having a nucleic acid sequence as shown in SEQ ID NO: 23, as well as polynucleotides having substantial nucleic acid identity with the nucleic acid sequences of SEQ ID NO: 23.

[0015] The present invention also provides non-human primate Fc receptors and non-human primate  $\beta$ -2 microglobulin. Fc polypeptides of the invention include those having an amino acid sequence shown in SEQ ID NOs: 9, 11, 15, 17, 18, 20, 29, and 64 as well as polypeptides having substantial amino acid sequence identity to the amino acid sequences of SEQ ID NOs 9, 11, 15, 17, 18, 20, 29, and 64 and useful fragments thereof.  $\beta$ -2 microglobulin polypeptides of the invention include those having an amino acid sequence shown in SEQ ID NO: 25, as well as polypeptides having substantial amino acid sequence identity to the amino acid sequence of SEQ ID NO: 25 and useful fragments thereof.

[0016] In another aspect the invention provides polynucleotide molecules encoding mature non-human primate Fc receptor polypeptides. The polynucleotides of the invention encode mature non-human primate Fc receptor polypeptides with an amino acid sequence of SEQ ID NO: 65, SEQ ID NO: 66, SEQ ID NO: 67, SEQ ID NO: 68, SEQ ID NO: 69, SEQ ID NO: 70, SEQ ID NO: 71, SEQ ID NO: 72 or fragments thereof. Fc receptor polynucleotide molecules of the invention include those molecules having a nucleic acid sequence as shown in SEQ ID NOs: 1, 3, 5, 7, 13, 22, 23 and 27, as well as polynucleotides having substantial nucleic acid identity with the nucleic acid sequences of SEQ ID NOs 1, 3, 5, 7, 13, 22, 23, and 27.

[0017] In another aspect of the invention, a method of obtaining a nucleic acid encoding a nonhuman primate Fc receptor is provided. The method comprises amplifying a nucleic acid from a nonhuman primate cell with a primer set comprising a forward and a reverse primer, wherein the primer sets are selected from the group consisting of SEQ ID NO:31 and SEQ ID NO:32, SEQ ID NO:33 and SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36, SEQ ID NO:37 and SEQ ID NO:38, SEQ ID NO:39 and SEQ ID NO:40, SEQ ID NO:41 and SEQ ID NO:42, SEQ ID NO:43 and SEQ ID NO:44, SEQ ID NO:45 and SEQ ID NO:46, SEQ ID NO:47 and SEQ ID NO:48, SEQ ID NO:49 and SEQ ID NO:50, SEQ ID NO:51 and SEQ ID NO:52, and SEQ ID NO:53 and SEQ ID NO:54; and isolating the amplified nucleic acid. The nonhuman primate cell is a preferably a cynomolgus spleen cell or a chimp spleen cell.

[0018] The invention includes variants, derivatives, and fusion proteins of the non-human primate Fc $\gamma$  receptor polypeptides and  $\beta$ -2 microglobulin. For example, the fusion

proteins of the invention include the non-human primate Fc $\gamma$  receptor polypeptides fused to heterologous protein or peptide that confers a desired function, i.e., purification, stability, or secretion. The fusion proteins of the invention can be produced, for example, from an expression construct containing a polynucleotide molecule encoding one of the polypeptides of the invention in frame with a polynucleotide molecule encoding the heterologous protein.

[0019] The invention also provides vectors, plasmids, expression systems, host cells, and the like, containing the polynucleotides of the invention. Several recombinant methods for the production of the polypeptides of the invention include expression of the polynucleotide molecules in cell free expression systems, in cellular hosts, in tissues, and in animal models, according to known methods.

[0020] The non-human primate Fc $\gamma$  receptors are useful in animal models for the evaluation of the therapeutic safety, efficacy and pharmacokinetics of agents, especially agents having a Fc region. A method of the invention involves contacting an agent with Fc receptor binding domain with a non-human primate Fc receptor polypeptide, preferably a mature soluble polypeptide, and determining the effect of contact on at least biological property of the Fc region containing molecule. A method of the invention involves contacting a cell expressing at least one non-human primate Fc $\gamma$  receptor polypeptide with an agent having a Fc region and determining whether the agent alters biological activity of the cell or is toxic to the cell. The invention also includes a method for screening variants of agents including an Fc region for the ability of such variants to bind to and activate FcRs. An example of such variants include antibodies that have amino acid substitutions at specific residues that may alter binding affinity for one or more Fc receptor classes.

[0021] Another example, of screening for agents with FcR binding domains includes identifying agents that have an altered affinity for a Fc $\gamma$  receptor having an ITAM region compared to a Fc $\gamma$  receptor having an ITIM region. In addition, the invention provides reagents, compositions, and methods that are useful identifying an agent that has an altered affinity for a Fc $\gamma$  receptor having an ITIM region, or for a method for identifying an agent with increased binding affinity for a Fc $\gamma$  receptor having an ITAM region.

[0022] These and various other features as well as advantages which characterize the invention will be apparent from a reading of the following detailed description and a review of the appended claims.

#### BRIEF DESCRIPTION OF THE FIGURES

[0023] FIG. 1A: FIG. 1A illustrates monomeric IgG subclass binding to human Fc $\gamma$ RI.

[0024] FIG. 1B: FIG. 1B illustrates monomeric IgG subclass binding to cynomolgus Fc $\gamma$ RI.

[0025] FIG. 2: FIG. 2 illustrates hexameric immune complex binding to cynomolgus Fc $\gamma$ RIIA.

[0026] FIG. 3A: FIG. 3A illustrates hexameric immune complex binding to human Fc $\gamma$ RIIB.

[0027] FIG. 3B: FIG. 3B illustrates hexameric immune complex binding to cynomolgus Fc $\gamma$ RIIB.

[0028] FIG. 4A: FIG. 4A illustrates hexameric immune complex binding to human Fc $\gamma$ RIIIA-F158.

[0029] FIG. 4B: FIG. 4B illustrates hexameric immune complex binding to human Fc $\gamma$ RIIIA-V158.

[0030] FIG. 4C: FIG. 4C illustrates hexameric immune complex binding to cynomolgus Fc $\gamma$ RIIIA.

[0031] FIG. 5: FIG. 5 illustrates hexameric immune complex binding of human IgG 1 variants to cynomolgus Fc $\gamma$ RIIA.

[0032] FIG. 6: FIG. 6 illustrates hexameric immune complex binding of human IgG variants to cynomolgus Fc $\gamma$ RIIB.

[0033] FIG. 7: FIG. 7 illustrates hexameric immune complex binding of human IgG variants to cynomolgus Fc $\gamma$ RIIIA.

[0034] FIG. 8: FIG. 8 illustrates concentration dependent monomeric IgG subclass binding to human FcRn.

[0035] FIG. 9: FIG. 9 illustrates concentration dependent monomeric IgG subclass binding to cynomolgus FcRn (S3).

[0036] FIG. 10: FIG. 10 illustrates concentration dependent monomeric IgG subclass binding to cynomolgus FcRn (N3).

#### IDENTIFICATION OF SEQUENCES AND SEQUENCE IDENTIFIERS

[0037]

SEQ ID NO.	DESCRIPTION	LOCATION	ACCESSION NO.
1	<i>Cynomolgus</i> DNA for a Fc $\gamma$ RI $\alpha$ -chain	Table 3	—
2	Human DNA for a Fc $\gamma$ RI $\alpha$ -chain	Table 3	GenBank L03418
3	<i>Cynomolgus</i> DNA for a Fc $\gamma$ RIIA	Table 5	—
4	Human DNA for a Fc $\gamma$ RIIA	Table 5	GenBank M28697
5	<i>Cynomolgus</i> DNA for a Fc $\gamma$ RIIB	Table 6	—
6	Human DNA for a Fc $\gamma$ RIIB	Table 6	GenBank X52473
7	<i>Cynomolgus</i> DNA for a Fc $\gamma$ RIIIA $\alpha$ -chain	Table 7	—
8	Human DNA for a Fc $\gamma$ RIIIA $\alpha$ -chain	Table 7	GenBank X52645
9	Amino acid sequence of a <i>cynomolgus</i> Fc $\gamma$ RI $\alpha$ -chain	Table 10	—
10	Amino acid sequence of a human Fc $\gamma$ RI $\alpha$ -chain	Table 10	GenBank P12314
11	Amino acid sequence of a <i>cynomolgus</i> Fc $\gamma$ RI/III gamma chain	Table 12	—
12	Amino acid sequence of a human Fc $\gamma$ RI/III gamma chain	Table 12	GenBank P30273

-continued

SEQ ID NO.	DESCRIPTION	LOCATION	ACCESSION NO.
13	DNA sequence for a <i>cynomolgus</i> gamma chain DNA	Table 4	—
14	DNA sequence for a human gamma chain DNA	Table 4	GenBank M33195
15	Amino acid sequence of a <i>cynomolgus</i> FcγRIIA	Table 11	—
16	Amino acid sequence of a human FcγRIIA	Table 11	GenBank P12318
17	Amino acid sequence of a chimp FcγRIIA	Table 11	—
18	Amino acid sequence of a <i>cynomolgus</i> FcγRIIB	Table 11	—
19	Amino acid sequence of a human FcγRIIB	Table 11	GenBank X52473
20	Amino acid sequence of a <i>cynomolgus</i> FcγRIIIA α-chain	Table 11	—
21	Amino acid sequence of a human FcγRIIIA α-chain	Table 11	GenBank P08637
22	DNA sequence for a chimp FcγRIIA	Table 5	—
23	<i>Cynomolgus</i> B-2 microglobulin DNA	Table 8	—
24	Human B-2 microglobulin DNA	Table 8	AB 021288
25	Amino acid sequence of <i>cynomolgus</i> B-2 microglobulin	Table 13	—
26	Amino acid sequence of human β-2 microglobulin	Table 13	P01884
27	<i>Cynomolgus</i> FcRn α -chain DNA	Table 9	—
28	Human FcRn α -chain DNA	Table 9	U12255
29	Amino acid sequence of <i>cynomolgus</i> FcRn α -chain (S3)	Table 14	—
30	Amino acid sequence of human FcRn α -chain	Table 14	U12255
31	<i>Cynomolgus</i> FcγRI full-length forward primer	Table 1	—
32	<i>Cynomolgus</i> FcγRI full-length reverse primer	Table 1	—
33	<i>Cynomolgus</i> FcγRI-H6-GST forward primer	Table 1	—
34	<i>Cynomolgus</i> FcγRI-H6-GST reverse primer	Table 1	—
35	<i>Cynomolgus</i> FcγRIIB full-length forward primer	Table 1	—
36	<i>Cynomolgus</i> FcγRIIB full-length reverse primer	Table 1	—
37	<i>Cynomolgus</i> FcγRIIB-H6-GST forward primer	Table 1	—
38	<i>Cynomolgus</i> FcγRIIB-H6-GST reverse primer	Table 1	—
39	<i>Cynomolgus</i> FcγRIIIA full-length forward primer	Table 1	—
40	<i>Cynomolgus</i> FcγRIIIA full-length reverse primer	Table 1	—
41	<i>Cynomolgus</i> FcγRIIIA-H6-GST forward primer	Table 1	—
42	<i>Cynomolgus</i> FcγRIIIA-H6-GST reverse primer	Table 1	—
43	<i>Cynomolgus</i> Fc gamma chain forward primer	Table 1	—
44	<i>Cynomolgus</i> Fc gamma chain reverse primer	Table 1	—
45	<i>Cynomolgus</i> β-2 Microglobulin forward primer	Table 1	—
46	<i>Cynomolgus</i> β-2 Microglobulin reverse primer	Table 1	—
47	<i>Cynomolgus</i> FcγRIIA full-length forward primer	Table 1	—
48	<i>Cynomolgus</i> FcγRIIA full-length reverse primer	Table 1	—
49	<i>Cynomolgus</i> FcγRIIA-H6-GST forward primer	Table 1	—
50	<i>Cynomolgus</i> FcγRIIA-H6-GST reverse primer	Table 1	—
51	<i>Cynomolgus</i> FcRn full-length forward primer	Table 1	—
52	<i>Cynomolgus</i> FcRn full-length reverse primer	Table 1	—
53	<i>Cynomolgus</i> FcRn-H6 forward primer	Table 1	—
54	<i>Cynomolgus</i> FcRn-H6 reverse primer	Table 1	—
55	PCR primer 0F1	Table 2	—
56	PCR primer 0R1	Table 2	—
57	PCR primer 0F2	Table 2	—
58	PCR primer 0F3	Table 2	—
59	PCR primer 0R2	Table 2	—
60	PCR primer 0F4	Table 2	—
61	PCR primer 0R3	Table 2	—
62	PCR primer 0F5	Table 2	—
63	PCR primer 0R4	Table 2	—
64	Amino acid sequence of <i>cynomolgus</i> FcRn α-chain (N3)	Table 14	—
65	Amino acid sequence of a mature <i>cynomolgus</i> FcγRI α-chain	Table 10	—
66	Amino acid sequence of a mature <i>cynomolgus</i> FcγRIIA	Table 11	—
67	Amino acid sequence of a mature chimp FcγRIIA	Table 11	—
68	Amino acid sequence of a mature <i>cynomolgus</i> FcγRIIB	Table 11	—
69	Amino acid sequence of a mature <i>cynomolgus</i> FcγRIIIA α-chain	Table 11	—
70	Amino acid sequence of a mature <i>cynomolgus</i> β-2 microglobulin	Table 13	—
71	Amino acid sequence of a mature <i>cynomolgus</i> FcγRn α-chain (S3)	Table 14	—
72	Amino acid sequence of a mature <i>cynomolgus</i> FcRn α-chain (N3)	Table 14	—

#### DETAILED DESCRIPTION OF THE INVENTION

[0038] The following definitions are provided to facilitate understanding of certain terms used frequently herein and are not meant to limit the scope of the present disclosure.

[0039] Throughout the present specification and claims, the numbering of the residues in an IgG heavy chain is that of the EU index as in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991),

expressly incorporated herein by reference. The "EU index as in Kabat" refers to the residue numbering of the human IgG1 EU antibody.

**[0040]** The term "amino acids" refers to any of the twenty naturally occurring amino acids as well as any modified amino acid sequences. Modifications may include natural processes such as posttranslational processing, or may include chemical modifications which are known in the art. Modifications include but are not limited to: phosphorylation, ubiquitination, acetylation, amidation, glycosylation, covalent attachment of flavin, ADP-ribosylation, cross linking, iodination, methylation, and alike.

**[0041]** The term "antibody" is used in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), chimeric antibodies, humanized antibodies, fully synthetic antibodies, and antibody fragments so long as they exhibit the desired biological activity.

**[0042]** The term "antisense" refers to polynucleotide sequences that are complementary to a target "sense" polynucleotide sequence.

**[0043]** The term "complementary" or "complementarity" refers to the ability of a polynucleotide in a polynucleotide molecule to form a base pair with another polynucleotide in a second polynucleotide molecule. For example, the sequence A-G-T is complementary to the sequence T-C-A. Complementarity may be partial, in which only some of the polynucleotides match according to base pairing, or complete, where all the polynucleotides match according to base pairing.

**[0044]** The term "expression" refers to transcription and translation occurring within a host cell. The level of expression of a DNA molecule in a host cell may be determined on the basis of either the amount of corresponding mRNA that is present within the cell or the amount of DNA molecule encoded protein produced by the host cell (Sambrook et al., 1989, *Molecular cloning: A Laboratory Manual*, 18.1-18.88).

**[0045]** The term "Fc region" is used to define a C-terminal region of an immunoglobulin heavy chain. Although the boundaries of the Fc region of an IgG heavy chain might vary slightly, the human IgG heavy chain Fc region stretches from amino acid residue at position Cys226 to the carboxyl-terminus. The term "Fc region-containing molecule" refers to an molecule, such as an antibody or immunoadhesin, which comprises an Fc region. The Fc region of an IgG comprises two constant domains, CH2 and CH3. The "CH2" domain of a human IgG Fc region (also referred to as "C $\gamma$ 2" domain) usually extends from amino acid 231 to amino acid 340. The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are interposed between the two CH2 domains of an intact native IgG molecule. Burton, *Molec. Immunol.* 22:161-206 (1985).

**[0046]** The term "Fc receptor" refers to a receptor that binds to the Fc region of an antibody or Fc region containing molecule. The preferred Fc receptor is a receptor which binds an IgG antibody (Fc $\gamma$ R) and includes receptors of the Fc $\gamma$ RI, Fc $\gamma$ RII, Fc $\gamma$ RIII, and FcRn subclasses, including allelic variants and alternatively spliced forms of these

receptors. The term "FcR polypeptide" is used to describe a polypeptide that forms a receptor that binds to the Fc region of an antibody or Fc region containing molecule. The term "Fc receptor polypeptide" also includes both the mature polypeptide and the polypeptide with the signal sequence. The term "Fc $\gamma$ R polypeptide" is used to describe a polypeptide that forms a receptor that binds to the Fc region of an IgG antibody or IgG Fc region containing molecule. For example, Fc $\gamma$ RI and Fc $\gamma$ RIII receptors each include a Fc receptor polypeptide  $\alpha$ -chain and a Fc receptor polypeptide homo or heterodimer of a  $\gamma$ -chain. FcRn receptors include an Fc receptor polypeptide alpha chain and a  $\beta$ -2 microglobulin. Typically, the  $\alpha$ -chains have the extracellular regions that bind to the Fc-region containing agent. FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991); Capel et al., *Immunomethods* 4:25-34 (1994); and de Haas et al., *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein.

**[0047]** The term "fragment" is used to describe a portion of an Fc receptor polypeptide or a nucleic acid encoding a portion of an Fc receptor polypeptide. The fragment is preferably capable of binding to a Fc region containing molecule. The structure of human Fc $\gamma$  $\alpha$ -chain of Fc $\gamma$ RI/III and Fc $\gamma$ RIIA or B has been characterized and includes a signal sequence, 2 or 3 extracellular C-2 Ig like domains; a transmembrane domain; and an intracellular cytoplasmic tail. Fragments of an Fc receptor  $\alpha$ -chain or Fc $\gamma$ RIIA or B include, but are not limited to, soluble Fc receptor polypeptides with one or more of the extracellular C-2 Ig like domains, the transmembrane domain, or intracellular domain of the Fc receptor polypeptides.

**[0048]** The term "binding domain" refers to the region of a polypeptide that binds to another molecule. In the case of an Fc receptor polypeptide or FcR, the binding domain can comprise a portion of a polypeptide chain thereof (e.g. the  $\alpha$ -chain thereof) which is responsible for binding an Fc region of an immunoglobulin or other Fc region containing molecule. One useful binding domain is the extracellular domain of an Fc receptor  $\alpha$ -chain polypeptide.

**[0049]** The term "fusion protein" is a polypeptide having two portions combined where each of the portions is a polypeptide having a different property. This property may be a biological property, such as activity *in vitro* or *in vivo*. The property may also be a simple chemical or physical property, such as binding to a target molecule, catalysis of a reaction etc. The two portions may be linked directly by a single peptide bond or through a peptide linker containing one or more amino acid residues. The fused polypeptide may be used, among other things, to determine the location of the fusion protein in a cell, enhance the stability of the fusion protein, facilitate the oligomerization of the protein, or facilitate the purification of the fusion protein. Examples of such fusion proteins include proteins expressed as fusion with a portion of an immunoglobulin molecule, proteins expressed as fusion proteins with a leucine zipper moiety, Fc receptors polypeptides fused to glutathione S-transferase, and Fc receptor polypeptides fused with one or more amino acids that serve to allow detection or purification of the receptor such as Gly6-His tag.

**[0050]** The term "homology" refers to a degree of complementarity or sequence identity between polynucleotides.

[0051] The term “host cell” or “host cells” refers to cells established in ex vivo culture. It is a characteristic of host cells discussed in the present disclosure that they be capable of expressing Fc receptors. Examples of suitable host cells useful for aspects of the present invention include, but are not limited to, insect and mammalian cells. Specific examples of such cells include SF9 insect cells (Summers and Smith, 1987, Texas Agriculture Experiment Station Bulletin, 1555), human embryonic kidney cells (293 cells), Chinese hamster ovary (CHO) cells (Puck et al., 1958, *Proc. Natl. Acad. Sci. USA* 60, 1275-1281), human cervical carcinoma cells (HELA) (ATCC CCL 2), human liver cells (Hep G2) (ATCC HB8065), human breast cancer cells (MCF-7) (ATCC HTB22), and human colon carcinoma cells (DLD-1) (ATCC CCL 221), Daudi cells (ATCC CRL-213), and the like.

[0052] The term “hybridization” refers to the pairing of complementary polynucleotides during an annealing period. The strength of hybridization between two polynucleotide molecules is impacted by the homology between the two molecules, stringency of the conditions involved, the melting temperature of the formed hybrid and the G:C ratio within the polynucleotides.

[0053] As used herein, the term “immunoadhesin” designates antibody-like molecules which combine the “binding domain” of a heterologous “adhesin” protein (e.g. a receptor, ligand or enzyme) with one or more immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of the adhesin amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site (antigen combining site) of an antibody (i.e. is “heterologous”) and an immunoglobulin constant domain sequence. The immunoglobulin constant domain sequence is preferably the Fc portion of an immunoglobulin.

[0054] “Immune complex” refers to the relatively stable structure which forms when at least one target molecule and at least one Fc region-containing polypeptide bind to one another forming a larger molecular weight complex. Examples of immune complexes are antigen-antibody aggregates and target molecule-immunoadhesin aggregates. Immune complex can be administered to a mammal, e.g. to evaluate clearance of the immune complex in the mammal or can be used to evaluate the binding properties of FcR or Fc receptor polypeptides.

[0055] The term “isolated” refers to a polynucleotide or polypeptide that has been separated or recovered from at least one contaminant of its natural environment. Contaminants of one natural environment are materials, which would interfere with using the polynucleotide or polypeptide therapeutically or in assays. Ordinarily, isolated polypeptides or polynucleotides are prepared by at least one purification step.

[0056] A “native sequence” polypeptide refers to a polypeptide having the same amino acid sequence as the corresponding polypeptide derived from nature. The term specifically encompasses naturally occurring truncated or secreted forms of the polypeptide, naturally occurring variant forms (e.g. alternatively spliced forms) and naturally occurring allelic variants. A “mature polypeptide” refers to a polypeptide that does not contain a signal peptide.

[0057] The term “nucleic acid sequence” refers to the order or sequence of deoxyribonucleotides along a strand of

deoxyribonucleic acid. The order of these deoxyribonucleotides determines the order of amino acids along a polypeptide chain. The deoxyribonucleotide sequence thus codes for the amino acid sequence.

[0058] The term “polynucleotide” refers to a linear sequence of nucleotides. The nucleotides are either a linear sequence of polyribonucleotides or polydeoxyribonucleotides, or a mixture of both. Examples of polynucleotides in the context of the present invention include—single and double stranded DNA, single and double stranded RNA, and hybrid molecules that have both mixtures of single and double stranded DNA and RNA. Further, the polynucleotides of the present invention may have one or more modified nucleotides.

[0059] The terms, “protein,” “peptide,” and “polypeptide” are used interchangeably to denote an amino acid polymer or a set of two or more interacting or bound amino acid polymers.

[0060] The term “purify,” or “purified” refers to a target protein that is free from at least 5-10% of the contaminating proteins. Purification of a protein from contaminating proteins can be accomplished through any number of well known techniques, including, ammonium sulfate or ethanol precipitation, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Various protein purification techniques are illustrated in Current Protocols in Molecular Biology, Ausubel et al., eds. (Wiley & Sons, New York, 1988, and quarterly updates).

[0061] The term “Percent (%) nucleic acid or amino acid sequence identity” describes the percentage of nucleic acid sequence or amino acid residues that are identical with amino acids in a reference polypeptide, after aligning the sequence and introducing gaps, if necessary to achieve the maximum sequence identity, and not considering any conservative substitutions as part of the sequence identity. For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

[0062] where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program’s alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. Preferably, % sequence identity can be determined by aligning the sequences manually and again multiplying 100 times the fraction X/Y, where X is the number of amino acids scored as identical matches by manual comparison and Y is the total number of amino acids in B. Further, the above described methods can also be used for purposes of determining % nucleic acid sequence identity. Alternatively, computer programs commonly employed for these purposes, such as the Gap program (Wisconsin Sequence Analysis

Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), that uses the algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.*, 2: 482-489 can be used.

[0063] Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained by manual alignment. However, the ALIGN-2 sequence comparison computer program can be used as described in WO 00/15796.

[0064] The term “stringency” refers to the conditions (temperature, ionic strength, solvents, etc) under which hybridization between polynucleotides occurs. A hybridization reaction conducted under high stringency conditions is one that will only occur between polynucleotide molecules that have a high degree of complementary base pairing (about 85% to 100% of sequence identity). Conditions for high stringency hybridization, for example, may include an overnight incubation at about 42° C. for about 2.5 hours in 6× SSC/0.1% SDS, followed by washing of the filters in 1.0× SSC at 65° C., 0.1% SDS. A hybridization reaction conducted under moderate stringency conditions is one that will occur between polynucleotide molecules that have an intermediate degree of complementary base pairing (about 50% to 84% identity).

[0065] As used herein the term “variant” means a polynucleotide or polypeptide with a sequence that differs from a native polynucleotide or polypeptide. Variants can include changes that result in amino acid substitutions, additions, and deletions in the resulting variant polypeptide when compared to a full length native sequence or a mature polypeptide sequence.

[0066] The term “vector,” “extra-chromosomal vector” or “expression vector” refers to a first piece of DNA, usually double-stranded, which may have inserted into it a second piece of DNA, for example a piece of heterologous DNA like the cDNA of cynomolgus FcγRI. Heterologous DNA is DNA that may or may not be naturally found in the host cell and includes additional copies of nucleic acid sequences naturally present in the host genome. The vector transports the heterologous DNA into a suitable host cell. Once in the host cell the vector may be capable of integrating into the host cell chromosomes. The vector may also contain the necessary elements to select cells containing the integrated DNA as well as elements to promote transcription of mRNA from the transfected DNA. Examples of vectors within the scope of the present invention include, but are not limited to, plasmids, bacteriophages, cosmids, retroviruses, and artificial chromosomes.

#### Modes of Carrying Out the Invention

[0067] The invention is based upon, among other things, the isolation and sequencing of nucleic acids encoding Fc receptor polypeptides from non-human primates, such as cynomolgus monkeys and chimps. In particular, the invention provides isolated polynucleotides encoding FcR polypeptides with an amino acid sequence of SEQ ID NO: 9, 11, 15, 17, 18, 20, 29, 64 or fragments thereof. The invention also provides isolated polynucleotides encoding mature FcR polypeptides with an amino acid sequence of SEQ ID NO: 65, 66, 67, 68, 69, 71 or 72, or fragments thereof. The invention also provides an isolated polynucle-

otide encoding β-2 microglobulin having an amino acid sequence of SEQ ID NO: 25 or SEQ ID NO: 70.

[0068] The cynomolgus monkey or chimp Fc receptor polynucleotides and polypeptides of the invention are useful for evaluation of binding of antibodies of any subclass (especially antibodies with prospective therapeutic utility) to cynomolgus or chimpanzee FcR polypeptides prior to in vivo evaluation in a primate. Evaluation could include testing binding to primate FcRs or Fc receptor polypeptides in an ELISA-format assay or to transiently- or stably-transfected human or primate cells (e.g. CHO, COS). Evaluation of the ability of a human antibody to bind to cynomolgus or other primate FcRs or Fc receptor polypeptides (either in an ELISA- or transfected cell format) could be used as a preliminary test prior to evaluation of pharmacokinetics/pharmacodynamics in vivo. Binding of antibodies or antibody variants to cynomolgus FcRn or FcRn polypeptides would be useful to identify antibodies or antibody variants that could have a longer half life in vivo. Binding of antibodies to FcRn correlates with a longer half life in vivo.

[0069] The primate FcRs or Fc receptor polypeptides could also be used to screen for variants (e.g. protein-sequence or carbohydrate) of primate or human IgG which exhibit either improved or reduced binding to these receptors or receptor polypeptides; such variants could then be evaluated in vivo in a primate model for altered efficacy of the antibody, e.g. augmentation or abrogation of IgG effector functions. In addition, soluble cynomolgus or chimpanzee Fc receptor polypeptides could be evaluated as therapeutics in primate models.

[0070] For example, in one aspect of the invention, a method is provided for identifying agents that selectively activate ITAM motifs in target Fc receptors while failing to activate ITIM motifs in other Fc receptors. Preferably these agents are antibodies and more preferably these agents are monoclonal antibodies. These identified agents may have uses in designing therapeutic antibodies which preferentially bind to and activate only ITAM-containing FcγR (i.e. not simultaneously engaging the inhibitory ITIM-containing receptors) which could thereby improve the cytotoxicity or phagocytosis ability of the therapeutic antibody or the ability of the therapeutic antibody to be internalized by antigen-presenting cells for increased immune system response against the target antigen.

[0071] Finally, the cynomolgus FcγR polynucleotides and polypeptides of the invention permit a more detailed analysis of FcγR-mediated molecular interactions. The amino acids in human IgG1 which interact with human FcγR have been mapped (Shields, R. L., Namenuk, A. K., Hong, K., Meng, Y. G., Rae, J., Briggs, J., Xie, D., Lai, J., Stadlen, A., Li, B., Fox, J. A., and Presta, L. G. (2001) *J. Biol. Chem.* 276, 6591-6604). Testing the binding of these same human IgG1 variants against cynomolgus FcγR can aid in mapping the interaction of specific amino acids in the human IgG1 with amino acids in the FcγR.

[0072] Within the application, unless otherwise stated, the techniques utilized may be found in any of several well-known references, such as: *Molecular Cloning: A Laboratory Manual* (Sambrook et al. (1989) *Molecular cloning: A Laboratory Manual*), *Gene Expression Technology* (Methods in Enzymology, Vol. 185, edited by D. Goeddel, 1991

Academic Press, San Diego, Calif.), "Guide to Protein Purification" in *Methods in Enzymology* (M. P. Deutscher, 3d., (1990) Academic Press, Inc.), *PCR Protocols: A Guide to Methods and Applications* (Innis et al. (1990) Academic Press, San Diego, Calif.), *Culture of Animal Cells: A Manual of Basic Technique*, 2<sup>nd</sup> ed. (R. I. Freshney (1987) Liss, Inc., New York, N.Y.), and *Gene Transfer and Expression Protocols*, pp 109-128, ed. E. J. Murray, The Humana Press Inc., Clifton, N.J.).

#### Polynucleotide Sequences

**[0073]** One aspect of the invention provides isolated nucleic acid molecules encoding Fc receptor polypeptides from cynomolgus monkeys and chimps. Due to the degeneracy of the genetic code, two DNA sequences may differ and yet encode identical amino acid sequences. The present invention thus provides isolated nucleic acid molecules comprising a polynucleotide sequence encoding cynomolgus FcR polypeptides, wherein the polynucleotide sequences encode a polypeptide with an amino acid sequence of SEQ ID NO: 9, or SEQ ID NO: 11, or SEQ ID NO: 15, or SEQ ID NO: 18, or SEQ ID NO: 20, or SEQ ID NO: 29, or SEQ ID NO: 64, or fragments thereof. The present invention also provides isolated nucleic acid molecules comprising a polynucleotide sequence encoding a chimp FcR polypeptide of the invention, wherein the polynucleotide sequence encodes a polypeptide with an amino acid sequence of SEQ ID NO: 17 or fragments thereof. The invention also provides for isolated nucleic acid molecules comprising a polynucleotide sequence encoding cynomolgus  $\beta$ -2 microglobulin with an amino acid sequence of SEQ ID NO: 25.

**[0074]** The present invention also provides isolated nucleic acid molecules comprising a polynucleotide sequence encoding mature nonprimate FcR polypeptides, wherein the polynucleotide sequences encode a polypeptide with an amino acid sequence of SEQ ID NO: 65, 66, 68, 67, 69, 70, 71, or 72.

**[0075]** The nucleotide sequences shown in the tables, in most instances, begin at the coding sequence for the signal sequence of the Fc receptor polypeptide.

**[0076]** Nucleotide sequences of the non-human primate receptors have been aligned with human sequences for FcR polypeptides or  $\beta$ -2 microglobulin to determine % sequence identity. Nucleotide sequences of primate and human proteins are aligned manually and differences in nucleotide or protein sequence noted. Percent identity is calculated as number of identical residues/number of total residues. When the sequences differ in the total number of residues, two values for percent identity are provided, using the two different numbers for total residues. Some nucleic acid sequences for human FcR are known to those of skill in the art and are identified by GenBank accession numbers.

**[0077]** In one embodiment, the invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a cynomolgus FcRI  $\alpha$ -chain. One example of a cynomolgus FcRI  $\alpha$ -chain has an amino acid sequence including the signal sequence as shown in Table 10 (SEQ. ID. NO: 9). The mature cynomolgus FcRI  $\alpha$ -chain has an amino acid sequence shown in Table 10 (SEQ ID NO: 65). An example of an isolated nucleic acid encoding a cynomolgus FcRI  $\alpha$ -chain is shown in Table 3 (SEQ ID NO: 1). A nucleic acid

sequence encoding a cynomolgus FcRI  $\alpha$ -chain has about 91% or 96% sequence identity when aligned with a human nucleic acid sequence (SEQ ID NO: 2) encoding a FcRI  $\alpha$ -chain as shown in Table 3 (GenBank Accession No. L03418).

**[0078]** In another embodiment, the invention provides an isolated nucleic acid comprising a polynucleotide sequence encoding a cynomolgus gamma chain of FcRI/III. An example of such a nucleic acid sequence is shown in Table 4 (SEQ ID NO: 13). An example of a cynomolgus gamma chain polypeptide is shown in Table 12 (SEQ ID NO: 11). A nucleic acid encoding a cynomolgus gamma chain has about 99% sequence identity when aligned with a human nucleic acid sequence (SEQ ID NO: 14) encoding a FcR gamma chain as shown in Table 4 (GenBank Accession No. M33195).

**[0079]** In another embodiment, the invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a cynomolgus FcRIIA. One example of cynomolgus FcRIIA has an amino acid sequence including the signal sequence as shown in Table 11 (SEQ. ID. NO: 15). The mature cynomolgus FcRIIA has an amino acid sequence as shown in Table 21 (SEQ ID NO: 66). An example of an isolated nucleic acid encoding a cynomolgus FcRIIA is shown in Table 5 (SEQ ID NO: 3). A nucleic acid sequence encoding a cynomolgus FcRIIA  $\alpha$ -chain has about 94% sequence identity when aligned with a human nucleic acid sequence (SEQ ID NO: 4) encoding a FcRIIA as shown in Table 5 (Genbank Accession No. M28697).

**[0080]** The invention also provides for isolated nucleic acids comprising a polynucleotide encoding FcR from chimps such as an isolated nucleic acid comprising a polynucleotide encoding a FcRIIA receptor. One example of a chimp FcRIIA has an amino acid sequence including the signal sequence as shown in Table 11 (SEQ. ID. NO: 17). The mature chimp FcRIIA has an amino acid sequence as shown in Table 11 (SEQ ID NO: 67). An example of an isolated nucleic acid encoding a chimp FcRIIA is shown in Table 5 (SEQ ID NO: 22). A nucleic acid sequence having a sequence of SEQ ID NO: 22 has about 99% sequence identity when aligned with a human nucleic acid sequence (SEQ ID NO: 4) encoding a FcRIIA as shown in Table 5 (GenBank Accession No. M28697).

**[0081]** In another embodiment, the invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a cynomolgus FcRIIB. One example of a cynomolgus FcRIIB has an amino acid sequence as shown in Table 11 (SEQ. ID. NO: 18). The mature cynomolgus FcRIIB has an amino acid sequence as shown in Table 22 (SEQ ID NO: 68). An example of an isolated nucleic acid encoding a cynomolgus FcRIIB is shown in Table 6 (SEQ ID NO: 5). A nucleic acid sequence encoding a cynomolgus FcRIIB has about 94% sequence identity when aligned with a human nucleic acid sequence (SEQ ID NO: 6) encoding a FcRIIB as shown in Table 6 (GenBank Accession No. X52473).

**[0082]** In another embodiment, the invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a cynomolgus FcRIIIA  $\alpha$ -chain. One example of a cynomolgus FcRIIIA has an amino acid sequence as shown in Table 11 (SEQ. ID. NO: 20). The mature cynomolgus FcRIIIA has an amino acid sequence as shown in

Table 23 (SEQ ID NO: 69). An example of an isolated nucleic acid encoding a cynomolgus FcγRIIIA α-chain is shown in Table 7 (SEQ ID NO: 7). A nucleic acid sequence cynomolgus FcγRIIIA α-chain has about 96% sequence identity when aligned with a human nucleic acid sequence (SEQ ID NO: 8) encoding a FcγRIIIA α-chain as shown in Table 7 (GenBank Accession No. X52645).

**[0083]** The invention also provides isolated nucleic acid molecules having a polynucleotide sequence encoding a cynomolgus Fc receptor (FcRn) α-chain. One example of a cynomolgus Fc receptor α-chain (S3) has an amino acid sequence of SEQ ID NO: 29 as shown in Table 14. An allele has been identified encoding a polypeptide with an amino acid sequence which differs from that of SEQ ID NO: 29 by a substitution of an asparagine for a serine at the third residue in the mature polypeptide. This polypeptide sequence has been designated SEQ ID NO: 64. The mature polypeptides of FcRn α-chain (S3) and FcRn α-chain (N3) have the amino acid sequences of SEQ ID NO: 71 and 72, respectively. An example of an isolated nucleic acid encoding a cynomolgus FcRn α-chain is SEQ ID NO: 27 shown in Table 9. A nucleic acid encoding a cynomolgus FcRn has about 97% sequence identity when aligned with a human sequence (SEQ ID NO: 28) encoding a human FcRn α-chain as shown in Table 9 (GenBank Accession No. U12255).

**[0084]** In another embodiment, the invention provides isolated nucleic acid molecules comprising a polynucleotide sequence encoding cynomolgus β-2 microglobulin. One example of a cynomolgus β-2 microglobulin has an amino acid sequence as shown in Table 13 (SEQ ID NO: 25). The mature P-2 microglobulin has a sequence as shown in Table 13 (SEQ ID NO: 70). An example of an isolated nucleic acid encoding a cynomolgus β-2 microglobulin is shown in Table 8 (SEQ ID NO: 23). A nucleic acid cynomolgus β-2 microglobulin has about 95% sequence identity when aligned with a human sequence (SEQ ID NO: 24) encoding β-2 microglobulin as shown in Table 8 (GenBank Accession No. AB021288).

**[0085]** The non-human primate nucleic acids of the invention include cDNA, chemically synthesized DNA, DNA isolated by PCR, and combinations thereof. RNA transcribed from cynomolgus or chimp cDNA is also encompassed by the invention. The cynomolgus DNA can be obtained using standard methods from tissues such as the spleen or liver and as described in the Examples below. The chimp FcγR DNA can be obtained using standard methods from tissues such as spleen or liver and as described in the Examples below.

**[0086]** In another aspect of the invention, a method of obtaining a nucleic acid encoding a nonhuman primate Fc receptor is provided. The method comprises amplifying a nucleic acid from a nonhuman primate cell with a primer set comprising a forward and a reverse primer, wherein the primer sets are selected from the group consisting of SEQ ID NO:31 and SEQ ID NO:32, SEQ ID NO:33 and SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36, SEQ ID NO:37 and SEQ ID NO:38, SEQ ID NO:39 and SEQ ID NO:40, SEQ ID NO:41 and SEQ ID NO:42, SEQ ID NO:43 and SEQ ID NO:44, SEQ ID NO:45 and SEQ ID NO:46, SEQ ID NO:47 and SEQ ID NO:48, SEQ ID NO:49 and SEQ ID NO:50, SEQ ID NO:51 and SEQ ID NO:52, and SEQ ID NO:53 and SEQ ID NO:54; and isolating the amplified

nucleic acid. The nonhuman primate cell is a preferably a cynomolgus spleen cell or a chimp spleen cell. Some of the primer sets provide for amplification of an extracellular fragment of the Fc receptor polypeptides fused to GlyHis-GST.

**[0087]** Fragments of the cynomolgus and chimp FcγR-encoding nucleic acid molecules described herein, as well as polynucleotides capable of hybridizing to such nucleic acid molecules, may be used in a number of ways including as a probe or as primers in a polymerase chain reaction (PCR). Such probes may be used, e.g., to detect the presence of FcγR polynucleotides in *in vitro* assays, as well as in Southern and Northern blots. Cell types expressing the FcγR may also be identified by the use of such probes. Such procedures are well known, and the skilled artisan will be able to choose a probe of a length suitable to the particular application. For PCR, 5' and 3' primers corresponding to the termini of the nucleic acid molecules are employed to isolate and amplify that sequence using conventional techniques. Fragments useful as probes are typically oligonucleotides about 18 to 20 nucleotides, including up to the full length of the polynucleotides encoding the FcγR. Fragments useful as PCR primers typically are oligonucleotides of 20 to 50 nucleotides.

**[0088]** Other useful fragments of the different cynomolgus FcγR polynucleotides are antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence capable of binding to a target FcγR mRNA (using a sense strand), or DNA (using an antisense strand) sequence.

**[0089]** Other useful fragments include polynucleotides that encode domains of a Fcγ receptor polypeptide. The fragments are preferably capable of binding to a Fc region containing molecule. One embodiment of a polynucleotide fragment is a fragment that encodes extracellular domains of a Fcγ receptor polypeptide in which the transmembrane and cytoplasmic domains have been deleted. Other domains of Fcγ receptors are identified in, for example, Table 10 and Table 11. Nucleic acid fragments encoding one or more polypeptide domains are included within the scope of the invention.

**[0090]** The invention also provides variant cynomolgus and chimp FcγR nucleic acid molecules as well as variant cynomolgus β-2 microglobulin nucleic acid molecules. Variant polynucleotides can include changes to the nucleic acid sequence that result in amino acid substitutions, additions, and deletions in the resultant variant polypeptide when compared to a native polypeptide, for instance SEQ ID NOs: 9, 11, 15, 17, 18, 20, 25, 29, or 64. The changes to the variant nucleic acid sequences can include changes to the nucleic acid sequence that result in replacement of an amino acid by a residue having similar physiochemical properties, such as substituting one aliphatic residue (Ile, Val, Leu, or Ala) for another, or substitutions between basic residues Lys and Arg, acidic residues Glu and Asp, amide residues Gln and Asn, hydroxyl residues Ser and Tyr, or aromatic residues Phe and Tyr. Variant polynucleotide sequences of the present invention are preferably at least about 95% identical, more preferably at least about 96% identical, more preferably at least about 97% or 98% identical, and most preferably at least about 99% identical, to a nucleic acid sequence encoding the full length native sequence, a polypeptide lacking a signal sequence, an extracellular domain of the polypeptide,



or a nucleic acid encoding a fragment of the Fc $\gamma$  receptor polypeptide or  $\beta$ -2 microglobulin of sequences of SEQ ID NOs: 1, 3, 5, 7, 23 or 27.

[0091] The percentage of sequence identity between the sequences and a variant sequence as discussed above may also be determined, for example, by comparing the variant sequence with a reference sequence using any of the computer programs commonly employed for this purpose, such as ALIGN 2 or by using manual alignment. Percent identity is calculated as [number of identical residues]/[number of total residues]. When the sequences differed in the total number of residues, two values for percent identity are provided, using the two different numbers for total residues.

[0092] Alterations of the cynomolgus monkey and chimp Fc $\gamma$ R polypeptides, and cynomolgus monkey  $\beta$ -2 microglobulin, nucleic acid and amino acid sequences may be accomplished by any of a number of known techniques. For example, mutations may be introduced at particular locations by procedures well known to the skilled artisan, such as oligonucleotide-directed mutagenesis, which is described by Walder et al., 1986, *Gene*, 42:133; Bauer et al., 1985, *Gene* 37:73; Craik, 1985, *BioTechniques*, 12:19; Smith et al., 1981, *Genetic Engineering: Principles and Methods*, Plenum Press; and U.S. Pat. No. 4,518,584 and U.S. Pat. No. 4,737,462.

[0093] The invention also provides cynomolgus and chimp Fc $\gamma$ R polypeptides, cynomolgus FcRn polypeptide,  $\beta$ -2 microglobulin nucleic acid molecules, or fragments and variants thereof, ligated to heterologous polynucleotides to encode fusion proteins. The heterologous polynucleotides can be ligated to the 3' or 5' end of the nucleic acid molecules of the invention, for example SEQ ID NOs: 1, 3, 5, 7, 13, 22, 25 or 27, to avoid interfering with the in-frame expression of the resultant cynomolgus and chimp Fc $\gamma$ R, cynomolgus FcRn, and  $\beta$ -2 microglobulin polypeptides. Alternatively, the heterologous polynucleotide can be ligated within the coding region of the nucleic acid molecule of the invention. Heterologous polynucleotides can encode a single amino acid, peptide, or polypeptides that provide for secretion, improved stability, or facilitate purification of the cynomolgus and chimp encoded polypeptides of the invention.

[0094] A preferred embodiment is a nucleic acid sequence encoding an extracellular domain of the  $\alpha$ -chain of Fc $\gamma$ RI, Fc $\gamma$ R or FcRn fused to Gly(His)<sub>6</sub>-gst tag or Fc $\gamma$ RIIA or IIB fused to Gly(His)<sub>6</sub>-gst tag obtained as described in Example 1. The Gly(His)<sub>6</sub>-gst tag provides for ease of purification of polypeptides encoded by the nucleic acid.

[0095] The cynomolgus and chimp Fc $\gamma$ R polypeptide and  $\beta$ -2 microglobulin nucleic acid molecules of the invention can be cloned into prokaryotic or eukaryotic host cells to express the resultant polypeptides of the invention. Any recombinant DNA or RNA method can be used to create the host cell that expresses the target polypeptides of the invention, including, but not limited to, transfection, transformation or transduction. Methods and vectors for genetically engineering host cells with the polynucleotides of the present invention, including fragments and variants thereof, are well known in the art, and can be found in *Current Protocols in Molecular Biology*, Ausubel et al., eds. (Wiley & Sons, New York, 1988, and updates). Vectors and host cells for use with the present invention are described in the Examples provided herein.

[0096] The invention also provides isolated nucleic acids comprising a polynucleotide encoding the mature Fc receptor polypeptide. The isolated nucleic acids can further comprise a nucleic acid sequence encoding a heterologous signal sequence. A heterologous signal sequence is one obtained from a polynucleotide encoding a polypeptide different than the native sequence non-human primate Fc receptor polypeptides of the invention. Heterologous signal sequences include signal sequences from human Fc receptor polypeptides as well as from polypeptides like tissue plasminogen activator.

#### Polypeptide Sequences

[0097] Another aspect of the invention is directed to FcR polypeptides from non-human primates such as cynomolgus monkeys and chimps. The Fc $\gamma$ R polypeptides include Fc $\gamma$ RI  $\alpha$ -chain, Fc $\gamma$ RIIA, Fc $\gamma$ RIIB, Fc $\gamma$ RIIA  $\alpha$ -chain, FcRn  $\alpha$ -chain, Fc $\gamma$ I/III  $\gamma$ -chain, and  $\beta$ -2 microglobulin. The polypeptides bind IgG antibody or other molecules having a Fc region. Some of the receptors are low affinity receptors which preferably bind to IgG antibody complexes. FcR polypeptides also mediate effector cell functions such as antibody dependent cellular cytotoxicity, induction of mediator release from the cell, uptake and destruction of antibody coated particles, and transport of immunoglobulins.

[0098] Amino acid sequences of the Fc $\gamma$ R polypeptides derived from cynomolgus monkeys and chimps are aligned with the amino acid sequences encoding human Fc $\gamma$ R polypeptides to determine the % of sequence identity with the human sequences. Amino acid sequences of primate and human proteins are aligned manually and differences in nucleotide or protein sequence noted. Percent identity is calculated as number of identical residues/number of total residues. When the sequences differ in the total number of residues, two values for percent identity are provided, using the two different numbers for total residues. Some amino acid sequences encoding human Fc $\gamma$ R polypeptides are known to those skilled in the art and are identified by GenBank Accession numbers.

[0099] The polypeptide sequences shown in the tables are numbered starting from the signal sequence or from the first amino acid of the mature protein. When the amino acid residues of the polypeptide are numbered starting from the signal sequence the numbers are identified by the number of the residue and a line. When the amino acid residues of the polypeptide are also numbered from the first amino acid of the mature human protein, the amino acid is designated by the number and A symbol. In Table 11, the first N terminal residue of the cynomolgus sequences is designated with an asterisk, but the numbering is still that corresponding to the mature human protein. The numbering of the amino acid residues of the FcR polypeptides is sequential.

[0100] The non-human primate receptors were also analyzed to compare the binding of the non-human primate Fc receptor polypeptides to various subclasses of human IgG and IgG variants to human Fc receptors. The binding to the subclasses also included binding to IgG4b. IgG4b is a form of IgG4, but has a change in the hinge region at amino acid residue 228 from serine to a proline. This change results in a molecule that is more stable than the native IgG4 due to increase formation of interchain disulfide bonds as described

in Angal, S., King, D. J., Bodmer, M. W., Turner, A., Lawson, D. G., Robert, G., Pedley B. and Adair, J. R (1993) A single amino acid substitution abolishes heterogeneity of chimeric—mouse/human (IgG4) antibody. *Molec. Immunol.* 30:105-108.

**[0101]** One embodiment of the invention is a cynomolgus FcγRI polypeptide. A cynomolgus FcγRI binds to IgG and other molecules having an Fc region, preferably human monomeric IgG. One example of an α-chain of a cynomolgus FcγRI is a polypeptide having a sequence of SEQ ID NO: 9. Based on the alignment with the human sequence, the mature cynomolgus FcγRI has a sequence of SEQ ID NO: 65. An extracellular fragment obtained as described in example 1 has an amino acid sequence of Δ1 to Δ269 as shown in table 10.

**[0102]** An alignment of the amino acid sequence α-chain of the FcγRI from human and cynomolgus monkeys is also shown in Table 10. The amino acid numbers shown below the amino acids with the symbol Δ are numbered from the start of the mature polypeptide not including the signal sequence. The numbers above the amino acid residues represent the numbering of the residues starting at the signal sequence. Each of the domains of the FcγRI α-chain are shown including signal sequence, extracellular domain 1, extracellular domain 2, extracellular domain 3, and the transmembrane and intracellular sequence. The alignment of a human sequence of SEQ ID NO: 10 (GenBank Accession No. P12314) with a cynomolgus FcγRI α-chain sequence starting from the signal sequence shows about a 90% or 94% sequence identity with the human sequence depending on whether the 3' extension present on the human sequence was used in the calculation.

**[0103]** This alignment of the cynomolgus sequence with the human sequence shows that the cynomolgus FcγRI α-chain has the same number of amino acids in the signal sequence, the three extracellular domains, and transmembrane domain as found in the human FcγRI sequence (Table 10). In contrast, the cynomolgus FcγRI α-chain intracellular domain is shorter than that of the human FcγRI α-chain by seventeen amino acids (Table 10). A cynomolgus FcγRI α-chain binds to human monomeric subclasses as follows: IgG3≧IgG1>IgG4b>>>IgG2, which is similar to that of the human FcγRI.

**[0104]** Fc receptors of the I and IIIA subclass are complex molecules including an α-chain complexed to either a homo or hetero dimer of a γ-chain. The invention also includes a cynomolgus FcR gamma chain. One example of a gamma chain polypeptide has an amino acid sequence of SEQ ID NO: 11 as shown in Table 12. When the cynomolgus gamma chain amino acid sequence is aligned with a human sequence for the gamma chain of SEQ ID NO: 12 (GenBank Accession No. P30273) it has about 99% sequence identity with the human sequence. The ITAM motif of the cynomolgus gamma chain is identical to that of the human gamma chain.

**[0105]** Another embodiment of the invention is a cynomolgus FcγRIIA. A cynomolgus FcγRIIA binds to immunoglobulins and other molecules having an Fc region, preferably immunoglobulins complexed to an antigen or each other. More preferably, the receptor binds a dimeric or hexameric immune complex of human Ig. One example of a cynomolgus FcγRIIA has an amino acid sequence of SEQ

ID NO: 15. The mature cynomolgus FcγRIIA has an amino acid sequence of SEQ ID NO: 66 (Table 21). an extracellular fragment obtained with the primers of example 1 has an amino acid sequence of Δ1 to Δ182 as shown in Table 21.

**[0106]** The cynomolgus FcγRIIA sequence was aligned with a human amino acid sequence of FcγRIIA as shown in Table 11 (SEQ ID NO: 16) (Accession No. P12318). In table 11, the amino acid numbers shown below the amino acids with the symbol A are numbered from the start of the mature human polypeptide not including the signal sequence. The numbers above the amino acid residues represent the numbering of the residues starting at the signal sequence. When the cynomolgus sequence is aligned with the human sequence it has about 87% or 89% sequence identity with the human sequence depending on whether the alignment starts with the MAMETQ sequence. This alignment shows that the cynomolgus FcγRIIA has fewer amino acids in the signal peptide sequence than found in the human FcγRIIA (Table 11). Cynomolgus FcγRIIA has about the same number of amino acids in the two extracellular domains, transmembrane domain, and intracellular domain as found in the human FcγRIIA sequence (Table 11). Notably, the cynomolgus FcγRIIA contains the identical two ITAM motifs as found in the human receptor (Table 11).

**[0107]** The cynomolgus FcγRIIA binds to hexameric complexes of subclasses IgG with the following binding pattern: IgG3=IgG2>IgG1>IgG4b, IgG4. A human FcγRIIA isoform with an arginine at the amino acid corresponding to the amino acid 131 (R131) binds hexameric IgG subclasses as follows: IgG3>IgG1>>>IgG2>IgG4. A human FcγRIIA isoform with a histidine at the amino acid corresponding to the amino acid 131 (H131) binds hexameric IgG subclasses as follows: IgG3≧IgG1=IgG2>>>IgG4. Cynomolgus FcγRIIA with an amino acid sequence of SEQ ID NO: 15 has H 131 and binds to human subclasses of IgG in a similar manner to those human Fc receptors with the H131 isoform variant. However, the cynomolgus Fc receptor binds IgG2 as efficiently as it binds IgG3.

**[0108]** Another embodiment of the invention is a chimp FcγRIIA. A chimp FcγRIIA binds to immunoglobulins and other molecules having an Fc region, preferably immunoglobulins complexed to an antigen or each other. Preferably the receptor binds a dimeric or hexameric immune complex of human Ig. One example of a chimp FcγRIIA has an amino acid sequence of SEQ ID NO: 17. Based on the alignment with the human sequence, the mature chimp FcγRIIA has an amino acid sequence of SEQ ID NO: 67.

**[0109]** The chimp FcγRIIA amino acid sequence was aligned starting with the signal sequence with a human sequence for FcγRIIA of SEQ ID NO: 16 as shown in Table 11 (Accession No. P12318). The alignment shows that when compared to the human sequence, the chimp sequence has about 97% sequence identity. This alignment also shows that the chimpanzee FcγRIIA has one less amino acid in the signal peptide sequence than found in the human FcγRIIA α-chain (Table 11). Chimpanzee FcγRIIA has the same number of amino acids in the two extracellular domains, transmembrane domain, and intracellular domain as found in the human FcγRIIA sequence (Table 11). Notably, the chimpanzee FcγRIIA contains the identical two ITAM motifs as found in the human and cynomolgus receptors (Table 11).

[0110] Another embodiment of the invention is a cynomolgus FcγRIIB. A cynomolgus FcγRIIB binds to immunoglobulins and other molecules having an Fc region, preferably immunoglobulins complexed to an antigen or each other. More preferably, the receptor binds a dimeric or hexameric immune complex of human Ig. One example of a cynomolgus FcγRIIB has an amino acid sequence of SEQ ID NO: 18. The mature cynomolgus FcγRIIB has an amino acid sequence of SEQ ID NO: 68 (Table 22). an extracellular fragment obtained with the primers of example 1 has an amino acid sequence of Δ1 to Δ184 as shown in table 22.

[0111] The cynomolgus FcγRIIB has about 92% sequence identity with a human amino acid sequence of FcγRIIB as shown in Table 11 (SEQ ID NO: 19) (Accession No. X52473). An alignment of the cynomolgus sequence with the human sequence shows that the cynomolgus FcγRIIB has about the same number of amino acids in the signal peptide, two extracellular domains, and transmembrane domain as found in the human FcγRIIB sequence (Table 11). The cynomolgus FcγRIIB has three amino acids inserted in the N-terminal portion of the intracellular domain (compared to human FcγRIIB) (Table 11). Notably, the cynomolgus FcγRIIB intracellular domain contains the identical ITIM motif as found in the human receptor (Table 11).

[0112] The cynomolgus FcγRIIB binds to hexameric complexes of subclasses IgG with the following binding pattern: IgG2≧IgG3>IgG1>IgG4b, IgG4. A human FcγRIIB binds hexameric IgG subclasses as follows: IgG3≧IgG1>IgG2>IgG4. The cynomolgus FcγRIIB binds IgG2 much more efficiently than the human FcγRIIB.

[0113] Another embodiment of the invention is a cynomolgus FcγRIIA. A cynomolgus receptor FcγRIIA binds to immunoglobulins and other molecules having an Fc region, preferably immunoglobulins complexed. Preferably, the receptor binds a dimeric or hexameric immune complex of human Ig. One example of an amino acid sequence of the α-chain of FcγRIIA is SEQ ID NO: 20. The mature cynomolgus FcγRIIA α-chain has a sequence of SEQ ID NO: 69 (Table 23). An extracellular fragment obtained using the primer as described in example 1 has an amino acid sequence of Δ1 to Δ187 as shown in Table 23.

[0114] The cynomolgus FcγRIIA α-chain sequence was aligned with a human amino acid sequence of FcγRIIA as shown in Table 11 (SEQ ID NO: 21) (Accession No. P08637). In table 11, the amino acid numbers shown below the amino acids with the symbol A are numbered from the start of the mature human polypeptide not including the signal sequence. The numbers above the amino acid residues represent the numbering of the residues starting at the signal sequence. The alignment with the human and cynomolgus FcγRIIA sequence shows the sequence has about 91% sequence identity to the human sequence. This alignment of the cynomolgus sequence with the human sequence shows that the cynomolgus FcγRIIA α-chain has about the same number of amino acids in the signal peptide, the two extracellular domains, the transmembrane domain, and intracellular domain as found in the human FcγRIIA sequence (Table 11). Neither the cynomolgus nor human intracellular domains contain an ITAM motif; the activating ITAM motif for human FcγRIIA is supplied by the associated γ-chain and the same situation most likely occurs in cynomolgus monkeys.

[0115] The cynomolgus FcγRIIA α-chain binds to hexameric complexes of subclasses IgG with the following binding pattern: IgG1>IgG3>>IgG2>IgG4b, IgG4. A human FcγRIIA isoform with a phenylalanine at the amino acid corresponding to the amino acid 158 (F158) binds hexameric IgG subclasses as follows: IgG3=IgG1>>>IgG2, IgG4. A human FcγRIIA isoform with a valine at the amino acid corresponding to the amino acid 158 (V158) binds hexameric IgG subclasses as follows: IgG1>IgG3>>>IgG2A, IgG4. Cynomolgus FcγRIIA with an amino acid sequence of SEQ ID NO: 20 has an isoleucine at amino acid position corresponding to amino acid 158 and binds human Ig subclasses similar to human FcγRIIA VI 58.

[0116] Human IgG1 binds to human FcγRIIA-V158 better than it does to human FcγRIIA-F158 (Koene, H. R., Kleijer, M., Algra, J., Roos, D., von dem Borne, E. G. K., and de Hass, M. (1997) *Blood* 90, 1109-1114; Wu, J., Edberg, J. C., Redecha, P. B., Bansal, V., Guyre, P. M., Coleman, K., Salmon, J. E., and Kimberly, R. P. (1997) *J. Clin. Invest.* 100, 1059-1070; Shields, R. L., Namenuk, A. K., Hong, K., Meng, Y. G., Rae, J., Briggs, J., Xie, D., Lai, J., Stadlen, A., Li, B., Fox, J. A., and Presta, L. G. (2001) *J. Biol. Chem.* 276, 6591-6604). In humans, the FcγRIIA-F158 allele predominates with approximately 90% of humans having at least one FcγRIIA-F158 allele (Lehrnbecher, T., Foster, C. B., Zhu, S., Leitman, S. F., Goldin, L. R., Huppi, K., and Chanock, S. J. (1999) *Blood* 94, 4220-4232). In addition, recent studies have begun to correlate specific disease states with the FcγRIIA polymorphic status of individuals (Wu, J., Edberg, J. C., Redecha, P. B., Bansal, V., Guyre, P. M., Coleman, K., Salmon, J. E., and Kimberly, R. P. (1997) *J. Clin. Invest.* 100, 1059-1070; Lehrnbecher, T., Foster, C. B., Zhu, S., Venzon, D., Steinberg, S. M., Wyvill, K., Metcalf, J. A., Cohen, S. S., Kovacs, J., Yarchan, R., Blauvelt, A., and Chanock, S. J. (2000) *Blood* 95, 2386-2390; Nieto, A., Caliz, R., Pascual, M., Mataran, L., Garcia, S., and Martin, J. (2000) *Arthritis & Rheumatism* 43, 735-739). Notably, the chimpanzee and cynomolgus FcγRIIA have valine and isoleucine, respectively, at position 158. The similarity of binding of the four human subclasses of IgG to cynomolgus FcγRIIA and human FcγRIIA-V158 (as opposed to human FcγRIIA-F158) suggests that evaluation of human antibodies in primate models should account for the primate model reflecting only a minority of humans with respect to binding to FcγRIIA receptors, i.e. FcγRIIA-V158/V158 homozygotes. For example, since human FcγRIIA-V158 exhibits superior antibody-dependent cellular cytotoxicity (ADCC) compared to human FcγRIIA-F158 (Shields, R. L., Namenuk, A. K., Hong, K., Meng, Y. G., Rae, J., Briggs, J., Xie, D., Lai, J., Stadlen, A., Li, B., Fox, J. A., and Presta, L. G. (2001) *J. Biol. Chem.* 276, 6591-6604), primate models may overestimate the efficacy of human antibody effector functions associated with FcγRIIA.

[0117] However, the binding patterns of human IgG subclasses to other cynomolgus FcRs, especially FcγRI, indicate that the non-human primates can be used as effective models to evaluate the safety, efficacy and pharmacokinetics of Fc region binding molecules.

[0118] The invention also provides for Fc receptor polypeptides identified as FcRn. Amino acid sequences of cynomolgus FcRn are shown in Table 14. In Table 14, the numbers shown below the amino acids and designated with

the signal  $\Delta$  are numbered from the start of the mature polypeptide. Two alleles were identified and are shown in Table 14. A cynomolgus FcRn  $\alpha$ -chain has an amino acid sequence of SEQ ID NO: 29 with a serine at residue 3 of the mature polypeptide. A cynomolgus FcRn  $\alpha$ -chain has a sequence of SEQ ID NO: 64 and has an asparagine at residue 3 of the mature polypeptide. The mature polypeptides of FcRn  $\alpha$ -chain S3 and FcRn  $\alpha$ -chain N3 have a sequence of SEQ ID NO: 71 and 72, respectively. An extracellular fragment of a FcRn as obtained using the primers as described in example 1 has an amino acid sequence of  $\Delta 1$  to  $\Delta 274$  as shown in table 14.

**[0119]** A sequence alignment of cynomolgus FcRn  $\alpha$ -chain sequences to human FcRn  $\alpha$ -chain (SEQ ID NO: 20) (GenBank Accession No. U12255) shows that the cynomolgus sequence is about 97% identical to the human sequence. Cynomolgus FcRn (S3) and FcRn (N3)  $\alpha$ -chains bind to subclasses of IgG with the following binding pattern: IgG3>>IgG4>IgG2>IgG1, which is similar to that of the human FcRn  $\alpha$ -chain.

**[0120]** The invention also includes cynomolgus  $\beta$ -2 microglobulin polypeptides. A cynomolgus  $\beta$ -2 microglobulin polypeptide has a sequence of SEQ ID NO: 25, Table 13. The mature  $\beta$ -2 microglobulin polypeptide has a sequence of SEQ ID NO: 70. When the cynomolgus  $\beta$ -2 microglobulin sequence is aligned with a human sequence for  $\beta$ -2 microglobulin (SEQ ID NO: 26; GenBank Accession No. P01884), it shows that the cynomolgus sequence has about 92% sequence identity to human  $\beta$ -2 microglobulin.

**[0121]** Variants, derivatives, fusion proteins, and fragments of the different cynomolgus and chimp FcR polypeptides that retain any of the biological activities of the FcRs, are also within the scope of the present invention. Note that one of ordinary skill in the art will readily be able to determine whether a variant, derivative, or fragment of a FcR polypeptide displays activity by subjecting the variant, derivative, or fragment to an immunoglobulin binding assay as described below in Example 3.

**[0122]** Derivatives of the different cynomolgus and chimp FcRs can be polypeptides modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, polyethylene glycol (PEG) groups, lipids, phosphate, acetyl groups and the like.

**[0123]** In another embodiment, the polypeptides of the invention include fragments of the polypeptides that lack a portion or all of the transmembrane and intracellular domains: e.g. amino acid residues of the mature polypeptide as follows: FcRI  $\alpha$ -chain amino acid residues 270-336 of SEQ ID NO: 65; FcRIIA amino acid residues 183 to 282 of SEQ ID NO: 66; chimp FcRIIA amino acid residues 172 to 281 of SEQ ID NO: 67; FcRIIB amino acid residues 185 to 252 of SEQ ID NO: 68; FcRIIA  $\alpha$ -chain amino acid residues 188 to 234 of SEQ ID NO: 69; or FcRn amino acid residues 275 to 342 of SEQ ID NO: 71 or SEQ ID NO: 72. A soluble FcR polypeptide may include a portion of the transmembrane domain and intracellular, as long as the polypeptide is secreted from the cell in which it is produced. Preferably, the fragments are capable of binding to an Fc region containing molecule.

**[0124]** Fragments of polypeptides also include one or more domain of the polypeptide identified in Table 10 or

Table 11, including signal peptide, domain 1, domain 2, domain 3, transmembrane/intracellular, or a cytoplasmic domain including the ITAM or ITIM motif. Exemplary fragments of the polypeptides also include soluble polypeptides having only domain 1, domain 2 and domain 3 amino acid sequences of the corresponding mature FcR polypeptides: e.g., amino acid residues  $\Delta 1$  to  $\Delta 269$  of cynomolgus FcRI (Table 10), amino acid residues  $\Delta 1$  to  $\Delta 182$  of cynomolgus FcRIIA (Table 21), amino acid residues  $\Delta 1$  to  $\Delta 184$  of cynomolgus FcRIIB (Table 22), amino acid residues  $\Delta 1$  to  $\Delta 187$  of cynomolgus FcRIIA (Table 23), and amino acids  $\Delta 1$  to  $\Delta 274$  of cynomolgus FcRn (Table 14).

**[0125]** Cynomolgus or chimp FcR variants within the scope of the invention may comprise conservatively substituted sequences, meaning that one or more amino acid residues of each polypeptide may be replaced by different residues that do not alter the secondary and/or tertiary structure of the polypeptide. Such substitutions may include the replacement of an amino acid by a residue having similar physicochemical properties, such as substituting one aliphatic residue (Ile, Val, Leu or Ala) for another, or substitution between basic residues Lys and Arg, acidic residues Glu and Asp, amide residues Gln and Asn, hydroxyl residues Ser and Tyr, or aromatic residues Phe and Tyr. Further information regarding making phenotypically silent amino acid exchanges may be found in Bowie et al., *Science* 247:1306-1310 (1990). Other variants which might retain substantially the biological activities of the proteins are those where amino acid substitutions have been made in areas outside functional regions of the protein.

**[0126]** The invention also provides variant cynomolgus and chimp FcR polypeptides. Variant polypeptide can include changes to the polypeptide sequence that result in the amino acid substitutions, additions, and deletions in the resultant variant polypeptide when compared to the native polypeptide, for instance SEQ ID NOs: 9, 15, 17, 18, 20, 25, 29, or 64. The changes to the variant polypeptide sequences can include changes to the nucleic acid sequence that result in replacement of an amino acid by a residue having similar physicochemical properties, such as substituting one aliphatic residue (Ile, Val, Leu, or Ala) for another, or substitutions between basic residues Lys and Arg, acidic residues Glu and Asp, amide residues Gln and Asn, hydroxyl residues Ser and Tyr, or aromatic residues Phe and Tyr. Variant polypeptide sequences of the present invention are preferably at least about 90% identical, more preferably at least about 91% identical, more preferably at least 92% or 93% identical, more preferably 94% identical, more preferably 95% or 96% identical, more preferably 97% or 98% identical, and most preferably at least about 99% identical, to a full length native sequence, a polypeptide lacking a signal sequence, an extracellular domain of the polypeptide, or a fragment of the Fc receptor or  $\beta$ -2 microglobulin of sequences of SEQ ID NOs: 9, 15, 17, 18, 20, 25, 29, or 64.

**[0127]** Another embodiment of the present invention are polypeptides of the invention fused to heterologous amino acids, peptides, or polypeptides. Such amino acids, peptides, or polypeptides, preferably facilitate purification of the polypeptide. Many of the available peptides used for such a function allow selective binding of the fusion protein to a binding partner. For example, the cynomolgus FcRI polypeptide, having a sequence as shown in SEQ ID NO:9, may be modified to comprise a peptide to form a fusion

protein which specifically binds to a binding partner, or peptide tag. Non-limiting examples of such peptide tags include the 6-His tag, Gly/His<sub>6</sub>/GST tag, thioredoxin tag, hemagglutinin tag, GlyIh156 tag, and OmpA signal sequence tag. Full length, variable and truncated polypeptides of the present invention may be fused to such heterologous amino acids, peptides, or polypeptides. For example, the transmembrane and intracellular domains of cynomolgus FcγRIA can be replaced by DNA encoding the Gly/His<sub>6</sub>/GST tag fused as His271. As will be understood by one of skill in the art, the binding partner which recognizes and binds to the peptide may be any molecule or compound including metal ions (e.g., metal affinity columns), antibodies, or fragments thereof, and any protein or peptide which binds the peptide, such as the FLAG tag. The polypeptides of the present invention can also be fused to the immunoglobulin constant domain of an antibody to form immunoadhesin molecules.

[0128] The polypeptides of the present invention are preferably provided in an isolated form, and preferably are purified. The polypeptides may be recovered and purified from recombinant cell cultures by well-known methods, including ammonium sulfate or ethanol precipitation, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. In a preferred embodiment, high performance liquid chromatography (HPLC) is employed for purification.

#### Vectors and Host Cells

[0129] The present invention also relates to vectors comprising the polynucleotide molecules of the invention, as well as host cell transformed with such vectors. Any of the polynucleotide molecules of the invention may be joined to a vector, which generally includes a selectable marker and an origin of replication, for propagation in a host. Host cells are genetically engineered to express the polypeptides of the present invention. The vectors include DNA encoding any of the polypeptides described above or below, operably linked to suitable transcriptional or translational regulatory sequences, such as those derived from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, mRNA ribosomal binding sites, and appropriate sequences which control transcription and translation. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the DNA encoding the target protein. Thus, a promoter nucleotide sequence is operably linked to a cynomolgus monkey or chimp FcγR DNA sequence, FcRn α-chain DNA sequence, or β-2 microglobulin DNA sequence if the promoter nucleotide sequence directs the transcription of the FcγR sequence.

[0130] Expression of non-human primate receptors of the invention can also be accomplished by removing the native nucleic acid encoding the signal sequence or replacing the native nucleic acid signal sequence with a heterologous signal sequence. Heterologous signal sequences include those from human Fc receptor polypeptides or other polypeptides, such as tissue plasminogen activator. Nucleic acids encoding signal sequences from heterologous sources are known to those of skill in the art.

[0131] Selection of suitable vectors to be used for the cloning of polynucleotide molecules encoding the target

polypeptides of this invention will depend upon the host cell in which the vector will be transformed, and, where applicable, the host cell from which the target polypeptide is to be expressed. Suitable host cells for expression of the polypeptides of the invention include prokaryotes, yeast, and higher eukaryotic cells, each of which is discussed below.

[0132] Expression of functional cynomolgus monkey or chimp FcγR polypeptides of the invention may require the genetic engineering of a host cell to contemporaneously express two or more polypeptide molecules. As was discussed previously, most FcγRs are complex molecules requiring the expression of both a IgG binding and a signal transducing polypeptide chain. The complex of two or more polypeptide chains forms the functional receptor. As such, for example, a host cell may be co-transfected with a first vector expressing the FcγRI α-chain, having a first selection marker, and a second vector expressing the FcγRI γ-chain, having a second selection marker. Only host cells that have acquired both vectors and are expressing both polypeptides would survive and express functional FcγRI. Other methods are envisioned for the co-transfection of multiple polypeptide chains into target host cells, including the linked expression of target polypeptides from the same vector.

[0133] The cynomolgus monkey or chimp FcγR, FcRn, or β-2 microglobulin polypeptides to be expressed in such host cells may also be fusion proteins which include regions from heterologous proteins. Such regions may be included to allow, e.g., secretion, improved stability, or facilitated purification of the polypeptide. For example, a sequence encoding an appropriate signal peptide can be incorporated into expression vectors. A DNA sequence for a signal peptide (secretory leader) may be fused in-frame to the target sequence so that target protein is translated as a fusion protein comprising the signal peptide. The DNA sequence for a signal peptide can replace the native nucleic acid encoding a signal peptide or in addition to the nucleic acid sequence encoding the native sequence signal peptide. A signal peptide that is functional in the intended host cell promotes extracellular secretion of the polypeptide. Preferably, the signal sequence will be cleaved from the target polypeptide upon secretion from the cell. Non-limiting examples of signal sequences that can be used in practicing the invention include the yeast I-factor and the honeybee melatin leader in Sf9 insect cells.

[0134] Suitable host cells for expression of target polypeptides of the invention include prokaryotes, yeast, and higher eukaryotic cells. Suitable prokaryotic hosts to be used for the expression of these polypeptides include bacteria of the genera *Escherichia*, *Bacillus*, and *Salmonella*, as well as members of the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. For expression in, e.g., *E. coli*, a target polypeptide may include an N-terminal methionine residue to facilitate expression of the recombinant polypeptide in a prokaryotic host. The N-terminal Met may optionally then be cleaved from the expressed polypeptide.

[0135] Expression vectors for use in prokaryotic hosts generally comprise one or more phenotypic selectable marker genes. Such genes generally encode, e.g., a protein that confers antibiotic resistance or that supplies an auxotrophic requirement. A wide variety of such vectors are readily available from commercial sources. Examples include pSPORT vectors, pGEM vectors (Promega),

pPROEX vectors (LTI, Bethesda, Md.), Bluescript vectors (Stratagene), and pQE vectors (Qiagen).

**[0136]** The cynomolgus monkey or chimp Fc $\gamma$ R, FcRn, or P-2 microglobulin, may also be expressed in yeast host cells from genera including *Saccharomyces*, *Pichia*, and *Kluveromyces*. Preferred yeast hosts are *S. cerevisiae* and *P. pastoris*. Yeast vectors will often contain an origin of replication sequence from a 2T yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Vectors replicable in both yeast and *E. coli* (termed shuttle vectors) may also be used. In addition to the above-mentioned features of yeast vectors, a shuttle vector will also include sequences for replication and selection in *E. coli*. Direct secretion of the target polypeptides expressed in yeast hosts may be accomplished by the inclusion of nucleotide sequence encoding the yeast I-factor leader sequence at the 5' end of the cynomolgus Fc $\gamma$ R-encoding nucleotide sequence.

**[0137]** Insect host cell culture systems may also be used for the expression of the polypeptides of the invention. In a preferred embodiment, the target polypeptides of the invention are expressed using a baculovirus expression system. Further information regarding the use of baculovirus systems for the expression of heterologous proteins in insect cells are reviewed by Luckow and Summers, *Bio/Technology* 6:47 (1988).

**[0138]** In another preferred embodiment, the cynomolgus Fc $\gamma$ R polypeptides are individually expressed in mammalian host cells. Non-limiting examples of suitable mammalian cell lines include the COS-7 line of monkey kidney cells (Gluzman et al., *Cell* 23:175 (1981)), Chinese hamster ovary (CHO) cells (Puck et al., *Proc. Natl. Acad. Sci. USA*, 60:1275-1281 (1958), CV-1 and human cervical carcinoma cells (HELA) (ATCC CCL 2). Preferably, HEK293 cells are used for expression of the target proteins of this invention.

**[0139]** The choice of a suitable expression vector for expression of the target polypeptides of the invention will of course depend upon the specific mammalian host cell to be used, and is within the skill of the ordinary artisan. Examples of suitable expression vectors include pcDNA3.1/Hygro (Invitrogen), 409, and pSVL (Pharmacia Biotech). A preferred vector for expression of the cynomolgus Fc $\gamma$ R polypeptides is pRK. Eaton, D. L., Wood, W. I., Eaton, D., Hass, P. E., Hollingshead, P., Wion, K., Mather, J., Lawn, R. M., Vohar, G. A., and Gorman, C. (1986) *Biochemistry* 25:8343-47. Expression vectors for use in mammalian host cells may include transcriptional and translational control sequences derived from viral genomes. Commonly used promoter sequences and enhancer sequences which may be used in the present invention include, but are not limited to, those derived from human cytomegalovirus (CMV), Adenovirus 2, Polyoma virus, and Simian virus 40 (SV40). Methods for the construction of mammalian expression vectors are disclosed, for example, in Okayama and Berg (*Mol. Cell. Biol.* 3:280 (1983)); Cosman et al. (*Mol. Immunol.* 23:935 (1986)) and Cosman et al. (*Nature* 312:768 (1984)).

#### Method of Evaluating Biological Properties, Safety and Efficacy of Fc Region Containing Molecules

**[0140]** One aspect of the invention includes a method for the evaluation of the pharmacokinetics/pharmacodynamics

of FcR binding molecules such as humanized antibodies with cynomolgus monkey or chimp Fc receptors prior to an in vivo evaluation in a primate. This aspect of the invention is based on the finding that cynomolgus and chimp FcR polypeptides have a high degree of sequence identity with human Fc receptor polypeptides and bind to IgG subclasses in a similar manner. Evaluations can include testing, for example, humanized antibodies of any subclass (especially antibodies with prospective therapeutic utility) on target Fc receptors of the invention in an ELISA-format assay or to transiently expressing cells.

**[0141]** A method of the invention involves evaluating the binding of a Fc region containing polypeptide or agent to cynomolgus or chimp Fc receptor polypeptide by contacting the Fc region containing molecule with a cynomolgus or chimp Fc receptor polypeptide. The cynomolgus or chimp Fc receptor polypeptide can be soluble or can be expressed as a membrane bound protein on transiently infected cells. Binding of the Fc region containing molecule to the cynomolgus or chimp Fc receptor polypeptide indicates that the Fc region containing molecule or polypeptide is suitable for in vivo evaluation in a primate. Binding to cynomolgus FcRn molecules provides an indication that Fc region containing molecule or polypeptide will have a longer half-life in vivo.

**[0142]** The invention also provides for screening variants of Fc region containing molecules such as antibody variants for their biological properties, safety, efficacy and pharmacokinetics. Antibody variants are typically altered at one or more residues and then the variants are analyzed for alteration in biological activities including altered binding affinity for Fc receptors. Screening for alterations in biological activities by variants may be tested both in vivo and in vitro. For example, receptor polypeptides of the present invention can be used in an ELISA-format assay or transiently infected cells. Antibody variants which bind to cynomolgus and/or chimp FcR polypeptides, such as the  $\alpha$ -chain of Fc $\gamma$ RII, Fc $\gamma$ RIII or FcRn or Fc $\gamma$ RIIA or Fc $\gamma$ RIIB, are variants that are suitable for in vivo evaluation in primates as a therapeutic agent.

**[0143]** Direct binding and binding affinity determination between the different Fc region containing molecules is preferably performed against soluble extracellular domains of cynomolgus Fc $\gamma$ R polypeptides. For example, the transmembrane domain and intracellular domain of a target Fc $\gamma$ R can be replaced by DNA encoding a Gly-His<sub>6</sub> tag or glutathione S-transferase (GST) (see Example 3). The Gly-His<sub>6</sub> tag or GST provide a convenient method for immobilizing the Fc binding region of the receptor to a solid support for identification and/or determination of binding affinities between the receptor and target antibody variant. Potential assays include ELISA-format assays, co-precipitation format assays, and column chromatographic format assays. Identified Fc region containing molecules should directly interact with the soluble cynomolgus Fc $\gamma$ R and have equivalent or greater binding affinities for the cynomolgus Fc $\gamma$ R, as compared to corresponding human Fc $\gamma$ R.

**[0144]** Another aspect of the invention provides methods of identifying agents that have altered binding to a cynomolgus Fc $\gamma$ R comprising an ITAM and/or ITIM region. A method of the invention involves identifying an agent that

has increased binding affinity for an FcR comprising an ITAM region and a decreased affinity for a FcR comprising an ITIM region.

**[0145]** Target agents include molecules that have a Fc region, preferably an antibody and more preferably an IgG antibody. If the target agent is an antibody it may be a variant antibody with an altered amino acids sequence compared to the native sequence of the antibody. Preferably variant antibodies have had amino acid substitutions in regions of the antibody that are involved in binding to Fc $\gamma$  receptor, including amino acids corresponding to amino acids 226 to 436 in a human IgG. Variant antibodies can be prepared using standard methods such as site specific oligonucleotide or PCR mediated methods as described previously. Examples of variant antibodies includes alanine variants of human IgG1, anti IgE E27 prepared as described in Shields et al., *J. Biol. Chem.* 276:6591 (2001).

**[0146]** Binding affinities of antibodies and/or variant antibodies are determined using standard methods as described in Shields et al., *J. Biol. Chem.* 276:6591 (2001) and in Examples 3-7 below. Binding affinities are preferably determined by binding to cells that express a Fc $\gamma$  receptor of the type being analyzed. However, binding affinities of antibodies or Fc region containing molecules can also be determined using soluble Fc $\gamma$  receptors or Fey receptors expressed on or secreted from a host cell.

**[0147]** A variant antibody that has an increased affinity for a cynomolgus Fc $\gamma$ RIIA compared with a human Fc $\gamma$ RIIA is an antibody that has a change in amino acid sequence at the position corresponding to amino acid 298 of human IgG1. One such variant has a change at that position from serine to alanine and is designated as S298A. Another variant antibody with a change at that position is designated as S298A/E333A/K334 which is a variant antibody with alanine in positions corresponding to amino acid 298, 333 and 334 of native sequence IgG1. These variants have increased binding affinity to a cynomolgus Fc $\gamma$ RIIA compared to a human Fc $\gamma$ RIIA.

**[0148]** In another method of the invention, target agents with altered binding affinity to a cynomolgus Fc $\gamma$ RIIB as compared to human Fc $\gamma$ RIIB are identified. The agents are preferably variants of native sequence antibodies. Binding affinities are determined as described above and as shown in the Examples below. Agents with enhanced binding to a Fc $\gamma$ RIIB may preferentially stimulate ITIM inhibitory functions. Agents with decreased affinity for a cynomolgus Fc $\gamma$ RIIB may have decreased stimulation of inhibitory function.

**[0149]** Variant antibodies that have decreased affinity for a cynomolgus Fc $\gamma$ RIIB compared to a human Fc $\gamma$ RIIB are: R255A, E258A, S37A, D280A and R301M.

**[0150]** Another embodiment of the invention involves the use of variant antibodies S298A or S298A/E333A/K334 to identify agents that can activate Fc $\gamma$  receptors comprising an ITAM while not engaging Fey receptors comprising an ITIM region.

**[0151]** Variant antibodies with S298A, and S292A/E333A/K334, have increased binding affinity to a cynomolgus Fc $\gamma$ RIIA, and decreased binding affinity to a cynomolgus Fc $\gamma$ RIIB. Such methods can be conducted in vivo or in vitro.

**[0152]** These methods are also useful for identifying the location of amino acid in native sequence antibodies that can be modified to increase binding of the antibody to FcR polypeptides, preferably human and cynomolgus Fc $\gamma$ R, comprising an ITAM region and/or to decrease binding affinity to Fc $\gamma$ R comprising an ITIM region. Modifications to the amino acid sequence at the identified locations can be prepared by standard methods.

**[0153]** Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

## EXAMPLES

### Example 1

#### Molecular Cloning of Cynomolgus and Chimp Fc Receptor DNA and $\beta$ -2 Microglobulins

#### **[0154]** Materials and Methods:

##### Cloning of Cynomolgus Monkey Fc $\gamma$ R

**[0155]** Since cynomolgus monkey DNA shares approximately 90% homology to human DNA, a series of PCR primers for each Fc $\gamma$ R was designed based on the sequence of the corresponding human receptor. Each sense primer starts at a site immediately 5' of the coding region or at the start of the coding region. The antisense primers were designed in the same way, i.e. immediately 3' of the C terminal stop codon or at the C terminal stop codon. Primers incorporated endonuclease restriction sites used to subclone PCR product into a pRK vector (Eaton et al.). The sequences of the primers are shown in Table 1.

TABLE 1

Restriction sites are underlined.	
Receptor	Cyno Fc $\gamma$ RI Full-Length
Forward Primer	CAGGTCAATCTCTAGACTCCACCGCTTGGAG (SEQ ID NO: 31)
Reverse Primer	GGTCAACTATAAGCTTGGACGGTCCAGATCGAT (SEQ ID NO: 32)
Restriction XbaI/HindIII Sites	
Receptor	Cyno Fc $\gamma$ RI-H6-GST
Forward Primer	CAGGTCAATCATCGATATGTGGTTCTTGACAGCT (SEQ ID NO: 33)
Reverse Primer	GGTCAACTATGCTAGCATGGTGATGATGGTGGTGCCAG ACAGGAGTTGGTA (SEQ ID NO: 34)
Restriction ClaI/NheI Sites	
Receptor	Cyno Fc $\gamma$ RIIB Full-Length
Forward Primer	CAGGTCAATCTCTAGAAATGGGAATCCTGTCATTCTT (SEQ ID NO: 35)
Reverse Primer	GGTCAACTATAAGCTTCTAAATACGGTTCTGGTC (SEQ ID NO: 36)
Restriction XbaI/HindIII Sites	

TABLE 1-continued

Restriction sites are underlined.	
Receptor	Cyno Fc $\gamma$ RIIB-H6-GST
Forward Primer	CAGGTCAATC <u>ATCGAT</u> ATGCTTCTGTGGACAGC (SEQ ID NO: 37)
Reverse Primer	GGTCAACTATGGT <u>GACCT</u> ATCGGTGAAGAGCTGC (SEQ ID NO: 38)
Restriction ClaI/BstEII Sites	
Receptor	Cyno Fc $\gamma$ RIIIA Full-Length
Forward Primer	CAGGTCAATCT <u>CTAGAA</u> TGTGGCAGCTGCTCCT (SEQ ID NO: 39)
Reverse Primer	TCAACTATA <u>AGCTT</u> ATGTTCAGAGATGCTGCTG (SEQ ID NO: 40)
Restriction XbaI/HindIII Sites	
Receptor	Cyno Fc $\gamma$ RIIIA-H6-GST
Forward Primer	CAGGTCAATCT <u>CTAGAA</u> TGTGGCAGCTGCTCCT (SEQ ID NO: 41)
Reverse Primer	GGTCAACTATGGT <u>CACCT</u> TGGTACCCAGGTGGAAA (SEQ ID NO: 42)
Restriction XbaI/BstEII Sites	
Receptor	Cyno Fc $\gamma$ Chain
Forward Primer	CAGGTCAATCATCGAT <u>GAATTC</u> CCACCATGATTCCAGC AGTGGTC (SEQ ID NO: 43)
Reverse Primer	GGTCAACTATA <u>AGCTT</u> CTACTGTGGTGGTTTCTCA (SEQ ID NO: 44)
Restriction EcoRI/HindIII Sites	
Receptor	Cyno $\beta$ -2 Microglobulin
Forward Primer	CAGGTCAATC <u>ATCGAT</u> TCGGGCCGAGATGTCT (SEQ ID NO: 45)
Reverse Primer	GGTCAACTAT <u>CTAGAT</u> TACATGTCTCGATCCCA (SEQ ID NO: 46)
Restriction ClaI/XbaI Sites	
Receptor	Cyno Fc $\gamma$ RIIA Full-Length
Forward Primer	CAGGTCAATCT <u>CTAGAA</u> TGTCTCAGAATGTATGTC (SEQ ID NO: 47)
Reverse Primer	GGTCAACTATA <u>AGCTT</u> TTAGTTATTACTGTTGTCATA (SEQ ID NO: 48)
Restriction XbaI/HindIII Sites	
Receptor	Cyno Fc $\gamma$ RIIA-H6-GST
Forward Primer	CAGGTCAATC <u>ATCGAT</u> ATGTCTCAGAATGTATGTC (SEQ ID NO: 49)

TABLE 1-continued

Restriction sites are underlined.	
Reverse Primer	GGTCAACTATGGT <u>GACCT</u> ATCGGTGAAGAGCTGC (SEQ ID NO: 50)
Restriction ClaI/BstEII Sites	
Receptor	Cyno FcRn Full-Length
Forward Primer	CAGGTCAATC <u>ATCGAT</u> AGGTCGTCCTCTCAGC (SEQ ID NO: 51)
Reverse Primer	GGTCAACTAT <u>GAATTC</u> TCGGAATGGCGGATGG (SEQ ID NO: 52)
Restriction ClaI/EcoRI Sites	
Receptor	Cyno FcRn-H6
Forward Primer	CAGGTCAATC <u>ATCGAT</u> AGGTCGTCCTCTCAGC (SEQ ID NO: 53)
Reverse Primer	GGTCAACTAT <u>GAATTC</u> ATGGTGATGATGGTGGTGGCAG GACTTGGCTGGAGTTTC (SEQ ID NO: 54)
Restriction ClaI/EcoRI Sites	

[0156] The cDNA for FcRs was isolated by reverse transcriptase-PCR (GeneAmp, PerkinElmer Life Sciences) of oligo(dT)-primed RNA from cynomolgus spleen cells using primers as shown in Table 1. The cDNA was subcloned into previously described pRK mammalian cell expression vectors, as described in Eaton et al., 1986, *Biochemistry*, 25:8343-8347. PCR reactions were set up using 200 ng of cDNA vector library from cynomolgus spleen and ExTaq Premix (Panvera, Madison, Wis.) according to the manufacturers instructions. After denaturation at 90° C. for 30 s, 25 cycles were run with annealing at 55° C. for 1 min, elongation at 72° C. for 3 min, and denaturation at 98° C. for 30 s. DNA bands migrating at the expected size (Fc $\gamma$ RI, Fc $\gamma$ RIIIA, FcRn, 1100 base pairs; Fc $\gamma$ RIIA, Fc $\gamma$ RIIB, 1000 base pairs; Fc $\gamma$  chain, 300 base pairs;  $\beta$ -2 microglobulin, 400 base pairs) were isolated, cloned into pRK vectors, then transformed into *Escherichia coli* XL 1-Blue (Stratagene, San Diego, Calif.). Individual clones were selected and double-stranded DNA for each was purified using Qiagen mini-prep DNA kits (cat. # 27106; Qiagen). DNA sequencing was performed on an Applied Biosystems model 377 sequencer using Big-Dye Terminator Cycle Sequencing kits (Applied Biosystems, Foster City, Calif.).

[0157] Initial PCR reactions for Fc $\gamma$ RIIA did not reveal a PCR product. To determine whether or not Fc $\gamma$ RIIA was present in cynomolgus monkeys, a sense primer was designed in a region conserved between human Fc $\gamma$ RIIA, human Fc $\gamma$ RIIB, and cynomolgus Fc $\gamma$ RIIB (OF1, Table 2). An antisense primer was designed based on the consensus sequence in the region encoding the ITAM of human Fc $\gamma$ RIIA (OR1, Table 2). Using these two PCR primers (OF1, OR1) and the PCR protocol described above, a PCR product of approximately 700 base pairs was obtained. The PCR band was isolated and subcloned into a pRK vector,



individual clones were isolated and sequenced as described above. Sequence analysis revealed that the fragment had 90% identity to human FcγRIIA.

[0158] In order to determine the DNA sequence at the 5' end of the receptor, a nested PCR reaction was utilized. For the first step of the nested PCR reaction, a sense PCR primer (OF2, Table 2) was designed to lay down on the pRK vector 5' of the vector cloning site. This primer was used in conjunction with reverse primer OR1. The PCR reaction was performed on the cDNA library as described above, the product was diluted 1:500 and 1 μL was used as a template for the second step of the nested PCR reaction. Due to the fact that primer OF2 would lay down on all members of the cDNA library (all members being cloned into separate pRK vectors), only a small quantity of PCR fragment was obtained and hence this was used as a template for amplification in the second step. The sense primer (OF3, Table 2) for the second step was designed to lay down on the pRK vector sequence 3' of OF2 and the reverse primer (OR2, Table 2) was based on partial sequence of FcγRIIA determined above. The second step of the nested PCR reaction revealed a band of approximately 600 base pairs. The band was isolated and individual clones were prepared and sequenced as described above.

[0159] The DNA sequence at the 3' end of the receptor was determined in a similar manner. An initial PCR reaction on the cDNA library was performed using the forward primer OF4, designed from the sequence of the FcγRIIA fragment, and the reverse primer OR3, designed to lay down in the pRK vector 3' from the end of the FcγRIIA. The resultant fragment was used as template for the second step of the nested PCR reaction. The second step used the forward primer OF5, designed from the sequence of the FcγRIIA fragment, and the reverse primer OR4, designed to lay down in the pRK vector 5' from primer OR3. The second step of the nested PCR reaction revealed a band of approximately 800 base pairs. The band was isolated and individual clones were sequenced as described above. PCR primers for the full length FcγRIIA were designed based on the information acquired from the nested PCR reactions. Full length FcγRIIA was cloned using the method described for all other receptors. The sequences of the primers described above are shown in Table 2.

TABLE 2

		(SEQ ID NO: 55)
OF1	CAGGTCAATCTCTAGACAGTGGTTCCACAATGG	
		(SEQ ID NO: 56)
OR1	GGTCAACTATAAGCTTAAGAGTCAGGTAGATGTTT	

TABLE 2-continued

		(SEQ ID NO: 57)
OF2	CAGGTCAATC TCTAGA ATACATAACCTTATGTATCAT	
		(SEQ ID NO: 58)
OF3	CAGGTCAATC TCTAGA TATAGAATAACATCCACTTTG	
		(SEQ ID NO: 59)
OR2	GGTCAACTAT AAGCTT CAGAGTCATGTAGCCG	
		(SEQ ID NO: 60)
OF4	CAGGTCAATC TCTAGA ATTCCACTGATCCTGTGAA	
		(SEQ ID NO: 61)
OR3	GGTCAACTAT AAGCTT GCTTTATTGTGAAATTGTG	
		(SEQ ID NO: 62)
OF5	CAGGTCAATC TCTAGA ACTTGGACGTCAAACGATT	
		(SEQ ID NO: 63)
OR4	GGTCAACTAT AAGCTT CTGCAATAACAAGTTGGG	

Example 2

Alignment of Nucleotide and Amino Acid Sequences of Cynomolgus, Chimp and Human FcγR

[0160] Nucleotide and amino acid sequences for FcR polypeptides from human, cynomolgus and chimps were aligned and % sequence identity calculated.

[0161] Nucleotide and amino acid sequences of primate and human proteins were aligned manually and differences in nucleotide or protein sequence noted. Percent identity was calculated as [number of identical residues]/[number of total residues]. When the sequences differed in the total number of residues, two values for percent identity are provided, using the two different numbers for total residues. Nucleotide sequences begin at the coding sequence for the signal sequence.

[0162] The alignment of nucleic acid sequences for human (SEQ ID NO: 2) and cynomolgus FcγRI α-chain (SEQ ID NO: 1) as shown in Table 3 below. The dots indicate locations of nucleotide sequence differences. An analysis of the % sequence identity shows that the human and cynomolgus nucleotide sequences encoding FcγRI α-chain have about 91% or 96% sequence identity depending on whether the nucleotides of 3' extensions are included in the calculation.

TABLE 3

Alignment of Human and Cynomolgus High-Affinity FcγRI DNA					
1030 matches in an overlap of 1074: 95.9% identity					
1030 matches in an overlap of 1128: 91.3% identity					
	10	20	30	40	50
Human	ATGTGGTCTCTTGACA	ACTCTGCTCCTTTGGG	TTCAGTTGATGGG	CAAGT	
.					
Cyno	ATGTGGTCTCTTGAC	AGCTCTGCTCCTTTGGG	TTCAGTTGATGGG	CAAGT	

TABLE 3-continued

Alignment of Human and Cynomolgus High-Affinity FcγRI DNA 1030 matches in an overlap of 1074: 95.9% identity 1030 matches in an overlap of 1128: 91.3% identity					
	60	70	80	90	100
Human	GGACACCACAAAGGCAGTGATCACTTTGCAGCCTCCATGGGTCAGCGTGT				
	•				
Cyno	GGATACCACAAAGGCAGTGATCACTTTGCAGCCTCCATGGGTCAGCGTGT				
	110	120	130	140	150
Human	TCCAAGAGGAAACCGTAACCTTGCACTGTGAGGTGCTCCATCTGCCTGGG				
	• • • • •				
Cyno	TCCAAGAGGAAACTGTAACCTTACAGTGTGAGGTGCCCCGTCTGCCTGGG				
	160	170	180	190	200
Human	AGCAGCTCTACACAGTGGTTTCTCAATGGCACAGCCACTCAGACCTCGAC				
	•				
Cyno	AGCAGCTCCACACAGTGGTTTCTCAATGGCACAGCCACTCAGACCTCGAC				
	210	220	230	240	250
Human	CCCCAGCTACAGAATCACCTCTGCCAGTGTCAATGACAGTGGTGAATACA				
	• •				
Cyno	TCCCAGCTACAGAATCACCTCTGCCAGTGTCAAGGACAGTGGTGAATACA				
	260	270	280	290	300
Human	GGTGCCAGAGAGGTCTCTCAGGGCGAAGTGACCCCATACAGCTGGAATC				
	•				
Cyno	GGTGCCAGAGAGGTCCCTCAGGGCGAAGTGACCCCATACAGCTGGAATC				
	310	320	330	340	350
Human	CACAGAGGCTGGCTACTACTGCAGGTCTCCAGCAGAGTCTTCACGGAAGG				
	• • •				
Cyno	CACAGAGACTGGCTACTACTGCAGGTATCCAGCAGAGTCTTCACAGAAGG				
	360	370	380	390	400
Human	AGAACCTCTGGCCTTGAGGTGTCATGCGTGGAAGGATAAGCTGGGTACA				
	•				
Cyno	AGAACCTCTGGCCTTGAGGTGTCATGCGTGGAAGGATAAGCTGGGTACA				
	410	420	430	440	450
Human	ATGTGCTTTACTATCGAAATGGCAAAGCCTTTAAGTTTTTCCACTGGAAT				
	• • • • •				
Cyno	ATGTGCTTTACTATCAAAATGGCAAAGCCTTTAAGTTTTTCTACCGGAAT				
	460	470	480	490	500
Human	TCTAACCTCACCATTCTGAAAACCAACATAAGTCACAATGGCACCTACCA				
	• • • • •				
Cyno	TCTCAACTCACCATTCTGAAAACCAACATAAGTCACAACGGCGCCTACCA				

TABLE 3-continued

Alignment of Human and Cynomolgus High-Affinity FcγRI DNA 1030 matches in an overlap of 1074: 95.9% identity 1030 matches in an overlap of 1128: 91.3% identity					
	510	520	530	540	550
Human	TTGCTCAGGCATGGGAAAGCATCGCTACACATCAGCAGGAATATCTGTCA				
	•			•	
Cyno	CTGCTCAGGCATGGGAAAGCATCGCTACACATCAGCAGGAGTATCTGTCA				
	560	570	580	590	600
Human	CTGTGAAAGAGCTATTTCCAGCTCCAGTGCTGAATGCATCTGTGACATCC				
				•	
Cyno	CTGTGAAAGAGCTATTTCCAGCTCCAGTGCTGAATGCATCCGTGACATCC				
	610	620	630	640	650
Human	CCACTCCTGGAGGGGAATCTGGTCACCCTGAGCTGTGAAACAAAGTTGCT				
	•				
Cyno	CCGCTCCTGGAGGGGAATCTGGTCACCCTGAGCTGTGAAACAAAGTTGCT				
	660	670	680	690	700
Human	CTTGCAGAGGCCTGGTTTGCAGCTTTACTTCTCCTTCTACATGGGCAGCA				
	••				
Cyno	TCTGCAGAGGCCTGGTTTGCAGCTTTACTTCTCCTTCTACATGGGCAGCA				
	710	720	730	740	750
Human	AGACCCTGCGAGGCAGGAACACATCCTCTGAATACCAAATACTAACTGCT				
		•			
Cyno	AGACCCTGCGAGGCAGGAACACGTCCTCTGAATACCAAATACTAACTGCT				
	760	770	780	790	800
Human	AGAAGAGAAGACTCTGGGTTTATACTGGTGCGAGGCTGCCACAGAGGATGG				
		•	••	• •	
Cyno	AGAAGAGAAGACTCTGGGTTTATACTGGTGCGAGGCCACCACAGAAGACGG				
	810	820	830	840	850
Human	AAATGTCCTTAAGCGCAGCCCTGAGTTGGAGCTTCAAGTGCTTGGCCTCC				
Cyno	AAATGTCCTTAAGCGCAGCCCTGAGTTGGAGCTTCAAGTGCTTGGCCTCC				
	860	870	880	890	900
Human	AGTTACCAACTCCTGTCTGGTTTCATGTCCTTTTCTATCTGGCAGTGGGA				
		•		•	
Cyno	AGTTACCAACTCCTGTCTGGCTTCATGTCCTTTTCTATCTGGTAGTGGGA				
	910	920	930	940	950
Human	ATAATGTTTTTAGTGAACACTGTTCTCTGGGTGACAATACGTAAAGAACT				
Cyno	ATAATGTTTTTAGTGAACACTGTTCTCTGGGTGACAATACGTAAAGAACT				

TABLE 3-continued

Alignment of Human and Cynomolgus High-Affinity FcγRI DNA 1030 matches in an overlap of 1074: 95.9% identity 1030 matches in an overlap of 1128: 91.3% identity					
	960	970	980	990	1000
Human	GAAAAGAAAGAAAAAGTGGGATTAGAAATCTCTTTGGATTCTGGTCATG				
		.	.	.	
Cyno	GAAAAGAAAGAAAAAGTGGGAATTAGAAATATCTTTGGATTCTGCTCATG				
	1010	1020	1030	1040	1050
Human	AGAAGAAGGTAATTTCCAGCCTTCAAGAAGACAGACATTTAGAAGAAGAG				
	.				
Cyno	AGAAGAAGGTAAC TTCCAGCCTTCAAGAAGACAGACATTTAGAAGAAGAG				
	1060	1070	1080	1090	1100
Human	CTGAAATGTCAGGAACAAAAAGAAGACAGCTGCAGGAAGGGGTGCACCG				
	..	.	.		
Cyno	CTGAAGAGTCAGGAACAAGAATAA				
	1110	1120			
Human	GAAGGAGCCCCAGGGGGCCACGTAGCAG 3' extension				

[0163] The Human DNA sequence shown in Table 3 has GenBank Accession No. L03418. Porges, A. J. Redecha, P. B., Doebele, R., Pan, L. C., Salmon, J. E. and Kimberly, R. P., *Novel Fc gamma receptor I family gene products in human mononuclear cells*, J. Clin Invest. 90, 2102-2109 (1992).

[0164] An alignment of nucleic acid sequences encoding human (SEQ ID NO: 14) and cynomolgus (SEQ ID NO: 13) gamma chain is shown in Table 4.

[0165] Analysis of the % sequence identity shows that the nucleic acid sequences encoding human and cynomolgus FcγRI/III gamma chain have about 99% identity.

TABLE 4

Alignment of Human and Cynomolgus Gamma-Chain DNA 258 matches in an overlap of 261: 98.9% identity					
	10	20	30	40	50
Human	ATGATTCCAGCAGTGGTCTTGCTCTTACTCCTTTTGGTTGAACAAGCAGC				
Cyno	ATGATTCCAGCAGTGGTCTTGCTCTTACTCCTTTTGGTTGAACAAGCAGC				
	60	70	80	90	100
Human	GGCCCTGGGAGAGCCTCAGCTCTGCTATATCCTGGATGCCATCCTGTTTC				
Cyno	GGCCCTGGGAGAGCCTCAGCTCTGCTATATCCTGGATGCCATCCTGTTTC				
	110	120	130	140	150
Human	TGTATGGAATTGTCCTCACCCTCCTCTACTGTCGACTGAAGATCCAAGTG				
Cyno	TGTATGGAATTGTCCTCACCCTCCTCTACTGTCGACTGAAGATCCAAGTG				
	160	170	180	190	200
Human	CGAAAGGCAGCTATAACCAGCTATGAGAAATCAGATGGTGTTTACACGGG				
	.				
Cyno	CGAAAGGCAGCTATAGCCAGCTATGAGAAATCAGATGGTGTTTACACGGG				

TABLE 4-continued

Alignment of Human and Cynomolgus Gamma-Chain DNA 258 matches in an overlap of 261: 98.9% identity					
	210	220	230	240	250
Human	CCTGAGCACCAGGAACCAGGAGACTTACGAGACTCTGAAGCATGAGAAAC				
		.	.		
Cyno	CCTGAGCACCAGGAACCAGGAAACTTATGAGACTCTGAAGCATGAGAAAC				
	260				
Human	CACCACAGTAG				
Cyno	CACCACAGTAG				

[0166] The DNA sequence for the human gamma chain as GenBank Accession No. M33195 J05285. Kuester, H., Thompson, H. and Kinet, J.-P., *Characterization and expression of the gene for the human receptor gamma subunit: Definition of a new gene family*, J. Biol. Chem. 265, 6448-6452 (1990).

[0167] An alignment of the human (SEQ ID NO: 4), chimp (SEQ ID NO: 22) and cynomolgus (SEQ ID NO: 3) nucleic acid sequence encoding FcγRIIA is shown in Table 5. An analysis of the % sequence identity shows that the human and cynomolgus sequences encoding FcγRIIA have about 94% sequence identity. A comparison of chimp and human sequences encoding FcγRIIA have about 99% sequence identity.

TABLE 5

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA DNA Human/Cyno 878 matches in an overlap of 933: 94.1% identity without one gap of three nucleotides 878 matches in an overlap of 936: 93.8% identity with one gap of three nucleotides Human/Chimp 924 matches in an overlap of 933: 99.0% identity without one gap of three nucleotides 924 matches in an overlap of 936: 98.7% identity with one gap of three nucleotides					
	10	20	30	40	50
Chimp	ATGTCTCAGAATGTATGTCCAGAAACCTGTGGCTGCTTCAACCATTGAC				
Human	ATGTCTCAGAATGTATGTCCAGAAACCTGTGGCTGCTTCAACCATTGAC				
		.	.		
Cyno	ATGTCTCAGAATGTATGTCCCGCAACCTGTGGCTGCTTCAACCATTGAC				
	60	70	80	90	100
Chimp	AGTTTTGCTGCTGCTGGCTTCTGCAGACAGTCAAGCT---GCTCCCCCAA				
				...	
Human	AGTTTTGCTGCTGCTGGCTTCTGCAGACAGTCAAGCTGCAGCTCCCCCAA				
				.	...
Cyno	AGTTTTGCTGCTGCTGGCTTCTGCAGACAGTCAAACT---GCTCCCCCGA				
	110	120	130	140	150
Chimp	AGGCTGTGCTGAAACTTGAGCCCCGTGGATCAACGTGCTCCAGGAGGAC				
Human	AGGCTGTGCTGAAACTTGAGCCCCGTGGATCAACGTGCTCCAGGAGGAC				
		.			.
Cyno	AGGCTGTGCTGAAACTCGAGCCCCGTGGATCAACGTGCTCCGGGAGGAC				

TABLE 5-continued

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA DNA					
Human/Cyno 878 matches in an overlap of 933: 94.1% identity without one gap of three nucleotides					
878 matches in an overlap of 936: 93.8% identity with one gap of three nucleotides					
Human/Chimp 924 matches in an overlap of 933: 99.0% identity without one gap of three nucleotides					
924 matches in an overlap of 936: 98.7% identity with one gap of three nucleotides					
	160	170	180	190	200
Chimp	TCTGTGACTCTGACATGCCGGGGGGCTCGCAGCCCTGAGAGCGACTCCAT				
		.			
Human	TCTGTGACTCTGACATGCCAGGGGGCTCGCAGCCCTGAGAGCGACTCCAT				
		.	..	.	.
Cyno	TCTGTGACTCTGACGTGCGGGGGCGCTCACAGCCCTGACAGCGACTCCAC				
	210	220	230	240	250
Chimp	TCAGTGGTTCCACAATGGGAATCTCATCCCCACCCACACGCAGCCCAGCT				
			.		
Human	TCAGTGGTTCCACAATGGGAATCTCATTCACCCACCCACACGCAGCCCAGCT				
			.	.	.
Cyno	TCAGTGGTTCCACAATGGGAATCGCATCCCCACCCACACAGCCCAGCT				
	260	270	280	290	300
Chimp	ACAGGTTCAAGGCCAACAACAATGACAGCGGGGAGTACACGTGCCAGACT				
Human	ACAGGTTCAAGGCCAACAACAATGACAGCGGGGAGTACACGTGCCAGACT				
			.	.	
Cyno	ACAGGTTCAAGGCCAACAACAATGATAGCGGGGAGTACAGGTGCCAGACT				
	310	320	330	340	350
Chimp	GGCCAGACCAGCCTCAGCGACCCTGTGCATCTGACTGTGCTTTCCGAATG				
Human	GGCCAGACCAGCCTCAGCGACCCTGTGCATCTGACTGTGCTTTCCGAATG				
	.		.	.	.
Cyno	GGCCGAGACCAGCCTCAGCGACCCTGTTCATCTGACTGTGCTTTCTGAGTG				
	360	370	380	390	400
Chimp	GCTGGTGCTCCAGACCCCTCACCTGGAGTTCCAGGAGGGAGAAACCATCG				
					.
Human	GCTGGTGCTCCAGACCCCTCACCTGGAGTTCCAGGAGGGAGAAACCATCA				
	.	.	.		
Cyno	GCTGGCGCTTCAGACCCCTCACCTGGAGTTCCGGGAGGGAGAAACCATCA				
	410	420	430	440	450
Chimp	TGCTGAGGTGCCACAGCTGGAAGGACAAGCCTCTGGTCAAGGTCACATTC				
Human	TGCTGAGGTGCCACAGCTGGAAGGACAAGCCTCTGGTCAAGGTCACATTC				
			.		
Cyno	TGCTGAGGTGCCACAGCTGGAAGGACAAGCCTCTGATCAAGGTCACATTC				

TABLE 5-continued

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA DNA					
Human/Cyno 878 matches in an overlap of 933: 94.1% identity without one gap of three nucleotides					
878 matches in an overlap of 936: 93.8% identity with one gap of three nucleotides					
Human/Chimp 924 matches in an overlap of 933: 99.0% identity without one gap of three nucleotides					
924 matches in an overlap of 936: 98.7% identity with one gap of three nucleotides					
	460	470	480	490	500
Chimp	TTCCAGAATGGAAAATCCCAGAAATTCTCCCATTTGGATCCCAACCTCTC				
			.	.	.
Human	TTCCAGAATGGAAAATCCCAGAAATTCTCCCGTTTGGATCCCACTTCTC				
		. . .	.	.	.
Cyno	TTCCAGAATGGAATAGCCAAGAAATTTTCCCATATGGATCCCAATTTCTC				
	510	520	530	540	550
Chimp	CATCCCACAAGCAAACCACAGTCACAGTGGTGATTACCACTGCACAGGAA				
Human	CATCCCACAAGCAAACCACAGTCACAGTGGTGATTACCACTGCACAGGAA				
Cyno	CATCCCACAAGCAAACCACAGTCACAGTGGTGATTACCACTGCACAGGAA				
	560	570	580	590	600
Chimp	ACATAGGCTACACGCTGTTCTCATCCAAGCCTGTGACCATCACTGTCCAA				
Human	ACATAGGCTACACGCTGTTCTCATCCAAGCCTGTGACCATCACTGTCCAA				
		. . .	.		
Cyno	ACATAGGCTACACACCATACTCATCCAAACCTGTGACCATCACTGTCCAA				
	610	620	630	640	650
Chimp	GCGCCAGCGTGGGCAGCTCTTCACCAATGGGGATCATTGTGGCTGTGGT				
	.	.	.		
Human	GTGCCAGCATGGGCAGCTCTTCACCAATGGGGATCATTGTGGCTGTGGT				
	.		.		
Cyno	GTGCCAGCGTGGGCAGCTCTTCACCGATGGGGATCATTGTGGCTGTGGT				
	660	670	680	690	700
Chimp	CATTGCGACTGCTGTAGCAGCCATTGTTGCTGCTGTAGTGGCCTTGATCT				
Human	CATTGCGACTGCTGTAGCAGCCATTGTTGCTGCTGTAGTGGCCTTGATCT				
	. . .	.			
Cyno	CACTGGGATTGCTGTAGCGGCCATTGTTGCTGCTGTAGTGGCCTTGATCT				
	710	720	730	740	750
Chimp	ACTGCAGGAAAAAGCGGATTTCAGCCAATTCCACTGATCCTGTGAAGGCT				
Human	ACTGCAGGAAAAAGCGGATTTCAGCCAATTCCACTGATCCTGTGAAGGCT				
Cyno	ACTGCAGGAAAAAGCGGATTTCAGCCAATTCCACTGATCCTGTGAAGGCT				
	760	770	780	790	800
Chimp	GCCCAATTTGAGCCACCTGGACGTCAAATGATTGCCATCAGAAAGAGACA				

TABLE 5-continued

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA DNA					
Human/Cyno 878 matches in an overlap of 933: 94.1% identity without one gap of three nucleotides					
878 matches in an overlap of 936: 93.8% identity with one gap of three nucleotides					
Human/Chimp 924 matches in an overlap of 933: 99.0% identity without one gap of three nucleotides					
924 matches in an overlap of 936: 98.7% identity with one gap of three nucleotides					
Human	GCCCAATTTGAGCCACCTGGACGTCAAATGATTGCCATCAGAAAGAGACA				
	. . . .				
Cyno	GCCCCGATTTGAGCCACTTGGACGTCAAACGATTGCCCTCAGAAAGAGACA				
	810	820	830	840	850
Chimp	ACTTGAAGAAACCAACAATGACTATGAAACAGCTGACGGCGGCTACATGA				
Human	ACTTGAAGAAACCAACAATGACTATGAAACAGCTGACGGCGGCTACATGA				
	.				
Cyno	ACTTGAAGAAACCAACAATGACTATGAAACAGCCGACGGCGGCTACATGA				
	860	870	880	890	900
Chimp	CTCTGAACCCAGGGCACCTACTGACGATGATAAAACATCTACCTGACT				
Human	CTCTGAACCCAGGGCACCTACTGACGATGATAAAACATCTACCTGACT				
	. .				
Cyno	CTCTGAACCCAGGGCACCTACTGATGATGATAGAAACATCTACCTGACT				
	910	920	930		
Chimp	CTTCCTCCCAACGACCATGTCAACAGTAATAACTAA				
Human	CTTCCTCCCAACGACCATGTCAACAGTAATAACTAA				
	. . .				
Cyno	CTTCTCCTCCCAACGACTATGACAACAGTAATAACTAA				

[0168] The sequence for the human FcγRIIA receptor has GenBank Accession No. M28697. Seki, T., *Identification of multiple isoforms of the low-affinity human IgG Fc receptor*, Immunogenetics 30, 5-12 (1989).

[0169] Alignment of the nucleic acid sequences encoding human (SEQ ID NO: 6) and cynomolgus (SEQ ID NO: 5) FcγRIIB is shown in Table 6.

[0170] Analysis of the % sequence identity shows that the human and cynomolgus sequences encoding FcγRIIB have about 94% identity.

TABLE 6

Alignment of Human and Cynomolgus Low-Affinity FcγRIIB DNA					
837 matches out of 885: 94.6% identity (without gap)					
837 matches out of 894: 93.6% identity (with gap)					
	10	20	30	40	50
Human	ATGGGAATCCTGTCATTCTTACCTGTCCTTGCCACTGAGAGTGACTGGGC				
	.				



TABLE 6-continued

Alignment of Human and Cynomolgus Low-Affinity FcγRIIB DNA 837 matches out of 885: 94.6% identity (without gap) 837 matches out of 894: 93.6% identity (with gap)				
Cyno	ATGGGAATCCTGTCATCTTACCTGTCCTTGCTACTGAGAGTGACTGGGC			
	60	70	80	90 100
Human	TGACTGCAAGTCCCCCAGCCTTGGGGTCATATGCTTCTGTGGACAGCTG			
		.	.	.
Cyno	TGACTGCAAGTCCTCCAGCCTTGGGGCCACATGCTTCTGTGGACAGCTG			
	110	120	130	140 150
Human	TGCTATTCTCGGCTCCTGTTGCTGGGACACCTGCAGCTCCCCCAAAGGCT			
				.
Cyno	TGCTATTCTCGGCTCCTGTTGCTGGGACACCTGCAGCTCCCCGAAGGCT			
	160	170	180	190 200
Human	GTGCTGAAACTCGAGCCCCAGTGGATCAACGTGCTCCAGGAGGACTCTGT			
		.		.
Cyno	GTGCTGAAACTCGAGCCCCGTGGATCAACGTGCTCCGGGAGGACTCTGT			
	210	220	230	240 250
Human	GACTCTGACATGCCGGGGGACTCACAGCCCTGAGAGCGACTCCATTCACT			
	.	.	.	.
Cyno	GACTCTGACGTGCGGGGGCGCTCACAGCCCTGACAGCGACTCCACTCAGT			
	260	270	280	290 300
Human	GGTTCCACAATGGGAATCTCATTCACCCACACGCAGCCAGCTACAGG			
		.		
Cyno	GGTTCCACAATGGGAATCTCATTCACCCACACGCAGCCAGCTACAGG			
	310	320	330	340 350
Human	TTCAAGGCCAACAACAATGACAGCGGGGAGTACACGTGCCAGACTGGCCA			
		.	.	.
Cyno	TTCAAGGCCAACAACAATGATAGCGGGGAGTACAGGTGCCAGACTGGCCG			
	360	370	380	390 400
Human	GACCAGCCTCAGCGACCCTGTGCATCTGACTGTGCTTCTGAGTGGCTGG			
		.		
Cyno	GACCAGCCTCAGCGACCCTGTTATCTGACTGTGCTTCTGAGTGGCTGG			
	410	420	430	440 450
Human	TGCTCCAGACCCCTCACCTGGAGTTCCAGGAGGGAGAAACCATCGTGCTG			
	.		.	.
Cyno	CGCTCCAGACCCCTCACCTGGAGTTCCGGGAGGGAGAAACCATCTTGCTG			
	460	470	480	490 500
Human	AGGTGCCACAGCTGGAAGGACAAGCCTCTGGTCAAGGTACATTCTTCCA			
		.		

TABLE 6-continued

Alignment of Human and Cynomolgus Low-Affinity FcγRIIB DNA 837 matches out of 885: 94.6% identity (without gap) 837 matches out of 894: 93.6% identity (with gap)				
Cyno	AGGTGCCACAGCTGGAAGGACAAGCCTCTGATCAAGGTCACATTCTTCCA			
	510	520	530	540 550
Human	GAATGGAAAATCCAAGAAATTTTCCCGTTCGGATCCCAACTTCTCCATCC			
	.	.	..	.
Cyno	GAATGGAAATATCCAAGAAATTTTCCCATATGAATCCCAACTTCTCCATCC			
	560	570	580	590 600
Human	CACAAGCAAACCCACAGTCACAGTGGTGATTACCACTGCACAGGAAACATA			
Cyno	CACAAGCAAACCCACAGTCACAGTGGTGATTACCACTGCACAGGAAACATA			
	610	620	630	640 650
Human	GGCTACACGCTGTACTCATCCAAGCCTGTGACCATCACTGTCCAAGCTCC			
	..	.		..
Cyno	GGCTACACACCATACTCATCCAACCTGTGACCATCACTGTCCAAGTGCC			
	660	670	680	690 700
Human	-----CAGCTCTTCACCGATGGGGATCATTGTGGCTGTGGTCACTG			
	*****	.		
Cyno	CAGCATGGGCAGCTCTTCACCGATAGGGATCATTGTGGCTGTGGTCACTG			
	710	720	730	740 750
Human	GGATTGCTGTAGCGGCCATTGTTGCTGCTGTAGTGGCCTTGATCTACTGC			
Cyno	GGATTGCTGTAGCGGCCATTGTTGCTGCTGTAGTGGCCTTGATCTACTGC			
	760	770	780	790 800
Human	AGGAAAAAGCGGATTTTCAGCCAATCCCACTAATCCTGATGAGGCTGACAA			
			.	
Cyno	AGGAAAAAGCGGATTTTCAGCCAATCCCACTAATCCTGACGAGGCTGACAA			
	810	820	830	840 850
Human	AGTTGGGGCTGAGAACACAATCACCTATTCACTTCTCATGCACCCGGATG			
			.	.
Cyno	AGTTGGGGCTGAGAACACAATCACCTATTCACTTCTCATGCATCCGGACG			
	860	870	880	
Human	CTCTGGAAGAGCCTGATGACCAGAACCGTATTTAG			
		.	.	
Cyno	CTCTGGAAGAGCCTGATGACCAAAACCGNGTTTAG			

[0171] The human sequence for FcγRIIB has GenBank Accession No. X52473. Engelhardt, W., Geerds, C. and Frey, J., *Distribution, inducibility and biological function of the cloned and expressed human beta Fc receptor II*, Eur. J. Immunol. 20 (6), 1367-1377 (1990)

[0172] Alignment of the nucleic acid sequences encoding a human (SEQ ID NO: 8) and cynomolgus (SEQ ID NO: 7) FcγRIIIA is shown in Table 7.

[0173] Analysis of the % sequence identity shows that the human and cynomolgus nucleic acid sequences encoding FcγRIIIA have about 96% identity.

TABLE 7

Alignment of Human and Cynomolgus Low-Affinity FcγRIIIA DNA 733 matches in an overlap of 765: 95.8% identity					
	10	20	30	40	50
Human	ATGTGGCAGCTGCTCCTCCCAACTGCTCTGCTACTTCTAGTTTCAGCTGG				
Cyno	ATGTGGCAGCTGCTCCTCCCAACTGCTCTGCTACTTCTAGTTTCAGCTGG				
	60	70	80	90	100
Human	CATGCGGACTGAAGATCTCCCAAGGCTGTGGTGTTCCTGGAGCCTCAAT				
	.				
Cyno	CATGCGGGCTGAAGATCTCCCAAGGCTGTGGTGTTCCTGGAGCCTCAAT				
	110	120	130	140	150
Human	GGTACAGGGTGCTCGAGAAGGACAGTGTGACTCTGAAGTGCCAGGGAGCC				
	.				
Cyno	GGTACAGGGTGCTCGAGAAGGACCGTGTGACTCTGAAGTGCCAGGGAGCC				
	160	170	180	190	200
Human	TACTCCCCTGAGGACAATTCCACACAGTGGTTTCACAATGAGAGCCTCAT				
	.				
Cyno	TACTCCCCTGAGGACAATTCCACACGGTGGTTTCACAATGAGAGCCTCAT				
	210	220	230	240	250
Human	CTCAAGCCAGGCCTCGAGCTACTTCATTGACGCTGCCACAGTCGACGACA				
	.		..	.	. .
Cyno	CTCAAGCCAGACCTCGAGCTACTTCATTGCTGCTGCCAGAGTCAACAACA				
	260	270	280	290	300
Human	GTGGAGAGTACAGGTGCCAGACAAACCTCTCCACCCTCAGTGACCCGGTG				
	. .				
Cyno	GTGGAGAGTACAGGTGCCAGACAAGCCTCTCCACACTCAGTGACCCGGTG				
	310	320	330	340	350
Human	CAGCTAGAAGTCCATATCGGCTGGCTATTGCTCCAGGCCCTCGGTGGGT				
	.		.		
Cyno	CAGCTGGAAGTCCATATCGGCTGGCTATTGCTCCAGGCCCTCGGTGGGT				
	360	370	380	390	400
Human	GTTCAAGGAGGAAGACCCTATTACCTGAGGTGTCACAGCTGGAAGAACA				
	..				
Cyno	GTTCAAGGAGGAAGAATCTATTACCTGAGGTGTCACAGCTGGAAGAACA				
	410	420	430	440	450
Human	CTGCTCTGCATAAGGTCACATATTTACAGAATGGCAAAGGCAGGAAGTAT				
	..	.			
Cyno	CTCTTCTGCATAAGGTCACGTATTTACAGAATGGCAAAGGCAGGAAGTAT				
	460	470	480	490	500

TABLE 7-continued					
Alignment of Human and Cynomolgus Low-Affinity FcγRIIIA DNA 733 matches in an overlap of 765: 95.8% identity					
Human	TTTCATCATAATTCTGACTTCTACATTCCAAAAGCCACACTCAAAGACAG				
	.				
Cyno	TTTCATCAGAATTCTGACTTCTACATTCCAAAAGCCACACTCAAAGACAG				
	510	520	530	540	550
Human	CGGCTCCTACTTCTGCAGGGGGCTTTTGGGAGTAAAAATGTGTCTTCAG				
		.	.		.
Cyno	CGGCTCCTACTTCTGCAGGGGACTTATTTGGGAGTAAAAATGTATCTTCAG				
	560	570	580	590	600
Human	AGACTGTGAACATCACCATCACTCAAGGTTTGGCAGTGTCAACCATCTCA				
		.		.	
Cyno	AGACTGTGAACATCACCATCACTCAAGATTGGCAGTGTCAATCATCTCA				
	610	620	630	640	650
Human	TCATTCTTTCCACCTGGGTACCAAGTCTCTTCTGCTTGGTGATGGTACT				
			.		
Cyno	TCATTCTTTCCACCTGGGTACCAAGTCTCTTCTGCTTGGTGATGGTACT				
	660	670	680	690	700
Human	CCTTTTTCAGTGGACACAGGACTATATTTCTCTGTGAAGACAAACATTC				
			.	.	.
Cyno	CCTTTTTCAGTGGACACAGGACTATATTTCTCTATGAAGAAAAGCATTC				
	710	720	730	740	750
Human	GAAGCTCAACAAGAGACTGGAAGGACCATAAATTAAATGGAGAAAGGAC				
	.	.	.		.
Cyno	CAAGCTCAACAAGGACTGGGAGGACCATAAATTAAATGGAGCAAGGAC				
	760				
Human	CCTCAAGACAAATGA				
Cyno	CCTCAAGACAAATGA				

[0174] The human sequence for FcγIII has GenBank Accession No. X52645 M31937). Ravetch, J. V. and Perussia, B., *Alternative membrane forms of Fc gamma RII-I(CD16) on human natural killer cells and neutrophils. Cell type-specific expression of two genes that differ in single nucleotide substitutions*, J. Exp. Med. 170 (2), 481-497 (1989).

[0175] Alignment of the nucleic acid sequences encoding a human (SEQ ID NO: 24) and cynomolgus (SEQ ID NO: 23) β-2 microglobulin is shown in Table 8.

[0176] Analysis of the % sequence identity shows that the human and cynomolgus nucleic acid sequences encoding β-2 microglobulin have about 95% identity.

TABLE 8					
Alignment of Human and Cynomolgus β2-Microglobulin DNA 341/360 = 94.7% identity					
	10	20	30	40	50
Human	ATGTCTCGCTCCGTGGCCTTAGCTGTGCTCGCGCTACTCTCTTTCTGG				
	.	.	.	.	

TABLE 8-continued

Alignment of Human and Cynomolgus $\beta$ 2-Microglobulin DNA 341/360 = 94.7% identity	
Cyno	ATGTCTCCCTCAGTGGCCTTAGCCGTGCTGGCGCTACTCTCTCTTTCTGG
	60 70 80 90 100
Human	CCTGGAGGCTATCCAGCGTACTCCAAAGATTAGGTTTACTCACGTCATC
	.
Cyno	CCTGGAGGCTATCCAGCGTACTCCAAAGATTAGGTTTACTCACGCCATC
	110 120 130 140 150
Human	CAGCAGAGAATGGAAAGTCAAATTTCTGAATTGCTATGTGTCTGGGTTT
	. . .
Cyno	CACCAGAGAATGGAAAGCCAAATTTCTGAATTGCTATGTGTCTGGATTT
	160 170 180 190 200
Human	CATCCATCCGACATTGAAGTTGACTTACTGAAGAATGGAGAGAGAATTGA
	. . . . .
Cyno	CATCCATCTGATATTGAAGTTGACTTACTGAAGAATGGAGAGAAAATGGG
	210 220 230 240 250
Human	AAAAGTGGAGCATTCAGACTTGTCTTTTCAGCAAGGACTGGTCTTTCTATC
	.
Cyno	AAAAGTGGAGCATTCAGACTTGTCTTTTCAGCAAGGACTGGTCTTTCTATC
	260 270 280 290 300
Human	TCTTGTAAGTACACTGAATTCACCCCACTGAAAAAGATGAGTATGCCTGC
	.
Cyno	TCTTGTAAGTACACTGAATTCACCCCACTGAAAAAGATGAGTATGCCTGC
	310 320 330 340 350
Human	CGTGTGAACCATGTGACTTTGTACAGCCCAAGATAGTTAAGTGGGATCG
	. . . . .
Cyno	CGTGTGAACCATGTGACTTTGTACAGGCCCAAGATAGTTAAGTGGGATCG
	360
Human	AGACATGTAA
Cyno	AGACATGTAA

[0177] The DNA sequence for the human  $\beta$ -2 microglobulin has GenBank Accession No. ABO21288. Matsumoto, K., Minamitani, T., *Human mRNA for beta 2-microglobulin*, DDBJ/EMBL/GenBank databases (1998).

[0178] Alignment of the nucleic acid sequences encoding a human (SEQ ID NO: 28) and cynomolgus (SEQ ID NO: 27) FcRn  $\alpha$ -chain is shown in Table 9.

[0179] Analysis of the % sequence identity shows that the human and cynomolgus nucleic acid sequences encoding FcRn  $\alpha$ -chain have about 97% identity.

TABLE 9

Alignment of Human and Cynomolgus FcRn $\alpha$ -Chain DNA 1062/1098 = 96.7% identity					
	10	20	30	40	50
Human	ATGGGGGTCCCGCGCCTCAGCCCTGGGCGCTGGGGCTCCTGCTCTTTCT				
	.				
Cyno	ATGAGGGTCCCGCGCCTCAGCCCTGGGCGCTGGGGCTCCTGCTCTTTCT				
	60	70	80	90	100
Human	CCTTCCTGGGAGCCTGGGCGCAGAAAGCCACCTCTCCCTCCTGTACCACC				
	. .				
Cyno	CCTGCCCCGGGAGCCTGGGCGCAGAAAGCCACCTCTCCCTCCTGTACCACC				
	110	120	130	140	150
Human	TTACCGCGGTGTCTCGCCTGCCCCGGGACTCCTGCCTTCTGGGTGTCC				
	. . .				
Cyno	TCACCGCGGTGTCTCGCCCCGGGACGCCTGCCTTCTGGGTGTCC				
	160	170	180	190	200
Human	GGCTGGCTGGGCCCGCAGCAGTACCTGAGCTACAATAGCCTGCGGGCGA				
	. . . .				
Cyno	GGCTGGCTGGGCCCGCAGCAGTACCTGAGCTACGACAGCCTGAGGGCCA				
	210	220	230	240	250
Human	GGCGGAGCCCTGTGGAGCTTGGGTCTGGGAAAACAGGTGTCCTGGTATT				
	.				
Cyno	GGCGGAGCCCTGTGGAGCTTGGGTCTGGGAAAACCAAGTGTCTGGTATT				
	260	270	280	290	300
Human	GGGAGAAAGAGACCACAGATCTGAGGATCAAGGAGAAGCTCTTTCTGGAA				
Cyno	GGGAGAAAGAGACCACAGATCTGAGGATCAAGGAGAAGCTCTTTCTGGAA				
	310	320	330	340	350
Human	GCTTTCAAAGCTTTGGGGGAAAAGTCCCTACACTCTGCAGGGCCTGCT				
	.				
Cyno	GCTTTCAAAGCTTTGGGGGAAAAGGCCCTACACTCTGCAGGGCCTGCT				
	360	370	380	390	400
Human	GGGCTGTGAACCTGGGCCCTGACAACACCTCGGTGCCACCGCCAAGTTCG				
	.				
Cyno	GGGCTGTGAACCTGAGCCCTGACAACACCTCGGTGCCACCGCCAAGTTCG				
	410	420	430	440	450
Human	CCCTGAACGGCGAGGAGTTCATGAATTTCGACCTCAAGCAGGGCACCTGG				
Cyno	CCCTGAACGGCGAGGAGTTCATGAATTTCGACCTCAAGCAGGGCACCTGG				
	460	470	480	490	500
Human	GGTGGGGACTGGCCCCGAGGCCCTGGCTATCAGTCAGCGGTGGCAGCAGCA				

TABLE 9-continued

Alignment of Human and Cynomolgus FcRn $\alpha$ -Chain DNA 1062/1098 = 96.7% identity	
Cyno	GGTGGGGACTGGCCCGAGGCCCTGGCTATCAGTCAGCGGTGGCAGCAGCA
	510 520 530 540 550
Human	GGACAAGGCGGCCAACAAGGAGCTCACCTTCCTGCTATTCTCCTGCCCGC
	•
Cyno	GGACAAGGCGGCCAACAAGGAGCTCACCTTCCTGCTATTCTCCTGCCCAC
	560 570 580 590 600
Human	ACCGCCTGCGGGAGCACCTGGAGAGGGGCCGCGAAACCTGGAGTGAAG
	• •
Cyno	ACCGGCTGCGGGAGCACCTGGAGAGGGCCGTGAAACCTGGAGTGAAG
	610 620 630 640 650
Human	GAGCCCCCTCCATGCGCCTGAAGGCCCGACCCAGCAGCCCTGGCTTTTC
	• •
Cyno	GAGCCCCCTCCATGCGCCTGAAGGCCCGACCCGGCAACCTGGCTTTTC
	660 670 680 690 700
Human	CGTGCTTACCTGCAGCGCCTTCCTTCTACCCCTCCGGAGCTGCAACTTC
	• •
Cyno	CGTGCTTACCTGCAGCGCCTTCCTTCTACCCCTCCGGAACCTGCAACTGC
	710 720 730 740 750
Human	GGTTCCTGCGGAATGGGCTGGCCGCTGGCACCGGCCAGGGTGACTTCGGC
	• • •
Cyno	GGTTCCTGCGGAATGGGATGGCCGCTGGCACCGGACAGGGCGACTTCGGC
	760 770 780 790 800
Human	CCCAACAGTGACGGATCCTTCCACGCCTCGTCGTCATAACAGTCAAAAG
	•
Cyno	CCCAACAGTGACGGCTCCTTCCACGCCTCGTCGTCATAACAGTCAAAAG
	810 820 830 840 850
Human	TGGCGATGAGCACCCTACTGCTGCATTGTGACGACGCGGGGCTGGCGC
	•
Cyno	TGGCGATGAGCACCCTACTGCTGCATCGTGACGACGCGGGGCTGGCGC
	860 870 880 890 900
Human	AGCCCCTCAGGGTGGAGCTGGAATCTCCAGCCAAGTCCTCCGTGCTCGTG
	• •
Cyno	AGCCCCTCAGGGTGGAGCTGGAACCTCCAGCCAAGTCCTCCGTGCTCGTG
	910 920 930 940 950
Human	GTGGGAATCGTCATCGGTGTCTTGCTACTCAGGCAGCGGCTGTAGGAGG
Cyno	GTGGGAATCGTCATCGGTGTCTTGCTACTCAGGCAGCGGCTGTAGGAGG

TABLE 9-continued

Alignment of Human and Cynomolgus FcRn α-Chain DNA 1062/1098 = 96.7% identity					
	960	970	980	990	1000
Human	AGCTCTGTTGTGGAGAAGGATGAGGAGTGGGCTGCCAGCCCCTTGGATCT				
Cyno	AGCTCTGTTGTGGAGAAGGATGAGGAGTGGGCTGCCAGCCCCTTGGATCT				
	1010	1020	1030	1040	1050
Human	CCCTTCGTGGAGACGACACCGGGTCCTCCTGCCACCCCAGGGGAGGCC				
	.	.	..	.	
Cyno	CCCTCCGTGGAGATGACACCGGGTCCTCCTGCCACCCCAGGGGAGGCC				
	1060	1070	1080	1090	
Human	CAGGATGCTGATTTGAAGGATGTAAATGTGATTCCAGCCACCGCCTGA				
		.	.	.	.
Cyno	CAGGATGCTGATTGGAAGGATATAAATGTGATCCAGCCACTGCCTGA				

[0180] The DNA sequence for the human FcRn α-chain has GenBank Accession No. U12255. Story, C. M., Mikulska, J., and Simister, N. E., *A major histocompatibility complex class I-like Fc receptor cloned from human placenta: Possible role in transfer of immunoglobulin G from mother to fetus*, J. Exp. Med. 180, 2377-2381 (1994).

[0181] An alignment of the amino acid sequences for human (SEQ ID NO: 10) and cynomolgus (SEQ ID NO: 9) FcγRI α-chain is shown in Table 10. As described previously, the α-chain of FcγRI has various domains, including a signal peptide, three extracellular C-2 Ig like domains, a transmembrane domain and an intracellular domain. The amino acid numbers shown below the amino acids with the symbol Δ are numbered from the start of the mature polypeptide not including the signal sequence. Based on the alignment with the human sequence, the mature cynomolgus

FcγRI has an amino acid sequences of residues Δ1 to Δ336 (SEQ ID NO: 65). The n-terminal sequence of cynomolgus sequences FcγRI may vary from that shown below. It would be within the skill in the art to express the nucleic acid sequence encoding the cynomolgus FcγRI sequence and identify the n-terminal sequence. An extracellular fragment of cynolomolgus FcγRI obtained using the primers of example 1 has an amino acid sequence of Δ1 to Δ269. Any numbers above the amino acid residues represent the numbering of the residues starting at the signal sequence.

[0182] Analysis of the % sequence identity shows that the amino acid sequences for human and cynomolgus FcγRI have about 90% identity when the 3' extension is taken into account and about 94% when the 3' extension is not included.

TABLE 10

Alignment of Human and Cynomolgus High-Affinity FcγRI					
Human	MWFLTTLLLWVPVDGQVDTTK				
	.				
Cyno	MWFLTALLLWVPVDGQVDTTK				
Domain 1					
Human	AVISLQPPWVSVFQEETVTLHCEVLHLPGSSSTQWFLNGTAT				
	.	.	..		
Cyno	AVITLQPPWVSVFQEETVTLQCEVPRLPGSSSTQWFLNGTAT				
	Δ	Δ	Δ	Δ	Δ
	1	10	20	30	40
		70	80	90	100
Human	QTSTPSYRITSASVNSGGEYRCQRLSGRSDPIQLEIHR				
	.	.			



TABLE 10-continued

Alignment of Human and Cynomolgus High-Affinity FcγRI				
Cyno	QTSTPSYRITSASVKDSGEYRCQRGPSGRSDPIQLEIHR			
	Δ	Δ	Δ	Δ
	50	60	70	80
Domain 2				
Human	GWLLQLQVSSRVFTEGEPLALRCHAWKDKLVYNVLYYRNGKAFKF			
	.		.	
Cyno	DWLLQLQVSSRVFTEGEPLALRCHAWKDKLVYNVLYYQNGKAFKF			
	Δ	Δ	Δ	Δ
	90	100	110	120
	150	160	170	180
Human	FHWNSNLTILKTNISHNGTYHCSGMGKHRYTSAGISVTVKELFP			
	..	.	.	.
Cyno	FYRNSQLTILKTNISHNGAYHCSGMGKHRYTSAGVSVTVKELFP			
	Δ	Δ	Δ	Δ
	130	140	150	160
Domain 3				
Human	APVLNASVTSPLLEGNLVTLSCETKLLQRPGLQLYFSFYMGSKTLRG			
Cyno	APVLNASVTSPLLEGNLVTLSCETKLLQRPGLQLYFSFYMGSKTLRG			
	Δ	Δ	Δ	Δ
	170	180	190	200
Human	RNTSSEYQILTARREDSGLYWCEAATEDGNVLKRSPLELQVLGLQLP			
	.	.		
Cyno	RNTSSEYQILTARREDSGFYWCEATTEDGNVLKRSPLELQVLGLQLP			
	Δ	Δ	Δ	Δ
	220	230	240	250
transmembrane/intracellular				
Human	TPVWFHVLFFYLAVGIMFLVNTVLWVTIRKELKRKKKWDLEISLDSGHE			
	.	.	.	.
Cyno	TPVWLHVLFFYLVGIMFLVNTVLWVTIRKELKRKKKWNLEISLDSAHE			
	Δ	Δ	Δ	Δ
	270	280	290	300
Human	KKVTSSLQEDRHLEELKCQEQKEQLQEGVHRKEPQGAT			
	.	.		
Cyno	KKVTSSLQEDRHLEELKSQEQE			
	Δ	Δ	Δ	Δ
	320	330	340	350

Human vs Cyno 335/357 = 93.8% identity without human 3' extension  
 335/374 = 89.6% identity with human 3' extension

[0183] The amino acid sequence for human FcγRI has Accession Nos.: P112314; P12315; EMBL; X14356; CAA32537.1. EMBL; X14355; CAA32536.1. PIR; S03018. PIR; S03019. PIR; A41357. PIR; B41357. HSSP; P12319; 1ALT. MIM; 146760; -. InterPro; IPR003006; -. Pfam; PF00047; Allen J. M., Seed B., Nucleic Acids Res. 16, 11824-11824, 1988, *Nucleotide sequence of three cDNAs for the human high affinity Fc receptor (FcRI)*; Allen J. M., Seed B., Science 243, 378-381, 1989, *Isolation and expression of functional high-affinity Fc receptor complementary DNAs*.

[0184] An alignment of amino acid sequences for human, cynomolgus, and chimp sequences for FcγRIIA (cynomolgus/SEQ ID NO: 15; human/SEQ ID NO: 16; chimp/SEQ ID NO: 17), FcγRIIB (cynomolgus/SEQ ID NO: 18; human/SEQ ID NO: 19), and FcγRIIIA (cynomolgus/SEQ ID NO: 20; human/SEQ ID NO: 21) is shown in Table 11.

[0185] The sequence is divided into domains as described previously: signal peptide, 3 extracellular C-2 like domains, and a transmembrane intracellular domain. In Table 11, the amino acid numbers shown below the amino acids with the symbol A are numbered from the start of the mature human polypeptide not including the signal sequence. The mature polypeptides for cynomolgus and chimp FcγRIIA, cynomolgus FcγRIIB, and cynomolgus FcγRIIIA start at the amino acid identified with the asterisk in Table 11 and are separately shown in Tables 21, 22, and 23, and are as follows:

[0186] 1) cynomolgus FcγRIIA amino acids Δ1 to Δ282 (SEQ ID NO: 66), N terminal sequence TAP- PKA (Table 21);

[0187] 2) chimp FcγRIIA amino Δ1 to Δ249 (SEQ ID NO: 67)(based on alignment with the human sequence);

[0188] 3) cynomolgus FcγRIIB amino acids Δ1 to Δ252 (SEQ ID NO: 68), N terminal sequence TPAAPP (table 22); and

[0189] 4) cynomolgus FcγRIIIA amino acids Δ1 to Δ234 (SEQ ID NO: 69), N terminal sequence EDLPKA (table 23).

[0190] In table 11, any numbers above the amino acid residues represent the numbering of the residues starting at the signal sequence. The asterisks in the table indicate the start of the n-terminal sequence for cynomolgus FcγRIIA, FcγRIIB, and FcγRIIIA.

[0191] Extracellular fragments of the Fc receptor polypeptides were obtained using the primers described in example 1. An extracellular fragment of FcγRIIA obtained using the primers of example 1 has an amino acid sequence of Δ1 to Δ182, as shown in table 21. An extracellular fragment of FcγRIIB obtained using the primers of example 1 has an amino acid sequence of Δ1 to Δ184, as shown in Table 22. An extracellular fragment of FcγRIIIA obtained using the primers of example 1 has an amino acid sequence of Δ1 to Δ187, as shown in Table 23.

[0192] Analysis of the % sequence identity shows the following:

[0193] 1) Chimp and human amino acid sequences for FcγRIIA have about 97% identity;

[0194] 2) Cynomolgus and human amino acid sequences for FcγRIIA have about 87% identity with MAMETQ (possible portion of signal peptide) and 89% identity without MAMETQ in the alignment;

[0195] 3) Cynomolgus and chimp amino acid sequences for FcγRIIA have about 87% identity including MAMETQ in the alignment and 89% without MAMETQ in the alignment;

[0196] 4) Cynomolgus and human amino acid sequences for FcγRIIB have about 92% identity; and

[0197] 5) Cynomolgus and human amino acid sequences for FcγRIIIA have about 91% identity.

TABLE 11

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA, FcγRIIB, FcγRIIIA

signal peptide	.....	.	..
IIA-human	-----MAMETQMSQNVCPRNLLWLLQPLTVLLLLASADSQAA		
IIA- chimp	-----MAMETQMSQNVCPRNLLWLLQPLTVLLLLASADSQA-		
IIA-cyno	-----MSQNVCPGNLWLLQPLTVLLLLASADSQT-		*
		.	
IIB-human	MGILSFLPVLATESDWADCKSPQPGHMLLWTAVLFLAPVAGTPA		
IIB-cyno	MGILSFLPVLATESDWADCKSSQPGHMLLWTAVLFLAPVAGTPA		*
		.	
IIIA-human			MWQLLLPTALLLVLSAGMRTE

TABLE 11-continued

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA,  
FcγRIIB, FcγRIIIA

IIIA-cyno	MWQLLLPTALLLLVSAGMRAE			
				Δ *
				1
Domain 1				
		.	.	.
IIA-human	APPKAVLKLEPPWINVLQEDSVTLTCQGARSPESDSIQWFHN			
IIA-chimp	APPKAVLKLEPPWINVLQEDSVTLTCRGARSPESDSIQWFHN			
IIA-cyno	APPKAVLKLEPPWINVLREDSVTLTCGGAHSPDSDSTQWFHN			
	Δ	Δ	Δ	Δ
	1	10	20	30
		.	.	.
IIB-human	APPKAVLKLEPPWINVLQEDSVTLTCRGTHSPESDSIQWFHN			
IIB-cyno	APPKAVLKLEPPWINVLREDSVTLTCGGAHSPDSDSTQWFHN			
		.		.
IIIA-human	DLPKAVVFLEPQWYRVLEKDSVTLKCQGAYSPEDNSTQWFHN			
IIIA-cyno	DLPKAVVFLEPQWYRVLEKDRVTLKCQGAYSPEDNSTRWFNH			
	Δ	Δ	Δ	Δ
	10	20	30	40
	.		.	.
IIA-human	GNLIPHTHTQPSYRFKANNNDSGEYTCQTGQTSLSDPVHLTVLSE			
IIA-chimp	GNLIPHTHTQPSYRFKANNNDSGEYTCQTGQTSLSDPVHLTVLSE			
IIA-cyno	GNRIPHTHTQPSYRFKANNNDSGEYRCQTRTSLSDPVHLTVLSE			
	Δ	Δ	Δ	Δ
	50	60	70	80
		.	.	
IIB-human	GNLIPHTHTQPSYRFKANNNDSGEYTCQTGQTSLSDPVHLTVLSE			
IIB-cyno	GNLIPHTHTQPSYRFKANNNDSGEYRCQTRTSLSDPVHLTVLSE			
	.	.	.	.
IIIA-human	ESLISSQASSYFIDAATVDDSGEYRCQTNLSTLSDPVQLEVHIG			
IIIA-cyno	ESLISSQTSSYFIAAARVNNSGEYRCQTSLSLSDPVQLEVHIG			
	Δ	Δ	Δ	Δ
	50	60	70	80
Domain 2		.	.	.
	.	.	.	.
IIA-human	WVLVQTPHLEFQEGETIMLRCHSWKDKPLVKVTFQNGKSQKFS			
IIA-chimp	WVLVQTPHLEFQEGETIVLRCHSWKDKPLVKVTFQNGKSQKFS			

TABLE 11-continued

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA,  
FcγRIIB, FcγRIIIA

IIA-cyno	WLALQTPHLEFREGETIMLRCHSWKDKPLIKVTFQNGIAKKFS				
	Δ	Δ	Δ	Δ	Δ
	90	100	110	120	130
	.	.	.	.	.
IIB-human	WVLQTPHLEFQEGETIVLRCHSWKDKPLVKVTFQNGKSKKFS				
IIB-cyno	WLALQTPHLEFREGETILLRCHSWKDKPLIKVTFQNGISKKFS				
		..	.		
IIIA-human	WLLQAPRWVFKEEDPIHLRCHSWKNTALHKVTYLQNGKGRKYF				
IIIA-cyno	WLLQAPRWVFKEEESIHLRCHSWKNTLLHKVTYLQNGKGRKYF				
	Δ	Δ	Δ	Δ	Δ
	90	100	110	120	130
	..	..		..	.
IIA-human	RLDPTFSIPQANHSHSGDYHCTGNIGYTLFSSKPVITITVQV				
IIA-chimp	HLDPNLSIPQANHSHSGDYHCTGNIGYTLFSSKPVITITVQA				
IIA-cyno	HMDPNFSIPQANHSHSGDYHCTGNIGYTPYSSKPVITITVQV				
	Δ	Δ	Δ	Δ	Δ
	131	140	150	160	170
	...			.	.
IIB-human	RSDPNFSIPQANHSHSGDYHCTGNIGYTLYSSKPVITITVQA				
IIB-cyno	HMNPHFSIPQANHSHSGDYHCTGNIGYTPYSSKPVITITVQV				
	.		.		
IIIA-human	HHNSDFYIPKATLKDSGSYFCRGLFGSKNVSETVNITITQ				
IIIA-cyno	HQNSDFYIPKATLKDSGSYFCRGLIGSKNVSETVNITITQ				
	Δ	Δ	Δ	Δ	
	140	150	158	170	
transmembrane/intracellular	.	.	...		
IIA-human	PSMGSSSPMGIIVAVVIATAVAIAVAIVVALIYCRKKRISANSTD				
IIA-chimp	PSVGSSSPVGIIVAVVIATAVAIAVAIVVALIYCRKKRISANSTD				
IIA-cyno	PSVGSSSPMGIIVAVVTGIATAVAIAVAIVVALIYCRKKRISANSTD				
	Δ	Δ	Δ	Δ	
	180	190	200	210	
	...	.			
IIB-human	P---SSSPMGIIVAVVTGIATAVAIAVAIVVALIYCRKKRISANPTN				
IIB-cyno	PSMGSSSPIGIIVAVVTGIATAVAIAVAIVVALIYCRKKRISANPTN				
	.	.		.	..
IIIA-human	GLAVSTISSFFPGYQVSFCLVMVLLFAVDTLGYFSVKTNIRSST				

TABLE 11-continued

Alignment of Human, Cynomolgus and Chimp Low-Affinity FcγRIIA,  
FcγRIIB, FcγRIIIA

IIIA-cyno	DLAVSSISSFFPPGYQVSFCLVMVLLFAVDLTGLYFSMKKSIPSSST				
	Δ	Δ	Δ	Δ	
	180	190	200	210	
	• • • •			ITAM motif	
IIA-human	PVKAAQFEPPGRQMIAIRKRQLEETNNDYETADGGYMTLNPRAPT				
IIA-chimp	PVKAAQFEPPGRQMIAIRKRQLEETNNDYETADGGYMTLNPRAPT				
IIA-cyno	PVKAARFEPLGRQTIALRKRQLEETNNDYETADGGYMTLNPRAPT				
	Δ	Δ	Δ	Δ	Δ
	220	230	240	250	260
				•	
IIB-human	PDEADKVGAEENTITYSLLMHPDALEEPDDQNRI				
IIB-cyno	PDEADKVGAEENTITYSLLMHPDALEEPDDQNRV				
		ITIM motif			
	• •				
IIIA-human	RDWKDHKFKWRKDPQDK				
IIIA-cyno	RDWEDHKFKWSKDPQDK				
	Δ	Δ			
	220	230			
		ITAM motif			
	• • •				
IIA-human	DDDKNIYLTLPNDHVNSNN				
IIA-chimp	DDDKNIYLTLPNDHVNSNN				
IIA-cyno	DDDRNIYLTLSFNDYDNSNN				
	Δ	Δ			
	270	280			

IIA chimp/human 308/317 = 97.2% identity  
 cyno/human 277/317 = 87.4% identity (+MAMETQ)  
 277/311 = 89.1% identity (-MAMETQ)  
 cyno/chimp 276/316 = 87.3% identity (+MAMETQ)  
 276/310 = 89.0% identity (-MAMETQ)  
 IIB cyno/human 270/294 = 91.8% identity  
 IIIA cyno/human 232/254 = 91.3% identity

[0198] The human amino acid sequence for FcRIIA has the following Accession Nos.: P12318; EMBL; M31932; AAA35827.1. EMBL; Y00644; CAA68672.1. EMBL; J03619; AAA35932.1. EMBL; A21604; CAA01563.1. PIR; A31932. PIR; J10118. PIR; S02297. PIR; S00477. PIR; S06946. HSSP; P12319; 1ALT. MIM; 146790; -. InterPro; IPR003006; -. Pfam; PF00047. Brooks D. G., Qiu W. Q., Luster A. D., Ravetch J. V., J. Exp. Med. 170, 1369-1385, 1989, *Structure and expression of human IgG FcRII(CD32). Functional heterogeneity is encoded by the alternatively spliced products of multiple genes*; Stuart S. G., Trounstein

M. L., Vaux D. J. T., Koch T., Martens C. L., Moore K. W., J. Exp. Med. 166, 1668-1684, 1987, *Isolation and expression of cDNA clones encoding a human receptor for IgG (Fc gamma RII)*; Hibbs M. L., Bonadonna L., Scott B. M., McKenzie I. F. C., Hogarth P. M., Proc. Natl. Acad. Sci. U.S.A. 85, 2240-2244, 1988, *Molecular cloning of a human immunoglobulin G Fc receptor*; Stengelin S., Stamenkovic I., Seed B., EMBO J. 7, 1053-1059, 1988, *Isolation of cDNAs for two distinct human Fc receptors by ligand affinity cloning*; Salmon J. E., Millard S., Schachter L. A., Arnett F. C., Ginzler E. M., Gourley M. F., Ramsey-Goldman R., Peterson M. G. E., Kimberly R. P., J. Clin. Invest. 97,

1348-1354, 1996, *Fc gamma RIIA alleles are heritable risk factors for lupus nephritis in African Americans*.

[0199] The human sequence for FcγRIIB has Accession No. X52473. Engelhardt, W., Geerds, C. and Frey, J., *Distribution, inducibility and biological function of the cloned and expressed human beta Fc receptor II*, Eur. J. Immunol. 20 (6), 1367-1377 (1990).

[0200] The human amino acid sequence for FcγRIIA has Accession Nos.: P08637; EMBL; X52645; CAA36870.1. EMBL; Z46222; CAA86295.1. PIR; JI.0107. MIM; 146740; -. InterPro; IPR003006; -. Pfam; PF00047; Ravetch J. V., Perussia B., J. Exp. Med. 170, 481497, 1989, *Alternative membrane forms of Fc gamma RIII(CD16) on human natural killer cells and neutrophils. Cell type-specific expression of two genes that differ in single nucleotide substitutions*; Gessner J. E., Grussenmeyer T., Kolanus W., Schmidt R. E., J. Biol. Chem. 270, 1350-1361, 1995, *The human low affinity immunoglobulin G Fc receptor III-A and III-B genes: Molecular characterization of the promoter regions*; de Haas M., Koene H. R., Kleijer M., de Vries E., Simsek S., van Tol M. J. D., Roos D., von dem Borne A. E. G. K., J. Immunol. 156, 3948-3955, 1996, *A triallelic Fc gamma receptor type IIIA polymorphism influences the binding of human IgG by NK cell Fc gamma RIIIA*; Koene H. R., Kleijer M., Algra J., Roos D., von dem Borne A. E. G. K., de Haas M., Blood 90, 1109-1114, 1997, *Fc gamma RIIIA-158V/F polymorphism influences the binding of IgG by natural killer cell Fc gamma RIIIA, independently of the Fc gamma RIIIA-48L/R/H phenotype*; Wu J., Edberg J. C., Redecha P. B., Bansal V., Guyre P. M., Coleman K., Salmon J. E., Kimberly R. P., J. Clin. Invest. 100, 1059-1070, 1997, *A novel polymorphism of Fc gamma RIIIA (CD16) alters receptor function and predisposes to autoimmune disease*.

TABLE 21

Sequence of Mature FcRIIA				
Domain 1				
TAPPKAVLKLEPPWINVLREDSVTLTTCGGAHSPDSSTQWFHN				
Δ	Δ	Δ	Δ	Δ
1	10	20	30	40
GNRIPTHQTQPSYRFKANNNDSGEYRCQTGRITSLSDPVHLTVLSE				
	Δ	Δ	Δ	Δ
	50	60	70	80
Domain 2				
WLALQTPHLEFREGETIMLRCHSWKDKPLIKVTFFQNGIAKKFS				
Δ	Δ	Δ	Δ	Δ
90	100	110	120	130
HMDPNFSIPQANSHSGDYHCTGNIGYTPYSSKPVTTITVQV				
	Δ	Δ	Δ	Δ
	140	150	160	170
Intracellular/transmembrane domain				
PSVGSSSPMGIIVAVVTGIAAIAAIVAAVVALIYCRKKRISANSTD				
Δ	Δ	Δ	Δ	

TABLE 21-continued

Sequence of Mature FcRIIA				
180	190	200	210	
ITAM				
PVKAARFEPLGRQTIALRKRLQLEETNNDYETADGGYMTLNPRAPT				
Δ	Δ	Δ	Δ	Δ
220	230	240	250	260
ITAM				
DDDRNIYLTLSPNDDYDNN				
Δ	Δ			
270	280			

[0201]

TABLE 22

Sequence of Mature FcγRIIB				
Domain 1				
TPAAPPKAVLKLEPPWINVLREDSVTLTTCGGAHSPDSSTQWFHN				
Δ	Δ	Δ	Δ	Δ
1	10	20	30	40
GNLIPHTHTQPSYRFKANNNDSGEYRCQTGRITSLSDPVHLTVLSE				
Δ	Δ	Δ	Δ	
50	60	70	80	
Domain 2				
WLALQTPHLEFREGETIMLRCHSWKDKPLIKVTFFQNGISKKFS				
Δ	Δ	Δ	Δ	Δ
90	100	110	120	130
HMNPNFSIPQANSHSGDYHCTGNIGYTPYSSKPVTTITVQV				
	Δ	Δ	Δ	Δ
	140	150	160	170
Transmembrane/intracellular				
PSMGSSSPIGIIIVAVVTGIAAIAAIVAAVVALIYCRKKRISANPTN				
Δ	Δ	Δ	Δ	
180	190	200	210	
ITIM motif				
PDEADKVGAEENTITYSLMHDPDALEEPDDQNRV				
Δ	Δ	Δ	Δ	
220	230	240	250	

[0202]

TABLE 23				
Sequence for Mature FcγRIIIA				
Domain 1				
EDLPKAVVFLEPQWYRVLEKDRVTLCQGAYSPEDNSTRWFHN				
Δ	Δ	Δ	Δ	Δ
1	10	20	30	40
ESLISSQTSSYFIAAARVNNSGEYRCQTSLSLSDPVQLEVHIG				
	Δ	Δ	Δ	Δ
50	60	70	80	
Domain 2				
WLLQAPRWVFKEEESIHLRCHSWKNTLLHKVTYLQNGKRKYF				
Δ	Δ	Δ	Δ	Δ
90	100	110	120	130
HQNSDFYIPKATLKDSGSYFCRGLIGSKNVSSETVNITITQ				
	Δ	Δ	Δ	Δ
140	150	160	170	
Transmembrane/intracellular				
DLAVSSISSFFPPGYQVSFCLVMVLLFAVDTGLYFSMKKSIPSST				
	Δ	Δ	Δ	Δ
180	190	200	210	
RDWEDHKFKWSKDPQDK				
Δ	Δ			
220	230			

[0203] An alignment of the nucleic acid sequence encoding the human (SEQ ID NO: 12) and cynomolgus (SEQ ID NO: 11) gamma chain of FcγRI/III is shown in Table 12.

[0204] Analysis of % sequence identity shows that the nucleic acid sequences encoding human and cynomolgus gamma chain FcγRI/III have about 99% identity.

TABLE 12				
Alignment of Human and Cynomolgus FcγRI/III				
Gamma-Chain				
	1	10		
Human	MIPAVVLLLLLLVEQAAA			
Cyno	MIPAVVLLLLLLVEQAAA			
	20	30	40	50
Human	LGEPQLCYILDAILFLYGIVLTLLYCRLKIQV			
Cyno	LGEPQLCYILDAILFLYGIVLTLLYCRLKIQV			
		60	70	80
Human	RKAAITSYEKSDGVY <u>TGL</u> STRNQET <u>YETL</u> KHEKPPQ			
	•			
Cyno	RKAAIASYEKSDGVY <u>TGL</u> STRNQET <u>YETL</u> KHEKPPQ			
	ITAM motif ITAM motif			
Cyno vs Human = 85/86 = 98.8% identity				

[0205] An amino acid sequence for human gamma chain has Accession Nos.: P30273; EMBL; M33195; AAA35828.1. EMBL; M33196; -. PIR; A35241. MIM; 147139; -. Kuester H., Thompson H., Kinet J.-P., J. Biol. Chem. 265, 6448-6452, 1990, *Characterization and expression of the gene for the human Fc receptor gamma subunit. Definition of a new gene family.*

[0206] An alignment of the amino acid sequences for human (SEQ ID NO: 26) and cynomolgus (SEQ ID NO: 25) β-2 microglobulin is shown in Table 13. The mature β-2 microglobulin has an amino acid sequence of amino acids Δ1 to Δ99 (SEQ ID NO: 70).

[0207] Analysis of the % sequence identity shows that the amino acid sequences for human and cynomolgus β-2 microglobulin have about 92% identity with no deletions or insertions.

TABLE 13	
Alignment of Human and Cynomolgus β2-Microglobulin	
Human	MSRSVALAVLALLSLSGLEA
•	
Cyno	MSPSVALAVLALLSLSGLEA
Human	IQRTPKIQVYSRHPAENGKSNFLNCYVSGFHPSDIEVDLLKNGERIEKVEHSD
• • •	

TABLE 13-continued

Alignment of Human and Cynomolgus $\beta$ 2-Microglobulin						
Cyno	IQRTPKIQVYSRHPPEKPNFLNCYVSGFHPSDIEVDLLKNGEKMKGKVEHSD					
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	1	10	20	30	40	50
Human	LSFSKDWFSYLLYYTEFTPEKDEYACRVNHVTLSPKIVKWDRDM					
	.					
Cyno	LSFSKDWFSYLLYYTEFTPEKDEYACRVNHVTLSPGRTVTKWDRDM					
	$\Delta$	$\Delta$	$\Delta$	$\Delta$		
	60	70	80	90		

[0208] Cyno vs Human 109/119=91.6% identity

[0209] The human amino acid sequence for  $\beta$ -2 microglobulin has Accession Nos.: P01884; EMBL; M17987; AAA51811.1. EMBL; M17986; AAA51811.1. EMBL; AB021288; BAA35182.1. EMBL; AF072097; AAD48083.1. EMBL; V00567; CAA23830.1. EMBL; M30683; AAA87972.1. EMBL; M30684; AAA88008.1. PIR; A02179. PIR; A28579. PDB; 1HLA. Guessow D., Rein R., Ginjaar I., Hochstenbach F., Seemann G., Kottman A., Ploegh H. L., *The human beta 2-microglobulin gene. Primary structure and definition of the transcriptional unit*, J. Immunol. 139, 3132-3138 (1987); Matsumoto K., Minamitani T., *Human mRNA for beta 2-microglobulin*, Medline: Embl/genbank/ddbj database (1998); Zhao Z., Huang X., Li N., Zhu X., Cao X., *A novel gene from human dendritic cell*, Embl/genbank/ddbj databases (1998); Rosa F., Berissi H., Weissenbach J., Maroteaux L., Fellous M., Revel M., *The beta-2-microglobulin mRNA in human Daudi cells has a mutated initiation codon but is still inducible by interferon*, EMBO J. 2, 239-243 (1983); Suggs S. V., Wallace R. B., Hirose T., Kawashima E. H., Itakura K., *Use of synthetic oligonucleotides as hybridization probes: isolation of cloned cDNA sequences for human beta 2-microglobulin*, Proc. Natl. Acad. Sci. USA 78, 6613-6617 (1981); Cunningham B. A., Wang J. L., Berggard I., Peterson P. A., *The complete amino acid sequence of beta 2-microglobulin*, Biochem. 12, 4811-4822 (1973); Lawlor D. A., Warren E., Ward F. E., Parham P., *Comparison of class I MHC alleles in human and*

*apes*, Immunol. Rev. 113, 147-185 (1990); Bjorkman P. J., Saper M. A., Samraoui B., Bennett W. S., Strominger J. L., Wiley D. C., *Structure of the human class I histocompatibility antigen, HLA-A2*, Nature 329, 506-512 (1987); Saper M. A., Bjorkman P. J., Wiley D. C., *Refined structure of the human histocompatibility antigen HLA-A2 at 2.6A resolution*, J. Mol. Biol. 219, 277-319 (1991); Collins E. J., Garboczi D. N., Karpusas M. N., Wiley D. C., *The three-dimensional structure of a class I major histocompatibility complex molecule missing the alpha 3 domain of the heavy chain*, Proc. Natl. Acad. Sci USA 92, 1218-1221 (1995).

[0210] An alignment of the amino acid sequences for human (SEQ ID NO: 30) and cynomolgus FcRn  $\alpha$ -chain (SEQ ID NO: 29) is shown in Table 14. Two alleles of cynomolgus FcRn were identified. One sequence is that of SEQ ID NO: 29 and has a serine at position 3 (S3) of the mature polypeptide. Another sequence is SEQ ID NO: 64 has an asparagine at position 3 (N3) in the mature polypeptide. The mature polypeptide of FcRnS3  $\alpha$ -chain has a sequence of amino acids  $\Delta$ 1 to  $\Delta$ 342 (SEQ ID NO: 71). The mature polypeptide of FcRnN3  $\alpha$ -chain has a sequence of  $\Delta$ 1 to  $\Delta$ 342 (SEQ ID NO: 72). An extracellular fragment of the FcRn prepared by the method of example 1, has an amino acid sequence of  $\Delta$ 1 to  $\Delta$ 274.

[0211] Analysis of the % sequence identity shows that the amino acid sequences for human and cynomolgus FcRn have about 97% identity with no deletions or insertions.

TABLE 14

Alignment of Human and Cynomolgus FcRn $\alpha$ -Chain 354/365 = 97% identity	
Signal	
Cyno	MRVPRPQPWALGLLLFLPLPGSLG
	.
Human	MGVPRPQPWALGLLLFLPLPGSLG
Extracellular Domain	
Cyno	AESHLSELLYHLTAVSSPAPGTPAFVWSGLGPGQYLSYDSLRLGQAEPCGA
CynoN3	N
	.



TABLE 14-continued

Alignment of Human and Cynomolgus FcRn $\alpha$ -Chain 354/365 = 97% identity					
Human	AESHLSLLYHLTAVSSPAPGTPAFWVSGWLGPPQYLSYNSLRGEAEPCGA				
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	10	20	30	40	50
Cyno	WVWENQVSWYWEKETTDLRIKEKLFLEAFKALGGKGPYTLQGGLGCELSP				
	•				
Human	WVWENQVSWYWEKETTDLRIKEKLFLEAFKALGGKGPYTLQGGLGCELGP				
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	60	70	80	90	100
Cyno	DNTSVPTAKFALNGEEFMNFDLKQGTWGGDWPEALAIQRWQQQDKAANK				
Human	DNTSVPTAKFALNGEEFMNFDLKQGTWGGDWPEALAIQRWQQQDKAANK				
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	110	120	130	140	150
Cyno	ELTFLLFSCPHRLREHLERGRGNLEWKEPPSMRLKARPGNPGFSVLTCSA				
	••				
Human	ELTFLLFSCPHRLREHLERGRGNLEWKEPPSMRLKARPPSPGFSVLTCSA				
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	160	170	180	190	200
Cyno	FSFYPPQLRFLRNGMAAGTGQDFGPNSDGSFHASSSLTVKSGDEHHY				
	•				
Human	FSFYPPQLRFLRNGLAAGTGQDFGPNSDGSFHASSSLTVKSGDEHHY				
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	210	220	230	240	250
Cyno	CCIVQHAGLAQPLRVELETPAKSS				
	•				
Human	CCIVQHAGLAQPLRVELESPAKSS				
	$\Delta$	$\Delta$			
	260	270			
Transmembrane/Intracellular					
Cyno	VLVVGIVIGVLLLTAAAVGGALLWRRMRSGLPAPWISLRGDDTGSLLETP				
	0				
Human	VLVVGIVIGVLLLTAAAVGGALLWRRMRSGLPAPWISLRGDDTGVLLETP				
	$\Delta$	$\Delta$	$\Delta$	$\Delta$	$\Delta$
	280	290	300	310	320
Cyno	GEAQDADSKDINVIPATA				
	• •				

TABLE 14-continued

Alignment of Human and Cynomolgus FcRn $\alpha$ -Chain 354/365 = 97% identity		
Human	GEAQDADLDKDVNVIPATA	
	$\Delta$	$\Delta$
	330	340

[0212] The human amino acid sequence for FcRn has Accession No.: U12255. Story C. M., Mikulska J., Simister N. E., A major histocompatibility complex class I-like Fc receptor cloned from human placenta: Possible role in transfer of immunoglobulin G from mother to fetus, *J. Exp. Med.* 180, 2377-2381 (1994).

## Example 3

Cynomolgus Fc $\gamma$ RI And Human Fc $\gamma$ RI Bind Human IgG Subclasses Equivalently

## [0213] Materials and Methods:

[0214] Human IgG2, IgG3, and IgG4 isotypes of E27 (IgG 1) were constructed by subcloning the appropriate heavy chain Fc cDNA from a human spleen cDNA library into a pRK vector containing the E27 variable heavy domain. All IgG subclasses and variants were expressed using the same E27  $\kappa$  light chain as described in Shields, R. L., Namenuk, A. K., Hong, K., Meng, Y. G., Rae, J., Briggs, J., Xie, D., Lai, J., Stadlen, A., Li, B., Fox, J. A., and Presta, L. G. (2001) *J. Biol. Chem.* 276:6591-6604 or U.S. Pat. No. 6,194,551.

[0215] Following cotransfection of heavy and light chain plasmids into 293 cells, IgG1, IgG2, IgG4 and variants were purified by protein A chromatography. IgG3 was purified using protein G chromatography. All protein preparations were analyzed using a combination of SDS-polyacrylamide gel electrophoresis, ELISA, and spectroscopy.

[0216] The cDNA for Human Fc $\gamma$ RI was isolated by reverse transcriptase-PCR (GeneAmp, PerkinElmer Life Sciences) of oligo(dT)-primed RNA from U937 cells using primers that generated a fragment encoding the  $\alpha$ -chain extra-cellular domain. Human Fc $\gamma$ R extracellular domains bound to Gly/6-His/GST fusions were prepared as described in Shields, R. L., Namenuk, A. K., Hong, K., Meng, Y. G., Rae, J., Briggs, J., Xie, D., Lai, J., Stadlen, A., Li, B., Fox, J. A., and Presta, L. G. (2001) *J. Biol. Chem.* 276:6591-6604 or U.S. Pat. No. 6,194,551. The cDNA was subcloned into previously described pRK mammalian cell expression vectors, as described in Eaton et al., 1986, *Biochemistry*, 25:8343-8347. The cDNA for cynomolgus Fc $\gamma$ RI was isolated as described in Example 1.

[0217] To facilitate the purification of the expressed human and cynomolgus Fc $\gamma$ RI, the transmembrane domain and intracellular domain of each were replaced by DNA encoding a Gly-His<sub>6</sub> tag and human glutathione S-transferase (GST). The GST sequence was obtained by PCR from the pGEX4T2 plasmid (Amersham Pharmacia Biotech) with NheI and XbaI restriction sites at the 5' and 3' ends, respectively. The expressed Fc $\gamma$ RI contained the extracellu-

lar domains of the  $\alpha$ -chain fused at His271 to Gly/His<sub>6</sub>/GST. Primers used to subclone the extracellular portion of the cynomolgus Fc $\gamma$ RI  $\alpha$ -chain are shown in Table 1.

[0218] The cynomolgus and human Fc $\gamma$ RI plasmids were transfected into human embryonic kidney 293 cells by calcium phosphate precipitation (Gorman, C. M., Gies, D. R., and McCray, G. (1990) *DNA Prot. Engineer. Tech.* 2, 3-10). Supernatants were collected 72 hours after conversion to serum-free PSO<sub>4</sub> medium supplemented with 10 mg/liter recombinant bovine insulin, 1 mg/liter human transferrin, and trace elements. Proteins were purified by nickel-nitrilotriacetic acid chromatography (Qiagen, Valencia, Calif.). Purified protein was analyzed through a combination of 4-20% SDS-polyacrylamide gel electrophoresis, ELISA, and amino acid analysis.

[0219] Standard enzyme-linked immunoabsorbent assays (ELISA) were performed in order to detect and quantify interactions between cynomolgus Fc $\gamma$ RI or human Fc $\gamma$ RI and human IgG1, IgG2, IgG3, or IgG4 (Table 15). ELISA plates (Nunc) were coated with 150 ng/well by adding 100  $\mu$ L of 1.5  $\mu$ g/ml stock solution cynomolgus Fc $\gamma$ RI or human Fc $\gamma$ RI in PBS for 48 hours at 4° C. After washing plates five times with wash buffer, (PBS, pH 7.4 containing 0.5% Tween-20), plates were blocked with 250  $\mu$ L of assay buffer (50 mM Tris-buffered saline, 0.05% Tween-20, 0.5% RIA-grade bovine serum albumin, 2 mM EDTA, pH 7.4) at 25° C. for 1 hours. Plates were washed five times with wash buffer.

[0220] Serial 3-fold dilutions of monomeric antibody (10.0-0.0045  $\mu$ g/ml) were added to plates and incubated for 2 hours. After washing plates five times with assay buffer, the detection reagent was added. Several different horseradish peroxidase (HRP)-conjugated reagents were used to detect the IgG-Fc $\gamma$ RI interaction, including: HRP-Protein G (Bio-Rad), goat HRP-anti-human IgG (Boehringer-Mannheim, Indianapolis, Ind.), and murine HRP-anti-human Kappa light chain. After incubation with detecting reagent at 25° C. for 90 minutes, plates were washed five times with wash buffer and 100  $\mu$ L of 0.4 mg/ml o-phenylenediamine dihydrochloride (Sigma, St. Louis, Mo.) was added. Absorbance at 490 nm was read using a Vmax plate reader (Molecular Devices, Mountain View, Calif.). Note that values reported in Table 15 are the mean+deviation relative to binding of human IgG1 at an IgG1 concentration of 0.370  $\mu$ g/ml. Titration plots for human IgG using murine HRP-anti-human Kappa light chain as detecting reagent are shown for cynomolgus Fc $\gamma$ RI (FIG. 1B) and human Fc $\gamma$ RI (FIG. 1A).

## [0221] Results and Discussion:

[0222] As illustrated in Table 15, the pattern of binding of cynomolgus Fc $\gamma$ RI and human Fc $\gamma$ RI to the four human IgG

subclasses was similar, regardless of the detection reagent. In each case, human or cynomolgus showed the highest level of binding to IgG3 and the lowest level of binding to IgG2. In particular, the pattern for both human and cynomolgus receptor-IgG interaction was  $\text{IgG3} \geq \text{IgG1} > \text{IgG4} > \text{IgG2}$ . Note that the data from the human Fc $\gamma$ RI-IgG binding interactions corresponds to data previously reported. Gessner et al, 1998, *Ann. Hematol.* 76:231-248; Deo et al., 1997, *Immunology Today* 18:127-135; Van de Winkel, 1993, *Immunology Today* 14:215-221.

TABLE 15

Subclass	Binding of monomeric human IgG subclasses to cynomolgus and human Fc $\gamma$ RI <sup>a</sup>			
	Cynomolgus Fc $\gamma$ RI		Human Fc $\gamma$ RI	
	ProtG <sup>b</sup>	anti-huIgG	anti-kappa	ProtG
E27IgG1	1.00	1.00	1.00	1.00
E27IgG2	0.13 $\pm$ 0.04	0.04, 0.04	0.11, 0.14	0.08, 0.08
E27IgG3	1.01 $\pm$ 0.06	1.22, 1.15	1.32, 1.37	1.14, 1.03
E27IgG4	0.52 $\pm$ 0.04	0.44, 0.45	0.60, 0.63	0.27, 0.27

<sup>a</sup>Detection reagents were HRP-conjugated Protein G (ProtG), HRP-conjugated murine anti-human IgG, heavy chain specific (anti-huIgG), or HRP-conjugated murine anti-human kappa light chain (anti-kappa). Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at 0.37  $\mu$ g/ml.

<sup>b</sup>Mean  $\pm$  S.D., n = 4.

[0223] As illustrated in **FIGS. 1A and 1B**, binding affinity of the human and cynomolgus Fc $\gamma$ RI is similar for each of the tested IgG subclasses. In both cases, human and cynomolgus receptors showed a markedly higher affinity for IgG3 and IgG1 as compared to the IgG4 and IgG2. **FIGS. 1A and 1B** also shows that the IgG subclass binding to Fc $\gamma$ RI is concentration-dependent and saturable.

[0224] This data illustrates that cynomolgus Fc $\gamma$ RI can replace human Fc $\gamma$ RI in the detection of IgG subclasses as human and cynomolgus reveal similar binding patterns of interaction with similar affinities for each IgG subclass.

#### Example 4

##### Cynomolgus Fc $\gamma$ RIIA Binds Human IgG2

#### [0225] Materials and Methods

[0226] ELISA assays analyzing human IgG subclass binding to cynomolgus Fc $\gamma$ RIIA were performed using essentially the methods as described in Example 3. However, because Fc $\gamma$ RIIA is a low-affinity Fc $\gamma$ R, hexameric complexes of each human IgG subclass was formed prior to addition to the Fc receptor. Hexameric complexes were formed by mixing the human IgG subclass with a human IgG at a 1:1 molar ratio. Liu, J., Lester, P., Builder, S., and Shire, S. J. (1995) *Biochemistry* 34:10474-10482. Preparation of the hexameric complexes and their use in Fc $\gamma$ RII and Fc $\gamma$ RIII assays were as described in Shields, R. L., Namenuk, A. K., Hong, K., Meng, Y. G., Rae, J., Briggs, J., Xie, D., Lai, J., Stadlen, A., Li, B., Fox, J. A., and Presta, L. G. (2001) *J. Biol. Chem.* 276:6591-6604. A plasmid encoding human Fc $\gamma$ RIIA(R131) can be readily prepared using the sequence information as described in GenBank or other published sources and see Warmerdam et al., 1991 *J. of Immunology* 147:1338-1343 and Clark et al., 1991 *J. of Immunology* 21:1911-1916.

#### [0227] Results and Discussion:

[0228] As illustrated by Table 16, the pattern of cynomolgus Fc $\gamma$ RIIA binding to hexameric complexes of the human IgG subclasses was  $\text{IgG3} = \text{IgG2} > \text{IgG1} > \text{IgG4}$ . Previous analysis of human IgG subclass binding to the two polymorphic human Fc $\gamma$ RIIA forms showed the pattern: human Fc $\gamma$ RIIA(R131)- $\text{IgG3} \geq \text{IgG1} > \text{IgG2} \geq \text{IgG4}$  and Fc $\gamma$ RIIA(H 131)- $\text{IgG3} \geq \text{IgG1} = \text{IgG2} > \text{IgG4}$ . Gessner et al, 1998, *Ann. Hematol.* 76:231-248; Deo et al., 1997, *Immunology Today* 18:127-135; Van de Winkel, 1993, *Immunology Today* 14:215-221. These binding patterns show that cynomolgus Fc $\gamma$ RIIA, which has a histidine at amino acid 131, is comparable to the human Fc $\gamma$ RIIA(H131), both of which bind human IgG2. In contrast, human Fc $\gamma$ RIIA(R131) has been reported to bind human IgG2 poorly. Note also that cynomolgus Fc $\gamma$ RIIA binds human IgG2 as efficiently as it binds human IgG3, a difference from the human Fc $\gamma$ RIIA(H 131) receptor.

TABLE 16

Subclass	Binding of hexameric complexes of human IgG subclasses to cynomolgus and human Fc $\gamma$ RIIA <sup>a</sup>		
	Cynomolgus Fc $\gamma$ RIIA		
	ProtG	anti-huIgG	anti-kappa
E27IgG1	1.00	1.00	1.00
E27IgG2	2.11	1.27	2.20 $\pm$ 0.93 <sup>b</sup>
E27IgG3	1.10	1.56	2.44 $\pm$ 0.47
E27IgG4	0.12	0.12	0.42 $\pm$ 0.18
	Human Fc $\gamma$ RIIA(H131)		
	E27IgG1	1.00	1.00
	E27IgG2	0.95	0.83
	E27IgG3	0.78	1.03
	E27IgG4	0.25	0.47
	Human Fc $\gamma$ RIIA(R131)		
	E27IgG1	1.00	1.00
	E27IgG2	0.63	0.40
	E27IgG3	1.17	1.14
	E27IgG4	0.59	0.44

<sup>a</sup>Detection reagents were HRP-conjugated Protein G (ProtG), HRP-conjugated murine anti-human IgG, heavy chain specific (anti-huIgG), or HRP-conjugated murine anti-human kappa light chain (anti-kappa). Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at 0.123  $\mu$ g/ml.

<sup>b</sup>Mean  $\pm$  SD, n = 3.

[0229] The binding of cynomolgus Fc $\gamma$ RIIA to each IgG subclass generally increased as the concentration of each antibody subclass increased (**FIG. 2**).

[0230] The data from table 16 and **FIG. 2** illustrates that cynomolgus Fc $\gamma$ RIIA binds human IgG2 and IgG3 with high efficiency and may be a preferable agent for use in detecting these human subclasses to either of the two human polymorphic forms of Fc $\gamma$ RIIA.

#### Example 5

##### Cynomolgus Fc $\gamma$ RIIB Binds Human IgG2

#### [0231] Materials and Methods

[0232] The methods used to detect Fc $\gamma$ RIIB binding to human IgG subclasses was essentially as shown in Examples 3 and 4. Plasmid encoding human Fc $\gamma$ RIIB is known and readily obtainable by those of skill in the art and see Kurucz

et al., 2000, *Immunol Lett* 75(1):33-40. Data reported in Table 17-represent the mean $\pm$ deviation relative to binding of human IgG1 at an IgG1 concentration of 0.370  $\mu$ g/ml.

**[0233] Results and Discussion:**

**[0234]** Table 17 illustrates the binding of hexameric complexes of the human IgG subclasses to human and cynomolgus Fc $\gamma$ RIIB. The binding pattern between the IgG subclasses and human Fc $\gamma$ RIIB is IgG3 $\geq$ IgG1>IgG2>IgG4 and between the IgG subclasses and cynomolgus Fc $\gamma$ RIIB is IgG2 $\geq$ IgG3>IgG1>IgG4. This binding pattern was the same for both human (**FIG. 3A**) and cynomolgus (**FIG. 3B**) over a range of IgG concentrations.

**[0235]** This data illustrates that cynomolgus Fc $\gamma$ RIIB has a stronger binding affinity for IgG2 than does human Fc $\gamma$ RIIB.

TABLE 17

Binding of Hexameric Complexes of Human IgG Subclasses to Cynomolgus and Human Fc $\gamma$ RIIB				
Subclass	Cynomolgus Fc $\gamma$ RIIB			Human Fc $\gamma$ RIIB
	ProtG <sup>b</sup>	anti-huIgG <sup>c</sup>	anti-kappa <sup>d</sup>	ProtG <sup>d</sup>
E27IgG1	1.00	1.00	1.00	1.00
E27IgG2	1.89 $\pm$ 0.37	1.26 $\pm$ 0.15	2.73 $\pm$ 1.00	0.43 $\pm$ 0.10
E27IgG3	1.25 $\pm$ 0.17	1.69 $\pm$ 0.20	2.99 $\pm$ 1.26	1.03 $\pm$ 0.13
E27IgG4	0.48 $\pm$ 0.11	0.58 $\pm$ 0.16	0.64 $\pm$ 0.21	0.23 $\pm$ 0.08

<sup>a</sup>Detection reagents were HRP-conjugated Protein G (ProtG), HRP-conjugated murine anti-human IgG, heavy chain specific (anti-huIgG), or HRP-conjugated murine anti-human kappa light chain (anti-kappa). Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at 0.37  $\mu$ g/ml.

<sup>b</sup>Mean  $\pm$  SD, n = 8.

<sup>c</sup>Mean  $\pm$  SD, n = 5.

<sup>d</sup>Mean  $\pm$  SD, n = 3.

**Example 6**

**Cynomolgus Fc $\gamma$ RIIA And Human Fc $\gamma$ RIIA-V158 Exhibit Equivalent Binding To Human IgG Subclasses**

**[0236] Materials and Methods:**

**[0237]** The methods used to detect Fc $\gamma$ RIIA binding to human IgG subclasses was essentially as shown in Examples 3 and 4. As described previously, a human DNA sequence for Fc $\gamma$ RIIA  $\alpha$ -chain is known and readily obtainable by those of skill in the art. Data reported in Table 18 represents the mean $\pm$ deviation relative to binding of human IgG1 at an IgG1 concentration of 0.370  $\mu$ g/ml.

**[0238] Results and Discussion:**

**[0239]** As illustrated in Table 18, cynomolgus Fc $\gamma$ RIIA and human Fc $\gamma$ RIIA-V 158 both bind human IgG subclasses with essentially the same pattern, IgG1>IgG3>IgG2 $\geq$ IgG4, as compared to human Fc $\gamma$ RIIA-F158, which binds with the pattern, IgG3=IgG1>>>IgG2=IgG4. The human Fc $\gamma$ RIIA-F158-human IgG subclass binding data is in agreement with previous reports. Gessner et al, 1998, *Ann. Hematol.* 76:231-248; Deo et al., 1997, *Immunology Today* 18:127-135; Van de Winkel, 1993, *Immunology Today* 14:215-221. **FIGS. 4A, 4B, and 4C** illustrate the binding pattern for human Fc $\gamma$ RIIA-F158, human Fc $\gamma$ RIIA-V158, and cynomolgus Fc $\gamma$ RIIA, respec-

tively, for increasing concentrations of each IgG subclass and indicate that the binding interactions are specific and concentration dependent and saturable.

**[0240]** The data illustrates that cynomolgus Fc $\gamma$ RIIA and human Fc $\gamma$ RIIA-V158 have equivalent binding interactions with the human IgG subclasses, and in particular that cynomolgus Fc $\gamma$ RIIA has preferred binding to the IgG2 subclass as compared to the human Fc $\gamma$ RIIA.

TABLE 18

Binding of Hexameric Complexes of Human IgG Subclasses to Cynomolgus and Human Fc $\gamma$ RIIA			
Subclass	Cynomolgus <sup>b</sup>	Human(F158) <sup>c</sup>	Human(V158) <sup>c</sup>
E27IgG1	1.00	1.00	1.00
E27IgG2	0.11 $\pm$ 0.02	0.06, 0.13	0.06, 0.03
E27IgG3	0.82 $\pm$ 0.08	0.75, 0.82	0.79, 0.82
E27IgG4	0.15 $\pm$ 0.04	0.06, 0.11	0.06, 0.04

<sup>a</sup>Detection reagent was HRP-conjugated Protein G. Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at 0.37  $\mu$ g/ml for cynomolgus Fc $\gamma$ RIIA and human Fc $\gamma$ RIIA(V158) and 1.11  $\mu$ g/ml for human Fc $\gamma$ RIIA(F158).

<sup>b</sup>Mean  $\pm$  SD, n = 4.

<sup>c</sup>Human(F158) and Human(V158) are polymorphic forms of human Fc $\gamma$ RIIA with phenylalanine or valine at receptor position 158.

**Example 7**

**Cynomolgus Fc $\gamma$ RIIA Binds Human IgG1 Variants S298A and S298A/E333A/K334A**

**[0241] Materials and Methods:**

**[0242]** Site-directed mutagenesis on E27 IgG1 was essentially as described in Shields et al., 2001, *J. Biol. Chem.*, 276:6591-6604. Briefly, site-directed mutagenesis was used to generate IgG1 variants in which a number of solvent-exposed residues in the CH2 and CH3 domains were individually altered to alanine. The alanine variants were D265A, S298A, S37A, R292A, D280A and S298A/E333A.

**[0243]** ELISA reactions were essentially as described in Examples 3-6, where IgG variants were incubated with the Fc receptors, rather than native IgG protein. Note that for the values provided in Table 19, human receptors are (Absorbance Variant/Absorbance Native IgG1) at 1  $\mu$ g/ml and for cynomolgus receptors, values are (Absorbance Variant/Absorbance Native IgG1) at 0.370  $\mu$ g/ml.

**[0244] Results and Discussion:**

**[0245]** As illustrated by Table 19 and **FIGS. 5-7**, the binding pattern of all IgG variants to cynomolgus Fc $\gamma$ RI was similar to that for human Fc $\gamma$ RI. With regard to IgG variant binding to cynomolgus Fc $\gamma$ RIIA, the pattern generally followed the same pattern for human polymorph Fc $\gamma$ RIIA(H131). (**FIG. 5**). As above, this likely reflects the fact that the cynomolgus Fc $\gamma$ RIIA has a histidine as residue 131. Note, however, that there were two notable exceptions, variant S298A and variant S298A/E333A/K334A had improved binding to the cynomolgus Fc $\gamma$ RIIA as compared to native human IgG1, and these same variants bound poorly to human Fc $\gamma$ RIIA.

**[0246]** Referring to Table 19 and **FIG. 6**, the pattern of variant IgG binding to cynomolgus Fc $\gamma$ RIIB exhibited several differences from the binding pattern for human

Fc $\gamma$ RIIB. In particular, variants R255A, E255A, E258A, S37A, D280A, and R301A bound the cynomolgus Fc $\gamma$ RIIB equivalently as they had native human IgG, whereas these same variants all exhibited improved binding to the human Fc $\gamma$ RIIB when compared to native human IgG.

[0247] Referring to Table 19 and FIG. 7, the binding pattern of the variant IgG to cynomolgus Fc $\gamma$ RIIA followed the binding pattern established for human polymorph Fc $\gamma$ RIIA-V 158, as compared to the binding pattern for human polymorph Fc $\gamma$ RIIA-F 158. This likely reflects the fact that the cynomolgus Fc $\gamma$ RIIA has a similar amino acid residue, isoleucine, at position 158 as does human Fc $\gamma$ RIIA-V158 (compared to the phenylalanine located in Fc $\gamma$ RIIA-F158).

[0248] Blocking the inhibitory signals (e.g., ITIM-containing Fc $\gamma$ RIIB) mediated by Fc receptors, which counterbalance the activating signals (e.g., ITAM-containing Fc $\gamma$ RI, Fc $\gamma$ RIIA, and Fc $\gamma$ RIIA) mediated by Fc receptors, may provide for improved therapeutic efficacy of antibodies. An unexpected result shown in Table 19 is that variants having S298A showed improved binding to cynomolgus Fc $\gamma$ RIIA, maintained native-like binding to cynomolgus Fc $\gamma$ RI and Fc $\gamma$ RIIA, and showed significantly decreased binding to cynomolgus Fc $\gamma$ RIIB. Two variants in particular, S298A and S298A/E333A/K334A may be used to selectively engage the activating ITAM-containing Fc receptors, while simultaneously not engaging the inhibitory ITIM-containing Fc $\gamma$ RIIB.

TABLE 19

Binding of Human E27 IgG1 Variants to Human and Cynomolgus Fc $\gamma$ R				
Variant	Fc $\gamma$ RI	Fc $\gamma$ RIIA	Fc $\gamma$ RIIB	Fc $\gamma$ RIIA
S239A				
Human	0.81 $\pm$ 0.09	0.73 $\pm$ 0.25	0.76 $\pm$ 0.36	0.26 $\pm$ 0.08
Cynomolgus	N/A	0.68 $\pm$ 0.04	N/A	N/A
R255A				
Human	0.99 $\pm$ 0.12	1.30 $\pm$ 0.20	1.59 $\pm$ 0.42	0.98 $\pm$ 0.18
Cynomolgus	0.85 $\pm$ 0.15	1.09 $\pm$ 0.07	0.80 $\pm$ 0.06	0.91 $\pm$ 0.08
E258A				
Human	1.18 $\pm$ 0.13	1.33 $\pm$ 0.22	1.65 $\pm$ 0.38	1.12 $\pm$ 0.12
Cynomolgus	0.91 $\pm$ 0.08	0.88 $\pm$ 0.05	0.99 $\pm$ 0.07	0.93 $\pm$ 0.11
D265A				
Human	0.16 $\pm$ 0.05	0.07 $\pm$ 0.01	0.13 $\pm$ 0.05	0.09 $\pm$ 0.06
Cynomolgus	N/A	0.05 $\pm$ 0.02	0.05	0.04 $\pm$ 0.01
S37A				
Human	1.09 $\pm$ 0.08	1.52 $\pm$ .22(R) 1.10 $\pm$ .12(H)	1.84 $\pm$ 0.43	1.05 $\pm$ 0.24
Cynomolgus	1.02 $\pm$ 0.09	1.23 $\pm$ 0.34	1.04 $\pm$ 0.30	0.88 $\pm$ 0.11
H268A				
Human	1.10 $\pm$ 0.11	1.21 $\pm$ .14(R) 0.97 $\pm$ .15(H)	1.44 $\pm$ 0.22	0.54 $\pm$ 0.12
Cynomolgus	1.02 $\pm$ 0.09	0.99 $\pm$ 0.07	1.20	0.86 $\pm$ 0.07
D280A				
Human	1.04 $\pm$ 0.08	1.34 $\pm$ 0.14	1.60 $\pm$ 0.31	1.09 $\pm$ 0.20
Cynomolgus	0.97 $\pm$ 0.08	1.45 $\pm$ 0.18	1.20 $\pm$ 0.11	0.99 $\pm$ 0.04
R292A				
Human	0.95 $\pm$ 0.05	0.27 $\pm$ 0.13	0.17 $\pm$ 0.07	0.89 $\pm$ 0.17
Cynomolgus	0.87 $\pm$ 0.08	0.80 $\pm$ 0.23	0.63 $\pm$ 0.06	0.90 $\pm$ 0.09
E293A				
Human	1.11 $\pm$ 0.07	1.08 $\pm$ 0.19	1.07 $\pm$ 0.20	0.31 $\pm$ 0.13
Cynomolgus	N/A	0.92 $\pm$ 0.07	N/A	N/A
S298A				
Human	1.11 $\pm$ 0.03	0.40 $\pm$ .15(R) 0.24 $\pm$ .08(H)	0.23 $\pm$ 0.13	1.34 $\pm$ 0.20(F) 1.07 $\pm$ .07(V)
Cynomolgus	1.06 $\pm$ 0.09	2.07 $\pm$ 0.30	0.20 $\pm$ 0.09	0.98 $\pm$ 0.13
R301M				
Human	1.06 $\pm$ 0.12	1.29 $\pm$ 0.17	1.56 $\pm$ 0.12	0.48 $\pm$ 0.21
Cynomolgus	1.00 $\pm$ 0.09	1.62 $\pm$ 0.30	1.27 $\pm$ 0.20	0.85 $\pm$ 0.08
P329A				
Human	0.48 $\pm$ 0.10	0.08 $\pm$ 0.02	0.12 $\pm$ 0.08	0.21 $\pm$ 0.03
Cynomolgus	N/A	0.21 $\pm$ 0.06	N/A	N/A
E333A				
Human	0.98 $\pm$ 0.15	0.92 $\pm$ 0.12	0.76 $\pm$ 0.11	1.27 $\pm$ 0.17
Cynomolgus	N/A	0.67 $\pm$ 0.09	N/A	N/A
K334A				
Human	1.06 $\pm$ 0.07	1.01 $\pm$ 0.15	0.90 $\pm$ 0.12	1.39 $\pm$ 0.19(F) 1.10 $\pm$ .07(V)
Cynomolgus	1.08 $\pm$ 0.09	0.92 $\pm$ 0.15	0.66 $\pm$ 0.14	1.00 $\pm$ 0.15
A339T				
Human	1.06 $\pm$ 0.04	1.09 $\pm$ 0.03	1.20 $\pm$ 0.03	1.34 $\pm$ 0.09
Cynomolgus	N/A	1.05 $\pm$ 0.02	N/A	N/A
S298A/E333A/K334A				
Human	N/A	0.35 $\pm$ 0.13	0.18 $\pm$ 0.08	1.51 $\pm$ 0.31(F) 1.11 $\pm$ .08(V)
Cynomolgus	1.19 $\pm$ 0.08	1.99 $\pm$ 0.24	0.12 $\pm$ 0.04	1.08 $\pm$ 0.15

## Example 8

## Cynomolgus FcRn And Human FcRn Bind Human IgG Subclasses Equivalently

**[0249]** Materials and Methods:

**[0250]** Human IgG2, IgG3, and IgG4 isotypes of E27 (IgG1) were constructed by subcloning the appropriate heavy chain Fc cDNA from a human spleen cDNA library into a pRK vector containing the E27 variable heavy domain. All IgG subclasses and variants were expressed using the same E27  $\kappa$  light chain.

**[0251]** Following cotransfection of heavy and light chain plasmids into 293 cells, IGG1, IgG2, IgG4 and variants were purified by protein A chromatography. IgG3 was purified using protein G chromatography. All protein preparations were analyzed using a combination of SDS-polyacrylamide gel electrophoresis, ELISA, and spectroscopy.

**[0252]** Herceptin™ IgG1 was essentially constructed as described in Coussens et al., 1985, *Science*, 230:1132-39. Herceptin™ IgG1 is a recombinant DNA-derived monoclonal antibody having an IgG1  $\kappa$  chain that contains a consensus amino acid framework with complementary-determining regions of a murine antibody (4D5) that binds HER2.

**[0253]** The cDNA for cynomolgus FcRn was isolated by reverse transcriptase-PCR (GeneAmp, PerkinElmer Life Sciences) of oligo(dT)-primed RNA from cynomolgus spleen cells using primers that generated a fragment encoding the  $\alpha$ -chain extra-cellular domain as described in Example 1. The cDNA was subcloned into previously described pRK mammalian cell expression vectors, as described in Eaton et al., 1986, *Biochemistry*, 25:8343-8347. Two DNA sequences were identified and confirmed that differed at base 77, one sequence had base G, giving Ser 3 in the mature polypeptide, and the other had base A giving Asparagine 3 in the mature polypeptide. The cDNA for cynomolgus FcRn (S3) and FcRn (N3) were isolated essentially as described in Example 1.

**[0254]** The cynomolgus and human FcRn plasmids were transfected into human embryonic kidney cells by calcium phosphate precipitation (Gorman, C. M., Gies, D. R., and McCray, G, 1990, *DNA Prot. Engineer. Tech.*, 2:3-10). Supernatants were collected 72 hours after conversion to serum-free  $\text{PSO}_4$  medium supplemented with 10 mg/liter recombinant bovine insulin, 1 mg/liter human transferrin, and trace elements. Proteins were purified using nickel nitrothiactic acid chromatography (Qiagen, Valencia, Calif.). Purified protein was analyzed through a combination of 4-20% SDS-polyacrylamide gel electrophoresis, ELISA, and amino acid analysis.

**[0255]** Standard enzyme-linked immunoabsorbent assays (ELISA) were performed in order to detect and quantify interactions between cynomolgus FcRn (S3), FcRn (N3) or human FcRn and human IgG1 (including herceptin IgG1), IgG2, IgG3, or IgG4 (table 20). ELISA plates (Nunc) were coated with 2  $\mu\text{g}/\text{ml}$  streptavidin (Zymed Laboratories Inc., South San Francisco, Calif.) in 50 mM carbonate buffer, pH 9.6, at 4° C. overnight. Plates were blocked with PBS, 0.5% BSA, 10 ppm Proclin 300 (Supelco, Bellefonte, Pa.), pH 7.2 at 25° C. for 1 h. FcRn-Gly-His<sub>6</sub> was biotinylated using a standard protocol with biotin-X—NHS (Research Organics,

Cleveland, Ohio) and bound to streptavidin coated plates at 2  $\mu\text{g}/\text{ml}$  in PBS, 0.5 BSA, 0.05% polysorbate-20 (sample buffer), pH 7.2 at 25° C. for 1 h. Plates were then rinsed with sample buffer, pH 6.0. Eight serial 2-fold dilutions of E27 standard or variants in sample buffer at pH 6.0 were incubated for 2 h. Plates were rinsed with sample buffer pH 6.0 and bound IgG was detected with peroxidase-conjugated goat F(ab')<sub>2</sub> anti-human IgG F(ab')<sub>2</sub> (Jackson ImmunoResearch) in pH 6.0 sample buffer using 3,3',5,5'-tetramethylbenzidine (Kirkegaard & Perry Laboratories, Gaithersburg, Md.) as substrate. Absorbance at 450 nm was read on a V<sub>max</sub> plate reader (Molecular Devices).

**[0256]** The data shown in Table 20 was plotted as saturation binding curves.

**[0257]** Results and Discussion:

**[0258]** As illustrated in Table 20 and corresponding FIGS. 8-10, the pattern of binding of cynomolgus FcRn (S3), FcRn (N3) and human FcRn to the four human IgG subclasses was similar. In each case, human and cynomolgus FcRns showed the highest level of binding to IgG3 and the lowest level of binding to IgG1. In particular, the pattern for both human and cynomolgus receptor-IgG interaction was IgG3>>IgG4>IgG2>IgG1. Note that the data from the human FcRn-IgG binding interactions corresponds to data previously reported. AP West Jr. and P. J. Bjorkman *Biochemistry* 39:9698 (2000).

**[0259]** In addition, the data illustrates that the binding affinity of the human and cynomolgus FcRns is similar for IgG1, IgG2, and IgG3, and is slightly stronger for IgG4, as compared to the human FcRn for IgG4. As illustrated graphically in FIGS. 8-10, binding of the human and cynomolgus FcRns to the human IgG subclasses is concentration-dependent and saturable.

TABLE 20

Binding of Human IgG Subclasses to Human FcRn				
Subclass	Cyno S3 <sup>a</sup>	Cyno N3 <sup>a</sup>	Human <sup>b</sup>	Human <sup>c</sup>
E27IgG1	1.00, 1.00	1.00, 1.00	1.00	1.00
E27IgG2	1.30, 1.15	1.49, 1.39	1.06 $\pm$ 0.10	0.93 $\pm$ 0.16
E27IgG3	3.82, 3.59	4.34, 3.97	5.60 $\pm$ 1.31	1.55 $\pm$ 0.45
E27IgG4	1.52, 1.44	1.59, 1.62	1.06 $\pm$ 0.23	0.95 $\pm$ 0.14

<sup>a</sup>Assay with NeutrAvidin coated on plate followed by FcRn-biotin, then sample and detection with HRP-conjugated goat anti-human F(ab')<sub>2</sub>. Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at [mAb] = 50 ng/ml for two assays. Cyno S3 and N3 differ only in the amino acid at position 3.

<sup>b</sup>Assay with NeutrAvidin coated on plate followed by FcRn-biotin, then sample and detection with HRP-conjugated goat anti-human F(ab')<sub>2</sub>. Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at [mAb] = 50 ng/ml for five assays. A second, separate lot of E27IgG1 showed a ratio of 0.81  $\pm$  0.03 (mean  $\pm$  S.D., n = 3) compared to the E27IgG1 used as standard.

<sup>c</sup>Assay with human IgG coated on the plate followed by sample, then FcRn-biotin and detection with HRP-conjugated streptavidin. Values are the ratio of OD<sub>490 nm</sub> (E27IgG subclass) to OD<sub>490 nm</sub> (E27IgG1) at [mAb] = 50 ng/ml for four assays. A second, separate lot of E27IgG1 showed ratios of 0.92 and 0.88 compared to the E27IgG1 used as standard.

**[0260]** This data illustrates that cynomolgus FcRn can replace human FcRn in the detection of human IgG subclasses as human and cynomolgus FcRn reveal similar binding patterns of interaction with similar affinities for each IgG subclass.

**[0261]** It will be clear that the invention is well adapted to attain the ends and advantages mentioned as well as those

inherent therein. While a presently preferred embodiment has been described for purposes of this disclosure, various changes and modifications may be made which are well within the scope of the invention. Numerous other changes may be made which will readily suggest themselves to those

skilled in the art and which are encompassed in the spirit of the invention disclosed herein and as defined in the appended claims.

[0262] All publications cited herein are hereby incorporated by reference.

---

SEQUENCE LISTING

---

<160> NUMBER OF SEQ ID NOS: 72

<210> SEQ ID NO 1  
<211> LENGTH: 1074  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1074)  
<223> OTHER INFORMATION: FcgammaRI alpha-chain

<400> SEQUENCE: 1

```
atgtggttct tgacagctct gtccttttgg gttccagttg atgggcaagt ggataccaca    60
aaggcagtga tcacttttga gcctccatgg gtcagcgtgt tccaagagga aactgtaacc    120
ttacagtgtg aggtgccccc tctgcctggg agcagctcca cacagtgggt tctcaatggc    180
acagccactc agacctcgac tcccagctac agaatacact ctgccagtgt caaggacagt    240
ggtgaataca ggtgccagag aggtccctca gggcgaagtg acccataca gctggaaatc    300
cacagagact ggctactact gcaggatatc agcagagtct tcacagaagg agaacctctg    360
gccttgagggt gtcattgatg gaaggataag ctggtgtaca atgtgcttta ctatcaaaat    420
ggcaaagcct ttaagttttt ctaccggaat tctcaactca ccattctgaa aaccaacata    480
agtcacaacg gcgcctacca ctgctcaggc atgggaaagc atcgctacac atcagcagga    540
gtatctgtca ctgtgaaaga gctatttcca gctccagtgc tgaatgcac cgtagacatcc    600
ccgctcctgg aggggaatct ggtcacctcg agctgtgaaa caaagttgct tctgcagagg    660
cctggtttgc agctttactt ctctctctac atgggcagca agacctgag aggcaggaac    720
acgtcctctg aataccaaat actaactgct agaagagaag actctggggt ttactgggtg    780
gaggccacca cagaagacgg aaatgtcctt aagcgcagcc ctgagtggga gcttcaagtg    840
cttgccctcc agttaccaac tctgtcttgg cttcatgtcc ttttctatct ggtagtggga    900
ataatgtttt tagtgaacac tgttctctgg gtgacaatac gtaaagaact gaaaagaaag    960
aaaaagtgga atttagaaat atctttggat tctgctcatg agaagaaggt aacttccagc   1020
cttcaagaag acagacattt agaagaagag ctgaagagtc aggaacaaga ataa        1074
```

<210> SEQ ID NO 2  
<211> LENGTH: 1128  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1128)  
<223> OTHER INFORMATION: FcgammaRI alpha-chain

<400> SEQUENCE: 2

```
atgtggttct tgacaactct gtccttttgg gttccagttg atgggcaagt ggacaccaca    60
aaggcagtga tcacttttga gcctccatgg gtcagcgtgt tccaagagga aaccgtaacc    120
```

## -continued

ttgcaactgtg aggtgctcca tctgcctggg agcagctcta cacagtgggt tctcaatggc	180
acagccactc agacctcgac cccagctac agaatacct ctgccagtgt caatgacagt	240
ggtgaataca ggtgccagag aggtctctca gggcgaagt accccatata gctggaaatc	300
cacagaggct ggctactact gcaggtctcc agcagagtct tcacggaagg agaacctctg	360
gccttgaggt gtcattgctg gaaggataag ctggtgtaca atgtgcttta ctatcgaaat	420
ggcaaagcct ttaagttttt ccaactggaat tctaacctca ccattctgaa aaccaacata	480
agtcacaatg gcacctacca ttgctcaggc atgggaaagc atcgctacac atcagcagga	540
atatctgtca ctgtgaaaga gctatttcca gctccagtgc tgaatgcatc tgtgacatcc	600
ccactcctgg aggggaatct ggtcaccctg agctgtgaaa caaagttgct cttgcagagg	660
cctggtttgc agctttactt ctccttctac atgggcagca agaccctgcg aggcaggaac	720
acatcctctg aataccaaat actaactgct agaagagaag actctgggtt atactggtgc	780
gaggctgcc cagaggatgg aaatgtcctt aagcgagcc ctgagttgga gcttcaagt	840
cttgccctcc agttaccaac tcctgtctgg ttctcatgtc tttctatctt ggcagtggga	900
ataatgtttt tagtgaacac tgttctctgg gtgacaatac gtaaagaact gaaaagaaag	960
aaaaagtggt atttagaaat ctctttggat tctggtcatg agaagaaggt aatttccagc	1020
cttcaagaag acagacattt agaagaagag ctgaaatgtc aggaacaaaa agaagaacag	1080
ctgcaggaag ggggtgcaccg gaaggagccc cagggggcca cgtagcag	1128

<210> SEQ ID NO 3  
 <211> LENGTH: 933  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(933)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 3

atgtctcaga atgtatgtcc cggcaacctg tggctgcttc aaccattgac agttttgctg	60
ctgtgtgctt ctgcagacag tcaaactgct ccccggaagg ctgtgctgaa actcgagccc	120
ccgtggatca acgtgctccg ggaggactct gtgactctga cgtgcggggg cgctcacagc	180
cctgacagcg actccactca gtggttccac aatgggaatc gcatccccac ccacacacag	240
cccagctaca ggttcaaggc caacaacaat gatagcgggg agtacagggt ccagactggc	300
cggaccagcc tcagcgaccc tgttcatctg actgtgcttt ctgagtggtt ggcgcttcag	360
acccctcacc tggagttccg ggaggagaa accatcatgc tgagggtcca cagctggaag	420
gacaagcctc tgatcaaggt cacattcttc cagaatgaa tagccaagaa attttcccat	480
atggatccca atttctccat cccacaagca aaccacagtc acagtgggtg ttaccactgc	540
acaggaaaca taggtacac accatactca tccaaacctg tgaccatcac tgtccaagt	600
cccagcgtgg gcagctcttc accgatgggg atcattgtgg ctgtggtcac tgggattgct	660
gtagcggcca ttgttgctgc tgtagtggcc ttgatctact gcaggaaaaa gcggatttca	720
gccaatccca ctgatcctgt gaaggctgcc cgatttgagc cacttgagcg tcaaacgatt	780
gccctcagaa agagacaact tgaagaaacc aacaatgact atgaaacagc cgacggcggc	840
tacatgactc tgaacccag ggcacctact gatgatgata gaaacatcta cctgactctt	900



## -continued

---

tctcccaacg actatgacaa cagtaataac taa 933

<210> SEQ ID NO 4  
 <211> LENGTH: 936  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(936)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 4

```

atgtctcaga atgtatgtcc cagaaacctg tggctgcttc aaccattgac agttttgctg    60
ctgctggcctt ctgcagacag tcaagctgca gctcccccac aggctgtgct gaaacttgag    120
ccccctgga tcaactgtct ccaggaggac tctgtgactc tgacatgccca gggggctcgc    180
agccctgaga gcgactccat tcagtgggtc cacaatggga atctcattcc cccccacacg    240
cagcccagct acaggttcaa ggccaacaac aatgacagcg gggagtacac gtgccagact    300
ggccagacca gcctcagcga ccctgtgcat ctgactgtgc tttccgaatg gctggtgctc    360
cagaccctc acctggagtt ccaggaggga gaaaccatca tgtgaggtg ccacagctgg    420
aaggacaagc ctctgttcaa ggtcacattc ttccagaatg gaaaatccca gaaattctcc    480
cgtttggatc ccaccttctc catcccaca gcaaaccaca gtcacagtgg tgattaccac    540
tgacagggaa acataggcta cacgctgttc tcatccaagc ctgtgacct cactgtccaa    600
gtgccagca tgggcagctc ttcaccaatg gggatcattg tggctgtggt cattgcgact    660
gctgtagcag ccattgttgc tgtgtagtgc gccttgatct actgcaggaa aaagcggatt    720
tcagccaatt ccactgatcc tgtgaaggct gcccaatttg agccacctgg acgtcaaatg    780
attgccatca gaaagagaca acttgaagaa accaacaatg actatgaaac agctgacggc    840
ggctacatga ctctgaaccc cagggcacct actgacgatg ataaaaacat ctacctgact    900
cttctccca acgaccatgt caacagtaat aactaa    936
  
```

<210> SEQ ID NO 5  
 <211> LENGTH: 885  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(885)  
 <223> OTHER INFORMATION: FcgammaRIIB  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (879)..(879)  
 <223> OTHER INFORMATION: n = a or g or c or t/u unknown or other

<400> SEQUENCE: 5

```

atgggaatcc tgtcattctt acctgtcctt gctactgaga gtgactgggc tgactgcaag    60
tcctcccagc ctgggggcca catgcttctg tggacagctg tgctattcct ggctcctgtt    120
gtggggacac ctgcagctcc cccgaaggct gtgctgaaac tcgagccccc gtggatcaac    180
gtgctccggg aggactctgt gactctgacg tgcgggggcg ctcacagccc tgacagcgac    240
tccactcagt ggttccacaa tgggaatctc atccccaccc acacgcagcc cagctacagg    300
ttcaaggcca acaacaatga tagcggggag tacaggtgcc agactggccg gaccagcctc    360
agcgaccctg ttcactctgac tgtgctttct gagtggctgg cgctccagac ccctcacctg    420
  
```

## -continued

---

```

gagttccggg agggagaaaac catcttgctg aggtgccaca gctggaagga caagcctctg 480
atcaagggtca cattcttcca gaatggaata tccaagaaat tttcccatat gaatcccaac 540
ttctccatcc cacaagcaaa ccacagtcac agtggtgatt accactgcac aggaaacata 600
ggctacacac catactcatc caaacctgtg accatcactg tccaagtgcc cagcatgggc 660
agctcttcac cgatagggat cattgtggct gtggtcactg ggattgctgt agcggccatt 720
gttgctgctg tagtggcctt gatctactgc aggaaaaagc ggatttcagc caatcccact 780
aatcctgacg aggtcgacaa agttggggct gagaacacaa tcacctattc acttctcatg 840
catccggacg ctctggaaga gcctgatgac caaaaccgng ttttag 885

```

```

<210> SEQ ID NO 6
<211> LENGTH: 876
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(876)
<223> OTHER INFORMATION: FcgammaRIIB

```

```

<400> SEQUENCE: 6

```

```

atgggaatcc tgtcattctt acctgtcctt gccactgaga gtgactgggc tgactgcaag 60
tccccccagc cttgggggtca tatgttcttg tggacagctg tgctattcct ggctcctgtt 120
gctgggacac ctgcagctcc cccaaaggct gtgctgaaac tcgagcccca gtggatcaac 180
gtgtccagg aggactctgt gactctgaca tgccggggga ctcacagccc tgagagcgac 240
tccattcagt ggttccacaa tgggaatctc attcccaccc acacgcagcc cagctacagg 300
ttcaaggcca acaacaatga cagcggggag tacacgtgcc agactggcca gaccagcctc 360
agcgaccctg tgcatctgac tgtgctttct gagtggctgg tgctccagac ccctcacctg 420
gagttccagg agggagaaaac catcgtgctg aggtgccaca gctggaagga caagcctctg 480
gtcaagggtca cattcttcca gaatggaaaa tccaagaaat tttcccgttc ggatcccaac 540
ttctccatcc cacaagcaaa ccacagtcac agtggtgatt accactgcac aggaaacata 600
ggctacacgc tgtactcatc caagcctgtg accatcactg tccaagctcc cagctcttca 660
ccgatgggga tcattgtggc tgtggtcact gggattgctg tagcggccat tgttgctgct 720
gtagtggcct tgatctactg caggaaaaag cggatttcag ccaatcccac taatcctgat 780
gaggctgaca aagttggggc tgagaacaca atcacctatt cacttctcat gcacccggat 840
gctctggaag agcctgatga ccagaaccgt atttag 876

```

```

<210> SEQ ID NO 7
<211> LENGTH: 765
<212> TYPE: DNA
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(765)
<223> OTHER INFORMATION: FcgammaRIIIA alpha-chain

```

```

<400> SEQUENCE: 7

```

```

atgtggcagc tgctcctccc aactgctctg ctacttctag tttcagctgg catgcgggct 60
gaagatctcc caaaggctgt ggtgttctct gagcctcaat ggtacagggt gctcgagaag 120
gaccgtgtga ctctgaagtg ccaggggagcc tactcccctg aggacaattc cacacgggtg 180

```

## -continued

---

```

tttcacaatg agagcctcat ctcaagccag acctcgagct acttcattgc tgetgccaga 240
gtcaacaaca gtggagagta cagggtgccag acaagcctct ccacactcag tgacccggtg 300
cagctggaag tccatatcgg ctggctattg ctccaggccc ctcggtgggt gttcaaggag 360
gaagaatcta ttcacctgag gtgtcacagc tggaagaaca ctcttctgca taaggtcacg 420
tatttacaga atggcaaagg caggaagtat ttcatcaga attctgactt ctacattcca 480
aaagccacac tcaaagacag cggctcctac ttctgcaggg gacttattgg gagtaaaaat 540
gtatcttcag agactgtgaa catcaccatc actcaagatt tggcagtgtc atccatctca 600
tcattctttc cacctgggta ccaagtctct ttctgcctgg tgatgggtact cctttttgca 660
gtggacacag gactatattt ctctatgaag aaaagcattc caagctcaac aagggactgg 720
gaggaccata aatttaaagt gagcaaggac cctcaagaca aatga 765

```

```

<210> SEQ ID NO 8
<211> LENGTH: 765
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(765)
<223> OTHER INFORMATION: FcgammaRIIIA alpha-chain

```

```

<400> SEQUENCE: 8

```

```

atgtggcagc tgetcctccc aactgctctg ctacttctag tttcagctgg catgcggtact 60
gaagatctcc caaagctgtg ggtgttcctg gagcctcaat ggtacagggt gtcgagagaag 120
gacagtgtga ctctgaagtg ccaggggagcc tactcccctg aggacaattc cacacagtgg 180
tttcacaatg agagcctcat ctcaagccag gcctcgagct acttcattga cgctgccaca 240
gtcgacgaca gtggagagta cagggtgccag acaaacctct ccaccctcag tgacccggtg 300
cagctagaag tccatatcgg ctggctgttg ctccaggccc ctcggtgggt gttcaaggag 360
gaagacccta ttcacctgag gtgtcacagc tggaagaaca ctgctctgca taaggtcaca 420
tatttacaga atggcaaagg caggaagtat ttcatcata attctgactt ctacattcca 480
aaagccacac tcaaagacag cggctcctac ttctgcaggg ggcttttttg gagtaaaaat 540
gtgtcttcag agactgtgaa catcaccatc actcaagggt tggcagtgtc aaccatctca 600
tcattctttc cacctgggta ccaagtctct ttctgcttgg tgatgggtact cctttttgca 660
gtggacacag gactatattt ctctgtgaag acaaacattc gaagctcaac aagagactgg 720
aaggaccata aatttaaagt gagaaaggac cctcaagaca aatga 765

```

```

<210> SEQ ID NO 9
<211> LENGTH: 357
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(357)
<223> OTHER INFORMATION: FcgammaRI <chain

```

```

<400> SEQUENCE: 9

```

```

Met Trp Phe Leu Thr Ala Leu Leu Trp Val Pro Val Asp Gly Gln
1           5           10          15
Val Asp Thr Thr Lys Ala Val Ile Thr Leu Gln Pro Pro Trp Val Ser
20          25          30

```

-continued

---

Val	Phe	Gln	Glu	Glu	Thr	Val	Thr	Leu	Gln	Cys	Glu	Val	Pro	Arg	Leu
	35						40					45			
Pro	Gly	Ser	Ser	Ser	Thr	Gln	Trp	Phe	Leu	Asn	Gly	Thr	Ala	Thr	Gln
	50					55				60					
Thr	Ser	Thr	Pro	Ser	Tyr	Arg	Ile	Thr	Ser	Ala	Ser	Val	Lys	Asp	Ser
65					70					75				80	
Gly	Glu	Tyr	Arg	Cys	Gln	Arg	Gly	Pro	Ser	Gly	Arg	Ser	Asp	Pro	Ile
				85				90						95	
Gln	Leu	Glu	Ile	His	Arg	Asp	Trp	Leu	Leu	Leu	Gln	Val	Ser	Ser	Arg
			100					105					110		
Val	Phe	Thr	Glu	Gly	Glu	Pro	Leu	Ala	Leu	Arg	Cys	His	Ala	Trp	Lys
		115					120					125			
Asp	Lys	Leu	Val	Tyr	Asn	Val	Leu	Tyr	Tyr	Gln	Asn	Gly	Lys	Ala	Phe
	130					135					140				
Lys	Phe	Phe	Tyr	Arg	Asn	Ser	Gln	Leu	Thr	Ile	Leu	Lys	Thr	Asn	Ile
145					150					155					160
Ser	His	Asn	Gly	Ala	Tyr	His	Cys	Ser	Gly	Met	Gly	Lys	His	Arg	Tyr
				165					170					175	
Thr	Ser	Ala	Gly	Val	Ser	Val	Thr	Val	Lys	Glu	Leu	Phe	Pro	Ala	Pro
		180						185					190		
Val	Leu	Asn	Ala	Ser	Val	Thr	Ser	Pro	Leu	Leu	Glu	Gly	Asn	Leu	Val
		195					200					205			
Thr	Leu	Ser	Cys	Glu	Thr	Lys	Leu	Leu	Leu	Gln	Arg	Pro	Gly	Leu	Gln
	210					215					220				
Leu	Tyr	Phe	Ser	Phe	Tyr	Met	Gly	Ser	Lys	Thr	Leu	Arg	Gly	Arg	Asn
225					230					235					240
Thr	Ser	Ser	Glu	Tyr	Gln	Ile	Leu	Thr	Ala	Arg	Arg	Glu	Asp	Ser	Gly
				245					250					255	
Phe	Tyr	Trp	Cys	Glu	Ala	Thr	Thr	Glu	Asp	Gly	Asn	Val	Leu	Lys	Arg
		260						265					270		
Ser	Pro	Glu	Leu	Glu	Leu	Gln	Val	Leu	Gly	Leu	Gln	Leu	Pro	Thr	Pro
		275					280					285			
Val	Trp	Leu	His	Val	Leu	Phe	Tyr	Leu	Val	Val	Gly	Ile	Met	Phe	Leu
	290					295					300				
Val	Asn	Thr	Val	Leu	Trp	Val	Thr	Ile	Arg	Lys	Glu	Leu	Lys	Arg	Lys
305					310					315					320
Lys	Lys	Trp	Asn	Leu	Glu	Ile	Ser	Leu	Asp	Ser	Ala	His	Glu	Lys	Lys
			325						330					335	
Val	Thr	Ser	Ser	Leu	Gln	Glu	Asp	Arg	His	Leu	Glu	Glu	Glu	Leu	Lys
		340						345						350	
Ser	Gln	Glu	Gln	Glu											
		355													

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 374

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: MISC\_FEATURE

&lt;222&gt; LOCATION: (1)..(374)

&lt;223&gt; OTHER INFORMATION: FcgammaRI alpha-chain

&lt;400&gt; SEQUENCE: 10

Met Trp Phe Leu Thr Thr Leu Leu Leu Trp Val Pro Val Asp Gly Gln

-continued

1	5	10	15
Val Asp Thr	Thr Lys Ala Val	Ile Ser Leu Gln Pro	Pro Trp Val Ser
	20	25	30
Val Phe Gln	Glu Glu Thr Val	Thr Leu His Cys	Glu Val Leu His Leu
	35	40	45
Pro Gly Ser	Ser Ser Thr Gln	Trp Phe Leu Asn	Gly Thr Ala Thr Gln
	50	55	60
Thr Ser Thr	Pro Ser Tyr Arg	Ile Thr Ser Ala	Ser Val Asn Asp Ser
	65	70	75
Gly Glu Tyr	Arg Cys Gln Arg	Gly Leu Ser Gly	Arg Ser Asp Pro Ile
	85	90	95
Gln Leu Glu	Ile His Arg Gly	Trp Leu Leu Leu	Gln Val Ser Ser Arg
	100	105	110
Val Phe Thr	Glu Gly Glu Pro	Leu Ala Leu Arg	Cys His Ala Trp Lys
	115	120	125
Asp Lys Leu	Val Tyr Asn Val	Leu Tyr Tyr Arg	Asn Gly Lys Ala Phe
	130	135	140
Lys Phe Phe	His Trp Asn Ser	Asn Leu Thr Ile	Leu Lys Thr Asn Ile
	145	150	155
Ser His Asn	Gly Thr Tyr His	Cys Ser Gly Met	Gly Lys His Arg Tyr
	165	170	175
Thr Ser Ala	Gly Ile Ser Val	Thr Val Lys Glu	Leu Phe Pro Ala Pro
	180	185	190
Val Leu Asn	Ala Ser Val Thr	Ser Pro Leu Leu	Glu Gly Asn Leu Val
	195	200	205
Thr Leu Ser	Cys Glu Thr Lys	Leu Leu Leu Gln	Arg Pro Gly Leu Gln
	210	215	220
Leu Tyr Phe	Ser Phe Tyr Met	Gly Ser Lys Thr	Leu Arg Gly Arg Asn
	225	230	235
Thr Ser Ser	Glu Tyr Gln Ile	Leu Thr Ala Arg	Arg Glu Asp Ser Gly
	245	250	255
Leu Tyr Trp	Cys Glu Ala Ala	Thr Glu Asp Gly	Asn Val Leu Lys Arg
	260	265	270
Ser Pro Glu	Leu Glu Leu Gln	Val Leu Gly Leu	Gln Leu Pro Thr Pro
	275	280	285
Val Trp Phe	His Val Leu Phe	Tyr Leu Ala Val	Gly Ile Met Phe Leu
	290	295	300
Val Asn Thr	Val Leu Trp Val	Thr Ile Arg Lys	Glu Leu Lys Arg Lys
	305	310	315
Lys Lys Trp	Asp Leu Glu Ile	Ser Leu Asp Ser	Gly His Glu Lys Lys
	325	330	335
Val Thr Ser	Ser Leu Gln Glu	Asp Arg His Leu	Glu Glu Glu Leu Lys
	340	345	350
Cys Gln Glu	Gln Lys Glu Glu	Gln Leu Gln Glu	Gly Val His Arg Lys
	355	360	365
Glu Pro Gln	Gly Ala Thr		
	370		

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 86

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Cynomolgus

-continued

---

```

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(86)
<223> OTHER INFORMATION: FcgammaRI/III gamma-chain

<400> SEQUENCE: 11

Met Ile Pro Ala Val Val Leu Leu Leu Leu Leu Val Glu Gln Ala
1           5           10           15

Ala Ala Leu Gly Glu Pro Gln Leu Cys Tyr Ile Leu Asp Ala Ile Leu
20          25          30

Phe Leu Tyr Gly Ile Val Leu Thr Leu Leu Tyr Cys Arg Leu Lys Ile
35          40          45

Gln Val Arg Lys Ala Ala Ile Ala Ser Tyr Glu Lys Ser Asp Gly Val
50          55          60

Tyr Thr Gly Leu Ser Thr Arg Asn Gln Glu Thr Tyr Glu Thr Leu Lys
65          70          75          80

His Glu Lys Pro Pro Gln
85

```

```

<210> SEQ ID NO 12
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(86)
<223> OTHER INFORMATION: FcgammaRI/III gamma-chain

<400> SEQUENCE: 12

Met Ile Pro Ala Val Val Leu Leu Leu Leu Leu Val Glu Gln Ala
1           5           10           15

Ala Ala Leu Gly Glu Pro Gln Leu Cys Tyr Ile Leu Asp Ala Ile Leu
20          25          30

Phe Leu Tyr Gly Ile Val Leu Thr Leu Leu Tyr Cys Arg Leu Lys Ile
35          40          45

Gln Val Arg Lys Ala Ala Ile Thr Ser Tyr Glu Lys Ser Asp Gly Val
50          55          60

Tyr Thr Gly Leu Ser Thr Arg Asn Gln Glu Thr Tyr Glu Thr Leu Lys
65          70          75          80

His Glu Lys Pro Pro Gln
85

```

```

<210> SEQ ID NO 13
<211> LENGTH: 261
<212> TYPE: DNA
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(261)
<223> OTHER INFORMATION: gamma chain

<400> SEQUENCE: 13

atgattccag cagtgtctct gctcttactc cttttggttg aacaagcagc ggccttgagg 60
gagcctcagc tctgctatat cctggatgcc atcctgtttc tgtatggaat tgcctccacc 120
ctcctctact gtcgactgaa gatccaagtg cgaaaggcag ctatagccag ctatgagaaa 180
tcagatgggt ttacacggg cctgagcacc aggaaccagg aaacttatga gactctgaag 240
catgagaaac caccacagta g                                     261

```

-continued

<210> SEQ ID NO 14  
 <211> LENGTH: 261  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(261)  
 <223> OTHER INFORMATION: gamma chain

<400> SEQUENCE: 14

```

atgattccag cagtgtctt gctcttactc cttttggttg aacaagcagc ggccctggga    60
gagcctcagc tctgctatat cctggatgcc atcctgtttc tgtatggaat tgtcctcacc    120
ctcctctact gtcgactgaa gatccaagtg cgaaaggcag ctataaccag ctatgagaaa    180
tcagatgggtg ttacacggg cctgagcacc aggaaccagg agacttacga gactctgaag    240
catgagaaac caccacagta g                                         261
  
```

<210> SEQ ID NO 15  
 <211> LENGTH: 310  
 <212> TYPE: PRT  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(310)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 15

```

Met Ser Gln Asn Val Cys Pro Gly Asn Leu Trp Leu Leu Gln Pro Leu
1          5          10          15

Thr Val Leu Leu Leu Leu Ala Ser Ala Asp Ser Gln Thr Ala Pro Pro
20         25         30

Lys Ala Val Leu Lys Leu Glu Pro Pro Trp Ile Asn Val Leu Arg Glu
35         40         45

Asp Ser Val Thr Leu Thr Cys Gly Gly Ala His Ser Pro Asp Ser Asp
50         55         60

Ser Thr Gln Trp Phe His Asn Gly Asn Arg Ile Pro Thr His Thr Gln
65         70         75         80

Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly Glu Tyr Arg
85         90         95

Cys Gln Thr Gly Arg Thr Ser Leu Ser Asp Pro Val His Leu Thr Val
100        105        110

Leu Ser Glu Trp Leu Ala Leu Gln Thr Pro His Leu Glu Phe Arg Glu
115        120        125

Gly Glu Thr Ile Met Leu Arg Cys His Ser Trp Lys Asp Lys Pro Leu
130        135        140

Ile Lys Val Thr Phe Phe Gln Asn Gly Ile Ala Lys Lys Phe Ser His
145        150        155        160

Met Asp Pro Asn Phe Ser Ile Pro Gln Ala Asn His Ser His Ser Gly
165        170        175

Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Pro Tyr Ser Ser Lys
180        185        190

Pro Val Thr Ile Thr Val Gln Val Pro Ser Val Gly Ser Ser Ser Pro
195        200        205

Met Gly Ile Ile Val Ala Val Val Thr Gly Ile Ala Val Ala Ala Ile
210        215        220
  
```

-continued

---

```

Val Ala Ala Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg Ile Ser
225                230                235                240

Ala Asn Ser Thr Asp Pro Val Lys Ala Ala Arg Phe Glu Pro Leu Gly
                245                250                255

Arg Gln Thr Ile Ala Leu Arg Lys Arg Gln Leu Glu Glu Thr Asn Asn
                260                265                270

Asp Tyr Glu Thr Ala Asp Gly Gly Tyr Met Thr Leu Asn Pro Arg Ala
                275                280                285

Pro Thr Asp Asp Asp Arg Asn Ile Tyr Leu Thr Leu Ser Pro Asn Asp
                290                295                300

Tyr Asp Asn Ser Asn Asn
305                310

```

```

<210> SEQ ID NO 16
<211> LENGTH: 317
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(317)
<223> OTHER INFORMATION: FcgammaRIIA

```

```

<400> SEQUENCE: 16

```

```

Met Ala Met Glu Thr Gln Met Ser Gln Asn Val Cys Pro Arg Asn Leu
1                5                10                15

Trp Leu Leu Gln Pro Leu Thr Val Leu Leu Leu Ala Ser Ala Asp
                20                25                30

Ser Gln Ala Ala Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro
                35                40                45

Trp Ile Asn Val Leu Gln Glu Asp Ser Val Thr Leu Thr Cys Gln Gly
                50                55                60

Ala Arg Ser Pro Glu Ser Asp Ser Ile Gln Trp Phe His Asn Gly Asn
                65                70                75                80

Leu Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn
                85                90                95

Asn Asp Ser Gly Glu Tyr Thr Cys Gln Thr Gly Gln Thr Ser Leu Ser
                100               105               110

Asp Pro Val His Leu Thr Val Leu Ser Glu Trp Leu Val Leu Gln Thr
                115               120               125

Pro His Leu Glu Phe Gln Glu Gly Glu Thr Ile Met Leu Arg Cys His
                130               135               140

Ser Trp Lys Asp Lys Pro Leu Val Lys Val Thr Phe Phe Gln Asn Gly
                145               150               155               160

Lys Ser Gln Lys Phe Ser Arg Leu Asp Pro Thr Phe Ser Ile Pro Gln
                165               170               175

Ala Asn His Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly
                180               185               190

Tyr Thr Leu Phe Ser Ser Lys Pro Val Thr Ile Thr Val Gln Val Pro
                195               200               205

Ser Met Gly Ser Ser Ser Pro Met Gly Ile Ile Val Ala Val Val Ile
                210               215               220

Ala Thr Ala Val Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr
                225               230               235               240

```



-continued

---

Cys Arg Lys Lys Arg Ile Ser Ala Asn Ser Thr Asp Pro Val Lys Ala  
                                   245                                  250                                  255

Ala Gln Phe Glu Pro Pro Gly Arg Gln Met Ile Ala Ile Arg Lys Arg  
                                   260                                  265                                  270

Gln Leu Glu Glu Thr Asn Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr  
                                   275                                  280                                  285

Met Thr Leu Asn Pro Arg Ala Pro Thr Asp Asp Asp Lys Asn Ile Tyr  
                                   290                                  295                                  300

Leu Thr Leu Pro Pro Asn Asp His Val Asn Ser Asn Asn  
                                   305                                  310                                  315

<210> SEQ ID NO 17  
 <211> LENGTH: 316  
 <212> TYPE: PRT  
 <213> ORGANISM: Chimp  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(316)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 17

Met Ala Met Glu Thr Gln Met Ser Gln Asn Val Cys Pro Arg Asn Leu  
 1                                  5                                  10                                  15

Trp Leu Leu Gln Pro Leu Thr Val Leu Leu Leu Leu Ala Ser Ala Asp  
                                   20                                  25                                  30

Ser Gln Ala Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro Trp  
                                   35                                  40                                  45

Ile Asn Val Leu Gln Glu Asp Ser Val Thr Leu Thr Cys Arg Gly Ala  
                                   50                                  55                                  60

Arg Ser Pro Glu Ser Asp Ser Ile Gln Trp Phe His Asn Gly Asn Leu  
                                   65                                  70                                  75                                  80

Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn  
                                   85                                  90                                  95

Asp Ser Gly Glu Tyr Thr Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp  
                                   100                                  105                                  110

Pro Val His Leu Thr Val Leu Ser Glu Trp Leu Val Leu Gln Thr Pro  
                                   115                                  120                                  125

His Leu Glu Phe Gln Glu Gly Glu Thr Ile Val Leu Arg Cys His Ser  
                                   130                                  135                                  140

Trp Lys Asp Lys Pro Leu Val Lys Val Thr Phe Phe Gln Asn Gly Lys  
                                   145                                  150                                  155                                  160

Ser Gln Lys Phe Ser His Leu Asp Pro Asn Leu Ser Ile Pro Gln Ala  
                                   165                                  170                                  175

Asn His Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr  
                                   180                                  185                                  190

Thr Leu Phe Ser Ser Lys Pro Val Thr Ile Thr Val Gln Ala Pro Ser  
                                   195                                  200                                  205

Val Gly Ser Ser Ser Pro Val Gly Ile Ile Val Ala Val Val Ile Ala  
                                   210                                  215                                  220

Thr Ala Val Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys  
                                   225                                  230                                  235                                  240

Arg Lys Lys Arg Ile Ser Ala Asn Ser Thr Asp Pro Val Lys Ala Ala  
                                   245                                  250                                  255

Gln Phe Glu Pro Pro Gly Arg Gln Met Ile Ala Ile Arg Lys Arg Gln

-continued

260	265	270
Leu Glu Glu Thr Asn Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr Met		
275	280	285
Thr Leu Asn Pro Arg Ala Pro Thr Asp Asp Asp Lys Asn Ile Tyr Leu		
290	295	300
Thr Leu Pro Pro Asn Asp His Val Asn Ser Asn Asn		
305	310	315
<210> SEQ ID NO 18		
<211> LENGTH: 294		
<212> TYPE: PRT		
<213> ORGANISM: Cynomolgus		
<220> FEATURE:		
<221> NAME/KEY: MISC_FEATURE		
<222> LOCATION: (1)..(294)		
<223> OTHER INFORMATION: FcgammaRIIB		
<400> SEQUENCE: 18		
Met Gly Ile Leu Ser Phe Leu Pro Val Leu Ala Thr Glu Ser Asp Trp		
1	5	10
Ala Asp Cys Lys Ser Ser Gln Pro Trp Gly His Met Leu Leu Trp Thr		
20	25	30
Ala Val Leu Phe Leu Ala Pro Val Ala Gly Thr Pro Ala Ala Pro Pro		
35	40	45
Lys Ala Val Leu Lys Leu Glu Pro Pro Trp Ile Asn Val Leu Arg Glu		
50	55	60
Asp Ser Val Thr Leu Thr Cys Gly Gly Ala His Ser Pro Asp Ser Asp		
65	70	75
Ser Thr Gln Trp Phe His Asn Gly Asn Leu Ile Pro Thr His Thr Gln		
85	90	95
Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly Glu Tyr Arg		
100	105	110
Cys Gln Thr Gly Arg Thr Ser Leu Ser Asp Pro Val His Leu Thr Val		
115	120	125
Leu Ser Glu Trp Leu Ala Leu Gln Thr Pro His Leu Glu Phe Arg Glu		
130	135	140
Gly Glu Thr Ile Leu Leu Arg Cys His Ser Trp Lys Asp Lys Pro Leu		
145	150	155
Ile Lys Val Thr Phe Phe Gln Asn Gly Ile Ser Lys Lys Phe Ser His		
165	170	175
Met Asn Pro Asn Phe Ser Ile Pro Gln Ala Asn His Ser His Ser Gly		
180	185	190
Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Pro Tyr Ser Ser Lys		
195	200	205
Pro Val Thr Ile Thr Val Gln Val Pro Ser Met Gly Ser Ser Ser Pro		
210	215	220
Ile Gly Ile Ile Val Ala Val Val Thr Gly Ile Ala Val Ala Ala Ile		
225	230	235
Val Ala Ala Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg Ile Ser		
245	250	255
Ala Asn Pro Thr Asn Pro Asp Glu Ala Asp Lys Val Gly Ala Glu Asn		
260	265	270
Thr Ile Thr Tyr Ser Leu Leu Met His Pro Asp Ala Leu Glu Glu Pro		
275	280	285

---

 -continued
 

---

Asp Asp Gln Asn Arg Val  
290

<210> SEQ ID NO 19  
<211> LENGTH: 291  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
<220> FEATURE:  
<221> NAME/KEY: MISC\_FEATURE  
<222> LOCATION: (1)..(291)  
<223> OTHER INFORMATION: FcgammaRIIB

<400> SEQUENCE: 19

Met Gly Ile Leu Ser Phe Leu Pro Val Leu Ala Thr Glu Ser Asp Trp  
1 5 10 15  
Ala Asp Cys Lys Ser Pro Gln Pro Trp Gly His Met Leu Leu Trp Thr  
20 25 30  
Ala Val Leu Phe Leu Ala Pro Val Ala Gly Thr Pro Ala Ala Pro Pro  
35 40 45  
Lys Ala Val Leu Lys Leu Glu Pro Gln Trp Ile Asn Val Leu Gln Glu  
50 55 60  
Asp Ser Val Thr Leu Thr Cys Arg Gly Thr His Ser Pro Glu Ser Asp  
65 70 75 80  
Ser Ile Gln Trp Phe His Asn Gly Asn Leu Ile Pro Thr His Thr Gln  
85 90 95  
Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asp Ser Gly Glu Tyr Thr  
100 105 110  
Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp Pro Val His Leu Thr Val  
115 120 125  
Leu Ser Glu Trp Leu Val Leu Gln Thr Pro His Leu Glu Phe Gln Glu  
130 135 140  
Gly Glu Thr Ile Val Leu Arg Cys His Ser Trp Lys Asp Lys Pro Leu  
145 150 155 160  
Val Lys Val Thr Phe Phe Gln Asn Gly Lys Ser Lys Lys Phe Ser Arg  
165 170 175  
Ser Asp Pro Asn Phe Ser Ile Pro Gln Ala Asn His Ser His Ser Gly  
180 185 190  
Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Leu Tyr Ser Ser Lys  
195 200 205  
Pro Val Thr Ile Thr Val Gln Ala Pro Ser Ser Ser Pro Met Gly Ile  
210 215 220  
Ile Val Ala Val Val Thr Gly Ile Ala Val Ala Ile Val Ala Ala  
225 230 235 240  
Val Val Ala Leu Ile Tyr Cys Arg Lys Lys Arg Ile Ser Ala Asn Pro  
245 250 255  
Thr Asn Pro Asp Glu Ala Asp Lys Val Gly Ala Glu Asn Thr Ile Thr  
260 265 270  
Tyr Ser Leu Leu Met His Pro Asp Ala Leu Glu Glu Pro Asp Asp Gln  
275 280 285  
Asn Arg Ile  
290

<210> SEQ ID NO 20  
<211> LENGTH: 254

---

-continued

---

<212> TYPE: PRT  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: MISC\_FEATURE  
<222> LOCATION: (1)..(254)  
<223> OTHER INFORMATION: FcgammaRIIIA  
  
<400> SEQUENCE: 20  
Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Leu Val Ser Ala  
1 5 10 15  
Gly Met Arg Ala Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro  
20 25 30  
Gln Trp Tyr Arg Val Leu Glu Lys Asp Arg Val Thr Leu Lys Cys Gln  
35 40 45  
Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Arg Trp Phe His Asn Glu  
50 55 60  
Ser Leu Ile Ser Ser Gln Thr Ser Ser Tyr Phe Ile Ala Ala Ala Arg  
65 70 75 80  
Val Asn Asn Ser Gly Glu Tyr Arg Cys Gln Thr Ser Leu Ser Thr Leu  
85 90 95  
Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln  
100 105 110  
Ala Pro Arg Trp Val Phe Lys Glu Glu Glu Ser Ile His Leu Arg Cys  
115 120 125  
His Ser Trp Lys Asn Thr Leu Leu His Lys Val Thr Tyr Leu Gln Asn  
130 135 140  
Gly Lys Gly Arg Lys Tyr Phe His Gln Asn Ser Asp Phe Tyr Ile Pro  
145 150 155 160  
Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Ile  
165 170 175  
Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln  
180 185 190  
Asp Leu Ala Val Ser Ser Ile Ser Ser Phe Phe Pro Pro Gly Tyr Gln  
195 200 205  
Val Ser Phe Cys Leu Val Met Val Leu Leu Phe Ala Val Asp Thr Gly  
210 215 220  
Leu Tyr Phe Ser Met Lys Lys Ser Ile Pro Ser Ser Thr Arg Asp Trp  
225 230 235 240  
Glu Asp His Lys Phe Lys Trp Ser Lys Asp Pro Gln Asp Lys  
245 250

<210> SEQ ID NO 21  
<211> LENGTH: 254  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
<220> FEATURE:  
<221> NAME/KEY: MISC\_FEATURE  
<222> LOCATION: (1)..(254)  
<223> OTHER INFORMATION: FcgammaRIIIA  
  
<400> SEQUENCE: 21

Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Leu Val Ser Ala  
1 5 10 15  
Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro  
20 25 30  
Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr Leu Lys Cys Gln

-continued

35	40	45
Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp Phe His Asn Glu		
50	55	60
Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile Asp Ala Ala Thr		
65	70	80
Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn Leu Ser Thr Leu		
	85	90
Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln		
	100	105
Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile His Leu Arg Cys		
	115	120
His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr Tyr Leu Gln Asn		
	130	135
Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp Phe Tyr Ile Pro		
145	150	155
Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Phe		
	165	170
Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln		
	180	185
Gly Leu Ala Val Ser Thr Ile Ser Ser Phe Phe Pro Pro Gly Tyr Gln		
	195	200
Val Ser Phe Cys Leu Val Met Val Leu Leu Phe Ala Val Asp Thr Gly		
	210	215
Leu Tyr Phe Ser Val Lys Thr Asn Ile Arg Ser Ser Thr Arg Asp Trp		
225	230	235
Lys Asp His Lys Phe Lys Trp Arg Lys Asp Pro Gln Asp Lys		
	245	250

<210> SEQ ID NO 22  
 <211> LENGTH: 933  
 <212> TYPE: DNA  
 <213> ORGANISM: Chimp  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(933)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 22

atgtctcaga atgtatgtcc cagaaacctg tggctgcttc aaccattgac agttttgctg	60
ctgctggcctt ctgcagacag tcaagctgct cccccaaagg ctgtgctgaa acttgagccc	120
ccgtggatca acgtgctcca ggaggactct gtgactctga catgccgggg ggctcgcagc	180
cctgagagcg actccattca gtggttccac aatgggaatc tcatccccac ccacacgcag	240
cccagctaca ggttcaaggc caacaacaat gacagcgggg agtacacgtg ccagactggc	300
cagaccagcc tcagcgaccc tgtgcatctg actgtgcttt ccgaatggct ggtgctccag	360
acccctcacc tggagttcca ggaggagaa accatcgtgc tgagggtgcca cagctggaag	420
gacaagcctc tgggtcaaggc cacattcttc cagaatggaa aatcccagaa attctcccat	480
ttggatccca acctctccat cccacaagca aaccacagtc acagtgggtga ttaccactgc	540
acaggaaaca taggctacac gctgtttctc tccaagcctg tgaccatcac tgtccaagcg	600
cccagcgtgg gcagctcttc accagtgggg atcattgtgg ctgtggatcat tgcgactgct	660
gtagcagcca ttgttgctgc tgtagtggcc ttgatctact gcaggaaaaa gcggatttca	720

## -continued

---

```

gccaatcca ctgatacctgt gaaggctgcc caatttgagc cacctggacg tcaaatgatt 780
gccatcagaa agagacaact tgaagaaacc aacaatgact atgaacagc tgacggcggc 840
tacatgactc tgaacccag ggcacctact gacgatgata aaaacatcta cctgactctt 900
cctcccaacg accatgtcaa cagtaataac taa 933

```

```

<210> SEQ ID NO 23
<211> LENGTH: 360
<212> TYPE: DNA
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(360)
<223> OTHER INFORMATION: B-2 microglobulin

```

```

<400> SEQUENCE: 23

```

```

atgtctccct cagtggcctt agccgtgctg gcgctactct ctctttcttg cctggaggct 60
atccagcgta ctccaaagat tcagggtttac tcacgccatc caccagagaa tggaaagcca 120
aatttcctga attgctatgt gtctggattt catccatctg atattgaagt tgacttactg 180
aagaatggag agaaaatggg aaaagtggag cattcagact tgtctttcag caaagactgg 240
tctttctatc tcttgtacta cactgaattc acccccaatg aaaaagatga gtatgcctgc 300
cgtgtgaacc atgtgacttt gtcagggccc aggacagtta agtgggatcg agacatgtaa 360

```

```

<210> SEQ ID NO 24
<211> LENGTH: 360
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(360)
<223> OTHER INFORMATION: B-2 microglobulin

```

```

<400> SEQUENCE: 24

```

```

atgtctcgct ccgtggcctt agctgtgctc gcgctactct ctctttcttg cctggaggct 60
atccagcgta ctccaaagat tcagggtttac tcacgtcatc cagcagagaa tggaaagtca 120
aatttcctga attgctatgt gtctgggttt catccatccg acattgaagt tgacttactg 180
aagaatggag agagaattga aaaagtggag cattcagact tgtctttcag caaggactgg 240
tctttctatc tcttgtacta cactgaattc accccactg aaaaagatga gtatgcctgc 300
cgtgtgaacc atgtgacttt gtcacagccc aagatagtta agtgggatcg agacatgtaa 360

```

```

<210> SEQ ID NO 25
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: Beta-2 microglobulin

```

```

<400> SEQUENCE: 25

```

```

Met Ser Pro Ser Val Ala Leu Ala Val Leu Ala Leu Leu Ser Leu Ser
1          5          10          15
Gly Leu Glu Ala Ile Gln Arg Thr Pro Lys Ile Gln Val Tyr Ser Arg
20          25          30
His Pro Pro Glu Asn Gly Lys Pro Asn Phe Leu Asn Cys Tyr Val Ser

```

-continued

35	40	45
Gly Phe His Pro Ser Asp Ile Glu Val Asp Leu Leu Lys Asn Gly Glu		
50	55	60
Lys Met Gly Lys Val Glu His Ser Asp Leu Ser Phe Ser Lys Asp Trp		
65	70	75 80
Ser Phe Tyr Leu Leu Tyr Tyr Thr Glu Phe Thr Pro Asn Glu Lys Asp		
	85	90 95
Glu Tyr Ala Cys Arg Val Asn His Val Thr Leu Ser Gly Pro Arg Thr		
	100	105 110
Val Lys Trp Asp Arg Asp Met		
115		

<210> SEQ ID NO 26  
 <211> LENGTH: 119  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(119)  
 <223> OTHER INFORMATION: Beta-2 microglobulin

<400> SEQUENCE: 26

Met Ser Arg Ser Val Ala Leu Ala Val Leu Ala Leu Ser Leu Ser	
1	5 10 15
Gly Leu Glu Ala Ile Gln Arg Thr Pro Lys Ile Gln Val Tyr Ser Arg	
	20 25 30
His Pro Ala Glu Asn Gly Lys Ser Asn Phe Leu Asn Cys Tyr Val Ser	
	35 40 45
Gly Phe His Pro Ser Asp Ile Glu Val Asp Leu Leu Lys Asn Gly Glu	
50	55 60
Arg Ile Glu Lys Val Glu His Ser Asp Leu Ser Phe Ser Lys Asp Trp	
65	70 75 80
Ser Phe Tyr Leu Leu Tyr Tyr Thr Glu Phe Thr Pro Thr Glu Lys Asp	
	85 90 95
Glu Tyr Ala Cys Arg Val Asn His Val Thr Leu Ser Gln Pro Lys Ile	
	100 105 110
Val Lys Trp Asp Arg Asp Met	
115	

<210> SEQ ID NO 27  
 <211> LENGTH: 1098  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1098)  
 <223> OTHER INFORMATION: FcRn alpha-chain

<400> SEQUENCE: 27

atgaggggtcc cgcggcctca gccctgggctg ctggggctcc tgctctttct cctgcccggg	60
agcctgggctg cagaaagcca cctctccctc ctgtaccacc tcaccgcggg gtccctcgccc	120
gccccgggga cgctgcctt ctgggtgtcc ggctggctgg gcccgagca gtacctgagc	180
tacgacagcc tgagggggcca ggcggagccc tgtggagctt gggctctggga aaaccaagtg	240
tcctggtatt gggagaaaga gaccacagat ctgaggatca aggagaagct ctttctggaa	300
gctttcaaa ctttgggggg aaaaggcccc tacactctgc agggcctgct gggctgtgaa	360

## -continued

---

```

ctgagccctg acaacacctc ggtgccacc gccagttcg ccctgaacgg cgaggagttc 420
atgaatttcg acctcaagca gggcacctgg ggtggggact ggcccagagg cctggctatc 480
agtcagcggg ggcagcagca ggacaaggcg gccacaagg agtcacactt cctgctattc 540
tcctgcccac accggctgcg ggagcacctg gagaggggcc gtggaaacct ggagtggaa 600
gagccccctt ccatgcgcct gaaggcccga cccggcaacc ctggcttttc cgtgcttacc 660
tgacgcgcct tctccttcta ccctccggaa ctgcaactgc ggttcctgcg gaatgggatg 720
gccgctggca ccggacaggg cgacttcggc cccaacagtg acggctcctt ccacgcctcg 780
tcgtcactaa cagtcaaaag tggcgatgag caccactact gctgcactgt gcagcacgcg 840
gggctggcgc agcccctcag ggtggagctg gaaactccag ccaagtcctc ggtgctcgtg 900
gtgggaatcg tcatcggtgt cttgtactc acggcagcgg ctgtaggagg agctctgttg 960
tggagaagga tgaggagtgg gctgccagcc ccttggatct ccctccgtgg agatgacacc 1020
gggtccctcc tgcccacccc gggggaggcc caggatgctg attcgaagga tataaatgtg 1080
atcccagcca ctgcctga 1098

```

```

<210> SEQ ID NO 28
<211> LENGTH: 1098
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1098)
<223> OTHER INFORMATION: FeRn alpha-chain

```

```

<400> SEQUENCE: 28

```

```

atgggggtcc cgcggcctca gccctgggcg ctggggctcc tgctctttct ccttcctggg 60
agcctgggcg cagaaagcca cctctccctc ctgtaccacc ttaccgcggg gtcctcgcct 120
gcccgggga ctcctgcctt ctgggtgtcc ggctggctgg gccgcagca gtacctgagc 180
tacaatagcc tcggggcgga ggcggagccc tgtggagctt gggctcggga aaaccagggtg 240
tcctggtatt ggagaaaga gaccacagat ctgaggatca aggagaagct ctttctggaa 300
gctttcaaag ctttggggg aaaaggctcc tacactctgc agggcctgct gggctgtgaa 360
ctgggccctg acaacacctc ggtgccacc gccagttcg ccctgaacgg cgaggagttc 420
atgaatttcg acctcaagca gggcacctgg ggtggggact ggcccagagg cctggctatc 480
agtcagcggg ggcagcagca ggacaaggcg gccacaagg agtcacactt cctgctattc 540
tcctgcccgc accgcctgcg ggagcacctg gagaggggcc gcggaaacct ggagtggaa 600
gagccccctt ccatgcgcct gaaggcccga cccagcagcc ctggcttttc cgtgcttacc 660
tgacgcgcct tctccttcta ccctccggag ctgcaacttc ggttcctgcg gaatgggctg 720
gccgctggca ccggccaggg tgacttcggc cccaacagtg acggatcctt ccacgcctcg 780
tcgtcactaa cagtcaaaag tggcgatgag caccactact gctgcattgt gcagcacgcg 840
gggctggcgc agcccctcag ggtggagctg gaatctccag ccaagtcctc cgtgctcgtg 900
gtgggaatcg tcatcggtgt cttgtactc acggcagcgg ctgtaggagg agctctgttg 960
tggagaagga tgaggagtgg gctgccagcc ccttggatct ccctccgtgg agacgacacc 1020
ggggtccctc tgcccacccc aggggaggcc caggatgctg atttgaagga tgtaaatgtg 1080
attccagcca ccgcctga 1098

```



-continued

---

```

<210> SEQ ID NO 29
<211> LENGTH: 365
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(365)
<223> OTHER INFORMATION: FcRn (S3)

<400> SEQUENCE: 29

Met Arg Val Pro Arg Pro Gln Pro Trp Ala Leu Gly Leu Leu Leu Phe
 1          5          10          15

Leu Leu Pro Gly Ser Leu Gly Ala Glu Ser His Leu Ser Leu Leu Tyr
 20          25          30

His Leu Thr Ala Val Ser Ser Pro Ala Pro Gly Thr Pro Ala Phe Trp
 35          40          45

Val Ser Gly Trp Leu Gly Pro Gln Gln Tyr Leu Ser Tyr Asp Ser Leu
 50          55          60

Arg Gly Gln Ala Glu Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val
 65          70          75          80

Ser Trp Tyr Trp Glu Lys Glu Thr Thr Asp Leu Arg Ile Lys Glu Lys
 85          90          95

Leu Phe Leu Glu Ala Phe Lys Ala Leu Gly Gly Lys Gly Pro Tyr Thr
100          105          110

Leu Gln Gly Leu Leu Gly Cys Glu Leu Ser Pro Asp Asn Thr Ser Val
115          120          125

Pro Thr Ala Lys Phe Ala Leu Asn Gly Glu Glu Phe Met Asn Phe Asp
130          135          140

Leu Lys Gln Gly Thr Trp Gly Gly Asp Trp Pro Glu Ala Leu Ala Ile
145          150          155          160

Ser Gln Arg Trp Gln Gln Asp Lys Ala Ala Asn Lys Glu Leu Thr
165          170          175

Phe Leu Leu Phe Ser Cys Pro His Arg Leu Arg Glu His Leu Glu Arg
180          185          190

Gly Arg Gly Asn Leu Glu Trp Lys Glu Pro Pro Ser Met Arg Leu Lys
195          200          205

Ala Arg Pro Gly Asn Pro Gly Phe Ser Val Leu Thr Cys Ser Ala Phe
210          215          220

Ser Phe Tyr Pro Pro Glu Leu Gln Leu Arg Phe Leu Arg Asn Gly Met
225          230          235          240

Ala Ala Gly Thr Gly Gln Gly Asp Phe Gly Pro Asn Ser Asp Gly Ser
245          250          255

Phe His Ala Ser Ser Ser Leu Thr Val Lys Ser Gly Asp Glu His His
260          265          270

Tyr Cys Cys Ile Val Gln His Ala Gly Leu Ala Gln Pro Leu Arg Val
275          280          285

Glu Leu Glu Thr Pro Ala Lys Ser Ser Val Leu Val Val Gly Ile Val
290          295          300

Ile Gly Val Leu Leu Leu Thr Ala Ala Ala Val Gly Gly Ala Leu Leu
305          310          315          320

Trp Arg Arg Met Arg Ser Gly Leu Pro Ala Pro Trp Ile Ser Leu Arg
325          330          335

```

-continued

---

Gly Asp Asp Thr Gly Ser Leu Leu Pro Thr Pro Gly Glu Ala Gln Asp  
                   340                  345                  350

Ala Asp Ser Lys Asp Ile Asn Val Ile Pro Ala Thr Ala  
           355                  360                  365

<210> SEQ ID NO 30  
 <211> LENGTH: 365  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(365)  
 <223> OTHER INFORMATION: FcRn alpha-chain

<400> SEQUENCE: 30

Met Gly Val Pro Arg Pro Gln Pro Trp Ala Leu Gly Leu Leu Leu Phe  
 1                  5                  10                  15

Leu Leu Pro Gly Ser Leu Gly Ala Glu Ser His Leu Ser Leu Leu Tyr  
           20                  25                  30

His Leu Thr Ala Val Ser Ser Pro Ala Pro Gly Thr Pro Ala Phe Trp  
           35                  40                  45

Val Ser Gly Trp Leu Gly Pro Gln Gln Tyr Leu Ser Tyr Asn Ser Leu  
           50                  55                  60

Arg Gly Glu Ala Glu Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val  
   65                  70                  75                  80

Ser Trp Tyr Trp Glu Lys Glu Thr Thr Asp Leu Arg Ile Lys Glu Lys  
           85                  90                  95

Leu Phe Leu Glu Ala Phe Lys Ala Leu Gly Gly Lys Gly Pro Tyr Thr  
           100                  105                  110

Leu Gln Gly Leu Leu Gly Cys Glu Leu Gly Pro Asp Asn Thr Ser Val  
           115                  120                  125

Pro Thr Ala Lys Phe Ala Leu Asn Gly Glu Glu Phe Met Asn Phe Asp  
           130                  135                  140

Leu Lys Gln Gly Thr Trp Gly Gly Asp Trp Pro Glu Ala Leu Ala Ile  
   145                  150                  155                  160

Ser Gln Arg Trp Gln Gln Gln Asp Lys Ala Ala Asn Lys Glu Leu Thr  
           165                  170                  175

Phe Leu Leu Phe Ser Cys Pro His Arg Leu Arg Glu His Leu Glu Arg  
           180                  185                  190

Gly Arg Gly Asn Leu Glu Trp Lys Glu Pro Pro Ser Met Arg Leu Lys  
           195                  200                  205

Ala Arg Pro Ser Ser Pro Gly Phe Ser Val Leu Thr Cys Ser Ala Phe  
           210                  215                  220

Ser Phe Tyr Pro Pro Glu Leu Gln Leu Arg Phe Leu Arg Asn Gly Leu  
   225                  230                  235                  240

Ala Ala Gly Thr Gly Gln Gly Asp Phe Gly Pro Asn Ser Asp Gly Ser  
           245                  250                  255

Phe His Ala Ser Ser Ser Leu Thr Val Lys Ser Gly Asp Glu His His  
           260                  265                  270

Tyr Cys Cys Ile Val Gln His Ala Gly Leu Ala Gln Pro Leu Arg Val  
           275                  280                  285

Glu Leu Glu Ser Pro Ala Lys Ser Ser Val Leu Val Val Gly Ile Val  
           290                  295                  300

Ile Gly Val Leu Leu Leu Thr Ala Ala Ala Val Gly Gly Ala Leu Leu

## -continued

305	310	315	320
Trp Arg Arg Met Arg Ser Gly Leu Pro Ala Pro Trp Ile Ser Leu Arg			
	325	330	335
Gly Asp Asp Thr Gly Val Leu Leu Pro Thr Pro Gly Glu Ala Gln Asp			
	340	345	350
Ala Asp Leu Lys Asp Val Asn Val Ile Pro Ala Thr Ala			
	355	360	365

<210> SEQ ID NO 31  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(33)  
 <223> OTHER INFORMATION: FcgammaRI - forward primer  
  
 <400> SEQUENCE: 31  
  
 caggtcaatc tctagactcc caccagcttg gag 33

<210> SEQ ID NO 32  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(33)  
 <223> OTHER INFORMATION: FcgammaRI - reverse primer  
  
 <400> SEQUENCE: 32  
  
 ggtcaactat aagcttggac ggtccagatc gat 33

<210> SEQ ID NO 33  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(34)  
 <223> OTHER INFORMATION: FcgammaRI-H6-GST - forward primer  
  
 <400> SEQUENCE: 33  
  
 caggtcaatc atcgatatgt ggttcttgac agct 34

<210> SEQ ID NO 34  
 <211> LENGTH: 51  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(51)  
 <223> OTHER INFORMATION: FcgammaRI-H6-GST - reverse primer  
  
 <400> SEQUENCE: 34  
  
 ggtcaactat gctagcatgg tgatgatggt ggtgccagac aggagttggt a 51

<210> SEQ ID NO 35  
 <211> LENGTH: 36  
 <212> TYPE: DNA  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(36)  
 <223> OTHER INFORMATION: FcgammaRIIB - forward primer

---

-continued

---

&lt;400&gt; SEQUENCE: 35

caggtcaatc tctagaatgg gaatcctgtc attctt

36

&lt;210&gt; SEQ ID NO 36

&lt;211&gt; LENGTH: 34

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Cynomolgus

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1)..(34)

&lt;223&gt; OTHER INFORMATION: FcgammaRIIB - reverse primer

&lt;400&gt; SEQUENCE: 36

ggtcaactat aagcttctaa atacggttct ggtc

34

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Cynomolgus

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1)..(33)

&lt;223&gt; OTHER INFORMATION: FcgammaRIIB-H6-GST - forward primer

&lt;400&gt; SEQUENCE: 37

caggtcaatc atcgatatgc ttctgtggac agc

33

&lt;210&gt; SEQ ID NO 38

&lt;211&gt; LENGTH: 34

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Cynomolgus

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1)..(34)

&lt;223&gt; OTHER INFORMATION: FcgammaRIIB-H6-GST - reverse primer

&lt;400&gt; SEQUENCE: 38

ggtcaactat ggtgacctat cggatgaagag ctgc

34

&lt;210&gt; SEQ ID NO 39

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Cynomolgus

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1)..(33)

&lt;223&gt; OTHER INFORMATION: FcgammaRIIIA - forward primer

&lt;400&gt; SEQUENCE: 39

caggtcaatc tctagaatgt ggcagctgct cct

33

&lt;210&gt; SEQ ID NO 40

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Cynomolgus

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1)..(33)

&lt;223&gt; OTHER INFORMATION: FcgammaRIIIA - reverse primer

&lt;400&gt; SEQUENCE: 40

tcaactataa gcttatgttc agagatgctg ctg

33

---

-continued

---

<210> SEQ ID NO 41  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(33)  
<223> OTHER INFORMATION: FcgammaRIIIA-H6-GST - forward primer

<400> SEQUENCE: 41

caggtcaatc tctagaatgt ggcagctgct cct 33

<210> SEQ ID NO 42  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(35)  
<223> OTHER INFORMATION: FcgammaRIIIA-H6-GST - reverse primer

<400> SEQUENCE: 42

ggtcaactat ggtcaccttg gtaccaggt ggaaa 35

<210> SEQ ID NO 43  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(45)  
<223> OTHER INFORMATION: Fc gamma - forward primer

<400> SEQUENCE: 43

caggtcaatc atcgatgaat tcccaccatg attccagcag tggtc 45

<210> SEQ ID NO 44  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(35)  
<223> OTHER INFORMATION: Fc gamma - reverse primer

<400> SEQUENCE: 44

ggtcaactat aagcttctac tgtggtggtt tctca 35

<210> SEQ ID NO 45  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(32)  
<223> OTHER INFORMATION: B-2 microglobulin - forward primer

<400> SEQUENCE: 45

caggtcaatc atcgattcgg gccgagatgt ct 32

<210> SEQ ID NO 46  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature

---

-continued

---

<222> LOCATION: (1)..(34)  
<223> OTHER INFORMATION: B-2 microglobulin - reverse primer  
  
<400> SEQUENCE: 46  
  
ggccaactat tctagattac atgtctcgat ccca 34  
  
<210> SEQ ID NO 47  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(35)  
<223> OTHER INFORMATION: FcgammaRIIA - forward primer  
  
<400> SEQUENCE: 47  
  
caggtcaatc tctagaatgt ctcagaatgt atgtc 35  
  
<210> SEQ ID NO 48  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(37)  
<223> OTHER INFORMATION: FcgammaRIIA - reverse primer  
  
<400> SEQUENCE: 48  
  
ggccaactat aagcttttag ttattactgt tgtcata 37  
  
<210> SEQ ID NO 49  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(35)  
<223> OTHER INFORMATION: FcgammaRIIA-H6-GST - forward primer  
  
<400> SEQUENCE: 49  
  
caggtcaatc atcgatatgt ctcagaatgt atgtc 35  
  
<210> SEQ ID NO 50  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(34)  
<223> OTHER INFORMATION: FcgammaRIIA-H6-GST - reverse primer  
  
<400> SEQUENCE: 50  
  
ggccaactat ggtgacccat cggatgaagag ctgc 34  
  
<210> SEQ ID NO 51  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(32)  
<223> OTHER INFORMATION: FcRn - forward primer  
  
<400> SEQUENCE: 51  
  
caggtcaatc atcgataggt cgtcctctca gc 32

---

-continued

---

<210> SEQ ID NO 52  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(32)  
<223> OTHER INFORMATION: FcRn - reverse primer

<400> SEQUENCE: 52

ggtcaactat gaattctcgg aatggcggat gg

32

<210> SEQ ID NO 53  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(32)  
<223> OTHER INFORMATION: FcRn-H6 - forward primer

<400> SEQUENCE: 53

cagggtcaatc atcgataggt cgtcctctca gc

32

<210> SEQ ID NO 54  
<211> LENGTH: 55  
<212> TYPE: DNA  
<213> ORGANISM: Cynomolgus  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(55)  
<223> OTHER INFORMATION: FcRn-H6 - reverse primer

<400> SEQUENCE: 54

ggtcaactat gaattcatgg tgatgatggt ggtgcgagga cttggctgga gtttc

55

<210> SEQ ID NO 55  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OF1

<400> SEQUENCE: 55

cagggtcaatc tctagacagt ggtccacaa tgg

33

<210> SEQ ID NO 56  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OR1

<400> SEQUENCE: 56

ggtcaactat aagcttaaga gtcaggtaga tgttt

35

<210> SEQ ID NO 57  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OF2

<400> SEQUENCE: 57

---

-continued

---

cagggtcaatc tctagaatac ataaccttat gtatcat 37

<210> SEQ ID NO 58  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OF3

<400> SEQUENCE: 58

cagggtcaatc tctagatata gaataacatc cactttg 37

<210> SEQ ID NO 59  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OR2

<400> SEQUENCE: 59

ggtcaactat aagcttcaga gtcattgtagc cg 32

<210> SEQ ID NO 60  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OF4

<400> SEQUENCE: 60

cagggtcaatc tctagaattc cactgatcct gtgaa 35

<210> SEQ ID NO 61  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCT primer OR3

<400> SEQUENCE: 61

ggtcaactat aagcttgctt tatttgtgaa atttgtg 37

<210> SEQ ID NO 62  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OF5

<400> SEQUENCE: 62

cagggtcaatc tctagaactt ggacgtcaaa cgatt 35

<210> SEQ ID NO 63  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: PCR primer OR4

<400> SEQUENCE: 63

ggtcaactat aagcttctgc aataaacaag ttggg 35



-continued

---

```

<210> SEQ ID NO 64
<211> LENGTH: 365
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(365)
<223> OTHER INFORMATION: FcRn (N3)

<400> SEQUENCE: 64

Met Arg Val Pro Arg Pro Gln Pro Trp Ala Leu Gly Leu Leu Leu Phe
 1             5             10             15

Leu Leu Pro Gly Ser Leu Gly Ala Glu Asn His Leu Ser Leu Leu Tyr
 20             25             30

His Leu Thr Ala Val Ser Ser Pro Ala Pro Gly Thr Pro Ala Phe Trp
 35             40             45

Val Ser Gly Trp Leu Gly Pro Gln Gln Tyr Leu Ser Tyr Asp Ser Leu
 50             55             60

Arg Gly Gln Ala Glu Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val
 65             70             75             80

Ser Trp Tyr Trp Glu Lys Glu Thr Thr Asp Leu Arg Ile Lys Glu Lys
 85             90             95

Leu Phe Leu Glu Ala Phe Lys Ala Leu Gly Gly Lys Gly Pro Tyr Thr
 100            105            110

Leu Gln Gly Leu Leu Gly Cys Glu Leu Ser Pro Asp Asn Thr Ser Val
 115            120            125

Pro Thr Ala Lys Phe Ala Leu Asn Gly Glu Glu Phe Met Asn Phe Asp
 130            135            140

Leu Lys Gln Gly Thr Trp Gly Gly Asp Trp Pro Glu Ala Leu Ala Ile
 145            150            155            160

Ser Gln Arg Trp Gln Gln Gln Asp Lys Ala Ala Asn Lys Glu Leu Thr
 165            170            175

Phe Leu Leu Phe Ser Cys Pro His Arg Leu Arg Glu His Leu Glu Arg
 180            185            190

Gly Arg Gly Asn Leu Glu Trp Lys Glu Pro Pro Ser Met Arg Leu Lys
 195            200            205

Ala Arg Pro Gly Asn Pro Gly Phe Ser Val Leu Thr Cys Ser Ala Phe
 210            215            220

Ser Phe Tyr Pro Pro Glu Leu Gln Leu Arg Phe Leu Arg Asn Gly Met
 225            230            235            240

Ala Ala Gly Thr Gly Gln Gly Asp Phe Gly Pro Asn Ser Asp Gly Ser
 245            250            255

Phe His Ala Ser Ser Ser Leu Thr Val Lys Ser Gly Asp Glu His His
 260            265            270

Tyr Cys Cys Ile Val Gln His Ala Gly Leu Ala Gln Pro Leu Arg Val
 275            280            285

Glu Leu Glu Thr Pro Ala Lys Ser Ser Val Leu Val Val Gly Ile Val
 290            295            300

Ile Gly Val Leu Leu Leu Thr Ala Ala Ala Val Gly Gly Ala Leu Leu
 305            310            315            320

Trp Arg Arg Met Arg Ser Gly Leu Pro Ala Pro Trp Ile Ser Leu Arg
 325            330            335

Gly Asp Asp Thr Gly Ser Leu Leu Pro Thr Pro Gly Glu Ala Gln Asp
 340            345            350

```

-continued

---

Ala Asp Ser Lys Asp Ile Asn Val Ile Pro Ala Thr Ala  
 355 360 365

<210> SEQ ID NO 65  
 <211> LENGTH: 336  
 <212> TYPE: PRT  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(336)  
 <223> OTHER INFORMATION: FcgammaRI alpha-chain

<400> SEQUENCE: 65

Ala Val Ile Thr Leu Gln Pro Pro Trp Val Ser Val Phe Gln Glu Glu  
 1 5 10 15  
 Thr Val Thr Leu Gln Cys Glu Val Pro Arg Leu Pro Gly Ser Ser Ser  
 20 25 30  
 Thr Gln Trp Phe Leu Asn Gly Thr Ala Thr Gln Thr Ser Thr Pro Ser  
 35 40 45  
 Tyr Arg Ile Thr Ser Ala Ser Val Lys Asp Ser Gly Glu Tyr Arg Cys  
 50 55 60  
 Gln Arg Gly Pro Ser Gly Arg Ser Asp Pro Ile Gln Leu Glu Ile His  
 65 70 75 80  
 Arg Asp Trp Leu Leu Gln Val Ser Ser Arg Val Phe Thr Glu Gly  
 85 90 95  
 Glu Pro Leu Ala Leu Arg Cys His Ala Trp Lys Asp Lys Leu Val Tyr  
 100 105 110  
 Asn Val Leu Tyr Tyr Gln Asn Gly Lys Ala Phe Lys Phe Phe Tyr Arg  
 115 120 125  
 Asn Ser Gln Leu Thr Ile Leu Lys Thr Asn Ile Ser His Asn Gly Ala  
 130 135 140  
 Tyr His Cys Ser Gly Met Gly Lys His Arg Tyr Thr Ser Ala Gly Val  
 145 150 155 160  
 Ser Val Thr Val Lys Glu Leu Phe Pro Ala Pro Val Leu Asn Ala Ser  
 165 170 175  
 Val Thr Ser Pro Leu Leu Glu Gly Asn Leu Val Thr Leu Ser Cys Glu  
 180 185 190  
 Thr Lys Leu Leu Gln Arg Pro Gly Leu Gln Leu Tyr Phe Ser Phe  
 195 200 205  
 Tyr Met Gly Ser Lys Thr Leu Arg Gly Arg Asn Thr Ser Ser Glu Tyr  
 210 215 220  
 Gln Ile Leu Thr Ala Arg Arg Glu Asp Ser Gly Phe Tyr Trp Cys Glu  
 225 230 235 240  
 Ala Thr Thr Glu Asp Gly Asn Val Leu Lys Arg Ser Pro Glu Leu Glu  
 245 250 255  
 Leu Gln Val Leu Gly Leu Gln Leu Pro Thr Pro Val Trp Leu His Val  
 260 265 270  
 Leu Phe Tyr Leu Val Val Gly Ile Met Phe Leu Val Asn Thr Val Leu  
 275 280 285  
 Trp Val Thr Ile Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asn Leu  
 290 295 300  
 Glu Ile Ser Leu Asp Ser Ala His Glu Lys Lys Val Thr Ser Ser Leu  
 305 310 315 320

-continued

---

Gln Glu Asp Arg His Leu Glu Glu Glu Leu Lys Ser Gln Glu Gln Glu  
                                   325                                  330                                  335

<210> SEQ ID NO 66  
 <211> LENGTH: 282  
 <212> TYPE: PRT  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(282)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 66

Thr Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro Trp Ile Asn  
 1                                  5                                  10                                  15

Val Leu Arg Glu Asp Ser Val Thr Leu Thr Cys Gly Gly Ala His Ser  
                                   20                                  25                                  30

Pro Asp Ser Asp Ser Thr Gln Trp Phe His Asn Gly Asn Arg Ile Pro  
                                   35                                  40                                  45

Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser  
                                   50                                  55                                  60

Gly Glu Tyr Arg Cys Gln Thr Gly Arg Thr Ser Leu Ser Asp Pro Val  
 65                                  70                                  75                                  80

His Leu Thr Val Leu Ser Glu Trp Leu Ala Leu Gln Thr Pro His Leu  
                                   85                                  90                                  95

Glu Phe Arg Glu Gly Glu Thr Ile Met Leu Arg Cys His Ser Trp Lys  
                                   100                                  105                                  110

Asp Lys Pro Leu Ile Lys Val Thr Phe Phe Gln Asn Gly Ile Ala Lys  
                                   115                                  120                                  125

Lys Phe Ser His Met Asp Pro Asn Phe Ser Ile Pro Gln Ala Asn His  
                                   130                                  135                                  140

Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Pro  
 145                                  150                                  155                                  160

Tyr Ser Ser Lys Pro Val Thr Ile Thr Val Gln Val Pro Ser Val Gly  
                                   165                                  170                                  175

Ser Ser Ser Pro Met Gly Ile Ile Val Ala Val Val Thr Gly Ile Ala  
                                   180                                  185                                  190

Val Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys Arg Lys  
                                   195                                  200                                  205

Lys Arg Ile Ser Ala Asn Ser Thr Asp Pro Val Lys Ala Ala Arg Phe  
                                   210                                  215                                  220

Glu Pro Leu Gly Arg Gln Thr Ile Ala Leu Arg Lys Arg Gln Leu Glu  
 225                                  230                                  235                                  240

Glu Thr Asn Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr Met Thr Leu  
                                   245                                  250                                  255

Asn Pro Arg Ala Pro Thr Asp Asp Asp Arg Asn Ile Tyr Leu Thr Leu  
                                   260                                  265                                  270

Ser Pro Asn Asp Tyr Asp Asn Ser Asn Asn  
                                   275                                  280

<210> SEQ ID NO 67  
 <211> LENGTH: 281  
 <212> TYPE: PRT  
 <213> ORGANISM: Chimp  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE

-continued

<222> LOCATION: (1)..(281)  
 <223> OTHER INFORMATION: FcgammaRIIA

<400> SEQUENCE: 67

```

Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro Trp Ile Asn Val
1          5          10          15
Leu Gln Glu Asp Ser Val Thr Leu Thr Cys Arg Gly Ala Arg Ser Pro
20          25          30
Glu Ser Asp Ser Ile Gln Trp Phe His Asn Gly Asn Leu Ile Pro Thr
35          40          45
His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn Asp Ser Gly
50          55          60
Glu Tyr Thr Cys Gln Thr Gly Gln Thr Ser Leu Ser Asp Pro Val His
65          70          75          80
Leu Thr Val Leu Ser Glu Trp Leu Val Leu Gln Thr Pro His Leu Glu
85          90          95
Phe Gln Glu Gly Glu Thr Ile Val Leu Arg Cys His Ser Trp Lys Asp
100         105         110
Lys Pro Leu Val Lys Val Thr Phe Phe Gln Asn Gly Lys Ser Gln Lys
115         120         125
Phe Ser His Leu Asp Pro Asn Leu Ser Ile Pro Gln Ala Asn His Ser
130         135         140
His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr Thr Leu Phe
145         150         155         160
Ser Ser Lys Pro Val Thr Ile Thr Val Gln Ala Pro Ser Val Gly Ser
165         170         175
Ser Ser Pro Val Gly Ile Ile Val Ala Val Val Ile Ala Thr Ala Val
180         185         190
Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys Arg Lys Lys
195         200         205
Arg Ile Ser Ala Asn Ser Thr Asp Pro Val Lys Ala Ala Gln Phe Glu
210         215         220
Pro Pro Gly Arg Gln Met Ile Ala Ile Arg Lys Arg Gln Leu Glu Glu
225         230         235         240
Thr Asn Asn Asp Tyr Glu Thr Ala Asp Gly Gly Tyr Met Thr Leu Asn
245         250         255
Pro Arg Ala Pro Thr Asp Asp Asp Lys Asn Ile Tyr Leu Thr Leu Pro
260         265         270
Pro Asn Asp His Val Asn Ser Asn Asn
275         280

```

<210> SEQ ID NO 68  
 <211> LENGTH: 252  
 <212> TYPE: PRT  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(252)  
 <223> OTHER INFORMATION: FcgammaaRIIB

<400> SEQUENCE: 68

```

Thr Pro Ala Ala Pro Pro Lys Ala Val Leu Lys Leu Glu Pro Pro Trp
1          5          10          15
Ile Asn Val Leu Arg Glu Asp Ser Val Thr Leu Thr Cys Gly Gly Ala
20          25          30

```

-continued

```

His Ser Pro Asp Ser Asp Ser Thr Gln Trp Phe His Asn Gly Asn Leu
 35          40          45

Ile Pro Thr His Thr Gln Pro Ser Tyr Arg Phe Lys Ala Asn Asn Asn
 50          55          60

Asp Ser Gly Glu Tyr Arg Cys Gln Thr Gly Arg Thr Ser Leu Ser Asp
 65          70          75          80

Pro Val His Leu Thr Val Leu Ser Glu Trp Leu Ala Leu Gln Thr Pro
      85          90          95

His Leu Glu Phe Arg Glu Gly Glu Thr Ile Leu Leu Arg Cys His Ser
 100          105          110

Trp Lys Asp Lys Pro Leu Ile Lys Val Thr Phe Phe Gln Asn Gly Ile
 115          120          125

Ser Lys Lys Phe Ser His Met Asn Pro Asn Phe Ser Ile Pro Gln Ala
 130          135          140

Asn His Ser His Ser Gly Asp Tyr His Cys Thr Gly Asn Ile Gly Tyr
 145          150          155          160

Thr Pro Tyr Ser Ser Lys Pro Val Thr Ile Thr Val Gln Val Pro Ser
      165          170          175

Met Gly Ser Ser Ser Pro Ile Gly Ile Ile Val Ala Val Val Thr Gly
 180          185          190

Ile Ala Val Ala Ala Ile Val Ala Ala Val Val Ala Leu Ile Tyr Cys
 195          200          205

Arg Lys Lys Arg Ile Ser Ala Asn Pro Thr Asn Pro Asp Glu Ala Asp
 210          215          220

Lys Val Gly Ala Glu Asn Thr Ile Thr Tyr Ser Leu Leu Met His Pro
 225          230          235          240

Asp Ala Leu Glu Glu Pro Asp Asp Gln Asn Arg Val
      245          250

```

```

<210> SEQ ID NO 69
<211> LENGTH: 234
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(234)
<223> OTHER INFORMATION: FcgammaRIIIA - Alpha chain

```

<400> SEQUENCE: 69

```

Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro Gln Trp Tyr Arg
 1          5          10          15

Val Leu Glu Lys Asp Arg Val Thr Leu Lys Cys Gln Gly Ala Tyr Ser
      20          25          30

Pro Glu Asp Asn Ser Thr Arg Trp Phe His Asn Glu Ser Leu Ile Ser
 35          40          45

Ser Gln Thr Ser Ser Tyr Phe Ile Ala Ala Ala Arg Val Asn Asn Ser
 50          55          60

Gly Glu Tyr Arg Cys Gln Thr Ser Leu Ser Thr Leu Ser Asp Pro Val
 65          70          75          80

Gln Leu Glu Val His Ile Gly Trp Leu Leu Gln Ala Pro Arg Trp
      85          90          95

Val Phe Lys Glu Glu Glu Ser Ile His Leu Arg Cys His Ser Trp Lys
 100          105          110

```

## -continued

---

```

Asn Thr Leu Leu His Lys Val Thr Tyr Leu Gln Asn Gly Lys Gly Arg
   115                               120                               125

Lys Tyr Phe His Gln Asn Ser Asp Phe Tyr Ile Pro Lys Ala Thr Leu
   130                               135                               140

Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Ile Gly Ser Lys Asn
  145                               150                               155                               160

Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln Asp Leu Ala Val
   165                               170                               175

Ser Ser Ile Ser Ser Phe Phe Pro Pro Gly Tyr Gln Val Ser Phe Cys
   180                               185                               190

Leu Val Met Val Leu Leu Phe Ala Val Asp Thr Gly Leu Tyr Phe Ser
   195                               200                               205

Met Lys Lys Ser Ile Pro Ser Ser Thr Arg Asp Trp Glu Asp His Lys
  210                               215                               220

Phe Lys Trp Ser Lys Asp Pro Gln Asp Lys
 225                               230

```

```

<210> SEQ ID NO 70
<211> LENGTH: 99
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(99)
<223> OTHER INFORMATION: Beta-2 microglobulin

```

```

<400> SEQUENCE: 70

```

```

Ile Gln Arg Thr Pro Lys Ile Gln Val Tyr Ser Arg His Pro Pro Glu
 1           5           10           15

Asn Gly Lys Pro Asn Phe Leu Asn Cys Tyr Val Ser Gly Phe His Pro
 20           25           30

Ser Asp Ile Glu Val Asp Leu Leu Lys Asn Gly Glu Lys Met Gly Lys
 35           40           45

Val Glu His Ser Asp Leu Ser Phe Ser Lys Asp Trp Ser Phe Tyr Leu
 50           55           60

Leu Tyr Tyr Thr Glu Phe Thr Pro Asn Glu Lys Asp Glu Tyr Ala Cys
 65           70           75           80

Arg Val Asn His Val Thr Leu Ser Gly Pro Arg Thr Val Lys Trp Asp
 85           90           95

Arg Asp Met

```

```

<210> SEQ ID NO 71
<211> LENGTH: 342
<212> TYPE: PRT
<213> ORGANISM: Cynomolgus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(342)
<223> OTHER INFORMATION: FcγRn alpha-chain (S3)

```

```

<400> SEQUENCE: 71

```

```

Ala Glu Ser His Leu Ser Leu Leu Tyr His Leu Thr Ala Val Ser Ser
 1           5           10           15

Pro Ala Pro Gly Thr Pro Ala Phe Trp Val Ser Gly Trp Leu Gly Pro
 20           25           30

Gln Gln Tyr Leu Ser Tyr Asp Ser Leu Arg Gly Gln Ala Glu Pro Cys
 35           40           45

```

-continued

---

Gly Ala Trp Val Trp Glu Asn Gln Val Ser Trp Tyr Trp Glu Lys Glu  
 50 55 60  
 Thr Thr Asp Leu Arg Ile Lys Glu Lys Leu Phe Leu Glu Ala Phe Lys  
 65 70 75 80  
 Ala Leu Gly Gly Lys Gly Pro Tyr Thr Leu Gln Gly Leu Leu Gly Cys  
 85 90 95  
 Glu Leu Ser Pro Asp Asn Thr Ser Val Pro Thr Ala Lys Phe Ala Leu  
 100 105 110  
 Asn Gly Glu Glu Phe Met Asn Phe Asp Leu Lys Gln Gly Thr Trp Gly  
 115 120 125  
 Gly Asp Trp Pro Glu Ala Leu Ala Ile Ser Gln Arg Trp Gln Gln Gln  
 130 135 140  
 Asp Lys Ala Ala Asn Lys Glu Leu Thr Phe Leu Leu Phe Ser Cys Pro  
 145 150 155 160  
 His Arg Leu Arg Glu His Leu Glu Arg Gly Arg Gly Asn Leu Glu Trp  
 165 170 175  
 Lys Glu Pro Pro Ser Met Arg Leu Lys Ala Arg Pro Gly Asn Pro Gly  
 180 185 190  
 Phe Ser Val Leu Thr Cys Ser Ala Phe Ser Phe Tyr Pro Pro Glu Leu  
 195 200 205  
 Gln Leu Arg Phe Leu Arg Asn Gly Met Ala Ala Gly Thr Gly Gln Gly  
 210 215 220  
 Asp Phe Gly Pro Asn Ser Asp Gly Ser Phe His Ala Ser Ser Ser Leu  
 225 230 235 240  
 Thr Val Lys Ser Gly Asp Glu His His Tyr Cys Cys Ile Val Gln His  
 245 250 255  
 Ala Gly Leu Ala Gln Pro Leu Arg Val Glu Leu Glu Thr Pro Ala Lys  
 260 265 270  
 Ser Ser Val Leu Val Val Gly Ile Val Ile Gly Val Leu Leu Leu Thr  
 275 280 285  
 Ala Ala Ala Val Gly Gly Ala Leu Leu Trp Arg Arg Met Arg Ser Gly  
 290 295 300  
 Leu Pro Ala Pro Trp Ile Ser Leu Arg Gly Asp Asp Thr Gly Ser Leu  
 305 310 315 320  
 Leu Pro Thr Pro Gly Glu Ala Gln Asp Ala Asp Ser Lys Asp Ile Asn  
 325 330 335  
 Val Ile Pro Ala Thr Ala  
 340

<210> SEQ ID NO 72  
 <211> LENGTH: 342  
 <212> TYPE: PRT  
 <213> ORGANISM: Cynomolgus  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (1)..(342)  
 <223> OTHER INFORMATION: FcgammaRn alpha-chain (N3)  
 <400> SEQUENCE: 72

Ala Glu Asn His Leu Ser Leu Leu Tyr His Leu Thr Ala Val Ser Ser  
 1 5 10 15  
 Pro Ala Pro Gly Thr Pro Ala Phe Trp Val Ser Gly Trp Leu Gly Pro  
 20 25 30

-continued

---

Gln	Gln	Tyr	Leu	Ser	Tyr	Asp	Ser	Leu	Arg	Gly	Gln	Ala	Glu	Pro	Cys
	35						40					45			
Gly	Ala	Trp	Val	Trp	Glu	Asn	Gln	Val	Ser	Trp	Tyr	Trp	Glu	Lys	Glu
	50					55					60				
Thr	Thr	Asp	Leu	Arg	Ile	Lys	Glu	Lys	Leu	Phe	Leu	Glu	Ala	Phe	Lys
	65				70					75				80	
Ala	Leu	Gly	Gly	Lys	Gly	Pro	Tyr	Thr	Leu	Gln	Gly	Leu	Leu	Gly	Cys
				85					90					95	
Glu	Leu	Ser	Pro	Asp	Asn	Thr	Ser	Val	Pro	Thr	Ala	Lys	Phe	Ala	Leu
		100						105					110		
Asn	Gly	Glu	Glu	Phe	Met	Asn	Phe	Asp	Leu	Lys	Gln	Gly	Thr	Trp	Gly
	115						120					125			
Gly	Asp	Trp	Pro	Glu	Ala	Leu	Ala	Ile	Ser	Gln	Arg	Trp	Gln	Gln	Gln
	130					135					140				
Asp	Lys	Ala	Ala	Asn	Lys	Glu	Leu	Thr	Phe	Leu	Leu	Phe	Ser	Cys	Pro
	145				150					155					160
His	Arg	Leu	Arg	Glu	His	Leu	Glu	Arg	Gly	Arg	Gly	Asn	Leu	Glu	Trp
			165						170					175	
Lys	Glu	Pro	Pro	Ser	Met	Arg	Leu	Lys	Ala	Arg	Pro	Gly	Asn	Pro	Gly
		180						185					190		
Phe	Ser	Val	Leu	Thr	Cys	Ser	Ala	Phe	Ser	Phe	Tyr	Pro	Pro	Glu	Leu
		195					200					205			
Gln	Leu	Arg	Phe	Leu	Arg	Asn	Gly	Met	Ala	Ala	Gly	Thr	Gly	Gln	Gly
	210					215					220				
Asp	Phe	Gly	Pro	Asn	Ser	Asp	Gly	Ser	Phe	His	Ala	Ser	Ser	Ser	Leu
	225				230					235					240
Thr	Val	Lys	Ser	Gly	Asp	Glu	His	His	Tyr	Cys	Cys	Ile	Val	Gln	His
			245						250					255	
Ala	Gly	Leu	Ala	Gln	Pro	Leu	Arg	Val	Glu	Leu	Glu	Thr	Pro	Ala	Lys
		260					265						270		
Ser	Ser	Val	Leu	Val	Val	Gly	Ile	Val	Ile	Gly	Val	Leu	Leu	Leu	Thr
		275					280					285			
Ala	Ala	Ala	Val	Gly	Gly	Ala	Leu	Leu	Trp	Arg	Arg	Met	Arg	Ser	Gly
	290					295					300				
Leu	Pro	Ala	Pro	Trp	Ile	Ser	Leu	Arg	Gly	Asp	Asp	Thr	Gly	Ser	Leu
	305				310					315					320
Leu	Pro	Thr	Pro	Gly	Glu	Ala	Gln	Asp	Ala	Asp	Ser	Lys	Asp	Ile	Asn
			325						330					335	
Val	Ile	Pro	Ala	Thr	Ala										
			340												

---

1-43. (cancelled)

44. A method for evaluating at least one biological property of an Fc region containing molecule comprising:

- contacting an isolated non-human primate Fc receptor polypeptide with an Fc region containing molecule; and
- determining the effect of the contact on at least one biological property of the Fc region containing molecule.

45. A method according to claim 44, wherein the Fc region containing molecule is an antibody.

46. A method according to claim 45, wherein the antibody is a humanized antibody.

47. A method according to claim 46, wherein the antibody is an antibody variant.

48. A method according to claim 47, wherein the non-human primate Fc receptor polypeptide is a soluble receptor.

49. A method according to claim 48, wherein the non-human primate receptor polypeptide is selected from the group consisting of FcγRI α-chain, FcγRIIA, FcγRIIB, FcγRIIA α-chain, FcRn α-chain and mixtures thereof.

50. A method according to claim 44, wherein the non-human primate receptor polypeptide is expressed on a cell.



51. A method according to claim 44, wherein the biological property is the binding affinity of the Fc region containing molecule for the non-human primate receptor polypeptide.

52. A method according to claim 44, wherein the biological property is the toxicity of the Fc region containing molecule.

53. A method according to claim 44, wherein the isolated non-human primate Fc receptor polypeptide is a FcRn  $\alpha$ -chain and the biological property is the half-life of the Fc region containing molecule.

54. A method according to claim 44, wherein the non-human primate Fc receptor polypeptide comprises an amino sequence of 1 to 265 of SEQ ID NO: 65.

55. A method according to claim 44, wherein the non-human primate Fc receptor polypeptide comprises an amino acid sequence of 1 to 172 of SEQ ID NO: 66.

56. A method according to claim 44, wherein the non-human primate Fc receptor polypeptide comprises an amino acid sequence of 1 to 174 of SEQ ID NO: 68.

57. A method according to claim 47, wherein the non-human primate receptor polypeptide comprises an amino acid sequence of amino acids 1 to 172 of SEQ ID NO: 69.

58. A method according to claim 44, wherein the non-human primate Fc receptor polypeptide comprises an amino acid sequence of amino acids 1 to 171 of SEQ ID NO: 67.

59. A method for evaluating at least one biological property of an Fc region containing molecule comprising:

- a) contacting a Fc region containing molecule with a cell transformed with an isolated nucleic acid encoding a nonhuman primate Fc receptor polypeptide; and
- b) determining the effect of the contact on at least one biological property of the Fc region containing molecule.

60. A method according to claim 59, wherein the Fc region containing molecule is an antibody or antibody variant.

61. A method according to claim 59, wherein the biological property is the binding affinity of the Fc region containing molecule for the non-human primate Fc receptor polypeptide.

62. A method according to claim 59, wherein the cell is transformed with at least two nucleic acids according to claim 1.

63. A method according to claim 62, wherein the nucleic acids comprise a nucleic acid that encodes a cynomolgus Fc $\gamma$ R1  $\alpha$ -chain of SEQ ID NO: 9 and a nucleic acid that encodes a cynomolgus Fc $\gamma$ R gamma chain of SEQ ID NO: 11.

64. A method according to claim 62, wherein the nucleic acids comprise a nucleic acid that encodes a cynomolgus Fc $\gamma$ RIII  $\alpha$ -chain of SEQ ID NO: 20 and a nucleic acid that encodes a cynomolgus Fc $\gamma$ R gamma chain of SEQ ID NO: 11.

65. A method according to claim 62, wherein the nucleic acids comprise a nucleic acid that encodes a cynomolgus Fc $\gamma$ R  $\alpha$ -chain of SEQ ID NO: 29 and a nucleic acid sequence that encodes a cynomolgus  $\beta$ -2 microglobulin of SEQ ID NO: 25.

66. A method for identifying an agent that has an increased affinity for at least one cynomolgus Fc receptor polypeptide with an ITAM region compared to human Fc receptor polypeptide comprising:

a) determining the binding affinity of the agent to at least one cynomolgus Fc receptor polypeptide associated a polypeptide with an ITAM region;

b) determining the binding affinity of the agent to the corresponding human Fc receptor polypeptide; and

c) selecting agents that have an increased affinity for the cynomolgus Fc $\gamma$  receptor polypeptide associated with a polypeptide with an ITAM region compared to the corresponding human Fc receptor.

67. A method according to claim 66, wherein the agent is an antibody.

68. A method according to claim 67, wherein the agent is an IgG antibody.

69. A method according to claim 67, wherein the Fc receptor polypeptide is selected from the group consisting of Fc $\gamma$ R1  $\alpha$ -chain, Fc $\gamma$ RIIA, Fc $\gamma$ RIIA  $\alpha$ -chain and mixtures thereof.

70. A method for identifying an agent that has an altered affinity for a cynomolgus Fc receptor polypeptide with an ITIM region compared to corresponding human Fc receptor polypeptide comprising:

a) determining a binding affinity for the agent to be at least one cynomolgus Fc $\gamma$ RIIB receptor polypeptide;

b) determining a binding affinity of the agent to corresponding human Fc $\gamma$ RIIB receptor polypeptide; and

c) selecting agents with altered affinity for a cynomolgus Fc $\gamma$ RIIB receptor polypeptide with an ITIM region compared to corresponding human Fc $\gamma$ RIIB polypeptide.

71. A method according to claim 70, wherein the agent is an antibody.

72. A method for identifying an agent with increased binding affinity for a cynomolgus Fc receptor polypeptide with an ITAM region and decreased affinity for a cynomolgus Fc receptor polypeptide with an ITIM region comprising:

a) determining a binding affinity of the agent for at least one cynomolgus Fc receptor polypeptide associated with an ITAM region and a binding affinity of the agent to the corresponding human Fc receptor polypeptide;

b) determining the binding affinity of the agent for at least one cynomolgus Fc receptor polypeptide with an ITIM region and a binding affinity of the agent for the corresponding human Fc receptor polypeptide; and

c) selecting an agent with enhanced binding for a cynomolgus Fc receptor polypeptide with an ITAM region and a decreased affinity for a cynomolgus Fc receptor polypeptide with an ITIM region compared to the corresponding human Fc receptor polypeptides.

73. A method according to claim 72, wherein the Fc $\gamma$  receptor with an ITAM region is an Fc $\gamma$  receptor IIA and the Fc $\gamma$  receptor with an ITIM region is a Fc $\gamma$  receptor IIB.

74. A method according to claim 73, wherein the agent is an antibody.

75-90. (cancelled)