



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2294355 C 2006/11/28

(11)(21) **2 294 355**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1997/07/01

(87) Date publication PCT/PCT Publication Date: 1998/12/30

(45) Date de délivrance/Issue Date: 2006/11/28

(85) Entrée phase nationale/National Entry: 1999/12/15

(86) N° demande PCT/PCT Application No.: EP 1997/003493

(87) N° publication PCT/PCT Publication No.: 1998/059248

(30) Priorité/Priority: 1997/06/20 (EPPCT/EP97/03287)

(51) Cl.Int./Int.Cl. *C12N 5/12* (2006.01),
C07K 16/44 (2006.01), *C07K 16/18* (2006.01),
C12P 21/08 (2006.01), *G01N 33/577* (2006.01),
G01N 33/92 (2006.01), *G01N 33/96* (2006.01)

(72) Inventeurs/Inventors:
HOLVOET, PAUL NOEL, BE;
COLLEN, DESIRE JOSE, BE

(73) Propriétaire/Owner:
LEUVEN RESEARCH & DEVELOPMENT VZW, BE


(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : DOSAGES, ANTICORPS ET ETALONS DE DETECTION DE LIPOPROTEINES DE BASSE DENSITE,
OXYDEES ET MODIFIEES PAR MDA

(54) Title: ASSAYS, ANTIBODIES, AND STANDARDS FOR DETECTION OF OXIDIZED AND MDA-MODIFIED LOW
DENSITY LIPOPROTEINS

(57) **Abrégé/Abstract:**

Immunoassays for malondialdehyde-modified low density lipoprotein (MDA-modified LDL) and oxidized low density lipoprotein (OxLDL), monoclonal antibodies (and the cell lines for them) for use in the assays, and a storage-stable standard (which may be used as a calibrator and/or control) are disclosed. MDA-modified LDL and OxLDL are implicated in atherosclerosis and its etiology.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/92, 33/96, C07K 16/44, G01N 33/577, C12P 21/08	A1	(11) International Publication Number: WO 98/59248 (43) International Publication Date: 30 December 1998 (30.12.98)
(21) International Application Number: PCT/EP97/03493 (22) International Filing Date: 1 July 1997 (01.07.97) (30) Priority Data: PCT/EP97/03287 20 June 1997 (20.06.97) WO (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): LEUVEN RESEARCH & DEVELOPMENT VZW [BE/BE]; Groot Begijnhof, Benedenstraat 59, B-3000 Leuven (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): HOLVOET, Paul, Noel [BE/BE]; Broekstraat 13A, B-3010 Kessel-Lo (BE). COLLEN, Désiré, José [BE/BE]; Schoonzichtlaan 20, B-3020 Winkelse-Herent (BE). (74) Agent: VAN SOMEREN, Petronella, Francisca, Hendrika, Maria; Arnold & Siedsma, Sweelinckplein 1, NL-2517 GK The Hague (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ASSAYS, ANTIBODIES, AND STANDARDS FOR DETECTION OF OXIDIZED AND MDA-MODIFIED LOW DENSITY LIPOPROTEINS (57) Abstract Immunoassays for malondialdehyde-modified low density lipoprotein (MDA-modified LDL) and oxidized low density lipoprotein (OxLDL), monoclonal antibodies (and the cell lines for them) for use in the assays, and a storage-stable standard (which may be used as a calibrator and/or control) are disclosed. MDA-modified LDL and OxLDL are implicated in atherosclerosis and its etiology.		

WO 98/59248

PCT/EP97/03493

**ASSAYS, ANTIBODIES, AND STANDARDS FOR DETECTION OF
OXIDIZED AND MDA-MODIFIED LOW DENSITY LIPOPROTEINS**

BACKGROUND

The present invention relates to assays, antibodies (particularly monoclonal antibodies), and standards for detection (i.e., determination of the presence and/or quantitation of the amount) of oxidized
5 low density lipoprotein (OxLDL) and malondialdehyde-modified low density lipoprotein (MDA-modified LDL) in samples, the samples typically being derived from body fluids or tissues.

Lipoproteins are multicomponent complexes of
10 protein and lipids. Each type of lipoprotein has a characteristic molecular weight, size, chemical composition, density, and physical role. The protein and lipid are held together by noncovalent forces.

Lipoproteins can be classified on the basis of
15 their density as determined by ultracentrifugation. Thus, four classes of lipoproteins can be distinguished: High Density Lipoproteins (HDL), Intermediate Density Lipoproteins (IDL), Low Density Lipoproteins (LDL), and Very Low Density Lipoproteins (VLDL).

20 The purified protein components of a lipoprotein particle are called apolipoproteins (apo). Each type of lipoprotein has a characteristic apolipoprotein composition. In LDL the prominent apolipoprotein protein is apo B-100. Apo B-100 is one of
25 the longest single chain polypeptides known and consists of 4536 amino acids. Of these amino acids the lysine residues or moieties (there are 356 such lysine residues or moieties) can be substituted or modified by aldehydes (e.g., malondialdehyde).

30 Oxidation of the lipids in LDL (whether in vitro, e.g., by copper-induced oxidation, or whether in vivo) results in the generation of reactive aldehydes, which can then interact with the lysine residues or moieties of apo B-100. The outcome of this lysine
35 substitution or modification is that the resulting OxLDL, which is also MDA-modified LDL, is no longer recognized

21766-897

2

by the LDL receptor at the surface of fibroblasts but by scavenger receptors at the surface of macrophages. At least 60 out of the 356 lysines (or lysine residues or moieties) of apo B-100 have to be substituted in order to be recognized by the scavenger receptors (see document number 1 of the documents listed near the end of this application). The uptake of such OxLDL by macrophages results in foam cell generation, which is considered to be an initial step in atherosclerosis.

Endothelial cells under oxidative stress (e.g., in acute myocardial infarction patients) and activated blood platelets also produce aldehydes, which interact with the lysine moieties in apo B-100, resulting in the generation of aldehyde-modified LDL that is also recognized by the scavenger receptors. However, the lipids in this aldehyde-modified LDL are not oxidized. Enzymatic activity in macrophages (e.g. myeloperoxidase) results in the oxidation of both the lipid and the protein moieties of LDL. All these pathways result in aldehyde-type modification of the protein moiety of LDL.

In vitro experiments and experiments in animal models have suggested that OxLDL and/or aldehyde-modified LDL may contribute to the progression of atherosclerosis by inducing endothelial dysfunction, foam cell generation, smooth muscle cell proliferation, and platelet activation (for review see document number 2). A positive correlation between the levels of autoimmune antibodies that cross-react with aldehyde-modified LDL and the progression of carotid atherosclerotic lesions in patients suggested that OxLDL and/or aldehyde-modified LDL might contribute to the progression of human atherosclerosis (see document 3).

However, the possibility that the autoimmune antibodies were directed against other aldehyde-modified proteins, e.g., albumin, could not be excluded. Therefore, the contribution of OxLDL and

aldehyde-modified LDL (whether or not resulting from oxidation of the lipid moiety) to human atherosclerosis may be able to be established when non-invasive tests that are specific for these substances (i.e., have high affinity for those substances in preference to other substances) become available.

Because the underlying mechanisms of oxidation of LDL may be different in different patient populations (e.g., in diabetes patients, chronic renal failure patients, heart transplant patients) and because at least some of the mechanisms may be independent of lipid oxidation, such tests should be specific for both OxLDL and aldehyde-modified LDL (e.g., MDA-modified LDL) and thus preferentially be based on the detection of conformational changes that specifically occur in the apo B-100 moiety of LDL following aldehyde-type substitution of lysine residues. In other words, there is a need for such non-invasive tests (i.e., assays) that are highly specific for the analytes of interest (i.e., MDA-modified LDL and OxLDL). There is also a need for antibodies that are specific for the analytes of interest. There is also a need for a stable standard (e.g., to be used as calibrator and/or control) for the assays.

SUMMARY OF THE INVENTION

An invention satisfying those needs and having other features and advantages that will be apparent to those skilled in the art has now been developed. The present invention provides antibody-based assays that are capable of specifically quantitating (quantifying) both OxLDL and aldehyde-modified LDL or MDA-modified LDL in samples, e.g., samples derived from body fluids (like plasma or serum) or tissues. The present invention also provides monoclonal antibodies useful in those assays and cell lines (hybridomas) that produce those antibodies. The present invention also provides a storage-stable standard, which can be used as a calibrator and as a control for the assays. Having such a standard is

21766-897

4

necessary for having reliable and reproducible and therefore useful assays.

According to the present invention, there is provided an immunological assay for the detection and/or
5 quantification of human MDA-modified LDL and human OxLDL in a sample derived from the body fluids or tissues of a human being, said assay comprising: (a) contacting the sample with a first antibody that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL and human OxLDL;
10 and (b) thereafter visualizing and/or quantifying a binding reaction between the first antibody and the MDA-modified LDL and OxLDL present in the sample; wherein the MDA-modified LDL and OxLDL for which the first antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contain at least 60 substituted
15 lysine moieties per apo B-100 moiety.

That assay may, for example, be a competitive assay, a sandwich assay, an immunohistochemical assay, etc. "competitive assays" are well-known and any competitive assay may be used in this invention provided it is within
20 the limitations of the invention and that the benefits of the invention can be achieved. "Sandwich assays" are well-known and any sandwich assay may be used in this invention provided it is within the limitations of the invention and that the benefits of the invention can be achieved.
25 "Immunohistochemical assays" are well-known and any immunohistochemical assay may be used in this invention provided it is within the limitations of the invention and that the benefits of the invention can be achieved.

According to the present invention, there is
30 further provided an immunological sandwich assay for the detection and/or quantification of human MDA-modified LDL in a sample derived

21766-897

4a

from the body fluids or tissues of a human being in which assay a first antibody that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL is bound to a substrate, said assay comprising: (a) contacting the sample
5 with the substrate having bound to it the first antibody under binding conditions so that human MDA-modified LDL in the sample will bind to the first antibody; (b) thereafter removing unbound sample from the substrate; (c) thereafter contacting the substrate with a second antibody that has an
10 affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL; and (d) thereafter visualizing and/or quantifying the MDA-modified LDL that was present in the sample; wherein the MDA-modified LDL for which the first antibody and the second antibody have an affinity of at least
15 about $5 \times 10^8 \text{ M}^{-1}$ contains at least 60 substituted lysine moieties per apo B-100 moiety.

21766-897

5

As used herein (including the claims), "high
15 affinity" means an affinity constant (association
constant) of at least about $5 \times 10^8 \text{ M}^{-1}$, desirably at least
about $1 \times 10^9 \text{ M}^{-1}$, preferably at least about $1 \times 10^{10} \text{ M}^{-1}$,
and most preferably of at least about $1 \times 10^{11} \text{ M}^{-1}$. As used
herein (including the claims), "low affinity" means an
20 affinity constant (association constant) of less than
about $1 \times 10^7 \text{ M}^{-1}$, desirably less than about $1 \times 10^6 \text{ M}^{-1}$, and
preferably less than about $1 \times 10^5 \text{ M}^{-1}$. Affinity constants
are determined in accordance with the appropriate method
described in Holvoet et al. (4).

25 The antibodies that can be used in this
invention will bind with MDA-modified LDL and/or OxLDL
whose apo B-100 moieties contain at least 60, desirably
at least about 90, more desirably at least about 120,
preferably at least about 180, more preferably at least
30 about 210, and most preferably at least about 240
substituted lysine residues per apo B-100 moiety. The
range of lysine substitution will generally be from 60 to
about 240 and preferably from about 120 to about 240
substituted lysine moieties per apo B-100 moiety.

35 Each new monoclonal antibody is highly specific
for a conformational epitope that is present when at
least about 60, preferably at least about 120 lysine
residues, are substituted and by virtue thereof can

distinguish various markers or indications related to atherosclerosis. Antibodies recognizing epitopes present when less than about 60 lysines are substituted or modified are less specific but are still useful (e.g., they may be used as the secondary antibody in a sandwich ELISA).

The preferred antibodies used herein are monoclonal antibodies mAb-4E6, mAb-1h11, and mAb-8A2. Their affinity constants for native LDL, MDA-modified LDL, and OxLDL are as follows:

<u>Antibody</u>	<u>Native LDL</u>	<u>MDA-modified LDL</u>	<u>OxLDL</u>
mAb-4E6	less than 1×10^6	3×10^{10}	2×10^{10}
15 mAb-1H11	less than 1×10^6	3×10^{10}	less than 1×10^6
mAb-8A2	5×10^9	1×10^{10}	1×10^{10}

In yet another aspect, the present invention concerns (a) monoclonal antibody mAb-4E6 produced by hybridoma Hyb4E6 deposited at the BCCM under deposit accession number LMBP 1660 CB on or about April 24, 1997, (b) monoclonal antibody mAb-8A2 produced by hybridoma Hyb8A2 deposited at the BCCM under deposit accession number LMBP 1661 CB on or about April 24, 1997, (c) hybridoma Hyb4E6 deposited at the BCCM under deposit accession number LMBP 1660 CB on or about April 24, 1997, and (d) hybridoma Hyb8A2 deposited at the BCCM under deposit accession number LMBP 1661 CB on or about April 24, 1997.

The antibodies used in the assays of this invention are preferably those two (i.e., mAb-4E6 and mAb-8A2) as well as mAb-1H11. The cell line for antibody mAb-1H11 is produced by hybridoma Hyb1H11, which was deposited at the BCCM under deposit accession number LMBP 1659 CB on or about April 24, 1997.

The BCCM is the Belgian Coordinated Collections of Microorganisms authorized by the "Budapest Treaty of

28 April 1977 on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure." Its address is c/o The University of Gent, K. Ledeganckstraat 35, B-9000 Gent, Belgium.

5 The assay may be of a type that is well-known, such as an Enzyme-Linked Immunosorbent Assay (ELISA). For example, in the case of a sandwich ELISA, mAb-4E6 (for MDA-modified LDL and OxLDL) or mAb-1H11 (for MDA-modified LDL) may be bound to a solid substrate and subsequently
10 contacted with a sample to be assayed. After removal of the sample, binding between the specific antibody and OxLDL and/or MDA-modified LDL captured out of the sample can be visualized and/or quantified by detection means. Detection means may be a labeled, less specific secondary
15 antibody that recognizes a different part of the apo B-100 moiety of the captured analyte (e.g., mAb-8A2).

 In the case of a competitive ELISA, a solid substrate coated with OxLDL or MDA-modified LDL may be contacted for a predetermined period of time with the
20 monoclonal antibody mAb-4E6 and a sample thought or known to contain OxLDL and/or MDA-modified LDL, after which period of time unbound antibody and sample are removed and a binding reaction between antibody and OxLDL and/or MDA-modified LDL bound to the substrate is visualized
25 and/or quantified. Quantification in a competitive ELISA is indirect because the binding between the antibody and the analyte in the sample is not measured but instead the amount of antibody that binds to the known amount of OxLDL or MDA-modified LDL that is coated on (bound to)
30 the substrate is measured. The more antibody bound to the known amount of OxLDL or MDA-modified LDL coated on the substrate, the less analyte there was in the sample.

21766-897

7a

According to the present invention, there is further provided a method of standardizing an assay for human MDA-modified LDL and human OxLDL by using as a calibrator or as a control a stable standard containing

5 MDA-modified LDL whose extent of substitution of its lysine moieties will remain essentially constant over normal periods of time during normal storage for biological materials, the MDA-modified LDL of said standard being made by contacting malondialdehyde with LDL at a predetermined

10 molar ratio of malondialdehyde to the apo B-100 moiety of the LDL, the standard containing an agent that reduces the ability of any metal ions present to catalyze oxidation of the LDL and/or an anti-oxidant.

“Over normal periods of time during normal
5 storage for biological materials” as used herein refers
to the time periods and conditions under which biological
materials to be used in assays and other laboratory work
are typically stored. Those conditions will typically
include low temperature and in appropriate cases
10 freezing, either with or without lyophilization.
Depending on the particular biological material, if the
material is stored under the appropriate temperature and
other conditions (e.g., lack of vibration or other
movement, proper humidity), the material may be stable
15 for at least three months, desirably for over a year,
preferably for over two years, and most preferably for
over three years.

The standard preferably contains an agent that
reduces the ability of any metal ions present to catalyze
20 oxidation of the LDL (e.g., a chelating agent, such as
EDTA) and/or one or more anti-oxidants (e.g., BHT and/or
Vitamin E). Preferably both the agent that reduces the
ability of any metal ions present to catalyze oxidation
of the LDL and the anti-oxidant are used. It has
25 surprisingly been found that when using an antibody that
is specific for both OxLDL and MDA-modified LDL, the
storage-stable standard of this invention (containing
MDA-modified LDL and not OxLDL) can be used. That
eliminates the need to try to formulate, store, and use a
30 stable standard containing OxLDL. OxLDL may continue to
oxidize under typical storage conditions, making using as
a standard a composition containing OxLDL difficult if
not almost impossible. EDTA will typically be used in
concentrations of 0.5 to 5 mM, preferably in
35 concentrations of 0.5 to 2 mM. BHT will typically be used
in concentrations of 5 to 50 μ M, preferably in
concentrations of 10 to 20 μ M. Vitamin E will typically
be used in concentrations of 5 to 50 μ M, preferably in

concentrations of 10 to 20 μ M. The standard may also contain anti-platelet agents and coagulation inhibitors.

It has been found that LDL that has been modified by treatment with MDA is highly stable. Such MDA-modified LDL (which is not oxidized, i.e., its lipid moiety is not oxidized) could be added to reference plasma samples and those samples could be frozen and thawed without increasing the extent of lysine substitution. Because the total number of lysine residues in all apo B-100 molecules is identical, a constant MDA/apo B-100 molar ratio in the reaction mixture will result in an identical number of substituted lysines in the MDA-modified LDL. In contrast, for example, metal-ion mediated oxidation of LDL ultimately results in a variable extent of lysine substitution because it depends on the oxidation sensitivity of the LDL preparation, which by itself depends on fatty acid composition and antioxidant content, which are highly variable even in healthy control individuals.

As described below, a correlation between the oxidation of LDL and the extent of post-transplant atherosclerosis in heart transplant patients was established using this invention. The relationship between endothelial injury and the modification of LDL was established in chronic renal failure patients that are at high risk for atherosclerotic cardiovascular disease. It was also demonstrated that endothelial injury is an initial step in atherosclerosis.

Based on the characteristics of the oxidatively modified LDL from the plasma of heart transplant and chronic renal failure patients, it was concluded that cell-mediated aldehyde modification independent of lipid oxidation was at least partially involved. This finding further supported the hypothesis that an assay for oxidatively modified LDL has to detect both OxLDL and aldehyde-modified LDL.

21766-897

10

According to the present invention, there is further provided a kit for conducting a sandwich assay for the determination of human OxLDL or human MDA-modified LDL or both in a sample derived from the body fluids or tissues of a human being, said kit comprising: (a) a substrate on which is bound a first antibody that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human OxLDL or human MDA-modified LDL or both, the OxLDL and MDA-modified LDL each having at least 60 substituted lysine moieties per apo B-100 moiety, and (b) a labeled antibody having an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human OxLDL that becomes bound to the first antibody during the assay or for human MDA-modified LDL that becomes bound to the first antibody during the assay or for both that become bound to the first antibody during the assay. Preferably the kit further comprises a reactive substance for reaction with the labeled antibody (e.g., an enzyme) to give an indication of the presence of the labeled antibody. Preferably the kit also comprises the stable standards, e.g., in the form of stable calibrators and/or stable controls. Thus, e.g., the bound antibody may be mAb-4E6 or mAb-1H11 and the labeled antibody may be mAb-8A2.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings are provided to help further describe the invention, which drawings are as follows:

Figure 1 illustrates the correlation between amounts of oxidized and MDA-modified LDL in coronary lesions in Watanabe heritable hyperlipidemic rabbits (A) and in miniature pigs (B).

Figure 2 illustrates the accumulation of OxLDL and MDA-modified LDL in coronary arteries of cardiac explants of

21766-897

10a

ischemic heart disease but not of dilated cardiomyopathy patients.

Figure 3 illustrates the inhibition of the binding of mAb-4E6 to immobilized OxLDL by native LDL, OxLDL and
5 MDA-modified LDL in solution.

Figure 4 illustrates a typical standard curve obtained with MDA-modified LDL in sandwich ELISA.

Figure 5 illustrates levels of OxLDL and aldehyde-modified LDL in posttransplant plasma samples of heart transplant patients with different extents of angiographically assessed coronary artery stenosis.

5 Figure 6 illustrates the correlation between plasma levels of OxLDL and aldehyde-modified LDL and titers of specific autoantibodies.

Figure 7 illustrates the correlation between plasma levels of OxLDL and aldehyde-modified LDL and of
10 von Willebrand factor antigen.

These drawings are provided for illustrative purposes only and should not be used to unduly limit the invention.

15 **DETAILED DESCRIPTION OF THE INVENTION**

The present invention will be further described in conjunction with the following examples, which are for illustrative purposes and which should not be used to unduly limit the invention.

20

EXAMPLES

EXAMPLE 1

Preparation and characterization of antibodies specific for OxLDL and for aldehyde-modified LDL

25 Balb/c mice were immunized by intravenous and intraperitoneal injection of either OxLDL or MDA-modified LDL. OxLDL was obtained by in vitro incubation of LDL (final apo B-100 concentration 700 µg/ml) with copper chloride (final concentration 640 µM) for 16 h at 37°C.
30 MDA-modified LDL was prepared by incubation of LDL (final apo B-100 concentration 700 µg/ml) with a 0.25 M MDA-solution for 3 h at 37°C. The numbers of substituted lysines, measured in the TBARS assay, was typically 210 per apo B-100 molecule for OxLDL and 240 for MDA-modified
35 LDL. Hybridomas were obtained by PEG induced fusion of spleen lymphocytes derived from immunized mice with P3-X63/Ag-6.5.3 myeloma cells according to standard techniques (4). The screening for hybridomas producing

21766-897

12

specific antibodies was performed with ELISA using microtiter plates coated with malondialdehyde-modified LDL or copper-oxidized LDL. 308 hybridomas were obtained after immunization of mice with either OxLDL (211) or 5 MDA-modified LDL (97). Hyb4E6 produced antibodies specific for both malondialdehyde-modified and copper-oxidized LDL (mAb-4E6), and Hyb1H11 produced antibodies specific for malondialdehyde-modified LDL (mAb-1H11) alone. The IgG fraction of the antibodies was purified by 10 affinity chromatography on protein A-Sepharose* and the affinity of the purified IgGs was determined in a solid phase radioimmunoassay and/or in ELISA. The K_d values of the monoclonal antibody mAb-4E6 were $< 10^6 \text{ M}^{-1}$ for native LDL and $> 10^9 \text{ M}^{-1}$ for malondialdehyde-modified LDL and 15 copper-oxidized LDL. The K_d values for the monoclonal antibody mAb-1H11 were $< 10^6 \text{ M}^{-1}$ for both LDL and OxLDL and $> 10^9 \text{ M}^{-1}$ for malondialdehyde-modified LDL. The K_d values for the monoclonal antibody mAb-8A2, obtained after immunization of mice with LDL, were $> 10^9 \text{ M}^{-1}$ for all LDL 20 forms. Delipidation of MDA-modified LDL and OxLDL resulted in a loss of the immunoreactivity of mAb-4E6, suggesting that it is directed against a conformational epitope in the protein moiety of oxidatively modified LDL.

25

EXAMPLE 2

Use of mAb-4E6 for the quantitation of OxLDL and MDA-
modified LDL in coronary lesions of Watanabe heritable
hyperlipidemic rabbits and miniature pigs on a
30 cholesterol rich diet

Coronary arteries were obtained from 2 and 5 month old Watanabe heritable hyperlipidemic rabbits (n=30) on normal chow or from miniature pigs (n=26) which were fed a diet enriched in cholesterol (4%), saturated 35 fat (14% beef tallow) and bile extract (1%) for 6 to 24 weeks.

Arterial specimens were submerged within 30 min after removal in PBS (pH 7.4) containing 4% sucrose, 20
*Trade-mark

μ M vitamin E and 10 μ M butylated hydroxytoluene as antioxidants, and 1 mM EDTA, snap-frozen in liquid nitrogen and stored at -80°C . Frozen 7 μ M sections were stained with hematoxylin and eosin and with oil red O or immunostained as described below. Morphometric parameters of atherosclerotic lesions were measured by planimetry using the Leica 2 Quantimet color image analyzer (Cambridge, UK). The area within the external elastic lamina, the internal elastic lamina and the lumen were measured. Media was defined as the area between the internal and external elastic lamina. Intima was defined as the area within the internal elastic lamina not occupied by vessel lumen.

Oxidized apo B-100 containing lipoproteins were detected with the specific monoclonal antibody mAb-4E6, alkaline-phosphatase conjugated rabbit-anti-mouse IgG antibodies and the fuchsin alkaline phosphate substrate system (Dako, Carpinteria, CA), and the absorbance was measured in the color image analyzer. Specificity of immunostaining was confirmed by inhibition of staining with excess of copper-oxidized LDL but not with native LDL or with malondialdehyde-modified albumin. The staining co-localized with that monoclonal antibody mAb-13F6, specific for apo B-100. Absorbance (approximately 10%) measured with excess copper-oxidized LDL was presumed to represent background staining.

Figure 1 illustrates the correlation between the levels of oxidized apo B-100 containing lipoproteins, i.e. OxLDL and MDA-modified LDL, in the lesions and the mean intimal area of coronary lesions in Watanabe hyperlipidemic rabbits (A) and in miniature pigs (B). Those data thus demonstrate a correlation between the accumulation of OxLDL and MDA-modified LDL and the progression of coronary atherosclerotic lesions in 2 different animal models. In Watanabe rabbits the progression of the lesions is due to the increase of LDL cholesterol associated with the heritable LDL receptor

deficiency, whereas the progression in miniature pigs is due to a diet-induced increase in LDL cholesterol.

EXAMPLE 3

5 Immunohistochemistry

1. Introduction

This example is a typical example of the use of the highly specific antibody mAb-4E6 in immunohistochemistry applied to human atherosclerotic lesions. In a
10 similar manner corresponding experiments may be performed, for which certain conditions can be adapted by the skilled person using his common knowledge in the field.

15 2. Material and methods

Coronary artery specimens, obtained at the time of transplantation from patients with ischemic heart disease (n= 7) or dilated cardiomyopathy (n= 7), were treated as described earlier (document 7). The specimens
20 were collected within 30 min after removal of the heart in PBS (pH 7.4) containing 4% sucrose, 20 μ M vitamin E and 10 μ M butylated hydroxytoluene as antioxidants, and 1 mM EDTA, and were stored at -80°C. Frozen 7 μ m thick sections were cut and stained with hematoxylin and eosin.
25 Six to 8 sections at a distance of 84 μ m were analyzed for each specimen to insure representative results. Duplicate slides were developed with monoclonal antibodies mAb-4E6, specific for oxidized LDL, PG-M1, specific for human macrophages, or 1A4, specific for
30 human smooth muscle α -actin (both from Dako SA, Glostrup, Denmark). Specificity of binding of mAb-4E6 was confirmed by its inhibition with OxLDL but not with native LDL.

3. Results

35 Coronary artery segments of 7 individuals with pretransplant dilated cardiomyopathy did not contain atherosclerotic lesions and the monoclonal antibody did not detect OxLDL and/or aldehyde-modified LDL in these

segments. Coronary artery segments of 7 patients with pretransplant ischemic heart disease all contained atherosclerotic lesions which contained OxLDL and/or aldehyde-modified LDL (Figure 2). This information is
5 sufficient to state that the antibody detects OxLDL in atherosclerotic lesions in a highly specific manner.

OxLDL was associated with macrophage foam cells (preferentially in lesions with < 50% stenosis), with smooth muscle foam cells and with the necrotic lipid core
10 (preferentially in lesions with > 50% stenosis).

Macrophages and smooth muscle cells were identified by immunostaining with specific monoclonal antibodies (5). These data supported the hypothesis that oxidation of LDL may be associated with the development of ischemic
15 coronary artery disease. The monoclonal antibody mAb-4E6 of the present invention that detected the immunoreactive material in the tissue sections was then further used in ELISA (cf. Example 4).

20 4. Legend to Figure 2

Light micrographs (a, c, e; x 40) and phase contrast micrographs (b, d, f; x 400) of representative left anterior descending coronary artery specimens of a patient with dilated cardiomyopathy (male; 40 years of
25 age) (a, b) and of a patient with ischemic heart disease (male; 57 years of age) (c-f). Tissue sections were immunostained with the monoclonal antibody mAb-4E6. Oxidized LDL was undetectable in the neointima of the first patient (a, b), but demonstrable in plaques of the
30 second patient. The oxidized LDL was associated with macrophage foam cells that infiltrated at the shoulder areas of fibrous plaques (c, d) and with smooth muscle foam cells in fibrous caps (e, f).

EXAMPLE 4Competitive ELISA

1. Introduction

According to the invention an ELISA was
 5 established for the quantitation of OxLDL and aldehyde-
 modified LDL in plasma. It was based on the inhibition of
 the binding of mAb-4E6 to the wells of microtiter plates
 coated with copper-oxidized LDL. This antibody was
 obtained as described in Example 1.

10

2. Material and methods

Standard OxLDL and aldehyde-modified LDL and
 plasma samples were diluted in PBS containing 1 mM EDTA,
 20 μ M vitamin E, 10 μ M butylated hydroxytoluene, 20 μ M
 15 dipyridamole and 15 mM theophylline to prevent in vitro
 LDL oxidation and platelet activation. Equal volumes of
 diluted purified mAb-4E6 solution (final concentration
 7.5 ng/ml) and of diluted standard solution (copper-
 oxidized LDL added as competing ligand at a final
 20 concentration ranging from 50 to 500 ng/ml) were mixed
 and incubated for 30 min at room temperature. Then 200 μ l
 aliquots of the mixtures were added to wells coated with
 MDA-modified LDL or OxLDL.

Samples were incubated for 2 h at room
 25 temperature. After washing, the wells were incubated for
 1h with horse-radish peroxidase conjugated rabbit IgG
 raised against mouse immunoglobulins and washed again.
 The peroxidase reaction was performed as described
 earlier (5) and the absorbance (A) was read at 492 nm.

30 Controls without competing ligand and blanks
 without antibody were included routinely. The percent
 inhibition of binding of mAb-4E6 to the immobilized
 ligand was calculated as:

$$35 \frac{A^{492nm}_{control} - A^{492nm}_{sample}}{A^{492nm}_{control} - A^{492nm}_{blank}}$$

and standard curves were obtained by plotting the
 percentage of inhibition vs the concentration of
 competing ligand.

The lower limit of detection was 0.020 mg/dl in undiluted human plasma. Intra- and interassay coefficients of variation were 10 and 12%, respectively. Standard OxLDL and aldehyde-modified LDL and plasma samples were diluted in PBS containing antioxidants and antiplatelet agents as described above.

3. Results

The specificity of mAb-4E6 for OxLDL and aldehyde-modified LDL is illustrated in Figure 3. 50% inhibition of binding of mAb-4E6 to immobilized OxLDL and aldehyde-modified LDL was obtained with 0.025 mg/dl copper-oxidized LDL and 25 mg/dl native LDL, respectively. The C_{50} value, i.e., the concentration that is required to obtain 50 % inhibition of antibody binding, increased from 2.5 mg/dl for MDA-modified LDL with 60 substituted lysine residues per apo B-100 molecule to 0.025 mg/dl for MDA-modified LDL with 240 substituted lysine residues per apo B-100 molecule (Figure 3). Copper-oxidation resulted in fragmentation of the apo B-100 moiety but did not abolish the binding of mAb-4E6 (Figure 3). 50-fold higher molar concentrations of MDA-modified albumin were required to obtain 50% inhibition (not shown), whereas up to 1,000-fold higher molar concentrations of MDA-modified lysine did not affect mAb-4E6 binding. OxLDL and aldehyde-modified LDL isolated from patient plasma had the same reactivity as MDA-modified LDL with 120 substituted lysines and as copper-oxidized LDL with 210 substituted lysines. Intra- and interassay coefficients of variation were 10 and 12%, respectively. When copper-oxidized LDL were added to human plasma at a final concentration of 0.25 and 2 mg/dl, respectively, recoveries were 95 and 105%, respectively.

4. Legend to Figure 3

Interaction of mAb-4E6 with competing ligands in solution. Copper-oxidized LDL (1 μ g/ml) was the plated

antigen. mAb-4E6 was added in the absence and in the presence of competing ligands: copper-oxidized LDL (∇), MDA-modified LDL with 240 ($\blacksquare$), 120 ($\diamond$), 90 (O) and 60 ($\bullet$) blocked or substituted or modified lysines per apo B-100, respectively, native LDL ($\blacktriangle$), and OxLDL and aldehyde-modified LDL ($\blacklozenge$) isolated from the plasma of severe chronic renal failure patients. Results are expressed as B/B_0 where B_0 is the amount of mAb-4E6 bound in the absence and B that amount bound in the presence of competing ligand.

EXAMPLE 5

Sandwich ELISA

1. Introduction

According to the invention a sandwich-type ELISA was established for the quantitation of OxLDL and aldehyde-modified LDL in plasma. It was based on the binding of immunoreactive material to the wells of microtiter plates coated with the monoclonal antibody mAb-4E6 and the detection of bound immunoreactive material with the use of the monoclonal antibody mAb-8A2 labeled with peroxidase. This version of the ELISA is more suited for use in the clinical laboratory because it overcomes the need to prepare standard solutions of in vitro oxidized and/or aldehyde-modified LDL which can only be kept at -4°C for a limited period of time, typically 2 weeks. MDA-modified LDL may be added to reference plasma and those standard preparations may be stored at -80°C for up to 1 year (see above).

2. Material and methods

Standard preparations and plasma samples were diluted in PBS containing antioxidants and antiplatelet agents as described above, 180 μl aliquots of 80-fold diluted plasma and of standard solutions containing between 10 and 0.01 nM of malondialdehyde-modified LDL were applied to the wells of microtiter plates coated with mAb-4E6 (200 μl of a 4 $\mu\text{g}/\text{ml}$ IgG solution) and

incubated for 2 h at room temperature. After washing, the wells were incubated for 1 h with horseradish peroxidase conjugated mAb-8A2, IgG (final IgG concentration 65 ng/ml) and washed again. The peroxidase reaction was performed as described above. The absorbance measured at 492 nm correlates with the log-value of the aldehyde-modified LDL concentration in the range between 1.5 nM and 0.3 nM.

3. Legend to Figure 4

Standard curves for the sandwich ELISA mAb-4E6 was the plated antibody. MDA-modified LDL was the ligand. Bound MDA-modified LDL was detected with mAb-8A2 conjugated to horse radish peroxidase. MDA-modified LDL was added to 8 different plasma samples to a final concentration of 100 nM and further diluted in buffer to final concentrations ranging from 2 to 0.2 nM.

EXAMPLE 6

Use of the ELISA in diagnosis of posttransplant coronary artery disease

1. Introduction

The ELISA of the present invention was used to study the association between plasma levels of OxLDL and aldehyde-modified LDL and posttransplant coronary artery disease.

2. Material and methods

2.1. Patients

The posttransplant study group contained 47 patients transplanted for dilated cardiomyopathy and 60 patients treated for ischemic heart disease. The clinical characteristics of these patients are summarized in Table 1. At the time of blood sampling, between 12 and 84 months after surgery, all patients were in a stable cardiac condition without evidence of acute rejection. From 14 patients (7 dilated cardiomyopathy and 7 ischemic heart disease patients) coronary arteries of cardiac

explants were isolated and studied by immunohistochemistry (as demonstrated in Example 3). Adequate information about smoking habits was available for 92 of the 107 patients (16 smokers and 76 non-smokers). There was no adequate information about smoking habits of donors. Blood samples of 53 non-smoking controls (25 males/28 females; age: 52 ± 1.3 years) without a history of atherosclerotic cardiovascular disease were obtained. The controls were matched for age, gender and levels of LDL cholesterol. They were selected from the laboratory and clinical staff.

2.2. Coronary angiography

Routine annual coronary angiograms were available for all posttransplant patients at the time of blood sampling. Coronary artery disease was independently assessed by two angiographers who were unaware of the OxLDL and aldehyde-modified LDL levels and was visually graded as follows:

- grade 0: normal coronary arteries
- grade I: minor abnormalities with < 50% stenosis of primary or secondary branches and normal left ventricular function
- grade II: $\geq 50\%$ stenosis of primary or secondary branches, or distal involvement with impaired left ventricular function.

It is well known that angiography systematically underestimates the extent of coronary intimal thickening in cardiac transplant recipients. This study therefore does not attempt to accurately quantify the coronary artery disease in our patients. Rather the subdivision in groups defined above relies on angiographic data that are easily distinguishable and that have been shown to correlate with histopathologic findings. Out of 107 patients, 46 patients had a normal coronary angiogram 3 years before and development of angiographic, coronary artery disease within a 3 year follow-up period was assessed in all these patients. The reference normal

coronary angiogram was the first post-operative angiogram in 18 patients, the second in 14 patients and the third in 14 patients.

The study was approved by the Institutional Review Board and the study subjects provided informed consent.

2.3. Blood sampling

Venous blood samples from patients and controls were collected on 0.1 vol of 0.1 M citrate, containing 1 mM EDTA, 20 μ M vitamin E, 10 μ M butylated hydroxytoluene, 20 μ M dipyridamole and 15 mM theophylline to prevent in vitro LDL oxidation and platelet activation. Blood samples were centrifuged at 3,000 g for 15 min at room temperature within 1 h of collection and stored at -20°C until the assays were performed.

2.4. Lipoproteins: isolation and modification

LDL were isolated from pooled sera of fasting normolipidemic donors by density gradient ultracentrifugation (document 6). Standard preparations of MDA-modified and copper-oxidized LDL were prepared as described elsewhere (7, 8) and were used as assay controls. Apo B-100 molecules of in vitro MDA-modified LDL (7) and of copper-oxidized LDL (8) contained on average 244, and 210 substituted lysines (out of a total of 356), respectively (5, 9). Whereas the extent of lysine-substitution in in vitro MDA-modified LDL and copper-oxidized LDL is very similar, the lipid moiety of the former is not oxidized. Specificity of the monoclonal antibody mAb-4E6 for both MDA-modified LDL and copper-oxidized LDL suggests that it depends on the extent of protein (lysine) modification only. All lipoprotein concentrations were therefore expressed in terms of protein. OxLDL and aldehyde-modified LDL isolated from the plasma of patients were characterized as described previously (5, 9).

2.5. Assays

Cholesterol and triglycerides were measured by enzymatic methods (Boehringer Mannheim, Meylon, France). Typing of major histocompatibility complex class I (HLA-B) and class II (HLA-DR) antigen was performed by the microlymphocytotoxicity technique.

The ELISA of the invention was used to detect OxLDL and aldehyde-modified LDL.

10 2.6. Statistical analysis

Controls and patients were compared by ANOVA test followed by nonparameteric Mann-Whitney or Dunnett's multiple comparison test on logarithmically transformed values, in the Instat^{*} V2.05a statistical program (Graph Pad Software, San Diego, CA). Non-quantitative parameters were compared by Chi-square analysis. OxLDL and aldehyde-modified LDL levels measured in 3 aliquots of the same plasma samples were compared in Friedman nonparametric repeated measures test. Logistic regression analysis, using the SAS software (SAS Institute Inc., USA), was performed to evaluate the correlation between angiographically assessed coronary artery stenosis (as dependent variable) and plasma levels of OxLDL and aldehyde-modified LDL, age and sex of recipients, age and sex of donors, pretransplant history of ischemic heart disease or dilated cardiomyopathy, duration of ischemia, length of follow up, number of rejections, number of HLA-mismatches, cytomegalovirus infection, hypertension (antihypertensive treatment), diabetes, treatment with lipid lowering drugs (statins or fibrates) and serum levels of LDL cholesterol, HDL cholesterol and triglycerides as independent variables. p-values of less than 0.05 were considered to indicate statistical significance. Logistic regression analysis was also performed to evaluate the correlation between plasma levels of OxLDL and aldehyde-modified LDL and development of coronary artery stenosis during a 3-year follow-up period.

*Trade-mark

3. Results

The correlation between OxLDL and aldehyde-modified LDL and coronary artery stenosis was evaluated in 47 patients transplanted for dilated cardiomyopathy and in 60 patients treated for ischemic heart disease. Analysis of clinical data for the two groups of heart transplant patients (Table 1) revealed no significant differences in age and gender of the recipients, age and gender of donors, duration of ischemia of the donor heart, number of rejection episodes, number of HLA-mismatches, frequency of Cytomegalovirus infections, hypertension or diabetes, and grade of coronary artery stenosis. Patients transplanted for ischemic heart disease were followed longer and received more frequently lipid lowering drugs (Table 1).

Analysis of the laboratory data (Table 2) revealed no significant differences in serum levels of triglycerides, HDL cholesterol and LDL cholesterol between groups of patients or between patients and controls. However, significant differences in levels of OxLDL and aldehyde-modified LDL were observed. Mean plasma levels of OxLDL and aldehyde-modified LDL were 1.3 ± 0.14 mg/dl in dilated cardiomyopathy patients ($p < 0.001$ vs controls) and 1.7 ± 0.13 mg/dl in ischemic heart disease patients ($p < 0.001$ vs controls and < 0.01 vs dilated cardiomyopathy patients) (Table 2). Plasma levels of OxLDL and aldehyde-modified LDL in control subjects matched for age, gender and serum levels of triglycerides, HDL cholesterol and LDL cholesterol were 0.60 ± 0.034 mg/dl ($n = 53$; $p < 0.001$ vs both transplanted dilated cardiomyopathy and ischemic heart disease patients).

Levels of OxLDL and aldehyde-modified LDL were not different in samples that were stored for 24 h to 4 months after collection, and up to four thawing and freezing cycles did not cause an increase of OxLDL and aldehyde-modified LDL levels. These findings indicated that the addition of EDTA, antioxidants and anti-platelet

agents adequately prevented the in vitro oxidation of LDL. In a subset of 87 consecutive plasma samples levels of OxLDL and/or aldehyde-modified LDL were measured in 3 separate aliquots on 3 different days. The levels were
5 1.30 $\pm$ 0.074 mg/dl, 1.48 $\pm$ 0.101 mg/dl and 1.46 $\pm$ 0.090 mg/dl, respectively. Friedman nonparametric repeated measures test revealed no significant differences.

Mean OxLDL and aldehyde-modified LDL levels were 1.2 $\pm$ 0.053 mg/dl (n= 79) in posttransplant samples
10 of patients with angiographically normal coronary arteries (grade 0), 2.1 $\pm$ 0.30 mg/dl in patients with grade I coronary artery stenosis (n= 18; p< 0.001 vs grade 0) and 3.2 $\pm$ 0.45 mg/dl in patients with grade II coronary artery stenosis (n= 10; p< 0.001 vs grade 0 and
15 p< 0.05 vs grade I) (Figure 5). Serum levels of LDL cholesterol, triglycerides and HDL cholesterol were very similar in patients with higher grade of coronary artery stenosis. Levels of OxLDL and aldehyde-modified LDL in plasma samples of patients transplanted for dilated
20 cardiomyopathy or ischemic heart disease, with the same grade of coronary artery stenosis, were similar: 1.1 $\pm$ 0.072 and 1.4 $\pm$ 0.079 mg/dl for grade 0 patients and 2.6 $\pm$ 0.60 and 2.4 $\pm$ 0.29 mg/dl for patients with higher grade of coronary artery stenosis. The number of patients
25 with elevated levels of OxLDL and aldehyde-modified LDL (> 1 mg/dl, i.e. mean levels of controls + 2 SD) were 43 (out of 60) in the subpopulation of patients with pretransplant ischemic heart disease and 21 (out of 47) in the subpopulation of patients with pretransplant
30 dilated cardiomyopathy. Forty-two out of 79 patients with angiographically normal coronary arteries had elevated levels of OxLDL and aldehyde-modified LDL. Elevated levels were detected in 12 (out of 18) patients with grade I and in all patients with grade II stenosis
35 (p=0.0046 for trend).

To allow further characterization of the immunoreactive material detected in the ELISA, LDL fractions were isolated from the plasma of all of 10

21766-897

25

patients with grade II coronary artery stenosis (18). These fractions retained $85 \pm 10 \%$ (mean $\pm$ SD) of the immunoreactive material, whereas no immunoreactive material migrated in the serum albumin position. OxLDL and aldehyde-modified LDL were isolated from isolated LDL fractions by ion-exchange chromatography on a mono Q-Sepharose* column with a recovery of 75%. The number of substituted lysines per apo B-100 molecule was 130 ± 10 for OxLDL and aldehyde-modified LDL compared to 5 ± 1 (p<0.001) for native LDL. The respective cholesterol/protein ratios were 3.3 ± 0.54 and 1.8 ± 0.36 (p<0.001). The levels of arachidonate and linoleate in OxLDL and aldehyde-modified LDL isolated from the plasma of these patients were 75 and 80 % lower than these in native LDL isolated from the same plasma samples. The inhibition curves obtained with OxLDL and aldehyde-modified LDL isolated from the plasma of heart transplant patients were superimposable with these obtained with in vitro oxidized LDL with the same extent of protein modification (120 substituted lysines per apo B-100 molecule) (Figure 3).

The protein/antigen ratio and the relative reactivity in the ELISA of OxLDL and aldehyde-modified LDL isolated from the plasma of these patients were similar to these of copper-oxidized or MDA-modified standard LDL preparations.

Logistic regression analysis (Table 3) identified 3 parameters that were significantly and independently correlated with posttransplant coronary artery stenosis including levels of OxLDL and aldehyde-modified LDL, length of follow up and donor age.

In contrast, pretransplant history of dilated cardiomyopathy or of ischemic heart disease, age and gender of recipients, gender of donors, duration of ischemia of the donor heart, extent of HLA-mismatch, number of rejections, hypertension, diabetes, and serum levels of LDL cholesterol, HDL cholesterol and triglycerides in recipients did not significantly

*Trade-mark

contribute to the individual variations in extent of coronary artery stenosis (Table 3).

Serum levels of LDL cholesterol, HDL cholesterol and triglycerides in patients were similar to these in controls (Table 2), so that higher grade of coronary artery stenosis was unlikely to depend on these variables in this study group. Fifty-six of the 107 transplant patients received lipid lowering drugs (46 with statins and 10 with fibrates) (Table 1), but the treatment with these drugs was not correlated with the incidence of angiographic graft vasculopathy (Table 3). Seventy-five (out of 107) patients were treated with calcium channel blockers. The plasma levels of OxLDL and aldehyde-modified LDL in these patients (1.53 ± 0.11 mg/dl) were very similar to these in non-treated patients ($1.74 \pm$ mg/dl) and treatment with these drugs was not correlated with the extent of coronary artery stenosis.

Development of coronary artery disease was observed in 12 out of 46 heart transplantation patients during a 3-year follow-up period. There were no differences in age and gender of recipients, age and gender of donors, duration of ischemia, extent of HLA mismatch, frequency of cytomegalovirus infections, hypertension and diabetes (Table 4) nor in serum levels of triglycerides, HDL cholesterol and LDL cholesterol (Table 5) between patients without and with development of coronary artery disease. However, levels of OxLDL and aldehyde-modified LDL were significantly elevated in patients with development of coronary artery disease (Table 5).

Logistic regression analysis revealed that plasma levels of OxLDL and aldehyde-modified LDL (Chi-square value=7.1; $p=0.0076$) and age of donor (Chi-square value=4.4; $p=0.035$) predicted the development of coronary artery disease in these patients. Three of these patients developed coronary artery disease in the first year, 3 in the second and 6 in the third year. The plasma levels of OxLDL and aldehyde-modified LDL were 3.9 ± 0.6

mg/dl, 2.0 ± 0.37 mg/dl and 1.2 ± 0.33 mg/dl, respectively. Although statistical analysis showed no correlation with gender, hypertension and Cytomegalovirus infection, 8 out of 12 of these patients were male, 5 hypertensive and had Cytomegalovirus infection.

4. Discussion

This demonstrates:

- 1) that cardiac explants of patients with ischemic heart disease, but not with dilated cardiomyopathy, contain oxidized LDL in macrophages and in smooth muscle cells in atheromatous plaques;
 - 2) that posttransplant coronary artery disease is associated with increased plasma levels of OxLDL and aldehyde-modified LDL both in patients transplanted for dilated cardiomyopathy or for ischemic heart disease, and
 - 3) that increased plasma levels of OxLDL and aldehyde-modified LDL correlate with the development of coronary artery stenosis.
- OxLDL and aldehyde-modified LDL levels in plasma samples of heart transplant patients without angiographically detectable coronary artery lesions were 2-fold higher than in plasma samples of control subjects without a history of atherosclerotic cardiovascular disease, who were matched for age, gender, and plasma levels of LDL cholesterol, HDL cholesterol and triglycerides. A further 2.7-fold increase was observed in posttransplant plasma samples of patients with pronounced coronary artery stenosis. These data suggest that elevated plasma levels of OxLDL and aldehyde-modified LDL may be an indicator of posttransplant coronary artery stenosis. Increased plasma levels of OxLDL and aldehyde-modified LDL correlated with the extent of coronary artery stenosis and also with its progression, suggesting that OxLDL and aldehyde-modified LDL may play a pathogenic role in the accelerated progression of coronary artery disease in heart transplant patients.

It has been suggested that posttransplant atherosclerosis results from a "response to injury" of the endothelium (10). The extent of ischemic injury in endomyocardial biopsies was indeed found to be a strong
5 predictor of the development of accelerated atherosclerosis (11-13). Endothelial injury may be induced by cellular delayed-type hypersensitivity immune responses elicited by class II histocompatibility (HLA) antigens on coronary artery endothelium (14), by
10 cytomegalovirus infection (15, 16), by cyclosporin (17) and by OxLDL and aldehyde-modified LDL (18) that may act synergistically with cyclosporin (19). In the present study, the extent of histoincompatibility between pairs of donors and recipients, the number of episodes of
15 rejection or Cytomegalovirus infection did not correlate with the grade of coronary artery stenosis, whereas OxLDL and aldehyde-modified LDL were significantly and independently correlated with posttransplant coronary artery disease. The observed association between the age
20 of the donor and the occurrence of coronary artery disease is in agreement with previous findings that coronary atherosclerosis in the donor heart predisposes to accelerated posttransplant coronary artery stenosis (20).

25 OxLDL and aldehyde-modified LDL were demonstrated in coronary arteries in cardiac explants of ischemic heart disease patients suggesting that OxLDL and aldehyde-modified LDL that accumulate in the arterial wall may contribute to the progression of coronary artery
30 stenosis. The cholesterol/protein ratio in OxLDL and aldehyde-modified LDL was very similar to that in LDL extracted from atherosclerotic lesions as described previously (21,22). A possible explanation is that at least part of the OxLDL and aldehyde-modified LDL is
35 released from the arterial wall. Previously, we have demonstrated that plaque rupture in acute myocardial infarction patients is associated with the release of oxidatively modified LDL (5).

In vitro data suggest that OxLDL and aldehyde-modified LDL may be linked to atherogenesis by a sequence of events (reviewed in 2,23). Endothelial cells exposed to OxLDL and aldehyde-modified LDL secrete adhesion molecules, chemoattractant proteins and colony-stimulating factors that enhance the infiltration, proliferation and accumulation of monocytes/macrophages in the arterial wall. Uptake of OxLDL and aldehyde-modified LDL by infiltrated macrophages may result in the generation of foam cells that produce oxygen radicals and thus further contribute to the oxidation of LDL. It has been demonstrated that OxLDL and aldehyde-modified LDL inhibit the migration of aortic endothelial cells in vitro, suggesting that OxLDL and aldehyde-modified LDL may limit the healing response of the endothelium after injury, and that basic fibroblast growth factor reverses the atherosclerosis associated impairment of human coronary angiogenesis-like responses in vitro (24,25). OxLDL and aldehyde-modified LDL may also contribute to rapidly progressing coronary atherosclerosis by inducing platelet adhesion, by decreasing the anticoagulant and fibrinolytic capacities of activated endothelium and by impairing vasodilation and inducing shear stress (2,23).

Increased intracellular levels of ferritin (26) or of alpha-tocopherol analogs (27) decreased the extent of endothelial injury elicited by OxLDL and aldehyde-modified LDL in vitro, whereas antioxidants protect against progression of atherosclerosis in experimental animals (reviewed in document 28).

In summary, the present example demonstrates that posttransplant atherosclerosis correlates with plasma levels of OxLDL and aldehyde-modified LDL.

5. Legend to the Figure 5

Plasma levels of OxLDL and aldehyde-modified LDL and angiographically assessed grade of coronary artery stenosis. Grade 0: normal coronary arteries; grade I: minor abnormalities with < 50% stenosis of primary or

secondary branches and normal left ventricular function; and grade II: $\geq 50\%$ stenosis of primary or secondary branches, or distal occlusions with impaired left ventricular function.

5

EXAMPLE 7Use of the ELISA in renal failure patients

1. Material and methods

1.1. Subjects

10 The patient population consisted of 20 mild chronic renal failure (MCRF) and 77 severe chronic renal failure patients: 21 on conservative treatment including dietary and antihypertensive treatment (SCRF), and 56 on a four-hour, three times a week hemodialysis schedule
15 (HEMO) for 66 months (95% CI, 50-82 months). All hemodialysis patients were given an oral polyvitamin preparation (Ol-Amine, La Meuse, Belgium) after hemodialysis, which contained only minute amounts of
20 of vitamin C). Controls and non-dialyzed patients did not receive routine prescriptions of vitamin supplements. The high frequency of atherosclerotic disease in these patients (Table 6) is in agreement with previously published data (29, 30). The diagnosis of atherosclerotic
25 heart disease, cerebrovascular disease and peripheral vascular disease was made after reviewing the patient files for a history of myocardial infarction, unstable angina or antianginal treatment, cerebrovascular accidents, transient ischemic attack or events related to
30 peripheral vascular disease such as ischemic ulcers, amputation or bypass surgery. Angiograms were available for only a few patients. No patients had evidence of unstable atherosclerotic disease at the time of blood sampling nor in the following days. A group of 27 healthy
35 volunteers (Table 6) without a history of renal disease or atherosclerotic vascular disease served as controls. Patients receiving lipid lowering drugs were excluded.

21766-897

31

The study was approved by the Institutional Review Board and the study subjects provided informed consent.

1.2. Blood samples

5 Venous blood samples from patients and controls were collected on 0.1 vol of 0.1 M citrate, containing 1 mM EDTA, 20 μ M vitamin E, 10 μ M butylated hydroxytoluene, 20 μ M dipyridamole and 15 mM theophylline to prevent in vitro LDL oxidation and in vitro platelet activation, 10 respectively. Blood samples were centrifuged at 3,000 g for 15 min at room temperature within 1 h of collection and stored at -20°C until the assays were performed.

1.3. Assays

15 Titers of autoantibodies against OxLDL and aldehyde-modified LDL and native LDL were measured according to Salonen et al. (3) as described in detail elsewhere (5). vWF antigen levels were measured in a sandwich-type ELISA based on a polyclonal rabbit anti- 20 human vWF antiserum (Dako, Glostrup, Denmark), horseradish peroxidase-conjugated rabbit anti-human vWF IgG (Dako) and o-phenylenediamine. Plasma levels of total cholesterol, HDL cholesterol and triglycerides were determined using standard enzymatic assays (Boehringer 25 Mannheim, Meylon, France). The LDL cholesterol levels were calculated using the Friedewald formula. For the patients not in hemodialysis, creatinine clearance rates were calculated from plasma creatinine levels using the Cockcroft and Gault formula (31).

30

1.4. Statistical analysis

Controls and patients were compared by ANOVA test followed by Dunnett's multiple comparison test, in the Instat* V2.05a statistical program (Graph Pad 35 Software, San Diego, CA). Correlation coefficients were calculated according to Spearman. Multiple regression analysis, using the SAS software (SAS Institute Inc., USA), was performed to study the relationship between *Trade-mark

OxLDL and aldehyde-modified LDL as dependent variable, and age, sex, hypertension (antihypertensive treatment), levels of triglycerides, HDL cholesterol, LDL cholesterol and creatinine clearance rates (marker of extent of renal failure) and levels of vWF (marker of endothelial injury) as independent variables.

2. Results

Mean plasma levels of OxLDL and aldehyde-modified LDL in controls were 0.59 mg/dl (95% CI, 0.52-0.66 mg/dl; n= 27), and were 2.7-fold higher in MCRF patients ($p < 0.01$ as determined by Dunnett's multiple comparison test), 3.1-fold higher in SCRF patients ($p < 0.001$), and 5.4-fold higher in HEMO patients ($p < 0.001$) (Table 7). OxLDL and aldehyde-modified LDL levels were inversely correlated with creatinine clearance rates ($r = -0.65$; $p < 0.001$; $n = 73$). HEMO patients were not included in this analysis because their plasma creatinine clearance cannot be determined adequately.

In a series of 14 hemodialyzed patients, levels of OxLDL and aldehyde-modified LDL were found to be very similar in fresh and in fresh frozen plasma samples. Three freezing and thawing cycles did not cause an increase of OxLDL and aldehyde-modified LDL, indicating that addition of antioxidants and antiplatelet agents prevented in vitro oxidation.

Plasma samples were obtained from 14 hemodialyzed patients on 3 consecutive days before the start of the dialysis procedure. The levels of OxLDL and aldehyde-modified LDL in these samples were similar: 3.4 ± 0.25 mg/dl, 3.2 ± 0.21 mg/dl and 3.5 ± 0.28 mg/dl, respectively. Furthermore, plasma samples were obtained during (after 2 h) and at the end (after 4h) of hemodialysis. Plasma levels of OxLDL and aldehyde-modified LDL were 4.0 ± 0.60 mg/dl and 4.7 ± 0.70 mg/dl ($p = \text{NS}$ vs before) as compared to 3.4 ± 0.25 mg/dl before the start of the dialysis procedure. Thus the

21766-897

33

hemodialysis procedure did not induce a significant increase in the OxLDL and aldehyde-modified LDL levels.

Adequate information about smoking habits was only available for controls (27 non-smokers) and for HEMO patients (12 smokers and 45 non-smokers). Levels of OxLDL and aldehyde-modified LDL were somewhat higher in smoking HEMO patients (3.6 mg/dl; 95% CI, 2.1-5.6 mg/dl) than in non-smoking HEMO patients (3.0 mg/dl; 95% CI, 2.5-3.6 mg/dl; $p = \text{NS}$). The plasma levels of OxLDL and aldehyde-modified LDL in hemodialyzed patients with a history of unstable atherosclerotic cardiovascular disease were 3.5 ± 0.40 mg/dl ($n = 30$) as compared to 2.8 ± 0.60 mg/dl ($n = 26$, $p = \text{NS}$) in hemodialyzed patients without a history of unstable atherosclerotic cardiovascular disease.

LDL fractions were isolated from the plasma of 10 controls, of 10 MCRF patients, of 10 SCRF patients and of 10 HEMO patients by gel filtration on a Superose 6HR 10/30 column, as described previously (5). $75 \pm 6\%$ (mean $\pm$ SD), $80 \pm 4\%$, $83 \pm 6\%$ and $79 \pm 5\%$ of the immunoreactive material was recovered in the LDL fractions. No immunoreactive material migrated in the serum albumin position. The inhibition curves obtained with the respective LDL fractions were parallel to those obtained with in vitro copper-oxidized or MDA-modified standard LDL preparations. OxLDL and aldehyde-modified LDL were isolated from isolated LDL fractions of 10 SCRF patients by ion-exchange chromatography on a mono Q-Sepharose* column with a recovery of 75%. Their physicochemical properties are summarized in Table 8. The levels of arachidonate of OxLDL and aldehyde-modified LDL isolated from these patients were reduced with 75%, whereas its linoleate levels were reduced with 80%. Thirty-seven % of the lysine residues of OxLDL were substituted with aldehydes. The inhibition curves obtained with OxLDL and aldehyde-modified LDL isolated from the plasma of chronic renal failure patients were parallel to these obtained with OxLDL and aldehyde-modified LDL that was obtained by in vitro oxidation of LDL that had been isolated from the

*Trade-mark

plasma of control subjects (Figure 3). The protein/antigen ratio and the relative reactivity in the ELISA of OxLDL and aldehyde-modified LDL isolated from the plasma of these patients were similar to these of copper-oxidized or MDA-modified standard LDL preparations (Table 8).

Titers of autoantibodies against OxLDL and aldehyde-modified LDL were 4.2 (95% CI, 4.0-4.4) in controls, were similar in MCRF and SCRF patients, but significantly increased in HEMO patients ($p < 0.001$) (Table 7). Autoantibody titers correlated with levels of OxLDL and aldehyde-modified LDL in SCRF patients ($r = 0.44$; $p = 0.047$) and in HEMO patients ($r = 0.37$; $p = 0.0055$) (Figure 6). No circulating autoantibodies against native LDL could be detected.

Levels of vWF were 100 percent in controls (95% CI, 90-110 percent), and were 1.5-fold higher in MCRF patients ($p = \text{NS}$ vs controls), 1.6-fold higher in SCRF patients ($p < 0.01$) and 2.1-fold higher ($p < 0.001$) in HEMO patients (Table 7). Levels of vWF were not significantly higher in smoking HEMO patients (250 percent; 95%, 150-340 percent; $n = 12$) than in non-smoking HEMO patients (220 percent; 95% CI, 190-260 percent; $n = 45$). Levels of vWF correlated with levels of OxLDL and aldehyde-modified LDL in MCRF patients ($r = 0.59$; $p < 0.0057$), in SCRF patients ($r = 0.69$; $p = 0.0006$) and in HEMO patients ($r = 0.62$; $p < 0.0001$) (Figure 7). In contrast, levels of vWF did not correlate with LDL cholesterol levels or with body weight.

Multiple regression analysis revealed that the extent of renal failure ($F = 14$; $p = 0.0004$) and the extent of endothelial injury ($F = 26$; $p = 0.0001$), but not age, sex, hypertension, triglyceride levels, HDL cholesterol or LDL cholesterol levels, accounted for a significant fraction of the variations in OxLDL and aldehyde-modified LDL levels (Table 9). Even when only subjects without evidence of ischemic atherosclerotic disease ($n = 53$) were included in the model ($R^2\text{-value} = 0.68$) only the extent of

renal failure ($F= 21$; $p= 0.0001$) and the extent of endothelial injury ($F= 14$; $p= 0.0006$) contributed significantly to the variations in OxLDL and aldehyde-modified LDL levels. No other variables contributed significantly to these variations after exclusion of subjects without evidence of ischemic atherosclerotic disease. When only subjects with evidence of ischemic atherosclerotic disease ($n= 15$) were included only the extent of endothelial injury ($F= 6.2$; $p= 0.047$; R^2 -value= 0.65) contributed to the variations in OxLDL and aldehyde-modified LDL levels. Exclusion of diabetic patients did not significantly change the data either. After exclusion of the extent of renal failure as an independent variable, multiple regression analysis revealed that hemodialysis ($F= 5.6$; $p= 0.021$; $n= 77$), LDL cholesterol levels ($F= 7.1$; $p= 0.0095$) and endothelial injury ($F= 35$; $p= 0.0001$) accounted for a significant fraction of the variation in OxLDL and aldehyde-modified LDL levels in severe chronic renal failure patients (Table 10).

3. Discussion

In vitro work and experimental animal data suggest that oxidized LDL (OxLDL and aldehyde-modified LDL) may contribute to the progression of atherosclerosis (reviewed in document 2), and OxLDL and aldehyde-modified LDL have been demonstrated in human atherosclerotic plaques (5). The immuno-assay of this invention identifies OxLDL and aldehyde-modified LDL (MDA-modified LDL) with ≥ 60 substituted lysines per apo B-100 molecule, which represents the threshold of substitution required for scavenger receptor mediated uptake (1). Increased levels of OxLDL and aldehyde-modified LDL have been measured by ELISA in the plasma of chronic renal failure patients.

Overall, 80 percent of the immunoreactive material isolated from the plasma of patients was recovered in the LDL fractions that were separated by gel

filtration. No immunoreactive material migrated in the albumin position. Inhibition curves obtained with the isolated OxLDL and aldehyde-modified LDL were parallel to these of in vitro copper-oxidized or MDA-modified LDL standard preparations and the protein/antigen ratio and the C_{50} value of the isolated OxLDL and aldehyde-modified LDL were identical to these of standard OxLDL and aldehyde-modified LDL preparations. These data suggested that increased immunoreactivity of OxLDL and aldehyde-
5 modified LDL fractions in plasma of these patients with the antibodies of this invention depended indeed on the higher extent of protein modification and not on changes in lipid composition as was previously observed with other antibodies (32). The increased electrophoretic
10 mobility, the increased lysine modification, the increased cholesterol/protein ratio, the decreased arachidonic acid and linoleate levels were very similar to these of modified LDL extracted from atherosclerotic lesions (21, 22). OxLDL and aldehyde-modified LDL induced
20 foam cell generation, suggesting that OxLDL and aldehyde-modified LDL were not "minimally modified" LDL.

Multiple regression analysis revealed that chronic renal failure and endothelial injury contributed significantly to the variation in OxLDL and aldehyde-
25 modified LDL levels even when patients with evidence of ischemic atherosclerotic disease were excluded. Indeed, 79.6% and 82.4% of the variations in OxLDL and aldehyde-modified LDL levels could be explained in these models. No patients had evidence of unstable atherosclerotic
30 disease at the time of blood sampling nor in the following days and exclusion of patients with a history of ischemic atherosclerotic disease did not affect the contribution of the extent of renal failure and of endothelial injury to the variations in OxLDL and
35 aldehyde-modified LDL.

LDL cholesterol levels in controls and patients were very similar and LDL cholesterol levels did not contribute to the variations in OxLDL and aldehyde-

modified LDL levels. Sutherland et al. (33) demonstrated that the lag time of conjugated diene formation, which is a measure for the sensitivity of LDL to in vitro oxidation, was similar in patients with chronic renal failure and in matched controls. The maximum rate and the extent of LDL oxidation were even lower in patients with renal disease than in controls, due to lower levels of linoleic acid and higher levels of oleic acid.

Furthermore, Schulz et al. (34) demonstrated that despite the fact that hemodialysis causes leukocyte activation, the in vitro LDL oxidation lag time was similar in renal patients and in healthy controls. It was concluded that the antioxidative defense of lipoproteins was preserved in renal failure and during dialysis.

In experimental models, antioxidants such as probucol and vitamin E were found to protect against glomerular injury (35, 36) and to slow atherogenic processes (28). Renal vasoconstriction caused by cholesterol feeding was corrected by probucol or by a thromboxane antagonist (35). Galle et al. (38) demonstrated that the inhibition of endothelium-dependent dilation induced by oxidized lipoprotein could be prevented by high density lipoproteins that are significantly decreased in hemodialyzed patients. In addition, minerals like selenium and nutrients such as coenzyme Q10 may minimize free radical generation and thus oxidative stress. Folic acid, vitamin B12 and vitamin B6 may be essential in the prevention of hyperhomocysteinemia that may contribute to the endothelial injury (39) and to oxidation of LDL (40) in these patients. A diet rich in mono-unsaturated fatty acids (oleic acid, resistant to oxidation) reduced the extent of endothelial injury in diabetes patients (41). Thus it is possible that dietary or pharmacological means may reduce OxLDL and aldehyde-modified LDL and von Willebrand factor in chronic renal failure and alleviate the enhanced generalized atherosclerosis in such patients.

After adjustment for the extent of renal failure, multiple regression analysis revealed that both LDL cholesterol levels and endothelial injury strongly contributed to the variations in OxLDL and aldehyde-modified LDL levels in severe chronic renal failure patients.

Hemodialysis results in platelet and leukocyte activation (42, 43), which generates oxygen radicals and aldehydes that may also contribute to oxidation of LDL. OxLDL and aldehyde-modified LDL may then contribute to thrombogenesis and atherogenesis by stimulating platelets (44). Because of the rather limited number of patients, subgroup analysis to further study the interaction between hemodialysis, oxidation of LDL and ischemic atherosclerotic disease could not be performed (45).

4. Legend to Figures 6 and 7

Figure 6. Correlation between plasma levels of OxLDL and aldehyde-modified LDL (log values) and titers of autoantibodies (log values): regression line for severe chronic renal failure patients, either on conservative treatment ($\blacktriangle$; ---) ($r=0.44$; $p=0.047$) or on hemodialysis ($\blacksquare$; —) ($r=0.37$; $p=0.0055$). No significant correlation was observed in controls and in mild chronic renal failure patients.

Figure 7. Correlation between plasma levels of OxLDL and aldehyde-modified LDL (log values) and of von Willebrand factor antigen (log values): regression line for mild chronic renal failure patients ($\bullet$; -.-.-) ($r=0.59$; $p=0.0057$) or for severe chronic renal failure patients either on conservative treatment ($\blacktriangle$; ---) ($r=0.69$; $p=0.0006$) or on hemodialysis ($\blacksquare$; —) ($r=0.62$; $p<0.00001$). No significant correlation was observed in controls.

EXAMPLE 8Preparation of reference-standard for use in immunological assays

1. Introduction

5 According to the invention it has been found that LDL that is modified by treatment with malondialdehyde (MDA) is highly stable. Furthermore, the extent of modification is highly reproducible. LDL modified with MDA in a particular ratio has an identical
10 number of substituted lysines and can therefore be used as a reference sample in immunological assays. This example shows the preparation of the standard.

2. Material and methods

15 MDA-modified LDL was added to control plasma (containing anti-oxidants and anti-platelet compounds and anti-coagulants) to a final concentration of 100 nM MDA modified apo B-100. Aliquots were frozen at -80°C. In 6 days were aliquots were thawed, diluted to final
20 concentrations ranging from 10 to 0.1 nM MDA-modified apo B-100 and analyzed in ELISA (4 dilution curves per day).

3. Results

25 The inter-assay variation coefficients of 10 subsequent sandwich ELISA's of this invention using 10 subsequent, independent MDA-modified LDL standard preparations of this invention are summarized in Table 11.

30 These data show that for concentrations of MDA-modified LDL ranging from 10 and 0.01 nM the inter-assay variation ranged from 7.6 to 16.9%.

ABBREVIATIONS

C_{50} :	concentration required to obtain 50% inhibition of antibody binding
MDA:	malondialdehyde
HEMO:	severe chronic renal failure patients on maintenance hemodialysis
MCRF:	mild chronic renal failure patients
SCRF:	severe chronic renal failure patients on conservative treatment
OxLDL:	oxidized low density lipoproteins.

Table 1: Clinical data of heart transplant patients

Characteristics	Heart transplant patients		
	Dilated cardiomyopathy		p-values
	(n= 47)	(n= 60)	
Age of recipient (yr)	54± 1.6	55±0.95	*NS
Gender of recipient (M/F)	41/6	53/7	*NS
Age of donor (yr)	29 ± 1.5	29 ± 1.4	**NS
Gender of donor (M/F)	31/16	44/16	*NS
Length of follow up (mo)	39±3.1	50±2.7	**0.008
Duration of ischemia (min)	130±7.0	140±5.3	**NS
No of HLA mismatches			
DR	1.5±0.09	1.4±0.08	**NS
B+DR	3.1±0.13	3.0±0.13	**NS
No of rejection episodes	0.38±0.13	0.25±0.06	**NS
Cytomegalovirus infection	26	43	*NS
Hypertension	37	53	*NS
Diabetes	4	3	*NS
Coronary artery disease			
Grade 0	39	40	*NS
Grade I	5	13	*NS
Grade II	3	7	*NS
Lipid lowering drugs	17	39	*0.004
Statins	13	33	*0.006
Fibrates	4	6	*NS
Calcium channel blockers	31	47	*NS

Data represent mean ± SEM or number of patients. * p-values determined by Chi-square test. ** p-values determined by Dunnnett's multiple comparison test. NS: not significant.

Table 2: Laboratory data of controls and heart transplant patients

Characteristics	Controls		Heart transplant patients			
			Dilated cardiomyopathy (DC)		Ischemic heart disease	
	(n= 27)	(n= 47)	p vs control	(n= 60)	p s control	p-valuesvs DC
Serum triglyderides (mg/dl)†	130±11	130±8.3	NS	140±7.0	NS	NS
HDL cholesterol (mg/dl)‡	44±2.1	54±2.5	NS	49±1.9	NS	NS
LDL cholesterol (mg/dl)‡	120±4.7	100±4.4	NS	110±3.3	NS	NS
Oxidized LDL (mg/dl)	0.59±0.036	1.3±0.14	<0.001	1.7±0.13	<0.001	<0.01

Data represent mean ± SEM. p-values determined by Dunnett's multiple comparison test. NS: not significant. † to convert values for serum triglycerides to millimoles per liter, multiply by 0.011. ‡ to convert values for serum cholesterol to millimoles per liter, multiply by 0.026.

Table 3: Logistic regression analysis of the relation between clinical-laboratory data and extent of coronary artery stenosis in heart transplant patients.

Independent variable	Chi-square value	p-value
Oxidised LDL	18	0.0001
Length of follow up	11	0.0008
Age of donor	3.9	0.047
Age of recipient	0.12	0.73
Sex of recipient	1.8	0.18
Sex of donor	0.025	0.88
History of pretransplant dilated cardiomyopathy (n= 47) or ischemic heart disease (n= 60)	0.0018	0.97
Duration of ischemia	0.25	0.62
No of HLA mismatches	1.6	0.20
No of rejection episodes	3.0	0.081
Cytomegalovirus infection	0.17	0.47
Hypertension	1.9	0.16
Diabetes	0.0016	0.97
Treatment with lipid lowering drugs		
Statins	1.1	0.30
Fibrates	0.12	0.73
Treatment with calcium channel blockers	0.16	0.49
Serum triglycerides	0.18	0.67
Serum HDL cholesterol	0.25	0.61
Serum LDL cholesterol	0.044	0.83

The data set contained 107 patients. Original cardiac disease was dilated cardiomyopathy in 47 and ischemic heart disease in 60 patients. Coronary artery stenosis was assessed angiographically. All quantitative parameters were transformed logarithmically to obtain a normal distribution for linear regression. Chi-square values were obtained after adjustment for all other variables.

Table 4: Clinical data of heart transplant patients without and with progression of coronary artery stenosis during a 3 years follow-up period.

Characteristics	Heart transplant patients		p-value
	Without progression (n= 34)	With progression (n= 12)	
Age of recipient (yr)	58 ± 1.4	60 ± 1.4	**NS
Gender of recipient (M/F)	21/14	11/1	*NS
Age of donor (yr)	25 ± 1.3	32 ± 3.8	**NS
Gender of donor (M/F)	27/7	10/2	*NS
Duration of ischemia (min)	130 ± 6.7	140 ± 11	**NS
No of HLA mismatches			
DR	1.2 ± 0.13	1.5 ± 0.15	**NS
B+DR	2.8 ± 0.21	3.2 ± 0.24	**NS
Cytomegalovirus infection	24	11	*NS
Hypertension	21	10	*NS
Diabetes	1	2	*NS

Data represent mean ± SEM or number of patients. *p-values determined by Chi-square analysis. **p-values determined by Dunnett's multiple comparison test. NS= not significant

Table 5: Laboratory data of heart transplant patients without and with progression of coronary artery stenosis during a 3 years follow-up period.

Characteristics	Heart transplantation patients		p-value
	Without progression (n=34)	With progression (n=12)	
Serum triglycerides (mg/dl)	130 ± 8.6	150 ± 14	NS
HDL cholesterol (mg/dl)	50 ± 2.7	49 ± 4.9	NS
LDL cholesterol (mg/dl)	110 ± 3.6	105 ± 8.7	NS
Oxidized LDL (mg/dl)	1.2 ± 0.069	2.6 ± 0.33	0.0005

Data represent means ± SEM. p-values determined by Dunnett's multiple regression comparison test. NS= not significant.

Table 6

Characteristics	Controls (n=27)	Mild chronic renal failure (n=20)	Severe chronic renal failure	
			non-dialysed (n=21)	hemodialysed (n=56)
Males/Females	12/15	11/9	7/14	33/23
Age (years)	54 (50-58)*	52 (44-60)*	55 (49-62)*	61 (58-65)*
Body weight (kg)	72 (69-76)*	73 (67-79)*	59 (53-65)*	65 (61-68)*
Creatinine clearance (ml/min)	110 (110-120)*	34 (29-39)*	8.4 (7-10)*	nd
Primary renal disease:				
Glomerulonephritis	-	4	3	11
Autosomal dominant polycystic kidney disease	-	2	6	10
Diabetes	-	1	4	6
Reflux-nephropathy	-	1	2	2
Chronic Interstitial Nephritis	-	2	2	9
Hypertensive nephropathy	-	2	0	2
Other ¹	-	6	1	9
Unknown	-	2	3	7
Hypertension	1	16	19	18
Atherosclerotic heart disease	-	6	7	24
Cerebrovascular accidents	-	0	3	9
Peripheral vascular disease	-	2	1	13

* Data represent means and 95% confidence intervals (between brackets).

¹ including: hereditary nephropathy, sarcoidosis, renal tuberculosis, thrombotic thrombocytopenic purpura, myeloma, traumatic loss, congenital urinary tract abnormalities.

nd: creatinine clearance rate cannot be determined adequately.

Table 7

Laboratory data of study subjects

	Controls		MCRF patients		SCRF patients		HEMO patients	
	(n=27)		(n=20)	p vs controls	(n=21)	p vs controls	(n=56)	p vs controls
Triglycerides (mg/dl)	120 (100-150)		150 (130-170)	NS	120 (100-140)	NS	130 (110-160)	NS
HDL cholesterol (mg/dl)	44 (39-48)		38 (33-42)	NS	44 (38-50)	NS	37 (35-40)	< 0.05
LDL cholesterol (mg/dl)	120 (110-130)		110 (100-130)	NS	110 (105-130)	NS	120 (110-130)	NS
Oxidized LDL (mg/dl)	0.59 (0.52-0.66)		1.6 (1.0-2.2)	< 0.01	1.8 (1.3-2.3)	< 0.001	3.2 (2.7-3.7)	< 0.001
Autoantibodies (titer)	4.2 (4.0-4.4)		4.7 (4.0-5.4)	NS	5.0 (4.2-5.8)	NS	6.6 (5.7-7.4)	< 0.001
vWF (percent)	100 (90-110)		150 (110-180)	NS	160 (130-190)	< 0.01	210 (180-240)	< 0.001

Data represent means and 95% confidence intervals (between brackets).

Table 8

Characteristics of native LDL and of OxLDL isolated from plasma of severe chronic renal failure patients

	Native LDL	OxLDL
Protein/antigen ratio	> 100	1.1
Reactivity with mAb-4E6 (C ₅₀ mg/dl)	25	0.02
Relative electrophoretic mobility	1	1.7
Malondialdehyde (mole/mole protein)	3	68
Substituted lysines per apo B-100	5	130
Cholesterol/protein ratio	1.8	3.3
Free cholesterol/cholesterol ester ratio	0.38	0.36
Phospholipid/protein ratio	1.7	1.6
Fatty acids (%)		
16:0	14	37
18:1	19	50
18:2	55	10
20:4	12	3

Data represent means of ten LDL preparations of chronic renal failure patients. Native LDL and OxLDL patients were separated by ion-exchange chromatography.

Table 9

Multiple regression analysis of the dependence of OxLDL on the extent of renal failure

Variable	F-value	p-value
Age	1.2	0.28
Sex	1.4	0.25
Hypertension	1.1	0.31
Triglycerides	1.5	0.23
HDL cholesterol	1.7	0.20
LDL cholesterol	0.99	0.32
Renal failure	14	0.0004

The data set contained 27 controls, 20 MCRF patients and 21 SCRF patients. F-values were obtained after adjustment for the other variables. Cockcroft creatinine clearance rates were used as a quantitative parameter for the extent of renal failure. All linear variables were logarithmically transformed to obtain normality for linear regression analysis. The multiple R^2 value of the multiple regression model was 0.634.

Table 10

Multiple regression analysis of the dependence of OxLDL on hemodialysis and LDL cholesterol levels in severe chronic renal failure patients

Variable	F-value	p-value
Age	0.31	0.58
Sex	0.19	0.66
Hypertension	0.01	0.95
HDL cholesterol	0.02	0.89
Triglycerides	3.7	0.060
Hemodialysis	5.6	0.021
LDL cholesterol	7.1	0.0095

The data set contained 21 SCRF and 56 HEMO patients. All linear variables were transformed logarithmically to obtain normality for linear regression analysis. The multiple R^2 value of the multiple regression model was 0.56.

Table 11. Inter-assay variation coefficients of sandwich ELISA using 10 subsequent, independent MDA-modified LDL preparations

Concentration (nM)	Inter-assay variations coefficients (%)
10	9.6
5	7.6
2.5	8.4
1.25	13.2
0.62	12.0
0.31	13.0
0.16	12.3
0.08	15.5
0.04	16.9
0.02	13.6
0.01	11.4

DOCUMENTS

1. Haberland MD, Fogelman AM, Edwards PA: Specificity of receptor-mediated recognition of malondialdehyde-modified low density lipoproteins. Proc Natl Acad Sci 1982;79:1712-1716.
2. Holvoet P, Collen D: Oxidized lipoproteins in atherosclerosis and thrombosis. FASEB J 1994;8:1279-8440.
3. Salonen JT, Ylä-Herttuala S, Yamamoto R, Butler S, Korpela H, Salonen R, Nyyssonen K, Palinski W, Witztum JL: Autoantibody against oxidized LDL and progression of carotid atherosclerosis. Lancet 339: 883-887, 1992.
4. Holvoet P, Perez G, Bernar H, Brouwers E, Vanloo B, Rosseneu M, Collen D: Stimulation with a monoclonal antibody (mAb4E4) of scavenger receptor-mediated uptake of chemically modified low density lipoproteins by THP-1 derived macrophages enhances foam cell generation. J Clin Invest 93: 89-98, 1994.
5. Holvoet P, Perez G, Zhao Z, Brouwers E, Bernar H, Collen D: Malondialdehyde-modified low density lipoproteins in patients with atherosclerotic disease. J Clin Invest 1995;95:2611-2619.
6. Havel RJ, Eder HA, Bragdon JH: The distribution and chemical composition of ultracentrifugally separated lipoproteins in human sera. J Clin Invest 1955;34:1345-1353.
7. Steinbrecher UP: Oxidation of low density lipoprotein results in derivatization of lysine residues of apolipoprotein B by lipid peroxide decomposition products. J Biol Chem 1987;262:3603-3608.
8. Sparrow CP, Parthasarathy S, Leake DS, Witztum JL, Steinberg D: Enzymatic modification of low density lipoprotein by purified lipoxxygenase plus phospholipase-A₂ mimic cell-mediated oxidative modification. J Lipid Res 1988; 29: 745-753.
9. Holvoet P, Donck J, Landeloos M, Brouwers E, Luijckens K, Arnout J, Lesaffre E, Vanrenterghem Y, Collen

D: Correlation between oxidized low density lipoproteins and von Willebrand factor in chronic renal failure.

Thromb Haemostas 1996;76:663-669.

10. Libby P, Salomon RN, Payne DD, Schoen FJ, Pober JS: Functions of vascular wall cells related to development of transplantation-associated coronary arteriosclerosis. Transplant Proc 1989;21:3677-3684.

11. Hruban RH, Beschorner WE, Baumgartner WA, Augustine SM, Ren H, Reitz BA, Hutchins GM: Accelerated arteriosclerosis in heart transplant recipients is associated with a T-lymphocyte-mediated endothelialitis. Am J Pathol 1990;137:871-882.

12. Rose EA, Smith CR, Petrossian GA, Barr ML, Reemtsma K: Humoral immune responses after cardiac transplantation: correlation with fatal rejection and graft atherosclerosis. Surgery 1989;106:203-208.

13. Tanaka H, Sukhova GK, Swanson SJ, Cybulsky MI, Schoen FJ, Libby P: Endothelial and smooth muscle cells express leukocyte adhesion molecules heterogeneously during acute rejection of rabbit cardiac allografts. Am J Pathol 1994;144:938-951.

14. Crisp SJ, Dunn MJ, Rose ML, Barbir M, Yacoub MH: Antiendothelial antibodies after heart transplantation: the accelerating factor in transplant-associated coronary artery disease? J Heart Lung Transplant 1994;13:81-92.

15. Grattan MT, Moreno-Cabral CE, Starnes VA, Oyer PE, Stinson EB, Shumway NE: Cytomegalovirus infection is associated with cardiac allograft rejection and atherosclerosis. JAMA 1989;261:3561-3566.

16. Koskinen P, Lemstrom K, Bruggeman C, Lautenschlager I, Hayry P: Acute cytomegalovirus infection induces a subendothelial inflammation (endothelialitis) in the allograft vascular wall. A possible linkage with enhanced allograft arteriosclerosis. Am J Pathol 1994;144:41-50.

17. Cartier R, Dagenais F, Hollmann C, Cambron H, Buluran J: Chronic exposure to cyclosporin affects

endothelial and smooth muscle reactivity in the rat aorta. Ann Thorac Surg 1994;58:789-794.

18. Chin JH, Azhar S, Hoffman BB: Inactivation of endothelial derived relaxing factor by oxidized lipoproteins. J.Clin.Invest. 1992;89:10-18.

19. Galle J, Schollmeyer P, Wanner C: Cyclosporin and oxidized low density lipoproteins synergistically potentiate vasoconstriction: influence of the endothelium. Eur Heart J 1993;14(Suppl):111-117.

20. Tuzcu EM, Hobbs RE, Rincon G, Bott-Silverman C, De Franco AC, Robinson K, McCarthy PM, Stewart RW, Guyer S, Nissen SE: Occult and frequent transmission of atherosclerotic coronary disease with cardiac transplantation. Insights from intravascular ultrasound. Circulation 1995;91:1706-1713.

21. Hoff HF, O'Neill: Lesion-derived low density lipoprotein and oxidized low density lipoprotein share a lability for aggregation, leading to enhanced macrophage degradation. Arterioscler Thromb 1991; 11:1209-1222.

22. Steinbrecher UP, Loughheed M: Scavenger receptor-independent stimulation of cholesterol esterification in macrophages by low density lipoprotein extracted from human aortic intima. Arterioscler Thromb 1992; 12:608-625

33. Steinberg D, Witztum JL: Lipoproteins and atherogenesis: current concepts. J Am Med Assoc 1990;264:3047-3052.

23. Ross R: The pathogenesis of atherosclerosis: a perspective for the 1990s. Nature 1993;362:801-809.

24. Murugesan G, Chisolm GM, Fox PL: Oxidized low density lipoprotein inhibits the migration of aortic endothelial cells in vitro. J Cell Biol 1993; 120: 1011-1019.

25. Chen CH, Nguyen HH, Weilbaecher D, Luo S, Gotto AM Jr, Henry PD: Basic growth factor reverses atherosclerotic impairment of human coronary angiogenesis-like responses in vitro. Atherosclerosis 1995; 116: 261-268.

26. Juckett MB, Balla J, Balla G, Jessurun J, Jacob HS, Vercellotti GM: Ferritin protects endothelial cells from oxidized low density lipoprotein in vitro. Am J Pathol 1995;147:782-789.

27. Mabile L, Fitoussi G, Periquet B, Schmitt A, Salvayre R, Negre-Salvayre A: Alpha-Tocopherol and trolox block the early intracellular events (TBARS and calcium rises) elicited by oxidized low density lipoproteins in cultured endothelial cells. Free Radic Biol Med 1995;19:177-187.

28. Steinberg D: Clinical trials of antioxidants in atherosclerosis: are we doing the right thing ? Lancet 1995;346:36-38.

29. Degoulet P, Legrain M, Reach I, Aime F, Devries C, Rojas P, Jacobs C: Mortality risk factors in patients treated by chronic hemodialysis. Nephron 31: 103-110, 1982.

30. Neff MS, Eiser AR, Slifkin RF, Baum M, Baez A, Gupta S, Amarga E: Patients surviving 10 years of hemodialysis. Am J Med 74: 996-1004, 1983.

31. Cockcroft DW, Gault MH: Prediction of creatinine clearance from serum creatinine. Nephron 16: 31-41, 1976.

32. Reade V, Tailleux A, Reade R, Harduin P, Cachera C, Tacquet A, Fruchart JC, Fievet C: Expression of apolipoprotein B epitopes in low density lipoproteins of hemodialyzed patients. Kidney Int 44: 1360-1365, 1993.

33. Sutherland WH, Walker RJ, Ball MJ, Stapley SA, Robertson MC: Oxidation of low density lipoproteins from patients with renal failure or renal transplants. Kidney Int 48: 227-236, 1995.

34. Schulz T, Schiffel H, Scheithe R, Hrboticky N, Lorenz R: Preserved antioxidative defense of lipoproteins in renal failure and during hemodialysis. Am J Kidney Dis 25: 564-571, 1995.

35. Keane WF, Mulcahy WS, Kasiske BL, Kim Y, O'Donnell MP: Hyperlipidemia and progressive renal disease. Kidney Int Suppl 39: S41-S48, 1991.

36. Trachtman H, Schwob N, Maesaka J, Valderrama E: Dietary supplementation ameliorates renal injury in chronic puromycin aminonucleoside nephropathy. *J Am Soc Nephrol* 5: 1811-1819, 1995.
37. Kaplan R, Aynedjian HS, Schlondorff D, Bank N: Renal vasoconstriction caused by short-term cholesterol feeding is corrected by thromboxane antagonist or probucol. *J Clin Invest* 86: 1707-1714, 1990.
38. Galle J, Bengen J, Schollmeyer P, Wanner C: Oxidized lipoprotein(a) inhibits endothelium-dependent dilation: prevention by high density lipoprotein. *Eur J Pharmacol* 265: 111-115, 1994.
39. Friedman JA, Dwyer JT: Hyperhomocysteinemia as a risk factor for cardiovascular disease in patients undergoing hemodialysis. *Nutr Rev* 53: 197-201, 1995.
40. McCully KS: Chemical pathology of homocysteine. I. Atherogenesis. *Ann Clin Lab Sci* 23: 477-493, 1993.
41. Rasmussen O, Thomsen C, Ingerslev J, Hermansen K: Decrease of von Willebrand factor levels after a high-monounsaturated fat diet in non-insulin-dependent diabetic subjects. *Metabolism* 43: 1406-1409, 1994.
42. Reverter JC, Escolar G, Sanz C, Cases A, Villamor N, Nieuwenhuis HK, Lopez J, Ordinas A: Platelet activation during hemodialysis measured through exposure of P-selectin: analysis by flow cytometric and ultrastructural techniques. *J Lab Clin Med* 124: 79-85, 1994.
43. Zwaginga JJ, Koomans HA, Sixma JJ, Rabelink TJ: Thrombus formation and platelet-vessel wall interaction in the nephrotic syndrome under flow conditions. *J Clin Invest* 93: 204-211, 1994.
44. Zhao B, Diehl JH, Harrachruprecht B, Winterhorst H: Oxidized LDL induces serotonin release from blood platelets. *Am J Hematol* 48: 285-287, 1995.
45. Pocock SJ: Subgroup analysis, in Pocock SJ (ed): *Clinical trial. A practical approach*. Wiley J & Sons, Chichester, 1993, p 211-218.

21766-897

57

CLAIMS:

1. An immunological assay for the detection and/or quantification of human MDA-modified LDL and human OxLDL in a sample derived from the body fluids or tissues of a human being, said assay comprising:
 - (a) contacting the sample with a first antibody that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL and human OxLDL; and
 - (b) thereafter visualizing and/or quantifying a binding reaction between the first antibody and the MDA-modified LDL and OxLDL present in the sample;wherein the MDA-modified LDL and OxLDL for which the first antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contain at least 60 substituted lysine moieties per apo B-100 moiety.
2. The assay of claim 1 in which the MDA-modified LDL and OxLDL for which the first antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contain at least about 90 substituted lysine moieties per apo B-100 moiety.
3. The assay of claim 1 in which the MDA-modified LDL and OxLDL for which the first antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contain at least about 120 substituted lysine moieties per apo B-100 moiety.
4. The assay of claim 1 in which the MDA-modified LDL and OxLDL for which the first antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contain at least about 210 substituted lysine moieties per apo B-100 moiety.
5. The assay of claim 1 in which the MDA-modified LDL and OxLDL for which the first antibody has an affinity of at

21766-897

58

least about $5 \times 10^8 \text{ M}^{-1}$ contain at least about 240 substituted lysine moieties per apo B-100 moiety.

6. The assay of any one of claims 1 to 5 which is a competitive assay.

5 7. The competitive assay of claim 6 in which MDA-modified LDL and/or OxLDL are bound to a substrate, comprising contacting the sample and the first antibody with the substrate having bound to it MDA-modified LDL and OxLDL.

8. The assay of any one of claims 1 to 5 which is a
10 sandwich assay in which the first antibody is bound to a substrate, comprising contacting the sample with the substrate having bound to it the first antibody.

9. The assay of claim 8 in which a second antibody is used and the second antibody has an affinity of at least
15 about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL and human OxLDL.

10. The assay of claim 9 in which the second antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human native LDL.

11. The assay of any one of claims 1 to 10 in which
20 the affinity constant of the first antibody for MDA-modified LDL and for OxLDL is at least about $1 \times 10^{10} \text{ M}^{-1}$.

12. The assay of any one of claims 1 to 11 in which the first antibody has an affinity of less than about $1 \times 10^7 \text{ M}^{-1}$ for human native LDL.

25 13. The assay of claim 12 in which the affinity constant of the first antibody for human native LDL is less than about $1 \times 10^6 \text{ M}^{-1}$.

14. The assay of any one of claims 1 to 13 in which the first antibody is the monoclonal antibody mAb-4E6

21766-897

59

produced by hybridoma Hyb4E6 deposited at the BCCM under deposit accession number LMBP 1660 CB on or about April 24, 1997.

15. The assay of any one of claims 9 to 10 in which
5 the affinity constant of the second antibody for MDA-modified LDL and for OxLDL is at least about $1 \times 10^{10} \text{ M}^{-1}$.

16. The assay of claim 15 in which the affinity constant of the second antibody for human native LDL is at least about $1 \times 10^9 \text{ M}^{-1}$.

10 17. The assay of claim 16 in which the second antibody is the monoclonal antibody mAb-8A2 produced by hybridoma Hyb8A2 deposited at the BCCM under deposit accession number LMBP 1661 CB on or about April 24, 1997.

18. The assay of any one of claims 1 to 17 which is an
15 immunohistochemical assay in which the sample is a tissue sample and it is contacted with the first antibody.

19. The assay of any one of claims 1 to 17 in which the sample is derived from the body fluids of a human being.

20. The assay of any one of claims 1 to 19 in which
20 the first antibody is capable of detecting 0.02 mg/dl of human MDA-modified LDL and human OxLDL in undiluted human plasma.

21. An immunological sandwich assay for the detection and/or quantification of human MDA-modified LDL in a sample
25 derived from the body fluids or tissues of a human being in which assay a first antibody that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL is bound to a substrate, said assay comprising:

21766-897

60

(a) contacting the sample with the substrate having bound to it the first antibody under binding conditions so that human MDA-modified LDL in the sample will bind to the first antibody;

5 (b) thereafter removing unbound sample from the substrate;

(c) thereafter contacting the substrate with a second antibody that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human MDA-modified LDL; and

10 (d) thereafter visualizing and/or quantifying the MDA-modified LDL that was present in the sample;

wherein the MDA-modified LDL for which the first antibody and the second antibody have an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contains at least 60 substituted lysine moieties per apo B-100 moiety.

22. The assay of claim 21 in which the MDA-modified LDL for which the first antibody and the second antibody have an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contains at least about 90 substituted lysine moieties per apo B-100 moiety.

23. The assay of claim 21 in which the MDA-modified LDL for which the first antibody and the second antibody have an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contains at least about 120 substituted lysine moieties per apo B-100 moiety.

24. The assay of claim 21 in which the MDA-modified LDL for which the first antibody and the second antibody have an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contains at least about 210 substituted lysine moieties per apo B-100 moiety.

21766-897

61

25. The assay of claim 21 in which the MDA-modified LDL for which the first antibody and the second antibody have an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ contains at least about 240 substituted lysine moieties per apo
5 B-100 moiety.

26. The assay of any one of claims 21 to 25 in which the first antibody also has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human OxLDL.

27. The assay of any one of claims 21 to 26 in which
10 the first antibody has an affinity of less than about $1 \times 10^7 \text{ M}^{-1}$ for human native LDL.

28. The assay of any one of claims 21 to 25 and 27 in which the first antibody has an affinity of less than about $1 \times 10^7 \text{ M}^{-1}$ for human OxLDL.

15 29. The assay of any one of claims 21 to 28 in which the second antibody has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human native LDL.

30. The assay of any one of claims 21 to 29 in which the affinity of the first antibody for MDA-modified LDL is
20 at least about $1 \times 10^{10} \text{ M}^{-1}$.

31. The assay of any one of claims 21 to 30 in which the affinity of the first antibody for human native LDL is less than about $1 \times 10^6 \text{ M}^{-1}$.

32. The assay of any one of claims 21 to 31 in which
25 the affinity of the second antibody for human native LDL is at least about $1 \times 10^9 \text{ M}^{-1}$.

33. The assay of any one of claims 21 to 25 in which the first antibody is the monoclonal antibody mAb-4E6 produced by hybridoma Hyb4E6 deposited at the BCCM under

21766-897

62

deposit accession number LMBP 1660 CB on or about April 24, 1997.

34. The assay of any one of claims 21 to 25 in which the first antibody is the monoclonal antibody mAb-1H11
5 produced by hybridoma Hyb1H11 deposited at the BCCM under deposit accession number LMBP 1659 CB on or about April 24, 1997.

35. The assay of any one of claims 21 to 25 in which the second antibody is the monoclonal antibody mAb-8A2
10 produced by hybridoma Hyb8A2 deposited at the BCCM under deposit accession number LMBP 1661 CB on or about April 24, 1997.

36. The assay of any one of claims 21 to 35 in which at least one antibody is capable of detecting 0.02 mg/dl of
15 human MDA-modified LDL in undiluted human plasma.

37. Monoclonal antibody mAb-4E6 produced by hybridoma Hyb4E6 deposited at the BCCM under deposit accession number LMBP 1660 CB on or about April 24, 1997.

38. Hybridoma Hyb4E6 deposited at the BCCM under
20 deposit accession number LMBP 1660 CB on or about April 24, 1997.

39. Monoclonal antibody mAb-8A2 produced by hybridoma Hyb8A2 deposited at the BCCM under deposit accession number LMBP 1661 CB on or about April 24, 1997.

25 40. Hybridoma Hyb8A2 deposited at the BCCM under deposit accession number LMBP 1661 CB on or about April 24, 1997.

41. A method of standardizing an assay for human MDA-modified LDL and human OxLDL by using as a calibrator or

21766-897

63

as a control a stable standard containing MDA-modified LDL whose extent of substitution of its lysine moieties will remain essentially constant over normal periods of time during normal storage for biological materials, the

5 MDA-modified LDL of said standard being made by contacting malondialdehyde with LDL at a predetermined molar ratio of malondialdehyde to the apo B-100 moiety of the LDL, the standard containing an agent that reduces the ability of any metal ions present to catalyze oxidation of the LDL and/or

10 an anti-oxidant.

42. The method of claim 41 wherein the standard contains both an agent that reduces the ability of any metal ions present to catalyze oxidation of the LDL and an anti-oxidant.

15 43. The method of any one of claims 41 to 42 wherein the agent that reduces the ability of any metal ions present to catalyze oxidation of the LDL is a chelating agent.

44. The method of claim 43 wherein the chelating agent is EDTA.

20 45. The method of any one of claims 41 to 44 wherein the anti-oxidant is selected from the group consisting of BHT (butylated hydroxytoluene) and Vitamin E.

46. The method of any one of claims 41 to 45 wherein the standard further comprises a physiological fluid.

25 47. The method of claim 46 wherein the physiological fluid is plasma.

48. The method of any one of claims 41 to 47 wherein the standard further comprises at least one anti-platelet coagulation compound and/or anti-coagulant.

21766-897

64

49. The method of any one of claims 41 to 48 wherein the standard is used as a calibrator.

50. The method of any one of claims 41 to 48 wherein the standard is used as a control.

5 51. A kit for conducting a sandwich assay for the determination of human OxLDL or human MDA-modified LDL or both in a sample derived from the body fluids or tissues of a human being, said kit comprising:

(a) a substrate on which is bound a first antibody
10 that has an affinity of at least about $5 \times 10^8 \text{ M}^{-1}$ for human OxLDL or human MDA-modified LDL or both, the OxLDL and MDA-modified LDL each having at least 60 substituted lysine moieties per apo B-100 moiety, and

(b) a labeled antibody having an affinity of at
15 least about $5 \times 10^8 \text{ M}^{-1}$ for human OxLDL that becomes bound to the first antibody during the assay or for human MDA-modified LDL that becomes bound to the first antibody during the assay or for both that become bound to the first antibody during the assay.

20 52. The kit of claim 51 further comprising a reactive substance for reaction with the labeled antibody to give an indication of the presence of the labeled antibody.

53. The kit of claim 52 wherein the reactive substance comprises an enzyme.

25 54. The kit of any one of claims 51 to 53 further comprising the calibrator of claim 49.

55. The kit of any one of claims 51 to 54 further comprising the control of claim 50.

21766-897

65

56. The kit of any one of claims 51 to 55 for conducting a sandwich assay for the determination of human OxLDL and human MDA-modified LDL in a sample derived from the body fluids or tissues of a human being.

5 57. The kit as claimed in any one of claims 51 to 55 for conducting a sandwich assay for the determination of human MDA-modified LDL in a sample derived from the body fluids or tissues of a human being.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS




FIG.1

2/6




FIG. 2A

SUBSTITUTE SHEET (RULE 26)

3/6







FIG.2B

SUBSTITUTE SHEET (RULE 26)


4/6



5/6



6/6

FIG. 6FIG. 7