



(11)

EP 3 134 499 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

29.01.2025 Bulletin 2025/05

(21) Application number: 15782790.8

(22) Date of filing: 23.04.2015

(51) International Patent Classification (IPC):

C11D 1/00 (2006.01)

C11D 11/00 (2006.01)

C11D 1/14 (2006.01)

C11D 1/66 (2006.01)

C11D 1/75 (2006.01)

(52) Cooperative Patent Classification (CPC):

C11D 11/00; C11D 1/146; C11D 1/662; C11D 1/75

(86) International application number:

PCT/US2015/027403

(87) International publication number:

WO 2015/164677 (29.10.2015 Gazette 2015/43)

(54) **CLEANING FORMULATIONS FOR CHEMICALLY SENSITIVE INDIVIDUALS: COMPOSITIONS AND METHODS**

REINIGUNGSFORMULIERUNGEN FÜR CHEMISCH EMPFINDLICHE PERSONEN:
ZUSAMMENSETZUNGEN UND VERFAHREN

FORMULATIONS DE NETTOYAGE POUR DES INDIVIDUS SENSIBLES AUX PRODUITS CHIMIQUES : COMPOSITIONS ET PROCÉDÉS

(84) Designated Contracting States:

AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR

(30) Priority: 23.04.2014 US 201461982887 P

(43) Date of publication of application:

01.03.2017 Bulletin 2017/09

(73) Proprietors:

- van Buskirk, Gregory
Danville, CA 94506 (US)
- Steinemann, Anne
Townsville, QLD 4810 (AU)
- Nair, Vinod S.
San Jose, CA 95120 (US)

(72) Inventors:

- van Buskirk, Gregory
Danville, CA 94506 (US)
- Steinemann, Anne
Townsville, QLD 4810 (AU)
- Nair, Vinod S.
San Jose, CA 95120 (US)

(74) Representative: **Bryers Intellectual Property Ltd**

Bristol & Bath Science Park
Dirac Crescent, Emerson's Green
Bristol, BS16 7FR (GB)

(56) References cited:

WO-A1-2009/131816 WO-A1-2010/141184
US-A1- 2002 034 553 US-A1- 2007 207 940
US-A1- 2011 098 376 US-A1- 2012 015 856
US-B2- 8 246 978

- DATABASE WPI Week 200908, Derwent World Patents Index; AN 2009-B28685, XP002777330
- R. A. GRAVELING ET AL: "A review of multiple chemical sensitivity", OCCUPATIONAL AND ENVIRONMENTAL MEDICINE, vol. 56, no. 2, 1 February 1999 (1999-02-01), pages 73 - 85, XP055441290, ISSN: 1351-0711, DOI: 10.1136/oem.56.2.73
- CARESS, SM ET AL.: "A Review of a Two-Phase Population Study of Multiple Chemical Sensitivities.", ENVIRON HEALTH PERSPECT., vol. 111, no. 12, September 2003 (2003-09-01), pages 1490 - 1497, XP055233679, Retrieved from the Internet <URL:<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1241652/pd/fehp0111-001490.pdf>> [retrieved on 20150623]

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES., BROOKHAVEN LANDFILL., 29 November 2005 (2005-11-29), pages 1 - 32, XP055400481, Retrieved from the Internet <URL:https://apps.health.ny.gov/statistics/environmental/public_health_tracking/tracker/documents/brookhavenlandfill_general_hc112905.pdf> [retrieved on 20150623]
- SUGAYA, N ET AL.: "Development of a headspace GC/MS analysis for carbonyl compounds (aldehydes and ketones) in household products after derivatization with o-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine.", ANAL SCI., vol. 20, 5 May 2004 (2004-05-05), pages 865 - 870, XP055233682, Retrieved from the Internet <URL:https://www.jstage.jst.go.jp/article/analsci/20/5/20_5_865/_pdf> [retrieved on 20150623]
- METCALFE, DD : "Introduction: What Are the Issues in Addressing the Allergenic Potential of Genetically Modified Foods.", ENVIRON HEALTH PERSPECT., vol. 111, 8 June 2003 (2003-06-08), pages 1110 - 1113, XP055233683, Retrieved from the Internet <URL:<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1241559/pdf/eihp0111-001110.pdf>> [retrieved on 20150623]
- PORTER, AW ET AL.: "Identification of opdA, a Gene Involved in Biodegradation of the Endocrine Disrupter Octylphenol.", APPL ENVIRON MICROBIOL., vol. 73, no. 22, November 2007 (2007-11-01), pages 7373 - 7379, XP055233685, Retrieved from the Internet <URL:<http://aem.asm.org/content/73/22/7373.full.pdf+html>> [retrieved on 20150623]

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Description

CROSS-REFERENCE TO RELATED APPLICATION

5 **[0001]** This application claims the benefit of U.S. Prov'l. Appl. Ser. No. 61/982,877 filed 23 April 2014, which U.S. Appl. No. 15/306,109, with filing date 23 April 2015 and patent number U.S. Pat. No. 10,767,137, also claims the benefit of.

BACKGROUND OF THE DISCLOSURE

10 FIELD OF THE DISCLOSURE

[0002] The present disclosure relates to cleaning compositions in general, and cleaning compositions well suited for those individuals, who experience multiple chemical sensitivities (MCS), in particular. Individuals with MCS are virtually unable to use commercially available cleaners. The instant disclosure concerns the selection of ingredients and methods for formulating and evaluating a series of cleaning products for use by any person, including individuals with MCS.

15 **[0003]** Cleaning product compositions that are suitable for cleaning clothing, dishware, countertops and other hard surfaces have been commercially prepared, marketed, and sold to consumers for over two hundred years. As cleaning technology progressed, environmental and safety issues sometimes lagged behind discoveries in cleaning efficacy. For example, in the late 1950's and early 1960's, it was found that synthetic surfactants that had supplanted natural soap products exhibited poor biodegradability, and were building up in waste water streams; streams laden with tenacious foam were widespread, and tremendous efforts were focused on finding alternatives. In the 1970's, certain builder compounds also came under scrutiny for their environmental impact, such as eutrophication on inland lakes and ponds. In the search for alternate builder materials, one candidate material, nitrilotriacetic acid, NTA, was found to be a very promising candidate. Fortunately, before it reached mass distribution, safety tests showed that it could transport heavy metals across 20 placental membranes, which was thought of as potentially harmful to developing fetuses.

25 **[0004]** These examples, among others, has led to attention being paid to safety on par with that of cleaning performance; in fact, attention to the issue has resulted in numerous very effective ingredients being removed from commerce. By and large, most cleaning ingredients currently in use have resolved many of the issues of the past, and now there is widespread 30 effort to understand the potential after-effects of cleaning ingredients as far as acute toxicity, chronic toxicity, carcinogenicity, mutagenicity, teratogenicity, and hormone disruption are now commonly looked at before ingredients are brought to market.

35 **[0005]** One further phenomenon that has received attention in the last few years is the effect of cleaning compositions on chemically sensitive individuals. In westernized countries, asthma and related atopic disorders such as eczema and hay fever are now major public health concerns, due to their high prevalence-approximately 20% of the people in the United States are estimated to be sufferers. Understandably, there is concern associated with significant ill health and high societal and healthcare costs. Multiple scientific studies have raised concerns about the potential for consumer products to cause or exacerbate asthma or asthma-like responses.

40 **[0006]** While the removal of dyes and fragrances from cleaning products have alleviated responses of some sensitive individuals, there are a considerable number of consumers who are not able to use commercially-available products for 45 reasons that until now have not been well-understood. These individuals are forced to use centuries-old cleaners such as vinegar and baking soda, products that are lacking in cleaning efficacy, but are a last resort for these individuals. Unfortunately, while the mechanism whereby these individuals become highly and multiply sensitized is not understood, when they do become sensitized, there is no known cure of reversal of debilitating responses. Products are therefore needed that are not only designed for these individuals, but for a general population that is possibly but unknowingly vulnerable to acquiring multiple chemical sensitivities.

50 **[0007]** In recent years, more and more products are being sold which claim to be "green", "environmentally friendly", "natural", "organic", "sustainable", etc., with the implication that such products contain ingredients that are bio-based, or at least have lower levels of petrochemical ingredients. While some of these products have been based on well-founded technology, the actions of some have caused environmental advocates as well as the media to warn against the 55 phenomenon of trying to promote a product's credentials through dubious claims as "greenwashing." Although some regulatory agencies, such as the EPA and FDA, provided regulations and standards for environmentally hazardous substances and food and drugs respectively, there is no similar agency that specifically covers cleaning products. In addition, none of these agencies have developed clear guidelines for the terms "natural", "green", "environmentally friendly" or the like. There are some organizations, which provide lists of approved natural components and standards for components based on standardized test methods which measure toxicity, biodegradability and other factors for determining the naturalness and environmental impact of a given product. However there is little guidance on issues like, use of "eco-hybrids" or "hybrid surfactants" that are comprised of both petroleum and plant based chemistries, which is contributes to the ongoing problem of "greenwashing".

[0008] There is perhaps a larger problem with the implication that no matter how "green" or "natural" a product might be, that such products may imply that they are safer for consumers than other mainstream products. While standards have been established to measure the degree of bio-basis of a product, the need for standards to better promote the safety of such products has received too little attention, much less been established. No organizations can certify the overall safety of consumer cleaning products, in particular towards consumers that suffer from multiple chemical sensitivities.

[0009] In addition to proximate effects of potentially deleterious ingredients, increasing attention has turned toward understanding conveyance of such chemicals from the household to the larger environment. Indeed it has been reported that the exhaust coming out of a dryer vent has detectable amounts of volatile organic compounds (VOCs) in all tested commercially available detergent products. See A.C. Steinemann, L.G. Gallagher, A.L. Davis, and I.C. MacGregor, "Chemical Emissions from Residential Dryer Vents During Use of Fragranced Laundry Products," *Air Quality, Atmosphere and Health*, 6 (2011) 151-156. VOCs from consumer products can migrate outdoors and thus impact outdoor air quality. According to California Air Resources Board 1990 statistics, some 265 tons of VOCs were released into California air from the use of consumer products each day. See B. Bridges, "Fragrance: Emerging Health and Environmental Concerns," *Flavour and Fragrance Journal*, 17 (2002) 361-371. This makes it difficult for a customer trying to make an environmentally conscious decision to purchase cleaning products that will not release harmful VOCs into the atmosphere.

[0010] In summary, cleaning products available in the market today do not explicitly address all aspects of consumer safety. While the vast majority of cleaning product manufacturers ensure that their products cause minimal acute and chronic toxicity problems, exposure to cleaning product chemicals has been associated with the development and exacerbation of asthma and related disorders. However, consumers who may desire to lessen their exposure to harmful chemicals by purchasing safe cleaning products are unable to do so because product ingredients are not fully disclosed on labels. Further, the ingredient profiles of cleaning products that are claimed to be green are remarkably similar to those not labeled green, causing confusion in the minds of consumers looking for safe cleaning products. Indeed, experts on indoor air quality have shown the presence of known carcinogens and hazardous air pollutants even in cleaning products that are free of fragrances and dyes.

25

DESCRIPTION OF THE RELATED ART

[0011] U.S. Pat. Nos. 6,973,362 and 7,096,084 to Long, et al., teach a method for evaluating chemical components based on their function in the product. The methods taught by Long require first the function of a given raw material in a product first be identified, and then a set of predetermined criteria be applied based on the function of the raw material, to determine the raw material's designated environmental class rating, which is then given an environmental grade of from 1-3. The problem with this method is that it requires an individual, burdensome analysis of each component of a composition to arrive at a final value for the composition as a whole. In addition, it requires that the individual components be analyzed by their function and one or more components in a composition may have multiple functions. Furthermore, this method requires knowledge of all the components, their percentages in the formulation and their functions in a given formulation, which makes testing products off the shelf impossible or impractical because the required information is often not readily available. The end result is that although this method provides a standardized method for measuring the environmental impact of a given chemical formulation, it too is burdensome and requires too much information about the components and their functions to make it practical for use in testing a wide range of compositions that are available on store shelves.

[0012] International Publications Nos. WO2007099294, WO2009024743, and WO2009024747 assigned to Reckitt Benckiser Group, plc, teach compositions for toilet cleaning and hard surface cleaning which are "environmentally acceptable," but the application does not clearly define what is meant by "environmentally acceptable". The publications merely teach cleaning compositions, which do not have high levels of volatile organic compounds or VOCs, and exclude certain acids, solvents, chelating agents and thickeners. While these applications teach certain "environmentally acceptable" compositions, it does not establish any criteria or test methods which could be used to determine if other compositions meet this criteria other than those compositions which may have the same exact ingredients as those taught in the application.

[0013] Similarly, U.S. Pat. Nos. 5,990,065 and 6,069,122 assigned to Procter & Gamble teach compositions for dishwashing detergents that contain natural surfactants and solvents, but they do not teach a method or criteria of determining whether a composition is "natural" or a means of measuring the natural components in a given composition. These patents merely teach a means of making a particular dishwashing composition that contains some natural ingredients.

[0014] US 2012/015856 A1 to Harthun, et al, concerns a "Solution Carbon Index" (SCI) which is analogous to Applicants' pMC criteria. However, unlike pMC and Applicants' formulations, Harthun's SCI is a *weighted average* of individual component Carbon Indexes. There is no recognition that *all* of their components must have a pMC of 80% or greater. Indeed, in the single example formulation (see Table I), Harthun includes Deriphaf[®] 160C, a sodium N-lauryl- β -iminodipropionate, which is described as having a carbon Index of 66, and Dehyphon[®] LS 36, an alcohol ethoxylate

propoxylate, which is described as having a Carbon Index of only 35. Accordingly, there is no guarantee that individual ingredients have a pMC above 80%, as required by Applicants.

5 [0015] Caress, SM, et al. discusses the prevalence, symptomatology, and etiology of Multiple Chemical Sensitivities. While they discuss broad categories of chemicals that have been shown to trigger a sensitivity response (e.g., fragrances, cleaning agents, solvents), they do not propose specific agents that may be responsible for such a response, nor do they propose any screening method, whereby one may identify chemicals that would be acceptable for use by those suffering from MCS. Indeed, one of the co-authors of Caress is a joint inventor of the instant application. At the time of publication of Caress in 2003, one would have been led to believe that the only course of action would be a broadscale avoidance of cleaning products.

10 [0016] US 2007/207940 to G. Fenyvesi, et al. discloses compositions containing a singular solvent, 1,3-propanediol. With the possible exception of additional bio-based glycols that may optionally be included, the inventors are silent as to the nature of additional surfactants, solvents, or builders. In fact, their examples are predominantly petrochemically-based, such that their pMC values would be much less than the threshold of Applicants' 80% criteria. For example: Carbopol™ 934, sodium xylene sulfonate, EO/PO block copolymer, sodium polyacrylate, linear sodium dodecylbenzene sulfonate, Tergitol® 15-S-7, triethanolamine, C12-13 linear alcohol EO-7, sodium laureth sulfate and Monoethanolamine-all with pMC's of zero-as well as cocamide DEA and cocamidopropyl betaine with pMCs of 76% and 64%, respectively. Furthermore, the inventors do not discuss the effects of any of these chemicals as far as persons with MCS.

15 [0017] Among their ingredients, CZ 20 0707-412 to Smidrkal, et al., cite the use of sodium fatty acids, which are soaps, and sodium carbonate, which is present in every sample. The inventors are silent as to the sourcing of alkyl polyglucosides, 20 and whether or not contaminants from the production thereof are present or absent. In addition, they are silent as to the use of solvents in their inventive compositions altogether and the only organic builders given in their examples are petrochemically-based-the sodium salt of a copolymer of acrylic acid and maleic acid. Further, the inventors are silent as to the necessity of limiting vapor pressures of the components.

25 [0018] WO 2009/131816 to Luciow, et al. describes methods for formulating a personal cleansing composition, containing a primary surfactant, a secondary amphoteric and/or nonionic surfactant, a humectant and a polymeric and/or quaternary conditioner are selected; each component of which has a natural and renewable resource origin. However, the inventors do not establish a minimum biological content, and numerus examples of their ingredients do not meet Applicants minimum criterion of 80% bio-basis for *each* ingredient, e.g., cocamidopropyl betaine (64% bio-based), sodium laureth sulfate (0% bio-based), tetrasodium EDTA (0% bio-based) and sodium benzoate (0% bio-based).

30 [0019] WO 2010/141184 A1 (Clorox-Coulter, et al.) discusses a method for measuring the percentage of natural ingredients of a cleaning composition using radiocarbon dating data, and give examples of compositions with high pMCs. However, inventors-among whom is co-inventor van Buskirk of the instant application-did not recognize that beyond the screening criterion for pMC, there are other criteria to meet in order to make such chemicals and formulated compositions suitable for those suffering from MCS., e.g., sourcing of chemicals such as alkyl polyglucoside, vapor pressures of 35 components, presence or absence of toxic chemicals in the product headspace, and the like.

40 [0020] R.A. Graveling, et al., "A Review of Multiple Chemical Sensitivity" surveys literature available up to their time of publication (1999) pertaining to MCS, without assigning specific causes. The authors acknowledge the possibility of physical causality, perhaps sensitization of the limbic (midbrain) system. The authors also found that the levels of sensitizers that elicit responses from those with MCS are well below those ordinarily regarded as toxic. The authors propose no specific agents that may be responsible for such a response, nor propose any screening method whereby one may identify chemicals that would be acceptable for use by those suffering from MCS.

SUMMARY OF THE DISCLOSURE

45 [0021] The present disclosure concerns a new scientific protocol for the formulation of cleaning products to minimize the triggering of asthma or other immunological responses in humans. In addition to improving the outlook for symptom-free cleaning, products generated according to the criteria described herein, while virtually non-petroleum based, are equivalent in performance to existing cleaning products on the market.

50 DEFINITIONS

[0022] In the present specification and claims, reference will be made to phrases and terms of art which are expressly defined for use herein as follows:

55 *Active ingredient* or *active material* refers to entities that contribute to the cleaning of stains and soils and/or disinfecting of fabrics or surfaces. A chemical mixture as procured from suppliers may be diluted with a solvent such as water, which serves no purpose in cleaning and/or disinfection; in such case, the active ingredient refers only to the portion of the chemical mixture that serves a purpose to clean and/or disinfect. This term does not generally include aesthetic ingredients such as fragrance materials, colorants, viscosity modifiers, preservatives, or the like.

[0023] *Biologically based carbon* or *bio-based carbon* is carbon derived from plant or animal sources that have lived up until the relatively recent past. It is distinguished from carbon derived from fossil sources such as coal, subterranean natural gas, oil or petroleum-based carbon. Bio-based carbon is characterized by the presence of radioactive ^{14}C , unlike fossil sources of carbon in which radioactive ^{14}C is depleted or entirely absent.

[0024] *Chemical allergy* describes the adverse health effects that may result when exposure to a chemical elicits an immune response in an individual. Chemical allergens produce reactions similar to allergens such as pollens, weeds, and dander, but appear to be generated when lower-molecular weight chemicals bind to carrier macromolecules. See M.H. Karol, O.T. Macina, and A. Cunningham, "Cell and molecular biology of chemical allergy," *Ann Allergy Asthma Immunol.* 87 (2001) 28-32.

[0025] *Cleaning composition* or *cleaning formulation* as used herein refers to a mixture of ingredients assembled together for the purpose of providing an aid to the removal of dirt, soil, grime, food waste, etc., from a surface or individual. A cleaning composition may be formulated for use in cleaning laundry, hard surfaces such as dishes, kitchen surfaces, bathrooms, glass, mirrors, etc., and may be comprised of both of active ingredients and aesthetic ingredients. A cleaning composition is distinguished from a product that is primarily a single cleaning active, such as a bar of soap. A cleaning composition is typically the product presented for sale to consumers.

[0026] *Greenwashing* as used herein refers to the practice of making or making a false, misleading, or inflated green marketing claims. This practice was expanded upon in December 2007 by the environmental marketing firm TerraChoice. See "The Six Sins of Greenwashing™," A 'Green Paper' by TerraChoice Environmental Marketing Inc. (November 2007); <http://www.sins of greenwashing.com/index6b90.pdf>.

[0027] *Headspace* or *headspace technology* as used herein concerns measurement and characterization of components present in the space above a particular composition or ingredient. Headspace analysis involves removing volatile compounds from the headspace surrounding an object or other material of interest using either an inert gas or by establishing a vacuum. The compounds are then trapped and analyzed with techniques such as gas chromatography, mass spectrometry or Carbon-13 NMR. (See, for example, en.wikipedia.org/wiki/Headspace_technology).

[0028] *Modern carbon* refers to carbon derived from modern life forms, either plant or animal. It is distinguished from carbon derived from fossil sources such as coal, subterranean natural gas, oil or petroleum-based carbon. It is characterized by presence of radioactive ^{14}C in its make-up, which is depleted in feedstocks sourced from fossil carbon.

[0029] *Product* refers to a cleaning composition or cleaning formulation offered for commercial sale. The term can be understood to be synonymous with cleaning composition or cleaning formulation.

[0030] *Renewable carbon source* or *renewably sourced carbon* is synonymous with modern carbon, and refers to carbon sourced from non-primitive or non-ancient sources, i.e., it is not derived from fossil sources, which is coal, subterranean natural gas, oil or petroleum-based carbon. Renewable carbon source or renewably sourced carbon derives from modern life forms, either plant or animal, and is labeled as renewable because it is relatively easily replenished relative to fossil carbon, which takes millennia if not eons to form. It is characterized by presence of radioactive ^{14}C in its make-up, which is depleted in feedstocks sourced from fossil carbon.

[0031] *Soap* as used herein refers to saponified animal fats and vegetable oils. Soap is understood to be distinguishable from synthetic surfactants, builders, pH adjusters, solvents, soil release agents, antimicrobials, enzymes and bleaching agents.

40 DETAILED DESCRIPTION OF THE DISCLOSURE

[0032] The instant disclosure concerns a multi-tiered approach to screening ingredients for suitability for use in cleaning products, formulating cleaning products that contain acceptable ingredients, and evaluating the resulting cleaning products thus formulated. As all cleaning products are combinations of raw materials, which individually may constitute mixtures, the chance of including undesirable chemicals in cleaning products is therefore high without an appropriate screening process.

MODERN CARBON-BASED INGREDIENTS

[0033] Radiocarbon dating and analysis is a commonly used process to date carbon-based artifacts and remains within the field of archeology. More recently, radiocarbon dating has been used for testing a variety of different products including but not limited to: personal care products, wipes, lubricants, plastics, cleaning products, gardening products, etc. The subject is discussed extensively in "Determining the Modern Carbon Content of Biobased Products Using Radiocarbon Analysis", by G.A. Norton and S.L. Devlin, from Iowa State University, published by *Bioresource Technology* 97 (2006) 2084-2090.

[0034] The article on determining modern carbon content describes the process of radiocarbon dating for the determination of bio-based content in a formulation. Several carbon isotopes are present in nature, ^{12}C , ^{13}C and ^{14}C . The ^{12}C is a stable isotope and the ^{14}C is an unstable isotope and undergoes radioactive decay. The ^{14}C is produced in the

atmosphere where it is oxidized to CO_2 and CO_2 is then absorbed by plants until the $^{12}\text{C}/^{14}\text{C}$ ratio in all living matter is essentially the same as that in the atmosphere. When something dies, it stops absorbing carbon and the amount of ^{14}C diminishes with time, as it naturally undergoes radioactive decay. The rate of decay for the ^{14}C is measurable and can be calculated. The decay rate for ^{14}C is slow, about 5730 years, relative to the movement of carbon through the food chain,

5 from plants to animals to bacteria. All carbon in biomass at earth's surface contains atmospheric levels of ^{14}C whereas petrochemical feedstock that has been dead and in the ground for millions of years will have little to no ^{14}C . Therefore, material derived from a recently living plant will have an abundance of ^{14}C that is approximately equal to that in the atmosphere, whereas petrochemical feedstocks will not have a ^{14}C signature.

[0035] By knowing the feedstocks of individual components of a molecule, one can estimate its amount of bio-based or modern carbon. For example, if all the component carbons of an ingredient are from plant- or animal-basis, it is deemed 100% bio-based or modern carbon; if only half of the component carbons are from bio-based or modern sources, while the other half of the component carbons are from non-modern sources such as coal, subterranean natural gas, oil or petroleum-based carbon, then the ingredient is 50% bio-based or modern carbon. This number, designated as Percent Modern Carbon (pMC), has been described by others as Biorenewable Carbon Index (BCI) or Renewable Carbon Index 15 (RCI), and is used synonymously herein. As long as one is knowledgeable about the source of all the carbons in the molecule of interest, that is, whether they are derived from modern carbon sources or non-modern carbon sources, one can estimate the Percent Modern Carbon (pMC) using Equation (1):

$$20 \text{ Estimated pMC} = \frac{\text{number of carbons from modern carbon sources}}{\text{total number of carbons from all sources}} \times 100\% \quad (1)$$

[0036] Alternatively, one can analyze for bio-based or modern carbon content, alternately termed Percent Modern Carbon (pMC), can be carried out by standard test methodology such as radiocarbon analysis, according to ASTM method 25 D6866-05, which relies on analyzing the sample for radioactive ^{14}C . Using ^{14}C analysis and calculations, one can determine or confirm the amount of carbon in a material from fossil carbon, which is coal, oil or petroleum-based carbon. By measuring the amount of radioactive carbon in a sample, the amount of modern carbon or bio-based carbon can be determined. As one can understand, the Percent Modern Carbon (pMC), Biorenewable Carbon Index (BCI) or Renewable Carbon Index (RCI) is a measure of the percent of modern or biobased carbon in an individual ingredient or in a 30 composition.

[0037] The Percent Modern Carbon (pMC), Biorenewable Carbon Index (BCI) or Renewable Carbon Index (RCI) only refers to the element, carbon, in the molecule or compound. Therefore, it is an index of the ratio of new, modern, bio-based carbon to "old", typically petrochemical-based carbon. pMC (as well as its synonymous terms BCI and RCI) does not refer to any other elements such as H, N, O, S, etc. that may be present in a compound. One complication in the calculation of 35 pMC is that inorganic carbon, such as that from the carbonates, would be included as "old" carbon, although it might originate from a "natural" mineral source. However, laboratories do have ways to deal with this complication experimentally and can account for mineral-based carbon. Materials with 100% modern carbon or bio-based carbon have no fossil carbon or petroleum based carbon and are considered carbon from renewable resources.

[0038] The radioactive carbon dating analysis that serves as the bases for pMC/BMI/RCI may be performed using 40 American Society of Testing Materials (ASTM) method D6866-05. ASTM D6866-05 describes various techniques for measuring radioactive carbon using 1) accelerator mass spectrometry (AMS), 2) benzene synthesis, or 3) carbon dioxide absorption, also known as the carbon dioxide cocktail method. For benzene synthesis or carbon dioxide absorption methods, a liquid scintillation counter (LSC) is used to detect byproducts of the ^{14}C decay process. When preparing a sample for radiocarbon analysis, the sample composition maybe dehydrated, to prepare the sample for testing. 45 Depending on the method used for radiocarbon analysis, the degree of uncertainty may vary slightly. Using ASTM method D6866-05, the degree of uncertainty is approximately 1 to 2%. Using an LSC, the degree of uncertainty reaches approximately $\pm 3\%$. When using the AMS method or the benzene synthesis method to measure ^{14}C , radioactive carbon count must be corrected for isotopic fractionation to obtain a corrected radiocarbon count. The carbon dioxide cocktail method does not require a correction for isotopic fractionation. The radioactive carbon dating process and analysis may be done for whole compositions or for individual components of compositions, and any combinations or variations thereof.

[0039] A method for determining the suitability of ingredients for use in the novel cleaning compositions described herein 50 involves performing an assessment of the bio-basis of the ingredient, either through the estimation means described above or by analytical data such as that described in ASTM D6866-05. It is preferable that ingredients used herein are predominantly, if not entirely, renewably sourced, i.e., biologically-based or bio-based, as well as readily and completely biodegradable. It has been found in the course of the present work that individuals with Multiple Chemical Sensitivities or MCS may tolerate ingredients with higher content of modern carbon better than ingredients high in content of non-modern carbon, such as petrochemicals. According to the instant disclosure, therefore, cleaning ingredients-and preferably all 55 formula ingredients-are selected to contain at least 80%, and more preferably at least 85% bio-based or modern carbon,

more preferably at least 90% bio-based or modern carbon, and most preferably 100% bio-based or modern carbon. It is preferred that the entire formulation be at least 90% bio-based or modern carbon, more preferably greater than about 95% bio-based or modern carbon, and most preferably greater than about 99% bio-based or modern carbon.

5 **SELECTION OF INGREDIENTS**

[0040] It has been determined in the course of the present work described herein that even ingredients that claim or analyze to be 100% bio-based can contain undesirable contaminants, such as low levels of residual petrochemical solvents, catalysts, or unsafe by-products. Hence, it is important to also analyze for materials that contain known hazardous volatile organic compounds (VOCs) and carcinogens, and/or that may contain potential "telltale" indicators for petrochemicals, such as phenyl derivatives. This is typically accomplished by conducting a headspace analysis of the ingredient under consideration for use in a particular cleaning formulation. Methods have been developed for this purpose, most specifically EPA Compendium Method TO-15, "Determination Of Volatile Organic Compounds (VOCs) In Air Collected In Specially-Prepared Canisters And Analyzed By Gas Chromatography/Mass Spectrometry (GC/MS)," EPA, 1999, and U.S. EPA Method TO-11A, "Determination of Formaldehyde in Ambient Air Using Adsorbent Cartridge Followed by High Performance Liquid Chromatography (HPLC)," EPA, 1999.

[0041] The analytes from such a headspace determination can be compared against authoritative lists of hazardous ingredients, such as can be found in the *Clean Air Act-Toxic and Flammable Substances for Accidental Release Prevention* list, the *Clean Air Act-Hazardous Air Pollutant* list, the *Comprehensive Environmental Response, Compensation, and Liability Act Hazardous Substance* list, the *Clean Water Act-Priority Pollutant* list, the *Emergency Planning & Community Right to Know Act Toxic Release Inventory Chemical* list, the *Federal Insecticide, Fungicide, and Rodenticide Act Registered Pesticide* list, the *Occupational Safety and Health Act-Air Contaminants* list, and/or the *Resource Conservation and Recovery Act Hazardous Constituents* list. A method for determining the suitability of an ingredient for use in novel cleaning compositions as described herein involves determining the level of VOCs that may be contributed to a final formulation by performing a headspace analysis on the ingredient. Another method for determining the suitability of a cleaning composition for use with individuals that exhibit MCS involves determining the level of VOCs in the headspace of the as-formulated cleaning composition. It may be understood that physiological responses may differ for each contaminant, and most preferably none of the analytes found in the headspace is to be found on the authoritative lists. Analytes that might be found on the authoritative lists should be present at levels below about 1000 µg/m³, more preferably below 10 µg/m³.

[0042] Finally, through working with individuals that exhibit MCS in the course of the instant work, it has been determined that it is advantageous that cleaning compositions essentially contain no active components that have a vapor pressure exceeding 0.1 mm Hg at 20°C.. Without being bound by theory, it is believed that moieties such as surfactant residues that have a chain length of greater than 8 carbon atoms are insufficiently reactive with proteins to form complexes. Alternatively, it is postulated that any conjugate having greater than an 8-carbon atom chain length that may form, are present in concentrations that are lower than a threshold level needed to trigger an immunogenic response. As such, it is preferred to formulate cleaning products that contain ingredients, especially surfactants with hydrophobic carbon chains that are essentially devoid of carbon chains of 8 or less. In other words, surfactants and other moieties having greater than 8-carbon chain lengths are especially preferred.

[0043] It is further advantageous for a cleaning composition to also be devoid of impurities that have a vapor pressure exceeding 0.1 mm Hg at 20°C. Impurity or impurities as used herein therefore refer to an ingredient that is not knowingly or intentionally desired to be incorporated into a cleaning composition of the instant disclosure by a formulator or other individual, as will be readily understood by one skilled in the relevant art. Yet another method for determining the suitability of ingredients for use in novel cleaning compositions as described herein involves selecting ingredients that contain no active component that has a vapor pressure exceeding 0.1 mm Hg at 20°C. Still another method for determining the suitability of ingredients for use in formulating the novel cleaning compositions described herein involves selecting ingredients that contain no impurities that have a vapor pressure greater than 0.1 mm Hg at 20°C. Marketing studies have confirmed that consumers associate fragrance substantivity with increased cleanliness. However, multiple scientific studies have implicated fragrances as being the culprit in exacerbating or causing deleterious health effects in susceptible individuals or entire segments of the population. It is therefore preferable to incorporate fragrances that are known to not cause deleterious effects. Without being bound by theory, the incorporation of optically active isomers of fragrance molecules in their naturally occurring form is favored as possibly having less adverse effects than their synthetic analogs.

55 **EVALUATION OF INGREDIENTS AND FORMULATIONS**

[0044] Once candidate ingredients are identified and tested as described above, they are evaluated for use in potential cleaning formulations using a blind study protocol. The blind studies used in the course of the instant work were comprised of a specially selected panel of volunteers. Volunteers diagnosed with both multiple chemical sensitivities (MCS) and

asthma have been found to be able to detect the presence of problematic chemicals, even at low levels. A panel comprised of just such individuals was used for many aspects of the studies conducted herein. While animals use olfactory-mediated defense systems to detect, locate and identify predators in their surrounding environment, it has been found that human subjects are similarly able to discriminate among negative odors accurately. See E.A. Krusemark and W. Li, "Enhanced

5 olfactory sensory perception of threat in anxiety: An event-related fMRI study," *Chemosensory Perception*, 5 (2012) 37-45.

[0045] In fact, people with MCS have demonstrated an ability to detect harmful chemicals at levels far lower than the rest of the population. In the course of the instant work, at least one individual with MCS was used to rank prospective ingredients for acceptability in cleaning formulations based upon sensory responses, which included olfactory as well as skin contact. Instrumental analyses were then implemented to correlate results with sensory ratings from the human panel, and to identify and/or quantify the chemicals detected and deemed to be potentially harmful to humans. Ingredients that were deemed acceptable by the human panel and the instrumental analyses were then used as raw materials for cleaning products described herein. It is believed that this level of pre-screening and testing represents a first in the world for consumer cleaning product formulations work, and has provided an unprecedented level of safety testing for consumer products. Accordingly, with the technology newly presented and described herein, a method for providing cleaning formulations for use by the general public and chemically-sensitized individuals, in particular, involves:

1. assessing the bio-basis of an ingredient or ingredients for use in a cleaning product;
2. evaluating the ingredient or ingredients from step 1. for acceptability by at least one individual who manifests multiple chemical sensitivities; and
3. formulating a cleaning product using the acceptable ingredient or ingredients from step 2; wherein each ingredient contains at least 80% pMC.

25 [0046] As a double-check on the safety of cleaning product formulation ingredients, they can be evaluated for the presence or absence of potentially harmful volatile organic carbon (VOC) compounds. In a recent publication it was found that 37 products emitted 156 different VOCs, with an average of 15 VOCs per product. Of these 156 VOCs, 42 VOCs are classified as toxic or hazardous under U.S. federal laws, and each product emitted at least one of these chemicals. See A. Steinemann, "Volatile Emissions from Common Consumer Products," *Air Quality, Atmosphere & Health*, March 2015.

30 Emissions of carcinogenic hazardous air pollutants (HAPs) from green fragranced products were not significantly different from regular fragranced products. The most common chemicals in fragranced products were terpenes which, interestingly, were not found to be present in fragrance-free formulations. Of the volatile ingredients found in the headspace of these products, fewer than 3% were disclosed on any product label or material safety data sheet (MSDS).

[0047] After the acceptance of ingredients is established via the methods identified above, cleaning products using these approved chemicals may then be formulated and evaluated for efficacy. As it is recognized that combinations of effects can cause antagonistic responses, evaluations of fully formulated products were then carried out via sensory evaluation and VOC analysis. This permits further evaluation of the suitability of product formulations and the ability to assess product performance as compared with existing cleaning products. This was done on a qualitative rating scale both for cleaning efficacy and for presumed safety.

40 [0048] Accordingly, another method for providing cleaning formulations for use by chemically-sensitized individuals in addition to the general public, involves:

1. assessing the bio-basis of an ingredient or ingredients for use in a cleaning product;
2. evaluating the ingredient or ingredients from step 1. for acceptability by at least one individual who manifests multiple chemical sensitivities;
3. formulating a cleaning product using the acceptable ingredient or ingredients from step 2; and
4. performing a VOC headspace analysis of the cleaning product formulated in step 3, wherein an analyte value of less than about 1000 $\mu\text{g}/\text{m}^3$ is regarded as acceptable.

[0049] Yet another method for providing cleaning products for use by chemically-sensitized individuals as well as for the general public, involves, in addition to steps 1. through 4. above, at least one of the steps of:

5. confirming the cleaning efficacy of a cleaning product formulated according to steps 1. through 4. above; and
6. evaluating the cleaning product formulated according to steps 1. through 4. above for acceptability for use by at least

one individual who manifests multiple chemical sensitivities.

[0050] A different method for providing cleaning products that are particularly well suited for use by chemically-sensitized individuals, includes:

- 5 1. assessing the bio-basis of an ingredient or ingredients for use in a cleaning product;
- 10 2. evaluating the ingredient or ingredients from step 1. for acceptability by at least one individual who manifests multiple chemical sensitivities;
- 15 3. formulating a cleaning product using the acceptable ingredient or ingredients from step 2; and
4. performing a headspace analysis of the cleaning product formulated in step 3; wherein assessing step 1. includes an analysis of the feedstock of the ingredient or ingredients according to ASTM method D6866-05 or one consistent therewith.

[0051] Still another method for providing cleaning products according to the instant disclosure includes any of assessing steps 1. above, further wherein the assessing is achieved by analysis according to or consistent with ASTM D6866-05.

[0052] Yet another method for providing cleaning products that are particularly well suited for use by chemically-sensitized individuals as well as the general public, includes:

- 20 1. assessing the bio-basis of an ingredient or ingredients for use in a cleaning product;
- 25 2. evaluating the ingredient or ingredients from step 1. for acceptability by at least one individual who manifests multiple chemical sensitivities;
3. formulating a cleaning product using the acceptable ingredient or ingredients from step 2; and
- 30 4. performing a headspace analysis of the cleaning product formulated in step 3; wherein assessing step 1. includes an analysis of the feedstock of the ingredient or ingredients by analysis that may be according to ASTM method D6866-05, a method that is consistent therewith, by consulting appropriate tabulated material, or by any combination of the foregoing.

[0053] Still another method for providing cleaning products that are particularly well suited for use by chemically-sensitized individuals as well as the general public, includes, in addition to any of steps 1. through 5. above, at least one of the following criteria:

- 35 a. the ingredient or ingredients of step 1. have a pMC of at least 80%, more preferably at least 85%, and most preferably at least 90%;
- 40 b. individual ingredients are evaluated for acceptability and deemed suitable by at least one individual who manifests multiple chemical sensitivities or MCS;
- 45 c. the cleaning formulation has a pMC of at least 90%, more preferably at least 95%, and most preferably at least 99%;
- d. the headspace analysis reveals less than 1000 $\mu\text{g}/\text{m}^3$ VOCs regulated by governmental bodies;
- 50 e. the cleaning formulation is deemed acceptable by at least one individual who manifests multiple chemical sensitivities or MCS.

[0054] A cleaning product according to the disclosure herein that is particularly well suited for use by chemically-sensitized individuals as well as the general public, includes: a composition comprising at least one ingredient that is a non-soap cleaning active, wherein the ingredient has a pMC of at least 80%, wherein a headspace analysis of the composition reveals the absence of phenyl compounds or their derivatives, wherein less than about 5% by weight of the ingredients have a vapor pressure that is above 0.1 mm Hg at 20°C, wherein the composition contains less than about 1% by weight of a fragrance material; wherein headspace analysis of the cleaning product reveals less than about 1000 $\mu\text{g}/\text{m}^3$ VOCs that are regulated by governmental bodies; and wherein the composition has less than 0.1% by weight of ingredients that have been demonstrated to cause adverse reactions in chemically-sensitive individuals.

CLEANING FORMULATION COMPONENTS/INGREDIENTS

[0055] Cleaning formulations are generally comprised of a mixture of ingredients, each of which serves a purpose in the removal of soils and stains. Generally, such formulations can include one or more of the following active ingredients: surfactants, builders, pH adjusters, solvents, soil release agents, antimicrobials, enzymes and bleaching agents. Such formulations often include ingredients that are more aesthetic in their function: fragrance materials, dyes and colorants, viscosity control agents, pearlizing and opacifying agents, brighteners, preservatives, etc. A discussion of the types and best practice for incorporation of these materials follows.

10 ANIONIC SURFACTANTS

[0056] Cleaning compositions according to the instant disclosure can contain an anionic surfactant. When an anionic surfactant is added to the compositions described herein, it can typically be added at a level from about 0.05% to about 15% by weight, preferably from about 0.05% to about 5% by weight, and more preferably from about 0.1% to about 1% by weight of the composition. It is preferred that anionic surfactants have alkyl chain lengths greater than 10. It is further preferred that they be sourced from bio-based materials rather than petrochemicals. While this largely eliminates phenyl derivatives, it is envisioned that these materials could also be sourced from bio-based materials. It is yet further preferred that these materials be devoid of contaminants such as 1,4-dioxane. While this largely eliminates ethoxylated derivatives, it is envisioned that these materials can be sourced with a bio-based source of ethylene oxide, and that the 1,4-dioxane contaminant can be scrupulously removed or avoided during production.

[0057] Anionic surfactants suitable for use in the formulations discussed herein include C₁₀-C₁₄ alkyl sulfates and ethoxysulfates (e.g., Stepanol WA-EXTRA from Stepan Company), C₁₀-C₁₈ alkyl sulfonates, C₁₀-C₁₄ linear or branched alkyl benzene sulfonates, and C₁₀-C₁₅ alkyl ethoxycarboxylate. Anionic surfactants may be paired with organic counterions or multivalent counterions in order to prevent interference with cationic species. Further examples of suitable surfactants are described in McCutcheon's Vol. 1: Emulsifiers and Detergents, North American Ed., McCutcheon Division, MC Publishing Co., 1995.

[0058] In the course of the instant work, it was found that a number of anionic surfactants containing ethylene oxide, either through petrochemical or bio-based sources, contained detectable levels of 1,4-dioxane as a contaminant. Such surfactants are to be scrupulously avoided, with preference given to anionic surfactants that have no detectable level of 1,4-dioxane.

[0059] Highly preferred materials anionic surfactants are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to air, moisture or sunlight.

35 NONIONIC SURFACTANTS

[0060] The compositions can contain a nonionic surfactant. When a nonionic surfactant is added to the composition, it can typically be added at a level from about 0.05% to about 30% by weight, preferably from about 0.05% to about 20% by weight, and more preferably from about 0.1% to about 10% by weight of the composition.

[0061] Nonionic surfactants that are suitable for use herein include alkyl polysaccharides, as disclosed in U.S. Pat. No. 4,565,647 to Llenado. Especially preferred are those nonionic surfactants that have a hydrophobic group containing from about 10 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms in addition to at least one hydrophilic saccharide group such as glucose. Most preferred are alkylpolysaccharides whose hydrophobic groups are sourced from bio-based materials such as coconut or palm oil, and whose hydrophilic groups are source from bio-based materials, an example of which is glucose sourced from corn. Certain alkylpolysaccharides that contain low levels of phenyl derivatives may also be acceptable for use with the formulations presented herein, providing the source of phenyl is bio-based.

[0062] Further suitable nonionic surfactants include addition products of fatty alcohols, fatty acids, and fatty amines (most preferably sourced from bio-based materials such as vegetable oils), coupled with alkoxylation agents such as ethylene oxide (EO), propylene oxide (PO), isopropylene oxide (IPO), or butylene oxide (BO), or a mixture thereof. While most alkylene oxide units are derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources in the future. Moreover, it was found that a number of such ingredients had detectable levels of 1,4-dioxane as a contaminant. Such sources of alcohol alkoxylates must be scrupulously avoided, with preference given to sources that have no detectable level of 1,4-dioxane. Any of the alkoxylated materials of the particular type described hereinafter can be used as the nonionic surfactant. Preferably, the nonionic surfactant is selected from the group consisting of primary and secondary alcohol ethoxylates as well as mixtures thereof. Nonionic surfactants may also contain a mixture of alcohol ethoxylates and propoxylates and mixtures thereof. Further examples of suitable surfactants

are described in McCutcheon's Vol. 1: Emulsifiers and Detergents, North American Ed., McCutcheon Division, MC Publishing Co., 1995.

[0063] Highly preferred nonionic surfactants are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices or fabrics into which they are introduced or applied, either during treatment followed by drying and/or curing, or after drying and/or curing followed by normal exposure to air, moisture or sunlight exposure.

AMPHOTERIC AND ZWITTERIONIC SURFACTANTS

[0064] The compositions of the present disclosure can contain amphoteric and/or zwitterionic surfactants. When an amphoteric or zwitterionic surfactant is added to a composition of the present disclosure, it can typically be added at a level from about 0.05% to about 30%, preferably from about 0.05% to about 20% by weight, and more preferably from about 0.1% to about 10% by weight of the composition.

[0065] Suitable amphoteric surfactants include amine oxides having the formula $(R_1)(R_2)(R_3)NO$ wherein each of R_1 , R_2 and R_3 is independently a saturated substituted or unsubstituted, linear or branched hydrocarbon chain containing from 1 to 30 carbon atoms. Preferred amine oxide surfactants that can be used herein include amine oxides having the formula $(R_1)(R_2)(R_3)NO$ wherein R_1 is a hydrocarbon chain having from 1 to 30 carbon atoms, preferably from 10 to 20, more preferably from 10 to 16, further preferably from 10 to 12, and wherein R_2 and R_3 are independently substituted or unsubstituted, linear or branched hydrocarbon chains comprising from 1 to 4 carbon atoms, preferably from 1 to 3 carbon atoms, and more preferably are methyl groups. R_1 may be a saturated substituted or unsubstituted, linear or branched hydrocarbon chain. Suitable amine oxides for use herein are, for instance, naturally derived $C_{12}-C_{16}$ amine oxides commercially available from Lonza Group and Stepan Company. It is especially preferred that the pendent alkyl groups R_2 and R_3 are derived from bio-based sources, such as wood alcohol.

[0066] Suitable zwitterionic surfactants for use with the formulations presented herein may contain both cationic and anionic hydrophilic groups on the same molecule at a relatively wide pH range. A typical cationic group is a quaternary ammonium group, although other positively charged groups like phosphonium, imidazolium and sulfonium groups can be used. Typical anionic hydrophilic groups are carboxylates and sulfonates, although other groups like sulfates, phosphonates, and the like can be used. A generic formula for some zwitterionic surfactants that can be used herein is $R_1-N'(R_2)(R_3)R_4X$, wherein R_1 is a hydrophobic group comprising from 10 to 30 carbon atoms; R_2 and R_3 are each C_1-C_4 alkyl, hydroxylalkyl or other substituted alkyl group which can also be joined to form ring structures with the N; R_4 is a moiety joining the cationic nitrogen atom to the hydrophilic group and is typically an alkylene, hydroxy alkylene, or polyalkoxy group containing from 1 to 10 carbon atoms; and X is the hydrophilic group which is preferably a carboxylate or sulfonate group. Preferred hydrophobic groups R_1 are bio-based alkyl groups containing from 10 to 24, preferably less than 18, and more preferably less than 16 carbon atoms. The hydrophobic group can contain unsaturation and/or substituents and/or linking groups such as aryl groups, amido groups, ester groups and the like. In general, the simple alkyl groups are preferred for cost and stability reasons. It is especially preferred if the pendent alkyl groups R_2 and R_3 could be derived from bio-based sources, such as methyl groups derived from bio-based sources such as wood alcohol. Examples of amphoteric surfactants include alkylamphoglycinates, and alkyl iminopropionate. Highly preferred zwitterionic surfactants include betaine and sulphobetaine surfactants, derivatives thereof or mixtures thereof. The betaine or sulphobetaine surfactants are preferred herein as they are particularly suitable for the cleaning of delicate materials, including fine fabrics such as silk, wool and other naturally derived textile materials. Betaine and sulphobetaine surfactants are also extremely mild to the skin and/or fabrics to be treated that come in contact with the user's skin.

[0067] Suitable betaine and sulphobetaine surfactants to be used herein include the betaine/sulphobetaine and betaine-like detergents wherein the molecule contains both basic and acidic groups which form an inner salt giving the molecule both cationic and anionic hydrophilic groups over a broad range of pH values. Some common examples of these detergents are described in U.S. Pat. No. 2,082,275 to Daimler, et al., U.S. Pat. No. 2,702,279 to Funderburk, et al., and U.S. Pat. No. 2,255,082 to Orthner, et al. Further examples of suitable surfactants are described in McCutcheon's Vol. 1: Emulsifiers and Detergents, North American Ed., McCutcheon Division, MC Publishing Co., 1995.

[0068] Highly preferred materials of this class of amphoteric and zwitterionic surfactants are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

CATIONIC SURFACTANTS

[0069] The compositions of the present disclosure can contain a cationic surfactant. When a cationic surfactant is added to the compositions disclosed herein, it can typically be added at a level from about 0.05% to about 30% by weight, preferably from about 0.05% to about 20% by weight, and more preferably from about 0.1% to about 10% by weight of the

composition.

[0070] The cationic surfactant can optionally be one or more fabric softener actives. Preferred fabric softening actives according to the present disclosure include amines and quaternized amines. The following are examples of preferred softener actives: N,N-di(tallowyl-oxy-ethyl)-N,N-dimethyl ammonium chloride; N,N-di(canolyl-oxy-ethyl)-N,N-dimethyl ammonium chloride; N,N-di(tallowyl-oxy-ethyl)-N-methyl, N-(2-hydroxyethyl) ammonium methyl sulfate; N,N-di(canolyl-oxy-ethyl)-N-methyl, N-(2-hydroxyethyl) ammonium methyl sulfate; N,N-di(tallowylamidoethyl)-N-methyl, N-(2-hydroxyethyl) ammonium methyl sulfate; N,N-di(2-tallowyloxy-2-oxo-ethyl)-N,N-dimethyl ammonium chloride; N,N-di(2-canolyloxy-2-oxo-ethyl)-N,N-dimethyl ammonium chloride; N,N-di(2-tallowyloxyethylcarbonyloxyethyl)-N,N-dimethyl ammonium chloride; N,N-di(2-canolyloxyethylcarbonyloxyethyl)-N,N-dimethyl ammonium chloride; N-(2-tallowyloxy-2-ethyl)-N-(2-tallowyloxy-2-oxo-ethyl)-N,N-dimethyl ammonium chloride; N-(2-canolyloxy-2-ethyl)-N-(2-canolyloxy-2-oxo-ethyl)-N,N-dimethyl ammonium chloride, N,N,N-tri(tallowyl-oxy-ethyl)-N-methyl ammonium chloride; N,N,N-tri(canolyl-oxy-ethyl)-N-methyl ammonium chloride; N-(2-tallowyloxy-2-oxoethyl)-N-(tallowyl)-N,N-dimethyl ammonium chloride; N-(2-canolyloxy-2-oxoethyl)-N-(canolyl)-N,N-dimethyl ammonium chloride; 1,2-ditallowyloxy-3-N,N,N-trimethylammoniopropane chloride; and 1,2-dicanolyloxy-3-N,N,N-trimethylammoniopropane chloride; and mixtures of the above actives. Particularly preferred is N,N-di(tallowyl-oxy-ethyl)-N,N-dimethyl ammonium chloride, where the tallow chains are at least partially unsaturated and N,N-di(canolyl-oxy-ethyl)-N,N-dimethyl ammonium chloride, N,N-di(tallowyl-oxy-ethyl)-N-methyl, N-(2-hydroxyethyl) ammonium methyl sulfate; N,N-di(canolyl-oxy-ethyl)-N-methyl, N-(2-hydroxyethyl) ammonium methyl sulfate; and mixtures thereof. Additional fabric softening agents useful herein are described in U.S. Pat. No. 5,643,865 to Mermelstein, *et al.*; U.S. Pat. No. 5,622,925 to de Buzzaccarini, *et al.*; U.S. Pat. No. 5,545,350 to Baker, *et al.*; U.S. Pat. No. 5,474,690 to Wahl, *et al.*; U.S. Pat. No. 5,417,868 to Turner, *et al.*; U.S. Pat. No. 4,661,269 to Trinh, *et al.*; U.S. Pat. No. 4,439,335 to Burns; U.S. Pat. No. 4,401,578 to Verbruggen; U.S. Pat. No. 4,308,151 to Cambre; U.S. Pat. No. 4,237,016 to Rudkin, *et al.*; U.S. Pat. No. 4,233,164 to Davis; U.S. Pat. No. 4,045,361 to Watt, *et al.*; U.S. Pat. No. 3,974,076 to Wiersema, *et al.*; U.S. Pat. No. 3,886,075 to Bernadino; U.S. Pat. No. 3,861,870 to Edwards, *et al.*; and European Patent Application publication No. 472,178, to Yamamura, *et al.*

[0071] Other suitable cationic surfactants include ethoxylated quaternary ammonium surfactants. Some preferred ethoxylated quaternary ammonium surfactants include PEG-5 cocoammonium methosulfate; PEG-15 cocoammonium chloride; PEG-15 oleoammonium chloride; and bis(polyethoxyethanol) tallow ammonium chloride. While these cationic surfactants are not preferred due to the ethylene oxide units usually being petrochemically-based, it is envisioned that the ethylene oxide units could also be bio-based. Further examples of suitable surfactants are described in McCutcheon's Vol. 1: Emulsifiers and Detergents, North American Ed., McCutcheon Division, MC Publishing Co., 1995.

[0072] The counterion to these cationic surfactants may be selected, without limitation, from the group consisting of fluoride, chloride, bromide, iodide, chlorite, chlorate, hydroxide, hypophosphite, phosphite, phosphate, carbonate, formate, acetate, lactate, and other carboxylates, oxalate, methyl sulfate, ethyl sulfate, benzoate, and salicylate, and the like. Highly preferred materials of this class of cationic surfactants and their counterions are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

BUILDERS AND pH ADJUSTERS

[0073] Builders are materials used to boost the performance of surfactants used for cleaning. Their best builder compounds react with multivalent cations, "softening" water by removing "hardness" ions (e.g., calcium and magnesium) that bind with surfactants, reducing their effectiveness. Moreover these hardness ions can react with stains, making them more difficult to remove. Some builders also modify solution pH to provide alkalinity, which aids cleaning (stain neutralization, saponification, surface modification). Further, some builders can disperse and/or suspend soils, due to their ability to modify the surface charge on the soils that come into solution.

[0074] Adjustment of pH may be carried out by including a small quantity of an acid in the formulation. Because no strong pH buffers need be present, only small amounts of acid may be required. The pH may be adjusted with inorganic or organic acids, for example hydrochloric acid or alternatively with monobasic or dibasic organic acids, such as acetic acid, maleic acid or in particular glycolic acid. Additional acids that can be used include, but are not limited to, methyl sulfonic, hydrochloric, sulfuric, phosphoric, citric, maleic, and succinic acids.

[0075] Adjustment of pH may be carried out by including a small quantity of a base in the formulation. Because no strong pH buffers need be present, only small amounts of base may be required. The pH may be adjusted with inorganic bases, including, but not limited to, alkali metal or alkaline earth metal salts of hydroxides, carbonates, bicarbonates, borates, sulfonates, phosphates, phosphonates and silicates. The pH may be adjusted with organic bases, including, but not limited to, salts of monocarboxylic acids, salts of dicarboxylic acids, salts of citric acid and other suitable organic acids with water soluble conjugate bases presented previously herein. The pH may be adjusted with organic bases such as the alkanolamines including methanol, ethanol and propanol amines, including dimethanol-, diethanol- and dipropanola-

mines, and including trimethanol-, triethanol- and tripropanolamines.

[0076] Highly preferred materials of this class of pH adjusters are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

SOLVENTS

[0077] The cleaning compositions can contain organic solvents that act as diluents, coupling agents, and to some extent aid cleaning. It is preferred that such solvents be bio-based, and while many solvents are typically obtained from petrochemical sources, it is envisioned that they could be derived from bio-based sources. Further preferred are solvents that do not appreciably contribute to VOCs.

[0078] Examples of organic solvents include, but are not limited to, C₁-C₆ alkanols, C₁-C₆ diols, C₁-C₁₀ alkyl ethers of alkylene glycols, C₃-C₂₄ alkylene glycol ethers, polyalkylene glycols, short chain carboxylic acids, short chain esters, isoparaffinic hydrocarbons, mineral spirits, alkyl aromatics, terpenes, terpene derivatives, terpenoids, terpenoid derivatives, formaldehyde, and pyrrolidones. Alkanols include, but are not limited to, methanol, ethanol, n-propanol, isopropanol, butanol, pentanol, and hexanol, and isomers thereof. Diols include, but are not limited to, methylene, ethylene, propylene and butylene glycols. Alkylene glycol ethers include, but are not limited to, ethylene glycol monopropyl ether, ethylene glycol monobutyl ether, ethylene glycol monohexyl ether, diethylene glycol monopropyl ether, diethylene glycol monobutyl ether, diethylene glycol monohexyl ether, propylene glycol methyl ether, propylene glycol ethyl ether, propylene glycol n-propyl ether, propylene glycol monobutyl ether, propylene glycol t-butyl ether, di- or tri-polypropylene glycol methyl or ethyl or propyl or butyl ether, acetate and propionate esters of glycol ethers. Short chain esters include, but are not limited to, glycol acetate, and cyclic or linear volatile methylsiloxanes. Water insoluble solvents such as isoparaffinic hydrocarbons, mineral spirits, alkyl aromatics, terpenoids, terpenoid derivatives, terpenes, and terpenes derivatives can be mixed with a water-soluble solvent when employed.

SOIL RELEASE AGENTS

[0079] The composition can include a soil release agent that is present from about 0% to about 5% by weight, preferably from about 0.05% to about 3% by weight, and more preferably from about 0.1% to about 2% by weight of the composition. Polymeric soil release agents useful in the present disclosure include co-polymeric blocks of terephthalate and polyethylene oxide or polypropylene oxide, and the like. A preferred soil release agent is a copolymer having blocks of terephthalate and polyethylene oxide. While most terephthalate and alkylene oxide units are derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. These polymers may be comprised of repeating units of ethylene terephthalate and polyethylene oxide terephthalate at a molar ratio of ethylene terephthalate units to polyethylene oxide terephthalate units from about 25:75 to about 35:65, and the polyethylene oxide terephthalate containing polyethylene oxide blocks having molecular weights from about 300 to about 2000. The molecular weight of this type of polymeric soil release agent can be in the range from about 5,000 to about 55,000. Suitable soil release agents are disclosed in U.S. Pat. Nos. 4,702,857 to Gosselink, 4,711,730 to Gosselink, et al., 4,713,194 to Gosselink; 4,877,896 to Maldonado, et al.; 4,956,447 Gosselink, et al.; and 4,749,596 to Po, et al. Especially desirable optional ingredients are polymeric soil release agents comprising block copolymers of polyalkylene terephthalate and polyoxyethylene terephthalate, and block copolymers of polyalkylene terephthalate and polyethylene glycol. The polyalkylene terephthalate blocks may preferably comprise ethylene and/or propylene groups. Many such soil release polymers are nonionic, for example, the nonionic soil release polymer described in U.S. Pat. No. 4,849,257 to Borcher, Sr., et al. The polymeric soil release agents useful in the present disclosure can include anionic and cationic polymeric soil release agents. Suitable anionic polymeric or oligomeric soil release agents are disclosed in U.S. Pat. No. 4,018,569 to Chang,. Other suitable polymers are disclosed in U.S. Pat. No. 4,808,086 to Evans, et al.

[0080] Highly preferred materials of this class of soil release polymers are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

ANTISTATIC AGENTS

[0081] The composition can include antistatic agents, which can be present at a level from about 0% to about 5% by weight, preferably from about 0.005% to about 5% by weight, more preferably from about 0.05% to about 2% by weight, and further preferably from about 0.2% to about 1% of the composition. While many of these compounds are derived from petrochemical sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources.

Preferred antistatic agents of the present disclosure include cationic surfactants, including quaternary ammonium compounds such as alkyl benzyl dimethyl ammonium chloride; dicoco quaternary ammonium chloride; coco dimethyl benzyl ammonium chloride; soya trimethyl quaternary ammonium chloride; hydrogenated tallow dimethyl benzyl ammonium chloride; and methyl dihydrogenated tallow benzyl ammonium chloride. Other preferred antistatic agents of the present disclosure are alkyl imidazolinium salts. Other preferred antistatic agents are the ion pairs of, e.g., anionic detergent surfactants and fatty amines, or quaternary ammonium derivatives thereof, e.g., those disclosed in U.S. Pat. No. 4,756,850 to Nayar. Other preferred antistatic agents are ethoxylated and/or propoxylated sugar derivatives; while most alkylene oxide units are derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. Preferred antistatic agents include monolauryl trimethyl ammonium chloride, hydroxycetyl hydroxyethyl dimethyl ammonium chloride (available from BASF Corporation under the trade name DEHYQUART E), and ethyl bis(polyethoxyethanol) alkyl ammonium ethyl sulfate (available from Evonik Corporation under the trade name VARIQUAT 66), polyethylene glycols, polymeric quaternary ammonium salts (such as those available from Rhodia Group under the MIRAPOL trade name), quaternized polyethyleneimines, vinylpyrrolidone/methacrylamidopropyl trimethylammonium chloride copolymer (available from Ashland Inc. under the trade name GAFQUAT HS-100), triethonium hydrolyzed collagen ethosulfate (available from Angene Chemical under the trade name QUAT-PRO E), and mixtures thereof.

[0082] Highly preferred materials of this class of antistatic agents are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

FRAGRANCE MATERIALS

[0083] While not preferred due to their propensity to induce untoward symptoms in sensitized individuals, it has been discovered that truly natural, bio-based fragrance materials may be added to the composition. It appears that preferred fragrance materials are comprised of extracts of natural products, upon which no additional functionalization reactions have been carried out. Further, preferred fragrance materials should not have been isolated in such a way as to introduce petrochemical solvents, which appear to further exacerbate symptoms of sensitization. Such materials may have been isolated by methods well-known to the industry such as extraction with suitable solvents, supercritical fluid extraction, steam distillation, rectification, and expression. It is also foreseen that by adding fragrance sources such as plant materials directly to the product, and relying on the product matrix itself to extract the desired fragrance notes, one can obtain desired fragrance notes.

[0084] The selection of the perfume or perfumes maybe based upon the application, the desired effect on the consumer, and preferences of the formulator. The perfume selected for use in the compositions and formulations of the present disclosure may contain ingredients with odor characteristics which are preferred in order to provide a fresh impression on the surface to which the composition is directed, for example, those which provide a fresh impression for fabrics. Such perfume may be preferably present at a level from about 0.01% to about 5% by weight, preferably from about 0.05% to about 3% by weight, and more preferably from about 0.1% to about 2% by weight of the total composition.

[0085] Preferably, the fragrance materials are mixtures comprising multiple ingredients selected from the group consisting of aromatic and aliphatic esters having molecular weights from about 130 to about 250; aliphatic and aromatic alcohols having molecular weights from about 90 to about 240; aliphatic ketones having molecular weights from about 150 to about 260; aromatic ketones having molecular weights from about 150 to about 270; aromatic and aliphatic lactones having molecular weights from about 130 to about 290; aliphatic aldehydes having molecular weights from about 140 to about 200; aromatic aldehydes having molecular weights from about 90 to about 230; aliphatic and aromatic ethers having molecular weights from about 150 to about 270; and condensation products of aldehydes and amines having molecular weights from about 180 to about 320; and mixtures thereof.

[0086] Highly preferred materials of this class of fragrances and perfumes are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

ANTIMICROBIALS AND PRESERVATIVES

[0087] Antimicrobials and/or preservatives can be used with the formulations presented herein. Typical concentrations for biocidal effectiveness of these compounds may range from about 0.001% to about 0.8% by weight, preferably from about 0.005% to about 0.3% by weight, and more preferably from about 0.01% to 0.2% by weight of the usage composition. The corresponding concentrations for the concentrated compositions are from about 0.003% to about 2% by weight, preferably from about 0.006% to about 1.2% by weight, and more preferably from about 0.1% to about 0.8% by weight of

the concentrated compositions.

[0088] Preservatives are especially preferred when organic compounds that are subject to microorganisms are added to the compositions of the present disclosure, especially when they are used in aqueous compositions. When such compounds are present, long term and even short-term storage stability of the compositions and formulations becomes an important issue since contamination by certain microorganisms with subsequent microbial growth often results in an unsightly and/or malodorous solution. Therefore, because microbial growth in these compositions and formulations is highly objectionable when it occurs, it is preferable to include a solubilized water-soluble, antimicrobial preservative, which is effective for inhibiting and/or regulating microbial growth in order to increase storage stability of the preferably clear and often aqueous compositions and formulations of the present disclosure.

[0089] Typical microorganisms that can be found in laundry products include bacteria, for example, *Bacillus thuringiensis* (cereus group) and *Bacillus sphaericus*, and fungi, for example, *Aspergillus ustus*. *Bacillus sphaericus* is one of the most numerous members of *Bacillus* species in soils. In addition, microorganisms such as *Escherichia coli* and *Pseudomonas aeruginosa* are found in some water sources, and can be introduced during the preparation of aqueous solutions of the present disclosure. It is preferable to use a broad spectrum preservative, for example, one that is effective on both bacteria (both Gram positive and Gram negative) and fungi. A limited spectrum preservative, for example, one that is only effective on a single group of microorganisms, for example, fungi, can be used in combination with a broad spectrum preservative or other limited spectrum preservatives with complimentary and/or supplementary activity. A mixture of broad-spectrum preservatives can also be used. Antimicrobial preservatives useful in the present disclosure can be biocidal compounds, that is, substances that kill microorganisms, or biostatic compounds, that is, substances that inhibit and/or regulate the growth of microorganisms.

[0090] Preferred antimicrobial preservatives include those that are water-soluble and are effective at low levels. While such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. In general, the water-soluble preservatives that may be used include organic sulfur compounds, halogenated compounds, cyclic organic nitrogen compounds, low molecular weight aldehydes, quaternary compounds, dehydroacetic acid, phenyl and phenoxy compounds, and mixtures thereof. Examples of preservatives useful with the formulations presented herein include, but are not limited to, the short chain alkyl esters of p-hydroxybenzoic acid (commonly known as parabens); N-(4-chlorophenyl)-N-(3,4-dichlorophenyl) urea (also known as 3,4,4-trichlorocarbanilide or triclocarban); 2,4,4-trichloro-2'-hydroxydiphenyl ether, commonly known as Triclosan®; a mixture of about 77% 5-chloro-2-methyl-4-isothiazolin-3-one and about 23% 2-methyl-4-isothiazolin-3-one, a broad spectrum preservative available from the Dow Chemical Company as a 1.5% aqueous solution under the trade name KATHON CG; 5-bromo-5-nitro-1,3-dioxane, available from BASF Corporation under the trade name BRONIDOX L; 2-bromo-2-nitropropane-1,3-diol, available from Dow Chemical Company under the trade name BRONOPOL; 1,1-hexamethylenebis(5-p-(chlorophenyl)biguanide) (commonly known as chlorhexidine) and its salts, for example, with acetic and gluconic acids; a 95:5 mixture of 1,3- bis(hydroxymethyl)-5,5-dimethyl-2,4-imidazolidinedione and 3-butyl-2-iodopropynyl carbamate, available from Lonza Group under the trade name GLYDANT Plus; N-[1,3-bis(hydroxymethyl)2,5-dioxo-4-imidazolidinyl]-N,N'-bis(hydroxymethyl) urea, commonly known as diazolidinyl urea, available from Ashland Inc. under the trade name GERMALL II; N,N'-methylenebis-[N'-[1-(hydroxymethyl)-2,5-dioxo-4-imidazolidinyl] urea] (commonly known as imidazolidinyl urea), available, for example, from 3V-Sigma under the trade name ABIOL, from Induchem USA, Inc. under the trade name UNICIDE U-13, and from Ashland Inc. under the trade name GERMALL 115; polymethoxy bicyclic oxazolidine, available from Ashland Inc. under the trade name NUOSEPT; formaldehyde; glutaraldehyde; polyaminopropyl biguanide under the trade name COSMOCIL CQ or MIKROKIL from Lonza Group; and mixtures thereof. In general, however, the preservative can be any organic preservative material that is appropriate for applying to a fabric. With respect to the embodiments presented herein, such preservative(s) will preferably not cause damage to a fabric appearance, for example, through discoloration, coloration, or bleaching of the fabric. If the antimicrobial preservative is included in the compositions and formulations of the present disclosure, it is preferably present in an effective amount, wherein an "effective amount" means a level sufficient to prevent spoilage or prevent growth of inadvertently added microorganisms for a specific period of time. Preferred levels of preservative are from about 0.0001% to about 0.5% by weight, more preferably from about 0.0002% to about 0.2% by weight, further preferably from about 0.0003% to about 0.1% by weight, of the composition. Optionally, the preservative can be used at a level that provides an antimicrobial effect on the treated fabrics.

[0091] The composition may suitably use an optional solubilized, water-soluble antimicrobial active, useful in providing protection against organisms that become attached to the treated material. The free, uncomplexed antimicrobial, e.g., antibacterial, active provides an optimum antibacterial performance. Sanitization of fabrics can be achieved by the compositions of the present disclosure containing, antimicrobial materials, e.g., antibacterial halogenated compounds, quaternary compounds, and phenolic compounds. Some of the more robust antimicrobial halogenated compounds which can function as disinfectants/sanitizers as well as finish product preservatives, and are useful in the compositions of the present disclosure include 1,1'-hexamethylene bis(5-(p-chlorophenyl)biguanide), commonly known as chlorhexidine, in addition to its salts, e.g., with hydrochloric, acetic and gluconic acids. The gluconate salt is highly water-soluble, at about

70% by weight in water, while the diacetate salt has a solubility of about 1.8% weight in water. When chlorhexidine is used as a sanitizer with the formulations discussed herein, it can typically be present at a level from about 0.001% to about 1.0% by weight, preferably from about 0.002% to about 0.3% by weight, and more preferably from about 0.01% to about 0.1% by weight of the usage composition. In some cases, a level from about 1% to about 2% by weight may be needed for virucidal

5 activity. Other useful biguanide compounds include COSMOCIL CQ, VANTOCII, IB, including poly (hexamethylene biguanide) hydrochloride. Other useful cationic antimicrobial agents include the bis-biguanide alkanes. Usable water soluble salts of the above are chlorides, bromides, sulfates, alkyl sulfonates such as methyl sulfonate and ethyl sulfonate, phenylsulfonates such as p-methylphenyl sulfonates, nitrates, acetates, gluconates, and the like. Non-limiting examples 10 of useful quaternary compounds include: (1) benzalkonium chlorides and/or substituted benzalkonium chlorides such as commercially available BARQUAT (available from Lonza), MAQUAT (available from Pilot Chemical), VARIQUAT (available from Evonik), and HYAMINE (available from Lonza); (2) dialkyl quaternary such as BARDAC products of Lonza, (3) N-(3-chloroallyl) hexaminium chlorides such as DOWICIDE and DOWICIL available from Dow; (4) benzethonium chloride such as HYAMINE 1622 from Lonza; (5) methylbenzethonium chloride represented by HYAMINE 10X supplied by Lonza, (6) cetylpyridinium chloride such as Cepacol chloride available from of Merrell Labs.

15 [0092] Preferred antimicrobial compounds for use herein include quaternary ammonium compounds containing alkyl or substituted alkyl groups, alkyl amide and carboxylic acid groups, ether groups, unsaturated alkyl groups, and cyclic quaternary ammonium compounds, which can be chlorides, dichlorides, bromides, methylsulphates, chlorophenates, cyclohexyl sulphamates or salts of the other acids. Among the useful cyclic quaternary ammonium compounds are the 20 following: alkylpyridinium chlorides and/or sulphates, the alkyl group being preferably cetyl, dodecyl or hexadecyl group; - alkylisoquinolyl chlorides and/or bromides, the alkyl group being preferably dodecyl group. Particularly suitable quaternary ammonium compounds for use herein include alkyltrimethylbenzyl ammonium chloride, octyl decyl dimethylammonium chloride, dioctyl dimethyl ammonium chloride, didecyl dimethyl ammonium chloride, alkyl dimethyl ammonium saccharinate, cetylpyridinium and mixtures thereof.

25 [0093] It is also envisioned that certain inorganic materials based on silver, copper, or clays materials such as Dragonite™ Halloysite clay (Applied Minerals, New York, NY) may be suitable for this purpose. Silver and copper materials may be embedded within the packaging matrix, so as to keep liquids contained therein preserved.

30 [0094] Highly preferred materials of this class of antimicrobials and preservatives are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

DYES AND COLORANTS

35 [0095] Colorants can be added to the formulations disclosed herein. As many people manifest sensitivity to synthetic dyes, they are not preferred. However, certain natural colorants such as chlorophyll may be suitable for incorporation herein. Pigments, which are insoluble colorants, may also be suitable for incorporation in the formulations described herein. Typical concentrations of these compounds may range from about 0.001% to about 0.8% by weight, preferably from about 0.005% to about 0.3% by weight, and more preferably from about 0.01% to 0.2% by weight of the composition.

40 [0096] Colorants and dyes, especially bluing agents, can be optionally added to the compositions of the present disclosure for visual appeal and performance impression. When colorants are used, they may be used at extremely low levels to avoid fabric staining.

45 [0097] Highly preferred materials of this class of dyes and colorants are those that do not effectively bind to or permanently dye or color fabrics treated by use of the compositions disclosed herein, nor cause any significant color change, nor impart any discoloration, such as graying, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

VISCOSITY CONTROL AGENTS

50 [0098] Optionally added viscosity control agents can be organic or inorganic in nature and may either lower or raise the viscosity of the formulation. While many such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. Examples of organic viscosity modifiers to lower viscosity are aryl carboxylates and sulfonates (for example including, but not limited to benzoate, 2-hydroxybenzoate, 2-aminobenzoate, benzenesulfonate, 2-hydroxybenzenesulfonate, 2-aminobenzenesulfonate), fatty acids and esters, fatty alcohols, and water-miscible solvents such as short chain alcohols. Examples of inorganic viscosity control agents are water-soluble ionizable salts. A wide variety of ionizable salts can be used. Examples of suitable salts are the halides and acetates of ammonium ion and the group IA and IIA metals of the Periodic Table of the Elements, for example, calcium chloride, lithium chloride, sodium chloride, potassium chloride, magnesium chloride, ammonium

chloride, sodium bromide, potassium bromide, calcium bromide, magnesium bromide, ammonium bromide, sodium iodide, potassium iodide, calcium iodide, magnesium iodide, ammonium iodide, sodium acetate, potassium acetate, or mixtures thereof. Calcium chloride is preferred. The ionizable salts are particularly useful during the process of mixing the ingredients to make the compositions herein, and later to obtain the desired viscosity. The amount of ionizable salts used

5 depends on the amount of active ingredients used in the compositions and can be adjusted according to the desire of the formulator. Typical levels of salts used to control the composition viscosity are from 0 to about 10% by weight, preferably from about 0.01% to about 6% by weight, and more preferably from about 0.02% to about 3% by weight of the composition.

[0099] Viscosity modifiers or thickening agents can be added to increase the ability of the compositions to stably suspend water-insoluble articles, for example, perfume microcapsules. Such materials include hydroxypropyl substituted 10 guar gum (such as that available from Rhodia Group under the trade name JAGUAR HP200), polyethylene glycol (such as that available from Dow Chemical Corporation under the trade name CARBOWAX 20M), hydrophobically modified hydroxyethylcellulose (such as that available from the Ashland Inc. under the trade name NATROSOL Plus), and/or 15 organophilic clays (for example, hectorite and/or bentonite clays such as those available from Elementis Specialties under the name BENTONE 27, 34 and 38 or from Eckart America under the trade name BENTOLITE L; and those described in U.S. Pat. No. 4,103,047 to Zaki, et al.). These viscosity raisers or thickeners can typically be used at levels from about 0.5% to about 30% by weight, preferably from about 1% to about 5% by weight, more preferably from about 1.5% to about 3.5% by weight, and further preferably from about 2% to about 3% by weight, of the composition.

[0100] Highly preferred materials of this class of thickeners and viscosity control and viscosity modifiers are those that do 20 not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

PEARLIZING AND OPACIFYING AGENTS

25 [0101] Examples of pearlizing or opacifying agents that can be added to the compositions disclosed herein include, but are not restricted to, glycol distearate, propylene glycol distearate, and glycol stearate. Some of these products are available from PMC Group under the KEMESTER trade name. While many such compounds are commonly derived from petrochemicals sources at present, and are as such not preferred, it is envisioned that they could be derived from bio-based sources at some future point.

[0102] Highly preferred materials of this class of pearlizing and opacifying agents are those that do bind to treated 30 fabrics, nor cause any significant color change nor impart any discoloration, such as whitening, graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

ANTIOXIDANTS AND SUNSCREEN MATERIALS

[0103] Examples of antioxidants that can be added to the compositions of herein are propyl gallate, available from 40 Eastman Chemical Products, Inc. under the trade names TENOX PG and TENOX S-1, and dibutylated hydroxytoluene, available from UOP Inc. under the trade name SUSTANE BHT. Also preferred are antioxidants for providing sun-fade protection for fabrics treated with composition of the present disclosure, such antioxidants being described in EP0773982. Preferred antioxidants include 2-(N-methyl-N-cocoamino)ethyl-3',5'-di-tert-butyl-4'-hydroxybenzoate; 2-(N, N-dimethyl-amino)ethyl-3',5'-di-tert-butyl-4'-hydroxybenzoate; 2-(N- methyl-N-cocoamino)ethyl-3',4',5'-trihydroxybenzoate; and mixtures thereof, more preferably 2-(N-methyl-N-cocoamino)ethyl-3',5'-di-tert-butyl-4'-hydroxy benzoate. Of these 45 compounds, the butylated derivatives are preferred in the compositions of the present disclosure because tri-hydroxybenzoates have a tendency to discolor upon exposure to light. While many such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources in the future. The antioxidant compounds of the present disclosure demonstrate light stability in the compositions of the 50 present disclosure. Light stable as used herein means that the antioxidant compounds disclosed herein do not discolor when exposed to either sunlight or simulated sunlight for approximately 2 to 60 hours at a temperature of from about 25°C to about 45°C. Antioxidant compounds and free radical scavengers can generally protect dyes from degradation by first preventing the generation of single oxygen and peroxy radicals, and thereafter terminating the degradation pathways. Not to be limited by theory, a general discussion of the mode of action for antioxidants and free radical scavengers is disclosed 55 in Kirk-Othmer Encyclopedia of Chemical Technology, Volume 3, pages 128 - 148, Third Edition (1978).

[0104] The formulations that are the subject of the instant disclosure may comprise an organic sunscreen. Suitable sunscreens can have UVA absorbing properties, UVB absorbing properties, or a combination of both. The formulations newly presented herein may preferably comprise a UVA absorbing sunscreen actives that absorb UV radiation having a

wavelength from about 320 nm to about 400 nm. Suitable UVA absorbing sunscreen actives include dibenzoylmethane derivatives, anthranilate derivatives such as methylanthranilate and homomethyl-1-N-acetylanthranilate, and mixtures thereof. Examples of dibenzoylmethane sunscreen actives are described in U.S. Pat. No. 4,387,089 to De Polo; and in Sunscreens: Development, Evaluation, and Regulatory Aspects edited by N.J. Lowe and N.A. Shaath, Marcel Dekker, Inc (1990). The UVA absorbing sunscreen active is preferably present in an amount to provide broad-spectrum UVA protection either independently, or in combination with, other UV protective actives that may be present in the composition. Preferred UVA sunscreen actives include dibenzoylmethane sunscreen actives and their derivatives. They include, but are not limited to, those selected from 2-methyldibenzoylmethane, 4-methyldibenzoylmethane, 4- isopropyldibenzoylmethane, 4-tert-butylbibenzoylmethane, 2,4-dimethyldibenzoylmethane, 2, 5-dimethyldibenzoylmethane, 4,4'-diisopropylbenzoylmethane, 4-(1,1-dimethylethyl)-4'- methoxydibenzoylmethane, 2-methyl-5-isopropyl-4'-methoxydibenzoylmethane, 2-methyl-5-tert-butyl-4'-methoxydibenzoylmethane, 2,4-dimethyl-4'-methoxydibenzoylmethane, 2,6-dimethyl-4'-tert-butyl-4'-methoxydibenzoylmethane, and mixtures thereof. Preferred dibenzoyl sunscreen actives include those selected from 4-(1,1-dimethylethyl)-4'-methoxydibenzoylmethane, 4-isopropyldibenzoylmethane, and mixtures thereof. A more preferred sunscreen active is 4-(1,1-dimethylethyl)-4'-methoxydibenzoylmethane, which is also known as butylethoxydibenzoylmethane or Avobenzone, is commercially available under the names of PARSOL 1789 from DSM Nutritional Products, LLC and EUSOLEX 9020 from EMD Chemicals Inc./Rona. The sunscreen 4-isopropyldibenzoylmethane, which is also known as isopropyldibenzoylmethane, is commercially available from EMD Chemicals Inc./Rona under the name of EUSOLEX 8020. The formulations of the instant disclosure may preferably further comprise a UVB sunscreen active that absorbs UV radiation having a wavelength of from about 290 nm to about 320 nm. The compositions may 5 preferably comprise an amount of the UVB sunscreen active that is safe and effective to provide UVB protection either independently, or in combination with, other UV protective actives that may be present in the compositions. The compositions preferably comprise from about 0.1% to about 16%, more preferably from about 0.1% to about 12%, and further preferably from about 0.5% to about 8% by weight, of UVB absorbing organic sunscreen. A wide variety of UVB 10 sunscreen actives are suitable for use herein. Non-limiting examples of such organic sunscreen actives are described in U.S. Pat. No 5,087,372 to Toyomot and U.S. Pat. Nos. 5,073,371 and 5,073,372 both to Turner, et al. Preferred UVB 15 sunscreen actives are selected from 2-ethylhexyl-2-cyano-3,3-diphenylacrylate (referred to as octocrylene), 2-phenylbenzimidazole-5-sulphonic acid (PBSA), cinnamates and their derivatives such as 2-ethylhexyl-p-methoxycinnamate and octyl-p-methoxycinnamate, TEA salicylate, octyldimethyl PABA, camphor derivatives and their derivatives, and mixtures thereof. Preferred organic sunscreen actives include 2-ethylhexyl-2-cyano-3,3-diphenylacrylate (commonly 20 named octocrylene), 2-phenyl-benzimidazole-5-sulphonic acid (PBSA), octyl-p-methoxycinnamate, and mixtures thereof. Salt and acid neutralized forms of the acidic sunscreens are also useful.

[0105] An agent may also be added to any of the formulations described in the present disclosure to stabilize the UVA sunscreen and to prevent it from photo-degrading on exposure to ultraviolet radiation and thereby maintaining its UVA protection efficacy. Wide ranges of compounds have been cited as providing these stabilizing properties and should be 25 chosen to compliment both the UVA sunscreen and the composition as a whole. Suitable stabilizing agents include, but are not limited to, those described in U.S. Pat. No. 5,972,316 to Robinson; U.S. Pat. No. 5,968,485 to Robinson; U.S. Pat. No. 5,935,556 to Tanner, et al.; and U.S. Pat. 5,827,508 Tanner, et al. Preferred examples of stabilizing agents for use in the present formulations disclosure herein include 2-ethylhexyl-2-cyano-3,3-diphenylacrylate (referred to as octocrylene), ethyl-2-cyano-3,3-diphenyl-acrylate-2-ethylhexyl-3,3-diphenylacrylate, ethyl-3,3-bis (4-methoxyphenyl)acrylate, and mixtures thereof.

[0106] Highly preferred materials of this class of antioxidants and sunscreen actives are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the 30 drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

[0107] The formulations of the present disclosure may preferably deposit from about 0.1 mg/g fabric to about 5 mg/g fabric of the sun-fade actives to reduce the sun fading of the fabric. Repeated treatment of fabric with formulations 35 presented herein, may result in higher deposition levels, which contributes even further to the sun-fading protection benefit.

50 DYE TRANSFER INHIBITORS AND DYE FIXATIVES

[0108] The formulations disclosed herein can comprise from about 0.001% to about 20% by weight, preferably from 45 about 0.5% preferably to about 10% by weight, and more preferably from about 1% to about 5% by weight of one or more dye transfer inhibitors or dye fixing agents.

[0109] Compositions and formulations of the present disclosure can contain ethoxylated amines, amphoteric, betaines, polymers such as polyvinylpyrrolidone, and other ingredients that inhibit dye transfer. While many such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that 55 they could be derived from bio-based sources. Optional dye fixing agents can be cationic, and based on quaternized

nitrogen compounds or on nitrogen compounds having a strong cationic charge that is formed in situ under the conditions of usage. Cationic fixatives are available under various trade names from several suppliers. Representative examples include: CROSCOLOR PMF (July 1981, Code No. 7894) and CROSCOLOR NOFF (January 1988, Code No. 8544) ex Crosfield; INDOSOL E-50 (February 27, 1984, Ref. No. 6008.35.84; polyethyleneamine-based) ex Sandoz; SANDOFIX

5 TPS, ex Sandoz, is a preferred dye fixative for use herein. Additional non-limiting examples include SANDOFIX SWE (a cationic resinous compound) from Sandoz, REWIN SRF, REWIN SRF-O and REWIN DWR Crochet-Beitlich GMBH; Tinofix ECO, Tinofix FRD and Solvent from Ciba-Geigy. Other cationic dye fixing agents are described in "After treatments for Improving the Fastness of Dyes on Textile Fibres", Christopher C. Cook, Rev. Prog. Coloration, Vol. XH, (1982). Dye fixing agents suitable for use in the formulations of the instant disclosure include ammonium compounds such as fatty acid-diamine condensates, inter alia, the hydrochloride, acetate, methosulphate and benzyl hydrochloride salts of diamine esters. Non-limiting examples include oleyldiethyl aminoethylamide, oleylmethyl diethylenediamine methosulphate, and monostearylethylene diaminotrimethylammonium methosulphate. In addition, the N-oxides of tertiary amines; derivatives of polymeric alkyldiamines, polyamine-cyanuric chloride condensates; and aminated glycerol dichlorhydrins are suitable for use as dye fixatives in the compositions of the presented herein.

10 15 [0110] Highly preferred materials of this class of dye transfer inhibitors and dye fixatives are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

20 CHLORINE SCAVENGERS

25 [0111] The compositions of the present disclosure may optionally comprise from about 0.01%, preferably from about 0.02%, more preferably from about 0.25% to about 15%, further preferably to about 10%, and yet more preferably to about 5% of a chlorine scavenger. In cases wherein the cation portion and the anion portion of the non-polymeric scavenger each react with chlorine, the amount of scavenger can be adjusted to fit the needs of the formulator. While many such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. Suitable chlorine scavengers include ammonium salts having the formula: [(R)₃R'N]X wherein each R is independently hydrogen, C₁-C₄ alkyl, C₁-C₄ substituted alkyl, and mixtures thereof; preferably R is hydrogen or methyl, more preferably hydrogen; R' is hydrogen C₁-C₁₀ alkyl, C₁-C₁₀ substituted alkyl, and mixtures thereof. Preferably R is hydrogen and X is a compatible anion. Non-limiting examples include chloride, bromide, citrate, and sulfate; preferably X is chloride. Non-limiting examples of preferred chlorine scavengers include ammonium chloride, ammonium sulfate, and mixtures thereof, preferably ammonium chloride. Other chlorine scavengers include reducing agents such as thiosulfate.

30 35 [0112] Highly preferred materials of this class of chlorine scavengers are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

40 WETTING AGENTS

45 [0113] The formulations and compositions disclosed herein may contain as an optional ingredient from about 0.005% to about 3.0% by weight, and more preferably from about 0.03% to 1.0% by weight of a wetting agent. Such wetting agents may be selected from polyhydroxy compounds. While many such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. Examples of water soluble polyhydroxy compounds that can be used as wetting agents in the compositions disclosed herein include glycerol, polyglycerols having a weight-average molecular weight from about 150 to about 800, and polyoxyethylene glycols and polyoxypropylene glycols having a weight-average molecular weight from about 200 to about 4000, preferably from about 200 to about 1000, and more preferably from about 200 to about 600. Polyoxyethylene glycols having a weight-average molecular weight from about 200 to about 600 are especially preferred. Mixtures of the above-described polyhydroxy compounds may also be used. A particularly preferred polyhydroxy compound is polyoxyethylene glycol having a weight-average molecular weight of about 400, available from Dow Chemical Corporation under the trade name PEG-400.

50 55 [0114] Highly preferred materials of this class of wetting agents are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

ELECTROLYTES

[0115] Suitable inorganic salts for use as an optional electrolyte in the present compositions include MgI_2 , $MgBr_2$, $MgCl_2$, $Mg(NO_3)_2$, $Mg_3(PO_4)_2$, $Mg_2P_2O_7$, $MgSO_4$, magnesium silicate, NaI , $NaBr$, $NaCl$, NaF , Na_3PO_4 , Na_2SO_3 , Na_2SO_4 , $NaNO_3$, $Na_4P_2O_5$, sodium silicate, sodium metasilicate, sodium tetrachloroaluminate, sodium tripolyphosphate (STPP), $Na_2S_3O_7$, sodium zirconate, CaF_2 , $CaCl_2$, $CaBr_2$, CaI_2 , $CaSO_4$, $Ca(NO_3)_2$, KI , KBr , KCl , KF , KNO_3 , KIO_3 , K_2SO_4 , K_2SO_3 , K_3PO_4 , $K_4(P_2O_7)$, potassium pyrosulfate, potassium pyrosulfite, LiI , $LiBr$, $LiCl$, LiF , $LiNO_3$, AlF_3 , $AlCl_3$, $AlBr_3$, AlI_3 , $Al_2(SO_4)_3$, $Al(PO_4)$, $Al(NO_3)_3$, aluminum silicate; including hydrates of these salts and including combinations of these salts or salts with mixed cations e.g. potassium aluminum $AlK(SO_4)_2$ and salts with mixed anions, e.g. potassium tetrachloroaluminate and sodium tetrafluoroaluminate. Salts incorporating cations from groups IIIa, IVa, Va, VIa, VIIa, VIII, Ib, and IIb on the periodic chart with atomic numbers greater than are also useful in reducing dilution viscosity but less preferred due to their tendency to change oxidation states and thus they can adversely affect the odor or color of the formulation or lower weight efficiency. Salts with cations from group Ia or IIa with atomic numbers greater than 20 as well as salts with cations from the lanthanide or actinide series are useful in reducing dilution viscosity, but less preferred due to lower weight efficiency or toxicity. Mixtures of above salts are also useful.

[0116] Also preferred are quaternary ammonium salts, quaternary alkyl ammonium salts, quaternary dialkyl ammonium salts, quaternary trialkyl ammonium salts and quaternary tetraalkyl ammonium salts wherein the alkyl substituent comprises a methyl, ethyl, propyl, butyl or higher C_5-C_{12} linear alkane radical, or combinations thereof. Organic salts useful with the compositions presented herein include magnesium, sodium, lithium, potassium, zinc, and aluminum salts of carboxylic acids, including formates, acetates, propionates, pelargonates, citrates, gluconates, lactates, and aromatic acids such as benzoates, phenolates, and substituted benzoates or phenolates, such as phenolates, salicylates, polyaromatic acids, terephthalates, and polyacids e.g. oxylates, adipates, succinates, benzenedicarboxylates and benzenetricarboxylates. Other useful organic salts include carbonates and/or hydrogen carbonate (HCO_3^{-1}) when the pH is targeted to be alkaline, alkyl and aromatic sulfates and sulfonates, e.g., sodium methyl sulfate, benzene sulfonates and derivatives such as xylene sulfonate, and amino acids.

[0117] Electrolytes can comprise mixed salts of the above single salts, salts neutralized with mixed cations such as potassium/sodium tartrate, partially neutralized salts such as sodium hydrogen tartrate or potassium hydrogen phthalate, and salts comprising one cation with mixed anions.

[0118] Highly preferred mixed salt materials comprising inorganic and organic electrolytes are those that do not cause any significant color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after a drying and/or curing step that may be followed by normal exposure to air, moisture or sunlight.

[0119] Generally, inorganic electrolytes are preferred over organic electrolytes for better weight efficiency and lower costs. Mixtures of inorganic and organic salts can be used. Typical levels of electrolyte in the present compositions can be less than about 10% by weight, preferably from about 0.5% to about 5% by weight, more preferably from about 0.75% to about 2.5% by weight, and further preferably from about 1% to about 2% by weight of the inventive composition.

ENZYMES

[0120] Additional desirable adjuncts may be enzymes (although it may be preferred to also include an enzyme stabilizer), including, but not limited to hydrolases, hydroxylases, cellulases, peroxidases, laccases, mannosases, amylases, lipases and proteases. Proteases are one especially preferred class of enzymes. Typical examples of proteases include Maxatase and Maxacal from Genencor International, Alcalase, Savinase, and Esperase, all available from Novozymes North America, Inc. See also U.S. Pat. No. 4,511,490 to Stanislowski, et al. Further suitable enzymes are amylases, which are carbohydrate-hydrolyzing enzymes. It may also be preferred to include mixtures of amylases and proteases. Suitable amylases include Termamyl from Novozymes, North America Inc, and Maxamyl from Genencor International Co. Still other suitable enzymes are cellulases, such as those described in U.S. Pat. No. 4,479,881 to Tai; U.S. Pat. No. 4,443,355 to Murata, et al.; U.S. Pat. No. 4,435,307 to Barbesgaard, et al.; and U.S. Pat. No. 3,983,082 to Ohya, et al. Yet other suitable enzymes are lipases, such as those described in U.S. Pat. No. 3,950,277 to Silver; U.S. Pat. No. 4,707,291 to Thorn, et al.; U.S. Pat. Nos. 5,296,161 and 5,030,240 both to Wiersema, et al.; and U.S. Pat. No. 5,108,457 to Poulose, et al. The hydrolytic enzyme may be present in an amount of about 0.01-5%, more preferably about 0.01-3%, and further preferably about 0.1-2% by weight of the detergent. Mixtures of any of the foregoing hydrolases are desirable, especially protease/amylase blends.

[0121] Highly preferred materials of this class of enzymes are those that do not cause any significant residual odor or color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

BLEACHING AGENTS


[0122] The compositions disclosed herein may optionally comprise from about 0.01%, preferably from about 0.02% by weight, more preferably from about 0.25% to about 15% by weight, further preferably to about 10% by weight, and yet more preferably to about 5% by weight of a bleaching agent. Suitable bleaching agents include peroxygen and peroxide-releasing compounds. Peroxygen compounds include alkali metal salts of percarbonate, perborate and peroxymonosulfate. Peroxide compounds, including hydrogen peroxide and compounds generating hydrogen peroxide in solution, peroxyacids and precursors to peroxyacids and peroxyimide acids, and metal based oxidants are also suitable. Suitable bleaching agents include preformed peracids and organic peroxides, including alkonyl and acyl peroxides such as tertiary butyl peroxide and benzoyl peroxide, and related alkonyl and acyl peroxide and superoxide derivatives of alkyls and arenes. Additionally, an appropriate bleach activator for the active oxygen source or peroxide may be present, such as those found in Arbogast, et al., U.S. Pat. Nos. 5,739,327 and 5,741,437; Alvarez, et al.; U.S. Pat. No. 5,814,242, Deline, et al.; U.S. Pat. No. 5,877,315; and U.S. Pat. No. 5,888,419 to Casella, et al., which relate to cyanonitrile derivatives; U.S. Pat. Nos. 4,959,187 and 4,778,816 to Fong, et al.; U.S. Pat. Nos. 5,112,514 and 5,002,691 to Bolkan, et al., and U.S. Pat. No. 5,269,962 to and Brodbeck, et al., which relate to alkanoyloxyacetyl derivatives; and U.S. Pat. Nos. 5,234,616, 5,130,045 and 5,130,044 to Mitchell, et al., all of which relate to alkanoyloxyphenyl sulfonates.

[0123] Highly preferred materials of this class of bleaching agents are those that do not cause any significant fabric damage or color change, nor impart any discoloration, such as graying or yellowing, to the matrices into which they are introduced, or to fabrics to which they may be applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

BRIGHTENERS

[0124] Optical brighteners, also referred to as fluorescent whitening agents or FWAs, have long been used to impart whitening to fabrics during the laundering process. These fluorescent materials act by absorbing ultraviolet wavelength of light and emitting visible light, generally in the color blue wavelength ranges. The FWAs settle out or deposit onto fabrics during the wash cycle. These include the stilbene, styrene, and naphthalene derivatives, which upon being impinged by ultraviolet light, emit or fluoresce light in the visible wavelength. While many such compounds are commonly derived from petrochemicals sources, and are as such not preferred, it is envisioned that they could be derived from bio-based sources. It is also envisioned that by being dyes, there are individuals with MCS that may not be able tolerate their presence, and as such natural ingredients such as pigments that possess the ability to fluoresce may be preferable.

[0125] FWAs or brighteners are useful for improving the appearance of fabrics, which have become dingy through repeated soilings and washings. Due to the cationic nature of the composition, it is preferred that the FWAs not be explicitly anionic but rather either nonionic; cationic; amphoteric; or neutralized, ion-paired moieties of anionic FWAs as described in Petrin, et al., U.S. Pat. No. 5,057,236. Preferred anionic FWAs for ion-pairing according to Petrin, et al., '236 are Blankophor BBH, RKH and BHC, from Blankophor GmbH & Co. KG; and Tinopal 5BMX-C, CBS-X and RBS, from BASF Corporation. Fluorescent whiteners most currently used in common laundry compositions generally fall into a category referred to in the art as diaminostilbene disulfonic acid-cyanuric chloride brighteners or DASC-brighteners. These compounds have the following general formula (I):



(I)

[0126] Examples of such DASC fluorescent whiteners include those sold by BASF Corporation under the trade name "Tinopal", which are substituted stilbene 2,2'-disulfonic acid products, e.g., disodium 4,4'-bis-((4-anilino-6-morpholino-1,3,5-triazin-2-yl)amino)stilbene-2,2'-disulfonate (sold as Tinopal AMS); disodium 4,4'-bis-((4-anilino-6-(N-2-hydroxyethyl-N-methyl amino)-1,3,5-triazin-2-yl)amino)stilbene-2,2'-disulfonate (sold as Tinopal 5BM); disodium 4,4'-bis-((4-anilino-6-(bis-(2-hydroxyethyl)amino)-1,3,5-triazin-2-yl)amino)stilbene-2,2'-disulfonate (sold as Tinopal UNPA). Another

example sold by Bayer Corporation is disodium 4,4'-bis-((4-anilino-6-methylamino)-1,3,5-triazin-2-yl)amino)stilbene-2,2'-disulfonate (sold as Phorwite HRS).

[0127] Examples of suitable FWAs can be found in U.K. Pat. Nos. 1,298,577; 2,076,011; 2,026,054; 2,026,566; 1,393,042; and U.S. Pat. No. 3,951,960 to Heath, et al., U.S. Pat. No. 4,298,290 to Barnes, et al., U.S. Pat. No. 5 3,993,659 to Meyer, U.S. Pat. No. 3,980,713 to Matsunaga, et al., and U.S. Pat. No. 3,627,758 to Weber, et al.,. See also, U.S. Pat. No. 4,900,468 to Mitchell, et al., column 5, line 66 to column 6, line 27.

[0128] As stated above, most preferred are cationic, nonionic, and amphoteric FWAs, such as those cited in U.S. Pat. Nos. 4,433,975, 4,432,886, 4,384,121, all to Meyer and U.S. Pat. No. 4,263,431 to Weber, et al. Further examples of suitable FWAs are described in McCutcheon's Vol. 2: Functional Materials, North American Ed., McCutcheon Division, 10 MC Publishing Co., 1995, and Encyclopedia of Chemical Technology, 11th volume, John Wiley & Sons, 1994. Other examples of fluorescent brightening materials suitable for use with the formulations presented herein may be found in U.S. Pat. No. 6,251,303 to Bawendi, et al.; U.S. Pat. No. 6,127,549 to Hao, et al.; U.S. Pat. No. 6,133,215 to Zeiger, et al.; U.S. Pat. No. 6,117,189 to Reinehr, et al.; U.S. Pat. No. 6,120,704 to Martini; and U.S. Pat. No. 6,162,869 to Sharma, et al.

[0129] Highly preferred materials of this class of brighteners are those that do not cause any significant color change, nor 15 impart any discoloration, such as graying or yellowing, to the fabrics to which they are applied, either during treatment followed by drying and/or curing, or after the drying and/or curing step followed by normal exposure to the elements, such as air, moisture or sunlight exposure.

EXAMPLES and STUDIES

[0130] Using the novel assessment protocols defined herein, additional cleaning product formulation guidelines can be 20 developed and promulgated and made available for cleaning product manufacturers. Following the processes disclosed and described herein, a series of products were formulated and tested against commercially available brands in each category, using standard industrial assay techniques. The approach taken to preparing the formulations described herein 25 is believed to be unique in that it links product safety, environmental stewardship and product performance. Customers, therefore, do not have to sacrifice product performance for safety and/or sustainability.

STUDY 1: EFFECT OF PETROLEUM-BASED INGREDIENTS

[0131] As surfactants make up the majority of cleaning product formulations, it is highly important that their contributions 30 be accounted for in individuals that experience MCS. It has been postulated that avoiding petrochemicals may be a first-tier approach in making safer cleaning products. Disappointingly, there are numerous products on the market that claim to be petrochemical-free, and yet are not acceptable to some who experience MCS. As such, a number of raw materials from typical source manufacturers were evaluated in the course of the instant work for their actual bio-renewable carbon. 35 Disappointingly, a number were found to contain hybrid surfactants of significant petrochemical content. The results are shown below in **TABLE 1**.

TABLE 1: Surfactants and Percent Modern Carbon

Surfactant	Percent Modern Carbon (pMC) ^a
Sodium alkylbenzene sulfonate	0%
Lauramine oxide	0%
Sodium lauryl sulfate	0%
C ₁₂₋₁₄ alcohol ethoxylate (7EO)	48%
Cocoamidopropyl betaine	64%
Cocoamidopropyl amine oxide	72%
Cocamide DEA	75%
Sodium coco ether (2EO) sulfate	75%
Cocodimethyl amine oxide	86%
C ₁₀₋₁₆ alkyl polyglucoside	100%
C ₈₋₁₀ alkyl polyglucoside	100%
Sodium coco sulfate	100%
Sodium octyl sulfate (from bio-basis)	100%

55 Note to **TABLE 1**:

^a Measure of the percent of modern or bio-based carbon in an ingredient or composition, as estimated from evaluation of feedstocks of component carbons, or determined by ASTM D6866-05.

[0132] As may be readily observed, many surfactants possess a significant portion, that is 25% or more by weight, of petrochemical contribution. In other words, despite being positioned as "natural" surfactants, their Biorenewable Carbon Index (and thus their biorenewable carbon content) is less than 80% by weight. As such, it has been found that surfactants that are preferred for use with the compositions described herein are those having a Biorenewable Carbon Index of at least 80%, that is, with a BCI of $\geq 80\%$.

STUDY 2: EFFECT OF TRACE CONTAMINANTS

[0133] In a second study, it was surprisingly found that certain surfactants that had a very high Biorenewable Carbon Index, even as high as 100%, could have adverse effects on certain individuals with MCS. Without being bound by theory, it is believed that this phenomenon is due to the non-exact nature of the BCI or RCI measurement ($\pm 3\%$ by weight) vis-à-vis the low amounts of contaminants, perhaps much less than 1% by weight, that are present in certain ingredients. To test this hypothesis, a number of chemically sensitive individuals assessed four types of alkyl polyglucosides, or APGs, from three manufacturers, all of which have an apparent BCI/RCI of 100%. Each of the APG candidates was rated on a 3-point index: Acceptable, Marginal, and Unacceptable. The Marginal and Unacceptable candidates were then analyzed for the presence of trace contaminants. Surprisingly, those candidates all contained detectable amounts of phenyl derivatives (toluene, acetophenone), apparently owing to the nature of the catalyst used during manufacture and the fact that it was perhaps not stripped out prior to distribution. Candidates that were determined to be Acceptable did not have such phenyl residue. The results are shown in **TABLE 2** below.

TABLE 2: Acceptability of Candidate Alkyl Polyglucosides

Surfactant	Percent Modern Carbon (pMC) ^a	Acceptability Rating
Commercial APG Product A	100%	Unacceptable
Commercial APG Product B	100%	Unacceptable
Commercial APG Product C	100%	Marginal
Commercial APG Product D	100%	Acceptable

Note to **TABLE 2**.

^a Measure of the percent of modern or bio-based carbon in an ingredient or composition, as estimated from evaluation of feedstocks of component carbons, or determined by ASTM D6866-05.

STUDY 3: EFFECT OF CARBON CHAIN LENGTH

[0134] In the manufacture of surfactants, it can be assumed that a small amount, albeit a significant one, of feedstock remains unreacted. This amount can often be less than 1% by weight, but to individuals who are sensitive to such species, the impact can be significant. Often, this can result in an olfactory response as lower-chain alcohols are quite odiferous, but to individuals with MCS, the impact is more significant. Without being bound by theory, it is believed that chemicals—especially surfactant residues—that have a chain length of eight carbon atoms or less may react with proteins to form complexes that trigger an immunogenic response. Chemicals that have a carbon chain length greater than eight carbon atoms may be insufficiently reactive with proteins to form complexes. Alternatively, it is postulated that any conjugates having greater than eight-carbon atom chain lengths that may form, are present in concentrations that are lower than a threshold level needed to trigger an immunogenic response. As such, a number of raw materials were evaluated by a number of chemically sensitive individuals for acceptability using the same scale as above. The results are presented in **TABLE 3** below.

TABLE 3: Effect of Carbon Chain Length of Surfactant Feedstock

Surfactant	Percent Modern Carbon (pMC) ^a	Average Carbon Chain Length	Acceptability Rating
Sodium octyl sulfate (from bio-basis)	100%	8	Unacceptable
C ₈₋₁₀ alkyl polyglucoside	100%	9	Unacceptable
C ₁₀₋₁₆ alkyl polyglucoside	100%	13	Acceptable-Unacceptable*

(continued)

Surfactant	Percent Modern Carbon (pMC) ^a	Average Carbon Chain Length	Acceptability Rating
Sodium coco sulfate	100%	12	Acceptable

Notes to **TABLE 3**

^a Measure of the percent of modern or bio-based carbon in an ingredient or composition, as estimated from evaluation of feedstocks of component carbons, or determined by ASTM D6866-05.

* Response received was dependent upon the source of alkyl polyglucoside, which in turn was found to depend on the level of trace contaminants, as discussed above.

STUDY 4: VOLATILE ORGANICS

[0135] As mentioned above, in an analysis of 37 commercial products, Steinemann (2015) found emissions of 156 different VOCs, with an average of 15 VOCs per product. Of these 156 VOCs, 42 VOCs were classified as toxic or hazardous under U.S. federal laws, and each product emitted at least one of these chemicals. Despite inferences, the emissions of hazardous air pollutants (HAPs) from so-called green fragranced products were not significantly different from non-green labeled fragranced products.

[0136] Without being bound to theory, the inventors believe that minimizing or eliminating sources of VOC can significantly reduce contaminants that may introduce hazardous air pollutants into consumer cleaning compositions. For example, the vapor pressure of ethanol at room temperature is about 45 mm Hg at 20°C, while that of 1,3-propanediol is 0.08 mm Hg, and that of glycerine is about 0.00018 mm Hg. Substitution of ethanol with other low-volatility solvents can therefore not only lower the overall vapor pressure of a product, and but may reduce the likelihood of untoward effects on persons that suffer from MCS.

EXAMPLE 1: Effect of Carbon Chain Length of Surfactant Feedstock

[0137] A laundry detergent was formulated in accordance with the guidelines presented above. Accordingly, 15.0 parts of alkyl polyglucoside (Triton CG-600, 50% active from Dow Chemical Company) were added to 63.1 parts deionized water with mixing, followed by 13.0 parts of sodium coco sulfate (Stepanol WA-Extra, 29% active from Stepan Company), 2.0 parts glycerine (Pricerene 9091 from Croda), 2.0 parts boric acid, 2.0 parts oleic acid (Acme-Hardesty Co), 1.0 part sodium gluconate, 1 part sodium hydroxide, 0.1 part protease (Novozymes), 0.1 part calcium chloride, 0.1 part sodium chloride, 0.05 parts amylase (Novozymes), and 0.05 parts preservative (Neolone M10, 10% active).

[0138] A portion of the resulting formulations, designated Sample F in **TABLE 4** below, was then submitted for evaluation and analysis versus a commercially available, safety-positioned, unfragranced liquid detergent product, labeled Sample E in **TABLE 4** below. The samples were analyzed via two different methods: U.S. EPA Compendium Method TO-15, "Determination Of Volatile Organic Compounds (VOCs) In Air Collected In Specially-Prepared Canisters And Analyzed By Gas Chromatography/Mass Spectrometry (GC/MS)," EPA, 1999, and U.S. EPA Method TO-11A, "Determination of Formaldehyde in Ambient Air Using Adsorbent Cartridge Followed by High Performance Liquid Chromatography (HPLC)," EPA, 1999. The second of the two analyses is specific for aldehydes such as formaldehyde and acetaldehyde.

[0139] **TABLE 4** summarizes findings for the two laundry detergent samples. Note that reported analyte values were above a threshold of 1000 µg/m³, thereby ensuring with some confidence that they are emitted from the products:

TABLE 4: HEADSPACE Analysis of Samples by EPA Method TO-15

Compound	CAS Number	Sample E Liquid Laundry Detergent (>1000 µg/m ³) ^{a,b}	Sample F Liquid Laundry Detergent (> 1000 µg/m ³) ^{a,c}
3,5-Dimethyl-1-hexene	7423-69-0	Y	N
Cyclotetradecane	295-17-0	Y	N
2-Phenoxyethanol	122-99-6	Y	N
Methyl octanoate	111-11-5	Y	N
Trichloromethane (chloroform)*	67-66-3	Y	N
11-Bromoundecanoic acid	2834-05-1	Y	N
Methyl undecanoate	1731-86-8	Y	N
2-Propenoic acid, 1,7,7-trimethylbicyclo[2.2.1]-hept-2-yl ester	5888-33-5	Y	Y

(continued)

Compound	CAS Number	Sample E Liquid Laundry Detergent (>1000 $\mu\text{g}/\text{m}^3$) ^{a,b}	Sample F Liquid Laundry Detergent (> 1000 $\mu\text{g}/\text{m}^3$) ^{a,c}
Isobutyl nonyl oxalate	Unknown	Y	Y
1-Undecanol	112-42-5	N	Y
2-Methyl-cyclopentanol	24070-77-7	N	Y
Pentadecane	629-62-9	N	Y

Notes to TABLE 4:

* Known hazardous substance, cancer risk; **bold** = classified as toxic or hazardous under U.S. federal law.

^aAbbreviations are used to indicate presence or absence of material in head space: Y = Yes; N = No.

^b Sample E: Commercially available liquid laundry detergent

^c Sample F: Laundry detergent formulated according to the instant disclosure, described in EXAMPLE 1, above.

[0140] The analytes were compared against eight Federal registers of potentially hazardous VOCs, per the study by Steinemann (2015). One analyte found in the commercial product, chloroform, is present on all registries; this is in agreement with the study Steinemann (2015), wherein every product analyzed had at least one such potentially hazardous VOC. Remarkably, chloroform was found to be absent in the liquid laundry detergent product prepared according to the instant specification. In fact, the product prepared according to the methods described herein did not have one chemical that appears on any of the registries of hazardous chemicals present at a level above 1000 $\mu\text{g}/\text{m}^3$. The results are summarized in Table 5 below.

25

30

35

40

45

50

55

TABLE 5: Headspace Analysis of Samples via EPA Method TO-15

	Compound	CAS Number	CAA-TFS ^a	CAA-HAP ^b	CERCLA ^c	CWA ^d	EPCRA ^e	FIFRA ^f	OSHA ^g	RCRA ^h
10	3,5-Dimethyl-1-hexene	7423-69-0	-	-	-	-	-	-	-	-
15	Cyclotetradecane	295-17-0	-	-	-	-	-	-	-	-
20	2-Phenoxyethanol	122-99-6	--	--	--	--	--	--	--	--
25	Methyl octanoate	111-11-5	--	--	--	--	--	--	--	--
30	Trichloromethane (chloroform) *	67-66-3	✓	✓	✓	✓	✓	✓	✓	✓
35	11-Bromoundecanoic acid	2834-05-1	--	--	--	--	--	--	--	--
40	Methyl undecanoate	1731-86-8	--	--	--	--	--	--	--	--
45	2-Propenoic acid, 1,7,7-trimethyl-bicyclo[2.2.1]hept-2-yl ester	5888-33-5	--	--	--	--	--	--	--	--
50	Isobutyl nonyl oxalate	Unknown	--	--	--	--	--	--	--	--
55	1-Undecanol	112-42-5	--	--	--	--	--	--	--	--
60	2-Methyl-cyclopentanol	24070-77-7	--	--	--	--	--	--	--	--
65	Pentadecane	629-62-9	--	--	--	--	--	--	--	--

Notes to TABLE 5

* Known hazardous substance, cancer risk; **bold** = classified as toxic or hazardous under U.S. federal law.

CAA-TFS^a: Clean Air Act—Toxic and Flammable Substances for Accidental

40

45

50

55

Release Prevention

CAA-HAP ^b :	Clean Air Act—Hazardous Air Pollutant
CERCLA ^c :	Comprehensive Environmental Response, Compensation, and Liability Act—Hazardous Substance
CWA ^d :	Clean Water Act—Priority Pollutant
EPCRA ^e :	The Emergency Planning & Community Right to Know Act—Toxic Release Inventory Chemical
FIFRA ^f :	Federal Insecticide, Fungicide, and Rodenticide Act—Registered Pesticide
OSHA ^g :	Occupational Safety and Health Act—Air Contaminants
RCRA ^h :	Resource Conservation and Recovery Act—Hazardous Constituents

[0141] The two specimens, Samples E and F were then analyzed via U.S. EPA Method TO-11A, which is specific for

aldehydes such as formaldehyde and acetaldehyde. Sample F, a composition prepared in accordance with the methods described in the instant specification, had significantly less acetaldehyde than commercial product Sample B, and is absent of formaldehyde down to the detection limit. Results are summarized below in **TABLE 6**.

5 **TABLE 6:** Headspace Analysis of Samples via EPA Method TO-11A

	<u>Sample</u>	Formaldehyde		Acetaldehyde	
		parts per billion (by volume)	µg/m ³	parts per billion (by volume)	µg/m ³
10	Commercial Liquid Detergent (Sample E) ^a	18.8	23.09	7.45	13.42
	Formulated Liquid Detergent (Sample F) ^b	(below detection limit)	(below detection limit)	3.07	5.53

15 **Notes to TABLE 6**

a Sample E: Commercially available liquid laundry detergent

b Sample F: Laundry detergent formulated according to the instant disclosure, described in EXAMPLE 1, above.

20 **STUDY 5: REPRESENTATIVE FORMULATIONS**

25 **[0142]** Based on the assessment criteria described herein, several cleaning formulas were generated in accordance with the methods described herein and found to be highly effective at cleaning. And yet, when evaluated by a panel of five individuals that have MCS, the formulas were found to be totally acceptable for use without deleterious physical effects. Representative formulas prepared and tested according to the instant specification are listed in **TABLE 7** and results summarized in **TABLE 8** below.

30 **TABLE 7: Representative Cleaning Formulas**

	Ingredient	Sample F Liquid laundry detergent	Sample G Liquid dishwashing detergent	Sample H All-Purpose cleaner	Sample I General bathroom cleaner
30	Sodium coco sulfate, 29% active	13.0%	45.0%	--	--
	Alkyl polyglucoside, 50% active	15.0%	9.0%	3.0%	5.0%
	Cocoamine oxide	--	12.0%	--	--
35	Glycerine	2.5%	3.5%	1.5%	--
	Boric acid	2.0%	--	--	--
	Citric acid	--	0.25%	--	4.0%
	Oleic acid	2.0%	--	--	--
40	Sodium gluconate	1.0%	--	--	--
	Sodium hydroxide	1.0%	--	--	--
	Potassium citrate	--	--	0.5%	--
	Protease	0.1%	--	--	--
45	Calcium chloride	0.1%	--	--	--
	Sodium chloride	0.1%	--	--	--
	Amylase	0.05%	--	--	--
	Preservative, 10% active	0.05%	0.05%	0.05%	0.05%
	Water	q.s.	q.s.	q.s.	q.s.
50	Percent Modern Carbon (pMC) ^a	99.97%	97.96%	99.86%	99.92%

55 Note to **TABLE 7**:

^a Measure of the percent of modern or bio-based carbon in an ingredient or composition, as estimated from evaluation of feedstocks of component carbons, or determined by ASTM D6866-05.

TABLE 8: Use Results of Representative Cleaning Formulas

Product	Sample F Liquid laundry detergent	Sample G Liquid dishwashing detergent	Sample H All-Purpose cleaner	Sample I General bathroom cleaner
5 Acceptable Performance (versus commercial products)	Yes	Yes	Yes	Yes
10 Acceptable Use (without ill effects)?	Yes	Yes	Yes	Yes
15	<p>[0143] A liquid laundry detergent that may be prepared according to the information presented herein and be well suited for use by chemically-sensitive individuals, contains: 1) 30% sodium coco sulfate, 29% active; 2) 15% alkyl polyglucoside, 50% active; 3) 2.5% glycerine; 4) 2.0% boric acid; 5) 2.0% oleic acid; 6) 1.0% sodium gluconate; 7) 1.0% sodium hydroxide; 8) 0.1% protease; 9) 0.1% calcium chloride; 10) 0.1% sodium chloride; 11) 0.05% amylase; 12) 0.05% preservative, 10% active; and the balance water, where all percents are understood to refer to weight percent. Furthermore, a liquid laundry detergent that may be prepared according to the methods presented herein and consistent with the above composition may be found to have a percent modern carbon (pMC) of approximately 99.97%.</p>			
20	<p>[0144] In another aspect, a dishwashing detergent that may be prepared according to the information presented herein and be well suited for use especially by chemically-sensitized individuals, contains: 1) 45.0% sodium coco sulfate, 29% active; 2) 9.0% alkyl polyglucoside, 50% active; 3) 12% cocamine oxide, 30%; 4) 3.5% glycerine; 5) 0.25% citric acid; 6) 0.05% preservative, 10% active; and the balance water, where all percents are understood to refer to weight percent. Furthermore, a dishwashing detergent that may be prepared according to the methods presented herein and consistent with the above composition may be found to have a percent modern carbon (pMC) of approximately 97.96%.</p>			
25	<p>[0145] In yet another aspect, an all-purpose cleaner that may be prepared according to the information presented herein and be found to be well suited for use especially by chemically-sensitized individuals, contains: 1) 3.0% alkyl polyglucoside, 50% active; 2) 1.5% glycerine; 3) 0.5% potassium citrate, 4) 0.05% preservative, 10% active; and the balance water, where all percents are understood to refer to weight percent. Furthermore, an all-purpose cleaner that may be prepared according to the methods presented herein and consistent with the above composition may be found to have a percent modern carbon (pMC) of approximately 99.86%.</p>			
30	<p>[0146] In still another aspect, a general bathroom cleaner that may be prepared according to the information presented herein and be found to be well suited for use especially by chemically-sensitized individuals, contains: 1) 5.0% alkyl polyglucoside, 50% active; 2) 4% citric acid; 3) 0.05% preservative, 10% active; and the balance water, where all percents are understood to refer to weight percent. Furthermore, a general bathroom cleaner that may be prepared according to the methods presented herein and consistent with the above composition may be found to have a percent modern carbon (pMC) of approximately 99.92%.</p>			
35	<p>[0147] It is to be noted that the foregoing samples and examples demonstrate the manner in which novel formulations and methods disclosed herein can be used to provide cleaning products that exhibit enhanced hypoallergenicity and can be generated from sustainable sources without sacrificing cleaning efficacy. The foregoing samples and examples demonstrate the manner in which the compositions and methods described herein provide screening for many recognized deleterious health effects without effectively sacrificing cleaning efficacy for the sake of sustainability of materials.</p>			
40	<p>[0148] The instant disclosure presents information that has been described in detail herein with reference to specific embodiments, methods and examples. However, these specific embodiments should not be construed as narrowing the scope of the formulations and methods described herein, but rather construed as illustrative examples. It is to be further understood that obvious embodiments, modifications and equivalents thereof are anticipated and are considered to be within the scope of the newly presented formulations and methods. The subject matter of the instant disclosure is further illustrated and described in the claims that follow.</p>			
45				

Claims

50

1. A cleaning composition suitable for use by chemically-sensitive individuals, comprising:

a. at least one surfactant chosen from the group comprising:

55

- 1) 0.05-15% by weight of an anionic surfactant;
- 2) 0.05-30% by weight of a nonionic surfactant; and
- 3) 0.05-30% by weight of at least one of an amphoteric and a zwitterionic surfactant;

- b. 0-10% by weight of at least one of a builder and a pH adjuster; and
- c. 0-95% by weight of at least one solvent;

wherein:

- 5 1) each of carbon-based items a. through c. has a pMC of at least 80%;
- 2) headspace analysis of the composition reveals the absence of phenyl compounds or their derivatives;
- 3) less than 5% by weight of each item has a vapor pressure that is above 0.1 mm Hg at 20°C;
- 4) the composition contains less than 1% by weight of a fragrance material; and
- 10 5) headspace analysis of the cleaning composition reveals the presence of less than 1000 µg/m³ of toxic or hazardous VOCs;

wherein the anionic surfactant is selected from the group consisting of C₁₀-C₁₄ alkyl sulfates and ethoxysulfates, C₁₀-C₁₈ alkyl sulfonates, C₁₀-C₁₄ linear or branched alkyl benzene sulfonates, C₁₀-C₁₅ alkyl ethoxycarboxylates and combinations of any of the foregoing;

15 wherein the nonionic surfactant is selected from the group consisting of alkylpolysaccharides and addition products of fatty alcohols, fatty acids, and fatty amines coupled with alkoxylating agents selected from the group comprising ethylene oxide (EO), propylene oxide (PO), isopropylene oxide (IPO), butylene oxide (BO), and combinations thereof;

20 wherein the amphoteric surfactant is selected from the group consisting of amine oxides having the formula (R₁)(R₂)(R₃)NO wherein R₁ is a hydrocarbon chain having from 1 to 30 carbon atoms, R₂ and R₃ are independently substituted or unsubstituted, C₁-C₄ linear or branched hydrocarbon chains, and the zwitterionic surfactant is selected from the group consisting of R₁-N'(R₂)(R₃)R₄X, wherein R₁ is a C₁₀-30 hydrophobic group; R₂ and R₃ are each C₁-C₄ alkyl, hydroxyalkyl or other substituted alkyl group which can also be joined to form ring structures with the N; R₄ is a C₁-10 moiety joining the cationic nitrogen atom to the hydrophilic group and is typically an alkylene, hydroxy alkylene, or polyalkoxy group, and X is a carboxylate group or a sulfonate group;

25 wherein the at least one builder or pH adjuster is selected from the group consisting of calcium and magnesium sequestrants; inorganic or organic monobasic or dibasic organic acids; salts of monocarboxylic acids, salts of dicarboxylic acids, salts of citric acid and other suitable organic acids with water soluble conjugate bases; organic bases such as alkanolamines, including methanol, ethanol and propanol amines, including dimethanol-, diethanol- and dipropanolamines, and including trimethanol-, triethanol- and tripropanolamines; and

30 wherein the at least one solvent is selected from the group consisting of organic solvents comprising C₁-C₆ alkanols, C₁-C₆ diols, C₁-C₁₀ alkyl ethers of alkylene glycols, C₃-C₂₄ alkylene glycol ethers, polyalkylene glycols, short chain carboxylic acids, short chain esters, isoparaffinic hydrocarbons, mineral spirits, alkyl aromatics, terpenes, terpene derivatives, terpenoids, terpenoid derivatives, formaldehyde, and pyrrolidones; water insoluble solvents; and water-soluble solvents.

2. The cleaning composition of claim 1, wherein:

- 40 a. the anionic surfactant is further selected from the group comprising sodium alkylbenzene sulfonate, sodium lauryl sulfate, sodium coco ether sulfate, sodium coco sulfate, and sodium octyl sulfate;
- b. the nonionic surfactant is further selected from the group comprising C₁₀-14 alcohol ethoxylates, C₁₀-16 alkyl polyglucosides, and C₈-10 alkyl polyglucosides;
- c. the at least one of an amphoteric and a zwitterionic surfactant is further selected from the group comprising C₁₂-C₁₆ amine oxides, cocamine oxide, lauramine oxide, cocamidopropyl betaine, cocamidopropyl amine oxide, cocamide DEA, alkylamphoglycinate, and alkyl iminopropionates;
- d. the at least one builder or pH adjuster is further selected from the group comprising acids selected from among acetic, maleic acid, glycolic acid, methyl sulfonic, hydrochloric, sulfuric, phosphoric, citric, succinic, boric and sodium gluconate; inorganic bases selected from among alkali metal and alkaline earth metal salts of hydroxides, borates, sulfonates, phosphates, phosphonates and silicates; organic bases selected from among alkanolamines including methanol, ethanol and propanol amines, dimethanol-, diethanol- and dipropanolamines, and trimethanol-, triethanol- and tripropanolamines; and salts of organic acids, including alkali metal and alkaline earth metal salts of citric and gluconic acid;
- e. the at least one solvent is further selected from the group comprising methanol, ethanol, n-propanol, isopropanol, butanol, pentanol, and hexanol, and isomers thereof; methylene, ethylene, propylene and butylene glycols; ethylene glycol monopropyl ether, ethylene glycol monobutyl ether, ethylene glycol monohexyl ether, diethylene glycol monopropyl ether, diethylene glycol monobutyl ether, diethylene glycol monohexyl ether, propylene glycol methyl ether, propylene glycol ethyl ether, propylene glycol n-propyl ether, propylene glycol monobutyl ether, propylene glycol t-butyl ether, di- or tri-polypropylene glycol methyl or ethyl or propyl or butyl

ether, acetate and propionate esters of glycol ethers; glycol acetate, and cyclic or linear volatile methylsiloxanes; and isoparaffinic hydrocarbons, mineral spirits, alkyl aromatics, terpenoids, terpenoid derivatives, terpenes, and terpene derivatives.

5 3. The cleaning composition of Claim 2, wherein:

- a. the anionic surfactant is selected from the group comprising sodium alkylbenzene sulfonate, sodium lauryl sulfate, sodium coco ether (2EO) sulfate, sodium coco sulfate, and sodium octyl sulfate;
- 10 b. the nonionic surfactant is selected from the group comprising C₁₂₋₁₄ alcohol ethoxylate, C₁₀₋₁₆ alkyl polyglucosides and C₈₋₁₀ alkyl polyglucosides;
- c. the amphoteric and zwitterionic surfactants are selected from the group comprising lauramine oxide, cocoamidopropyl betaine, cocoamidopropylamine oxide, cocoamide DEA, and cocodimethyl amine oxide;
- 15 d. the at least one builder or pH adjuster is selected from the group comprising boric acid, sodium gluconate, sodium hydroxide, and citric acid; and
- e. the at least one solvent is selected from the group comprising glycerine and water.

4. The cleaning composition of any of Claims 1-3, wherein each of the ingredients comprising the composition contains less than 1% by weight of trace contaminants, wherein the trace contaminants are selected from the group comprising phenyl derivatives and lower-chain alcohols.

20 5. The cleaning composition of Claim 4, wherein:

- a. the phenyl derivatives further comprise toluene and acetophenone; and
- b. the lower-chain alcohols further comprise alcohols having a chain length of eight carbon atoms or less.

25 6. The cleaning composition of any of Claims 1-3, further wherein the anionic surfactant contains no detectable level of 1,4-dioxane.

7. A method for producing the cleaning composition of Claim 1, comprising the steps of:

- 30 a. assessing the bio-basis of any of the carbon-based components a. through c. for use in the cleaning composition by determining the percent modern carbon (pMC) for each component, wherein a component having a pMC of at least 80% is deemed acceptable;
- b. formulating a cleaning composition using the acceptable component or components from step a; and
- 35 c. performing a headspace analysis for the cleaning composition produced in step b.

8. The method of Claim 7, further wherein each of the ingredients comprising the composition contains less than 1% by weight of trace contaminants, wherein the trace contaminants are selected from the group comprising phenyl derivatives and lower-chain alcohols, and a headspace analysis shows a level of less than 1000 µg/m³ of VOCs comprising the group of 1,4-dioxane, chloroform, formaldehyde and acetaldehyde.

40 9. The method of Claim 8, wherein:

- 45 a. the phenyl derivatives further comprise toluene and acetophenone;
- b. the lower-chain alcohols further comprise alcohols having a chain length of eight carbon atoms or less; and
- c. the anionic surfactant contains no detectable level of 1,4-dioxane.

10. The method of Claim 9, wherein:

- 50 a. the anionic surfactant is selected from the group comprising sodium alkylbenzene sulfonate, sodium lauryl sulfate, sodium coco ether (2EO) sulfate, sodium coco sulfate and sodium octyl sulfate;
- b. the nonionic surfactant is selected from the group comprising C₁₂₋₁₄ alcohol ethoxylate, C₁₀₋₁₆ alkyl polyglucosides and C₈₋₁₀ alkyl polyglucosides;
- c. the amphoteric and zwitterionic surfactants are selected from the group comprising lauramine oxide, cocoamidopropyl betaine, cocoamidopropylamine oxide, cocoamide DEA, and cocodimethyl amine oxide;
- 55 d. the at least one builder or pH adjuster is selected from the group comprising boric acid, sodium gluconate, sodium hydroxide, citric acid; and
- e. the at least one solvent is selected from the group comprising glycerine and water.

11. The cleaning composition of Claim 3, wherein:

- a. the anionic surfactant is sodium coco sulfate;
- b. the nonionic surfactant is alkyl polyglucoside;
- 5 c. the amphoteric or zwitterionic surfactant is cocamine oxide;
- d. the builder or pH adjuster is selected from the group comprising boric acid, citric acid, sodium gluconate, sodium hydroxide, and potassium citrate; and
- e. the solvent is glycerine.

10 12. The method of Claim 10, wherein:

- a. the anionic surfactant is sodium coco sulfate;
- b. the nonionic surfactant is alkyl polyglucoside;
- 15 c. the amphoteric or zwitterionic surfactant is cocamine oxide;
- d. the builder or pH adjuster is selected from the group comprising boric acid, citric acid, sodium gluconate, sodium hydroxide, and potassium citrate; and
- e. the solvent is glycerine.

20 13. The cleaning composition of Claim 6, wherein the composition is deemed suitable for use by at least one individual who manifests multiple chemical sensitivities.

14. The method of Claim 7, further comprising the step of:

- d. evaluating the suitability of the cleaning composition for use by at least one individual who manifests multiple chemical sensitivities.

25

Patentansprüche

30 1. Reinigungszusammensetzung, die für die Verwendung durch chemikaliensensitive Personen geeignet ist, umfassend:

- a. mindestens ein Tensid, ausgewählt aus der Gruppe umfassend:

- 35 1) 0,05-15 Gew.-% eines anionischen Tensids;
- 2) 0,05-30 Gew.-% eines nichtionischen Tensids; und
- 3) 0,05-30 Gew.-% mindestens eines von einem amphoteren und einem zwitterionischen Tensid;

- b. 0-10 Gew.-% mindestens eines Builders und eines Mittels zur Einstellung des pH-Werts; und
- c. 0-95 Gew.-% mindestens eines Lösungsmittels;

40

wobei:

- 45 1) jeder der kohlenstoffbasierten Posten a. bis c. einen pMC-Wert von mindestens 80 % aufweist;
- 2) die Kopfraumanalyse der Zusammensetzung die Abwesenheit von Phenylverbindungen oder Derivaten davon zeigt;
- 3) weniger als 5 Gew.-% jedes Postens einen Dampfdruck von mehr als 0,1 mm Hg bei 20 °C aufweist;
- 4) die Zusammensetzung weniger als 1 Gew.-% eines Duftstoffs enthält; und
- 5) die Kopfraumanalyse der Reinigungszusammensetzung das Vorliegen von weniger als 1000 µg/m³ toxischen oder gefährlichen VOCs zeigt;

50 wobei das anionische Tensid aus der Gruppe bestehend aus C₁₀-C₁₄-Alkylsulfaten und -Ethoxysulfaten, C₁₀-C₁₈-Alkylsulfonaten, linearen oder verzweigten C₁₀-C₁₄-Alkylbenzolsulfonaten, C₁₀-C₁₅-Alkylethoxycarboxylaten und Kombinationen von beliebigen der obigen Substanzen ausgewählt ist; wobei das nichtionische Tensid aus der Gruppe bestehend aus Alkylpolysacchariden und Additionsprodukten von Fettalkoholen, Fettsäuren und Fettaminen, die mit Alkoxylierungsmitteln aus der Gruppe umfassend Ethylenoxid (EO), Propylenoxid (PO), Isopropylenglycid (IPO), Butylenoxid (BO) und Kombinationen davon gekuppelt sind, ausgewählt ist; wobei das amphotere Tensid aus der Gruppe bestehend aus Aminoxiden mit der Formel (R₁)(R₂)(R₃)NO ausgewählt sind, wobei R₁ für eine Kohlenwasserstoffkette mit 1 bis 30 Kohlenstoffatomen steht, R₂ und R₃ unabhängig für substituierte oder unsubstituierte lineare oder verzweigte C₁-C₄-Kohlenwasserstoffketten ste-

hen, und das zwitterionische Tensid aus der Gruppe bestehend aus $R_1-N'(R_2)(R_3)R_4X$ ausgewählt ist, wobei R_1 für eine hydrophobe C_{10-30} -Gruppe steht; R_2 und R_3 jeweils für C_1-C_4 -Alkyl, Hydroxyalkyl oder eine andere substituierte Alkylgruppe, die auch mit dem N zu Ringstrukturen verknüpft sein kann, stehen; R_4 für eine C_{1-10} -Gruppierung steht, die das kationische Stickstoffatom mit der hydrophilen Gruppe verknüpft und typischerweise eine Alkylen-, Hydroxyalkylen- oder Polyalkoxygruppe ist, und X für eine Carboxylatgruppe oder eine Sulfonatgruppe steht; wobei der mindestens eine Builder oder das mindestens eine Mittel zur Einstellung des pH-Werts aus der Gruppe bestehend aus Calcium- und Magnesium-Sequestermitteln; anorganischen oder organischen einbasigen oder zweibasigen organischen Säuren; Salzen von Monocarbonsäuren, Salzen von Dicarbonsäuren, Salzen von Citronensäure und anderen geeigneten organischen Säuren mit wasserlöslichen konjugierten Basen; organischen Basen wie Alkanolaminen einschließlich Methanol-, Ethanol- und Propanolaminen einschließlich Dimethanol-, Diethanol- und Dipropanolaminen und einschließlich Trimethanol-, Triethanol- und Tripropanolaminen ausgewählt ist; und wobei das mindestens eine Lösungsmittel aus der Gruppe bestehend aus organischen Lösungsmitteln, umfassend C_1-C_6 -Alkanole, C_1-C_6 -Diole, C_1-C_6 -Triole, C_1-C_{10} -Alkylether von Alkylenglykolen, C_3-C_{24} -Alkylenglykolether, Polyalkylenglykole, kurzkettige Carbonsäuren, kurzkettige Ester, isoparaffinische Kohlenwasserstoffe, Testbenzin, Alkylaromaten, Terpene, Terpenderivate, Terpenoide, Terpenoiderivate, Formaldehyd und Pyrrolidone; wasserunlöslichen Lösungsmitteln und wasserlöslichen Lösungsmitteln ausgewählt ist.

2. Reinigungszusammensetzung nach Anspruch 1, wobei:

- a. das anionische Tensid ferner aus der Gruppe umfassend Natriumalkylbenzolsulfonat, Natriumlaurylsulfat, Natriumcocoethersulfat, Natriumcocosulfat und Natriumoctylsulfat ausgewählt ist;
- b. das nichtionische Tensid ferner aus der Gruppe umfassend $C_{10-C_{14}}$ -Alkoholethoxylate, $C_{10-C_{16}}$ -Alkylpolyglucoside und C_8-C_{10} -Alkylpolyglucoside ausgewählt ist;
- c. das mindestens eine von einem amphoteren und einem zwitterionischen Tensid ferner aus der Gruppe umfassend $C_{12-C_{16}}$ -Aminoxide, Cocoaminoxid, Lauraminoxid, Cocoamidopropylbetain, Cocoamidopropylaminoxid, Cocoamid-DEA, Alkylamphoglycinate und Alkyliminopropionate ausgewählt ist;
- d. der mindestens eine Builder oder das mindestens eine Mittel zur Einstellung des pH-Werts ferner aus der Gruppe umfassend Säuren, ausgewählt aus Essigsäure, Maleinsäure, Glykolsäure, Methylsulfonsäure, Chlorwasserstoffsäure, Schwefelsäure, Phosphorsäure, Citronensäure, Bernsteinsäure, Borsäure und Natriumgluconat; anorganischen Basen, ausgewählt aus Alkalimetall- und Erdalkalimetallsalzen von Hydroxiden, Boraten, Sulfonaten, Phosphaten, Phosphonaten und Silikaten; organischen Basen, ausgewählt aus Alkanolaminen einschließlich Methanol-, Ethanol- und Propanolaminen, Dimethanol-, Diethanol- und Dipropanolaminen und Trimethanol-, Triethanol- und Tripropanolaminen; und Salzen von organischen Säuren einschließlich Alkalimetall- und Erdalkalimetallsalzen von Citronensäure und Gluconsäure ausgewählt ist;
- e. das mindestens eine Lösungsmittel ferner aus der Gruppe umfassend Methanol, Ethanol, n-Propanol, Isopropanol, Butanol, Pentanol und Hexanol und Isomere davon; Methylen-, Ethylen-, Propylen- und Butylenglykole; Ethylenglykolmonopropylether, Ethylenglykolmonobutylether, Ethylenglykolmonohexylether, Diethylenglykolmonopropylether, Diethylenglykolmonobutylether, Diethylenglykolmonohexylether, Propylenglykolmethylether, Propylenglykolethylether, Propylenglykol-n-propylether, Propylenglykolmonobutylether, Propylenglykol-t-butylether, Di- oder Tripropylethylmethylether oder -ethylether oder -propylether oder -butylether, Acetat- und Propionatester von Glykolethern; Glykolacetat und cyclische oder lineare flüchtige Methylsiloxane und isoparaffinische Kohlenwasserstoffe, Testbenzin, Alkylaromate, Terpenoide, Terpenoiderivate, Terpene und Terpenderivate ausgewählt ist.

3. Reinigungszusammensetzung nach Anspruch 2, wobei:

- a. das anionische Tensid aus der Gruppe umfassend Natriumalkylbenzolsulfonat, Natriumlaurylsulfat, Natrium-cocoether(2EO)sulfat, Natriumcocosulfat und Natriumoctylsulfat ausgewählt ist;
- b. das nichtionische Tensid aus der Gruppe umfassend $C_{12-C_{14}}$ -Alkoholethoxylat, $C_{10-C_{16}}$ -Alkylpolyglucoside und C_8-C_{10} -Alkylpolyglucoside ausgewählt ist;
- c. das amphotere und zwitterionische Tensid aus der Gruppe umfassend Lauraminoxid, Cocoamidopropylbetain, Cocoamidopropylaminoxid, Cocoamid-DEA und Cocodimethylaminoxid ausgewählt sind;
- d. der mindestens eine Builder oder das mindestens eine Mittel zur Einstellung des pH-Werts aus der Gruppe umfassend Borsäure, Natriumgluconat, Natriumhydroxid und Citronensäure ausgewählt ist; und
- e. das mindestens eine Lösungsmittel aus der Gruppe umfassend Glycerin und Wasser ausgewählt ist.

4. Reinigungszusammensetzung nach einem der Ansprüche 1-3, wobei jeder der Bestandteile, die die Zusammen-

setzung ausmachen, weniger als 1 Gew.-% Spurenverunreinigungen umfasst, wobei die Spurenverunreinigungen aus der Gruppe bestehend aus Phenylderivaten und kürzerkettigen Alkoholen ausgewählt sind.

5. Reinigungszusammensetzung nach Anspruch 4, wobei:

5 a. die Phenylivate ferner Toluol und Acetophenon umfassen; und
b. die kürzerkettigen Alkohole ferner Alkohole mit einer Kettenlänge von acht Kohlenstoffatomen oder weniger umfassen.

10 6. Reinigungszusammensetzung nach einem der Ansprüche 1-3, ferner wobei das anionische Tensid kein nachweisbares Niveau von 1,4-Dioxan enthält.

7. Verfahren zur Herstellung der Reinigungszusammensetzung nach Anspruch 1, das folgende Schritte umfasst:

15 a. Beurteilen der Biobasis von beliebigen der kohlenstoffbasierten Komponenten a. bis c. für die Verwendung in der Reinigungszusammensetzung durch Bestimmung des prozentualen Gehalts an modernem Kohlenstoff (percent modern carbon, pMC) für jede Komponente, wobei eine Komponente mit einem pMC-Wert von mindestens 80 % als annehmbar erachtet wird;

20 b. Formulieren einer Reinigungszusammensetzung unter Verwendung der annehmbaren Komponente bzw. der annehmbaren Komponenten aus Schritt a; und

c. Durchführen einer Kopfraumanalyse für die in Schritt b. hergestellte Reinigungszusammensetzung.

25 8. Verfahren nach Anspruch 7, ferner wobei jede der Bestandteile, die die Zusammensetzung ausmachen, weniger als 1 Gew.-% Spurenverunreinigungen umfasst, wobei die Spurenverunreinigungen aus der Gruppe bestehend aus Phenylderivaten und kürzerkettigen Alkoholen ausgewählt sind, und eine Kopfraumanalyse ein Niveau von weniger als 1000 µg/m³ VOCs, umfassend die Gruppe 1,4-Dioxan, Chloroform, Formaldehyd und Acetaldehyd, zeigt.

9. Verfahren nach Anspruch 8, wobei:

30 a. die Phenylivate ferner Toluol und Acetophenon umfassen;
b. die kürzerkettigen Alkohole ferner Alkohole mit einer Kettenlänge von acht Kohlenstoffatomen oder weniger umfassen; und
c. das anionische Tensid kein nachweisbares Niveau von 1,4-Dioxan enthält.

35 10. Verfahren nach Anspruch 9, wobei:

a. das anionische Tensid aus der Gruppe umfassend Natriumalkylbenzolsulfonat, Natriumlaurylsulfat, Natrium-cocoether(2EO)sulfat, Natriumcocosulfat und Natriumoctylsulfat ausgewählt ist;
40 b. das nichtionische Tensid aus der Gruppe umfassend C₁₂-C₁₄-Alkoholethoxylat, C₁₀-C₁₆-Alkylpolyglucoside und C₈-C₁₀-Alkylpolyglucoside ausgewählt ist;
c. das amphotere und zwitterionische Tensid aus der Gruppe umfassend Lauraminoxid, Cocoamidopropylbetaïn, Cocoamidopropylaminoxid, Cocoamid-DEA und Cocodimethylaminoxid ausgewählt sind;
45 d. der mindestens eine Builder oder das mindestens eine Mittel zur Einstellung des pH-Werts aus der Gruppe umfassend Borsäure, Natriumgluconat, Natriumhydroxid und Citronensäure ausgewählt ist; und
e. das mindestens eine Lösungsmittel aus der Gruppe umfassend Glycerin und Wasser ausgewählt ist.

11. Reinigungszusammensetzung nach Anspruch 3, wobei:

50 a. es sich bei dem anionischen Tensid um Natriumcocosulfat handelt;
b. es sich bei dem nichtionischen Tensid um Alkylpolyglucosid handelt;
c. es sich bei dem amphoteren oder zwitterionischen Tensid um Cocoaminoxid handelt;
d. der Builder oder das Mittel zur Einstellung des pH-Werts aus der Gruppe umfassend Borsäure, Citronensäure, Natriumgluconat, Natriumhydroxid und Kaliumcitrat ausgewählt ist; und
55 e. es sich bei dem Lösungsmittel um Glycerin handelt.

12. Verfahren nach Anspruch 10, wobei:

a. es sich bei dem anionischen Tensid um Natriumcocosulfat handelt;

- b. es sich bei dem nichtionischen Tensid um Alkylpolyglucosid handelt;
- c. es sich bei dem amphoteren oder zwitterionischen Tensid um Cocoaminoxid handelt;
- d. der Builder oder das Mittel zur Einstellung des pH-Werts aus der Gruppe umfassend Borsäure, Citronensäure, Natriumgluconat, Natriumhydroxid und Kaliumcitrat ausgewählt ist; und
- e. es sich bei dem Lösungsmittel um Glycerin handelt.

13. Reinigungszusammensetzung nach Anspruch 6, wobei die Zusammensetzung als für die Verwendung durch mindestens eine Person mit multipler Chemikaliensensitivität geeignet erachtet wird.

10 14. Verfahren nach Anspruch 7, das ferner den folgenden Schritt umfasst:

d. Evaluieren der Eignung der Reinigungszusammensetzung für die Verwendung durch mindestens eine Person mit multipler Chemikaliensensitivität.

15 Revendications

1. Composition de nettoyage appropriée pour une utilisation par des personnes sensibles à des produits chimiques comprenant :

20 a. au moins un tensioactif choisi dans le groupe comprenant :

- 1) 0,05 à 15 % en poids d'un tensioactif anionique ;
- 2) 0,05 à 30 % en poids d'un tensioactif non ionique ; et
- 3) 0,05 à 30 % en poids d'au moins un tensioactif amphotère et un tensioactif zwitterionique ;

25 b. 0 à 10 % en poids d'au moins un parmi un adjuvant et un agent d'ajustement du pH ; et
c. 0 à 95 % en poids d'au moins un solvant ;

1) chacun des éléments à base de carbone a. à c. ayant un pMC d'au moins 80 % ;

2) une analyse de l'espace de tête de la composition révélant l'absence de composés de phényle ou leurs dérivés ;

3) moins de 5 % en poids de chaque élément ayant une pression de vapeur qui est supérieure à 0,1 mm Hg à 20 °C ;

4) la composition contenant moins de 1 % en poids d'une matière de fragrances ; et
5) une analyse de l'essence de tête de la composition de nettoyage révélant la présence

5) une analyse de l'espace de tête de la composition de nettoyage révélant la présence de moins de 1 000 µg/m³ de VOC toxiques ou dangereux ;

le tensioactif anionique étant choisi dans le groupe constitué par des sulfates et éthoxysulfates d'alkyle en C₁₀-C₁₄, des sulfonates d'alkyle en C₁₀-C₁₈, des sulfonates d'alkylbenzène linéaire ou ramifié en C₁₀-C₁₄ et des éthoxycarboxylates d'alkyle en C₁₀-C₁₅ et des combinaisons de quelconques des précédents ;

le tensioactif non ionique étant choisi dans le groupe constitué par des alkylpolysaccharides et des produits d'addition d'alcools gras, d'acides gras et d'amines grasses couplé(e)s avec des agents d'alcoxylation choisis dans le groupe constitué par l'oxyde d'éthylène (EO), l'oxyde de propylène (PO), l'oxyde d'isopropylène (IPO), l'oxyde de butylène (BO) et des combinaisons correspondantes ;

le tensioactif amphotère étant choisi dans le groupe constitué par des oxydes d'amine ayant la formule $(R_1)(R_2)R_3NO$, R_1 étant une chaîne hydrocarbonée ayant de 1 à 30 atomes de carbone, R_2 et R_3 étant indépendamment des chaînes hydrocarbonées linéaires ou ramifiées en C_6-C_{12} substituées ou non substituées, et le tensioactif

des chaînes hydrocarbonées linéaires ou ramifiées en C₁-C₄, substituées ou non substituées, et le tensioactif zwitterionique étant choisi dans le groupe constitué par R₁-N'(R₂)(R₃)R₄X, R₁ étant un groupe hydrophobe en C₁₀₋₃₀ ; R₂ et R₃ étant chacun alkyle en C₁-C₄, hydroxyalkyle ou un autre groupe alkyle substitué qui peuvent également être joints pour former des structures cycliques avec le N ; R₄ étant un groupement C₁₋₁₀ joignant l'atome d'azote cationique au groupe hydrophile et étant typiquement un groupe alkylène, hydroxy alkylène ou polyalcoxy, et X étant un groupe carboxylate ou un groupe sulfonate :

polyacrylique et A étant un groupe carbonylato ou un groupe carbonato ; l'au moins un adjuvant ou agent d'ajustement du pH étant choisi dans le groupe constitué par des séquestrants du calcium et du magnésium ; des acides organiques monobasiques ou dibasiques inorganiques ou organiques ; et au moins un agent de conservation dans le groupe constitué par les additifs cités dans la liste ci-dessous :

des sels d'acides monocarboxyliques, des sels d'acides dicarboxyliques, des sels d'acide citrique et d'autres acides organiques appropriés avec des bases conjuguées solubles dans l'eau ; des bases organiques telles que des alcanolamines, y compris des méthanolamines, des éthanolamines et des propanolamines, y compris des diméthanolamines, des diéthanolamines et des dipropanolamines, et y compris des triméthanolamines, des triéthanolamines et des tripropanolamines ; et

l'au moins un solvant étant choisi dans le groupe constitué par des solvants organiques comprenant des C₁-C₆

5 alcanols, des C₁-C₆ diols, des C₁-C₆ triols, des éthers de C₁-C₁₀ alkyle d'alkyléneglycols, des éthers de C₃-C₂₄ alkyléneglycols, des polyalkyléneglycols, des acides carboxyliques à chaîne courte, des esters à chaîne courte, des hydrocarbures isoparaffiniques, des essences minérales, des composés alkylaromatiques, des terpènes, des dérivés de terpènes, des terpénoïdes, des dérivés de terpénoïdes, le formaldéhyde et des pyrrolidones ; des solvants insolubles dans l'eau ; et des solvants solubles dans l'eau.

2. Composition de nettoyage selon la revendication 1,

10 a. le tensioactif anionique étant en outre choisi dans le groupe comprenant un sulfonate d'alkylbenzène de sodium, le laurylsulfate de sodium, un sulfate d'éther de coco de sodium, un sulfate de coco de sodium et le sulfate d'octyle de sodium ;
b. le tensioactif non ionique étant en outre choisi dans le groupe comprenant des éthoxylates d'alcools en C₁₀-C₁₄, des C₁₀₋₁₆-alkylpolyglucosides et des C₈₋₁₀-alkylpolyglucosides ;
15 c. l'au moins un parmi un tensioactif amphotère et un tensioactif zwitterionique étant en outre choisi dans le groupe constitué par des oxydes d'amines en C₁₂₋₁₆, l'oxyde de cocamine, l'oxyde de lauramine, la cocamido-propylbétaïne, l'oxyde de cocamidopropylamine, le cocamide DEA, des alkylamphoglycinates, et des alkyliminopropionates ;
d. l'au moins un adjuvant ou agent d'ajustement du pH étant en outre choisi dans le groupe comprenant des acides choisis parmi l'acide acétique, l'acide maléique, l'acide glycolique, l'acide méthylsulfonique, l'acide chlorhydrique, l'acide sulfurique, l'acide phosphorique, l'acide citrique, l'acide succinique, l'acide borique et le gluconate de sodium ; les bases inorganiques choisies parmi les sels de métaux alcalins et de métaux alcalino-terreux d'hydroxydes, de borates, de sulfonates, de phosphates, de phosphonates et de silicates ; les bases organiques choisies parmi les alcanolamines, y compris les méthanolamines, les éthanolamines et les propanolamines, les diméthanolamines, les diéthanolamines et les dipropanolamines, et les triméthanolamines, les triéthanolamines et les tripropanolamines ; et les sels d'acides organiques, y compris les sels de métaux alcalins et de métaux alcalino-terreux d'acide citrique et gluconique ;
e. l'au moins un solvant étant en outre choisi dans le groupe comprenant le méthanol, l'éthanol, le n-propanol, l'isopropanol, le butanol, le pentanol et l'hexanol, ainsi que leurs isomères ; les méthylènes, éthylène, propylène et butylène glycols ; l'éther monopropylique d'éthylène glycol, l'éther monobutylique d'éthylène glycol, l'éther monohexylique d'éthylène glycol, l'éther monopropylique de diéthylène glycol, l'éther monobutylique de diéthylène glycol, l'éther monohexylique de diéthylène glycol, l'éther méthylique de propylène glycol, l'éther éthylique de propylène glycol, l'éther n-propylique de propylène glycol, l'éther monobutylique de propylène glycol, l'éther t-butylque de propylène glycol, l'éther méthylique ou éthylique ou propylique ou butylique de di- ou tri-polypropylène glycol, les esters d'acétate et de propionate d'éthers de glycol ; l'acétate de glycol et les méthylsiloxanes volatils cycliques ou linéaires ; et les hydrocarbures isoparaffiniques, les essences minérales, les alkylaromatiques, les terpénoïdes, les dérivés terpénoïdes, les terpènes et les dérivés de terpènes.

3. Composition de nettoyage selon la revendication 2,

40 a. le tensioactif anionique étant choisi dans le groupe comprenant un sulfonate d'alkylbenzène de sodium, le laurylsulfate de sodium, le sulfate d'éther (2EO) de coco de sodium, un sulfate de coco de sodium et le sulfate d'octyle de sodium ;
b. le tensioactif non ionique étant choisi dans le groupe comprenant un éthoxylate d'alcool en C₁₂₋₁₄, des C₁₀₋₁₆-alkylpolyglucosides et des C₈₋₁₀-alkylpolyglucosides ;
45 c. les tensioactifs amphotères et zwitterioniques étant choisis dans le groupe comprenant l'oxyde de lauramine, la cocamidopropylbétaïne, l'oxyde de cocamidopropylamine, le cocamide DEA, et l'oxyde de cocodiméthylamine ;
d. l'au moins un adjuvant ou agent d'ajustement du pH étant choisi dans le groupe comprenant l'acide borique, le gluconate de sodium, l'hydroxyde de sodium et l'acide citrique ; et
e. l'au moins un solvant étant choisi dans le groupe comprenant la glycérine et l'eau.

50 5. Composition de nettoyage selon l'une quelconque des revendications 1 à 3, chacun des ingrédients comprenant la composition contenant moins de 1 % en poids de contaminants à l'état de traces, les contaminants à l'état de traces étant choisis dans le groupe comprenant les dérivés de phényle et les alcools à chaîne inférieure.

55 5. Composition de nettoyage selon la revendication 4,

a. les dérivés de phényle comprenant en outre le toluène et l'acétophénone ; et
b. les alcools à chaîne inférieure comprenant en outre des alcools ayant une longueur de chaînes de huit atomes

de carbone ou moins.

6. Composition de nettoyage selon l'une quelconque des revendications 1 à 3, le tensioactif anionique ne contenant pas de taux détectable de 1,4-dioxane.

5 7. Procédé pour la production de la composition de nettoyage selon la revendication 1, comprenant les étapes de :

10 a. l'évaluation de la base biologique de l'un quelconque des composants à base de carbone a. à c. pour une utilisation dans la composition de nettoyage par détermination du pourcentage de carbone moderne (pMC) pour chaque composant, un composant ayant un pMC d'au moins 80 % étant jugé acceptable ;
b. la formulation d'une composition de nettoyage en utilisant le composant ou les composants acceptables de l'étape a. ; et
c. la réalisation d'une analyse de l'espace de tête pour la composition de nettoyage produite dans l'étape b.

15 8. Procédé selon la revendication 7, en outre chacun des ingrédients composant la composition contenant moins de 1 % en poids de contaminants à l'état de traces, les contaminants à l'état de traces étant choisis dans le groupe comprenant les dérivés de phényle et les alcools à chaîne inférieure, et une analyse de l'espace de tête présentant un taux inférieur à 1 000 µg/m³ de COV comprenant le groupe de 1,4-dioxane, chloroforme, formaldéhyde et acétaldéhyde.

20 9. Procédé selon la revendication 8,

25 a. les dérivés de phényle comprenant en outre le toluène et l'acétophénone ;
b. les alcools à chaîne inférieure comprenant en outre des alcools ayant une longueur de chaînes de huit atomes de carbone ou moins ; et
c. le tensioactif anionique ne contenant pas de taux détectable de 1,4-dioxane.

10. Procédé selon la revendication 9,

30 a. le tensioactif anionique étant choisi dans le groupe comprenant un sulfonate d'alkylbenzène de sodium, le laurylsulfate de sodium, le sulfate d'éther (2EO) de coco de sodium, un sulfate de coco de sodium et le sulfate d'octyle de sodium ;
b. le tensioactif non ionique étant choisi dans le groupe comprenant un éthoxylate d'alcool en C₁₂-14, des C₁₀-16-alkylpolyglucosides et des C₈-10-alkylpolyglucosides ;
35 c. les tensioactifs amphotères et zwitterioniques étant choisis dans le groupe comprenant l'oxyde de lauramine, la cocamidopropylbétaïne, l'oxyde de cocamidopropylamine, le cocamide DEA, et l'oxyde de cocodiméthylamine ;
d. l'au moins un adjuvant ou agent d'ajustement du pH étant choisi dans le groupe comprenant l'acide borique, le gluconate de sodium, l'hydroxyde de sodium et l'acide citrique ; et
e. l'au moins un solvant étant choisi dans le groupe comprenant la glycérine et l'eau.

40 11. Composition de nettoyage selon la revendication 3,

45 a. le tensioactif anionique étant un sulfate de coco de sodium ;
b. le tensioactif non ionique étant un alkylpolyglucoside ;
c. le tensioactif amphotère ou zwitterionique étant l'oxyde de cocamine ;
d. l'adjuvant ou l'agent d'ajustement du pH étant choisi dans le groupe comprenant l'acide borique, l'acide citrique, le gluconate de sodium, l'hydroxyde de sodium et le citrate de potassium ; et
e. le solvant étant la glycérine.

50 12. Procédé selon la revendication 10,

55 a. le tensioactif anionique étant un sulfate de coco de sodium ;
b. le tensioactif non ionique étant un alkylpolyglucoside ;
c. le tensioactif amphotère ou zwitterionique étant l'oxyde de cocamine ;
d. l'adjuvant ou l'agent d'ajustement du pH étant choisi dans le groupe comprenant l'acide borique, l'acide citrique, le gluconate de sodium, l'hydroxyde de sodium et le citrate de potassium ; et
e. le solvant étant la glycérine.

13. Composition de nettoyage selon la revendication 6, la composition étant jugée appropriée pour une utilisation par au moins une personne qui manifeste plusieurs sensibilités à des produits chimiques.

14. Procédé selon la revendication 7, comprenant en outre l'étape de :
5 d. évaluation de l'adéquation de la composition de nettoyage pour une utilisation par au moins une personne qui manifeste plusieurs sensibilités à des produits chimiques.

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61982877 B [0001]
- US 30610915 [0001]
- US 10767137 B [0001]
- US 6973362 B [0011]
- US 7096084 B, Long [0011]
- WO 2007099294 A [0012]
- WO 2009024743 A [0012]
- WO 2009024747 A [0012]
- US 5990065 A [0013]
- US 6069122 A [0013]
- US 2012015856 A1, Harthun [0014]
- US 2007207940 A, G. Fenyvesi [0016]
- CZ 200707412, Smidrkal [0017]
- WO 2009131816 A, Luciow [0018]
- WO 2010141184 A1, Clorox-Coulter [0019]
- US 4565647 A [0061]
- US 2082275 A, Daimler [0067]
- US 2702279 A, Funderburk [0067]
- US 2255082 A, Orthner [0067]
- US 5643865 A, Mermelstein [0070]
- US 5622925 A, de Buzzaccarini [0070]
- US 5545350 A, Baker [0070]
- US 5474690 A, Wahl [0070]
- US 5417868 A, Turner [0070]
- US 4661269 A, Trinh [0070]
- US 4439335 A, Burns [0070]
- US 4401578 A, Verbruggen [0070]
- US 4308151 A, Cambre [0070]
- US 4237016 A, Rudkin [0070]
- US 4233164 A, Davis [0070]
- US 4045361 A, Watt [0070]
- US 3974076 A, Wiersema [0070]
- US 3886075 A, Bernadino [0070]
- US 3861870 A, Edwards [0070]
- EP 472178 A, Yamamura [0070]
- US 4702857 A, Gosselink [0079]
- US 4711730 A, Gosselink [0079]
- US 4713194 A, Gosselink [0079]
- US 4877896 A, Maldonado [0079]
- US 4956447 A, Gosselink [0079]
- US 4749596 A, Po [0079]
- US 4849257 A, Borchers, Sr. [0079]
- US 4018569 A, Chang [0079]
- US 4808086 A, Evans [0079]
- US 4756850 A, Nayar [0081]
- US 4103047 A, Zaki [0099]
- EP 0773982 A [0103]
- US 4387089 A, De Polo [0104]
- US 5087372 A, Toyomot [0104]
- US 5073371 A [0104]
- US 5073372 A, Turner [0104]
- US 5972316 A, Robinson [0105]
- US 5968485 A, Robinson [0105]
- US 5935556 A, Tanner [0105]
- US 5827508 A, Tanner [0105]
- US 4511490 A, Stanislowski [0120]
- US 4479881 A, Tai [0120]
- US 4443355 A, Murata [0120]
- US 4435307 A, Barbesgaard [0120]
- US 3983082 A, Ohya [0120]
- US 3950277 A, Silver [0120]
- US 4707291 A, Thorn [0120]
- US 5296161 A [0120]
- US 5030240 A, Wiersema [0120]
- US 5108457 A, Poulose [0120]
- US 5739327 A, Arbogast [0122]
- US 5741437 A [0122]
- US 5814242 A, Alvarez [0122]
- US 5877315 A, Deline [0122]
- US 5888419 A, Casella [0122]
- US 4959187 A [0122]
- US 4778816 A, Fong [0122]
- US 5112514 A [0122]
- US 5002691 A, Bolkan [0122]
- US 5269962 A, Brodbeck [0122]
- US 5234616 A [0122]
- US 5130045 A [0122]
- US 5130044 A, Mitchell [0122]
- US 5057236 A, Petrin [0125]
- GB 1298577 A [0127]
- GB 2076011 A [0127]
- GB 2026054 A [0127]
- GB 2026566 A [0127]
- GB 1393042 A [0127]
- US 3951960 A, Heath [0127]
- US 4298290 A, Barnes [0127]
- US 3993659 A, Meyer [0127]
- US 3980713 A, Matsunaga [0127]
- US 3627758 A, Weber [0127]
- US 4900468 A, Mitchell [0127]
- US 4433975 A [0128]
- US 4432886 A [0128]
- US 4384121 A, Meyer [0128]
- US 4263431 A, Weber [0128]
- US 6251303 B, Bawendi [0128]
- US 6127549 A, Hao [0128]
- US 6133215 A, Zeiger [0128]
- US 6117189 A, Reinehr [0128]

- US 6120704 A, Martini [0128]
- US 6162869 A, Sharma [0128]

Non-patent literature cited in the description

- **A.C. STEINEMANN ; L.G. GALLAGHER ; A.L. DAVIS ; I.C. MACGREGOR.** Chemical Emissions from Residential Dryer Vents During Use of Fragranced Laundry Products. *Air Quality, Atmosphere and Health*, 2011, vol. 6, 151-156 [0009]
- **B. BRIDGES.** Fragrance: Emerging Health and Environmental Concerns. *Flavour and Fragrance Journal*, 2002, vol. 17, 361-371 [0009]
- **R.A. GRAVELING et al.** A Review of Multiple Chemical Sensitivity, 1999 [0020]
- **M.H. KAROL ; O.T. MACINA ; A. CUNNINGHAM.** Cell and molecular biology of chemical allergy. *Ann Allergy Asthma Immunol.*, 2001, vol. 87, 28-32 [0024]
- **G.A. NORTON ; S.L. DEVLIN.** Determining the Modern Carbon Content of Biobased Products Using Radiocarbon Analysis. *Bioresource Technology*, 2006, vol. 97, 2084-2090 [0033]
- **E.A. KRUSEMARK ; W. LI.** Enhanced olfactory sensory perception of threat in anxiety: An event-related fMRI study. *Chemosensory Perception*, 2012, vol. 5, 37-45 [0044]
- **A. STEINEMANN.** Volatile Emissions from Common Consumer Products. *Air Quality, Atmosphere & Health*, March 2015 [0046]
- **MCCUTCHEON.** Emulsifiers and Detergents. MC Publishing Co., 1995, vol. 1 [0057] [0062] [0067] [0071]
- **KIRK-OTHMER.** Encyclopedia of Chemical Technology. 1978, vol. 3, 128-148 [0103]
- Sunscreens: Development, Evaluation, and Regulatory Aspects. Marcel Dekker, Inc, 1990 [0104]
- **MCCUTCHEON.** Functional Materials. MC Publishing Co., 1995, vol. 2 [0128]
- Encyclopedia of Chemical Technology. John Wiley & Sons, 1994 [0128]