



(22) Date de dépôt/Filing Date: 1999/12/06
(41) Mise à la disp. pub./Open to Public Insp.: 2000/06/15
(62) Demande originale/Original Application: 2 351 580
(30) Priorités/Priorities: 1998/12/07 (GB9826882.4);
1999/03/04 (GB9904934.8)

(51) Cl.Int./Int.Cl. *C07D 498/08* (2006.01)

(71) Demandeur/Applicant:
NOVARTIS AG, CH

(72) Inventeurs/Inventors:
NAVARRO, FRANCOIS, FR;
PETIT, SAMUEL, FR;
STONE, GUY, CH

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : FORME CRISTALLINE DE LA 40-O-(2-HYDROXYETHYL)RAPAMYCINE ET METHODE DE PREPARATION
CONNEXE

(54) Title: CRYSTALLINE FORM OF 40-O- (2-HYDROXY-ETHYL) RAPAMYCIN AND PROCESS FOR PREPARATION
THEREOF

(57) **Abrégé/Abstract:**

The invention provides the compound 40-O- (2-hydroxy-ethyl) rapamycin in crystalline non-solvable form as well as a process for preparing 40-O- (2-hydroxy-ethyl)-rapamycin in substantially pure form, comprising crystallizing 40-O- (2-hydroxy-ethyl)-rapamycin from a crystal bearing medium and recovering the crystals thus obtained.



21489-9705D

CRYSTALLINE FORM OF 40-O-(2-HYDROXY-ETHYL) RAPAMYCIN AND
PROCESS FOR PREPARATION THEREOF

Abstract

The invention provides the compound 40-O-(2-
5 hydroxy-ethyl) rapamycin in crystalline non-solvable form as
well as a process for preparing 40-O-(2-hydroxy-ethyl)-
rapamycin in substantially pure form, comprising
crystallizing 40-O-(2-hydroxy-ethyl)-rapamycin from a
crystal bearing medium and recovering the crystals thus
10 obtained.

21489-9705D

1

**CRYSTALLINE FORM OF 40-O-(2-HYDROXY-ETHYL) RAPAMYCIN AND
PROCESS FOR PREPARATION THEREOF**

This is a divisional application of Canadian Patent Application No. 2,351,580 filed December 6, 1999.

5 The present invention relates to the stabilization of a pharmaceutically active ingredient sensitive to oxidation, e.g. a poly-ene macrolide, preferably a poly-ene macrolide having immunosuppressant properties, particularly rapamycins.

10 The subject matter of this divisional application is directed to a crystalline form of 40-O-(2-hydroxyethyl) rapamycin and to a process for preparation thereof.

 The subject matter of the parent application has been restricted to a process for stabilizing a poly-ene
15 macrolide and to a solid mixture comprising the poly-ene macrolide and an antioxidant. However, it should be understood that the expression "the invention" and the like as used herein, encompasses the subject matter of both the parent and this divisional application.

20 The handling and storage particularly in the bulk form of pharmaceutically active ingredients which are sensitive to oxidation is difficult. Special handling is necessary and often the oxidation-sensitive ingredient is stored in air-tight packaging under protective gas.
25 Substantial amounts of stabilizers are added during the formulating process of such pharmaceutically active ingredients.

21489-9705D

1a

Poly-ene macrolides have satisfactory stability properties. However, it has now been found that their stability to oxygen may substantially be improved by the addition of a stabilizer, e.g. an antioxidant, during their
5 isolation step.

According to one aspect of the invention of the parent application, there is provided a solid mixture comprising a poly-ene macrolide and an antioxidant wherein the poly-ene macrolide is selected from the group consisting
10 of rapamycin, a 16-O-substituted rapamycin, and a 40-O-substituted rapamycin and wherein the antioxidant is present in a catalytic amount.

According to another aspect of the invention of the parent application, there is provided a process for
15 stabilizing a poly-ene macrolide selected from the group consisting of rapamycin, a 16-O-substituted rapamycin, and a 40-O-substituted rapamycin, said process comprising:
(a) dissolving pure rapamycin, pure 16-O or pure 40-O-substituted rapamycin in an inert solvent; (b) adding
20 an antioxidant to the resulting solution in an amount up to 1% based on the rapamycin, the 16-O or the 40-O-substituted rapamycin weight, and (c) isolating the resulting mixture of the rapamycin or the 16-O or 40-O-substituted rapamycin and the antioxidant.

25 According to still another aspect of the invention of the parent application, there is provided a process for stabilizing a compound selected from the group consisting of 40-O-(2-hydroxy)ethyl-rapamycin, 40-O-(3-hydroxy)propyl-rapamycin, and 40-O-[2-(2-hydroxy)ethoxy]ethyl-rapamycin
30 comprising: a. dissolving the compound in an inert solvent; b. adding an antioxidant to the resulting solution in an amount of up to 1% based on the compound weight; and

21489-9705D

1b

c. isolating the resulting mixture of the compound and the antioxidant.

According to yet another aspect of the invention of the parent application, there is provided a mixture in solid form comprising a compound selected from the group consisting of 40-O-(2-hydroxy)ethyl-rapamycin, 40-O-(3-hydroxy)propyl-rapamycin, and 40-O-[2-(2-hydroxy)ethoxy]ethyl-rapamycin, and an antioxidant wherein the antioxidant is present in an amount of up to 1% based on the compound weight.

According to a further aspect of the invention of the parent application, there is provided a mixture in solid form comprising a compound selected from the group consisting of 40-O-(2-hydroxy)ethyl-rapamycin, 40-O-(3-hydroxy)propyl-rapamycin, and 40-O-[2-(2-hydroxy)ethoxy]ethyl-rapamycin, and an antioxidant wherein the antioxidant is present in an amount of up to 1% based on the compound weight, wherein the compound is prepared by a process comprising: a. dissolving the compound in an inert solvent; b. adding an antioxidant to the resulting solution in an amount of up to 1% based on the compound weight; and c. isolating the resulting mixture of the compound and the antioxidant.

According to one aspect of the invention of the present divisional application, there is provided a process for preparing 40-O-(2-hydroxy-ethyl)-rapamycin in substantially pure form, comprising crystallizing 40-O-(2-hydroxy-ethyl)-rapamycin from a crystal bearing medium and recovering the crystals thus obtained. According to another aspect of the invention, there is provided the compound obtained by the process.

21489-9705D

1c

According to the invention, there is provided

1. A process for stabilizing a poly-ene macrolide comprising adding an antioxidant to the purified macrolide, preferably at the commencement of its isolation step.

5 This process is particularly useful for the production of a stabilized poly-ene macrolide in bulk. The amount of antioxidant may conveniently be up to 1%, more preferably from 0.01 to 0.5% (based on the weight of the macrolide). Such a small amount is referred to hereinafter
10 as a catalytic amount.

As alternatives to the above the present invention also provides:

2. A mixture, e.g. a bulk mixture, comprising a poly-ene macrolide and an anti-oxidant, preferably a
15 catalytic amount thereof, preferably in solid form.

The mixture may be in particulate form e.g. crystallized or amorphous form. It may be in a sterile or substantially sterile condition, e.g. in a condition suitable for pharmaceutical use.

21489-9705D

- 2 -

3. Use of a mixture as defined above in 2. in the manufacture of a pharmaceutical composition.

Examples of poly-enes macrolides are e.g. molecules comprising double bonds, preferably conjugated double bonds, for example such having antibiotic and/or immunosuppressant properties, e.g. macrolides comprising a lactam or lactone bond and their derivatives, e.g. compounds which have a biological activity qualitatively similar to that of the natural macrolide, e.g. chemically substituted macrolides. Suitable examples include e.g. rapamycins and ascomycins. A preferred poly-ene macrolide is a macrolide comprising at least 2 conjugated double bonds, e.g. 3 conjugated double bonds.

Rapamycin is a known lactam macrolide produceable, for example by Streptomyces hygroscopicus. The structure of rapamycin is given in Kessler, H. et al.; 1993; Helv. Chim. Acta, 76 : 117. Rapamycin has antibiotic and immunosuppressant properties. Derivatives of rapamycin are known, e.g. 16-O-substituted rapamycins, for example as disclosed in WO 94/02136 and WO 96/41807, 40-O-substituted rapamycins, for example as disclosed in WO 94/09010, WO 92/05179, WO 95/14023, 94/02136, WO 94/02385 and WO 96/13273, all of which being incorporated herein by reference. Preferred rapamycin derivatives are e.g. rapamycins wherein the hydroxy in position 40 of formula A illustrated at page 1 of WO 94/09010 is replaced by -OR wherein R is hydroxyalkyl, hydroxyalkoxyalkyl, acylaminoalkyl or aminoalkyl, e.g. 40-O-(2-hydroxy)ethyl-rapamycin, 40-O-(3-hydroxy)propyl-rapamycin, and 40-O-[2-(2-hydroxy)ethoxy]ethyl-rapamycin.

Ascomycins, of which FK-506 and ascomycin are the best known, form another class of lactam macrolides, many of which have potent immunosuppressive and anti-inflammatory activity. FK506 is a lactam macrolide produced by Streptomyces tsukubaensis. The structure of FK506 is given in the Appendix to the Merck Index, 11th ed. (1989) as item A5. Ascomycin is described e.g. in USP 3,244,592. Ascomycin, FK506, other naturally occurring macrolides having a similar biological activity and their derivatives, e.g. synthetic analogues and derivatives are termed collectively "Ascomycins". Examples of synthetic analogues or derivatives are e.g. halogenated ascomycins, e.g. 33-epi-chloro-33-desoxy-ascomycin such as disclosed in EP-A-427,680, tetrahydropyran derivatives, e.g. as disclosed in EP-A-626,385.

21489-9705D

- 3 -

Particularly preferred macrolides are rapamycin and 40-O-(2-hydroxy)ethyl-rapamycin.

Preferred antioxidants are for example 2,6-di-tert.-butyl-4-methylphenol (hereinafter BHT), vitamin E or C, BHT being particularly preferred.

A particularly preferred mixture of the invention is a mixture of rapamycin or 40-O-(2-hydroxy)ethyl-rapamycin and 0.2% (based on the weight of the macrolide) of antioxidant, preferably BHT.

The antioxidant may be added to the poly-ene macrolide at the commencement of the isolation steps, preferably the final isolation step, more preferably just prior to the final precipitation step. The macrolide is preferably in a purified state. It may be dissolved in an inert solvent and the antioxidant is added to the resulting solution, followed by a precipitation step of the stabilized macrolide, e.g. in an amorphous form or in the form of crystals. Preferably the mixture of the invention is in amorphous form.

The resulting stabilized macrolide exhibits surprisingly an improved stability to oxidation and its handling and storage, e.g. in bulk form prior to its further processing for example into a galenic composition, become much easier. It is particularly interesting for macrolides in amorphous form.

The macrolide stabilized according to the invention may be used as such for the production of the desired galenic formulation. Such formulations may be prepared according to methods known in the art, comprising the addition of one or more pharmaceutically acceptable diluent or carrier, including the addition of further stabilizer if required.

Accordingly there is further provided:

4. A pharmaceutical composition comprising, as active ingredient, a stabilized mixture as disclosed above, together with one or more pharmaceutically acceptable diluent or carrier.

The composition of the invention may be adapted for oral, parenteral, topical (e.g. on the skin), ocular, nasal or inhalation (e.g. pulmonary) administration. A preferred

21489-9705D

- 4 -

composition is one for oral administration, preferably a water-free composition when the active ingredient is a lactone macrolide.

The pharmaceutical compositions of the invention may comprise further excipients, e.g. a lubricant, a disintegrating agent, a surfactant, a carrier, a diluent, a flavor enhancer, etc. It may be in liquid form, e.g. solutions, suspensions or emulsions such as a microemulsions, e.g. as disclosed in USP 5,536,729, or in solid form, e.g. capsules, tablets, dragées, powders (including micronized or otherwise reduced particulates), solid dispersions, granulates, etc., e.g. as disclosed in WO 97/03654, the contents of which being incorporated herein by reference, or semi-solid forms such as ointments, gels, creams and pastes. They are preferably adapted to be in a form suitable for oral administration. Preferably they are in solid form. The pharmaceutical compositions of the invention may be prepared according to known methods, by mixing the macrolide stabilized according to the invention with the additional ingredients under stirring; the ingredients may be milled or ground and if desired compressed, e.g. into tablets.

This invention is particularly interesting for rapamycin compositions in liquid or solid form. A particularly preferred composition is a solid dispersion, e.g. comprising a stabilized rapamycin according to the invention and a carrier medium, e.g. a water-soluble polymer such as hydroxypropylmethylcellulose, e.g. as disclosed in WO 97/03654.

The compositions of the invention are useful for the indications as known for the macrolide they contain at e.g. known dosages. For example, when the macrolide has immunosuppressant properties, e.g. rapamycin or a rapamycin derivative, the composition may be useful e.g. in the treatment or prevention of organ or tissue acute or chronic allo- or xeno-transplant rejection, autoimmune diseases or inflammatory conditions, asthma, proliferative disorders, e.g. tumors, or hyperproliferative vascular disorders, preferably in the prevention or treatment of transplant rejection.

The amount of macrolide and of the composition to be administered depend on a number of factors, e.g. the active ingredient used, the conditions to be treated, the duration of the treatment etc. For e.g. rapamycin or 40-O-(2-hydroxy)ethyl-rapamycin, a suitable daily dosage form for oral administration comprise from 0.1 to 10 mg, to be administered once or in divided form.

21489-9705D

- 5 -

In another aspect, this invention also provides 40-O-(2-hydroxy)ethyl-rapamycin in a crystalline form, particularly in a substantially pure form. Preferably the crystal form is characterized by the absence or substantial absence of any solvent component; it is in non-solvate form.

40-O-(2-hydroxy)ethyl-rapamycin in crystalline form belongs to the monoclinic sytem. The resulting crystals have a m.p. of 146°-147°C, especially 146.5°C. To assist identification of the new crystalline form, X-ray diffraction analysis data are provided. The conditions under which these data are obtained are as follows:

Temperature	293(2)K
Wavelength	1.54178 Å
Space group	P2 ₁
Unit cell dimensions	
a	14.378.(2) Å
b	11.244(1) Å
c	18.310(2) Å
β	108.58(1)°
Volume	2805.8(6) Å ³
Z	2
Density (calculated)	1.134 g/cm ³
Absorption coefficient	0.659 mm ⁻¹
F(000)	1040
Crystal size	0.59x0.11x0.03 mm
θ range for data collection	2.55 to 57.20°
Reflections collected	4182
Independent reflections	4037 [R(int) = 0.0341]
Intensity decay	32%
Refinement method	Full-matrix least-squares on F ²
Data/restraints/parameters	3134/1/613
Goodness-of-fit on F ²	1.055
Final R indices [I>2 sigma(I)]	R ₁ =0.0574, wR ₂ =0.1456
Largest diff. peak and hole	0.340 and -0.184 e/Å ³

21489-9705D

- 6 -

40-O-(2-hydroxy)ethyl-rapamycin in crystalline form may be prepared by dissolving the amorphous compound in a solvent e.g. ethyl acetate and adding an aliphatic hydrocarbon C_nH_{2n+2} ($n=5, 6$ or 7). After addition of the hydrocarbon, the resulting mixture may be warmed e.g. at a temperature of 25 to 50°C , e.g. up to 30 - 35°C . Storing of the resulting mixture may conveniently take place at a low temperature, e.g. below 25°C , preferably from 0 to 25°C . The crystals are filtered and dried. Heptane is preferred as an aliphatic hydrocarbon. If desired, nucleation procedures may be commenced e.g. by sonication or seeding.

The present invention also provides a process for purifying 40-O-(2-hydroxy)ethyl-rapamycin comprising crystallizing 40-O-(2-hydroxy)ethyl-rapamycin from a crystal bearing medium, e.g. as disclosed above, and recovering the crystals thus obtained. The crystal bearing medium may include one or more components in addition to those recited above. A particularly suitable crystal bearing medium has been found to be one comprising ca. 2 parts ethyl acetate and ca. 5 parts aliphatic hydrocarbon, e.g. heptane.

40-O-(2-hydroxy)ethyl-rapamycin in crystalline form has been found to possess in vitro and in vivo immunosuppressive activity comparable to that of the amorphous form. In the localized GvHD, maximal inhibition (70-80%) of lymph node swelling is achieved with a dosage of 3 mg with 40-O-(2-hydroxy)ethyl-rapamycin in crystalline form.

40-O-(2-hydroxy)ethyl-rapamycin may be useful for the same indications as known for the amorphous compound, e.g. to prevent or treat acute and chronic allo- or xeno-transplant rejection, autoimmune diseases or inflammatory conditions, asthma, proliferative disorders, e.g. tumors, or hyperproliferative vascular disorders, e.g. as disclosed in WO 94/09010 or in WO 97/35575, the contents thereof being incorporated herein by reference. In general, satisfactory results are obtained on oral administration at dosages on the order of from 0.05 to 5 or up to 20 mg/kg/day, e.g. on the order of from 0.1 to 2 or up to 7.5 mg/kg/day administered once or, in divided doses 2 to 4x per day. Suitable daily dosages for patients are thus on the order of up to 10 mg., e.g. 0.1 to 10 mg.

40-O-(2-hydroxy)ethyl-rapamycin in crystalline form may be administered by any conventional route, e.g. orally, for example tablets or capsules, or nasally or pulmonary (by inhalation). It may be administered as the sole active ingredient or together with other

21489-9705D

- 7 -

drugs, e.g. immunosuppressive and/or immunomodulatory and/or anti-inflammatory agents, e.g. as disclosed in WO 94/09010.

In accordance with the foregoing, the present invention also provides:

5. A method for preventing or treating acute or chronic allo- or xeno-transplant rejection, autoimmune diseases or inflammatory conditions, asthma, proliferative disorders, or hyperproliferative vascular disorders, in a subject in need of such treatment, which method comprises administering to said subject a therapeutically effective amount of 40-O-(2-hydroxy)ethyl-rapamycin in crystalline form;
6. 40-O-(2-hydroxy)ethyl-rapamycin in crystalline form for use as a pharmaceutical; e.g. in a method as disclosed above;
7. A pharmaceutical composition comprising 40-O-(2-hydroxy)ethyl-rapamycin in crystalline form together with a pharmaceutically acceptable diluent or carrier therefor;
8. A kit or package for use in immunosuppression or inflammation, including a pharmaceutical composition as disclosed above and a pharmaceutical composition comprising an immunosuppressant or immunomodulatory drug or an anti-inflammatory agent.

The following examples illustrate the invention without limiting it.

Example 1: Crystallisation

0.5 g amorphous 40-O-(2-hydroxy)ethyl-rapamycin is dissolved in 2.0 ml ethyl acetate at 40°C. 5.0 ml heptane is added and the solution becomes "milky". After warming to 30°C, the solution becomes clear again. Upon cooling to 0°C and with scratching an oil falls out of the solution. The test tube is closed and stored at 10°C overnight. The resulting white voluminous solid is then filtered and washed with 0.5 ml of a mixture of ethyl acetate/hexane (1:2.5) and the resulting crystals are dried at 40°C under 5 mbar for 16 hours. 40-O-(2-hydroxy)ethyl-rapamycin in crystalline form having a m.p. of 146.5°C is thus obtained.

Crystallisation at a larger scale may be performed as follows:

21489-9705D

- 8 -

250 g amorphous 40-O-(2-hydroxy)ethyl-rapamycin is dissolved in 1.0 l ethyl acetate under argon with slow stirring. This solution is heated at 30°C and then during 45 minutes, 1.5 l heptane is added dropwise. 0.25 g of seed crystals prepared as disclosed above are added under the same conditions in portions. The mixture is further stirred at 30°C over a period of 2 hours and the crystallisation mixture is cooled to 25°C over 1 hour and then to 10°C for 30 minutes and filtered. The crystals are washed with 100 ml of a mixture ethyl acetate/hexane (2:3). Subsequent drying is performed at 50°C and ca 5 mbar. m.p. 146.5°C

IR in KBr: 3452, 2931, 1746, 1717, 1617, 1453, 1376, 1241, 1191, 1163, 1094, 1072,
1010, 985, 896 cm⁻¹

Single X-ray structure with coordinates are indicated in Figures 1 to 3 below.

Example 2: Production of stabilized 40-O-(2-hydroxy)ethyl-rapamycin

100g 40-O-(2-hydroxy)ethyl-rapamycin are dissolved in 600l abs. ethanol. After addition of 0.2g BHT, the resulting solution is added dropwise with stirring to 3.0 l water within 1 hour. The resulting suspension is stirred for an additional 30 minutes. Filtration with subsequent washing (3x200 ml water/ethanol at a v/v ratio of 5:1) results in a moist white product which is further dried under vacuum (1mbar) at 30°C for 48 hours. The resulting dried product contains 0.2% (w/w) BHT.

The resulting product shows improved stability on storage. The sum of by-products and degradation products in % after 1 week storage are as follows:

Compound	50°C in open flask
Ex. 2 (0.2% BHT)	1.49
Without BHT	>10

The procedure of above Example may be repeated but using, as active ingredient, rapamycin.

21489-9705D

9

CLAIMS:

1. A process for preparing 40-O-(2-hydroxy-ethyl)-rapamycin in substantially pure form, comprising crystallizing 40-O-(2-hydroxy-ethyl)-rapamycin from a crystal bearing medium and recovering the crystals thus obtained.
2. The compound 40-O-(2-hydroxy-ethyl)-rapamycin prepared according to the process defined in claim 1 which is in crystalline non-solvate form.
- 10 3. The compound 40-O-(2-hydroxy-ethyl)-rapamycin prepared according to the process defined in claim 1 having a crystal lattice as follows: $a=14.37 \text{ \AA}$, $b=11.24 \text{ \AA}$, $c=18.31 \text{ \AA}$, the volume being 2805 \AA^3 .

FETHERSTONHAUGH & CO.

OTTAWA, CANADA

PATENT AGENTS