




US 20210292393A1

(19) **United States**(12) **Patent Application Publication**
WESTENDORF et al.(10) **Pub. No.: US 2021/0292393 A1**(43) **Pub. Date: Sep. 23, 2021**(54) **ANTI-CORONAVIRUS ANTIBODIES AND METHODS OF USE****as represented by the Secretary,
Department of Health and Human
Servi, Bethesda, MD (US)**(71) Applicants: **AbCellera Biologics Inc.**, Vancouver
(CA); **The United States of America,**
as represented by the Secretary,
Department of Health and Human
Servi, Bethesda, MD (US)(21) Appl. No.: **17/116,588**(22) Filed: **Dec. 9, 2020****Related U.S. Application Data**(72) Inventors: **Kathryn WESTENDORF**, Vancouver
(CA); **Stefanie ZENTELIS**, Vancouver
(CA); **Krithika MUTHURAMAN**,
Toronto (CA); **Kevin JEPSON**,
Vancouver (CA); **Ester FALCONER**,
Vancouver (CA); **John MASCOLA**,
Bethesda, MD (US); **Barney**
GRAHAM, Bethesda, MD (US);
Kizzmekia CORBETT, Bethesda, MD
(US); **Julie LEDGERWOOD**,
Bethesda, MD (US); **Lingshu WANG**,
Bethesda, MD (US); **Olubukola**
ABIONA, Bethesda, MD (US); **Wei**
SHI, Bethesda, MD (US); **Wing-pui**
KONG, Bethesda, MD (US); **Yi**
ZHANG, Bethesda, MD (US); **Bryan**
Edward JONES, San Diego, CA (US);
Denisa FOSTER, San Diego, CA (US);
Julian DAVIES, La Jolla, CA (US);
Qing CHAI, San Diego, CA (US);
Christopher Carl FRYE, Bargserville,
IN (US); **Ganapathy**
GOPALRATHNAM, Fishers, IN (US);
Jörg HENDLE, San Diego, CA (US);
John Michael SAUDER, Carlsbad, CA
(US); **Jeffrey Streetman BOYLES**,
Indianapolis, IN (US); **Anna**
PUSTILNIK, San Diego, CA (US)(60) Provisional application No. 63/116,483, filed on Nov.
20, 2020, provisional application No. 63/085,042,
filed on Sep. 29, 2020, provisional application No.
63/080,351, filed on Sep. 18, 2020, provisional ap-
plication No. 63/036,089, filed on Jun. 8, 2020,
provisional application No. 63/030,530, filed on May
27, 2020, provisional application No. 63/010,999,
filed on Apr. 16, 2020, provisional application No.
62/987,313, filed on Mar. 9, 2020.**Publication Classification**(51) **Int. Cl.**
C07K 16/10 (2006.01)
A61K 39/215 (2006.01)
G01N 33/50 (2006.01)
A61K 47/68 (2017.01)
A61K 39/00 (2006.01)
(52) **U.S. Cl.**
CPC **C07K 16/10** (2013.01); **A61K 39/215**
(2013.01); **A61K 2039/505** (2013.01); **A61K**
47/6801 (2017.08); **G01N 33/5091** (2013.01)(57) **ABSTRACT**Antibodies that bind SARS-CoV spike protein, SARS-
CoV-2 spike protein, and methods of using same for treating
or preventing conditions associated with SARS or COVID-
19 and for detecting SARS-CoV or SARS-CoV-2.**Specification includes a Sequence Listing.**(73) Assignees: **AbCellera Biologics Inc.**, Vancouver
(CA); **The United States of America,****SARS-CoV-2 S-2P binding to 555 Fab**

$$K_D = 1.45 \times 10^{-9} \text{ M}$$

$$k_a = 2.91 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$$

$$k_d = 4.23 \times 10^{-4} \text{ s}^{-1}$$

[illegible]

www.ck12.org

[illegible]

www.ck12.org


ID	824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																									824-C1 Fold on- including PA1-823 His																								
----	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

FIGURE 1

[illegible]

www.pearsoned.com

Figure 2A



Fab555




Figure 2B






Figure 2C

Figure 3A

Fab447

**Figure 3B**






Figure 3C

Figure 4A

Fab483

**Figure 4B**





Figure 4C

Figure 5A



Fab419




Figure 5B

Fab 419





Figure 5C

Figure 6A



Fab388






Figure 6B



Fab388

Figure 6C






Figure 7

Figure 8A




Figure 8B

Fab494




Figure 8C




Figure 9A




Figure 9B




Figure 9C




Figure 9D




Figure 9E





Figure 9F






Figure 10A




Figure 10B

- ◆ 50 mg/kg IgG1 Control
- 1 mg/kg LY-CoV555
- ★ 2.5 mg/kg LY-CoV555
- ◇ 15 mg/kg LY-CoV555
- ✱ 50 mg/kg LY-CoV555

- * q-val < 0.05, 1 mg/kg
- * q-val < 0.05, 2.5 mg/kg
- ◇ q-val < 0.05, 15 mg/kg
- ✱ q-val < 0.05, 25 mg/kg




Figure 10C




Figure 10D

- ◆ 50 mg/kg IgG1 Control
- 1 mg/kg LY-CoV555
- ★ 2.5 mg/kg LY-CoV555
- ◆ 15 mg/kg LY-CoV555
- ▼ 50 mg/kg LY-CoV555

- * q-val < 0.05, 1 mg/kg
- * q-val < 0.05, 2.5 mg/kg
- * q-val < 0.05, 15 mg/kg
- * q-val < 0.05, 25 mg/kg




Figure 10E




Figure 10F




Figure 10G




Figure 10H

ANTI-CORONAVIRUS ANTIBODIES AND METHODS OF USE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] Priority is claimed to U.S. Provisional Applications Nos. 62/987,313 filed 9 Mar. 2020; 63/010,999 filed 16 Apr. 2020; 63/030,530 filed 27 May 2020; 63/036,089 filed 8 Jun. 2020; 63/080,351 filed 18 Sep. 2020; 63/085,042 filed 29 Sep. 2020; and 63/116,483 filed 20 Nov. 2020, each of which is hereby incorporated by reference in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] This invention was made with government support under D18AC00002 awarded by the Defense Advanced Research Projects Agency. The government has certain rights in the invention.

SEQUENCE LISTING

[0003] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on 4 Dec. 2020, is named 27050015WO01120420SEQLST25.txt and is 4,552,494 bytes in size.

TECHNICAL FIELD OF THE INVENTION

[0004] This disclosure generally relates to the fields of medicine, immunology, and infectious disease. More specifically, the disclosure relates to anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) and antibody-like molecules and, in particular, to human antibodies, antibody fragments, and nucleic-acid-vectored versions thereof, and methods for treating coronavirus infections, methods for prophylaxis against coronavirus infections, and immunoassays for the detection of coronaviruses.

BACKGROUND OF THE INVENTION

[0005] Past decades have seen yearly new or reemerging viral outbreaks including Severe Acute Respiratory Syndrome coronavirus (SARS-CoV), West Nile, Ebola, Middle Eastern Respiratory Syndrome Coronavirus (MERS-CoV), Zika, and pandemic influenza. The risk of a pandemic is multiplied by growing populations and urbanization, climate change, global travel, and civil conflict. Recently, the 2019 novel coronavirus (SARS-CoV-2) outbreak has led the World Health Organization (WHO) to declare a global health emergency. These viral outbreaks have important consequences on human societies, creating a huge burden on healthcare systems and having important repercussions on the economy.

[0006] Pandemic outbreaks present a serious risk to global security and trade. For example, the 1918 H1N1 pandemic influenza virus (Spanish flu) claimed an estimated 50 million lives in a matter of months. The World Bank estimates that a pandemic to that scale would cost about 5% global gross domestic product (GDP), or about \$3 trillion USD. The more recent 2009 H1N1 pandemic is estimated to have infected 11-21% of the world's population (80% under the age of 65) and claimed between 151,700 to 575,400 lives. In non-pandemic years, the WHO estimates three to five million

cases of severe influenza, resulting in 250,000 to 500,000 global deaths. The propensity for rapid genetic change, antigenic diversity, and the breadth of potential hosts gives influenza type A significant pandemic potential. The rapid mutation of surface antigens allows the virus to evade the immune response, requiring an annual prediction of circulating strains to include in seasonal vaccines.

[0007] A universal flu vaccine that could provide long-term and cross-strain protection has so far been elusive, and while there are candidates in preclinical and clinical trials, it will likely be several years before one becomes available. Less predictable than seasonal drifts, genetic reassortments (antigenic shift) of hemagglutinin or neuraminidase in zoonotic hosts can generate novel strains of influenza virus that can transfer to humans and cause a fast-spreading pandemic across an immunologically naïve population. The California 2009 H1N1 strain was discovered to have originated in swine, and both the Spanish flu and Hong Kong 1957 pandemic strains are thought to have originated from an avian source. Surveillance of rapidly-shifting virulence factors in domesticated and wild zoonotic hosts is currently not possible, adding to the urgent need for a rapid response strategy to emerging influenza pandemics. The availability of a large collection of human antibodies against viral families that constitute a present or future threat would accelerate response to outbreaks by allowing for rapid identification of antibodies with properties suitable for diagnostics, prophylaxis, or therapeutics.

[0008] Pandemic Potential of Coronaviruses

[0009] Coronaviruses are a large family of viruses that infect mammals and birds, with four endemic strains that circulate commonly in humans, including human coronavirus 229E, OC43, NL63 and HKU1. These endemic strains generally cause mild flu-like symptoms, but can also cause more serious pneumonia in vulnerable populations. However, when new coronaviruses jump from zoonotic hosts (bats, birds, camels, etc.) to humans, they can result in a much more severe respiratory disease that can spread quickly through the population. Examples of such outbreaks include SARS (Severe Acute Respiratory Syndrome) in 2003, caused by SARS-associated coronavirus (SARS-CoV) which resulted in an outbreak that infected over 8,000 people worldwide and was ~10% fatal, and MERS (Middle Eastern Respiratory Syndrome) in 2012, caused by MERS-CoV, which to date has infected over 2,500 people with approximately 35% fatalities.

[0010] At the time of this application, a new coronavirus, named SARS-CoV-2, has become a serious global outbreak causing a severe respiratory disease referred to as COVID-19, and looks certain to become a global pandemic. Following the first reported cases in Wuhan, China in late 2019, there are presently over 100,000 confirmed cases globally, resulting in over 3,500 deaths. SARS-Cov-2 is now the most serious coronavirus outbreak in history. Unlike MERS-CoV and SARS-CoV, SARS-CoV-2 has spread rapidly on a global scale, with confirmed cases in more than 90 countries to date. In response to this outbreak, there is an urgent need for antibody-based therapeutics, prophylactics, and diagnostics to treat, prevent, and detect the disease. Each of these applications demand antibodies with specific properties including epitope recognition, affinity, ease of manufacturing, solubility, and specificity. Furthermore, for use in therapeutic applications, human antibodies are required. The availability of a large and diverse library of human antibodies

ies with specificity to coronaviruses would be a valuable resource for rapid response to emerging coronavirus outbreaks, allowing for the rapid screening and selection of antibodies suitable for therapeutic, prophylactic, and diagnostics uses.

[0011] Antibodies as a Countermeasure Against Viral Threats

[0012] Although vaccines are the cheapest and most effective protective countermeasures, their use is limited against viruses with high antigenic drift, antigenic shift, and against newly emerging viral strains from zoonotic hosts to which humans have little or no herd immunity. After vaccination, it takes several days post-immunization for the immune system to generate protective immunity—this is often impractical in pandemic situations where human efforts must be rapidly deployed in outbreak areas and immediate action is needed to prevent or treat infections. Passive immunization with antiviral monoclonal antibodies (mAbs) has emerged as a viable strategy to protect at-risk populations from seasonal epidemics and fast-moving pandemics.

[0013] The challenges associated with deploying such an approach at scale has prompted efforts to isolate neutralizing antibodies from convalescent patients that can be used as prophylactics or therapeutics. Several groups have shown it is feasible to discover and deliver highly potent mAbs in very short timeframes. During the 2014-2015 MERS-CoV outbreaks, two groups isolated highly potent MERS-specific mAbs, produced them recombinantly in gram quantities and tested them in animal models in a relatively short timeframe: Corti and colleagues went from immortalized B-cell screening from a single human convalescent donor to the production of prophylactically protective mAbs in four months. Pascal and colleagues identified potent fully human mAbs from immunized transgenic mice within several weeks, using both hybridoma and B-cell sorting methods. Likewise, during the 2015-2016 Ebola epidemic, several groups rapidly generated potent Ebola virus GP-specific neutralizing mAbs from memory B-cells of convalescent human donors. In all these cases, manufacturing antibody proteins in sufficient quantities for deployment at large scale was a limiting factor to generate an effective countermeasure in a timely manner.

[0014] Although multiple antiviral mAbs are in preclinical and clinical development, there is only one approved antiviral mAb on the market (palivizumab from MedImmune for respiratory syncytial virus (RSV). While mAbs have experienced large success in other indications such as oncology and immune disorders, obstacles in antiviral mAb discovery largely originate from two factors: First, the relative high cost of mAb production and demanding administration protocols can make this approach difficult to apply on a global scale. Second, only a small proportion of infected individuals generate broad and potent neutralizing responses to viruses with high antigenic variability (i.e. HIV, Ebola, Lassa, and influenza). These rare mAbs generally represent a small fraction of B cells in infected individuals, which makes them very difficult to find.

[0015] The hybridoma method has historically been the main driver of antibody discovery, but it is very inefficient for finding rare mAbs. In silico display technologies overcome some of hybridoma's inefficiencies, but mAb panels discovered this way bypass nature's rigorous selection process and as a consequence often suffer from low diversity and poor developability. In fact, the recent burst in rare mAb

discovery (i.e. against HIV, RSV, HMPV, Lassa, and HCMV) comes from high-throughput single B cell approaches that can directly screen patient samples. This high-throughput interrogation of natural immune-reservoirs can identify rare cross-reactive and potent mAbs, which can reduce dosing frequency and, with it, the unit cost of treatment. Molecular engineering can additionally improve both potency and cross-reactivity, and extend mAb half-life. Recent advances in nucleic acid-based mAb delivery have shown promise to decrease both costs and dosing frequencies. Clinical studies have demonstrated that administration of mRNA-encoded antibodies against Chikungunya virus could generate neutralizing titers without toxicity in human patients. These can furthermore be produced at commercial scale with high purity, and formulated to facilitate storage and shipping, which are all important factors in deploying mAbs rapidly and globally.

[0016] Pandemic Response

[0017] As part of our participation in the Pandemic Prevention Platform (P3) program of the US Defense Advanced Research Projects Agency (DARPA), we aim to build an ultra-rapid response pipeline for the discovery and delivery of field-ready therapeutic countermeasures in response to any viral outbreaks. Under P3, in a simulated pandemic scenario, we used microfluidic high-throughput single B cell screening to discover mAbs against pandemic 2009 H1N1 influenza directly from a human donor. We designed machine learning tools to automate high-confidence selection of mAbs during screening, rapidly sequenced hundreds of H1-reactive mAbs, and downselected candidates for preclinical testing using a bioinformatic pipeline and data visualization software. We developed a high-throughput and automated cloning and expression strategy to generate recombinant mAbs for neutralization assays, then administered the top neutralizers encoded as plasmid DNA (DMAbs) in a lethal challenge mouse model and showed 100% protection against a 20xLD₅₀ dose of pandemic H1N1 influenza virus. This demonstration of discovery to successful pre-clinical testing spanned only fifty-five days, showing that DMABs can be effective at neutralizing pandemic viral strains in preclinical disease models, and that de novo discovery and development of therapeutic mAbs as countermeasures against a pandemic virus is possible within a rapid response timeline.

BRIEF SUMMARY OF THE INVENTION

[0018] The present disclosure relates to methods, compositions of matter, and articles of manufacture that may be used in the diagnosis, monitoring, and treatment or prevention of SARS-CoV and SARS-CoV-2-linked diseases such as COVID-19. To that end, provided herein is a large library of fully human antibodies against the spike (S) protein of coronaviruses, e.g., SARS-CoV-2. The sequences of the heavy chain and light chain variable regions (VH and VL) of these antibodies were originally identified from a convalescent patient following infection with a coronavirus, converted to full-length human IgG1 isotype (e.g., IgG1m3 allotype), and the recombinant versions of these antibodies were subsequently produced and characterized as described herein. Therefore, these antibodies are recombinant in nature. The antibodies provided herein each binds to the spike protein of SARS-CoV-2 virus, and some of the antibodies cross-react with the spike protein of one or more other coronaviruses.

[0019] In one embodiment, the library is stored as a collection of electronic DNA sequences of the heavy and light chain variable regions of each member of the library. In another embodiment the library is stored as DNA molecules that encode the heavy and light chain variable regions of each member in the library. In another embodiment, the library is stored as antibody protein molecules for rapid testing. In another embodiment, the library is stored as nucleic acid-encoded versions of the antibody proteins, in mRNA or DNA formats and expression vectors suitable for direct administration to humans as nucleic acids instead of proteins.

[0020] The library of one thousand and forty five anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) (e.g., anti-SARS-CoV-2 antibodies) annotated by internal designations antibodies 258 to 577 and 589 to 1587 are disclosed herein and are set forth in SEQ ID NOs: 1-5316. Nucleic acids encoding the heavy chain variable regions (VH) of these antibodies are set forth in odd numbered sequences in SEQ ID NOs: 1-660, 1321-2750, and 4181-4748 detailed herein. Nucleic acids encoding the light chain variable regions (VL) of these antibodies are set forth in even numbered sequences in SEQ ID NOs: 1-660, 1321-2750, and 4181-4748 detailed herein. Amino acid sequences of the heavy chain variable regions (VH) of these antibodies are set forth in odd numbered sequences in SEQ ID NOs: 661-1320, 2751-4180, and 4749-5316 detailed herein. Amino acid sequences of the light chain variable regions of these antibodies are set forth in even numbered sequences of SEQ ID NOs: 661-1320, 2751-4180, and 4749-5316 detailed herein. As one of skill in the art will appreciate, the four sequences pertaining to any particular antibody will be separated by sequences pertaining to other antibodies. For example, the SEQ ID NOs assigned to the first antibody (designated 258) are SEQ ID NOs: 1, 2, 661 and 662 while the SEQ ID NOs assigned to the second antibody (designated 259) are SEQ ID NOs: 3, 4, 663 and 664 and so on all the way to the one thousand three hundred and twenty ninth antibody (designated 1587), which is assigned SEQ ID NOs: 4747, 4748, 5315, and 5316.

[0021] In some embodiments, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof comprises (a) a heavy chain variable region comprising residues 31-35 for CDR-H1, residues 50-65 for CDR-H2, and residues 95-102 for CDR-H3; and (b) a light chain variable region comprising residues 24-34 for CDR-L1, residues 50-56 for CDR-L2, and residues 89-97 for CDR-L3; wherein the CDR numbering is according to Kabat.

[0022] In some embodiments, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof comprises: (a) a heavy chain variable region comprising residues 26-32 for CDR-H1, residues 50-58 for CDR-H2, and residues 95-102 for CDR-H3; and (b) a light chain variable region comprising residues 24-34 for CDR-L1, residues 50-56 of SEQ ID NO: 62 for CDR-L2, and residues 89-97 of SEQ ID NO: 62 for CDR-L3; wherein the CDR numbering is according to Chothia.

[0023] In some embodiments, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof comprises: (a) a heavy chain variable region comprising residues 30-35 for CDR-H1, residues 47-58 for CDR-H2, and residues 93-101 for CDR-H3; and (b) a light chain variable region comprising residues 30-36

for CDR-L1, residues 46-55 for CDR-L2, and for CDR-L3; wherein the CDR numbering is according to MacCallum.

[0024] In some embodiments, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof comprises a heavy chain variable region having an amino acid sequence that is at least 60%, 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to one of the heavy chain variable region sequences explicitly disclosed in SEQ ID NOs: 660-1320 and 2751-4180 and a light chain variable region having an amino acid sequence that is at least 60%, 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to one of the light chain variable region sequences explicitly disclosed in SEQ ID NOs: 660-1320 and 2751-4180.

[0025] In some embodiments, members of the antibody library are specific to SARS-CoV-2. In some embodiments, the antibodies are known to cross-react against SARS-CoV and SARS-CoV-2 spike proteins. In some embodiments, the antibodies are specific to SARS-CoV spike protein.

[0026] In some embodiments, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof is a neutralizing antibody. In other embodiments, the antibody or antigen-binding fragment thereof is a depleting antibody.

[0027] In some embodiments, the antibody or antigen-binding fragment thereof comprises at least one amino acid substitution. In particular embodiments, the at least one amino acid substitution is a conservative substitution. In some embodiments, the at least one amino acid substitution is a substitution of an amino acid for a non-genetically encoded amino acid or a synthetic amino acid.

[0028] In some embodiments, the antibody or antigen-binding fragment thereof is conjugated to an immunomodulator, a cytokine, a cytotoxic agent, a chemotherapeutic agent, a diagnostic agent, an antiviral agent, an antimicrobial agent, or a drug.

[0029] In some embodiments, the antibody or antigen-binding fragment thereof is formulated as a pharmaceutical composition. In some embodiments, the pharmaceutical composition may comprise one or more pharmaceutically acceptable carriers, diluents, or excipients. In particular embodiments, the antibody or antigen-binding fragment thereof may be conjugated to an immunomodulator, a cytokine, a cytotoxic agent, a chemotherapeutic agent, a diagnostic agent, an antiviral agent, an antimicrobial agent, or a drug prior to formulation.

[0030] The inventions disclosed herein also encompass isolated nucleic acids encoding part or all of the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) and antigen-binding fragments thereof disclosed herein. In some embodiments, the foregoing nucleic acids may be incorporated into a vector. In some embodiments, the foregoing nucleic acids may be incorporated into a host cell or a vector then into a host cell.

[0031] The inventions disclosed herein also encompass methods of identifying a SARS-CoV-2-infected cell comprising contacting a cell with an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof conjugated to a detectable agent and detecting specific binding of the antibody or antigen-binding fragment thereof to the cell.

[0032] The inventions disclosed herein also encompass methods of using an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or antigen-binding fragment thereof

as a test reagent for the development and production of a vaccine (e.g., an inactivated virus) against a SARS-CoV-2 disease (such as COVID-19).

[0033] The inventions disclosed herein also encompass methods of diagnosing a SARS-CoV-2 infection in a patient comprising contacting a sample obtained from a patient with a SARS-CoV-2 antibody or antigen-binding fragment thereof conjugated to a detectable agent and detecting specific binding of the antibody or antigen-binding fragment thereof to a SARS-CoV-2 antigen present in the sample.

[0034] The inventions disclosed herein also encompass methods of treating a SARS-CoV-2-linked disease (SARS) or SARS-CoV-2-linked disease (such as COVID-19) comprising administering to a patient a therapeutically effective amount of a SARS-CoV-2 antibody or antigen-binding fragment thereof. Prior to administration, the antibody or antigen-binding fragment thereof may be conjugated (for example, to an anti-viral agent) or formulated as a pharmaceutical composition.

[0035] The inventions disclosed herein also encompass articles of manufacture useful for diagnosing or treating a SARS-CoV-2-linked disease comprising a receptacle comprising a SARS-CoV-2 antibody or antigen-binding fragment thereof, or antibody conjugate, or pharmaceutical composition as well as instructional materials for using the same to treat or diagnose the SARS-CoV-2-linked disease.

[0036] The inventions disclosed herein also encompass processes for producing a SARS-CoV-2 antibody comprising cultivating a host cell under conditions such that the antibody is expressed and recovering the expressed antibody.

[0037] The foregoing is a summary and thus contains, by necessity, simplifications, generalizations, and omissions of detail; consequently, those skilled in the art will appreciate that the summary is illustrative only and is not intended to be in any way limiting. Other aspects, features, and advantages of the methods, compositions and/or devices and/or other subject matter described herein will become apparent in the teachings set forth herein. The summary is provided to introduce a selection of concepts in a simplified form that are further described below in the Detailed Description. This summary is not intended to identify key features or essential features of the claimed subject matter, nor is it intended to be used as an aid in determining the scope of the claimed subject matter.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0038] FIG. 1 is a heat map generated from epitope binning experiments using CARTERRA® Epitope analysis software. Binding signals were normalized to the Ag-only signal average equivalent to one RU (relative unit). A threshold window (0.9 RU to 1.1 RU) was used to classify analytes into 3 categories, i.e. blockers (analytes with a binding signal under the lower limit threshold), sandwichers (analytes with a binding signal over the higher limit threshold) and ambiguous (analytes with a signal falling between the lower and higher limit thresholds). The software automatically clusters like-behaved mAbs into a heat map. B=Blockers (2 mAbs that block each other and likely compete for the same epitope; medium gray). NB=Non-blockers (2 mAbs that do not block each other and likely

bind 2 different epitopes, or “sandwich”; white). A=Ambiguous (light gray). Self-vs-self interaction is highlighted in dark gray.

[0039] FIG. 2A shows the binding kinetics curve of Fab 555 to SARS-CoV-2 spike protein. FIG. 2B shows representative negative stain electron microscopy images of Fab 555 in complex with the trimeric SARS-CoV-2 spike protein. FIG. 2C shows a three-dimensional reconstruction of the complex of Fab 555 and SARS-CoV-2 spike protein within the class-averaged image density; with the peptide regions corresponding to the epitope information derived from HDXMS experiments labeled dark grey.

[0040] FIG. 3A shows the binding kinetics curve of Fab 447 to SARS-CoV-2 spike protein. FIG. 3B shows representative negative stain electron microscopy images of Fab 447 in complex with the trimeric SARS-CoV-2 spike protein. FIG. 3C shows a three-dimensional reconstruction of the complex of Fab 447 and SARS-CoV-2 spike protein within the class-averaged image density.

[0041] FIG. 4A shows the binding kinetics curve of Fab 483 to SARS-CoV-2 spike protein. FIG. 4B shows representative negative stain electron microscopy images of Fab 483 in complex with the trimeric SARS-CoV-2 spike protein. FIG. 4C shows a three-dimensional reconstruction of the complex of Fab 483 and SARS-CoV-2 spike protein within the class-averaged image density.

[0042] FIG. 5A shows the binding kinetics curve of Fab 419 to SARS-CoV-2 spike protein. FIG. 5B shows representative negative stain electron microscopy images of Fab 419 in complex with the trimeric SARS-CoV-2 spike protein. FIG. 5C shows a three-dimensional reconstruction of the complex of Fab 419 and SARS-CoV-2 spike protein within the class-averaged image density.

[0043] FIG. 6A shows the binding kinetics curve of Fab 388 to SARS-CoV-2 spike protein. FIG. 6B shows representative negative stain electron microscopy images of Fab 388 in complex with the trimeric SARS-CoV-2 spike protein. FIG. 6C shows a three-dimensional reconstruction of the complex of Fab 388 and SARS-CoV-2 spike protein within the class-averaged image density.

[0044] FIG. 7 shows the relative binding site of Fabs 555, 447, 483, 419, 388 on SARS-CoV-2 spike protein.

[0045] FIG. 8A shows the binding kinetics curve of Fab 494 to SARS-CoV-2 spike protein. FIG. 8B shows representative negative stain electron microscopy images of Fab 494 in complex with the trimeric SARS-CoV-2 spike protein. FIG. 8C shows a three-dimensional reconstruction of the complex of Fab 494 and SARS-CoV-2 spike protein within the class-averaged image density.

[0046] FIG. 9A is a ribbon diagram of Fab 555/481/488: RBD domain complex structured as determined by X-ray crystallography. FIG. 9B shows the crystal structure of the RBD-Fab555 complex superimposed with the ACE2 receptor from a structure of the RBD-ACE2 complex (PDB ID: 6M0J). FIG. 9C is a zoomed-in view of key atomic interactions at the interface of the Fab 555 light chain and the RBD of SARS-CoV-2 spike protein. FIG. 9D is a zoomed-in view of key atomic interactions at the interface of the Fab 555 heavy chain and the RBD of SARS-CoV-2 spike protein. FIG. 9E shows the cryo-EM structure of the Fab 555-spike protein complex low-pass filtered to 8 Å resolution and shown at low threshold in order to visualize all three Fabs. FIG. 9F is a high-resolution cryo-EM map of the Fab555-spike protein complex.

[0047] FIGS. 10A-10H show the effect of mAb 555 (aka LY-CoV555) on the viral loads in rhesus macaques challenged with SARS-CoV-2. 24 hours prior to viral challenge, Rhesus macaques were administered varying amounts of LY-CoV555 as a single IV dose. Viral loads in the BAL (FIGS. 10A and 10B), throat swabs (FIGS. 10C and 10D), nasal swabs (FIGS. 10E and 10F) or lung tissue (FIGS. 10G and 10H) were assessed over the course of 6 days post-inoculation by measuring genomic RNA (FIGS. 10A, 10C, 10E, 10G) or subgenomic mRNA (FIGS. 10B, 10D, 10F, 10H) by qRT-PCR. FIGS. 10A-10F: Values represent the mean and standard error of the mean for 3 or 4 animals. FIGS. 10G and 10H: bars represent the mean of 3 or 4 animals. Samples below the lower limit of quantification (LLOQ) were designated a value of % LLOQ. LLOQ=50 copies for genomes or sg mRNA.

DETAILED DESCRIPTION OF THE INVENTION

[0048] While the present invention may be embodied in many different forms, disclosed herein are specific illustrative embodiments thereof that exemplify the principles of the invention. It should be emphasized that the present invention is not limited to the specific embodiments illustrated. Moreover, any section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

[0049] Unless otherwise defined herein, scientific, and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. More specifically, as used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a protein” includes a plurality of proteins; reference to “a cell” includes mixtures of cells, and the like. In addition, ranges provided in the specification and appended claims include both end points and all points between the end points. Therefore, a range of 2.0 to 3.0 includes 2.0, 3.0, and all points between 2.0 and 3.0.

[0050] Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclature used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art.

EXEMPLARY EMBODIMENTS OF THE INVENTION

[0051] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding

fragment thereof that specifically binds to a SARS-CoV-2 antigen, SARS-CoV-2 viral particle, or SARS-CoV-2-infected cell, wherein the antibody or antigen-binding fragment thereof comprises:

- [0052]** (a) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 661 and three CDRs of a light chain variable region set forth as SEQ ID NO: 662; or
- [0053]** (b) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 663 and three CDRs of a light chain variable region set forth as SEQ ID NO: 664; or
- [0054]** (c) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 665 and three CDRs of a light chain variable region set forth as SEQ ID NO: 666; or
- [0055]** (d) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 667 and three CDRs of a light chain variable region set forth as SEQ ID NO: 668; or
- [0056]** (e) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 669 and three CDRs of a light chain variable region set forth as SEQ ID NO: 670; or
- [0057]** (f) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 671 and three CDRs of a light chain variable region set forth as SEQ ID NO: 672; or
- [0058]** (g) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 673 and three CDRs of a light chain variable region set forth as SEQ ID NO: 674; or
- [0059]** (h) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 675 and three CDRs of a light chain variable region set forth as SEQ ID NO: 676; or
- [0060]** (i) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 677 and three CDRs of a light chain variable region set forth as SEQ ID NO: 678; or
- [0061]** (j) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 679 and three CDRs of a light chain variable region set forth as SEQ ID NO: 680; or
- [0062]** (k) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 681 and three CDRs of a light chain variable region set forth as SEQ ID NO: 682; or
- [0063]** (l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 683 and three CDRs of a light chain variable region set forth as SEQ ID NO: 684; or
- [0064]** (m) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 685 and three CDRs of a light chain variable region set forth as SEQ ID NO: 686; or
- [0065]** (n) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 687 and three CDRs of a light chain variable region set forth as SEQ ID NO: 688; or
- [0066]** (o) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 689 and three CDRs of a light chain variable region set forth as SEQ ID NO: 690; or
- [0067]** (p) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 691 and three CDRs of a light chain variable region set forth as SEQ ID NO: 692; or
- [0068]** (q) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 693 and three CDRs of a light chain variable region set forth as SEQ ID NO: 694; or
- [0069]** (r) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 695 and three CDRs of a light chain variable region set forth as SEQ ID NO: 696; or
- [0070]** (s) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 697 and three CDRs of a light chain variable region set forth as SEQ ID NO: 698; or
- [0071]** (t) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 699 and three CDRs of a light chain variable region set forth as SEQ ID NO: 700; or

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible][illegible]

[illegible]

[illegible]

[illegible]

[illegible][illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible][illegible]

[illegible]

[illegible]

[illegible][illegible]

[illegible]

[illegible]

[illegible][illegible]

[illegible][illegible]

[1373] (v-l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5283 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5284; or

[1374] (w-l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5285 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5286; or

[1375] (x-l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5287 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5288; or

[1376] (y-l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5289 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5290; or

[1377] (z-l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5291 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5292; or

[1378] (a-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5293 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5294; or

[1379] (b-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5295 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5296; or

[1380] (c-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5297 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5298; or

[1381] (d-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5299 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5300; or

[1382] (e-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5301 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5302; or

[1383] (f-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5303 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5304; or

[1384] (g-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5305 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5306; or

[1385] (h-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5307 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5308; or

[1386] (i-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5309 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5310; or

[1387] (j-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5311 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5312; or

[1388] (k-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5313 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5314; or

[1389] (l-li) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 5315 and three CDRs of a light chain variable region set forth as SEQ ID NO: 5316.

[1390] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof, wherein the antibody or fragment thereof comprises: (a) CDR-H1 comprising residues 31-35 of the heavy chain variable region (VH), CDR-H2 comprising residues 50-65 of the VH, and CDR-H3 comprising residues 95-102 of the VH; and (b) CDR-L1 comprising residues 24-34 of the light chain variable region (VL), CDR-L2 comprising residues 50-56 of the VL, and CDR-L3 comprising residues 89-97 of the VL; and wherein the CDR numbering is according to Kabat.

[1391] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding

fragment thereof, wherein the antibody or fragment thereof comprises: (a) CDR-H1 comprising residues 26-32 of the VH, CDR-H2 comprising residues 50-58 of the VH, and CDR-H3 comprising residues 95-102 of the VH; and (b) CDR-L1 comprising residues 24-34 of the VL, CDR-L2 comprising residues 50-56 of the VL, and CDR-L3 comprising residues 89-97 of the VL; and wherein the CDR numbering is according to Chothia.

[1392] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof, wherein the antibody or fragment thereof comprises: (a) CDR-H1 comprising residues 30-35 of the VH, CDR-H2 comprising residues 47-58 of the VH, and CDR-H3 comprising residues 93-101 of the VH; and (b) CDR-L1 comprising residues 30-36 of the VL, CDR-L2 comprising residues 46-55 of the VL, and CDR-L3 comprising the residues 89-96 of the VL; and wherein the CDR numbering is according to MacCallum.

[1393] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof, wherein the antibody or fragment thereof comprises a heavy chain variable region having an amino acid sequence that is at least 60% identical to a heavy chain variable region sequence comprising three CDRs of the heavy chain variable region set forth in odd-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316 and a light chain variable region having an amino acid sequence that is at least 60% identical to one of the light chain variable region sequences comprising three CDRs of a corresponding light chain variable region set forth in even-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316.

[1394] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof, wherein the antibody or fragment thereof comprises (a) the VH set forth as SEQ ID NO: 1225 and the VL set forth as SEQ ID NO: 1226; or (b) the VH set forth as SEQ ID NO: 729 and the VL set forth as SEQ ID NO: 730; or (c) the VH set forth as SEQ ID NO: 763 and the VL set forth as SEQ ID NO: 764; or (d) the VH set forth as SEQ ID NO: 873 and the VL set forth as SEQ ID NO: 874; or (e) the VH set forth as SEQ ID NO: 891 and the VL set forth as SEQ ID NO: 892; or (f) the VH set forth as SEQ ID NO: 921 and the VL set forth as SEQ ID NO: 922; or (g) the VH set forth as SEQ ID NO: 961 and the VL set forth as SEQ ID NO: 962; or (h) the VH set forth as SEQ ID NO: 973 and the VL set forth as SEQ ID NO: 974; or (i) the VH set forth as SEQ ID NO: 979 and the VL set forth as SEQ ID NO: 980; or (j) the VH set forth as SEQ ID NO: 983 and the VL set forth as SEQ ID NO: 984; or (k) the VH set forth as SEQ ID NO: 1029 and the VL set forth as SEQ ID NO: 1030; or (l) the VH set forth as SEQ ID NO: 1035 and the VL set forth as SEQ ID NO: 1036; or (m) the VH set forth as SEQ ID NO: 1039 and the VL set forth as SEQ ID NO: 1040; or (n) the VH set forth as SEQ ID NO: 1069 and the VL set forth as SEQ ID NO: 1070; or (o) the VH set forth as SEQ ID NO: 1103 and the VL set forth as SEQ ID NO: 1104; or (p) the VH set forth as SEQ ID NO: 1107 and the VL set forth as SEQ ID NO: 1108; or (q) the VH set forth as SEQ ID NO: 1111 and the VL set forth as SEQ ID NO: 1112; or (r) the VH set forth as SEQ ID NO: 1121 and the VL set forth as SEQ ID NO: 1122; or (s) the VH set forth as SEQ ID NO: 1133 and the VL set forth as SEQ ID NO: 1134; or (t) the VH set forth as SEQ ID NO: 1157 and the VL set forth as SEQ ID NO: 1158; or (u) the VH set forth as SEQ ID NO: 1243 and

the VL set forth as SEQ ID NO: 1244; or (v) the VH set forth as SEQ ID NO: 1251 and the VL set forth as SEQ ID NO: 1252; or (w) the VH set forth as SEQ ID NO: 1255 and the VL set forth as SEQ ID NO: 1256; or (x) the VH set forth as SEQ ID NO: 1269 and the VL set forth as SEQ ID NO: 1270.

[1395] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof, wherein the antibody or fragment thereof comprises: (a) a heavy chain comprising SEQ ID NO: 5319 and a light chain comprising SEQ ID NO: 5320; or (b) a heavy chain comprising SEQ ID NO: 5321 and a light chain comprising SEQ ID NO: 5322; or (c) a heavy chain comprising SEQ ID NO: 5323 and a light chain comprising SEQ ID NO: 5324; or (d) a heavy chain comprising SEQ ID NO: 5325 and a light chain comprising SEQ ID NO: 5326; or (e) a heavy chain comprising SEQ ID NO: 5327 and a light chain comprising SEQ ID NO: 5328; or (f) a heavy chain comprising SEQ ID NO: 5329 and a light chain comprising SEQ ID NO: 5330; or (g) a heavy chain comprising SEQ ID NO: 5331 and a light chain comprising SEQ ID NO: 5332; or (h) a heavy chain comprising SEQ ID NO: 5333 and a light chain comprising SEQ ID NO: 5334; or (i) a heavy chain comprising SEQ ID NO: 5335 and a light chain comprising SEQ ID NO: 5336; or (j) a heavy chain comprising SEQ ID NO: 5337 and a light chain comprising SEQ ID NO: 5338; or (k) a heavy chain comprising SEQ ID NO: 5339 and a light chain comprising SEQ ID NO: 5340; or (l) a heavy chain comprising SEQ ID NO: 5341 and a light chain comprising SEQ ID NO: 5342; or (m) a heavy chain comprising SEQ ID NO: 5343 and a light chain comprising SEQ ID NO: 5344; or (n) a heavy chain comprising SEQ ID NO: 5345 and a light chain comprising SEQ ID NO: 5346; or (o) a heavy chain comprising SEQ ID NO: 5347 and a light chain comprising SEQ ID NO: 5348; (p) a heavy chain comprising SEQ ID NO: 5349 and a light chain comprising SEQ ID NO: 5350; or (q) a heavy chain comprising SEQ ID NO: 5351 and a light chain comprising SEQ ID NO: 5352; or (r) a heavy chain comprising SEQ ID NO: 5353 and a light chain comprising SEQ ID NO: 5354; or (s) a heavy chain comprising SEQ ID NO: 5355 and a light chain comprising SEQ ID NO: 5356; or (t) a heavy chain comprising SEQ ID NO: 5357 and a light chain comprising SEQ ID NO: 5358; or (u) a heavy chain comprising SEQ ID NO: 5359 and a light chain comprising SEQ ID NO: 5360; or (v) a heavy chain comprising SEQ ID NO: 5361 and a light chain comprising SEQ ID NO: 5362; or (w) a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364; or (x) a heavy chain comprising SEQ ID NO: 5365 and a light chain comprising SEQ ID NO: 5366.

[1396] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof, wherein the antibody or fragment thereof comprises a heavy chain variable region having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316 and a light chain variable region having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316.

[1397] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding

fragment thereof described in the preceding paragraphs of this section that specifically binds to a SARS-CoV spike protein.

[1398] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that specifically binds to a SARS-CoV-2 spike protein.

[1399] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that is cross-reactive with the SARS-CoV spike protein and SARS-CoV-2 spike protein.

[1400] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that is a neutralizing antibody.

[1401] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that is a depleting antibody.

[1402] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that binds an epitope comprising one or more of the following residues of the SARS-COV-2 spike protein: Y351, Y449, N450, L452, L455, F456, T470, I472, N481, G482, V483, E484, G485, F486, Y489, F490, L492, Q493, S494, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In particular embodiments, the epitope comprises two or more of the listed residues, three or more of the listed residues, or five or more of the listed residues.

[1403] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that binds an epitope comprising one or more of the following residues of the SARS-COV-2 spike protein: R403, D405, R408, Q409, T415, G416, K417, D420, Y421, Y453, L455, F456, R457, K458, S459, N460, Y473, Q474, A475, G476, F486, N487, Y489, Q493, S494, Y495, G496, Q498, T500, N501, G502, Y505, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In particular embodiments, the epitope comprises two or more of the listed residues, three or more of the listed residues, or five or more of the listed residues.

[1404] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that binds the SARS-CoV-2 spike protein, wherein the antibody binds an epitope comprising one or more of the following residues of SARS-COV-2 spike protein: R403, T415, G416, K417, D420, Y421, L455, F456, R457, K458, S459, N460, Y473, Q474, A475, G476, S477, F486, N487, Y489, N501, G502, Y505, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In particular embodiments, the epitope comprises two or more of the listed residues, three or more of the listed residues, or five or more of the listed residues.

[1405] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that has an IgG1 isotype.

[1406] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding

fragment thereof described in the preceding paragraphs of this section that comprises an Fc region comprising N-glycoside-linked sugar chains bound to the Fc region, wherein the sugar chains do not contain fucose.

[1407] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that comprises at least one amino acid substitution. In particular embodiments, the at least one amino acid substitution is a conservative substitution. In particular embodiments, the at least one amino acid substitution is a substitution of an amino acid for a non-genetically encoded amino acid or a synthetic amino acid.

[1408] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that is conjugated to an immunomodulator, a cytokine, a cytotoxic agent, a chemotherapeutic agent, a diagnostic agent, an antiviral agent, an antimicrobial agent, or a drug.

[1409] In particular embodiments, the inventions disclosed herein encompass an antibody conjugate comprising an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that is conjugated to an immunomodulator, a cytokine, a cytotoxic agent, a chemotherapeutic agent, a diagnostic agent, an antiviral agent, an antimicrobial agent, or a drug.

[1410] In particular embodiments, the inventions disclosed herein encompass a pharmaceutical composition comprising an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section or an antibody conjugate comprising the same, and one or more pharmaceutically acceptable carriers, diluents, or excipients. In particular embodiments, the pharmaceutical composition comprises at least one additional antibody that binds the SARS-CoV-2 spike protein. In particular embodiments, the pharmaceutical composition comprises histidine, sodium chloride, sucrose, and polysorbate 80. In particular embodiments, the pharmaceutical composition has a pH of about 6.0. In particular embodiments, the pharmaceutical composition comprises 5 mM histidine, 50 mM NaCl, 6% sucrose, and 0.05% polysorbate 80 and has a pH of about 6.0.

[1411] In particular embodiments, the concentration of the antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section that is in the pharmaceutical composition is about 35 mg/mL to about 125 mg/mL. In particular embodiments, the concentration of the antibody or antigen-binding fragment thereof in the pharmaceutical composition is about 35 mg/mL or about 125 mg/mL.

[1412] In particular embodiments, the inventions disclosed herein encompass an isolated nucleic acid encoding a heavy chain variable region having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316. In particular embodiments, the inventions disclosed herein encompass an isolated nucleic acid encoding a light chain variable region having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316. In particular embodiments, the inventions disclosed herein encompass an isolated nucleic acid encoding a heavy chain having an amino acid sequence that is identical to the heavy chain variable region set forth in

odd-numbered SEQ ID NOs 531-5366. In particular embodiments, the inventions disclosed herein encompass an isolated nucleic acid encoding a light chain having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 531-5366.

[1413] In particular embodiments, the inventions disclosed herein encompass a vector comprising at least one of (a) an isolated nucleic acid encoding a heavy chain variable region having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316; and (b) an isolated nucleic acid encoding a light chain variable region having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316. In particular embodiments, the inventions disclosed herein encompass a vector comprising at least one of (a) an isolated nucleic acid encoding a heavy chain having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 531-5366; and (b) an isolated nucleic acid encoding a light chain having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 531-5366.

[1414] In particular embodiments, the inventions disclosed herein encompass a host cell comprising at least one of (a) an isolated nucleic acid encoding a heavy chain variable region having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316; and (b) an isolated nucleic acid encoding a light chain variable region having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316. In particular embodiments, the inventions disclosed herein encompass a host cell comprising at least one of (a) an isolated nucleic acid encoding a heavy chain having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 531-5366; and (b) an isolated nucleic acid encoding a light chain having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 531-5366.

[1415] In particular embodiments, the inventions disclosed herein encompass a host cell comprising a vector comprising at least one of (a) an isolated nucleic acid encoding a heavy chain variable region having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316; and (b) an isolated nucleic acid encoding a light chain variable region having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 661-1320, 2751-4180, and 4749-5316. In particular embodiments, the inventions disclosed herein encompass a host cell comprising a vector comprising at least one of (a) an isolated nucleic acid encoding a heavy chain having an amino acid sequence that is identical to the heavy chain variable region set forth in odd-numbered SEQ ID NOs 531-5366; and (b) an isolated nucleic acid encoding a light chain having an amino acid sequence that is identical to the light chain variable region set forth in even-numbered SEQ ID NOs 531-5366.

[1416] In particular embodiments, the inventions disclosed herein encompass a process for producing an antibody or antigen-binding fragment thereof described in the

preceding paragraphs of this section comprising (a) culturing the host cell described in the preceding paragraphs under conditions such that the antibody or antigen-binding fragment thereof is expressed; and (b) recovering the expressed antibody or antigen-binding fragment thereof.

[1417] In particular embodiments, the inventions disclosed herein encompass an article of manufacture useful for diagnosing or treating a SARS-CoV or SARS-CoV-2-linked disease comprising a receptacle comprising the antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section, or an antibody conjugate comprising the same, or a pharmaceutical composition comprising the same and instructional materials for using the same to treat or diagnose the SARS-CoV or SARS-CoV-2-linked disease.

[1418] In particular embodiments, the inventions disclosed herein encompass a method of identifying a SARS-CoV or SARS-CoV-2-infected cell comprising: (a) contacting a cell with the an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section, which is conjugated to a detectable agent; and (b) detecting specific binding of the antibody or antigen-binding fragment thereof to the cell.

[1419] In particular embodiments, the inventions disclosed herein encompass a method of diagnosing a SARS-CoV or SARS-CoV-2 infection in a patient comprising: (a) contacting a sample obtained from a patient with the antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section, which is conjugated to a detectable agent; and (b) detecting specific binding of the antibody or antigen-binding fragment thereof to a SARS-CoV or SARS-CoV-2 antigen present in the sample.

[1420] In particular embodiments, the inventions disclosed herein encompass a method of treating or preventing a SARS-CoV or SARS-CoV-2-linked disease comprising administering to a patient a therapeutically effective amount of the antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section, or an antibody conjugate comprising the same, or a pharmaceutical composition comprising the same.

[1421] In particular embodiments, the inventions disclosed herein encompass a method of preventing or treating COVID-19 comprising: (a) contacting a sample obtained from a patient with an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section, which is conjugated to a detectable agent; (b) detecting specific binding of the antibody or antigen-binding fragment thereof to a SARS-CoV-2 antigen present in the sample; and (c) administering to the patient a therapeutically effective amount of the an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section or a pharmaceutical composition comprising the same.

[1422] In particular embodiments, the inventions disclosed herein encompass methods of prophylaxis and therapy described in this section wherein the antibody or antigen-binding fragment thereof is administered or administrable to the patient intravenously or subcutaneously at about 700 mg to about 7000 mg. In particular embodiments, the antibody is administered to the patient intravenously or subcutaneously at about 700 mg. In some embodiments, in methods of prophylaxis and therapy described in this section, the antibody or antigen-binding fragment thereof is administered or administrable to the patient intravenously or subcutaneously at about 35 mg to about 700 mg.

[1423] In particular embodiments, the inventions disclosed herein encompass methods of prophylaxis and therapy described in this section wherein the method further comprises administering to the patient another antibody or antigen-binding fragment thereof that binds the SARS-CoV-2 S protein.

[1424] In particular embodiments, the inventions disclosed herein encompass methods of prophylaxis and therapy described in this section wherein the patient has mild to moderate COVID-19. In particular embodiments, the inventions disclosed herein encompass methods of prophylaxis and therapy described in this section wherein the patient is at risk for contracting COVID-19. In particular embodiments, the inventions disclosed herein encompass methods of prophylaxis and therapy described in this section wherein the patient is at high risk for progressing to severe COVID-19 or hospitalization.

[1425] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section or a pharmaceutical composition comprising the same for use in the diagnosis, theragnosis, treatment and/or prophylaxis of a SARS-CoV or SARS-CoV-2-related disease or symptom.

[1426] In particular embodiments, the inventions disclosed herein encompass an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section or a pharmaceutical composition comprising the same for use in the manufacture of a medicament for diagnosis, theragnosis, treatment and/or prophylaxis of a SARS-CoV or SARS-CoV-2-related disease or symptom.

[1427] In particular embodiments, the inventions disclosed herein encompass a method of testing an anti-coronavirus vaccine, comprising (a) contacting a sample of an anti-coronavirus vaccine with an antibody or antigen-binding fragment thereof described in the preceding paragraphs of this section, conjugated to a detectable agent; and (b) detecting specific binding of the antibody or antigen-binding fragment thereof to the anti-coronavirus vaccine present in the sample; wherein the anti-coronavirus vaccine comprises a coronavirus subunit or fragment thereof.

[1428] Anti-Coronavirus Antibodies (e.g., Anti-SARS-CoV-2 Antibodies)

[1429] The present disclosure is directed to anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) and their use in the diagnosis, theragnosis, treatment and/or prophylaxis of SARS-CoV and SARS-CoV-2 infections and related symptoms (i.e., SARS and COVID-19 coronavirus disease). As used herein, the term “anti-coronavirus antibody(ies)” describes antibodies that specifically recognize, bind to, or otherwise associate with one or more antigens present on the SARS-COV and/or SARS-CoV-2 spike protein (i.e., antibodies that are specific for one or the other protein or that are cross-reactive are encompassed). Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be used alone or in conjunction with a wide variety of other compounds or biological response modifiers. In other selected embodiments, two or more discrete anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be used in combination to provide enhanced therapeutic effects or may be used to fabricate multispecific constructs. Accordingly, provided are also compositions comprising two, three, or more discrete anti-coronavirus

antibodies. The anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) herein are recombinant in nature.

[1430] Provided herein are antibodies or antigen-binding fragments thereof that specifically bind to a SARS-CoV-2 antigen, SARS-CoV-2 viral particle, or SARS-CoV-2-infected cell (“anti-SARS-CoV-2 antibodies or antigen binding fragments”). In some embodiments, the anti-SARS-CoV-2 antibodies or antigen binding fragments specifically bind the spike (S) protein of SARS-CoV-2. The terms “bind” and “binds” as used herein are intended to mean, unless indicated otherwise, the ability of a protein or molecule to form a chemical bond or attractive interaction with another protein or molecule, which results in proximity of the two proteins or molecules as determined by common methods known in the art.

[1431] The cryo-electron microscopy structure of the SARS-CoV-2 S protein has been described recently, and shows the SARS-CoV-2 S protein is a trimeric class I fusion protein that exists in a metastable pre-fusion conformation and undergoes a substantial structural rearrangement to fuse the viral membrane with the host cell membrane (Wrapp et al., *Science* 367, 1260-1263, 2020). This process is triggered when the receptor-binding domain (RBD) of the S1 subunit of the SARS-CoV-2 S protein binds to the host cell receptor angiotensin-converting enzyme 2 (ACE2); and receptor binding destabilizes the pre-fusion trimer, resulting in shedding of the S1 subunit and transition of the S2 subunit to a stable post-fusion conformation (Wrapp et al., *Science* 367, 1260-1263, 2020).

[1432] The amino acid sequence of SARS-CoV-2 spike (S) protein has been described before, for example, GenBank Accession No: YP_009724390.1 provides such an exemplary sequence, as shown below:

(SEQ ID NO: 5317)

```
MFVFLVLLPLVSSQCVNLTRTQLPPAYTNSFTRGVYPDKVFRSSVLHS
TQDLFLPFFSNVTWFHAIHVGSTNGTKRFDNPVLPFNDGVYFASTEKSN
IRGWIFGTTLDSTQLSLIVNNATNVVVKVCEFCNDPFLGVYHKNNK
SWMESEFRVYSSANNCTFEYVSQPPLMDLEGKQGNFKNLREFVFNIDGY
FKIYSKHPTINLVRDLPGQFSALEPLVDLPIGINITRFQTLALHRSYLT
PGDSSSGWTAGAAAYVGYLQPRFTLLKYNENGTITDAVDCALDPLSETK
CTLSFTVEKGIYQTSNFRVQPTESIVRFPNITNLCPPGEVFNATRFASV
YAWNKRKISNCVADYSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSF
VIRGDEVQRQIAPGQTGKIADYNYKLDDFTGCVIAWNSNNLDSKVGNYN
YLRLFRKSNLKPFRDISTEIQAGSTPCNGVEGFNCFYPLQSYGFQPT
NGVGYQPYRVVLSFELLHAPATVCGPKKSTNLVKNKCNFNFNGLTGTG
VLTESNKKFLPFQFGRIADTTDAVRDPQTLIILDITPCSFGGVSVITP
GTNTSNQVAVLYQDVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCL
IGAETHVNSYECIDIPIGAGICASYQTQNSPRRARSVASQSIIAYTMSLG
AENSVAYSNNSIAIPTNFTISVTEILPVSMTKTSVDCTMYICGDSTEC
NLLQYGSFCTQLNRALTGIAVEQDKNTQEVFAQVKQIYKTPPIKDFGGF
NFSQILPDPSKPSKRSFIEDLLFNKVTLDAGFIKQYGDCLGDIARDLI
```

-continued

```
CAQKFNGLTVLPPLLTDEMIQYTSALLAGTITSGWTFGAGAALQIPFAM
QMAYRFNGIGVTQNVLYENQKLIANQFNSAIGKIQDSLSSSTASALGKLQD
VVNQNAQALNTLVKQLSSNFGAISSVLNDILSRDLKVEAEVQIDRLITGR
LQSLQTYVTQQLIRAAEIRASANLAATKMECVLGQSKRVDFCGKGYHLM
SFPQSAAPHGVFLVHTVYVPAQEKNTTAPAICHDKGAHFPREGVFSVNGT
HWFVTQRNFYEPQIITDNTFVSGNCDVVIGIVNNTVYDPLQPELDSFKE
ELDKYFKNHTSPDVLGDISGINASVVNIQKEIDRLNEVAKNLNESLIKL
QELGKYEQYIKWPWYIWLGFIAGLIAIVMVTIMLCMTSCCSCCLKGCCSC
GSCCKFDEDDSEPVLLKGVKLHYT.
```

[1433] The receptor-binding domain (RBD) comprises amino acids 329 to 520 of the SARS-CoV-2 S protein, e.g., amino acids 329 to 520 of SEQ ID NO: 5317, or a variant comprising one or more mutations; the N-terminal domain (NTD) comprises amino acids 16-328 of the SARS-CoV-2 S protein, e.g., amino acids 16-328 of SEQ ID NO: 5317, or a variant comprising one or more mutations; S2 domain comprises amino acids 686-1213 of the SARS-CoV-2 S protein, e.g., amino acids 686-1213 of SEQ ID NO: 5317, or a variant comprising one or more mutations (see Wrapp et al., *Science* 367, 1260-1263, 2020).

[1434] The amino acid sequence of SARS-CoV spike protein has also been described before, for example, UniProt ID: P59594, as shown below:

(SEQ ID NO: 5318)

```
MFIFLLFLTLTSGSDLDRCTTFDDVQAPNYTQHTSSMRGVYPDEIFRSD
TLYLTDQLFLPFPYSNVTGFHTINHTFGNPVIFPKDGIYFAATEKSNVVRG
WVFGDTMNNKSQSVIINNSTNVVIRACNPFELCDNPPFAVSKPMGTQHTMT
IFDNAFNCTFEYISDAFSLDVSEKSGNFKHLREFVFNKDGFLVYKGYQ
PIDVVRDLPSGFNTLKPFIKPLGINITNFRILATFSPAQDIWGTSA
YFVGYLKPFTFMLKYDENGITIDAVIDCSQNPLAELKCSVKSFEIDKGIYQ
TSNFRVVPBGDVVRFPNITNLCPPGEVFNATKFPSSVYAWERKKISNCVAD
YSVLVNSTFFSTFKCYGVSATKLNDLCFSNVYADSFVVGDDVRQIAPGQ
TGVIADYNYKLDDFMGCVLAWNTRNIDATSTGNVNYKYRLRHGKLRF
ERDISNVFSPDGKPCPTPALNCYWPLNDYGFYTTTGIGYQPYRVVLSF
ELLNAPATVCGPKLSTDLIKQCVNFNENGLTGTGLTPSSKRFQPFQF
GRDVSDFDTSVRDPKTSEILDISPCSFGGVSVITPGTNASSEVAVLYQDV
NCTDVSTAIHADQLTPAWRIYSTGNVFTQAGCLIGAETHVDSYECIDIP
IGAGICASYHTVSLRSTSQKSIVAYTMSLGADSSIAYSNNTIAIPTNFS
ISITTEVMPVSMKTSVDCNMYICGDSTECANLLQYGSFCTQLNRALSG
IAAEQDRNTREVFAQVKQMYKPTPLKYFGGFNSQILPDPLKPTKRSFIE
DLLFNKVTLDAGFMKQYGECLGDINARDLCAQKFNGLTVLPPLLTDDM
IAAYTAALVSGTATAGWTFGAGAALQIPFAMQMAYRFNGIGVTQNVLYEN
QKQIANQFNKAIQSIESLTTTSTALGKLQDVVNQNAQALNTLVKQLSSN
FGAISSVLNDILSRDLKVEAEVQIDRLITGRQLSLQTYVTQQLIRAAEIR
```

-continued

ASANLAATKMSECVLGGQSKRVDFCGKGYHLMSFPQAAPHGVVFLHVTYVP
SQERNFTTAPAIHECKAYFPREGVVFVFNGTSWFITQRNFFSPQIITTDN
TFVSGNCDVVIGIINNTVYDPLQPELKSFKELDKYFKNHTSPDVLGDGI
SGINASVVNIQKEIDRLNEVAKNLNLNLSLIDLQELGKYEQYIKWPVYVWL
FIAGLIAIVMTILLCCMTSCCCLKGACSCGCKCFDEDDSEPVLGKVK
LHYT.

[1435] In some embodiments, provided herein are anti-SARS-CoV-2 antibodies having internal designations 258 to 577 and 589 to 1587 as described herein. In some embodiments, provided herein are anti-SARS-CoV-2 antibodies or antigen binding fragments that comprise the VH and/or the VL of any one of antibodies 258 to 577 and 589 to 1587. In some embodiments, provided herein are anti-SARS-CoV-2 antibodies or antigen binding fragments that comprise a VH domain having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% identity to the VH domain of any one of antibodies 258 to 577 and 589 to 1587, or a VL domain having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% identity to the VH domain of any one of antibodies 258 to 577 and 589 to 1587. In some embodiments, provided herein are anti-SARS-CoV-2 antibodies or antigen binding fragments that comprise a VH domain having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% identity to the VH domain of any one of antibodies 258 to 577 and 589 to 1587, and a VL domain having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% identity to the VH domain of any one of antibodies 258 to 577 and 589 to 1587, preferably wherein these homologous VH and VL domains correspond to those of the same antibody.

[1436] In some embodiments, provided herein are anti-SARS-CoV-2 antibodies or antigen binding fragments that comprise a VH domain comprising the same three CDRs as comprised in the VH domain of any one of antibodies 258 to 577 and 589 to 1587 and/or comprises a VL domain comprising the same three CDRs as the VL domain of any one of antibodies 258 to 577 and 589 to 1587, wherein the CDRs are defined by Kabat, Chothia or MacCallum numbering.

[1437] In some embodiments, the anti-SARS-CoV-2 antibodies or antigen binding fragments provided herein are neutralizing antibodies. In particular embodiments, neutralizing anti-SARS-CoV-2 antibodies or antigen binding fragments comprise: (1) three CDRs in the heavy chain variable region (VH) set forth as SEQ ID NO: 729 and three CDRs in the light chain variable region (VL) set forth as SEQ ID NO: 730; (2) three CDRs in the VH set forth as SEQ ID NO: 763 and three CDRs in the VL set forth as SEQ ID NO: 764; (3) three CDRs in the VH set forth as SEQ ID NO: 873 and three CDRs in the VL set forth as SEQ ID NO: 874; (4) three CDRs in the VH set forth as SEQ ID NO: 891 and three CDRs in the VL set forth as SEQ ID NO: 892; (5) three CDRs in the VH set forth as SEQ ID NO: 921 and three CDRs in the VL set forth as SEQ ID NO: 922; (6) three CDRs in the VH set forth as SEQ ID NO: 961 and three CDRs in the VL set forth as SEQ ID NO: 962; (7) three CDRs in the VH set forth as SEQ ID NO: 973 and three CDRs in the VL set forth as SEQ ID NO: 974; (8) three

CDRs in the VH set forth as SEQ ID NO: 979 and three CDRs in the VL set forth as SEQ ID NO: 980; (9) three CDRs in the VH set forth as SEQ ID NO: 983 and three CDRs in the VL set forth as SEQ ID NO: 984; (10) three CDRs in the VH set forth as SEQ ID NO: 1029 and three CDRs in the VL set forth as SEQ ID NO: 1030; (11) three CDRs in the VH set forth as SEQ ID NO: 1035 and three CDRs in the VL set forth as SEQ ID NO: 1036; (12) three CDRs in the VH set forth as SEQ ID NO: 1039 and three CDRs in the VL set forth as SEQ ID NO: 1040; (13) three CDRs in the VH set forth as SEQ ID NO: 1069 and three CDRs in the VL set forth as SEQ ID NO: 1070; (14) three CDRs in the VH set forth as SEQ ID NO: 1103 and three CDRs in the VL set forth as SEQ ID NO: 1104; (15) three CDRs in the VH set forth as SEQ ID NO: 1107 and three CDRs in the VL set forth as SEQ ID NO: 1108; (16) three CDRs in the VH set forth as SEQ ID NO: 1111 and three CDRs in the VL set forth as SEQ ID NO: 1112; (17) three CDRs in the VH set forth as SEQ ID NO: 1121 and three CDRs in the VL set forth as SEQ ID NO: 1122; (18) three CDRs in the VH set forth as SEQ ID NO: 1133 and three CDRs in the VL set forth as SEQ ID NO: 1134; (19) three CDRs in the VH set forth as SEQ ID NO: 1157 and three CDRs in the VL set forth as SEQ ID NO: 1158; (20) three CDRs in the VH set forth as SEQ ID NO: 1225 and three CDRs in the VL set forth as SEQ ID NO: 1226; (21) three CDRs in the VH set forth as SEQ ID NO: 1243 and three CDRs in the VL set forth as SEQ ID NO: 1244; (22) three CDRs in the VH set forth as SEQ ID NO: 1251 and three CDRs in the VL set forth as SEQ ID NO: 1252; (23) three CDRs in the VH set forth as SEQ ID NO: 1255 and three CDRs in the VL set forth as SEQ ID NO: 1256; or (24) three CDRs in the VH set forth as SEQ ID NO: 1269 and three CDRs in the VL set forth as SEQ ID NO: 1270; wherein the CDRs are defined by Kabat, Chothia or MacCallum numbering.

[1438] In other embodiments, neutralizing anti-SARS-CoV-2 antibodies or antigen binding fragments comprise: (1) the VH set forth as SEQ ID NO: 729 and the VL set forth as SEQ ID NO: 730; (2) the VH set forth as SEQ ID NO: 763 and the VL set forth as SEQ ID NO: 764; (3) the VH set forth as SEQ ID NO: 873 and the VL set forth as SEQ ID NO: 874; (4) the VH set forth as SEQ ID NO: 891 and the VL set forth as SEQ ID NO: 892; (5) the VH set forth as SEQ ID NO: 921 and the VL set forth as SEQ ID NO: 922; (6) the VH set forth as SEQ ID NO: 961 and the VL set forth as SEQ ID NO: 962; (7) the VH set forth as SEQ ID NO: 973 and the VL set forth as SEQ ID NO: 974; (8) the VH set forth as SEQ ID NO: 979 and the VL set forth as SEQ ID NO: 980; (9) the VH set forth as SEQ ID NO: 983 and the VL set forth as SEQ ID NO: 984; (10) the VH set forth as SEQ ID NO: 1029 and the VL set forth as SEQ ID NO: 1030; (11) the VH set forth as SEQ ID NO: 1035 and the VL set forth as SEQ ID NO: 1036; (12) the VH set forth as SEQ ID NO: 1039 and the VL set forth as SEQ ID NO: 1040; (13) the VH set forth as SEQ ID NO: 1069 and the VL set forth as SEQ ID NO: 1070; (14) the VH set forth as SEQ ID NO: 1103 and the VL set forth as SEQ ID NO: 1104; (15) the VH set forth as SEQ ID NO: 1107 and the VL set forth as SEQ ID NO: 1108; (16) the VH set forth as SEQ ID NO: 1111 and the VL set forth as SEQ ID NO: 1112; (17) the VH set forth as SEQ ID NO: 1121 and the VL set forth as SEQ ID NO: 1122; (18) the VH set forth as SEQ ID NO: 1133 and the VL set forth as SEQ ID NO: 1134; (19) the VH set forth as SEQ ID NO:

1157 and the VL set forth as SEQ ID NO: 1158; (20) the VH set forth as SEQ ID NO: 1225 and the VL set forth as SEQ ID NO: 1226; (21) the VH set forth as SEQ ID NO: 1243 and the VL set forth as SEQ ID NO: 1244; (22) the VH set forth as SEQ ID NO: 1251 and the VL set forth as SEQ ID NO: 1252; (23) the VH set forth as SEQ ID NO: 1255 and the VL set forth as SEQ ID NO: 1256; or (24) the VH set forth as SEQ ID NO: 1269 and the VL set forth as SEQ ID NO: 1270.

[1439] In some embodiments, the anti-SARS-CoV-2 antibodies or antigen binding fragments provided herein are antibodies that block SARS-CoV-2 binding to ACE2, i.e., ACE2 blockers. In particular embodiments, ACE2-blocking anti-SARS-CoV-2 antibodies or antigen binding fragments comprise: (i) three CDRs in the heavy chain variable region (VH) set forth as SEQ ID NO: 729 and three CDRs in the light chain variable region (VL) set forth as SEQ ID NO: 730; (ii) three CDRs in the VH set forth as SEQ ID NO: 891 and three CDRs in the VL set forth as SEQ ID NO: 892; (iii) three CDRs in the VH set forth as SEQ ID NO: 961 and three CDRs in the VL set forth as SEQ ID NO: 962; (iv) three CDRs in the VH set forth as SEQ ID NO: 979 and three CDRs in the VL set forth as SEQ ID NO: 980; (v) three CDRs in the VH set forth as SEQ ID NO: 1039 and three CDRs in the VL set forth as SEQ ID NO: 1040; (vi) three CDRs in the VH set forth as SEQ ID NO: 1103 and three CDRs in the VL set forth as SEQ ID NO: 1104; (vii) three CDRs in the VH set forth as SEQ ID NO: 1107 and three CDRs in the VL set forth as SEQ ID NO: 1108; (viii) three CDRs in the VH set forth as SEQ ID NO: 1111 and three CDRs in the VL set forth as SEQ ID NO: 1112; (ix) three CDRs in the VH set forth as SEQ ID NO: 1121 and three CDRs in the VL set forth as SEQ ID NO: 1122; (x) three CDRs in the VH set forth as SEQ ID NO: 1133 and three CDRs in the VL set forth as SEQ ID NO: 1134; (xi) three

CDRs in the VH set forth as SEQ ID NO: 1157 and three CDRs in the VL set forth as SEQ ID NO: 1158; (xii) three CDRs in the VH set forth as SEQ ID NO: 1143 and three CDRs in the VL set forth as SEQ ID NO: 1144; (xiii) three CDRs in the VH set forth as SEQ ID NO: 1251 and three CDRs in the VL set forth as SEQ ID NO: 1252; or (xiv) three CDRs in the VH set forth as SEQ ID NO: 1255 and three CDRs in the VL set forth as SEQ ID NO: 1256; wherein the CDRs are defined by Kabat, Chothia or MacCallum numbering.

[1440] In other embodiments, ACE2-blocking anti-SARS-CoV-2 antibodies or antigen binding fragments comprise: (i) the VH set forth as SEQ ID NO: 729 and the VL set forth as SEQ ID NO: 730; (ii) the VH set forth as SEQ ID NO: 891 and the VL set forth as SEQ ID NO: 892; (iii) the VH set forth as SEQ ID NO: 961 and the VL set forth as SEQ ID NO: 962; (iv) the VH set forth as SEQ ID NO: 979 and the VL set forth as SEQ ID NO: 980; (v) the VH set forth as SEQ ID NO: 1039 and the VL set forth as SEQ ID NO: 1040; (vi) the VH set forth as SEQ ID NO: 1103 and the VL set forth as SEQ ID NO: 1104; (vii) the VH set forth as SEQ ID NO: 1107 and the VL set forth as SEQ ID NO: 1108; (viii) the VH set forth as SEQ ID NO: 1111 and the VL set forth as SEQ ID NO: 1112; (ix) the VH set forth as SEQ ID NO: 1121 and the VL set forth as SEQ ID NO: 1122; (x) the VH set forth as SEQ ID NO: 1133 and the VL set forth as SEQ ID NO: 1134; (xi) the VH set forth as SEQ ID NO: 1157 and the VL set forth as SEQ ID NO: 1158; (xii) the VH set forth as SEQ ID NO: 1143 and the VL set forth as SEQ ID NO: 1144; (xiii) the VH set forth as SEQ ID NO: 1251 and the VL set forth as SEQ ID NO: 1252; or (xiv) the VH set forth as SEQ ID NO: 1255 and the VL set forth as SEQ ID NO: 1256.

[1441] The amino acid sequences of the antibodies described in the foregoing paragraphs are provided below:

292-VH

(SEQ ID NO: 729)

EVQLVESGGGLIQQPGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTYYADSVKGRFTISR

DNSKNTLYLQMNSLRAEDTAVYYCARDLQGGGPGWGQGLTVTVSS

292-VL

(SEQ ID NO: 730)

AIQMTQSPSSLSASVGVDRVTITCRASQGIKNDLWYQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDTF

LTISLQPEDFATYYCLQDYNYPRTFGQGTKVEIK

309-VH

(SEQ ID NO: 763)

QVQLVESGGGVVQPGSLRLSCAASGFTFSSYAMHWVRQAPGKLEWVAIVSYDGSNKYYADSVKGRFT

ISRDNKNTLYLQMNSLRAEDTAVYYCARPYSGSYQSYFDYWGQGLTVTVSS

309-VL

(SEQ ID NO: 764)

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLIYGASSRATGIPDRFSGSGSGTDTF

LTISRLEPEDFAVYYCQYQGSPLTFGGGTKVEIK

364-VH

(SEQ ID NO: 873)

EVQLVESGGGLVQPGSLRLSCTASGFTFGDYAMSWVRQAPGKLEWVGFIRSKAYGGTTEYAASVKGRF

TISRDDSKSIAYLQMNSLKTEDTAVYYCTRFIDYDIWGSYRYTTLFDYWGQGLTVTVSS

- continued

364-VL

(SEQ ID NO: 874)

DIQMTQSPSTLSASVGDRTTITCRASQSISSWLA WYQQKPGKAPKLLIYKASSLES GVP SRFS GSGSGTEFTL

TISSLQPD FAYYCQQYNSYSYTTGGTKLEIK

373-VH

(SEQ ID NO: 891)

QVQLQESGPGLVKPS ETL SLTCTVSGGSISSYWSWIRQPPGKGLEWIGYIYSGSTNYNPSL KSRVTISVDT

SKNQFSLKLS SVTAADTAVYYCARGPDYDFWSGYFYGMDVWGQGT TTVTVSS

373-VL

(SEQ ID NO: 892)

EIVLTQSPGTL SLSPGERATL SCRASQSVSSSYLA WYQQKPGQAPRLLIYGASSRATGIPDRFS GSGSGTDFT

LTISRLEPEDFAVYYCQQYGS SLTFGGTKVEIK

388-VH

(SEQ ID NO: 921)

QVQLVESGGGVVQGRSLRLSCAASGFTFNSYAIHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFTI

SRDNSKNTLYLQMNSLRAEDTAVYYCARGRGYRSYFDYWGQGT LTVTVSS

388-VL

(SEQ ID NO: 922)

EIVLTQSPGTL SLSPGERATL SCRASQSVSSSYLA WYQQKPGQAPRLLIYGASSRATGIPDRFS GSGSGTDFT

LTISRLEPEDFAVYYCQQYGS SPNTFGGTKLEIK

408-VH

(SEQ ID NO: 961)

QVQLVESGGGVVQGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFT

ISRDNSRNTLYLQMNSLRAEDTAVYYCAKGADTPHYSYDGLSVGYYYYGMDVWGQGT TTVTVSS

408-VL

(SEQ ID NO: 962)

DIQMTQSPSSLSASVGDRTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSFQSGVPSRFS GSGSGTDFTL

TISSLQPEDFAYYCQQYSAPFTFGGTKVDIK

414-VH

(SEQ ID NO: 973)

QVQLVQSGAEVKKPGASVKVCKASGYTFTSYMHVVRQAPGQGLEWMGIINPSGGSTSYAQKFGGRVT

MTRDTSSTSTVYMESSLRSED TAVYYCARDPPGRDFWSGYF GAPDY YYYGMDVWGQGT TTVTVSS

414-VL

(SEQ ID NO: 974)

DIQMTQSPSSLSASVGDRTTITCRASQGISNYLAWFQQKPGKAPKSLIYAASSLQSGVPSKFSGSGSGTDFTL

TISSLQPEDFAYYCQQYNSYPYTFGGTKLEIK

417-VH

(SEQ ID NO: 979)

QLQLQESGPGLVKPS ETL SLTCTVSGGSISSSYWGWIRQPPGKGLEWIGSIYSGSTYYNPSL KSRVTISV

DTSKNQFSLKLS SVTAADTAVYYCARAPIMITFGGVTGHFDYWGQGT LTVTVSS

417-VL

(SEQ ID NO: 980)

EIVLTQSPGTL SLSPGERATL SCRASQSVSSSYLA WYQQKPGQAPRLLIYGASSRATGIPDRFS GSGSGTDFT

LTISRLEPEDFAVYYCQQYGS SPLTFGGTKVEIK

419-VH

(SEQ ID NO: 983)

QVQLVQSGAEVKKPGASVKVCKASGYTFTSYMHVVRQAPGQGLEWMGIINPSGGSTSYAQKFGGRVT

MTRDTSSTSTVYMESSLRSED TAVYYCARDWTQSSGYDY YGLDVWGQGT TTVTVSS

419-VL

(SEQ ID NO: 984)

SYELTQPPSVSVSPGQTARITCSGDALPKQYAYWYQQKPGQAPVVVIYKDSERPSGIPERFSGSSSGTTVTLT

ISGVQAEDEADYYCQ SADSSGTYVVFGGTKLTVL

- continued

442 - VH

(SEQ ID NO: 1029)

QVQLVESGGGVVQPGRSLRLSSAASGFTFSSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVRGRFTI

SRDNSKNTLYLQMNSLRAEDTAVYYCARPKGGSYSDAFDIWGQGTMTVTVSS

442 - VL

(SEQ ID NO: 1030)

EIVLTQSPGTLSTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT

LTISRLEPEDFAVYYCQQYGSSPQSFGPGTKVDIK

445 - VH

(SEQ ID NO: 1035)

QVQLVQSGAEVKKPGASVKVCKASGYTFTSYMHWVRQAPGQGLEWMGIINPSGGSTSYAQKFGGRVT

MTRDTSTSTVMELSSLRSEDTAVYYCARDPTEVGATSEYYYYGMDVWGQGTITVTVSS

445 - VL

(SEQ ID NO: 1036)

SYELTQPPSVSVSPGQTARITCSGDALPKQYAYWYQQKPGQAPVLVIYKDSERPSGIPERFSGSSSGTTVTLT

ISGVQAEDADYYCQSDSSGTYVVFSGGKLTVL

447 - VH

(SEQ ID NO: 1039)

EVQLVESGGGLVQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGSTYYADSVKGRFTIS

RDNSKNTLYLQMNSLRAEDTAVYYCARDKSSGSGPWGQGTITVTVSS

447 - VL

(SEQ ID NO: 1040)

AIQMTQSPSSLSASVGDRTTITCRASQGIKNDLWYQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFT

LTISLQPEDFATYYCLQDYNPRTFGQGTKEIK

462 - VH

(SEQ ID NO: 1069)

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVIWDGNNKYYADSVKGRF

TISRDNKNTLYLQMNSLRAEDTAVYYCAKDPTWFGELPSYYYYGMDVWGQGTITVTVSS

462 - VL

(SEQ ID NO: 1070)

DIQMTQSPSTLSASVGDRTTITCRASQISSWLAWYQQKPGKAPKLLIYKASSLESQVPSRFRSGSGSGTEFTL

TISSLQPDDEFATYYCQYNSYPITFGQGTREIK

479 - VH

(SEQ ID NO: 1103)

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSYISSSSSTIYADSVKGRFTISR

DNAKNTLYLQMNSLRAEDTAVYYCARDLGARTPWDIVVPAAMDYWGQGTITVTVSS

479 - VL

(SEQ ID NO: 1104)

EIVLTQSPGTLSTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT

LTISRLEPEDFAVYYCQQYGRSPNTFGQGTKEIK

481 - VH

(SEQ ID NO: 1107)

EVQLVESGGGLIQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTFYADSVKGRFTISR

DNSKNTLYLQMNSLRAEDTAVYYCAREVAGTYDYWGQGTITVTVSS

481 - VL

(SEQ ID NO: 1108)

DIQMTQSPSSVSASVGDRTTITCRASQISSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFT

LTISLQPEDFATYYCQANSFPGGTFGPGTKVDIK

483 - VH

(SEQ ID NO: 1111)

QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWIRQPPGKGLEWIGYIYSGSTNYNPISLKSRTISVDI

SKNQFSLKLSSVTAADTAVYYCARAPEEKSSEIGELVGWGWFDPWGQGTITVTVSS

- continued

483-VL

(SEQ ID NO: 1112)

SYELTQPPSVSVSPGQTASIIICSGDKLGDKYACWYQQKPGQSPVLVIYQDSKRPSGIPERFSGSNSGNTATLT

ISGTQAMDEADYYCQAWDSSTVVFGGGKLTVL

488-VH

(SEQ ID NO: 1121)

EVQLVESGGGLIQPGGSLRLSCAASGLTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTYYADSVKGRFTISR

DNSKNTLYLQMNSLRAEDTAVYYCARSPYGGNSWGQGLTVTVSS

488-VL

(SEQ ID NO: 1122)

DIQMTQSPSSLSASVGDRVITITCQASQDISNYLNWYQQKPGKAPKLLIYDASNLETGVPSRFRSGSGSGTDFT

FTISLQPEDFATYYCQYDNLPITFGQGRLEIK

494-VH

(SEQ ID NO: 1133)

EVQLVESGGGLVQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTFYADSVKGRFTISR

DNSKNTLYLQMNSLRAEDTAVYYCARDSDQLLDYWGQGLTVTVSS

494-VL

(SEQ ID NO: 1134)

DIQLTQSPSFLSASVGDRVITITCRASQGISSYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFRSGSGSGTEFTL

TISSLQPEDFATYYCQQLNSYPPTFGPGTKVDIK

506-VH

(SEQ ID NO: 1157)

QITLKESGPTLVKPTQTTLTCTFSGFSLSTSGVGWIRQPPGKALEWLALIYDDDKRYSPLKSRLTITK

DTSKNQVLTMTNMDPVDATYYCAHSLSSIFDYWGQGLTVTVSS

506-VL

(SEQ ID NO: 1158)

QSALTQPASVSGSPGQSITISCTGTSSDVGDYNYVSWYQQHPGKAPKLMIEVSNRPSGVNRFSGSKSGNT

ASLTISGLQAEDEADYYCSSYTSSSTVFGGKLTVL

540-VH

(SEQ ID NO: 1225)

EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGHIKSKTDGGTTDYAAPVKGR

FTISRDDSNTLYLQMNSLKTEDTAVYYCTREPYFDYWGQGLTVTVSS

540-VL

(SEQ ID NO: 1226)

DIQMTQSPSTLSASVGDRVITITCRASQSISSWLAWYQQKPGKAPKLLIYKASSLESQVPSRFRSGSGSGTEFTL

TISSLQPDDEFATYYCQYNSYRYTFGQGTKLEIK

549-VH

(SEQ ID NO: 1243)

EVQLVESGGGLIQPGGSLRLSCAASGLTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTYYADSVKGRFTISR

DNSKNTLYLQMNSLRAEDTAVYYCARSPYGGNSWGQGLTVTVSS

549-VL

(SEQ ID NO: 1244)

DIQMTQSPSSLSASVGDRVITITCRTSYIYNWYQQKPGKAPKFLIYAASSFQNGVPSRFRSGSGSGTDFTF

TISSLQPEDFATYYCQGYSTPLTFGGGTKVEIK

553-VH

(SEQ ID NO: 1251)

QVQLVQSGAEVKKPGASVKVCKASGYTFTSYGISWVRQAPGQGLEWMGWISAYNGNTNYAQKLQGRV

TMTTDTSTSTAYMELRSLRSDDTAVYYCARDRGYAATFGVFDYWGQGLTVTVSS

553-VL

(SEQ ID NO: 1252)

DIQMTQSPSSLSASVGDRVITITCRASQSISSYLWYQQKPGKAPKLLISAASSLQSGVPSRFRSGSGSGTDFTL

TISSLQPEDFATYYCQSYSTAFTFGGGTKVDIK

-continued

555-VH

(SEQ ID NO: 1255)

QVQLVQSGAEVKKPGSSVKVCKASGGTFSNYAISWVRQAPGQGLEWMGRIIPILGIANYAQKPFQGRVTIT

ADKSTSTAYMELSSLRSEDTAVYYCARGYYEARHYYYYYAMDVWGQTAVTVSS

555-VL

(SEQ ID NO: 1256)

DIQMTQSPSSLSASVGDRTTITCRASQSISSYLSWYQQKPKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTL

TITSLQPEDFATYYCQSYSTPRTFGQGTKVEIK

562-VH

(SEQ ID NO: 1269)

QVQLVESGGGVVQPGRLRLSCAASGFTFSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFT

ISRDNKNTLYLQMNSLRAEDTAVYYCARASSGGYQGPDPWGQGLTVTVSS

562-VL

(SEQ ID NO: 1270)

QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVSNRPSGVSNRPSGSKSGNT

ASLTISGLQAEDEADYYCSYSSSTLLYVFGTGKTVTL

[1442] In some embodiments, the anti-SARS-CoV-2 antibodies provided herein have IgG1 isotype. In some embodiments, the anti-SARS-CoV-2 antibodies provided herein have IgG1m3 allotype.

[1443] In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5319 and a light chain comprising SEQ ID NO: 5320. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5321 and a light chain comprising SEQ ID NO: 5322. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5322 and a light chain comprising SEQ ID NO: 5323. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5327 and a light chain comprising SEQ ID NO: 5328. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5329 and a light chain comprising SEQ ID NO: 5330. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5331 and a light chain comprising SEQ ID NO: 5332. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5333 and a light chain comprising SEQ ID NO: 5334. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5337 and a light chain comprising SEQ ID NO: 5338. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5339 and a light chain comprising SEQ ID NO: 5340.

In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5341 and a light chain comprising SEQ ID NO: 5342. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5343 and a light chain comprising SEQ ID NO: 5344. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5345 and a light chain comprising SEQ ID NO: 5346. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5349 and a light chain comprising SEQ ID NO: 5350. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5353 and a light chain comprising SEQ ID NO: 5354. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5355 and a light chain comprising SEQ ID NO: 5356. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5357 and a light chain comprising SEQ ID NO: 5358. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5359 and a light chain comprising SEQ ID NO: 5360. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5361 and a light chain comprising SEQ ID NO: 5362. In some embodiments, provided herein is an antibody comprising a heavy chain comprising SEQ ID NO: 5365 and a light chain comprising SEQ ID NO: 5366.

[1444] The amino acid sequences of the antibodies described in the preceding paragraph are provided below:

292 Heavy chain full length sequence

(SEQ ID NO: 5319)

EVQLVESGGGLIQPGSLRLSCAASGFTVSSNYMSWVRQAPGKLEWVSVIYSGGSTYYADSVKGRFTISR

DNSKNILYLQMNSLRAEDTAVYYCARDLQGGGPGWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAAL

GCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDK

RVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV

- continued

HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSR

EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFC

SVMHEALHNHYTQKSLSLSPGK

292 Light chain full length sequence

(SEQ ID NO: 5320)

AIQMTQSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPKAPKLLIYAASSLQSGVPSRSESGSGTDFT

LTISLQPEDFATYYCLQDYNYPRTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK

VQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

309 Heavy chain full length sequence

(SEQ ID NO: 5321)

QVQLVESGGGVVQPGSRRLSCAASGFTFSSYAMHWVRQAPGKLEWVAVISYDGSNKYYADSVKGRFT

ISRDNKNTLYLQMNSLRAEDTAVYYCARPYSGSYQSYFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTS

GGTAALGCLVKDYFPEPVTWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPS

NTKVDKRVPEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY

VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV

YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQ

GNVFSCSVMHEALHNHYTQKSLSLSPGK

309 Light chain full length sequence

(SEQ ID NO: 5322)

EIVMTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRESGSGSGTDFT

LTISRLEPEDFAVYYCQQYGSPLTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK

VQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

364 Heavy chain full length sequence

(SEQ ID NO: 5323)

EVQLVESGGGLVQPGSRRLSCTASGFTFGDYAMSWFRQAPGKLEWVGFIKSKAYGGTTEYAASVKGRF

TISRDDSKSIAYLQMNSLKTEDTAVYYCTRFGIDYDIWGSYRYTTLFDYWGQGLTVTVSSASTKGPSVFP

LAPSSKSTSGGTAALGCLVKDYFPEPVTWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYI

CNVNHKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP

EVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK

GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT

VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

364 Light chain full length sequence

(SEQ ID NO: 5324)

DIQMTQSPSTLSASVGDRVTITCRASQSISSWLAWYQQKPKAPKLLIYKASSLESQVPSRFSGSGSGTEFTL

TISSLQPDDEFATYYCQQYNSYSYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV

QWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

373 Heavy chain full length sequence

(SEQ ID NO: 5325)

QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWIRQPPGKLEWIGIYYSGSTNYNPSLKSRTISVDT

SKNQFSLKLSVTAADTAVYYCARGPDYDFWGSYFYGMDVWGQGTITVTVSSASTKGPSVFPLAPSSKST

SGGTAALGCLVKDYFPEPVTWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLTQTYICNVNHKPS

NTKVDKRVPEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY

VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV

YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDGSGSFFLYSKLTVDKSRWQQ

GNVFSCSVMHEALHNHYTQKSLSLSPGK

-continued

373 Light chain full length sequence

(SEQ ID NO: 5326)

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT
LTISRLEPEDFAVYYCQQYGSSLTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV
QWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

388 Heavy chain full length sequence

(SEQ ID NO: 5327)

QVQLVESGGGVVQPGRSLRLSCAASGFTFNSYAIHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFTI
SRDNSKNTLYLQMNSLRAEDTAVYYCARGRGGYRSYFDYWGGTLVTVSSASTKGPSVFPLAPSSKSTSG
GTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNT
KVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVD
LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGN
VFSCSVMHEALHNHYTQKSLSLSPGK

388 Light chain full length sequence

(SEQ ID NO: 5328)

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT
LTISRLEPEDFAVYYCQQYGSSPNTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

408 Heavy chain full length sequence

(SEQ ID NO: 5329)

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFT
ISRDNSRNTLYLQMNSLRAEDTAVYYCAKGADTPHYSYGYDFLSVGYYYYGMDVWGGTQTVTVSSASTKG
PSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGT
QTYICNVNHKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS
HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLY
SLKTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

408 Light chain full length sequence

(SEQ ID NO: 5330)

DIQMTQSPSSLSASVGDRVTITCRASQISISYLNWYQQKPGKAPKLLIYAASSFQSGVPSRFSRSGSGTDFTL
TISSLQPEDFATYYCQQYSAPFTFGPGTKVDIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV
QWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

414 Heavy chain full length sequence

(SEQ ID NO: 5331)

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWRQAPGGGLEWMGIINPSGGSTSYAQKFQGRVT
MTRDTSTSTVMELSSLRSEDITAVYYCARDPPGRDFWSGYFYGAPDYIIYYGMDVWGGTQTVTVSSAST
KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSS
LGTQTYICNVNHKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV
DVSHEDPEVKFNWYVDGVEVHNAKTKPREEVYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFF
LYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

414 Light chain full length sequence

(SEQ ID NO: 5332)

DIQMTQSPSSLSASVGDRVTITCRASQISNYLAWFQQKPGKAPKSLIYAASSLQSGVPSKFSRSGSGTDFTL
TISSLQPEDFATYYCQQYNSYPYTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV
QWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

- continued

417 Heavy chain full length sequence

(SEQ ID NO: 5333)

QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYYWGWRQPPGKGLEWIGSIYYSGSTYYNPSLKSRTISV
DTSKNQFSLKLSSTAADTAVYYCARAPIMITFGGVTGHFDYWGGTLVTVSSASTKGPSVFPLAPSSKSTS
GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPS
NTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY
VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV
YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQ
GNVFSCSVMHEALHNHYTQKSLSLSPGK

417 Light chain full length sequence

(SEQ ID NO: 5334)

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT
LTISRLEPEDFAVYYCQQYGSSPLTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

419 Heavy chain full length sequence

(SEQ ID NO: 5335)

QVQLVQSGAEVKKPGASVKVSFKASGYTFTSYMHVWRQAPQGQLEWMIINPSGGSTSYAQKFQGRVT
MTRDTSSTVYMESSLRSIEDTAVYYCARDWTQSSGYDYYYGLDVWGQGTITVTVSSASTKGPSVFPLAPS
SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVN
HKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF
NMYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR
EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKS
RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

419 Light chain full length sequence

(SEQ ID NO: 5336)

SYELTQPPSVSVSPGQTARITCSGDALPKQYAYWYQQKPGQAPVVVIYKDSERPSGIPERFSGSSSGTTVTLT
ISGVQAEDEADYNQCSADSSGTYVVFVGGGKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGA
VTVAWKADSSPVKAGVETTPSKSQSNNKYAASSYLSLTPEQWKSRSYSQCQVTHEGSTVEKTVAPTECS

442 Heavy chain full length sequence

(SEQ ID NO: 5337)

QVQLVESGGGVVQPGSRSLRLSSAASGFTFSSYAMHWVRWAPGKLEWVAVISYDGSNKYYADSVRGRFTI
SRDNSKNTLYLQMNSLRAEDTAVYYCARPKGGSYDAFDIWQGTMTVTVSSASTKGPSVFPLAPSSKSTS
GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPS
NTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY
VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV
YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQ
GNVFSCSVMHEALHNHYTQKSLSLSPGK

442 Light chain full length sequence

(SEQ ID NO: 5338)

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT
LTISRLEPEDFAVYYCQQYSSPQSFPGTKVDIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

- continued

445 Heavy chain full length sequence

(SEQ ID NO: 5339)

QVQLVQSGAEVKKPGASVKVCKASGYTFTSYMHWRQAPGQGLEWMGIINPSGGSTSYAQKFQGRVT
MTRDTSTSTVYMELSSLRSEDVAVYCARDPTEVGATSEYYYYGMDVWGQGTITVSSASTKGPSVFPLA
PSSKSTSGGTAALGCLVKDFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICN
VNHKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV
KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPAPIEKTISKAKGQ
PREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVD
KSRWQQGNVFCSCVMHEALHNHYTQKSLSKSPGK

445 Light chain full length sequence

(SEQ ID NO: 5340)

SYELTOPPSVSVSPGQTARITCSGDALPKQYAYWYQKPGQAPVLVIYKDSERPSGIPERFSGSSSGTTVTLT
ISGVQAEDEADYYCQADSSGTYVVFVGGGKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGA
VTVAWKADSSPVKAGVETTPSKSQSNNKYAASSYLSLTPEQWKSIIIRSYSCQVTIIEGSTVEKTVAPTECS

447 Heavy chain full length sequence

(SEQ ID NO: 5341)

EVQLVESGGGLVQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTYYADSVKGRFTIS
RDNSKNTLYLQMNSLRAEDTAVYYCARDKSSGSGPWGQGTITVSSASTKGPSVFPLAPSSKSTSGGTAA
LGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVD
KRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE
VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPAPIEKTISKAKGQPREPQVYTLPP
SREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV
FCSCVMHEALHNHYTQKSLSLSPGK

447 Light chain full length sequence

(SEQ ID NO: 5342)

AIQMTQSPSSLSASVGDRVTITCRASQGIKNDLWYQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFT
LTISSLQPEDFATYYCLQDYNYPRTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

462 Heavy chain full length sequence

(SEQ ID NO: 5343)

QVQLVESGGGVVQPGSRSLRLSCAASGFTFSSYGMHWRQAPGKGLEWVAVIWDGNNKYYADSVKGRF
TISRDNKNTLYLQMNSLRAEDTAVYYCAKDPTWFGELPSYYYYGMDVWGQGTITVSSASTKGPSVFPL
APSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPE
VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPAPIEKTISKAKG
QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTV
DKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

462 Light chain full length sequence

(SEQ ID NO: 5344)

DIQMTQSPSTLSASVGDRVTITCRASQSISSWLAWYQQKPGKAPKLLIYKASSLESQVPSRFRSGSGSGTEFTL
TISLQPDDEFATYYCQQYNSYPPIITFGQTRLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

-continued

479 Heavy chain full length sequence

(SEQ ID NO: 5345)

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSIISSSSSTIYYADSVKGRFTISR
DNAKNSLYLQMNSLRAEDTAVYYCARDLGARTPWDIVVPAAMDYWGQGLTVTVSSASTKGPSVFPLAP
SSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV
NHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVK
FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDK
SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

479 Light chain full length sequence

(SEQ ID NO: 5346)

EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFT
LTISRLEPEDFAVYYCQQYGRSPNTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

481 Heavy chain full length sequence

(SEQ ID NO: 5347)

EVQLVESGGGLIQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSIIYSGGSTFYADSVKGRFTISR
DNSKNTLYLQMNSLRAEDTAVYYCAREVAGTYDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAAL
GCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDK
RVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV
HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSR
EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSC
SVMHEALHNHYTQKSLSLSPGK

481 Light chain full length sequence

(SEQ ID NO: 5348)

DIQMTQSPSSVSASVGRVTTICTRASQGISWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFT
LTISLQPEDFATYYCQQANSFPGGTFGPGTKVDIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
KVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

483 Heavy chain full length sequence

(SEQ ID NO: 5349)

QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIRQPPGKGLEWIGYIIYSGSTNYNPSLKSRTISVDT
SKNQFSLKLSVTAADTAVYYCARAPEEKSSEIGELVGWGWFDWPGQGLTVTVSSASTKGPSVFPLAPSSK
STSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHK
PSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ
VYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK

483 Light chain full length sequence

(SEQ ID NO: 5350)

SYELTQPPSVSVSPGQTASIIICSGDKLGDKYACWYQQKPGQSPVLVIYQDSKRPSGIPERFSGNSGNTATLT
ISGTQAMDEADYYCQAWDSSTVVFGGGKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAV
TVAWKADSSPVKAGVETTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS

488 Heavy chain full length sequence

(SEQ ID NO: 5351)

EVQLVESGGGLIQPGGSLRLSCAASGLTVSSNYMSWVRQAPGKGLEWVSIIYSGGSTYYADSVKGRFTISR
DNSKNTLYLQMNSLRAEDTAVYYCARSPYGGNSWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALG

- continued

CLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKR
VEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE
EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSS
VMHEALHNHYTQKSLSLSPGK

488 Light chain full length sequence

(SEQ ID NO: 5352)

DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPKAPKLLIYDASNLETGVPSTRFSGSGSGTDFT
FTISSLQPEDIATYYCQQYDNLPITFGQGTREIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV
QWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

494 Heavy chain full length sequence

(SEQ ID NO: 5353)

EVQLVESGGGLVQPQGSRLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTFYADSVKGRFTISR
DNSKNTLYLQMNSLRAEDTAVYYCARDSDQLLDYWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAA
LGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD
KRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE
VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPP
SREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV
SCSMHEALHNHYTQKSLSLSPGK

494 Light chain full length sequence

(SEQ ID NO: 5354)

DIQLTQSPSFLSASVGDRVTITCRASQGISSYLAWYQQKPKAPKLLIYAASLTQSGVPSRFSGSGSGTEFTL
TISSLQPEDFATYYCQQLNSYPPTFGPGTKVDIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

506 Heavy chain full length sequence

(SEQ ID NO: 5355)

QITLKESGPTLVKPTQTTLTCTFSGFSLSTSGVGVGWIRQPPGKALEWLALIYDDDKRYSPLKSRLTITK
DTSKNQVVLTMNMPDVTATYYCAHSLSSIIFYWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAAAL
GCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDK
RVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV
HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSR
EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFC
SVMHEALHNHYTQKSLSLSPGK

506 Light chain full length sequence

(SEQ ID NO: 5356)

QSALTQPASVSGSPGQSITISCTGTSSDVGDYNYVSWYQQHPGKAPKLMIEVSNRPSGVSNRFSGSGSGNT
ASLTISGLQAEDADYYCSSYTSSSTVFGGGTKLTVLGQPKAAFSVTLFPPSSEELQANKATLVCLISDFYPG
AVTVAWKADSSPVKAGVETTPPSKQSNKYYASSYLSLTPEQWKSHRYSYSCQVTHEGSTVEKTVAPTECS

- continued

540 Heavy chain full length sequence

(SEQ ID NO: 5357)

EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGHIKSKTDGGTTDYAAPVKGR
FTISRDDSNTLYLQMNLSKTEDTAVYYCTREPYFYFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGT
AALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTK
VDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG
VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL
PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN
VFSCSVMEALHNHYTQKSLSLSPGK

540 Light chain full length sequence

(SEQ ID NO: 5358)

DIQMTQSPSTLSASVGDRVTITCRASQSISSWLAWYQQKPGKAPKLLIYKASSLESQVPSRFSGSGSGTEFTL
TISSLQPDDEFATYYCQQYNSRYRTPGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK
VQWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

549 Heavy chain full length sequence

(SEQ ID NO: 5359)

EVQLVESGGGLIQPGGSLRLSCAASGLTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTYYADSVKGRFTISR
DNSKNTLYLQMNLSRAEDTAVYYCARSYPGGNSWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALG
CLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKR
VEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE
EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSS
VMHEALHNHYTQKSLSLSPGK

549 Light chain full length sequence

(SEQ ID NO: 5360)

DIQMTQSPSSLSASVGDRVTITCRTSQTIYNLYNWYQQKPGKAPFLIYAASSFQNGVPSRFSGSGSGTDFTF
TISSLQPEDFATYYCQQGYSTPLTFGGGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV
QWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

553 Heavy chain full length sequence

(SEQ ID NO: 5361)

QVQLVQSGAEVKKPGASVKVCKASGYTFTSYGISWVRQAPGQGLEWMGWISAYNGNTNYAQKLQGRV
TMTTDTSTSTAYMELRSLRSDDTAVYYCARDRGYAATFGVFDYWGQGLTVTVSSASTKGPSVFPLAPSSK
STSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHK
PSNTKVDKRVKPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ
VYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMEALHNHYTQKSLSLSPGK

553 Light chain full length sequence

(SEQ ID NO: 5362)

DIQMTQSPSSLSASVGDRVTITCRASQSISSLNWYQQKPGKAPLLISAASSLQSGVPSRFSGSGSGTDFTL
TISSLQPEDFATYYCQQSYSTAFTPGPGTKVDIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV
QWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

555 Heavy chain full length sequence

(SEQ ID NO: 5363)

QVQLVQSGAEVKKPGSSVKVCKASGGTFSNYAISWVRQAPGQGLEWMGRIIPILGIANYAQKFQGRVTIT
ADKSTSTAYMELSSLRSEDATVYYCARGYYEARHYYYYYAMDVWGQTAVTVSSASTKGPSVFPLAPSS

- continued

KSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYINCVDH

KPSNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN

WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPRE

PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKSR

WQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

555 Light chain full length sequence

(SEQ ID NO: 5364)

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLSWYQQKPKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTL

TITSLQPEDFATYYCQQSYSTPRTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKV

QWKVDNALQSGNSQESVTEQDSKDSSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

562 Heavy chain full length sequence

(SEQ ID NO: 5365)

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGRFT

ISRDNSKNTLYLMNLSRAEDTAVYYCARSSGGYQGPDPWGQGLTVTVSSASTKGPSVFPLAPSSKSTS

GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYINCVDH

NPKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY

VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV

YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSLKTVDKSRWQQ

GNVFSCSVMHEALHNHYTQKSLSLSPGK

562 Light chain full length sequence

(SEQ ID NO: 5366)

QSALTQPASVSGSPGQSITISCTGTSSDVGGINVSWYQQHPGKAPKLMIDVSNRPSGVSNRFSGSGSGNT

ASLTISGLQAEDADYYCSTSSSTLLYVFGTGTQVTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDF

YPGAFTVAWAKDSSPVKAGVETTTSPSKQSNKYYAASSYLSLTPEQWKSRSYSCQVTHEGSTVEKTVAPT

ECS

Fc region with ACT5 mutations

(SEQ ID NO: 5367)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVP

SSSLGTQTYINCVDHNSKNTKVDKRVPEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV

VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA

PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG

SFFLYSLKTVDKSRWQQGNVFSCSVLHEALHAHYTRKELSLSPGK

[1445] In particular embodiments, the foregoing antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364. In some embodiments, the foregoing antibody comprises a heavy chain comprising SEQ ID NO: 5335 and a light chain comprising SEQ ID NO: 5336. In some embodiments the foregoing antibody comprises a heavy chain comprising SEQ ID NO: 5347 and a light chain comprising SEQ ID NO: 5348. In some embodiments, the foregoing antibody comprises a heavy chain comprising SEQ ID NO: 5351 and a light chain comprising SEQ ID NO: 5352. In some embodiments, the foregoing antibody comprises a heavy chain comprising SEQ ID NO: 5325 and a light chain comprising SEQ ID NO: 5326.

[1446] Also provided herein are antibodies that bind SARS-CoV-2 spike protein, wherein the antibodies bind an epitope comprising one or more residues within amino acids

434-444 and 459-495 of the SARS-CoV-2 spike protein, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In some embodiments, provided herein are antibodies that bind SARS-CoV-2 spike protein, wherein the antibodies bind an epitope comprising two, three, four, five or more residues within amino acids 434-444 and 459-495 of the SARS-CoV-2 spike protein, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In some embodiments, the epitope is determined by HDX-MS (hydrogen-deuterium exchange mass spectrometry).

[1447] Also provided herein are antibodies that bind SARS-CoV-2 spike protein, wherein the antibodies bind an epitope comprising one or more of the following residues of SARS-CoV-2 spike protein: Y351, Y449, N450, L452, L455, F456, T470, I472, N481, G482, V483, E484, G485, F486, Y489, F490, L492, Q493, S494, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In

some embodiments, the epitope comprises two or more (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19) of the listed residues. In some embodiments, the epitope comprises three or more of the listed residues. In some embodiments, the epitope comprises four or more of the listed residues. In some embodiments, the epitope comprises five or more of the listed residues. In some embodiments, the epitope is determined by X-ray crystallography.

[1448] Also provided herein are antibodies that bind SARS-CoV-2 spike protein, wherein the antibodies bind an epitope comprising one or more of the following residues of SARS-CoV-2 spike protein: R403, D405, R408, Q409, T415, G416, K417, D420, Y421, Y453, L455, F456, R457, K458, S459, N460, Y473, Q474, A475, G476, F486, N487, Y489, Q493, S494, Y495, G496, Q498, T500, N501, G502, Y505, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In some embodiments, the epitope comprises two or more (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32) of the listed residues. In some embodiments, the epitope comprises three or more of the listed residues. In some embodiments, the epitope comprises four or more of the listed residues. In some embodiments, the epitope comprises five or more of the listed residues. In some embodiments, the epitope is determined by X-ray crystallography.

[1449] Also provided herein are antibodies that bind SARS-CoV-2 spike protein, wherein the antibodies bind an epitope comprising one or more of the following residues of SARS-CoV-2 spike protein: R403, T415, G416, K417, D420, Y421, L455, F456, R457, K458, S459, N460, Y473, Q474, A475, G476, S477, F486, N487, Y489, N501, G502, Y505, wherein the amino acid residue positions correspond to SEQ ID NO: 5317. In some embodiments, the epitope comprises two or more (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23) of the listed residues. In some embodiments, the epitope comprises three or more of the listed residues. In some embodiments, the epitope comprises four or more of the listed residues. In some embodiments, the epitope comprises five or more of the listed residues. In some embodiments, the epitope is determined by X-ray crystallography.

[1450] The term “epitope” refers to the amino acid residues, of an antigen, that are bound by an antibody. An epitope can be a linear epitope, a conformational epitope, or a hybrid epitope. An epitope can be determined according to different experimental techniques, also called “epitope mapping techniques.” It is understood that the determination of an epitope may vary based on the different epitope mapping techniques used and may also vary with the different experimental conditions used, e.g., due to the conformational changes or cleavages of the antigen induced by specific experimental conditions. Epitope mapping techniques are known in the art (e.g., Rockberg and Nilvebrant, *Epitope Mapping Protocols: Methods in Molecular Biology*, Humana Press, 3rd ed. 2018), including, but not limited to, X-ray crystallography, nuclear magnetic resonance (NMR) spectroscopy, electron microscopy, site-directed mutagenesis, species swap mutagenesis, alanine-scanning mutagenesis, hydrogen-deuterium exchange (HDX), and cross-blocking assays.

[1451] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or

antigen-binding fragment thereof is selected from an antibody or antigen-binding fragment described herein. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof described herein. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies selected from antibodies 258 to 577 and 589 to 1587. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one of the antibodies neutralize SARS-CoV-2. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies selected from antibodies 292, 309, 364, 373, 388, 408, 414, 417, 419, 442, 445, 447, 462, 479, 481, 483, 488, 494, 506, 540, 549, 553, 555, and 562.

[1452] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one antibody or antigen-binding fragment thereof blocks SARS-CoV-2 binding to ACE2. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof is selected from antibodies 292, 373, 408, 417, 447, 479, 481, 483, 488, 494, 506, 549, 553, and 555.

[1453] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof binds the RBD of the SARS-CoV-2 S protein. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof is an antibody selected from 292, 447, 481, 488, 494, 506, 549, 553, and 555. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one of the anti-SARS-CoV-2 antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364.

[1454] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof binds the N-terminal domain (NTD) of the SARS-CoV-2 S protein. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof is an antibody selected from antibodies 417, 419, and 479.

[1455] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof binds the S2 domain/subunit of the SARS-CoV-2 S protein. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof is an antibody selected from antibodies 309, 364, 373, 388, 442, 462, 540, and 562.

[1456] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof that bind different epitopes of the SARS-CoV-2 S protein. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising a first anti-SARS-CoV-2 antibody or antigen-binding fragment thereof that binds a first epitope in the receptor-binding domain (RBD) of the SARS-CoV-2 S protein, and a second anti-SARS-CoV-2 antibody or antigen-binding fragment thereof that binds a second epitope of the SARS-CoV-2 S protein, wherein the second epitope is different from the first epitope.

[1457] Other suitable anti-SARS-CoV-2 antibodies that can be used in the composition include the antibodies described in Shi, R., et al., A human neutralizing antibody targets the receptor binding site of SARS-CoV-2. *Nature* 2020, <https://doi.org/10.1038/s41586-020-2381-y>, e.g., CB6 or CA1. According to the authors, the sequences of CA1 and CB6 antibodies have been deposited in GenBank with the accession codes MT470194-MT470197, with MT470194 and MT470195 encoding the light and heavy chain of CA1; and MT470196 and MT470197 encoding the light and heavy chain of CB6.

[1458] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one of the anti-SARS-CoV-2 antibody is an anti-SARS-CoV-2 antibody described herein, and the other antibody is CA1 or CB6. In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one of the anti-SARS-CoV-2 antibody is 555, and the other antibody is CB6. 555 is also known as bamlanivimab; and CB6 is also known as etesevimab.

[1459] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof, wherein at least one antibody or antigen-binding fragment thereof binds the RBD of the SARS-CoV-2 S protein, and at least one antibody or antigen-binding fragment thereof binds the NTD or S2 domain of the SARS-CoV-2 S protein.

[1460] In some embodiments, provided herein are compositions (e.g., pharmaceutical compositions) comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof that are both ACE2-blocking and bind to an epitope in the receptor-binding domain (RBD) or the N-terminal domain (NTD) of the SARS-CoV-2 S protein. In particular embodiments, at least one antibody or antigen-

binding fragment thereof comprises: (i) three CDRs in the VH set forth as SEQ ID NO: 729 and three CDRs in the VL set forth as SEQ ID NO: 730; (ii) three CDRs in the VH set forth as SEQ ID NO: 779 and three CDRs in the VL set forth as SEQ ID NO: 780; (iii) three CDRs in the VH set forth as SEQ ID NO: 1039 and three CDRs in the VL set forth as SEQ ID NO: 1040; (iv) three CDRs in the VH set forth as SEQ ID NO: 1107 and three CDRs in the VL set forth as SEQ ID NO: 1108; (v) three CDRs in the VH set forth as SEQ ID NO: 1121 and three CDRs in the VL set forth as SEQ ID NO: 1122. (vi) three CDRs in the VH set forth as SEQ ID NO: 1133 and three CDRs in the VL set forth as SEQ ID NO: 1134; (vii) three CDRs in the VH set forth as SEQ ID NO: 1157 and three CDRs in the VL set forth as SEQ ID NO: 1158; (viii) three CDRs in the VH set forth as SEQ ID NO: 1243 and three CDRs in the VL set forth as SEQ ID NO: 1244; (ix) three CDRs in the VH set forth as SEQ ID NO: 1251 and three CDRs in the VL set forth as SEQ ID NO: 1252; or (ix) three CDRs in the VH set forth as SEQ ID NO: 1255 and three CDRs in the VL set forth as SEQ ID NO: 1256; wherein the CDRs are defined by Kabat, Chothia or MacCallum numbering.

[1461] In other embodiments, at least one antibody or antigen-binding fragment thereof that is both ACE2-blocking and binds to an epitope in the receptor-binding domain (RBD) or the N-terminal domain (NTD) of the SARS-CoV-2 S protein comprises: (i) the VH set forth as SEQ ID NO: 729 and the VL set forth as SEQ ID NO: 730; (ii) the VH set forth as SEQ ID NO: 779 and the VL set forth as SEQ ID NO: 780; (iii) the VH set forth as SEQ ID NO: 1039 and the VL set forth as SEQ ID NO: 1040; (iv) the VH set forth as SEQ ID NO: 1107 and the VL set forth as SEQ ID NO: 1108; (v) the VH set forth as SEQ ID NO: 1121 and the VL set forth as SEQ ID NO: 1122; (vi) the VH set forth as SEQ ID NO: 1133 and the VL set forth as SEQ ID NO: 1134; (vii) the VH set forth as SEQ ID NO: 1157 and the VL set forth as SEQ ID NO: 1158; (viii) the VH set forth as SEQ ID NO: 1243 and the VL set forth as SEQ ID NO: 1244; (ix) the VH set forth as SEQ ID NO: 1251 and the VL set forth as SEQ ID NO: 1252; or (x) the VH set forth as SEQ ID NO: 1255 and the VL set forth as SEQ ID NO: 1256.

[1462] In particular embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) are neutralizing antibodies. In still other embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) are depleting antibodies. Moreover, as with the aforementioned fusion constructs, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be conjugated, linked, or otherwise associated with selected cytotoxic agents, polymers, biological response modifiers or the like to provide passive immunotherapy with various (and optionally multiple) mechanisms of action.

[1463] Donors who have been infected with (and have recovered from) SARS-CoV-2 are an ideal source of immune cells to discover, test, develop and manufacture antibodies as therapeutic treatments, prophylactic countermeasures, and diagnostic reagents to rapidly address emerging and recurring viral threats. Such antibodies can be discovered from a blood sample drawn from a donor that has been infected with SARS-CoV-2, a minimum of 2 weeks, e.g., a minimum of 7-8 weeks, prior to the blood draw. As such, the earliest confirmed patients, and even index patients can be a source of therapeutic, prophylactic, and diagnostic antibodies, using a minimally-invasive blood draw. The risk

of viral escape from antigenic drift can be mitigated by surveillance of the infected population and identification of cross-reactive antibodies that recognize several viral strains. However, a rapid response that significantly shortens the drug discovery process, encompassing discovery, testing and manufacture of therapeutic, prophylactic, and diagnostic antibodies from index or early patients is key to quickly preventing emerging viral threats from becoming a threat to global health, and financial and political stability.

[1464] Those skilled in the art will appreciate that anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be used in conjugated or unconjugated form. That is, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) may be associated with or conjugated to (e.g. covalently or non-covalently) pharmaceutically active compounds, biological response modifiers, diagnostic moieties, or biocompatible modifiers. In this respect it will be understood that such conjugates may comprise peptides, polypeptides, proteins, fusion proteins, nucleic acid molecules, small molecules, mimetic agents, synthetic drugs, inorganic molecules, organic molecules, and radioisotopes. Moreover, a conjugate may be covalently or non-covalently linked to the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) in various molar ratios depending, at least in part, on the method used to effect the conjugation.

[1465] As used herein, the term “antibody” or “immunoglobulin” are used interchangeably and in the broadest sense and cover both intact molecules and immunologically-reactive fragments thereof. These terms cover, for example, synthetic antibodies, monoclonal antibodies, oligoclonal or polyclonal antibodies, multiclinal antibodies, recombinantly produced antibodies, intrabodies, multispecific antibodies, bispecific antibodies, monovalent antibodies, multivalent antibodies, human antibodies, humanized antibodies, chimeric antibodies, CDR-grafted antibodies, primatized antibodies, Fab fragments, F(ab') fragments, F(ab')₂ fragments, F(ab)c fragments, single-chain FvFc (scFvFc), single-chain Fvs (scFv), Dabs, nanobodies, anti-idiotypic (anti-Id) antibodies, and any other immunologically-reactive/antigen-binding fragments thereof.

[1466] Antibodies are grouped into five distinct classes that can be distinguished biochemically, and, depending on the amino acid sequence of the constant domain of their heavy chains, can readily be assigned to the appropriate class. For historical reasons, the major classes of intact antibodies are termed IgA, IgD, IgE, IgG, and IgM. In humans, the IgG and IgA classes may be further divided into recognized subclasses (isotypes), i.e., IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 depending on structure and certain biochemical properties. It will be appreciated that the IgG isotypes in humans are named in order of their abundance in serum with IgG1 being the most abundant.

[1467] While all five classes of antibodies (i.e. IgA, IgD, IgE, IgG, and IgM) and all isotypes (i.e., IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2), as well as variations thereof, are within the scope of the present disclosure, preferred embodiments belonging to the IgG class are discussed in some detail solely for the purposes of illustration. For example, in some embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein have IgG1 isotype. It will be understood that such disclosure is, however, merely demonstrative of exemplary compositions and methods and is not in any way limiting.

[1468] In this respect, human IgG immunoglobulins comprise two identical light polypeptide chains of molecular weight approximately 23,000 Daltons, and two identical heavy chains of molecular weight 53,000-70,000 depending on the isotype. Heavy-chain constant domains that correspond to the different classes of antibodies are denoted by the corresponding lower case Greek letter α , δ , ϵ , γ , and μ , respectively. The light chains of the antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains. Those skilled in the art will appreciate that the subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

[1469] The four chains are joined by disulfide bonds in a “Y” configuration wherein the light chains bracket the heavy chains starting at the mouth of the “Y” and continuing through the variable region to the dual ends of the “Y”. Each light chain is linked to a heavy chain by one covalent disulfide bond while two disulfide linkages in the hinge region join the heavy chains. The respective heavy and light chains also have regularly spaced intrachain disulfide bridges the number of which may vary based on the isotype of IgG.

[1470] Each heavy chain has at one end a variable region (V_H) followed by a number of constant regions. Each light chain has a variable region at one end (V_L) and a constant region at its other end; the constant region of the light chain is aligned with the first constant region of the heavy chain, and the light chain variable region is aligned with the variable region of the heavy chain. The variable regions of both the light (V_L) and heavy (V_H) chain portions determine antigen recognition and specificity. Conversely, the constant regions of the light chain (CL) and the heavy chain (C_H1 , C_H2 or C_H3) confer and regulate important biological properties such as secretion, transplacental mobility, circulation half-life, complement binding, and the like. By convention, the numbering of the constant region regions increases as they become more distal from the antigen binding site or amino-terminus of the antibody. Thus, the amino or N-terminus of the antibody comprises the variable region and the carboxy or C-terminus comprises the constant region. Thus, the C_H3 and C_L regions actually comprise the carboxy-terminus of the heavy and light chain, respectively.

[1471] Portions of the variable regions of both the heavy chain and light chain differ extensively in sequence among immunoglobulins and these hypervariable sites largely define the binding and specificity of a particular antibody. These hypervariable sites manifest themselves in three segments, known as complementarity determining regions (CDRs). The more highly conserved portions of variable regions flanking the CDRs are termed framework regions (FRs). More specifically, in naturally occurring monomeric IgG antibodies, the six CDRs present on each arm of the antibody are short, non-contiguous sequences of amino acids that are specifically positioned to form the antigen-binding site as the antibody assumes its three-dimensional configuration in vivo or in vitro. CDRs encompass amino acid residues identified using any sequence or structure-based method or nomenclature system known in the art and as described below.

[1472] By way of example, CDRs may be defined using the nomenclature described by Kabat et al. (1991, NIH Publication 91-3242, National Technical Information Ser-

vice, Springfield, Va.), specifically, residues 31-35 (CDR-H1), 50-65 (CDR-H2), and 95-102 (CDR-H3) in the heavy chain variable region and residues 24-34 (CDR-L1), 50-56 (CDR-L2), and 89-97 (CDR-L3) in the light chain variable region.

[1473] By way of example, CDRs may also be defined using the nomenclature described by Chothia et al. (*J. Mol. Biol.* 196:901-917 (1987); *Nature* 342, pp. 877-883 (1989)), specifically, residues 26-32 (CDR-H1), 50-58 (CDR-H2), and 95-102 (CDR-H3) in the heavy chain variable region and residues 23-34 (CDR-L1), 50-56 (CDR-L2), and 89-97 (CDR-L3) in the light chain variable region.

[1474] By way of example, CDRs may also be defined using the nomenclature described by MacCallum et al. (*J. Mol. Biol.* 262:732-745 (1996), specifically, residues 30-35 (CDR-H1), 47-58 (CDR-H2), and 93-101 (CDR-H3) in the heavy chain variable region and residues 30-36 (CDR-L1), 46-55 (CDR-L2), and 89-96 (CDR-L3) in the light chain variable region.

[1475] CDRs vary considerably from antibody to antibody (and by definition will not exhibit homology with the Kabat consensus sequences). Maximal alignment of framework residues frequently requires the insertion of spacer residues in the numbering system, to be used for the Fv region. In addition, the identity of certain individual residues at any given Kabat site number may vary from antibody chain to antibody chain due to interspecies or allelic divergence.

[1476] One skilled in the art could readily define, identify derive and/or enumerate the CDRs as defined by Kabat et al., Chothia et al. or MacCallum et al. for each respective antibody heavy and light chain sequence set forth herein. Accordingly, each of the subject CDRs and antibodies comprising CDRs defined by all such nomenclature are expressly included within the scope of the instant disclosure.

[1477] The framework regions comprise the remainder of the heavy and light chain variable regions and are thus comprised of a non-contiguous sequence between about 100-120 amino acids in length. For example, using the nomenclature of Kabat et al., framework region 1 corresponds to the region of the variable region encompassing amino acids 1-30; framework region 2 corresponds to the region of the variable region encompassing amino acids 36-49; framework region 3 corresponds to the region of the variable region encompassing amino acids 66-94, and framework region 4 corresponds to the region of the variable region from amino acids 103 to the end of the variable region. Similarly, using the definition of CDRs by Chothia et al. (e.g., CDR-L1 23-34, CDR-L2 50-56, CDR-L3 89-97; CDR-H1 26-32, CDR-H2 50-58, CDR-H3 95-102) or MacCallum et al., the framework region boundaries are separated by the respective CDR termini as described above.

[1478] The framework regions show less inter-molecular variability in amino acid sequence and largely adopt a β -sheet conformation and the CDRs form loops which connect, and in some cases form part of, the β -sheet structure. Thus, these framework regions act to form a scaffold that provides for positioning the six CDRs in correct orientation by inter-chain, non-covalent interactions. The antigen-binding site formed by the positioned CDRs defines a surface complementary to the epitope on the immunoreactive antigen. This complementary surface promotes the non-covalent binding of the antibody to the immunoreactive antigen epitope.

[1479] All or part of the heavy and light chain variable regions may be recombined or engineered using standard recombinant and expression techniques to provide improve one or more properties of the resultant antibody. That is, the heavy or light chain variable region from a first antibody (or any portion thereof) may be mixed and matched with any selected portion of the heavy or light chain variable region from a second antibody. For example, in one embodiment, the entire light chain variable region comprising the three light chain CDRs of a first antibody may be paired with the entire heavy chain variable region comprising the three heavy chain CDRs of a second antibody. Moreover, in other embodiments, individual heavy and light chain CDRs derived from various antibodies may be mixed and matched to provide the desired antibody having optimized characteristics. Thus, an exemplary antibody may comprise three light chain CDRs from a first antibody, two heavy chain CDRs derived from a second antibody and a third heavy chain CDR from a third antibody.

[1480] With the aforementioned structural considerations in mind, those skilled in the art will appreciate that the antibodies of the present invention may comprise any one of a number of functional embodiments. In this respect, compatible antibodies may comprise any immunoreactive antibody (as the term is defined herein) that provides the desired physiological response in a subject. While any of the disclosed antibodies may be used in conjunction with the present teachings, certain embodiments of the invention will comprise chimeric, humanized, or human monoclonal antibodies or immunoreactive fragments thereof. Yet other embodiments may, for example, comprise homogeneous or heterogeneous multimeric constructs, Fc variants and conjugated or glycosylationally-altered antibodies. Moreover, it will be understood that such configurations are not mutually exclusive and that compatible individual antibodies may comprise one or more of the functional aspects disclosed herein. For example, a compatible antibody may comprise a single chain diabody with humanized variable regions or a fully human full length IgG3 antibody with Fc modifications that alter the glycosylation pattern to modulate serum half-life. Other exemplary embodiments are readily apparent to those skilled in the art and may easily be discernable as being within the scope of the invention.

[1481] Antibodies produced by naïve libraries (either natural or synthetic) can be of moderate affinity (K_a of about 10^6 to 10^7 M^{-1}), but affinity maturation can also be mimicked in vitro by constructing and reselecting from secondary libraries as described in the art. For example, mutations can be introduced at random in vitro by using error-prone polymerase.

[1482] Additionally, affinity maturation can be performed by randomly mutating one or more CDRs, e.g. using PCR with primers carrying random sequence spanning the CDR of interest, in selected individual Fv clones and screening for higher affinity clones. Another approach is to recombine the VH or VL regions selected by phage display with repertoires of naturally occurring variable region variants obtained from unimmunized donors and screen for higher affinity in several rounds of chain reshuffling. This technique allows the production of antibodies and antibody fragments with a dissociation constant K_a (k_{off}/k_{on}) of about 10^{-9} M or less.

[1483] Regardless of the type of antibody (e.g. chimeric, humanized, etc.), those skilled in the art will appreciate that immunoreactive or antigen-binding fragments of the same

may also be used. In the broadest sense, an antibody fragment comprises at least a portion of an intact antibody (e.g. a naturally occurring immunoglobulin). More particularly the term “fragment” refers to a part or portion of an antibody or antibody chain comprising fewer amino acid residues than an intact or complete antibody or antibody chain. As used herein, the term “antigen-binding fragment” refers to a polypeptide fragment of an immunoglobulin or antibody that binds antigen or competes with intact antibody (i.e., with the intact antibody from which they were derived) for antigen binding (i.e., specific binding). As used herein, antigen-binding fragments included an antibody light chain (V_L), an antibody heavy chain (V_H), a single chain antibody (scFv), a F(ab')₂ fragment, a Fab fragment, an Fd fragment, an Fv fragment, single region antibody fragments, diabodies, linear antibodies, single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

[1484] Those skilled in the art will also appreciate that antibody fragments can be obtained via chemical or enzymatic treatment of an intact or complete modulator (e.g., antibody or antibody chain) or by recombinant means. In this regard, while various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized de novo either chemically or by using recombinant DNA methodology.

[1485] By way of example, papain digestion of antibodies produces two identical antigen-binding fragments, called Fab fragments, each with a single antigen-binding site, and a residual Fc fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen. The Fab fragment also contains the constant region of the light chain and the first constant region (C_{H1}) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy-chain C_H1 region including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant regions bear at least one free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[1486] By way of further example, an Fv fragment is an antibody fragment that contains a complete antigen recognition and binding site. This region is made up of a dimer of one heavy and one light chain variable region in tight association, which can be covalent in nature, for example in scFv. It is in this configuration that the three CDRs of each variable region interact to define an antigen binding site on the surface of the V_H - V_L dimer. Collectively, the six CDRs or a subset thereof confer antigen binding specificity to the antibody. However, even a single variable region (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although usually at a lower affinity than the entire binding site.

[1487] In some embodiments an anti-coronavirus antibody fragment (e.g., anti-SARS-CoV-2 antibody fragment), for example, is one that comprises the Fc region, retains at least one of the biological functions normally associated with the Fc region when present in an intact antibody, such as FcRn binding, antibody half-life modulation, ADCC function and complement binding. In one embodiment, an antibody frag-

ment is a monovalent antibody that has an in vivo half-life substantially similar to an intact antibody. For example, such an antibody fragment may comprise on antigen binding arm linked to an Fc sequence capable of conferring in vivo stability to the fragment.

[1488] Those skilled in the art will also appreciate that anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be monovalent or multivalent (e.g., bivalent, trivalent, etc.). As used herein, the term “valency” refers to the number of potential binding sites associated with an antibody. Each target binding site specifically binds one target molecule or specific position or locus on a target molecule. When an antibody of the instant invention comprises more than one target binding site (multivalent), each target binding site may specifically bind the same or different molecules (e.g., may bind to different ligands or different antigens, or different epitopes or positions on the same antigen). In some embodiment, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein will be monovalent in that each binding site of the molecule will specifically bind to a single position or epitope. In other embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein will be multivalent in that they comprise more than one binding site and the different binding sites specifically associate with more than a single position or epitope. In such cases the multiple epitopes may be present on a virion or infected cell displaying viral antigen or a single epitope may be present on one molecule while a second, different epitope may be present on another molecule or surface.

[1489] As alluded to above, multivalent antibodies may immunospecifically bind to different epitopes of the desired target molecule or may immunospecifically bind to both the target molecule as well as a heterologous epitope, such as a heterologous polypeptide or solid support material. While some anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein only bind two antigens (i.e. are bispecific antibodies), antibodies with additional specificities such as tri-specific antibodies are also encompassed. Examples of bispecific antibodies include, without limitation, those with one arm directed against a SARS-CoV or SARS-CoV-2 spike protein and the other arm directed against any other antigen. Methods for making bispecific antibodies are known in the art. Traditional production of full-length bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities. Other more sophisticated compatible multispecific constructs and methods of their fabrication are also known.

[1490] In yet other embodiments, antibody variable regions with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant region sequences. The fusion preferably is with an immunoglobulin heavy chain constant region, comprising at least part of the hinge, C_{H2} , and/or C_{H3} regions. In one example, the first heavy-chain constant region (CH1) containing the site necessary for light chain binding is present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains

used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when, the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

[1491] In one embodiment of this approach, bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation.

[1492] According to another approach known in the art, a pair of antibody molecules can be engineered to maximize the percentage of heterodimers that are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 region of an antibody constant region. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory cavities of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[1493] Bispecific antibodies also include cross-linked or heteroconjugate antibodies. Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents and techniques are well known in the art.

[1494] In addition to various modifications, substitutions, additions or deletions to the variable or binding regions of anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein, those skilled in the art will appreciate that selected embodiments may also comprise substitutions or modifications of the constant region (i.e. the Fc region). More particularly, it is contemplated that anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may contain one or more additional amino acid residue substitutions, mutations and/or modifications, which result in a compound with preferred characteristics including, but not limited to: altered pharmacokinetics, increased serum half-life, increase binding affinity, reduced immunogenicity, increased production, altered Fc ligand binding, enhanced or reduced ADCC or CDC activity, altered glycosylation and/or disulfide bonds and modified binding specificity.

[1495] As used herein, the term “Fc region” defines a C-terminal region of an immunoglobulin heavy chain, including native sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accord-

ingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. A functional Fc region possesses an effector function of a native sequence Fc region. Exemplary effector functions include C1q binding; CDC; Fc receptor binding; ADCC; phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor; BCR), etc. Such effector functions generally require the Fc region to be combined with a binding region (e.g., an antibody variable region) and can be assessed using various assays as disclosed, for example, in definitions herein.

[1496] As used herein, the term “Fc receptor” or FcR describes a receptor that binds to the Fc region of an antibody. In some embodiments, an FcR is a native human FcR. In some embodiments, an FcR is one that binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of those receptors. FcγII receptors include FcγRIIA (an activating receptor) and FcγRIIB (an inhibiting receptor), which have similar amino acid sequences that differ primarily in the cytoplasmic regions thereof.

[1497] Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic region. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic region. Methods of measuring binding to FcRn are known in the art.

[1498] As used herein, “complement dependent cytotoxicity” or CDC refers to the lysing of a target cell in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule, an antibody for example, complexed with a cognate antigen. To assess complement activation, a CDC assay may be performed. Further, antibody-dependent cell-mediated cytotoxicity or ADCC refers to a form of cytotoxicity in which secreted Ig bound onto Fc receptors (FcRs) present on certain cytotoxic cells (e.g., Natural Killer (NK) cells, neutrophils, and macrophages) enables these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. Specific high-affinity IgG antibodies directed to the target arm cytotoxic cells and are absolutely required for such killing. Lysis of the target cell is extracellular, requires direct cell-to-cell contact, and does not involve complement.

[1499] Variant anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein that have altered FcR binding affinity or ADCC activity are those that have either enhanced or diminished FcR binding activity and/or ADCC activity compared to a parent or unmodified antibody or to a modulator comprising a native sequence Fc region. A variant antibody that displays increased binding to an FcR binds at least one FcR with better affinity than the parent or unmodified antibody or to a modulator comprising a native sequence Fc region. A variant antibody that displays decreased binding to an FcR, binds at least one FcR with worse affinity than the parent or unmodified antibody or to a modulator comprising a native sequence Fc region. Such variants which display decreased binding to an FcR may possess little or no appreciable binding to an FcR, e.g., 0-20% binding to the FcR compared to a native sequence

IgG Fc region, e.g. as determined by techniques well known in the art. In some embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein have enhanced ADCC activities.

[1500] As to FcRn, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein can also encompass Fc variants with modifications to the constant region that provide half-lives (e.g., serum half-lives) in a mammal, preferably a human, of greater than 5 days, greater than 10 days, greater than 15 days, preferably greater than 20 days, greater than 25 days, greater than 30 days, greater than 35 days, greater than 40 days, greater than 45 days, greater than 2 months, greater than 3 months, greater than 4 months, or greater than 5 months. The increased half-lives of the antibodies (or Fc containing molecules) of the present invention in a mammal, preferably a human, results in a higher serum titer of antibodies or antibody fragments in the mammal, and thus, reduces the frequency of the administration of antibodies or antibody fragments and/or reduces the concentration of antibodies or antibody fragments to be administered. Antibodies having increased in vivo half-lives can be generated by techniques known to those of skill in the art. For example, antibodies with increased in vivo half-lives can be generated by modifying (e.g., substituting, deleting, or adding) amino acid residues identified as involved in the interaction between the Fc region and the FcRn receptor. Binding to human FcRn in vivo and serum half-life of human FcRn high affinity binding polypeptides can be assayed, e.g., in transgenic mice or transfected human cell lines expressing human FcRn, or in primates to which the polypeptides with a variant Fc region are administered.

[1501] In some embodiments, the anti-SARS-CoV-2 antibodies disclosed herein can encompass an Fc region with modifications that improve their half-lives (e.g., serum half-lives) in a human. Studies have shown that some Fc mutation that enhances FcRn binding results in increased binding to rheumatoid factor (RF), whereas some Fc mutation combinations enhance FcRn binding and prolong antibody half-life without increased binding to RF, e.g., N434A/Y436T/Q438R/S440E (ACT1), N434A/Y436V/Q438R/S440E (ACT2), M428L/N434A/Y436T/Q438R/S440E (ACT3), M428L/N434A/Y436V/Q438R/S440E (ACT4), M428L/N434A/Q438R/S440E (ACT5) (Maeda, et al., MABS 2017, 9(5): 844-853, positions numbered according to EU Index numbering). In some embodiments, the anti-SARS-CoV-2 antibodies disclosed herein comprise an Fc region comprising any set of mutations selected from ACT1, ACT2, ACT3, ACT4, or ACT5. In some embodiments, the anti-SARS-CoV-2 antibodies disclosed herein comprise an Fc region comprising the ACT5 mutations: M428L/N434A/Q438R/S440E (positions numbered according to EU Index numbering). For example, the anti-SARS-CoV-2 antibodies disclosed herein can comprise an Fc region comprising SEQ ID NO: 5367.

[1502] In still other embodiments, glycosylation patterns or compositions of the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are modified. More particularly, preferred embodiments may comprise one or more engineered glycoforms, i.e., an altered glycosylation pattern or altered carbohydrate composition that is covalently attached to a molecule comprising an Fc region. Engineered glycoforms may be useful for a variety of purposes, including but not limited to enhancing or reducing effector function, increasing the affinity of the antibody for

a target antigen or facilitating production of the antibody. In cases where reduced effector function is desired, it will be appreciated that the antibody may be engineered to express in an aglycosylated form. Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the antibody sequence. That is, one or more amino acid substitutions can be made that result in elimination of one or more variable region framework glycosylation sites to thereby eliminate glycosylation at that site. Conversely, enhanced effector functions or improved binding may be imparted to the Fc containing molecule by engineering in one or more additional glycosylation sites.

[1503] Additionally, or alternatively, an Fc variant can be made that has an altered glycosylation composition, such as a hypofucosylated antibody having reduced amounts of fucosyl residues or an antibody having increased bisecting GlcNAc structures. These and similar altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Engineered glycoforms may be generated by any method known to one skilled in the art, for example by using engineered or variant expression strains, by co-expression with one or more enzymes (for example N-acetylglucosaminyltransferase III (GnTII 1)), by expressing a molecule comprising an Fc region in various organisms or cell lines from various organisms or by modifying carbohydrate(s) after the molecule comprising Fc region has been expressed.

[1504] In some embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein have reduced fucosylation. In some embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein comprise an Fc region comprising N-glycoside-linked sugar chains bound to the Fc region, wherein the sugar chains do not contain fucose. In some embodiments, the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein have increased ADCC activities, compared to the same antibodies comprising N-glycoside-linked sugar chains that comprise fucose.

[1505] Characterization of Anti-Coronavirus Antibodies

[1506] No matter how obtained or which of the aforementioned forms the antibody modulator takes (e.g., humanized, human, etc.), anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein exhibit one or more desirable characteristics. Thus, anti-coronavirus antibody-producing cells (e.g., human B cells, hybridomas or yeast colonies) may be selected, cloned and further screened for these desirable characteristics including, for example, robust growth, high antibody production and, as discussed in more detail below, desirable antibody characteristics. Hybridomas can be expanded in vivo in syngeneic animals, in animals that lack an immune system, e.g., nude mice, or in cell culture in vitro. Methods of selecting, cloning and expanding hybridomas and/or colonies, each of which produces a discrete antibody species, are well known to those of ordinary skill in the art.

[1507] For example, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be characterized by their ability to neutralize virions via the recognition of viral surface antigens essential for receptor binding and/or entry into host cells. Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may also be characterized by their ability to eliminate infected cells displaying viral antigens at their surface through complement-dependent cytotoxicity (CDC), antibody-dependent cellular phagocytosis (ADCP),

or antibody-dependent cell-mediated cytotoxicity (ADCC). Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may also be characterized by their ability to directly impact viral propagation via antibody-dependent, cell-mediated virus inhibition (ADCVI). Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may also be characterized by their epitope specificity or a number of different physical characteristics including, e.g., binding affinities, melting temperature (T_m), and isoelectric points.

[1508] In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are neutralizing antibodies. As used herein, the term “neutralizing antibody” refers to an antibody that binds to or interacts with a virion and prevents binding or association of the virion with a host cell and/or entry into a host cell or acts as an egress inhibitor insofar as the antibody may not appear to be a neutralizing antibody in a conventional *in vitro* neutralization assay, but the antibody still inhibits propagation of the viral infection.

[1509] Examples of neutralization assays include conventional neutralization assays based on the inhibition of a virus cytopathic effect (CPE) on cells in culture. For example, neutralization may be tested by reducing or blocking formation of CPE in cells infected with SARS-CoV or SARS-CoV-2. Virus and antibody may be premixed before addition to cells, followed by measuring blocking of virus entry. Hemagglutinin inhibition (HI) may be tested *in vitro* and can detect the blocking of a virus’ ability to bind to red blood cells. Antibodies that block the sialic acid receptor binding site will neutralize virus binding to cells, thereby blocking infection. Neutralization assays can also detect blocking of virus egress, as in the case of neuraminidase inhibitors like TAMIFLU®. In some embodiments, the neutralization assay can be a pseudoneutralization assay, for example, as described in the Examples below. In some embodiments, the neutralization assay can be a live virus neutralization assay, for example, as described in the Examples below.

[1510] In other embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are depleting antibodies. As used herein, the term “depleting antibody” refers to an antibody binds to or associates a SARS-CoV-2 antigen on or near the surface of an infected cell and induces, promotes or causes the death, incapacitation or elimination of the cell (e.g., by complement-dependent cytotoxicity (CDC), antibody-dependent cellular cytotoxicity (ADCC), or antibody-dependent cellular phagocytosis (ADCP)). Preferably a depleting antibody will be able to remove, incapacitate, eliminate, or kill at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, or 99% of infected cells in a defined cell population.

[1511] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may also be characterized by their epitope specificity. Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein will associate with, or bind to, discrete epitopes or determinants presented by the selected target(s). As used herein, the term “epitope” refers to that portion of the target antigen capable of being recognized and specifically bound by a particular antibody. Epitopes can be formed both from contiguous amino acids and noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained upon protein denaturing, whereas epitopes formed by tertiary folding are typically lost upon protein denaturing. An epitope typically

includes at least 3, and more usually, at least 5 to 10 amino acids in a unique spatial conformation. More specifically, the skilled artisan will appreciate the term epitope includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor or otherwise interacting with a molecule. Epitopic determinants generally consist of chemically active surface groupings of molecules such as amino acids or carbohydrate or sugar side chains and generally have specific three-dimensional structural characteristics, as well as specific charge characteristics. Additionally, an epitope may be linear or conformational. In a linear epitope, all of the points of interaction between the protein and the interacting molecule (such as an antibody) occur linearly along the primary amino acid sequence of the protein. In a conformational epitope, the points of interaction occur across amino acid residues on the protein that are linearly separated from one another.

[1512] Once a desired epitope on an antigen is determined, it is possible to generate antibodies to that epitope, e.g., by immunizing with a peptide comprising the epitope using techniques known in the art. Alternatively, during the discovery process, the generation and characterization of antibodies may elucidate information about desirable epitopes. From this information, it is then possible to competitively screen antibodies for binding to the same epitope. An approach to achieve this is to conduct competition studies to find antibodies that competitively bind with one another, i.e. the antibodies compete for binding to the antigen. A high throughput process for binning antibodies based upon their cross-competition is known in the art.

[1513] As used herein, the term “binning” refers to a method to group antibodies based on their antigen binding characteristics. The grouping is somewhat arbitrary, depending on how different the observed binding patterns of the antibodies tested. Thus, while the technique is a useful tool for categorizing antibodies, the bins do not always directly correlate with epitopes and such initial determinations of epitope binding should be further confirmed by other art recognized methodology.

[1514] With this caveat one can determine whether a selected primary antibody (or fragment thereof) binds to the same epitope or cross competes for binding with a second antibody by using methods known in the art. In one embodiment, one exposes virions or one or more infected cell(s) to a first antibody under saturating conditions and then measures the ability of a second antibody to bind to the same population of virions or cell(s). If the second antibody is able to bind, then the second antibody binds to a different epitope than the first antibody. However, if the second antibody is not able to bind, then the second antibody binds to the same epitope, an overlapping epitope, or an epitope that is in close proximity to the epitope bound by the first antibody. As known in the art, the desired data can be obtained using solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay, surface plasmon resonance, bio-layer interferometry, or flow cytometric methodology.

[1515] As used herein, the term “compete” means competition between antibodies as determined by an assay in which the antibody or immunologically-reactive fragment under test prevents or inhibits specific binding of a reference antibody to a common antigen. Typically, such an assay involves the use of purified antigen bound to a solid surface or cells bearing either of these, an unlabeled test immuno-

globulin and a labeled reference immunoglobulin. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur. Additional details regarding methods for determining competitive binding are provided in the Examples herein. Usually, when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 40%, 45%, 50%, 55%, 60%, 65%, 70% or 75%. In some instance, binding is inhibited by at least 80%, 85%, 90%, 95%, or 97% or more.

[1516] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may also be characterized by their binding affinity. In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) specifically bind to a target antigen expressed on a virion or infected cell, i.e., the dissociation constant K_d (k_{off}/k_{on}) is $\leq 10^{-8}$ M. The antibody specifically binds it antigen with high affinity when the K_d is $\leq 5 \times 10^{-9}$ M, and with very high affinity when the K_d is $\leq 5 \times 10^{-10}$ M. In particular embodiments, the antibody has a K_d of $\leq 10^{-9}$ M and an off-rate of about 1×10^{-4} /sec. In other embodiments, the off-rate is $\leq 1 \times 10^{-5}$ /sec. In other embodiments, the antibodies will bind with a K_d of between about 10^{-8} M and 10^{-10} M, and in yet other embodiments, antibodies bind with a $K_d \leq 2 \times 10^{-10}$ M. Still other selected embodiments comprise antibodies that have a dissociation constant or K_d (k_{off}/k_{on}) of less than 10^{-2} M, less than 5×10^{-2} M, less than 10^{-3} M, less than 5×10^{-3} M, less than 10^{-4} M, less than 5×10^{-4} M, less than 10^{-5} M, less than 5×10^{-5} M, less than 10^{-6} M, less than 5×10^{-6} M, less than 10^{-7} M, less than 5×10^{-7} M, less than 10^{-8} M, less than 5×10^{-8} M, less than 10^{-9} M, less than 5×10^{-9} M, less than 10^{-10} M, less than 5×10^{-10} M, less than 10^{-11} M, less than 5×10^{-11} M, less than 10^{-12} M, less than 5×10^{-12} M, less than 10^{-13} M, less than 5×10^{-13} M, less than 10^{-14} M, less than 5×10^{-14} M, less than 10^{-15} M, or less than 5×10^{-15} M.

[1517] In another embodiment, an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) that specifically binds to its antigen (e.g., on a virion or infected cell displaying a viral antigen) has an association rate constant or k_{on} rate ((Ab)+ antigen (Ag) $\xrightarrow{k_{on}}$ Ab-Ag) of at least 10^5 M $^{-1}$ s $^{-1}$, at least 2×10^5 M $^{-1}$ s $^{-1}$, at least 5×10^5 M $^{-1}$ s $^{-1}$, at least 10^6 M $^{-1}$ s $^{-1}$, at least 5×10^6 M $^{-1}$ s $^{-1}$, at least 10^7 M $^{-1}$ s $^{-1}$, at least 5×10^7 M $^{-1}$ s $^{-1}$, or at least 10^8 M $^{-1}$ s $^{-1}$.

[1518] In another embodiment, an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) that specifically binds to its antigen (e.g., on a virion or infected cell displaying a viral antigen) has a dissociation rate constant or k_{off} rate ((Ab)+ antigen (Ag) $\xleftarrow{k_{off}}$ Ab-Ag) of less than 10^{-1} s $^{-1}$, less than 5×10^{-1} s $^{-1}$, less than 10^{-2} s $^{-1}$, less than 5×10^{-2} s $^{-1}$, less than 10^{-3} s $^{-1}$, less than 5×10^{-3} s $^{-1}$, less than 10^{-4} s $^{-1}$, less than 5×10^{-4} s $^{-1}$, less than 10^{-5} s $^{-1}$, less than 5×10^{-5} s $^{-1}$, less than 10^{-6} s $^{-1}$, less than 5×10^{-6} s $^{-1}$, less than 10^{-7} s $^{-1}$, less than 5×10^{-7} s $^{-1}$, less than 10^{-8} s $^{-1}$, less than 5×10^{-8} s $^{-1}$, less than 10^{-9} s $^{-1}$, less than 5×10^{-9} s $^{-1}$ or less than 10^{-10} s $^{-1}$.

[1519] In another embodiment, an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) that specifically

binds to its antigen (e.g., on a virion or infected cell displaying a viral antigen) will have an affinity constant or K_a (k_{on}/k_{off}) of at least 10^2 M $^{-1}$, at least 5×10^2 M $^{-1}$, at least 10^3 M $^{-1}$, at least 5×10^3 M $^{-1}$, at least 10^4 M $^{-1}$, at least 5×10^4 M $^{-1}$, at least 10^5 M $^{-1}$, at least 5×10^5 M $^{-1}$, at least 10^6 M $^{-1}$, at least 5×10^6 M $^{-1}$, at least 10^7 M $^{-1}$, at least 5×10^7 M $^{-1}$, at least 10^8 M $^{-1}$, at least 5×10^8 M $^{-1}$, at least 10^9 M $^{-1}$, at least 5×10^9 M $^{-1}$, at least 10^{10} M $^{-1}$, at least 5×10^{10} M $^{-1}$, at least 10^{11} M $^{-1}$, at least 5×10^{11} M $^{-1}$, at least 10^{12} M $^{-1}$, at least 5×10^{12} M $^{-1}$, at least 10^{13} M $^{-1}$, at least 5×10^{13} M $^{-1}$, at least 10^{14} M $^{-1}$, at least 5×10^{14} M $^{-1}$, at least 10^{15} M $^{-1}$ or at least 5×10^{15} M $^{-1}$.

[1520] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may also be characterized by their thermal stability as reflected by their respective melting point (Tm). The Tm of the Fab region of an antibody can be a good indicator of the thermal stability of an antibody and may further provide an indication of the shelf life. Tm is merely the temperature of 50% unfolding for a given region or sequence. A lower Tm indicates more aggregation/less stability, whereas a higher Tm indicates less aggregation/more stability. Thus, antibodies or fragments or derivatives having higher Tm are preferable. Moreover, using art-recognized techniques it is possible to alter the composition of antibodies or regions thereof to increase or optimize molecular stability. Thermal melting temperatures (Tm) of a protein region (e.g., a Fab region) can be measured using any standard method known in the art, e.g., by differential scanning calorimetry.

[1521] In some embodiments, the Fab region of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein has a Tm value higher than at least 50° C., 55° C., 60° C., 65° C., 70° C., 75° C., 80° C., 85° C., 90° C., 95° C., 100° C., 105° C., 110° C., 115° C. or 120° C. In another embodiment, the Fab region of an anti-coronavirus antibody disclosed herein has a Tm value higher than at least about 50° C., about 55° C., about 60° C., about 65° C., about 70° C., about 75° C., about 80° C., about 85° C., about 90° C., about 95° C., about 100° C., about 105° C., about 110° C., about 115° C. or about 120° C.

[1522] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may also be characterized by their isoelectric point (pI), which is generally defined as the pH at which a polypeptide carries no net charge. It is known in the art that protein solubility is typically lowest when the pH of the solution is equal to the isoelectric point (pi) of the protein. Therefore, it is possible to optimize solubility by altering the number and location of ionizable residues in the antibody to adjust the pI. For example, the pI of a polypeptide can be manipulated by making the appropriate amino acid substitutions (e.g., by substituting a charged amino acid such as a lysine, for an uncharged residue such as alanine). Without wishing to be bound by any particular theory, amino acid substitutions of an antibody that result in changes of the pI of said antibody may improve solubility and/or the stability of the antibody. One skilled in the art would understand which amino acid substitutions would be most appropriate for a particular antibody to achieve a desired p. The pI of a protein may be determined by a variety of methods including but not limited to, isoelectric focusing and various computer algorithms.

[1523] In some embodiments, the pI of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein is higher than about 6.5, about 7.0, about 7.5, about

8.0, about 8.5, or about 9.0. In another embodiment, the pI of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein is higher than 6.5, 7.0, 7.5, 8.0, 8.5, or 9.0. In yet another embodiment, substitutions resulting in alterations in the pI of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein will not significantly diminish its binding affinity. As discussed in more detail below, it is specifically contemplated that the substitution(s) of the Fc region that result in altered binding to FcγR may also result in a change in the pI. In a preferred embodiment, substitution(s) of the Fc region are specifically chosen to effect both the desired alteration in FcγR binding and any desired change in pI. As used herein, the pI value is defined as the pI of the predominant charge form.

[1524] Nucleic Acids and Anti-Coronavirus Antibody Expression

[1525] Provided herein are also nucleic acids encoding a heavy chain or light chain, or a VH or VL, of the novel anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein, and vectors comprising one or more such nucleic acids. The terms “nucleic acid” or “polynucleotide”, as used interchangeably herein, refer to polymers of nucleotides, including single-stranded and/or double-stranded nucleotide-containing molecules, such as DNA, cDNA and RNA molecules, incorporating native, modified, and/or analogs of, nucleotides. Polynucleotides of the present disclosure may also include substrates incorporated therein, for example, by DNA or RNA polymerase or a synthetic reaction. In some embodiments, provided herein are nucleic acids encoding a heavy chain or light chain of the anti-SARS-CoV-2 antibodies described herein, e.g., any one of Abs 258-1329. In some embodiments, provided herein are nucleic acids encoding a VH or VL of the anti-SARS-CoV-2 antibodies described herein, e.g., any one of Abs 258-1329. In some embodiments, provided herein are nucleic acids encoding a VH or VL comprising any one of SEQ ID NOs: 661-1320, 2751-4180, and 4749-5316. In some embodiments, provided herein are nucleic acids comprising a sequence from any one of SEQ ID NOs: 1-660, 1321-2750, and 4181-4748.

[1526] In some embodiments, provided herein are nucleic acids encoding a heavy chain or light chain of the anti-SARS-CoV-2 antibodies described herein. In some embodiments, provided herein are nucleic acids encoding a heavy chain comprising SEQ ID NO: 5363, a light chain comprising SEQ ID NO: 5364, or both a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364. In some embodiments, provided herein are nucleic acids encoding a heavy chain comprising SEQ ID NO: 5335, a light chain comprising SEQ ID NO: 5336, or both a heavy chain comprising SEQ ID NO: 5335 and a light chain comprising SEQ ID NO: 5336. In some embodiments, provided herein are nucleic acids encoding a heavy chain comprising SEQ ID NO: 5347, a light chain comprising SEQ ID NO: 5348, or both a heavy chain comprising SEQ ID NO: 5347 and a light chain comprising SEQ ID NO: 5348. In some embodiments, provided herein are nucleic acids encoding a heavy chain comprising SEQ ID NO: 5351, a light chain comprising SEQ ID NO: 5352, or both a heavy chain comprising SEQ ID NO: 5351 and a light chain comprising SEQ ID NO: 5352. In some embodiments, provided herein are nucleic acids encoding a heavy chain comprising SEQ ID NO: 5325, a light chain comprising

SEQ ID NO: 5326, or both a heavy chain comprising SEQ ID NO: 5325 and a light chain comprising SEQ ID NO: 5326.

[1527] DNA encoding the anti-coronaviruses antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding antibody heavy and light chains). Isolated and subcloned hybridoma cells (or phage or yeast derived colonies) may serve as a preferred source of such DNA. More particularly, the isolated DNA (which may be modified) can be used to clone constant and variable region sequences for the manufacture of antibodies.

[1528] One exemplary method entails extraction of RNA from the selected cells, conversion to cDNA, and amplification by PCR using antibody specific primers. Suitable primers are well known in the art, and as exemplified herein, are readily available from numerous commercial sources. It will be appreciated that, to express a recombinant human or non-human antibody isolated by screening of a combinatorial library, the DNA encoding the antibody is cloned into a recombinant expression vector and introduced into host cells including mammalian cells, insect cells, plant cells, yeast, and bacteria. In yet other embodiments, the modulators are introduced into and expressed by simian COS cells, NS0 cells, Chinese Hamster Ovary (CHO) cells or myeloma cells that do not otherwise produce the desired construct. As will be discussed in more detail below, transformed cells expressing the desired modulator may be grown up in relatively large quantities to provide clinical and commercial supplies of the fusion construct or immunoglobulin.

[1529] Whether the nucleic acid encoding an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein is obtained or derived from phage display technology, yeast libraries, hybridoma based technology, synthetically, or from commercial sources, it should be understood that the inventions disclosed herein encompass nucleic acid molecules and sequences encoding anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies), fusion proteins, or antigen-binding fragments or derivatives thereof. The inventions disclosed herein further encompass nucleic acids or nucleic acid molecules (e.g., polynucleotides) that hybridize under high stringency, or alternatively, under intermediate or lower stringency hybridization conditions (e.g., as defined below), to polynucleotides complementary to nucleic acids having a polynucleotide sequence that encodes a modulator of the invention or a fragment or variant thereof. The term nucleic acid molecule or isolated nucleic acid molecule, as used herein, is intended to include at least DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA. Moreover, the present invention comprises any vehicle or construct, incorporating such modulator encoding polynucleotide including, without limitation, vectors, plasmids, host cells, cosmid or viral constructs.

[1530] As used herein, the term “isolated nucleic acid” refers to a nucleic acid that was (i) amplified in vitro, for example by polymerase chain reaction (PCR), (ii) recombinantly produced by cloning, (iii) purified, for example by cleavage and gel-electrophoretic fractionation, or (iv) synthesized, for example by chemical synthesis. An isolated nucleic acid is a nucleic acid that is available for manipulation by recombinant DNA techniques.

[1531] More specifically, nucleic acids that encode anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein, polynucleotides sufficient for use as hybridization probes, PCR primers or sequencing primers for identifying, analyzing, mutating or amplifying a polynucleotide encoding a polypeptide, anti-sense nucleic acids for inhibiting expression of a polynucleotide, and complementary sequences of the foregoing are also encompassed herein. Such nucleic acids can be any length. They can be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 1,500, 3,000, 5,000 or more nucleotides in length, and/or can comprise one or more additional sequences, for example, regulatory sequences, and/or be part of a larger nucleic acid, for example, a vector. These nucleic acids can be single-stranded or double-stranded and can comprise RNA and/or DNA nucleotides, and artificial variants thereof (e.g., peptide nucleic acids).

[1532] As indicated, the invention further provides nucleic acids that hybridize to other nucleic acids under particular hybridization conditions. Methods for hybridizing nucleic acids are well known in the art. For example, a moderately stringent hybridization condition uses a prewashing solution containing 5× sodium chloride/sodium citrate (SSC), 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization buffer of about 50% formamide, 6×SSC, and a hybridization temperature of 55° C. (or other similar hybridization solutions, such as one containing about 50% formamide, with a hybridization temperature of 42° C.), and washing conditions of 60° C., in 0.5×SSC, 0.1% SDS. A stringent hybridization condition hybridizes in 6×SSC at 45° C., followed by one or more washes in 0.1×SSC, 0.2% SDS at 68° C. One of skill in the art can manipulate the hybridization and/or washing conditions to increase or decrease the stringency of hybridization such that nucleic acids comprising nucleotide sequences that are at least 65, 70, 75, 80, 85, 90, 95, 98 or 99% identical to each other typically remain hybridized to each other. More generally, for the purposes of the instant disclosure the term “substantially identical” with regard to a nucleic acid sequence refers to a sequence of nucleotides exhibiting at least about 85%, or 90%, or 95%, or 97% sequence identity to the reference nucleic acid sequence. The basic parameters affecting the choice of hybridization conditions and guidance for devising suitable conditions are well known, and can be readily determined by those having ordinary skill in the art based on, for example, the length and/or base composition of the nucleic acid.

[1533] It will further be appreciated that nucleic acids may be present alone or in combination with other nucleic acids, which may be homologous or heterologous. In preferred embodiments, a nucleic acid is functionally linked to expression control sequences that may be homologous or heterologous with respect to that nucleic acid. In this context, the term “homologous” means that a nucleic acid is also functionally linked to the expression control sequence naturally and the term “heterologous” means that a nucleic acid is not functionally linked to the expression control sequence naturally.

[1534] A nucleic acid, such as a nucleic acid expressing RNA and/or protein or peptide, and an expression control sequence are functionally linked to one another, if they are covalently linked to one another in such a way that expression or transcription of the nucleic acid is under the control or under the influence of said expression control sequence.

If the nucleic acid is to be translated into a functional protein, then, with an expression control sequence functionally linked to a coding sequence, induction of the expression control sequence results in transcription of the nucleic acid, without causing a frame shift in the coding sequence or the coding sequence not being capable of being translated into the desired protein or peptide. As used herein, the term “expression control sequence” includes promoters, ribosome binding sites, enhancers and other control elements that regulate transcription of a gene or translation of mRNA. In particular embodiments, the expression control sequences can be regulated. The exact structure of expression control sequences may vary as a function of the species or cell type, but generally comprises 5'-untranscribed and 5'- and 3'-untranslated sequences which are involved in initiation of transcription and translation, respectively, such as TATA box, capping sequence, CAAT sequence, and the like. More specifically, 5'-untranscribed expression control sequences comprise a promoter region that includes a promoter sequence for transcriptional control of the functionally linked nucleic acid. Expression control sequences may also comprise enhancer sequences or upstream activator sequences.

[1535] As used herein, the term “promoter” or “promoter region” relates to a nucleic acid sequence which is located upstream (5') to the nucleic acid sequence being expressed and controls expression of the sequence by providing a recognition and binding site for RNA-polymerase. The promoter region may include further recognition and binding sites for further factors that are involved in the regulation of transcription of a gene. A promoter may control the transcription of a prokaryotic or eukaryotic gene. Furthermore, a promoter may be inducible and may initiate transcription in response to an inducing agent or may be constitutive if transcription is not controlled by an inducing agent. A gene that is under the control of an inducible promoter is not expressed or only expressed to a small extent if an inducing agent is absent. In the presence of the inducing agent the gene is switched on or the level of transcription is increased. This is mediated, in general, by binding of a specific transcription factor.

[1536] Promoters for use in the production of anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein include promoters for SP6, T3 and T7 polymerase, human U6 RNA promoter, CMV promoter, and artificial hybrid promoters thereof (e.g. CMV) where a part or parts are fused to a part or parts of promoters of genes of other cellular proteins such as e.g. human GAPDH (glyceraldehyde-3-phosphate dehydrogenase), and may include (an) additional intron(s).

[1537] As used herein, the term “expression” is used in its most general meaning and comprises the production of RNA or of RNA and protein/peptide. It also comprises partial expression of nucleic acids. Furthermore, expression may be carried out transiently or stably.

[1538] In a preferred embodiment, a nucleic acid molecule present in a vector, where appropriate with a promoter, which controls expression of the nucleic acid. As used herein, the term “vector” is used here in its most general meaning and comprises any intermediary vehicle for a nucleic acid that enables the nucleic acid, for example, to be introduced into prokaryotic and/or eukaryotic cells and, where appropriate, to be integrated into a genome. Vectors of this kind are preferably replicated and/or expressed in the

cells. Vectors may comprise plasmids, phagemids, bacteriophages or viral genomes. As used herein, the term “plasmid” generally relates to a construct of extrachromosomal genetic material, usually a circular DNA duplex, which can replicate independently of chromosomal DNA. In a preferred embodiment, the expression vector is a glutamine synthetase (GS) expression vector.

[1539] It will be appreciated by those of skill in the art, that many conventional techniques in molecular biology, microbiology, and recombinant DNA technology are optionally used. Such conventional techniques relate to vectors, host cells and recombinant methods as defined herein. In addition, essentially any polynucleotide (including, e.g., labeled or biotinylated polynucleotides) can be obtained from any of a variety of commercial sources.

[1540] The inventions disclosed herein also encompass recombinant host cells allowing recombinant expression of antibodies of the invention or portions thereof. Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein produced by expression in such recombinant host cells are referred to herein as recombinant antibodies. The inventions disclosed herein also encompass progeny cells of such host cells, and anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) produced by the same. In a preferred embodiment, the host cell is a Chinese Hamster Ovary (CHO) cell.

[1541] As used herein, the term “recombinant host cell” or “host cell” means a cell into which a recombinant expression vector has been introduced. It should be understood that recombinant host cell and host cell mean not only the particular subject cell but also the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term host cell as used herein. Such cells may comprise a vector as described above.

[1542] The inventions disclosed herein also encompass methods for making anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein. According to one embodiment, such a method comprises culturing a cell transfected or transformed with a vector as described above, and isolating the antibody.

[1543] As indicated above, expression of an antibody preferably comprises expression vector(s) containing a polynucleotide that encodes the anti-coronavirus antibody. Methods that are well known to those skilled in the art can be used to construct expression vectors comprising antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Particular embodiments provide replicable vectors comprising a nucleotide sequence encoding an anti-coronavirus antibody disclosed herein operably linked to a promoter. In preferred embodiments, such vectors may include a nucleotide sequence encoding the heavy chain of an antibody molecule (or fragment thereof), a nucleotide sequence encoding the light chain of an antibody (or fragment thereof), or both the heavy and light chain.

[1544] Using art recognized molecular biology techniques and current protein expression methodology, substantial quantities of the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be produced.

More specifically, nucleic acid molecules encoding such antibodies may be integrated into well-known and commercially available protein production systems comprising various types of host cells to provide preclinical, clinical, or commercial quantities of the desired pharmaceutical product. In preferred embodiments the nucleic acid molecules encoding the antibodies are engineered into vectors or expression vectors that provide for efficient integration into the selected host cell and subsequent high expression levels of the antibody.

[1545] Preferably nucleic acid molecules encoding anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein and vectors comprising these nucleic acid molecules can be used for transfection of a suitable mammalian, plant, bacterial or yeast host cell though it will be appreciated that prokaryotic systems may also be used. Transfection can be by any known method for introducing polynucleotides into a host cell. Methods for the introduction of heterologous polynucleotides into mammalian cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be introduced into mammalian cells by viral vectors. Methods of transforming mammalian cells are well known in the art. Methods of transforming plant cells are also well known in the art, including, e.g., agrobacterium-mediated transformation, biolistic transformation, direct injection, electroporation, and viral transformation. Methods of transforming bacterial and yeast cells are also well known in the art.

[1546] Moreover, the host cell may be co-transfected with two expression vectors of the invention, for example, the first vector encoding a heavy chain polypeptide and the second vector encoding a light chain polypeptide. The two vectors may contain identical selectable markers that enable substantially equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain is preferably placed before the heavy chain to avoid an excess of toxic free heavy chain. The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

[1547] Varieties of host-expression vector systems, many commercially available, may be used to express anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein. Such host-expression systems represent vehicles by which the coding sequences of interest may be expressed and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express a molecule of the invention in situ. Such systems include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*, *Streptomyces*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing modulator coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transfected with recombinant yeast expression vectors containing modulator coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing modulator coding sequences; plant cell systems (e.g., *Nicotiana*, *Arabidopsis*, duckweed, corn, wheat, potato, etc.) infected with recom-

binant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transfected with recombinant plasmid expression vectors (e.g., Ti plasmid) containing modulator coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

[1548] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable.

[1549] In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) may be used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The coding sequences may be cloned individually into non-essential regions (for example, the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example, the polyhedrin promoter).

[1550] In mammalian host cells, a number of viral-based expression systems may be used to introduce the desired nucleotide sequence. In cases where an adenovirus is used as an expression vector, the coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E 1 or E3) will result in a recombinant virus that is viable and capable of expressing the molecule in infected hosts. Specific initiation signals may also be required for efficient translation of inserted coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, and the like. Thus, compatible mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC). These include, inter alia, Chinese hamster ovary (CHO) cells, NS0 cells, SP2 cells, HEK-293T cells, 293 Freestyle cells (Life Technologies), NIH-3T3 cells, HeLa cells, baby hamster kidney (BHK) cells, African green monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), A549 cells, and a number of other cell lines.

[1551] For long-term, high-yield production of recombinant proteins stable expression is preferred. Accordingly, cell lines that stably express the selected modulator may be engineered using standard art recognized techniques. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Fol-

lowing the introduction of the foreign DNA, engineered cells may be allowed to grow for 1 to 2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compositions that interact directly or indirectly with the molecule.

[1552] A number of selection systems are well known in the art and may be used including, but not limited to, the herpes simplex virus thymidine kinase, hypoxanthineguanine phosphoribosyltransferase, and adenine phosphoribosyltransferase genes can be employed in tk-, hgprrt- or aprt-cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate; gpt, which confers resistance to mycophenolic acid; neo, which confers resistance to the aminoglycoside G-418; and hygromycin, which confers resistance to hygromycin.

[1553] Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone. It will be appreciated by those of skill in the art that one particularly preferred method of establishing a stable, high yield cell line comprises the glutamine synthetase gene expression system (the GS system) which provides an efficient approach for enhancing expression under certain conditions.

[1554] In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function and/or purification of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. As known in the art, appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the expressed polypeptide. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product are particularly effective. Accordingly, some preferred mammalian host lines include, but are not limited to, CHO, VERY, BHK, HeLa, COS, NS0, MDCK, 293, 3T3, and W138. Depending on the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) and the selected production system, those of skill in the art may easily select and optimize appropriate host cells for efficient expression of the antibody.

[1555] Those of skill in the art will appreciate that anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be chemically synthesized using techniques known in the art. For example, a peptide corresponding to a polypeptide fragment of the invention can be synthesized by use of a peptide synthesizer. If desired, non-genetically encoded amino acids or synthetic amino acids can be substituted or added into a polypeptide sequence.

[1556] Representative non-genetically encoded amino acids include but are not limited to 2-amino adipic acid;

3-aminoadipic acid; P-aminopropionic acid; 2-aminobutyric acid; 4-aminobutyric acid (piperidinic acid); 6-aminocaproic acid; 2-aminoheptanoic acid; 2-aminoisobutyric acid; 3-aminoisobutyric acid; 2-aminopimelic acid; 2,4-diaminobutyric acid; desmosine; 2,2'-diaminopimelic acid; 2,3-diaminopropionic acid; N-ethylglycine; N-ethylasparagine; hydroxylysine; allo-hydroxylysine; 3-hydroxyproline; 4-hydroxyproline; isodesmosine; allo-isoleucine; N-methylglycine (sarcosine); N-methylisoleucine; N-methylvaline; norvaline; norleucine; and ornithine.

[1557] Representative synthetic amino acids include, for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-im-benzylhistidine.

[1558] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein also can be produced transgenically through the generation of a mammal or plant that is transgenic for the immunoglobulin heavy and light chain sequences (or fragments or derivatives or variants thereof) of interest and production of the desired compounds in a recoverable form. For example, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies), can be produced in, and recovered from, e.g., the milk of goats, cows, or other mammals. In some embodiments, non-human transgenic animals that comprise human immunoglobulin loci are immunized with SARS-CoV-2 virions or an immunogenic portion thereof, as described above.

[1559] Non-human transgenic animals or plants may be produced by introducing one or more nucleic acid molecules encoding an anti-coronavirus antibody into the animal or plant by standard transgenic techniques. The transgenic cells used for making the transgenic animal can be embryonic stem cells or somatic cells or a fertilized egg. The transgenic non-human organisms can be chimeric, nonchimeric heterozygotes, and nonchimeric homozygotes. In some embodiments, the transgenic non-human animals have a targeted disruption and replacement by a targeting construct that encodes, for example, a heavy chain and/or a light chain of interest. While anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be produced in any transgenic animal, particularly preferred embodiments include mice, rats, sheep, pigs, goats, cattle, and horses. In particular embodiments, the non-human transgenic animal expresses the desired pharmaceutical product in blood, milk, urine, saliva, tears, mucus, and other bodily fluids from which it is readily obtainable using art recognized purification techniques.

[1560] It is likely that modulators, including antibodies, expressed by different cell lines or in transgenic animals will have different glycosylation patterns from each other. However, the instant inventions encompass anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) encoded by the nucleic acid molecules provided herein, or comprising the amino acid sequences provided herein, regardless of the glycosylation state of the molecule, and more generally, regardless of the presence or absence of post-translational modification(s). The instant inventions also encompass anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies)

that are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, etc. Various post-translational modifications are also encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of prokaryotic host cell expression. Moreover, as set forth in the text and below the anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may also be modified with a detectable label, such as an enzymatic, fluorescent, radioisotopic or affinity label to allow for their detection and isolation.

[1561] Once anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein have been produced by recombinant expression or any one of the other techniques disclosed herein, they may be purified by any method known in the art for purification of immunoglobulins, or more generally by any other standard technique for the purification of proteins. In this respect the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) may be isolated. As used herein, an "isolated anti-coronavirus antibody" is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses of the antibody and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. Isolated anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) include antibodies in situ within recombinant cells because at least one component of the antibody's natural environment will not be present.

[1562] When using recombinant techniques, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the desired molecule is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, may be removed, for example, by centrifugation or ultrafiltration. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

[1563] Compositions comprising anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein prepared from cells can be purified using, for example, hydroxyapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc region that is present in the selected construct. Protein A can be used to purify antibodies that are based on human IgG1, IgG2 or IgG4 heavy chains. Protein

G is recommended for all mouse isotypes and for human IgG3. The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically-stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_H3 region, the BAKERBOND ABX™ resin is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin, SEPHAROSE® chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE and ammonium sulfate precipitation are also available depending on the antibody to be recovered. In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are purified, at least in part, using Protein A or Protein G affinity chromatography.

[1564] By way of illustration of the foregoing, cDNA sequences encoding an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) heavy chain and light chain may be cloned and engineered into a glutamine synthetase (GS) expression vector. The engineered immunoglobulin expression vector may then be stably transfected into CHO cells. As one skilled in the art will appreciate, mammalian expression of antibodies will result in glycosylation, typically at highly conserved N-glycosylation sites in the Fc region. Stable clones may be verified for expression of an antibody specifically binding to SARS-CoV spike protein or SARS-CoV-2 spike protein. Positive clones may be expanded into serum-free culture medium for antibody production in bioreactors. Medium, into which an antibody has been secreted, may be purified by conventional techniques. For example, the medium may be conveniently applied to a Protein A or G SEPHAROSE® FF column that has been equilibrated with a compatible buffer, such as phosphate buffered saline. The column is washed to remove nonspecific binding components. The bound antibody is eluted, for example, by pH gradient and antibody fractions are detected, such as by SDS-PAGE, and then pooled. The antibody may be concentrated and/or sterile filtered using common techniques. Soluble aggregate and multimers may be effectively removed by common techniques, including size exclusion, hydrophobic interaction, ion exchange, or hydroxyapatite chromatography. The product may be immediately frozen, for example, at -70°C. , or may be lyophilized.

[1565] Anti-Coronavirus Antibody Conjugates

[1566] Once anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein have been purified, they may be linked with, fused to, conjugated to (e.g., covalently or non-covalently) or otherwise associated with diagnostic moieties or biocompatible modifiers. As used herein the term “conjugate” means any molecule associated with an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein regardless of the method of association. In this respect it will be understood that such conjugates may comprise peptides, polypeptides, proteins, polymers, nucleic acid molecules, small molecules, mimetic agents, synthetic drugs, inorganic molecules, organic molecules and radioisotopes. Moreover, as indicated above the selected conjugate may be covalently or non-covalently

linked to the antibody and exhibit various molar ratios depending, at least in part, on the method used to effect the conjugation.

[1567] In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be conjugated or associated with proteins, polypeptides or peptides that impart selected characteristics (e.g., biotoxins, biomarkers, purification tags, etc.). In particular embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a heterologous protein or polypeptide wherein the polypeptide comprises at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids. The construct does not necessarily need to be directly linked, but may occur through linker sequences. For example, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be used to target heterologous polypeptides to virions or infected cells, either in vitro or in vivo, by fusing or conjugating the antibodies to other antibodies specific for other antigens. Moreover, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) that are fused or conjugated to heterologous polypeptides may also be used in in vitro immunoassays and may be compatible with purification methodology known in the art.

[1568] In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be conjugated or otherwise associated with biocompatible modifiers that may be used to adjust, alter, improve, or moderate antibody properties. For example, antibodies or fusion constructs with increased in vivo half-lives can be generated by attaching relatively high molecular weight polymer molecules such as commercially available polyethylene glycol (PEG) or similar biocompatible polymers. Those skilled in the art will appreciate that PEG may be obtained in many different molecular weight and molecular configurations that can be selected to impart specific properties to the antibody (e.g. the half-life may be tailored). PEG can be attached to modulators or antibody fragments or derivatives with or without a multifunctional linker either through site-specific conjugation of the PEG to the N- or C-terminus of antibodies or via epsilon-amino groups present on lysine residues. Linear or branched polymer derivatization that results in minimal loss of biological activity may be used. The degree of conjugation can be closely monitored by SDS-PAGE and mass spectrometry to ensure optimal conjugation of PEG molecules to antibody molecules. Unreacted PEG can be separated from antibody-PEG conjugates by, e.g., size exclusion or ion-exchange chromatography. In a similar manner, the disclosed modulators can be conjugated to albumin in order to make the antibody or antibody fragment more stable in vi-o or have a longer half-life in vivo. The techniques are well known in the art. Other biocompatible conjugates are evident to those of ordinary skill and may readily be identified and utilized in accordance with the teachings herein.

[1569] In other embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are conjugated to a diagnostic or detectable agent, marker or reporter which may be a biological molecule (e.g., a peptide or nucleotide), a small molecule, fluorophore, or radioisotope. Labeled modulators can be useful for monitoring the development or progression of SARS-CoV-2 infection or as

part of a clinical testing procedure to determine the efficacy of a particular therapy including anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein (i.e. theragnostics), or to determine a future course of treatment. Such markers or reporters may also be useful in purifying anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein.

[1570] Diagnosis and detection can be accomplished by coupling the modulator to detectable substances including, but not limited to, various enzymes comprising for example horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as but not limited to streptavidin/biotin and avidin/biotin; fluorescent materials, such as but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent materials, such as but not limited to, luminol; bioluminescent materials, such as but not limited to, luciferase, luciferin, and aequorin; radioactive materials, such as but not limited to iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{115}In , ^{113}In , ^{112}In , ^{111}In), and technetium (^{99}Tc), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F), ^{153}Sm , ^{177}Lu , ^{59}Gd , ^{49}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru , ^{68}Ge , ^{57}Co , ^{65}Zn , ^{85}Sr , ^{32}P , ^{153}Gd , ^{169}Yb , ^{51}Cr , ^{54}Mn , ^{75}Se , ^{113}Sn , and ^{117}Te ; positron emitting metals using various positron emission tomographies, noradioactive paramagnetic metal ions, and molecules that are radiolabeled or conjugated to specific radioisotopes. In such embodiments appropriate detection methodology is well known in the art and readily available from numerous commercial sources.

[1571] In other embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein can be fused to marker sequences, such as a peptide or fluorophore to facilitate purification or diagnostic procedures such as immunohistochemistry or FACs. In some embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector, among others, many of which are commercially available. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin “HA” tag, which corresponds to an epitope derived from the influenza hemagglutinin protein and the “flag” tag.

[1572] In yet other embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein can be conjugated to an immunomodulator, cytokine, cytotoxic agent, chemotherapeutic agent, antiviral agent, antimicrobial agent, or other drug. In some embodiments, the antibodies are conjugated to an antiviral agent.

[1573] Diagnostic Methods Using Anti-Coronavirus Antibodies

[1574] The inventions disclosed herein also encompass in vitro or in vivo methods for detecting, diagnosing or monitoring coronavirus infections and methods of screening cells from a patient to identify coronavirus infected cells, including cells from a patient who is currently infected with SARS-CoV-2, or cells from a patient who is recovered from a past SARS-CoV-2 infection. Such methods include identifying an individual infected with coronavirus for treatment, monitoring progression of a coronavirus infection comprising contacting the patient or a sample obtained from a patient with one or more anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein, and detect-

ing the presence or absence, or level of association of the antibody to a coronavirus antigen in the sample. In a particularly preferred embodiment, one or more anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be used to detect and quantify coronavirus levels in a patient sample (e.g., plasma or blood). Association with a coronavirus antigen in the sample likely denotes that the individual may be effectively treated with one or more anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein. The methods may further comprise a step of comparing the level of binding to a control. Other diagnostic or theragnostic methods compatible with the teachings herein are well known in the art and can be practiced using commercial materials such as dedicated reporting systems.

[1575] Exemplary compatible assay methods include radioimmunoassays, enzyme immunoassays, competitive-binding assays, fluorescent immunoassay, immunoblot assays, Western Blot analysis, flow cytometry assays, and ELISA assays. More generally detection of coronavirus in a biological sample may be accomplished using any art-known assay. Compatible in vivo theragnostics or diagnostics may comprise art recognized imaging or monitoring techniques such as magnetic resonance imaging (MRI), computerized tomography (e.g. CAT scan), positron tomography (e.g., PET scan) radiography, ultrasound, etc. Those skilled in the art will readily be able to recognize and implement appropriate detection, monitoring or imaging techniques (often comprising commercially available sources) based on the etiology, pathological manifestation, or clinical progression of the disorder.

[1576] In another embodiment, the invention provides a method of analyzing coronavirus infection progression and/or pathogenesis in vivo.

[1577] In another aspect, and as discussed in more detail below, the inventions disclosed herein also encompass kits for detecting, monitoring, or diagnosing a coronavirus infection, identifying an individual having a coronavirus infection for possible treatment or monitoring progression (or regression) of the infection in a patient, wherein the kit comprises an anti-coronavirus antibody as described herein, and reagents for detecting the effect of the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) on a sample from the patient.

[1578] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein and cells, cultures, populations and compositions comprising the same, including progeny thereof, can also be used to screen for or identify compounds or agents (e.g., drugs) that affect a function or activity of coronavirus virions or coronavirus infected cells or progeny thereof by binding to an antigen present on the surface of the virion or infected cell. The inventions disclosed herein therefore encompass systems and methods for evaluation or identification of a compound or agent that can affect a function or activity of the coronavirus virus. Such compounds and agents can be drug candidates that are screened for the treatment of coronavirus infection, for example. In one embodiment, a system or method comprises coronavirus virions and/or coronavirus infected cells and a compound or agent (e.g., drug), wherein the virions/cells and compound or agent (e.g., drug) are in contact with each other.

[1579] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may also be used as a

reagent to test a vaccine, such as an inactivated virus, live-attenuated vaccine, or recombinant subunit vaccine. In some embodiments, such anti-SARS-CoV-2 antibodies or antigen binding fragments comprise: (i) three CDRs in the heavy chain variable region (VH) set forth as SEQ ID NO: 661 and three CDRs in the light chain variable region (VL) set forth as SEQ ID NO: 662; (ii) three CDRs in the VH set forth as SEQ ID NO: 727 and three CDRs in the VL set forth as SEQ ID NO: 728; (iii) three CDRs in the VH set forth as SEQ ID NO: 761 and three CDRs in the VL set forth as SEQ ID NO: 762; (iv) three CDRs in the VH set forth as SEQ ID NO: 867 and three CDRs in the VL set forth as SEQ ID NO: 868; (v) three CDRs in the VH set forth as SEQ ID NO: 1123 and three CDRs in the VL set forth as SEQ ID NO: 1124; (vi) three CDRs in the VH set forth as SEQ ID NO: 1167 and three CDRs in the VL set forth as SEQ ID NO: 1168; (vii) three CDRs in the VH set forth as SEQ ID NO: 1267 and three CDRs in the VL set forth as SEQ ID NO: 1268; or (viii) three CDRs in the VH set forth as SEQ ID NO: 1313 and three CDRs in the VL set forth as SEQ ID NO: 1314; wherein the CDRs are defined by Kabat, Chothia or MacCallum numbering. In other embodiments, such anti-SARS-CoV-2 antibodies or antigen binding fragments comprise (i) a heavy chain variable region (VH) set forth as SEQ ID NO: 661 and a light chain variable region (VL) set forth as SEQ ID NO: 662; (ii) a VH set forth as SEQ ID NO: 727 and a VL set forth as SEQ ID NO: 728; (iii) a VH set forth as SEQ ID NO: 761 and a VL set forth as SEQ ID NO: 762; (iv) a VH set forth as SEQ ID NO: 867 and a VL set forth as SEQ ID NO: 868; (v) a VH set forth as SEQ ID NO: 1123 and a VL set forth as SEQ ID NO: 1124; (vi) a VH set forth as SEQ ID NO: 1167 and a VL set forth as SEQ ID NO: 1168; (vii) a VH set forth as SEQ ID NO: 1267 and a VL set forth as SEQ ID NO: 1268; or (viii) a VH set forth as SEQ ID NO: 1313 and a VL set forth as SEQ ID NO: 1314.

[1580] The amino acid sequences of the antibodies described in the foregoing paragraphs are provided below:

258-VH (SEQ ID NO: 661)
 QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWIRQPGKGLEWIGR
 IYTSGSTNYPNLSKSRVTMSVDTSKNQFSLKLSVTAADTAVYYCAAGYG
 SIDYWGQGLTVTVSS

258-VL (SEQ ID NO: 662)
 DIVMTQSPPLSLPVTGPGEPAISCRSSQSLHNSGYNLYDWYKQPGQSPQ
 LLIIYLGSNRAGVPRDFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTP
 RTFGQGTKLEIK

291-VH (SEQ ID NO: 727)
 QVQLVESGGGVVQPGRLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAV
 ISYDGSNKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARAS
 GGSYFGGMDVWGQGTITVTVSS

-continued

291-VL (SEQ ID NO: 728)
 QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMI
 YEVSNRPSGVSNRPSGSKSGNTASLTISGLQAEDEADYYCSSYTSSTLY
 VFGTGTKVTVL

308-VH (SEQ ID NO: 761)
 QLQLQESGPGLVKPSSETLSLTCTVSGGSISSSYHWGWIROPKGLLEWI
 GSIYYSGSTYYNPSLKSRTISVDTSKNQFSLKLSVTAADTAVYYCAGL
 RVVITFGGVIPKGGAFDIWGQGTMTVTVSS

308-VL (SEQ ID NO: 762)
 QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMI
 YDVSNRPSGVSNRPSGSKSGNTASLTISGLQAEDEADYYCSSYTSSTTV
 FGGGTKLTVL

361-VH (SEQ ID NO: 867)
 QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYVSWIRQHPGKGLLEWI
 GYIYYSGSTYYNPSLKSRTISVDTSKNQFSLKLSVTAADTAVYYCATT
 MVRGVIRLDHYGMDVWGQGTITVTVSS

361-VL (SEQ ID NO: 868)
 QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMI
 YEVSNRPSGVSNRPSGSKSGNTASLTISGLQAEDEADYYCSSYTSSTLL
 FGTGKTKVTVL

DIQMTQSPSSLSASVGRVTITCQASQGISNYLWYQQKPGKAPKLLIYD
 ASNLETGVPSPRPSGSGSGTDFTFTISSLPEDIAITYCQQYDNLPIITFGQ
 GTRLEIK

489-VH (SEQ ID NO: 1123)
 QVQLVESGGGVVQPGRLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAV
 ISYDGSNKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARAG
 SGNYYNWPDWVGQGLTVTVSS

489-VL (SEQ ID NO: 1124)
 EIVMTQSPATLSVSPGERATLSCRASQTVSSNLYWYQQKPGQAPRLLIYG
 ASTRATGIPARFSGSGSGTEFTLTISSLQSEDFAVYYCQQYNNWPPYTFG
 QGTKLEIK

511-VH (SEQ ID NO: 1167)
 QLQLQESGPGLVKPSSETLSLTCTVSGGSISSSYHWGWIROPKGLLEWI
 GSIYYSGSTYYNPSLKSRTISVDTSKNQFSLKLSVTAADTAVYYCASE
 KVDFWSGGPYGMDVWGQGTITVTVSS

-continued

511-VL
(SEQ ID NO: 1168)
QSALTQPASVSGSPGQSITISCTGTSSDVGSYNYVSWYQQHPGKAPKLM
YEVSNRPSGVSNRFSGSKSGNTASLTISGLQAEADYYCSTYSISTLV
FGGGTKLTVL

561-VH
(SEQ ID NO: 1267)
QVQLVESGGGVVQPGSRSLRLSCAASGFTFSSYAMHWVRQAPGKLEWVAV
ISYDGSNKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVVYCARPL
SGSYRSAFDIWGQGTMTVTVSS

561-VL
(SEQ ID NO: 1268)
EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIYG
ASTRATGIPARFSGSGSGTEFTLTISLQSEDAVYYCQQYNNWPRTFG
QGTKVEIK

585-VH
(SEQ ID NO: 1313)
QVQLVESGGGVVQPGSRSLRLSCAASGFTFATYAMHWVRQAPGKLEWVAV
ISHDGSNKHYADSVKGRFTIRDNSKNTLYLQMNSLRAEGTAIYYCARESL
EAAAPPFDYWGQGTLLTVTVSS

585-VL
(SEQ ID NO: 1314)
SYELTQPPSVSVSPGQTATIIICSGDKLGEKYASWYQQKPGQSPALVIYQD
RKRPSGIPERFSGSNSGNTATLTISGTQAMDEADYYCQAWDSSNSVVFVG
GTKLTVP

[1581] Exemplary activity or function that can be modulated include changes in cell morphology, expression of a marker, differentiation or de-differentiation, maturation, proliferation, viability, apoptosis or cell death neuronal progenitor cells or progeny thereof.

[1582] Methods of screening and identifying agents and compounds include those suitable for high-throughput screening, which include arrays of cells (e.g., microarrays) positioned or placed, optionally at pre-determined locations or addresses. High-throughput robotic or manual handling methods can probe chemical interactions and determine levels of expression of many genes in a short period of time. Techniques have been developed that utilize molecular signals (e.g., fluorophores) and automated analyses that process information at a very rapid rate.

[1583] Such screening methods (e.g., high-throughput) can identify active agents and compounds rapidly and efficiently. For example, cells can be positioned or placed (pre-seeded) on a culture dish, tube, flask, roller bottle or plate (e.g., a single multi-well plate or dish such as an 8, 16, 32, 64, 96, 384 and 1536 multi-well plate or dish), optionally at defined locations, for identification of potentially therapeutic molecules. Libraries that can be screened include, for example, small molecule libraries, phage display libraries, fully human antibody yeast display libraries, siRNA libraries, and adenoviral transfection vectors.

[1584] Pharmaceutical Compositions and Therapeutic Uses

[1585] Provided herein are methods of treating or preventing a SARS-Co-V or SARS-Co-V-2-linked disease (e.g.,

COVID-19) by administering to a patient a therapeutically effective amount of one or more anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein, or a pharmaceutical composition comprising one or more (e.g., two or three) anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) described herein.

[1586] As used interchangeably herein, “treatment” or “treating” or “treat” refers to all processes wherein there may be a slowing, interrupting, arresting, controlling, stopping, alleviating or ameliorating symptoms or complications, or reversing of the progression of the disorders or disease disclosed herein, e.g., SARS-CoV-2 viral infection or COVID-19 disease, but does not necessarily indicate a total elimination of all disease or disorder symptoms.

[1587] As used herein, “prevention”, “prevent”, and/or “preventing”, which are used interchangeably herein, refers to the prophylactic treatment of a disease or disorder, or delaying the onset or progression of the disease or disorder, e.g., SARS-CoV-2 viral infection or COVID-19 disease.

[1588] In some embodiments, methods of treating or preventing a SARS-Co-V or SARS-Co-V-2-linked disease (e.g., COVID-19) comprise administering to a patient a pharmaceutical composition comprising one or more (e.g., two or three) anti-SARS-CoV-2 antibodies described herein. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies or antigen-binding fragments thereof that bind different epitopes of the SARS-CoV-2 S protein. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one of the antibodies is selected from antibodies 258 to 577 and 589 to 1587. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one of the antibodies is 555. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one of the antibodies neutralize SARS-CoV-2. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one antibody or antigen-binding fragment thereof blocks SARS-CoV-2 binding to ACE2. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one antibody or antigen-binding fragment thereof binds the RBD of the SARS-CoV-2 S protein. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one antibody or antigen-binding fragment thereof binds the NTD domain of the SARS-CoV-2 S protein. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one antibody or antigen-binding fragment

thereof binds the S2 domain of the SARS-CoV-2 S protein. In some embodiments, such methods comprise administering to a patient a pharmaceutical composition comprising two or three anti-SARS-CoV-2 antibodies, wherein at least one antibody or antigen-binding fragment thereof binds the RBD of the SARS-CoV-2 S protein and at least one antibody or antigen-binding fragment thereof binds the NTD or S2 domain of the SARS-CoV-2 S protein.

[1589] In some embodiments, provided herein are methods of treating or preventing a SARS-CoV-2-linked disease (e.g., COVID-19) comprising administering to a patient an antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364, or a pharmaceutical composition comprising such an antibody. In some embodiments, such methods further comprise administering to the patient another antibody that binds the SARS-CoV-2 S protein, e.g., CB-6. In some embodiments, provided herein are methods of treating or preventing a SARS-CoV-2-linked disease (e.g., COVID-19) comprise administering to a patient a pharmaceutical composition comprising an antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364, and at least one additional antibody that binds SARS-CoV-2 (e.g., CB6).

[1590] In some embodiments, provided herein are methods of preventing COVID-19 comprising administering to a patient who is at risk for contracting COVID-19, an antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364, or a pharmaceutical composition comprising such an antibody. In some embodiments, such methods further comprise administering to the patient another antibody that binds the SARS-CoV-2 S protein, e.g., CB-6.

[1591] In some embodiments, provided herein are methods of reducing COVID-19 related hospitalization or Emergency Room (ER) visit of a patient having COVID-19 by administering to the patient a therapeutically effective amount of an anti-SARS-CoV-2 antibody described herein or a pharmaceutical composition comprising such an antibody. In some embodiments, provided herein are methods of reducing COVID-19 related hospitalization or ER visit of a patient having COVID-19 by administering to the patient a therapeutically effective amount of an antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364, or a pharmaceutical composition comprising such an antibody. In some embodiments, provided herein are methods of reducing COVID-19 related hospitalization or ER visit of a patient having COVID-19 by administering to the patient a therapeutically effective amount of an antibody comprises a heavy chain comprising SEQ ID NO: 5363 and a light chain comprising SEQ ID NO: 5364, and another anti-SARS-CoV-2 antibody (e.g., an antibody that binds the SARS-CoV-2 S protein, e.g., CB6).

[1592] In some embodiments, provided herein are methods of treating or preventing COVID-19 comprising: contacting a sample obtained from a patient with an antibody or antigen-binding fragment thereof described herein, conjugated to a detectable agent; detecting specific binding of the antibody or antigen-binding fragment thereof to a SARS-CoV-2 antigen present in the sample; and administering to the patient a therapeutically effective amount of an antibody or antigen-binding fragment thereof described herein or a

pharmaceutical composition comprising such an antibody or antigen-binding fragment thereof.

[1593] In some embodiments, the patient has moderate to severe COVID-19, but is not hospitalized. In some embodiments, the patient has mild to moderate COVID-19. For example, mild COVID-19 patients can include individuals who have any of various signs and symptoms, e.g., fever, cough, sore throat, malaise, headache, muscle pain, without shortness of breath, dyspnea, or abnormal imaging. Moderate COVID-19 patients can include individuals who have evidence of lower respiratory disease by clinical assessment or imaging and a saturation of oxygen (SaO₂) greater than (>)93 percent (%) on room air at sea level. In some embodiments, the patient is at risk for contracting COVID-19. In some embodiments, the patient has a positive SARS-CoV-2 viral testing result. In some embodiments, the patient is an adult or pediatric patient who is 12 years of age and older and weigh at least 40 kilograms (kg). In some embodiments, the patient is at high risk for progressing to severe COVID-19 and/or hospitalization, e.g., the patient (i) is 65 years of age or older (≥ 65); (ii) has a body mass index (BMI) of 35 or greater (≥ 35); (iii) has chronic kidney disease; (iv) has diabetes; (v) has immunosuppressive disease, (vi) is receiving immunosuppressive treatment; (vii) is 55 years of age or older (≥ 55) and has cardiovascular disease, hypertension, chronic obstructive pulmonary disease, or other chronic respiratory disease; or (viii) is 12-17 years of age and have a BMI $\geq 85\%$ for their age and gender, or sickle cell disease, congenital or acquired heart disease, neurodevelopmental disorders (e.g., cerebral palsy), a medical-related technological dependence (e.g., tracheostomy, gastrostomy, or positive pressure ventilation not related to COVID-19), or asthma, reactive airway or other chronic respiratory disease that requires daily medication for control. In some embodiments, the patient has mild to moderate COVID-19 and the patient is at high risk for progressing to severe COVID-19 and/or hospitalization, e.g., the patient (i) is 65 years of age or older (≥ 65); (ii) has a body mass index (BMI) of 35 or greater (≥ 35); (iii) has chronic kidney disease; (iv) has diabetes; (v) has immunosuppressive disease, (vi) is receiving immunosuppressive treatment; (vii) is 55 years of age or older (≥ 55) and has cardiovascular disease, hypertension, chronic obstructive pulmonary disease, or other chronic respiratory disease; or (viii) is 12-17 years of age and have a BMI $\geq 85\%$ for their age and gender, or sickle cell disease, congenital or acquired heart disease, neurodevelopmental disorders (e.g., cerebral palsy), a medical-related technological dependence (e.g., tracheostomy, gastrostomy, or positive pressure ventilation not related to COVID-19), or asthma, reactive airway or other chronic respiratory disease that requires daily medication for control.

[1594] Also provided are anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) or antigen-binding fragments or pharmaceutical compositions comprising one or more (e.g., two or three) anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) or antigen-binding fragments, for use in therapy. In some embodiments anti-SARS-CoV-2 antibodies or antigen-binding fragments or pharmaceutical compositions comprise one or more (e.g., two or three) anti-SARS-CoV-2 antibodies or antigen-binding fragments, for use in the treatment or prevention of COVID-19. Further provided herein are uses of anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) or antigen-binding fragments

described herein in the manufacture of a medicament for the treatment or prevention of COVID-19.

[1595] Depending on the form of anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody), mode of intended delivery, and numerous other variables, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be formulated as desired using art recognized techniques. Various pharmaceutically acceptable carriers, which include vehicles, adjuvants, and diluents, are readily available from numerous commercial sources. Moreover, an assortment of pharmaceutically acceptable auxiliary substances, such as pH adjusting and buffering agents, tonicity adjusting agents, stabilizers, wetting agents, and the like, are also available. Certain non-limiting exemplary carriers include saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof.

[1596] In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be administered to a patient neat or with a minimum of additional components. In other embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be formulated to contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries that are well known in the art and are relatively inert substances that facilitate administration or which aid processing of the active compounds into preparations that are pharmaceutically optimized for delivery. For example, an excipient can give form or consistency or act as a diluent to improve the pharmacokinetics of the antibody. Suitable excipients include but are not limited to stabilizing agents, wetting, and emulsifying agents, salts for varying osmolality, encapsulating agents, buffers, and skin penetration enhancers.

[1597] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be formulated for enteral, parenteral, or topical administration. Indeed, all three types of formulation may be used simultaneously to achieve systemic administration of the active ingredient. Excipients as well as formulations for parenteral and non-parenteral drug delivery are known in the art. Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts. In addition, suspensions of the active compounds as appropriate for oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions may contain substances that increase the viscosity of the suspension and include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers. Liposomes can also be used to encapsulate the agent for delivery into the cell.

[1598] Suitable formulations for enteral administration include hard or soft gelatin capsules, pills, tablets, including coated tablets, elixirs, suspensions, syrups or inhalations and controlled release forms thereof.

[1599] In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be adsorbed onto red blood cells to facilitate preferential delivery to the lungs preventing a shortened half-life through processing in the liver and spleen and providing a higher concentration in the lungs.

[1600] In general, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be adminis-

tered in vivo, to a subject in need thereof, by various routes, including, but not limited to, oral, intravenous, intra-arterial, subcutaneous, parenteral, intranasal, intramuscular, intracardiac, intraventricular, intratracheal, buccal, rectal, intraperitoneal, intradermal, topical, transdermal, and intrathecal, or otherwise by implantation or inhalation. Compositions may be formulated into preparations in solid, semi-solid, liquid, or gaseous forms; including, but not limited to, tablets, capsules, powders, granules, ointments, solutions, suppositories, enemas, injections, inhalants, and aerosols. The appropriate formulation and route of administration may be selected according to the intended application and therapeutic regimen.

[1601] In some embodiments, the pharmaceutical compositions described herein comprise one or more anti-SARS-CoV-2 antibodies. In some embodiments, the pharmaceutical compositions further comprise one or more of the following excipients: histidine, sodium chloride, sucrose, polysorbate 80. In some embodiments, the pharmaceutical compositions comprise at least one anti-SARS-CoV-2 antibody (e.g., 555), histidine, sodium chloride, sucrose, polysorbate 80. In some embodiments, the pharmaceutical compositions have a pH of about 6.0. In some embodiments, the pharmaceutical composition comprises at least one anti-SARS-CoV-2 antibody (e.g., 555), 5 mM histidine, 50 mM NaCl, 6% sucrose, and 0.05% polysorbate 80 and has a pH of about 6.0. In some embodiments, the anti-SARS-CoV-2 antibody concentration in the pharmaceutical composition is about 10 mg/mL to about 150 mg/mL. In some embodiments, the anti-SARS-CoV-2 antibody concentration in the pharmaceutical composition is about 35 mg/mL to about 125 mg/mL. In some embodiments, the anti-SARS-CoV-2 antibody concentration in the pharmaceutical composition is about 35 mg/mL. In some embodiments, the anti-SARS-CoV-2 antibody concentration in the pharmaceutical composition is about 125 mg/mL.

[1602] Similarly, the particular dosage regimen, i.e., dose, timing, and repetition, will depend on the particular individual and that individual's medical history. Empirical considerations such as pharmacokinetics (e.g., half-life, clearance rate, etc.) will contribute to the determination of the dosage. Frequency of administration may be determined and adjusted over the course of therapy, and is based on reducing the number of hyperproliferative or neoplastic cells, including tumor initiating cells, maintaining the reduction of such neoplastic cells, reducing the proliferation of neoplastic cells, or delaying the development of metastasis. Alternatively, sustained continuous release formulations of a subject therapeutic composition may be appropriate. Various formulations and devices for achieving sustained release are known in the art.

[1603] Pharmaceutical compositions are administered in therapeutically effective amount for treatment or prophylaxis of a coronavirus infection. As used herein, the term "therapeutically effective amount" means that amount of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or pharmaceutical composition comprising the same that will elicit the biological or medical response in a subject that is sought by a medical doctor or other clinician. In particular, with regard to viral infections and proliferation of virus, a "therapeutically effective amount" is intended to include an amount sufficient to achieve one or more of the following effects: (i) reduction or amelioration the severity of a viral infection, viral disease or a symptom associated

therewith; (ii) reduction in the duration of a viral infection, viral disease, or a symptom associated therewith; (iii) prevention of the progression of a viral infection, viral disease, or a symptom associated therewith; (iv) regression of a viral infection, viral disease, or a symptom associated therewith; (v) prevention of the development or onset of a viral infection, viral disease, or a symptom associated therewith; (vi) prevention of the recurrence of a viral infection, viral disease, or a symptom associated therewith; (vii) reduction or prevention of the spread of coronavirus from one cell to another cell, one tissue to another tissue, or one organ to another organ; (viii) prevention or reduction of the spread/transmission of coronavirus from one subject to another subject; (ix) reduction in organ failure associated with a viral infection or viral disease; (x) reduction in the hospitalization of a subject; (xi) reduction in the hospitalization length; (xii) an increase in the survival of a subject with coronavirus infection or a disease associated therewith; (xiii) elimination of a coronavirus infection or a disease associated therewith; (xiv) inhibition or reduction in viral replication; (xv) inhibition or reduction in the binding or fusion of virions to a host cell(s); (xvi) inhibition or reduction in the entry of virions into a host cell(s); (xvii) inhibition or reduction of replication of the viral genome; (xviii) inhibition or reduction in the synthesis of viral proteins; (xix) inhibition or reduction in the assembly of viral particles; (xx) inhibition or reduction in the release of viral particles from a host cell(s); (xxi) reduction in viral titer, (xxii) the reduction in the number of symptoms associated with coronavirus infection or viral disease; (xxiii) enhancement, improvement, supplementation, complementation, or augmentation of the prophylactic or therapeutic effect(s) of another therapy; (xxiv) prevention of the onset or progression of a secondary infection associated with a viral infection; (xxv) prevention of the onset or diminution of the severity of another disease occurring secondary to a coronavirus infection; and/or (xxvi) change in the immune response coronavirus infection including cytokines, chemokines, complement, cellular responses, etc.

[1604] In some embodiments, a therapeutically effective amount of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or pharmaceutical composition comprising the same has a beneficial effect but does not cure a viral infection or a disease associated therewith. In certain embodiments, therapy may encompass the administration of multiple doses of an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) or pharmaceutical composition comprising the same at a certain frequency to achieve an amount of the therapy that has a prophylactic and/or therapeutic effect.

[1605] Readily observable symptoms associated with coronavirus and other viral infections include fever, cough and/or sore throat, runny or stuffy nose, headache and/or body aches, chills, fatigue, generalized weakness, nausea, and vomiting and/or diarrhea.

[1606] A therapeutically effective amount is typically dependent on the weight of the subject being treated, his or her physical condition, the extensiveness of the condition to be treated, and the age of the subject being treated. In general, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be administered in an amount in the range of about 10 ng/kg body weight to about 100 mg/kg body weight per dose. In certain embodiments, antibodies may be administered in an amount in the range of

about 50 µg/kg body weight to about 5 mg/kg body weight per dose. In other embodiments, antibodies may be administered in an amount in the range of about 100 µg/kg body weight to about 10 mg/kg body weight per dose. In other embodiments, antibodies may be administered in an amount in the range of about 100 µg/kg body weight to about 20 mg/kg body weight per dose. In other embodiments, antibodies may be administered in an amount in the range of about 0.5 mg/kg body weight to about 20 mg/kg body weight per dose. In other embodiments, antibodies may be administered in a dose of at least about 100 µg/kg body weight, at least about 250 µg/kg body weight, at least about 750 µg/kg body weight, at least about 3 mg/kg body weight, at least about 5 mg/kg body weight, or at least about 10 mg/kg body weight.

[1607] In some embodiments, an anti-SARS-CoV-2 antibody or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 100 mg to about 10,000 mg. In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 700 mg to about 7000 mg (e.g., about 700 mg, about 750 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, about 1000 mg, about 1100 mg, about 1200 mg, about 1300 mg, about 1400 mg, about 1500 mg, about 1600 mg, about 1700 mg, about 1800 mg, about 1900 mg, about 2000 mg, about 2100 mg, about 2200 mg, about 2300 mg, about 2400 mg, about 2500 mg, about 2600 mg, about 2700 mg, about 2800 mg, about 2900 mg, about 3000 mg, about 3100 mg, about 3200 mg, about 3300 mg, about 3400 mg, about 3500 mg, about 3600 mg, about 3700 mg, about 3800 mg, about 3900 mg, about 4000 mg, about 4100 mg, about 4200 mg, about 4300 mg, about 4400 mg, about 4500 mg, about 4600 mg, about 4700 mg, about 4800 mg, about 4900 mg, about 5000 mg, about 5100 mg, about 5200 mg, about 5300 mg, about 5400 mg, about 5500 mg, about 5600 mg, about 5700 mg, about 5800 mg, about 5900 mg, about 6000 mg, about 6100 mg, about 6200 mg, about 6300 mg, about 6400 mg, about 6500 mg, about 6600 mg, about 6700 mg, about 6800 mg, about 6900 mg, about 7000 mg). In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 700 mg, 1400 mg, 2800 mg, 4200 mg, 5600 mg, or 7000 mg. In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 35 mg to about 700 mg (e.g., about 35 mg, about 70 mg, about 105 mg, about 140 mg, about 150 mg, about 175 mg, about 210 mg, about 245 mg, about 280 mg, about 315 mg, about 350 mg, 385 mg, 420 mg, 455 mg, 490 mg, 525 mg, 560 mg, 595 mg, 630 mg, 665 mg, or 700 mg). In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 70 mg, 140 mg, 150 mg, 175 mg, 210 mg, 280 mg, 350 mg, 420 mg, 490 mg, 560 mg, or 630 mg. In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 700 mg. In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or

a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 350 mg. In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 175 mg. In some embodiments, an anti-SARS-CoV-2 antibody (e.g., 555) or a pharmaceutical composition comprising such an antibody, is administered intravenously or subcutaneously to a patient at a dose of about 150 mg.

[1608] Other dosing regimens may be predicated on Body Surface Area (BSA) calculations. As is well known in the art, a patient's BSA is calculated using the patient's height and weight and provides a measure of a subject's size as represented by the surface area of his or her body. In some embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are administered in dosages from 10 mg/m² to 800 mg/m². In other embodiments, antibodies are administered in dosages from 50 mg/m² to 500 mg/m² and even more preferably at dosages of 100 mg/m², 150 mg/m², 200 mg/m², 250 mg/m², 300 mg/m², 350 mg/m², 400 mg/m² or 450 mg/m².

[1609] Escalation for an individual patient can occur at the discretion of a clinician in the absence of any clinically significant occurrence that the clinician might reasonably believe would present an undue safety risk for the patient, such as, for example, Grade ≥ 3 non-hematologic toxicity, Grade ≥ 3 nausea, vomiting or diarrhea uncontrolled by maximum antiemetic/anti-diarrhea therapy, Grade 4 neutropenia lasting >7 days in the absence of growth factor support, Grade 3 or 4 neutropenia of any duration accompanied with fever $\geq 38.5^\circ$ C. and/or systemic infection, or other Grade ≥ 4 hematologic toxicity.

[1610] Anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein are usually administered to the patient on multiple occasions. An exemplary treatment regimen entails administration once per every two weeks, once a month, or once every 3 to 6 months. For example, patients can receive the antibody (e.g., as an intravenous formulation) once every four weeks as a cycle, for example every twenty-eight days. The dosing frequency can be adjusted depending on the pharmacokinetic profile of the antibody in the patient. For example, the half-life of the antibody may warrant a two week frequency of dosing. In some methods, two or more antibodies with different binding specificities may be administered simultaneously, in which case the dosage of each antibody administered falls within the ranges indicated. Intervals between single dosages can be weekly, monthly, or yearly. Intervals can also be irregular depending upon levels of antibody in the blood and other clinical indicia. In some methods, the dosage is adjusted to achieve a plasma antibody concentration of about 1-1000 $\mu\text{g/mL}$ or about 25-300 $\mu\text{g/mL}$. Alternatively, antibodies can be administered as a sustained release formulation, in which case less frequent administration is required. Antibodies may be administered to the patient for at least 9 months, at least 12 months, or for a longer period of time to achieve a desired result.

[1611] Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, human antibodies show the longest half-life, followed by humanized antibodies, chimeric antibodies, and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or

therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, until a partial or complete response is achieved, and/or until the patient shows lessening or amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

[1612] The duration of a therapeutic regimen depends on the disease being treated, the age and condition of the patient, the stage and type of the patient's disease, how the patient responds to the treatment, etc. A clinician can observe the therapy's effects closely and make any adjustments as needed. When agents are used in combination, the two or more therapeutic agents are administered simultaneously or sequentially in any order, i.e., an antibody disclosed herein is administered prior to administering a second therapeutic agent, concurrently with a second therapeutic agent, or subsequent to administration of a second therapeutic agent. For example, a combination therapy may be performed by administering a first therapeutic agent prior to (e.g., 1 minute, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concurrently with, or subsequent to (e.g., 1 minute, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) administering a second therapeutic agent.

[1613] The dosage, frequency, and mode of administration of each component of a combination therapy can be controlled independently. For example, one therapeutic agent may be administered orally three times per day, while the second therapeutic agent may be administered intramuscularly once per day. Combination therapy may be given in on-and-off cycles that include rest periods. The compounds may also be admixed or otherwise formulated together such that one administration delivers both therapeutic agents. In this case, each therapeutic agent is generally present in an amount of 1-95% by weight of the total weight of the composition. Alternatively, therapeutic agents can be formulated separately and in individual dosage amounts. Combinations of therapeutic agents for treatment can be provided as components of a pharmaceutical pack.

[1614] Preferably, combination therapies elicit a synergistic therapeutic effect, i.e., an effect greater than the sum of their individual effects or therapeutic outcomes, such as those described above. For example, a synergistic therapeutic effect may be an effect of at least about two-fold greater than sum of the therapeutic effects elicited by the single agents of a given combination, or at least about five-fold greater, or at least about ten-fold greater, or at least about twenty-fold greater, or at least about fifty-fold greater, or at least about one hundred-fold greater. A synergistic therapeutic effect may also be observed as an increase in therapeutic effect of at least 10% compared to the sum of the therapeutic effects elicited by the single agents of a given combination, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or more. A synergistic effect is

also an effect that permits reduced dosing of therapeutic agents when they are used in combination.

[1615] Articles of Manufacture

[1616] The inventions disclosed herein also encompass pharmaceutical packs and kits comprising one or more containers and comprising one or more doses of an anti-coronavirus (e.g., anti-SARS-CoV-2 antibody) disclosed herein. In certain embodiments, a unit dosage is provided wherein the unit dosage contains a predetermined amount of a composition comprising, for example, an anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) disclosed herein, with or without one or more additional agents. For other embodiments, such a unit dosage is supplied in single-use prefilled syringe for injection. In still other embodiments, the composition contained in the unit dosage may comprise saline, sucrose, or the like; a buffer, such as phosphate, or the like, and/or be formulated within a stable and effective pH range. Alternatively, in certain embodiments, the composition may be provided as a lyophilized powder that may be reconstituted upon addition of an appropriate liquid, for example, sterile water. In certain preferred embodiments, the composition comprises one or more substances that inhibit protein aggregation, including, but not limited to, sucrose and arginine. Any label on, or associated with, the container(s) indicates that the enclosed composition is used for diagnosis or treatment.

[1617] The present invention also provides kits for producing single-dose or multi-dose administration units of an anti-coronavirus antibody (e.g., an anti-SARS-CoV-2 antibody) disclosed herein and, optionally, one or more other diagnostic or therapeutic agents. The kit comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits will generally contain in a suitable container a pharmaceutically acceptable formulation of the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibodies) and, optionally, one or more other diagnostic or therapeutic agents in the same or different containers. The kits may also contain other pharmaceutically acceptable formulations, either for diagnosis or combined therapy. Such kits may also provide appropriate reagents to conjugate the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibodies) with the other diagnostic or therapeutic agent(s).

[1618] More specifically the kits may have a single container that contains the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody), with or without additional components, or they may have distinct containers for each desired agent. Where combined therapeutics are provided for conjugation, a single solution may be pre-mixed, either in a molar equivalent combination, or with one component in excess of the other. Alternatively, the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) and any optional diagnostic or therapeutic agent of the kit may be maintained separately within distinct containers prior to administration to a patient. The kits may also comprise a second/third container means for containing a sterile, pharmaceutically acceptable buffer or other diluent such as bacteriostatic

water for injection (BWFI), phosphate-buffered saline (PBS), Ringer's solution and dextrose solution.

[1619] When the components of the kit are provided in one or more liquid solutions, the liquid solution is preferably an aqueous solution, with a sterile aqueous solution being particularly preferred. However, the components of the kit may be provided as dried powder(s). When reagents or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container.

[1620] As indicated briefly above the kits may also contain a means by which to administer the antibody and any optional components to the patient, e.g., one or more needles or syringes, or even an eye dropper, pipette, or other such like apparatus, from which the formulation may be injected or introduced into the patient. Such kits will also typically include a means for containing the vials, or such like, and other component in close confinement for commercial sale, such as, e.g., injection or blow-molded plastic containers into which the desired vials and other apparatus are placed and retained. Any label or package insert indicates that the anti-coronavirus antibody (e.g., anti-SARS-CoV-2 antibody) composition is used for treating cancer, for example colorectal cancer.

[1621] In other preferred embodiments, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) disclosed herein may be used in conjunction with, or comprise, diagnostic or therapeutic devices useful in the diagnosis or treatment of proliferative disorders. For example, in one embodiment, anti-coronavirus antibodies (e.g., anti-SARS-CoV-2 antibodies) may be combined with certain diagnostic devices or instruments that may be used to detect, monitor, quantify or profile cells or marker compounds involved in the etiology or manifestation a coronavirus infection.

EXAMPLES

[1622] The following examples have been included to illustrate aspects of the inventions disclosed herein. In light of the present disclosure and the general level of skill in the art, those of skill appreciate that the following examples are intended to be exemplary only and that numerous changes, modifications, and alterations may be employed without departing from the scope of the disclosure.

Example 1

[1623] Isolation and Generation of Monoclonal Antibodies

[1624] Blood samples were obtained from convalescent human donors with either a confirmed SARS-CoV or SARS-CoV-2 infection. The samples were enriched for antibody-secreting B-cells and screened using single cell secretion assays, to enrich for antibodies that bind to recombinant SARS-CoV spike protein and/or SARS-CoV-2 spike protein using two different miniaturized single cell secretion assays: a multiplexed bead-based assay and a live-cell assay. The live cell assay is an ideal strategy to rapidly express and screen unknown antigens during a pandemic, without the need for recombinant protein expression and in-depth knowledge of the target.

[1625] For the bead assay, unique antibody sequences were confirmed to bind the screening target (SARS-CoV-2 full length spike) using a multiplexed bead assay on high

throughput flow cytometry. Different optically encoded bead types were conjugated to one of the following unique antigens: full length spike of SARS-CoV-2, MERS, SARS-CoV, HKU1, WIV1 or the S1 subunit of SARS-CoV-2 spike. Purified antibodies were incubated with the multiplexed beads, and negative control beads conjugated to BSA-His or FoldOn-Hisat either 50 nM, 10 nM or 2 nM antibody concentration for 30 minutes at room temperature. Beads were washed and binding was detected by using a fluorescently labeled anti-human secondary antibody. Fluorescence was measured using high throughput plate-based flow cytometry. Benchmark antibodies identified to SARS-CoV were used as positive controls due to similarity in spike sequences between SARS-CoV and SARS-CoV-2; human IgG isotype and an irrelevant antibody were used as negative controls. Multiplexed beads conjugated to full-length trimeric SARS-CoV spike protein and full-length trimeric SARS-CoV-2 spike protein enriched for antibodies that were either monospecific to SARS-CoV, SARS-CoV-2, or cross-reactive to both.

[1626] For the live cell assay, unique antibody sequences were confirmed to bind the screening target (SARS-CoV-2 full length spike) using high throughput flow cytometry. CHO cells were transiently transfected to express the full length spike protein of either wild type or mutant (V483A, D614G, V367F) SARS-CoV-2 on the cell surface. Full length spike protein sequences of either wildtype or reported viral mutants (V483A, D614G, V467F). SARS-CoV-2 and GFP inserts were cloned into the vector backbone pCDNA3.1. Suspension CHO cells were transiently transfected with the plasmid using electroporation. Full length native conformation spike protein expression was confirmed by testing with benchmark antibodies discovered against SARS-CoV that target different stalk and head domains using flow cytometry. Western blot was performed with a whole cell and plasma membrane isolate to confirm full length protein expression on the cell surface.

[1627] Purified antibodies were incubated with the readout cells, and an untransfected control CHO line either at 50 nM, 10 nM or 2 nM antibody concentration for 30 minutes at 4° C. CHO cells were washed and binding was detected by using a fluorescently labeled anti-human secondary antibody. Fluorescence was measured using high throughput plate-based flow cytometry. Benchmark antibodies identified to SARS-CoV were used as positive controls due to similarity in spike sequences between SARS-CoV and SARS-CoV-2; human IgG isotype and an irrelevant antibody were used as negative controls. Median fluorescence intensity of each antibody was normalized over the median fluorescence intensity of the human isotype control for respective antigens. The median fold over isotype values

from different validation experiments were plotted. Mean and standard deviation was calculated where applicable and the values plotted as a column bar graph with median fold over isotype on Y-axis and the different antibodies represented along the X-axis. Antibody values greater than 5-fold over isotype were considered as binders. The cut-off value was determined based on the binding to the negative controls.

[1628] Individual B-cells from chambers identified as having positive binding events (hits) were recovered from the microfluidic device, from were generated next-generation sequencing (NGS) libraries of the antibody genes from the recovered single cells and sequenced them using the MiSeq platform (Illumina, USA).

[1629] After determining the V_H and V_L sequences of the antigen-reactive B cells, they were converted to full-length IgG1 antibody sequences. The full-length heavy chain sequences of IgG1m3 allotype were constructed based on the V_H sequences, and the full-length light chain kappa or lambda sequences were constructed based on the V_L sequences. Full-length heavy chain and light chain sequences were cloned into expression vectors and recombinantly expressed.

[1630] Table 1-1 shows the results of a bead assay described above demonstrating which exemplary anti-coronavirus antibodies bind to wild-type SARS-CoV-2. Columns A-F: 50 nM SARS-CoV-2. Columns G and H: 10 nM SARS-CoV-2. Column I: 2 nM SARS-CoV-2. Blank cells: data not available.

[1631] Table 1-2 shows the results of a live cell assay described above demonstrating which exemplary anti-coronavirus antibodies bind to wild-type SARS-CoV-2. Columns A-E: 50 nM SARS-CoV-2. Column F and G: 10 nM SARS-CoV-2. Column H: 2 nM SARS-CoV-2. Blank cells: data not available.

[1632] Table 1-3 shows the results of a live cell assay described above demonstrating which exemplary anti-coronavirus antibodies bind to wild-type SARS-CoV-2. Columns A-E: 50 nM SARS-CoV-2 V483A mutant. Column F: 50 nM SARS-CoV-2 V367F mutant. Column G: 50 nM SARS-CoV-2 D641G mutant. Blank cells: data not available.

[1633] Table 1-4 shows the results of a bead assay described above demonstrating which exemplary anti-coronavirus antibodies cross-react with SARS. Columns A-E: 50 nM SARS. Columns F and G: 10 nM SARS. Column H: 2 nM SARS. Blank cells: data not available.

[1634] Table 1-5 shows the results of a bead assay described above demonstrating which exemplary anti-coronavirus antibodies cross-react with WIV1. Columns A-E: 50 nM WIV1. Columns F and G: 10 nM WIV1. Column H: 2 nM WIV1. Blank cells: data not available.

TABLE 1-1

Binding to Wild-Type SARS-CoV-2 (Bead Assay)									
Antibody ID	A	B	C	D	E	F	G	H	I
258		654.17		282.10				336.60	
259				274.62					
260				724.13					
261	542.68	997.13		619.94			160.26	444.66	47.44
262		372.32		64.07				20.60	
263				722.50					
264				597.90					
265				1283.39					

TABLE 1-1-continued

Binding to Wild-Type SARS-CoV-2 (Bead Assay)									
Antibody ID	A	B	C	D	E	F	G	H	I
266				472.98					
267				312.20					
268	297.76	0.99		513.25			111.48	0.98	30.29
269		480.28		191.05				188.11	
270				304.73					
271				448.53					
272		1352.02		678.91				852.12	
273				923.69					
274		943.76		127.21	1.18			392.61	
275	345.95	665.27		437.69			180.61	348.03	74.79
276				747.77					
277	548.05	867.71		506.71			320.53	482.79	121.04
278	119.65	556.13		321.43			14.80	184.75	3.29
279		3.09		1.36				1.11	
280	219.33	615.18		293.51			111.79	374.85	42.46
281				570.04					
283		232.15		250.46				57.46	
284				714.03					
285	265.30	542.28		405.29			74.32	222.19	14.36
286	227.89	228.10		312.20			64.27	48.44	9.93
287	711.63	1017.56		573.05			473.50	548.16	205.64
288				495.15					
290				1.16					
291		757.28		381.70				413.53	
292	1010.17	1494.67	969.44	921.34	972.31		699.57	1093.49	419.66
294				1232.58					
295				598.78					
296		1.09		1.15				1.08	
297		925.83		513.34				596.03	
298		1385.48		748.89				842.30	
299				1.43					
300				721.20					
301				865.03					
302	532.02	920.84		571.43			133.45	387.79	17.39
303				340.89					
304		1.22		1.17				1.09	
305		470.00		448.76				130.16	
306				338.09					
307		1188.27		753.87				572.61	
308		757.76		346.18				398.14	
309	557.73	856.65	558.04	473.88	578.48		344.16	518.05	146.21
310		250.07		151.80				83.28	
311		1014.32		527.12				506.43	
312		717.25		370.51				244.35	
313		656.13		344.72				328.21	
314		1086.51		657.54				424.26	
315				592.29					
316				441.70					
317				890.13					
318		839.38		409.72				528.53	
319				434.71					
320				1203.87					
321		253.61		59.23				55.51	
322		110.56		312.59				8.84	
323				994.30					
324				1.13					
325		455.25		135.46				140.57	
326				192.92					
327				413.06					
328				1.20					
329				496.63					
330		366.17		312.31				70.57	
331				850.47					
332				1.22					
333				1.08					
334		294.16		136.76				98.63	
335	1.06	1.08		77.65	38.96	105.22	1.19	0.93	0.99
336		679.19		105.22					
337				406.31				117.98	
338		562.82		756.03					
339				155.20				206.77	
340				873.86					
				546.84					

TABLE 1-1-continued

Binding to Wild-Type SARS-CoV-2 (Bead Assay)									
Antibody ID	A	B	C	D	E	F	G	H	I
341	1.29	1.22		1.17			1.19	1.00	1.09
342				745.63					
343				1.20					
344				403.95					
345				561.95					
346	274.63	358.12		193.49			90.67	98.99	19.23
347				598.50					
348				869.29					
349				118.34					
350		1288.69		812.79				938.17	
351				1.06					
352		1211.33		396.07				364.14	
353	596.85	945.10		578.48			441.27	605.76	234.76
354	2.03	95.02		8.56			1.17	16.26	1.06
355		708.91		514.32				309.09	
356	754.87	1083.64		637.2.9			499.03	608.86	198.83
357		262.21		248.50				82.13	
358	282.03	415.72		345.87			71.19	147.18	14.26
359	389.89	681.77		326.29			172.29	336.71	74.66
360	178.08	341.26		268.03			57.69	116.28	16.60
361		618.28		207.68				217.28	
362				503.30					
363				1032.16					
364	250.11	94.54	155.45	714.43	567.82		57.20	18.17	19.03
365				532.61					
366				552.94					
367				1093.21					
368		987.26		472.10				544.93	
369	359.05	648.54		327.14			194.80	323.95	82.65
370		992.98		467.27				570.72	
371		555.38		386.49				226.34	
372				822.92					
373	367.94	1062.17	709.69	754.81	866.40		176.12	617.72	111.39
374	100.62	104.26		90.13			24.41	24.06	5.54
375				287.29					
377		1.05		0.86				1.01	
378				525.80					
379				648.43					
380				1.15					
381	349.53	806.74		495.13			190.64	368.50	69.68
382	0.93	1.05		1.07			0.95	1.07	0.80
383	308.53	691.74		310.52			126.22	277.71	34.36
384	423.68	875.57		567.48			290.88	560.26	140.75
385				601.75					
386	1.05	1.04		446.12	464.11		1.15	1.03	0.84
387		598.74		316.10				220.09	
388	456.60	924.88	627.20	496.15	596.86		298.68	623.40	188.99
389		355.21		205.18				98.57	
390		636.70		341.28				317.60	
391		615.70		142.50				160.54	
392				509.20					
393	263.50	704.89		397.18			118.09	335.73	40.96
394		1340.33		752.37				832.36	
395				591.97					
396				553.32					
397		407.53		231.47				122.95	
398				648.01					
399		556.95		171.02				157.94	
400		1202.88		748.10				510.98	
401		418.06		201.06				186.92	
402	485.92	1356.49		701.68			327.63	936.11	171.64
403	257.07	762.96		302.07			87.60	353.99	24.44
404		363.12		198.49				115.92	
405				486.64					
407				784.82					
408	601.28	1196.34	847.10	861.47	998.68		267.62	497.99	95.29
409				552.36					
410	141.97	592.17		249.30			23.40	178.63	2.66
411	0.78	1.05		576.30	747.57		0.81	1.03	0.82
412				689.58					
413		969.16		366.69				416.00	
414	374.45	658.47	567.00	667.48	610.89		203.16	443.30	92.27
415	409.68	962.17		434.95			261.42	539.48	127.87

TABLE 1-1-continued

Binding to Wild-Type SARS-CoV-2 (Bead Assay)									
Antibody ID	A	B	C	D	E	F	G	H	I
416				110.41					
417	421.91	961.35	589.51	460.10	578.51		234.88	506.90	101.29
418				753.99					
419		3941.02	1016.73	768.38	1016.40			1171.12	
420		149.82		189.44				25.22	
421				505.29					
422				1249.68					
423				1186.89					
426	0.79	1.04		550.97	555.80		0.82	0.98	0.92
427				494.44					
428	145.80	342.30		128.38			40.41	92.27	7.50
429	502.90	1086.89		598.82			382.83	697.73	244.98
430	120.41	101.30		293.74			14.63	10.33	1.97
431				714.53					
432	0.97	0.72		0.96			1.02	0.88	0.90
433		1325.24		393.10				493.87	
434				1.33					
435				605.81					
436	7.84	39.28		11.70			1.15	2.00	1.04
437				591.26					
438	457.94	1135.88		758.99			215.69	472.36	68.86
439				226.66					
440		693.42		350.78				88.25	
441		52.86		129.79				1.31	
442	415.35	961.18	528.51	427.47	501.23		266.41	518.68	119.97
443				1.06					
444		0.97		0.91				0.90	
445		1554.93	805.30	668.49	839.72			771.63	
446				435.44					
447		1690.11	1010.32	1018.92	1127.60			1413.52	
448		779.34		371.76				229.43	
449		678.26		441.01				195.25	
450		1.30		0.89				1.11	
451		395.86		197.42				129.42	
452				853.50					
453	263.82	587.52		418.12	532.22		131.98	252.17	46.46
454				1094.42					
455		533.89		333.27				175.30	
456				595.49					
457	1.04	1.28		564.89	631.84		0.98	1.03	0.93
458		218.68		318.89				56.46	
459	459.63	911.52		494.38			298.31	483.23	132.85
460		732.42		422.97				458.44	
461		1762.41		762.70				1006.02	
462		1215.76	596.85	719.50	728.93			720.21	
463		918.84		447.44				570.90	
464		1.27		0.99				1.04	
465		1164.65		171.45				359.00	
466	0.94	1.12		679.43	836.59		0.88	1.01	0.78
467		724.17		378.10				214.72	
468		1892.20		818.27				1154.67	
469		729.88		240.83				125.22	
470				1178.02					
471				770.02					
472		98.26		57.38				22.33	
473	283.41	656.91		493.49			122.90	488.36	45.27
474				1.47					
475				682.90					
476				901.67					
477				846.89					
478	425.63	801.68		500.81			266.86	406.41	117.86
479		930.19	583.15	479.93	631.87			568.45	
480				689.79					
481		1421.56	859.59	872.99	932.50			986.47	
482		871.96		587.30				369.20	
483		970.31	662.14	417.38	627.17			572.28	
484	377.79	736.80		487.79			195.35	324.87	72.89
485				509.90					
486	339.44	725.73		461.05			195.81	375.98	83.03
487				661.24					
488	727.67	1441.72	903.25	848.16	982.36		431.29	751.45	157.90
489		832.67		460.46				483.69	
490		1.04		285.49	409.18			1.10	

TABLE 1-1-continued

Binding to Wild-Type SARS-CoV-2 (Bead Assay)									
Antibody ID	A	B	C	D	E	F	G	H	I
491	422.62	864.59		492.61	598.95		262.46	435.65	104.91
492				663.54					
493		1009.77		334.97				381.21	
494		921.31	680.18	614.14	721.68			409.53	
495		1085.27		383.85				530.14	
496				1086.41					
497				629.60					
498				497.11					
499		1122.21		386.00				431.73	
500	318.49	630.75		481.37			182.05	291.84	70.75
501		1638.91		769.93				507.57	
502	550.86	1144.38		681.93			282.78	554.06	87.31
503		151.56		790.27				8.59	
504				708.99					
505				657.81					
506		2283.31	1111.78	823.97	1028.39			1197.23	
507				571.25					
508	120.54	401.03		123.83			19.72	147.75	3.09
509				496.12					
510				0.95					
511		835.54		327.79				303.14	
512				547.64					
513		691.44		307.73				224.48	
514				1.60					
515		1461.98		691.55				736.01	
516				1.48					
517				433.64					
518	371.72	782.60		443.17			171.94	400.93	71.95
519	56.56	122.97		74.58			4.04	26.55	1.13
520		461.48		201.34				126.55	
521		49.21		34.05				2.06	
522				433.00					
523				30.76					
524		1.51		1.24				1.08	
525		533.28		419.16				180.43	
526	197.27	524.47		138.20			31.28	114.67	5.24
527				148.09					
528	105.01	123.50		308.96			14.03	285.35	2.44
529	1.06	1.34		12.21			0.92	22.74	0.80
530	150.10	295.90		151.58			54.36	149.84	11.72
531				380.55					
532	817.45	1566.59		999.97			663.17	1052.24	368.40
533				23.40					
534		695.90		218.50				404.28	
535		1425.49		948.76				1098.37	
536				643.45					
537				786.79					
538				607.35					
539		1951.09	983.89	799.27	990.16			1179.41	
540		769.66	470.19	345.97	485.40			310.25	
541				1.24					
542				597.71					
543				802.50					
544				639.22					
545				938.88					
546				1161.05					
547				605.82					
548				642.50					
549		1350.94	901.67	1.04	1.15			656.49	
550				1.53					
551				649.34					
552		1197.75	607.47	517.18	660.30			483.39	
553		412.96	306.84	457.47	546.27			110.66	
554		821.54		437.86				364.56	
555		1852.09	605.08	1113.30	1233.03			1379.32	
556				625.23					
557		765.62		538.53				362.93	
558		964.14		373.03				410.26	
559		1183.02		0.97	1.15			477.48	
560				991.42					
561		788.14		572.82				421.78	
562		742.77	466.07	402.17	491.67			365.23	
563				512.96					

TABLE 1-1-continued

Binding to Wild-Type SARS-CoV-2 (Bead Assay)								
Antibody ID	A	B	C	D	E	F	G	H
564				522.61				
565		1.29		0.97				1.11
566				613.29				
567		233.98		153.96				68.44
568		156.72		53.37				25.21
569				478.07				
570		355.29		188.82				130.30
571				337.90				
572		0.98		471.28	567.62			1.07
573		0.96		295.02	256.90			0.94
574		227.18		172.40				68.46
575		831.67		562.18				331.22
576				565.54				
577		0.93		1.06				0.94
579		8.84		34.80				1.69
580				771.66				
581				0.99				
582				676.11				
583				717.61				
584		155.46		66.16				8.93
585		650.62		311.18				349.37
586				1.18				
587		77.27		1.31				12.82
588				1.46				

TABLE 1-2

Binding to Wild-Type SARS-CoV-2 (Live Cell Assay)								
Antibody ID	A	B	C	D	E	F	G	H
258		11.99					9.66	
259				2.58				
260				10.26				
261	42.06	31.24				28.40	21.75	10.65
262		1.49					0.89	
263				22.48				
264				10.26				
265				21.45				
266				10.28				
267				2.81				
268	12.60	1.13				7.83	0.98	4.75
269		13.32					9.09	
270				1.87				
271				7.98				
272		24.07					14.42	
273				16.95				
274		21.26		1.55			12.33	
275	11.71	8.94				8.65	6.23	6.29
276				9.24				
277	12.15	10.20				10.21	8.09	8.64
278	25.82	27.27				18.90	16.52	7.17
279		8.59					3.84	
280	33.46	25.01				26.29	18.55	15.29
281				24.33				
283		9.63					7.62	
284				30.25				
285	36.19	31.07				26.21	20.43	16.70
286	12.38	8.02				10.10	4.90	8.86
287	38.18	30.60				29.26	21.95	12.75
288				24.27				
290				0.98				
291		16.68					15.08	
292	23.77	16.28	17.03	18.57		19.75	14.73	10.99
294				22.24				
295				9.67				
296		1.06					1.12	
297		11.72					10.67	
298		23.17					17.94	
299				2.11				

TABLE 1-2-continued

Binding to Wild-Type SARS-CoV-2 (Live Cell Assay)								
Antibody ID	A	B	C	D	E	F	G	H
300				9.71				
301				29.71				
302	46.22	31.31				17.25	8.93	5.13
303				9.59				
304		1.09					1.12	
305		27.18					15.62	
386				2.57				
307		11.40					5.38	
308		17.66					15.99	
309	11.93	9.67	8.52	9.04		10.55	8.16	7.92
310		12.34					9.68	
311		18.34					10.39	
312		9.75					3.94	
313		9.31					7.31	
314		13.00					3.95	
315				8.85				
316				10.68				
317				11.23				
318		22.75					17.94	
319				10.40				
320				20.66				
321		7.38					2.54	
322		4.94					1.65	
323				20.62				
324				9.44				
325		1.14					1.08	
326				10.11				
327				9.54				
328				1.60				
329				9.41				
330		20.60					12.08	
331				20.08				
332				1.75				
333				1.2.7				
334		7.68					5.56	
335	1.28	0.97		6.41	7.43	1.26	0.98	1.24
335				7.43				
336		18.45					5.57	
337				2.08				
338		9.48					7.51	
339				21.25				
340				33.17				
341	1.20	1.04				1.15	0.98	1.17
342				4.69				
343				1.12				
344				7.55				
345				11.71				
346	6.41	4.41				2.55	1.66	1.53
347				7.81				
348				19.61				
349				1.43				
350		53.42					46.10	
351				1.06				
352		18.45					3.63	
353	16.51	12.15				12.17	8.68	10.36
354	9.83	7.81				6.61	4.70	3.87
355		7.00					1.75	
356	42.33	31.24				29.98	22.58	12.63
357		7.08					5.2.6	
358	13.57	10.65				10.75	8.15	8.99
359	11.79	9.74				9.95	7.43	6.54
360	10.31	7.78				6.94	4.91	3.72
361		9.35					7.51	
362				9.45				
363				16.75				
364	30.75	6.57	19.95	26.68		19.82	1.96	7.22
365				10.70				
366				8.97				
367				26.48				
368		22.03					15.87	
369	11.08	8.80				8.86	6.37	5.56
370		12.33					9.66	
371		1.30					0.97	

TABLE 1-2-continued

Binding to Wild-Type SARS-CoV-2 (Live Cell Assay)								
Antibody ID	A	B	C	D	E	F	G	H
372				30.17				
373	62.79	27.08	19.81	18.43		34.64	15.12	16.55
374	2.59	1.19				1.67	0.85	1.62
375				5.57				
377		0.83					0.78	
378				25.34				
379				1.63				
380				1.06				
381	50.53	24.54				35.01	10.81	14.16
382	1.20	0.78				1.20	0.69	1.27
383	3.83	1.61				1.88	1.00	1.52
384	22.52	10.96				16.83	7.96	14.30
385				10.46				
386	1.26	0.81		8.70		1.15	0.71	1.13
387		6.22					2.13	
388	30.71	12.40	9.24	9.33		19.36	8.74	17.36
389		1.13					0.72	
390		0.96					0.73	
391		9.53					2.91	
392				1.30				
393	24.09	10.95				16.29	7.55	10.45
394		22.14					15.91	
395				7.80				
396				3.58				
397		8.87					5.24	
398				8.33				
399		20.67					9.23	
400		22.67					10.64	
401		8.35					7.68	1
402	44.67	24.74				36.54	17.12	20.05
403	21.33	9.50				15.66	7.34	10.23
404		7.20					5.05	
405				9.70				
407				2.66				
408	34.15	12.70	9.22	12.09		11.01	3.45	4.17
409				13.84				
410	9.01	5.00				2.75	1.33	1.66
411	1.49	0.77		9.87		1.34	0.66	1.37
412				22.64				
413		1.31					1.11	
414	52.46	11.67	29.15	26.17		49.60	15.52	29.37
415	26.51	8.21				18.91	8.01	16.32
416				1.58				
417	6.14	2.15	2.43	1.91		3.64	1.45	2.89
418				10.87				
419		22.69	25.75	24.97			20.28	
420		3.95					1.99	
421				9.23				
422				19.62				
423				14.56				
426	2.17	1.37		28.78		2.00	0.98	2.08
427				7.71				
428	12.60	4.03				4.93	1.94	2.57
429	26.33	10.61				19.31	9.26	17.17
430	22.37	8.29				16.88	7.40	15.03
431				10.72				
432	1.25	0.83				0.94	0.81	1.12
433		12.40					5.14	
434				1.02				
435				7.58				
436	2.44	1.67				1.13	0.87	1.18
437				8.00				
438	27.82	13.19				15.98	11.33	5.97
439				8.20				
440		13.80					3.53	
441		9.54					2.76	
442	13.93	8.33	8.58	8.39		9.95	7.98	8.24
443				1.18				
444		1.37					1.09	
445		18.08	14.73	14.59			9.16	
446				7.99				
447		16.90	15.76	16.77			17.63	
448		6.22					2.45	

TABLE 1-2-continued

Binding to Wild-Type SARS-CoV-2 (Live Cell Assay)								
Antibody ID	A	B	C	D	E	F	G	H
449		23.27					19.58	
450		1.05					0.85	
451		7.40					6.12	
452				26.81				
453	12.56	7.65		9.98		9.58	7.71	5.44
454				29.60				
455		7.87					8.10	
456				9.25				
457	1.28	0.90		14.83		1.10	0.88	1.04
458		7.35					4.19	
459	13.07	8.75				9.87	8.14	8.23
460		1.13					0.97	
461		11.29					4.47	
462		15.89	12.71	13.17			10.13	
463		11.71					9.98	
464		0.87					0.83	
465		22.04					9.46	
466	1.14	0.89		1.91		0.90	0.85	1.00
467		8.20					6.50	
468		22.65					17.21	
469		5.29					1.43	
470				20.69				
471				3.03				
472		1.05					0.91	
473	43.89	28.68				31.30	23.32	24.10
474				1.49				
475				32.23				
476				18.19				
477				30.73				
478	17.42	8.81				10.65	7.13	8.31
479		2.29	1.47	1.28			1.42	
480				9.77				
481		15.65	15.58	18.13			14.00	
482		4.48					2.31	
483		26.11	30.69	26.79			20.44	
484	20.01	9.73				11.41	8.15	9.01
485				1.37				
486	13.61	8.35				9.20	7.06	7.04
487				1.76				
488	32.82	12.66	24.35	23.96		17.37	11.34	6.50
489		11.32					8.74	
490		1.07		1.28			1.08	
491	14.25	8.47		9.96		10.06	7.86	7.40
492				9.25				
493		1.26					1.10	
494		14.22	11.72	12.40			7.29	
495		1.23					0.83	
496				19.70				
497				28.32				
498				1.69				
499		1.35					0.90	
500	13.37	8.58				9.94	7.56	7.70
501		32.38					4.87	
502	28.56	18.49				18.98	14.68	7.68
503		1.87					1.22	
504				2.78				
505				1.55				
506		30.07	30.76	22.32			18.96	
507				1.51				
508	7.41	2.94				2.50	1.11	1.48
509				10.19				
510				1.18				
511		9.99					7.72	
512				30.25				
513		8.42					7.37	
514				1.04				
515		10.37					3.93	
516				3.31				
517				10.32				
518	16.49	8.65				10.43	7.25	8.31
519	11.35	5.82				5.56	3.78	2.99
520		8.90					7.29	
521		2.42					0.86	

TABLE 1-2-continued

Binding to Wild-Type SARS-CoV-2 (Live Cell Assay)								
Antibody ID	A	B	C	D	E	F	G	H
522				9.18				
523				5.18				
524		0.58					0.55	
525		8.70					7.47	
526	10.16	5.86				3.45	2.03	2.17
527				10.36				
528	23.21	11.94				8.72	5.31	2.99
529	1.65	0.78				1.37	0.74	1.47
530	11.80	5.76				3.97	2.24	2.02
531				7.45				
532	40.22	15.17				27.74	20.08	14.53
533				4.31				
534		12.13					5.22	
535		50.14					48.91	
536				2.36				
537				4.42				
538				1.96				
539		24.95	16.77	21.19			12.22	
540		8.05	8.45	8.15			7.25	
541				1.10				
542				7.52				
543				13.30				
544				2.48				
545		1		20.57				
546				19.29				
547				10.64				
548				9.26				
549		14.91	1.17	1.16			11.73	
550				1.12				
551				2.48				
552		4.37	3.42	4.76			1.58	
553		0.85	1.21	1.47			0.78	
554		8.67					7.21	
555		13.84	27.55	28.72			16.18	
556				7.59				
557		10.02					7.10	
558		9.48					8.06	
559		15.91		1.38			8.40	
560				22.81				
561		10.44					7.44	
562		17.30	7.64	8.57			15.43	
563				9.67				
564				10.78				
565		1.31					1.11	
566				31.54				
567		8.03					5.10	
568		8.70					3.85	
569				8.78				
570		16.53					14.76	
571				8.30				
572		0.72		9.54			0.67	
573		1.09		21.55			1.09	
574		7.47					5.06	
575		24.84					21.10	
576				8.75				
577		1.26					1.14	
579		1.61					1.16	
580				11.74				
581				36.09				
582				19.63				
583				8.54				
584		3.72					1.24	
585		16.33					13.84	
586				1.18				
587		5.47					2.08	
588				32.67				

TABLE 1-3

Reactivity with SARS-CoV-2 mutants							
Anti-body ID	A	B	C	D	E	F	G
258	23.72		10.33			42.93	6.88
259			1.83				
260			11.04				
261	91.23		41.86			59.94	87.01
262	1.44		1.63			1.08	1.24
263			17.27				
264			11.10				
265			14.53				
266			10.84				
267			3.03				
268	1.11		11.35			8.90	8.18
269	22.84		11.58			25.35	4.83
270			2.45				
271			8.53				
272	88.35		40.81			40.41	79.72
273			16.00				
274	65.52		4.35	1.01		31.84	61.17
275	24.06		12.79			38.40	6.84
276			11.11				
277	26.19		11.67			41.89	9.35
278	76.99		27.61			61.95	70.91
279	11.80		8.20			13.40	2.37
280	66.12		30.06			42.78	68.35
281			20.29				
283	30.58		16.31			36.22	8.17
284			22.02				
285	67.60		36.55			48.91	84.33
286	8.64		11.86			28.16	6.17
287	88.80		38.05			60.87	87.37
288			18.73				
290			1.51				
291	31.06		16.58			33.92	8.03
292	47.28	13.15	25.85	12.92		23.61	42.77
294			18.44				
295			9.93				
296	0.68		1.27			1.01	0.95
297	27.96		12.88			44.70	8.63
298	69.72		49.21			35.71	66.84
299			2.18				
300			10.19				
301			21.31				
302	37.89		24.64			26.48	45.35
303			10.02				
304	1.00		0.66			0.99	0.97
305	86.93		58.08			41.01	78.88
306			2.89				
307	13.85		8.22			8.47	9.73
308	30.41		15.62			33.36	7.93
309	26.61	10.37	11.69	10.70		41.22	8.01
310	20.36		13.44			28.06	5.35
311	56.69		30.52			29.57	59.66
312	24.72		13.46			11.26	24.51
313	30.28		14.80			35.14	8.14
314	42.80		51.60			19.62	30.43
315			8.39				
316			10.43				
317			10.32				
318	34.78		17.97			40.26	9.92
319			10.57				
320			19.72				
321	15.85		4.03			3.81	11.75
322	2.29		4.95			1.54	3.07
323			18.97				
324			24.90				
325	2.31		1.17			1.65	1.11
326			10.16				
327			10.39				
328			4.00				
329			10.62				
330	55.05		37.16			22.41	56.53
331			18.09				

TABLE 1-3-continued

Reactivity with SARS-CoV-2 mutants							
Anti-body ID	A	B	C	D	E	F	G
332			1.93				
333			1.19				
334	23.68		12.99			29.66	6.38
335	1.00		9.08	7.26	6.07	1.78	1.78
335			6.07				
336	31.95		34.03			13.90	29.45
337			2.25				
338	28.71		13.51			33.19	7.81
339			18.23				
340			26.27				
341	1.06		1.01			2.18	2.22
342			4.88				
343			1.22				
344			8.15				
345			9.41				
346	9.53		5.63			7.05	8.57
347			8.57				
348			14.18				
349			1.65				
350	99.30		26.95			59.37	90.02
351			0.90				
352	19.28		18.13			1.08	0.94
353	29.41		18.82			45.27	8.84
354	17.09		9.80			33.17	5.07
355	16.81		12.11			8.08	20.27
356	95.46		36.82			60.38	90.79
357	24.02		12.06			28.02	5.37
358	27.31		13.63			38.39	8.74
359	24.40		10.06			38.77	6.88
360	17.97		10.00			33.90	4.92
361	30.83		14.44			28.28	8.06
362			10.28				
363			15.05				
364	9.44	17.75	35.04	21.95		2.12	2.22
365			11.22				
366			9.74				
367			16.30				
368	68.21		39.13			31.55	64.35
369	21.18		9.35			36.25	6.12
370	34.51		16.97			35.28	9.47
371	2.93		2.27			3.85	1.61
372			18.09				
373	59.91	18.35	27.26	15.41		42.79	59.37
374	2.38		1.22			3.10	1.11
375			3.67				
377	1.05		1.09			0.99	0.92
378			14.35				
379			1.30				
380			1.27				
381	62.74		29.43			43.56	67.87
382	0.96		0.88			0.97	0.77
383	3.84		1.80			11.38	6.38
384	30.14		15.65			44.85	8.74
385			8.16				
386	1.08		11.73	10.06		1.20	0.80
387	15.80		11.04			6.69	15.92
388	28.17	10.64	12.43	10.46		43.43	8.68
389	3.16		1.24			2.29	1.52
390	2.65		1.36			3.43	1.15
391	36.73		12.45			18.77	35.65
392			1.02				
393	25.98		13.62			39.34	7.44
394	70.93		44.22			37.83	63.40
395			8.92				
396			3.34				
397	29.46		17.73			32.81	7.97
398			9.00				
399	64.80		25.35			30.56	70.12
400	65.26		51.70			37.37	66.79
401	30.99		14.52			35.51	8.76
402	71.22		24.90			46.90	67.78

TABLE 1-3-continued

Anti-body ID	Reactivity with SARS-CoV-2 mutants						
	A	B	C	D	E	F	G
403	23.69		10.33			35.65	7.47
404	23.52		12.58			27.36	6.02
405			7.51				
407			2.28				
408	39.16	9.44	22.27	12.51		29.52	36.22
409			9.21				
410	8.75		6.87			7.64	17.91
411	1.07		16.28	7.86		1.06	0.79
412			13.12				
413	3.75		1.67			3.41	1.32
414	75.18	20.88	35.05	23.48		49.20	78.55
415	29.15		10.95			42.39	8.14
416		1.09					
417	7.38	2.18	2.13	2.04		7.71	2.12
418			11.97				
419	86.33	21.62	52.30	21.43		40.72	70.63
420	12.38		13.73			7.59	2.50
421			9.33				
422			15.24				
423			15.26				
426	1.52		32.58	23.81		1.37	1.17
427			5.61				
428	8.00		4.93			14.45	1.51
429	26.33		12.20			40.90	8.50
430	26.48		11.04			38.30	9.17
431			11.65				
432	1.04		0.82			1.06	0.88
433	28.73		19.75			12.34	23.05
434			1.34				
435			7.93				
436	1.70		0.85			2.31	1.00
437			4.71				
438	73.72		26.10			43.96	71.20
439			9.01				
440	17.38		16.55			7.17	15.81
441	12.42		9.04			6.24	14.86
442	25.79	10.40	10.26	10.31		38.14	7.29
443			0.93				
444	1.32		2.50			1.10	1.09
445	58.93	13.58	28.73	13.67		30.82	54.86
446			6.16				
447	51.18	11.25	21.65	11.18		35.27	43.98
448	14.84		6.99			5.79	13.34
449	82.69		51.64			36.87	68.06
450	0.93		1.27			1.01	0.95
451	23.37		13.85			29.93	6.29
452			0.97				
453	25.63		10.63	7.84		39.35	7.41
454			19.73				
455	30.58		14.24			32.32	8.24
456			8.16				
457	1.01		22.86	14.46		1.14	0.75
458	22.72		14.68			27.44	6.23
459	26.02		11.69			40.82	8.07
460	2.05		1.63			2.42	1.07
461	30.03		24.42			6.89	27.84
462	50.03	9.72	18.76	9.59		42.75	54.75
463	38.25		9.56			43.84	8.36
464	1.04		1.00			1.04	0.86
465	75.70		24.73			37.29	68.30
466	1.15		3.81	1.94		1.10	0.75
467	27.23		16.17			30.57	7.54
468	70.65		53.95			34.57	57.63
469	6.59		10.41			3.33	5.13
470			7.63				
471			2.75				
472	2.35		2.49			3.19	1.22
473	85.60		40.48			52.91	85.20
474			1.05				
475			20.18				
476			14.06				

TABLE 1-3-continued

Anti-body ID	Reactivity with SARS-CoV-2 mutants						
	A	B	C	D	E	F	G
477			23.93				
478	24.54		9.85			38.85	6.68
479	5.19	1.72	1.26	1.30		4.31	1.54
480			10.68				
481	67.86	12.60	22.27	14.26		46.55	62.21
482	12.36		4.37			2.89	14.35
483	103.79	22.62	55.72	21.95		48.21	98.05
484	28.39		11.13			43.00	8.15
485			1.11				
486	23.47		10.13			37.05	6.79
487			1.28				
488	75.86	22.75	34.95	22.44		52.57	79.42
489	36.73		8.61			40.68	7.95
490	1.15		1.13	1.52		1.14	1.03
491	26.57		11.57	7.97		38.81	7.45
492			7.71				
493	4.63		2.26			4.82	1.72
494	44.93	9.51	32.44	9.93		33.28	55.86
495	3.47		1.65			3.82	1.43
496			12.62				
497			22.21				
498			0.95				
499	4.39		2.60			1.18	4.30
500	25.40		12.06			39.05	6.98
501	38.82		58.69			29.96	46.12
502	75.66		33.69			51.78	77.33
503	2.13		56.95			1.64	2.24
504			2.47				
505			1.00				
506	84.86	21.28	55.34	19.71		42.26	69.46
507			1.12				
508	3.16		1.73			2.71	3.99
509			11.41				
510			1.06				
511	30.09		15.44			31.27	8.27
512			17.78				
513	29.60		14.98			35.81	8.11
514			1.39				
515	20.24		22.33			9.43	21.56
516			2.61				
517			11.46				
518	27.07		11.02			39.74	7.56
519	16.48		6.13			30.76	4.93
520	27.90		14.55			31.20	7.60
521	2.72		5.78			1.68	2.68
522			6.75				
523			3.76				
524	0.99		1.25			1.12	0.85
525	28.03		18.43			32.40	7.12
526	8.86		2.82			9.69	6.76
577			7.97				
528	19.27		9.66			31.23	3.71
529	1.19		2.02			0.98	0.77
530	11.91		3.47			8.39	8.68
531			8.14				
537	82.82		37.09			45.13	74.93
533			4.86				
534	53.33		17.02			42.90	55.16
535	108.81		32.53			64.49	93.16
536			2.59				
537			3.74				
538			1.48				
539	62.31	17.22	43.72	18.37		27.05	52.98
540	29.98	9.07	15.43	9.58		30.28	8.41
541			1.60				
542			8.41				
543			8.92				
544			1.58				
545			14.34				
546			18.95				
547			11.36				

TABLE 1-3-continued

Reactivity with SARS-CoV-2 mutants							
Anti-body ID	A	B	C	D	E	F	G
548			10.05				
549	84.96	16.67	0.64	1.50		55.32	82.00
550			1.46				
551			1.70				
552	14.18	2.77	11.30	3.46		13.19	23.53
553	2.42	1.20	1.20	1.57		2.72	1.20
554	25.25		11.74			38.34	7.49
555	62.72	19.48	28.50	20.07		42.59	68.10
556			7.35				
557	23.76		12.51			36.28	7.35
558	30.48		16.05			33.78	8.19
559	49.04		1.20	1.10		25.42	49.47
560			14.80				
561	26.03		13.15			40.78	8.07
562	35.88	9.85	9.23	10.13		42.03	8.13
563			10.47				
564			11.65				
565	1.05		0.94			0.96	0.93
566			18.74				
567	21.30		10.73			28.64	5.41
568	21.69		10.32			8.40	20.99

TABLE 1-3-continued

Reactivity with SARS-CoV-2 mutants							
Anti-body ID	A	B	C	D	E	F	G
569			10.37				
570	36.80		9.05			43.77	8.96
571			6.77				
572	0.99		11.72	10.88		1.08	0.86
573	1.17		23.89	17.83		1.05	0.94
574	19.94		13.15			27.47	5.03
575	90.42		57.96			51.44	102.62
576			10.04				
577	1.68		1.63			1.92	0.99
579	2.02		1.32			1.95	1.89
580			8.07				
581			60.55				
582			12.97				
583			11.39				
584	1.34		0.96			0.99	0.92
585	29.09		15.22			33.00	7.49
586			1.53				
587	10.99		1.93			8.33	2.38
588			44.93				

TABLE 1-4

Cross-reactivity with SARS								
Antibody ID	A	B	C	D	E	F	G	H
258		84.75					42.13	
259				1.03				
260				107.52				
261	0.89	1.09				0.93	1.03	0.91
262		0.87					0.88	
263				1.13				
264				55.94				
265				1.09				
266				33.42				
267				16.93				
268	25.86	1.17				10.35	1.02	2.84
269		55.38					33.51	
270				1.34				
271				27.37				
272		1.18					1.04	
273				1.23				
274		1.13		1.11			1.08	
275	36.29	64.76				16.13	32.17	7.84
276				117.48				
277	40.33	77.67				23.59	45.94	9.94
278	0.96	37.87				1.03	10.66	0.98
279		1.27					1.02	
280	1.03	1.09				1.00	1.05	1.00
281				1.10				
283		1.10					1.18	
284				1.20				
285	1.01	1.03				1.06	0.99	1.01
286	22.12	22.73				7.62	6.49	1.88
287	0.93	1.12				0.92	1.05	0.91
288				1.00				
290				1.25				
291		58.79					31.31	
292	0.89	1.09	0.96	1.14		0.96	1.05	0.92
294				1.07				
295				1.24				
296		0.99					1.08	
297		102.18					57.83	
298		1.13					1.11	
299				1.13				
300				74.40				
301				1.45				

TABLE 1-4-continued

Cross-reactivity with SARS								
Antibody ID	A	B	C	D	E	F	G	H
302	1.06	2.73				1.08	1.34	1.06
303				21.75				
304		1.13					1.09	
305		1.09					1.04	
306				1.39				
307		1.21					1.03	
308		48.28					20.77	
309	60.50	90.38	49.83	48.65		37.72	60.05	20.77
310		5.18					1.69	
311		1.07					0.99	
312		1.11					1.04	
313		4.93					1.51	
314		1.06					1.05	
315				1.07				
316				28.80				
317				1.14				
318		198.29					130.24	
319				30.64				
320				989.19				
321		1.14					1.01	
322		0.95					0.93	
323				1.29				
324				1.14				
325		1.12					1.03	
326				11.52				
327				33.63				
328				1.22				
329				2.46				
330		1.10					1.11	
331				1.34				
332				0.99				
333				1.21				
334		12.47					6.78	
335	1.06	1.04		5.85	10.23	1.08	0.96	1.10
335				10.23				
336		1.09					1.08	
337				439.36				
338		84.79					36.61	
339				1.27				
340				1.26				
341	1.02	1.11				1.13	0.97	1.12
342				1.16				
343				1.19				
344				9.62				
345				1.09				
346	1.09	1.02				1.13	0.99	1.07
347				51.45				
348				1.16				
349				1.12				
350		0.94					1.05	
351				1.08				
352		0.92					1.08	
353	38.44	73.63				27.63	42.98	15.85
354	1.20	1.90				1.08	1.20	1.04
355		0.88					0.95	
356	1.05	1.17				1.09	1.11	1.12
357		11.83					4.75	
358	22.46	29.76				5.87	10.02	1.77
359	1.04	3.15				1.20	1.26	1.08
360	1.17	1.03				1.18	1.03	1.09
361		64.67					39.66	
362				38.15				
363				1.09				
364	1.08	0.96	1.00	1.11		1.18	0.98	1.09
365				38.59				
366				1.23				
367				1.08				
368		1.09					1.09	
369	1.58	2.10				1.19	1.11	1.11
370		200.26					132.64	
371		1.21					0.98	
372				1.08				
373	1.78	7.32	3.18	1.54		0.90	1.90	0.71

TABLE 1-4-continued

Cross-reactivity with SARS								
Antibody ID	A	B	C	D	E	F	G	H
374	0.80	1.21				0.78	1.17	0.70
375				1.22				
377		1.14					0.98	
378				0.99				
379				5.90				
380				1.21				
381	0.89	1.16				0.92	1.20	0.72
382	0.91	1.06				0.91	1.06	0.72
383	196.09	523.45				59.24	162.39	10.35
384	64.70	158.27				52.52	106.14	21.89
385				59.56				
386	0.99	1.01		9.91		1.11	1.03	0.81
387		1.02					1.00	
388	33.07	87.45	29.94	33.98		16.78	53.32	13.05
389		0.94					0.86	
390		1.03					1.02	
391		0.98					1.04	
392				1.08				
393	6.30	32.99				2.45	12.72	1.09
394		1.21					1.12	
395				1.16				
396				1.36				
397		1.30					1.18	
398				140.88				
399		1.03					1.00	
400		0.91					1.00	
401		20.74					10.07	
402	0.67	9.28				0.69	2.25	0.70
403	18.79	89.59				7.39	46.48	2.24
404		1.04					1.04	
405				80.94				
407				1.07				
408	0.75	1.21	1.08	1.12		0.79	1.23	0.83
409				1.03				
410	0.73	1.18				0.77	1.17	0.80
411	0.79	0.99		1.05		0.78	1.05	0.79
412				1.09				
413		1.72					1.11	
414	0.64	0.67	0.99	1.01		0.69	0.93	0.77
415	58.68	143.01				34.50	86.00	17.24
416				18.99				
417	0.82	2.52	1.26	1.12		0.72	1.28	0.76
418				66.78				
419		1.55	1.01	1.12			1.19	
420		1.48					1.10	
421				1.19				
422				1.19				
423				1.23				
426	0.77	1.06		1.16		0.81	1.06	0.91
427				1.14				
428	15.32	49.44				6.70	23.25	2.62
429	51.46	113.99				38.18	73.33	27.68
430	9.09	24.99				1.79	4.22	1.09
431				49.88				
432	0.92	0.66				0.95	0.90	0.86
433		1.28					1.03	
434				1.36				
435				1.15				
436	0.99	1.32				0.96	1.05	0.99
437				0.94				
438	0.97	1.30				0.92	1.13	0.89
439				1.23				
440		1.37					1.01	
441		1.08					1.00	
442	16.74	54.82	23.05	21.17		11.57	31.73	5.61
443				1.03				
444		1.00					1.00	
445		1.24	1.06	1.17			1.00	
446				6.58				
447		1.15	1.05	1.08			1.09	
448		1.46					1.07	
449		1.55					1.16	
450		1.49					1.19	

TABLE 1-4-continued

Cross-reactivity with SARS								
Antibody ID	A	B	C	D	E	F	G	H
451		9.59					2.09	
452				1.09				
453	33.13	94.86		67.35		16.96	39.48	5.95
454				1.10				
455		61.42					10.23	
456				676.27				
457	1.00	1.25		1.87		0.92	1.09	0.96
458		1.03					1.01	
459	52.25	121.73				35.53	67.90	20.46
460		1.27					1.05	
461		1.35					1.05	
462		1.15	1.05	1.06			1.17	
463		1.21					1.08	
464		1.43					1.12	
465		1.49					1.13	
466	0.88	1.09		1.01		0.89	1.04	0.86
467		24.35					7.32	
468		1.51					1.13	
469		1.43					1.00	
470				1.09				
471				104.04				
472		1.05					1.02	
473	0.99	1.19				0.87	1.04	0.88
474				1.10				
475				1.18				
476				1.04				
477				1.30				
478	41.47	73.13				24.94	41.14	10.35
479		1.15	1.07	1.06			1.11	
480				53.06				
481		1.16	1.05	1.05			1.02	
482		1.07					1.01	
483		1.01	1.05	1.08			1.03	
484	49.43	109.95				24.38	53.91	10.05
485				1.12				
486	15.95	38.62				6.35	15.17	2.25
487				7.25				
488	0.91	1.14	1.14	1.06		0.90	1.17	0.86
489		43.18					16.09	
490		1.12		1.10			1.04	
491	36.79	87.68		39.31		19.94	41.36	8.41
492				87.12				
493		1.38					1.07	
494		0.92	1.20	1.10			0.96	
495		693.74					205.82	
496				1.15				
497				1.23				
498				1.31				
499		1.51					1.16	
500	1.33	2.20				0.95	1.18	0.85
501		1.54					1.17	
502	0.83	1.06				0.82	1.10	0.78
503		1.12					1.02	
504				344.63				
505				1.04				
506		1.47	1.09	0.98			1.17	
507				1.07				
508	0.89	0.76				0.89	0.93	0.83
509				38.76				
510				1.03				
511		120.39					64.81	
512				1.11				
513		77.65					35.16	
514				1.15				
515		1.29					1.08	
516				1.03				
517				26.51				
518	38.75	90.18				15.26	40.14	7.54
519	1.04	1.77				0.85	1.08	0.86
520		17.66					10.65	
521		1.46					1.16	
527				1.21				
523				1.11				

TABLE 1-4-continued

Antibody ID	Cross-reactivity with SARS							
	A	B	C	D	E	F	G	H
524		1.44					1.11	
525		1.66					1.25	
526	1.17	3.35				0.87	1.46	0.80
527				31.12				
528	18.25	35.31				2.75	8.12	1.23
529	0.93	1.24				0.86	1.21	0.82
530	1.62	2.96				0.92	3.42	0.86
531				27.60				
532	0.82	1.03				0.80	1.00	0.82
533				1.23				
534		0.88					0.99	
535		0.98					1.07	
536				928.04				
537				1.04				
538				17.64				
539		1.47	1.04	1.04			1.13	
540		105.72	58.27	60.70			59.08	
541				1.20				
542				100.69				
543				1.03				
544				1.05				
545				1.13				
546				1.18				
547				48.01				
548				18.10				
549		1.08	1.00	1.39			1.02	
550				1.41				
551				5.06				
552		124.24	48.20	125.49			6.53	
553		1.09	1.07	1.08			1.06	
554		27.98					8.45	
555		1.47	1.01	1.07			1.24	
556				1.09				
557		6.36					1.72	
558		67.62					15.01	
559		1.33		1.07			1.05	
560				1.14				
561		55.84					27.28	
562		17.86	6.09	2.57			5.67	
563				35.37				
564				37.28				
565		1.27					1.08	
566				0.97				
567		1.74					1.13	
568		0.99					1.07	
569				3.31				
570		6.92					2.42	
571				1.28				
572		0.97		30.96			1.04	
573		0.94		1.18			0.92	
574		1.05					1.01	
575		0.98					1.06	
576				6.87				
577		0.92					0.95	
579		1.07					1.08	
580				1.12				
581				1.14				
582				1.13				
583				115.04				
584		1.17					1.10	
585		1.17					0.96	
586				1.36				
587		2.84					1.20	
588				1.40				

TABLE 1-5

Cross-reactivity with WIV1								
Antibody ID	A	B	C	D	E	F	G	H
258		135.44					58.99	
259				1.03				
260				220.46				
261	1.13	1.76				0.93	1.12	0.97
262		0.98					0.96	
263				1.11				
264				159.96				
265				1.12				
266				96.47				
267				1.32				
268	38.33	1.11				6.61	1.01	1.51
269		47.77					14.12	
270				1.33				
271				34.07				
272		1.32					1.14	
273				1.28				
274	1.09	1.30					119	
275	141.39	209.04				65.98	91.80	26.97
276				259.79				
277	158.96	211.28				69.95	96.81	26.46
278	1.18	59.86				1.05	13.91	1.01
279		1.16					1.05	
280	1.12	1.17				1.11	1.10	1.02
281				1122.43				
283		1.22					1.18	
284				1.10				
285	1.07	1.08				1.10	0.97	1.02
286	38.69	34.32				8.88	7.86	1.91
287	1.13	1.35				0.99	1.08	0.97
288				1.03				
290				1.22				
291		143.48					64.83	
292	1.08	1.33	1.05	1.12		1.03	1.17	1.00
294				1.39				
295				1.29				
296		0.99					1.08	
297		242.40					112.89	
298		1.23					1.15	
299				1.14				
300				243.80				
301				1.41				
302	1.15	2.81				1.08	1.20	1.11
303				36.60				
304		1.01					1.04	
305		1.22					1.11	
306				1.27				
307		1.27					1.04	
308		66.37					20.95	
309	174.39	250.30	166.19	177.69		89.72	126.19	38.21
310		52.71					20.81	
311		1.11					1.14	
312		1.19					1.10	
313		8.58					1.92	
314		1.24					1.12	
315				1.09				
316				93.76				
317				1.06				
318		290.62					166.40	
319				96.20				
320				736.12				
321		1.04					1.03	
32/		0.97					0.91	
323				1.34				
324				1.08				
325		1.30					1.10	
326				8.94				
327				87.70				
328				1.18				
329				1.69				
330		1.29					1.20	
331				1.25				
332				1.04				
333				1.18				

TABLE 1-5-continued

Cross-reactivity with WIV1								
Antibody ID	A	B	C	D	E	F	G	H
334		51.31					15.99	
335				6.68				
335	1.08	1.06		3.52	6.68	1.09	0.97	1.09
336		1.15					1.18	
337				280.42				
338		111.87					32.46	
339				1.24				
340				1.25				
341	1.18	1.07				1.17	0.95	1.10
342				1.17				
343				1.17				
344				16.59				
345				1.03				
346	1.14	1.07				1.17	1.03	1.15
347				134.18				
348				1.39				
349				1.06				
350		1.13					1.05	
351				1.12				
352		1.16					1.06	
353	159.56	250.40				96.91	133.73	45.78
354	1.20	1.51				1.09	1.19	1.10
355		0.96					1.01	
356	1.28	1.50				1.15	1.23	1.18
357		15.57					2.42	
358	51.74	67.79				8.83	16.91	2.03
359	1.16	3.70				1.17	1.27	1.14
360	1.21	1.11				1.17	1.08	1.11
361		114.31					30.73	
362				119.44				
363				1.14				
364	751.63	466.05	579.87	973.16		408.62	117.53	221.32
365				128.16				
366				1.23				
367				1.20				
368		1.32					1.20	
369	1.49	2.13				1.24	1.16	1.14
370		364.65					191.57	
371		1.27					1.09	
372				1.10				
373	2.06	9.84	6.08	1.74		0.89	1.83	0.75
374	0.89	1.32				0.82	1.21	0.73
375				398.74				
377		1.14					1.06	
378				272.12				
379				1.29				
380				1.16				
381	1.07	1.31				0.98	1.26	0.79
382	0.99	1.04				0.94	1.08	0.75
383	35.23	74.56				5.97	16.67	1.30
384	172.09	339.57				88.43	180.47	39.44
385				161.00				
386	1.12	1.06		19.61		1.18	1.05	0.83
387		1.22					1.02	
388	129.20	269.76	163.98	160.52		58.29	133.80	32.50
389		1.09					1.00	
390		1.13					1.07	
391		1.16					1.15	
392				1.04				
393	22.77	79.52				5.23	25.33	1.34
394		1.41					1.30	
395				1.18				
396				1.35				
397		1.50					1.21	
398				212.69				
399		1.24					1.12	
400		1.07					1.05	
401		75.19					28.91	
402	0.82	5.36				0.74	1.54	0.77
403	34.33	143.46				7.03	46.14	1.61
404		1.16					1.02	
405				106.45				
407				1.05				

TABLE 1-5-continued

Cross-reactivity with WIV1								
Antibody ID	A	B	C	D	E	F	G	H
408	1.08	2.03	1.51	1.78		0.86	1.48	0.88
409				1.14				
410	0.87	1.29				0.81	1.26	0.81
411	0.85	1.07		1.09		0.81	1.09	0.82
412				1.14				
413		1.60					1.14	
414	0.75	0.72	1.10	1.11		0.72	0.98	0.82
415	139.01	331.46				67.79	176.30	30.30
416				7.85				
417	0.99	2.49	1.57	1.36		0.83	1.29	0.79
418				251.53				
419		2.02	1.23	1.24			1.26	
420		1.66					1.15	
421				1.26				
422				1.71				
423				1.19				
426	0.82	1.01		1.16		0.82	1.05	0.94
427				1.11				
428	23.66	66.90				3.55	14.17	1.18
429	154.39	300.98				72.84	155.20	30.62
430	8.77	13.17				1.36	2.01	1.03
431				207.17				
432	0.94	0.64				0.94	0.91	0.94
433		1.44					1.15	
434				1.29				
435				1.17				
436	1.01	1.24				0.95	1.09	1.03
437				0.91				
438	1.05	1.34				0.94	1.18	0.97
439				1.17				
440		1.52					1.02	
441		1.14					0.97	
442	88.16	208.67	110.59	102.71		45.40	105.95	19.52
443				1.04				
444		1.13					0.98	
445		1.53	1.14	1.26			1.05	
446				4.63				
447		1.28	1.07	1.26			1.21	
448		1.59					1.11	
449		1172.33					324.64	
450		1.62					1.10	
451		122.04					34.74	
452				1.13				
453	62.35	143.21		119.96		21.75	46.89	5.99
454				1.38				
455		103.30					11.43	
456				16.69				
457	1.01	1.19		2.36		0.93	1.03	1.01
458			1.15				0.99	
459	143.64	295.91				78.05	150.10	35.62
460		1.16					1.05	
461		1.59					1.02	
462		1.19	1.09	1.06			1.25	
463		1.37					1.25	
464		1.56					1.03	
465		1.87					1.23	
466	0.92	0.98		1.07		0.90	1.01	0.84
467		52.61					7.06	
468		1.67					1.14	
469		1.59					1.15	
470				1.13				
471				53.28				
472		1.23					1.08	
473	0.97	1.17				0.87	1.04	0.91
474				1.07				
475				1.19				
476				1.09				
477				1.45				
478	85.64	144.18				34.13	48.28	9.71
479		1.27	1.16	1.27			1.10	
480				217.74				
481		1.16	1.12	1.23			1.16	
482		1.21					1.14	

TABLE 1-5-continued

Cross-reactivity with WIV1								
Antibody ID	A	B	C	D	E	F	G	H
483		1.17	1.10	1.07				1.00
484	91.14	178.22				32.22	61.46	8.95
485				1.08				
486	46.38	102.09				13.69	32.84	3.46
487				1.23				
488	1.21	1.69	1.48	1.42		0.94	1.26	0.96
489		81.60					25.93	
490		1.07		1.14			1.03	
491	119.09	253.69		144.60		59.16	116.69	24.00
492				175.72				
493		1.86					1.22	
494		1.14	1.15	1.19			0.96	
495		259.73					46.61	
496				1.12				
497				1.23				
498				1.23				
499		1.87					1.26	
500	1.38	2.32				0.92	1.21	0.91
501		1.72					1.24	
502	184.46	201.31				19.08	21.84	1.33
503		1.17					1.08	
504				275.76				
505				1.03				
506		1.70	1.14	1.10			1.19	
507				1.12				
508	0.99	1.68				0.89	1.03	0.89
509				116.54				
510				0.94				
511		243.32					72.04	
512				1.13				
513	213.97					56.68		
514				1.14				
515		1.42					1.11	
516				1.05				
517				91.64				
518	104.25	208.45				34.85	91.39	13.99
519	1.05	1.54				0.86	1.16	0.86
520		41.49					7.26	
521		1.59					1.10	
522				1.19				
523				1.08				
524		1.60					1.15	
525		1.89					1.24	
526	1.13	3.01				0.88	1.32	0.86
527				25.96				
528	6.20	6.34				1.11	1.61	0.96
529	0.95	1.16				0.85	1.20	0.85
530	1.76	2.72				0.89	4.02	0.90
531				80.74				
532	1.17	1.46				0.94	1.21	0.91
533				1.30				
534		1.36					1.11	
535		1.20					1.14	
536				515.71				
537				1.05				
538				12.14				
539		1.78	1.10	1.19			1.13	
540		200.24	109.04	116.13			64.75	
541				1.17				
542				179.91				
543				1.10				
544				1.30				
545				1.26				
546				2.57				
547				162.78				
548				75.23				
549		1.60	1.41	1.14			1.23	
550				1.42				
551				149.48				
552		8.82	3.07	5.85			1.47	

TABLE 1-5-continued

Antibody ID	Cross-reactivity with WIV1							
	A	B	C	D	E	F	G	H
553		1.18	1.10	1.23			1.14	
554		44.29					8.90	
555		3.87	1.09	2.83			1.93	
556				1.15				
557		6.31					1.79	
558		139.72					28.48	
559		1.41		1.00			1.06	
560				1.18				
561		141.25					56.86	
562		25.48	13.18	2.20			5.98	
563				153.89				
564				124.07				
565		1.34					1.19	
566				1.04				
567		2.11					1.24	
568		1.13					1.06	
569				5.63				
570		74.58					28.89	
571				1.15				
572		0.94		141.13			1.11	
573		0.93		12.07			0.96	
574		1.18					1.09	
575		631.60					159.82	
576				19.42				
577		0.88					0.92	
579		1.01					1.07	
580				1.13				
581				1.07				
582				1.09				
583				254.57				
584		1.11					1.12	
585		1.03					1.07	
586				1.36				
587		2.28					1.21	
588				1.28				

Example 2

[1635] Expression and Purification of Antibodies

[1636] Antibodies were expressed and purified essentially as follows. An appropriate host cell, such as HEK 293 or CHO, was either transiently or stably transfected with an expression system for secreting antibodies using an optimal predetermined HIC:LC vector ratio (such as 1:3 or 1-2) or a single vector system encoding both the HC and the LC. Clarified media, into which the antibody had been secreted, was purified using any of many commonly-used techniques. For example, the medium may be conveniently applied to a MABSELECT® column (GE Healthcare), or KappaSelect column (GE Healthcare) for Fab fragment, that has been equilibrated with a compatible buffer, such as phosphate buffered saline (pH 7.4).

[1637] The column was washed to remove nonspecific binding components. The bound antibody was eluted, for example, by pH gradient (such as 20 mM Tris buffer, pH 7.0 to 10 mM sodium citrate buffer, pH 3.0, or phosphate buffered saline pH 7.4 to 100 mM glycine buffer, pH 3.0). Antibody fractions were detected, such as by SDS-PAGE, and then were pooled. Further purification was deemed optional, depending on intended use.

[1638] Populations of each antibody were concentrated and/or sterile filtered using common techniques. Soluble

aggregate and multimers are effectively removed by common techniques, including size exclusion, hydrophobic interaction, ion exchange, multimodal, or hydroxyapatite chromatography. The purity of the antibody after these chromatography steps was between about 95% to about 99%. The product was either refrigerated, immediately frozen at -70°C ., or was lyophilized.

[1639] Various physical and expression-related characteristics particular exemplary antibodies were determined using methodologies known in the art (see Tables 2-1 through 2-5 below).

[1640] Antibody encoding plasmid DNA was transfected into Expi293-F™ cells (GIBCO) to produce antibodies using the manufacturer's recommended protocol with some modifications. Expi293-F™ cells were aliquoted into 24 well plates and incubated with shaking at 37°C . with 8% CO_2 and 85% humidity. Heavy and light chain plasmids were pooled at a 1:1 heavy to light chain mass, mixed with Expi293Fectamine, and added dropwise to cells in each well. Expi293F Enhancer I and II were added 20 hr post-transfection and cell supernatants were harvested 4 d post transfection. Antibody titers were measured by biolayer interferometry on an Octet HTX instrument (ForteBio). See Table 2-1. Expression volume was 4 mL.

TABLE 2-1

Expression levels (N = 2) of monoclonal antibodies in expiHEK293 cells.						
Antibody ID	Expression level ($\mu\text{g/mL}$) replica 1	Expression amount (μg) replica 1	Expression level ($\mu\text{g/mL}$) replica 2	Expression amount (μg) replica 2	Average expression level ($\mu\text{g/mL}$) $\pm$ SD	Average expression amount (μg) $\pm$ SD
258	108.8	391.7	135.5	487.8	122.2 $\pm$ 13.3	439.8 $\pm$ 48
261	181.6	653.8	189.4	681.8	185.5 $\pm$ 3.9	667.8 $\pm$ 14
262	86	309.6	104.2	375.1	95.1 $\pm$ 9.1	342.4 $\pm$ 32.8
268	25.9	93.2	24.6	88.6	25.3 $\pm$ 0.6	90.9 $\pm$ 2.3
269	90.1	324.4	105.7	380.5	97.9 $\pm$ 7.8	352.5 $\pm$ 28.1
272	19.1	68.8	19.5	70.2	19.3 $\pm$ 0.2	69.5 $\pm$ 0.7
274	100.4	361.4	104.1	374.8	102.3 $\pm$ 1.8	368.1 $\pm$ 6.7
275	196.3	706.7	223.5	804.6	209.9 $\pm$ 13.6	755.7 $\pm$ 49
277	51	183.6	73.3	263.9	62.2 $\pm$ 11.2	223.8 $\pm$ 40.1
278	90.8	326.9	97	349.2	93.9 $\pm$ 3.1	338.1 $\pm$ 11.2
279	81.5	293.4	90.5	325.8	86 $\pm$ 4.5	309.6 $\pm$ 16.2
280	21.6	77.8	22	79.2	21.8 $\pm$ 0.2	78.5 $\pm$ 0.7
283	57.3	206.3	60.6	218.2	59 $\pm$ 1.7	212.3 $\pm$ 5.9
285	132.4	476.6	145.6	524.2	139 $\pm$ 6.6	500.4 $\pm$ 23.8
286	28	100.8	31.4	113	29.7 $\pm$ 1.7	106.9 $\pm$ 6.1
287	67.2	241.9	68.7	247.3	68 $\pm$ 0.8	244.6 $\pm$ 2.7
291	375.3	1351.1	461.8	1662.5	418.6 $\pm$ 43.3	1506.8 $\pm$ 155.7
292	217.1	781.6	229.7	826.9	223.4 $\pm$ 6.3	804.3 $\pm$ 22.7
296	95.7	344.5	90.1	324.4	92.9 $\pm$ 2.8	334.5 $\pm$ 10.1
297	40	144	39.1	140.8	39.6 $\pm$ 0.4	142.4 $\pm$ 1.6
298	59.1	212.8	78.5	282.6	68.8 $\pm$ 9.7	247.7 $\pm$ 34.9
302	52.2	187.9	58.8	211.7	55.5 $\pm$ 3.3	199.8 $\pm$ 11.9
304	123.1	443.2	127.8	460.1	125.5 $\pm$ 2.4	451.7 $\pm$ 8.5
305	79.2	285.1	91.6	329.8	85.4 $\pm$ 6.2	307.5 $\pm$ 22.4
307	44.8	161.3	46.7	168.1	45.8 $\pm$ 1	164.7 $\pm$ 3.4
308	134.9	485.6	139.4	501.8	137.2 $\pm$ 2.3	493.7 $\pm$ 8.1
309	143.9	518	164.1	590.8	154 $\pm$ 10.1	554.4 $\pm$ 36.4
310	118.4	426.2	113.7	409.3	116.1 $\pm$ 2.4	417.8 $\pm$ 8.4
311	105.7	380.5	120.8	434.9	113.3 $\pm$ 7.6	407.7 $\pm$ 27.2
312	44.6	160.6	53.8	193.7	49.2 $\pm$ 4.6	177.2 $\pm$ 16.6
313	142.2	511.9	200.6	722.2	171.4 $\pm$ 29.2	617.1 $\pm$ 105.2
314	2.6	9.5	2.9	10.3	2.8 $\pm$ 0.2	9.9 $\pm$ 0.4
318	59.8	215.3	61.6	221.8	60.7 $\pm$ 0.9	218.6 $\pm$ 3.3
321	11.9	42.8	12.7	45.7	12.3 $\pm$ 0.4	44.3 $\pm$ 1.5
322	164.6	592.6	176.8	636.5	170.7 $\pm$ 6.1	614.6 $\pm$ 22
325	119	428.4	125.3	451.1	122.2 $\pm$ 3.2	439.8 $\pm$ 11.4
330	139.6	502.6	135.1	486.4	137.4 $\pm$ 2.3	494.5 $\pm$ 8.1
334	36	129.6	38.6	139	37.3 $\pm$ 1.3	134.3 $\pm$ 4.7
335	1.4	5.1	1.4	5.1	1.4 $\pm$ 0	5.1 $\pm$ 0
336	45.8	164.9	49	176.4	47.4 $\pm$ 1.6	170.7 $\pm$ 5.8
338	87.4	314.6	110.9	399.2	99.2 $\pm$ 11.8	356.9 $\pm$ 42.3
341	8	28.7	8.5	30.6	8.3 $\pm$ 0.3	29.7 $\pm$ 1
346	31.8	114.5	33.4	120.2	32.6 $\pm$ 0.8	117.4 $\pm$ 2.9
350	75.4	271.4	99.4	357.8	87.4 $\pm$ 12	314.6 $\pm$ 43.2
352	72.3	260.3	72.5	261	72.4 $\pm$ 0.1	260.7 $\pm$ 0.3
353	144	518.4	130.8	470.9	137.4 $\pm$ 6.6	494.7 $\pm$ 23.8
354	70.4	253.4	67.3	242.3	68.9 $\pm$ 1.6	247.9 $\pm$ 5.6
355	8.8	31.8	8.7	31.3	8.8 $\pm$ 0.1	31.6 $\pm$ 0.3
356	88.9	320	102.8	370.1	95.9 $\pm$ 7	345.1 $\pm$ 25.1
357	193.8	697.7	198.3	713.9	196.1 $\pm$ 2.3	705.8 $\pm$ 8.1
358	136.7	492.1	159	572.4	147.9 $\pm$ 11.2	532.3 $\pm$ 40.2
359	196.5	707.4	201.4	725	199 $\pm$ 2.5	716.2 $\pm$ 8.8
360	126.6	455.8	146.4	527	136.5 $\pm$ 9.9	491.4 $\pm$ 35.6
361	70.6	254.2	72.3	260.3	71.5 $\pm$ 0.9	257.3 $\pm$ 3.1
364	29.1	104.8	31.3	112.7	30.2 $\pm$ 1.1	108.8 $\pm$ 4
368	67	241.2	68.6	247	67.8 $\pm$ 0.8	244.1 $\pm$ 2.9
369	84.9	305.6	99.6	358.6	92.3 $\pm$ 7.3	332.1 $\pm$ 26.5
370	56.3	202.7	71	255.6	63.7 $\pm$ 7.3	229.2 $\pm$ 26.5
371	89.2	321.1	104.5	376.2	96.9 $\pm$ 7.7	348.7 $\pm$ 27.6
373	56.4	203	70.6	254.2	63.5 $\pm$ 7.1	228.6 $\pm$ 25.6
374	111	399.6	125.3	451.1	118.2 $\pm$ 7.2	425.4 $\pm$ 25.8
377	108.1	389.2	133.1	479.2	120.6 $\pm$ 12.5	434.2 $\pm$ 45
381	142	511.2	151.9	546.8	147 $\pm$ 5	529 $\pm$ 17.8
382	44.8	161.3	46.9	168.8	45.9 $\pm$ 1.1	165.1 $\pm$ 3.8
383	49.6	178.6	51.2	184.3	50.4 $\pm$ 0.8	181.5 $\pm$ 2.9
384	33.5	120.6	35.7	128.5	34.6 $\pm$ 1.1	121.6 $\pm$ 4
386	2	7.1	2.2	7.7	2.1 $\pm$ 0.1	7.4 $\pm$ 0.3
387	66.6	239.8	84.9	305.6	75.8 $\pm$ 9.1	272.7 $\pm$ 32.9
388	136.1	490	158	568.8	147.1 $\pm$ 11	529.4 $\pm$ 39.4
389	62.7	225.7	58.7	211.3	60.7 $\pm$ 2	218.5 $\pm$ 7.2

TABLE 2-1-continued

Expression levels (N = 2) of monoclonal antibodies in expiHEK293 cells.						
Antibody ID	Expression level (μg/mL) replica 1	Expression amount (μg) replica 1	Expression level (μg/mL) replica 2	Expression amount (μg) replica 2	Average expression level (μg/mL) ± SD	Average expression amount (μg) ± SD
390	116.8	420.5	112.7	405.7	114.8 ± 2.1	413.1 ± 7.4
391	50.9	183.2	51.6	185.8	51.3 ± 0.4	184.5 ± 1.3
393	202.8	730.1	197.7	711.7	200.3 ± 2.6	720.9 ± 9.2
394	79.5	286.2	80.3	289.1	79.9 ± 0.4	287.7 ± 1.5
397	64.4	231.8	87.1	313.6	75.8 ± 11.4	272.7 ± 40.9
399	131.9	474.8	180.1	648.4	156 ± 24.1	561.6 ± 86.8
400	320.9	1155.2	383.2	1379.5	352.1 ± 31.2	1267.4 ± 112.2
401	19.8	71.3	24.1	86.8	22 ± 2.2	79.1 ± 7.8
402	46.9	168.8	55	198	51 ± 4.1	183.4 ± 14.6
403	173.2	623.5	172.3	620.3	172.8 ± 0.4	621.9 ± 1.6
404	26.6	95.8	29.6	106.6	28.1 ± 1.5	101.2 ± 5.4
408	93.8	337.7	101.6	365.8	97.7 ± 3.9	351.8 ± 14.1
410	118.8	427.7	131.7	474.1	125.3 ± 6.5	450.9 ± 23.2
411	1	3.7	1.2	4.2	1.1 ± 0.1	4 ± 0.3
413	149.9	539.6	134.5	484.2	142.2 ± 7.7	511.9 ± 27.7
414	59.7	214.9	67.8	244.1	63.8 ± 4.1	229.5 ± 14.6
415	192.2	691.9	200.2	720.7	196.2 ± 4	706.3 ± 14.4
417	78.5	282.6	84.3	303.5	81.4 ± 2.9	293.1 ± 10.5
419	75.9	273.2	76.7	276.1	76.3 ± 0.4	274.7 ± 1.5
420	3.1	11.2	3.3	11.8	3.2 ± 0.1	11.5 ± 0.3
426	1.8	6.5	1.9	6.7	1.9 ± 0	6.6 ± 0.1
428	57	205.2	64	230.4	60.5 ± 3.5	217.8 ± 12.6
429	90.7	326.5	100.6	362.2	95.7 ± 5	344.4 ± 17.9
430	19.6	70.6	19.5	70.2	19.6 ± 0.1	70.4 ± 0.2
432	16	57.6	16.6	59.8	16.3 ± 0.3	58.7 ± 1.1
433	383.4	1380.2	426.2	1534.3	404.8 ± 21.4	1457.3 ± 77.1
436	54.7	196.9	63.5	228.6	59.1 ± 4.4	212.8 ± 15.9
438	11.3	40.7	12.1	43.6	11.7 ± 0.4	42.2 ± 1.5
440	74.1	266.8	81.2	292.3	77.7 ± 3.6	279.6 ± 12.8
441	168.6	607	165.3	595.1	167 ± 1.6	601.1 ± 5.9
442	126.1	454	143.3	515.9	134.7 ± 8.6	485 ± 31
444	54.6	196.6	60.5	217.8	57.6 ± 3	207.2 ± 10.6
445	105.7	380.5	115.4	415.4	110.6 ± 4.9	398 ± 17.5
447	113.9	410	126.5	455.4	120.2 ± 6.3	432.7 ± 22.7
448	72.5	261	85.4	307.4	79 ± 6.5	284.2 ± 23.2
449	139	500.4	193.4	696.2	166.2 ± 27.2	598.3 ± 97.9
450	7.1	25.6	7.6	27.4	7.4 ± 0.3	26.5 ± 0.9
451	150.1	540.4	155	558	152.6 ± 2.5	549.2 ± 8.8
453	117.9	424.4	109	392.4	113.5 ± 4.5	408.4 ± 16
455	123.9	446	156.7	564.1	140.3 ± 16.4	505.1 ± 59.1
457	1.1	3.9	1.3	4.5	1.2 ± 0.1	4.2 ± 0.3
458	157.9	568.4	160.7	578.5	159.3 ± 1.4	573.5 ± 5.1
459	68.9	248	87.8	316.1	78.4 ± 9.5	282.1 ± 34.1
460	140.6	506.2	142.6	513.4	141.6 ± 1	509.8 ± 3.6
461	62.6	225.4	81.4	293	72 ± 9.4	259.2 ± 33.8
462	44.2	159.1	53.1	191.2	48.7 ± 4.5	175.2 ± 16.1
463	143.7	517.3	156.9	564.8	150.3 ± 6.6	541.1 ± 23.8
464	115	414	131.6	473.8	123.3 ± 8.3	443.9 ± 29.9
465	105	378	146	525.6	125.5 ± 20.5	451.8 ± 73.8
466	1.5	5.3	1.7	6.1	1.6 ± 0.1	5.7 ± 0.4
467	142.1	511.6	166.2	598.3	154.2 ± 12.1	555 ± 43.4
468	169.2	609.1	160.8	578.9	165 ± 4.2	594 ± 15.1
469	49.2	177.1	54.2	195.1	51.7 ± 2.5	186.1 ± 9
472	107.1	385.6	106.1	382	106.6 ± 0.5	383.8 ± 1.8
473	203.9	734	237.7	855.7	220.8 ± 16.9	794.9 ± 60.9
478	325.5	1171.8	291.4	1049	308.5 ± 17.1	1110.4 ± 61.4
479	93.3	335.9	101.6	365.8	97.5 ± 4.2	350.9 ± 15
481	58.2	209.5	73.3	263.9	65.8 ± 7.5	236.7 ± 27.2
482	33	118.8	32.7	117.7	32.9 ± 0.1	118.3 ± 0.5
483	81.1	292	76.3	274.7	78.7 ± 2.4	283.4 ± 8.7
484	28.8	103.7	27.4	98.6	2.1 ± 0.7	101.2 ± 2.6
486	246.9	888.8	292.5	1053	269.7 ± 22.8	970.9 ± 82.1
488	53.3	191.9	53.2	191.5	53.3 ± 0	191.7 ± 0.2
489	91.4	329	105.6	380.2	98.5 ± 7.1	354.6 ± 25.6
490	0	0	0.4	1.3	0.2 ± 0.2	0.7 ± 0.7
491	272.9	982.4	284.7	1024.9	278.8 ± 5.9	1003.7 ± 21.3
493	97.9	352.4	135.5	487.8	116.7 ± 18.8	420.1 ± 67.7
494	22.8	82.1	20.3	73.1	21.6 ± 1.3	77.6 ± 4.5
495	85.2	306.7	123.8	445.7	104.5 ± 19.3	376.2 ± 69.5
499	95.2	342.7	111	399.6	103.1 ± 7.9	371.2 ± 28.5

TABLE 2-1-continued

Expression levels (N = 2) of monoclonal antibodies in expiHEK293 cells.						
Antibody ID	Expression level (μg/mL) replica 1	Expression amount (μg) replica 1	Expression level (μg/mL) replica 2	Expression amount (μg) replica 2	Average expression level (μg/mL) ± SD	Average expression amount (μg) ± SD
500	65.8	236.9	72.6	261.4	69.2 ± 3.4	249.2 ± 12.3
501	122.4	440.6	170.5	613.8	146.5 ± 24.1	527.2 ± 86.6
502	38.7	139.3	47.5	171	43.1 ± 4.4	155.2 ± 15.9
503	67.7	243.7	61.4	221	64.6 ± 3.2	232.4 ± 11.4
506	134.7	484.9	143.8	517.7	139.3 ± 4.6	501.3 ± 16.4
508	53.1	191.2	55.8	200.9	54.5 ± 1.4	196.1 ± 4.9
511	65.2	234.7	66	237.6	65.6 ± 0.4	236.2 ± 1.5
513	217.7	783.7	245.4	883.4	231.6 ± 13.9	833.6 ± 49.9
515	56.9	204.8	60.7	218.5	58.8 ± 1.9	211.7 ± 6.8
518	95.3	343.1	110.3	397.1	102.8 ± 7.5	370.1 ± 27
519	288.1	1037.2	247.5	891	267.8 ± 20.3	964.1 ± 73.1
520	89.7	322.9	133.3	479.9	111.5 ± 21.8	401.4 ± 78.5
521	64.9	233.6	61.4	221	63.2 ± 1.8	227.3 ± 6.3
524	140.4	505.4	206.4	743	173.4 ± 33	624.2 ± 118.8
525	137.7	495.7	127.5	459	132.6 ± 5.1	477.4 ± 18.4
526	83.9	302	78.1	281.2	81 ± 2.9	291.6 ± 10.4
528	84.5	304.2	85.8	308.9	85.2 ± 0.6	306.6 ± 2.3
529	2.4	8.8	2.6	9.2	2.5 ± 0.1	9 ± 0.2
530	69.8	251.3	70.3	253.1	70.1 ± 0.3	257.7 ± 0.9
532	23.5	84.6	25.1	90.4	24.3 ± 0.8	87.5 ± 2.9
534	115.3	415.1	113.8	409.7	114.6 ± 0.8	412.4 ± 2.7
535	154.9	557.6	170.1	612.4	162.5 ± 7.6	585 ± 27.4
539	104.5	376.2	98.7	355.3	101.6 ± 2.9	365.8 ± 10.5
540	61	219.6	71.9	258.8	66.5 ± 5.5	239.2 ± 19.6
549	35.1	126.4	36.4	131	35.8 ± 0.6	128.7 ± 2.3
552	113.8	409.7	113.3	407.9	113.6 ± 0.3	408.8 ± 0.9
553	3.9	14.2	3.8	13.7	3.9 ± 0.1	14 ± 0.3
554	163.7	589.3	179.8	647.3	171.8 ± 8.1	618.3 ± 29
555	21.6	77.8	20.9	75.2	21.3 ± 0.4	76.5 ± 1.3
557	336.9	1212.8	358.4	1290.2	347.7 ± 10.8	1251.5 ± 38.7
558	397.9	1432.4	397.9	1432.4	397.9 ± 0	1432.4 ± 0
559	226.6	815.8	236.7	852.1	231.7 ± 5.1	834 ± 18.2
561	262.2	943.9	275	990	268.6 ± 6.4	967 ± 23.1
562	163.6	589	166.4	599	165 ± 1.4	594 ± 5
565	46.8	168.5	46.5	167.4	46.7 ± 0.1	168 ± 0.5
567	92.7	333.7	108.2	389.5	100.5 ± 7.8	361.6 ± 27.9
568	53.6	193	63.6	229	58.6 ± 5	211 ± 18
570	99.7	358.9	101	363.6	100.4 ± 0.6	361.3 ± 2.4
572	1.4	4.9	1.6	5.6	1.5 ± 0.1	5.3 ± 0.4
573	39.4	141.8	40.9	147.2	40.2 ± 0.8	144.5 ± 2.7
574	149.8	539.3	189.3	681.5	169.6 ± 19.8	610.4 ± 71.1
575	98.9	356	104.8	377.3	101.9 ± 3	366.7 ± 10.7
577	65.6	236.2	59.8	215.3	62.7 ± 2.9	225.8 ± 10.5
579	29.5	106.2	29.9	107.6	29.7 ± 0.2	106.9 ± 0.7
584	57.9	208.4	62	223.2	60 ± 2.1	215.8 ± 7.4
585	99.2	357.1	111	399.6	105.1 ± 5.9	378.4 ± 21.3
587	19.6	70.6	19.3	69.5	19.5 ± 0.2	70.1 ± 0.5

[1641] Antibody encoding plasmid DNA was transfected into Expi293-FTM cells (GIBCO) For protein purification, 8 μL of a 25% slurry solution of magnetic protein A beads (GenScript) were added to 30 μg of expressed mAb and incubated overnight with shaking at 37° C. The next day, beads were washed and protein eluted with 480 μL of 100 mM glycine pH 2.0. Elution fractions were neutralized to pH 7.0 and dried onto filter plates. The dried mAb was resuspended in buffer to a final concentration of approximately 1

mg/mL and sterile filtered. Purified mAbs were quantified by bio-layer interferometry using an Octet HTX instrument (forteBIO). Purity of mAbs was analyzed by denaturing capillary electrophoresis (CE) SDS page using a LabChip instrument (Perkin Elmer) according to the manufacturers protocol. The data were analyzed using the LabChip GX Reviewer Software (Perkin Elmer). See Tables 2-2 and 2-3. No value available where empty.

TABLE 2-2

Purification yields (N = 2) after being buffer exchanged into phosphate buffered saline.						
Antibody ID	Concentration (µg/mL) replica 1	Amount (µg) replica 1	Concentration (µg/mL) replica 2	Amount (µg) replica 2	Average concentration (µg/mL) ± SD	Average amount (µg) ± SD
258	1852	253	1819	236	1836 ± 17	245 ± 9
261	2412	330	3585	465	2999 ± 587	397 ± 68
262	1057	142	1593	204	1325 ± 268	173 ± 31
268	264	36	134	17	199 ± 65	27 ± 10
269	1516	204	1009	129	1263 ± 254	166 ± 37
272	314	42	318	41	316 ± 2	42 ± 1
274	1195	160	2029	260	1612 ± 417	210 ± 50
275	2589	354	4231	549	3410 ± 821	451 ± 98
277	422	58	1286	167	854 ± 432	112 ± 55
278	1479	202	1311	170	1395 ± 84	186 ± 16
279	1277	171	1381	177	1329 ± 52	174 ± 3
280	281	38	305	40	293 ± 12	39 ± 1
283	996	134	1271	163	1133 ± 138	148 ± 15
285	2210	302	2041	265	2125 ± 84	284 ± 19
286	445	61	99	13	272 ± 173	37 ± 24
287	1127	154	583	76	855 ± 272	115 ± 39
291	4277	574	4559	585	4418 ± 141	580 ± 5
292	2589	354	3552	461	3070 ± 482	407 ± 54
296	1638	220	1124	144	1381 ± 257	182 ± 38
297	544	74	690	89	617 ± 73	82 ± 7
298	918	123	1451	186	1185 ± 266	155 ± 31
302	1066	146	843	109	954 ± 111	127 ± 18
304	1908	261	1933	251	1920 ± 13	256 ± 5
305	632	85	1254	161	943 ± 311	123 ± 38
307	614	84	730	95	672 ± 58	89 ± 6
308	2080	279	1880	241	1980 ± 100	260 ± 19
309	2325	318	2893	375	2609 ± 284	346 ± 29
310	2406	323	1763	226	2084 ± 321	274 ± 48
311	1809	243	1812	232	1810 ± 2	237 ± 5
312	442	59	726	93	584 ± 142	76 ± 17
313	2443	328	3228	414	2835 ± 393	371 ± 43
314	8	1	8	1	8 ± 0	1 ± 0
318	1059	142	816	105	938 ± 122	124 ± 19
321	172	23	153	20	162 ± 9	22 ± 2
322	2401	322	2841	364	2621 ± 220	343 ± 21
325	2164	291	2018	259	2091 ± 73	275 ± 16
330	1920	258	2261	290	2091 ± 170	274 ± 16
334	621	83	650	83	636 ± 14	83 ± 0
335	0	0	0	0	0 ± 0	0 ± 0
336	602	81	862	111	732 ± 130	96 ± 15
338	982	132	1423	182	1202 ± 221	157 ± 25
341	39	5	17	2	28 ± 11	4 ± 2
346	554	76	706	27	380 ± 174	51 ± 24
350	1370	184	1332	171	1351 ± 19	178 ± 7
352	824	111	1261	162	1043 ± 218	136 ± 26
353	1885	258	2166	281	2026 ± 140	269 ± 12
354	1052	144	889	115	971 ± 82	129 ± 14
355	118	16	125	16	122 ± 3	16 ± 0
356	944	129	1292	168	1118 ± 174	149 ± 20
357	3560	478	3451	443	3506 ± 55	461 ± 18
358	7356	322	2867	372	2611 ± 256	347 ± 25
359	2861	391	3390	440	36 ± 264	416 ± 25
360	1637	224	2625	341	2131 ± 494	282 ± 59
361	1113	149	1057	136	1085 ± 28	143 ± 7
364	86	12	213	28	149 ± 64	20 ± 8
368	1273	171	1018	131	1145 ± 127	151 ± 20
369	985	135	1428	185	1207 ± 222	160 ± 25
370	923	124	942	121	932 ± 10	122 ± 1
371	1058	142	1780	228	1419 ± 361	185 ± 43
373	938	128	778	101	858 ± 80	115 ± 14
374	1926	763	2356	306	2141 ± 215	285 ± 21
377	1708	229	2337	300	2023 ± 314	265 ± 35
381	1755	240	2199	285	1977 ± 222	262 ± 23
382	759	104	862	112	811 ± 51	108 ± 4
383	885	121	914	119	900 ± 14	120 ± 1
384	433	59	506	66	469 ± 37	63 ± 3
386	0	0	0	0	0 ± 0	0 ± 0
387	862	116	1395	179	1128 ± 267	147 ± 32
388	1867	255	2929	380	2398 ± 531	318 ± 62
389	1149	154	1021	131	1085 ± 64	143 ± 12

TABLE 2-2-continued

Purification yields (N = 2) after being buffer exchanged into phosphate buffered saline.						
Antibody ID	Concentration (µg/mL) replica 1	Amount (µg) replica 1	Concentration (µg/mL) replica 2	Amount (µg) replica 2	Average concentration (µg/mL) ± SD	Average amount (µg) ± SD
390	2065	277	7067	265	2066 ± 1	271 ± 6
391	1020	137	1000	128	1010 ± 10	132 ± 4
393	2614	357	2548	331	2581 ± 33	344 ± 13
394	1632	219	1340	172	1486 ± 146	196 ± 74
397	1035	139	1477	189	1256 ± 221	164 ± 25
399	2668	358	2534	325	260 ± 67	342 ± 17
400	4241	569	5563	713	4902 ± 661	641 ± 72
401	209	28	280	36	244 ± 36	32 ± 4
402	879	120	773	100	826 ± 53	110 ± 10
403	1939	265	2308	299	2124 ± 184	282 ± 17
404	363	49	448	57	406 ± 42	53 ± 4
408	896	123	809	105	853 ± 44	114 ± 9
410	1866	255	2162	281	2014 ± 148	268 ± 13
411	0	0	0	0	0 ± 0	0 ± 0
413	2729	366	2010	258	2370 ± 360	312 ± 54
414	394	54	475	62	434 ± 41	58 ± 4
415	2606	356	2926	380	2766 ± 160	368 ± 12
417	1464	200	1247	162	1356 ± 109	181 ± 19
419	1625	218	1316	169	1471 ± 155	194 ± 25
420	17	2	19	2	18 ± 1	2 ± 0
426	0	0	0	0	0 ± 0	0 ± 0
428	1064	146	871	113	968 ± 97	129 ± 16
429	1306	178	1443	187	1374 ± 69	183 ± 4
430	205	28	240	31	222 ± 18	29 ± 2
432	256	35	222	29	239 ± 17	32 ± 3
433	4217	566	2740	351	3478 ± 738	459 ± 108
436	1117	153	962	125	1039 ± 77	139 ± 14
438	148	70	163	21	156 ± 7	21 ± 0
440	1715	230	1192	153	1453 ± 261	192 ± 39
441	1637	220	2282	293	1959 ± 323	256 ± 37
442	2331	319	2230	289	2280 ± 50	304 ± 15
444	763	102	974	125	868 ± 106	114 ± 11
445	1892	254	1630	209	1761 ± 131	231 ± 22
447	1599	219	1885	245	1742 ± 143	232 ± 13
448	1326	178	1139	146	1233 ± 94	162 ± 16
449	2391	321	3216	412	2804 ± 412	367 ± 46
450	80	11	66	0	73 ± 7	10 ± 1
451	2382	320	2657	341	2520 ± 137	330 ± 11
453	1683	230	1713	222	1698 ± 15	226 ± 4
455	2229	299	2560	328	2394 ± 166	314 ± 14
457	0	0	0	0	0 ± 0	0 ± 0
458	2698	362	2736	351	2717 ± 19	357 ± 6
459	1311	179	1587	206	1449 ± 138	193 ± 13
460	2155	289	1854	238	2005 ± 151	264 ± 26
461	1164	156	1515	194	1340 ± 175	175 ± 19
462	700	96	795	103	747 ± 48	99 ± 4
463	2173	297	1963	255	2068 ± 105	276 ± 21
464	1958	263	1513	194	1735 ± 222	228 ± 34
465	1849	248	2070	266	1959 ± 111	257 ± 9
466	0	0	0	0	0 ± 0	0 ± 0
467	450	60	483	62	466 ± 17	61 ± 1
468	2384	320	2152	276	2268 ± 116	298 ± 22
469	935	126	1025	131	980 ± 45	128 ± 3
472	1557	209	1846	237	1702 ± 144	223 ± 14
473	1327	181	1623	211	1475 ± 148	196 ± 15
478	2841	388	4098	532	3469 ± 629	460 ± 72
479	1488	203	1657	215	1573 ± 84	209 ± 6
481	958	131	1179	153	1068 ± 111	142 ± 11
482	436	59	456	59	446 ± 10	59 ± 0
483	1559	209	895	115	1227 ± 332	162 ± 47
484	411	56	389	51	400 ± 11	54 ± 3
486	2997	410	4063	527	3530 ± 533	468 ± 59
488	894	122	840	109	867 ± 27	116 ± 7
489	1241	170	1209	157	1225 ± 16	163 ± 6
490	0	0	0	0	0 ± 0	0 ± 0
491	2699	369	3986	517	3343 ± 643	443 ± 74
493	1645	221	2159	277	1902 ± 257	249 ± 28
494	252	34	305	39	279 ± 27	36 ± 3
495	1700	228	1736	223	1718 ± 18	226 ± 3
499	1584	213	1930	247	1757 ± 173	230 ± 17

TABLE 2-2-continued

Purification yields (N = 2) after being buffer exchanged into phosphate buffered saline.						
Antibody ID	Concentration (µg/mL) replica 1	Amount (µg) replica 1	Concentration (µg/mL) replica 2	Amount (µg) replica 2	Average concentration (µg/mL) ± SD	Average amount (µg) ± SD
500	947	129	1089	141	1018 ± 71	135 ± 6
501	2219	298	2206	283	2212 ± 6	290 ± 7
502	360	49	666	86	513 ± 153	68 ± 18
503	1280	172	502	64	891 ± 389	118 ± 54
506	2079	279	1848	237	1963 ± 115	258 ± 21
508	954	130	580	75	767 ± 187	103 ± 28
511	1290	173	901	116	1095 ± 194	145 ± 29
513	3893	523	4089	524	3991 ± 98	523 ± 1
515	860	116	1082	139	971 ± 111	127 ± 12
518	966	132	1687	219	1327 ± 360	176 ± 43
519	2527	345	3749	486	338 ± 611	416 ± 0
520	1263	170	2202	282	1733 ± 469	226 ± 56
521	1186	159	898	115	1042 ± 144	137 ± 22
524	2218	298	2596	333	2407 ± 189	315 ± 18
525	2211	297	2073	266	2142 ± 69	281 ± 15
526	1410	193	988	128	1199 ± 211	160 ± 32
528	1357	185	1189	154	1273 ± 84	170 ± 16
529	17	2	13	2	15 ± 2	2 ± 0
530	1092	149	1111	144	1101 ± 10	147 ± 3
532	259	35	247	32	253 ± 6	34 ± 2
534	1515	207	1284	167	1399 ± 115	187 ± 20
535	2033	273	1559	200	1796 ± 237	236 ± 36
539	1762	237	1465	188	1613 ± 148	212 ± 24
540	973	131	1214	156	1093 ± 121	143 ± 13
549	578	79	493	64	535 ± 42	71 ± 7
552	2264	304	1833	235	2048 ± 215	269 ± 34
553	37	5	22	3	29 ± 7	4 ± 1
554	1990	272	2118	275	2054 ± 64	273 ± 2
555	140	19	122	16	131 ± 9	18 ± 2
557	3462	473	4758	617	4110 ± 648	545 ± 72
558	5285	710	5164	662	5225 ± 61	686 ± 24
559	2984	401	3322	426	3153 ± 169	413 ± 13
561	2440	334	3954	513	3197 ± 757	423 ± 90
562	2714	364	2464	316	2589 ± 125	340 ± 24
565	804	108	595	76	699 ± 104	92 ± 16
567	636	85	1867	239	1252 ± 615	162 ± 77
568	953	128	968	124	961 ± 7	126 ± 2
570	1536	206	1337	171	1436 ± 99	189 ± 18
572	0	0	0	0	0 ± 0	0 ± 0
573	0	0	0	0	0 ± 0	0 ± 0
574	2233	300	3209	412	2721 ± 488	356 ± 56
575	897	120	1435	184	1166 ± 269	152 ± 32
577	1185	159	670	86	927 ± 257	123 ± 37
579	55	35	239	31	247 ± 8	33 ± 2
584	996	136	792	103	894 ± 102	120 ± 17
585	1503	202	2011	258	1757 ± 254	230 ± 28
587	168	23	138	18	153 ± 15	21 ± 3

TABLE 2-3

Purity (%) and apparent molecular weight (MW) (N = 1) of monoclonal antibodies using capillary electrophoresis (CE) SDS PAGE.		
Antibody ID	Purity (%)	Apparent MW (kDa)
258	100	191
261	86	182
262		
268		
269	100	200
272		
274		
275	97	175
277		
278	95	195
279	100	197
280		

TABLE 2-3-continued

Purity (%) and apparent molecular weight (MW) (N = 1) of monoclonal antibodies using capillary electrophoresis (CE) SDS PAGE.		
Antibody ID	Purity (%)	Apparent MW (kDa)
283	90	191
285	100	192
286		
287	100	196
291	100	197
292	100	185
296		
297		
298	96	187
302	98	180
304	100	177
305	100	192

TABLE 2-3-continued

Purity (%) and apparent molecular weight (MW) (N = 1) of monoclonal antibodies using capillary electrophoresis (CE) SDS PAGE.		
Antibody ID	Purity (%)	Apparent MW (kDa)
307		
308	93	196
309	94	174
310	90	193
311	95	180
312		
313	91	182
314		
318	100	197
321		
322	100	182
325	91	178
330	100	186
334		
335		
336		
338	93	191
341		
346		
350	90	193
352	92	199
353	100	174
354	99	176
355		
356	100	175
357	94	190
358	100	173
359	100	177
360	100	171
361	96	199
364		
368	93	184
369	100	176
370	100	197
371	100	198
373	100	179
374	100	181
377		
381	100	179
382	100	175
383	100	171
384		
386		
387	100	194
388	100	175
389	100	194
398	100	191
391	100	193
393	100	178
394	100	195
397	100	195
399	95	195
400		
401		
402	100	175
403	94	171
404		
408	100	180
410	100	172
411		
413	100	194
414		
415	100	173
417	100	170
419	100	194
420		
426		
428	100	177
429	100	183
430		
432		
433	100	184

TABLE 2-3-continued

Purity (%) and apparent molecular weight (MW) (N = 1) of monoclonal antibodies using capillary electrophoresis (CE) SDS PAGE.		
Antibody ID	Purity (%)	Apparent MW (kDa)
436	100	191
438		
440	90	186
441	100	189
442	91	182
444		
445	97	198
447	100	179
448	97	193
449	100	199
450		
451	100	197
453	100	189
455	92	194
457		
458	100	192
459		
460	100	194
461	90	198
462	100	189
463	100	188
464	91	192
465	94	188
466		
467		
468	100	194
469	100	196
472	100	191
473	100	191
478	100	185
479	100	182
481	100	184
482		
483		
484		
486	100	177
488	100	182
489	100	187
490		
491	100	175
493	100	194
494		
495	95	191
499	100	188
500	100	188
501	95	198
502		
503		
506	94	195
508	98	179
511		
513	100	199
515	88	195
518	100	180
519	100	183
520	86	195
521		
524	92	192
525	84	192
526	100	186
528	100	185
529		
530	100	184
532		
534	98	195
535	90	193
539	100	200
540	100	182
549		
552	100	196
553		
554	100	190

TABLE 2-3-continued

Purity (%) and apparent molecular weight (MW) (N = 1) of monoclonal antibodies using capillary electrophoresis (CE) SDS PAGE.		
Antibody ID	Purity (%)	Apparent MW (kDa)
555		
557	100	190
558	100	196
559	93	188
561	100	189
562	100	196
565		
567	98	178
568	100	181
570	91	197
572		
573		
574	100	182
575	100	179
577		
579		
584	100	189
585	100	189
587		

[1642] The melting temperature (T_m) of mAbs was assessed by differential scanning fluorimetry (DSF) using a Thermal Cycler instrument (Bio-Rad)). 6 μ L of mAb solution at 350 μ g/mL was mixed with 6 μ L of a 19 $\times$ concentrated SYPROTM Orange solution (Thermo Fisher). Thermal unfolding as assessed by a change in fluorescence was measured at a starting temperature of 25° C. and increased to 95° C. in 0.5° C./min increments. Data was analyzed and melting curves integrated using the Bio-Rad CFX Maestro software. The T_m was defined as the local minimum taken from the derivative of the melting curve. See Table 2-4. No value available where empty.

TABLE 2-4

Melting point (T _m) (N = 1) of monoclonal antibodies in the differential scanning fluorimetry (DSF) assay.	
Antibody ID	T _m (° C.)
258	67.5
261	67
262	67
268	
269	67
272	
274	
275	67
277	
278	67.5
279	67
280	
283	66
285	67
286	
287	67
291	67
292	62.5
296	65.5
297	
298	66
302	67
304	65.5
305	67.5
387	
308	67.5
389	66.5

TABLE 2-4-continued

Melting point (T _m) (N = 1) of monoclonal antibodies in the differential scanning fluorimetry (DSF) assay.	
Antibody ID	T _m (° C.)
318	66.5
311	66.5
312	
313	67
314	
318	61.5
321	
322	66.5
325	66.5
330	66.5
334	
335	
336	
338	66.5
341	
346	
350	67
352	64.5
353	66.5
354	67
355	
356	67
357	67
358	67
359	66.5
368	66.5
361	67
364	
368	67
369	67
378	63
371	67
373	67
374	67
377	64
381	67
382	66
383	66
384	
386	
387	66.5
388	67
389	66
390	65.5
391	64
393	67
394	66.5
397	66
399	66.5
400	
401	
402	67.5
403	68
404	
408	67
418	67
411	
413	64.5
414	
415	67
417	67
419	66.5
420	
426	
428	67
429	67
430	
432	
433	66.5
436	67.5
438	
440	60

TABLE 2-4-continued

Melting point (Tm) (N = 1) of monoclonal antibodies in the differential scanning fluorimetry (DSF) assay.	
Antibody ID	Tm (° C.)
441	59.5
442	67
444	
445	67
447	67
448	63.5
449	67
450	
451	65.5
453	66
455	65.5
457	
458	
459	67
460	65
461	66.5
462	65.5
463	67
464	66
465	66.5
466	
467	
468	67
469	66
472	62
473	67.5
478	67
479	67
481	61
482	
483	66
484	
486	67
488	67
489	67
498	
491	67
493	66.5
494	
495	62.5
499	66
580	67
501	65.5
502	
503	
586	67
508	67
511	
513	67
515	66
518	67
519	67
520	65
521	
524	67
525	67
526	67
528	67
529	
538	67
532	
534	67
535	65.5
539	67
540	66.5
549	
552	65.5
553	
554	67.5
555	
557	67.5
558	67

TABLE 2-4-continued

Melting point (Tm) (N = 1) of monoclonal antibodies in the differential scanning fluorimetry (DSF) assay.	
Antibody ID	Tm (° C.)
559	67
561	67
562	66.5
565	
567	67
568	67.5
578	66.5
572	
573	
574	67
575	67
577	
579	
584	67
585	63.5
587	

[1643] Percent aggregation and polydispersity of mAbs was assessed by dynamic light scattering (DLS) on a DYNAPRO® Plate Reader III instrument (Wyatt). DLS of individual samples (7 µL of mAb at varying concentrations (0.5-2 mg/mL)) was acquired at 20° C. with 5×5 s acquisitions per sample. Data was analyzed in the Dynamics software (Wyatt) using the regularization algorithm. Percent polydispersity and percent mass of soluble mAbs were calculated for the size range of 2-10 nm. See Table 2-5. No value available where empty.

TABLE 2-5

Amount of soluble antibody (%) and poly dispersity of soluble antibody (%) determined by dynamic light scattering (DLS).		
Antibody ID	Amount soluble antibody (%)	Polydispersity of soluble antibody (%)
258	99	24.7
261	96.3	12.6
262		
268		
269	99.3	36.2
272		
274	96.2	36.3
275	100	10.4
277		
278	57.5	11.1
279	97.7	41.9
280	100	18.3
283	97.4	18.7
285	86.1	18.3
286		
287	96.4	14.7
291	99.5	14.6
292	99.7	24.4
296		
297	99.9	30.9
298	97.7	24.2
302	99.9	6.6
304		
305	96.2	23.1
307		
308	91.2	13.4
309	100	24
310	93.2	12.4
311	93.8	35.2
312		
313	94.9	30.1
314		

TABLE 2-5-continued

Amount of soluble antibody (%) and poly dispersity of soluble antibody (%) determined by dynamic light scattering (DLS).		
Antibody ID	Amount soluble antibody (%)	Polydispersity of soluble antibody (%)
318		
321		
322		
325	98.2	37.6
330	93.3	24
334		
335		
336		
338	99	17.6
341		
346		
350	99.9	23.3
352	95.6	23.3
353	100	15
354	99.9	8.7
355		
356	100	18.3
357	96.6	11.1
358	98.8	28.1
359	100	13.9
368	100	17.6
361		
364		
368		
369	100	16.3
378		
371	96.1	38.6
373	99.8	16.2
374	98.1	23.1
377	97.2	39.7
381	99.3	18.2
382	100	12.5
383		
384		
386		
387	99.2	15
388	99.8	15.6
389		
390	99.9	17.1
391		
393	100	16.6
394	99.5	23.6
397	100	39.2
399	99.7	30
480	85.2	33.1
401		
402	99.5	5.8
483	99.2	23.1
404		
408	100	18.1
418	99.9	28.4
411		
413	98.5	27.3
414		
415	100	30.9
417	97.5	22.4
419	100	20.9
420		
426		
428	98.7	27.7
429	100	15.6
430		
432		
433	99.7	32.7
436	99.5	25.7
438		
440		
441	100	23.5
442	99.8	9.6
444		
445	98.3	36

TABLE 2-5-continued

Amount of soluble antibody (%) and poly dispersity of soluble antibody (%) determined by dynamic light scattering (DLS).		
Antibody ID	Amount soluble antibody (%)	Polydispersity of soluble antibody (%)
447	99.7	10.4
448		
449	97.9	15
450		
451	92.5	39.8
453	100	24.4
455	87.2	16.8
457		
458	99.9	22.8
459	100	28.6
460	97.9	25.3
461	98.5	20.6
462	100	18.7
463	100	7
464	99.8	18.8
465	98.3	35
466		
467		
468	100	48.3
469		
472	99.9	32.7
473	99.9	5.3
478	100	20.5
479		
481	76.3	19.1
482		
483		
484		
486	100	6.6
488	68.5	25.5
489		
498		
491	99.1	30.3
493	99.8	11.6
494		
495	99.7	23.7
499	99.2	24.2
580	100	10.6
501	99.1	29.2
502		
503		
506	97.8	21.4
508	59.8	11.1
511		
513	97.7	21.7
515		
518	100	16.7
519	99.4	11.4
520	99.9	17.7
521		
524	95.8	31
525	100	10.8
526	96.7	39.8
528	94	36.1
529		
530	100	22.1
532		
534	96	45.2
535	100	19.6
539	98.6	39.7
548	98.1	30.9
549		
552	100	28.5
553		
554	100	18.8
555		
557	100	10.9
558	97.8	16.2
559	95	21.4
561	100	11.2
562	99	18.7

TABLE 2-5-continued

Amount of soluble antibody (%) and poly dispersity of soluble antibody (%) determined by dynamic light scattering (DLS).		
Antibody ID	Amount soluble antibody (%)	Polydispersity of soluble antibody (%)
565		
567	92.1	32.7
568	98.2	39.4
570	99.6	34.3
572		
573		
574	99.7	17.5
575	96.5	24.2
577		
579		
584		
585	99.9	18.6
587		

Example 3

[1644] Antibody Characterization Using High-Throughput Surface Plasmon Resonance

[1645] High-throughput surface plasmon resonance capture kinetic experiments were performed on an CATERRA® LSA™ instrument equipped with an HC-30M chip type. Purified monoclonal antibodies obtained from HEK cells were immobilized on a chip by direct coupling. The chip surface was activated and mAbs diluted to either 10 ug/mL or 1 µg/mL were printed onto the chip surface for 10 min. The chip surface was quenched, followed by washing. Relevant benchmarks and negative control mAbs were also printed on the chip surface.

[1646] A 3-fold dilution series of the antigen of interest starting at 300 nM is sequentially injected onto the chip surface. For each concentration, the Ag was injected for 5 min (association phase), followed by running buffer injection for 15 min (dissociation phase). Regeneration cycles were performed between each dilution series. The data were analyzed using the CATERRA® Kinetics analysis software and fit globally to a 1:1 Langmuir binding model to determine apparent association (k_a) and dissociation (k_d) kinetic rate constants and binding affinity constants (KD). See Table 3-1.

[1647] Epitope binning experiments for mAbs coupled to chip were performed using the CATERRA® LSA instrument. Samples were prepared by mixing each mAb in 10-20-fold molar excess with the Ag (1:1 freshly prepared mix of 100 ug/mL mAb and 40 nM Ag). Each Ag-mAb premix was injected sequentially over the chip surface for 4 min (association phase to ligand previously printed onto chip), followed by a running buffer injection for 2 min (dissociation phase). Regeneration cycles were performed between samples. An Ag-only injection (20 nM concentration) was performed as a reference. The data were analyzed using the CATERRA® Epitope analysis software for heat map generation (see FIG. 1).

[1648] Antibody blocking competition experiments were performed to test binding of antibodies to specific domains of SARS-CoV-2 spike protein. Benchmark antibodies with known binding domains were used in competition experiments with identical set up described above. Binding to a specific spike protein domain was determined by analyzing data for antibodies that competed with established bench-

marks. Competition with a known antibody benchmark indicated binding to the same domain. See Table 3-2). C indicates competition, C* indicates symmetric competition (competition observed regardless of which of the two antibodies was bound to the chip and which was soluble), NC indicates no competition, and blanks indicate no available data.

TABLE 3-1

Binding Kinetics			
Antibody ID	k_a (M ⁻¹ s ⁻¹)	k_d (s ⁻¹)	K_D (M)
258	8.40E+04	8.36E-05	9.96E-10
261	1.29E+05	6.95E-04	5.37E-09
269	6.09E+04	4.44E-04	7.29E-09
274	3.01E+04	8.54E-04	2.84E-08
275	1.01E+05	4.80E-05	4.73E-10
277	3.05E+05	6.24E-05	2.04E-10
278	2.01E+05	8.62E-04	4.78E-09
279	3.73E+04	1.95E-03	5.23E-08
283	1.28E+05	2.69E-05	2.11E-10
285	1.57E+05	2.74E-04	1.75E-09
291	4.64E+05	5.43E-05	1.17E-10
297	7.98E+04	1.27E-04	1.60E-09
298	5.97E+05	1.34E-05	2.25E-11
302	1.05E+05	7.63E-05	7.26E-10
304	8.73E+04	2.74E-04	3.14E-09
305	7.26E+04	1.00E-03	1.38E-08
307	1.01E-05	1.71E-03	1.70E-08
308	1.39E+05	6.89E-05	4.95E-10
309	5.96E+05	2.73E-05	4.58E-11
310	1.21E+05	6.43E-05	5.32E-10
311	2.13E+04	4.07E-04	1.91E-08
313	5.02E+05	4.89E-05	9.75E-11
318	1.52E+05	1.70E-03	1.11E-08
322	1.90E+04	4.01E-04	2.11E-08
325	2.09E+04	1.54E-04	7.36E-09
330	7.65E+04	7.30E-04	9.54E-09
336	1.90E+05	2.10E-04	1.11E-09
338	5.02E+04	4.43E-05	8.82E-10
350	2.45E+05	1.08E-04	4.42E-10
352	3.69E+04	5.00E-04	1.35E-08
353	6.80E+05	4.09E-05	6.01E-11
354	4.11E+04	1.79E-03	4.36E-08
356	2.65E+04	9.66E-05	3.65E-09
357	7.80E+04	6.53E-05	8.37E-10
358	2.89E+05	4.87E-04	1.69E-09
359	1.51E+05	6.67E-05	4.41E-10
360	1.04E+05	1.45E-04	1.40E-09
361	3.80E+04	2.92E-05	7.69E-10
368	3.81E+04	4.86E-04	1.27E-08
369	3.16E+05	6.36E-05	2.01E-10
370	1.78E+05	9.86E-04	5.55E-09
371	1.72E+05	6.29E-05	3.66E-10
373	5.12E+04	6.85E-04	1.34E-08
374	4.96E+04	7.86E-05	1.58E-09
381	3.11E+04	2.07E-04	6.66E-09
382	1.20E+05	1.34E-03	1.12E-08
383	9.18E+04	7.09E-05	7.72E-10
387	2.16E+04	8.08E-04	3.74E-08
388	6.10E+05	1.39E-05	2.28E-11
389	4.50E+05	1.00E-05	2.22E-11
390	1.29E+05	1.97E-05	1.53E-10
391	2.70E+04	1.00E-05	3.70E-10
393	9.41E+04	2.95E-05	3.13E-10
394	2.66E+04	3.52E-05	1.32E-09
397	5.91E+05	1.00E-05	1.69E-11
399	3.17E+04	1.00E-05	3.15E-10
400	4.88E+05	6.24E-04	1.28E-09
402	3.04E+04	6.48E-05	2.14E-09
403	3.40E+04	5.97E-05	1.76E-09
408	3.86E+04	1.00E-05	2.59E-10
410	1.61E+05	3.29E-04	2.05E-09
413	2.96E+05	1.00E-05	3.38E-11
415	8.02E+05	1.00E-05	1.25E-11

TABLE 3-1-continued

Binding Kinetics			
Antibody ID	k_a (M ⁻¹ s ⁻¹)	k_d (s ⁻¹)	K_D (M)
495	2.16E+05	1.55E-05	7.18E-11
499	6.33E+04	1.88E-05	2.97E-10
500	2.66E+05	1.14E-04	4.30E-10
501	2.98E+04	2.61E-04	8.78E-09
502	1.08E+05	2.69E-04	2.49E-09
506	4.01E+05	2.10E-04	5.23E-10
511	2.42E+05	2.15E-04	8.87E-10
513	2.63E+05	2.62E-05	9.96E-11
515	3.02E+04	1.34E-03	4.42E-08
518	4.01E+05	2.75E-05	6.85E-11
519	1.41E+05	2.38E-04	1.69E-09
520	4.36E+04	1.00E-05	2.29E-10
521	3.02E+04	2.99E-04	9.90E-09
524	7.86E+04	2.72E-03	3.46E-08
525	3.67E+05	3.94E-05	1.07E-10
526	2.89E+04	4.20E-04	1.45E-08
528	5.60E+05	2.85E-04	5.09E-10
530	9.54E+04	2.09E-04	2.19E-09
534	2.48E+04	1.00E-05	4.04E-10
535	1.05E+05	4.15E-05	3.95E-10
539	9.26E+04	5.13E-04	5.54E-09
540	9.25E+04	1.00E-05	1.08E-10
552	2.58E+05	1.01E-05	3.93E-11
554	2.71E+05	3.50E-05	1.29E-10
557	2.33E+05	3.35E-05	1.44E-10
558	3.78E+05	8.94E-05	2.37E-10
559	1.96E+04	6.31E-04	3.22E-08
561	2.90E+05	4.25E-05	1.46E-10
562	5.48E+05	1.16E-05	2.11E-11
567	1.83E+05	4.62E-04	2.52E-09
570	2.20E+05	1.00E-05	4.54E-11
574	1.32E+05	3.00E-04	2.28E-09
575	1.33E+05	7.52E-04	5.65E-09
584	6.55E+04	8.67E-04	1.32E-08
585	1.85E+05	4.59E-05	2.48E-10

TABLE 3-2

[illegible]

TABLE 3-2-continued

Epitope Binning												
Antibody ID	Positive controls										Negative controls	
	S1 ND				S1 RBD		S2					
	S652-19	S652-22	S652-102	S652-118	S652-109	S652-115	S652-112	S652-103	S652-116	S652-123	8203-C1 (foldon)	PA1-983 (anti-His)
356	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
357	NC	NC	NC	NC	NC	NC	C*	C*	C*	NC	NC	NC
358	NC	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC
359	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
360	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
361	NC	NC	NC	NC	NC	NC	C*	C	C	NC	NC	NC
369	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
370			NC		NC		C	C	C			NC
381	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
388	NC	NC	NC	NC	NC	NC	C	C	NC	NC	NC	NC
389	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
390	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
391	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
393	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
394	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
397	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
400	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
402	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
408			NC		C		NC	NC	NC			NC
413	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
415	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
417	C	C	C*	C	NC	NC	NC	NC	NC	NC	NC	NC
419			NC		NC		NC	NC	NC			NC
428	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
429	NC	NC	NC	NC	NC	NC	C*	C*	C*	NC	NC	NC
441	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
442			NC		NC		NC	NC	NC			NC
447	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
449	C	NC	NC	C	NC	NC	NC	NC	NC	NC	NC	NC
453	NC	NC	NC	NC	NC	NC	C*	C*	C*	NC	NC	NC
455	NC	NC	NC	NC	NC	NC	C*	C*	C*	NC	NC	NC
458	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
459			NC		NC		NC	NC	NC			NC
460			C		NC		NC	NC	NC			NC
461	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
462			NC		NC		NC	NC	NC			NC
463	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
464	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
465			NC		NC		NC	NC	NC			NC
469	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
473	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
478	NC	NC	NC	NC	NC	NC	C*	C*	C*	NC	NC	NC
479	C	C	C *	C	NC	NC	NC	NC	NC	NC	NC	NC
481	C	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
483	C	C	NC	C	NC	NC	NC	NC	NC	NC	NC	NC
486	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
488	NC	NC	NC	NC	C	NC	NC	NC	NC	NC	NC	NC
489	NC	NC	NC	NC	NC	NC	C	C*	C*	NC	NC	NC
491	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
493	C	NC	C*	C	NC	NC	NC	NC	NC	NC	NC	NC
495	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
499	C	NC	C*	C	NC	NC	NC	NC	NC	NC	NC	NC
500	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
502	C	NC	NC	C	NC	NC	NC	NC	NC	NC	NC	NC
506	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
511	NC	NC	NC	NC	NC	NC	C*	C	C*	NC	NC	NC
513	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
518	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
519			NC		NC		NC	NC	NC			NC
520			NC		NC		C	C	C			NC
525	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC

TABLE 3-2-continued

Epitope Binning												
Antibody ID	Positive controls									Negative controls		
	S1 ND			S1 RBD			S2					
	S652-19	S652-22	S652-102	S652-118	S652-109	S652-115	S652-112	S652-103	S652-116	S652-123	8203-C1 (foldon)	PA1-983 (anti-His)
528	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
530	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
534			NC		NC		NC	NC	NC			NC
535			NC		NC		NC	NC	NC			NC
539	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
540	NC	NC	C	NC	NC	NC	C*	C*	C*	NC	NC	NC
552	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
554	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
557	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC	NC
558	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
561	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC
562	C	NC	C	C	NC	NC	NC	C	C	NC	NC	NC
567	NC	NC	NC	NC	NC	NC	C	C*	C*	NC	NC	C
574	NC	NC	NC	NC	NC	NC	C	C	C	NC	NC	NC
575	C	NC	C	C	NC	NC	NC	NC	NC	NC	NC	NC
585	NC	NC	NC	NC	NC	NC	C	NC	NC	NC	NC	NC

Example 4

[1649] Epitope Binning, ACE2 Blocking and Binding Kinetics of Selected mAbs

[1650] A CARTERRA® LSA™ instrument, a fully integrated HT-SPR™ (high throughput surface plasmon resonance) system, was used for premix epitope binning, ACE2 blocking assay, and binding kinetics measurement of selected anti-SARS-CoV2 mAbs colony-cloned from CHO cells. The assays were performed according to the manufacturer's brochure. The reagents used in the experiments are shown in Table 4-1.

TABLE 4-1

Reagents		
Name	Provider	Catalog No.
Recombinant Human ACE2	R&D Systems	933-ZN-010
2019 nCoV Spike Protein (RBD, His tag)	Sino Biological	40592-V08H
SARS-CoV-2 (2019-nCoV) Spike protein (S2 ECD, His Tag)	Sino Biological	40590-V08B
NCP-CoV (2019-nCoV) Spike Protein (S1 Subunit, His Tag)	Sino Biological	40591-V08H
nCoV-5 Spike Protein Trimer	NIH	N/A
Dextran NSB Reducer	GE Health Sciences	BR100691
Heparin Sodium Salt	Fisher	BP2425

[1651] The instrument uses a multi-channel buffer of 25 mM MES at pH 5.5, and a single-channel buffer of 1×HBSEPP+. HC30M chips were used and the array preparation was performed according to the CARTERRA® protocol, including chip activation for 7 minutes in activation buffer (100 µL ECD+100 µL sNHS+100 µL 25 mM MES pH 5.5),

coupling to samples in 10 mM acetate (pH 4.0) for 10 minutes, and deactivation for 7 minutes in 1M ethanolamine (pH 8.5).

[1652] For premix epitope binning, 20 nM of SARS-CoV-2 spike protein was premixed with 200 nM of the mAb diluted in 1×HBSEPP with 0.1 mg/mL BSA, and incubated for a minimum of 3 hours. The complex of spike protein/mAb was then tested for binding to the immobilized mAbs on the prepared HC30M chips, with association for 5 minutes and dissociation for 1 minute. Regeneration was performed in 20 mM glycine (pH 2.0) with 1M NaCl for 45 seconds twice.

[1653] To test the mAbs' ability to block ACE2, 20 nM of SARS-CoV-2 spike protein was premixed with 200 nM of the His-tagged ACE2 (ACE2-His) diluted in 1×HBSEPP with 0.5M NaCl, 1% BSA, 1× Dextran, and 2 mg/mL Heparin, and incubated for about 12 hours. The complex of spike protein/ACE2-His was then tested for binding to the immobilized mAbs on the prepared HC30M chips, with association for 5 minutes and dissociation for 1 minute. Regeneration is performed in 20 mM glycine pH 2.0 with 1M NaCl for 30 seconds twice.

[1654] To test binding kinetics, the prepared HC30M chips were tested against SARS-CoV-2 spike protein with titration beginning at 200 nM with three-fold serial dilutions, with association for 5 minutes and dissociation for 20 minutes. Regeneration was performed in 20 mM glycine (pH 2.0) with 1M NaCl for 45 seconds twice. For the domain-specific protein (RBD, S2 or S1 subunit), the tested concentrations were 500 nM, 250 nM, and 125 nM. The test was performed at either 25° C. or 37° C.

[1655] Tables 4-2 and 4-3 summarize the epitope bin, binding kinetics, neutralization and ACE2-blocking activities of selected mAbs, including 292, 309, 364, 373, 388, 408, 414, 417, 419, 442, 445, 447, 462, 479, 481, 483, 488, 494, 506, 540, 549, 553, 555, and 562.

TABLE 4-2

Epitope bins, binding kinetics, neutralization, and ACE2-blocking activities of selected mAbs								
mAb ID	Epitope Bin	Ace2-block?	K_D (M)-25° C.	K_d (M)-25° C.	K_D (M)-37° C.	K_d (M)-37° C.	S1 K_D (M)-25° C.	RBD K_D (M)-25° C.
292	1	Y	6.08E-11	3.09E-05	7.86E-11	8.64E-05	1.18E-07	6.78E-09
309	2	N	1.58E-11	8.34E-06	3.31E-11	3.08E-05		
364	3	N	7.61E-09	2.80E-04	2.33E-11	1.00E-05		
373	2	Y	1.07E-08	4.81E-04	3.70E-11	1.00E-05		
388	2	N	1.63E-11	8.61E-06	4.47E-11	4.71E-05		
408	2	Y	1.74E-11	1.03E-06	5.58E-11	1.24E-05	1.71E-07	
414	N/A	N	7.93E-09	1.52E-04	5.21E-09	8.84E-04		
417	1	Y	2.14E-11	1.02E-05	1.96E-10	1.03E-04	7.50E-07	
419	2	N	3.78E-10	1.04E-05	1.25E-09	7.75E-04	2.35E-07	
442	4	N	1.76E-08	1.74E-04	1.01E-09	2.95E-05		
445	2	N	1.20E-09	1.90E-05	4.17E-10	1.27E-04	6.13E-07	
447	Y		1.10E-10	3.74E-05	2.60E-10	1.62E-04	1.48E-07	4.38E-09
462	5	N	7.58E-09	1.95E-04	4.62E-09	3.42E-04		
479	6	Y	5.59E-11	1.61E-05	2.86E-11	1.00E-05	4.20E-07	
481	1	Y	1.18E-10	2.62E-05	2.34E-10	8.99E-05	1.70E-07	1.41E-08
483	6	Y	5.27E-11	1.48E-05	2.33E-10	7.95E-05		
488	1	Y	5.26E-11	4.15E-06	4.76E-10	7.61E-05	3.65E-08	1.1E-08
494	1	Y	8.35E-11	1.38E-05	2.25E-10	9.08E-05	1.89E-07	1.35E-08
506	1	Y	4.91E-11	2.06E-05	6.93E-11	7.68E-05	9.09E-08	1.47E-08
540	4	N	1.22E-10	1.18E-05	9.05E-10	1.18E-04		
549	1	Y	2.57E-13	2.03E-08	5.00E-10	7.27E-05	4.14E-08	1.39E-08
553	1	Y	1.31E-10	9.50E-06	3.57E-10	6.70E-05	1.53E-07	
555	7	Y	2.44E-11	1.63E-05	5.77E-11	6.34E-05	7.97E-09	4.13E-09
562	4	N	1.73E-10	4.97E-05	1.38E-10	5.77E-05		

TABLE 4-3

Epitope Bins	
Epitope Bin	mAb Properties
1	ACE2 blocking, RBD or NTD binder
2	S1 or S2 (not RBD) binder, some may block ACE2
3	S2 binder
4	S2 binder
5	S2 binder
6	S1 or S2 (not RBD), but ACE2 blocking
7	RBD Binder (bins with VRC RBD Benchmark), ACE2 blocking

Example 5

[1656] Biophysical Characterization of Selected mAbs

[1657] Materials and Methods

[1658] mAbs were subjected to several biophysical characterizations including analytical size exclusion chromatography (SEC-HPLC), hydrophobic interaction chromatography-high performance liquid chromatography (HIC-HPLC), heparin chromatography (Heparin-HPLC), cross-interaction chromatography (CIC), and affinity-capture self-interaction nanoparticle spectroscopy (AC-SINS). See Table 5-1.

[1659] SEC-HPLC

[1660] 4 μ g of samples were injected onto a Waters BEH200 SEC, 4.6 $\times$ 150 mm, 1.7 μ m column (Waters Cat #186005225). A flow rate of 0.3 mL/min with the running buffer containing 50 mM sodium phosphate, pH 6.8, 0.3M NaCl, 0.005% sodium azide was used. UV absorbance was monitored at 280 nm using an Agilent 1260 HPLC. Retention time (RT) of main peak and percentage of monomer are reported.

[1661] HIC-HPLC

[1662] 20 μ g IgG samples (1 mg/mL) were diluted 1:1 with 2 $\times$ Buffer A concentrate (2 M ammonium sulfate, 0.1 M sodium phosphate at pH 6.8) to achieve a final ammonium sulfate concentration of 1 M before analysis. A TSK-GEL® butyl-NPR (4.6 mm ID $\times$ 10 cm, 2.5 μ m, Tosoh #42168) column was used with a linear gradient (0-100% buffer B) of mobile phase A (1M ammonium sulfate, 50 mM sodium phosphate, pH 6.8) and mobile phase B solution (50 mM sodium phosphate, pH 6.8) over 23 min at a flow rate of 1 mL/min with UV absorbance monitoring at 280 nm. Retention time of main peak is reported.

[1663] Heparin-HPLC

[1664] 20 μ g of IgG samples (1 mg/mL) in PBS was injected onto a POROS™ Heparin 50 μ m (4.6 $\times$ 50 mm, 0.8 mL, Thermo Scientific #4333412) column. Flow was kept at 1.5 mL/min with an initial linear gradient from 0% buffer B to 40% buffer B in 6 minutes then up to 60% B in 2 minutes, followed by a 1 minute gradient increase to 100% B. Gradient was kept at 100% for an additional minute to remove any remaining protein. Retention time of main peak is reported.

[1665] CIC

[1666] CIC was performed as described previously (Jacobs, S. A., et al., *Pharm. Res.* 27, 65-71, 2010). In brief, the CIC column was prepared by coupling ~30 mg of human serum polyclonal antibodies (I4506; Sigma) to a 1-mL HITRAP® column (17-0716-01; GE Healthcare), followed by quenching with ethanolamine. The blank column was prepared in the same except without human serum IgGs. Approximately 20 μ g of each antibody was tested at a flow rate of 0.2 mL/min using 10 mM sodium citrate, 10 mM NaCl, pH 6.5 as a mobile phase on an Agilent 1260 series HPLC system. Retention times obtained by both IgG and blank columns were used to calculate k'. In addition, due to

peak tailing in some samples, peak width at 50% height is also obtained to monitor non-specific interaction of test antibodies.

[1667] AC-SINS

[1668] The AC-SINS assay was performed as described previously with modifications (Wu, J. et al., *Protein big. Des. Sel.* 28, 403-14, 2015). In short, gold nanoparticles (15705; Ted Pella Inc.) were coated with 100% capturing anti-human goat IgG Fc (109-005-008; Jackson ImmunoResearch). The conjugation reaction was quenched with 1 µg/mL polyethylene glycol (PEG) and eluted into 0.25× PBS. PEG was added to the conjugated gold mixture at a concentration of 0.2 µg/mL prior to use. The antibodies of interest were then incubated with the coated gold particles for 1 h at room temperature and the wavelength shift was measured using Tecan M1000 Pro Plate Reader within the range of 475-625 nm, in increments of 1 nm. Test antibodies were diluted in either PBS or 10 mM histidine, pH 6.0 prior to incubation. Delta plasma wavelength shift in comparison with buffer control is reported. The self-interacting antibodies show a higher wavelength shift away from the buffer controls.

All assays were run in duplicate. Benchmark mAb S652-118 and immunized mouse serum were used as positive controls, using seven titration points to calculate IC₅₀, IC₈₀ and IC₉₀ values (see Tables 6-2 and 6-3). Sigmoidal curves were generated and IC₅₀ and IC₈₀ values were calculated as the concentration of antibody required to neutralize 50% and 80% of the virus, respectively (see Table 6-4). Antibodies for which a curve could not be fit are indicated as having an IC₅₀ value of >50 µg/mL.

TABLE 6-1

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 µg/mL	10 µg/mL	1 µg/mL	0.1 µg/mL	0.01 µg/mL
258		6	33	3	11
259	0		0		
260		12	16	23	
261		9	7	-6	12
262		16	8	8	

TABLE 5-1

Biophysical properties of selected mAbs										
Antibody ID	aSEC RT (min)	aSEC % monomer	HIC RT (min)	HepSO4 RT (min)	CIC RT (min) (IgG)	CIC RT (min) (Blank)	CIC k'	CIC Peak Width 50%	AC-SINS H6N30 (Delta pwl)	AC-SINS PBS (Delta pwl)
291	4	92.7	3	2.0	4.8	4.3	0.09	1.7	-3.0	0.0
309	4.1	87.3	5.3	2.1	4.6	4.3	0.07	1.5	-3.0	0.0
364	6.4	79.5*	18.6	2.2	6.6	5.6	0.2	3.5	3.0	3.0
373	4.6	90.3	17.0	1.3	5.6	5.8	-0.04	2.5	0.0	1.0
388	4.1	91.6	1.9	1.3	4.9	4.4	0.13	1.4	0.0	1.0
408	4.9	92	15.2	2.2	5.2	4.8	0.1	2.1	1.0	3.0
414	8.2	97.4	25.9	1.9	7.5	6.1	0.23	3.7	5.0	8.0
417	4.8	96	11.7	1.6	4.6	4.4	0.05	1.5	-3.0	0.5
419	4	94	9.1	0.4	5.3	4.7	0.12	2.2	21.0	2.3
442	4.1	90.4	5.1	2.0	4.6	4.3	0.06	1.5	-3.0	0.0
445	3.9	88.4	6.5	0.4	4.7	4.5	0.06	1.5	7.0	0.0
447	4.1	92.6	2.2	2.3	4.9	4.4	0.12	1.6	-3.0	0.0
462	4.3	93.6	9.1	1.8	4.7	4.4	0.05	1.5	-1.0	1.0
479	5.5	75.8	8.5	1.7	4.6	4.4	0.05	1.5	-3.0	0.8
481	4.2	91.3	9.4	1.6	4.7	4.4	0.09	1.7	-3.0	0.0
483	3.9	94.4	5.7	0.3	4.5	4.3	0.03	1.4	-0.5	0.0
488	4	92.6	5.7	1.4	4.7	4.5	0.04	1.6	-2.0	0.0
494	4	90.7	5.9	1.5	4.6	4.3	0.05	1.6	-3.0	0.0
506	4	88.4	6.4	1.5	4.3	4.4	-0.02	1.4	-3.0	0.0
540	4	92	1.8	2.4	4.8	4.3	0.1	1.5	-2.0	0.3
549	4.2	89.1	13.8	2.1	5.2	4.7	0.1	2.7	N/A	N/A
553	4.1	85.3	7.2	2.1	4.7	4.4	0.08	1.5	-3.0	1.0
555	5	97.8	10.7	2.5	5.1	4.5	0.12	2.2	0.5	1.0
562	4.1	91.9	6.6	1.8	4.6	4.4	0.05	1.5	-3.0	0.0

Example 6

[1669] Pseudoneutralization Assay

[1670] Pseudoneutralization assays were performed using four concentrations of antibodies. Antibody dilutions were premixed with pseudotyped lentivirus expressing the SARS-COV-2 full-length spike protein and incubated with 293T cells transiently expressing human ACE2 receptor for 72 hours. Cells were lysed and viral entry into cells was quantified by Promega Luciferase assay kit using a plate reader. % neutralization at each concentration was calculated by comparison to virus only (0% neutralization) and cell only (100% neutralization) control wells (see Table 6-1).

TABLE 6-1-continued

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 µg/mL	10 µg/mL	1 µg/mL	0.1 µg/mL	0.01 µg/mL
263	0	0	12	5	
264	0	26	41	42	
265	5	0	20	22	
266	23	18	31	38	
267	9	0	0	0	
268		33	-1	-2	2

TABLE 6-1-continued

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 $\mu\text{g/mL}$	10 $\mu\text{g/mL}$	1 $\mu\text{g/mL}$	0.1 $\mu\text{g/mL}$	0.01 $\mu\text{g/mL}$
269		7	10	-4	-4
270	0	0	0	0	
271	10	6	22	29	
272		-3	-6	-8	-4
273	16	0	0	0	
274		5	5	25	13
274	38	8	3	0	
275		13	-12	-2	-5
276	44	50	10	17	
277		-1	7	18	22
278		-6	-13	-17	-27
279		18	20	10	6
280		18	-4	-12	-19
281	9	0	0	5	
282		4	18	26	31
283		10	15	21	22
284	0	0	0	0	
285		1	-15	-32	-16
287		7	19	-8	-3
288	0	0	0	0	
289	25	0	0	18	
290	0	0	19	14	
291		24	16	-1	5
292		60	1	-2	6
293		-10	15	-7	-3
294	83	80	56	37	
295	33	5	0	29	
296		10	7	-15	0
297		-17	7	15	11
298		15	2	29	18
299	36	13	9	34	
300	40	14	0	13	
301	0	0	0	2	
302		24	10	38	23
303	0	0	0	0	
304		-11	6	12	16
305		16	6	7	12
306	0	0	0	0	
308		19	13	-14	2
309		-19	-10	32	30
310		9	2	5	-14
311		-15	-7	39	33
312		10	11	-1	18
313		8	10	-3	13
314		-1	4	-16	4
315	21	22	22	0	
316	36	32	35	25	
317	39	30	28	20	
318		26	-12	11	13
319	45	39	34	36	
320	96	93	68	22	
321		13	7	-2	34
322		18	3	-17	-20
323	0	0	0	16	
324	47	44	33	33	
325		24	14	5	10
326	31	24	21	0	
327	26	22	15	12	
328	0	0	0	0	
329	0	0	20	10	
330		23	22	18	16
331	0	0	0	1	
332	0	0	3	0	
333	0	0	0	0	
334		20	18	7	14
335		-19	-7	33	33
335	14	0	0	0	
336		14	19	12	16
337	28	0	11	11	
338		26	22	11	18
339	0	0	24	7	

TABLE 6-1-continued

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 $\mu\text{g/mL}$	10 $\mu\text{g/mL}$	1 $\mu\text{g/mL}$	0.1 $\mu\text{g/mL}$	0.01 $\mu\text{g/mL}$
340	0	0	0	6	
341		-25	-23	37	23
342	0	0	21	17	
343	0	8	17	27	
344	27	32	38	24	
345	37	0	0	0	
347	12	32	24	16	
348	41	19	17	0	
349	0	8	0	0	
350		17	9	-10	-23
351	37	31	1	0	
352		22	16	16	19
353		1	16	-1	-4
354		10	21	9	4
355		3	11	15	-3
356		10	6	-9	2
357		13	6	9	2
358		19	15	3	17
359		14	20	2	-14
360		25	13	-12	2
361		17	1	2	6
362	0	0	0	0	
363	0	0	0	2	
364		19	10	-5	11
365	26	17	5	21	
366	21	0	13	24	
367	15	2	0	0	
368		10	24	20	21
370		8	23	5	7
371		7	21	-2	-4
372	5	0	0	0	
373		60	37	25	15
374		23	9	-3	21
375	15	15	9	35	
376		-4	-4	-2	2
377		15	-5	-6	-3
378	0	0	0	0	
379	0	0	13	16	
380	30	10	29	40	
381		10	19	24	27
382		27	34	20	22
383		0	-3	-7	-5
384		28	17	13	23
385	0	0	0	0	
387		12	14	17	15
388		19	32	49	36
389		-12	2	-4	-7
390		-4	3	8	8
391		-2	-10	-3	-5
392	0	0	10	10	
393		10	21	40	25
394		5	5	4	-2
395	14	0	31	34	
396	35	19	33	34	
397		-1	-5	11	6
398	27	17	26	43	
399		-8	-13	23	7
400		-6	-7	2	7
401		0	4	9	4
402		27	41	13	23
403		22	12	-23	-7
404		-14	-10	-27	-22
405	23	22	15	22	
406		7	10	23	32
407	19	25	41	34	
408		35	14	5	-1
409	25	17	33	13	
410		5	-37	10	7
412	93	99	22	23	
413		5	0	-8	5
414		32	36	26	41

TABLE 6-1-continued

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 $\mu\text{g/mL}$	10 $\mu\text{g/mL}$	1 $\mu\text{g/mL}$	0.1 $\mu\text{g/mL}$	0.01 $\mu\text{g/mL}$
415		19	30	-6	24
416	0	0	16	19	
417		55	32	29	37
418	3	13	0	3	
419		36	35	5	5
420		3	-2	7	24
421	26	24	0	0	
422	91	87	82	18	
423	40	48	30	5	
424		8	13	12	29
425	41	47	35	17	
426		21	24	-3	9
427	0	0	18	34	
428		-11	-4	-14	26
429		-19	-3	-76	-1
431	46	45	33	16	
432		8	-6	1	-15
433		14	-9	-6	18
434	35	44	36	18	
435	2	0	0	47	
436		11	19	28	33
437	0	0	0	0	
438		-11	-7	4	4
439	25	17	0	0	
440		11	12	-3	7
441		-17	8	13	24
442		33	9	27	26
443	31	0	0	16	
444		0	4	24	21
445		38	12	9	-9
446	30	10	0	6	
447		87	22	15	35
448		19	14	6	-1
449		-4	-12	1	2
450		15	-12	14	4
451		5	-13	-10	-21
452	40	0	3	17	
453		-12	-6	-18	-19
453	32	7	0	0	
454	37	17	25	15	
455		14	2	20	3
456	0	0	0	0	
457		-10	-13	-20	-13
458		-29	-14	-18	-17
459		7	10	-18	-28
460		16	3	21	32
461		-1	4	23	13
462		40	9	6	38
463		18	16	16	38
464		-6	-10	-2	11
465		-8	2	10	13
467		3	9	20	23
468		13	7	22	9
469		-1	-10	1	7
470	15	0	15	9	
471	41	0	0	10	
472		-8	-6	12	10
473		-11	9	15	-4
474		0	0	0	
475	0	0	9	22	
476	0	0	0	0	
477	0	0	7	9	
478		17	8	-13	-17
479		30	11	24	43
480	38	27	27	39	
481		55	17	10	36
482		0	15	6	2
483		31	43	5	11
484		10	28	8	8
485	30	16	25	13	
486		13	20	-13	-5

TABLE 6-1-continued

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 $\mu\text{g/mL}$	10 $\mu\text{g/mL}$	1 $\mu\text{g/mL}$	0.1 $\mu\text{g/mL}$	0.01 $\mu\text{g/mL}$
487	24	9	15	0	
488		74	19	39	25
489		-5	15	12	32
491		11	-4	34	18
491	54	26	4	0	
492	36	21	0	0	
493		0	3	-2	-7
494		13	19	23	10
495		17	14	8	6
496	38	26	6	0	
497	15	0	15	26	
498	34	20	21	4	
499		-5	2	-1	5
500		-29	-45	-17	-4
501		23	27	17	13
503		-4	0	14	9
504	47	4	0	0	
505	11	8	0	0	
506		38	21	-1	-15
507	12	0	21	16	
508		-8	-6	-14	3
509	59	23	36	23	
510	20	15	19	63	
511		16	11	4	8
512	24	25	23	15	
513		12	10	7	-6
514	46	38	1	4	
515		5	-11	43	22
516	44	45	40	24	
517	0	0	0	0	
518		-1	-12	-6	7
519		5	9	9	14
520		-1	2	-7	1
521		15	16	4	11
522	27	38	35	57	
523	31	37	42	0	
524		13	18	29	25
525		8	24	20	33
526		6	0	7	22
527	13	14	19	0	
528		-3	-17	-30	-27
529		8	7	4	-8
530		-7	-11	-30	-11
531	14	0	27	7	
533	3	0	0	0	
534		13	17	3	8
535		30	21	-1	12
536	39	0	29	3	
537	0	32	21	11	
538	33	9	31	4	
539		31	20	10	13
540		15	18	22	12
541	21	0	0	0	
542	6	24	0	0	
543	63	13	18	18	
544	15	0	0	0	
545	24	0	0	0	
546	50	25	0	0	
547	0	0	10	17	
548	25	9	0	0	
549		84	55	25	27
550	38	1	0	0	
551	57	0	0	15	
552		41	6	19	28
553		26	28	8	6
554		-1	-1	4	10
555		88	47	-5	7
556	0	0	0	7	
557		4	13	-2	1
558		9	-10	3	7
559		9	5	45	27

TABLE 6-1-continued

Four-point Pseudoneutralization screen of monoclonal antibodies % Neutralization					
Antibody ID	50 $\mu\text{g/mL}$	10 $\mu\text{g/mL}$	1 $\mu\text{g/mL}$	0.1 $\mu\text{g/mL}$	0.01 $\mu\text{g/mL}$
559	34	17	42	26	
560	0	0	18	8	
561		-7	12	-7	-11
562		18	8	6	-19
563	18	11	5	0	
564	13	0	0	0	
565		16	20	24	24
566	11	0	0	4	
567		5	25	18	15
568		-4	6	5	-2
569	15	15	5	0	
570		-2	-10	-5	-17
571	36	3	33	35	
573		-11	-7	-15	-10
574		-21	4	3	15
575		13	-1	-2	2
576	3	17	17	0	
577		9	2	9	9
578		0	0	0	
579		14	4	18	1
580	0	0	0	0	
581	13	20	25	0	
582	11	0	0	0	
583	0	0	21	19	
584		9	30	25	26
585		4	-5	42	17
586	0	0	0	0	
587		1	-9	3	18
588	0	0	24	52	

TABLE 6-2

Positive Control	Concentration ($\mu\text{g/mL}$)	% Neutralization
S652-118	50.0	65.94712289
	12.5	56.6591417
	3.125	35.52770257
	0.78125	14.37639388
	0.1953125	-3.015215894
	0.048828125	41.2447797
	0.012207031	-4.806124096

TABLE 6-3

Positive Control	Dilution	% Neutralization
Moose serum	50	91.76866015
	200	67.11624723
	800	11.53270188
	3200	-7.777682412
	12800	-9.670886593
	51200	-6.058915489
	204800	7.678774689

TABLE 6-4

IC ₅₀ and IC ₈₀ values for select anti-coronavirus antibodies			
Antibody ID	IC ₅₀ ($\mu\text{g/mL}$)	IC ₈₀ ($\mu\text{g/mL}$)	IC ₉₀ ($\mu\text{g/mL}$)
292	>50	>50	
309	>50	>50	
364	>50	>50	
373	>50	>50	
388	>50	>50	
408	>50	>50	
414	>50	>50	
417	>50	>50	
419	>50	>50	
442	>50	>50	
445	>50	>50	
447	>50	>50	
462	>50	>50	
479	>50	>50	
481	17.16	>50	
483	>50	>50	
488	3.52	>50	
494	>50	>50	
506	>50	>50	
539	>50	>50	
540	>50	>50	
549	>50	>50	
552	>50	>50	
553	>50	>50	
555	0.38	1.40	
562	>50	>50	
Controls			
S652-118	31.01	>50	n/a
Mouse serum	78	58	51

Example 7

[1671] Epitope Mapping of Anti-SARS-COV-2 Antibody 555 and Other Anti-SARS-COV-2 Antibodies by Hydrogen Deuterium Exchange Mass Spectrometry (HDX-MS)

[1672] Hydrogen deuterium exchange with mass spectrometry (HDX-MS) was performed in order to determine where the exemplified antibody (mAb 555) binds the SARS COV-2 spike protein. Peptide identification for SARS Cov-2 spike was performed on a Waters SYNAPT™ G2Si (Waters Corporation) instrument using 5 μg of SARS COV-2 spike at zero exchange (0.1 $\times$ PBS in H₂O) using nepenthesin II (Nep II) for digestion. The mass spectrometer was set in HDMSe (Mobility ESI+ mode); in a mass acquisition range of m/z 255.00-1950.00; with a scan time of 0.4 s. Data was processed using PLGS 2.3.2 (Waters Corporation). For the exchange experiments, the complex of human SARS COV-2 spike with mAb 555 was prepared at the molar ratio of 1:1.2 in 10 mM sodium phosphate buffer, pH 7.4 containing 150 mM NaCl (1 $\times$ PBS buffer). The experiment was initiated by adding 25 μL of D₂O buffer containing 0.1 $\times$ PBS to 2.5 μL of spike (1 mg/mL) or spike-mAb 555 complex at 15° C. for various amounts of time (0 s, 10 s, 30 s, 2 min, 10 min and 120 min) using a custom Tecan sample preparation system (Espada et al., *J. Am. Soc. Mass Spectrom.*, 2019, 30:2580-2583). The reaction was quenched using equal volume of was 0.32 M TCEP, 3 M guanidine HCl, 0.1M phosphate pH 2.5 for two minutes at 4° C. and immediately frozen at -70° C. The sample injection system is comprised of a UR3 robot, a LEAP PAL3 HDX autosampler, and a HPLC system interfaced with a Waters SYNAPT® G2-Si (Waters Corporation) (modified from Espada et al.). The LC mobile phases

consisted of water (A) and acetonitrile (B), each containing 0.2% formic acid. Each sample was thawed using 50 μ L of 0.32 M TCEP, 1.5 M guanidine HCl, 0.1 M phosphate pH 2.5, for 1 min and injected on to a Nep II column for digestion at 4° C. with mobile phase A at the flow rate of 250 L/min for 2.5 minutes. The resulting peptides were trapped on a Waters BEH VANGUARD™ Pre-column at 4° C., and chromatographically separated using a Waters ACQUITY UPLC™ BEH C18 analytical column at 4° C. with a flow rate of 200 μ L/min and a gradient of 3%-85% mobile phase B over 7 minutes, and directed into mass spectrometer for mass analysis. The SYNAPT® G2-Si was calibrated with Glu-fibrinopeptide prior to use. Mass spectra were acquired over the m/z range of 255 to 1950 in HDMS mode, with the lock mass m/z of 556.2771 (Leucine Enkephalin). The relative deuterium incorporation for each peptide was determined by processing the MS data for deuterated samples along with the undeuterated control using the identified peptide list in DYNAMX™ 3.0 (Waters Corporation). The free and bound states of SARS Cov-2 spike were compared for deuterium incorporation differences.

[1673] The results are shown in Table 7.1. Sequence coverage for the SARS Cov-2 spike was 63.5%. Decrease in deuterium uptake between SARS Cov-2 spike plus mAb 555 complex vs. SARS CoV-2 spike alone was observed between residues 459-495 (SNLKPERISTEIQAGSTPCNGVEGFNCYFPLQSY; SEQ ID NO:5380) and residues 434-444 (IAWNSNNLDSK; SEQ ID NO: 5379), pointing to the probable epitope region. These data suggest mAb 555 has an epitope comprising one or more residues within amino acids 434-444 and 459-495 of the SARS-CoV-2 spike protein. The epitope region of mAb 555 correlate to the ACE2 binding site in SARS-CoV-2 spike RBD region (Yan et al., 2020, Science. 367(6485):1444-1448.)

TABLE 7.1

Peptides at which decreased exchange was observed in SARS Cov-2 spike plus mAb 555 complex as compared to SARS Cov-2 spike alone (overlapping peptides observed).		
Residues	Peptide sequence	SEQ ID NO
459-473	SNLKPPERDISTEIQ	5368
467-473	DISTEIQ	5369
468-473	ISTEIQ	5370
472-486	IYQAGSTPCNGVEGF	5371
472-488	IYQAGSTPCNGVEGFNC	5372
474-486	QAGSTPCNGVEGF	5373
474-488	QAGSTPCNGVEGFNC	5374
487-495	NCYFPLQSY	5375
489-495	YFPLQSY	5376
434-439	IAWNSN	5377
434-441	IAWNSNNL	5378
434-444	IAWNSNNLDSK	5379

[1674] HDX-MS were also performed for additional anti-SARS-COV-2 antibodies. The results are summarized in Table 7.2.

TABLE 7.2

Summary of HDX-MS data for additional anti-SARS-COV-2 antibodies. Sequence ranges of the SARS-COV-2 spike protein exhibiting protection are shown; the number of peptides for each sequence range is indicated in parentheses.

MAB	Regions showing protection
NTD Binders	
364	136-144 (2), 71-178 (3), 242-264 (3)
373	136-144 (2), 171-179 (2), 242-264 (3)
417	92-102 (4), 136-144 (2), 171-179(3) 242-264(3)
419	136-144 (2), 242-265 (4)
483	136-144 (2) 242-265 (4)
RBD Binders	
292	472-495 (6)
447	467-513 (9)
462	467-489 (7),
481	467-490 (6) -496-513(2)
488	417-421 (2), 433-444 (2), 467-513 (14)
506	433-455 (4), 496-513(2)
Undetermined/other binders	
408	136-143 (2), 307-318 (2), 621-636 (3)
479	980-1006 (5) 1179-1186 (1)
540	960-1007 (10)

Example 8

[1675] Epitope Mapping of Anti-SARS-COV-2 Fabs by Negative-Stain Electron Microscopy (nsEM)

[1676] Anti-SARS-CoV-2 antibodies were produced by expression in Chinese hamster ovary (CHO) cells, and purified using standard antibody purification techniques. The Fab portions of the antibodies were generated by proteolytic digestion using papain, followed by removal of un-cleaved protein using standard chromatography techniques.

[1677] SARS-CoV-2 spike ectodomain was diluted to 0.04 mg/mL in 2 mM Tris pH 8.0, 200 mM NaCl, 0.02% NaN₃ (dilution buffer) in the presence of 10-fold excess Fab, and incubated on ice for either 10 seconds (Fab 555), 2 minutes (Fab 447), 5 minutes (Fabs 309, 488, 562, 506, 442, 417, 462, 292, 540, 479, 553, 388), or 20 minutes (Fabs 447, 483, 419, 494). CF400-Cu grids (Electron Microscopy Sciences) were plasma cleaned for 30 seconds in a SOLARUS®950 plasma cleaner (Gatan) with a 4:1 ratio of O₂/H₂. 4.8 μ L of the protein sample was applied to the grid and allowed to incubate for 30 sec. The grid was then washed twice with dilution buffer stained with methylamine tungstate (NANO-W, Nanoprobes). Grids were then imaged using an FEI TALOS™ TEM (Thermo Scientific) and a Ceta 16 M detector. Micrographs were collected manually using TIA v4.14 software at a magnification of $\times 92,000$, corresponding to a pixel size of 1.63 Å/pixel. CTF estimation and particle picking were performed in cisTEM. 2D classification was performed in either cisTEM (Grant, et al., 2018. "CisTEM, User-Friendly Software for Single-Particle Image Processing," Edited by Edward H Egelman. *ELife* 7 (March): e35383) or CRYOSPARC™ (Punjani, *Nature Methods* 14 (3): 290-96, 2017), and ab initio reconstruction and refinement for 3D maps were performed in CRYOSPARC™.

[1678] Binding kinetics were determined based on bio-layer interferometry. 50 nM 2X-Strep-tagged SARS-CoV-2 ectodomain in BLI buffer (10 mM HEPES pH 7.5, 150 mM NaCl, 3 mM EDTA, 0.05% Tween 20 and 1 mg/mL BSA) was immobilized onto a streptavidin biosensor (FortéBio) for 600s using an Octet RED96e (FortéBio). The biosensor was then dipped into 100 nM Fab (diluted in BLI buffer), and the association signal was measured for 600 sec. Following this, the biosensor was dipped into BLI buffer to measure the dissociation signal for 600 sec. Data were reference-subtracted and fit to a 1:1 binding model using Octet Data Analysis Software v11.1 (FortéBio).

[1679] The nsEM results and binding kinetics for Fabs 555, 447, 494, 483, 419 and 388 are shown in FIGS. 2-8. The other Fabs were not seen by nsEM. The nsEM imaging and three dimensional reconstruction of the complex confirms that Fab 555 binds the RBD domain of the SARS-CoV-2 spike protein in an orientation that would directly interfere with the known ACE2-spike protein interaction (Yan, et al., 2020, *Science* 367(6485): 1444-1448).

Example 9

[1680] Live Virus Neutralization Assay of Anti-SARS-CoV-2 Antibodies

[1681] The efficacy of some anti-SARS-CoV-2 antibodies (mAbs 555, 419, 481, 488, 373) is measured by detecting the neutralization of infectious SARS-CoV-2 virus in a dose-response mode using cultured Vero E6 cells. These cells are known to be highly susceptible to infection by SARS-CoV-2. The assays were developed and performed at three independent laboratories by modifying previously published methods for use with SAR-CoV-2.

[1682] Assays at Lab 1 and Lab 2 were conducted using natural virus produced by infecting cultured Vero E6 cells with the SARS-CoV-2 clinical isolate USA/WA/1/2020 (BEI resources number NR52281) and incubating at 37° C. until cytopathology was evident (typically 48-72 hours). Expansion was limited to only 1-2 passages in cell culture to retain integrity of the original viral sequence. The virus stock was quantified by standard plaque assay, and aliquots were stored at -80° C. A freshly-thawed aliquot is used for each neutralization experiment.

[1683] Assays at Lab 2 were also conducted using the Italy-INMI1 isolate of SARS-CoV-2 (European Virus Archive—Global, ref #008V-03893), similarly grown in Vero E6 cells, titrated by plaque assay and stored frozen.

[1684] Assays at Lab 3 were performed using a modified version of the USA/WA/1/2020 isolate in which a non-essential gene (ORF7) was replaced by the NANOLUC® luciferase reporter gene (Promega). This technology was previously described for SARS-CoV and MERS-CoV (Sheahan, et al., *Sci. Transl. Med.* 2017; 9 eaal 3653).

[1685] Briefly, 20-140 plaque-forming units of virus were pre-incubated with serial dilutions of anti-SARS-CoV-2 antibodies (8-10 points per curve) for 1 hour at 37° C., inoculated onto monolayers of Vero E6 cells, and incubated at 37° C. to allow the non-neutralized virus to replicate.

[1686] The inhibition of replication resulting from virus neutralization by mAb555 was detected by the following methods:

[1687] Decreased production of viral nucleocapsid protein (NP) as detected using murine monoclonal antibody to SARS-CoV-2 NP with standard immunostaining techniques (Lab 1 method).

[1688] Protection against virus-induced cell death as detected by standard plaque assay (Lab 2 method).

[1689] Decreased signal from the inserted luciferase reporter gene (Lab 3 method).

[1690] Each sample was tested using 2-3 replicates per antibody dilution. Either mouse or human convalescent serum was used as a positive control. Percent neutralization was calculated relative to the signals produced by an IgG isotype antibody control and a no-virus control, and the data were plotted using nonlinear regression with a four-parameter fit analysis (GraphPad Prism v8.0.0). Overall estimates for IC₅₀ and IC₉₉ were made using a meta-analysis with all data from the 4 laboratories using a random effects model (Berkey, et al., *Stat Med.* 1995; 14(4):395-411) and the R package metafor (Viechtbauer, *J. Stat. Software* 2010; 36(3): 1-48).

[1691] The results of the neutralization assays for mAbs 555, 419, 481, 488, and 373 are presented in Table 9.1 and Table 9.2. Taken together, mAb 555 is shown to be a potent inhibitor of virus infectivity, with an estimated IC₅₀=0.03 µg/mL (95% CI: 0.01 µg/mL, 0.12 µg/mL) and IC₉₉=0.43 µg/mL (95% CI: 0.13 µg/mL, 1.43 µg/mL) for the USA/WA/1/2020 clinical isolate, and an estimated IC₅₀=0.05 µg/mL (95% CI: 0.04 µg/mL, 0.05 µg/mL) and IC₉₉=1.42 µg/mL (95% CI: 0.94 µg/mL, 2.13 µg/mL) for the Italy-INMI1 clinical isolate. IC₉₉ was calculated for mAb 555 only.

TABLE 9.1

Results of Neutralization Assays of USA/WA/1/2020 isolate by mAbs 555, 419, 481, 488, 373		
Study Location	Detection Method	IC ₅₀ µg/mL (95% CI)
mAb 555		
Lab 1	Detection of viral nucleoprotein	0.16 (0.15, 0.17)
Lab 2	Plaque reduction	0.02 (0.01, 0.02)
Lab 3-Run #1	Luciferase reporter	0.04 (0.01, 0.23)
Lab 3-Run #2	Luciferase reporter	0.01 (0.00, 0.03)
mAb 555 Overall		0.03 (0.01, 0.12)
mAb 419		
Lab 1	Detection of viral nucleoprotein	0.65 (0.56, 0.75)
Lab 2	Plaque reduction	1.54 (0.30, 7.80)
Lab 3	Luciferase reporter	0.61 (0.40, 0.95)
mAb 419 Overall		0.65 (0.57, 0.74)
mAb 481		
Lab 1	Detection of viral nucleoprotein	3.18 (2.28, 4.44)
Lab 2	Plaque reduction	N/A
Lab 3	Luciferase reporter	6.03 (4.75, 7.66)
mAb 481 Overall		4.43 (2.37, 8.29)
mAb 488		
Lab 1	Detection of viral nucleoprotein	1.69 (0.34, 8.52)
Lab 2	Plaque reduction	0.86 (0.47, 1.59)
Lab 3	Luciferase reporter	0.94 (0.73, 1.22)
mAb 488 Overall		0.94 (0.74, 1.19)
mAb 373		
Lab 1	Detection of viral nucleoprotein	17.4 (4.8, 62.9)
Lab 2	Plaque reduction	N/A
Lab 3	Luciferase reporter	12.8 (3.2, 50.9)
mAb 373 Overall		15.1 (5.9, 38.6)

TABLE 9.2

Results of Neutralization Assays of Italy-INMI1 isolate by mAb 555		
Study Location	Detection Method	IC ₅₀ , ug/mL (95% CI)
Lab 2	Plaque reduction	0.05 (0.04, 0.05)

TABLE 9.3

Results of Neutralization Assays of other anti-SARS-CoV2 mAbs				
MAb	Lab 1 live virus IC ₅₀	Lab3 live virus IC ₅₀	Lab2 PRNT IC ₅₀ (WA isolate)	Lab2 PRNT IC ₅₀ (Italy isolate)
297	1.25 µg/mL	9.9 µg/mL	—	—
294	ND	ND	—	—
309	>20 µg/mL	>100 µg/mL	—	—
320	ND	ND	—	—
364	10 µg/mL	>100 µg/mL	—	—
388	>>40 µg/mL	>100 µg/mL	—	—
408	1.25 µg/mL	1.2 µg/mL	—	—
412	ND	ND	—	—
414	>20 µg/mL	>100 µg/mL	—	—
417	>40 µg/mL	>100 µg/mL	>100 µg/mL	—
442	>40 µg/mL	>100 µg/mL	>100 µg/mL	—
445	ND	ND	—	—
447	1.25 µg/mL	1.8 µg/mL	11.2 µg/mL	—
462	10 µg/mL	>100 µg/mL	—	—
479	>40 µg/mL	>100 µg/mL	—	—
483	0.63 µg/mL	27 µg/mL	>100 µg/mL	—
494	2.5 µg/mL	23.6 µg/mL	—	—
506	1.25 µg/mL	2 µg/mL	—	—
540	>40 µg/mL	>100 µg/mL	—	—
549	ND	ND	—	—
552	ND	ND	—	—
553	20 µg/mL	>100 µg/mL	—	—
562	>40 µg/mL	>100 µg/mL	—	—

Example 10

[1692] Binding of Anti-SARS-CoV-2 Antibody 555 to SARS-CoV-2 Spike Protein with Known Mutations

[1693] To identify mutations arising in the viral population that might impair binding and neutralization by mAb 555, an in-house viral surveillance bioinformatics workflow was established. Sequences of SARS-CoV-2 were downloaded from the GISAID database (Elbe, S., and Buckland-Merrett, G., 2017, *Global Challenges*, 1: 33-46) every 4 days as full-length DNA sequences, which were then processed via custom MATLAB® (MathWorks) scripts to align the spike sequences and extract mutational information with respect to a reference spike protein sequence from the strain hCoV-19/Wuhan/IVDC-HB-01/2019 (EPI_ISL_402119). Optional patient metadata generated by bioinformatic tools Nextstrain (Hadfield et al., 2018, *Bioinformatics*; 34(23), 4121-4123) were used to supplement the sequence data. Sequences were discarded if they contained >5% ambiguous bases of the spike protein, had <80% DNA identity to the reference spike, or contained multiple inserted or deleted bases, all of which indicate sequencing errors rather than antigenic drift. The MATLAB® scripts parsed the filtered data to summarize the frequency of mutation, the codons of the mutated residues, potential mutations, duration in circulation, and locations of strain isolation. For a spike mutation to be considered for in-depth binding characterization with mAb

555, the following threshold was used: mutations must appear >5 times, be isolated from >1 locations, be circulating for >13 days, and reside within the receptor binding domain (RBD, residues 329-520).

[1694] Mutations that have arisen in the receptor binding domain of SARS-CoV-2 variants are listed in Table 10.1, along with the number of occurrences. The RBD mutations are rare and collectively appear in 0.5% of the deposited sequences.

[1695] Binding experiments were carried out with full-length SARS-CoV-2 spike protein. Suspension CHO cells were transiently transfected with the plasmid using electroporation. Full length spike protein expression (using the original Wuhan reference sequence) was confirmed by testing with benchmark antibodies discovered against SARS-CoV that target different stalk and head domains, using flow cytometry. Furthermore, western blot was performed with a whole cell and plasma membrane isolate to confirm full length protein expression on the cell surface. mAb 555 was confirmed to bind the screening target (SARS-CoV-2 full length spike) using high throughput flow cytometry. CHO cells were transiently transfected to express the full length spike protein of either the reference sequence or mutants of SARS-CoV-2 spike protein on the cell surface. mAb 555 was incubated with the readout cells, and an untransfected control CHO line either at 50 nM antibody concentration for 30 minutes at 4° C. CHO cells were washed, and binding was detected by using a fluorescently labeled anti-human secondary antibody. Fluorescence was measured using high throughput plate-based flow cytometry. Benchmark antibodies identified to SARS were used as positive control due to similarity in spike sequences between SARS and SARS-CoV-2; human IgG isotype and an irrelevant antibody were used as negative controls. Median fluorescence intensity of each antibody was normalized over the median fluorescence intensity of the human isotype control for respective antigens. Antibody values greater than 5-fold over isotype were considered as binders. The cut-off value was determined based on the binding to the negative controls.

[1696] mAb 555 was capable of binding to three of the mutations tested (Table 10.2). Of the mutations that have been observed, only two mutated residues (G476S and V483A) reside within the epitope for mAb 555. The first mutation of these to arise, V483A, was incorporated into an isolated receptor binding domain, and the binding affinity of this reagent to mAb 555 was tested as described above. In this experiment, mAb555 bound both the reference receptor binding domain and the mutated domain containing the V483A with similar affinities (Table 10.3).

TABLE 10.1

Mutation frequencies within the receptor binding domain of the SARS-CoV-2 spike protein (residues 329-520) that have been observed within the detection threshold, based on the GISAID database as of 28 Apr. 2020.	
Mutation	Number of occurrences (from a total of >10000 sequence entries)
V367F	15
Q414E	7
G476S	9

TABLE 10.1-continued

Mutation frequencies within the receptor binding domain of the SARS-CoV-2 spike protein (residues 329-520) that have been observed within the detection threshold, based on the GISAID database as of 28 Apr. 2020.		
Mutation	Number of occurrences (from a total of >10000 sequence entries)	
V483A	24	
A520S	6	

TABLE 10.2

Binding of mAb 555 to mutant forms of the full length SARS-CoV-2 spike protein by high-throughput flow cytometry		
Protein	Number of measurements	Range of determined fluorescence signal, relative to isotype control
Negative control (untransfected cells)	2	0.64-0.9 fold over control
Reference Spike protein	3	13.8-28.7 fold over control
V367F	3	18.4-42.6 fold over control
G476S	2	36.7-46.9 fold over control
V483A	4	19.5-62.7 fold over control

TABLE 10.3

Effect of the V483A mutation on the binding affinity of mAb 555, measured by surface plasmon resonance (CARTERRA ® LSA)	
Receptor binding domain	Binding affinity (K_D)
Reference Spike protein	1.96×10^{-9} M
V483A	4.41×10^{-9} M

Example 11

[1697] Nucleic Acid Sequence of Encoding Genes and Translated Amino Acid Sequence of Mab 555

[1698] Non-coding sequences are italicized. DNA coding sequence for the mskappa ss is underlined with a solid line. DNA coding sequence for the variable domain is in bold font. DNA coding sequence for the human kappa constant domain is in normal font. DNA coding sequence for the translational stop codons are italicized and underlined with a solid line.

[1699] DNA Sequence for Encoding the mAb 555 HC

(SEQ ID NO: 5381)

```

1  AAGCTTGCTC GAGCCACCAT GGAGACAGAC ACACTCCTGC TATGGGTACT GCTGCTCTGG
61  GTTCCAGGAT CCACTGGACA GGTGCAGCTG GTGCAGTCTG GGGCTGAGGT GAAGAAGCCT
121 GGGTCCTCGG TGAAGGTCTC CTGCAAGGCT TCTGGAGGCA CCTTCAGCAA CTATGCTATC
181 AGCTGGGTGC GACAGGCCCC TGGACAAGGG CTTGAGTGGA TGGGAAGGAT CATCCCTATC
241 CTTGGTATAG CAAACTACGC ACAGAAAGTTC CAGGGCAGAG TCACGATTAC CGCGGACAAA
301 TCCACGAGCA CAGCCTACAT GGAGCTGAGC AGCCTGAGAT CTGAGGACAC GGCCGTGTAT
361 TACTGTGCGA GAGGTTACTA CGAAGCGAGG CATTACTACT ACTACTACGC TATGGACGTC
421 TGGGGCCAAG GGACCGCGGT CACCGTCTCC TCAGCCTCCA CCAAGGGCCC ATCGGTCTTC
481 CCCCTGGCAC CCTCCTCCAA GAGCACCTCT GGGGGCACAG CGGCCCTGGG CTGCCTGGTC
541 AAGGACTACT TCCCCGAACC GGTGACGGTG TCGTGGAATC CAGGCGCACT GACCAGCGGC
601 GTGCACACCT TCCCGGCTGT CCTACAGTCC TCAGGACTCT ACTCCCTCAG CAGCGTGGTG
661 ACCGTGCCCT CCAGCAGCTT GGGCACCCAG ACCTACATCT GCAACGTGAA TCACAAGCCC
721 AGCAACACCA AGGTGGACAA GAGAGTTGAG CCCAAATCTT GTGAGAAAAC TCACACATGC
781 CCACCGTGCC CAGCACCTGA ACTCCTGGGG GGACCGTCAG TCTTCTCTTT CCCCCAAAA
841 CCCAAGGACA CCCTCATGAT CTCCCGGACC CCTGAGGTCA CATGCGTGGT GGTGGACGTG
901 AGCCACGAAG ACCCTGAGGT CAAGTTCAAC TGGTATGTGG ACGGCGTGGA GGTGCATAAT
961 GCCAAGACAA AGCCGCGGGA GGAGCAGTAC AACAGCACGT ACCGTGTGGT CAGCGTCTTC
1021 ACCGTCTTGC ACCAAGACTG GCTGAATGGC AAGGAGTACA AGTGAAGGT CTCCAAGAAA
1081 GCCCTCCCAG CCCCATCGA GAAAACCATC TCCAAAGCCA AAGGGCAGCC CCGAGAACCA
1141 CAGGTGTACA CCCTGCCCCC ATCCCGGGAG GAGATGACCA AGAACCAAGT CAGCCTGACC
1201 TGCCTGGTCA AAGGCTTCTA TCCCAGCGAC ATCGCCGTGG AGTGGGAGAG CAATGGGCAG
1261 CCGGAGAACA ACTACAAGAC CACGCCTCCC GTGCTGGACT CCGACGGCTC CTTCTTCTTC

```

-continued

1321 TATTCCAAGC TCACCGTGGA CAAGAGCAGG TGGCAGCAGG GGAACGTCTT CTCATGCTCC

1381 GTGATGCATG AGGCTOTGCA CAACCACTAC ACGCAGAAGA GCCTCTCCCT GTCTCCGGGC

1441 AAATGATAGG TTTAAACCGA ATTC

[1700] Non-coding sequences are italicized. DNA coding for mskappa ss is underlined with a solid line. DNA coding for the variable domain of anti-SARS-COV-2 HC is in bold font. DNA coding sequence for the human IgG1 constant

domain is in normal font. DNA coding sequence for the translational stop codons are italicized and underlined with a solid line.

[1701] DNA Sequence for Encoding the mAb 555 LC

(SEQ ID NO: 5382)

```

1  AAGCTTGCTC GAGCCACCAT GGAGACAGAC ACACTCCTGC TATGGGTACT GCTGCTCTGG
-----
61  GTTCCAGGAT CTACTGGCGA CATCCAGATG ACCCAGTCTC CATCCTCCCT GTCTGCATCT
-----
121  GTAGGAGACA GAGTCACCAT CACTTGCCGG GCAAGTCAGA GCATTAGCAG CTATTTAAGT
181  TGGTATCAGC AGAAACCAGG GAAAGCCCTT AAGCTCCTGA TCTATGCTGC ATCCAGTTTG
241  CAAAGTGGGG TCCCATCAAG GTTCAGTGGC AGTGGATCTG GGACAGATTT CACTCTCACC
301  ATCACCAGTC TGCAACCTGA AGATTTTGCA ACTTACTACT GTCAACAGAG TTACAGTACC
361  CCTCGCACGT TCGGCCAAGG GACCAAGGTG GAAATCAAAA GAACTGTGGC GGCGCCATCT
421  GTCTTCATCT TCCGCCATC TGATGAGCAG TTGAAATCCG GAACTGCCTC TGTGTGTGTC
481  CTGCTGAATA ACTTCTATCC CAGAGAGGCC AAAGTACAGT GGAAGGTGGA TAACGCCCTC
541  CAATCGGGTA ACTCCAGGA GAGTGTACA GAGCAGGACA GCAAGGACAG CACCTACAGC
601  CTCAGCAGCA CCCTGACGCT GAGCAAAGCA GACTACGAGA AACACAAAGT CTACGCCTGC
661  GAAGTCACCC ATCAGGGCCT GAGCTCGCCC GTCACAAAGA GCTTCAACAG GGGAGAGTGC
721  TAATAGGTTT AAACCGAATT C

```

[1702] Non-coding sequences are italicized. DNA coding sequence for the mskappa ss is underlined with a solid line. DNA coding sequence for the variable domain is in bold font. DNA coding sequence for the human kappa constant domain is in normal font. DNA coding sequence for the translational stop codons are italicized and underlined with a solid line.

[1703] Deduced Mature Amino Acid Sequence for mAb 555 HC (SEQ ID NO: 5363)

```

1  QVQLVQSGAE VKKPGSSVKV SCKASGGTFS NYAISNVRQA PGQGLEWMR IIPILGIANY
61  AQKFQGRVTI TADKSTSTAY MELSSLRSED TAVYYCARGY YEARHYYYYY AMDVWGQTA
121  VTVSSASTKG PSVFPLAPSS KSTSGGTAAL GCLVKDYFPE PVTGWSNSGA LTSGVHTFPA
181  VLQSSGLYSL SSVVTVPSST LGTQTYICNV NHKPSNTKVD KRVEPKSCDK THTCPPCPAP
-----
241  ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV VVDVSHEDPE VKFNWYVDGV EVHNAKTKPR
301  EEQYNSTYRV VSVLTVLHQD WLNGKEYKCK VSNKALPAPI EKTISKAKGQ PREPQVYTLF

```

-continued

361 PSREEMTKNQ VSLTCLVKGF YPSDIAVEWE SNGQPENNYK TTPPVLDSDG SFFLYSKLTV

421 DKSRWQQGNV FSCSVMHEAL HNHYTQKSL S LSPGK

[1704] The variable domain is in bold font and the human IgG1 G1m3 constant chain is in normal font. The IgG1 hinge region is underlined. The CDR sequences are shown using borders.

[1705] Deduced Mature Amino Acid Sequence for mAb 555 LC (SEQ ID NO: 5364)

1 DIQMTQSPSS LSASVGRDVT ITC**RASQ**SI**S** SYLS**NY**Q**Q**KP GKAPKLLI**YA** ASS**LQ**S**SV**PS
61 **RF**SGSGSGTD FTLTITSLQP **EDF**AT**Y**Q**Q** SYST**PRT**FGQ G**TK**VEIKRTV AAPSVFI**FP**P
121 SDEQLKSGTA SVVCLLN**NY**F P**RE**AKVQWKV DNALQSGNSQ **ESVTE**QDSKD STYSL**S**STLT
181 LSKADYEKHK VYACEVTHQG LSSPVTKSFN RGEC

[1706] The variable domain is in bold font and the human kappa constant domain is in normal font. CDR sequences are shown using borders.

TABLE 11-2-continued

TABLE 11-1

mAb 555-HC and LC CDR sequences		
Sequence	SEQ ID NO	
CDR-H1 KASGGIFSNYAIS	5383	
CDR-H2 RIIPILGIANYAQKFQ Q	5384	
CDR-H3 ARGYYEARHHYYYYAMDV	5385	
CDR-L1 RASQSISSYLS	5386	
CDR-L2 YAASSLQ S	5387	
CDR-L3 QQSYST PRT	5388	

[1707] The CDR sequences of other exemplary antibodies are shown below in Table 11-2.

TABLE 11-2

	SEQ ID NO	
Internal Designation No.: 258		
CDR-H1 TVSGGSISSYYWS	5389	
CDR-H2 RIYTSGSTNYNPSLKS	5390	
CDR-H3 AAGYGSIDY	5391	
CDR-L1 RSSQSLHNSNGYNYLD	5392	
CDR-L2 YLGSNRAS	5393	
CDR-L3 MQALQTPRT	5394	

	SEQ ID NO	
Internal Designation No.: 291		
CDR-H1 AASGFTFSSYAMH	5395	
CDR-H2 VISYDGSNKYYADSVKG	5396	
CDR-H3 ARASGGSYFGGMDV	5397	
CDR-L1 TGTSSDVGGYNYVS	5398	
CDR-L2 YEVSNRPS	5399	
CDR-L3 SSYTSSSTLYV	5400	
Internal Designation No.: 292		
CDR-H1 AASGFTVSSNYMS	5401	
CDR-H2 VIYSGGSTYYADSVKG	5402	
CDR-H3 ARDLQGGGGP	5403	
CDR-L1 RASQGIRNDLG	5404	
CDR-L2 YAASSLQ S	5405	
CDR-L3 LQDYNYPRT	5406	
Internal Designation No.: 308		
CDR-H1 SVSGGSISSSYHWG	5407	
CDR-H2 SIYSGSTYYNPSLKS	5408	
CDR-H3 AGLRVVITFGGVIPKGGAFDI	5409	
CDR-L1 TGTSSDVGGYNYVS	5410	
CDR-L2 YDVSNRPS	5411	
CDR-L3 SSYTSSSTVV	5412	

TABLE 11-2-continued

		SEQ ID NO
<u>Internal Designation No.: 309</u>		
CDR-H1	AASGFTFSSYAMH	5413
CDR-H2	VISYDGSNKYYADSVKG	5414
CDR-H3	ARPYSGSYQSYFDY	5415
CDR-L1	RASQSVSSSYLA	5416
CDR-L2	YGASSRAT	5417
CDR-L3	QQYGSSPLT	5418
<u>Internal Designation No.: 361</u>		
CDR-H1	TVSGGSISSGGYYWS	5419
CDR-H2	YIYYSGSTYYNPSLKS	5420
CDR-H3	ATTMVRGVIRLDHYGMDV	5421
CDR-L1	TGTSSDVGGYNYVS	5422
CDR-L2	YEVSNRPS	5423
CDR-L3	SSYTSSSTLL	5424
<u>Internal Designation No.: 364</u>		
CDR-H1	TASGFTFGDYAMS	5425
CDR-H2	FIRSKAYGGTTEYAASVKG	5426
CDR-H3	TRFGIDYDIWGSYRYTTLFDY	5427
CDR-L1	RASQSISSWLA	5428
CDR-L2	YKASSLES	5429
CDR-L3	QQYNSYSYT	5430
<u>Internal Designation No.: 373</u>		
CDR-H1	TVSGGSISSYYWS	5431
CDR-H2	YIYYSGSTNYNPSLKS	5432
CDR-H3	ARGPDYYDFWSGYFYGMVDV	5433
CDR-L1	RASQSVSSSYLA	5434
CDR-L2	YGASSRAT	5435
CDR-L3	QQYGSSSLT	5436
<u>Internal Designation No.: 388</u>		
CDR-H1	AASGFTFNYSYAIH	5437
CDR-H2	VISYDGSNKYYADSVKG	5438
CDR-H3	ARGRGGYRSYFDY	5439
CDR-L1	RASQSVSSSYLA	5440
CDR-L2	YGASSRAT	5441
CDR-L3	QQYGSSPNT	5442

TABLE 11-2-continued

		SEQ ID NO
<u>Internal Designation No.: 408</u>		
CDR-H1	AASGFTFSSYAMH	5443
CDR-H2	VISYDGSNKYYADSVKG	5444
CDR-H3	AKGADTPHYSYDFLSVGYYYGMDV	5445
CDR-L1	RASQSISSYLN	5446
CDR-L2	YAASSFQS	5447
CDR-L3	QQSYSAPPT	5448
<u>Internal Designation No.: 414</u>		
CDR-H1	KASGYTFTSYMH	5449
CDR-H2	IINPSGGSTSYAQKFQG	5450
CDR-H3	ARDPPGRDFWSGYTGAPDYYYGMDV	5451
CDR-L1	RASQGISNYLA	5452
CDR-L2	YAASSLQS	5453
CDR-L3	QQYNSYPYT	5454
<u>Internal Designation No.: 417</u>		
CDR-H1	TVSGGSISSSSYYWG	5455
CDR-H2	SIYYSGSTYYNPSLKS	5456
CDR-H3	ARAPIMITFGGVTGHFDY	5457
CDR-L1	RASQSVSSSYLA	5458
CDR-L2	YGASSRAT	5459
CDR-L3	QQYGSSPLT	5460
<u>Internal Designation No.: 419</u>		
CDR-H1	KASGYTFTSYMH	5461
CDR-H2	IINPSGGSTSYAQKFQG	5462
CDR-H3	ARDWTQSSGYDYYYGLDV	5463
CDR-L1	SGDALPKQYAY	5464
CDR-L2	YKDSERPS	5465
CDR-L3	QSADSSGTYVV	5466
<u>Internal Designation No.: 442</u>		
CDR-H1	AASGFTFSSYAMH	5467
CDR-H2	VISYDGSNKYYADSVRG	5468
CDR-H3	ARPKGGSYSDAFDI	5469

TABLE 11-2-continued

		SEQ ID NO
CDR-L1	RASQSVSSSYLA	5470
CDR-L2	YGASSRAT	5471
CDR-L3	QQYGSSPQS	5472
<u>Internal Designation No.: 445</u>		
CDR-H1	KASGYTFTSYMH	5473
CDR-H2	IINPSGGSTSYAQKFQG	5474
CDR-H3	ARDPTEVGATSEYYYYGMDV	5475
CDR-L1	SGDALPKQYAY	5476
CDR-L2	YKDSEKPS	5477
CDR-L3	QSADSSGTYV	5478
<u>Internal Designation No.: 447</u>		
CDR-H1	AASGFTVSSNYMS	5479
CDR-H2	VIYSGGSTYYADSVKG	5480
CDR-H3	ARDKSSGSGP	5481
CDR-L1	RASQGIRNDLG	5482
CDR-L2	YAASSLQS	5483
CDR-L3	LQDYNYPRT	5484
<u>Internal Designation No.: 462</u>		
CDR-H1	AASGETFSSYGMH	5485
CDR-H2	VIWYDGNKYYADSVKG	5486
CDR-H3	AKDPTWFGELPSYYYYGMDV	5487
CDR-L1	RASQSISSWLA	5488
CDR-L2	YKASSLES	5489
CDR-L3	QQYNSYPIT	5490
<u>Internal Designation No.: 479</u>		
CDR-H1	AASGFTFSSYSMN	5491
CDR-H2	YISSSSSTIYYADSVKG	5492
CDR-H3	ARDLGARTPWDIVVPAAMDY	5493
CDR-L1	RASQSVSSSYLA	5494
CDR-L2	YGASSRAT	5495
CDR-L3	QQYGRSPNT	5496

TABLE 11-2-continued

		SEQ ID NO
<u>Internal Designation No.: 481</u>		
CDR-H1	AASGFTVSSNYMS	5497
CDR-H2	VIYSGGSTFYADSVKG	5498
CDR-H3	AREVAGTYDY	5499
CDR-L1	RASQGISSWLA	5500
CDR-L2	YAASSLQS	5501
CDR-L3	QQANSFPGGT	5502
<u>Internal Designation No.: 483</u>		
CDR-H1	TVSGGSISSYYWS	5503
CDR-H2	YIYSGSTNYNPSLKS	5504
CDR-H3	ARAPEEKSSEIGELVGWGFDP	5505
CDR-L1	SGDKLGDKYAC	5506
CDR-L2	YQDSKRPS	5507
CDR-L3	QAWDSSTVV	5508
<u>Internal Designation No.: 488</u>		
CDR-H1	AASGLYVSSNYMS	5509
CDR-H2	VIYSGGSTYYADSVKG	5510
CDR-H3	ARSPYGGNS	5511
CDR-L1	QASQDISNYLN	5512
CDR-L2	YDASNLET	5513
CDR-L3	QQYDNLPIIT	5514
<u>Internal Designation No.: 489</u>		
CDR-H1	AASGFTFSSYAMH	5515
CDR-H2	VISYDGSNKYYADSVKG	5516
CDR-H3	ARAGSGNYNWFDP	5517
CDR-L1	RASQTVSSNLV	5518
CDR-L2	YGASTRAT	5519
CDR-L3	QQYNNWPPYT	5520
<u>Internal Designation No.: 494</u>		
CDR-H1	AASGFTVSSNYMS	5521
CDR-H2	VIYSGGSTFYADSVKG	5522
CDR-H3	ARDSGDQLLDY	5523
CDR-L1	RASQGISSYLA	5524

TABLE 11-2-continued

		SEQ ID NO
CDR-L2	YAASTLQS	5525
CDR-L3	QQLNSYPPFT	5526
<u>Internal Designation No.: 506</u>		
CDR-H1	TFSGFSLSGSGVGVG	5527
CDR-H2	LIYWDDDKRYSPLKS	5528
CDR-H3	AHHSLSIFDY	5529
CDR-L1	TGTSSDVGDYNYVS	5530
CDR-L2	YEVSNRPS	5531
CDR-L3	SSYTSSSTV	5532
<u>Internal Designation No.: 511</u>		
CDR-H1	TVSGGSISSSYYWG	5533
CDR-H2	SIYYSGSTYYNPSLKS	5534
CDR-H3	ASEKVDWFSGGPYGM DV	5535
CDR-L1	TGTSSDVGSYNYVS	5536
CDR-L2	YEVSNRPS	5537
CDR-L3	SSYTSISTLV	5538
<u>Internal Designation No.: 540</u>		
CDR-H1	AASGFTFSNAWMS	5539
CDR-H2	HIKSKTDGGTTDYAAPVKG	5540
CDR-H3	TREPYYFDY	5541
CDR-L1	RASQSISSWLA	5542
CDR-L2	YKASSLES	5543
CDR-L3	QQYNSRYT	5544
<u>Internal Designation No.: 549</u>		
CDR-H1	AASGLTVSSNYMS	5545
CDR-H2	VIYSGGSTYYADSVKG	5546
CDR-H3	ARSPYGGNS	5547
CDR-L1	RTSQTIYNYLN	5548
CDR-L2	YAASSFQN	5549
CDR-L3	QQGYSTPLT	5550
<u>Internal Designation No.: 553</u>		
CDR-H1	KASGYTFTSYGIS	5551
CDR-H2	WISAYNGNTNYAQKLQG	5552
CDR-H3	ARDRGYAATFGVFDY	5553

TABLE 11-2-continued

		SEQ ID NO
CDR-L1	RASQSISSYLN	5554
CDR-L2	SAASSLQS	5555
CDR-L3	QQSYSTAF T	5556
<u>Internal Designation No.: 561</u>		
CDR-H1	AASGFTFSSYAMH	5557
CDR-H2	VISYDGSNKYYADSVKG	5558
CDR-H3	ARPLSGSYRSAFDI	5559
CDR-L1	RASQSVSSNLA	5560
CDR-L2	YGASTRAT	5561
CDR-L3	QQYNNWPPRT	5562
<u>Internal Designation No.: 562</u>		
CDR-H1	AASGFTFSSYAMH	5563
CDR-H2	VISYDGSNKYYADSVKG	5564
CDR-H3	ARASSGGYQGPFDP	5565
CDR-L1	TGTSSDVGGYNYVS	5566
CDR-L2	YDVSNRPS	5567
CDR-L3	SSYTSSTLLYV	5568
<u>Internal Designation No.: 585</u>		
CDR-H1	AASGFTFATYAMH	5569
CDR-H2	LISHDGSNKHYADSVKG	5570
CDR-H3	ARESLEAAAPPFDY	5571
CDR-L1	SGDKLGKEYAS	5572
CDR-L2	YQDRKRPS	5573
CDR-L3	QAWDSSNSVV	5574

Example 12

[1708] X-Ray Crystallography Analysis of Selected Anti-SARS-COV-2 Antibodies

[1709] X-ray crystallography analysis of anti-SARS-CoV-2 Fabs 555, 481 and 488 (all RBD binders) are performed as follows. A 12 mg/mL solution of 555 Fab-RBD complex was set up in vapor diffusion sitting drops at a ratio of 1:1 with a well solution of 100 mM sodium acetate pH 4.6 and 20% PEG 10 K. Crystals appeared within two days and were harvested on the 3rd day after the set up. 11.5 mg/mL solution of 481 Fab-RBD complex was set up in vapor diffusion sitting drops at a ratio of 1:1 with a well solution of 100 mM Tri-Sodium Citrate pH=5.8, and 14% PEG 4 K, and 10% 2-Propanol. Crystals appeared within two days and were harvested on the 4th day after the set up. 11.7 mg/mL solution of 488 Fab-RBD complex was set up in vapor diffusion sitting drops at a ratio of 1:1 with a well solution of 100 mM HEPES pH=7.7, and 8% PEG 3350, and 200

mM L-Proline. Crystals appeared within one day and grew to their full size within two days. They were harvested on the 3rd day after the set up. Crystals were flash-frozen in liquid nitrogen following 1-minute incubation in cryoprotectant solution containing 25% glycerol in mother liquor.

[1710] Diffraction data were collected at Lilly Research Laboratories Collaborative Access Team (LRL-CAT) and beamline at Sector 31 of the Advanced Photon Source at Argonne National Laboratory, Chicago, Ill. Crystals stored in liquid nitrogen were mounted on a goniometer equipped with an Oxford Cryosystems Cryostat maintained at a temperature of 100° K. The wavelength used was 0.9793 Å collecting 900 diffraction images at a 0.2 degree oscillation angle and 0.12 seconds exposure time on a Pilatus3 S 6M detector at a distance of 392 mm. The diffraction data were indexed and integrated using auto PROC/XDS and merged and scaled in AIMLESS from the CCP4 suite (Vonnrhein, *Acta Crystallogr D Biol Crystallogr* 67, 293-302, 2011; Kabsch, *Acta crystallographica. Section D, Biological crystallography* 66, 133-144, 2010; Evans, *Acta Crystallogr D Biol Crystallogr* 69, 1204-1214, 2013; Winn, *Acta crystallographica. Section D, Biological crystallography* 67, 235-242, 2011).

[1711] Non-isomorphous data readily yielded initial structures by Molecular Replacement using for the Fab portion crystal structures from the proprietary Eli Lilly structure database and for the SARS2 spike RBD the public domain structure with the access code 6yla [Huo et al. 2020; manuscript in preparation]. The initial structure coordinates for each dataset were further refined using Refmac5 (CCP4) applying isotropic temperature factors. Model building was performed with Coot (CCP4) and final structure validation with MolProbity (Chen, et al., *Acta Crystallogr D Biol Crystallogr* 66: 12-21, 2010) and CCP4 validation tools.

[1712] The crystal structures of Fabs 481, 488 and 555, in complex with the SARS-CoV-2 spike protein RBD are shown in FIG. 9A. Fabs 481 and 488 bind in a nearly identical fashion almost completely obscuring the ACE2 binding site. This binding site is expected to be accessible in the “up” conformation of the RBD. Fab 555 binds to a different epitope on RBD that is slightly overlapping with the ACE2 binding site (FIG. 9B). The epitope of Fab 555 (FIGS. 9C and 9D) is predicted to be fully accessible on both the “up” and the “down” conformation of the RBD. This prediction is confirmed by high resolution cryo-EM imaging in which complexes of three 555 Fabs bound to a single SARS-COV-2 spike protein are observed in both the up and down conformation (FIGS. 9E and 9F).

[1713] Detailed epitope analysis for Fabs 555, 481 and 488 are performed as follows. Molecular operating environment (MOE) software from Chemical Computing Group (CCG), is used to measure contact surface areas (using the molecular contact surface analysis SVL script) and generate figures. CCP4 contact software is used to identify the number of van der Waals and hydrogen bonding interactions (numbers in parentheses, respectively, in Tables 12.1, 12.2 and 12.3). A distance cutoff of 3.5 Å (angstroms) was used for hydrogen bonding interactions, and 4.5 Å for van der Waals contacts. SARS-COV-2 spike protein residues with at least one atom that is within 4.5 Å to any residue of the antibody/Fab is considered part of the epitope for the antibody/Fab.

[1714] As shown in Table 12.1, the epitope of Fab/mAb 555 comprises one or more of the following amino acid

residues of SARS-COV-2 spike protein: Y351, Y449, N450, L452, L455, F456, T470, I472, N481, G482, V483, E484, G485, F486, Y489, F490, L492, Q493, S494 (the residue positions correspond to SEQ ID NO: 5317).

[1715] As shown in Table 12.2, the epitope of Fab/mAb 481 comprises one or more of the following amino acid residues of SARS-COV-2 spike protein: R403, D405, R408, Q409, T415, G416, K417, D420, Y421, Y453, L455, F456, R457, K458, S459, N460, Y473, Q474, A475, G476, F486, N487, Y489, Q493, S494, Y495, G496, Q498, T500, N501, G502, Y505 (the residue positions correspond to SEQ ID NO: 5317).

[1716] As shown in Table 12.3, the epitope of Fab/mAb 481 comprises one or more of the following amino acid residues of SARS-COV-2 spike protein: R403, T415, G416, K417, D420, Y421, L455, F456, R457, K458, S459, N460, Y473, Q474, A475, G476, S477, F486, N487, Y489, N501, G502, Y505 (the residue positions correspond to SEQ ID NO: 5317).

TABLE 12.1

Interactions between SARS-COV-2 spike protein residues and Fab 555 residues		
Residues of SARS-COV-2 spike protein	Residues of Fab 555: HC; LC	% dSA*
Y351	L55 (1)	36.2
Y449	S30 (1), N31 (15)	49.3
N450	K74 (1)	13.4
L452	L55 (3)	53.3
L455	A103 (1)	47
F456	A103 (2)	25.8
T470	L55 (1), I57 (5)	56.6
I472	R50 (1)	18.6
N481	T94 (4)	34.4
G482	R50 (2, 1), N59 (7, 2)	55.5
V483	W47 (2), R50 (1), N59 (2)	87.2
E484	R50 (7, 3), I52 (2), Y101 (6), Y110 (14), R96 (5, 1)	98.6
G485	Y110 (11), S91 (1), Y32 (5), Y92 (3)	97.8
F486	Y32 (17), Y92 (28, 1)	65.9
Y489	Y110 (2), Y32 (2)	37.6
F490	I52 (5), I57 (2), L55 (2), Y101 (8)	90.7
L492	L55 (1)	72.2
Q493	A103 (4, 1), E102 (15), R104 (7, 1)	81.4
S494	E102 (4, 2), N31 (1, 1)	65.2
Total:	224 (211 vdW, 13 H-bonds)	583 Å ²

*% dSA is percentage of the sidechain total surface area, which indicates the change in solvent exposed surface area. For each residue of the Fab, the number of specific atom-atom contacts is shown in parentheses, including both van der Waal (vdW), and hydrogen bonds (Hbond).

TABLE 12.2

Interactions between SARS-COV-2 spike protein residues and Fab 481 residues		
Residues of SARS-COV-2 spike protein	Residues of Fab 481: HC; LC	% dSA*
R403	F94 (2), N92 (7, 2); S93 (1)	56.6
D405	F94 (1)	10.9
R408	F94 (3)	10.6
Q409	F94 (2)	21
T415	S56 (5), F58 (8)	49
G416	Y52 (3), F58 (5)	100
K417	Y33 (5), Y52 (8)	44.9
D420	Y52 (2), S56 (4, 2)	59.7
Y421	Y33 (4), Y52 (8), P53 (4), G54 (6, 1)	73.3

TABLE 12.2-continued

Interactions between SARS-COV-2 spike protein residues and Fab 481 residues		
Residues of SARS-COV-2 spike protein	Residues of Fab 481: HC; LC	% ASA*
Y453	W32 (2)	27.9
L455	Y33 (10, 1), A100 (3)	75.7
F456	Y33 (4), V99 (1)	50.8
R457	P53 (4)	19.7
K458	S31 (4), S31 (4), G54 (3), P53 (3, 1)	33.2
S459	G54 (1)	12.3
N460	G54 (6)	36
Y473	S31 (9, 1), P53 (2)	47.8
Q474	S31 (1)	11.1
A475	F27 (3), T28 (6, 1), S31 (3), N32 (6, 1), R97 (1)	96.2
G476	F27 (2), T28 (4)	52.8
F486	V2 (3), G26 (2), R97 (3), Y105 (6)	31.9
N487	G26 (2, 1), F27 (4), R97 (3, 2)	63.4
Y489	R97 (3, 1), V99 (6)	52.8
Q493	A100 (2), W32 (4)	42.7
S494	W32 (3)	27.3
Y495	W32 (4, 1)	30.7
G496	S30 (2, 1)	41.3
Q498	S30 (12, 1), S67 (3, 2)	56.8
T500	G28 (3, 1), G68 (1)	27.2
N501	G28 (12), I29 (1), S30 (3, 1)	53.9
G502	G28 (4, 2), Q27 (7)	88
Y505	G28 (2), I2 (5), I29 (2), N92 (11), Q90 (5, 1), S93 (2)	66.8
Total	vdW: 272; H-bonds: 24	815 Å ²

*% ASA is percentage of the sidechain total surface area, which indicates the change in solvent exposed surface area. For each residue of the Fab, the number of specific atom-atom contacts is shown in parentheses, including both van der Waal (vdW), and hydrogen bonds (Hbond).

TABLE 12.3

Interactions between SARS-COV-2 spike protein residues and Fab 488 residues		
Residues of SARS-COV-2 spike protein	Residues of Fab 488: HC; LC	% ASA*
R403	D92 (3, 2), N93 (2, 1)	46.7
T415	S56 (4), Y58 (8, 2)	46
G416	Y52 (2), Y58 (4, 1)	89.6
K417	Y33 (5), Y52 (6), Y100 (8, 1)	57.4
D420	Y52 (2), S56 (4, 2)	61
Y421	Y33 (3), Y52 (8), S53 (5, 1), S4 (6, 1)	72.7
L455	Y33 (6, 1), Y100 (11)	68.3
F456	Y33 (4), G101 (3), Y100 (4)	46.4
R457	S53 (2, 1)	23.2
K458	T28 (1), S30 (6, 2), S31 (5, 1), S53 (8, 1), G54 (3)	47
S459	G54 (3)	16.3
N460	G54 (7)	37.9
Y473	S31 (9, 1), S53 (1)	45.8
Q474	S31 (1)	12.1
A475	L27 (3), T28 (6, 1), S31 (3), N32 (7, 1), R97 (1)	96.8
G476	L27 (2), T28 (6, 1)	56.3
S477	T28 (1)	12.9
F486	V2 (2), R97 (2)	21.3
N487	G26 (2, 1), L27 (3), R97 (3, 2)	60.9
Y489	N32 (1), R97 (3, 2), G101 (5), G102 (2, 1)	52.1
N501	D28 (4)	23.1

TABLE 12.3-continued

Interactions between SARS-COV-2 spike protein residues and Fab 488 residues		
Residues of SARS-COV-2 spike protein	Residues of Fab 488: HC; LC	% ASA*
G502	D28 (6, 1)	64.6
Y505	D28 (9), I29 (8), S30 (5), Y32 (5), D92 (5, 1), N93 (6, 1)	72
Total	vdW: 244; H-bonds: 30	671 Å ²

*% ASA is percentage of the sidechain total surface area, which indicates the change in solvent exposed surface area. For each residue of the Fab, the number of specific atom-atom contacts is shown in parentheses, including both van der Waal (vdW), and hydrogen bonds (Hbond).

Example 13

[1717] Rhesus Macaque Model of SARS-CoV-2 Infection

[1718] To assess the ability of mAb 555 (also known as LY-CoV555) to protect from viral challenge, rhesus macaques were intravenously (IV) administered with 1 mg/kg, 2.5 mg/kg, 15 mg/kg, or 50 mg/kg of mAb 555 or 50 mg/kg of an IgG1 control antibody 24 hours prior to virus challenge. Rhesus macaques were inoculated intranasally and intratracheally with a total of 1.1×10^5 TCID₅₀ of SARS-CoV-2 (USA-WA1/2020) and were monitored by twice daily cage-side observations, and respiratory exams throughout the study. Respiratory and clinical signs of disease in the macaques were limited, and gross necropsy observations collected at Day 6 were mild. Bronchoalveolar lavage (BAL) fluid, nasal and throat swabs were collected on Days 1, 3, and 6 after viral challenge (study Day 0). Viral genomes (gRNA) and subgenomic RNA (sg mRNA), indicative of active viral replication, were detectable in BAL, throat swabs, and nasal swabs for all animals following intranasal and intratracheal inoculation with SARS-CoV-2 (FIGS. 10A-10H).

[1719] Substantial reduction in viral loads was demonstrated in the upper and lower respiratory tracts of all LY-CoV555-treated groups (FIGS. 10A, 10C, 10E, 10G and Table 13.1). In the BAL and throat swabs, LY-CoV555 treatment resulted in 1.5 to 5-log reduction in gRNA 1-day post inoculation, with significance (q-value <0.05) at 15 mg/kg (BAL) and 2.5, 15, and 50 mg/kg (throat). The 2.5, 15, and 50 mg/kg doses of LY-CoV555 demonstrated significant reductions (q-value <0.05) in BAL and nasal gRNA on Days 3 and 6. Consistent with the BAL gRNA, Day 6 lung tissue also demonstrated significant (q-value <0.05) reduction in gRNA at doses of LY-CoV555 of 2.5 mg/kg and above. Overall reductions in gRNA were dose-related, with maximal protection observed at doses of 2.5 mg/kg and above. In addition to viral load determination, BAL samples from Days 3 and 6 were subjected to whole genome sequencing by next generation sequencing. Of the samples with sufficient coverage to identify genomic variants (N=12), no sequence alterations were detected in the LY-CoV555 or RBD domains compared to the reference (Table 13.1).

[1720] Subgenomic RNA analysis demonstrated profound impact of LY-CoV555 treatment on viral replication (FIGS. 10B, 10D, 10F, 10H and Table 13.1). In Day 1 BAL, throat, and nasal samples, 1-4 log reduction in sgRNA was seen for all LY-CoV555 treated groups, with significance (q-value <0.05) observed in all groups in the throat, and at 1, 2.5 and

15 or 50 mg/kg groups in BAL and nasal samples. Notably, by Day 3 sgRNA in the BAL and nasal samples was undetectable in animals dosed with 2.5-50 mg/kg LY-CoV555, with significant 3-4 log reductions observed relative to control (q-value <0.05). At Day 6, significant reduction of sgRNA in the lung tissue was demonstrated (q-value<0.05), at 2.5 through 50 mg/kg dose levels. Overall, these data show potent neutralization of SARS-CoV-2 in a non-human primate model by mAb 555.

[1721] To support dose-response evaluation of the model, serum concentrations of LY-CoV555 were determined evaluated by ELISA. LY-CoV555 demonstrated sustained serum concentrations after IV administration, consistent with expected pharmacokinetics for human IgG in a non-human primate model (Table 13.2). The dose responsive concentrations of serum LY-CoV555 were consistent with the dose-related reductions in viral loads in the lungs, throat, and nasal passages. Importantly, serum concentrations of LY-CoV555 at doses of 2.5 mg/kg and higher were associated with maximal protection in this Rhesus macaques infection model, and clearly demonstrate potent in vivo activity at low mg/kg dose levels. The delayed impact on viral loads in nasal swabs could reflect slower distribution of antibody into the nasal epithelial lining fluid versus the lung or throat.

TABLE 13.1

Statistical analyses for impact of mAb 555 (LY-CoV555) on viral loads in SARS-CoV-2 challenged Rhesus macaques.											
		BAL		Throat Swab		Nasal Swab		Right Lung		Left Lung	
		Genomes q-val	sg mRNA q-val	Genomes q-val	sg mRNA q-val	Genomes q-val	sg mRNA q-val	Genomes q-val	sg mRNA q-val	Genomes q-val	sg mRNA q-val
Day 1	1 mg/kg	0.159	0.037	0.084	0.008	0.207	0.023				
	2.5 mg/kg	0.069	0.038	0.005	0.002	0.186	0.010				
	15 mg/kg	0.008	0.007	0.001	0.002	0.454	0.090				
	50 mg/kg	0.258	0.194	0.002	0.002	0.210	0.008				
Day 3	1 mg/kg	0.022	0.010	0.730	0.921	0.069	0.005				
	2.5 mg/kg	0.008	0.008	0.075	0.921	0.008	0.002				
	15 mg/kg	0.007	0.015	0.794	0.094	0.031	0.002				
	50 mg/kg	0.019	0.012	0.814	0.727	0.005	0.002				
Day 6	1 mg/kg	0.192	0.674	0.250	0.921	0.027	0.578	0.137	0.010	0.137	0.080
	2.5 mg/kg	0.007	0.317	0.036	0.921	0.005	0.418	0.005	0.004	0.015	0.028
	15 mg/kg	0.011	0.317	0.037	0.794	0.038	0.607	0.002	0.005	0.005	0.037
	50 mg/kg	0.028	0.455	0.054	0.864	0.002	0.603	0.005	0.008	0.006	0.031

q-values in bold represent values <0.05, and indicate statistical significance.

TABLE 13-2

Serum total human IgG concentrations and AUC0-6days following intravenous administration of mAb 555 or control IgG to Rhesus Macaques in SARS-CoV-2 challenge model.						
		Serum concentration (mcg/mL)				
		Day 0*	Day 1	Day 3	Day 6	AUC _{0-Day6} (mcg*hr/mL)
Group 1: 50 mg/kg IgG1 control (N = 4)	mean	667	348	164	88	57500
	stdev	115	71	52	32	8360
Group 2: 1 mg/kg LY-CoV555 (N = 4)	mean	15	13	10	8	1920
	stdev	3	3	2	1	356
Group 3: 2.5 mg/kg LY-CoV555 (N = 4)	mean	38	30	21	15	4310
	stdev	14	11	6	3	1380
Group 4: 15 mg/kg LY-CoV555 (N = 3)	mean	276	215	145	98	30900
	stdev	37	14	21	13	3190
Group 5: 50 mg/kg LY-CoV555 (N = 3)	mean	679	539	376	258	77800
	stdev	101	61	47	78	12000

Abbreviations: AUC0-Day6 = area under the concentration time curve. N = number of animals per group.

*Day of viral challenge.

[1722] Methods for Rhesus Macaques Studies

[1723] The rhesus macaque model of SARS-CoV-2 infection was conducted according to the method of Chandrasekar et al., Science 2020. This study was approved by the Institutional Animal Care and Use Committee of BioQual Inc. in accordance with the animal welfare requirements and accreditations. Housing and handling of the

animals was performed in accordance with the standards of the AAALAC International's reference resource: the eighth edition of the *Guide for the Care and Use of Laboratory animals*, Animal Welfare Act as amended, and the 2015 reprint of the Public Health Service (PHS) Policy on Human Care and Use of Laboratory Animals. Handling of samples and animals was compliance with the Biosafety in Microbiological and Biomedical Laboratories (BMBL), 5th edition (Centers for Disease Control). Naïve female rhesus macaques of Indian origin (purpose bred, *Macaca mulatta* from PrimGen 8-12 years of age) were administered at 1, 2.5, 15 or 50 mg/kg of LY-CoV555 or 50 mg/kg of an IgG1 control antibody by slow intravenous bolus (N=3 or 4 animals per group). On study day 0 (one day following antibody administration), monkeys received a viral challenge of 1.1×10^5 PFU SARS-CoV-2 USA-WA1/2020 in 2 mL volume administered divided as 0.5 mL/nose (IN) and 1.0 mL intratracheally (IT). Live phase parameters were monitored pre-study through necropsy (Day 6). COVID specific observations were collected daily in conscious animals to monitor overall health and welfare and determine the need for veterinary intervention and/or euthanasia. COVID observations were scored on a scale of 0 to 10 and included measures of respiratory rate and dyspnea, overall appearance, activity, and responsiveness. Clinical observations were assessed cage-side twice daily and included evaluations of overall animal appearance, fecal consistency, and appetite. Body weights and rectal body temperatures were measured daily in anesthetized animals. At termination on

detection in the assay, a value of 25 ($\frac{1}{2}$ the limit of quantitation) was used for calculations. Serum samples were also collected for determination of LYCoV-555 concentrations by total human IgG ELISA assay.

Example 14

[1725] Characterization of Additional Anti-SARS-CoV2 Antibodies

[1726] Additional anti-SARS-CoV2 antibodies were characterized using the methods described in Examples 3-6, 9, and 10. The spike protein binding, ACE2 blocking, and binding affinity determination were performed using methods described in Example 4. The pseudoneutralization assay was performed using methods described in Example 6. Competitive binding assays were performed using methods similar to the epitope binning assays described in Example 4. The results are shown in Table 14-1.

[1727] The live virus neutralization assay was performed at Lab 4 using cultured Vero E6 cells infected with the SARS-CoV-2 virus clinical isolate USA/WA/1/2020. Pre-mixed solution of the SARS-CoV-2 virus and the tested anti-SARS-CoV-2 antibody were added to the Vero E6 cells and incubated at 37° C. for 24 hours to allow the non-neutralized SARS-CoV-2 virus to replicate. Decreased production of viral nucleocapsid protein as detected using standard immunostaining techniques. IC_{50} were calculated and the results are shown in Table 14-1.

TABLE 14-1

Activities of selected anti-SARS-CoV2 mAbs								
mAb ID	Spike Domain Bound	ACE2 Blocking?	Binding Affinity to WT Spike	Binding Affinity to D614G Spike	Pseudo-neutralization IC_{50} (μg/mL)	Live Virus Neutralization IC_{50} (μg/mL)	Compete with 555 for binding?	Compete with 488 for binding?
923	RBD	Yes	35 pM	22 pM	0.093	N/A	No	Yes
933	RBD	Yes	276 pM	29 nM	0.022	N/A	No	Yes
966	RBD	No	1.8 nM	1.3 nM	0.021	>10	No	No
989	RBD	No	104 pM	39 pM	0.19	1.5	No	No
1000	RBD	Yes	92 pM	16 pM	0.006	0.013	Yes	Yes
1009	RBD	Yes	168 pM	34 pM	0.17	N/A	No	No
1012	RBD	Yes	139 pM	27 pM	0.006	0.028	Yes	Yes
1015	RBD	Yes	96 pM	20 pM	0.011	0.032	Yes	No
1027	RBD	Yes	205 pM	43 pM	0.01	0.005	Yes	Yes
1037	RBD	Yes	126 pM	23 pM	0.025	0.009	No	No
1075	NTD	No	93 pM	23 pM	0.039	0.011	N/A	N/A
1081	RBD	No	93 pM	41 pM	0.012	4.6	No	No
1091	RBD	No	54 pM	19 pM	0.016	0.074	Yes	No
1098	RBD	No	173 pM	22 pM	0.087	N/A	No	No
1118	RBD	Yes	223 pM	44 pM	0.026	N/A	Yes	Yes
1130	NTD	Yes	94 pM	40 pM	0.114	2.3	N/A	N/A
1193	NTD	Yes	178 pM	41 pM	0.69	0.31	N/A	N/A
1203	RBD	Yes	302 pM	79 pM	0.02	N/A	No	Yes

study Day 6, macroscopic observations in the lung were evaluated by a board-certified veterinary pathologist.

[1724] Bronchioalveolar lavage (BAL), nasal and oral swabs were collected on days 1, 3 and 6, and lung tissue samples (at necropsy, day 6) were collected to assess genomic and subgenomic viral RNA via qRT PCR, conducted as reported (Chandrashekar et al, Science 2020). The lower limit of detection for genomic and sub-genomic RNA copies was 50. Where values were below the lower limit of

[1728] While this invention has been disclosed with reference to particular embodiments, it is apparent that other embodiments and variations of the inventions disclosed herein can be devised by others skilled in the art without departing from the true spirit and scope thereof. The appended claims include all such embodiments and equivalent variations.

SEQUENCE LISTING

The patent application contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<https://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US20210292393A1>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

1. An antibody or antigen-binding fragment thereof that specifically binds to a SARS-CoV-2 spike protein, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) three CDRs of a heavy chain variable region (VH) set forth as SEQ ID NO: 729 and three CDRs of a light chain variable region (VL) set forth as SEQ ID NO: 730; or
- (b) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 763 and three CDRs of a light chain variable region set forth as SEQ ID NO: 764; or
- (c) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 873 and three CDRs of a light chain variable region set forth as SEQ ID NO: 874; or
- (d) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 891 and three CDRs of a light chain variable region set forth as SEQ ID NO: 892; or
- (e) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 921 and three CDRs of a light chain variable region set forth as SEQ ID NO: 922; or
- (f) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 961 and three CDRs of a light chain variable region set forth as SEQ ID NO: 962; or
- (g) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 973 and three CDRs of a light chain variable region set forth as SEQ ID NO: 974; or
- (h) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 979 and three CDRs of a light chain variable region set forth as SEQ ID NO: 980; or
- (i) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 983 and three CDRs of a light chain variable region set forth as SEQ ID NO: 984; or
- (e) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1029 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1030; or
- (k) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1035 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1036; or
- (l) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1039 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1040; or
- (m) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1069 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1070; or
- (n) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1103 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1104; or
- (o) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1107 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1108; or

- (p) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1111 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1112; or
- (q) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1121 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1122; or
- (r) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1133 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1134; or
- (s) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1157 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1158; or
- (t) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1225 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1226; or
- (u) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1243 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1244; or
- (v) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1251 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1252; or
- (w) three CDRs of a heavy chain variable region set forth as SEQ ID NO: 1269 and three CDRs of a light chain variable region set forth as SEQ ID NO: 1270.

2. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody or fragment thereof comprises:

- (a) CDR-H1 comprising residues 31-35 of the VH, CDR-H2 comprising residues 50-65 of the VH, and CDR-H3 comprising residues 95-102 of the VH; and
- (b) CDR-L1 comprising residues 24-34 of the VL, CDR-L2 comprising residues 50-56 of the VL, and CDR-L3 comprising residues 89-97 of the VL; and

wherein the CDR numbering is according to Kabat.

3. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody or fragment thereof comprises:

- (a) CDR-H1 comprising residues 26-32 of the VH, CDR-H2 comprising residues 50-58 of the VH, and CDR-H3 comprising residues 95-102 of the VH; and
- (b) CDR-L1 comprising residues 24-34 of the VL, CDR-L2 comprising residues 50-56 of the VL, and CDR-L3 comprising residues 89-97 of the VL; and

wherein the CDR numbering is according to Chothia.

4. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody or fragment thereof comprises:

- (a) CDR-H1 comprising residues 30-35 of the VH, CDR-H2 comprising residues 47-58 of the VH, and CDR-H3 comprising residues 93-101 of the VH; and

- (b) CDR-L1 comprising residues 30-36 of the VL, CDR-L2 comprising residues 46-55 of the VL, and CDR-L3 comprising the residues 89-96 of the VL; and wherein the CDR numbering is according to MacCallum.

5. The antibody or antigen-binding fragment thereof of claim 1, comprising a heavy chain variable region having an amino acid sequence that is at least 90% identical to the heavy chain variable region sequence set forth in (a)-(x) and a light chain variable region having an amino acid sequence that is at least 90% identical to one of the corresponding light chain variable region sequences set forth in the same (a)-(x).

6. The antibody or antigen-binding fragment thereof of claim 1, wherein the antibody or antigen-binding fragment thereof comprises:

- (a) a heavy chain variable region set forth as SEQ ID NO: 729 and a light chain variable region set forth as SEQ ID NO: 730; or
- (b) a heavy chain variable region set forth as SEQ ID NO: 763 and a light chain variable region set forth as SEQ ID NO: 764; or
- (c) a heavy chain variable region set forth as SEQ ID NO: 873 and a light chain variable region set forth as SEQ ID NO: 874; or
- (d) a heavy chain variable region set forth as SEQ ID NO: 891 and a light chain variable region set forth as SEQ ID NO: 892; or
- (e) a heavy chain variable region set forth as SEQ ID NO: 921 and a light chain variable region set forth as SEQ ID NO: 922; or
- (f) a heavy chain variable region set forth as SEQ ID NO: 961 and a light chain variable region set forth as SEQ ID NO: 962; or
- (g) a heavy chain variable region set forth as SEQ ID NO: 973 and a light chain variable region set forth as SEQ ID NO: 974; or
- (h) a heavy chain variable region set forth as SEQ ID NO: 979 and a light chain variable region set forth as SEQ ID NO: 980; or
- (i) a heavy chain variable region set forth as SEQ ID NO: 983 and a light chain variable region set forth as SEQ ID NO: 984; or
- (j) a heavy chain variable region set forth as SEQ ID NO: 1029 and a light chain variable region set forth as SEQ ID NO: 1030; or
- (k) a heavy chain variable region set forth as SEQ ID NO: 1035 and a light chain variable region set forth as SEQ ID NO: 1036; or
- (l) a heavy chain variable region set forth as SEQ ID NO: 1039 and a light chain variable region set forth as SEQ ID NO: 1040; or
- (m) a heavy chain variable region set forth as SEQ ID NO: 1069 and a light chain variable region set forth as SEQ ID NO: 1070; or
- (n) a heavy chain variable region set forth as SEQ ID NO: 1103 and a light chain variable region set forth as SEQ ID NO: 1104; or
- (o) a heavy chain variable region set forth as SEQ ID NO: 1107 and a light chain variable region set forth as SEQ ID NO: 1108; or
- (p) a heavy chain variable region set forth as SEQ ID NO: 1111 and a light chain variable region set forth as SEQ ID NO: 1112; or

- (q) a heavy chain variable region set forth as SEQ ID NO: 1121 and a light chain variable region set forth as SEQ ID NO: 1122; or
- (r) a heavy chain variable region set forth as SEQ ID NO: 1133 and a light chain variable region set forth as SEQ ID NO: 1134; or
- (s) a heavy chain variable region set forth as SEQ ID NO: 1157 and a light chain variable region set forth as SEQ ID NO: 1158; or
- (t) a heavy chain variable region set forth as SEQ ID NO: 1225 and a light chain variable region set forth as SEQ ID NO: 1226; or
- (u) a heavy chain variable region set forth as SEQ ID NO: 1243 and a light chain variable region set forth as SEQ ID NO: 1244; or
- (v) a heavy chain variable region set forth as SEQ ID NO: 1251 and a light chain variable region set forth as SEQ ID NO: 1252; or
- (w) a heavy chain variable region set forth as SEQ ID NO: 1269 and a light chain variable region set forth as SEQ ID NO: 1270.

7. The antibody of claim 1, wherein the antibody comprises:

- (a) a heavy chain comprising SEQ ID NO: 5319 and a light chain comprising SEQ ID NO: 5320; or
- (b) a heavy chain comprising SEQ ID NO: 5321 and a light chain comprising SEQ ID NO: 5322; or
- (c) a heavy chain comprising SEQ ID NO: 5323 and a light chain comprising SEQ ID NO: 5324; or
- (d) a heavy chain comprising SEQ ID NO: 5325 and a light chain comprising SEQ ID NO: 5326; or
- (e) a heavy chain comprising SEQ ID NO: 5327 and a light chain comprising SEQ ID NO: 5328; or
- (f) a heavy chain comprising SEQ ID NO: 5329 and a light chain comprising SEQ ID NO: 5330; or
- (g) a heavy chain comprising SEQ ID NO: 5331 and a light chain comprising SEQ ID NO: 5332; or
- (h) a heavy chain comprising SEQ ID NO: 5333 and a light chain comprising SEQ ID NO: 5334; or
- (i) a heavy chain comprising SEQ ID NO: 5335 and a light chain comprising SEQ ID NO: 5336; or
- (j) a heavy chain comprising SEQ ID NO: 5337 and a light chain comprising SEQ ID NO: 5338; or
- (k) a heavy chain comprising SEQ ID NO: 5339 and a light chain comprising SEQ ID NO: 5340; or
- (l) a heavy chain comprising SEQ ID NO: 5341 and a light chain comprising SEQ ID NO: 5342; or
- (m) a heavy chain comprising SEQ ID NO: 5343 and a light chain comprising SEQ ID NO: 5344; or
- (n) a heavy chain comprising SEQ ID NO: 5345 and a light chain comprising SEQ ID NO: 5346; or
- (o) a heavy chain comprising SEQ ID NO: 5347 and a light chain comprising SEQ ID NO: 5348; or
- (p) a heavy chain comprising SEQ ID NO: 5349 and a light chain comprising SEQ ID NO: 5350; or
- (g) a heavy chain comprising SEQ ID NO: 5351 and a light chain comprising SEQ ID NO: 5352; or
- (r) a heavy chain comprising SEQ ID NO: 5353 and a light chain comprising SEQ ID NO: 5354; or
- (s) a heavy chain comprising SEQ ID NO: 5355 and a light chain comprising SEQ ID NO: 5356; or
- (t) a heavy chain comprising SEQ ID NO: 5357 and a light chain comprising SEQ ID NO: 5358; or

- (u) a heavy chain comprising SEQ ID NO: 5359 and a light chain comprising SEQ ID NO: 5360; or
- (v) a heavy chain comprising SEQ ID NO: 5361 and a light chain comprising SEQ ID NO: 5362; or
- (w) a heavy chain comprising SEQ ID NO: 5365 and a light chain comprising SEQ ID NO: 5366.
- 8.-15.** (canceled)
- 16.** The antibody of claim **1**, wherein the antibody has an IgG1m3 allotype.
- 17.** The antibody of claim **16**, wherein the antibody comprises an Fc region comprising N-glycoside-linked sugar chains bound to the Fc region, wherein the sugar chains do not contain fucose.
- 18.-23.** (canceled)
- 24.** The antibody or antigen-binding fragment thereof of claim **1**, which comprises at least one amino acid substitution.
- 25.** The antibody or antigen-binding fragment thereof of claim **24**, wherein the at least one amino acid substitution is a conservative substitution.
- 26.** The antibody or antigen-binding fragment thereof of claim **24**, wherein the at least one amino acid substitution is a substitution of an amino acid for a non-genetically encoded amino acid or a synthetic amino acid.
- 27.** The antibody or antigen-binding fragment thereof of claim **1**, which is conjugated to an immunomodulator, a cytokine, a cytotoxic agent, a chemotherapeutic agent, a diagnostic agent, an antiviral agent, an antimicrobial agent, or a drug.
- 28.** An antibody conjugate comprising the antibody or antigen-binding fragment thereof of claim **1** conjugated to an immunomodulator, a cytokine, a cytotoxic agent, a chemotherapeutic agent, a diagnostic agent, an antiviral agent, an antimicrobial agent, or a drug.
- 29.** A pharmaceutical composition comprising the antibody or antigen-binding fragment thereof of claim **1**, and one or more pharmaceutically acceptable carriers, diluents, or excipients.
- 30.** The pharmaceutical composition of claim **29**, wherein the pharmaceutical composition comprises at least one additional antibody that binds the SARS-CoV-2 spike protein.
- 31.** The pharmaceutical composition of claim **29**, wherein the pharmaceutical composition further comprises histidine, sodium chloride, sucrose, and polysorbate 80.
- 32.** The pharmaceutical composition of claim **29**, wherein the pharmaceutical composition has a pH of about 6.0.
- 33.** The pharmaceutical composition of claim **29**, wherein the pharmaceutical composition comprises 5 mM histidine, 50 mM NaCl, 6% sucrose, and 0.05% polysorbate 80 and has a pH of about 6.0.
- 34.** The pharmaceutical composition of claim **29**, wherein the antibody concentration in the pharmaceutical composition is about 35 mg/mL to about 125 mg/mL.
- 35.-37.** (canceled)
- 38.** A nucleic acid encoding the heavy chain, the light chain, or both the heavy chain and the light chain, of the antibody of claim **1**.
- 39.** A vector comprising the nucleic acid of claim **38**.
- 40.** A host cell comprising the nucleic acid of claim **38**.
- 41.** A process for producing an antibody comprising:
 - (a) cultivating the host cell of claim **40** under conditions such that the antibody is expressed; and
 - (b) recovering the expressed antibody.
- 42.** An article of manufacture useful for diagnosing or treating a SARS-Co-V or SARS-Co-V-2-linked disease comprising a receptacle comprising the antibody or antigen-binding fragment thereof of claim **1** and instructional materials for using the same to treat or diagnose the SARS-Co-V or SARS-Co-V-2-linked disease.
- 43.** A method of identifying a SARS-Co-V or SARS-CoV-2-infected cell comprising:
 - (a) contacting a cell with the antibody or antigen-binding fragment thereof of claim **1**, which is conjugated to a detectable agent; and
 - (b) detecting specific binding of the antibody or antigen-binding fragment thereof to the cell.
- 44.** A method of diagnosing a SARS-Co-V or SARS-CoV-2 infection in a patient comprising:
 - (a) contacting a sample obtained from a patient with the antibody or antigen-binding fragment thereof of claim **1**, which is conjugated to a detectable agent; and
 - (b) detecting specific binding of the antibody or antigen-binding fragment thereof to a SARS-Co-V or SARS-CoV-2 antigen present in the sample.
- 45.** A method of treating or preventing a SARS-Co-V or SARS-Co-V-2-linked disease comprising administering to a patient a therapeutically effective amount of the antibody or antigen-binding fragment thereof of claim **1**.
- 46.** A method of treating or preventing COVID-19 comprising administering to a patient a therapeutically effective amount of the antibody or antigen-binding fragment thereof of claim **1**.
- 47.** A method of treating or preventing COVID-19 comprising:
 - (a) contacting a sample obtained from a patient with the antibody or antigen-binding fragment thereof of claim **1**, conjugated to a detectable agent;
 - (b) detecting specific binding of the antibody or antigen-binding fragment thereof to a SARS-CoV-2 antigen present in the sample; and
 - (c) administering to the patient a therapeutically effective amount of the antibody or antigen-binding fragment thereof of claim **1**.
- 48.** The method of claim **46**, wherein the antibody is administered to the patient intravenously or subcutaneously at about 700 mg to about 7000 mg.
- 49.** (canceled)
- 50.** The method of claim **46**, wherein the method further comprises administering to the patient another antibody that binds the SARS-CoV-2 S protein.
- 51.-56.** (canceled)
- 57.** A method of testing an anti-coronavirus vaccine, comprising:
 - (a) contacting a sample of an anti-coronavirus vaccine with the antibody or antigen-binding fragment thereof of claim **1**, conjugated to a detectable agent; and
 - (b) detecting specific binding of the antibody or antigen-binding fragment thereof to the anti-coronavirus vaccine present in the sample;
 wherein the anti-coronavirus vaccine comprises a coronavirus subunit or fragment thereof.