

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/071212 A2

(43) International Publication Date

8 May 2014 (08.05.2014)

WIPO | PCT

(51) International Patent Classification:

*C07K 16/24* (2006.01) *A61K 9/00* (2006.01)  
*C07K 16/28* (2006.01) *A61K 9/19* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/US2013/068110

(22) International Filing Date:

1 November 2013 (01.11.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/721,364 1 November 2012 (01.11.2012) US  
61/794,231 15 March 2013 (15.03.2013) US

(71) Applicant: **ABBVIE INC.** [US/US]; 1 North Waukegan Road, North Chicago, Illinois 60064 (US).

(72) Inventors: **SIEDLER, Michael**; Brueckenkopfstrasse 1/1, 69120 Heidelberg (DE). **KUMAR, Vineet**; 1325 Skiles Blvd., West Chester, Pennsylvania 19382 (US). **CHARI, Ravi**; 2 Clara Street, Apt. 2, Worcester, MA 01606 (US). **SALUJA, Sonal**; 322 Madison Place, Shrewsbury, MA 01545 (US). **PATIL, Vishwesh**; 72 Newport Street, Arlington, MA 02476 (US).

(74) Agents: **COWLES, Cristin, H.** et al.; McCarter & English, LLP, 265 Franklin Street, Boston, MA 02110 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: STABLE DUAL VARIABLE DOMAIN IMMUNOGLOBULIN PROTEIN FORMULATIONS




Fig. 1

(57) Abstract: The disclosure provides stable aqueous formulations comprising an Aqueous Stable Dual Variable Domain Immuno-globulin (AS-DVD-Ig) protein. The disclosure also provides stable lyophilized formulations comprising a Lyophilized Stable Dual Variable Domain Immunoglobulin (LS-DVD-Ig) protein.

WO 2014/071212 A2

## **STABLE DUAL VARIABLE DOMAIN IMMUNOGLOBULIN PROTEIN FORMULATIONS**

### **RELATED APPLICATIONS**

This application claims the benefit of priority to US Provisional Appln. No. 61/721364, filed on November 1, 2012. This application also claims the benefit of priority to US Provisional Appln. No. 61/794231, filed on March 15, 2013. The contents of both the priority applications are hereby incorporated by reference.

### **BACKGROUND OF THE INVENTION**

A basic principle of pharmaceutical protein formulations is that certain instabilities, *e.g.*, chemical instability and physical instability, must be overcome. Chemical instabilities often lead to the modification of the protein through bond formation or cleavage. Examples of problems associated with chemical instability include deamidation, racemization, hydrolysis, oxidation, beta elimination and disulfide exchange. While physical instabilities do not lead to covalent changes in proteins, they are just as problematic and difficult to overcome. Physical instabilities involve changes in the higher order structure (secondary and above) of proteins, which can result in denaturation, adsorption to surfaces, aggregation, and/or precipitation (Manning *et al.* (1989) *Pharm. Res.* 6:903). For therapeutic proteins, chemical and physical instabilities can create significant challenges in formulating the protein for delivery to a patient. Aggregation is often considered the most common type of physical instability. For example, exposure to hydrophobic interfaces fosters physical instability by alignment of protein molecules at the interface, unfolding the protein and maximizing exposure of hydrophobic residues to air, and initiating aggregation.

Highly concentrated protein formulations, especially those in liquid form, are often desirable for therapeutic purposes since they allow for dosages with smaller volumes, and provide for the possibility of subcutaneous delivery. The development of high protein concentration formulations, however, presents many challenges. For example, a high protein concentration often results in increased protein aggregation, insolubility and degradation (for review, see Shire *et al.* (2004) *J. Pharm. Sci.* 93:1390).

To date, the majority of approved therapeutic proteins are antibodies. The development of commercially viable antibody pharmaceutical formulations has not, however, been straightforward despite the fact that antibodies generally have the same structure (see Wang *et al.* (2007) *J. Pharm. Sci.* 96:1). Concentration dependent aggregation is considered one of the greatest challenges in formulating antibodies (see Shire *et al.* (2004) *J. Pharm. Sci.* 93:1390).

Dual Variable Domain Immunoglobulin (DVD-Ig<sup>TM</sup>) proteins are multivalent binding proteins that are engineered to combine the function and specificity of two monoclonal antibodies into one molecular entity (See Wu *et al.*, US Patent No. 7,612,181). Given the multivalent nature of DVD-Ig proteins, these molecules hold tremendous promise as therapeutics. However, DVD-Ig proteins present a new formulation challenge given their much larger size (approximately 200 kDa) and complexity, compared to antibodies.

## SUMMARY OF THE INVENTION

While the majority of Dual Variable Domain Immunoglobulin (DVD-Ig<sup>TM</sup>) proteins are prone to destabilization (*e.g.*, aggregation) in aqueous formulations, a subset of DVD-Ig proteins can be stably formulated. Some DVD-Ig proteins in the aqueous state suffer from certain problems, such as aggregation and/or fragmentation of the DVD-Ig protein monomer. Unpredictably, a subset of DVD-Ig proteins can be stably formulated in the aqueous state even at high concentrations. Such stable DVD-Ig proteins are referred to herein as Aqueous Stable Dual Variable Domain Immunoglobulin proteins or “AS-DVD-Ig” proteins.

In certain embodiments, the disclosure provides stable aqueous formulations comprising AS-DVD-Ig proteins, including high concentration AS-DVD-Ig formulations. AS-DVD-Ig proteins are a subpopulation of DVD-Ig proteins characterized by their ability to remain stable, *e.g.*, in concentrations 50 mg/ml or greater, during storage (*e.g.*, exhibit an aggregation increase of less than 3 % (determined by size exclusion chromatography (SEC)) following accelerated storage (40°C) in an aqueous formulation at a concentration of at least 50 mg/ml). In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 15% loss in relative percentage monomers as determined by size exclusion chromatography (SEC) when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 °C. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about

10% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 °C. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 5% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 °C. In certain embodiments, an AS-DVD-Ig protein is characterized as having less than 10% aggregation as determined by SEC when formulated in a citrate phosphate buffer at a concentration of at least 50 mg/ml following 14 days of storage at 40°C. In certain embodiments, the AS-DVD-Ig protein is characterized as having 6% or less aggregation as determined by SEC following 14 days of storage at 40°C, wherein the AS-DVD-Ig protein at a concentration of at least 50 mg/ml is stored in a citrate phosphate buffer or a histidine buffer. In certain embodiments, the AS-DVD-Ig protein is characterized as having 5% or less aggregation as determined by SEC following 14 days of storage at 40°C, wherein the AS-DVD-Ig protein at a concentration of at least 50 mg/ml is stored in a citrate phosphate buffer or a histidine buffer. In certain embodiments, the AS-DVD-Ig protein is characterized as having 4% or less aggregation as determined by SEC following 14 days of storage at 40°C, wherein the AS-DVD-Ig protein at a concentration of at least 50 mg/ml is stored in a citrate phosphate buffer or a histidine buffer. In certain embodiments, the AS-DVD-Ig protein is characterized as having 3% or less aggregation as determined by SEC following 14 days of storage at 40°C, wherein the AS-DVD-Ig protein at a concentration of at least 50 mg/ml is stored in a citrate phosphate buffer or a histidine buffer. In certain embodiments, the AS-DVD-Ig protein is characterized as having 2% or less aggregation as determined by SEC following 14 days of storage at 40°C, wherein the AS-DVD-Ig protein at a concentration of at least 50 mg/ml is stored in a citrate phosphate buffer or a histidine buffer. In certain embodiments, the AS-DVD-Ig protein is characterized as having 1% or less aggregation as determined by SEC following 14 days of storage at 40°C, wherein the AS-DVD-Ig protein at a concentration of at least 50 mg/ml is stored in a citrate phosphate buffer or a histidine buffer. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having about a 10% relative (rel.) peak area or less change in monomers at about 40 °C after 21 days of storage at a concentration of about 100 mg/ml in an aqueous formulation at a pH between about 5.5 to about 6.5. In other embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having about a 1% rel. peak area or less change in monomers at about 5 °C after 21 days of storage at a concentration of about 100 mg/ml at a pH between about 5.5 to about 6.5 in an aqueous

formulation. Examples of AS-DVD-Ig proteins that may be included in the formulations of the disclosure include, but are not limited to, an AS-DVD-Ig protein having binding specificity for IL4 and IL13; IL1 $\alpha$  and IL1 $\beta$ ; TNF $\alpha$  and IL17; and DLL4 and VEGF.

In certain embodiments, the formulation comprises less than about 10% aggregate AS-DVD-Ig protein. In certain embodiments, the formulation comprises less than about 9% aggregate AS-DVD-Ig protein. In certain embodiments, the formulation comprises less than about 8% aggregate AS-DVD-Ig protein. In certain embodiments, the formulation comprises less than about 7% aggregate AS-DVD-Ig protein. In certain embodiments, the formulation comprises less than about 6% aggregate AS-DVD-Ig protein. In certain embodiments, the formulation comprises less than about 5% aggregate AS-DVD-Ig protein. In certain embodiments, the formulation comprises less than about 4% aggregate AS-DVD-Ig protein. In a further embodiment, the formulation comprises less than about 3% aggregate AS-DVD-Ig protein. Aggregation may be determined by SEC analysis.

In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having a about 10% relative (rel.) peak area or less change in monomers at about 40 °C after 21 days of storage at a concentration of about 100 mg/ml in an aqueous formulation at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having a about 1% rel. peak area or less change in monomers at about 5 °C after 21 days of storage at a concentration of about 100 mg/ml at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0, in an aqueous formulation.

In certain embodiments, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein and a buffer. For example, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein and a buffer having a molarity of about 5 to about 50 mM, wherein the formulation has a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 6.5. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 15% loss in relative percentage monomers as determined by size exclusion chromatography (SEC) when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 degrees C. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 10% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at

about 40 degrees C. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 5% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 degrees C. In certain embodiments, the formulation comprises about 6% or less aggregation as determined by SEC analysis. In other embodiment, the formulation comprises about 5% or less aggregation as determined by SEC analysis. In certain embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having about a 10% relative (rel.) peak area or less change in monomers at about 40 °C after 21 days of storage at a concentration of about 100 mg/ml in an aqueous formulation at a pH between about 5.5 to about 6.5. In other embodiments, the AS-DVD-Ig protein is characterized as a DVD-Ig protein having about a 1% rel. peak area or less change in monomers at about 5 °C after 21 days of storage at a concentration of about 100 mg/ml at a pH between about 5.5 to about 6.5 in an aqueous formulation. In certain embodiments, the formulation comprises about 1 to about 250 mg/ml, about 1 to 200 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, or about 80 to about 105 mg/ml of the AS-DVD-Ig protein. In certain embodiments, the buffer used in the formulation is acetate, histidine, glycine, arginine, phosphate, citrate, or citrate / phosphate buffer. In certain embodiments, the molarity of the buffer ranges from 5 to 50 mM, *e.g.*, 10 to 20 mM.

In certain embodiments, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein, a buffer and a surfactant. For example, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein, a buffer having a molarity of about 5 to about 50 mM, and a surfactant, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the formulation comprises about 1 to about 250 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, or about 80 to about 105 mg/ml of the AS-DVD-Ig protein. In certain embodiments, the surfactant is a polysorbate, *e.g.*, polysorbate 80 or polysorbate 20. In certain embodiments, the concentration of polysorbate in the formulation is about 0.001% to about 1%, about 0.005% to about 0.05%, about 0.01% to about 0.05%, or about 0.1%. In certain embodiments, the concentration of polysorbate 80 or polysorbate 20 in the formulation is about 0.005% to about 0.02%. In certain embodiments, the buffer used in the

formulation is acetate, histidine, glycine, arginine, phosphate, citrate, or citrate / phosphate buffer. In certain embodiments, the molarity of the buffer ranges from 5 to 50 mM, *e.g.*, 10 to 20 mM.

In certain embodiments, the disclosure includes an aqueous formulation comprising an AS-DVD-Ig protein, a buffer, and a polyol. For example, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein, a buffer having a molarity of about 5 to about 50 mM, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the formulation comprises about 1 to about 250 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, or about 80 to about 105 mg/ml of the AS-DVD-Ig protein. In certain embodiments, the polyol is sorbitol. In certain embodiments, the concentration of sorbitol in the formulation is about 20 to about 60 mg/ml sorbitol, about 25 to about 55 mg/ml, about 30 to about 50 mg/ml, or about 35 to about 45 mg/ml. In certain embodiments, the polyol is sucrose. In certain embodiments, the concentration of sucrose in the formulation is about 60 to about 100 mg/ml, about 65 to about 95 mg/ml, about 70 to about 90 mg/ml, or about 75 to about 85 mg/ml. In certain embodiments, the polyol is mannitol. In certain embodiments, the concentration of mannitol in the formulation is about 10 to about 100 mg/ml, or about 20 to about 80, about 20 to about 70, about 30 to about 60, or about 30 to about 50 mg/ml. In certain embodiments, the buffer used in the formulation is acetate, histidine, glycine, arginine, phosphate, citrate, or citrate / phosphate buffer. In certain embodiments, the molarity of the buffer ranges from 5 to 50 mM, *e.g.*, 10 to 20 mM.

In certain embodiments, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein, a buffer, a polyol, and a surfactant. For example, the disclosure provides an aqueous formulation comprising an AS-DVD-Ig protein, a buffer having a molarity of about 5 to about 50 mM, a surfactant, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the formulation comprises about 1 to about 250 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, or about 80 to about 105 mg/ml of the AS-DVD-Ig. In certain embodiments, the polyol is sorbitol. In certain embodiment, the sorbitol concentration in the formulation is about 20 to about 60 mg/ml, about 25 to about 55 mg/ml, about 30 to about 50 mg/ml, or about 35 to about 45 mg/ml. In certain embodiments, the

polyol is sucrose. In certain embodiments, the concentration of sucrose in the formulation is about 60 to about 100 mg/ml, about 65 to about 95 mg/ml, about 70 to about 90 mg/ml, or about 75 to about 85 mg/ml. In certain embodiments, the polyol is mannitol. In certain embodiments, the concentration of mannitol in the formulation is about 10 to about 100 mg/ml, or about 20 to about 80, about 20 to about 70, about 30 to about 60, or about 30 to about 50 mg/ml. In certain embodiments, the surfactant is a polysorbate, *e.g.*, polysorbate 80 or polysorbate 20. In certain embodiments, the concentration of polysorbate in the formulation is about 0.001% to about 1%, about 0.005% to about 0.05%, about 0.01% to about 0.05%, or about 0.1%. In certain embodiments, the concentration of polysorbate 80 or polysorbate 20 in the formulation is about 0.005% to about 0.02%. In certain embodiments, the buffer used in the formulation is acetate, histidine, glycine, arginine, phosphate, citrate, or citrate / phosphate buffer. In certain embodiments, the molarity of the buffer ranges from 5 to 50 mM, *e.g.*, 10 to 20 mM.

In certain embodiment, the disclosure provides a formulation comprising an AS-DVD-Ig protein, a polyol (*e.g.*, sorbitol, mannitol, or sucrose), a buffer (*e.g.*, acetate, histidine, glycine, arginine, phosphate, citrate, or citrate / phosphate), and a surfactant (*e.g.*, a polysorbate), wherein said formulation has a pH of about 5 to about 7, and wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 15% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of about 60 mg/ml following 14 days storage at about 40 degrees C. In certain embodiments, the polysorbate is polysorbate 80 or polysorbate 20. In certain embodiments, the concentration of polysorbate 80 or polysorbate 20 is about 0.005% to about 0.02%.

In certain embodiments, the disclosure provides a formulation comprising an AS-DVD-Ig protein, a polyol, histidine buffer, and a polysorbate, wherein said formulation has a pH of about 5 to about 7, and wherein the AS-DVD-Ig protein is characterized as having 6% aggregation or less as determined by SEC, where the AS-DVD-Ig protein is formulated in a citrate phosphate buffer or histidine buffer at a concentration of at least 60 mg/ml, following 14 days storage at 40° C.

In certain embodiments, the disclosure provides a formulation comprising an AS-DVD-Ig protein and a buffer having a molarity of about 5 to about 50 mM, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 10% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of about 60 mg/ml, following 14 days storage at about 40

degrees C, and the formulation has a pH of 4.5 to 7.5. In certain embodiments, the formulation further comprises a surfactant, a polyol, or combinations thereof.

The disclosure is also based, in part, on the surprising discovery that while the majority of DVD-Ig<sup>TM</sup> proteins are prone to destabilization in a lyophilized state, a subset of DVD-Ig proteins are able to be stably formulated in a lyophilized form. Such stable DVD-Ig proteins are referred to herein as Lyophilized Stable Dual Variable Domain Immunoglobulin proteins or “LS-DVD-Ig” proteins. Formulations wherein the DVD-Ig protein was in the lyophilized state suffered from certain problems such as aggregation and/or fragmentation of the DVD-Ig protein monomer. Unpredictably, a subset of DVD-Ig proteins can be stably formulated in the lyophilized state even at high concentrations.

In certain embodiments, the disclosure is a lyophilized formulation comprising a Lyophilized-Stable DVD Immunoglobulin (LS-DVD-Ig) protein, wherein when said formulation is reconstituted said formulation comprises about 1 to about 100 mg/ml of the LS-DVD-Ig protein, about 10 to about 50 mM of a buffer, a polyol, about 0.01 to about 0.2 mg/ml of a polysorbate, and has a pH of about 5 to about 7, *e.g.*, about 5.5 to about 6.5. In certain embodiments, the LS-DVD-Ig protein has more than 10% rel. peak area change in monomers observed, following accelerated storage at a pH between about 5.5 and about 6.5 in an aqueous formulation for 21 days at about 40 °C, when formulated at a concentration over about 100 mg/ml.

In certain embodiments, the LS-DVD-Ig protein has more than about 10% rel. peak area change in monomers observed, following accelerated storage at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0, in an aqueous formulation for 21 days at about 40 °C, when formulated at a concentration of about 100 mg/ml or more.

In certain embodiments, the disclosure provides a lyophilized formulation prepared by lyophilizing an aqueous formulation comprising a buffer have a molarity (*e.g.*, histidine, succinate or citrate and/or phosphate) of about 5 to about 50 mM, a surfactant, and a polyol (*e.g.*, mannitol, sorbitol, sucrose, or trehalose), wherein the formulation has a pH of about 4.5 to about 7.5, *e.g.*, about 5.5 to about 6.5.

In certain embodiments, the disclosure provides LS-DVD-Ig proteins or AS-DVD-Ig proteins that are stable in formulations having a pH of about 4.5 to about 7.5. In certain embodiments, the formulation has a pH of 5 to 6.5. In certain embodiments, the formulation

has a pH of about 5.7 to about 6.3. In certain embodiments, the formulation of the disclosure has a pH of about 5.5 to about 6.5. In certain embodiments, the formulation of the disclosure has a pH of 5.8 to about 6.2, or a pH of 6.

In certain embodiments, the disclosure provides a formulation comprising an AS-DVD-Ig protein or an LS-DVD-Ig protein, a polyol, histidine buffer, and a polysorbate, wherein said formulation has a pH of about 5 to about 7, and wherein the AS-DVD-Ig protein or an LS-DVD-Ig protein is characterized as having 15% aggregation or less as determined by SEC, where the AS-DVD-Ig protein or an LS-DVD-Ig protein is formulated in a citrate phosphate buffer or histidine buffer at a concentration of at least 60 mg/ml, following 14 days storage at 40°C. Such formulations may be either in a lyophilized or an aqueous state, as AS-DVD-Ig protein or an LS-DVD-Ig protein identified as having 15% aggregation or less as determined by SEC, where the AS-DVD-Ig protein or an LS-DVD-Ig protein is formulated in a citrate phosphate buffer or a histidine buffer at a concentration of at least 60 mg/ml, following 14 days storage at 40°C.

One advantage of the compositions of the disclosure is that the AS-DVD-Ig protein or LS-DVD-Ig protein can be stably formulated in liquid form at a high concentration. In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein has a concentration of about 1 to about 200 mg/ml. In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein has a concentration of about 20 to about 100 mg/ml. In certain embodiments, the formulation comprises about 1 to about 250 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, or about 80 to about 105 mg/ml of the AS-DVD-Ig protein or LS-DVD-Ig.

Examples of buffers that may be used in the formulations of the disclosure include, but are not limited to, acetate, histidine, glycine, arginine, phosphate, and citrate. In certain embodiments, the molarity of the buffer in the formulation is about 5 to about 50 mM. In certain embodiments, the buffer molarity is about 10 mM to about 20 mM.

Examples of polyols that may be used in the formulations of the disclosure include, but are not limited to, sorbitol, mannitol, and sucrose. In certain embodiments, the polyol is sorbitol. In certain embodiments, about 30 to about 50 mg/ml of sorbitol is used in the formulation. In certain embodiments, the polyol is sucrose. In certain embodiments, about

70 to about 90 mg/ml of sucrose is used in the formulation. In a further embodiment, the polyol is mannitol. In certain embodiments, about 30 to about 50 mg/ml of mannitol is used in the formulation.

Examples of surfactants that may be used in the formulations of the disclosure include, but are not limited to, polysorbates and poloxamers. In certain embodiments, the surfactant is a polysorbate, examples of which are polysorbate 80 and polysorbate 20. Other examples include poloxamer Pluronic F-68, albumin, lecithin, and cyclodextrins. In certain embodiments, the polysorbate has a concentration of about 0.05 mg/ml to about 2mg/ml. In a further embodiment, the polysorbate has a concentration of about 0.01 to about 0.2 mg/ml.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises first and second polypeptide chains, each independently comprising VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first variable domain; VD2 is a second variable domain; C is a constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site. In a further embodiment, the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, and wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 does not comprise an Fc region; n is 0 or 1, wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site. In certain embodiments, the two first polypeptide chains and two second polypeptide chains, wherein the binding protein comprises four functional target binding sites. In certain embodiments, X1 is not CL.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises a polypeptide chain comprising VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first variable domain, VD2 is a second variable domain, C is a constant domain, X1 represents an amino acid or polypeptide, X2 represents an Fc region and n is 0 or 1.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein used in the compositions and methods of the disclosure comprises four polypeptide chains, wherein two polypeptide chains comprise VD1-(X1) $n$ -VD2-C-(X2) $n$ , wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and two polypeptide chains comprise VD1-(X1) $n$ -VD2-C-(X2) $n$ , wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 does not comprise an Fc region; and n is 0 or 1; and wherein said four polypeptide chains of said binding protein form four functional antigen binding sites.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises a polypeptide chain wherein the polypeptide chain comprises VD1-(X1) $n$ -VD2-C-(X2) $n$ , wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; and n is 0 or 1. In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises a polypeptide chain, wherein the polypeptide chain comprises VD1-(X1) $n$ -VD2-C-(X2) $n$ , wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a linker with the proviso that it is not a CH1 or CL; X2 does not comprise an Fc region; and n is 0 or 1. In a further embodiment, (X1) $n$  on the heavy and/or light chain is (X1)0 and/or (X2) $n$  on the heavy and/or light chain is (X2)0.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises first and second polypeptide chains, wherein the first polypeptide chain comprises a first VD1-(X1) $n$ -VD2-C-(X2) $n$ , wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a first linker with the proviso that it is not CH2; X2 is an Fc region; n is 0 or 1; and wherein the second polypeptide chain comprises a second VD1-(X1) $n$ -VD2-C-(X2) $n$ , wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a second linker with the proviso that it is not CH1 or CL; X2 does not comprise an Fc region; and n is 0 or 1.

In certain embodiments, the VD1 of the first polypeptide chain and the VD1 of the second polypeptide chain are from different first and second parent antibodies, respectively, or binding portions thereof. In certain embodiments, the VD2 of the first polypeptide chain and the VD2 of the second polypeptide chain are from different first and second parent antibodies, respectively, or binding portions thereof. In certain embodiments, the first and the second parent antibodies bind different epitopes on the same target or different targets. In certain embodiments, the first parent antibody or binding portion thereof binds the first target with a potency that is different from the potency with which the second parent antibody or binding portion thereof binds the second target. In certain embodiments, the first parent antibody or binding portion thereof binds the first target with an affinity different from the affinity with which the second parent antibody or binding portion thereof binds the second target.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprise, two first polypeptide chains and two second polypeptide chains.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises first and second polypeptide chains, each independently comprising VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first variable domain; VD2 is a second variable domain; C is a constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, and wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site. IN a further embodiment, the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, and wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 does not comprise an Fc region; n is 0 or 1, wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

In certain embodiments, the formulation of the disclosure comprises a DVD-Ig protein comprising a heavy or light chain amino acid sequence as set forth in Table 58 or 65.

In certain embodiments, the formulation of the disclosure comprises a DVD-Ig protein comprising a heavy or light chain variable region amino acid sequence as set forth in Table 58 or 65 (SEQ ID NOS: 28-75). Alternatively, the formulation of the disclosure comprises a DVD-Ig protein comprising CDRs as set forth in the heavy or light chain variable region amino acid sequences as set forth in Table 58 or 65 (SEQ ID NOS: 28-75).

In certain embodiments, the formulation of the disclosure comprises an anti-TNF/IL-17 DVD-Ig protein. In certain embodiments, the anti-TNF/IL-17 DVD-Ig protein comprises a heavy and light chain sequences having an amino acid sequence as set forth in SEQ ID NOS: 62 and 63, respectively.

In certain embodiments, the formulation of the disclosure comprises an anti-IL1 $\alpha$ /IL1 $\beta$  DVD-Ig. In certain embodiments, the anti-IL1 $\alpha$ /IL1 $\beta$  DVD-Ig protein comprises a heavy and light chain sequences having an amino acid sequence as set forth in SEQ ID NOS: 66 and 67, respectively.

In certain embodiments, the DVD-Ig protein used in the formulation of the disclosure binds one of the following target combinations (in either target order): CD20/CD80, VEGF/Her2, TNF/RANKL, TNF/DKK, CD20/RANKL, DLL4/PLGF, TNF/SOST (S2), IL-9(S2)/IgE, IL-12/IL-18, TNF/IL-17, TNF/PGE2, IL1 $\alpha$ /IL1 $\beta$ , or DLL4/VEGF.

In certain embodiments, the formulation of the disclosure is a pharmaceutical formulation, including a pharmaceutical aqueous formulation or a pharmaceutical lyophilized formulation.

Also included in the disclosure are methods of making and using AS-DVD-Ig protein or LS-DVD-Ig protein formulations.

In certain embodiments, the formulations of the disclosure are used for treating a disorder in a subject.

A further embodiment of the disclosure is a method of identifying either an AS-DVD-Ig protein or an LS-DVD-Ig protein. Such methods include aggregation testing (*e.g.*, by SEC analysis) following accelerated storage (*e.g.*, 14 days at about 40 degrees C) of a liquid formulation comprising the DVD-Ig protein, a citrate/phosphate buffer, and a high concentration of DVD-Ig protein (*e.g.*, about 50 mg/ml or greater).

## BRIEF DESCRIPTION OF DRAWINGS

**Figure 1** shows a graphic description of a comparison of DSC profiles of an IgG1 antibody (Briakinumab) to that of a DVD-Ig protein (TNF/PGE2; DVD-B) showing the difference in three vs. four domain unfolding, respectively. The sample composition of the DVD-Ig solution used was 1 mg/ml DVD-Ig protein, 1 mM ionic strength Histidine pH 6, 1 °C/minute scan rate.

**Figure 2A** shows a graphic description of serum stability of various DVD-Ig proteins.

**Figure 2B** shows the domain orientation concept for different variable domain combinations.

**Figure 3** shows a graphic description of the correlation between pharmacokinetic parameters of different DVD-Ig proteins and high molecular weight (HMW) aggregate formation.

**Figure 4** provides a graphic description of a thermodynamic comparison of 14 monoclonal antibodies and 16 DVD-Ig proteins showing the melting temperatures of the molecules relative to the number of molecules (y axis) for each temperature. The mean onset melting temperature for the antibodies was  $59.2\text{ °C} \pm 3.4\text{ °C}$ . The mean onset melting temperature for the DVD-Ig proteins was  $53.6\text{ °C} \pm 3.6\text{ °C}$ .

**Figure 5** graphically depicts the molar ellipticity in the DVD-Ig proteins as measured using near UV-CD scans between 250-320 nm wavelengths.

## DETAILED DESCRIPTION

### I. Definitions

The term "multivalent binding protein" is used to denote a binding protein comprising two or more target binding sites. The multivalent binding protein may be engineered to have the three or more antigen binding sites, and is generally not a naturally occurring antibody.

The term "multispecific binding protein" refers to a binding protein capable of binding two or more related or unrelated targets. An example of a multivalent binding protein is a Dual Variable Domain (DVD) binding protein, such as a DVD-Ig<sup>TM</sup>. In certain embodiments, DVD binding proteins comprise two or more antigen binding sites and are tetravalent or multivalent binding proteins. DVDs may be monospecific, *i.e.*, capable of binding one target, or multispecific, *i.e.*, capable of binding two or more targets.

The term "Dual Variable Domain Immunoglobulin" or "DVD-Ig<sup>TM</sup>" or "DVD-Ig

protein" refers to a DVD binding protein comprising two heavy chain DVD polypeptides and two light chain DVD polypeptides. Each half of a DVD-Ig comprises a heavy chain DVD polypeptide and a light chain DVD polypeptide, and two target binding sites. Each binding site comprises a heavy chain variable domain and a light chain variable domain with a total of 6 CDRs involved in target binding. Each variable domain (VD) in a DVD-Ig protein may be obtained from one or more "parent" monoclonal antibodies (mAbs) that bind one or more desired antigens or epitopes. In certain embodiments, the resulting DVD-Ig molecule retains activities of both parental mAbs. The term "DVD-Ig protein" is inclusive of the terms AS-DVD-Ig protein and LS-DVD-Ig protein described below.

The term "Aqueous Stable Dual Variable Domain Immunoglobulin" or "AS-DVD-Ig" or "AS-DVD-Ig protein" refers to a subset of DVD-Ig proteins that have low aggregation or a low change in monomer content due to physical degradation following stability tests at a given temperature and time period, *e.g.*, 5 °C or 40 °C for 14 to 21 days, at a concentration ranging from about 1 to about 100 mg/ml (*e.g.*, about 1-10 mg/ml or about 50-100 mg/ml) and at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0. Different stability tests may be used to define an AS-DVD-Ig. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has about 1% relative peak area or less change in monomers as determined by SEC analysis at about 5 °C after 21 days of storage at a concentration of about 50 mg/ml to about 100 mg/ml and at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0, in a buffered aqueous formulation. In certain embodiments, an AS-DVD-Ig protein has 10% relative (rel.) peak area or less change in monomers as determined by SEC analysis following accelerated storage for 21 days at about 40 °C at a concentration of about 50 mg/ml to about 100 mg/ml and at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0, in a buffered aqueous formulation. In certain embodiments, stability testing for AS-DVD-Ig proteins may be performed by testing the stability, *e.g.*, loss of monomers, of a solution having a DVD-Ig protein concentration of 50 to 100 mg/ml in a citrate phosphate buffer or histidine buffer at a pH between 5.0 – 6.5, *e.g.*, about 5.5 to about 6.0.

The term "aqueous formulation" refers to a liquid solution in which the solvent is water. In certain embodiments, the term "aqueous formulation" refers to a liquid formulation in which the solvent is water wherein the formulation was not previously lyophilized (*i.e.*, does not result from reconstitution of a lyophilized formulation).

The term “Lyophilized Stable Dual Variable Domain Immunoglobulin” or “LS-DVD-Ig” or “LS-DVD-Ig protein” refers to a subset of DVD-Ig proteins that have low aggregation or elevated levels of change in monomers (*i.e.*, loss of monomers) in the liquid state. Different stability tests may be used to define an LS-DVD-Ig. In certain embodiments, an LS-DVD-Ig protein has more than 10% rel. peak area change in monomers observed, following accelerated storage, *e.g.*, 21 days at about 40 °C, when formulated at a concentration of about 50 to 100 mg/ml at a pH between about 5.0 and about 6.5, *e.g.*, about 5.5 to about 6.0, in an aqueous formulation. DVD-Ig proteins may be tested in aqueous formulations containing citrate and phosphate buffer, or histidine buffer. Change in monomers can be determined according to methods known in the art, including, but not limited to, SEC. In certain embodiments, the accelerated storage conditions include storing the DVD-Ig protein in the absence of light at about 40 °C. In certain embodiments, an LS-DVD-Ig protein has 50% rel. peak area or less change in monomers as determined by SEC analysis following accelerated storage for 14 days at about 40 °C, where the LS-DVD-Ig protein is formulated at a concentration of at least 50 mg/ml in a citrate phosphate buffer in an aqueous formulation. In certain embodiments, stability testing for LS-DVD-Ig proteins may be performed by testing the stability, *e.g.*, loss of monomers, of a solution having a DVD-Ig protein concentration of about 50 mg/ml to 100 mg/ml) in a citrate phosphate buffer or histidine buffer at a pH between about 5.0 – about 6.5, *e.g.*, about 5.5 to about 6.0.

The term “pharmaceutical formulation” refers to preparations that are in such a form as to permit the biological activity of the active ingredients to be effective and, therefore, may be administered to a subject for therapeutic use. In certain embodiments, the disclosure provides an aqueous pharmaceutical formulation comprising an AS-DVD-Ig. In other embodiments, the disclosure provides a lyophilized pharmaceutical formulation comprising an LS-DVD-Ig.

A “stable” formulation is one in which the DVD-Ig protein therein essentially retains its physical stability and/or chemical stability and/or biological activity upon storage. Various analytical techniques for measuring protein stability are available in the art and are reviewed in, *e.g.*, Peptide and Protein Drug Delivery, pp. 247-301, Vincent Lee Ed., Marcel Dekker, Inc., New York, N.Y., Pubs. (1991) and Jones (1993) Adv. Drug Delivery Rev. 10: 29-90. In certain embodiments, the stability of the DVD-Ig protein is determined according to the percentage of monomer protein in the solution, with a low percentage of degraded (*e.g.*,

fragmented) and/or aggregated protein. For example, an aqueous formulation comprising a stable DVD-Ig protein may include at least 95% monomer DVD-Ig protein, *e.g.*, AS-DVD-Ig protein. Alternatively, an aqueous formulation of the disclosure may include no more than 5% aggregate and/or degraded DVD-Ig protein, *e.g.*, AS-DVD-Ig protein.

A DVD-Ig protein "retains its physical stability" in a pharmaceutical formulation if it shows substantially no signs of aggregation, precipitation and/or denaturation upon visual examination of color and/or clarity, or as measured by UV light scattering or by size exclusion chromatography. In certain embodiments of the disclosure, a stable aqueous formulation is a formulation having less than about 10% aggregation, and less than about 5% AS-DVD-Ig protein aggregation in the formulation.

A DVD-Ig protein "retains its chemical stability" in a pharmaceutical formulation if the chemical stability at a given time is such that the DVD-Ig protein is considered to still retain its biological activity as defined below. Chemical stability can be assessed by detecting and quantifying chemically altered forms of the DVD-Ig. Chemical alteration may involve size modifications (*e.g.*, clipping) which can be evaluated using size exclusion chromatography, SDS-PAGE and/or matrix-assisted laser desorption ionization / time of flight mass spectrometry (MALDI/TOF MS), for example. Other types of chemical alteration include charge alteration (*e.g.*, occurring as a result of deamidation), which can be evaluated by, *e.g.*, ion-exchange chromatography.

A DVD-Ig protein "retains its biological activity" in a pharmaceutical formulation, if the protein in a pharmaceutical formulation is biologically active for its intended purpose. For example, biological activity of a DVD-Ig protein is retained if the biological activity of the DVD-Ig protein in the pharmaceutical formulation is within about 30%, about 20%, or about 10% (within the errors of the assay) of the biological activity exhibited at the time the pharmaceutical formulation was prepared (*e.g.*, as determined in an antigen binding assay).

The term "surfactant", as used herein, refers to organic substances having amphipathic structures; namely, they are composed of groups of opposing solubility tendencies, typically an oil-soluble hydrocarbon chain and a water-soluble ionic group. Surfactants can be classified, depending on the charge of the surface-active moiety, into anionic, cationic, and nonionic surfactants. Surfactants are often used as wetting, emulsifying, solubilizing, and dispersing agents for various pharmaceutical compositions and preparations of biological materials. Examples of suitable surfactants include, but are not limited to,

sodium lauryl sulfate, polysorbates such as polyoxyethylene sorbitan monooleate, monolaurate, monopalmitate, monostearate or another ester of polyoxyethylene sorbitan (*e.g.*, the commercially available Tweens<sup>TM</sup>, such as, Tween<sup>TM</sup> 20 and Tween<sup>TM</sup> 80 (ICI Speciality Chemicals)), sodium dioctylsulfosuccinate (DOSS), lecithin, stearyl alcohol, cetostearyl alcohol, cholesterol, polyoxyethylene ricin oil, polyoxyethylene fatty acid glycerides, poloxamers (*e.g.*, Pluronics F68<sup>TM</sup> and F108<sup>TM</sup>, which are block copolymers of ethylene oxide and propylene oxide); polyoxyethylene castor oil derivatives or mixtures thereof. In certain embodiments, a formulation of the disclosure comprises Polysorbate 20, Polysorbate 40, Polysorbate 60, or Polysorbate 80.

The term "tonicity modifier" or "tonicity agent" refers to a compound that can be used to adjust the tonicity of a liquid formulation. Examples of tonicity modifiers include glycerin, lactose, mannitol, dextrose, sodium chloride, magnesium sulfate, magnesium chloride, sodium sulfate, sorbitol, trehalose, sucrose, raffinose, maltose and others known to those of ordinary skill in the art.

The term "polyol" refers to a substance with multiple hydroxyl groups, and includes sugars (reducing and nonreducing sugars), sugar alcohols and sugar acids. In certain embodiments, polyols have a molecular weight that is less than about 600 kD (*e.g.*, in the range from about 120 to about 400 kD). A "reducing sugar" is one that contains a free aldehyde or ketone group and can reduce metal ions or react covalently with lysine and other amino groups in proteins. A "nonreducing sugar" is one that lacks a free aldehyde or ketone group and is not oxidised by mild oxidising agents such as Fehling's or Benedict's solutions. Examples of reducing sugars are fructose, mannose, maltose, lactose, arabinose, xylose, ribose, rhamnose, galactose and glucose. Nonreducing sugars include sucrose, trehalose, sorbose, melezitose and raffinose. Mannitol, xylitol, erythritol, threitol, sorbitol and glycerol are examples of sugar alcohols. As to sugar acids, these include L-gluconate and metallic salts thereof. The polyol may also act as a tonicity agent. In certain embodiments of the disclosure, one ingredient of the formulation is sorbitol in a concentration of about 10 to about 70 mg/ml. In a particular embodiment of the disclosure, the concentration of sorbitol is about 30 to about 50 mg/ml. In certain embodiments, the concentration of sucrose is about 60 to about 100 mg/ml. In a particular embodiment of the disclosure, the concentration of sucrose is about 70 to about 90 mg/ml.

The term "buffer" refers to a buffered solution that resists changes in pH by the action

of its acid-base conjugate components. A buffer used in this disclosure has a pH in the range from about 4.5 to about 7.5. Examples of buffers that will control the pH in this range include acetate (*e.g.*, sodium acetate), succinate (such as sodium succinate), gluconate, methionine, imidazole, histidine, glycine, arginine, citrate, phosphate, citrate and phosphate, Tris, and other organic acid buffers. In certain embodiments, the buffer used in the formulation of the disclosure is histidine, glycine, arginine, acetate, citrate, and/or phosphate buffered saline (PBS).

A "reconstituted" formulation is one which has been prepared by dissolving a lyophilized protein formulation in a diluent such that the protein is dispersed in the reconstituted formulation. The reconstituted formulation is suitable for administration (*e.g.* parenteral administration) to a patient to be treated with the protein of interest (*e.g.*, LS-DVD-Ig).

A "diluent" of interest herein is one which is pharmaceutically acceptable (safe and non-toxic for administration to a human) and is useful for the preparation of a liquid formulation, such as a formulation reconstituted after lyophilization. Exemplary diluents include sterile water, bacteriostatic water for injection (BWFI), a pH buffered solution (*e.g.* phosphate-buffered saline), sterile saline solution, Ringer's solution or dextrose solution. In an alternative embodiment, diluents can include aqueous solutions of salts and/or buffers.

A "therapeutically effective amount" or "effective amount" of a binding protein refers to an amount effective in the prevention or treatment of a disorder for the treatment of which the antibody is effective.

The term "disorder" refers to any condition that would benefit from treatment with the formulations of the disclosure. This includes chronic and acute disorders or diseases including those pathological conditions that predispose the subject to the disorder in question.

The term "treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those patients in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented.

The terms "parenteral administration" and "administered parenterally" means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and

intrasternal injection and infusion. The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and is subject to metabolism and other like processes, for example, subcutaneous administration.

The term "antibody" broadly refers to an immunoglobulin (Ig) molecule, generally comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains, or any functional fragment, mutant, variant, or derivative thereof, that retains the essential target binding features of an Ig molecule.

In a full-length antibody, each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. Immunoglobulin molecules can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY) and class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass.

The term "Fc region" means the C-terminal region of an immunoglobulin heavy chain, which may be generated by papain digestion of an intact antibody. The Fc region may be a native sequence Fc region or a variant sequence Fc region. The Fc region of an immunoglobulin generally comprises two constant domains, a CH2 domain and a CH3 domain, and optionally comprises a CH4 domain.

The term "antigen-binding portion" refers to one or more fragments of a binding protein that specifically binds to a target or an antigen. Such embodiments may be monospecific, or may be bispecific, dual specific, or multi-specific (may specifically bind two or more different antigens).

A "functional antigen binding site" of a binding protein is one that is capable of binding a target antigen. The antigen binding affinity of the functional antigen binding site is

not necessarily as strong as the parent antibody from which the antigen binding site is derived, but the ability to bind antigen must be measurable using a known method for evaluating antibody binding to an antigen. Moreover, the antigen binding affinity of each of the functional antigen binding sites of a multivalent binding protein need not be quantitatively the same.

The term "linker" denotes polypeptides comprising two or more amino acid residues joined by peptide bonds that are used to link one or more antigen binding portions. Such linker polypeptides are well known in the art (see, *e.g.*, Holliger *et al.* (1993) Proc. Natl. Acad. Sci. USA 90:6444-6448; Poljak *et al.* (1994) Structure 2:1121-1123).

An "immunoglobulin constant domain" refers to a heavy or light chain constant domain. Human heavy chain and light chain (*e.g.*, IgG) constant domain amino acid sequences are known in the art.

The term "monoclonal antibody" refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different epitopes, each monoclonal antibody is directed against a single epitope on the antigen. The modifier "monoclonal" is not to be construed as requiring production of the antibody by any particular method.

The term "human antibody" includes antibodies that have variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies may include amino acid residues not encoded by human germline immunoglobulin sequences (*e.g.*, mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo). However, the term "human antibody" is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The term "chimeric antibody" means antibodies that comprise heavy and light chain variable region sequences from one species and constant region sequences from another species, such as antibodies having murine heavy and light chain variable regions linked to human constant regions.

The term "CDR-grafted antibody" means antibodies that comprise heavy and light chain variable region sequences from one species but in which the sequences of one or more of the CDR regions of their VH and/or VL are replaced with the CDR sequences of another species, such as antibodies having human heavy and light chain variable regions in which one or more of the murine CDRs (*e.g.*, CDR3) has been replaced with murine CDR sequences.

The term "humanized antibody" means an antibody that comprises heavy and light chain variable region sequences from a non-human species (*e.g.*, a mouse) but in which at least a portion of the VH and/or VL sequence has been altered to be more "human-like", *i.e.*, more similar to human germline variable sequences. One type of humanized antibody comprises non-human CDR sequences and human framework sequences.

The term "CDR" means the complementarity determining region within antibody variable sequences. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The term "CDR set" as used herein refers to a group of three CDRs that occur in a single variable region capable of binding the target. The exact boundaries of these CDRs have been defined differently according to different systems.

The terms "Kabat numbering", "Kabat definitions and "Kabat labeling" are used interchangeably herein. These terms refer to a system of numbering amino acid residues that are more variable (*i.e.*, hypervariable) than other amino acid residues in the heavy and light chain variable regions of an antibody, or an antigen binding portion thereof (Kabat *et al.* (1971) Ann. NY Acad. Sci. 190:382-391 and Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the hypervariable region generally ranges from amino acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region generally ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

Chothia and coworkers (Chothia and Lesk (1987) J. Mol. Biol. 196:901-917 and Chothia *et al.* (1989) Nature 342:877-883) found that certain sub-portions within Kabat CDRs adopt nearly identical peptide backbone conformations, despite having great diversity at the level of amino acid sequence. These sub-portions were designated as L1, L2 and L3 or H1, H2 and H3 where the "L" and the "H" designates the light chain and the heavy chains

regions, respectively. These regions may be referred to as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (1995) FASEB J. 9:133-139 and MacCallum (1996) J. Mol. Biol. 262(5):732-45. Still other CDR boundary definitions may not strictly follow one of the above systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems, although embodiments use Kabat or Chothia defined CDRs.

The term "framework" or "framework sequence" refers to the remaining sequences of a variable region minus the CDRs. Because the exact definition of a CDR sequence can be determined by different systems, the meaning of a framework sequence is subject to correspondingly different interpretations. The six CDRs (CDR-H1, -H2, and -H3 of the heavy chain and CDR-L1, -L2, and -L3 of the light chain) also divide the framework regions on the light chain and the heavy chain into four sub-regions (FR1, FR2, FR3 and FR4) on each chain, in which CDR1 is positioned between FR1 and FR2, CDR2 between FR2 and FR3, and CDR3 between FR3 and FR4. The term "activity" includes activities such as the binding specificity and binding affinity of a DVD-Ig protein for two or more antigens.

The term "epitope" includes any polypeptide determinant capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and/or specific charge characteristics. In certain embodiments, an epitope is a region of an antigen that is bound by an antibody or multispecific binding protein.

The term "surface plasmon resonance" or "SPR", refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIACore system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, N.J.). For further descriptions, see Jonsson *et al.* (1993) Ann. Biol. Clin. 51:19-26; Jonsson *et al.* (1991) Biotechniques 11:620-627; Jonsson *et al.* (1995) J. Mol. Recognit. 8:125-131; and Johnsson *et al.* (1991) Anal. Biochem. 198:268-277.

The term "K<sub>on</sub>" refers to the on rate constant for association of a binding protein to the antigen to form the binding protein /antigen complex as is known in the art.

The term "K<sub>off</sub>" refers to the off rate constant for dissociation of a binding protein from the binding protein/antigen complex as is known in the art.

The term "K<sub>d</sub>" refers to the dissociation constant of a particular antibody-antigen interaction as is known in the art.

The terms "specific binding", "specifically binding" or "specifically binds", as used herein, in reference to the interaction of a binding protein with a second chemical species, mean that the interaction is dependent upon the presence of a particular structure (e.g., an antigenic determinant or epitope) on the chemical species; for example, a DVD-Ig protein recognizes and binds to a specific protein structure rather than to proteins generally. If a DVD-Ig protein is specific for epitope "A", the presence of a molecule containing epitope A (or free, unlabeled A), in a reaction containing labeled "A" and the DVD-Ig protein, will reduce the amount of labeled A bound to the DVD-Ig protein.

## **II. Dual Variable Domain Immunoglobulin (DVD-Ig) Proteins for Use in Formulations of the Disclosure**

The disclosure pertains to formulations, and uses thereof, of DVD-Ig proteins, particularly those identified as AS-DVD-Ig protein or LS-DVD-Ig protein (described in more detail below).

### **A. General DVD-Ig Protein Structure**

In certain embodiments, the DVD-Ig protein used in the formulations and methods of the disclosure comprises a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first variable domain, VD2 is a second variable domain, C is a constant domain, X1 represents an amino acid or polypeptide, X2 represents an Fc region and n is 0 or 1.

In certain embodiments, a DVD-Ig protein contains two polypeptide chains, wherein a first polypeptide chain comprises VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and a second polypeptide chain comprises VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first

light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 does not comprise an Fc region; and n is 0 or 1; and wherein said two polypeptide chains of said binding protein form two functional antigen binding sites.

In certain embodiments, a DVD-Ig protein contains four polypeptide chains, wherein two polypeptide chains comprise VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and two polypeptide chains comprise VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 does not comprise an Fc region; and n is 0 or 1; and wherein said four polypeptide chains of said binding protein form four functional antigen binding sites.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises a polypeptide chain wherein the polypeptide chain comprises VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; and n is 0 or 1. In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises a polypeptide chain, wherein the polypeptide chain comprises VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a linker with the proviso that it is not a CH1 or CL; X2 does not comprise an Fc region; and n is 0 or 1. In a further embodiment, (X1)n on the heavy and/or light chain is (X1)0 and/or (X2)n on the heavy and/or light chain is (X2)0.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises first and second polypeptide chains, wherein the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a first linker with the proviso that it is not CH2; X2 is an Fc region; n is 0 or 1; and wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain

constant domain; X1 is a second linker with the proviso that it is not CH1 or CL; X2 does not comprise an Fc region; and n is 0 or 1.

In certain embodiments, the VD1 of the first polypeptide chain and the VD1 of the second polypeptide chain are from different first and second parent antibodies, respectively, or binding portions thereof. In certain embodiments, the VD2 of the first polypeptide chain and the VD2 of the second polypeptide chain are from different first and second parent antibodies, respectively, or binding portions thereof. In certain embodiments, the first and the second parent antibodies bind different epitopes on the same target or different targets. In certain embodiments, the first parent antibody or binding portion thereof binds the first target with a potency that is different from the potency with which the second parent antibody or binding portion thereof binds the second target. In certain embodiments, the first parent antibody or binding portion thereof binds the first target with an affinity different from the affinity with which the second parent antibody or binding portion thereof binds the second target.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises, two first polypeptide chains and two second polypeptide chains.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises first and second polypeptide chains, each independently comprising VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first variable domain; VD2 is a second variable domain; C is a constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site. In a further embodiment, the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, and wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 does not comprise an Fc region; n is 0 or 1, wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

In certain embodiments, the AS-DVD-Ig protein or LS-DVD-Ig protein comprises first and second polypeptide chains, each independently comprising VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first variable domain; VD2 is a second variable domain; C is a constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, and wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site. In a further embodiment, the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first heavy chain variable domain; VD2 is a second heavy chain variable domain; C is a heavy chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 is an Fc region; n is 0 or 1, and wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n, wherein VD1 is a first light chain variable domain; VD2 is a second light chain variable domain; C is a light chain constant domain; X1 is a linker with the proviso that it is not CH1; X2 does not comprise an Fc region; n is 0 or 1, wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

Examples of DVD-Ig proteins are described in US Patent No. 7,612,181, which is incorporated by reference herein.

Examples of DVD-Ig proteins that may be used in the methods and compositions of the disclosure are described below, including Tables 58 and 65. Further examples of DVD-Ig proteins that may be used in the methods and compositions of the disclosure are described in SEQ ID NOs: 28 to 75.

#### B. Generation of DVD-Ig Proteins

The variable domains of a DVD-Ig protein can be obtained from parent antibodies, including polyclonal and monoclonal antibodies capable of binding targets of interest. These antibodies may be naturally occurring or may be generated by recombinant technology. Examples of antibodies that may be used in making DVD-Ig proteins include chimeric antibodies, human antibodies, and humanized antibodies. Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including, for example, the use of hybridoma, recombinant, and phage display technologies, or any combination thereof. Monoclonal antibodies may also be produced by immunizing a non-human animal

comprising some, or all, of the human immunoglobulin locus with an antigen of interest, such as, for example, XENOMOUSE™ transgenic mouse, an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production. Methods of generating DVD-Ig proteins are described in US Patent No. 7,612,181, the teachings of which are incorporated by reference herein. DVD-Ig proteins used in the compositions and methods of the disclosure may be made from antibodies capable of binding specific targets and well known in the art. These include, but are not limited to an anti-TNF antibody (U.S. Patent No. 6,258,562), anti-IL-12 and or anti-IL-12p40 antibody (U.S. Patent No. 6,914,128); anti-IL-18 antibody (US Patent Publication No. 20050147610), as well as anti-C5, anti-CBL, anti-CD147, anti-gp120, anti-VLA4, anti-CD11a, anti-CD18, anti-VEGF, anti-CD40L, anti-Id, anti-ICAM-1, anti-CXCL13, anti-CD2, anti-EGFR, anti-TGF-beta 2, anti-E-selectin, anti-Fact VII, anti-Her2/neu, anti-F gp, anti-CD11/18, anti-CD14, anti-ICAM-3, anti-CD80, anti-CD4, anti-CD3, anti-CD23, anti-beta2-integrin, anti-alpha4beta7, anti-CD52, anti-HLA DR, anti-CD22, anti-CD20, anti-MIF, anti-CD64 (FcR), anti-TCR alpha beta, anti-CD2, anti-Hep B, anti-CA 125, anti-EpCAM, anti-gp120, anti-CMV, anti-gpIIbIIIa, anti-IgE, anti-CD25, anti-CD33, anti-HLA, anti-VNRintegrin, anti-IL-1alpha, anti-IL-1beta, anti-IL-1 receptor, anti-IL-2 receptor, anti-IL-4, anti-IL4 receptor, anti-IL5, anti-IL-5 receptor, anti-IL-6, anti-IL-8, anti-IL-9, anti-IL-13, anti-IL-13 receptor, anti-IL-17, and anti-IL-23 antibodies (see Presta (2005) *J. Allergy Clin. Immunol.* 116:731-6 and Clark "Antibodies for Therapeutic Applications," Department of Pathology, Cambridge University, UK (2000), published online at M. Clark's home page at the website for the Department of Pathology, Cambridge University.

Parent monoclonal antibodies may also be selected from various therapeutic antibodies approved for use, in clinical trials, or in development for clinical use. Such therapeutic antibodies include, but are not limited to: rituzimab (RITUXAN™ Biogen Idec, Genentech/Roche) (see for example U.S. Patent No. 5,736,137) a chimeric anti-CD20 antibody approved to treat non-Hodgkin's lymphoma; ofatumumab (HUMAX-CD20™ Genmab, GlaxoSmithKlein) (described in U.S. Patent 5,500,362) an anti-CD20 antibody approved to treat chronic lymphocytic leukemia that is refractory to fludarabine and alemtuzumab; AME-133v (Mentrik Biotech) an anti-CD20 antibody; veltuzumab (hA20) (Immunomedics) an anti-CD20 antibody; HumaLYM (Intracel); PRO70769 (Genentech/Roche) (PCT/US2003/040426) an anti-CD20 antibody; trastuzumab

(HERCEPTIN™ Genentech/Roche) (described in U.S. Patent No. 5,677,171) a humanized anti-Her2/neu antibody approved to treat breast cancer; pertuzumab (rhuMab-2C4, OMNITARG™ Genentech/Roche) (described in U.S. Patent No. 4,753,894); cetuximab (ERBITUX™ Imclone) (described in U.S. Patent No. 4,943,533; PCT WO 96/40210) a chimeric anti-EGFR antibody approved to treat colorectal and head and neck cancer; panitumumab (ABX-EGF VECTIBIX® Amgen) (described in U.S. Patent No. 6,235,883) an anti-EGFR antibody approved to treat colorectal cancer; zalutumumab (HUMAX-EGFR™ Genmab) (described in U.S. Patent Application Serial No. 10/172,317) an anti-EGFR antibody; EMD55900 (Mab 425 Merck) an anti-EGFR antibody; EMD62000 and EMD72000 (Mab 425 Merck) anti-EGFR antibodies (described in U.S. Patent No. 5,558,864; Murthy *et al.* (1987) *Arch. Biochem. Biophys.* 252(2):549-60; Rodeck *et al.* (1987) *J. Cell. Biochem.* 35(4):315-20; Kettleborough *et al.* (1991) *Protein Eng.* 4(7):773-83; ICR62 (Institute of Cancer Research) an anti-EGFR antibody (described in PCT Publication No. WO 95/20045; Modjtahedi *et al.* (1993) *J. Cell. Biophys.* 22(1-3):129-46; Modjtahedi *et al.* (1993) *Br. J. Cancer* 67(2):247-53; Modjtahedi *et al.* (1996) *Br. J. Cancer* 73(2):228-35; Modjtahedi *et al.* (2003) *Int. J. Cancer* 105(2):273-80); nimotuzumab (TheraCIM hR3, THERALOC® YM Biosciences, Oncoscience AG) (described in U.S. Patent No. 5,891,996; U.S. Patent No. 6,506,883; Mateo *et al.* (1997) *Immunotechnol.* 3(1):71-81) an anti-EGFR antibody; ABT-806 (Ludwig Institute for Cancer Research, Memorial Sloan-Kettering) (Jungbluth *et al.* (2003) *Proc. Natl. Acad. Sci. USA* 100(2):639-44) an anti-EGFR antibody; KSB-102 (KS Biomedix); MR1-1 (IVAX, National Cancer Institute) (PCT Publication No. WO 0162931A2) an anti-EGFRvIII antibody; SC100 (Scancell) (PCT Publication No. WO 01/88138) an anti-EGFR antibody; alemtuzumab (CAMPATH™ Genzyme/Sanofi) an anti-CD52 antibody approved to treat B-cell chronic lymphocytic leukemia; muromonab-CD3 (Orthoclone OKT3™ Johnson and Johnson) an anti-CD3 antibody approved to treat organ transplant rejection; ibritumomab tiuxetan (ZEVALIN™ Spectrum Pharmaceuticals) an anti-CD20 antibody approved to treat non-Hogkins Lymphoma; gemtuzumab ozogamicin (hP67.6 MYLOTARG™ Pfizer) an anti-CD33 antibody conjugated to calicheamicin; alefacept (AMEVIVE™ Astellas Pharma) an anti-CD2 LFA-3 Fc fusion; abciximab (REOPRO™ Centocor Ortho Biotech Products, Lilly) a chimeric human-mouse anti-glycoprotein IIb/IIIa receptor and anti-vitronectic  $\alpha_v\beta_3$  receptor antibody approved as an adjunct to percutaneous coronary intervention to prevent cardiac ishemia; basiliximab (SIMULECT™ Novartis) an

anti-CF25 antibody approved to treat organ transplant rejection; palivizumab (SYNAGIS™ Medimmune) an antibody to the A antigenic site of F protein of RSV approved to treat RSV infection; infliximab (REMICADE™ Janssen Biotech) an anti-TNF $\alpha$  antibody approved to treat Crohn's disease, ulcerative colitis, arthritis, ankylosing spondylitis, psoriatic arthritis, and plaque psoriasis; adalimumab (HUMIRA™ AbbVie) an anti-TNF $\alpha$  antibody approved to treat rheumatoid arthritis, juvenile idiopathic arthritis, psoriatic arthritis, ankylosing spondylitis, Crohn's disease, ulcerative colitis, plaque psoriasis; CDP571 (HUMICADE™ Celltech, Biogen IDEC) an anti-TNF $\alpha$  antibody ; etanercept (ENBREL™ Amgen, Pfizer) an anti-TNF $\alpha$  Fc fusion antibody approved to treat rheumatoid arthritis, juvenile idiopathic arthritis, psoriatic arthritis, ankylosing spondylitis, plaque psoriasis; certolizumab pegol (CIMZIA)UCB Pharma) an anti-TNF $\alpha$  antibody approved to treat rheumatoid arthritis and Crohn's disease; ustekinumab (STELARA Janssen Biotech) a human anti-p40 subunit of IL-12 and IL-23 antibody approved to treat plaque psoriasis; galilimomab (ABX-CBL Abgenix) a mouse anti-CD147 antibody; ABX-IL8 (Abgenix) an anti-IL8 antibody ; ABX-MA1 (Abgenix) an anti-MUC18 antibody ; pemtumomab (Theragyn, R1549, 90Y-muHMFG1Antisoma) a mouse anti-MUC1-Yttrium 90 antibody conjugate; Therex (R1550 Antisoma) an anti-MUC1 antibody; AngioMab (muBC-1, AS1405 Antisoma); HuBC-1 (Antisoma); Thioplatin (AS1407 Antisoma); natalizumab (TYSABRI® Biogen Idec, Elan) an anti- $\alpha$ 4 integrin antibody approved to treat multiple sclerosis and Crohn's disease; VLA-1 (Santarus) a humanized anti-VLA-1 antibody ; LTBR mAb (Biogen Idec) an anti-lymphotoxin  $\beta$  receptor antibody; lerdelimumab (CAT-152 Cambridge Antibody Technology/Abbott) an anti-TGF- $\beta$ 2 antibody ; briakinumab (AbbVie) an anti-IL-12 and 23 antibody ; metelimumab (CAT-192 Cambridge Antibody Technology, Genzyme) an anti-TGF $\beta$ 1 antibody ; bertilimumab (CAT-213, iCO-008 Cambridge Antibody Technology, iCo Therapeutics, Immune Pharmaceuticals) an anti-eotaxin1 antibody ; belimumab (BENLYSTA® GlaxoSmithKline) an anti-B lymphocyte stimulator protein antibody approved to treat systemic lupus erythematosus; mapatumumab (HGS-ETR1 Cambridge Antibody Technology, Human Genome Sciences) an anti-TRAIL-R1 antibody; bevacizumab (AVASTIN™ Genentech/Roche) an anti-VEGF antibody approved to treat metastatic colorectal cancer, non-squamous non-small cell lung cancer, glioblastoma, metastatic renal cell cancer; anti-HER3/EGFR antibody (Genentech/Roche); an Anti-Tissue Factor antibody (Genentech/Roche); omalizumab (XOLAIR™ Genentech/Roche, Novartis) an anti-IgE

antibody approved to treat severe allergic asthma; efalizumab (RAPTIVA™ Genentech/Roche, Merck Serono) an anti-CD11a antibody; MLN-02 (Millenium, Genentech/Roche) an anti- $\alpha$ 4 $\beta$ 7 integrin antibody; zanolimumab (HUMAX CD4™ Emergent BioSolutions) an anti-CD4 antibody ; HUMAX -IL15™ (AMG-714 Genmab, Amgen) an anti-IL15 antibody ; HuMax-IL8 (HUMAX -Inflam™, MDX-018 Genmab, Cormorant Pharmaceuticals) an anti-IL8 antibody; HUMAX™ -Cancer, (Genmab, Medarex, Oxford GlycoSciences) an anti-Heparanase I antibody; HUMAX™ -Lymphoma (Genmab) an anti-IL8 antibody; HUMAX™ -TAC (Genmab) an anti-IL-2R $\alpha$ , CD25 antibody ; daratumumab (HuMax®-CD38, Genmab, Janssen Biotech) an anti-CD38 antibody; toralizumab (IDE-131 Biogen Idec) an anti-CD40L antibody ; clenolimimab (IDE-151 Biogen Idec) an anti-CD4 antibody ; glaiximab (IDE-114 Biogen Idec) an anti-CD80 antibody ; lumilixmab (IDE-152 Biogen Idec) an anti-CD23 ; anti-macrophage migration factor (MIF) antibodies (Biogen Idec, Taisho Pharmaceutical); mitumomab (BEC2 Imclone) a mouse anti-idiotypic antibody ; IMC-1C11 (Imclone) a chimeric anti-VEGFR2 antibody; DC101 (Imclone) murine anti-VEGFR2 antibody;; anti-VE cadherin antibody (Imclone); labetuzumab (CEA-CIDE™ Immunomedics) an anti-carcinoembryonic antigen antibody ; epratuzumab (LYMPHOCIDE™ Immunomedics) an anti-CD22 antibody ; yttrium (<sup>90</sup>Y) tacatuzumab tetraxetan (AFP-Cide® Immunomedics) an anti- $\alpha$ fetoprotein antibody; milatuzumab (MyelomaCide® Immunomedics) an anti-CF74 antibody; LeukoCide® (Immunomedics); ProstaCide® (Immunomedics); ipilimumab (Yervoy™, MDX-010 Bristol-Myers Squibb) an anti-CTLA4 antibody approved to treat melanoma; iratumumab (MDX-060 Medarex) an anti-CD30 antibody ; MDX-070 (Medarex) an anti-prostate specific membrane antigen; OSIDEM™ (IDM-1 Medarex, Immuno-Designed Molecules) an anti-Her2 antibody; HUMAX™ -CD4, an anti-CD4 antibody being developed by Medarex and Genmab; HuMax-IL15, an anti-IL15 antibody being developed by Medarex and Genmab; golimumab (SIMPONI™ Janssen Biotech) an anti-TNF $\alpha$  antibody approved to treat rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis; ustekinumab (STELARA®, CINTO 1275 Janssen Biotech) an anti-IL-12 antibody approved to treat plaque psoriasis; MOR101 and MOR102 (MorphoSys) anti-intercellular adhesion molecule-1 (ICAM-1) (CD54) antibodies; MOR201 (MorphoSys) an anti-fibroblast growth factor receptor 3 antibody ; visilizumab (NUVION™ PDL BioPharma) an anti-CD3 antibody ; fontolizumab (HUZAF™ PDL BioPharma) an anti-INF $\gamma$  antibody ; volociximab (M200 PDL BioPharma, Biogen Idec) an anti- $\alpha$ 5 $\beta$ 1 integrin

antibody ; SMART<sup>®</sup> IL-12 (PDL BioPharma) an anti-IL-12 ; ING-1 (Xoma) an anti-Ep-CAM antibody; omalizumab (XOLAIR<sup>™</sup> Genentech/Roche, Novartis) an anti-IgE antibody approved to treat allergic asthma; MLN01 (Xoma) an anti- $\beta$  integrin antibody; and tocilizumab (ACTEMRA<sup>™</sup> Genentech/Roche) an anti-IL6 antibody approved to treat rheumatoid arthritis and systemic juvenile idiopathic arthritis.

### C. Construction of DVD-Ig Proteins

A DVD-Ig protein is formed by combining two heavy chain DVD polypeptides and two light chain DVD polypeptides. The dual variable domain immunoglobulin (DVD-Ig) heavy chain comprises two heavy chain variable domains (VH) linked in tandem, directly or by a linker, followed by the constant domain CH1 and Fc region. The dual variable domain immunoglobulin (DVD-Ig) light chain is designed such that two light chain variable domains (VL) from the two parent mAbs are linked in tandem, directly or via a linker, followed by the light chain constant domain (CL) (see Figure 1A of U.S. Patent No. 7,612,181, incorporated by reference herein). Methods of making DVD-Ig proteins are also described in U.S. Patent No. 7,612,181, incorporated by reference herein.

The variable domains of the DVD-Ig protein can be obtained using recombinant DNA techniques from a parent antibody generated by any one of the methods described above. In certain embodiments, the variable domain is a CDR grafted or a humanized variable heavy or light chain domain. In certain embodiments, the variable domain is a human heavy or light chain variable domain.

The linker sequence may be a single amino acid or a polypeptide sequence. Examples of linker sequences that may be used to link variable domains include, but are not limited to, AKTTPKLEEGEFSEAR (SEQ ID NO:1); AKTTPKLEEGEFSEARV (SEQ ID NO:2); AKTTPKLGG (SEQ ID NO:3); SAKTTPKLGG (SEQ ID NO:4); SAKTTP (SEQ ID NO:5); RADAAP (SEQ ID NO:6); RADAAPTVS (SEQ ID NO:7); RADAAAAGGPGS (SEQ ID NO:8); RADAAAA(G<sub>4S</sub>).<sub>4</sub> (SEQ ID NO:9), SAKTTPKLEEGEFSEARV (SEQ ID NO:10); ADAAP (SEQ ID NO:11); ADAAPTVSIFPP (SEQ ID NO:12); TVAAP (SEQ ID NO:13); TVAAPSVFIFPP (SEQ ID NO:14); QPKAAP (SEQ ID NO:15); QPKAAPSVTLFPP (SEQ ID NO:16); AKTTPP (SEQ ID NO:17); AKTTPPSVTPLAP (SEQ ID NO:18); AKTTAP (SEQ ID NO:19); AKTTAPSVDPLAP (SEQ ID NO:20); ASTKGP (SEQ ID NO:21); ASTKGPSVFPLAP (SEQ ID NO:22); GGGGSGGGSGGGGS (SEQ ID NO:23);

GENKVEYAPALMALS (SEQ ID NO:24); GPAKELTPLKEAKVS (SEQ ID NO:25); GHEAAAVMQVQYPAS (SEQ ID NO:26); and GGGGSGGGS (SEQ ID NO: 27). Other examples of linkers are described in U.S. Patent Publication No. 20100226923, incorporated by reference herein. The choice of linker sequences may be determined based on crystal structure analysis of several antibody Fab molecules. There is a natural flexible linkage between the variable domain and the CH1/CL constant domain in Fab or antibody molecular structure. This natural linkage comprises approximately 10-12 amino acid residues, contributed by 4-6 residues from C-terminus of V domain and 4-6 residues from the N-terminus of the CL or CH1 domain. DVD-Ig proteins of the disclosure were generated using N-terminal 5-6 amino acid residues, or 11-12 amino acid residues, of CL or CH1 as the linker in the light chain and the heavy chain of the DVD-Ig protein, respectively. The N-terminal residues of the CL or the CH1 domains, particularly the first 5-6 amino acid residues, adopt a loop conformation without strong secondary structure, and therefore can act as flexible linkers between the two variable domains. The N-terminal residues of the CL or CH1 domains are natural extensions of the variable domains, as they are part of the Ig sequences, and therefore immunogenicity potentially arising from the linkers or junctions is minimized. Further examples of linkers are described in Table 65 (see underlined amino acids).

Other linker sequences may include a sequence of any length of the CL or CH1 domain but not all residues of a CL/CH1 domain; for example the first 5-12 amino acid residues of the CL or CH1 domain; the light chain linkers can be from C $\kappa$  or C $\lambda$ ; and the heavy chain linkers can be derived from CH1 of any isotype, including C $\gamma$ 1, C $\gamma$ 2, C $\gamma$ 3, C $\gamma$ 4, C $\alpha$ 1, C $\alpha$ 2, C $\delta$ , C $\epsilon$ , and C $\mu$ . Linker sequences may also be derived from other proteins such as Ig-like proteins, (e.g., TCR, FcR, KIR); G/S based sequences (e.g., G4S repeats); hinge region-derived sequences; and other natural sequences from proteins.

In certain embodiments, a constant domain is linked to the two linked variable domains using recombinant DNA techniques. For example, a sequence comprising linked heavy chain variable domains is linked to a heavy chain constant domain and sequence comprising linked light chain variable domains is linked to a light chain constant domain. In certain embodiments, the constant domains are a human heavy chain constant domain and a human light chain constant domain, respectively. In certain embodiments, the DVD-Ig heavy chain is further linked to an Fc region. The Fc region may comprise a native Fc region sequence, or a variant Fc region sequence. In certain embodiments, the Fc region is a human

Fc region. For example, the Fc region comprises an Fc region from an IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, or IgD.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure is a dual-specific tetravalent binding protein. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds CD20 and CD80. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds VEGF and HER2. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds TNF and RANKL. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds TNF and DKK. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds CD20 and RANKL. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds DLL4 and PLGF. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds DLL4 and VEGF. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds TNF and SOST. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds IL-9 and IgE. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds IL-12 and IL-18. An example of an IL-12 and IL-18 DVD-Ig protein is described in U.S. Patent No. 7,612,181. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds TNF and IL-17. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds TNF and PGE2. Examples of PGE2 DVD-Ig proteins are provided in U.S. Patent Publication No. 20100074900. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds IL-1 $\alpha$  and IL-1 $\beta$ . An example of an IL-1 $\alpha$  and IL-1 $\beta$  DVD-Ig protein is described in U.S. Patent No. 7,612,181. In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure binds IL-4 and IL-1. An example of an IL-4 and IL-13 DVD-Ig protein is described in U.S. Publication No. 20100226923. The amino acid and nucleic acid sequences of the DVD-Ig proteins described in the aforementioned patents and patent applications are incorporated by reference herein. Amino acid sequences of DVD-Ig proteins that may be used in the methods and compositions of the disclosure are described in SEQ ID NOs: 28-75.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of

the disclosure specifically binds CD20/CD80, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 30 and 31. In certain embodiments, the anti-CD20/CD80 DVD-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 30 and 31. In certain embodiments, the anti-CD20/CD80 DVD-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 30 and 31.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds CD80/CD20, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 32 and 33. In certain embodiments, the anti-CD80/CD20 DVD-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 32 and 33. In certain embodiments, the anti-CD80/CD20 DVD-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 32 and 33.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds VEGF/HER2, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 34 and 35. In certain embodiments, the anti-VEGF/HER2-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 34 and 35. In certain embodiments, the anti-VEGF/HER2 DVD-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 34 and 35.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds HER2/VEGF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 36 and 37. In certain embodiments, the anti-HER2/VEGF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 36 and 37. In certain embodiments, the anti-HER2/VEGF DVD-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 36 and 37.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds TNF/RANKL, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 38 and 39. In certain embodiments, the anti-TNF/RANKL-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 38 and

39. In certain embodiments, the anti-TNF/RANKL DVD-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 38 and 39.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds RANKL/TNF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 40 and 41. In certain embodiments, the anti-RANKL/TNF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 40 and 41. In certain embodiments, the anti-RANKL/TNF DVD-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 40 and 41.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds TNF/DKK, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 42 and 43. In certain embodiments, the anti-TNF/DKK-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 42 and 43. In certain embodiments, the anti-TNF/DKK-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 42 and 43.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds DKK/TNF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 44 and 45. In certain embodiments, the anti-DKK/TNF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 44 and 45. In certain embodiments, the anti-DKK/TNF-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 44 and 45.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds CD20/RANKL, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 46 and 47. In certain embodiments, the anti-CD20/RANKL-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 46 and 47. In certain embodiments, the anti-CD20/RANKL-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 46 and 47.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds RANKL/CD20, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 48 and 49. In certain embodiments, the anti-RANKL/CD20-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 48 and 49. In certain embodiments, the anti-RANKL/CD20-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 48 and 49.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds DLL4/PLGF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 50 and 51. In certain embodiments, the anti-DLL4/PLGF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 50 and 51. In certain embodiments, the anti-DLL4/PLGF-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 50 and 51.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds PLGF/DLL4, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 52 and 53. In certain embodiments, the anti-PLGF/DLL4-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 52 and 53. In certain embodiments, the anti-PLGF/DLL4-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 52 and 53.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds TNF/SOST (S2), and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 54 and 55. In certain embodiments, the anti-TNF/SOST (S2)-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 54 and 55. In certain embodiments, the anti-TNF/SOST (S2)-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 54 and 55.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds SOST (S2)/TNF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 56 and 57. In certain embodiments, the anti-SOST (S2)/TNF-Ig protein comprises amino acid sequences

corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 56 and 57. In certain embodiments, the anti-SOST (S2)/TNF-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 56 and 57.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds IL-9 (S2)/IgE, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 58 and 59. In certain embodiments, the anti-IL-9 (S2)/IgE-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 58 and 59. In certain embodiments, the anti-IL-9 (S2)/IgE-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 58 and 59.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds IgE/IL-9 (S2), and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 60 and 61. In certain embodiments, the anti-IgE/IL-9 (S2)-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 60 and 61. In certain embodiments, the anti-IgE/IL-9 (S2)-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 60 and 61.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds TNF/IL-17, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 62 and 63. In certain embodiments, the anti-TNF/IL-17-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 62 and 63. In certain embodiments, the anti-TNF/IL-17-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 62 and 63.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds TNF/PGE2, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 64 and 65. In certain embodiments, the anti-TNF/PGE2-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 64 and 65. In certain embodiments, the anti-TNF/PGE2-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 64 and 65.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of

the disclosure specifically binds IL-1 $\alpha$ /IL-1 $\beta$ , and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 66 and 67. In certain embodiments, the anti-IL-1 $\alpha$ /IL-1 $\beta$ -Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 66 and 67. In certain embodiments, the anti-IL-1 $\alpha$ /IL-1 $\beta$ -Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 66 and 67.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds DLL4/VEGF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 28 and 29. In certain embodiments, the anti-DLL4/VEGF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 28 and 29. In certain embodiments, the anti-DLL4/VEGF-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 28 and 29.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds DLL4/VEGF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 72 and 73. In certain embodiments, the anti-DLL4/VEGF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 72 and 73. In certain embodiments, the anti-DLL4/VEGF-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 72 and 73.

In certain embodiments, the DVD-Ig protein used in the methods and compositions of the disclosure specifically binds IL12/IL18, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 70 and 71. In certain embodiments, the anti-IL12/IL18-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 70 and 71. In certain embodiments, the anti-IL12/IL18-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 70 and 71.

#### D. Expression of DVD-Ig Proteins

DVD-Ig proteins of the present disclosure may be produced by any of a number of techniques known in the art. For example, expression from host cells, wherein expression vector(s) encoding the DVD heavy and DVD light chains is (are) transfected into a host cell

by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, *e.g.*, electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like.

Mammalian host cells for expressing the recombinant antibodies of the disclosure include Chinese Hamster Ovary (CHO cells) (including dhfr-CHO cells, described in Urlaub and Chasin (1980) Proc. Natl. Acad. Sci. USA 77:4216-4220, used with a DHFR selectable marker, *e.g.*, as described in Kaufman and Sharp (1982) J. Mol. Biol. 159:601-621) and DG44 or DUXB11 cells (Urlaub *et al.* (1986) Som. Cell Molec. Genet. 12:555; Haynes *et al.* (1983) Nuc. Acid. Res. 11:687-706; Lau *et al.* (1984) Mol. Cell. Biol. 4:1469-1475), NS0 myeloma cells, monkey kidney line (*e.g.*, CVI and COS, such as a COS 7 cell), SP2 cells, human embryonic kidney (HEK) cells, such as a HEK-293 cell, Chinese hamster fibroblast (*e.g.*, R1610), human cervical carcinoma (*e.g.*, HELA), murine fibroblast (*e.g.*, BALBc/3T3), murine myeloma (P3x63-Ag3.653; NS0; SP2/O), hamster kidney line (*e.g.*, HAK), murine L cell (*e.g.*, L-929), human lymphocyte (*e.g.*, RAJI), human kidney (*e.g.*, 293 and 293T). Host cell lines are typically commercially available (*e.g.*, from BD Biosciences, Lexington, Ky.; Promega, Madison, Wis.; Life Technologies, Gaithersburg, Md.) or from the American Type Culture Collection (ATCC, Manassas, Va.).

When recombinant expression vectors encoding DVD-Ig proteins are introduced into mammalian host cells, the DVD-Ig proteins are produced by culturing the host cells for a period of time sufficient to allow for expression of the DVD-Ig proteins in the host cells or secretion of the DVD-Ig proteins into the culture medium in which the host cells are grown. DVD-Ig proteins can be recovered from the culture medium using standard protein purification methods.

In an exemplary system for recombinant expression of DVD-Ig proteins, a recombinant expression vector encoding both the DVD-Ig heavy chain and the DVD-Ig light chain is introduced into dhfr-CHO cells by calcium phosphate-mediated transfection. Within the recombinant expression vector, the DVD-Ig heavy and light chain cDNAs are each operatively linked to CMV enhancer/AdMLP promoter regulatory elements to drive high levels of transcription of the cDNAs. The recombinant expression vector also carries cDNA encoding DHFR, which allows for selection of CHO cells that have been transfected with the vector using methotrexate selection/amplification. The selected transformant host cells are

cultured to allow for expression of the DVD-Ig heavy and light chains and intact DVD-Ig protein is recovered from the culture medium. Standard molecular biology techniques are used to prepare the recombinant expression vector, transfect the host cells, select for transformants, culture the host cells and recover the DVD-Ig protein from the culture medium. Still further, the disclosure provides a method of synthesizing a DVD-Ig protein of the disclosure by culturing a host cell of the disclosure in a suitable culture medium until a DVD-Ig protein of the disclosure is synthesized. The method can further comprise isolating the DVD-Ig protein from the culture medium. An important feature of DVD-Ig protein is that it can be produced and purified in a similar way as a conventional antibody.

E. Methods for Identifying Aqueous Stable DVD-Ig (AS-DVD-Ig) Proteins and Lyophilized Stable DVD-Ig (LS-DVD-Ig) Proteins

An unexpected and surprising finding is that a certain subset of DVD-Ig proteins (referred to as AS-DVD-Ig protein and LS-DVD-Ig proteins) are stable – even at high concentrations - in aqueous formulations, while a large number of DVD-Ig proteins are unstable and prone to aggregation. In addition, while the majority of DVD-Ig proteins have been found not to be stable in a lyophilized state, a certain subset of DVD-Ig proteins (referred to as LS-DVD-Ig proteins) are stable and can be successfully lyophilized using the formulations of the disclosure. Notably, DVD-Ig proteins identified as AS-DVD-Ig proteins are also LS-DVD-Ig proteins given that AS-DVD-Ig proteins are generally more stable than LS-DVD-Ig proteins. In addition, a non-AS-DVD Ig protein can be an LS DVD-Ig protein. The distinction between the two subpopulations may be based on the level of aggregation, freeze/thaw characteristics, or monomer content following storage, as described in the below assays. Generally, an increase in aggregation indicates a decrease in monomer content.

Thus, in certain embodiments, the disclosure comprises a method for distinguishing between AS-DVD-Ig proteins and non-AS-DVD-Ig proteins. The disclosure also comprises a method for distinguishing between LS-DVD-Ig proteins and non-LS-DVD-Ig proteins. In another alternative, the disclosure provides methods for distinguishing AS-DVD-Ig proteins from LS-DVD-Ig proteins. Following identification, AS-DVD-Ig protein and LS-DVD-Ig proteins may be successfully formulated in the compositions of the disclosure, while non-AS-DVD-Ig protein and non-LS-DVD-Ig proteins fail to remain stable in such formulations and are prone to aggregation and/or loss of monomer content. Generally, an increase in

aggregation indicates a decrease in monomer content.

In certain embodiments, a freeze/thaw (F/T) (e.g., - 80 °C / 30 °C) test may be used to identify DVD-Ig proteins that are AS-DVD-Ig proteins or LS-DVD-Ig proteins. Such a method relies upon determining the percentage of high molecular weight (HMW) species in a solution having a high concentration of DVD-Ig protein (e.g., 100 mg/ml). In one embodiment, an AS-DVD-Ig protein or an LS-DVD-Ig protein is defined as a DVD-Ig protein that shows 1% or less increase in high molecular weight (HMW) species (aggregates) (e.g., 0.5% or less aggregate) or less than a 1 % increase (e.g., 0.5%) in change in relative % monomer as determined by SEC following a F/T cycle. A F/T cycle includes freezing the DVD-Ig protein solution, thawing the solution, optionally repeating, and testing the thawed solution for aggregate levels using, e.g., SEC analysis. Freeze/thaw testing may be performed on a solution comprising a DVD-Ig protein at a concentration of about 1 mg/ml to about 10 mg/ml. In certain embodiments, the F/T testing includes measuring aggregation or alternatively change in monomer content by SEC after 4 freeze/thaw cycles from about 30 °C to – 80 °C. If testing results in more than about a 1 % increase in relative percent aggregates or change in relative % monomer after 4 cycles, then the DVD-Ig protein would be considered a non- LS DVD Ig.

In certain embodiments, the unfolding temperature of a DVD-Ig protein may be used to determine whether the DVD-Ig protein is an AS-DVD-Ig protein or an LS-DVD-Ig protein. As described in Figure 4, a DVD-Ig protein that shows a temperature of unfolding of less than about 45 °C is generally not characterized as an AS-DVD-Ig protein or an LS-DVD-Ig protein (“A” as described in Figure 4). In certain other embodiments, a DVD-Ig protein that shows a temperature of unfolding of about 45-50 °C is unlikely to be an AS-DVD-Ig protein but may be an LS-DVD-Ig protein (“B” as described in Figure 4). In certain other embodiments, a DVD-Ig protein that shows a temperature of unfolding that is higher than 50 °C is generally considered to be an AS-DVD-Ig protein and an LS-DVD-Ig protein (“C” as described in Figure 4). Tests to determine DVD-Ig protein unfolding temperatures are known in the art (see also Example 1 describing thermodynamic testing) and are generally performed at a pH of between about 5 and about 7, e.g., about 5.5 to about 6.5.

In order to determine whether a DVD-Ig protein is an AS-DVD-Ig protein or an LS-DVD-Ig, storage testing of the solution can be performed. For example, storage testing may be performed at 5 °C or 40 °C for 14 to 21 days at a DVD-Ig protein in solution at a

concentration ranging from 1 to 100 mg/ml, *e.g.*, about 50-100 mg/ml.

Testing at temperatures greater than ambient temperatures, *e.g.*, 40 °C, are often referred to as accelerated conditions. In certain embodiments, the accelerated storage conditions include storing the DVD-Ig protein in the absence of light at 40 °C. In certain embodiments, testing is based on a solution's DVD-Ig protein aggregation levels at a high temperature (*e.g.*, 40 °C) and a high concentration (*e.g.*, 50 mg/ml) as determined by SEC. For example, the DVD-Ig protein may be formulated at a concentration of at least about 50 mg/ml, *e.g.*, 50 to 100 mg/ml, in an aqueous formulation using a citrate phosphate buffer or a histidine buffer, and stored under accelerated conditions. Accelerated conditions may include temperatures higher than room temperature, *e.g.*, storage temperatures of about 35 to about 45 °Celsius (C), for extended periods of time, *e.g.*, about 10 to about 21 days. In certain embodiments, the accelerated storage conditions used to screen for an AS-DVD-Ig protein or LS-DVD-Ig protein are 14 days of storage at a temperature of 40 °C at a DVD-Ig protein concentration of 50 mg/ml or greater, *e.g.*, about 60 mg/ml or 50-100 mg/ml. Following accelerated storage testing at a concentration of 50 mg/ml or greater, *e.g.* 50-100 mg/ml, the solution may be tested for signs of DVD-Ig protein aggregation or change in monomer content.

In certain embodiments, DVD-Ig proteins may be tested to determine whether they are an AS-DVD-Ig protein or an LS-DVD-Ig protein in buffered solutions (*e.g.*, histidine or citrate / phosphate buffer) having a pH of 5.0 to 6.5, *e.g.*, a pH of about 6, and a concentration of about 50 mg/ml to about 100 mg/ml.

Notably, lower levels of DVD-Ig protein concentration (*e.g.*, 1 mg/ml) may also be used to test the protein, wherein lower levels of aggregate would be expected for an AS-DVD-Ig protein or an LS-DVD-Ig. For example, an AS-DVD-Ig protein is a DVD-Ig that has 3% or less aggregation when stored at about 40 °C after 21 days at a concentration of 1 mg/ml in an aqueous formulation.

Protein aggregation may be determined according to methods known in the art, including, but not limited to, Size Exclusion Chromatography (SEC). Generally, an increase in aggregation indicates a decrease in monomer content, which can also be determined using SEC analysis.

In certain embodiments, the DVD-Ig protein is considered an AS-DVD-Ig protein if the solution has 10% or less aggregation of the DVD-Ig protein as determined by Size

Exclusion Chromatography (SEC) analysis following accelerated storage at a concentration of 1-100 mg/ml, preferably about 50 mg/ml to about 100 mg/ml. In certain embodiments, the DVD-Ig protein is considered an AS-DVD-Ig protein if the solution has 6% or less aggregation of the DVD-Ig protein as determined by SEC analysis following accelerated storage at a concentration of about 50 mg/ml to about 100 mg/ml. In certain embodiments, the DVD-Ig protein is considered an AS-DVD-Ig protein if the DVD-Ig protein has less than 10%, alternatively less than 9%, less than 8 %, less than 7%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, or less than 1% aggregation as determined by SEC analysis following accelerated storage at a concentration of about 50 mg/ml to about 100 mg/ml. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig that has less than 8% aggregation following 14 days of accelerated storage (at, for example, about 40 °C). In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig that has 6% or less aggregation following 14 days of accelerated storage (at, for example, about 40 °C). In a further embodiment, an AS-DVD-Ig protein is defined as a DVD-Ig that has less than 5% aggregation following 14 days of accelerated storage (at, for example, about 40 °C). In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig that has 10% or less aggregation at about 40 °C after 21 days of storage at a concentration of 100 mg/ml in an aqueous formulation or has 10% or less aggregate following accelerated storage after 14 days at about 40 °C, when formulated at a concentration of 50 mg/ml or more in an aqueous formulation.

In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has 10% or less aggregation at about 40 °C after 21 days of storage at a concentration of 100 mg/ml in an aqueous formulation or, alternatively, a DVD-Ig protein that has 1% or less aggregation at about 5 °C after 21 days of storage at a concentration of 100 mg/ml in an aqueous formulation. Alternatively, an AS-DVD-Ig protein is a DVD-Ig protein that has 1.5% or less aggregation at about 5 °C after 21 days of storage at a concentration of 1 mg/ml in an aqueous formulation or 3% or less aggregation at about 40 °C after 21 days of storage at a concentration of 1 mg/ml in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has less than 10% aggregate formed following accelerated storage after 14 days at about 40 °C, when formulated at a concentration over 50 mg/ml in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has less than 10% aggregation following

14 days of accelerated storage (at, for example, about 40 °C). In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has less than 8% aggregation following 14 days of accelerated storage (at, for example, about 40 °C). In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has 6% or less aggregation following 14 days of accelerated storage (at, for example, about 40 °C). In a further embodiment, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has less than 5% aggregation following 14 days of accelerated storage (at, for example, about 40 °C). Aggregation can be determined according to methods known in the art, including, but not limited to, size exclusion chromatography (SEC). In certain embodiments, the accelerated storage conditions include storing the DVD-Ig protein in the absence of light at 40 °C. DVD-Ig proteins may be tested in aqueous formulations containing citrate and phosphate buffer, or histidine buffer. In certain embodiments, an AS-DVD-Ig protein has 10% or less aggregation as determined by SEC analysis following accelerated storage for 21 days at about 40 °C, where the AS-DVD-Ig protein is formulated at a concentration of 50 to 100 mg/ml in a citrate phosphate buffer or histidine buffer in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein has less than 6% aggregation as determined by SEC analysis following accelerated storage for 14 days at about 40 °C, where the AS-DVD-Ig protein is formulated at a concentration of at least 50 mg/ml in a citrate phosphate buffer or histidine buffer in an aqueous formulation.

While percent aggregation may be used to determine whether aggregation is present following accelerated storage, monomer content of the DVD-Ig protein solution may also be used. Generally, an increase in percent aggregation indicates a decrease in monomer content. Alternatively, a DVD-Ig protein may be considered an AS-DVD-Ig protein if the protein has 6% or less monomer loss (determined by SEC) after 14 days at 40 °C or 3% or less monomer loss (determined by SEC) after 7 days at 40 °C in a solution having a concentration of 50 mg/ml or more DVD-Ig protein at pH about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0 in 15 mM histidine. Monomer content may be used under any testing conditions, including, but not limited to, storage at 40 °C and/or at a pH of about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0.

In certain embodiments, an AS-DVD-Ig protein is a DVD-Ig protein that has about 10% relative (rel.) peak area or less change in monomers at about 40 °C after 21 days of storage at a concentration of about 100 mg/ml in an aqueous formulation or, alternatively, a

DVD-Ig protein that has about 1% rel. peak area or less change in monomers at about 5 °C after 21 days of storage at a concentration of about 100 mg/ml and at a pH between about 5.5 to about 6.5 in an aqueous formulation. Alternatively, an AS-DVD-Ig protein is a DVD-Ig protein that has about 1.5% rel. peak area or less change in monomers at about 5 °C after 21 days of storage at a concentration of about 1 mg/ml in an aqueous formulation or about 3% rel. peak area or less change in monomers at about 40 °C after 21 days of storage at a concentration of 1 mg/ml and at a pH between about 5.5 to about 6.5 in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has a change in monomers less than about 10% rel. peak area following accelerated storage after 14 days at about 40 °C, when formulated at a concentration over about 50 mg/ml and at a pH between about 5.5 and about 6.5 in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has less than 8% rel. peak area change in monomers following 14 days of accelerated storage (at, for example, about 40 °C) when formulated at a concentration over 60 mg/ml and at a pH between 5.5 – 6.5 in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has 6% rel. peak area or less change in monomers following 14 days of accelerated storage (at, for example, about 40 °C). In a further embodiment, an AS-DVD-Ig protein is defined as a DVD-Ig protein that has less than 5% rel. peak area change in monomers following 14 days of accelerated storage (at, for example, about 40 °C).

In certain embodiments, an AS-DVD-Ig protein has 10% rel. peak area or less change in monomers as determined by SEC analysis following accelerated storage for 21 days at about 40 °C, where the AS-DVD-Ig protein is formulated at a concentration of 50 to 100 mg/ml in a citrate phosphate buffer or histidine buffer at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0, in an aqueous formulation. In certain embodiments, an AS-DVD-Ig protein has less than 6% rel. peak area change in monomers as determined by SEC analysis following accelerated storage for 14 days at about 40 °C, where the AS-DVD-Ig protein is formulated at a concentration of at least 50 mg/ml in a citrate phosphate buffer or histidine buffer in an aqueous formulation.

In another alternative, AS-DVD-Ig proteins are identified based on a solution's stability aggregation and/or monomer content at a low temperature (*e.g.*, 5° C) and a high concentration (*e.g.*, 50 mg/ml) of DVD-Ig as determined by SEC. For example, a solution containing 50 mg/ml of an AS-DVD-Ig protein at a pH of about 5.0 to about 6.5, *e.g.*, about

5.5 to about 6.0 in 15 mM histidine may have 1% or less monomer (determined by SEC) loss after 7 days at 5 °C (determined by SEC). In another example, a solution containing 50 mg/ml of an AS-DVD-Ig protein at a pH of about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0 in 15 mM histidine may have 2% or less monomer loss after 14 days at 5 °C. Alternatively, an AS-DVD-Ig has 1% or less aggregation at about 5 °C after 21 days of storage at a concentration of 100 mg/ml in an aqueous formulation, or 1.5% or less aggregation at about 5°C after 21 days of storage at a concentration of 1 mg/ml in an aqueous formulation. In certain embodiments, monomer loss is determined at a pH of about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0.

In certain embodiments, the test solution conditions described herein also contain 0.02% (w/v) sodium azide as a bacteriostatic.

In certain embodiments, the DVD-Ig protein is considered an LS-DVD-Ig protein if the solution has 15% or less aggregation of the DVD-Ig protein as determined by Size Exclusion Chromatography (SEC) analysis. In certain embodiments, the LS-DVD-Ig protein is considered an LS-DVD-Ig protein if the DVD-Ig protein has 15% or less, alternatively less than 14%, less than 13%, less than 12%, less than 11%, less than aggregation as determined by SEC analysis.

In certain embodiments, an LS-DVD-Ig protein is defined as having less than 15% aggregate formed, following accelerated storage, *e.g.*, 14 days of accelerated storage at about 40 °C, when formulated at a concentration over 50 mg/ml in an aqueous formulation. In certain embodiments, an LS-DVD-Ig protein has 15% or less aggregation following 14 days of accelerated storage at, for example, 40 °C. In certain embodiments, an LS-DVD-Ig protein has less than 14% aggregation following 14 days of accelerated storage, at, for example, about 40 °C. In a further embodiment, an LS-DVD-Ig protein has less than 13% aggregation following 14 days of accelerated storage, at, for example, about 40 °C. Alternatively, an LS-DVD-Ig protein is defined as a DVD-Ig protein that has 1% or less aggregation following 4 freeze thaw cycles. DVD-Ig proteins may be tested in aqueous formulations containing citrate and phosphate buffer, or histidine buffer. In certain embodiments, an LS-DVD-Ig protein has 15% or less aggregation as determined by SEC analysis following accelerated storage for 14 days at about 40 °C, where the LS-DVD-Ig protein is formulated at a concentration of at least 50 mg/ml in a citrate phosphate buffer in an aqueous formulation. As described above, aggregation can be determined according to

methods known in the art, including, but not limited to, SEC.

In certain embodiments, an LS-DVD-Ig protein has more than 10% rel. peak area change in monomers observed, following accelerated storage, *e.g.*, 21 days at about 40 °C, when formulated at a concentration of about 50 to 100 mg/ml at a pH between about 5.0 and about 6.5, *e.g.*, about 5.5 to about 6.0, in an aqueous formulation. In certain embodiments, an LS-DVD-Ig protein has 50% rel. peak area or less change in monomers as determined by SEC analysis following accelerated storage for 14 days at about 40 °C, where the LS-DVD-Ig protein is formulated at a concentration of at least 50 mg/ml in a citrate phosphate buffer in an aqueous formulation. In certain embodiments, an LS-DVD-Ig protein has 20 % rel. peak area or less change in monomers following 21 days of accelerated storage at about 40 °C, when formulated at a concentration over 100 mg/ml at a pH between about 5.0 to about 6.5, *e.g.*, about 5.5 to about 6.0 in an aqueous formulation. In certain embodiments, an LS-DVD-Ig protein has less than 18% rel. peak area change in monomers following 14 days of accelerated storage, at, for example, about 40 °C. In a further embodiment, an LS-DVD-Ig protein has less than 13% rel. peak area change in monomers following 14 days of accelerated storage, at, for example, about 40 °C. Alternatively, an LS-DVD-Ig protein is defined as a DVD-Ig protein that has 1% rel. peak area or less change in monomers following 4 freeze thaw cycles. Alternatively, an LS-DVD-Ig protein is defined as a DVD-Ig protein that has 4% rel. peak area or less change in monomers following 7 days at about 25 °C at a concentration between 1-100 mg/mL, *e.g.*, about 1 to 10 mg/ml or about 50 to 100 mg/ml, in aqueous solution at the most stable pH. Alternatively, an LS-DVD-Ig protein is defined as a DVD-Ig protein that has 1 % rel. peak area or less change in monomers following 7 days at about 5 °C in aqueous solution at the most stable

Notably, the assays described herein used to determine whether a DVD-Ig protein is an AS-DVD-Ig protein or an LS-DVD-Ig protein are not exclusive, meaning that, for example, an AS-DVD-Ig may be characterized as having an unfolding temperature of at least 50 °C and also have about a 1% relative peak area or less change in monomers at about 5 °C after 21 days of storage at a concentration of about 50 to about 100 mg/ml and at a pH between about 5.5 to about 6.5 in an aqueous, buffered formulation. In certain embodiments, an AS-DVD-Ig protein has 10% rel. peak area or less change in monomers as determined by SEC analysis following accelerated storage for 21 days at about 40 °C and has an unfolding temperature of at least 50 °C.

Once the DVD-Ig protein is identified as being an AS-DVD-Ig protein or LS-DVD-Ig protein according to the aforementioned tests, the AS-DVD-Ig protein or LS-DVD-Ig protein can be stably formulated. Further identification of AS-DVD-Ig protein and LS-DVD-Ig proteins is described below in Example 4.

### **III. Aqueous Stable Dual Variable Domain Immunoglobulin (AS-DVD-Ig) Formulations of the Disclosure**

The disclosure provides stable aqueous formulations comprising AS-DVD-Ig proteins. The present disclosure features aqueous formulations having improved properties as compared to art-recognized formulations, in that AS-DVD-Ig proteins can be stably formulated, even at high concentrations.

Thus, the disclosure is based, at least in part, on the discovery that a subpopulation of DVD-Ig proteins can be stably formulated in an aqueous formulation having a pH of about 4.5 to about 7.5, and containing a buffer, a surfactant, and/or a polyol. These “Aqueous Stable DVD-Ig proteins” are referred to as AS-DVD-Ig proteins and can be identified using an accelerated storage assay where the DVD-Ig protein is formulated in a liquid form at a concentration greater than 50 mg/ml, *e.g.*, 50 mg/ml to 100 mg/ml (see also Section II.E. above).

In certain embodiments, the aqueous formulation of the disclosure has a pH of about 4.5 to about 7.5. As described in the working examples, pH was found to have an impact on the stability of the AS-DVD-Ig protein in a buffered formulation. In certain embodiments, the pH of the formulation containing the AS-DVD-Ig protein ranges from about 4.5 to about 7.5; alternatively, the pH of the AS-DVD-Ig protein formulation ranges from about 5.0 to about 7.0; alternatively the pH may range from about 5 to about 6.5; alternatively the pH of the formulation may range from about 5.5 to about 6.5. In a further embodiment, the pH ranges from about 5.8 to about 6.2. The ranges intermediate to the aforementioned pH values, *e.g.*, about 5.6 to about 6.4, are also intended to be part of the disclosure. Ranges of values using a combination of any of the aforementioned values as upper / lower limits are also intended to be included, *e.g.*, a pH range of about 5.5 to about 6.2. In certain embodiments, the pH of the formulation of the disclosure is about 6.0.

In certain embodiments, the aqueous formulation of the disclosure includes an AS-DVD-Ig protein and a buffer. Examples of buffers that may be used in the formulation of the

disclosure include, but are not limited to, acetate, histidine, glycine, arginine, phosphate, Tris, and citrate. The molarity of the buffer used in the formulation of the disclosure may range from about 1 to about 50 mM. In certain embodiments, the aqueous formulation of the disclosure has a buffer with a molarity of about 5 to about 50 mM. Alternatively, the molarity of the buffer is about 10 to about 20 mM.

In certain embodiments of the disclosure, the buffer system comprises about 1 to about 200 mM histidine (*e.g.*, about 2 to about 100 mM; about 5 to about 70 mM; about 5 to about 60 mM; about 5 to about 50 mM; about 10 to about 40 mM, about 10 to about 30 mM, or about 10 to about 20 mM) with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In certain embodiments, the buffer system of the disclosure comprises about 15 mM histidine with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5.

In certain embodiments, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 40 mM) glycine with a pH of about 4.5 to about 7.5. In a particular embodiment, the buffer system comprises glycine at a concentration of about 20 mM. In a more particular embodiment, the buffer system comprises glycine at a concentration of about 20 mM, and glycerol at a concentration of about 20 to about 30 mg/ml, *e.g.*, about 26 mg/ml, with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5.

In certain embodiments, the buffer system comprises about 1 to about 50 mM acetate (*e.g.*, about 5 to about 50 mM, about 2 to about 40 mM; about 5 to about 30 mM; or about 2 to about 15 mM) with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises acetate at a concentration of about 2 to about 15 mM.

In certain embodiments of the disclosure, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM, about 2 to about 40 mM; about 5 to about 30 mM; or about 2 to about 15 mM) arginine with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises arginine at a concentration of about 15 mM.

In still another embodiment of the disclosure, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM) citrate with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the

buffer system comprises citrate at a concentration of about 15 mM.

In still another embodiment of the disclosure, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM) phosphate with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises phosphate at a concentration of about 10 mM. In a one embodiment, the buffer system comprises phosphate at a concentration of about 10 mM, and sodium chloride at a concentration of about 125 mM.

In certain embodiments, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM) Tris with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises Tris at a concentration of about 2 to about 10 mM.

In yet another embodiment, the buffer system comprises phosphate and citrate, *e.g.*, phosphate (*e.g.*, sodium hydrogen phosphate) at a concentration of about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM, about 5 to about 10 mM), and citrate (citric acid) at a concentration of about 1 to about 50 mM (*e.g.*, about 5 to about 10 mM). In a particular embodiment, the buffer system comprises phosphate at a concentration of about 5 mM and citrate (citric acid) at a concentration of about 5 mM. In certain embodiments, the buffer system comprises phosphate at a concentration of about 10 mM and citrate (citric acid) at a concentration of about 10 mM.

In addition to the buffer, a polyol may be added to the aqueous formulation, *e.g.*, for added stability. The polyol may be added to the formulation in an amount that may vary with respect to the desired isotonicity of the formulation. In certain embodiments, the aqueous formulation is isotonic.

Examples of polyols that may be used in the aqueous formulations of the disclosure include, but are not limited to, sorbitol, mannitol, and sucrose fructose, mannose, maltose, lactose, arabinose, xylose, ribose, rhamnose, galactose and glucose. Nonreducing sugars include sucrose, trehalose, sorbose, melezitose, raffinose, mannitol, xylitol, erythritol, threitol, sorbitol and glycerol. The amount of polyol added may also vary with respect to the molecular weight of the polyol. For example, a lower amount of a monosaccharide (*e.g.*, mannitol) may be added, compared to a disaccharide (*e.g.*, trehalose).

In certain embodiments, the concentration of a polyol such as sorbitol is about 30 to about 50 mg/ml. In a one embodiment, the composition comprises about 20 to about 60

mg/ml sorbitol, about 25 to about 55 mg/ml, about 30 to about 50 mg/ml, about 35 to about 45 mg/ml, and ranges in between, *e.g.*, about 33 to about 48 mg/ml of sorbitol.

In certain embodiments, sucrose has a concentration of about 70 to about 90 mg/ml. In certain embodiments, the composition comprises about 60 to about 100 mg/ml sucrose, about 65 to about 95 mg/ml, about 70 to about 90 mg/ml, about 75 to about 85 mg/ml, and ranges in between, *e.g.*, about 72 to about 84 mg/ml of sucrose.

In certain embodiments, the polyol is mannitol. In certain embodiments, the composition comprises about 10 to about 100 mg/ml, or about 20 to about 80, about 20 to about 70, about 30 to about 60, about 30 to about 50 mg/ml of mannitol, for example, about 10, about 20, about 30, about 40, about 50, about 60, about 70, about 80, about 90, or about 100 mg/ml.

In certain embodiments, the aqueous formulation of the disclosure includes an AS-DVD-Ig, a buffer having a molarity of about 5 to about 50 mM, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5.

In addition to the buffer, a surfactant may be added to the aqueous formulations, *e.g.*, for added stability. Exemplary surfactants include nonionic detergents such as polysorbates (*e.g.*, polysorbates 20, 80) or poloxamers (*e.g.*, poloxamer 188). In certain embodiments, the amount of surfactant added is such that it reduces aggregation of the formulated AS-DVD-Ig protein and/or minimizes the formation of particulates in the formulation and/or reduces adsorption.

In certain embodiments, the aqueous formulation contains the detergent polysorbate 80 or Tween 80. Tween 80 is a term used to describe polyoxyethylene (20) sorbitan monooleate. In certain embodiments, the formulation contains about 0.001 to about 1% polysorbate 80, or about 0.005 and about 0.05% polysorbate 80, for example, about 0.001%, about 0.005, about 0.01%, about 0.05%, or about 0.1% polysorbate 80. In certain embodiments, about 0.01% polysorbate 80 is found in the formulation of the disclosure.

In certain embodiments, the aqueous formulation of the disclosure includes an AS-DVD-Ig, a buffer having a molarity of about 5 to about 50 mM, and a surfactant, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the surfactant is a polysorbate, *e.g.*, polysorbate 80 or polysorbate 20. In certain embodiments, the polysorbate has a concentration of about 0.005% to about 0.02%.

In certain embodiments, the aqueous formulation of the disclosure includes an AS-DVD-Ig, a buffer having a molarity of about 5 to about 50 mM, a surfactant, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the formulation includes an AS-DVD-Ig, a buffer (*e.g.*, histidine), a polysorbate, *e.g.*, polysorbate 80, and a sugar alcohol, *e.g.*, mannitol or sorbitol. In certain embodiments, the formulation includes an AS-DVD-Ig, a buffer (*e.g.*, histidine), a polysorbate, *e.g.*, polysorbate 80, and a non-reducing sugar, *e.g.*, sucrose.

One advantage of the aqueous formulation of the disclosure is that high concentrations of AS-DVD-Ig proteins may be stably maintained in an aqueous solution. Thus, in certain embodiments, the formulations of the disclosure comprise a high protein concentration, including, for example, a protein concentration greater than about 10 mg/ml, greater than about 20 mg/ml, greater than about 30 mg/ml, greater than about 40 mg/ml, greater than about 50 mg/ml, greater than about 100 mg/ml, greater than about 110 mg/ml, greater than about 120 mg/ml, greater than about 130 mg/ml, greater than about 140 mg/ml, greater than about 150 mg/ml, greater than about 160 mg/ml, greater than about 170 mg/ml, greater than about 180 mg/ml, greater than about 190 mg/ml, or greater than about 200 mg/ml.

In various embodiments of the disclosure, the concentration of the AS-DVD-Ig protein in the aqueous formulation is about 0.1-250 mg/ml, about 0.5-220 mg/ml, about 1-210 mg/ml, about 5-200 mg/ml, about 10-195 mg/ml, about 15-190 mg/ml, about 20-185 mg/ml, about 25-180 mg/ml, about 30-175 mg/ml, about 35-170 mg/ml, about 40-165 mg/ml, about 45-160 mg/ml, about 50-155 mg/ml, about 55-150 mg/ml, about 60-145 mg/ml, about 65-140 mg/ml, about 70-135 mg/ml, about 75-130 mg/ml, about 80-125 mg/ml, about 85-120 mg/ml, about 90-H5 mg/ml, about 95-110 mg/ml, about 95-105 mg/ml, or about 100 mg/ml. Ranges intermediate to the above recited concentrations, *e.g.*, about 31-174 mg/ml, are also intended to be part of this disclosure. For example, ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included.

The present disclosure features aqueous formulations having improved properties as compared to art-recognized formulations. For example, the formulations of the disclosure have an AS-DVD-Ig protein aggregation level of less than 7% aggregate, less than 6% aggregate, or less than 5% aggregate.

In certain embodiments, the AS-DVD-Ig protein used in the aqueous formulation of the disclosure is an anti-TNF/IL-17 DVD-Ig protein having heavy and light chain sequences having an amino acid sequence as set forth in SEQ ID NOs: 62 and 63, respectively.

In certain embodiments, the aqueous formulation of the disclosure comprises an anti-TNF/IL-17 DVD-Ig protein (*e.g.*, an anti-TNF/IL-17 DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 62 and 63; or an anti-TNF/IL-17 DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 62 and 63; or DVD-A), histidine or acetate buffer, and sucrose or sorbitol, wherein the formulation has a pH of about 5 to about 6 and wherein the protein concentration is about 50 mg/ml or more. In further embodiments, the aqueous formulation has a pH ranging from about 5 to about 5.5, *e.g.*, about 5.2. In a further embodiment, the aqueous formulation comprises a surfactant, *e.g.*, a polysorbate. Ranges for the recited buffers, excipients, and pH are described above.

In certain embodiments, the AS-DVD-Ig protein is an anti-IL1 $\alpha$ /IL-1 $\beta$  DVD-Ig comprising an anti-IL1 $\alpha$ /IL-1 $\beta$  DVD-Ig protein having a heavy and light chain sequences having an amino acid sequence as set forth in SEQ ID NOs: 66 and 67, respectively.

In certain embodiments, the aqueous formulation of the disclosure comprises an anti-IL1 $\alpha$ /IL-1 $\beta$  DVD-Ig protein (*e.g.*, an anti-IL1 $\alpha$ /IL-1 $\beta$  DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 66 and 67; or an anti-TNF/IL-17 DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 66 and 67; or DVD-C), histidine buffer, and sucrose or sorbitol, wherein the formulation has a pH of about 5 to about 6 and wherein the protein concentration is about 50 mg/ml or more. In further embodiments, the aqueous formulation has a pH ranging from about 5 to about 5.5, *e.g.*, about 5.4. In a further embodiment, the aqueous formulation comprises a surfactant, *e.g.*, a polysorbate. Ranges for the recited buffers, excipients, and pH are described above.

In certain embodiments, the aqueous formulation of the disclosure comprises an anti-DLL4/VEGF DVD-Ig protein (*e.g.*, an anti-DLL4/VEGF DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and

light chain CDRs set forth in SEQ ID NOs: 28 and 29 or SEQ ID NOs: 72 and 73; or an anti-DLL4/VEGF DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 28 and 29 or SEQ ID NOs: 72 and 73), histidine buffer, and sucrose or arginine, wherein the formulation has a pH of about 5 to about 6 and wherein the protein concentration is about 20 mg/mL to about 50 mg/ml. In further embodiments, the aqueous formulation has a pH ranging from about 5.5 to about 6. In a further embodiment, the aqueous formulation comprises a surfactant, *e.g.*, a polysorbate. Ranges for the recited buffers, excipients, and pH are described above.

#### **IV. Lyophilized Stable Dual Variable Domain Immunoglobulin (LS-DVD-Ig) Protein Formulations of the Disclosure**

The disclosure further provides stable lyophilized formulations comprising LS-DVD-Ig proteins. Thus, the disclosure is based, at least in part, on the discovery that a subpopulation of DVD-Ig proteins can be stably formulated in a lyophilized formulation having a pH of about 4.5 to about 7.5, and containing a buffer, a surfactant, and/or a polyol. These “Lyophilized Stable DVD-Ig proteins” or “LS-DVD-Ig proteins” can be identified using an accelerated storage assay (described above) where the DVD-Ig protein is formulated in a liquid form at a concentration greater than 50 mg/ml.

The below-mentioned features of the lyophilized formulations of the disclosure may refer to the solution prior to lyophilization or, alternatively, the reconstituted formulation, for example, where liquid concentrations (mg/ml) are referenced.

In certain embodiments, the lyophilized formulation of the disclosure has a pH of about 4.5 to about 7.5. In certain embodiments, the pH of the formulation containing the LS-DVD-Ig protein ranges from about 4.5 to about 7.5; alternatively, the pH of the LS-DVD-Ig protein formulation ranges from about 5.0 to about 7.0; alternatively the pH may range from about 5 to about 6.5; alternatively the pH of the formulation may range from about 5.5 to about 6.5. In a further embodiment, the pH ranges from about 5.8 to about 6.2. The ranges intermediate to the aforementioned pH values, *e.g.*, about 5.6 to about 6.4, are also intended to be part of the disclosure. Ranges of values using a combination of any of the aforementioned values as upper / lower limits are also intended to be included, *e.g.*, a pH

range of about 5.5 to about 6.2. In certain embodiments, the pH of the formulation of the disclosure is about 6.0.

In certain embodiments, the lyophilized formulation of the disclosure includes an LS-DVD-Ig protein and a buffer. Examples of buffers that may be used in the formulation of the disclosure include, but are not limited to, acetate, histidine, glycine, arginine, phosphate, Tris, and citrate. The molarity of the buffer used in the formulation of the disclosure may range from about 1 to about 50 mM. In certain embodiments, the aqueous formulation of the disclosure has a buffer with a molarity of about 5 to about 50 mM. Alternatively, the molarity of the buffer is about 10 to about 20 mM.

In certain embodiments of the disclosure, the buffer system comprises about 1 to about 200 mM histidine (*e.g.*, about 2 to about 100 mM; about 5 to about 70 mM; about 5 to about 60 mM; about 5 to about 50 mM; about 10 to about 40 mM, about 10 to about 30 mM, or about 10 to about 20 mM) with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In certain embodiments, the buffer system of the disclosure comprises about 15 mM histidine with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5.

In certain embodiments, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 40 mM) glycine with a pH of about 4.5 to about 7.5. In a particular embodiment, the buffer system comprises glycine at a concentration of about 20 mM. In a more particular embodiment, the buffer system comprises glycine at a concentration of about 20 mM, and glycerol at a concentration of about 20 to about 30 mg/ml, *e.g.*, about 26 mg/ml, with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5.

In certain embodiments, the buffer system comprises about 1 to about 50 mM acetate (*e.g.*, about 5 to about 50 mM, about 2 to about 40 mM; about 5 to about 30 mM; or about 2 to about 15 mM) with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises acetate at a concentration of about 2 to about 15 mM.

In certain embodiments of the disclosure, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM, about 2 to about 40 mM; about 5 to about 30 mM; or about 2 to about 15 mM) arginine with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer

system comprises arginine at a concentration of about 15 mM.

In still another embodiment of the disclosure, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM) citrate with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises citrate at a concentration of about 15 mM.

In still another embodiment of the disclosure, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM) phosphate with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises phosphate at a concentration of about 10 mM. In a one embodiment, the buffer system comprises phosphate at a concentration of about 10 mM, and sodium chloride at a concentration of about 125 mM.

In certain embodiments, the buffer system comprises about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM) Tris with a pH of about 4.5 to about 7.5, *e.g.*, a pH of about 5 to about 7, or a pH of about 5.5 to about 6.5. In a particular embodiment, the buffer system comprises Tris at a concentration of about 2 to about 10 mM.

In yet another embodiment, the buffer system comprises phosphate and citrate, *e.g.*, phosphate (*e.g.*, sodium hydrogen phosphate) at a concentration of about 1 to about 50 mM (*e.g.*, about 5 to about 50 mM, about 5 to about 10 mM), and citrate (citric acid) at a concentration of about 1 to about 50 mM (*e.g.*, about 5 to about 10 mM). In a particular embodiment, the buffer system comprises phosphate at a concentration of about 5 mM and citrate (citric acid) at a concentration of about 5 mM. In certain embodiments, the buffer system comprises phosphate at a concentration of about 10 mM and citrate (citric acid) at a concentration of about 10 mM.

In addition to the buffer, a polyol may be added to the formulation, *e.g.*, for added stability. The polyol may be added to the formulation in an amount that may vary with respect to the desired isotonicity of the formulation. In certain embodiments, the lyophilized formulation is isotonic upon reconstitution.

Examples of polyols that may be used in the lyophilized formulations of the disclosure include, but are not limited to, mannitol, sucrose, trehalose and raffinose. The amount of polyol added may also vary with respect to the molecular weight of the polyol. For example, a lower amount of a monosaccharide (*e.g.*, mannitol) may be added, compared to a disaccharide (*e.g.*, trehalose).

In certain embodiments, the concentration of a polyol such as sorbitol is about 30 to about 50 mg/ml. In a one embodiment, the composition comprises about 20 to about 60 mg/ml sorbitol, about 25 to about 55 mg/ml, about 30 to about 50 mg/ml, about 35 to about 45 mg/ml, and ranges in between, *e.g.*, about 33 to about 48 mg/ml of sorbitol.

In certain embodiments, sucrose has a concentration of about 70 to about 90 mg/ml. In certain embodiments, the composition comprises about 60 to about 100 mg/ml sucrose, about 65 to about 95 mg/ml, about 70 to about 90 mg/ml, about 75 to about 85 mg/ml, and ranges in between, *e.g.*, about 72 to about 84 mg/ml of sucrose.

In certain embodiments, the polyol is mannitol. In certain embodiments, the composition comprises about 10 to about 100 mg/ml, or about 20 to about 80, about 20 to about 70, about 30 to about 60, about 30 to about 50 mg/ml of mannitol, for example, about 10, about 20, about 30, about 40, about 50, about 60, about 70, about 80, about 90, or about 100 mg/ml.

In certain embodiments, the lyophilized formulation of the disclosure includes an AS-DVD-Ig, a buffer having a molarity of about 5 to about 50 mM, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5.

In addition to the buffer, a surfactant may be added to the lyophilized formulations, *e.g.*, for added stability. Exemplary surfactants include nonionic detergents such as polysorbates (*e.g.*, polysorbates 20, 60, 80,) or poloxamers (*e.g.*, poloxamer 188). In certain embodiments, the amount of surfactant added is such that it reduces aggregation of the formulated LS-DVD-Ig protein and/or minimizes the formation of particulates in the formulation and/or reduces adsorption.

In certain embodiments, the lyophilized formulation contains the detergent polysorbate 80 or Tween 80. Tween 80 is a term used to describe polyoxyethylene (20) sorbitan monooleate. In certain embodiments, the formulation contains about 0.001 to about 0.1% polysorbate 80, or about 0.005 and about 0.05%, 20 polysorbate 80, for example, about 0.001, about 0.005, about 0.01, about 0.05, or about 0.1% polysorbate 80. In certain embodiments, about 0.01% polysorbate 80 is found in the formulation of the disclosure.

In certain embodiments, the lyophilized formulation of the disclosure includes an LS-DVD-Ig, a buffer having a molarity of about 5 to about 50 mM, and a surfactant, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the surfactant is a polysorbate, *e.g.*, polysorbate 80 or polysorbate 20. In certain embodiments, the polysorbate

has a concentration of about 0.005% to about 0.02%.

In certain embodiments, the lyophilized formulation of the disclosure includes an LS-DVD-Ig, a buffer having a molarity of about 5 to about 50 mM, a surfactant, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5. In certain embodiments, the formulation includes an LS-DVD-Ig, a buffer (*e.g.*, histidine), a polysorbate (*e.g.*, polysorbate 80), and a sugar alcohol (*e.g.*, mannitol or sorbitol). In certain embodiments, the formulation includes an LS-DVD-Ig, a buffer (*e.g.*, histidine), a polysorbate, *e.g.*, polysorbate 80, and a non-reducing sugar, *e.g.*, sucrose.

The lyophilized formulation described herein is initially made as a "pre-lyophilized formulation," which is the formulation prior to the lyophilization process. The amount of protein present in the pre-lyophilized formulation is determined taking into account the desired dose volumes, mode(s) of administration etc. For example, the starting concentration of an LS-DVD-Ig protein can be from about 2 mg/ml to about 50 mg/ml.

In certain embodiments, the DVD-Ig protein used in the lyophilized formulations of the disclosure specifically binds TNF/IL-17, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 62 and 63. In certain embodiments, the anti-TNF/IL-17-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 62 and 63. In certain embodiments, the anti-TNF/IL-17-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 62 and 63.

In certain embodiments, the lyophilized formulation of the disclosure comprises an anti-TNF/IL-17 DVD-Ig protein ((*e.g.*, an anti-TNF/IL-17 DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 62 and 63; or an anti-TNF/IL-17 DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 62 and 63; or DVD-A), histidine or acetate buffer, and sucrose or sorbitol, where the formulation has a pH of about 5 to about 6 and wherein the protein concentration is greater than about 50 mg/ml upon reconstitution..

In certain embodiments, the DVD-Ig protein used in the lyophilized formulations of the disclosure specifically binds IL1 $\alpha$ /IL-1 $\beta$  and comprises a heavy and a light chain

sequence comprising an amino acid sequence as set forth in SEQ ID NOs: 66 and 67, respectively. In certain embodiments, the DVD-Ig protein used in the lyophilized formulations specifically binds IL-1 $\alpha$ /IL-1 $\beta$ , and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 66 and 67. In certain embodiments, the anti-IL-1 $\alpha$ /IL-1 $\beta$ -Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 66 and 67. In certain embodiments, the anti-IL-1 $\alpha$ /IL-1 $\beta$ -Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 66 and 67.

In certain embodiments, the lyophilized formulation of the disclosure comprises an anti-anti-IL1 $\alpha$ /IL1 $\beta$  DVD-Ig protein (*e.g.*, an anti-IL1 $\alpha$ /IL1 $\beta$  DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 66 and 67; or an anti-TNF/IL-17 DVD-Ig protein comprising a heavy and a light chain comprising amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 66 and 67; or DVD-C), histidine buffer, and sucrose or sorbitol, where the formulation has a pH of about 5 to about 6 and wherein the protein concentration is greater than about 50 mg/ml upon reconstitution. In certain embodiments, the aqueous formulation has a pH ranging from about 5 to about 5.5, *e.g.*, about 5.4.

In certain embodiments, the DVD-Ig protein used in the lyophilized formulations of the disclosure specifically binds TNF/PGE2, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 64 and 65. In certain embodiments, the anti-TNF/PGE2-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 64 and 65. In certain embodiments, the anti-TNF/PGE2-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 64 and 65.

In certain embodiments, the lyophilized formulation of the disclosure comprises a DVD-Ig protein that specifically binds DLL4/VEGF, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 28 and 29 or SEQ ID NOs: 72 and 73. In certain embodiments, the anti-DLL4/VEGF-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 28 and 29 or SEQ ID NOs: 72 and 73. In certain embodiments, the anti-

DLL4/VEGF-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 28 and 29 or SEQ ID NOs: 72 and 73.

In certain embodiments, the lyophilized formulation of the disclosure comprises a DVD-Ig protein that specifically binds IL12/IL18, and comprises amino acid sequences corresponding to the heavy and light chain CDRs set forth in SEQ ID NOs: 70 and 71. In certain embodiments, the anti-IL12/IL18-Ig protein comprises amino acid sequences corresponding to the heavy and light chain variable regions set forth in SEQ ID NOs: 70 and 71. In certain embodiments, the anti-IL12/IL18-Ig protein comprises the amino acid sequences corresponding to the heavy and light chains set forth in SEQ ID NOs: 70 and 71.

Lyophilization may be performed according to methods known in the art. Many different freeze-dryers are available for this purpose such as Hull50<sup>TM</sup> (Hull, USA) or GT20.TM. (Leybold-Heraeus, Germany) freeze-dryers. Freeze-drying is accomplished by freezing the formulation and subsequently subliming ice from the frozen content at a temperature suitable for primary drying. Under this condition, the product temperature is below the eutectic point or the collapse temperature of the formulation. Typically, the shelf temperature for the primary drying will range from about -30 to 25°C. (provided the product remains frozen during primary drying) at a suitable pressure, ranging typically from about 50 to 250 mTorr. The formulation, size and type of the container holding the sample (*e.g.*, glass vial) and the volume of liquid will mainly dictate the time required for drying, which can range from a few hours to several days (*e.g.* 40-60 hours). Optionally, a secondary drying stage may also be performed depending upon the desired residual moisture level in the product. The temperature at which the secondary drying is carried out ranges from about 0-40°C, depending primarily on the type and size of container and the type of protein employed. For example, the shelf temperature throughout the entire water removal phase of lyophilization may be from about 15-30°C (*e.g.*, about 20°C). The time and pressure required for secondary drying will be that which produces a suitable lyophilized cake, dependent, *e.g.*, on the temperature and other parameters. The secondary drying time is dictated by the desired residual moisture level in the product and typically takes at least about 5 hours (*e.g.* 10-15 hours). The pressure may be the same as that employed during the primary drying step. Freeze-drying conditions can be varied depending on the formulation and vial size.

Prior to administration to the patient, the lyophilized formulation is reconstituted with a pharmaceutically acceptable diluent such that the protein concentration in the reconstituted formulation is at least about 2 mg/ml, for example from about 2 mg/ml to about 100 mg/ml, alternatively from about 10 mg/ml to about 90 mg/ml, alternatively from about 30 mg/ml to about 50 mg/ml. Such high protein concentrations in the reconstituted formulation are considered to be particularly useful where subcutaneous delivery of the reconstituted formulation is intended. However, for other routes of administration, such as intravenous administration, lower concentrations of the protein in the reconstituted formulation may be desired (for example from about 2-50 mg/ml, or from about 3-40 mg/ml protein in the reconstituted formulation). In certain embodiments, the protein concentration in the reconstituted formulation is significantly higher than that in the pre-lyophilized formulation. Reconstitution generally takes place at a temperature of about 25.degree C to ensure complete hydration, although other temperatures may be employed as desired. The time required for reconstitution will depend, *e.g.*, on the type of diluent, amount of excipient(s) and protein. Exemplary diluents include sterile water, bacteriostatic water for injection (BWFI), a pH buffered solution (*e.g.* phosphate-buffered saline), sterile saline solution, Ringer's solution or dextrose solution. The diluent optionally contains a preservative. Exemplary preservatives have been described above, with aromatic alcohols such as benzyl or phenol alcohol being the preferred preservatives. The amount of preservative employed is determined by assessing different preservative concentrations for compatibility with the protein and preservative efficacy testing. For example, if the preservative is an aromatic alcohol (such as benzyl alcohol), it can be present in an amount from about 0.1-2.0% and preferably from about 0.5-1.5%, but most preferably about 1.0-1.2%.

## **V.      Uses of the Disclosure**

The formulations of the disclosure may be used both therapeutically, *i.e.*, *in vivo*, or as reagents for *in vitro* or *in situ* purposes. The methods of the disclosure may also be used to make a water-based formulation having characteristics that are advantageous for therapeutic use. The aqueous formulation may be used as a pharmaceutical formulation to treat a disorder in a subject.

The formulation of the disclosure may be used to treat any disorder for which the

therapeutic protein is appropriate for treating. A "disorder" is any condition that would benefit from treatment with the protein. This includes chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question. In the case of an anti-TNF DVD-Ig, a therapeutically effective amount of the DVD-Ig protein may be administered to treat an autoimmune disease, such as rheumatoid arthritis, an intestinal disorder, such as Crohn's disease, a spondyloarthropathy, such as ankylosing spondylitis, or a skin disorder, such as psoriasis. In the case of an anti-IL-12 DVD-Ig, a therapeutically effective amount of the DVD-Ig protein may be administered to treat a neurological disorder, such as multiple sclerosis, or a skin disorder, such as psoriasis. Other examples of disorders in which the formulation of the disclosure may be used to treat include cancer, including breast cancer, leukemia, lymphoma, and colon cancer.

The term "subject" is intended to include living organisms, *e.g.*, prokaryotes and eukaryotes. Examples of subjects include mammals, *e.g.*, humans, dogs, cows, horses, pigs, sheep, goats, cats, mice, rabbits, rats, and transgenic non-human animals. In specific embodiments of the disclosure, the subject is a human.

The term "treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder, as well as those in which the disorder is to be prevented.

The aqueous formulation may be administered to a mammal, including a human, in need of treatment in accordance with known methods of administration. Examples of methods of administration include parenteral, subcutaneous, intramuscular, intravenous, intraarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitory, intracelial, intracerebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, and transdermal.

The appropriate dosage ("therapeutically effective amount") of the protein will depend, for example, on the condition to be treated, the severity and course of the condition, whether the protein is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the protein, the type of protein used, and the discretion of the attending physician. The protein is administered to the patient at one time or

over a series of treatments and may be administered to the patient at any time from diagnosis onwards. The protein may be administered as the sole treatment or in conjunction with other drugs or therapies useful in treating the condition in question.

Actual dosage levels of the AS-DVD-Ig protein or LS-DVD-Ig protein, the active ingredient, in the pharmaceutical formulation of this disclosure may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

The selected dosage level will depend upon a variety of factors including the activity of the AS-DVD-Ig protein or LS-DVD-Ig protein found in the formulation, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

In certain embodiments of the disclosure, the dosage of the AS-DVD-Ig protein in the formulation is about 1 to about 250 mg. In certain embodiments, the dosage of the AS-DVD-Ig protein in the formulation is about 30 to about 140 mg, about 40 to about 120 mg, about 50 to about 110 mg, about 60 to about 100 mg, or about 70 to about 90 mg. In a further embodiment, the composition includes an AS-DVD-Ig protein dosage of about 1, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240 or 250 mg.

In certain embodiments of the disclosure, the dosage of the LS-DVD-Ig protein in the formulation (upon reconstitution) is about 1 to about 250 mg. In a further embodiment, the dosage of the LS-DVD-Ig in the formulation is about 30 to about 140 mg, about 40 to about 120 mg, about 50 to about 110 mg, about 60 to about 100 mg, or about 70 to about 90 mg. In a further embodiment, the composition includes an LS-DVD-Ig dosage of about 1, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240 or 250 mg.

The formulations of the disclosure overcome common problems known in formulation development, including the problem of protein aggregation often associated with high concentrations of protein, particularly complex proteins such as DVD-Ig proteins. Thus,

in certain embodiments, the formulations of the disclosure provide a new means by which high levels of this new therapeutic protein format may be administered to a patient.

## EXAMPLES

The Examples presented herein describe formulations containing dual variable domain immunoglobulin (DVD-Ig) proteins that have unexpected stability characteristics. The experiments were surprising in that certain DVD-Ig proteins, referred to as Aqueous Stable DVD-Ig (AS-DVD-Ig) proteins or Lyophilized Stable DVD-Ig (LS-DVD-Ig) proteins, were stable in aqueous or lyophilized formulations, respectively, whereas other DVD-Ig proteins showed aggregation and instability when similarly formulated. The experiments exemplify methods for identifying AS-DVD-Ig proteins and LS-DVD-Ig proteins, as well as stable formulations thereof.

## Materials and Methods

The methods described herein were used in experiments performed to assess and monitor the stability of DVD-Ig proteins in aqueous and lyophilized formulations.

### *General Methods*

DVD-Ig protein formulations were tested for general quality parameters (*e.g.*, pH), parameters of physical stability (*e.g.*, clarity, color, particle contamination and purity), and parameters of chemical stability (*e.g.*, deamidation, oxidation, and general chemical stability). Exemplary tests included tests for visible particulate contamination, for light obscuration particle count for subvisible particles, and for purity such as size exclusion high pressure liquid chromatography (also referred to herein as size exclusion chromatography (SEC)) and ion exchange chromatography (IEC).

Particulate contamination (*e.g.*, visible particles) was determined by visual inspection. Subvisible particles were monitored by light obscuration assays according to the United States Pharmacopeia (USP). The physicochemical stability of formulations was assessed by SEC, which allows for the detection of fragments and aggregates. To monitor chemical stability, SEC (for the detection of fragments and hydrolysis) and IEC were performed.

*DVD-Ig Proteins Tested*

The DVD-Ig proteins that were tested in the Examples provided herein are listed in Table 1. The sequences of the DVD-Ig proteins described in Table 1 are provided in Table 65.

Table 1: Dual Variable Domain Immunoglobulin (DVD-Ig) Proteins and Their Targets

DVD-Ig Protein Name	Targets
DVD 5	CD20/CD80
DVD 6	CD80/CD20
DVD 37	VEGF/HER2
DVD 38	HER2/VEGF
DVD 53	TNF/RANKL
DVD 54	RANKL/TNF
DVD 65	TNF/DKK
DVD 66	DKK/TNF
DVD 165	CD20/RANKL
DVD 166	RANKL/CD20
DVD 257	DLL4/PLGF
DVD 258	PLGF/DLL4
DVD 277	TNF/SOST (S2)
DVD 278	SOST(S2)/TNF
DVD 281	IL-9(S2)/IgE
DVD 282	IgE/IL-9(S2)
IL12IL18	IL-12/IL-18
DVD-A	TNF/IL-17
DVD-B	TNF/PGE2
DVD-C	IL-1 $\alpha$ /IL-1 $\beta$

DVD-Ig protein as starting material was provided following purification and was > 95 % monomeric form.

*Cation Exchange HPLC Methods*

Cation exchange HPLC, a form of IEC, was used to determine the identity and purity of the DVD-Ig protein formulations. The assay was performed with the parameters detailed below.

A Dionex ProPac® WCX-10 analytical column (Dionex Corp., Sunnyvale, CA),

combined with a Dionex WCX-10G guard column (Dionex Corp., Sunnyvale, CA), was run with upper column pressure being limited to 210 bar. The mobile phase A consisted of 10 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 6.0. This buffer was created by dissolving 4.97 g anhydrous disodium hydrogen phosphate in approximately 3300 mL Milli-Q water, adjusting the pH to 7.0 using 1 M phosphoric acid, increasing the buffer volume to 3500 mL with Milli-Q water and filtering the solution through a membrane filter. The mobile phase B consisted of 10 mM Na<sub>2</sub>HPO<sub>4</sub>, 500 mM NaCl, pH 6.0. This buffer was created by dissolving 2.56 g anhydrous disodium hydrogen phosphate in approximately 1500 ml Milli-Q water, adjusting the pH to 6.0 using 1 M phosphoric acid, increasing the buffer volume to 1800 ml with Milli-Q water and filtering the solution through a membrane filter. A summary of the cation exchange HPLC methods is described in Table 2.

Table 2: Summary of Cation Exchange HPLC Methods

Item	Description/Operating Conditions	
Guard column	ProPac WCX-10G, 4.0 x 50 mm	
Column	ProPac WCX-10, 4.0 x 250 mm	
Mobile phase A <sup>*</sup>	10 mM disodium hydrogen phosphate, pH 6.0	
Mobile phase B <sup>*</sup>	10 mM disodium hydrogen phosphate/500 mM sodium chloride, pH 6.0	
Gradient	Binary Gradient	
	Time (minute)	Mobile Phase B%
	0.01	15.0
	30.00	30.0
	32.00	100.0
	37.00	100.0
	39.00	15.0
	44.00	15.0
	44.10	Stop
Flow rate	1.0 ml/minute	
Detector wavelength	280 nm	
Autosampler temperature	Nominal 4°C	
Column oven temperature	35°C	
Sample load	100 µL/100 µg	
Run time	44.0 minutes	

For the IL12IL18 and DVD 66 DVD-Ig proteins, the mobile phases used were changed as follows:

Mobile phase A: 10 mM MES, pH 5.6; and

Mobile phase B: 10 mM MES + 500 mM NaCl, pH 5.6.

For IL1 $\alpha$ IL $\beta$ , the mobile phases used were changed as follows:

Mobile phase A: 20 mM Phosphate, pH 8.0; and

Mobile phase B: 20 mM Phosphate + 500 mM NaCl, pH 8.0.

**Table 3. Gradient for IL1 $\alpha$ IL $\beta$**

Time (minutes)	% B
0	7
5	7
25	25
27	100
32	100
34	7
37	7

Similar versions of IEC were used for various other DVD-Ig proteins.

#### *Size Exclusion HPLC Methods*

Size exclusion HPLC was used to determine the purity of DVD-Ig protein solutions. The assay was performed as outlined below.

A TSK gel guard (VWR Scientific, Bridgeport, NJ; cat. no. 08543, 6.0 mm x 4.0 cm, 7  $\mu$ m), was combined with a TSK gel G3000SW (VWR Scientific, Bridgeport, NJ; cat. no. 08541, 7.8 mm x 30 cm, 5  $\mu$ m) and run with an upper column pressure limit of 70 bar. The mobile phase consisted of 100 mM Na<sub>2</sub>HPO<sub>4</sub> / 200 mM Na<sub>2</sub>SO<sub>4</sub>, pH 7.0. This buffer was created by dissolving 49.68 g anhydrous disodium hydrogen phosphate and 99.44 g anhydrous sodium sulfate in approximately 3300 ml Milli-Q water, adjusting the pH to 7.0 using 1 M phosphoric acid, increasing the buffer volume to 3500 ml with Milli-Q water and filtering the solution through a membrane filter.

The experimental parameters were listed as follows:

Flow rate: 0.3 ml/minute;

Injection volume (equivalent to 20 µg sample): 20 µl;

Column temperature: room temperature;

Autosampler temperature: 2 to 8°C;

Run time: 50 minute;

Elution: isocratic gradient.

Detection was performed using a diode array detector using a 214 nm wavelength (> 0.1 min peak width and 8 nm band width) and a 360 nm reference wavelength (100 nm band width).

Test samples were injected in duplicate. Purity was determined by comparing the area of DVD-Ig protein peak to the total area of all 214 nm absorbing components in the sample, excluding buffer-related peaks. High molecular weight aggregates and antibody fragments were resolved from intact DVD-Ig protein using this method.

#### *Freeze/Thaw Assays*

The stability of DVD-Ig protein solutions was measured using repeated freeze/thaw assays. The DVD-Ig proteins were frozen at -80°C and then thawed at 30°C in a water bath and the resulting solutions were analyzed for aggregation by SEC and/or for subvisible particle formation by light obscuration assays.

#### *Accelerated and Real Time Storage Stability Studies*

The pH and storage temperature of formulations are two important factors influencing protein stability during long-term storage of protein liquid and lyophilisate formulations. To assess the impact of these factors, the protein formulations were exposed to short-term storage at elevated temperatures, *e.g.*, 40°C, (accelerated storage) in order to mimic long term storage and quickly gain insight in the formulation feasibility for long-term storage at lower temperatures (*e.g.*, 2-8°C). To assess the real time storage stability, the samples were also kept at 2-8°C.

#### *Light Obscuration Assays*

Light obscuration assays were performed to measure the insoluble particulate content of aggregating DVD-Ig protein solutions. Light obscuration measurement equipment (particle counter, model syringe, Klotz Bad Liebenzell, Germany, series S20037) was

equipped with laminar air hood (Thermo Electron Corp., Asheville, NC; Model No. ULT2586-9-A40) to minimize foreign particle contamination during measurements. Light obscuration analysis was performed as follows. A 3.5 ml sample was placed in a 5 ml round-bottom tube under laminar air flow conditions. Measurements were performed according to manufacturer's specifications in n=3 mode (0.8 mL per single measurement), after an initial 0.8 ml rinse.

#### *Differential Scanning Calorimetry (DSC)*

Prior to DSC analysis, DVD-Ig proteins were dialyzed into a suitable buffer system using Slide-A-Lyzer Cassettes (Thermo Fisher Scientific, Waltham, MA). This buffer system (e.g., 5 mM phosphate/5 mM citrate was also used as a reference/blank for the DSC measurement. The antibody was analyzed at 1-2 mg/ml. An automated VP-Capillary DSC System (MicroCal, Piscataway, NJ) was used. Unfolding of the molecules was studied by applying a 1°C/minute scan rate over a 25°C - 95°C temperature range. Other measurement parameters were as follows: fitting period: 16 seconds; pre-scan wait: 10 minutes; feedback mode: none.

#### *Air/Liquid Interface Shaking Studies*

Shaking studies were conducted at a concentration of 1 mg/ml in 6R glass vials on an HS 260 IKA shaker (Wilmington, NC) at a speed of 150 rpm (revolutions per minute). The optical density of samples was evaluated following shaking for various periods. Similarly, SEC was also done for samples pulled at various time points.

#### *PEG Solubility*

PEG 3000 was used for solubility studies. A 50 % w/v solution of PEG 3000 was prepared in water. Small aliquots of this solution were added to a stock solution of protein in buffer at 0.5 mg/ml concentration. The total volume required at the time first signs of precipitation originated was noted down.

#### *Real Solubility*

For real solubility evaluations, the DVD-Ig protein was concentrated and stored overnight at 5°C. The solution was visually inspected for precipitates, phase separation,

turbidity, etc. The supernatant (or both phases) was checked for dissolved concentration.

#### *Near UV-CD*

The near UV-CD scans were taken at 1 mg/ml concentrations using 2 ml vial fill on a Jasco spectrometer (JASCO Analytical Instruments, Easton, MD) between 250 and 320 nm. The scan rate was 50 nm/minute and an average of 3 scans was taken. The spectrometer was allowed to equilibrate by turning on the lamp before data acquisition.

#### *ATR-FTIR Spectrometry*

FTIR scans were taken at 1 mg/ml concentrations using 10  $\mu$ l solutions on a Bruker ATR-FTIR (Bruker Optics, Billerica, MA). The scans were collected between 400-4000  $\text{cm}^{-1}$  and area normalized and second derivatized before being curve fitted using Origin software (OriginLab, Northampton, MA).

#### *Light Scattering*

Light scattering studies were done on a Malvern zetasizer (Malvern Instruments Ltd., Worcestershire, UK) using a backscattering angle of 173°. Toluene was used as a standard and a buffer (*e.g.*, acetate, histidine, and Tris) was used as a blank. An automatic mode was used.

#### *Dynamic Scanning Fluorimetry (DSF)*

DSF was employed to assess the propensity of a protein to unfold. The technique involved the addition of dye Sypro Orange to the protein samples. This fluorescent dye is sensitive to hydrophobic surfaces and shows increased fluorescence in such environments. The sample with dye was then heated and the fluorescence signal as a function of temperature was monitored. As the temperature increased, the protein started to unfold and exposed its hydrophobic interior. This lead to dye binding to this region and a greater fluorescence signal. The temperature at which the signal begins to increase is the onset temperature ( $T_{on}$ ). Proteins that have less intrinsic stability are more prone to unfold and have lower  $T_{on}$  values than proteins with greater intrinsic stability. DSF also provided a high throughput tool for rapid screening of clones in a 96 well format and eliminated potential limitations of larger quantities of samples and longer run times in DSC. 6  $\mu$ l of the 0.4X SYPRO Orange dye

(Invitrogen, Carlsbad, CA) was added to 27  $\mu$ l of the DVD-Ig protein solution. The scan rate was 1°C/minute and scans were taken from 25-75°C.

### *Lyophilization Methods*

The DVD-Ig proteins were lyophilized according to standard methods and the stability of the resulting lyophilisates were investigated.

### *Sample Preparation and Lyophilization*

The vials were stoppered with autoclaved and dried lyo stoppers. Afterwards the vials were lyophilized with a freeze dryer. A typical cycle is shown below in Table 4. The samples were reconstituted to a 100 mg/ml DVD-Ig protein solution.

Table 4: Typical Freeze-Dry Cycle Used To Lyophilize DVD-Ig Proteins

Step		Shelf	Time /	Step	Pressure
		temperature [°C]	Step time [min]		
0	Start	20	0:00:00	0	1000
1	Loading	20	0:00:00	0	1000
2	Freezing I (ramp)	0	0:20:00	20	1000
3	Freezing I	0	2:10:00	130	1000
4	Freezing II (ramp)	-45	1:20:00	80	1000
5	Freezing II	-45	3:00:00	180	1000
6	Adjust vacuum	-45	1:00:00	60	0,066
7	Primary drying I (ramp)	-25	1:00:00	60	0,066
8	Primary drying I	-25	70:00:00	4200	0,066
9	Secondary drying II (ramp)	25	2:00:00	120	0,066
10	Secondary drying II	25	0:15:00	15	0,036
11	Secondary drying II	25	8:00:00	480	0,036
12	Holding step (ramp)	5	0:30:00	30	0,036
13	Holding step	5	0:00:00	0	0,036
14	Venting N2 atmosphere	5	0:00:00	0	500
	Total			5375	

## I. STABILITY OF DVD-Ig PROTEINS

Examples 1-3 demonstrate that DVD-Ig proteins are less stable, *e.g.*, aggregating more easily and having a lower melting temperature, than antibodies due to the increased structural complexity of DVD-Ig proteins.

### EXAMPLE 1. Thermodynamic Comparison of DVD-Ig Proteins and Antibodies

An experiment was performed to determine the stability of a DVD-Ig protein in comparison to an antibody. Differential scanning calorimetry (DSC) of an IgG1 molecule (a monoclonal antibody, mAb) and a DVD-Ig was performed to determine the differences in the thermodynamic properties of the two molecules. Specifically, DSC was performed to compare the thermodynamic properties of an exemplary antibody (Briakinumab, an anti-IL12 monoclonal antibody) to those of a DVD-Ig protein (TNF/PGE2; DVD-B). Formulation information and DSC conditions are provided below in Example 2. Comparison of the DSC profiles of Briakinumab to those of TNF/PGE2 (DVD-B) shows the differences in three versus four domain unfolding (Figure 1). It is also clear from Figure 1 that the thermal unfolding of the DVD-Ig protein started earlier than that of the antibody, which indicates that the overall thermodynamic stability (or the intrinsic stability) of the DVD-Ig molecule is lower than that of the antibody. It should be noted that while the DVD-Ig protein in Figure 1 (DVD-B) had a DSC profile indicating an initial unfolding at about 50 °C, this same DVD-Ig protein was characterized as an LS-DVD-Ig protein given results from further stability testing.

The thermodynamic stability (*e.g.*, onset of unfolding as determined by dynamic scanning calorimetry) of mAbs is in general approximately about 5 °C higher compared to the panel of DVD-Ig proteins listed in Table 1. Although there is generally no direct correlation between the temperature of unfolding and the aggregation propensity of a group of DVD-Ig proteins, the values can be used as a screen to assess the stability of a given group of DVD-Ig proteins. Therefore, the onset temperature of unfolding can be utilized to identify AS-DVD-Ig proteins or LS-DVD-Ig proteins. As described in Figure 4, the overall distribution shows that DVD-Ig proteins generally have a lower unfolding temperature in comparison to antibodies. Figure 4 provides the compiled results from 16 different DVD-Ig proteins and 14 different antibodies. The assays performed in the studies described in Figure 4 are similar to those described above and in Figure 1.

## EXAMPLE 2. Impact of pH on the Stability of DVD-Ig proteins in Solution

The thermodynamic stability (intrinsic stability) of DVD-Ig proteins formulated in solutions having a pH of 4, 6, or 8 was evaluated using differential scanning calorimetry (DSC). All formulations had a DVD-Ig protein concentration of 1 mg/ml in 5 mM citrate/5mM phosphate buffer. Heating was performed at a scan rate of 1 °C/minute. Results showing the impact of pH on the stability of multiple DVD-Ig proteins are provided in Table 5 below. Tm1–Tm4 described in Table 5 represent the thermal melting/unfolding temperatures of various domains, *e.g.*, CH1, CH2 and CH3 domains, etc. The stabilities of two antibodies (Adalimumab and Briakinumab) are also described for comparison.

Table 5: Impact of pH on Stability of DVD-Ig Protein Formulations with pH 4, 6, or 8, as Assessed by Differential Scanning Calorimetry (DSC)

DVD-Ig	pH	Tm1 °C	Tm2 °C	Tm3 °C	Tm4 °C	Onset °C
5	4	62.3	63.9	73.9	80.7	45
6	4	61.5	69.5	75.4	82.2	45
37	4	61.4	66.7	77.1	82.7	52
38	4	68.4	70.4	75.8	81.6	56
53	4	58.6	67.9	76.5	82.7	46
54	4	66	67.9	75.2	82.6	52.5
65	4	61.3	69	73.9	81.3	52.5
66	4	65.5	67.6	74.7	81.9	55
165	4	63	67.4	75.4	82.1	47.5
166	4	67.3	71.7	74.9	82.4	55
257	4	66.2	69.2	76.6	83.5	52.5
258	4	68.2	69.6	78.8	83.7	55
277	4	61.4	64.8	75.3	82.6	50
278	4	55.8	67.5	76.1	82.8	45
281	4	65.8	68.4	79.1	83.1	55
282	4	73.1	75	77.4	83.8	57.5
5	6	61.9	65.1	76	81.8	50
6	6	61.6	69.8	76.2	81.9	48
37	6	60.4	67.5	77.2	83.1	51
38	6	69.07	70.3	75.9	82	57.5
53	6	58.3	67.05	76.4	82	49
54	6	65.8	67.6	74.9	82	56
65	6	61.2	67.3	73.4	82.4	51.5
66	6	65.3	67.5	74.8	81.9	55
165	6	62.6	67.3	75.7	82.1	50
166	6	67.3	71	74	82	54
257	6	66.6	69.2	75.8	82.9	57
258	6	67.8	70	77.5	79.5	55

DVD-Ig	pH	Tm1 °C	Tm2 °C	Tm3 °C	Tm4 °C	Onset °C
277	6	61.4	65.8	74.7	82.23	52.5
278	6	56.6	66.7	75.8	82.3	45
281	6	65.8	67.8	78.6	81.7	53
282	6	69.8	75	77.9	83.4	57
5	8	65.16	68.35	77	82.48	45
6	8	62	69.07	73.08	82.61	45
38	8	68.4	70.7	74.8	82.8	50
53	8	57.6	68.7	75	83	40
54	8	65.5	67.6	74.5	82.8	52.5
65	8	61	69.6	72.5	82.8	48
66	8	64.8	67.1	72.4	82.9	52.5
165	8	63.9	68.6	74.1	82.6	47.5
166	8	64.3	69.3	74	82.7	45
257	8	67.2	69.3	76.7	83.6	55
258	8	69.5	71.4	78.1	83.9	53
277	8	60.18	68.98	74.5	83	50
278	8	52.8	68.8	75.8	83.02	44
281	8	65.2	67.1	78.3	83.2	51
282	8	71.9	74.5	77.4	83.9	55
mAbs						
Adalimumab	6	72	76	84	NA	62
Briakinumab	6	69	76	83	NA	59

As shown in Table 5, DVD-Ig proteins in general have unfolding onset temperatures of greater than about 50 °C, and the melting temperatures are therefore slightly lower than those of antibodies and other stable proteins. Example 2 shows that the onset temperature for Briakinumab and Adalimumab is around 60 °C at pH 6, whereas, for DVD-Ig proteins, the average is around 53 °C.

The data also shows that the melting temperatures of DVD-Ig proteins are higher at pH 6 and pH 8 than at pH 4. Thus, pH affects the physico-chemical stability of DVD-Ig proteins, and stability appears best at approximately pH 6.

To further assess the impact of solution pH on the stability of DVD-Ig protein formulations during long-term storage, DVD-Ig protein formulations were analyzed using SEC before being subjected to storage (T0) or after being subjected to 3 months of accelerated storage (T3m). Storage stability of the DVD-Ig proteins in solutions (1 mg/ml DVD-Ig protein in 5 mM citrate/5mM phosphate buffer with the presence of 80 mg/ml sucrose) formulated at pH of 4, 6, or 8 was evaluated. For accelerated storage, samples were filled into sterile vials (approx. 500 µl each) and stored under controlled conditions (in temperature chambers and in the absence of light) at 40°C. The percentage of DVD-Ig

protein monomers (Mon) aggregates (Agg), and fragments (Frag) was determined using SEC and the results are presented in Table 6.

**Table 6: Impact of pH on the Stability of 1 mg/ml DVD-Ig Protein Solutions Before and After Accelerated Storage**

DVD-Ig	pH	Mon/T0	Mon/T3m	Agg/T0	Agg/T3m	Frag/T0	Frag/T3m
5	4	91.14	57.51	5	0.79	3.85	41.6
6	4	97.38	62.46	2.29	4.11	0.31	33.4
38	4	94.9	56.52	3.58	22.02	1.5	21.44
53	4	97.61	28.11	0.99	53.94	1.38	17.93
54	4	96.57	53.89	1.93	15.2	1.48	30.89
65	4	94.46	50.35	1.33	26.21	4.21	23.42
66	4	97.99	58.39	0.9	17.72	1.1	23.87
165	4	96.2	68.38	2.25	3.18	1.53	28.42
166	4	97.09	66.3	0.76	0.66	2.14	30.55
258	4	98.61	45.42	0.46	30.56	0.91	24
277	4	97.05	61.26	1.55	11.48	1.38	27.23
278	4	98.33	45.58	0.9	28.13	0.76	26.27
5	6	90.46	81.75	7.02	3.04	2.51	12.77
6	6	97.02	85.44	2.78	2.71	0.18	11.82
38	6	95.31	87.89	3.48	5.48	1.2	6.61
53	6	97.75	86.68	1.02	6.21	1.22	7.08
54	6	96.48	88.03	2.07	5.37	1.43	6.59
65	6	94.89	87.13	0.89	5.78	4.21	7.06
66	6	97.99	88.93	0.9	5.13	1.09	5.91
165	6	96.21	86.31	2.24	5.84	1.53	7.82
166	6	98.6	89.37	1.39	0.92	0	7.38
258	6	98.98	84.76	0.2	9.41	0.81	5.81
277	6	97.48	83.96	1.48	5.11	1.03	10.91
278	6	98.65	79.81	0.78	9.52	0.56	10.65
5	8	90.21	67.86	7.26	4.51	2.51	22.75
6	8	97.36	76.33	2.44	8.89	0.18	14.76
38	8	95.09	80.11	3.47	9.37	1.42	10.5
53	8	97.81	79.63	0.96	9.82	1.22	10.53
54	8	96.69	81.13	1.93	9.44	1.37	9.41
65	8	93.12	79.5	1.27	9.88	5.59	10.59
66	8	97.69	81.37	0.99	9.17	1.31	9.44
165	8	96.57	71.02	2.03	18.85	1.39	10.11
166	8	97.23	79.76	0.76	1.92	1.99	11.3
258	8	98.67	84.22	0.24	5.91	1.07	9.84
277	8	97.06	70.08	1.72	16.13	1.2	13.78
278	8	98.18	43.34	0.98	45.59	0.83	11.04

The results in Table 6 showed that DVD-Ig proteins can be initially formulated in a pH range from pH 4 to pH 8 and that the initial values of the freshly prepared samples at T0 do not show major differences. However, following accelerated storage, the DVD-Ig proteins

at pH 4 and pH 8 showed significant loss in monomer and a corresponding increase in aggregates or fragmentation.

The stability of the tested DVD-Ig proteins was highest at around pH 6. All of the DVD-Ig proteins tested (including DVD 5, DVD 6, DVD 38, DVD 53, DVD 54, DVD 65, DVD 66, DVD 165, DVD 166, DVD 258, DVD 277, and DVD 278) had a greater percentage of monomers and a lower percentage of fragments at pH 6 than at pH 4 or at pH 8.

Nine of the twelve DVD-Ig proteins tested showed a lower percentage of aggregates at pH 6 than at pH 4, and for the three DVD-Ig proteins that showed the reverse pattern, the difference in the percentage of aggregates at pH 6 and pH 4 was very small (difference of less than 2.7%). Also, eleven of the twelve DVD-Ig proteins tested showed a lower percentage of aggregates at pH 6 than at pH 8. Thus, accelerated storage resulted in increased aggregate formation. However, the increase was less than anticipated, particularly at pH 6.

#### *Impact of Solution pH on the Storage Stability of IL12IL18 DVD-Ig Protein Formulations*

To assess the impact of solution pH on the stability of DVD-Ig protein formulations during storage, DVD-Ig protein formulations with solution pH of 4, 6, or 8 were analyzed using SEC and IEC before being subjected to storage (T0) or after being subjected to 4 days (4d), 7 days (7d), or 21 days (21d) of storage at 5°C, 40°C, or 50°C (See Tables 6, 7 and 8). The solutions evaluated had an IL12-IL18 DVD-Ig concentration of 2 mg/ml and were in a buffer of 10 mM citrate and 10 mM phosphate. Samples were filled into sterile vials (approx. 500 µl each) and stored under controlled conditions (in temperature chambers and in the absence of light). The percentage of DVD-Ig protein monomers, aggregates, and fragments was determined using SEC (see Table 7) and the numbers of main, acidic and basic species were assessed using IEC (see Table 8).

Table 7: Storage Stability of IL12-IL18 DVD-Ig Protein Formulations with pH 4, 6, or 8 as Measured Using SEC

	Monomer			Aggregates			Fragments		
	pH 4	pH 6	pH 8	pH 4	pH 6	pH 8	pH 4	pH 6	pH 8
T0	97.64	97.61	97.84	1.35	1.36	1.23	1.01	1.03	0.94
Storage at 5°C									
4d, 5°C	97.98	97.64	97.95	1.07	1.31	1.19	0.95	1.06	0.87
7d, 5°C	97.68	98.02	97.8	1.25	1.17	1.08	1.07	0.81	1.11
21d, 5°C	98.13	97.81	97.64	1.23	1.17	1.08	0.88	0.96	1.28

Accelerated Storage at 40°C									
4d, 40°C	97.21	97.32	96	0.56	1.4	1.68	2.24	1.28	2.32
7d, 40°C	96.15	97.32	94.57	0.71	1.35	1.96	3.15	1.33	3.48
21d, 40°C	91.99	96.39	91.31	1.16	1.49	3.15	6.85	2.12	5.54
Accelerated Storage at 50°C									
4d, 50°C	90.2	97.07	91.73	5.54	1.46	2.78	4.26	1.47	5.49
7d, 50°C	83.19	96.35	81.32	9.66	1.42	10.68	7.15	2.24	8.2
21d, 50°C	52.98	91.36	46.57	33.1	4.17	33.35	13.91	4.47	17.18

The SEC data show that the IL12IL18 DVD-Ig protein formulations were most stable at pH 6. At pH 6, the stored IL12IL18 DVD-Ig protein formulations generally showed >95% monomers and <2% aggregates. Even under accelerated storage conditions of 50°C for 21 days, the formulation retained >90% monomers and <5% aggregates. IL12IL18 DVD-Ig formulations were more stable at pH 6 than pH 4 and pH 8, particularly in the longer duration and higher temperature storage conditions (*e.g.*, in the 21 day, 50°C condition). According to these results the IL12/IL18 DVD-Ig protein would be considered an AS-DVD-Ig protein given the stability results following storage at 50 °C.

Table 8: Storage Stability of IL12IL18 DVD-Ig Protein Formulations with pH 4, 6, or 8 as Measured Using IEC

	Main Species			Acidic			Basic		
	pH 4	pH 6	pH 8	pH 4	pH 6	pH 8	pH 4	pH 6	pH 8
T0	69.98	71.6	69.13	14.96	15.21	18.12	15.06	13.19	12.75
4d, 5°C	70.32	71	68.46	15.08	15.54	18.84	14.6	13.46	12.75
7d, 5°C	69.74	70.69	67.14	15.43	15.91	19.71	14.83	13.41	13.15
21d, 5°C	69.67	71.32	66.78	15.65	16.26	20.95	14.68	12.42	12.26
4d, 40°C	56.43	68.09	43.52	21.14	18.73	45.51	22.33	13.19	10.97
7d, 40°C	49.1	65.11	26.55	26.08	22.03	59.93	24.83	12.86	13.53
4d, 50°C	36.58	57.09	25.88	29.47	29.48	48.23	33.95	13.43	25.89

The IEC data for the IL12IL18 DVD-Ig protein formulations at pH 4, 6 and 8 show that the chemical degradation is dependent on the pH and the storage temperature. The highest chemical stability was observed for samples stored at 5°C. The minor changes in stability detected after 21 days indicated that the IL12IL18 DVD-Ig protein was stable at this storage temperature, which is considered common for commercial biotherapeutics. Although the differences in stability were small, the highest chemical stability was observed for formulations at pH 6. The lack of chemical stability was even more pronounced at the elevated storage temperatures of 40 °C and 50 °C.

### EXAMPLE 3. DVD-Ig Proteins Aggregate More Easily Than Monoclonal Antibodies

The impact of shaking on the aggregation of antibodies versus DVD-Ig proteins was examined. Shaking is a stress that can lead to the aggregation of molecules. The susceptibility of a DVD-Ig protein, TNF/PGE2 (DVD-B), to aggregation following shaking was compared with a monoclonal antibody, Briakinumab, using solutions having a protein concentration of 1 mg/ml (solutions at pH 6, 10 mM citrate/10 mM phosphate) in 6R vials. The 6R vials were filled with samples of 5 ml of the protein solution and shaken on an HS 260 IKA shaker (Wilmington, NC) at a speed of 150 revolutions per minute (rpm) for various lengths of time (0, 5, 24, 48, or 96 hours). The samples were checked for optical density at 500 nm, which provides a measurement of the turbidity of the solutions. Higher turbidity indicates greater aggregation and less stability. The results are shown in Table 9, revealing that the DVD-Ig protein aggregates more readily than the monoclonal antibody.

Table 9: Impact of Shaking on the OD<sub>500</sub> of 1 mg/ml Solutions of TNF / PGE2 DVD-Ig Protein and Briakinumab

Time (h)	Optical Density (OD) at 500nM	
	Briakinumab	TNF / PGE2 (DVD-B)
0	0	-0.0095
5	0.001	0.01175
24	0.01	0.0949
48	0.025	0.295
96	0.045	0.58

As indicated by OD<sub>500</sub> measurements which show turbidity of the solution, shaking caused the DVD-Ig protein to form more sub-visible and visible aggregates than the monoclonal antibody. Thus, after 48 hours of shaking, the DVD-Ig protein was more prone to colloidal instability and, therefore, less stable than the monoclonal antibody. The greater formation of visible aggregates of DVD-Ig protein compared with monoclonal antibody indicates that the DVD-Ig protein is generally less stable than the monoclonal antibody against shear stress in solution. Also, these results suggest that not all DVD-Ig proteins are as stable at pH 6 as a monoclonal antibody.

## II. ASSAY FOR IDENTIFYING AN AQUEOUS STABLE DVD-IG PROTEIN (AS-DVD-IG)

### EXAMPLE 4. DVD-Ig Proteins Can be Characterized as “Aqueous Stable” or “Aqueous Non-Stable”

As discussed above, the inventors found that formulations wherein the DVD-Ig protein was in the aqueous state suffered from certain problems such as aggregation and/or fragmentation of the DVD-Ig protein monomer. The inventors conducted experiments and discovered, surprisingly and unpredictably, that a subset of DVD-Ig proteins can be stably formulated in the aqueous state even at high concentrations. The following example describes an SEC study showing that, surprisingly, DVD-Ig proteins can be characterized as either aqueous stable, *e.g.*, the DVD-Ig protein shows low change in percent rel. peak area in monomers or aqueous non-stable, *e.g.*, the DVD-Ig protein is prone to aggregation and/or fragmentation. Notably, many of the DVD-Ig proteins tested were found to be aqueous non-stable or lyophilized non-stable. Due to the structural complexity of DVD-Ig proteins and the prominence of hydrophobic interactions at high concentrations, it was not expected that DVD-Ig proteins would be stable in formulations at high concentrations.

To assess the impact of storage temperature during accelerated or long-term storage of protein liquid formulations on protein stability, various DVD-Ig proteins were exposed to short-term storage at elevated temperatures in order to quickly gain insight in the formulation feasibility for long-term storage at lower temperatures (*e.g.*, 2-8°C).

#### *Stability Screen at a High Concentration for 14 Days*

DVD-Ig protein formulations with concentrations of 60 mg/ml were analyzed using SEC before being subjected to storage (T=0) or after being subjected to 14 days of accelerated storage (T=14days) (Table 10). Storage stability of the DVD-Ig proteins in solution (60 mg/ml, 10 mM citrate/10mM phosphate buffer with 80 mg/ml sucrose) was evaluated at 40 °C. After defined storage periods, samples were pulled and the impact of storage time on DVD-Ig protein stability was evaluated. Briefly, samples were filled into sterile vials (approx. 500 µL each) and stored under controlled conditions (in temperature chambers and in the absence of light) at 40 °C. At predefined points of time, samples of prepared solutions were pulled for analysis according to the sample pull scheme. The percentages of DVD-Ig protein monomers (Mon), aggregates (Agg), and fragments (Frag)

were determined using SEC, and the results are presented in Table 10.

Table 10: Impact of High Concentration on the Storage Stability of DVD-Ig Protein Solutions

DVD-Ig	Mon/T0	Mon/T14d	Agg/T0	Agg/T14d	Frag/T0	Frag/T14d
5	79.1	73.28	19.12	24.26	1.77	2.44
6	84.01	89.46	12.69	9.81	3.29	0.71
37	92.85	74.35	6.34	23.42	0.8	2.21
65	95.4	92.79	1.31	5.39	3.28	1.79
66	96.56	94.49	1.08	3.88	2.35	1.6
165	90.17	64.09	8.99	33.44	0.83	2.45
166	98.17	94.93	1.09	3.11	0.72	1.93
257	94.83	76.47	4.77	21.78	0.39	1.72
277	97.46	85.06	1.77	13.19	0.76	1.73
278	98.45	73.01	1.06	25.69	0.48	1.27
281	93.92	68.01	2.78	30.5	3.29	1.46
282	98.34	95.39	1.03	3.05	0.62	1.54

Surprisingly, as shown in Table 10, the inventors discovered that a subset of the DVD-Ig proteins tested was stable when formulated in the aqueous state. Ten of the twelve DVD-Ig proteins tested (DVD 5, DVD 6, DVD 37, DVD 65, DVD 66, DVD 166, DVD 257, DVD 277, DVD 278, and DVD 282) showed less than 26% aggregate formation and had greater than 73% monomers following 14 days of accelerated storage. Five of the DVD-Ig proteins tested (DVD 6, DVD 65, DVD 66, DVD 166, and DVD 282) showed aggregate formation of less than 10%, and three of these (DVD 66, DVD 166, and DVD 282) showed aggregate formation of less than 5%.

As described above, certain DVD-Ig proteins (“Aqueous Stable DVD-Ig” proteins or “AS-DVD-Ig” proteins) remain stable (*e.g.*, less than 6 % relative (rel.) peak area change in monomers or less than 10% relative (rel.) peak area change in monomers as determined by SEC) following accelerated storage in 14 days at 40 °C, even when formulated at high concentration (*e.g.*, concentrations of 60 mg/ml or higher). The majority of DVD-Ig proteins tended to aggregate during accelerated storage (non-AS-DVD-Ig proteins), as would be expected based on the general structure of DVD-Ig proteins and the stability studies described in Examples 1-3. Thus, in certain embodiments, the cutoff for separating the AS-DVD-Ig proteins from the non-AS-DVD-Ig proteins was taken as the formation of 10 % net aggregates or less in 14 days at 40 °C when stored at > 50 mg/ml at pH 6, as seen in the above non-limiting example where five of the twelve DVD-Ig proteins tested showed

aggregate formation at this level. In certain embodiments, the cutoff for separating the AS-DVD-Ig proteins and the non-AS-DVD-Ig proteins was taken as the formation of 6 % net aggregates or less in 14 days at 40 °C when stored at > 50 mg/ml at pH 6, as seen in the above non-limiting example where four of the twelve DVD-Ig proteins tested showed aggregate formation at this level.

Many of DVD-Ig proteins do not show low aggregation, *e.g.*, 1% or less aggregation at 5 °C after 21 days or 10% or less aggregation at 40 °C following 21 days of storage. For example, in an assay which examined monomer loss after 7 days in a solution having a TNF/IL13 DVD-Ig protein concentration of 50 mg/ml at either 4 °C or 40 °C, many of DVD-Ig proteins showed an increase in monomer loss as determined by SEC. In some cases the amount of monomer loss was negative as the monomer level increased in these cases (*e.g.*, some of the aggregates dissociated and formed back monomer and hence the apparent decrease in loss). A third experiment tested TNF/SOST DVD-Ig proteins in a solution having a DVD-Ig protein concentration of 50 mg/ml at 4 °C. As in the experiment relating to TNF/IL13 DVD-Ig proteins, many of the DVD-Ig proteins showed an increase in monomer loss (determined by SEC).

Notably, the above assays can also be used to distinguish Lyophilized-Stable DVD-Immunoglobulin (LS-DVD-Ig) proteins. The cutoff for separating the LS-DVD-Ig proteins and the non-LS-DVD-Ig proteins was taken as the formation of 15 % net or less aggregates in 14 days at 40 °C when stored at > 50 mg/ml at pH 6. Thus, DVD-Ig proteins tested in the above assay that result in less than 10% or less than 6% aggregation or less than 10 % or less than 6 % relative (rel.) peak area change in monomers are considered both AS-DVD-Ig protein and LS-DVD-Ig proteins, and DVD-Ig proteins resulting in less than 15% aggregation or less than 15 % relative (rel.) peak area change in monomers are considered LS-DVD-Ig proteins. Both AS-DVD-Ig protein and LS-DVD-Ig proteins represent only a percentage of the overall DVD-Ig proteins tested.

### III. STABILITY OF NON-AS-DVD-IG PROTEINS IN FORMULATIONS

The following examples provide data showing the stability of non-AS-DVD-Ig proteins (which fail the aggregation test, *i.e.*, show more than, for example, 6% aggregation, described in Example 4 above), in formulations, in comparison to AS-DVD-Ig proteins (described in Sections IV to VIII).

**EXAMPLE 5: Impact of Concentration on the Storage Stability of an Exemplary Non-AS-DVD-Ig Protein**

To assess the impact of protein concentration on long term storage stability, formulations of an exemplary non-AS-DVD-Ig protein with concentrations of 1, 2, 5, 10, 25, 50, and 75 mg/ml were subjected to storage for 14 days at 40°C. The formulations had a pH of 6 and were in 15mM histidine buffer alone. The samples were filled into sterile vials (approx. 500 µl each) and stored under controlled conditions (in temperature chambers and in the absence of light). The samples were analyzed using SEC to determine the percentage of aggregates following storage. The resulting data is provided in Table 11.

Table 11: The Total Percent Aggregates in the DVD-B protein as Measured Using SEC Following Storage at 40°C for 14 Days

Concentration of DVD-B	Aggregates %
1 mg/ml	0
2 mg/ml	0.2
5 mg/ml	3.64
10 mg/ml	6.76
25 mg/ml	18.02
50 mg/ml	32.82
75 mg/ml	48.28

The data in Table 11 indicate that under the tested conditions (*i.e.*, pH 6, 15mM histidine buffer) DVD-B becomes unstable, namely, a high proportion of aggregates form after 14 days of storage at 40°C when high concentrations are reached. The percentage of aggregates formed exceeded 18% at a concentration of 25 mg/ml or more. Thus, DVD-B, a non-AS-DVD-Ig protein, was not stable in histidine buffer as evidenced by increased aggregation during storage.

**EXAMPLE 6: Impact of pH, Ionic Strength, and Concentration on the Storage Stability of an Exemplary Non-AS-DVD-Ig Protein**

To assess the impact of pH, ionic strength, and concentration on the storage stability of a DVD-Ig protein in solution, various formulations of DVD-B (5 mg/ml and 100 mg/ml) were evaluated at 40°C and 5°C. After defined storage periods, samples were pulled and the

impact of storage time on DVD-Ig protein stability was evaluated. The following buffers were used: acetate for pH 4.5, histidine for pH 6 and Tris for pH 8. A 2 mM concentration of buffer was used for 1 mM ionic strength solutions and a 10 mM concentration of buffer for 20 and 100 mM ionic strength solutions (sodium chloride was used to further maintain ionic strength). Samples were filled into sterile vials, approx. 500 µl each, and stored under controlled conditions in a temperature chamber and in the absence of light. After 3 months at 5°C (5C, 3m) or 21 days at 40°C (40C, 21d), samples of prepared solutions were analyzed using SEC. The numbers of net aggregates measured using SEC are presented in Table 12. Tables 13 and 14 further show that the addition of different stabilizers/excipients (e.g., sucrose and Tween80) did not result in significant improvement in percent monomer remaining after defined time points (results were obtained using the methodology described above). The negative values of net aggregate values at the initial time points for the low concentration samples indicate also the initial formation of reversible aggregates.

Table 12: Impact of Storage of DVD-B Protein Under Various Formulation Conditions on the Amount of Aggregates Formed as Measured By SEC

DVD-Ig Protein Concentration and Storage Condition	pH	Added Ionic strength	Aggregate (Net)
DVD-B, 100 mg/ml, 5°C, 3m	4.5	1	49.51
		20	54.79
		100	56.64
	6	0	11.68
		1	6.6
		20	9.4
		100	7.85
	8	1	5.66
		20	8.08
		100	7.87
DVD-B, 5 mg/ml, 40°C, 21d	6	0	-6.56
		1	-3.3
		20	10.79
		100	9.99
	6	0	73.37
DVD-B, 100 mg/ml, 40°C, 21d	6	1	70.39
		20	77.39
		100	65.4

Table 13: Polyol and Polysorbate Has Little to No Effect on Stability of DVD-B at 1 mg/ml in Histidine Buffer, pH 6

	Sample No.	Monomer	Aggregate	Fragment	AUC
Histidine Buffer					
T0		98.97	0.34	0.67	79787
T4, 40°C	1	98.13	0.82	1.04	78552
	2	97.98	0.87	1.13	78622
	Average	98.055	0.845	1.085	78587
T7, 40°C	1	97.45	1.15	1.39	77836
	2	97.42	1.2	1.36	78137
	Average	97.435	1.175	1.375	77986.5
T21, 40°C	1	92.21	4.34	3.44	170875
	2	93.19	3.7	3.1	72149
	Average	92.7	4.02	3.27	121512
T4, 50°C	1	94.06	4.15	1.77	39698
	2	93.37	4.44	2.18	43002
	Average	93.715	4.295	1.975	41350
T7, 50°C	1	93.09	3.9	2.99	26451
	2	91.95	4.81	3.22	30158
	Average	92.52	4.355	3.105	28304.5
30 mM Hist, 80 mg/ml Sucrose, 0.02 % Tween 80					
T0		97.5	1.61	0.88	62902
T21, 40°C	1	95.19	1.47	2.95	79362
	2	95.56	1.67	3.13	79445
	Average	95.375	1.57	3.04	79403.5
T4, 50°C	1	94.94	3.38	1.66	80742
	2	94.98	3.3	1.62	79436
	Average	94.96	3.34	1.64	80089
T7, 50°C	1	91.06	6.71	2.21	79672
	2	91.01	6.6	2.37	79820
	Average	91.035	6.655	2.29	79746

The addition of polyol to the formulation resulted in a slight improvement in monomer content and a decrease in the levels of aggregates of DVD-B at 1mg/ml.

Table 14: Polysorbate Has Little to No Effect on Stability of DVD-B at 100 mg/ml in Histidine Buffer, pH 6

Buffer: 15 mM Histidine				
	Monomer	Aggregate	Fragment	AUC
T0	96.26	2.43	1.3	73681
T7 (Vial1)	41.4	56.1	2.34	64692
T7 (Vial2)	42.5	55.2	2.14	63246
T7 (Avg.)	41.95	55.65	2.24	63969
T21(Vial1)	37.2	60.03	2.76	52389
T21(Vial2)	38.8	58.05	3.13	50722

T21(Avg.)	38	59.04	2.945	51555.5
Buffer: 15 mM Histidine + 0.02 % Tween 80				
	<b>Monomer</b>	<b>Aggregate</b>	<b>Fragment</b>	<b>AUC</b>
T0	96.22	2.43	1.33	72007
T7 (Vial1)	42.9	54.8	2.2	65403
T7 (Vial2)	47	50.8	2.09	58048
T7 (Avg.)	44.95	52.8	2.145	61725.5
T21(Vial1)	40.32	54.54	5.12	32321
T21(Vial2)	38.38	55.9	5.7	30927
T21(Avg.)	39.35	55.22	5.41	31624

The addition of polyol to the formulation resulted in a slight improvement in monomer content and a decrease in the levels of aggregates of DVD-B at 100mg/ml.

Table 15: Polyol or Surfactant Does Not Improve Stability (1 mg/ml) of DVD-B

pH 6 Formulation	Monomer (%)			Aggregate (%)			Fragment (%)		
	T0	1M	3M	T0	1M	3M	T0	1M	3M
15 mM Na Phos.	96.45	90.03	80.83	1.41	5	9.09	2.12	4.95	10.06
15 mM Na Cit.	96.51	90.72	85.3	1.44	5.3	6.57	2.04	3.97	8.11
15 mM Na Succ.	96.06	87.41	78.17	1.53	5.46	8.24	2.39	7.11	13.57
15 mM Na Acet.	96.14	89.62	81.8	1.48	4.76	7.52	2.36	5.6	10.66
15 mM Arg.	96.12	92.48	85.72	1.65	3.39	5.59	2.21	4.11	8.68
15 mM Hist.	96.42	91.82	81.6	1.29	3.85	6.39	2.28	4.32	12
Self Buff.	96.03	88.79	81.44	1.57	3.91	4.18	2.38	7.29	14.37
UB 10 mg/ml Mannitol	95.86	90.18	84.08	2.01	5.74	8.09	2.11	4.07	7.81
UB 10 mg/ml Sorbitol	96.49	89.36	84.09	1.38	6.56	7.68	2.11	4.07	8.21
UB 10 mg/ml Sucrose	96.12	90.03	84.26	1.58	5.92	7.62	2.28	4.03	8.11
UB 10 mg/ml Trehalose	96.34	89.97	84.31	1.5	5.98	7.59	2.14	4.03	8.09
UB 2.5% Gly	96.43	87.22		1.42	8.59		2.13	4.17	
UB 15 mM $(\text{NH}_4)_2\text{SO}_4$	96.69	90.92	85.76	1.22	4.97	5.68	2	4.1	8.55
UB 20 mM NaCl	96.44	90.35	84.36	1.41	5.5	6.74	2.13	4.13	8.88
UB 200 mM NaCl	96.37	91.85	85.2	1.52	3.24	3.62	2.09	4.89	11.16

UB = citrate / phosphate buffer

The data show that although various solution conditions were analyzed, DVD-B was not very stable even at pH 6 (see, for example Table 13). Also, at pH 6, the ionic strength did not show a consistent relationship with net aggregate formation. The poor stability is

indicated by the formation of high amounts of aggregates even in the 5°C storage condition. Furthermore, the addition of a polyol and/or a surfactant did not improve aggregation of DVD-B (see Tables 13, 14, and 15).

As described above in Examples 1 to 3, many DVD-Ig proteins are intrinsically unstable. However, surprisingly, certain DVD-Ig proteins can be characterized as being stable, as described in Example 4. The experiments described in the Examples below demonstrate that AS-DVD-Ig proteins can unexpectedly be stably formulated, even at high concentrations, despite the differences in amino acid sequence. The below examples stand in contrast to Examples 5 and 6, which show the failure of non-AS-DVD-Ig proteins to be formulated.

#### **IV. AS-DVD-IG PROTEINS ARE STABLE IN FORMULATIONS CONTAINING A BUFFER AT PH RANGE OF 4.5-7.5**

##### **EXAMPLE 7: Effect of Buffer Concentrations on the Stability of DVD-Ig Proteins**

The concentration of a buffer, *e.g.*, histidine, is one of the important factors that may influence protein stability during accelerated/long-term storage of protein liquid formulations. To assess the impact, the protein was exposed to short-term storage at elevated and real time temperatures in order to quickly gain insight into stable formulations for long-term storage at lower temperatures (*e.g.*, 2-8°C).

Storage stability of DVD-Ig proteins in solution was evaluated at 40°C and 5°C. After defined storage periods, samples were pulled and the impact of storage time on DVD-Ig protein stability was evaluated. The concentrations of histidine that were evaluated include 0, 5, 15, 50, and 200 mM.

Samples were filled into sterile vials (approx. 500 µl each) and stored under controlled conditions (in temperature chambers and in the absence of light). After 7 days and 21 days, samples of the prepared solutions were analyzed using SEC and IEC.

**Table 16: Impact of Storage of Various DVD-Ig Proteins at Low and High Concentrations under Various Histidine Concentrations on SEC**

T0									
DVD-A, pH 5.2, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-A, pH 5.2, 75 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	1.89	97.21	0.88	N	0 Histidine	2.63	96.39	0.97	N
5 mM Histidine	0.93	98.19	0.86	N	5 mM Histidine	0.52	98.27	1.19	N
15 mM Histidine	1.41	97.64	0.94	N	15 mM Histidine	1.69	97.38	0.92	N
50 mM Histidine	1.83	96.84	1.31	N	50 mM Histidine	2.01	96.86	1.11	N
200 mM Histidine	2.09	96.96	0.94	N	200 mM Histidine	2.27	96.78	0.94	N
40C, 7d									
DVD-A, pH 5.2, 1 mg/ml	Agg	Mon	Frag		DVD-A, pH 5.2, 75 mg/ml	Agg	Mon	Frag	
0 Histidine	1.55	96.4	2.03	N	0 Histidine	10.52	87.77	1.7	N
5 mM Histidine	1.19	97.17	1.62	N	5 mM Histidine	1.26	95.01	3.71	N
15 mM Histidine	2.47	95.49	2.03	N	15 mM Histidine	13.76	83.35	2.88	N
50 mM Histidine	2.31	95.42	2.25	N	50 mM Histidine	16.62	80.82	2.54	N
200 mM Histidine	2.14	95.92	1.92	N	200 mM Histidine	7.47	89.68	2.83	Y
40C, 21d									
DVD-A, pH 5.2, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-A, pH 5.2, 75 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine					0 Histidine	15.77	82.2	2.02	Y
5 mM Histidine	2.52	95.3	2.16	N	5 mM Histidine	14.07	83.8	2.12	N
15 mM Histidine	3.39	93.85	2.74	N	15 mM Histidine	18.07	79.49	2.42	N
50 mM Histidine	2.74	94.4	2.84	N	50 mM Histidine	13.41	83.71	2.86	Y
200 mM Histidine	1.9	94.91	3.18	N	200 mM Histidine	8.35	87.66	3.97	Y
5C, 21d									
DVD-A, pH 5.2, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-A, pH 5.2, 75 mg/ml	Agg	Mon	Frag	
0 Histidine	1.24	97.62	1.12	N	0 Histidine	1.78	97.16	1.04	N
5 mM Histidine	1	97.86	1.12	N	5 mM Histidine	1.74	97.33	0.91	N
15 mM Histidine	1.08	97.43	1.47	N	15 mM Histidine	2.01	96.77	1.21	N
50 mM Histidine	1.25	97.36	1.37	N	50 mM Histidine	2.31	96.45	1.22	N
200 mM Histidine	1.61	97.08	1.3	N	200 mM Histidine				N
T0									
DVD-C, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-C, pH 5.4, 100 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	2.31	96.35	1.33	N	0 Histidine	3.05	95.18	1.75	N
5 mM Histidine	1.82	96.64	1.52	N	5 mM Histidine	2.55	95.84	1.6	N
15 mM Histidine	1.83	96.53	1.63	N	15 mM Histidine	2.2	96.1	1.68	N
50 mM Histidine	1.87	96.67	1.44	N	50 mM Histidine	1.8	96.54	1.64	N
200 mM Histidine	2.17	96.14	1.68	N	200 mM Histidine	1.7	96.49	1.8	N
40C, 7d									
DVD-C, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-C, pH 5.4, 100 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	2.41	92.11	5.47	N	0 Histidine	3.18	94.49	2.31	N
5 mM Histidine	1.62	95.93	2.43	N	5 mM Histidine	2.65	95.2	2.13	N
15 mM Histidine	1.49	95.9	2.59	N	15 mM Histidine	2.46	94.96	2.57	N
50 mM Histidine	1.38	96.16	2.45	N	50 mM Histidine	2.33	95.16	2.49	N
200 mM	1.47	96.12	2.39	N	200 mM Histidine	1.82	95.19	2.97	N

Histidine									
40C, 21d									
DVD-C, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-C, pH 5.4, 100 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	1.38	95.5	3.11	N	0 Histidine	3.45	93.64	2.89	N
5 mM Histidine	1.5	95.39	3.1	N	5 mM Histidine	2.8	94.27	2.91	N
15 mM Histidine	1.26	96.01	2.71	N	15 mM Histidine	2.55	94.56	2.87	N
50 mM Histidine	1.34	95.83	2.82	N	50 mM Histidine	2.26	94.9	2.83	N
200 mM Histidine	1.38	95.72	2.88	N	200 mM Histidine	2.84	93.29	3.86	N
5C, 21d									
DVD-C, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	DVD-C, pH 5.4, 100 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	2.08	96.12	1.78	N	0 Histidine	2.77	94.77	2.44	N
5 mM Histidine	1.56	96.72	1.71	N	5 mM Histidine	2.35	95.43	2.2	N
15 mM Histidine	1.27	97.08	1.63	N	15 mM Histidine	2.18	95.42	2.38	N
50 mM Histidine	1.4	96.67	1.91	N	50 mM Histidine	1.95	96.07	1.96	N
200 mM Histidine	1.59	96.62	1.77	N	200 mM Histidine	1.95	96.1	1.94	N
T0									
IL12IL18, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	IL12IL18, pH 5.4, 150 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	2.95	95.23	1.8	N	0 Histidine	3.48	94.22	2.29	N
5 mM Histidine	1.92	96.28	1.79	N	5 mM Histidine	4.64	93.26	2.09	N
15 mM Histidine	2.16	95.71	2.12	N	15 mM Histidine	4.33	93.53	2.12	N
50 mM Histidine	2.07	96	1.91	N	50 mM Histidine	3.93	94.2	1.88	N
200 mM Histidine	2.46	95.65	1.87	N	200 mM Histidine	3.55	94.48	1.96	N
40C, 7d									
IL12IL18, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	IL12IL18, pH 5.4, 150 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	2.88	93.82	3.29	N	0 Histidine	4.71	91.46	3.82	N
5 mM Histidine	1.32	95.82	2.85	N	5 mM Histidine	5.79	90.44	3.75	N
15 mM Histidine	1.02	96.12	2.85	N	15 mM Histidine	4.87	91.81	3.3	N
50 mM Histidine	1.57	95.23	3.19	N	50 mM Histidine	8.58	87.7	3.71	N
200 mM Histidine	1.43	95.56	3	N	200 mM Histidine	6.78	89.81	3.4	N
40C, 21d									
IL12IL18, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	IL12IL18, pH 5.4, 150 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	1.51	94.69	3.79	N	0 Histidine	5.18	91.21	3.6	N
5 mM Histidine	1.08	95.87	3.04	N	5 mM Histidine	6.76	89.67	3.56	N
15 mM Histidine	1.07	95.85	3.06	N	15 mM Histidine	5.7	91.19	3.09	N
50 mM Histidine	1.2	95.56	3.32	N	50 mM Histidine	12.87	83.35	3.76	N
200 mM Histidine	1.42	95.42	3.15	N	200 mM Histidine	7.21	89.88	2.89	N
5C, 21d									
IL12IL18, pH 5.4, 1 mg/ml	Agg	Mon	Frag	Precipitation*	IL12IL18, pH 5.4, 150 mg/ml	Agg	Mon	Frag	Precipitation*
0 Histidine	1.71	96.03	2.24	N	0 Histidine	5.96	91.35	2.67	N
5 mM Histidine	1.25	96.44	2.3	N	5 mM Histidine	6.94	90.97	2.07	N
15 mM Histidine	1.4	96.13	2.46	N	15 mM Histidine	5.69	92.19	2.1	N
50 mM Histidine	1.78	96.03	2.18	N	50 mM Histidine	10.07	88.13	1.78	N
200 mM Histidine	2.1	95.7	2.19	N	200 mM Histidine	1.32	95.56	3.11	N

\* Precipitation indicates if insoluble visible aggregates were observed (Yes or No)

**Table 17: Impact of Storage of Various DVD-Ig Proteins at Low and High Concentrations Under Various Histidine Concentrations on IEC**

T0							
DVD-A, pH 5.2, 1 mg/ml	Acidic	Main	Basic	DVD-A, pH 5.2, 75 mg/ml	Acidic	Main	Basic
0 Histidine	15.66	49.3	35.02	0 Histidine	19.71	50.72	29.55
5 mM Histidine	13.15	50.9	35.93	5 mM Histidine	19.11	50.8	30.08
15 mM Histidine	15.11	46.74	38.13	15 mM Histidine	18.58	44.52	36.88
50mM Histidine	13.27	52.38	34.33	50mM Histidine	15.69	52.06	32.24
200 mM Histidine	14.55	52.01	33.42	200 mM Histidine	16.79	45.36	37.84
40C, 7d							
DVD-A, pH 5.2, 1 mg/ml	Acidic	Main	Basic	DVD-A, pH 5.2, 75 mg/ml	Acidic	Main	Basic
0 Histidine	24.97	40.98	34.03	0 Histidine	21.7	43.6	34.68
5 mM Histidine	22.21	44.35	33.43	5 mM Histidine	28.31	43.99	27.69
15 mM Histidine	19.79	44.91	35.29	15 mM Histidine	21.21	41.34	37.44
50mM Histidine	18.19	46.11	35.68	50mM Histidine	18.42	42.84	38.73
200 mM Histidine	18.08	48.08	33.83	200 mM Histidine	16.71	42.09	41.19
5C, 21d							
DVD-A, pH 5.2, 1 mg/ml	Acidic	Main	Basic	DVD-A, pH 5.2, 75 mg/ml	Acidic	Main	Basic
0 Histidine	15.66	46.47	37.86	0 Histidine	16.41	48.16	35.41
5 mM Histidine	18.76	49.37	31.85	5 mM Histidine	16.19	49.44	34.36
15 mM Histidine	14.34	52.8	32.85	15 mM Histidine	16.91	45.77	37.3
50mM Histidine	15.06	50.33	34.59	50mM Histidine	17.17	47.45	35.37
200 mM Histidine	14.49	48.8	36.7	200 mM Histidine	15.77	46.18	38.03
T0							
DVD-C, pH 5.4, 1 mg/ml				DVD-C, pH 5.4, 100 mg/ml			
0 Histidine	26.28	59.4	14.3	0 Histidine	24.53	66.44	9.01
5 mM Histidine	23.01	69.42	7.56	5 mM Histidine	24.17	65.96	9.86
15 mM Histidine	22.05	67.79	10.14	15 mM Histidine	23.67	67.77	8.55
50mM Histidine	21.9	69.09	9	50mM Histidine	22.71	67.65	9.63
200 mM Histidine	20.99	70.74	8.26	200 mM Histidine			
40C, 7d							
DVD-C, pH 5.4, 1 mg/ml				DVD-C, pH 5.4, 100 mg/ml			
0 Histidine	34.05	55.74	10.2	0 Histidine	28.68	61.69	9.61
5 mM Histidine	31.26	59.07	9.65	5 mM Histidine	29.05	61.57	9.37
15 mM Histidine	27.8	61.86	10.33	15 mM Histidine	29.2	61.25	9.54
50mM Histidine	25.97	59.33	14.49	50mM Histidine	27.43	61.86	10.69
200 mM Histidine	26.35	62.28	11.36	200 mM Histidine	28.04	59.58	12.37
5C, 21d							
DVD-C, pH 5.4, 1 mg/ml				DVD-C, pH 5.4, 100 mg/ml			
0 Histidine	25.05	65.7	9.23	0 Histidine	23.43	66.85	9.71
5 mM Histidine	23.66	66.17	10.15	5 mM Histidine	22.3	68.58	9.11
15 mM Histidine	22.11	69.96	8.91	15 mM Histidine	23.39	67.31	9.29
50mM Histidine	21.79	68.96	9.23	50mM Histidine	23.67	67.96	8.35
200 mM Histidine	21.96	70.23	7.8	200 mM Histidine	22.6	69.68	7.65

T0							
IL12IL18, pH 5.4, 1 mg/ml				IL12IL18, pH 5.4, 150 mg/ml			
0 Histidine	30.24	51.83	17.92	0 Histidine	31.05	50.74	18.2
5 mM Histidine	29.11	54.71	16.16	5 mM Histidine	30.3	53.53	16.16
15 mM Histidine	28.36	57.23	14.39	15 mM Histidine	30.02	53.9	16.07
50mM Histidine	29.15	53.29	17.55	50mM Histidine	28.42	51.31	20.26
200 mM Histidine	35.59	52.31	12.09	200 mM Histidine	28.85	55.24	15.9
40C, 7d							
IL12IL18, pH 5.4, 1 mg/ml				IL12IL18, pH 5.4, 150 mg/ml			
0 Histidine	37.24	45.3	17.45	0 Histidine	34.95	44.85	20.19
5 mM Histidine	36	47.57	16.42	5 mM Histidine	31.94	45.7	22.34
15 mM Histidine	36.17	50.01	13.81	15 mM Histidine	32.39	45.54	22.05
50mM Histidine	35.39	49.12	15.47	50mM Histidine	37	41.76	21.23
200 mM Histidine	38.48	45.03	16.47	200 mM Histidine	30.86	47.34	21.78
5C, 21d							
IL12IL18, pH 5.4, 1 mg/ml				IL12IL18, pH 5.4, 150 mg/ml			
0 Histidine	30.72	51.83	17.44	0 Histidine	30.91	50.49	18.59
5 mM Histidine	30.25	51	18.74	5 mM Histidine	29.14	50.47	20.37
15 mM Histidine	30.07	56.33	13.58	15 mM Histidine	28.45	50.95	20.58
50mM Histidine	30.51	55.5	13.97	50mM Histidine	27.91	52.66	19.42
200 mM Histidine	42.9	49	8.08	200 mM Histidine	27.96	48.84	22.19

Tables 16 and 17 show that the amount of monomer remaining at different time points and at the formation of insoluble aggregates indicates that histidine concentrations in the range of about 5 to about 50 mM provided optimum stability. A concentration of about 200 mM histidine resulted in the formation of insoluble aggregates in some cases (see indication of precipitation in Table 16). 0 mM histidine formulations showed enhanced aggregation as indicated by formation of insoluble and soluble aggregates in some cases (in some cases soluble aggregates were higher at 0 than that at 5 mM histidine concentrations). Secondly, the pH is expected to be well maintained for longer storage times in formulations containing histidine. ,

The error in IEC measurements is usually higher (2-3 % variation) compared to SEC measurements with the same formulation. Hence taking that into account, no significant differences were observed within formulations as assayed by IEC indicating that the chemical stability was not as affected by the molarity of histidine, hence the chemical stability is independent of the buffer concentration (generally, in contrast to aggregation propensity).

**Example 8: Example of the Stability of an AS-DVD Ig Protein (DVD-CAnti-IL1 alpha/beta DVD-Ig Protein) in Solution**

An anti-IL1 alpha/beta DVD-Ig protein (DVD-C) was assessed for stability over time at both 100 mg/ml and 1 mg/ml in different buffers, at different pHs and at different temperatures. Buffers that were tested at 100 mg/ml of DVD-C included 15 mM acetate pH 4; 15 mM acetate pH 5; 15 mM histidine pH 5.5; 15 mM succinate pH 5.5; 15 mM histidine pH 6.0; Water (no buffer) pH 6.0; 15 mM citrate pH 6.0; 15 mM histidine pH 6.5; and 15 mM Tris pH 8.0. Buffers that were tested at 1 mg/ml of DVD-C included 10 mM citrate + 10 mM phosphate buffer at pH 3, 4, 5, 6, 7, or 8.

The samples were stored at 50°C, 40°C, 25°C, and 5°C. At certain time points, samples were pulled and evaluated for stability. Physical stability was evaluated by size exclusion chromatography (SE-HPLC or SEC), including % aggregate, % monomer, % fragment, and total species recovered were quantitated. Chemical stability was evaluated by weak cation exchange chromatography (IEX-HPLC or IEC), including % acidic, % main, and % basic species quantitated.

Tables 18 and 19 describe stability of DVD-C at 100 mg/ml and 1 mg/ml, respectively, in various buffers and pH. In both Tables 17 and 18, size exclusion chromatography (SEC) data and ion-exchange chromatography (IEC) data is displayed. Formulation and abbreviation keys are given below each table.

**Table 18. Stability at Various Temperatures of DVD-C at 100 mg/ml in Different Buffers and at Different pHs.**

Time	Temp(°C)	Formulation	pH	SEC data				IEX data		
				% HMW	% M	% LMW	AUC	% Acidic	% Main	% Basic
T0	---	ace	4	0.88	98.03	1.09	45941	19.65	71.68	8.67
T0	---	ace	5	0.94	98.19	0.87	45789	19.79	71.60	8.61
T0	---	his	5.5	0.98	98.15	0.86	48085	20.40	71.90	7.70
T0	---	succ	5.5	1.04	97.96	1.00	49317	19.70	71.54	8.76
T0	---	his	6	1.15	97.93	0.92	48468	20.68	70.94	8.38
T0	---	water	6	1.87	97.27	0.86	48356	18.95	72.35	8.70
T0	---	citrate	6	1.38	97.54	1.09	44457	19.35	72.37	8.28

Time	Temp(°C)	Formulation	pH	SEC data				IEX data		
				% HMW	% M	% LMW	AUC	% Acidic	% Main	% Basic
T0	---	his	6.5	1.16	97.93	0.91	47102	21.17	70.45	8.38
T0	---	tris	8	2.40	96.65	0.95	53889	21.71	69.12	9.17
T7d	40	ace	4	0.92	97.03	2.05	49124	19.16	71.61	9.23
T7d	40	ace	5	1.12	97.31	1.57	49077	21.62	71.09	7.29
T7d	40	his	5.5	1.17	97.27	1.56	49662	19.48	73.60	6.92
T7d	40	succ	5.5	1.46	97.06	1.48	48821	23.18	69.51	7.31
T7d	40	his	6	1.72	96.96	1.32	42872	20.99	73.36	5.64
T7d	40	water	6	2.48	96.22	1.30	37817	21.67	72.02	6.31
T7d	40	citrate	6	1.79	96.83	1.37	43022	21.70	71.52	6.78
T7d	40	his	6.5	2.08	96.65	1.27	48176	23.68	70.89	5.43
T7d	40	tris	8	4.58	93.87	1.55	48306	40.86	52.70	6.44
T1mo	40	ace	4	1.32	94.29	4.38	52518	34.78	32.07	33.16
T1mo	40	ace	5	1.64	95.36	3.00	53762	36.48	52.11	11.41
T1mo	40	his	5.5	1.61	95.38	3.01	52489	33.12	55.52	11.36
T1mo	40	succ	5.5	2.25	94.73	3.01	51508	40.94	46.88	12.19
T1mo	40	his	6	2.81	94.53	2.66	52229	34.46	56.00	9.54
T1mo	40	water	6	3.66	93.75	2.58	52285	33.46	56.53	10.01
T1mo	40	citrate	6	2.67	94.91	2.42	51968	36.54	52.80	10.67
T1mo	40	his	6.5	3.61	93.69	2.71	50276	39.06	51.20	9.74
T1mo	40	tris	8	7.49	85.24	7.28	53081	52.90	15.51	31.58
T1mo	25	ace	4	0.98	97.30	1.72	56026	23.39	59.02	17.59
T1mo	25	ace	5	1.15	97.31	1.54	55264	22.44	69.01	8.55
T1mo	25	his	5.5	1.16	97.41	1.43	54356	22.05	69.43	8.51
T1mo	25	succ	5.5	1.42	97.07	1.51	52417	23.00	67.63	9.37
T1mo	25	his	6	1.75	96.74	1.51	52220	23.80	67.67	8.52
T1mo	25	water	6	2.66	95.94	1.40	51198	23.68	67.35	8.98
T1mo	25	citrate	6	1.91	96.59	1.49	51137	22.41	68.14	9.45
T1mo	25	his	6.5	1.93	96.58	1.49	50594	25.93	65.50	8.56
T1mo	25	tris	8	4.35	93.87	1.78	51644	39.99	50.14	9.87

Time	Temp(°C)	Formulation	pH	SEC data				IEX data		
				% HMW	% M	% LMW	AUC	% Acidic	% Main	% Basic
T3mo	40	ace	4	2.19	85.66	12.15	54546	45.24	29.61	25.15
T3mo	40	ace	5	2.69	89.70	7.61	63493	52.09	31.52	16.39
T3mo	40	his	5.5	2.57	89.79	7.64	64361	48.68	35.18	16.14
T3mo	40	succ	5.5	3.53	88.89	7.58	57855	60.00	24.73	15.28
T3mo	40	his	6	4.08	90.70	5.21	60062	50.85	37.10	12.04
T3mo	40	water	6	4.80	90.50	4.70	58908	47.55	43.39	9.06
T3mo	40	citrate	6	3.87	91.38	4.75	55505	56.39	30.85	12.76
T3mo	40	his	6.5	5.56	89.08	5.37	56243	58.77	32.28	8.95
T3mo	40	tris	8	14.20	75.16	10.64	59362	63.76	22.12	14.12
T3mo	25	ace	4	1.15	96.39	2.46	60821	25.24	61.82	12.94
T3mo	25	ace	5	1.29	96.85	1.86	53513	23.62	66.47	9.91
T3mo	25	his	5.5	1.34	96.84	1.82	56041	23.05	67.70	9.25
T3mo	25	succ	5.5	1.69	96.36	1.95	53657	27.27	61.25	11.48
T3mo	25	his	6	2.04	96.23	1.73	51684	26.17	63.16	10.67
T3mo	25	water	6	2.81	95.49	1.70	52442	25.01	64.07	10.91
T3mo	25	citrate	6	2.08	96.11	1.80	51993	24.48	64.34	11.18
T3mo	25	his	6.5	2.26	96.01	1.72	50988	30.52	59.06	10.42
T3mo	25	tris	8	5.61	91.87	2.51	52070	59.46	31.54	8.99
T3mo	5	ace	4	0.85	98.01	1.14	50571	18.16	71.76	10.09
T3mo	5	ace	5	1.06	97.91	1.04	49173	18.44	71.54	10.02
T3mo	5	his	5.5	1.33	97.65	1.02	51947	19.97	70.28	9.75
T3mo	5	succ	5.5	1.32	97.59	1.09	50358	19.12	70.88	10.00
T3mo	5	his	6	2.18	96.79	1.03	49875	22.34	67.47	10.19
T3mo	5	water	6	8.21	90.70	1.09	48448	18.89	67.56	13.55
T3mo	5	citrate	6	2.20	96.65	1.15	48907	19.08	70.77	10.16
T3mo	5	his	6.5	1.86	97.12	1.02	47895	22.91	67.10	9.99
T3mo	5	tris	8	5.93	93.02	1.05	49015	24.67	63.29	12.04

Formulation and abbreviation key:

ace = 15 mM acetate

his = 15 mM histidine

succ = 15 mM succinate

water = formulated in water by dialysis

citrate = 15 mM citrate

tris = 15 mM TRIS

% HMW = percentage of high molecular weight species quantitated by SEC

% M = percentage of monomer quantitated by SEC

% LMW = percentage of low molecular weight species (fragments) quantitated by SEC

AUC = total integrated area of the SEC curve

% acidic = percentage of acidic species relative to the main species quantitated by IEC

% main = percentage of main species quantitated by IEC

% basic = percentage of basic species relative to the main species quantitated by IEX

T0 = time zero

T7d = 7 days of storage

T1mo = 1 month of storage

T3mo = 3 months of storage

Table 19. Stability at various temperatures of DVD-C at 1.0 mg/ml in 10 mM citrate + 10 mM phosphate buffer and at different pHs.

Time	Temp(°C)	pH	SEC data				IEX data		
			% HMW	% M	% LMW	AUC	% Acidic	% Main	% Basic
0	---	3	ND	ND	ND	ND	ND	ND	ND
0	---	4	0.78	97.97	1.25	106805	10.14	73.38	16.48
0	---	5	1.07	97.71	1.21	104165	9.80	73.84	16.35
0	---	6	1.23	97.62	1.15	99838	10.80	73.52	15.69
0	---	7	1.37	97.43	1.21	98566	10.45	73.79	15.76
0	---	8	1.48	97.30	1.22	93914	10.33	74.41	15.26
7d	40	3	ND	ND	ND	ND	ND	ND	ND
7d	40	4	0.70	94.51	4.79	99177	38.59	41.06	20.35
7d	40	5	1.09	97.43	1.48	105735	24.92	57.07	18.01
7d	40	6	1.30	97.16	1.54	101459	21.48	62.95	15.57
7d	40	7	1.53	96.38	2.09	94903	16.60	69.48	13.93
7d	40	8	2.00	95.74	2.25	95090	50.88	36.05	13.07
7d	50	3	ND	ND	ND	ND	ND	ND	ND
7d	50	4	ND	ND	ND	ND	ND	ND	ND

Time	Temp(°C)	pH	SEC data				IEX data		
			% HMW	% M	% LMW	AUC	% Acidic	% Main	% Basic
7d	50	5	2.72	93.91	3.37	102195	49.52	32.87	17.61
7d	50	6	2.23	95.55	2.22	99767	37.88	47.12	15.00
7d	50	7	1.67	94.59	3.74	96751	54.67	32.09	13.24
7d	50	8	1.82	94.13	4.05	85005	61.11	13.48	25.41
21d	5	3	ND	ND	ND	ND	ND	ND	ND
21d	5	4	0.01	96.54	3.45	105701	11.64	73.47	14.89
21d	5	5	0.00	96.29	3.71	102947	11.40	73.19	15.41
21d	5	6	0.01	96.33	3.66	98166	11.56	73.70	14.74
21d	5	7	0.01	96.24	3.75	96077	17.02	67.39	15.60
21d	5	8	0.01	95.63	4.36	91507	17.87	68.99	13.15
21d	25	3	ND	ND	ND	ND	ND	ND	ND
21d	25	4	0.08	95.89	4.02	107054	32.05	49.45	18.51
21d	25	5	0.07	96.85	3.08	104609	20.27	62.85	16.88
21d	25	6	0.05	96.86	3.09	100308	18.61	66.21	15.18
21d	25	7	0.08	96.40	3.52	97790	18.83	67.03	14.14
21d	25	8	0.07	95.82	4.12	92666	41.03	45.81	13.16
21d	40	3	ND	ND	ND	ND	ND	ND	ND
21d	40	4	0.09	88.58	11.33	79690	61.27	18.98	19.74
21d	40	5	0.20	95.13	4.67	105902	46.17	37.31	16.52
21d	40	6	0.23	95.96	3.81	101455	31.13	54.02	14.85
21d	40	7	0.33	94.32	5.35	98972	53.39	34.34	12.27
21d	40	8	0.37	93.05	6.57	91576	59.60	15.00	25.40
21d	50	3	ND	ND	ND	ND	ND	ND	ND
21d	50	4	ND	ND	ND	ND	ND	ND	ND
21d	50	5	0.20	91.14	8.66	90993	56.65	10.23	33.13
21d	50	6	0.34	93.57	6.09	100229	66.40	21.90	11.70
21d	50	7	0.54	89.92	9.54	91695	50.18	26.86	22.97
21d	50	8	0.47	87.11	12.41	67323	47.09	17.82	35.09
3mo	40	3	ND	ND	ND	ND	ND	ND	ND

Time	Temp(°C)	pH	SEC data				IEX data		
			% HMW	% M	% LMW	AUC	% Acidic	% Main	% Basic
3mo	40	4	27.59	40.51	31.91	61759	NR	NR	NR
3mo	40	5	5.12	85.68	9.19	119476	NR	NR	NR
3mo	40	6	2.79	91.55	5.67	113431	NR	NR	NR
3mo	40	7	3.40	88.33	8.27	106409	NR	NR	NR
3mo	40	8	5.41	80.86	13.73	98142	NR	NR	NR
3mo	25	3	ND	ND	ND	ND	ND	ND	ND
3mo	25	4	1.10	91.64	7.26	109681	53.69	23.99	22.32
3mo	25	5	1.24	96.19	2.57	106230	37.33	41.02	21.65
3mo	25	6	1.56	96.65	1.79	102674	27.46	53.25	19.29
3mo	25	7	1.85	95.59	2.56	100526	17.69	65.85	16.46
3mo	25	8	2.37	94.80	2.83	95500	65.32	20.08	14.59
3mo	5	3	ND	ND	ND	ND	ND	ND	ND
3mo	5	4	0.83	97.90	1.28	104383	18.59	60.63	20.78
3mo	5	5	1.14	97.86	1.00	101554	16.91	58.78	24.30
3mo	5	6	1.31	97.75	0.95	97897	17.38	63.16	19.46
3mo	5	7	1.54	97.43	1.03	96067	18.48	63.33	18.19
3mo	5	8	1.67	97.27	1.06	92532	20.44	61.36	18.21
6.5mo	25	3	ND	ND	ND	ND	ND	ND	ND
6.5mo	25	4	1.67	85.69	12.64	7753	70.60	10.11	19.28
6.5mo	25	5	1.35	94.61	4.04	7865	50.36	28.97	20.67
6.5mo	25	6	1.69	95.80	2.51	7719	36.64	46.73	16.63
6.5mo	25	7	2.12	94.15	3.74	7523	55.68	28.89	15.44
6.5mo	25	8	2.71	93.02	4.26	7155	68.76	16.22	15.01
6.5mo	5	3	ND	ND	ND	ND	ND	ND	ND
6.5mo	5	4	0.76	97.78	1.46	7508	19.30	59.53	21.17
6.5mo	5	5	1.07	97.92	1.00	7315	17.96	60.34	21.70
6.5mo	5	6	1.38	97.75	0.87	6969	19.99	60.32	19.69
6.5mo	5	7	1.61	97.38	1.02	6859	19.46	62.30	18.24
6.5mo	5	8	1.71	97.17	1.11	6501	21.06	59.64	19.30

## Abbreviation key:

% HMW = percentage of high molecular weight species quantitated by SEC  
 % M = percentage of monomer quantitated by SEC  
 % LMW = percentage of low molecular weight species (fragments) quantitated by SEC

AUC = total integrated area of the SEC curve

% acidic = percentage of acidic species relative to the main species quantitated by IEX

% main = percentage of main species quantitated by IEX

% basic = percentage of basic species relative to the main species quantitated by IEX

ND = no species detected

NR = assay not performed

0 = time zero

7d = 7 days of storage

21d = 1 month of storage

3mo = 3 months of storage

6.5mo = 6.5 months of storage

The molecule completely degraded during dialysis at pH 3. No species were detected by SEC or IEC after dialysis at this condition at time zero. Also, storage at 50°C at pH 4 yielded the same result. Both results are indicated by ND.

In addition, the thermal stability of DVD-C was assessed by differential scanning calorimetry (DSC). The thermal stability was evaluated at 1.0 mg/ml of the molecule formulated in 10 mM citrate + 10 mM phosphate buffer at pH 4, 5, 6, 7, or 8. A higher onset temperature of unfolding or higher domain midpoint temperature of unfolding means greater thermal stability. The thermal stability at different pHs is often correlated with the long-term stability of the molecule formulated at those pHs. Therefore, the DSC data can help identify the pHs at which the molecule is most and least stable.

Table 20. Differential Scanning Calorimetry Data of DVD-C at 1.0 mg/ml in 10 mM Citrate + 10 mM Phosphate Buffer and at Different pHs\*

pH	Onset (°C)	Tm1 (°C)	Tm2 (°C)	Tm3 (°C)	Tm4 (°C)
4	49.2	64.4	67.3	74.2	78.8
5	55.0	68.1	69.9	76.2	82.0
6	54.8	67.5	69.4	75.7	83.1

7	55.0	67.2	69.0	74.4	82.7
8	54.4	67.1	68.8	73.8	82.5

\* Numbered Tm values indicate the midpoint of the unfolding transitions.

As described above, the stability of DVD-C was tested in a number of different buffers and pHs, when stored at three different temperatures (40°C, 25°C, or 5°C). The concentration of DVD-C ranged from 1 mg/ml to 100 mg/ml. At 100 mg/ml, buffers included acetate, histidine, succinate, citrate, and tris. DVD-C was also formulated in plain water. The pH of the formulations ranged from 4 to 8. At 1 mg/ml, the protein was formulated in citrate-phosphate buffer with the pH ranging from 3 to 8. Once formulated, the samples of the DVD-Ig proteins were stored at the aforementioned temperatures. At specific time points, aliquots were taken and assessed for physical stability by SEC and chemical stability by weak cation exchange (WCX).

Overall, the data indicated DVD-C is stable except at pH 3 and pH 4. Specifically, the data suggest that physical stability of DVD-C is greatest at a pH near 5.5 and that chemical stability is highest at a pH near 6.0. Histidine and succinate were determined to be appropriate buffers for these pHs.

In addition, the thermal stability of DVD-C was assessed by differential scanning calorimetry (DSC), as described in Table 18. The thermal stability was evaluated at 1.0 mg/ml of DVD-C formulated in citrate-phosphate buffer at pHs 4 to 8. A higher onset temperature of unfolding (Ton) or higher domain midpoint temperature of unfolding (Tm) means greater thermal stability. Thermal stability is likely correlated with the long-term stability of the DVD-Ig protein. The data indicate similar thermal stability in the pH range of 5 to 8.

#### **EXAMPLE 9: Effect of a Polyol on the Stability of AS-DVD-Ig Proteins**

Dynamic scanning fluorescence (DSF) was employed to assess the propensity of a protein to unfold. The impact of polyols, *e.g.*, sucrose and sorbitol, was investigated in order to assess the effect of these polyols on the stability of the protein in solution. DVD-A, DVD-C and an IL12/IL18 DVD-Ig protein were used as exemplary AS-DVD-Ig proteins. As shown in Table 21 below, in general, AS-DVD-Ig proteins at various concentrations (1, 100,

150 mg/ml) are stable with the presence of sucrose (e.g., 40-160 mg/ml) and sorbitol (e.g., 20-80 mg/ml). Moreover, an increase in the concentration of sorbitol and sucrose provided resulted in a slight increased stability. Hence addition of sugars to a buffer (e.g., 15 mM histidine) containing protein formulation enhances the stability of AS-DVD-Ig proteins slightly

Table 21: Impact Of Polyols On The Thermodynamic Stability Of Various AS-DVD-Ig Proteins As Assessed By Dynamic Scanning Fluorescence At pH 6

<b>IL12IL18 DVD-Ig Protein Conc.</b>	<b>Onset of Unfolding (°C)</b>
1mg/ml, No Sorbitol	62.8
1mg/ml, 20 mg/ml Sorbitol	63.2
1mg/ml, 40 mg/ml Sorbitol	63.1
1mg/ml, 80 mg/ml Sorbitol	63.5
150 mg/ml, No Sorbitol	57.5
150 mg/ml, 20 mg/ml Sorbitol	57.6
150 mg/ml, 40 mg/ml Sorbitol	57.6
150 mg/ml, 80 mg/ml Sorbitol	58.5
1mg/ml, No Sucrose	62.8
1mg/ml, 40 mg/ml Sucrose	63.1
1mg/ml, 80 mg/ml Sucrose	64.1
1mg/ml, 160 mg/ml Sucrose	64.3
150 mg/ml, No Sucrose	57.5
150 mg/ml, 40 mg/ml Sucrose	58.2
150 mg/ml, 80 mg/ml Sucrose	58.3
150 mg/ml, 160 mg/ml Sucrose	58.3
DVD-C	
1mg/ml, No Sorbitol	62
1mg/ml, 20 mg/ml Sorbitol	62.3
1mg/ml, 40 mg/ml Sorbitol	62
1mg/ml, 80 mg/ml Sorbitol	62.1
100 mg/ml, No Sorbitol	47
100 mg/ml, 20 mg/ml Sorbitol	47.5
100 mg/ml, 40 mg/ml Sorbitol	47
100 mg/ml, 80 mg/ml Sorbitol	48.1
1mg/ml, No Sucrose	62
1mg/ml, 40 mg/ml Sucrose	62.1
1mg/ml, 80 mg/ml Sucrose	62
1mg/ml, 160 mg/ml Sucrose	62.2
100 mg/ml, No Sucrose	47
100 mg/ml, 40 mg/ml Sucrose	46.9
100 mg/ml, 80 mg/ml Sucrose	48.1
100 mg/ml, 160 mg/ml Sucrose	48
DVD-A	

1mg/ml, No Sorbitol	55.8
1mg/ml, 20 mg/ml Sorbitol	56
1mg/ml, 40 mg/ml Sorbitol	56
1mg/ml, 80 mg/ml Sorbitol	56.1
75 mg/ml, No Sorbitol	49.5
75 mg/ml, 20 mg/ml Sorbitol	50
75 mg/ml, 40 mg/ml Sorbitol	50.5
75 mg/ml, 80 mg/ml Sorbitol	50.2
1mg/ml, No Sucrose	55.8
1mg/ml, 40 mg/ml Sucrose	56
1mg/ml, 80 mg/ml Sucrose	56.4
1mg/ml, 160 mg/ml Sucrose	56.1
75 mg/ml, No Sucrose	49.5
75 mg/ml, 40 mg/ml Sucrose	49.5
75 mg/ml, 80 mg/ml Sucrose	50
75 mg/ml, 160 mg/ml Sucrose	51.1

As evidenced by the above data in Table 21, polyols (*e.g.*, sorbitol and sucrose) can improve stability of AS-DVD-Ig proteins in solution in a broad range (20 to 160 mg/ml sucrose and 20 to 80/ mg/ml sorbitol). Furthermore, the data reveal a decrease in the onset of unfolding, hence a decrease in the thermodynamic stability, with increasing DVD-Ig protein concentration. These data correlate with the observed instability, such as aggregation observed at high concentration liquid formulations.

## V. AS-DVD-IG PROTEINS ARE STABLE IN FORMULATIONS CONTAINING A BUFFER AND A POLYOL AT PH RANGE OF 4.5-7.5

### EXAMPLE 10: Effect of A Buffer on the Storage Stability of DVD-A in A Formulation Containing A Polyol

To assess the effect of a buffer on the storage stability of DVD-A in different buffers, the stability of the formulations was assessed before storage (T0) or after 7 days (7d) or 21 days (21d) of storage at 40°C (accelerated storage). DVD-A (85 mg/ml), an AS-DVD-Ig protein, was formulated in various buffers (15 mM citrate, 15 mM histidine, 15 mM arginine, 15 mM acetate, or water) at a pH of 5.2. Samples were filled into sterile vials (approx. 500 µl each) and stored under controlled conditions in temperature chambers and in the absence of light. The samples were analyzed using SEC and the results are provided in Tables 22 and 23.

Table 22: Effect of Buffer Type on Storage Stability of DVD-A As Measured By SEC

Formulation	Time Point	Mon	Agg	Frag
15 mM acetate, 80 mg/ml sucrose, pH 4.0	T0	95.96	2.75	1.28
	7d, 40C	71.84	24.93	3.21
	21d, 40C	66.62	26.91	6.45
15 mM acetate, 80 mg/ml sucrose, pH 5.2	T0	96.06	2.78	1.14
	7d, 40C	89.19	8.76	2.03
	21d, 40C	90.93	4.41	4.65
15 mM arginine, 80 mg/ml sucrose, pH 5.2	T0	95.84	3.07	1.08
	7d, 40C	89.62	8.54	1.82
	21d, 40C	81.03	12.41	6.54
15 mM citrate, 80 mg/ml sucrose, pH 5.2	T0	95.72	3.19	1.08
	7d, 40C	90.54	6.96	2.48
	21d, 40C	87.39	7.74	4.86
15 mM histidine, 80 mg/ml sucrose, pH 5.2	T0	96	2.9	1.09
	7d, 40C	91.04	6.91	2.03
	21d, 40C	87.58	8.63	3.78
Water, 80 mg/ml sucrose, pH 5.2	T0	94.89	4.04	1.05
	7d, 40C	88.4	9.76	1.82
	21d, 40C	81.41	14.39	4.18

Table 23: Effect of Buffer Type on Storage Stability of DVD-A As Measured By IEC

Formulation	Time Point	Main	Acidic	Basic
15 mM acetate, 80 mg/ml sucrose, pH 4.0	T0	56.88	9.43	33.68
	7d, 40C	46.91	18.74	34.33
	21d, 40C	34.62	24.82	40.54
15 mM acetate, 80 mg/ml sucrose, pH 5.2	T0	57.2	9.45	33.34
	7d, 40C	47.61	18.74	33.64
	21d, 40C	35.29	33.35	31.35
15 mM arginine, 80 mg/ml sucrose, pH 5.2	T0	63.44	9.93	26.62
	7d, 40C	49.22	19.57	31.2
	21d, 40C	37.87	25.3	36.81
15 mM citrate, 80 mg/ml sucrose, pH 5.2	T0	59.59	7.88	35.51
	7d, 40C	45.92	20.82	33.24
	21d, 40C	31.97	31.81	36.21
15 mM histidine, 80 mg/ml sucrose, pH 5.2	T0	58.29	8.18	33.52
	7d, 40C	44.97	16.84	38.17
	21d, 40C	36.33	24.88	38.77
Water, 80 mg/ml sucrose, pH 5.2	T0	57.05	9.53	33.41
	7d, 40C	48.23	19.96	37.15
	21d, 40C	37.69	25.15	37.15

SEC results provided in Table 22 show that a pH 5.2 histidine formulation compared to the pH 4 histidine formulation had the lowest level of aggregates, although citrate and acetate buffers showed low levels of aggregates as well. Although citrate and acetate showed

less soluble aggregates as measured by SEC, the solutions were visibly turbid indicating formation of insoluble aggregates. The visual turbidity was not significant, however, given the SEC measurements. Overall, citrate and acetate formulations were only slightly less stable than the histidine formulations. Given that the noise/error in IEC measurement is usually higher, no significant differences in the chemical stability were observed within the formulations presented in Table 23. The polyol only formulation was also slightly less stable as compared to the histidine formulation. Given the overall stability characteristics of proteins identified as AS-DVD-Ig proteins, the slight differences observed in the SEC and IEC analysis of Tables 22 and 23 showed that each of the tested buffers at pH 5.2 was stable. Thus, the differences observed between the tested buffers at pH 5.2 indicated that each could be used to provide stable formulations for AS-DVD-Ig proteins, including those at high concentrations.

## **VI. AS-DVD-IG PROTEINS ARE STABLE IN FORMULATIONS CONTAINING A BUFFER, A POLYOL, AND A SURFACTANT AT PH RANGE OF 4.5-7.5**

### **EXAMPLE 11: AS-DVD-Ig Proteins Are Stable in A Range Of Histidine Formulations Containing Surfactants and Sugars**

The following example describes the impact of pH, buffers, and excipients (including surfactants and polyols) on the physico-chemical stability of DVD-Ig proteins at low and high concentration formulations during accelerated / real time stability testing. In order to demonstrate that DVD-Ig proteins have distinctly different protein properties compared to monoclonal antibodies, five different formulation conditions that are widely known and used to maintain the stability of monoclonal antibodies were tested.

Using size exclusion chromatography (SEC) and ion exchange chromatography (IEC), the storage stability of low concentration (1 mg/ml) and higher concentration (100 mg/ml) AS-DVD-Ig protein formulations was evaluated following three storage conditions: no storage (T0), 1 month at a controlled temperature of 5 °C (1m, 5C), and 1 month at a controlled temperature of 40 °C (1m, 40C). Formulations with varying pH (a range of 5.25 to 7.2 was selected), buffers, and excipients were tested, according to the following conditions:

- 1) pH 5.25 and pH 6, 15 mM histidine, 80 mg/ml sucrose, 0.01% Tween 80;
- 2) pH 6, 15 mM histidine, 40 mg/ml sorbitol, 0.01% Tween 80;
- 3) PBS (10 mM phosphate, 125 mM NaCl) at pH 6 and 7.2;
- 4) 20 mM glycine, 26 mg/ml glycerol pH 6; and
- 5) Water, 0.01% Tween 80 pH 5 and 6.

The results are presented in Tables 24-27 below. The data show that not all DVD-Ig proteins are stable in all tested pH and formulation conditions. DVD-B formed high amounts of aggregates under all the solution conditions tested and were classified as non-AS-DVD-Ig proteins (in accordance with the assay presented in Example 4). All the other DVD-Ig proteins (previously selected as being AS-DVD-Ig proteins) behaved well and were stable.

Sucrose, sorbitol, glycerol, and glycine were used to evaluate the effect of these excipients. Tween 80 (polysorbate 80), a surfactant that provides stabilization against shaking stress, was also used to evaluate its impact on the stability of high concentration solutions. The impact of salt concentration was evaluated by varying the ionic strength using sodium chloride.

In general, a formulation at pH 6 or pH 5.2 in a histidine buffer was effective for all AS-DVD-Ig proteins. Both sorbitol and sucrose each improved stability. Sucrose resulted in the formation of slightly less monomer after defined time points. The presence of salt resulted in increased instability (defined as less monomer remaining), especially in the case of the IL12/IL18 DVD-Ig protein of Table 24 at 100 mg/mL and to a lesser extent at 1 mg/mL, as shown in Table 27. The presence of glycerol and glycine resulted in less monomer remaining following shelf stability as compared to other formulations. For example, according to Table 24, the IL12IL18 DVD-Ig protein was slightly more stable at pH 6 in the presence of sucrose than sorbitol. The data also demonstrate that the formulations with either none or very little ionic strength showed comparable stability over time.

SEC data showed the physical stability of the DVD-Ig proteins, wherein the rate of formation of aggregates and/or fragments was evaluated (see Table 24 & 26). As shown in Table 26 all molecules showed no significant tendency to aggregation at low concentrations. IEC data is an indicator of the chemical stability of a DVD-Ig protein. Deamidation, for example, results in formation of acidic species (conversion of the main species to acidic species). Generation of positively charged variants would lead to an increase in the basic

species. Formation of any of the two acidic or basic species indicates instability, as the formation of these two species results in an overall decrease in the % of main species (see Table 25 & 27).

The results in Tables 24 to 27 suggest that an AS-DVD-Ig protein is stable in formulations comprising a surfactant alone, *e.g.*, 0.01% Tween 80 at pH 5 - 6.

Table 24: 100 mg/ml SEC Data for Various Stored Formulations

DVD-Ig	Formulation	Time Point, temp	Mon	Agg	Frag
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	96.53	1.1	2.36
		1m, 5C	95.97	1.75	2.26
		1m, 40C	92.58	2.86	4.55
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	96.31	1.15	2.52
		1m, 5C	96	1.71	2.27
		1m, 40C	92.11	3.5	4.38
IL12IL18	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	96.61	1.25	2.13
		1m, 5C	95.71	1.78	2.5
		1m, 40C	91.62	4.04	4.33
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	89.61	10.38	0
		1m, 5C	89.48	10.51	0
		1m, 40C	85.85	9.96	4.18
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	88.58	11.41	0
		1m, 5C	86.8	13.19	0
		1m, 40C	83.09	12.68	4.21
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	93.08	0.93	5.98
		1m, 5C	92.59	1.67	5.72
		1m, 40C	85.25	11.47	3.27
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	93.34	0.85	5.79
		1m, 5C	92.45	1.71	5.82
		1m, 40C	86.15	10.69	3.14
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	97.57	0.62	1.8
		1m, 5C	97.6	0.92	1.47
		1m, 40C	91.65	5	3.34
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	97.41	0.71	1.87
		1m, 5C	97.69	1.07	1.22
		1m, 40C	92.68	4.3	3.01
66	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	97.65	0.61	1.73
		1m, 5C	97.52	1.04	1.42
		1m, 40C	91.53	3.98	4.47
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 6	T0	96.13	1.57	2.28
		1m, 5C	94.2	3.34	2.44
		1m, 40C	88.48	7.3	4.2
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 7.2	T0	96.07	1.66	2.26
		1m, 5C	94.64	2.91	2.43
		1m, 40C	87.71	8.03	4.25
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 6	T0	96.24	2.11	1.64
		1m, 5C	94.66	3.85	1.48
		1m, 40C	41.53	53.68	4.77

DVD-Ig	Formulation	Time Point, temp	Mon	Agg	Frag
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 7.2	T0	95.75	2.45	1.79
		1m, 5C	95.11	3.62	1.26
		1m, 40C	33.96	60.56	5.46
IL12IL18	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	T0	96.48	1.27	2.24
		1m, 5C	95.06	2.07	2.86
		1m, 40C	90.33	4.7	4.95
DVD-B	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	T0	96.29	1.88	1.81
		1m, 5C	95.93	2.62	1.43
		1m, 40C	23.32	73.53	3.14
IL12IL18	Water, 0.01% Tween 80, pH 5.0	T0	95.06	2.55	2.37
		1m, 5C	94.12	2.97	2.89
		1m, 40C	90.99	4.93	4.07
IL12IL18	Water, 0.01% Tween 80, pH 6.0	T0	94.59	2.88	2.52
		1m, 5C	94.22	3.28	2.48
		1m, 40C	90.9	4.82	4.26
5	Water, 0.01% Tween 80, pH 5.0	T0	65.33	31.34	3.31
		1m, 5C	17.02	79.12	3.84
		1m, 40C	82.54	13.41	4.03
5	Water, 0.01% Tween 80, pH 6.0	T0	43.43	53.33	3.23
		1m, 5C	17.53	79.15	3.3
		1m, 40C	74.6	21.38	4.03

Table 25: 100 mg/ml IEC Data for Various Stored Formulations

DVD-Ig	Formulation	Time Point, temp	Main	Acidic	Basic
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	58.32	27.98	13.68
		1m, 5C	57.54	26.72	15.73
		1m, 40C	44.91	37.65	17.43
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	58.63	27.84	13.51
		1m, 5C	56.89	29.15	13.94
		1m, 40C	41.47	44.38	14.14
IL12IL18	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	58.1	28.09	13.79
		1m, 5C	60.53	27.48	11.98
		1m, 40C	41.02	44.69	14.29
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	42.99	11.66	45.34
		1m, 5C	42.53	10.26	47.19
		1m, 40C	34.99	25.55	39.45
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	41.03	11.84	47.12
		1m, 5C	41.73	10.91	47.34
		1m, 40C	36.18	25.94	37.86
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	49.64	39.02	11.33
		1m, 5C	45.48	38.77	15.73
		1m, 40C	51.17	30.64	18.18
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	50	39.1	10.88
		1m, 5C	49.31	38.95	11.72
		1m, 40C	29.83	52.65	17.5
66	15 mM Histidine, 80 mg/ml	T0	63.78	24.78	11.43

DVD-Ig	Formulation	Time Point, temp	Main	Acidic	Basic
	Sucrose, 0.01 % Tween, pH 5.25	1m, 5C	60.03	25.27	14.69
		1m, 40C	42.13	41.93	15.93
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	61.5	26.21	12.27
		1m, 5C	59.06	25.85	15.07
		1m, 40C	42.31	43.47	41.2
66	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	57.18	24.54	18.26
		1m, 5C	61.9	26.5	11.59
		1m, 40C	41.24	45.29	13.46
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 6	T0	58.2	28.1	13.69
		1m, 5C	57.43	27.35	15.21
		1m, 40C	40.68	42.04	17.27
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 7.2	T0	54.51	27.99	17.48
		1m, 5C	53.79	28.08	18.11
		1m, 40C	27.08	58.25	14.65
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 6	T0	52.94	27.32	19.72
		1m, 5C	54.38	27.19	18.42
		1m, 40C	29.14	38.63	32.21
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 7.2	T0	27.93	52.59	19.46
		1m, 5C	54.07	26.57	19.35
		1m, 40C	19.79	48.61	31.59
IL12IL18	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	T0	57.95	28.99	13.05
		1m, 5C	57.85	29.95	12.18
		1m, 40C	37.55	47.49	14.94
DVD-B	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	T0	30.77	51	18.22
		1m, 5C	50	33.32	16.67
		1m, 40C	27.41	58.52	14.05
IL12IL18	Water, 0.01% Tween 80, pH 5.0	T0	56.03	27.31	16.64
		1m, 5C	57.67	28.06	14.26
		1m, 40C	44.97	41.34	13.68
IL12IL18	Water, 0.01% Tween 80, pH 6.0	T0	58.4	28.57	13.02
		1m, 5C	58.93	26.36	14.7
		1m, 40C	45.37	39.4	15.21
5	Water, 0.01% Tween 80, pH 5.0	T0	46.39	19.48	34.12
		1m, 5C	16.75	8.54	74.69
		1m, 40C	44.57	36.64	18.78
5	Water, 0.01% Tween 80, pH 6.0	T0	32.31	14.89	52.78
		1m, 5C	16.32	8.27	75.4
		1m, 40C	39.7	35.23	25.06

Table 26: 1 mg/ml SEC Data for Various Stored Formulations

DVD-Ig	Formulation	Time Point, temp	Mon	Agg	Frag
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	97.87	0.73	1.39
		1m, 5C	97.83	0.57	1.59
		1m, 40C	95.55	0.74	3.7
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	98.04	0.32	1.62
		1m, 5C	97.95	0.24	1.79

DVD-Ig	Formulation	Time Point, temp	Mon	Agg	Frag
		1m, 40C	95.83	0.58	3.58
IL12IL18	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	96.2	0.56	3.22
		1m, 5C	96.41	0.47	3.1
		1m, 40C	94.21	0.66	5.12
		T0	92.05	7.94	0
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	1m, 5C	99.23	0.76	0
		1m, 40C	95.52	0.53	3.94
		T0	93.59	6.4	0
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	1m, 5C	99.33	0.66	0
		1m, 40C	95.3	0.59	4.1
		T0	94.74	0.51	4.73
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	1m, 5C	94.34	0.49	5.15
		1m, 40C	93.06	0.91	6.02
		T0	94.33	0.37	5.29
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	1m, 5C	93.98	0.36	5.65
		1m, 40C	92.71	0.87	6.41
		T0	98.76	0.51	0.72
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	1m, 5C	98.59	0.51	0.89
		1m, 40C	96.37	1.02	2.59
		T0	98.82	0.2	0.96
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	1m, 5C	98.9	0.16	0.92
		1m, 40C	96.82	0.66	2.51
		T0	97.87	0.29	1.83
66	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	1m, 5C	97.3	0.23	2.45
		1m, 40C	95.5	1.02	3.48
		T0	95.38	1.54	3.07
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 6	1m, 5C	95.29	1.54	3.15
		1m, 40C	91.87	1.95	6.17
		T0	94.36	2.73	2.89
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 7.2	1m, 5C	93.78	2.99	3.22
		1m, 40C	86.45	3.64	9.9
		T0	95.82	1.66	2.5
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 6	1m, 5C	95.54	1.53	2.91
		1m, 40C	91.39	2.31	6.29
		T0	94.87	2.43	2.68
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 7.2	1m, 5C	95.01	2.3	2.68
		1m, 40C	88.42	3.24	8.32
		T0	96.48	0.36	3.14
IL12IL18	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	1m, 5C	96.51	0.29	3.18
		1m, 40C	93.82	0.4	5.77
		T0	96.05	1.33	2.6
DVD-B	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	1m, 5C	95.47	1.15	3.37
		1m, 40C	94.14	1.22	4.63
		T0	89.16	2.42	8.41
IL12IL18	Water, 0.01% Tween 80, pH 5.0	1m, 5C	95.1	1.86	3.02
		1m, 40C	89.69	4.11	6.18
		T0	94.24	2.88	2.86
IL12IL18	Water, 0.01% Tween 80, pH 6.0	1m, 5C	94.89	2.6	2.49
		1m, 40C	90.66	3.38	5.94

DVD-Ig	Formulation	Time Point, temp	Mon	Agg	Frag
5	Water, pH 5.0	T0	69.49	26.36	4.13
		1m, 5C	68.87	24.88	6.23
		1m, 40C	91.35	3.6	5.08
5	Water, pH 6.0	T0	49.15	46.65	4.19
		1m, 5C	51.96	43.27	4.76
		1m, 40C	92.56	2.45	4.97

Table 27: 1 mg/ml IEC Data for the Various Stored Formulations

DVD-Ig	Formulation	Time Point, temp	Main	Acidic	Basic
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	61.83	24	14.14
		1m, 5C	61.76	26.77	11.46
		1m, 40C	41.58	39.69	18.72
IL12IL18	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	61.31	26.12	12.55
		1m, 5C	59.67	25.3	15.02
		1m, 40C	42.98	45.85	11.15
IL12IL18	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	60.81	25.86	13.31
		1m, 5C	59.07	27.12	13.8
		1m, 40C	38.22	47.55	14.21
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	41.13	10.3	48.56
		1m, 5C	46.96	10.38	42.64
		1m, 40C	36	24.57	39.42
6	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	44.43	9.92	45.63
		1m, 5C	46.33	11.55	42.11
		1m, 40C	37.38	26.09	36.52
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	50.6	39.58	9.81
		1m, 5C	49.57	37.75	11.66
		1m, 40C	33.38	50.27	16.33
65	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	51.74	38.73	9.52
		1m, 5C	53.21	36.58	10.2
		1m, 40C	35.11	49.96	14.92
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 5.25	T0	59.89	25.96	14.14
		1m, 5C	58.1	25.05	16.84
		1m, 40C	45.43	37.21	17.34
66	15 mM Histidine, 80 mg/ml Sucrose, 0.01 % Tween, pH 6	T0	58.18	24.86	16.94
		1m, 5C	61.69	24.94	13.36
		1m, 40C	44.83	41.39	13.77
66	15 mM Histidine, 40 mg/ml Sorbitol, 0.01 % Tween, pH 6	T0	61.49	24.96	13.53
		1m, 5C	60.34	24.96	16.69
		1m, 40C	38.32	50.92	10.75
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 6	T0	58.52	26.18	15.29
		1m, 5C	58.98	27.17	13.84
		1m, 40C	37.54	47.49	14.96
IL12IL18	10 mM Phosphate, 125 mM NaCl, pH 7.2	T0	63.15	23.13	13.71
		1m, 5C	58.58	26.07	15.33
		1m, 40C	16.92	73.33	9.74
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 6	T0	54.38	27.35	18.25
		1m, 5C	56.48	27.42	16.09

DVD-Ig	Formulation	Time Point, temp	Main	Acidic	Basic
		1m, 40C	30.99	50.68	18.32
DVD-B	10 mM Phosphate, 125 mM NaCl, pH 7.2	T0	54.86	30	15.12
		1m, 5C	52.66	31.63	15.69
		1m, 40C	8.64	60.51	30.84
		T0	57.58	27.8	14.6
IL12IL18	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	1m, 5C	58.02	29.5	12.46
		1m, 40C	28.8	60.9	10.29
		T0	53.05	30.46	16.48
TNFPGE 2	20 mM Glycine, 26 mg/ml Glycerol, pH 6.0	1m, 5C	44.21	33.97	21.8
		1m, 40C	5.52	81.17	13.3
		T0	57.76	26.51	15.71
IL12IL18	Water, 0.01% Tween 80, pH 5.0	1m, 5C	58.44	26.13	15.41
		1m, 40C	36.16	44.81	19.02
		T0	59.24	26.09	14.66
IL12IL18	Water, 0.01% Tween 80, pH 6.0	1m, 5C	58.66	26.96	14.36
		1m, 40C	35.97	44.16	19.85
		T0	51.73	19.4	28.85
5	Water, 0.01% Tween 80, pH 5.0	1m, 5C	48.63	20.14	31.21
		1m, 40C	44.52	41.73	13.73
		T0	34.34	14	51.62
5	Water, 0.01% Tween 80, pH 6.0	1m, 5C	32.27	15.38	47.33
		1m, 40C	42.03	44.28	13.67

**EXAMPLE 12. Impact of Storage on the Stability of an AS-DVD-Ig Protein, (DVD-C), In Various Formulations**

The pH and the storage temperature of a protein formulation are two important factors influencing protein stability during accelerated/long-term storage. To assess the impact of these factors, the DVD-Ig protein was exposed to short-term storage at elevated and real time temperatures in order to gain insight into the formulation feasibility of long-term storage at lower temperatures (e.g., 2-8°C).

The storage stability of DVD-C in solution (100 mg/ml) was evaluated in formulations at 40°C. After defined storage periods, samples were pulled and the impact of storage time on DVD-Ig protein stability was evaluated. Samples were filled into sterile vials (approx. 500 µl each) and stored under controlled conditions, in a temperature chamber and in the absence of light. At predefined points of time, samples of prepared solutions were pulled for analysis according to the sample pull scheme. The resulting data is provided in Tables 28 and 29.

Table 28: Impact of Storage of DVD-C at 100 mg/ml Concentrations in Various Conditions As Measured By SEC

Formulation	Time Point	Mon	Agg	Frag
15 mM acetate 80 mg/ml sucrose 0.02 % Tween 80 pH 5	T0	97.66	1.09	1.20
	7d, 40°C	96.32	1.70	1.98
	1m, 40°C	94.18	2.43	3.39
15 mM histidine 80 mg/ml sucrose 0.02 % Tween 80 pH 6	T0	97.66	1.091	1.20
	7d, 40°C	95.92	2.09	1.99
	1m, 40°C	93.64	2.72	3.64

Table 29: Impact of Storage of DVD-C at 100 mg/ml Concentrations in Various Conditions As Measured By IEC

Formulation	Time Point	Main	Acidic	Basic
15 mM acetate 80 mg/ml sucrose 0.02 % Tween 80 pH 5	T0	73.84	9.80	16.35
	T2m, 5°C	69.18	10.88	19.95
15 mM histidine 80 mg/ml sucrose 0.02 % Tween 80 pH 6	T0	73.84	9.80	16.35
	T2m, 5°C	67.99	10.59	21.42

The data provided in Tables 28 and 29 show that DVD-C was very stable (compared to some unstable DVD-Ig proteins, for example, in Example 5) in that only minimal loss in monomer levels occurred during the test storage conditions and hence would be classified as an AS-DVD-Ig protein.

#### **EXAMPLE 13: Effect of a Surfactant on the Stability of AS-DVD-Ig Proteins in Buffer and Polyol Containing Formulations**

It is generally beneficial to set a formulation pH more than 1 unit from the protein's isoelectric point (pI). The more a formulation pH approximates the pI, generally, the more the overall surface of the protein is regarded as uncharged, thus contributing to protein-protein attraction of non-polar groups, and thus enhancing non-covalent aggregation and instability. Shaking and stirring foster physical instability, creating hydrophobic air/water interfaces, which result in alignment of protein molecules at these interfaces, and eventually result in aggregation. Given that air is more hydrophobic than water, the interface between air and liquid is deemed to be a denaturing surface at which aggregation, especially of (partially) unfolded proteins can originate. The effective air-water interface can be increased by shaking or stirring.

In the following example, the effect of various concentrations of a surfactant (*e.g.*, Tween 80) on the instability of exemplary DVD-Ig proteins was evaluated. The study was

done in the absence or presence of the polyol sucrose. The data presented in Tables 30 and 31 compare various surfactant concentrations (0 to 2 mg/ml) in a histidine buffer at pH 5.2 or 5.4 with and without sucrose (80 mg/ml). Results of turbidity measurements show that a surfactant (Tween 80) in a concentration range of 0.05 mg/ml – 2 mg/ml provided stability against shear/ denaturation stress to AS-DVD-Ig proteins in general. The turbidity increased upon lowering the surfactant concentration to 0.01 mg/ml. Similar observations were made for AS-DVD-Ig proteins in the presence of sucrose. All studies were conducted at 15 mM histidine at a DVD-Ig protein concentration of 1 mg/ml. The corresponding SEC analysis showed that, in general, a significant loss in rel. % monomer for samples that contained 0.05 - 0.1 mg/mL Tween80, indicating that the amount of Tween 80 that yielded the most stable formulations is between 0.5 mg/mL and 2 mg/mL. Turbidity measurements at 500 nm using UV are listed in Table 30 and the SEC analysis of the samples is listed in Table 31. Ranges of pH were also tested in the formulations described below.

Table 30: Effect of Tween on the Stability of Various DVD-Ig Proteins as Assessed by Air/Liquid Interface Denaturation Study/Shaking Study\* by measurements at 500 nm using UV

DVD Form:	DVD-A, 15 mM Histidine, pH 5.2, 80 mg/ml Sucrose			DVD-C, 15 mM Histidine, pH 5.4, 80 mg/ml Sucrose			IL12IL18, 15 mM Histidine, pH 5.4, 80 mg/ml Sucrose		
Tween 80	<b>T0</b>	<b>T24h</b>	<b>T96h</b>	<b>T0</b>	<b>T24h</b>	<b>T96h</b>	<b>T0</b>	<b>T24h</b>	<b>T96h</b>
0	0.11	0.07	0.55	0.085	0.08	0.107	0.051	0.051	0.051
2 mg/ml	0.069	0.018	0.004	0.085	0.038	0.003	0.069	0.072	0.017
0.5 mg/ml	0.014	0.014	0.002	0.025	0.036	0.004	0.09	0.006	0.008
0.1 mg/ml	0.061	0.02	0.001	0.043	0.007	0.002	0.055	0.011	0.0109
0.05 mg/ml	0.018	0.006	0.001	0.019	0.008	0.002	0.022	0.008	0.0208
0.01 mg/ml	0.007	0.007	1.03	0.03	0.033	0.22	0.057	0.081	0.082

	DVD-A, 15 mM Histidine, pH 5.2			DVD-C, 15 mM Histidine, pH 5.4			IL12IL18, 15 mM Histidine, pH 5.4		
Tween 80	<b>T0</b>	<b>T24h</b>	<b>T96h</b>	<b>T0</b>	<b>T24h</b>	<b>T96h</b>	<b>T0</b>	<b>T24h</b>	<b>T96h</b>
0	0.045	0.038	0.29	0.015	0.045	0.038	0.046	0.046	0.038
2 mg/ml	0.049	0.024	0.004	0.015	0.02	0.003	0.097	0.03	0.018

0.5 mg/ml	0.016	0.058	0.004	0.03	0.019	0.003	0.049	0.031	0.013
0.1 mg/ml	0.005	0.005	0.002	0.004	0.005	0.002	0.012	0.021	0.017
0.05 mg/ml	0.004	0.011	0.002	0.005	0.001	0.009	0.005	0.015	0.017
0.01 mg/ml	0.005	0.03	0.66	0.01	0.015	0.022	0.05	0.022	0.041

\* The OD 500 was measured using UV

Table 31: Effect of Tween on the Stability of Various DVD-Ig Proteins As Assessed By Air/Liquid Interface Denaturation Study/Shaking Study\*

Tween	DVD Form: DVD-A, 15 mM Histidine, pH 5.2, 80 mg/ml Sucrose								DVD Form: DVD-C, 15 mM Histidine, pH 5.4, 80 mg/ml Sucrose								
	T0				T96h				T0				T96h				
	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	
0	1.7	97.25	1.03	73550	1.35	97.55	1.08	39881	1.96	96.79	1.23	64166	1.35	97.55	1.08	39881	
2	1.67	97.58	0.74	759057	2.33	96.67	0.99	74778	2.16	96.4	1.43	72649	2.33	96.67	0.99	74778	
0.5	1.76	97.43	0.8	75635	2.35	96.42	1.21	74782	2	96.66	1.33	71044	2.35	96.42	1.21	74782	
0.1	1.65	97.66	0.68	73158	3.85	95.06	1.07	72816	2.09	96.43	1.47	70277	3.85	95.06	1.07	72816	
0.05	1.66	97.61	0.71	77474	18.86	80.36	0.76	76431	2.01	96.56	1.41	700701	18.86	80.36	0.76	76431	
0.01	1.68	97.5	0.81	77410	9.32	83.25	7.41	11267	2.11	96.54	1.33	70397	9.32	83.25	7.41	11267	
Tween	DVD Form: IL12IL18, 15 mM Histidine, pH 5.4, 80 mg/ml Sucrose								DVD Form: DVD-A, 15 mM Histidine, pH 5.2								
	T0				T96h				T0				T96h				
	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	
80	7.15	91.18	1.66	76420	6.11	92.04	1.83	72538	1.59	97.58	0.81	76481	1.09	98.02	0.87	60575	
mg/ml	0	7.29	90.64	2.05	77867	6.09	91.94	1.96	76447	1.55	97.65	0.79	85152	1.63	97.49	0.87	84633
	2	7.04	91.4	1.54	77164	6.14	91.98	1.87	75844	1.65	97.64	0.7	88543	na	na	na	na
	0.5	7.08	91.23	1.68	72875	6.4	90.9	2.68	71645	1.69	97.57	0.73	85244	1.61	97.66	0.72	87583
	0.1	7.03	91.3	1.65	69962	8.8	86.44	4.75	72941	1.72	97.57	0.7	86031	2.22	96.67	1.09	86215
	0.05	6.81	91.57	1.61	77682	12.77	85.19	2.03	70341	1.63	97.63	0.72	81730	28.99	69.37	1.63	15513
Tween	DVD Form: DVD-C, 15 mM Histidine, pH 5.4								DVD Form: IL12IL18, 15 mM Histidine, pH 5.4								
	T0				T96h				T0				T96h				
	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	Agg	Mon	Frag	AUC	
80	2.05	96.56	1.37	71752	1.71	97.3	0.98	67869	6.9	91.47	1.61	75651	6.34	91.6	2.05	73693	
mg/ml	0	2.08	96.63	1.28	75988	2.03	96.58	1.37	75354	6.77	91.76	1.46	75316	6.43	91.61	1.95	75143
	2	2.1	96.44	1.45	77324	2	96.49	1.5	76488	6.88	91.57	1.53	75439	6.12	91.84	2.03	74397
	0.5	2.11	96.33	1.55	77859	1.86	96.59	1.54	76809	7.02	91.18	1.78	74981	7.06	90.81	2.11	74698
	0.1	2.21	96.28	1.5	79134	2.38	96.09	1.52	77153	7.04	91.37	1.57	76385	5.61	92.44	1.93	75375
	0.05	2.08	96.44	1.47	76726	2.95	95.52	1.51	74056	6.66	91.69	1.64	65218	10.76	87.29	1.93	62074

\* The SEC data corresponds to 0 and 96 h shaking samples of Table 30.

### Example 14: Effect of Surfactant Concentration on the Stability of IL12IL18 DVD-Ig Protein as Measured in Shaking Studies

The following example describes the effect of different concentrations (a range of concentration) of polysorbate 80 and poloxamer on the shaking stability of an IL12IL18 DVD-Ig protein at concentrations of 1 mg/mL at pH 6 (15 mM histidine + 80 mg/mL sucrose) as measured using optical density at two different wavelengths of 350 and 500 nm. Samples were taken after 0, 24, 48, 120, and 240 hours.

Table 32: Effect of Surfactants on Formulation with Buffer and Polyol

		IL12IL18 1mg/ml; 15 mM Histidine + 80 mg/mL sucrose pH 6										
Surfactant	Shake Time (H)	OD500 nm					OD350 nm					
		0	24	48	120	240	0	24	48	120	240	
Tween 80 (mg/ml)	0	0.003	0.06	0.085	0.137	0.387	0.01	0.097	0.135	0.22	0.574	
	0.01	0.0036	0.029	0.011	0.065	0.108	0.01	0.01	0.03	0.167	0.227	
	0.05	0.0056	0.003	0.01	0.087	0.113	0.013	0.01	0.027	0.17	0.22	
	0.1	0.0036	0.002	0.0022	0.067	0.004	0.01	0.01	0.009	0.135	0.011	
	0.5	0.003	0.002	0.002	0.04	0.001	0.01	0.009	0.009	0.086	0.009	
	2	0.035	0.002	0.0025	0.0025	0.002	0.062	0.012	0.013	0.014	0.013	
		IL12IL18 1mg/ml; 15 mM Histidine + 80 mg/mL sucrose pH 6										
		500 nM					350 nM					
Surfactant	Shake Time (H)	0	24	48	120	240	0	24	48	120	240	
Poloxamer (% w/v)	0	0.003	0.035	0.044	0.236	0.226	0.01	0.06	0.08	0.4	0.374	
	0.01	0.004	0.004	0.007	0.013	0.0126	0.01	0.012	0.019	0.031	0.034	
	0.05	0.003	0.014	0.002	0.027	0.002	0.01	0.02	0.011	0.009	0.011	
	0.1	0.003	0.035	0.0037	0.0038	0.003	0.01	0.01	0.012	0.012	0.012	
	0.5	0.003	0.003	0.003	0.003	0.004	0.01	0.01	0.011	0.011	0.013	
	2	0.0036	0.003	0.003	0.005	0.004	0.011	0.01	0.01	0.013	0.013	

As described in Table 32, the addition of surfactants increased the shaking stability of the IL12/IL18 DVD-Ig protein. Polysorbate concentration in the range of 0.05-2 mg/mL was determined to be the most effective for stability of the IL12/IL18 DVD-Ig protein, and poloxamer was determined to be most effective in the range of 0.1-2 % w/v. DVD-B was also tested and showed similar stability when tested in this particular shaking assay.

**Example 15: Effect of Polyol (Sucrose and Sorbitol) Concentration on the Stability of IL12IL18 and DVD-B DVD-Ig Protein**

The following example shows the effect of polyols sucrose and sorbitol in the presence of polysorbate 80 on the stability of the IL12IL18 DVD-Ig protein as measured at 3mg/mL, at pH 6 by intrinsic fluorescence using an automated high throughput instrument Optim-1000 from Avacta (York, UK) as described in the methods section. 9  $\mu$ l MCA's were used for the study. Thermal scans were obtained using a scan rate of 1°C/minute and scans were taken from 25-75°C. All samples were freshly prepared from stock solutions for the sucrose and sorbitol excipients.

Table 33: Effect of polyols on 3 mg/ml IL12IL18 and DVD-B DVD-Ig protein formulations with 15 mM His buffer at pH 6 and polysorbate

Sucrose (mg/ml)	Sorbitol (mg/ml)	Tween	DSF (°C) IL12IL18	DSF (°C) DVD-B
0		0.01%	60	54
10		0.01%	61	55
40		0.01%	61	55
70		0.01%	60	56
100		0.01%	60	57
	0	0.01%	61	53
	10	0.01%	62	54
	20	0.01%	61	55
	40	0.01%	62	55
	60	0.01%	62	56

Based on the data in Table 33, the stability of the DVD-B Ig protein increased with an increase in the concentration of either polyol (sucrose or sorbitol). It was determined that 100 mg/mL for sucrose and 60 mg/mL for sorbitol was about the maximum concentrations

that would achieve osmolality and hence were investigated. Sucrose in the range of 60-100 mg/mL was determined to be effective, while sorbitol in the range of 20-60 mg/mL was effective for stability. In contrast, the results shown in Table 33 for the IL12/IL18 DVD-Ig protein did not show an improvement in the stability as measured by DSF.

**Example 16: Impact of pH and Histidine Concentration (at pH 6) on the Shelf Stability of DVD-B and IL12IL18 DVD-Ig Protein at 100mg/ml**

The following examples show the effect of histidine concentration (ranging from 0 – 200 mM) on the shelf stability of the DVD-B and the IL12IL18 DVD-Ig protein as measured using size exclusion chromatography (SEC). Samples were prepared and storage stability of the DVD-Ig proteins in solution at 100 mg/mL was evaluated at 5°C and 40 °C. After defined storage periods, samples were pulled and the impact of storage time on DVD-Ig protein stability was evaluated. Briefly, samples were filled into sterile vials (approx. 500 µL each) and stored under controlled conditions (in temperature chambers and in the absence of light) at 40 °C. At predefined points of time, samples of prepared solutions were pulled for analysis according to the sample pull scheme.

Table 34: Effect of pH on Stability of IL12IL18 DVD-Ig Protein and DVD-B DVD-Ig Protein at 100 mg/mL in Solution

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
N/A	IL12-18	0	0 mM His pH 6+ tween+ sucrose	3.08	95.12	1.8
N/A	IL12-18	0	5 mM His pH 6+ tween+ sucrose	2.8	95.18	2.02
N/A	IL12-18	0	10 mM His pH 6+ tween+ sucrose	2.92	95.06	2.02
N/A	IL12-18	0	50 mM His pH 6+ tween+ sucrose	2.94	94.93	2.13
N/A	IL12-18	0	200 mM His pH 6+ tween+ sucrose	3.11	94.63	2.25
N/A	IL12-18	0	15 mM Ace pH 4.50+ tween+ sucrose	2.88	95.06	2.06
N/A	IL12-18	0	15 mM His pH 6+ tween+ sucrose	2.54	95.4	2.06
N/A	IL12-18	0	15 mM Pho pH 7.40+ tween+ sucrose	3.16	95.53	1.31
N/A	DVDB	0	0 mM His pH 6+ tween+ sucrose	3.96	94.2	1.84
N/A	DVDB	0	5 mM His pH 6+ tween+ sucrose	4.55	93.66	1.79
N/A	DVDB	0	10 mM His pH 6+ tween+ sucrose	4.61	93.64	1.75
N/A	DVDB	0	50 mM His pH 6+ tween+ sucrose	5.52	92.6	1.88
N/A	DVDB	0	200 mM His pH 6+ tween+ sucrose	4.94	93.24	1.83
N/A	DVDB	0	15 mM Ace pH 4.50+ tween+ sucrose	11.19	86.86	1.95
N/A	DVDB	0	15 mM His pH 6+ tween+ sucrose	5.92	92.9	1.18
N/A	DVDB	0	15 mM Pho pH 7.40+ tween+ sucrose	3.26	94.9	1.73
5	IL12-18	7	0 mM His pH 6+ tween+ sucrose	2.82	94.98	2.19
5	IL12-18	7	5 mM His pH 6+ tween+ sucrose	2.53	95.36	2.1

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
5	IL12-18	7	10 mM His pH 6+ tween+ sucrose	2.63	95.9	1.48
5	IL12-18	7	50 mM His pH 6+ tween+ sucrose	2.7	95.57	1.73
5	IL12-18	7	200 mM His pH 6+ tween+ sucrose	3.03	95.08	1.9
5	IL12-18	7	15 mM Ace pH 4.50+ tween+ sucrose	2.23	95.73	2.05
5	IL12-18	7	15 mM His pH 6+ tween+ sucrose	2.38	95.46	2.16
5	IL12-18	7	15 mM Pho pH 7.40+ tween+ sucrose	2.76	94.66	2.56
5	DVDB	7	0 mM His pH 6+ tween+ sucrose	4.76	93.43	1.82
5	DVDB	7	5 mM His pH 6+ tween+ sucrose	5.41	92.75	1.85
5	DVDB	7	10 mM His pH 6+ tween+ sucrose	5.15	93.11	1.74
5	DVDB	7	50 mM His pH 6+ tween+ sucrose	6.44	91.64	1.92
5	DVDB	7	200 mM His pH 6+ tween+ sucrose	5.99	92.27	1.74
5	DVDB	7	15 mM Ace pH 4.50+ tween+ sucrose	14.69	83.4	1.91
5	DVDB	7	15 mM His pH 6+ tween+ sucrose	7.97	90.15	1.87
5	DVDB	7	15 mM Pho pH 7.40+ tween+ sucrose	4.06	94.24	1.69
5	IL12-18	21	0 mM His pH 6+ tween+ sucrose	4.01	91.56	4.43
5	IL12-18	21	5 mM His pH 6+ tween+ sucrose	3.08	92.37	4.55
5	IL12-18	21	10 mM His pH 6+ tween+ sucrose	2.95	92.69	4.35
5	IL12-18	21	50 mM His pH 6+ tween+ sucrose	2.86	92.71	4.43
5	IL12-18	21	200 mM His pH 6+ tween+ sucrose	3.28	92.22	4.5
5	IL12-18	21	15 mM Ace pH 4.50+ tween+ sucrose	2.18	93.18	4.65
5	IL12-18	21	15 mM His pH 6+ tween+ sucrose	2.42	92.36	5.22
5	IL12-18	21	15 mM Pho pH 7.40+ tween+ sucrose	3.95	91.84	4.2
5	DVDB	21	0 mM His pH 6+ tween+ sucrose	5.21	91.17	3.62
5	DVDB	21	5 mM His pH 6+ tween+ sucrose	6.68	89.81	3.51
5	DVDB	21	10 mM His pH 6+ tween+ sucrose	6.26	89.85	3.89
5	DVDB	21	50 mM His pH 6+ tween+ sucrose	8.13	87.88	3.99
5	DVDB	21	200 mM His pH 6+ tween+ sucrose	8.29	87.66	4.06
5	DVDB	21	15 mM Ace pH 4.50+ tween+ sucrose	19.23	77.03	3.73
5	DVDB	21	15 mM His pH 6+ tween+ sucrose	9.54	86.75	3.71
5	DVDB	21	15 mM Pho pH 7.40+ tween+ sucrose	6.25	89.65	4.09
40	IL12-18	7	0 mM His pH 6+ tween+ sucrose	2.56	95.01	2.44
40	IL12-18	7	5 mM His pH 6+ tween+ sucrose	2.06	95.45	2.5
40	IL12-18	7	10 mM His pH 6+ tween+ sucrose	1.93	95.59	2.48
40	IL12-18	7	50 mM His pH 6+ tween+ sucrose	1.82	95.3	2.92
40	IL12-18	7	200 mM His pH 6+ tween+ sucrose	2.18	95.04	2.78
40	IL12-18	7	15 mM Ace pH 4.50+ tween+ sucrose	1.97	95.27	2.76
40	IL12-18	7	15 mM His pH 6+ tween+ sucrose	1.85	95.61	2.54
40	IL12-18	7	15 mM Pho pH 7.40+ tween+ sucrose	4.35	92.57	3.08
40	DVDB	7	0 mM His pH 6+ tween+ sucrose	44.35	54.38	1.28
40	DVDB	7	5 mM His pH 6+ tween+ sucrose	51.66	46.62	1.72
40	DVDB	7	10 mM His pH 6+ tween+ sucrose	50.76	47.45	1.79
40	DVDB	7	50 mM His pH 6+ tween+ sucrose	53.53	44.78	1.69
40	DVDB	7	200 mM His pH 6+ tween+ sucrose	43.21	54.95	1.84
40	DVDB	7	15 mM Ace pH 4.50+ tween+ sucrose	64.17	33.34	2.49
40	DVDB	7	15 mM His pH 6+ tween+ sucrose	50.51	46.87	2.5
40	DVDB	7	15 mM Pho pH 7.40+ tween+ sucrose	36.56	60.79	2.65
40	IL12-18	21	0 mM His pH 6+ tween+ sucrose	4.09	90.71	5.19
40	IL12-18	21	5 mM His pH 6+ tween+ sucrose	3.16	91.26	5.58
40	IL12-18	21	10 mM His pH 6+ tween+ sucrose	3.15	91.78	5.07

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
40	IL12-18	21	50 mM His pH 6+ tween+ sucrose	2.53	91.99	5.48
40	IL12-18	21	200 mM His pH 6+ tween+ sucrose	2.88	91.52	5.61
40	IL12-18	21	15 mM Ace pH 4.50+ tween+ sucrose	2.52	91.59	5.89
40	IL12-18	21	15 mM His pH 6+ tween+ sucrose	2.29	91.97	5.74
40	IL12-18	21	15 mM Pho pH 7.40+ tween+ sucrose	5.35	88.56	6.09
40	DVDB	21	0 mM His pH 6+ tween+ sucrose	55.03	39.46	5.51
40	DVDB	21	5 mM His pH 6+ tween+ sucrose	57.67	37.17	5.16
40	DVDB	21	10 mM His pH 6+ tween+ sucrose	55.56	39.33	5.11
40	DVDB	21	50 mM His pH 6+ tween+ sucrose	58.62	36.27	5.11
40	DVDB	21	200 mM His pH 6+ tween+ sucrose	56.6	37.74	5.66
40	DVDB	21	15 mM Ace pH 4.50+ tween+ sucrose	67.47	26.81	5.71
40	DVDB	21	15 mM His pH 6+ tween+ sucrose	63.14	31.52	5.34
40	DVDB	21	15 mM Pho pH 7.40+ tween+ sucrose	49.52	44.7	5.78

Table 34 also shows the effect of pH range from 4.5-7.4 (pH 4.5 is 15 mM Acetate, pH 6 is 15 mM Histidine and pH 7.4 is 15 mM Phosphate) on the stability of the two DVD-Ig proteins. The data indicate that the stability of IL12IL18 DVD-Ig protein, an AS-DVD-Ig protein, is maintained between the pH of about 4.5 to about 7.4, while the non-AS DVD-Ig protein DVD-B is unstable over pH 4.5 as indicated by the much higher level of aggregate formation over time. The amount of the buffering agent histidine seems to have almost no effect on stability, indicating that the stability issues for non-AS DVD-Ig proteins can be mitigated by formulation parameters, as might have been the case for monoclonal antibodies.

The data indicates that the stability of IL12IL18 DVD-Ig protein is maintained between the pH range of 4.5-7.4, while DVD-B is unstable at either pH 4.5 or pH 7.4 in solution. The stability of the two DVD-Ig proteins show similar stability profiles, however, between the 0-200 mM histidine concentration range at pH 6 given the above conditions in solution.

#### **Example 17: Effect of Citrate Concentration at pH 6 on the Shelf Stability of DVD-B and IL12IL18 DVD-Ig Proteins at 100 mg/ml**

The following example describes the impact of citrate buffer concentration (ranging from 0 – 100 mM) on the shelf stability of the DVD-B and IL12/IL18 DVD-Ig protein. Samples were prepared and storage stability of the DVD-Ig proteins in solution at 100 mg/mL was evaluated at 5°C and 40 °C. After defined storage periods, samples were pulled

and the impact of storage time on DVD-Ig protein stability was evaluated. Briefly, samples were filled into sterile vials (approx. 500  $\mu$ L each) and stored under controlled conditions (in temperature chambers and in the absence of light) at 40 °C. At predefined points of time, samples of prepared solutions were pulled for analysis according to the sample pull scheme.

Table 35: Effect of pH on Stability of DVD-B and IL12IL18 DVD-Ig Proteins

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
	IL12/18	0	0 mM Cit pH 6	3.2	94.98	1.82
	IL12/18	0	5 mM Cit pH 6	3.3	94.68	2.2
	IL12/18	0	10 mM Cit pH 6	3.06	94.76	2.18
	IL12/18	0	50 mM Cit pH 6	2.52	95.28	2.19
	IL12/18	0	100 mM Cit pH 6	2.82	95.05	2.13
	IL12/18	0	0 mM Cit pH 6+ Sucrose+tween	2.73	95.11	2.16
	IL12/18	0	5 mM Cit pH 6+ Sucrose+tween	2.47	95.55	1.97
	IL12/18	0	10 mM Cit pH 6+ Sucrose+tween	2.67	95.1	2.23
	IL12/18	0	50 mM Cit pH 6+ Sucrose+tween	2.57	95.37	2.05
	IL12/18	0	100 mM Cit pH 6+ Sucrose+tween	2.55	95.24	2.21
5	IL12/18	7	0 mM Cit pH 6	2.91	94.51	2.58
5	IL12/18	7	5 mM Cit pH 6	3.16	94.76	2.08
5	IL12/18	7	10 mM Cit pH 6	2.77	94.77	2.46
5	IL12/18	7	50 mM Cit pH 6	3.16	94.67	2.17
5	IL12/18	7	100 mM Cit pH 6	2.84	95.81	1.36
5	IL12/18	7	0 mM Cit pH 6+ Sucrose+tween	2.58	95.21	2.21
5	IL12/18	7	5 mM Cit pH 6+ Sucrose+tween	2.98	94.84	2.18
5	IL12/18	7	10 mM Cit pH 6+ Sucrose+tween	2.78	94.82	2.4
5	IL12/18	7	50 mM Cit pH 6+ Sucrose+tween	2.87	94.79	2.35
5	IL12/18	7	100 mM Cit pH 6+ Sucrose+tween	2.77	94.97	2.25
5	IL12/18	21	0 mM Cit pH 6	3.22	94.21	2.57
5	IL12/18	21	5 mM Cit pH 6	3.64	95.31	1.05
5	IL12/18	21	10 mM Cit pH 6	3.4	93.9	2.7
5	IL12/18	21	50 mM Cit pH 6	3.66	94.53	1.81
5	IL12/18	21	100 mM Cit pH 6	3.3	94.47	2.23
5	IL12/18	21	0 mM Cit pH 6+ Sucrose+tween	2.93	94.44	2.68
5	IL12/18	21	5 mM Cit pH 6+ Sucrose+tween	3.19	94.71	2.1
5	IL12/18	21	10 mM Cit pH 6+ Sucrose+tween	3.2	94.54	2.26
5	IL12/18	21	50 mM Cit pH 6+ Sucrose+tween	3.12	94.51	2.37
5	IL12/18	21	100 mM Cit pH 6+ Sucrose+tween	2.88	94.94	2.19
40	IL12/18	7	0 mM Cit pH 6	3.06	94.81	2.12
40	IL12/18	7	5 mM Cit pH 6	3.4	94.33	2.27
40	IL12/18	7	10 mM Cit pH 6	3.28	94.47	2.24
40	IL12/18	7	50 mM Cit pH 6	3.81	94.13	2.06
40	IL12/18	7	100 mM Cit pH 6	3.55	94.07	2.38
40	IL12/18	7	0 mM Cit pH 6+ Sucrose+tween	2.7	96.87	0.44
40	IL12/18	7	5 mM Cit pH 6+ Sucrose+tween	2.84	94.61	2.55
40	IL12/18	7	10 mM Cit pH 6+ Sucrose+tween	2.98	94.61	2.4
40	IL12/18	7	50 mM Cit pH 6+ Sucrose+tween	2.88	94.84	2.27
40	IL12/18	7	100 mM Cit pH 6+ Sucrose+tween	2.96	94.64	2.4

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
40	IL12/18	21	0 mM Cit pH 6	4.48	91.27	4.26
40	IL12/18	21	5 mM Cit pH 6	4.92	90.99	4.1
40	IL12/18	21	10 mM Cit pH 6	4.62	91.2	4.18
40	IL12/18	21	50 mM Cit pH 6	4.69	91.53	3.78
40	IL12/18	21	100 mM Cit pH 6	4.24	92.23	3.52
40	IL12/18	21	0 mM Cit pH 6+ Sucrose+tween	3.92	91.87	4.21
40	IL12/18	21	5 mM Cit pH 6+ Sucrose+tween	3.92	92.05	4.03
40	IL12/18	21	10 mM Cit pH 6+ Sucrose+tween	4.01	91.63	4.36
40	IL12/18	21	50 mM Cit pH 6+ Sucrose+tween	3.66	92.52	3.81
40	IL12/18	21	100 mM Cit pH 6+ Sucrose+tween	3.78	91.89	4.33
5		3 Month	0 mM Cit pH 6	2.53	94.36	3.11
5		3 Month	5 mM Cit pH 6	2.89	94.33	2.77
5	IL12/18	3 Month	10 mM Cit pH 6	3	94.35	2.64
5	IL12/18	3 Month	50 mM Cit pH 6	3.78	93.83	2.39
5	IL12/18	3 Month	100 mM Cit pH 6	3.58	94.45	1.98
5	IL12/18	3 Month	0 mM Cit pH 6+ Sucrose+tween	2.94	94.71	2.35
5	IL12/18	3 Month	5 mM Cit pH 6+ Sucrose+tween	3.58	94.29	2.13
5	IL12/18	3 Month	10 mM Cit pH 6+ Sucrose+tween	2.95	95.16	1.89
5	IL12/18	3 Month	50 mM Cit pH 6+ Sucrose+tween	3.44	94.58	1.98
5	IL12/18	3 Month	100 mM Cit pH 6+ Sucrose+tween	3.24	94.54	2.21
5	IL12/18	6 Month	0 mM Cit pH 6	6.4	90.99	2.61
5	IL12/18	6 Month	5 mM Cit pH 6	6.18	91.58	2.24
5	IL12/18	6 Month	10 mM Cit pH 6	6.04	92.12	1.84
5	IL12/18	6 Month	50 mM Cit pH 6	5.72	92.9	1.38
5	IL12/18	6 Month	100 mM Cit pH 6	5.98	92.86	1.16
5	IL12/18	6 Month	0 mM Cit pH 6+ Sucrose+tween	4.98	94.02	1
5	IL12/18	6 Month	5 mM Cit pH 6+ Sucrose+tween	5.47	92.96	1.5
5	IL12/18	6 Month	10 mM Cit pH 6+ Sucrose+tween	4.51	94.56	0.9
5	IL12/18	6 Month	50 mM Cit pH 6+ Sucrose+tween	4.58	93.44	1.9
5	IL12/18	6 Month	100 mM Cit pH 6+ Sucrose+tween	4.63	93.54	1.82
5	IL12/18	6 Month	0 mM Cit pH 6	6.4	90.99	2.61
	DVD-B	0	0 mM Cit pH 6	2.6	95.63	1.77
		0	5 mM Cit pH 6	2.6	95.49	1.87
	DVD-B	0	10 mM Cit pH 6	2.41	95.88	1.72
	DVD-B	0	50 mM Cit pH 6	1.76	96.62	1.62
	DVD-B	0	100 mM Cit pH 6	1.34	97	1.66
	DVD-B	0	0 mM Cit pH 6+ Sucrose+tween	1.54	97.31	1.15
	DVD-B	0	5 mM Cit pH 6+ Sucrose+tween	1.52	96.74	1.73
	DVD-B	0	10 mM Cit pH 6+ Sucrose+tween	1.5	96.8	1.71
	DVD-B	0	50 mM Cit pH 6+ Sucrose+tween	1.5	96.74	1.75
	DVD-B	0	100 mM Cit pH 6+ Sucrose+tween	1.11	97.16	1.73
5	DVD-B	7	0 mM Cit pH 6	2.61	95.54	1.85
5	DVD-B	7	5 mM Cit pH 6	3.01	95.37	1.62
5	DVD-B	7	10 mM Cit pH 6	3.2	95.12	1.68
5	DVD-B	7	50 mM Cit pH 6	2.85	95.5	1.64
5	DVD-B	7	100 mM Cit pH 6	2.14	96.22	1.64

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
5	DVD-B	7	0 mM Cit pH 6+ Sucrose+tween	2.32	95.95	1.73
5	DVD-B	7	5 mM Cit pH 6+ Sucrose+tween	2.86	95.47	1.66
5	DVD-B	7	10 mM Cit pH 6+ Sucrose+tween	2.71	95.61	1.68
5	DVD-B	7	50 mM Cit pH 6+ Sucrose+tween	2.75	95.62	1.64
5	DVD-B	7	100 mM Cit pH 6+ Sucrose+tween	1.97	96.35	1.68
5	DVD-B	21	0 mM Cit pH 6	3.33	94.44	2.23
5	DVD-B	21	5 mM Cit pH 6	4.97	92.72	2.31
5	DVD-B	21	10 mM Cit pH 6	5.52	92.56	1.92
5	DVD-B	21	50 mM Cit pH 6	4.88	93.82	1.3
5	DVD-B	21	100 mM Cit pH 6	3.7	94.76	1.54
5	DVD-B	21	0 mM Cit pH 6+ Sucrose+tween	3	95.28	1.72
5	DVD-B	21	5 mM Cit pH 6+ Sucrose+tween	4.88	93.06	2.06
5	DVD-B	21	10 mM Cit pH 6+ Sucrose+tween	5	93.41	1.59
5	DVD-B	21	50 mM Cit pH 6+ Sucrose+tween	5.01	93.14	1.84
5	DVD-B	21	100 mM Cit pH 6+ Sucrose+tween	3.77	94.38	1.85
40	DVD-B	7	0 mM Cit pH 6	51.75	46.63	1.62
40	DVD-B	7	5 mM Cit pH 6	58.47	39.22	2.31
40	DVD-B	7	10 mM Cit pH 6	58.39	38.8	2.8
40	DVD-B	7	50 mM Cit pH 6	47.99	49.69	2.32
40	DVD-B	7	100 mM Cit pH 6	39.78	58.14	2.09
40	DVD-B	7	0 mM Cit pH 6+ Sucrose+tween	47.61	49.83	2.56
40	DVD-B	7	5 mM Cit pH 6+ Sucrose+tween	56.88	40.48	2.64
40	DVD-B	7	10 mM Cit pH 6+ Sucrose+tween	46.46	50.97	2.57
40	DVD-B	7	50 mM Cit pH 6+ Sucrose+tween	45.57	52.03	2.4
40	DVD-B	7	100 mM Cit pH 6+ Sucrose+tween	34.2	63.53	2.27
40	DVD-B	21	0 mM Cit pH 6	60.28	36.18	3.54
40	DVD-B	21	5 mM Cit pH 6	n/a	n/a	n/a
40	DVD-B	21	10 mM Cit pH 6	n/a	n/a	n/a
40	DVD-B	21	50 mM Cit pH 6	n/a	n/a	n/a
40	DVD-B	21	100 mM Cit pH 6	n/a	n/a	n/a
40	DVD-B	21	0 mM Cit pH 6+ Sucrose+tween	56.73	39.59	3.67
40	DVD-B	21	5 mM Cit pH 6+ Sucrose+tween	59.87	36.82	3.3
40	DVD-B	21	10 mM Cit pH 6+ Sucrose+tween	53.36	43.59	3.05
40	DVD-B	21	50 mM Cit pH 6+ Sucrose+tween	51.27	45.72	3.01
40	DVD-B	21	100 mM Cit pH 6+ Sucrose+tween	44.11	52.84	3.05
5	DVD-B	3 Month	0 mM Cit pH 6	7.25	90.86	1.9
5	DVD-B	3 Month	5 mM Cit pH 6	13.89	84.29	1.82
5	DVD-B	3 Month	10 mM Cit pH 6	18.99	79.33	1.68
5	DVD-B	3 Month	50 mM Cit pH 6	16.07	82.2	1.73
5	DVD-B	3 Month	100 mM Cit pH 6	12.87	85.6	1.52
5	DVD-B	3 Month	0 mM Cit pH 6+ Sucrose+tween	7.43	90.74	1.83
5	DVD-B	3 Month	5 mM Cit pH 6+ Sucrose+tween	15.08	83.23	1.69
5	DVD-B	3 Month	10 mM Cit pH 6+ Sucrose+tween	16.03	82.02	1.95
5	DVD-B	3 Month	50 mM Cit pH 6+ Sucrose+tween	17.09	81.17	1.74
5	DVD-B	3 Month	100 mM Cit pH 6+ Sucrose+tween	10.89	87.64	1.46
5	DVD-B	6 Month	0 mM Cit pH 6	15.5	82.31	2.2
5	DVD-B	6 Month	5 mM Cit pH 6	27.45	71.13	1.43
5	DVD-B	6 Month	10 mM Cit pH 6	34.37	63.99	1.63
5	DVD-B	6 Month	50 mM Cit pH 6	29.11	69.6	1.29

°C	Sample	Time (D)	Formulation	Agg	Mon	Frag
5	DVD-B	6 Month	100 mM Cit pH 6			
5	DVD-B	6 Month	0 mM Cit pH 6+ Sucrose+tween	15.24	83.22	1.54
5	DVD-B	6 Month	5 mM Cit pH 6+ Sucrose+tween	27.28	71.19	1.53
5	DVD-B	6 Month	10 mM Cit pH 6+ Sucrose+tween	27.69	70.8	1.51
5	DVD-B	6 Month	50 mM Cit pH 6+ Sucrose+tween	27.18	70.78	2.04
5	DVD-B	6 Month	100 mM Cit pH 6+ Sucrose+tween	21.8	76.46	1.74

The two DVD-Ig proteins in Table 35 showed different stability profiles between the 0-100 mM citrate concentrations at pH 6. While AS-DVD-Ig protein IL12IL18 showed a high stability with only a minor increase in aggregates overtime, especially at 5 °C, the opposite was true for the non AS-DVD-Ig protein DVD-B. This suggests that a high concentration liquid formulation would be feasible for the IL12IL18 DVD-Ig protein but not for DVD-B under those conditions.

## VIII. STABLE LYOPHILIZED DVD-IG (LS-DVD-IG) PROTEIN FORMULATIONS

Examples 18 and 19 describe the stability of LS-DVD-Ig proteins in the lyophilized form. Example 18 describes surprising results that demonstrate freeze/ thaw (F/T) stability of LS-DVD-Ig proteins. Example 19 describes studies showing stable lyophilized formulations containing LS-DVD-Ig proteins. Freezing is the first step in lyophilization and hence molecules that do not have freeze thaw stability are susceptible to instability during lyophilization.

### EXAMPLE 18: Impact of Solution pH on the Stability of DVD-Ig Proteins Subjected to Repeated Freeze/Thaw Cycles

The freeze thaw behavior of DVD-Ig proteins at a protein concentration of 1 mg/ml in 5 mM citrate/5mM phosphate buffer was evaluated by cycling the protein solution up to 2 times between the frozen state and the liquid state at pH 4, pH 6, and pH 8. Freezing was performed using temperature controlled -80°C freezer, and thawing was performed using a 30°C temperature controlled water bath. Samples were pulled after the second freeze/thaw (F/T) cycle and analyzed by SEC. Table 36 shows the effect of freeze/thaw processing on the amount of monomer (Mon) of remaining and the amount of fragments (Frag) and aggregates (Agg) formed in the samples formulated at these pH levels.

**Table 36: Numbers Of DVD-Ig Protein Monomers, Aggregates, And Fragments as Determined by SEC Before and After Repeated Freeze/Thaw Cycles of DVD-Ig Protein Formulations with Solution pH Values of 4, 6, or 8**

DVD-Ig	pH	Mon/T0	Agg/T0	Frag/T0	Mon/T2	Agg/T2	Frag/T2
5	4	95.06	0.44	4.49	93.57	1.52	4.89
6	4	95.26	1.21	3.52	94.94	1.72	3.33
37	4	97.32	1.89	0.78	96.64	2.48	0.86
38	4	94.46	3.81	1.72	94.91	3.55	1.52
53	4	97.47	1	1.52	97.41	1.06	1.52
54	4	96.06	1.87	2.06	95.87	1.85	2.26
65	4	94.44	1.2	4.35	93.82	1.15	5.01
66	4	98.01	0.91	1.07	97.78	1.03	1.18
165	4	96.62	2.17	1.2	95.45	2.34	2.19
166	4	97.74	0.91	1.34	97.53	0.91	1.54
257	4	97.63	1.8	0.56	96.73	2.73	0.53
258	4	98.84	0.19	0.95	96.73	2.28	0.97
277	4	95.53	1.43	3.03	95.45	1.6	2.93
278	4	95.75	1.56	2.68	95.4	1.99	2.59
281	4	98.72	0.63	0.63	97.47	1.83	0.68
282	4	98.63	0.61	0.74	97.88	1.38	0.72
5	6	94.8	3.13	2.06	94.96	3.37	1.67
6	6	95.79	1.99	2.2	93.71	4.02	2.26
37	6	96.73	2.52	0.74	94.88	4.42	0.68
38	6	95.01	3.74	1.23	95.5	3.29	1.2
53	6	97.56	1	1.43	97.51	1.08	1.4
54	6	95.86	2.07	2.06	95.68	2.18	2.13
65	6	94.17	1.09	4.72	94.13	1.16	4.7
66	6	97.99	0.85	1.15	98.03	0.83	1.13
165	6	96.86	2.12	1.01	96.03	2.38	1.57
166	6	97.75	0.91	1.33	97.82	0.9	1.26
257	6	97.51	1.97	0.51	97.31	2.19	0.49
258	6	99.07	0.18	0.73	98.52	0.76	0.7
277	6	96.74	1.79	1.46	96.64	1.92	1.43
278	6	97.65	1.22	1.12	97.62	1.32	1.04
281	6	95.67	0.92	3.4	95.7	0.98	3.3
282	6	98.55	0.83	0.61	98.49	0.89	0.6
5	8	93.23	3.86	2.89	93.7	3.56	2.73
6	8	95.35	1.93	2.7	94.3	2.94	2.74
37	8	94.88	4.27	0.84	95.64	3.54	0.8
38	8	95.52	3.14	1.32	96.09	2.61	1.28
53	8	97.56	1.04	1.39	97.66	1.02	1.31
54	8	95.77	1.91	2.3	95.91	1.9	2.17
65	8	94.1	1.14	4.74	94.29	1.14	4.56
66	8	97.84	0.95	1.2	97.74	0.95	1.3
165	8	96.63	2.23	1.12	96.07	2.32	1.59
166	8	97.55	0.9	1.54	97.77	0.85	1.37
257	8	97.12	2.19	0.67	96.62	2.7	0.67

DVD-Ig	pH	Mon/T0	Agg/T0	Frag/T0	Mon/T2	Agg/T2	Frag/T2
258	8	98.81	0.28	0.9	98.71	0.39	0.89
277	8	95.48	2.23	2.27	95.52	2.24	2.22
278	8	95.94	1.83	2.21	96.08	1.67	2.23
281	8	95.61	1	3.38	95.94	0.98	3.06
282	8	98.28	1.04	0.67	98.26	1.05	0.67

DVD 5, DVD 6, DVD 37, DVD 38, DVD 53, DVD 54, DVD 65, DVD 66, DVD 165, DVD 166, DVD 257, DVD 258, DVD 277, DVD 278, DVD 281, and DVD 282 demonstrated stability after being subjected to repeated freeze thaw cycles. These data indicate that DVD-Ig proteins that are formulated in a pH range of about 4 to about 8 remain stable after repeated F/T processing. The high stability of the DVD-Ig protein formulations tested (all showed greater than 93% monomer content and 11/16 formulations showed greater than 95% monomer content) was unexpected, because DVD-Ig proteins are much more complex than IgGs. Complex molecules such as DVD-Ig proteins would be expected to aggregate and fragment easily when exposed to freezing and thawing.

*Impact of Solution pH On the Stability of DVD-B Subjected to Repeated Freeze/Thaw Cycles*

The freeze thaw behavior of DVD-B at a protein concentration of 2 mg/ml in 10 mM citrate/10mM phosphate buffer was evaluated by cycling the protein solution up to 4 times between the frozen state and the liquid state at pH 4-9. Freezing was performed by means of a temperature controlled -80°C freezer, and thawing was performed by means of a 30°C temperature controlled water bath. Samples were pulled after each freeze/thaw (F/T) cycle and analyzed by light obscuration and SEC. Table 37 shows the effect of freeze/thaw processing on the number of sub-visible particles formed as determined using light obscuration measurements at various pH values.

Table 37: Numbers of Subvisible Particles Per mL as Determined by Light Obscuration Assays After 0-4 Freeze/Thaw (F/T) Cycles of DVD-B Formulations with a pH Value of 4, 5, 6, 7, 8, Or 9

(A) Numbers of particles $\geq$ 1 micron in size.						
Number of F/T cycles	pH = 4	pH = 5	pH = 6	pH = 7	pH = 8	pH = 9
0	39.16	55	44	41	55	46
1	334.5	1291	38127	31896	28444	25970

2	6728.6	7935	80064	61592	58562	46863
3	13658	18733	128448	95775	89934	67225
4	5702	37930	175024	132768	120339	89389

(B) Numbers of particles  $\geq$  10 microns in size.

Number of F/T cycles	pH = 4	pH = 5	pH = 6	pH = 7	pH = 8	pH = 9
0	1.33	1.5	2.6	2.6	5.5	1.83
1	15.16	3.16	28	45	27.16	16.83
2	207.5	62	142	165	148	134
3	342	136	440	740	773	375
4	572	269	2418	4099	4537	1425

(C) Numbers of particles  $\geq$  25 microns in size.

Number of F/T cycles	pH = 4	pH = 5	pH = 6	pH = 7	pH = 8	pH = 9
0	0.33	0.33	0.16	0.5	1.16	0.16
1	3.3	0.5	3.83	2	4.33	1.33
2	40.16	10.83	2.33	6.33	1.16	3.16
3	71.16	6.67	13.5	39.83	39.33	5.33
4	118.83	67.63	198.67	476.16	641.5	142.83

The results of the light obscuration assays show that the numbers of particles formed by DVD-B formulations with a pH of 4 to 9 was low. The numbers of particles formed increased with increasing pH and were at a maximum at around the pI of the molecule (pI 8.5). However, with only one exception, the protein solutions tested satisfied the requirements of the International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH) guidelines, which requires less than 600 particles of size 25 microns or higher per ml.

Table 38 shows SEC measurements of the stability of DVD-B after freeze/thaw processing. These measurements include the percentage of monomers, aggregates, and fragments, as well as the area under the curve (AUC).

Table 38: Stability Of DVD-B as Determined by SEC After 0-4 Freeze/Thaw Cycles of DVD-B Formulations with Solution pH Values of 4, 5, 6, 7, 8, Or 9

(A) SEC measurements of solutions not subjected to F/T cycles				
	Mon	Agg	Frag	AUC
pH 4 Vial 1	96.38	1.44	2.16	76861
pH 4 Vial 2	95.61	1.56	2.82	76123
pH 4 Mean	95.995	1.5	2.49	76492
pH 5 Vial 1	96.1	1.57	2.32	73704

pH 5 Vial 2	96.16	1.55	2.28	74196
pH 5 Mean	96.13	1.56	2.3	73950
pH 6 Vial 2	95.09	1.69	3.2	77475
pH 7 Vial 2	95.61	1.81	2.56	75863
pH 8 Vial 2	95.84	1.87	2.27	74943
pH 9 Vial 1	95.66	2.11	2.21	82103
pH 9 Vial 2	95.43	2.14	2.41	81843
pH 9 Mean	95.545	2.125	2.31	81973

## (B) SEC measurements after one F/T cycle

pH 4 Vial 1	96.37	1.68	1.94	74694
pH 4 Vial 2	95.39	1.74	2.86	76213
pH 4 Mean	95.88	1.71	2.4	75453.5
pH 5 Vial 1	96.18	1.82	1.98	71511
pH 5 Vial 2	96.08	1.6	2.31	76345
pH 5 Mean	96.13	1.71	2.145	73928
pH 6 Vial 1	96.37	1.6	2.02	75448
pH 6 Vial 2	95.24	1.6	3.14	77009
pH 6 Mean	95.805	1.6	2.58	76228.5
pH 7 Vial 1	95.88	1.86	2.24	75821
pH 7 Vial 2	95.28	1.96	2.74	76470
pH 7 Mean	95.58	1.91	2.49	76145.5
pH 8 Vial 1	95.64	2.01	2.34	74757
pH 8 Vial 2	95.59	2.08	2.31	74270
pH 8 Mean	95.81536	1.777143	2.3925	75306.68
pH 9 Vial 1	95.62	2.15	2.22	81981
pH 9 Vial 2	95.57	2.13	2.29	81516
pH 9 Mean	95.595	2.14	2.255	81748.5

## (C) SEC measurements after two F/T cycles

pH 4 Vial 1	95.75	1.98	2.25	75402
pH 4 Vial 2	95.29	1.98	2.71	74567
pH 4 Mean	95.52	1.98	2.48	74984.5
pH 5 Vial 1	96.07	1.85	2.06	71454
pH 5 Vial 2	95.92	1.88	2.19	71525
pH 5 Mean	95.995	1.865	2.125	71489.5
pH 6 Vial 1	96.33	1.58	2.08	74666
pH 6 Vial 2	95.43	1.62	2.93	75080
pH 6 Mean	95.88	1.6	2.505	74873
pH 7 Vial 1	95.77	1.87	2.34	74813
pH 7 Vial 2	95.72	1.84	2.42	74584
pH 7 Mean	95.745	1.855	2.38	74698.5
pH 8 Vial 1	95.72	2.02	2.24	73460
pH 8 Vial 2	95.78	2	2.21	68769
pH 8 Mean	95.75	2.01	2.225	71114.5
pH 9 Vial 1	95.58	2.1	2.3	73356
pH 9 Vial 2	95.57	2.12	2.3	77337
pH 9 Mean	95.575	2.11	2.3	75346.5

## (D) SEC measurements after three F/T cycles

pH 4 Vial 1	95.62	2.14	2.22	72539
-------------	-------	------	------	-------

pH 4 Vial 2	95.01	2.23	2.75	72480
pH 4 Mean	95.315	2.185	2.485	72509.5
pH 5 Vial 1	95.9	1.99	2.09	68689
pH 5 Vial 2	95.8	1.98	2.21	65381
pH 5 Mean	95.85	1.985	2.15	67035
pH 6 Vial 2	95.55	1.63	2.81	71195
pH 7 Vial 1	96.03	1.84	2.12	70617
pH 7 Vial 2	95.77	1.86	2.35	71405
pH 7 Mean	95.9	1.85	2.235	71011
pH 8 Vial 1	95.86	2.04	2.08	69364
pH 8 Vial 2	95.59	2.12	2.28	69308
pH 8 Mean	95.725	2.08	2.18	69336
pH 9 Vial 1	95.03	2.42	2.54	77679
pH 9 Vial 2	95.59	2.22	2.18	76113
pH 9 Mean	95.31	2.32	2.36	76896
(E) SEC measurements after four F/T cycles				
pH 4 Vial 1	94.38	2.53	3.08	53902
pH 4 Vial 2	94.08	2.4	3.5	72943
pH 4 Mean	94.23	2.465	3.29	63422.5
pH 5 Vial 1	96.01	1.88	2.1	66962
pH 5 Vial 2	95.85	1.96	2.18	67786
pH 5 Mean	95.93	1.92	2.14	67374
pH 6 Vial 1	96.26	1.62	2.11	70371
pH 6 Vial 2	95.41	1.67	2.9	71086
pH 6 Mean	95.835	1.645	2.505	70728.5
pH 7 Vial 1	95.87	1.82	2.29	67317
pH 7 Vial 2	96.03	1.81	2.15	67869
pH 7 Mean	95.95	1.815	2.22	67593
pH 8 Vial 1	95.62	2.07	2.3	69629
pH 8 Vial 2	95.31	2.15	2.52	69401
pH 8 Mean	95.465	2.11	2.41	69515
pH 9 Vial 1	95.44	2.27	2.28	77495
pH 9 Vial 2	95.58	2.18	2.22	73373
pH 9 Mean	95.51	2.225	2.25	75434

The results in Table 38 indicate that the DVD-Ig protein does not form significant amounts of aggregates even after 4 F/T cycles. The good F/T stability is surprising as it was anticipated that due to their higher structural complexity DVD-Ig proteins would be more prone to undergo degradation at the liquid / ice interface or even cold denaturation upon multiple freeze /thaw cycles. This is a common observation for proteins and many other DVD-Ig proteins show a significant amount of aggregation upon freeze/thawing. Since freezing is the initial, and a critical, step of the lyophilizing process, the data suggests that DVD-B can be lyophilized and hence converted into a stable biotherapeutic dosage form.

The results in Table 37 indicate that the DVD-Ig protein does not form significant

aggregates even after 4 F/T cycles. The good F/T stability was surprising as it was anticipated that the stability might not be as good as was observed given the intrinsic lack of stability and increased structural complexity relative to an antibody.

#### **EXAMPLE 19: Impact of Lyophilization on the Stability of DVD-Ig Proteins**

Aggregates may form during the process of lyophilization as well as later on during shelf stability of the solid protein. The aggregates formed during lyophilization are generally measured following immediate reconstitution.

Storage stability of DVD-Ig proteins was evaluated for prolonged periods of time at controlled temperature conditions. After defined storage periods, samples were pulled and the impact of storage time and storage temperature on the stability of lyophilized DVD-Ig proteins was evaluated by size exclusion chromatography (SEC) and ion exchange chromatography (IEC). Three DVD-Ig proteins were studied: DVD-B (TNF-PGE2), DVD-A (TNF-IL17), and DVD-C (IL1 $\alpha$ -IL1 $\beta$ ). Of the three, DVD-A and DVD-C are AS-DVD-Ig proteins. DVD-B is a non-AS-DVD-Ig protein, but was identified as an LS-DVD-Ig protein as it (as well as LS-DVD-Ig proteins DVD-A and DVD-C) was found to be stable in a lyophilized formulation. The formulations were lyophilized in solutions as shown in Table 39 in a formulation containing a buffer, a polyol, and a surfactant.

Table 39: Composition of Lyophilized Formulations

Component	DVD-A <b>55 mg/ml</b>	DVD-B <b>55 mg/ml</b>	DVD-C <b>48 mg/ml</b>
Histidine	2.33 (15mM)	2.33 (15mM)	2.33 (15mM)
Sucrose	46 (4.6%)	46 (4.6%)	46 (4.6%)
Polysorbate 80	0.2 (0.02%)	0.2 (0.02%)	0.1 (0.01%)
0.1 M HCl	q.s. (pH 5.25)	q.s. (pH 5.25)	q.s. (pH 6.0)

Table 40 describes the percentages of monomers, aggregates, and fragments that were measured using SEC before storage (T0) or following storage of the lyophilized formulations at either 5°C or 40°C for the given storage periods.

Table 40: Stability of Stored Lyophilized Formulations as Assessed Using SEC

DVD-Ig	Storage Condition	Monomer	Aggregate	Fragment
DVD-A	T0	94.22	5.09	0.67
	1 Week 40°C	94.12	5.23	0.64
	4 Week 40°C	94.41	5.14	0.43
	3m 40°C	93.31	6.24	0.43
	3m 40°C + 3h RT	93.17	6.38	0.43
DVD-B	T0	96.59	2.11	1.28
	1 Week 40°C	96.11	2.59	1.28
	4 Week 40°C	95.44	3.19	1.36
	4 Week 5°C	96.47	2.2	1.31
	4 Week 40°C + 3h RT	95.35	3.29	1.34
DVD-C	T0	96.55	2.94	0.5
	1 Week 40°C	97.165	1.59	1.23
	1m 40°C	96.88	1.93	1.18
	3m 5°C	97.42	1.36	1.21

The SEC data in Table 40 shows that lyophilized LS-DVD-Ig proteins remain stable for periods of up to 3 months of storage because they show high percentages of monomers and low percentages of aggregates and fragments. After accelerated storage for 3 months at 40°C and 3 hours of reconstitution time, lyophilized DVD-A had more than 93% monomers, less than 6.4% aggregates, and only about 0.4% fragments. After 4 weeks of accelerated storage at 40°C, lyophilized DVD-B had more than 95% monomers and only about 3.2% aggregates and about 1.4% fragments. After 1 month of accelerated storage at 40°C, lyophilized DVD-C had approximately 97% monomers, 2% aggregates, and 1% fragments. Thus, lyophilized formulations of DVD-A, DVD-B, and DVD-C showed long term stability, as assessed by SEC.

Table 41 below provides data regarding the stability of stored formulations measured using IEC. The impact of chemical stability was not significant as observed from formation of acidic and basic species with time. Therefore, all tested molecules do not show significant chemical degradation in the lyophilized state. Storage stability of lyophilized DVD-Ig proteins was evaluated at 40°C and 5°C. After defined storage periods, the lyophilized samples were pulled and the impact of storage time on DVD-Ig protein stability was evaluated. Samples were reconstituted with water for injection and upon complete reconstitution analyzed using SEC and IEC.

Table 41: Stability of Stored Lyophilized Formulations as Assessed Using IEC

DVD-Ig	Storage condition	Main	Acidic	Basic	X
DVD-A	T0	73.28	3.6	20.78	2.32
	1 Week 40°C	72.59	3.38	21.91	2.1
	4 Week 40°C	70.73	3.6	23.44	2.21
	3m 40°C	71.75	3.84	22	2.39
	3m 40°C + 3h RT	71.8	3.71	22.03	2.44
DVD-B	T0	48.84	38.54	12.6	
	1 Week 40°C	48.18	38.72	13.09	
	4 Week 40°C	45.87	37.6	16.53	
	4 Week 5°C	48.17	38.16	13.56	
	4 Week 40°C + 3h RT	45.7	37.79	16.5	
DVD-C	1 Week 40°C	54.82	28.82	16.35	
	1m 40°C	53.61	28.84	17.53	
	3m 5°C	55.43	28.57	15.99	

The impact of chemical stability seen above in Table 40 was not significant as observed from formation of acidic and basic species with time.

Table 42 below provides the reconstitution times of the stored lyophilized formulations. One basic criterion for characterizing the feasibility of a lyophilized dosage form is the time it takes for a lyophilized sample to be converted in a homogenous, clear liquid after reconstitution with a solvent, usually water for injection. Ideally, the reconstitution procedure takes less than 10 minutes. As shown in Table 41 all tested samples showed complete reconstitution after a maximum of 5 minutes. This indicates that AS-DVD-Ig proteins (e.g., DVD-Ig proteins A & C) can be formulated as stable lyophilized formulations. Moreover, the above examples show that LS-DVD-Ig proteins - which are not stable in liquid formulations - can be stabilized using lyophilization (e.g., DVD-B).

Table 42: Reconstitution Time (RT) Of the Lyophilized Formulations (With Water for Injection)

DVD-Ig	Storage condition	RT
DVD-A	T0	30s
	T1 week 40°C	26s
	T4 week 40°C	48s
	T3 month 40°C	300s
DVD-B	T0	48s
	T1 week 40°C	60s
	T4 week 5°C	69s
	T4 week 40°C	89s

DVD-Ig	Storage condition	RT
DVD-C	T0	47s
	T1 week 40°C	76s
	3m 5°C	62s
	1m 40°C	53s

## IX. CHARACTERIZATION OF DVD-IG PROTEINS

Examples 20-23 provide further characterization of DVD-Ig proteins generally.

### Example 20: Solubility Assessment of DVD-Ig Proteins Using PEG 3000

Polyethylene glycols (PEG) are often used as “crowders,” which force the protein in a smaller amount of water, because some of the available solvent is bound to or occupied by the crowder, typically a polymer. PEGs are used to assess the true solubility of a protein by utilizing micro amounts of the protein material available at early stages of development. In general, the greater the amount of PEG is required to induce precipitation, the greater is the anticipated true solubility of the protein in solution.

The following studies were carried out using small aliquots of PEG solution (50% w/v) added to a stock solution of protein (0.5 mg/ml) in a buffer (5 mM citrate and 5 mM phosphate) at pH 6. Table 43 shows the data for various DVD-Ig proteins.

Table 43. Amount of PEG 3000 Required To Induce Precipitation in a 0.5 mg/ml Protein Solution of DVD-Ig Proteins

DVD-Ig	% PEG 3000 required to induce precipitation
5	10
6	9.37
37	10
38	9.33
53	8.12
54	9.37
65	7.5
66	7.5
165	10.625
166	11.875
257	9.37
258	10.62
277	12.5
278	13.75

281	8.12
282	7.5

These data show that the amount of PEG 3000 required to induce precipitation of DVD-Ig proteins is typical of highly soluble proteins (*i.e.*, those proteins with a true solubility that exceeds about 100 mg/ml). Although the PEG precipitation assay is a standard assay in antibody formulation assessment to provide information about its solubility, it would not be sufficient to predict whether a DVD-Ig protein would be classified as AS or LS or even non LS, indicating the complexity of DVD-Ig proteins and the challenging formulation efforts compared to monoclonal antibodies.

**Example 21: Assessment of the Tertiary Structure of DVD-Ig Proteins Using Near UV CD Scans**

The structure of a protein is one of the important factors influencing protein stability during accelerated/long-term storage of protein liquid and lyophilizate formulations. To assess the tertiary structure of the DVD-Ig proteins, near UV CD scan between 250-320 nm was taken on a Jasco Spectrometer with a scan rate of 50 nm/minute at a concentration of 1mg/ml. An average of 3 scans/ DVD-Ig protein was taken. The pH used in the study was 6 in 5 mM citrate and 5 mM phosphate conditions. UV CD scans of monoclonal antibodies were also obtained for comparison. The data presented in Figure 5 indicate that the DVD-Ig proteins have folded structures as indicated by the presence of featured ellipticity profiles in the 250-320 nm region.

The DVD-Ig ellipticity values presented in Figure 5 are similar to those observed for known stable monoclonal antibodies. Therefore, both the DVD-Ig proteins and the mAbs are expected to show similar storage stability. However, LS-DVD-Ig proteins are actually less stable than the mAbs. Therefore, UV-CD data alone cannot be used to accurately predict the stability issues observed for LS-DVD-Ig proteins.

**Example 22: Assessment of the Secondary Structure of DVD-Ig Proteins Using ATR-FTIR**

To assess the secondary structure of DVD-Ig proteins, the second derivative, area normalized FTIR scans taken on an ATR-FTIR instrument from Bruker (Tensor 27) in the

region 1600-1700 cm<sup>-1</sup> were curve fitted and the various peaks were analyzed and added up to get total % beta sheet structure in the molecule. Especially, peaks such as that at 1638 cm<sup>-1</sup>, which are an indicator of the beta sheet arrangement, were taken into account. This is because beta sheets are known to comprise the majority of the secondary structure of mAbs. Therefore it is expected that DVD-Ig proteins which are derived from mAbs should also contain mostly beta sheets as their secondary structure composition. Any deviation from this may indicate that the overall structural integrity of the molecule is compromised. The studies were done in 5 mM citrate/5 mM phosphate buffer at a pH of 4, 6, or 8. The concentration of the DVD-Ig protein was 1mg/ml. The total percentage of beta structure was assessed for 16 DVD-Ig proteins (see Table 44).

Table 44: The Total % Beta Structure in DVD-Ig Proteins Measured Using ATR-FTIR

DVD-Ig	pH 6	pH 4	pH 8
5	89.8	86.5	95.8
6	94.1	78.2	92.3
37	89.6	85.9	90.9
38	96.5	92	95.6
53	NA	NA	96.4
54	97.4	87.7	96.8
65	95.2	95.8	94.8
66	94.6	94.2	94.2
165	NA	NA	95.2
166	95.1	90	95.2
257	95.8	89.6	90.8
258	95.6	88.3	96.6
277	93.6	85.3	95
278	93.8	77.4	94.3
281	94.4	84.4	85
282	93	84.7	84.6

The results presented in Table 45 show that all of the 16 DVD-Ig proteins studied have a folded secondary structure that is composed primarily of beta elements. The proportion of beta elements ranged from about 85% to about 97% which is similar to that observed for standard antibodies. This was unexpected since DVD-Ig proteins are engineered molecules and are assumed to have a lower percentage of beta sheets as compared to mAbs (although they still are expected to contain beta sheets as their predominant secondary structure). These findings suggest that secondary structure composition is not a factor for the overall lower

storage stability of DVD-Ig proteins as compared to antibodies. The proportion of beta elements ranged from about 85% to about 97%.

### EXAMPLE 23: Impact of Ionic Strength and pH on the Second Virial Coefficient of DVD-B Formulations

Second virial coefficient ( $B_{22}$ ) is a thermodynamic parameter and an indicator of the protein-protein attractive or repulsive interactions in solutions. A positive value indicates repulsive interactions and a negative value indicates attractive interactions. Repulsive interactions usually translate into better long term storage. The scattered light intensity is related to the molecular weight and  $B_{22}$  by the following equation.

$$\frac{Kc}{R_\theta} = \frac{1}{M_w} + 2B_{22}c + B_{222}c^2 + \dots$$

Where K is optical constant and is given by

$$K = \frac{[2\pi n \left(\frac{dn}{dc}\right)]^2}{N_A \lambda^4}$$

$R_\theta$  is the excess Rayleigh ratio, a measure of light scattered by the solute, n is the solvent refractive index,  $dn/dc$  is the refractive index increment of the solute,  $N_A$  is the Avogadro's number, and  $\lambda$  is the wavelength of the incident light. Since for most dilute solutions, higher order virial coefficients have negligible values, the following equation (Debye) is used to obtain the second virial coefficients.

$$\frac{Kc}{R_\theta} = \frac{1}{M_w} + 2B_{22}c$$

The scattered intensities were measured on a Malvern Zetasizer Nano. The second virial coefficient values were all positive and indicate that DVD-Ig proteins behave as typical protein molecules with respect to this calculation at least under dilute conditions. The buffers used were acetate for pH 4.5, histidine for pH 6 and Tris for pH 8. 2 mM concentration of buffer was used for 1 mM ionic strength solutions and 10 mM for 20 and 100 mM ionic strength solutions. The rest of the ionic strength was maintained by sodium chloride. The results are shown in Tables 33 and 34. The values of the second virial coefficients were higher on average at pH 4.5 and pH 6.0 than at pH 8.0, suggesting that

DVD-B would store better at pH 4.5 or pH 6.0 than at pH 8.0. Also, the values of the second virial coefficients were higher at lower ionic strength, suggesting that lower ionic strength may also be associated with stability of TNF-PGE2 DVD-Ig protein.

$D_s$  is the self-diffusion coefficient of the molecule at infinite dilution.  $k_d$  is a parameter describing the interaction between the molecules in solution. A positive value for  $k_d$  indicates intermolecular repulsion and vice versa.

Table 45: Virial Coefficient Values for DVD-B protein at 0 mM Ionic Strength

<u>pH</u>	$B_{22} \times 10^{-4}$ (mol mL/gm <sup>2</sup> )	$B_{222} \times 10^{-2}$ (mol mL <sup>2</sup> /gm <sup>3</sup> )	$B_{2222} \times 10^{-1}$ (mol mL <sup>3</sup> /gm <sup>4</sup> )	$B_{22222} \times 10^2$ (mol mL <sup>4</sup> /gm <sup>5</sup> )	$M_w (X 1000)$
6.0	136 ± 5.77	-273.8 ± 16.6	3376.1 ± 339	-221 ± 35.8	189 ± 14

Table 46: Second Virial Coefficients Under Various Conditions for DVD-B Protein

<u>pH</u>	<u>Ionic Strength (mM)</u>	$D_s \times 10^{-7}$ (cm <sup>2</sup> /s) <sup>*</sup>	$k_d$ (mL/gm) <sup>†</sup>	$B_{22} \times 10^{-4}$ (mol mL/gm <sup>2</sup> ) <sup>‡</sup>	$M_w (X 1000)$ <sup>§</sup>
4.5	1	3.42 ± 0.04	434.90 ± 15.31	26.09 ± 0.82	195 ± 1
	20	3.66 ± 0.004	5.96 ± 0.32	5.08 ± 0.03	168 ± 2.89
	100	3.60 ± 0.03	-13.33 ± 0.88	3.09 ± 0.07	160 ± 3.38
6	1	3.41 ± 0.06	379.28 ± 51.34	14.71 ± 1.09	183 ± 0.57
	20	3.75 ± 0.02	-11.04 ± 1.37	3.37 ± 0.06	163 ± 2.35
	100	3.70 ± .002	-23.05 ± 0.02	2.51 ± 0.19	170 ± 3.40
8	1	3.81 ± 0.01	-4.74 ± 0.06	3.50 ± 0.05	155 ± 0.78
	20	3.73 ± 0.02	-27.31 ± 0.29	2.14 ± 0.03	162 ± 5.08
	100	3.66 ± 0.03	-25.33 ± 0.90	2.34 ± 0.07	164 ± 4.59

As mentioned, if  $B_{22}$  and  $kD$  are positive in sign and have a large magnitude, it is an indication of strong repulsive interactions among the DVD-Ig proteins in solution. This is an ideal condition for the long term storage stability of a DVD-Ig protein liquid formulation because there is a lowered probability of DVD-Ig proteins encountering each other in solution to form aggregates. The above data suggest that low ionic strength and low pH as part of a formulation will result in greater long term DVD-Ig protein stability than a formulation with high ionic strength and high pH. Also, since ionic strength and pH

contribute highly to the  $B_{22}$  value, it indicates that electrostatic repulsion in general comprises a significant portion of the  $B_{22}$  value.

With respect to comparing DVD-Ig proteins and antibodies, it may be that a more positive  $B_{22}$  value is required for a DVD-Ig protein formulation to have a similar long term stability profile as that of an antibody formulation. This is because DVD-Ig proteins are larger than antibodies and the hard-sphere contribution to repulsion (and  $B_{22}$ ) is greater for DVD-Ig proteins. Therefore, if both DVD-Ig proteins and antibodies have the same electrostatic repulsions contributing to  $B_{22}$ , the  $B_{22}$  term for DVD-Ig proteins will be more positive to reflect the same degree of protein-protein repulsion in solution.

#### **EXAMPLE 24: Pharmacokinetic Study of DVD-Ig Proteins**

The following example describes pharmacokinetic studies of various DVD-Ig proteins.

##### *Variable Domain Combination and Orientation Impacts Serum Stability*

As described in Figure 2, certain variable domain combinations provide a more stable DVD-Ig protein than other combinations. Figure 2A describes the serum stability for certain DVD-Ig proteins in two different domain orientations, *i.e.*, “outer / inner” and “inner / outer”. For example, the DVD-Ig protein TNF/SOST has a high molecular weight (HMW) % of about 16% in the “outer / inner” orientation, but has about a 75% HMW in the opposite orientation. The domain orientation concepts are set forth in Figure 2B. The DVD-Ig proteins in Figure 2 include short/short linker combinations and are huIgG1 isotypes.

##### *In Vitro Pharmacokinetic Study*

The pharmacokinetic (PK) properties of various biologic therapeutics were assessed following 4 mg/kg single intravenous doses in male Sprague-Dawley rats. Blood samples were collected throughout the 28 day studies. Serum samples were analyzed using an MSD assay employing anti-human Ig capture and Sulfo-Tag labeled goat anti-human IgG antibody for chemiluminescent detection. Pharmacokinetic parameters for each animal were calculated using WinNonlin software by non-compartmental analysis.

Figure 3 describes results from a pharmacokinetic study using rats. The study

examined the correlation between clearance (CL) and  $T \frac{1}{2}$  with high molecular weight (HMW) aggregate formation *in vitro* in rat serum. As shown in Figure 3, DVD-Ig proteins with less than 10% HMW aggregate in vitro are more likely to have low (< 0.3 ml / hour / kg) clearance and long (> 10 days) half life. The outliers included DVD 257 and DVD 258. The DLL4/VEGF DVD-Ig protein recognized the rat target, and the PK was affected by TMD. DVD 037 (VEGF/HER2; SEQ ID NOs: 34 and 35) showed bad *in vitro* properties, but good PK *in vivo*. The amino acid sequences for the heavy and light chains of DLL4/VEGF DVD-Ig protein are provided in SEQ ID NOs: 68 and 69 of Table 65.

#### **EXAMPLE 25: Viscosity Study of DVD-Ig Proteins**

The following example describes viscosity studies for an exemplary AS-DVD-Ig protein (DVD-A).

Viscosity was measured on m-VROC low volume viscometer from Rheosense (Redwood, CA). m-VROC measure viscosity from the pressure drop of a test liquid as it flows through a rectangular slit. As the test liquid is pumped to flow through the flow channel, pressure is measured at increasing distance from the inlet. Plot of the straight line in the pressure vs. position of the sensor is proportional to the viscosity.

The instrument was evacuated beforehand to minimize the usage of material and subsequently recover the material. Air was hence used to clean the instrument before a sample measurement was made. An initial flow rate of 40  $\mu$ l/minute – 200  $\mu$ l/minute was used to obtain the required pressure differential. Saturation of the pressure chamber quickly stabilizes the viscosity reading, and twenty microliters of sample achieved stabilization. Once the instrument has been primed with the sample, less than 5 microliters of additional sample was enough to give a stable second reading. A total of less than thirty microliters (< 35 microliters) of sample was thus enough to give readings in triplicate.

Viscosity of all samples was also measured on a rolling ball viscometer from Anton Paar (X, X). 1.8 mm capillary was used for samples of viscosity range 2-70 cP and 1.6 mm capillary was used for samples of minimal viscosity (less than 2 cP). The instrument was pre-calibrated and run at any of the various possible angles (70°, 50° and/or 40°).

Viscosity of DVDA was determined in histidine formulations having different molarity (*i.e.*, 0 mM to 30 mM histidine) and pH (*i.e.*, pH 4.8 to pH 8.3) in various DVD-A concentrations. Results from the measurements are provided below in Tables 47 to 49.

Table 47: Viscosity Measurements of 34 mg/ml DVD-A at Various pH and Histidine Molarity

DVDA 34mg/ml

<b>Anton Paar Viscosity readings (mPa.s.)</b>					
	<b>pH 4.8</b>	<b>pH 5.4</b>	<b>pH 6</b>	<b>pH 6.6</b>	<b>pH 8.3</b>
<b>0 Mm Histidine</b>	1.83	2.9	5.22	Na	na
<b>5 Mm Histidine</b>	1.59	1.81	2.46	3.86	1.83
<b>30 Mm Histidine</b>	1.33	1.71	2.03	3.5	1.97
<b>Rheosense Viscosity measurements</b>					
	<b>pH 4.8</b>	<b>pH 5.4</b>	<b>pH 6</b>	<b>pH 6.6</b>	<b>pH 8.3</b>
<b>0 Mm Histidine</b>					
1	3.07	4.11	6.03	Na	2.35
2	3.04	4.04	5.84	Na	2.36
3	2.93	4.15	6.08	Na	2.40
<b>Average Viscosity</b>	<b>3.01</b>	<b>4.10</b>	<b>5.98</b>	<b>Na</b>	<b>2.37</b>
<b>5 Mm Histidine</b>					
1	1.58	1.73	2.56	3.87	2.22
2	1.52	1.71	2.55	3.87	2.12
3	1.59	1.72	2.53	3.86	2.13
<b>Average Viscosity</b>	<b>1.56</b>	<b>1.72</b>	<b>2.55</b>	<b>3.86</b>	<b>2.16</b>
<b>30 Mm Histidine</b>					
1	1.37	1.57	1.88	3.34	2.84
2	1.39	1.58	1.91	3.37	2.93
3	1.33	1.58	1.86	3.50	2.73
<b>Average Viscosity</b>	<b>1.36</b>	<b>1.58</b>	<b>1.88</b>	<b>3.40</b>	<b>2.83</b>

Table 48: Viscosity Measurements of 15 mg/ml DVD-A at 0 Mm Histidine at pH 4.8 To 8.3

DVDA 15mg/ml 0mM					
	pH 4.8	pH 5.4	pH 6	pH 6.6	pH 8.3
<b>0 Mm Histidine</b>					
1	1.66	1.30	1.28	Na	1.85
2	1.64	1.34	1.31	Na	1.82
3	1.64	1.30	1.29	Na	1.80
<b>Average Viscosity</b>	<b>1.65</b>	<b>1.31</b>	<b>1.29</b>	<b>Na</b>	<b>1.82</b>

Table 49: Viscosity Measurements of Various Concentrations of DVD-A at Various pH And Histidine Molarity

DVDA					
		5mM		30mM	
		Anton Paar	Rheosense	Anton Paar	Rheosense
pH4.8 (95mg/ml)	1	4.63	6.924	5.29	4.179
	2	4.67	6.925	5.3	4.241
	Average	4.65	6.9245	5.295	4.21
Ph5.4 (77mg/ml)	1	10.45	10.09	6.3	6.027
	2	10.5	10.1	6.29	6.016
	Average	10.475	10.095	6.295	6.0215
pH6 (47.8mg/ml)	1	6.23	5.721	3.643	3.34
	2	6.3	5.695	3.648	3.36
	Average	6.265	5.708	3.6455	3.35
	5mM		30mM		

The results described above in Tables 47-49 show the impact of ionic strength, pH and protein concentration on the viscosity of the DVD-Ig protein solutions. DVD-A (SEQ ID NOs: 62 and 63) is an AS-DVD-Ig protein and the results above show that the viscosity values can be modulated by formulation means to accommodate a syringeable liquid formulation at high concentrations that would be appropriate for pharmaceutical compositions and in-vivo use. Since DVD-Ig proteins have not only a much higher molecular weight of about 200kD compared to standard monoclonal antibodies of about 150 kD, their structure is also more complex. Therefore, it was surprising that the viscosity of mAbs and DVD-Ig is relatively similar and is even at high concentration below about 10 cPs.

## EXAMPLE 26: Thermal Stability of DVD-Ig Proteins

The following example describes results from three different tests examining thermal stability of DVD-Ig proteins, including an exemplary AS-DVD-Ig protein and an exemplary LS-DVD-Ig protein, versus monoclonal antibodies, such as Adalimumab.

### *Dynamic Scanning Fluorescence*

An automated high throughput instrument Optim-1000 from Avacta (York, UK) was used for the study. 9 microliter micro cubic arrays (MCAs) were used for the study. For preparation of stock samples, 3 microliter Sypro orange (Invitrogen, Cambridge, MA) was added to 27 microliter sample solution in order to obtain a final 1X concentration of the dye. The dye is available as 5000X commercial product, although any dye would be suitable. Thermal scans were obtained from 26°C to 95°C at a scan rate of 60°C/hour. Baseline was fitted for linearity and the first point (the temperature) whose inclusion decreased the  $R^2$  below 0.95 was taken as the onset temperature. Repeat studies confirmed that the variation in onset temperatures was less than 5 %.

### *Intrinsic Fluorescence*

Tryptophan fluorescence was used to evaluate the unfolding temperatures. Hitachi FL-4500 instrument from Hitachi (Tokyo, Japan) was used for the study. The temperatures were maintained using a water bath. The temperature in the cuvette was monitored using a thermocouple and a temperature monitor CSi32 from Omega Inc. (Stamford, CT). A front surface triangular quartz cuvette from VWR (MA) was used as this minimized the inner filter effects and hence resulted in strong emission signals. An excitation wavelength of 295 nm was used. Emission was monitored between 328-338 nm. Although the  $\lambda_{max}$  was observed at 332 nm, the intensity was monitored at 335 nm for comparison. The thermal scans were obtained from 30°C to 70°C at a scan rate of 78°C/hour. Baseline was fitted for linearity and the first point (the temperature) whose inclusion decreased the  $R^2$  below 0.95 was taken as the onset temperature. Repeat studies confirmed that the variation in onset temperatures was less than 5 %. The increased scan rate did not significantly affect the onset temperatures.

*Differential Scanning Calorimetry (DSC)*

DSC was used to characterize the thermodynamic stability of the proteins under various solution conditions. An automated cap DSC instrument from Microcal (Northampton, MA) was used. The thermal scans were obtained from 25°C to 65°C at a scan rate of 60°C/hour. Since, aggregation and precipitation that follows unfolding in high concentration samples can lead to blocking of the cap DSC cells which than become rather difficult to clean, the scans were obtained only until  $\approx$  5°C beyond the onset temperature to prevent any such occurrence. A prescan equilibration thermostat of 10 minutes was applied before each scan. A corresponding buffer scan was taken immediately following the sample scan. The difference in onset was less than 2°C between repeat scans. Baseline was fitted for linearity and the first point (the temperature) whose inclusion decreased the  $R^2$  below 0.95 was taken as the onset temperature. Repeat studies confirmed that the variation in onset temperatures was less than 5 %.

Results from the study are provided in Table 50.

Table 50: Results from Thermal Stability Studies Comparing DVD-Ig Proteins to mAbs

	Intrinsic fluorescence		Extrinsic fluorescence		DSC	
	1 mg/mL	75 mg/mL	1 mg/mL	75 mg/mL	1 mg/mL	75 mg/mL
<b>mAb1</b>	66	64	63	57	59	57
<b>mAb2</b>	74	69	73	68	63	59
<b>mAb3</b>	65	62	65	63	59	55
<b>mAb4</b>	72	67	64	58	61	59
<b>mAb5</b>	71	64	66	61	59	56
<b>IL12/IL18</b>	69	67	64	61	62	59
<b>DVD-B</b> <b>TNF/PGE2</b>	56	53	53	49	54	50

The results described in Table 50 show the impact of protein concentration on the thermal stability of the protein solution. DVD1 (IL12IL18), an AS DVD and DVD2 (also referred to herein as DVD-B), an LS-DVD-Ig protein, and other mAbs all show that protein concentration only has a slight impact on the thermal stability of the protein. So the feasibility of a liquid high concentration formulation may be independent of the impact of protein concentration on the thermal stability of the protein; however, high concentration liquid formulations present other well-known types of instabilities, *e.g.*, shelf instabilities. Generally, as described in Examples 1-3, DVD-Ig proteins have a lower melting temperature than antibodies. In some instances, *e.g.*, DVD1, similar melting temperatures are observed, but generally this is not the case.

## **X. ANTI-DLL4 / ANTI-VEGF DVD-IG FORMULATIONS (AQUEOUS AND LYOPHILIZED)**

### **EXAMPLE 27: Preformulation Characterization of Anti-DLL4/Anti-VEGF DVD-Ig Protein h1A11.1SL-Av**

The storage stability (5°C) and accelerated stability (40°C) of an anti-DLL4/anti-VEGF DVD (h1A11.1-SL-Av, Table 41) was evaluated in the formulations and protein concentrations listed below. Stability was evaluated by size exclusion chromatography (SEC) and % aggregate, % monomer, % fragment, and total species recovered were quantitated. Overall, the formulations cover a pH range of 5 to 7 and a protein concentration range of 1.0 to 118 mg/ml.

At 5°C and 40°C temperatures and at protein concentrations of 50, 30, and 10 mg/ml, formulations were: 15 mM acetate pH 5; 15 mM phosphate pH 7; 30 mM acetate, 80 mg/ml sucrose, 0.02% Tween 80 at pH 5; 30 mM histidine, 80 mg/ml sucrose, 0.02% Tween 80 at pH 6; PBS (phosphate buffered saline). All formulations contained 0.02% sodium azide to prevent microbial growth during storage. At 5°C and 40°C temperatures and at protein concentrations of 60, 50, 30, and 10 mg/ml, the formulation was 15 mM histidine pH 6 (also containing 0.02% sodium azide to prevent microbial growth during storage). At 5°C and at a protein concentration of 118 mg/ml, the formulation was 15 mM histidine pH 6 (also containing 0.02% sodium azide to prevent microbial growth during storage). At 40°C and at a protein concentration of 1.0 mg/ml, the formulations were 10 mM citrate and 10 mM phosphate at pHs 5, 6, and 7. Formulations with protein were filtered to remove possible microbes.

Freeze-thaw stability was performed by subjecting the protein in formulation to four cycles of freezing at -80°C for at least 20 hours and thawing in a 30°C water bath. The formulations that were tested for freeze-thaw stability are listed below. Stability was evaluated by SE-HPLC and % aggregate, % monomer, % fragment, and total species recovered were quantitated. The formulations were 15 mM histidine pH 6 at 60 mg/ml protein (also containing 0.02% sodium azide to prevent microbial growth) and 10 mM citrate and 10 mM phosphate at pHs 5, 6, and 7 and 1.0 mg/ml protein (filtered to remove possible microbes).

Finally, differential scanning calorimetry to measure thermal stability was performed on the protein in 10 mM citrate and 10 mM phosphate buffer at pHs 5, 6, and 7 and 1.0 mg/ml protein. The onset temperature of unfolding and the midpoint temperatures of unfolding (Tm) of each protein domain were quantitated.

Table 51: Accelerated Stability at 40°C of H1a11.1-SL-Av at Different Concentrations and In Different Buffers, Excipients, and pHs

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	2.71	96.31	0.98	53058
50, 30, 10	T0	---	ace	5	2.89	96.08	1.03	48033
50, 30, 10	T0	---	his	6	2.81	96.23	0.96	46995
50, 30, 10	T0	---	phos	7	2.91	96.09	1.00	52571
50, 30, 10	T0	---	ace-suc-tw	5	2.54	96.50	0.96	50185
50, 30, 10	T0	---	his-suc-tw	6	2.37	96.62	1.01	50771
50, 30, 10	T0	---	PBS	7	2.90	96.08	1.01	49170
50	T7d	40	ace	5	5.19	93.32	1.49	49028
30	T7d	40	ace	5	3.86	94.68	1.47	48171
10	T7d	40	ace	5	2.60	95.97	1.43	48379
50	T7d	40	his	6	5.25	93.46	1.29	47731
30	T7d	40	his	6	4.13	94.58	1.29	46684
10	T7d	40	his	6	2.73	95.84	1.42	46877
50	T7d	40	phos	7	9.02	89.52	1.46	53429
30	T7d	40	phos	7	6.11	92.40	1.49	51923
10	T7d	40	phos	7	3.94	94.57	1.49	53098
50	T7d	40	ace-suc-tw	5	5.42	92.85	1.73	50373
30	T7d	40	ace-suc-tw	5	4.07	94.06	1.87	48768
10	T7d	40	ace-suc-tw	5	2.66	95.20	2.14	49396
50	T7d	40	his-suc-tw	6	3.44	95.02	1.54	50040
30	T7d	40	his-suc-tw	6	4.16	94.14	1.70	48715
10	T7d	40	his-suc-tw	6	2.86	95.24	1.90	49871
50	T7d	40	PBS	7	8.13	90.28	1.60	49207
30	T7d	40	PBS	7	5.82	92.55	1.63	48853
10	T7d	40	PBS	7	3.62	94.82	1.56	48166
50	T21d	40	ace	5	6.65	90.83	2.51	48536

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
30	T21d	40	ace	5	4.55	92.91	2.54	48520
10	T21d	40	ace	5	2.71	94.70	2.59	48395
50	T21d	40	his	6	7.01	90.71	2.27	46729
30	T21d	40	his	6	4.69	93.10	2.21	46687
10	T21d	40	his	6	2.77	94.93	2.30	46866
50	T21d	40	phos	7	13.39	83.83	2.78	52244
30	T21d	40	phos	7	9.38	87.76	2.86	53556
10	T21d	40	phos	7	4.77	92.32	2.91	52536
50	T21d	40	ace-suc-tw	5	6.37	90.34	3.30	48268
30	T21d	40	ace-suc-tw	5	4.27	91.91	3.82	47211
10	T21d	40	ace-suc-tw	5	2.26	93.02	4.72	46322
50	T21d	40	his-suc-tw	6	6.84	89.82	3.34	47140
30	T21d	40	his-suc-tw	6	4.60	91.90	3.50	47416
10	T21d	40	his-suc-tw	6	2.67	93.66	3.67	48166
50	T21d	40	PBS	7	12.13	84.81	3.06	49845
30	T21d	40	PBS	7	8.09	88.78	3.13	48108
10	T21d	40	PBS	7	4.20	92.63	3.17	48803

Buffer key (all buffers contain 0.02% sodium azide to prevent microbial growth):

**ace** = 15 mM acetate pH 5; **his** = 15 mM histidine pH 6; **phos** = 15 mM phosphate pH

7

**ace-suc-tw** = 30 mM acetate, 80 mg/ml sucrose, 0.02% Tw80

**his-suc-tw** = 30 mM histidine, 80 mg/ml sucrose, 0.02% Tw80

**PBS** = phosphate buffered saline

Table 52. Storage Stability at 5°C of H1a11.1-SL-Av at Different Concentrations and In Different Buffers, Excipients, and pHs (Buffer Key Same as In Table 53)

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	2.71	96.31	0.98	53058
50, 30, 10	T0	---	ace	5	2.89	96.08	1.03	48033
50, 30, 10	T0	---	his	6	2.81	96.23	0.96	46995
50, 30, 10	T0	---	phos	7	2.91	96.09	1.00	52571
50, 30, 10	T0	---	ace-suc-tw	5	2.54	96.50	0.96	50185
50, 30, 10	T0	---	his-suc-tw	6	2.37	96.62	1.01	50771
50, 30, 10	T0	---	PBS	7	2.90	96.08	1.01	49170
50	T7d	5	ace	5	2.96	95.99	1.05	49118
30	T7d	5	ace	5	2.74	96.21	1.06	48434
10	T7d	5	ace	5	2.62	96.23	1.15	48915
50	T7d	5	his	6	2.93	95.87	1.20	47967
30	T7d	5	his	6	2.75	96.06	1.19	47182
10	T7d	5	his	6	2.55	96.31	1.13	47395
50	T7d	5	phos	7	3.15	95.64	1.21	53843

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
30	T7d	5	phos	7	3.10	95.76	1.14	53372
10	T7d	5	phos	7	2.91	95.96	1.13	53269
50	T7d	5	ace-suc-tw	5	2.75	96.13	1.12	50236
30	T7d	5	ace-suc-tw	5	2.62	96.11	1.27	50026
10	T7d	5	ace-suc-tw	5	2.56	96.18	1.26	49290
50	T7d	5	his-suc-tw	6	2.84	96.10	1.07	50129
30	T7d	5	his-suc-tw	6	2.58	96.19	1.23	49272
10	T7d	5	his-suc-tw	6	2.64	96.08	1.28	50926
50	T7d	5	PBS	7	3.26	95.59	1.15	49502
30	T7d	5	PBS	7	3.07	95.64	1.29	49724
10	T7d	5	PBS	7	2.83	95.87	1.29	49563
50	T21d	5	ace	5	2.57	95.76	1.67	49722
30	T21d	5	ace	5	2.37	96.03	1.60	48882
10	T21d	5	ace	5	2.22	96.09	1.69	49255
50	T21d	5	his	6	2.63	95.63	1.74	44884
30	T21d	5	his	6	2.42	95.95	1.62	47510
10	T21d	5	his	6	2.19	96.08	1.73	47015
50	T21d	5	phos	7	3.06	94.96	1.98	53449
30	T21d	5	phos	7	2.69	95.46	1.85	52938
10	T21d	5	phos	7	2.35	95.84	1.81	52703
50	T21d	5	ace-suc-tw	5	2.25	95.76	1.99	50960
30	T21d	5	ace-suc-tw	5	2.08	95.90	2.02	49042
10	T21d	5	ace-suc-tw	5	1.97	95.84	2.19	49851
50	T21d	5	his-suc-tw	6	2.24	95.62	2.14	49983
30	T21d	5	his-suc-tw	6	2.09	95.86	2.05	48813
10	T21d	5	his-suc-tw	6	1.97	95.83	2.19	49984
50	T21d	5	PBS	7	2.84	95.07	2.09	50641
30	T21d	5	PBS	7	2.27	95.62	2.12	48441
10	T21d	5	PBS	7	1.99	95.94	2.07	48978
50	T10mo	5	his	6	8.05	91.04	0.91	45552
30	T10mo	5	his	6	5.81	93.29	0.90	46607
10	T10mo	5	his	6	3.62	95.46	0.92	46207
50	T10mo	5	his-suc-tw	6	8.08	90.26	1.67	45430
30	T10mo	5	his-suc-tw	6	5.98	92.43	1.58	42967
10	T10mo	5	his-suc-tw	6	3.95	94.25	1.80	42567

Table 53. Storage Stability at 5°C, Accelerated Stability at 40°C, and Freeze-Thaw Stability Of H1a11.1-SL-Av at Different Concentrations and In Different Buffers and pHs

Protein Concentration (mg/ml)	time/FT	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
1	T0	---	cit-phos	5	7.07	92.14	0.80	46824
1	T8d	40	cit-phos	5	2.23	96.39	1.38	47090
1	T22d	40	cit-phos	5	7.10	89.62	3.28	47956
1	FT2	---	cit-phos	5	7.91	90.75	1.34	46502
1	FT4	---	cit-phos	5	7.41	92.18	0.41	52181

Protein Concentration (mg/ml)	time/FT	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
1	T0	---	cit-phos	6	7.17	92.33	0.50	45809
1	T8d	40	cit-phos	6	2.56	96.03	1.42	46783
1	T22d	40	cit-phos	6	5.79	91.73	2.48	47401
1	FT2	---	cit-phos	6	7.14	91.48	1.38	45256
1	FT4	---	cit-phos	6	7.09	92.56	0.34	45004
1	T0	---	cit-phos	7	6.82	92.67	0.51	47025
1	T8d	40	cit-phos	7	2.52	95.95	1.53	48080
1	T22d	40	cit-phos	7	5.52	91.58	2.90	48706
1	FT2	---	cit-phos	7	7.23	91.52	1.25	46732
1	FT4	---	cit-phos	7	7.15	92.49	0.36	46561
60 and 118	T0	---	his	6	8.03	91.15	0.82	43528
60	T7d	40	his	6	7.17	91.76	1.07	45333
60	T21d	40	his	6	15.77	82.13	2.10	44729
60	T7d	5	his	6	3.83	95.32	0.86	46774
60	T26d	5	his	6	7.14	92.56	0.30	63982
118	T5mo	5	his	6	12.82	86.65	0.53	55869
60	T5mo	5	his	6	9.46	90.03	0.51	64573
60	FT2	---	his	6	6.71	92.59	0.70	42259
60	FT4	---	his	6	6.33	93.62	0.05	41054

Key:

**FT** = freeze thaw

**FT2** = analysis after two cycles of freeze and thaw; freezing at -80°C and thawing in a 30°C water bath

**FT4** = analysis after four cycles of freeze and thaw; freezing at -80°C and thawing in a 30°C water bath

**cit-phos** = 10 mM citrate and 10 mM phosphate

**his** = 15 mM histidine and 0.02% sodium azide (azide for preventing microbial growth)

Table 54. Differential Scanning Calorimetry Data of H1a11.1-SL-Av at 1 mg/ml in 10 mM Citrate + 10 mM Phosphate at Different pHs

pH	Onset (°C)	Tm1 (°C)	Tm2 (°C)	Tm3 (°C)	Tm4 (°C)
5	55	68.2	68.86	75.56	81.18
6	58	69.04	70.47	75.24	82.04
7	59	69.52	70.94	74.44	82.06

Overall, the data in Tables 51 to 54 suggest that h1A11.1-SL-Av has a degradation profile typically observed for stable monoclonal antibodies. The aggregation and fragmentation is greater at elevated temperatures (40°C) than at lower temperatures (5°C).

At 5°C, there is no apparent increase in aggregation or fragmentation after 21 days of storage. At 40°C, the amount of degradation was higher than those values typically observed for stable mAbs. However, it was not so great as to preclude it from further development. Overall, this DVD-Ig protein qualified as an AS-DVD-Ig protein based on its stability.

#### **EXAMPLE 28: Formulation Selection for Anti-DLL4/Anti-VEGF DVD-Ig Protein**

Materials and Methods. The stability of anti-DLL4/anti-VEGF DVD-Ig h1A11.1-SL-Av protein was evaluated in the five formulations listed in Table 55. All formulations were prepared in 15 mM histidine buffer. Formulations F1 to F4 were prepared at 50 mg/ml protein concentration. In these formulations, the pH ranged from 5.5 to 6.0, polysorbate 80 concentration ranged from 0 to 0.05% w/v, sucrose concentration ranged from 0 to 7.5% w/v, and arginine concentration ranged from 0 to 1% w/v. Formulation F4 was prepared in 15 mM histidine buffer at pH 6.0 without any stabilizers and served as a study control for the 50 mg/ml liquid formulation stability assessment. In addition, one formulation was prepared at 25 mg/ml protein concentration at pH 6.0 (Formulation F5). In this formulation, the polysorbate 80 concentration was 0.025% w/v and sucrose concentration was 3.8% w/v.

Table 55. Formulation Composition Description

Formulation Identifier	anti-DLL4 / anti-VEGF DVD-Ig Concentration (mg/mL)	Buffer	pH	Polysorbate 80 (Tween 80) (% w/v)	Sucrose (% w/v)	Arginine (% w/v)
F1	50	15 mM Histidine	6.0	0.05	7.5	0
F2	50	15 mM Histidine	5.5	0.05	7.5	0
F3	50	15 mM Histidine	6.0	0.05	7.5	1
F4	50	15 mM Histidine	6.0	0	0	0
F5	25	15 mM Histidine	6.0	0.025	3.8	0

In the above formulations, 15 mM histidine buffer was selected because it provides adequate buffering capacity to maintain the target formulation pH. Sucrose was evaluated as a stabilizer against freeze-thaw stress (cryoprotectant) and lyophilization process-induced stress (lyoprotectant). Polysorbate 80 (surfactant) and arginine were added to potentially stabilize the formulation against aggregates and particulates formation.

Freeze-thaw and Liquid Formulation Stability Testing. The stability of all liquid formulations was evaluated after three cycles of freeze/thaw (F/T) stress, and after 1 month storage at -80, 5, 25 and 40° C. Stability was tested by a broad panel of analytical assays including Visual appearance, % Aggregates by Size Exclusion Chromatography (SE-HPLC), Charge heterogeneity by Cation Exchange Chromatography (CEX-HPLC), Fragmentation by reduced SDS-Capillary Electrophoresis (CE-SDS), Sub-visible particles by Micro Flow Imaging (MFI) and DLL4/ VEGF binding potency using ELISA.

The freeze/thaw and liquid stability testing results are provided in Table 56.

Table 56. Freeze-Thaw and Liquid Formulation Stability Results

Formulation Identifier	Time Points	Visual Appearance	% Aggregates by SEC-HPLC	CEX-HPLC			% Purity	Sub-visible Particles by MFI			Binding Potency by ELISA	
				% Acidic region	% Main peak	% Basic region		≥ 2 μm/mL	≥ 10 μm/mL	≥ 25 μm/mL	DLL4	VEGF
F1	T0	EFVP	1.0	21.6	61.7	16.7	97.7	3333	5	0	93	113
	3FT	EFVP	1.1	21.5	61.8	16.6	97.7	2388	50	5	NP	NP
	1M at -80°C	EFVP	1.1	21.0	62.1	16.8	97.8	1364	15	5	NP	NP
	1M at 5°C	EFVP	2.0	21.2	62.4	16.3	97.8	1064	0	0	NP	NP
	1M at 25°C	EFVP	4.3	23.5	62.2	14.2	97.4	1559	5	5	NP	NP
	1M at 40°C	EFVP	7.2	35.8	43.0	21.2	95.0	1219	15	0	94	104
F2	T0	EFVP	1.3	21.4	61.8	16.8	97.5	1589	15	0	93	113
	3FT	EFVP	1.3	21.4	61.8	16.9	97.6	435	5	5	NP	NP
	1M at -80°C	EFVP	1.4	21.0	62.0	17.0	97.8	315	0	0	NP	NP
	1M at 5°C	EFVP	2.1	20.9	62.3	16.8	97.8	230	5	5	NP	NP
	1M at 25°C	EFVP	4.4	22.8	61.6	15.6	97.4	1149	5	0	NP	NP
	1M at 40°C	EFVP	7.9	33.8	40.9	25.3	95.1	655	5	0	95	97
F3	T0	EFVP	1.1	21.5	61.7	16.7	97.6	784	0	0	93	113
	3FT	EFVP	1.1	21.3	61.8	16.9	97.5	490	10	0	NP	NP
	1M at -80°C	EFVP	1.1	21.0	61.9	17.1	97.9	250	0	0	NP	NP
	1M at 5°C	EFVP	2.0	20.8	62.0	17.2	97.8	225	5	0	NP	NP
	1M at 25°C	EFVP	4.9	21.5	58.3	20.2	97.3	834	10	0	NP	NP
	1M at 40°C	EFVP	11.3	32.1	42.3	25.6	95.0	1464	5	0	97	101
F4	T0	TMTc	1.2	21.6	61.8	16.6	97.4	23707	370	5	93	113
	3FT	TMTc	1.5	21.4	61.8	16.8	97.4	105467	5906	30	NP	NP
	1M at -	TMTc	1.2	21.1	62.0	16.9	97.8	42024	1329	60	NP	NP

Formulation Identifier	Time Points	Visual Appearance	% Aggregates by SEC-HPLC	CEX-HPLC			% Purity	Sub-visible Particles by MFI			Binding Potency by ELISA	
				% Acidic region	% Main peak	% Basic region		≥ 2 $\mu\text{m}/\text{mL}$	≥ 10 $\mu\text{m}/\text{mL}$	≥ 25 $\mu\text{m}/\text{mL}$	DLL4	VEGF
	80°C											
	1M at 5°C	EFVP	2.0	21.3	62.3	16.5	97.8	1189	0	0	NP	NP
	1M at 25°C	EFVP	4.5	23.5	61.8	14.7	97.2	89299	3053	165	NP	NP
	1M at 40°C	EFVP	7.7	34.9	44.2	20.9	94.8	61754	5051	670	101	97
F5	T0	EFVP	0.9	21.6	61.6	16.7	97.7	2808	5	5	93	113
	3FT	EFVP	1.2	21.5	61.8	16.8	97.6	1949	0	0	NP	NP
	1M at -80°C	EFVP	1.1	21.0	62.2	16.8	97.8	270	5	0	NP	NP
	1M at 5°C	EFVP	1.5	21.2	61.9	16.8	97.8	709	10	0	NP	NP
	1M at 25°C	EFVP	2.6	23.5	62.0	14.6	97.2	944	15	0	NP	NP
	1M at 40°C	EFVP	3.9	37.0	44.4	18.6	95.0	974	20	0	92	95

Key. EFVP: Essentially Free of Visible Particles, TMTC: Too Many To Count, NP: Not Performed

As seen in the above table, Freeze-thaw stress resulted in the formation of visible particles and significantly higher sub-visible particle counts in Formulation F4 that was formulated without any stabilizers (e.g., polysorbate 80, sucrose, and arginine). Relative to the other formulations, this formulation also showed a trend of higher sub-visible particle counts after 1 month storage at 25 and 40°C. Formulation F5 with 25 mg/mL protein concentration showed significantly lower aggregation relative to the 50 mg/mL formulations over 1 month storage at 5, 25 and 40°C.

Lyophilized Formulation Stability Testing. The stability of select formulations was also evaluated after the formulations were lyophilized. The lyophilized drug product stability was assessed for all sucrose-containing formulations (F1, F2, F3, and F5). Stability was assessed after 2 weeks storage at 55°C. Stability was tested by a broad panel of analytical assays including Visual appearance (before and after reconstitution), Reconstitution time, % Aggregates by Size Exclusion Chromatography (SE-HPLC), Charge heterogeneity by Cation Exchange Chromatography (CEX-HPLC), Fragmentation by reduced SDS-Capillary Electrophoresis (CE-SDS), Sub-visible particles by Micro Flow Imaging (MFI), and Water Content by Karl Fischer titration.

The lyophilized formulation stability testing results are provided in Table 57. Reconstitution time for all evaluated formulations was approximately 1 minute. A slight increase in aggregation by SEC and % basic region by CEX was observed for all formulations under the stressed storage condition of 55°C. Minimal changes were observed in all other measured product stability attributes.

Table 57. Lyophilized Formulation Stability Results

Formulation Identifier	Time	Visual Appearance		% Aggreg by SEC-HPLC	CEX-HPLC			% Purity by R CE-SDS	Sub-visible Particles by MFI		
		Before Recon	After Recon		% Acidic region	% Main peak	% Basic region		≥ 2 μm/mL	≥ 10 μm/mL	≥ 25 μm/mL
<b>F1</b>	T0	White to off-white cake	EFVP	1.1	21.3	61.9	16.8	97.6	749	20	10
	2 weeks at 55°C	White to off-white cake	EFVP	1.6	20.6	58.1	21.3	97.5	1639	15	0
<b>F2</b>	T0	White to off-white cake	EFVP	1.3	21.2	61.8	17.0	97.6	1254	15	0
	2 weeks at 55°C	White to off-white cake	EFVP	2.0	20.4	57.8	21.8	97.6	1609	10	0
<b>F3</b>	T0	White to off-white cake	EFVP	1.1	21.2	61.9	16.9	97.6	719	5	0
	2 weeks at 55°C	White to off-white cake	EFVP	1.4	20.8	59.5	19.8	97.5	475	10	5
<b>F5</b>	T0	White to off-white cake	EFVP	1.0	21.3	61.8	16.9	97.5	844	35	5
	2 weeks at 55°C	White to off-white cake	EFVP	1.5	20.5	58.0	21.5	97.7	270	5	0

Key. EFVP = Essentially Free of Visible Particles; NP = Not Performed

The sequence of the anti-DLL4 / anti-VEGF DVD-Ig protein H1A11.1-SL-Av is set forth in Table 58 (DVD-Ig protein described in Examples 28 and 29).

Table 58: Full Length Sequence For DVD h1A11.1-SL-Av

Name	Sequence
	123456789012345678901234567890123456789012345678901234567890
h1A11.1-SL-Av light chain	DIQMTQSPSSLSASVGDRVTITC <b>RASEDIYSNLAWYQQKPGKAPKLLIYDTNNLADGVPS</b> RFSGSGSGTDFLTISLQPEDFATYYC <b>QQYNNYPPTFGQGTKLEIKR</b> <u><b>TVAAPSVFIFPP</b></u> DIQMTQSPSSLSASVGDRVTITC <b>SASQDISNYLNWYQQKPGKAPKVL</b> IY <u><b>FTSSLHSGVPS</b></u> RFSGSGSGTDFLTISLQPEDFATYYC <b>QQYSTVPWTFGQGTKVEIKR</b> <u><b>TVAAPSVFIFPP</b></u> <b>SDEQLKSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLT</b> <b>LSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</b> (SEQ ID NO: 28)
h1A11.1-SL-Av heavy chain	EVQLVESGGGLVQPGGSLRLSCAASGFTFS <b>NFPMA</b> WVRQAPGKGLEWVAT <b>ISSSDGTTYY</b> <b>RDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARGYYNSPFAY</b> WGQGTLVTVSS <u><b>AS</b></u> <u><b>TKGP</b></u> EVQLVESGGGLVQPGGSLRLSCAASGTYFFT <b>NYGMN</b> WVRQAPGKGLEWVG <b>WINTYTG</b> <b>EPTYAADFKRRTFSLDTSKSTAYLQMNSLRAEDTAVYYCAKYPHYYGSSH</b> WYFDVWGQG TLTVVSS <u><b>ASTKGPSVFPLAPSSKTSGGTAALGCLVKDYFPEPVTVWSNGALTSGVHTF</b></u> <b>PAVLIQSSGLYSLSSVVTVPSSSLGQTQYICNVNHKP</b> SNTKVDKKVEPKSCDKTHTCPPCP <b>APEAAGGPSPVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVFKFNWYVDGVEVHNAKTK</b> <b>PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISAKGQPREPQVYT</b> <b>LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGOPENNYKTPVLDSDGSFFLYSKL</b> <b>TVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSPGK</b> (SEQ ID NO: 29)

Table 58 provides the full-length heavy and light chain sequences for DVD h1A11.1-SL-Av. Linker sequences are underlined, while CDRs and constant region sequences are in bold.

#### EXAMPLE 29: Extended Preformulation Characterization of Second Anti-DLL4/Anti-VEGF DVD-Ig Protein (h1A11.1-LS-Av)

Extended preformulation characterization on anti-DLL4/-antiVEGF DVD-Ig proteins was performed to explore how different formulations conditions impact the stability of the DVD-Ig proteins. Data for h1A11.1-LS-Av is presented in Tables 59 and 60. The storage stability (5°C) and accelerated stability (40°C) of the DVD-Ig protein was evaluated in the formulations and protein concentrations listed below. Stability was evaluated by SEC and % aggregate, % monomer, % fragment, and total species recovered were quantitated. Overall, the formulations cover a pH range of 5 to 7 and a protein concentration range of 10 to 50 mg/ml.

At 5°C and 40°C temperatures and at concentrations of 50, 30, and 10 mg/ml the following formulations were evaluated: 15 mM acetate pH 5, 15 mM histidine pH 6, 15 mM phosphate pH 7, 30 mM acetate, 80 mg/ml sucrose, 0.02% Tween 80 at pH 5, 30 mM

histidine, 80 mg/ml sucrose, 0.02% Tween 80 at pH 6, and PBS (phosphate buffered saline). All formulations contained 0.02% sodium azide to prevent microbial growth during storage.

Table 59. Accelerated Stability at 40°C of h1A11.1-LS-Av

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	0.21	98.42	1.36	56054
50, 30, 10	T0	---	ace	5	0.28	98.41	1.31	56381
50, 30, 10	T0	---	his	6	0.46	98.23	1.31	54316
50, 30, 10	T0	---	phos	7	0.74	97.86	1.40	53212
50, 30, 10	T0	---	ace-suc-tw	5	0.24	98.16	1.60	56244
50, 30, 10	T0	---	his-suc-tw	6	0.30	98.11	1.59	54076
50, 30, 10	T0	---	PBS	7	0.52	98.05	1.43	50085
50	T7d	40	ace	5	1.63	96.74	1.63	55563
30	T7d	40	ace	5	1.13	97.24	1.62	55194
10	T7d	40	ace	5	0.84	97.49	1.67	55029
50	T7d	40	his	6	2.00	96.62	1.38	53566
30	T7d	40	his	6	1.17	97.46	1.38	52443
10	T7d	40	his	6	0.60	98.00	1.40	53812
50	T7d	40	phos	7	4.31	94.02	1.67	52934
30	T7d	40	phos	7	2.85	95.46	1.69	52663
10	T7d	40	phos	7	1.20	97.11	1.69	52411
50	T7d	40	ace-suc-tw	5	1.10	96.23	2.66	54837
30	T7d	40	ace-suc-tw	5	0.77	96.40	2.83	52474
10	T7d	40	ace-suc-tw	5	0.43	96.39	3.17	50855
50	T7d	40	his-suc-tw	6	1.69	96.27	2.05	53017
30	T7d	40	his-suc-tw	6	1.14	96.84	2.02	52153
10	T7d	40	his-suc-tw	6	0.59	97.30	2.11	52208
50	T7d	40	PBS	7	2.77	95.30	1.93	51623
30	T7d	40	PBS	7	1.73	96.28	1.99	49973
10	T7d	40	PBS	7	0.78	97.25	1.97	50851
50	T21d	40	ace	5	3.66	94.30	2.04	55920
30	T21d	40	ace	5	2.56	95.33	2.10	54188
10	T21d	40	ace	5	1.85	96.00	2.15	55213
50	T21d	40	his	6	4.14	94.28	1.58	54807
30	T21d	40	his	6	2.67	95.79	1.54	53071
10	T21d	40	his	6	1.59	96.82	1.58	54053
50	T21d	40	phos	7	8.52	89.32	2.16	53273
30	T21d	40	phos	7	5.58	92.54	1.89	53162
10	T21d	40	phos	7	3.01	94.89	2.10	52747
50	T21d	40	ace-suc-tw	5	4.12	93.78	2.10	56278
30	T21d	40	ace-suc-tw	5	2.93	94.94	2.13	55481
10	T21d	40	ace-suc-tw	5	1.99	95.75	2.26	54696
50	T21d	40	his-suc-tw	6	4.94	93.21	1.85	54034
30	T21d	40	his-suc-tw	6	n/a	n/a	n/a	n/a
10	T21d	40	his-suc-tw	6	2.00	96.30	1.70	52686
50	T21d	40	PBS	7	8.44	89.65	1.90	51697

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
30	T21d	40	PBS	7	5.54	92.43	2.03	50282
10	T21d	40	PBS	7	2.89	95.05	2.06	51580

Buffer key (all buffers contain 0.02% sodium azide to prevent microbial growth): **ace** = 15 mM acetate pH 5; **his** = 15 mM histidine pH 6; **phos** = 15 mM phosphate pH 7; **ace-suc-tw** = 30 mM acetate, 80 mg/ml sucrose, 0.02% Tween80; **his-suc-tw** = 30 mM histidine, 80 mg/ml sucrose, 0.02% Tween80; **PBS** = phosphate buffered saline

Table 60. Storage Stability At 5°C of h1A11.1-LS-Av

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	0.21	98.42	1.36	56054
50, 30, 10	T0	---	ace	5	0.28	98.41	1.31	56381
50, 30, 10	T0	---	his	6	0.46	98.23	1.31	54316
50, 30, 10	T0	---	phos	7	0.74	97.86	1.40	53212
50, 30, 10	T0	---	ace-suc-tw	5	0.24	98.16	1.60	56244
50, 30, 10	T0	---	his-suc-tw	6	0.30	98.11	1.59	54076
50, 30, 10	T0	---	PBS	7	0.52	98.05	1.43	50085
50	T7d	5	ace	5	0.18	98.17	1.64	57599
30	T7d	5	ace	5	0.16	98.21	1.64	55889
10	T7d	5	ace	5	0.13	98.17	1.70	53289
50	T7d	5	his	6	0.18	98.14	1.68	55742
30	T7d	5	his	6	0.12	98.06	1.82	53603
10	T7d	5	his	6	0.13	98.07	1.80	53505
50	T7d	5	phos	7	0.23	97.72	2.05	54355
30	T7d	5	phos	7	0.18	97.77	2.04	53561
10	T7d	5	phos	7	0.13	97.72	2.15	53151
50	T7d	5	ace-suc-tw	5	0.09	97.40	2.51	57158
30	T7d	5	ace-suc-tw	5	0.08	97.43	2.49	55025
10	T7d	5	ace-suc-tw	5	0.08	97.34	2.58	53882
50	T7d	5	his-suc-tw	6	0.10	97.48	2.43	55272
30	T7d	5	his-suc-tw	6	0.08	97.63	2.29	52763
10	T7d	5	his-suc-tw	6	0.05	97.41	2.53	52903
50	T7d	5	PBS	7	0.12	97.31	2.58	51698
30	T7d	5	PBS	7	0.09	97.24	2.67	50144
10	T7d	5	PBS	7	0.08	97.28	2.64	50428
50	T21d	5	ace	5	0.87	98.45	0.68	57706
30	T21d	5	ace	5	0.80	98.55	0.65	56566
10	T21d	5	ace	5	0.83	98.47	0.70	54226
50	T21d	5	his	6	1.05	98.29	0.66	55911
30	T21d	5	his	6	0.92	98.40	0.68	54225

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
10	T21d	5	his	6	0.90	98.41	0.70	54128
50	T21d	5	phos	7	1.25	98.09	0.66	54980
30	T21d	5	phos	7	1.20	98.11	0.69	53903
10	T21d	5	phos	7	1.01	98.29	0.69	53271
50	T21d	5	ace-suc-tw	5	0.92	98.36	0.72	61574
30	T21d	5	ace-suc-tw	5	0.89	98.39	0.72	55532
10	T21d	5	ace-suc-tw	5	0.83	98.46	0.71	55841
50	T21d	5	his-suc-tw	6	1.00	98.27	0.73	55484
30	T21d	5	his-suc-tw	6	0.92	98.37	0.70	53335
10	T21d	5	his-suc-tw	6	0.82	98.49	0.69	53736
50	T21d	5	PBS	7	1.49	97.79	0.71	52405
30	T21d	5	PBS	7	1.29	98.02	0.70	51284
10	T21d	5	PBS	7	1.12	98.18	0.70	51377

The buffer key for Table 59 is the same as in Table 60.

### Example 30: Formulation Studies of Additional DLL4-VEGF DVD-Ig Proteins

#### h1A11.1.A6-LS-Av and h1A11.1.A6-SL-Av

Extended preformulation characterization on additional DLL4/VEGF DVD-Ig proteins was performed to explore how different formulations conditions impact the stability. Data for h1A11.1.A6-LS-Av and h1A11.1.A6-SL-Av DLL4/VEGF DVD-Ig proteins are presented below. These DVD-Ig proteins proved to be unstable and failed the screening criteria to be considered AS-DVD-Ig proteins.

The storage stability (5°C) and accelerated stability (40°C) of h1A11.1.A6-LS-Av and h1A11.1.A6-SL-Av were evaluated in the formulations and protein concentrations listed below. Stability was evaluated by SEC and % aggregate, % monomer, % fragment, and total species recovered were quantitated. Overall, the formulations cover a pH range of 5 to 7 and a protein concentration range of 10 to 50 mg/ml.

At 5°C and 40°C temperatures and at 50, 30, and 10 mg/ml, the following conditions were tested: 15 mM acetate pH 5; 15 mM histidine pH 6; 15 mM phosphate pH 7; 30 mM acetate, 80 mg/ml sucrose, 0.02% Tween 80 at pH 5; 30 mM histidine, 80 mg/ml sucrose,

0.02% Tween 80 at pH 6; and PBS (phosphate buffered saline). All formulations contained 0.02% sodium azide to prevent microbial growth during storage

Overall, the data provided in Tables 61-63 suggests the two DVD-Ig proteins have an atypical degradation profile not observed for stable monoclonal antibodies. The aggregation rate was actually greater at 5°C than at 40°C.

At 5°C, there is a rapid increase in aggregation after 21 days of storage. At 40°C, the amount of degradation also increases, but not at the rate observed at 5°C. In both cases, the aggregation is concentration dependent.

Overall, these DVD-Ig proteins failed the screen based on their 5°C instability and are examples of non-AS-DVD-Ig proteins.

**Table 61. Accelerated Stability at 40°C of H1a11.1.A6-LS-Av at Different Concentrations and In Different Buffers, Excipients, and pHs**

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	0.66	98.64	0.70	48425
50, 30, 10	T0	---	ace	5	1.24	98.05	0.72	50944
50, 30, 10	T0	---	his	6	1.49	97.74	0.76	46462
50, 30, 10	T0	---	phos	7	1.76	97.67	0.58	43322
50, 30, 10	T0	---	ace-suc-tw	5	1.66	97.69	0.65	57038
50, 30, 10	T0	---	his-suc-tw	6	1.62	97.87	0.52	54299
50, 30, 10	T0	---	PBS	7	2.22	97.25	0.53	48116
50	T7d	40	ace	5	4.40	94.52	1.08	43246
30	T7d	40	ace	5	2.76	96.17	1.08	46266
10	T7d	40	ace	5	1.54	97.41	1.05	52178
50	T7d	40	his	6	5.11	94.02	0.87	50471
30	T7d	40	his	6	3.17	95.97	0.87	40875
10	T7d	40	his	6	1.60	97.52	0.88	48759
50	T7d	40	phos	7	12.24	86.69	1.07	44327
30	T7d	40	phos	7	7.81	91.09	1.11	42284
10	T7d	40	phos	7	3.46	95.42	1.12	44023
50	T7d	40	ace-suc-tw	5	5.46	93.48	1.07	63100
30	T7d	40	ace-suc-tw	5	3.42	95.55	1.03	59243
10	T7d	40	ace-suc-tw	5	1.83	97.03	1.14	60361
50	T7d	40	his-suc-tw	6	4.90	94.14	0.96	56706

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
30	T7d	40	his-suc-tw	6	3.65	95.50	0.84	53538
10	T7d	40	his-suc-tw	6	1.78	97.31	0.91	55447
50	T7d	40	PBS	7	14.78	84.08	1.14	58511
30	T7d	40	PBS	7	9.20	89.56	1.24	46917
10	T7d	40	PBS	7	4.08	94.73	1.19	49978
50	T21d	40	ace	5	5.58	92.75	1.67	53248
30	T21d	40	ace	5	3.59	94.67	1.74	51038
10	T21d	40	ace	5	1.90	96.38	1.72	50617
50	T21d	40	his	6	6.25	92.37	1.37	46908
30	T21d	40	his	6	3.82	94.77	1.41	46075
10	T21d	40	his	6	1.93	96.61	1.47	47827
50	T21d	40	phos	7	13.12	84.98	1.90	40311
30	T21d	40	phos	7	9.11	88.98	1.91	42037
10	T21d	40	phos	7	4.05	93.96	1.99	43009
50	T21d	40	ace-suc-tw	5	6.45	91.73	1.82	56752
30	T21d	40	ace-suc-tw	5	4.21	93.92	1.87	56330
10	T21d	40	ace-suc-tw	5	2.11	95.88	2.00	57757
50	T21d	40	his-suc-tw	6	6.69	91.83	1.49	54091
30	T21d	40	his-suc-tw	6	4.22	94.30	1.48	52932
10	T21d	40	his-suc-tw	6	2.09	96.45	1.47	54118
50	T21d	40	PBS	7	15.74	82.23	2.03	52080
30	T21d	40	PBS	7	9.98	87.82	2.19	47188
10	T21d	40	PBS	7	4.71	93.04	2.25	47400

Buffer key (all buffers contain 0.02% sodium azide to prevent microbial growth):

**ace** = 15 mM acetate pH 5; **his** = 15 mM histidine pH 6; **phos** = 15 mM phosphate pH

7

**ace-suc-tw** = 30 mM acetate, 80 mg/ml sucrose, 0.02% Tw80

**his-suc-tw** = 30 mM histidine, 80 mg/ml sucrose, 0.02% Tw80

**PBS** = phosphate buffered saline

Table 62. Storage Stability at 5°C of H1a11.1.A6-LS-Av at Different Concentrations and In Different Buffers, Excipients, and pHs (Buffer Key Same as In Table 63)

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	0.66	98.64	0.70	48425

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
50, 30, 10	T0	---	ace	5	1.24	98.05	0.72	50944
50, 30, 10	T0	---	his	6	1.49	97.74	0.76	46462
50, 30, 10	T0	---	phos	7	1.76	97.67	0.58	43322
50, 30, 10	T0	---	ace-suc-tw	5	1.66	97.69	0.65	57038
50, 30, 10	T0	---	his-suc-tw	6	1.62	97.87	0.52	54299
50, 30, 10	T0	---	PBS	7	2.22	97.25	0.53	48116
50	T7d	5	ace	5	4.68	94.74	0.58	54446
30	T7d	5	ace	5	2.77	96.62	0.62	51286
10	T7d	5	ace	5	1.75	97.62	0.63	51444
50	T7d	5	his	6	7.46	91.88	0.66	46683
30	T7d	5	his	6	4.83	94.49	0.68	46282
10	T7d	5	his	6	2.36	96.95	0.69	47844
50	T7d	5	phos	7	1.84	97.46	0.71	34008
30	T7d	5	phos	7	1.89	97.48	0.63	36992
10	T7d	5	phos	7	2.05	97.30	0.65	41962
50	T7d	5	ace-suc-tw	5	4.74	94.65	0.61	57937
30	T7d	5	ace-suc-tw	5	N/A	N/A	N/A	N/A
10	T7d	5	ace-suc-tw	5	1.95	97.41	0.64	58618
50	T7d	5	his-suc-tw	6	7.97	91.45	0.59	54735
30	T7d	5	his-suc-tw	6	4.85	94.49	0.66	53379
10	T7d	5	his-suc-tw	6	2.34	97.01	0.65	54187
50	T7d	5	PBS	7	5.68	93.65	0.67	46544
30	T7d	5	PBS	7	5.20	94.13	0.67	45219
10	T7d	5	PBS	7	3.56	95.76	0.68	47653
50	T21d	5	ace	5	9.33	89.97	0.70	52020
30	T21d	5	ace	5	4.25	95.04	0.70	51223
10	T21d	5	ace	5	1.95	97.36	0.69	50950
50	T21d	5	his	6	19.35	79.71	0.94	44105
30	T21d	5	his	6	9.91	89.29	0.80	46096
10	T21d	5	his	6	3.11	96.15	0.73	47777
50	T21d	5	phos	7	1.65	97.49	0.86	25059
30	T21d	5	phos	7	1.49	97.75	0.76	29723
10	T21d	5	phos	7	2.00	97.09	0.90	38517
50	T21d	5	ace-suc-tw	5	9.41	89.79	0.79	56438
30	T21d	5	ace-suc-tw	5	4.72	94.50	0.79	56230
10	T21d	5	ace-suc-tw	5	2.13	97.01	0.86	58579
50	T21d	5	his-suc-tw	6	20.57	78.32	1.11	53114
30	T21d	5	his-suc-tw	6	10.17	88.99	0.85	53155
10	T21d	5	his-suc-tw	6	3.20	95.94	0.86	54028
50	T21d	5	PBS	7	5.65	93.38	0.98	34294
30	T21d	5	PBS	7	2.99	96.04	0.96	36457

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
10	T21d	5	PBS	7	5.14	93.95	0.91	46566

Table 63. Accelerated Stability at 40°C of H1a11.1.A6-SL-Av at Different Concentrations and In Different Buffers, Excipients, and pHs

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	1.19	97.75	1.06	58988
50, 30, 10	T0	---	ace	5	1.23	97.85	0.92	47138
50, 30, 10	T0	---	his	6	1.30	97.84	0.87	44711
3.4	T0	---	phos	7	0.86	98.00	1.14	57373
50, 30, 10	T0	---	ace-suc-tw	5	1.30	97.85	0.85	52129
50, 30, 10	T0	---	his-suc-tw	6	1.28	97.85	0.87	47563
6.6	T0	---	PBS	7	1.35	97.76	0.89	71146
50	T7d	40	ace	5	7.66	90.88	1.46	48331
30	T7d	40	ace	5	4.05	94.43	1.52	45731
10	T7d	40	ace	5	1.45	97.05	1.51	47455
50	T7d	40	his	6	8.68	90.23	1.09	45897
30	T7d	40	his	6	4.74	94.15	1.11	44807
10	T7d	40	his	6	1.62	97.07	1.31	45567
3.4	T7d	40	phos	7	2.04	96.51	1.45	56002
50	T7d	40	ace-suc-tw	5	9.10	89.27	1.62	52696
30	T7d	40	ace-suc-tw	5	5.23	93.14	1.64	50591
10	T7d	40	ace-suc-tw	5	1.91	96.09	2.00	51790
50	T7d	40	his-suc-tw	6	8.55	90.18	1.27	48458
30	T7d	40	his-suc-tw	6	4.82	93.19	1.99	46963
10	T7d	40	his-suc-tw	6	1.78	96.78	1.45	46676
6.6	T7d	40	PBS	7	4.53	93.83	1.64	70277
50	T21d	40	ace	5	8.27	89.20	2.53	47139
30	T21d	40	ace	5	4.41	93.03	2.56	45779
10	T21d	40	ace	5	1.60	95.68	2.72	46794
50	T21d	40	his	6	9.26	88.56	2.18	44423
30	T21d	40	his	6	5.19	92.86	1.96	43874
10	T21d	40	his	6	1.74	95.71	2.55	45249
3.4	T21d	40	phos	7	2.43	95.14	2.42	54476
50	T21d	40	ace-suc-tw	5	9.45	87.66	2.89	51846
30	T21d	40	ace-suc-tw	5	5.35	91.28	3.37	50456

Protein Concentration (mg/ml)	time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
10	T21d	40	ace-suc-tw	5	1.94	94.29	3.78	50414
50	T21d	40	his-suc-tw	6	8.72	89.02	2.26	46410
30	T21d	40	his-suc-tw	6	5.08	92.63	2.29	43210
10	T21d	40	his-suc-tw	6	1.95	95.68	2.37	46539
6.6	T21d	40	PBS	7	4.71	92.43	2.86	68811

Buffer key (all buffers contain 0.02% sodium azide to prevent microbial growth):

**ace** = 15 mM acetate pH 5; **his** = 15 mM histidine pH 6; **phos** = 15 mM phosphate pH

7

**ace-suc-tw** = 30 mM acetate, 80 mg/ml sucrose, 0.02% Tw80

**his-suc-tw** = 30 mM histidine, 80 mg/ml sucrose, 0.02% Tw80

**PBS** = phosphate buffered saline

Table 64. Storage Stability at 5°C of H1a11.1.A6-SL-Av at Different Concentrations and In Different Buffers, Excipients, and pHs

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
---	pre-dialysis	---	---	---	1.19	97.75	1.06	58988
50, 30, 10	T0	---	ace	5	1.23	97.85	0.92	47138
50, 30, 10	T0	---	his	6	1.30	97.84	0.87	44711
3.4	T0	---	phos	7	0.86	98.00	1.14	57373
50, 30, 10	T0	---	ace-suc-tw	5	1.30	97.85	0.85	52129
50, 30, 10	T0	---	his-suc-tw	6	1.28	97.85	0.87	47563
6.6	T0	---	PBS	7	1.35	97.76	0.89	71146
50	T7d	5	ace	5	3.52	95.40	1.08	46507
30	T7d	5	ace	5	2.35	96.55	1.10	40267
10	T7d	5	ace	5	1.46	97.48	1.07	47761
50	T7d	5	his	6	3.48	95.24	1.27	46306
30	T7d	5	his	6	3.63	95.34	1.03	46581
10	T7d	5	his	6	2.01	97.01	0.98	45843
3.4	T7d	5	phos	7	1.42	97.41	1.17	52150
50	T7d	5	ace-suc-tw	5	3.97	94.99	1.04	53439
30	T7d	5	ace-suc-tw	5	2.65	96.35	1.00	52119
10	T7d	5	ace-suc-tw	5	1.60	97.23	1.17	52681

Protein Concentration (mg/ml)	Time	temp (°C)	buffer	pH	% Aggregate	% Monomer	% Fragment	Total Area
50	T7d	5	his-suc-tw	6	4.81	94.10	1.09	48136
30	T7d	5	his-suc-tw	6	3.38	95.49	1.13	48266
10	T7d	5	his-suc-tw	6	1.88	96.90	1.22	47966
6.6	T7d	5	PBS	7	2.25	96.78	0.97	68563
50	T21d	5	ace	5	7.82	92.06	0.11	47402
30	T21d	5	ace	5	4.22	95.68	0.10	45736
10	T21d	5	ace	5	1.77	98.13	0.09	47900
50	T21d	5	his	6	7.36	92.48	0.17	47111
30	T21d	5	his	6	6.51	93.34	0.15	43911
10	T21d	5	his	6	2.92	96.98	0.10	45446
3.4	T21d	5	phos	7	0.05	99.53	0.42	45856
50	T21d	5	ace-suc-tw	5	9.13	90.73	0.14	52424
30	T21d	5	ace-suc-tw	5	4.79	95.07	0.15	51575
10	T21d	5	ace-suc-tw	5	1.98	97.87	0.15	51870
50	T21d	5	his-suc-tw	6	9.50	90.30	0.20	46588
30	T21d	5	his-suc-tw	6	6.20	93.64	0.16	46023
10	T21d	5	his-suc-tw	6	2.61	97.27	0.12	47285
6.6	T21d	5	PBS	7	3.12	96.76	0.13	62856

Buffer key (all buffers contain 0.02% sodium azide to prevent microbial growth):

**ace** = 15 mM acetate pH 5; **his** = 15 mM histidine pH 6; **phos** = 15 mM phosphate pH

7

**ace-suc-tw** = 30 mM acetate, 80 mg/ml sucrose, 0.02% Tw80

**his-suc-tw** = 30 mM histidine, 80 mg/ml sucrose, 0.02% Tw80

**PBS** = phosphate buffered saline

Overall, the data in Tables 61 to 64 suggest the two DVD-Ig proteins have an atypical degradation profile not observed for stable monoclonal antibodies. The aggregation rate is actually greater at 5°C than at 40°C. At 5°C, there is a rapid increase in aggregation after 21 days of storage. At 40°C, the amount of degradation also increases, but not at the rate observed at 5°C. In both cases, the aggregation is concentration dependent. Overall, these DVD-Ig proteins did not qualify as AS-DVD-Ig proteins based on their 5°C instability and would be considered non-aqueous stable.

## Sequences

Amino acid sequences of the heavy and light chains for other DVD-Ig proteins described herein are provided below in Table 65. Linker sequences are underlined, while

CDR sequences are in bold. “L” below represents the light chain, and “H” below represents the heavy chain.

Table 65: DVD-Ig protein sequences

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			123456789 <b>0</b> 123456789 <b>0</b> 123456789 <b>0</b> 1234567890
DVD005H CD20/CD80	AB001VH	AB007VH	<p>QVQLQQPGAEVLVKGASVVKMSCKASGYTFT<b>SYNMH</b>WVKQT      PGRGLEWIG<b>AIYPGNGDTSYNQKFKG</b>KATLTADKSSSTAY      MQLSSLTSEDAVYYCAR<b>STYYGGDWYFNV</b>WGAGTTVTVS  <b>A</b><u><b>STKGP</b></u>QVQLQESGPGLVKPSETLSLTCAVSGGSIS<b>GGY</b>  <b>GWG</b>WIRQPPGKGLEWIG<b>SFYSSSGNTYYNP</b>SLKSQVTIST      DTSKNQFSLKLNSTMADTAVYYCVRDRLF<b>SVVGMVYNNW</b>  <b>FDV</b>WGPGVLTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS      WNSGALTSGVHTFPAVLQSSGLYSLSSVTPSSSLGTQT      YICNVNHPKNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG      PSVFLFPPPKDTLmisRTPEVTCVVVDVSHEDPEVKFNW      YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK      EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE      MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV      LDSDGSFFLYSKLTVDKSRWQQGNVFSCSVHEALHNHT      QKSLSLSPGK (<b>SEQ ID NO: 30</b>)</p>
DVD005L CD20/CD80	AB001VL	AB007VL	<p>QIVLSQSPAILSPSPGEKVTMTC<b>RASSSVSYI</b>HWFQQKPG      SSPKPW<b>IYATSNILASGVPV</b>RFSGSGSGTSYSLTISRVEAE      DAATYYC<b>QQWTSNPPT</b>FGGGTKLEIKR<b>TVAAP</b>ESALTQPP      SVSGAPGQKVТИС<b>TGSTSNI</b>GGYDLH<b>WYQQLPGTAPKLL</b>  <b>IYDINKRPSGI</b>SDRFSGSKSGTAASLAITGLQTEDEADYY  <b>CQSYDSSLNAQV</b>FGGGTRLTVLG</p> <p>TVAAPSFIGFPPSDEQLKSGTASVVCLNNFYPREAKVQW      KVDNALQSGNSQESVTEQDSKDSTYSLSTTLKADYEK      HKVYACEVTHQGLSSPVTKSFNRGEC (<b>SEQ ID NO: 31</b>)</p>
DVD006H CD80/CD20	AB007VH	AB001VH	<p>QVQLQESGPGLVKPSETLSLTCAVSGGSIS<b>GGY</b><b>GWG</b>WIRQ      PPGKGLEWIG<b>SFYSSSGNTYYNP</b>SLKSQVTISTDTSKNQF      SLKLNSTMADTAVYYCVRDRLF<b>SVVGMVYNNW</b><b>FDV</b>WGPG      VLTVSS<b>A</b><u><b>STKGP</b></u>QVQLQQPGAEVLVKGASVVKMSCKASGY      TFT<b>SYNMH</b>WVKQT<sup>PGRGLEWIG</sup><b>AIYPGNGDTSYNQKFKG</b>      ATL TADKSSSTAYMQLSSLTSEDAVYYCAR<b>STYYGGDWY</b>  <b>FNV</b>WGAGTTVTVSA</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS      WNSGALTSGVHTFPAVLQSSGLYSLSSVTPSSSLGTQT      YICNVNHPKNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG      PSVFLFPPPKDTLmisRTPEVTCVVVDVSHEDPEVKFNW      YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK      EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE      MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV      LDSDGSFFLYSKLTVDKSRWQQGNVFSCSVHEALHNHT      QKSLSLSPGK (<b>SEQ ID NO: 32</b>)</p>
DVD006L	AB007VL	AB001VL	ESALTQPPSVSGAPGQKVТИС <b>TGSTSNI</b> GGYDLH <b>WYQQL</b> PGTAPKLLIY <b>DINKRPSGI</b> SDRFSGSKSGTAASLAITGLQ

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
CD80/CD20			<p>TEDEADYYC<b>QSYDSSLNAQVFGGGTRLTVLGQPKAAPQIV</b>  <b>LSQSPAILSPSPGEKVTMTCRASSSVSYIHWFQQKPGSSP</b>  <b>KPWIYATSNLASGVPVRSGSGSGTSYSLTISRVEAEDAA</b>  <b>TYYC<b>QOWTSNPPTFGGGTKLEIKR</b></b></p> <p><b>QPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVA</b>  <b>WKADSSPVKAGVETTPSKQSNNKYAASSYLSLTPEQWKS</b>  <b>HRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 33)</b></p>
DVD037H VEGF/HER2	AB014VH	AB004VH	<p>EVQLVESGGGLVQPGGSLRLSCAASGYTFT<b>NYGMN</b>WVRQA  <b>PGKGLEWVGWINTYTGEPTYAADFKRRTFSLDTSKSTAY</b>  <b>LQMNLSRAEDTAVYYCAKYPHYYGSSHWFYFDVWGQGTLVT</b>  <b>VSS<b>ASTKGPEVQLVESGGGLVQPGGSLRLSCAASGFNIKD</b></b>  <b>TYIHWRQAPGKGLEWVARIYPTNGYTRYADSVKGRFTIS</b>  <b>ADTSKNTAYLQMNLSRAEDTAVYYCSR<b>WGGDGFYAMDY</b>WG</b>  <b>QGTLTVVSS</b></p> <p><b>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS</b>  <b>WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT</b>  <b>YICNVNHPNSNTKVDKKVEPKSCDKHTCPCPAPEAAGG</b>  <b>PSVFLFPPPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW</b>  <b>YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK</b>  <b>EYKCKVSNKALPAPIEKTISAKGQPQPREPVYTLPPSREE</b>  <b>MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV</b>  <b>LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHT</b>  <b>QKSLSLSPGK (SEQ ID NO: 34)</b></p>
DVD037L VEGF/HER2	AB014VL	AB004VL	<p>DIQMTQSPSSLSASVGDRVTITC<b>SASQDISNYLN</b>WYQQKP  <b>GTKVLIYFTSSLHSGVPSRFSGSGSGTDFTLTISSLQP</b>  <b>EDFATYYC<b>QQYSTVPWTFGQGTKVEIKRTVAAP</b>DIQMTQS</b>  <b>PSSLSASVGDRVTITC<b>RASQDVNTAVA</b>WYQQKPGKAPKLL</b>  <b>IYS<b>SASFYSGVPSRFSGSRSGTDFTLTISSLQPEDFATYY</b></b>  <b>CQQHYTTPPTFGQGTKVEIKR</b></p> <p><b>TVAAPSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQW</b>  <b>KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTSKADYEK</b>  <b>HKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 35)</b></p>
DVD038H Her2/VEGF	AB004VH	AB014VH	<p>EVQLVESGGGLVQPGGSLRLSCAASGFNIKD<b>TYIHWRQ</b>A  <b>PGKGLEWVARIYPTNGYTRYADSVKGRFTISADTSKNTAY</b>  <b>LQMNLSRAEDTAVYYCSR<b>WGGDGFYAMDY</b>WGQGTLTVSS</b>  <b><b>ASTKGPEVQLVESGGGLVQPGGSLRLSCAASGYTFT<b>NYGM</b></b>  <b>WVWRQAPGKGLEWVG<b>WINTYTGEPTYAADFKRRTFSLDT</b></b>  <b>SKSTAYLQMNLSRAEDTAVYYCAKYPHYYGSSHWFYFDVWG</b>  <b>QGTLTVVSS</b></b></p> <p><b>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS</b>  <b>WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT</b>  <b>YICNVNHPNSNTKVDKKVEPKSCDKHTCPCPAPEAAGG</b>  <b>PSVFLFPPPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW</b>  <b>YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK</b>  <b>EYKCKVSNKALPAPIEKTISAKGQPQPREPVYTLPPSREE</b>  <b>MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV</b>  <b>LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHT</b>  <b>QKSLSLSPGK (SEQ ID NO: 36)</b></p>
DVD038L	AB004VL	AB014VL	DIQMTQSPSSLSASVGDRVTITC <b>RASQDVNTAVA</b> WYQQKP

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
Her2/VEGF			<p>GKAPKLLIY<b>SASF</b>FLYSGVPSRFSGSRSGTDFTLTISLQP      EDFATYYC<b>QQ</b>HYTTPPTFGQGTVKVEIKR<b>TVAAP</b>DIQMTQS      PSSLSASVGDRVITIC<b>SASQD</b>ISNYLNWYQQKPGKAPKVL      IY<b>FTSS</b>LHSGVPSRFSGSGSGTDFTLTISLQPEDFATYY  <b>CQQY</b>STVWPWTFGQGTVKVEIKR</p> <p>TVAAPSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQW      KVDNALQSGNSQESVTEQDSKDSTYSLSSTLSKADYEK      HKVYACEVTHQGLSSPVTKSFRGEC (<b>SEQ ID NO:37</b>)</p>
DVD053H TNF/RANKL	AB017VH	AB018VH	<p>EVQLVESGGGLVQPGRSLRLSCAASGFTFDD<b>DYAMH</b>WVRQA      PGKGLEWVS<b>AIT</b>WNSGHIDYADSV<b>EGRFT</b>ISRDNAKNSLY      LQMNSLRAEDTAVYYCAK<b>VSYL</b>STASSLDYWGQGTLVTVS  <b>SASTKGP</b>EVQLLESGGGLVQPGGSLRLSCAASGFTFSS<b>SYA</b>  <b>MS</b>WVRQAPGKGLEWVS<b>GIT</b>GGGSTYYADSV<b>KGRFT</b>ISRD      NSKNTLYLQMNSLRAEDTAVYYCAK<b>DPGTTVIMSWFDP</b>WG      QGTLTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS      WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT      YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG      PSVFLFPPPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNW      YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK      EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE      MTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTPPV      LDSDGSSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHT      QKSLSLSPGK (<b>SEQ ID NO:38</b>)</p>
DVD053L TNF/RANKL	AB017VL	AB018VL	<p>DIQMTQSPSSLSASVGDRVITIC<b>RASQGIRNYL</b>AWYQQK      GKAPKLLIY<b>AASTLQSGVPS</b>RFSGSGSGTDFTLTISLQP      EDVATYYC<b>QRYNRA</b>PTFGQGTVKVEIKR<b>TVAAP</b>EIVLTQS      PGTLSLSPGERATLSCRAS<b>QSVRGRY</b>LAWYQQKPGQAPRL      LIY<b>GASSRATGIPDR</b>FGSGSGTDFLTISRLEPEDFAVF      YC<b>QQY</b>GSSPRTFGQGTVKVEIKR</p> <p>TVAAPSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQW      KVDNALQSGNSQESVTEQDSKDSTYSLSSTLSKADYEK      HKVYACEVTHQGLSSPVTKSFRGEC (<b>SEQ ID NO:39</b>)</p>
DVD054H RANKL/TNF	AB018VH	AB017VH	<p>EVQLLESGGGLVQPGGSLRLSCAASGFTFSS<b>SYAMS</b>WVRQA      PGKGLEWVS<b>GIT</b>GGGSTYYADSV<b>KGRFT</b>ISRDNSKNTLY      LQMNSLRAEDTAVYYCAK<b>DPGTTVIMSWFDP</b>WGQGTLTV  <b>SSASTKGP</b>EVQLVESGGGLVQPGRSLRLSCAASGFTFDD<b>Y</b>  <b>AMH</b>WVRQAPGKGLEWVS<b>AIT</b>WNSGHIDYADSV<b>EGRFT</b>ISR      DNAKNSLYLQMNSLRAEDTAVYYCAK<b>VSYL</b>STASSLDYWG      QGTLTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS      WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT      YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG      PSVFLFPPPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNW      YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK      EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE      MTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTPPV      LDSDGSSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHT      QKSLSLSPGK (<b>SEQ ID NO:40</b>)</p>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
DVD054L RANKL/TNF	AB018VL	AB017VL	<p>EIVLTQSPGTLSPGERATLSCR<b>RASQSVRGRYLA</b>WYQQKPGQAPRLLIY<b>GASSRATGIPD</b>RFSGSGSGTDFLTISRLEPEDFAVFYC<b>QQYQGSSPRT</b>FGQGTKVEIKR<b>TVAAP</b>DIQMTQSPSSLASAVGDRVITIC<b>RASQGIRNYLA</b>WYQQKPGKAPKL LIY<b>AASTLQSGVP</b>SRFSGSGSGTDFLTISLQPEDVATY<b>YCQRYNRAPYT</b>FGQGTKVEIKR</p> <p>TVAAPSFIGPPSDEQLKSGTASVVCCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTKADYEKHKVYACEVTHQGLSSPVTKSFNRCG (SEQ ID NO: 41)</p>
DVD065H TNF/ DKK	AB017VH	AB023VH	<p>EVQLVESGGGLVQPGRSLRLSCAASGFTFD<b>DYAMH</b>WVRQAPGKGLEW<b>VAITWNSGHIDYAD</b>SVEGRTFTISRDNAKNSLYLQMNSLRAEDTAVYYCAK<b>VSYLSTASSLDY</b>WGQGTLVTVSS<b>SASTKGP</b>EVQLVESGGGLVQPGANSLKLSCAASGFTFSD<b>YAM</b>MAWVRQSPKKGLEW<b>VAIIYDGSSYYRDSVKGRFTISRD</b>NAKSTLYLQMDSLRSEDTATYYCAT<b>GLGIATDYFDY</b>WGQGVLTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTPSSSLGTQTYICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGGPSVFLFPPPKDTLmisRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNFSCSVMHEALHNHTQKSLSLSPGK (SEQ ID NO: 42)</p>
DVD065L TNF/ DKK	AB017VL	AB023VL	<p>DIQMTQSPSSLSASVGDRVITIC<b>RASQGIRNYLA</b>WYQQKPGKAPKL LIY<b>AASTLQSGVP</b>SRFSGSGSGTDFLTISLQPEDVATY<b>YCQRYNRAPYT</b>FGQGTKVEIKR<b>TVAAP</b>DIRMTQSPASLSASLGETVNI<b>ECLASEDIYSD</b>WYQQKPGKSPQLLIYNANSLQNGVP<b>SRFSGSGSGTQYS</b>LSKINSLQSEDVATYFC<b>QQYNNYPPT</b>FGGGTKELEK</p> <p>TVAAPSFIGPPSDEQLKSGTASVVCCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTKADYEKHKVYACEVTHQGLSSPVTKSFNRCG (SEQ ID NO: 43)</p>
DVD066H DKK/ TNF	AB023VH	AB017VH	<p>EVQLVESGGGLVQPGRSLRLSCAASGFTFSD<b>YAM</b>WVRQSPKKGLEW<b>VAIIYDGSSYYRDSVKGRFTISRD</b>NAKSTLYLQMDSLRSEDTATYYCAT<b>GLGIATDYFDY</b>WGQGVLTVSS<b>ASTKGP</b>EVQLVESGGGLVQPGRSLRLSCAASGFTFD<b>DYAM</b>WVRQAPGKGLEW<b>VAITWNSGHIDYAD</b>SVEGRTFTISRDNAKNSLYLQMNSLRAEDTAVYYCAK<b>VSYLSTASSLDY</b>WGQGVLTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTPSSSLGTQTYICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPE<b>AAGG</b>PSVFLFPPPKDTLmisRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNFSCSVMHEALHNHTQKSLSLSPGK (SEQ ID NO: 44)</p>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
DVD066L DKK/TNF	AB023VL	AB017VL	<p>QKSLSLSPGK (<b>SEQ ID NO: 44</b>)</p> <p>DIRMTQSPASLSASLGETVNIECLASEDIYSDLAWYQQKP GKSPQLLIY<b>NANSI</b>QNGVPSRFSGSGSGTQYSLKINSLQS EDVATYFC<b>QOYNN</b>PPTFGGGTKLELKR<b>TVAAP</b>DIQMTQS PSSLSASAVGDRVТИTCRAS<b>QGIRNYL</b>AWYQQKPGKAPKLL IYAASTL<b>QSGVPS</b>RFSGSGSGTDFTLTISSLQPEDVATYY <b>CQRYNRAPY</b>TFGQGTKVEIKR</p> <p>TVAAPSFIGFPPSDEQLKSGTASVVCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYLSSTLTSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC (<b>SEQ ID NO: 45</b>)</p>
DVD165H CD20/RANKL	AB001VH	AB018VH	<p>QVQLQQPGAEVLVPGASVMSCKASGYTFT<b>SYNMH</b>WVKQT PGRGLEWIG<b>AIY</b>PGNGDTS<b>YNQ</b>KFKGKATLTADKSSSTAY MQLSSLTSEDSAVYYCAR<b>STYYGGDWYFNV</b>WGAGTTVTVS <b>A</b><b>ASTKGP</b>EVQVLLSFGGGLVQPGGSLRLSCAASGFTFS<b>SYA</b> <b>MS</b>WVRQAPGKGLEWVSG<b>GITGSGG</b>STYY<b>ADSVKG</b>RFTISRD NSKNTLYL<b>Q</b>MNSLRAEDTAVYYCAK<b>DPGTTVIM</b>S<b>WFDP</b>WG QGTLTVVSS</p> <p>ASTKGPSVFLAPSSKSTSGGTAAALGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT YICNVNHPNSNTKVDKVEPKSCDKTHTCPCPAPEAAGG PSVFLFPPKPKDLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPPV LSDGSFFLYSKLTVDKSRWQQGNVFS<b>SCVM</b>HEALHNHT QKSLSLSPGK (<b>SEQ ID NO: 46</b>)</p>
DVD165L CD20/RANKL	AB001VL	AB018VL	<p>QIVLSQSPAILSPSPGEKVTMTCRASSSVSY<b>I</b>HWFQQKPG SSPKPWIY<b>ATSNL</b>ASGPVRFSGSGSGT<b>SY</b>SLTISRVEAE DAATYYC<b>QO</b>WTSPNPTFGGGTKLEIKR<b>TVAAP</b>EIVLTQSP GTLSLSPGERATLSC<b>RASQ</b>SVRGRY<b>I</b>AWYQQKPGQAPRLL <b>IY</b><b>GASSRATGIPD</b>RFSGSGSGTDFTLTISRLEPEDFAVFY <b>CQY</b>QGSSPRTFGQGTKVEIKR</p> <p>TVAAPSFIGFPPSDEQLKSGTASVVCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYLSSTLTSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC (<b>SEQ ID NO: 47</b>)</p>
DVD166H RANKL/CD20	AB018VH	AB001VH	<p>EVQLLESFGGGLVQPGGSLRLSCAASGFTFS<b>SYA</b>MSWVRQA PGKGLEWVSG<b>GITGSGG</b>STYY<b>ADSVKG</b>RFTISRDNSKNTLY L<b>Q</b>MNSLRAEDTAVYYCAK<b>DPGTTVIM</b>S<b>WFDP</b>WGQGTLTV <b>SS</b><b>A</b><b>STKGP</b>QVQLQQPGAEVLVPGASVMSCKASGYTFT<b>SY</b> <b>NM</b>WVKQTGGRGLEWIG<b>AIY</b>PGNGDTS<b>YNQ</b>KFKGKATLT DKSSSTAYMQLSSLTSEDSAVYYCAR<b>STYYGGDWYFNV</b>WG AGTTTVVSA</p> <p>ASTKGPSVFLAPSSKSTSGGTAAALGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT YICNVNHPNSNTKVDKVEPKSCDKTHTCPCPAPEAAGG PSVFLFPPKPKDLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPPV</p>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
			LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK ( <b>SEQ ID NO:48</b> )
DVD166L RANKL/CD20	AB018VL	AB001VL	<p>EIVLTQSPGTLSSLPGERATLSC<b>RASQSVRGRYLA</b>WYQQKPGQAPRLLIY<b>GASSRATGIP</b>DRSGSGSGTDFLTISRLPEDFAFYC<b>QQYGSSPRT</b>FGQGKVEIKR<b>TVAAP</b>QIVLSQSPAILSPSPGEKVMT<b>CRASSSVSYI</b>HWFQQKPGSSPKPW<b>IYATSNLASGVPVRF</b>SGSGSGTSYSLTISRVEAEDAATYY<b>CQQWTSNPPT</b>FGGGTKEIKR</p> <p>TVAAPSFIGFPPSDEQLKSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTSKADYEKHKVYACEVTHQGLSSPVTKSFRGEC (<b>SEQ ID NO:49</b>)</p>
DVD257H DLL4/PLGF	DLL4H	PLGFH	<p>EVQLVESGGGLVQPGGSLRLSCAASGFTFT<b>DNWISWVR</b>QAPGKGLEWVG<b>YISPNSGFTYYADSVKGRFT</b>ISADTSKNTAYLQMNSLRAEDTAVYYCARD<b>DNFGGYFDY</b>WGQGTLTVSS<b>AS</b> <b>TKGP</b>QVQLQQSGAELVKPGASVKISCKASGYTF<b>DYYIN</b>WVKLAPGQGLEWIG<b>WIYPGSGNTKYNEKF</b>KGATLTIDTSSSTAYMQLSSLTSEDTAVYFCVRD<b>SPFFDY</b>WGQGTLTVSS (<b>SEQ ID NO:50</b>)</p>
DVD257L DLL4/PLGF	DLL4L	PLGFL	<p>DIQMTQSPSSLSASVGDRVITIC<b>RASQDVSTAVA</b>WYQQPKGKAPKLLIY<b>SASFYSGVPSRF</b>SGSGSGTDFLTISLQPEDFATTYYC<b>QQSYTGT</b>VTFGQGKVEIKR<b>TVAAP</b>DIVLTQSPDSLAVSLGERVTMNC<b>KSSQSLNSGMRKSFLA</b>WYQQKPGQSPKLLIY<b>WASTRESGV</b>PDRTGSGSGTDFLTISLQPEDFATTYYC<b>QQSYTGT</b>VTFGQGKVEIKR</p> <p>(<b>SEQ ID NO:51</b>)</p>
DVD258H PLGF/DLL4	PLGFH	DLL4H	<p>QVQLQQSGAELVKPGASVKISCKASGYTF<b>DYYIN</b>WVKLAPGQGLEWIG<b>WIYPGSGNTKYNEKF</b>KGATLTIDTSSSTAYMQLSSLTSEDTAVYFCVRD<b>SPFFDY</b>WGQGTLTVSS<b>ASTK</b> <b>GP</b>EVQLVESGGGLVQPGGSLRLSCAASGFTFT<b>DNWISWVR</b>QAPGKGLEWVG<b>YISPNSGFTYYADSVKGRFT</b>ISADTSKNTAYLQMNSLRAEDTAVYYCARD<b>DNFGGYFDY</b>WGQGTLTVSS (<b>SEQ ID NO:52</b>)</p>
DVD258L PLGF/DLL4	PLGFL	DLL4L	<p>DIVLTQSPDSLAVSLGERVTMNC<b>KSSQSLNSGMRKSFLA</b>WYQQKPGQSPKLLIY<b>WASTRESGV</b>PDRTGSGSGTDFLTISLQPEDFATTYYC<b>QQSYTGT</b>VTFGQGKVEIKR<b>TVAAP</b>DIQMTQSPSSLSASVGDRVITIC<b>RASQDVSTAVA</b>WYQQKPGKAPKLLIY<b>SASFYSGVPSRF</b>SGSGSGTDFLTISLQPEDFATTYYC<b>QQSYTGT</b>VTFGQGKVEIKR</p> <p>(<b>SEQ ID NO:53</b>)</p>
DVD277H TNF/SOST (S2)	AB017VH	AB050VH	<p>EVQLVESGGGLVQPGRSLRLSCAASGFTFD<b>DYAMH</b>WVRQAPGKGLEWVS<b>AITWNSGHIDYAD</b>SVEGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAK<b>VSYLSTASSLDY</b>WGQGTLTVSS<b>ASTKGP</b>EVQLQQSGPELMKPGASVKMSCKASGYTF<b>DYN</b></p>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			<b>MH</b> WMKQNQGKSLEWIGE <b>INPNSGGSGYNQKFKGKATLTV</b> D <b>KSSSTAYMELRSLTSEDSA</b> YYCAR <b>LGYYGNYEDWYFDVW</b> <b>GAGTTTVSS</b>  <b>ASTKGPSVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVS</b> <b>WNSGALTSGVHTFPAVI</b> QSSGLYSLSSVTVPSSSLGTQT <b>YICNVNHPNSNTKVDKVK</b> EPKSCDKTHTCPPCPAPEAAGG <b>PSVFLFPPKPKDTLMISRTPEV</b> TCVVVDVSHEDPEVKFNW <b>YVDGVEVHN</b> AKTKPREEQYNSTYRVSVLTVLHQDWLNGK <b>EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREE</b> <b>MTKNQVS</b> LTCLVKGFYPSDIAVEWESNGQPENNYKTPPV <b>LSDGSFFLYSKLTVDKSRWQQGNF</b> SCSVMHEALHNHYT <b>QKSLSLSPGK (SEQ ID NO: 54)</b>
DVD277L TNF / SOST (S2)	AB017VL	AB050VL	<b>DIQMTQSPSSLSASVGDRT</b> IT <b>C</b> RASQ <b>GIRNYLA</b> WYQQKP <b> GKAPKLLI</b> Y <b>A</b> ASTL <b>QSGVP</b> S <b>R</b> FSGSGSGTDFTLTISSLQP <b>EDVATYYC</b> Q <b>R</b> YN <b>R</b> APY <b>T</b> FG <b>Q</b> G <b>T</b> K <b>V</b> E <b>I</b> K <b>R</b> <b>T</b> VAAP <b>D</b> L <b>Q</b> MT <b>Q</b> T <b>TSSLSASLGDRT</b> IT <b>C</b> RASQ <b>D</b> IS <b>NYLN</b> WYQQKPDGT <b>V</b> K <b>L</b> <b>I</b> F <b>Y</b> T <b>STLQSGVP</b> S <b>R</b> FSGSGSGT <b>N</b> YSLT <b>I</b> T <b>N</b> LE <b>Q</b> DDA <b>A</b> TY <b>F</b> <b>C</b> Q <b>G</b> DT <b>L</b> P <b>Y</b> TFGG <b>G</b> T <b>K</b> LE <b>I</b> K <b>R</b>  <b>T</b> VAAP <b>S</b> VF <b>I</b> F <b>PPS</b> DE <b>Q</b> LK <b>S</b> GT <b>A</b> SV <b>V</b> CL <b>L</b> NN <b>F</b> P <b>E</b> AK <b>Q</b> W <b>K</b> VD <b>N</b> AL <b>Q</b> SG <b>N</b> Q <b>E</b> SV <b>T</b> EQ <b>D</b> SK <b>D</b> ST <b>Y</b> SL <b>S</b> ST <b>L</b> TS <b>K</b> AD <b>Y</b> <b>E</b> K <b>V</b> Y <b>A</b> CE <b>V</b> TH <b>Q</b> GL <b>S</b> SP <b>V</b> TK <b>S</b> FN <b>R</b> GE <b>C</b> (SEQ ID NO: 55)
DVD278H SOST (S2) / TNF	AB050VH	AB017VH	<b>EVQLQQSGPELMKPGASV</b> KMSCKASGY <b>T</b> FT <b>D</b> YN <b>MH</b> WMKQN <b>QGKSLEWIGE</b> <b>INPNSGGSGYNQKFKGKATLTV</b> D <b>KSSSTAY</b> <b>MELRSLTSEDSA</b> YYCAR <b>LGYYGNYEDWYFDV</b> WGAGTTV <b>VSS</b> <b>ASTKG</b> P <b>E</b> QL <b>V</b> ES <b>GG</b> GL <b>V</b> Q <b>P</b> GR <b>S</b> LR <b>L</b> SCA <b>A</b> SG <b>F</b> T <b>DD</b> <b>YAMH</b> W <b>V</b> R <b>Q</b> AP <b>G</b> K <b>GLE</b> W <b>V</b> <b>S</b> <b>A</b> <b>I</b> <b>T</b> <b>W</b> <b>N</b> <b>S</b> <b>G</b> <b>H</b> <b>I</b> <b>D</b> <b>Y</b> <b>A</b> <b>D</b> <b>S</b> <b>V</b> <b>E</b> <b>G</b> <b>R</b> <b>F</b> <b>T</b> <b>IS</b> <b>RDNAKNS</b> LY <b>L</b> Q <b>M</b> NS <b>L</b> R <b>A</b> ED <b>T</b> <b>A</b> <b>V</b> <b>Y</b> <b>C</b> <b>A</b> <b>K</b> <b>V</b> <b>S</b> <b>Y</b> <b>L</b> <b>S</b> <b>T</b> <b>A</b> <b>S</b> <b>L</b> <b>D</b> <b>Y</b> <b>W</b> <b>Q</b> <b>G</b> <b>T</b> <b>L</b> <b>V</b> <b>T</b> <b>V</b> <b>S</b>  <b>ASTKGPSVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVS</b> <b>WNSGALTSGVHTFPAVI</b> QSSGLYSLSSVTVPSSSLGTQT <b>YICNVNHPNSNTKVDKVK</b> EPKSCDKTHTCPPCPAPEAAGG <b>PSVFLFPPKPKDTLMISRTPEV</b> TCVVVDVSHEDPEVKFNW <b>YVDGVEVHN</b> AKTKPREEQYNSTYRVSVLTVLHQDWLNGK <b>EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREE</b> <b>MTKNQVS</b> LTCLVKGFYPSDIAVEWESNGQPENNYKTPPV <b>LSDGSFFLYSKLTVDKSRWQQGNF</b> SCSVMHEALHNHYT <b>QKSLSLSPGK (SEQ ID NO: 56)</b>
DVD278L SOST (S2) / TNF	AB050VL	AB017VL	<b>DLQMTQTTSSLSASLGDRT</b> IT <b>C</b> RASQ <b>D</b> IS <b>NYLN</b> WYQQKP <b>DGT</b> <b>V</b> <b>K</b> <b>L</b> <b>I</b> <b>F</b> <b>Y</b> <b>T</b> <b>STLQSGVP</b> S <b>R</b> FSGSGSGT <b>N</b> YSLT <b>I</b> T <b>N</b> LE <b>Q</b> <b>DDA</b> <b>A</b> <b>T</b> <b>Y</b> <b>F</b> <b>C</b> <b>Q</b> <b>G</b> <b>D</b> <b>T</b> <b>L</b> <b>P</b> <b>Y</b> TFGG <b>G</b> T <b>K</b> LE <b>I</b> K <b>R</b> <b>T</b> VAAP <b>D</b> I <b>Q</b> MT <b>Q</b> S <b>PSSLSASVGDRT</b> IT <b>C</b> RASQ <b>GIRNYLA</b> WYQQKPG <b>K</b> <b>A</b> <b>P</b> <b>K</b> <b>L</b> <b>I</b> <b>Y</b> <b>A</b> <b>STLQSGVP</b> S <b>R</b> FSGSGSGTDFTLTISSLQP <b>E</b> <b>D</b> <b>V</b> <b>A</b> <b>T</b> <b>Y</b> <b>C</b> <b>Q</b> <b>R</b> <b>Y</b> <b>N</b> <b>R</b> <b>A</b> <b>P</b> <b>Y</b> TF <b>Q</b> <b>G</b> <b>T</b> <b>K</b> <b>V</b> <b>E</b> <b>I</b> <b>R</b>  <b>T</b> VAAP <b>S</b> VF <b>I</b> F <b>PPS</b> DE <b>Q</b> LK <b>S</b> GT <b>A</b> SV <b>V</b> CL <b>L</b> NN <b>F</b> P <b>E</b> AK <b>Q</b> W <b>K</b> VD <b>N</b> AL <b>Q</b> SG <b>N</b> Q <b>E</b> SV <b>T</b> EQ <b>D</b> SK <b>D</b> ST <b>Y</b> SL <b>S</b> ST <b>L</b> TS <b>K</b> AD <b>Y</b> <b>E</b> K <b>V</b> Y <b>A</b> CE <b>V</b> TH <b>Q</b> GL <b>S</b> SP <b>V</b> TK <b>S</b> FN <b>R</b> GE <b>C</b> (SEQ ID NO: 57)
DVD281H IL-	AB056VH	AB053VH	<b>QVQLQQSGAELM</b> KPGASV <b>K</b> LS <b>C</b> AT <b>G</b> Y <b>T</b> FT <b>G</b> <b>SWIE</b> W <b>I</b> K <b>Q</b> R <b>PGHGLEWIG</b> Q <b>I</b> L <b>P</b> GS <b>G</b> S <b>A</b> <b>Y</b> <b>Y</b> <b>N</b> <b>E</b> <b>K</b> <b>F</b> <b>K</b> <b>G</b> <b>K</b> <b>A</b> <b>F</b> <b>T</b> <b>A</b> <b>D</b> <b>T</b> <b>S</b> <b>S</b> <b>K</b> <b>T</b> <b>V</b> <b>Y</b> <b>I</b> <b>O</b> <b>L</b> <b>I</b> <b>S</b> <b>L</b> <b>T</b> <b>E</b> <b>D</b> <b>S</b> <b>A</b> <b>I</b> <b>Y</b> <b>Y</b> <b>C</b> <b>A</b> <b>R</b> <b>E</b> <b>D</b> <b>N</b> <b>Y</b> <b>G</b> <b>S</b> <b>S</b> <b>L</b> <b>A</b> <b>Y</b> <b>W</b> <b>G</b> <b>O</b> <b>G</b> <b>T</b> <b>L</b> <b>L</b> <b>T</b> <b>V</b> <b>S</b> <b>A</b>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
9 (S2) / IgE			<p><b>ASTKGPEVQLVESGGGLVQPGGSLRLSCAVSGYSITSGYS</b>  <b>WNWIRQAPGKLEWVA</b><b>SITYDGSTNYNP</b><b>SVKG</b><b>RITISRDD</b>  <b>SKNTFY</b><b>LQMNSLRAEDTAVYYC</b><b>ARGSHYFGHWF</b><b>AVWGQG</b>  <b>TLTVSS</b></p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS    WNSGALTSGVHTFPALQSSGLYSLSSVTPSSSLGTQT    YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG    PSVFLFPPPKDTLMSRTPEVTCVVVDVSHEDPEVKFNW    YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK    EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE    MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV    LDSDGSFFFLYSLTVVDKSRWQQGNVFSCSVMHEALHNHT    QKSLSLSPGK (<b>SEQ ID NO: 58</b>)</p>
DVD281L IL-9 (S2) / IgE	AB056VL	AB053VL	<p>DILLTQSPAILSVPGERVSFSC<b>RASQSIGTNIH</b><b>WYQQRT</b>    NGSPLLK<b>YASESISIGIPS</b><b>RFSGGGSGTDF</b><b>TL</b><b>SINSVES</b>    EDIADYYC<b>QQSNNWPLT</b><b>FGAGT</b><b>KLELKR</b><b>TVAAP</b><b>D</b><b>IQLTQS</b>    PSSLSASAVGDRVITIC<b>RASQSV</b><b>DYDGDSY</b><b>MN</b><b>WYQQKPGKA</b>  <b>PKL</b><b>LIYAASYLESGVPS</b><b>RFSGSGSGTDF</b><b>TL</b><b>TISSLQ</b><b>PEDF</b>    ATYYC<b>QQSHE</b><b>DPYT</b><b>FGQGT</b><b>KVEIKR</b></p> <p>TVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQW    KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLKADYEK    HKVYACEVTHQGLSSPVTKSFRGEC (<b>SEQ ID NO: 59</b>)</p>
DVD282H IgE/IL-9 (S2)	AB053VH	AB056VH	<p>EVQLVESGGGLVQPGGSLRLSCAVSGYSIT<b>SGYS</b><b>WNWIRQ</b>    APGKGLEWVA<b>SITYDGSTNYNP</b><b>SVKG</b><b>RITISRDD</b><b>SKNTFY</b>    LQMNSLRAEDTAVYYC<b>ARGSHYFGHWF</b><b>AVWGQG</b><b>TLTVS</b>  <b>SASTKG</b><b>PQVQLQQSGAELMKPGASV</b><b>KL</b><b>SCKATG</b><b>YFTFT</b><b>GSW</b>  <b>IEWIKQRP</b><b>GHGLEWIG</b><b>QILPGSGS</b><b>SAYYNE</b><b>FKG</b><b>KATFTAD</b>    TSSKTVYIQLISLTTEDSAYYCAR<b>EDNYGSSSLAY</b><b>WGQG</b>  <b>TLLTVSA</b></p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS    WNSGALTSGVHTFPALQSSGLYSLSSVTPSSSLGTQT    YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG    PSVFLFPPPKDTLMSRTPEVTCVVVDVSHEDPEVKFNW    YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK    EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE    MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV    LDSDGSFFFLYSLTVVDKSRWQQGNVFSCSVMHEALHNHT    QKSLSLSPGK (<b>SEQ ID NO: 60</b>)</p>
DVD282L IgE/IL-9 (S2)	AB053VL	AB056VL	<p>DIQLTQSPSSLSASAVGDRVITIC<b>RASQSV</b><b>DYDGDSY</b><b>MN</b><b>WY</b>    QQKPGKAP<b>KL</b><b>LIYAASYLESGVPS</b><b>RFSGSGSGTDF</b><b>TL</b><b>TIS</b>    SLQPEDFATYYC<b>QQSHE</b><b>DPYT</b><b>FGQGT</b><b>KVEIKR</b><b>TVAAP</b><b>D</b>    LTQSPAILSVPGERVSFSC<b>RASQSIGTNIH</b><b>WYQQRT</b><b>NGS</b>    PRLLK<b>YASESISIGIPS</b><b>RFSGGGSGTDF</b><b>TL</b><b>SINSVE</b><b>EDI</b>    ADYYC<b>QQSNNWPLT</b><b>FGAGT</b><b>KLELKR</b></p> <p>TVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQW    KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLKADYEK    HKVYACEVTHQGLSSPVTKSFRGEC (<b>SEQ ID NO: 61</b>)</p>
TNF/IL-17H (DVD-A)	TNFH	IL-17H	EVQLVESGGGLVQPGRSLRLSCAASGFTFD <b>DYAMH</b> <b>WVRQA</b> PGKGLEWVA <b>AITWNSGHIDYADSVEG</b> <b>RFT</b> <b>ISRDNA</b> <b>NSLY</b>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			<p>1234567890123456789012345678901234567890</p> <p>LQMNSLRAEDTAVYYCAK<b>VSYLS</b>TASSLDYWGQGTLVTVS  <u>GGGGSGGGGS</u>EVQLVQSGAEVKKPGSSVKVSCKASGGSF  <b>GGY</b>GIGWVRQAPGQGLEWMGGITPFFGFADY<b>AQK</b>FQGRVT  ITADESITAYMELSGLTSDDTAVYYCAR<b>DPNEFWNGYY</b>  <b>THDFDS</b>WGQGTTVTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS  WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT  YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPELLGG  PSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW  YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK  EYKCKVSNKALPAPIEKTI<b>SKAKGQPREPQVYTLPPSRDE</b>  LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV  LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHT  QKSLSLSPGK  <b>(SEQ ID NO: 62)</b></p>
TNF / IL-17L (DVD-A)	TNFL	IL-17L	<p>DIQMTQSPSSLSASVGDRVTITC<b>RASQ</b>GIRNYLAWYQQKP  GKAPKLLI<b>Y</b><b>A</b>STL<b>Q</b>SGVPSRFSGSGSGTDFTLTISLQP  EDVATYYC<b>Q</b>RYNRA<b>P</b>YTFGQGTKEIKR<b>GGGGSGGSE</b>I  VLTQSPDFQSVPKEKVIT<b>ICRASQD</b>IGSELH<b>WY</b>QQKPDQ  PPKLLI<b>KY</b><b>A</b>SH<b>ST</b>SGVPSRFSGSGSGTDFTLTINGLEAD  AGTYYC<b>H</b><b>Q</b>TD<b>S</b>LP<b>Y</b>TFGPGTKV<b>DI</b>KR</p> <p>TVAAPS<b>V</b>FIFPPSDEQLKSGTASVVC<b>LNNF</b>YPREAKV<b>QW</b>  KVDNALQSGNSQESVTEQDSKD<b>STY</b>SL<b>S</b>TL<b>S</b>TL<b>K</b>AD<b>Y</b>  HKVYACEVTHQGLSSPVT<b>K</b>SFNR<b>G</b>E<b>C</b> <b>(SEQ ID NO: 63)</b></p>
TNF / PGE2H (DVD-B)	TNFH	PGE2	<p>EVQLVEGGGLVQPG<b>R</b>SLRLSCAASGFTFD<b>DYAMH</b>WVRQA  PGKGLEW<b>V</b>S<b>A</b>IT<b>WNSGH</b>IDI<b>YAD</b>SVEGRFT<b>I</b>SRDN<b>A</b>KN<b>S</b>LY  LQMNSLRAEDTAVYYCAK<b>VSYLS</b>TASSLDYWGQGTLVTVS  <b>SASTKGP</b>EVQLVQSGAEVKKPGASVKVSCKASGY<b>TFT</b><b>KY</b>W  <b>LG</b>WVRQAPGQGLEWM<b>GDI</b>Y<b>PGY</b>DY<b>THY</b>NE<b>KFD</b>RV<b>T</b>LT<b>D</b>  T<b>S</b>T<b>S</b>TAYMEL<b>R</b>SL<b>R</b>S<b>DDT</b>AVYY<b>C</b>AR<b>SDGS</b><b>S</b>TY<b>W</b>GQGTLV<b>T</b>  V<b>S</b>  <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS  WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT  YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPELLGG  PSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW  YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK  EYKCKVSNKALPAPIEKTI<b>SKAKGQPREPQVYTLPPSRDE</b>  LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV  LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHT  QKSLSLSPGK  <b>(SEQ ID NO: 64)</b></p> </p>
TNF / PGE2L (DVD-B)	TNF	PGE2	<p>DIQMTQSPSSLSASVGDRVTITC<b>RASQ</b>GIRNYLAWYQQKP  GKAPKLLI<b>Y</b><b>A</b>STL<b>Q</b>SGVPSRFSGSGSGTDFTLTISLQP  EDVATYYC<b>Q</b>RYNRA<b>P</b>YTFGQGTKEIKR<b>TVAAP</b>DVLMTQT  PLSLPVTP<b>G</b>EP<b>A</b>SI<b>C</b>T<b>S</b><b>S</b><b>Q</b><b>N</b><b>I</b><b>V</b><b>H</b><b>S</b><b>N</b><b>G</b><b>N</b><b>T</b><b>Y</b><b>L</b><b>E</b><b>W</b><b>Y</b><b>L</b><b>Q</b><b>K</b><b>P</b><b>Q</b>  SPQLLI<b>Y</b><b>K</b><b>V</b><b>S</b><b>N</b><b>R</b><b>F</b><b>S</b><b>G</b><b>V</b><b>P</b><b>D</b><b>R</b><b>F</b><b>S</b><b>G</b><b>S</b><b>G</b><b>T</b><b>D</b><b>F</b><b>L</b><b>K</b><b>I</b><b>S</b><b>R</b><b>V</b><b>E</b><b>A</b>  VGYYCF<b>Q</b><b>V</b><b>S</b><b>H</b><b>V</b><b>P</b><b>Y</b><b>T</b><b>F</b><b>G</b><b>G</b><b>T</b><b>K</b><b>V</b><b>E</b><b>I</b><b>R</b></p>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
			TVAAPSVFIFPPSDEQLKSGTASVVC <sub>LLNNF</sub> YPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYSLSSTL <sub>SKADYE</sub> K HKVYACEVTHQGLSSPVTKSFNRGEC ( <b>SEQ ID NO: 65</b> )
IL-1a/IL-1bH (DVD-C)	IL-1aH	IL-1bH	EVQLVESGGVVQPGRLSRLSCSASGFI <sub>FSRYDMS</sub> WVRQA PGKGLEWVA <sub>I</sub> SHGGAGT <sub>Y</sub> Y <sub>P</sub> DSVKG <sub>R</sub> FT <sub>I</sub> SRDN <sub>S</sub> KNTLF LQMDSLRPEDTG <sub>V</sub> YFC <sub>A</sub> GGV <sub>T</sub> KG <sub>Y</sub> FD <sub>V</sub> WGQG <sub>T</sub> PTV <sub>S</sub> <u>A</u> <u>STKGP</u> QVQLVESGGVVQPGRLSRL <sub>S</sub> TASGFT <sub>F</sub> SMFGV <sub>H</sub> WVRQAPGKGLEWVA <sub>A</sub> V <sub>S</sub> Y <sub>D</sub> GSN <sub>K</sub> Y <sub>Y</sub> AE <sub>S</sub> V <sub>K</sub> G <sub>R</sub> FT <sub>I</sub> SRDN <sub>S</sub> KNILFLQMDSLRLED <sub>T</sub> AVYYC <sub>A</sub> R <sub>G</sub> PK <sub>V</sub> V <sub>I</sub> P <sub>A</sub> PLAH <sub>W</sub> GQG TLVT <sub>F</sub> S  ASTKGPSVFPLAPSSK <sub>S</sub> TSGGTAALGCLV <sub>K</sub> DYF <sub>PEP</sub> PTV <sub>S</sub> WNSGALTSGVHTFP <sub>A</sub> V <sub>L</sub> QSSGLY <sub>S</sub> LSSV <sub>V</sub> TV <sub>P</sub> SS <sub>L</sub> GT <sub>Q</sub> T YICNVN <sub>H</sub> KPSNTKVDKK <sub>V</sub> EP <sub>K</sub> SCDK <sub>T</sub> H <sub>T</sub> C <sub>PP</sub> CP <sub>A</sub> PEAAGG PSVFLF <sub>PP</sub> PK <sub>D</sub> TL <sub>M</sub> ISRTPEV <sub>T</sub> CV <sub>V</sub> VD <sub>V</sub> SHED <sub>E</sub> PEV <sub>K</sub> FN <sub>W</sub> YVDGVEVHNAK <sub>T</sub> KP <sub>R</sub> EEQ <sub>Y</sub> N <sub>ST</sub> Y <sub>R</sub> V <sub>V</sub> S <sub>V</sub> L <sub>T</sub> V <sub>L</sub> H <sub>Q</sub> D <sub>W</sub> L <sub>N</sub> G <sub>K</sub> EYKCKVSNK <sub>A</sub> L <sub>P</sub> APIE <sub>K</sub> T <sub>I</sub> S <sub>K</sub> A <sub>G</sub> Q <sub>P</sub> REP <sub>Q</sub> V <sub>Y</sub> T <sub>L</sub> PP <sub>S</sub> REE MTKNQVSLT <sub>C</sub> L <sub>V</sub> K <sub>G</sub> F <sub>Y</sub> PS <sub>D</sub> I <sub>A</sub> V <sub>E</sub> W <sub>E</sub> S <sub>N</sub> G <sub>Q</sub> P <sub>E</sub> NNY <sub>K</sub> T <sub>T</sub> PP <sub>V</sub> LSDGSFF <sub>FL</sub> Y <sub>S</sub> K <sub>L</sub> T <sub>V</sub> D <sub>K</sub> S <sub>R</sub> W <sub>Q</sub> Q <sub>G</sub> N <sub>V</sub> F <sub>S</sub> C <sub>S</sub> V <sub>M</sub> HEALHN <sub>Y</sub> T QKSLSLSPGK ( <b>SEQ ID NO: 66</b> )
IL-1a/IL-1bL (DVD-C)	IL-1aL	IL-1bL	DIQMTQSPSSLSASVGDR <sub>V</sub> T <sub>I</sub> C <sub>R</sub> AS <sub>G</sub> N <sub>I</sub> H <sub>N</sub> Y <sub>L</sub> T <sub>W</sub> Y <sub>Q</sub> Q <sub>T</sub> P GKAPK <sub>L</sub> LI <sub>Y</sub> NA <sub>K</sub> T <sub>L</sub> AD <sub>G</sub> V <sub>P</sub> S <sub>R</sub> F <sub>S</sub> G <sub>S</sub> G <sub>S</sub> T <sub>D</sub> Y <sub>T</sub> FT <sub>I</sub> SS <sub>L</sub> Q <sub>P</sub> EDIATYYC <sub>Q</sub> H <sub>F</sub> W <sub>S</sub> I <sub>P</sub> Y <sub>T</sub> F <sub>G</sub> Q <sub>G</sub> T <sub>K</sub> L <sub>Q</sub> I <sub>T</sub> R <sub>T</sub> <u>TVAAP</u> DIQMTQS PSSVSASVGDR <sub>V</sub> T <sub>I</sub> C <sub>R</sub> AS <sub>Q</sub> GI <sub>S</sub> SS <sub>W</sub> L <sub>A</sub> W <sub>Y</sub> Q <sub>Q</sub> K <sub>P</sub> G <sub>K</sub> AP <sub>K</sub> L <sub>L</sub> <u>I</u> Y <sub>E</sub> AS <sub>N</sub> LET <sub>G</sub> V <sub>P</sub> S <sub>R</sub> F <sub>S</sub> G <sub>S</sub> G <sub>S</sub> D <sub>F</sub> LT <sub>I</sub> SS <sub>L</sub> Q <sub>P</sub> E <sub>D</sub> F <sub>A</sub> TY <sub>Y</sub> <u>C</u> <sub>Q</sub> Q <sub>T</sub> <sub>S</sub> <sub>F</sub> <sub>L</sub> <sub>L</sub> S <sub>F</sub> GG <sub>G</sub> T <sub>K</sub> VE <sub>H</sub> K <sub>R</sub>  TVAAPSVFIFPPSDEQLKSGTASVVC <sub>LLNNF</sub> YPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYSLSSTL <sub>SKADYE</sub> K HKVYACEVTHQGLSSPVTKSFNRGEC ( <b>SEQ ID NO: 67</b> )
DLL4/VEGFH C(h1A11.1. A6-LS-Av)			EVQLVESGGLVQPGGSLRLSCAASGFT <sub>R</sub> <u>HFPMA</u> WVRQA PGKGLEWVA <sub>T</sub> I <sub>S</sub> <sub>S</sub> DA <sub>W</sub> P <sub>S</sub> Y <sub>R</sub> DSVKG <sub>R</sub> FT <sub>I</sub> SRDN <sub>A</sub> K <sub>N</sub> SLY LQMN <sub>S</sub> LRAED <sub>T</sub> AVYYC <sub>S</sub> R <sub>G</sub> Y <sub>Y</sub> N <sub>S</sub> P <sub>F</sub> A <sub>Y</sub> WGQG <sub>T</sub> L <sub>V</sub> T <sub>V</sub> S <sub>S</sub> <u>A</u> <u>S</u> <sub>T</sub> KGP <sub>S</sub> V <sub>F</sub> PL <sub>A</sub> P <sub>E</sub> V <sub>L</sub> Q <sub>E</sub> S <sub>G</sub> GG <sub>L</sub> VQPGGSLRLSCAASG <sub>Y</sub> T <sub>F</sub> T <sub>N</sub> Y <sub>G</sub> M <sub>N</sub> W <sub>V</sub> RQAPGKGLEWVG <sub>W</sub> INT <sub>T</sub> Y <sub>G</sub> E <sub>P</sub> TY <sub>A</sub> AD <sub>F</sub> K <sub>R</sub> R <sub>F</sub> F <sub>S</sub> L <sub>D</sub> T <sub>S</sub> K <sub>S</sub> T <sub>A</sub> Y <sub>L</sub> Q <sub>M</sub> N <sub>S</sub> LRAED <sub>T</sub> AVYYC <sub>A</sub> K <sub>Y</sub> PH <sub>Y</sub> Y <sub>G</sub> SS <sub>H</sub> W <sub>Y</sub> <u>F</u> <sub>D</sub> <sub>V</sub> WGQG <sub>T</sub> L <sub>V</sub> T <sub>V</sub> S <sub>S</sub>  ASTKGPSVFPLAPSSK <sub>S</sub> TSGGTAALGCLV <sub>K</sub> DYF <sub>PEP</sub> PTV <sub>S</sub> WNSGALTSGVHTFP <sub>A</sub> V <sub>L</sub> QSSGLY <sub>S</sub> LSSV <sub>V</sub> TV <sub>P</sub> SS <sub>L</sub> GT <sub>Q</sub> T YICNVN <sub>H</sub> KPSNTKVDKK <sub>V</sub> EP <sub>K</sub> SCDK <sub>T</sub> H <sub>T</sub> C <sub>PP</sub> CP <sub>A</sub> PEAAGG PSVFLF <sub>PP</sub> PK <sub>D</sub> TL <sub>M</sub> ISRTPEV <sub>T</sub> CV <sub>V</sub> VD <sub>V</sub> SHED <sub>E</sub> PEV <sub>K</sub> FN <sub>W</sub> YVDGVEVHNAK <sub>T</sub> KP <sub>R</sub> EEQ <sub>Y</sub> N <sub>ST</sub> Y <sub>R</sub> V <sub>V</sub> S <sub>V</sub> L <sub>T</sub> V <sub>L</sub> H <sub>Q</sub> D <sub>W</sub> L <sub>N</sub> G <sub>K</sub> EYKCKVSNK <sub>A</sub> L <sub>P</sub> APIE <sub>K</sub> T <sub>I</sub> S <sub>K</sub> A <sub>G</sub> Q <sub>P</sub> REP <sub>Q</sub> V <sub>Y</sub> T <sub>L</sub> PP <sub>S</sub> REE MTKNQVSLT <sub>C</sub> L <sub>V</sub> K <sub>G</sub> F <sub>Y</sub> PS <sub>D</sub> I <sub>A</sub> V <sub>E</sub> W <sub>E</sub> S <sub>N</sub> G <sub>Q</sub> P <sub>E</sub> NNY <sub>K</sub> T <sub>T</sub> PP <sub>V</sub> LSDGSFF <sub>FL</sub> Y <sub>S</sub> K <sub>L</sub> T <sub>V</sub> D <sub>K</sub> S <sub>R</sub> W <sub>Q</sub> Q <sub>G</sub> N <sub>V</sub> F <sub>S</sub> C <sub>S</sub> V <sub>M</sub> HEALHN <sub>Y</sub> T QKSLSLSPGK* ( <b>SEQ ID NO: 68</b> )
DLL4/VEGFL C(h1A11.1. A6-LS-Av)			DIQMTQSPSSLSASVGDR <sub>V</sub> T <sub>I</sub> C <sub>R</sub> A <sub>S</sub> E <sub>D</sub> I <sub>Y</sub> S <sub>N</sub> L <sub>A</sub> W <sub>Y</sub> Q <sub>Q</sub> K <sub>P</sub> GKAPK <sub>L</sub> LI <sub>Y</sub> DT <sub>NN</sub> LAD <sub>G</sub> V <sub>P</sub> S <sub>R</sub> F <sub>S</sub> G <sub>S</sub> G <sub>S</sub> T <sub>D</sub> FT <sub>L</sub> TI <sub>S</sub> L <sub>Q</sub> <sub>P</sub> EDFATYYC <sub>Q</sub> Q <sub>Y</sub> NN <sub>Y</sub> P <sub>P</sub> T <sub>F</sub> Q <sub>G</sub> Q <sub>T</sub> K <sub>L</sub> E <sub>I</sub> K <sub>R</sub> <u>TVAAP</u> DIQMTQS PSSLSASVGDR <sub>V</sub> T <sub>I</sub> C <sub>S</sub> A <sub>S</sub> Q <sub>D</sub> I <sub>S</sub> NY <sub>L</sub> W <sub>Y</sub> Q <sub>Q</sub> K <sub>P</sub> G <sub>K</sub> AP <sub>K</sub> V <sub>L</sub>

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			1234567890123456789012345678901234567890
			<b>IYFTSSLHSGVPSRFSGSGSGTDFTLTISSLQPEDFATYY CQQYSTVPWTFGQGTKVEIKR</b>  TVAAPSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTKADYEK HKVYACEVTHQGLSSPVTKSFRGEC* (SEQ ID NO: 69)
DLL4/VEGFH C (h1A11.1. A6-SL-Av)			EVQLVESGGGLVQPGGSLRLSCAASGFTFR <b>HFPMAWVRQA</b> PGKGLEWA <b>T</b> <b>I</b> <b>S</b> <b>S</b> <b>D</b> <b>A</b> <b>W</b> <b>P</b> <b>S</b> <b>Y</b> <b>R</b> <b>D</b> <b>S</b> <b>V</b> <b>K</b> <b>G</b> <b>R</b> <b>F</b> <b>T</b> <b>I</b> <b>S</b> <b>R</b> <b>D</b> <b>N</b> <b>A</b> <b>K</b> <b>N</b> <b>S</b> <b>L</b> LQMNSLRAEDTAVYYCSRG <b>Y</b> <b>Y</b> <b>N</b> <b>S</b> <b>P</b> <b>F</b> <b>A</b> <b>Y</b> <b>W</b> <b>G</b> <b>Q</b> <b>G</b> <b>T</b> <b>L</b> <b>V</b> <b>T</b> <b>V</b> <b>S</b> <b>S</b> <b>A</b> <b>S</b> <b>T</b> <b>K</b> <b>G</b> <b>P</b> <b>E</b> <b>V</b> <b>Q</b> <b>L</b> <b>V</b> <b>E</b> <b>S</b> <b>G</b> <b>G</b> <b>L</b> <b>V</b> <b>Q</b> <b>P</b> <b>G</b> <b>G</b> <b>S</b> <b>L</b> <b>R</b> <b>L</b> <b>S</b> <b>C</b> <b>A</b> <b>A</b> <b>S</b> <b>G</b> <b>Y</b> <b>T</b> <b>F</b> <b>T</b> <b>N</b> <b>Y</b> <b>G</b> <b>M</b> <b>N</b> <b>W</b> VRQAPGKGLEWVGWINTYT <b>G</b> <b>E</b> <b>P</b> <b>T</b> <b>Y</b> <b>A</b> <b>A</b> <b>D</b> <b>F</b> <b>K</b> <b>R</b> <b>R</b> <b>F</b> <b>T</b> <b>E</b> <b>S</b> <b>L</b> <b>D</b> <b>T</b> <b>S</b> STAYLQMNSLRAEDTAVYYCAK <b>Y</b> <b>P</b> <b>H</b> <b>Y</b> <b>G</b> <b>S</b> <b>S</b> <b>H</b> <b>W</b> <b>Y</b> <b>F</b> <b>D</b> <b>V</b> <b>W</b> <b>Q</b> <b>G</b> TLTVSS  ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG PSVFLFPPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGK (SEQ ID NO: 74)
DLL4/VEGFL C (h1A11.1. A6-SL-Av)			DIQMTQSPSSLSASVGDRVITIC <b>R</b> <b>A</b> <b>S</b> <b>E</b> <b>D</b> <b>I</b> <b>Y</b> <b>S</b> <b>N</b> <b>L</b> <b>A</b> <b>W</b> <b>Y</b> <b>Q</b> <b>Q</b> <b>K</b> GKAPKLLIY <b>D</b> <b>T</b> <b>N</b> <b>N</b> <b>L</b> <b>A</b> <b>D</b> <b>G</b> <b>V</b> <b>P</b> <b>S</b> <b>R</b> <b>F</b> <b>S</b> <b>G</b> <b>S</b> <b>G</b> <b>T</b> <b>D</b> <b>F</b> <b>T</b> <b>L</b> <b>T</b> <b>I</b> <b>S</b> <b>L</b> <b>Q</b> EDFATYYC <b>Q</b> <b>Q</b> <b>Y</b> <b>N</b> <b>Y</b> <b>P</b> <b>P</b> <b>T</b> <b>F</b> <b>G</b> <b>Q</b> <b>G</b> <b>T</b> <b>K</b> <b>L</b> <b>E</b> <b>I</b> <b>K</b> <b>R</b> <b>T</b> <b>V</b> <b>A</b> <b>A</b> <b>P</b> <b>S</b> <b>V</b> <b>F</b> <b>I</b> <b>F</b> <b>P</b> DIQMTQSPSSLSASVGDRVITIC <b>S</b> <b>A</b> <b>S</b> <b>Q</b> <b>D</b> <b>I</b> <b>S</b> <b>N</b> <b>Y</b> <b>L</b> <b>N</b> <b>W</b> <b>Y</b> <b>Q</b> <b>Q</b> <b>K</b> GKAPKVLIY <b>F</b> <b>T</b> <b>S</b> <b>S</b> <b>L</b> <b>H</b> <b>S</b> <b>G</b> <b>V</b> <b>P</b> <b>S</b> <b>R</b> <b>F</b> <b>S</b> <b>G</b> <b>S</b> <b>G</b> <b>T</b> <b>D</b> <b>F</b> <b>T</b> <b>L</b> <b>T</b> <b>I</b> <b>S</b> <b>L</b> <b>Q</b> EDFATYYC <b>Q</b> <b>Q</b> <b>Y</b> <b>S</b> <b>T</b> <b>V</b> <b>P</b> <b>W</b> <b>T</b> <b>F</b> <b>G</b> <b>Q</b> <b>G</b> <b>T</b> <b>K</b> <b>V</b> <b>E</b> <b>I</b> <b>K</b>  TVAAPSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTKADYEK HKVYACEVTHQGLSSPVTKSFRGEC (SEQ ID NO: 75)
IL12/IL18H			EVTLRESPALVKPTQTLTCTFSGFSLSK <b>S</b> <b>V</b> <b>M</b> <b>G</b> <b>V</b> <b>W</b> IR QPPGKALEWLA <b>H</b> <b>I</b> <b>Y</b> <b>W</b> <b>DD</b> <b>D</b> <b>K</b> <b>Y</b> <b>Y</b> <b>N</b> <b>P</b> <b>S</b> <b>L</b> <b>K</b> <b>S</b> <b>R</b> <b>L</b> <b>T</b> <b>I</b> <b>S</b> <b>K</b> <b>D</b> <b>T</b> <b>S</b> <b>K</b> <b>N</b> <b>Q</b> VLTMTNMDPVDATAYYCAR <b>R</b> <b>G</b> <b>I</b> <b>R</b> <b>S</b> <b>A</b> <b>M</b> <b>D</b> <b>Y</b> <b>W</b> <b>Q</b> <b>G</b> <b>T</b> <b>T</b> <b>V</b> <b>V</b> <b>S</b> <b>A</b> <b>S</b> <b>T</b> <b>K</b> <b>G</b> <b>P</b> <b>E</b> <b>V</b> <b>Q</b> <b>L</b> <b>V</b> <b>Q</b> <b>S</b> <b>G</b> <b>T</b> <b>E</b> <b>V</b> <b>K</b> <b>K</b> <b>P</b> <b>G</b> <b>E</b> <b>S</b> <b>L</b> <b>K</b> <b>I</b> <b>S</b> <b>C</b> <b>K</b> <b>G</b> <b>S</b> <b>G</b> <b>T</b> <b>V</b> <b>T</b> <b>S</b> <b>Y</b> <b>W</b> GWRQMPGKGLEWMG <b>F</b> <b>I</b> <b>Y</b> <b>P</b> <b>G</b> <b>D</b> <b>S</b> <b>E</b> <b>T</b> <b>R</b> <b>Y</b> <b>S</b> <b>P</b> <b>T</b> <b>F</b> <b>Q</b> <b>G</b> <b>Q</b> <b>V</b> <b>T</b> <b>I</b> <b>S</b> <b>A</b> <b>D</b> <b>K</b> FNTAFLQWSSLKASDTAMYYCAR <b>V</b> <b>G</b> <b>S</b> <b>G</b> <b>W</b> <b>P</b> <b>Y</b> <b>T</b> <b>F</b> <b>D</b> <b>I</b> <b>W</b> <b>Q</b> <b>G</b> T MVTSS  ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQT YICNVNHPNSNTKVDKKVEPKSCDKTHCPCPAPEAAGG PSVFLFPPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV LSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGK (SEQ ID NO: 70)
IL12/IL18 LC			DIVMTQSPDSLAVSLGERATINC <b>K</b> <b>A</b> <b>S</b> <b>Q</b> <b>S</b> <b>V</b> <b>S</b> <b>N</b> <b>D</b> <b>V</b> <b>A</b> WYQQKP GQPPKLLIY <b>Y</b> <b>A</b> <b>S</b> <b>N</b> <b>R</b> <b>Y</b> <b>T</b> <b>G</b> <b>V</b> <b>P</b> <b>D</b> <b>R</b> <b>F</b> <b>S</b> <b>G</b> <b>S</b> <b>G</b> <b>T</b> <b>D</b> <b>F</b> <b>T</b> <b>L</b> <b>T</b> <b>I</b> <b>S</b> <b>L</b> <b>Q</b> A

DVD-Ig Heavy or Light Chain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence, with Line Break Between Variable and Constant Regions
			<p>1234567890123456789012345678901234567890</p> <p>EDVAVYYC<b>QQDYN</b>SPWTFGGGTKVEIKR<b>TVAAP</b>EIVMTQS  PATLSVSPGERATLSC<b>RASE</b><b>SISSN</b>LA<sup>WY</sup>QQKPGQAPRLF  <b>IY</b><b>TASTR</b><b>ATDIP</b>ARFSGSGSGTEFTLTISLQSEDFAVYY  <b>CQQYNNWP</b><b>SIT</b>FGQGTRLEIKR</p> <p>TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW  KVDNALQSGNSQESVTEQDSKDSTYSLSSTL<sup>SKADYEK</sup>  HKVYACEVTHQGLSSPVTKSFNRGEC (<b>SEQ ID NO: 71</b>)</p>
DLL4/VEGF HC h1A11.1- LS-AV			<p>EVQLVESGGGLVQPGGSLRLSCAASGFTFS<b>NFPMA</b>WVRQA  PGKGLEWVAT<b>TISSDGTYYRDSVKG</b>RFTISRDNAKNSLY  LQMNSLRAEDTA<sup>V</sup>Y<sup>C</sup><b>ARGYYNSPFAY</b>WGQGTLVTVSS<b>AS</b>  <b>TKG</b><b>PSVF</b><b>PLAP</b>EVQLVESGGGLVQPGGSLRLSCAASGYTF  <b>TNYGMN</b>WVRQAPGKGLEWVG<b>WINTY</b><b>TGEPTY</b><b>AADF</b><b>KRRF</b>  FSLDTSKSTAYLQMNSLRAEDTA<sup>V</sup>Y<sup>C</sup><b>AKY</b><b>PHYYGSSH</b><b>WY</b>  <b>FDV</b>WGQGTLVTVSS</p> <p>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS  WNSGALTSGVHTFP<sup>A</sup>VLQSSGLYSLSSVTV<sup>P</sup>SSSLGTQT  YICNVN<sup>H</sup>KPSNTKV<sup>D</sup>KKVEPKSCDKTHC<sup>P</sup>CP<sup>C</sup>PAPEAAGG  PSVFLFPPPKD<sup>T</sup>LMISRTPEVTCVVVDVSHEDPEV<sup>K</sup>FNW  YVDGVEVHNA<sup>K</sup>TKP<sup>R</sup>EEQYN<sup>S</sup>TYRVV<sup>S</sup>VL<sup>T</sup>VLHQDWLN<sup>G</sup>  EYKCKVSNKALP<sup>A</sup>PIEKT<sup>I</sup>SKAKGQ<sup>P</sup>REPQV<sup>T</sup>L<sup>P</sup>SREE  MTKNQVSLTCLVKGFYPSDIAVEWESNGQ<sup>P</sup>ENNYKT<sup>PP</sup>V  LDSDGSFFFLYSKLTVD<sup>K</sup>SRWQQGNVFSC<sup>S</sup>VMHEALHN<sup>Y</sup>T  Q<sup>K</sup>SLSLSPGK (<b>SEQ ID NO: 72</b>)</p>
DLL4/VEGF LC h1A11.1- LS-AV			<p>DIQMTQSPSSLSASVGDRVTITC<b>RASEDIY</b>SNLA<sup>WY</sup>QQKPGKAPKLLIY<b>DTNNLADGVP</b>SRFSGSGSGTDF<sup>T</sup>L<sup>T</sup>ISLQPEFATYYC<b>QQYNNY</b>PPTFGQG<sup>T</sup>KLEIKR<b>TVAAP</b>DIQMTQS  PSSLSASVGDRVTITC<b>SASQD</b><b>ISNYLN</b>WYQQKPGKAPKVL<b>IY</b><b>FTSS</b><b>LHSGVP</b>SRFSGSGSGTDF<sup>T</sup>L<sup>T</sup>ISLQPEDFATYYC<b>QQY</b><b>STVP</b><b>WTF</b>FGQG<sup>T</sup>KVEIKR</p> <p>TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW  KVDNALQSGNSQESVTEQDSKDSTYSLSSTL<sup>SKADYEK</sup>  HKVYACEVTHQGLSSPVTKSFNRGEC (<b>SEQ ID NO: 73</b>)</p>

### **Incorporation by Reference**

The contents of all cited references (including literature references, patents, patent applications, and websites) that may be cited throughout this application are hereby expressly incorporated by reference in their entirety, as are the references cited therein. The practice of the present disclosure will employ, unless otherwise indicated, conventional techniques of immunology, molecular biology and cell biology, which are well known in the art.

### **Equivalents**

The disclosure may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting of the disclosure described herein. Scope of the disclosure is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are therefore intended to be embraced herein.

## CLAIMS

1. An aqueous formulation comprising an Aqueous Stable DVD-Ig (AS-DVD-Ig) protein and a buffer having a molarity of about 5 to about 50 mM, wherein the formulation has a pH of about 4.5 to about 7.5.
2. The aqueous formulation of claim 1, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 15% loss in relative percentage monomers as determined by size exclusion chromatography (SEC) when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 degrees C.
3. The aqueous formulation of claim 1, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 10% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 degrees C.
4. The aqueous formulation of claim 1, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 5% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of at least about 50 mg/ml, following 14 days storage at about 40 degrees C.
5. The aqueous formulation of claim 1, wherein the formulation comprises about 6% or less aggregation as determined by SEC analysis.
6. The aqueous formulation of claim 1, wherein the formulation comprises about 5% or less aggregation as determined by SEC analysis.
7. The aqueous formulation of any one of claims 1-6, wherein the formulation has a pH of about 5 to about 6.5.
8. The aqueous formulation of any one of claims 1-7, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having about a 10% relative (rel.) peak area or less change in monomers at about 40 °C after 21 days of storage at a concentration of about 100 mg/ml in an aqueous formulation at a pH between about 5.5 to about 6.5.
9. The aqueous formulation of any one of claims 1-8, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having about a 1% rel. peak area or less change in

monomers at about 5 °C after 21 days of storage at a concentration of about 100 mg/ml at a pH between about 5.5 to about 6.5 in an aqueous formulation.

10. The aqueous formulation of any one of claims 1-9, further comprising at least one component selected from a surfactant, a polyol, and combinations thereof.

11. The aqueous formulation of claim 10, wherein the polyol is selected from the group consisting of sorbitol, mannitol, and sucrose.

12. The aqueous formulation of claim 11, wherein the polyol is mannitol and wherein the concentration of mannitol is selected from the group consisting of about 10 to about 100 mg/ml, about 20 to about 80, about 20 to about 70, about 30 to about 60, and about 30 to about 50 mg/ml.

13. The aqueous formulation of claim 11, wherein the polyol is sorbitol.

14. The aqueous formulation of claim 13, wherein the concentration of sorbitol is selected from the group consisting of about 20 to about 60 mg/ml, about 25 to about 55 mg/ml, about 30 to about 50 mg/ml, and about 35 to about 45 mg/ml.

15. The aqueous formulation of claim 11, wherein the polyol is sucrose.

16. The aqueous formulation of claim 15, wherein the concentration of sucrose is selected from the group consisting of about 60 to about 100 mg/ml, about 65 to about 95 mg/ml, about 70 to about 90 mg/ml, and about 75 to about 85 mg/ml.

17. The aqueous formulation of any one of claims 10-16, wherein the surfactant is a polysorbate.

18. The aqueous formulation of claim 17, wherein the concentration of polysorbate is selected from the group consisting of about 0.001% to about 1%, about 0.005% to about 0.05%, about 0.005% to about 0.02%, about 0.01% to about 0.05%, and about 0.1%.

19. The aqueous formulation of claim 17 or claim 18, wherein the polysorbate is polysorbate 80 or polysorbate 20.

20. The aqueous formulation of claim 19, wherein the polysorbate 80 or the polysorbate 20 has a concentration of about 0.005% to about 0.02%.

21. The formulation of any one of claims 1-20, wherein the buffer is selected from the group consisting of acetate, histidine, glycine, arginine, phosphate, and citrate.

22. The aqueous formulation of any one of claims 1-21, wherein the molarity of the buffer ranges from 10 to 20 mM.

23. The aqueous formulation of any one of claims 1-22, wherein the AS-DVD-Ig protein has a concentration of about 1 to about 200 mg/ml.

24. The aqueous formulation of claim 23, wherein the AS-DVD-Ig protein has a concentration selected from the group consisting of about 20 to about 100 mg/ml, about 1 to about 250 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, and about 80 to about 105 mg/ml of the AS-DVD-Ig protein.

25. An aqueous formulation comprising a Aqueous Stable DVD-Ig (AS-DVD-Ig) protein, a buffer having a molarity of about 5 to about 50 mM, a surfactant, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5.

26. The aqueous formulation of claim 25, wherein the polyol is selected from the group consisting of sorbitol, mannitol, and sucrose.

27. The aqueous formulation of claim 26, wherein the polyol is sorbitol.

28. The aqueous formulation of claim 27, wherein the concentration of sorbitol is about 30 to about 50 mg/ml.

29. The aqueous formulation of claim 26, wherein the polyol is sucrose.

30. The aqueous formulation of claim 29, wherein the sucrose has a concentration of about 70 to about 90 mg/ml.

31. The aqueous formulation of any one of claims 25-30, wherein the buffer is selected from the group consisting of acetate, histidine, glycine, arginine, phosphate, and citrate.

32. The aqueous formulation of any one of claims 25-31, wherein the molarity of the buffer ranges from about 10 to about 20 mM.

33. The aqueous formulation of any one of claims 25-32, wherein the AS-DVD-Ig protein has a concentration of about 1 to about 200 mg/ml.

34. The aqueous formulation of any one of claims 25-32, wherein the AS-DVD-Ig protein has a concentration selected from the group consisting of about 20 to about 100 mg/ml, about 1 to about 250 mg/ml, about 10 to about 230 mg/ml, about 20 to about 210 mg/ml, about 30 to about 190 mg/ml, about 40 to about 170 mg/ml, about 50 to about 150 mg/ml, about 60 to about 130 mg/ml, about 70 to about 110 mg/ml, and about 80 to about 105 mg/ml of the AS-DVD-Ig protein.

35. A formulation comprising an AS-DVD-Ig protein, a polyol, buffer, and a surfactant, wherein said formulation has a pH of about 5 to about 7, and wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 15% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of about 60 mg/ml following 14 days storage at about 40 degrees C.

36. The formulation of claim 35, wherein the polyol is selected from the group consisting of sorbitol, mannitol, and sucrose.

37. The formulation of claim 35 or 36, wherein the buffer is selected from the group consisting of acetate, histidine, glycine, arginine, phosphate, and citrate.

38. The formulation of any one of claims 35-38, wherein the surfactant is a polysorbate.

39. The formulation of claim 38, wherein the polysorbate is polysorbate 80 or polysorbate 20.

40. The formulation of claim 39, wherein the polysorbate 80 or the polysorbate 20 has a concentration of about 0.005% to about 0.02%.

41. A formulation comprising an AS-DVD-Ig protein, a polyol, histidine buffer, and a polysorbate, wherein said formulation has a pH of about 5 to about 7, and wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 15% aggregation as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of about 60 mg/ml following 14 days storage at about 40 degrees C.

42. The formulation of any one of claims 35-41, which is an aqueous formulation.

43. The formulation of any one of claims 35-41, which is a lyophilized formulation.

44. The formulation of any one of claims 1-43, wherein the AS-DVD-Ig protein comprises first and second polypeptide chains, each independently comprising VD1-(X1)<sub>n</sub>-VD2-C-(X2)<sub>n</sub>, wherein

VD1 is a first variable domain;

VD2 is a second variable domain;

C is a constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

n is 0 or 1,

wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

45. The formulation of claim 44, wherein the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein

VD1 is a first heavy chain variable domain;

VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

n is 0 or 1, and

wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n,

wherein

VD1 is a first light chain variable domain;

VD2 is a second light chain variable domain;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

n is 0 or 1,

wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

46. The formulation of claim 44 or 45, comprising two first polypeptide chains and two second polypeptide chains, wherein the binding protein comprises four functional target binding sites.

47. The formulation of any one of claims 44-46, wherein X1 is not CL.

48. The formulation of any one of claims 44-47, wherein the AS-DVD-Ig protein comprises three CDRs from the variable light or heavy chain amino acid sequences as set forth in SEQ ID NOS: 28 to 75.

49. The formulation of any one of claims 44-48, wherein the AS-DVD-Ig protein comprises a light or heavy chain amino acid sequence as set forth in SEQ ID NOS: 28 to 75.

50. The formulation of any one of claims 44-49, wherein the AS-DVD-Ig protein has a binding specificity selected from the group consisting of CD20/CD80, VEGF/Her2, TNF/RANKL, TNF/DKK, CD20/RANKL, DLL4/PLGF, TNF/SOST, IL-9/IgE, IL-12/IL-18, TNF/IL-17, TNF/PGE2, IL1 $\alpha$ /IL1 $\beta$ , and DLL4/VEGF.

51. The formulation of any one of claims 44-49, wherein the AS-DVD-Ig protein has a binding specificity selected from the group consisting of IL4/IL13, IL1 $\alpha$ /IL1 $\beta$ , and TNF $\alpha$  / IL17.

52. The formulation of claim 50, wherein the AS-DVD-Ig protein has a binding specificity for TNF and IL-17.

53. The formulation of claim 52, wherein the TNF $\alpha$  / IL17 specific AS-DVD-Ig protein is DVD A (SEQ ID NOS: 62 and 63).

54. The formulation of claim 50, wherein the AS-DVD-Ig protein has a binding specificity for IL1 $\alpha$ /IL1 $\beta$ .

55. The formulation of claim 54, wherein the IL1 $\alpha$ /IL1 $\beta$  specific AS-DVD-Ig protein is DVD C (SEQ ID NOS: 66 and 67).

56. The formulation of claim 50, wherein the AS-DVD-Ig protein has a binding specificity for IL-12/IL-18.

57. The formulation of any one of claims 1 to 56, wherein the formulation is a pharmaceutical formulation.

58. A pharmaceutical composition comprising an Aqueous Stable DVD-Ig (AS-DVD-Ig) protein and a buffer having a molarity of about 5 to about 50 mM, wherein the AS-DVD-Ig protein is characterized as a DVD-Ig protein having less than about 10% loss in relative percentage monomers as determined by SEC when formulated in a histidine or citrate phosphate buffer at a concentration of about 60 mg/ml, following 14 days storage at about 40 degrees C, and the formulation has a pH of 4.5 to 7.5.

59. The pharmaceutical composition of claim 58, further comprising at least one component selected from a surfactant, a polyol, and combinations thereof.

60. A method of treating a disorder, comprising administering the pharmaceutical composition of claim 58 or 59, such that the disorder is treated.

61. A lyophilized formulation comprising a Lyophilized-Stable DVD-Ig (LS-DVD-Ig) protein, wherein when said formulation is reconstituted, it comprises about 1 to about 100 mg/ml of the LS-DVD-Ig protein, about 10 to about 50 mM of a buffer, a polyol, about 0.01 to about 0.2 mg/ml of a polysorbate, and has a pH of about 5 to about 7.

62. The lyophilized formulation of claim 61, wherein the formulation has a pH of about 5.5 to about 6.5.

63. The lyophilized formulation of claim 61 or 62, wherein the LS-DVD-Ig protein has more than 10% rel. peak area change in monomers observed, following accelerated storage at a pH between about 5.5 and about 6.5 in an aqueous formulation for 21 days at about 40 °C, when formulated at a concentration over about 100 mg/ml.

64. The lyophilized formulation of any one of claims 61-63, wherein the LS-DVD-Ig protein comprises first and second polypeptide chains, each independently comprising VD1-(X1)n-VD2-C-(X2)n, wherein

VD1 is a first variable domain;

VD2 is a second variable domain;

C is a constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

n is 0 or 1,

wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

65. The lyophilized formulation of claim 64, wherein the first polypeptide chain comprises a first VD1-(X1)n-VD2-C-(X2)n, wherein

VD1 is a first heavy chain variable domain;

VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

n is 0 or 1, and

wherein the second polypeptide chain comprises a second VD1-(X1)n-VD2-C-(X2)n,

wherein

VD1 is a first light chain variable domain;

VD2 is a second light chain variable domain;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

n is 0 or 1,

wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site.

66. The lyophilized formulation of claim 64 or 65, comprising two first polypeptide chains and two second polypeptide chains, wherein the binding protein comprises four functional target binding sites.

67. The lyophilized formulation of any one of claims 64-66, wherein X1 is not CL.

68. The lyophilized formulation of any one of claims 64-67, wherein the LS-DVD-Ig protein comprises three CDRs from the variable light or heavy chain amino acid sequences as set forth in SEQ ID NOs: 28 to 75.

69. The lyophilized formulation of any one of claims 63-66, wherein the LS-DVD-Ig protein comprises a light or heavy chain amino acid sequence as set forth in SEQ ID NOs: 28 to 75.

70. The lyophilized formulation of any one of claims claims 61-69, wherein the LS-DVD-Ig protein has a binding specificity selected from the group consisting of CD20/CD80, VEGF/Her2, TNF/RANKL, TNF/DKK, CD20/RANKL, DLL4/PLGF, TNF/SOST, IL-9/IgE, IL-12/IL-18, TNF/IL-17, TNF/PGE2, IL1 $\alpha$ /IL1 $\beta$ , or DLL4/VEGF.

71. The lyophilized formulation of claim 70, wherein the LS-DVD-Ig protein has a binding specificity for TNF and IL-17.

72. The lyophilized formulation of claim 70, wherein the LS-DVD-Ig protein has a binding specificity for IL1 $\alpha$ /IL1 $\beta$ .

73. The lyophilized formulation of claim 70, wherein the LS-DVD-Ig protein has a binding specificity for IL-12/IL-18.

74. A lyophilized formulation prepared by lyophilizing an aqueous formulation comprising a LS-DVD-Ig protein, a buffer have a molarity of about 5 to about 50 mM, a surfactant, and a polyol, wherein the formulation has a pH of about 4.5 to about 7.5.

75. The lyophilized formulation of claim 74, wherein the buffer is selected from the group consisting of histidine, succinate, and citrate and/or phosphate.

76. The lyophilized formulation of claim 74 to 75, wherein the polyol is selected from the group consisting of mannitol, sorbitol, sucrose, and trehalose.

1/6




Fig. 1

2/6




Figure 2A

3/6



Domain orientation concept

Figure 2B

4/6




Figure 3

5/6




Fig. 4

6/6




Fig. 5