

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 December 2005 (29.12.2005)

PCT

(10) International Publication Number
WO 2005/123748 A1

(51) International Patent Classification⁷: **C07D 498/08**,
A61K 31/4386, A61P 9/06

(74) Agent: **ASTRAZENECA**; Global Intellectual Property,
S-151 85 Södertälje (SE).

(21) International Application Number:
PCT/SE2005/000891

(22) International Filing Date: 13 June 2005 (13.06.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0401539-2 15 June 2004 (15.06.2004) SE

(71) Applicant (for all designated States except US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BJÖRE, Annika** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **BONN, Peter** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **GRAN, Ulrik** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **KAJANUS, Johan** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **OLSSON, Christina** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). **PONTÉN, Fritiof** [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL OXABISPIDINE COMPOUNDS AND THEIR USE IN THE TREATMENT OF CARDIAC ARRHYTHMIAS



(I)

(57) Abstract: There is provided compounds of formula (I), wherein R¹, R², R³, R⁴, R⁴¹ to R⁴⁶, A, B and G have meanings given in the description, which are useful in the prophylaxis and in the treatment of arrhythmias, in particular atrial and ventricular arrhythmias.

Novel oxabispidine compounds and their use in the treatment of cardiac arrhythmias.

Field of the Invention

5 This invention relates to novel pharmaceutically useful compounds, in particular compounds which are useful in the treatment of cardiac arrhythmias.

Background and Prior Art

10

Cardiac arrhythmias may be defined as abnormalities in the rate, regularity, or site of origin of the cardiac impulse or as disturbances in conduction which causes an abnormal sequence of activation. Arrhythmias may be classified clinically by means of the presumed site of origin (i.e. as supraventricular, including atrial and atrioventricular, arrhythmias and ventricular arrhythmias) and/or by means of rate (i.e. bradyarrhythmias (slow) and tachyarrhythmias (fast)).

In the treatment of cardiac arrhythmias, the negative outcome in clinical trials (see, for example, the outcome of the Cardiac Arrhythmia Suppression Trial (CAST) reported in *New England Journal of Medicine*, 321, 406 (1989)) with “traditional” antiarrhythmic drugs, which act primarily by slowing the conduction velocity (class I antiarrhythmic drugs), has prompted drug development towards compounds which selectively delay cardiac repolarization, thus prolonging the QT interval. Class III antiarrhythmic drugs may be defined as drugs which prolong the trans-membrane action potential duration (which can be caused by a block of outward K^+ currents or from an increase of inward ion currents) and refractoriness, without affecting cardiac conduction.

One of the key disadvantages of hitherto known drugs which act by delaying repolarization (class III or otherwise) is that they all are known to exhibit a unique form of proarrhythmia known as *torsades de pointes* (turning of points), which may, on occasion be fatal. From the point of view of safety, the minimisation of this phenomenon (which has also been shown to be exhibited as a result of administration of non-cardiac drugs such as phenothiazines, tricyclic antidepressants, antihistamines and antibiotics) is a key problem to be solved in the provision of effective antiarrhythmic drugs.

Antiarrhythmic drugs based on bispidines (3,7-diazabicyclo[3.3.1]nonanes), are known from *inter alia* international patent applications WO 91/07405 and WO 99/31100, European patent applications 306 871, 308 843 and 655 228 and US patents 3,962,449, 4,556,662, 4,550,112, 4,459,301 and 5,468,858, as well as journal articles including, *inter alia*, *J. Med. Chem.* 39, 2559, (1996), *Pharmacol. Res.*, 24, 149 (1991), *Circulation*, 90, 2032 (1994) and *Anal. Sci.* 9, 429, (1993).


Certain oxabispidine compounds are disclosed as chemical curiosities in *Chem. Ber.*, 96, 2872 (1963). The use of certain other oxabispidine compounds in the treatment of cardiac arrhythmias is disclosed in WO 01/28992. Methods for the preparation of such oxabispidine compounds are disclosed in WO 02/28863, WO 02/28864, WO 02/83690 and WO 02/83691. Oxabispidine compounds in which one or both of the *N*-atoms bears a substituent that includes an "in-chain" sulfonamide group are neither disclosed nor suggested.

We have surprisingly found that a novel group of oxabispidine-based compounds exhibit electrophysiological activity and are therefore expected to be useful in the treatment of cardiac arrhythmias. The novel group of

oxabispidine-based compounds has advantageous properties compared to compounds of the prior art, such as enhanced potency, enhanced selectivity, and/or reduction of total clearance. These advantageous properties can distinguish the use of such compounds as pharmaceutical agents by lowering the daily clinical dose, lengthening the duration of action, and / or improving the side effect profile.

Disclosure of the Invention

According to the invention there is provided compounds of formula I,



wherein

R^1 represents C_{1-12} alkyl (which alkyl group is optionally substituted by one or more groups selected from halo, cyano, nitro, aryl, Het¹, $-N(R^{5a})R^6$, $-C(O)R^{5b}$, $-OR^{5c}$, $-C(O)XR^7$, $-C(O)N(R^{8a})R^{5d}$, $-OC(O)N(R^{8b})R^{5e}$, $-S(O)_2R^{9a}$, $-S(O)_2N(R^{9b})R^{9c}$ and $-N(R^{9b})S(O)_2R^{9d}$) or R^1 represents $-C(O)XR^7$, $-C(O)N(R^{8a})R^{5d}$ or $-S(O)_2R^{9a}$;

R^{5a} represents H or C_{1-6} alkyl (which latter group is optionally substituted by one or more substituents selected from $-OH$, halo, cyano, nitro, $-S(O)_2N(R^{9b})R^{9c}$ and $-N(R^{9b})S(O)_2R^{9d}$);

R^{5b} to R^{5e} independently represent, at each occurrence when used herein, H, C_{1-6} alkyl (which latter group is optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, Het², -S(O)₂N(R^{9b}) R^{9c} and -N(R^{9b})S(O)₂ R^{9d}), aryl or Het³, or R^{5d} or R^{5e} , together
 5 with, respectively, R^{8a} or R^{8b} , may represent C_{3-6} alkylene (which alkylene group is optionally interrupted by an O atom and/or is optionally substituted by one or more C_{1-3} alkyl groups);

R^6 represents H, C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, -S(O)₂N(R^{9b}) R^{9c}
 10 and -N(R^{9b})S(O)₂ R^{9d}), aryl, -C(O) R^{10a} , -C(O)OR^{10b} or -C(O)N(H) R^{10c} or -S(O)₂ R^{10d} ;

R^{10a} , R^{10b} , R^{10c} and R^{10d} independently represent C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, or R^{10a} represents H;

15 R^7 represents, at each occurrence when used herein, C_{1-12} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, C_{1-6} alkoxy, Het⁴, -S(O)₂N(R^{9b}) R^{9c} and -N(R^{9b})S(O)₂ R^{9d});

R^{8a} and R^{8b} independently represent H, C_{1-12} alkyl, C_{1-6} alkoxy (which latter two groups are optionally substituted by one or more substituents selected
 20 from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy, -S(O)₂N(R^{9b}) R^{9c} and -N(R^{9b})S(O)₂ R^{9d}), -D-aryl, -D-aryloxy, -D-Het⁵, -D-N(H)C(O) R^{11a} , -D-S(O)₂ R^{12a} , -D-C(O) R^{11b} , -D-C(O)OR^{12b}, -D-C(O)N(R^{11c}) R^{11d} , or R^{8a} or R^{8b} , together with, respectively, R^{5d} or R^{5e} , may represent C_{3-6} alkylene (which alkylene group is optionally interrupted
 25 by an O atom and/or is optionally substituted by one or more C_{1-3} alkyl groups);

R^{11a} to R^{11d} independently represent H, C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, or R^{11c} and R^{11d} together represent C_{3-6} alkylene;

R^{12a} and R^{12b} independently represent C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro and aryl) or aryl;

D represents, at each occurrence when used herein, a direct bond or C_{1-6} alkylene;

X represents O or S;

R^{9a} represents, at each occurrence when used herein, C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, $-S(O)_2N(R^{9b})R^{9c}$ and $-N(R^{9b})S(O)_2R^{9d}$) or aryl;

R^{9b} represents, at each occurrence when used herein, H or C_{1-6} alkyl;

R^{9c} and R^{9d} independently represent, at each occurrence when used herein, C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl and Het⁶), aryl or Het⁷, or R^{9c} represents H;

15

R^2 represents H, halo, C_{1-6} alkyl, $-OR^{13}$, $-E-N(R^{14})R^{15}$ or, together with R^3 , represents =O;

R^3 represents H, C_{1-6} alkyl or, together with R^2 , represents =O;

R^{13} represents H, C_{1-6} alkyl, -E-aryl, -E-Het⁸, $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or $-C(O)N(R^{17a})R^{17b}$;

20

R^{14} represents H, C_{1-6} alkyl, -E-aryl, -E-Het⁹, $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$, $-[C(O)]_pN(R^{17a})R^{17b}$ or $-C(NH)NH_2$;

R^{15} represents H, C_{1-6} alkyl, -E-aryl or $-C(O)R^{16d}$;

R^{16a} to R^{16d} independently represent, at each occurrence when used herein, C_{1-6} alkyl (optionally substituted by one or more substituents selected from halo, aryl and Het¹⁰), aryl, Het¹¹, or R^{16a} and R^{16d} independently represent H; R^{17a} and R^{17b} independently represent, at each occurrence when used herein, H or C_{1-6} alkyl (optionally substituted by one or more substituents selected

25

from halo, aryl and Het¹²), aryl, Het¹³, or together represent C₃₋₆ alkylene, optionally interrupted by an O atom;

E represents, at each occurrence when used herein, a direct bond or C₁₋₄ alkylene;

5 p represents 1 or 2;

Het¹ to Het¹³ independently represent five- to twelve-membered heterocyclic groups containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, which groups are optionally substituted by one or more
 10 substituents selected from -OH, oxo, halo, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, aryl, aryloxy, -N(R^{18a})R^{18b}, -C(O)R^{18c}, -C(O)OR^{18d}, -C(O)N(R^{18e})R^{18f}, -N(R^{18g})C(O)R^{18h}, -S(O)₂N(R¹⁸ⁱ)R^{18j} and -N(R^{18k})S(O)₂R^{18l};

R^{18a} to R^{18l} independently represent C₁₋₆ alkyl, aryl or R^{18a} to R^{18k}
 15 independently represent H;

A represents a direct bond, -J-, -J-N(R^{19a})-, -J-S(O)₂N(R^{19b})-, -J-N(R^{19c})S(O)₂- or -J-O- (in which latter four groups, -J is attached to the oxabispidine ring nitrogen);

20 B represents -Z-{[C(O)]_aC(H)(R^{20a})_b}-, -Z-[C(O)]_cN(R^{20b})-, -Z-N(R^{20c})S(O)₂-, -Z-S(O)₂N(R^{20d})-, -Z-S(O)_n-, -Z-O- (in which latter six groups, Z is attached to the carbon atom bearing R² and R³), -N(R^{20e})-Z-, -N(R^{20f})S(O)₂-Z-, -S(O)₂N(R^{20g})-Z- or -N(R^{20h})C(O)O-Z- (in which latter
 25 four groups, Z is attached to the phenyl or pyridyl group that is optionally substituted by R⁴);

J represents C₁₋₆ alkylene optionally interrupted by -S(O)₂N(R^{19d})- or -N(R^{19e})S(O)₂- and/or optionally substituted by one or more substituents selected from -OH, halo and amino;

Z represents a direct bond or C₁₋₄ alkylene, optionally interrupted by -N(R²⁰ⁱ)S(O)₂- or -S(O)₂N(R^{20j})-;

a, b and c independently represent 0 or 1;

n represents 0, 1 or 2;

5 R^{19a} to R^{19e} independently represent, at each occurrence when used herein, H or C₁₋₆ alkyl;

R^{20a} represents H or, together with a single R⁴ substituent at a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, R^{20a} represents C₂₋₄ alkylene optionally interrupted or
10 terminated by O, S, N(H) or N(C₁₋₆ alkyl);

R^{20b} represents H, C₁₋₆ alkyl or, together with a single R⁴ substituent at a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, R^{20b} represents C₂₋₄ alkylene;

R^{20c} to R^{20j} independently represent, at each occurrence when used herein, H
15 or C₁₋₆ alkyl;

G represents CH or N;

R⁴ represents one or more optional substituents selected from -OH, cyano,
20 halo, nitro, C₁₋₆ alkyl (optionally terminated by -N(H)C(O)OR^{21a}), C₁₋₆ alkoxy, -N(R^{22a})R^{22b}, -C(O)R^{22c}, -C(O)OR^{22d}, -C(O)N(R^{22e})R^{22f}, -N(R^{22g})C(O)R^{22h}, -N(R²²ⁱ)C(O)N(R^{22j})R^{22k}, -N(R^{22m})S(O)₂R^{21b}, -S(O)₂N(R²²ⁿ)R^{22o}, -S(O)₂R^{21c}, -OS(O)₂R^{21d} and aryl and an R⁴ substituent in a position on the phenyl or pyridyl group that is
25 *ortho*- to the position at which the group B is attached may

- (i) together with R^{20a}, represent C₂₋₄ alkylene optionally interrupted or terminated by O, S or N(H) or N(C₁₋₆ alkyl), or
- (ii) together with R^{20b}, represent C₂₋₄ alkylene;

R^{21a} to R^{21d} independently represent C₁₋₆ alkyl;

R^{22a} and R^{22b} independently represent H, C_{1-6} alkyl or together represent C_{3-6} alkylene, resulting in a four- to seven-membered nitrogen-containing ring;
 R^{22c} to R^{22o} independently represent H or C_{1-6} alkyl; and

5 R^{41} to R^{46} independently represent H or C_{1-3} alkyl;

wherein each aryl and aryloxy group, unless otherwise specified, is optionally substituted;

provided that

10 (a) at least one of the following is the case

(i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group),

(ii) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,

15 (iii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,

(iv) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or

(v) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;

(b) when A represents $-J-N(R^{19a})-$, $-J-N(R^{19c})S(O)_2-$ or $-J-O-$, then:

20 (i) J does not represent C_1 alkylene or 1,1- C_{2-6} alkylene; and

(ii) B does not represent $-N(R^{20b})-$, $-N(R^{20c})S(O)_2-$, $-S(O)_n-$, $-O-$, $-N(R^{20f})S(O)_2-Z-$ or $-N(R^{20h})C(O)O-Z-$ when R^2 and R^3 do not together represent $=O$; and

(c) when R^2 represents $-OR^{13}$ or $-N(R^{14})(R^{15})$, then:

25 (i) A does not represent $-J-N(R^{19a})-$, $-J-N(R^{19c})S(O)_2-$ or $-J-O-$; and

(ii) B does not represent $-N(R^{20b})-$, $-N(R^{20c})S(O)_2-$, $-S(O)_n-$, $-O-$, $-N(R^{20f})S(O)_2-Z-$ or $-N(R^{20h})C(O)O-Z-$;

or a pharmaceutically acceptable derivative thereof.

For the avoidance of doubt it is to be understood that where in this specification a group is qualified by 'hereinbefore defined', 'defined hereinbefore' or 'defined above' the said group encompasses the first
5 occurring and broadest definition as well as each and all of the particular definitions for that group.

Unless otherwise specified, alkyl groups and alkoxy groups as defined herein may be straight-chain or, when there is a sufficient number (i.e. a
10 minimum of three) of carbon atoms be branched-chain, and/or cyclic. Further, when there is a sufficient number (i.e. a minimum of four) of carbon atoms, such alkyl and alkoxy groups may also be part cyclic/acyclic. Such alkyl and alkoxy groups may also be saturated or, when there is a
15 sufficient number (i.e. a minimum of two) of carbon atoms, be unsaturated and/or interrupted by one or more oxygen and/or sulfur atoms. Unless otherwise specified, alkyl and alkoxy groups may also be substituted by one or more halo, and especially fluoro, atoms.

Unless otherwise specified, alkylene groups as defined herein may be
20 straight-chain or, when there is a sufficient number (i.e. a minimum of two) of carbon atoms, be branched-chain. Such alkylene chains may also be saturated or, when there is a sufficient number (i.e. a minimum of two) of carbon atoms, be unsaturated and/or interrupted by one or more oxygen and/or sulfur atoms. Unless otherwise specified, alkylene groups may also
25 be substituted by one or more halo atoms.

The term "aryl", when used herein, includes C₆₋₁₀ aryl groups such as phenyl, naphthyl and the like. The term "aryloxy", when used herein includes C₆₋₁₀ aryloxy groups such as phenoxy, naphthoxy and the like. For

the avoidance of doubt, aryloxy groups referred to herein are attached to the rest of the molecule *via* the O-atom of the oxy-group. Unless otherwise specified, aryl and aryloxy groups may be substituted by one or more substituents including -OH, halo, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, -N(R^{22a})R^{22b}, -C(O)R^{22c}, -C(O)OR^{22d}, -C(O)N(R^{22e})R^{22f},
 5 -N(R^{22g})C(O)R^{22h}, -N(R^{22m})S(O)₂R^{21b}, -S(O)₂N(R²²ⁿ)R^{22o}, -S(O)₂R^{21c}, and/or -OS(O)₂R^{21d} (wherein R^{21b} to R^{21d} and R^{22a} to R^{22o} are as hereinbefore defined). When substituted, aryl and aryloxy groups are preferably substituted by between one and three substituents.

10

The term "halo", when used herein, includes fluoro, chloro, bromo and iodo.

Het (Het¹, Het², Het³, Het⁴, Het⁵, Het⁶, Het⁷, Het⁸, Het⁹, Het¹⁰, Het¹¹, Het¹² and Het¹³) groups that may be mentioned include those containing 1 to 4
 15 heteroatoms (selected from the group oxygen, nitrogen and/or sulfur) and in which the total number of atoms in the ring system are between five and twelve. Het (Het¹, Het², Het³, Het⁴, Het⁵, Het⁶, Het⁷, Het⁸, Het⁹, Het¹⁰, Het¹¹, Het¹² and Het¹³) groups may be fully saturated, wholly aromatic, partly aromatic and/or bicyclic in character. Heterocyclic groups that may
 20 be mentioned include 1-azabicyclo[2.2.2]octanyl, benzimidazolyl, benzisoxazolyl, benzodioxanyl, benzodioxepanyl, benzodioxolyl, benzofuranyl, benzofurazanyl, benzomorpholinyl, 2,1,3-benzoxadiazolyl, benzoxazinonyl, benzoxazolidinyl, benzoxazolyl, benzopyrazolyl, benzo[*e*]pyrimidine, 2,1,3-benzothiadiazolyl, benzothiazolyl, benzothienyl, benzotriazolyl, chromanyl,
 25 chromenyl, cinnolinyl, 2,3-dihydrobenzimidazolyl, 2,3-dihydrobenzo[*b*]-furanyl, 1,3-dihydrobenzo[*c*]furanyl, 2,3-dihydropyrrolo[2,3-*b*]pyridinyl, dimethylisoxazolyl, dioxanyl, furanyl, hexahydropyrimidinyl, hydantoinyl, imidazolyl, imidazo[1,2-*a*]pyridinyl, imidazo[2,3-*b*]thiazolyl, indolyl, isoquinolinyl, isoxazolyl, maleimido, morpholinyl, oxadiazolyl, 1,3-

oxazinanyl, oxazolyl, phthalazinyl, piperazinyl, piperidinyl, purinyl, pyranyl, pyrazinyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrrolidinonyl, pyrrolidinyl, pyrrolinyl, pyrrolo[2,3-*b*]pyridinyl, pyrrolo[5,1-*b*]pyridinyl, pyrrolo[2,3-*c*]pyridinyl, pyrrolyl, quinazolinyl, quinolinyl, sulfolanyl, 3-sulfolenyl,
5 4,5,6,7-tetrahydrobenz-imidazolyl, 4,5,6,7-tetrahydrobenzopyrazolyl, 5,6,7,8-tetrahydrobenzo[*e*]-pyrimidine, tetrahydrofuranyl, tetrahydropyranyl, 3,4,5,6-tetrahydro-pyridinyl, 1,2,3,4-tetrahydropyrimidinyl, 3,4,5,6-tetrahydropyrimidinyl, thiadiazolyl, thiazolidinyl, thiazolyl, thienyl, thieno[5,1-*c*]pyridinyl, thiochromanyl, triazolyl, 1,3,4-triazolo[2,3-
10 *b*]pyrimidinyl and the like.

Values of Het¹ that may be mentioned include 2,3-dihydrobenzo[*b*]furanyl, furanyl, imidazolyl, isoxazolyl, pyridinyl and thiazolyl.

15 Values of Het⁷ that may be mentioned include imidazolyl and isoxazolyl.

Substituents on Het (Het¹, Het², Het³, Het⁴, Het⁵, Het⁶, Het⁷, Het⁸, Het⁹, Het¹⁰, Het¹¹, Het¹² and Het¹³) groups may, where appropriate, be located on any atom in the ring system including a heteroatom. The point of
20 attachment of Het (Het¹, Het², Het³, Het⁴, Het⁵, Het⁶, Het⁷, Het⁸, Het⁹, Het¹⁰, Het¹¹, Het¹² and Het¹³) groups may be *via* any atom in the ring system including (where appropriate) a heteroatom, or an atom on any fused carbocyclic ring that may be present as part of the ring system. Het (Het¹, Het², Het³, Het⁴, Het⁵, Het⁶, Het⁷, Het⁸, Het⁹, Het¹⁰, Het¹¹, Het¹² and Het¹³)
25 groups may also be in the N- or S-oxidised form.

Pharmaceutically acceptable derivatives include salts and solvates. Salts which may be mentioned include acid addition salts.

Pharmaceutically acceptable derivatives also include, at the oxabispidine or (when G represents N) pyridyl nitrogens, C₁₋₄ alkyl quaternary ammonium salts and N-oxides, provided that when a N-oxide is present:

- (a) no Het (Het¹, Het², Het³, Het⁴, Het⁵, Het⁶, Het⁷, Het⁸, Het⁹, Het¹⁰,
5 Het¹¹, Het¹² and Het¹³) group contains an unoxidised S-atom; and/or
(b) n does not represent 0 when B represents -Z-S(O)_n-.

The compounds of the invention may exhibit tautomerism. All tautomeric forms and mixtures thereof are included within the scope of the invention.

10

The compounds of the invention may also contain one or more asymmetric carbon atoms and may therefore exhibit optical and/or diastereoisomerism. Diastereoisomers may be separated using conventional techniques, e.g. chromatography or fractional crystallisation. The various stereoisomers may
15 be isolated by separation of a racemic or other mixture of the compounds using conventional, e.g. fractional crystallisation or HPLC, techniques. Alternatively the desired optical isomers may be made by reaction of the appropriate optically active starting materials under conditions which will not cause racemisation or epimerisation, or by derivatisation, for example with a
20 homochiral acid followed by separation of the diastereomeric esters by conventional means (e.g. HPLC, chromatography over silica). All stereoisomers are included within the scope of the invention.

Abbreviations are listed at the end of this specification.

25

Preferred values of each variable group are as follows. Such values may be used where appropriate with any of the values, definitions, claims, aspects or embodiments defined hereinbefore or hereinafter. In particular, each may be used as an individual limitation on the broadest definition of formula (I).

Preferred compounds of the invention include those in which:

- R^1 represents C_{1-8} alkyl (which alkyl group is optionally substituted by one or more groups selected from halo, aryl (which latter group is optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), $-C(O)R^{22c}$ and $-S(O)_2R^{21c}$), Het¹, $-N(R^{5a})R^6$, $-C(O)R^{5b}$, $-OR^{5c}$, $-C(O)N(R^{8a})R^{5d}$, $-OC(O)N(R^{8b})R^{5e}$, $-S(O)_2R^{9a}$, $-S(O)_2N(R^{9b})R^{9c}$ and $-N(H)S(O)_2R^{9d}$) or R^1 represents $-C(O)OR^7$, $-C(O)N(R^{8a})R^{5d}$ or $-S(O)_2R^{9a}$;
- R^{5a} represents H or C_{1-5} alkyl;
- R^{5b} to R^{5e} independently represent, at each occurrence when used herein, H, C_{1-6} alkyl (which latter group is optionally substituted by one or more substituents selected from cyano, nitro and optionally substituted aryl), aryl (which latter group is optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, $N(R^{22a})R^{22b}$ (in which latter group R^{22a} and R^{22b} together represent C_{3-6} alkylene), C_{1-5} alkyl and C_{1-5} alkoxy (which latter two groups are optionally substituted by one or more halo atoms)), Het³, or R^{5d} , together with R^{8a} , represents C_{4-5} alkylene (which alkylene group is optionally interrupted by an O atom);
- R^6 represents H, C_{1-6} alkyl, optionally substituted aryl $-C(O)R^{10a}$, $-C(O)OR^{10b}$ or $-C(O)N(H)R^{10c}$;
- R^{10a} and R^{10b} independently represent C_{1-5} alkyl (optionally substituted by one or more substituents selected from halo and optionally substituted aryl) or optionally substituted aryl;
- R^{10c} represents C_{1-4} alkyl;
- R^7 represents C_{1-6} alkyl (optionally substituted by one or more substituents selected from halo, optionally substituted aryl, C_{1-4} alkoxy and Het⁴);

- R^{8a} and R^{8b} independently represent H, C_{1-6} alkyl (which latter group is optionally substituted by one or more substituents selected from halo, cyano and nitro), -D-(optionally substituted aryl), -D-(optionally substituted aryloxy), -D-Het⁵, -D-N(H)C(O) R^{11a} , -D-C(O) R^{11b} , or R^{8a} , together with R^{5d}
- 5 represents C_{4-5} alkylene (which alkylene group is optionally interrupted by an O atom);
- R^{11a} and R^{11d} independently represent C_{1-4} alkyl (optionally substituted by one or more substituents selected from halo, cyano, nitro and optionally substituted aryl) or optionally substituted aryl;
- 10 D represents, at each occurrence when used herein, a direct bond or C_{1-4} alkylene;
- R^{9a} represents C_{1-6} alkyl (optionally substituted by one or more halo groups) or optionally substituted aryl;
- R^{9b} represents H or C_{1-3} alkyl;
- 15 R^{9c} and R^{9d} independently represent, at each occurrence when used herein, C_{1-5} alkyl (optionally substituted by one or more substituents selected from halo, optionally substituted aryl and Het⁶), optionally substituted aryl or Het⁷, or R^{9c} represents H;
- R^2 represents H, halo, C_{1-3} alkyl, -OR¹³, -N(H) R^{14} or, together with R^3 ,
- 20 represents =O;
- R^3 represents H, C_{1-3} alkyl or, together with R^2 , represents =O;
- R^{13} represents H, C_{1-4} alkyl, -E-(optionally substituted aryl), or -E-Het⁸;
- R^{14} represents H, C_{1-6} alkyl, -E-(optionally substituted aryl), -C(O) R^{16a} , -C(O)OR^{16b}, -S(O)₂ R^{16c} , -C(O)N(R^{17a}) R^{17b} or -C(NH)NH₂;
- 25 R^{16a} to R^{16c} independently represent C_{1-6} alkyl, or R^{16a} represents H;
- R^{17a} and R^{17b} independently represent H or C_{1-4} alkyl;
- E represents, at each occurrence when used herein, a direct bond or C_{1-2} alkylene;

Het¹ and Het³ to Het⁸ independently represent four- to ten-membered heterocyclic groups containing one to four heteroatoms selected from oxygen, nitrogen and/or sulfur, which groups are optionally substituted by one or more substituents selected from -OH, oxo, halo, cyano, C₁₋₄ alkyl, C₁₋₄ alkoxy, phenyl, -N(H)R^{18a}, -C(O)R^{18c}, -N(H)C(O)R^{18h} and -N(H)S(O)₂R^{18j};

R^{18a}, R^{18c}, R^{18h} and R^{18j} independently represent C₁₋₄ alkyl or optionally substituted aryl or R^{18a}, R^{18c} and R^{18h} independently represent H;

A represents C₁₋₄ alkylene optionally substituted by one or more substituents selected from OH and amino, -C₁₋₃ *n*-alkylene-S(O)₂N(H)- or -C₂₋₃ *n*-alkylene-N(H)S(O)₂- (in which latter two groups, alkylene is attached to the oxabispidine ring nitrogen);

B represents -Z-, -Z-N(H)-, -Z-C(O)N(R^{20b})-, -Z-N(R^{20c})S(O)₂-, -Z-S(O)₂N(R^{20d})-, -Z-S(O)₂-, -Z-O- (in which latter six groups, Z is attached to the carbon atom bearing R² and R³);

Z represents a direct bond or C₁₋₄ alkylene;

R^{20b} represents H, C₁₋₄ alkyl or, together with a single R⁴ substituent at a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, R^{20b} represents C₂₋₄ alkylene;

R^{20c} and R^{20d} independently represent H or C₁₋₃ alkyl;

when G represents N, G is in the *ortho*- or, in particular, the *para*-position relative to the point of attachment of B;

when G represents N, R⁴ is absent (i.e. represents H) or represents a single cyano group;

R⁴ represents one or more substituents selected from -OH, cyano, halo, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, -C(O)N(R^{22e})R^{22f}, and -N(R^{22m})S(O)₂-C₁₋₄ alkyl, or an R⁴ substituent in a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached may, together with R^{20b}, represent C₂₋₄ alkylene;

R^{21c} represents C_{1-4} alkyl;

R^{22c} , R^{22e} , R^{22f} and R^{22m} independently represent H or C_{1-4} alkyl;

R^{41} to R^{46} independently represent H;

optional substituents on aryl and aryloxy groups are, unless otherwise stated,

- 5 one or more substituents selected from halo, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), $-N(H)S(O)_2R^{21b}$ and $-S(O)_2N(H)R^{22o}$.

More preferred compounds of the invention include those in which:

- 10 R^1 represents straight- or branched-chain or part cyclic/acyclic C_{1-6} alkyl, which alkyl group is optionally interrupted by oxygen and/or substituted by:
(i) one or more halo or OR^{5c} groups; and/or (ii) one group selected phenyl (which latter group is optionally substituted by one or more (e.g. one to three) substituents selected from halo, cyano, C_{1-3} alkyl, C_{1-3} alkoxy (which
15 latter two groups are optionally substituted by one or more halo (e.g. fluoro) atoms), $-C(O)-C_{1-3}$ alkyl and $-S(O)_2-C_{1-4}$ alkyl), Het^1 , $-C(O)R^{5b}$, $-N(H)R^6$, $-C(O)N(R^{8a})R^{5d}$, $-OC(O)N(H)R^{8b}$, $-S(O)_2-C_{1-4}$ alkyl, $-S(O)_2N(H)R^{9c}$ and $-N(H)S(O)_2R^{9d}$ or R^1 represents $-C(O)OR^7$, $-C(O)N(R^{8a})R^{5d}$ or $-S(O)_2-C_{1-5}$ alkyl;
- 20 Het^1 represents a four- (e.g. five-) to ten-membered heterocyclic group containing one to three heteroatoms selected from oxygen, nitrogen and/or sulfur, which group is optionally substituted by one or more (e.g. one to three) substituents selected from halo, C_{1-3} alkyl, C_{1-3} alkoxy and $-C(O)-C_{1-4}$ alkyl;
- 25 R^{5b} , R^{5c} and R^{5d} independently represent H, C_{1-5} alkyl, phenyl (which latter group is optionally substituted by one or more substituents selected from $-OH$, halo, cyano, pyrrolidin-1-yl, C_{1-4} alkyl and C_{1-5} alkoxy (which latter group is optionally substituted by one or more halo (e.g. fluoro) atoms)) or Het^3 ;

Het³ represents a five- to ten-membered heterocyclic group containing one or two heteroatoms selected from oxygen and nitrogen, which group is optionally substituted by one or more substituents selected from oxo, C₁₋₂ alkyl and -C(O)-C₁₋₄ alkyl;

- 5 R⁶ represents H, C₁₋₄ alkyl, phenyl (which latter group is optionally substituted by one or more cyano groups) or -C(O)O-C₁₋₅ alkyl;

R⁷ represents C₁₋₅ alkyl optionally substituted by Het⁴;

- Het⁴ represents a five- to ten-membered heterocyclic group containing one or two heteroatoms selected from oxygen and nitrogen, which group is
10 optionally substituted by one or more substituents selected from C₁₋₂ alkyl and -C(O)-C₁₋₄ alkyl;

R^{8a} and R^{8b} independently represent H, C₁₋₅ alkyl or -D-(phenyl), the phenyl part of which latter group is optionally substituted by one or more (e.g. one to three) substituents selected from halo, C₁₋₃ alkyl and C₁₋₃ alkoxy;

- 15 D represents C₁₋₃ alkylene (e.g. CH₂ or C(CH₃)₂);

- R^{9c} and R^{9d} independently represent C₁₋₄ alkyl (optionally substituted by one or more substituents selected from halo, phenyl (which latter group is optionally substituted by one or more substituents selected from halo, C₁₋₂ alkyl and C₁₋₂ alkoxy (which latter two groups are optionally substituted
20 by one or more halo (e.g. fluoro) atoms)) and Het⁶); phenyl (which latter group is optionally substituted by one or more substituents selected from halo, C₁₋₃ alkyl and C₁₋₃ alkoxy (which latter two groups are optionally substituted by one or more halo (e.g. fluoro) atoms)) or Het⁷;

- Het⁶ and Het⁷ independently represent four- (e.g. five-) to ten-membered
25 heterocyclic groups containing one to three heteroatoms selected from oxygen, nitrogen and/or sulfur, which groups are optionally substituted by one or more (e.g. one to three) substituents selected from halo, C₁₋₃ alkyl and C₁₋₃ alkoxy;

R² represents H, -OR¹³ or -N(H)R¹⁴;

R³ represents H;

R¹³ represents H or phenyl (which latter group is optionally substituted by one or more substituents selected from cyano and C₁₋₂ alkoxy);

R¹⁴ represents H phenyl (which latter group is optionally substituted by one or more cyano groups) or -C(O)O-C₁₋₅ alkyl;

A represents C₁₋₃ *n*-alkylene;

B represents -Z-, -N(H)S(O)₂-, -S(O)₂N(H)-, -Z-N(H)-, -Z-C(O)N(R^{20b})-, -Z-S(O)₂-, -Z-O- (in which latter four groups, Z is attached to the carbon atom bearing R² and R³);

Z represents a direct bond or C₁₋₃ alkylene;

R^{20b}, together with a single R⁴ substituent at a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, represents C₂₋₃ alkylene;

G represents CH;

R⁴ represents one or two cyano or halo (e.g. fluoro) groups in the *ortho*- and/or, particularly, the *para*-position relative to the point of attachment of the group B, or alternatively, when B represents -Z-C(O)N(R^{20b})-,

- (i) an R⁴ substituent in a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached may, together with R^{20b}, represent C₂₋₃ alkylene, and
- (ii) R⁴ may further represent a nitro group in the *para*-position relative to the point of attachment of the group B.

Particularly preferred compounds of the invention include those in which:

- R¹ represents straight- or branched-chain or part cyclic/acyclic C₁₋₆ alkyl, which alkyl group is optionally interrupted by oxygen and/or substituted by:
- (i) one or more halo or OR^{5c} groups; and/or
 - (ii) one group selected phenyl (which latter group is optionally substituted by one or more (e.g. one to three) substituents selected from halo, cyano, C₁₋₂ alkyl, C₁₋₂ alkoxy (which

latter two groups are optionally substituted by one or more halo (e.g. fluoro) atoms), $-C(O)-C_{1-2}$ alkyl and $-S(O)_2-C_{1-2}$ alkyl), Het^1 , $-C(O)R^{5b}$, $-N(H)R^6$, $-C(O)N(H)R^{8a}$, $-OC(O)N(H)-C_{1-4}$ alkyl, $-S(O)_2-C_{1-4}$ alkyl, $-S(O)_2N(H)-C_{1-4}$ alkyl and $-N(H)S(O)_2R^{9d}$;

- 5 Het^1 represents a five- or six-membered heterocyclic group containing one or two heteroatoms selected from oxygen, nitrogen and/or sulfur, which group is optionally substituted by one or more (e.g. one or two) substituents selected from halo (e.g. chloro), C_{1-2} alkyl and C_{1-2} alkoxy;

R^{5b} and R^{5c} independently represent phenyl optionally substituted by one to
10 three substituents selected from halo, cyano, C_{1-2} alkyl and C_{1-2} alkoxy;

R^6 represents H, C_{1-4} alkyl or $-C(O)O-C_{1-5}$ alkyl;

R^{8a} represents C_{1-4} alkyl (e.g. *tert*-butyl) or $-D$ -(phenyl);

- R^{9d} represents C_{1-4} alkyl (optionally substituted by one or more substituents selected from halo and phenyl (which latter group is optionally substituted
15 by one or more substituents selected from halo (e.g. chloro), C_{1-2} alkyl and C_{1-2} alkoxy (which latter two groups are optionally substituted by one or more halo (e.g. fluoro) atoms))), phenyl (which latter group is optionally substituted by one or more substituents selected from halo (e.g. fluoro), C_{1-2} alkyl and C_{1-2} alkoxy (which latter two groups are optionally substituted by
20 one or more halo (e.g. fluoro) atoms)) or Het^7 ;

Het^7 represents a five- or six-membered heterocyclic group containing one nitrogen atom and optionally containing one or two further heteroatoms selected from oxygen, nitrogen and/or sulfur, which heterocyclic group is optionally substituted by one to three substituents selected from halo and
25 C_{1-2} alkyl;

A represents $-(CH_2)_{1-2}-$;

B represents $-Z-$, $-N(H)S(O)_2-$ (in which latter group, $-N(H)$ is attached to the carbon atom bearing R^2 and R^3), $-C(O)N(R^{20b})-$ (in which latter group,

-C(O) is attached to the carbon atom bearing R^2 and R^3) or -Z-O- (in which latter group, Z is attached to the carbon atom bearing R^2 and R^3);

Z represents a direct bond or $-(CH_2)_{1-2}-$;

R^{20b} , together with a single R^4 substituent at a position on the phenyl or
 5 pyridyl group that is *ortho*- to the position at which the group B is attached, represents $-(CH_2)_{2-3}-$;

R^4 represents a cyano or fluoro group in the *para*-position relative to the point of attachment of the group B, and, when B represents $-C(O)N(R^{20b})-$, R^4 may also, together with R^{20b} , represent $-(CH_2)_{2-3}-$.

10

Especially preferred compounds of the invention include those in which:

R^1 represents straight- or branched-chain C_{1-3} alkyl substituted by OR^{5c} , phenyl (which latter group is optionally substituted by one or two substituents selected from halo (e.g. fluoro or chloro), cyano, methyl,
 15 methoxy (which latter two groups are optionally substituted by one to three fluoro atoms), $-C(O)CH_3$ and $-S(O)_2CH_3$), Het^1 , $-C(O)R^{5b}$, $-N(H)C(O)O-C_{3-4}$ alkyl, $-C(O)N(H)R^{8a}$, $-OC(O)N(H)-C_{3-4}$ alkyl, $-S(O)_2N(H)-C_{3-4}$ alkyl or $-N(H)S(O)_2R^{9d}$;

Het^1 represents an aromatic five- or six-membered heterocyclic group
 20 containing one or two heteroatoms selected from oxygen, nitrogen and/or sulfur, which group is optionally substituted by one or two substituents selected from chloro, methyl and methoxy;

R^{5b} and R^{5c} independently represent phenyl optionally substituted by one or two substituents selected from cyano, methyl and methoxy;

25 R^{8a} represents *tert*-butyl, CH_2 -phenyl or $C(CH_3)_2$ -phenyl;

R^{9d} represents C_{1-4} alkyl (which latter group is optionally substituted by one or more fluoro atoms), $(CH_2)_{1-2}$ -phenyl (the phenyl part of which latter group is optionally substituted by one to three substituents selected from chloro, methyl and methoxy (which latter two groups are optionally

substituted by one or more fluoro atoms)), phenyl (optionally substituted by one or more substituents selected from fluoro, methyl and methoxy (which latter two groups are optionally substituted by one or more fluoro atoms)) or Het⁷;

- 5 Het⁷ represents a five-membered heterocyclic group containing one nitrogen atom and optionally containing one further heteroatom selected from oxygen, nitrogen and sulfur, which heterocyclic group is optionally substituted by one to three methyl groups;

R² represents H or -OH;

- 10 A represents -CH₂-;

B represents -CH₂-, -N(H)S(O)₂- (in which latter group, -N(H) is attached to the carbon atom bearing R² and R³) or -(CH₂)₀₋₁-O- (in which latter group, -CH₂- is attached to the carbon atom bearing R² and R³).

- 15 Compounds of formula I that are more preferred still include those in which:

R¹ represents

- (i) C₁₋₃ *n*-alkyl substituted by phenyl (which latter group is optionally substituted by one to three substituents selected from methyl and methoxy (which latter two groups are optionally substituted by one to
20 three (e.g. by two) fluoro atoms)) or phenoxy (the phenyl part of which latter group is optionally substituted by cyano),

(ii) -N(H)C(O)O-*tert*-butyl,

(iii) -S(O)₂N(H)-*tert*-butyl or

(iv) -N(H)S(O)₂R^{9d};

- 25 R^{9d} represents

- (a) methyl (optionally substituted by one to three fluoro atoms),
(b) isopropyl,
(c) *n*-butyl,


- (d) $-(CH_2)$ -phenyl (the phenyl part of which latter group is optionally substituted by one or two substituents selected from chloro, methyl and trifluoromethyl),
- (e) phenyl (which latter group is optionally substituted by one or two substituents selected from fluoro, methoxy and trifluoromethoxy) or
- (f) imidazolyl (e.g. imidazol-4-yl) or isoxazolyl (e.g. isoxazol-4-yl), which latter two groups are optionally substituted by one or two methyl groups.
- Particular values of each variable group are as follows. Such values may be used where appropriate with any of the values, definitions, claims, aspects or embodiments defined hereinbefore or hereinafter. In particular, each may be used as an individual limitation on the broadest definition of formula (I).
- R^1 represents C_{2-3} alkyl, which alkyl group is substituted by at least one - $S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group;
- R^{9b} represents H or C_{1-3} alkyl;
- R^{9c} and R^{9d} each independently represent hydrogen, C_{1-6} alkyl (optionally substituted by one or more halo groups), aryl (optionally substituted by one or more halo, cyano, methoxy, fluomethoxy, difluoromethoxy or trifluoromethoxy groups) or Het⁷ (such as imidazolyl or isoxazolyl);
- R^2 and R^3 each independently represent hydrogen, or hydroxy;
- A represents a direct bond or C_{1-3} alkylene;
- B represents a direct bond, C_{1-3} alkylene or C_{1-3} alkoxy (in which the oxygen is attached to the phenyl group that is optionally substituted with R^4);
- G represents carbon;
- R^{41} to R^{46} represents hydrogen; and
- R^4 represents one or more optional substituents selected from cyano and/or halo (such as fluoro) and an R^4 substituent is in a position on the phenyl


group that is *ortho*- and/or *para*- to the position at which the group B is attached.

Particular compounds of the invention include, for example, compounds of the Formula I, or pharmaceutically-acceptable salts thereof, wherein, unless
5 otherwise stated, each of each variable group has any of the meanings defined hereinbefore or in paragraphs (a) to (d) hereinafter :-


(a) R¹ represents a C₂-C₅ alkyl group, (such as ethyl, propyl, butyl, propyl or pentyl), which is optionally substituted by one or more sulfonamide groups
10 selected from 4-cyanobenzenesulfonamide, propane-2-sulfonamide, 1-phenylmethanesulfonamide, propane-1-sulfonamide, 4-fluorobenzenesulfonamide; benzenesulfonamide, 2,4-difluorobenzenesulfonamide, methane-sulfonamide, 3-chloro-1-phenylmethanesulphonamide, trifluoromethane-sulfonamide, 1-methyl-1H-imidazole-4-sulfonamide, butane-1-sulfonamide,
15 1-[4-(trifluoromethyl)phenyl]methanesulfonamide, 3,5-dimethylisoxazole-4-sulfonamide, 2-(trifluoromethoxy)benzenesulfonamide, [2-(trifluoromethoxy)phenyl]methanesulfonamide, 2,3-dihydro-1-benzofuran-5-sulfonamide, 2-cyanobenzenesulfonamide, 4-methoxybenzenesulfonamide, 5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonamide, 3-fluorobenzene-sulphonamide, 5-methylisoxazole-4-sulfonamide, 3-cyanobenzene-sulfonamide, 4-cyano-1-phenylmethanesulphonamide, or 2-fluorobenzene-sulphonamide;


(b) R⁴¹ to R⁴⁶ are hydrogen;



(c) the group  represents methylene, ethylene, propylene,
25 butylene, or 2-hydroxypropylene, optionally terminated or interrupted

with an oxygen atom and/ or optionally interrupted with a $-\text{SO}_2\text{-NH-}$ or $-\text{NH-SO}_2\text{-}$ group;



- (d) the group  represents phenyl, 4-cyanophenyl, 3,4-bis(difluoromethoxy)phenyl, 4-fluorophenyl, 4-(difluoromethoxy)-phenyl, 2-fluorophenyl, 2,5-difluorophenyl, 3,4-difluorophenyl, 4-cyano-2-fluorophenyl, 3,4-difluorophenyl, 2-cyanophenyl, 3-fluorophenyl, 4-cyano-2,6-difluorophenyl, 2,6-difluorophenyl, 3-cyanophenyl, 4-chlorophenyl, 4-(trifluoromethyl)phenyl, 2,4-difluorophenyl, 2-(trifluoromethoxy)phenyl, 3-chlorophenyl, or 4-methoxyphenyl.

Preferred compounds of the invention include the compounds of the Examples disclosed hereinafter. In this respect, preferred compounds of the invention that might be mentioned include:

- (i) N-(2-{7-[(2S)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)propane-2-sulfonamide;
- (ii) N-(2-{7-[(2S)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- (iii) N-(tert-butyl)-3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propane-1-sulfonamide;
- (iv) tert-butyl {2-[7-(2-[(4-fluorophenyl)sulfonyl]amino)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl}carbamate;
- (v) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-fluorobenzenesulfonamide;
- (vi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;

- (vii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)propane-2-sulfonamide;
- (viii) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyanobenzenesulfonamide;
- 5 (ix) N-[2-(7-{2-[3,4-bis(difluoromethoxy)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyanobenzenesulfonamide;
- (x) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-
- 10 [3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- (xii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xiii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- 15 (xiv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- (xvi) 4-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-
- 20 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xvii) 1-(3-chlorophenyl)-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- (xviii) 4-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (xix) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1,1,1-trifluoromethanesulfonamide;
- (xx) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;

- (xxi) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-methyl-1H-imidazole-4-sulfonamide;
- (xxii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)butane-1-sulfonamide;
- 5 (xxiii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-methyl-1H-imidazole-4-sulfonamide;
- (xxiv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-[4-(trifluoromethyl)phenyl]methane-sulfonamide;
- 10 (xxv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (xxvi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-[4-(trifluoromethyl)phenyl]methane-sulfonamide;
- 15 (xxvii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (xxviii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-2-(trifluoromethoxy)benzenesulfonamide;
- (xxix) 1-(3-chlorophenyl)-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-
- 20 diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- (xxx) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxi) 4-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl}benzenesulfonamide;
- 25 (xxxii) 4-cyano-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl}benzenesulfonamide;
- (xxxiii) 4-cyano-N-(2-{7-[4-(difluoromethoxy)benzyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (xxxiv) 4-cyano-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxv) 4-cyano-N-(2-{7-[2-(4-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 5 (xxxvi) 4-cyano-N-(2-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxvii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (xxxviii) 5-chloro-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,3-dimethyl-1H-pyrazole-4-sulfonamide;
- 10 (xxxix) 2-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xl) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-methoxybenzenesulfonamide;
- 15 (xli) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (xlii) 5-chloro-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,3-dimethyl-1H-pyrazole-4-sulfonamide;
- 20 (xliii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-methoxybenzenesulfonamide;
- (xliv) 4-cyano-N-(2-{7-[3-(2-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (xlv) 4-cyano-N-(2-{7-[3-(2,5-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlv) 4-cyano-N-(2-{7-[3-(3,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (xlvii) 4-cyano-N-(2-{7-[2-(2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlviii) 4-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 5 (xlix) 4-cyano-N-[2-(7-{2-[(4-fluorobenzyl)oxy]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- (l) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (II) 4-cyano-N-(2-{7-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (lii) 4-cyano-N-{2-[7-(2-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (liii) 4-cyano-N-(2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (liv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- (lv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- (lvi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-5-methylisoxazole-4-sulfonamide;
- 20 (lvii) N-{2-[7-(1-benzofuran-3-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (lviii) 4-cyano-N-{2-[7-(1H-indol-3-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 25 (lix) 4-cyano-N-(2-{7-[(1-methyl-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lx) 4-cyano-N-(2-{7-[(5-fluoro-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (lxi) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-4-fluorobenzenesulfonamide;
- (lxii) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-2,4-difluorobenzenesulfonamide;
- 5 (lxiii) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (lxiv) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-1-phenylmethanesulfonamide;
- (lxv) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
- 10 [3.3.1]non-3-yl}propyl)-1-phenylmethanesulfonamide;
- (lxvi) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (lxvii) 4-cyano-N-(2-{7-[(2-methyl-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (lxviii) 4-cyano-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxix) 4-cyano-N-(2-{7-[3-(3-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxx) 4-cyano-N-(2-{7-[3-(4-cyano-2-fluorophenoxy)propyl]-9-oxa-3,7-
- 20 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxi) 4-cyano-N-(2-{7-[3-(4-cyano-2,6-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxii) 4-cyano-N-(2-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (lxxiii) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)propane-2-sulfonamide;
- (lxxiv) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-1-phenylmethanesulfonamide;

- (lxxv) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxvi) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-3,5-dimethylisoxazole-4-sulfonamide;
- 5 (lxxvii) N-{3-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxviii) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxix) N-{2-[7-(1,3-benzoxazol-2-ylmethyl)-9-oxa-3,7-diazabicyclo-
- 10 [3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (lxxx) 3-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxxi) 3-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (lxxxii) 3-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (lxxxiii) 2-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (lxxxiv) 2-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-
- 20 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxxv) N-(4-cyanobenzyl)-2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl]ethanesulfonamide;
- (lxxxvi) N-(4-cyanobenzyl)-2-[7-(2-cyanobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl]ethanesulfonamide;
- 25 (lxxxvii) N-(4-cyanobenzyl)-N-[(3,5-dimethylisoxazol-4-yl)methyl]-2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethanesulfonamide;
- (lxxxviii) 4-cyano-N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;

- (lxxxix) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-4-fluorobenzenesulfonamide;
- (xc) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2-fluorobenzenesulfonamide;
- 5 (xci) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2-fluorobenzenesulfonamide;
- (xcii) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2,4-difluorobenzene-sulfonamide;
- 10 (xciii) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (xciv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xcv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- 15 (xcvi) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (xcvii) 3-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 20 (xcviii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xcix) N-(2-{7-[2-(4-chlorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyanobenzenesulfonamide;
- 25 (c) 4-cyano-N-[2-(7-{2-[4-(trifluoromethyl)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- (ci) 4-cyano-N-(2-{7-[2-(2,6-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (cii) 4-cyano-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (ciii) 4-cyano-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 5 (civ) 4-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- (cv) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyano-N-methylbenzenesulfonamide;
- (cvi) 4-cyano-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- 10 (cvii) 4-cyano-N-methyl-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (cviii) 4-cyano-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 15 (cix) 4-cyano-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cx) N-(4-cyanobenzyl)-2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (cxi) N-(4-cyanobenzyl)-2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- 20 (cxii) N-(4-cyanobenzyl)-2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (cxiii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N,3,5-trimethylisoxazole-4-sulfonamide;
- 25 (cxiv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- (cxv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;

- (cxvi) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- (cxvii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- 5 (cxviii) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-fluorobenzenesulfonamide;
- (cxix) 4-fluoro-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (cxx) N-(2-{7-[2-(4-cyanophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- 10 (cxxi) 4-fluoro-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxxii) 4-cyano-N-[2-(7-{2-[4-(difluoromethoxy)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- 15 (cxxiii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluoro-N-methylbenzenesulfonamide;
- (cxxiv) 4-cyano-N-(2-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 20 (cxxv) 4-cyano-N-(2-{7-[2-(2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxvi) N-(2-{7-[2-(1,2-benzisoxazol-3-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyanobenzenesulfonamide;
- (cxxvii) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-N-methylbenzenesulfonamide;
- 25 (cxxviii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methyl-2,3-dihydro-1-benzofuran-5-sulfonamide;

- (cxxxix) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]-non-3-yl}ethyl)-N-methyl-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (cxxx) 4-cyano-N-{2-[7-(4-cyano-2-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 5 (cxxxixi) 4-cyano-N-{2-[7-(4-cyano-2-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- (cxxxixii) 4-cyano-N-[2-(7-{2-[4-(difluoromethoxy)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-N-methylbenzenesulfonamide;
- (cxxxixiii) 4-fluoro-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (cxxxixiv) 4-fluoro-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxxxixv) 4-fluoro-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 15 (cxxxixvi) N-{2-[7-(1,2-benzisoxazol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl}-4-cyanobenzenesulfonamide;
- (cxxxixvii) 4-cyano-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxxixviii) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 20 (cxxxixix) 4-cyano-N-{3-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzenesulfonamide;
- (cxli) 4-cyano-N-{3-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzenesulfonamide;
- 25 (cxlii) 4-cyano-N-(3-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxliii) 4-cyano-N-(3-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;

- (cxliii) 4-cyano-N-(3-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxliv) 4-cyano-N-(3-{7-[2-(2,6-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 5 (cxlv) 4-cyano-N-(3-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlvi) N-[3-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propyl]-4-cyanobenzenesulfonamide;
- (cxlvii) 4-cyano-N-(3-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 10 (cxlviii) 4-cyano-N-(3-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlix) 4-cyano-N-(2-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 15 (cl) 4-cyano-N-(2-{7-[3-(2,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cli) 4-cyano-N-(2-{7-[2-(2,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (clii) 4-cyano-N-(2-{7-[3-(2,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 20 (cliii) N-(2-{7-[2-(4-chlorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyano-N-methylbenzenesulfonamide; and
- (cliv) 4-Cyano-N-(2-{7-[2-(2,4-difluoro-phenyl)-ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-ethyl)-N-methyl-benzenesulfonamide;


25

or a pharmaceutically acceptable salt thereof.

Preparation

According to the invention there is also provided a process for the preparation of compounds of formula I which comprises:

(a) reaction of a compound of formula II,



wherein R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as hereinbefore defined, with a compound of formula III,




III

wherein L^1 represents a leaving group such as halo, alkanesulfonate, perfluoroalkanesulfonate, arenesulfonate, $-OC(O)XR^7$, imidazole or $R^{23}O-$ (wherein R^{23} represents, for example, C_{1-10} alkyl or aryl, which groups are optionally substituted by one or more halo or nitro groups) and X, R^1 and R^7 are as hereinbefore defined, for example at between room and reflux temperature in the presence of a suitable base (e.g. triethylamine, potassium carbonate or a bicarbonate, such as sodium bicarbonate) and an appropriate solvent (e.g. dichloromethane, chloroform, acetonitrile, *N,N*-dimethylformamide, THF, toluene, water, a lower alkyl alcohol (e.g. ethanol) or mixtures thereof);

(b) for compounds of formula I in which R^1 represents C_{1-12} alkyl substituted by one or more substituents as defined above in respect of R^1 , which

37

substituent(s) is/include a $-N(R^{9b})S(O)_2R^{9d}$ group, reaction of a compound of formula IV,




wherein R^{1a} represents C_{1-12} alkylene, which group is optionally substituted by one or more substituents as defined above in respect of R^1 , and R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as hereinbefore defined, with a compound of formula V,



wherein L^2 represents a suitable leaving group such as halo and R^{9d} is as hereinbefore defined, for example under conditions that are known to those skilled in the art (e.g. at ambient temperature (such as from 15 to 30°C) in the presence of a suitable base (such as triethylamine, potassium carbonate or sodium hydrogencarbonate) and an appropriate solvent (such as DCM, $CHCl_3$, acetonitrile, DMF, THF, toluene, or mixtures thereof);

15

(c) for compounds of formula I in which R^1 represents C_{1-12} alkyl substituted by one or more substituents as defined above in respect of R^1 , which substituent(s) is/include a $-S(O)_2N(R^{9b})R^{9c}$ group, reaction of a compound of formula II, as hereinbefore defined, with a compound of formula VA,




20

wherein L^1 , R^{1a} , R^{9b} and R^{9c} are as hereinbefore defined, for example under conditions that are known to those skilled in the art (e.g. at ambient temperature

to reflux in the presence of a suitable base (such as triethylamine, potassium carbonate or sodium hydrogencarbonate) and an appropriate solvent (such as DCM, CHCl_3 , acetonitrile, DMF, THF, toluene, or mixtures thereof);

5

(d) for compounds of formula I in which R^1 represents $-\text{C}(\text{O})\text{XR}^7$ or $-\text{C}(\text{O})\text{N}(\text{R}^{8a})\text{R}^{5d}$, reaction of a compound of formula VI,



wherein R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B, G and L^1 are as hereinbefore defined,
10 with a compound of formula VII,



wherein R^{24} represents $-\text{XR}^7$ or $-\text{N}(\text{R}^{8a})\text{R}^{5d}$ and R^{5d} , R^7 , R^{8a} and X are as hereinbefore defined, for example under similar conditions to those described hereinbefore (process step (a));

15


(e) for compounds of formula I in which R^1 represents $-\text{C}(\text{O})\text{N}(\text{H})\text{R}^{8a}$, reaction of a compound of formula II, as hereinbefore defined, with a compound of formula VIII,



20 wherein R^{8a} is as hereinbefore defined, for example at between 0°C and reflux temperature in the presence of an appropriate organic solvent (e.g.


dichloromethane), or *via* solid phase synthesis under conditions known to those skilled in the art;

(f) reaction of a compound of formula IX,



5

wherein R^1 and R^{41} to R^{46} are as hereinbefore defined, with a compound of formula X,




wherein L^3 represents a leaving group such as halo, alkanesulfonate (e.g. mesylate), perfluoroalkanesulfonate or arenesulfonate (e.g. 2- or 4-nitrobenzenesulfonate, toluenesulfonate or benzenesulfonate) and R^2 , R^3 , R^4 , A, B and G are as hereinbefore defined, for example at elevated temperature (e.g. between 35°C and reflux temperature) in the presence of a suitable base (e.g. triethylamine or potassium carbonate) and an appropriate organic solvent (e.g. acetonitrile, dichloromethane, chloroform, dimethylsulfoxide, *N,N*-dimethylformamide, a lower alkyl alcohol (e.g. ethanol), isopropyl acetate or mixtures thereof);

(g) for compounds of formula I in which B represents $-Z-N(R^{20c})S(O)_2-$, reaction of a compound of formula IX, as hereinbefore defined, with a compound of formula XA,


20

40




wherein R^2 , R^3 , R^4 , A, G, Z and L^3 are as hereinbefore defined, for example under conditions described in respect of process (f) above;

- 5 (h) for compounds of formula I in which A represents CH_2 and R^2 represents $-OH$ or $-N(H)R^{14}$, reaction of a compound of formula IX, as hereinbefore defined, with a compound of formula XI,




- wherein Y represents O or $N(R^{14})$ and R^3 , R^4 , R^{14} , B and G are as
 10 hereinbefore defined, for example at elevated temperature (e.g. $60^\circ C$ to reflux) in the presence of a suitable solvent (e.g. a lower alkyl alcohol (e.g. IPA), acetonitrile, or a mixture of a lower alkyl alcohol and water);

- (i) for compounds of formula I in which B represents $-Z-O-$, reaction of a
 15 compound of formula XII,




wherein R^1 , R^2 , R^3 , R^{41} to R^{46} , A and Z are as hereinbefore defined, with a compound of formula XIII,



- 5 wherein R^4 and G are as hereinbefore defined, for example under Mitsunobu-type conditions e.g. at between ambient (e.g. 25°C) and reflux temperature in the presence of a tertiary phosphine (e.g. tributylphosphine or triphenylphosphine), an azodicarboxylate derivative (e.g. diethylazodicarboxylate or 1,1'-(azodicarbonyl)dipiperidine) and an
- 10 appropriate organic solvent (e.g. dichloromethane or toluene);


(j) for compounds of formula I in which G represents N and B represents -Z-O-, reaction of a compound of formula XII, as hereinbefore defined, with a compound of formula XIV,



15

wherein R^4 and L^3 are as hereinbefore defined, for example at between 10°C and reflux temperature in the presence of a suitable base (e.g. sodium hydride) and an appropriate solvent (e.g. *N,N*-dimethylformamide);


(k) for compounds of formula I in which R^2 represents $-OR^{13}$, in which R^{13} represents C_{1-6} alkyl, $-E$ -aryl or $-E$ -Het⁸, reaction of a compound of formula I in which R^2 represents OH with a compound of formula XV,



XV

- 5 wherein R^{13a} represents C_{1-6} alkyl, $-E$ -aryl or $-E$ -Het⁸ and E and Het⁸ are as hereinbefore defined, for example under Mitsunobu-type conditions (e.g. as described hereinbefore in process step (h));


- (l) for compounds of formula I in which R^2 represents $-OR^{13}$, in which R^{13} represents C_{1-6} alkyl, $-E$ -aryl or $-E$ -Het⁸, reaction of a compound of formula XVI,



XVI

- wherein R^1 , R^3 , R^4 , R^{41} to R^{46} , A , B , G and L^3 are as hereinbefore defined, with a compound of formula XV, as hereinbefore defined, for example at
 15 between ambient (e.g. 25°C) and reflux temperature, under Williamson-type conditions (i.e. in the presence of an appropriate base (e.g. KOH or NaH) and a suitable organic solvent (e.g. dimethylsulfoxide or *N,N*-dimethylformamide)) (the skilled person will appreciate that certain compounds of formula XVI (e.g. those in which L^3 represents halo) may also be regarded as
 20 compounds of formula I as hereinbefore defined);

(m) for compounds of formula I in which R^2 represents $-E-NH_2$, reduction of a compound of formula XVII,



wherein R^1 , R^3 , R^4 , R^{41} to R^{46} , A, B, E and G are as hereinbefore defined, for example by hydrogenation at a suitable pressure in the presence of a suitable catalyst (e.g. palladium on carbon) and an appropriate solvent (e.g. a water-ethanol mixture);

(n) for compounds of formula I in which R^2 represents $-E-N(R^{14})R^{15}$, wherein R^{14} represents C_{1-6} alkyl, $-E$ -aryl $-E$ -Het⁹, $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$ or $-C(O)N(R^{17a})R^{17b}$, reaction of a compound of formula I in which R^2 represents $-E-N(H)R^{15}$ with a compound of formula XVIII,



wherein R^{14a} represents C_{1-6} alkyl, $-E$ -aryl $-E$ -Het⁹, $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$ or $-C(O)N(R^{17a})R^{17b}$, and R^{16a} , R^{16b} , R^{16c} , R^{17a} , R^{17b} , Het⁹, E and L^1 are as hereinbefore defined, for example under conditions described hereinbefore (process step (a));


(o) for compounds of formula I in which R^2 represents $-E-N(R^{15})C(O)N(H)R^{17a}$, reaction of a compound of formula I in which R^2 represents $-E-N(H)R^{15}$ with a compound of formula XIX,



wherein R^{17a} is as hereinbefore defined, for example under conditions described hereinbefore (process step (e));

(p) for compounds of formula I in which R^2 represents
 5 $-E-N(H)[C(O)]_2NH_2$, reaction of a compound of formula I in which R^2 represents $-E-NH_2$ with oxalic acid diamide, for example at between -10 and $25^\circ C$ in the presence of a suitable coupling agent (e.g. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide), an appropriate activating agent (e.g. 1-hydroxybenzotriazole), a suitable base (e.g. triethylamine) and a reaction-
 10 inert solvent (e.g. *N,N*-dimethylformamide);

(q) for compounds of formula I in which R^2 represents $-E-N(H)C(NH)NH_2$, reaction of a compound of formula I in which R^2 represents $-E-NH_2$ with a compound of formula XX,



or an *N*-protected derivative thereof, wherein R^{23} is as hereinbefore defined, for example at between room and reflux temperature, optionally in the presence of a suitable solvent (e.g. toluene) and/or an appropriate acidic catalyst (e.g. acetic acid at, for example, 10 mol%);

20

(r) for compounds of formula I in which R^2 represents $-OR^{13}$, in which R^{13} represents $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or $-C(O)N(R^{17a})R^{17b}$, reaction of a compound of formula I in which R^2 represents $-OH$ with a compound of formula XXI,



wherein R^{13b} represents $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or $-C(O)N(R^{17a})R^{17b}$, L^4 represents a leaving group such as halo, *p*-nitrophenoxy, $-OC(O)R^{16a}$, $-OC(O)OR^{16b}$, $-OH$ or imidazole and R^{16a} , R^{16b} , R^{17a} and R^{17b} are as hereinbefore defined, for example at between $-10^\circ C$ and reflux temperature

in the presence of a suitable base (e.g. triethylamine, pyridine or potassium carbonate), an appropriate organic solvent (e.g. THF, dichloromethane or acetonitrile) and (where appropriate) a suitable coupling agent (e.g. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide);

5

- (s) for compounds of formula I in which R^2 represents H or -OH and R^3 represents H, reduction of a compound of formula I in which R^2 and R^3 together represent =O, in the presence of a suitable reducing agent and under appropriate reaction conditions; for example, for formation of compounds of formula I in which R^2 represents OH, reduction may be performed under mild reaction conditions in the presence of e.g. sodium borohydride and an appropriate organic solvent (e.g. THF); for formation of compounds of formula I in which R^2 represents OH, wherein the compound is enantiomerically enriched (or is a single enantiomer) at the chiral centre to which R^2 is attached, reduction may be performed enzymatically (for example under conditions known to those skilled in the art, such as in the presence of horse liver alcohol dehydrogenase and NADPH) or by hydrogenation in the presence of a suitable solution-phase (homogeneous) catalyst under conditions known to those skilled in the art; and for formation of compounds of formula I in which R^2 represents H, reduction may be performed either under Wolff-Kischner conditions known to those skilled in the art or by activating the relevant C=O group using an appropriate agent (such as tosylhydrazine) in the presence of a suitable reducing agent (e.g. sodium borohydride or sodium cyanoborohydride) and an appropriate organic solvent (e.g. a lower (e.g. C_{1-6}) alkyl alcohol);


25

- (t) for compounds of formula I in which R^2 represents halo, substitution of a corresponding compound of formula I in which R^2 represents -OH, using an


46

appropriate halogenating agent (e.g. for compounds in which R^2 represents fluoro, reaction with (diethylamino)sulfur trifluoride);

- (u) for compounds of formula I in which R^2 and R^3 represent H, A represents -J- and B represents $-N(R^{20e})-Z-$ (wherein $-N(R^{20e})$ is attached to the carbon atom bearing R^2 and R^3), reaction of a compound of formula XXII,




wherein R^1 , R^{20e} , R^{41} to R^{46} and J are as hereinbefore defined, with a compound of formula XXIII,



10

wherein R^4 , G, Z and L^3 are as hereinbefore defined, for example at elevated temperature (e.g. 40°C to reflux) in the presence of a suitable organic solvent (e.g. acetonitrile);


- (v) for compounds of formula I in which A represents $-(CH_2)_2-$ and R^2 and R^3 together represent $=O$, reaction of a compound of formula IX, as hereinbefore defined, with a compound of formula XXIV,



wherein B, G and R⁴ are as hereinbefore defined, for example at between room and reflux temperature in the presence of a suitable base (e.g. triethylamine, potassium carbonate or tetrabutylammonium hydroxide) and an appropriate organic solvent (e.g. a lower alkyl (e.g. C₁₋₆) alcohol);


5

(w) for compounds of formula I in which R¹ represents -C(O)XR⁷, -C(O)N(R^{8a})R^{5d} or -S(O)₂R^{9a}, reaction of a compound of formula XXV,



wherein R^{1b} represents -C(O)XR⁷, -C(O)N(R^{8a})R^{5d} or -S(O)₂R^{9a} and R^{5d}, R⁷, R^{8a}, R^{9a}, R⁴¹ to R⁴⁶ and L³ are as hereinbefore defined, with a compound of formula XXVI,

10



wherein R², R³, R⁴, A, B and G are as hereinbefore defined, for example at between room and reflux temperature in the presence of a suitable base (e.g. sodium hydrogencarbonate or potassium carbonate) and an appropriate organic solvent (e.g. acetonitrile);

15

(x) for compounds of formula I which are oxabispidine-nitrogen N-oxide derivatives, oxidation of the corresponding oxabispidine nitrogen of a corresponding compound of formula I, in the presence of a suitable oxidising agent (e.g. *m*CPBA), for example at 0°C in the presence of a suitable organic solvent (e.g. dichloromethane);

20

(y) for compounds of formula I which are C₁₋₄ alkyl quaternary ammonium salt derivatives, in which the alkyl group is attached to a oxabispidine nitrogen, reaction, at the oxabispidine nitrogen, of a corresponding compound of formula I with a compound of formula XXVII,



wherein R²⁵ represents C₁₋₄ alkyl and L⁵ is a leaving group such as halo, alkanesulfonate or arenesulfonate, for example at room temperature in the presence of an appropriate organic solvent (e.g. *N,N*-dimethylformamide), followed by purification (using e.g. HPLC) in the presence of a suitable counter-ion provider (e.g. NH₄OAc);


10

(z) conversion of one R⁴ substituent to another using techniques well known to those skilled in the art; or

15 (aa) introduction of one or more (further) R⁴ substituents to the aromatic ring using techniques well known to those skilled in the art (e.g. chlorination).

(ab) for compounds of formula I in wherein R¹ represents C₁₋₁₂ alkylene, which group is optionally substituted by one or more substituents as defined


20 above in respect of R¹, reaction of a compound of formula II




II

wherein R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as hereinbefore defined, with the appropriate aldehyde, for example under conditions that are known to those skilled in the art (e.g. at room temperature, such as from 15 to 30°C) in the presence of a reducing agent (such as sodium cyanoborohydride, sodium triacetoxyborohydride, or similar compounds) and an appropriate solvent (such as 1,2-dichloroethane, dichloroethane, methanol, ethanol or mixtures thereof);

(ac) for compounds of formula I in wherein A represents -J-, -J-N(R^{19a})-, -J-S(O)₂N(R^{19b})-, -J-N(R^{19c})S(O)₂- or -J-O- (in which latter four groups, -J is attached to the oxabispidine ring nitrogen), reaction of a compound with formula VII



wherein R^1 and R^{41} to R^{46} are as hereinbefore defined, with a compound of formula XXIII



wherein R^2 , R^4 , B and G are as hereinbefore defined and the C₀₋₅ alkylene group is optionally substituted by one or more substituents as defined above in

respect of J, for example under conditions that are known to those skilled in the art (e.g. at room temperature, such as from 15 to 30°C) in the presence of a reducing agent (such as sodium cyanoborohydride, sodium triacetoxymethylborohydride, or similar hydride donating compounds) and an appropriate solvent (such as 1,2-dichloroethane, dichloroethane, methanol, ethanol or mixtures thereof); or

(ad) deprotection of a protected derivative of a compound of formula I as defined above.

Compounds of formulae II, IV, IX, X, XI, XII, XVI, XVII, XXII and XXV may be prepared according to or by analogy with the procedures described or referred to in WO 01/28992, WO 02/28863, WO 02/28864, WO 02/83690 and WO 02/83691, the disclosures of which documents are hereby incorporated by reference.

Compounds of formula VA may be prepared by reaction of a compound of formula XXVIII,



wherein R^{9b} and R^{9c} are as hereinbefore defined, with a compound of formula XXIX,




wherein L¹, L² and R^{1a} are as hereinbefore defined, for example under conditions known to those skilled in the art (e.g. those described in respect of process (b) above).

Compounds of formula XA may be prepared by reaction of a compound of formula XXX,



wherein A, L² and L³ are as hereinbefore defined, with a compound of formula XXXI,



XXXI

wherein R⁴, G and Z are as hereinbefore defined, for example under
5 conditions known to those skilled in the art (e.g. those described in respect of
process (b) above).

Compounds of formulae III, V, VII, VIII, XIII, XIV, XV, XVIII, XIX, XX,
XXI, XXIII, XXIV, XXVI, XXVII, XXVIII, XXIX, XXX, XXXI and
10 derivatives thereof, are either commercially available, are known in the
literature, or may be obtained either by analogy with the processes described
herein, or by conventional synthetic procedures, in accordance with standard
techniques, from readily available starting materials using appropriate
reagents and reaction conditions.

15

Substituents on the aryl (e.g. phenyl), and (if appropriate) heterocyclic,
group(s) in compounds defined herein may be converted to other claimed
substituents using techniques well known to those skilled in the art. For
example, hydroxy may be converted to alkoxy, phenyl may be halogenated to
20 give halophenyl, nitro may be reduced to give amino, halo may be displaced
by cyano, etc.

The skilled person will also appreciate that various standard substituent or
functional group interconversions and transformations within certain
25 compounds of formula I will provide other compounds of formulae I. For

example, carbonyl may be reduced to hydroxy or alkylene, and hydroxy may be converted to halo.

The compounds of the invention may be isolated from their reaction mixtures
5 using conventional techniques.

It will be appreciated by those skilled in the art that, in the process described above, the functional groups of intermediate compounds may be, or may need to be, protected by protecting groups.

10

Functional groups which it is desirable to protect include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl and diarylalkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl and alkylcarbonyl groups (e.g. methyl-
15 and ethylcarbonyl groups). Suitable protecting groups for amino include benzyl, sulfonamido (e.g. benzenesulfonamido), *tert*-butyloxycarbonyl, 9-fluorenyl-methoxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for amidino and guanidino include benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C₁₋₆ alkyl or benzyl esters.

20

The protection and deprotection of functional groups may take place before or after any of the reaction steps described hereinbefore. Protecting groups may be removed in accordance with techniques which are well known to those skilled in the art and as described hereinafter.

25

The use of protecting groups is fully described in "Protective Groups in Organic Chemistry", edited by J.W.F. McOmie, Plenum Press (1973), and "Protective Groups in Organic Synthesis", 3rd edition, T.W. Greene & P.G.M. Wutz, Wiley-Interscience (1999).

Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative, and, on some occasions, more convenient, manner, the individual process steps mentioned herein may be performed in a different order, and/or the individual reactions may be performed at a different stage in the overall route (i.e. substituents may be added to and/or chemical transformations performed upon, different intermediates to those associated hereinbefore with a particular reaction). This will depend *inter alia* on factors such as the nature of other functional groups present in a particular substrate, the availability of key intermediates and the protecting group strategy (if any) to be adopted. Clearly, the type of chemistry involved will influence the choice of reagent that is used in the said synthetic steps, the need, and type, of protecting groups that are employed, and the sequence for accomplishing the synthesis.

15

It will also be appreciated by those skilled in the art that, although certain protected derivatives of compounds of formula I, which may be made prior to a final deprotection stage, may not possess pharmacological activity as such, they may be administered parenterally or orally and thereafter metabolised in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". Moreover, certain compounds of formula I may act as prodrugs of other compounds of formula I.

20

All prodrugs of compounds of formula I are included within the scope of the invention.

25

Some of the intermediates referred to hereinbefore are novel. According to a further aspect of the invention there is thus provided:

(a) a compound of formula II, as hereinbefore defined, or a protected derivative thereof, provided that

- (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- 5 (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;

(b) a compound of formula IV, as hereinbefore defined, or a protected derivative thereof, provided that

- 10 (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;

15 (c) a compound of formula VI, as hereinbefore defined, or a protected derivative thereof, provided that

- (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- 20 (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;

(d) a compound of formula IX, as hereinbefore defined, or a protected derivative thereof, provided that R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group);

25

(e) a compound of formula X, as hereinbefore defined, or a protected derivative thereof, provided that

- (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,

- (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;
- (f) a compound of formula XI, as hereinbefore defined, or a protected
 5 derivative thereof, provided that
- (i) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (ii) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;
- (g) a compound of formula XII, as hereinbefore defined, or a protected
 10 derivative thereof, provided that
- (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group),
- (ii) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$, and/or
- 15 (iii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$;
- (h) a compound of formula XVII, as hereinbefore defined, or a protected derivative thereof, provided that
- (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or
 20 $-N(R^{9b})S(O)_2R^{9d}$ group),
- (ii) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (iii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- (iv) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- 25 (v) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;
- (i) a compound of formula XXII, as hereinbefore defined, or a protected derivative thereof, provided that

- (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group), and/or
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$;
- 5 (j) a compound of formula XXIII, as hereinbefore defined, or a protected derivative thereof, provided that
 - (i) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
 - (ii) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;
- 10 (k) a compound of formula XXIV, as hereinbefore defined, or a protected derivative thereof, provided that
 - (i) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
 - (ii) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$; and
- 15 (l) a compound of formula XXVI, as hereinbefore defined, or a protected derivative thereof, provided that
 - (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
 - (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
 - (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
 - 20 (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$.

Medical and pharmaceutical use

Compounds of the invention are useful because they possess pharmacological activity. They are therefore indicated as pharmaceuticals.

5

Thus, according to a further aspect of the invention there is provided the compounds of the invention for use as pharmaceuticals.

In particular, the compounds of the invention exhibit myocardial
10 electrophysiological activity, for example as demonstrated in the test described below.

The compounds of the invention are thus expected to be useful in both the prophylaxis and the treatment of arrhythmias, and in particular atrial and
15 ventricular arrhythmias.

The compounds of the invention are thus indicated in the treatment or prophylaxis of cardiac diseases, or in indications related to cardiac diseases, in which arrhythmias are believed to play a major role, including ischaemic heart
20 disease, sudden heart attack, myocardial infarction, heart failure, cardiac surgery and thromboembolic events.

In the treatment of arrhythmias, compounds of the invention have been found to selectively delay cardiac repolarization and increase refractoriness.

25

According to a further aspect of the invention, there is provided a method of treatment of an arrhythmia which method comprises administration of a therapeutically effective amount of a compound of the invention to a person suffering from, or susceptible to, such a condition.

Pharmaceutical preparations

The compounds of the invention will normally be administered orally,
5 subcutaneously, intravenously, intraarterially, transdermally, intranasally, by
inhalation, or by any other parenteral route, in the form of pharmaceutical
preparations comprising the active ingredient either as a free base or a non-
toxic organic or inorganic acid addition salt, in a pharmaceutically acceptable
dosage form. Depending upon the disorder and patient to be treated, as well
10 as the route of administration, the compositions may be administered at
varying doses.

The compounds of the invention may also be combined with any other drugs
useful in the treatment of arrhythmias and/or other cardiovascular disorders.
15 In particular the compounds of the invention may be combined with an anti-
coagulant.

When used herein, the term "*an anticoagulant*" includes references to one a
substance selected from the group consisting of aspirin, warfarin,
20 enoxaparin, heparin, low molecular weight heparin, cilostazol, clopidogrel,
ticlopidine, tirofiban, abciximab, dipyridamole, plasma protein fraction,
human albumin, low molecular weight dextran, hetastarch, reteplase,
alteplase, streptokinase, urokinase, dalteparin, filgrastin, immunoglobulin,
ginkgolide B, hirudins, foropafant, roceparant, bivalirudin, dermatan sulfate
25 mediolanum, eptilibatide, tirofiban, thrombomodulin, abcxmab, low
molecular weight dermatan sulfate-opocrin, eptacog alfa, argatroban,
fondaparinux sodium, tifacogin, lepirudin, desirudin, OP2000, roxifiban,
parnaparin sodium, human hemoglobin (Hemosol), bovine hemoglobin
(Biopure), human hemoglobin (Northfield), antithrombin III, RSR 13,

heparin-oral (Emisphere) transgenic antithrombin III, H37695, enoxaparin sodium, mesoglycan, CTC 111, bivalirudin, and any derivatives and/or combinations thereof.

- 5 Particular anticoagulants that may be mentioned include aspirin and warfarin.

The term "*an anticoagulant*" also includes references to thrombin inhibitors.

- 10 Thrombin inhibitors that may be mentioned include low molecular weight thrombin inhibitors. The term "*low molecular weight thrombin inhibitors*" will be understood by those skilled in the art, and includes references to any composition of matter (e.g. chemical compound) that inhibits thrombin to an experimentally determinable degree (as determined by *in vivo* and/or *in vitro* tests), and which possesses a molecular weight of below about 2,000,
15 preferably below about 1,000.

Preferred low molecular weight thrombin inhibitors include low molecular weight peptide-based, amino acid-based, and/or peptide analogue-based, thrombin inhibitors, as well as derivatives thereof.

20

- The term "*low molecular weight peptide-based, amino acid-based, and/or peptide analogue-based, thrombin inhibitors*" will be well understood by one skilled in the art to include references to low molecular weight thrombin inhibitors with one to four peptide linkages, and includes those described in
25 the review paper by Claesson in *Blood Coagul. Fibrin.* 5, 411 (1994), as well as those disclosed in

US Patent No 4,346,078, International Patent Applications WO 93/11152, WO 93/18060, WO 93/05069, WO 94/20467, WO 94/29336, WO 95/35309, WO 95/23609, WO 96/03374, WO 96/06832, WO 96/06849, WO 96/25426,

WO 96/32110, WO 97/01338, WO 97/02284, WO 97/15190, WO 97/30708,
WO 97/40024, WO 97/46577, WO 98/06740, WO 97/49404, WO 97/11693,
WO 97/24135, WO 97/47299, WO 98/01422, WO 98/57932, WO 99/29664,
WO 98/06741, WO 99/37668, WO 99/37611, WO 98/37075, WO 99/00371,
5 WO 99/28297, WO 99/29670, WO 99/40072, WO 99/54313, WO 96/31504,
WO 00/01704 and WO 00/08014; and European Patent Applications 648
780, 468 231, 559 046, 641 779, 185 390, 526 877, 542 525, 195 212, 362
002, 364 344, 530 167, 293 881, 686 642, 669 317, 601 459 and 623 596,
the disclosures in all of which documents are hereby incorporated by
10 reference.

In the present application, derivatives of thrombin inhibitors include
chemical modifications, such as esters, prodrugs and metabolites, whether
active or inactive, and pharmaceutically acceptable salts and solvates, such
15 as hydrates, of any of these, and solvates of any such salt.

Preferred low molecular weight peptide-based thrombin inhibitors include
those known collectively as the "gatrans". Particular gatrans which may be
mentioned include HOOC-CH₂-(R)Cha-Pic-Nag-H (known as inogatran)
20 and HOOC-CH₂-(R)Cgl-Aze-Pab-H (known as melagatran) (see
International Patent Application WO 93/11152 and WO 94/29336,
respectively, and the lists of abbreviations contained therein).

International Patent Application WO 97/23499 discloses a number of
25 compounds which have been found to be useful as prodrugs of thrombin
inhibitors. Said prodrugs have the general formula



wherein R^a represents H, benzyl or C_{1-10} alkyl, R^b (which replaces one of the hydrogen atoms in the amidino unit of Pab-H) represents OH, $OC(O)R^c$ or $C(O)OR^d$, R^c represents C_{1-17} alkyl, phenyl or 2-naphthyl and R^d represents C_{1-12} alkyl, phenyl, C_{1-3} alkylphenyl, or 2-naphthyl. Preferred compounds
5 include $R^aOOC-CH_2-(R)Cgl-Aze-Pab-OH$, wherein R^a represents benzyl or C_{1-10} alkyl, e.g. ethyl or isopropyl, especially $EtOOC-CH_2-(R)Cgl-Aze-Pab-OH$. The active thrombin inhibitors themselves are disclosed in WO 94/29336.

10 According to a further aspect of the invention there is thus provided a pharmaceutical formulation including a compound of the invention in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

Suitable daily doses of the compounds of the invention in therapeutic
15 treatment of humans are about 0.005 to 50.0 mg/kg body weight at oral administration and about 0.005 to 15.0 mg/kg body weight at parenteral administration. Preferable ranges of daily doses of the compounds of the invention in therapeutic treatment of humans are about 0.005 to 20.0 mg/kg body weight at oral administration and about 0.005 to 10.0 mg/kg body weight
20 at parenteral administration.

The compounds of the invention have the advantage that they are effective against cardiac arrhythmias.

25 Compounds of the invention may also have the advantage that they may be more efficacious than, be less toxic than, have a broader range of activity (including exhibiting any combination of class I, class II, class III and/or class IV activity (especially class I and/or class IV activity in addition to class III activity)) than, be more potent than, be longer acting than, produce

fewer side effects (including a lower incidence of proarrhythmias such as *torsades de pointes*) than, be more easily absorbed than, or that they may have other useful pharmacological properties over, compounds known in the prior art.

5

Biological Tests

Test A

Primary Electrophysiological Effects In Anaesthetised Guinea Pigs

10 Guinea pigs weighing between 500 and 1000 g were used. The animals were housed for at least one week before the experiment and had free access to food and tap water during that period.

Anaesthesia was induced by an intraperitoneal injection of pentobarbital (50
15 to 60 mg/kg) and catheters were introduced into one carotid artery (for blood pressure recording and blood sampling) and into one jugular vein (for drug infusions). Needle electrodes were placed on the limbs for recording of ECGs (lead II). A thermistor was placed in the rectum and the animal was placed on a heating pad, set to a rectal temperature of between 37.5 and 38.5°C.

20

A tracheotomy was performed and the animal was artificially ventilated with room air by use of a small animal ventilator, set to keep blood gases within the normal range for the species. In order to reduce autonomic influences both vagi were cut in the neck, and 0.5 mg/kg of propranolol was given
25 intravenously, 15 minutes before the start of the experiment.

The left ventricular epicardium was exposed by a left-sided thoracotomy, and a custom-designed suction electrode for recording of the monophasic action potential (MAP) was applied to the left ventricular free wall. The electrode

was kept in position as long as an acceptable signal could be recorded, otherwise it was moved to a new position. A bipolar electrode for pacing was clipped to the left atrium. Pacing (1 ms duration, twice the diastolic threshold) was performed with a custom-made constant current stimulator.

5 The heart was paced at a frequency just above the spontaneous sinus rate during 30 s every fifth minute throughout the study.

The MAP signal, the blood pressure signal and the lead II ECG were collected (the sampling frequency was 1000 Hz and each sampling period 10 s) on a

10 personal computer during the last 10 s of each 30 s pacing sequence and the last 10 s of the following min of sinus rhythm. The signals were processed using a custom-designed computer program (PharmLab v 4.0).

The test procedure consisted of two basal control recordings, 3 minutes apart,

15 during both pacing and sinus rhythm. After the second control recording, the first dose of the test substance was infused in a volume of 0.2 mL/kg into the jugular vein catheter for 30 seconds. Three minutes later, pacing was started and a new recording was made. Five minutes after the previous dose, the next dose of test substance was administered. Six to ten

20 consecutive doses were given during each experiment.

Data analysis

Of the numerous variables measured in this analysis, three were selected as the most important for comparison and selection of active compounds. The

25 three variables selected were the MAP duration at 75 percent repolarization during pacing, the atrio-ventricular (AV) conduction time (defined as the interval between the atrial pace pulse and the start of the ventricular MAP) during pacing, and the heart rate (defined as the RR interval during sinus rhythm). Systolic and diastolic blood pressure were measured in order to

judge the haemodynamic status of the anaesthetised animal. Further, the ECG was checked for arrhythmias and/or morphological changes.

The mean of the two control recordings was set to zero and the effects recorded after consecutive doses of test substance were expressed as percentage changes from this value. By plotting these percentage values against the cumulative dose administered before each recording, it was possible to construct dose-response curves. In this way, each experiment generated three dose-response curves, one for MAP duration, one for AV-conduction time and one for the sinus frequency (RR interval). A mean curve of all experiments performed with a test substance was calculated, and potency values were derived from the mean curve. All dose-response curves in these experiments were constructed by linear connection of the data points obtained. The cumulative dose prolonging the MAP duration by 10% from the baseline was used as an index to assess the class III electrophysiological potency of the agent under investigation (D_{10}).

Test B

Rb⁺-efflux assay for detection of HERG channel blockers

20

The human ether-a-go-go related gene (HERG) encodes the voltage-gated K⁺ channel underlying the cardiac rapid delayed rectifier current I_{Kr} . The IC₅₀ value for HERG channel blockade was determined using a high throughput functional assay based on depolarisation-induced Rb⁺-efflux from Chinese hamster ovary cells stably expressing the HERG-channel.

25

Cells were grown in Ham F12 (Life Technologies 31765-027) supplemented with 10% FBS and 0.6 mg/mL hygromycin B and were routinely passaged twice-weekly. For experimental studies, cells were plated at a density of

15,000 cells/well in Falcon, 384-well tissue culture-treated black-walled clear-bottomed plates and were thereafter incubated overnight at 37°C in a cell culture incubator.

- 5 Following incubating overnight, cell plates were washed and a Rb⁺-Load buffer (a physiological buffer containing Rb⁺) was added. Cell plates were then incubated for 3 hours and were thereafter washed. Following this wash, the test compounds were added. The cell plates were then incubated for another 10 minutes and, following this incubation period, external K⁺
- 10 concentration was increased in order to depolarize the cells and activate HERG channels. After a ten minute exposure period to the increased K⁺ concentration, supernatants were transferred to new microplates for subsequent determination of Rb⁺ content, using Atomic Absorption Spectrometry analysis.

15

- The basal Rb⁺ efflux (content of Rb⁺ (mg/L) in supernatants of wells receiving only wash buffer) was defined as 100% inhibition and the stimulated Rb⁺ efflux (content of Rb⁺ (mg/L) in supernatants of wells exposed only to increased external potassium concentration) was defined as
- 20 0% inhibition.

Compound activity was expressed as:

$$100 \times \left[1 - \frac{A - B}{C - B} \right]$$

A: Rb⁺ content in wells receiving test compound +increased external K⁺.

25 B: Basal Rb⁺ efflux.

C: Stimulated Rb⁺ efflux.

The invention is illustrated by way of the following examples.

Examples

General Experimental Procedures

- 5 Mass spectra were recorded on one of the following instruments: MUX(8)-LCT, ZQ Massspectrometer and Quattro micro, all from Waters Micromass.

LC-MS:

- Separation was performed using Agilent 1100 Series Modules or Waters
10 1525 pump on a ACT (Advanced Chromatography Technologies) ACE C8
3 x 50 mm 3 μ m with gradient elution.

Samples were injected using Waters 2700 Sample Manager.

Mobile phases:

Generic gradients were applied from 5% to 95% acetonitrile.

- 15 Buffers containing 10 mM ammonium acetate or 5 mM ammonium formate
/ 5 mM formic acid were used.

- The mass spectra were recorded with a Waters ZQ2000 or Waters ZMD
equipped with an electrospray interface, switching positive and negative
ionization mode. UV spectra were collected by a Agilent 1100 PDA or
20 Waters 2996 DAD and the evaporative light scattering (ELS) signal by a
Sedere Sedex 55 or 75.

Data collection and evaluation were performed using the MassLynx
software.

25

^1H NMR and ^{13}C NMR measurements were performed on a BRUKER ACP
300 and Varian 300, 400, 500 and 600 Mercury, Unity plus and Inova
spectrometers, operating at ^1H frequencies of 300, 400, 500 and 600 MHz

respectively and at ^{13}C frequencies of 75.4, 100.6, 125.7 and 150.9 MHz respectively.

Rotamers may or may not be denoted in spectra depending upon ease of interpretation of spectra. Unless otherwise stated, chemical shifts are given in ppm with the solvent as internal standard.

Synthesis of Intermediates

The following intermediates were not commercially available, and were therefore prepared by the methods described below.

Preparation A

4-[(2S)-3-[7-(2-Aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy]benzonitrile

tert-Butyl 2-{7-[(2S)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethylcarbamate (2.9 g, 6.56 mmol; see WO 01/28992) was dissolved in DCM (30 mL). Trifluoroacetic acid (22.2 g, 194.7 mmol) was added and the solution was stirred for 2 h. The mixture was concentrated under reduced pressure, redissolved in toluene and acetonitrile and concentrated *in vacuo* again. The product mixture was dissolved in acetonitrile (50 mL) and solid K_2CO_3 (9.7 g, 10.2 mmol) was added. The mixture was filtered and concentrated under reduced pressure. The mixture was then redissolved in a mixture of EtOAc and a saturated solution of NaHCO_3 . The mixture was extracted twice with DCM, the combined organic extracts dried (Na_2SO_4) and evaporated to afford 1.6 g (68.7 %) of the title compound.

Preparation B

4-{3-[7-(2-Aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propoxy}-benzonitrile, hydrochloride salt

(i) (2-{7-[3-(4-Cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)carbamic acid *tert*-butyl ester

4-[3-(9-Oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propoxy]benzonitrile (6.5 g, 0.0226 mol; see WO 01/28992) and *tert*-butyl 2-bromoethylcarbamate (6 g, 0.0271 mol; see WO 01/28992) were dissolved in 80 mL of dry acetonitrile. Anhydrous potassium carbonate (4.68 g, 0.034 mol) was added and the mixture was stirred at 60°C overnight. A further portion of *tert*-butyl 2-bromoethylcarbamate (2.53 g, 0.0113 mol) was added and the reaction heated at 65°C overnight. The reaction mixture was then filtered and the filtrate concentrated under reduced pressure. The product was purified by column chromatography using methanol-chloroform as eluent to give the sub-title compound (4.5 g) as a yellow solid.

(ii) 4-{3-[7-(2-Aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-propoxy}benzonitrile, hydrochloride salt
(2-{7-[3-(4-Cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-ethyl)carbamic acid *tert*-butyl ester (4.5 g, 0.0104 mol; see step (i) above) was added to a solution of HCl in dioxane. The reaction was stirred at room temperature for 2 h, after which the reaction mixture was filtered and the resultant solid washed with dioxane, then diethyl ether and dried to give the title compound (3.89 g) as a white solid.

25

Preparation C

{2-[9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl} *tert*-butylcarbamate, hydrochloride salt

(i) 2-Bromoethyl *tert*-butylcarbamate

To a cooled solution of triphosgene (76 g, 0.256 mol, 0.75 eq.) in dry DCM (150 mL) was added, dropwise at 0°C over a period of 3 hrs under a N₂ atmosphere, a mixture of *tert*-butylamine (25 g, 0.3418 mol, 1.0 eq.), DMAP (2.08g, 0.017 mol, 0.05 eq.) and triethylamine (67.1 g, 0.6836 mol, 2.0 eq.) in dry DCM (100 mL). After addition of the mixture was complete, the reaction mixture was stirred at RT for 12 hrs. The reaction mixture was cooled to 0°C and 2-bromoethanol (106.8 g, 0.854 mol, 2.5 eq.) was added. The reaction mixture was then stirred at RT for 6 hrs, before being diluted with water and extracted with DCM. The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, and then concentrated under reduced pressure. The resulting residue was distilled under reduced pressure to yield carbamate the sub-title compound as a colourless liquid. Yield: 28 g.

(ii) {2-[7-Benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl} *tert*-butylcarbamate

A mixture of 3-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]nonane dihydrochloride (8 g, 0.0275 mol, 1.0 eq.; see WO 01/28992), 2-bromoethyl *tert*-butylcarbamate (9.23 g, 0.0412 g, 1.5 eq.) and fused K₂CO₃ (18.96 g, 0.1375 mol, 5.0 eq.) in dry acetonitrile (120 mL) was stirred at 70°C overnight under a N₂ atmosphere. The reaction mixture was cooled to RT, filtered and concentrated under reduced pressure. The resulting residue was purified by column chromatography over silica gel (using 92:8 DCM : methanol as final eluent) to yield the sub-title compound (5 g, 43%).

(iii) {2-[9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl} *tert*-butylcarbamate, hydrochloride salt

{2-[7-Benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl} *tert*-butyl-carbamate (5 g; see step (ii) above) was dissolved in dry dioxane (50 mL) and the resulting solution cooled to 0°C. To this solution was added, dropwise with continuous stirring, 100 mL of dry dioxane saturated with HCl. The reaction mixture was stirred at RT for 1 hr. Dioxane was decanted carefully and the hygroscopic solid precipitate was washed with dry ether twice and then dried under vacuum to yield the dihydrochloride salt as a white solid. This salt was dissolved in dry methanol (100 mL), to which was then added 10% palladium on carbon. The resulting mixture was stirred under a H₂ atmosphere for 12 hrs. The reaction mixture was filtered through Celite®, the filtrate concentrated under reduced pressure and the resulting solid dried under high vacuum to provide the title compound as a pale yellow solid. Yield: 3.8 g.

Preparation D

1-[3-(9-Oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propanoyl]indoline-5-carbonitrile, hydrochloride salt

(i) 5-Bromo-2,3-dihydro-1H-indole

NaBH₃CN (19.3 g, 0.3061 mol) was added in 3 portions to a cooled (15°C) solution of 5-bromoindole (20 g, 0.10 mol) in glacial acetic acid (500 mL) and stirred for 2 h at this same temperature. The reaction mixture was quenched with water, cooled and basified with a sodium hydroxide pellet. It was then extracted with diethyl ether. The organic layer was washed with water and brine, dried over sodium sulfate, concentrated and purified by column chromatography over silica gel to give the sub-title compound as a pale yellow solid. Yield: 20 g.

(ii) 5-Bromo-1-(trifluoroacetyl)indoline

5-Bromo-2,3-dihydro-1*H*-indole (20 g, 0.1010 mol; see step (i) above) was taken in (100 mL) of DCM and cooled to 0°C. Trifluoroacetic anhydride (42.42 g, 0.2020 mol) was added dropwise to the reaction mixture, which was then stirred for 30 min. The reaction mixture was quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over sodium sulfate, concentrated and purified by column chromatography over silica gel to give the sub-title compound as a white solid. Yield: 28 g.

10 (iii) 1-(2,2,2-Trifluoroacetyl)-2,3-dihydro-1*H*-indole-5-carbonitrile
1-(5-Bromo-2,3-dihydro-1*H*-indol-1-yl)-2,2,2-trifluoroethanone (19 g, 0.0646 mol; see step (ii) above) was dissolved in DMF (50 mL). CuCN (8.68 g, 0.0969 mol) was then added and the reaction mixture was stirred at 160°C for 4 days, before being cooled to RT, diluted with water and
15 extracted with ethyl acetate. The organic layer was washed with water and brine, dried over sodium sulfate then concentrated under reduced pressure. The resulting residue was purified by column chromatography to give the sub-title compound as a white solid. Yield: 13 g.

20 (iv) 2,3-Dihydro-1*H*-indole-5-carbonitrile
1-(2,2,2-Trifluoroacetyl)-2,3-dihydro-1*H*-indole-5-carbonitrile (13 g, 0.0542 mol; see step (iii) above) was dissolved in THF (100 mL), to which 1% NaOH was then added. After stirring for 5 h at RT, the reaction mixture was extracted with ethyl acetate. The organic layer was washed with water
25 and brine, dried over sodium sulfate and solvent was evaporated to provide the sub-title compound as a pale yellow solid. Yield: 8 g.

(v) 1-(3-Chloropropionyl)-2,3-dihydro-1*H*-indole-5-carbonitrile

To a solution of 2,3-dihydro-1*H*-indole-5-carbonitrile (8 g, 0.0556 mol; see step (iv) above) in DCM (50 mL) at 0°C was added *N,N*-dimethylaniline (13.5 g, 0.1112 mol) followed by 2-chloropropionyl chloride (8.46 g, 0.0667 mol). The reaction mixture was stirred for 3 h before being diluted
5 with water and extracted with DCM. The organic layer was washed with water and brine, dried over sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography over silica gel to yield the sub-title compound as a pale yellow solid. Yield: 10.5 g.

10

(vi) *tert*-Butyl 7-[3-oxo-3-(5-cyano-2,3-dihydro-1*H*-indol-1-yl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate
1-(3-Chloropropionyl)-2,3-dihydro-1*H*-indole-5-carbonitrile (3.07 g, 0.031 mol; see step (v) above), 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-
15 carboxylic acid *tert*-butyl ester (3 g, 0.031 mol; see WO 01/28992) and potassium carbonate (6.43 g, 0.0465 mol) were taken in dry acetonitrile (30 mL) and stirred at 60°C overnight under a nitrogen atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to provide a crude product. This product was then purified by column
20 chromatography using methanol in DCM as eluent to yield the sub-title compound as pale yellow solid. Yield: 5 g.

(vii) 1-[3-(9-Oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propanoyl]indoline-5-carbonitrile, hydrochloride salt
25 *tert*-Butyl 7-[3-oxo-3-(5-cyano-2,3-dihydro-1*H*-indol-1-yl)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate (5 g, 0.017 mol; see step (vi) above) was taken up in 50 mL of dioxane that was saturated with HCl gas. The mixture was then stirred for 1 h under nitrogen atmosphere at RT. The

resulting solid was filtered and washed with ether and then dried under vacuum to yield 4.42 g of the title compound as an off white solid.

Preparation E

5 4-Cyano-N-[2-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzene-sulfonamide

(i) N-(2-Bromoethyl)-4-cyanobenzenesulfonamide

To an ice cooled solution of 2-bromoethylamine hydrogenbromide (12.2 g,
10 0.0595 mol) in dry dichloromethane (100 mL) was added triethylamine and the mixture stirred for 15 min. 4-Cyanobenzenesulfonyl chloride (10 g, 0.0486 mol) in dichloromethane was added, dropwise. Stirring was continued at RT for 1 h. The reaction mixture was diluted with dichloromethane, washed with water, followed by brine, then dried over
15 sodium sulfate. Solvents were evaporated and the residue crystallized from petroleum ether to give 12.5 g of the sub-title compound as a pale yellow solid.

(ii) 7-[2-(4-Cyanobenzenesulfonylamino)ethyl]-9-oxa-3,7-diazabicyclo-
20 [3.3.1]nonane-3-carboxylic acid *tert*-butyl ester

A suspension of N-(2-bromoethyl)-4-cyanobenzenesulfonamide (5 g, 0.017 mol; see step (i) above), 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (3.96 g, 0.0173 mol; see WO 01/28992) and potassium carbonate (3.6 g, 0.025 mol) in acetonitrile (50 mL) was stirred at
25 60°C overnight. The reaction mixture was filtered and the solvent concentrated under reduced pressure to yield the crude product, which was purified by column chromatography (using methanol in chloroform as the eluent) to yield (5 g) of the sub-title compound as pale yellow solid.

(iii) 4-Cyano-N-[2-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzene-sulfonamide

7-[2-(4-Cyanobenzenesulfonylamino)ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]-nonane-3-carboxylic acid *tert*-butyl ester (5 g; see step (ii) above) was taken
5 up in 30 mL of dioxane that was saturated with HCl gas. The mixture was stirred for 1 h before the dioxane was decanted off and the remaining solid was washed with diethyl ether, yielding the HCl salt of the title compound. This salt was then taken up in DCM-NaHCO₃ (200 mL, 1:1) and the resulting mixture stirred for 1 h at RT. The compound was extracted with
10 dichloromethane, washed with water, brine and dried over sodium sulfate. Solvent evaporation under reduced pressure yielded 3.4 g of the title compound as white solid.

Preparation F

4-{2-[7-(2-Aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}-benzonitrile, trihydrochloride salt

(i) (2-{7-[2-(4-Cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)carbamic acid *tert*-butyl ester, diacetate salt
[2-(9-Oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]carbamic acid *tert*-butyl
20 ester, 2,4,6-trimethylbenzenesulfonic acid salt (10.1 g, 21.3 mmol; see WO 02/83690) was mixed with 4-(2-bromoethoxy)benzonitrile (5.37 g, 23.7 mmol) and K₂CO₃ (7.44 g, 74.6 mmol) in acetonitrile (200 mL). The mixture was refluxed overnight before further K₂CO₃ (2.88 g, 21 mmol) was added and the mixture refluxed for an additional 8 h. The reaction mixture
25 was filtered and evaporated. The product obtained thereby was purified by chromatography (kromasil C8, 100X500 mm, CH₃CN/NH₄OAc-buffer with 0.2% formic acid; Gradient 40-80% acetonitrile). The product was freeze-dried and the diacetate was isolated. (10.8 g, 94.5% yield.)

(ii) 4-{2-[7-(2-Aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethoxy}-benzonitrile, trihydrochloride salt

- Hydrochloric acid (140 mmol, 35 mL of a 4 M solution in dioxane) was added to an ice-cooled slurry of the diacetate salt of (2-{7-[2-(4-
- 5 cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)carbamic acid *tert*-butyl ester (10.75 g, 20 mmol; see step (i) above) in dioxane (50 mL). The reaction mixture was stirred at RT overnight. Another 65 mL (260 mmol) of 4 M HCl solution was added, after which the reaction mixture was stirred for three hours and then evaporated. The residue was
- 10 dissolved in water and washed with diethyl ether. Freeze-drying of the aqueous phase gave 8.3 g (97%) the title compound as a white solid.
- ¹H NMR (D₂O): δ 2.78 (m, 4 H), 3.22 (m, 4 H), 3.65 (m, 2 H), 3.71 (t, *J* = 4.6 Hz, 2 H), 3.91 (m, 2 H), 4.36 (bs, 2 H), 4.57 (t, *J* = 4.6 Hz, 2 H), 7.16 (d, *J* = 9 Hz), 7.79 (d, *J* = 9 Hz).
- 15 ¹³C NMR (D₂O): δ 33.2, 52.4, 53.0, 53.4, 53.9, 59.2, 63.7, 101.4, 113.1, 117.6, 132.3, 158.5.

Preparation G

4-(2-Bromoethyl)-1,2-bis(difluoromethoxy)benzene

- 20 (i) 3,4-Bis(difluoromethoxy)benzaldehyde
- 3,4-Dihydroxybenzaldehyde (20 g, 0.144 mol) and potassium carbonate (40 g, 0.2898 mol) were taken up in dry DMF (1 L) and cooled to -78°C. A Dewar condenser was fitted and freon gas (CHClF₂) was passed through the reaction mixture for 1 h. The reaction mixture was stirred at 85°C
- 25 overnight. Solvent was evaporated under reduced pressure and the residue purified by column chromatography over silica gel (using petroleum ether in ethyl acetate as eluent) to yield the sub-title compound as a pale yellow liquid. Yield: 11 g.

(ii) Methyl triphenylphosphonium iodide

To a solution of triphenylphosphine (30 g, 0.114 mol) in benzene (200 mL) was added methyl iodide (11 mL, 0.1756 mol) dropwise, and the solution stirred at room temperature for 15 min under a nitrogen atmosphere. The
5 sub-title compound formed as a colourless solid, which was filtered and then dried. Yield: 30 g.

(iii) 1,2-Bis(difluoromethoxy)-4-vinylbenzene

Potassium *tert*-butoxide (6.7 g, 0.0683 mol) was taken in dry THF (25 mL)
10 and stirred at 0°C under a nitrogen atmosphere for 15 min. Methyl triphenylphosphonium iodide (27.6 g, 0.0683 mol; see step (ii) above) was then added, followed by 3,4-bis(difluoromethoxy)benzaldehyde (8 g, 0.034 mol; see step (i) above). The reaction mixture was then stirred for 1 h at RT, before being quenched with water. The organic solvents were
15 evaporated under reduced pressure and the aqueous layer extracted with dichloromethane. Evaporation of the solvents followed, by chromatography of the residue over silica gel (using petroleum ether in ethyl acetate as eluent) afforded the sub-title compound as a pale yellow liquid. Yield: 6.5 g

20 (iv) 2-[3,4-Bis(difluoromethoxy)phenyl]ethanol

To a solution of 1,2-bis(difluoromethoxy)-4-vinylbenzene (10 g, 0.0431 mol; see step (iii) above) in 25 mL of THF was added, dropwise, borane-methyl sulfide complex (10 mmol; 10 mL of a 1 M solution in THF). The resulting mixture was stirred at 60°C for 1 h, before being cooled to RT.
25 Excess borane was decomposed by adding few drops of water and the resultant mixture refluxed for 1 h with 2 M NaOH (100 mL) 50% H₂O₂ (87 mL). The mixture was then extracted with dichloromethane, washed with water and then brine, and dried over sodium sulfate. Solvent

evaporation under reduced pressure gave the sub-title compound as a pale yellow liquid. Yield: 6 g.

(v) 4-(2-Bromoethyl)-1,2-bis(difluoromethoxy)benzene

- 5 2-[3,4-Bis(difluoromethoxy)phenyl]ethanol (6 g, 0.024 mol; see step (iv) above) and triphenylphosphine (12.6 g, 0.048 mol) were taken up in dichloromethane (100 mL) and cooled to 0°C. CBr₄ (15.9 g, 0.048 mol) in dichloromethane was added dropwise, and the mixture stirred at RT overnight under a nitrogen atmosphere. The reaction mixture was filtered,
10 the filtrate concentrated under reduced pressure and the residue purified by column chromatography over silica gel (using 20% ethyl acetate in petroleum ether as eluent) to give the title compound as a yellow liquid. Yield 5.2 g.

15 Preparation H

4-Fluoro-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide

(i) N-(2-Bromo-ethyl)-4-fluoro-benzenesulfonamide

- 4-Fluoro benzene sulfonyl chloride (8 g, 0.0411 mol) was added dropwise at
20 0 °C to a solution of 2-bromoethylamine hydrogen bromide (10.27 g, 0.0501 mol) and triethyl amine (14.3 ml, 0.1029 mol) in dry dichloromethane (100 ml) and stirred at room temperature for 1 h under nitrogen atmosphere. The reaction mixture was diluted with water and extracted with ethyl acetate. Organic layer was washed with water and brine and dried over sodium
25 sulfate. Solvent evaporation under reduced pressure afforded the sub-title product as a liquid. Yield: 11g

(ii) 7-[2-(4-Fluoro-benzenesulfonylamino)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

A suspension of N-(2-Bromo-ethyl)-4-fluoro-benzenesulfonamide (7 g, 0.0179 mol; see step (i) above) and 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (11.2 g, 0.159 mol; see WO 01/28992) and K₂CO₃ (9.7 g, 0.0798 mol) in dry acetonitrile (70 ml) was stirred at 60 °C overnight under nitrogen atmosphere. The reaction mixture was partitioned between brine and ethyl acetate. Organic layer was washed with brine and dried over sodium sulfate. Solvent evaporation under reduced pressure followed by column chromatography over silica gel using 7 % ethyl acetate in petroleum ether as eluent afforded the sub-title compound as a solid.

Yield: 4g

(iii) 4-Fluoro-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide, hydrochloride salt

HCl in dioxane (50 ml) was added to a solution of 7-[2-(4-Fluoro-benzenesulfonylamino)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (4 g; see step (ii) above) in dry dioxane (50 ml) at RT and the reaction mixture was stirred at RT for 1 h. Dioxane was decanted, precipitated solid was washed with dry diethyl ether (three times) and dried under vacuum to give the title compound as pale yellow solid. Yield :3.55g

Preparation I

4-Cyano-N-methyl-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide, hydrochloride salt

(i) 4-Cyano-N-methyl-N-(2-oxo-ethyl)-benzenesulfonamide

4-Cyanobenzenesulfonyl chloride (6 g, 0.0297 mol) in 50 ml of dry DCM was added at 0 °C to a solution of N-methylethanolamine (2.67 g, 0.0356 mol) and triethylamine (7.49 g, 0.0742 mol) in dry dichloromethane (75 ml) under nitrogen atmosphere and stirred at RT for 2 h. The reaction mixture

was quenched with water and extracted with dichloromethane. Organic layer was washed with water and brine and dried over sodium sulfate. Solvent evaporation under reduced pressure afforded crude product as solid. This solid was then washed with petroleum ether (three times) and dried under vacuum to give the sub-title compound (5.18 g) as white solid.

(ii) 4-Cyano-N-methyl-N-(2-oxo-ethyl)-benzenesulfonamide

Dry DMSO (5.03 g, 0.0465 mol) was added to a solution of oxalyl chloride (4.1 g, 0.0323 mol) in dry DCM (100 ml) and stirred at RT for 30 min under nitrogen atmosphere.

4-Cyano-N-methyl-N-(2-oxo-ethyl)-benzenesulfonamide (5.17 g, 0.0215 mol; see step (i) above) in dry DCM (30 ml) was added at RT and stirring continued at RT for 2 h. The reaction mixture was quenched with triethylamine (10.85 g) at -78 °C and slowly warmed to RT. 10 % Citric acid was added and the compound was extracted with dichloro methane. The combined organic layer was washed with water and brine and dried over sodium sulfate. Solvent evaporated under reduced pressure to give the sub-title compound (4.97 g) as yellow oil. This was taken for next step without further purification.

(iii) 7-{2-[(4-Cyano-benzenesulfonyl)-methyl-amino]-ethyl}-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

A mixture of step (ii) product 4-Cyano-N-methyl-N-(2-oxo-ethyl)-benzenesulfonamide (4.97 g), and 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (4.95 g; see WO 01/28992) and acetic acid (1.7 ml) in dry dichloromethane (50 ml) was stirred at RT for 3 h and then cooled to 0 °C. NaBH₃CN (1.96 g, 0.0312 mol) was added and stirring continued at RT overnight under nitrogen atmosphere. The reaction mixture was quenched with water, extracted with dichloromethane, washed with

water and brine and dried over sodium sulfate. Solvent evaporated under reduced pressure and the residue was purified by column chromatography over silica gel using 28 % ethyl acetate in petroleum ether eluent to give the sub-title compound (2.2 g) as a liquid.

5

(iv): 4-Cyano-N-methyl-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide, hydrochloride salt

Step (iii) product 7-{2-[(4-Cyano-benzenesulfonyl)-methyl-amino]-ethyl}-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester (2.2 g) was dissolved in dioxane (10 ml) and to which HCl in dioxane (saturated, 20 ml) was added drop by drop at RT under nitrogen atmosphere. Solvent was decanted, the precipitated solid was washed with dry diethyl ether (three times) and dried under vacuum to yield the title compound (2 g) as white powder.

15

Preparation J

4-{2-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-benzonitrile

(i) (3-{7-[2-(4-Cyano-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-propyl)-carbamic acid tert-butyl ester

A suspension of 4-[2-(9-Oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethoxy]-benzonitrile (6.5 g, 0.02 mol, see WO 01/28992), (3-Bromo-propyl)-carbamic acid tert-butyl ester (6.0 g, 0.025 mol) and dry K₂CO₃ (13.08 g, 0.0946 mol) in 40 ml of dry acetonitrile was stirred at 60 °C overnight under N₂ atmosphere. The reaction mixture was filtered through celite and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 20 % ethyl acetate in petroleum ether as eluent to yield sub-title compound (3.5 g) as a liquid.

25

(ii) 4-{2-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-benzonitrile

Dioxane (15 ml, saturated with HCl gas) was added to a solution of (3-{7-[2-(4-Cyano-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-propyl)-carbamic acid tert-butyl ester (3.5 g, from step (i) above) in 5 ml of dry dioxane and stirred for 1 h at RT under nitrogen atmosphere. Dioxane was decanted, precipitated solid was filtered and washed with dry diethyl ether (3 times) and dried under vacuum to give HCl salt of the title compound (1.7 g) as white powder.

10

Preparation K

4-{3-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-propoxy}-benzonitrile hydrochloric acid salt

(i) (3-{7-[3-(4-Cyano-phenoxy)-propyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-propyl)-carbamic acid tert-butyl ester

15

The sub-title compound was made in analogy with preparation J (i) above using 4-[3-(9-Oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-propoxy]-benzonitrile in place of 4-[2-(9-Oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethoxy]-benzonitrile (see WO 01/28992).

20

(ii) 4-{3-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-propoxy}-benzonitrile hydrochloric acid salt

(3-{7-[3-(4-Cyano-phenoxy)-propyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-propyl)-carbamic acid tert-butyl ester (3.5 g) from step (i) above was taken in 25 ml of dioxane (saturated with HCl gas) and stirred at room temperature for 30 min under nitrogen atmosphere. Dioxane was decanted, the solid was washed with diethyl ether (3 times) and dried under vacuum to yield (2.8 g) of the title compound as white powder.

25

Preparation L

4-Cyano-N-[3-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-propyl]-benzenesulfonamide hydrochloric salt

(i) N-(3-Bromo-propyl)-4-cyano-benzenesulfonamide:

- 5 4-Cyanobenzenesulfonyl chloride (5 g, 0.0248 mol) was added dropwise to a solution of 3-bromopropylamine hydrobromide (6.61 g, 0.0302 mol) and triethylamine (6.26 g, 0.062 mol) in dry dichloromethane (50 ml) at 0 °C and stirred at room temperature overnight under nitrogen atmosphere. The reaction was quenched with water and extracted with dichloromethane. The
- 10 organic layer was washed with water, brine and dried over sodium sulfate. Solvent evaporation under reduced pressure afforded the sub-title compound (6.39 g) as a solid. This was used in the next step without further purification

- 15 (ii) 7-[3-(4-Cyano-benzenesulfonylamino)-propyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

- A suspension of N-(3-Bromo-propyl)-4-cyano-benzenesulfonamide (6.39g, 0.021 mol; see step (i) above), 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (4.32 g, 0.0189 mol see WO 01/28992) and
- 20 dry K₂CO₃ (10.16 g, 0.0735 mol) in dry acetonitrile (70 ml) was stirred at 60 °C overnight under N₂ atmosphere. The reaction mixture was filtered through celite and filtrate was concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 50 % ethyl acetate in petroleum ether, as eluent to give the sub-title compound
- 25 (4.29 g) as a solid.

(iii) 4-Cyano-N-[3-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-propyl]-benzenesulfonamide hydrochloric acid salt

7-[3-(4-Cyano-benzenesulfonylamino)-propyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (4.29 g, see step (ii) above) was taken in 20 ml of dioxane, 40 ml of diethylether (saturated with HCl gas) was added and stirred for 1 h at RT under nitrogen atmosphere. Solvent
5 was decanted. The solid was washed with dry diethylether (4 times) and dried under vacuum to give the title compound (5.3 g) as a white solid.

Preparation M

4-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-ylmethyl]-
10 benzonitrile hydrochloric acid salt

(i) 7-(4-Cyano-benzyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester

A suspension of 4-bromomethylbenzonitrile (10 g, 0.054 mol), 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (11.5 g, 0.0439
15 mol; see WO 01/28992) and dry K₂CO₃ (21.2 g, 0.153 mol) in 200 ml of dry acetonitrile was stirred at 60 °C overnight under N₂ atmosphere. The reaction mixture was filtered through celite and filtrate was concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 10 % ethyl acetate in petroleum ether
20 as eluent to yield (10 g) of the sub-title compound as colorless liquid.

(ii) 4-(9-Oxa-3,7-diaza-bicyclo[3.3.1]non-3-ylmethyl)-benzonitrile, hydrochloride

7-(4-Cyanobenzyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid
25 *tert*-butyl ester from step (i) above (10 g) was taken in 15 ml of dioxane (saturated with HCl gas) and stirred for 1 h at RT under nitrogen atmosphere. Dioxane was decanted, the precipitated solid was filtered, washed with dry diethyl ether (4 times) and dried under vacuum to give HCl

salt of the sub-title compound (7.5 g) as a powder. This was directly taken for next step without further purification.

(iii) {3-[7-(4-Cyano-benzyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-

5 propyl}-carbamic acid tert-butyl ester

A suspension of 4-(9-Oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl-methyl)-benzonitrile hydrochloride (6.8 g, 0.021 mol; from step (ii) above), (3-Bromopropyl)-carbamic acid tert-butyl ester (3.93 g, 0.017 mol) and dry K_2CO_3 (8.24 g, 0.059 mol) in 27 ml of dry acetonitrile was stirred at 60 °C
10 overnight under N_2 atmosphere. The reaction mixture was filtered through celite and filtrate was concentrated under reduced pressure to give (7.5 g) of the sub-title compound as pale a liquid.

(iv) 4-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-ylmethyl]-

15 benzonitrile hydrochloride

{3-[7-(4-Cyano-benzyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-propyl}-carbamic acid tert-butyl ester (7 g from step (iii) above) was taken in 50 ml of dioxane (saturated with HCl gas) and stirred for 1 h at RT under nitrogen atmosphere. Dioxane was decanted, the precipitated solid was filtered and
20 washed with dry diethylether (4 times) and dried under vacuum to give the title compound (6.5 g) as a powder.

Preparation N

4-{2-[7-(2-Amino-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-

25 3-fluoro-benzonitrile hydrochloric acid salt

(i) 4-Bromo-2-fluorophenol

Bromine (68.7 ml, 1.339 mol) dissolved in acetic acid (300 ml) was added drop by drop to a cooled solution of 2-fluorophenol (150 g, 1.339 mol) in acetic acid (1300 ml) and the reaction mixture was stirred at room

temperature overnight. The reaction mixture was quenched with aq. sodium bisulfite solution and extracted with dichloromethane. The organic layer was washed with water and brine and dried over sodium sulfate. Solvent evaporation under reduced pressure afforded 4-bromo-2-fluorophenol (210 g) as a liquid. This was directly taken for next step without further purification.

(ii) 4-Bromo-2-fluoro-1-methoxybenzene

Methyl iodide (182.1 ml, 1.319 mol) was added at 0 °C to a well stirred suspension of 4-bromo-2-fluorophenol (210 g, 1.099 mol, from step (i) above) and K₂CO₃ (303.92 g, 2.19 mol) in dry acetone (1.7 L) and stirring continued at 60 °C for two days under nitrogen atmosphere. The reaction mixture was filtered and the solvent was concentrated under reduced pressure to yield 4-bromo-2-fluoro-anisole (225 g) as a liquid. This was directly taken for next step without further purification.

(iii) 3-Fluoro-4-methoxybenzonitrile

A mixture of 4-Bromo-2-fluoro-1-methoxybenzene (107 g, 0.5244 mol, from step (ii) above), CuCN (70.4 g, 0.7866 mol) in dry DMF (150 ml) was stirred at 120 °C overnight. The reaction mixture was cooled to room temperature, diluted with water and extracted with ethyl acetate. Organic layer was washed with water and brine and dried over sodium sulfate. Solvent evaporation under reduced pressure followed by column chromatography over silica gel using 3% ethyl acetate in petroleum ether gave the sub-title compound (24.4 g) as a solid.

(iv) 3-Fluoro-4-hydroxy-benzonitrile

BBr₃ (23 ml, 0.242 mol) was added to 3-Fluoro-4-methoxy-benzonitrile (24.4 g, 0.16 mol) in dichloromethane (200 ml) at -78 °C and stirring

continued overnight at room temperature. Another portion of BBr_3 (23 ml, 0.242 mol) was added at -78°C and stirring continued at RT for 2 days under nitrogen atmosphere. The reaction mixture was quenched with ice water and extracted with dichloromethane. Organic layer was washed with
5 water and brine, and dried over sodium sulfate. Solvent evaporation under reduced pressure gave 20 g of the sub-title compound as a solid. This was taken for next step without further purification.

(v) 4-(2-Bromo-ethoxy)-3-fluorobenzonitrile

10 A suspension of 3-fluoro-4-hydroxybenzonitrile (20 g, 0.1459 mol, from step (iv) above), anhydrous. K_2CO_3 (40.33 g, 0.2918 mol) and 1,2-dibromoethane (76.8 ml, 0.8754 mol) in dry DMF (150 ml) was stirred at 60°C for 5 days under nitrogen atmosphere. The reaction mixture was filtered through celite and solvent evaporated under reduced pressure. The residue was
15 purified by column chromatography over silica gel using 2 % ethyl acetate in petroleum ether, as eluent to yield the sub-title compound (21.6 g) as a solid.

(vi) 7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-

20 bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

A suspension of 4-(2-Bromo-ethoxy)-3-fluoro-benzonitrile (21.6 g, 0.0885 mol, from step (v) above), 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester (21.1 g, 0.07965 mol; see WO 01/28992) and dry K_2CO_3 (48.9 g, 0.354 mol) in 200 ml of dry acetonitrile was stirred at
25 60°C for five days under N_2 atmosphere. The reaction mixture was filtered through celite and filtrate was concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 27 % ethyl acetate in petroleum ether, as eluent to yield the sub-title compound (20.5 g) as a solid.

(vii) 3-Fluoro-4-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethoxy]-benzonitrile hydrochloride hydrochloric acid salt

7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-

- 5 bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester (20.5 g, from step (vi) above) was taken in 20 ml of dioxane and added 100 ml of dioxane (saturated with HCl gas) and stirred for 1 h at RT under nitrogen atmosphere. Solvent was decanted and the precipitated solid was washed with dry diethylether (4 times) and dried under vacuum to give HCl salt of
10 the sub-title compound (21 g) as a solid.

(viii) (2-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-carbamic acid tert-butyl ester

- A suspension of intermediate 3-fluoro-4-[2-(9-oxa-3,7-diaza-
15 bicyclo[3.3.1]non-3-yl)-ethoxy]-benzonitrile hydrochloride (5 g, 0.0137 mol, from step (vii) above), and *tert*-butyl 2-bromoethylcarbamate (3.98 g, 0.0178 mol, see WO 01/28992) and potassium carbonate (7.57 g, 0.0548 mol) in dry acetonitrile (100 ml) was stirred at 60 °C overnight under nitrogen atmosphere. The reaction mixture was filtered through celite and
20 filtrate was concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 5 % methanol in dichloromethane as eluent to give the sub-title compound (4.6 g) as an oil.

(viv) 4-{2-[7-(2-Amino-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile trifluoroacetic acid salt

- 25 (2-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-carbamic acid tert-butyl ester (4.6 g from step (viii) above) was taken in dry dichloromethane (20 ml) and cooled to 0 °C. Trifluoro acetic acid (75 ml) was added drop by drop and the reaction

mixture was stirred at RT for 2 h under nitrogen atmosphere. Solvent and trifluoroacetic acid were evaporated under reduced pressure to give the TFA salt of the sub-title compound (7.6 g) as an oil.

- 5 (x) 4-{2-[7-(2-Amino-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile hydrochloric acid salt
4-{2-[7-(2-Amino-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-
3-fluoro-benzonitrile trifluoroacetic acid salt (8.3g,12.3mmol, from step
(viv) above) was dissolved in dioxane (15 ml). HCl (15 ml of a 4M solution
10 in dioxane) was added whilst stirring resulting in the precipitation of the
HCl-salt. The salt was filtered and dried, dissolved in water and freeze
dried. Yield: 3.6 g of the title compound as a solid.

Preparation O

- 15 4-{2-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile hydrochloric acid salt

(i) (3-Bromo-propyl)-carbamic acid tert-butyl ester

- (Boc)₂O (13.5 g, 0.062 mol) was added drop by drop at 0 °C to a solution of
3-bromo propyl amine hydrobromide (15 g, 0.0688 mol) and triethyl amine
20 (13.89 g, 0.1376 mol) in dry dichloromethane (125 ml) under nitrogen
atmosphere. The reaction mixture was stirred at RT for 1 h and quenched
with water. Organic layer was washed with water and brine and dried over
sodium sulfate. Solvent evaporation under reduced pressure followed by
purification over silica gel using 5 % ethyl acetate in pet ether as eluent
25 afforded the sub-title compound (11.6 g) as yellow oil.

(ii) (3-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-propyl)-carbamic acid ter-butyl ester

A suspension of 3-Fluoro-4-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethoxy]-benzonitrile hydrochloride hydrochloric acid salt (5 g, 0.0137 mol, from step N (vii) above), (3-Bromo-propyl)-carbamic acid tert-butyl ester (4.24 g, 0.0178 mol, from step (i) above) and potassium carbonate (7.57 g, 0.0548 mol) in dry acetonitrile (100 ml) was stirred at 60 °C overnight under nitrogen atmosphere. The reaction mixture was filtered through celite and solvent concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 5 % methanol in dichloromethane as eluent to give the sub-title compound (5.8 g) as yellow oil.

(iii) 4-{2-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile TFA-salt
(3-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-propyl)-carbamic acid isopropyl ester (5.8 g, from step (ii) above) was taken in dry dichloromethane (25 ml) and cooled to 0 °C. Trifluoroacetic acid (75 ml) was added drop by drop and the reaction mixture was stirred at RT for 2 h under nitrogen atmosphere. Solvent and trifluoroacetic acid were evaporated under reduced pressure to give the TFA salt of the sub-title compound (9 g) as an oil.

(iv) 4-{2-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile hydrochloric acid salt
4-{2-[7-(3-Amino-propyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile TFA-salt (7.79g, 11.3 mmol, from step (iii) above) was dissolved in dioxane (15 ml). HCl (14 ml of a 4M solution in dioxane) was added. The HCl-salt precipitated, the dioxane was decanted off and to the solid was added dioxane (10 ml) followed by HCl (4 ml in dioxane). The solvent was decanted off and the solid was dried under vacuum. The

solid was dissolved in water (5ml) and freeze dried giving 2.98 g of the title compound

Preparation P

5 2-(9-Oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethanesulfonic acid 4-cyano-benzylamide

(i) 2-Chloro-ethanesulfonic acid 4-cyano-benzylamide

Chloroethane sulfonyl chloride (4.8 g, 0.029 mol) was added drop by drop to solution of 4-aminomethyl benzonitrile methan sulphonic acid salt acid (7 g, 0.029 mol) and pyridine (24 ml) in dry dichloromethane (25 ml) at -5 °C under nitrogen atmosphere. The reaction mixture was stirred at RT for 2 days and then partitioned between water and dichloromethane. The organic layer was washed with water and brine and dried over sodium sulfate. Solvent was evaporated under reduced pressure and the residue was purified by column chromatography over silica gel using 1 % methanol in chloroform as eluent to give mixture of the sub-title compound and ethenesulfonic acid 4-cyano-benzylamide (3.4 g) as a solid. This was directly taken for next step without further purification.

20 (ii) 7-[2-(4-Cyano-benzylsulfamoyl)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

A suspension of step (i) intermediate (3.4 g, 0.013 mol), 9-oxa-3,7-diaza-bicyclo[3.3.1]nonane-3-carboxylic acid *tert*-butyl ester; (3.49 g, 0.013 mol see WO 01/28992) and potassium carbonate (4.54 g, 0.033 mol) in dry acetonitrile (50 ml) was stirred at 60 °C overnight under nitrogen atmosphere. The reaction mixture was filtered and solvent concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 1 % methanol in dichloromethane as eluent to give the sub-title compound (2.2 g) as white solid.

(iii) 7-[2-(4-Cyano-benzylsulfamoyl)-ethyl]-9-oxa-3,7-diaza-
bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester

7-[2-(4-Cyano-benzylsulfamoyl)-ethyl]-9-oxa-3,7-diaza-

- 5 bicyclo[3.3.1]nonane-3-carboxylic acid tert-butyl ester (2.2 g, from step (ii) above) was taken in dry dioxane (5 ml) and was added 15 ml of dioxane (saturated with HCl gas) and stirred at RT for 30 min. The precipitated solid was filtered, washed with dry diethylether (three times) and finally dried under vacuum to give the title compound (1.96 g) as a solid.

10

Preparation Q

4-(2-Bromoethyl)-1,2-difluorobenzene

(i) 1,2-Difluoro-4-(2-methoxy-vinyl)-benzene

- Potassium *tert* butoxide (11 g, 0.098 mol) was added to methoxymethyl
15 triphenyl phosphonium chloride (31.12 g, 0.091 mol) in 150 ml of dry THF at -30 °C under argon atmosphere and stirred for 1 h. 3,4-difluorobenzaldehyde (10 g, 0.07 mol) in 60 ml of THF was added dropwise at the same temperature and stirring continued for 3 h at room temperature. The reaction mixture was treated with 100 ml water, 150 ml petroleum ether
20 and filtered through celite. Organic layer was separated and aq. layer was extracted with petroleum ether (2 X 100 ml). Combined organic layer was washed with brine and dried over sodium sulfate. Solvent evaporation under reduced pressure followed by purification over silica gel using petroleum ether as eluent afforded (5.5 g) of the sub-title compound as a colorless
25 liquid.

(ii) (3,4-Difluoro-phenyl)-acetaldehyde

To a solution of 1,2-difluoro-4-(2-methoxy-vinyl)-benzene (5.5 g, 0.032mol, from step (i) above) in 80 ml acetone was added 66 ml of 3 (M) HCl and

stirred at 45 °C for 4 h. Acetone was evaporated under reduced pressure and the residue was

partitioned between water and ether. The organic layer was washed with NaHCO₃, brine and dried over sodium sulfate. Solvent evaporation under
5 reduced pressure afforded (5.5 g) of the sub-title compound as a liquid. This was directly taken for next step without further purification.

(iii) 2-(3,4-Difluoro-phenyl)-ethanol

NaBH₄ (1.46 g, 0.038 mol) was added at 0 °C to a solution (3,4-difluoro-
10 phenyl)-acetaldehyde (5.5 g 0.35 mol, from step (ii) above) in 90 ml ethanol and stirred at RT overnight. The solvent was evaporated under reduced pressure and the residue was partitioned between 50 ml of 1 N HCl and 50 ml of diethylether. Organic layer was washed with water, dried over sodium sulfate and solvent evaporated under reduced pressure to give (5.5 g) of the
15 sub-title compound as a liquid. This was directly taken for next step without further purification.

(iv) 4-(2-Bromo-ethyl)-1,2-difluorobenzene

A mixture of 2-(3,4-difluoro-phenyl)-ethanol (5 g, 0.032 mol, from step (iii)
20 above), 51.84 ml (49 %, 0.64 mol) of HBr in acetic acid and 5 ml of sulfuric acid was stirred at 100 °C for 4 h. The reaction mixture was poured into ice and extracted with diethylether. Organic layer was washed with NaHCO₃, water and dried over sodium sulfate. Solvent evaporation under reduced pressure followed by purification (three times) over silica gel using
25 petroleum ether as eluent afforded (1.8 g) of the title compound as a liquid.

Preparation R

4-Bromomethyl-3-fluoro-benzonitrile

(i) (4-Bromo-2-fluoro-phenyl)-methanol

To a solution of 4-bromo-2-fluorobenzaldehyde (25 g, 0.123 mol) in methanol (250 ml) was added NaBH_4 (7.02 g, 0.1847 mol) and the reaction mixture was stirred at RT for 2 h. The reaction mixture was diluted with 25 ml of water and solvent evaporated under reduced pressure. The compound
5 was extracted with ethyl acetate, organic layer was washed with water and brine and dried over sodium sulfate. Solvent evaporation under reduced pressure afforded the sub-title compound (25 g) as an oil. This was directly taken for next step without further purification.

10 (ii) 3-Fluoro-4-hydroxymethylbenzonitrile

A suspension of (4-bromo-2-fluoro-phenyl)-methanol (25 g, 0.122 mol, from step (i) above), $\text{Zn}(\text{CN})_2$ (9.7 g, 0.085 mol) and $\text{Pd}(\text{PPh}_3)_4$ (7.04 g, 0.006 mol) in dry DMF (150 ml) was stirred at 90 °C overnight under nitrogen atmosphere. The reaction mixture was filtered and solvent
15 concentrated under reduced pressure. The residue was purified by column chromatography over silica gel using 25 % ethyl acetate in petroleum ether, as eluent to the sub-title compound (13 g) as white solid.

(iii) 4-Bromomethyl-3-fluorobenzonitrile

20 Carbon tetrabromide (5.7 g, 0.172 mol) in dry dichloromethane (150 ml) as added to a solution of step 3-fluoro-4-hydroxymethylbenzonitrile (13 g, 0.086 mol, from step (ii) above) and triphenyl phosphine (45 g, 0.172 mol) in dry dichloromethane (50 ml) and stirred at RT overnight under nitrogen atmosphere. The reaction was quenched with water, extracted with
25 dichloromethane, washed with water and brine and dried over sodium sulfate. Solvent evaporated under reduced pressure and the residue was purified by column chromatography over silica gel using 10 % ethyl acetate in petroleum ether as eluent to give the title compound (8 g) as an off-white solid.

Preparation S2-(2-Bromo-ethoxy)-1,3-difluorobenzene

A suspension of 2,6-difluoro phenol (18 g, 0.385 mol), 1,2-dibromoethane
5 (71.6 ml, 0.83 mol) and K_2CO_3 (47.86 g, 0.3463 mol) in dry acetonitrile
(190 ml) was stirred at 60 °C overnight under nitrogen atmosphere. The
reaction mixture was filtered and solvent concentrated under reduced
pressure. The residue was purified by column chromatography over silica
gel using 7 % ethyl acetate in petroleum ether as eluent to give the title
10 compound (20.8 g) as colorless oil.

Preparation T4-(3-Bromo-propoxy)-3,5-difluorobenzonitrile

A mixture of 2,6-difluoro-4-cyanophenol (6 g, 0.038 mol), 1,3-
15 dibromopropane (46.9 g, 0.23 mol) and dry potassium carbonate (8 g, 0.058
mol) in dry DMF (50 ml) was stirred at RT overnight under nitrogen
atmosphere. Solvent was removed under reduced pressure and the residue
was purified by column chromatography over silica gel using 4 % ethyl
acetate in petroleum ether as eluent to yield (9.6 g) of the title compound as a
20 liquid.

Preparation U4-(3-Bromo-propoxy)-3-fluorobenzonitrile(i) 3-Fluoro-4-(3-hydroxy-propoxy)-benzonitrile

25 A mixture of 4-cyano-2-fluoro phenol (5 g, 0.0365 mol; from prep N(iv)
above), 3-bromopropanol (10.15 g, 0.073 mol) and potassium carbonate (7.6
g, 0.054 mol) in dry acetone (50 ml) was stirred at 60 °C overnight under
nitrogen atmosphere. The reaction mixture was filtered and the solvent was
concentrated under reduced pressure. The residue was purified by column

chromatography over silica gel using 5 % ethyl acetate in petroleum ether to yield (5.5 g) of the sub-title compound as a solid.

(ii) 4-(3-Bromopropoxy)-3-fluorobenzonitrile

- 5 PBr₃ (8.14 g, 0.03 mol) was added drop by drop at 0 °C to a solution of 3-fluoro-4-(3-hydroxypropoxy)-benzonitrile (5.5 g, 0.028 mol, from step (i) above in carbon tetrachloride (50 ml) and stirred at room temperature overnight under nitrogen atmosphere. The reaction was quenched with water, and extracted with dichloromethane. Organic layer was washed with
10 water, brine, dried over sodium sulfate and solvent evaporated under reduced pressure. The residue was purified by column chromatography over silica gel using 6 % ethyl acetate in petroleum ether as eluent to yield (4.3 g) of the title compound as a liquid.

15 Preparation U

Methanesulfonic acid 2-(2,6-difluoro-phenyl)-ethyl ester

(i) methyl-2,6-difluoro phenyl acetate

- H₂SO₄ (conc., 0.75 ml) was added to a solution of (2,6-difluoro-phenyl)-acetic acid (14 g, 0.0814 mol) in dry methanol (160 ml) and stirred at room
20 temperature overnight. Solvent evaporated under reduced pressure and the residue was partitioned between water and diethyl ether. Organic layer was washed water, brine and dried over sodium sulfate. Solvent evaporation under reduced pressure yielded 16 g of methyl-2, 6-difluoro phenyl acetate as pale yellow liquid. This was directly taken for next step without further
25 purification.

(ii) 2-(2,6-Difluoro-phenyl)-ethanol

LAH (4.7 g, 0.129 mol) in dry diethylether (400 ml) was added at 0 °C to a stirred suspension of methyl-2, 6-difluoro phenyl acetate (16 g, 0.086 mol,

from step (i) above in dry diethylether (80 ml) and stirred at room temperature for 2 h under nitrogen atmosphere. The reaction mixture was quenched with water, NaOH and filtered. The aq. Layer was again acidified with 1.5 N HCl and extracted with diethylether. Organic layer was washed
5 with water, brine and dried over sodium sulfate. Solvent evaporation under reduced pressure afforded (13.5 g) of 2-(2,6-difluoro-phenyl)ethanol as aliquid. This was directly taken for next step without further purification.

(iii) Methanesulfonic acid 2-(2,6-difluoro-phenyl)-ethyl ester

10 Methanesulfonyl chloride (10.76 g, 0.094 mol) was added at 0 °C to a solution of 2-(2,6-difluoro phenyl)ethanol (13.5 g, 0.085 mol, from step (ii) above) and triethylamine (12.93 g, 0.128 mol) in dry dichloromethane (135 ml) and stirred for 3 h under nitrogen atmosphere. The reaction was quenched with water, extracted with dichloromethane, dried over sodium
15 sulfate and solvent evaporated under reduced pressure to give (18 g) of the title compound as an oil.

Preparation V

1-(2-Bromoethyl)-4-trifluoromethylbenzene

20 (i) 1-(2-Methoxy-vinyl)-4-trifluoromethylbenzene

Potassium *tert* butoxide (13.46 g, 0.120 mol) was added to methoxymethyl triphenyl phosphonium chloride (38.30 g, 0.112 mol) in 150 ml dry THF at -30 °C under argon atmosphere and stirred for 1 h. 4-
trifluoromethylbenzaldehyde (15.0 g, 0.086 mol) in 60 ml THF was added
25 dropwise at the same temperature and stirring continued for 3 h at room temperature. The reaction mixture was treated with 100 ml water, 150 ml petroleum ether and filtered through celite. Organic layer was separated and the aq. layer was extracted with petroleum ether (2 X 100 ml). Combined organic layer was washed with brine and dried over sodium sulfate. Solvent

evaporation under reduced pressure followed by purification over silica gel using petroleum ether as eluent afforded (10.5 g) of the sub-title compound as a colorless liquid. This was directly taken for next step without further purification.

5

(ii) (4-Trifluoromethyl-phenyl)-acetaldehyde

To a solution of 1-(2-methoxy-vinyl)-4-trifluoromethylbenzene (10.5 g, 0.052 mol, from step (i) above) in 158 ml acetone was added 107.3 ml 3 M HCl and stirred at 45 °C for 4 h. Acetone was evaporated under reduced pressure and the residue was partitioned between water and ether. Organic layer was washed with NaHCO₃, brine and dried over sodium sulfate. Solvent evaporation under reduced pressure afforded the sub-title compound (9.4 g) as pale yellow liquid. This was directly taken for next step without further purification.

15

(iii) 2-(4-Trifluoromethylphenyl)-ethanol

NaBH₄ (2 g, 0.055 mol) was added at 0 °C to a solution of 4-trifluoromethyl-phenyl)-acetaldehyde (9.4 g, 0.05 mol, from step (ii) above) in 154 ml ethanol and stirred at RT overnight. The solvent was evaporated under reduced pressure and the residue was partitioned between 50 ml 1 N HCl and 50 ml diethyl ether. Organic layer was washed with water, dried over sodium sulfate and the solvent evaporated under reduced pressure to give 10 g of the sub-title compound as a liquid. This was directly taken for next step without further purification.

25

(iv) 1-(2-Bromo-ethyl)-4-trifluoromethylbenzene

A mixture of step 2-(4-trifluoromethylphenyl)-ethanol (10 g, 0.053 mol, from step (iii) above), 86 ml (1.06 M) HBr in acetic acid and 10 ml sulfuric acid was stirred at 100 °C for 4 h. The reaction mixture was poured into ice

and extracted with diethylether. The organic layer was washed with NaHCO_3 , water and dried over sodium sulfate. Solvent evaporation under reduced pressure followed by purification over silica gel using petroleum ether as eluent afforded 5 g of the title compound as a liquid.

5

Preparation W

Methanesulfonic acid 2-(4-difluoromethoxy-phenyl)-ethyl ester

(i) 1-Difluoromethoxy-4-((E,Z)-2-methoxy-vinyl)-benzene (E/Z ratio ~3:2)

A suspension of methoxymethyl triphenylphosphonium chloride (6.69 g, 19.5 mmol) in THF (200 mL) was cooled to ca. -20°C . *n*-BuLi (2.5 M in hexanes, 7.2 mL, 18 mmol) was added in portions over 5 min and the orange-red reaction mixture was stirred for an additional 15 min. 4-difluoromethoxybenzaldehyde (2.58 g, 15.0 mmol) in THF (50 mL) was added at -10°C and the reaction mixture was stirred at this temperature for 30 min and then at room temperature for 19 h. Water (100 mL) was added and the mixture was concentrated under reduced pressure. The remaining aqueous phase was extracted with ethyl acetate (2 x 100 mL) and the combined organic phase was washed with water (2 x 100 mL) and brine (100 mL), dried over Na_2SO_4 , and concentrated *in vacuo*. The crude product was purified by three times repeated chromatography on silica gel. (HorizonTM, flash system from BiotageTM. Column: Flash 40+M, 40 x 150 mm. Eluent: 3% ethyl acetate in heptane. Combination of pure fractions and concentration *in vacuo* afforded 1.60 g (53%) of the title compound as a colourless oil.

¹H-NMR (500 MHz, CDCl_3): δ 7.56 (2H, minor isomer), 7.20 (2H, major isomer), 7.03 (d, 2H, minor isomer), 7.02 (d, 2H, major isomer), 7.00 (d, 1H, major isomer), 6.47 (t, 1H, minor isomer), 6.47 (t, 1H, major isomer), 6.14 (d, 1H, minor isomer), 5.78 (d, 1H, major isomer), 5.20 (d, 1H, minor isomer), 3.78 (s, 3H, minor isomer), 3.69 (s, 3H, major isomer)

(ii) (4-Difluoromethoxyphenyl)-acetaldehyde

To a solution of 1-difluoromethoxy-4-((E,Z)-2-methoxyvinyl)-benzene (**E/Z**
5 **ratio ~ 3:2**,) (1.58 g, 7.9 mmol, from step (i) above) in acetone (20 mL)
was added 3 M HCl (3.7 mL, 11.1 mmol) and the solution was stirred at 45
°C for 4 h. After concentration under reduced pressure, water (25 mL) was
added to the residue and the aqueous phase was extracted with diethylether
(2 x 50 mL). The combined organic phase was washed with saturated
10 aqueous NaHCO₃ (50 mL) and brine (50 mL), dried over Na₂SO₄, and
concentrated *in vacuo* to give 1.32 g of crude product as a yellow oil with a
purity of ca. 80%. The crude product was used directly in the subsequent
reduction step without further purification.

1H-NMR (500 MHz, CDCl₃): δ 9.76 (t, 1H), 7.21 (d, 2H), 7.13 (d, 2H), 6.51
15 (t, 1H), 3.70 (d, 2H)

(iii) 2-(4-Difluoromethoxyphenyl)-ethanol

A solution of (4-difluoromethoxyphenyl)-acetaldehyde 1.22 g, 6 mmol;
from step (ii) above) in ethanol (20 mL) was cooled to 0 °C. Sodium
20 borohydride (0.27 g, 7.2 mmol) was added and the resulting reaction
mixture was stirred at 0 °C for 2 h and then at room temperature for 16 h.
After concentration under reduced pressure, the residue was diluted with
diethylether (50 mL) and 1 M HCl (50 mL). The phases were separated and
the organic phase was washed with water (50 mL) and brine (50 mL), dried
25 over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by
two times repeated chromatography on silica gel. (HorizonTM, flash system
from BiotageTM. Column: Flash 40+M, 40 x 150 mm. Eluent: Gradients
from 25 to 40% ethyl acetate in heptane. Concentration *in vacuo* afforded
0.62 g (50%) of the sub-title compound as a light yellow oil.

¹H-NMR (500 MHz, CDCl₃): δ 7.22 (d, 2H), 7.07 (d, 2H), 6.48 (t, 1H), 3.88–3.80 (m, 2H), 2.85 (t, 2H), 1.55 (m, br, 1H)

(iv) Methanesulfonic acid 2-(4-difluoromethoxyphenyl)-ethyl ester

5 A solution of 2-(4-difluoromethoxyphenyl)-ethanol (0.582 g, 3.09 mmol, from step (iii) above) in methylene chloride (8 mL) was cooled to 0 °C. Triethylamine (0.52 mL, 3.7 mmol) was added followed by a solution of methanesulfonyl chloride (0.26 mL, 3.4 mmol) in methylene chloride (2 mL) and the reaction mixture was stirred at 0 °C for 90 min. The reaction
10 mixture was diluted with methylene chloride (40 mL) and the organic phase was washed with water (2 x 30 mL) and brine (30 mL), dried over Na₂SO₄, and concentrated *in vacuo* to give 0.78 g (95%) of the title compound as a colourless oil.

¹H-NMR (500 MHz, CDCl₃): δ 7.23 (d, 2H), 7.09 (d, 2H), 6.49 (t, 1H), 4.40
15 (t, 2H), 3.05 (t, 2H), 2.89 (s, 3H)

Preparation X

1-(3-Bromopropoxy)-2-fluorobenzene

A mixture of 2-fluorophenol (2.24 g, 20.0 mmol), 1,3-dibromopropane (20.3
20 mL, 200 mmol), and potassium carbonate (2.76 g, 20.0 mmol) in acetonitril (30 mL) was heated at 80 °C overnight. The reaction mixture was filtered and the filtrate was concentrated *in vacuo* to give 4.70 g of the title compound as a light yellow oil with a purity of ca. 80%. The product was used in the subsequent reaction step without further purification.

25

¹H-NMR (500 MHz, CDCl₃): δ 7.11–7.03 (m, 2H), 6.99 (m, 1H), 6.91 (m, 1H), 4.18 (t, 2H), 3.64 (t, 2H), 2.39–2.31 (m, 2H)

Preparation Y

2-(3-Bromo-propoxy)-1,4-difluoro-benzene

A mixture of 2,5-difluorophenol (2.60 g, 20.0 mmol), 1,3-dibromopropane (20.3 mL, 200 mmol), and potassium carbonate (2.76 g, 20.0 mmol) in acetonitril (30 mL) was heated at 80 °C overnight. The reaction mixture was
5 filtered and the filtrate was concentrated *in vacuo* to give 4.82 g of the title compound as a light yellow oil with a purity of ca. 80%. The product was used in the subsequent reaction step without further purification.

¹H-NMR (500 MHz, CDCl₃): δ 7.01 (m, 1H), 6.72 (m, 1H), 6.59 (m, 1H),
10 4.15 (t, 2H), 3.62 (t, 2H), 2.39–2.31 (m, 2H)

Preparation Z4-(3-Bromo-propoxy)-1,2-difluoro-benzene

A mixture of 3,4-difluorophenol (2.60 g, 20.0 mmol), 1,3-dibromopropane
15 (20.3 mL, 200 mmol), and potassium carbonate (2.76 g, 20.0 mmol) in acetonitril (30 mL) was heated at 75 °C overnight. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The crude product was purified by chromatography on silica gel. (HorizonTM, flash system from BiotageTM. Column: Flash 40+M, 40 x 150 mm. Eluent: 3% ethyl acetate in
20 heptane. Concentration *in vacuo* afforded 4.35 g of the title compound as a colourless oil with a purity of ca. 90%.

¹H-NMR (500 MHz, CDCl₃): δ 7.06 (dd, 1H), 6.73 (m, 1H), 6.60 (m, 1H),
4.05 (t, 2H), 3.59 (t, 2H), 2.33–2.27 (m, 2H)

25

Preparation AA1-(3-Bromopropoxy)-3-fluorobenzene

A mixture of 3-fluorophenol (2.24 g, 20.0 mmol), 1,3-dibromopropane (20.3 mL, 200 mmol), and potassium carbonate (2.76 g, 20.0 mmol) in acetonitril

(30 mL) was heated at 80 °C overnight. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The crude product was purified by chromatography on silica gel. (HorizonTM, flash system from BiotageTM. Column: Flash 40+M, 40 x 150 mm. Eluent: 3% ethyl acetate in heptane, 5 Concentration *in vacuo* afforded 3.68 g of the title compound as a colourless oil with a purity of ca. 75%. The product was used in the subsequent reaction step without further purification.

1H-NMR (500 MHz, CDCl₃): δ 7.25 (m, 1H), 6.75–6.63 (m, 2H), 4.12 (t, 10 2H), 3.62 (t, 2H), 2.38–2.31 (m, 2H)

Preparation AB

Methanesulfonic acid 2-(4-fluoro-benzyloxy)-ethyl ester

(i) 2-(4-Fluorobenzyloxy)-ethanol

15 To a suspension of sodium hydride (1.9 g of a 60% suspension in mineral oil, washed with heptane) in THF (20 mL) was added ethylene glycol (19.9 g, 0.32 mol) in portions over 15 min. The reaction mixture was stirred at room temperature for 2 h and then heated to 80 °C. Benzylbromide (6.05 g, 32.0 mmol) in THF (30 mL) was added and the reaction mixture was stirred 20 at 80 °C for 3.5 h. After cooling to room temperature, saturated aqueous NH₄Cl (75 mL) was added and the phases were separated. The water phase was extracted with diethylether (3 x 75 mL) and the combined organic phase was washed with saturated aqueous NH₄Cl (75 mL) and brine (75 mL), dried over MgSO₄ and concentrated *in vacuo* to afford 6.83 g (95%) of the 25 sub-title compound as an oil with a purity of ca. 90%.

1H-NMR (500 MHz, CDCl₃): δ 7.33–7.28 (m, 2H), 7.06–7.00 (m, 2H), 4.51 (s, 2H), 3.77–3.72 (m, 2H), 3.60–3.55 (m, 2H), 2.23 (m, 1H)

(ii) Methanesulfonic acid 2-(4-fluoro-benzyloxy)-ethyl ester

A solution of 2-(4-fluorobenzyloxy)-ethanol (5.11 g, 30.0 mmol, from step (i) above) in methylene chloride (60 mL) was cooled to -10°C . Triethylamine (5.0 mL, 36 mmol) was added followed by methanesulfonyl chloride (3.78 g, 33.0 mmol) and the reaction mixture was stirred at -5°C for 2 h. The reaction mixture was diluted with methylene chloride (200 mL) and the organic phase was washed with water (2 x 75 mL) and brine (75 mL), dried over MgSO_4 , and concentrated *in vacuo* to give 7.16 g (96%) of the title compound as a yellow oil.

$^1\text{H-NMR}$ (500 MHz, CDCl_3): δ 7.33–7.28 (m, 2H), 7.07–7.01 (m, 2H), 4.54 (s, 2H), 4.41–4.37 (m, 2H), 3.76–3.71 (m, 2H), 3.09 (s, 3H)

Preparation AC

4-{[7-(2-aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}benzonitrile, hydrochloride salt.

(i) *tert*-butyl{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}carbamate

Potassium carbonate (3.82 g, 27.6 mmol) was added to a mixture of *tert*-butyl [2-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]carbamate (5.00 g, 18.4 mmol) and 4-(bromomethyl)benzonitrile (3.61 g, 18.4 mmol) in acetonitrile (150 mL). The mixture was heated to reflux over night, cooled to room temperature, filtered and evaporated. The residue was dissolved in dichloromethane (100 mL), washed with water and dried over MgSO_4 .

Purification by preparative HPLC gave 7.04 g (78 %) of the sub.title compound as a salt with one equivalent of acetic acid and one equivalent of formic acid.

¹H NMR (500 MHz, CD₃OD) δ 8.49 (1 H, bs), 7.77 (2 H, d), 7.58 (2 H, d), 4.15 (2 H, m), 3.75 (2 H, s), 3.69 (2 H, m), 3.46 (2 H, t), 3.33 (2 H, m), 3.16 (2 H, t) 3.09 (2 H, m), 2.76 (2 H, m), 1.96 (3 H, s), 1.44 (9 H, s).

5 4-{[7-(2-aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]methyl}benzonitrile, hydrochloride salt.

A saturated solution of HCl in ethyl acetate (200 mL) was added to an ice-cooled slurry of the acetic and formic acid salt of *tert*-butyl {2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}carbamate (7.00
10 g, 14.2 mmol). The reaction mixture was stirred at room temperature for 3 hours after which it was filtered and the resultant solid was washed with ethyl acetate. Drying *in vacuo* gave 6.12 g (96 %) of the title compound with one equivalent of residual ethyl acetate.

15 ¹H NMR (500 MHz, D₂O) δ 7.95 (2 H, d), 7.71 (2 H, d), 4.48 (2H, s), 4.31 (2 H, m), 3.69 (2 H, m), 3.52 (2 H, m), 3.20 (4 H, m), 2.77 (4 H, m).

Preparation AD

Methanesulfonic acid 2-benzo[d]isoxazol-3-yl-ethyl ester

20 (i) 2-Benzo[d]isoxazol-3-yl-ethanol

To a solution of benzo[d]isoxazol-3-yl-acetic acid (0.86 g, 4.85 mmol) in anhydrous THF (40 mL) was added dropwise borane dimethylsulfide complex (0.58 mL, 10 M, 5.8 mmol) at 0°C. The mixture was allowed to warm to room temperature and was stirred over night. MeOH (10 mL) was
25 carefully added and when the gas evolution ceased, the mixture was heated under gentle reflux for 4 hours. The reaction mixture was concentrated *in vacuo* and the remainder was dissolved in dichloromethane (20 mL). An aqueous solution of Na₂CO₃ (20 mL) was added and the aqueous layer was extracted with dichloromethane (3 x 20 mL). The combined organic layers

were dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by chromatography on silica gel which afforded 537 mg (67.8%) of the title compound.

^{13}C NMR (125.7 MHz, CDCl_3) δ 163.06, 156.92, 130.27, 123.65, 121.98,
5 121.65, 110.10, 60.34, 29.01

(ii) Methanesulfonic acid 2-benzo[d]isoxazol-3-yl-ethyl ester

To a solution of 2-benzo[d]isoxazol-3-yl-ethanol (0.537 g, 3.29 mmol, from step (i) above) and triethylamine (0.55 mL, 3.95 mmol) in dichloromethane
10 (20 mL) was added methanesulfonyl chloride (0.31 mL, 3.95 mmol) at 0°C . The reaction mixture was stirred for 1 hour and was then extracted with water (2 x 20 mL) and brine (20 mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*, which afforded 818 mg (98%) of the title compound.

15 ^1H NMR (500 MHz, CDCl_3) δ 7.73 (1H, d), 7.61 (2H, m), 7.37 (1H, m), 4.73 (2H, t), 3.49 (2H, t), 3.01 (3H, s)

Preparation AE

Methanesulfonic acid 2-(2-fluoro-phenyl)-ethyl ester

20 To a solution of 2-(2-fluorophenyl)-ethanol (4.00 g, 28.5 mmol) and triethylamine (6.36 mL, 45.7 mmol) in dichloromethane (15 mL) was added methanesulfonyl chloride (2.88 mL, 37.1 mmol) at -5°C . The mixture was stirred for 4 hours and was allowed to slowly warm to room temperature. The mixture was diluted with dichloromethane and extracted with 1 M HCl
25 and water. The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by column chromatography, using dichloromethane as eluent, to afford 5.88 g (94.4%) the title compound.

^1H NMR (400 MHz, CDCl_3) δ 7.28-7.22 (2H, m), 7.12-7.02 (2H, m), 4.42 (2H, t), 3.10 (2H, t), 2.88 (3H, s)

Examples

Example 1

5 *N*-(2-{7-[(2*S*)-3-(4-Cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide
4-{(2*S*)-3-[7-(2-Aminoethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]-2-hydroxypropoxy}benzonitrile (0.096 g, 0.28 mmol; see step Preparation A above) was dissolved in a 1:2 ratio of DCM:acetonitrile (3 mL).
10 Triethylamine (0.17 g, 1.68 mmol) was added, followed by phenylmethanesulfonyl chloride (0.068 g, 0.36 mmol). The mixture was stirred at room temperature overnight with K₂CO₃ (0.38 g, 2.77 mmol) to ensure that the free base of triethylamine was in the reaction mixture. The reaction mixture was filtered and the filtrate concentrated under reduced
15 pressure. Purification by preparative HPLC gave 0.1 g (72%) of the title compound.

¹H NMR (400 MHz, CD₃OD) δ 7.64-7.66 (2H, d), 7.36-7.43 (5H, m), 7.05-7.07 (2H, m), 4.41 (3H, m), 4.17 (2H, m), 4.08 (2H, m), 3.62-3.7 (2H, m), 3.12-3.4 (8H, m), 2.72-2.86 (2H, m), 2.46-2.59 (2H, m).

20

Example 2

N-(2-{7-[2-(3,4-Bis(difluoromethoxy)phenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyanobenzenesulfonamide

A mixture of 4-cyano-*N*-[2-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-benzenesulfonamide (0.3 g, 7.5 mmol; see Preparation E above), 4-(2-bromo-ethyl)-1,2-bis(difluoromethoxy)benzene (0.26 g, 9 mmol; see Preparation G above) and potassium carbonate (0.41 g, 30 mmol) was stirred at 60°C overnight under a nitrogen atmosphere. The reaction mixture was filtered and solvent was evaporated under reduced pressure. The

residue was purified by column chromatography, using methanol in chloroform as eluent, to yield the title compound as a white solid. Yield: 0.23 g.

^{13}C NMR (75 MHz) δ 144.5, 142.31 140.52, 132.78, 127.51, 126.91, 122.53,
5 122.34, 119.23, 117.29, 116.02, 115.76, 112.29, 68.14, 67.50, 66.72, 61.22,
56.34, 55.35, 54.48, 53.33, 39.07, 38.74, 31.39

Example 3

4-Cyano-N-(2-{7-[2-(4-difluoromethoxy-phenyl)-ethyl]-9-oxa-3,7-diaza- 10 bicyclo[3.3.1]non-3-yl}-ethyl)-benzenesulfonamide

A mixture of 4-cyano-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide dihydrochloride (0.164 g, 0.40 mmol, see prep E above),
methanesulfonic acid 2-(4-difluoromethoxy-phenyl)-ethyl ester (0.128 g,
15 0.48 mmol, from Prep W above), potassium carbonate (0.210 g, 1.52 mmol),
and water (0.1 mL) in acetonitrile (4 mL) was heated by microwave irradiation (160 °C, 15 min). Solid material was filtered off and the filtrate was loaded onto a cation-exchange column (SCX-2, IsoluteTM, 2 g/15 mL).
The column was washed with a solution of methylene
20 chloride/acetonitrile/methanol (2:1:1, 40 mL) and eluted with 20% methanol
saturated with ammonia in methylene chloride (30 mL). The filtrate was concentrated *in vacuo* and the crude product was purified by chromatography on silica gel using methanol in chloroform as eluent.
Concentration *in vacuo* afforded 37 mg (18%) of the title compound as an
25 yellow oil.

^1H -NMR (500 MHz, CDCl_3): δ 8.00 (d, 2H), 7.78 (d, 2H), 7.30 (d, 2H),
7.03 (d, 2H), 6.47 (t, 1H), 4.15–3.70 (m, broad, 2H), 3.70–2.25 (m, 16H)

¹³C-NMR (125 MHz, CDCl₃): δ 149.8, 144.9, 137.3, 133.0, 130.3, 127.7, 119.8, **117.5, 116.1, 116.1** (t), 68.2, 62.2, 56.8, 54.7, 53.6, 39.4, 32.1.

Example 4

5 4-Cyano-N-(2-{7-[2-(4-difluoromethoxy-phenyl)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-N-methyl-benzenesulfonamide

A mixture of 4-cyano-N-methyl-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide dihydrochloride (0.169 g, 0.40 mmol, prep. I above), methanesulfonic acid 2-(4-difluoromethoxy-phenyl)-ethyl ester
10 (0.107 g, 0.40 mmol, prep W above), potassium carbonate (0.177 g, 1.28 mmol), and water (0.1 mL) in acetonitrile (4 mL) was heated in an oil bath (60 °C) for 20 h. Solid material was filtered off and the filtrate was loaded onto a cation-exchange column (SCX-2, IsoluteTM, 2 g/15 mL). The column was washed with a solution of methylene chloride/acetonitrile/methanol
15 (2:1:1, 40 mL) and eluted with 20% methanol saturated with ammonia in methylene chloride (30 mL). The filtrate was concentrated *in vacuo* and the crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane eluent. Concentration *in vacuo* afforded 118 mg (57%) of the title compound as a solid white material.

20 ¹H-NMR (500 MHz, CDCl₃): δ 7.91 (d, 2H), 7.81 (d, 2H), 7.20 (d, 2H), 7.01 (d, 2H), 6.48 (t, 1H), 3.88 (m, br, 2H), 3.21–3.14 (m, 2H), 2.98 (s, 3H), 2.91–2.79 (m, 4H), 2.75–2.67 (m, 2H), 2.58–2.40 (m, 8H)

¹³C-NMR (125 MHz, CDCl₃): δ 149.5, 142.6, 137.9, 133.0, 130.1, 127.9,
25 119.6, **117.4, 116.3, 116.1** (t), 68.4, 61.1, 58.3, 56.4, 46.7, 35.8, 32.3

Example 5

N-{2-[7-(2-Benzo[d]isoxazol-3-yl-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethyl}-4-cyano-benzenesulfonamide

To 4-cyano-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide dihydrochloride (0.123 g, 0.30 mmol, from prep E above), methanesulfonic acid 2-benzo[d]isoxazol-3-yl-ethyl ester (0.076 g, 0.315 mmol, from prep AD above) and potassium carbonate (0.145 g, 1.05 mmol; from Prep) was added acetonitrile (4 mL) and water (0.1 mL). The mixture was heated by microwave irradiation (10 minutes, 160°C) and was then filtered and evaporated. The crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane as eluent, which afforded 60 mg (41.5%) of the title compound.

¹H NMR (500 MHz, CDCl₃) δ 8.00 (2H, d), 7.85 (1H, d), 7.78 (2H, d), 7.55 (2H, m), 7.31 (1H, m), 3.91 (2H, m), 3.39 (2H, m), 3.12-2.47 (14H, m, broad)

Example 6

4-Cyano-N-(2-{7-[2-(2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-N-methyl-benzenesulfonamide

To 4-cyano-N-methyl-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide dihydrochloride (0.127 g, .30 mmol, prep I above), 1-(2-bromo-ethoxy)-2-fluoro-benzene (0.069 g, 0.315 mmol;) and potassium carbonate (0.145 g, 1.05 mmol) was added acetonitrile (4 mL) and water (0.1 mL). The mixture was heated by microwave irradiation (10 minutes, 160°C) and was then filtered and evaporated. The crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane as eluent, which afforded 85 mg (58.0%) of the title compound.

¹³C NMR (125.7 MHz, CDCl₃) δ 153.90, 151.95, 147.02, 142.81, 133.06, 128.02, 124.64, 124.61, 121.54, 117.60, 116.53, 116.38, 116.33, 115.36, 68.52, 67.57, 58.46, 58.27, 56.96, 56.44, 46.66, 35.93

Example 7

2-{7-[2-(3-Fluoro-phenyl)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethanesulfonic acid 4-cyano-benzylamide

5 To 2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethanesulfonic acid 4-cyano-benzylamide dihydrochloride (0.127 g, 0.30 mmol; prep P above), 1-(2-bromo-ethyl)-3-fluoro-benzene (0.064 g, 0.315 mmol) and potassium carbonate (0.145 g, 1.05 mmol) was added acetonitrile (4 mL) and water (0.1 mL). The mixture was stirred for 20 minutes before it was heated by
10 microwave irradiation (10 minutes, 160°C) and was then filtered and evaporated. The crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane as eluent, which afforded 43 mg (30.3%) of the title compound.

¹³C NMR (125.7 MHz, CDCl₃) δ 163.90, 161.94, 144.35, 141.66, 132.65,
15 130.03, 129.96, 128.11, 123.86, 118.70, 115.12, 114.95, 113.51, 113.34, 111.76, 68.34, 60.85, 56.85, 56.41, 54.40, 47.82, 47.47, 31.35

Example 8

4-Cyano-N-{2-[7-(1-methyl-1H-indol-3-ylmethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethyl}-benzenesulfonamide

A mixture of 4-cyano-N-[2-(9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl)-ethyl]-benzenesulfonamide (0.101 g, 0.30 mmol) and 1-methyl-1H-indole-3-carbaldehyde (0.081 g, 0.51 mmol) in 1,2-dichloroethane (4 mL) was agitated for 1.5 hours at room temperature. Sodium triacetoxyborohydride
25 (0.216 g, 1.02 mmol) was added and resulting mixture was agitated over night at room temperature. An aqueous solution of Na₂CO₃ (3 mL) was added and the aqueous layer was extracted with dichloromethane (3 x 4 mL). The combined organic layers were concentrated *in vacuo* and the crude product was purified by chromatography on silica gel using methanol

saturated with ammonia in dichloromethane as eluent, which afforded 76 mg (52.8%) of the title compound.

¹³C NMR (125.7 MHz, CDCl₃) δ 144.59, 136.86, 132.87, 130.73, 129.14, 127.65, 121.86, 119.68, 119.10, 117.56, 116.05, 109.75, 68.56, 56.97,
5 55.04, 54.68, 54.19, 39.40, 32.93

Example 9

N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide

10 4-{2-[7-(2-Amino-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}benzonitrile tri hydrochloride salt (1 g, 2.35 mmol; Prep F) and 3,5-dimethylisoxazole-4-sulfonyl chloride (0.597g, 3 mmol) were dissolved in DCM (25 mL). Triethylamine (1.95 mL, 3 mmol) was added and the reaction mixture was stirred at room temperature overnight under a nitrogen
15 atmosphere. The reaction mixture was concentrated and then redissolved in DCM (25 mL) and transferred to a phase separator. The DCM layer was washed with water (30 mL) and the layers were separated. The retained water was extracted with DCM (25 mL) and the layers were separated. The combined DCM fractions were concentrated under reduced pressure.
20 Purification was performed by precipitation from a mixture of DCM, EtOAc and IPA to give a solid material (0.2 g, 17%).

¹H NMR (400 MHz, CDCl₃) δ 7.58-7.60 (2H, d), 6.98-7.00 (2H, d), 4.30-4.33 (2H, m), 3.89 (2H, m), 2.98-3.01 (4H, m), 2.69-2.81 (8H, m), 2.64 (3H, s), 2.44-2.46 (2H, t), 2.39 (3H, s).

25 ¹³C NMR (125 MHz, CDCl₃) δ 173.11, 162.15 157.69, 134.21, 119.43, 116.30, 115.55, 104.18, 68.62, 67.01, 58.45, 57.50, 54.96, 54.22, 38.97, 12.81, 10.91.

Example 10

N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N,3,5-trimethylisoxazole-4-sulfonamide

N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide (43 mg, 0.09 mmol, from
5 example 9 above) and Cs₂CO₃ (86 mg, 0.26 mmol) were suspended in acetouitrile (2 mL). Iodomethane (17 µl, 0.26 mmol) was added and the reaction mixture was stirred at room temperature overnight under a nitrogen
amosphere. More iodomethane (17 µl, 0.26 mmol) was added. The reaction
mixture was left to stir for a further 20 h at room temperature overnight
10 under a nitrogen atmosphere. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure. The crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane as eluent, which afforded 39mg (88%) of the title compound.

15 ¹H NMR (500 MHz, CDCl₃) δ 7.59-7.61 (2H, d), 6.96-6.98 (2H, d), 4.12-4.15 (2H, m), 3.90 (2H, m), 3.23-3.26 (2H, m), 2.98 (3H, s), 2.86-2.92 (4H, m), 2.68-2.75 (4H, m), 2.63 (3H, s), 2.49-2.56 (4H, m), 2.40 (3H, s).

Example 11

20 N-(2-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-2,4-difluoro-benzenesulfonamide
4-{2-[7-(2-Amino-ethyl)-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl]-ethoxy}-3-fluoro-benzonitrile hydrochloride salt (0.266 g, 0.6 mmol; prep N above) and 2,4-difluorobenzenesulfonyl chloride (0.170 g, 0.80 mmol) were
25 dissolved in DCM (6 mL). Triethylamine (0.416 mL, 3 mmol) was added and the reaction mixture was stirred at room temperature for 20 h under a nitrogen amosphere. The reaction mixture was concentrated and then redissolved in DCM (3 mL) and transferred to a phase separator. The DCM layer was washed with water (3 mL) and the layers were separated. The

retained water was extracted with DCM (3 mL) and the layers were separated. The combined DCM fractions were concentrated under reduced pressure. The crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane as eluent, which
5 afforded 173mg (56%) of the title compound.

¹H NMR (500 MHz, CD₃OD) δ 7.89-7.94 (1H, m), 7.46-7.50 (2H, m), 7.10-7.26 (3H, m), 4.42-4.45 (2H, t), 3.86 (2H, m), 3.05-3.09 (4H, m), 2.78-2.82 (4H, m), 2.60-2.65 (4H, m), 2.34-2.37 (2H, m).

10

Example 12

N-(2-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-2,4-difluoro-N-methyl-benzenesulfonamide

Cyanomethylenetri-n-butylphosphorane (90%, 0.158g, 0.59 mmol) was
15 weighed into a microwave vial. N-(2-{7-[2-(4-Cyano-2-fluoro-phenoxy)-ethyl]-9-oxa-3,7-diaza-bicyclo[3.3.1]non-3-yl}-ethyl)-2,4-difluoro-benzenesulfonamide (100 mg, 0.2 mmol, from example 11 above) and THF (2 mL) were added. Methanol (16 µl, 0.39 mmol) was then added and the reaction was heated with microwave irradiation at 160 °C for 15 min.
20 MeOH (1 mL) was added and the reaction was heated with microwave irradiation for a further 10 min in order to consume excess reagent.

The reaction mixture was concentrated and the crude product was purified by chromatography on silica gel using methanol saturated with ammonia in dichloromethane as eluent. The product isolated could be purified further by
25 recrystallisation from MeOH to give a solid (55 mg, 53%).

¹H NMR (500 MHz, CDCl₃) δ 7.85-7.90 (1H, m), 7.42-7.44 (1H, m), 7.34-7.37 (1H, m), 7.09-7.13 (1H, m), 6.93-7.02 (2H, m), 4.23-4.26 (2H, m), 3.88-3.90 (2H, m), 3.28-3.30 (2H, m), 2.99 (3H, s), 2.89-2.95 (4H, m), 2.77-2.79 (2H, m), 2.70-2.72 (2H, m), 2.48-2.54 (4H, m).

Example 13

The following compounds were prepared, from appropriate intermediates (such as those described hereinbefore), according to or by analogy with

5 methods described herein:

- (i) N-(2-{7-[(2S)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)propane-2-sulfonamide;
- (ii) N-(tert-butyl)-3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-
10 diazabicyclo[3.3.1]non-3-yl}propane-1-sulfonamide;
- (iii) tert-butyl {2-[7-(2-{[(4-fluorophenyl)sulfonyl]amino}ethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl]carbamate;
- (iv) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-fluorobenzenesulfonamide;
- 15 (v) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- (vi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)propane-2-sulfonamide;
- (vii) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-
20 cyanobenzenesulfonamide;
- (viii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (ix) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- 25 (x) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xi) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;

- (xii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xiii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- 5 (xiv) 4-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xv) 1-(3-chlorophenyl)-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- (xvi) 4-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (xvii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1,1,1-trifluoromethanesulfonamide;
- (xviii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- 15 (xix) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-methyl-1H-imidazole-4-sulfonamide;
- (xx) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)butane-1-sulfonamide;
- (xxi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-methyl-1H-imidazole-4-sulfonamide;
- 20 (xxii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-[4-(trifluoromethyl)phenyl]methanesulfonamide;
- (xxiii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-1-[4-(trifluoromethyl)phenyl]methanesulfonamide;
- 25 (xxiv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;

- (xxv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-2-(trifluoromethoxy)benzenesulfonamide;
- (xxvi) 1-(3-chlorophenyl)-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- 5 (xxvii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxviii) 4-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl}benzenesulfonamide;
- (xxix) 4-cyano-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo-
- 10 [3.3.1]non-3-yl}ethyl}benzenesulfonamide;
- (xxx) 4-cyano-N-(2-{7-[4-(difluoromethoxy)benzyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxi) 4-cyano-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (xxxii) 4-cyano-N-(2-{7-[2-(4-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxiii) 4-cyano-N-(2-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxiv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
- 20 [3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (xxxv) 5-chloro-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,3-dimethyl-1H-pyrazole-4-sulfonamide;
- (xxxvi) 2-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-
- 25 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxvii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-4-methoxybenzenesulfonamide;
- (xxxviii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;

- (xxxix) 5-chloro-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,3-dimethyl-1H-pyrazole-4-sulfonamide;
- (xl) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-methoxybenzenesulfonamide;
- 5 (xli) 4-cyano-N-(2-{7-[3-(2-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlii) 4-cyano-N-(2-{7-[3-(2,5-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (xliii) 4-cyano-N-(2-{7-[3-(3,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xliv) 4-cyano-N-(2-{7-[2-(2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlv) 4-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (xlv) 4-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlvii) 4-cyano-N-[2-(7-{2-[(4-fluorobenzyl)oxy]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- (xlviii) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 20 (xlviii) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlix) 4-cyano-N-{2-[7-(2-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (l) 4-cyano-N-(2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (l) 4-cyano-N-(2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (li) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- (lii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;

- (liii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)-5-methylisoxazole-4-sulfonamide;
- (liv) N-{2-[7-(1-benzofuran-3-yl)methyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl}-4-cyanobenzenesulfonamide;
- 5 (lv) 4-cyano-N-{2-[7-(1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl}benzenesulfonamide;
- (lvi) 4-cyano-N-(2-{7-[(5-fluoro-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lvii) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
10 [3.3.1]non-3-yl}propyl)-4-fluorobenzenesulfonamide;
- (lviii) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-2,4-difluorobenzenesulfonamide;
- (lix) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 15 (lx) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl}-1-phenylmethanesulfonamide;
- (lxi) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-1-phenylmethanesulfonamide;
- (lxii) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
20 [3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (lxiii) 4-cyano-N-(2-{7-[(2-methyl-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxiv) 4-cyano-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (lxv) 4-cyano-N-(2-{7-[3-(3-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxvi) 4-cyano-N-(2-{7-[3-(4-cyano-2-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (lxvii) 4-cyano-N-(2-{7-[3-(4-cyano-2,6-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxviii) 4-cyano-N-(2-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 5 (lxix) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)propane-2-sulfonamide;
- (lxx) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-1-phenylmethanesulfonamide;
- (lxxi) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-
- 10 [3.3.1]non-3-yl}propyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxii) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxiii) N-{3-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl}-3,5-dimethylisoxazole-4-sulfonamide;
- 15 (lxxiv) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxv) N-{2-[7-(1,3-benzoxazol-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (lxxvi) 3-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-
- 20 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxvii) 3-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxviii) 3-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 25 (lxxix) 2-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (lxxx) 2-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (lxxxix) N-(4-cyanobenzyl)-2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethanesulfonamide;
- (lxxxii) N-(4-cyanobenzyl)-2-[7-(2-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethanesulfonamide;
- 5 (lxxxiii) N-(4-cyanobenzyl)-N-[(3,5-dimethylisoxazol-4-yl)methyl]-2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (lxxxiv) 4-cyano-N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 10 (lxxxv) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-4-fluorobenzenesulfonamide;
- (lxxxvi) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2-fluorobenzenesulfonamide;
- (lxxxvii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2-fluorobenzenesulfonamide;
- 15 (lxxxviii) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2,4-difluorobenzene-sulfonamide;
- (lxxxix) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 20 (xc) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xci) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- 25 (xcii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (xciii) 3-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (xciv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xcv) N-(2-{7-[2-(4-chlorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyanobenzenesulfonamide;
- 5 (xcvi) 4-cyano-N-[2-(7-{2-[4-(trifluoromethyl)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- (xcvii) 4-cyano-N-(2-{7-[2-(2,6-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xcviii) 4-cyano-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (xcix) 4-cyano-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (c) 4-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- 15 (ci) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyano-N-methylbenzenesulfonamide;
- (cii) 4-cyano-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- (ciii) 4-cyano-N-methyl-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 20 (civ) 4-cyano-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cv) 4-cyano-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 25 (cvi) N-(4-cyanobenzyl)-2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (cvii) N-(4-cyanobenzyl)-2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;

- (cviii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- (cix) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- 5 (cx) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- (cxi) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-fluorobenzenesulfonamide;
- 10 (cxii) 4-fluoro-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (cxiii) N-(2-{7-[2-(4-cyanophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (cxiv) 4-fluoro-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (cxv) 4-cyano-N-(2-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxvi) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-N-methylbenzenesulfonamide;
- 20 (cxvii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methyl-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (cxviii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methyl-2,3-dihydro-1-benzofuran-5-sulfonamide;
- 25 (cxix) 4-cyano-N-{2-[7-(4-cyano-2-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (cxx) 4-cyano-N-{2-[7-(4-cyano-2-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;

- (cxxi) 4-fluoro-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxxii) 4-fluoro-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 5 (cxxiii) 4-fluoro-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (cxxiv) N-{2-[7-(1,2-benzisoxazol-3-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (cxxv) 4-cyano-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 10 (cxxvi) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxvii) 4-cyano-N-{3-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzenesulfonamide;
- 15 (cxxviii) 4-cyano-N-{3-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzenesulfonamide;
- (cxxix) 4-cyano-N-(3-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxxx) 4-cyano-N-(3-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 20 (cxxxii) 4-cyano-N-(3-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxxxiii) 4-cyano-N-(3-{7-[2-(2,6-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 25 (cxxxiv) 4-cyano-N-(3-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxxxv) N-[3-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propyl]-4-cyanobenzenesulfonamide;

- (cxxxv) 4-cyano-N-(3-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxxxvi) 4-cyano-N-(3-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 5 (cxxxvii) 4-cyano-N-(2-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxxviii) 4-cyano-N-(2-{7-[3-(2,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxxix) 4-cyano-N-(2-{7-[2-(2,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (cxl) 4-cyano-N-(2-{7-[3-(2,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxli) N-(2-{7-[2-(4-chlorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyano-N-methylbenzenesulfonamide; and
- 15 (cxlii) 4-Cyano-N-(2-{7-[2-(2,4-difluoro-phenyl)-ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-ethyl)-N-methyl-benzenesulfonamide;

Example 14

Title compounds of the above Examples were tested in Test A above and were
20 found to exhibit D₁₀ values of more than 5.5.

Example 15

Title compounds of the above Examples were tested in Test B above and were
found to exhibit pIC₅₀ values of greater than 4.5. Indeed the compounds of
25 Examples 1 and 9 were found to have pIC₅₀ values of 5.72 and 5.49,
respectively.

Abbreviations


	Ac	=	acetyl
	API	=	atmospheric pressure ionisation (in relation to MS)
	aq.	=	aqueous
	br	=	broad (in relation to NMR)
5	Bt	=	benzotriazole
	<i>t</i> -BuOH	=	<i>tert</i> -butanol
	CI	=	chemical ionisation (in relation to MS)
	<i>m</i> CPBA	=	<i>meta</i> -chloroperoxybenzoic acid
	d	=	doublet (in relation to NMR)
10	DBU	=	diazabicyclo[5.4.0]undec-7-ene
	DCM	=	dichloromethane
	dd	=	doublet of doublets (in relation to NMR)
	DMAP	=	4-dimethylaminopyridine
	DMF	=	<i>N,N</i> -dimethylformamide
15	DMSO	=	dimethylsulfoxide
	EDC	=	1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide
	Et	=	ethyl
	EtOAc	=	ethyl acetate
	eq.	=	equivalents
20	ES	=	electrospray (in relation to MS)
	FAB	=	fast atom bombardment (in relation to MS)
	FBS	=	foetal bovine serum
	h	=	hour(s)
	HCl	=	hydrochloric acid
25	HEPES	=	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
	HPLC	=	high performance liquid chromatography
	IMS	=	industrial methylated spirits
	IPA	=	<i>iso</i> -propyl alcohol (propan-2-ol)
	m	=	multiplet (in relation to NMR)

	Me	=	methyl
	MeCN	=	acetonitrile
	MeOH	=	methanol
	min.	=	minute(s)
5	m.p.	=	melting point
	MS	=	mass spectroscopy
	NADPH	=	nicotinamide adenine dinucleotide phosphate, reduced form
	OAc	=	acetate
10	Pd/C	=	palladium on carbon
	q	=	quartet (in relation to NMR)
	RT	=	room temperature
	s	=	singlet (in relation to NMR)
	t	=	triplet (in relation to NMR)
15	TEA	=	triethylamine
	THF	=	tetrahydrofuran
	tlc	=	thin layer chromatography

Prefixes *n*-, *s*-, *i*-, *t*- and *tert*- have their usual meanings: normal, secondary,
20 *iso*, and tertiary.

Claims

1. A compound of formula I,



5

wherein

R¹ represents C₁₋₁₂ alkyl (which alkyl group is optionally substituted by one or more groups selected from halo, cyano, nitro, aryl, Het¹, -N(R^{5a})R⁶,
 10 -C(O)R^{5b}, -OR^{5c}, -C(O)XR⁷, -C(O)N(R^{8a})R^{5d}, -OC(O)N(R^{8b})R^{5e}, -S(O)₂R^{9a},
 -S(O)₂N(R^{9b})R^{9c} and -N(R^{9b})S(O)₂R^{9d}) or R¹ represents -C(O)XR⁷,
 -C(O)N(R^{8a})R^{5d} or -S(O)₂R^{9a};

R^{5a} represents H or C₁₋₆ alkyl (which latter group is optionally substituted by
 15 one or more substituents selected from -OH, halo, cyano, nitro,
 -S(O)₂N(R^{9b})R^{9c} and -N(R^{9b})S(O)₂R^{9d});

R^{5b} to R^{5e} independently represent, at each occurrence when used herein, H,
 C₁₋₆ alkyl (which latter group is optionally substituted by one or more
 substituents selected from -OH, halo, cyano, nitro, aryl, Het²,
 20 -S(O)₂N(R^{9b})R^{9c} and -N(R^{9b})S(O)₂R^{9d}), aryl or Het³, or R^{5d} or R^{5e}, together
 with, respectively, R^{8a} or R^{8b}, may represent C₃₋₆ alkylene (which alkylene

group is optionally interrupted by an O atom and/or is optionally substituted by one or more C₁₋₃ alkyl groups);

R⁶ represents H, C₁₋₆ alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, -S(O)₂N(R^{9b})R^{9c} and -N(R^{9b})S(O)₂R^{9d}), aryl, -C(O)R^{10a}, -C(O)OR^{10b} or -C(O)N(H)R^{10c} or -S(O)₂R^{10d};

R^{10a}, R^{10b}, R^{10c} and R^{10d} independently represent C₁₋₆ alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, or R^{10a} represents H;

R⁷ represents, at each occurrence when used herein, C₁₋₁₂ alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, C₁₋₆ alkoxy, Het⁴, -S(O)₂N(R^{9b})R^{9c} and -N(R^{9b})S(O)₂R^{9d});

R^{8a} and R^{8b} independently represent H, C₁₋₁₂ alkyl, C₁₋₆ alkoxy (which latter two groups are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C₁₋₄ alkyl, C₁₋₄ alkoxy, -S(O)₂N(R^{9b})R^{9c} and -N(R^{9b})S(O)₂R^{9d}), -D-aryl, -D-aryloxy, -D-Het⁵, -D-N(H)C(O)R^{11a}, -D-S(O)₂R^{12a}, -D-C(O)R^{11b}, -D-C(O)OR^{12b}, -D-C(O)N(R^{11c})R^{11d}, or R^{8a} or R^{8b}, together with, respectively, R^{5d} or R^{5e}, may represent C₃₋₆ alkylene (which alkylene group is optionally interrupted by an O atom and/or is optionally substituted by one or more C₁₋₃ alkyl groups);

R^{11a} to R^{11d} independently represent H, C₁₋₆ alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro and aryl), aryl, or R^{11c} and R^{11d} together represent C₃₋₆ alkylene;

R^{12a} and R^{12b} independently represent C₁₋₆ alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro and aryl) or aryl;

D represents, at each occurrence when used herein, a direct bond or C₁₋₆ alkylene;

X represents O or S;

R^{9a} represents, at each occurrence when used herein, C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl, $-S(O)_2N(R^{9b})R^{9c}$ and $-N(R^{9b})S(O)_2R^{9d}$) or aryl;

R^{9b} represents, at each occurrence when used herein, H or C_{1-6} alkyl;

R^{9c} and R^{9d} independently represent, at each occurrence when used herein, C_{1-6} alkyl (optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, aryl and Het⁶), aryl or Het⁷, or R^{9c} represents H;

10

R^2 represents H, halo, C_{1-6} alkyl, $-OR^{13}$, $-E-N(R^{14})R^{15}$ or, together with R^3 , represents =O;

R^3 represents H, C_{1-6} alkyl or, together with R^2 , represents =O;

R^{13} represents H, C_{1-6} alkyl, -E-aryl, -E-Het⁸, $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or $-C(O)N(R^{17a})R^{17b}$;

R^{14} represents H, C_{1-6} alkyl, -E-aryl, -E-Het⁹, $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$, $-[C(O)]_pN(R^{17a})R^{17b}$ or $-C(NH)NH_2$;

R^{15} represents H, C_{1-6} alkyl, -E-aryl or $-C(O)R^{16d}$;

R^{16a} to R^{16d} independently represent, at each occurrence when used herein, C_{1-6} alkyl (optionally substituted by one or more substituents selected from halo, aryl and Het¹⁰), aryl, Het¹¹, or R^{16a} and R^{16d} independently represent H;

R^{17a} and R^{17b} independently represent, at each occurrence when used herein, H or C_{1-6} alkyl (optionally substituted by one or more substituents selected from halo, aryl and Het¹²), aryl, Het¹³, or together represent C_{3-6} alkylene,

optionally interrupted by an O atom;

E represents, at each occurrence when used herein, a direct bond or C_{1-4} alkylene;

p represents 1 or 2;

Het¹ to Het¹³ independently represent five- to twelve-membered heterocyclic groups containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, which groups are optionally substituted by one or more substituents selected from -OH, oxo, halo, cyano, nitro,
 5 C₁₋₆ alkyl, C₁₋₆ alkoxy, aryl, aryloxy, -N(R^{18a})R^{18b}, -C(O)R^{18c}, -C(O)OR^{18d},
 -C(O)N(R^{18e})R^{18f}, -N(R^{18g})C(O)R^{18h}, -S(O)₂N(R¹⁸ⁱ)R^{18j} and
 -N(R^{18k})S(O)₂R^{18l};

R^{18a} to R^{18l} independently represent C₁₋₆ alkyl, aryl or R^{18a} to R^{18k} independently represent H;

10

A represents a direct bond, -J-, -J-N(R^{19a})-, -J-S(O)₂N(R^{19b})-,
 -J-N(R^{19c})S(O)₂- or -J-O- (in which latter four groups, J is attached to the
 oxabispidine ring nitrogen);

B represents -Z-[[C(O)]_aC(H)(R^{20a})]_b-, -Z-[C(O)]_cN(R^{20b})-,
 15 -Z-N(R^{20c})S(O)₂-, -Z-S(O)₂N(R^{20d})-, -Z-S(O)_n-, -Z-O- (in which latter six
 groups, Z is attached to the carbon atom bearing R² and R³), -N(R^{20e})-Z-,
 -N(R^{20f})S(O)₂-Z-, -S(O)₂N(R^{20g})-Z- or -N(R^{20h})C(O)O-Z- (in which latter
 four groups, Z is attached to the phenyl or pyridyl group that is optionally
 substituted by R⁴);

20 J represents C₁₋₆ alkylene optionally interrupted by -S(O)₂N(R^{19d})- or
 -N(R^{19e})S(O)₂- and/or optionally substituted by one or more substituents
 selected from -OH, halo and amino;

Z represents a direct bond or C₁₋₄ alkylene, optionally interrupted by
 -N(R²⁰ⁱ)S(O)₂- or -S(O)₂N(R^{20j})-;

25 a, b and c independently represent 0 or 1;

n represents 0, 1 or 2;

R^{19a} to R^{19e} independently represent, at each occurrence when used herein,
 H or C₁₋₆ alkyl;

R^{20a} represents H or, together with a single R^4 substituent at a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, R^{20a} represents C_{2-4} alkylene optionally interrupted or terminated by O, S, N(H) or N(C_{1-6} alkyl);

- 5 R^{20b} represents H, C_{1-6} alkyl or, together with a single R^4 substituent at a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, R^{20b} represents C_{2-4} alkylene;

R^{20c} to R^{20j} independently represent, at each occurrence when used herein, H or C_{1-6} alkyl;

10

G represents CH or N;

- R^4 represents one or more optional substituents selected from -OH, cyano, halo, nitro, C_{1-6} alkyl (optionally terminated by $-N(H)C(O)OR^{21a}$),
 15 C_{1-6} alkoxy, $-N(R^{22a})R^{22b}$, $-C(O)R^{22c}$, $-C(O)OR^{22d}$, $-C(O)N(R^{22e})R^{22f}$,
 $-N(R^{22g})C(O)R^{22h}$, $-N(R^{22i})C(O)N(R^{22j})R^{22k}$, $-N(R^{22m})S(O)_2R^{21b}$,
 $-S(O)_2N(R^{22n})R^{22o}$, $-S(O)_2R^{21c}$, $-OS(O)_2R^{21d}$ and aryl

and an R^4 substituent in a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached may

- 20 (i) together with R^{20a} , represent C_{2-4} alkylene optionally interrupted or terminated by O, S or N(H) or N(C_{1-6} alkyl), or
 (ii) together with R^{20b} , represent C_{2-4} alkylene;

R^{21a} to R^{21d} independently represent C_{1-6} alkyl;

- R^{22a} and R^{22b} independently represent H, C_{1-6} alkyl or together represent C_{3-6}
 25 alkylene, resulting in a four- to seven-membered nitrogen-containing ring;

R^{22c} to R^{22o} independently represent H or C_{1-6} alkyl; and

R^{41} to R^{46} independently represent H or C_{1-3} alkyl;

wherein each aryl and aryloxy group, unless otherwise specified, is optionally substituted;

provided that

- 5 (a) at least one of the following is the case
 - (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group),
 - (ii) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
 - 10 (iii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
 - (iv) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
 - (v) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$;
- (b) when A represents $-J-N(R^{19a})-$, $-J-N(R^{19c})S(O)_2-$ or $-J-O-$, then:
 - 15 (i) J does not represent C_1 alkylene or 1,1- C_{2-6} alkylene; and
 - (ii) B does not represent $-N(R^{20b})-$, $-N(R^{20c})S(O)_2-$, $-S(O)_n-$, $-O-$, $-N(R^{20f})S(O)_2-Z-$ or $-N(R^{20h})C(O)O-Z-$ when R^2 and R^3 do not together represent $=O$; and
- (c) when R^2 represents $-OR^{13}$ or $-N(R^{14})(R^{15})$, then:
 - 20 (i) A does not represent $-J-N(R^{19a})-$, $-J-N(R^{19c})S(O)_2-$ or $-J-O-$; and
 - (ii) B does not represent $-N(R^{20b})-$, $-N(R^{20c})S(O)_2-$, $-S(O)_n-$, $-O-$, $-N(R^{20f})S(O)_2-Z-$ or $-N(R^{20h})C(O)O-Z-$;

or a pharmaceutically acceptable derivative thereof.

25

2. A compound as claimed in Claim 1, wherein R^1 represents C_{1-8} alkyl (which alkyl group is optionally substituted by one or more groups selected from halo, aryl (which latter group is optionally substituted by one or more substituents selected from $-OH$, halo, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy

(which latter two groups are optionally substituted by one or more halo atoms), $-C(O)R^{22c}$ and $-S(O)_2R^{21c}$, Het^1 , $-N(R^{5a})R^6$, $-C(O)R^{5b}$, $-OR^{5c}$, $-C(O)N(R^{8a})R^{5d}$, $-OC(O)N(R^{8b})R^{5e}$, $-S(O)_2R^{9a}$, $-S(O)_2N(R^{9b})R^{9c}$ and $-N(H)S(O)_2R^{9d}$ or R^1 represents $-C(O)OR^7$, $-C(O)N(R^{8a})R^{5d}$ or $-S(O)_2R^{9a}$;

5

3. A compound as claimed in Claim 1 or Claim 2, wherein R^{5a} represents H or C_{1-5} alkyl and/or R^{5b} to R^{5e} independently represent, at each occurrence when used herein, H, C_{1-6} alkyl (which latter group is optionally substituted by one or more substituents selected from cyano, nitro and optionally substituted aryl), aryl (which latter group is optionally substituted by one or more substituents selected from $-OH$, halo, cyano, nitro, $N(R^{22a})R^{22b}$ (in which latter group R^{22a} and R^{22b} together represent C_{3-6} alkylene), C_{1-5} alkyl and C_{1-5} alkoxy (which latter two groups are optionally substituted by one or more halo atoms)), Het^3 , or R^{5d} , together with R^{8a} , represents C_{4-5} alkylene (which alkylene group is optionally interrupted by an O atom);

4. A compound as claimed in any one of the preceding claims, wherein, wherein R^6 represents H, C_{1-6} alkyl, optionally substituted aryl $-C(O)R^{10a}$, $-C(O)OR^{10b}$ or $-C(O)N(H)R^{10c}$.

5. A compound as claimed in any one of the preceding claims, wherein R^{10a} and R^{10b} independently represent C_{1-5} alkyl (optionally substituted by one or more substituents selected from halo and optionally substituted aryl) or optionally substituted aryl.

6. A compound as claimed in any one of the preceding claims, wherein R^{10c} represents C_{1-4} alkyl.

7. A compound as claimed in any one of the preceding claims, wherein R^7 represents C_{1-6} alkyl (optionally substituted by one or more substituents selected from halo, optionally substituted aryl, C_{1-4} alkoxy and Het⁴).
- 5 8. A compound as claimed in any one of the preceding claims, wherein R^{8a} and R^{8b} independently represent H, C_{1-6} alkyl (which latter group is optionally substituted by one or more substituents selected from halo, cyano and nitro), -D-(optionally substituted aryl), -D-(optionally substituted aryloxy), -D-Het⁵, -D-N(H)C(O) R^{11a} , -D-C(O) R^{11b} , or R^{8a} , together with R^{5d}
10 represents C_{4-5} alkylene (which alkylene group is optionally interrupted by an O atom).
9. A compound as claimed in any one of the preceding claims, wherein R^{11a} and R^{11d} independently represent C_{1-4} alkyl (optionally substituted by
15 one or more substituents selected from halo, cyano, nitro and optionally substituted aryl) or optionally substituted aryl.
10. A compound as claimed in any one of the preceding claims, wherein D represents, at each occurrence when used herein, a direct bond or C_{1-4}
20 alkylene.
11. A compound as claimed in any one of the preceding claims, wherein R^{9a} represents C_{1-6} alkyl (optionally substituted by one or more halo groups) or optionally substituted aryl.
25
12. A compound as claimed in any one of the preceding claims, wherein R^{9b} represents H or C_{1-3} alkyl.

13. A compound as claimed in any one of the preceding claims, wherein R^{9c} and R^{9d} independently represent, at each occurrence when used herein, C_{1-5} alkyl (optionally substituted by one or more substituents selected from halo, optionally substituted aryl and Het^6), optionally substituted aryl or Het^7 , or R^{9c} represents H;
14. A compound as claimed in any one of the preceding claims, wherein R^2 represents H, halo, C_{1-3} alkyl, $-OR^{13}$, $-N(H)R^{14}$ or, together with R^3 , represents $=O$.
15. A compound as claimed in any one of the preceding claims, wherein R^3 represents H, C_{1-3} alkyl or, together with R^2 , represents $=O$.
16. A compound as claimed in any one of the preceding claims, wherein R^{13} represents H, C_{1-4} alkyl, $-E$ -(optionally substituted aryl), or $-E-Het^8$.
17. A compound as claimed in any one of the preceding claims, wherein R^{14} represents H, C_{1-6} alkyl, $-E$ -(optionally substituted aryl), $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$, $-C(O)N(R^{17a})R^{17b}$ or $-C(NH)NH_2$.
18. A compound as claimed in any one of the preceding claims, wherein R^{16a} to R^{16c} independently represent C_{1-6} alkyl, or R^{16a} represents H.
19. A compound as claimed in any one of Claims 1 to 18, wherein R^{17a} and R^{17b} independently represent H or C_{1-4} alkyl.
20. A compound as claimed in any one of the preceding claims, wherein E represents, at each occurrence when used herein, a direct bond or C_{1-2} alkylene.

21. A compound as claimed in any one of the preceding claims, wherein Het¹ and Het³ to Het⁸ independently represent four- to ten-membered heterocyclic groups containing one to four heteroatoms selected from oxygen, nitrogen and/or sulfur, which groups are optionally substituted by one or more substituents selected from -OH, oxo, halo, cyano, C₁₋₄ alkyl, C₁₋₄ alkoxy, phenyl, -N(H)R^{18a}, -C(O)R^{18c}, -N(H)C(O)R^{18h} and -N(H)S(O)₂R^{18j};
22. A compound as claimed in any one of the preceding claims, wherein R^{18a}, R^{18c}, R^{18h} and R^{18j} independently represent C₁₋₄ alkyl or optionally substituted aryl or R^{18a}, R^{18c} and R^{18h} independently represent H.
23. A compound as claimed in any one of the preceding claims, wherein A represents C₁₋₄ alkylene optionally substituted by one or more substituents selected from OH and amino, -C₁₋₃ *n*-alkylene-S(O)₂N(H)- or -C₂₋₃ *n*-alkylene-N(H)S(O)₂- (in which latter two groups, alkylene is attached to the oxabispidine ring nitrogen).
24. A compound as claimed in any one of the preceding claims, wherein B represents -Z-, -Z-N(H)-, -Z-C(O)N(R^{20b})-, -Z-N(R^{20c})S(O)₂-, -Z-S(O)₂N(R^{20d})-, -Z-S(O)₂-, -Z-O- (in which latter six groups, Z is attached to the carbon atom bearing R² and R³).
25. A compound as claimed in any one of the preceding claims, wherein Z represents a direct bond or C₁₋₄ alkylene.
26. A compound as claimed in any one of the preceding claims, wherein R^{20b} represents H, C₁₋₄ alkyl or, together with a single R⁴ substituent at a

position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached, R^{20b} represents C₂₋₄ alkylene.

27. A compound as claimed in any one of the preceding claims, wherein
5 R^{20c} and R^{20d} independently represent H or C₁₋₃ alkyl.

28. A compound as claimed in any one of the preceding claims, wherein, when G represents N, G is in the *ortho*- or *para*-position relative to the point of attachment of B.

29. A compound as claimed in any one of the preceding claims, wherein, when G represents N, R⁴ is absent or represents a single cyano group.

30. A compound as claimed in any one of Claims 1 to 28, wherein R⁴,
15 represents one or more substituents selected from -OH, cyano, halo, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, -C(O)N(R^{22e})R^{22f}, and -N(R^{22m})S(O)₂-C₁₋₄ alkyl, or an R⁴ substituent in a position on the phenyl or pyridyl group that is *ortho*- to the position at which the group B is attached may, together with R^{20b}, represent C₂₋₄ alkylene.

31. A compound as claimed in any one of Claims 1 to 28 and 30, wherein
20 R^{21c} represents C₁₋₄ alkyl and R^{22c}, R^{22e}, R^{22f} and R^{22m} independently represent H or C₁₋₄ alkyl.

32. A compound as claimed in any one of the preceding claims, wherein
25 R⁴¹ to R⁴⁶ independently represent H.

33. A compound as claimed in any one of the preceding claims, wherein optional substituents on aryl and aryloxy groups are, unless otherwise stated,

one or more substituents selected from halo, cyano, nitro, C₁₋₄ alkyl, C₁₋₄ alkoxy (which latter two groups are optionally substituted by one or more halo atoms), -N(H)S(O)₂R^{21b} and -S(O)₂N(H)R^{22o}.

- 5 34. A compound as claimed in any one of the preceding claims, wherein alkyl groups and alkoxy groups may be, unless otherwise specified:
- (i) straight- or branched-chain or cyclic or part cyclic/acyclic;
 - (ii) saturated or unsaturated;
 - (iii) interrupted by one or more oxygen atoms; and/or
 - 10 (iv) substituted by one or more fluoro or chloro atoms.
35. A compound as claimed in Claim 1 which is:
- (i) N-(2-{7-[(2S)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)propane-2-sulfonamide;
 - 15 (ii) N-(2-{7-[(2S)-3-(4-cyanophenoxy)-2-hydroxypropyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
 - (iii) N-(tert-butyl)-3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propane-1-sulfonamide;
 - (iv) tert-butyl {2-[7-(2-{[(4-fluorophenyl)sulfonyl]amino}ethyl)-9-oxa-
20 3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}carbamate;
 - (v) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-fluorobenzenesulfonamide;
 - (vi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
 - 25 (vii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)propane-2-sulfonamide;
 - (viii) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyanobenzenesulfonamide;

- (ix) N-[2-(7-{2-[3,4-bis(difluoromethoxy)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyanobenzenesulfonamide;
- (x) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 5 (xi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- (xii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xiii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- 10 (xiv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (xv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- 15 (xvi) 4-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xvii) 1-(3-chlorophenyl)-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- (xviii) 4-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 20 (xix) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,1,1-trifluoromethanesulfonamide;
- (xx) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- 25 (xxi) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-methyl-1H-imidazole-4-sulfonamide;
- (xxii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)butane-1-sulfonamide;

- (xxiii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)-1-methyl-1H-imidazole-4-sulfonamide;
- (xxiv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)-1-[4-(trifluoromethyl)phenyl]methane-
5 sulfonamide;
- (xxv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (xxvi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)-1-[4-(trifluoromethyl)phenyl]methane-
10 sulfonamide;
- (xxvii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (xxviii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)-2-(trifluoromethoxy)benzenesulfonamide;
- 15 (xxix) 1-(3-chlorophenyl)-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-
diazabicyclo[3.3.1]non-3-yl}ethyl)methanesulfonamide;
- (xxx) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxi) 4-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo-
20 [3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (xxxii) 4-cyano-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo-
[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (xxxiii) 4-cyano-N-(2-{7-[4-(difluoromethoxy)benzyl]-9-oxa-3,7-
diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (xxxiv) 4-cyano-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-
diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxv) 4-cyano-N-(2-{7-[2-(4-fluorophenoxy)ethyl]-9-oxa-3,7-
diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (xxxvi) 4-cyano-N-(2-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xxxvii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- 5 (xxxviii) 5-chloro-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,3-dimethyl-1H-pyrazole-4-sulfonamide;
- (xxxix) 2-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (xl) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-methoxybenzenesulfonamide;
- (xli) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (xlii) 5-chloro-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1,3-dimethyl-1H-pyrazole-4-sulfonamide;
- 15 (xliii) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-methoxybenzenesulfonamide;
- (xliv) 4-cyano-N-(2-{7-[3-(2-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 20 (xlv) 4-cyano-N-(2-{7-[3-(2,5-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlvi) 4-cyano-N-(2-{7-[3-(3,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (xlvii) 4-cyano-N-(2-{7-[2-(2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (xlviii) 4-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;

- (xlix) 4-cyano-N-[2-(7-{2-[(4-fluorobenzyl)oxy]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- (l) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 5 (li) 4-cyano-N-(2-{7-[2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lii) 4-cyano-N-{2-[7-(2-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (liii) 4-cyano-N-(2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (liv) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- (lv) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- 15 (lvi) N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-5-methylisoxazole-4-sulfonamide;
- (lvii) N-{2-[7-(1-benzofuran-3-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (lviii) 4-cyano-N-{2-[7-(1H-indol-3-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 20 (lix) 4-cyano-N-(2-{7-[(1-methyl-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lx) 4-cyano-N-(2-{7-[(5-fluoro-1H-indol-3-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 25 (lxi) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-4-fluorobenzenesulfonamide;
- (lxii) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2,4-difluorobenzenesulfonamide;

- (*lxiii*) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (*lxiv*) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-1-phenylmethanesulfonamide;
- 5 (*lxv*) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-1-phenylmethanesulfonamide;
- (*lxvi*) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (*lxvii*) 4-cyano-N-(2-{7-[(2-methyl-1H-indol-3-yl)methyl]-9-oxa-3,7-
10 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*lxviii*) 4-cyano-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*lxix*) 4-cyano-N-(2-{7-[3-(3-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (*lxx*) 4-cyano-N-(2-{7-[3-(4-cyano-2-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*lxxi*) 4-cyano-N-(2-{7-[3-(4-cyano-2,6-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*lxxii*) 4-cyano-N-(2-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-
20 diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*lxxiii*) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)propane-2-sulfonamide;
- (*lxxiv*) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-1-phenylmethanesulfonamide;
- 25 (*lxxv*) N-(3-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (*lxxvi*) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo-[3.3.1]non-3-yl}propyl)-3,5-dimethylisoxazole-4-sulfonamide;

- (lxxvii) N-{3-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}-3,5-dimethylisoxazole-4-sulfonamide;
- (lxxviii) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-3,5-dimethylisoxazole-4-sulfonamide;
- 5 (lxxix) N-{2-[7-(1,3-benzoxazol-2-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (lxxx) 3-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxxi) 3-cyano-N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 10 (lxxxii) 3-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (lxxxiii) 2-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 15 (lxxxiv) 2-cyano-N-(2-{7-[3-(4-cyanophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (lxxxv) N-(4-cyanobenzyl)-2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethanesulfonamide;
- (lxxxvi) N-(4-cyanobenzyl)-2-[7-(2-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethanesulfonamide;
- 20 (lxxxvii) N-(4-cyanobenzyl)-N-[(3,5-dimethylisoxazol-4-yl)methyl]-2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (lxxxviii) 4-cyano-N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 25 (lxxxix) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-4-fluorobenzenesulfonamide;
- (xc) N-(3-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2-fluorobenzenesulfonamide;

- (*xcii*) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2-fluorobenzenesulfonamide;
- (*xciii*) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-2,4-difluorobenzene-sulfonamide;
- (*xciv*) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (*xcv*) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3-fluorobenzenesulfonamide;
- (*xcvi*) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-3,5-dimethylisoxazole-4-sulfonamide;
- (*xcvii*) 3-cyano-N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*xcviii*) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*xcix*) N-(2-{7-[2-(4-chlorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyanobenzenesulfonamide;
- (*c*) 4-cyano-N-[2-(7-{2-[4-(trifluoromethyl)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- (*ci*) 4-cyano-N-(2-{7-[2-(2,6-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*cii*) 4-cyano-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (*ciii*) 4-cyano-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;

- (civ) 4-cyano-N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- (cv) N-[2-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-4-cyano-N-methylbenzenesulfonamide;
- 5 (cvi) 4-cyano-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- (cvii) 4-cyano-N-methyl-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- (cviii) 4-cyano-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 10 (cix) 4-cyano-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cx) N-(4-cyanobenzyl)-2-{7-[(3,5-dimethylisoxazol-4-yl)methyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- 15 (cxi) N-(4-cyanobenzyl)-2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (cxii) N-(4-cyanobenzyl)-2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethanesulfonamide;
- (cxiii) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N,3,5-trimethylisoxazole-4-sulfonamide;
- 20 (cxiv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-1-phenylmethanesulfonamide;
- (cxv) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,3-dihydro-1-benzofuran-5-sulfonamide;
- 25 (cxvi) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;
- (cxvii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluorobenzenesulfonamide;

- (cxviii) N-{2-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-fluorobenzenesulfonamide;
- (cxix) 4-fluoro-N-{2-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 5 (cxx) N-(2-{7-[2-(4-cyanophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-fluorobenzenesulfonamide;
- (cxxi) 4-fluoro-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxxii) 4-cyano-N-[2-(7-{2-[4-(difluoromethoxy)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]benzenesulfonamide;
- 10 (cxxiii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-2,4-difluoro-N-methylbenzenesulfonamide;
- (cxxiv) 4-cyano-N-(2-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 15 (cxxv) 4-cyano-N-(2-{7-[2-(2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxvi) N-(2-{7-[2-(1,2-benzisoxazol-3-yl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyanobenzenesulfonamide;
- 20 (cxxvii) N-(3-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)-N-methylbenzenesulfonamide;
- (cxxviii) N-(2-{7-[2-(4-cyano-2-fluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methyl-2,3-dihydro-1-benzofuran-5-sulfonamide;
- 25 (cxxix) N-(2-{7-[2-(4-cyanophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methyl-2,3-dihydro-1-benzofuran-5-sulfonamide;
- (cxxx) 4-cyano-N-{2-[7-(4-cyano-2-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;


- (cxxxix) 4-cyano-N-{2-[7-(4-cyano-2-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-N-methylbenzenesulfonamide;
- (cxxxix) 4-cyano-N-[2-(7-{2-[4-(difluoromethoxy)phenyl]ethyl}-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)ethyl]-N-methylbenzenesulfonamide;
- 5 (cxxxix) 4-fluoro-N-(2-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxxxix) 4-fluoro-N-(2-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cxxxix) 4-fluoro-N-{2-[7-(2-phenylethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}benzenesulfonamide;
- 10 (cxxxix) N-{2-[7-(1,2-benzisoxazol-3-ylmethyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]ethyl}-4-cyanobenzenesulfonamide;
- (cxxxix) 4-cyano-N-(2-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 15 (cxxxix) 4-cyano-N-(2-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cxxxix) 4-cyano-N-{3-[7-(4-cyanobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzenesulfonamide;
- (cxli) 4-cyano-N-{3-[7-(4-fluorobenzyl)-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl]propyl}benzenesulfonamide;
- 20 (cxli) 4-cyano-N-(3-{7-[2-(4-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlii) 4-cyano-N-(3-{7-[2-(3-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- 25 (cxlii) 4-cyano-N-(3-{7-[2-(2-fluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlii) 4-cyano-N-(3-{7-[2-(2,6-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;

- (cxlv) 4-cyano-N-(3-{7-[2-(3,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlvi) N-[3-(7-benzyl-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl)propyl]-4-cyanobenzenesulfonamide;
- 5 (cxlvii) 4-cyano-N-(3-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlviii) 4-cyano-N-(3-{7-[2-(2,6-difluorophenoxy)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}propyl)benzenesulfonamide;
- (cxlix) 4-cyano-N-(2-{7-[3-(4-fluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- 10 (cl) 4-cyano-N-(2-{7-[3-(2,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-N-methylbenzenesulfonamide;
- (cli) 4-cyano-N-(2-{7-[2-(2,4-difluorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- 15 (clii) 4-cyano-N-(2-{7-[3-(2,4-difluorophenoxy)propyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)benzenesulfonamide;
- (cliii) N-(2-{7-[2-(4-chlorophenyl)ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}ethyl)-4-cyano-N-methylbenzenesulfonamide; or
- (cliv) 4-Cyano-N-(2-{7-[2-(2,4-difluoro-phenyl)-ethyl]-9-oxa-3,7-diazabicyclo[3.3.1]non-3-yl}-ethyl)-N-methyl-benzenesulfonamide;
- 20

or a pharmaceutically acceptable salt thereof

36. A pharmaceutical formulation including a compound as defined in any one of Claims 1 to 35 in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.
- 25

37. A pharmaceutical formulation for use in the prophylaxis or the treatment of an arrhythmia, comprising a compound as defined in any one of Claims 1 to 35.
- 5 38. A compound as defined in any one of Claims 1 to 35 for use as a pharmaceutical.
39. A compound as defined in any one of Claims 1 to 35 for use in the prophylaxis or the treatment of an arrhythmia.
- 10 40. The use of a compound as defined in any one of Claims 1 to 35 as active ingredient for the manufacture of a medicament for use in the prophylaxis or the treatment of an arrhythmia.
- 15 41. The use as claimed in Claim 40, wherein the arrhythmia is an atrial or a ventricular arrhythmia.
42. A method of prophylaxis or treatment of an arrhythmia which method comprises administration of a therapeutically effective amount of a compound as defined in any one of Claims 1 to 35 to a person suffering from, or susceptible to, such a condition.
- 20 43. A compound according to any one of claims 1 -35 in combination with any other drugs for use as a medicament for the treatment of arrhythmias or cardiovascular disorders.
- 25 44. A process for the preparation of a compound of formula I as defined in Claim 1, which process comprises:
- (a) reaction of a compound of formula II,




wherein R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as defined in Claim 1, with a compound of formula III,



5 wherein L^1 represents a leaving group;

(b) for compounds of formula I in which R^1 represents C_{1-12} alkyl substituted by one or more substituents as defined in Claim 1 in respect of R^1 , which substituent(s) is/include a $-N(R^{9b})S(O)_2R^{9d}$ group, reaction of a compound of formula IV,




10

wherein R^{1a} represents C_{1-12} alkylene, which group is optionally substituted by one or more substituents as defined in Claim 1 in respect of R^1 , and R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G as defined in Claim 1, with a compound of formula V,




wherein L^2 represents a leaving group;

(c) for compounds of formula I in which R^1 represents C_{1-12} alkyl substituted by one or more substituents as defined above in respect of R^1 , which substituent(s) is/include a $-S(O)_2N(R^{9b})R^{9c}$ group, reaction of a compound of formula II, as defined above, with a compound of formula VA,



wherein L^1 and R^{1a} are as defined above and R^{9b} and R^{9c} are as defined in Claim 1;

(d) for compounds of formula I in which R^1 represents $-C(O)XR^7$ or $-C(O)N(R^{8a})R^{5d}$, reaction of a compound of formula VI,




10

wherein R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as defined in Claim 1 and L^1 is as defined above, with a compound of formula VII,



wherein R^{24} represents $-XR^7$ or $-N(R^{8a})R^{5d}$ and R^{5d} , R^7 , R^{8a} and X are as defined in Claim 1;

(e) for compounds of formula I in which R^1 represents $-C(O)N(H)R^{8a}$, reaction of a compound of formula II, as defined above, with a compound of formula VIII,




wherein R^{8a} is as defined in Claim 1;


(f) reaction of a compound of formula IX,

20

153




wherein R^1 and R^{41} to R^{46} as defined in Claim 1, with a compound of formula X,



- 5 wherein L^3 represents a leaving group and R^2 , R^3 , R^4 , A, B and G are as defined in Claim 1;

(g) for compounds of formula I in which B represents $-Z-N(R^{20c})S(O)_2-$, reaction of a compound of formula IX, as defined above, with a compound of formula XA,




10

wherein R^2 , R^3 , R^4 , A, G and Z are as defined in Claim 1, and L^3 is as defined above;

(h) for compounds of formula I in which A represents CH_2 and R^2 represents $-OH$ or $-N(H)R^{14}$, reaction of a compound of formula IX, as defined above,


- 15 with a compound of formula XI,

154




wherein Y represents O or N(R¹⁴) and R³, R⁴, R¹⁴, B and G are as defined in Claim 1;

- (i) for compounds of formula I in which B represents -Z-O-, reaction of a
5 compound of formula XII,




wherein R¹, R², R³, R⁴¹ to R⁴⁶, A and Z are as defined in Claim 1, with a compound of formula XIII,




- 10 wherein R⁴ and G are as defined in Claim 1;

(j) for compounds of formula I in which G represents N and B represents -Z-O-, reaction of a compound of formula XII, as defined above, with a compound of formula XIV,



- 15 wherein R⁴ is as defined in Claim 1 and L³ is as defined above;


(k) for compounds of formula I in which R^2 represents $-OR^{13}$, in which R^{13} represents C_{1-6} alkyl, $-E$ -aryl or $-E$ -Het⁸, reaction of a compound of formula I in which R^2 represents OH with a compound of formula XV,



XV

- 5 wherein R^{13a} represents C_{1-6} alkyl, $-E$ -aryl or $-E$ -Het⁸ and E and Het⁸ are as defined in Claim 1;

(l) for compounds of formula I in which R^2 represents $-OR^{13}$, in which R^{13} represents C_{1-6} alkyl, $-E$ -aryl or $-E$ -Het⁸, reaction of a compound of formula XVI,




XVI

10

wherein R^1 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as defined in Claim 1 and L^3 is as defined above, with a compound of formula XV, as defined above;

(m) for compounds of formula I in which R^2 represents $-E-NH_2$, reduction of a compound of formula XVII,



XVII

15

wherein R^1 , R^3 , R^4 , R^{41} to R^{46} , A, B, E and G are as defined in Claim 1;

(n) for compounds of formula I in which R^2 represents $-E-N(R^{14})R^{15}$, wherein R^{14} represents C_{1-6} alkyl, $-E$ -aryl, $-E$ -Het⁹, $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$ or $-C(O)N(R^{17a})R^{17b}$, reaction of a compound of formula I in which R^2 represents $-E-N(H)R^{15}$ with a compound of formula XVIII,



wherein R^{14a} represents C_{1-6} alkyl, $-E$ -aryl, $-E$ -Het⁹, $-C(O)R^{16a}$, $-C(O)OR^{16b}$, $-S(O)_2R^{16c}$ or $-C(O)N(R^{17a})R^{17b}$, and R^{16a} , R^{16b} , R^{16c} , R^{17a} , R^{17b} , Het⁹ and E are as defined in Claim 1 and L^1 is as defined above;

(o) for compounds of formula I in which R^2 represents $-E-N(R^{15})C(O)N(H)R^{17a}$, reaction of a compound of formula I in which R^2 represents $-E-N(H)R^{15}$ with a compound of formula XIX,



wherein R^{17a} is as defined in Claim 1;

(p) for compounds of formula I in which R^2 represents $-E-N(H)[C(O)]_2NH_2$, reaction of a compound of formula I in which R^2 represents $-E-NH_2$ with oxalic acid diamide;

(q) for compounds of formula I in which R^2 represents $-E-N(H)C(NH)NH_2$, reaction of a compound of formula I in which R^2 represents $-E-NH_2$ with a compound of formula XX,



or an *N*-protected derivative thereof, wherein R^{23} is as defined above;

(r) for compounds of formula I in which R^2 represents $-OR^{13}$, in which R^{13} represents $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or $-C(O)N(R^{17a})R^{17b}$, reaction of a compound of formula I in which R^2 represents $-OH$ with a compound of formula XXI,




wherein R^{13b} represents $-C(O)R^{16a}$, $-C(O)OR^{16b}$ or $-C(O)N(R^{17a})R^{17b}$, L^4 represents a leaving group and R^{16a} , R^{16b} , R^{17a} and R^{17b} are as defined in Claim 1;

157


(s) for compounds of formula I in which R^2 represents H or -OH and R^3 represents H, reduction of a compound of formula I in which R^2 and R^3 together represent =O;

(t) for compounds of formula I in which R^2 represents halo, substitution of a
 5 corresponding compound of formula I in which R^2 represents -OH, using a halogenating agent;

(u) for compounds of formula I in which R^2 and R^3 represent H, A represents -J- and B represents $-N(R^{20e})-Z-$ (wherein $-N(R^{20e})$ is attached to the carbon atom bearing R^2 and R^3), reaction of a compound of formula XXII,




wherein R^1 , R^{20e} , R^{41} to R^{46} and J are as defined in Claim 1, with a compound of formula XXIII,



wherein R^4 , G and Z are as defined in Claim 1 and L^3 is as defined above;


15 (v) for compounds of formula I in which A represents $-(CH_2)_2-$ and R^2 and R^3 together represent =O, reaction of a compound of formula IX, as defined above, with a compound of formula XXIV,




158

wherein B, G and R⁴ are as defined in Claim 1;

(w) for compounds of formula I in which R¹ represents -C(O)XR⁷, -C(O)N(R^{8a})R^{5d} or -S(O)₂R^{9a}, reaction of a compound of formula XXV,



- 5 wherein R^{1b} represents -C(O)XR⁷, -C(O)N(R^{8a})R^{5d} or -S(O)₂R^{9a} and R^{5d}, R⁷, R^{8a}, R^{9a} and R⁴¹ to R⁴⁶ are as defined in Claim 1 and L³ is as defined above, with a compound of formula XXVI,



wherein R², R³, R⁴, A, B and G as defined in Claim 1;

- 10 (x) for compounds of formula I which are oxabispidine-nitrogen *N*-oxide derivatives, oxidation of the corresponding oxabispidine nitrogen of a corresponding compound of formula I, in the presence of an oxidising agent;


- (y) for compounds of formula I which are C₁₋₄ alkyl quaternary ammonium salt derivatives, in which the alkyl group is attached to a oxabispidine nitrogen, reaction, at the oxabispidine nitrogen, of a corresponding compound of formula I with a compound of formula XXVII,



wherein R²⁵ represents C₁₋₄ alkyl and L⁵ is a leaving group;


- 20 (z) conversion of one R⁴ substituent to another;
- (aa) introduction of one or more (further) R⁴ substituents to the aromatic ring;
- (ab) for compounds of formula I in wherein R¹ represents C₁₋₁₂ alkylene,

which group is optionally substituted by one or more substituents as defined above in respect of R^1 , reaction of a compound of formula II




- 5 wherein R^2 , R^3 , R^4 , R^{41} to R^{46} , A, B and G are as hereinbefore defined, with the appropriate aldehyde, for example under conditions that are known to those skilled in the art (e.g. at room temperature, such as from 15 to 30°C) in the presence of a reducing agent (such as sodium cyanoborohydride, sodium triacetoxymethylborohydride, or similar compounds) and an appropriate solvent
- 10 (such as 1,2-dichloroethane, dichloroethane, methanol, ethanol or mixtures thereof);

- (ac) for compounds of formula I in wherein A represents -J-, -J-N(R^{19a})-, -J-S(O)₂N(R^{19b})-, -J-N(R^{19c})S(O)₂- or -J-O- (in which latter four groups, -J is attached to the oxabispidine ring nitrogen), reaction of a compound with
- 15 formula VII



160

wherein R^1 and R^{41} to R^{46} are as hereinbefore defined, with a compound of formula XXIII



5

wherein R^2 , R^4 , B and G are as hereinbefore defined and the C_{0-5} alkylene group is optionally substituted by one or more substituents as defined above in respect of J, for example under conditions that are known to those skilled in the art (e.g. at room temperature, such as from 15 to 30°C) in the presence of

10 a reducing agent (such as sodium cyanoborohydride, sodium triacetoxymethylborohydride, or similar hydride donating compounds) and an appropriate solvent (such as 1,2-dichloroethane, dichloroethane, methanol, ethanol or mixtures thereof); or

(ad) deprotection of a protected derivative of a compound of formula I as

15 defined above.

45. A compound of formula II, as defined in Claim 44, or a protected derivative thereof, provided that

- (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- 20 (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20g})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$.

46. A compound of formula IV, as defined in Claim 44, or a protected derivative thereof, provided that

- (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- 5 (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$.

47. A compound of formula VI, as defined in Claim 44, or a protected derivative thereof, provided that

- (i) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- (iii) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- 15 (iv) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$.

48. A compound of formula IX, as defined in Claim 44, or a protected derivative thereof, provided that R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group).

49. A compound of formula XII, as defined in Claim 44, or a protected derivative thereof, provided that

- (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group),
- 25 (ii) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$, and/or
- (iii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$.

50. A compound of formula XVII, as defined in Claim 44, or a protected derivative thereof, provided that

- (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group),
- (ii) A represents $-J-S(O)_2N(R^{19b})-$ or $-J-N(R^{19c})S(O)_2-$,
- (iii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$,
- (iv) B represents $-Z-N(R^{20c})S(O)_2-$, $-Z-S(O)_2N(R^{20d})-$, $-N(R^{20f})S(O)_2-Z-$ or $-S(O)_2N(R^{20g})-Z-$ and/or
- (v) Z is interrupted by $-N(R^{20i})S(O)_2-$ or $-S(O)_2N(R^{20j})-$.

51. A compound of formula XXII, as defined in Claim 44, or a protected derivative thereof, provided that

- (i) R^1 represents C_{1-12} alkyl (which alkyl group is substituted by one or more groups including at least one $-S(O)_2N(R^{9b})R^{9c}$ and/or $-N(R^{9b})S(O)_2R^{9d}$ group), and/or
- (ii) J is interrupted by $-S(O)_2N(R^{19d})-$ or $-N(R^{19e})S(O)_2-$.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE 2005/000891

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 498/08, A61K 31/4386, A61P 9/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CHEM. ABS DATA, WPI DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 0128992 A2 (ASTRAZENECA AB), 26 April 2001 (26.04.2001)	1-51
	--	
A	WO 0228863 A1 (ASTRAZENECA AB), 11 April 2002 (11.04.2002)	1-51
	-- -----	

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

15 Sept 2005

Date of mailing of the international search report

20-09-2005

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Carolina Gómez Lagerlöf/EÖ

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/000891

WO	0128992	A2	26/04/2001	AP	200202471	D	00/00/0000
				AT	292134	T	15/04/2005
				AU	769513	B	29/01/2004
				AU	1069101	A	30/04/2001
				BG	106588	A	31/01/2003
				BR	0014804	A	11/06/2002
				CA	2386910	A	26/04/2001
				CN	1399639	A,T	26/02/2003
				CZ	20021328	A	11/09/2002
				DE	60019158	D	00/00/0000
				DK	1235831	T	27/06/2005
				EA	5319	B	24/02/2005
				EE	200200203	A	16/06/2003
				EP	1235831	A,B	04/09/2002
				SE	1235831	T3	
				EP	1559717	A	03/08/2005
				HR	20020336	A	29/02/2004
				HU	0204085	A	28/04/2003
				IL	149108	D	00/00/0000
				JP	2003512352	T	02/04/2003
				MX	PA02003848	A	23/10/2002
				NO	20021811	A	06/06/2002
				NZ	518156	A	28/05/2004
				PL	354970	A	22/03/2004
				SE	9903759	D	00/00/0000
				SK	5062002	A	06/11/2002
				TR	200201059	T	00/00/0000
				US	6559143	B	06/05/2003
				US	20030153564	A	14/08/2003
				ZA	200202575	A	02/07/2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/000891

WO	0228863	A1	11/04/2002	AU	9250301	A	15/04/2002
				AU	9250401	A	15/04/2002
				BR	0114366	A	12/08/2003
				BR	0114367	A	09/12/2003
				CA	2421776	A	11/04/2002
				CA	2422810	A	11/04/2002
				CN	1468244	A,T	14/01/2004
				CN	1468245	A,T	14/01/2004
				CZ	294217	B	13/10/2004
				CZ	20030923	A	13/08/2003
				EE	200300131	A	16/06/2003
				EP	1330461	A	30/07/2003
				EP	1330462	A	30/07/2003
				HU	0302288	A	28/10/2003
				IL	154802	D	00/00/0000
				IL	154803	D	00/00/0000
				JP	2004510775	T	08/04/2004
				JP	2004510776	T	08/04/2004
				MX	PA03002679	A	24/06/2003
				MX	PA03002759	A	28/07/2003
				NO	20031413	A	22/05/2003
				NO	20031414	A	26/05/2003
				NZ	524573	A	26/11/2004
				NZ	524574	A	26/11/2004
				PL	364047	A	13/12/2004
				SK	3862003	A	07/10/2003
				US	20040039199	A	26/02/2004
				US	20050171100	A	04/08/2005
				WO	0228864	A	11/04/2002
				ZA	200301757	A	05/02/2004
				ZA	200302202	A	28/06/2004
				AP	200202471	D	00/00/0000
				AT	274154	T	15/09/2004
				AT	292134	T	15/04/2005
				AU	769513	B	29/01/2004
				AU	1069101	A	30/04/2001
				AU	4702901	A	03/12/2001
				BG	106588	A	31/01/2003
				BR	0014804	A	11/06/2002
				BR	0111046	A	15/04/2003
				CA	2386910	A	26/04/2001
				DE	60019158	D	00/00/0000
				DE	60105022	D,T	18/08/2005
				DK	1235831	T	27/06/2005
				EA	5319	B	24/02/2005
				EE	200200203	A	16/06/2003
				EP	1235831	A,B	04/09/2002
				SE	1235831	T3	

INTERNATIONAL SEARCH REPORT

Information on patent family members

31/08/2005

International application No.

PCT/SE 2005/000891

EP	1290361 A,B	12/03/2003
HR	20020336 A	29/02/2004
HU	0204085 A	28/04/2003
IL	149108 D	00/00/0000
JP	2003512352 T	02/04/2003
MX	PA02003848 A	23/10/2002
NO	20021811 A	06/06/2002
NZ	518156 A	28/05/2004
PL	354970 A	22/03/2004
SE	515575 C	03/09/2001
SE	0001994 A	03/09/2001
SK	5062002 A	06/11/2002
US	20030093981 A	22/05/2003
WO	0190604 A	29/11/2001

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2005/000891

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 4 2
because they relate to subject matter not required to be searched by this Authority, namely:
See next sheet.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2005/000891

Box II.1

Claim 42 relates to a method of treatment of the human or animal body by surgery or by therapy, as well as diagnostic methods /Rule 39.1(iv). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compounds.