



Office de la Propriété

Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2477751 A1 2003/10/09

(21) 2 477 751

(12) **DEMANDE DE BREVET CANADIEN**  
**CANADIAN PATENT APPLICATION**

(13) A1


(86) Date de dépôt PCT/PCT Filing Date: 2003/03/28  
(87) Date publication PCT/PCT Publication Date: 2003/10/09  
(85) Entrée phase nationale/National Entry: 2004/08/30  
(86) N° demande PCT/PCT Application No.: EP 2003/003263  
(87) N° publication PCT/PCT Publication No.: 2003/082285  
(30) Priorité/Priority: 2002/04/03 (02007568.5) EP

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> A61K 31/46, A61K 31/415, A61K 31/135,  
A61P 37/08  
(71) **Demandeur/Applicant:**  
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,  
DE  
(72) **Inventeur/Inventor:**  
SEKO, NORITAKA, JP  
(74) **Agent:** FETHERSTONHAUGH & CO.

(54) Titre : COMBINAISONS D'EPINASTINE, DE BELLADONE ET DE PSEUDOEPHEDRINE EN TANT QUE  
NOUVELLES FORMULATIONS PHARMACEUTIQUES  
(54) Title: COMBINATIONS OF EPINASTINE, BELLADONNA AND PSEUDOEPHEDRINE AS NEW PHARMACEUTICAL  
FORMULATIONS

(57) **Abrégé/Abstract:**

The present invention relates to novel oral pharmaceutical compositions comprising as pharmaceutically active compounds a combination of an antihistaminic-effective amount of epinastine or a pharmaceutically acceptable salt thereof, an anticholinergic amount of Belladonna alkaloids (Belladonna) or a pharmaceutically acceptable salt thereof and of a decongestant-effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof. Optionally, the formulation may comprise methylephedrine (methylephrine) in a decongestant-effective amount or a pharmaceutically acceptable salt thereof. The formulation further comprises suitable pharmaceutically acceptable carriers or excipients. Another aspect of the present invention relates to methods for the preparation of these compositions and methods of using them in the treatment of allergic diseases and/or disorders. In particular the inventive composition is useful in the treatment of saisonal allergic rhinitis and saisonal allergic conjunctivitis.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number  
WO 03/082285 A1(51) International Patent Classification<sup>7</sup>: A61K 31/46,  
A61P 37/08 // (A61K 31/46, 31:415, 31:135)CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,  
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number: PCT/EP03/03263

(22) International Filing Date: 28 March 2003 (28.03.2003)

(25) Filing Language: English

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(26) Publication Language: English

(30) Priority Data:  
02007568.5 3 April 2002 (03.04.2002) EP

## Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(71) Applicant (for all designated States except US):  
BOEHRINGER INGELHEIM INTERNATIONAL  
GMBH [DE/DE]; Binger Str. 173, 55216 INGELHEIM  
(DE).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventor; and  
(75) Inventor/Applicant (for US only): SEKO, Noritaka  
[JP/JP]; 3-2-10 Shibutani, Ikeda, Osaka 563-0028 (JP).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

WO 03/082285 A1

(54) Title: COMBINATIONS OF EPINASTINE, BELLADONNA AND PSEUDOEPHEDRINE AS NEW PHARMACEUTICAL FORMULATIONS

(57) Abstract: The present invention relates to novel oral pharmaceutical compositions comprising as pharmaceutically active compounds a combination of an antihistaminic-effective amount of epinastine or a pharmaceutically acceptable salt thereof, an anticholinergic amount of Belladonna alkaloids (Belladonna) or a pharmaceutically acceptable salt thereof and of a decongestant-effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof. Optionally, the formulation may comprise methylephedrine (methylephrine) in a decongestant-effective amount or a pharmaceutically acceptable salt thereof. The formulation further comprises suitable pharmaceutically acceptable carriers or excipients. Another aspect of the present invention relates to methods for the preparation of these compositions and methods of using them in the treatment of allergic diseases and/or disorders. In particular the inventive composition is useful in the treatment of saisonal allergic rhinitis and saisonal allergic conjunctivitis.

## Combinations of epinastine, Belladonna and pseudoephedrine as new pharmaceutical formulations

The present invention relates to novel oral pharmaceutical compositions comprising

5 as pharmaceutically active compounds a combination of an antihistaminic-effective amount of epinastine or a pharmaceutically acceptable salt thereof, an anticholinergic amount of Belladonna alkaloids (Belladonna) or a pharmaceutically acceptable salt thereof and of a decongestant-effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof. Optionally, the formulation may

10 comprise methylephedrine (methylephrine) or a pharmaceutically acceptable salt thereof in a decongestant-effective amount. The formulation further comprises suitable pharmaceutically acceptable carriers or excipients. Another aspect of the present invention relates to methods for the preparation of these compositions and methods of using them in the treatment of allergic diseases and/or disorders. In

15 particular the inventive composition is useful in the treatment of saisonal allergic rhinitis and saisonal allergic conjunctivitis.

### Background of the invention

Saisonal allergic rhinitis (SAR) and saisonal allergic conjunctivitis (SAC) are allergic

20 driven diseases with a specific symptomatology.

SAR is characterised by sneezing, itching, blocked nose („congested nose“) and runny nose, while SAC is characterised by eye itching, red eye and sensation of foreign body. Both allergic reactions may occur separately of each other or at the

25 same time.

An adequate systemic symptomatological treatment of SAR and SAC should address all the symptoms.

30 From the state of the art there is not known any suitable substance able to deal with all the symptoms.

2

It is known that H1 antihistamine will deal with the histamine driven symptoms such as sneezing or itching. H1 antihistamine may also have an effect on runny noses or red eyes but to a lesser grade. Due to this fact they are not the first choice substances to treat the latter. Additionally, H1 antihistamines are unable treat 5 blocked noses.

From JP-A 10298107 it is known that hyperergasis of respiratory tract secretion may be suppressed by compositions comprising H1 antihistamines and anticholinergics.

10 However, the aforementioned combination is only useful to treat some of the symptoms related to allergic disorders as SAR or SAC.

In particular these formulations or combinations are not suited to treat the issue of blocked noses or red eyes.

15

It now was found that the combination of epinastine, an H1 antihistaminic agent, Belladonna alkaloids and pseudoephedrine successfully treat all the aforementioned symptoms of SAR or SAC.

20

#### Objective of the present invention

It is one objective of the present invention to treat the symptoms of seasonal allergic rhinitis, i.e. sneezing, itching, blocked nose and runny nose.

25 It is another objective of the present invention to treat the symptoms of seasonal conjunctivitis, i.e. eye itching, red eyes and sensation of foreign bodies.

It is another objective to develop one medication to treat the symptoms of both diseases, SAR and SAC simultaneously.

30

Another objective is to develop a suitable pharmaceutical formulation for treating allergic congestion of the Eustachian tubes and / or other diseases from allergic origin deserving the administration of antihistamine and decongestant drugs.

Another objective of the present invention is the treatment of common cold and in the symptomatic relief associated with cough, cold and flu symptoms.

5 Still another objective of the present invention is to overcome the disadvantages of the medications known in the art in the treatment of SAR and/or SAC.

Among these objectives, the developments of medications to treat SAR and/or SAC are preferred.

10

#### Description of the invention

The present invention solves the problem of insufficient treatment of SAR and/or SAC by providing a pharmaceutical formulation comprising an antihistaminic-effective amount of epinastine or a pharmaceutically acceptable salt thereof, an

15 anticholinergic amount of Belladonna alkaloids or a pharmaceutically acceptable salt thereof and of a decongestant-effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof. Optionally, the formulation may additionally comprise methylephedrine or a pharmaceutically acceptable salt thereof in a decongestant-effective amount. Further ingredients of the formulation of the present

20 invention may be pharmaceutically acceptable carriers or excipients.

The term Belladonna alkaloids is commonly used in pharmaceutics. The exact method of their winning and the active ingredients of this mixture of alkaloids can be taken from the Deutsches Arzneibuch 9 (DAB 9), Volume 2, pages 932 to 944,

25 Wissenschaftliche Verlagsgesellschaft Stuttgart mbH; Govi-Verlag GmbH, Frankfurt. These pages 932 to 944 are herewith incorporated by reference. Belladonna alkaloids are won as an extract of the plant *Atropa Belladonna*, i.e. an extract of the leafs and/or the root.

The main component of the Belladonna alkaloids is atropin. Atropine itself comprises

30 L-(-)-hyoscyamine and its racemate which develops by drying. Other alkaloids found in Belladonna are L-(-)-hyoscine (L-(-)-scopolamine), N-Oxides of hyoscine and/or hyoscamine, atropamine, belladonnine and optionally nicotine, N-methylpyrrolidine, N-methylpyrrolidine, pyridin, cuskygrine and further alkaloids. The names of the

alkaloids as written above are taken from the German textbook DAB 9, referred to above. In case of ambiguities the names shall be taken directly from the textbook, page 934.

5

Preferably in the context of the present invention the mixture of the above named alkaloids are taken. However, the invention is not limited to the use of this exact mixture. In fact, any mixture or any single alkaloid of the designated alkaloids extracted from *Atropa Belladonna* can be used. In particular, the invention comprises

10 atropin or L-(-)-hyoscyamin alone without the other named alkaloids. In the context of the present invention, the term belladonna alkaloids preferably stands mainly for hyoscyamin and scopolamine as major components in extract of belladonna roots and/or leaves. These anticholinergic alkaloids have analgesic-antispasmodic action and inhibitory action of secretion. Extract of *Datura* can also be selected as a  
15 substitute for belladonna alkaloids.

Also the above mentioned active ingredients are the preferred ones and as a consequence thereof the formulation preferably does not contain any further active ingredients, the formulation of the present invention is not limited to these active  
20 ingredients alone. As an additional active compound the compositions according to the invention may optionally contain one or several compounds selected from the group consisting of mucolitic and analgesic-antipyretic compounds and vitamins. Preferred mucolitic ingredients are selected from bromhexine and ambroxol. Preferred analgesic-antipyretic compounds are selected from paracetamol and  
25 ibuprofen. Preferred vitamins are selected from vitamin B2, B6 and C. Preferably a leukotriene antagonist is not present.

In a preferred embodiment the present invention relates to an oral pharmaceutical composition. Due to the short-lasting effects of pseudoephedrine and Belladonna  
30 and - relatively to this – the long-lasting effect of epinastine it is of advantage to have a sustained release of Belladonna and the decongestant effective amount of pseudoephedrine and/or methylephrine and an immediate release of an antihistaminic effective amount of epinastine.

The preferred dosage forms are tablets or capsules.

Concerning the application via a tablet, in the context of the present the invention a

5 bilayer tablet is preferred wherein a first layer A provides for the sustained release of Belladonna and pseudoephedrine, which comprises a decongestant effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof and a anticholinergic amount of Belladonna or a pharmaceutically acceptable salt thereof.

A second layer B provides for the immediate release of epinastine and comprises an

10 antihistaminic effective amount of epinastine or a pharmaceutically acceptable salt thereof. In case the formulation contains additionally methylephedrine or one of its pharmaceutically acceptable salts, the appropriate amount thereof is present in layer A, already comprising pseudoephedrine.

15 Both layers A or B may further comprise pharmaceutically acceptable excipients and/or carriers.

The bilayer tablet according to the invention may additionally contain a tablet coating C consisting of pharmaceutically acceptable excipients, which mask the bitter taste

20 of one of the active compounds.

In a preferred embodiment of the inventive bilayer tablet layer A comprises a decongestant effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof and optionally methylephedrine or a pharmaceutical acceptable salt thereof in a matrix of a swellable hydrophilic polymer which provides a sustained release profile in a period of 3 to 24, preferably 6 to 18, most preferably about 12 hours.

In another application form the inventive composition may be formulated as a

30 capsule. Such a capsule can provide the active ingredients either instantly or some of them are provided instantly and others are provided in a sustained manner. As outlined above it is preferred to formulate the active ingredients pseudoephedrine (or its salts) and Belladonna alkaloids (or its salts thereof) as well as the optionally used

6

methylephrine (or its salts) as a sustained releases form and epinastine or its salts as immediate release form.

Preferably the capsules are made of materials that at least partially can be digested  
 5 by humans. Such capsules f.e. are disclosed in EP 0143524. The latter discloses a two-part capsule of material which is easily digestible by humans.

EP 0460921 describes capsules of chitosan and starch, grain powder, oligosaccharides, methacrylic acid-methylacrylate, methacrylic acid-ethylacrylate,  
 10 hydroxypropylmethylcelluloseacetate, -succinate or -phthaleate.

GB 938828 discloses capsules comprising water-soluble gelatine, methylcellulose, Polyvinylalkohol or water-soluble non-toxic thermoplasts.

15 EP 0 606 486 B1 discloses capsules being composed of hydroxypropylmethylcellulose, methylcellulose, hydroxypropylcellulose, starch, hydroxypropylstarch, and sodium alginate.

JP 2002-525412A discloses capsules being composed of pullulan.  
 20 Principally all theses capsules can be take for the present invention, preferred are gelatine-capsules, in particular hard-gelatine capsules. Other preferred capsules are made of starch or of a cellulose-derivative like hydroxypropylmethylcellulose and pullulan. Pullulan is a neutral simple polysaccharide produced from cultured *Aureobasidium pullulans*. It has a structure of chains of repeated  $\alpha$ -1.6 bondage of maltotriose composed of three glucoses in  $\alpha$ -1.4 bondage.  
 25 It is listed in Japanese Pharmaceutical Excipients(JPE).

Preferred standard capsules have the following physical characteristics:

Size	5	4	3	2	1	0
------	---	---	---	---	---	---

Body-Volume [ml]	0.13	0.21	0.28	0.37	0.49	0.68
------------------	------	------	------	------	------	------

Among them capsule-size of 1 or 2 are preferred.

According to the invention the term pharmaceutically acceptable salts stands for acid addition salts of the active compounds pseudoephedrine, epinastine, Belladonna alkaloids and/or methylephedrine. These acid addition salts can be formed with inorganic acids like hydrochloric acid, hydrobromic acid or sulfuric acid or with 5 organic acids as for instance oxalic acid, fumaric acid or methansulfonic acid.

Epinastine is preferably used as its hydrochloric acid addition salt. Pseudoephedrine and also methylephedrine are preferably used as the hydrochlorides or the sulfates. Within the present invention the hydrochloride-salts for the latter two compounds are most preferred.

10

The release of pseudoephedrine and optionally methylephedrine takes place over 3 to 24, preferably 6 to 24, most preferably about 12 to 24 hours. The preferred dose regimen is a „once a day application”, regardless of how the formulation is applied.

15 The concentration range of pseudoephedrine salt plus methylephedrine salt in the compositions according to the invention is between 5 and 240 mg daily, preferably 10 to 200 mg daily, more preferably 20 to 150 mg daily.

If methylephedrine or a salt thereof is present, what is preferred, both compounds

20 pseudoephedrine and methylephedrine are preferably present in the formulation in the same amount, i.e. amount w/w. Thus, to reach a total amount of pseudoephedrine plus methylephedrine (their salts respectively) of e.g. 78 mg daily each of the two compounds is present in an amount of 39 mg daily, for a total amount of 60 mg daily, each compound is present in an amount of 30 mg daily.

25

The concentration range of epinastine salt in the compositions according to the invention is between 2 and 20 mg daily, preferably 5 to 15 mg daily, more preferably 7.5 to 12.5 mg daily.

30 The concentration range of Belladonna alkaloids in the compositions according to the invention is between 0.05 and 4.0 mg daily, preferably between 0.05 and 2.0 mg daily, more preferably 0.1 to 1.5 mg daily, most preferably between 0.2 and 0.6 mg daily.

In case of a bilayer tablet, each layer is in contact with each other in a portion of their surface, but provides independent release profiles for both active substances mentioned before.

5 The sustained release layer A comprises beside the active ingredient(s) a swellable hydrophilic polymer.

Typical swellable hydrophilic polymers include cellulose ethers such as methylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, carboxymethylcellulose and

10 carboxyethylcellulose or mixtures thereof. The use of hydroxypropylmethylcellulose (HPMC) is preferred. Particularly useful are the HPMC polymers HPMC USP2910 and USP2208 like for instance Methocel E5. E4M, E15M, K15M, and K100M supplied by the Dow Chemical Company. In the aforementioned abbreviations the designation "E" refers to USP2910 whereas "K" refers to USP2208. The number 15 designation refers to the viscosity in a 2% aqueous solution (e.g. 5 designates a viscosity of 5 cps; 15M designates a viscosity of 15000 cps).

The excipients that could be optionally used in the sustained release layer A are insoluble polymers, soluble or insoluble fillers, antiadherents, coloring agents,

20 lubricants and additional binders. Typical fillers are for example lactose, microcrystalline cellulose, dibasic calcium phosphate and cornstarch. Examples of antiadherents, which are used to prevent tablets from sticking to the tablet press, are colloidal silicon dioxide and talc. Magnesium stearate, talc and stearic acid are typical lubricants. Typical binders are povidone, and cornstarch.

25

The immediate release matrix layer B comprises beside the active ingredient different combinations of excipients. The excipients that could be optionally used in the immediate release layer B are insoluble polymers, soluble or insoluble fillers, antiadherents, lubricants, coloring agents, disintegrants and additional binders.

30 Typical fillers are for example lactose, microcrystalline cellulose, dibasic calcium phosphate and cornstarch. Examples of antiadherents, which are used to prevent tablets from sticking to the tablet press, are colloidal silicon dioxide and talc. Typical disintegrants are crospovidone, sodium starch glycolate and crosscarmellose

sodium. Typical coloring agents are selected from FD&C red 40 HT Aluminum lake, 2-hydroxy-1.1'-azonaphthalene-3.6.4'-trisulfonic acid trisodium salt, erythrosine, iron oxides, 1-(4-sulpho-1-naphthylazo)-2-naphthol-6.8-disulphonic acid trisodium salt, 2',4',5',7'-tetrabromo-4.5.6.7-tetrachloro-fluorescein disodium salt, 2.4.5.7-Tetraiodo-5 3.6-dihydroxyxanthene-9-spiro-1'-(4',5',6',7'-tetrachloro-3'H-isobenzofuran-3'one dipotassium salt, trisodium 3-carboxy-5-hydroxy-1-p-sulphophenyl-4-p-sulfophenylazopyrazole, 6-hydroxy-5-((4-sulphonphenyl)azo-2-naphthalenesulphonic acid disodium salt and optionally aluminium lakes thereof. Magnesium stearate, talc and stearic acid are typical lubricants. Typical binders are povidone, and cornstarch.

10

Water and ethanol are examples of volatile components which can be used in the manufacture process of both layers to granulate powders. These volatile components are removed during processing and therefore do not appear in the finished product.

15

The tablet coating is optional since the presence of it does not modify significantly the release rates of the active substances present in the core layers. The presence of the coating is preferred because it masks the bitter taste of one of the active substances and enhances the properties of dosage form. Because of that a lot 20 different coatings with different polymers, and plasticizers and other excipients could be used with the condition of not modifying significantly the release profile of the active substances present in the core tablet. A typical coating comprises a polymer such as hydroxypropylmethylcellulose and a plasticizer such as polyethylene glycol. Optional excipients could be added to the coating like antifoaming agents and 25 opacifying agents. Example of an antifoaming agent is silicone. Examples of opacifying agents are Titanium dioxide, talc and aluminum lake dyes.

The inventive formulation also can be applied via a tablet comprising sustained release and non-sustained release granules or a capsule comprising the same.

30

In case of such a tablet, non-sustained release granules and sustained release granules, which are coated with a sustained release film are mixed with suitable excipients and then they are compressed as a tablet.

Similarly non-sustained release granules and sustained release granules which are coated with sustained release film are mixed 1:9 to 9:1, preferably 3:7 to 7:3 are filled into a capsule or are compressed into tablet.

5

A non-sustained release granule comprises an amount of epinastine or a pharmaceutically acceptable salt thereof. Optionally it may comprises a portion of the total amount of belladonna alkaloids or a pharmaceutically acceptable salt thereof and a portion of the total amount of pseudoephedrine or a pharmaceutically acceptable salt thereof and optionally a portion of the total amount of 10 methylephedrine or a pharmaceutically acceptable salt thereof, if necessary.

A sustained release granule comprises either a portion or the total amount of belladonna alkaloids or a pharmaceutically acceptable salt thereof, pseudoephedrine 15 or a pharmaceutically acceptable salt thereof and optionally methylephedrine or a pharmaceutically acceptable salt thereof.

Preferably the non-sustained release granules contain only epinastine or a pharmaceutically acceptable salt thereof as active ingredient while the sustained 20 release granules comprise the remaining active ingredients.

Any compounds conventionally used as a sustained-release coat can be used for the purpose of this invention. Specific examples which can be given include water insoluble polymers such as ethylcellulose, aminoalkyl methacrylate copolymer 25 polyvinyl acetate, polyvinyl chloride, polyethylene, and the like; intestinally soluble polymers such as cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, carboxymethylcellulose, styrene acrylic acid copolymer, methacrylic acid copolymer, maleic anhydrous acid copolymer, shellac, and the like; paraffin waxes 30 such as paraffin, microcrystalline wax, and the like; higher alcohols, preferably saturated and unsaturated C<sub>6</sub> – C<sub>26</sub>-alcohols, preferred unbranched and unsubstituted, such as stearyl alcohol, cetyl alcohol, and the like; esters of higher fatty acids, preferably saturated and unsaturated C<sub>6</sub> – C<sub>26</sub>-acids, preferred

## 11

unbranched and unsubstituted, such as glycerol esters of fatty acids, hydrogenated oils, carnauba wax, beeswax, Japan (haze) wax, and the like; and higher fatty acids as defined above such as stearic acid, palmitic acid, myristic acid, behenic acid, and the like (or the sodium, calcium or magnesium salts of these higher fatty acids).

5 These excipients may be used solely or mixed. The coating amount is preferably 10% to 50% for granules.

Furthermore, the excipients that could be optionally used in sustained release film are water soluble polymers, sugar alcohols, plasticizers, titanium oxide, talc, coloring agents and so on. Typical water soluble polymers and sugar alcohols are

10 hydroxypropyl methylcellulose, hydroxypropylcellulose, methylcellulose, polyvinylpyrrolidone, polyethylene glycol. Typical plasticizers are glycerine fatty acid ester, triethyl citrate, propylene glycol, triacetin.

For any of the inventive application forms, bilayer tablet, tablet or capsule any of the  
15 aforesaid ingredients can be taken, if appropriate.

In the context of the present invention capsules and tablets comprising sustained release and non-sustained release granules are preferred.

20 The invention will be further described by the following examples. These examples disclose certain preferred embodiments of the invention. The methods of manufacturing the compositions according to the invention like for instance granulation, tablet compression, tablet-coating etc. are well known to the person skilled in the art. Those skilled in the art will appreciate that various changes,  
25 modifications and substitutions can be made therein without departing from the spirit of the invention. Accordingly, it is intended that the invention be not limited to the following explicitly disclosed examples.

Example N°1 - CompositionCore**A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K 15 M PRCR *	198
Lactose Monohydrate	104.8
Microcrystalline cellulose	106
Colloidal silicon dioxide	1.65
Magnesium Stearate	2.75
Povidone	16.5
<b>Total first layer</b>	<b>550</b>

5 **B. Second layer**

<u>Layer Epinastine</u>	<b>mg per 2 tablets (daily)</b>
Epinastine HCl	10
FD&C red 40 HT Aluminum lake (allura red AC)	0.38
Microcrystalline cellulose	70
Lactose Monohydrate	154.62
Povidone	12.5
Magnesium Stearate	2.5
<b>Total second layer</b>	<b>250</b>

<b>Total core</b>	<b>800</b>
-------------------	------------



**C. Coating**

<b>Film Coating</b>	<b>mg per 2 tablets (daily)</b>
Methocel E5	15
Polyethylene Glycol 6000	1.97
Silicone antifoam S184	0.03
Total film coating	1.7
<b>Total Film coated tablet</b>	<b>817</b>

\* PR means Premium grade and CR means Controlled Released grade.

5

Method of ManufactureA. First layer:

- A1. Dissolve povidone in a hydroalcoholic mixture;
- A2. Blend pseudoephedrine hydrochloride, methylephedrine hydrochloride, Belladonna alkaloids, a portion of the microcrystalline cellulose, lactose and Methocel K15M for 5-30 minutes in a suitable mixer.
- A3. Use alcoholic or hydroalcoholic solution prepared previously in step A1 to granulate the powder mix of step A2.
- A4. Dry and mill the granulation from step A3, using suitable size screen.
- A5. Blend the screened granulation with a portion of the microcrystalline cellulose and colloidal silicon dioxide for 3-15 minutes.
- A6. Add magnesium stearate and blend for 3-15 minutes.

B Second layer:

- B1. Pass through a suitable screen Epinastine HCl, Allura red AC (FD & C red 40 HT) aluminum lake and microcrystalline cellulose. Blend for 5-30 minutes in a suitable mixer.
- B2. Add lactose and povidone. Blend for 60 minutes 15-120 minutes in a suitable mixer.
- B3. Add magnesium stearate. Blend for 3-20 minutes in a suitable mixer.

C. Compression:

Compress A and B into a suitable bilayer tableting machine in suitable size tablets.

5 D. Coating

D1. Dissolve Methocel E5 and Polyethylene Glycol in suitable amount of water.

D2. Dissolve silicone antifoam in suitable amount of isopropilic alcohol.

D3. Add D2. to D1. and mix.

D4. Coat tablets with the Methocel E5 /Polyethylene glycol solution from step D3. in  
10 a suitable coater.

Example N° 2 - CompositionCore**A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<u>mg per 2 tablets (daily)</u>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K 15 M PRCR *	198
Lactose Monohydrate	126.2
Microcrystalline cellulose	100
Colloidal silicon dioxide	2.75
Magnesium Stearate	2.75
<b>Total first layer</b>	<b>450</b>

15

**B. Second layer**

<u>Layer Epinastine</u>	<u>mg per 2 tablets (daily)</u>

16

Epinastine HCl	10
Lactose Monohydrate	168.4
Microcrystalline cellulose	70
Punceau 4R red aluminum lake	0.35
Magnesium Stearate	1.25
Total second layer	250

<b>Total core</b>	<b>700</b>
-------------------	------------

### C. Coating

<b>Film Coating</b>	<b>mg per 2 tablets (daily)</b>
Methocel E5	4.42
Polyethylene Glycol 6000	2.72
Talc	8.76
Titanium dioxide	1.1
Total film coating	17

<b>Total Film coated tablet</b>	<b>817</b>
---------------------------------	------------

5

\* PR means Premium grade and CR means Controlled Released grade.

### Method of Manufacture

#### A. First layer:

10 A1. Blend pseudoephedrine hydrochloride, methylephedrine hydrochloride, Belladonna alkaloids, microcrystalline cellulose, lactose, colloidal silicon dioxide and HPMC K15M for 5-30 minutes in a suitable mixer.

A2. Add magnesium stearate and blend for 3-15 minutes.

15 B. Second layer:

17

- B1. Pass through a suitable screen Epinastine HCl, and microcrystalline cellulose. Blend for 5-30 minutes in a suitable mixer.
- B2. Add lactose. Blend for 60 minutes 15-120 minutes in a suitable mixer.
- B3. Add magnesium stearate. Blend for 3-20 minutes in a suitable mixer.

5

#### C. Compression:

Compress A and B into a suitable bilayer tableting machine in suitable size tablets.

#### D. Coating

- D1. Dissolve Methocel E5 and Polyethylene Glycol in suitable amount of water.
- D2. Add Titanium Dioxide and Talc in suitable amount of water and mix
- D3. Add D2. to D1. And mix.
- D4. Coat tablets with the Methocel E5 /Polyethylene glycol solution from step D3. in a suitable coater.

15

#### **Example N° 3**

##### Core

##### **A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K4M PRCR	247.5
Lactose Monohydrate	165.7
Talc	11
Magnesium Stearate	5.5
<b>Total first layer</b>	<b>550</b>

20

\* PR means Premium grade and CR means Controlled Released grade.

Second layer and coating are identical to example 2; the manufacture method was conducted analogously to the method outlined in example 2;

**Example N° 4**Core**A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15 M PRCR	198
Lactose Monohydrate	99.2
Microcrystalline cellulose	99.5
Colloidal silicon dioxide	2.75
Povidone	27.5
Magnesium stearate	2.75
<b>Total</b>	<b>550</b>

\* PR means Premium grade and CR means Controlled Released grade.

5

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 5**10 Core**A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	330
Lactose	83.2

20

Talc	11
Magnesium Stearate	5.5
<b>Total</b>	<b>550</b>

\* CR means Controlled Released grade.

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

5

### Example N° 6

#### Core

#### A. First layer

<u>Layer pseudoephedrine, Belladonna,</u> <u>Methylephedrine</u>	mg per 2 tablets (daily)
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	275
Microcrystalline Cellulose	138.2
Talc	11
Magnesium Stearate	5.5
Ethanol	sq.
<b>Total</b>	<b>550</b>

\* CR means Controlled Released grade.

10

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 7**Core**A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	215
Dibasic Calcium phosphate	108.2
Ethylcelullose	40
Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>500</b>

\* CR means Controlled Released grade.

5

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 8**Core**A. First layer**

<u>Layer pseudoephedrine, Belladonna,</u> <u>Methylephedrine</u>	mg per 2 tablets (daily)
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	137.5
Methocel K100M CR	137.5
Lactose	138.2
Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>550</b>

\* CR means Controlled Released grade.

5

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 9**10 Core**A. First layer**

<u>Layer pseudoephedrine, Belladonna,</u> <u>Methylephedrine</u>	mg per 2 tablets (daily)
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K100M CR	275
Lactose	138.2

23

Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>550</b>

\* CR means Controlled Released grade.

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

5

### Example N° 10

#### Core

#### A. First layer

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<u>mg per 2 tablets (daily)</u>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	206.2
Methocel K100M CR	68.8
Lactose	138.2
Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>550</b>

\* CR means Controlled Released grade.

10

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 11**Core**A. First layer**

<u>Layer pseudoephedrine, Belladonna, Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	235
Dibasic Calcium phosphate	108.2
Ethylcellulose	20
Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>500</b>

\* CR means Controlled Released grade.

5

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 12****Core****A. First layer**

<u>Layer pseudoephedrine, Belladonna,</u> <u>Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	255
Lactose	39.7
Microcrystalline Cellulose	68.5
Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>500</b>

\* CR means Controlled Released grade.

5

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

**Example N° 13**

10 **Core**

**A. First layer**

<u>Layer pseudoephedrine, Belladonna,</u> <u>Methylephedrine</u>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	60
Methylephedrine hydrochloride	60
Belladonna	0.3
Methocel K15M CR	255
Dibasic calcium phosphate	108.2

Talc	11
Magnesium Stearate	5.5
Ethanol	s.q.
<b>Total</b>	<b>500</b>

\* CR means Controlled Released grade.

Second layer and coating are identical to example 1; the manufacture method was conducted analogously to the method outlined in example 1;

5

#### Example N° 14

##### a) Non-sustained release granules: 2 capsules

<b>Non-sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Hydroxypropylcellulose	3.5
Sucrose	475.35
<b>Total</b>	<b>549</b>

10

##### b) Sustained release granules: 2 capsules

<b>Sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Hydroxypropylcellulose	4
Sucrose	90.85
Methacrylic acid copolymer, type B	40.6
Glycerol esters of fatty acids	3.1
Talc	1.3

<b>Total</b>	<b>200 mg</b>
--------------	---------------

### Capsulation

non-sustained relaes granules	549
sustained relaes granules	200
Talc	1
<b>Total</b>	<b>750</b>

### Method of Manufacture

5    A. Non-sustained release granules

    A1. Dissolve hydroxypropylcellulose in ethanol.

    A2. Blend epinastine hydrochloride and pseudoephedrine hydrochloride, methylephedrine hydrochloride and belladonna in a suitable mixer and pulverize the powder mix.

10    A3. Produce spherical granules by spraying the solution prepared previously in step A1 over sucrose introducing the powder mix obtained from step A2.

    A4. Dry and pass through granules from step A3 with suitable size screen to produce non-sustained release granules.

15    B. Sustained release granules

    B1. Dissolve hydroxypropylcellulose in ethanol.

    B2. Blend pseudoephedrine hydrochloride, methylephedrine hydrochloride and belladonna in a suitable mixer.

    B3. Produce spherical granules by spraying the solution prepared previously in step B1 over sucrose introducing the powder mix obtained from step B2.

20    B4. Dry and pass through granules from step B3 with suitable size screen

    B5. Dissolve Methacrylic acid copolymer, type B in ethanol and mix with glycerol esters of fatty acids and talc.

    B6. Coat the granules obtained from step B4 with the solution prepared previously in step B5 to produce sustained release granules.

25    C. Capsulation

    C1. Mix non-sustained release granules and sustained release granules with talc.

C2. Fill the mixture obtained from step C1 into capsules

**Example N° 15**

**a) Non-sustained release granules: 2 capsules**

<b>Non-sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Hydroxypropylcellulose	3.5
Sucrose	475.35
<b>Total</b>	<b>549</b>

5

**b) Sustained release granules: 2 capsules**

<b>Sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Hydroxypropylcellulose	4
Sucrose	90.85
Ethylcellulose	38.75
Hydroxypropylmethycellulose 2910	1
Glycerol esters of fatty acids	2.25
Talc	3
<b>Total</b>	<b>200</b>

**Capsulation**

non-sustained relaes granules	549
sustained relaes granules	200
Talc	1

<b>Total</b>	750
--------------	-----

The manufacture method was conducted analogously to the method outlined in example 14.

## 5 Example N° 16

### a) Non-sustained release granules: 2 capsules

<b>Non-sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	310.1
<b>Total</b>	<b>380.25</b>

### b) Sustained release granules: 2 capsules

<b>Sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	114.85
Ammonio methacrylate copolymer	31.5
Ethylcellulose	7.875
Glycerol esters of fatty acids	1.641
Talc	2.734
<b>Total</b>	<b>218.75</b>

### Capsulation

non-sustained relaes granules	380.25
sustained relaes granules	218.75

30

Talc	1
<b>Total</b>	<b>600</b>

The manufacture method was conducted analogously to the method outlined in example 14.

## 5 Example N° 17

### a) Non-sustained release granules: 2 capsules

<b>Non-sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	310.1
<b>Total</b>	<b>380.25</b>

### b) Sustained release granules: 2 capsules

<b>Sustained release granules</b>	<b>mg per 2 capsules (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	114.86
Ammonio methacrylate copolymer	35.83
Ethylcellulose	3.54
Glycerol esters of fatty acids	1.64
Talc	2.73
<b>Total</b>	<b>218.75</b>

31

non-sustained relaes granules	380.25
sustained relaes granules	218.75
Talc	1
<b>Total</b>	<b>600</b>

The manufacture method was conducted analogously to the method outlined in example 14.

5

### Example N° 18 Tablet

#### a) Non-sustained release granules

<b>Non-sustained release granules</b>	<b>mg per 2 tablets (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Hydroxypropylcellulose	12.5
Microcrystalline cellulose	154.85
Lactose	12.5
<b>Total</b>	<b>250</b>

10

#### b) Sustained release granules

<b>Sustained release granules</b>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Hydroxypropylcellulose	4
Sucrose	90.85

Methacrylic acid copolymer, type B	30.45
Magnesium stearate	10.15
Glycerol esters of fatty acids	3.1
Talc	1.3
<b>Total</b>	<b>200</b>

### Compression

non-sustained release granules	250
sustained release granules	200.
Microcrystalline cellulose	126
Croscarmellose sodium	12
Talc	6
Magnesium stearate	6
<b>Total</b>	<b>600</b>

### Method of Manufacture

#### 5 A. Non-sustained release granules

- A1. Dissolve hydroxypropylcellulose in ethanol.
- A2. Blend epinastine hydrochloride and pseudoephedrine hydrochloride, methylephedrine hydrochloride, belladonna, microcrystalline cellulose and lactose in a suitable mixer and knead the mixture with the solution from step 10 A1.
- A3. Dry and pass through granules obtained from step A2 with suitable size screen to produce non-sustained release granules

#### B. Sustained release granules

- B1. Dissolve hydroxypropylcellulose in ethanol.
- B2. Blend pseudoephedrine hydrochloride, methylephedrine hydrochloride and belladonna in a suitable mixer.
- B3. Produce spherical granules by spraying the solution prepared previously in step B1 over sucrose introducing the powder mix obtained from step B2.
- B4. Dry and pass through granules from step B3 with suitable size screen

33

B5. Dissolve Methacrylic acid copolymer, type B in ethanol and mix with glycerol esters of fatty acids, magnesium stearate and talc.

B6. Coat the granules obtained from step B4 with the solution prepared previously in step B5 to produce sustained release granules.

5

C. Compression:

C1. Mix non-sustained release granules and sustained release granules with microcrystalline cellulose, croscarmellose sodium, talc and magnesium stearate.

10 C2. Compress the mixture into a suitable tabletting machine in suitable size tablets.

**Example N° 19 Tablet**

**a) Non-sustained release granules:**

<b>Non-sustained release granules</b>	<b>mg per 2 tablets (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	161.1
<b>Total</b>	<b>231.25</b>

15

**b) Sustained release granules:**

<b>Sustained release granules</b>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	114.85
Ammonio methacrylate copolymer	31.5
Ethylcellulose	7.875
Glycerol esters of fatty acids	1.641
Talc	2.734

<b>Total</b>	<b>218.75</b>
--------------	---------------

### Compression

non-sustained relaes granules	231.25
sustained relaes granules	218.75
Microcrysatalline cellulose	126
Croscarmellose sodium	12
Talc	6
Magnesium stearate	6
<b>Total</b>	<b>600</b>

The manufacture method was conducted analogously to the method outlined in  
 5 example 18.

### Example N° 20 Tablet

#### a) Non-sustained release granules:

<b>Non-sustained release granules</b>	<b>mg per 2 tablets (daily)</b>
Epinastine hydrochloride	10
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	161.1
<b>Total</b>	<b>231.25</b>

10

#### b) Sustained release granules:

<b>Sustained release granules</b>	<b>mg per 2 tablets (daily)</b>
Pseudoephedrine hydrochloride	30
Methylephedrine hydrochloride	30
Belladonna	0.15
Sucrose	114.86
Ammonio methacrylate copolymer	35.83

Ethylcellulose	3.54
Glycerol esters of fatty acids	1.64
Talc	2.73
<b>Total</b>	<b>218.75</b>

### Compression

non-sustained relaes granules	231.25
sustained relaes granules	218.75
Microcrysataline cellulose	126
Coscarmellose sodium	12
Talc	6
Magnesium stearate	6
<b>Total</b>	<b>600</b>

The manufacture method was conducted analogously to the method outlined in  
5 example 18.

### Examples No° 20 to 40:

10 The same as examples No° 1 to 20 but the methylephrine is displaced by the same amount of pseudoephedrine, i.e. the amount of pseudoephedrine is doubled.

Claims

- 1) Oral pharmaceutical compositions comprising as pharmaceutically active compounds a combination of a) an antihistaminic-effective amount of epinastine or a pharmaceutically acceptable salt thereof, b) an anticholinergic amount of Belladonna alkaloids or a pharmaceutically acceptable salt thereof and c) a decongestant-effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof optionally in combination with methylephrine or a pharmaceutically acceptable salt thereof and further comprising pharmaceutically acceptable carriers or excipients.
- 2) Oral pharmaceutical composition according to claim 1, characterised in that the Belladonna alkaloids comprise at least one alkaloid of the group being selected of atropin, L-(-)-hyoscyamin, L-(-)-hyoscin, N-Oxides of hyoscin and/or hyoscamin, atropamin, belladonnin and optionally nicotin, N-methylpyrrolin, N-methylpyrrolidin, pyridin, cuskygrin.
- 3) Oral pharmaceutical composition according to claim 1 or 2, characterised in that all active ingredients are formulated for instant release.
- 4) Oral pharmaceutical composition according to any of claims 1 to 3, characterised in that epinastine or a pharmaceutically acceptable salt thereof is formulated for instant release and at least a portion of the other active ingredients, i.e. Belladonna alkaloids or a pharmaceutically acceptable salt thereof, pseudoephedrine or a pharmaceutically acceptable salt thereof and optionally methylephrine, are formulated for sustained release.
- 5) Oral pharmaceutical composition according to claim 4, characterised in that the total amounts of Belladonna alkaloids or a pharmaceutically acceptable salt thereof, pseudoephedrine or a pharmaceutically acceptable salt thereof and optionally methylephrine, are formulated for sustained release.

- 6) Oral pharmaceutical composition according to any of claims 1 to 5, characterised in that the formulation contains methylephrine or a pharmaceutically acceptable salt thereof.
- 5 7) Oral pharmaceutical composition according to any of claims 1 to 6, characterised in that the concentration range of epinastine or a pharmaceutically acceptable salt thereof is between 2 mg and 20 mg daily, the concentration range of the Belladonna alkaloids or salt pharmaceutically acceptable salt thereof is between 0.05 and 4.0 mg daily and the concentration 10 range of pseudoephedrine plus methylephrine or the corresponding pharmaceutically acceptable salts thereof is between 5 and 240 mg daily.
- 15 8) Oral pharmaceutical composition according to claim 7, characterised in that the concentrations of pseudoephedrine and methylephrine are of the same amount.
- 9) Oral pharmaceutical composition according to any of claims 1 to 8, characterised in that it represents a bilayer tablet.
- 20 10) Bilayer tablet according to claim 9, wherein a first layer A, providing for the sustained release of Belladonna alkaloids, pseudoephedrine and optionally methylephrine or the corresponding pharmaceutical salts of the named active ingredients, comprises the amounts of these compounds according to claim 1 and wherein a second layer B, providing for the immediate release of epinastine, comprises an antihistaminic effective amount of epinastine or a 25 pharmaceutically acceptable salt thereof.
- 30 11) Bilayer tablet according to one of claims 9 or 10, characterised in that layer A comprises 60 mg pseudoephedrine hydrochloride, 60 mg methylephrine-hydrochloride and 0.3 mg Belladonna alkaloids and layer B comprises 10 mg epinastine-HCl or a tablet in that any of the amounts are divided by 2.

- 12) Bilayer tablet according to any of claims 9 to 11, characterised in that it additionally contains a tablet coating C consisting of pharmaceutically acceptable excipients.
- 5 13) Bilayer tablet according to any of claims 9 to 12, characterised in that layer A comprises a decongestant effective amount of pseudoephedrine or a pharmaceutically acceptable salt thereof and methylephrine or a pharmaceutically acceptable salt thereof and an anticholinergic amount of the Belladonna alkaloids or a salt thereof in a matrix of a swellable hydrophilic 10 polymer.
- 14) Capsule comprising an oral formulation according to one of claims 1 to 9.
- 15) Capsule according to claim 14, characterised in that the material of the capsules comprises a compound being selected from the group of chitosan and starch, grain powder, oligosaccharides, methacrylic acid-methylacrylate, methacrylic acid-ethylacrylate, hydroxypropylmethylcelluloseacetate, -succinate or -phthaleate, polyvinylalcohol, water-soluble non-toxic thermoplasts, hydroxypropylmethyl-cellulose, methylcellulose, hydroxypropylcellulose, 20 hydroxypropylstarch, sodium alginate, gelatine, hard-gelatine and pullulan.
- 16) Capsule according to claim 14 or 15, characterised in that the ingredients are formulated as sustained release and non-sustained release granules.
- 25 17) Capsule according to claim 16 in combination with claim 4 or 5.
- 18) Capsule according to claim 14, 15 or 16, characterised in that the non-sustained granules are coated by a water insoluble polymers, intestinally soluble polymers, paraffin waxes, higher alcohols, higher fatty acids and/or 30 higher fatty acid esters.

- 19) Tablet comprising an oral formulation according to one of claims 1 to 9, characterised in that the ingredients are formulated as granules which are compressed to a tablet.
- 5 20) Tablet according claim 19, characterised in that the the ingredients are formulated as sustained release and non-sustained release granules.
- 21) Tablet according to claim 20 in combination with claim 4 or 5.
- 10 22) Tablet according to claim 19, 20 or 21, characterised in that that the non-sustained granules are coated by a water insoluble polymers, intestinally soluble polymers, paraffin waxes, higher alcohols, higher fatty acids and/or higher fatty acid esters.
- 15 23) Use of a pharmaceutical composition according to one of claims 1 to 9, a bilayer tablet according to any of claims 10 to 13, a capsule according to any of claims 14 to 18 or a tablet accroding to any of claims 19 to 21 for the treatment of saisonal allergic rhinitis, saisonal allergic conjunctivitis, allergic rhinitis, allergic congestion of the Eustachian tubes and / or other diseases from allergic origin deserving the administration of antihistamine and decongestant drugs, in the treatment of for instance common cold and in the symptomatic relief associated with cough, cold and flu symptoms.
- 20 24) Use according claim 22 for the treatment of seasonal allergic rhinitis and/or seasonal allergic conjunctivitis.
- 25