



(86) Date de dépôt PCT/PCT Filing Date: 2008/08/13

(87) Date publication PCT/PCT Publication Date: 2009/02/19

(45) Date de délivrance/Issue Date: 2015/11/24

(85) Entrée phase nationale/National Entry: 2010/02/10

(86) N° demande PCT/PCT Application No.: JP 2008/064520

(87) N° publication PCT/PCT Publication No.: 2009/022702

(30) Priorité/Priority: 2007/08/13 (JP2007-210804)

(51) Cl.Int./Int.Cl. *C07D 493/04* (2006.01)

(72) Inventeurs/Inventors:

GOTO, KIMIHIKO, JP;
YAMAMOTO, KAZUMI, JP;
SAKAI, MASAYO, JP;
MITOMI, MASAOKI, JP;
ANDO, TAKASHI, JP


(73) Propriétaire/Owner:

BASF SE, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PROCEDE POUR PRODUIRE DES DERIVES DE PYRIPYROPENE ET DES PRODUITS INTERMEDIAIRES
DE PRODUCTION DE CELUI-CI

(54) Title: PROCESS FOR PRODUCING PYRIPYROPENE DERIVATIVES AND INTERMEDIATES FOR THE
PRODUCTION THEREOF



(57) **Abrégé/Abstract:**


Disclosed is a process for producing compound C represented by formula C: [Chemical formula 1] (see formula C) wherein R' represents straight chain, branched chain, or cyclic C₂₋₆ alkylcarbonyl, wherein R_{1b} is used as a protective group for hydroxyl at the 7-position of compound C. R_{1b} represents formyl; optionally substituted straight chain C₁₋₄ alkylcarbonyl; optionally substituted benzyl; group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; C₁₋₆ alkyloxy-C₁₋₆ alkyl optionally substituted by halogen atom; C₁₋₆ alkylthio-C₁₋₆ alkyl optionally substituted by halogen atom; straight chain, branched chain, or cyclic C₁₋₄ alkyl optionally substituted by halogen atom, provided that, when alkyl in the C₁₋₄ alkyl group is of a branched chain or cyclic type, the alkyl group is C₃₋₄ alkyl; C₂₋₆ alkenyl optionally substituted by halogen atom; C₂₋₆ alkynyl optionally substituted by halogen atom; or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group. The process can produce pyripyropene derivatives that have acyloxy groups at the 1- and 11-positions and a hydroxyl group at the 7-position and are useful as insect pest control agents at a high yield.



ABSTRACT

Disclosed is a process for producing compound C represented by formula C:

5 [Chemical formula 1]



wherein R' represents straight chain, branched chain, or cyclic C₂₋₆ alkylcarbonyl, wherein R_{1b} is used as a protective group for hydroxyl at the 7-position of compound C. R_{1b} represents

10 formyl; optionally substituted straight chain C₁₋₄ alkylcarbonyl; optionally substituted benzyl; group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; C₁₋₆ alkyloxy-C₁₋₆ alkyl optionally substituted by

15 halogen atom; C₁₋₆ alkylthio-C₁₋₆ alkyl optionally substituted by halogen atom; straight chain, branched chain, or cyclic C₁₋₄ alkyl optionally substituted by halogen atom, provided that, when alkyl in the C₁₋₄ alkyl group is of a branched chain or cyclic type, the alkyl group is C₃₋₄ alkyl; C₂₋₆ alkenyl optionally

20 substituted by halogen atom; C₂₋₆ alkynyl optionally substituted by halogen atom; or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group. The process can produce pyripyropene derivatives that have acyloxy groups at the 1- and 11-positions and a hydroxyl group at the

25 7-position and are useful as insect pest control agents at a high yield.

20375-1007

1

SPECIFICATION

PROCESS FOR PRODUCING PYRIPYROPENE DERIVATIVES AND
INTERMEDIATES FOR THE PRODUCTION THEREOF

5

[0001]

BACKGROUND OF THE INVENTION

15

[0002]

Field of Invention

20

The present invention relates to a process for producing pyripyropene derivatives useful as pest control agents and more specifically relates to a process for producing pyripyropene derivatives that have acyloxy groups at the 1- and 11-positions and a hydroxyl group at the 7-position thereof.

[0003]

Background Art

25

Pyripyropene derivatives having acyloxy at 1- and 11-positions and hydroxyl at the 7-position thereof are compounds that have control effects against pests, as described in WO 2006/129714.

[0004]

30

WO 2006/129714 and Japanese Patent Application Laid-Open No. 259569/1996 disclose a process for producing pyripyropene derivatives having acyloxy at 1- and 11-positions and hydroxyl at the 7-position thereof. According to the production process, the pyripyropene derivatives are purified or isolated from a plurality of products produced by nonselective hydrolysis of acyloxy using a 1,7,11-triacyloxy compound as a

35

starting compound. This production process, however, suffers

20375-1007

2

from problems such as low yield and unsuitability for quantity synthesis.

[0005]

Further, Japanese Patent Application Laid-Open No. 259569/1996 describes the use of a combination of protective groups for the synthesis of pyripyropene derivatives, and Journal of Antibiotics Vol. 49, No. 11, p. 1149, 1996, Bioorganic Medicinal Chemistry Letter Vol. 5, No. 22, p. 2683, 1995, Japanese Patent Application Laid-Open No. 269065/1996, and WO 2008/013336 disclose a synthesis example that introduces acyl into the 7-position by utilizing a protective group. These documents, however, do not disclose a specific process that utilizes a protective group in the production of pyripyropene derivatives that have acyloxy at the 1- and 11-positions and hydroxyl at the 7-position thereof.

SUMMARY OF THE INVENTION

[0006]

The present inventors have found that pyripyropene derivatives having acyloxy groups at the 1- and 11-positions and a hydroxyl group at the 7-position thereof can be produced at high yield from pyripyropene A (Japanese Patent Application Laid-Open No. 259569/1996; Bioorganic Medicinal Chemistry Letter Vol. 5, No. 22, p. 2683, 1995; and WO2004/060065) which is obtained as a naturally occurring substance, by using a proper protective group. The present invention has been made based on such finding.


[0007]

The present invention relates to a process for producing pyripyropene derivatives useful as pest control agents and to provide compounds as intermediates for the production of pyripyropene derivatives.

[0008]

According to a first aspect of the present invention, there is provided a process for producing compound C represented by formula C:


[Chemical formula 1]



wherein R' represents straight chain, branched chain, or
 5 cyclic C₂₋₆ alkylcarbonyl, provided that, when the alkyl moiety in
 the alkylcarbonyl group is of a branched chain or cyclic type, R'
 represents C₃₋₆ alkylcarbonyl, the process comprising the steps
 of:


(a₁) hydrolyzing acetyl at the 7-position of compound A1
 10 represented by formula A1:

[Chemical formula 2]



wherein Ac represents acetyl,
 15 with a base to selectively deacylate compound A1, then
 protecting hydroxyl at the 7-position to give compound B1
 represented by formula B1:

[Chemical formula 3]



B1

wherein

Ac is as defined above,


R_{1a} represents optionally substituted straight chain C_{2-4} alkylcarbonyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; C_{1-6} alkyloxy- C_{1-6} alkyl optionally substituted by halogen atom; C_{1-6} alkylthio- C_{1-6} alkyl optionally substituted by halogen atom; straight chain, branched chain, or cyclic C_{1-4} alkyl optionally substituted by halogen atom, provided that, when alkyl in the C_{1-4} alkyl group is of a branched chain or cyclic type, the alkyl group represents C_{3-4} alkyl; C_{2-6} alkenyl optionally substituted by halogen atom; C_{2-6} alkynyl optionally substituted by halogen atom; optionally substituted benzyl; or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group,

wherein, in R_{1a} , the substituent optionally possessed by alkylcarbonyl is selected from the group consisting of halogen atoms, C_{1-4} alkyloxy, C_{1-4} haloalkyloxy, C_{1-4} alkylcarbonyl, C_{1-4} haloalkylcarbonyl, C_{1-4} alkylcarbonyloxy, and C_{1-4} haloalkylcarbonyloxy, and the substituent optionally possessed by the heterocyclic group and benzyl is selected from the group consisting of halogen atoms, C_{1-4} alkyl, C_{1-4} alkyloxy, C_{1-4} haloalkyloxy, C_{1-4} alkylthio, C_{1-4} haloalkyl, C_{1-4} alkylcarbonyl, C_{1-4} haloalkylcarbonyl, C_{1-4} alkylcarbonyloxy, C_{1-4} haloalkylcarbonyloxy, nitro, and cyano,

then hydrolyzing acetyl at the 1- and 11-positions of

compound B1 with a base to deacylate compound B1 and thus to give compound Fa represented by formula Fa:

[Chemical formula 4]




Fa

5

wherein R_{1a} is as defined above, or
(a₂) hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A1 or compound A4' represented by formula A4':

[Chemical formula 5]

10




A4'

wherein A₁', A₇', and A₁₁', which may be the same or different, represent acetyl or propionyl, provided that A₁', A₇', and A₁₁' do not simultaneously represent acetyl,

15 with a base to deacylate compound A1 or A4' and then protecting hydroxyl at the 1- and 11-positions to give compound D represented by formula D:


[Chemical formula 6]

6




wherein two R_2 s together represent a group selected from groups represented by formulae D-1, D-2, D-3, and D-4:
[Chemical formula 7]


5




D-1



D-2



D-3




D-4

wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X_s , which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5,

10

then protecting hydroxyl at the 7-position of compound D to give compound E represented by formula E:

[Chemical formula 8]



15


wherein

R_{1b} represents formyl; optionally substituted straight chain C₁₋₄ alkylcarbonyl; optionally substituted benzyl; group - SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; C₁₋₆ alkyloxy-C₁₋₆ alkyl optionally substituted by halogen atom; C₁₋₆ alkylthio-C₁₋₆ alkyl optionally substituted by halogen atom; straight chain, branched chain, or cyclic C₁₋₄ alkyl optionally substituted by halogen atom, provided that, when alkyl in the C₁₋₄ alkyl group is of a branched chain or cyclic type, the alkyl group is C₃₋₄ alkyl; C₂₋₆ alkenyl optionally substituted by halogen atom; C₂₋₆ alkynyl optionally substituted by halogen atom; or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group,

wherein, in R_{1b}, the substituent optionally possessed by alkylcarbonyl is selected from the group consisting of halogen atoms, C₁₋₄ alkyloxy, C₁₋₄ haloalkyloxy, C₁₋₄ alkylcarbonyl, C₁₋₄ haloalkylcarbonyl, C₁₋₄ alkylcarbonyloxy, and C₁₋₄ haloalkylcarbonyloxy, and the substituent optionally possessed by the heterocyclic group and benzyl is selected from the group consisting of halogen atoms, C₁₋₄ alkyl, C₁₋₄ alkyloxy, C₁₋₄ haloalkyloxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, C₁₋₄ alkylcarbonyl, C₁₋₄ haloalkylcarbonyl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ haloalkylcarbonyloxy, nitro, and cyano, and



R₂ is as defined above, and further removing the protective groups at the 1- and 11-positions of compound E to give compound Fb represented by formula Fb:

[Chemical formula 9]



wherein R_{1b} is as defined above, and
 (b) acylating hydroxyl at the 1- and 11-positions of compound
 Fa or Fb with an acylating agent corresponding to contemplated
 5 R' to give compound B2a or B2b represented by formula B2a or
 B2b:

[Chemical formula 10]




10 wherein R_{1a} , R_{1b} , and R' are as defined above,
 and then removing the protective group at the 7-position of
 compound B2a or compound B2b.

[0009]

According to a second aspect of the present invention,
 15 there is provided a process for producing compound B2a
 represented by formula B2a:


[Chemical formula 11]



wherein R_{1a} is as defined above and R' represents cyclic C_{3-6} alkylcarbonyl, the process comprising:

5 hydrolyzing acetyl at the 7-position of compound A1 described above with a base to selectively deacylate compound A1, then protecting hydroxyl at the 7-position to give compound B1 represented by formula B1:

[Chemical formula 12]



10


wherein Ac and R_{1a} are as defined above,

then hydrolyzing acetyl at the 1- and 11-positions of compound B1 with a base to deacylate compound B1 and thus to give compound Fa represented by formula Fa:

15

[Chemical formula 13]

10



wherein R_{1a} is as defined above,


and then acylating hydroxyl at the 1- and 11-positions of compound Fa with an acylating agent corresponding to
5 contemplated R' .

[0010]

According to a third aspect of the present invention, there is provided a process for producing compound B2b represented by formula B2b described above wherein R'
10 represents cyclic C_{3-6} alkylcarbonyl, the process comprising:

hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A4 represented by formula A4:

[Chemical formula 14]



15

wherein A_1 , A_7 , and A_{11} , which may be the same or different, represent acetyl or propionyl,

with a base to deacylate compound A4, then protecting hydroxyl at the 1- and 11-positions to give compound D, then
20 protecting hydroxyl at the 7-position of compound D to give compound E described above, further removing the protective

group at the 1- and 11-positions of compound E to give compound Fb described above, and then acylating hydroxyl at the 1- and 11-positions of compound Fb with an acylating agent corresponding to R'.

5 [0011]

According to a fourth aspect of the present invention, there is provided a process for producing compound C represented by formula C wherein R' represents cyclic C₃₋₆ alkylcarbonyl. The process comprises acylating hydroxyl at the 1- and 11-positions of compound Fb described above with an acylating agent corresponding to R' to give compound B2b and then removing the protective group at the 7-position of compound B2b.

[0012]

15 According to a fifth aspect of the present invention, there is provided a process for producing compound C described above. The process comprises hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A4 with a base to deacylate compound A4, then protecting hydroxyl at the 1- and 11-positions to give compound D, then protecting hydroxyl at the 7-position of compound D to give compound E, further removing the protective groups at the 1- and 11-positions of compound E to give compound Fb, then acylating hydroxyl at the 1- and 11-positions of compound Fb with an acylating agent corresponding to R' to give compound B2b, and then removing the protective group at the 7-position of compound B2b.


[0013]

30 According to a sixth aspect of the present invention, there is provided a compound that has acyloxy at the 1- and 11-positions and hydroxyl at the 7-position and is useful as an intermediate for the production of pyripyropene derivatives. The compound is represented by formula B2b:

[Chemical formula 15]

20375-1007

12



wherein R_{1b} is as defined above; and R' represents cyclic C_{3-6} alkylcarbonyl.


[0013a]

5

In particular aspects the invention relates to:

[1] A process for producing compound C represented by formula C:

[Chemical formula 1]



wherein R' represents cyclic C_{3-6} alkylcarbonyl, the process comprising


10 the steps of:

20375-1007

12a

(a₁) hydrolyzing acetyl at the 7-position of compound A1 represented by formula A1:

[Chemical formula 2]




A1

5

wherein Ac represents acetyl,

with a base to selectively deacylate compound A1, then protecting hydroxyl at the 7-position to give compound B1 represented by formula B1:

[Chemical formula 3]



B1

10

wherein

Ac is as defined above,


20375-1007

12b

R_{1a} represents group $-\text{SiR}_3\text{R}_4\text{R}_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group,

- 5 then hydrolyzing acetyl at the 1- and 11-positions of compound B1 with a base to deacylate compound B1 and thus to give compound Fa represented by formula Fa:


[Chemical formula 4]



- 10 wherein R_{1a} is as defined above, or

(a₂) hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A1 or compound A4' represented by formula A4':

[Chemical formula 5]




20375-1007

12c

wherein A_1' , A_7' , and A_{11}' , which may be the same or different, represent acetyl or propionyl, provided that A_1' , A_7' , and A_{11}' do not simultaneously represent acetyl,

5 with a base to deacylate compound A1 or A4' and then protecting hydroxyl at the 1- and 11-positions to give compound D represented by formula D:


[Chemical formula 6]




wherein two R_2 s together represent a group selected from groups represented by formulae D-1, D-2, D-3, and D-4:

10


[Chemical formula 7]




D-1



D-2



D-3



D-4


wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X_s , which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5,

20375-1007

12d

then protecting hydroxyl at the 7-position of compound D to give compound E represented by formula E:

[Chemical formula 8]



5 wherein.

R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, and


10 R_2 is as defined above,

and further removing the protective groups at the 1- and 11-positions of compound E to give compound Fb represented by formula Fb:

20375-1007

12e


[Chemical formula 9]



wherein R_{1b} is as defined above, and

- (b) acylating hydroxyl at the 1- and 11-positions of compound Fa or Fb
 5 with an acylating agent corresponding to contemplated R' to give compound B2a or B2b represented by formula B2a or B2b:

[Chemical formula 10]



wherein R_{1a} , R_{1b} , and R' are as defined above,

- 10 and then removing the protective group at the 7-position of compound B2a or compound B2b.

[2] The process according to [1], wherein R_{1b} represents acetyl; chloroacetyl; or group $-\text{SiR}_3\text{R}_4\text{R}_5$ optionally substituted by halogen atom wherein R_3 ,

20375-1007


12f

R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl.

[3] The process according to [1], wherein two R_2 s together represent a group represented by formula D-1 or D-2:

5

[Chemical formula 11]




wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X s, which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5.

10

[4] A process for producing compound C represented by formula C:

[Chemical formula 23]




wherein R' represents cyclic C_{3-6} alkylcarbonyl, the process comprising:

20375-1007

12g

acylating hydroxyl at the 1- and 11-positions of compound Fb represented by formula Fb:

[Chemical formula 24]




5 wherein

R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group,

10 with an acylating agent corresponding to R' to give compound B2b represented by formula B2b:

[Chemical formula 25]



20375-1007

12h


wherein R_{1b} and R' are as defined above,

and then removing the protective group at the 7-position of compound B2b.

[5] A process for producing compound C represented by formula C:

5

[Chemical formula 26]




wherein R' represents cyclic C_{3-6} alkylcarbonyl, the process comprising:

hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A4 represented by formula A4:

10

[Chemical formula 27]




20375-1007

12i

wherein A_1 , A_7 , and A_{11} , which may be the same or different, represent acetyl or propionyl,


with a base to deacylate compound A4, then protecting hydroxyl at the 1- and 11-positions to give compound D represented by formula D:

5 [Chemical formula 28]



wherein two R_2 s together represent a group selected from groups represented by formulae D-1, D-2, D-3, and D-4:

[Chemical formula 29]



10


wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X_s , which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5,

15 then protecting hydroxyl at the 7-position of compound D to give compound E represented by formula E:

20375-1007

12j

[Chemical formula 30]




wherein

- R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, and

R_2 is as defined above,

- and further removing the protective groups at the 1- and 11-positions of compound E to give compound Fb represented by formula Fb:

[Chemical formula 31]




20375-1007

12k

wherein R_{1b} is as defined above,

then acylating hydroxyl at the 1- and 11-positions of compound Fb with an acylating agent corresponding to R' to give compound B2b represented by B2b:

[Chemical formula 32]



5


wherein R_{1b} and R' are as defined above,

and then removing the protective group at the 7-position of compound B2b.

[6] A compound represented by formula B2b:

10

[Chemical formula 33]



wherein.

20375-1007

121

R_{1b} represents acetyl; chloroacetyl; group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, and

5 R' represents cyclic C₃₋₆ alkylcarbonyl.

[0014]

The present invention allows for producing pyripyropene derivatives that have acyloxy groups at the 1- and 11-positions and a hydroxyl group at the 7-position and are useful as pest control agents at high yield.

10 DETAILED DESCRIPTION OF THE INVENTION

[0015]

The term "halogen" as used herein means fluorine, chlorine, bromine, or iodine.

[0016]

15 The terms "alkyl", "alkenyl", or "alkynyl" as used herein as a substituent or a part of a substituent means alkyl, alkenyl, or alkynyl that is of a straight chain, branched chain, or cyclic type or a type of a combination thereof unless otherwise specified.

[0017]

20 The symbol "C_{a-b}" attached to a substituent as used herein means that the number of carbon atoms contained in the substituent as used herein is a to b. Further, "C_{a-b}" in "C_{a-b} alkylcarbonyl" means that the number of carbon atoms in the alkyl moiety excluding the carbon atoms in the carbonyl moiety is a to b.

20375-1007

12m

[0018]

The term "haloalkyl" as used herein means alkyl substituted by at least one halogen atom. Likewise, the terms

"haloalkyloxy," "haloalkylcarbonyl," and "haloalkylcarbonyloxy" respectively mean alkyloxy substituted by at least one halogen atom, alkylcarbonyl substituted by at least one halogen atom, and alkylcarbonyloxy substituted by at least one halogen atom.

5 [0019]

Specific examples of the straight chain, branched chain, or cyclic C₂₋₆ alkylcarbonyl group, represented by R', wherein, when the alkyl moiety in the C₂₋₆ alkylcarbonyl group is of a branched chain or cyclic type, the alkyl moiety is C₃₋₆ alkylcarbonyl, include cyclopropanecarbonyl and propionyl. The alkylcarbonyl group is preferably cyclic C₃₋₆ alkylcarbonyl, more preferably cyclopropanecarbonyl.

[0020]

Specific examples of the group -SiR₃R₄R₅, wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl, represented by R_{1a} and R_{1b} include trimethylsilyl, triethylsilyl, triisopropylsilyl, tert-butyl dimethylsilyl, and tert-butyl diphenylsilyl. The group -SiR₃R₄R₅ is optionally substituted, and such substituents include halogen atoms. In the group -SiR₃R₄R₅, preferably, all of R₃, R₄, and R₅ represent straight chain or branched chain C₁₋₆ alkyl, that is, the group -SiR₃R₄R₅ is preferably alkyl silyl, more preferably tert-butyl dimethylsilyl.

[0021]

Specific examples of the straight chain, branched chain, or cyclic C₁₋₄ alkyl group, represented by R_{1a} and R_{1b}, wherein, when alkyl in the C₁₋₄ alkyl group is of a branched chain or cyclic type, the alkyl group is C₃₋₄ alkyl, include methyl, ethyl, propyl, i-propyl, cyclopropyl, n-butyl, i-butyl, and tert-butyl. The alkyl group is optionally substituted, and such substituents include halogen atoms.

[0022]

Specific examples of the C₂₋₆ alkenyl group represented by R_{1a} and R_{1b} include vinyl, (1- or 2-)propenyl, (1-, 2-, or 3-)butenyl, (1-, 2-, 3-, or 4-)pentenyl, and (1-, 2-, 3-, 4-, or 5-)hexenyl. The alkenyl group is optionally substituted, and such

substituents include halogen atoms.

[0023]

Specific examples of the C₂₋₆ alkynyl group represented by R_{1a} and R_{1b} include ethynyl, (1- or 2-)propynyl, (1-, 2-, or 3-
5)butynyl, (1-, 2-, 3-, or 4-)pentynyl, and (1-, 2-, 3-, 4-, or 5-)hexynyl. The alkynyl group is optionally substituted, and such substituents include halogen atoms.

[0024]

Specific examples of the saturated or unsaturated five- or
10 six-membered heterocyclic group represented by R_{1a} and R_{1b} include tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, and tetrahydrothiofuranyl. The heterocyclic group is optionally substituted, and such substituents include halogen atoms, C₁₋₄ alkyl, C₁₋₄ alkyloxy, C₁₋₄ haloalkyloxy, C₁₋₄
15 alkylthio, C₁₋₄ haloalkyl, C₁₋₄ alkylcarbonyl, C₁₋₄ haloalkylcarbonyl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ haloalkylcarbonyloxy, nitro, and cyano. The heterocyclic group is preferably tetrahydropyranyl.

[0025]

Specific examples of the straight chain C₂₋₄ alkylcarbonyl
20 group represented by R_{1a} include propionyl, propylcarbonyl, and n-butylcarbonyl. The alkylcarbonyl group is optionally substituted, and such substituents include halogen atoms, C₁₋₄ alkyloxy, C₁₋₄ haloalkyloxy, C₁₋₄ alkylcarbonyl, C₁₋₄
25 haloalkylcarbonyl, C₁₋₄ alkylcarbonyloxy, and C₁₋₄ haloalkylcarbonyloxy.

[0026]

Specific examples of the straight chain C₁₋₄ alkylcarbonyl
30 group represented by R_{1b} include acetyl, propionyl, propylcarbonyl, and n-butylcarbonyl. The alkylcarbonyl group is optionally substituted, and such substituents include halogen atoms, C₁₋₄ alkyloxy, C₁₋₄ haloalkyloxy, C₁₋₄ alkylcarbonyl, C₁₋₄ haloalkylcarbonyl, C₁₋₄ alkylcarbonyloxy, and C₁₋₄ haloalkylcarbonyloxy.

35 [0027]

The C₁₋₆ alkyloxy-C₁₋₆ alkyl group represented by R_{1a} and

R_{1b} is optionally substituted, and such substituents include halogen atoms.

[0028]

The C₁₋₆ alkylthio-C₁₋₆ alkyl group represented by R_{1a} and
5 R_{1b} is optionally substituted, and such substituents include halogen atoms.

[0029]

The benzyl group represented by R_{1a} and R_{1b} is optionally substituted, and such substituents include halogen atoms, C₁₋₄
10 alkyl, C₁₋₄ alkyloxy, C₁₋₄ haloalkyloxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, C₁₋₄ alkylcarbonyl, C₁₋₄ haloalkylcarbonyl, C₁₋₄ alkylcarbonyloxy, C₁₋₄ haloalkylcarbonyloxy, nitro, and cyano.

[0030]

Preferably, R_{1a} represents group -SiR₃R₄R₅ optionally
15 substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group, more preferably group -SiR₃R₄R₅ wherein R₃, R₄, and R₅ each
20 independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, still more preferably group -SiR₃R₄R₅ wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; or
25 tetrahydropyranyl, most preferably tert-butyldimethylsilyl or tetrahydropyranyl.

[0031]


Preferably, R_{1b} represents acetyl, chloroacetyl, an optionally substituted saturated or unsaturated five- or six-
30 membered heterocyclic group, or group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl, more preferably acetyl, chloroacetyl, or group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄,
35 and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl, still more preferably acetyl,

chloroacetyl, or tert-butyldimethylsilyl, most preferably acetyl or chloroacetyl.

[0032]

The substituent represented by combining two R₂s
5 together is preferably a group represented by formula D-1 or D-2:

[Chemical formula 16]



10 wherein Y₁ represents a hydrogen atom or C₁₋₄ alkyl; Xs, which may be the same or different, represent a hydrogen atom, C₁₋₄ alkoxy, or nitro; and n is 0 to 5, more preferably isopropylidene, benzylidene, or p-methoxybenzylidene. According to another
15 embodiment, the substituent represented by combining two R₂s together is preferably D-1, more preferably isopropylidene.

[0033]

Preferably, A₁, A₇, and A₁₁ each represent acetyl.

[0034]

According to a preferred embodiment of the present
20 invention, in the process according to the first aspect of the present invention or the process according to the fifth aspect of the present invention, R' represents cyclic C₃₋₆ alkylcarbonyl.

[0035]

According to another preferred embodiment of the
25 present invention, in the process according to the first aspect of the present invention, R_{1a} represents group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl; or an optionally substituted saturated or
30 unsaturated five- or six-membered heterocyclic group.


[0036]

According to still another preferred embodiment of the present invention, in the process according to the first aspect of the present invention or the process according to the third aspect of the present invention, R_{1b} represents acetyl, chloroacetyl, or group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl.

[0037]

According to a further preferred embodiment of the present invention, in the process according to the first aspect of the present invention or the process according to the third aspect of the present invention, two R_2 s together represent a group represented by formula D-1 or D-2:

[Chemical formula 17]



wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X_s , which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5.

[0038]

According to a still further preferred embodiment of the present invention, in the process according to the second aspect of the present invention, R_{1a} represents optionally substituted straight chain C_{2-4} alkylcarbonyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group.

[0039]

According to another preferred embodiment of the present invention, in the compound according to the sixth

aspect of the present invention, R_{1b} represents acetyl, chloroacetyl, group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl, or an
5 optionally substituted saturated or unsaturated five- or six-membered heterocyclic group; and R' represents cyclic C_{3-6} alkylcarbonyl.

[0040]

According to still another preferred embodiment of the
10 present invention, in the process, R' in formulae B2a, B2b, and C represents propionyl or cyclopropanecarbonyl.

[0041]

According to a further preferred embodiment of the present invention, in the process, the contemplated compound
15 is produced through compound Fa wherein, in formula B1, Fa, or B2a, R_{1a} represents optionally substituted straight chain or branched chain alkylsilyl or an optionally substituted saturated or unsaturated five- or six-membered heterocyclic group.

[0042]

20 According to a still further preferred embodiment of the present invention, in the process, the contemplated compound is produced through compound Fb wherein R_{1b} in formula E, Fb, or B2b represents acetyl, chloroacetyl, or optionally substituted straight chain or branched chain alkylsilyl.

25 According to another preferred embodiment of the present invention, in the process, the contemplated compound is produced through compound Fb wherein R_2 in formula D or E is a group represented by formula D-3.


According to a more preferred embodiment of the present
30 invention, the contemplated compound is produced through compounds D, E, Fb, and B2b wherein R_2 in formula D or E represents a group represented by formula D-3; R_{1b} in formula E, Fb, or B2b represents acetyl, chloroacetyl, or optionally substituted straight chain or branched chain alkylsilyl; and R' in
35 formulae B2b and C represents cyclopropanecarbonyl.

[0043]

According to another aspect of the present invention, there is provided a process for producing compound C represented by formula C:

[Chemical formula 18]

5




wherein R' represents cyclic C₃₋₆ alkylcarbonyl, the process comprising:

hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A4 represented by formula A4:

10

[Chemical formula 19]



wherein A₁, A₇, and A₁₁, which may be the same or different, represent acetyl or propionyl


15

with a base to deacylate compound A4, then protecting hydroxyl at the 1- and 11-positions to give compound D represented by formula D:

[Chemical formula 20]

20


20



wherein two R_2 s together represent a group represented by formula D-1:

[Chemical formula 21]


5



wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl, then protecting hydroxyl at the 7-position of compound D to give compound E represented by formula E:

10

[Chemical formula 22]




wherein R_{1b} represents acetyl or chloroacetyl and R_2 is as defined above,


15

further removing the protective groups at the 1- and 11-positions of compound E to give compound Fb represented by formula Fb:

[Chemical formula 23]



then acylating hydroxyl at the 1- and 11-positions of
 compound Fb with an acylating agent corresponding to R' to
 5 give compound B2b represented by formula B2b:
 [Chemical formula 24]



wherein R_{1b} and R' are as defined above,
 10 and then removing the protective group at the 7-position
 of compound B2b.


[0044]

The present invention will be described in detail
 according to the following scheme.

15

[0045]

[Chemical formula 25]



[0046]

In the scheme, Ac, R_{1a}, R_{1b}, A₁, A₇, A₁₁, and R₂ are as defined above; R' represents straight chain, branched chain, or cyclic C₂₋₆ alkylcarbonyl wherein, when the alkyl moiety in the

C₂₋₆ alkylcarbonyl group is of a branched chain or cyclic type, the alkyl moiety is C₃₋₆ alkylcarbonyl.

[0047]

The product in each step may be used in a next step
5 without post treatment.

[0048]

1-1: Production of compound A3 from compound A1

Compound A1 can be produced by processes described, for example, in Japanese Patent Application Laid-Open No.
10 184158/1994, WO2004/060065, Japanese Patent Application Laid-Open No. 259569/1996, or Bioorganic Medicinal Chemistry Letter Vol. 5, No. 22, p. 2683.

[0049]

Solvents usable in the step of producing compound A3
15 from compound A1 include alcohol solvents having 1 to 4 carbon atoms such as methanol, ether solvents such as diethyl ether, diisopropyl ether, tetrahydrofuran, and dioxane, aprotic polar organic solvents such as N,N-dimethylformamide, dimethylsulfoxide, N,N-dimethylacetamide, and acetonitrile,
20 halogenated solvents such as dichloromethane and chloroform, or water, and mixed solvents composed of two or more of these solvents.

[0050]

Bases usable herein include inorganic bases such as
25 sodium carbonate, potassium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, potassium hydroxide, sodium hydride, potassium hydride, sodium cyanide, potassium cyanide, magnesium hydroxide, calcium hydroxide, lithium hydroxide, and barium hydroxide,
30 alkali metals such as sodium methoxide, sodium ethoxide, and potassium tert-butoxide, alkoxides of alkaline earth metals, or organic bases such as 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine, diisopropylethylamine, pyridine, hydrazine, and guanidine.
35 Preferred are 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, sodium carbonate, potassium

carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, potassium hydroxide, sodium methoxide, and potassium tert-butoxide. Particularly preferred are 1,8-diazabicyclo[5.4.0]undeca-7-ene and potassium tert-butoxide.

[0051]

The amount of the base used is preferably 0.01 to 1.2 equivalents based on the amount of compound A1. The reaction temperature is preferably -20°C to 50°C. The reaction time is preferably 0.5 hr to seven days.

[0052]

1-2: Production of compound B1 from compound A3

In the step of producing compound B1 from compound A3, hydroxyl at the 7-position can be protected by using a halide of R_{1a} represented by R_{1a}-Hal, wherein Hal represents halogen atom, an acid anhydride of R_{1a}, or a mixed acid anhydride of R_{1a}, corresponding to contemplated R_{1a}, or 3,4-dihydropyran in the presence of a base, in the presence of an acid, or in the absence of a base and an acid, or using a condensing agent such as dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, carbonyldiimidazole, dipyridyl disulfide, diimidazolyl disulfide, 1,3,5-trichlorobenzoyl chloride, 1,3,5-trichlorobenzoyl anhydride, PyBop, or PyBrop.

[0053]

This step may be carried out in the absence or presence of a solvent. Solvents usable herein include ketone solvents such as acetone and diethyl ketone, ether solvents such as diethyl ether, diisopropyl ether, and tetrahydrofuran, ester solvents such as ethyl acetate and butyl acetate, aprotic polar organic solvents such as N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, and acetonitrile, polar organic solvents such as pyridine, halogenated hydrocarbon solvents such as dichloromethane and chloroform, or aromatic hydrocarbon solvents such as toluene, and mixed solvents composed of two or more of these solvents.

[0054]

Bases usable herein include, for example, sodium carbonate, potassium carbonate, sodium hydride, potassium tert-butoxide, sodium methoxide, sodium ethoxide, pyridine, 5 dimethylaminopyridine, imidazole, 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine, or diisopropylethylamine.

[0055]

Acids usable herein include, for example, p- 10 toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, 10-camphorsulfonic acid, hydrochloric acid, or sulfuric acid.

[0056]

The reaction temperature is preferably -20°C to 50°C. 15 The reaction time is preferably 0.5 hr to four days.

[0057]

1-3: Production of compound Fa from compound B1

Solvents usable in the step of producing compound Fa from compound B1 include alcohol solvents having 1 to 4 carbon 20 atoms such as methanol, ether solvents such as diethyl ether, diisopropyl ether, tetrahydrofuran, and dioxane, aprotic polar organic solvents such as N,N-dimethylformamide, dimethylsulfoxide, N,N-dimethylacetamide, and acetonitrile, halogenated solvents such as dichloromethane and chloroform, 25 or water, and mixed solvents composed of two or more of these solvents.

[0058]

Bases usable herein include inorganic bases such as sodium carbonate, potassium carbonate, sodium hydrogen 30 carbonate, potassium hydrogen carbonate, sodium hydroxide, potassium hydroxide, sodium hydride, potassium hydride, sodium cyanide, potassium cyanide, magnesium hydroxide, calcium hydroxide, lithium hydroxide, and barium hydroxide, alkali metals such as sodium methoxide, sodium ethoxide, and 35 potassium tert-butoxide, alkoxides of alkaline earth metals, or organic bases such as 1,8-diazabicyclo[5.4.0]undeca-7-ene,

1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine, diisopropylethylamine, pyridine, hydrazine, and guanidine. Preferred are 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, sodium carbonate, potassium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, and potassium hydroxide. Particularly preferred is potassium carbonate.

[0059]

The amount of the base used is preferably 0.01 to 10 equivalents based on the amount of compound B1. The reaction temperature is preferably -20°C to 50°C. The reaction time is preferably 0.5 to 48 hr.

[0060]

2-1: Production of compound A2 from compound A4

Compound A4 and compound A4' may be naturally occurring substances obtained by processes described, for example, in Japanese Patent Application Laid-Open No. 184158/1994, WO94/09147, and Japanese Patent Application Laid-Open No. 239385/1996. Alternatively, for example, derivatives obtained by a process described, for example, in Japanese Patent Application Laid-Open No. 259569/1996.

[0061]

Solvents usable in the step of producing compound A2 from compound A4 include alcohol solvents having 1 to 4 carbon atoms such as methanol, ether solvents such as diethyl ether, diisopropyl ether, tetrahydrofuran, and dioxane, aprotic polar organic solvents such as N,N-dimethylformamide, dimethylsulfoxide, N,N-dimethylacetamide, and acetonitrile, halogenated solvents such as dichloromethane and chloroform, or water, and mixed solvents composed of two or more of these solvents.

[0062]

Bases usable herein include inorganic bases such as sodium carbonate, potassium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, potassium hydroxide, sodium hydride, potassium hydride,

sodium cyanide, potassium cyanide, magnesium hydroxide, calcium hydroxide, lithium hydroxide, and barium hydroxide, alkali metals such as sodium methoxide, sodium ethoxide, and potassium tert-butoxide, alkoxides of alkaline earth metals, or
5 organic bases such as 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine, diisopropylethylamine, pyridine, hydrazine, and guanidine. Preferred are 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, sodium carbonate, potassium
10 carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, and potassium hydroxide. Particularly preferred is potassium carbonate.

[0063]

The amount of the base used is preferably 0.01 to 10
15 equivalents based on the amount of compound A4. The reaction temperature is preferably -20°C to 50°C. The reaction time is preferably 0.5 to 48 hr.

[0064]

According to this step, compound A2 can be likewise
20 produced from compound A1 or compound A4' (compounds which are the same as compound A4, except that A₁, A₇, and A₁₁ in compound A4 are A₁', A₇', and A₁₁', respectively).

[0065]

2-2: Production of compound E through compound D from
25 compound A2

Solvents usable in the step of producing compound D from compound A2 include ketone solvents such as acetone and diethyl ketone, ether solvents such as diethyl ether, diisopropyl ether, dioxane, and tetrahydrofuran, ester solvents such as
30 ethyl acetate and butyl acetate, aprotic polar organic solvents such as N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, and acetonitrile, halogenated hydrocarbon solvents such as dichloromethane and chloroform, or aromatic hydrocarbon solvents such as toluene, and mixed solvents
35 composed of two or more of these solvents.

[0066]

Hydroxyl at 1- and 11-positions can be protected by using, for example, dimethoxypropane, acetone, optionally substituted benzaldehyde or a dimethyl acetal form thereof, 2-methoxypropene, 2-ethoxypropene, phosgene, triphosgene, 5 trichloroacetyl chloride, p-nitrobenzyloxycarbonyl chloride, or carbonyldiimidazole corresponding to contemplated R₂. Further, preferably, an acid catalyst such as p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, 10-camphorsulfonic acid, hydrogen fluoride, 10 hydrochloric acid, hydrogen bromide, sulfuric acid, iodine, iron chloride, tin chloride, zinc chloride, aluminum chloride, trimethylchlorosilane, trimethylsilyltriflate, or 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (preferably pyridinium p-toluenesulfonate and p-toluenesulfonic acid) is used in an 15 amount of 0.001 to 20 equivalents, more preferably 0.01 to 5 equivalents, still more preferably 0.01 to 0.04 equivalent, based on compound A2.

[0067]

The reaction temperature is preferably -20°C to 50°C, 20 more preferably room temperature to 50°C. The reaction time is preferably 0.5 to 48 hr.

[0068]

More specifically, for example, compounds D and E which are compounds wherein R₂ represents D-1 can be produced by 25 allowing a reaction to proceed using a protective group introducing reagent such as dimethoxypropane, 2-methoxypropene, or 2-ethoxypropene in the presence of an acid catalyst such as p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, or 10-camphorsulfonic acid in an amount of 0.001 to 20 equivalents, 30 preferably 0.01 to 5 equivalents, more preferably 0.01 to 0.04 equivalent based on compound A2, or by allowing a reaction to proceed using p-toluenesulfonic acid, pyridinium p-toluenesulfonate, sulfuric acid, or copper sulfate in an amount 35 of 0.001 to 20 equivalents based on compound A2 in an acetone solvent.

[0069]

Compounds D and E which are compounds wherein R_2 represents D-2 can be produced by allowing a reaction to proceed using an optionally substituted benzaldehyde or a dimethyl acetal form thereof in the presence of an acid catalyst
5 such as p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, or zinc chloride in an amount of 0.001 to 20 equivalents based on compound A2.

[0070]

10 Next, the step of producing compound E from compound D can be carried out in the absence or presence of a solvent. Solvents usable herein include ketone solvents such as acetone, diethyl ketone, ether solvents such as diethyl ether, diisopropyl ether, dioxane, and tetrahydrofuran, ester solvents such as
15 ethyl acetate and butyl acetate, aprotic polar organic solvents such as N,N-dimethylformamide, dimethylsulfoxide, N,N-dimethylacetamide, and acetonitrile, polar organic solvents such as pyridine, halogenated hydrocarbon solvents such as dichloromethane and chloroform, or aromatic hydrocarbon
20 solvents such as toluene, and mixed solvents composed of two or more of these solvents.

[0071]

A protective group corresponding to R_{1b} can be introduced into hydroxyl at the 7-position by using a halide of
25 R_{1b} represented by R_{1b} -Hal, R_{1b} OH, R_{1b} Cl, $(R_{1b})_2O$, a mixed acid anhydride of R_{1b} , or 3,4-dihydropyran in the presence of a base, in the presence of an acid, or in the absence of a base and an acid, or using a condensing agent such as dicyclohexylcarbodiimide,
30 dimethylaminopropyl)carbodiimide 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, carbonyldiimidazole, dipyridyl disulfide, diimidazolyl disulfide, 1,3,5-trichlorobenzoyl chloride, 1,3,5-trichlorobenzoyl anhydride, PyBop, or PyBrop.

[0072]

35 Bases usable herein include, for example, sodium carbonate, potassium carbonate, sodium hydride, potassium

tert-butoxide, sodium methoxide, sodium ethoxide, pyridine, dimethylaminopyridine, imidazole, 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine, or diisopropylethylamine.

5 [0073]

Acids usable herein include, for example, p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, 10-camphorsulfonic acid, hydrochloric acid, or sulfuric acid.

10 [0074]

The reaction temperature is preferably -20°C to 50°C. The reaction time is preferably 0.5 hr to seven days.

[0075]

15 More specifically, for example, a compound wherein R_{1b} represents straight chain C_{1-4} alkylcarbonyl optionally substituted by halogen atom, for example, acetyl or chloroacetyl can be produced by allowing a reaction to proceed using $R_{1b}Cl$ or $(R_{1b})_2O$ in an amount of 1 to 20 equivalents based on compound D and pyridine, dimethylaminopyridine, or
20 triethylamine as a base in an amount of 0.1 to 20 equivalents based on compound D in the absence of a solvent or in tetrahydrofuran, dichloromethane, N,N-dimethylformamide, or pyridine or in a mixed solvent composed of two or more of these solvents at -20°C to 50°C.

25 [0076]

A compound wherein R_{1b} represents group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl can be produced by allowing a reaction to
30 proceed using a halide of R_{1b} in an amount of 1 to 10 equivalents based on compound D and imidazole as a base in an amount of 1 to 10 equivalents based on compound D in dichloromethane, chloroform, N,N-dimethylformamide, N,N-dimethylacetamide, or a mixed solvent composed of two or
35 more of these solvents at -20°C to 50°C.

[0077]

2-3: Production of compound Fb from compound E

Solvents usable in the step of producing compound Fb from compound E include alcohol solvents having 1 to 4 carbon atoms such as methanol, ketone solvents such as acetone and diethyl ketone, ether solvents such as diethyl ether, diisopropyl ether, tetrahydrofuran, and dioxane, aprotic polar organic solvents such as N,N-dimethylformamide, dimethylsulfoxide, N,N-dimethylacetamide, and acetonitrile, halogenated solvents such as dichloromethane and chloroform, or water, and mixed solvents composed of two or more of these solvents.

[0078]

In the deprotection of the R₂ moiety, an organic acid such as acetic acid, trifluoroacetic acid, trifluoroacetic anhydride, hydrogen fluoride, hydrochloric acid, hydrogen bromide, sulfuric acid, p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, or 10-camphorsulfonic acid, or a hydrogenation catalyst such as boron chloride, magnesium bromide, dinitro zinc, bismuth chloride, cerium chloride, iron chloride, tin chloride, zinc chloride, aluminum chloride, palladium-carbon, or palladium hydroxide can be used, depending upon the type of the protective group, in an amount of 0.01 to 20 equivalents based on compound E.

[0079]

The reaction temperature is preferably -20°C to 50°C. The reaction time is preferably 0.5 to 48 hr.

[0080]

More specifically, for example, when R₂ in compound E represents D-1, compound Fb can be produced by allowing a reaction to proceed using 0.01 to 20 equivalents, based on compound E, of hydrochloric acid, acetic acid, p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, or dinitro zinc, bismuth chloride (preferably hydrochloric acid, acetic acid, or pyridinium p-toluenesulfonate) in water, methanol, tetrahydrofuran, dichloromethane, chloroform, N,N-dimethylformamide, acetonitrile, or acetic acid or a mixed solvent composed of two or more of these solvents

at -20°C to 50°C, preferably at room temperature to 40°C.

[0081]

When R₂ represents D-2, compound Fb can be produced by allowing a reaction to proceed using 0.01 to 20 equivalents, based on compound E, of 10-camphorsulfonic acid in water, methanol, tetrahydrofuran, or chloroform or a mixed solvent composed of two or more of these solvents at -20°C to 50°C, preferably at room temperature to 40°C.

[0082]

10 3: Production of compound B2a from compound Fa and production of compound B2b from compound Fb

The step of producing compound B2a from compound Fa and the step of producing compound B2b from compound Fb can be carried out in the absence or presence of a solvent. Solvents usable herein include ketone solvents such as acetone and diethyl ketone, ether solvents such as diethyl ether, diisopropyl ether, and tetrahydrofuran, ester solvents such as ethyl acetate and butyl acetate, aprotic polar organic solvents such as N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, and acetonitrile, halogenated hydrocarbon solvents such as dichloromethane and chloroform, or aromatic hydrocarbon solvents such as toluene, and mixed solvents composed of two or more of these solvents.

[0083]

25 Group R' can be introduced into the 1- and 11-position by using R'OH, R'Cl, (R')₂O, or a mixed acid anhydride corresponding to contemplated R' in the presence or absence of a base or using a condensing agent such as dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, carbonyldiimidazole, dipyrityl disulfide, diimidazolyl disulfide, 1,3,5-trichlorobenzoyl chloride, 1,3,5-trichlorobenzoyl anhydride, PyBop, or PyBrop.

[0084]

35 Bases usable herein include, for example, sodium carbonate, potassium carbonate, sodium hydride, potassium

tert-butoxide, sodium methoxide, sodium ethoxide, pyridine, 4-dimethylaminopyridine, imidazole, 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine, and diisopropylethylamine (preferably
5 pyridine).

[0085]

The reaction temperature is preferably -20°C to 50°C, more preferably 0°C to 30°C. The reaction time is preferably 0.5 to 48 hr.

10 [0086]

4: Production of compound C from compound B2a or compound B2b

Solvents usable in the step of producing compound C from compound B2a or B2b include alcohol solvents having 1 to
15 4 carbon atoms such as methanol, ether solvents such as diethyl ether, diisopropyl ether, tetrahydrofuran, and dioxane, aprotic polar organic solvents such as N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, and acetonitrile, halogenated hydrocarbon solvents such as dichloromethane and
20 chloroform, aromatic hydrocarbon solvents such as toluene, or water, and mixed solvents composed of two or more of these solvents.

[0087]

The deprotection of R_{1b} in compound B2b can be carried
25 out depending upon the type of the protective group. For example, when R_{1b} represents formyl, acetyl, or chloroacetyl, for example, an inorganic base such as sodium carbonate, potassium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium hydroxide, potassium hydroxide,
30 sodium hydride, potassium hydride, sodium cyanide, potassium cyanide, magnesium hydroxide, calcium hydroxide, lithium hydroxide, or barium hydroxide, an alkali metal such as sodium methoxide, sodium ethoxide, or potassium tert-butoxide, an alkoxide of an alkaline earth metal, an organic base such as
35 1,8-diazabicyclo[5.4.0]undeca-7-ene, 1,5-diazabicyclo[4.3.0]nona-5-ene, triethylamine,

diisopropylethylamine, pyridine, hydrazine, or guanidine, preferably sodium methoxide, sodium hydroxide, sodium carbonate, or sodium hydrogen carbonate can be used as a base in an amount of 0.01 to 10 equivalents, preferably 0.1 to 2 equivalents, based on compound B2b.

[0088]

When R_{1b} represents C_{1-6} alkyloxy- C_{1-6} alkyl optionally substituted by halogen atom, C_{1-6} alkylthio- C_{1-6} alkyl optionally substituted by halogen atom, straight chain, branched chain, or cyclic C_{1-4} alkyl optionally substituted by halogen atom, C_{2-6} alkenyl optionally substituted by halogen atom, C_{2-6} alkynyl optionally substituted by halogen atom, an optionally substituted saturated or unsaturated five- or six-membered heterocyclic ring, optionally substituted benzyl, or group - $SiR_3R_4R_5$ optionally substituted by halogen atom, for example, an organic acid such as p-toluenesulfonic acid, p-toluenesulfonic acid monohydrate, pyridinium p-toluenesulfonate, hydrogen fluoride-pyridine, trihydrogen fluoride-triethylamine, acetic acid, acetic acid chloride, trifluoroacetic acid, trifluoroacetic anhydride, hydrogen fluoride, hydrochloric acid, hydrogen bromide, sulfuric acid, thiophenol, or 10-camphorsulfonic acid, a hydrogenation catalyst such as boron chloride, boron bromide, magnesium bromide, cerium chloride, copper chloride, copper sulfate, lithium chloride, iron chloride, tin chloride, zinc chloride, zinc bromide, aluminum chloride, titanium chloride, palladium-carbon, palladium hydroxide, or palladium chloride, trimethylchlorosilane, trimethyliodosilane, trimethylsilyltriflate, or 2,3-dichloro-5,6-dicyano-1,4-benzoquinone may be used in an amount of 0.1 to 10 equivalents based on compound B2b.

[0089]

The reaction temperature is preferably -20°C to 50°C , more preferably 0°C to room temperature. The reaction time is preferably 0.5 hr to seven days.

[0090]

According to this step, compound C can be likewise produced from compound B2a.

EXAMPLES

[0091]

The present invention is further illustrated by the following Examples that are not intended as a limitation of the invention.

[0092]

Example 1:

Synthesis of 7-deacetylpyripyropene A

Pyripyropene A (30 mg) was dissolved in an 80% aqueous methanol solution (2 mL). 1,8-Diazabicyclo[5.4.0]-undeca-7-ene (9 mg) was added to the solution, and the mixture was stirred at room temperature for 1.5 hr. Acetic acid was added to the reaction solution to stop the reaction. The solvent was then removed by evaporation under the reduced pressure, water was added to the residue, and the mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with saturated brine and was dried over anhydrous magnesium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product of 7-deacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, acetone : hexane = 1 : 1) to give 7-deacetylpyripyropene A (17 mg, yield 61%). The results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-7 described in Japanese Patent Application Laid-Open No. 259569/1996.

[0093]

Example 2:

Synthesis of 7-O-tert-butyldimethylsilyl-7-deacetylpyripyropene A

7-Deacetylpyripyropene A (30 mg) synthesized by the process described in Example 1 was dissolved in N,N-dimethylformamide (5 mL), and imidazole (113 mg) and tert-butyldimethylchlorosilane (250 mg) were added to the solution. The mixture was stirred at room temperature for 24 hr. The

reaction solution was then poured into water, and the mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with saturated brine and was dried over anhydrous magnesium sulfate, and the solvent was then removed by evaporation under the reduced pressure to give a crude product of 7-O-tert-butyldimethylsilyl-7-deacetylpyripyropene A (470 mg).

[0094]

ESI-MS: m/z 656 ($M+H$)⁺;

¹H-NMR (CDCl₃): δ 0.11 (3H, s), 0.16 (3H, s), 0.90 (3H, s), 0.96 (9H, s), 1.30-1.38 (1H, m), 1.32-1.37 (1H, m), 1.41 (3H, s), 1.60 (3H, s), 1.61-1.69 (2H, m), 1.77-1.92 (1H, m), 2.05 (6H, s), 2.15 (1H, m), 2.89 (1H, d, $J = 2.4$ Hz), 3.64-3.70 (2H, m), 3.73 (1H, d, $J = 11.6$ Hz), 3.83 (1H, d, $J = 11.6$ Hz), 4.78 (1H, dd, $J = 4.8, 11.2$ Hz), 4.99 (1H, m), 6.36 (1H, s), 7.42 (1H, dd, $J = 4.8, 8.0$ Hz), 8.11 (1H, d, $J = 8.0$ Hz), 8.70 (1H, d, $J = 4.4$ Hz), 9.00 (1H, d, $J = 2.0$ Hz).

[0095]

Example 3:

Synthesis of 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A

7-O-tert-butyldimethylsilyl-7-deacetylpyripyropene A (470 mg) produced in Example 2 was dissolved in an 88% aqueous methanol solution (40 mL). Potassium carbonate (307 mg) was added to the solution, and the mixture was stirred at room temperature for 19.5 hr. The solvent was removed by evaporation under the reduced pressure. Water and ethyl acetate were added to the residue. The solid remaining undissolved was collected by filtration to give 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A (65 mg). The mother liquid was extracted with ethyl acetate. The ethyl acetate layer was then washed with saturated brine and was dried over anhydrous magnesium sulfate. The solvent was removed by evaporation under the reduced pressure to give 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A (235 mg). Thus, 300 mg in total (yield in two stages from Example

2: 95%) of 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A was obtained.

[0096]

ESI-MS: m/z 572 ($M+H$)⁺;

5 ¹H-NMR (CD₃OD): δ 0.08 (3H, s), 0.13 (3H, s), 0.64 (3H, s), 0.90 (9H, s), 1.19 (1H, dt, J = 3.6, 12.8 Hz), 1.31 (3H, s), 1.33-1.36 (2H, m), 1.48 (1H, t, J = 12.0 Hz), 1.53 (3H, s), 1.62-1.80 (3H, m), 1.99-2.03 (1H, m), 3.16 (1H, d, J = 10.8 Hz), 3.44 (1H, d, J = 10.8 Hz), 3.56 (1H, dd, J = 4.8, 11.6 Hz),
10 3.76 (1H, dd, J = 5.2, 11.2 Hz), 4.86 (1H, d, J = 3.2 Hz), 6.47 (1H, s), 7.47 (1H, ddd, J = 0.8, 4.8, 8.0 Hz), 8.17 (1H, dt, J = 2.0, 8.4 Hz), 8.55 (1H, dd, J = 2.0, 4.8 Hz), 8.91 (1H, dd, J = 0.8, 2.4 Hz).

[0097]

15 Example 4:

Synthesis of 7-O-tert-butyldimethylsilyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A (57 mg) synthesized by the process
20 described in Example 3 was dissolved in N,N-dimethylformamide (2 mL). Pyridine (0.5 mL) was added to the solution at 0°C, the mixture was stirred at that temperature for 30 min, and cyclopropanecarbonyl chloride (62 mg) was added thereto. The mixture was stirred at that temperature for 3 hr, and the
25 reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate, and the solvent was then removed by evaporation under the reduced pressure to give a crude product (87 mg) of 7-O-tert-butyldimethylsilyl-1,11-O-dicyclopropanecarbonyl-1,7,11-
30 trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 30 : 1) to give 7-O-tert-butyldimethylsilyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (67 mg, yield 95%).
35

[0098]

ESI-MS: m/z 708 ($M+H$)⁺;

¹H-NMR (CDCl₃): δ 0.11 (3H, s), 0.15 (3H, s), 0.85-0.88 (4H, m), 0.91 (3H, s), 0.96 (9H, s), 0.92-1.01 (4H, m), 1.25-1.36 (1H, m), 1.42 (3H, s), 1.45-1.47 (1H, m), 1.53-1.65 (5H, m),
5 1.58 (3H, s), 1.80-1.93 (2H, m), 2.12-2.16 (1H, m), 2.81 (1H, d, J = 2.0 Hz), 3.65 (1H, d, J = 12.0 Hz), 3.70 (1H, m), 3.91 (1H, d, J = 11.6 Hz), 4.81 (1H, dd, J = 4.8, 11.6 Hz), 4.98 (1H, m), 6.36 (1H, s), 7.41 (1H, dd, J = 4.8, 8.0 Hz), 8.10 (1H, dt, J = 2.0, 8.4 Hz), 8.69 (1H, dd, J = 1.6, 4.8 Hz), 9.00 (1H, d, J =
10 2.0 Hz).

[0099]

Example 5:

Synthesis of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

15 7-O-tert-butyldimethylsilyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (100 mg) synthesized by the process described in Example 4 was dissolved in tetrahydrofuran (1.5 mL). Pyridine (0.6 mL) and hydrogen fluoride/pyridine (0.9 mL) were added to the solution
20 at 0°C. The mixture was stirred at that temperature for 4 days, an aqueous sodium hydrogen carbonate solution was then added thereto, and the mixture was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then
25 removed by evaporation under the reduced pressure to give a crude product (92 mg) of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 20 :
30 1) to give 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (79 mg, yield 95%).

[0100]

ESI-MS: m/z 594 ($M+H$)⁺;

¹H-NMR (CDCl₃): δ 0.85-0.88 (4H, m), 0.92 (3H, s), 0.96-1.01 (4H, m), 1.35 (1H, dt, J = 4.0, 12.6 Hz), 1.42 (3H, s), 1.45-1.50 (2H, m), 1.56-1.63 (3H, m), 1.66 (3H, s), 1.79-1.93 (3H,

m), 2.14 (1H, m), 2.17 (1H, d, J = 3.6 Hz), 2.85 (1H, d, J = 2.0 Hz), 3.74 (1H, d, J = 12.0 Hz), 3.78-3.82 (1H, m), 3.86 (1H, d, J = 11.6 Hz), 4.82 (1H, dd, J = 5.2, 11.6 Hz), 4.99 (1H, m), 6.52 (1H, s), 7.42 (1H, dd, J = 4.8, 8.0 Hz), 8.11 (1H, dt, J = 1.9, 8.1 Hz), 8.70 (1H, dd, J = 1.6, 4.8 Hz), 9.00 (1H, d, J = 2.0 Hz).

[0101]

Example 6:

Synthesis of 7-deacetyl-7-O-tetrahydropyranylpripyropene A

10 7-Deacetylpyripyropene A (500 mg) produced by the process described in Example 1 was dissolved in dichloromethane (10 mL), and 3,4-dihydropyran (372 mg) and pyridinium p-toluenesulfonate (348 mg) were added to the solution. The mixture was stirred at room temperature for 73.5
15 hr. The reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous magnesium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product (742 mg) of 7-
20 deacetyl-7-O-tetrahydropyranylpripyropene A. The results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-44 described in Japanese Patent Application Laid-Open No. 259569/1996.

[0102]

25 Example 7:

Synthesis of 1,7,11-trideacetyl-7-O-tetrahydropyranylpripyropene A

7-Deacetyl-7-O-tetrahydropyranylpripyropene A (742 mg) produced in Example 6 was dissolved in a 66% aqueous
30 methanol solution (9 mL). Potassium carbonate (511 mg) was added to the solution, and the mixture was stirred at room temperature for 4 hr. Water was added thereto, and the solid remaining undissolved was collected by filtration to give 1,7,11-trideacetyl-7-O-tetrahydropyranylpripyropene A (453 mg, yield
35 in two stages from Example 6: 90%).

[0103]

ESI-MS: m/z 542 ($M+H$)⁺.

[0104]

Example 8:

5 Synthesis of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetyl-7-O-tetrahydropyranylpyripyropene A

1,7,11-Trideacetyl-7-O-tetrahydropyranylpyripyropene A (450 mg) synthesized by the process described in Example 7 was dissolved in N,N-dimethylformamide (6 mL). Pyridine (3 mL) was added to the solution at 0°C, and the mixture was
10 stirred at that temperature for 10 min. Cyclopropanecarbonyl chloride (525 mg) was added thereto. The mixture was stirred at that temperature for one hr. The reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried
15 over anhydrous magnesium sulfate, and the solvent was then removed by evaporation under the reduced pressure to give a crude product (800 mg) of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetyl-7-O-tetrahydropyranylpyripyropene A.

[0105]

20 ESI-MS: m/z 678 ($M+H$)⁺.

[0106]

Example 9:

Synthesis of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

25 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetyl-7-O-tetrahydropyranylpyripyropene A (800 mg) produced in Example 8 was dissolved in methanol (8 mL), and p-toluenesulfonic acid monohydrate (142 mg) was added to the solution at 0°C. The mixture was stirred at that temperature for 21.5 hr. An
30 aqueous sodium hydrogen carbonate solution was then added thereto. Methanol was removed by evaporation under the reduced pressure, and the residue was extracted with ethyl acetate. The ethyl acetate layer was washed with saturated brine and was dried over anhydrous magnesium sulfate. The
35 solvent was then removed by evaporation under the reduced pressure to give a crude product (570 mg) of 1,11-O-

dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by chromatography on silica gel (Mega Bond Elut (Varian), acetone : hexane = 3 : 5) to give 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (346 mg, yield in two stages from Example 8: 70%). ESI-MS data and ¹H-NMR data of the compound were in agreement with those in Example 5.

[0107]

Example 10:

10 Synthesis of 1,7,11-trideacetylpyripyropene A

Pyripyropene A (1 g) was dissolved in a 66% aqueous methanol solution (15 mL). Potassium carbonate (355 mg) was added to the solution, and the mixture was stirred at room temperature for 20 hr. The solvent was removed by evaporation under the reduced pressure. Ethyl acetate and water were added thereto, and crystals remaining undissolved were collected by filtration to give 1,7,11-trideacetylpyripyropene A (737 mg, yield 94%). The results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-3 described in Japanese Patent Application Laid-Open No. 259569/1996.

[0108]

Example 11:

25 Synthesis of 1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A

1,7,11-Trideacetylpyripyropene A (200 mg) synthesized by the process described in Example 10 was dissolved in N,N-dimethylformamide (2 mL). Acetone dimethyl acetal (456 mg) and pyridinium p-toluenesulfonate (550 mg) were added to the solution. The mixture was stirred at room temperature for 25.5 hr. The reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with a saturated aqueous sodium hydrogen carbonate solution and saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product of 1,7,11-

trideacetyl-1,11-O-isopropylidenepyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 10 : 1) to give 1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A (171 mg, yield 79%). The results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-16 described in Japanese Patent Application Laid-Open No. 269065/1996.

[0109]

10 Example 12:

Synthesis of 7-O-tert-butyldimethylsilyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A

1,7,11-Trideacetyl-1,11-O-isopropylidenepyripyropene A (168 mg) synthesized by the process described in Example 11 was dissolved in N,N-dimethylformamide (2 mL). Imidazole (92 mg) and tert-butyldimethylchlorosilane (204 mg) were added to the solution. The mixture was stirred at room temperature for 22 hr, and the reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product (193 mg) of 7-O-tert-butyldimethylsilyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 20 : 1) to give 7-O-tert-butyldimethylsilyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A (187 mg, yield 90%).

30 [0110]

ESI-MS: m/z 612 (M+H)⁺;

¹H-NMR (CDCl₃): δ 0.11 (3H, s), 0.16 (3H, s), 0.96 (9H, s), 1.03 (1H, m), 1.10 (3H, s), 1.33 (1H, dt, J = 3.6, 12.8 Hz), 1.40 (3H, s), 1.43 (3H, s), 1.44 (3H, s), 1.39-1.44 (1H, m), 1.55-1.58 (2H, m), 1.58 (3H, s), 1.64 (1H, q, J = 12.0 Hz), 1.81 (1H, dq, J = 3.6, 12.8 Hz), 2.20 (1H, dt, J = 3.2, 12.8 Hz),

2.81 (1H, d, J = 1.6 Hz), 3.42 (1H, d, J = 10.8 Hz), 3.51 (1H, d, J = 10.4 Hz), 3.50-3.53 (1H, m), 3.72 (1H, dd, J = 4.8, 11.2 Hz), 4.97 (1H, m), 6.35 (1H, s), 7.41 (1H, dd, J = 4.8, 8.0 Hz), 8.10 (1H, dt, J = 1.6, 8.0 Hz), 8.69 (1H, dd, J = 1.6, 4.8 Hz),
5 9.00 (1H, d, J = 2.0 Hz).

[0111]

Example 13:

Synthesis of 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A

10 7-O-tert-butyldimethylsilyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A (116 mg) synthesized by the process described in Example 12 was dissolved in tetrahydrofuran (1 mL), and 63% acetic acid (4 mL) was added to the solution at 0°C. The mixture was stirred at room
15 temperature for 24 hr. An aqueous sodium hydrogen carbonate solution was added thereto, and the mixture was extracted with chloroform. The chloroform layer was washed with a saturated aqueous sodium hydrogen carbonate solution and saturated brine and was dried over anhydrous sodium sulfate. The
20 solvent was then removed by evaporation under the reduced pressure to give a crude product (101 mg) of 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform :
25 methanol = 10 : 1) to give 7-O-tert-butyldimethylsilyl-1,7,11-trideacetylpyripyropene A (91 mg, yield 84%).

[0112]

ESI-MS: m/z 572 (M+H)⁺;

¹H-NMR (CD₃OD): δ 0.08 (3H, s), 0.13 (3H, s), 0.64 (3H, s),
30 0.90 (9H, s), 1.19 (1H, dt, J = 3.6, 12.8 Hz), 1.31 (3H, s), 1.33-1.36 (2H, m), 1.48 (1H, t, J = 12.0 Hz), 1.53 (3H, s), 1.62-1.80 (3H, m), 1.99-2.03 (1H, m), 3.16 (1H, d, J = 10.8 Hz), 3.44 (1H, d, J = 10.8 Hz), 3.56 (1H, dd, J = 4.8, 11.6 Hz), 3.76 (1H, dd, J = 5.2, 11.2 Hz), 4.86 (1H, d, J = 3.2 Hz), 6.47
35 (1H, s), 7.47 (1H, ddd, J = 0.8, 4.8, 8.0 Hz), 8.17 (1H, dt, J = 2.0, 8.4 Hz), 8.55 (1H, dd, J = 2.0, 4.8 Hz), 8.91 (1H, dd, J =

0.8, 2.4 Hz).

[0113]

Example 14:

Synthesis of 1,11-dideacetyl-1,11-O-isopropylidenepyripyropene

5 A

1,7,11-Trideacetyl-1,11-O-isopropylidenepyripyropene A (100 mg) synthesized by the process described in Example 11 was dissolved in dichloromethane (2 mL). Triethylamine (61 mg), 4-dimethylaminopyridine (7 mg), and acetic anhydride (26 mg) were added to the solution. The mixture was stirred at room temperature for 4 hr. The reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate, and the solvent was then removed by evaporation under the reduced pressure to give a crude product (120 mg) of 1,11-dideacetyl-1,11-O-isopropylidenepyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 20 : 1) to give 1,11-dideacetyl-1,11-O-isopropylidenepyripyropene A (103 mg, yield 95%). The results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-43 described in Japanese Patent Application Laid-Open No. 269065/1996.

[0114]

25 Example 15:

Synthesis of 1,11-dideacetylpyripyropene A

1,11-Dideacetyl-1,11-O-isopropylidenepyripyropene A (99 mg) synthesized by the process described in Example 14 was dissolved in tetrahydrofuran (1.2 mL) and methanol (2.4 mL), and pyridinium p-toluenesulfonate (185 mg) was added to the solution. The mixture was stirred at room temperature for 30 hr, triethylamine was then added thereto, and the solvent was removed by evaporation under the reduced pressure. Chloroform and water were added to the residue, and the mixture was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous

sodium sulfate, and the solvent was removed by evaporation under the reduced pressure to give a crude product (85 mg) of 1,11-dideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 60 : 1) to give 1,11-dideacetylpyripyropene A (64 mg, yield 72%). The results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-5 described in Japanese Patent Application Laid-Open No. 259569/1996.

10 [0115]

Example 15a:

Synthesis of 1,11-dideacetylpyripyropene A

1,7,11-Trideacetylpyripyropene A (9.67 g) was suspended in N,N-dimethylformamide (48 mL). Acetone dimethyl acetal (6.61 g) and p-toluenesulfonic acid monohydrate (0.08 g) were added to the suspension, and the mixture was stirred at 38 to 41°C for 4 hr. 4-Dimethylaminopyridine (0.08 g) was added thereto, and the mixture was removed by evaporation under the reduced pressure for 1.5 hr. The residue was cooled to 0°C. Triethylamine (2.57 g) and acetic anhydride (2.37 g) were added to the cooled solution, and the mixture was stirred at that temperature for 16 hr. Water (96 g) was then added to the reaction solution, and the mixture was adjusted to pH 7.17 by the addition of 5% hydrochloric acid. The precipitated light yellow powder was collected by filtration and was washed twice with water (20 g). The crude product thus obtained was suspended with methanol (48 mL), 15% hydrochloric acid (4.7 g) was added to the suspension, and the mixture was stirred at 25 to 27°C for 2 hr. Water (33 mL) was added thereto, and the insolubles were filtered, followed by adjustment of pH to 4.41 by the addition of a 5% aqueous sodium hydroxide solution. Further, water (31 mL) was added thereto. The precipitated light yellow powder was collected by filtration and was washed twice with a 30% aqueous methanol solution (20 mL). The washed powder was dried at 40°C for 23 hr to give 8.62 g of 1,11-dideacetylpyripyropene A. The results of measurement by

¹H-NMR showed that the ¹H-NMR data were in agreement with those of the compound produced in Example 15.

[0116]

Example 16:

5 Synthesis of 1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A

1,11-Dideacetylpyripyropene A (61 mg) synthesized by the process described in Example 15 was dissolved in N,N-dimethylformamide (1.2 mL). Pyridine (0.3 mL) was added to
10 the solution at 0°C, the mixture was stirred at that temperature for 10 min, cyclopropanecarbonyl chloride (77 mg) was added thereto, and the mixture was stirred at that temperature for 1.5 hr. The reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed
15 with saturated brine and was dried over anhydrous magnesium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product (97 mg) of 1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A. The crude product was purified by preparative thin-layer column
20 chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 20 : 1) to give 1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A (73 mg, yield 93%).

[0117]

ESI-MS: m/z 636 (M+H)⁺;

25 ¹H-NMR (CDCl₃): δ 0.84-0.89 (4H, m), 0.89 (3H, s), 0.90-1.06 (4H, m), 1.37 (1H, dt, J = 3.8, 13.2 Hz), 1.45 (3H, s), 1.53 (1H, d, J = 4.0 Hz), 1.55-1.67 (4H, m), 1.70 (3H, s), 1.79-1.87 (2H, m), 1.89-1.94 (2H, m), 2.14-2.18 (1H, m), 2.16 (3H, s), 2.97 (1H, d, J = 2.0 Hz), 3.77 (2H, s), 4.81 (1H, dd, J = 4.8, 11.7
30 Hz), 5.00 (1H, m), 5.02 (1H, dd, J = 5.0, 11.4 Hz), 6.46 (1H, s), 7.40 (1H, dd, J = 4.9, 8.0 Hz), 8.09 (1H, dt, J = 1.9, 8.1 Hz), 8.68 (1H, dd, J = 1.6, 4.8 Hz), 9.00 (1H, d, J = 2.0 Hz).

[0118]

Example 16a:

35 Synthesis of 1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A

1,11-Dideacetylpyripyropene A (25.76 g) was suspended in ethyl acetate (130 mL), pyridine (15.84 g) was added to the suspension. The mixture was cooled to 10 to 15°C. Cyclopropanecarbonyl chloride (15.70 g) was added dropwise thereto, and the mixture was stirred at 25 to 30°C for 3 hr. The reaction solution was again cooled to 10 to 15°C, and water (50 mL) was added dropwise thereto. The mixture was adjusted to pH 2.59 by the addition of 5 N-hydrochloric acid, followed by separation. The organic layer was washed with 5% sodium bicarbonate water (50 mL) and 10% brine (50 mL) in that order. The ethyl acetate solution thus obtained was removed by evaporation under the reduced pressure and was further replaced with methanol to adjust the liquid volume to about 130 mL. Water (130 mL) was added dropwise thereto. The precipitated light yellow powder was collected by filtration, was washed twice with a 50% aqueous methanol solution (40 mL), and was dried at 40°C for 23 hr to give 30.80 g of 1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A. The results of measurement by ¹H-NMR showed that the ¹H-NMR data were in agreement with those of the compound produced in Example 16.

[0119]

Example 17:

Synthesis of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A (67 mg) synthesized by the process described in Example 16 was dissolved in a 95% aqueous methanol solution (0.07 mL). Sodium carbonate (22 mg) was added to the solution at 0°C. The mixture was stirred at that temperature for 4 days. Acetic acid was then added thereto. Methanol was removed by evaporation under the reduced pressure, and the residue was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude

product (74 mg) of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 10 : 1) to give 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (47 mg, yield 76%). ESI-MS data and ¹H-NMR data of the compound were in agreement with those of the compound produced in Example 5.

[0120]

10 Example 17a:

Synthesis of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

1,11-O-dicyclopropanecarbonyl-1,11-dideacetylpyripyropene A (30.00 g) was suspended in a mixed liquid composed of water (20 mL) and methanol (190 mL), and the mixture was cooled to 0 to 5°C. A 1 M methanol solution (4.49 mL) of sodium methoxide was added thereto, and the mixture was stirred at that temperature for 23 hr. 1.2% hydrochloric acid (20 mL) was added to the reaction solution, and the mixture was filtered through a 0.5-μm filter, followed by evaporation under the reduced pressure to adjust the liquid volume to about 90 mL. A mixed liquid (120 mL) of methanol/water = 2/1 was added to the residue to adjust the liquid volume to about 150 mL. Further, a mixed liquid (120 mL) of methanol/water = 2/1 was added thereto to adjust the liquid volume to about 180 mL. The mixture was stirred at room temperature for one hr, was then cooled to 5°C, and was stirred for 17 hr. The precipitated light yellow powder was collected by filtration, was washed twice with a 30% aqueous methanol solution (50 mL), and was dried at 40°C for 22 hr to give 23.82 g of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A. The results of measurement by ¹H-NMR showed that the ¹H-NMR data were in agreement with those of the compound produced in Example 17.

35 [0121]

Example 18:

Synthesis of 7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A

1,7,11-Trideacetyl-1,11-O-isopropylidenepyripyropene A (100 mg) synthesized by the process described in Example 11 was dissolved in tetrahydrofuran (2 mL), and triethylamine (61 mg) and chloroacetic anhydride (103 mg) were added to the solution. The mixture was stirred at room temperature for 3.5 hr, and the reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product (118 mg) of 7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 20 : 1) to give 7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A (80 mg, yield 70%).

[0122]

ESI-MS: m/z 574 ($M+H$)⁺;
¹H-NMR (CDCl₃): δ 1.11 (3H, s), 1.16 (1H, dd, J = 2.4, 12.6 Hz), 1.33-1.41 (1H, m), 1.44 (3H, s), 1.45 (3H, s), 1.52 (1H, d, J = 4.0 Hz), 1.58-1.65 (1H, m), 1.62 (3H, s), 1.70 (3H, s), 1.66-1.75 (2H, m), 1.77-1.86 (1H, m), 2.22 (1H, m), 2.90 (1H, d, J = 2.0 Hz), 3.48 (2H, s), 3.54 (1H, dd, J = 3.6, 12.0 Hz), 4.19 (2H, d, J = 4.0 Hz), 5.00 (1H, m), 5.09 (1H, dd, J = 5.6, 11.6 Hz), 6.45 (1H, s), 7.41 (1H, dd, J = 4.8, 8.0 Hz), 8.10 (1H, dt, J = 1.6, 8.0 Hz), 8.70 (1H, dd, J = 1.6, 4.8 Hz), 9.02 (1H, d, J = 1.6 Hz).

[0123]

Example 19:

Synthesis of 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A

7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-isopropylidenepyripyropene A (35 mg) synthesized by the process described in Example 18 was dissolved in tetrahydrofuran (0.6 mL) and methanol (1.2 mL), and

pyridinium p-toluenesulfonate (61 mg) was added to the solution. The mixture was stirred at room temperature for 31 hr, and the reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was
5 washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product (30 mg) of 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column
10 chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 10 : 1) to give 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A (24 mg, yield 74%).

[0124]

ESI-MS: m/z 534 (M+H)⁺;

15 ¹H-NMR (CD₃OD): δ 0.74 (3H, s), 1.32 (1H, m), 1.44 (3H, s), 1.54 (2H, m), 1.69-1.75 (2H, m), 1.75 (3H, s), 1.79-1.86 (1H, m), 1.91-1.94 (1H, m), 2.12 (1H, m), 3.26 (1H, d, J = 11.6 Hz), 3.52 (1H, d, J = 10.8 Hz), 3.67 (1H, dd, J = 5.2, 11.6 Hz), 4.33 (2H, d, J = 2.4 Hz), 4.98 (1H, m), 5.15 (1H, dd, J = 5.2, 11.6
20 Hz), 6.79 (1H, s), 7.55 (1H, dd, J = 4.8, 8.0 Hz), 8.28 (1H, dt, J = 2.4, 8.0 Hz), 8.62 (1H, dd, J = 1.6, 4.8 Hz), 9.02 (1H, d, J = 2.4 Hz).

[0125]

Example 20:

25 Synthesis of 7-O-chloroacetyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A (21 mg) synthesized by the process described in Example 19 was dissolved in N,N-dimethylformamide (1.2 mL), and pyridine (0.3
30 mL) and cyclopropanecarbonyl chloride (25 mg) were added to the solution at 0°C. The mixture was stirred at that temperature for 2.5 hr. The reaction solution was then poured into water, and the mixture was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried
35 over anhydrous magnesium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a

crude product (37 mg) of 7-O-chloroacetyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 30 : 1) to give 7-O-chloroacetyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (16 mg, yield 58%).

[0126]

ESI-MS: m/z 670 (M+H)⁺;

10 ¹H-NMR (CDCl₃): δ 0.85-0.90 (4H, m), 0.91 (3H, s), 0.96-1.08 (4H, m), 1.38 (1H, dt, J = 4.0, 12.6 Hz), 1.45 (3H, s), 1.54-1.67 (5H, m), 1.72 (3H, s), 1.81-1.95 (3H, m), 2.17 (1H, m), 2.89 (1H, d, J = 1.6 Hz), 3.78 (2H, s), 4.17 (2H, d, J = 2.8 Hz), 4.82 (1H, dd, J = 4.8, 11.6 Hz), 5.01 (1H, m), 5.09 (1H, dd, J = 5.2, 11.6 Hz), 6.45 (1H, s), 7.41 (1H, dd, J = 4.8, 8.0 Hz), 8.10 (1H, dt, J = 1.6, 8.0 Hz), 8.69 (1H, dd, J = 1.6, 4.8 Hz), 9.02 (1H, d, J = 1.6 Hz).

[0127]

Example 21:

20 Synthesis of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A

7-O-chloroacetyl-1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (14 mg) synthesized by the process described in Example 20 was dissolved in a 95% aqueous methanol solution (1.4 mL). Sodium hydrogen carbonate (1.9 mg) was then added to the solution. The mixture was stirred at room temperature for 3 hr. Acetic acid was added thereto, and methanol was then removed by evaporation under the reduced pressure to give a crude product of 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by preparative thin-layer column chromatography (Merck silica gel 60F₂₅₄ 0.5 mm, chloroform : methanol = 10 : 1) to give 1,11-O-dicyclopropanecarbonyl-1,7,11-trideacetylpyripyropene A (12 mg, yield 94%). ESI-MS data and ¹H-NMR data of the compound were in agreement with those of the compound produced in Example 5.

[0128]

Example 22:Synthesis of 1,11-O-benzylidene-1,7,11-trideacetylpyripyropene A

5 1,7,11-Trideacetylpyripyropene A (1.0 g) synthesized by the process described in Example 10 was dissolved in N,N-dimethylformamide (10 mL). Pyridinium p-toluenesulfonate (2.75 g) and benzaldehyde dimethyl acetal (3.3 g) were added to the solution. The mixture was stirred at room temperature
10 for 5 hr. The reaction solution was then poured into water, and the mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with saturated brine and was dried over anhydrous sodium sulfate, and the solvent was then removed by evaporation under the reduced pressure to give a
15 crude product of 1,11-O-benzylidene-1,7,11-trideacetylpyripyropene A. The crude product was purified by chromatography on silica gel (Mega Bond Elut (Varian), acetone : chloroform = 1 : 10) to give 1,11-O-benzylidene-1,7,11-trideacetylpyripyropene A (887 mg, yield 74%). The
20 results of measurement by ESI-MS and ¹H-NMR showed that the compound was PR-93 described in Japanese Patent Application Laid-Open No. 269065/1996.

[0129]

Example 23:

25 Synthesis of 1,11-O-benzylidene-7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A

 1,11-O-benzylidene-1,7,11-trideacetylpyripyropene A (1.0 g) synthesized by the process described in Example 22 was dissolved in pyridine (2.5 mL). Chloroacetic anhydride (206
30 mg) was added to the solution at 0°C. The mixture was stirred at that temperature for 1.5 hr, and the reaction solution was then poured into water. The mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The
35 solvent was then removed by evaporation under the reduced pressure to give a crude product of 1,11-O-benzylidene-7-O-

chloroacetyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by chromatography on silica gel (Mega Bond Elut (Varian), acetone : hexane = 1 : 100) to give 1,11-O-benzylidene-7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A
5 (359 mg, yield 72%).

[0130]

ESI-MS: m/z 622 ($M+H$)⁺;

¹H-NMR (CDCl₃): δ 1.19-1.22 (1H, m), 1.25 (3H, s), 1.41 (1H, m), 1.48 (3H, s), 1.53-1.56 (1H, m), 1.70 (3H, s), 1.70-1.84
10 (3H, m), 1.95 (1H, m), 2.27 (1H, m), 2.88 (1H, d, J = 1.6 Hz), 3.49 (1H, d, J = 10.4 Hz), 3.50-3.53 (1H, m), 3.89 (1H, d, J = 10.4 Hz), 4.20 (2H, d, J = 3.2 Hz), 5.02 (1H, m), 5.12 (1H, dd, J = 5.2, 11.6 Hz), 5.54 (1H, s), 6.46 (1H, s), 7.33-7.43 (4H, m), 7.51 (2H, dd, J = 1.6, 8.0 Hz), 8.11 (1H, dt, J = 2.0, 8.0 Hz),
15 8.70 (1H, dd, J = 1.6, 4.8 Hz), 9.02 (1H, d, J = 2.0 Hz).

[0131]

Example 24:

Synthesis of 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A

1,11-O-benzylidene-7-O-chloroacetyl-1,7,11-
20 trideacetylpyripyropene A (10 mg) synthesized by the process described in Example 23 was dissolved in chloroform (1 mL) and methanol (9 mL), and 10-camphorsulfonic acid (3 mg) was added to the solution. The mixture was stirred at room temperature for 5 days. The reaction solution was then poured
25 into water, and the mixture was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product of 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A.
30 The crude product was purified by chromatography on silica gel (Mega Bond Elut (Varian), acetone : chloroform = 1 : 10) to give 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A (8 mg, yield 100%). ESI-MS data and ¹H-NMR data of the compound were in agreement with those of the compound produced in
35 Example 19.

[0132]

Example 25:Synthesis of 1,7,11-trideacetyl-1,11-O-p-methoxybenzylidenepyripyropene A

1,7,11-Trideacetylpyripyropene A (1.0 g) synthesized by
5 the process described in Example 10 was dissolved in N,N-
dimethylformamide (22 mL). Pyridinium p-toluenesulfonate
(2.76 g) and p-methoxybenzaldehyde dimethyl acetal (0.4 g)
were added to the solution. The mixture was stirred at room
10 temperature for 4 hr. The reaction solution was then poured into
water, and the mixture was extracted with ethyl acetate. The
ethyl acetate layer was washed with saturated brine and was
dried over anhydrous sodium sulfate. The solvent was removed
by evaporation under the reduced pressure to give a crude
15 product of 1,7,11-trideacetyl-1,11-O-p-
methoxybenzylidenepyripyropene A. The crude product was
purified by chromatography on silica gel (Mega Bond Elut
(Varian), acetone : chloroform = 1 : 10) to give 1,7,11-
trideacetyl-1,11-O-p-methoxybenzylidenepyripyropene A (520
20 mg, yield 41%). The results of measurement by ESI-MS and
¹H-NMR showed that the compound was PR-124 described in
Japanese Patent Application Laid-Open No. 269065/1996.

[0133]

Example 26:Synthesis of 7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-p-methoxybenzylidenepyripyropene A

1,7,11-Trideacetyl-1,11-O-p-methoxybenzylidene-
pyripyropene A (100 mg) synthesized by the process described
in Example 25 was dissolved in tetrahydrofuran (2 mL).
Triethylamine (50 mg) and chloroacetic anhydride (60 mg) were
30 added to the solution. The mixture was stirred at room
temperature for 2.5 hr, and the reaction solution was then
poured into water and was extracted with chloroform. The
chloroform layer was washed with saturated brine and was dried
over anhydrous sodium sulfate. The solvent was then removed
35 by evaporation under the reduced pressure to give a crude
product of 7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-p-

methoxybenzylidene-pyripyropene A. The crude product was recrystallied from methanol and ethyl acetate, and the recrystallized product was further purified by chromatography on silica gel (Mega Bond Elut (Varian), acetone : chloroform = 1 : 30) to give 7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-p-methoxybenzylidenepyripyropene A (83 mg, yield 75%).

[0134]

ESI-MS: m/z 652 ($M+H$)⁺;

¹H-NMR (CDCl₃): δ 1.20 (1H, m), 1.24 (3H, s), 1.40 (1H, m), 1.47 (3H, s), 1.54 (1H, d, $J = 3.6$ Hz), 1.72 (3H, s), 1.66-1.80 (3H, m), 1.93 (1H, m), 2.26 (1H, m), 2.87 (1H, d, $J = 2.0$ Hz), 3.47 (1H, d, $J = 10.0$ Hz), 3.47-3.51 (1H, m), 3.80 (3H, s), 3.87 (1H, d, $J = 10.4$ Hz), 4.20 (2H, d, $J = 3.2$ Hz), 5.01 (1H, m), 5.12 (1H, dd, $J = 5.6, 11.6$ Hz), 5.50 (1H, s), 6.46 (1H, s), 6.90 (2H, d, $J = 8.8$ Hz), 7.41 (1H, dd, $J = 4.8, 12.0$ Hz), 7.43 (2H, d, $J = 8.8$ Hz), 8.10 (1H, dt, $J = 2.0, 8.4$ Hz), 8.70 (1H, dd, $J = 2.4, 4.8$ Hz), 9.02 (1H, d, $J = 2.0$ Hz).

[0135]

Example 27:

Synthesis of 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A
7-O-chloroacetyl-1,7,11-trideacetyl-1,11-O-p-methoxybenzylidenepyripyropene A (30 mg) synthesized by the process described in Example 26 was dissolved in chloroform (5 mL) and methanol (1 mL), and 10-camphorsulfonic acid (3 mg) was added to the solution. The mixture was stirred at room temperature for 5 days, and the reaction solution was then poured into water and was extracted with chloroform. The chloroform layer was washed with saturated brine and was dried over anhydrous sodium sulfate. The solvent was then removed by evaporation under the reduced pressure to give a crude product of 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A. The crude product was purified by chromatography on silica gel (Mega Bond Elut (Varian), acetone : chloroform = 1 : 5) to give 7-O-chloroacetyl-1,7,11-trideacetylpyripyropene A (14 mg, yield 69%). ESI-MS data and ¹H-NMR data of the compound were in agreement with those of the compound produced in Example 19.


20375-1007

56

CLAIMS:

1. A process for producing compound C represented by formula C:


[Chemical formula 1]



- 5 wherein R' represents cyclic C₃₋₆ alkylcarbonyl, the process comprising the steps of:

(a₁) hydrolyzing acetyl at the 7-position of compound A1 represented by formula A1:

[Chemical formula 2]




wherein Ac represents acetyl,

20375-1007

57

with a base to selectively deacylate compound A1, then protecting hydroxyl at the 7-position to give compound B1 represented by formula B1:

[Chemical formula 3]



5 wherein

Ac is as defined above,


R_{1a} represents group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered
 10 heterocyclic group,

then hydrolyzing acetyl at the 1- and 11-positions of compound B1 with a base to deacylate compound B1 and thus to give compound Fa represented by formula Fa:

20375-1007

58


[Chemical formula 4]



wherein R_{1a} is as defined above, or

(a₂) hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A1 or
 5 compound A4' represented by formula A4':

[Chemical formula 5]




wherein A₁', A₇', and A₁₁', which may be the same or different,
 represent acetyl or propionyl, provided that A₁', A₇', and A₁₁' do not simultaneously
 10 represent acetyl,

with a base to deacylate compound A1 or A4' and then protecting
 hydroxyl at the 1- and 11-positions to give compound D represented by formula D:

20375-1007

59


[Chemical formula 6]




wherein two R_2 s together represent a group selected from groups represented by formulae D-1, D-2, D-3, and D-4:

5


[Chemical formula 7]




D-1



D-2



D-3



D-4

wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X s, which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5,


10

then protecting hydroxyl at the 7-position of compound D to give compound E represented by formula E:

20375-1007

60

[Chemical formula 8]




wherein

R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, and

R_2 is as defined above,

and further removing the protective groups at the 1- and 11-positions of compound E to give compound Fb represented by formula Fb:

[Chemical formula 9]




wherein R_{1b} is as defined above, and

20375-1007

61

(b) acylating hydroxyl at the 1- and 11-positions of compound Fa or Fb with an acylating agent corresponding to contemplated R' to give compound B2a or B2b represented by formula B2a or B2b:

[Chemical formula 10]



5

wherein R_{1a}, R_{1b}, and R' are as defined above,


and then removing the protective group at the 7-position of compound B2a or compound B2b.

2. The process according to claim 1, wherein R_{1b} represents acetyl;
 10 chloroacetyl; or group -SiR₃R₄R₅ optionally substituted by halogen atom wherein R₃, R₄, and R₅ each independently represent straight chain or branched chain C₁₋₆ alkyl or phenyl.
3. The process according to claim 1, wherein two R₂s together represent a group represented by formula D-1 or D-2:

20375-1007

62

[Chemical formula 11]




wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X_s , which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0

5 to 5.

4. A process for producing compound C represented by formula C:

[Chemical formula 23]




wherein R' represents cyclic C_{3-6} alkylcarbonyl, the process comprising:

10 acylating hydroxyl at the 1- and 11-positions of compound Fb represented by formula Fb:

20375-1007

63

[Chemical formula 24]




wherein

- R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group,

with an acylating agent corresponding to R' to give compound B2b represented by formula B2b:

10

[Chemical formula 25]



wherein R_{1b} and R' are as defined above,


20375-1007

64

and then removing the protective group at the 7-position of compound B2b.

5. A process for producing compound C represented by formula C:

[Chemical formula 26]




5

wherein R' represents cyclic C₃₋₆ alkylcarbonyl, the process comprising:

hydrolyzing acyl at the 1-, 7-, and 11-positions of compound A4 represented by formula A4:

[Chemical formula 27]



10


wherein A₁, A₇, and A₁₁, which may be the same or different, represent acetyl or propionyl,

20375-1007

65


with a base to deacylate compound A4, then protecting hydroxyl at the 1- and 11-positions to give compound D represented by formula D:

[Chemical formula 28]




- 5 wherein two R_2 s together represent a group selected from groups represented by formulae D-1, D-2, D-3, and D-4:


[Chemical formula 29]




D-1



D-2



D-3



D-4


- 10 wherein Y_1 represents a hydrogen atom or C_{1-4} alkyl; X s, which may be the same or different, represent a hydrogen atom, C_{1-4} alkoxy, or nitro; and n is 0 to 5,

then protecting hydroxyl at the 7-position of compound D to give compound E represented by formula E:

20375-1007

66

[Chemical formula 30]




wherein

- R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent
- 5 straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, and

 R_2 is as defined above,

- and further removing the protective groups at the 1- and 11-positions of
- 10 compound E to give compound Fb represented by formula Fb:

[Chemical formula 31]


wherein R_{1b} is as defined above,

20375-1007

67

then acylating hydroxyl at the 1- and 11-positions of compound Fb with an acylating agent corresponding to R' to give compound B2b represented by B2b:

[Chemical formula 32]



5


wherein R_{1b} and R' are as defined above,

and then removing the protective group at the 7-position of compound B2b.

6.

A compound represented by formula B2b:

[Chemical formula 33]



10

wherein


20375-1007

68

R_{1b} represents acetyl; chloroacetyl; group $-SiR_3R_4R_5$ optionally substituted by halogen atom wherein R_3 , R_4 , and R_5 each independently represent straight chain or branched chain C_{1-6} alkyl or phenyl; or a saturated or unsaturated five- or six-membered heterocyclic group, and

5

R' represents cyclic C_{3-6} alkylcarbonyl.



C