



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2035384 C 2003/07/08

(11)(21) 2 035 384

(12) BREVET CANADIEN
CANADIAN PATENT

(13) C

(22) Date de dépôt/Filing Date: 1991/01/31

(41) Mise à la disp. pub./Open to Public Insp.: 1991/08/02

(45) Date de délivrance/Issue Date: 2003/07/08

(30) Priorités/Priorities: 1990/02/01 (P 40 02 897.6) DE;
1990/02/09 (P 40 03 880.7) DE

(51) Cl.Int.⁵/Int.Cl.⁵ C12N 15/13, G01N 33/53

(72) Inventeurs/Inventors:

LITTLE, MELVYN, DE;
BREITLING, FRANK BERTHOLD, DE;
SEEHAUS, THOMAS, DE;
DUBEL, STEFAN, DE;
KLEWINGHAUS, IRIS, DE

(73) Propriétaire/Owner:

DADE BEHRING MARBURG GMBH, DE

(74) Agent: BERESKIN & PARR

(54) Titre : PREPARATION ET UTILISATION DE BANQUES GENOMIQUES D'ANTICORPS SYNTHETIQUES
("BIBLIOTHEQUES D'ANTICORPS SYNTHETIQUES")

(54) Title: PREPARATION AND USE OF GENE BANKS OF SYNTHETIC HUMAN ANTIBODIES ("SYNTHETIC HUMAN-ANTIBODY LIBRARIES")

(57) Abrégé/Abstract:

The invention relates to the preparation and use of gene banks of synthetic human antibodies (huAb) or parts of antibodies which contain the antigen-binding domain. Starting from a huAb framework in a suitable vector, the hypervariable regions of the antibody cDNA are formed by almost "randomly" combined oligonucleotides. Relatively conserved amino acids in the hypervariable regions have here been taken account of in the choice of appropriate nucleotides during the oligonucleotide synthesis and the ratio of the nucleotides used is likewise chosen such that a nonsense codon is to be expected at most in every 89th position. Expression of this synthetic huAb cDNA in microbial expression systems, e.g. in E. coli in the vector pFMT which is described below, thus makes a synthetic huAb library with a comprehensive repertoire for screening using selected antigens available in vitro.

BEHRINGWERKE AKTIENGESELLSCHAFT HOE 90/B 006J - Ma 834
Dr. Lp/rd

Abstract of the disclosure

5 Preparation and use of gene banks of synthetic human
 antibodies ("synthetic human-antibody libraries")

The invention relates to the preparation and use of gene banks of synthetic human antibodies (huAb) or parts of antibodies which contain the antigen-binding domain.

10 Starting from a huAb framework in a suitable vector, the hypervariable regions of the antibody cDNA are formed by almost "randomly" combined oligonucleotides. Relatively conserved amino acids in the hypervariable regions have here been taken account of in the choice of appropriate 15 nucleotides during the oligonucleotide synthesis and the ratio of the nucleotides used is likewise chosen such that a nonsense codon is to be expected at most in every 89th position. Expression of this synthetic huAb cDNA in microbial expression systems, e.g. in E. coli in the vector pFMT which is described below, thus makes a 20 synthetic huAb library with a comprehensive repertoire for screening using selected antigens available in vitro.

5 Preparation and use of gene banks of synthetic human
 antibodies ("synthetic human-antibody libraries")

The invention relates to the preparation and use of gene banks of synthetic human antibodies (huAb) or parts of antibodies which contain the antigen-binding domain.
10 Starting from a huAb framework in a suitable vector, the hypervariable regions of the antibody cDNA are formed by almost "randomly" combined oligonucleotides. Relatively conserved amino acids in the hypervariable regions have here been taken account of in the choice of appropriate nucleotides during the oligonucleotide synthesis and the ratio of the nucleotides used is likewise chosen such that a nonsense codon is to be expected at most in every 89th position. Expression of this synthetic huAb cDNA in microbial expression systems, e.g. in *E. coli* in the vector pFMT which is described below, thus makes a
15 synthetic huAb library with a comprehensive repertoire for screening using selected antigens available in vitro.
20

The human or mammalian immune system comprises an estimated number of between 10^6 and 10^8 different antibodies.
25 This number of antibodies seems to be sufficient to cause an immune reaction of the body both against all naturally occurring antigens and against artificial antigens. If it is furthermore taken into account that often different antibodies react with the same antigen, the repertoire of antibodies that are really different would rather be in
30 the region from 10^6 to 10^7 .

Up to now specific antibodies have always been obtained starting from an immunization with the particular antigen, for example injection of the antigen into the body or in vitro incubation of spleen cells with this antigen.
35 In the case of polyclonal antibodies, the immunoglobulins

can then be isolated from the serum and the specific antibodies can be isolated therefrom, e.g. by absorption methods. Monoclonal antibodies are isolated from the cell supernatants or from the cell lysate of spleen tumor 5 cells (hybridoma cells) which have been fused with individual B lymphocytes and cloned. The abovementioned methods are unsuitable in particular for the preparation of specific human antibodies or human monoclonal anti-bodies.

10 The present invention therefore has the object of developing a generally usable method for generating specific human monoclonal antibodies (huMAbs) or parts of antibodies, which contain synthetic hypervariable domains.

15 It has now been found that by using almost randomly synthesized oligonucleotides coding for the three hypervariable regions of each variable part of heavy or light chains (called CDR1, 2 and 3, CDR meaning complementary determining region) synthetic human gene banks can be 20 generated. The synthesized antibody DNA was then preferably ligated into an antibody expression vector especially constructed for this purpose, namely the vector pFMT, preferably after amplification using the polymerase chain reaction (PCR).

25 The oligonucleotides which are used for the synthesis of the variable domains of heavy and light chains are compiled in Tab. 1. Set A here contains fewer limitations than set B. The limitations below the synthesis of the hypervariable regions (see CDR regions in Tab. 4) being 30 random were made in H3, H4, H6, L2, L3 and L5 in set A in order firstly to allow for positions in the sequence for certain conserved amino acids, secondly to reduce the number of stop codons, and thirdly to incorporate a new restriction site.

35 (a) In order to reduce the probability of the stop codon

occurring, only half the amount of the three other nucleotides was allowed for T at the first position of each codon and A was omitted at the third position of each codon, in each case. As a statistical average, only every 89th codon will thus be a stop codon.

5

- (b) For the 2nd codon in the CDR1 region of the light chain, only those nucleotides were allowed which code for the amino acids V, A or G.
- 10 (c) Likewise, only those combinations coding for V, I or M were allowed for codon No. 10 in the CDR1 region of the light chain and for codon No. 4 in the CDR1 region of the heavy chains.
- 15 (d) In the CDR3 region of the light chain, only those nucleotides coding for the amino acid glutamine were allowed for codon No. 1.
- (e) In the CDR2 region of the heavy chain, only those nucleotides coding for the amino acid tyrosine were allowed for codon No. 11.
- 20 (f) An A was advantageously incorporated at the third position of the last codon in the CDR2 region of the heavy chain in order to introduce a restriction site for MluI.

25

The random nature of these oligonucleotides was preferably limited even further in those positions where predominantly one or few amino acids occur (set B in Tab. 1, the limitations here are based on the tables by Kabat et al. (1987), Sequences of Proteins of Immunological Interest-U.S. Dept. of Health and Human Services, U.S. Government Printing Offices). A list of the corresponding nucleotides and brief notes on the codon combinations are compiled in Tab. 1 and in the notes for Tab. 1.

30

After ligation of equimolar amounts of the oligonucleotides H1 to H7 and L1 to L5, these are ligated into the pretreated expression vector pFMT. Preferably, a PCR step using the primers H1 and H8, or L1 and L6 should be carried out beforehand in order to amplify the amount of DNA. After producing suitable restriction sites at the ends of the antibody DNA using appropriate restriction enzymes, the DNA is ligated into the antibody expression vector pFMT in the same manner as above (see examples).

The expression pFMT makes possible the expression of antibody cDNA and the subsequent secretion of the expression products in bacteria (E.coli). The antibody operon of the plasmid contains the sequences of the variable parts of both the heavy and light chain of an antibody. Suitable leader sequences from the amino terminal part of a bacterial protein makes secretion of the antibody parts possible. The leader sequences are cleaved off by a bacterial enzyme during the secretion. During the secretion of the antibody cDNA products, the light and heavy chains of the antibody (with or without an adjacent constant domain) become associated. This results in the formation of an antibody or antibody fragment which, in either case, contains a functional antigen binding site. Similar constructs for individual antibodies have also been described by other authors (Better et al. (1988), Science 240, 1041, and Skerrras & Plückthun (1988), Science 240, 1038).

In the synthetic human-antibody library formed by the expression in, for example, E. coli, the desired human antibodies or antibody parts are found by screening bacterial clones using the selected antigen. In a preferred embodiment, an additional sequence which codes for a marker peptide, for example a TAG sequence, is incorporated so that the expression products can be detected in a simple way using established monoclonal antibodies against the marker peptide (Wehland et al. (1984), EMBO J. 3, 1295).

The abovementioned exemplary formulations and the examples below shall be understood as illustrating but not restricting the invention.

5 The invention therefore relates to gene banks of synthetic huAb or antigen-binding parts thereof, obtained by means of (1) cDNA for the hypervariable regions generated on a random basis, where the random sequences are limited by clauses (a) to (e) set A or in accordance with Tab. 1, set B, (2) preferably a subsequent 10 amplification step of these random sequences and (3) ligation of the said cDNA into a suitable expression vector, preferably pFMT, an additional coding sequence for a marker peptide being incorporated in a preferred embodiment.

15 The invention also relates to a process for the separation of the abovementioned gene banks, and the process and the use thereof for the isolation of clones which secrete specific antibodies or antigen-binding parts thereof.

20 Finally, the invention is explained in detail in the examples and contained in the patent claims.

Examples:

Example 1: Preparation of an antibody expression vector

25 The plasmid pKK233-2 (Amann and Brosius, (1985) Gene 40, and Straus and Gilbert (1985) Proc. Natl. Acad. Sci. 82, 2014) was chosen as base vector for the construction of the antibody expression vector (Fig. 1).

30 Before the incorporation of the antibody operon, the plasmid was cut with SalI and BamHI, the ends were filled in with Klenow polymerase and ligated. By doing so, the two restriction sites and the DNA between them were

deleted.

Additionally, the plasmid was cleaved with HindIII, the ends were filled in with Klenow polymerase and ligated using BamHI linkers. By this procedure, the HindIII restriction site was removed and a BamHI site inserted. The antibody DNA was inserted into this modified plasmid. A simplified structure of the antibody operon coding for a dicistronic antibody mRNA is shown in Tab. 2. In order to make possible the secretion of the antibody, the leader sequence of the bacterial enzyme pectate lyase was used. The leader sequence of this enzyme has already been used for the expression and secretion of a chimeric murine/human antibody (Fab fragment, Better et al., loc. cit.), and of the variable region of a "humanized" antibody (Ward et al., loc. cit.; Huse et al., loc. cit.). DNA for the first leader sequence (P_1 , upstream of the heavy chain), and the sequence for a second ribosome binding site (RBS) and a second leader sequence (P_2 upstream of the light chain) were synthesized from several oligonucleotides (Tab. 3).

Antibody cDNAs which code for the variable regions of the heavy and light chains of a human antibody (HuVhlys or HuVlllys; Riechmann et al., (1988) J. Mol. Biol. 203, 825) were obtained from Dr. G. Winter (Cambridge, UK). The restriction sites HindIII (HuVhlys) and EcoRV (HuVlllys) were introduced to make possible the insertion of the antibody cDNA into the expression vector. Further restriction sites for BanII (HuVhlys) and BstEII or KpnI (HuVlllys) were introduced to exchange hypervariable regions en bloc. At the end of the HuVhlys cDNA sequence a stop signal was incorporated. A BanII site in the light chain was removed. These alterations were carried out by means of site directed mutagenesis in the bacteriophage M13mp18 (Zoller and Smith, Meth. Enzymol. 100, 468-500). The sequence of the completed antibody DNA is shown in Tab. 4.

For the insertion of the leader sequence P_1 (Tab. 3) the modified plasmid pKK233-2 was digested using the restriction enzymes NcoI and PstI, and P_1 was inserted in between these sites (pKK233-2- P_1). Further cloning steps, 5 apart from the last step, were carried out using the plasmid pUC18. The reason is that the presence of individual parts of the antibody operon in the expression vector adversely influences the growth of the bacterial host.

10 Before the cloning in pUC18, its BamHI restriction site had to be removed. After digesting with BamHI, the single-stranded ends were filled in using the Klenow fragment and were religated. This modified plasmid was then digested using PstI and HindIII, and P_2 plus RBS was 15 ligated in between the restriction sites (pUC18- P_2). During this process, the original HindIII restriction site of the plasmid disappears and a new HindIII restriction site is incorporated. pUC18- P_2 was then digested using PstI and HindIII, and the DNA of the heavy chain 20 (PstI-HindIII insert from M13) was ligated into these two sites (pUC18-HP₂). This plasmid was then digested using EcoRV and BamHI, and the DNA of the light chain (EcoRV- 25 BamHI insert from M13) was ligated in (pUC18-HP₂L).

The PstI-BamHI insert was then recloned in pUC18 after 25 the restriction sites for HindIII, BanII and KpnI therein had previously been removed. The HindIII restriction site was removed as above for pKK233-2, the religation taking place without an insertion of BamHI linkers, however. Subsequently, the resulting plasmid was digested using 30 SmaI and BanII, and, after filling in the protruding ends by means of T4 DNA polymerase, religated. The insertion of the PstI-BamHI restriction fragment results in pUC-HP₂L. In a preferred embodiment, a Tag sequence was additionally inserted in the BanII and HindIII restriction sites (Tab. 3). The Tag sequence encodes the recognition 35 sequence Glu-Gly-Glu-Glu-Phe of the monoclonal antibody Y1 1/2 (Wehland et al., (1984), EMBO J. 3, 1295). Because

of this peptide marker the expression product of the resulting plasmid pUC-HP₂L is readily detectable.

For the insertion of HP₂L or HTP₂L in the expression vector, the two plasmids were cut using PstI and BamHI, 5 and the PstI-BamHI HP₂L insert from pUC-HP₂L or the HTP₂L insert from pUC-HTP₂L was ligated into the modified plasmid pKK233-2-P₁ into these two restriction sites. A diagrammatic representation of the completed expression vector pFMT is shown in Tab. 5.

10 **Example 2: Synthesis of antibody DNA containing random sequences in hypervariable regions**

The synthesized oligonucleotides for the synthesis of the variable parts of antibody DNA are compiled in Tab. 1. For the synthesis of the hypervariable regions almost 15 random nucleotide sequences were used. Limitations on the random nature are illustrated in Tab. 1. Two different sets of oligonucleotides were synthesized. In set A the hypervariable regions are predominantly random apart from those few positions where almost exclusively certain 20 amino acids occur. In set B, the random nature of the nucleotide sequences in those positions where predominantly one or few amino acids occur was additionally limited.

25 The oligonucleotides were purified by HPLC chromatography or polyacrylamide gel electrophoresis, and then 5'-phosphorylated.

Example 3: Ligation of the synthetic oligonucleotides

The oligonucleotides in Tab. 1 were ligated together 30 stepwise on an antibody DNA template. For this purpose, large amounts (about 1 mg) of single-stranded M13mp=18 DNA containing the antibody DNA inserts were isolated. In order to separate the antibody DNA from the vector, the inserts were made double-stranded on the two ends using

two appropriate oligonucleotides and were digested using the enzymes PstI and HindIII (heavy chain) or using EcoRV and BamHI (light chain). The antibody DNA was then purified using agar gel electrophoresis.

5 On these DNA templates, first only three oligonucleotides were ligated: H1, pH2 and pH3 (heavy chain), and L1, pL2 and pL3 (light chain), H1 and L1 having been marked first with ^{32}P at their 5' end ("p" designates 5'-phosphorylated). Amounts of 100pmol of each oligonucleotide were used. The hybridized oligonucleotides were purified on 2% agarose gels and analyzed on a sequencing gel. The amount was determined by a radioactivity measurement. Equimolar amounts of pH4 and pH5 (heavy chain), and pL4 and pL5 (light chain) were then ligated onto the already ligated 10 oligonucleotides on each particular template. These DNAs were then purified as in the preceding step and the procedure was repeated up to the purification step, using equimolar amounts of pH6 and pH7. Finally, the ligated 15 oligonucleotides were purified by means of a denaturing polyacrylamide gel and preferably amplified using the polymerase chain reaction (PCR). Alternatively or in order to avoid losses caused by the last purification step, the oligonucleotides were amplified using PCR directly after the last ligation step. The primers H1 and 20 H8 (heavy chain), and L1 and L6 (light chain) were used under standard conditions for the PCR. Amplified template DNA was digested selectively using KpnI (light chain) or using AluI (heavy chain). Where appropriate, a second 25 amplification step using the PCR was subsequently carried out.

Example 4: Insertion of the antibody DNA into the expression plasmid

35 The synthesized antibody DNA was cut using the restriction enzymes PstI and BanIII (heavy chain), and BstEII and KpnI (light chain). The bands having the expected molecular weight were purified by agar gel

electrophoresis, precipitated using ethanol and then, in two steps (first the DNA of the light chain and then the DNA of the heavy chain), ligated into the pUC-HP₂L (see above) which had been cut and purified in the same way. 5 The HP₂L insert was then ligated into the restriction sites PstI and BamHI of the plasmids pKK233-2-P₁ (see Example 1). An analogous way was used for the HTP₂L fragment. The antibody library is therefore established 10 in the antibody expression plasmid (Tab. 6). The reason for intermediate cloning in pUC is that the presence of individual parts of the antibody operon in the expression vector has an adverse influence on the growth of the bacterial host (see above also).

15 **Example 5: Expression and screening of antibodies in
E. coli**

Competent E. coli are transfected with pFMT plasmids containing the inserted antibody-DNA library, grown on agarose plates and then incubated using nitrocellulose filters coated with the desired antigen. After removing 20 non-specifically bound antibodies, the active clones are identified with a labeled antibody against the human immunoglobulins secreted from E. coli. In the preferred embodiment, the antibody YL 1/2 which is directed against the Tag sequence is used for this purpose.

Legend for Fig. 1:

Restriction map of the expression vector pKK233-2 (Amann and Brosius, loc. cit.).

Ptrc denotes hybrid tryptophan lac promoter

5 RBS denotes ribosome binding site

rrnB denotes ribosomal RNA B operon

5S denotes gene for 5S RNA

Before cloning antibody DNA in the expression vector, the following alterations were carried out:

- 10 1) The SalI and EcoRI restriction sites were removed together with the DNA between them.
- 2) The HindIII restriction site was converted to a BamHI restriction site.

TAB. 1

Oligonucleotide for the synthesis of a library of anti-body DNA (variable parts)

Set A

H1 5' CCAGGTCCAAC TGCA GGAG AGCG GTCC AGGT CTT GTGAGACCTAG3'

H2 5' CCAGACCC TGAGC CTGAC CTGAC CCGT G3'

H3 5' TGTCTGGCTTCACCTTCAGC | T1/2 TT | CTTT1/2TTTGGGTGCCAGCCACCTGGAC3'
 C C | C CC | A CC CC
 | A AG | G AA AG
 | G G | 3 GG G

H4 5' GAGGTCTTGACTGGATTGGT | T1/2TT | TAT | T1/2TT | T1/2TACCGGTGACAATGCTGGTAGAC3'
 | C CC | | C CC | C C
 | A AG | | A AG | A A
 | G G | 10 | G G | 5 G G

H5 5' ACCAGCAAGAACCAAGTTCA GGCTGCGTCTCAGCAGCGTGACAGC3'

H6 5' CGCCGACACCGCGGTCTACTACTGTGCCGC | T1/2 TT | TGGGGTCAGGGCT3'
 | C CC |
 | A AG |
 | G G | 10

H7 5' CCCTCGTCACAGTCTCCTCA3'

H8 5' CTGTGACQAGGCTGCCCTGACCCCA3'

L1 5' QGCCCAAGCGTGGGTGACAGG3'

L2 5' GTGACCATCACCTGTT1/2TTGTT | T1/2TT | CTTT1/2TTTGGTAACAGCAGAACCCAGGT3'
 C CC CC | C CC | A CC CC
 A AG GA | A AG | G AA AG
 G G G | G | 7 GG G

L3 5' AAGGCTCCAAAGCTGCTGATCTAC | T1/2TT | GGTGTGCCAAGCCGTTTCAGCGGTAGCGGT3'
 | C CC |
 | A AG |
 | G G | 7

L4 5' AGCGGTACGGACTTCACCTTCACCATCAGCAGCCTCCAGCCAGAGGAC3'

L5 5' ATCGCCACCTACTACTGCCAG | T1/2TT | TTGGCCAAAGGTAC3'
 | C CC |
 | A AG |
 | G G | 8

L6 5' CCACCTTGGTACCTTGGCCGAA3'

Set B

H1, H2, H5, H7, H8 and L1, L4 and L6 are identical to those in set A.

H3 5' TGTCTGGCTTCACCTTCAGC AC10%T95% TT20%C TC20%G T5% TA20% T2% T5% T70%
 C C GA45%A5% C10% GG80% A85% G80% C28%A75%G30%
 G45% A70% G10% A50%G20%
 G20%

TGGGTGCCAGCCACCTGGAC3'

H4 5' GAGGTCTTGAGTGGATTGGT T14.5%TT AT90%C T14.5%TT T5% T10%T70% T14.5%TT
 C28.5%CC G10% C28.5%CC C70%C80%G30% C28.5%CC
 A28.5%AG A28.5%AG A20%A10% A28.5%AG
 G28.5%G G28.5%G G5% G28.5%G

AAT A15%A20%C ACT AT16%A A70%C10%C TAT A80%C20%C CCC AC10%C T40%TC C5% T5% G
 GG G85%G80% GA C80% G30%A70% A G20%A60% GAA A40%A C20% A90%A90%
 G A4% G20% G20% G50% G40% G5% G5%

A20%A10%T CGCGTQACAATGCTGGTAGAC3'
 G80%G90%

H6 5' CGCCGACACCGCGGTCTACTACTGTGCGCGC T1/2 TT GC25%C TATTGGGTCAGGGCT3'
 C CC A75%
 A AG
 G G8

L2 5' GTGACCATCACCTGT CAA T30%CG AGT C75%AA T30%T10%C C10%T40%T90%
 AG G70% A10% C10%C30% A50%C10%A10%
 G15% A30%A60% G40%A10%
 G30% G40%

A70%T20%T70% C5% T5% T90% T90%T10%C C70%TA20% AC40%T TGGTAACAGCAGAAGCCAGGT3'
 G30%A40%A30% A90%C20%A10% C2% C5% A20% G80% QA40%
 G40% G5% A40% A6% A85% G10% G20%
 G35% G2%

L3 5' AAGGCTCCAAAGCTGCTGATCTAC T14.5%TT A40%T20%T AC5% C AC20%T70% CTA
 C28.5%CC G60%C70% A10% A60%A30% G
 A28.5%AG A7% G85% G20%
 G28.5%G G3%

C20%C70%C20% T70%CT GGTGTGCCAAGCCGTTTCAGCGGTAGCGGT3'
 G80%A30%A80% C15%
 A15%

L5 5' ATCGCCACCTACTACTGCTCT10%A CT20%T10% T60%C30%G T35%T5% T T15%T5% C
 A90% A80%G90% A10%G70% C5% C20% C10%C20%
 G30% A40%A50% A60%A75%
 G20%G25% G15%



T14.5%TT CT15%C90% T14.5%TT ACGTTGGCCAAGGTAC
 C28.5%CC C70%A10% C28.5%CC
 A28.5%AG A15% A28.5%AG
 G28.5%G G28.5%G

Notes for Tab. 1

The random nature of the oligonucleotides of set B was limited in a manner which generates approximately the relevant amount of frequent amino acids in each position of the hypervariable regions (in accordance with the tables of Kabat et al, loc. cit.). In this strategy the number of expected stop codons was also reduced even further. In contrast with the oligonucleotides in set A, a restriction site for MluI was not introduced.

TAB. 2CONSTRUCTION OF THE VECTOR PFM1 FOR THE EXPRESSION AND
SECRETION OF ANTIBODIES IN BACTERIA

DNA OF THE VARIABLE DOMAIN OF A HUMAN LYSOZYME ANTIBODY



P/O: promoter/operator, RBS: ribosome binding site, P2: leader sequence of pectate lyase, VH: variable domain of the heavy chain, VL: variable domain of the light chain

TAB. 3

Sequences of the leader sequences P1 and P2 in the antibody operon, and of the Tag sequences

P1

Leader sequence of pectate lyase (P1)

M K Y L L P T A A A G L L L A A Q P A M A Q V Q L Q
 CATGAAATAACCTCTTGCCTACGGCAGCCGCTGGCTTGCTGGCAGCTCAGCCGGCATGGCGCAAGTTCAGCTGCA(G)
 PstI

P2

RBS

Leader sequence of pectate lyase (P2)

M K Y L L P T A A A
 (C)TGCAGCCAAGCTTGAATTCAAAAGAGGAGAAATTAACTCCATGAAGTACTTACTGCCGACCGCTGCAGCG
 PstI HindIII
 G L L L L A A Q P A M A D I
 GGTCTCCTGCTGTTGGCGCTCAGCCGGCTATGGCTGATATCGGATCCCAGCT
 EcoRV BamHI

The nucleotides in parentheses are the adjacent nucleotides of the plasmid

The leader sequences were synthesized by hybridization of the following oligonucleotides.

P1

- a. 5'CATGAAATAACCTCTTGCCTACGGCAGCCGCTGGCTTG3'
- b. 5'TTAACTCCATGAAGTACTTACTGCCGACCGCTGC3'
- c. 3'ACGTCGGTTCGAACTTAAGTTAACTCCCTTAATTGAGGTACTTCATGAATGACGGCTGGCGACGCCGCCAGAGGA
 CGACAACCGCCGAGTCGGCCGATAACCGACTATAGCCTAGGTGCA5'
- d. 5'GCTCAGCCGGCTATGGCTGATATCGGATCC3'
- e. 5'GCGGGTCTCCTGCTGTTGGCG3'

The Tag sequences were synthesized by hybridization of the following sequences:

- a. 5'CCTTAGTCACAGTATCCTCAGAAGGTGAAGAATTCTA3'
- b. 5'AGCTTACAATTCTCACCTCTGAGGACTGTGACTAAGGAGCC3'

TAB. 4

Nucleotide sequences of antibody DNA


a) Heavy chain (variable domain), HuVhlys HindIII.....

1 10
 G V H S Q V Q L Q E S G P G L V R
 CTCTCCACAGGTGTCCACTCCAGGTCCAACTGCAGGAGAGCGGTCCAGGTCTTGAGA
 PstI
 20 30 CDR1
 P S Q T L S L T C T V S G F T F S /G//Y//G/
 CCTAGCCAGACCCCTGAGCCTGACCTGCACCGTGTCTGGCTTCACCTTCAGCGGCTATGGT
 BspMI
 50
 /V /N /W V R Q P P G R G L E W I G /M/ I / W / G /
 GTAAACTGGGTGAGACAGCCACCTGGACGAGGTCTTGAGTGGATTGGAATGATTGGGGT
 CDR2 60 70
 /D /G /N /T /D /Y /N /S /A /L /K /S R V T M L V D T
 GATGGAAACACAGACTATAATTCAGCTCTCAAATCCAGAGTGACAATGCTGGTAGACACC
 80 90
 S K N Q F S L R L S S V T A A D T A V Y
 AGCAAGAACCCAGTTCAGCCTGAGACTCAGCAGCGTACAGCCGCCACACCGCGGTCTAT
 SacII
 100 CDR3 110
 Y C A R E /R /D /Y /R /L /D /Y W G Q G S L V T
 TATTGTGCAAGAGAGAGAGATTATAGGCTTGACTACTGGGGCTCCCTCGTCACA
 BanII
 V S S Stop
 GTCTCCTCATAAGCTTCTTACAACCTCTCTTCTATTCAGCTAA.....BamHI
 HindIII

b) Light chain (variable domain), Hu Vllys HindIII.....

1 10
 G V H S D I Q N T Q S P S S L S A
 CTCTCCACAGGTGTCCACTCCGATATCCAGATGACCCAGAGCCCAAGCAGCCTGAGCGCC
 EcoRV
 20 30 CDR1
 S V G D R V T I T C R / A / S / G / N / I / H / N / Y / L
 AGCGTGGGTGACAGGGTGACCATCACCTGTAGAGCCAGCGGTAAACATCCACAACACTACCTG
 BstEII
 40 50 CDR2
 /A / W Y Q Q K P G K A P K L L I Y / Y / T / T / T
 GCTTGGTACCAAGCAGAAGCCAGGTAAAGGCTCCAAAGCTGCTGATCTACTACACCACCA
 60 70
 /L / A / D G V P S R F S G S G S G T D F T F
 CTGGCTGACGGTGTGCCAAGCAGATTCAAGCTGCGTAGCGGTAGCGGTACCGACTTCACCTC
 80 90 CDR3
 T I S S L Q P E D I A T Y Y C / Q / H / F / W / S
 ACCATCAGCAGCCTCCAGCCAGAGGACATGCCACCTACTACTGCCAGCACTCTGGAGC
 100
 /T / P / R / T F G Q G T K V E I K R . E STOP
 ACCCCAAAGGGACGTTGGCCCAAGGTACCAAGGTGGAAATCAAACGTGAGTAGAATTAAAC
 KpnI
 TTTGCTTCTCAGTTGGATCC
 BamHI

The Antibody Expression Plasmid pFMT




There is an RBS in the plasmid upstream of the heavy chain part but is not drawn in here.

2035384

- 19 -

TAB. 6

Insertion of the antibody vector pEMT



- 20 -

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A synthetic human antibody-DNA library obtained by:
 - 5 (1) synthesizing almost random sequences for the antibody hypervariable regions, wherein (a) relatively conserved amino acids in the hypervariable regions have been taken into account in the choice of appropriate nucleotides during the oligonucleotide synthesis and (b) the ratio of the nucleotides used is chosen such that a nonsense codon is to be 10 expected at most in every 89th position;
 - (2) inserting said almost random sequences for the hypervariable regions into a human antibody framework; and
 - (3) incorporating said almost random sequences combined with said human antibody framework into an expression vector.
 - 15 2. The synthetic human antibody-DNA library as claimed in claim 1, wherein the synthesized almost random sequences for the hypervariable regions are amplified before the incorporation in the expression vector.
 - 20 3. The synthetic human antibody-DNA library as claimed in claim 1, wherein said almost random sequences are incorporated into antibody variable regions derived from HuVhlys or HuVlls.
 4. The synthetic human antibody-DNA library as claimed in claim 2, - 25 wherein said almost random sequences are incorporated into antibody variable regions derived from HuVhlys or HuVlls.
 - 5. The synthetic human antibody-DNA library of claim 1 wherein the DNA is transfected into a microbial expression system.
 - 30 6. The synthetic human antibody-DNA library of claim 2 wherein the DNA is transfected into a microbial expression system.

- 21 -

7. The synthetic human antibody-DNA library of claim 3 wherein the DNA is transfected into a microbial expression system.
- 5 8. The synthetic human antibody-DNA library of claim 4 wherein the DNA is transfected into a microbial expression system.
9. The synthetic human antibody-DNA library of claim 5 wherein the microbial expression system is bacterial.
- 10 10. The synthetic human antibody-DNA library of claim 6 wherein the microbial expression system is bacterial.
11. The synthetic human antibody-DNA library of claim 7 wherein the microbial expression system is bacterial.
- 15 12. The synthetic human antibody-DNA library of claim 8 wherein the microbial expression system is bacterial.
- 20 13. The synthetic human antibody-DNA library of claim 9 wherein the bacteria are *E. coli*.
14. The synthetic human antibody-DNA library of claim 10 wherein the bacteria are *E. coli*.
- 25 15. The synthetic human antibody-DNA library of claim 11 wherein the bacteria are *E. coli*.
16. The synthetic human antibody-DNA library of claim 12 wherein the bacteria are *E. coli*.
- 30

- 22 -

17. A process for preparing a synthetic human antibody-DNA library, which comprises:

(1) synthesizing almost random sequences for the antibody hypervariable regions, wherein

5 (a) relatively conserved amino acids in the hypervariable regions have been taken into account in the choice of appropriate nucleotides during the oligonucleotide synthesis and

(b) the ratio of the nucleotides used is chosen such that a nonsense codon is to be expected at most in every 89th position;

10 (2) inserting said almost random sequences for the hypervariable regions into a human antibody framework; and

(3) incorporating said almost random sequences combined with said human antibody framework into an expression vector.

15 18. The process for preparing a synthetic human antibody-DNA library as claimed in claim 17, wherein the synthesized random sequences are amplified before incorporation in an expression vector.

19. The process for preparing a synthetic human antibody-DNA library as 20 claimed in claim 17, wherein the almost random sequences are incorporated into antibody variable regions derived from HuVhlys or HuVlls.

20. The process for preparing a synthetic human antibody-DNA library as claimed in claim 18, wherein the almost random sequences are incorporated 25 into antibody variable regions derived from HuVhlys or HuVlls.

21. The process as claimed in claim 17, wherein said expression vector further comprises DNA coding for a marker peptide and the desired clones are identified using antibodies against the marker peptide.

- 23 -

22. The process as claimed in claim 21, wherein said marker peptide is the TAG sequence and the desired clones are identified using the antibody YL 1/2.

5 23. The process as claimed in claim 18, wherein said expression vector further comprises DNA coding for a marker peptide and the desired clones are identified using antibodies against the marker peptide.

10 24. The process as claimed in claim 23, wherein said marker peptide is the TAG sequence and the desired clones are identified using the antibody YL1/2.

25. The process for preparing a synthetic human antibody-DNA library of claim 17 wherein the variable regions are incorporated into pFMT.

15 26. The process for preparing a synthetic human antibody-DNA library of claim 18 wherein the variable regions are incorporated into pFMT.

20 27. The process for preparing a synthetic human antibody-DNA library of claim 17 wherein the expression vector is transfected into a microbial expression system.

28. The process of claim 27 wherein the microbial expression system is bacterial.

25 29. The process of claim 28 wherein the bacteria are *E. coli*.

30 30. The process for preparing a synthetic human antibody-DNA library of claim 18 wherein the expression vector is transfected into a microbial expression system.

31. The process of claim 30 wherein the microbial expression system is bacterial.

- 24 -

32. The process of claim 31 wherein the bacteria are *E. coli*.

33. A process for obtaining clones secreting specific human antibodies
5 comprising the steps of:

- (a) screening synthetic human antibody-DNA libraries using specific antigens, and
- (b) isolating said clones which secrete the desired specific human antibodies,

10 wherein said specific human antibody-DNA library is obtained by:

- (1) synthesizing almost random sequences for the antibody hypervariable regions, wherein
 - (a) relatively conserved amino acids in the hypervariable regions have been taken into account in the choice of appropriate nucleotides during the 15 oligonucleotide synthesis and
 - (b) the ratio of the nucleotides used is chosen such that a nonsense codon is to be expected at most in every 89th position;
- (2) inserting said almost random sequences for the hypervariable regions into a human antibody framework; and

20 (3) incorporating said almost random sequences combined with said human antibody framework into an expression vector.

34. The process for obtaining clones secreting specific human antibodies
25 as claimed in claim 33, wherein the process of obtaining the specific human antibody-DNA library further comprises the step of amplifying the almost random sequences of the hypervariable regions before incorporation into the expression vector.

30 35. The process for obtaining clones secreting specific human antibodies as claimed in claim 33, wherein the almost random sequences are incorporated into antibody variable regions derived from HuVhlys or HuVlls.

- 25 -

36. The process for obtaining clones secreting specific human antibodies as claimed in claim 34, wherein the almost random sequences are incorporated into antibody variable regions derived from HuVhlys or HuVlls.

5

37. The process as claimed in claim 33, wherein said expression vector further comprises DNA coding for a marker peptide and the desired clones are identified using antibodies against the marker peptide.

10 38. The process as claimed in claim 37, wherein said marker peptide is the TAG sequence and the desired clones are identified using the antibody YL 1/2.

15 39. The process as claimed in claim 34, wherein said expression vector further comprises DNA coding for a marker peptide and the desired clones are identified using antibodies against the marker peptide.

20 40. The process as claimed in claim 39, wherein said marker peptide is the TAG sequence and the desired clones are identified using the antibody YL 1/2.

41. The process for obtaining clones secreting specific human antibodies of claim 33 wherein the variable regions are incorporated into pFMT.

25 42. The process for obtaining clones secreting specific human antibodies of claim 34 wherein the variable regions are incorporated into pFMT.

30 43. The process for obtaining clones secreting specific human antibodies of claim 33 wherein the expression vector is transfected into a microbial expression system.

- 26 -

44. The process of claim 43 wherein the microbial expression system is bacterial.

45. The process of claim 44 wherein the bacteria are *E. coli*.

5

46. The process for obtaining clones secreting specific human antibodies of claim 34 wherein the expression vector is transfected into a microbial expression system.

10 47. The process of claim 46 wherein the microbial expression system is bacterial.

48. The process of claim 47 wherein the bacteria are *E. coli*.

15 49. The synthetic human antibody-DNA library as claimed in claim 1 wherein the expression vector further comprises DNA coding for a marker peptide.

20 50. The synthetic human antibody-DNA library as claimed in claim 49 wherein the marker peptide is the TAG sequence.

51. The synthetic human antibody-DNA library as claimed in claim 2 wherein the expression vector further comprises DNA coding for a marker peptide.

25

52. The synthetic human antibody-DNA library as claimed in claim 51 wherein the marker peptide is the TAG sequence.

30 53. The synthetic human antibody-DNA library as claimed in claim 2 wherein the variable regions are incorporated into pFMT.

- 27 -

54. The synthetic human antibody-DNA library as claimed in claim 53
wherein the variable regions are incorporated into pFMT.

3035384




FIG. 1

By: Rogers, Breckin & Par.