

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 July 2006 (27.07.2006)

PCT

(10) International Publication Number
WO 2006/077601 A2

(51) International Patent Classification:
A61M 5/315 (2006.01)

(74) Agent: WEBB, Cynthia; Webb & Associates, P.O. Box 2189, 76121 Rehovot (IL).

(21) International Application Number:
PCT/IL2006/000096

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 24 January 2006 (24.01.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/645,605 24 January 2005 (24.01.2005) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): YEDA RESEARCH & DEVELOPMENT CO. LTD. [IL/IL]; The Weizmann Institute of Science, P.O. Box 95, 76100 Rehovot (IL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): SHAI, Yechiel [IL/IL]; Bar Simantov Street 70, 56333 Yehud (IL). COHEN, Irun, R. [IL/IL]; 11 Hankin Street, 76354 Rehovot (IL). QUINTANA, Francisco, J. [AR/AR]; Teniente B Matienzo, 1831 1H Capital Federal, 1426 (AR). GERBER, Doron [IL/IL]; 10 Kalisher Street, 46497 Herzliya (IL).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HIV-1 gp41 FUSION PEPTIDES FOR IMMUNOMODULATION

(57) Abstract: The present invention provides pharmaceutical compositions and methods for prevention or treatment of autoimmune diseases and other T cell mediated inflammatory diseases and conditions, which comprise as an active ingredient an effective quantity of a peptide derived from HIV gp41 fusion peptide domain or fragments, analogs, homologs and derivatives thereof. The invention further provides novel peptides derived from HIV gp41 fusion peptide domain, useful in the treatment of T cell mediated pathologies.



WO 2006/077601 A2

HIV-1 gp41 FUSION PEPTIDES FOR IMMUNOMODULATION**FIELD OF THE INVENTION**

The present invention provides novel uses of peptides derived from the HIV gp41 fusion peptide domain, in methods for prevention or treatment of autoimmune and other T cell-mediated pathologies which comprise administering to a subject an effective quantity of an HIV gp41 fusion peptide or fragments, homologs and derivatives thereof. Certain novel fragments of the HIV gp41 fusion peptide useful in the methods of the present invention are claimed as such.

BACKGROUND OF THE INVENTION

Human immunodeficiency virus (HIV) infection confounds the immune response. Untreated HIV infection usually leads to a state of general immunosuppression, the acquired immune deficiency syndrome (AIDS), and susceptibility to otherwise innocuous opportunistic infections. However, to establish a successful infection and replicate, the virus has to evade immune control, a task that HIV accomplishes by using a broad array of mechanisms, recently reviewed (Johnson and Desrosiers, 2002). Of particular interest is the inhibition of the CD4⁺ T-cell activity directed to HIV itself (Rosenberg *et al.*, 1997; Norris and Rosenberg, 2001); anti-HIV CD4⁺ T cells are required to establish a CD8⁺ T-cell response capable of controlling the virus (Altfeld and Rosenberg, 2000).

HIV infection of target cells requires fusion of the viral membrane with the cellular membrane; this process is catalyzed by the product of the env gene, the envelope glycoprotein gp160. Mature gp160 is composed of two non-covalently associated subunits - gp120 and gp41 (Wyatt and Sodroski, 1998). Following the interaction of gp120 with membrane receptors on the target cell, the gp41 subunit plays a critical role in virus entry into the target cell. Several functional domains have been identified previously in gp41 (Figure 1). The N-terminal hydrophobic fusion domain, the fusion peptide (FP), is thought to play a central role in membrane fusion (Figure 1). Indeed, a mutant FP with a single amino acid (aa) substitution, V2E, shows less fusogenic activity than wild type FP (Kliger *et al.*, 1997). The first 16 aa of FP inserts into the target cell membrane, and the C20 region inserts into the virus membrane (Peisajovich and Shai, 2003; Suarez *et al.*, 2000). The N36 and C34 peptides contain heptad repeats that form a six-helix bundle linker (Chan *et al.*, 1997) that brings the viral and target membranes into close proximity. Fusion can be

inhibited by a peptide corresponding to the C terminal heptad repeat, DP178 (amino acids 638-673 of the HIV-1_{LAI} gp41 protein); this peptide is a potent inhibitor of HIV infection, and has been recently approved for human use (Lawless *et al.*, 1996). HIV gp41-derived peptides useful for inhibiting viral infection were disclosed, for example, in U.S. Patent
5 Nos. 5,464,933, 6,133,418, 6,093,794 (directed to DP178 and to DP107, a peptide corresponding to amino acids 558-595 of the HIV_{LAI} gp41, and analogs thereof) and 5,840,843 (directed to corresponding to amino acid residues 600-862, or a portion thereof comprising the sequence corresponding to amino acid residues 637-666 of the envelope glycoprotein of HIV-1_{IIIB}).

10 With regard to the interaction of FP with the T-cell membrane, Cladera *et al* reported that a synthetic peptide encoding the 16 N-terminal aa of FP shows a heterogeneous distribution on the membrane of the Jurkat T-cell line (Cladera *et al.*, 2001). Prominent among the membrane domains of responding T cells is the immune synapse. The immune synapse is the cluster of transmembrane molecules which ensures specific
15 interaction between antigen specific T cells and antigen presenting cells. The immune synapse includes the TCR and the CD4 molecules and other key molecules involved in T-cell activation (Davis and Dustin, 2004; Huppa *et al.*, 2003). Immune synapse function is required for complete T cell activation (Huppa *et al.*, 2003). None of the background art, however, discloses or suggests that the FP domain of HIV-1 gp41 may localize to the
20 immune synapse and regulate T cell activation.

While the normal immune system is closely regulated, aberrations in immune responses are not uncommon. In some instances, the immune system functions inappropriately and reacts to a component of the host as if it were, in fact, foreign. Such a response results in an autoimmune disease, in which the host's immune system attacks the
25 host's own tissue. T cells, as the primary regulators of the immune system, directly or indirectly affect such autoimmune pathologies.

T cell-mediated inflammatory diseases refers to any condition in which an inappropriate T cell response is a component of the disease. This includes both diseases mediated directly by T cells, and also diseases in which an inappropriate T cell response
30 contributes to the production of abnormal antibodies.

Numerous diseases are believed to result from autoimmune mechanisms. Prominent among these are rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, Type I diabetes, myasthenia gravis, pemphigus vulgaris. Autoimmune diseases affect

millions of individuals worldwide and the cost of these diseases, in terms of actual treatment expenditures and lost productivity, is measured in billions of dollars annually.

T cells also play a major role in the rejection for organ transplantation or graft versus host disease by bone marrow (hematopoietic stem cell) transplantation. Regulation
5 of such immune responses is therefore therapeutically desired.

Traditional reagents and methods used to attempt to regulate an immune response in a patient also result in unwanted side effects and have limited effectiveness. For example, immunosuppressive reagents (e. g., cyclosporin A, azathioprine, and prednisone) used to treat patients with autoimmune diseases also suppress the patient's entire immune
10 response, thereby increasing the risk of infection, and can cause toxic side effects to non-lymphoid tissues. Due to the medical importance of immune regulation and the inadequacies of existing immunopharmacological reagents, reagents and methods to regulate specific parts of the immune system have been the subject of study for many years.

15 In some autoimmune diseases the relevant autoantigens are known and can therefore be used for specific therapies. For example, methods for inducing immunological tolerance and/or protective immunity to a specific autoantigen have been disclosed for example by WO 01/12222, WO 97/02016 and WO 01/30378 among many others.

Other components of immune responses such as cytokines and adhesion molecules
20 have also been a target for developing immunomodulatory agents, as disclosed, for example by WO 01/57056, US 6,316,420, WO 04/002500 and WO 00/63251 among many others.

Peptides based on TCR derived sequences, for disrupting TCR function presumably by interfering with assembly have also been disclosed (WO 96/22306, WO 97/47644).
25 However, these peptides were demonstrated to be effective at concentrations about 100 fold higher than the peptides of the present invention, thus having a substantially higher potential for toxicity and side effects.

A method of treating or inhibiting symptoms of an autoimmune disease by administering a sub-immunogenic amount of an antigen more immunoreactive with
30 alloimmune-immunogen-absorbed (AIA) serum as compared to nonimmune serum of the same species was disclosed in US 5,230,887. One putative antigen, based on its purported serological cross reactivity with MHC Class II antigens, was suggested to be intact gp41 of HIV. The alleged cross reactivity resides in a C-terminal peptide (Golding *et al.*, 1989).

WO 89/09785 is directed to peptide sequences capable of inhibiting HIV-induced cell fusion or cytopathic syncytia formation, which correspond to a hydrophobic domain located at the amino terminus of gp41 of HIV-1 and the amino terminus of gp40 of HIV-2. The disclosure stipulates that these peptides could have a D-isomer rather than the L-isomer at the amino terminus of the peptide and/or the first two amino acids of the peptide, though no specific embodiment of any peptide comprising a D-amino acid is disclosed. The '785 publication discloses inhibition of HIV-induced fusion and syncytia formation using a family of related peptides, all comprising at least the first three amino acids of gp41 of the BH10 strain; specific peptides correspond to amino acid residues 1-3, a peptide corresponding to amino acid residues 1-6, and a peptide comprising amino acid residues 1-6 in an altered order.

WO 2005/060350 of some of the inventors of the present invention, published after the priority date of the present invention, discloses membrane binding diastereomeric peptides comprising amino acid sequences corresponding to a fragment of a transmembrane protein, wherein at least two amino acid residues of the diastereomeric peptides being in a D-isomer configuration, useful in inhibiting fusion membrane protein events, including specifically viral replication and transmission. The '350 publication discloses, inter alia, the use of diastereomeric peptides corresponding to amino acids 512 to 544 of HIV-1 LAV1 gp41 for inhibiting membrane fusion processes.

There exists a long-felt need for more effective means of curing or ameliorating T cell mediated inflammatory or autoimmune diseases and ameliorating T cell mediated pathologies. The development of new immunosuppressive agents capable of selectively inhibiting the activation of T lymphocytes with minimal side effects is therefore desirable.

SUMMARY OF THE INVENTION

The present invention discloses for the first time novel uses for peptides derived from the gp41 fusion peptide domain (FP) of HIV or fragments, homologs and derivatives thereof effective in preventing or treating T cell mediated pathologies, including but not limited to inflammatory diseases, autoimmunity and graft rejection. Certain novel active fragments of the gp41 fusion peptide domain (FP) of HIV particularly useful in these methods are claimed as such.

The present invention is based, in part, on the unexpected discovery that the isolated fusion peptide (FP) of the HIV-1 gp41 molecule has therapeutic properties towards T cell mediated inflammatory autoimmune diseases. FP is known in the art to function

together with other gp41 domains to mediate virion fusion with host cells. It is now disclosed for the first time that FP co-localizes with the TCR and CD4 molecules in the T cell membrane and is now shown to inhibit T-cell activation *in vitro* and *in vivo*. Surprisingly, it was discovered that FP (SEQ ID NO:1) specifically inhibited antigen-specific T-cell proliferation and cytokine secretion while T-cell activation by non specific
5 activators, such as mitogenic antibodies, was not affected. Notably, FP inhibited the activation of arthritogenic T cells and adjuvant arthritis *in vivo* in animal models of these diseases. In addition, FP was found to be non-immunogenic *in vivo*, in a sequence and structure dependent manner uncorrelated with its ability to inhibit cell-cell fusion. These
10 unexpected discoveries disclose a novel function of gp41 fusion peptide domain having novel applications for the treatment of T cell mediated pathologies.

Unexpectedly, it was herein discovered that an FP fragment corresponding to amino acid residues 5-13 (SEQ ID NO:407) of SEQ ID NO:1, retain the ability of FP to inhibit antigen-specific T cell proliferation to a greater extent than an FP fragment
15 corresponding to amino acid residues 1-8 (SEQ ID NO:406). The present invention is further based on the unexpected discovery that diastereomeric peptides corresponding to FP or partial sequences thereof inhibit inflammation despite the disruption of the secondary structure of the peptide.

Thus, the present invention provides novel uses for the isolated fusion peptide
20 derived from gp41 of HIV and its fragments, analogs, mutants, variants, conjugates, derivatives and salts, in modulating T cell immunity. The present invention thus relates to the use of both known peptides such as full-length FP (SEQ ID NO:1) its diastereomeric derivative IFFA (SEQ ID NO:6), and the FP mutant V2E (SEQ ID NO:2), as well as peptides not previously described in the art. The invention further provides novel
25 fragments, analogs and variants of FP, as detailed below.

In one aspect, the present invention is directed to the use of an isolated peptide derived from HIV gp41 fusion peptide domain or fragments, analogs, mutants, variants, conjugates, derivatives and salts thereof for the treatment of T cell mediated pathologies, including, but not limited to inflammatory diseases, autoimmune diseases and graft
30 rejection.

According to certain particular embodiments, the disease is a T cell-mediated autoimmune disease including but not limited to: multiple sclerosis, autoimmune neuritis, systemic lupus erythematosus (SLE), psoriasis, Type I diabetes (IDDM), Sjogren's disease,

thyroid disease, myasthenia gravis, sarcoidosis, autoimmune uveitis, inflammatory bowel disease (Crohn's and ulcerative colitis), autoimmune hepatitis or rheumatoid arthritis.

The present invention is particularly exemplified herein below by the animal disease model of adjuvant arthritis (AA), a T cell mediated inflammatory autoimmune
5 disease that serves as an experimental model for rheumatoid arthritis. This model is intended as a non-limitative example used for illustrative purposes of the principles of the invention.

In another particular embodiment, the T cell mediated pathology is selected from the group consisting of allograft rejection and graft-versus-host disease.

10 According to particular embodiments the peptide is a fragment derived from the fusion peptide domain of the gp41 protein of HIV-1. In one preferable embodiment, the peptide has an amino acid sequence as set forth in SEQ ID NO:1 (see Table 1). According to alternative embodiments the peptide is the V2E variant (SEQ ID NO:2; see Table 1). According to certain preferred embodiments, the peptide is an HIV-1 gp41 fusion peptide
15 variant according to Table 2, having an amino acid sequence as set forth in any one of SEQ ID NOS:7-198. According to other preferred embodiments, fusion peptide is an HIV-1 gp41 fusion peptide fragment according to Table 2, having an amino acid sequence as set forth in any one of SEQ ID NOS:199-405. In another particular embodiment, the peptide is an HIV-1 gp41 fusion peptide fragment corresponding to amino acids 1-8 of SEQ ID
20 NO:1, herein designated FP₁₋₈ (SEQ ID NO:406; see Table 1). In another particular embodiment, the peptide is an HIV-1 gp41 fusion peptide fragment corresponding to amino acids 5-13 of SEQ ID NO:1, herein designated FP₅₋₁₃ (SEQ ID NO:407; see Table 1). In another particular embodiment, the fusion peptide is a variant of FP₅₋₁₃ having an amino acid sequence as set forth in any one of SEQ ID NOS:409-414 (see Table 2). It is
25 noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2, 7-407 and 409-414 and longer peptides comprising these sequences are within the scope of the present invention.

In some embodiments, the peptide comprises both D and L amino acids. In another particular embodiment, the peptide is a diastereomeric peptide corresponding to the full
30 length FP, herein designated IFFA (see Table 1) having an amino acid sequence as set forth in SEQ ID NO:6. In another particular embodiment, the peptide is a diastereomeric peptide corresponding to FP₅₋₁₃, herein designated FP_{5-13 A6} (see Table 1) having an amino acid sequence as set forth in SEQ ID NO:408.

In certain embodiment, the peptide includes analogs, variants, derivatives and conjugates of FP₅₋₁₃ capable of inhibiting T cell activation as set forth in formula (I) herein:



5 wherein:

X_1 represents an N-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence alanine-valine-glycine, or may be absent;

10 AA_1 , AA_2 , AA_6 and AA_9 each independently represent an alanine or glycine amino acid residue;

AA_3 represents a phenylalanine, isoleucine, leucine, valine or methionine amino acid residue;

15 AA_4 , AA_5 , AA_7 and AA_8 each independently represent a phenylalanine, isoleucine, leucine, valine, methionine or serine amino acid residue, wherein no more than two amino acid residues of AA_4 , AA_5 , AA_7 and AA_8 are identical, with the exception that any three of AA_4 , AA_5 , AA_7 and AA_8 may be leucine amino acid residues;

20 X_2 represents a C-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence valine-glutamine-alanine, or may be absent;

wherein each amino acid can be of either L or D form and the peptide is no more than 30 amino acid residues in length.

25 In one currently preferred embodiment, the peptide does not contain more than one serine residue.

In another aspect, the invention provides methods for treating or preventing the symptoms of a disease or disorder related to an inappropriate or detrimental T cell response, comprising administering to an individual in need thereof a therapeutically
30 effective amount of a pharmaceutical composition comprising as an active ingredient an

isolated peptide derived from HIV gp41 fusion peptide domain or fragments, analogs, variants, conjugates, derivatives and salts thereof.

In one embodiment, the HIV is HIV-1. In another embodiment, the peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids.

In another embodiment, the peptide is a peptide capable of inhibiting T cell activation as set forth in formula (I), as detailed above.


Another aspect of the present invention is a method of inhibiting T-cell activation, wherein said method comprises administering to an individual in need thereof a therapeutically effective amount of a pharmaceutical composition comprising as an active ingredient an isolated peptide derived from HIV gp41 fusion peptide domain or fragments, analogs, variants, conjugates, derivatives and salts thereof.

In one embodiment, the HIV is HIV-1. In another embodiment, the peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, derivatives and salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

In another embodiment, the peptide is a peptide capable of inhibiting T cell activation as set forth in formula (I), as detailed above.

Other aspects of the present invention are directed to novel peptides derived from HIV gp41 fusion peptide domain, and pharmaceutical compositions comprising same.

Thus, in another aspect, there is provided a peptide capable of inhibiting T cell activation having the formula (I):



wherein:

X_1 represents an N-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence alanine-valine-glycine, or may be absent;

AA_1 , AA_2 , AA_6 and AA_9 each independently represent an alanine or glycine amino acid residue;

AA₃ represents a phenylalanine, isoleucine, leucine, valine or methionine amino acid residue;

5 AA₄, AA₅, AA₇ and AA₈ each independently represent a phenylalanine, isoleucine, leucine, valine, methionine or serine amino acid residue, wherein no more than two amino acid residues of AA₄, AA₅, AA₇ and AA₈ are identical, with the exception that any three of AA₄, AA₅, AA₇ and AA₈ may be leucine amino acid residues;

10 X₂ represents a C-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence valine-glutamine-alanine, or may be absent;

and wherein each amino acid can be of either L or D form and the peptide is no more than 30 amino acid residues in length.

15 In one currently preferred embodiment, the peptide does not contain more than one serine residue. In another embodiment, the peptide is FP₅₋₁₃, having an amino acid sequence as set forth in SEQ ID NO:407. In another embodiment, the peptide having an amino acid sequence as set forth in any one of SEQ ID NOS:409-414.

20 In another embodiment, the peptide comprises both L and D amino acids. In one particular embodiment, the peptide is FP₅₋₁₃ A₆, having an amino acid sequence as set forth in SEQ ID NO:408.

According to one aspect, the present invention provides pharmaceutical compositions comprising as an active ingredient a peptide of formula (I) and salts thereof, and a pharmaceutically acceptable carrier or diluent.

25 These and other embodiments of the present invention will become apparent in conjunction with the figures, description and claims that follow.

BRIEF DESCRIPTION OF THE DRAWINGS

30 **Figure 1** depicts the functional domains of HIV-1 gp41 ectodomain. gp41 becomes active after gp120 binds to surface receptors; FP inserts into the membrane of the target cell; C20 inserts into the membrane of the virion; N36 and C34 form a six-helix

bundle “spring” that brings the membranes into apposition; and ISU is immunosuppressive.

Figure 2 demonstrates the co-localization of FP with the CD4 and TCR molecules in the T-cell immune synapse. FP, V2E or AMP peptides were conjugated to rhodamine (Rho) and used to study peptide binding to the membranes of activated T cells, in combination with FITC-labeled antibodies to CD4 or TCR. (a) Distribution of Rho-labeled FP or AMP in activated T cells. (b) Co-localization of FP-Rho with the CD4 and TCR molecules. (c) Co-localization of V2E-Rho with the TCR. (d) FRET between FP-Rho or V2E-Rho and FITC-labeled antibodies to TCR.

Figure 3 presents FP inhibition of the T-cell response to Mt. LNC from Mt immunized rats were activated *in vitro* with the Mt176-90 peptide (a, c and e) or PPD (b, d and f) in the presence of FP (black), V2E (gray) or p277 (white) and the proliferative responses (a and b), IFN γ secretion (c, d) and IL-10 secretion (e, f) were assayed. Similar results were obtained in at least three additional experiments.

Figure 4 contains a graph showing that FP acts on the T cells and not on the APC in the immune synapse. A2b T cells or APC were separately pre-incubated with FP (black), V2E (gray), or p277 (white) for 2 hours, and washed. The treated T cells were mixed with untreated APC, and the treated APC were mixed with untreated A2b, and the proliferation of the A2b T-cells upon stimulation with Mt176-90 was assayed.

Figure 5 demonstrates that FP does not inhibit T-cell activation induced by PMA/ionomycin or antibodies to CD3. A2b T cells were stimulated with PMA/ionomycin (a) or antibodies to CD3 (b) in the presence of FP (black) or p277 (white), and T cell proliferation was studied.

Figure 6 illustrates inhibition of AA by FP. AA was induced by immunization to Mt in oil, mixed with FP (diamonds), V2E (triangles), p277 (hollow squares) or PBS (full squares). Arthritis was scored every two or three days starting at day 10 (a); the leg swelling was measured at day 26 (b); the DTH response to PPD was measured at day 16 (c); and IFN γ secretion was measured at day 26 upon stimulation of LNC with HSP71 or Mt176-90 (d).

Figure 7 provides a schematic representation of the working hypotheses for FP in HIV infection and in immunotherapy.

Figure 8 demonstrates FP inhibition of T cell immunity to FP *in vivo*. LNC from rats immunized to Mt in oil, mixed with FP (triangles), V2E (crosses), IFFA (squares) or PBS (diamonds) were incubated *in vitro* with FP, V2E, IFFA, or PBS, respectively, and IFN γ secretion was assayed. Similar results were obtained in at least three additional experiments.

Figure 9 demonstrates that FP-derived fragments and diastereomeric peptides inhibit antigen-specific T cell proliferation. A-C. Proliferative responses of LNC from Mt immunized mice activated *in vitro* with the MOG₃₅₋₅₅ peptide in the presence or absence of FP₅₋₁₃ (A), FP₁₋₈ (B) or FP_{5-13A6} (C). D. Proliferative responses of LNC from Mt immunized rats activated *in vitro* with PPD in the presence or absence of IFFA.

Figure 10 demonstrates in-vivo inhibition of Adjuvant Arthritis in rats by IFFA. AA was induced by immunization to Mt in oil, mixed with IFFA peptide (circles) or PBS (diamonds). Arthritis was scored every two or three days starting at day 10; standard error was under 10%.

Figure 11 demonstrates in-vivo inhibition of DTH by dermal treatment with FP and IFFA.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a novel process for controlling T cell activity. It is now shown for the first time that a composition containing HIV gp41 fusion peptide (FP) is an effective therapeutic reagent for treating T cell-mediated diseases.

Specifically, the present invention relates to the use of the isolated fusion peptide of a human virus in order to inhibit T-cell activation. Preferably the virus is the human immune deficiency virus. More preferably the virus is HIV-1. The isolated fusion peptide is used for preventing or ameliorating T cell-mediated pathologies, such as autoimmune diseases, inflammatory diseases, graft rejection and allergy.

The present invention is based in part on the unexpected discovery that HIV-1 gp41 fusion peptide domain (FP) is able to suppress antigen-specific T cell activation, as will be described in more detail herein.

Without wishing to be bound by a particular mechanism of action it is postulated that the peptides of this invention localize to the immune synapse, thereby inhibiting T cell activation.

Figure 7 provides a schematic representation of the postulated functions of FP, whereby FP serves two functions in HIV infection and provides a new tool for immunotherapy. *HIV Infection* schematically illustrates two effects of FP on HIV infection. Insertion of FP into the T-cell immune synapse facilitates fusion and infection, while it down-regulates specific T-cell immunity to HIV epitopes. *Immunotherapy* extends the down-regulation by FP to a T-cell mediated immune response under circumstances where the T-cell mediated immune response is deleterious, e.g., a response directed towards self antigens, or graft rejection. Thus, FP itself or an active fragment thereof can serve as a new immunotherapeutic agent.

10 **T cell activation**

T lymphocytes (T cells) are one of a variety of distinct cell types involved in an immune response. The activity of T cells is regulated by antigen, presented to a T cell in the context of a major histocompatibility complex (MHC) molecule. The T cell receptor (TCR) then binds to the MHC-antigen complex. Once antigen is complexed to MHC, the MHC-antigen complex is bound by a specific TCR on a T cell, thereby altering the activity of that T cell.

Proper activation of T lymphocytes by antigen-presenting cells requires stimulation not only of the TCR, but the combined and coordinated engagement of its co-receptors. Most TCR co-receptors bind cell-surface ligands and are concentrated in areas of cell-cell contact, forming what has been termed an immune synapse. Synapse formation has been associated with the induction of antigen-specific T cell proliferation, cytokine production and lytic granule release, and its function was determined necessary for complete T cell activation (Davis and Dustin, 2004; Huppa *et al.*, 2003).

25 **T cell mediated pathologies**

In one aspect, the present invention provides a method for treating a T cell mediated pathology. The term "T-cell mediated pathology" refers to any condition in which an inappropriate or detrimental T cell response is a component of the etiology or pathology of a disease or disorder. The term is intended to include both diseases directly mediated by T cells, and also diseases in which an inappropriate or detrimental T cell response contributes to the production of abnormal antibodies, as well as graft rejection.

In one embodiment of the invention, the composition is useful for treating a T cell-mediated autoimmune disease, including but not limited to: multiple sclerosis, autoimmune neuritis, systemic lupus erythematosus (SLE), psoriasis, Type I diabetes (IDDM), Sjogren's

disease, thyroid disease, myasthenia gravis, sarcoidosis, autoimmune uveitis, inflammatory bowel disease (Crohn's and ulcerative colitis) or autoimmune hepatitis, rheumatoid arthritis.

In other embodiments the composition is useful for treating a T cell-mediated inflammatory disease, including but not limited to: inflammatory or allergic diseases such as asthma, hypersensitivity lung diseases, hypersensitivity pneumonitis, delayed-type hypersensitivity, interstitial lung disease (ILD) (e.g., idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis or other inflammatory diseases); scleroderma; psoriasis (including T-cell mediated psoriasis); dermatitis (including atopic dermatitis and eczematous dermatitis), iritis, conjunctivitis, keratoconjunctivitis, cutaneous lupus erythematosus, scleroderma, vaginitis, proctitis, drug eruptions, allergic encephalomyelitis, acute necrotizing hemorrhagic encephalopathy, idiopathic bilateral progressive sensorineural hearing loss, aplastic anemia, pure red cell anemia, idiopathic thrombocytopenia, polychondritis, Graves ophthalmopathy and primary biliary cirrhosis.

In other embodiments, the composition is useful for treating graft rejection, including allograft rejection or graft-versus-host disease.

Peptides and Peptide Based Pharmaceutical Compositions

The present invention is based, in part, on the surprising discovery that HIV fusion peptide (FP) and fragments and derivatives thereof inhibit T cell activation. Unexpectedly, it was further discovered, that an isolated FP fragment corresponding to amino acid residues 5-13 of FP, and diastereomeric derivative thereof, and, to a lesser extent, an isolated fragment corresponding to amino acid residues 1-8 of FP, retain the ability of FP to inhibit T cell antigen-dependent proliferation.

Thus, The present invention provides, in a first aspect, a peptide capable of inhibiting T cell activation of the formula (I):



wherein:

X_1 represents an N-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence alanine-valine-glycine, or may be absent;

AA₁, AA₂, AA₆ and AA₉ each independently represent an alanine or glycine amino acid residue;

AA₃ represents a phenylalanine, isoleucine, leucine, valine or methionine amino acid residue;

5 AA₄, AA₅, AA₇ and AA₈ each independently represent a phenylalanine, isoleucine, leucine, valine, methionine or serine amino acid residue, wherein no more than two amino acid residues of AA₄, AA₅, AA₇ and AA₈ are identical, with the exception that any three of AA₄, AA₅, AA₇ and AA₈ may be leucine amino acid residues;

10 X₂ represents a C-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence valine-glutamine-alanine, or may be absent;

and wherein each amino acid can be of either L or D form and the peptide is no more
15 than 30 amino acid residues in length.

Hydrophobicity is generally defined with respect to the partition of an amino acid between a nonpolar solvent and water. Hydrophobic amino acids are those acids which show a preference for the nonpolar solvent. Relative hydrophobicity of amino acids can be expressed on a hydrophobicity scale on which glycine has the value 0.5. On such a scale,
20 amino acids which have a preference for water have values below 0.5 and those that have a preference for nonpolar solvents have a value above 0.5. As used herein, the term "hydrophobic amino acid" refers to an amino acid that, on the hydrophobicity scale has a value greater or equal to 0.5, in other words, has a tendency to partition in the nonpolar solvent which is at least equal to that of glycine. Examples of naturally occurring
25 hydrophobic amino acids are aliphatic amino acids alanine, glycine, isoleucine, leucine, methionine, proline, and valine, and aromatic amino acids tryptophan, phenylalanine, and tyrosine. These amino acids confer hydrophobicity as a function of the length of aliphatic and size of aromatic side chains, when found as residues within a protein.

In a currently preferred embodiment, the peptide does not contain more than one
30 serine residue. In one embodiment, the peptide is FP₅₋₁₃, having an amino acid sequence as set forth in SEQ ID NO:407 (GALFLGFLG). Other exemplary embodiments of the generic structure of formula (I) are isolated peptides selected from the group consisting of: GAVFLGFLG (SEQ ID NO:409), GAMFLGFLG (SEQ ID NO:410), GAVLLGFLG

(SEQ ID NO:411), GAFFLGFLG (SEQ ID NO:412), GAMIFGFLG (SEQ ID NO:413) and GALLFGFLG (SEQ ID NO:414).

In another embodiment, the peptide is a diastereomeric peptide, i.e. a peptide comprising both L and D amino acids. The diastereomeric peptides may be advantageous
5 over all L- or all D-amino acid peptides having the same amino acid sequence because of their higher water solubility and lower susceptibility to proteolytic degradation. Such characteristics endow the diastereomeric peptides with higher efficacy and higher bioavailability than those of the all L or all D-amino acid peptides comprising the same amino acid sequence. In one particular embodiment, the peptide is FP₅₋₁₃ A₆, having an
10 amino acid sequence as set forth in SEQ ID NO:408 (GALFLGFLG, the D amino acid is bold and underlined) .

The peptide of formula (I) is preferably less than 30 amino acids in length. It is to be understood that longer peptides, e.g. up to 50 amino acids in length may also be used for the treatment of T cell mediated pathologies according to the invention. However, shorter
15 peptides are preferable, in one embodiment, for being easier to manufacture. In another currently preferred embodiment, the peptide is no more than 20 amino acids in length. In another currently preferred embodiment, the peptide is no more than 10 amino acids in length. Thus, in certain embodiments, the optional N and C termini, i.e. X₁ and X₂, are up to 20, preferably up to 10, and more preferably 5 or in other embodiments up to 3 amino
20 acids in length or may be absent.

The amino acid sequences of X₁ and X₂ may comprise sequences corresponding to the flanking regions of FP, so long as the peptide is not a peptide previously described in the art. It is noted, that the peptides provided by formula (I) are not intended to include known peptides such as the full length FP (SEQ ID NO:1) and diastereomeric derivatives
25 corresponding to the full length sequence, such as IFFA (SEQ ID NO:6) and other diastereomeric derivatives corresponding to the full length sequence disclosed in WO2005/060350, the known V2E mutant (SEQ ID NO:2) disclosed by Kliger et al. (1997), the peptide corresponding to amino acids 1-16 of SEQ ID NO:1, having an amino acid sequence as set forth in SEQ ID NO:225, disclosed by Martin et al. (1992), and the
30 peptides comprising amino acids 1-3 of SEQ ID NO:1 disclosed by WO 89/09785.

In alternate embodiments, X₁ and X₂ may comprise sequences not derived from FP, so long as they retain the required level of hydrophobicity.

According to another aspect, the present invention is directed to compositions comprising the isolated fusion peptide derived from gp41 of HIV as well as analogs, mutants, variants conjugates, and derivatives thereof.

According to particular embodiments the fusion peptide is derived from the gp41 protein of HIV-1. According to alternative embodiments the fusion peptide is the V2E variant. According to certain preferred embodiments, the fusion peptide has an amino acid sequence as set forth in SEQ ID NO:1. In other preferred embodiments, the fusion peptide has an amino acid sequence as set forth in SEQ ID NO:2. In other preferred embodiments, the fusion peptide is an HIV-1 gp41 fusion peptide variant according to Table 2, having an amino acid sequence as set forth in any one of SEQ ID NOS:7-198. Databases of various HIV strains and variants are available (see, for example, <http://www.hiv.lanl.gov/content/index>). According to other preferred embodiments, the fusion peptide is an HIV-1 gp41 fusion peptide fragment according to Table 2, having an amino acid sequence as set forth in any one of SEQ ID NOS:199-405. According to other preferred embodiments, the fusion peptide is an HIV-1 gp41 fusion peptide fragment according to Table 2, having an amino acid sequence as set forth in any one of SEQ ID NOS:406-407 and 409-414. In other preferred embodiments, the fusion peptide comprises both L and D amino acid residues. In another preferred embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:6 and 408.

It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention. The fusion peptide fragments according to the present invention are preferably 5-50 amino acids in length, more preferably 10-36 amino acids in length.

In other preferred embodiments, the compositions comprise a peptide of formula (I), as detailed above.

The peptides of the invention may be synthesized using the procedures described in detail in the Examples. However, other methods known in the art, including, but not limited to, solid phase (e.g. Boc or f-Moc chemistry) as well as solution phase synthesis methods, may be used for synthesizing the peptides of the invention.

The amino acid residues described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the peptide retains the desired functional property.

It should be understood that a fusion peptide need not be identical to the amino acid sequence of the peptide of the invention, so long as it includes the required sequence and is able to function as the peptide of the invention as described herein.

The present invention encompasses any analog, derivative, and conjugate
5 containing the FP of the invention, so long as the peptide is capable of inhibiting T cell activation. Thus, the present invention encompasses peptides containing non-natural amino acid derivatives or non-protein side chains.

The term "analog" includes any peptide having an amino acid sequence substantially identical to one of the sequences specifically shown herein in which one or
10 more residues have been conservatively substituted with a functionally similar residue and which displays the abilities as described herein. Examples of conservative substitutions include the substitution of one non-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between
15 glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that such peptide
20 displays the requisite inhibitory function on T cells as specified herein.

The term derivative includes any chemical derivative of the peptide of the invention having one or more residues chemically derivatized by reaction of side chains or functional groups. Such derivatized molecules include, for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl
25 groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-im-benzylhistidine. Also included as chemical derivatives are those peptides, which
30 contain one or more naturally occurring amino acid derivatives of the twenty standard amino acid residues. For example: 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for

histidine; homoserine may be substituted or serine; and ornithine may be substituted for lysine.

In addition, a peptide derivative can differ from the natural sequence of the peptides of the invention by chemical modifications including, but are not limited to, terminal-NH₂ acylation, acetylation, or thioglycolic acid amidation, and by terminal-carboxyamidation, 5 e.g., with ammonia, methylamine, and the like. Peptides can be either linear, cyclic or branched and the like, which conformations can be achieved using methods well known in the art.

Thus, for example, the "blocking groups" represented in formula (I) by X₁ and X₂ 10 are chemical groups that are routinely used in the art of peptide chemistry to confer biochemical stability and resistance to digestion by exopeptidases. Suitable N-terminal protecting groups include, for example, C₁₋₅ alkanoyl groups such as acetyl; other exemplary blocking groups include, without limitation, t-butyloxycarbonyl, methyl, succinyl, methoxysuccinyl, suberyl, adipyl, azeloyl, dansyl, benzyloxycarbonyl, 15 fluorenylmethoxycarbonyl, methoxyaselayl, methoxyadipyl, methoxysuberyl, and 2,3-dinitrophenyl groups. Also suitable as N-terminal protecting groups are amino acid analogs lacking the amino function. Suitable C-terminal protecting groups include groups which form ketones or amides at the carbon atom of the C-terminal carboxyl, or groups which form esters at the oxygen atom of the carboxyl. Ketone and ester-forming groups include 20 alkyl groups, particularly branched or unbranched C₁₋₅ alkyl groups, e.g., methyl, ethyl, and propyl groups, while amide-forming groups include amino functions such as primary amine, or alkylamino functions, e.g., mono- C₁₋₅ alkylamino and di-C₁₋₅ alkylamino groups such as methylamino, ethylamino, dimethylamino, diethylamino, methylethylamino and the like. Other exemplary blocking groups may include, without limitation, C₃₋₈ cycloalkyl 25 group such as cyclopentyl, cyclohexyl, C₆₋₁₂ aryl group such as phenyl and α -naphthyl, phenyl-C₁₋₂ alkyl group such as benzyl, phenethyl or C₇₋₁₄ aralkyl group, C₁₋₂ alkyl group such as α -naphthyl methyl group, and additionally, pivaloyloxymethyl group which is generally used as an oral bioavailable ester. Amino acid analogs are also suitable for protecting the C-terminal end of the present compounds, for example, decarboxylated 30 amino acid analogues such as agmatine.

Peptides of the present invention also include any peptide having one or more additions and/or deletions of residues relative to the sequence of the fusion peptide of the invention, so long as the requisite inhibitory activity is maintained.

Addition of amino acid residues may be performed at either terminus of the peptides of the invention for the purpose of providing a "linker" by which the peptides of this invention can be conveniently bound to a carrier. Such linkers are usually of at least one amino acid residue and can be of 40 or more residues, more often of 1 to 10 residues.

5 Typical amino acid residues used for linking are tyrosine, cysteine, lysine, glutamic and aspartic acid, or the like.

A peptide of the present invention may be coupled to or conjugated with another protein or polypeptide to produce a conjugate. Such a conjugate may have advantages over the peptide used alone. For example, a peptide of the invention may be conjugated to an
10 antigen involved in a T cell mediated pathology. Without wishing to be bound by any theory or mechanism of action, vaccination with such a conjugate may result in reduced T cell activation to the conjugated antigen, and thereby induce a tolerogenic immune response to said disease target antigen. The peptides can be conjugated directly via an amide bond, synthesized as a dual ligand peptide, or joined by means of a linker moiety as
15 is well known in the art to which the present invention pertains.

In another embodiment, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of an isolated fusion peptide of the invention and a pharmaceutically acceptable carrier.

A pharmaceutical composition useful in the practice of the present invention
20 typically contains a peptide of the invention formulated into the pharmaceutical composition as a neutralized pharmaceutically acceptable salt form. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide), which are formed with inorganic acids, such as for example, hydrochloric or phosphoric acid, or with organic acids such as acetic, oxalic, tartaric, and the like.

25 Suitable bases capable of forming salts with the peptides of the present invention include, but are not limited to, inorganic bases such as sodium hydroxide, ammonium hydroxide, potassium hydroxide and the like; and organic bases such as mono-, di- and tri-alkyl and aryl amines (e.g. triethylamine, diisopropyl amine, methyl amine, dimethyl amine and the like) and optionally substituted ethanolamines (e.g. ethanolamine,
30 diethanolamine and the like).

The preparation of pharmaceutical compositions, which contain peptides or polypeptides as active ingredients is well known in the art. Typically, such compositions are prepared as indictable, either as liquid solutions or suspensions, however, solid forms,

which can be suspended or solubilized prior to injection, can also be prepared. The preparation can also be emulsified. The active therapeutic ingredient is mixed with inorganic and/or organic carriers, which are pharmaceutically acceptable and compatible with the active ingredient. Carriers are pharmaceutically acceptable excipients (vehicles) comprising more or less inert substances when added to a pharmaceutical composition to confer suitable consistency or form to the composition. Suitable carriers are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents and pH buffering agents, which enhance the effectiveness of the active ingredient.

Therapeutic Use

In one aspect, the present invention provides the use of pharmaceutical compositions comprising the gp41 fusion peptide domain (FP), effective in preventing or treating T Cell mediated pathologies.

One aspect of the present invention is a method of treating an autoimmune disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising an isolated fusion peptide of the invention or an active fragment thereof. In one embodiment, the HIV is HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

Another aspect of the present invention is a method of treating an autoimmune disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising a peptide capable of inhibiting T cell activation according to formula (I) or salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

Another aspect of the present invention is a method of preventing the symptoms of an autoimmune disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising an isolated fusion peptide of the invention. In one embodiment, the HIV is

HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

Another aspect of the present invention is a method of preventing the symptoms of an autoimmune disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising a peptide capable of inhibiting T cell activation according to formula (I) or salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

One aspect of the present invention is a method of treating a T-cell mediated inflammatory disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising an isolated fusion peptide of the invention. In one embodiment, the HIV is HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

Another aspect of the present invention is a method of treating a T-cell mediated inflammatory disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising a peptide capable of inhibiting T cell activation according to formula (I) or salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

Another aspect of the present invention is a method of preventing the symptoms of a T-cell mediated inflammatory disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising an isolated fusion peptide of the invention. In one embodiment, the HIV is HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the

fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

Another aspect of the present invention is a method of preventing the symptoms of a T-cell mediated inflammatory disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising a peptide capable of inhibiting T cell activation according to formula (I) or salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

One aspect of the present invention is a method of treating or preventing the symptoms of graft rejection or graft versus host disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising an isolated fusion peptide of the invention. In one embodiment, the HIV is HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

Another aspect of the present invention is a method of a method of treating or preventing the symptoms of graft rejection or graft versus host disease, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising a peptide capable of inhibiting T cell activation according to formula (I) or salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

One aspect of the present invention is a method of inhibiting T-cell activation, wherein said method comprises administering to an individual in need of said treatment a therapeutically effective amount of a pharmaceutical composition comprising an isolated fusion peptide of the invention. In one embodiment, the HIV is HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID

NOS:1, 2 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

Another aspect of the present invention is a method of inhibiting T-cell activation, wherein said method comprises administering to an individual in need of said treatment a
5 therapeutically effective amount of a pharmaceutical composition comprising a peptide capable of inhibiting T cell activation according to formula (I) or salts thereof. In another embodiment, the peptide comprises both D and L amino acids.

In another aspect, the invention is directed to the use of a fusion peptide of the invention for the preparation of a pharmaceutical composition for treating or preventing T
10 cell mediated pathologies. In one embodiment, the HIV is HIV-1. In another embodiment, the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof. In another embodiment, the fusion peptide comprises both D and L amino acids. It is noted that both shorter active fragments derived from the peptides denoted as SEQ ID NOS:1, 2
15 and 6-414 and longer peptides comprising these sequences are within the scope of the present invention.

In another aspect, the invention is directed to the use of a peptide capable of inhibiting T cell activation according to formula (I) for the preparation of a pharmaceutical composition for treating or preventing T cell mediated pathologies.

20 In another aspect, the pharmaceutical composition can be delivered by a variety of means including intravenous, intramuscularly, infusion, oral, intranasal, intraperitoneal, subcutaneous, rectal, topical, or into other regions, such as into synovial fluids. However delivery of the composition transdermally is also contemplated, such by diffusion via a transdermal patch. For oral ingestion it is possible to prepare peptide analogs or specific
25 peptide formulations having improved oral bioavailability and enhanced resistance to degradation as are known in the art.

The composition is administered in a manner compatible with the dosage formulation, and in a therapeutically effective amount. The quantity to be administered depends on the subject to be treated, capacity of the subject's blood hemostatic system to
30 utilize the active ingredient, and the degree of inhibition of T cell activation or T cell mediated pathology desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual.

A therapeutically effective amount of a peptide of the invention is an amount that when administered to a patient is capable of inhibiting T cell activation. Assays for detecting the activity of the peptides of the invention may include, but are not limited to, inhibition of T cell antigen-specific proliferation, inhibition of T cell antigen-specific secretion of cytokines such as IFN- γ and IL-10, and inhibition of in vivo disease models including, but not limited to adjuvant arthritis and DTH, as described in the Examples. However, other methods for detecting the inhibition of antigen-specific T cell activation are well known in the art, and may be used for assessing the activity of the peptides of the invention. Preferably, a therapeutically effective amount of a peptide of the present invention is an amount that reduces (inhibits) T cell activation by at least 10 percent, more preferably by at least 50 percent, and most preferably by at least 90 percent, when measured in an in vitro assay or in an in vivo assay. Preferably, a pharmaceutical composition is useful for inhibiting a T cell mediated pathology in a patient as described further herein. In this embodiment, a therapeutically effective amount is an amount that when administered to a patient is sufficient to inhibit, preferably to eradicate, a T cell mediated pathology. A preferred single dose of fusion peptide is from about 5 μ g to about 50 mg per kg of body weight, preferably from about 50 μ g to about 5 mg per kg of body weight, and more preferably from about 0.125 mg to about 2 mg per kg of body weight. Typically, the physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention. The fusion peptides of the invention may be administered, for example, as daily or weekly administrations of single doses as described above.

Methods of treating a disease according to the invention may include administration of the pharmaceutical compositions of the present invention as a single active agent, or in combination with additional methods of treatment. The methods of treatment of the invention may be in parallel to, prior to, or following additional methods of treatment.

30

The following examples are presented in order to more fully illustrate some embodiments of the invention. They should, in no way be construed, however, as limiting the broad scope of the invention.

EXAMPLES

Peptide Synthesis. The peptides used in this work (presented in Table 1 below) were synthesized using a solid phase method as previously described (Kliger *et al.*, 1997, Merrifield *et al.*, 1982). The synthetic peptides were purified (>98% homogeneity) by reverse-phase HPLC on a C4 column using a linear gradient of 20-60% acetonitrile in 0.05% TFA, for 60 min. The peptides were subjected to amino acid analysis and mass spectrometry to confirm their composition. Unless stated otherwise, stock solutions of concentrated peptides were maintained in DMSO to avoid aggregation of the peptides prior to use. The final concentration of DMSO in each experiment (5% v/v) had no effect on the system under investigation.

Table 1. Peptides' designation and sequence.

Designation	Amino Acid Sequence	Start/End	Protein	SEQ ID NO:
FP (full length)	AVGIGALFLGFLGAAGSTMGARSM ^L TLTVQARQ	512-535	gp41	1
V2E	A EGIGALFLGFLGAAGSTMGARSM ^L TLTVQARQ	512-535	gp41	2
AMP	LLKLLKKLLKKLLKL	--	Syn.	3
p277	VLGGGCALLRCIPALDSLTPANED	437-460	HSP60	4
Mt176-90	EESNTFGLQLELTEG	176-190	HSP65	5
IFFA	AVG I GAL F LG F LG A AGSTMGARSM ^L TLTVQARQ	--	Syn.	6
FP ₁₋₈	AVGIGALF	512-519	gp41	406
FP ₅₋₁₃	GALFLGFLG	516-524	gp41	407
FP ₅₋₁₃ A6	G ALFLGFLG	--	Syn.	408

Start and End positions are designated according to the HIV-1 HXB2 strain gp160 sequence. The Valine to Glutamic acid mutation at position 2 of the FP is presented in bold face. D-amino acids are presented in bold face and underlined.

gp41, HIV-1 glycoprotein gp41.

AMP, synthetic amphipathic peptide (Papo *et al.*, 2002).

HSP60, human 60 kDa heat shock protein.

HSP65, *M. tuberculosis* 65 kDa heat shock protein.

Syn., synthetic sequence

Fluorescent Labeling of Peptides. The resin-bound peptides were treated with 4-chloro-7-nitrobenz-2-oxa-1,3-diazole fluoride (NBD-F) or 5-carboxytetramethylrhodamine, succinimidyl ester (5-TAMRA, SE (Rhodamine-SE),

respectively. The NBD-F and Rhodamine-SE fluorescent probes were purchased from Molecular Probes (City, State, Country). The reaction with NBD-F took place in DMF, and the reaction with Rhodamine in DMF containing 2% diisopropylethylamine as described previously (Gerber and Shai, 2000). The fluorescent probes were used in an excess of 2
5 equivalents, leading to the formation of resin bound N-terminal NBD or Rhodamine peptides. After 1 h, the resins were washed thoroughly with DMF and then with methylene chloride. The resin was dried under nitrogen flow and then cleaved for 3hr with TFA 95%, H₂O 2.5% and Triethylsilane 2.5%. The fluorescently-labeled peptides were purified by RP-HPLC (reverse phase high-performance liquid chromatography) on a C4 Bio-Rad
10 semi-preparative column (250x10 mm, 300Å pore-size, 5-µm particle size) using a gradient of 20%-60% of acetonitrile/water (both containing 0.05% TFA) for 60 min. The purified peptides were shown to be homogeneous (>98%) by analytical RP-HPLC.

Cell lines, antigens and adjuvants. The CD4⁺ T cell clone A2b 21 reacts with the 180-188 epitope of the 65 kDa heat shock protein (HSP65) of *M. tuberculosis* (Mt), this
15 epitope is contained in the peptide Mt176-90 used herein (van Eden et al., 1988).

Mt Strain H37Ra and incomplete Freund's adjuvant (IFA) were purchased from Difco (Detroit, MI, USA). Tuberculin purified protein derivative (PPD) was provided by the Statens Serum institute (Copenhagen, Denmark). Purified recombinant 71 kDa heat shock protein (HSP71) was generously provided by Prof. Ruurd van der Zee (Institute of
20 Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht, The Netherlands). PMA, ionomycin, ovalbumin (OVA) and Concanavalin A (Con A) were purchased from Sigma (Rehovot, Israel).

Co-localization of peptides with membrane molecules. A2b cells were fixed with 4% para-formaldehyde for 15 min on ice and washed with PBS. The cells were then treated
25 with 2% BSA in PBS at room temperature to block unspecific binding. After 30 min the cells were divided into aliquots containing 50,000 cells per 100 µl and either αTCR-FITC or αCD4-FITC were added (1:100) for two hours. The rhodamine-labeled FP or V2E peptides were added during the last 5 min of incubation at a final concentration of 0.5-1 µM. The cells were then washed with PBS and deposited onto a glass slide. The labeled
30 cell samples were observed under a fluorescence confocal microscope. FITC excitation was set at 488 nm, with the laser set at 20% power to minimize bleaching of the fluorophore. Fluorescence was recorded from 505-525 nm. Rhodamine excitation was set

at 543 nm, with the laser set at 5% power. Fluorescence data were collected from 560 nm and up.

Fluorescence energy transfer (FRET) between FITC (donor label) and Rhodamine (acceptor label) was detected by the increase in FITC fluorescence in a spot where the Rhodamine probe was bleached. Bleaching was achieved by point excitation at 543 nm for 6 seconds with the laser set to 100%. To verify that the increase in FITC fluorescence was not due to auto-fluorescence, bleaching was performed first using the 488 nm laser and only then at 543 nm. No signal was observed in either 505-525 nm or above 560 nm, eliminating the possibility of auto-fluorescence.

T-cell proliferation. T-cell proliferation assays were performed using either lymph node cells (LNC) or the A2b T cell line, which reacts with the Mt176-90 peptide. Popliteal and inguinal LNC were removed 26 days after the injection of Mt in incomplete Freund's adjuvant (IFA), when strong T cell responses to PPD and Mt176-90 are detectable (Quintana *et al.*, 2002). LNC were cultured at a concentration of 2×10^5 cells per well; 5×10^4 A2b T cells were stimulated in the presence of irradiated 5×10^5 thymic antigen presenting cells (APC) per well, prepared as previously described (van Eden *et al.*, 1985). The cells were plated in quadruplicates in 200 μ l round bottom microtiter wells (Costar Corp., Cambridge, USA), with or without antigen, in the presence of various concentrations of the peptides under study. For some experiments, the cells were activated with immobilized anti-CD3 antibodies or PMA/ionomycin as described (Wang *et al.*, 2002). Cultures were incubated for 72 hr at 37°C in a humidified atmosphere of 7.5% CO₂. T-cell responses were detected by the incorporation of [methyl-3H]-thymidine (Amersham, Buckinghamshire, UK; 1 μ Ci/well), added during the last 18 hr of incubation. The results of T cell proliferation experiments are shown as the % of inhibition of the T cell proliferation triggered by the antigen in the absence of HIV or control peptides.

Cytokine assays. Supernatants were collected after 72 hr of stimulation, and rat IL-10 and IFN γ were quantified by enzyme-linked immunosorbent assay (ELISA) using Pharmingen's OPTEIA kit (Pharmingen, San Diego, USA) as described (Quintana *et al.*, 2002). When needed, cytokine levels are expressed as percentage of cytokine inhibition relative to cytokine levels when no peptide is present. Otherwise, the cytokines are shown as pg/ml. The lower limits of detection for the experiments described in this paper were 15 pg/ml for IL-10 and IFN γ . Cytokine amounts were calculated based on calibration curves constructed using recombinant cytokines as standards.

Animals. Three- month old female Lewis rats were used in our experiments, raised and maintained under pathogen-free conditions in the Animal Breeding Center of the Weizmann Institute of Science. The experiments were performed under the supervision and guidelines of the Animal Welfare Committee.

5 **Induction and Assessment of Adjuvant Arthritis (AA).** To test the effect of FP on T-cell activation *in vivo*, AA was used as a model system. AA was induced by injecting 50 μ l of Mt suspended in IFA (0.5 mg/ml) at the base of the tail. At the time of AA induction, each rat also received 100 μ g of FP or control peptide, or PBS dissolved in 50 μ l of IFA and mixed with Mt/IFA used to induce AA. The day of AA induction was
10 designated as day 0. Disease severity was assessed by direct observation of all 4 limbs in each animal. A relative score between 0 and 4 was assigned to each limb, based on the degree of joint inflammation, redness and deformity; thus the maximum possible score for an individual animal was 16 (Quintana *et al.*, 2002). The mean AA score ($\pm$ SEM) is shown for each experimental group. Arthritis was also quantified by measuring hind limb
15 diameter with a caliper. Measurements were taken on the day of the induction of AA and 26 days later (at the peak of AA); the results are presented as the mean $\pm$ SEM of the difference between the two values for all the animals in each group. The person who scored the disease was blinded to the identity of the groups.

Delayed type hypersensitivity (DTH). Twenty μ l of PPD (0.5 mg/ml in PBS)
20 were injected intradermally into the pinna of the right ear on day 16 after AA induction; 20 μ l of sterile PBS were injected in the left ear as control. The thickness of the ear was measured 48 hr later using a vernier caliper and expressed as the difference between the right and the left ear.

Example 1: FP co-localizes with CD4 and TCR in the IS

25 The distribution of FP on the membrane of activated T cells was studied using FP conjugated to Rhodamine (FP-Rho) or NBD (FP-NBD). Rather than uniformly labeling the T-cell membrane, both FP-Rho (Figure 2A) and FP-NBD inserted themselves into a particular membrane domain. This concentrated distribution contrasted with a control membrane active amphipathic peptide (AMP-Rho) that demonstrated a uniform
30 distribution on the cell membrane (Figure 2A).

 On the membrane of CD4⁺ T cells, the IS domain is marked by the TCR, CD3 and CD4 molecules, among other components (Huppa and Davis, 2003). Figure 2B shows the localization of the CD4 and TCR molecules at the IS membrane domain. Note that the FP

conjugates co-localized with the CD4 and TCR molecules (Figure 2B). Both FP-Rho (Figure 2B) and FP-NBD showed the same co-localization; hence distribution to the IS was not limited to a particular fluorophore. In addition, a rhodamine-labeled mutant of FP, V2E (V2E-Rho), showed a similar co-localization with CD4 and the TCR (Figure 2C).

5 The co-localization of FP and its V2E mutant was confirmed with the TCR and CD4 receptors by fluorescence energy transfer (FRET) (Figure 2D). Using a 543nm laser, a spot on the cell membrane was bleached, thereby reducing the fluorescence of the FP-Rho or V2E-Rho conjugates at that particular spot. A significant increase in the fluorescence of the labeled TCR was observed due to FRET between the labeled, bleached
10 FP (acceptor) and a TCR-specific FITC-conjugated antibody (donor). The average R_0 is under 50 Å for the FITC/Rho pair (Heidecker *et al.*, 1995), therefore, the detection of FRET (Figure 2D) suggests that the donor and acceptor molecules must be less than 50 Å apart in the membrane. These results suggest that FP binds to the T cell membrane, and preferentially co-localizes closely with the CD4 and TCR molecules at the IS.

15 **Example 2: FP interferes with T-cell activation *in vitro***

To determine whether the insertion of FP into the IS interferes with T-cell activation, the T-cell response of LNC of Mt immunized rats to the Mt antigen PPD or to the Mt176-90 peptide was studied; these antigens are known to induce strong proliferative responses and cytokine release from T cells in the draining LNC of AA rats (Quintana *et al.*, 2003; Quintana *et al.*, 2002).
20 *al.*, 2003; Quintana *et al.*, 2002).

Figures 3A and 3B show that FP and the V2E mutant peptide inhibited the T-cell proliferative responses to PPD and to Mt176-90 in a dose-dependent manner. Moreover, the inhibitory effect of V2E was lower than that of FP, suggesting that inhibition is sequence specific and that critical molecular interactions were perturbed by the V to E
25 substitution in V2E (see Table 1). The FP peptide, and to a lesser extent V2E, also inhibited in a dose-dependent manner the secretion of IFN γ and IL-10 triggered by stimulation with PPD or the Mt176-90 peptide (Figures 3 C-F). The inhibitory effects of the FP and V2E peptides on antigen-triggered proliferation and cytokine release were not due to cell death, since cells incubated with FP, V2E or the control peptide p277 showed
30 the same survival in culture.

Example 3: FP acts on the T cells and not on the APC

The inhibitory effects of FP on T-cell activation were further studied using a rat CD4⁺ T cell clone, A2b, which proliferates and secretes IFN γ upon stimulation with the

Mt176-90 peptide (Quintana *et al.*, 2003; Quintana *et al.*, 2002). Activation of A2b by Mt176-90 in the presence of FP led to decreased cell proliferation (Figure 4) and less IFN γ release.

To define the cell targeted by FP in the inhibition of T cell activation, the A2b T cells or the APC were pre-incubated separately with FP, before mixing the cells together with the Mt176-90 peptide. Pre-incubation of the APC had no effect on A2b proliferation (Figure 4). However, pre-incubation of the A2b T cells with the FP, and not with the control peptide p277, led to a significant inhibition of T cell proliferation (Figure 4). Thus, FP inhibits T cell activation by directly acting on the T cells rather than on the APC.

Example 4: FP does not inhibit T-cell activation induced by PMA/ionomycin or antibodies to CD3

To learn whether FP also can inhibit T-cell activation other than that induced by APC presentation of specific antigen, the effect of FP on T-cell activation induced by PMA/ionomycin (wherein white histograms represent cell receiving 0.5 mg/ml ionomycin and black histograms represent cells receiving 1 mg/ml ionomycin) or a mitogenic monoclonal antibody to CD3 was tested. FP did not inhibit the activation of A2b T cells by either PMA/ionomycin (Figure 5A) or mitogenic anti-CD3 (Figure 5B). These findings indicate that FP specifically interferes with T-cell activation induced by the recognition of the MHC-peptide complex presented by the APC.

Example 5: FP inhibits T-cell immunity *in vivo*

As a test for the inhibitory effects of FP on the activation of specific T cells *in vivo*, the effects of FP on AA were studied. The immunization of Lewis rats with Mt in oil triggers AA, an experimental autoimmune disease driven by Mt-specific T cells cross-reactive with self-antigens (Holoshitz *et al.*, 1984; Holoshitz *et al.*, 1983). Mt176-90-specific T cells are detectable upon induction of AA (Anderton *et al.*, 1994); indeed the A2b T-cell clone cross-reacts with cartilage and mediates AA (van Eden *et al.*, 1985). Since FP inhibited the T-cell response of primed LNC and of clone A2b to PPD and Mt17-90 *in vitro* (Figures 3 and 4), the effects of FP on the *in vivo* activation of the T cells that drive AA were investigated. FP administered with the antigen at the time of AA induction led to a significantly milder arthritis, both in terms of clinical score (Figure 6A) and ankle swelling (Figure 6B). The control peptides p277 or V2E did not inhibit AA. The mean maximum score was 13.7 ± 0.3 in the control-treated rats, compared to 7.3 ± 0.7 in the FP-treated rats ($p < 0.05$ for both test groups compared to the control group).

The activity of T cells that mediate AA can also be detected *in vivo* by studying the delayed type hypersensitivity (DTH) response to PPD (van Eden *et al.*, 1985). The DTH response to PPD 16 days after AA induction in rats treated with peptide FP, V2E or p277 was studied. Figure 6C shows that the administration of FP led to a 35% reduction in the DTH response to PPD, while the inhibition caused by treatment with the V2E or the p277 peptides was 25% and 10%, respectively.

The T cells driving AA manifest a Th1 phenotype; they secrete relatively large amounts of IFN γ upon activation with Mt antigens such as HSP71 or Mt176-90 (Quintana *et al.*, 2003; Quintana *et al.*, 2002). In contrast, the control of AA by various treatments is usually accompanied by a decreased Th1 response (Quintana *et al.*, 2003; Quintana *et al.*, 2002, Tanaka *et al.*, 1999). Note that LNC from FP-treated rats (black histograms) showed reduced secretion of IFN γ upon stimulation with Mycobacterial antigens HSP71 or Mt176-90 (also designated Mt180), while treatment with V2E (gray histograms) or p277 (white histograms) affected IFN γ secretion only slightly (Figure 6D).

Taken together, these results indicate that the FP can interfere *in vivo* with T-cell activation induced by specific antigens. This interference led to milder AA (Figures 6A and 6B), decreased DTH reactivity (Figure 6C), and lower IFN γ secretion in response to Mt antigens (Figure 6D).

Example 6: FP inhibits T cell immunity to FP *in vivo*.

Rats were injected with FP, a control peptide (V2E or IFFA) or PBS in the presence of Mt and IFA. Twenty six days later, LNC were collected and incubated *ex vivo* with FP, V2E, IFFA, or PBS, respectively, and their IFN- γ secretion level was determined. As can be seen in Figure 8, LNC from FP injected rats could not be activated by FP *ex vivo*, as their level of IFN- γ secretion was compatible with that of non-activated LNC (incubated with PBS). These results indicate low immunogenicity of FP. However, LNC from V2E injected rats showed moderate *ex vivo* activation with V2E (about five fold higher than that of FP), and LNC from rats injected with the IFFA peptide, a diastereomer derivative of FP in which four D-amino acid residues were incorporated (SEQ ID NO:6), were highly reactive to IFFA *ex vivo* (about ten fold more than FP). Thus, the immunosuppressive effect of FP is sequence and structure specific. Similar results were obtained when the experiment was repeated with peptides corresponding to the 16 N-terminal amino acids of FP, V2E and IFFA (data not shown).

The incorporation of D amino acids in the IFFA peptide, which was herein shown to substantially increase its immunogenicity, does not affect its ability to inhibit gp41-mediated membrane fusion (Gerber *et al.*, 2004). Hence, the immnuosuppressive ability of FP, disclosed herein for the first time, is not merely a reflection its known fusogenic ability, but is rather a newly-identified function residing in the peptide.

Example 7: FP-derived fragments and diastereomeric peptides interfere with T-cell activation *in vitro*.

The effect of the FP-derived peptides FP₅₋₁₃, FP₁₋₈, FP_{5-13A6} and IFFA (see Table 1) on antigen-specific T cell proliferation was examined. T-cell proliferation assays were performed using popliteal and inguinal lymph node cells (LNC) removed 26 days after the injection of Mt in incomplete Freund's adjuvant (IFA), as described above. In some experiments, mice LNC were cultured with or without the MOG 35-55 peptide antigen (0.5 µg/ml, Figure 8A; 0.25 µg/ml, Figures 8B-C) in the presence of various concentrations of the peptides under study. In other experiments, rat LNC were cultured with or without PPD antigen (25 µg/ml, Figure 8D) in the presence of various concentrations of the peptides under study. The results of T cell proliferation experiments are shown as the % of inhibition of the T cell proliferation triggered by the antigen in the absence of HIV or control peptides.

As can be seen in Figure 8, all the examined peptides inhibited T cell proliferation in a dose-dependant manner, wherein the fragment corresponding to amino acids 5-13 and the diastereomeric peptide thereof inhibited T cell proliferation to a greater extent than the fragment corresponding to amino acids 1-8 of FP

Example 8: FP-derived diastereomeric peptide inhibits T-cell immunity *in vivo*.

To further examine the effect of diastereomeric FP derived peptides on T cell activation *in vivo*, AA and DTH models were used.

AA was induced in rats as indicated above; at the time of AA induction, each rat also received 100 µg of IFFA or PBS dissolved in 50 µl of IFA and mixed with Mt/IFA used to induce AA. Disease severity was assessed as described above.

As can be seen in Figure 10, the administration of IFFA resulted in milder arthritis, determined by AA clinical score.

In other experiments, DTH response to oxazolone in the presence of FP and IFFA was measured. Female Balb/c mice (5 mice per group) were sensitized to the shaved

abdominal skin with 100 microliter of 2% oxazolone dissolved in acetone/olive oil (4:1 vol/vol) applied topically, and 5 days later challenged with 20 microliter of 0.5% oxazolone in acetone/olive oil, 10 microliter administered to each side of the ear. One hour after stimulation, FP, IFFA (100 µg in 40 µl DMSO) or DMSO were administered to each side of the ear. A constant area of the ear was measured immediately before challenge and 24 h after challenge.

As can be seen in Figure 11, both FP and IFFA inhibited the DTH reaction.

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without undue experimentation and without departing from the generic concept, and, therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

References

1. Johnson, W.E. & Desrosiers, R.C. *Annu. Rev. Med.* **53**, 499-518 (2002).
2. Rosenberg, E.S. et al. *Science* **278**, 1447-1450 (1997).
3. Norris, P.J. & Rosenberg, E.S. *Aids* **15 Suppl 2**, 16-21 (2001).
4. Altfeld, M. & Rosenberg, E.S. *Curr. Opin. Immunol.* **12**, 375-380 (2000).
5. Wyatt, R. & Sodroski, J. *Science* **280**, 1884-1888 (1998).
6. Kliger, Y. et al. *J. Biol. Chem.* **272**, 13496-13505 (1997).
7. Peisajovich, S.G. & Shai, Y. *Biochim. Biophys. Acta* **1614**, 122-129 (2003).
8. Suarez, T., Nir, S., Goni, F.M., Saez-Cirion, A. & Nieva, J.L. *FEBS Lett* **477**, 145-149 (2000).
9. Chan, D.C., Fass, D., Berger, J.M. & Kim, P.S. *Cell* **89**, 263-273 (1997).
10. Lawless, M.K. et al. *Biochemistry* **35**, 13697-13708. (1996).
11. Cladera, J., Martin, I. & O'Shea, P. *Embo. J.* **20**, 19-26 (2001).
12. Davis, D.M. & Dustin, M.L. *Trends Immunol.* **25**, 323-327 (2004).

13. Huppa, J.B., Gleimer, M., Sumen, C. & Davis, M.M. *Nat. Immunol.* **4**, 749-755 (2003).
14. *Nat. Rev. Immunol.* **3**, 973-983 (2003).
15. Heidecker, M., Yan-Marriott, Y. & Marriott, G. *Biochemistry* **34**, 11017-11025 (1995).
16. Quintana, F.J., Carmi, P., Mor, F. & Cohen, I.R. *J. Immunol.* **171**, 3533-3541 (2003).
17. Quintana, F.J., Carmi, P., Mor, F. & Cohen, I.R. *J. Immunol.* **169**, 3422-3428 (2002).
18. Holoshitz, J., Matitiau, A. & Cohen, I.R. *J. Clin. Invest.* **73**, 211-215. (1984).
19. Holoshitz, J., Naparstek, Y., Ben-Nun, A. & Cohen, I.R. *Science* **219**, 56-58 (1983).
20. Anderton, S.M., van der Zee, R., Noordzij, A. & van Eden, W. *J. Immunol.* **152**, 3656-3664. (1994).
21. van Eden, W. et al. *Proc. Natl. Acad. Sci. USA* **82**, 5117-5120 (1985).
22. Tanaka, S. et al. *J. Immunol.* **163**, 5560-5565. (1999).
23. Wang, X.M. et al. *Clin. Immunol.* **105**, 199-207 (2002).
24. Merrifield, R.B., Vizioli, L.D. & Boman, H.G. *Biochemistry* **21**, 5020-5031 (1982).
25. Gerber, D. & Shai, Y. *J. Biol. Chem.* **275**, 23602-23607 (2000).
26. van Eden, W. et al. *Nature* **331**, 171-173 (1988).
27. Papo, N., Oren, Z., Pag, U., Sahl, H.G. & Shai, Y. *J. Biol. Chem.* **277**, 33913-33921 (2002).
28. Gerber D, Pritsker M, Gunther-Ausborn S, Johnson B, Blumenthal R, Shai Y. *J Biol Chem.* **279**, 48224-30 (2004).
29. Edelhoch, H. *Biochemistry* **6**, 1948-1954 (1967).
30. Martin et al. *Biochem Biophys Acta.* **1445**, 124-133 (1992).

Table 2 – fusion peptide variants and active fragments.

Amino acid sequence	SEQ ID NO:
AVGIGALFLGFLGAAGSTMGARSMTLTVQARQL	1
VGIGALFLGFLGAAGSTMGAASMTLTVQARQL	7
VGLGAVFLGFLGAAGSTMGAASMTLTVQARQL	8
AAGIGAVLLGFLGAAGSTMGAASMTLTVQARQL	9
AAGIGAVLPGFLGAAGSTMGAASMTLTVQARQL	10
AAGIGAVLPGFLGAARSTMGAASMTLTVQARQL	11
AAGLGAVFLGFLGAAGSTMGAASMTLTVQARQL	12
AIGIGAMFLGFLGAAGSTMGAASITLTVQARQL	13
AIGIGAVFIGFLGAAGSTMGAASITLTVQARQL	14
AIGIGAVFLGFLGAAGSTMGAASITLTVQARQL	15
AIGIGAVFLGFLGAAGSTMGAASMTLTVQARQL	16
AIGIGAVFLGFLGTAGSTMGAASITLTVQARQL	17
AIGIGAVVLGFLGTAGSTMGAASITLTVQARQL	18
AIGLGAAFLGFLGAAGSTMGAASLTLTVQARQL	19
AIGLGAAFLGFLGAAGSTMGAASMTLTVQARQL	20
AIGLGAAFLGFLGAAGSTMGCASMTLTVQARQL	21
AIGLGAAFLGFLGAAGSTMGVASMTLTVQARQL	22
AIGLGAALLGFLGAAGSTMGAASMTLTVQARQL	23
AIGLGALFLGFLGAAGSTMGAASLTLTVQARQL	24
AIGLGALFLGFLGAAGSTMGAASMTLTVQARQV	25
AIGLGAMFLGFLGAAGSTMGAASLTLTVQARQL	26
AIGLGAMFLGFLGAAGSTMGAASMTLTVQARQL	27
AIGLGAMFLGFLGAAGSTMGAASMTLTVQARQV	28
AIGLGAMFLGFLGAAGSTMGAASVTLTVQARQL	29
AIGLGAMFLGFLGAAGSTMGARSMTLTVQARQL	30
AIGLGAMFLGFLGAAGSTMGARSVTLTVQARQL	31
AIGLGAMFLGFLGTAGSTMGAASLTLTVQARQL	32
AIGLGAVFIGFLGAAGSTMGAASITLTVQARQL	33
AIGLGAVFLGFLGAAGSTMGAASLTLTVQARQL	34
AIGLGAVFLGFLGAAGSTMGAASMTLTVQARQV	35
AIGLGAVFXGFLGAAGSTMGAASMTLTVQARQL	36
AMGIGAVFLGFLGAAGSTMGAASITLTVQARQL	37
AMGIGAVLLGFLGAPGSTMGAASMTLTVQARQL	38
AVGIGAALLGFLGAAGSTMGAASMTLTVQARQL	39
AVGIGAFFLGFLGAAGSTMGAASITLTVQARQL	40
AVGIGAFVLGFLGAAGSTMGAASITLTVQARQL	41
AVGIGALFLGFLAAAGSTMGAASITLTVQARQL	42
AVGIGALFLGFLAAAGSTMGARSITLTVQARQL	43
AVGIGALFLGFLGAAGSAMGAASMTLTVQARQL	44
AVGIGALFLGFLGAAGSTMGAASITLTVQARQL	45
AVGIGALFLGFLGAAGSTMGAASLTLTVQARQL	46
AVGIGALFLGFLGAAGSTMGAASMTLTVQARLL	47
AVGIGALFLGFLGAAGSTMGAASMTLTVQARQ	48

AVGIGALFLGFLGAAGSTMGAASMTLTVQARQL	49
AVGIGALFLGFLGAAGSTMGAASVTLTVQARQL	50
AVGIGALFLGFLGAAGSTMGCTSMTLTVQARQL	51
AVGIGALFLGFLGPAGSTMGAASITLTVQARQL	52
AVGIGALFLGFLGTAGSTMGAASITLTVQARQL	53
AVGIGALFLGFLGTAGSTMGAASLTLTVQARQL	54
AVGIGALIFGFLGAAGSTMGAASITLTVQARQL	55
AVGIGALIFGFLGAAGSTMGAASLTLTVQARQL	56
AVGIGALLFGFLGAAGSTMGAASMTLTVQARQL	57
AVGIGALLFGFLGAAGSTMGAASVTLTVQARQL	58
AVGIGALLFGSLGAAGSTMGAASMTLTVQARLL	59
AVGIGALLIGFLGAAGSTMGAASMTLTVQARQL	60
AVGIGALLLGFLGAAGSTMGAASLTLTVQARQL	61
AVGIGALLLGFLGAAGSTMGAASVTLTVQARQL	62
AVGIGALLVGFLGAAGSTMGAASMTLTVQARQL	63
AVGIGALLVGFLGTAGSTMGAASITLTVQARQL	64
AVGIGALSLGFLGAAGSTMGAASMTLTVQARLL	65
AVGIGALVFGFLGAAGSTMGAASITLTVQARQL	66
AVGIGAMFLGFLGAAGSTMGAASITLTVQARQL	67
AVGIGAMFLGFLGAAGSTMGAASLTLTVQARQL	68
AVGIGAMFLGFLGAAGSTMGAASMTLTVQARQL	69
AVGIGAMFLGFLGAAGSTMGAASMTLTVQARQL	70
AVGIGAMFLGFLGAAGSTMGAASVTLTVQARQL	71
AVGIGAMFLGFLGMAGSTMGAASITLTVQARQL	72
AVGIGAMFLGILGAAGSTMGAASITLTVQARQL	73
AVGIGAMFLGILGTAGSAMGAASMTLTVQARQL	74
AVGIGAMIFGFLGAAGSTMGAASITLTVQARQL	75
AVGIGAMIFGFLGAAGSTMGAASLTLTVQARQL	76
AVGIGAMIFGFLGAAGSTMGAASVTLTVQARQL	77
AVGIGAMIFGFLGAARSTMGAASITLTVQARQL	78
AVGIGAMIFGFLGAPGSTMGAASITLTVQARQL	79
AVGIGAMIFGFSGAAGSTMGAASITLTVQARQL	80
AVGIGAMILGFLGAAGSTMGAASITLTVQARQL	81
AVGIGAMLFGFLGAAGSTMGAASITLTVQARQL	82
AVGIGAMLFGFLGAAGSTMGAASMTLTVQARQL	83
AVGIGAVFFGFLGAAGSTMGAASITLTVQARQL	84
AVGIGAVFIGFLGAAGSTLGAASMTLTVQARQL	85
AVGIGAVFIGFLGAAGSTMGAASITLTVQARQL	86
AVGIGAVFIGFLGAAGSTMGAASMTLTVQARQL	87
AVGIGAVFIGFLGAAGSTMGAASVTLTVQARQL	88
AVGIGAVFIGFLSAAGSTMGAASITLTVQARQL	89
AVGIGAVFLGFLAAAGSSMGAASMTLTVQARQL	90
AVGIGAVFLGFLAAAGSTMGAASITLTVQARQL	91
AVGIGAVFLGFLAAAGSTMGAASITLTVQARQV	92
AVGIGAVFLGFLATAGSTMGAASITLTVQARQL	93
AVGIGAVFLGFLGAAGSTMGAASVTLTVQARQL	94

AVGIGAVFLGFLGAAGSTMGAASITLTIQARQL	95
AVGIGAVFLGFLGAAGSTMGAASITLTVQARQL	96
AVGIGAVFLGFLGAAGSTMGAASLTLTVQARQL	97
AVGIGAVFLGFLGAAGSTMGAASMTLTVQARLL	98
AVGIGAVFLGFLGAAGSTMGAASMTLTVQARQL	99
AVGIGAVFLGFLGAAGSTMGAASMTLTVQARRL	100
AVGIGAVFLGFLGAAGSTMGAASMTVTVQARQL	101
AVGIGAVFLGFLGAAGSTMGAASMTLTVQARQL	102
AVGIGAVFLGFLGAAGSTMGAASVTLTVQARQL	103
AVGIGAVFLGFLGAAGSTMGARSMTLTVQARLL	104
AVGIGAVFLGFLGAAGSTMGATSITLTVQARQL	105
AVGIGAVFLGFLGAAGSTMGAVSITLTVQARQL	106
AVGIGAVFLGFLGAEGSTMGAASLTLTVQARQL	107
AVGIGAVFLGFLGFAGSTMGAASVTLTVQARQL	108
AVGIGAVFLGFLGVAGSTMGAASITLTVQARQL	109
AVGIGAVFLGFLGVAGSTMGAASMTLTVQARQL	110
AVGIGAVFLGILGAAGSTMGAASITLTVQARQL	111
AVGIGAVFLGSLGAAGSTMGAASITLTVQARQL	112
AVGIGAVFLRFLGAAGSTMGAASITLTVQARQL	113
AVGIGAVFVGFLGAAGSTMGAASITLTVQARQL	114
AVGIGAVIFGFLGAAGSTMGAASITLTVQARQL	115
AVGIGAVIFGFLGAAGSTMGAASLTLTVQARQL	116
AVGIGAVILGFLGAAGSTMGAASITLTVQARQL	117
AVGIGAVLFGFLGAAGSTMGAASITLTVQARQL	118
AVGIGAVLFGFLGAAGSTMGAASITLTVQARQV	119
AVGIGAVLFGFLGAAGSTMGAASLTLTVQARQL	120
AVGIGAVLIGFLGAAGSTMGAASITLTVQARQL	121
AVGIGAVLLGFLGAAGSTMGAASITLTVQARQL	122
AVGIGAVLLGFLGAAGSTMGAASLTLTVQARQL	123
AVGIGAVLLGFLGAAGSTMGAASMTLTVQARQL	124
AVGIGAVLLGFLGAAGSTMGAASVTLTVQARQL	125
AVGIGAVLLGFLGAAGSTMGVASMTLTVQARQL	126
AVGIGAVLLGFLGTAGSTMGAASITLTVQARQL	127
AVGIGAVLLGFLGTAGSTMGAASMTLTVQARQL	128
AVGIGAVLVGFLGAAGSTMGAASITLTVQARQL	129
AVGIGAVSLGFLGAAGSTMGAASMTLTVQARLL	130
AVGIGIMIFGFLGAAGSTMGAASITLTVQARQL	131
AVGIGIMIFGFLGAARSTMGAASITLTVQARQL	132
AVGIGTMIFGFLGAAGSTMGAASITLTVQARQL	133
AVGIGVLLGFLGAAGSTMGAASMTLTVQARQL	134
AVGIGVVFFGFLGAAGSTMGAASITLTVQARQL	135
AVGLAAVFFGFLGAAGSTMGAASITLTVQARQL	136
AVGLEAVFLGFLGAAGSTMGAASVTLTVQARQL	137
AVGLGAAFLGFLGAAGSTMGAASITLTVQARQL	138
AVGLGAFFLGFLGAAGSTMGAASITLTVQARQL	139
AVGLGAFFLGFLGVAGSTMGAASITLTVQARQL	140

AVGLGAI FIGFLGAAGSTMGAASITLTVQARQL	141
AVGLGALFFGFLGAAGSTMGAASITLTVQARQL	142
AVGLGALFIGFLGAAGSTMGAASITLTVQARQL	143
AVGLGALFIGFLGAAGSTMGAASMTLTVQARQL	144
AVGLGALFIGFLGAAGSTMGAASVTLTVQARQL	145
AVGLGALFLGFLGAAGSTMGAASITLTVQARQL	146
AVGLGALFLGFLGAAGSTMGAASLTLTVQARQL	147
AVGLGALFLGFLGAAGSTMGAASMTLTVQARQL	148
AVGLGALFLGFLGAAGSTMGAASVTLTVQARQL	149
AVGLGALFLGFLGAAGSTMGARSMTLTVQARQL	150
AVGLGALFLGFLGGAGSTMGAASLTLTVQARQL	151
AVGLGALLFGFLGAAGSTMGAASITLTVQARQL	152
AVGLGAMFIGFLGAAGSTMGAASVTLTVQARQL	153
AVGLGAMFLGFLGAAGSTMGAASITLTVQARQL	154
AVGLGAMFLGFLGAAGSTMGAASLTLTVQARQL	155
AVGLGAMFLGFLGAAGSTMGAASMTLTVQARQL	156
AVGLGAMFLGFLGAAGSTMGAASVTLTVQARQL	157
AVGLGAMIFGFLGAAGSTMGAASLTLTVQARQL	158
AVGLGAVFFGFLGAAGSTMGAASITLTVQARQL	159
AVGLGAVFIGFLGAAGSTMGAASITLTVQARQL	160
AVGLGAVFIGFLGAAGSTMGAASMTLTVQARQL	161
AVGLGAVFIGFLGAAGSTMGAASVTLTVQARQL	162
AVGLGAVFLEFLGAAGSTMGAASVTLTVQARQL	163
AVGLGAVFLGFLGAAGSTMGAASITLTVQARQL	164
AVGLGAVFLGFLGAAGSTMGAASLTLTVQARQL	165
AVGLGAVFLGFLGAAGSTMGAASMTLTVQARQL	166
AVGLGAVFLGFLGAAGSTMGAASVTLTVQARQL	167
AVGLGAVFLGFLGAAGSTMGARSITLTVQARQL	168
AVGLGAVFLGFLGTAGSTMGAASITLTVQARQL	169
AVGLGAVFLGSLGAAGSTMGAASITLTVQARQL	170
AVGLGAVFQGFLGAAGSTMGAASITLTVQARQL	171
AVGLGAVIFGFLGAAGSTMGAASITLTVQARQL	172
AVGLGAVLFGFLGAAGSTMGAASITLTVQARQL	173
AVGLGAVLLGFLGAAGSTMGAASITLTVQARQL	174
AVGLGAVLLGFLGTAGSTMGAASITLTVQARQL	175
AVGLGVAFLGFLGAAGSTMGAASITLTVQARQL	176
AVGMAAVFFGFLGAAGSTMGAASITLTVQARQL	177
AVGMAAVFIGFLGAAGSTMGAASITLTVQARQL	178
AVGMAAVFLGFLGTAGSTMGAASLTLTVQARQL	179
AVGMGAFFLGFLGAAGSTMGAASITLTVQARQL	180
AVGMGAFFLGFLGAAGSTMGAASLTLTVQARQL	181
AVGMGALFLGFLGAAGSTMGAASITLTVQARQL	182
AVGMGALFLGFLGAAGSTMGAASLTLTVQARQL	183
AVGMGALFLGFLGAAGSTMGAASMTLTVQARQL	184
AVGMGALFLGFLGAAGSTMGAASVTLTVQARQL	185
AVGMGALFLGFLGAAGSTMGAVSMTLTVQARQL	186

AVGMGALFLGFLGTAGSTMGAVSMTLTVQARQL	187
AVGMGALFLGFLSAAGSTMGAASITLTVQARQL	188
AVGMGAMFLGFLAAAGSTMGAASLTLTVQARQL	189
AVGMGAMILGFLSAAGSTMGAASITLTVQARQL	190
AVGMGASFLGFLGAAGSTMGAASITLTVQARQL	191
AVGMGAVFLGFLGAAGSTMGAASITLTVQARQL	192
AVGMGAVFLGFLSAAGSTMGAASITLTVQARQL	193
AVGMGAVLLGFLGAAGSTMGAASITLTVQARQL	194
AVGVGALFLGFLSAAGSTMGAASITLTVQARQL	195
AVGVGALLIGFLGAAGSTMGAASMTLTVQARQL	196
AVGVGAMIFGFLGAAGSTMGAASITLTVQARQL	197
AVGVGAMILGFLGAAGSTMGAASITLTVQARQL	198
VGIGALFLGFLGAAG	199
VGLGAVFLGFLGAAG	200
AAGIGAVLLGFLGAAG	201
AAGIGAVLPGFLGAAG	202
AAGIGAVLPGFLGAAR	203
AAGLGAVFLGFLGAAG	204
AIGIGAMFLGFLGAAG	205
AIGIGAVFIGFLGAAG	206
AIGIGAVFLGFLGAAG	207
AIGIGAVFLGFLGTAG	208
AIGIGAVVLGFLGTAG	209
AIGLGAFLGFLGAAG	210
AIGLGAALLGFLGAAG	211
AIGLGALFLGFLGAAG	212
AIGLGAMFLGFLGAAG	213
AIGLGAMFLGFLGAAG	214
AIGLGAMFLGFLGTAG	215
AIGLGAVFIGFLGAAG	216
AIGLGAVFLGFLGAAG	217
AIGLGAVFXGFLGAAG	218
AMGIGAVFLGFLGAAG	219
AMGIGAVLLGFLGAPG	220
AVGIGAALLGFLGAAG	221
AVGIGAFFLGAAG	222
AVGIGAFVLGFLGAAG	223
AVGIGALFLGFLAAAG	224
AVGIGALFLGFLGAAG	225
AVGIGALFLGFLGPAG	226
AVGIGALFLGFLGTAG	227
AVGIGALIFGFLGAAG	228
AVGIGALLFGFLGAAG	229
AVGIGALLFGSLGAAG	230
AVGIGALLIGFLGAAG	231
AVGIGALLLGFLGAAG	232

AVGIGALLVGFLGAAG	233
AVGIGALLVGFLGTAG	234
AVGIGALSLGFLGAAG	235
AVGIGALVFGFLGAAG	236
AVGIGAMFLGFLGAAG	237
AVGIGAMFLGFLGMAG	238
AVGIGAMFLGILGAAG	239
AVGIGAMFLGILGTAG	240
AVGIGAMIFGFLGAAG	241
AVGIGAMIFGFLGAAR	242
AVGIGAMIFGFLGAPG	243
AVGIGAMIFGFSGAAG	244
AVGIGAMILGFLGAAG	245
AVGIGAMLFGFLGAAG	246
AVGIGAVFFGFLGAAG	247
AVGIGAVFIGFLGAAG	248
AVGIGAVFIGFLSAAG	249
AVGIGAVFLGFLAAAG	250
AVGIGAVFLGFLATAG	251
AVGIGAVFLGFLGAAG	252
AVGIGAVFLGFLGAEG	253
AVGIGAVFLGFLGFAG	254
AVGIGAVFLGFLGVAG	255
AVGIGAVFLGILGAAG	256
AVGIGAVFLGSLGAAG	257
AVGIGAVFLRFLGAAG	258
AVGIGAVFVGFLGAAG	259
AVGIGAVIFGFLGAAG	260
AVGIGAVILGFLGAAG	261
AVGIGAVLFGFLGAAG	262
AVGIGAVLIGFLGAAG	263
AVGIGAVLLGFLGAAG	264
AVGIGAVLLGFLGTAG	265
AVGIGAVLVGFLGAAG	266
AVGIGAVSLGFLGAAG	267
AVGIGIMIFGFLGAAG	268
AVGIGIMIFGFLGAAR	269
AVGIGTMIFGFLGAAG	270
AVGIGVLLLGFLGAAG	271
AVGIGVVFFGFLGAAG	272
AVGLAAVFFGFLGAAG	273
AVGLEAVFLGFLGAAG	274
AVGLGAAFLGFLGAAG	275
AVGLGAFFLGFLGAAG	276
AVGLGAFFLGFLGVAG	277
AVGLGAIFIGFLGAAG	278

AVGLGALFFGFLGAAG	279
AVGLGALFIGFLGAAG	280
AVGLGALFLGFLGAAG	281
AVGLGALFLGFLGGAG	282
AVGLGALLFGFLGAAG	283
AVGLGAMFIGFLGAAG	284
AVGLGAMFLGFLGAAG	285
AVGLGAMIFGFLGAAG	286
AVGLGAVFFGFLGAAG	287
AVGLGAVFIGFLGAAG	288
AVGLGAVFLEFLGAAG	289
AVGLGAVFLGFLGAAG	290
AVGLGAVFLGFLGTAG	291
AVGLGAVFLGSLGAAG	292
AVGLGAVFQGFLGAAG	293
AVGLGAVIFGFLGAAG	294
AVGLGAVLFGFLGAAG	295
AVGLGAVLLGFLGAAG	296
AVGLGAVLLGFLGTAG	297
AVGLGVAFLGFLGAAG	298
AVGMAAVFFGFLGAAG	299
AVGMAAVFIGFLGAAG	300
AVGMAAVFLGFLGTAG	301
AVGMGAFFLGFLGAAG	302
AVGMGALFLGFLGAAG	303
AVGMGALFLGFLGTAG	304
AVGMGALFLGFLSAAG	305
AVGMGAMFLGFLAAAG	306
AVGMGAMILGFLSAAG	307
AVGMGASFLGFLGAAG	308
AVGMGAVFLGFLGAAG	309
AVGMGAVFLGFLSAAG	310
AVGMGAVLLGFLGAAG	311
AVGVGALFLGFLSAAG	312
AVGVGALLIGFLGAAG	313
AVGVGAMIFGFLGAAG	314
AVGVGAMILGFLGAAG	315
VGIGALFLGFL	316
VGLGAVFLGFL	317
AAGIGAVLLGFL	318
AAGIGAVLPGFL	319
AAGLGAVFLGFL	320
AIGIGAMFLGFL	321
AIGIGAVFIGFL	322
AIGIGAVFLGFL	323
AIGIGAVVLGFL	324

AIGLGAAFLGFL	325
AIGLGAALLGFL	326
AIGLGALFLGFL	327
AIGLGAMFLGFL	328
AIGLGAVFIGFL	329
AIGLGAVFLGFL	330
AIGLGAVFXGFL	331
AMGIGAVFLGFL	332
AMGIGAVLLGFL	333
AVGIGAALLGFL	334
AVGIGAFFLGFL	335
AVGIGAFVLGFL	336
AVGIGALFLGFL	337
AVGIGALIFGFL	338
AVGIGALLFGFL	339
AVGIGALLFGSL	340
AVGIGALLIGFL	341
AVGIGALLLGFL	342
AVGIGALLVGFL	343
AVGIGALSLGFL	344
AVGIGALVFGFL	345
AVGIGAMFLGFL	346
AVGIGAMFLGIL	347
AVGIGAMIFGFL	348
AVGIGAMIFGFS	349
AVGIGAMILGFL	350
AVGIGAMLFGFL	351
AVGIGAVFFGFL	352
AVGIGAVFIGFL	353
AVGIGAVFLGFL	354
AVGIGAVFLGIL	355
AVGIGAVFLGSL	356
AVGIGAVFLRFL	357
AVGIGAVFVGFL	358
AVGIGAVIFGFL	359
AVGIGAVILGFL	360
AVGIGAVLFGFL	361
AVGIGAVLIGFL	362
AVGIGAVLLGFL	363
AVGIGAVLVGFL	364
AVGIGAVSLGFL	365
AVGIGIMIFGFL	366
AVGIGTMIFGFL	367
AVGIGVLLLGL	368
AVGIGVVFFGFL	369
AVGLAAVFFGFL	370

AVGLEAVFLGFL	371
AVGLGAAFLGFL	372
AVGLGAFFLGFL	373
AVGLGAIFIGFL	374
AVGLGALFFGFL	375
AVGLGALFIGFL	376
AVGLGALFLGFL	377
AVGLGALLFGFL	378
AVGLGAMFIGFL	379
AVGLGAMFLGFL	380
AVGLGAMIFGFL	381
AVGLGAVFFGFL	382
AVGLGAVFIGFL	383
AVGLGAVFLEFL	384
AVGLGAVFLGFL	385
AVGLGAVFLGSL	386
AVGLGAVFQGFL	387
AVGLGAVIFGFL	388
AVGLGAVLFGFL	389
AVGLGAVLLGFL	390
AVGLGVAFLGFL	391
AVGMAAVFFGFL	392
AVGMAAVFIGFL	393
AVGMAAVFLGFL	394
AVGMGAFFLGFL	395
AVGMGALFLGFL	396
AVGMGAMFLGFL	397
AVGMGAMILGFL	398
AVGMGASFLGFL	399
AVGMGAVFLGFL	400
AVGMGAVLLGFL	401
AVGVGALFLGFL	402
AVGVGALLIGFL	403
AVGVGAMIFGFL	404
AVGVGAMILGFL	405
AVGIGALF	406
GALFLGFLG	407
GAVFLGFLG	409
GAMFLGFLG	410
GAVLLGFLG	411
GAFFLGFLG	412
GAMIFGFLG	413
GALLFGFLG	414

CLAIMS

1. A method of treating or preventing the symptoms of a disease or disorder related to an inappropriate or detrimental T cell response, comprising administering to an individual in need thereof a therapeutically effective amount of a pharmaceutical composition comprising as an active ingredient an isolated peptide derived from HIV gp41 fusion peptide domain or fragments, analogs, variants, conjugates, derivatives and salts thereof.
2. The method of claim 1, wherein the HIV is HIV-1.
3. The method of claim 1 wherein the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof.
4. The method of claim 1, wherein the fusion peptide has the amino acid sequence AVGIGALFLGFLGAAGSTMGARSMTLTVQARQL (SEQ ID NO:1).
5. The method of claim 1, wherein the fusion peptide has the amino acid sequence AEGIGALFLGFLGAAGSTMGARSMTLTVQARQL (SEQ ID NO:2).
6. The method of claim 1, wherein the fusion peptide has the amino acid sequence AVGIGALFLGFLGAAGSTMGARSMTLTVQARQL, wherein the underlined amino acid residues at positions 3, 6, 9 and 11 are of the "D" isomer configuration (SEQ ID NO:6).
7. The method of claim 1, wherein the fusion peptide has the amino acid sequence AVGIGALF (SEQ ID NO:406).
8. The method of claim 1, wherein the fusion peptide has the amino acid sequence GALFLGFLG (SEQ ID NO:407).
9. The method of claim 1, wherein the fusion peptide has the amino acid sequence GALFLGFLG, wherein the underlined amino acid residue at position 2 is of the "D" isomer configuration (SEQ ID NO:408).
10. The method of claim 1, wherein the fusion peptide is selected from the group consisting of: GAVFLGFLG (SEQ ID NO:409), GAMFLGFLG (SEQ ID NO:410), GAVLLGFLG (SEQ ID NO:411), GAFFLGFLG (SEQ ID NO:412), GAMIFGFLG (SEQ ID NO:413) and GALLFGFLG (SEQ ID NO:414).

11. The method of claim 1 wherein the fusion peptide comprises both D and L amino acids.
12. The method of claim 1 wherein the T cell pathology is an autoimmune disease.
13. The method of claim 12, wherein the autoimmune disease is selected from the group consisting of: multiple sclerosis, rheumatoid arthritis, juvenile rheumatoid arthritis, autoimmune neuritis, systemic lupus erythematosus, psoriasis, Type I diabetes, Sjogren's disease, thyroid disease, myasthenia gravis, sarcoidosis, autoimmune uveitis, inflammatory bowel disease (Crohn's and ulcerative colitis), and autoimmune hepatitis.
14. The method of claim 8 wherein the T cell pathology is an inflammatory disease.
15. The method of claim 8 wherein the T cell pathology is selected from graft rejection and graft versus host disease.
16. A method of treating or preventing the symptoms of a disease or disorder related to an inappropriate or detrimental T cell response, comprising administering to an individual in need thereof a therapeutically effective amount of a pharmaceutical composition comprising as an active ingredient a peptide, or salt thereof, capable of inhibiting T cell activation having the formula (I):
- $$X_1-AA_1-AA_2-AA_3-AA_4-AA_5-AA_6-AA_7-AA_8-AA_9-X_2 \quad (I)$$
- wherein:
- X₁ represents an N-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence alanine-valine-glycine, or may be absent;
- AA₁, AA₂, AA₆ and AA₉ each independently represent an alanine or glycine amino acid residue;
- AA₃ represents a phenylalanine, isoleucine, leucine, valine or methionine amino acid residue;
- AA₄, AA₅, AA₇ and AA₈ each independently represent a phenylalanine, isoleucine, leucine, valine, methionine or serine amino acid residue, wherein no more than two amino acid residues of AA₄, AA₅, AA₇ and AA₈ are identical, with

the exception that any three of AA₄, AA₅, AA₇ and AA₈ may be leucine amino acid residues;

5 X₂ represents a C-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence valine-glutamine-alanine, or may be absent;

and wherein each amino acid can be of either L or D form and the peptide is no more than 30 amino acid residues in length.

10 17. The method of claim 16 wherein the fusion peptide comprises both D and L amino acids.

18. The method of claim 16 wherein said peptide does not contain more than one serine residue.

19. The method of claim 16 wherein the T cell pathology is an autoimmune disease.

15 20. The method of claim 19, wherein the autoimmune disease is selected from the group consisting of: multiple sclerosis, rheumatoid arthritis, juvenile rheumatoid arthritis, autoimmune neuritis, systemic lupus erythematosus, psoriasis, Type I diabetes, Sjogren's disease, thyroid disease, myasthenia gravis, sarcoidosis, autoimmune uveitis, inflammatory bowel disease (Crohn's and ulcerative colitis), and autoimmune hepatitis.

20 21. The method of claim 16 wherein the T cell pathology is an inflammatory disease.

22. The method of claim 16 wherein the T cell pathology is selected from graft rejection and graft versus host disease.

25 ~~23. A method of inhibiting T-cell activation comprising administering to an individual in need thereof a therapeutically effective amount of a pharmaceutical composition comprising as an active ingredient an isolated peptide derived from HIV gp41 fusion peptide domain or fragments, analogs, variants, conjugates, derivatives and salts thereof.~~

24. The method of claim 23 wherein the HIV is HIV-1.

30 25. The method of claim 23 wherein the fusion peptide has an amino acid sequence as set forth in any one of SEQ ID NOS:1, 2, and 6-414 or fragments, analogs, variants, conjugates, derivatives and salts thereof.

26. The method of claim 23 wherein the fusion peptide comprises both D and L amino acids.

27. A method of inhibiting T cell activation comprising administering to an individual in need thereof a therapeutically effective amount of a pharmaceutical composition comprising as an active ingredient a peptide, or salt thereof, capable of inhibiting T cell activation having the formula (I):



wherein:

10 X_1 represents an N-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence alanine-valine-glycine, or may be absent;

AA_1 , AA_2 , AA_6 and AA_9 each independently represent an alanine or glycine amino acid residue;

15 AA_3 represents a phenylalanine, isoleucine, leucine, valine or methionine amino acid residue;

AA_4 , AA_5 , AA_7 and AA_8 each independently represent a phenylalanine, isoleucine, leucine, valine, methionine or serine amino acid residue, wherein no more than two amino acid residues of AA_4 , AA_5 , AA_7 and AA_8 are identical, with the exception that any three of AA_4 , AA_5 , AA_7 and AA_8 may be leucine amino acid residues;


20 X_2 represents a C-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence valine-glutamine-alanine, or may be absent;

and wherein each amino acid can be of either L or D form and the peptide is no more than 30 amino acid residues in length.

28. The method of claim 27 wherein the fusion peptide comprises both D and L amino acids.

30 29. The method of claim 27 wherein said peptide does not contain more than one serine residue.

30. A peptide capable of inhibiting T cell activation having the formula (I):



wherein:

5 X_1 represents an N-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide having the sequence alanine-valine-glycine, or may be absent;

AA_1 , AA_2 , AA_6 and AA_9 each independently represent an alanine or glycine amino acid residue;

10 AA_3 represents a phenylalanine, isoleucine, leucine, valine or methionine amino acid residue;

AA_4 , AA_5 , AA_7 and AA_8 each independently represent a phenylalanine, isoleucine, leucine, valine, methionine or serine amino acid residue, wherein no more than two amino acid residues of AA_4 , AA_5 , AA_7 and AA_8 are identical, with
15 the exception that any three of AA_4 , AA_5 , AA_7 and AA_8 may be leucine amino acid residues;

X_2 represents a C-terminal blocking group, an amino acid sequence of up to about 20 amino acid residues in length wherein at least 50% of the amino acid residues are hydrophobic and wherein said sequence does not comprise a peptide
20 having the sequence valine-glutamine-alanine, or may be absent;

and wherein each amino acid can be of either L or D form and the peptide is no more than 30 amino acid residues in length.

31. The peptide of claim 30 wherein said peptide does not contain more than one serine residue.

25 32. The peptide of claim 30 having the amino acid sequence GALFLGFLG (SEQ ID NO:407).

33. The peptide of claim 30 wherein said peptide is selected from the group consisting of: GAVFLGFLG (SEQ ID NO:409), GAMFLGFLG (SEQ ID NO:410), GAVLLGFLG (SEQ ID NO:411), GAFFLGFLG (SEQ ID NO:412), GAMIFGFLG (SEQ ID NO:413) and GALLFGFLG (SEQ ID NO:414).
30

34. The peptide of claim 30, wherein said peptide comprises both "L" and "D" amino acids.
35. The peptide of claim 34 having the amino acid sequence GALFLGFLG, wherein the underlined amino acid residue at position 2 is of the "D" isomer configuration (SEQ
5 ID NO:408).
36. A pharmaceutical compositions comprising the peptide of claim 30 and a pharmaceutically acceptable carrier or diluent.




Figure 1

2/13




Figure 2A




Figure 2B



Figure 2C




Figure 2D

3/13




Figure 3A




Figure 3B




Figure 3C

4/13




Figure 3D




Figure 3E




Figure 3F

5/13




Figure 4

6/13




Figure 5A




Figure 5B

7/13




Figure 6A




Figure 6B

8/13




Figure 6C





Figure 6D

9/13

HIV Infection



Immunotherapy




Figure 7

10/13




Figure 8

11/13




Figure 9A




Figure 9B




Figure 9C

12/13




Figure 9D

13/13




Figure 10




Figure 11

yeda 047 pct new.ST25.txt
SEQUENCE LISTING

<110> Yeda Research and Development Co. Ltd.
Shai, Yechiel
Cohen, Irun R.
Quintana, Francisco J.
Gerber, Doron J.

<120> HIV-1 gp41 Fusion Peptides for Immunomodulation

<130> YEDA 047 PCT

<160> 414

<170> PatentIn version 3.3

<210> 1

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 1

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Arg Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 2

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 2

Ala Glu Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Arg Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 3

<211> 15

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic peptide

yeda 047 pct new.ST25.txt

<400> 3

Leu Leu Lys Leu Leu Lys Lys Leu Leu Lys Lys Leu Leu Lys Leu
1 5 10 15

<210> 4

<211> 24

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of human 60 kDa heat shock protein

<400> 4

Val Leu Gly Gly Gly Cys Ala Leu Leu Arg Cys Ile Pro Ala Leu Asp
1 5 10 15

Ser Leu Thr Pro Ala Asn Glu Asp
20

<210> 5

<211> 15

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of M. tuberculosis 65 kDa heat shock protein

<400> 5

Glu Glu Ser Asn Thr Phe Gly Leu Gln Leu Glu Leu Thr Glu Gly
1 5 10 15

<210> 6

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic peptide

<220>

<221> MISC_FEATURE

<222> (4)..(4)

<223> D-Ile

<220>

<221> MISC_FEATURE

<222> (8)..(8)

<223> D-Phe

<220>

<221> MISC_FEATURE

<222> (11)..(11)

<223> D-Phe

<220>

<221> MISC_FEATURE

<222> (14)..(14)

<223> D-Ala

<400> 6

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Arg Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 7
 <211> 32
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 7

Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly Ser
 1 5 10 15

Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln Leu
 20 25 30

<210> 8
 <211> 32
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 8

Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly Ser
 1 5 10 15

Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln Leu
 20 25 30

<210> 9
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 9

Ala Ala Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 10
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 10

Ala Ala Gly Ile Gly Ala Val Leu Pro Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 11
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 11

Ala Ala Gly Ile Gly Ala Val Leu Pro Gly Phe Leu Gly Ala Ala Arg
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 12
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 12

Ala Ala Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 13
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 13

Ala Ile Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 14
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 14

Ala Ile Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 15
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 15

Ala Ile Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 16

yeda 047 pct new.ST25.txt

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 16

Ala Ile Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 17

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 17

Ala Ile Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 18

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 18

Ala Ile Gly Ile Gly Ala Val Val Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 19

<211> 33

<212> PRT

<213> Artificial sequence yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 19

Ala Ile Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 20

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 20

Ala Ile Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 21

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 21

Ala Ile Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Cys Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 22

<211> 33

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 22

Ala Ile Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Val Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 23

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 23

Ala Ile Gly Leu Gly Ala Ala Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 24

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 24

Ala Ile Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 25

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 25

Ala Ile Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Val

<210> 26

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 26

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 27

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 27

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 28

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 28

yeda 047 pct new.ST25.txt

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Val

<210> 29
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 29

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 30
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 30

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Arg Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 31
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 31

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 Page 10

yeda 047 pct new.ST25.txt

1 5 10 15

Ser Thr Met Gly Ala Arg Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 32
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 32

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 33
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 33

Ala Ile Gly Leu Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 34
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 34

Ala Ile Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

yeda 047 pct new.ST25.txt

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 35
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 35

Ala Ile Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Val

<210> 36
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<220>
 <221> misc_feature
 <222> (9)..(9)
 <223> Xaa can be any naturally occurring amino acid

<400> 36

Ala Ile Gly Leu Gly Ala Val Phe Xaa Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 37
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 37

yeda 047 pct new.ST25.txt

Ala Met Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 38
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 38

Ala Met Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Pro Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 39
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 39

Ala Val Gly Ile Gly Ala Ala Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Ala Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 40
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 40

Ala Val Gly Ile Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly

```
yeda 047 pct new.ST25.txt
```

1 5 yeda 047 pct new.S125.txt 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln

20 25 30

Leu

<210>	41
<211>	33
<212>	PRT
<213>	Artificial sequence

```
<220>
<223> Peptides derived of HIV-1 gp41
```

<400> 41

Ala Val Gly Ile Gly Ala Phe Val Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210>	42
<211>	33
<212>	PRT
<213>	Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 42

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Ala Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210>	43
<211>	33
<212>	PRT
<213>	Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 43

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Ala Ala Ala Gly
1 5 10 15

yeda 047 pct new.ST25.txt

Ser Thr Met Gly Ala Arg Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 44
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 44

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Ala Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 45
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 45

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 46
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 46

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln

20 yeda 047 pct new.ST25.txt
25 30

Leu

<210> 47
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 47

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Leu
20 25 30

Leu

<210> 48
<211> 32
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 48

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

<210> 49
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 49

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 50
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41
 <400> 50

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 51
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41
 <400> 51

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Cys Thr Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 52
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41
 <400> 52

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Pro Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 53

yeda 047 pct new.ST25.txt

<211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 53

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 54
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 54

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 55
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 55

Ala Val Gly Ile Gly Ala Leu Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 56
 <211> 33
 <212> PRT

<213> Artificial sequence yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 56

Ala Val Gly Ile Gly Ala Leu Ile Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 57

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 57

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 58

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 58

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 59

<211> 33

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 59

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Ser Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Leu
 20 25 30

Leu

<210> 60

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 60

Ala Val Gly Ile Gly Ala Leu Leu Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 61

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 61

Ala Val Gly Ile Gly Ala Leu Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 62

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 62

Ala Val Gly Ile Gly Ala Leu Leu Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 63

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 63

Ala Val Gly Ile Gly Ala Leu Leu Val Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 64

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 64

Ala Val Gly Ile Gly Ala Leu Leu Val Gly Phe Leu Gly Thr Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 65

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 65

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Ala Leu Ser Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Leu
 20 25 30

Leu

<210> 66
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 66

Ala Val Gly Ile Gly Ala Leu Val Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 67
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 67

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 68
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 68

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly

yeda 047 pct new.ST25.txt
 1 5 10 15
 Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 69
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 69

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Ala Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 70
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 70

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 71
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 71

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

yeda 047 pct new.ST25.txt

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 72
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 72

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Met Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 73
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 73

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Ile Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 74
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 74

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Ile Leu Gly Thr Ala Gly
 1 5 10 15

Ser Ala Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln

20 yeda 047 pct new.ST25.txt
25 30

Leu

<210> 75
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 75

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 76
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 76

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 77
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 77

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

yeda 047 pct new.ST25.txt

Leu

<210> 78
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 78

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Arg
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 79
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 79

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Pro Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 80
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 80

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Ser Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 81
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 81

Ala Val Gly Ile Gly Ala Met Ile Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 82
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 82

Ala Val Gly Ile Gly Ala Met Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 83
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 83

Ala Val Gly Ile Gly Ala Met Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 84
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41
 <400> 84

Ala Val Gly Ile Gly Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 85
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41
 <400> 85

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Leu Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 86
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41
 <400> 86

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 87

yeda 047 pct new.ST25.txt

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 87

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 88

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 88

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 89

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 89

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 90

<211> 33

<212> PRT

<213> Artificial sequence yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 90

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Ala Ala Ala Gly
1 5 10 15

Ser Ser Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 91

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 91

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Ala Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 92

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 92

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Ala Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Val

<210> 93

<211> 33

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 93

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Ala Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 94

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 94

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ala Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 95

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 95

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Ile Gln Ala Arg Gln
 20 25 30

Leu

<210> 96

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 96

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 97

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 97

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 98

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 98

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Leu
 20 25 30

Leu

<210> 99

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 99

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 100
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 100

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Arg
 20 25 30

Leu

<210> 101
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 101

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Val Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 102
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 102

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly

Leu

<400> 103

Leu

<400> 104

Leu

<400> 105

Page 34

yeda 047 pct new.ST25.txt

Ser Thr Met Gly Ala Thr Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 106
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 106

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Val Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 107
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 107

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Glu Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 108
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 108

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Phe Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln

20 yeda 047 pct new.ST25.txt
25 30

Leu

<210> 109
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 109

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Val Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 110
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 110

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Val Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 111
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 111

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Ile Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

yeda 047 pct new.ST25.txt

Leu

<210> 112
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 112

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Ser Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 113
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 113

Ala Val Gly Ile Gly Ala Val Phe Leu Arg Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 114
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 114

Ala Val Gly Ile Gly Ala Val Phe Val Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 115
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 115

Ala Val Gly Ile Gly Ala Val Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 116
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 116

Ala Val Gly Ile Gly Ala Val Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 117
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 117

Ala Val Gly Ile Gly Ala Val Ile Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 118
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 118

Ala Val Gly Ile Gly Ala Val Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 119
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 119

Ala Val Gly Ile Gly Ala Val Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Val

<210> 120
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 120

Ala Val Gly Ile Gly Ala Val Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 121

yeda 047 pct new.ST25.txt

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 121

Ala Val Gly Ile Gly Ala Val Leu Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 122

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 122

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 123

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 123

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 124

<211> 33

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 124

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 125

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 125

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 126

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 126

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Val Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 127

<211> 33

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 127

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 128

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 128

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 129

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 129

Ala Val Gly Ile Gly Ala Val Leu Val Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 130

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 130

Ala Val Gly Ile Gly Ala Val Ser Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Leu
20 25 30

Leu

<210> 131

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 131

Ala Val Gly Ile Gly Ile Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 132

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 132

Ala Val Gly Ile Gly Ile Met Ile Phe Gly Phe Leu Gly Ala Ala Arg
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 133

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 133

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Thr Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 134
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 134

Ala Val Gly Ile Gly Val Leu Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Ala Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 135
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 135

Ala Val Gly Ile Gly Val Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 136
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 136

Ala Val Gly Leu Ala Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 Page 44

yeda 047 pct new.ST25.txt 15
 1 5 10
 Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 137
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 137

Ala Val Gly Leu Glu Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 138
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 138

Ala Val Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 139
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 139

Ala Val Gly Leu Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

yeda 047 pct new.ST25.txt

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 140
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 140

Ala Val Gly Leu Gly Ala Phe Phe Leu Gly Phe Leu Gly Val Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 141
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 141

Ala Val Gly Leu Gly Ala Ile Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 142
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 142

Ala Val Gly Leu Gly Ala Leu Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 Page 46

20 yeda 047 pct new.ST25.txt
25 30

Leu

<210> 143
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 143

Ala Val Gly Leu Gly Ala Leu Phe Ile Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 144
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 144

Ala Val Gly Leu Gly Ala Leu Phe Ile Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 145
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 145

Ala Val Gly Leu Gly Ala Leu Phe Ile Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

yeda 047 pct new.ST25.txt

Leu

<210> 146
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 146

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 147
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 147

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 148
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 148

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 149
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 149

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 150
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 150

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Arg Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 151
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 151

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Gly Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 152
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 152

Ala Val Gly Leu Gly Ala Leu Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 153
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 153

Ala Val Gly Leu Gly Ala Met Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 154
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 154

Ala Val Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 155

yeda 047 pct new.ST25.txt

<211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 155

Ala Val Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 156
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 156

Ala Val Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 157
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 157

Ala Val Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 158
 <211> 33
 <212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 158

Ala Val Gly Leu Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 159

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 159

Ala Val Gly Leu Gly Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 160

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 160

Ala Val Gly Leu Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 161

<211> 33

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 161

Ala Val Gly Leu Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 162

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 162

Ala Val Gly Leu Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 163

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 163

Ala Val Gly Leu Gly Ala Val Phe Leu Glu Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 164

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 164

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 165

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 165

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 166

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 166

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 167

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 167

yeda 047 pct new.ST25.txt

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 168
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 168

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Arg Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 169
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 169

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 170
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 170

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Ser Leu Gly Ala Ala Gly

yeda 047 pct new.ST25.txt
 1 5 10 15
 Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 171
 <211> 33
 <212> PRT
 <213> Artificial sequence
 <220>
 <223> Peptides derived of HIV-1 gp41
 <400> 171

Ala Val Gly Leu Gly Ala Val Phe Gln Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15
 Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 172
 <211> 33
 <212> PRT
 <213> Artificial sequence
 <220>
 <223> Peptides derived of HIV-1 gp41
 <400> 172

Ala Val Gly Leu Gly Ala Val Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15
 Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 173
 <211> 33
 <212> PRT
 <213> Artificial sequence
 <220>
 <223> Peptides derived of HIV-1 gp41
 <400> 173

Ala Val Gly Leu Gly Ala Val Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

yeda 047 pct new.ST25.txt

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 174
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 174

Ala Val Gly Leu Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 175
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 175

Ala Val Gly Leu Gly Ala Val Leu Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 176
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 176

Ala Val Gly Leu Gly Val Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 Page 57

20 yeda 047 pct new.ST25.txt
25 30

Leu

<210> 177
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 177

Ala Val Gly Met Ala Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 178
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 178

Ala Val Gly Met Ala Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

Leu

<210> 179
<211> 33
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 179

Ala Val Gly Met Ala Ala Val Phe Leu Gly Phe Leu Gly Thr Ala Gly
1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
20 25 30

yeda 047 pct new.ST25.txt

Leu

<210> 180
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 180

Ala Val Gly Met Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 181
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 181

Ala Val Gly Met Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 182
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 182

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 183
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 183

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 184
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 184

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 185
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 185

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Val Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

yeda 047 pct new.ST25.txt

<210> 186
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 186

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Val Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 187
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 187

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Val Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 188
 <211> 33
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 188

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 189

yeda 047 pct new.ST25.txt

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 189

Ala Val Gly Met Gly Ala Met Phe Leu Gly Phe Leu Ala Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 190

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 190

Ala Val Gly Met Gly Ala Met Ile Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 191

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> ~~Peptides derived of HIV-1 gp41~~

<400> 191

Ala Val Gly Met Gly Ala Ser Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 192

<211> 33

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 192

Ala	Val	Gly	Met	Gly	Ala	Val	Phe	Leu	Gly	Phe	Leu	Gly	Ala	Ala	Gly
1				5					10					15	

Ser	Thr	Met	Gly	Ala	Ala	Ser	Ile	Thr	Leu	Thr	Val	Gln	Ala	Arg	Gln
			20					25					30		

Leu

<210> 193

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 193

Ala	Val	Gly	Met	Gly	Ala	Val	Phe	Leu	Gly	Phe	Leu	Ser	Ala	Ala	Gly
1				5					10					15	

Ser	Thr	Met	Gly	Ala	Ala	Ser	Ile	Thr	Leu	Thr	Val	Gln	Ala	Arg	Gln
			20					25					30		

Leu

<210> 194

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 194

Ala	Val	Gly	Met	Gly	Ala	Val	Leu	Leu	Gly	Phe	Leu	Gly	Ala	Ala	Gly
1				5					10					15	

Ser	Thr	Met	Gly	Ala	Ala	Ser	Ile	Thr	Leu	Thr	Val	Gln	Ala	Arg	Gln
			20					25					30		

Leu

<210> 195

<211> 33

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 195

Ala Val Gly Val Gly Ala Leu Phe Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 196

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 196

Ala Val Gly Val Gly Ala Leu Leu Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Met Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 197

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 197

Ala Val Gly Val Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 198

<211> 33

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 198

Ala Val Gly Val Gly Ala Met Ile Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

Ser Thr Met Gly Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln
 20 25 30

Leu

<210> 199

<211> 15

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 199

Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 200

<211> 15

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 200

Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 201

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 201

Ala Ala Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 202

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 202

Ala Ala Gly Ile Gly Ala Val Leu Pro Gly Phe Leu Gly Ala Ala Gly

yeda 047 pct new.ST25.txt

Ala Ile Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 208
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 208

Ala Ile Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Thr Ala Gly
1 5 10 15

<210> 209
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 209

Ala Ile Gly Ile Gly Ala Val Val Leu Gly Phe Leu Gly Thr Ala Gly
1 5 10 15

<210> 210
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 210

Ala Ile Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 211
<211> 16
<212> PRT
<213> Artificial

<220>
<223> Peptides derived of HIV-1 gp41

<400> 211

Ala Ile Gly Leu Gly Ala Ala Leu Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 212
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 212

Ala Ile Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 213

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 213

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 214

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 214

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 215

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 215

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Thr Ala Gly
1 5 10 15

<210> 216

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 216

Ala Ile Gly Leu Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 217

<211> 16

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 217

Ala Ile Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 218

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<220>

<221> misc_feature

<222> (9)..(9)

<223> Xaa can be any naturally occurring amino acid

<400> 218

Ala Ile Gly Leu Gly Ala Val Phe Xaa Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 219

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 219

Ala Met Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 220

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 220

Ala Met Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Pro Gly
 1 5 10 15

<210> 221

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 221

Ala Val Gly Ile Gly Ala Ala Leu Leu Gly Phe Leu Gly Ala Ala Gly

yeda 047 pct new.ST25.txt

1	5	10	15
---	---	----	----

<210> 222
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 222

Ala Val Gly Ile Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 223
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 223

Ala Val Gly Ile Gly Ala Phe Val Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 224
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 224

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Ala Ala Ala Gly
1 5 10 15

<210> 225
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 225

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 226
<211> 16
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 226

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Pro Ala Gly
 1 5 10 15

<210> 227
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 227

Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

<210> 228
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 228

Ala Val Gly Ile Gly Ala Leu Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 229
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 229

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 230
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 230

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Ser Leu Gly Ala Ala Gly
 1 5 10 15

<210> 231
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 231

Ala Val Gly Ile Gly Ala Leu Leu Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 232

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 232

Ala Val Gly Ile Gly Ala Leu Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 233

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 233

Ala Val Gly Ile Gly Ala Leu Leu Val Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 234

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 234

Ala Val Gly Ile Gly Ala Leu Leu Val Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

<210> 235

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 235

Ala Val Gly Ile Gly Ala Leu Ser Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 236

<211> 16

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 236

Ala Val Gly Ile Gly Ala Leu Val Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 237

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 237

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 238

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 238

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Met Ala Gly
 1 5 10 15

<210> 239

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 239

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Ile Leu Gly Ala Ala Gly
 1 5 10 15

<210> 240

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 240

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Ile Leu Gly Thr Ala Gly
 1 5 10 15

<210> 241

<211> 16

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 241

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 242

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 242

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Arg
 1 5 10 15

<210> 243

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 243

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Pro Gly
 1 5 10 15

<210> 244

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 244

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Ser Gly Ala Ala Gly
 1 5 10 15

<210> 245

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 245

Ala Val Gly Ile Gly Ala Met Ile Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 246

yeda 047 pct new.ST25.txt

<211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 246

Ala Val Gly Ile Gly Ala Met Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 247
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 247

Ala Val Gly Ile Gly Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 248
 <211> 16
 <212> PRT
 <213> Artificial

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 248

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 249
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 249

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

<210> 250
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 250

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Ala Ala Ala Gly
 1 5 10 15

yeda 047 pct new.ST25.txt

<210> 251
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 251

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Ala Thr Ala Gly
 1 5 10 15

<210> 252
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 252

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 253
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 253

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Glu Gly
 1 5 10 15

<210> 254
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 254

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Phe Ala Gly
 1 5 10 15

<210> 255
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 255

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu Gly Val Ala Gly

1	5	10	15
<210>	256		
<211>	16		
<212>	PRT		
<213>	Artificial sequence		
<220>			
<223>	Peptides derived of HIV-1 gp41		
<400>	256		
Ala Val Gly Ile Gly Ala Val Phe Leu Gly Ile Leu Gly Ala Ala Gly			
1	5	10	15
<210>	257		
<211>	16		
<212>	PRT		
<213>	Artificial sequence		
<220>			
<223>	Peptides derived of HIV-1 gp41		
<400>	257		
Ala Val Gly Ile Gly Ala Val Phe Leu Gly Ser Leu Gly Ala Ala Gly			
1	5	10	15
<210>	258		
<211>	16		
<212>	PRT		
<213>	Artificial sequence		
<220>			
<223>	Peptides derived of HIV-1 gp41		
<400>	258		
Ala Val Gly Ile Gly Ala Val Phe Leu Arg Phe Leu Gly Ala Ala Gly			
1	5	10	15
<210>	259		
<211>	16		
<212>	PRT		
<213>	Artificial sequence		
<220>			
<223>	Peptides derived of HIV-1 gp41		
<400>	259		
Ala Val Gly Ile Gly Ala Val Phe Val Gly Phe Leu Gly Ala Ala Gly			
1	5	10	15
<210>	260		
<211>	16		
<212>	PRT		
<213>	Artificial sequence		
<220>			
<223>	Peptides derived of HIV-1 gp41		
<400>	260		

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Ala Val Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 261
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 261

Ala Val Gly Ile Gly Ala Val Ile Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 262
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 262

Ala Val Gly Ile Gly Ala Val Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 263
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 263

Ala Val Gly Ile Gly Ala Val Leu Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 264
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 264

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 265
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 265

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu Gly Thr Ala Gly
1 5 10 15

<210> 266

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 266

Ala Val Gly Ile Gly Ala Val Leu Val Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 267

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 267

Ala Val Gly Ile Gly Ala Val Ser Leu Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 268

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 268

Ala Val Gly Ile Gly Ile Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
1 5 10 15

<210> 269

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 269

Ala Val Gly Ile Gly Ile Met Ile Phe Gly Phe Leu Gly Ala Ala Arg
1 5 10 15

<210> 270

<211> 16

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 270

Ala Val Gly Ile Gly Thr Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 271

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 271

Ala Val Gly Ile Gly Val Leu Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 272

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 272

Ala Val Gly Ile Gly Val Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 273

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 273

Ala Val Gly Leu Ala Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 274

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 274

Ala Val Gly Leu Glu Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 275

<211> 16

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 275

Ala Val Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 276

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 276

Ala Val Gly Leu Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 277

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 277

Ala Val Gly Leu Gly Ala Phe Phe Leu Gly Phe Leu Gly Val Ala Gly
 1 5 10 15

<210> 278

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 278

Ala Val Gly Leu Gly Ala Ile Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 279

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 279

Ala Val Gly Leu Gly Ala Leu Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 280

yeda 047 pct new.ST25.txt

<211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 280

Ala Val Gly Leu Gly Ala Leu Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 281
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 281

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 282
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 282

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu Gly Gly Ala Gly
 1 5 10 15

<210> 283
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 283

Ala Val Gly Leu Gly Ala Leu Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 284
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 284

Ala Val Gly Leu Gly Ala Met Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

yeda 047 pct new.ST25.txt

<210> 285
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 285

Ala Val Gly Leu Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 286
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 286

Ala Val Gly Leu Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 287
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 287

Ala Val Gly Leu Gly Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 288
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 288

Ala Val Gly Leu Gly Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 289
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 289

Ala Val Gly Leu Gly Ala Val Phe Leu Glu Phe Leu Gly Ala Ala Gly

1	5	yeda 047	pct new.ST25.txt	10	15
<210>	290				
<211>	16				
<212>	PRT				
<213>	Artificial sequence				
<220>					
<223>	Peptides derived of HIV-1 gp41				
<400>	290				
Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly					
1	5			10	15
<210>	291				
<211>	16				
<212>	PRT				
<213>	Artificial sequence				
<220>					
<223>	Peptides derived of HIV-1 gp41				
<400>	291				
Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu Gly Thr Ala Gly					
1	5			10	15
<210>	292				
<211>	16				
<212>	PRT				
<213>	Artificial sequence				
<220>					
<223>	Peptides derived of HIV-1 gp41				
<400>	292				
Ala Val Gly Leu Gly Ala Val Phe Leu Gly Ser Leu Gly Ala Ala Gly					
1	5			10	15
<210>	293				
<211>	16				
<212>	PRT				
<213>	Artificial sequence				
<220>					
<223>	Peptides derived of HIV-1 gp41				
<400>	293				
Ala Val Gly Leu Gly Ala Val Phe Gln Gly Phe Leu Gly Ala Ala Gly					
1	5			10	15
<210>	294				
<211>	16				
<212>	PRT				
<213>	Artificial sequence				
<220>					
<223>	Peptides derived of HIV-1 gp41				
<400>	294				

yeda 047 pct new.ST25.txt

Ala Val Gly Leu Gly Ala Val Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 295
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 295

Ala Val Gly Leu Gly Ala Val Leu Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 296
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 296

Ala Val Gly Leu Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 297
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 297

Ala Val Gly Leu Gly Ala Val Leu Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

<210> 298
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 298

Ala Val Gly Leu Gly Val Ala Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 299
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 299

Ala Val Gly Met Ala Ala Val Phe Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 300

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 300

Ala Val Gly Met Ala Ala Val Phe Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 301

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 301

Ala Val Gly Met Ala Ala Val Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

<210> 302

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 302

Ala Val Gly Met Gly Ala Phe Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 303

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 303

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 304

<211> 16

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 304

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Gly Thr Ala Gly
 1 5 10 15

<210> 305

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 305

Ala Val Gly Met Gly Ala Leu Phe Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

<210> 306

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 306

Ala Val Gly Met Gly Ala Met Phe Leu Gly Phe Leu Ala Ala Ala Gly
 1 5 10 15

<210> 307

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 307

Ala Val Gly Met Gly Ala Met Ile Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

<210> 308

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 308

Ala Val Gly Met Gly Ala Ser Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 309

<211> 16

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 309

Ala Val Gly Met Gly Ala Val Phe Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 310

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 310

Ala Val Gly Met Gly Ala Val Phe Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

<210> 311

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 311

Ala Val Gly Met Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 312

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 312

Ala Val Gly Val Gly Ala Leu Phe Leu Gly Phe Leu Ser Ala Ala Gly
 1 5 10 15

<210> 313

<211> 16

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 313

Ala Val Gly Val Gly Ala Leu Leu Ile Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 314

yeda 047 pct new.ST25.txt

<211> 16
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 314
 Ala Val Gly Val Gly Ala Met Ile Phe Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 315
 <211> 16
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 315
 Ala Val Gly Val Gly Ala Met Ile Leu Gly Phe Leu Gly Ala Ala Gly
 1 5 10 15

<210> 316
 <211> 11
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 316
 Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu
 1 5 10

<210> 317
 <211> 11
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 317
 Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu
 1 5 10

<210> 318
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 318
 Ala Ala Gly Ile Gly Ala Val Leu Leu Gly Phe Leu
 1 5 10

yeda 047 pct new.ST25.txt

<210> 319
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 319

Ala Ala Gly Ile Gly Ala Val Leu Pro Gly Phe Leu
1 5 10

<210> 320
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 320

Ala Ala Gly Leu Gly Ala Val Phe Leu Gly Phe Leu
1 5 10

<210> 321
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 321

Ala Ile Gly Ile Gly Ala Met Phe Leu Gly Phe Leu
1 5 10

<210> 322
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 322

Ala Ile Gly Ile Gly Ala Val Phe Ile Gly Phe Leu
1 5 10

<210> 323
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 323

Ala Ile Gly Ile Gly Ala Val Phe Leu Gly Phe Leu
Page 90

1 5 yeda 047 pct new.ST25.txt
10

<210> 324
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 324

Ala Ile Gly Ile Gly Ala Val Val Leu Gly Phe Leu
1 5 10

<210> 325
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 325

Ala Ile Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu
1 5 10

<210> 326
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 326

Ala Ile Gly Leu Gly Ala Ala Leu Leu Gly Phe Leu
1 5 10

<210> 327
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 327

Ala Ile Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu
1 5 10

<210> 328
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 328

yeda 047 pct new.ST25.txt

Ala Ile Gly Leu Gly Ala Met Phe Leu Gly Phe Leu
 1 5 10

<210> 329
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 329

Ala Ile Gly Leu Gly Ala Val Phe Ile Gly Phe Leu
 1 5 10

<210> 330
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 330

Ala Ile Gly Leu Gly Ala Val Phe Leu Gly Phe Leu
 1 5 10

<210> 331
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<220>
 <221> misc_feature
 <222> (9)..(9)
 <223> Xaa can be any naturally occurring amino acid

<400> 331

Ala Ile Gly Leu Gly Ala Val Phe Xaa Gly Phe Leu
 1 5 10

<210> 332
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 332

Ala Met Gly Ile Gly Ala Val Phe Leu Gly Phe Leu
 1 5 10

<210> 333

yeda 047 pct new.ST25.txt

<211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 333
 Ala Met Gly Ile Gly Ala Val Leu Leu Gly Phe Leu
 1 5 10

<210> 334
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 334
 Ala Val Gly Ile Gly Ala Ala Leu Leu Gly Phe Leu
 1 5 10

<210> 335
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 335
 Ala Val Gly Ile Gly Ala Phe Phe Leu Gly Phe Leu
 1 5 10

<210> 336
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 336
 Ala Val Gly Ile Gly Ala Phe Val Leu Gly Phe Leu
 1 5 10

<210> 337
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 337
 Ala Val Gly Ile Gly Ala Leu Phe Leu Gly Phe Leu
 1 5 10

yeda 047 pct new.ST25.txt

<210> 338
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 338

Ala Val Gly Ile Gly Ala Leu Ile Phe Gly Phe Leu
 1 5 10

<210> 339
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 339

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Phe Leu
 1 5 10

<210> 340
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 340

Ala Val Gly Ile Gly Ala Leu Leu Phe Gly Ser Leu
 1 5 10

<210> 341
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 341

Ala Val Gly Ile Gly Ala Leu Leu Ile Gly Phe Leu
 1 5 10

<210> 342
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 342

Ala Val Gly Ile Gly Ala Leu Leu Leu Gly Phe Leu
 Page 94

1 5 yeda 047_pct new.ST25.txt
10

<210> 343
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 343

Ala Val Gly Ile Gly Ala Leu Leu Val Gly Phe Leu
1 5 10

<210> 344
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 344

Ala Val Gly Ile Gly Ala Leu Ser Leu Gly Phe Leu
1 5 10

<210> 345
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 345

Ala Val Gly Ile Gly Ala Leu Val Phe Gly Phe Leu
1 5 10

<210> 346
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 346

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu
1 5 10

<210> 347
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 347

yeda 047 pct new.ST25.txt

Ala Val Gly Ile Gly Ala Met Phe Leu Gly Ile Leu
1 5 10

<210> 348
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 348

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Leu
1 5 10

<210> 349
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 349

Ala Val Gly Ile Gly Ala Met Ile Phe Gly Phe Ser
1 5 10

<210> 350
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 350

Ala Val Gly Ile Gly Ala Met Ile Leu Gly Phe Leu
1 5 10

<210> 351
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 351

Ala Val Gly Ile Gly Ala Met Leu Phe Gly Phe Leu
1 5 10

<210> 352
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 352

Ala Val Gly Ile Gly Ala Val Phe Phe Gly Phe Leu
1 5 10

<210> 353

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 353

Ala Val Gly Ile Gly Ala Val Phe Ile Gly Phe Leu
1 5 10

<210> 354

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 354

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Phe Leu
1 5 10

<210> 355

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 355

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Ile Leu
1 5 10

<210> 356

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 356

Ala Val Gly Ile Gly Ala Val Phe Leu Gly Ser Leu
1 5 10

<210> 357

<211> 12

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 357

Ala Val Gly Ile Gly Ala Val Phe Leu Arg Phe Leu
 1 5 10

<210> 358

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 358

Ala Val Gly Ile Gly Ala Val Phe Val Gly Phe Leu
 1 5 10

<210> 359

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 359

Ala Val Gly Ile Gly Ala Val Ile Phe Gly Phe Leu
 1 5 10

<210> 360

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 360

Ala Val Gly Ile Gly Ala Val Ile Leu Gly Phe Leu
 1 5 10

<210> 361

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 361

Ala Val Gly Ile Gly Ala Val Leu Phe Gly Phe Leu
 1 5 10

<210> 362

<211> 12

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 362

Ala Val Gly Ile Gly Ala Val Leu Ile Gly Phe Leu
 1 5 10

<210> 363

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 363

Ala Val Gly Ile Gly Ala Val Leu Leu Gly Phe Leu
 1 5 10

<210> 364

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 364

Ala Val Gly Ile Gly Ala Val Leu Val Gly Phe Leu
 1 5 10

<210> 365

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 365

Ala Val Gly Ile Gly Ala Val Ser Leu Gly Phe Leu
 1 5 10

<210> 366

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 366

Ala Val Gly Ile Gly Ile Met Ile Phe Gly Phe Leu
 1 5 10

<210> 367

yeda 047 pct new.ST25.txt

<211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 367
 Ala Val Gly Ile Gly Thr Met Ile Phe Gly Phe Leu
 1 5 10

<210> 368
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 368
 Ala Val Gly Ile Gly Val Leu Leu Leu Gly Phe Leu
 1 5 10

<210> 369
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 369
 Ala Val Gly Ile Gly Val Val Phe Phe Gly Phe Leu
 1 5 10

<210> 370
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 370
 Ala Val Gly Leu Ala Ala Val Phe Phe Gly Phe Leu
 1 5 10

<210> 371
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 371
 Ala Val Gly Leu Glu Ala Val Phe Leu Gly Phe Leu
 1 5 10

yeda 047 pct new.ST25.txt

<210> 372
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 372

Ala Val Gly Leu Gly Ala Ala Phe Leu Gly Phe Leu
 1 5 10

<210> 373
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 373

Ala Val Gly Leu Gly Ala Phe Phe Leu Gly Phe Leu
 1 5 10

<210> 374
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 374

Ala Val Gly Leu Gly Ala Ile Phe Ile Gly Phe Leu
 1 5 10

<210> 375
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 375

Ala Val Gly Leu Gly Ala Leu Phe Phe Gly Phe Leu
 1 5 10

<210> 376
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Peptides derived of HIV-1 gp41

<400> 376

Ala Val Gly Leu Gly Ala Leu Phe Ile Gly Phe Leu
 Page 101

1 5 yeda 047 pct new.ST25.txt
10

<210> 377
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 377

Ala Val Gly Leu Gly Ala Leu Phe Leu Gly Phe Leu
1 5 10

<210> 378
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 378

Ala Val Gly Leu Gly Ala Leu Leu Phe Gly Phe Leu
1 5 10

<210> 379
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 379

Ala Val Gly Leu Gly Ala Met Phe Ile Gly Phe Leu
1 5 10

<210> 380
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 380

Ala Val Gly Leu Gly Ala Met Phe Leu Gly Phe Leu
1 5 10

<210> 381
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 381

yeda 047 pct new.ST25.txt

Ala Val Gly Leu Gly Ala Met Ile Phe Gly Phe Leu
1 5 10

<210> 382
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 382

Ala Val Gly Leu Gly Ala Val Phe Phe Gly Phe Leu
1 5 10

<210> 383
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 383

Ala Val Gly Leu Gly Ala Val Phe Ile Gly Phe Leu
1 5 10

<210> 384
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 384

Ala Val Gly Leu Gly Ala Val Phe Leu Glu Phe Leu
1 5 10

<210> 385
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

<400> 385

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Phe Leu
1 5 10

<210> 386
<211> 12
<212> PRT
<213> Artificial sequence

<220>
<223> Peptides derived of HIV-1 gp41

yeda 047 pct new.ST25.txt

<400> 386

Ala Val Gly Leu Gly Ala Val Phe Leu Gly Ser Leu
1 5 10

<210> 387

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 387

Ala Val Gly Leu Gly Ala Val Phe Gln Gly Phe Leu
1 5 10

<210> 388

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 388

Ala Val Gly Leu Gly Ala Val Ile Phe Gly Phe Leu
1 5 10

<210> 389

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 389

Ala Val Gly Leu Gly Ala Val Leu Phe Gly Phe Leu
1 5 10

<210> 390

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 390

Ala Val Gly Leu Gly Ala Val Leu Leu Gly Phe Leu
1 5 10

<210> 391

<211> 12

<212> PRT

<213> Artificial sequence

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 391

Ala Val Gly Leu Gly Val Ala Phe Leu Gly Phe Leu
 1 5 10

<210> 392

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 392

Ala Val Gly Met Ala Ala Val Phe Phe Gly Phe Leu
 1 5 10

<210> 393

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 393

Ala Val Gly Met Ala Ala Val Phe Ile Gly Phe Leu
 1 5 10

<210> 394

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 394

Ala Val Gly Met Ala Ala Val Phe Leu Gly Phe Leu
 1 5 10

<210> 395

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 395

Ala Val Gly Met Gly Ala Phe Phe Leu Gly Phe Leu
 1 5 10

<210> 396

<211> 12

<212> PRT

yeda 047 pct new.ST25.txt

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 396

Ala	Val	Gly	Met	Gly	Ala	Leu	Phe	Leu	Gly	Phe	Leu
1				5					10		

<210> 397

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 397

Ala	Val	Gly	Met	Gly	Ala	Met	Phe	Leu	Gly	Phe	Leu
1				5					10		

<210> 398

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 398

Ala	Val	Gly	Met	Gly	Ala	Met	Ile	Leu	Gly	Phe	Leu
1				5					10		

<210> 399

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 399

Ala	Val	Gly	Met	Gly	Ala	Ser	Phe	Leu	Gly	Phe	Leu
1				5					10		

<210> 400

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Peptides derived of HIV-1 gp41

<400> 400

Ala	Val	Gly	Met	Gly	Ala	Val	Phe	Leu	Gly	Phe	Leu
1				5					10		

<210> 401

yeda 047 pct new.ST25.txt

<211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 401
 Ala Val Gly Met Gly Ala Val Leu Leu Gly Phe Leu
 1 5 10

<210> 402
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 402
 Ala Val Gly Val Gly Ala Leu Phe Leu Gly Phe Leu
 1 5 10

<210> 403
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 403
 Ala Val Gly Val Gly Ala Leu Leu Ile Gly Phe Leu
 1 5 10

<210> 404
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 404
 Ala Val Gly Val Gly Ala Met Ile Phe Gly Phe Leu
 1 5 10

<210> 405
 <211> 12
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Peptides derived of HIV-1 gp41

 <400> 405
 Ala Val Gly Val Gly Ala Met Ile Leu Gly Phe Leu
 1 5 10

yeda 047 pct new.ST25.txt

<210> 406
<211> 8
<212> PRT
<213> Artificial

<220>
<223> Peptides derived of HIV-1 gp41

<400> 406

Ala Val Gly Ile Gly Ala Leu Phe
1 5

<210> 407
<211> 9
<212> PRT
<213> Artificial

<220>
<223> Peptides derived of HIV-1 gp41

<400> 407

Gly Ala Leu Phe Leu Gly Phe Leu Gly
1 5

<210> 408
<211> 9
<212> PRT
<213> Artificial

<220>
<223> synthetic peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Ala

<400> 408

Gly Ala Leu Phe Leu Gly Phe Leu Gly
1 5

<210> 409
<211> 9
<212> PRT
<213> Artificial

<220>
<223> Peptides derived of HIV-1 gp41

<400> 409

Gly Ala Val Phe Leu Gly Phe Leu Gly
1 5

<210> 410
<211> 9
<212> PRT
<213> Artificial

yeda 047 pct new.ST25.txt

<220>

<223> Peptides derived of HIV-1 gp41

<400> 410

Gly Ala Met Phe Leu Gly Phe Leu Gly
1 5

<210> 411

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Peptides derived of HIV-1 gp41

<400> 411

Gly Ala Val Leu Leu Gly Phe Leu Gly
1 5

<210> 412

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Peptides derived of HIV-1 gp41

<400> 412

Gly Ala Phe Phe Leu Gly Phe Leu Gly
1 5

<210> 413

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Peptides derived of HIV-1 gp41

<400> 413

Gly Ala Met Ile Phe Gly Phe Leu Gly
1 5

<210> 414

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Peptides derived of HIV-1 gp41

<400> 414

Gly Ala Leu Leu Phe Gly Phe Leu Gly
1 5