



(51) International Patent Classification:

C12N 9/22 (2006.01) C12N 15/10 (2006.01)
C12N 15/113 (2010.01) C12N 15/90 (2006.01)

(21) International Application Number:

PCT/US2020/018432

(22) International Filing Date:

14 February 2020 (14.02.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/805,868	14 February 2019 (14.02.2019)	US
62/805,878	14 February 2019 (14.02.2019)	US
62/805,899	14 February 2019 (14.02.2019)	US
62/874,414	15 July 2019 (15.07.2019)	US

(71) Applicant: **METAGENOMI IP TECHNOLOGIES, LLC** [US/US]; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US).

(72) Inventors: **THOMAS, Brian**; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US). **BROWN, Christopher**; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US). **KANTOR, Rose**; 515 Highland Ave., San Mateo, California 94401 (US). **DEVOTO, Au-**

dra; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US). **BUTTERFIELD, Cristina**; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US). **ALEXANDER, Lisa**; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US). **GOLTSMAN, Daniela S. A.**; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US). **LIU, Jason**; 5980 Horton Street, Suite 600, Emeryville, California 94608 (US).

(74) Agent: **ALEMOZAFAR, Ali R.**; Wilson Sonsini, 650 Page Mill Road, Palo Alto, California 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

(54) Title: ENZYMES WITH RUVB DOMAINS




FIG. 1

(57) Abstract: The present disclosure provides for endonuclease enzymes having distinguishing domain features, as well as methods of using such enzymes or variants thereof.

UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

ENZYMES WITH RUVB DOMAINS**CROSS-REFERENCE**

[0001] This application claims the benefit of U.S. Provisional Application No. 62/805,868, filed February 14, 2019 and titled "MG1 ENZYMES WITH RUVB DOMAINS", U.S. Provisional Application No. 62/874,414, filed July 15, 2019 and titled "MG1 ENZYMES WITH RUVB DOMAINS", U.S. Provisional Application No. 62/805,878, filed February 14, 2019 and titled "MG2 ENZYMES CONTAINING RUVB DOMAINS", and U.S. Provisional Application No. 62/805,899, filed February 14, 2019 and titled "MG3 ENZYMES WITH RUVB DOMAINS", each of which is entirely incorporated herein by reference.

BACKGROUND

[0002] Cas enzymes along with their associated Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) guide ribonucleic acids (RNAs) appear to be a pervasive (~45% of bacteria, ~84% of archaea) component of prokaryotic immune systems, serving to protect such microorganisms against non-self nucleic acids, such as infectious viruses and plasmids by CRISPR-RNA guided nucleic acid cleavage. While the deoxyribonucleic acid (DNA) elements encoding CRISPR RNA elements may be relatively conserved in structure and length, their CRISPR-associated (Cas) proteins are highly diverse, containing a wide variety of nucleic acid-interacting domains. While CRISPR DNA elements have been observed as early as 1987, the programmable endonuclease cleavage ability of CRISPR/Cas complexes has only been recognized relatively recently, leading to the use of recombinant CRISPR/Cas systems in diverse DNA manipulation and gene editing applications.

SEQUENCE LISTING

[0003] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on February 13, 2020, is named 55921-703_601_SL.txt and is 23,363,113 bytes in size.

SUMMARY

[0004] In some aspects, the present disclosure provides for an engineered nuclease system, comprising: (a) an endonuclease comprising a RuvC_III domain and an HNH domain, wherein the endonuclease is derived from an uncultivated microorganism, wherein the endonuclease is a class 2, type II Cas endonuclease; and (b) an engineered guide ribonucleic acid structure configured to form a complex with the endonuclease comprising: (i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and (ii) a tracr

ribonucleic acid sequence configured to bind to the endonuclease. In some embodiments, the RuvC_III domain comprises a sequence with at least 70%, at least 75%, at least 80% or at least 90% sequence identity to any one of SEQ ID NOs: 1827-3637.

[0005] In some aspects, the present disclosure provides for an engineered nuclease system comprising: (a) an endonuclease comprising a RuvC_III domain having at least 75% sequence identity to any one of SEQ ID NOs: 1827-3637; and (b) an engineered guide ribonucleic acid structure configured to form a complex with the endonuclease comprising: (i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and (ii) a tracr ribonucleic acid sequence configured to bind to the endonuclease.

[0006] In some aspects, the present disclosure provides for an engineered nuclease system comprising: (a) an endonuclease configured to bind to a protospacer adjacent motif (PAM) sequence comprising SEQ ID NOs: 5512-5537, wherein the endonuclease is a class 2, type II Cas endonuclease; and (b) an engineered guide ribonucleic acid structure configured to form a complex with the endonuclease comprising: (i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and (ii) a tracr ribonucleic acid sequence configured to bind to the endonuclease.

[0007] In some embodiments, the endonuclease is derived from an uncultivated microorganism. In some embodiments, the endonuclease has not been engineered to bind to a different PAM sequence. In some embodiments, the endonuclease is not a Cas9 endonuclease, a Cas14 endonuclease, a Cas12a endonuclease, a Cas12b endonuclease, a Cas 12c endonuclease, a Cas12d endonuclease, a Cas12e endonuclease, a Cas13a endonuclease, a Cas13b endonuclease, a Cas13c endonuclease, or a Cas 13d endonuclease. In some embodiments, the endonuclease has less than 80% identity to a Cas9 endonuclease. In some embodiments, the endonuclease further comprises an HNH domain. In some embodiments, the tracr ribonucleic acid sequence comprises a sequence with at least 80% sequence identity to about 60 to 90 consecutive nucleotides selected from any one of SEQ ID NOs: 5476-5511 and SEQ ID NO: 5538.

[0008] In some aspects, the present disclosure provides for an engineered nuclease system comprising, (a) an engineered guide ribonucleic acid structure comprising: (i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and (ii) a tracr ribonucleic acid sequence configured to bind to an endonuclease, wherein the tracr ribonucleic acid sequence comprises a sequence with at least 80% sequence identity to about 60 to 90 consecutive nucleotides selected from any one of SEQ ID NOs: 5476-5511 and SEQ ID NO: 5538; and (b) a class 2, type II Cas endonuclease configured to bind to the engineered guide ribonucleic acid. In some embodiments, the endonuclease is configured to bind to a protospacer adjacent motif (PAM) sequence selected from the group comprising SEQ ID NOs: 5512-5537.

[0009] In some embodiments, the engineered guide ribonucleic acid structure comprises at least two ribonucleic acid polynucleotides. In some embodiments, the engineered guide ribonucleic acid structure comprises one ribonucleic acid polynucleotide comprising the guide ribonucleic acid sequence and the tracr ribonucleic acid sequence.

[0010] In some embodiments, the guide ribonucleic acid sequence is complementary to a prokaryotic, bacterial, archaeal, eukaryotic, fungal, plant, mammalian, or human genomic sequence. In some embodiments, the guide ribonucleic acid sequence is 15-24 nucleotides in length. In some embodiments, the endonuclease comprises one or more nuclear localization sequences (NLSs) proximal to an N- or C-terminus of the endonuclease. In some embodiments, the NLS comprises a sequence selected from SEQ ID NOs: 5597-5612.

[0011] In some embodiments, the engineered nuclease system further comprises a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least 20 nucleotides 5' to the target deoxyribonucleic acid sequence, a synthetic DNA sequence of at least 10 nucleotides, and a second homology arm comprising a sequence of at least 20 nucleotides 3' to the target sequence. In some embodiments, the first or second homology arm comprises a sequence of at least 40, 80, 120, 150, 200, 300, 500, or 1,000 nucleotides.

[0012] In some embodiments, the system further comprises a source of Mg^{2+} .

[0013] In some embodiments, the endonuclease and the tracr ribonucleic acid sequence are derived from distinct bacterial species within a same phylum. In some embodiments, the endonuclease is derived from a bacterium belonging to a genus *Dermabacter*. In some embodiments, the endonuclease is derived from a bacterium belonging to Phylum Verrucomicrobia, Phylum Candidatus Peregrinibacteria, or Phylum Candidatus Melainabacteria. In some embodiments, the endonuclease is derived from a bacterium comprising a 16S rRNA gene having at least 90% identity to any one of SEQ ID NOs: 5592-5595.

[0014] In some embodiments, the HNH domain comprises a sequence with at least 70% or at least 80% identity to any one of SEQ ID NOs: 5638-5460. In some embodiments, the endonuclease comprises SEQ ID NOs: 1-1826 or a variant thereof having at least 55% identity thereto. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 1827-1830 or SEQ ID NOs: 1827-2140.

[0015] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 3638-3641 or SEQ ID NOs: 3638-3954. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs:

5615-5632. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 1-4 or SEQ ID NOs: 1-319.

[0016] In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5461-5464, SEQ ID NOs: 5476-5479, or SEQ ID NOs: 5476-5489. In some embodiments, the guide RNA structure comprises an RNA sequence predicted to comprise a hairpin consisting of a stem and a loop, wherein the stem comprises at least 10, at least 12 or at least 14 base-paired ribonucleotides, and an asymmetric bulge within 4 base pairs of the loop.

[0017] In some embodiments, the endonuclease is configured to bind to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5512-5515 or SEQ ID NOs: 5527-5530.

[0018] In some embodiments: (a) the endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1827; (b) the guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5461 or SEQ ID NO: 5476; and (c) the endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5512 or SEQ ID NO: 5527. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1828; (b) the guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5462 or SEQ ID NO: 5477; and (c) the endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5513 or SEQ ID NO: 5528. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1829; (b) the guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5463 or SEQ ID NO: 5478; and (c) the endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5514 or SEQ ID NO: 5529. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1830; (b) the guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5464 or SEQ ID NO: 5479; and (c) the endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5515 or SEQ ID NO: 5530.

[0019] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2141-2142 or SEQ ID NOs: 2141-2241. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 3955-3956 or SEQ ID NOs: 3955-4055. In some embodiments, the endonuclease comprises at least 1,

at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5632-5638. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 320-321 or SEQ ID NOs: 320-420. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5465, SEQ ID NOs: 5490-5491 or SEQ ID NOs: 5490-5494. In some embodiments, the guide RNA structure comprises a tracr ribonucleic acid sequence comprising a hairpin comprising at least 8, at least 10, or at least 12 base-paired ribonucleotides. In some embodiments, the endonuclease is configured to bind to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5516 and SEQ ID NOs: 5531. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2141; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5490; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5531. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2142; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5465 or SEQ ID NO: 5491; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5516.

[0020] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2245-2246. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4059-4060. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5639-5648. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 424-425. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5498-5499 and SEQ ID NO: 5539. In some embodiments, the guide RNA structure comprises a guide ribonucleic acid sequence predicted to comprise a hairpin with an uninterrupted base-paired region comprising at least 8 nucleotides of a guide ribonucleic acid sequence and at least 8 nucleotides of a tracr ribonucleic acid sequence, and wherein the tracr ribonucleic acid sequence comprises, from 5' to 3', a first hairpin and a second hairpin, wherein the first hairpin has a longer stem than the second hairpin.

[0021] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2242-2244 or SEQ ID NOs: 2247-2249. In some embodiments, the endonuclease comprises a sequence at least 70%,

80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4056-4058 and SEQ ID NOs 4061-4063. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5639-5648. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 421-423 or SEQ ID NOs: 426-428. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5466-5467, SEQ ID NOs: 5495-5497, SEQ ID NO: 5500-5502, and SEQ ID NO: 5539. In some embodiments, the guide RNA structure comprises a guide ribonucleic acid sequence predicted to comprise a hairpin with an uninterrupted base-paired region comprising at least 8 nucleotides of a guide ribonucleic acid sequence and at least 8 nucleotides of a tracr ribonucleic acid sequence, and wherein the tracr ribonucleic acid sequence comprises, from 5' to 3', a first hairpin and a second hairpin, wherein the first hairpin has a longer stem than the second hairpin. In some embodiments, the endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5517-5518 or SEQ ID NOs: 5532-5534. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2247; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5500; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5517 or SEQ ID NO: 5532. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2248; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5501; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5518 or SEQ ID NOs: 5533. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2249; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5502; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5534.

[0022] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2253 or SEQ ID NOs: 2253-2481. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4067 or SEQ ID NOs: 4067-4295. In some embodiments, the endonuclease comprises a peptide motif according to SEQ ID NO: 5649. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 432 or SEQ ID NOs: 432-660. In some embodiments, the guide RNA structure comprises a

sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5468 or SEQ ID NO: 5503. In some embodiments, the endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5519. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2253; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5468 or SEQ ID NO: 5503; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5519.

[0023] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2482-2489. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4296-4303. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of or SEQ ID NOs: 661-668. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of or SEQ ID NOs: 2490-2498. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4304-4312. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 669-677. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5504.

[0024] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2499 or SEQ ID NOs: 2499-2750. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4313 or SEQ ID NOs: 4313-4564. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5650-5667. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 678 or SEQ ID NOs: 678-929. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5469 or SEQ ID NO: 5505. In some embodiments, the endonuclease is configured to binding to a PAM comprising SEQ ID NOs: 5520 or SEQ ID NOs: 5535. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2499; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5469 or SEQ ID NO: 5505; and (c) the

endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5520 or SEQ ID NO: 5535.

[0025] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2751 or SEQ ID NOs: 2751-2913. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4565 or SEQ ID NOs: 4565-4727. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5668-5678. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 930 or SEQ ID NOs: 930-1092. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5470 or SEQ ID NOs: 5506. In some embodiments, the endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5521 or SEQ ID NOs: 5536. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2751; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5470 or SEQ ID NO: 5506; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5521 or SEQ ID NO: 5536.

[0026] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2914 or SEQ ID NOs: 2914-3174. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4728 or SEQ ID NOs: 4728-4988. In some embodiments, the endonuclease comprises at least 1, at least 2, or at least 3 peptide motifs selected from the group consisting of SEQ ID NOs: 5676-5678. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1093 or SEQ ID NOs: 1093-1353. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5471, SEQ ID NO: 5507, and SEQ ID NOs: 5540-5542. In some embodiments, the guide RNA structure comprises a tracr ribonucleic acid sequence predicted to comprise at least two hairpins comprising less than 5 base-paired ribonucleotides. In some embodiments, the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5522. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2914; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5471 or SEQ

ID NO: 5507; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5522.

[0027] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 3175 or SEQ ID NOs: 3175-3330. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4989 or SEQ ID NOs: 4989-5146. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5679-5686. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1354 or SEQ ID NOs: 1354-1511. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5472 or SEQ ID NOs: 5508. In some embodiments, the endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NO: 5523 or SEQ ID NO: 5537. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3175; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5472 or SEQ ID NO: 5508; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5523 or SEQ ID NO: 5537.

[0028] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 3331 or SEQ ID NOs: 3331-3474. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5147 or SEQ ID NOs: 5147-5290. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5674-5675 and SEQ ID NOs: 5687-5693. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1512 or SEQ ID NOs: 1512-1655. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5473 or SEQ ID NO: 5509. In some embodiments, the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5524. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3331; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5473 or SEQ ID NO: 5509; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5524.

[0029] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 3475 or SEQ ID NOs: 3475-3568. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5291 or SEQ ID NOs: 5291-5389. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5694-5699. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1656 or SEQ ID NOs: 1656-1755. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5474 or SEQ ID NO: 5510. In some embodiments, the endonuclease is configured to binding to a PAM comprising SEQ ID NOs: 5525. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3475; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5474 or SEQ ID NO: 5510; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5525.

[0030] In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 3569 or SEQ ID NOs: 3569-3637. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5390 or SEQ ID NOs: 5390-5460. In some embodiments, the endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5700-5717. In some embodiments, the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1756 or SEQ ID NOs: 1756-1826. In some embodiments, the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5475 or SEQ ID NOs: 5511. In some embodiments, the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5526. In some embodiments: (a) the endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3569; (b) the guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5475 or SEQ ID NO: 5511; and (c) the endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5526. In some embodiments, the sequence identity is determined by a BLASTP, CLUSTALW, MUSCLE, MAFFT, or Smith-Waterman homology search algorithm. In some embodiments, the sequence identity is determined by the BLASTP homology search algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[0031] In some aspects, the present disclosure provides for an engineered guide ribonucleic acid polynucleotide comprising: (a) a DNA-targeting segment comprising a nucleotide sequence that is complementary to a target sequence in a target DNA molecule; and (b) a protein-binding segment comprising two complementary stretches of nucleotides that hybridize to form a double-stranded RNA (dsRNA) duplex, wherein the two complementary stretches of nucleotides are covalently linked to one another with intervening nucleotides, and wherein the engineered guide ribonucleic acid polynucleotide is configured to forming a complex with an endonuclease comprising a RuvC_III domain having at least 75% sequence identity to any one of SEQ ID NOs: 1827-3637 and targeting the complex to the target sequence of the target DNA molecule. In some embodiments, the DNA-targeting segment is positioned 5' of both of the two complementary stretches of nucleotides.

[0032] In some embodiments: (a) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5476-5479 or SEQ ID NOs: 5476-5489; (b) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of (SEQ ID NOs: 5490-5491 or SEQ ID NOs: 5490-5494) and SEQ ID NO: 5538; (c) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5498-5499; (d) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5495-5497 and SEQ ID NOs: 5500-5502; (e) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5503; (f) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5504; (g) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NOs: 5505; (h) protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5506; (i) protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5507; (j) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5508; (k) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5509; (l) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5510; or (m) the protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5511.

[0033] In some embodiments: (a) the guide ribonucleic acid polynucleotide comprises an RNA sequence comprising a hairpin comprising a stem and a loop, wherein the stem comprises at least 10, at least 12, or at least 14 base-paired ribonucleotides, and an asymmetric bulge within 4 base pairs of the loop; (b) the guide ribonucleic acid polynucleotide comprises a tracr ribonucleic acid sequence predicted to comprise a hairpin comprising at least 8, at least 10, or at least 12 base-paired ribonucleotides; (c) the guide ribonucleic acid polynucleotide comprises a guide ribonucleic acid sequence predicted to comprise a hairpin with an uninterrupted base-paired region comprising at least 8 nucleotides of a guide ribonucleic acid sequence and at least 8 nucleotides of a tracr ribonucleic acid sequence, and wherein the tracr ribonucleic acid sequence comprises, from 5' to 3', a first hairpin and a second hairpin, wherein the first hairpin has a longer stem than the second hairpin; or (d) the guide ribonucleic acid polynucleotide comprises a tracr ribonucleic acid sequence predicted to comprise at least two hairpins comprising less than 5 base-paired ribonucleotides.

[0034] In some aspects, the present disclosure provides for a deoxyribonucleic acid polynucleotide encoding any of the engineered guide ribonucleic acid polynucleotides described herein.

[0035] In some aspects, the present disclosure provides for a nucleic acid comprising an engineered nucleic acid sequence optimized for expression in an organism, wherein the nucleic acid encodes a class 2, type II Cas endonuclease comprising a RuvC_III domain and an HNH domain, and wherein the endonuclease is derived from an uncultivated microorganism.

[0036] In some aspects, the present disclosure provides for a nucleic acid comprising an engineered nucleic acid sequence optimized for expression in an organism, wherein the nucleic acid encodes an endonuclease comprising a RuvC_III domain having at least 70% sequence identity to any one of SEQ ID NOs: 1827-3637. In some embodiments, the endonuclease comprises an HNH domain having at least 70% or at least 80% sequence identity to any one of SEQ ID NOs: 3638-5460. In some embodiments, the endonuclease comprises SEQ ID NOs: 5572-5591 or a variant thereof having at least 70% sequence identity thereto. In some embodiments, the endonuclease comprises a sequence encoding one or more nuclear localization sequences (NLSs) proximal to an N- or C-terminus of the endonuclease. In some embodiments, the NLS comprises a sequence selected from SEQ ID NOs: 5597-5612.

[0037] In some embodiments, the organism is prokaryotic, bacterial, eukaryotic, fungal, plant, mammalian, rodent, or human. In some embodiments, the organism is *E. coli*, and: (a) the nucleic acid sequence has at least 70%, 80%, or 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5572-5575; (b) the nucleic acid sequence has at least 70%, 80%, or 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5576-

5577; (c) the nucleic acid sequence has at least 70%, 80%, or 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5578-5580; (d) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5581; (e) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5582; (f) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5583; (g) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5584; (h) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5585; (i) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5586; or (j) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5587. In some embodiments, the organism is human, and: (a) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5588 or SEQ ID NO: 5589; or (b) the nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5590 or SEQ ID NO: 5591.

[0038] In some aspects, the present disclosure provides for a vector comprising a nucleic acid sequence encoding a class 2, type II Cas endonuclease comprising a RuvC_III domain and an HNH domain, wherein the endonuclease is derived from an uncultivated microorganism.

[0039] In some aspects, the present disclosure provides for a vector comprising the any of the nucleic acids described herein. In some embodiments, the vector further comprises a nucleic acid encoding an engineered guide ribonucleic acid structure configured to form a complex with the endonuclease comprising: (a) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and (b) a tracr ribonucleic acid sequence configured to binding to the endonuclease. In some embodiments, the vector is a plasmid, a minicircle, a CELiD, an adeno-associated virus (AAV) derived virion, or a lentivirus.

[0040] In some aspects, the present disclosure provides for a cell comprising any of the vectors described herein.

[0041] In some aspects, the present disclosure provides for a method of manufacturing an endonuclease, comprising cultivating any of the cells described herein.

[0042] In some aspects, the present disclosure provides for a method for binding, cleaving, marking, or modifying a double-stranded deoxyribonucleic acid polynucleotide, comprising: (a) contacting the double-stranded deoxyribonucleic acid polynucleotide with a class 2, type II Cas endonuclease in complex with an engineered guide ribonucleic acid structure configured to bind to the endonuclease and the double-stranded deoxyribonucleic acid polynucleotide; (b) wherein the double-stranded deoxyribonucleic acid polynucleotide comprises a protospacer adjacent motif (PAM); and (c) wherein the PAM comprises a sequence selected from the group consisting of SEQ ID NOs: 5512-5526 or SEQ ID NOs: 5527-5537. In some embodiments, the double-stranded deoxyribonucleic acid polynucleotide comprises a first strand comprising a sequence

complementary to a sequence of the engineered guide ribonucleic acid structure and a second strand comprising the PAM. In some embodiments, the PAM is directly adjacent to the 3' end of the sequence complementary to the sequence of the engineered guide ribonucleic acid structure.

[0043] In some embodiments, the class 2, type II Cas endonuclease is not a Cas9 endonuclease, a Cas14 endonuclease, a Cas12a endonuclease, a Cas12b endonuclease, a Cas 12c endonuclease, a Cas12d endonuclease, a Cas12e endonuclease, a Cas13a endonuclease, a Cas13b endonuclease, a Cas13c endonuclease, or a Cas 13d endonuclease. In some embodiments, the class 2, type II Cas endonuclease is derived from an uncultivated microorganism. In some embodiments, the double-stranded deoxyribonucleic acid polynucleotide is a eukaryotic, plant, fungal, mammalian, rodent, or human double-stranded deoxyribonucleic acid polynucleotide.

[0044] In some embodiments: (a) the PAM comprises a sequence selected from the group consisting of SEQ ID NOs: 5512-5515 and SEQ ID NOs: 5527-5530; (b) the PAM comprises SEQ ID NO: 5516 or SEQ ID NO: 5531; (c) the PAM comprises SEQ ID NO: 5539; (d) the PAM comprises SEQ ID NO: 5517 or SEQ ID NO: 5518; (e) the PAM comprises SEQ ID NO: 5519; (f) the PAM comprises SEQ ID NO: 5520 or SEQ ID NO: 5535; (g) the PAM comprises SEQ ID NO: 5521 or SEQ ID NO: 5536; (h) the PAM comprises SEQ ID NO: 5522; (i) the PAM comprises SEQ ID NO: 5523 or SEQ ID NO: 5537; (j) the PAM comprises SEQ ID NO: 5524; (k) the PAM comprises SEQ ID NO: 5525; or (l) the PAM comprises SEQ ID NO: 5526.

[0045] In some aspects, the present disclosure provides for a method of modifying a target nucleic acid locus, the method comprising delivering to the target nucleic acid locus any of the engineered nuclease systems described herein, wherein the endonuclease is configured to form a complex with the engineered guide ribonucleic acid structure, and wherein the complex is configured such that upon binding of the complex to the target nucleic acid locus, the complex modifies the target nucleic locus. In some embodiments, modifying the target nucleic acid locus comprises binding, nicking, cleaving, or marking the target nucleic acid locus. In some embodiments, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). In some embodiments, the target nucleic acid comprises genomic DNA, viral DNA, viral RNA, or bacterial DNA. In some embodiments, the target nucleic acid locus is *in vitro*. In some embodiments, the target nucleic acid locus is within a cell. In some embodiments, the cell is a prokaryotic cell, a bacterial cell, a eukaryotic cell, a fungal cell, a plant cell, an animal cell, a mammalian cell, a rodent cell, a primate cell, or a human cell.

[0046] In some embodiments, delivering the engineered nuclease system to the target nucleic acid locus comprises delivering the nucleic acid of any of claims 135-140 or the vector of any of claims 142-146. In some embodiments, delivering the engineered nuclease system to the target nucleic acid locus comprises delivering a nucleic acid comprising an open reading frame

encoding the endonuclease. In some embodiments, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. In some embodiments, the engineered nuclease system to the target nucleic acid locus comprises delivering a capped mRNA containing the open reading frame encoding the endonuclease. In some embodiments, the engineered nuclease system to the target nucleic acid locus comprises delivering a translated polypeptide. In some embodiments, the engineered nuclease system to the target nucleic acid locus comprises delivering a deoxyribonucleic acid (DNA) encoding the engineered guide ribonucleic acid structure operably linked to a ribonucleic acid (RNA) pol III promoter. In some embodiments, the endonuclease induces a single-stranded break or a double-stranded break at or proximal to the target locus.

[0047] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0048] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0049] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings (also “Figure” and “FIG.” herein), of which:

[0050] FIGURE 1 depicts typical organizations of CRISPR/Cas loci of different classes and types.

[0051] FIGURE 2 depicts the architecture of a natural Class2/Type II crRNA/tracrRNA pair, compared to a hybrid sgRNA wherein both are joined.

[0052] FIGURE 3 depicts schematics showing organization of CRISPR loci encoding enzymes from the MG1 family.

[0053] FIGURE 4 depicts schematics showing organization of CRISPR loci encoding enzymes from the MG2 family.

[0054] FIGURE 5 depicts schematics showing organization of CRISPR loci encoding enzymes from the MG3 family.

[0055] FIGURE 6 depicts a structure-based alignment of an enzyme of the present disclosure (MG1-1) versus Cas9 from *Staphylococcus aureus* (SEQ ID NO:5613).

[0056] FIGURE 7 depicts a structure-based alignment of an enzyme of the present disclosure (MG2-1) versus Cas9 from *Staphylococcus aureus* (SEQ ID NO:5613).

[0057] FIGURE 8 depicts a structure-based alignment of an enzyme of the present disclosure (MG3-1) versus Cas9 from *Actinomyces naeslundii* (SEQ ID NO: 5614).

[0058] FIGURES 9A, 9B, 9C, 9D, 9E, 9F, 9G, and 9H depicts a structure-based alignment of MG1 family enzymes MG1-1 through MG1-6 (SEQ ID NOs: 5, 6, 9, 1, 2, and 3).

[0059] FIGURE 10 depicts *in vitro* cleavage of DNA by MG1-4 in complex with its corresponding sgRNA containing targeting sequences of varying lengths.

[0060] FIGURE 11 depicts *in cell* cleavage of *E. coli* genomic DNA using MG1-4 along with its corresponding sgRNA. Shown are dilution series of cells transformed with MG1-4 along with target or non-target spacer (top); bottom panel shows the data quantitated, where the left bar represents non-target sgRNA and the right bar represents target sgRNA.

[0061] FIGURE 12 depicts *in cell* indel formation generated by transfection of HEK cells with MG1-4 or MG1-6 constructs described in Example 11 alongside their corresponding sgRNAs containing various different targeting sequences targeting various locations in the human genome.

[0062] FIGURE 13 depicts *in vitro* cleavage of DNA by MG3-6 in complex with its corresponding sgRNA containing targeting sequences of varying lengths.

[0063] FIGURE 14 depicts *in cell* cleavage of *E. coli* genomic DNA using MG3-7 along with its corresponding sgRNA. Shown are dilution series of cells transformed with MG3-7 along with target or non-target spacer (top); bottom panel shows the data quantitated, where the left bar represents non-target sgRNA and the right bar represents target sgRNA.

[0064] FIGURE 15 depicts *in cell* indel formation generated by transfection of HEK cells with MG3-7 constructs described in Example 13 alongside their corresponding sgRNAs containing various different targeting sequences targeting various locations in the human genome.

[0065] FIGURE 16 depicts *in vitro* cleavage of DNA by MG15-1 in complex with its corresponding sgRNA containing targeting sequences of varying lengths.

[0066] FIGURES 17, 18, 19, and 20 depict agarose gels showing the results of PAM vector library cleavage in the presence of TXTL extracts containing various MG family nucleases and

their corresponding tracrRNAs or sgRNAs.

[0067] FIGURES 21, 22, 23, 24, 25 and 26 depict predicted structures (predicted e.g., as in Example 7) of corresponding sgRNAs of MG enzymes described herein.

[0068] FIGURES 27, 28, 29, 30, 31, 32 and 33 depict seqLogo representations of PAM sequences derived via NGS as described herein (e.g. as described in Example 6).

[0069] FIGURE 34 depicts in cell cleavage of *E. coli* genomic DNA using MG2-7 along with its corresponding sgRNA. Shown are dilution series of cells transformed with MG2-7 along with target or non-target spacer (top); bottom panel shows the data quantitated, where the right bar represents non-target sgRNA and the left bar represents target sgRNA.

[0070] FIGURE 35 depicts in cell cleavage of *E. coli* genomic DNA using MG14-1 along with its corresponding sgRNA. Shown are dilution series of cells transformed with MG14-1 along with target or non-target spacer (top); bottom panel shows the data quantitated, where the right bar represents non-target sgRNA and the left bar represents target sgRNA.

[0071] FIGURE 36 depicts in cell cleavage of *E. coli* genomic DNA using MG15-1 along with its corresponding sgRNA. Shown are dilution series of cells transformed with MG15-1 along with target or non-target spacer (top); bottom panel shows the data quantitated, where the right bar represents non-target sgRNA and the left bar represents target sgRNA.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

[0072] The Sequence Listing filed herewith provides exemplary polynucleotide and polypeptide sequences for use in methods, compositions and systems according to the disclosure. Below are exemplary descriptions of sequences therein.

[0073] MG1

[0074] SEQ ID NOs: 1-319 show the full-length peptide sequences of MG1 nucleases.

[0075] SEQ ID NOs: 1827-2140 show the peptide sequences of RuvC_III domains of MG1 nucleases above.

[0076] SEQ ID NOs: 3638-3955 show the peptide of HNH domains of MG1 nucleases above.

[0077] SEQ ID NOs: 5476-5479 show the nucleotide sequences of MG1 tracrRNAs derived from the same loci as MG1 nucleases above (e.g., same loci as SEQ ID NO:1-4, respectively).

[0078] SEQ ID NOs: 5461-5464 show the nucleotide sequences of sgRNAs engineered to function with an MG1 nuclease (e.g., SEQ ID NO:1-4, respectively), where Ns denote nucleotides of a targeting sequence.

[0079] SEQ ID NOs: 5572-5575 show nucleotide sequences for *E. coli* codon-optimized coding sequences for MG1 family enzymes (SEQ ID NOs: 1-4).

[0080] SEQ ID NOs: 5588-5589 show nucleotide sequences for human codon-optimized coding sequences for MG1 family enzymes (SEQ ID NOs: 1 and 3).

[0081] SEQ ID NOs: 5616-5632 show peptide motifs characteristic of MG1 family enzymes.

[0082] MG2

[0083] SEQ ID NOs: 320-420 show the full-length peptide sequences of MG2 nucleases.

[0084] SEQ ID NOs: 2141-2241 show the peptide sequences of RuvC_III domains of MG2 nucleases above.

[0085] SEQ ID NOs: 3955-4055 show the peptide of HNH domains of MG2 nucleases above.

[0086] SEQ ID NOs: 5490-5494 show the nucleotide sequences of MG2 tracrRNAs derived from the same loci as MG2 nucleases above (e.g., same loci as SEQ ID NOs: 320, 321, 323, 325, and 326, respectively).

[0087] SEQ ID NO: 5465 shows the nucleotide sequence of an sgRNA engineered to function with an MG2 nuclease (e.g., SEQ ID NO: 321 above).

[0088] SEQ ID NOs: 5572-5575 show nucleotide sequences for *E. coli* codon-optimized coding sequences for MG2 family enzymes.

[0089] SEQ ID NOs: 5631-5638 show peptide sequences characteristic of MG2 family enzymes.

[0090] MG3

[0091] SEQ ID NOs: 421-431 show the full-length peptide sequences of MG3 nucleases.

[0092] SEQ ID NOs: 2242-2251 show the peptide sequences of RuvC_III domains of MG3 nucleases above.

[0093] SEQ ID NOs: 4056-4066 show the peptide of HNH domains of MG3 nucleases above.

[0094] SEQ ID NOs: 5495-5502 show the nucleotide sequences of MG3 tracrRNAs derived from the same loci as MG3 nucleases above (e.g., same loci as SEQ ID NOs: 421-428, respectively).

[0095] SEQ ID NOs: 5466-5467 show the nucleotide sequence of sgRNAs engineered to function with an MG3 nuclease (e.g., SEQ ID NOs: 421 - 423).

[0096] SEQ ID NOs: 5578-5580 show nucleotide sequences for *E. coli* codon-optimized coding sequences for MG3 family enzymes.

[0097] SEQ ID NOs: 5639-5648 show peptide sequences characteristic of MG3 family enzymes.

[0098] MG4

[0099] SEQ ID NOs: 432-660 show the full-length peptide sequences of MG4 nucleases.

[00100] SEQ ID NOs: 2253-2481 show the peptide sequences of RuvC_III domains of MG4 nucleases above.

[00101] SEQ ID NOs: 4067-4295 show the peptide of HNH domains of MG4 nucleases above.

[00102] SEQ ID NO: 5503 shows the nucleotide sequences of an MG4 tracrRNA derived from the same loci as MG4 nucleases above.

[00103] SEQ ID NO: 5468 shows the nucleotide sequence of sgRNAs engineered to function with an MG4 nuclease.

[00104] SEQ ID NO: 5649 shows a peptide sequence characteristic of MG4 family enzymes.

[00105] MG6

[00106] SEQ ID NOs: 661-668 show the full-length peptide sequences of MG6 nucleases.

[00107] SEQ ID NOs: 2482-2489 show the peptide sequences of RuvC_III domains of MG6 nucleases above.

[00108] SEQ ID NOs: 4296-4303 show the peptide of HNH domains of MG3 nucleases above.

[00109] MG7

[00110] SEQ ID NOs: 669-677 show the full-length peptide sequences of MG7 nucleases.

[00111] SEQ ID NOs: 2490-2498 show the peptide sequences of RuvC_III domains of MG7 nucleases above.

[00112] SEQ ID NOs: 4304-4312 show the peptide of HNH domains of MG3 nucleases above.

[00113] SEQ ID NO: 5504 shows the nucleotide sequence of an MG7 tracrRNA derived from the same loci as MG7 nucleases above.

[00114] MG14

[00115] SEQ ID NOs: 678-929 show the full-length peptide sequences of MG14 nucleases.

[00116] SEQ ID NOs: 2499-2750 show the peptide sequences of RuvC_III domains of MG14 nucleases above.

[00117] SEQ ID NOs: 4313-4564 show the peptide of HNH domains of MG14 nucleases above.

[00118] SEQ ID NO: 5505 shows the nucleotide sequences of MG14 tracrRNA derived from the same loci as MG14 nucleases above.

[00119] SEQ ID NO: 5581 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG14 family enzyme.

[00120] SEQ ID NOs: 5650-5667 show peptide sequences characteristic of MG14 family enzymes.

[00121] MG15

[00122] SEQ ID NOs: 930-1092 show the full-length peptide sequences of MG15 nucleases.

[00123] SEQ ID NOs: 2751-2913 show the peptide sequences of RuvC_III domains of MG15 nucleases above.

[00124] SEQ ID NOs: 4565-4727 show the peptide of HNH domains of MG15 nucleases above.

[00125] SEQ ID NO: 5506 shows the nucleotide sequences of MG15 tracrRNA derived from the same loci as MG15 nucleases above.

[00126] SEQ ID NOs: 5470 shows the nucleotide sequence of an sgRNA engineered to function with an MG15 nuclease.

[00127] SEQ ID NO: 5582 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG15 family enzyme.

[00128] SEQ ID NOs: 5668-5675 show peptide sequences characteristic of MG15 family enzymes.

[00129] MG16

[00130] SEQ ID NOs: 1093-1353 show the full-length peptide sequences of MG16 nucleases.

[00131] SEQ ID NOs: 2914-3174 show the peptide sequences of RuvC_III domains of MG16 nucleases above.

[00132] SEQ ID NOs: 4728-4988 show the peptide of HNH domains of MG16 nucleases above.

[00133] SEQ ID NOs: 5507 show the nucleotide sequences of an MG16 tracrRNA derived from the same loci as MG3 nucleases above.

[00134] SEQ ID NOs: 5471 shows the nucleotide sequence of sgRNAs engineered to function with an MG16 nuclease.

[00135] SEQ ID NO: 5583 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG16 family enzyme.

[00136] SEQ ID NOs: 5676-5678 show peptide sequences characteristic of MG16 family enzymes.

[00137] MG18

[00138] SEQ ID NOs: 1354-1511 show the full-length peptide sequences of MG18 nucleases.

[00139] SEQ ID NOs: 3175-3330 show the peptide sequences of RuvC_III domains of MG18 nucleases above.

[00140] SEQ ID NOs: 4989-5146 show the peptide of HNH domains of MG18 nucleases above.

[00141] SEQ ID NO: 5508 shows the nucleotide sequences of MG18 tracrRNA derived from the same loci as MG18 nucleases above.

[00142] SEQ ID NOs: 5472 shows the nucleotide sequence of an sgRNA engineered to function with an MG18 nuclease.

[00143] SEQ ID NO: 5584 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG18 family enzyme.

[00144] SEQ ID NOs: 5679-5686 show peptide sequences characteristic of MG18 family enzymes.

[00145] MG21

[00146] SEQ ID NOs: 1512-1655 show the full-length peptide sequences of MG21 nucleases.

[00147] SEQ ID NOs: 3331-3474 show the peptide sequences of RuvC_III domains of MG21 nucleases above.

[00148] SEQ ID NOs: 5147-5290 show the peptide of HNH domains of MG21 nucleases above.

[00149] SEQ ID NOs: 5509 show the nucleotide sequence of an MG21 tracrRNA derived from the same loci as MG21 nucleases above.

[00150] SEQ ID NOs: 5473 shows the nucleotide sequence of an sgRNA engineered to function with an MG21 nuclease.

[00151] SEQ ID NO: 5585 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG21 family enzyme.

[00152] SEQ ID NOs: 5687-5692 and 5674-5675 show peptide sequences characteristic of MG21 family enzymes.

[00153] MG22

[00154] SEQ ID NOs: 1656-1755 show the full-length peptide sequences of MG22 nucleases.

[00155] SEQ ID NOs: 3475-3568 show the peptide sequences of RuvC_III domains of MG22 nucleases above.

[00156] SEQ ID NOs: 5291-5389 show the peptide of HNH domains of MG22 nucleases above.

[00157] SEQ ID NO: 5510 show the nucleotide sequence of an MG22 tracrRNA derived from the same loci as MG22 nucleases above.

[00158] SEQ ID NOs: 5474 shows the nucleotide sequence of an sgRNAs engineered to function with an MG22 nuclease.

[00159] SEQ ID NO: 5586 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG22 family enzyme.

[00160] SEQ ID NOs: 5694-5699 show peptide sequences characteristic of MG22 family enzymes.

[00161] MG23

[00162] SEQ ID NOs: 1756-1826 show the full-length peptide sequences of MG23 nucleases.

[00163] SEQ ID NOs: 3569-3637 show the peptide sequences of RuvC_III domains of MG23 nucleases above.

[00164] SEQ ID NOs: 5390-5460 show the peptide of HNH domains of MG23 nucleases above.

[00165] SEQ ID NO: 5511 shows the nucleotide sequences of an MG23 tracrRNA derived from the same loci as MG23 nucleases above.

[00166] SEQ ID NOs: 5475 shows the nucleotide sequence of an sgRNA engineered to function with an MG23 nuclease.

[00167] SEQ ID NO: 5587 shows a nucleotide sequence for an *E. coli* codon-optimized coding sequences for an MG23 family enzyme.

[00168] SEQ ID NOs: 5700-5717 show peptide sequences characteristic of MG23 family enzymes.

DETAILED DESCRIPTION

[00169] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

[00170] The practice of some methods disclosed herein employ, unless otherwise indicated, techniques of immunology, biochemistry, chemistry, molecular biology, microbiology, cell biology, genomics and recombinant DNA. See for example Sambrook and Green, *Molecular Cloning: A Laboratory Manual*, 4th Edition (2012); the series *Current Protocols in Molecular Biology* (F. M. Ausubel, et al. eds.); the series *Methods In Enzymology* (Academic Press, Inc.), *PCR 2: A Practical Approach* (M.J. MacPherson, B.D. Hames and G.R. Taylor eds. (1995)), Harlow and Lane, eds. (1988) *Antibodies, A Laboratory Manual*, and *Culture of Animal Cells: A Manual of Basic Technique and Specialized Applications*, 6th Edition (R.I. Freshney, ed. (2010)) (which is entirely incorporated by reference herein).

[00171] As used herein, the singular forms “a”, “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise. Furthermore, to the extent that the terms “including”, “includes”, “having”, “has”, “with”, or variants thereof are used in either the detailed description and/or the claims, such terms are intended to be inclusive in a manner similar to the term “comprising”.

[00172] The term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. For example, “about” can mean within one or more than one standard deviation, per the practice in the art. Alternatively, “about” can mean a range of up to 20%, up to 15%, up to 10%, up to 5%, or up to 1% of a given value.

[00173] As used herein, a “cell” generally refers to a biological cell. A cell may be the basic structural, functional and/or biological unit of a living organism. A cell may originate from any organism having one or more cells. Some non-limiting examples include: a prokaryotic cell, eukaryotic cell, a bacterial cell, an archaeal cell, a cell of a single-cell eukaryotic organism, a protozoa cell, a cell from a plant (e.g., cells from plant crops, fruits, vegetables, grains, soy bean, corn, maize, wheat, seeds, tomatoes, rice, cassava, sugarcane, pumpkin, hay, potatoes, cotton, cannabis, tobacco, flowering plants, conifers, gymnosperms, ferns, clubmosses, hornworts, liverworts, mosses), an algal cell, (e.g., *Botryococcus braunii*, *Chlamydomonas reinhardtii*, *Nannochloropsis gaditana*, *Chlorella pyrenoidosa*, *Sargassum patens* C. Agardh, and the like),

seaweeds (e.g., kelp), a fungal cell (e.g., a yeast cell, a cell from a mushroom), an animal cell, a cell from an invertebrate animal (e.g., fruit fly, cnidarian, echinoderm, nematode, etc.), a cell from a vertebrate animal (e.g., fish, amphibian, reptile, bird, mammal), a cell from a mammal (e.g., a pig, a cow, a goat, a sheep, a rodent, a rat, a mouse, a non-human primate, a human, etc.), and etcetera. Sometimes a cell is not originating from a natural organism (e.g., a cell can be a synthetically made, sometimes termed an artificial cell).

[00174] The term “nucleotide,” as used herein, generally refers to a base-sugar-phosphate combination. A nucleotide may comprise a synthetic nucleotide. A nucleotide may comprise a synthetic nucleotide analog. Nucleotides may be monomeric units of a nucleic acid sequence (e.g., deoxyribonucleic acid (DNA) and ribonucleic acid (RNA)). The term nucleotide may include ribonucleoside triphosphates adenosine triphosphate (ATP), uridine triphosphate (UTP), cytosine triphosphate (CTP), guanosine triphosphate (GTP) and deoxyribonucleoside triphosphates such as dATP, dCTP, dITP, dUTP, dGTP, dTTP, or derivatives thereof. Such derivatives may include, for example, [α S]dATP, 7-deaza-dGTP and 7-deaza-dATP, and nucleotide derivatives that confer nuclease resistance on the nucleic acid molecule containing them. The term nucleotide as used herein may refer to dideoxyribonucleoside triphosphates (ddNTPs) and their derivatives. Illustrative examples of dideoxyribonucleoside triphosphates may include, but are not limited to, ddATP, ddCTP, ddGTP, ddITP, and ddTTP. A nucleotide may be unlabeled or detectably labeled, such as using moieties comprising optically detectable moieties (e.g., fluorophores). Labeling may also be carried out with quantum dots. Detectable labels may include, for example, radioactive isotopes, fluorescent labels, chemiluminescent labels, bioluminescent labels and enzyme labels. Fluorescent labels of nucleotides may include but are not limited fluorescein, 5-carboxyfluorescein (FAM), 2'7'-dimethoxy-4'5-dichloro-6-carboxyfluorescein (JOE), rhodamine, 6-carboxyrhodamine (R6G), N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX), 4-(4'dimethylaminophenylazo) benzoic acid (DABCYL), Cascade Blue, Oregon Green, Texas Red, Cyanine and 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS). Specific examples of fluorescently labeled nucleotides can include [R6G]dUTP, [TAMRA]dUTP, [R110]dCTP, [R6G]dCTP, [TAMRA]dCTP, [JOE]ddATP, [R6G]ddATP, [FAM]ddCTP, [R110]ddCTP, [TAMRA]ddGTP, [ROX]ddTTP, [dR6G]ddATP, [dR110]ddCTP, [dTAMRA]ddGTP, and [dROX]ddTTP available from Perkin Elmer, Foster City, Calif; FluoroLink DeoxyNucleotides, FluoroLink Cy3-dCTP, FluoroLink Cy5-dCTP, FluoroLink Fluor X-dCTP, FluoroLink Cy3-dUTP, and FluoroLink Cy5-dUTP available from Amersham, Arlington Heights, Ill.; Fluorescein-15-dATP, Fluorescein-12-dUTP, Tetramethyl-rodamine-6-dUTP, IR770-9-dATP, Fluorescein-12-ddUTP, Fluorescein-12-UTP, and Fluorescein-15-2'-dATP available from Boehringer Mannheim, Indianapolis, Ind.; and

Chromosome Labeled Nucleotides, BODIPY-FL-14-UTP, BODIPY-FL-4-UTP, BODIPY-TMR-14-UTP, BODIPY-TMR-14-dUTP, BODIPY-TR-14-UTP, BODIPY-TR-14-dUTP, Cascade Blue-7-UTP, Cascade Blue-7-dUTP, fluorescein-12-UTP, fluorescein-12-dUTP, Oregon Green 488-5-dUTP, Rhodamine Green-5-UTP, Rhodamine Green-5-dUTP, tetramethylrhodamine-6-UTP, tetramethylrhodamine-6-dUTP, Texas Red-5-UTP, Texas Red-5-dUTP, and Texas Red-12-dUTP available from Molecular Probes, Eugene, Oreg. Nucleotides can also be labeled or marked by chemical modification. A chemically-modified single nucleotide can be biotin-dNTP. Some non-limiting examples of biotinylated dNTPs can include, biotin-dATP (e.g., bio-N6-ddATP, biotin-14-dATP), biotin-dCTP (e.g., biotin-11-dCTP, biotin-14-dCTP), and biotin-dUTP (e.g., biotin-11-dUTP, biotin-16-dUTP, biotin-20-dUTP).

[00175] The terms “polynucleotide,” “oligonucleotide,” and “nucleic acid” are used interchangeably to generally refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof, either in single-, double-, or multi-stranded form. A polynucleotide may be exogenous or endogenous to a cell. A polynucleotide may exist in a cell-free environment. A polynucleotide may be a gene or fragment thereof. A polynucleotide may be DNA. A polynucleotide may be RNA. A polynucleotide may have any three-dimensional structure and may perform any function. A polynucleotide may comprise one or more analogs (e.g., altered backbone, sugar, or nucleobase). If present, modifications to the nucleotide structure may be imparted before or after assembly of the polymer. Some non-limiting examples of analogs include: 5-bromouracil, peptide nucleic acid, xeno nucleic acid, morpholinos, locked nucleic acids, glycol nucleic acids, threose nucleic acids, dideoxynucleotides, cordycepin, 7-deaza-GTP, fluorophores (e.g., rhodamine or fluorescein linked to the sugar), thiol containing nucleotides, biotin linked nucleotides, fluorescent base analogs, CpG islands, methyl-7-guanosine, methylated nucleotides, inosine, thiouridine, pseudouridine, dihydrouridine, queuosine, and wyosine. Non-limiting examples of polynucleotides include coding or non-coding regions of a gene or gene fragment, loci (locus) defined from linkage analysis, exons, introns, messenger RNA (mRNA), transfer RNA (tRNA), ribosomal RNA (rRNA), short interfering RNA (siRNA), short-hairpin RNA (shRNA), micro-RNA (miRNA), ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, cell-free polynucleotides including cell-free DNA (cfDNA) and cell-free RNA (cfRNA), nucleic acid probes, and primers. The sequence of nucleotides may be interrupted by non-nucleotide components.

[00176] The terms “transfection” or “transfected” generally refer to introduction of a nucleic acid into a cell by non-viral or viral-based methods. The nucleic acid molecules may be gene

sequences encoding complete proteins or functional portions thereof. See, e.g., Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, 18.1-18.88.

[00177] The terms “peptide,” “polypeptide,” and “protein” are used interchangeably herein to generally refer to a polymer of at least two amino acid residues joined by peptide bond(s). This term does not connote a specific length of polymer, nor is it intended to imply or distinguish whether the peptide is produced using recombinant techniques, chemical or enzymatic synthesis, or is naturally occurring. The terms apply to naturally occurring amino acid polymers as well as amino acid polymers comprising at least one modified amino acid. In some cases, the polymer may be interrupted by non-amino acids. The terms include amino acid chains of any length, including full length proteins, and proteins with or without secondary and/or tertiary structure (e.g., domains). The terms also encompass an amino acid polymer that has been modified, for example, by disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, oxidation, and any other manipulation such as conjugation with a labeling component. The terms “amino acid” and “amino acids,” as used herein, generally refer to natural and non-natural amino acids, including, but not limited to, modified amino acids and amino acid analogues. Modified amino acids may include natural amino acids and non-natural amino acids, which have been chemically modified to include a group or a chemical moiety not naturally present on the amino acid. Amino acid analogues may refer to amino acid derivatives. The term “amino acid” includes both D-amino acids and L-amino acids.

[00178] As used herein, the “non-native” can generally refer to a nucleic acid or polypeptide sequence that is not found in a native nucleic acid or protein. Non-native may refer to affinity tags. Non-native may refer to fusions. Non-native may refer to a naturally occurring nucleic acid or polypeptide sequence that comprises mutations, insertions and/or deletions. A non-native sequence may exhibit and/or encode for an activity (e.g., enzymatic activity, methyltransferase activity, acetyltransferase activity, kinase activity, ubiquitinating activity, etc.) that may also be exhibited by the nucleic acid and/or polypeptide sequence to which the non-native sequence is fused. A non-native nucleic acid or polypeptide sequence may be linked to a naturally-occurring nucleic acid or polypeptide sequence (or a variant thereof) by genetic engineering to generate a chimeric nucleic acid and/or polypeptide sequence encoding a chimeric nucleic acid and/or polypeptide.

[00179] The term “promoter”, as used herein, generally refers to the regulatory DNA region which controls transcription or expression of a gene and which may be located adjacent to or overlapping a nucleotide or region of nucleotides at which RNA transcription is initiated. A promoter may contain specific DNA sequences which bind protein factors, often referred to as transcription factors, which facilitate binding of RNA polymerase to the DNA leading to gene

transcription. A ‘basal promoter’, also referred to as a ‘core promoter’, may generally refer to a promoter that contains all the basic necessary elements to promote transcriptional expression of an operably linked polynucleotide. Eukaryotic basal promoters typically, though not necessarily, contain a TATA-box and/or a CAAT box.

[00180] The term “expression”, as used herein, generally refers to the process by which a nucleic acid sequence or a polynucleotide is transcribed from a DNA template (such as into mRNA or other RNA transcript) and/or the process by which a transcribed mRNA is subsequently translated into peptides, polypeptides, or proteins. Transcripts and encoded polypeptides may be collectively referred to as “gene product.” If the polynucleotide is derived from genomic DNA, expression may include splicing of the mRNA in a eukaryotic cell.

[00181] As used herein, “operably linked”, “operable linkage”, “operatively linked”, or grammatical equivalents thereof generally refer to juxtaposition of genetic elements, e.g., a promoter, an enhancer, a polyadenylation sequence, etc., wherein the elements are in a relationship permitting them to operate in the expected manner. For instance, a regulatory element, which may comprise promoter and/or enhancer sequences, is operatively linked to a coding region if the regulatory element helps initiate transcription of the coding sequence. There may be intervening residues between the regulatory element and coding region so long as this functional relationship is maintained.

[00182] A “vector” as used herein, generally refers to a macromolecule or association of macromolecules that comprises or associates with a polynucleotide and which may be used to mediate delivery of the polynucleotide to a cell. Examples of vectors include plasmids, viral vectors, liposomes, and other gene delivery vehicles. The vector generally comprises genetic elements, e.g., regulatory elements, operatively linked to a gene to facilitate expression of the gene in a target.

[00183] As used herein, “an expression cassette” and “a nucleic acid cassette” are used interchangeably generally to refer to a combination of nucleic acid sequences or elements that are expressed together or are operably linked for expression. In some cases, an expression cassette refers to the combination of regulatory elements and a gene or genes to which they are operably linked for expression.

[00184] A “functional fragment” of a DNA or protein sequence generally refers to a fragment that retains a biological activity (either functional or structural) that is substantially similar to a biological activity of the full-length DNA or protein sequence. A biological activity of a DNA sequence may be its ability to influence expression in a manner known to be attributed to the full-length sequence.

[00185] As used herein, an “engineered” object generally indicates that the object has been

modified by human intervention. According to non-limiting examples: a nucleic acid may be modified by changing its sequence to a sequence that does not occur in nature; a nucleic acid may be modified by ligating it to a nucleic acid that it does not associate with in nature such that the ligated product possesses a function not present in the original nucleic acid; an engineered nucleic acid may be synthesized in vitro with a sequence that does not exist in nature; a protein may be modified by changing its amino acid sequence to a sequence that does not exist in nature; an engineered protein may acquire a new function or property. An “engineered” system comprises at least one engineered component.

[00186] As used herein, “synthetic” and “artificial” are used interchangeably to refer to a protein or a domain thereof that has low sequence identity (e.g., less than 50% sequence identity, less than 25% sequence identity, less than 10% sequence identity, less than 5% sequence identity, less than 1% sequence identity) to a naturally occurring human protein. For example, VPR and VP64 domains are synthetic transactivation domains.

[00187] The term “tracrRNA” or “tracr sequence”, as used herein, can generally refer to a nucleic acid with at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% sequence identity and/or sequence similarity to a wild type exemplary tracrRNA sequence (e.g., a tracrRNA from *S. pyogenes* *S. aureus*, etc or SEQ ID NOs: 5476-5511). tracrRNA can refer to a nucleic acid with at most about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% sequence identity and/or sequence similarity to a wild type exemplary tracrRNA sequence (e.g., a tracrRNA from *S. pyogenes* *S. aureus*, etc). tracrRNA may refer to a modified form of a tracrRNA that can comprise a nucleotide change such as a deletion, insertion, or substitution, variant, mutation, or chimera. A tracrRNA may refer to a nucleic acid that can be at least about 60% identical to a wild type exemplary tracrRNA (e.g., a tracrRNA from *S. pyogenes* *S. aureus*, etc) sequence over a stretch of at least 6 contiguous nucleotides. For example, a tracrRNA sequence can be at least about 60% identical, at least about 65% identical, at least about 70% identical, at least about 75% identical, at least about 80% identical, at least about 85% identical, at least about 90% identical, at least about 95% identical, at least about 98% identical, at least about 99% identical, or 100 % identical to a wild type exemplary tracrRNA (e.g., a tracrRNA from *S. pyogenes* *S. aureus*, etc) sequence over a stretch of at least 6 contiguous nucleotides. Type II tracrRNA sequences can be predicted on a genome sequence by identifying regions with complementarity to part of the repeat sequence in an adjacent CRISPR array.

[00188] As used herein, a “guide nucleic acid” can generally refer to a nucleic acid that may hybridize to another nucleic acid. A guide nucleic acid may be RNA. A guide nucleic acid may be DNA. The guide nucleic acid may be programmed to bind to a sequence of nucleic acid site-specifically. The nucleic acid to be targeted, or the target nucleic acid, may comprise nucleotides.

The guide nucleic acid may comprise nucleotides. A portion of the target nucleic acid may be complementary to a portion of the guide nucleic acid. The strand of a double-stranded target polynucleotide that is complementary to and hybridizes with the guide nucleic acid may be called the complementary strand. The strand of the double-stranded target polynucleotide that is complementary to the complementary strand, and therefore may not be complementary to the guide nucleic acid may be called noncomplementary strand. A guide nucleic acid may comprise a polynucleotide chain and can be called a “single guide nucleic acid.” A guide nucleic acid may comprise two polynucleotide chains and may be called a “double guide nucleic acid.” If not otherwise specified, the term “guide nucleic acid” may be inclusive, referring to both single guide nucleic acids and double guide nucleic acids. A guide nucleic acid may comprise a segment that can be referred to as a “nucleic acid-targeting segment” or a “nucleic acid-targeting sequence.” A nucleic acid-targeting segment may comprise a sub-segment that may be referred to as a “protein binding segment” or “protein binding sequence” or “Cas protein binding segment”.

[00189] The term “sequence identity” or “percent identity” in the context of two or more nucleic acids or polypeptide sequences, generally refers to two (e.g., in a pairwise alignment) or more (e.g., in a multiple sequence alignment) sequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence over a local or global comparison window, as measured using a sequence comparison algorithm. Suitable sequence comparison algorithms for polypeptide sequences include, e.g., BLASTP using parameters of a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment for polypeptide sequences longer than 30 residues; BLASTP using parameters of a wordlength (W) of 2, an expectation (E) of 1000000, and the PAM30 scoring matrix setting gap costs at 9 to open gaps and 1 to extend gaps for sequences of less than 30 residues (these are the default parameters for BLASTP in the BLAST suite available at <https://blast.ncbi.nlm.nih.gov>); CLUSTALW with parameters of ; the Smith-Waterman homology search algorithm with parameters of a match of 2, a mismatch of -1, and a gap of -1; MUSCLE with default parameters; MAFFT with parameters retree of 2 and maxiterations of 1000; Novafold with default parameters; HMMER hmmalign with default parameters.

[00190] As used herein, the term “RuvC_III domain” generally refers to a third discontinuous segment of a RuvC endonuclease domain (the RuvC nuclease domain being comprised of three discontinuous segments, RuvC_I, RuvC_II, and RuvC_III). A RuvC domain or segments thereof can generally be identified by alignment to known domain sequences, structural alignment to proteins with annotated domains, or by comparison to Hidden Markov Models (HMMs) built

based on known domain sequences (e.g., Pfam HMM PF18541 for RuvC_III).

[00191] As used herein, the term “HNH domain” generally refers to an endonuclease domain having characteristic histidine and asparagine residues. An HNH domain can generally be identified by alignment to known domain sequences, structural alignment to proteins with annotated domains, or by comparison to Hidden Markov Models (HMMs) built based on known domain sequences (e.g., Pfam HMM PF01844 for domain HNH).

[00192] *Overview*

[00193] The discovery of new Cas enzymes with unique functionality and structure may offer the potential to further disrupt deoxyribonucleic acid (DNA) editing technologies, improving speed, specificity, functionality, and ease of use. Relative to the predicted prevalence of Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) systems in microbes and the sheer diversity of microbial species, relatively few functionally characterized CRISPR/Cas enzymes exist in the literature. This is partly because a huge number of microbial species may not be readily cultivated in laboratory conditions. Metagenomic sequencing from natural environmental niches that represent large numbers of microbial species may offer the potential to drastically increase the number of new CRISPR/Cas systems known and speed the discovery of new oligonucleotide editing functionalities. A recent example of the fruitfulness of such an approach is demonstrated by the 2016 discovery of CasX/CasY CRISPR systems from metagenomic analysis of natural microbial communities.

[00194] CRISPR/Cas systems are RNA-directed nuclease complexes that have been described to function as an adaptive immune system in microbes. In their natural context, CRISPR/Cas systems occur in CRISPR (clustered regularly interspaced short palindromic repeats) operons or loci, which generally comprise two parts: (i) an array of short repetitive sequences (30-40bp) separated by equally short spacer sequences, which encode the RNA-based targeting element; and (ii) ORFs encoding the Cas encoding the nuclease polypeptide directed by the RNA-based targeting element alongside accessory proteins/enzymes. Efficient nuclease targeting of a particular target nucleic acid sequence generally requires both (i) complementary hybridization between the first 6-8 nucleic acids of the target (the target seed) and the crRNA guide; and (ii) the presence of a protospacer-adjacent motif (PAM) sequence within a defined vicinity of the target seed (the PAM usually being a sequence not commonly represented within the host genome). Depending on the exact function and organization of the system, CRISPR-Cas systems are commonly organized into 2 classes, 5 types and 16 subtypes based on shared functional characteristics and evolutionary similarity.

[00195] Class I CRISPR-Cas systems have large, multisubunit effector complexes, and comprise Types I, III, and IV.

[00196] Type I CRISPR-Cas systems are considered of moderate complexity in terms of components. In Type I CRISPR-Cas systems, the array of RNA-targeting elements is transcribed as a long precursor crRNA (pre-crRNA) that is processed at repeat elements to liberate short, mature crRNAs that direct the nuclease complex to nucleic acid targets when they are followed by a suitable short consensus sequence called a protospacer-adjacent motif (PAM). This processing occurs via an endoribonuclease subunit (Cas6) of a large endonuclease complex called Cascade, which also comprises a nuclease (Cas3) protein component of the crRNA-directed nuclease complex. Cas I nucleases function primarily as DNA nucleases.

[00197] Type III CRISPR systems may be characterized by the presence of a central nuclease, known as Cas10, alongside a repeat-associated mysterious protein (RAMP) that comprises Csm or Cmr protein subunits. Like in Type I systems, the mature crRNA is processed from a pre-crRNA using a Cas6-like enzyme. Unlike type I and II systems, type III systems appear to target and cleave DNA-RNA duplexes (such as DNA strands being used as templates for an RNA polymerase).

[00198] Type IV CRISPR-Cas systems possess an effector complex that consists of a highly reduced large subunit nuclease (csf1), two genes for RAMP proteins of the Cas5 (csf3) and Cas7 (csf2) groups, and, in some cases, a gene for a predicted small subunit; such systems are commonly found on endogenous plasmids.

[00199] Class II CRISPR-Cas systems generally have single-polypeptide multidomain nuclease effectors, and comprise Types II, V and VI.

[00200] Type II CRISPR-Cas systems are considered the simplest in terms of components. In Type II CRISPR-Cas systems, the processing of the CRISPR array into mature crRNAs does not require the presence of a special endonuclease subunit, but rather a small trans-encoded crRNA (tracrRNA) with a region complementary to the array repeat sequence; the tracrRNA interacts with both its corresponding effector nuclease (e.g. Cas9) and the repeat sequence to form a precursor dsRNA structure, which is cleaved by endogenous RNase III to generate a mature effector enzyme loaded with both tracrRNA and crRNA. Cas II nucleases are known as DNA nucleases. Type 2 effectors generally exhibit a structure consisting of a RuvC-like endonuclease domain that adopts the RNase H fold with an unrelated HNH nuclease domain inserted within the folds of the RuvC-like nuclease domain. The RuvC-like domain is responsible for the cleavage of the target (e.g., crRNA complementary) DNA strand, while the HNH domain is responsible for cleavage of the displaced DNA strand.

[00201] Type V CRISPR-Cas systems are characterized by a nuclease effector (e.g. Cas12) structure similar to that of Type II effectors, comprising a RuvC-like domain. Similar to Type II, most (but not all) Type V CRISPR systems use a tracrRNA to process pre-crRNAs into mature

crRNAs; however, unlike Type II systems which requires RNase III to cleave the pre-crRNA into multiple crRNAs, type V systems are capable of using the effector nuclease itself to cleave pre-crRNAs. Like Type-II CRISPR-Cas systems, Type V CRISPR-Cas systems are again known as DNA nucleases. Unlike Type II CRISPR-Cas systems, some Type V enzymes (e.g., Cas12a) appear to have a robust single-stranded nonspecific deoxyribonuclease activity that is activated by the first crRNA directed cleavage of a double-stranded target sequence.

[00202] Type VI CRISPR-Cas systems have RNA-guided RNA endonucleases. Instead of RuvC-like domains, the single polypeptide effector of Type VI systems (e.g. Cas13) comprises two HEPN ribonuclease domains. Differing from both Type II and V systems, Type VI systems also appear to not need a tracrRNA for processing of pre-crRNA into crRNA. Similar to type V systems, however, some Type VI systems (e.g., C2C2) appear to possess robust single-stranded nonspecific nuclease (ribonuclease) activity activated by the first crRNA directed cleavage of a target RNA.

[00203] Because of their simpler architecture, Class II CRISPR-Cas have been most widely adopted for engineering and development as designer nuclease/genome editing applications.

[00204] One of the early adaptations of such a system for in vitro use can be found in Jinek et al. (Science. 2012 Aug 17;337(6096):816-21, which is entirely incorporated herein by reference). The Jinek study first described a system that involved (i) recombinantly-expressed, purified full-length Cas9 (e.g., a Class II, Type II Cas enzyme) isolated from *S. pyogenes* SF370, (ii) purified mature ~42 nt crRNA bearing a ~20 nt 5' sequence complementary to the target DNA sequence desired to be cleaved followed by a 3' tracr-binding sequence (the whole crRNA being in vitro transcribed from a synthetic DNA template carrying a T7 promoter sequence); (iii) purified tracrRNA in vitro transcribed from a synthetic DNA template carrying a T7 promoter sequence, and (iv) Mg²⁺. Jinek later described an improved, engineered system wherein the crRNA of (ii) is joined to the 5' end of (iii) by a linker (e.g., GAAA) to form a single fused synthetic guide RNA (sgRNA) capable of directing Cas9 to a target by itself (compare top and bottom panel of FIGURE 2).

[00205] Mali et al. (Science. 2013 Feb 15; 339(6121): 823–826.), which is entirely incorporated herein by reference, later adapted this system for use in mammalian cells by providing DNA vectors encoding (i) an ORF encoding codon-optimized Cas9 (e.g., a Class II, Type II Cas enzyme) under a suitable mammalian promoter with a C-terminal nuclear localization sequence (e.g., SV40 NLS) and a suitable polyadenylation signal (e.g., TK pA signal); and (ii) an ORF encoding an sgRNA (having a 5' sequence beginning with G followed by 20 nt of a complementary targeting nucleic acid sequence joined to a 3' tracr-binding sequence, a linker, and the tracrRNA sequence) under a suitable Polymerase III promoter (e.g., the U6 promoter) .

[00206] *MGI Enzymes*

[00207] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 1827-2140. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 1827-2140. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 1827-2140. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 1827-1831. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 1827-1831. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 1827-1831. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to SEQ ID NO: 1827. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to SEQ ID NO: 1828. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to SEQ ID NO: 1829. In some cases, the endonuclease may comprise a RuvC_III domain having at least about

70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to SEQ ID NO: 1830. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to SEQ ID NO: 1831.

[00208] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3638-3955. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3638-3955. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3638-3955. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3638-3955. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3638-3955. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3638-3955. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3638-3641. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3638-3641. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3638-3641. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3638. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3638. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3638. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3639. In some

cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3639. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3639. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3640. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3640. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3640. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3641. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3641. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3641.

[00209] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1-6 or 9-319. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1-6 or 9-319. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1-4. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1-4. In some cases, the endonuclease may comprise a peptide motif substantially identical to any one of SEQ ID NOs: 5615, 5616, or 5617.

[00210] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said

endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 1-6 or 9-319, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1-319. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 below, or a combination thereof:

[00211] Table 1: Example NLS Sequences that can be used with Cas Effectors According to the Disclosure

Source	NLS amino acid sequence	SEQ ID NO:
SV40	PKKKRKV	5593
nucleoplasmin bipartite NLS	KRPAATKKAGQAKKKK	5594
c-myc NLS	PAAKRVKLD	5595
c-myc NLS	RQRRNELKRSP	5596
hRNPA1 M9 NLS	NQSSNFGPMKGGNFGGRSSGPYGGGGQYFA KPRNQGGY	5597
Importin-alpha IBB domain	RMRIZFKNKGKDTAELRRRRVEVSVELRKAK KDEQILKRRNV	5598
Myoma T protein	VSRKRPRP	5599
Myoma T protein	PPKKARED	5600
p53	PQPKKKPL	5601
mouse c-abl IV	SALIKKKKKMAP	5602
influenza virus NS1	DRLRR	5603
influenza virus NS1	PKQKKRK	5604
Hepatitis virus delta antigen	RKLKKKIKKL	5605
mouse Mx1 protein	REKKKFLKRR	5606
human poly(ADP-ribose) polymerase	KRKGDEVGDGVDEVAKKKS	5607
steroid hormone receptor (human) glucocorticoid	RKCLQAGMNLEARKTKK	5608

[00212] In some cases, the endonuclease may be recombinant (e.g., cloned, expressed, and purified by a suitable method such as expression in *E. coli* followed by epitope-tag purification). In some cases, the endonuclease may be derived from a bacterium with a 16S rRNA gene having at least about 90% identity to any one of SEQ ID NOs: 5592-5595. The endonuclease may be

derived from a species having a 16S rRNA gene at least about 80%, at least about 82%, at least about 83%, at least about 84%, at least about 85%, at least about 86%, at least about 87%, at least about 88%, at least about 89%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 5592-5595. The endonuclease may be derived from a species having a 16S rRNA gene substantially identical to any one of SEQ ID NOs: 5592-5595. The endonuclease may be derived from a bacterium belonging to the Phylum Verrucomicrobia or the Phylum Candidatus Peregrinibacteria.

[00213] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00214] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00215] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5476-5489. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least

about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5476-5489. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5476-5489. The tracrRNA may comprise any of SEQ ID NOs: 5476-5489.

[00216] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to any one of SEQ ID NOs: 5461-5464. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 5461-5464. The sgRNA may comprise a sequence substantially identical to any one of SEQ ID NOs: 5461-5464.

[00217] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00218] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus, may modify the target nucleic acid locus. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus

may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00219] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 1827-2140. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to any of SEQ ID NOs: 5572-5575 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 5572-5575. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00220] In some cases, the present disclosure may provide for an expression cassette comprising the system disclosed herein, or the nucleic acid described herein. In some cases, the expression cassette or nucleic acid may be supplied as a vector. In some cases, the expression cassette, nucleic acid, or vector may be supplied in a cell. In some cases, the cell is a cell of a bacterium with a 16S rRNA gene having at least about 90% (e.g., at least about 99%) identity to any one of SEQ ID NOs: 5592-5595.

[00221] *MG2 Enzymes*

[00222] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2141-2241. In some cases, the endonuclease may comprise a RuvC_III

domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2141-2241. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2141-2142. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2141-2142. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2141-2142. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2141-2142.

[00223] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3955-4055. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3955-4055. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3955-4055. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 3955-3956. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 3955-3956. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 3955-3956.

[00224] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at

least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 320-420. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 320-420. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 320-321. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 320-321.

[00225] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 320-420, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 320-420. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof.

[00226] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00227] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprise a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA

tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00228] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5490-5494. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5490-5494. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5490-5494. The tracrRNA may comprise any of SEQ ID NOs: 5490-5494.

[00229] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5465. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5465. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5465.

[00230] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00231] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00232] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2141-2241. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to any of SEQ ID NOs: 5576-5577 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 5576-5577. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid

(RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00233] *MG3 Enzymes*

[00234] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2242-2251. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2242-2251. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2242-2251. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2242-2244. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2242-2244. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2242-2244.

[00235] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4056-4066. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4056-4066. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4056-4066. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4056-4058. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about

91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4056-4058. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4056-4058.

[00236] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 421-431. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 421-431. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 421-423. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 421-423.

[00237] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 421-431, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 421-431. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof.

[00238] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00239] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00240] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5495-5502. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5495-5502. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of any one of SEQ ID NOs: 5495-5502. The tracrRNA may comprise any of SEQ ID NOs: 5495-5502.

[00241] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to any one of SEQ ID NOs: 5466-5467. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs:

5466-5467. The sgRNA may comprise a sequence substantially identical to any one of SEQ ID NOs: 5466-5467.

[00242] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00243] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00244] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2242-2251. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to any of SEQ ID NOs: 5578-5580 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about

70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 5578-5580. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00245] *MG4 Enzymes*

[00246] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2253-2481. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2253-2481. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2253-2481. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2253-2481. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2253-2481. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2253-2481.

[00247] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4067-4295. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4067-4295. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4067-4295. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4067-4295. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4067-4295. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4067-4295.

[00248] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 432-660. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 432-660. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 432-660. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 432-660.

[00249] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 432-660, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 432-660. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at

least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00250] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00251] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00252] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5503. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5503. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least

about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5503. The tracrRNA may comprise SEQ ID NO: 5503.

[00253] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5468. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5468. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5468.

[00254] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00255] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00256] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a

RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2253-2481. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00257] *MG6 Enzymes*

[00258] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2482-2489. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2482-2489. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2482-2489.

[00259] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4296-4303. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4296-4303. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4056-4066.

[00260] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at

least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 661-668. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 661-668.

[00261] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 661-668, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 661-668. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00262] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00263] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00264] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence.

[00265] In some cases, the system above may comprise two different guide RNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00266] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00267] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2482-2489. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40,

PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00268] *MG7 Enzymes*

[00269] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2490-2498. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2490-2498. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2490-2498. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2490-2498. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2490-2498. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2490-2498.

[00270] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4304-4312. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99%

identical to any one of SEQ ID NOs: 4304-4312. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4304-4312. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4304-4312. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 4304-4312. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4304-4312.

[00271] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 669-677. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 669-677. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 669-677. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 669-677.

[00272] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 669-677, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 669-677. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00273] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00274] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00275] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5504. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5504. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5504. The tracrRNA may comprise SEQ ID NO: 5504.

[00276] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded

DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00277] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00278] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2490-2498. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00279] *MG14 Enzymes*

[00280] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2499-2750. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2499-2750. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2499-2750. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2499-2750. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2499-2750. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2499-2750.

[00281] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4313-4564. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4313-4564. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4313-4564. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4313-4564. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ

ID NOs: 4067-4295. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4313-4564.

[00282] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 678-929. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 678-929. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 678-929. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 678-929.

[00283] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 678-929, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 678-929. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof.

[00284] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00285] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5'

targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00286] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5505. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5505. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5505. The tracrRNA may comprise SEQ ID NO: 5505.

[00287] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5469. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5469. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5469.

[00288] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at

least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00289] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00290] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2499-2750. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NO: 5581 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5581. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease.

The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00291] *MG15 Enzymes*

[00292] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2751-2913. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2751-2913. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2751-2913. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2751-2913. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2751-2913. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2751-2913.

[00293] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4565-4727. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4565-4727. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4565-4727. The endonuclease may

comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4565-4727. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4565-4727. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4565-4727.

[00294] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 930-1092. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 930-1092. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 930-1092. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 930-1092.

[00295] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 930-1092, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 930-1092. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00296] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of

3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00297] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00298] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5506. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5506. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5506. The tracrRNA may comprise SEQ ID NO: 5506.

[00299] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5470. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least

about 98%, or at least about 99% identity to SEQ ID NO: 5470. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5470.

[00300] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00301] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00302] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2751-2913. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NO: 5582 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least

about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5582. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00303] *MG16 Enzymes*

[00304] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 2914-3174. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2914-3174. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 2914-3174. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 2914-3174. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 2914-3174. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 2914-3174.

[00305] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4728-4988. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4728-4988. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4728-4988. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4728-4988. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4728-4988. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4728-4988.

[00306] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1093-1353. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1093-1353. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1093-1353. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1093-1353.

[00307] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 1093-1353, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1093-1353. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at

least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00308] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00309] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00310] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5507. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5507. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least

about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5507. The tracrRNA may comprise SEQ ID NO: 5507.

[00311] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5471. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5471. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5471.

[00312] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00313] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00314] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a

RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 2914-3174. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NO: 5583 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5583. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00315] *MG18 Enzymes*

[00316] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 3175-3300. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3175-3300. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 3175-3300. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 3175-3300. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at

least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3175-3300. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 3175-3300.

[00317] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4989-5146. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4989-5146. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4989-5146. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 4989-5146. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 4989-5146. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 4989-5146.

[00318] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1354-1511. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1354-1511. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1354-1511. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1354-1511.

[00319] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs:

1354-1511, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1354-1511. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00320] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00321] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprise a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00322] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5508. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least

about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5508. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5508. The tracrRNA may comprise SEQ ID NO: 5508.

[00323] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5472. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5472. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5472.

[00324] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00325] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The

target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00326] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 3175-3300. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NOs: 5584 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NOs: 5584. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00327] *MG21 Enzymes*

[00328] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 3331-3474. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3331-3474. In some cases, the

endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 3331-3474. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 3331-3474. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3331-3474. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 3331-3474.

[00329] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 5147-5290. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 5147-5290. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 5147-5290. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 5147-5290. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 5147-5290. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 5147-5290.

[00330] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1512-1655. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1512-1655. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about

98%, or at least about 99% identity to any one of SEQ ID NOs: 1512-1655. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1512-1655.

[00331] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 1512-1655, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1512-1655. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00332] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00333] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00334] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90)

consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90)

consecutive nucleotides of SEQ ID NO: 5509. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5509. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5509. The tracrRNA may comprise SEQ ID NO: 5509.

[00335] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5473. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5473. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5473.

[00336] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00337] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids

encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00338] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 3331-3474. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NOs: 5585 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NOs: 5585. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00339] *MG22 Enzymes*

[00340] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 3475-3568. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about

30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3475-3568. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 3475-3568. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 3475-3568. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3475-3568. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 3475-3568.

[00341] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 5291-5389. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 5291-5389. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 5291-5389. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 5291-5389. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 5291-5389. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 5291-5389.

[00342] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1656-1755. In some

cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1656-1755. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1656-1755. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1656-1755.

[00343] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 432-660, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1656-1755. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof:

[00344] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00345] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5' targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic

acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00346] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5510. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5510. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5510. The tracrRNA may comprise SEQ ID NO: 5510.

[00347] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5474. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5474. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5474.

[00348] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00349] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one

synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00350] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 3475-3568. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NOs: 5586 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NOs: 5586. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease. The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

[00351] *MG23 Enzymes*

[00352] In one aspect, the present disclosure provides for an engineered nuclease system comprising (a) an endonuclease. In some cases, the endonuclease is a Cas endonuclease. In some cases, the endonuclease is a Type II, Class II Cas endonuclease. The endonuclease may comprise a RuvC_III domain, wherein said RuvC_III domain has at least about 70% sequence identity to any one of SEQ ID NOs: 3569-3637. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the RuvC_III domain has at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3569-3637. In some cases, the endonuclease may comprise a RuvC_III domain, wherein the substantially identical to any one of SEQ ID NOs: 3569-3637. The endonuclease may comprise a RuvC_III domain having at least about 70% sequence identity to any one of SEQ ID NOs: 3569-3637. In some cases, the endonuclease may comprise a RuvC_III domain having at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% identity to any one of SEQ ID NOs: 3569-3637. In some cases, the endonuclease may comprise a RuvC_III domain substantially identical to any one of SEQ ID NOs: 3569-3637.

[00353] The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 5390-5460. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ ID NOs: 5390-5460. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 5390-5460. The endonuclease may comprise an HNH domain having at least about 70% identity to any one of SEQ ID NOs: 5390-5460. In some cases, the endonuclease may comprise an HNH domain having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identical to any one of SEQ

ID NOs: 5390-5460. The endonuclease may comprise an HNH domain substantially identical to any one of SEQ ID NOs: 5390-5460.

[00354] In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1756-1826. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1756-1826. In some cases, the endonuclease may comprise a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1756-1826. In some cases, the endonuclease may be substantially identical to any one of SEQ ID NOs: 1756-1826.

[00355] In some cases, the endonuclease may comprise a variant having one or more nuclear localization sequences (NLSs). The NLS may be proximal to the N- or C-terminus of said endonuclease. The NLS may be appended N-terminal or C-terminal to any one of SEQ ID NOs: 1756-1826, or to a variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to any one of SEQ ID NOs: 1756-1826. The NLS may be an SV40 large T antigen NLS. The NLS may be a c-myc NLS. The NLS can comprise a sequence with at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99% identity to any one of SEQ ID NOs: 5593-5608. The NLS can comprise a sequence substantially identical to any one of SEQ ID NOs: 5593-5608. The NLS can comprise any of the sequences in Table 1 or a combination thereof.

[00356] In some cases, sequence identity may be determined by the BLASTP, CLUSTALW, MUSCLE, MAFFT, Novafold, or Smith-Waterman homology search algorithm. The sequence identity may be determined by the BLASTP algorithm using parameters of a wordlength (W) of 3, an expectation (E) of 10, and using a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

[00357] In some cases, the system above may comprise (b) at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease bearing a 5' targeting region complementary to a desired cleavage sequence. In some cases, the 5'

targeting region may comprises a PAM sequence compatible with the endonuclease. In some cases, the 5' most nucleotide of the targeting region may be G. In some cases, the 5' targeting region may be 15-23 nucleotides in length. The guide sequence and the tracr sequence may be supplied as separate ribonucleic acids (RNAs) or a single ribonucleic acid (RNA). The guide RNA may comprise a crRNA tracrRNA binding sequence 3' to the targeting region. The guide RNA may comprise a tracrRNA sequence preceded by a 4-nucleotide linker 3' to the crRNA tracrRNA binding region. The sgRNA may comprise, from 5' to 3': a non-natural guide nucleic acid sequence capable of hybridizing to a target sequence in a cell; and a tracr sequence. In some cases, the non-natural guide nucleic acid sequence and the tracr sequence are covalently linked.

[00358] In some cases, the tracr sequence may have a particular sequence. The tracr sequence may have at least about 80% to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of a natural tracrRNA sequence. The tracr sequence may have at least about 80% sequence identity to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5511. In some cases, the tracrRNA may have at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to at least about 60-90 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5511. In some cases, the tracrRNA may be substantially identical to at least about 60-100 (e.g., at least about 60, at least about 65, at least about 70, at least about 75, at least about 80, at least about 85, or at least about 90) consecutive nucleotides of SEQ ID NO: 5511. The tracrRNA may comprise SEQ ID NO: 5511.

[00359] In some cases, the at least one engineered synthetic guide ribonucleic acid (sgRNA) capable of forming a complex with the endonuclease may comprise a sequence having at least about 80% identity to SEQ ID NO: 5475. The sgRNA may comprise a sequence having at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 5475. The sgRNA may comprise a sequence substantially identical to SEQ ID NO: 5475.

[00360] In some cases, the system above may comprise two different sgRNAs targeting a first region and a second region for cleavage in a target DNA locus, wherein the second region is 3' to the first region. In some cases, the system above may comprise a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at

least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 5' to the first region, a synthetic DNA sequence of at least about 10 nucleotides, and a second homology arm comprising a sequence of at least about 20 (e.g., at least about 40, 80, 120, 150, 200, 300, 500, or 1kb) nucleotides 3' to the second region.

[00361] In another aspect, the present disclosure provides a method for modifying a target nucleic acid locus of interest. The method may comprise delivering to the target nucleic acid locus any of the non-natural systems disclosed herein, including an enzyme and at least one synthetic guide RNA (sgRNA) disclosed herein. The enzyme may form a complex with the at least one sgRNA, and upon binding of the complex to the target nucleic acid locus of interest, may modify the target nucleic acid locus of interest. Delivering the enzyme to said locus may comprise transfecting a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise electroporating a cell with the system or nucleic acids encoding the system. Delivering the nuclease to said locus may comprise incubating the system in a buffer with a nucleic acid comprising the locus of interest. In some cases, the target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The target nucleic acid locus may comprise genomic DNA, viral DNA, viral RNA, or bacterial DNA. The target nucleic acid locus may be within a cell. The target nucleic acid locus may be in vitro. The target nucleic acid locus may be within a eukaryotic cell or a prokaryotic cell. The cell may be an animal cell, a human cell, bacterial cell, archaeal cell, or a plant cell. The enzyme may induce a single or double-stranded break at or proximal to the target locus of interest.

[00362] In cases where the target nucleic acid locus may be within a cell, the enzyme may be supplied as a nucleic acid containing an open reading frame encoding the enzyme having a RuvC_III domain having at least about 75% (e.g., at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%) identity to any one of SEQ ID NOs: 3569-3637. The deoxyribonucleic acid (DNA) containing an open reading frame encoding said endonuclease may comprise a sequence substantially identical to SEQ ID NOs: 5587 or at variant having at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NOs: 5587. In some cases, the nucleic acid comprises a promoter to which the open reading frame encoding the endonuclease is operably linked. The promoter may be a CMV, EF1a, SV40, PGK1, Ubc, human beta actin, CAG, TRE, or CaMKIIa promoter. The endonuclease may be supplied as a capped mRNA containing said open reading frame encoding said endonuclease.

The endonuclease may be supplied as a translated polypeptide. The at least one engineered sgRNA may be supplied as deoxyribonucleic acid (DNA) containing a gene sequence encoding said at least one engineered sgRNA operably linked to a ribonucleic acid (RNA) pol III promoter. In some cases, the organism may be eukaryotic. In some cases, the organism may be fungal. In some cases, the organism may be human.

EXAMPLES

Example 1. –Metagenomic analysis for new proteins

[00363] Metagenomic samples were collected from sediment, soil and animal. Deoxyribonucleic acid (DNA) was extracted with a Zymobiomics DNA mini-prep kit and sequenced on an Illumina HiSeq® 2500. Samples were collected with consent of property owners. Additional raw sequence data from public sources included animal microbiomes, sediment, soil, hot springs, hydrothermal vents, marine, peat bogs, permafrost, and sewage sequences. Metagenomic sequence data was searched using Hidden Markov Models generated based on known Cas protein sequences including type II Cas effector proteins. Novel effector proteins identified by the search were aligned to known proteins to identify potential active sites. This metagenomic workflow resulted in delineation of the MG1, MG2, MG3, MG4, MG6, MG14, MG15, MG16, MG18, MG21, MG22, and MG23 families of class II, type II CRISPR endonucleases described herein.

Example 2A. – Discovery of an MG1 Family of CRISPR systems

[00364] Analysis of the data from the metagenomic analysis of Example 1 revealed a new cluster of previously undescribed putative CRISPR systems initially comprising six members (MG1-1, MG1-2, MG1-3, MG1-4, MG1-5, and MG1-6 recorded as SEQ ID NOs: 5, 6, 1, 2, and 3 respectively). This family is characterized by an enzyme bearing HNH and RuvC domains. The RuvC domains of this family have a RuvC_III portion having low homology to previously described Cas9 family members. Although the initial family members have a maximum of 56.8% identity among them, all 6 enzymes exhibit a divergent RuvC_III portion of the RuvC domain and bear the common motif of RHHALDAMV (SEQ ID NO:5615), KHHALDAMC (SEQ ID NO:5616), or KHHALDAIC (SEQ ID NO:5617). These motifs are not found in other described Cas9-like enzymes. The corresponding protein and nucleic acid sequences for these new enzymes and their relevant subdomains are presented in the sequence listing. Putative tracrRNA sequences were identified based on their location relative to the other genes and are presented as SEQ ID NOs: 5476-5479. The enzyme systems appear to derive from the Phylum Verrucomicrobia, the Phylum Candidatus Peregrinibacteria, or the Phylum Candidatus Melainabacteria based on the sequences of 16S rRNAs from genome bins containing the

CRISPR systems. The 16S rRNA sequences are presented as SEQ ID NOs: 5592-5596). A detailed domain-level alignment of the CRISPR system sequences together calling out the features described by Shmakov et al. (Mol Cell. 2015 Nov 5;60(3):385-97), which is entirely incorporated by reference) is depicted in FIGURES 9A, 9B, 9C, 9D, 9E, 9F, 9G, and 9H. A comparison of MG1-1, 1-2, and 1-3 versus additional proprietary protein datasets revealed additional protein sequences with similar architecture, presented as SEQ NOs: 7-319. These MG1 protein sequences led to the discovery of additional MG1 motifs as shown in SEQ ID NOs: 5618-5632.

Example 2B. – Discovery of an MG2 Family of CRISPR systems

[00365] Analysis of data from the metagenomic analysis of Example 1 revealed a new cluster of previously undescribed putative CRISPR systems comprising six members (MG2-1, MG2-2, MG2-3, MG2-5, and MG2-6). The corresponding protein and nucleic acid sequences for these new enzymes and exemplary subdomains are presented as SEQ ID NOs: 320, 322-325. Based on their location relative to the other genes, putative tracrRNA sequences were identified in the operon and are presented as SEQ ID NOs: 5490, 5492-5494, and 5538. A detailed domain-level alignment of these sequences versus Cas9 as outlined in Shmakov et al. (Mol Cell. 2015 Nov 5;60(3):385-97.), is depicted in FIGURE 7.

[00366] A comparison of MG2-1, MG2-2, MG2-3, MG2-5, and MG2-6 versus additional proprietary protein datasets revealed additional protein sequences with similar architecture, presented as SEQ NOs: 321 and 326-420. Motifs commonly found in MG2 family members are presented as SEQ ID NOs: 5631-5638.

Example 2C. – Discovery of an MG3 Family of CRISPR systems

[00367] Analysis of the data from the metagenomic analysis of Example 1 revealed a new previously undescribed putative CRISPR system: MG3-1. The corresponding amino acid sequences for this new enzyme and its exemplary subdomains are presented as SEQ ID NOs: 424, 2245, and 4059. Based on proximity to the other elements in the operon, a putative tracrRNA containing sequence was identified and is included as SEQ ID NO: 5498. A detailed domain-level alignment of the sequence versus Cas9 from *Actinomyces naeshundii* is depicted in FIGURE 8.

[00368] A comparison of MG3-1 versus additional proprietary protein datasets revealed additional protein sequences with similar architecture, presented as SEQ NOs: 421-423, 425-431.

Example 2D. – Discovery of MG4, 7, 14, 15, 16, 18, 21, 22, 23 Families of CRISPR systems

[00369] Analysis of the data from the metagenomic analysis of Example 1 revealed new clusters of previously undescribed putative CRISPR systems comprising 9 families of one member each (MG 4-5, MG7-2, MG14-1, MG15-1, MG16-2, MG18-1, MG21-1, MG22-1, MG23-1). The corresponding protein and nucleic acid sequences for these new enzymes and their exemplary subdomains are presented as SEQ ID NOs: 432, 669, 678, 930, 1093, 1354, 1512, 1656, 1756. Based on proximity to the other elements in the operon, a putative tracr containing sequence was identified for each family. These sequences are presented in the sequence listing as SEQ ID NOs: 5503-5511, respectively.

[00370] A comparison of MG 4-5, MG7-2, MG14-1, MG15-1, MG16-2, MG18-1, MG21-1, MG22-1, MG23-1 versus additional proprietary protein datasets revealed additional protein sequences with similar architecture, presented as SEQ NOs: 433-660, 670-677, 679-929, 931-1092, 1094-1353, 1355-1511, 1513-1655, 1657-1755, and 1757-1826. Motifs common to the nucleases of these sets of CRISPR systems are presented as SEQ ID NO: 5649 for MG4; SEQ ID NOs: 5650-5667 for MG14; 5668-5675 for MG15; SEQ ID NOs: 5676-5678 for MG16; SEQ ID NOs: 5679-5686 for MG18; SEQ ID NOs: 5687-5693 and SEQ ID NOs: 5674-5675 for MG21; SEQ ID NOs: 5694-5699 for MG22; and SEQ ID NOs: 5700-5717 for MG23.

Example 3.-Prophetic--Determination of Protospacer-Adjacent Motif.

[00371] Experiments are performed as in any of the examples in Karvelis et al. Methods. 2017 May 15;121-122:3-8, which is entirely incorporated by reference herein, to identify the protospacer adjacent motif (PAM) sequence specificity for the novel enzymes described herein to allow for optimal synthetic sequence targeting.

[00372] In one example (in-vivo screen), cells bearing plasmids encoding any of the enzymes described herein and protospacer-targeting guide RNA are co-transformed with a plasmid library containing an antibiotic resistance gene, and a protospacer sequence flanked by a randomized PAM sequence. Plasmids containing functional PAMs are cleaved by the enzyme, leading to cell death. Deep-sequencing of the enzyme cleavage-resistant plasmid pool isolated from the surviving cells displays a set of depleted plasmids that contain functional cleavage-permitting PAMs.

[00373] In another example (in vitro screen), PAM library in the form of DNA plasmid or concatemeric repeats is subjected to cleavage by the RNP complex (e.g., including the enzyme, tracrRNA and crRNA or the enzyme and hybrid sgRNA) assembled in vitro or in cell lysates. Resulting free DNA ends from successful cleavage events are captured by adapter ligation, followed by the PCR amplification of the PAM-sided products. Amplified library of functional PAMs is subjected to deep sequencing and PAMs licensing DNA cleavage are identified.

Example 4.-Prophetic—Use of synthetic CRISPR system as described herein in a mammalian cell for genome editing

[00374] DNA/RNA sequences encoding (i) an ORF encoding codon-optimized enzyme under a cell-compatible promoter with a cell-compatible C-terminal nuclear localization sequence (e.g., SV40 NLS in the case of human cells) and a suitable polyadenylation signal (e.g., TK pA signal in the case of human cells); and (ii) an ORF encoding an sgRNA (having a 5' sequence beginning with G followed by 20 nt of a complementary targeting nucleic acid sequence targeting genomic DNA followed by a corresponding compatible PAM identified via Example 3 and a 3' tracr-binding sequence, a linker, and the tracrRNA sequence) under a suitable Polymerase III promoter (e.g., the U6 promoter in mammalian cells) are prepared. In some embodiments, these sequences are prepared on the same or separate plasmid vectors, which are transfected via a suitable technique into eukaryotic cells. In some embodiments, these sequences are prepared as separate DNA sequences, which are transfected or microinjected into cells. In some embodiments, these sequences are prepared as synthesized RNAs or in-vitro transcribed RNAs which are transfected or microinjected into cells. In some embodiments, these sequences are translated into proteins and transfected or microinjected into cells.

[00375] Whichever transfection method is selected, (i) and (ii) are introduced into cells. A period of incubation is allowed to pass so that the enzyme and/or sgRNA can be transcribed and/or translated into active form. After the incubation period, genomic DNA in the vicinity of the targeting sequence is analyzed (e.g., by sequencing). An indel is introduced into the genomic DNA in the vicinity of the targeting sequence as a result of enzyme-mediated cleavage and non-homologous end joining.

[00376] In some embodiments, (i) and (ii) are introduced into cells with a third repair nucleotide that encodes regions of the genome flanking the cleavage site of sizes 25 bp or larger, which will facilitate homology directed repair. Containing within these flanking sequences may be a single base pair mutation, a functional gene fragment, a foreign or native gene for expression, or several genes composing a biochemical pathway.

Example 5.-Prophetic—Use of synthetic CRISPR system as described herein in vitro

[00377] Any of the enzymes described herein are cloned into a suitable E. coli expression plasmid containing a purification tag and are recombinantly expressed in E. coli and purified using the recombinant tag. RNAs comprising a 5' G followed by a 20 nt targeting sequence and PAM sequence, a tracrRNA binding region of a compatible crRNA, a GAAA linker, and a compatible tracrRNA are synthesized by suitable solid-phase RNA synthesis methods.

Recombinant enzymes and sgRNA are combined in a suitable cleavage buffer containing Mg²⁺ (e.g., 20 mM HEPES pH 7.5, 100 mM KCl, 5 mM MgCl₂, 1 mM DTT, 5% glycerol) and the reaction is initiated by introducing a target DNA including a sequence complementary to the targeting sequence and PAM sequence. Cleavage of the DNA is monitored by a suitable assay (e.g., agarose gel electrophoresis followed by ethidium bromide staining (or similarly acting DNA-intercalating agent) and UV visualization).

Example 6.-(General protocol) PAM Sequence identification/confirmation for the endonucleases described herein

[00378] PAM sequences were determined by sequencing plasmids containing randomly-generated PAM sequences that could be cleaved by putative endonucleases expressed in an *E. coli* lysate-based expression system (myTXTL, Arbor Biosciences). In this system, an *E. coli* codon optimized nucleotide sequence was transcribed and translated from a PCR fragment under control of a T7 promoter. A second PCR fragment with a tracr sequence under a T7 promoter and a minimal CRISPR array composed of a T7 promoter followed by a repeat-spacer-repeat sequence was transcribed in the same reaction. Successful expression of the endonuclease and tracr sequence in the TXTL system followed by CRISPR array processing provided active *in vitro* CRISPR nuclease complexes.

[00379] A library of target plasmids containing a spacer sequence matching that in the minimal array followed by 8N mixed bases (putative PAM sequences) was incubated with the output of the TXTL reaction. After 1-3 hr, the reaction was stopped and the DNA was recovered via a DNA clean-up kit, e.g., Zymo DCC, AMPure XP beads, QiaQuick etc. Adapter sequences were blunt-end ligated to DNA with active PAM sequences that had been cleaved by the endonuclease, whereas DNA that had not been cleaved was inaccessible for ligation. DNA segments comprising active PAM sequences were then amplified by PCR with primers specific to the library and the adapter sequence. The PCR amplification products were resolved on a gel to identify amplicons that corresponded to cleavage events. The amplified segments of the cleavage reaction were also used as template for preparation of an NGS library. Sequencing this resulting library, which was a subset of the starting 8N library, revealed the sequences which contain the correct PAM for the active CRISPR complex. For PAM testing with a single RNA construct, the same procedure was repeated except that an *in vitro* transcribed RNA was added along with the plasmid library and the tracr/minimal CRISPR array template was omitted. For endonucleases where NGS libraries were prepared, seqLogo (see e.g., Huber et al. Nat Methods. 2015 Feb;12(2):115-21) representations were constructed and are presented in Figures 27, 38, 29, 30, 31, 32, 33, 34, and 35. The seqLogo module used to construct these representations takes the

position weight matrix of a DNA sequence motif (e.g. a PAM sequence) and plots the corresponding sequence logo as introduced by Schneider and Stephens (see e.g. Schneider et al. Nucleic Acids Res. 1990 Oct 25;18(20):6097-100. The characters representing the sequence in the seqLogo representations have been stacked on top of each other for each position in the aligned sequences (e.g. PAM sequences). The height of each letter is proportional to its frequency, and the letters have been sorted so the most common one is on top.

Example 7.-(General protocol) RNA Folding of tracrRNA and sgRNA structures

[00380] Folded structures of guide RNA sequences at 37°C were computed using the method of Andronescu et al. Bioinformatics. 2007 Jul 1;23(13):i19-28, which is incorporated by reference herein in its entirety. Predicted structures of exemplary sgRNAs described herein are presented in Figures 21, 22, 23, 24, 25, and 26.

Example 8.-(General protocol) In vitro cleavage efficiency of MG CRISPR Complexes

[00381] Endonucleases were expressed as His-tagged fusion proteins from an inducible T7 promoter in a protease deficient *E. coli* B strain. Cells expressing the His-tagged proteins were lysed by sonication and the His-tagged proteins were purified by Ni-NTA affinity chromatography on a HisTrap FF column (GE Lifescience) on an AKTA Avant FPLC (GE Lifescience). The eluate was resolved by SDS-PAGE on acrylamide gels (Bio-Rad) and stained with InstantBlue Ultrafast coomassie (Sigma-Aldrich). Purity was determined using densitometry of the protein band with ImageLab software (Bio-Rad). Purified endonucleases were dialyzed into a storage buffer composed of 50 mM Tris-HCl, 300 mM NaCl, 1 mM TCEP, 5% glycerol; pH 7.5 and stored at -80°C.

[00382] Target DNAs containing spacer sequences and PAM sequences (determined e.g., as in Example 6) were constructed by DNA synthesis. A single representative PAM was chosen for testing when the PAM had degenerate bases. The target DNAs comprised 2200 bp of linear DNA derived from a plasmid via PCR amplification with a PAM and spacer located 700 bp from one end. Successful cleavage resulted in fragments of 700 and 1500 bp. The target DNA, *in vitro* transcribed single RNA, and purified recombinant protein were combined in cleavage buffer (10 mM Tris, 100 mM NaCl, 10 mM MgCl₂) with an excess of protein and RNA and incubated for 5 minutes to 3 hours, usually 1 hr. The reaction was stopped via addition of RNase A and incubation at 60 minutes. The reaction was then resolved on a 1.2% TAE agarose gel and the fraction of cleaved target DNA is quantified in ImageLab software.

Example 9.-(General protocol) Testing of Genome Cleavage Activity of MG CRISPR Complexes in *E. coli*

[00383] *E. coli* lacks the capacity to efficiently repair double-stranded DNA breaks. Thus, cleavage of genomic DNA can be a lethal event. Exploiting this phenomenon, endonuclease activity was tested in *E. coli* by recombinantly expressing an endonuclease and a tracrRNA in a target strain with spacer/target and PAM sequences integrated into its genomic DNA.

[00384] In this assay, the PAM sequence is specific for the endonuclease being tested as determined by the methods described in Example 6. sgRNA sequences were determined based upon the sequence and predicted structure of the tracrRNA. Repeat-anti-repeat pairings of 8-12 bp (generally 10bp) were chosen, starting from the 5' end of the repeat. The remaining 3' end of the repeat and 5' end of the tracrRNA were replaced with a tetraloop. Generally, the tetraloop was GAAA, but other tetraloops can be used, particularly if the GAAA sequence is predicted to interfere with folding. In these cases, a TTCG tetraloop was used.

[00385] Engineered strains with PAM sequences integrated into their genomic DNA were transformed with DNA encoding the endonuclease. Transformants were then made chemocompetent and transformed with 50 ng of single guide RNAs either specific to the target sequence ("on target"), or non-specific to the target ("non target"). After heat shock, transformations were recovered in SOC for 2 hrs at 37°C. Nuclease efficiency was then determined by a 5-fold dilution series grown on induction media. Colonies were quantified from the dilution series in triplicate.

Example 10.-(General protocol) Testing of Genome Cleavage Activity of MG CRISPR Complexes in Mammalian Cells

[00386] To show targeting and cleavage activity in mammalian cells, the MG Cas effector protein sequences were tested in two mammalian expression vectors: (a) one with a C-terminal SV40 NLS and a 2A-GFP tag, and (b) one with no GFP tag and two SV40 NLS sequences, one on the N-terminus and one on the C-terminus. In some instances, nucleotide sequences encoding the endonucleases were codon-optimized for expression in mammalian cells.

[00387] The corresponding single guide RNA sequence (sgRNA) with targeting sequence attached is cloned into a second mammalian expression vector. The two plasmids are cotransfected into HEK293T cells. 72 hr after co-transfection of the expression plasmid and a sgRNA targeting plasmid into HEK293T cells, the DNA is extracted and used for the preparation of an NGS-library. Percent NHEJ is measured via indels in the sequencing of the target site to demonstrate the targeting efficiency of the enzyme in mammalian cells. At least 10 different target sites were chosen to test each protein's activity.

Example 11.- Characterization of MG1 family members**[00388] PAM Specificity, tracrRNA/sgRNA Validation**

[00389] The targeted endonuclease activity of MG1 family endonuclease systems was confirmed using the myTXTL system described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG1-4 (dual guide: see gel 1, lane 3, single guide: see gel 6 lane 2), MG1-5 (gel 2 lane 10), MG1-6 (dual guide: see gel 5 lane 6, single guide see: gel 6 lane 5), and MG1-7 (dual guide: see gel 3 lane 13, single guide: see gel 3 lane 2) (protein SEQ ID NOs: 1-4, respectively). Sequencing the PCR products revealed active PAM sequences for these enzymes as shown in Table 2.

[00390] Table 2: PAM sequence specificities and related data for MG1 enzymes

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG1-4	1	nRRRAA	5527	5476	5461	nRRR	5512
MG1-5	2	nnnnCC	5528	5477	5462	nnnnYY	5513
MG1-6	3	nnRRWC	5529	5478	5463	nnRRAY	5514
MG1-7	4	nRRRAA	5530	5479	5464	nRRRAAG	5515

[00391] Synthetic single guide RNAs (sgRNAs) were designed based on the sequences and predicted structures of the tracrRNAs and are presented as SEQ ID NOs: 5461-5464. The PAM sequence screen of Example 6 was repeated with the sgRNAs. The results of this experiment are also presented in Table 2, which reveals that PAM specificity changed slightly when using sgRNAs.

[00392] Targeted endonuclease activity In Vitro

[00393] *In vitro* activity of the MG1-4 endonuclease system (protein SEQ ID NO: 1 with sgRNA SEQ ID NO: 5461) on a target DNA with a PAM sequence CAGGAAGG was verified using the method of Example 8. The single guide sequence reported above (SEQ ID NO: 5461) was used, with varying spacer/targeting sequence lengths from 18-24 nt replacing the Ns of the sequence. The results are shown in Figure 10, wherein the left panel shows a gel demonstrating DNA cleavage by MG1-4 in combination with corresponding single guide sgRNAs having different targeting sequence lengths (18-24nt), and the right panel shows the same data quantified as a bar

graph. The data demonstrated that targeting sequences from 18-24 nucleotides were functional with the MG1-4/sgRNA system.

[00394] *Targeted endonuclease activity in bacterial cells*

[00395] *In vivo* activity of the MG1-4 endonuclease system (protein SEQ ID NO: 1, sgRNA SEQ ID NO: 5461) was tested with the PAM sequence CAGGAAGG as in Example 9. Transformed *E. coli* were plated in serial dilution, and the results (showing *E. coli* serial dilutions in the left panel and quantitated growth in the right panel) are presented in Figure 11. A substantial reduction in the growth of *E. coli* expressing on target sgRNA compared to *E. coli* expressing non-target sgRNA indicates that genomic DNA was specifically cleaved by the endonuclease in *E. coli* cells.

[00396] *Targeted endonuclease activity in mammalian cells*

[00397] The method of Example 10 was used to demonstrate targeting and cleavage activity in mammalian cells. Open reading frames encoding the MG1-4 (protein SEQ ID NO: 5527) and MG1-6 (protein SEQ ID NO: 5529) sequences were cloned into 2 mammalian expression vectors, one with a C-terminal SV40 NLS and a 2A-GFP tag (*E. coli* MG-BB) and one with no GFP tag and 2 NLS sequences, one on the N-terminus and one on the C-terminus (*E. coli* pMG5-BB). For MG1-6, the open reading frame was additionally codon-optimized for mammalian expression (SEQ ID NO: 5589) and cloned into the 2-NLS plasmid backbone (MG-16hs). The results of this experiment are shown in Figure 12. The endonuclease expression vectors were cotransfected into HEK293T cells with a second vector for expressing a sgRNA (e.g., SEQ ID NOs: 5512 or 5515) with a tracr sequence specific for the endonuclease and a guide sequence selected from Tables 3-4. 72 hr after co-transfection the DNA was extracted and used for the preparation of an NGS-library. Cleavage activity was detected by the appearance of internal deletions (NHEJ remnants) proximal to the sequence of the target site. Percent NHEJ was measured via indels in the sequencing of the target site to demonstrate the targeting efficiency of the enzyme in mammalian cells and is presented in Figure 12.

[00398] **Table 3: MG1-4 mammalian targeting sequences**

MG1-4 Target ID	MG1-4 Targeting Sequence	Targeting sequence SEQ ID NO:	Targeted Gene
1	aatatgtagctgtttgggaggt	5543	VEGFA
2	ctagggggcgctcggccaccac	5544	VEGFA
3	tggctaaagagggaatgggctt	5545	VEGFA
4	cacaccccggtctggctaaag	5546	VEGFA
5	tcggaggagccgtgtccgcgc	5547	VEGFA

6	gcggaaccacggctcctccgaag	5548	VEGFA
7	gtacaaacggcagaagctggag	5549	EMX1
8	gaggaagggcctgagtcgagca	5550	EMX1
9	aaggcaaacatcctgataatgg	5551	Apolipoprotein

[00399] Table 4: MG1-6 mammalian targeting sequences

MG1-6 Target ID	MG1-6 Targeting Sequence	Targeting sequence SEQ ID NO:	Targeted Gene
1	tcttttagccagagccggggtgt	5552	VEGFA
2	tggaccccctatttctgacctc	5553	VEGFA
3	atgggagcccttcttcttctgc	5554	EMX1
4	tgccacgaagcaggccaatggg	5555	EMX1
5	tggtgtctgttgagggtgcta	5556	HBB-R01
6	gggcaggttggtatcaaggta	5557	HBB-R01
7	aggtgctgacgtagtagtgct	5558	FANCF
8	gccctactccgctttcacctt	5559	FANCF
9	aatgtatgctggctttaagg	5560	IVS40
10	gctcctttggctaggggaagtgt	5561	IVS40

Example 12.- Characterization of MG2 family members**[00400] PAM Specificity, tracrRNA/sgRNA Validation**

[00401] The targeted endonuclease activity of MG2 family members was confirmed in the myTXTL system as described in Example 6. Results of this assay are shown in Figures 17-20. In the assay shown in Figures 17-20, active proteins that successfully cleave the library result in a band around 170 bp in the gel. Amplification products were observed for MG2-1 (see gel 2 lane 11 and gel 4 lane 6) and MG2-7 (see gel 11 lane 10) (SEQ ID NOs: 320 and 321, respectively). Sequencing the PCR products revealed active PAM sequences in Table 5 below:

[00402] Table 5: PAM sequence specificities and related data for MG2 enzymes

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG2-1	320	nRCGT A	5531	5490	N/A	N/A	N/A
MG2-7	321	N/A	N/A	5491	5465	NNNRTA	5516

[00403] Targeted endonuclease activity in bacterial cells

[00404] *In vivo* activity of the MG2-7 endonuclease system with a sgRNA (endonuclease SEQ ID NO: 321; sgRNA SEQ ID NO: 5465) and an AGCGTAAG PAM sequence was confirmed using the method described in Example 9. Transformed *E. coli* were plated in serial dilution, and the results (showing *E. coli* serial dilutions in the left panel and quantitated growth in the right panel) are presented in Figure 34. A substantial reduction in the growth of *E. coli* expressing on target sgRNA compared to *E. coli* expressing non-target sgRNA indicates that genomic DNA was specifically cleaved by the MG1-4 endonuclease in *E. coli* cells.

[00405] Example 13.- Characterization of MG3 family members

[00406] *PAM Specificity, tracrRNA/sgRNA Validation*

[00407] The targeted endonuclease activity of MG3 family members was confirmed using the myTXTL system as described in Example 6 using tracr sequences and CRISPR arrays. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG3-6 (dual guide: see gel 2 lane 8; single guide: see gel 3 lane 3), MG3-7 (dual guide: see gel 2 lane 3, single guide: see gel 3 lane 4), and MG3-8 (dual guide: see gel 9 lane 5) (SEQ ID NOs: 421, 422, and 423, respectively). Sequencing the PCR products revealed active PAM sequences in Table 6 below:

[00408] Table 6: PAM sequence specificities and related data for MG3 enzymes

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG3-6	421	nnRGGTT	5532	5500	5466	nnGGG	5517
MG3-7	422	nnRnYAY	5533	5501	5467	nnGnTnT	5518
MG3-8	423	nnRGGTT	5534	5502	N/A	N/A	N/A

[00409] Synthetic single guide RNAs (sgRNAs) were designed based on the sequences and predicted structures of the tracrRNAs and are presented as SEQ ID NOs: 5466-5467. The PAM sequence screen of Example 6 was repeated with the sgRNAs. The results of this experiment are also presented in Table 6, which reveals that PAM specificity changed slightly when using sgRNAs.

[00410] *Targeted endonuclease activity In Vitro*

[00411] *In vitro* activity of the MG3-6 (endonuclease SEQ ID NO: 421) was verified with the PAM sequence GTGGGTTA using the method of Example 8. The single guide sequence

reported above (SEQ ID NO: 5466) was used, with varying spacer/targeting sequence lengths from 18-24 nt replacing the Ns of the sequence. The results are shown in Figure 13, wherein the top panel shows a gel demonstrating DNA cleavage by MG3-6 in combination with different sgRNAs having different targeting sequence lengths (18-24nt), and the bottom panel shows the same data quantified as a bar graph. The data demonstrated that targeting sequences from 18-24 nucleotides were functional with the MG3-6/sgRNA system.

[00412] *Targeted endonuclease activity in bacterial cells*

[00413] *In vivo* activity of the MG3-7 endonuclease system (protein SEQ ID NO: 422; sgRNA SEQ ID NO: 5467) was tested with the PAM sequence TGGACCTG using the method of Example 9. Transformed *E. coli* were plated in serial dilution, and the results (showing *E. coli* serial dilutions in the top panel and quantitated growth in the bottom panel) are presented in Figure 14. A substantial reduction in the growth of *E. coli* expressing on target sgRNA compared to *E. coli* expressing non-target sgRNA indicates that genomic DNA was being specifically cleaved by the MG3-7 endonuclease system.

[00414] *Targeted endonuclease activity in mammalian cells*

[00415] The method of Example 10 was used to demonstrate targeting and cleavage activity in mammalian cells. Open reading frames encoding MG3-7 (protein SEQ ID NO: 422) was cloned into 2 mammalian expression vectors, one with a C-terminal SV40 NLS and a 2A-GFP tag (*E. coli* MG-BB) and one with no GFP tag and 2 NLS sequences, one on the N-terminus and one on the C-terminus (*E. coli* pMG5-BB). The endonuclease expression vectors were cotransfected into HEK293T cells with a second vector for expressing the sgRNA above with a guide sequence selected from Table 7. The results of this experiment are shown in Figure 12. 72 hr after co-transfection DNA was extracted and used for the preparation of an NGS-library. Cleavage activity was detected by the appearance of internal deletions (NHEJ remnants) in the vicinity of the target site. Results are presented in Figure 15.

[00416] The target site which were encoded on the sgRNA plasmids are shown in Table 7 below.

[00417] **Table 7: MG3-7 mammalian targeting sequences**

MG3-7 Target ID	MG3-7 Targeting Sequence	Targeting sequence SEQ ID NO:	Targeted Gene
1	cccctatttctgacctccaaa	5563	VEGFA
2	tgtggtccagaaccggaggac	5564	EMX1
3	ggccctgggcaggttggtatca	5565	HBB-R01
4	tccttaaacctgtcttgaacc	5566	HBB-R01
5	ctgactcctgaggagaagtctg	5567	HBB-R01
6	tccgagcttctggcggtctcaa	5568	FANCF
7	tatcatttcgcggatgttcaa	5569	FANCF
8	tcgggcagagggtgcatcacct	5570	Apolipoprotein

9	ataataagcagaacttttagtg	5571	Fibrinogen
10	gttttcttagttattaatttc	5572	Fibrinogen

Example 13.- Characterization of MG4 family members

[00418] PAM Specificity, tracrRNA/sgRNA Validation

[00419] The targeted endonuclease activity of MG4 family endonuclease systems was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for, MG4-2 (dual guide: see gel2 lane 9, single guide: see gel 10 lane 7) (SEQ ID NO: 432). Sequencing the PCR products revealed active PAM sequences shown in Table 8 below.

Table 8: PAM sequence specificities and related data for MG4 enzymes

Enzyme	Enzyme protein SEQ ID NO	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG4-5	432	5503	5468	nnCCR	5519

Example 14.- Characterization of MG14 family members

[00420] PAM Specificity, tracrRNA/sgRNA Validation

[00421] The targeted endonuclease activity of MG14 family members (was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG14-1 (dual guide: see gel 1 lane 4, single guide: see gel 3 lane 8) (SEQ ID NO: 678). Sequencing the PCR products revealed active PAM sequence specificities shown in Table 9 below.

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM determined	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM determined	Synthetic (single guide) PAM SEQ ID NO:
MG14-1	678	NNNNGGT A	5535	5505	5469	NNNGG RTA	5520

[00422] Targeted endonuclease activity in bacterial cells

[00423] *In vivo* activity of the MG14-1 endonuclease system with a sgRNA (endonuclease SEQ ID NO: 678; sgRNA SEQ ID NO: 5469) and a GGCGGGGA PAM sequence was confirmed using the method described in Example 9. Transformed *E. coli* were plated in serial dilution, and the results (showing *E. coli* serial dilutions in the left panel and quantitated growth in the right panel) are presented in Figure 35. A substantial reduction in the growth of *E. coli* expressing on target sgRNA compared to *E. coli* expressing non-target sgRNA indicates that genomic DNA was specifically cleaved by the MG1-4 endonuclease in *E. coli* cells.

Example 15.- Characterization of MG15 family members

[00424] *PAM Specificity, tracrRNA/sgRNA Validation*

[00425] The targeted endonuclease activity of MG15 family members was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG15-1 (dual guide: see gel 7 lane 7, single guide: see gel 3 lane 9) (SEQ ID NO: 930). Sequencing the PCR products revealed active PAM sequence specificities detailed in Table 10 below.

[00426] Table 10:

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG15-1	930	nnnnC	5536	5506	5470	nnnnC	5521

[00427] *In Vitro Activity*

[00428] *In vitro* activity of the MG15-1 endonuclease system (protein SEQ ID NO: 930; sgRNA SEQ ID NO: 5470) was tested with the PAM sequence GGGTCAA using the method of Example 8. The single guide sequence reported above (SEQ ID NO: 5470) was used, with varying spacer/targeting sequence lengths from 18-24 nt (replacing the Ns of the sequence). The results are shown in Figure 16, wherein the top panel shows a gel demonstrating DNA cleavage by MG15-1 in combination with different sgRNAs having different targeting sequence lengths (18-24nt), and the bottom panel shows the same data quantified as a bar graph. The data demonstrated that targeting sequences from 18-24 nucleotides were functional with the MG15-1/sgRNA system.

[00429] *Targeted endonuclease activity in bacterial cells*

[00430] *In vivo* activity of the MG15-1 endonuclease system with a sgRNA (endonuclease SEQ ID NO: 930; sgRNA SEQ ID NO: 5470) and a GGGTCAAA PAM sequence was confirmed using the method described in Example 9. Transformed *E. coli* were plated in serial dilution, and the results (showing *E. coli* serial dilutions in the left panel and quantitated growth in the right panel) are presented in Figure 35. A substantial reduction in the growth of *E. coli* expressing on target sgRNA compared to *E. coli* expressing non-target sgRNA indicates that genomic DNA was specifically cleaved by the MG1-4 endonuclease in *E. coli* cells.

Example 16.- Characterization of MG16 family members

[00431] *PAM Specificity, tracrRNA/sgRNA Validation*

[00432] The targeted endonuclease activity of MG16 family members was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG16-2 (see gel 11, lane 17) (SEQ ID NO: 1093). Sequencing the PCR products revealed active PAM sequence specificities detailed in Table 11 below.

[00433] Table 11:

Enzyme	Enzyme protein SEQ ID NO	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG16-2	1093	5471	nRTnCC	5522

Example 17.- Characterization of MG18 family members

[00434] *PAM Specificity, tracrRNA/sgRNA Validation*

[00435] The targeted endonuclease activity of MG18 family members was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG18-1 (dual guide: see gel 9 lane 9, single guide: see gel 11 lane 12) (SEQ ID NO: 1354). Sequencing the PCR products revealed active PAM sequence specificities detailed in Table 12 below.

[00436] Table 12:

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG18-1	1354	nRWART	5537	5508	5472	nnnRRT	5523

Example 18.- Characterization of MG21 family members

[00437] PAM Specificity, tracrRNA/sgRNA Validation

[00438] The targeted endonuclease activity of MG21 family was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG21-1 (see gel 11 lane 2) (SEQ ID NO: 1512). Sequencing the PCR products revealed active PAM sequence specificities detailed in Table 13 below.

[00439] Table 13:

Enzyme	Enzyme protein SEQ ID NO	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG21-1	1512	5473	nnRnR	5524

Example 19.- Characterization of MG22 family members

[00440] PAM Specificity, tracrRNA/sgRNA Validation

[00441] The targeted endonuclease activity of MG22 family members was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. In the assay shown Figures 17-20, active proteins that successfully cleave the library result in a band around 170 bp in the gel. Amplification products were observed for MG22-1 (see gel 11 lane 3) (protein SEQ ID NO: 1656). Sequencing the PCR products revealed active PAM sequence specificities detailed in Table 14 below.

[00442] Table 14:

Enzyme	Enzyme protein SEQ ID NO	Native (dual guide) PAM determined	PAM SEQ ID NO:	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM determined	Synthetic (single guide) PAM SEQ ID NO:
MG22-1	1656	N/A	N/A	5510	5474	nnRCnT	5525

Example 20.- Characterization of MG23 family members

[00443] PAM Specificity, tracrRNA/sgRNA Validation

[00444] The targeted endonuclease activity of MG23 family members was confirmed using the myTXTL system as described in Example 6. In this assay, PCR amplification of cleaved target plasmids yields a product that migrates at approximately 170 bp in the gel, as shown in Figures 17-20. Amplification products were observed for MG23-1 (see gel 11 lane 4) (SEQ ID NO: 1756). Sequencing the PCR products revealed active PAM sequences specificities for these enzymes detailed Table 15 below.

[00445] Table 15:

Enzyme	Enzyme protein SEQ ID NO	tracrRNA SEQ ID NO:	sgRNA SEQ ID NO:	Synthetic (single guide) PAM	Synthetic (single guide) PAM SEQ ID NO:
MG23-1	1756	5511	5475	nRRA	5526

[00446] Systems of the present disclosure may be used for various applications, such as, for example, nucleic acid editing (e.g., gene editing), binding to a nucleic acid molecule (e.g., sequence-specific binding). Such systems may be used, for example, for addressing (e.g., removing or replacing) a genetically inherited mutation that may cause a disease in a subject, inactivating a gene in order to ascertain its function in a cell, as a diagnostic tool to detect disease-causing genetic elements (e.g. via cleavage of reverse-transcribed viral RNA or an amplified DNA sequence encoding a disease-causing mutation), as deactivated enzymes in combination with a probe to target and detect a specific nucleotide sequence (e.g. sequence encoding antibiotic resistance in bacteria), to render viruses inactive or incapable of infecting host cells by targeting viral genomes, to add genes or amend metabolic pathways to engineer organisms to produce valuable small molecules, macromolecules, or secondary metabolites, to establish a gene drive element for evolutionary selection, to detect cell perturbations by foreign small molecules and nucleotides as a biosensor.

[00447] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

WHAT IS CLAIMED IS:

1. An engineered nuclease system, comprising:
 - (a) an endonuclease comprising a RuvC_III domain and an HNH domain, wherein said endonuclease is derived from an uncultivated microorganism, wherein said endonuclease is a class 2, type II Cas endonuclease; and
 - (b) an engineered guide ribonucleic acid structure configured to form a complex with said endonuclease comprising:
 - (i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and
 - (ii) a tracr ribonucleic acid sequence configured to bind to said endonuclease.
2. The engineered nuclease system of claim 1, wherein said RuvC_III domain comprises a sequence with at least 70%, at least 75%, at least 80% or at least 90% sequence identity to any one of SEQ ID NOs: 1827-3637.
3. An engineered nuclease system comprising:
 - (a) an endonuclease comprising a RuvC_III domain having at least 75% sequence identity to any one of SEQ ID NOs: 1827-3637; and
 - (b) an engineered guide ribonucleic acid structure configured to form a complex with said endonuclease comprising:
 - (i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and
 - (ii) a tracr ribonucleic acid sequence configured to bind to said endonuclease.
4. An engineered nuclease system comprising:
 - (a) an endonuclease configured to bind to a protospacer adjacent motif (PAM) sequence comprising SEQ ID NOs: 5512-5537, wherein said endonuclease is a class 2, type II Cas endonuclease; and

(b) an engineered guide ribonucleic acid structure configured to form a complex with said endonuclease comprising:

(i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and

(ii) a tracr ribonucleic acid sequence configured to bind to said endonuclease.

5. The engineered nuclease system of claim 4, wherein said endonuclease is derived from an uncultivated microorganism.

6. The engineered nuclease system of any one of claims 4-5, wherein said endonuclease has not been engineered to bind to a different PAM sequence.

7. The engineered nuclease system of claim 4, wherein said endonuclease is not a Cas9 endonuclease, a Cas14 endonuclease, a Cas12a endonuclease, a Cas12b endonuclease, a Cas 12c endonuclease, a Cas12d endonuclease, a Cas12e endonuclease, a Cas13a endonuclease, a Cas13b endonuclease, a Cas13c endonuclease, or a Cas 13d endonuclease.

8. The engineered nuclease system of claim 4, wherein said endonuclease has less than 80% identity to a Cas9 endonuclease.

9. The engineered nuclease system of any one of claims 3-8, wherein said endonuclease further comprises an HNH domain.

10. The engineered nuclease system of any one of claims 1-9, wherein said tracr ribonucleic acid sequence comprises a sequence with at least 80% sequence identity to about 60 to 90 consecutive nucleotides selected from any one of SEQ ID NOs: 5476-5511 and SEQ ID NO: 5538.

11. An engineered nuclease system comprising,

(a) an engineered guide ribonucleic acid structure comprising:

(i) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and

- (ii) a tracr ribonucleic acid sequence configured to bind to an endonuclease,

wherein said tracr ribonucleic acid sequence comprises a sequence with at least 80% sequence identity to about 60 to 90 consecutive nucleotides selected from any one of SEQ ID NOs: 5476-5511 and SEQ ID NO: 5538; and

- (b) a class 2, type II Cas endonuclease configured to bind to said engineered guide ribonucleic acid.

12. The engineered nuclease system of any of claims 1-3 or 11, wherein said endonuclease is configured to bind to a protospacer adjacent motif (PAM) sequence selected from the group comprising SEQ ID NOs: 5512-5537.

13. The engineered nuclease system of any one of claims 1-11, wherein said engineered guide ribonucleic acid structure comprises at least two ribonucleic acid polynucleotides.

14. The engineered nuclease system of any one of claims 1-11, wherein said engineered guide ribonucleic acid structure comprises one ribonucleic acid polynucleotide comprising said guide ribonucleic acid sequence and said tracr ribonucleic acid sequence.

15. The engineered nuclease system of any one of claims 1-14, wherein said guide ribonucleic acid sequence is complementary to a prokaryotic, bacterial, archaeal, eukaryotic, fungal, plant, mammalian, or human genomic sequence.

16. The engineered nuclease system of any one of claims 1-15, wherein said guide ribonucleic acid sequence is 15-24 nucleotides in length.

17. The engineered nuclease system of any one of claims 1-16, wherein said endonuclease comprises one or more nuclear localization sequences (NLSs) proximal to an N- or C-terminus of said endonuclease.

18. The engineered nuclease system of any one of claims 1-17, wherein said NLS comprises a sequence selected from SEQ ID NOs: 5597-5612.

19. The engineered nuclease system of any one of claims 1-18, further comprising

a single- or double-stranded DNA repair template comprising from 5' to 3': a first homology arm comprising a sequence of at least 20 nucleotides 5' to said target deoxyribonucleic acid sequence, a synthetic DNA sequence of at least 10 nucleotides, and a second homology arm comprising a sequence of at least 20 nucleotides 3' to said target sequence.

20. The engineered nuclease system of claim 19, wherein said first or second homology arm comprises a sequence of at least 40, 80, 120, 150, 200, 300, 500, or 1,000 nucleotides.

21. The engineered nuclease system of any one of claims 1-20, wherein said system further comprises a source of Mg^{2+} .

22. The engineered nuclease system of any one of claims 1-21, wherein said endonuclease and said target ribonucleic acid sequence are derived from distinct bacterial species within a same phylum.

23. The engineered nuclease system of any one of claims 1-22, wherein said endonuclease is derived from a bacterium belonging to a genus *Dermabacter*.

24. The engineered nuclease system of any one of claims 1-22, wherein said endonuclease is derived from a bacterium belonging to Phylum Verrucomicrobia, Phylum Candidatus Peregrinibacteria, or Phylum Candidatus Melainabacteria.

25. The engineered nuclease system of any one of claims 1-22, wherein said endonuclease is derived from a bacterium comprising a 16S rRNA gene having at least 90% identity to any one of SEQ ID NOs: 5592-5595.

26. The engineered nuclease system of any one of claims 1-25, wherein said HNH domain comprises a sequence with at least 70% or at least 80% identity to any one of SEQ ID NOs: 5638-5460.

27. The engineered nuclease system of any one of claims 1-26, wherein said endonuclease comprises SEQ ID NOs: 1-1826 or a variant thereof having at least 55% identity thereto.

28. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 1827-1830 or SEQ ID NOs: 1827-2140.

29. The engineered nuclease system of any one of claims 1-28, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 3638-3641 or SEQ ID NOs: 3638-3954.

30. The engineered nuclease system of any one of claims 1-29, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5615-5632.

31. The engineered nuclease system of any one of claims 1-30, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 1-4 or SEQ ID NOs: 1-319.

32. The engineered nuclease system of any one of claims 1-31, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5461-5464, SEQ ID NOs: 5476-5479, or SEQ ID NOs: 5476-5489.

33. The engineered nuclease system of any one of claims 1-32, wherein said guide RNA structure comprises an RNA sequence predicted to comprise a hairpin consisting of a stem and a loop, wherein the stem comprises at least 10, at least 12 or at least 14 base-paired ribonucleotides, and an asymmetric bulge within 4 base pairs of the loop.

34. The engineered nuclease system of any one of claims 1-33, wherein said endonuclease is configured to bind to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5512-5515 or SEQ ID NOs: 5527-5530.

35. The engineered nuclease system of any one of claims 1-34, wherein:

- a) said endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1827;
- b) said guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5461 or SEQ ID NO: 5476; and

- c) said endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5512 or SEQ ID NO: 5527.

36. The engineered nuclease system of any one of claims 1-34, wherein:

- a) said endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1828;
- b) said guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5462 or SEQ ID NO: 5477; and
- c) said endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5513 or SEQ ID NO: 5528.

37. The engineered nuclease system of any one of claims 1-34, wherein:

- a) said endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1829;
- b) said guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5463 or SEQ ID NO: 5478; and
- c) said endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5514 or SEQ ID NO: 5529.

38. The engineered nuclease system of any one of claims 1-34, wherein:

- a) said endonuclease comprises a sequence at least 70%, at least 80%, or at least 90% identical to SEQ ID NO: 1830;
- b) said guide RNA structure comprises a sequence at least 70%, at least 80%, or at least 90% identical to at least one of SEQ ID NO: 5464 or SEQ ID NO: 5479; and
- c) said endonuclease is configured to bind to a PAM comprising SEQ ID NO: 5515 or SEQ ID NO: 5530.

39. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2141-2142 or SEQ ID NOs: 2141-2241.

40. The engineered nuclease system of any one of claims 1-27 or claim 39, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 3955-3956 or SEQ ID NOs: 3955-4055.

41. The engineered nuclease system of any one of claims 1-27 or claims 39-40, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5632-5638.

42. The engineered nuclease system of any one of claims 1-27 or claims 39-41, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 320-321 or SEQ ID NOs: 320-420.

43. The engineered nuclease system of any one of claims 1-27 or claims 39-42, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5465, SEQ ID NOs: 5490-5491 or SEQ ID NOs: 5490-5494.

44. The engineered nuclease system of any one of claims 1-27 or claims 39-43, wherein said guide RNA structure comprises a tracr ribonucleic acid sequence comprising a hairpin comprising at least 8, at least 10, or at least 12 base-paired ribonucleotides.

45. The engineered nuclease system of any one of claims 1-27 or claims 39-44, wherein said endonuclease is configured to bind to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5516 and SEQ ID NOs: 5531.

46. The engineered nuclease system of any one of claims 1-27 or claim 39-45, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2141;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5490; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5531.

47. The engineered nuclease system of any one of claims 1-27 or claim 39-45, wherein

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2142;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5465 or SEQ ID NO: 5491; and

- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5516.

48. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2245-2246.

49. The engineered nuclease system of any one of claims 1-27 or claim 48, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4059-4060.

50. The engineered nuclease system of any one of claims 1-27 or claims 48-49, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5639-5648.

51. The engineered nuclease system of any one of claims 1-27 or claims 48-50, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 424-425.

52. The engineered nuclease system of any one of claims 1-27 or claims 48-51, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5498-5499 and SEQ ID NO: 5539.

53. The engineered nuclease system of any one of claims 1-27 or claims 48-52, wherein said guide RNA structure comprises a guide ribonucleic acid sequence predicted to comprise a hairpin with an uninterrupted base-paired region comprising at least 8 nucleotides of a guide ribonucleic acid sequence and at least 8 nucleotides of a tracr ribonucleic acid sequence, and wherein said tracr ribonucleic acid sequence comprises, from 5' to 3', a first hairpin and a second hairpin, wherein said first hairpin has a longer stem than said second hairpin.

54. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2242-2244 or SEQ ID NOs: 2247-2249.

55. The engineered nuclease system of any one of claims 1-27 or claim 54, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4056-4058 and SEQ ID NOs 4061-4063.

56. The engineered nuclease system of any one of claims 1-27 or claims 54-55, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5639-5648.

57. The engineered nuclease system of any one of claims 1-27 or claims 54-56, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 421-423 or SEQ ID NOs: 426-428.

58. The engineered nuclease system of any one of claims 1-27 or claims 54-57, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5466-5467, SEQ ID NOs: 5495-5497, SEQ ID NO: 5500-5502, and SEQ ID NO: 5539.

59. The engineered nuclease system of any one of claims 1-27 or claims 54-58, wherein said guide RNA structure comprises a guide ribonucleic acid sequence predicted to comprise a hairpin with an uninterrupted base-paired region comprising at least 8 nucleotides of a guide ribonucleic acid sequence and at least 8 nucleotides of a tracr ribonucleic acid sequence, and wherein said tracr ribonucleic acid sequence comprises, from 5' to 3', a first hairpin and a second hairpin, wherein said first hairpin has a longer stem than said second hairpin.

60. The engineered nuclease system of any one of claims 1-27 or claims 54-59, wherein said endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5517-5518 or SEQ ID NOs: 5532-5534.

61. The engineered nuclease system of any one of claims 1-27 or claims 54-60, wherein:
- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2247;
 - b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5500; and
 - c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5517 or SEQ ID NO: 5532.

62. The engineered nuclease system of any one of claims 1-27 or claims 54-60, wherein:
- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2248;
 - b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5501; and
 - c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5518 or SEQ ID NOs: 5533.
63. The engineered nuclease system of any one of claims 1-27 or claims 54-60, wherein:
- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2249;
 - b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5502; and
 - c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5534.
64. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2253 or SEQ ID NOs: 2253-2481.
65. The engineered nuclease system of any one of claims 1-27 or claim 64, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4067 or SEQ ID NOs: 4067-4295.
66. The engineered nuclease system of any one of claims 1-27 or claims 64-65, wherein said endonuclease comprises a peptide motif according to SEQ ID NO: 5649.
67. The engineered nuclease system of any one of claims 1-27 or claims 64-66, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 432 or SEQ ID NOs: 432-660.
68. The engineered nuclease system of any one of claims 1-27 or claims 64-67, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5468 or SEQ ID NO: 5503.

69. The engineered nuclease system of any one of claims 1-27 or claims 64-68, wherein said endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5519.

70. The engineered nuclease system of any one of claims 1-27 or claims 64-69, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2253;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5468 or SEQ ID NO: 5503; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5519.

71. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 2482-2489.

72. The engineered nuclease system of any one of claims 1-27 or claim 71, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4296-4303.

73. The engineered nuclease system of any one of claims 1-27 or claims 71-72, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of or SEQ ID NOs: 661-668.

74. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of or SEQ ID NOs: 2490-2498.

75. The engineered nuclease system of any one of claims 1-27 or claim 74, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 4304-4312.

76. The engineered nuclease system of any one of claims 1-27 or claims 74-75, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 669-677.

77. The engineered nuclease system of any one of claims 1-27 or claims 74-76, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5504.

78. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2499 or SEQ ID NOs: 2499-2750.

79. The engineered nuclease system of any one of claims 1-27 or claim 78, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4313 or SEQ ID NOs: 4313-4564.

80. The engineered nuclease system of any one of claims 1-27 or claims 78-79, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5650-5667.

81. The engineered nuclease system of any one of claims 1-27 or claims 78-80, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 678 or SEQ ID NOs: 678-929

82. The engineered nuclease system of any one of claims 1-27 or claims 78-81, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5469 or SEQ ID NO: 5505.

83. The engineered nuclease system of any one of claims 1-27 or claims 78-82, wherein said endonuclease is configured to binding to a PAM comprising SEQ ID NOs: 5520 or SEQ ID NOs: 5535.

84. The engineered nuclease system of any one of claims 1-27 or claims 78-83, wherein:
a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2499;

- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5469 or SEQ ID NO: 5505; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5520 or SEQ ID NO: 5535.

85. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2751 or SEQ ID NOs: 2751-2913.

86. The engineered nuclease system of any one of claims 1-27 or claim 85, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4565 or SEQ ID NOs: 4565-4727.

87. The engineered nuclease system of any one of claims 1-27 or claims 85-86, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5668-5678.

88. The engineered nuclease system of any one of claims 1-27 or claims 85-87, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 930 or SEQ ID NOs: 930-1092.

89. The engineered nuclease system of any one of claims 1-27 or claims 85-88, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5470 or SEQ ID NOs: 5506.

90. The engineered nuclease system of any one of claims 1-27 or claims 85-89, wherein said endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NOs: 5521 or SEQ ID NOs: 5536.

91. The engineered nuclease system of any one of claims 1-27 or claims 85-90, wherein:
- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2751;
 - b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5470 or SEQ ID NO: 5506; and

- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5521 or SEQ ID NO: 5536.

92. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 2914 or SEQ ID NOs: 2914-3174.

93. The engineered nuclease system of any one of claims 1-27 or claim 92, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4728 or SEQ ID NOs: 4728-4988.

94. The engineered nuclease system of any one of claims 1-27 or claims 92-93, wherein said endonuclease comprises at least 1, at least 2, or at least 3 peptide motifs selected from the group consisting of SEQ ID NOs: 5676-5678.

95. The engineered nuclease system of any one of claims 1-27 or claims 92-94, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1093 or SEQ ID NOs: 1093-1353.

96. The engineered nuclease system of any one of claims 1-27 or claims 92-95, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5471, SEQ ID NO: 5507, and SEQ ID NOs: 5540-5542.

97. The engineered nuclease system of any one of claims 1-27 or claims 92-96, wherein said guide RNA structure comprises a tracr ribonucleic acid sequence predicted to comprise at least two hairpins comprising less than 5 base-paired ribonucleotides.

98. The engineered nuclease system of any one of claims 1-27 or claims 92-97, wherein said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5522.

99. The engineered nuclease system of any one of claims 1-27 or claims 92-98, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 2914;

- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5471 or SEQ ID NO: 5507; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5522.

100. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 3175 or SEQ ID NOs: 3175-3330.

101. The engineered nuclease system of any one of claims 1-27 or claim 100, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 4989 or SEQ ID NOs: 4989-5146.

102. The engineered nuclease system of any one of claims 1-27 or claims 100-101, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5679-5686.

103. The engineered nuclease system of any one of claims 1-27 or claims 100-102, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1354 or SEQ ID NOs: 1354-1511.

104. The engineered nuclease system of any one of claims 1-27 or claims 100-103, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5472 or SEQ ID NOs: 5508.

105. The engineered nuclease system of any one of claims 1-27 or claims 100-104, wherein said endonuclease is configured to binding to a PAM comprising a sequence selected from the group consisting of SEQ ID NO: 5523 or SEQ ID NO: 5537.

106. The engineered nuclease system of any one of claims 1-27 or claims 100-105, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3175;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5472 or SEQ ID NO: 5508; and

- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5523 or SEQ ID NO: 5537.

107. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 3331 or SEQ ID NOs: 3331-3474.

108. The engineered nuclease system of any one of claims 1-27 or claim 107, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NOs: 5147 or SEQ ID NOs: 5147-5290.

109. The engineered nuclease system of any one of claims 1-27 or claims 107-108, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5674-5675 and SEQ ID NOs: 5687-5693.

110. The engineered nuclease system of any one of claims 1-27 or claims 107-109, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1512 or SEQ ID NOs: 1512-1655.

111. The engineered nuclease system of any one of claims 1-27 or claims 107-110, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5473 or SEQ ID NO: 5509.

112. The engineered nuclease system of any one of claims 1-27 or claims 107-111, wherein said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5524.

113. The engineered nuclease system of any one of claims 1-27 or claims 107-112, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3331;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5473 or SEQ ID NO: 5509; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5524.

114. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 3475 or SEQ ID NOs: 3475-3568.

115. The engineered nuclease system of any one of claims 1-27 or claim 114, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5291 or SEQ ID NOs: 5291-5389.

116. The engineered nuclease system of any one of claims 1-27 or claims 114-115, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5694-5699.

117. The engineered nuclease system of any one of claims 1-27 or claims 114-116, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1656 or SEQ ID NOs: 1656-1755.

118. The engineered nuclease system of any one of claims 1-27 or claims 114-117, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5474 or SEQ ID NO: 5510.

119. The engineered nuclease system of any one of claims 1-27 or claims 114-118, wherein said endonuclease is configured to binding to a PAM comprising SEQ ID NOs: 5525.

120. The engineered nuclease system of any one of claims 1-27 or claims 114-119, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3475;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5474 or SEQ ID NO: 5510; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5525.

121. The engineered nuclease system of any one of claims 1-27, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 3569 or SEQ ID NOs: 3569-3637.

122. The engineered nuclease system of any one of claims 1-27 or claim 121, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 5390 or SEQ ID NOs: 5390-5460.

123. The engineered nuclease system of any one of claims 1-27 or claims 121-122, wherein said endonuclease comprises at least 1, at least 2, at least 3, at least 4, or at least 5 peptide motifs selected from the group consisting of SEQ ID NOs: 5700-5717.

124. The engineered nuclease system of any one of claims 1-27 or claims 121-123, wherein said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to a sequence selected from the group consisting of SEQ ID NO: 1756 or SEQ ID NOs: 1756-1826.

125. The engineered nuclease system of any one of claims 1-27 or claims 121-124, wherein said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5475 or SEQ ID NOs: 5511.

126. The engineered nuclease system of any one of claims 1-27 or claims 121-125, wherein said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5526.

127. The engineered nuclease system of any one of claims 1-27 or claims 121-126, wherein:

- a) said endonuclease comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 3569;
- b) said guide RNA structure comprises a sequence at least 70%, 80%, or 90% identical to SEQ ID NO: 5475 or SEQ ID NO: 5511; and
- c) said endonuclease is configured to binding to a PAM comprising SEQ ID NO: 5526.

128. The engineered nuclease system of any one of claims 1-127, wherein said sequence identity is determined by a BLASTP, CLUSTALW, MUSCLE, MAFFT, or Smith-Waterman homology search algorithm.

129. The engineered nuclease system of claim 128, wherein said sequence identity is determined by said BLASTP homology search algorithm using parameters of a wordlength (W)

of 3, an expectation (E) of 10, and a BLOSUM62 scoring matrix setting gap costs at existence of 11, extension of 1, and using a conditional compositional score matrix adjustment.

130. An engineered guide ribonucleic acid polynucleotide comprising:

- a) a DNA-targeting segment comprising a nucleotide sequence that is complementary to a target sequence in a target DNA molecule; and
- b) a protein-binding segment comprising two complementary stretches of nucleotides that hybridize to form a double-stranded RNA (dsRNA) duplex,

wherein said two complementary stretches of nucleotides are covalently linked to one another with intervening nucleotides, and

wherein said engineered guide ribonucleic acid polynucleotide is configured to forming a complex with an endonuclease comprising a RuvC_III domain having at least 75% sequence identity to any one of SEQ ID NOs: 1827-3637 and targeting said complex to said target sequence of said target DNA molecule.

131. The engineered guide ribonucleic acid polynucleotide of claim 130, wherein said DNA-targeting segment is positioned 5' of both of said two complementary stretches of nucleotides.

132. The engineered guide ribonucleic acid polynucleotide of any of claims 130-131, wherein:

- a) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5476-5479 or SEQ ID NOs: 5476-5489;
- b) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of (SEQ ID NOs: 5490-5491 or SEQ ID NOs: 5490-5494) and SEQ ID NO: 5538;
- c) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5498-5499;
- d) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5495-5497 and SEQ ID NOs: 5500-5502;

- e) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5503;
- f) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5504;
- g) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NOs: 5505;
- h) protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5506;
- i) protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5507;
- j) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5508;
- k) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5509;
- l) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5510; or
- m) said protein binding segment comprises a sequence having at least 70%, at least 80%, or at least 90% identity to SEQ ID NO: 5511.

133. The engineered guide ribonucleic acid polynucleotide of any of claim 130-132, wherein:

- a) said guide ribonucleic acid polynucleotide comprises an RNA sequence comprising a hairpin comprising a stem and a loop, wherein said stem comprises at least 10, at least 12, or at least 14 base-paired ribonucleotides, and an asymmetric bulge within 4 base pairs of the loop;
- b) said guide ribonucleic acid polynucleotide comprises a tracr ribonucleic acid sequence predicted to comprise a hairpin comprising at least 8, at least 10, or at least 12 base-paired ribonucleotides;
- c) said guide ribonucleic acid polynucleotide comprises a guide ribonucleic acid sequence predicted to comprise a hairpin with an uninterrupted base-paired region comprising at least 8 nucleotides of a guide ribonucleic acid sequence and at least 8 nucleotides of a tracr ribonucleic acid sequence, and wherein said tracr ribonucleic acid sequence comprises, from 5' to 3', a first hairpin and a second hairpin, wherein said first hairpin has a longer stem than said second hairpin; or

- d) said guide ribonucleic acid polynucleotide comprises a tracr ribonucleic acid sequence predicted to comprise at least two hairpins comprising less than 5 base-paired ribonucleotides.

134. A deoxyribonucleic acid polynucleotide encoding the engineered guide ribonucleic acid polynucleotide of any one of claims 130-133.

135. A nucleic acid comprising an engineered nucleic acid sequence optimized for expression in an organism, wherein said nucleic acid encodes a class 2, type II Cas endonuclease comprising a RuvC_III domain and an HNH domain, and wherein said endonuclease is derived from an uncultivated microorganism.

136. A nucleic acid comprising an engineered nucleic acid sequence optimized for expression in an organism, wherein said nucleic acid encodes an endonuclease comprising a RuvC_III domain having at least 70% sequence identity to any one of SEQ ID NOs: 1827-3637.

137. The nucleic acid of any one of claims 135-136, wherein said endonuclease comprises an HNH domain having at least 70% or at least 80% sequence identity to any one of SEQ ID NOs: 3638-5460.

138. The nucleic acid of any one of claims 135-137, wherein said endonuclease comprises SEQ ID NOs: 5572-5591 or a variant thereof having at least 70% sequence identity thereto.

139. The nucleic acid of any one of claims 135-138, wherein said endonuclease comprises a sequence encoding one or more nuclear localization sequences (NLSs) proximal to an N- or C-terminus of said endonuclease.

140. The nucleic acid of claim 139, wherein said NLS comprises a sequence selected from SEQ ID NOs: 5597-5612.

141. The nucleic acid of any one of claims 135-140, wherein said organism is prokaryotic, bacterial, eukaryotic, fungal, plant, mammalian, rodent, or human.

142. The nucleic acid of claim 141, wherein said organism is *E. coli*, and wherein:

- a) said nucleic acid sequence has at least 70%, 80%, or 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5572-5575;
- b) said nucleic acid sequence has at least 70%, 80%, or 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5576-5577;
- c) said nucleic acid sequence has at least 70%, 80%, or 90% identity to a sequence selected from the group consisting of SEQ ID NOs: 5578-5580;
- d) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5581;
- e) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5582;
- f) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5583;
- g) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5584;
- h) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5585;
- i) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5586; or
- j) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5587.

143. The nucleic acid of claim 141, wherein said organism is human, and wherein:

- a) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5588 or SEQ ID NO: 5589; or
- b) said nucleic acid sequence has at least 70%, 80%, or 90% identity to SEQ ID NO: 5590 or SEQ ID NO: 5591.

144. A vector comprising a nucleic acid sequence encoding a class 2, type II Cas endonuclease comprising a RuvC_III domain and an HNH domain, wherein said endonuclease is derived from an uncultivated microorganism.

145. A vector comprising the nucleic acid of any of claims 135-143.

146. The vector of any of claims 144-145, further comprising a nucleic acid encoding an engineered guide ribonucleic acid structure configured to form a complex with said endonuclease comprising:

- a) a guide ribonucleic acid sequence configured to hybridize to a target deoxyribonucleic acid sequence; and
- b) a tracr ribonucleic acid sequence configured to binding to said endonuclease.

147. The vector of any of claims 144-146, wherein the vector is a plasmid, a minicircle, a CELiD, an adeno-associated virus (AAV) derived virion, or a lentivirus.

148. A cell comprising the vector of any of claims 144-147.

149. A method of manufacturing an endonuclease, comprising cultivating said cell of claim 146.

150. A method for binding, cleaving, marking, or modifying a double-stranded deoxyribonucleic acid polynucleotide, comprising:

- (a) contacting said double-stranded deoxyribonucleic acid polynucleotide with a class 2, type II Cas endonuclease in complex with an engineered guide ribonucleic acid structure configured to bind to said endonuclease and said double-stranded deoxyribonucleic acid polynucleotide;
- (b) wherein said double-stranded deoxyribonucleic acid polynucleotide comprises a protospacer adjacent motif (PAM); and
- (c) wherein said PAM comprises a sequence selected from the group consisting of SEQ ID NOs: 5512-5526 or SEQ ID NOs: 5527-5537.

151. A method of claim 149, wherein said double-stranded deoxyribonucleic acid polynucleotide comprises a first strand comprising a sequence complementary to a sequence of said engineered guide ribonucleic acid structure and a second strand comprising said PAM.

152. A method of claim 151, wherein said PAM is directly adjacent to the 3' end of said sequence complementary to said sequence of said engineered guide ribonucleic acid structure.

153. A method of any one of claims 149-152, wherein said class 2, type II Cas endonuclease is not a Cas9 endonuclease, a Cas14 endonuclease, a Cas12a endonuclease, a Cas12b endonuclease, a Cas 12c endonuclease, a Cas12d endonuclease, a Cas12e endonuclease, a Cas13a endonuclease, a Cas13b endonuclease, a Cas13c endonuclease, or a Cas 13d endonuclease.

154. A method of any one of claims 149-153, wherein said class 2, type II Cas endonuclease is derived from an uncultivated microorganism.

155. A method of any one of claims 149-154, wherein said double-stranded deoxyribonucleic acid polynucleotide is a eukaryotic, plant, fungal, mammalian, rodent, or human double-stranded deoxyribonucleic acid polynucleotide.

156. A method of any one of claims 149-155, wherein:

- a) said PAM comprises a sequence selected from the group consisting of SEQ ID NOs: 5512-5515 and SEQ ID NOs: 5527-5530;
- b) said PAM comprises SEQ ID NO: 5516 or SEQ ID NO: 5531;
- c) said PAM comprises SEQ ID NO: 5539;
- d) said PAM comprises SEQ ID NO: 5517 or SEQ ID NO: 5518;
- e) said PAM comprises SEQ ID NO: 5519;
- f) said PAM comprises SEQ ID NO: 5520 or SEQ ID NO: 5535;
- g) said PAM comprises SEQ ID NO: 5521 or SEQ ID NO: 5536;
- h) said PAM comprises SEQ ID NO: 5522;
- i) said PAM comprises SEQ ID NO: 5523 or SEQ ID NO: 5537;
- j) said PAM comprises SEQ ID NO: 5524;
- k) said PAM comprises SEQ ID NO: 5525; or
- l) said PAM comprises SEQ ID NO: 5526.

157. A method of modifying a target nucleic acid locus, said method comprising delivering to said target nucleic acid locus said engineered nuclease system of any one of claims 1-129, wherein said endonuclease is configured to form a complex with said engineered guide ribonucleic acid structure, and wherein said complex is configured such that upon binding of said complex to said target nucleic acid locus, said complex modifies said target nucleic locus.

158. A method of claim 156, wherein modifying said target nucleic acid locus comprises binding, nicking, cleaving, or marking said target nucleic acid locus.

159. A method of any of claims 156-158, wherein said target nucleic acid locus comprises deoxyribonucleic acid (DNA) or ribonucleic acid (RNA).

160. A method of claim 159, wherein said target nucleic acid comprises genomic DNA, viral DNA, viral RNA, or bacterial DNA.

161. A method of any one of claims 156-160, wherein said target nucleic acid locus is *in vitro*.

162. A method of any one of claims 156-160, wherein said target nucleic acid locus is within a cell.

163. A method of claim 162, wherein said cell is a prokaryotic cell, a bacterial cell, a eukaryotic cell, a fungal cell, a plant cell, an animal cell, a mammalian cell, a rodent cell, a primate cell, or a human cell.

164. A method of any one of claims 162-163, wherein delivering said engineered nuclease system to said target nucleic acid locus comprises delivering the nucleic acid of any of claims 135-140 or the vector of any of claims 142-146.

165. A method of any one of claims 162-163, wherein delivering said engineered nuclease system to said target nucleic acid locus comprises delivering a nucleic acid comprising an open reading frame encoding said endonuclease.

166. A method of claim 164, wherein said nucleic acid comprises a promoter to which said open reading frame encoding said endonuclease is operably linked.

167. A method of any one of claims 162-163, wherein delivering said engineered nuclease system to said target nucleic acid locus comprises delivering a capped mRNA containing said open reading frame encoding said endonuclease.

168. A method of any one of claims 162-163, wherein delivering said engineered nuclease system to said target nucleic acid locus comprises delivering a translated polypeptide.

169. A method of any one of claims 162-163, wherein delivering said engineered nuclease system to said target nucleic acid locus comprises delivering a deoxyribonucleic acid (DNA) encoding said engineered guide ribonucleic acid structure operably linked to a ribonucleic acid (RNA) pol III promoter.

170. A method of any one of claims 156-169, wherein said endonuclease induces a single-stranded break or a double-stranded break at or proximal to said target locus.






FIG. 1

Cas9 programmed by crRNA:tracrRNA duplex**Cas9 programmed by single chimeric RNA****FIG. 2**




FIG. 3




FIG. 4




FIG. 5

6 / 43




FIG. 6

7 / 43




FIG. 8




FIG. 9A




FIG. 9D




FIG. 9E




FIG. 9F




FIG. 10




FIG. 11




FIG. 12





FIG. 13




FIG. 14




FIG. 15





FIG. 16







FIG. 17



Gel 4



Gel 5



Gel 6

FIG. 18






FIG. 19

27 / 43



Gel 10



Gel 11

FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25






FIG. 26


MG1-4 Native PAM



MG1-4 Synthetic PAM



MG1-5 Native PAM



MG1-5 Synthetic PAM






FIG. 27


MG1-6 Native PAM



MG1-6 Synthetic PAM



MG1-7 Native PAM



MG2-1 Native PAM






FIG. 28

36 / 43


MG2-7 Synthetic PAM



MG3-6 Native PAM



MG3-6 Synthetic PAM



MG3-7 Native PAM






FIG. 29

37 / 43


MG3-7 Synthetic PAM



MG3-8 Native PAM



MG4-5 Synthetic PAM



MG14-1 Native PAM






FIG. 30

38 / 43


MG14-1 Synthetic PAM



MG15-1 Native PAM



MG15-1 Synthetic PAM



MG16-2 Synthetic PAM






FIG. 31

39 / 43


MG18-1 Native PAM



MG18-1 Synthetic PAM



MG21-1 Synthetic PAM



MG22-1 Synthetic PAM




FIG. 32

MG23-1 Synthetic PAM




FIG. 33




FIG. 34





FIG. 35




FIG. 36

A. CLASSIFICATION OF SUBJECT MATTER**C12N 9/22(2006.01)i, C12N 15/113(2010.01)i, C12N 15/10(2006.01)i, C12N 15/90(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N 9/22; C12N 15/00; C12N 5/00; C12N 15/113; C12N 15/10; C12N 15/90

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: RuvC III domain, HNH domain, uncultivated microorganism, guide RNA, tracr RNA, protospacer adjacent motif (PAM), CAS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JINEK, M. et al. 'A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity', Science, 17 August 2012, Vol. 337, pages 816-821 abstract, pages 816, 817, 820, figure 1, 2	1-8, 11, 12, 130-132 , 135-137, 144, 150
A	GASIUNAS, G. et al. 'Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria', PNAS, 25 September 2012, Vol. 109, No. 39, pages E2579-E2586 the whole document	1-8, 11, 12, 130-132 , 135-137, 144, 150
A	SHMAKOV, S. et al. 'Discovery and functional characterization of diverse class 2 CRISPR-Cas systems', Molecular Cell, 5 November 2015, Vol. 60, pages 1-13 the whole document	1-8, 11, 12, 130-132 , 135-137, 144, 150
A	SHMAKOV, S. et al. 'Diversity and evolution of class 2 CRISPR-Cas systems', Nature Reviews Microbiology, March 2017, Vol. 15, pages 169-182 the whole document	1-8, 11, 12, 130-132 , 135-137, 144, 150
A	US 8889418 B2 (THE BROAD INSTITUTE, INC. et al.) 18 November 2014 the whole document	1-8, 11, 12, 130-132 , 135-137, 144, 150



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

30 June 2020 (30.06.2020)

Date of mailing of the international search report

30 June 2020 (30.06.2020)

Name and mailing address of the ISA/KR

International Application Division

Korean Intellectual Property Office


189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

HEO, Joo Hyung

Telephone No. +82-42-481-8150



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2020/018432

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 8889418 B2	18/11/2014	AU 2013-359212 A1	30/07/2015
		AU 2013-359212 B2	19/01/2017
		AU 2014-281026 A1	28/01/2016
		AU 2014-281026 A9	16/06/2016
		AU 2014-281026 B2	28/05/2020
		AU 2016-244241 A1	03/11/2016
		AU 2016-244241 B2	20/06/2019
		AU 2017-202542 A1	11/05/2017
		AU 2017-202542 B2	11/07/2019
		AU 2019-201608 A1	04/04/2019
		BR 112015013784 A2	11/07/2017
		CA 2894684 A1	19/06/2014
		CN 105209621 A	30/12/2015
		DK 2898075 T3	27/06/2016
		EP 2898075 A1	29/07/2015
		EP 2898075 B1	09/03/2016
		ES 2576126 T3	05/07/2016
		HK 1207119 A1	22/01/2016
		JP 2016-182140 A	20/10/2016
		JP 2016-501532 A	21/01/2016
		JP 2018-143253 A	20/09/2018
		JP 6395765 B2	26/09/2018
		KR 10-2015-0105634 A	17/09/2015
		PL 2898075 T3	30/09/2016
		PT 2898075 E	16/06/2016
		SG 10201801969T A	27/04/2018
		SG 11201504519T A	30/07/2015
		US 2014-0186919 A1	03/07/2014
		US 2014-0242700 A1	28/08/2014
		US 2014-0273234 A1	18/09/2014
		US 2014-0335620 A1	13/11/2014
		US 2017-0198269 A1	13/07/2017
		US 8865406 B2	21/10/2014
		US 8895308 B1	25/11/2014
		WO 2014-093635 A1	19/06/2014
		WO 2014-093635 A9	24/07/2014

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2020/018432

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 20, 129, 140, 142, 143, 149, 151, 152, 158, 160, 163, 166
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims 20, 129, 140, 142, 143, 149, 151, 152, 158, 160, 163 and 166 refer to an unsearchable claim which is not drafted in accordance with PCT Rule 6.4(a).

3. ☒ Claims Nos.: 9, 10, 13-19, 21-128, 133, 134, 138, 139, 141, 145-148, 153-157, 159, 161, 162, 164, 165, 167-170
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.