



(22) Date de dépôt/Filing Date: 2006/11/17
(41) Mise à la disp. pub./Open to Public Insp.: 2007/06/22
(30) Priorité/Priority: 2005/12/22 (US60/753,453)

(51) Cl.Int./Int.Cl. *A01H 4/00* (2006.01),
A01H 7/00 (2006.01), *A01N 3/00* (2006.01),
C12N 5/02 (2006.01), *C12N 5/04* (2006.01)

(71) Demandeur/Applicant:
WEYERHAEUSER COMPANY, US

(72) Inventeur/Inventor:
JAMRUSZKA-LEWIS, AMY M., US

(74) Agent: SMART & BIGGAR

(54) Titre : METHODES POUR CONSERVER DES EMBRYONS SOMATIQUES DE CONIFERES
(54) Title: METHODS FOR STORING CONIFER SOMATIC EMBRYO GERMINANTS

(57) **Abrégé/Abstract:**

In one aspect, the present invention provides methods of storing conifer somatic embryo germinants germinated on a sterile germination medium for delayed transplanting into a growth medium. The methods of the invention comprise the steps of: (a) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to six months, said germinants comprising a visible, well-defined epicotyl and radicle; and (b) placing the germinants in water for a time greater than about 1 hour at a temperature below about 24°C prior to transplantation into growth medium.



ABSTRACT OF THE DISCLOSURE

In one aspect, the present invention provides methods of storing conifer somatic embryo germinants germinated on a sterile germination medium for delayed transplanting into a growth medium. The methods of the invention comprise the steps of: (a) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to six months, said germinants comprising a visible, well-defined epicotyl and radicle; and (b) placing the germinants in water for a time greater than about 1 hour at a temperature below about 24°C prior to transplantation into growth medium.

METHODS FOR STORING CONIFER SOMATIC EMBRYO GERMINANTS

FIELD OF THE INVENTION

The present invention relates to methods for storing conifer somatic embryo
5 germinants.

BACKGROUND OF THE INVENTION

The demand for coniferous trees, such as pines and firs, to make wood products
continues to increase. One proposed solution to meeting this high demand is to identify
individual trees that possess desirable characteristics, such as a rapid rate of growth, and
10 produce numerous, genetically identical, clones of the superior trees by somatic cloning.

Somatic cloning is the process of producing plant embryos, *in vitro*, from plant cells
that are not zygotes. These clones can be cultivated to yield stands, or whole forests, of
conifer trees that possess the desirable characteristic(s). One method for somatically
cloning trees utilizes *in vitro* treatment of isolated, living, conifer tissue under conditions
15 that promote formation of conifer somatic embryos, and then whole plants from the treated
tissue. An explant such as an immature seed or the embryo from an immature seed is
placed on a gelled initiation medium. This medium will usually contain plant growth
hormones from the groups known as auxins and cytokinins. If initiation is successful a
gelatinous mass containing multiple immature embryos will be generated in several weeks.
20 This mass is then removed and subcultured on a maintenance and multiplication medium
which may be gelled, liquid, or some combination of these. Embryos from maintenance
medium may then be placed on a development medium that normally lacks the auxins and
cytokinins but may instead include the hormone abscisic acid. The somatic embryos
develop into cotyledonary embryos with a size and morphology that closely resembles their
25 mature zygotic counterparts. The embryos are then placed on a germination medium on
which radicle and epicotyl elongation occur over a period of several weeks. The germinants
are then removed and placed in soil for further development into plantlets. After a period of
greenhouse growth they may be outplanted. Alternatively, the embryos removed from the
development medium may be placed into manufactured seeds; e.g., as shown in Carlson et
30 al., U.S. Patent No. 5,236,469.

A continuing problem, however, has been that conversion percentage from somatic
embryos to plants growing in soil has frequently been lower than desired. In addition, a

logistical problem exists for handling large numbers of conifer somatic embryos that are not packaged into manufactured seeds. These embryos are individually removed from the sterile development medium and placed on a germination medium. After an appropriate time the germinants are placed in a potting soil mixture for further growth. Typically many
5 hundreds of thousands of germinants are potted at once in clonal field tests. The germinants are very tender and susceptible to damage by disease organisms such as fungi. In addition, it is preferable to outplant germinants that have similar dimensions in clonal field tests in order to obtain reliable results regarding the characteristics of various clones. However, a logistical challenge exists for producing a population of germinants having similar
10 dimensions at the time of outplanting.

There is therefore a continuing need for a method for storing conifer germinants that reduces the risk of contamination with microbes, is amenable to large scale production, and results in a high conversion rate after transplantation.

15 SUMMARY OF THE INVENTION

In one aspect, the present invention provides methods of storing conifer somatic embryo germinants germinated on a sterile germination medium for delayed transplanting into a growth medium. The methods of the invention comprise the steps of: (a) placing the germinants while still on sterile germination medium into a cold environment in which the
20 temperature is in the range of 0.5°C to 10°C for a time period up to 6 months, said germinants comprising a visible, well-defined epicotyl and radicle; and (b) placing the germinants in water for a time greater than about 1 hour at a temperature below about 24°C prior to transplantation into growth medium.

In another aspect, the present invention provides methods of producing and storing
25 germinants prior to transplantation into growth medium. The methods of this aspect of the invention comprise the steps of: (a) culturing conifer somatic embryos on sterile germination medium for a sufficient period of time to produce germinants, said germinants comprising a visible, well-defined epicotyl and radicle; (b) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in
30 the range of 0.5°C to 10°C for a time period up to 6 months, and (c) placing the germinants in water at a temperature below about 24°C for a time greater than about 1 hour prior to transplanting the stored germinants into growth medium.

In another aspect, the present invention provides methods of increasing the conversion percentage from somatic embryos to plants growing in growth medium. In accordance with this aspect of the invention the methods comprise the steps of: (a) culturing conifer somatic embryos on sterile germination medium for a sufficient period of time to
5 produce germinants, said germinants comprising a visible, well-defined epicotyl and radicle; (b) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to 6 months; (c) placing the germinants in water for a time greater than about 1 hour at a temperature below about 24°C; and (d) removing the germinants from the water and
10 transplanting them into growth medium for further growth.

In yet another aspect, the present invention provides methods of producing a synchronized population of somatic embryo germinants in preparation for transplantation. In accordance with this aspect of the invention the methods comprise the steps of: (a) culturing somatic embryos on sterile germination medium for a sufficient period of time to
15 produce germinants having a desired dimension; (b) selecting a plurality of germinants having the desired dimension; (c) storing the selected germinants while still on sterile germination medium in a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to 6 months; and (d) placing the stored germinants in water for a time greater than about 1 hour at a temperature below about 24°C prior to
20 planting the selected germinants into growth medium.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The following examples merely illustrate the best mode now contemplated for practicing the invention, but should not be construed to limit the invention.

As used herein, the term "cotyledonary embryo" means an embryo that possesses
25 one or more cotyledons.

As used herein, the term "somatic embryo" refers to a plant embryo that developed *in vitro* from a plant cell, or tissue, that is not a zygote.

As used herein, the term "embryogenic tissue" refers to any tissue, derived from a conifer which is capable of producing one or more conifer cotyledonary somatic embryos,
30 including, for example, conifer embryonal suspensor masses.

As used herein the term "germinant" refers to an immature plant that possesses a well developed radicle and cotyledonary structure with a growing epicotyl, both readily

apparent to the naked eye, and ready for planting in soil. For example, the germinants typically have an epicotyl of about 10mm or greater.

As used herein, the term "converted embryo" is an embryo that has germinated and been established as a plant growing in soil.

5 As used herein, the term "synchronized population of germinants" refers to a stock of germinants having a target specification at outplanting. The germinants may be cultured to a desired target specification, such as having a desired dimension, (e.g., an epicotyl stem length of at least 10 mm).

10 Unless stated otherwise, all concentration values that are expressed as percentages are weight per volume percentages.

The present invention provides methods for storing conifer somatic embryo germinants. The methods each include the step of placing the germinants while still on germination medium into a cold environment for a time period of up to 6 months (such as, for example, from at least 2 weeks to 24 weeks, or such as from 4 weeks to 10 weeks),
15 followed by the step of placing the germinants in water for a time greater than about 1 hour up to from one to several weeks (such as, for example from several hours up to one week) prior to transplantation into a growth medium. It has been observed that conifer germinants may be stored up to 4 weeks or more prior to planting in growth medium when stored according to the methods of the invention without loss of post-transplant vigor.

20 The storage methods of the invention are particularly advantageous in that they make it possible to produce a synchronized population of germinants in preparation for transplantation. The synchronized population of germinants are produced by culturing germinants to a desired target specification, and storing the germinants in a cold environment on sterile germination medium as they become available over a period of days
25 or weeks. The stored germinants are then placed into water for a time greater than about 1 hour at a temperature below about 24°C prior to planting. Consequently, the storage methods of the invention allow for a longer embryo production window (e.g. several weeks to several months) in which to obtain a desired population of germinants, which can then be outplanted within a narrow time frame (e.g. one day up to two weeks), or during a winter
30 dormant period, thereby resulting in a synchronized population at outplanting. The ability to synchronize the production of germinants in preparation for transplantation is particularly useful in the context of planting clonal field tests.

The methods of the present invention can be used to store conifer somatic embryo germinants from any conifer species, such as members of the family Pinacea, including members of the genus *Pinus* (e.g., loblolly pine (*Pinus taeda*)), or such as members of the genus *Pseudotsuga* (e.g., Douglas-fir (*Pseudotsuga menziesii*)).

5 The germinants may be produced by any method that yields conifer somatic embryo germinants. For example, germinants may be produced by first culturing conifer somatic embryos by known protocols, such as the methods described in U.S. Patent Nos. 6,134,830 and 5,563,061. After the embryos have reached a mature cotyledonary stage they are then placed on a germination medium under environmental conditions of 21°C to 27°C for a
10 sufficient period of time for an epicotyl and radicle to develop. This time period may range between 4 and 14 weeks, depending on the particular species and genotype. More typically it is between 8 and 12 weeks. A suitable light/dark photoperiod may be used, such as a 24 hour photoperiod.

The germination medium typically has no exogenous hormones, a lowered nitrogen
15 content and a reduced level of osmoticants. The germination medium typically contains agar and is sterilized prior to use. The germinants may be germinated under suitable conditions on solid germination media contained in any suitable container, such as in Petri dishes stored in germination boxes.

An example of a suitable germination medium for Douglas-fir (*Pseudotsuga
20 menziesii*) is provided below in TABLE 1.

TABLE 1: *Pseudotsuga menziesii* BM_G Germination Media

Constituent	Concentration, mg/L
NH ₄ NO ₃	206.3
KNO ₃	1170.0
CaCl ₂ .6H ₂ O	220.0
KH ₂ PO ₄	85.0
MgSO ₄ .7H ₂ O	185.0
MnSO ₄ .H ₂ O	8.45
ZnSO ₄ .7H ₂ O	4.30
CuSO ₄ .5H ₂ O	0.013
FeSO ₄ .7H ₂ O	13.93
Na ₂ EDTA	18.63

H ₃ BO ₃	3.10
NaMoO ₄ .2H ₂ O	0.125
CoCl ₂ .6H ₂ O	0.0125
KI	0.42
myo-Inositol	100.0
Thiamine.HCL	1.00
Nicotinic acid	0.50
Pyridoxine.HCL	0.50
Glycine	2.00
L-Glutamine	450.0
Sugar	20,000
pH	5.7
Activated charcoal	2500
Tissue culture agar	8000

In accordance with the methods of the invention, once developed, the germinants, while still on the sterile germination medium, are placed into a cold environment in which the temperature is in the range of about 0.5°C to about 10°C, preferably from about 4°C to about 8°C. Preferably the container comprising the germinantes on the germination media is placed into the cold storage environment without any manipulation of the germinants required.

The germinants may be stored in the cold environment until transplantation into growth medium for at least two weeks up to about 6 months (such as from 1 week to 20 weeks, or more preferably from about 2 weeks to about 4 weeks) without loss of post-transplant vigor, as described in EXAMPLES 1 and 2.

After the desired time of storage on germination medium and prior to transplant, the germinants are removed from the germination medium and placed in deionized water in sealed dishes. In some embodiments of the method, the germinants are transferred under sterile conditions, such as under a laminar flow hood. In other embodiments of the method, the germinants are transferred under non-sterile conditions. The germinants are stored in water for a time greater than about 1 hour at a temperature below about 24°C. Preferred temperatures are below 10°C, such as from about 0.5°C to about 5°C. Improved rates of

post transplant survival have been found for germinants stored in water for a period greater than 1 hour to about three weeks, as described in EXAMPLE 2.

The germinants stored in accordance with the method of the invention are eventually transferred to growth medium for further growth which eventually may result in conifer trees. The growth medium may be any suitable medium, such as potting soil, or a mixture of peat, vermiculate and perlite. The potted germinants are typically grown in a greenhouse and maintained at 24°C for a 16 hour light period and at 18°C for an 8 hour dark period. The plants may be compared and rated for survival percentage as described in EXAMPLES 1 and 2.

10 In another aspect, the present invention provides methods of increasing the conversion percentage from somatic embryos to plants growing in growth medium. The method according to this aspect of the invention comprise the steps of: (a) culturing conifer somatic embryos on sterile germination medium for a sufficient period of time to produce germinants, said germinants comprising a visible, well-defined epicotyl and radicle; 15 (b) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to 6 months; (c) placing the germinants in water for a time greater than about 1 hour at a temperature below about 24°C; and (d) removing the germinants from the water and transplanting them into growth medium for further growth. The methods of this aspect of 20 the invention can be used to increase the conversion percentage from somatic embryos to plants growing in growth medium for any conifer species, such as members of the family Pinacea, including members of the genus *Pinus* (e.g., loblolly pine (*Pinus taeda*)), or such as members of the genus *Pseudotsuga* (e.g., Douglas-fir (*Pseudotsuga menziesii*)).

In some embodiments, the germinants are stored on sterile germination media for a 25 time period from about 1 week up to 6 months, such as from 2 weeks to 20 weeks, or more preferably from about 2 weeks to about 4 weeks. In some embodiments, the germinants that have been pre-stored on germination media and then stored in water for a time period from about 1 hour up to 2 weeks, such as from about 1 week to about 2 weeks, more preferably about 1 week.

30 The following examples merely illustrate the best mode now contemplated for practicing the invention, but should not be construed to limit the invention.

EXAMPLE 1

This Example describes the results of an experiment demonstrating that a two-step storage treatment on Douglas-fir germinants increases the sterile storage period without causing detrimental effects to greenhouse survival.

5 Methods and Materials:

Germinants from ten different genotypes of Douglas-fir somatic embryos were produced up to the development stage using the methods described in Gupta, U.S. Patent No. 5,563,061 and Welty, U.S. Patent No. 6,134,830. Following the development stage, the cotyledonary embryos were placed on an agar gelled sterile germination medium, BM6, in
10 Petri dishes for 4-12 weeks under environmental conditions of 21°C to 27°C until good epicotyl development was apparent (e.g., an epicotyl stem length of about 10mm or greater, such as at least from 10mm to 20mm, such as at least 12mm, at least 15mm or greater). BM6 medium is a basal medium lacking growth hormones which has been modified by reducing sucrose, myo-inositol and organic nitrogen.

15 Once good epicotyl development was apparent, the closed germination containers containing the germinants on sterile germination medium were moved directly into a cooler at a temperature of about 4°C and stored in the dark for various amount of time up to 6 months. At various time points after storage on germination media, the germinants were removed from the germination media and placed into Petri dishes that were half-filled with
20 Millipore or Nanopure water and then sealed with parafilm. The roots of the germinants were immersed in water while the shoots floated on the surface. The germinants in water were transferred to a cold environment at a temperature of about 4°C for a period of time up to 4 weeks.

Although the germinants were stored under sterile conditions up to this point, it was
25 determined that the transfer, under a laminar flow hood, from the germination media to the Petri dishes containing water could result in desiccation of the germinants. Consequently, the transfer from germination media to water storage was not made under sterile conditions.

Germinants from ten different genotypes of Douglas-fir were treated as follows. Group I control samples were germinated on sterile germination media, then removed from
30 the germination media and transplanted immediately into potting media without a period of storage. Group II germinants were germinated on sterile germination media in germination boxes, and the germination boxes were then moved into a dark cooler at 4°C for 1 week

followed by one week in water storage. Group III germinants were germinated on sterile germination media in germination boxes, the boxes were moved into a dark cooler at 4°C for 2 weeks, followed by one week in water storage. Upon removal from storage, the Petri dishes and boxes were inspected for signs of contamination. No contamination was found during the following experiments. The germinants were transplanted into potting media and the percent survival was measured at 8 weeks post transplant. The potting media consisted of equal parts of peat, vermiculate and perlite. The percent survival of the germinants treated under the various conditions are shown below in TABLE 2.

10 TABLE 2: Douglas-fir Percent Survival Post-transplant

Time held in storage	Douglas-fir Genotype (percent survival 8 weeks post-transplant)										Treatment Average	Paired Test
	7306	7435	3637	5022	5008	5010	5002	7667	663	7744		
Control (no storage)	64.3	22.6	67.1	60.5	46.1	37.3	32.3	91.7	53.9	89.2	56.5	0.813
1 week on Germination media + 1 week in H ₂ O	63.5	13	51.4	67.2	41.1	40.5	48.1	100.0	51.6	90.5	56.7	
2 weeks on Germination media + 1 week in H ₂ O	72.6	21.7	59.3	72.3	48.6	39.4	36.3	96.3	38.8	88.0	57.3	

Results:

As shown above in TABLE 2, greenhouse survival post transplant for germinants stored on germination media followed with one week of storage in water was at least as good, and in some cases better than that observed for water storage alone.

Conclusion:

The method of long term storage (e.g., 2 to 4 weeks, and likely up to 6 months) of Douglas-fir germinants on germination media followed by short term storage (e.g., 1 to 2 weeks) in water results in improved survival rates for transplants and provides a convenient method by which to accumulate germinants prior to transplant. This method provides several advantages including increased survival post-transplant, convenient storage of large numbers of germinants, and a reduced risk of microbe contamination prior to transplantation.

EXAMPLE 2

This Example describes experiments that measure the post-transplant survival of various loblolly pine genotypes after storage of germinants in water for various lengths of time as compared to the post-transplant survival of germinants that were stored on germination media followed with one week of storage in water.

Methods and Materials:*Germinants Stored in Water*

A control experiment was carried out to determine the effect of storage in water for various time periods on the post-transplant survival of loblolly pine clones. Germinants from three different loblolly pine genotypes were produced up to the germination stage using the methods described in Gupta, U.S. Patent No. 5,563,061 and Welty, U.S. Patent No. 6,134,830. The germinants for each clone were removed from germination media and placed into Petri dishes that were half filled with Millipore or Nanopure water and then sealed with parafilm. The Petri dishes were stacked in a labeled crisper (1 genotype to crisper) and then put into a dark walk-in cooler (approximately 4°C) for the length of time indicated below in TABLE 3 prior to transplant. A control batch of germinants from each clone was transplanted in potting medium immediately without storage in water. The potting medium consisted of equal parts of peat, vermiculite, and perlite. The potted germinants were held for further growth in a greenhouse maintained at 24°C for a 16 hour light period and at 18°C for an 8 hour dark period. After 8 weeks or more the plants were rated for survival percentage. The results are shown below in TABLE 3.

Germinants Stored on Germination Media followed by Short Term Storage in Water

Germinants from different loblolly pine genotypes were produced up to the germination stage using the methods described in Gupta, U.S. Patent No. 5,563,061 and Welty, U.S. Patent No. 6,134,830. The germinants from the different genotypes were stored on sterile germination media in the dark at 4°C for various time periods, as shown in TABLE 4, then the germinants were removed from germination media and placed into Petri dishes that were half-filled with Millipore or Nanopure water and then sealed with parafilm and stored at 4°C for a week prior to transplantation. The results are shown in TABLE 4 and TABLE 5.

10

TABLE 3: Loblolly Pine Survival Post-Transplant After Storage in Water

Time in Water Storage	Loblolly Pine Genotype (percent survival 8 or more weeks post transplant)				3 Clone Average	4 Clone Average	Pr >t Paired test (3 or 4 genotypes)
	55	56	68	75			
Control (not stored)	-	99.1	96	88.6	94.6	-	0.966
1 week	-	98.6	93.8	91.1	94.5	-	
Control (not stored)	94.4	99.1	96	86.5	93.9	94.0	0.164
2 weeks	88.9	96.7	84.3	50.8	77.3	80.2	
Control (not stored)	-	98.8	96	87	93.9	-	0.860
3 weeks	-	96.9	96	89.7	94.2	-	
Control (not stored)	93.6	97.8	96.2	88.5	94.2	94.0	0.728
4 weeks	95.7	96.5	98.0	83.3	92.6	93.4	

Control (not stored)	-	98.3	96	86.6	93.6	-	0.147
5 weeks	-	96.9	84	71.9	84.3	-	

TABLE 4: Loblolly Pine Survival Post-transplant After Storage on Germination Media Followed by Storage in Water

Time held in Storage	Loblolly Pine Genotype (percent survival 8 or more weeks post transplant)			3-Clone Average	Pr > t Paired test
	55	56	75		
Control= 0 days on germination media + 21 days in H ₂ O	100	95.5	79.1	91.5	-
14 days on germination media + 7 days in H ₂ O	100	93.4	77.6	90.3	0.195
21 days on germination media + 7 days in H ₂ O	97.3	98.4	82.1	92.6	0.628
Control= 0 days on germination media + 21 days in H ₂ O	100	100	94.6	98.2	-
28 days on germination media + 7 days in H ₂ O	100	92.5	91.3	94.6	0.239

5

TABLE 5: Loblolly Pine Survival Post-transplant After Storage on Germination Media Followed by Storage in Water

Time held in Storage	Loblolly Pine Genotype										Treatment Average	Treatment Average
	202	205	212	213	214	215	226	228	241	243	(10 genos)	(4 genos)
Control- No Storage	93.8	100	96.7	96.6	75	97.7	98	100	100	93.8	95.2	97.8
2 weeks on Germ media + 1 week in H ₂ O	-	-	100	96.3	-	-	97.7	100	-	-	-	98.5 ¹
4 weeks on Germ media + 1 week in H ₂ O	89.3	100	95.9	98	89.8	100	93.6	98.1	93.3	96.3	95.4 ²	96.4 ³

¹Paired t-test comparing 2-week Box Storage against the control (4 genos) (Pr>t) = 0.50

²Paired t-test comparing 4-week Box Storage against the control (10 genos) (Pr>t) = 0.89

³Paired t-test comparing 4-week Box Storage against the control (4 genos) (Pr>t) = 0.32

5

Results:

As shown above in TABLE 2, the results indicate that short term water storage (e.g., 1 week) increases the post-transplant survival of loblolly pine germinants, however longer term water storage (e.g., 2 to 4 weeks or longer) decreases the percent survival in the majority of clone genotypes (3/4) tested.

10

As shown above in TABLE 3 and TABLE 4, longer term storage on germination media (e.g., from 2 to 4 weeks or longer) followed by short term (e.g., 1 week) water storage resulted in post-transplant survival of loblolly pine germinants at survival rates at least as good as, and in some cases better than, germinants that were planted directly into potting media with no storage.

15

Conclusion

The method of long term storage (e.g., 2 to 4 weeks up to 6 months) of loblolly pine germinants on germination media followed by short term storage (e.g., 1 to 2 weeks) in water results in improved survival rates for transplants and provides a convenient method by which to accumulate germinants prior to transplant.

While the preferred embodiment of the invention has been illustrated and described, it will be appreciated that various changes can be made therein without departing from the spirit and scope of the invention.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A method of storing conifer somatic embryo germinants germinated on a sterile germination medium for delayed transplanting into a growth medium, comprising the
5 steps of:
 - (a) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to 6 months, said germinants comprising a visible, well-defined epicotyl and radicle; and
 - 10 (b) placing the germinants in water for a time greater than about 1 hour at a temperature below about 24°C prior to transplantation into growth medium.
2. The method of Claim 1 further comprising the step of planting at least a portion of the germinants into growth medium for further growth.
3. The method of Claim 1, wherein the germinants are stored in water for a time
15 from about one week to about two weeks.
4. The method of Claim 3, wherein the germinants are stored on germination medium for a time period from about one week to about four weeks.
5. The method of Claim 1, wherein the water temperature is in the range of about 0.5°C to about 5°C.
- 20 6. The method of Claim 1 in which the germinants are Douglas-fir.
7. The method of Claim 1 in which the germinants are loblolly pine.
8. A method of producing and storing germinants prior to transplantation into growth medium, comprising the steps of:
 - (a) culturing conifer somatic embryos on sterile germination medium for
25 a sufficient period of time to produce germinants, said germinants comprising a visible, well-defined epicotyl and radicle;

(b) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to six months, and

(c) placing the germinants in water at a temperature below about 24°C
5 for a time greater than about one hour prior to planting the stored germinants into growth medium.

9. The method of Claim 8, further comprising the step of planting the germinants into growth medium for further growth.

10. The method of Claim 8 in which the germinants are Douglas-fir.

10 11. The method of Claim 8 in which the germinants are loblolly pine.

12. A method of increasing the conversion percentage from somatic embryos to plants growing in growth medium, the method comprising:

(a) culturing somatic embryos on sterile germination medium for a sufficient period of time to produce germinants, said germinants comprising a visible, well-
15 defined epicotyl and radicle;

(b) placing the germinants while still on sterile germination medium into a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to six months;

(c) placing the germinants in water for a time greater than about 1 hour at
20 a temperature below about 24°C; and

(d) removing the germinants from the water and planting them into a growth medium for further growth.

13. The method of Claim 12, wherein the germinants are stored in water for a time from about one week to about two weeks.

25 14. The method of Claim 13, wherein the germinants are stored on germination medium for a term period from about one week to about four weeks.

15. The method of Claim 12 in which the germinants are Douglas-fir.

16. The method of Claim 12 in which the germinants are loblolly pine.

17. A method of producing a synchronized population of somatic embryo germinants in preparation for transplantation, comprising:

5 (a) culturing somatic embryos on sterile germination medium for a sufficient period of time to produce germinants having a desired dimension,

(b) selecting a plurality of germinants having the desired dimension;

(c) storing the selected germinants while still on sterile germination medium in a cold environment in which the temperature is in the range of 0.5°C to 10°C for a time period up to six months; and

10 (d) placing the stored germinants in water for a time greater than about 1 hour at a temperature below about 24°C prior to planting the selected germinants into growth medium.

18. The method of Claim 17, further comprising the step of planting the germinants in growth medium.

15 19. The method of Claim 17 wherein the desired dimension is an epicotyl stem length of at least 10mm.

20. The method of Claim 18, wherein the germinants are planted in a clonal field test.