

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3826632 T3**

(12) **Oversættelse af
europæisk patentskrift**

- (51) Int.Cl.: **A 61 K 31/4045 (2006.01)** **A 61 P 25/18 (2006.01)** **A 61 P 25/22 (2006.01)**
A 61 P 25/24 (2006.01) **A 61 P 25/30 (2006.01)** **C 07 D 209/16 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-07-11**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2022-05-18**
- (86) Europæisk ansøgning nr.: **20731372.7**
- (86) Europæisk indleveringsdag: **2020-06-02**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-06-02**
- (86) International ansøgning nr.: **EP2020065244**
- (87) Internationalt publikationsnr.: **WO2020245133**
- (30) Prioritet: **2019-06-03 GB 201907871**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Small Pharma Ltd, 6-8 Bonhill Street, London EC2A 4BX, Storbritannien**
- (72) Opfinder: **RANDS, Peter, c/o Small Pharma Ltd 6-8 Bonhill Street, London EC2A 4BX, Storbritannien**
JOEL, Zelah, c/o Small Pharma Ltd 6-8 Bonhill Street, London EC2A 4BX, Storbritannien
BENWAY, Tiffanie, c/o Small Pharma Ltd 6-8 Bonhill Street, London EC2A 4BX, Storbritannien
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **TERAPEUTISKE SAMMENSÆTNINGER, DER OMFATTER DEUTEREREDE ELLER DELVIST DEUTEREREDE N,N-DIMETHYLTRYPTAMINFORBINDELSER**
- (56) Fremdragne publikationer:
WO-A1-2019/081764
WO-A1-2021/089872
SIMON D. BRANDT ET AL: "Microwave-accelerated synthesis of psychoactive deuterated N , N -dialkylated-[alpha] , [alpha] , [beta] , [beta] -d 4]-tryptamines", JOURNAL OF LABELLED COMPOUNDS AND RADIOPHARMACEUTICALS, vol. 51, no. 14, 1 November 2008 (2008-11-01), pages 423-429, XP055724013, GB ISSN: 0362-4803, DOI: 10.1002/jlcr.1557
MORRIS ET AL.: "Indolealkylamine metabolism: Synthesis of deuterated indolealkylamines as metabolic probes", JOURNAL OF LABELLED COMPOUNDS & RADIOPHARMACEUTICALS, vol. 33, no. 6, 1993, pages 455-465, XP009522345, DOI: 10.1002/jlcr.2580330603 cited in the application
BEATON J M ET AL: "A comparison of the behavioral effects of proteo- and deuterio-N, N-dimethyltryptamine", PHARMACOLOGY BIOCHEMISTRY AND BEHAVIOR, ELSEVIER, US, vol. 16, no. 5, 1 May 1982 (1982-05-01), pages 811-814, XP023806754, ISSN: 0091-3057, DOI: 10.1016/0091-3057(82)90240-4
WALKER R W ET AL: "Gas chromatographic-mass spectrometric isotope dilution assay for N,N-dimethyltryptamine in human plasma", BIOCHEMICAL MEDICINE, ACADEMIC PRESS, SAN DIEGO, CA, US, vol. 8, no. 1, 1 August 1973 (1973-08-01), pages 105-113, XP026443695, ISSN: 0006-2944

Fortsættes ...

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates to compositions comprising: (a) an α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; (b) an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; and (c) *N,N*-dimethyltryptamine, or a pharmaceutically acceptable salt thereof; wherein the composition comprises 50% or more by weight of the α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof, and the α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof.

[0002] In particular, in a preferred aspect of the foregoing, the present invention relates to compositions comprising a combination of *N,N*-dimethyltryptamine and 2% or more by weight of a deuterated *N,N*-dimethyltryptamine compound selected from α,α -dideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine. In additional and alternative aspects of the foregoing, compositions of the present invention comprise a combination of *N,N*-dimethyltryptamine and 2% or more by weight of a partially deuterated *N,N*-dimethyltryptamine compound selected from α,β,β -trideutero-*N,N*-dimethyltryptamine, α,β -dideutero-*N,N*-dimethyltryptamine, and α -deutero-*N,N*-dimethyltryptamine. Methods of synthesising compositions of the present invention, and such compositions for use in treating psychiatric or psychocognitive disorders, such as major depressive disorder, are also provided.

BACKGROUND OF THE INVENTION

[0003] Classical psychedelics have shown preclinical and clinical promise in treating psychiatric disorders (Carhart-Harris and Goodwin (2017), The Therapeutic Potential of Psychedelic Drugs: Past, Present and Future, *Neuropsychopharmacology* 42, 2105-2113). In particular, psilocybin has demonstrated significant improvement in a range of depression and anxiety rating scales in randomised double blind studies (Griffiths et al. (2016), Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: a randomised double-blind trial, *Journal of Psychopharmacology* 30(12), 1181-1197).

[0004] *N,N*-dimethyltryptamine (DMT) is also understood to hold therapeutic value as a short-acting psychedelic, however its duration of action (under 20 minutes) is so short as to limit effective therapy. Administration protocols have been developed to extend the immersive psychedelic experience of DMT (Gallimore and Strassman (2016), A model for the application of target-controlled intravenous infusion for a prolonged immersive DMT psychedelic experience, *Frontiers in Pharmacology*, 7:211). However, these protocols carry risk of toxic buildup in patients who are poor metabolisers of DMT (for further discussion see Strassman et al (1994), Dose response study of *N,N*-dimethyltryptamine in humans, *Arch Gen Psychiatry* 51,

85).

[0005] $\alpha,\alpha,\beta,\beta$ -Tetradeutero-*N,N*-dimethyltryptamine is known to exhibit a kinetic isotope effect which bestows a significant difference on its *in vivo* pharmacokinetic profile as compared with *N,N*-dimethyltryptamine. Substitution of hydrogen with a deuterium at an sp^3 carbon centre is known to give rise to a 'kinetic isotope effect' by virtue of the difference in bond strength between a CH and a CD bond. First demonstrated in 1982 (Barker et al. (1982), Comparison of the brain levels of *N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine following intraperitoneal injection, *Biochemical Pharmacology*, 31(15), 2513-2516), the half-life of $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine in the rodent brain is suggestive that administration of $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine alone would maintain a patient in DMT space for longer than therapeutically essential.

[0006] WO 2019/081764 A1 describes a combination product for the treatment and/or prevention of psychiatric and/or neurological disorders, which comprises (i) a compound, for example *N,N*-dimethyltryptamine or $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine, which promotes neurogenesis and has hallucinogenic and/or psychedelic side effects, and (ii) a 5-HT_{2A} receptor antagonist which alleviates and/or removes the hallucinogenic and/or psychedelic side effects caused by the first compound.

[0007] Beaton et al. *Pharmacol. Biochem. Behav.* 1982, 16 811-814 describes a comparison of the behavioural effects of *N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine at dose levels of 2.5 and 5.0 mg/kg in rats.

SUMMARY OF THE INVENTION

[0008] The references to methods of treatment in this description are to be interpreted as references to the compounds, pharmaceutical compositions and medicaments of the present invention for use in a method for treatment of the human (or animal) body by therapy.

[0009] The present invention is based, in part, upon the ability to apply knowledge of the kinetic isotope effect exhibited by $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine in order to modify, controllably, the pharmacokinetic profile of *N,N*-dimethyltryptamine, thereby permitting more flexible therapeutic application. In particular, by providing individual drug substance compositions comprising *mixtures* of *N,N*-dimethyltryptamine and deuterated *N,N*-dimethyltryptamine analogues, in particular *N,N*-dimethyltryptamine comprising at least one deuterium atom at the alpha position (i.e. attached to the carbon atom to which the dimethylamino moiety is attached) and, according to certain embodiments, one or two deuterium atoms at the beta position, the present invention provides compositions and methods which enable a finely tuned single dose to maintain a patient in full dissociation from the external world, referred to herein as 'DMT space', for a therapeutically optimised duration without relying on infusion protocols or combination therapy with monoamine oxidase inhibitors

in the clinic. The present invention provides a clinically applicable solution which reduces clinical complexity and increases clinical flexibility in the administration of DMT-assisted psychotherapy.

[0010] Moreover, we have observed a quantifiable relationship between the extent of deuteration, and by proxy the H:D ratio of input reducing agent in synthetic methods disclosed herein, and the effect on potentiation of the metabolic half-life of the parent compound. Such technical effect may be used to quantifiably increase the precision with which compositions comprising pluralities of *N,N*-dimethyltryptamine compounds (that is to say comprising more than one type of compound selected from *N,N*-dimethyltryptamine and its deuterated analogues, in particular those deuterated at the alpha and optionally beta positions, or pharmaceutically acceptable salts of these) may be prepared.

[0011] Viewed from a first aspect, therefore, the invention provides a composition comprising: (a) an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; (b) an α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; and (c) *N,N*-dimethyltryptamine, or a pharmaceutically acceptable salt thereof; wherein the composition comprises 50% or more by weight of the α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof, and the α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof.

[0012] Viewed from a second aspect, the invention provides a composition as defined according to the first aspect for use in therapy.

[0013] Viewed from a third aspect, the invention provides a composition as defined according to the first aspect for use in a method of treating a psychocognitive disorder in a patient.

[0014] Further aspects and embodiments of the present invention will be evident from the discussion that follows below.

BRIEF DESCRIPTION OF THE FIGURES

[0015]

Fig. 1. depicts the predicted pharmacokinetic profile of partially deuterated DMT compared to DMT and fully deuterated DMT. Predicted A) plasma concentration and B) brain tissue concentration, showing the extended half-life of partially deuterated DMT. Hashed area depicts effect site concentrations (>60 ng/mL) that are experienced as full dissociation from the external world, referred to as 'DMT space'.

Fig. 2 plots calculated *in vitro* half-life for DMT and 6 deuterated-containing compositions described in Example 4. A) Linear regression analysis. The r^2 value for half-life is 0.754; where

the slope was found to be significantly different to zero, $p=0.01$. B) Half-life of deuterated analogues as a percent change from (undeuterated) DMT (dashed line).

Fig. 3 *In vitro* intrinsic clearance for DMT and 6 deuterium-containing compositions described in Example 4. A) Linear regression analysis. The r^2 value for intrinsic clearance is 0.7648; where the slope was found to be significantly different to zero, $p=0.01$. B) Intrinsic clearance of deuterated analogues as a percent change from (undeuterated) DMT (dashed line).

DETAILED DESCRIPTION OF THE INVENTION

[0016] Throughout this specification, one or more aspect of the invention may be combined with one or more features described in the specification to define distinct embodiments of the invention.

[0017] References herein to a singular of a noun encompass the plural of the noun, and vice-versa, unless the context implies otherwise.

[0018] Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

[0019] The present invention provides a composition comprising: (a) an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; (b) an α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; and (c) *N,N*-dimethyltryptamine, or a pharmaceutically acceptable salt thereof; wherein the composition comprises 50% or more by weight of the α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof, and the α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof.

[0020] A deuterium atom is a hydrogen atom with an additional neutron.

[0021] Frequently herein, the terms α,α -dideutero-*N,N*-dimethyltryptamine compounds and α -protio, α -deutero-*N,N*-dimethyltryptamine compounds are referred to as deuterated (or fully deuterated) *N,N*-dimethyltryptamine and partially deuterated *N,N*-dimethyltryptamine respectively. A deuterated (or fully deuterated) *N,N*-dimethyltryptamine thus refers strictly to an *N,N*-dimethyltryptamine compound with both protons at the α position substituted with deuterium atoms. The term partially deuterated *N,N*-dimethyltryptamine strictly refers to an *N,N*-dimethyltryptamine compound in which one of the two protons at the α position is substituted with a deuterium atom. A deuterated *N,N*-dimethyltryptamine compound of the present invention is any *N,N*-dimethyltryptamine compound substituted with two deuterium

atoms at the α position, and a partially deuterated *N,N*-dimethyltryptamine compound of the present invention is any *N,N*-dimethyltryptamine compound with one hydrogen atom and one deuterium atom at the α position.


[0022] Accordingly, the invention provides a composition comprising: (a) an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; (b) an α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof; and (c) *N,N*-dimethyltryptamine, or a pharmaceutically acceptable salt thereof; wherein the composition comprises 50% or more by weight of the α,α -dideutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof, and the α -protio, α -deutero-*N,N*-dimethyltryptamine compound, or a pharmaceutically acceptable salt thereof.

[0023] *N,N*-dimethyltryptamine and all its deuterated analogues freely form addition salts with anionic counterions. Throughout the specification, an *N,N*-dimethyltryptamine compound (in particular *N,N*-dimethyltryptamine, α,α -dideutero-*N,N*-dimethyltryptamine compounds and α -protio, α -deutero-*N,N*-dimethyltryptamine compounds) refers equally to any pharmaceutically acceptable salt, e.g. the fumarate salt.

[0024] Typically, acidic reagents may be used to prepare salts, in particular pharmaceutically acceptable salts, of *N,N*-dimethyltryptamine compounds. Examples of suitable acidic reagents are selected from the group consisting of fumaric acid, hydrochloric acid, tartaric acid, citric acid, hydrobromic acid, sulfuric acid, phosphoric acid, acetic acid, maleic acid, lactic acid, tartaric acid and gluconic acid. Often, where in the form of salts, *N,N*-dimethyltryptamine compounds, in particular as the compounds of the invention, in the compositions of the invention or otherwise used according to the various aspects of the present invention, are fumarate hydrochloride, tartrate or citrate salts, in particular fumarate salts.

[0025] The compounds present in the composition of the first aspect of the invention (and other, as appropriate), may thus be present as *N,N*-dimethyltryptamine compounds, in particular *N,N*-dimethyltryptamine or deuterated or partially deuterated *N,N*-dimethyltryptamine compounds as defined herein, in free base or salt form (such as the salts described herein), optionally as solvates (e.g. hydrates) thereof.

[0026] According to particular embodiments of the invention, α,α -dideutero-*N,N*-dimethyltryptamine compounds (fully deuterated compounds) comprise 0, 1 or 2 deuterium atoms at the β position but, other than the presence of 2, 3 or 4 deuterium atoms, are identical to the compound *N,N*-dimethyltryptamine. According to these particular and other particular embodiments of the invention, α -protio, α -deutero-*N,N*-dimethyltryptamine compounds (partially deuterated compounds) may also comprise 0, 1 or 2 deuterium atoms at the β position but, other than the presence of 1, 2 or 3 deuterium atoms, are identical to the compound *N,N*-dimethyltryptamine. The six fully deuterated and partially deuterated compounds according to these particular embodiments of the invention are depicted in the following table:



[0027] The composition of the first aspect comprises 50% or more by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises 60% or more by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises 75% or more by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises up to 90% by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises up to 95% by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises up to 96% by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises up to 97% by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds. In preferred embodiments of the first aspect, the composition comprises up to 98% by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds.

[0028] Accordingly, it will be understood from the foregoing that, according to particular embodiments of the first aspect of the invention, in particular those embodiments discussed in the following eight paragraphs, the composition comprises between 50% and 90%, 50% and 95%, 50% and 96%, 50% and 97%, 50% and 98%; 60% and 90%, 60% and 95%, 60% and 96%, 60% and 97%, 60% and 98%; or 75% and 90%, 75% and 95%, 75% and 96%, 75% and 97%, 75% and 98%, by weight of the deuterated and partially deuterated *N,N*-dimethyltryptamine compounds.

[0029] In embodiments of the first aspect, the composition comprises a combination of *N,N*-dimethyltryptamine, an α -protio, α -deutero-*N,N*-dimethyltryptamine compound and an α,α -dideutero-*N,N*-dimethyltryptamine compound selected from α,α -dideutero-*N,N*-dimethyltryptamine, α,α,β -trideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine, for example a combination of *N,N*-dimethyltryptamine, an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, and α,α -dideutero-*N,N*-dimethyltryptamine, a combination of *N,N*-dimethyltryptamine, an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, and $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine or a combination of *N,N*-dimethyltryptamine, an α -protio, α -deutero-*N,N*-dimethyltryptamine compound, α,α -dideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine.

[0030] In preferred embodiments of the first aspect, the composition comprises a combination of *N,N*-dimethyltryptamine and 2% or more by weight of one or more deuterated *N,N*-dimethyltryptamine compounds selected from α,α -dideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine.

[0031] In other embodiments of the first aspect, the composition comprises a combination of *N,N*-dimethyltryptamine and 2% or more by weight of one or more deuterated *N,N*-dimethyltryptamine compounds selected from α,α -dideutero-*N,N*-dimethyltryptamine, α,α,β -trideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine.

[0032] In preferred embodiments of the first aspect, the composition comprises 2% or more by weight of $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine.

[0033] In preferred embodiments of the first aspect, the composition comprises 2% or more by weight of α,α -dideutero-*N,N*-dimethyltryptamine.

[0034] In certain embodiments of the first aspect, the composition comprises 2% or more by weight of $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine and 2% or more by weight of α,α -dideutero-*N,N*-dimethyltryptamine.

[0035] It will be understood that, within the first aspect of the invention, wherever a composition comprises 2% or more by weight of one or more deuterated *N,N*-dimethyltryptamine compounds, that such compositions may comprise up to 95% by weight of one or more deuterated *N,N*-dimethyltryptamine compounds.

[0036] In preferred embodiments of the first aspect, the composition comprises or further comprises, for example further comprises, one or more partially deuterated *N,N*-dimethyltryptamine compounds selected from α,β,β -trideutero-*N,N*-dimethyltryptamine, α,β -dideutero-*N,N*-dimethyltryptamine, and α -deutero-*N,N*-dimethyltryptamine. Preferably, the partially deuterated *N,N*-dimethyltryptamine compound is α -deutero-*N,N*-dimethyltryptamine.

[0037] In preferred embodiments within the first aspect, the one or more partially deuterated *N,N*-dimethyltryptamine compound comprises up to 50% by weight of the total composition.

[0038] The compositions of the invention comprise *N,N*-dimethyltryptamine.

[0039] According to specific embodiments, the compositions of the present invention consist essentially of the claimed *N,N*-dimethyltryptamine compounds (i.e. compounds selected from *N,N*-dimethyltryptamine and its deuterated analogues as defined in the claims, or pharmaceutically acceptable salts of these). By the composition consisting essentially of said *N,N*-dimethyltryptamine compounds is meant that the composition may comprise additional components (other than said *N,N*-dimethyltryptamine compounds) but that the presence of these additional components will not materially affect the essential characteristics of the composition. In particular, compositions consisting essentially of said *N,N*-dimethyltryptamine compounds will not comprise material amounts of other pharmaceutically active substances (i.e. material amounts of other drug substances).

[0040] Even more typically, the compositions of the present invention are as defined in claim 2.

[0041] The composition of the present invention preferably comprise from 5% to 95% by weight of a deuterated *N,N*-dimethyltryptamine compound selected from α,α -dideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetraeutero-*N,N*-dimethyltryptamine.

[0042] Outside the scope of the present claims, the present disclosure provides a composition obtainable by the reduction of 2-(3-indolyl)-*N,N*-dimethylglyoxalamide with a reducing agent consisting essentially of up to 90 mol% of lithium aluminium hydride and 10 mol% or more of lithium aluminium deuteride; and a composition obtainable by the reduction of 2-(3-indolyl)-*N,N*-dimethylacetamide with a reducing agent consisting essentially of up to 90 mol% of lithium aluminium hydride and 10 mol% or more of lithium aluminium deuteride. The reducing agent may be dissolved or suspended in a liquid medium. Typically, owing to strong reactivity with water and protic solvents such as alcohols, although available in solid (powdered) form, lithium aluminium hydride (or deuteride) are often manipulated in dried, aprotic solvents such as ethers, often under an inert atmosphere. The skilled person is well aware of such precautions and appropriate protocols.

[0043] It will be understood that the present disclosure thus provides a composition obtainable by the reduction of a composition comprising one or more of 2-(3-indolyl)-*N,N*-dimethylglyoxalamide and of 2-(3-indolyl)-*N,N*-dimethylacetamide with a reducing agent consisting essentially of up to 90 molar % of lithium aluminium hydride and 10 molar % or more of lithium aluminium deuteride, optionally dissolved or suspended in a liquid medium.

[0044] Preferably, the reducing agent as described in the immediately preceding paragraph, consists essentially of 85 molar % lithium aluminium hydride and 15 molar % lithium aluminium deuteride, or consists essentially of 80 molar % lithium aluminium hydride and 20 molar % of lithium aluminium deuteride, or consists essentially of 25 molar % lithium aluminium hydride and 75 molar % lithium aluminium deuteride, or consists essentially of 30 molar % lithium aluminium hydride and 70 molar % lithium aluminium deuteride, or consists essentially of 65

molar % lithium aluminium hydride and 35 molar % lithium aluminium deuteride, or consists essentially of 60 molar % lithium aluminium hydride and 40 molar % lithium aluminium deuteride, or consists essentially of 55 molar % lithium aluminium hydride and 45 molar % lithium aluminium deuteride, or consists essentially of 50 molar % lithium aluminium hydride and 50 molar % lithium aluminium deuteride, or consists essentially of 45 molar % lithium aluminium hydride and 55 molar % of lithium aluminium deuteride, or consists essentially of 40 molar % by weight of lithium aluminium hydride and 60 molar % of lithium aluminium deuteride, or consists essentially of 35 molar % of lithium aluminium hydride and 65 molar % lithium aluminium deuteride, or consists essentially of 30 molar % lithium aluminium hydride and 70 molar % lithium aluminium deuteride, or consists essentially of 25 molar % lithium aluminium hydride and 75 molar % lithium aluminium deuteride, or consists essentially of 20 molar % lithium aluminium hydride and 80 molar % lithium aluminium deuteride, or consists essentially of 15 molar % by weight of lithium aluminium.

[0045] According to particular non-claimed aspects, by reciting that the reducing agent consists essentially of lithium aluminium hydride and lithium aluminium deuteride is meant that the reducing agent may comprise additional components but that the presence of these components will not materially affect the essential characteristics of the reducing agent (in particular stability and reductive propensity).

[0046] Outside the scope of the claimed invention, the present disclosure provides a pharmaceutical composition comprising a composition as defined herein, in combination with a pharmaceutically acceptable excipient.

[0047] The pharmaceutical composition comprises a composition as defined herein in combination with one or more pharmaceutically acceptable excipients. Suitable pharmaceutical compositions can be prepared by the skilled person, with examples of pharmaceutically acceptable excipients including but not being limited to those described in Gennaro et. al., Remington: The Science and Practice of Pharmacy, 20th Edition, Lippincott, Williams and Wilkins, 2000 (specifically part 5: pharmaceutical manufacturing). Suitable excipients are also described in the Handbook of Pharmaceutical Excipients, 2nd Edition; Editors A. Wade and P. J. Weller, American Pharmaceutical Association, Washington, The Pharmaceutical Press, London, 1994.

[0048] The pharmaceutical compositions may be compressed into solid dosage units, such as tablets, or be processed into capsules or suppositories. By means of pharmaceutically suitable liquids the compounds can also be prepared in the form of a solution, suspension, emulsion, or as a spray. For making dosage units, including tablets, the use of conventional additives such as fillers, colorants, polymeric binders and the like is contemplated. In general, any pharmaceutically acceptable additive can be used.

[0049] Suitable fillers with which the pharmaceutical compositions can be prepared and administered include lactose, starch, cellulose and derivatives thereof, and the like, or mixtures thereof used in suitable amounts. For parenteral administration, aqueous suspensions, isotonic

saline solutions and sterile injectable solutions may be used, containing pharmaceutically acceptable dispersing agents and/or wetting agents, such as propylene glycol or butylene glycol.

[0050] The present disclosure also provides a pharmaceutical composition, in combination with packaging material suitable for the composition, the packaging material including instructions for the use of the pharmaceutical composition.

[0051] The compositions of the invention are useful in therapy and may be administered to a patient in need thereof. As used herein, the term 'patient' preferably refers to a human patient, but may also refer to a domestic mammal. The term does not encompass laboratory mammals.

[0052] In accordance with the third aspect of the invention, there is provided a composition as defined according to the first aspect for use in a method of treating a psychocognitive disorder in a patient.

[0053] As used herein the term psychocognitive disorder encompasses psychiatric disorders which may be associated with one or more cognitive impairment. As used herein the term 'psychiatric disorder' is a clinically significant behavioural or psychological syndrome or pattern that occurs in an individual and that is associated with present distress (e.g., a painful symptom) or disability (i.e., impairment in one or more important areas of functioning) or with a significantly increased risk of suffering death, pain, disability, or an important loss of freedom.

[0054] Diagnostic criteria for psychiatric and psychocognitive disorders referred to herein are provided in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, (DSM-5).

[0055] As used herein the term 'obsessive-compulsive disorder' is defined by the presence of either obsessions or compulsions, but commonly both. The symptoms can cause significant functional impairment and/or distress. An obsession is defined as an unwanted intrusive thought, image or urge that repeatedly enters the person's mind. Compulsions are repetitive behaviours or mental acts that the person feels driven to perform. Typically obsessive-compulsive disorder (OCD) manifests as one or more obsession which drives adoption of a compulsion. For example, an obsession with germs may drive a compulsion to clean. A compulsion can either be overt and observable by others, such as checking that a door is locked, or a covert mental act that cannot be observed, such as repeating a certain phrase in one's mind.

[0056] As used herein the term 'depressive disorder' includes major depressive disorder, persistent depressive disorder, bipolar disorder, bipolar depression, and depression in terminally ill patients.

[0057] As used herein the term 'major depressive disorder' (MDD, also referred to as major depression or clinical depression) is defined as the presence of five or more of the following

symptoms over a period of two-weeks or more (also referred to herein as a 'major depressive episode'), most of the day, nearly every day:

- depressed mood, such as feeling sad, empty or tearful (in children and teens, depressed mood can appear as constant irritability);
- significantly reduced interest or feeling no pleasure in all or most activities;
- significant weight loss when not dieting, weight gain, or decrease or increase in appetite (in children, failure to gain weight as expected);
- insomnia or increased desire to sleep;
- either restlessness or slowed behaviour that can be observed by others;
- fatigue or loss of energy;
- feelings of worthlessness, or excessive or inappropriate guilt;
- trouble making decisions, or trouble thinking or concentrating;
- recurrent thoughts of death or suicide, or a suicide attempt.

[0058] At least one of the symptoms must be either a depressed mood or a loss of interest or pleasure.

[0059] Persistent depressive disorder, also known as dysthymia, is defined as a patient exhibiting the following two features:

1. A. has depressed mood for most the time almost every day for at least two years. Children and adolescents may have irritable mood, and the time frame is at least one year.
2. B. While depressed, a person experiences at least two of the following symptoms:
 - Either overeating or lack of appetite.
 - Sleeping too much or having difficulty sleeping.
 - Fatigue, lack of energy.
 - Poor self-esteem.
 - Difficulty with concentration or decision making.

[0060] As used herein the term 'treatment resistant depression' describes MDD which fails to achieve an adequate response to an adequate treatment with standard of care therapy.

[0061] As used herein 'bipolar disorder' also known as manic-depressive illness, is a disorder that causes unusual shifts in mood, energy, activity levels, and the ability to carry out day-to-day tasks.

[0062] There are two defined sub-categories of bipolar disorder; all of them involve clear changes in mood, energy, and activity levels. These moods range from periods of extremely "up," elated, and energised behaviour (known as manic episodes, and defined further below)

to very sad, "down," or hopeless periods (known as depressive episodes). Less severe manic periods are known as hypomanic episodes.

[0063] Bipolar I Disorder- defined by manic episodes that last at least 7 days, or by manic symptoms that are so severe that the person needs immediate hospital care. Usually, depressive episodes occur as well, typically lasting at least 2 weeks. Episodes of depression with mixed features (having depression and manic symptoms at the same time) are also possible.

[0064] Bipolar II Disorder- defined by a pattern of depressive episodes and hypomanic episodes, but not the full-blown manic episodes described above.

[0065] As used herein 'bipolar depression' is defined as an individual who is experiencing depressive symptoms with a previous or coexisting episode of manic symptoms, but does not fit the clinical criteria for bipolar disorder.

[0066] As used herein the term 'anxiety disorder' includes generalised anxiety disorder, phobia, panic disorder, social anxiety disorder, and post-traumatic stress disorder.

[0067] 'Generalised anxiety disorder' (GAD) as used herein means a chronic disorder characterised by long-lasting anxiety that is not focused on any one object or situation. Those suffering from GAD experience non-specific persistent fear and worry, and become overly concerned with everyday matters. GAD is characterised by chronic excessive worry accompanied by three or more of the following symptoms: restlessness, fatigue, concentration problems, irritability, muscle tension, and sleep disturbance.

[0068] 'Phobia' is defined as a persistent fear of an object or situation the affected person will go to great lengths to avoid, typically disproportional to the actual danger posed. If the feared object or situation cannot be avoided entirely, the affected person will endure it with marked distress and significant interference in social or occupational activities.

[0069] A patient suffering a from a 'panic disorder' is defined as one who experiences one or more brief attack (also referred to as a panic attack) of intense terror and apprehension, often marked by trembling, shaking, confusion, dizziness, nausea, and/or difficulty breathing. A panic attack is defined as a fear or discomfort that abruptly arises and peaks in less than ten minutes.

[0070] 'Social anxiety disorder' is defined as an intense fear and avoidance of negative public scrutiny, public embarrassment, humiliation, or social interaction. Social anxiety often manifests specific physical symptoms, including blushing, sweating, and difficulty speaking.

[0071] 'Post-traumatic stress disorder' (PTSD) is an anxiety disorder that results from a traumatic experience. Post-traumatic stress can result from an extreme situation, such as combat, natural disaster, rape, hostage situations, child abuse, bullying, or even a serious

accident. Common symptoms include hypervigilance, flashbacks, avoidant behaviours, anxiety, anger and depression.

[0072] As used herein the term 'substance abuse' means a patterned use of a drug in which the user consumes the substance in amounts or with methods which are harmful to themselves or others.

[0073] As used herein the term 'an avolition disorder' refers to a disorder which includes as a symptom the decrease in motivation to initiate and perform self-directed purposeful activities.

[0074] In preferred embodiments of the third aspect of the present invention, the psychocognitive disorder is selected from (i) an obsessive compulsive disorder, (ii) a depressive disorder, (iii) an anxiety disorder, (iv) substance abuse, and (v) an avolition disorder.

[0075] According to particular embodiments of the third aspect of the present invention, the depressive disorder is major depressive disorder. According to still more particular embodiments, the major depressive disorder is treatment-resistant major depressive disorder.



[0076] Compositions comprising partially deuterated *N,N*-dimethyltryptamine compounds of the present invention can be synthesised following the reaction schemes (synthetic schemes) provided in Scheme 1 and Scheme 2 below. The chemistry depicted in the schemes was reported by PE Morris and C Chiao (Journal of Labelled Compounds And Radiopharmaceuticals, Vol. XXXIII, No. 6, 455-465 (1993)). Compositions comprising partially deuterated *N,N*-dimethyltryptamine compounds of the present invention can also be synthesised following the synthetic scheme depicted in Scheme 3 and Example 4 below.

[0077] The relative proportions of *N,N*-dimethyltryptamine against deuterated *N,N*-dimethyltryptamine compounds and partially deuterated *N,N*-dimethyltryptamine compounds may be controlled by varying the ratio of lithium aluminium hydride and lithium aluminium deuteride in the reducing agent. Relative proportions may further be varied by adding one or more of *N,N*-dimethyltryptamine, α,α -dideutero-*N,N*-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetradeutero-*N,N*-dimethyltryptamine to the compositions described hereinabove.



[0078] A particular advantage of the present disclosure is that the reductions described herein allow particularly high purities to be obtained, without the necessity for subsequent chromatographic purification (e.g. column chromatography), thereby increasing the efficiency through which compositions of the invention may be prepared. Moreover, the ability to avoid the use of chromatography in order to achieve high purities makes scale up more efficient and therefore cost-effective.

[0079] Identification of the compositions resultant from the reduction step in Schemes 1 and 2 may be achieved, if desired, by chromatographic separation of the components of the mixtures by conventional means at the disposal of the skilled person in combination with spectroscopic

and/or mass spectrometric analysis.



Scheme 1



Scheme 2

[0080] Alternative compositions of the present invention are obtainable by mixing N,N -dimethyltryptamine, obtainable by Scheme 1 or Scheme 2 when the reducing agent is exclusively lithium aluminium hydride, with a deuterated N,N -dimethyltryptamine compound obtainable from Scheme 1 or Scheme 2 when the reducing agent is exclusively lithium aluminium deuteride.

[0081] The compositions described hereinabove may be further modified by adding one or more deuterated or partially deuterated N,N -dimethyltryptamine compounds. Stocks of such deuterated or partially deuterated N,N -dimethyltryptamine compounds may be obtained, for example, from the chromatographic separation described above.

[0082] Whilst identification of the compositions resultant from the reduction described herein may be achieved by chromatographic separation of the components of the mixtures, in

combination with spectroscopic and/or mass spectrometric analysis, a particular benefit of the present disclosure is that, according to particular embodiments, there may be no necessity to do so. This is because, over and above the purities achievable in accordance with the present disclosure we have as alluded to above recognised that there is a quantifiable relationship between the extent of deuteration (or in other words the quantity or proportion of deuterium in the *N,N*-dimethyltryptamine compounds in the compositions of the present invention) and the metabolic half-life of the resultant composition. The extent of deuteration may be controlled through the amount of deuterium-containing reducing agent used in the method of the disclosure, through which (according to particular embodiments) the compositions of the invention may be obtained, and thus control exercised, in a predictable way, over potentiation of the metabolic half-life of the parent compound (undeuterated *N,N*-dimethyltryptamine).

[0083] In particular, as detailed in Example 4 and related Figures 2 and 3, we have demonstrated that increasing deuterium enrichment at the α -carbon of *N,N*-dimethyltryptamine increases metabolic stability, leading to a decrease in clearance and longer half-life. A linear relationship exists between molecular weight and half-life, in particular when the input reducing agent for production of the deuterium-enriched *N,N*-dimethyltryptamine-containing compositions of this invention comprise LiAlH_4 and LiAlD_4 with ratio between about 1 : 2.5 and about 2.5 : 1.

[0084] Such types of composition constitute specific embodiments of the first aspect of the invention. According to these specific embodiments, the composition consists essentially of a mixture of *N,N*-dimethyltryptamine and both of α -deutero-*NN*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine, the composition optionally being in the form of a pharmaceutically acceptable salt, wherein the mean molecular weight of *N,N*-dimethyltryptamine, α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine present in the composition is from 188.28 to 190.28.

[0085] As used herein, mean molecular weight means the weighted average of molecular weights of the of the *N,N*-dimethyltryptamine and one or both of α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine, as measured by an appropriate mass spectroscopic technique, for example LC-MS SIM (selected-ion monitoring), ignoring any weight contribution by formation of pharmaceutically acceptable salts, where applicable.

[0086] It will be understood that providing compositions with such specific mean molecular weights can be achieved by those skilled in the art through the teachings herein, in particular by adjusting the relative proportions of lithium aluminium hydride and lithium aluminium deuteride in the reductions described herein.

[0087] In this context, by reciting that the composition consists essentially of the mixture of *N,N*-dimethyltryptamine and both of α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine means that the composition may comprise additional components to these but that the presence of such additional components will not materially affect the essential characteristics of the composition. In particular, the composition will not comprise material

quantities of other pharmaceutically active compounds, including other *N,N*-dimethyltryptamine compounds. Thus material quantities of other deuterated *N,N*-dimethyltryptamine compounds, in particular β -deutero-*N,N*-dimethyltryptamine compounds and β,β -dideutero-*N,N*-dimethyltryptamine compounds, such as β -deutero-*N,N*-dimethyltryptamine and β,β -dideutero-*N,N*-dimethyltryptamine and β -deutero-*N,N*-dimethyltryptamine compounds and β,β -dideutero-*N,N*-dimethyltryptamine compounds having respectively one or two deuterium atoms in place of hydrogen atoms at the α position are not present in compositions of such embodiments.

[0088] In other words, and alternatively put, the compositions according to these specific embodiments constitute a drug substance comprising a biologically active ingredient consisting essentially of a mixture of *N,N*-dimethyltryptamine, α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine, wherein the biologically active ingredient has a mean molecular weight from 188.28 to 190.28 and wherein the drug substance is optionally in the form of a pharmaceutically acceptable salt.

[0089] It will be understood that the compositions according to these specific embodiments comprise said α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine in amounts greater than found in isotopically unenriched *N,N*-dimethyltryptamine. It will also be understood that the greater the proportion of α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine in these specific embodiments, the higher the mean molecular weight of the composition.

[0090] According to more specific embodiments, the mean molecular weight of *N,N*-dimethyltryptamine, α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine present in the composition is from 188.9 to 189.7, for example 188.90 to 189.70.

[0091] According to still more specific embodiments described herein, including compositions in which the mean molecular weight of *N,N*-dimethyltryptamine, α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine present in the composition is from 188.9 to 189.7, for example 188.90 to 189.70, the composition optionally is in the form of a pharmaceutically acceptable salt, by which it will be understood that the *N,N*-dimethyltryptamine, α -deutero-*N,N*-dimethyltryptamine and α,α -dideutero-*N,N*-dimethyltryptamine present in the composition are present in pharmaceutically acceptable salt form. Such salts may be as described elsewhere herein and, according to yet more specific embodiments, the composition is in the form of a fumarate salt.

EXAMPLES

Example 1:

[0092] 2-(3-indolyl)-*N,N*-dimethylacetamide is synthesised as described in Morris and Chiao (1993), Journal of labelled compounds and radiopharmaceuticals, Vol. XXXIII, No. 6. 0.063g LiAlH₄ and 0.278g LiAlD₄ (20:80 molar ratio of LiAlH₄:LiAlD₄) are suspended in 25 ml dry Et₂O with magnetic stirring. 0.785g of 2-(3-indolyl)-*N,N*-dimethylacetamide is dissolved in 300mL and added dropwise to the stirred suspension. The reagents are heated under reflux for 3 hours, then cooled in an ice bath and quenched dropwise with water. The resulting mixture is filtered, dried, and the solvents removed under vacuum, to yield approx. 0.5g of a composition consisting of 3 molar % *N,N*-dimethyltryptamine, 28 molar % α -deutero-*N,N*-dimethyltryptamine and 69 molar % of α,α -dideutero-*N,N*-dimethyltryptamine.

Example 2:

[0093] 0.173g LiAlH₄ and 0.156g LiAlD₄ (55:45 molar ratio of LiAlH₄:LiAlD₄) are suspended in 25 ml dry Et₂O with magnetic stirring. 0.785g of 2-(3-indolyl)-*N,N*-dimethylacetamide is dissolved in 300mL and added dropwise to the stirred suspension. The reagents are heated under reflux for 3 hours, then cooled in an ice bath and quenched dropwise with water. The resulting mixture is filtered, dried, and the solvents removed under vacuum, to yield approx. 0.5g of a composition consisting of 24 molar % *N,N*-dimethyltryptamine, 50 molar % α -deutero-*N,N*-dimethyltryptamine and 26 molar % of α,α -dideutero-*N,N*-dimethyltryptamine.

Reference Example 3:


[0094] 0.293g LiAlH₄ and 0.035g LiAlD₄ (90:10 molar ratio of LiAlH₄:LiAlD₄) are suspended in 25 ml dry Et₂O with magnetic stirring. 0.785g of 2-(3-indolyl)-*N,N*-dimethylacetamide is dissolved in 300mL and added dropwise to the stirred suspension. The reagents are heated under reflux for 3 hours, then cooled in an ice bath and quenched dropwise with water. The resulting mixture is filtered, dried, and the solvents removed under vacuum, to yield approx. 0.5g of a composition consisting of 76 molar % *N,N*-dimethyltryptamine, 22 molar % α -deutero-*N,N*-dimethyltryptamine and 2 molar % of α,α -dideutero-*N,N*-dimethyltryptamine.

Example 4: use of human hepatocytes to assess the *in vitro* intrinsic clearance of deuterated DMT analogue blends relative to DMT

[0095] *In vitro* determination of intrinsic clearance is a valuable model for predicting *in vivo* hepatic clearance. The liver is the main organ of drug metabolism in the body, containing both phase I and phase II drug metabolising enzymes, which are present in the intact cell.

Synthesis of samples

[0096] N,N-DMT 220.9 g (as free base) was prepared as N,N-DMT fumarate, using the chemistry depicted in Scheme 3. An additional 4-6 g of six partially deuterated mixtures were also produced using modified conditions.



Scheme 3

Synthesis of DMT

Stage 1: coupling of indole-3-acetic acid and dimethylamine

[0097] To a 5 L vessel under N₂ was charged indole-3-acetic acid (257.0 g, 1.467 mol), hydroxybenzotriazole (HOBt, ~20% wet) (297.3 g, 1.760 mol) and dichloromethane (2313mL) to give a milky white suspension. 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC.HCl, 337.5 g, 1.760 mol) was then charged portion-wise over 5 minutes at 16-22 °C. The reaction mixture was stirred for 2 hours at ambient temperature before 2M dimethylamine in THF (1100 mL, 2.200 mol) was charged dropwise over 20 minutes at 20-30 °C. The resultant solution was stirred at ambient temperature for 1 hour where HPLC indicated 1.1% indole-3-acetic acid and 98.1% target product referred to as Stage 1). The reaction mixture was then charged with 10% K₂CO₃ (1285 mL) and stirred for 5 minutes. The layers were separated, and the upper aqueous layer extracted with dichloromethane (643 mL x 2). The organic extracts were combined and washed with saturated brine (643 mL). The organic extracts were then dried over MgSO₄, filtered and concentrated *in vacuo* at 45 °C. This provided 303.1 g of crude Stage 1 as an off-white sticky solid. The crude material was then subjected to a slurry in *tert*-butyl methyl ether (TBME, 2570 mL) at 50 °C for 2 hours before

being cooled to ambient temperature, filtered and washed with TBME (514 mL x 2). The filter-cake was then dried *in vacuo* at 50 °C to afford Stage 1 266.2 g (yield=90%) as an off-white solid in a purity of 98.5 % by HPLC and >95 % by NMR.

Stage 2: preparation of DMT

[0098] To a 5 L vessel under N₂ was charged Stage 1 (272.5 g, 1.347 mol) and tetrahydrofuran (THF, 1363mL) to give an off-white suspension. 2.4 M LiAlH₄ in THF (505.3 mL, 1.213mol) was then charged dropwise over 35 minutes at 20-56 °C to give an amber solution. The solution was heated to 60 °C for 2 hours where HPLC indicated Stage 1 ND, target product bracket referred to as Stage 2, 92.5%), Impurity 1 (2.6%), Impurity 2 (1.9%). The complete reaction mixture was cooled to ambient temperature and then charged to a solution of 25% Rochelle's salts (aq) (2725 mL) dropwise over 30 minutes at 20-30 °C. The resultant milky white suspension was allowed to stir at 20-25 °C for 1 hour after which the layers were separated and the upper organic layer washed with saturated brine (681 mL). The organic layer was then dried over MgSO₄, filtered and concentrated *in vacuo* at 45 °C. The resultant crude oil was subjected to an azeotrope from ethanol (545mL x 2). This provided 234.6g (yield=92%) of Stage 2 in a purity of 95.0% by HPLC and >95% by NMR.

Stage 3a (i)-(iii): preparation of seed crystals of DMT fumarate

[0099]

1. (i) Stage 2 (100 mg) was taken up in 8 volumes of isopropyl acetate and warmed to 50 °C before charging fumaric acid (1 equivalent) as a solution in ethanol. The flask was then allowed to mature at 50 °C for 1 hour before cooling to room temperature and stirring overnight, resulting in a white suspension. The solids were isolated by filtration and dried for 4 hours at 50 °C to provide 161 mg of product (> 99% yield). Purity by HPLC was determined to be 99.5% and by NMR to be > 95%.
2. (ii) Substitution of isopropyl acetate for isopropyl alcohol in method (i) afforded a white suspension after stirring overnight. The solids were isolated by filtration and dried for 4 hours at 50 °C to provide 168 mg of product (> 99% yield). Purity by HPLC was determined to be 99.8% and by NMR to be > 95%.

[0100] Substitution of isopropyl acetate for tetrahydrofuran in method (i) afforded a white suspension after stirring overnight. The solids were isolated by filtration and dried for 4 hours at 50 °C to provide 161 mg of product (> 99% yield). Purity by HPLC was determined to be 99.4% and by NMR to be > 95%.

[0101] Analysis by x-ray powder diffraction, showed the products of each of methods 9i) to (iii)

to be the same, which was labelled Pattern A.

Stage 3b: preparation of DMT fumarate

[0102] To a 5 L flange flask under N₂ was charged fumaric acid (152.7 g, 1.315 mol) and Stage 2 (248.2 g, 1.315 mol) as a solution in ethanol (2928 mL). The mixture was heated to 75 °C to give a dark brown solution. The solution was polish filtered into a preheated (80 °C) 5 L jacketed vessel. The solution was then cooled to 70 °C and seeded with Pattern A (0.1 wt%), the seed was allowed to mature for 30 minutes before cooling to 0 °C at a rate of 5 °C/hour. After stirring for an additional 4 hours at 0 °C, the batch was filtered and washed with cold ethanol (496 mL x 2) and then dried at 50 °C overnight. This provided 312.4 g (yield=78%) of Stage 3 in a purity of 99.9% by HPLC and >95% by NMR. XRPD: Pattern A.

Synthesis of deuterated mixtures of DMT compounds

[0103] A modified synthesis at stage 2 using solid LiAlH₄/LiAlD₄ mixtures was adopted, using 1.8 equivalents of LiAlH₄/LiAlD₄ versus 0.9 equivalents using the process described above for undeuterated DMT.

[0104] Six deuteration reactions were performed.

Representative synthesis of a deuterated mixture (using 1:1 LiAlH₄ : LiAlD₄) of DMT compounds

[0105] To a 250 mL 3-neck flask under N₂ was charged LiAlH₄ (1.013 g, 26.7 mmol), LiAlD₄ (1.120 g, 26.7 mmol) and THF (100 mL). The resultant suspension was stirred for 30 minutes before stage 1 (6 g, 29.666 mmol) was charged portion-wise over 15 minutes at 20-40 °C. The reaction mixture was then heated to reflux (66 °C) for 2 hours where HPLC indicated no stage 1 remained. The mixture was cooled to 0 °C and quenched with 25% Rochelle's salts (aq) (120 mL) over 30 minutes at <30 °C. The resultant milky suspension was stirred for 1 hour and then allowed to separate. The lower aqueous layer was removed and the upper organic layer washed with saturated brine (30mL). The organics were then dried over MgSO₄, filtered and concentrated *in vacuo*. This provided 4.3 g of crude material. The crude was then taken up in ethanol (52 mL) and charged with fumaric acid (2.66 g, 22.917 mmol) before heating to 75 °C. The resultant solution was allowed to cool to ambient temperature overnight before further cooling to 0-5 °C for 1 hour. The solids were isolated by filtration and washed with cold ethanol (6.5 mL x 2). The filtercake was dried at 50 °C overnight to provided 5.7 g (yield=63%) of product in a purity of 99.9% by HPLC and >95% by NMR.

Assessment of extents of deuteration

[0106] This was achieved by LCMS-SIM (SIM = single ion monitoring), the analysis giving a separate ion count for each mass for the three deuterated *N,N*-dimethyltryptamine compounds (*N,N*-dimethyltryptamine (D0), α -deutero-*N,N*-dimethyltryptamine (D1) and α,α -dideutero-*N,N*-dimethyltryptamine (D2)) at the retention time for *N,N*-dimethyltryptamine. The percentage of each component was then calculated from these ion counts.

[0107] For example, %D0 = $[D0/(D0 + D1 + D2)] \times 100$.

HPLC Parameters

[0108]

System:	Agilent 1100/1200 series liquid chromatograph or equivalent		
Column:	Triart Phenyl; 150 x 4.6mm, 3.0 μ m particle size (Ex: YMC, Part number: TPH12S03-1546PTH)		
Mobile phase A:	Water: Trifluoroacetic acid (100:0.05%)		
Mobile phase B:	Acetonitrile : Trifluoroacetic acid (100:0.05%)		
Gradient:	Time	%A	%B
	0	95	5
	13	62	38
	26	5	95
	30.5	5	95
	31	95	5
Flow rate:	1.0 mL/min		
Stop time:	31 minutes	Post runtime:	4 minutes
Injection volume:	5 μ L	Wash vial:	N/A
Column temperature:	30°C combined		
Wavelength:	200 nm, (4 nm)	Reference:	N/A

Mass spectrometry parameters

[0109]

System:		Agilent 6100 series Quadrupole LC-MS or equivalent				
Drying gas flow:		12.0 L/min		Drying gas temp.:		350°C
Nebuliser pressure:		35 psig				
Fragmentor:		110		Gain:		1.00
Cpd	RT	RRT	Conc	Diluent	Detection	Mass
D0	10.64	1.00	0.30 mg/ml	CH ₃ CN:H ₂ O (50:50)	(+) SIM	189.10 m/z
D1	10.64	1.00	0.30 mg/ml	CH ₃ CN:H ₂ O (50:50)	(+) SIM	190.10 m/z
D2	10.64	1.00	0.30 mg/ml	CH ₃ CN:H ₂ O (50:50)	(+) SIM	191.10 m/z
MS-SIM range is the target mass ± 0.1 m/z						

[0110] The data for the six deuterated reactions are tabulated in Table 1 below:

Mixture No. (LiAlH ₄ :LiAlD ₄ ratio)	Input (stage 1)	Output stage 3 (yield)	Purity by HPLC	Purity by NMR	Deuteration %		
					Do	D ₁	D ₂
1 (SPL026) (0:1)	5g	5.3g (65%)	99.7%	>95%	0.7%	2.7%	96.6%
2 (1:1)	6g	5.699g (63%)	99.9%	>95%	30.0%	48.3%	21.7%
3 (1:2)	5g	4.206g (52%)	99.9%	>95%	16.5%	46.8%	36.8%
4 (1:3)	5g	5.558g (68%)	99.8%	>95%	9.3%	41.5%	49.2%
5 (2:1)	5g	4.218g (52%)	99.9%	>95%	47.5%	41.3%	11.2%
*6 (3:1)	5g	5.0g (62%)	99.4%	>95%	57.5%	35.3%	7.4%
* reference compound							

***In vitro* intrinsic clearance of DMT (SPL026) and 6 deuterated compound blends**

[0111] *In vitro* determination of intrinsic clearance is a valuable model for predicting *in vivo* hepatic clearance. The liver is the main organ of drug metabolism in the body, containing both phase I and phase II drug metabolising enzymes, which are present in the intact cell.

Aim

[0112] To use human hepatocytes to assess the *in vitro* intrinsic clearance of deuterated DMT analogue blends relative to DMT.

Description of the experiment

[0113] Human (mixed gender) hepatocytes pooled from 10 donors (0.545 million cells/mL) were used to investigate the *in vitro* intrinsic clearance of DMT and 6 deuterated analogues.

[0114] A concentration of 5 μ M was used for all test compounds, as well as sumatriptan, serotonin, benzylamine controls. This concentration was chosen in order to maximise the signal-to-noise ratio, while remaining under the Michaelis constant (K_m) for the monoamine oxidase enzyme (MAO). Diltiazem and diclofenac controls were used at a laboratory-validated concentration of 1 μ M.

[0115] Test compounds were mixed with the hepatocyte suspension within a 96-well plate and incubated for up to 60 minutes at 37 °C. The suspension was continuously agitated. At 7 time points, small aliquots were withdrawn, and the test compound/blend concentration therein was measured by LC-MS/MS. The time points measured were 2, 4, 8, 15, 30, 45 and 60 minutes.

[0116] The following LC-MS/MS conditions were used for the analysis:

Instrument:	Thermo TSQ Quantiva with Thermo Vanquish UPLC system
Column:	Luna Omega 2.1x50 mm 2.6 μ m
Solvent A:	H ₂ O + 0.1 % formic acid
Solvent B:	Acetonitrile + 0.1 % formic acid
Flow rate:	0.8 ml/min
Injection vol:	1 μ l
Column temp:	65 °C

[0117] Gradient:

Time (mins)	% Solvent B
0.00	5.0
0.90	75.0
1.36	99.0
1.36	5.0
1.80	5.0

[0118] MS parameters:

Positive ion spray voltage:	4000 V
Vaporiser temperature:	450 °C
Ion transfer tube temp:	365 °C
Sheath gas:	54
Aux gas:	17
Sweep gas:	1
Dwell time	8 ms

[0119] MRM transitions:

- D0 = mass to charge ratio 189.14 > 58.16.
- D1 = mass to charge ratio 190.14 > 59.17.
- D2 = mass to charge ratio 191.14 > 60.17.

[0120] The MRM transitions were determined from a preliminary analysis of DMT samples containing either no deuterium (for D0 transition), or high levels of either D1 or D2 deuteration (for the D1 and D2 transitions respectively).

[0121] The resulting concentration-time profile was then used to calculate intrinsic clearance (CL_{int}) and half-life (t_{1/2}). To do this, the MS peak area or MS peak area/IS response of each analyte is plotted on a natural log scale on the y axis versus time (min) of sampling on the X axis. The slope of this line is the elimination rate constant. This is converted to a half-life by -ln(2)/slope. Intrinsic clearance is calculated from the slope/elimination rate constant and the formula is CL_{int} = (-1000*slope)/cell density in 1E6 cells/ml, to give units of microlitre/min/million cells.

Results

[0122] Intrinsic clearance and half-life values were calculated for DMT and the 6 deuterated mixtures described above. These data were weighted dependent on the ratio of D0, D1 and D2 to give an overall intrinsic clearance and half-life value for each compound blend (Table 2).

Table 2 *In vitro* intrinsic clearance and calculated half-life of DMT and 6 deuterated mixtures

Compound name or Mixture No (per Table 1)	LiAlH ₄ :LiAlD ₄ input ratio	D ₀ : D ₁ : D ₂ output ratio	Molecular weight	Intrinsic clearance (μL/min/million cells)	Half-life (min)
DMT (SPL026)	1:0	100:0:0	188.269	13.77	92.39
1	0:1	0.7 : 2.7 : 96.6	190.240	7.15	178.79
2	1:1	30.0:48.3:21.7	189.192	10.46	125.80
3	1:2	16.5 : 46.8 : 36.8	189.669	9.36	140.43
4	1:3	9.3 : 41.5 : 49.2	189.676	11.14	116.84
5	2:1	47.5:41.3:11.2	188.910	10.99	119.61
*6	3:1	57.4 : 35.3 : 7.4	188.961	13.64	95.04

* reference compound

[0123] Data were fitted with a linear model using regression analysis, which revealed that deuterium enrichment at the α-carbon of DMT decreases intrinsic clearance linearly with increasing molecular weight (MW), therefore enabling manufacture of DMT drug substances with half-lives which can be accurately predicted in the range identified.

[0124] Mixture 1, which contains 96.6% D₂-DMT, sees the biggest change, with the intrinsic clearance rate almost halved compared to undeuterated-DMT (Fig. 3), nearly doubling the half-life (Fig. 2). Intermediate blends of deuteration (Mixtures 2 to 5) decreased intrinsic clearance in a manner correlated with molecular weight (Fig. 3).

Conclusion

[0125] These data demonstrate that increasing deuterium enrichment at the α-carbon of DMT increases metabolic stability, leading to a decrease in clearance and longer half-life. A linear relationship exists between MW and half-life, in particular when the input reducing agent for production of the deuterium enriched DMT-containing drug substance by methods of the present disclosure comprise LiAlH₄ and LiAlD₄ with ratio between 1 : 2.5 and 2.5: 1.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2019081764A1](#) [0006]

Non-patent literature cited in the description

- **CARHART-HARRISGOODWIN**The Therapeutic Potential of Psychedelic Drugs: Past, Present and Future *Neuropsychopharmacology*, 2017, vol. 42, 2105-2113 [0003]
- **GRIFFITHS et al.**Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: a randomised double-blind trial *Journal of Psychopharmacology*, 2016, vol. 30, 121181-1197 [0003]
- **GALLIMORESTRASSMAN**A model for the application of target-controlled intravenous infusion for a prolonged immersive DMT psychedelic experience *Frontiers in Pharmacology*, 2016, vol. 7, 211- [0004]
- **STRASSMAN et al.**Dose response study of N,N-dimethyltryptamine in humans *Arch Gen Psychiatry*, 1994, vol. 51, 85- [0004]
- **BARKER et al.**Comparison of the brain levels of N,N-dimethyltryptamine and $\alpha,\alpha,\beta,\beta$ -tetra deuterio-N,N-dimethyltryptamine following intraperitoneal injection *Biochemical Pharmacology*, 1982, vol. 31, 152513-2516 [0005]
- **BEATON et al.** *Pharmacol. Biochem. Behav.*, 1982, vol. 16, 811-814 [0007]
- **GENNARO**Remington: The Science and Practice of Pharmacy *Lippincott, Williams and Wilkins* 20000000 [0047]
- Handbook of Pharmaceutical Excipients *American Pharmaceutical Association, Washington, The Pharmaceutical Press* 19940000 [0047]
- Diagnostic and Statistical Manual of Mental Disorders [0054]
- **PE MORRIS CHIAO** *Journal of Labelled Compounds And Radiopharmaceuticals*, 1993, vol. XXXIII, 6455-465 [0076]
- **MORRIS CHIAO** *Journal of labelled compounds and radiopharmaceuticals*, 1993, vol. XXXIII, 6 [0092]

Patentkrav

1. Sammensætning, der omfatter: (a) en α,α -dideutero-*N,N*-dimethyltryptaminforbindelse, eller et farmaceutisk acceptabelt salt deraf; (b) en α -protio-, α -deutero-*N,N*-dimethyltryptaminforbindelse, eller et farmaceutisk acceptabelt salt deraf; og (c) *N,N*-dimethyltryptamin, eller et farmaceutisk acceptabelt salt deraf;
- 5 hvor sammensætningen omfatter 50 % eller mere efter α,α -dideutero-*N,N*-dimethyltryptaminforbindelsens vægt, eller et farmaceutisk acceptabelt salt deraf, og α -protio-, α -deutero-*N,N*-dimethyltryptaminforbindelsen, eller et farmaceutisk acceptabelt salt deraf.
- 10
2. Sammensætning ifølge krav 1, der navnlig består af:
- (a) α,α -dideutero-*N,N*-dimethyltryptamin, eller et farmaceutisk acceptabelt salt deraf;
- (b) α -protio-, α -deutero-*N,N*-dimethyltryptamin, eller et farmaceutisk acceptabelt salt deraf; og
- 15 (c) *N,N*-dimethyltryptamin, eller et farmaceutisk acceptabelt salt deraf;
- hvor middelmolekylvægten af α -protio-, α -deutero-*N,N*-dimethyltryptamin, α,α -dideutero-*N,N*-dimethyltryptamin, og *N,N*-dimethyltryptamin, til stede i sammensætningen er fra 188,90 til 190,28.
- 20
3. Sammensætning ifølge krav 1 eller krav 2, der omfatter 60 % eller mere efter α,α -dideutero-*N,N*-dimethyltryptaminforbindelsens vægt, eller et farmaceutisk acceptabelt salt deraf, og α -protio-, α -deutero-*N,N*-dimethyltryptaminforbindelsen eller et farmaceutisk acceptabelt salt deraf.
- 25
4. Sammensætning ifølge et hvilket som helst af kravene 1 til 3, hvor α,α -dideutero-*N,N*-dimethyltryptaminforbindelsen er α,α -dideutero-*N,N*-dimethyltryptamin, eller et farmaceutisk acceptabelt salt deraf.
- 30
5. Sammensætning ifølge et hvilket som helst af kravene 1, 3 og 4 omfattende en α -protio-, α -deutero-*N,N*-dimethyltryptaminforbindelse valgt fra gruppen bestående af α,β,β -trideutero-*N,N*-dimethyltryptamin-, α,β -dideutero-*N,N*-dimethyltryptamin, og α -protio-, α -deutero-*N,N*-dimethyltryptamin, og farmaceutisk acceptable salte deraf.
6. Sammensætning ifølge et hvilket som helst af kravene 1 til 5, der omfatter α -protio-, α -

deutero-*N,N*-dimethyltryptamin, eller et farmaceutisk acceptabelt salt deraf.

7. Sammensætning ifølge et hvilket som helst af kravene 1 til 6 med en renhed, der er større end eller svarer til 99 % ved HPLC, fortrinsvis en renhed, der er større end eller svarer til 99,9 % ved hjælp af HPLC.
8. Sammensætning ifølge et hvilket som helst af kravene 1 til 7, hvor α -protio-, α -deutero-*N,N*-dimethyltryptamin- og α,α -dideutero-*N,N*-dimethyltryptaminforbindelserne, og *N,N*-dimethyltryptamin, er i form af et farmaceutisk acceptabelt salt, fortrinsvis hvor det farmaceutisk acceptable salt er et fumaratsalt.
9. Sammensætning ifølge et hvilket som helst af kravene 1 til 7, hvor α -protio-, α -deutero-*N,N*-dimethyltryptamin- og α,α -dideutero-*N,N*-dimethyltryptaminforbindelserne, og *N,N*-dimethyltryptamin, er i form af et fumarat salt.
10. Sammensætning ifølge et hvilket som helst af kravene 1 til 9 til anvendelse i terapi.
11. Sammensætning ifølge et hvilket som helst af kravene 1 til 9 til anvendelse i behandling af en psykokognitiv lidelse hos en patient.
12. Sammensætning til anvendelse ifølge krav 11, hvor den psykokognitive lidelse vælges fra gruppen bestående af (i) en obsessiv-kompulsiv lidelse, (ii) en depressiv lidelse, (iii) en angstlidelse, (iv) stofmisbrug og (v) en avolitionslidelse.
13. Sammensætning til anvendelse ifølge krav 12, hvor lidelsen er svær depressiv lidelse, fortrinsvis hvor lidelsen er behandlingsresistent depression.
14. Sammensætning ifølge et hvilket som helst af kravene 1 til 9 til anvendelse i dimethyltryptamin-assisteret psykoterapi.

DRAWINGS




Fig. 1




Fig. 2




Fig. 3